



# ODE classification


Nasser M. Abbasi

May 7, 2017    compiled on — Sunday May 07, 2017 at 12:34 AM



system of first\_order ODE's





# 1 examples

---

## 1.1 second order, constant coeff.

### 1.1.1 second order, constant coeff. homogeneous

second order, constant coeff. homogeneous, one root repeated

$$y'' - 2y' + 1 = 0$$

Let  $y = Ae^{rx}$  and plug into the above and simplify, we obtain the characteristic equation

$$\begin{aligned}r^2 - 2r + 1 &= 0 \\(r - 1)^2 &= 0 \\r &= 1\end{aligned}$$

Repeated root. Hence the two L.I. basis solutions are

$$\begin{aligned}y_1 &= e^x \\y_2 &= xe^x\end{aligned}$$

And the homogeneous solution is

$$\begin{aligned}y_h &= c_1y_1 + c_2y_2 \\&= c_1e^x + c_2xe^x\end{aligned}$$

second order, constant coeff. homogeneous, two real distinct roots

$$y'' + y' - 2y = 0$$

Let  $y = Ae^{rx}$  and plug into the above and simplify, we obtain the characteristic equation

$$\begin{aligned}r^2 + r - 2 &= 0 \\(r - 1)(r + 2) &= 0 \\r_1 &= 1 \\r_2 &= -2\end{aligned}$$

Hence the L.I. basis solutions are

$$\begin{aligned}y_1 &= e^x \\y_2 &= e^{-2x}\end{aligned}$$

And the homogeneous solution is

$$\begin{aligned}y_h &= c_1y_1 + c_2y_2 \\&= c_1e^x + c_2e^{-2x}\end{aligned}$$

**second order, constant coeff. homogeneous, two complex conjugate roots**

$$y'' - 6y' + 13y = 0$$

Let  $y = Ae^{rx}$  and plug into the above and simplify, we obtain the characteristic equation

$$r^2 - 6r + 13 = 0$$

Whose roots are

$$r_1 = 3 + 2i$$

$$r_2 = 3 - 2i$$

Hence the L.I. basis solutions are

$$y_1 = e^{(3+2i)x}$$

$$y_2 = e^{(3-2i)x}$$

the homogeneous solution is

$$\begin{aligned} y_h &= c_1 y_1 + c_2 y_2 \\ &= c_1 e^{(3+2i)x} + c_2 e^{(3-2i)x} \end{aligned}$$

This can be converted to real basis using Euler relation which results in

$$\begin{aligned} y_h &= C_1 e^{3x} \cos 2x + C_2 e^3 \sin 2x \\ &= e^{3x} (C_1 \cos 2x + C_2 \sin 2x) \end{aligned}$$

### 1.1.2