

# My Spring 2008 courses at California State University, Fullerton

Nasser M. Abbasi

spring 2008      Compiled on October 14, 2025 at 5:14pm      [public]

math 504 simulation

Also audited few lecture in physics 555A. Here is one problem I did there

## 1 Problem

Show that the recurrence formula

$$C_q = -\frac{2(k-q)}{q(q+2l+1)(k+l)}C_{q-1} \quad (1)$$

can be written as

$$C_q = (-1)^q \left( \frac{2}{k+l} \right)^q \frac{(k-1)!}{(k-q-1)!} \frac{(2l+1)!}{q!(q+2l+1)!} C_0 \quad (2)$$

## 2 Solution

Proof by induction on  $q$ . For  $q = 1$ , equation (1) becomes

$$C_1 = -\frac{2(k-1)}{(2l+2)(k+l)}C_0$$

and equation (2) becomes

$$\begin{aligned} C_1 &= (-1) \left( \frac{2}{k+l} \right) \frac{(k-1)!}{(k-2)!} \frac{(2l+1)!}{(2l+2)!} C_0 \\ &= -\frac{2(k-1)}{(k+l)(2l+2)!} C_0 \end{aligned}$$

Hence it is true for  $q = 1$ . Now *assume* it is true for  $q = n$ , in otherwords, assume that

$$C_n = -\frac{2(k-n)}{n(n+2l+1)(k+l)}C_{n-1} \quad (3)$$

implies

$$C_n = (-1)^n \left( \frac{2}{k+l} \right)^n \frac{(k-1)!}{(k-n-1)!} \frac{(2l+1)!}{n!(n+2l+1)!} C_0 \quad (4)$$

Now for the induction step. we need to show that it is true for  $n+1$ , i.e. given (4) is true, we need to show that, by replacing  $n$  by  $n+1$  in the above, that

$$C_{n+1} = -\frac{2(k-(n+1))}{(n+1)((n+1)+2l+1)(k+l)}C_n \quad (5)$$

implies

$$\begin{aligned}
C_{n+1} &= (-1)^{n+1} \left( \frac{2}{k+l} \right)^{n+1} \frac{(k-1)!}{(k-(n+1)-1)!} \frac{(2l+1)!}{(n+1)! ((n+1)+2l+1)!} C_0 \\
&= (-1)^{n+1} \left( \frac{2}{k+l} \right)^{n+1} \frac{(k-1)!}{(k-n-2)!} \frac{(2l+1)!}{(n+1)! (n+2l+2)!} C_0
\end{aligned} \tag{6}$$

We start with (5), and replace the  $C_n$  term with what we assumed to be true from (4), hence (5) can be rewritten as

$$C_{n+1} = -\frac{2(k-(n+1))}{(n+1)((n+1)+2l+1)(k+l)} \overbrace{\left[ (-1)^n \left( \frac{2}{k+l} \right)^n \frac{(k-1)!}{(k-n-1)!} \frac{(2l+1)!}{n! (n+2l+1)!} C_0 \right]}^{C_n \text{ from (4)}}$$

Simplify the above leads to

$$C_{n+1} = (-1)^{n+1} \left( \frac{2}{k+l} \right)^{n+1} \frac{(k-1)!}{(k-n-2)!} \frac{(2l+1)!}{n! (n+2l+2)!} C_0$$

Which is (6). Therefore, the relationship is true for any  $n$ . QED