

Computer algebra independent integration tests

1-Algebraic-functions/1.1-Binomial-products/1.1.2-Quadratic/1.1.2.5-
 $a+b-x^2-p-c+d-x^2-q-e+f-x^2-r$

Nasser M. Abbasi

September 20, 2021

Compiled on September 20, 2021 at 2:22am

Contents

1	Introduction	3
2	detailed summary tables of results	17
3	Listing of integrals	29
4	Appendix	137

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Performance	8
1.4	list of integrals that has no closed form antiderivative	10
1.5	list of integrals solved by CAS but has no known antiderivative	11
1.6	list of integrals solved by CAS but failed verification	12
1.7	Timing	12
1.8	Verification	13
1.9	Important notes about some of the results	13
1.10	Design of the test system	15

This report gives the result of running the computer algebra independent integration test. Elementary Algebraic integrals version.

The download section below contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [30]. This is test number [9].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3.1 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.3.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.45 on Linux. (via sagemath 9.3)
5. Fricas 1.3.7 on Linux (via sagemath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sagemath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)
9. IntegrateAlgebraic under Mathematica 12.3.1 on windows 10. https://github.com/stblake/algebraic_integration. September 15, 2021 version.

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (30)	0.00 (0)
Mathematica	100.00 (30)	0.00 (0)
Maple	100.00 (30)	0.00 (0)
Fricas	93.33 (28)	6.67 (2)
Giac	86.67 (26)	13.33 (4)
Mupad	73.33 (22)	26.67 (8)
Maxima	73.33 (22)	26.67 (8)
Sympy	70.00 (21)	% 30.00 (9)
IntegrateAlgebraic	26.67 (8)	73.33 (22)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	96.67	0.00	3.33	0.00
Giac	80.00	6.67	0.00	13.33
Fricas	73.33	20.00	0.00	6.67
Maxima	73.33	0.00	0.00	26.67
Maple	56.67	43.33	0.00	0.00
Sympy	40.00	30.00	0.00	30.00
IntegrateAlgebraic	20.00	6.67	0.00	73.33
Mupad	N/A	73.33	0.00	26.67

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	0	0.00 %	0.00 %	0.00 %
Fricas	2	0.00 %	100.00 %	0.00 %
IntegrateAlgebraic	22	100.00 %	0.00 %	0.00 %
Giac	4	0.00 %	0.00 %	100.00 %
Maxima	8	100.00 %	0.00 %	0.00 %
Sympy	9	88.89 %	11.11 %	0.00 %
Mupad	8	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS


1.3 Performance


The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.19	170.77	1.00	161.00	1.00
Mathematica	0.23	170.73	1.01	154.00	1.00
Maple	0.01	520.63	3.36	302.50	1.61
Maxima	1.28	199.55	1.09	180.50	1.09
Fricas	2.49	550.32	3.18	440.00	3.16
Sympy	4.91	325.67	1.83	304.00	1.76
Giac	1.46	228.58	1.34	198.50	1.28
Mupad	0.52	220.00	1.19	192.50	1.09
IntegrateAlgebraic	1.20	464.50	2.77	163.00	1.27

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used columns from the above table.





1.4 list of integrals that has no closed form antiderivative

{

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

IntegrateAlgebraic {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {29}

IntegrateAlgebraic {}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using sagemath (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be indentified by looking at the output of the integration in each section for Maxima. The exception message will indicate of the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/XCAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error Exception raised: NotImplementedError

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the builtin system function LeafSize is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user slelievre at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^{2/2}$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
The following field present only in Rubi and Mathematica Tables
13. integer. 1 if result was verified or 0 if not verified.
The following fields present only in Rubi Tables
14. integer. Number of rules used.
15. integer. Integrand leaf size.
16. real number. Ratio of field 14 over field 15
17. integer. 1 if result was verified or 0 if not verified.
18. String of form "{n,n,...}" which is list of the rules used by Rubi

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	18
2.2	Detailed conclusion table per each integral for all CAS systems	21
2.3	Detailed conclusion table specific for Rubi results	27

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	19
2.1.2	Mathematica	19
2.1.3	Maple	19
2.1.4	Maxima	19
2.1.5	FriCAS	19
2.1.6	Sympy	19
2.1.7	Giac	20
2.1.8	Mupad	20
2.1.9	IntegrateAlgebraic	20

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 30 }

B grade: { }

C grade: { 29 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 15, 16, 17, 18, 19 }

B grade: { 13, 14, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 }

C grade: { }

F grade: { }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 }

B grade: { }

C grade: { }

F grade: { 23, 24, 25, 26, 27, 28, 29, 30 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 23, 24, 25 }

B grade: { 8, 15, 22, 26, 27, 30 }

C grade: { }

F grade: { 28, 29 }

2.1.6 Sympy

A grade: { 1, 2, 3, 4, 6, 8, 9, 10, 11, 16, 17, 18 }

B grade: { 5, 7, 12, 13, 14, 15, 19, 20, 21 }

C grade: { }

F grade: { 22, 23, 24, 25, 26, 27, 28, 29, 30 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 27, 28 }

B grade: { 29, 30 }

C grade: { }

F grade: { 23, 24, 25, 26 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 }

C grade: { }

F grade: { 23, 24, 25, 26, 27, 28, 29, 30 }

2.1.9 IntegrateAlgebraic

A grade: { 23, 24, 25, 26, 28, 30 }

B grade: { 27, 29 }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, Mathematica was abbreviated to MMA and IntegrateAlgebraic to I.A.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	172	172	172	176	175	218	236	210	182	0
N.S.	1	1.00	1.00	1.02	1.02	1.27	1.37	1.22	1.06	0.00
time (sec)	N/A	0.192	0.087	0.000	0.668	1.101	0.101	0.308	0.102	0.000
Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	130	130	130	135	134	165	173	161	143	0
N.S.	1	1.00	1.00	1.04	1.03	1.27	1.33	1.24	1.10	0.00
time (sec)	N/A	0.135	0.070	0.001	0.754	1.073	0.092	0.294	0.838	0.000
Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	94	94	96	94	93	114	121	114	99	0
N.S.	1	1.00	1.02	1.00	0.99	1.21	1.29	1.21	1.05	0.00
time (sec)	N/A	0.083	0.037	0.001	0.566	1.172	0.082	0.281	0.047	0.000
Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	56	56	56	53	52	62	63	66	54	0
N.S.	1	1.00	1.00	0.95	0.93	1.11	1.12	1.18	0.96	0.00
time (sec)	N/A	0.037	0.013	0.002	0.554	0.630	0.070	0.335	0.048	0.000
Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	81	81	72	119	74	191	206	80	108	0
N.S.	1	1.00	0.89	1.47	0.91	2.36	2.54	0.99	1.33	0.00
time (sec)	N/A	0.080	0.054	0.006	1.237	1.423	0.618	0.310	0.115	0.001

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	108	108	95	163	101	318	190	95	95	0
N.S.	1	1.00	0.88	1.51	0.94	2.94	1.76	0.88	0.88	0.00
time (sec)	N/A	0.085	0.065	0.012	1.205	1.443	1.366	0.363	0.167	0.001
Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	130	130	130	175	145	471	246	135	136	0
N.S.	1	1.00	1.00	1.35	1.12	3.62	1.89	1.04	1.05	0.00
time (sec)	N/A	0.108	0.083	0.010	1.431	1.341	3.133	0.458	0.968	0.001
Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	B	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	171	171	171	210	194	642	313	184	176	0
N.S.	1	1.00	1.00	1.23	1.13	3.75	1.83	1.08	1.03	0.00
time (sec)	N/A	0.163	0.104	0.010	1.570	1.467	6.658	0.343	0.177	0.001
Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	226	226	226	237	236	289	304	283	233	0
N.S.	1	1.00	1.00	1.05	1.04	1.28	1.35	1.25	1.03	0.00
time (sec)	N/A	0.214	0.085	0.001	0.634	1.164	0.112	0.241	0.857	0.000
Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	158	158	158	169	168	202	216	202	158	0
N.S.	1	1.00	1.00	1.07	1.06	1.28	1.37	1.28	1.00	0.00
time (sec)	N/A	0.167	0.064	0.001	0.614	0.871	0.099	0.357	0.069	0.000
Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	94	94	96	101	100	114	121	120	99	0
N.S.	1	1.00	1.02	1.07	1.06	1.21	1.29	1.28	1.05	0.00
time (sec)	N/A	0.081	0.031	0.000	0.765	1.048	0.082	0.288	0.047	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	142	142	115	243	160	366	347	178	203	0
N.S.	1	1.00	0.81	1.71	1.13	2.58	2.44	1.25	1.43	0.00
time (sec)	N/A	0.208	0.071	0.006	1.351	1.195	1.038	0.294	0.857	0.001
Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	164	164	134	299	186	552	483	195	257	0
N.S.	1	1.00	0.82	1.82	1.13	3.37	2.95	1.19	1.57	0.00
time (sec)	N/A	0.232	0.099	0.013	1.939	1.382	2.827	0.327	0.167	0.001
Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	207	207	183	397	236	777	400	238	243	0
N.S.	1	1.00	0.88	1.92	1.14	3.75	1.93	1.15	1.17	0.00
time (sec)	N/A	0.239	0.130	0.013	2.012	1.426	11.984	0.449	0.976	0.001
Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	B	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	240	240	242	360	304	1024	486	311	303	0
N.S.	1	1.00	1.01	1.50	1.27	4.27	2.02	1.30	1.26	0.00
time (sec)	N/A	0.280	0.177	0.013	2.007	1.118	42.072	0.304	0.976	0.001
Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	310	310	310	339	326	409	423	401	335	0
N.S.	1	1.00	1.00	1.09	1.05	1.32	1.36	1.29	1.08	0.00
time (sec)	N/A	0.387	0.130	0.003	0.859	1.209	0.126	0.376	0.116	0.000
Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	226	226	226	244	239	289	304	289	233	0
N.S.	1	1.00	1.00	1.08	1.06	1.28	1.35	1.28	1.03	0.00
time (sec)	N/A	0.216	0.088	0.001	0.836	1.223	0.111	0.347	0.846	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	130	130	130	149	146	165	173	173	143	0
N.S.	1	1.00	1.00	1.15	1.12	1.27	1.33	1.33	1.10	0.00
time (sec)	N/A	0.135	0.055	0.001	0.841	0.848	0.093	0.353	0.815	0.000
Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	A	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	227	227	179	401	266	586	508	307	312	0
N.S.	1	1.00	0.79	1.77	1.17	2.58	2.24	1.35	1.37	0.00
time (sec)	N/A	0.371	0.108	0.007	2.029	1.004	1.492	0.494	0.849	0.001
Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	242	242	176	475	296	834	661	321	389	0
N.S.	1	1.00	0.73	1.96	1.22	3.45	2.73	1.33	1.61	0.00
time (sec)	N/A	0.399	0.128	0.015	1.978	1.239	4.749	0.370	0.977	0.001
Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	A	A	B	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	291	291	219	589	343	1102	865	371	495	0
N.S.	1	1.00	0.75	2.02	1.18	3.79	2.97	1.27	1.70	0.00
time (sec)	N/A	0.416	0.168	0.015	1.944	1.556	26.162	0.387	0.223	0.001
Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	A	B	F(-1)	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	348	348	295	735	416	1422	0	447	444	0
N.S.	1	1.00	0.85	2.11	1.20	4.09	0.00	1.28	1.28	0.00
time (sec)	N/A	0.450	0.219	0.014	2.389	1.381	0.000	0.368	1.187	0.001
Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	A	F	F(-2)	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	128	128	124	1942	0	777	0	0	-1	151
N.S.	1	1.00	0.97	15.17	0.00	6.07	0.00	0.00	-0.01	1.18
time (sec)	N/A	0.142	0.265	0.033	0.000	4.882	0.000	0.000	0.000	0.385

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	A	F	F(-2)	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	304	304	265	1541	0	1718	0	0	-1	266
N.S.	1	1.00	0.87	5.07	0.00	5.65	0.00	0.00	-0.00	0.88
time (sec)	N/A	0.299	0.366	0.038	0.000	20.972	0.000	0.000	0.000	0.669
Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	A	F	F(-2)	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	166	166	150	1052	0	1111	0	0	-1	175
N.S.	1	1.00	0.90	6.34	0.00	6.69	0.00	0.00	-0.01	1.05
time (sec)	N/A	0.105	0.339	0.016	0.000	6.224	0.000	0.000	0.000	0.355
Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	B	F	F(-2)	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	91	91	88	646	0	737	0	0	-1	114
N.S.	1	1.00	0.97	7.10	0.00	8.10	0.00	0.00	-0.01	1.25
time (sec)	N/A	0.048	0.116	0.011	0.000	4.140	0.000	0.000	0.000	0.245
Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	B	F	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	49	49	49	306	0	241	0	74	-1	103
N.S.	1	1.00	1.00	6.24	0.00	4.92	0.00	1.51	-0.02	2.10
time (sec)	N/A	0.020	0.011	0.011	0.000	1.559	0.000	0.363	0.000	0.110
Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	F(-1)	F	A	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	122	122	113	782	0	0	0	173	-1	239
N.S.	1	1.00	0.93	6.41	0.00	0.00	0.00	1.42	-0.01	1.96
time (sec)	N/A	0.111	0.210	0.031	0.000	0.000	0.000	0.524	0.000	0.427
Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	C	B	F	F(-1)	F	B	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD	NO
size	203	203	531	1865	0	0	0	479	-1	2522
N.S.	1	1.00	2.62	9.19	0.00	0.00	0.00	2.36	-0.00	12.42
time (sec)	N/A	0.265	2.639	0.036	0.000	0.000	0.000	26.049	0.000	6.798

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad	I.A.
grade	A	A	A	B	F	B	F	B	F	A
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD	Yes
size	113	113	191	1622	0	513	0	336	-1	146
N.S.	1	1.00	1.69	14.35	0.00	4.54	0.00	2.97	-0.01	1.29
time (sec)	N/A	0.119	0.808	0.029	0.000	5.765	0.000	3.395	0.000	0.623

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [24] had the largest ratio of [.2667]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	2	1	1.00	24	0.042
2	A	2	1	1.00	24	0.042
3	A	2	1	1.00	24	0.042
4	A	2	1	1.00	22	0.045
5	A	3	3	1.00	24	0.125
6	A	3	3	1.00	24	0.125
7	A	3	3	1.00	24	0.125
8	A	4	4	1.00	24	0.167
9	A	2	1	1.00	26	0.038
10	A	2	1	1.00	26	0.038
11	A	2	1	1.00	24	0.042
12	A	4	3	1.00	26	0.115
13	A	4	4	1.00	26	0.154
14	A	4	3	1.00	26	0.115
15	A	4	3	1.00	26	0.115
16	A	2	1	1.00	26	0.038
17	A	2	1	1.00	26	0.038
18	A	2	1	1.00	24	0.042
19	A	5	3	1.00	26	0.115
20	A	5	4	1.00	26	0.154
21	A	5	4	1.00	26	0.154
22	A	5	3	1.00	26	0.115
23	A	6	6	1.00	28	0.214
24	A	14	8	1.00	30	0.267
25	A	9	7	1.00	30	0.233
26	A	5	5	1.00	28	0.179
27	A	2	2	1.00	21	0.095
28	A	5	3	1.00	30	0.100
29	A	7	5	1.00	30	0.167
30	A	4	4	1.00	28	0.143

Chapter 3

Listing of integrals

Local contents

3.1	$\int (a + bx^2)(c + dx^2)(e + fx^2)^4 dx$	30
3.2	$\int (a + bx^2)(c + dx^2)(e + fx^2)^3 dx$	33
3.3	$\int (a + bx^2)(c + dx^2)(e + fx^2)^2 dx$	36
3.4	$\int (a + bx^2)(c + dx^2)(e + fx^2) dx$	39
3.5	$\int \frac{(a+bx^2)(c+dx^2)}{e+fx^2} dx$	41
3.6	$\int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^2} dx$	44
3.7	$\int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^3} dx$	47
3.8	$\int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^4} dx$	51
3.9	$\int (a + bx^2)(c + dx^2)^2(e + fx^2)^3 dx$	55
3.10	$\int (a + bx^2)(c + dx^2)^2(e + fx^2)^2 dx$	58
3.11	$\int (a + bx^2)(c + dx^2)^2(e + fx^2) dx$	61
3.12	$\int \frac{(a+bx^2)(c+dx^2)^2}{e+fx^2} dx$	64
3.13	$\int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^2} dx$	68
3.14	$\int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^3} dx$	72
3.15	$\int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^4} dx$	76
3.16	$\int (a + bx^2)(c + dx^2)^3(e + fx^2)^3 dx$	80
3.17	$\int (a + bx^2)(c + dx^2)^3(e + fx^2)^2 dx$	83
3.18	$\int (a + bx^2)(c + dx^2)^3(e + fx^2) dx$	86
3.19	$\int \frac{(a+bx^2)(c+dx^2)^3}{e+fx^2} dx$	89
3.20	$\int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^2} dx$	93
3.21	$\int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^3} dx$	97
3.22	$\int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^4} dx$	101

3.23	$\int \frac{(a+bx^2)\sqrt{c+dx^2}}{e+fx^2} dx$	105
3.24	$\int \frac{(a+bx^2)^3}{(c+dx^2)\sqrt{e+fx^2}} dx$	109
3.25	$\int \frac{(a+bx^2)^2}{(c+dx^2)\sqrt{e+fx^2}} dx$	114
3.26	$\int \frac{a+bx^2}{(c+dx^2)\sqrt{e+fx^2}} dx$	118
3.27	$\int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx$	122
3.28	$\int \frac{1}{(a+bx^2)(c+dx^2)\sqrt{e+fx^2}} dx$	125
3.29	$\int \frac{1}{(a+bx^2)^2(c+dx^2)\sqrt{e+fx^2}} dx$	128
3.30	$\int \frac{a+bx^2}{\sqrt{c+dx^2}(e+fx^2)^2} dx$	133

$$3.1 \quad \int (a + bx^2)(c + dx^2)(e + fx^2)^4 dx$$

Optimal. Leaf size=172

$$\frac{1}{3}e^3x^3(4acf+ade+bce)+\frac{1}{5}e^2x^5(2af(3cf+2de)+be(4cf+de))+\frac{1}{11}f^3x^{11}(adf+bcf+4bde)+\frac{1}{9}f^2x^9(af(cf+4de)+2$$

Rubi [A] time = 0.19, antiderivative size = 172, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {521}

$$\frac{1}{5}e^2x^5(2af(3cf+2de)+be(4cf+de))+\frac{1}{3}e^3x^3(4acf+ade+bce)+\frac{1}{11}f^3x^{11}(adf+bcf+4bde)+\frac{1}{9}f^2x^9(af(cf+4de)+2be(2cf+3de))+\frac{2}{7}efx^7(af(2cf+3de)+be(3cf+2de))+ace^4x+\frac{1}{13}bdf^4x^{13}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^4,x]

[Out] a*c*e^4*x + (e^3*(b*c*e + a*d*e + 4*a*c*f)*x^3)/3 + (e^2*(2*a*f*(2*d*e + 3*c*f) + b*e*(d*e + 4*c*f))*x^5)/5 + (2*e*f*(a*f*(3*d*e + 2*c*f) + b*e*(2*d*e + 3*c*f))*x^7)/7 + (f^2*(a*f*(4*d*e + c*f) + 2*b*e*(3*d*e + 2*c*f))*x^9)/9 + (f^3*(4*b*d*e + b*c*f + a*d*f)*x^11)/11 + (b*d*f^4*x^13)/13

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)(e + fx^2)^4 dx &= \int (ace^4 + e^3(bce + ade + 4acf)x^2 + e^2(2af(2de + 3cf) + be(de + 4cf)) \\ &= ace^4x + \frac{1}{3}e^3(bce + ade + 4acf)x^3 + \frac{1}{5}e^2(2af(2de + 3cf) + be(de + 4cf)) \end{aligned}$$

Mathematica [A] time = 0.09, size = 172, normalized size = 1.00

$$\frac{1}{3}e^3x^3(4acf+ade+bce)+\frac{1}{5}e^2x^5(2af(3cf+2de)+be(4cf+de))+\frac{1}{11}f^3x^{11}(adf+bcf+4bde)+\frac{1}{9}f^2x^9(af(cf+4de)+2be(2cf+3de))+\frac{2}{7}efx^7(af(2cf+3de)+be(3cf+2de))+ace^4x+\frac{1}{13}bdf^4x^{13}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^4,x]

[Out] a*c*e^4*x + (e^3*(b*c*e + a*d*e + 4*a*c*f)*x^3)/3 + (e^2*(2*a*f*(2*d*e + 3*c*f) + b*e*(d*e + 4*c*f))*x^5)/5 + (2*e*f*(a*f*(3*d*e + 2*c*f) + b*e*(2*d*e + 3*c*f))*x^7)/7 + (f^2*(a*f*(4*d*e + c*f) + 2*b*e*(3*d*e + 2*c*f))*x^9)/9 + (f^3*(4*b*d*e + b*c*f + a*d*f)*x^11)/11 + (b*d*f^4*x^13)/13

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)(e + fx^2)^4 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^4,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^4, x]

fricas [A] time = 1.10, size = 218, normalized size = 1.27

$$\frac{1}{13}x^{13}f^4db + \frac{4}{11}x^{11}f^3adb + \frac{1}{11}x^{11}f^4cb + \frac{1}{11}x^{11}f^4da + \frac{2}{3}x^9f^2db + \frac{4}{9}x^9f^2cb + \frac{4}{9}x^9f^2da + \frac{1}{9}x^9f^2ca + \frac{4}{7}x^7f^2db + \frac{6}{7}x^7f^2cb + \frac{6}{7}x^7f^2da + \frac{4}{7}x^7f^2ca + \frac{1}{5}x^5f^2db + \frac{4}{5}x^5f^2cb + \frac{4}{5}x^5f^2da + \frac{6}{5}x^5f^2ca + \frac{1}{3}x^3f^2db + \frac{1}{3}x^3f^2cb + \frac{4}{3}x^3f^2da + \frac{4}{3}x^3f^2ca + xe^4ca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^4,x, algorithm="fricas")

[Out] $\frac{1}{13}x^{13}f^4d*b + \frac{4}{11}x^{11}f^3e*d*b + \frac{1}{11}x^{11}f^4c*b + \frac{1}{11}x^{11}f^4d*a + \frac{2}{3}x^9f^2e^2*d*b + \frac{4}{9}x^9f^3e*c*b + \frac{4}{9}x^9f^3e*d*a + \frac{1}{9}x^9f^4c*a + \frac{4}{7}x^7f^2e^3*d*b + \frac{6}{7}x^7f^2e^2*c*b + \frac{6}{7}x^7f^2e^2*d*a + \frac{4}{7}x^7f^3e*c*a + \frac{1}{5}x^5f^4d*b + \frac{4}{5}x^5f^3e*c*b + \frac{4}{5}x^5f^3e*d*a + \frac{6}{5}x^5f^2e^2*c*a + \frac{1}{3}x^3f^3e^4*c*b + \frac{1}{3}x^3f^3e^4*d*a + \frac{4}{3}x^3f^3e^3*c*a + x*e^4*c*a$

giac [A] time = 0.31, size = 210, normalized size = 1.22

$$\frac{1}{13}bdf^4x^{13} + \frac{1}{11}bcf^4x^{11} + \frac{1}{11}adf^4x^{11} + \frac{4}{11}bdf^3x^{11}e + \frac{1}{9}acf^4x^9 + \frac{4}{9}bcf^3x^9e + \frac{4}{9}adf^3x^9e + \frac{2}{3}bdf^2x^9e^2 + \frac{4}{7}a*c*f^3*x^7*e + \frac{6}{7}b*c*f^2*x^7*e^2 + \frac{6}{7}a*d*f^2*x^7*e^2 + \frac{4}{7}bdf^2x^7e^2 + \frac{6}{5}acf^2x^5e^2 + \frac{4}{5}bcf^2x^5e^2 + \frac{4}{5}adf^2x^5e^2 + \frac{1}{3}bde^3e^4 + \frac{4}{3}acf^3e^3 + \frac{1}{3}bcc^3e^4 + \frac{1}{3}ad^3e^4 + acxe^4$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^4,x, algorithm="giac")

[Out] $\frac{1}{13}b*d*f^4*x^{13} + \frac{1}{11}b*c*f^4*x^{11} + \frac{1}{11}a*d*f^4*x^{11} + \frac{4}{11}b*d*f^3*x^{11}e + \frac{1}{9}a*c*f^4*x^9 + \frac{4}{9}b*c*f^3*x^9e + \frac{4}{9}a*d*f^3*x^9e + \frac{2}{3}b*d*f^2*x^9e^2 + \frac{4}{7}a*c*f^3*x^7*e + \frac{6}{7}b*c*f^2*x^7*e^2 + \frac{6}{7}a*d*f^2*x^7*e^2 + \frac{4}{7}b*d*f^2*x^7*e^2 + \frac{4}{7}b*d*f*x^7*e^3 + \frac{6}{5}a*c*f^2*x^5*e^2 + \frac{4}{5}b*c*f*x^5*e^3 + \frac{4}{5}a*d*f*x^5*e^3 + \frac{1}{3}b*d*x^5*e^4 + \frac{4}{3}a*c*f*x^3*e^3 + \frac{1}{3}b*c*x^3*e^4 + \frac{1}{3}a*d*x^3*e^4 + a*c*x*e^4$

maple [A] time = 0.00, size = 176, normalized size = 1.02

$$\frac{bd^4f^4x^{13}}{13} + \frac{(4bde^3 + (ad+bc)f^4)x^{11}}{11} + \frac{(acf^4 + 6bd^2e^2 + 4(ad+bc)e^2f^3)x^9}{9} + ac^4x + \frac{(4ace^3 + 4bde^3f + 6(ad+bc)e^2f^2)x^7}{7} + \frac{(6ac^2e^2 + bde^4 + 4(ad+bc)e^3f)x^5}{5} + \frac{(4ac^3f + (ad+bc)e^4)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^4,x)

[Out] $\frac{1}{13}b*d*f^4*x^{13} + \frac{1}{11}((a*d+b*c)*f^4 + 4*b*d*e*f^3)*x^{11} + \frac{1}{9}((a*c*f^4 + 4*(a*d+b*c)*e*f^3 + 6*b*d*e^2*f^2)*x^9 + \frac{1}{7}((4*a*c*e*f^3 + 6*(a*d+b*c)*e^2*f^2 + 4*b*d*e^3*f)*x^7 + \frac{1}{5}((6*a*c*e^2*f^2 + 4*(a*d+b*c)*e^3*f + b*d*e^4)*x^5 + \frac{1}{3}((4*a*c*e^3*f + (a*d+b*c)*e^4)*x^3 + a*c*e^4*x$

maxima [A] time = 0.67, size = 175, normalized size = 1.02

$$\frac{1}{13}bdf^4x^{13} + \frac{1}{11}(4bde^3 + (bc+ad)f^4)x^{11} + \frac{1}{9}(6bde^2f^2 + acf^4 + 4(bc+ad)e^2f^3)x^9 + \frac{2}{7}(2bde^3f + 2ace^3 + 3(bc+ad)e^2f^2)x^7 + ace^4x + \frac{1}{5}(bde^4 + 6ace^2f^2 + 4(bc+ad)e^3f)x^5 + \frac{1}{3}(4ace^3f + (bc+ad)e^4)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^4,x, algorithm="maxima")

[Out] $\frac{1}{13}b*d*f^4*x^{13} + \frac{1}{11}((4*b*d*e*f^3 + (b*c + a*d)*f^4)*x^{11} + \frac{1}{9}((6*b*d*e^2*f^2 + a*c*f^4 + 4*(b*c + a*d)*e*f^3)*x^9 + \frac{2}{7}((2*b*d*e^3*f + 2*a*c*e*f^3 + 3*(b*c + a*d)*e^2*f^2)*x^7 + a*c*e^4*x + \frac{1}{5}((b*d*e^4 + 6*a*c*e^2*f^2 + 4*(b*c + a*d)*e^3*f)*x^5 + \frac{1}{3}((4*a*c*e^3*f + (b*c + a*d)*e^4)*x^3$

mupad [B] time = 0.10, size = 182, normalized size = 1.06

$$x^3 \left(\frac{ad^4}{3} + \frac{bce^4}{3} + \frac{4ace^3f}{3} \right) + x^{11} \left(\frac{adf^4}{11} + \frac{bcf^4}{11} + \frac{4bde^3f^3}{11} \right) + x^5 \left(\frac{bd^4}{5} + \frac{4ad^2e^2f}{5} + \frac{4bce^3f}{5} + \frac{6ace^2f^2}{5} \right) + x^9 \left(\frac{acf^4}{9} + \frac{4adef^3}{9} + \frac{4bce^3f}{9} + \frac{2bd^2e^2f^2}{3} \right) + \frac{2efx^7(2acf^2 + 2bd^2e^2 + 3adef + 3bcef)}{7} + ac^4x + \frac{bdf^4x^{13}}{13}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)*(e + f*x^2)^4,x)


```
[Out] x^3*((a*d*e^4)/3 + (b*c*e^4)/3 + (4*a*c*e^3*f)/3) + x^11*((a*d*f^4)/11 + (b*c*f^4)/11 + (4*b*d*e*f^3)/11) + x^5*((b*d*e^4)/5 + (4*a*d*e^3*f)/5 + (4*b*c*e^3*f)/5 + (6*a*c*e^2*f^2)/5) + x^9*((a*c*f^4)/9 + (4*a*d*e*f^3)/9 + (4*b*c*e*f^3)/9 + (2*b*d*e^2*f^2)/3) + (2*e*f*x^7*(2*a*c*f^2 + 2*b*d*e^2 + 3*a*d*e*f + 3*b*c*e*f))/7 + a*c*e^4*x + (b*d*f^4*x^13)/13
```

sympy [A] time = 0.10, size = 236, normalized size = 1.37

$$ace^4x + \frac{bdf^4x^{13}}{13} + x^{11}\left(\frac{adf^4}{11} + \frac{bcf^4}{11} + \frac{4bde^3f}{11}\right) + x^9\left(\frac{acf^4}{9} + \frac{4ade^3f}{9} + \frac{4bcef^3}{9} + \frac{2bd^2f^2}{3}\right) + x^7\left(\frac{4ace^3f}{7} + \frac{6ade^2f^2}{7} + \frac{6bce^2f^2}{7} + \frac{4bde^3f}{7}\right) + x^5\left(\frac{6ace^2f^2}{5} + \frac{4ade^3f}{5} + \frac{4bce^3f}{5} + \frac{bde^4}{5}\right) + x^3\left(\frac{4ace^3f}{3} + \frac{ade^4}{3} + \frac{bce^4}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)*(f*x**2+e)**4,x)
```

```
[Out] a*c*e**4*x + b*d*f**4*x**13/13 + x**11*(a*d*f**4/11 + b*c*f**4/11 + 4*b*d*e*f**3/11) + x**9*(a*c*f**4/9 + 4*a*d*e*f**3/9 + 4*b*c*e*f**3/9 + 2*b*d*e**2*f**2/3) + x**7*(4*a*c*e*f**3/7 + 6*a*d*e**2*f**2/7 + 6*b*c*e**2*f**2/7 + 4*b*d*e**3*f/7) + x**5*(6*a*c*e**2*f**2/5 + 4*a*d*e**3*f/5 + 4*b*c*e**3*f/5 + b*d*e**4/5) + x**3*(4*a*c*e**3*f/3 + a*d*e**4/3 + b*c*e**4/3)
```

$$3.2 \quad \int (a + bx^2)(c + dx^2)(e + fx^2)^3 dx$$

Optimal. Leaf size=130

$$\frac{1}{3}e^2x^3(3acf+ade+bce)+\frac{1}{9}f^2x^9(adf+bcf+3bde)+\frac{1}{7}fx^7(af(cf+3de)+3be(cf+de))+\frac{1}{5}ex^5(3af(cf+de)+be(3cf+de))$$

Rubi [A] time = 0.13, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {521}

$$\frac{1}{3}e^2x^3(3acf+ade+bce)+\frac{1}{9}f^2x^9(adf+bcf+3bde)+\frac{1}{7}fx^7(af(cf+3de)+3be(cf+de))+\frac{1}{5}ex^5(3af(cf+de)+be(3cf+de))+ace^3x+\frac{1}{11}bdf^3x^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^3,x]

[Out] a*c*e^3*x + (e^2*(b*c*e + a*d*e + 3*a*c*f)*x^3)/3 + (e*(3*a*f*(d*e + c*f) + b*e*(d*e + 3*c*f))*x^5)/5 + (f*(3*b*e*(d*e + c*f) + a*f*(3*d*e + c*f))*x^7)/7 + (f^2*(3*b*d*e + b*c*f + a*d*f)*x^9)/9 + (b*d*f^3*x^11)/11

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)(e + fx^2)^3 dx &= \int (ace^3 + e^2(bce + ade + 3acf)x^2 + e(3af(de + cf) + be(de + 3cf))x^4 + f(bce + ade + 3acf)x^6 + e(3af(de + cf) + be(de + 3cf))x^8 + f^2(3bde + bcf + 3ade)x^{10} + bdf^3x^{12}) dx \\ &= ace^3x + \frac{1}{3}e^2(bce + ade + 3acf)x^3 + \frac{1}{5}e(3af(de + cf) + be(de + 3cf))x^5 + \frac{1}{7}f(bce + ade + 3acf)x^7 + \frac{1}{9}e(3af(de + cf) + be(de + 3cf))x^9 + \frac{1}{11}bdf^3x^{11} \end{aligned}$$

Mathematica [A] time = 0.07, size = 130, normalized size = 1.00

$$\frac{1}{3}e^2x^3(3acf+ade+bce)+\frac{1}{9}f^2x^9(adf+bcf+3bde)+\frac{1}{7}fx^7(af(cf+3de)+3be(cf+de))+\frac{1}{5}ex^5(3af(cf+de)+be(3cf+de))+ace^3x+\frac{1}{11}bdf^3x^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^3,x]

[Out] a*c*e^3*x + (e^2*(b*c*e + a*d*e + 3*a*c*f)*x^3)/3 + (e*(3*a*f*(d*e + c*f) + b*e*(d*e + 3*c*f))*x^5)/5 + (f*(3*b*e*(d*e + c*f) + a*f*(3*d*e + c*f))*x^7)/7 + (f^2*(3*b*d*e + b*c*f + a*d*f)*x^9)/9 + (b*d*f^3*x^11)/11

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)(e + fx^2)^3 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^3, x]

fricas [A] time = 1.07, size = 165, normalized size = 1.27

$$\frac{1}{11}x^{11}f^3db+\frac{1}{3}x^9f^2edb+\frac{1}{9}x^9f^3cb+\frac{1}{9}x^9f^3da+\frac{3}{7}x^7f^2db+\frac{3}{7}x^7f^2eb+\frac{3}{7}x^7f^2eda+\frac{1}{7}x^7f^3ca+\frac{1}{5}x^5e^3db+\frac{3}{5}x^5f^2eb+\frac{3}{5}x^5f^2da+\frac{3}{5}x^5f^2eca+\frac{1}{3}x^3e^3cb+\frac{1}{3}x^3e^3da+x^3f^2ca+xe^3ca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^3,x, algorithm="fricas")

[Out] $1/11*x^{11}*f^3*d*b + 1/3*x^9*f^2*e*d*b + 1/9*x^9*f^3*c*b + 1/9*x^9*f^3*d*a + 3/7*x^7*f*e^2*d*b + 3/7*x^7*f^2*e*c*b + 3/7*x^7*f^2*e*d*a + 1/7*x^7*f^3*c*a + 1/5*x^5*e^3*d*b + 3/5*x^5*f*e^2*c*b + 3/5*x^5*f*e^2*d*a + 3/5*x^5*f^2*e*c*a + 1/3*x^3*e^3*c*b + 1/3*x^3*e^3*d*a + x^3*f*e^2*c*a + x*e^3*c*a$

giac [A] time = 0.29, size = 161, normalized size = 1.24

$$\frac{1}{11} b d f^3 x^{11} + \frac{1}{9} b c f^3 x^9 + \frac{1}{9} a d f^3 x^9 + \frac{1}{3} b d f^2 x^9 e + \frac{1}{7} a c f^3 x^7 + \frac{3}{7} b c f^2 x^7 e + \frac{3}{7} a d f^2 x^7 e + \frac{3}{7} b d f x^7 e^2 + \frac{3}{5} a c f^2 x^5 e + \frac{3}{5} b c f x^5 e^2 + \frac{3}{5} a d f x^5 e^2 + \frac{1}{5} b d x^5 e^3 + a c f x^3 e^2 + \frac{1}{3} b c x^3 e^3 + \frac{1}{3} a d x^3 e^3 + a c x e^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^3,x, algorithm="giac")

[Out] $1/11*b*d*f^3*x^{11} + 1/9*b*c*f^3*x^9 + 1/9*a*d*f^3*x^9 + 1/3*b*d*f^2*x^9*e + 1/7*a*c*f^3*x^7 + 3/7*b*c*f^2*x^7*e + 3/7*a*d*f^2*x^7*e + 3/7*b*d*f*x^7*e^2 + 3/5*a*c*f^2*x^5*e + 3/5*b*c*f*x^5*e^2 + 3/5*a*d*f*x^5*e^2 + 1/5*b*d*x^5*e^3 + a*c*f*x^3*e^2 + 1/3*b*c*x^3*e^3 + 1/3*a*d*x^3*e^3 + a*c*x*e^3$

maple [A] time = 0.00, size = 135, normalized size = 1.04

$$\frac{b d f^3 x^{11}}{11} + \frac{(3 b d e f^2 + (a d + b c) f^3) x^9}{9} + \frac{(a c f^3 + 3 b d e^2 f + 3 (a d + b c) e f^2) x^7}{7} + a c e^3 x + \frac{(3 a c e f^2 + b d e^3 + 3 (a d + b c) e^2 f) x^5}{5} + \frac{(3 a c e^2 f + (a d + b c) e^3) x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^3,x)

[Out] $1/11*b*d*f^3*x^{11} + 1/9*((a*d+b*c)*f^3 + 3*b*d*e*f^2)*x^9 + 1/7*(a*c*f^3 + 3*(a*d+b*c)*e*f^2 + 3*b*d*e^2*f)*x^7 + 1/5*(3*a*c*e*f^2 + 3*(a*d+b*c)*e^2*f + b*d*e^3)*x^5 + 1/3*(3*a*c*e^2*f + (a*d+b*c)*e^3)*x^3 + a*c*e^3*x$

maxima [A] time = 0.75, size = 134, normalized size = 1.03

$$\frac{1}{11} b d f^3 x^{11} + \frac{1}{9} (3 b d e f^2 + (b c + a d) f^3) x^9 + \frac{1}{7} (3 b d e^2 f + a c f^3 + 3 (b c + a d) e f^2) x^7 + a c e^3 x + \frac{1}{5} (b d e^3 + 3 a c e f^2 + 3 (b c + a d) e^2 f) x^5 + \frac{1}{3} (3 a c e^2 f + (b c + a d) e^3) x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^3,x, algorithm="maxima")

[Out] $1/11*b*d*f^3*x^{11} + 1/9*(3*b*d*e*f^2 + (b*c + a*d)*f^3)*x^9 + 1/7*(3*b*d*e^2*f + a*c*f^3 + 3*(b*c + a*d)*e*f^2)*x^7 + a*c*e^3*x + 1/5*(b*d*e^3 + 3*a*c*e*f^2 + 3*(b*c + a*d)*e^2*f)*x^5 + 1/3*(3*a*c*e^2*f + (b*c + a*d)*e^3)*x^3$

mupad [B] time = 0.84, size = 143, normalized size = 1.10

$$x^5 \left(\frac{b d e^3}{5} + \frac{3 a c e f^2}{5} + \frac{3 a d e^2 f}{5} + \frac{3 b c e^2 f}{5} \right) + x^7 \left(\frac{a c f^3}{7} + \frac{3 a d e f^2}{7} + \frac{3 b c e f^2}{7} + \frac{3 b d e^2 f}{7} \right) + x^3 \left(\frac{a d e^3}{3} + \frac{b c e^3}{3} + a c e^2 f \right) + x^9 \left(\frac{a d f^3}{9} + \frac{b c f^3}{9} + \frac{b d e f^2}{3} \right) + a c e^3 x + \frac{b d f^3 x^{11}}{11}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)*(e + f*x^2)^3,x)

[Out] $x^5*((b*d*e^3)/5 + (3*a*c*e*f^2)/5 + (3*a*d*e^2*f)/5 + (3*b*c*e^2*f)/5) + x^7*((a*c*f^3)/7 + (3*a*d*e*f^2)/7 + (3*b*c*e*f^2)/7 + (3*b*d*e^2*f)/7) + x^9*(3*((a*d*e^3)/3 + (b*c*e^3)/3 + a*c*e^2*f) + x^9*((a*d*f^3)/9 + (b*c*f^3)/9 + (b*d*e*f^2)/3) + a*c*e^3*x + (b*d*f^3*x^{11})/11$

sympy [A] time = 0.09, size = 173, normalized size = 1.33

$$a c e^3 x + \frac{b d f^3 x^{11}}{11} + x^9 \left(\frac{a d f^3}{9} + \frac{b c f^3}{9} + \frac{b d e f^2}{3} \right) + x^7 \left(\frac{a c f^3}{7} + \frac{3 a d e f^2}{7} + \frac{3 b c e f^2}{7} + \frac{3 b d e^2 f}{7} \right) + x^5 \left(\frac{3 a c e f^2}{5} + \frac{3 a d e^2 f}{5} + \frac{3 b c e^2 f}{5} + \frac{b d e^3}{5} \right) + x^3 \left(a c e^2 f + \frac{a d e^3}{3} + \frac{b c e^3}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)*(d*x**2+c)*(f*x**2+e)**3,x)`

[Out] $a*c*e**3*x + b*d*f**3*x**11/11 + x**9*(a*d*f**3/9 + b*c*f**3/9 + b*d*e*f**2/3) + x**7*(a*c*f**3/7 + 3*a*d*e*f**2/7 + 3*b*c*e*f**2/7 + 3*b*d*e**2*f/7) + x**5*(3*a*c*e*f**2/5 + 3*a*d*e**2*f/5 + 3*b*c*e**2*f/5 + b*d*e**3/5) + x**3*(a*c*e**2*f + a*d*e**3/3 + b*c*e**3/3)$

$$3.3 \quad \int (a + bx^2)(c + dx^2)(e + fx^2)^2 dx$$

Optimal. Leaf size=94

$$\frac{1}{7}fx^7(adf+bcf+2bde)+\frac{1}{5}x^5(af(cf+2de)+be(2cf+de))+\frac{1}{3}ex^3(2acf+ade+bce)+ace^2x+\frac{1}{9}bdf^2x^9$$

Rubi [A] time = 0.08, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {521}

$$\frac{1}{7}fx^7(adf+bcf+2bde)+\frac{1}{5}x^5(af(cf+2de)+be(2cf+de))+\frac{1}{3}ex^3(2acf+ade+bce)+ace^2x+\frac{1}{9}bdf^2x^9$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^2,x]

[Out] a*c*e^2*x + (e*(b*c*e + a*d*e + 2*a*c*f)*x^3)/3 + ((a*f*(2*d*e + c*f) + b*e*(d*e + 2*c*f))*x^5)/5 + (f*(2*b*d*e + b*c*f + a*d*f)*x^7)/7 + (b*d*f^2*x^9)/9

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)(e + fx^2)^2 dx &= \int (ace^2 + e(bce + ade + 2acf)x^2 + (af(2de + cf) + be(de + 2cf))x^4 + \\ &= ace^2x + \frac{1}{3}e(bce + ade + 2acf)x^3 + \frac{1}{5}(af(2de + cf) + be(de + 2cf))x^5 + \end{aligned}$$

Mathematica [A] time = 0.04, size = 96, normalized size = 1.02

$$\frac{1}{5}x^5(acf^2 + 2adef + 2bcef + bde^2) + \frac{1}{7}fx^7(adf + bcf + 2bde) + \frac{1}{3}ex^3(2acf + ade + bce) + ace^2x + \frac{1}{9}bdf^2x^9$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^2,x]

[Out] a*c*e^2*x + (e*(b*c*e + a*d*e + 2*a*c*f)*x^3)/3 + ((b*d*e^2 + 2*b*c*e*f + 2*a*d*e*f + a*c*f^2)*x^5)/5 + (f*(2*b*d*e + b*c*f + a*d*f)*x^7)/7 + (b*d*f^2*x^9)/9

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)(e + fx^2)^2 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2)^2, x]

fricas [A] time = 1.17, size = 114, normalized size = 1.21

$$\frac{1}{9}x^9 f^2 db + \frac{2}{7}x^7 f e db + \frac{1}{7}x^7 f^2 cb + \frac{1}{7}x^7 f^2 da + \frac{1}{5}x^5 e^2 db + \frac{2}{5}x^5 f e cb + \frac{2}{5}x^5 f e da + \frac{1}{5}x^5 f^2 ca + \frac{1}{3}x^3 e^2 cb + \frac{1}{3}x^3 e^2 da + \frac{2}{3}x^3 f e ca + x e^2 ca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^2,x, algorithm="fricas")

[Out] $\frac{1}{9}x^9 f^2 d b + \frac{2}{7}x^7 f e d b + \frac{1}{7}x^7 f^2 c b + \frac{1}{7}x^7 f^2 d a + \frac{1}{5}x^5 e^2 d b + \frac{2}{5}x^5 f e c b + \frac{2}{5}x^5 f e d a + \frac{1}{5}x^5 f^2 c a + \frac{1}{3}x^3 e^2 c b + \frac{1}{3}x^3 e^2 d a + \frac{2}{3}x^3 f e c a + x e^2 c a$

giac [A] time = 0.28, size = 114, normalized size = 1.21

$$\frac{1}{9}b d f^2 x^9 + \frac{1}{7}b c f^2 x^7 + \frac{1}{7}a d f^2 x^7 + \frac{2}{7}b d f x^7 e + \frac{1}{5}a c f^2 x^5 + \frac{2}{5}b c f x^5 e + \frac{2}{5}a d f x^5 e + \frac{1}{5}b d x^5 e^2 + \frac{2}{3}a c f x^3 e + \frac{1}{3}b c x^3 e^2 + \frac{1}{3}a d x^3 e^2 + a c x e^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^2,x, algorithm="giac")

[Out] $\frac{1}{9}b d f^2 x^9 + \frac{1}{7}b c f^2 x^7 + \frac{1}{7}a d f^2 x^7 + \frac{2}{7}b d f x^7 e + \frac{1}{5}a c f^2 x^5 + \frac{2}{5}b c f x^5 e + \frac{2}{5}a d f x^5 e + \frac{1}{5}b d x^5 e^2 + \frac{2}{3}a c f x^3 e + \frac{1}{3}b c x^3 e^2 + \frac{1}{3}a d x^3 e^2 + a c x e^2$

maple [A] time = 0.00, size = 94, normalized size = 1.00

$$\frac{b d f^2 x^9}{9} + \frac{(2 b d e f + (a d + b c) f^2) x^7}{7} + a c e^2 x + \frac{(a c f^2 + b d e^2 + 2(a d + b c) e f) x^5}{5} + \frac{(2 a c e f + (a d + b c) e^2) x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^2,x)

[Out] $\frac{1}{9}b d f^2 x^9 + \frac{1}{7}((a d + b c) f^2 + 2 b d e f) x^7 + \frac{1}{5}(a c f^2 + 2(a d + b c) e f + b d e^2) x^5 + \frac{1}{3}(2 a c e f + (a d + b c) e^2) x^3 + a c e^2 x$

maxima [A] time = 0.57, size = 93, normalized size = 0.99

$$\frac{1}{9}b d f^2 x^9 + \frac{1}{7}(2 b d e f + (b c + a d) f^2) x^7 + \frac{1}{5}(b d e^2 + a c f^2 + 2(b c + a d) e f) x^5 + a c e^2 x + \frac{1}{3}(2 a c e f + (b c + a d) e^2) x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e)^2,x, algorithm="maxima")

[Out] $\frac{1}{9}b d f^2 x^9 + \frac{1}{7}(2 b d e f + (b c + a d) f^2) x^7 + \frac{1}{5}(b d e^2 + a c f^2 + 2(b c + a d) e f) x^5 + a c e^2 x + \frac{1}{3}(2 a c e f + (b c + a d) e^2) x^3$

mupad [B] time = 0.05, size = 99, normalized size = 1.05

$$x^5 \left(\frac{a c f^2}{5} + \frac{b d e^2}{5} + \frac{2 a d e f}{5} + \frac{2 b c e f}{5} \right) + x^3 \left(\frac{a d e^2}{3} + \frac{b c e^2}{3} + \frac{2 a c e f}{3} \right) + x^7 \left(\frac{a d f^2}{7} + \frac{b c f^2}{7} + \frac{2 b d e f}{7} \right) + a c e^2 x + \frac{b d f^2 x^9}{9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)*(e + f*x^2)^2,x)

[Out] $x^5 \left(\frac{a c f^2}{5} + \frac{b d e^2}{5} + \frac{2 a d e f}{5} + \frac{2 b c e f}{5} \right) + x^3 \left(\frac{a d e^2}{3} + \frac{b c e^2}{3} + \frac{2 a c e f}{3} \right) + x^7 \left(\frac{a d f^2}{7} + \frac{b c f^2}{7} + \frac{2 b d e f}{7} \right) + a c e^2 x + \frac{b d f^2 x^9}{9}$

sympy [A] time = 0.08, size = 121, normalized size = 1.29

$$a c e^2 x + \frac{b d f^2 x^9}{9} + x^7 \left(\frac{a d f^2}{7} + \frac{b c f^2}{7} + \frac{2 b d e f}{7} \right) + x^5 \left(\frac{a c f^2}{5} + \frac{2 a d e f}{5} + \frac{2 b c e f}{5} + \frac{b d e^2}{5} \right) + x^3 \left(\frac{2 a c e f}{3} + \frac{a d e^2}{3} + \frac{b c e^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)*(f*x**2+e)**2,x)
```

```
[Out] a*c*e**2*x + b*d*f**2*x**9/9 + x**7*(a*d*f**2/7 + b*c*f**2/7 + 2*b*d*e*f/7)
+ x**5*(a*c*f**2/5 + 2*a*d*e*f/5 + 2*b*c*e*f/5 + b*d*e**2/5) + x**3*(2*a*c
*e*f/3 + a*d*e**2/3 + b*c*e**2/3)
```

3.4 $\int (a + bx^2)(c + dx^2)(e + fx^2) dx$

Optimal. Leaf size=56

$$\frac{1}{5}x^5(adf + bcf + bde) + \frac{1}{3}x^3(acf + ade + bce) + acex + \frac{1}{7}bdfx^7$$

Rubi [A] time = 0.04, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.045$, Rules used = {521}

$$\frac{1}{5}x^5(adf + bcf + bde) + \frac{1}{3}x^3(acf + ade + bce) + acex + \frac{1}{7}bdfx^7$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)*(e + f*x^2),x]

[Out] a*c*e*x + ((b*c*e + a*d*e + a*c*f)*x^3)/3 + ((b*d*e + b*c*f + a*d*f)*x^5)/5 + (b*d*f*x^7)/7

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)(e + fx^2) dx &= \int (ace + (bce + ade + acf)x^2 + (bde + bcf + adf)x^4 + bdfx^6) dx \\ &= acex + \frac{1}{3}(bce + ade + acf)x^3 + \frac{1}{5}(bde + bcf + adf)x^5 + \frac{1}{7}bdfx^7 \end{aligned}$$

Mathematica [A] time = 0.01, size = 56, normalized size = 1.00

$$\frac{1}{5}x^5(adf + bcf + bde) + \frac{1}{3}x^3(acf + ade + bce) + acex + \frac{1}{7}bdfx^7$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)*(e + f*x^2),x]

[Out] a*c*e*x + ((b*c*e + a*d*e + a*c*f)*x^3)/3 + ((b*d*e + b*c*f + a*d*f)*x^5)/5 + (b*d*f*x^7)/7

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)(e + fx^2) dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2),x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)*(e + f*x^2), x]

fricas [A] time = 0.63, size = 62, normalized size = 1.11

$$\frac{1}{7}x^7fdb + \frac{1}{5}x^5edb + \frac{1}{5}x^5fcb + \frac{1}{5}x^5fda + \frac{1}{3}x^3ecb + \frac{1}{3}x^3eda + \frac{1}{3}x^3fca + xeca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e),x, algorithm="fricas")

[Out] $\frac{1}{7}bdfx^7 + \frac{1}{5}bcfx^5 + \frac{1}{5}adfx^5 + \frac{1}{5}bdx^5e + \frac{1}{3}acfx^3 + \frac{1}{3}bcx^3e + \frac{1}{3}adx^3e + acex$

giac [A] time = 0.34, size = 66, normalized size = 1.18

$$\frac{1}{7}bdfx^7 + \frac{1}{5}bcfx^5 + \frac{1}{5}adfx^5 + \frac{1}{5}bdx^5e + \frac{1}{3}acfx^3 + \frac{1}{3}bcx^3e + \frac{1}{3}adx^3e + acex$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e),x, algorithm="giac")

[Out] $\frac{1}{7}b*d*f*x^7 + \frac{1}{5}b*c*f*x^5 + \frac{1}{5}a*d*f*x^5 + \frac{1}{5}b*d*x^5*e + \frac{1}{3}a*c*f*x^3 + \frac{1}{3}b*c*x^3*e + \frac{1}{3}a*d*x^3*e + a*c*x*e$

maple [A] time = 0.00, size = 53, normalized size = 0.95

$$\frac{bdfx^7}{7} + \frac{(bde + (ad + bc)f)x^5}{5} + acex + \frac{(acf + (ad + bc)e)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)*(f*x^2+e),x)

[Out] $\frac{1}{7}b*d*f*x^7 + \frac{1}{5}((a*d+b*c)*f+b*d*e)*x^5 + \frac{1}{3}((a*c*f+(a*d+b*c)*e)*x^3 + a*c*e*x$

maxima [A] time = 0.55, size = 52, normalized size = 0.93

$$\frac{1}{7}bdfx^7 + \frac{1}{5}(bde + (bc + ad)f)x^5 + acex + \frac{1}{3}(acf + (bc + ad)e)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)*(f*x^2+e),x, algorithm="maxima")

[Out] $\frac{1}{7}b*d*f*x^7 + \frac{1}{5}(b*d*e + (b*c + a*d)*f)*x^5 + a*c*e*x + \frac{1}{3}((a*c*f + (b*c + a*d)*e)*x^3$

mupad [B] time = 0.05, size = 54, normalized size = 0.96

$$\frac{bdfx^7}{7} + \left(\frac{adf}{5} + \frac{bcf}{5} + \frac{bde}{5}\right)x^5 + \left(\frac{acf}{3} + \frac{ade}{3} + \frac{bce}{3}\right)x^3 + acex$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)*(e + f*x^2),x)

[Out] $x^3((a*c*f)/3 + (a*d*e)/3 + (b*c*e)/3) + x^5((a*d*f)/5 + (b*c*f)/5 + (b*d*e)/5) + a*c*e*x + (b*d*f*x^7)/7$

sympy [A] time = 0.07, size = 63, normalized size = 1.12

$$acex + \frac{bdfx^7}{7} + x^5\left(\frac{adf}{5} + \frac{bcf}{5} + \frac{bde}{5}\right) + x^3\left(\frac{acf}{3} + \frac{ade}{3} + \frac{bce}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)*(f*x**2+e),x)

[Out] $a*c*e*x + b*d*f*x**7/7 + x**5*(a*d*f/5 + b*c*f/5 + b*d*e/5) + x**3*(a*c*f/3 + a*d*e/3 + b*c*e/3)$

$$3.5 \quad \int \frac{(a+bx^2)(c+dx^2)}{e+fx^2} dx$$

Optimal. Leaf size=81

$$\frac{(be-af)(de-cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e} f^{5/2}} - \frac{x(-2adf-3bcf+3bde)}{3f^2} + \frac{dx(a+bx^2)}{3f}$$

Rubi [A] time = 0.08, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {528, 388, 205}

$$-\frac{x(-2adf-3bcf+3bde)}{3f^2} + \frac{(be-af)(de-cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e} f^{5/2}} + \frac{dx(a+bx^2)}{3f}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2))/(e + f*x^2), x]

[Out] -((3*b*d*e - 3*b*c*f - 2*a*d*f)*x)/(3*f^2) + (d*x*(a + b*x^2))/(3*f) + ((b*e - a*f)*(d*e - c*f)*ArcTan[(Sqrt[f]*x)/Sqrt[e]]/(Sqrt[e]*f^(5/2)))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(b*(n*(p+q+1)+1)), x] + Dist[1/(b*(n*(p+q+1)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-1)*Simp[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p+q+1)+1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)(c+dx^2)}{e+fx^2} dx &= \frac{dx(a+bx^2)}{3f} + \frac{\int \frac{-a(de-3cf)-(3bde-3bcf-2adf)x^2}{e+fx^2} dx}{3f} \\ &= -\frac{(3bde-3bcf-2adf)x}{3f^2} + \frac{dx(a+bx^2)}{3f} + \frac{((be-af)(de-cf)) \int \frac{1}{e+fx^2} dx}{f^2} \\ &= -\frac{(3bde-3bcf-2adf)x}{3f^2} + \frac{dx(a+bx^2)}{3f} + \frac{(be-af)(de-cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e} f^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 72, normalized size = 0.89

$$\frac{(be - af)(de - cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e} f^{5/2}} + \frac{x(adf + bcf - bde)}{f^2} + \frac{bdx^3}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2))/(e + f*x^2), x]

[Out] ((-(b*d*e) + b*c*f + a*d*f)*x)/f^2 + (b*d*x^3)/(3*f) + ((b*e - a*f)*(d*e - c*f)*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(Sqrt[e]*f^(5/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)}{e + fx^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2), x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2), x]

fricas [A] time = 1.42, size = 191, normalized size = 2.36

$$\left[\frac{2bdef^2x^3 - 3(bde^2 + acf^2 - (bc + ad)ef)\sqrt{-ef} \log\left(\frac{fx^2 - \sqrt{-ef}x - e}{fx^2 + e}\right) - 6(bde^2f - (bc + ad)ef^2)x}{6ef^3}, \frac{bdef^2x^3 + 3(bde^2 + acf^2 - (bc + ad)ef)\sqrt{ef} \arctan\left(\frac{\sqrt{ef}x}{e}\right) - 3(bde^2f - (bc + ad)ef^2)x}{3ef^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e), x, algorithm="fricas")

[Out] [1/6*(2*b*d*e*f^2*x^3 - 3*(b*d*e^2 + a*c*f^2 - (b*c + a*d)*e*f)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 6*(b*d*e^2*f - (b*c + a*d)*e*f^2)*x)/(e*f^3), 1/3*(b*d*e*f^2*x^3 + 3*(b*d*e^2 + a*c*f^2 - (b*c + a*d)*e*f)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) - 3*(b*d*e^2*f - (b*c + a*d)*e*f^2)*x)/(e*f^3)]

giac [A] time = 0.31, size = 80, normalized size = 0.99

$$\frac{(acf^2 - bcfe - adfe + bde^2) \arctan\left(\sqrt{f}xe^{\left(-\frac{1}{2}\right)}\right)e^{\left(-\frac{1}{2}\right)}}{f^{\frac{5}{2}}} + \frac{bdf^2x^3 + 3bcf^2x + 3adf^2x - 3bdfxe}{3f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e), x, algorithm="giac")

[Out] (a*c*f^2 - b*c*f*e - a*d*f*e + b*d*e^2)*arctan(sqrt(f)*x*e^(-1/2))*e^(-1/2)/f^(5/2) + 1/3*(b*d*f^2*x^3 + 3*b*c*f^2*x + 3*a*d*f^2*x - 3*b*d*f*x*e)/f^3

maple [A] time = 0.01, size = 119, normalized size = 1.47

$$\frac{bdx^3}{3f} + \frac{ac \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}} - \frac{ade \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef} f} - \frac{bce \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef} f} + \frac{bd e^2 \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef} f^2} + \frac{adx}{f} + \frac{bcx}{f} - \frac{bdex}{f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)/(f*x^2+e), x)

[Out] $1/3/f*x^3*b*d+1/f*a*d*x+1/f*b*c*x-1/f^2*b*d*e*x+1/(e*f)^{(1/2)}*\arctan(f*x/(e*f)^{(1/2)})*a*c-1/f/(e*f)^{(1/2)}*\arctan(f*x/(e*f)^{(1/2)})*a*d*e-1/f/(e*f)^{(1/2)}*\arctan(f*x/(e*f)^{(1/2)})*b*c*e+1/f^2/(e*f)^{(1/2)}*\arctan(f*x/(e*f)^{(1/2)})*b*d*e^2$

maxima [A] time = 1.24, size = 74, normalized size = 0.91

$$\frac{(bde^2 + acf^2 - (bc + ad)ef) \arctan\left(\frac{fx}{\sqrt{ef}}\right) + bdfx^3 - 3(bde - (bc + ad)f)x}{\sqrt{ef} f^2} + \frac{bdfx^3 - 3(bde - (bc + ad)f)x}{3f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e),x, algorithm="maxima")`

[Out] $(b*d*e^2 + a*c*f^2 - (b*c + a*d)*e*f)*\arctan(f*x/\sqrt{e*f})/(\sqrt{e*f}*f^2) + 1/3*(b*d*f*x^3 - 3*(b*d*e - (b*c + a*d)*f)*x)/f^2$

mupad [B] time = 0.11, size = 108, normalized size = 1.33

$$x \left(\frac{ad + bc}{f} - \frac{bde}{f^2} \right) + \frac{bdx^3}{3f} + \frac{\operatorname{atan}\left(\frac{\sqrt{f}x(a f - b e)(c f - d e)}{\sqrt{e}(a c f^2 + b d e^2 - a d e f - b c e f)}\right) (a f - b e)(c f - d e)}{\sqrt{e} f^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((a + b*x^2)*(c + d*x^2))/(e + f*x^2),x)`

[Out] $x*((a*d + b*c)/f - (b*d*e)/f^2) + (b*d*x^3)/(3*f) + (\operatorname{atan}((f^{(1/2)}*x*(a*f - b*e)*(c*f - d*e))/(e^{(1/2)}*(a*c*f^2 + b*d*e^2 - a*d*e*f - b*c*e*f)))* (a*f - b*e)*(c*f - d*e))/(e^{(1/2)}*f^{(5/2)})$

sympy [B] time = 0.62, size = 206, normalized size = 2.54

$$\frac{bdx^3}{3f} + x \left(\frac{ad}{f} + \frac{bc}{f} - \frac{bde}{f^2} \right) - \frac{\sqrt{-\frac{1}{ef^5}}(af - be)(cf - de) \log\left(-\frac{ef^2 \sqrt{-\frac{1}{ef^5}}(af - be)(cf - de)}{acf^2 - adef - bcef + bde^2} + x\right)}{2} + \frac{\sqrt{-\frac{1}{ef^5}}(af - be)(cf - de) \log\left(\frac{ef^2 \sqrt{-\frac{1}{ef^5}}(af - be)(cf - de)}{acf^2 - adef - bcef + bde^2} + x\right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)*(d*x**2+c)/(f*x**2+e),x)`

[Out] $b*d*x**3/(3*f) + x*(a*d/f + b*c/f - b*d*e/f**2) - \sqrt{-1/(e*f**5)}*(a*f - b*e)*(c*f - d*e)*\log(-e*f**2*\sqrt{-1/(e*f**5)}*(a*f - b*e)*(c*f - d*e)/(a*c*f**2 - a*d*e*f - b*c*e*f + b*d*e**2) + x)/2 + \sqrt{-1/(e*f**5)}*(a*f - b*e)*(c*f - d*e)*\log(e*f**2*\sqrt{-1/(e*f**5)}*(a*f - b*e)*(c*f - d*e)/(a*c*f**2 - a*d*e*f - b*c*e*f + b*d*e**2) + x)/2$

$$3.6 \quad \int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^2} dx$$

Optimal. Leaf size=108

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(3de-cf)-af(cf+de))}{2e^{3/2}f^{5/2}} - \frac{x(a+bx^2)(de-cf)}{2ef(e+fx^2)} + \frac{bx(3de-cf)}{2ef^2}$$

Rubi [A] time = 0.08, antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {526, 388, 205}

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(3de-cf)-af(cf+de))}{2e^{3/2}f^{5/2}} - \frac{x(a+bx^2)(de-cf)}{2ef(e+fx^2)} + \frac{bx(3de-cf)}{2ef^2}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^2,x]

[Out] (b*(3*d*e - c*f)*x)/(2*e*f^2) - ((d*e - c*f)*x*(a + b*x^2))/(2*e*f*(e + f*x^2)) - ((b*e*(3*d*e - c*f) - a*f*(d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(2*e^(3/2)*f^(5/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*b*n*(p+1)), x] + Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)*Simp[c*(b*e*n*(p+1) + b*e - a*f) + d*(b*e*n*(p+1) + (b*e - a*f)*(n*q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^2} dx &= -\frac{(de-cf)x(a+bx^2)}{2ef(e+fx^2)} - \frac{\int \frac{-a(de+cf)-b(3de-cf)x^2}{e+fx^2} dx}{2ef} \\ &= \frac{b(3de-cf)x}{2ef^2} - \frac{(de-cf)x(a+bx^2)}{2ef(e+fx^2)} - \frac{(be(3de-cf)-af(de+cf)) \int \frac{1}{e+fx^2} dx}{2ef^2} \\ &= \frac{b(3de-cf)x}{2ef^2} - \frac{(de-cf)x(a+bx^2)}{2ef(e+fx^2)} - \frac{(be(3de-cf)-af(de+cf)) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{2e^{3/2}f^{5/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 95, normalized size = 0.88

$$-\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(3de - cf) - af(cf + de))}{2e^{3/2}f^{5/2}} + \frac{x(be - af)(de - cf)}{2ef^2(e + fx^2)} + \frac{bdx}{f^2}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^2,x]

[Out] (b*d*x)/f^2 + ((b*e - a*f)*(d*e - c*f)*x)/(2*e*f^2*(e + f*x^2)) - ((b*e*(3*d*e - c*f) - a*f*(d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(2*e^(3/2)*f^(5/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)}{(e + fx^2)^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^2, x]

fricas [A] time = 1.44, size = 318, normalized size = 2.94

$$\frac{4bd^2f^2x^3 + (3bde^3 - acef^2 - (bc + ad)d^2f + (3bd^2f - acf^2 - (bc + ad)ef^2)x^2)\sqrt{ef}\log\left(\frac{x^2 - 2\sqrt{ef}x + e}{f^2 + ex}\right) + 2(3bde^3f + acef^3 - (bc + ad)d^2f^2)x + 2bd^2f^2x^3 - (3bde^3 - acef^2 - (bc + ad)d^2f + (3bd^2f - acf^2 - (bc + ad)ef^2)x^2)\sqrt{ef}\arctan\left(\frac{\sqrt{ef}x}{e}\right) + (3bd^2f + acef^3 - (bc + ad)d^2f)x}{4(e^2fx^2 + e^2f^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^2,x, algorithm="fricas")

[Out] [1/4*(4*b*d*e^2*f^2*x^3 + (3*b*d*e^3 - a*c*e*f^2 - (b*c + a*d)*e^2*f + (3*b*d*e^2*f - a*c*f^3 - (b*c + a*d)*e*f^2)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) + 2*(3*b*d*e^3*f + a*c*e*f^3 - (b*c + a*d)*e^2*f^2)*x)/(e^2*f^4*x^2 + e^3*f^3), 1/2*(2*b*d*e^2*f^2*x^3 - (3*b*d*e^3 - a*c*e*f^2 - (b*c + a*d)*e^2*f + (3*b*d*e^2*f - a*c*f^3 - (b*c + a*d)*e*f^2)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) + (3*b*d*e^3*f + a*c*e*f^3 - (b*c + a*d)*e^2*f^2)*x)/(e^2*f^4*x^2 + e^3*f^3)]

giac [A] time = 0.36, size = 95, normalized size = 0.88

$$\frac{bdx}{f^2} + \frac{(acf^2 + bcfe + adfe - 3bde^2)\arctan\left(\sqrt{f}xe^{\left(-\frac{1}{2}\right)}\right)e^{\left(-\frac{3}{2}\right)}}{2f^{\frac{5}{2}}} + \frac{(acf^2x - bcfxe - adfxe + bdx e^2)e^{(-1)}}{2(fx^2 + e)f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^2,x, algorithm="giac")

[Out] b*d*x/f^2 + 1/2*(a*c*f^2 + b*c*f*e + a*d*f*e - 3*b*d*e^2)*arctan(sqrt(f)*x*e^(-1/2))*e^(-3/2)/f^(5/2) + 1/2*(a*c*f^2*x - b*c*f*x*e - a*d*f*x*e + b*d*x*e^2)*e^(-1)/((f*x^2 + e)*f^2)

maple [A] time = 0.01, size = 163, normalized size = 1.51

$$\frac{acx}{2(fx^2 + e)e} + \frac{ac\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{2\sqrt{ef}e} - \frac{adx}{2(fx^2 + e)f} + \frac{ad\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{2\sqrt{ef}f} - \frac{bcx}{2(fx^2 + e)f} + \frac{bc\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{2\sqrt{ef}f} + \frac{bdex}{2(fx^2 + e)f^2} - \frac{3bde\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{2\sqrt{ef}f^2} + \frac{bdx}{f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^2,x)

[Out] b*d/f^2*x+1/2/e*x/(f*x^2+e)*a*c-1/2/f*x/(f*x^2+e)*a*d-1/2/f*x/(f*x^2+e)*b*c+1/2/f^2*e*x/(f*x^2+e)*b*d+1/2/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c+1/2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d+1/2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c-3/2/f^2*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d

maxima [A] time = 1.20, size = 101, normalized size = 0.94

$$\frac{(bde^2 + acf^2 - (bc + ad)ef)x}{2(ef^3x^2 + e^2f^2)} + \frac{bdx}{f^2} - \frac{(3bde^2 - acf^2 - (bc + ad)ef) \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{2\sqrt{ef}ef^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^2,x, algorithm="maxima")

[Out] 1/2*(b*d*e^2 + a*c*f^2 - (b*c + a*d)*e*f)*x/(e*f^3*x^2 + e^2*f^2) + b*d*x/f^2 - 1/2*(3*b*d*e^2 - a*c*f^2 - (b*c + a*d)*e*f)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*e*f^2)

mupad [B] time = 0.17, size = 95, normalized size = 0.88

$$\frac{bdx}{f^2} + \frac{\operatorname{atan}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right) (acf^2 - 3bde^2 + adef + bcef)}{2e^{3/2}f^{5/2}} + \frac{x(acf^2 + bde^2 - adef - bcef)}{2e(f^3x^2 + ef^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^2,x)

[Out] (b*d*x)/f^2 + (atan((f^(1/2)*x)/e^(1/2))*(a*c*f^2 - 3*b*d*e^2 + a*d*e*f + b*c*e*f))/(2*e^(3/2)*f^(5/2)) + (x*(a*c*f^2 + b*d*e^2 - a*d*e*f - b*c*e*f))/(2*e*(e*f^2 + f^3*x^2))

sympy [A] time = 1.37, size = 190, normalized size = 1.76

$$\frac{bdx}{f^2} + \frac{x(acf^2 - adef - bcef + bde^2)}{2e^2f^2 + 2ef^3x^2} - \frac{\sqrt{-\frac{1}{e^3f^5}}(acf^2 + adef + bcef - 3bde^2) \log\left(-e^2f^2\sqrt{-\frac{1}{e^3f^5}} + x\right)}{4} + \frac{\sqrt{-\frac{1}{e^3f^5}}(acf^2 + adef + bcef - 3bde^2) \log\left(e^2f^2\sqrt{-\frac{1}{e^3f^5}} + x\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)/(f*x**2+e)**2,x)

[Out] b*d*x/f**2 + x*(a*c*f**2 - a*d*e*f - b*c*e*f + b*d*e**2)/(2*e**2*f**2 + 2*e*f**3*x**2) - sqrt(-1/(e**3*f**5))*(a*c*f**2 + a*d*e*f + b*c*e*f - 3*b*d*e**2)*log(-e**2*f**2*sqrt(-1/(e**3*f**5)) + x)/4 + sqrt(-1/(e**3*f**5))*(a*c*f**2 + a*d*e*f + b*c*e*f - 3*b*d*e**2)*log(e**2*f**2*sqrt(-1/(e**3*f**5)) + x)/4

$$3.7 \quad \int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^3} dx$$

Optimal. Leaf size=130

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(3cf+de)+be(cf+3de))}{8e^{5/2}f^{5/2}} - \frac{x(be(cf+3de)-af(3cf+de))}{8e^2f^2(e+fx^2)} - \frac{x(a+bx^2)(de-cf)}{4ef(e+fx^2)^2}$$

Rubi [A] time = 0.11, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {526, 385, 205}

$$-\frac{x(be(cf+3de)-af(3cf+de))}{8e^2f^2(e+fx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(3cf+de)+be(cf+3de))}{8e^{5/2}f^{5/2}} - \frac{x(a+bx^2)(de-cf)}{4ef(e+fx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^3,x]

[Out] -((d*e - c*f)*x*(a + b*x^2))/(4*e*f*(e + f*x^2)^2) - ((b*e*(3*d*e + c*f) - a*f*(d*e + 3*c*f))*x)/(8*e^2*f^2*(e + f*x^2)) + ((b*e*(3*d*e + c*f) + a*f*(d*e + 3*c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(8*e^(5/2)*f^(5/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)(c + dx^2)}{(e + fx^2)^3} dx &= -\frac{(de - cf)x(a + bx^2)}{4ef(e + fx^2)^2} - \frac{\int \frac{-a(de+3cf)-b(3de+cf)x^2}{(e+fx^2)^2} dx}{4ef} \\ &= -\frac{(de - cf)x(a + bx^2)}{4ef(e + fx^2)^2} - \frac{(be(3de + cf) - af(de + 3cf))x}{8e^2 f^2 (e + fx^2)} + \frac{(be(3de + cf) + af(de + 3cf))}{8e^2 f^2} \\ &= -\frac{(de - cf)x(a + bx^2)}{4ef(e + fx^2)^2} - \frac{(be(3de + cf) - af(de + 3cf))x}{8e^2 f^2 (e + fx^2)} + \frac{(be(3de + cf) + af(de + 3cf))}{8e^2 f^2} \end{aligned}$$

Mathematica [A] time = 0.08, size = 130, normalized size = 1.00

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(3cf + de) + be(cf + 3de))}{8e^{5/2}f^{5/2}} + \frac{x(af(3cf + de) + be(cf - 5de))}{8e^2 f^2 (e + fx^2)} + \frac{x(be - af)(de - cf)}{4ef^2 (e + fx^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^3,x]

[Out] ((b*e - a*f)*(d*e - c*f)*x)/(4*e*f^2*(e + f*x^2)^2) + ((b*e*(-5*d*e + c*f) + a*f*(d*e + 3*c*f))*x)/(8*e^2*f^2*(e + f*x^2)) + ((b*e*(3*d*e + c*f) + a*f*(d*e + 3*c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(8*e^(5/2)*f^(5/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)}{(e + fx^2)^3} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^3, x]

fricas [A] time = 1.34, size = 471, normalized size = 3.62

$$\frac{2(3ab^2f^2 - 3acdf^2 - (bc + ad^2f^2))x^4 + (3abd^2 + 3acdf^2 + (bc + ad^2f^2)^2 + 2(3abd^2 - 5acdf^2 - (bc + ad^2f^2)^2))\sqrt{e}\arctan\left(\frac{\sqrt{f}x}{\sqrt{e}}\right) + 2(3abd^2f - 5acdf^2 + (bc + ad^2f^2)^2)}{16(f^2f^2 + 2ef^2 + f^2)^2} - \frac{(3ab^2f^2 - 3acdf^2 - (bc + ad^2f^2)^2 - (3abd^2 + 3acdf^2 + (bc + ad^2f^2)^2))\sqrt{e}\arctan\left(\frac{\sqrt{f}x}{\sqrt{e}}\right) + (3abd^2f - 5acdf^2 + (bc + ad^2f^2)^2)}{8(f^2f^2 + 2ef^2 + f^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^3,x, algorithm="fricas")

[Out] [-1/16*(2*(5*b*d*e^3*f^2 - 3*a*c*e*f^4 - (b*c + a*d)*e^2*f^3)*x^3 + (3*b*d*e^4 + 3*a*c*e^2*f^2 + (b*c + a*d)*e^3*f + (3*b*d*e^2*f^2 + 3*a*c*f^4 + (b*c + a*d)*e*f^3)*x^4 + 2*(3*b*d*e^3*f + 3*a*c*e*f^3 + (b*c + a*d)*e^2*f^2)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) + 2*(3*b*d*e^4*f - 5*a*c*e^2*f^3 + (b*c + a*d)*e^3*f^2)*x)/(e^3*f^5*x^4 + 2*e^4*f^4*x^2 + e^5*f^3), -1/8*((5*b*d*e^3*f^2 - 3*a*c*e*f^4 - (b*c + a*d)*e^2*f^3)*x^3 - (3*b*d*e^4 + 3*a*c*e^2*f^2 + (b*c + a*d)*e^3*f + (3*b*d*e^2*f^2 + 3*a*c*f^4 + (b*c + a*d)*e*f^3)*x^4 + 2*(3*b*d*e^3*f + 3*a*c*e*f^3 + (b*c + a*d)*e^2*f^2)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) + (3*b*d*e^4*f - 5*a*c*e^2*f^3 + (b*c + a*d)*e^3*f^2)*x)/(e^3*f^5*x^4 + 2*e^4*f^4*x^2 + e^5*f^3)]

giac [A] time = 0.46, size = 135, normalized size = 1.04

$$\frac{(3acf^2 + bcfe + adfe + 3bde^2)\arctan\left(\sqrt{f}xe^{\left(\frac{-1}{2}\right)}\right)e^{\left(\frac{-5}{2}\right)}}{8f^{\frac{5}{2}}} + \frac{(3acf^3x^3 + bcf^2x^3e + adf^2x^3e - 5bdfx^3e^2 + 5acf^2xe - bcfxe^2 - adfxe^2 - 3bdxe^3)e^{(-2)}}{8(fx^2 + e)^2f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^3,x, algorithm="giac")

[Out] $\frac{1}{8}*(3*a*c*f^2 + b*c*f*e + a*d*f*e + 3*b*d*e^2)*\arctan(\sqrt{f}*x*e^{(-1/2)})*e^{(-5/2)}/f^{(5/2)} + \frac{1}{8}*(3*a*c*f^3*x^3 + b*c*f^2*x^3*e + a*d*f^2*x^3*e - 5*b*d*f*x^3*e^2 + 5*a*c*f^2*x*e - b*c*f*x*e^2 - a*d*f*x*e^2 - 3*b*d*x*e^3)*e^{(-2)}/((f*x^2 + e)^2*f^2)$

maple [A] time = 0.01, size = 175, normalized size = 1.35

$$\frac{3ac \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{8\sqrt{ef} e^2} + \frac{ad \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{8\sqrt{ef} ef} + \frac{bc \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{8\sqrt{ef} ef} + \frac{3bd \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{8\sqrt{ef} f^2} + \frac{\frac{(3ac f^2 + adef + bcef - 5bd e^2)x^3}{8e^2 f} + \frac{(5ac f^2 - adef - bcef - 3bd e^2)x}{8e f^2}}{(f x^2 + e)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^3,x)

[Out] $(1/8*(3*a*c*f^2+a*d*e*f+b*c*e*f-5*b*d*e^2)/e^2/f*x^3+1/8*(5*a*c*f^2-a*d*e*f-b*c*e*f-3*b*d*e^2)/f^2/e*x)/(f*x^2+e)^2+3/8/e^2/(e*f)^{(1/2)}*\arctan(1/(e*f)^{(1/2)}*f*x)*a*c+1/8/e/f/(e*f)^{(1/2)}*\arctan(1/(e*f)^{(1/2)}*f*x)*a*d+1/8/e/f/(e*f)^{(1/2)}*\arctan(1/(e*f)^{(1/2)}*f*x)*b*c+3/8/f^2/(e*f)^{(1/2)}*\arctan(1/(e*f)^{(1/2)}*f*x)*b*d$

maxima [A] time = 1.43, size = 145, normalized size = 1.12

$$\frac{(5 b d e^2 f - 3 a c f^3 - (b c + a d) e f^2) x^3 + (3 b d e^3 - 5 a c e f^2 + (b c + a d) e^2 f) x}{8 (e^2 f^4 x^4 + 2 e^3 f^3 x^2 + e^4 f^2)} + \frac{(3 b d e^2 + 3 a c f^2 + (b c + a d) e f) \arctan\left(\frac{f x}{\sqrt{e f}}\right)}{8 \sqrt{e f} e^2 f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^3,x, algorithm="maxima")

[Out] $-1/8*((5*b*d*e^2*f - 3*a*c*f^3 - (b*c + a*d)*e*f^2)*x^3 + (3*b*d*e^3 - 5*a*c*e*f^2 + (b*c + a*d)*e^2*f)*x)/(e^2*f^4*x^4 + 2*e^3*f^3*x^2 + e^4*f^2) + 1/8*(3*b*d*e^2 + 3*a*c*f^2 + (b*c + a*d)*e*f)*\arctan(f*x/\sqrt{e*f})/(\sqrt{e*f}*e^2*f^2)$

mupad [B] time = 0.97, size = 136, normalized size = 1.05

$$\frac{\operatorname{atan}\left(\frac{\sqrt{f} x}{\sqrt{e}}\right) (3 a c f^2 + 3 b d e^2 + a d e f + b c e f)}{8 e^{5/2} f^{5/2}} - \frac{x (3 b d e^2 - 5 a c f^2 + a d e f + b c e f)}{8 e f^2} - \frac{x^3 (3 a c f^2 - 5 b d e^2 + a d e f + b c e f)}{8 e^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^3,x)

[Out] $(\operatorname{atan}((f^{(1/2)}*x)/e^{(1/2)})*(3*a*c*f^2 + 3*b*d*e^2 + a*d*e*f + b*c*e*f))/(8*e^{(5/2)}*f^{(5/2)}) - ((x*(3*b*d*e^2 - 5*a*c*f^2 + a*d*e*f + b*c*e*f))/(8*e*f^2) - (x^3*(3*a*c*f^2 - 5*b*d*e^2 + a*d*e*f + b*c*e*f))/(8*e^2*f))/(e^2 + f^2*x^4 + 2*e*f*x^2)$

sympy [B] time = 3.13, size = 246, normalized size = 1.89

$$\frac{\sqrt{-\frac{1}{e f^5}} (3 a c f^2 + a d e f + b c e f + 3 b d e^2) \log\left(-e^3 f^2 \sqrt{-\frac{1}{e f^5}} + x\right)}{16} + \frac{\sqrt{-\frac{1}{e f^5}} (3 a c f^2 + a d e f + b c e f + 3 b d e^2) \log\left(e^3 f^2 \sqrt{-\frac{1}{e f^5}} + x\right)}{16} + \frac{x^3 (3 a c f^3 + a d e f^2 + b c e f^2 - 5 b d e^2 f) + x (5 a c e f^2 - a d e^2 f - b c e^2 f - 3 b d e^3)}{8 e^4 f^2 + 16 e^3 f^3 x^2 + 8 e^2 f^4 x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)/(f*x**2+e)**3,x)

[Out] $-\sqrt{-1/(e**5*f**5)}*(3*a*c*f**2 + a*d*e*f + b*c*e*f + 3*b*d*e**2)*\log(-e**3*f**2*\sqrt{-1/(e**5*f**5)} + x)/16 + \sqrt{-1/(e**5*f**5)}*(3*a*c*f**2 + a$

```
*d*e*f + b*c*e*f + 3*b*d*e**2)*log(e**3*f**2*sqrt(-1/(e**5*f**5)) + x)/16 +  
(x**3*(3*a*c*f**3 + a*d*e*f**2 + b*c*e*f**2 - 5*b*d*e**2*f) + x*(5*a*c*e*f  
**2 - a*d*e**2*f - b*c*e**2*f - 3*b*d*e**3))/(8*e**4*f**2 + 16*e**3*f**3*x*  
*2 + 8*e**2*f**4*x**4)
```

$$3.8 \quad \int \frac{(a+bx^2)(c+dx^2)}{(e+fx^2)^4} dx$$

Optimal. Leaf size=171

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(5cf+de)+be(cf+de))}{16e^{7/2}f^{5/2}} + \frac{x(af(5cf+de)+be(cf+de))}{16e^3f^2(e+fx^2)} - \frac{x(3be(cf+de)-af(5cf+de))}{24e^2f^2(e+fx^2)^2} - \frac{x(a+bx^2)(de-cf)}{6ef(e+fx^2)^3}$$

Rubi [A] time = 0.16, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {526, 385, 199, 205}

$$\frac{x(af(5cf+de)+be(cf+de))}{16e^3f^2(e+fx^2)} - \frac{x(3be(cf+de)-af(5cf+de))}{24e^2f^2(e+fx^2)^2} + \frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(5cf+de)+be(cf+de))}{16e^{7/2}f^{5/2}} - \frac{x(a+bx^2)(de-cf)}{6ef(e+fx^2)^3}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^4, x]

[Out] -((d*e - c*f)*x*(a + b*x^2))/(6*e*f*(e + f*x^2)^3) - ((3*b*e*(d*e + c*f) - a*f*(d*e + 5*c*f))*x)/(24*e^2*f^2*(e + f*x^2)^2) + ((b*e*(d*e + c*f) + a*f*(d*e + 5*c*f))*x)/(16*e^3*f^2*(e + f*x^2)) + ((b*e*(d*e + c*f) + a*f*(d*e + 5*c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(16*e^(7/2)*f^(5/2))

Rule 199

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p + 1))/(a*b*n*(p + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)}{(e + fx^2)^4} dx &= -\frac{(de - cf)x(a + bx^2)}{6ef(e + fx^2)^3} - \frac{\int \frac{-a(de+5cf)-3b(de+cf)x^2}{(e+fx^2)^3} dx}{6ef} \\
&= -\frac{(de - cf)x(a + bx^2)}{6ef(e + fx^2)^3} - \frac{(3be(de + cf) - af(de + 5cf))x}{24e^2f^2(e + fx^2)^2} + \frac{(be(de + cf) + af(de + 5cf))}{8e^2f^2} \\
&= -\frac{(de - cf)x(a + bx^2)}{6ef(e + fx^2)^3} - \frac{(3be(de + cf) - af(de + 5cf))x}{24e^2f^2(e + fx^2)^2} + \frac{(be(de + cf) + af(de + 5cf))}{16e^3f^2(e + fx^2)} \\
&= -\frac{(de - cf)x(a + bx^2)}{6ef(e + fx^2)^3} - \frac{(3be(de + cf) - af(de + 5cf))x}{24e^2f^2(e + fx^2)^2} + \frac{(be(de + cf) + af(de + 5cf))}{16e^3f^2(e + fx^2)}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 171, normalized size = 1.00

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(5cf + de) + be(cf + de))}{16e^{7/2}f^{5/2}} + \frac{x(af(5cf + de) + be(cf + de))}{16e^3f^2(e + fx^2)} + \frac{x(af(5cf + de) + be(cf - 7de))}{24e^2f^2(e + fx^2)^2} + \frac{x(be - af)(de - cf)}{6ef^2(e + fx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^4,x]

[Out] ((b*e - a*f)*(d*e - c*f)*x)/(6*e*f^2*(e + f*x^2)^3) + ((b*e*(-7*d*e + c*f) + a*f*(d*e + 5*c*f))*x)/(24*e^2*f^2*(e + f*x^2)^2) + ((b*e*(d*e + c*f) + a*f*(d*e + 5*c*f))*x)/(16*e^3*f^2*(e + f*x^2)) + ((b*e*(d*e + c*f) + a*f*(d*e + 5*c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(16*e^(7/2)*f^(5/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)}{(e + fx^2)^4} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^4,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^4, x]

fricas [B] time = 1.47, size = 642, normalized size = 3.75

fricas

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^4,x, algorithm="fricas")

[Out] [1/96*(6*(b*d*e^3*f^3 + 5*a*c*e*f^5 + (b*c + a*d)*e^2*f^4)*x^5 - 16*(b*d*e^4*f^2 - 5*a*c*e^2*f^4 - (b*c + a*d)*e^3*f^3)*x^3 - 3*(b*d*e^5 + 5*a*c*e^3*f^2 + (b*d*e^2*f^3 + 5*a*c*f^5 + (b*c + a*d)*e*f^4)*x^6 + (b*c + a*d)*e^4*f + 3*(b*d*e^3*f^2 + 5*a*c*e*f^4 + (b*c + a*d)*e^2*f^3)*x^4 + 3*(b*d*e^4*f + 5*a*c*e^2*f^3 + (b*c + a*d)*e^3*f^2)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 6*(b*d*e^5*f - 11*a*c*e^3*f^3 + (b*c + a*d)*e^4*f^2)*x)/(e^4*f^6*x^6 + 3*e^5*f^5*x^4 + 3*e^6*f^4*x^2 + e^7*f^3), 1/48*(3*(b*d*e^3*f^3 + 5*a*c*e*f^5 + (b*c + a*d)*e^2*f^4)*x^5 - 8*(b*d*e^4*f^2 - 5*a*c

$$*e^2*f^4 - (b*c + a*d)*e^3*f^3)*x^3 + 3*(b*d*e^5 + 5*a*c*e^3*f^2 + (b*d*e^2*f^3 + 5*a*c*f^5 + (b*c + a*d)*e*f^4)*x^6 + (b*c + a*d)*e^4*f + 3*(b*d*e^3*f^2 + 5*a*c*e*f^4 + (b*c + a*d)*e^2*f^3)*x^4 + 3*(b*d*e^4*f + 5*a*c*e^2*f^3 + (b*c + a*d)*e^3*f^2)*x^2)*\sqrt{e*f}*\arctan(\sqrt{e*f}*x/e) - 3*(b*d*e^5*f - 11*a*c*e^3*f^3 + (b*c + a*d)*e^4*f^2)*x)/(e^4*f^6*x^6 + 3*e^5*f^5*x^4 + 3*e^6*f^4*x^2 + e^7*f^3)]$$

giac [A] time = 0.34, size = 184, normalized size = 1.08

$$\frac{(5acf^2 + bcfe + adfe + bde^2)\arctan\left(\sqrt{f}xe^{\left(-\frac{1}{2}\right)}\right)e^{\left(-\frac{3}{2}\right)}}{16f^{\frac{5}{2}}} + \frac{(15acf^4x^5 + 3bcf^3x^5e + 3adf^3x^5e + 3bdf^2x^5e^2 + 40acf^3x^3e + 8bcf^2x^3e^2 + 8adf^2x^3e^2 - 8bdfx^3e^3 + 33acf^2e^2 - 3bcfx^2e^3 - 3adfx^2e^3 - 3bdxe^4)e^{(-3)}}{48(fx^2 + e)^3f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^4,x, algorithm="giac")

[Out] 1/16*(5*a*c*f^2 + b*c*f*e + a*d*f*e + b*d*e^2)*arctan(sqrt(f)*x*e^(-1/2))*e^(-7/2)/f^(5/2) + 1/48*(15*a*c*f^4*x^5 + 3*b*c*f^3*x^5*e + 3*a*d*f^3*x^5*e + 3*b*d*f^2*x^5*e^2 + 40*a*c*f^3*x^3*e + 8*b*c*f^2*x^3*e^2 + 8*a*d*f^2*x^3*e^2 - 8*b*d*f*x^3*e^3 + 33*a*c*f^2*x*e^2 - 3*b*c*f*x*e^3 - 3*a*d*f*x*e^3 - 3*b*d*x*e^4)*e^(-3)/((f*x^2 + e)^3*f^2)

maple [A] time = 0.01, size = 210, normalized size = 1.23

$$\frac{5ac \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef} e^3} + \frac{ad \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef} e^2f} + \frac{bc \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef} e^2f} + \frac{bd \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef} e f^2} + \frac{(5ac f^2 + adef + bcef + bde^2)x^5}{16e^3} + \frac{(5ac f^2 + adef + bcef - bde^2)x^3}{6e^2f} + \frac{(11ac f^2 - adef - bcef - bde^2)x}{16e f^2} + \frac{1}{(f x^2 + e)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^4,x)

[Out] (1/16*(5*a*c*f^2+a*d*e*f+b*c*e*f+b*d*e^2)/e^3*x^5+1/6*(5*a*c*f^2+a*d*e*f+b*c*e*f-b*d*e^2)/e^2/f*x^3+1/16*(11*a*c*f^2-a*d*e*f-b*c*e*f-b*d*e^2)/e/f^2*x)/(f*x^2+e)^3+5/16/e^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c+1/16/e^2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d+1/16/e^2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c+1/16/e/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d

maxima [A] time = 1.57, size = 194, normalized size = 1.13

$$\frac{3(bde^2f^2 + 5acf^4 + (bc + ad)ef^3)x^5 - 8(bde^3f - 5acef^3 - (bc + ad)e^2f^2)x^3 - 3(bde^4 - 11ace^2f^2 + (bc + ad)e^3f)x}{48(e^3f^5x^6 + 3e^4f^4x^4 + 3e^5f^3x^2 + e^6f^2)} + \frac{(bde^2 + 5acf^2 + (bc + ad)ef)\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef}e^3f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)/(f*x^2+e)^4,x, algorithm="maxima")

[Out] 1/48*(3*(b*d*e^2*f^2 + 5*a*c*f^4 + (b*c + a*d)*e*f^3)*x^5 - 8*(b*d*e^3*f - 5*a*c*e*f^3 - (b*c + a*d)*e^2*f^2)*x^3 - 3*(b*d*e^4 - 11*a*c*e^2*f^2 + (b*c + a*d)*e^3*f)*x)/(e^3*f^5*x^6 + 3*e^4*f^4*x^4 + 3*e^5*f^3*x^2 + e^6*f^2) + 1/16*(b*d*e^2 + 5*a*c*f^2 + (b*c + a*d)*e*f)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*e^3*f^2)

mupad [B] time = 0.18, size = 176, normalized size = 1.03

$$\frac{x^5(5ac f^2 + bde^2 + adef + bcef)}{16e^3} - \frac{x(bde^2 - 11acf^2 + adef + bcef)}{16ef^2} + \frac{x^3(5ac f^2 - bde^2 + adef + bcef)}{6e^2f} + \frac{\operatorname{atan}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(5ac f^2 + bde^2 + adef + bcef)}{16e^{7/2}f^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2))/(e + f*x^2)^4,x)

[Out] ((x^5*(5*a*c*f^2 + b*d*e^2 + a*d*e*f + b*c*e*f))/(16*e^3) - (x*(b*d*e^2 - 11*a*c*f^2 + a*d*e*f + b*c*e*f))/(16*e*f^2) + (x^3*(5*a*c*f^2 - b*d*e^2 + a

$$\frac{d*ef + b*c*ef)}{(6*e^2*f)))/(e^3 + f^3*x^6 + 3*e^2*f*x^2 + 3*e*f^2*x^4) +$$

$$(\text{atan}((f^{(1/2)}*x)/e^{(1/2)})*(5*a*c*f^2 + b*d*e^2 + a*d*ef + b*c*ef))/(16*e^{(7/2)}*f^{(5/2)})$$

sympy [A] time = 6.66, size = 313, normalized size = 1.83

$$\frac{\sqrt{-\frac{1}{2f}} (5acf^2 + adef + bce^2 + bde^2) \log\left(-e^4 f^2 \sqrt{-\frac{1}{2f}} + x\right) + \sqrt{-\frac{1}{2f}} (5acf^2 + adef + bce^2 + bde^2) \log\left(e^4 f^2 \sqrt{-\frac{1}{2f}} + x\right)}{32} + \frac{x^2 (15acf^4 + 3adef^3 + 3bce^2 f^2 + 3bde^2 f^2) + x^3 (40acef^3 + 8ade^2 f^2 + 8bce^2 f^2 - 8bde^2 f) + x (33ace^2 f^2 - 3ade^3 f - 3bce^3 f - 3bde^4)}{48e^6 f^2 + 144e^4 f^2 x^2 + 144e^4 f^4 x^4 + 48e^3 f^5 x^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)/(f*x**2+e)**4,x)

[Out] $-\sqrt{-1/(e**7*f**5)}*(5*a*c*f**2 + a*d*ef + b*c*ef + b*d*e**2)*\log(-e**4*f**2*\sqrt{-1/(e**7*f**5)} + x)/32 + \sqrt{-1/(e**7*f**5)}*(5*a*c*f**2 + a*d*ef + b*c*ef + b*d*e**2)*\log(e**4*f**2*\sqrt{-1/(e**7*f**5)} + x)/32 + (x**5*(15*a*c*f**4 + 3*a*d*ef**3 + 3*b*c*ef**3 + 3*b*d*e**2*f**2) + x**3*(40*a*c*ef**3 + 8*a*d*e**2*f**2 + 8*b*c*e**2*f**2 - 8*b*d*e**3*f) + x*(33*a*c*e**2*f**2 - 3*a*d*e**3*f - 3*b*c*e**3*f - 3*b*d*e**4))/(48*e**6*f**2 + 144*e**5*f**3*x**2 + 144*e**4*f**4*x**4 + 48*e**3*f**5*x**6)$

$$3.9 \quad \int (a + bx^2)(c + dx^2)^2 (e + fx^2)^3 dx$$

Optimal. Leaf size=226

$$\frac{1}{9}fx^9(adf(2cf + 3de) + b(c^2f^2 + 6cdef + 3d^2e^2)) + \frac{1}{7}x^7(af(c^2f^2 + 6cdef + 3d^2e^2) + be(3c^2f^2 + 6cdef + d^2e^2))$$

Rubi [A] time = 0.21, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {521}

$$\frac{1}{9}fx^9(adf(2cf + 3de) + b(c^2f^2 + 6cdef + 3d^2e^2)) + \frac{1}{7}x^7(af(c^2f^2 + 6cdef + 3d^2e^2) + be(3c^2f^2 + 6cdef + d^2e^2)) + \frac{1}{5}ax^5(a(3c^2f^2 + 6cdef + d^2e^2) + bce(3cf + 2de)) + \frac{1}{3}ce^2x^3(3acf + 2ade + bce) + \frac{1}{11}d^2x^{11}(adf + 2bcf + 3bde) + ac^2e^3x + \frac{1}{13}bd^2fx^{13}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^3,x]

[Out] a*c^2*e^3*x + (c*e^2*(b*c*e + 2*a*d*e + 3*a*c*f)*x^3)/3 + (e*(b*c*e*(2*d*e + 3*c*f) + a*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^5)/5 + ((a*f*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2) + b*e*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^7)/7 + (f*(a*d*f*(3*d*e + 2*c*f) + b*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2))*x^9)/9 + (d*f^2*(3*b*d*e + 2*b*c*f + a*d*f)*x^11)/11 + (b*d^2*f^3*x^13)/13

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^2 (e + fx^2)^3 dx &= \int (ac^2e^3 + ce^2(bce + 2ade + 3acf)x^2 + e(bce(2de + 3cf) + a(d^2e^2 + 6cde^2 + 3c^2e^2))) dx \\ &= ac^2e^3x + \frac{1}{3}ce^2(bce + 2ade + 3acf)x^3 + \frac{1}{5}e(bce(2de + 3cf) + a(d^2e^2 + 6cde^2 + 3c^2e^2))x^5 \end{aligned}$$

Mathematica [A] time = 0.08, size = 226, normalized size = 1.00

$$\frac{1}{9}fx^9(adf(2cf + 3de) + b(c^2f^2 + 6cdef + 3d^2e^2)) + \frac{1}{7}x^7(af(c^2f^2 + 6cdef + 3d^2e^2) + be(3c^2f^2 + 6cdef + d^2e^2)) + \frac{1}{5}ax^5(a(3c^2f^2 + 6cdef + d^2e^2) + bce(3cf + 2de)) + \frac{1}{3}ce^2x^3(3acf + 2ade + bce) + \frac{1}{11}d^2x^{11}(adf + 2bcf + 3bde) + ac^2e^3x + \frac{1}{13}bd^2fx^{13}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^3,x]

[Out] a*c^2*e^3*x + (c*e^2*(b*c*e + 2*a*d*e + 3*a*c*f)*x^3)/3 + (e*(b*c*e*(2*d*e + 3*c*f) + a*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^5)/5 + ((a*f*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2) + b*e*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^7)/7 + (f*(a*d*f*(3*d*e + 2*c*f) + b*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2))*x^9)/9 + (d*f^2*(3*b*d*e + 2*b*c*f + a*d*f)*x^11)/11 + (b*d^2*f^3*x^13)/13

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^2 (e + fx^2)^3 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^3, x]

fricas [A] time = 1.16, size = 289, normalized size = 1.28

$$\frac{1}{13}x^{13}f^3db + \frac{2}{11}x^{11}f^2adfb + \frac{2}{11}x^{11}f^3dcb + \frac{1}{11}x^{11}f^3dca + \frac{1}{3}x^9f^2db + \frac{2}{9}x^9f^2adcb + \frac{1}{9}x^9f^2cb + \frac{1}{3}x^9f^2adca + \frac{2}{9}x^7f^2db + \frac{2}{9}x^7f^2adcb + \frac{2}{9}x^7f^2cb + \frac{2}{9}x^7f^2adca + \frac{6}{7}x^5f^2db + \frac{6}{7}x^5f^2adcb + \frac{6}{7}x^5f^2cb + \frac{6}{7}x^5f^2adca + \frac{1}{7}x^3f^2db + \frac{2}{5}x^3f^2adcb + \frac{2}{5}x^3f^2cb + \frac{1}{5}x^3f^2adca + \frac{3}{5}x^3f^2db + \frac{1}{3}x^3f^2adca + x^3f^2ca + x^3f^2cb$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^3,x, algorithm="fricas")

[Out] $1/13*x^{13}*f^3*d^2*b + 3/11*x^{11}*f^2*e*d^2*b + 2/11*x^{11}*f^3*d*c*b + 1/11*x^{11}*f^3*d^2*a + 1/3*x^9*f^2*e*d^2*b + 2/3*x^9*f^2*e*d*c*b + 1/9*x^9*f^3*c^2*b + 1/3*x^9*f^2*e*d^2*a + 2/9*x^9*f^3*d*c*a + 1/7*x^7*e^3*d^2*b + 6/7*x^7*f^2*e*d*c*b + 3/7*x^7*f^2*e*c^2*b + 3/7*x^7*f^2*e*d^2*a + 6/7*x^7*f^2*e*d*c*a + 1/7*x^7*f^3*c^2*a + 2/5*x^5*e^3*d*c*b + 3/5*x^5*f^2*e*c^2*b + 1/5*x^5*e^3*d^2*a + 6/5*x^5*f^2*e*d*c*a + 3/5*x^5*f^2*e*c^2*a + 1/3*x^3*e^3*c^2*b + 2/3*x^3*e^3*d*c*a + x^3*f^2*e^2*c^2*a + x^3*e^3*c^2*a$

giac [A] time = 0.24, size = 283, normalized size = 1.25

$$\frac{1}{13}bdf^3x^{13} + \frac{2}{11}bcd^2f^3x^{11} + \frac{1}{11}ad^2f^3x^{11} + \frac{3}{11}bdf^3x^{11} + \frac{1}{9}b^2d^2f^3x^9 + \frac{2}{9}ad^2f^3x^9 + \frac{2}{9}bcd^2f^3x^9 + \frac{1}{3}ad^2f^3x^9 + \frac{1}{3}bdf^3x^9 + \frac{1}{3}b^2d^2f^3x^7 + \frac{2}{9}ad^2f^3x^7 + \frac{2}{9}bcd^2f^3x^7 + \frac{6}{7}ad^2f^3x^7 + \frac{6}{7}bdf^3x^7 + \frac{6}{7}b^2d^2f^3x^5 + \frac{1}{7}ad^2f^3x^5 + \frac{2}{5}ad^2f^3x^5 + \frac{3}{5}b^2d^2f^3x^5 + \frac{6}{5}ad^2f^3x^5 + \frac{2}{5}bcd^2f^3x^5 + \frac{1}{5}ad^2f^3x^5 + \frac{1}{3}b^2d^2f^3x^3 + \frac{2}{3}ad^2f^3x^3 + \frac{2}{3}bcd^2f^3x^3 + ad^2f^3x^3 + \frac{1}{3}b^2d^2f^3x^3 + \frac{2}{3}ad^2f^3x^3 + ad^2x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^3,x, algorithm="giac")

[Out] $1/13*b*d^2*f^3*x^{13} + 2/11*b*c*d*f^3*x^{11} + 1/11*a*d^2*f^3*x^{11} + 3/11*b*d^2*f^2*x^{11}*e + 1/9*b*c^2*f^3*x^9 + 2/9*a*c*d*f^3*x^9 + 2/3*b*c*d*f^2*x^9*e + 1/3*a*d^2*f^2*x^9*e + 1/3*b*d^2*f*x^9*e^2 + 1/7*a*c^2*f^3*x^7 + 3/7*b*c^2*f^2*x^7*e + 6/7*a*c*d*f^2*x^7*e + 6/7*b*c*d*f*x^7*e^2 + 3/7*a*d^2*f*x^7*e^2 + 1/7*b*d^2*x^7*e^3 + 3/5*a*c^2*f^2*x^5*e + 3/5*b*c^2*f*x^5*e^2 + 6/5*a*c*d*f*x^5*e^2 + 2/5*b*c*d*x^5*e^3 + 1/5*a*d^2*x^5*e^3 + a*c^2*f*x^3*e^2 + 1/3*b*c^2*x^3*e^3 + 2/3*a*c*d*x^3*e^3 + a*c^2*x*e^3$

maple [A] time = 0.00, size = 237, normalized size = 1.05

$$\frac{bd^2f^3x^{13}}{13} + \frac{(3bd^2ef^2 + (ad^2 + 2bcd)f^2)x^{11}}{11} + \frac{(3bd^2ef^2 + 3(ad^2 + 2bcd)ef^2 + (2acd + b^2c^2)f^2)x^9}{9} + \frac{(a^2f^3 + bd^2e^3 + 3(ad^2 + 2bcd)e^2f + 3(2acd + b^2c^2)ef^2)x^7}{7} + \frac{(3a^2ef^2 + (ad^2 + 2bcd)e^3 + 3(2acd + b^2c^2)e^2f)x^5}{5} + \frac{(3a^2e^2f + (2acd + b^2c^2)e^3)x^3}{3} + \frac{(3a^2e^2f + (2acd + b^2c^2)e^3)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^3,x)

[Out] $1/13*b*d^2*f^3*x^{13} + 1/11*((a*d^2+2*b*c*d)*f^3+3*b*d^2*e*f^2)*x^{11} + 1/9*((2*a*c*d+b*c^2)*f^3+3*(a*d^2+2*b*c*d)*e*f^2+3*b*d^2*e^2*f)*x^9 + 1/7*(a*c^2*f^3+3*(2*a*c*d+b*c^2)*e*f^2+3*(a*d^2+2*b*c*d)*e^2*f+b*d^2*e^3)*x^7 + 1/5*(3*a*c^2*e*f^2+3*(2*a*c*d+b*c^2)*e^2*f+(a*d^2+2*b*c*d)*e^3)*x^5 + 1/3*(3*a*c^2*e^2*f+(2*a*c*d+b*c^2)*e^3)*x^3 + a*c^2*e^3*x$

maxima [A] time = 0.63, size = 236, normalized size = 1.04

$$\frac{1}{13}b^2f^3x^{13} + \frac{1}{11}(3bd^2ef^2 + (2bcd + ad^2)f^2)x^{11} + \frac{1}{9}(3bd^2ef^2 + 3(2bcd + ad^2)ef^2 + (b^2c^2 + 2acd)f^2)x^9 + \frac{1}{7}(bd^2e^3 + ac^2f^3 + 3(2bcd + ad^2)e^2f + 3(b^2c^2 + 2acd)ef^2)x^7 + \frac{1}{5}(3a^2ef^2 + (2bcd + ad^2)e^3 + 3(b^2c^2 + 2acd)e^2f)x^5 + \frac{1}{3}(3a^2e^2f + (b^2c^2 + 2acd)e^3)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^3,x, algorithm="maxima")

[Out] $1/13*b*d^2*f^3*x^{13} + 1/11*(3*b*d^2*e*f^2 + (2*b*c*d + a*d^2)*f^3)*x^{11} + 1/9*(3*b*d^2*e^2*f + 3*(2*b*c*d + a*d^2)*e*f^2 + (b*c^2 + 2*a*c*d)*f^3)*x^9 + 1/7*(b*d^2*e^3 + a*c^2*f^3 + 3*(2*b*c*d + a*d^2)*e^2*f + 3*(b*c^2 + 2*a*c*d)*e*f^2)*x^7 + a*c^2*e^3*x + 1/5*(3*a*c^2*e*f^2 + (2*b*c*d + a*d^2)*e^3 + 3*(b*c^2 + 2*a*c*d)*e^2*f)*x^5 + 1/3*(3*a*c^2*e^2*f + (b*c^2 + 2*a*c*d)*e^3)*x^3$

mupad [B] time = 0.86, size = 233, normalized size = 1.03

$$x^5 \left(\frac{3bc^2e^2f}{5} + \frac{3ac^2ef^2}{5} + \frac{2bcd^2e^2}{5} + \frac{6acde^2f}{5} + \frac{ad^2e^2}{5} \right) + x^9 \left(\frac{bc^2f^3}{9} + \frac{2bcde^2f^2}{9} + \frac{2acd^2f^3}{9} + \frac{bd^2e^2f}{3} + \frac{ad^2ef^2}{3} \right) + x^7 \left(\frac{3bc^2ef^2}{7} + \frac{ac^2f^3}{7} + \frac{6bcd^2ef}{7} + \frac{6acde^2f^2}{7} + \frac{bd^2e^2}{7} + \frac{3ad^2e^2f}{7} \right) + \frac{bd^2f^3x^{11}}{13} + \frac{c^2x^3(3acf+2ade+bce)}{3} + \frac{df^2x^{11}(adf+2bcf+3bde)}{11} + ad^2e^2x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^3,x)

[Out] $x^5 \left(\frac{(a*d^2*e^3)}{5} + \frac{(2*b*c*d*e^3)}{5} + \frac{(3*a*c^2*e*f^2)}{5} + \frac{(3*b*c^2*e^2*f)}{5} + \frac{(6*a*c*d*e^2*f)}{5} \right) + x^9 \left(\frac{(b*c^2*f^3)}{9} + \frac{(2*a*c*d*f^3)}{9} + \frac{(a*d^2*e*f^2)}{3} + \frac{(b*d^2*e^2*f)}{3} + \frac{(2*b*c*d*e*f^2)}{3} \right) + x^7 \left(\frac{(a*c^2*f^3)}{7} + \frac{(b*d^2*e^3)}{7} + \frac{(3*a*d^2*e^2*f)}{7} + \frac{(3*b*c^2*e*f^2)}{7} + \frac{(6*a*c*d*e*f^2)}{7} + \frac{(6*b*c*d*e^2*f)}{7} \right) + \frac{(b*d^2*f^3*x^{13})}{13} + \frac{(c*e^2*x^3*(3*a*c*f + 2*a*d*e + b*c*e))}{3} + \frac{(d*f^2*x^{11}*(a*d*f + 2*b*c*f + 3*b*d*e))}{11} + a*c^2*e^3*x$

sympy [A] time = 0.11, size = 304, normalized size = 1.35

$$ac^2e^2x + \frac{bd^2f^3x^{13}}{13} + x^{11} \left(\frac{ad^2f^3}{11} + \frac{2bcd^2f^2}{11} + \frac{3bd^2ef^2}{11} \right) + x^9 \left(\frac{2acd^2f^3}{9} + \frac{ad^2ef^2}{3} + \frac{bc^2f^3}{9} + \frac{2bcd^2ef^2}{9} + \frac{bd^2e^2f}{3} \right) + x^7 \left(\frac{ac^2f^3}{7} + \frac{6acde^2f^2}{7} + \frac{3ad^2ef}{7} + \frac{3bc^2ef^2}{7} + \frac{6bcd^2ef}{7} + \frac{bd^2e^2}{7} \right) + x^5 \left(\frac{3ac^2ef^2}{5} + \frac{6acde^2f}{5} + \frac{ad^2e^3}{5} + \frac{3bc^2ef}{5} + \frac{2bcd^2e^2}{5} \right) + x^3 \left(ac^2ef + \frac{2acd^2e^2}{3} + \frac{bc^2e^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2*(f*x**2+e)**3,x)

[Out] $a*c**2*e**3*x + b*d**2*f**3*x**13/13 + x**11*(a*d**2*f**3/11 + 2*b*c*d*f**3/11 + 3*b*d**2*e*f**2/11) + x**9*(2*a*c*d*f**3/9 + a*d**2*e*f**2/3 + b*c**2*f**3/9 + 2*b*c*d*e*f**2/3 + b*d**2*e**2*f/3) + x**7*(a*c**2*f**3/7 + 6*a*c*d*e*f**2/7 + 3*a*d**2*e**2*f/7 + 3*b*c**2*e*f**2/7 + 6*b*c*d*e**2*f/7 + b*d**2*e**3/7) + x**5*(3*a*c**2*e*f**2/5 + 6*a*c*d*e**2*f/5 + a*d**2*e**3/5 + 3*b*c**2*e**2*f/5 + 2*b*c*d*e**3/5) + x**3*(a*c**2*e**2*f + 2*a*c*d*e**3/3 + b*c**2*e**3/3)$

$$3.10 \quad \int (a + bx^2)(c + dx^2)^2 (e + fx^2)^2 dx$$

Optimal. Leaf size=158

$$\frac{1}{7}x^7(2adf(cf + de) + b(c^2f^2 + 4cdef + d^2e^2)) + \frac{1}{5}x^5(a(c^2f^2 + 4cdef + d^2e^2) + 2bce(cf + de)) + \frac{1}{9}dfx^9(adf +$$

Rubi [A] time = 0.17, antiderivative size = 158, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {521}

$$\frac{1}{7}x^7(2adf(cf + de) + b(c^2f^2 + 4cdef + d^2e^2)) + \frac{1}{5}x^5(a(c^2f^2 + 4cdef + d^2e^2) + 2bce(cf + de)) + \frac{1}{9}dfx^9(adf + 2b(cf + de)) + \frac{1}{3}cex^3(2a(cf + de) + bce) + ac^2e^2x + \frac{1}{11}bd^2f^2x^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^2,x]

[Out] a*c^2*e^2*x + (c*e*(b*c*e + 2*a*(d*e + c*f))*x^3)/3 + ((2*b*c*e*(d*e + c*f) + a*(d^2*e^2 + 4*c*d*e*f + c^2*f^2))*x^5)/5 + ((2*a*d*f*(d*e + c*f) + b*(d^2*e^2 + 4*c*d*e*f + c^2*f^2))*x^7)/7 + (d*f*(a*d*f + 2*b*(d*e + c*f))*x^9)/9 + (b*d^2*f^2*x^11)/11

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^2 (e + fx^2)^2 dx &= \int (ac^2e^2 + ce(bce + 2a(de + cf))x^2 + (2bce(de + cf) + a(d^2e^2 + 4cde) \\ &= ac^2e^2x + \frac{1}{3}ce(bce + 2a(de + cf))x^3 + \frac{1}{5}(2bce(de + cf) + a(d^2e^2 + 4cde) \end{aligned}$$

Mathematica [A] time = 0.06, size = 158, normalized size = 1.00

$$\frac{1}{7}x^7(2adf(cf + de) + b(c^2f^2 + 4cdef + d^2e^2)) + \frac{1}{5}x^5(a(c^2f^2 + 4cdef + d^2e^2) + 2bce(cf + de)) + \frac{1}{9}dfx^9(adf + 2b(cf + de)) + \frac{1}{3}cex^3(2a(cf + de) + bce) + ac^2e^2x + \frac{1}{11}bd^2f^2x^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^2,x]

[Out] a*c^2*e^2*x + (c*e*(b*c*e + 2*a*(d*e + c*f))*x^3)/3 + ((2*b*c*e*(d*e + c*f) + a*(d^2*e^2 + 4*c*d*e*f + c^2*f^2))*x^5)/5 + ((2*a*d*f*(d*e + c*f) + b*(d^2*e^2 + 4*c*d*e*f + c^2*f^2))*x^7)/7 + (d*f*(a*d*f + 2*b*(d*e + c*f))*x^9)/9 + (b*d^2*f^2*x^11)/11

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^2 (e + fx^2)^2 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^2, x]

fricas [A] time = 0.87, size = 202, normalized size = 1.28

$$\frac{1}{11}x^{11}f^2d^2b + \frac{2}{9}x^9fcd^2b + \frac{2}{9}x^9f^2dcb + \frac{1}{9}x^9f^2d^2a + \frac{1}{7}x^7f^2d^2b + \frac{4}{7}x^7fcdcb + \frac{1}{7}x^7f^2c^2b + \frac{2}{7}x^7fcd^2a + \frac{2}{7}x^7f^2dca + \frac{2}{5}x^5f^2dcb + \frac{2}{5}x^5fcd^2b + \frac{1}{5}x^5f^2d^2a + \frac{4}{5}x^5fcdca + \frac{1}{5}x^5f^2c^2a + \frac{1}{3}x^3f^2c^2b + \frac{2}{3}x^3fcdca + \frac{2}{3}x^3fcd^2a + x^2c^2a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^2,x, algorithm="fricas")

$$\begin{aligned} \text{[Out]} & 1/11*x^{11}*f^2*d^2*b + 2/9*x^9*f*e*d^2*b + 2/9*x^9*f^2*d*c*b + 1/9*x^9*f^2*d^2*a \\ & + 1/7*x^7*e^2*d^2*b + 4/7*x^7*f*e*d*c*b + 1/7*x^7*f^2*c^2*b + 2/7*x^7*f*e*d^2*a \\ & + 2/7*x^7*f^2*d*c*a + 2/5*x^5*e^2*d*c*b + 2/5*x^5*f*e*c^2*b + 1/5*x^5*e^2*d^2*a \\ & + 4/5*x^5*f*e*d*c*a + 1/5*x^5*f^2*c^2*a + 1/3*x^3*e^2*c^2*b + 2/3*x^3*e^2*d*c*a \\ & + 2/3*x^3*f*e*c^2*a + x*e^2*c^2*a \end{aligned}$$

giac [A] time = 0.36, size = 202, normalized size = 1.28

$$\frac{1}{11}bd^2f^2x^{11} + \frac{2}{9}bcd^2fx^9 + \frac{1}{9}ad^2f^2x^9 + \frac{2}{9}bd^2fx^9e + \frac{1}{7}bd^2f^2x^7 + \frac{2}{7}acd^2fx^7 + \frac{4}{7}bcd^2fx^7e + \frac{2}{7}ad^2fx^7e + \frac{1}{7}bd^2fx^7e + \frac{1}{5}ac^2f^2x^5 + \frac{2}{5}bd^2fx^5e + \frac{4}{5}acd^2fx^5e + \frac{2}{5}bcd^2fx^5e + \frac{1}{5}ad^2fx^5e + \frac{2}{3}ac^2fx^3e + \frac{1}{3}bd^2fx^3e + \frac{2}{3}acd^2fx^3e + ac^2xe^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^2,x, algorithm="giac")

$$\begin{aligned} \text{[Out]} & 1/11*b*d^2*f^2*x^{11} + 2/9*b*c*d*f^2*x^9 + 1/9*a*d^2*f^2*x^9 + 2/9*b*d^2*f*f*x^9 \\ & *e + 1/7*b*c^2*f^2*x^7 + 2/7*a*c*d*f^2*x^7 + 4/7*b*c*d*f*f*x^7*e + 2/7*a*d^2*f*f*x^7 \\ & *e + 1/7*b*d^2*x^7*e^2 + 1/5*a*c^2*f^2*x^5 + 2/5*b*c^2*f*f*x^5*e + 4/5*a*c*d*f*f*x^5 \\ & *e + 2/5*b*c*d*x^5*e^2 + 1/5*a*d^2*x^5*e^2 + 2/3*a*c^2*f*f*x^3*e + 1/3*b*c^2*x^3*e^2 \\ & + 2/3*a*c*d*x^3*e^2 + a*c^2*x*e^2 \end{aligned}$$

maple [A] time = 0.00, size = 169, normalized size = 1.07

$$\frac{bd^2f^2x^{11}}{11} + \frac{(2bd^2ef + (ad^2 + 2bcd)f^2)x^9}{9} + \frac{(bd^2e^2 + 2(ad^2 + 2bcd)ef + (2acd + b^2c^2)f^2)x^7}{7} + ac^2e^2x + \frac{(ac^2f^2 + (ad^2 + 2bcd)e^2 + 2(2acd + b^2c^2)ef)x^5}{5} + \frac{(2a^2ef + (2acd + b^2c^2)e^2)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^2,x)

$$\begin{aligned} \text{[Out]} & 1/11*b*d^2*f^2*x^{11} + 1/9*((a*d^2 + 2*b*c*d)*f^2 + 2*b*d^2*e*f)*x^9 + 1/7*((2*a*c*d \\ & + b*c^2)*f^2 + 2*(a*d^2 + 2*b*c*d)*e*f + b*d^2*e^2)*x^7 + 1/5*(a*c^2*f^2 + 2*(2*a*c*d + \\ & b*c^2)*e*f + (a*d^2 + 2*b*c*d)*e^2)*x^5 + 1/3*(2*a*c^2*e*f + (2*a*c*d + b*c^2)*e^2)*x \\ & ^3 + a*c^2*e^2*x \end{aligned}$$

maxima [A] time = 0.61, size = 168, normalized size = 1.06

$$\frac{1}{11}bd^2f^2x^{11} + \frac{1}{9}(2bd^2ef + (2bcd + ad^2)f^2)x^9 + \frac{1}{7}(bd^2e^2 + 2(2bcd + ad^2)ef + (bc^2 + 2acd)f^2)x^7 + ac^2e^2x + \frac{1}{5}(ac^2f^2 + (2bcd + ad^2)e^2 + 2(bc^2 + 2acd)ef)x^5 + \frac{1}{3}(2ac^2ef + (bc^2 + 2acd)e^2)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e)^2,x, algorithm="maxima")

$$\begin{aligned} \text{[Out]} & 1/11*b*d^2*f^2*x^{11} + 1/9*(2*b*d^2*e*f + (2*b*c*d + a*d^2)*f^2)*x^9 + 1/7*(\\ & b*d^2*e^2 + 2*(2*b*c*d + a*d^2)*e*f + (b*c^2 + 2*a*c*d)*f^2)*x^7 + a*c^2*e^2 \\ & *x + 1/5*(a*c^2*f^2 + (2*b*c*d + a*d^2)*e^2 + 2*(b*c^2 + 2*a*c*d)*e*f)*x^5 \\ & + 1/3*(2*a*c^2*e*f + (b*c^2 + 2*a*c*d)*e^2)*x^3 \end{aligned}$$

mupad [B] time = 0.07, size = 158, normalized size = 1.00

$$x^5 \left(\frac{2bc^2ef}{5} + \frac{ac^2f^2}{5} + \frac{2bcd^2e^2}{5} + \frac{4acdef}{5} + \frac{ad^2e^2}{5} \right) + x^7 \left(\frac{bc^2f^2}{7} + \frac{4bcdef}{7} + \frac{2acd^2f^2}{7} + \frac{bd^2e^2}{7} + \frac{2ad^2ef}{7} \right) + \frac{bd^2f^2x^{11}}{11} + ac^2e^2x + \frac{cx^3(2acf + 2ade + bce)}{3} + \frac{dfx^9(adf + 2bcf + 2bde)}{9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^2*(e + f*x^2)^2,x)

$$\begin{aligned} \text{[Out]} & x^5*((a*c^2*f^2)/5 + (a*d^2*e^2)/5 + (2*b*c*d*e^2)/5 + (2*b*c^2*e*f)/5 + (4 \\ & *a*c*d*e*f)/5) + x^7*((b*c^2*f^2)/7 + (b*d^2*e^2)/7 + (2*a*c*d*f^2)/7 + (2* \end{aligned}$$

$$\frac{a*d^2*e*f}{7} + \frac{(4*b*c*d*e*f)}{7} + \frac{(b*d^2*f^2*x^{11})}{11} + a*c^2*e^2*x + \frac{(c*e*x^3*(2*a*c*f + 2*a*d*e + b*c*e))}{3} + \frac{(d*f*x^9*(a*d*f + 2*b*c*f + 2*b*d*e))}{9}$$

sympy [A] time = 0.10, size = 216, normalized size = 1.37

$$ac^2e^2x + \frac{bd^2f^2x^{11}}{11} + x^9\left(\frac{ad^2f^2}{9} + \frac{2bcd^2f^2}{9} + \frac{2bd^2ef^2}{9}\right) + x^7\left(\frac{2acd^2f^2}{7} + \frac{2ad^2ef^2}{7} + \frac{bc^2f^2}{7} + \frac{4bcdef^2}{7} + \frac{bd^2e^2}{7}\right) + x^5\left(\frac{ac^2f^2}{5} + \frac{4acdef^2}{5} + \frac{ad^2e^2}{5} + \frac{2bc^2ef^2}{5} + \frac{2bcd^2e^2}{5}\right) + x^3\left(\frac{2ac^2ef^2}{3} + \frac{2acde^2}{3} + \frac{bc^2e^2}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2*(f*x**2+e)**2,x)

[Out] a*c**2*e**2*x + b*d**2*f**2*x**11/11 + x**9*(a*d**2*f**2/9 + 2*b*c*d*f**2/9 + 2*b*d**2*e*f/9) + x**7*(2*a*c*d*f**2/7 + 2*a*d**2*e*f/7 + b*c**2*f**2/7 + 4*b*c*d*e*f/7 + b*d**2*e**2/7) + x**5*(a*c**2*f**2/5 + 4*a*c*d*e*f/5 + a*d**2*e**2/5 + 2*b*c**2*e*f/5 + 2*b*c*d*e**2/5) + x**3*(2*a*c**2*e*f/3 + 2*a*c*d*e**2/3 + b*c**2*e**2/3)

$$3.11 \quad \int (a + bx^2)(c + dx^2)^2(e + fx^2) dx$$

Optimal. Leaf size=94

$$\frac{1}{7}dx^7(adf+2bcf+bde)+\frac{1}{5}x^5(ad(2cf+de)+bc(cf+2de))+\frac{1}{3}cx^3(acf+2ade+bce)+ac^2ex+\frac{1}{9}bd^2fx^9$$

Rubi [A] time = 0.08, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {521}

$$\frac{1}{7}dx^7(adf + 2bcf + bde) + \frac{1}{5}x^5(ad(2cf + de) + bc(cf + 2de)) + \frac{1}{3}cx^3(acf + 2ade + bce) + ac^2ex + \frac{1}{9}bd^2fx^9$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2),x]

[Out] a*c^2*e*x + (c*(b*c*e + 2*a*d*e + a*c*f)*x^3)/3 + ((b*c*(2*d*e + c*f) + a*d*(d*e + 2*c*f))*x^5)/5 + (d*(b*d*e + 2*b*c*f + a*d*f)*x^7)/7 + (b*d^2*f*x^9)/9

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^2(e + fx^2) dx &= \int (ac^2e + c(bce + 2ade + acf)x^2 + (bc(2de + cf) + ad(de + 2cf))x^4 + d(bcd^2e + 2acdf + 2bcde)x^6 + ac^2ex + \frac{1}{3}c(bce + 2ade + acf)x^3 + \frac{1}{5}(bc(2de + cf) + ad(de + 2cf))x^5 + \frac{1}{7}cd^2fx^7) dx \\ &= ac^2ex + \frac{1}{3}c(bce + 2ade + acf)x^3 + \frac{1}{5}(bc(2de + cf) + ad(de + 2cf))x^5 + \frac{1}{7}cd^2fx^7 \end{aligned}$$

Mathematica [A] time = 0.03, size = 96, normalized size = 1.02

$$\frac{1}{5}x^5(2acdf + ad^2e + bc^2f + 2bcde) + \frac{1}{7}dx^7(adf + 2bcf + bde) + \frac{1}{3}cx^3(acf + 2ade + bce) + ac^2ex + \frac{1}{9}bd^2fx^9$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2),x]

[Out] a*c^2*e*x + (c*(b*c*e + 2*a*d*e + a*c*f)*x^3)/3 + ((2*b*c*d*e + a*d^2*e + b*c^2*f + 2*a*c*d*f)*x^5)/5 + (d*(b*d*e + 2*b*c*f + a*d*f)*x^7)/7 + (b*d^2*f*x^9)/9

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^2(e + fx^2) dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2),x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^2*(e + f*x^2), x]

fricas [A] time = 1.05, size = 114, normalized size = 1.21

$$\frac{1}{9}x^9fd^2b + \frac{1}{7}x^7ed^2b + \frac{2}{7}x^7fdcb + \frac{1}{7}x^7fd^2a + \frac{2}{5}x^5edcb + \frac{1}{5}x^5fc^2b + \frac{1}{5}x^5ed^2a + \frac{2}{5}x^5fdca + \frac{1}{3}x^3ec^2b + \frac{2}{3}x^3edca + \frac{1}{3}x^3fc^2a + xec^2a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e),x, algorithm="fricas")

[Out] $\frac{1}{9}x^9fd^2b + \frac{1}{7}x^7ed^2b + \frac{2}{7}x^7fdcb + \frac{1}{7}x^7fd^2a + \frac{2}{5}x^5edcb + \frac{1}{5}x^5fc^2b + \frac{1}{5}x^5ed^2a + \frac{2}{5}x^5fdca + \frac{1}{3}x^3ec^2b + \frac{2}{3}x^3edca + \frac{1}{3}x^3fc^2a + xec^2a$

giac [A] time = 0.29, size = 120, normalized size = 1.28

$$\frac{1}{9}bd^2fx^9 + \frac{2}{7}bcdfx^7 + \frac{1}{7}ad^2fx^7 + \frac{1}{7}bd^2x^7e + \frac{1}{5}bc^2fx^5 + \frac{2}{5}acdfx^5 + \frac{2}{5}bcdx^5e + \frac{1}{5}ad^2x^5e + \frac{1}{3}ac^2fx^3 + \frac{1}{3}bc^2x^3e + \frac{2}{3}acdx^3e + ac^2xe$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e),x, algorithm="giac")

[Out] $\frac{1}{9}b*d^2*f*x^9 + \frac{2}{7}b*c*d*f*x^7 + \frac{1}{7}a*d^2*f*x^7 + \frac{1}{7}b*d^2*x^7*e + \frac{1}{5}b*c^2*f*x^5 + \frac{2}{5}a*c*d*f*x^5 + \frac{2}{5}b*c*d*x^5*e + \frac{1}{5}a*d^2*x^5*e + \frac{1}{3}a*c^2*f*x^3 + \frac{1}{3}b*c^2*x^3*e + \frac{2}{3}a*c*d*x^3*e + a*c^2*x*e$

maple [A] time = 0.00, size = 101, normalized size = 1.07

$$\frac{bd^2fx^9}{9} + \frac{(bd^2e + (ad^2 + 2bcd)f)x^7}{7} + ac^2ex + \frac{((ad^2 + 2bcd)e + (2acd + bc^2)f)x^5}{5} + \frac{(ac^2f + (2acd + bc^2)e)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e),x)

[Out] $\frac{1}{9}b*d^2*f*x^9 + \frac{1}{7}*((a*d^2 + 2*b*c*d)*f + b*d^2*e)*x^7 + \frac{1}{5}*((2*a*c*d + b*c^2)*f + (a*d^2 + 2*b*c*d)*e)*x^5 + \frac{1}{3}*(a*c^2*f + (2*a*c*d + b*c^2)*e)*x^3 + a*c^2*e*x$

maxima [A] time = 0.76, size = 100, normalized size = 1.06

$$\frac{1}{9}bd^2fx^9 + \frac{1}{7}(bd^2e + (2bcd + ad^2)f)x^7 + \frac{1}{5}((2bcd + ad^2)e + (bc^2 + 2acd)f)x^5 + ac^2ex + \frac{1}{3}(ac^2f + (bc^2 + 2acd)e)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2*(f*x^2+e),x, algorithm="maxima")

[Out] $\frac{1}{9}b*d^2*f*x^9 + \frac{1}{7}*(b*d^2*e + (2*b*c*d + a*d^2)*f)*x^7 + \frac{1}{5}*((2*b*c*d + a*d^2)*e + (b*c^2 + 2*a*c*d)*f)*x^5 + a*c^2*e*x + \frac{1}{3}*(a*c^2*f + (b*c^2 + 2*a*c*d)*e)*x^3$

mupad [B] time = 0.05, size = 99, normalized size = 1.05

$$x^5\left(\frac{ad^2e}{5} + \frac{bc^2f}{5} + \frac{2acdf}{5} + \frac{2bcde}{5}\right) + x^3\left(\frac{ac^2f}{3} + \frac{bc^2e}{3} + \frac{2acde}{3}\right) + x^7\left(\frac{ad^2f}{7} + \frac{bd^2e}{7} + \frac{2bcd f}{7}\right) + ac^2ex + \frac{bd^2fx^9}{9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^2*(e + f*x^2),x)

[Out] $x^5*((a*d^2*e)/5 + (b*c^2*f)/5 + (2*a*c*d*f)/5 + (2*b*c*d*e)/5) + x^3*((a*c^2*f)/3 + (b*c^2*e)/3 + (2*a*c*d*e)/3) + x^7*((a*d^2*f)/7 + (b*d^2*e)/7 + (2*b*c*d*f)/7) + a*c^2*e*x + (b*d^2*f*x^9)/9$

sympy [A] time = 0.08, size = 121, normalized size = 1.29

$$ac^2ex + \frac{bd^2fx^9}{9} + x^7\left(\frac{ad^2f}{7} + \frac{2bcd f}{7} + \frac{bd^2e}{7}\right) + x^5\left(\frac{2acdf}{5} + \frac{ad^2e}{5} + \frac{bc^2f}{5} + \frac{2bcde}{5}\right) + x^3\left(\frac{ac^2f}{3} + \frac{2acde}{3} + \frac{bc^2e}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)**2*(f*x**2+e),x)
```

```
[Out] a*c**2*e*x + b*d**2*f*x**9/9 + x**7*(a*d**2*f/7 + 2*b*c*d*f/7 + b*d**2*e/7)
+ x**5*(2*a*c*d*f/5 + a*d**2*e/5 + b*c**2*f/5 + 2*b*c*d*e/5) + x**3*(a*c**
2*f/3 + 2*a*c*d*e/3 + b*c**2*e/3)
```


$$3.12 \quad \int \frac{(a+bx^2)(c+dx^2)^2}{e+fx^2} dx$$

Optimal. Leaf size=142

$$\frac{x(5adf(3de-5cf)-b(8c^2f^2-25cdef+15d^2e^2))}{15f^3} - \frac{(be-af)(de-cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e}f^{7/2}} - \frac{x(c+dx^2)(-5adf-}{15f^2}$$

Rubi [A] time = 0.21, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {528, 388, 205}

$$\frac{x(5adf(3de-5cf)-b(8c^2f^2-25cdef+15d^2e^2))}{15f^3} - \frac{x(c+dx^2)(-5adf-4bcf+5bde)}{15f^2} - \frac{(be-af)(de-cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e}f^{7/2}} + \frac{bx(c+dx^2)^2}{5f}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2), x]

[Out] -((5*a*d*f*(3*d*e - 5*c*f) - b*(15*d^2*e^2 - 25*c*d*e*f + 8*c^2*f^2))*x)/(15*f^3) - ((5*b*d*e - 4*b*c*f - 5*a*d*f)*x*(c + d*x^2))/(15*f^2) + (b*x*(c + d*x^2)^2)/(5*f) - ((b*e - a*f)*(d*e - c*f)^2*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(Sqrt[e]*f^(7/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(b*(n*(p+q+1)+1)), x] + Dist[1/(b*(n*(p+q+1)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-1)*Simp[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p+q+1)+1, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^2}{e + fx^2} dx &= \frac{bx(c + dx^2)^2}{5f} + \frac{\int \frac{(c+dx^2)(-c(be-5af)+(-5bde+4bcf+5adf)x^2)}{e+fx^2} dx}{5f} \\
&= -\frac{(5bde - 4bcf - 5adf)x(c + dx^2)}{15f^2} + \frac{bx(c + dx^2)^2}{5f} + \frac{\int \frac{c(be(5de-7cf)-5af(de-3cf))-(5adf)}{e}}{5f} \\
&= -\frac{(5adf(3de - 5cf) - b(15d^2e^2 - 25cdef + 8c^2f^2))x}{15f^3} - \frac{(5bde - 4bcf - 5adf)x(c + dx^2)}{15f^2} \\
&= -\frac{(5adf(3de - 5cf) - b(15d^2e^2 - 25cdef + 8c^2f^2))x}{15f^3} - \frac{(5bde - 4bcf - 5adf)x(c + dx^2)}{15f^2}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 115, normalized size = 0.81

$$-\frac{(be - af)(de - cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e} f^{7/2}} + \frac{x(adf(2cf - de) + b(de - cf)^2)}{f^3} + \frac{dx^3(adf + 2bcf - bde)}{3f^2} + \frac{bd^2x^5}{5f}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2), x]

[Out] ((b*(d*e - c*f)^2 + a*d*f*(-(d*e) + 2*c*f))*x)/f^3 + (d*(-(b*d*e) + 2*b*c*f + a*d*f)*x^3)/(3*f^2) + (b*d^2*x^5)/(5*f) - ((b*e - a*f)*(d*e - c*f)^2*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(Sqrt[e]*f^(7/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^2}{e + fx^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2), x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2), x]

fricas [A] time = 1.19, size = 366, normalized size = 2.58

$$\frac{[6bd^2f^2x^5 - 10(bd^2f^2 - 2bcd + ad^2f^2)x^4 + 15(bd^2f^2 - ac^2f^2 - 2bcd + ad^2f^2)x^3 + (bc^2 + 2acd)f^2\sqrt{e}\log\left(\frac{c^2 + \sqrt{e}cx}{-2cd + ad^2f^2}\right) + 30(bd^2f^2 - 2bcd + ad^2f^2)(bc^2 + 2acd)f^2]x^5 - 15(bd^2f^2 - ac^2f^2 - 2bcd + ad^2f^2)(bc^2 + 2acd)f^2\sqrt{e}\arctan\left(\frac{\sqrt{e}x}{\sqrt{e}}\right) + 15(bd^2f^2 - 2bcd + ad^2f^2)(bc^2 + 2acd)f^2]x^4}{15f^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e), x, algorithm="fricas")

[Out] [1/30*(6*b*d^2*e*f^3*x^5 - 10*(b*d^2*e^2*f^2 - (2*b*c*d + a*d^2)*e*f^3)*x^3 + 15*(b*d^2*e^3 - a*c^2*f^3 - (2*b*c*d + a*d^2)*e^2*f + (b*c^2 + 2*a*c*d)*e*f^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) + 30*(b*d^2*e^3*f - (2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x)/(e*f^4), 1/15*(3*b*d^2*e*f^3*x^5 - 5*(b*d^2*e^2*f^2 - (2*b*c*d + a*d^2)*e*f^3)*x^3 - 15*(b*d^2*e^3 - a*c^2*f^3 - (2*b*c*d + a*d^2)*e^2*f + (b*c^2 + 2*a*c*d)*e*f^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) + 15*(b*d^2*e^3*f - (2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x)/(e*f^4)]

giac [A] time = 0.29, size = 178, normalized size = 1.25

$$\frac{(ac^2f^3 - bc^2f^2e - 2acdf^2e + 2bcdfe^2 + ad^2fe^2 - bd^2e^3) \arctan\left(\sqrt{f}xe^{\left(\frac{1}{2}\right)}\right)e^{\left(-\frac{1}{2}\right)}}{f^{\frac{7}{2}}} + \frac{3bd^2f^4x^5 + 10bcdfe^4x^3 + 5ad^2f^4x^3 - 5bd^2f^3x^3e + 15bc^2f^4x + 30acdf^4x - 30bcdfe^3xe - 15ad^2f^3xe + 15bd^2f^2xe^2}{15f^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e),x, algorithm="giac")

[Out] (a*c^2*f^3 - b*c^2*f^2*e - 2*a*c*d*f^2*e + 2*b*c*d*f*e^2 + a*d^2*f*e^2 - b*d^2*e^3)*arctan(sqrt(f)*x*e^(-1/2))*e^(-1/2)/f^(7/2) + 1/15*(3*b*d^2*f^4*x^5 + 10*b*c*d*f^4*x^3 + 5*a*d^2*f^4*x^3 - 5*b*d^2*f^3*x^3*e + 15*b*c^2*f^4*x + 30*a*c*d*f^4*x - 30*b*c*d*f^3*x*e - 15*a*d^2*f^3*x*e + 15*b*d^2*f^2*x*e^2)/f^5

maple [A] time = 0.01, size = 243, normalized size = 1.71

$$\frac{bd^2x^5}{5f} + \frac{ad^2x^3}{3f} + \frac{2bcdx^3}{3f} - \frac{bd^2ex^3}{3f^2} + \frac{a^2e \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}} - \frac{2acde \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}f} + \frac{ad^2e^2 \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}f^2} - \frac{b^2e \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}f} + \frac{2bcd e^2 \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}f^2} - \frac{bd^2e^3 \arctan\left(\frac{fx}{\sqrt{ef}}\right)}{\sqrt{ef}f^3} + \frac{2acdx}{f} - \frac{ad^2ex}{f^2} + \frac{bc^2x}{f} - \frac{2bcdex}{f^2} + \frac{bd^2e^2x}{f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e),x)

[Out] 1/5/f*b*d^2*x^5+1/3/f*x^3*a*d^2+2/3/f*x^3*b*c*d-1/3/f^2*x^3*b*d^2*e+2/f*a*c*d*x-1/f^2*a*d^2*e*x+1/f*b*c^2*x-2/f^2*b*c*d*e*x+1/f^3*b*d^2*e^2*x+1/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2-2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d*e+1/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^2*e^2-1/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2*e+2/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d*e^2-1/f^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^2*e^3

maxima [A] time = 1.35, size = 160, normalized size = 1.13

$$\frac{(bd^2e^3 - ac^2f^3 - (2bcd + ad^2)e^2f + (bc^2 + 2acd)ef^2) \arctan\left(\frac{fx}{\sqrt{ef}}\right) + 3bd^2f^2x^5 - 5(bd^2ef - (2bcd + ad^2)f^2)x^3 + 15(bd^2e^2 - (2bcd + ad^2)ef + (bc^2 + 2acd)f^2)x}{\sqrt{ef}f^3} + \frac{3bd^2f^2x^5 - 5(bd^2ef - (2bcd + ad^2)f^2)x^3 + 15(bd^2e^2 - (2bcd + ad^2)ef + (bc^2 + 2acd)f^2)x}{15f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e),x, algorithm="maxima")

[Out] -(b*d^2*e^3 - a*c^2*f^3 - (2*b*c*d + a*d^2)*e^2*f + (b*c^2 + 2*a*c*d)*e*f^2)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*f^3) + 1/15*(3*b*d^2*f^2*x^5 - 5*(b*d^2*e*f - (2*b*c*d + a*d^2)*f^2)*x^3 + 15*(b*d^2*e^2 - (2*b*c*d + a*d^2)*e*f + (b*c^2 + 2*a*c*d)*f^2)*x)/f^3

mupad [B] time = 0.86, size = 203, normalized size = 1.43

$$x^3 \left(\frac{ad^2 + 2bcd}{3f} - \frac{bd^2e}{3f^2} \right) + x \left(\frac{bc^2 + 2adc}{f} - \frac{e \left(\frac{ad^2 + 2bcd}{f} - \frac{bd^2e}{f^2} \right)}{f} \right) + \frac{bd^2x^5}{5f} + \frac{\operatorname{atan}\left(\frac{\sqrt{f}x(af-be)(cf-de)^2}{\sqrt{e}(-b^2ef^2+ac^2f^3+2bcd e^2f-2acde f^2-bd^2e^3+ad^2e^2f)} \right) (af-be)(cf-de)^2}{\sqrt{e}f^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2),x)

[Out] x^3*((a*d^2 + 2*b*c*d)/(3*f) - (b*d^2*e)/(3*f^2)) + x*((b*c^2 + 2*a*c*d)/f - (e*((a*d^2 + 2*b*c*d)/f - (b*d^2*e)/f^2))/f + (b*d^2*x^5)/(5*f) + (atan((f^(1/2)*x*(a*f - b*e)*(c*f - d*e)^2)/(e^(1/2)*(a*c^2*f^3 - b*d^2*e^3 + a*d^2*e^2*f - b*c^2*e*f^2 - 2*a*c*d*e*f^2 + 2*b*c*d*e^2*f)))*(a*f - b*e)*(c*f - d*e)^2)/(e^(1/2)*f^(7/2))

sympy [B] time = 1.04, size = 347, normalized size = 2.44

$$\frac{bd^2x^5}{5f} + x^3 \left(\frac{ad^2}{3f} + \frac{2bcd}{3f} - \frac{bd^2e}{3f^2} \right) + x \left(\frac{2acd}{f} - \frac{ad^2e}{f^2} + \frac{bc^2}{f} - \frac{2bcde}{f^2} + \frac{bd^2e^2}{f^3} \right) - \frac{\sqrt{\frac{1}{ef}}(af-be)(cf-de)^2 \log\left(\frac{ef^3 \sqrt{\frac{1}{ef}}(af-be)(cf-de)^2}{a^2f^3 - 2acde f^2 + ad^2e^2f - bc^2ef^2 + 2bcd e^2f - bd^2e^3 + x} \right)}{2} + \frac{\sqrt{\frac{1}{ef}}(af-be)(cf-de)^2 \log\left(\frac{ef^3 \sqrt{\frac{1}{ef}}(af-be)(cf-de)^2}{a^2f^3 - 2acde f^2 + ad^2e^2f - bc^2ef^2 + 2bcd e^2f - bd^2e^3 + x} \right)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2/(f*x**2+e),x)

```
[Out] b*d**2*x**5/(5*f) + x**3*(a*d**2/(3*f) + 2*b*c*d/(3*f) - b*d**2*e/(3*f**2))
+ x*(2*a*c*d/f - a*d**2*e/f**2 + b*c**2/f - 2*b*c*d*e/f**2 + b*d**2*e**2/f
**3) - sqrt(-1/(e*f**7))*(a*f - b*e)*(c*f - d*e)**2*log(-e*f**3*sqrt(-1/(e*
f**7)))*(a*f - b*e)*(c*f - d*e)**2/(a*c**2*f**3 - 2*a*c*d*e*f**2 + a*d**2*e*
*2*f - b*c**2*e*f**2 + 2*b*c*d*e**2*f - b*d**2*e**3) + x)/2 + sqrt(-1/(e*f*
*7))*(a*f - b*e)*(c*f - d*e)**2*log(e*f**3*sqrt(-1/(e*f**7)))*(a*f - b*e)*(c
*f - d*e)**2/(a*c**2*f**3 - 2*a*c*d*e*f**2 + a*d**2*e**2*f - b*c**2*e*f**2
+ 2*b*c*d*e**2*f - b*d**2*e**3) + x)/2
```

$$3.13 \quad \int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^2} dx$$

Optimal. Leaf size=164

$$\frac{(de - cf) \tan^{-1} \left(\frac{\sqrt{f}x}{\sqrt{e}} \right) (be(5de - cf) - af(cf + 3de))}{2e^{3/2}f^{7/2}} - \frac{dx(be(15de - 13cf) - 3af(3de - cf))}{6ef^3} + \frac{dx(c + dx^2)(5be - 3af)}{6ef^2}$$

Rubi [A] time = 0.23, antiderivative size = 164, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {526, 528, 388, 205}

$$\frac{(de - cf) \tan^{-1} \left(\frac{\sqrt{f}x}{\sqrt{e}} \right) (be(5de - cf) - af(cf + 3de))}{2e^{3/2}f^{7/2}} + \frac{dx(c + dx^2)(5be - 3af)}{6ef^2} - \frac{dx(be(15de - 13cf) - 3af(3de - cf))}{6ef^3} - \frac{x(c + dx^2)^2 (be - af)}{2ef(e + fx^2)}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^2,x]

[Out] -(d*(b*e*(15*d*e - 13*c*f) - 3*a*f*(3*d*e - c*f))*x)/(6*e*f^3) + (d*(5*b*e - 3*a*f)*x*(c + d*x^2))/(6*e*f^2) - ((b*e - a*f)*x*(c + d*x^2)^2)/(2*e*f*(e + f*x^2)) + ((d*e - c*f)*(b*e*(5*d*e - c*f) - a*f*(3*d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(2*e^(3/2)*f^(7/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^2} dx &= -\frac{(be - af)x(c + dx^2)^2}{2ef(e + fx^2)} - \frac{\int \frac{(c+dx^2)(-c(be+af)-d(5be-3af)x^2)}{e+fx^2} dx}{2ef} \\
&= \frac{d(5be - 3af)x(c + dx^2)}{6ef^2} - \frac{(be - af)x(c + dx^2)^2}{2ef(e + fx^2)} - \frac{\int \frac{c(be(5de-3cf)-3af(de+cf))+d(be(15de-13cf)-3af(3de-cf))x}{e+fx^2} dx}{6ef^2} \\
&= -\frac{d(be(15de - 13cf) - 3af(3de - cf))x}{6ef^3} + \frac{d(5be - 3af)x(c + dx^2)}{6ef^2} - \frac{(be - af)x(c + dx^2)^2}{2ef(e + fx^2)} \\
&= -\frac{d(be(15de - 13cf) - 3af(3de - cf))x}{6ef^3} + \frac{d(5be - 3af)x(c + dx^2)}{6ef^2} - \frac{(be - af)x(c + dx^2)^2}{2ef(e + fx^2)}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 134, normalized size = 0.82

$$\frac{(de - cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(5de - cf) - af(cf + 3de))}{2e^{3/2}f^{7/2}} - \frac{x(be - af)(de - cf)^2}{2ef^3(e + fx^2)} + \frac{dx(adf + 2bcf - 2bde)}{f^3} + \frac{bd^2x^3}{3f^2}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^2,x]

[Out] (d*(-2*b*d*e + 2*b*c*f + a*d*f)*x)/f^3 + (b*d^2*x^3)/(3*f^2) - ((b*e - a*f)*(d*e - c*f)^2*x)/(2*e*f^3*(e + f*x^2)) + ((d*e - c*f)*(b*e*(5*d*e - c*f) - a*f*(3*d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(2*e^(3/2)*f^(7/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^2, x]

fricas [A] time = 1.38, size = 552, normalized size = 3.37

1/12*(4*b*d^2*e^2*f^3*x^5 - 4*(5*b*d^2*e^3*f^2 - 3*(2*b*c*d + a*d^2)*e^2*f^3)*x^3 - 3*(5*b*d^2*e^4 + a*c^2*e*f^3 - 3*(2*b*c*d + a*d^2)*e^3*f + (b*c^2 + 2*a*c*d)*e^2*f^2 + (5*b*d^2*e^3*f + a*c^2*f^4 - 3*(2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 6*(5*b*d^2*e^4*f - a*c^2*e*f^4 - 3*(2*b*c*d + a*d^2)*e^3*f^2 + (b*c^2 + 2*a*c*d)*e^2*f^3)*x)/(e^2*f^5*x^2 + e^3*f^4), 1/6*(2*b*d^2*e^2*f^3*x^5 - 2*(5*b*d^2*e^3*f^2 - 3*(2*b*c*d + a*d^2)*e^2*f^3)*x^3 + 3*(5*b*d^2*e^4 + a*c^2*e*f^3 - 3*(2*b*c*d + a*d^2)*e^3*f + (b*c^2 + 2*a*c*d)*e^2*f^2 + (5*b*d^2*e^3*f + a*c^2*f^4 - 3*(2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) - 3*(5*b*d^2*e^4*f -

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^2,x, algorithm="fricas")

[Out] [1/12*(4*b*d^2*e^2*f^3*x^5 - 4*(5*b*d^2*e^3*f^2 - 3*(2*b*c*d + a*d^2)*e^2*f^3)*x^3 - 3*(5*b*d^2*e^4 + a*c^2*e*f^3 - 3*(2*b*c*d + a*d^2)*e^3*f + (b*c^2 + 2*a*c*d)*e^2*f^2 + (5*b*d^2*e^3*f + a*c^2*f^4 - 3*(2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 6*(5*b*d^2*e^4*f - a*c^2*e*f^4 - 3*(2*b*c*d + a*d^2)*e^3*f^2 + (b*c^2 + 2*a*c*d)*e^2*f^3)*x)/(e^2*f^5*x^2 + e^3*f^4), 1/6*(2*b*d^2*e^2*f^3*x^5 - 2*(5*b*d^2*e^3*f^2 - 3*(2*b*c*d + a*d^2)*e^2*f^3)*x^3 + 3*(5*b*d^2*e^4 + a*c^2*e*f^3 - 3*(2*b*c*d + a*d^2)*e^3*f + (b*c^2 + 2*a*c*d)*e^2*f^2 + (5*b*d^2*e^3*f + a*c^2*f^4 - 3*(2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) - 3*(5*b*d^2*e^4*f -

$$a*c^2*e*f^4 - 3*(2*b*c*d + a*d^2)*e^3*f^2 + (b*c^2 + 2*a*c*d)*e^2*f^3)*x)/(e^2*f^5*x^2 + e^3*f^4)]$$

giac [A] time = 0.33, size = 195, normalized size = 1.19

$$\frac{(a^2 f^3 + b c^2 f^2 e + 2 a c d f^2 e - 6 b c d f e^2 - 3 a d^2 f e^2 + 5 b d^2 e^3) \arctan\left(\sqrt{f} x e^{\left(\frac{-1}{2}\right)}\right) e^{\left(\frac{-3}{2}\right)} + (a^2 f^3 x - b c^2 f^2 x e - 2 a c d f^2 x e + 2 b c d f x e^2 + a d^2 f x e^2 - b d^2 x e^3) e^{(-1)} + \frac{b d^2 f^4 x^3 + 6 b c d f^4 x + 3 a d^2 f^4 x - 6 b d^2 f^3 x e}{3 f^6}}{2 f^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^2,x, algorithm="giac")

[Out] 1/2*(a*c^2*f^3 + b*c^2*f^2*e + 2*a*c*d*f^2*e - 6*b*c*d*f*e^2 - 3*a*d^2*f*e^2 + 5*b*d^2*e^3)*arctan(sqrt(f)*x*e^(-1/2))*e^(-3/2)/f^(7/2) + 1/2*(a*c^2*f^3*x - b*c^2*f^2*x*e - 2*a*c*d*f^2*x*e + 2*b*c*d*f*x*e^2 + a*d^2*f*x*e^2 - b*d^2*x*e^3)*e^(-1)/((f*x^2 + e)*f^3) + 1/3*(b*d^2*f^4*x^3 + 6*b*c*d*f^4*x + 3*a*d^2*f^4*x - 6*b*d^2*f^3*x*e)/f^6

maple [B] time = 0.01, size = 299, normalized size = 1.82

$$\frac{b d^2 x^3}{3 f^2} + \frac{a c^2 x}{2(f x^2 + e) e} + \frac{a^2 \arctan\left(\frac{f x}{\sqrt{e}}\right)}{2 \sqrt{e} e} - \frac{a c d x}{(f x^2 + e) f} + \frac{a c d \arctan\left(\frac{f x}{\sqrt{e}}\right)}{\sqrt{e} f} + \frac{a d^2 e x}{2(f x^2 + e) f^2} - \frac{3 a d^2 e \arctan\left(\frac{f x}{\sqrt{e}}\right)}{2 \sqrt{e} f^2} - \frac{b c^2 x}{2(f x^2 + e) f} + \frac{b c^2 \arctan\left(\frac{f x}{\sqrt{e}}\right)}{2 \sqrt{e} f} + \frac{b c d e x}{(f x^2 + e) f^2} - \frac{3 b c d e \arctan\left(\frac{f x}{\sqrt{e}}\right)}{\sqrt{e} f^2} - \frac{b d^2 e^2 x}{2(f x^2 + e) f^3} + \frac{5 b d^2 e^2 \arctan\left(\frac{f x}{\sqrt{e}}\right)}{2 \sqrt{e} f^3} + \frac{a d^2 x}{f^2} + \frac{2 b c d x}{f^2} - \frac{2 b d^2 e x}{f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^2,x)

[Out] 1/3*d^2/f^2*x^3*b+d^2/f^2*a*x+2*d/f^2*b*c*x-2*d^2/f^3*b*e*x+1/2/e*x/(f*x^2+e)*a*c^2-1/f*x/(f*x^2+e)*a*c*d+1/2/f^2*e*x/(f*x^2+e)*a*d^2-1/2/f*x/(f*x^2+e)*b*c^2+1/f^2*e*x/(f*x^2+e)*b*c*d-1/2/f^3*e^2*x/(f*x^2+e)*b*d^2+1/2/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2+1/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d-3/2/f^2*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^2+1/2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2-3/f^2*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d+5/2/f^3*e^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^2

maxima [A] time = 1.94, size = 186, normalized size = 1.13

$$\frac{(b d^2 e^3 - a c^2 f^3 - (2 b c d + a d^2) e^2 f + (b c^2 + 2 a c d) e f^2) x}{2(e f^4 x^2 + e^2 f^3)} + \frac{b d^2 f x^3 - 3(2 b d^2 e - (2 b c d + a d^2) f) x}{3 f^3} + \frac{(5 b d^2 e^3 + a c^2 f^3 - 3(2 b c d + a d^2) e^2 f + (b c^2 + 2 a c d) e f^2) \arctan\left(\frac{f x}{\sqrt{e}}\right)}{2 \sqrt{e} e f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^2,x, algorithm="maxima")

[Out] -1/2*(b*d^2*e^3 - a*c^2*f^3 - (2*b*c*d + a*d^2)*e^2*f + (b*c^2 + 2*a*c*d)*e*f^2)*x/(e*f^4*x^2 + e^2*f^3) + 1/3*(b*d^2*f*x^3 - 3*(2*b*d^2*e - (2*b*c*d + a*d^2)*f)*x)/f^3 + 1/2*(5*b*d^2*e^3 + a*c^2*f^3 - 3*(2*b*c*d + a*d^2)*e^2*f + (b*c^2 + 2*a*c*d)*e*f^2)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*e*f^3)

mupad [B] time = 0.17, size = 257, normalized size = 1.57

$$x \left(\frac{a d^2 + 2 b c d}{f^2} - \frac{2 b d^2 e}{f^3} \right) + \frac{b d^2 x^3}{3 f^2} + \frac{x(-b c^2 e f^2 + a c^2 f^3 + 2 b c d e^2 f - 2 a c d e f^2 - b d^2 e^3 + a d^2 e^2 f)}{2 e(f^4 x^2 + e f^3)} + \frac{\operatorname{atan}\left(\frac{\sqrt{f} x (c f - d e) (a c f^2 - 5 b d^2 e^2 + 3 a d e f + b c e f)}{\sqrt{e} (b c^2 e f^2 + a c^2 f^3 - 6 b c d e^2 f + 2 a c d e f^2 + 5 b d^2 e^3 - 3 a d^2 e^2 f)}\right) (c f - d e) (a c f^2 - 5 b d^2 e^2 + 3 a d e f + b c e f)}{2 e^{3/2} f^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^2,x)

[Out] x*((a*d^2 + 2*b*c*d)/f^2 - (2*b*d^2*e)/f^3) + (b*d^2*x^3)/(3*f^2) + (x*(a*c^2*f^3 - b*d^2*e^3 + a*d^2*e^2*f - b*c^2*e*f^2 - 2*a*c*d*e*f^2 + 2*b*c*d*e^2*f))/(2*e*(e*f^3 + f^4*x^2)) + (atan((f^(1/2)*x*(c*f - d*e)*(a*c*f^2 - 5*b*d*e^2 + 3*a*d*e*f + b*c*e*f)))/(e^(1/2)*(a*c^2*f^3 + 5*b*d^2*e^3 - 3*a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 - 6*b*c*d*e^2*f)))*(c*f - d*e)*(a*c*f^2 - 5*b*d*e^2 + 3*a*d*e*f + b*c*e*f))/(2*e^(3/2)*f^(7/2))

sympy [B] time = 2.83, size = 483, normalized size = 2.95

$$\frac{bx^3}{3f^2} + x \left(\frac{af^2}{f^2} + \frac{2bcd}{f^2} - \frac{2bd^2e}{f^3} \right) + \frac{x(ac^2f^3 - 2acdef^2 + ad^2e^2f - bc^2ef^2 + 2bcd^2f - bd^2e^2)}{2f^3 + 2ef^2} \sqrt{-\frac{4}{27}} (cf - de) (acf^2 + 3adef + bcf - 5bd^2) \log \left(\frac{cf^2 \sqrt{-\frac{4}{27}} (cf - de) (acf^2 + 3adef + bcf - 5bd^2)}{27f^3 + 27cd^2f^2 - 3d^2ef^2 + 27e^2f^2 + 27d^2e^2} + x \right) + \frac{x(ac^2f^3 - 2acdef^2 + ad^2e^2f - bc^2ef^2 + 2bcd^2f - bd^2e^2)}{2f^3 + 2ef^2} \sqrt{-\frac{4}{27}} (cf - de) (acf^2 + 3adef + bcf - 5bd^2) \log \left(\frac{cf^2 \sqrt{-\frac{4}{27}} (cf - de) (acf^2 + 3adef + bcf - 5bd^2)}{27f^3 + 27cd^2f^2 - 3d^2ef^2 + 27e^2f^2 + 27d^2e^2} + x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2/(f*x**2+e)**2,x)

[Out] b*d**2*x**3/(3*f**2) + x*(a*d**2/f**2 + 2*b*c*d/f**2 - 2*b*d**2*e/f**3) + x*(a*c**2*f**3 - 2*a*c*d*e*f**2 + a*d**2*e**2*f - b*c**2*e*f**2 + 2*b*c*d*e**2*f - b*d**2*e**3)/(2*e**2*f**3 + 2*e*f**4*x**2) - sqrt(-1/(e**3*f**7))*(c*f - d*e)*(a*c*f**2 + 3*a*d*e*f + b*c*e*f - 5*b*d*e**2)*log(-e**2*f**3*sqrt(-1/(e**3*f**7))*(c*f - d*e)*(a*c*f**2 + 3*a*d*e*f + b*c*e*f - 5*b*d*e**2)/(a*c**2*f**3 + 2*a*c*d*e*f**2 - 3*a*d**2*e**2*f + b*c**2*e*f**2 - 6*b*c*d*e**2*f + 5*b*d**2*e**3) + x)/4 + sqrt(-1/(e**3*f**7))*(c*f - d*e)*(a*c*f**2 + 3*a*d*e*f + b*c*e*f - 5*b*d*e**2)*log(e**2*f**3*sqrt(-1/(e**3*f**7))*(c*f - d*e)*(a*c*f**2 + 3*a*d*e*f + b*c*e*f - 5*b*d*e**2)/(a*c**2*f**3 + 2*a*c*d*e*f**2 - 3*a*d**2*e**2*f + b*c**2*e*f**2 - 6*b*c*d*e**2*f + 5*b*d**2*e**3) + x)/4

$$3.14 \quad \int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^3} dx$$

Optimal. Leaf size=207

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(-c^2f^2 - 6cdef + 15d^2e^2) - af(3c^2f^2 + 2cdef + 3d^2e^2))}{8e^{5/2}f^{7/2}} + \frac{dx(be(15de - cf) - 3af(cf + de))}{8e^2f^3}$$

Rubi [A] time = 0.24, antiderivative size = 207, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {526, 388, 205}

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(-c^2f^2 - 6cdef + 15d^2e^2) - af(3c^2f^2 + 2cdef + 3d^2e^2))}{8e^{5/2}f^{7/2}} - \frac{x(c+dx^2)(be(5de - cf) - af(3cf + de))}{8e^2f^2(e+fx^2)} + \frac{dx(be(15de - cf) - 3af(cf + de))}{8e^2f^3} - \frac{x(c+dx^2)^2(be - af)}{4ef(e+fx^2)^2}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^3, x]

[Out] (d*(b*e*(15*d*e - c*f) - 3*a*f*(d*e + c*f))*x)/(8*e^2*f^3) - ((b*e - a*f)*x*(c + d*x^2)^2)/(4*e*f*(e + f*x^2)^2) - ((b*e*(5*d*e - c*f) - a*f*(d*e + 3*c*f))*x*(c + d*x^2))/(8*e^2*f^2*(e + f*x^2)) - ((b*e*(15*d^2*e^2 - 6*c*d*e*f - c^2*f^2) - a*f*(3*d^2*e^2 + 2*c*d*e*f + 3*c^2*f^2))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(8*e^(5/2)*f^(7/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*b*n*(p+1)), x] + Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)*Simp[c*(b*e*n*(p+1) + b*e - a*f) + d*(b*e*n*(p+1) + (b*e - a*f)*(n*q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^3} dx &= -\frac{(be - af)x(c + dx^2)^2}{4ef(e + fx^2)^2} - \frac{\int \frac{(c+dx^2)(-c(be+3af)-d(5be-af)x^2)}{(e+fx^2)^2} dx}{4ef} \\
&= -\frac{(be - af)x(c + dx^2)^2}{4ef(e + fx^2)^2} - \frac{(be(5de - cf) - af(de + 3cf))x(c + dx^2)}{8e^2f^2(e + fx^2)} + \frac{\int \frac{-c(af(de-3cf)-d(5be-af)x^2)}{(e+fx^2)^2} dx}{8e^2f^2} \\
&= \frac{d(be(15de - cf) - 3af(de + cf))x}{8e^2f^3} - \frac{(be - af)x(c + dx^2)^2}{4ef(e + fx^2)^2} - \frac{(be(5de - cf) - af(de - cf))x}{8e^2f^2(e + fx^2)} \\
&= \frac{d(be(15de - cf) - 3af(de + cf))x}{8e^2f^3} - \frac{(be - af)x(c + dx^2)^2}{4ef(e + fx^2)^2} - \frac{(be(5de - cf) - af(de - cf))x}{8e^2f^2(e + fx^2)}
\end{aligned}$$

Mathematica [A] time = 0.13, size = 183, normalized size = 0.88

$$-\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(-c^2f^2 - 6cdef + 15d^2e^2) - af(3c^2f^2 + 2cdef + 3d^2e^2))}{8e^{5/2}f^{7/2}} + \frac{x(de - cf)(be(9de - cf) - af(3cf + 5de))}{8e^2f^3(e + fx^2)} - \frac{x(be - af)(de - cf)^2}{4ef^3(e + fx^2)^2} + \frac{bd^2x}{f^3}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^3,x]

[Out] (b*d^2*x)/f^3 - ((b*e - a*f)*(d*e - c*f)^2*x)/(4*e*f^3*(e + f*x^2)^2) + ((d*e - c*f)*(b*e*(9*d*e - c*f) - a*f*(5*d*e + 3*c*f))*x)/(8*e^2*f^3*(e + f*x^2)^2) - ((b*e*(15*d^2*e^2 - 6*c*d*e*f - c^2*f^2) - a*f*(3*d^2*e^2 + 2*c*d*e*f + 3*c^2*f^2))*ArcTan[Sqrt[f]*x/Sqrt[e]]/(8*e^(5/2)*f^(7/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^3} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^3, x]

fricas [A] time = 1.43, size = 777, normalized size = 3.75

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^3,x, algorithm="fricas")

[Out] [1/16*(16*b*d^2*e^3*f^3*x^5 + 2*(25*b*d^2*e^4*f^2 + 3*a*c^2*e*f^5 - 5*(2*b*c*d + a*d^2))*e^3*f^3 + (b*c^2 + 2*a*c*d)*e^2*f^4)*x^3 + (15*b*d^2*e^5 - 3*a*c^2*e^2*f^3 - 3*(2*b*c*d + a*d^2))*e^4*f - (b*c^2 + 2*a*c*d)*e^3*f^2 + (15*b*d^2*e^3*f^2 - 3*a*c^2*f^5 - 3*(2*b*c*d + a*d^2))*e^2*f^3 - (b*c^2 + 2*a*c*d)*e*f^4)*x^4 + 2*(15*b*d^2*e^4*f - 3*a*c^2*e*f^4 - 3*(2*b*c*d + a*d^2))*e^3*f^2 - (b*c^2 + 2*a*c*d)*e^2*f^3)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f))*x - e)/(f*x^2 + e) + 2*(15*b*d^2*e^5*f + 5*a*c^2*e^2*f^4 - 3*(2*b*c*d + a*d^2))*e^4*f^2 - (b*c^2 + 2*a*c*d)*e^3*f^3)*x)/(e^3*f^6*x^4 + 2*e^4*f^5*x^2 + e^5*f^4), 1/8*(8*b*d^2*e^3*f^3*x^5 + (25*b*d^2*e^4*f^2 + 3*a*c^2*e*f^5 -

$$5*(2*b*c*d + a*d^2)*e^3*f^3 + (b*c^2 + 2*a*c*d)*e^2*f^4)*x^3 - (15*b*d^2*e^5 - 3*a*c^2*e^2*f^3 - 3*(2*b*c*d + a*d^2)*e^4*f - (b*c^2 + 2*a*c*d)*e^3*f^2 + (15*b*d^2*e^3*f^2 - 3*a*c^2*f^5 - 3*(2*b*c*d + a*d^2)*e^2*f^3 - (b*c^2 + 2*a*c*d)*e*f^4)*x^4 + 2*(15*b*d^2*e^4*f - 3*a*c^2*e*f^4 - 3*(2*b*c*d + a*d^2)*e^3*f^2 - (b*c^2 + 2*a*c*d)*e^2*f^3)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) + (15*b*d^2*e^5*f + 5*a*c^2*e^2*f^4 - 3*(2*b*c*d + a*d^2)*e^4*f^2 - (b*c^2 + 2*a*c*d)*e^3*f^3)*x)/(e^3*f^6*x^4 + 2*e^4*f^5*x^2 + e^5*f^4)]$$

giac [A] time = 0.45, size = 238, normalized size = 1.15

$$\frac{bd^2x}{f^3} + \frac{(3ac^2f^3 + bc^2f^2e + 2acd^2f^2 + 6bcd^2f^2 + 3ad^2f^2 - 15bd^2e^2) \arctan\left(\sqrt{fx} \left(\frac{x}{e}\right)^{\frac{1}{2}}\right)}{8f^2} + \frac{(3ac^2f^4x^3 + bc^2f^3x^2e + 2acd^2f^3x^2e - 10bcd^2f^3x^2e - 5ad^2f^2x^2e^2 + 9bd^2f^2x^2e + 5ac^2f^2xe - bc^2f^2xe^2 - 2acd^2f^2xe^2 - 6bcd^2f^2xe^2 - 3ad^2f^2xe^2 + 7bd^2xe^4)e^{-2}}{8(fx^2 + e)^2f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^3,x, algorithm="giac")

[Out] $b*d^2*x/f^3 + 1/8*(3*a*c^2*f^3 + b*c^2*f^2*e + 2*a*c*d*f^2*e + 6*b*c*d*f*e^2 + 3*a*d^2*f*e^2 - 15*b*d^2*e^3)*arctan(sqrt(f)*x*e^{(-1/2)})*e^{(-5/2)}/f^{(7/2)} + 1/8*(3*a*c^2*f^4*x^3 + b*c^2*f^3*x^3*e + 2*a*c*d*f^3*x^3*e - 10*b*c*d*f^2*x^3*e^2 - 5*a*d^2*f^2*x^3*e^2 + 9*b*d^2*f*x^3*e^3 + 5*a*c^2*f^3*x*e - b*c^2*f^2*x*e^2 - 2*a*c*d*f^2*x*e^2 - 6*b*c*d*f*x*e^3 - 3*a*d^2*f*x*e^3 + 7*b*d^2*x*e^4)*e^{(-2)}/((f*x^2 + e)^2*f^3)$

maple [B] time = 0.01, size = 397, normalized size = 1.92

$$\frac{3ac^2f^3}{8(fx^2+e)^2} + \frac{acd^2}{4(fx^2+e)^2} + \frac{5ad^2e^2}{8(fx^2+e)^2} + \frac{b^2x^2}{8(fx^2+e)^2} + \frac{5bcd^2}{4(fx^2+e)^2} + \frac{9bd^2e^2}{8(fx^2+e)^2} + \frac{5a^2x}{8(fx^2+e)^2} + \frac{ad^2x}{4(fx^2+e)^2} + \frac{3ad^2x}{8(fx^2+e)^2} + \frac{b^2x}{8(fx^2+e)^2} + \frac{3bd^2x}{4(fx^2+e)^2} + \frac{7bd^2x}{8(fx^2+e)^2} + \frac{3ac^2 \arctan\left(\frac{fx}{e}\right)}{8\sqrt{ef}e^2} + \frac{acd \arctan\left(\frac{fx}{e}\right)}{4\sqrt{ef}e} + \frac{3ad^2 \arctan\left(\frac{fx}{e}\right)}{8\sqrt{ef}e^2} + \frac{b^2 \arctan\left(\frac{fx}{e}\right)}{8\sqrt{ef}e} + \frac{3bcd \arctan\left(\frac{fx}{e}\right)}{4\sqrt{ef}e^2} + \frac{3bd^2 \arctan\left(\frac{fx}{e}\right)}{8\sqrt{ef}e^2} + \frac{15bd^2 \arctan\left(\frac{fx}{e}\right)}{8\sqrt{ef}e^2} + \frac{bd^2x}{f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^3,x)

[Out] $b*d^2/f^3*x + 3/8*f/(f*x^2+e)^2/e^2*x^3*a*c^2 + 1/4/(f*x^2+e)^2/e*x^3*a*c*d - 5/8/f/(f*x^2+e)^2*x^3*a*d^2 + 1/8/(f*x^2+e)^2/e*x^3*b*c^2 - 5/4/f/(f*x^2+e)^2*x^3*b*c*d + 9/8/f^2/(f*x^2+e)^2*x^3*b*d^2*e + 5/8/(f*x^2+e)^2/e*x*a*c^2 - 1/4/f/(f*x^2+e)^2*a*c*d*x - 3/8/f^2/(f*x^2+e)^2*a*d^2*e*x - 1/8/f/(f*x^2+e)^2*b*c^2*x - 3/4/f^2/(f*x^2+e)^2*b*c*d*e*x + 7/8/f^3/(f*x^2+e)^2*b*d^2*e^2*x + 3/8/e^2/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*a*c^2 + 1/4/f/e/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*a*c*d + 3/8/f^2/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*a*d^2 + 1/8/f/e/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*b*c^2 + 3/4/f^2/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*b*c*d - 15/8/f^3*e/(e*f)^{(1/2)}*arctan(1/(e*f)^{(1/2)}*f*x)*b*d^2$

maxima [A] time = 2.01, size = 236, normalized size = 1.14

$$\frac{bd^2x}{f^3} + \frac{(9bd^2e^3f + 3ac^2f^4 - 5(2bcd + ad^2)e^2f^2 + (bc^2 + 2acd)ef^3)x^3 + (7bd^2e^4 + 5ac^2ef^3 - 3(2bcd + ad^2)e^2f - (bc^2 + 2acd)e^2f^2)x}{8(e^2f^3x^4 + 2e^2f^4x^2 + e^4f^3)} - \frac{(15bd^2e^3 - 3ac^2f^3 - 3(2bcd + ad^2)e^2f - (bc^2 + 2acd)ef^2) \arctan\left(\frac{fx}{\sqrt{e}}\right)}{8\sqrt{ef}e^2f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^3,x, algorithm="maxima")

[Out] $b*d^2*x/f^3 + 1/8*((9*b*d^2*e^3*f + 3*a*c^2*f^4 - 5*(2*b*c*d + a*d^2)*e^2*f^2 + (b*c^2 + 2*a*c*d)*e*f^3)*x^3 + (7*b*d^2*e^4 + 5*a*c^2*e*f^3 - 3*(2*b*c*d + a*d^2)*e^3*f - (b*c^2 + 2*a*c*d)*e^2*f^2)*x)/(e^2*f^5*x^4 + 2*e^3*f^4*x^2 + e^4*f^3) - 1/8*(15*b*d^2*e^3 - 3*a*c^2*f^3 - 3*(2*b*c*d + a*d^2)*e^2*f - (b*c^2 + 2*a*c*d)*e*f^2)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*e^2*f^3)$

mupad [B] time = 0.98, size = 243, normalized size = 1.17

$$\frac{\operatorname{atan}\left(\frac{\sqrt{fx}}{\sqrt{e}}\right) (bc^2ef^2 + 3ac^2f^3 + 6bcd^2ef + 2acde^2f - 15bd^2e^3 + 3ad^2e^2f)}{8e^{5/2}f^{7/2}} - \frac{x(b^2e^2f^2 - 5a^2f^3 + 6bcd^2f + 2acde^2f - 7bd^2e^2 + 3ad^2e^2f) - x^3(bc^2ef^3 + 3ac^2f^4 - 10bcd^2f^2 + 2acde^2f + 9bd^2e^3f - 5ad^2e^2f^2)}{8e^{5/2}f^{7/2}} + \frac{bd^2x}{f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^3,x)

```
[Out] (atan((f^(1/2)*x)/e^(1/2))*(3*a*c^2*f^3 - 15*b*d^2*e^3 + 3*a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 + 6*b*c*d*e^2*f))/(8*e^(5/2)*f^(7/2)) - ((x*(3*a*d^2*e^2*f - 7*b*d^2*e^3 - 5*a*c^2*f^3 + b*c^2*e*f^2 + 2*a*c*d*e*f^2 + 6*b*c*d*e^2*f))/(8*e) - (x^3*(3*a*c^2*f^4 - 5*a*d^2*e^2*f^2 + b*c^2*e*f^3 + 9*b*d^2*e^3*f - 10*b*c*d*e^2*f^2 + 2*a*c*d*e*f^3))/(8*e^2))/(e^2*f^3 + f^5*x^4 + 2*e*f^4*x^2) + (b*d^2*x)/f^3
```

sympy [B] time = 11.98, size = 400, normalized size = 1.93

$$\frac{b d^2 x}{f^3} - \frac{\sqrt{-\frac{1}{e f^5}} (3 a c^2 f^3 + 2 a c d e f^2 + 3 a d^2 e^2 f + b c^2 e f^2 + 6 b c d e^2 f - 15 b d^2 e^3) \log\left(-\frac{e^2 f^3}{\sqrt{-\frac{1}{e f^5}}} + x\right)}{16} + \frac{\sqrt{-\frac{1}{e f^5}} (3 a c^2 f^3 + 2 a c d e f^2 + 3 a d^2 e^2 f + b c^2 e f^2 + 6 b c d e^2 f - 15 b d^2 e^3) \log\left(\frac{e^2 f^3}{\sqrt{-\frac{1}{e f^5}}} + x\right)}{16} + \frac{x^3 (3 a c^2 f^4 - 5 a d^2 e^2 f^2 + b c^2 e f^3 + 9 b d^2 e^3 f - 10 b c d e^2 f^2 + 2 a c d e f^3)}{8 e^2} + \frac{b d^2 x}{f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)**2/(f*x**2+e)**3,x)
```

```
[Out] b*d**2*x/f**3 - sqrt(-1/(e**5*f**7))*(3*a*c**2*f**3 + 2*a*c*d*e*f**2 + 3*a*d**2*e**2*f + b*c**2*e*f**2 + 6*b*c*d*e**2*f - 15*b*d**2*e**3)*log(-e**3*f**3*sqrt(-1/(e**5*f**7)) + x)/16 + sqrt(-1/(e**5*f**7))*(3*a*c**2*f**3 + 2*a*c*d*e*f**2 + 3*a*d**2*e**2*f + b*c**2*e*f**2 + 6*b*c*d*e**2*f - 15*b*d**2*e**3)*log(e**3*f**3*sqrt(-1/(e**5*f**7)) + x)/16 + (x**3*(3*a*c**2*f**4 + 2*a*c*d*e*f**3 - 5*a*d**2*e**2*f**2 + b*c**2*e*f**3 - 10*b*c*d*e**2*f**2 + 9*b*d**2*e**3*f) + x*(5*a*c**2*e*f**3 - 2*a*c*d*e**2*f**2 - 3*a*d**2*e**3*f - b*c**2*e**2*f**2 - 6*b*c*d*e**3*f + 7*b*d**2*e**4))/(8*e**4*f**3 + 16*e**3*f**4*x**2 + 8*e**2*f**5*x**4)
```

$$3.15 \quad \int \frac{(a+bx^2)(c+dx^2)^2}{(e+fx^2)^4} dx$$

Optimal. Leaf size=240

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)\left(af(5c^2f^2 + 2cdef + d^2e^2) + be(c^2f^2 + 2cdef + 5d^2e^2)\right)}{16e^{7/2}f^{7/2}} - \frac{x\left(af(-15c^2f^2 + 4cdef + 3d^2e^2) + be\right)}{48e^3f^3(e+fx^2)}$$

Rubi [A] time = 0.28, antiderivative size = 240, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {526, 385, 205}

$$\frac{x(af(-15c^2f^2 + 4cdef + 3d^2e^2) + be(-3c^2f^2 - 4cdef + 15d^2e^2))}{48e^3f^3(e+fx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(5c^2f^2 + 2cdef + d^2e^2) + be(c^2f^2 + 2cdef + 5d^2e^2))}{16e^{7/2}f^{7/2}} - \frac{x(c+dx^2)(de(af+5be) - cf(5af+be))}{24e^2f^2(e+fx^2)^2} - \frac{x(c+dx^2)^2(be-af)}{6ef(e+fx^2)^3}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^4,x]

[Out] -((b*e - a*f)*x*(c + d*x^2)^2)/(6*e*f*(e + f*x^2)^3) - ((d*e*(5*b*e + a*f) - c*f*(b*e + 5*a*f))*x*(c + d*x^2))/(24*e^2*f^2*(e + f*x^2)^2) - ((a*f*(3*d^2*e^2 + 4*c*d*e*f - 15*c^2*f^2) + b*e*(15*d^2*e^2 - 4*c*d*e*f - 3*c^2*f^2))*x)/(48*e^3*f^3*(e + f*x^2)) + ((b*e*(5*d^2*e^2 + 2*c*d*e*f + c^2*f^2) + a*f*(d^2*e^2 + 2*c*d*e*f + 5*c^2*f^2))*ArcTan[$\frac{\sqrt{f}x}{\sqrt{e}}$]/(16*e^(7/2)*f^(7/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*c - a*d)*x*(a + b*x^n)^(p+1))/(a*b*n*(p+1)), x] - Dist[(a*d - b*c*(n*(p+1) + 1))/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && (LtQ[p, -1] || ILtQ[1/n + p, 0])

Rule 526

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(a*b*n*(p+1)), x] + Dist[1/(a*b*n*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1)*Simp[c*(b*e*n*(p+1) + b*e - a*f) + d*(b*e*n*(p+1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^4} dx &= -\frac{(be - af)x(c + dx^2)^2}{6ef(e + fx^2)^3} - \frac{\int \frac{(c+dx^2)(-c(be+5af)-d(5be+af)x^2)}{(e+fx^2)^3} dx}{6ef} \\
&= -\frac{(be - af)x(c + dx^2)^2}{6ef(e + fx^2)^3} - \frac{(de(5be + af) - cf(be + 5af))x(c + dx^2)}{24e^2 f^2 (e + fx^2)^2} + \frac{\int \frac{c(de(5be+af)+}{24e^2 f^2 (e + fx^2)^2} \\
&= -\frac{(be - af)x(c + dx^2)^2}{6ef(e + fx^2)^3} - \frac{(de(5be + af) - cf(be + 5af))x(c + dx^2)}{24e^2 f^2 (e + fx^2)^2} - \frac{(af(3d^2 e^2 + 4cde - cf^2))x}{24e^2 f^2 (e + fx^2)^2} \\
&= -\frac{(be - af)x(c + dx^2)^2}{6ef(e + fx^2)^3} - \frac{(de(5be + af) - cf(be + 5af))x(c + dx^2)}{24e^2 f^2 (e + fx^2)^2} - \frac{(af(3d^2 e^2 + 4cde - cf^2))x}{24e^2 f^2 (e + fx^2)^2}
\end{aligned}$$

Mathematica [A] time = 0.18, size = 242, normalized size = 1.01

$$\frac{\tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(af(5c^2f^2 + 2cdef + d^2e^2) + be(c^2f^2 + 2cdef + 5d^2e^2))}{16e^{7/2}f^{7/2}} + \frac{x(af(5c^2f^2 + 2cdef + d^2e^2) + be(c^2f^2 + 2cdef - 11d^2e^2))}{16e^3f^3(e + fx^2)} + \frac{x(de - cf)(be(13de - cf) - af(5cf + 7de))}{24e^2f^3(e + fx^2)^2} - \frac{x(be - af)(de - cf)^2}{6ef^3(e + fx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^4, x]

[Out] $-\frac{1}{6} * ((b * e - a * f) * (d * e - c * f) ^ 2 * x) / (e * f ^ 3 * (e + f * x ^ 2) ^ 3) + ((d * e - c * f) * (b * e * (13 * d * e - c * f) - a * f * (7 * d * e + 5 * c * f)) * x) / (24 * e ^ 2 * f ^ 3 * (e + f * x ^ 2) ^ 2) + ((b * e * (-11 * d ^ 2 * e ^ 2 + 2 * c * d * e * f + c ^ 2 * f ^ 2) + a * f * (d ^ 2 * e ^ 2 + 2 * c * d * e * f + 5 * c ^ 2 * f ^ 2)) * x) / (16 * e ^ 3 * f ^ 3 * (e + f * x ^ 2)) + ((b * e * (5 * d ^ 2 * e ^ 2 + 2 * c * d * e * f + c ^ 2 * f ^ 2) + a * f * (d ^ 2 * e ^ 2 + 2 * c * d * e * f + 5 * c ^ 2 * f ^ 2)) * \text{ArcTan}[\text{Sqrt}[f] * x / \text{Sqrt}[e]]) / (16 * e ^ (7 / 2) * f ^ (7 / 2))$

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^2}{(e + fx^2)^4} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^4, x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^4, x]

fricas [B] time = 1.12, size = 1024, normalized size = 4.27

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^4, x, algorithm="fricas")

[Out] $[-\frac{1}{96} * (6 * (11 * b * d ^ 2 * e ^ 4 * f ^ 3 - 5 * a * c ^ 2 * e * f ^ 6 - (2 * b * c * d + a * d ^ 2) * e ^ 3 * f ^ 4 - (b * c ^ 2 + 2 * a * c * d) * e ^ 2 * f ^ 5) * x ^ 5 + 16 * (5 * b * d ^ 2 * e ^ 5 * f ^ 2 - 5 * a * c ^ 2 * e ^ 2 * f ^ 5 + (2 * b * c * d + a * d ^ 2) * e ^ 4 * f ^ 3 - (b * c ^ 2 + 2 * a * c * d) * e ^ 3 * f ^ 4) * x ^ 3 + 3 * (5 * b * d ^ 2 * e ^ 6 + 5 * a * c ^ 2 * e ^ 3 * f ^ 3 + (2 * b * c * d + a * d ^ 2) * e ^ 5 * f + (b * c ^ 2 + 2 * a * c * d) * e ^ 4 * f ^ 2 + (5 * b * d ^ 2 * e ^ 3 * f ^ 3 + 5 * a * c ^ 2 * f ^ 6 + (2 * b * c * d + a * d ^ 2) * e ^ 2 * f ^ 4 + (b * c ^ 2 + 2 * a * c * d) * e * f ^ 5) * x ^ 6 + 3 * (5 * b * d ^ 2 * e ^ 4 * f ^ 2 + 5 * a * c ^ 2 * e * f ^ 5 + (2 * b * c * d + a * d ^ 2) * e ^ 3 * f ^ 3 + (b * c ^ 2 + 2 * a * c * d) * e ^ 2 * f ^ 4) * x ^ 4 + 3 * (5 * b * d ^ 2 * e ^ 5 * f + 5 * a * c ^ 2 * e ^ 2 * f ^ 4 + ($

$$2*b*c*d + a*d^2)*e^{4*f^2} + (b*c^2 + 2*a*c*d)*e^{3*f^3}*x^2)*\sqrt{-e*f})*\log((f*x^2 - 2*\sqrt{-e*f}*x - e)/(f*x^2 + e)) + 6*(5*b*d^2*e^{6*f} - 11*a*c^2*e^{3*f^4} + (2*b*c*d + a*d^2)*e^{5*f^2} + (b*c^2 + 2*a*c*d)*e^{4*f^3})*x)/(e^{4*f^7}*x^6 + 3*e^{5*f^6}*x^4 + 3*e^{6*f^5}*x^2 + e^{7*f^4}), -1/48*(3*(11*b*d^2*e^{4*f^3} - 5*a*c^2*e*f^6 - (2*b*c*d + a*d^2)*e^{3*f^4} - (b*c^2 + 2*a*c*d)*e^{2*f^5})*x^5 + 8*(5*b*d^2*e^{5*f^2} - 5*a*c^2*e^{2*f^5} + (2*b*c*d + a*d^2)*e^{4*f^3} - (b*c^2 + 2*a*c*d)*e^{3*f^4})*x^3 - 3*(5*b*d^2*e^6 + 5*a*c^2*e^{3*f^3} + (2*b*c*d + a*d^2)*e^{5*f} + (b*c^2 + 2*a*c*d)*e^{4*f^2} + (5*b*d^2*e^{3*f^3} + 5*a*c^2*f^6 + (2*b*c*d + a*d^2)*e^{2*f^4} + (b*c^2 + 2*a*c*d)*e*f^5)*x^6 + 3*(5*b*d^2*e^{4*f^2} + 5*a*c^2*e*f^5 + (2*b*c*d + a*d^2)*e^{3*f^3} + (b*c^2 + 2*a*c*d)*e^{2*f^4})*x^4 + 3*(5*b*d^2*e^{5*f} + 5*a*c^2*e^{2*f^4} + (2*b*c*d + a*d^2)*e^{4*f^2} + (b*c^2 + 2*a*c*d)*e^{3*f^3})*x^2)*\sqrt{e*f})*\arctan(\sqrt{e*f}*x/e) + 3*(5*b*d^2*e^{6*f} - 11*a*c^2*e^{3*f^4} + (2*b*c*d + a*d^2)*e^{5*f^2} + (b*c^2 + 2*a*c*d)*e^{4*f^3})*x)/(e^{4*f^7}*x^6 + 3*e^{5*f^6}*x^4 + 3*e^{6*f^5}*x^2 + e^{7*f^4})]$$

giac [A] time = 0.30, size = 311, normalized size = 1.30

$$\frac{(5a^2f^3 + b^2f^2 + 2acdfe + 2bcdf^2 + ad^2f^2 + 5bd^2e) \arctan\left(\frac{f}{\sqrt{e}}\right) + \frac{(15a^2f^2 + 3b^2f^2 + 6acdfe + 6bcdf^2 + 3ad^2f^2 - 33bd^2e - 40ac^2f^2 + 8bc^2f^2 + 16acdfe - 16bcdf^2 - 8ad^2f^2 - 40bd^2e + 33ac^2f^2 - 3b^2f^2 - 6acdfe - 6bcdf^2 - 3ad^2f^2 - 15bd^2e)f^3}{48(f^2 + e)^3}}{16f^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^4,x, algorithm="giac")

[Out] 1/16*(5*a*c^2*f^3 + b*c^2*f^2*e + 2*a*c*d*f^2*e + 2*b*c*d*f*e^2 + a*d^2*f*e^2 + 5*b*d^2*e^3)*arctan(sqrt(f)*x*e^(-1/2))*e^(-7/2)/f^(7/2) + 1/48*(15*a*c^2*f^5*x^5 + 3*b*c^2*f^4*x^5*e + 6*a*c*d*f^4*x^5*e + 6*b*c*d*f^3*x^5*e^2 + 3*a*d^2*f^3*x^5*e^2 - 33*b*d^2*f^2*x^5*e^3 + 40*a*c^2*f^4*x^3*e + 8*b*c^2*f^3*x^3*e^2 + 16*a*c*d*f^3*x^3*e^2 - 16*b*c*d*f^2*x^3*e^3 - 8*a*d^2*f^2*x^3*e^3 - 40*b*d^2*f*x^3*e^4 + 33*a*c^2*f^3*x*e^2 - 3*b*c^2*f^2*x*e^3 - 6*a*c*d*f^2*x*e^3 - 6*b*c*d*f*x*e^4 - 3*a*d^2*f*x*e^4 - 15*b*d^2*x*e^5)*e^(-3)/((f*x^2 + e)^3*f^3)

maple [A] time = 0.01, size = 360, normalized size = 1.50

$$\frac{5a^2 \arctan\left(\frac{f}{\sqrt{e}}\right) + acd \arctan\left(\frac{f}{\sqrt{e}}\right) + ad^2 \arctan\left(\frac{f}{\sqrt{e}}\right) + bc^2 \arctan\left(\frac{f}{\sqrt{e}}\right) + bcd \arctan\left(\frac{f}{\sqrt{e}}\right) + 5bd^2 \arctan\left(\frac{f}{\sqrt{e}}\right) + \frac{(5a^2f^3 + 2acdf^2 + b^2f^2 + 2bcdfe + 2bd^2e - 11bd^2e)f^3}{16\sqrt{ef}e^3} + \frac{(5a^2f^2 + 2acdf^2 - 4bd^2e + b^2f^2 - 2bcdfe - 5bd^2e)f^3}{16\sqrt{ef}e^3} + \frac{(11a^2f^2 - 2acdf^2 - 4bd^2e - 2bcdfe - 5bd^2e)f^3}{16e^3}}{16\sqrt{ef}e^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^4,x)

[Out] (1/16*(5*a*c^2*f^3+2*a*c*d*e*f^2+a*d^2*e^2*f+b*c^2*e*f^2+2*b*c*d*e^2*f-11*b*d^2*e^3)/e^3/f*x^5+1/6*(5*a*c^2*f^3+2*a*c*d*e*f^2-a*d^2*e^2*f+b*c^2*e*f^2-2*b*c*d*e^2*f-5*b*d^2*e^3)/e^2/f^2*x^3+1/16*(11*a*c^2*f^3-2*a*c*d*e*f^2-a*d^2*e^2*f-b*c^2*e*f^2-2*b*c*d*e^2*f-5*b*d^2*e^3)/f^3/e*x)/(f*x^2+e)^3+5/16/e^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2+1/8/e^2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d+1/16/e/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^2+1/16/e^2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2+1/8/e/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d+5/16/f^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^2

maxima [A] time = 2.01, size = 304, normalized size = 1.27

$$\frac{3(11bd^2e^2f^2 - 5a^2f^3 - (2bcd + ad^2)e^2f^2 - (bc^2 + 2acd)ef^2)x^5 + 8(5bd^2ef^2 - 5ac^2ef^2 + (2bcd + ad^2)e^2f^2 - (bc^2 + 2acd)e^2f^2)x^3 + 3(5bd^2e^3 - 11a^2e^2f^3 + (2bcd + ad^2)ef^2 + (bc^2 + 2acd)e^2f^2)x + \frac{(5bd^2e^3 + 5a^2f^3 + (2bcd + ad^2)e^2f^2 + (bc^2 + 2acd)ef^2) \arctan\left(\frac{f}{\sqrt{e}}\right)}{16\sqrt{ef}e^3}}{48(e^2f^2x^6 + 3e^2f^2x^4 + 3e^2f^2x^2 + e^2f^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^2/(f*x^2+e)^4,x, algorithm="maxima")

[Out] -1/48*(3*(11*b*d^2*e^{3*f^2} - 5*a*c^2*f^5 - (2*b*c*d + a*d^2)*e^{2*f^3} - (b*c^2 + 2*a*c*d)*e*f^4)*x^5 + 8*(5*b*d^2*e^{4*f} - 5*a*c^2*e*f^4 + (2*b*c*d + a*d^2)*e^{3*f^2} - (b*c^2 + 2*a*c*d)*e^{2*f^3})*x^3 + 3*(5*b*d^2*e^5 - 11*a*c^2*e^{2*f^3} + (2*b*c*d + a*d^2)*e^{4*f} + (b*c^2 + 2*a*c*d)*e^{3*f^2})*x)/(e^{3*f^6}*x

$$e^6 + 3e^4 f^5 x^4 + 3e^5 f^4 x^2 + e^6 f^3) + 1/16(5bd^2 e^3 + 5ac^2 f^3 + (2b^2 c^2 d + a^2 d^2) e^2 f + (b^2 c^2 + 2a^2 c^2 d) e f^2) \arctan(fx/\sqrt{ef}) / (\sqrt{ef} e^3 f^3)$$

mupad [B] time = 0.98, size = 303, normalized size = 1.26

$$\frac{\frac{b^3(b^2e^2f^2+5a^2e^2f^2-2bcde^2f+2acde^2f-5b^2e^2-a^2e^2f)}{6e^2f^2} - \frac{x(b^2e^2f^2-11a^2e^2f^2+2bcde^2f+2acde^2f+5b^2e^2+a^2e^2f)}{16ef^2} + \frac{b^3(b^2e^2f^2+5a^2e^2f^2+2bcde^2f+2acde^2f-11b^2e^2+a^2e^2f)}{16ef^2}}{e^3+3e^2fx^2+3ef^2x^4+f^3x^6} + \frac{\operatorname{atan}\left(\frac{\sqrt{f}}{\sqrt{e}}\right)(bc^2ef^2+5a^2ef^3+2bcd^2e^2f+2acde^2f+5bd^2e^3+a^2d^2e^2f)}{16e^{7/2}f^{7/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^2)/(e + f*x^2)^4,x)

[Out] ((x^3*(5*a*c^2*f^3 - 5*b*d^2*e^3 - a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 - 2*b*c*d*e^2*f))/(6*e^2*f^2) - (x*(5*b*d^2*e^3 - 11*a*c^2*f^3 + a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 + 2*b*c*d*e^2*f))/(16*e*f^3) + (x^5*(5*a*c^2*f^3 - 11*b*d^2*e^3 + a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 + 2*b*c*d*e^2*f))/(16*e^3*f))/(e^3 + f^3*x^6 + 3*e^2*f*x^2 + 3*e*f^2*x^4) + (atan((f^(1/2)*x)/e^(1/2))*(5*a*c^2*f^3 + 5*b*d^2*e^3 + a*d^2*e^2*f + b*c^2*e*f^2 + 2*a*c*d*e*f^2 + 2*b*c*d*e^2*f))/(16*e^(7/2)*f^(7/2))

sympy [B] time = 42.07, size = 486, normalized size = 2.02

$$\frac{\sqrt{\frac{1}{77}(5a^2f^2+2abd^2+af^2f^2+b^2f^2+2abd^2f+2bd^2f^2+2bd^2f^2)\log\left(\frac{f}{\sqrt{77}}+1\right)} + \sqrt{\frac{1}{77}(5a^2f^2+2abd^2+af^2f^2+b^2f^2+2abd^2f+2bd^2f^2+2bd^2f^2)\log\left(\frac{f}{\sqrt{77}}+1\right)}}{\frac{e^3+3e^2fx^2+3ef^2x^4+f^3x^6}{32}} + \frac{e^3(5a^2f^3+5abd^2f^3+3a^2d^2f^3+3bd^2f^3-33bd^2f^3)+e^3(43a^2f^3+16abd^2f^3-30bd^2f^3+30bd^2f^3-13abd^2f^3-40bd^2f^3)+e^3(33a^2d^2f^3-6abd^2f^3-30bd^2f^3-13abd^2f^3)}{48e^{7/2}f^{7/2}+144e^{5/2}f^{5/2}x^2+144e^{3/2}f^{3/2}x^4+48e^{1/2}f^{1/2}x^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**2/(f*x**2+e)**4,x)

[Out] -sqrt(-1/(e**7*f**7))*(5*a*c**2*f**3 + 2*a*c*d*e*f**2 + a*d**2*e**2*f + b*c**2*e*f**2 + 2*b*c*d*e**2*f + 5*b*d**2*e**3)*log(-e**4*f**3*sqrt(-1/(e**7*f**7)) + x)/32 + sqrt(-1/(e**7*f**7))*(5*a*c**2*f**3 + 2*a*c*d*e*f**2 + a*d**2*e**2*f + b*c**2*e*f**2 + 2*b*c*d*e**2*f + 5*b*d**2*e**3)*log(e**4*f**3*sqrt(-1/(e**7*f**7)) + x)/32 + (x**5*(15*a*c**2*f**5 + 6*a*c*d*e*f**4 + 3*a*d**2*e**2*f**3 + 3*b*c**2*e*f**4 + 6*b*c*d*e**2*f**3 - 33*b*d**2*e**3*f**2) + x**3*(40*a*c**2*e*f**4 + 16*a*c*d*e**2*f**3 - 8*a*d**2*e**3*f**2 + 8*b*c**2*e**2*f**3 - 16*b*c*d*e**3*f**2 - 40*b*d**2*e**4*f) + x*(33*a*c**2*e**2*f**3 - 6*a*c*d*e**3*f**2 - 3*a*d**2*e**4*f - 3*b*c**2*e**3*f**2 - 6*b*c*d*e**4*f - 15*b*d**2*e**5))/(48*e**6*f**3 + 144*e**5*f**4*x**2 + 144*e**4*f**5*x**4 + 48*e**3*f**6*x**6)

$$3.16 \quad \int (a + bx^2)(c + dx^2)^3 (e + fx^2)^3 dx$$

Optimal. Leaf size=310

$$\frac{3}{11}dfx^{11} (adf(cf + de) + b(c^2f^2 + 3cdef + d^2e^2)) + \frac{3}{5}cex^5 (a(c^2f^2 + 3cdef + d^2e^2) + bce(cf + de)) + \frac{1}{3}c^2e^2x^3$$

Rubi [A] time = 0.39, antiderivative size = 310, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {521}

$$\frac{3}{11}dfx^{11} (adf(cf + de) + b(c^2f^2 + 3cdef + d^2e^2)) + \frac{3}{5}cex^5 (a(c^2f^2 + 3cdef + d^2e^2) + bce(cf + de)) + \frac{1}{3}c^2e^2x^3$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^3,x]

[Out] a*c^3*e^3*x + (c^2*e^2*(b*c*e + 3*a*(d*e + c*f))*x^3)/3 + (3*c*e*(b*c*e*(d*e + c*f) + a*(d^2*e^2 + 3*c*d*e*f + c^2*f^2))*x^5)/5 + ((3*b*c*e*(d^2*e^2 + 3*c*d*e*f + c^2*f^2) + a*(d^3*e^3 + 9*c*d^2*e^2*f + 9*c^2*d*e*f^2 + c^3*f^3))*x^7)/7 + ((3*a*d*f*(d^2*e^2 + 3*c*d*e*f + c^2*f^2) + b*(d^3*e^3 + 9*c*d^2*e^2*f + 9*c^2*d*e*f^2 + c^3*f^3))*x^9)/9 + (3*d*f*(a*d*f*(d*e + c*f) + b*(d^2*e^2 + 3*c*d*e*f + c^2*f^2))*x^11)/11 + (d^2*f^2*(a*d*f + 3*b*(d*e + c*f))*x^13)/13 + (b*d^3*f^3*x^15)/15

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^3 (e + fx^2)^3 dx &= \int (ac^3e^3 + c^2e^2(bce + 3a(de + cf))x^2 + 3ce(bce(de + cf) + a(d^2e^2 + 3cde^2 + 3c^2de^2 + d^3e^3))x^4 + \\ &= ac^3e^3x + \frac{1}{3}c^2e^2(bce + 3a(de + cf))x^3 + \frac{3}{5}ce(bce(de + cf) + a(d^2e^2 + 3cde^2 + 3c^2de^2 + d^3e^3))x^5 + \dots \end{aligned}$$

Mathematica [A] time = 0.13, size = 310, normalized size = 1.00

$$\frac{3}{11}dfx^{11} (adf(cf + de) + b(c^2f^2 + 3cdef + d^2e^2)) + \frac{3}{5}cex^5 (a(c^2f^2 + 3cdef + d^2e^2) + bce(cf + de)) + \frac{1}{3}c^2e^2x^3$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^3,x]

[Out] a*c^3*e^3*x + (c^2*e^2*(b*c*e + 3*a*(d*e + c*f))*x^3)/3 + (3*c*e*(b*c*e*(d*e + c*f) + a*(d^2*e^2 + 3*c*d*e*f + c^2*f^2))*x^5)/5 + ((3*b*c*e*(d^2*e^2 + 3*c*d*e*f + c^2*f^2) + a*(d^3*e^3 + 9*c*d^2*e^2*f + 9*c^2*d*e*f^2 + c^3*f^3))*x^7)/7 + ((3*a*d*f*(d^2*e^2 + 3*c*d*e*f + c^2*f^2) + b*(d^3*e^3 + 9*c*d^2*e^2*f + 9*c^2*d*e*f^2 + c^3*f^3))*x^9)/9 + (3*d*f*(a*d*f*(d*e + c*f) + b*(d^2*e^2 + 3*c*d*e*f + c^2*f^2))*x^11)/11 + (d^2*f^2*(a*d*f + 3*b*(d*e + c*f))*x^13)/13 + (b*d^3*f^3*x^15)/15

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^3 (e + fx^2)^3 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^3, x]

fricas [A] time = 1.21, size = 409, normalized size = 1.32

$\frac{1}{15} b^3 f^3 x^{15} + \frac{3}{13} b^2 c d^2 f^3 x^{13} + \frac{1}{13} a d^3 f^3 x^{13} + \frac{3}{13} b^2 c d^2 f^2 x^{11} e + \frac{3}{11} b^2 c^2 d f^3 x^{11} + \frac{3}{11} a^2 c d^2 f^3 x^{11} + \frac{9}{11} b^2 c^2 d^2 f^2 x^{11} e + \frac{3}{11} a^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e^2 + \frac{1}{9} b^2 c^3 f^3 x^9 + \frac{1}{3} a^2 c^2 d^2 f^3 x^9 + b^2 c^2 d^2 f^2 x^9 e + a^2 c^2 d^2 f^2 x^9 e + b^2 c^2 d^2 f^2 x^9 e^2 + \frac{1}{3} a^2 d^3 f^2 x^9 e^2 + \frac{1}{7} a^2 c^3 f^3 x^7 + \frac{1}{9} b^2 d^3 x^9 e^3 + \frac{3}{7} b^2 c^3 f^2 x^7 e + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e + \frac{9}{7} b^2 c^2 d^2 f^2 x^7 e^2 + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e^2 + \frac{3}{7} b^2 c^2 d^2 x^7 e^3 + \frac{1}{7} a^2 d^3 x^7 e^3 + \frac{3}{5} a^2 c^3 f^2 x^5 e + \frac{3}{5} b^2 c^3 f^2 x^5 e^2 + \frac{9}{5} a^2 c^2 d^2 f^2 x^5 e^2 + \frac{3}{5} b^2 c^2 d^2 x^5 e^3 + \frac{3}{5} a^2 c^2 d^2 x^5 e^3 + a^2 c^3 f^2 x^3 e^2 + \frac{1}{3} b^2 c^3 x^3 e^3 + a^2 c^2 d^2 x^3 e^3 + a^2 c^3 x^3 e^3$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^3,x, algorithm="fricas")

[Out] $\frac{1}{15} b^3 f^3 x^{15} + \frac{3}{13} b^2 c d^2 f^3 x^{13} + \frac{1}{13} a d^3 f^3 x^{13} + \frac{3}{13} b^2 c d^2 f^2 x^{11} e + \frac{3}{11} b^2 c^2 d f^3 x^{11} + \frac{3}{11} a^2 c d^2 f^3 x^{11} + \frac{9}{11} b^2 c^2 d^2 f^2 x^{11} e + \frac{3}{11} a^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e^2 + \frac{1}{9} b^2 c^3 f^3 x^9 + \frac{1}{3} a^2 c^2 d^2 f^3 x^9 + b^2 c^2 d^2 f^2 x^9 e + a^2 c^2 d^2 f^2 x^9 e + b^2 c^2 d^2 f^2 x^9 e^2 + \frac{1}{3} a^2 d^3 f^2 x^9 e^2 + \frac{1}{7} a^2 c^3 f^3 x^7 + \frac{1}{9} b^2 d^3 x^9 e^3 + \frac{3}{7} b^2 c^3 f^2 x^7 e + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e + \frac{9}{7} b^2 c^2 d^2 f^2 x^7 e^2 + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e^2 + \frac{3}{7} b^2 c^2 d^2 x^7 e^3 + \frac{1}{7} a^2 d^3 x^7 e^3 + \frac{3}{5} a^2 c^3 f^2 x^5 e + \frac{3}{5} b^2 c^3 f^2 x^5 e^2 + \frac{9}{5} a^2 c^2 d^2 f^2 x^5 e^2 + \frac{3}{5} b^2 c^2 d^2 x^5 e^3 + \frac{3}{5} a^2 c^2 d^2 x^5 e^3 + a^2 c^3 f^2 x^3 e^2 + \frac{1}{3} b^2 c^3 x^3 e^3 + a^2 c^2 d^2 x^3 e^3 + a^2 c^3 x^3 e^3$

giac [A] time = 0.38, size = 401, normalized size = 1.29

$\frac{1}{15} b^3 f^3 x^{15} + \frac{3}{13} b^2 c d^2 f^3 x^{13} + \frac{1}{13} a d^3 f^3 x^{13} + \frac{3}{13} b^2 c d^2 f^2 x^{11} e + \frac{3}{11} b^2 c^2 d f^3 x^{11} + \frac{3}{11} a^2 c d^2 f^3 x^{11} + \frac{9}{11} b^2 c^2 d^2 f^2 x^{11} e + \frac{3}{11} a^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e^2 + \frac{1}{9} b^2 c^3 f^3 x^9 + \frac{1}{3} a^2 c^2 d^2 f^3 x^9 + b^2 c^2 d^2 f^2 x^9 e + a^2 c^2 d^2 f^2 x^9 e + b^2 c^2 d^2 f^2 x^9 e^2 + \frac{1}{3} a^2 d^3 f^2 x^9 e^2 + \frac{1}{7} a^2 c^3 f^3 x^7 + \frac{1}{9} b^2 d^3 x^9 e^3 + \frac{3}{7} b^2 c^3 f^2 x^7 e + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e + \frac{9}{7} b^2 c^2 d^2 f^2 x^7 e^2 + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e^2 + \frac{3}{7} b^2 c^2 d^2 x^7 e^3 + \frac{1}{7} a^2 d^3 x^7 e^3 + \frac{3}{5} a^2 c^3 f^2 x^5 e + \frac{3}{5} b^2 c^3 f^2 x^5 e^2 + \frac{9}{5} a^2 c^2 d^2 f^2 x^5 e^2 + \frac{3}{5} b^2 c^2 d^2 x^5 e^3 + \frac{3}{5} a^2 c^2 d^2 x^5 e^3 + a^2 c^3 f^2 x^3 e^2 + \frac{1}{3} b^2 c^3 x^3 e^3 + a^2 c^2 d^2 x^3 e^3 + a^2 c^3 x^3 e^3$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^3,x, algorithm="giac")

[Out] $\frac{1}{15} b^3 d^3 f^3 x^{15} + \frac{3}{13} b^2 c d^2 f^3 x^{13} + \frac{1}{13} a d^3 f^3 x^{13} + \frac{3}{13} b^2 c d^2 f^2 x^{11} e + \frac{3}{11} b^2 c^2 d f^3 x^{11} + \frac{3}{11} a^2 c d^2 f^3 x^{11} + \frac{9}{11} b^2 c^2 d^2 f^2 x^{11} e + \frac{3}{11} a^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e + \frac{3}{11} b^2 d^3 f^2 x^{11} e^2 + \frac{1}{9} b^2 c^3 f^3 x^9 + \frac{1}{3} a^2 c^2 d^2 f^3 x^9 + b^2 c^2 d^2 f^2 x^9 e + a^2 c^2 d^2 f^2 x^9 e + b^2 c^2 d^2 f^2 x^9 e^2 + \frac{1}{3} a^2 d^3 f^2 x^9 e^2 + \frac{1}{7} a^2 c^3 f^3 x^7 + \frac{1}{9} b^2 d^3 x^9 e^3 + \frac{3}{7} b^2 c^3 f^2 x^7 e + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e + \frac{9}{7} b^2 c^2 d^2 f^2 x^7 e^2 + \frac{9}{7} a^2 c^2 d^2 f^2 x^7 e^2 + \frac{3}{7} b^2 c^2 d^2 x^7 e^3 + \frac{1}{7} a^2 d^3 x^7 e^3 + \frac{3}{5} a^2 c^3 f^2 x^5 e + \frac{3}{5} b^2 c^3 f^2 x^5 e^2 + \frac{9}{5} a^2 c^2 d^2 f^2 x^5 e^2 + \frac{3}{5} b^2 c^2 d^2 x^5 e^3 + \frac{3}{5} a^2 c^2 d^2 x^5 e^3 + a^2 c^3 f^2 x^3 e^2 + \frac{1}{3} b^2 c^3 x^3 e^3 + a^2 c^2 d^2 x^3 e^3 + a^2 c^3 x^3 e^3$

maple [A] time = 0.00, size = 339, normalized size = 1.09

$\frac{b^3 d^3 f^3 x^{15}}{15} + \frac{3 b^2 c d^2 f^3 x^{13} + (a d^3 + 3 b^2 c d^2) f^3 x^{13}}{13} + \frac{3 b^2 c^2 d f^3 x^{11} + (3 a^2 c d^2 + 3 b^2 c^2 d) f^3 x^{11}}{11} + \frac{9 b^2 c^2 d^2 f^2 x^{11} e + (3 a^2 d^3 + 3 b^2 d^3) f^2 x^{11} e + 3 (3 a^2 c d^2 + 3 b^2 c^2 d) f^2 x^{11} e^2}{11} + \frac{b^2 c^3 f^3 x^9 + (a^2 c^2 d^2 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2) x^9}{9} + \frac{b^2 c^2 d^2 f^2 x^9 e + (a^2 c^2 d^2 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2) x^9 e + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^9 e^2}{9} + \frac{b^2 c^3 f^3 x^7 + (a^2 c^3 + 3 b^2 c^3) f^3 x^7}{7} + \frac{b^2 d^3 x^9 e^3 + (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^9 e^3 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^9 e^3}{9} + \frac{3 b^2 c^3 f^2 x^7 e + (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^7 e + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^7 e^2}{7} + \frac{3 b^2 c^2 d^2 x^7 e^3 + (a^2 d^3 + 3 b^2 c^2 d) f^2 x^7 e^3 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^7 e^3}{7} + \frac{3 b^2 c^2 d^2 x^5 e^3 + (a^2 c^2 d + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2) x^5 e^3 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^5 e^3}{5} + \frac{a^2 c^3 f^2 x^3 e^2 + (3 a^2 c^3 + 3 b^2 c^3) f^2 x^3 e^2 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^3 e^2}{3} + \frac{b^2 c^3 x^3 e^3 + (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^3 e^3 + 3 (3 a^2 c^2 d + 3 b^2 c^2 d) f^2 x^3 e^3}{3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^3,x)

[Out] $\frac{1}{15} b^3 d^3 f^3 x^{15} + \frac{1}{13} ((a d^3 + 3 b^2 c d^2) f^3 + 3 b^2 d^3 e f^2) x^{13} + \frac{1}{11} ((3 a^2 c d^2 + 3 b^2 c^2 d) f^3 + 3 (a d^3 + 3 b^2 c d^2) e f^2 + 3 b^2 d^3 e^2 f) x^{11} + \frac{1}{9} ((3 a^2 c^2 d + b^2 c^3) f^3 + 3 (3 a^2 c d^2 + 3 b^2 c^2 d) e f^2 + 3 (a d^3 + 3 b^2 c d^2) e^2 f + 3 b^2 d^3 e^3) x^9 + \frac{1}{7} (a^2 c^3 f^3 + 3 (3 a^2 c^2 d + b^2 c^3) e f^2 + 3 (3 a^2 c d^2 + 3 b^2 c^2 d) e^2 f + (a d^3 + 3 b^2 c d^2) e^3) x^7 + \frac{1}{5} (3 a^2 c^3 e f^2 + 3 (3 a^2 c^2 d + b^2 c^3) e^2 f + (3 a^2 c d^2 + 3 b^2 c^2 d) e^3) x^5 + \frac{1}{3} (3 a^2 c^3 e^2 f + (3 a^2 c^2 d + b^2 c^3) e^3) x^3 + a^2 c^3 e^3 x$

maxima [A] time = 0.86, size = 326, normalized size = 1.05

$\frac{1}{15} b^3 d^3 f^3 x^{15} + \frac{1}{13} (3 a^2 c d^2 + 3 b^2 c^2 d) f^3 x^{13} + \frac{1}{11} (3 a^2 c d^2 + 3 b^2 c^2 d) f^3 x^{11} + \frac{1}{9} (3 a^2 c^2 d + 3 b^2 c^3) f^3 x^9 + \frac{1}{7} (3 a^2 c^3 + 3 (3 a^2 c^2 d + b^2 c^3) f^2) x^7 + \frac{1}{5} (3 a^2 c^3 e f^2 + 3 (3 a^2 c^2 d + b^2 c^3) e^2 f + (3 a^2 c d^2 + 3 b^2 c^2 d) e^3) x^5 + \frac{1}{3} (3 a^2 c^3 e^2 f + (3 a^2 c^2 d + b^2 c^3) e^3) x^3 + a^2 c^3 e^3 x$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^3,x, algorithm="maxima")

[Out] $1/15*b*d^3*f^3*x^{15} + 1/13*(3*b*d^3*e*f^2 + (3*b*c*d^2 + a*d^3)*f^3)*x^{13} + 3/11*(b*d^3*e^2*f + (3*b*c*d^2 + a*d^3)*e*f^2 + (b*c^2*d + a*c*d^2)*f^3)*x^{11} + 1/9*(b*d^3*e^3 + 3*(3*b*c*d^2 + a*d^3)*e^2*f + 9*(b*c^2*d + a*c*d^2)*e*f^2 + (b*c^3 + 3*a*c^2*d)*f^3)*x^9 + a*c^3*e^3*x + 1/7*(a*c^3*f^3 + (3*b*c*d^2 + a*d^3)*e^3 + 9*(b*c^2*d + a*c*d^2)*e^2*f + 3*(b*c^3 + 3*a*c^2*d)*e*f^2)*x^7 + 3/5*(a*c^3*e*f^2 + (b*c^2*d + a*c*d^2)*e^3 + (b*c^3 + 3*a*c^2*d)*e^2*f)*x^5 + 1/3*(3*a*c^3*e^2*f + (b*c^3 + 3*a*c^2*d)*e^3)*x^3$

mupad [B] time = 0.12, size = 335, normalized size = 1.08

$x^{\frac{15}{15}} \left(\frac{b^3 d^3 f^3}{15} + \frac{3 a c^3 e^3}{15} + \frac{3 b^2 d^2 e f^2}{15} + \frac{9 a c^2 d^2 e f^2}{15} + \frac{3 a c^2 d^2 e^2 f}{15} \right) + x^{11} \left(\frac{3 b^2 d^3 e f^2}{11} + \frac{9 b c^2 d^2 e f^2}{11} + \frac{3 a c^2 d^2 e f^2}{11} + \frac{3 b^2 d^2 e^2 f}{11} + \frac{3 a d^3 e f^2}{11} \right) + x^9 \left(\frac{3 b^2 d^3 e^2 f}{9} + \frac{9 a c^2 d^2 e f^2}{9} + \frac{9 a c^2 d^2 e^2 f}{9} + \frac{3 b c^2 d^2 e f^2}{9} + \frac{3 a d^3 e^2 f}{9} \right) + x^7 \left(\frac{b^3 d^3 e^3}{7} + \frac{9 a c^3 e^3}{7} + \frac{9 a c^2 d^2 e f^2}{7} + \frac{9 a c^2 d^2 e^2 f}{7} + \frac{3 b c^2 d^2 e f^2}{7} \right) + x^5 \left(\frac{3 a c^3 e f^2}{5} + \frac{3 a c^3 e^2 f}{5} + \frac{3 b c^2 d^2 e f^2}{5} + \frac{3 a c^2 d^2 e^2 f}{5} + \frac{3 a c^2 d^2 e^3}{5} \right) + x^3 \left(\frac{3 a^2 c^3 e^2 f}{3} + \frac{3 a^2 c^3 e^3}{3} + \frac{3 b c^2 d^2 e^2 f}{3} + \frac{3 a c^2 d^2 e^3}{3} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^3,x)

[Out] $x^5*((3*a*c*d^2*e^3)/5 + (3*b*c^2*d*e^3)/5 + (3*a*c^3*e*f^2)/5 + (3*b*c^3*e^2*f)/5 + (9*a*c^2*d*e^2*f)/5) + x^{11}*((3*a*c*d^2*f^3)/11 + (3*b*c^2*d*f^3)/11 + (3*a*d^3*e*f^2)/11 + (3*b*d^3*e^2*f)/11 + (9*b*c*d^2*e*f^2)/11) + x^7*((a*c^3*f^3)/7 + (a*d^3*e^3)/7 + (3*b*c*d^2*e^3)/7 + (3*b*c^3*e*f^2)/7 + (9*a*c*d^2*e^2*f)/7 + (9*a*c^2*d*e*f^2)/7 + (9*b*c^2*d*e^2*f)/7) + x^9*((b*c^3*f^3)/9 + (b*d^3*e^3)/9 + (a*c^2*d*f^3)/3 + (a*d^3*e^2*f)/3 + a*c*d^2*e*f^2 + b*c*d^2*e^2*f + b*c^2*d*e*f^2) + (b*d^3*f^3*x^{15})/15 + (c^2*e^2*x^3*(3*a*c*f + 3*a*d*e + b*c*e))/3 + (d^2*f^2*x^{13}*(a*d*f + 3*b*c*f + 3*b*d*e))/13 + a*c^3*e^3*x$

sympy [A] time = 0.13, size = 423, normalized size = 1.36

$a^3 c^3 e^3 x^5 + b^3 d^3 f^3 x^{15} + x^{13} (a^3 d^3 f^3 + 3 b^2 c^2 d^2 e f^2 + 3 b^2 d^3 e^2 f + 3 a^2 c^3 e^2 f) + x^{11} (3 a^2 c^3 e f^2 + 3 a^2 c^3 e^2 f + 3 a^2 c^2 d^2 e f^2 + 3 a^2 c^2 d^2 e^2 f) + x^9 (a^3 c^3 f^3 + a^3 d^3 e^3 + a^2 c^2 d^2 e f^2 + b^2 c^2 d^2 e f^2 + b^2 c^2 d^2 e^2 f) + x^7 (a^3 c^3 f^3 + 9 a^2 c^2 d^2 e f^2 + 9 a^2 c^2 d^2 e^2 f + 3 b^2 c^2 d^2 e f^2 + 3 b^2 c^2 d^2 e^2 f) + x^5 (a^3 c^3 e f^2 + a^3 c^3 e^2 f + 3 b^2 c^2 d^2 e f^2 + 3 a^2 c^2 d^2 e^2 f + 3 a^2 c^2 d^2 e^3) + x^3 (a^2 c^3 e^2 f + a^2 c^3 e^3 + b^2 c^2 d^2 e^2 f + b^2 c^2 d^2 e^3)$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3*(f*x**2+e)**3,x)

[Out] $a*c**3*e**3*x + b*d**3*f**3*x**15/15 + x**13*(a*d**3*f**3/13 + 3*b*c*d**2*f**3/13 + 3*b*d**3*e*f**2/13) + x**11*(3*a*c*d**2*f**3/11 + 3*a*d**3*e*f**2/11 + 3*b*c**2*d*f**3/11 + 9*b*c*d**2*e*f**2/11 + 3*b*d**3*e**2*f/11) + x**9*(a*c**2*d*f**3/3 + a*c*d**2*e*f**2 + a*d**3*e**2*f/3 + b*c**3*f**3/9 + b*c**2*d*e*f**2 + b*c*d**2*e**2*f + b*d**3*e**3/9) + x**7*(a*c**3*f**3/7 + 9*a*c**2*d*e*f**2/7 + 9*a*c*d**2*e**2*f/7 + a*d**3*e**3/7 + 3*b*c**3*e*f**2/7 + 9*b*c**2*d*e**2*f/7 + 3*b*c*d**2*e**3/7) + x**5*(3*a*c**3*e*f**2/5 + 9*a*c**2*d*e**2*f/5 + 3*a*c*d**2*e**3/5 + 3*b*c**3*e**2*f/5 + 3*b*c**2*d*e**3/5) + x**3*(a*c**3*e**2*f + a*c**2*d*e**3 + b*c**3*e**3/3)$

$$3.17 \quad \int (a + bx^2)(c + dx^2)^3 (e + fx^2)^2 dx$$

Optimal. Leaf size=226

$$\frac{1}{9}dx^9 (adf(3cf + 2de) + b(3c^2f^2 + 6cdef + d^2e^2)) + \frac{1}{7}x^7 (ad(3c^2f^2 + 6cdef + d^2e^2) + bc(c^2f^2 + 6cdef + 3d^2e^2))$$

Rubi [A] time = 0.22, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.038$, Rules used = {521}

$$\frac{1}{9}dx^9 (adf(3cf + 2de) + b(3c^2f^2 + 6cdef + d^2e^2)) + \frac{1}{7}x^7 (ad(3c^2f^2 + 6cdef + d^2e^2) + bc(c^2f^2 + 6cdef + 3d^2e^2)) + \frac{1}{5}cx^5 (a(c^2f^2 + 6cdef + 3d^2e^2) + bce(2cf + 3de)) + \frac{1}{3}c^2cx^3(2acf + 3ade + bce) + \frac{1}{11}d^2fx^{11}(adf + 3bcf + 2bde) + ac^3e^2x + \frac{1}{13}bd^2fx^{13}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^2,x]

[Out] a*c^3*e^2*x + (c^2*e*(b*c*e + 3*a*d*e + 2*a*c*f)*x^3)/3 + (c*(b*c*e*(3*d*e + 2*c*f) + a*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2))*x^5)/5 + ((b*c*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2) + a*d*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^7)/7 + (d*(a*d*f*(2*d*e + 3*c*f) + b*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^9)/9 + (d^2*f*(2*b*d*e + 3*b*c*f + a*d*f)*x^11)/11 + (b*d^3*f^2*x^13)/13

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^3 (e + fx^2)^2 dx &= \int (ac^3e^2 + c^2e(bce + 3ade + 2acf)x^2 + c(bce(3de + 2cf) + a(3d^2e^2 + 6cdef + d^2e^2))x^4 + \\ &= ac^3e^2x + \frac{1}{3}c^2e(bce + 3ade + 2acf)x^3 + \frac{1}{5}c(bce(3de + 2cf) + a(3d^2e^2 + 6cdef + d^2e^2))x^5 + \dots \end{aligned}$$

Mathematica [A] time = 0.09, size = 226, normalized size = 1.00

$$\frac{1}{9}dx^9 (adf(3cf + 2de) + b(3c^2f^2 + 6cdef + d^2e^2)) + \frac{1}{7}x^7 (ad(3c^2f^2 + 6cdef + d^2e^2) + bc(c^2f^2 + 6cdef + 3d^2e^2)) + \frac{1}{5}cx^5 (a(c^2f^2 + 6cdef + 3d^2e^2) + bce(2cf + 3de)) + \frac{1}{3}c^2cx^3(2acf + 3ade + bce) + \frac{1}{11}d^2fx^{11}(adf + 3bcf + 2bde) + ac^3e^2x + \frac{1}{13}bd^2fx^{13}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^2,x]

[Out] a*c^3*e^2*x + (c^2*e*(b*c*e + 3*a*d*e + 2*a*c*f)*x^3)/3 + (c*(b*c*e*(3*d*e + 2*c*f) + a*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2))*x^5)/5 + ((b*c*(3*d^2*e^2 + 6*c*d*e*f + c^2*f^2) + a*d*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^7)/7 + (d*(a*d*f*(2*d*e + 3*c*f) + b*(d^2*e^2 + 6*c*d*e*f + 3*c^2*f^2))*x^9)/9 + (d^2*f*(2*b*d*e + 3*b*c*f + a*d*f)*x^11)/11 + (b*d^3*f^2*x^13)/13

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^3 (e + fx^2)^2 dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^2, x]

fricas [A] time = 1.22, size = 289, normalized size = 1.28

$$\frac{1}{13}x^{13}f^2b + \frac{2}{11}x^{11}fadfb + \frac{3}{11}x^{11}f^2dcb + \frac{1}{11}x^{11}f^2dca + \frac{1}{9}x^9f^2dcb + \frac{2}{3}x^9fadfb + \frac{1}{3}x^9fadca + \frac{2}{9}x^9fadcb + \frac{1}{3}x^9fadca + \frac{1}{3}x^9fadca + \frac{2}{7}x^7fadcb + \frac{6}{7}x^7fadcb + \frac{1}{7}x^7fadcb + \frac{1}{7}x^7fadcb + \frac{6}{7}x^7fadcb + \frac{3}{7}x^7fadcb + \frac{3}{5}x^5fadcb + \frac{6}{5}x^5fadcb + \frac{1}{5}x^5fadcb + \frac{1}{5}x^5fadcb + \frac{1}{3}x^3fadcb + x^2dca + \frac{2}{3}fadca + x^2dca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^2,x, algorithm="fricas")

[Out] $1/13*x^{13}*f^2*d^3*b + 2/11*x^{11}*f*e*d^3*b + 3/11*x^{11}*f^2*d^2*c*b + 1/11*x^{11}*f^2*d^3*a + 1/9*x^9*e^2*d^3*b + 2/3*x^9*f*e*d^2*c*b + 1/3*x^9*f^2*d*c^2*b + 2/9*x^9*f*e*d^3*a + 1/3*x^9*f^2*d^2*c*a + 3/7*x^7*e^2*d^2*c*b + 6/7*x^7*f*e*d*c^2*b + 1/7*x^7*f^2*c^3*b + 1/7*x^7*e^2*d^3*a + 6/7*x^7*f*e*d^2*c*a + 3/7*x^7*f^2*d*c^2*a + 3/5*x^5*e^2*d*c^2*b + 2/5*x^5*f*e*c^3*b + 3/5*x^5*e^2*d^2*c*a + 6/5*x^5*f*e*d*c^2*a + 1/5*x^5*f^2*c^3*a + 1/3*x^3*e^2*c^3*b + x^3*e^2*d*c^2*a + 2/3*x^3*f*e*c^3*a + x*e^2*c^3*a$

giac [A] time = 0.35, size = 289, normalized size = 1.28

$$\frac{1}{13}bd^3f^2x^{13} + \frac{2}{11}bdf^2x^{11} + \frac{1}{11}ad^3f^2x^{11} + \frac{2}{11}bd^3f^2x^{11} + \frac{1}{9}bd^3f^2x^9 + \frac{1}{3}ad^3f^2x^9 + \frac{2}{3}bd^3f^2x^9 + \frac{1}{3}ad^3f^2x^9 + \frac{1}{3}ad^3f^2x^9 + \frac{2}{7}bd^3f^2x^7 + \frac{6}{7}bd^3f^2x^7 + \frac{1}{7}bd^3f^2x^7 + \frac{1}{7}bd^3f^2x^7 + \frac{6}{7}bd^3f^2x^7 + \frac{3}{7}bd^3f^2x^7 + \frac{3}{5}bd^3f^2x^5 + \frac{6}{5}bd^3f^2x^5 + \frac{1}{5}bd^3f^2x^5 + \frac{1}{5}bd^3f^2x^5 + \frac{1}{3}bd^3f^2x^3 + x^2dca + \frac{2}{3}fadca + x^2dca$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^2,x, algorithm="giac")

[Out] $1/13*b*d^3*f^2*x^{13} + 3/11*b*c*d^2*f^2*x^{11} + 1/11*a*d^3*f^2*x^{11} + 2/11*b*d^3*f*x^{11}*e + 1/3*b*c^2*d*f^2*x^9 + 1/3*a*c*d^2*f^2*x^9 + 2/3*b*c*d^2*f*x^9*e + 2/9*a*d^3*f*x^9*e + 1/9*b*d^3*x^9*e^2 + 1/7*b*c^3*f^2*x^7 + 3/7*a*c^2*d*f^2*x^7 + 6/7*b*c^2*d*f*x^7*e + 6/7*a*c*d^2*f*x^7*e + 3/7*b*c*d^2*x^7*e^2 + 1/7*a*d^3*x^7*e^2 + 1/5*a*c^3*f^2*x^5 + 2/5*b*c^3*f*x^5*e + 6/5*a*c^2*d*f*x^5*e + 3/5*b*c^2*d*x^5*e^2 + 3/5*a*c*d^2*x^5*e^2 + 2/3*a*c^3*f*x^3*e + 1/3*b*c^3*x^3*e^2 + a*c^2*d*x^3*e^2 + a*c^3*x*e^2$

maple [A] time = 0.00, size = 244, normalized size = 1.08

$$\frac{bd^3f^2x^{13}}{13} + \frac{(2bd^3ef + (ad^3 + 3bcd^2)f^2)x^{11}}{11} + \frac{(bd^3e^2 + 2(ad^3 + 3bcd^2)ef + (3acd^2 + 3bc^2d)f^2)x^9}{9} + a^2c^2x + \frac{((ad^3 + 3bcd^2)e^2 + 2(3acd^2 + 3bc^2d)ef + (3a^2d + bc^2)f^2)x^7}{7} + \frac{(a^2e^2 + (3acd^2 + 3bc^2d)e^2 + 2(3a^2d + bc^2)ef)x^5}{5} + \frac{(2a^2ef + (3a^2d + bc^2)e^2)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^2,x)

[Out] $1/13*b*d^3*f^2*x^{13} + 1/11*((ad^3 + 3b*c*d^2)*f^2 + 2*b*d^3*e*f)*x^{11} + 1/9*((3*a*c*d^2 + 3*b*c^2*d)*f^2 + 2*(ad^3 + 3b*c*d^2)*e*f + b*d^3*e^2)*x^9 + 1/7*((3*a*c^2*d + b*c^3)*f^2 + 2*(3*a*c*d^2 + 3*b*c^2*d)*e*f + (ad^3 + 3b*c*d^2)*e^2)*x^7 + 1/5*(a*c^3*f^2 + 2*(3*a*c^2*d + b*c^3)*e*f + (3*a*c*d^2 + 3*b*c^2*d)*e^2)*x^5 + 1/3*(2*a*c^3*e*f + (3*a*c^2*d + b*c^3)*e^2)*x^3 + a*c^3*e^2*x$

maxima [A] time = 0.84, size = 239, normalized size = 1.06

$$\frac{1}{13}bd^3f^2x^{13} + \frac{1}{11}(2bd^3ef + (3bcd^2 + ad^3)f^2)x^{11} + \frac{1}{9}(bd^3e^2 + 2(3bcd^2 + ad^3)ef + 3(bc^2d + acd^2)f^2)x^9 + \frac{1}{7}((3bcd^2 + ad^3)e^2 + 6(bc^2d + acd^2)ef + (bc^3 + 3acd^2)f^2)x^7 + a^2c^2x + \frac{1}{5}(a^2e^2 + 3(bc^2d + acd^2)e^2 + 2(bc^3 + 3acd^2)ef)x^5 + \frac{1}{3}(2a^2ef + (bc^3 + 3acd^2)e^2)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e)^2,x, algorithm="maxima")

[Out] $1/13*b*d^3*f^2*x^{13} + 1/11*(2*b*d^3*e*f + (3*b*c*d^2 + a*d^3)*f^2)*x^{11} + 1/9*(b*d^3*e^2 + 2*(3*b*c*d^2 + a*d^3)*e*f + 3*(b*c^2*d + a*c*d^2)*f^2)*x^9 + 1/7*((3*b*c*d^2 + a*d^3)*e^2 + 6*(b*c^2*d + a*c*d^2)*e*f + (b*c^3 + 3*a*c^2*d)*f^2)*x^7 + a*c^3*e^2*x + 1/5*(a*c^3*f^2 + 3*(b*c^2*d + a*c*d^2)*e^2 + 2*(b*c^3 + 3*a*c^2*d)*e*f)*x^5 + 1/3*(2*a*c^3*e*f + (b*c^3 + 3*a*c^2*d)*e^2)*x^3$

mupad [B] time = 0.85, size = 233, normalized size = 1.03

$$x^5 \left(\frac{2bc^2ef}{5} + \frac{ac^3f^2}{5} + \frac{3b^2cd^2}{5} + \frac{6ac^2def}{5} + \frac{3acd^2e^2}{5} \right) + x^9 \left(\frac{b^2d^2f^2}{3} + \frac{2bcd^2ef}{3} + \frac{acd^2f^2}{3} + \frac{bd^3e^2}{9} + \frac{2ad^2ef}{9} \right) + x^7 \left(\frac{b^2f^2}{7} + \frac{6b^2def}{7} + \frac{3ac^2d^2f^2}{7} + \frac{3bcd^2e^2}{7} + \frac{6acd^2ef}{7} + \frac{ad^3e^2}{7} \right) + \frac{bd^3f^2x^{11}}{13} + \frac{c^2ex^3(2acf+3ade+bce)}{3} + \frac{d^2fx^{11}(adf+3bcf+2bde)}{11} + ac^3e^2x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^3*(e + f*x^2)^2,x)

[Out] $x^5 \left(\frac{a^3c^3f^2}{5} + \frac{2b^2c^3e^2f}{5} + \frac{3a^2c^2d^2e^2}{5} + \frac{3b^2c^2d^2e^2}{5} + \frac{6a^2c^2d^2e^2f}{5} \right) + x^9 \left(\frac{b^2d^3e^2}{9} + \frac{2a^2d^3e^2f}{9} + \frac{a^2c^2d^2f^2}{3} + \frac{b^2c^2d^2f^2}{3} + \frac{2b^2c^2d^2e^2f}{3} \right) + x^7 \left(\frac{a^2d^3e^2}{7} + \frac{b^2c^3f^2}{7} + \frac{3a^2c^2d^2f^2}{7} + \frac{3b^2c^2d^2e^2}{7} + \frac{6a^2c^2d^2e^2f}{7} + \frac{6b^2c^2d^2e^2f}{7} \right) + \frac{b^2d^3f^2x^{13}}{13} + \frac{c^2e^2x^3(2acf+3ade+bce)}{3} + \frac{d^2f^2x^{11}(adf+3bcf+2bde)}{11} + a^2c^3e^2x$

sympy [A] time = 0.11, size = 304, normalized size = 1.35

$$ac^3e^2x + \frac{bd^3f^2x^{13}}{13} + x^{11} \left(\frac{ad^3f^2}{11} + \frac{3bcd^2f^2}{11} + \frac{2bd^3ef}{11} \right) + x^9 \left(\frac{acd^2f^2}{3} + \frac{2ad^3ef}{9} + \frac{bc^2d^2f^2}{3} + \frac{2bcd^2ef}{3} + \frac{bd^3e^2}{9} \right) + x^7 \left(\frac{3ac^2d^2f^2}{7} + \frac{6acd^2ef}{7} + \frac{ad^3e^2}{7} + \frac{bc^2f^2}{7} + \frac{6bc^2def}{7} + \frac{3bcd^2e^2}{7} \right) + x^5 \left(\frac{ac^3f^2}{5} + \frac{6ac^2def}{5} + \frac{3acd^2e^2}{5} + \frac{2bc^3ef}{5} + \frac{3bc^2de^2}{5} \right) + x^3 \left(\frac{2ac^3ef}{3} + ac^2de^2 + \frac{bc^3e^2}{3} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3*(f*x**2+e)**2,x)

[Out] $a^2c^3e^2x + b^2d^3f^2x^{13}/13 + x^{11} \left(\frac{ad^3f^2}{11} + \frac{3b^2cd^2f^2}{11} + \frac{2b^2d^3ef}{11} \right) + x^9 \left(\frac{acd^2f^2}{3} + \frac{2ad^3ef}{9} + \frac{bc^2d^2f^2}{3} + \frac{2bcd^2ef}{3} + \frac{bd^3e^2}{9} \right) + x^7 \left(\frac{3a^2c^2d^2f^2}{7} + \frac{6a^2acd^2ef}{7} + \frac{ad^3e^2}{7} + \frac{bc^2f^2}{7} + \frac{6bc^2def}{7} + \frac{3bcd^2e^2}{7} \right) + x^5 \left(\frac{a^2c^3f^2}{5} + \frac{6a^2ac^2def}{5} + \frac{3a^2acd^2e^2}{5} + \frac{2b^2c^3ef}{5} + \frac{3b^2c^2de^2}{5} \right) + x^3 \left(\frac{2ac^3ef}{3} + a^2de^2 + \frac{bc^3e^2}{3} \right)$

$$3.18 \quad \int (a + bx^2)(c + dx^2)^3 (e + fx^2) dx$$

Optimal. Leaf size=130

$$\frac{1}{3}c^2x^3(acf+3ade+bce)+\frac{1}{9}d^2x^9(adf+3bcf+bde)+\frac{1}{7}dx^7(ad(3cf+de)+3bc(cf+de))+\frac{1}{5}cx^5(3ad(cf+de)+bc(cf+3de))+ac^3ex+\frac{1}{11}bd^3fx^{11}$$

Rubi [A] time = 0.13, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.042$, Rules used = {521}

$$\frac{1}{3}c^2x^3(acf+3ade+bce)+\frac{1}{9}d^2x^9(adf+3bcf+bde)+\frac{1}{7}dx^7(ad(3cf+de)+3bc(cf+de))+\frac{1}{5}cx^5(3ad(cf+de)+bc(cf+3de))+ac^3ex+\frac{1}{11}bd^3fx^{11}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2), x]

[Out] a*c^3*e*x + (c^2*(b*c*e + 3*a*d*e + a*c*f)*x^3)/3 + (c*(3*a*d*(d*e + c*f) + b*c*(3*d*e + c*f))*x^5)/5 + (d*(3*b*c*(d*e + c*f) + a*d*(d*e + 3*c*f))*x^7)/7 + (d^2*(b*d*e + 3*b*c*f + a*d*f)*x^9)/9 + (b*d^3*f*x^11)/11

Rule 521

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_))^(r_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x^n)^p*(c + d*x^n)^q*(e + f*x^n)^r, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IGtQ[r, 0]

Rubi steps

$$\begin{aligned} \int (a + bx^2)(c + dx^2)^3 (e + fx^2) dx &= \int (ac^3e + c^2(bce + 3ade + acf)x^2 + c(3ad(de + cf) + bc(3de + cf))x^4 \\ &\quad + ac^3ex + \frac{1}{3}c^2(bce + 3ade + acf)x^3 + \frac{1}{5}c(3ad(de + cf) + bc(3de + cf))x^5 \\ &\quad + \frac{1}{7}d(3bcf + bde) + \frac{1}{9}d^2x^9(adf + 3bcf + bde) + \frac{1}{7}dx^7(ad(3cf + de) + 3bc(cf + de)) + \frac{1}{5}cx^5(3ad(cf + de) + bc(cf + 3de)) + ac^3ex + \frac{1}{11}bd^3fx^{11} \end{aligned}$$

Mathematica [A] time = 0.06, size = 130, normalized size = 1.00

$$\frac{1}{3}c^2x^3(acf+3ade+bce)+\frac{1}{9}d^2x^9(adf+3bcf+bde)+\frac{1}{7}dx^7(ad(3cf+de)+3bc(cf+de))+\frac{1}{5}cx^5(3ad(cf+de)+bc(cf+3de))+ac^3ex+\frac{1}{11}bd^3fx^{11}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2), x]

[Out] a*c^3*e*x + (c^2*(b*c*e + 3*a*d*e + a*c*f)*x^3)/3 + (c*(3*a*d*(d*e + c*f) + b*c*(3*d*e + c*f))*x^5)/5 + (d*(3*b*c*(d*e + c*f) + a*d*(d*e + 3*c*f))*x^7)/7 + (d^2*(b*d*e + 3*b*c*f + a*d*f)*x^9)/9 + (b*d^3*f*x^11)/11

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a + bx^2)(c + dx^2)^3 (e + fx^2) dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2), x]

[Out] IntegrateAlgebraic[(a + b*x^2)*(c + d*x^2)^3*(e + f*x^2), x]

fricas [A] time = 0.85, size = 165, normalized size = 1.27

$$\frac{1}{11}x^{11}fd^3b+\frac{1}{9}x^9ad^3b+\frac{1}{3}x^9fd^2cb+\frac{1}{9}x^9fd^3a+\frac{3}{7}x^7ed^2cb+\frac{3}{7}x^7fd^2b+\frac{1}{7}x^7ed^3a+\frac{3}{7}x^7fd^2ca+\frac{3}{5}x^5edc^2b+\frac{1}{5}x^5fc^3b+\frac{3}{5}x^5ad^2ca+\frac{3}{5}x^5fd^2a+\frac{1}{3}x^3ec^3b+x^3adc^2a+\frac{1}{3}x^3fc^3a+xec^3a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e),x, algorithm="fricas")

[Out] $\frac{1}{11}x^{11}f*d^3*b + \frac{1}{9}x^9*e*d^3*b + \frac{1}{3}x^9*f*d^2*c*b + \frac{1}{9}x^9*f*d^3*a + \frac{3}{7}x^7*e*d^2*c*b + \frac{3}{7}x^7*f*d*c^2*b + \frac{1}{7}x^7*e*d^3*a + \frac{3}{7}x^7*f*d^2*c*a + \frac{3}{5}x^5*e*d*c^2*b + \frac{1}{5}x^5*f*c^3*b + \frac{3}{5}x^5*e*d^2*c*a + \frac{3}{5}x^5*f*d*c^2*a + \frac{1}{3}x^3*e*c^3*b + x^3*e*d*c^2*a + \frac{1}{3}x^3*f*c^3*a + x*e*c^3*a$

giac [A] time = 0.35, size = 173, normalized size = 1.33

$$\frac{1}{11}bd^3fx^{11} + \frac{1}{3}bcd^2fx^9 + \frac{1}{9}ad^3fx^9 + \frac{1}{9}bd^3x^9e + \frac{3}{7}bc^2dfx^7 + \frac{3}{7}acd^2fx^7 + \frac{3}{7}bcd^2x^7e + \frac{1}{7}ad^3x^7e + \frac{1}{5}bc^3fx^5 + \frac{3}{5}ac^2dfx^5 + \frac{3}{5}bc^2dx^5e + \frac{3}{5}acd^2x^5e + \frac{1}{3}ac^3fx^3 + \frac{1}{3}bc^3x^3e + ac^2dx^3e + ac^3xe$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e),x, algorithm="giac")

[Out] $\frac{1}{11}b*d^3*f*x^{11} + \frac{1}{3}b*c*d^2*f*x^9 + \frac{1}{9}a*d^3*f*x^9 + \frac{1}{9}b*d^3*x^9*e + \frac{3}{7}b*c^2*d*f*x^7 + \frac{3}{7}a*c*d^2*f*x^7 + \frac{3}{7}b*c*d^2*x^7*e + \frac{1}{7}a*d^3*x^7*e + \frac{1}{5}b*c^3*f*x^5 + \frac{3}{5}a*c^2*d*f*x^5 + \frac{3}{5}b*c^2*d*x^5*e + \frac{3}{5}a*c*d^2*x^5*e + \frac{1}{3}a*c^3*f*x^3 + \frac{1}{3}b*c^3*x^3*e + a*c^2*d*x^3*e + a*c^3*x^3*e$

maple [A] time = 0.00, size = 149, normalized size = 1.15

$$\frac{bd^3fx^{11}}{11} + \frac{(bd^3e + (ad^3 + 3bcd^2)f)x^9}{9} + \frac{((ad^3 + 3bcd^2)e + (3acd^2 + 3bc^2d)f)x^7}{7} + ac^3ex + \frac{((3acd^2 + 3bc^2d)e + (3a^2d + bc^3)f)x^5}{5} + \frac{(ac^3f + (3a^2d + bc^3)e)x^3}{3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e),x)

[Out] $\frac{1}{11}b*d^3*f*x^{11} + \frac{1}{9}*((a*d^3 + 3*b*c*d^2)*f + b*d^3*e)*x^9 + \frac{1}{7}*((3*a*c*d^2 + 3*b*c^2*d)*f + (a*d^3 + 3*b*c*d^2)*e)*x^7 + \frac{1}{5}*((3*a*c^2*d + b*c^3)*f + (3*a*c*d^2 + 3*b*c^2*d)*e)*x^5 + \frac{1}{3}*(a*c^3*f + (3*a*c^2*d + b*c^3)*e)*x^3 + a*c^3*e*x$

maxima [A] time = 0.84, size = 146, normalized size = 1.12

$$\frac{1}{11}bd^3fx^{11} + \frac{1}{9}(bd^3e + (3bcd^2 + ad^3)f)x^9 + \frac{1}{7}((3bcd^2 + ad^3)e + 3(bc^2d + acd^2)f)x^7 + ac^3ex + \frac{1}{5}(3(bc^2d + acd^2)e + (bc^3 + 3ac^2d)f)x^5 + \frac{1}{3}(ac^3f + (bc^3 + 3ac^2d)e)x^3$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3*(f*x^2+e),x, algorithm="maxima")

[Out] $\frac{1}{11}b*d^3*f*x^{11} + \frac{1}{9}*(b*d^3*e + (3*b*c*d^2 + a*d^3)*f)*x^9 + \frac{1}{7}*((3*b*c*d^2 + a*d^3)*e + 3*(b*c^2*d + a*c*d^2)*f)*x^7 + a*c^3*e*x + \frac{1}{5}*(3*(b*c^2*d + a*c*d^2)*e + (b*c^3 + 3*a*c^2*d)*f)*x^5 + \frac{1}{3}*(a*c^3*f + (b*c^3 + 3*a*c^2*d)*e)*x^3$

mupad [B] time = 0.81, size = 143, normalized size = 1.10

$$x^5\left(\frac{bc^3f}{5} + \frac{3acd^2e}{5} + \frac{3ac^2df}{5} + \frac{3bc^2de}{5}\right) + x^7\left(\frac{ad^3e}{7} + \frac{3acd^2f}{7} + \frac{3bc^2de}{7} + \frac{3bc^2df}{7}\right) + x^3\left(\frac{ac^3f}{3} + \frac{bc^3e}{3} + ac^2de\right) + x^9\left(\frac{ad^3f}{9} + \frac{bd^3e}{9} + \frac{bcd^2f}{3}\right) + ac^3ex + \frac{bd^3fx^{11}}{11}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)*(c + d*x^2)^3*(e + f*x^2),x)

[Out] $x^5*((b*c^3*f)/5 + (3*a*c*d^2*e)/5 + (3*a*c^2*d*f)/5 + (3*b*c^2*d*e)/5) + x^7*((a*d^3*e)/7 + (3*a*c*d^2*f)/7 + (3*b*c*d^2*e)/7 + (3*b*c^2*d*f)/7) + x^3*((a*c^3*f)/3 + (b*c^3*e)/3 + a*c^2*d*e) + x^9*((a*d^3*f)/9 + (b*d^3*e)/9 + (b*c*d^2*f)/3) + a*c^3*e*x + (b*d^3*f*x^{11})/11$

sympy [A] time = 0.09, size = 173, normalized size = 1.33

$$ac^3ex + \frac{bd^3fx^{11}}{11} + x^9\left(\frac{ad^3f}{9} + \frac{bcd^2f}{3} + \frac{bd^3e}{9}\right) + x^7\left(\frac{3acd^2f}{7} + \frac{ad^3e}{7} + \frac{3bc^2df}{7} + \frac{3bcd^2e}{7}\right) + x^5\left(\frac{3ac^2df}{5} + \frac{3acd^2e}{5} + \frac{bc^3f}{5} + \frac{3bc^2de}{5}\right) + x^3\left(\frac{ac^3f}{3} + ac^2de + \frac{bc^3e}{3}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)**3*(f*x**2+e),x)
```

```
[Out] a*c**3*e*x + b*d**3*f*x**11/11 + x**9*(a*d**3*f/9 + b*c*d**2*f/3 + b*d**3*e/9) + x**7*(3*a*c*d**2*f/7 + a*d**3*e/7 + 3*b*c**2*d*f/7 + 3*b*c*d**2*e/7) + x**5*(3*a*c**2*d*f/5 + 3*a*c*d**2*e/5 + b*c**3*f/5 + 3*b*c**2*d*e/5) + x**3*(a*c**3*f/3 + a*c**2*d*e + b*c**3*e/3)
```

$$3.19 \quad \int \frac{(a+bx^2)(c+dx^2)^3}{e+fx^2} dx$$

Optimal. Leaf size=227

$$\frac{x(c+dx^2)(7adf(5de-9cf)-b(24c^2f^2-63cdef+35d^2e^2))}{105f^3} + \frac{x(7adf(33c^2f^2-40cdef+15d^2e^2)-b(-48c^3f^3-280cd^2e^2f+105d^3e^3))}{105f^3}$$

Rubi [A] time = 0.37, antiderivative size = 227, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {528, 388, 205}

$$\frac{x(c+dx^2)(7adf(5de-9cf)-b(24c^2f^2-63cdef+35d^2e^2))}{105f^3} + \frac{x(7adf(33c^2f^2-40cdef+15d^2e^2)-b(231c^2def^2-48c^3f^3-280cd^2e^2f+105d^3e^3))}{105f^3} - \frac{x(c+dx^2)^2(-7adf-6bcf+7bde)}{35f^2} + \frac{(bc-af)(de-cf)^3 \tan^{-1}\left(\frac{\sqrt{e}}{\sqrt{f}}\right)}{\sqrt{e}f^{9/2}} + \frac{bx(c+dx^2)^3}{7f}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2), x]

[Out] ((7*a*d*f*(15*d^2*e^2 - 40*c*d*e*f + 33*c^2*f^2) - b*(105*d^3*e^3 - 280*c*d^2*e^2*f + 231*c^2*d*e*f^2 - 48*c^3*f^3))*x)/(105*f^4) - ((7*a*d*f*(5*d*e - 9*c*f) - b*(35*d^2*e^2 - 63*c*d*e*f + 24*c^2*f^2))*x*(c + d*x^2))/(105*f^3) - ((7*b*d*e - 6*b*c*f - 7*a*d*f)*x*(c + d*x^2)^2)/(35*f^2) + (b*x*(c + d*x^2)^3)/(7*f) + ((b*e - a*f)*(d*e - c*f)^3*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(Sqrt[e]*f^(9/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 528

Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p+1)*(c + d*x^n)^q)/(b*(n*(p+q+1)+1)), x] + Dist[1/(b*(n*(p+q+1)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-1)*Simp[c*(b*e - a*f + b*e*n*(p+q+1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p+q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p+q+1)+1, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^3}{e + fx^2} dx &= \frac{bx(c + dx^2)^3}{7f} + \frac{\int \frac{(c+dx^2)^2(-c(be-7af)+(-7bde+6bcf+7adf)x^2)}{e+fx^2} dx}{7f} \\
&= -\frac{(7bde - 6bcf - 7adf)x(c + dx^2)^2}{35f^2} + \frac{bx(c + dx^2)^3}{7f} + \frac{\int \frac{(c+dx^2)(c(be(7de-11cf)-7af)(c+dx^2)+(-7bde+6bcf+7adf)x^2)}{e+fx^2} dx}{7f} \\
&= -\frac{(7adf(5de - 9cf) - b(35d^2e^2 - 63cdef + 24c^2f^2))x(c + dx^2)}{105f^3} - \frac{(7bde - 6bcf - 7adf)(c + dx^2)^2}{105f^3} \\
&= \frac{(7adf(15d^2e^2 - 40cdef + 33c^2f^2) - b(105d^3e^3 - 280cd^2e^2f + 231c^2def^2 - 48cd^2ef^2 + 24c^2f^3))x(c + dx^2)}{105f^4} - \frac{(7bde - 6bcf - 7adf)(c + dx^2)^2}{105f^3} \\
&= \frac{(7adf(15d^2e^2 - 40cdef + 33c^2f^2) - b(105d^3e^3 - 280cd^2e^2f + 231c^2def^2 - 48cd^2ef^2 + 24c^2f^3))x(c + dx^2)}{105f^4} - \frac{(7bde - 6bcf - 7adf)(c + dx^2)^2}{105f^3}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 179, normalized size = 0.79

$$\frac{x(adf(3c^2f^2 - 3cdef + d^2e^2) - b(de - cf)^3)}{f^4} + \frac{dx^3(adf(3cf - de) + b(3c^2f^2 - 3cdef + d^2e^2))}{3f^3} + \frac{d^2x^5(adf + 3bcf - bde)}{5f^2} + \frac{(be - af)(de - cf)^3 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)}{\sqrt{e}f^{9/2}} + \frac{bd^3x^7}{7f}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2), x]

[Out] ((-(b*(d*e - c*f)^3) + a*d*f*(d^2*e^2 - 3*c*d*e*f + 3*c^2*f^2))*x)/f^4 + (d*(a*d*f*(-(d*e) + 3*c*f) + b*(d^2*e^2 - 3*c*d*e*f + 3*c^2*f^2))*x^3)/(3*f^3) + (d^2*(-(b*d*e) + 3*b*c*f + a*d*f)*x^5)/(5*f^2) + (b*d^3*x^7)/(7*f) + ((b*e - a*f)*(d*e - c*f)^3*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(Sqrt[e]*f^(9/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^3}{e + fx^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2), x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2), x]

fricas [A] time = 1.00, size = 586, normalized size = 2.58

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e), x, algorithm="fricas")

[Out] [1/210*(30*b*d^3*e*f^4*x^7 - 42*(b*d^3*e^2*f^3 - (3*b*c*d^2 + a*d^3)*e*f^4)*x^5 + 70*(b*d^3*e^3*f^2 - (3*b*c*d^2 + a*d^3)*e^2*f^3 + 3*(b*c^2*d + a*c*d^2)*e*f^4)*x^3 - 105*(b*d^3*e^4 + a*c^3*f^4 - (3*b*c*d^2 + a*d^3)*e^3*f + 3*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 210*(b*d^3*e^4*f - (3*b*c*d^2 + a*d^3)*e^3*f^2 + 3*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x)/(e*f^5), 1/105*(15*b*d^3*e*f^4*x^7 - 21*(b*d^3*e^2*f^3 - (3*b*c*d^2 + a*d^3)*e*f^4)*x^5 + 35*(b*d^3*e^3*f^2 - (3*b*c*d^2 + a*d^3)*e^2*f^3 + 3*(b*c^2*d + a*c*d^2)*e*f^4)*x^3 - 105*(b*d^3*e^4 + a*c^3*f^4 - (3*b*c*d^2 + a*d^3)*e^3*f + 3*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 210*(b*d^3*e^4*f - (3*b*c*d^2 + a*d^3)*e^3*f^2 + 3*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x)/(e*f^5)]

$$2*d + a*c*d^2)*e*f^4)*x^3 + 105*(b*d^3*e^4 + a*c^3*f^4 - (3*b*c*d^2 + a*d^3)*e^3*f + 3*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*\text{sqrt}(e*f)*\text{arctan}(\text{sqrt}(e*f)*x/e) - 105*(b*d^3*e^4*f - (3*b*c*d^2 + a*d^3)*e^3*f^2 + 3*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x)/(e*f^5)]$$

giac [A] time = 0.49, size = 307, normalized size = 1.35

$$\frac{(a^2 f^4 - b^2 f^4 - 3 a c^2 d^2 f^2 + 3 b c^2 d^2 f^2 - 3 b c^2 d^2 f^2 - a d^3 f^2 + b d^3 f^2) \arctan\left(\sqrt{f} \frac{x}{e}\right) + \frac{15 b d^3 f^2 + 63 b c d^3 f^2 + 21 a d^3 f^2 - 21 b d^3 f^2 + 105 b c^2 d^2 f^2 + 105 a c^2 d^2 f^2 - 105 b c^2 d^2 f^2 - 35 a d^3 f^2 + 35 b d^3 f^2 + 105 b c^2 d^2 f^2 + 105 a c^2 d^2 f^2 - 315 b c^2 d^2 f^2 - 315 a c^2 d^2 f^2 + 105 a d^3 f^2 - 105 b d^3 f^2}{105 f^5}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e),x, algorithm="giac")

[Out] (a*c^3*f^4 - b*c^3*f^3*e - 3*a*c^2*d*f^3*e + 3*b*c^2*d*f^2*e^2 + 3*a*c*d^2*f^2*e^2 - 3*b*c*d^2*f*e^3 - a*d^3*f*e^3 + b*d^3*e^4)*arctan(sqrt(f)*x*e^(-1/2))/f^(9/2) + 1/105*(15*b*d^3*f^6*x^7 + 63*b*c*d^2*f^6*x^5 + 21*a*d^3*f^6*x^5 - 21*b*d^3*f^5*x^5*e + 105*b*c^2*d*f^6*x^3 + 105*a*c*d^2*f^6*x^3 - 105*b*c*d^2*f^5*x^3*e - 35*a*d^3*f^5*x^3*e + 35*b*d^3*f^4*x^3*e^2 + 105*b*c^3*f^6*x + 315*a*c^2*d*f^6*x - 315*b*c^2*d*f^5*x*e - 315*a*c*d^2*f^5*x*e + 315*b*c*d^2*f^4*x*e^2 + 105*a*d^3*f^4*x*e^2 - 105*b*d^3*f^3*x*e^3)/f^7

maple [A] time = 0.01, size = 401, normalized size = 1.77

$$\frac{b d^3 x^2 + a d^3 x^2 - 3 b c d^2 x^2 + 3 b c d^2 x^2 - 3 b c d^2 x^2 - a d^3 x^2 + b d^3 x^2}{27 f^2} + \frac{3 b c d^2 x^2}{57 f^2} + \frac{b d^3 x^2}{57 f^2} + \frac{a d^3 x^2}{57 f^2} + \frac{a d^3 x^2}{57 f^2} + \frac{b c^2 d x^2}{3 f^2} + \frac{b c^2 d x^2}{3 f^2} + \frac{b c^2 d x^2}{3 f^2} + \frac{b c^2 d x^2}{3 f^2} + \frac{a^2 \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{3 a^2 d \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{3 a c^2 d^2 \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{a d^3 \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{b c^2 d \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{3 b c^2 d^2 \arctan\left(\frac{x}{\sqrt{f}}\right)}{\sqrt{f}} + \frac{3 a c^2 d x}{f} + \frac{3 a c^2 d x}{f} + \frac{a d^3 x}{f} + \frac{b c^2 x}{f} + \frac{3 b c^2 d x}{f} + \frac{3 b c^2 d x}{f} + \frac{3 b c^2 d x}{f} + \frac{3 b c^2 d x}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e),x)

[Out] 1/7/f*b*d^3*x^7+1/5/f*x^5*a*d^3+3/5/f*x^5*b*c*d^2-1/5/f^2*x^5*b*d^3*e+1/f*x^3*a*c*d^2-1/3/f^2*x^3*a*d^3*e+1/f*x^3*b*c^2*d-1/f^2*x^3*b*c*d^2*e+1/3/f^3*x^3*b*d^3*e^2+3/f*a*c^2*d*x-3/f^2*a*c*d^2*e*x+1/f^3*a*d^3*e^2*x+1/f*b*c^3*x-3/f^2*b*c^2*d*e*x+3/f^3*b*c*d^2*e^2*x-1/f^4*b*d^3*e^3*x+1/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^3-3/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2*d*e+3/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d^2*e^2-1/f^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^3*e^3-1/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^3*e+3/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2*d*e^2-3/f^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d^2*e^3+1/f^4/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^3*e^4

maxima [A] time = 2.03, size = 266, normalized size = 1.17

$$\frac{(b^2 d^4 + a^2 f^4 - (3 b c^2 d + a d^3) f^2 + 3 (b c^2 d + a c d^2) f^2 - (b c^3 + 3 a c^2 d) e f^2) \arctan\left(\frac{f x}{\sqrt{e}}\right) + \frac{15 b d^3 f^2 x^7 - 21 (b d^3 e f^2 - (3 b c d^2 + a d^3) f^2) x^5 + 35 (b d^3 e f^2 - (3 b c d^2 + a d^3) f^2) + 3 (b c^2 d + a c d^2) f^2 x^3 - 105 (b d^3 e^3 - (3 b c^2 d + a d^3) e^2 f + 3 (b c^2 d + a c d^2) e f^2 - (b c^3 + 3 a c^2 d) f^2) x}{105 f^4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e),x, algorithm="maxima")

[Out] (b*d^3*e^4 + a*c^3*f^4 - (3*b*c*d^2 + a*d^3)*e^3*f + 3*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*arctan(f*x/sqrt(e*f))/(sqrt(e*f)*f^4) + 1/105*(15*b*d^3*f^3*x^7 - 21*(b*d^3*e*f^2 - (3*b*c*d^2 + a*d^3)*f^3)*x^5 + 35*(b*d^3*e^2*f - (3*b*c*d^2 + a*d^3)*e*f^2 + 3*(b*c^2*d + a*c*d^2)*f^3)*x^3 - 105*(b*d^3*e^3 - (3*b*c*d^2 + a*d^3)*e^2*f + 3*(b*c^2*d + a*c*d^2)*e*f^2 - (b*c^3 + 3*a*c^2*d)*f^3)*x)/f^4

mupad [B] time = 0.85, size = 312, normalized size = 1.37

$$x \left(\frac{b c^2 + 3 a d e^2}{f} + \frac{e \left(\frac{a d^2 + 3 b c d^2}{f} - \frac{b d^3 e}{f^2} \right) - \frac{3 c d (a d + b c)}{f}}{f} \right) + x^5 \left(\frac{a d^3 + 3 b c d^2}{5 f} - \frac{b d^3 e}{5 f^2} \right) - x^3 \left(\frac{e \left(\frac{a d^2 + 3 b c d^2}{f} - \frac{b d^3 e}{f^2} \right) - c d (a d + b c)}{3 f} \right) + \frac{b d^3 x^7}{7 f} + \frac{\text{atan}\left(\frac{\sqrt{f} x (e f - b) (c f - d e)^3}{\sqrt{e} (-b c^2 e f^2 + a c^2 f^2 + 3 b c^2 d e f^2 - 3 a c^2 d e f^2 - 3 b c^2 d e f^2 + 3 a c d^2 e^2 f^2 + b d^3 e^2 - a d^3 e^2 f)}\right) (a f - b e) (c f - d e)^3}{\sqrt{e} f^{9/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2),x)

[Out] $x \cdot \frac{(b \cdot c^3 + 3 \cdot a \cdot c^2 \cdot d) / f + (e \cdot ((e \cdot ((a \cdot d^3 + 3 \cdot b \cdot c \cdot d^2) / f - (b \cdot d^3 \cdot e) / f^2)) / f - (3 \cdot c \cdot d \cdot (a \cdot d + b \cdot c)) / f)) / f}{f} + x^5 \cdot \frac{(a \cdot d^3 + 3 \cdot b \cdot c \cdot d^2) / (5 \cdot f) - (b \cdot d^3 \cdot e) / (5 \cdot f^2)}{f^2} - x^3 \cdot \frac{(e \cdot ((a \cdot d^3 + 3 \cdot b \cdot c \cdot d^2) / f - (b \cdot d^3 \cdot e) / f^2)) / (3 \cdot f) - (c \cdot d \cdot (a \cdot d + b \cdot c)) / f + (b \cdot d^3 \cdot x^7) / (7 \cdot f) + (\operatorname{atan}((f^{1/2} \cdot x \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3) / (e^{1/2} \cdot (a \cdot c^3 \cdot f^4 + b \cdot d^3 \cdot e^4 - a \cdot d^3 \cdot e^3 \cdot f - b \cdot c^3 \cdot e \cdot f^3 - 3 \cdot a \cdot c^2 \cdot d \cdot e \cdot f^3 - 3 \cdot b \cdot c \cdot d^2 \cdot e^3 \cdot f + 3 \cdot a \cdot c \cdot d^2 \cdot e^2 \cdot f^2 + 3 \cdot b \cdot c^2 \cdot d \cdot e^2 \cdot f^2))) \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3}{e^{1/2} \cdot f^{9/2}}$

sympy [B] time = 1.49, size = 508, normalized size = 2.24

$$\frac{b^3 d^2}{2f} + x^5 \left(\frac{a^3}{5f} + \frac{3acd^2}{5f} + \frac{bd^3e}{5f^2} \right) + x^3 \left(\frac{acd}{f} - \frac{ad^3e}{3f^2} + \frac{b^2d}{f} - \frac{bcd^2e}{f^2} + \frac{bd^3e}{3f^3} \right) + x \left(\frac{3ad^2d}{f} - \frac{3acd^2e}{f^2} + \frac{ad^3e}{f^2} + \frac{b^2}{f} - \frac{3bc^2de}{f^2} + \frac{3acd^2e}{f^3} - \frac{bd^3e}{f^4} \right) \cdot \frac{\sqrt{\frac{1}{f^9} (ef - be)(cf - de)^3 \log\left(\frac{e^3 \sqrt{\frac{1}{2f} (ef - be)(cf - de)^3}}{2ef^3 - 3ac^2d^2f^2 - 3acd^2e^2f - 2d^3e^3f^2 - 3bc^2d^2f - 3cd^2e^2f - 3ad^2e^3f + x}\right)}}{2} + \frac{\sqrt{\frac{1}{f^9} (ef - be)(cf - de)^3 \log\left(\frac{e^3 \sqrt{\frac{1}{2f} (ef - be)(cf - de)^3}}{2ef^3 - 3ac^2d^2f^2 - 3acd^2e^2f - 2d^3e^3f^2 - 3bc^2d^2f - 3cd^2e^2f - 3ad^2e^3f + x}\right)}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3/(f*x**2+e),x)

[Out] $b \cdot d^3 \cdot x^7 / (7 \cdot f) + x^5 \cdot (a \cdot d^3 / (5 \cdot f) + 3 \cdot b \cdot c \cdot d^2 / (5 \cdot f) - b \cdot d^3 \cdot e / (5 \cdot f^2)) + x^3 \cdot (a \cdot c \cdot d^2 / f - a \cdot d^3 \cdot e / (3 \cdot f^2) + b \cdot c \cdot d^2 / f - b \cdot c \cdot d^2 \cdot e / f^2 + b \cdot d^3 \cdot e^2 / (3 \cdot f^3)) + x \cdot (3 \cdot a \cdot c \cdot d^2 / f - 3 \cdot a \cdot c \cdot d^2 \cdot e / f^2 + a \cdot d^3 \cdot e^2 / f^3 + b \cdot c \cdot d^2 / f - 3 \cdot b \cdot c \cdot d^2 \cdot e / f^2 + 3 \cdot b \cdot c \cdot d^2 \cdot e^2 / f^3 - b \cdot d^3 \cdot e^3 / f^4) - \sqrt{-1/(e \cdot f^9)} \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3 \cdot \log(-e \cdot f^4 \cdot \sqrt{-1/(e \cdot f^9)}) \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3 / (a \cdot c^3 \cdot f^4 - 3 \cdot a \cdot c^2 \cdot d \cdot e \cdot f^3 + 3 \cdot a \cdot c \cdot d^2 \cdot e^2 \cdot f^2 - a \cdot d^3 \cdot e^3 \cdot f - b \cdot c^3 \cdot e \cdot f^3 + 3 \cdot b \cdot c^2 \cdot d \cdot e^2 \cdot f^2 - 3 \cdot b \cdot c \cdot d^2 \cdot e^3 \cdot f + b \cdot d^3 \cdot e^4) + x) / 2 + \sqrt{-1/(e \cdot f^9)} \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3 \cdot \log(e \cdot f^4 \cdot \sqrt{-1/(e \cdot f^9)}) \cdot (a \cdot f - b \cdot e) \cdot (c \cdot f - d \cdot e)^3 / (a \cdot c^3 \cdot f^4 - 3 \cdot a \cdot c^2 \cdot d \cdot e \cdot f^3 + 3 \cdot a \cdot c \cdot d^2 \cdot e^2 \cdot f^2 - a \cdot d^3 \cdot e^3 \cdot f - b \cdot c^3 \cdot e \cdot f^3 + 3 \cdot b \cdot c^2 \cdot d \cdot e^2 \cdot f^2 - 3 \cdot b \cdot c \cdot d^2 \cdot e^3 \cdot f + b \cdot d^3 \cdot e^4) + x) / 2$

$$3.20 \quad \int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^2} dx$$

Optimal. Leaf size=242

$$\frac{dx(5af(3c^2f^2 - 22cdef + 15d^2e^2) - be(81c^2f^2 - 190cdef + 105d^2e^2))}{30ef^4} - \frac{(de - cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(7de - cf) - cf)}{2e^{3/2}f^{9/2}}$$

Rubi [A] time = 0.40, antiderivative size = 242, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {526, 528, 388, 205}

$$\frac{dx(5af(3c^2f^2 - 22cdef + 15d^2e^2) - be(81c^2f^2 - 190cdef + 105d^2e^2))}{30ef^4} - \frac{(de - cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(7de - cf) - cf)}{2e^{3/2}f^{9/2}} + \frac{dx(c + dx^2)^2(7be - 5af)}{10ef^2} - \frac{dx(c + dx^2)(be(35de - 33cf) - 5af(5de - 3cf))}{30ef^3} - \frac{x(c + dx^2)^3(be - af)}{2ef(e + fx^2)}$$

Antiderivative was successfully verified.

[In] Int[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^2,x]

[Out] -(d*(5*a*f*(15*d^2*e^2 - 22*c*d*e*f + 3*c^2*f^2) - b*e*(105*d^2*e^2 - 190*c*d*e*f + 81*c^2*f^2))*x)/(30*e*f^4) - (d*(b*e*(35*d*e - 33*c*f) - 5*a*f*(5*d*e - 3*c*f))*x*(c + d*x^2))/(30*e*f^3) + (d*(7*b*e - 5*a*f)*x*(c + d*x^2)^2)/(10*e*f^2) - ((b*e - a*f)*x*(c + d*x^2)^3)/(2*e*f*(e + f*x^2)) - ((d*e - c*f)^2*(b*e*(7*d*e - c*f) - a*f*(5*d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]]/(2*e^(3/2)*f^(9/2))

Rule 205

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 388

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 526

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] :> -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]

Rule 528

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] :> Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^2} dx &= \frac{(be - af)x(c + dx^2)^3}{2ef(e + fx^2)} - \frac{\int \frac{(c+dx^2)^2(-c(be+af)-d(7be-5af)x^2)}{e+fx^2} dx}{2ef} \\
&= \frac{d(7be - 5af)x(c + dx^2)^2}{10ef^2} - \frac{(be - af)x(c + dx^2)^3}{2ef(e + fx^2)} - \frac{\int \frac{(c+dx^2)(c(be(7de-5cf)-5af(de+c))}{e+fx^2} dx}{2ef} \\
&= -\frac{d(be(35de - 33cf) - 5af(5de - 3cf))x(c + dx^2)}{30ef^3} + \frac{d(7be - 5af)x(c + dx^2)^2}{10ef^2} - \frac{\int \frac{(c+dx^2)(c(be(7de-5cf)-5af(de+c))}{e+fx^2} dx}{2ef} \\
&= -\frac{d(5af(15d^2e^2 - 22cdef + 3c^2f^2) - be(105d^2e^2 - 190cdef + 81c^2f^2))x}{30ef^4} - \frac{d(be(35de - 33cf) - 5af(5de - 3cf))x(c + dx^2)}{30ef^3} \\
&= -\frac{d(5af(15d^2e^2 - 22cdef + 3c^2f^2) - be(105d^2e^2 - 190cdef + 81c^2f^2))x}{30ef^4} - \frac{d(be(35de - 33cf) - 5af(5de - 3cf))x(c + dx^2)}{30ef^3}
\end{aligned}$$

Mathematica [A] time = 0.13, size = 176, normalized size = 0.73

$$\frac{d^2x^3(adf + 3bcf - 2bde)}{3f^3} - \frac{(de - cf)^2 \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right)(be(7de - cf) - af(cf + 5de))}{2e^{3/2}f^{9/2}} + \frac{x(be - af)(de - cf)^3}{2ef^4(e + fx^2)} + \frac{dx(adf(3cf - 2de) + 3b(de - cf)^2)}{f^4} + \frac{bd^3x^5}{5f^2}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^2,x]

[Out] (d*(3*b*(d*e - c*f)^2 + a*d*f*(-2*d*e + 3*c*f))*x)/f^4 + (d^2*(-2*b*d*e + 3*b*c*f + a*d*f)*x^3)/(3*f^3) + (b*d^3*x^5)/(5*f^2) + ((b*e - a*f)*(d*e - c*f)^3*x)/(2*e*f^4*(e + f*x^2)) - ((d*e - c*f)^2*(b*e*(7*d*e - c*f) - a*f*(5*d*e + c*f))*ArcTan[(Sqrt[f]*x)/Sqrt[e]]/(2*e^(3/2)*f^(9/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^2} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^2,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^2, x]

fricas [A] time = 1.24, size = 834, normalized size = 3.45

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^2,x, algorithm="fricas")

[Out] [1/60*(12*b*d^3*e^2*f^4*x^7 - 4*(7*b*d^3*e^3*f^3 - 5*(3*b*c*d^2 + a*d^3))*e^2*f^4)*x^5 + 20*(7*b*d^3*e^4*f^2 - 5*(3*b*c*d^2 + a*d^3))*e^3*f^3 + 9*(b*c^2*d + a*c*d^2)*e^2*f^4)*x^3 + 15*(7*b*d^3*e^5 - a*c^3*e*f^4 - 5*(3*b*c*d^2 + a*d^3))*e^4*f + 9*(b*c^2*d + a*c*d^2)*e^3*f^2 - (b*c^3 + 3*a*c^2*d)*e^2*f^3 + (7*b*d^3*e^4*f - a*c^3*f^5 - 5*(3*b*c*d^2 + a*d^3))*e^3*f^2 + 9*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x^2)*sqrt(-e*f)*log((f*x^2

- 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) + 30*(7*b*d^3*e^5*f + a*c^3*e*f^5 - 5*(3*b*c*d^2 + a*d^3)*e^4*f^2 + 9*(b*c^2*d + a*c*d^2)*e^3*f^3 - (b*c^3 + 3*a*c^2*d)*e^2*f^4)*x)/(e^2*f^6*x^2 + e^3*f^5), 1/30*(6*b*d^3*e^2*f^4*x^7 - 2*(7*b*d^3*e^3*f^3 - 5*(3*b*c*d^2 + a*d^3)*e^2*f^4)*x^5 + 10*(7*b*d^3*e^4*f^2 - 5*(3*b*c*d^2 + a*d^3)*e^3*f^3 + 9*(b*c^2*d + a*c*d^2)*e^2*f^4)*x^3 - 15*(7*b*d^3*e^5 - a*c^3*e*f^4 - 5*(3*b*c*d^2 + a*d^3)*e^4*f + 9*(b*c^2*d + a*c*d^2)*e^3*f^2 - (b*c^3 + 3*a*c^2*d)*e^2*f^3 + (7*b*d^3*e^4*f - a*c^3*f^5 - 5*(3*b*c*d^2 + a*d^3)*e^3*f^2 + 9*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) + 15*(7*b*d^3*e^5*f + a*c^3*e*f^5 - 5*(3*b*c*d^2 + a*d^3)*e^4*f^2 + 9*(b*c^2*d + a*c*d^2)*e^3*f^3 - (b*c^3 + 3*a*c^2*d)*e^2*f^4)*x)/(e^2*f^6*x^2 + e^3*f^5)]

giac [A] time = 0.37, size = 321, normalized size = 1.33

$$\frac{(a^2 f^4 + b^2 f^4 + 3 a^2 d f^2 - 9 a d^2 f^2 + 15 b d^2 f^2 + 5 a d^3 f^2 - 7 b d^3 f^2) \arctan\left(\sqrt{f x^2 + e}\right) f^{\frac{3}{2}}}{2 f^{\frac{3}{2}}} + \frac{(a^2 f^4 x - b^2 f^4 x - 3 a^2 d f^2 x + 3 b^2 d f^2 x + 3 a d^3 f^2 x - 3 b d^3 f^2 x - a d^4 f^2 x + b d^4 f^2 x^{-1})}{2(f^2 + e)^{\frac{3}{2}}} + \frac{3 b d^3 f^2 + 15 b a d^2 f^2 + 5 a d^3 f^2 - 10 b d^3 f^2 x + 45 b^2 d f^2 x + 45 a d^3 f^2 x - 90 a d^3 f^2 x - 30 a d^3 f^2 x + 45 b d^3 f^2 x^2}{15 f^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^2,x, algorithm="giac")

[Out] 1/2*(a*c^3*f^4 + b*c^3*f^3*e + 3*a*c^2*d*f^3*e - 9*b*c^2*d*f^2*e^2 - 9*a*c*d^2*f^2*e^2 + 15*b*c*d^2*f*e^3 + 5*a*d^3*f*e^3 - 7*b*d^3*e^4)*arctan(sqrt(f)*x*e^(-1/2))*e^(-3/2)/f^(9/2) + 1/2*(a*c^3*f^4*x - b*c^3*f^3*x*e - 3*a*c^2*d*f^3*x*e + 3*b*c^2*d*f^2*x*e^2 + 3*a*c*d^2*f^2*x*e^2 - 3*b*c*d^2*f*x*e^3 - a*d^3*f*x*e^3 + b*d^3*x*e^4)*e^(-1)/((f*x^2 + e)*f^4) + 1/15*(3*b*d^3*f^8*x^5 + 15*b*c*d^2*f^8*x^3 + 5*a*d^3*f^8*x^3 - 10*b*d^3*f^7*x^3*e + 45*b*c*d^2*f^8*x + 45*a*c*d^2*f^8*x - 90*b*c*d^2*f^7*x*e - 30*a*d^3*f^7*x*e + 45*b*d^3*f^6*x*e^2)/f^10

maple [B] time = 0.02, size = 475, normalized size = 1.96

$$\frac{b^2 d^3}{3 f^2} + \frac{a d^3}{3 f^2} + \frac{b c d^2}{3 f^2} + \frac{2 a d^2 c}{3 f^2} + \frac{a^2 c}{2(f^2 + e)} + \frac{a^2 d \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{3 a c d x}{2(f^2 + e) f} + \frac{3 a c d \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{3 a c d c x}{2(f^2 + e) f^2} + \frac{9 a c d c \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{a d^2 c x}{2(f^2 + e) f^2} + \frac{5 a d^2 c \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{b^2 c}{2(f^2 + e)} + \frac{b^2 d \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{3 b^2 d x}{2(f^2 + e) f^2} + \frac{9 b^2 d c \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{3 b c d c x}{2(f^2 + e) f^2} + \frac{15 b c d c \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{b d^2 c x}{2(f^2 + e) f^2} + \frac{7 b d^2 c \arctan\left(\frac{a x}{\sqrt{f x^2 + e}}\right)}{2 \sqrt{f} e} + \frac{3 a d^2 c}{f^2} + \frac{2 a d^2 c x}{f^2} + \frac{3 b c d c x}{f^2} + \frac{6 b c d c x}{f^2} + \frac{3 b d^2 c x}{f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^2,x)

[Out] 3*d^2/f^2*a*c*x+d^2/f^2*x^3*b*c+3/2/f^2*e*x/(f*x^2+e)*b*c^2*d-3/2/f^3*e^2*x/(f*x^2+e)*b*c*d^2-9/2/f^2*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d^2-9/2/f^2*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2*d+15/2/f^3*e^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d^2+3/2/f^2*e*x/(f*x^2+e)*a*c*d^2-1/2/f^3*e^2*x/(f*x^2+e)*a*d^3+1/2/f^4*e^3*x/(f*x^2+e)*b*d^3+3/2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2*d+5/2/f^3*e^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^3-7/2/f^4*e^3/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^3-6*d^2/f^3*b*c*e*x-3/2/f*x/(f*x^2+e)*a*c^2*d+1/5*d^3/f^2*b*x^5+1/3*d^3/f^2*x^3*a-2/3*d^3/f^3*x^3*b*e+1/2/e*x/(f*x^2+e)*a*c^3+1/2/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^3+1/2/f/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^3+3*d/f^2*b*c^2*x+3*d^3/f^4*b*e^2*x-1/2/f*x/(f*x^2+e)*b*c^3-2*d^3/f^3*a*e*x

maxima [A] time = 1.98, size = 296, normalized size = 1.22

$$\frac{(b^2 d^4 + a^2 f^4 - (3 b c d^2 + a d^3) e^2 f + 3 (b c^2 d + a c d^2) e^2 f^2 - (b c^3 + 3 a c^2 d) e f^3) x}{2(e f^2 x^2 + e^2 f^4)} + \frac{3 b d^3 f^2 x^5 - 5(2 b d^3 e f - (3 b c d^2 + a d^3) f^2) x^3 + 15(3 b c d^2 - 2(3 b c d^2 + a d^3) e f + 3(b c^2 d + a c d^2) f^2) x}{15 f^4} - \frac{(7 b d^3 e^4 - a^2 f^4 - 5(3 b c d^2 + a d^3) e^2 f^2 + 9(b c^2 d + a c d^2) e^2 f^2 - (b c^3 + 3 a c^2 d) e f^3) \arctan\left(\frac{f x}{\sqrt{e f^2}}\right)}{2 \sqrt{e f^4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^2,x, algorithm="maxima")

[Out] 1/2*(b*d^3*e^4 + a*c^3*f^4 - (3*b*c*d^2 + a*d^3)*e^3*f + 3*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*x/(e*f^5*x^2 + e^2*f^4) + 1/15*(3*b*d^3*f^2*x^5 - 5*(2*b*d^3*e*f - (3*b*c*d^2 + a*d^3)*f^2)*x^3 + 15*(3*b*d^3*e^2 - 2*(3*b*c*d^2 + a*d^3)*e*f + 3*(b*c^2*d + a*c*d^2)*f^2)*x)/f^4 - 1/2*(7*b*d^3*e^4 - a*c^3*f^4 - 5*(3*b*c*d^2 + a*d^3)*e^3*f + 9*(b*c^2*d + a*c*d^2)*e^2*f^2 - (b*c^3 + 3*a*c^2*d)*e*f^3)*x/(e*f^5*x^2 + e^2*f^4)

$\wedge 2) * e^2 * f^2 - (b * c^3 + 3 * a * c^2 * d) * e * f^3) * \arctan(f * x / \sqrt{e * f}) / (\sqrt{e * f}) * e * f^4)$

mupad [B] time = 0.98, size = 389, normalized size = 1.61

$$x^2 \left(\frac{a d^3 + 3 b c d^2 - 2 b d^3 e}{3 f^2} - x \left(\frac{2 c \left(\frac{c^2 - 3 a d^2}{f^2} - \frac{2 d^2}{f^2} \right)}{f} + \frac{b d^3 e}{f^4} - \frac{3 c d (a d + b c)}{f^2} \right) + \frac{b d^3 e^2}{5 f^2} + \frac{x (-b c^2 e f^3 + a c^2 f^4 + 3 b c^2 d e^2 f^2 - 3 a c^2 d e f^3 - 3 b c d^2 e^2 f + 3 a c d^2 e^2 f^2 + b d^3 e^3 - a d^3 e^2 f)}{2 e (f^2 + e f)} \right) + \frac{\operatorname{atan}\left(\frac{\sqrt{f-d} \sqrt{e f^2 + 5 a d e f + b c e f}}{\sqrt{b c^2 f^2 + a c^2 f^2 + 3 b c^2 d e f^2 + 3 a c^2 d e f^3 + 3 b c d^2 e^2 f^2 + 3 a c d^2 e^2 f^3 + 3 b d^3 e^3 - a d^3 e^2 f}}\right)}{2 e^{3/2} f^{3/2}} (e f - d e)^2 (a e f^2 - 7 b d e^2 + 5 a d e f + b c e f)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^2,x)

[Out] $x^3 * ((a * d^3 + 3 * b * c * d^2) / (3 * f^2) - (2 * b * d^3 * e) / (3 * f^3)) - x * ((2 * e * ((a * d^3 + 3 * b * c * d^2) / f^2 - (2 * b * d^3 * e) / f^3)) / f + (b * d^3 * e^2) / f^4 - (3 * c * d * (a * d + b * c)) / f^2) + (b * d^3 * x^5) / (5 * f^2) + (x * (a * c^3 * f^4 + b * d^3 * e^4 - a * d^3 * e^3 * f - b * c^3 * e * f^3 - 3 * a * c^2 * d * e * f^3 - 3 * b * c * d^2 * e^3 * f + 3 * a * c * d^2 * e^2 * f^2 + 3 * b * c^2 * d * e^2 * f^2)) / (2 * e * (e * f^4 + f^5 * x^2)) + (\operatorname{atan}((f^{1/2}) * x * (c * f - d * e)^2 * (a * c * f^2 - 7 * b * d * e^2 + 5 * a * d * e * f + b * c * e * f)) / (e^{1/2} * (a * c^3 * f^4 - 7 * b * d^3 * e^4 + 5 * a * d^3 * e^3 * f + b * c^3 * e * f^3 + 3 * a * c^2 * d * e * f^3 + 15 * b * c * d^2 * e^3 * f - 9 * a * c * d^2 * e^2 * f^2 - 9 * b * c^2 * d * e^2 * f^2))) * (c * f - d * e)^2 * (a * c * f^2 - 7 * b * d * e^2 + 5 * a * d * e * f + b * c * e * f)) / (2 * e^{3/2} * f^{9/2})$

sympy [B] time = 4.75, size = 661, normalized size = 2.73

$$\frac{b^2 d^2}{5 f^2} + x \left(\frac{b d^3}{f^2} + \frac{b c^2}{f^2} \right) + \left(\frac{2 a c^2}{3 f^2} + \frac{2 b c^2}{f^2} + \frac{2 b d^2}{f^2} + \frac{2 b c^2}{f^2} \right) + \frac{(a^2 f^2 - 3 a^2 b d f^2 + 3 a b^2 c f^2 - a d^2 f^2 - b c^2 f^2 + 3 b c^2 d f^2 - 3 a b c^2 f + b d^3 f^2)}{2 e f^2 + 2 e^2 f} + \frac{\sqrt{e f} (e f - d e) (a c f^2 + b c f - 7 b d e^2) \log\left(\frac{f^2 \sqrt{e f} (e f^2 + 5 a d e f + b c e f - 7 b d e^2)}{2 e^{3/2} f^{3/2} (e f^2 + 5 a d e f + b c e f - 7 b d e^2)}\right)}{4} + \frac{\sqrt{e f} (e f - d e) (a c f^2 + b c f - 7 b d e^2) \log\left(\frac{f^2 \sqrt{e f} (e f^2 + 5 a d e f + b c e f - 7 b d e^2)}{2 e^{3/2} f^{3/2} (e f^2 + 5 a d e f + b c e f - 7 b d e^2)}\right)}{4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3/(f*x**2+e)**2,x)

[Out] $b * d ** 3 * x ** 5 / (5 * f ** 2) + x ** 3 * (a * d ** 3 / (3 * f ** 2) + b * c * d ** 2 / f ** 2 - 2 * b * d ** 3 * e / (3 * f ** 3)) + x * (3 * a * c * d ** 2 / f ** 2 - 2 * a * d ** 3 * e / f ** 3 + 3 * b * c ** 2 * d / f ** 2 - 6 * b * c * d ** 2 * e / f ** 3 + 3 * b * d ** 3 * e ** 2 / f ** 4) + x * (a * c ** 3 * f ** 4 - 3 * a * c ** 2 * d * e * f ** 3 + 3 * a * c * d ** 2 * e ** 2 * f ** 2 - a * d ** 3 * e ** 3 * f - b * c ** 3 * e * f ** 3 + 3 * b * c ** 2 * d * e ** 2 * f ** 2 - 3 * b * c * d ** 2 * e ** 3 * f + b * d ** 3 * e ** 4) / (2 * e ** 2 * f ** 4 + 2 * e * f ** 5 * x ** 2) - \operatorname{sqrt}(-1 / (e ** 3 * f ** 9)) * (c * f - d * e) ** 2 * (a * c * f ** 2 + 5 * a * d * e * f + b * c * e * f - 7 * b * d * e ** 2) * \log(-e ** 2 * f ** 4 * \operatorname{sqrt}(-1 / (e ** 3 * f ** 9)) * (c * f - d * e) ** 2 * (a * c * f ** 2 + 5 * a * d * e * f + b * c * e * f - 7 * b * d * e ** 2)) / (a * c ** 3 * f ** 4 + 3 * a * c ** 2 * d * e * f ** 3 - 9 * a * c * d ** 2 * e ** 2 * f ** 2 + 5 * a * d ** 3 * e ** 3 * f + b * c ** 3 * e * f ** 3 - 9 * b * c ** 2 * d * e ** 2 * f ** 2 + 15 * b * c * d ** 2 * e ** 3 * f - 7 * b * d ** 3 * e ** 4) + x) / 4 + \operatorname{sqrt}(-1 / (e ** 3 * f ** 9)) * (c * f - d * e) ** 2 * (a * c * f ** 2 + 5 * a * d * e * f + b * c * e * f - 7 * b * d * e ** 2) * \log(e ** 2 * f ** 4 * \operatorname{sqrt}(-1 / (e ** 3 * f ** 9)) * (c * f - d * e) ** 2 * (a * c * f ** 2 + 5 * a * d * e * f + b * c * e * f - 7 * b * d * e ** 2)) / (a * c ** 3 * f ** 4 + 3 * a * c ** 2 * d * e * f ** 3 - 9 * a * c * d ** 2 * e ** 2 * f ** 2 + 5 * a * d ** 3 * e ** 3 * f + b * c ** 3 * e * f ** 3 - 9 * b * c ** 2 * d * e ** 2 * f ** 2 + 15 * b * c * d ** 2 * e ** 3 * f - 7 * b * d ** 3 * e ** 4) + x) / 4$

3.21
$$\int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^3} dx$$

Optimal. Leaf size=291

$$\frac{dx \left(3af \left(-3c^2f^2 - 4cdef + 15d^2e^2 \right) - be \left(3c^2f^2 - 100cdef + 105d^2e^2 \right) \right)}{24e^2f^4} + \frac{(de - cf) \tan^{-1} \left(\frac{\sqrt{f}x}{\sqrt{e}} \right) \left(be \left(-c^2f^2 - 10cdef + 15d^2e^2 \right) - 3af \left(c^2f^2 + 2cdef + 5d^2e^2 \right) \right)}{8e^2f^2(e+fx^2)}$$

Rubi [A] time = 0.42, antiderivative size = 291, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 26, number of rules / integrand size = 0.154, Rules used = {526, 528, 388, 205}

$$\frac{dx \left(3af \left(-3c^2f^2 - 4cdef + 15d^2e^2 \right) - be \left(3c^2f^2 - 100cdef + 105d^2e^2 \right) \right)}{24e^2f^4} + \frac{(de - cf) \tan^{-1} \left(\frac{\sqrt{f}x}{\sqrt{e}} \right) \left(be \left(-c^2f^2 - 10cdef + 15d^2e^2 \right) - 3af \left(c^2f^2 + 2cdef + 5d^2e^2 \right) \right)}{8e^2f^2(e+fx^2)} + \frac{x \left(e + dx^2 \right) \left(be(7de - cf) - 3af(cf + de) \right)}{8e^2f^2(e+fx^2)} + \frac{dx \left(e + dx^2 \right) \left(be(35de - 3cf) - 3af(3cf + 5de) \right)}{24e^2f^3} - \frac{x \left(e + dx^2 \right)^3 \left(be - af \right)}{4ef \left(e + fx^2 \right)^2}$$

Antiderivative was successfully verified.

```
[In] Int[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^3,x]
```

```
[Out] (d*(3*a*f*(15*d^2*e^2 - 4*c*d*e*f - 3*c^2*f^2) - b*e*(105*d^2*e^2 - 100*c*d*e*f + 3*c^2*f^2))*x)/(24*e^2*f^4) + (d*(b*e*(35*d*e - 3*c*f) - 3*a*f*(5*d*e + 3*c*f))*x*(c + d*x^2))/(24*e^2*f^3) - ((b*e - a*f)*x*(c + d*x^2)^3)/(4*e*f*(e + f*x^2)^2) - ((b*e*(7*d*e - c*f) - 3*a*f*(d*e + c*f))*x*(c + d*x^2)^2)/(8*e^2*f^2*(e + f*x^2)) + ((d*e - c*f)*(b*e*(35*d^2*e^2 - 10*c*d*e*f - c^2*f^2) - 3*a*f*(5*d^2*e^2 + 2*c*d*e*f + c^2*f^2))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(8*e^(5/2)*f^(9/2))
```

Rule 205

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 526

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e - a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^3} dx &= -\frac{(be - af)x(c + dx^2)^3}{4ef(e + fx^2)^2} - \frac{\int \frac{(c+dx^2)^2(-c(be+3af)-d(7be-3af)x^2)}{(e+fx^2)^2} dx}{4ef} \\
&= -\frac{(be - af)x(c + dx^2)^3}{4ef(e + fx^2)^2} - \frac{(be(7de - cf) - 3af(de + cf))x(c + dx^2)^2}{8e^2f^2(e + fx^2)} + \frac{\int \frac{(c+dx^2)(-c(be+3af)-d(7be-3af)x^2)}{(e+fx^2)^2} dx}{4ef} \\
&= \frac{d(be(35de - 3cf) - 3af(5de + 3cf))x(c + dx^2)}{24e^2f^3} - \frac{(be - af)x(c + dx^2)^3}{4ef(e + fx^2)^2} - \frac{(be(7de - cf) - 3af(de + cf))x(c + dx^2)^2}{8e^2f^2(e + fx^2)} \\
&= \frac{d(3af(15d^2e^2 - 4cdef - 3c^2f^2) - be(105d^2e^2 - 100cdef + 3c^2f^2))x}{24e^2f^4} + \frac{d(be(35de - 3cf) - 3af(5de + 3cf))x(c + dx^2)}{24e^2f^3} - \frac{(be - af)x(c + dx^2)^3}{4ef(e + fx^2)^2} \\
&= \frac{d(3af(15d^2e^2 - 4cdef - 3c^2f^2) - be(105d^2e^2 - 100cdef + 3c^2f^2))x}{24e^2f^4} + \frac{d(be(35de - 3cf) - 3af(5de + 3cf))x(c + dx^2)}{24e^2f^3} - \frac{(be - af)x(c + dx^2)^3}{4ef(e + fx^2)^2}
\end{aligned}$$

Mathematica [A] time = 0.17, size = 219, normalized size = 0.75

$$\frac{(de - cf) \tan^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e}}\right) (be(-c^2f^2 - 10cdef + 35d^2e^2) - 3af(c^2f^2 + 2cdef + 5d^2e^2))}{8e^{5/2}f^{9/2}} + \frac{d^2x(adf + 3bcf - 3bde)}{f^4} - \frac{x(de - cf)^2(be(13de - cf) - 3af(cf + 3de))}{8e^2f^4(e + fx^2)} + \frac{x(be - af)(de - cf)^3}{4ef^4(e + fx^2)^2} + \frac{bd^3x^3}{3f^3}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^3,x]

[Out] (d^2*(-3*b*d*e + 3*b*c*f + a*d*f)*x)/f^4 + (b*d^3*x^3)/(3*f^3) + ((b*e - a*f)*(d*e - c*f)^3*x)/(4*e*f^4*(e + f*x^2)^2) - ((d*e - c*f)^2*(b*e*(13*d*e - c*f) - 3*a*f*(3*d*e + c*f))*x)/(8*e^2*f^4*(e + f*x^2)) + ((d*e - c*f)*(b*e*(35*d^2*e^2 - 10*c*d*e*f - c^2*f^2) - 3*a*f*(5*d^2*e^2 + 2*c*d*e*f + c^2*f^2))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(8*e^(5/2)*f^(9/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^3} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^3,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^3, x]

fricas [A] time = 1.56, size = 1102, normalized size = 3.79

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^3,x, algorithm="fricas")

[Out] [1/48*(16*b*d^3*e^3*f^4*x^7 - 16*(7*b*d^3*e^4*f^3 - 3*(3*b*c*d^2 + a*d^3)*e^3*f^4)*x^5 - 2*(175*b*d^3*e^5*f^2 - 9*a*c^3*e*f^6 - 75*(3*b*c*d^2 + a*d^3)*e^4*f^3 + 45*(b*c^2*d + a*c*d^2)*e^3*f^4 - 3*(b*c^3 + 3*a*c^2*d)*e^2*f^5)*x^3 - 3*(35*b*d^3*e^6 + 3*a*c^3*e^2*f^4 - 15*(3*b*c*d^2 + a*d^3)*e^5*f + 9*(b*c^2*d + a*c*d^2)*e^4*f^2 + (b*c^3 + 3*a*c^2*d)*e^3*f^3 + (35*b*d^3*e^4*f

$\wedge 2 + 3*a*c^3*f^6 - 15*(3*b*c*d^2 + a*d^3)*e^3*f^3 + 9*(b*c^2*d + a*c*d^2)*e^2*f^4 + (b*c^3 + 3*a*c^2*d)*e*f^5)*x^4 + 2*(35*b*d^3*e^5*f + 3*a*c^3*e*f^5 - 15*(3*b*c*d^2 + a*d^3)*e^4*f^2 + 9*(b*c^2*d + a*c*d^2)*e^3*f^3 + (b*c^3 + 3*a*c^2*d)*e^2*f^4)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) - 6*(35*b*d^3*e^6*f - 5*a*c^3*e^2*f^5 - 15*(3*b*c*d^2 + a*d^3)*e^5*f^2 + 9*(b*c^2*d + a*c*d^2)*e^4*f^3 + (b*c^3 + 3*a*c^2*d)*e^3*f^4)*x)/(e^3*f^7*x^4 + 2*e^4*f^6*x^2 + e^5*f^5), 1/24*(8*b*d^3*e^3*f^4*x^7 - 8*(7*b*d^3*e^4*f^3 - 3*(3*b*c*d^2 + a*d^3)*e^3*f^4)*x^5 - (175*b*d^3*e^5*f^2 - 9*a*c^3*e*f^6 - 75*(3*b*c*d^2 + a*d^3)*e^4*f^3 + 45*(b*c^2*d + a*c*d^2)*e^3*f^4 - 3*(b*c^3 + 3*a*c^2*d)*e^2*f^5)*x^3 + 3*(35*b*d^3*e^6 + 3*a*c^3*e^2*f^4 - 15*(3*b*c*d^2 + a*d^3)*e^5*f + 9*(b*c^2*d + a*c*d^2)*e^4*f^2 + (b*c^3 + 3*a*c^2*d)*e^3*f^3 + (35*b*d^3*e^4*f^2 + 3*a*c^3*f^6 - 15*(3*b*c*d^2 + a*d^3)*e^3*f^3 + 9*(b*c^2*d + a*c*d^2)*e^2*f^4 + (b*c^3 + 3*a*c^2*d)*e*f^5)*x^4 + 2*(35*b*d^3*e^5*f + 3*a*c^3*e*f^5 - 15*(3*b*c*d^2 + a*d^3)*e^4*f^2 + 9*(b*c^2*d + a*c*d^2)*e^3*f^3 + (b*c^3 + 3*a*c^2*d)*e^2*f^4)*x^2)*sqrt(e*f)*arctan(sqrt(e*f)*x/e) - 3*(35*b*d^3*e^6*f - 5*a*c^3*e^2*f^5 - 15*(3*b*c*d^2 + a*d^3)*e^5*f^2 + 9*(b*c^2*d + a*c*d^2)*e^4*f^3 + (b*c^3 + 3*a*c^2*d)*e^3*f^4)*x)/(e^3*f^7*x^4 + 2*e^4*f^6*x^2 + e^5*f^5)]$

giac [A] time = 0.39, size = 371, normalized size = 1.27

$$\frac{(3ac^2f^2 + bc^2f + 3ac^2d^2 + 9ac^2d^2 + 9ad^2f^2 - 45bdf^2 - 15ad^2f^2 + 35bd^2f^2)\arctan\left(\frac{\sqrt{f^2+e}}{f}\right)}{8f^3} - \frac{(3ac^2f^2 + bc^2f^2 + 3ac^2d^2 + 9ac^2d^2 - 15ad^2f^2 + 27bd^2f^2 + 9ad^2f^2 + 5ac^2f^2 - 13bd^2f^2 - bc^2f^2 - 3ac^2d^2 - 9ac^2d^2 - 9ad^2f^2 + 25bd^2f^2 + 7ad^2f^2 - 11bd^2f^2)e^{\frac{3}{2}}}{8(f^2 + e)^{\frac{3}{2}}} + \frac{bd^2f^2 + 9bd^2f + 3ad^2f + 9bd^2d}{3f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^3,x, algorithm="giac")

[Out] 1/8*(3*a*c^3*f^4 + b*c^3*f^3*e + 3*a*c^2*d*f^3*e + 9*b*c^2*d*f^2*e^2 + 9*a*c*d^2*f^2*e^2 - 45*b*c*d^2*f*e^3 - 15*a*d^3*f*e^3 + 35*b*d^3*e^4)*arctan(sqrt(f)*x*e^(-1/2))*e^(-5/2)/f^(9/2) + 1/8*(3*a*c^3*f^5*x^3 + b*c^3*f^4*x^3*e + 3*a*c^2*d*f^4*x^3*e - 15*b*c^2*d*f^3*x^3*e^2 - 15*a*c*d^2*f^3*x^3*e^2 + 27*b*c*d^2*f^2*x^3*e^3 + 9*a*d^3*f^2*x^3*e^3 + 5*a*c^3*f^4*x*e - 13*b*d^3*f*x^3*e^4 - b*c^3*f^3*x*e^2 - 3*a*c^2*d*f^3*x*e^2 - 9*b*c^2*d*f^2*x*e^3 - 9*a*c*d^2*f^2*x*e^3 + 21*b*c*d^2*f*x*e^4 + 7*a*d^3*f*x*e^4 - 11*b*d^3*x*e^5)*e^(-2)/((f*x^2 + e)^2*f^4) + 1/3*(b*d^3*f^6*x^3 + 9*b*c*d^2*f^6*x + 3*a*d^3*f^6*x - 9*b*d^3*f^5*x*e)/f^9

maple [B] time = 0.02, size = 589, normalized size = 2.02

$$\frac{3ac^2f^2}{8(f^2 + e)^{\frac{3}{2}}} - \frac{3ac^2f^2 + bc^2f + 3ac^2d^2 + 9ac^2d^2 + 9ad^2f^2 - 45bdf^2 - 15ad^2f^2 + 35bd^2f^2}{8(f^2 + e)^{\frac{3}{2}}} \arctan\left(\frac{\sqrt{f^2+e}}{f}\right) + \frac{bd^2f^2 + 9bd^2f + 3ad^2f + 9bd^2d}{3f^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^3,x)

[Out] -45/8/f^3*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c*d^2+27/8/f^2/(f*x^2+e)^2*x^3*b*c*d^2*e-9/8/f^2/(f*x^2+e)^2*a*c*d^2*e*x-9/8/f^2/(f*x^2+e)^2*b*c^2*d*e*x+21/8/f^3/(f*x^2+e)^2*b*c*d^2*e^2*x+3/8/f/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^2*d+3/8/(f*x^2+e)^2/e*x^3*a*c^2*d-15/8/f/(f*x^2+e)^2*x^3*b*c^2*d-13/8/f^3/(f*x^2+e)^2*x^3*b*d^3*e^2-3/8/f/(f*x^2+e)^2*a*c^2*d*x+7/8/f^3/(f*x^2+e)^2*a*d^3*e^2*x-11/8/f^4/(f*x^2+e)^2*b*d^3*e^3*x+9/8/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d^2-15/8/f^3*e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*d^3+1/8/f/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^3+9/8/f^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2*d+35/8/f^4*e^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*d^3+3/8*f/(f*x^2+e)^2/e^2*x^3*a*c^3-15/8/f/(f*x^2+e)^2*x^3*a*c*d^2+9/8/f^2/(f*x^2+e)^2*x^3*a*d^3*e+d^3/f^3*a*x+1/3*d^3/f^3*x^3*b-3*d^3/f^4*b*e*x-1/8/f/(f*x^2+e)^2*b*c^3*x+3/8/e^2/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c^3+3*d^2/f^3*b*c*x+1/8/(f*x^2+e)^2/e*x^3*b*c^3+5/8/(f*x^2+e)^2/e*x*a*c^3

maxima [A] time = 1.94, size = 343, normalized size = 1.18

$$\frac{(13bd^2f - 3ac^2f - 9(3bd^2 + ad^2)f^2 + 15(bc^2d + ad^2)f^2 - (bc^3 + 3ac^2d)f^3 + (11bd^2f^2 - 5ac^2f^2 - 7(3bd^2 + ad^2)f^2 + 9(bc^2d + ad^2)f^2 + (bc^3 + 3ac^2d)f^2))x}{8(e^2f^2 + 2e^2f^2 + e^2f^2)} - \frac{bd^2f^2 - 3(3bd^2 - (3bd^2 + ad^2))f}{3f^2} + \frac{(35bd^2 + 3ac^2f^2 - 15(3bd^2 + ad^2)f^2 + 9(bc^2d + ad^2)f^2 + (bc^3 + 3ac^2d)f^2)\arctan\left(\frac{e}{\sqrt{f^2+e}}\right)}{8\sqrt{f^2+e}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^3,x, algorithm="maxima")

[Out]
$$-1/8*((13*b*d^3*e^4*f - 3*a*c^3*f^5 - 9*(3*b*c*d^2 + a*d^3)*e^3*f^2 + 15*(b*c^2*d + a*c*d^2)*e^2*f^3 - (b*c^3 + 3*a*c^2*d)*e*f^4)*x^3 + (11*b*d^3*e^5 - 5*a*c^3*e*f^4 - 7*(3*b*c*d^2 + a*d^3)*e^4*f + 9*(b*c^2*d + a*c*d^2)*e^3*f^2 + (b*c^3 + 3*a*c^2*d)*e^2*f^3)*x)/(e^2*f^6*x^4 + 2*e^3*f^5*x^2 + e^4*f^4) + 1/3*(b*d^3*f*x^3 - 3*(3*b*d^3*e - (3*b*c*d^2 + a*d^3)*f)*x)/f^4 + 1/8*(35*b*d^3*e^4 + 3*a*c^3*f^4 - 15*(3*b*c*d^2 + a*d^3)*e^3*f + 9*(b*c^2*d + a*c*d^2)*e^2*f^2 + (b*c^3 + 3*a*c^2*d)*e*f^3)*\arctan(f*x/\sqrt{e*f})/(\sqrt{e*f})*e^2*f^4)$$

mupad [B] time = 0.22, size = 495, normalized size = 1.70

$$\frac{\frac{d^3 b^2 x^2 + 2 a d^2 b x + 2 a^2 d}{e^2 f^2 + 2 e f + e^2} + \frac{3 b d^2 c}{f^2} + \frac{b d^2 a}{3 f^2} + \frac{\arctan\left(\frac{\sqrt{e} (b x + d) \sqrt{e f}}{\sqrt{e f^2 + 2 e f + e^2}}\right) (c f - d) (b c^2 f^2 + 3 a c^2 f + 10 b c d e f + 6 a c d e^2 - 35 b d^2 e + 15 a d^2 f)}{8 b^2 c^2 f^2}}{e^2 f^2 + 2 e f + e^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^3,x)

[Out]
$$\left(\frac{(x^3(3*a*c^3*f^5 + 9*a*d^3*e^3*f^2 + b*c^3*e*f^4 - 13*b*d^3*e^4*f + 3*a*c^2*d*e*f^4 - 15*a*c*d^2*e^2*f^3 + 27*b*c*d^2*e^3*f^2 - 15*b*c^2*d*e^2*f^3))}{(8*e^2)} - \frac{(x(11*b*d^3*e^4 - 5*a*c^3*f^4 - 7*a*d^3*e^3*f + b*c^3*e*f^3 + 3*a*c^2*d*e*f^3 - 21*b*c*d^2*e^3*f + 9*a*c*d^2*e^2*f^2 + 9*b*c^2*d*e^2*f^2))}{(8*e)}\right)/(e^2*f^4 + f^6*x^4 + 2*e*f^5*x^2) + x*((a*d^3 + 3*b*c*d^2)/f^3 - (3*b*d^3*e)/f^4) + (b*d^3*x^3)/(3*f^3) + (\operatorname{atan}((f^{1/2})x*(c*f - d*e)*(3*a*c^2*f^3 - 35*b*d^2*e^3 + 15*a*d^2*e^2*f + b*c^2*e*f^2 + 6*a*c*d*e*f^2 + 10*b*c*d*e^2*f)))/(e^{1/2}*(3*a*c^3*f^4 + 35*b*d^3*e^4 - 15*a*d^3*e^3*f + b*c^3*e*f^3 + 3*a*c^2*d*e*f^3 - 45*b*c*d^2*e^3*f + 9*a*c*d^2*e^2*f^2 + 9*b*c^2*d*e^2*f^2))*(c*f - d*e)*(3*a*c^2*f^3 - 35*b*d^2*e^3 + 15*a*d^2*e^2*f + b*c^2*e*f^2 + 6*a*c*d*e*f^2 + 10*b*c*d*e^2*f))/(8*e^{5/2}*f^{9/2})$$

sympy [B] time = 26.16, size = 865, normalized size = 2.97

$$\frac{\sqrt{e} (c f - d e) \sqrt{e f} \operatorname{atan}\left(\frac{\sqrt{e} (b x + d) \sqrt{e f}}{\sqrt{e f^2 + 2 e f + e^2}}\right) (c f - d e) (3 a c^2 f^3 - 35 b d^2 e^3 + 15 a d^2 e^2 f + b c^2 e f^2 + 6 a c d e f^2 + 10 b c d e^2 f)}{8 b^2 c^2 f^2} + \frac{b d^3 x^3}{3 f^3} + \frac{x (a d^3 + 3 b c d^2)}{f^3} - \frac{3 b d^3 e}{f^4} - \frac{\sqrt{-1/(e f^2 + 2 e f + e^2)} (c f - d e) (3 a c^2 f^3 + 6 a c d e f^2 + 15 a d^2 e^2 f + b c^2 e f^2 + 10 b c d e^2 f - 35 b d^2 e^3) \log(-e f^2 + 2 e f + e^2)}{8 e^2 f^4 + 8 e f^6 x^4 + 2 e^3 f^5 x^2} + \frac{\sqrt{-1/(e f^2 + 2 e f + e^2)} (c f - d e) (3 a c^2 f^3 + 6 a c d e f^2 + 15 a d^2 e^2 f + b c^2 e f^2 + 10 b c d e^2 f - 35 b d^2 e^3) \log(e f^2 + 2 e f + e^2)}{8 e^2 f^4 + 8 e f^6 x^4 + 2 e^3 f^5 x^2} + \frac{x (5 a c^3 e f^4 - 3 a c^2 d e^2 f^3 - 9 a c d^2 e^3 f^2 + 7 a d^3 e^4 f - b c^3 e^2 f^3 - 9 b c^2 d e^3 f^2 + 21 b c d^2 e^4 f - 11 b d^3 e^5)}{8 e^4 f^4 + 16 e^3 f^5 x^2 + 8 e^2 f^6 x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3/(f*x**2+e)**3,x)

[Out]
$$b*d**3*x**3/(3*f**3) + x*(a*d**3/f**3 + 3*b*c*d**2/f**3 - 3*b*d**3*e/f**4) - \sqrt{-1/(e**5*f**9)}*(c*f - d*e)*(3*a*c**2*f**3 + 6*a*c*d*e*f**2 + 15*a*d**2*e**2*f + b*c**2*e*f**2 + 10*b*c*d*e**2*f - 35*b*d**2*e**3)*\log(-e**3*f**4*\sqrt{-1/(e**5*f**9)}*(c*f - d*e)*(3*a*c**2*f**3 + 6*a*c*d*e*f**2 + 15*a*d**2*e**2*f + b*c**2*e*f**2 + 10*b*c*d*e**2*f - 35*b*d**2*e**3)/(3*a*c**3*f**4 + 3*a*c**2*d*e*f**3 + 9*a*c*d**2*e**2*f**2 - 15*a*d**3*e**3*f + b*c**3*e*f**3 + 9*b*c**2*d*e**2*f**2 - 45*b*c*d**2*e**3*f + 35*b*d**3*e**4) + x)/16 + \sqrt{-1/(e**5*f**9)}*(c*f - d*e)*(3*a*c**2*f**3 + 6*a*c*d*e*f**2 + 15*a*d**2*e**2*f + b*c**2*e*f**2 + 10*b*c*d*e**2*f - 35*b*d**2*e**3)*\log(e**3*f**4*\sqrt{-1/(e**5*f**9)}*(c*f - d*e)*(3*a*c**2*f**3 + 6*a*c*d*e*f**2 + 15*a*d**2*e**2*f + b*c**2*e*f**2 + 10*b*c*d*e**2*f - 35*b*d**2*e**3)/(3*a*c**3*f**4 + 3*a*c**2*d*e*f**3 + 9*a*c*d**2*e**2*f**2 - 15*a*d**3*e**3*f + b*c**3*e*f**3 + 9*b*c**2*d*e**2*f**2 - 45*b*c*d**2*e**3*f + 35*b*d**3*e**4) + x)/16 + (x**3*(3*a*c**3*f**5 + 3*a*c**2*d*e*f**4 - 15*a*c*d**2*e**2*f**3 + 9*a*d**3*e**3*f**2 + b*c**3*e*f**4 - 15*b*c**2*d*e**2*f**3 + 27*b*c*d**2*e**3*f**2 - 13*b*d**3*e**4*f) + x*(5*a*c**3*e*f**4 - 3*a*c**2*d*e**2*f**3 - 9*a*c*d**2*e**3*f**2 + 7*a*d**3*e**4*f - b*c**3*e**2*f**3 - 9*b*c**2*d*e**3*f**2 + 21*b*c*d**2*e**4*f - 11*b*d**3*e**5))/(8*e**4*f**4 + 16*e**3*f**5*x**2 + 8*e**2*f**6*x**4)$$

3.22
$$\int \frac{(a+bx^2)(c+dx^2)^3}{(e+fx^2)^4} dx$$

Optimal. Leaf size=348

$$\frac{dx \left(be \left(-3c^2 f^2 - 10cdef + 105d^2 e^2 \right) - af \left(15c^2 f^2 + 14cdef + 15d^2 e^2 \right) \right)}{48e^3 f^4} - \frac{x \left(c + dx^2 \right) \left(be \left(-3c^2 f^2 - 8cdef + 35d^2 e^2 \right) - af \left(15c^2 f^2 + 14cdef + 15d^2 e^2 \right) \right)}{48e^3 f^3 \left(e + fx^2 \right)}$$

Rubi [A] time = 0.45, antiderivative size = 348, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 26, number of rules / integrand size = 0.115, Rules used = {526, 388, 205}

$$\frac{x(c+dx^2)(be(-3c^2f^2-8cdef+35d^2e^2)-af(15c^2f^2+14cdef+15d^2e^2))}{48e^3f^3(e+fx^2)} + \frac{dx(be(-3c^2f^2-10cdef+105d^2e^2)-af(15c^2f^2+14cdef+15d^2e^2))}{48e^3f^4} - \frac{\tan^{-1}\left(\frac{dx}{\sqrt{e}}\right)(be(-3c^2df^2-c^2f^2-15af^2f+35d^2e^2)-af(3c^2df^2+5c^2f^2+3af^2f+5d^2e^2))}{16e^{9/2}f^2} - \frac{x(c+dx^2)(be(7de-cf)-af(5cf+de))}{24e^2f^2(e+fx^2)} - \frac{x(c+dx^2)(be-af)}{6ef(e+fx^2)}$$

Antiderivative was successfully verified.

```
[In] Int[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^4,x]
[Out] (d*(b*e*(105*d^2*e^2 - 10*c*d*e*f - 3*c^2*f^2) - a*f*(15*d^2*e^2 + 14*c*d*e*f + 15*c^2*f^2))*x)/(48*e^3*f^4) - ((b*e - a*f)*x*(c + d*x^2)^3)/(6*e*f*(e + f*x^2)^3) - ((b*e*(7*d*e - c*f) - a*f*(d*e + 5*c*f))*x*(c + d*x^2)^2)/(24*e^2*f^2*(e + f*x^2)^2) - ((b*e*(35*d^2*e^2 - 8*c*d*e*f - 3*c^2*f^2) - a*f*(5*d^2*e^2 + 4*c*d*e*f + 15*c^2*f^2))*x*(c + d*x^2))/(48*e^3*f^3*(e + f*x^2)) - ((b*e*(35*d^3*e^3 - 15*c*d^2*e^2*f - 3*c^2*d*e*f^2 - c^3*f^3) - a*f*(5*d^3*e^3 + 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 + 5*c^3*f^3))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(16*e^(7/2)*f^(9/2))
```

Rule 205

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 388

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p + 1))/(b*(n*(p + 1) + 1)), x] - Dist[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]
```

Rule 526

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(a*b*n*(p + 1)), x] + Dist[1/(a*b*n*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^(q - 1)*Simp[c*(b*e*n*(p + 1) + b*e - a*f) + d*(b*e*n*(p + 1) + (b*e - a*f)*(n*q + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && GtQ[q, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^4} dx &= -\frac{(be - af)x(c + dx^2)^3}{6ef(e + fx^2)^3} - \frac{\int \frac{(c + dx^2)^2(-c(be + 5af) - d(7be - af)x^2)}{(e + fx^2)^3} dx}{6ef} \\
&= -\frac{(be - af)x(c + dx^2)^3}{6ef(e + fx^2)^3} - \frac{(be(7de - cf) - af(de + 5cf))x(c + dx^2)^2}{24e^2f^2(e + fx^2)^2} + \frac{\int \frac{(c + dx^2)(c^2 - d^2x^2)}{(e + fx^2)^3} dx}{6ef} \\
&= -\frac{(be - af)x(c + dx^2)^3}{6ef(e + fx^2)^3} - \frac{(be(7de - cf) - af(de + 5cf))x(c + dx^2)^2}{24e^2f^2(e + fx^2)^2} - \frac{(be(35d^2e^2 - 10cdef - 3c^2f^2))x}{48e^3f^4} \\
&= \frac{d(be(105d^2e^2 - 10cdef - 3c^2f^2) - af(15d^2e^2 + 14cdef + 15c^2f^2))x}{48e^3f^4} - \frac{(be - af)x}{6ef} \\
&= \frac{d(be(105d^2e^2 - 10cdef - 3c^2f^2) - af(15d^2e^2 + 14cdef + 15c^2f^2))x}{48e^3f^4} - \frac{(be - af)x}{6ef}
\end{aligned}$$

Mathematica [A] time = 0.22, size = 295, normalized size = 0.85

$$\frac{x(de - cf)(be(-c^2f^2 - 4cdef + 29d^2e^2) - af(5c^2f^2 + 8cdef + 11d^2e^2))}{16e^3f^4(e + fx^2)} - \frac{\tan^{-1}\left(\frac{\sqrt{fx}}{\sqrt{e}}\right)(be(-c^3f^3 - 3c^2def^2 - 15cd^2e^2f + 35d^3e^3) - af(5c^3f^3 + 3c^2def^2 + 3cd^2e^2f + 5d^3e^3))}{16e^{7/2}f^{9/2}} - \frac{x(de - cf)^2(be(19de - cf) - af(5cf + 13de))}{24e^2f^4(e + fx^2)^2} + \frac{x(be - af)(de - cf)^3}{6ef^4(e + fx^2)^3} + \frac{bf^2x}{f^4}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^4,x]

[Out] (b*d^3*x)/f^4 + ((b*e - a*f)*(d*e - c*f)^3*x)/(6*e*f^4*(e + f*x^2)^3) - ((d*e - c*f)^2*(b*e*(19*d*e - c*f) - a*f*(13*d*e + 5*c*f))*x)/(24*e^2*f^4*(e + f*x^2)^2) + ((d*e - c*f)*(b*e*(29*d^2*e^2 - 4*c*d*e*f - c^2*f^2) - a*f*(11*d^2*e^2 + 8*c*d*e*f + 5*c^2*f^2))*x)/(16*e^3*f^4*(e + f*x^2)) - ((b*e*(35*d^3*e^3 - 15*c*d^2*e^2*f - 3*c^2*d*e*f^2 - c^3*f^3) - a*f*(5*d^3*e^3 + 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 + 5*c^3*f^3))*ArcTan[(Sqrt[f]*x)/Sqrt[e]])/(16*e^(7/2)*f^(9/2))

IntegrateAlgebraic [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)(c + dx^2)^3}{(e + fx^2)^4} dx$$

Verification is not applicable to the result.

[In] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^4,x]

[Out] IntegrateAlgebraic[((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^4, x]

fricas [B] time = 1.38, size = 1422, normalized size = 4.09

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^4,x, algorithm="fricas")

[Out] [1/96*(96*b*d^3*e^4*f^4*x^7 + 6*(77*b*d^3*e^5*f^3 + 5*a*c^3*e*f^7 - 11*(3*b*c*d^2 + a*d^3)*e^4*f^4 + 3*(b*c^2*d + a*c*d^2)*e^3*f^5 + (b*c^3 + 3*a*c^2*d)*e^2*f^6)*x^5 + 16*(35*b*d^3*e^6*f^2 + 5*a*c^3*e^2*f^6 - 5*(3*b*c*d^2 + a

```
*d^3)*e^5*f^3 - 3*(b*c^2*d + a*c*d^2)*e^4*f^4 + (b*c^3 + 3*a*c^2*d)*e^3*f^5
)*x^3 + 3*(35*b*d^3*e^7 - 5*a*c^3*e^3*f^4 - 5*(3*b*c*d^2 + a*d^3)*e^6*f - 3
*(b*c^2*d + a*c*d^2)*e^5*f^2 - (b*c^3 + 3*a*c^2*d)*e^4*f^3 + (35*b*d^3*e^4*
f^3 - 5*a*c^3*f^7 - 5*(3*b*c*d^2 + a*d^3)*e^3*f^4 - 3*(b*c^2*d + a*c*d^2)*e
^2*f^5 - (b*c^3 + 3*a*c^2*d)*e*f^6)*x^6 + 3*(35*b*d^3*e^5*f^2 - 5*a*c^3*e*f
^6 - 5*(3*b*c*d^2 + a*d^3)*e^4*f^3 - 3*(b*c^2*d + a*c*d^2)*e^3*f^4 - (b*c^3
+ 3*a*c^2*d)*e^2*f^5)*x^4 + 3*(35*b*d^3*e^6*f - 5*a*c^3*e^2*f^5 - 5*(3*b*c
*d^2 + a*d^3)*e^5*f^2 - 3*(b*c^2*d + a*c*d^2)*e^4*f^3 - (b*c^3 + 3*a*c^2*d)
*e^3*f^4)*x^2)*sqrt(-e*f)*log((f*x^2 - 2*sqrt(-e*f)*x - e)/(f*x^2 + e)) + 6
*(35*b*d^3*e^7*f + 11*a*c^3*e^3*f^5 - 5*(3*b*c*d^2 + a*d^3)*e^6*f^2 - 3*(b*
c^2*d + a*c*d^2)*e^5*f^3 - (b*c^3 + 3*a*c^2*d)*e^4*f^4)*x)/(e^4*f^8*x^6 + 3
*e^5*f^7*x^4 + 3*e^6*f^6*x^2 + e^7*f^5), 1/48*(48*b*d^3*e^4*f^4*x^7 + 3*(77
*b*d^3*e^5*f^3 + 5*a*c^3*e*f^7 - 11*(3*b*c*d^2 + a*d^3)*e^4*f^4 + 3*(b*c^2*
d + a*c*d^2)*e^3*f^5 + (b*c^3 + 3*a*c^2*d)*e^2*f^6)*x^5 + 8*(35*b*d^3*e^6*f
^2 + 5*a*c^3*e^2*f^6 - 5*(3*b*c*d^2 + a*d^3)*e^5*f^3 - 3*(b*c^2*d + a*c*d^2
)*e^4*f^4 + (b*c^3 + 3*a*c^2*d)*e^3*f^5)*x^3 - 3*(35*b*d^3*e^7 - 5*a*c^3*e^
3*f^4 - 5*(3*b*c*d^2 + a*d^3)*e^6*f - 3*(b*c^2*d + a*c*d^2)*e^5*f^2 - (b*c^
3 + 3*a*c^2*d)*e^4*f^3 + (35*b*d^3*e^4*f^3 - 5*a*c^3*f^7 - 5*(3*b*c*d^2 + a
*d^3)*e^3*f^4 - 3*(b*c^2*d + a*c*d^2)*e^2*f^5 - (b*c^3 + 3*a*c^2*d)*e*f^6)*
x^6 + 3*(35*b*d^3*e^5*f^2 - 5*a*c^3*e*f^6 - 5*(3*b*c*d^2 + a*d^3)*e^4*f^3 -
3*(b*c^2*d + a*c*d^2)*e^3*f^4 - (b*c^3 + 3*a*c^2*d)*e^2*f^5)*x^4 + 3*(35*b
*d^3*e^6*f - 5*a*c^3*e^2*f^5 - 5*(3*b*c*d^2 + a*d^3)*e^5*f^2 - 3*(b*c^2*d +
a*c*d^2)*e^4*f^3 - (b*c^3 + 3*a*c^2*d)*e^3*f^4)*x^2)*sqrt(e*f)*arctan(sqrt
(e*f)*x/e) + 3*(35*b*d^3*e^7*f + 11*a*c^3*e^3*f^5 - 5*(3*b*c*d^2 + a*d^3)*e
^6*f^2 - 3*(b*c^2*d + a*c*d^2)*e^5*f^3 - (b*c^3 + 3*a*c^2*d)*e^4*f^4)*x)/(e
^4*f^8*x^6 + 3*e^5*f^7*x^4 + 3*e^6*f^6*x^2 + e^7*f^5)]
```

giac [A] time = 0.37, size = 447, normalized size = 1.28

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^4,x, algorithm="giac")

```
[Out] b*d^3*x/f^4 + 1/16*(5*a*c^3*f^4 + b*c^3*f^3*e + 3*a*c^2*d*f^3*e + 3*b*c^2*d
*f^2*e^2 + 3*a*c*d^2*f^2*e^2 + 15*b*c*d^2*f*e^3 + 5*a*d^3*f*e^3 - 35*b*d^3*
e^4)*arctan(sqrt(f)*x*e^(-1/2))*e^(-7/2)/f^(9/2) + 1/48*(15*a*c^3*f^6*x^5 +
3*b*c^3*f^5*x^5*e + 9*a*c^2*d*f^5*x^5*e + 9*b*c^2*d*f^4*x^5*e^2 + 9*a*c*d^
2*f^4*x^5*e^2 - 99*b*c*d^2*f^3*x^5*e^3 - 33*a*d^3*f^3*x^5*e^3 + 40*a*c^3*f^
5*x^3*e + 87*b*d^3*f^2*x^5*e^4 + 8*b*c^3*f^4*x^3*e^2 + 24*a*c^2*d*f^4*x^3*e
^2 - 24*b*c^2*d*f^3*x^3*e^3 - 24*a*c*d^2*f^3*x^3*e^3 - 120*b*c*d^2*f^2*x^3*
e^4 - 40*a*d^3*f^2*x^3*e^4 + 33*a*c^3*f^4*x*e^2 + 136*b*d^3*f*x^3*e^5 - 3*b
*c^3*f^3*x*e^3 - 9*a*c^2*d*f^3*x*e^3 - 9*b*c^2*d*f^2*x*e^4 - 9*a*c*d^2*f^2*
x*e^4 - 45*b*c*d^2*f*x*e^5 - 15*a*d^3*f*x*e^5 + 57*b*d^3*x*e^6)*e^(-3)/((f*
x^2 + e)^3*f^4)
```

maple [B] time = 0.01, size = 735, normalized size = 2.11

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^4,x)

```
[Out] -15/16/f^3/(f*x^2+e)^3*b*c*d^2*e^2*x+3/16/f/e^2/(e*f)^(1/2)*arctan(1/(e*f)^(
1/2)*f*x)*a*c^2*d+3/16/f^2/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*a*c*d^2
+3/16/f^2/e/(e*f)^(1/2)*arctan(1/(e*f)^(1/2)*f*x)*b*c^2*d-5/2/f^2/(f*x^2+e)
^3*x^3*b*c*d^2*e-3/16/f^2/(f*x^2+e)^3*a*c*d^2*e*x-3/16/f^2/(f*x^2+e)^3*b*c^
2*d*e*x+3/16*f/(f*x^2+e)^3/e^2*x^5*a*c^2*d+29/16/f^2/(f*x^2+e)^3*x^5*b*d^3*
e+5/6*f/(f*x^2+e)^3/e^2*x^3*a*c^3-1/2/f/(f*x^2+e)^3*x^3*a*c*d^2-5/6/f^2/(f*
```


$$x^2+e)^3x^3ad^3e+3/16/(fx^2+e)^3/ex^5b^2c^2d+1/2/(fx^2+e)^3/ex^3ac^2d+3/16/(fx^2+e)^3/ex^5ac^2d^2+19/16/f^4/(fx^2+e)^3b^2d^3e^3x+15/16/f^3/(ef)^{(1/2)}\arctan(1/(ef)^{(1/2)}fx)bc^2d-35/16/f^4e/(ef)^{(1/2)}\arctan(1/(ef)^{(1/2)}fx)b^2d^3+1/16/f/e^2/(ef)^{(1/2)}\arctan(1/(ef)^{(1/2)}fx)bc^3-1/2/f/(fx^2+e)^3x^3b^2c^2d+17/6/f^3/(fx^2+e)^3x^3b^2d^3e^2-3/16/f/(fx^2+e)^3ac^2d^2x-5/16/f^3/(fx^2+e)^3ad^3e^2x+5/16f^2/(fx^2+e)^3/ex^5ac^3+1/16f/(fx^2+e)^3/ex^2x^5b^2c^3-33/16/f/(fx^2+e)^3x^5b^2c^2d+b^2d^3/f^4x+5/16/e^3/(ef)^{(1/2)}\arctan(1/(ef)^{(1/2)}fx)ac^3+5/16/f^3/(ef)^{(1/2)}\arctan(1/(ef)^{(1/2)}fx)ad^3-11/16/f/(fx^2+e)^3x^5ad^3-1/16/f/(fx^2+e)^3b^2c^3x+1/6/(fx^2+e)^3/ex^3b^2c^3+11/16/(fx^2+e)^3/exac^3$$

maxima [A] time = 2.39, size = 416, normalized size = 1.20

$$\frac{b^2x^3(29b^2d^3e^3f^5+5ac^2f^5-11(3b^2d+ad^2)f^4+3(b^2d+ad^2)f^3+(b^2+3ac^2d)f^2)+8(17b^2d^2f+5ac^2f^2-5(3b^2d+ad^2)f^2-3(b^2d+ad^2)f^2)+3(19b^2d^2+11ac^2d^2-5(3b^2d+ad^2)f^2-3(b^2d+ad^2)f^2)-(b^2+3ac^2d)f^2}{48(f^2+3ef^2+3e^2f^2+e^3f^2)} \cdot \frac{(35b^2d^3e^3f^4-5ac^3f^4-5(3b^2d+ad^2)f^3-3(b^2d+ad^2)f^2-(b^2+3ac^2d)f^2)\arctan\left(\frac{fx}{\sqrt{ef}}\right)}{16\sqrt{ef}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^3/(f*x^2+e)^4,x, algorithm="maxima")

[Out] $b^2d^3x/f^4 + 1/48*(3*(29*b^2d^3e^4f^2 + 5*a*c^3f^6 - 11*(3*b^2c^2d^2 + a*d^3)*e^3f^3 + 3*(b^2c^2d + a*c^2d^2)*e^2f^4 + (b^2c^3 + 3*a*c^2d)*e*f^5)*x^5 + 8*(17*b^2d^3e^5f + 5*a*c^3e*f^5 - 5*(3*b^2c^2d^2 + a*d^3)*e^4f^2 - 3*(b^2c^2d + a*c^2d^2)*e^3f^3 + (b^2c^3 + 3*a*c^2d)*e^2f^4)*x^3 + 3*(19*b^2d^3e^6 + 11*a*c^3e^2f^4 - 5*(3*b^2c^2d^2 + a*d^3)*e^5f - 3*(b^2c^2d + a*c^2d^2)*e^4f^2 - (b^2c^3 + 3*a*c^2d)*e^3f^3)*x)/(e^3f^7x^6 + 3e^4f^6x^4 + 3e^5f^5x^2 + e^6f^4) - 1/16*(35*b^2d^3e^4 - 5*a*c^3f^4 - 5*(3*b^2c^2d^2 + a*d^3)*e^3f - 3*(b^2c^2d + a*c^2d^2)*e^2f^2 - (b^2c^3 + 3*a*c^2d)*e*f^3)*\arctan(f*x/\sqrt{e*f})/(\sqrt{e*f}*e^3f^4)$

mupad [B] time = 1.19, size = 444, normalized size = 1.28

$$\frac{b^2x^3(29b^2d^3e^3f^5+5ac^2f^5-11(3b^2d+ad^2)f^4+3(b^2d+ad^2)f^3+(b^2+3ac^2d)f^2)+8(17b^2d^2f+5ac^2f^2-5(3b^2d+ad^2)f^2-3(b^2d+ad^2)f^2)+3(19b^2d^2+11ac^2d^2-5(3b^2d+ad^2)f^2-3(b^2d+ad^2)f^2)-(b^2+3ac^2d)f^2}{48(f^2+3ef^2+3e^2f^2+e^3f^2)} \cdot \frac{\arctan\left(\frac{fx}{\sqrt{ef}}\right)(b^2c^3f^4+5ac^2f^4+3b^2c^2d^2f^2+3ac^2d^2f+15b^2c^2d^2f+3ac^2d^2f-35b^2d^3e^4+5ad^3e^4)}{16\sqrt{ef}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)*(c + d*x^2)^3)/(e + f*x^2)^4,x)

[Out] $((x^3*(5*a*c^3f^5 - 5*a*d^3e^3f^2 + b^2c^3e*f^4 + 17*b^2d^3e^4f + 3*a*c^2d*e*f^4 - 3*a*c^2d^2e^2f^3 - 15*b^2c^2d^2e^3f^2 - 3*b^2c^2d^2e^2f^3))/(6*e^2) + (x^5*(5*a*c^3f^6 - 11*a*d^3e^3f^3 + 29*b^2d^3e^4f^2 + b^2c^3e*f^5 + 3*a*c^2d*e*f^5 + 3*a*c^2d^2e^2f^4 - 33*b^2c^2d^2e^3f^3 + 3*b^2c^2d^2e^2f^4))/(16*e^3) - (x*(5*a*d^3e^3f - 19*b^2d^3e^4 - 11*a*c^3f^4 + b^2c^3e*f^3 + 3*a*c^2d*e*f^3 + 15*b^2c^2d^2e^3f + 3*a*c^2d^2e^2f^2 + 3*b^2c^2d^2e^2f^2))/(16*e))/(e^3f^4 + f^7x^6 + 3e*f^6x^4 + 3e^2f^5x^2) + (b^2d^3x)/f^4 + (\operatorname{atan}((f^{1/2})x)/e^{1/2})*(5*a*c^3f^4 - 35*b^2d^3e^4 + 5*a*d^3e^3f + b^2c^3e*f^3 + 3*a*c^2d*e*f^3 + 15*b^2c^2d^2e^3f + 3*a*c^2d^2e^2f^2 + 3*b^2c^2d^2e^2f^2))/(16*e^{(7/2)}*f^{(9/2)})$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)*(d*x**2+c)**3/(f*x**2+e)**4,x)

[Out] Timed out

$$3.23 \quad \int \frac{(a+bx^2)\sqrt{c+dx^2}}{e+fx^2} dx$$

Optimal. Leaf size=128

$$\frac{\tanh^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c+dx^2}}\right)(-2adf - bcf + 2bde)}{2\sqrt{d}f^2} + \frac{(be - af)\sqrt{de - cf} \tanh^{-1}\left(\frac{x\sqrt{de - cf}}{\sqrt{e}\sqrt{c+dx^2}}\right)}{\sqrt{e}f^2} + \frac{bx\sqrt{c + dx^2}}{2f}$$

Rubi [A] time = 0.14, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {528, 523, 217, 206, 377, 208}

$$\frac{\tanh^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c+dx^2}}\right)(-2adf - bcf + 2bde)}{2\sqrt{d}f^2} + \frac{(be - af)\sqrt{de - cf} \tanh^{-1}\left(\frac{x\sqrt{de - cf}}{\sqrt{e}\sqrt{c+dx^2}}\right)}{\sqrt{e}f^2} + \frac{bx\sqrt{c + dx^2}}{2f}$$

Antiderivative was successfully verified.

```
[In] Int[((a + b*x^2)*Sqrt[c + d*x^2])/(e + f*x^2), x]
```

```
[Out] (b*x*Sqrt[c + d*x^2])/(2*f) - ((2*b*d*e - b*c*f - 2*a*d*f)*ArcTanh[(Sqrt[d]*x)/Sqrt[c + d*x^2]])/(2*Sqrt[d]*f^2) + ((b*e - a*f)*Sqrt[d*e - c*f]*ArcTanh[(Sqrt[d*e - c*f]*x)/(Sqrt[e]*Sqrt[c + d*x^2])])/(Sqrt[e]*f^2)
```

Rule 206

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 208

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 217

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 377

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 523

```
Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)]), x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 528

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(f*x*(a + b*x^n)^(p + 1)*(c + d*x^n)^q)/(b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
```

$a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /;$ FreeQ[{
 $a, b, c, d, e, f, n, p}, x]$ && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)\sqrt{c + dx^2}}{e + fx^2} dx &= \frac{bx\sqrt{c + dx^2}}{2f} + \frac{\int \frac{-c(be-2af)+(-2bde+bcf+2adf)x^2}{\sqrt{c+dx^2}(e+fx^2)} dx}{2f} \\ &= \frac{bx\sqrt{c + dx^2}}{2f} + \frac{((be - af)(de - cf)) \int \frac{1}{\sqrt{c+dx^2}(e+fx^2)} dx}{f^2} - \frac{(2bde - bcf - 2adf) \int \frac{1}{\sqrt{c+dx^2}} dx}{2f^2} \\ &= \frac{bx\sqrt{c + dx^2}}{2f} + \frac{((be - af)(de - cf)) \text{Subst}\left(\int \frac{1}{e-(de-cf)x^2} dx, x, \frac{x}{\sqrt{c+dx^2}}\right)}{f^2} - \frac{(2bde - bcf - 2adf) \int \frac{1}{\sqrt{c+dx^2}} dx}{2f^2} \\ &= \frac{bx\sqrt{c + dx^2}}{2f} - \frac{(2bde - bcf - 2adf) \tanh^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c+dx^2}}\right)}{2\sqrt{d}f^2} + \frac{(be - af)\sqrt{de - cf} \tanh^{-1}\left(\frac{\sqrt{d}x}{\sqrt{c+dx^2}}\right)}{\sqrt{e}f^2} \end{aligned}$$

Mathematica [A] time = 0.27, size = 124, normalized size = 0.97

$$\frac{\log\left(\sqrt{d}\sqrt{c+dx^2}+dx\right)(2adf+bcf-2bde)}{\sqrt{d}} - \frac{2(be-af)\sqrt{cf-de}\tan^{-1}\left(\frac{x\sqrt{cf-de}}{\sqrt{e}\sqrt{c+dx^2}}\right)}{\sqrt{e}} + bfx\sqrt{c+dx^2}}{2f^2}$$

Antiderivative was successfully verified.

[In] Integrate[((a + b*x^2)*Sqrt[c + d*x^2])/(e + f*x^2), x]

[Out] (b*f*x*Sqrt[c + d*x^2] - (2*(b*e - a*f)*Sqrt[-(d*e) + c*f]*ArcTan[(Sqrt[-(d*e) + c*f]*x)/(Sqrt[e]*Sqrt[c + d*x^2])])/Sqrt[e] + ((-2*b*d*e + b*c*f + 2*a*d*f)*Log[d*x + Sqrt[d]*Sqrt[c + d*x^2]])/Sqrt[d])/(2*f^2)

IntegrateAlgebraic [A] time = 0.38, size = 151, normalized size = 1.18

$$\frac{\log\left(\sqrt{c + dx^2} - \sqrt{d}x\right)(-2adf - bcf + 2bde)}{2\sqrt{d}f^2} + \frac{(be - af)\sqrt{cf - de}\tan^{-1}\left(\frac{-fx\sqrt{c+dx^2} + \sqrt{d}e + \sqrt{d}fx^2}{\sqrt{e}\sqrt{cf-de}}\right)}{\sqrt{e}f^2} + \frac{bx\sqrt{c + dx^2}}{2f}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[((a + b*x^2)*Sqrt[c + d*x^2])/(e + f*x^2), x]

[Out] (b*x*Sqrt[c + d*x^2])/(2*f) + ((b*e - a*f)*Sqrt[-(d*e) + c*f]*ArcTan[(Sqrt[d]*e + Sqrt[d]*f*x^2 - f*x*Sqrt[c + d*x^2])/(Sqrt[e]*Sqrt[-(d*e) + c*f])])/ (Sqrt[e]*f^2) + ((2*b*d*e - b*c*f - 2*a*d*f)*Log[-(Sqrt[d]*x) + Sqrt[c + d*x^2]])/ (2*Sqrt[d]*f^2)

fricas [A] time = 4.88, size = 777, normalized size = 6.07

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^(1/2)/(f*x^2+e), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(d*x^2 + c)*b*d*f*x - (2*b*d*e - (b*c + 2*a*d)*f)*sqrt(d)*log(-2*d*x^2 - 2*sqrt(d*x^2 + c)*sqrt(d)*x - c) - (b*d*e - a*d*f)*sqrt((d*e - c

$$f)/e)*\log(((8*d^2*e^2 - 8*c*d*e*f + c^2*f^2)*x^4 + c^2*e^2 + 2*(4*c*d*e^2 - 3*c^2*e*f)*x^2 - 4*(c*e^2*x + (2*d*e^2 - c*e*f)*x^3)*\sqrt{d*x^2 + c}*\sqrt{(d*e - c*f)/e})/(f^2*x^4 + 2*e*f*x^2 + e^2)))/(d*f^2), 1/4*(2*\sqrt{d*x^2 + c})*b*d*f*x + 2*(2*b*d*e - (b*c + 2*a*d)*f)*\sqrt{-d}*\arctan(\sqrt{-d}*x/\sqrt{d*x^2 + c})) - (b*d*e - a*d*f)*\sqrt{(d*e - c*f)/e}*\log(((8*d^2*e^2 - 8*c*d*e*f + c^2*f^2)*x^4 + c^2*e^2 + 2*(4*c*d*e^2 - 3*c^2*e*f)*x^2 - 4*(c*e^2*x + (2*d*e^2 - c*e*f)*x^3)*\sqrt{d*x^2 + c}*\sqrt{(d*e - c*f)/e})/(f^2*x^4 + 2*e*f*x^2 + e^2)))/(d*f^2), 1/4*(2*\sqrt{d*x^2 + c})*b*d*f*x - 2*(b*d*e - a*d*f)*\sqrt{-d}*e*\arctan(1/2*((2*d*e - c*f)*x^2 + c*e)*\sqrt{d*x^2 + c}*\sqrt{-d}*e/(d*e - c*f)/e)/((d^2*e - c*d*f)*x^3 + (c*d*e - c^2*f)*x)) - (2*b*d*e - (b*c + 2*a*d)*f)*\sqrt{d}*\log(-2*d*x^2 - 2*\sqrt{d*x^2 + c}*\sqrt{d}*x - c))/(d*f^2), 1/2*(\sqrt{d*x^2 + c})*b*d*f*x + (2*b*d*e - (b*c + 2*a*d)*f)*\sqrt{-d})*\arctan(\sqrt{-d}*x/\sqrt{d*x^2 + c}) - (b*d*e - a*d*f)*\sqrt{-d}*e*\arctan(1/2*((2*d*e - c*f)*x^2 + c*e)*\sqrt{d*x^2 + c}*\sqrt{-d}*e/(d^2*e - c*d*f)*x^3 + (c*d*e - c^2*f)*x)))/(d*f^2)]$$

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)*(d*x^2+c)^(1/2)/(f*x^2+e),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,x);OUTPUT:

maple [B] time = 0.03, size = 1942, normalized size = 15.17

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)*(d*x^2+c)^(1/2)/(f*x^2+e),x)

[Out] $\frac{1}{2}b*x*(d*x^2+c)^{1/2}/f+1/2*b/f*c/d^{1/2}*\ln(d^{1/2}*x+(d*x^2+c)^{1/2})+1/2/(-e*f)^{1/2}*((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2}*a-1/2/(-e*f)^{1/2}/f*((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2}*b*e+1/2/f*d^{1/2}*\ln((d*(-e*f)^{1/2}/f+d*(x-(-e*f)^{1/2}/f))/d^{1/2}+((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})*a-1/2/f^2*d^{1/2}*\ln((d*(-e*f)^{1/2}/f+d*(x-(-e*f)^{1/2}/f))/d^{1/2}+((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})*b*e-1/2/(-e*f)^{1/2}/((c*f-d*e)/f)^{1/2}*\ln((2*(c*f-d*e)/f+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+2*((c*f-d*e)/f)^{1/2})*((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})/(x-(-e*f)^{1/2}/f))*c*a+1/2/(-e*f)^{1/2}/f/((c*f-d*e)/f)^{1/2}*\ln((2*(c*f-d*e)/f+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+2*((c*f-d*e)/f)^{1/2})*((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})/(x-(-e*f)^{1/2}/f))*d*e*a-1/2/(-e*f)^{1/2}/f^2/((c*f-d*e)/f)^{1/2}*\ln((2*(c*f-d*e)/f+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+2*((c*f-d*e)/f)^{1/2})*((x-(-e*f)^{1/2}/f)^2*d+2*d*(-e*f)^{1/2}/f*(x-(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})/(x-(-e*f)^{1/2}/f))*d*e^2*b-1/2/(-e*f)^{1/2}*((x+(-e*f)^{1/2}/f)^2*d-2*d*(-e*f)^{1/2}/f*(x+(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})*a+1/2/(-e*f)^{1/2}/f*((x+(-e*f)^{1/2}/f)^2*d-2*d*(-e*f)^{1/2}/f*(x+(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})*b*e+1/2/f*d^{1/2}*\ln((-d*(-e*f)^{1/2}/f+d*(x+(-e*f)^{1/2}/f))/d^{1/2}+((x+(-e*f)^{1/2}/f)^2*d-2*d*(-e*f)^{1/2}/f*(x+(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2})*a-1/2/f^2*d^{1/2}*\ln((-d*(-e*f)^{1/2}/f+d*(x+(-e*f)^{1/2}/f))/d^{1/2}+((x+(-e*f)^{1/2}/f)^2*d-2*d*(-e*f)^{1/2}/f*(x+(-e*f)^{1/2}/f)+(c*f-d*e)/f)^{1/2}))*b*e+1/2/(-e*f)^{1/2}/((c*f-d*e)/f)^{1/2}*$

```
ln((2*(c*f-d*e)/f-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+2*((c*f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+(c*f-d*e)/f)^(1/2))/(x+(-e*f)^(1/2)/f))*c*a-1/2/(-e*f)^(1/2)/f/((c*f-d*e)/f)^(1/2)*ln((2*(c*f-d*e)/f-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+2*((c*f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+(c*f-d*e)/f)^(1/2))/(x+(-e*f)^(1/2)/f))*c*b*e-1/2/(-e*f)^(1/2)/f/((c*f-d*e)/f)^(1/2)*ln((2*(c*f-d*e)/f-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+2*((c*f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+(c*f-d*e)/f)^(1/2))/(x+(-e*f)^(1/2)/f))*d*e*a+1/2/(-e*f)^(1/2)/f^2/((c*f-d*e)/f)^(1/2)*ln((2*(c*f-d*e)/f-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+2*((c*f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*d*(-e*f)^(1/2)/f*(x+(-e*f)^(1/2)/f)+(c*f-d*e)/f)^(1/2))/(x+(-e*f)^(1/2)/f))*d*e^2*b
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)\sqrt{dx^2 + c}}{fx^2 + e} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)*(d*x^2+c)^(1/2)/(f*x^2+e),x, algorithm="maxima")
```

```
[Out] integrate((b*x^2 + a)*sqrt(d*x^2 + c)/(f*x^2 + e), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)\sqrt{dx^2 + c}}{fx^2 + e} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a + b*x^2)*(c + d*x^2)^(1/2))/(e + f*x^2),x)
```

```
[Out] int(((a + b*x^2)*(c + d*x^2)^(1/2))/(e + f*x^2), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)\sqrt{c + dx^2}}{e + fx^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x**2+a)*(d*x**2+c)**(1/2)/(f*x**2+e),x)
```

```
[Out] Integral((a + b*x**2)*sqrt(c + d*x**2)/(e + f*x**2), x)
```

$$3.24 \quad \int \frac{(a+bx^2)^3}{(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=304

$$\frac{b(8a^2f^2 - 8abef + 3b^2e^2) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{8df^{5/2}} - \frac{b^2x\sqrt{e+fx^2}(bc-ad)}{2d^2f} - \frac{3b^2x\sqrt{e+fx^2}(be-2af)}{8df^2} + \frac{b^2x(a+bx^2)\sqrt{e+fx^2}}{4df}$$

Rubi [A] time = 0.30, antiderivative size = 304, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 8, integrand size = 30, number of rules / integrand size = 0.267, Rules used = {545, 416, 388, 217, 206, 523, 377, 205}

$$\frac{b(8a^2f^2 - 8abef + 3b^2e^2) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{8df^{5/2}} - \frac{b^2x\sqrt{e+fx^2}(bc-ad)}{2d^2f} - \frac{3b^2x\sqrt{e+fx^2}(be-2af)}{8df^2} + \frac{b^2x(a+bx^2)\sqrt{e+fx^2}}{4df} + \frac{b(bc-ad)(be-2af) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{2d^2f^{3/2}} - \frac{(bc-ad)^3 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d^3\sqrt{de-cf}} + \frac{b(bc-ad)^2 \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d^3\sqrt{f}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^3/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] -(b^2*(b*c - a*d)*x*Sqrt[e + f*x^2])/(2*d^2*f) - (3*b^2*(b*e - 2*a*f)*x*Sqrt[e + f*x^2])/(8*d*f^2) + (b^2*x*(a + b*x^2)*Sqrt[e + f*x^2])/(4*d*f) - ((b*c - a*d)^3*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*d^3*Sqrt[d*e - c*f]) + (b*(b*c - a*d)^2*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(d^3*Sqrt[f]) + (b*(b*c - a*d)*(b*e - 2*a*f)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(2*d^2*f^(3/2)) + (b*(3*b^2*e^2 - 8*a*b*e*f + 8*a^2*f^2)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(8*d*f^(5/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 416

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(d*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q-1))/(b*(n*(p+q)+1)),

$x] + \text{Dist}[1/(b*(n*(p + q) + 1)), \text{Int}[(a + b*x^n)^p*(c + d*x^n)^{(q - 2)}*\text{Simp}[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q - 1) + 1))*x^n, x], x] /; \text{FreeQ}\{a, b, c, d, n, p\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[q, 1] \&\& \text{NeQ}[n*(p + q) + 1, 0] \&\& !\text{IGtQ}[p, 1] \&\& \text{IntBinomialQ}[a, b, c, d, n, p, q, x]$

Rule 523

$\text{Int}[(e + f*x^n)/(a + b*x^n)*\text{Sqrt}[c + d*x^n], x_Symbol] := \text{Dist}[f/b, \text{Int}[1/\text{Sqrt}[c + d*x^n], x], x] + \text{Dist}[(b*e - a*f)/b, \text{Int}[1/(a + b*x^n)*\text{Sqrt}[c + d*x^n], x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x]$

Rule 545

$\text{Int}[(c + d*x^2)^q*(e + f*x^2)^r/(a + b*x^2), x_Symbol] := \text{Dist}[d/b, \text{Int}[(c + d*x^2)^{(q - 1)}*(e + f*x^2)^r, x], x] + \text{Dist}[(b*c - a*d)/b, \text{Int}[(c + d*x^2)^{(q - 1)}*(e + f*x^2)^r/(a + b*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e, f, r\}, x \&\& \text{GtQ}[q, 1]$

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^3}{(c + dx^2)\sqrt{e + fx^2}} dx &= \frac{b \int \frac{(a+bx^2)^2}{\sqrt{e+fx^2}} dx}{d} + \frac{(-bc + ad) \int \frac{(a+bx^2)^2}{(c+dx^2)\sqrt{e+fx^2}} dx}{d} \\ &= \frac{b^2x(a + bx^2)\sqrt{e + fx^2}}{4df} - \frac{(b(bc - ad)) \int \frac{a+bx^2}{\sqrt{e+fx^2}} dx}{d^2} + \frac{(bc - ad)^2 \int \frac{a+bx^2}{(c+dx^2)\sqrt{e+fx^2}} dx}{d^2} \\ &= -\frac{b^2(bc - ad)x\sqrt{e + fx^2}}{2d^2f} - \frac{3b^2(be - 2af)x\sqrt{e + fx^2}}{8df^2} + \frac{b^2x(a + bx^2)\sqrt{e + fx^2}}{4df} + \dots \\ &= -\frac{b^2(bc - ad)x\sqrt{e + fx^2}}{2d^2f} - \frac{3b^2(be - 2af)x\sqrt{e + fx^2}}{8df^2} + \frac{b^2x(a + bx^2)\sqrt{e + fx^2}}{4df} + \dots \\ &= -\frac{b^2(bc - ad)x\sqrt{e + fx^2}}{2d^2f} - \frac{3b^2(be - 2af)x\sqrt{e + fx^2}}{8df^2} + \frac{b^2x(a + bx^2)\sqrt{e + fx^2}}{4df} + \dots \end{aligned}$$

Mathematica [A] time = 0.37, size = 265, normalized size = 0.87

$$\frac{8b(3a^2d^2 - 3abcd + b^2c^2) \tanh^{-1}\left(\frac{\sqrt{fx}}{\sqrt{e+fx^2}}\right) + 4b^2de(bc - 3ad) \tanh^{-1}\left(\frac{\sqrt{fx}}{\sqrt{e+fx^2}}\right) - 4b^2dx\sqrt{e+fx^2}(bc - 3ad) + \frac{8(ad - bc)^3 \tan^{-1}\left(\frac{x\sqrt{de - cf}}{\sqrt{c}\sqrt{e+fx^2}}\right) + 3b^3d^2e\left(e \tanh^{-1}\left(\frac{\sqrt{fx}}{\sqrt{e+fx^2}}\right) - \sqrt{fx}\sqrt{e+fx^2}\right) + 2b^3d^2x^3\sqrt{e+fx^2}}{8d^3}}{\sqrt{f}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^3/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ((-4*b^2*d*(b*c - 3*a*d)*x*Sqrt[e + f*x^2])/f + (2*b^3*d^2*x^3*Sqrt[e + f*x^2])/f + (8*(-(b*c) + a*d)^3*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*Sqrt[d*e - c*f]) + (4*b^2*d*(b*c - 3*a*d)*e*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/f^(3/2) + (8*b*(b^2*c^2 - 3*a*b*c*d + 3*a^2*d^2)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/Sqrt[f] + (3*b^3*d^2*e*(-(Sqrt[f]*x)*Sqrt[e + f*x^2]) + e*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]]))/f^(5/2))/(8*d^3)

IntegrateAlgebraic [A] time = 0.67, size = 266, normalized size = 0.88

$$\frac{\log(\sqrt{e+fx^2}-\sqrt{f}x)(-24a^2bd^2f^2+24ab^2cd^2f^2+12ab^2d^2ef-8b^3c^2f^2-4b^3c^2d^2e^2)}{8d^3f^2} + \frac{(-a^3d^3+3a^2bcd^2-3ab^2c^2d+b^3c^3)\tan^{-1}\left(\frac{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}{\sqrt{c}\sqrt{de-f}}\right)}{\sqrt{c}d^3\sqrt{de-f}} + \frac{\sqrt{e+fx^2}(12ab^2d^2fx-4b^3c^2fx-3b^3dex+2b^3d^2fx^3)}{8d^2f^2}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[(a + b*x^2)^3/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] (Sqrt[e + f*x^2]*(-3*b^3*d*e*x - 4*b^3*c*f*x + 12*a*b^2*d*f*x + 2*b^3*d*f*x^3))/(8*d^2*f^2) + ((b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*ArcTan[(c*Sqrt[f] + d*Sqrt[f]*x^2 - d*x*Sqrt[e + f*x^2])/(Sqrt[c]*Sqrt[d*e - c*f])])/(Sqrt[c]*d^3*Sqrt[d*e - c*f]) + ((-3*b^3*d^2*e^2 - 4*b^3*c*d*e*f + 12*a*b^2*d^2*e*f - 8*b^3*c^2*f^2 + 24*a*b^2*c*d*f^2 - 24*a^2*b*d^2*f^2)*Log[-(Sqrt[f]*x) + Sqrt[e + f*x^2]])/(8*d^3*f^(5/2))

fricas [A] time = 20.97, size = 1718, normalized size = 5.65

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^3/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")

[Out] [1/16*(4*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(-c*d*e + c^2*f)*f^3*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + (3*b^3*c*d^3*e^3 + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e^2*f + 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 6*a^2*b*c*d^3)*e*f^2 - 8*(b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2)*f^3)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*x^2 + e)*sqrt(f)*x - e) + 2*(2*(b^3*c*d^3*e*f^2 - b^3*c^2*d^2*f^3)*x^3 - (3*b^3*c*d^3*e^2*f + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e*f^2 - 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2)*f^3)*x)*sqrt(f*x^2 + e))/(c*d^4*e*f^3 - c^2*d^3*f^4), -1/16*(8*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(c*d*e - c^2*f)*f^3*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x)) - (3*b^3*c*d^3*e^3 + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e^2*f + 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 6*a^2*b*c*d^3)*e*f^2 - 8*(b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2)*f^3)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*x^2 + e)*sqrt(f)*x - e) - 2*(2*(b^3*c*d^3*e*f^2 - b^3*c^2*d^2*f^3)*x^3 - (3*b^3*c*d^3*e^2*f + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e*f^2 - 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2)*f^3)*x)*sqrt(f*x^2 + e))/(c*d^4*e*f^3 - c^2*d^3*f^4), 1/8*(2*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(-c*d*e + c^2*f)*f^3*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2)) - (3*b^3*c*d^3*e^3 + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e^2*f + 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 6*a^2*b*c*d^3)*e*f^2 - 8*(b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2)*f^3)*sqrt(-f)*arctan(sqrt(-f)*x/sqrt(f*x^2 + e)) + (2*(b^3*c*d^3*e*f^2 - b^3*c^2*d^2*f^3)*x^3 - (3*b^3*c*d^3*e^2*f + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e*f^2 - 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2)*f^3)*x)*sqrt(f*x^2 + e))/(c*d^4*e*f^3 - c^2*d^3*f^4), -1/8*(4*(b^3*c^3 - 3*a*b^2*c^2*d + 3*a^2*b*c*d^2 - a^3*d^3)*sqrt(c*d*e - c^2*f)*f^3*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x)) + (3*b^3*c*d^3*e^3 + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e^2*f + 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2 + 6*a^2*b*c*d^3)*e*f^2 - 8*(b^3*c^4 - 3*a*b^2*c^3*d + 3*a^2*b*c^2*d^2)*f^3)*sqrt(-f)*arctan(sqrt(-f)*x/sqrt(f*x^2 + e)) - (2*(b^3*c*d^3*e*f^2 - b^3*c^2*d^2*f^3)*x^3 - (3*b^3*c*d^3*e^2*f + (b^3*c^2*d^2 - 12*a*b^2*c*d^3)*e*f^2 - 4*(b^3*c^3*d - 3*a*b^2*c^2*d^2)*f^3)*x)*sqrt(f*x^2 + e))/(c*d^4*e*f^3 - c^2*d^3*f^4)]

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^3/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,x):;OUTPUT:
```

maple [B] time = 0.04, size = 1541, normalized size = 5.07

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^3/(d*x^2+c)/(f*x^2+e)^(1/2),x)
```

```
[Out] 1/4*b^3/d*x^3/f*(f*x^2+e)^(1/2)-3/8*b^3/d*e/f^2*x*(f*x^2+e)^(1/2)+3/8*b^3/d
*e^2/f^(5/2)*ln(x*f^(1/2)+(f*x^2+e)^(1/2))+3/2*b^2/d*x/f*(f*x^2+e)^(1/2)*a-
1/2*b^3/d^2*x/f*(f*x^2+e)^(1/2)*c-3/2*b^2/d*e/f^(3/2)*ln(x*f^(1/2)+(f*x^2+e)
^(1/2))*a+1/2*b^3/d^2*e/f^(3/2)*ln(x*f^(1/2)+(f*x^2+e)^(1/2))*c+3*b/d*a^2*
ln(x*f^(1/2)+(f*x^2+e)^(1/2))/f^(1/2)-3*b^2/d^2*a*c*ln(x*f^(1/2)+(f*x^2+e)
^(1/2))/f^(1/2)+b^3/d^3*c^2*ln(x*f^(1/2)+(f*x^2+e)^(1/2))/f^(1/2)-1/2/(-c*d)
^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d+2*f*(-c*d)^(1/2)/d*(x-1/d*(-c
*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)*((x-1/d*(-c*d)^(1/2))^2*f+2*f*(-c*d)^(1/2)
)/d*(x-1/d*(-c*d)^(1/2))-c*f-d*e)/d^(1/2))/(x-1/d*(-c*d)^(1/2)))*a^3+3/2
/d/(-c*d)^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d+2*f*(-c*d)^(1/2)/d*
(x-1/d*(-c*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)*((x-1/d*(-c*d)^(1/2))^2*f+2*f*(-c
*d)^(1/2)/d*(x-1/d*(-c*d)^(1/2))-c*f-d*e)/d^(1/2))/(x-1/d*(-c*d)^(1/2))
)*a^2*c*b-3/2/d^2/(-c*d)^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d+2*f*
(-c*d)^(1/2)/d*(x-1/d*(-c*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)*((x-1/d*(-c*d)^(
1/2))^2*f+2*f*(-c*d)^(1/2)/d*(x-1/d*(-c*d)^(1/2))-c*f-d*e)/d^(1/2))/(x-1/
d*(-c*d)^(1/2)))*a*b^2*c^2+1/2/d^3/(-c*d)^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2
*(c*f-d*e)/d+2*f*(-c*d)^(1/2)/d*(x-1/d*(-c*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)
*((x-1/d*(-c*d)^(1/2))^2*f+2*f*(-c*d)^(1/2)/d*(x-1/d*(-c*d)^(1/2))-c*f-d*e)
/d^(1/2))/(x-1/d*(-c*d)^(1/2)))*b^3*c^3+1/2/(-c*d)^(1/2)/(-c*f-d*e)/d^(
1/2)*ln((-2*(c*f-d*e)/d-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c*d)^(1/2))+2*(-c*f-d*
e)/d^(1/2)*((x+1/d*(-c*d)^(1/2))^2*f-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c*d)^(1/2)
))-c*f-d*e)/d^(1/2))/(x+1/d*(-c*d)^(1/2)))*a^3-3/2/d/(-c*d)^(1/2)/(-c*f-
d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c*d)^(1/2))+2*
(-c*f-d*e)/d^(1/2)*((x+1/d*(-c*d)^(1/2))^2*f-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c
*d)^(1/2))-c*f-d*e)/d^(1/2))/(x+1/d*(-c*d)^(1/2)))*a^2*c*b+3/2/d^2/(-c*d)
^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c
*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)*((x+1/d*(-c*d)^(1/2))^2*f-2*f*(-c*d)^(1/2)
)/d*(x+1/d*(-c*d)^(1/2))-c*f-d*e)/d^(1/2))/(x+1/d*(-c*d)^(1/2)))*a*b^2*c
^2-1/2/d^3/(-c*d)^(1/2)/(-c*f-d*e)/d^(1/2)*ln((-2*(c*f-d*e)/d-2*f*(-c*d)
^(1/2)/d*(x+1/d*(-c*d)^(1/2))+2*(-c*f-d*e)/d^(1/2)*((x+1/d*(-c*d)^(1/2))^
2*f-2*f*(-c*d)^(1/2)/d*(x+1/d*(-c*d)^(1/2))-c*f-d*e)/d^(1/2))/(x+1/d*(-c*
d)^(1/2)))*b^3*c^3
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^3}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^3/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((b*x^2 + a)^3/((d*x^2 + c)*sqrt(f*x^2 + e)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(bx^2 + a)^3}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*x^2)^3/((c + d*x^2)*(e + f*x^2)^(1/2)), x)`

[Out] `int((a + b*x^2)^3/((c + d*x^2)*(e + f*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^3}{(c + dx^2)\sqrt{e + fx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*x**2+a)**3/(d*x**2+c)/(f*x**2+e)**(1/2), x)`

[Out] `Integral((a + b*x**2)**3/((c + d*x**2)*sqrt(e + f*x**2)), x)`

$$3.25 \quad \int \frac{(a+bx^2)^2}{(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=166

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d^2\sqrt{de-cf}} - \frac{b(bc-ad) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d^2\sqrt{f}} - \frac{b(be-2af) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{2df^{3/2}} + \frac{b^2x\sqrt{e+fx^2}}{2df}$$

Rubi [A] time = 0.11, antiderivative size = 166, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.233$, Rules used = {545, 388, 217, 206, 523, 377, 205}

$$\frac{(bc-ad)^2 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d^2\sqrt{de-cf}} - \frac{b(bc-ad) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d^2\sqrt{f}} - \frac{b(be-2af) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{2df^{3/2}} + \frac{b^2x\sqrt{e+fx^2}}{2df}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)^2/((c + d*x^2)*Sqrt[e + f*x^2]), x]

[Out] (b^2*x*Sqrt[e + f*x^2])/(2*d*f) + ((b*c - a*d)^2*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*d^2*Sqrt[d*e - c*f]) - (b*(b*c - a*d)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(d^2*Sqrt[f]) - (b*(b*e - 2*a*f)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(2*d*f^(3/2))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 388

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Simp[(d*x*(a + b*x^n)^(p+1))/(b*(n*(p+1)+1)), x] - Dist[(a*d - b*c*(n*(p+1)+1))/(b*(n*(p+1)+1)), Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p+1)+1, 0]

Rule 523

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] :> Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d

, e, f, n}, x]

Rule 545

Int[(((c_) + (d_.)*(x_)^2)^(q_)*((e_) + (f_.)*(x_)^2)^(r_))/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[d/b, Int[(c + d*x^2)^(q - 1)*(e + f*x^2)^r, x], x] + Dist[(b*c - a*d)/b, Int[((c + d*x^2)^(q - 1)*(e + f*x^2)^r)/(a + b*x^2), x], x] /; FreeQ[{a, b, c, d, e, f, r}, x] && GtQ[q, 1]

Rubi steps

$$\begin{aligned} \int \frac{(a + bx^2)^2}{(c + dx^2)\sqrt{e + fx^2}} dx &= \frac{b \int \frac{a+bx^2}{\sqrt{e+fx^2}} dx}{d} + \frac{(-bc + ad) \int \frac{a+bx^2}{(c+dx^2)\sqrt{e+fx^2}} dx}{d} \\ &= \frac{b^2x\sqrt{e + fx^2}}{2df} - \frac{(b(bc - ad)) \int \frac{1}{\sqrt{e+fx^2}} dx}{d^2} + \frac{(bc - ad)^2 \int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx}{d^2} - \frac{(b(bc - ad)) \int \frac{1}{\sqrt{e+fx^2}} dx}{d^2} \\ &= \frac{b^2x\sqrt{e + fx^2}}{2df} - \frac{(b(bc - ad)) \text{Subst}\left(\int \frac{1}{1-fx^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{d^2} + \frac{(bc - ad)^2 \text{Subst}\left(\int \frac{1}{c-dx^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{d^2} \\ &= \frac{b^2x\sqrt{e + fx^2}}{2df} + \frac{(bc - ad)^2 \tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d^2\sqrt{de-cf}} - \frac{b(bc - ad) \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d^2\sqrt{f}} - \frac{b(bc - ad)}{d^2\sqrt{f}} \end{aligned}$$

Mathematica [A] time = 0.34, size = 150, normalized size = 0.90

$$\frac{2f(bc-ad)^2 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right) + b^2\sqrt{c}dx\sqrt{e+fx^2}\sqrt{de-cf}}{\sqrt{c}f\sqrt{de-cf}} - \frac{b \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)(-4adf+2bcf+bde)}{f^{3/2}}}{2d^2}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)^2/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ((b^2*Sqrt[c]*d*Sqrt[d*e - c*f]*x*Sqrt[e + f*x^2] + 2*(b*c - a*d)^2*f*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*f*Sqrt[d*e - c*f]) - (b*(b*d*e + 2*b*c*f - 4*a*d*f)*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/f^(3/2))/(2*d^2)

IntegrateAlgebraic [A] time = 0.36, size = 175, normalized size = 1.05

$$\frac{(-a^2d^2 + 2abcd - b^2c^2) \tan^{-1}\left(\frac{c\sqrt{f-dx}\sqrt{e+fx^2}+d\sqrt{f}x^2}{\sqrt{c}\sqrt{de-cf}}\right) + \log(\sqrt{e+fx^2} - \sqrt{f}x)(-4abdf + 2b^2cf + b^2de)}{\sqrt{c}d^2\sqrt{de-cf}} + \frac{b^2x\sqrt{e+fx^2}}{2d^2f^{3/2}} + \frac{b^2x\sqrt{e+fx^2}}{2df}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[(a + b*x^2)^2/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] (b^2*x*Sqrt[e + f*x^2])/(2*d*f) + (((-b^2*c^2) + 2*a*b*c*d - a^2*d^2)*ArcTan[(c*Sqrt[f] + d*Sqrt[f]*x^2 - d*x*Sqrt[e + f*x^2])/(Sqrt[c]*Sqrt[d*e - c*f])])/(Sqrt[c]*d^2*Sqrt[d*e - c*f]) + ((b^2*d*e + 2*b^2*c*f - 4*a*b*d*f)*Log[-(Sqrt[f]*x) + Sqrt[e + f*x^2]])/(2*d^2*f^(3/2))

fricas [A] time = 6.22, size = 1111, normalized size = 6.69

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")
[Out] [-1/4*((b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(-c*d*e + c^2*f)*f^2*log(((d^2*e
^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 -
4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x
^4 + 2*c*d*x^2 + c^2)) - 2*(b^2*c*d^2*e*f - b^2*c^2*d*f^2)*sqrt(f*x^2 + e)*
x + (b^2*c*d^2*e^2 + (b^2*c^2*d - 4*a*b*c*d^2)*e*f - 2*(b^2*c^3 - 2*a*b*c^2
*d)*f^2)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*x^2 + e)*sqrt(f)*x - e)/(c*d^3*e*
f^2 - c^2*d^2*f^3), 1/4*(2*(b^2*c^2 - 2*a*b*c*d + a^2*d^2)*sqrt(c*d*e - c^2
*f)*f^2*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2
+ e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x)) + 2*(b^2*c*d^2*e*f
- b^2*c^2*d*f^2)*sqrt(f*x^2 + e)*x - (b^2*c*d^2*e^2 + (b^2*c^2*d - 4*a*b*c
*d^2)*e*f - 2*(b^2*c^3 - 2*a*b*c^2*d)*f^2)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*
x^2 + e)*sqrt(f)*x - e)/(c*d^3*e*f^2 - c^2*d^2*f^3), -1/4*((b^2*c^2 - 2*a*
b*c*d + a^2*d^2)*sqrt(-c*d*e + c^2*f)*f^2*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2
*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3
- c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2))
- 2*(b^2*c*d^2*e*f - b^2*c^2*d*f^2)*sqrt(f*x^2 + e)*x - 2*(b^2*c*d^2*e^2 +
(b^2*c^2*d - 4*a*b*c*d^2)*e*f - 2*(b^2*c^3 - 2*a*b*c^2*d)*f^2)*sqrt(-f)*ar
ctan(sqrt(-f)*x/sqrt(f*x^2 + e))/(c*d^3*e*f^2 - c^2*d^2*f^3), 1/2*((b^2*c^
2 - 2*a*b*c*d + a^2*d^2)*sqrt(c*d*e - c^2*f)*f^2*arctan(1/2*sqrt(c*d*e - c^
2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (
c*d*e^2 - c^2*e*f)*x)) + (b^2*c*d^2*e*f - b^2*c^2*d*f^2)*sqrt(f*x^2 + e)*x
+ (b^2*c*d^2*e^2 + (b^2*c^2*d - 4*a*b*c*d^2)*e*f - 2*(b^2*c^3 - 2*a*b*c^2*d
)*f^2)*sqrt(-f)*arctan(sqrt(-f)*x/sqrt(f*x^2 + e))/(c*d^3*e*f^2 - c^2*d^2*
f^3)]
```

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,x):;OUTPUT:
```

maple [B] time = 0.02, size = 1052, normalized size = 6.34

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x)
[Out] 1/2*b^2*x*(f*x^2+e)^(1/2)/d/f-1/2*b^2/d*e/f^(3/2)*ln(f^(1/2)*x+(f*x^2+e)^(1
/2))+2*b/d*a*ln(f^(1/2)*x+(f*x^2+e)^(1/2))/f^(1/2)-b^2/d^2*c*ln(f^(1/2)*x+(
f*x^2+e)^(1/2))/f^(1/2)-1/2/(-c*d)^(1/2)/(-c*f-d*e)/d)^(1/2)*ln((2*(-c*d)^(
1/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^(1/2)*((x-(-c*d
)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x
-(-c*d)^(1/2)/d)*a^2+1/d/(-c*d)^(1/2)/(-c*f-d*e)/d)^(1/2)*ln((2*(-c*d)^(1
/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^(1/2)*((x-(-c*d
)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x-
(-c*d)^(1/2)/d)*a*b*c-1/2/d^2/(-c*d)^(1/2)/(-c*f-d*e)/d)^(1/2)*ln((2*(-c*d
)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^(1/2)*((x-(-c
*d)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/
(x-(-c*d)^(1/2)/d)*b^2*c^2+1/2/(-c*d)^(1/2)/(-c*f-d*e)/d)^(1/2)*ln((-2*(-
c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^(1/2)*((x+
(-c*d)^(1/2)/d)^2*f-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2)
```

$$\frac{((x+(-c*d)^{(1/2)}/d))*a^{-2}-1/d/(-c*d)^{(1/2)}/(-(c*f-d*e)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*f-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-(c*f-d*e)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*a*b*c+1/2/d^2/(-c*d)^{(1/2)}/(-(c*f-d*e)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*f-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-(c*f-d*e)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))*b^2*c^2$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(bx^2 + a)^2}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)^2/((d*x^2 + c)*sqrt(f*x^2 + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(bx^2 + a)^2}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)^2/((c + d*x^2)*(e + f*x^2)^(1/2)),x)

[Out] int((a + b*x^2)^2/((c + d*x^2)*(e + f*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + bx^2)^2}{(c + dx^2)\sqrt{e + fx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)**2/(d*x**2+c)/(f*x**2+e)**(1/2),x)

[Out] Integral((a + b*x**2)**2/((c + d*x**2)*sqrt(e + f*x**2)), x)

$$3.26 \quad \int \frac{a+bx^2}{(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=91

$$\frac{b \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d\sqrt{f}} - \frac{(bc-ad) \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d\sqrt{de-cf}}$$

Rubi [A] time = 0.05, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$, Rules used = {523, 217, 206, 377, 205}

$$\frac{b \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d\sqrt{f}} - \frac{(bc-ad) \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] Int[(a + b*x^2)/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] -(((b*c - a*d)*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*d*Sqrt[d*e - c*f])) + (b*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/(d*Sqrt[f])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 523

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] :> Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{a + bx^2}{(c + dx^2)\sqrt{e + fx^2}} dx &= \frac{b \int \frac{1}{\sqrt{e+fx^2}} dx}{d} + \frac{(-bc + ad) \int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx}{d} \\ &= \frac{b \operatorname{Subst}\left(\int \frac{1}{1-fx^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{d} + \frac{(-bc + ad) \operatorname{Subst}\left(\int \frac{1}{c-(-de+cf)x^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{d} \\ &= -\frac{(bc - ad) \tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}d\sqrt{de-cf}} + \frac{b \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{d\sqrt{f}} \end{aligned}$$

Mathematica [A] time = 0.12, size = 88, normalized size = 0.97

$$\frac{(ad-bc) \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}} + \frac{b \tanh^{-1}\left(\frac{\sqrt{f}x}{\sqrt{e+fx^2}}\right)}{\sqrt{f}}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] (((-(b*c) + a*d)*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*Sqrt[d*e - c*f]) + (b*ArcTanh[(Sqrt[f]*x)/Sqrt[e + f*x^2]])/Sqrt[f])/d

IntegrateAlgebraic [A] time = 0.25, size = 114, normalized size = 1.25

$$\frac{(bc - ad) \tan^{-1}\left(\frac{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}{\sqrt{c}\sqrt{de-cf}}\right)}{\sqrt{c}d\sqrt{de-cf}} - \frac{b \log(\sqrt{e + fx^2} - \sqrt{f}x)}{d\sqrt{f}}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[(a + b*x^2)/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ((b*c - a*d)*ArcTan[(c*Sqrt[f] + d*Sqrt[f]*x^2 - d*x*Sqrt[e + f*x^2])/(Sqrt[c]*Sqrt[d*e - c*f])])/(Sqrt[c]*d*Sqrt[d*e - c*f]) - (b*Log[-(Sqrt[f]*x) + Sqrt[e + f*x^2]])/(d*Sqrt[f])

fricas [B] time = 4.14, size = 737, normalized size = 8.10

$\frac{c\sqrt{de-cf} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right) + (bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right)}{4(d^2-cf)^2} - \frac{(bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right) + (bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right)}{2(d^2-cf)^2} - \frac{(bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right) + (bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right)}{2(d^2-cf)^2} - \frac{(bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right) + (bc-ad)\sqrt{f} \log\left(\frac{(c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2)\sqrt{c}\sqrt{de-cf}}{c\sqrt{f}-dx\sqrt{e+fx^2}+d\sqrt{f}x^2}\right)}{2(d^2-cf)^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(-c*d*e + c^2*f)*(b*c - a*d)*f*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2)) + 2*(b*c*d*e - b*c^2*f)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*x^2 + e)*sqrt(f)*x - e)/(c*d^2*e*f - c^2*d*f^2), -1/2*(sqrt(c*d*e - c^2*f)*(b*c - a*d)*f*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x)) - (b*c*d*e - b*c^2*f)*sqrt(f)*log(-2*f*x^2 - 2*sqrt(f*x^2 + e)*sqrt(f)*x - e)/(c*d^2*e*f - c^2*d*f^2), 1/4*(sqrt(-c*d*e + c^2*f)*(b*c - a*d)*f*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2)) -

$4*(b*c*d*e - b*c^2*f)*sqrt(-f)*arctan(sqrt(-f)*x/sqrt(f*x^2 + e))/(c*d^2*e*f - c^2*d*f^2), -1/2*(sqrt(c*d*e - c^2*f)*(b*c - a*d)*f*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x)) + 2*(b*c*d*e - b*c^2*f)*sqrt(-f)*arctan(sqrt(-f)*x/sqrt(f*x^2 + e))/(c*d^2*e*f - c^2*d*f^2]$

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x):;OUTPUT:

maple [B] time = 0.01, size = 646, normalized size = 7.10

$$\frac{d \ln\left(\frac{\sqrt{2c-d}\sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}}}{\frac{2c-d}{d}}\right)}{2\sqrt{cd}\sqrt{\frac{2c-d}{d}}} + \frac{d \ln\left(\frac{\sqrt{2c-d}\sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}}}{\frac{2c-d}{d}}\right)}{2\sqrt{cd}\sqrt{\frac{2c-d}{d}}} + \frac{bc \ln\left(\frac{\sqrt{2c-d}\sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}}}{\frac{2c-d}{d}}\right)}{2\sqrt{cd}\sqrt{\frac{2c-d}{d}}} - \frac{bc \ln\left(\frac{\sqrt{2c-d}\sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}} \sqrt{\frac{2ax+2cx+d}{2c-d}}}{\frac{2c-d}{d}}\right)}{2\sqrt{cd}\sqrt{\frac{2c-d}{d}}} + \frac{b \ln(\sqrt{f}x + \sqrt{fx^2+e})}{d\sqrt{f}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x)

[Out] b/d*ln(f^(1/2)*x+(f*x^2+e)^(1/2))/f^(1/2)-1/2/(-c*d)^(1/2)/(-(c*f-d*e)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x-(-c*d)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x-(-c*d)^(1/2)/d)*a+1/2/(-c*d)^(1/2)/d/(-(c*f-d*e)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x-(-c*d)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x-(-c*d)^(1/2)/d))*b*c+1/2/(-c*d)^(1/2)/(-(c*f-d*e)/d)^(1/2)*ln((-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x+(-c*d)^(1/2)/d)^2*f-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x+(-c*d)^(1/2)/d))*a-1/2/(-c*d)^(1/2)/d/(-(c*f-d*e)/d)^(1/2)*ln((-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x+(-c*d)^(1/2)/d)^2*f-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x+(-c*d)^(1/2)/d))*b*c

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + a}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)/((d*x^2 + c)*sqrt(f*x^2 + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{bx^2 + a}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)/((c + d*x^2)*(e + f*x^2)^(1/2)),x)

[Out] int((a + b*x^2)/((c + d*x^2)*(e + f*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{a + bx^2}{(c + dx^2)\sqrt{e + fx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)/(d*x**2+c)/(f*x**2+e)**(1/2), x)

[Out] Integral((a + b*x**2)/((c + d*x**2)*sqrt(e + f*x**2)), x)

$$3.27 \quad \int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=49

$$\frac{\tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}}$$

Rubi [A] time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {377, 205}

$$\frac{\tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] Int[1/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])]/(Sqrt[c]*Sqrt[d*e - c*f])

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx &= \text{Subst}\left(\int \frac{1}{c-(-de+cf)x^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right) \\ &= \frac{\tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 49, normalized size = 1.00

$$\frac{\tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])]/(Sqrt[c]*Sqrt[d*e - c*f])

IntegrateAlgebraic [B] time = 0.11, size = 103, normalized size = 2.10

$$\frac{\tan^{-1}\left(\frac{d\sqrt{f}x^2}{\sqrt{c}\sqrt{de-cf}} - \frac{dx\sqrt{e+fx^2}}{\sqrt{c}\sqrt{de-cf}} + \frac{\sqrt{c}\sqrt{f}}{\sqrt{de-cf}}\right)}{\sqrt{c}\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[1/((c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] -(ArcTan[(Sqrt[c]*Sqrt[f])/Sqrt[d*e - c*f] + (d*Sqrt[f]*x^2)/(Sqrt[c]*Sqrt[d*e - c*f]) - (d*x*Sqrt[e + f*x^2])/(Sqrt[c]*Sqrt[d*e - c*f])]/(Sqrt[c]*Sqrt[d*e - c*f]))

fricas [B] time = 1.56, size = 241, normalized size = 4.92

$$\left[\frac{\sqrt{-cde + c^2f} \log\left(\frac{(d^2e^2 - 8cdef + 8c^2f^2)x^4 + c^2e^2 - 2(3cde^2 - 4c^2ef)x^2 - 4((de - 2cf)x^3 - cex)\sqrt{-cde + c^2f}\sqrt{fx^2 + e}}{d^2x^4 + 2cdx^2 + c^2}\right)}{4(cde - c^2f)}, \frac{\arctan\left(\frac{\sqrt{cde - c^2f}((de - 2cf)x^2 - ce)\sqrt{fx^2 + e}}{2((cdef - c^2f^2)x^3 + (cde^2 - c^2ef)x)}\right)}{2\sqrt{cde - c^2f}} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")

[Out] [-1/4*sqrt(-c*d*e + c^2*f)*log(((d^2*e^2 - 8*c*d*e*f + 8*c^2*f^2)*x^4 + c^2*e^2 - 2*(3*c*d*e^2 - 4*c^2*e*f)*x^2 - 4*((d*e - 2*c*f)*x^3 - c*e*x)*sqrt(-c*d*e + c^2*f)*sqrt(f*x^2 + e))/(d^2*x^4 + 2*c*d*x^2 + c^2))/(c*d*e - c^2*f), 1/2*arctan(1/2*sqrt(c*d*e - c^2*f)*((d*e - 2*c*f)*x^2 - c*e)*sqrt(f*x^2 + e)/((c*d*e*f - c^2*f^2)*x^3 + (c*d*e^2 - c^2*e*f)*x))/sqrt(c*d*e - c^2*f)]

giac [A] time = 0.36, size = 74, normalized size = 1.51

$$\frac{\sqrt{f} \arctan\left(\frac{(\sqrt{f}x - \sqrt{fx^2 + e})^2 d + 2cf - de}{2\sqrt{-c^2f^2 + cdfe}}\right)}{\sqrt{-c^2f^2 + cdfe}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")

[Out] -sqrt(f)*arctan(1/2*((sqrt(f)*x - sqrt(f*x^2 + e))^2*d + 2*c*f - d*e)/sqrt(-c^2*f^2 + c*d*f*e))/sqrt(-c^2*f^2 + c*d*f*e)

maple [B] time = 0.01, size = 306, normalized size = 6.24

$$\frac{\ln\left(\frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)f - \frac{2(cf-de)}{d} + 2\sqrt{\frac{cf-de}{d}}\sqrt{\left(x - \frac{\sqrt{-cd}}{d}\right)^2 f + \frac{2\sqrt{-cd}\left(x - \frac{\sqrt{-cd}}{d}\right)f - cf-de}{d}}}{x - \frac{\sqrt{-cd}}{d}}\right)}{2\sqrt{-cd}\sqrt{\frac{cf-de}{d}}}\right) + \frac{\ln\left(\frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)f - \frac{2(cf-de)}{d} + 2\sqrt{\frac{cf-de}{d}}\sqrt{\left(x + \frac{\sqrt{-cd}}{d}\right)^2 f - \frac{2\sqrt{-cd}\left(x + \frac{\sqrt{-cd}}{d}\right)f - cf-de}{d}}}{x + \frac{\sqrt{-cd}}{d}}\right)}{2\sqrt{-cd}\sqrt{\frac{cf-de}{d}}}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*x^2+c)/(f*x^2+e)^(1/2),x)

[Out] -1/2/(-c*d)^(1/2)/(-(c*f-d*e)/d)^(1/2)*ln((2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x-(-c*d)^(1/2)/d)^2*f+2*(-c*d)^(1/2)*(x-(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x-(-c*d)^(1/2)/d)+1/2/(-c*d)^(1/2)/(-(c*f-d*e)/d)^(1/2)*ln((-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-2*(c*f-d*e)/d+2*(-(c*f-d*e)/d)^(1/2)*((x+(-c*d)^(1/2)/d)^2*f-2*(-c*d)^(1/2)*(x+(-c*d)^(1/2)/d)/d*f-(c*f-d*e)/d)^(1/2))/(x+(-c*d)^(1/2)/d)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((d*x^2 + c)*sqrt(f*x^2 + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\left\{ \begin{array}{ll} \frac{\operatorname{atan}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{fx^2+e}}\right)}{\sqrt{-c(cf-de)}} & \text{if } 0 < de - cf \\ \frac{\ln\left(\frac{\sqrt{c(fx^2+e)+x}\sqrt{cf-de}}{\sqrt{c(fx^2+e)-x}\sqrt{cf-de}}\right)}{2\sqrt{c(cf-de)}} & \text{if } de - cf < 0 \\ \int \frac{1}{(dx^2+c)\sqrt{fx^2+e}} dx & \text{if } de - cf \notin \mathbb{R} \vee cf = de \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c + d*x^2)*(e + f*x^2)^(1/2)),x)

[Out] piecewise(0 < - c*f + d*e, atan((x*(- c*f + d*e)^(1/2))/(c^(1/2)*(e + f*x^2)^(1/2)))/(-c*(c*f - d*e)^(1/2), - c*f + d*e < 0, log(((c*(e + f*x^2)^(1/2) + x*(c*f - d*e)^(1/2))/((c*(e + f*x^2)^(1/2) - x*(c*f - d*e)^(1/2)))/(2*(c*(c*f - d*e)^(1/2))), ~in(- c*f + d*e, 'real') | c*f == d*e, int(1/((c + d*x^2)*(e + f*x^2)^(1/2)), x))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c + dx^2)\sqrt{e + fx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*x**2+c)/(f*x**2+e)**(1/2),x)

[Out] Integral(1/((c + d*x**2)*sqrt(e + f*x**2)), x)

$$3.28 \quad \int \frac{1}{(a+bx^2)(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=122

$$\frac{b \tan^{-1}\left(\frac{x\sqrt{be-af}}{\sqrt{a}\sqrt{e+fx^2}}\right)}{\sqrt{a}(bc-ad)\sqrt{be-af}} - \frac{d \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}(bc-ad)\sqrt{de-cf}}$$

Rubi [A] time = 0.11, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {532, 377, 205}

$$\frac{b \tan^{-1}\left(\frac{x\sqrt{be-af}}{\sqrt{a}\sqrt{e+fx^2}}\right)}{\sqrt{a}(bc-ad)\sqrt{be-af}} - \frac{d \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}(bc-ad)\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)*(c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] (b*ArcTan[(Sqrt[b*e - a*f]*x)/(Sqrt[a]*Sqrt[e + f*x^2])]/(Sqrt[a]*(b*c - a*d)*Sqrt[b*e - a*f]) - (d*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])]/(Sqrt[c]*(b*c - a*d)*Sqrt[d*e - c*f]))

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 532

Int[1/(((a_) + (b_.)*(x_)^2)*((c_) + (d_.)*(x_)^2)*Sqrt[(e_) + (f_.)*(x_)^2]), x_Symbol] := Dist[b/(b*c - a*d), Int[1/((a + b*x^2)*Sqrt[e + f*x^2]), x], x] - Dist[d/(b*c - a*d), Int[1/((c + d*x^2)*Sqrt[e + f*x^2]), x], x] /; FreeQ[{a, b, c, d, e, f}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a+bx^2)(c+dx^2)\sqrt{e+fx^2}} dx &= \frac{b \int \frac{1}{(a+bx^2)\sqrt{e+fx^2}} dx}{bc-ad} - \frac{d \int \frac{1}{(c+dx^2)\sqrt{e+fx^2}} dx}{bc-ad} \\ &= \frac{b \text{Subst}\left(\int \frac{1}{a-(-be+af)x^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{bc-ad} - \frac{d \text{Subst}\left(\int \frac{1}{c-(-de+cf)x^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{bc-ad} \\ &= \frac{b \tan^{-1}\left(\frac{\sqrt{be-af}x}{\sqrt{a}\sqrt{e+fx^2}}\right)}{\sqrt{a}(bc-ad)\sqrt{be-af}} - \frac{d \tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}(bc-ad)\sqrt{de-cf}} \end{aligned}$$

Mathematica [A] time = 0.21, size = 113, normalized size = 0.93

$$\frac{b \tan^{-1}\left(\frac{x\sqrt{be-af}}{\sqrt{a}\sqrt{e+fx^2}}\right)}{\sqrt{a}\sqrt{be-af}} - \frac{d \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}\sqrt{de-cf}}$$

$$bc - ad$$

Antiderivative was successfully verified.

[In] Integrate[1/((a + b*x^2)*(c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] ((b*ArcTan[(Sqrt[b*e - a*f]*x)/(Sqrt[a]*Sqrt[e + f*x^2])])/(Sqrt[a]*Sqrt[b*e - a*f]) - (d*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*Sqrt[d*e - c*f]))/(b*c - a*d)

IntegrateAlgebraic [A] time = 0.43, size = 239, normalized size = 1.96

$$\frac{b \tan^{-1}\left(\frac{b\sqrt{f}x^2}{\sqrt{a}\sqrt{be-af}} - \frac{bx\sqrt{e+fx^2}}{\sqrt{a}\sqrt{be-af}} + \frac{\sqrt{a}\sqrt{f}}{\sqrt{be-af}}\right)}{\sqrt{a}(bc - ad)\sqrt{be - af}} - \frac{d\sqrt{de - cf} \tan^{-1}\left(\frac{d\sqrt{f}x^2}{\sqrt{c}\sqrt{de-cf}} - \frac{dx\sqrt{e+fx^2}}{\sqrt{c}\sqrt{de-cf}} + \frac{\sqrt{c}\sqrt{f}}{\sqrt{de-cf}}\right)}{\sqrt{c}(bc - ad)(cf - de)}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[1/((a + b*x^2)*(c + d*x^2)*Sqrt[e + f*x^2]),x]

[Out] -((b*ArcTan[(Sqrt[a]*Sqrt[f])/Sqrt[b*e - a*f] + (b*Sqrt[f]*x^2)/(Sqrt[a]*Sqrt[b*e - a*f]) - (b*x*Sqrt[e + f*x^2])/(Sqrt[a]*Sqrt[b*e - a*f])])/(Sqrt[a]*(b*c - a*d)*Sqrt[b*e - a*f])) - (d*Sqrt[d*e - c*f]*ArcTan[(Sqrt[c]*Sqrt[f])/Sqrt[d*e - c*f] + (d*Sqrt[f]*x^2)/(Sqrt[c]*Sqrt[d*e - c*f]) - (d*x*Sqrt[e + f*x^2])/(Sqrt[c]*Sqrt[d*e - c*f])])/(Sqrt[c]*(b*c - a*d)*(-(d*e) + c*f))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [A] time = 0.52, size = 173, normalized size = 1.42

$$-f^{\frac{3}{2}} \left(\frac{b \arctan\left(\frac{(\sqrt{f}x - \sqrt{fx^2 + e})^2 b + 2af - be}{2\sqrt{-a^2 f^2 + abfe}}\right)}{\sqrt{-a^2 f^2 + abfe}(bcf - adf)} - \frac{d \arctan\left(\frac{(\sqrt{f}x - \sqrt{fx^2 + e})^2 d + 2cf - de}{2\sqrt{-c^2 f^2 + cdfe}}\right)}{\sqrt{-c^2 f^2 + cdfe}(bcf - adf)} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")

[Out] -f^(3/2)*(b*arctan(1/2*((sqrt(f)*x - sqrt(f*x^2 + e))^2*b + 2*a*f - b*e)/sqrt(-a^2*f^2 + a*b*f*e))/(sqrt(-a^2*f^2 + a*b*f*e)*(b*c*f - a*d*f)) - d*arctan(1/2*((sqrt(f)*x - sqrt(f*x^2 + e))^2*d + 2*c*f - d*e)/sqrt(-c^2*f^2 + c*d*f*e))/(sqrt(-c^2*f^2 + c*d*f*e)*(b*c*f - a*d*f)))

maple [B] time = 0.03, size = 782, normalized size = 6.41

$$\frac{b^2 d \ln\left(\frac{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + 2\sqrt{\frac{af}{b}}\sqrt{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + \frac{af}{b}}}{x\sqrt{\frac{af}{b}}}\right)}{2\sqrt{-ab}(\sqrt{-cd}b + \sqrt{-ab}d)(-\sqrt{-cd}b + \sqrt{-ab}d)\sqrt{\frac{af}{b}}}$$

$$+ \frac{b^2 d \ln\left(\frac{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + 2\sqrt{\frac{af}{b}}\sqrt{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + \frac{af}{b}}}{x\sqrt{\frac{af}{b}}}\right)}{2\sqrt{-ab}(\sqrt{-cd}b + \sqrt{-ab}d)(-\sqrt{-cd}b + \sqrt{-ab}d)\sqrt{\frac{af}{b}}}$$

$$+ \frac{b d^2 \ln\left(\frac{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + 2\sqrt{\frac{af}{b}}\sqrt{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + \frac{af}{b}}}{x\sqrt{\frac{af}{b}}}\right)}{2(\sqrt{-cd}b + \sqrt{-ab}d)(-\sqrt{-cd}b + \sqrt{-ab}d)\sqrt{-cd}\sqrt{\frac{af}{b}}}$$

$$- \frac{b d^2 \ln\left(\frac{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + 2\sqrt{\frac{af}{b}}\sqrt{\frac{2\sqrt{a}\sqrt{-\frac{af}{b}}}{b} + \frac{2\sqrt{c}\sqrt{-\frac{cf}{d}}}{d} + \frac{af}{b}}}{x\sqrt{\frac{af}{b}}}\right)}{2(\sqrt{-cd}b + \sqrt{-ab}d)(-\sqrt{-cd}b + \sqrt{-ab}d)\sqrt{-cd}\sqrt{\frac{af}{b}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x)`

[Out] $\frac{1}{2} b d^2 / ((-a b)^{1/2} d + (-c d)^{1/2} b) / ((-a b)^{1/2} d - (-c d)^{1/2} b) / (-c d)^{1/2} / (-c f - d e) / d^{1/2} * \ln((2 * (-c d)^{1/2} * (x - (-c d)^{1/2} / d) / d * f - 2 * (c f - d e) / d + 2 * (-c f - d e) / d)^{1/2} * ((x - (-c d)^{1/2} / d)^2 * f + 2 * (-c d)^{1/2} * (x - (-c d)^{1/2} / d) / d * f - (c f - d e) / d)^{1/2} / (x - (-c d)^{1/2} / d) + 1/2 * b^2 * d / (-a b)^{1/2} / ((-a b)^{1/2} d + (-c d)^{1/2} b) / ((-a b)^{1/2} d - (-c d)^{1/2} b) / (-a f - b e) / b^{1/2} * \ln((-2 * (a f - b e) / b - 2 * f * (-a b)^{1/2} / b * (x + (-a b)^{1/2} / b) + 2 * (-a f - b e) / b)^{1/2} * ((x + (-a b)^{1/2} / b)^2 * f - 2 * f * (-a b)^{1/2} / b * (x + (-a b)^{1/2} / b) - (a f - b e) / b)^{1/2} / (x + (-a b)^{1/2} / b) - 1/2 * b * d^2 / ((-a b)^{1/2} d + (-c d)^{1/2} b) / ((-a b)^{1/2} d - (-c d)^{1/2} b) / (-c d)^{1/2} / (-c f - d e) / d^{1/2} * \ln((-2 * (-c d)^{1/2} * (x + (-c d)^{1/2} / d) / d * f - 2 * (c f - d e) / d + 2 * (-c f - d e) / d)^{1/2} * ((x + (-c d)^{1/2} / d)^2 * f - 2 * (-c d)^{1/2} * (x + (-c d)^{1/2} / d) / d * f - (c f - d e) / d)^{1/2} / (x + (-c d)^{1/2} / d) - 1/2 * b^2 * d / (-a b)^{1/2} / ((-a b)^{1/2} d + (-c d)^{1/2} b) / ((-a b)^{1/2} d - (-c d)^{1/2} b) / (-a f - b e) / b^{1/2} * \ln((-2 * (a f - b e) / b + 2 * f * (-a b)^{1/2} / b * (x - (-a b)^{1/2} / b) + 2 * (-a f - b e) / b)^{1/2} * ((x - (-a b)^{1/2} / b)^2 * f + 2 * f * (-a b)^{1/2} / b * (x - (-a b)^{1/2} / b) - (a f - b e) / b)^{1/2} / (x - (-a b)^{1/2} / b)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b x^2 + a)(d x^2 + c) \sqrt{f x^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x^2+a)/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/((b*x^2 + a)*(d*x^2 + c)*sqrt(f*x^2 + e)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(b x^2 + a)(d x^2 + c) \sqrt{f x^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((a + b*x^2)*(c + d*x^2)*(e + f*x^2)^(1/2)),x)`

[Out] `int(1/((a + b*x^2)*(c + d*x^2)*(e + f*x^2)^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + b x^2)(c + d x^2) \sqrt{e + f x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*x**2+a)/(d*x**2+c)/(f*x**2+e)**(1/2),x)`

[Out] `Integral(1/((a + b*x**2)*(c + d*x**2)*sqrt(e + f*x**2)), x)`

$$3.29 \quad \int \frac{1}{(a+bx^2)^2(c+dx^2)\sqrt{e+fx^2}} dx$$

Optimal. Leaf size=203

$$\frac{b(4a^2df - 2abcf - 3abde + b^2ce) \tan^{-1}\left(\frac{x\sqrt{be-af}}{\sqrt{a}\sqrt{e+fx^2}}\right)}{2a^{3/2}(bc-ad)^2(be-af)^{3/2}} + \frac{b^2x\sqrt{e+fx^2}}{2a(a+bx^2)(bc-ad)(be-af)} + \frac{d^2 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}(bc-ad)^2\sqrt{de-cf}}$$

Rubi [A] time = 0.27, antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {546, 377, 205, 527, 12}

$$\frac{b(4a^2df - 2abcf - 3abde + b^2ce) \tan^{-1}\left(\frac{x\sqrt{be-af}}{\sqrt{a}\sqrt{e+fx^2}}\right)}{2a^{3/2}(bc-ad)^2(be-af)^{3/2}} + \frac{b^2x\sqrt{e+fx^2}}{2a(a+bx^2)(bc-ad)(be-af)} + \frac{d^2 \tan^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{c}\sqrt{e+fx^2}}\right)}{\sqrt{c}(bc-ad)^2\sqrt{de-cf}}$$

Antiderivative was successfully verified.

[In] Int[1/((a + b*x^2)^2*(c + d*x^2)*Sqrt[e + f*x^2]), x]

[Out] (b^2*x*Sqrt[e + f*x^2])/(2*a*(b*c - a*d)*(b*e - a*f)*(a + b*x^2)) + (b*(b^2*c*e - 3*a*b*d*e - 2*a*b*c*f + 4*a^2*d*f)*ArcTan[(Sqrt[b*e - a*f]*x)/(Sqrt[a]*Sqrt[e + f*x^2])])/(2*a^(3/2)*(b*c - a*d)^2*(b*e - a*f)^(3/2)) + (d^2*ArcTan[(Sqrt[d*e - c*f]*x)/(Sqrt[c]*Sqrt[e + f*x^2])])/(Sqrt[c]*(b*c - a*d)^2*Sqrt[d*e - c*f])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 205

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]*ArcTan[x/Rt[a/b, 2]])/a, x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 377

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 527

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 546

Int[((c_) + (d_.)*(x_)^2)^(q_)*((e_) + (f_.)*(x_)^2)^(r_)/((a_) + (b_.)*(x_)^2), x_Symbol] := Dist[b^2/(b*c - a*d)^2, Int[((c + d*x^2)^(q+2)*(e + f*x^2)^r)/(a + b*x^2), x], x] - Dist[d/(b*c - a*d)^2, Int[(c + d*x^2)^q*(e + f*x^2)^r*(2*b*c - a*d + b*d*x^2), x], x] /; FreeQ[{a, b, c, d, e, f, r}, x] && LtQ[q, -1]

Rubi steps

$$\int \frac{1}{(a + bx^2)^2 (c + dx^2) \sqrt{e + fx^2}} dx = -\frac{b \int \frac{-bc+2ad+bdx^2}{(a+bx^2)^2 \sqrt{e+fx^2}} dx}{(bc - ad)^2} + \frac{d^2 \int \frac{1}{(c+dx^2) \sqrt{e+fx^2}} dx}{(bc - ad)^2}$$

$$= \frac{b^2 x \sqrt{e + fx^2}}{2a(bc - ad)(be - af)(a + bx^2)} + \frac{d^2 \text{Subst}\left(\int \frac{1}{c - (-de+cf)x^2} dx, x, \frac{x}{\sqrt{e+fx^2}}\right)}{(bc - ad)^2} + \dots$$

$$= \frac{b^2 x \sqrt{e + fx^2}}{2a(bc - ad)(be - af)(a + bx^2)} + \frac{d^2 \tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c} \sqrt{e+fx^2}}\right)}{\sqrt{c}(bc - ad)^2 \sqrt{de - cf}} + \frac{b(b^2 ce - 3abd)}{\dots}$$

$$= \frac{b^2 x \sqrt{e + fx^2}}{2a(bc - ad)(be - af)(a + bx^2)} + \frac{d^2 \tan^{-1}\left(\frac{\sqrt{de-cf}x}{\sqrt{c} \sqrt{e+fx^2}}\right)}{\sqrt{c}(bc - ad)^2 \sqrt{de - cf}} + \frac{b(b^2 ce - 3abd)}{\dots}$$

$$= \frac{b^2 x \sqrt{e + fx^2}}{2a(bc - ad)(be - af)(a + bx^2)} + \frac{b(b^2 ce - 3abde - 2abcf + 4a^2 df) \tan^{-1}\left(\frac{x}{\sqrt{e+fx^2}}\right)}{2a^{3/2}(bc - ad)^2 (be - af)^{3/2}}$$

Mathematica [C] time = 2.64, size = 531, normalized size = 2.62

$$\frac{bx \sqrt{e+fx^2} (bc-ad) \left(-30fx^2 \sqrt{\frac{a^2(e+fx^2)(bc-af)}{a^2(a+bx^2)^2}} - 45e \sqrt{\frac{a^2(e+fx^2)(bc-af)}{a^2(a+bx^2)^2}} + 16fx^2 \sqrt{\frac{a(e+fx^2)}{a(a+bx^2)}} \left(\frac{d^2(bc-af)}{a(a+bx^2)} \right)^{5/2} {}_2F_1\left(2, 3, \frac{7}{2}, \frac{(bc-af)x^2}{a(a+bx^2)}\right) + 16e \sqrt{\frac{a(e+fx^2)}{a(a+bx^2)}} \left(\frac{d^2(bc-af)}{a(a+bx^2)} \right)^{5/2} {}_2F_1\left(2, 3, \frac{7}{2}, \frac{(bc-af)x^2}{a(a+bx^2)}\right) + 30fx^2 \sin^{-1}\left(\sqrt{\frac{d^2(bc-af)}{a(a+bx^2)}}\right) + 45e \sin^{-1}\left(\sqrt{\frac{d^2(bc-af)}{a(a+bx^2)}}\right) \right) - 30bd \tan^{-1}\left(\frac{x \sqrt{bc-af}}{\sqrt{c} \sqrt{e+fx^2}}\right) + 30a^2 \tan^{-1}\left(\frac{x \sqrt{bc-af}}{\sqrt{c} \sqrt{e+fx^2}}\right)}{e^2 (a+bx^2)^2 \left(\frac{d^2(bc-af)}{a(a+bx^2)}\right)^{3/2} \sqrt{\frac{a(e+fx^2)}{a(a+bx^2)}}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/((a + b*x^2)^2*(c + d*x^2)*Sqrt[e + f*x^2]),x]
```

```
[Out] ((-30*b*d*ArcTan[(Sqrt[b*e - a*f])*x]/(Sqrt[a]*Sqrt[e + f*x^2]))/(Sqrt[a]*Sqrt[b*e - a*f]) + (30*d^2*ArcTan[(Sqrt[d*e - c*f])*x]/(Sqrt[c]*Sqrt[e + f*x^2]))/(Sqrt[c]*Sqrt[d*e - c*f]) + (b*(b*c - a*d)*x*Sqrt[e + f*x^2]*(-45*e*Sqrt[(a*(b*e - a*f)*x^2*(e + f*x^2))/(e^2*(a + b*x^2)^2)] - 30*f*x^2*Sqrt[(a*(b*e - a*f)*x^2*(e + f*x^2))/(e^2*(a + b*x^2)^2)] + 45*e*ArcSin[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))]] + 30*f*x^2*ArcSin[Sqrt[((b*e - a*f)*x^2)/(e*(a + b*x^2))]] + 16*e*((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(5/2)*Sqrt[(a*(e + f*x^2))/(e*(a + b*x^2))]*Hypergeometric2F1[2, 3, 7/2, ((b*e - a*f)*x^2)/(e*(a + b*x^2))] + 16*f*x^2*((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(5/2)*Sqrt[(a*(e + f*x^2))/(e*(a + b*x^2))]*Hypergeometric2F1[2, 3, 7/2, ((b*e - a*f)*x^2)/(e*(a + b*x^2))]))/(e^2*((b*e - a*f)*x^2)/(e*(a + b*x^2)))^(3/2)*(a + b*x^2)^2*Sqrt[(a*(e + f*x^2))/(e*(a + b*x^2))]))/(30*(b*c - a*d)^2)
```

IntegrateAlgebraic [B] time = 6.80, size = 2522, normalized size = 12.42

Result too large to show

Antiderivative was successfully verified.

```
[In] IntegrateAlgebraic[1/((a + b*x^2)^2*(c + d*x^2)*Sqrt[e + f*x^2]),x]
```

```
[Out] (-((e*Sqrt[f]*x^2)/a) - (f^(3/2)*x^4)/a + (e*x*Sqrt[e + f*x^2])/(2*a) + (f*x^3*Sqrt[e + f*x^2])/a)/((a + b*x^2)*(c + d*x^2)*(e + f*x^2)*(e + 2*f*x^2 - 2*Sqrt[f]*x*Sqrt[e + f*x^2])) + ((c*d*e*Sqrt[f])/(2*a*(b*c - a*d)*(d*e - c*f)) - (d^2*e*Sqrt[f]*x^2)/(2*a*(b*c - a*d)*(d*e - c*f)) + (c*d*f^(3/2)*x^2)/(a*(b*c - a*d)*(d*e - c*f)) + (d^2*e*x*Sqrt[e + f*x^2])/(2*a*(b*c - a*d)*(d*e - c*f)) - (c*d*f*x*Sqrt[e + f*x^2])/(a*(b*c - a*d)*(d*e - c*f)) + (d^2*e^2*ArcTan[(a*Sqrt[f] + b*Sqrt[f]*x^2 - b*x*Sqrt[e + f*x^2])/(Sqrt[a]*Sqrt[e + f*x^2])])/(2*a*(b*c - a*d)*(d*e - c*f)))
```

$$\begin{aligned} & [b*e - a*f]]) / (2 * \text{Sqrt}[a] * (b*c - a*d) * \text{Sqrt}[b*e - a*f] * (d*e - c*f)) + (d^3 * e \\ & ^2 * x^2 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * \text{Sqrt}[a] * c * (b * c - a * d) * \text{Sqrt}[b * e - a * f] * (d * e - c * f)) + (d \\ & ^2 * e * f * x^2 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * (b * c - a * d) * \text{Sqrt}[b * e - a * f] * (d * e - c * f)) + (d \\ & ^3 * e * f * x^4 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * c * (b * c - a * d) * \text{Sqrt}[b * e - a * f] * (d * e - c * f)) - \\ & (d^2 * e * \text{Sqrt}[f] * x * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * (b * c - a * d) * \text{Sqrt}[b * e - a * f] * (d * e - c * f)) - (d^3 * e * \text{Sqrt}[f] * x^3 * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * c * (b * c - a * d) * \text{Sqrt}[b * e - a * f] * (d * e - c * f)) / ((c + d * x^2) * (e + 2 * f * x^2 - 2 * \text{Sqrt}[f] * x * \text{Sqrt}[e + f * x^2])) + (-1/2 * (e^2 * f^(3/2)) / (a * (- (b * e) + a * f) * (- (d * e) + c * f)) - (e * f^(5/2) * x^2) / (2 * a * (- (b * e) + a * f) * (- (d * e) + c * f)) + (e * f^2 * x * \text{Sqrt}[e + f * x^2]) / (2 * a * (- (b * e) + a * f) * (- (d * e) + c * f)) - (b * e^3 * f^2 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * \text{Sqrt}[a] * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) + (\text{Sqrt}[a] * e^2 * f^3 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) - (3 * b * e^2 * f^3 * x^2 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * \text{Sqrt}[a] * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) + (3 * \text{Sqrt}[a] * e * f^4 * x^2 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) - (b * e * f^4 * x^4 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) + (\text{Sqrt}[a] * f^5 * x^4 * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / ((b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) + (b * e^2 * f^(5/2) * x * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) - (\text{Sqrt}[a] * e * f^(7/2) * x * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / ((b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) + (b * e * f^(7/2) * x^3 * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (\text{Sqrt}[a] * (b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) - (\text{Sqrt}[a] * f^(9/2) * x^3 * \text{Sqrt}[e + f * x^2] * \text{ArcTan}[(a * \text{Sqrt}[f] + b * \text{Sqrt}[f] * x^2 - b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / ((b * e - a * f)^(3/2) * (- (b * e) + a * f) * (- (d * e) + c * f)) / ((e + f * x^2) * (e + 2 * f * x^2 - 2 * \text{Sqrt}[f] * x * \text{Sqrt}[e + f * x^2])) - \text{ArcTan}[(\text{Sqrt}[a] * \text{Sqrt}[f]) / \text{Sqrt}[b * e - a * f] + (b * \text{Sqrt}[f] * x^2) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f]) - (b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (2 * a^(3/2) * c * \text{Sqrt}[b * e - a * f]) + (\text{Sqrt}[a] * d^2 * \text{ArcTan}[(\text{Sqrt}[a] * \text{Sqrt}[f]) / \text{Sqrt}[b * e - a * f] + (b * \text{Sqrt}[f] * x^2) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f]) - (b * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[a] * \text{Sqrt}[b * e - a * f])]) / (c * (b * c - a * d)^2 * \text{Sqrt}[b * e - a * f]) - (d^2 * \text{ArcTan}[(\text{Sqrt}[c] * \text{Sqrt}[f]) / \text{Sqrt}[d * e - c * f] + (d * \text{Sqrt}[f] * x^2) / (\text{Sqrt}[c] * \text{Sqrt}[d * e - c * f]) - (d * x * \text{Sqrt}[e + f * x^2]) / (\text{Sqrt}[c] * \text{Sqrt}[d * e - c * f])]) / (\text{Sqrt}[c] * (b * c - a * d)^2 * \text{Sqrt}[d * e - c * f]) \end{aligned}$$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [B] time = 26.05, size = 479, normalized size = 2.36

$$\frac{1}{2} \left(\frac{2d^2 \arctan\left(\frac{\sqrt{f}x - \sqrt{f^2+e}}{2\sqrt{cf+adf}}\right)}{\sqrt{b^2c^2f^2 - 2abcdf^2 + a^2d^2f^2}} + \frac{(2ab^2cf - 4a^2bdf - b^3ce + 3ab^2de) \arctan\left(\frac{\sqrt{f}x - \sqrt{f^2+e}}{2\sqrt{cf+adf}}\right)}{(a^2b^2c^2f^3 - 2a^3bcd^2f^2 + a^4b^2f^3 - ab^3c^2f^2e + 2a^2b^2cd^2f^2e - a^3bd^2f^2e)\sqrt{-a^2f^2 + abfe}} + \frac{2(2(\sqrt{f}x - \sqrt{f^2+e})^2 abf - (\sqrt{f}x - \sqrt{f^2+e})^2 b^2e + b^2d^2)}{(a^2bcf^3 - a^3df^3 - ab^2cf^2e + a^2bd^2f^2e)(\sqrt{f}x - \sqrt{f^2+e})^4 b + 4(\sqrt{f}x - \sqrt{f^2+e})^2 af - 2(\sqrt{f}x - \sqrt{f^2+e})^2 bc + be^2} \right) f^{\frac{1}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="giac")

[Out]
$$-1/2*(2*d^2*\arctan(1/2*((\sqrt{f}*x - \sqrt{f*x^2 + e})^2*d + 2*c*f - d*e)/\sqrt{-c^2*f^2 + c*d*f*e}))/((b^2*c^2*f^2 - 2*a*b*c*d*f^2 + a^2*d^2*f^2)*\sqrt{-c^2*f^2 + c*d*f*e}) + (2*a*b^2*c*f - 4*a^2*b*d*f - b^3*c*e + 3*a*b^2*d*e)*\arctan(1/2*((\sqrt{f}*x - \sqrt{f*x^2 + e})^2*b + 2*a*f - b*e)/\sqrt{-a^2*f^2 + a*b*f*e}))/((a^2*b^2*c^2*f^3 - 2*a^3*b*c*d*f^3 + a^4*d^2*f^3 - a*b^3*c^2*f^2*e + 2*a^2*b^2*c*d*f^2*e - a^3*b*d^2*f^2*e)*\sqrt{-a^2*f^2 + a*b*f*e}) + 2*(2*(\sqrt{f}*x - \sqrt{f*x^2 + e})^2*a*b*f - (\sqrt{f}*x - \sqrt{f*x^2 + e})^2*b^2*e + b^2*e^2)/((a^2*b*c*f^3 - a^3*d*f^3 - a*b^2*c*f^2*e + a^2*b*d*f^2*e)*((\sqrt{f}*x - \sqrt{f*x^2 + e})^4*b + 4*(\sqrt{f}*x - \sqrt{f*x^2 + e})^2*a*f - 2*(\sqrt{f}*x - \sqrt{f*x^2 + e})^2*b*e + b*e^2)))f^{(5/2)}$$

maple [B] time = 0.04, size = 1865, normalized size = 9.19

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x)

[Out]
$$-1/2*b^2*d^4/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-c*d)^{(1/2)}/(-c*f-d*e)/d)^{(1/2)}*\ln((2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^{(1/2)}*((x-(-c*d)^{(1/2)}/d)^2*f+2*(-c*d)^{(1/2)}*(x-(-c*d)^{(1/2)}/d)/d*f-(c*f-d*e)/d)^{(1/2)})/(x-(-c*d)^{(1/2)}/d))-3/4*b^3*d^3/(-a*b)^{(1/2)}/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-a*f-b*e)/b)^{(1/2)}*\ln((-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x+(-a*b)^{(1/2)}/b)^2*f-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x+(-a*b)^{(1/2)}/b))+1/4*b^4*d^2/(-a*b)^{(1/2)}/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-a*f-b*e)/b)^{(1/2)}*\ln((-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x+(-a*b)^{(1/2)}/b)^2*f-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x+(-a*b)^{(1/2)}/b))*c-1/4*b^2*d/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(a*f-b*e)/(x-(-a*b)^{(1/2)}/b))*((x-(-a*b)^{(1/2)}/b)^2*f+2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)}+1/4*b*d/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)*f*(-a*b)^{(1/2)}/(a*f-b*e)/(-a*f-b*e)/b)^{(1/2)}*\ln((2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x-(-a*b)^{(1/2)}/b)^2*f+2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x-(-a*b)^{(1/2)}/b))+1/2*b^2*d^4/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-c*d)^{(1/2)}/(-c*f-d*e)/d)^{(1/2)}*\ln((-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-2*(c*f-d*e)/d+2*(-c*f-d*e)/d)^{(1/2)}*((x+(-c*d)^{(1/2)}/d)^2*f-2*(-c*d)^{(1/2)}*(x+(-c*d)^{(1/2)}/d)/d*f-(c*f-d*e)/d)^{(1/2)})/(x+(-c*d)^{(1/2)}/d))-1/4*b^2*d/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(a*f-b*e)/(x+(-a*b)^{(1/2)}/b))*((x+(-a*b)^{(1/2)}/b)^2*f-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)}-1/4*b*d/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)*f*(-a*b)^{(1/2)}/(a*f-b*e)/(-a*f-b*e)/b)^{(1/2)}*\ln((-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x+(-a*b)^{(1/2)}/b)^2*f-2*(-a*b)^{(1/2)}*(x+(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x+(-a*b)^{(1/2)}/b))+3/4*b^3*d^3/(-a*b)^{(1/2)}/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-a*f-b*e)/b)^{(1/2)}*\ln((2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x-(-a*b)^{(1/2)}/b)^2*f+2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x-(-a*b)^{(1/2)}/b))-1/4*b^4*d^2/(-a*b)^{(1/2)}/a/((-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-(-c*d)^{(1/2)}*b+(-a*b)^{(1/2)}*d)^2/(-a*f-b*e)/b)^{(1/2)}*\ln((2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-2*(a*f-b*e)/b+2*(-a*f-b*e)/b)^{(1/2)}*((x-(-a*b)^{(1/2)}/b)^2*f+2*(-a*b)^{(1/2)}*(x-(-a*b)^{(1/2)}/b)/b*f-(a*f-b*e)/b)^{(1/2)})/(x-(-a*b)^{(1/2)}/b))*c$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(bx^2 + a)^2(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x^2+a)^2/(d*x^2+c)/(f*x^2+e)^(1/2),x, algorithm="maxima")

[Out] integrate(1/((b*x^2 + a)^2*(d*x^2 + c)*sqrt(f*x^2 + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(bx^2 + a)^2(dx^2 + c)\sqrt{fx^2 + e}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a + b*x^2)^2*(c + d*x^2)*(e + f*x^2)^(1/2)),x)

[Out] int(1/((a + b*x^2)^2*(c + d*x^2)*(e + f*x^2)^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a + bx^2)^2(c + dx^2)\sqrt{e + fx^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*x**2+a)**2/(d*x**2+c)/(f*x**2+e)**(1/2),x)

[Out] Integral(1/((a + b*x**2)**2*(c + d*x**2)*sqrt(e + f*x**2)), x)

$$3.30 \quad \int \frac{a+bx^2}{\sqrt{c+dx^2}(e+fx^2)^2} dx$$

Optimal. Leaf size=113

$$\frac{x\sqrt{c+dx^2}(be-af)}{2e(e+fx^2)(de-cf)} - \frac{(acf-2ade+bce)\tanh^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{e}\sqrt{c+dx^2}}\right)}{2e^{3/2}(de-cf)^{3/2}}$$

Rubi [A] time = 0.12, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {527, 12, 377, 208}

$$\frac{x\sqrt{c+dx^2}(be-af)}{2e(e+fx^2)(de-cf)} - \frac{(acf-2ade+bce)\tanh^{-1}\left(\frac{x\sqrt{de-cf}}{\sqrt{e}\sqrt{c+dx^2}}\right)}{2e^{3/2}(de-cf)^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Int[(a + b*x^2)/(Sqrt[c + d*x^2]*(e + f*x^2)^2), x]
```

```
[Out] ((b*e - a*f)*x*Sqrt[c + d*x^2])/(2*e*(d*e - c*f)*(e + f*x^2)) - ((b*c*e - 2*a*d*e + a*c*f)*ArcTanh[(Sqrt[d*e - c*f]*x)/(Sqrt[e]*Sqrt[c + d*x^2])])/(2*e^(3/2)*(d*e - c*f)^(3/2))
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 208

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-(a/b), 2]*ArcTanh[x/Rt[-(a/b), 2]])/a, x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 377

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 527

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := -Simp[((b*e - a*f)*x*(a + b*x^n)^(p+1)*(c + d*x^n)^(q+1))/(a*n*(b*c - a*d)*(p+1)), x] + Dist[1/(a*n*(b*c - a*d)*(p+1)), Int[(a + b*x^n)^(p+1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(n*(p+q+2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{a + bx^2}{\sqrt{c + dx^2} (e + fx^2)^2} dx &= \frac{(be - af)x\sqrt{c + dx^2}}{2e(de - cf)(e + fx^2)} + \frac{\int \frac{-bce + 2ade - acf}{\sqrt{c + dx^2} (e + fx^2)} dx}{2e(de - cf)} \\
&= \frac{(be - af)x\sqrt{c + dx^2}}{2e(de - cf)(e + fx^2)} - \frac{(bce - 2ade + acf) \int \frac{1}{\sqrt{c + dx^2} (e + fx^2)} dx}{2e(de - cf)} \\
&= \frac{(be - af)x\sqrt{c + dx^2}}{2e(de - cf)(e + fx^2)} - \frac{(bce - 2ade + acf) \operatorname{Subst}\left(\int \frac{1}{e - (de - cf)x^2} dx, x, \frac{x}{\sqrt{c + dx^2}}\right)}{2e(de - cf)} \\
&= \frac{(be - af)x\sqrt{c + dx^2}}{2e(de - cf)(e + fx^2)} - \frac{(bce - 2ade + acf) \tanh^{-1}\left(\frac{\sqrt{de - cf} x}{\sqrt{e} \sqrt{c + dx^2}}\right)}{2e^{3/2}(de - cf)^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.81, size = 191, normalized size = 1.69

$$\frac{x(be - af) \left(f(c + dx^2) - \frac{(e + fx^2)(2de - cf) \tanh^{-1}\left(\sqrt{\frac{x^2(de - cf)}{e(c + dx^2)}}\right)}{e\sqrt{\frac{x^2(de - cf)}{e(c + dx^2)}}} \right)}{2ef\sqrt{c + dx^2} (e + fx^2) (de - cf)} + \frac{b \tanh^{-1}\left(\frac{x\sqrt{de - cf}}{\sqrt{e} \sqrt{c + dx^2}}\right)}{\sqrt{e} f \sqrt{de - cf}}$$

Antiderivative was successfully verified.

[In] Integrate[(a + b*x^2)/(Sqrt[c + d*x^2]*(e + f*x^2)^2), x]

[Out] ((b*e - a*f)*x*(f*(c + d*x^2) - ((2*d*e - c*f)*(e + f*x^2)*ArcTanh[Sqrt[((d*e - c*f)*x^2)/(e*(c + d*x^2))]])/(e*Sqrt[((d*e - c*f)*x^2)/(e*(c + d*x^2))]))/(2*e*f*(d*e - c*f)*Sqrt[c + d*x^2]*(e + f*x^2)) + (b*ArcTanh[(Sqrt[d*e - c*f]*x)/(Sqrt[e]*Sqrt[c + d*x^2])])/(Sqrt[e]*f*Sqrt[d*e - c*f])

IntegrateAlgebraic [A] time = 0.62, size = 146, normalized size = 1.29

$$\frac{\sqrt{cf - de}(-acf + 2ade - bce) \tan^{-1}\left(\frac{-fx\sqrt{c + dx^2} + \sqrt{d}e + \sqrt{d}fx^2}{\sqrt{e}\sqrt{cf - de}}\right)}{2e^{3/2}(de - cf)^2} + \frac{x\sqrt{c + dx^2}(be - af)}{2e(e + fx^2)(de - cf)}$$

Antiderivative was successfully verified.

[In] IntegrateAlgebraic[(a + b*x^2)/(Sqrt[c + d*x^2]*(e + f*x^2)^2), x]

[Out] ((b*e - a*f)*x*Sqrt[c + d*x^2])/(2*e*(d*e - c*f)*(e + f*x^2)) + (Sqrt[-(d*e) + c*f]*(-(b*c*e) + 2*a*d*e - a*c*f)*ArcTan[(Sqrt[d]*e + Sqrt[d]*f*x^2 - f*x*Sqrt[c + d*x^2])/(Sqrt[e]*Sqrt[-(d*e) + c*f])])/(2*e^(3/2)*(d*e - c*f)^2)

fricas [B] time = 5.77, size = 513, normalized size = 4.54

$$\frac{4(bd^3 + acf^2 - (bc + ad)e^2)\sqrt{dx^2 + c} - (acf + (bc - 2ad)e^2 + (ac^2 + (bc - 2ad)ef)^2)\sqrt{dx^2 + c} \operatorname{atan}\left(\frac{(b\delta^2 - 8bd^2f + c^2\delta^2 + 4(ad^2 - 3c^2f)^2 + 4((2d - cf)^2 + ac)\sqrt{d^2 - c})\sqrt{dx^2 + c}}{f^2 + 2cf + d^2}\right) - 2(bd^3 + acf^2 - (bc + ad)e^2)\sqrt{dx^2 + c} + (acf + (bc - 2ad)e^2 + (ac^2 + (bc - 2ad)ef)^2)\sqrt{-d^2 + cf} \operatorname{arctan}\left(\frac{\sqrt{-d^2 + cf}(2d - f)^2 + c\sqrt{dx^2 + c}}{2((b\delta^2 - 8bd^2f + c^2\delta^2 + 4(ad^2 - 3c^2f)^2 + 4((2d - cf)^2 + ac)\sqrt{d^2 - c})}\right)}{8(b\delta^2 - 2cd^2f + c^2\delta^2 + (b\delta^2 - 2cd^2f + c^2\delta^2)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(f*x^2+e)^2/(d*x^2+c)^(1/2), x, algorithm="fricas")

[Out] [1/8*(4*(b*d*e^3 + a*c*e*f^2 - (b*c + a*d)*e^2*f)*sqrt(d*x^2 + c)*x - (a*c*e*f + (b*c - 2*a*d)*e^2 + (a*c*f^2 + (b*c - 2*a*d)*e*f)*x^2)*sqrt(d*e^2 - c

```
*e*f)*log(((8*d^2*e^2 - 8*c*d*e*f + c^2*f^2)*x^4 + c^2*e^2 + 2*(4*c*d*e^2 -
3*c^2*e*f)*x^2 + 4*((2*d*e - c*f)*x^3 + c*e*x)*sqrt(d*e^2 - c*e*f)*sqrt(d*
x^2 + c))/(f^2*x^4 + 2*e*f*x^2 + e^2)))/(d^2*e^5 - 2*c*d*e^4*f + c^2*e^3*f^
2 + (d^2*e^4*f - 2*c*d*e^3*f^2 + c^2*e^2*f^3)*x^2), 1/4*(2*(b*d*e^3 + a*c*e
*f^2 - (b*c + a*d)*e^2*f)*sqrt(d*x^2 + c)*x + (a*c*e*f + (b*c - 2*a*d)*e^2
+ (a*c*f^2 + (b*c - 2*a*d)*e*f)*x^2)*sqrt(-d*e^2 + c*e*f)*arctan(1/2*sqrt(-
d*e^2 + c*e*f)*((2*d*e - c*f)*x^2 + c*e)*sqrt(d*x^2 + c)/((d^2*e^2 - c*d*e*
f)*x^3 + (c*d*e^2 - c^2*e*f)*x)))/(d^2*e^5 - 2*c*d*e^4*f + c^2*e^3*f^2 + (d
^2*e^4*f - 2*c*d*e^3*f^2 + c^2*e^2*f^3)*x^2)]
```

giac [B] time = 3.40, size = 336, normalized size = 2.97

$$\frac{(ac\sqrt{d}f + bc\sqrt{d}e - 2ad^2e)\arctan\left(\frac{(\sqrt{dx - \sqrt{d^2+c}})^2 f - cf + 2de}{2\sqrt{d}f - d^2e}\right) - (\sqrt{dx - \sqrt{d^2+c}})^2 ac\sqrt{d}f - (\sqrt{dx - \sqrt{d^2+c}})^2 bc\sqrt{d}fe - 2(\sqrt{dx - \sqrt{d^2+c}})^2 ad^3fe - ac^2\sqrt{d}f^2 + 2(\sqrt{dx - \sqrt{d^2+c}})^2 bd^2e^2 + bc^2\sqrt{d}fe}{2\sqrt{d}f - d^2e}(cfe - de^2) - \frac{((\sqrt{dx - \sqrt{d^2+c}})^4 f - 2(\sqrt{dx - \sqrt{d^2+c}})^2 cf + 4(\sqrt{dx - \sqrt{d^2+c}})^2 de + c^2 f)(cfe - dfe^2)}{(\sqrt{dx - \sqrt{d^2+c}})^4 f - 2(\sqrt{dx - \sqrt{d^2+c}})^2 cf + 4(\sqrt{dx - \sqrt{d^2+c}})^2 de + c^2 f}(cfe - dfe^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(f*x^2+e)^2/(d*x^2+c)^(1/2),x, algorithm="giac")

```
[Out] -1/2*(a*c*sqrt(d)*f + b*c*sqrt(d)*e - 2*a*d^(3/2)*e)*arctan(1/2*((sqrt(d)*x
- sqrt(d*x^2 + c))^2*f - c*f + 2*d*e)/sqrt(c*d*f*e - d^2*e^2))/sqrt(c*d*f
*e - d^2*e^2)*(c*f*e - d*e^2) - ((sqrt(d)*x - sqrt(d*x^2 + c))^2*a*c*sqrt(
d)*f^2 - (sqrt(d)*x - sqrt(d*x^2 + c))^2*b*c*sqrt(d)*f*e - 2*(sqrt(d)*x - s
qrt(d*x^2 + c))^2*a*d^(3/2)*f*e - a*c^2*sqrt(d)*f^2 + 2*(sqrt(d)*x - sqrt(d
*x^2 + c))^2*b*d^(3/2)*e^2 + b*c^2*sqrt(d)*f*e)/(((sqrt(d)*x - sqrt(d*x^2 +
c))^4*f - 2*(sqrt(d)*x - sqrt(d*x^2 + c))^2*c*f + 4*(sqrt(d)*x - sqrt(d*x^
2 + c))^2*d*e + c^2*f)*(c*f^2*e - d*f*e^2))
```

maple [B] time = 0.03, size = 1622, normalized size = 14.35

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*x^2+a)/(f*x^2+e)^2/(d*x^2+c)^(1/2),x)

```
[Out] 1/4/e/(c*f-d*e)/(x-(-e*f)^(1/2)/f)*((x-(-e*f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*
(x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2)*a-1/4/f/(c*f-d*e)/(x-(-e*f)^(1/2)/
f)*((x-(-e*f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*((x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/
f)^(1/2)*b-1/4/e/f*(-e*f)^(1/2)*d/(c*f-d*e)/((c*f-d*e)/f)^(1/2)*ln((2*(-e*f)
)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/f+2*((c*f-d*e)/f)^(1/2)*((x-(-e*
f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2))/
(x-(-e*f)^(1/2)/f))*a+1/4/f^2*(-e*f)^(1/2)*d/(c*f-d*e)/((c*f-d*e)/f)^(1/2)*l
n((2*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/f+2*((c*f-d*e)/f)^(1/2
))*((x-(-e*f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f
)^(1/2))/((x-(-e*f)^(1/2)/f))*b-1/4/e/(-e*f)^(1/2)/((c*f-d*e)/f)^(1/2)*ln((2
*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/f+2*((c*f-d*e)/f)^(1/2)*((
x-(-e*f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1
/2))/((x-(-e*f)^(1/2)/f))*a-1/4/(-e*f)^(1/2)/f/((c*f-d*e)/f)^(1/2)*ln((2*(-e
*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/f+2*((c*f-d*e)/f)^(1/2)*((x-(-
e*f)^(1/2)/f)^2*d+2*(-e*f)^(1/2)*(x-(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2)
)/(x-(-e*f)^(1/2)/f))*b+1/4/e/(c*f-d*e)/(x+(-e*f)^(1/2)/f)*((x+(-e*f)^(1/2)/
f)^2*d-2*(-e*f)^(1/2)*(x+(-e*f)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2)*a-1/4/f/(c*
f-d*e)/(x+(-e*f)^(1/2)/f)*((x+(-e*f)^(1/2)/f)^2*d-2*(-e*f)^(1/2)*(x+(-e*f)
)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2)*b+1/4/e/f*(-e*f)^(1/2)*d/(c*f-d*e)/((c*f-d*
e)/f)^(1/2)*ln((-2*(-e*f)^(1/2)*(x+(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/f+2*((c*
f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*(-e*f)^(1/2)*(x+(-e*f)^(1/2)/f)*d
/f+(c*f-d*e)/f)^(1/2))/((x+(-e*f)^(1/2)/f))*a-1/4/f^2*(-e*f)^(1/2)*d/(c*f-d*
e)/((c*f-d*e)/f)^(1/2)*ln((-2*(-e*f)^(1/2)*(x+(-e*f)^(1/2)/f)*d/f+2*(c*f-d*
e)/f+2*((c*f-d*e)/f)^(1/2)*((x+(-e*f)^(1/2)/f)^2*d-2*(-e*f)^(1/2)*(x+(-e*f)
)^(1/2)/f)*d/f+(c*f-d*e)/f)^(1/2))/((x+(-e*f)^(1/2)/f))*b+1/4/e/(-e*f)^(1/2)/
((c*f-d*e)/f)^(1/2)*ln((-2*(-e*f)^(1/2)*(x+(-e*f)^(1/2)/f)*d/f+2*(c*f-d*e)/
```


$$f+2*((c*f-d*e)/f)^{(1/2)}*((x+(-e*f)^{(1/2)}/f)^2*d-2*(-e*f)^{(1/2)}*(x+(-e*f)^{(1/2)}/f)*d/f+(c*f-d*e)/f)^{(1/2)})/(x+(-e*f)^{(1/2)}/f))*a+1/4/(-e*f)^{(1/2)}/f/((c*f-d*e)/f)^{(1/2)}*\ln((-2*(-e*f)^{(1/2)}*(x+(-e*f)^{(1/2)}/f)*d/f+2*(c*f-d*e)/f+2*((c*f-d*e)/f)^{(1/2)}*((x+(-e*f)^{(1/2)}/f)^2*d-2*(-e*f)^{(1/2)}*(x+(-e*f)^{(1/2)}/f)*d/f+(c*f-d*e)/f)^{(1/2)})/(x+(-e*f)^{(1/2)}/f))*b$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{bx^2 + a}{\sqrt{dx^2 + c} (fx^2 + e)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x^2+a)/(f*x^2+e)^2/(d*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*x^2 + a)/(sqrt(d*x^2 + c)*(f*x^2 + e)^2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{bx^2 + a}{\sqrt{dx^2 + c} (fx^2 + e)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2)/((c + d*x^2)^(1/2)*(e + f*x^2)^2),x)

[Out] int((a + b*x^2)/((c + d*x^2)^(1/2)*(e + f*x^2)^2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{a + bx^2}{\sqrt{c + dx^2} (e + fx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*x**2+a)/(f*x**2+e)**2/(d*x**2+c)**(1/2),x)

[Out] Integral((a + b*x**2)/(sqrt(c + d*x**2)*(e + f*x**2)**2), x)

Chapter 4

Appendix

Local contents

4.1	Download section	138
4.2	Listing of Grading functions	138

4.1 Download section

The following zip files contain the raw integrals used in this test.

This is a subset of Rubi test suite thanks to Albert Rich, which includes only the algebraic integrals with elementray optimal antiderivatives. It also includes a subset of a test file provided thanks to Sam Blake.

Mathematica format Mathematica_syntax_CAS_integration_elementary_version.zip

Maple and Mupad format Maple_syntax_CAS_integration_elementary_version.zip

Sympy format SYMPY_syntax_CAS_integration_elementary_version.zip

Sage math format SAGE_syntax_CAS_integration_elementary_version.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
    If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
      If[LeafCount[result]<=2*LeafCount[optimal],
        "A",
        "B"],
      "C"],
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      "C",
      "F"]]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
```

```

(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=

```

```
MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]
```

```
AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]
```

4.2.2 Maple grading function

```
# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#                    if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#                    see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
    debug:=false;

    leaf_count_result:=leafcount(result);
    #do NOT call ExpnType() if leaf size is too large. Recursion problem
    if leaf_count_result > 500000 then
        return "B";
    fi;

    leaf_count_optimal:=leafcount(optimal);

    ExpnType_result:=ExpnType(result);
    ExpnType_optimal:=ExpnType(optimal);

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;

```

```

if is_contains_complex(result) then
  if is_contains_complex(optimal) then
    if debug then
      print("both result and optimal complex");
    fi;
    #both result and optimal complex
    if leaf_count_result<=2*leaf_count_optimal then
      return "A";
    else
      return "B";
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C";
  end if
else # result do not contain complex
  # this assumes optimal do not as well
  if debug then
    print("result do not contain complex, this assumes optimal do
not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B";
  end if
end if
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function

```

```

# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`') or type(expn,'`*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

```



```

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
  #sympy do not have leaf count function. This is approximation
  return round(1.7*count_ops(expr))

def is_sqrt(expr):
  if isinstance(expr,Pow):
    if expr.args[1] == Rational(1,2):
      return True
    else:
      return False
  else:
    return False

def is_elementary_function(func):
  return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                 asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                 asinh,acosh,atanh,acoth,asech,acsch
                 ]

def is_special_function(func):
  return func in [ erf,erfc,erfi,
                 fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                 gamma,loggamma,digamma,zeta,polylog,LambertW,
                 elliptic_f,elliptic_e,elliptic_pi,exp_polar
                 ]

def is_hypergeometric_function(func):
  return func in [hyper]

def is_appell_function(func):
  return func in [appellf1]

def is_atom(expn):

```

```

try:
    if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
        return True
    else:
        return False

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'^+^') or
    type(expn,'*^')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
    expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType
    ,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:
        m1 = max(map(expnType, list(expn.args)))
        return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

#main function

```

```

def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```

4.2.4 SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False

```

```

else:
    return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M',
        hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
    sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):

```

```

        return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens")
:
            return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
        return False

    except AttributeError as error:
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],
Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer
)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0])) #max(2,expnType(
expn.args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
isinstance(expn,Add) or isinstance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn)))
    elif is_elementary_function(expn.operator()): #is_elementary_function(expn
.func)
        return max(3,expnType(expn.operands()[0]))
    elif is_special_function(expn.operator()): #is_special_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(4,m1) #max(4,m1)
    elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
        return max(5,m1) #max(5,m1)
    elif is_appell_function(expn.operator()):
        m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))

```

```

    return max(6,m1)    #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands()))    #max(map(expnType, list(
expn.args)))
    return max(8,m1)    #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug: print ("Enter grade_antiderivative for sagemath")

    leaf_count_result  = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

    expnType_result  = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```