

Computer Algebra Independent Integration Tests

Summer 2023 edition

0-Independent-test-suites/4-Charlwood-Problems

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 10:13am

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	39
4	Appendix	295

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [50]. This is test number [4].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (50)	0.00 (0)
Fricas	100.00 (50)	0.00 (0)
Rubi	96.00 (48)	4.00 (2)
Giac	82.00 (41)	18.00 (9)
Maple	66.00 (33)	34.00 (17)
Maxima	52.00 (26)	48.00 (24)
Sympy	38.00 (19)	62.00 (31)
Mupad	24.00 (12)	76.00 (38)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	86.000	4.000	6.000	4.000
Mathematica	80.000	8.000	12.000	0.000
Fricas	60.000	36.000	4.000	0.000
Giac	52.000	30.000	0.000	18.000
Maxima	42.000	6.000	4.000	48.000
Maple	34.000	16.000	16.000	34.000
Sympy	30.000	6.000	2.000	62.000
Mupad	0.000	24.000	0.000	76.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Rubi	2	100.00	0.00	0.00
Giac	9	100.00	0.00	0.00
Maple	17	100.00	0.00	0.00
Maxima	24	100.00	0.00	0.00
Sympy	31	70.97	29.03	0.00
Mupad	38	0.00	100.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.10
Mathematica	0.21
Giac	0.30
Maxima	0.32
Maple	0.61
Mupad	0.65
Fricas	0.73
Sympy	4.41

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	44.32	1.48	31.00	0.98
Rubi	66.25	1.25	40.00	1.00
Fricas	84.74	1.58	50.50	1.35
Giac	88.37	1.61	50.00	1.41
Mathematica	112.46	1.58	45.50	1.00
Maple	152.39	2.27	54.00	1.15
Maxima	220.77	3.04	27.00	0.88
Mupad	223.25	2.82	69.00	1.98

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {5, 48}

Mathematica {45}

Maple {18, 33, 35}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v1.0

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	36

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 6, 7, 8, 9, 10, 11, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 43, 44, 46, 47, 48, 49, 50 }

B grade { 4, 42 }

C grade { 5, 12, 13 }

F normal fail { 3, 45 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 6, 7, 9, 10, 11, 14, 16, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 38, 39, 40, 42, 43, 44, 45, 46, 48, 49, 50 }

B grade { 15, 17, 21, 41 }

C grade { 5, 8, 12, 13, 37, 47 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 2, 6, 7, 9, 19, 24, 27, 32, 34, 36, 37, 38, 39, 40, 43, 46, 48 }

B grade { 1, 4, 8, 42, 44, 47, 49, 50 }

C grade { 3, 5, 12, 18, 28, 31, 33, 35 }

F normal fail { 10, 11, 13, 14, 15, 16, 17, 20, 21, 22, 23, 25, 26, 29, 30, 41, 45 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 5, 9, 10, 11, 12, 13, 14, 18, 19, 20, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 39, 46, 48 }

B grade { 4, 6, 7, 15, 16, 17, 21, 22, 37, 38, 40, 41, 42, 43, 44, 47, 49, 50 }

C grade { 8, 45 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 9, 16, 18, 19, 20, 24, 25, 27, 28, 31, 32, 33, 34, 35, 36, 37, 46, 48 }

B grade { 7, 40, 43 }

C grade { 8, 30 }

F normal fail { 4, 5, 6, 10, 11, 12, 13, 14, 15, 17, 21, 22, 23, 26, 29, 38, 39, 41, 42, 44, 45, 47, 49, 50 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 2, 6, 10, 14, 15, 16, 17, 18, 19, 20, 21, 22, 24, 25, 26, 28, 33, 34, 35, 36, 40, 43, 47, 48, 49, 50 }

B grade { 1, 4, 7, 8, 12, 13, 27, 29, 30, 31, 32, 37, 41, 42, 44 }

C grade { }

F normal fail { 3, 5, 9, 11, 23, 38, 39, 45, 46 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 4, 9, 12, 19, 24, 30, 37, 40, 46, 47, 48, 50 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 3, 5, 6, 7, 8, 10, 11, 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 25, 26, 27, 28, 29, 31, 32, 33, 34, 35, 36, 38, 39, 41, 42, 43, 44, 45, 49 }

F(-2) exception fail { }

Sympy

A grade { 1, 2, 10, 18, 19, 20, 24, 25, 31, 33, 34, 35, 36, 46, 48 }

B grade { 28, 37, 40 }

C grade { 16 }

F normal fail { 3, 6, 7, 8, 9, 14, 17, 21, 22, 26, 27, 29, 30, 32, 38, 39, 41, 42, 43, 45, 47, 49 }

F(-1) timeout fail { 4, 5, 11, 12, 13, 15, 23, 44, 50 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	52	93	58	54	102	272	0
N.S.	1	1.00	1.02	1.82	1.14	1.06	2.00	5.33	0.00
time (sec)	N/A	0.023	0.018	0.146	0.277	0.263	3.722	0.281	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	16	15	15	12	15	0
N.S.	1	1.00	1.00	0.94	0.88	0.88	0.71	0.88	0.00
time (sec)	N/A	0.021	0.006	0.213	0.307	0.241	0.084	0.296	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	A	A	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	0	106	252	4	49	0	0	0
N.S.	1	0.00	1.54	3.65	0.06	0.71	0.00	0.00	0.00
time (sec)	N/A	0.000	0.700	1.098	0.907	23.673	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	B	F	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	332	188	413	0	451	0	235	666
N.S.	1	3.42	1.94	4.26	0.00	4.65	0.00	2.42	6.87
time (sec)	N/A	0.497	0.273	0.179	0.000	0.255	0.000	0.399	1.504

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	C	F	A	F(-1)	F	F(-1)
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	289	159	312	0	33	0	0	0
N.S.	1	6.42	3.53	6.93	0.00	0.73	0.00	0.00	0.00
time (sec)	N/A	0.382	0.806	2.597	0.000	0.302	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	56	74	64	0	88	0	79	0
N.S.	1	1.00	1.32	1.14	0.00	1.57	0.00	1.41	0.00
time (sec)	N/A	0.052	0.084	0.259	0.000	0.308	0.000	0.301	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	12	27	30	0	28	0
N.S.	1	1.00	1.00	0.80	1.80	2.00	0.00	1.87	0.00
time (sec)	N/A	0.029	0.012	0.576	0.205	0.306	0.000	0.299	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	C	C	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	137	99	1604	4065	226	0	495	0
N.S.	1	1.00	0.72	11.71	29.67	1.65	0.00	3.61	0.00
time (sec)	N/A	0.145	0.201	1.602	0.781	0.300	0.000	0.336	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	70	36	60	32	0	0	60
N.S.	1	1.00	1.71	0.88	1.46	0.78	0.00	0.00	1.46
time (sec)	N/A	0.020	0.249	0.297	0.289	0.258	0.000	0.000	0.420

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	A	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	44	62	38	0	0	28	56	46	0
N.S.	1	1.41	0.86	0.00	0.00	0.64	1.27	1.05	0.00
time (sec)	N/A	0.540	0.169	0.000	0.000	0.246	0.383	0.276	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	68	68	64	0	0	43	0	0	0
N.S.	1	1.00	0.94	0.00	0.00	0.63	0.00	0.00	0.00
time (sec)	N/A	0.112	0.038	0.000	0.000	0.262	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	C	F	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	269	226	471	0	191	0	364	661
N.S.	1	1.91	1.60	3.34	0.00	1.35	0.00	2.58	4.69
time (sec)	N/A	0.596	0.132	0.096	0.000	0.268	0.000	0.291	1.375

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	F	F	A	F(-1)	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	152	286	2175	0	0	200	0	373	0
N.S.	1	1.88	14.31	0.00	0.00	1.32	0.00	2.45	0.00
time (sec)	N/A	0.340	3.092	0.000	0.000	0.266	0.000	0.341	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	45	51	44	0	0	63	0	57	0
N.S.	1	1.13	0.98	0.00	0.00	1.40	0.00	1.27	0.00
time (sec)	N/A	0.094	0.034	0.000	0.000	0.259	0.000	0.290	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	34	34	76	0	0	62	0	36	0
N.S.	1	1.00	2.24	0.00	0.00	1.82	0.00	1.06	0.00
time (sec)	N/A	0.026	0.073	0.000	0.000	0.282	0.000	0.285	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	B	C	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	22	22	22	0	18	56	78	18	0
N.S.	1	1.00	1.00	0.00	0.82	2.55	3.55	0.82	0.00
time (sec)	N/A	0.016	0.020	0.000	0.262	0.275	20.249	0.280	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	32	32	89	0	0	58	0	36	0
N.S.	1	1.00	2.78	0.00	0.00	1.81	0.00	1.12	0.00
time (sec)	N/A	0.028	0.054	0.000	0.000	0.257	0.000	0.269	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	43	43	43	89	41	32	42	62	0
N.S.	1	1.00	1.00	2.07	0.95	0.74	0.98	1.44	0.00
time (sec)	N/A	0.030	0.017	0.080	0.272	0.261	2.687	0.282	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	28	21	34	34	48	35	174
N.S.	1	1.00	1.00	0.75	1.21	1.21	1.71	1.25	6.21
time (sec)	N/A	0.008	0.015	0.336	0.181	0.228	0.901	0.270	0.144

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	26	0	22	22	20	22	0
N.S.	1	1.00	1.00	0.00	0.85	0.85	0.77	0.85	0.00
time (sec)	N/A	0.027	0.016	0.000	0.227	0.241	3.249	0.284	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	38	38	85	0	0	138	0	38	0
N.S.	1	1.00	2.24	0.00	0.00	3.63	0.00	1.00	0.00
time (sec)	N/A	0.039	0.053	0.000	0.000	0.259	0.000	0.285	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	70	94	88	0	0	110	0	52	0
N.S.	1	1.34	1.26	0.00	0.00	1.57	0.00	0.74	0.00
time (sec)	N/A	0.095	0.063	0.000	0.000	0.255	0.000	0.296	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	48	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.83	0.00	0.00	0.00
time (sec)	N/A	0.102	0.026	0.000	0.000	0.256	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	41	37	36	35	31	36	27
N.S.	1	1.00	0.75	0.67	0.65	0.64	0.56	0.65	0.49
time (sec)	N/A	0.043	0.016	0.052	0.190	0.234	2.143	0.263	0.398

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	26	26	26	0	22	22	20	22	0
N.S.	1	1.00	1.00	0.00	0.85	0.85	0.77	0.85	0.00
time (sec)	N/A	0.031	0.015	0.000	0.321	0.244	3.568	0.279	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	78	78	119	0	0	115	0	122	0
N.S.	1	1.00	1.53	0.00	0.00	1.47	0.00	1.56	0.00
time (sec)	N/A	0.219	0.041	0.000	0.000	0.246	0.000	0.347	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	25	35	35	39	0	73	0
N.S.	1	1.00	0.64	0.90	0.90	1.00	0.00	1.87	0.00
time (sec)	N/A	0.041	0.023	0.073	0.274	0.241	0.000	0.280	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	B	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	54	15	23	29	23	0
N.S.	1	1.00	1.00	3.18	0.88	1.35	1.71	1.35	0.00
time (sec)	N/A	0.020	0.012	0.645	0.268	0.252	0.435	0.284	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	57	57	77	0	0	81	0	104	0
N.S.	1	1.00	1.35	0.00	0.00	1.42	0.00	1.82	0.00
time (sec)	N/A	0.059	0.045	0.000	0.000	0.250	0.000	0.298	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	C	A	F	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	45	45	45	0	387	69	0	108	37
N.S.	1	1.00	1.00	0.00	8.60	1.53	0.00	2.40	0.82
time (sec)	N/A	0.035	0.026	0.000	0.442	0.258	0.000	0.284	0.031

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	33	56	22	47	19	54	0
N.S.	1	1.00	1.14	1.93	0.76	1.62	0.66	1.86	0.00
time (sec)	N/A	0.040	0.019	0.677	0.272	0.249	3.830	0.278	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	20	19	22	0	40	0
N.S.	1	1.00	1.00	0.95	0.90	1.05	0.00	1.90	0.00
time (sec)	N/A	0.034	0.011	0.229	0.272	0.252	0.000	0.291	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	34	34	27	119	27	27	29	28	0
N.S.	1	1.00	0.79	3.50	0.79	0.79	0.85	0.82	0.00
time (sec)	N/A	0.026	0.014	0.083	0.281	0.248	1.369	0.293	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	21	29	29	33	26	58	0
N.S.	1	1.00	0.64	0.88	0.88	1.00	0.79	1.76	0.00
time (sec)	N/A	0.028	0.019	0.069	0.275	0.244	2.951	0.266	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	25	25	35	34	15	15	22	22	0
N.S.	1	1.00	1.40	1.36	0.60	0.60	0.88	0.88	0.00
time (sec)	N/A	0.020	0.031	0.388	0.202	0.245	16.148	0.303	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	40	39	25	41	41	44	0
N.S.	1	1.00	1.18	1.15	0.74	1.21	1.21	1.29	0.00
time (sec)	N/A	0.029	0.013	0.064	0.295	0.268	2.595	0.272	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	46	14	24	29	46	27	13
N.S.	1	1.00	2.88	0.88	1.50	1.81	2.88	1.69	0.81
time (sec)	N/A	0.017	0.074	0.154	0.274	0.255	9.243	0.275	0.220

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	22	0	42	0	0	0
N.S.	1	1.00	1.00	0.96	0.00	1.83	0.00	0.00	0.00
time (sec)	N/A	0.022	0.300	1.543	0.000	0.269	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	19	0	18	0	0	0
N.S.	1	1.00	1.00	0.83	0.00	0.78	0.00	0.00	0.00
time (sec)	N/A	0.021	0.262	1.048	0.000	0.271	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	39	24	82	93	105	36	55
N.S.	1	1.00	1.77	1.09	3.73	4.23	4.77	1.64	2.50
time (sec)	N/A	0.053	0.033	0.606	0.284	0.264	0.659	0.304	0.386

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	42	42	87	0	0	146	0	90	0
N.S.	1	1.00	2.07	0.00	0.00	3.48	0.00	2.14	0.00
time (sec)	N/A	0.118	0.067	0.000	0.000	0.268	0.000	0.321	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	A	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	59	45	78	0	54	0	92	0
N.S.	1	2.11	1.61	2.79	0.00	1.93	0.00	3.29	0.00
time (sec)	N/A	0.137	0.028	0.773	0.000	0.279	0.000	0.509	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	55	37	565	186	0	50	0
N.S.	1	1.00	1.62	1.09	16.62	5.47	0.00	1.47	0.00
time (sec)	N/A	0.039	0.044	0.220	0.370	0.262	0.000	0.293	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	50	65	67	0	63	0	67	0
N.S.	1	1.28	1.67	1.72	0.00	1.62	0.00	1.72	0.00
time (sec)	N/A	0.034	0.058	0.888	0.000	0.298	0.000	0.328	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	F	C	F	F	F(-1)
verified	N/A	N/A	No	N/A	TBD	TBD	TBD	TBD	TBD
size	337	0	552	0	0	337	0	0	0
N.S.	1	0.00	1.64	0.00	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.000	2.848	0.000	0.000	0.272	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	58	78	67	52	87	0	78
N.S.	1	1.00	0.75	1.01	0.87	0.68	1.13	0.00	1.01
time (sec)	N/A	0.176	0.088	4.497	0.277	0.247	0.501	0.000	0.256

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	120	120	95	508	0	298	0	92	413
N.S.	1	1.00	0.79	4.23	0.00	2.48	0.00	0.77	3.44
time (sec)	N/A	0.098	0.165	0.171	0.000	0.261	0.000	0.284	1.102

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	No	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	37	31	28	26	22	29	27	40
N.S.	1	1.19	1.00	0.90	0.84	0.71	0.94	0.87	1.29
time (sec)	N/A	0.009	0.092	0.041	0.317	0.253	9.144	0.286	0.802

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	138	0	60	0	34	0
N.S.	1	1.00	1.00	4.76	0.00	2.07	0.00	1.17	0.00
time (sec)	N/A	0.030	0.025	0.194	0.000	0.261	0.000	0.326	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	106	106	210	0	205	0	111	455
N.S.	1	1.00	1.00	1.98	0.00	1.93	0.00	1.05	4.29
time (sec)	N/A	0.084	0.177	0.096	0.000	0.275	0.000	0.345	1.153

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [1] had the largest ratio of [1.3999999999999999]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.00	5	1.400
2	A	2	2	1.00	15	0.133
3	F	0	0	N/A	0.000	N/A
4	B	32	13	3.42	14	0.929
5	C	5	4	6.42	19	0.210
6	A	7	7	1.00	13	0.538
7	A	4	4	1.00	13	0.308
8	A	9	7	1.00	14	0.500
9	A	7	7	1.00	12	0.583
10	A	5	5	1.41	19	0.263
11	A	7	8	1.00	29	0.276
12	C	40	15	1.91	14	1.071
13	C	32	16	1.88	27	0.593
14	A	9	11	1.13	18	0.611
15	A	3	4	1.00	24	0.167
16	A	3	4	1.00	12	0.333
17	A	3	4	1.00	22	0.182
18	A	4	4	1.00	15	0.267
19	A	4	4	1.00	13	0.308
20	A	2	3	1.00	23	0.130
21	A	5	6	1.00	17	0.353
22	A	7	8	1.34	15	0.533
23	A	4	9	1.00	25	0.360
24	A	5	5	1.00	27	0.185

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	2	3	1.00	23	0.130
26	A	18	11	1.00	27	0.407
27	A	3	3	1.00	17	0.176
28	A	2	2	1.00	13	0.154
29	A	7	6	1.00	17	0.353
30	A	5	5	1.00	15	0.333
31	A	4	4	1.00	15	0.267
32	A	2	2	1.00	17	0.118
33	A	5	5	1.00	13	0.385
34	A	3	3	1.00	15	0.200
35	A	2	2	1.00	13	0.154
36	A	5	5	1.00	13	0.385
37	A	2	2	1.00	11	0.182
38	A	2	2	1.00	24	0.083
39	A	2	2	1.00	24	0.083
40	A	4	5	1.00	10	0.500
41	A	6	7	1.00	12	0.583
42	B	5	5	2.11	13	0.385
43	A	4	4	1.00	13	0.308
44	A	4	4	1.28	15	0.267
45	F	0	0	N/A	0.000	N/A
46	A	13	10	1.00	12	0.833
47	A	12	8	1.00	12	0.667
48	A	6	6	1.19	18	0.333
49	A	4	4	1.00	14	0.286
50	A	6	6	1.00	14	0.429

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \arcsin(x) \log(x) dx$	41
3.2	$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$	47
3.3	$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx$	50
3.4	$\int \log(1 + x\sqrt{1+x^2}) dx$	54
3.5	$\int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx$	67
3.6	$\int \tan(x)\sqrt{1+\tan^4(x)} dx$	72
3.7	$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx$	77
3.8	$\int \sqrt{2+2\tan(x)+\tan^2(x)} dx$	81
3.9	$\int \arctan(\sqrt{-1+\sec(x)}) \sin(x) dx$	92
3.10	$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx$	97
3.11	$\int \frac{x \log(1+x^2) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx$	101
3.12	$\int \arctan(x + \sqrt{1-x^2}) dx$	106
3.13	$\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx$	116
3.14	$\int \frac{\arcsin(x)}{1+\sqrt{1-x^2}} dx$	127
3.15	$\int \frac{\log(x+\sqrt{1+x^2})}{(1-x^2)^{3/2}} dx$	132
3.16	$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx$	136
3.17	$\int \frac{\log(x+\sqrt{-1+x^2})}{(1+x^2)^{3/2}} dx$	140
3.18	$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx$	144
3.19	$\int \frac{\sqrt{1+x^3}}{x} dx$	148
3.20	$\int \frac{x \log(x+\sqrt{-1+x^2})}{\sqrt{-1+x^2}} dx$	152

3.21	$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx$	155
3.22	$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx$	159
3.23	$\int \frac{x \arctan(x) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx$	164
3.24	$\int \frac{x \log(1+\sqrt{1-x^2})}{\sqrt{1-x^2}} dx$	169
3.25	$\int \frac{x \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx$	173
3.26	$\int \frac{x \log(x+\sqrt{1-x^2})}{\sqrt{1-x^2}} dx$	176
3.27	$\int \frac{\log(x)}{x^2 \sqrt{1-x^2}} dx$	183
3.28	$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx$	187
3.29	$\int \frac{\arctan(x)}{x^2 \sqrt{1-x^2}} dx$	191
3.30	$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx$	196
3.31	$\int \frac{\arctan(x)}{x^2 \sqrt{1+x^2}} dx$	201
3.32	$\int \frac{\arcsin(x)}{x^2 \sqrt{1-x^2}} dx$	205
3.33	$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx$	208
3.34	$\int \frac{\log(x)}{x^2 \sqrt{1+x^2}} dx$	212
3.35	$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx$	216
3.36	$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx$	219
3.37	$\int \frac{\sin(x)}{1+\sin^2(x)} dx$	224
3.38	$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx$	228
3.39	$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx$	232
3.40	$\int \frac{\log(\sin(x))}{1+\sin(x)} dx$	236
3.41	$\int \log(\sin(x)) \sqrt{1+\sin(x)} dx$	241
3.42	$\int \frac{\sec(x)}{\sqrt{-1+\sec^4(x)}} dx$	246
3.43	$\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx$	251
3.44	$\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx$	256
3.45	$\int \sqrt{-\sqrt{-1+\sec(x)} + \sqrt{1+\sec(x)}} dx$	260
3.46	$\int x \arctan(x)^2 \log(1+x^2) dx$	266
3.47	$\int \arctan(x\sqrt{1+x^2}) dx$	272
3.48	$\int -\arctan(\sqrt{x-\sqrt{1+x}}) dx$	279
3.49	$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx$	283
3.50	$\int \arctan(x\sqrt{1-x^2}) dx$	287

3.1 $\int \arcsin(x) \log(x) dx$

Optimal result	41
Rubi [A] (verified)	41
Mathematica [A] (verified)	43
Maple [B] (verified)	43
Fricas [A] (verification not implemented)	44
Sympy [A] (verification not implemented)	44
Maxima [A] (verification not implemented)	45
Giac [B] (verification not implemented)	45
Mupad [F(-1)]	46

Optimal result

Integrand size = 5, antiderivative size = 51

$$\int \arcsin(x) \log(x) dx = -2\sqrt{1-x^2} + \operatorname{arctanh}(\sqrt{1-x^2}) - x \arcsin(x)(1 - \log(x)) + \sqrt{1-x^2} \log(x)$$

[Out] $\operatorname{arctanh}((-x^2+1)^{(1/2)})-x*\arcsin(x)*(1-\ln(x))-2*(-x^2+1)^{(1/2)}+\ln(x)*(-x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 1.400$, Rules used = {4715, 267, 2434, 272, 52, 65, 212}

$$\int \arcsin(x) \log(x) dx = -x \arcsin(x) + x \arcsin(x) \log(x) + \operatorname{arctanh}(\sqrt{1-x^2}) - 2\sqrt{1-x^2} + \sqrt{1-x^2} \log(x)$$

[In] `Int[ArcSin[x]*Log[x],x]`

[Out] `-2*Sqrt[1 - x^2] - x*ArcSin[x] + ArcTanh[Sqrt[1 - x^2]] + Sqrt[1 - x^2]*Log[x] + x*ArcSin[x]*Log[x]`

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ`

$[m, 0] \&\& (!\text{IntegerQ}[n] \mid\mid (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0])) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

$\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

Rule 267

$\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p+1)}/(b*n*(p+1)), x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{EqQ}[m, n - 1] \&\& \text{NeQ}[p, -1]$

Rule 272

$\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n] - 1)}(a + b*x)^p, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

Rule 2434

$\text{Int}[(a_.) + \text{Log}[(c_.)(x_)^{(n_.)}]*(b_.)]*(P_x_.)*(F_)[(d_.)*((e_.) + (f_.)(x_))]^{(m_.)}, x_Symbol] \rightarrow \text{With}[\{u = \text{IntHide}[P_x*F[d*(e + f*x)]^m, x]\}, \text{Dist}[a + b*\text{Log}[c*x^n], u, x] - \text{Dist}[b*n, \text{Int}[\text{Dist}[1/x, u, x], x], x]] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{PolynomialQ}[P_x, x] \&\& \text{IGtQ}[m, 0] \&\& \text{MemberQ}[\{\text{ArcSin}, \text{ArcCos}, \text{ArcSinh}, \text{ArcCosh}\}, F]$

Rule 4715

$\text{Int}[(a_.) + \text{ArcSin}[(c_.)(x_)]*(b_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcSin}[c*x])^n, x] - \text{Dist}[b*c*n, \text{Int}[x*((a + b*\text{ArcSin}[c*x])^{(n-1)})/\text{Sqrt}[1 - c^2*x^2]), x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{GtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \sqrt{1-x^2} \log(x) + x \arcsin(x) \log(x) - \int \left(\frac{\sqrt{1-x^2}}{x} + \arcsin(x) \right) dx \\
&= \sqrt{1-x^2} \log(x) + x \arcsin(x) \log(x) - \int \frac{\sqrt{1-x^2}}{x} dx - \int \arcsin(x) dx \\
&= -x \arcsin(x) + \sqrt{1-x^2} \log(x) + x \arcsin(x) \log(x) \\
&\quad - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1-x}}{x} dx, x, x^2 \right) + \int \frac{x}{\sqrt{1-x^2}} dx \\
&= -2\sqrt{1-x^2} - x \arcsin(x) + \sqrt{1-x^2} \log(x) \\
&\quad + x \arcsin(x) \log(x) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1-xx}} dx, x, x^2 \right) \\
&= -2\sqrt{1-x^2} - x \arcsin(x) + \sqrt{1-x^2} \log(x) \\
&\quad + x \arcsin(x) \log(x) + \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x^2} \right) \\
&= -2\sqrt{1-x^2} - x \arcsin(x) + \text{arctanh}(\sqrt{1-x^2}) + \sqrt{1-x^2} \log(x) + x \arcsin(x) \log(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.02

$$\begin{aligned}
\int \arcsin(x) \log(x) dx &= -2\sqrt{1-x^2} + x \arcsin(x)(-1 + \log(x)) \\
&\quad + \left(-1 + \sqrt{1-x^2} \right) \log(x) + \log \left(1 + \sqrt{1-x^2} \right)
\end{aligned}$$

[In] Integrate[ArcSin[x]*Log[x],x]

[Out] -2*Sqrt[1 - x^2] + x*ArcSin[x]*(-1 + Log[x]) + (-1 + Sqrt[1 - x^2])*Log[x] + Log[1 + Sqrt[1 - x^2]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(45) = 90.

Time = 0.15 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.82

method	result
default	$\frac{2 \arcsin(x) \tan\left(\frac{\arcsin(x)}{2}\right) \ln\left(\frac{2 \tan\left(\frac{\arcsin(x)}{2}\right)}{1 + \tan^2\left(\frac{\arcsin(x)}{2}\right)}\right) - 2\left(\tan^2\left(\frac{\arcsin(x)}{2}\right)\right) \ln\left(\frac{2 \tan\left(\frac{\arcsin(x)}{2}\right)}{1 + \tan^2\left(\frac{\arcsin(x)}{2}\right)}\right) - 2 \arcsin(x) \tan\left(\frac{\arcsin(x)}{2}\right) - 4}{1 + \tan^2\left(\frac{\arcsin(x)}{2}\right)}$

```
[In] int(arcsin(x)*ln(x),x,method=_RETURNVERBOSE)
```

```
[Out] 2*(arcsin(x)*tan(1/2*arcsin(x))*ln(2*tan(1/2*arcsin(x))/(1+tan(1/2*arcsin(x))^2))-tan(1/2*arcsin(x))^2*ln(2*tan(1/2*arcsin(x))/(1+tan(1/2*arcsin(x))^2))-arcsin(x)*tan(1/2*arcsin(x))-2)/(1+tan(1/2*arcsin(x))^2)-ln(1+tan(1/2*arcsin(x))^2)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.06

$$\int \arcsin(x) \log(x) dx = x \arcsin(x) \log(x) - x \arcsin(x) + \sqrt{-x^2 + 1}(\log(x) - 2) + \frac{1}{2} \log(\sqrt{-x^2 + 1} + 1) - \frac{1}{2} \log(\sqrt{-x^2 + 1} - 1)$$

```
[In] integrate(arcsin(x)*log(x),x, algorithm="fricas")
```

```
[Out] x*arcsin(x)*log(x) - x*arcsin(x) + sqrt(-x^2 + 1)*(log(x) - 2) + 1/2*log(sqrt(-x^2 + 1) + 1) - 1/2*log(sqrt(-x^2 + 1) - 1)
```

Sympy [A] (verification not implemented)

Time = 3.72 (sec) , antiderivative size = 102, normalized size of antiderivative = 2.00

$$\int \arcsin(x) \log(x) dx = x \log(x) \arcsin(x) - x \arcsin(x) + \sqrt{1 - x^2} \log(x) - \sqrt{1 - x^2} - \begin{cases} -\frac{x}{\sqrt{-1 + \frac{1}{x^2}}} - \operatorname{acosh}\left(\frac{1}{x}\right) + \frac{1}{x\sqrt{-1 + \frac{1}{x^2}}} & \text{for } \frac{1}{|x^2|} > 1 \\ \frac{ix}{\sqrt{1 - \frac{1}{x^2}}} + i \operatorname{asin}\left(\frac{1}{x}\right) - \frac{i}{x\sqrt{1 - \frac{1}{x^2}}} & \text{otherwise} \end{cases}$$

```
[In] integrate(asin(x)*ln(x),x)
```

```
[Out] x*log(x)*asin(x) - x*asin(x) + sqrt(1 - x**2)*log(x) - sqrt(1 - x**2) - Piecewise((-x/sqrt(-1 + x**(-2))) - acosh(1/x) + 1/(x*sqrt(-1 + x**(-2))), 1/Asin(x**2) > 1), (I*x/sqrt(1 - 1/x**2) + I*asin(1/x) - I/(x*sqrt(1 - 1/x**2))), True))
```

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.14

$$\int \arcsin(x) \log(x) dx = (x \log(x) - x) \arcsin(x) + \sqrt{-x^2 + 1} \log(x) - 2\sqrt{-x^2 + 1} + \log\left(\frac{2\sqrt{-x^2 + 1}}{|x|} + \frac{2}{|x|}\right)$$

[In] integrate(arcsin(x)*log(x),x, algorithm="maxima")

[Out] (x*log(x) - x)*arcsin(x) + sqrt(-x^2 + 1)*log(x) - 2*sqrt(-x^2 + 1) + log(2*sqrt(-x^2 + 1)/abs(x) + 2/abs(x))

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(42) = 84.

Time = 0.28 (sec) , antiderivative size = 272, normalized size of antiderivative = 5.33

$$\int \arcsin(x) \log(x) dx = x \arcsin(x) \log(x) + \sqrt{-x^2 + 1} \log(x) - \frac{2x \arcsin(x)}{(\sqrt{-x^2 + 1} + 1) \left(\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1 \right)} + \frac{x^2 \log(\sqrt{-x^2 + 1} + 1)}{(\sqrt{-x^2 + 1} + 1)^2 \left(\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1 \right)} + \frac{\log(\sqrt{-x^2 + 1} + 1)}{\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1} - \frac{x^2 \log(|x|)}{(\sqrt{-x^2 + 1} + 1)^2 \left(\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1 \right)} - \frac{\log(|x|)}{\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1} + \frac{2x^2}{(\sqrt{-x^2 + 1} + 1)^2 \left(\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1 \right)} - \frac{2}{\frac{x^2}{(\sqrt{-x^2 + 1} + 1)^2} + 1}$$

[In] integrate(arcsin(x)*log(x),x, algorithm="giac")

[Out] x*arcsin(x)*log(x) + sqrt(-x^2 + 1)*log(x) - 2*x*arcsin(x)/((sqrt(-x^2 + 1) + 1)*(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1)) + x^2*log(sqrt(-x^2 + 1) + 1)/((sqrt(-x^2 + 1) + 1)^2*(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1)) + log(sqrt(-x^2 + 1) + 1)/(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1) - x^2*log(abs(x))/((sqrt(-x^2 + 1) + 1)

)^2*(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1)) - log(abs(x))/(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1) + 2*x^2/((sqrt(-x^2 + 1) + 1)^2*(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1)) - 2/(x^2/(sqrt(-x^2 + 1) + 1)^2 + 1)

Mupad [F(-1)]

Timed out.

$$\int \arcsin(x) \log(x) dx = \int \operatorname{asin}(x) \ln(x) dx$$

[In] int(asin(x)*log(x),x)

[Out] int(asin(x)*log(x), x)

3.2 $\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$

Optimal result	47
Rubi [A] (verified)	47
Mathematica [A] (verified)	48
Maple [A] (verified)	48
Fricas [A] (verification not implemented)	48
Sympy [A] (verification not implemented)	49
Maxima [A] (verification not implemented)	49
Giac [A] (verification not implemented)	49
Mupad [F(-1)]	49

Optimal result

Integrand size = 15, antiderivative size = 17

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

[Out] x-arcsin(x)*(-x^2+1)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {4767, 8}

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

[In] Int[(x*ArcSin[x])/Sqrt[1 - x^2],x]

[Out] x - Sqrt[1 - x^2]*ArcSin[x]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 4767

Int[((a_.) + ArcSin[(c_.)*(x_)]*(b_.))^(n_.)*(x_)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x^2)^(p + 1)*((a + b*ArcSin[c*x])^n/(2*e*(p + 1))), x] + Dist[b*(n/(2*c*(p + 1)))*Simp[(d + e*x^2)^p/(1 - c^2*x^2)^p], Int[(1 - c^2*x^2)^(p + 1/2)*(a + b*ArcSin[c*x])^(n - 1), x], x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= -\sqrt{1-x^2} \arcsin(x) + \int 1 dx \\ &= x - \sqrt{1-x^2} \arcsin(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

[In] Integrate[(x*ArcSin[x])/Sqrt[1 - x^2],x]

[Out] x - Sqrt[1 - x^2]*ArcSin[x]

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.94

method	result	size
default	$x - \arcsin(x) \sqrt{-x^2 + 1}$	16

[In] int(x*arcsin(x)/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] x-arcsin(x)*(-x^2+1)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2 + 1} \arcsin(x) + x$$

[In] integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -sqrt(-x^2 + 1)*arcsin(x) + x

Sympy [A] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = x - \sqrt{1-x^2} \arcsin(x)$$

[In] integrate(x*asin(x)/(-x**2+1)**(1/2),x)

[Out] x - sqrt(1 - x**2)*asin(x)

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1} \arcsin(x) + x$$

[In] integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(-x^2 + 1)*arcsin(x) + x

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1} \arcsin(x) + x$$

[In] integrate(x*arcsin(x)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] -sqrt(-x^2 + 1)*arcsin(x) + x

Mupad [F(-1)]

Timed out.

$$\int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx = \int \frac{x \arcsin(x)}{\sqrt{1-x^2}} dx$$

[In] int((x*asin(x))/(1 - x^2)^(1/2),x)

[Out] int((x*asin(x))/(1 - x^2)^(1/2), x)

3.3 $\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	50
Rubi [F]	50
Mathematica [A] (verified)	51
Maple [C] (verified)	51
Fricas [A] (verification not implemented)	52
Sympy [F]	52
Maxima [A] (verification not implemented)	52
Giac [F]	53
Mupad [F(-1)]	53

Optimal result

Integrand size = 18, antiderivative size = 69

$$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx = \frac{(\sqrt{x} + 3\sqrt{1+x})\sqrt{-x + \sqrt{x}\sqrt{1+x}}}{4\sqrt{2}} - \left(\frac{3}{8} + x\right)\arcsin(\sqrt{x} - \sqrt{1+x})$$

[Out] $-(3/8+x)*\arcsin(x^{(1/2)}-(1+x)^{(1/2)})+1/8*(x^{(1/2)}+3*(1+x)^{(1/2)})*(-x+x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*2^{(1/2)}$

Rubi [F]

$$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx = \int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx$$

[In] `Int[-ArcSin[Sqrt[x] - Sqrt[1 + x]], x]`

[Out] `-(x*ArcSin[Sqrt[x] - Sqrt[1 + x]]) + Defer[Subst][Defer[Int][Sqrt[1 - x^2 + x*Sqrt[-1 + x^2]], x], x, Sqrt[1 + x]]/Sqrt[2]`

Rubi steps

$$\begin{aligned} \text{integral} &= -x \arcsin(\sqrt{x} - \sqrt{1+x}) + \int \frac{\sqrt{-x + \sqrt{x}\sqrt{1+x}}}{2\sqrt{2}\sqrt{1+x}} dx \\ &= -x \arcsin(\sqrt{x} - \sqrt{1+x}) + \frac{\int \frac{\sqrt{-x + \sqrt{x}\sqrt{1+x}}}{\sqrt{1+x}} dx}{2\sqrt{2}} \\ &= -x \arcsin(\sqrt{x} - \sqrt{1+x}) + \frac{\text{Subst}\left(\int \sqrt{1 - x^2 + x\sqrt{-1 + x^2}} dx, x, \sqrt{1+x}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.70 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.54

$$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx = \frac{1}{8}(\sqrt{x} + 3\sqrt{1+x})\sqrt{-2x + 2\sqrt{x}\sqrt{1+x}} - x \arcsin(\sqrt{x} - \sqrt{1+x}) - \frac{3}{8} \arctan\left(\frac{\sqrt{-2x + 2\sqrt{x}\sqrt{1+x}}}{-\sqrt{x} + \sqrt{1+x}}\right)$$

[In] Integrate[-ArcSin[Sqrt[x] - Sqrt[1 + x]],x]

[Out] ((Sqrt[x] + 3*Sqrt[1 + x])*Sqrt[-2*x + 2*Sqrt[x]*Sqrt[1 + x]])/8 - x*ArcSin[Sqrt[x] - Sqrt[1 + x]] - (3*ArcTan[Sqrt[-2*x + 2*Sqrt[x]*Sqrt[1 + x]]/(-Sqrt[x] + Sqrt[1 + x]))/8

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 1.10 (sec) , antiderivative size = 252, normalized size of antiderivative = 3.65

method	result
default	$-\frac{(2 \arcsin(\sqrt{x}-\sqrt{1+x})+i)\left(2i\sqrt{-2x+2\sqrt{x}\sqrt{1+x}}\sqrt{1+x}-2i\sqrt{-2x+2\sqrt{x}\sqrt{1+x}}\sqrt{x}-4\sqrt{x}\sqrt{1+x}+4x+1\right)}{32} - \frac{(2i\sqrt{-2x+2\sqrt{x}\sqrt{1+x}})}{32}$

[In] int(-arcsin(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)

[Out]
$$-1/32*(2*\arcsin(x^{(1/2)}-(1+x)^{(1/2)})+I)*(2*I*(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*(1+x)^{(1/2)}-2*I*(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*x^{(1/2)}-4*x^{(1/2)}*(1+x)^{(1/2)}+4*x+1)-1/32*(2*I*(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*x^{(1/2)}-2*I*(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*(1+x)^{(1/2)}+4*x-4*x^{(1/2)}*(1+x)^{(1/2)}+1)*(-I+2*\arcsin(x^{(1/2)}-(1+x)^{(1/2)}))-1/4*(2*x^{(1/2)}*(1+x)^{(1/2)}+2*x+1)*(2*I*x^{(1/2)}*(1+x)^{(1/2)}-2*I*x-(1+x)^{(1/2)}*(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}+(-2*x+2*x^{(1/2)}*(1+x)^{(1/2)})^{(1/2)}*x^{(1/2)}+\arcsin(x^{(1/2)}-(1+x)^{(1/2)})-I)$$

Fricas [A] (verification not implemented)

none

Time = 23.67 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.71

$$\int -\arcsin\left(\sqrt{x}-\sqrt{1+x}\right) dx = \frac{1}{8}(8x+3)\arcsin\left(\sqrt{x+1}-\sqrt{x}\right) + \frac{1}{8}\sqrt{2\sqrt{x+1}\sqrt{x}-2x}\left(3\sqrt{x+1}+\sqrt{x}\right)$$

[In] integrate(-arcsin(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")

[Out] 1/8*(8*x + 3)*arcsin(sqrt(x + 1) - sqrt(x)) + 1/8*sqrt(2*sqrt(x + 1)*sqrt(x) - 2*x)*(3*sqrt(x + 1) + sqrt(x))

Sympy [F]

$$\int -\arcsin\left(\sqrt{x}-\sqrt{1+x}\right) dx = -\int \operatorname{asin}\left(\sqrt{x}-\sqrt{x+1}\right) dx$$

[In] integrate(-asin(x**(1/2)-(1+x)**(1/2)),x)

[Out] -Integral(asin(sqrt(x) - sqrt(x + 1)), x)

Maxima [A] (verification not implemented)

none

Time = 0.91 (sec) , antiderivative size = 4, normalized size of antiderivative = 0.06

$$\int -\arcsin\left(\sqrt{x}-\sqrt{1+x}\right) dx = \frac{1}{2}\pi x$$

[In] integrate(-arcsin(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")

[Out] 1/2*pi*x

Giac [F]

$$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx = \int -\arcsin(-\sqrt{x+1} + \sqrt{x}) dx$$

[In] integrate(-arcsin(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")

[Out] integrate(-arcsin(-sqrt(x + 1) + sqrt(x)), x)

Mupad [F(-1)]

Timed out.

$$\int -\arcsin(\sqrt{x} - \sqrt{1+x}) dx = \int \operatorname{asin}(\sqrt{x+1} - \sqrt{x}) dx$$

[In] int(asin((x + 1)^(1/2) - x^(1/2)),x)

[Out] int(asin((x + 1)^(1/2) - x^(1/2)), x)

3.4 $\int \log \left(1 + x\sqrt{1 + x^2} \right) dx$

Optimal result	54
Rubi [B] (verified)	55
Mathematica [A] (verified)	60
Maple [B] (verified)	61
Fricas [B] (verification not implemented)	62
Sympy [F(-1)]	63
Maxima [F]	63
Giac [B] (verification not implemented)	63
Mupad [B] (verification not implemented)	65

Optimal result

Integrand size = 14, antiderivative size = 97

$$\int \log \left(1 + x\sqrt{1 + x^2} \right) dx = -2x + \sqrt{2(1 + \sqrt{5})} \arctan \left(\sqrt{-2 + \sqrt{5}}(x + \sqrt{1 + x^2}) \right) \\ - \sqrt{2(-1 + \sqrt{5})} \operatorname{arctanh} \left(\sqrt{2 + \sqrt{5}}(x + \sqrt{1 + x^2}) \right) \\ + x \log \left(1 + x\sqrt{1 + x^2} \right)$$

[Out] -2*x+x*ln(1+x*(x^2+1)^(1/2))-arctanh((x+(x^2+1)^(1/2))*(2+5^(1/2))^(1/2))*(-2+2*5^(1/2))^(1/2)+arctan((x+(x^2+1)^(1/2))*(-2+5^(1/2))^(1/2))*(2+2*5^(1/2))^(1/2)

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 332 vs. $2(97) = 194$.

Time = 0.50 (sec) , antiderivative size = 332, normalized size of antiderivative = 3.42, number of steps used = 32, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$, Rules used = {2628, 6874, 267, 1144, 209, 213, 1265, 838, 721, 1107, 1180, 1261, 713}

$$\begin{aligned} \int \log(1 + x\sqrt{1+x^2}) dx = & \sqrt{\frac{2}{5}}(\sqrt{5}-1) \arctan\left(\sqrt{\frac{2}{\sqrt{5}-1}}\sqrt{x^2+1}\right) \\ & + \sqrt{\frac{2}{5(\sqrt{5}-1)}} \arctan\left(\sqrt{\frac{2}{\sqrt{5}-1}}\sqrt{x^2+1}\right) \\ & + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\ & - \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\ & - \sqrt{\frac{2}{5}(1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{x^2+1}\right) \\ & + \sqrt{\frac{2}{5(1+\sqrt{5})}} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{x^2+1}\right) \\ & + \sqrt{\frac{1}{10}(\sqrt{5}-1)} \operatorname{arctanh}\left(\sqrt{\frac{2}{\sqrt{5}-1}}x\right) \\ & + 2\sqrt{\frac{1}{5}(\sqrt{5}-2)} \operatorname{arctanh}\left(\sqrt{\frac{2}{\sqrt{5}-1}}x\right) \\ & + x \log(\sqrt{x^2+1}x+1) - 2x \end{aligned}$$

[In] Int[Log[1 + x*Sqrt[1 + x^2]],x]

[Out] -2*x - Sqrt[(1 + Sqrt[5])/10]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x] + 2*Sqrt[(2 + Sqrt[5])/5]*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x] + Sqrt[2/(5*(-1 + Sqrt[5]))]*ArcTan[Sqrt[2/(-1 + Sqrt[5])]*Sqrt[1 + x^2]] + Sqrt[(2*(-1 + Sqrt[5]))/5]*ArcTan[Sqrt[2/(-1 + Sqrt[5])]*Sqrt[1 + x^2]] + 2*Sqrt[(-2 + Sqrt[5])/5]*ArcTanh[Sqrt[2/(-1 + Sqrt[5])]*x] + Sqrt[(-1 + Sqrt[5])/10]*ArcTanh[Sqrt[2/(-1 + Sqrt[5])]*x] + Sqrt[2/(5*(1 + Sqrt[5]))]*ArcTanh[Sqrt[2/(1 + Sqrt[5])]*Sqrt[1 + x^2]] - Sqrt[(2*(1 + Sqrt[5]))/5]*ArcTanh[Sqrt[2/(1 + Sqrt[5])]*Sqrt[1 + x^2]] + x*Log[1 + x*Sqrt[1 + x^2]]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 267

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]

Rule 713

Int[Sqrt[(d_) + (e_)*(x_)]/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[2*e, Subst[Int[x^2/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0]

Rule 721

Int[1/(Sqrt[(d_) + (e_)*(x_)]*((a_) + (b_)*(x_) + (c_)*(x_)^2)), x_Symbol] := Dist[2*e, Subst[Int[1/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0]

Rule 838

Int[(((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_)))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[g*((d + e*x)^m/(c*m)), x] + Dist[1/c, Int[(d + e*x)^(m - 1)*(Simp[c*d*f - a*e*g + (g*c*d - b*e*g + c*e*f)*x, x]/(a + b*x + c*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && FractionQ[m] && GtQ[m, 0]

Rule 1107

Int[((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(-1), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[c/q, Int[1/(b/2 - q/2 + c*x^2), x], x] - Dist[c/q, Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[b^2 - 4*a*c]

Rule 1144

Int[((d_)*(x_))^(m_)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[(d^2/2)*(b/q + 1), Int[(d*x)^(m - 2)/(b/2

+ q/2 + c*x^2), x], x] - Dist[(d^2/2)*(b/q - 1), Int[(d*x)^(m - 2)/(b/2 - q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b^2 - 4*a*c, 0] && GeQ[m, 2]

Rule 1180

Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[e/2 + (2*c*d - b*e)/(2*q), Int[1/(b/2 - q/2 + c*x^2), x], x] + Dist[e/2 - (2*c*d - b*e)/(2*q), Int[1/(b/2 + q/2 + c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]

Rule 1261

Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 2628

Int[Log[u_], x_Symbol] :> Simp[x*Log[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[u, x]

Rule 6874

Int[u_, x_Symbol] :> With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \text{integral} &= x \log \left(1 + x\sqrt{1+x^2} \right) - \int \frac{x(1+2x^2)}{x+x^3+\sqrt{1+x^2}} dx \\
 &= x \log \left(1 + x\sqrt{1+x^2} \right) - \int \left(\frac{x}{x+x^3+\sqrt{1+x^2}} + \frac{2x^3}{x+x^3+\sqrt{1+x^2}} \right) dx \\
 &= x \log \left(1 + x\sqrt{1+x^2} \right) - 2 \int \frac{x^3}{x+x^3+\sqrt{1+x^2}} dx - \int \frac{x}{x+x^3+\sqrt{1+x^2}} dx
 \end{aligned}$$

$$\begin{aligned}
&= x \log \left(1 + x\sqrt{1+x^2} \right) \\
&\quad - 2 \int \left(1 - \frac{x}{\sqrt{1+x^2}} + \frac{1-x^2}{-1+x^2+x^4} - \frac{x\sqrt{1+x^2}}{-1+x^2+x^4} + \frac{x^3\sqrt{1+x^2}}{-1+x^2+x^4} \right) dx \\
&\quad - \int \left(\frac{x}{\sqrt{1+x^2}} + \frac{x^2}{-1+x^2+x^4} - \frac{x^3\sqrt{1+x^2}}{-1+x^2+x^4} \right) dx \\
&= -2x + x \log \left(1 + x\sqrt{1+x^2} \right) + 2 \int \frac{x}{\sqrt{1+x^2}} dx - 2 \int \frac{1-x^2}{-1+x^2+x^4} dx \\
&\quad + 2 \int \frac{x\sqrt{1+x^2}}{-1+x^2+x^4} dx - 2 \int \frac{x^3\sqrt{1+x^2}}{-1+x^2+x^4} dx \\
&\quad - \int \frac{x}{\sqrt{1+x^2}} dx - \int \frac{x^2}{-1+x^2+x^4} dx + \int \frac{x^3\sqrt{1+x^2}}{-1+x^2+x^4} dx \\
&= -2x + \sqrt{1+x^2} + x \log \left(1 + x\sqrt{1+x^2} \right) + \frac{1}{2} \text{Subst} \left(\int \frac{x\sqrt{1+x}}{-1+x+x^2} dx, x, x^2 \right) \\
&\quad + \frac{1}{10} (-5 + \sqrt{5}) \int \frac{1}{\frac{1}{2} - \frac{\sqrt{5}}{2} + x^2} dx - \frac{1}{10} (5 + \sqrt{5}) \int \frac{1}{\frac{1}{2} + \frac{\sqrt{5}}{2} + x^2} dx \\
&\quad - \frac{1}{5} (-5 + 3\sqrt{5}) \int \frac{1}{\frac{1}{2} - \frac{\sqrt{5}}{2} + x^2} dx + \frac{1}{5} (5 + 3\sqrt{5}) \int \frac{1}{\frac{1}{2} + \frac{\sqrt{5}}{2} + x^2} dx \\
&\quad + \text{Subst} \left(\int \frac{\sqrt{1+x}}{-1+x+x^2} dx, x, x^2 \right) - \text{Subst} \left(\int \frac{x\sqrt{1+x}}{-1+x+x^2} dx, x, x^2 \right) \\
&= -2x - \sqrt{\frac{1}{10} (1 + \sqrt{5})} \arctan \left(\sqrt{\frac{2}{1 + \sqrt{5}}} x \right) \\
&\quad + 2\sqrt{\frac{1}{5} (2 + \sqrt{5})} \arctan \left(\sqrt{\frac{2}{1 + \sqrt{5}}} x \right) \\
&\quad + 2\sqrt{\frac{1}{5} (-2 + \sqrt{5})} \operatorname{arctanh} \left(\sqrt{\frac{2}{-1 + \sqrt{5}}} x \right) \\
&\quad + \sqrt{\frac{1}{10} (-1 + \sqrt{5})} \operatorname{arctanh} \left(\sqrt{\frac{2}{-1 + \sqrt{5}}} x \right) + x \log \left(1 + x\sqrt{1+x^2} \right) \\
&\quad + \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1+x} (-1+x+x^2)} dx, x, x^2 \right) \\
&\quad + 2 \text{Subst} \left(\int \frac{x^2}{-1-x^2+x^4} dx, x, \sqrt{1+x^2} \right) \\
&\quad - \text{Subst} \left(\int \frac{1}{\sqrt{1+x} (-1+x+x^2)} dx, x, x^2 \right)
\end{aligned}$$

$$\begin{aligned}
&= -2x - \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + 2\sqrt{\frac{1}{5}(-2+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{1}{10}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad + x \log(1+x\sqrt{1+x^2}) - 2\operatorname{Subst}\left(\int \frac{1}{-1-x^2+x^4} dx, x, \sqrt{1+x^2}\right) \\
&\quad + \frac{1}{5}(5-\sqrt{5}) \operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}+\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right) \\
&\quad + \frac{1}{5}(5+\sqrt{5}) \operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}-\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right) \\
&\quad + \operatorname{Subst}\left(\int \frac{1}{-1-x^2+x^4} dx, x, \sqrt{1+x^2}\right) \\
&= -2x - \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{2}{5}(-1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{-1+\sqrt{5}}}\sqrt{1+x^2}\right) \\
&\quad + 2\sqrt{\frac{1}{5}(-2+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{1}{10}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad - \sqrt{\frac{2}{5}(1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{1+x^2}\right) + x \log(1+x\sqrt{1+x^2}) \\
&\quad + \frac{\operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}-\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right)}{\sqrt{5}} - \frac{\operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}+\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right)}{\sqrt{5}} \\
&\quad - \frac{2\operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}-\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right)}{\sqrt{5}} + \frac{2\operatorname{Subst}\left(\int \frac{1}{-\frac{1}{2}+\frac{\sqrt{5}}{2}+x^2} dx, x, \sqrt{1+x^2}\right)}{\sqrt{5}}
\end{aligned}$$

$$\begin{aligned}
&= -2x - \sqrt{\frac{1}{10}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + 2\sqrt{\frac{1}{5}(2+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{2}{5(-1+\sqrt{5})}} \arctan\left(\sqrt{\frac{2}{-1+\sqrt{5}}}\sqrt{1+x^2}\right) \\
&\quad + \sqrt{\frac{2}{5}(-1+\sqrt{5})} \arctan\left(\sqrt{\frac{2}{-1+\sqrt{5}}}\sqrt{1+x^2}\right) \\
&\quad + 2\sqrt{\frac{1}{5}(-2+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{1}{10}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right) \\
&\quad + \sqrt{\frac{2}{5(1+\sqrt{5})}} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{1+x^2}\right) \\
&\quad - \sqrt{\frac{2}{5}(1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{1+x^2}\right) + x \log(1+x\sqrt{1+x^2})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.94

$$\begin{aligned}
\int \log(1+x\sqrt{1+x^2}) dx &= -2x + \frac{(5+\sqrt{5}) \arctan\left(\sqrt{\frac{2}{1+\sqrt{5}}}x\right)}{\sqrt{10}(1+\sqrt{5})} \\
&\quad + \sqrt{\frac{1}{2}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{1}{2} + \frac{\sqrt{5}}{2}}\sqrt{1+x^2}\right) \\
&\quad - \frac{(-5+\sqrt{5}) \operatorname{arctanh}\left(\sqrt{\frac{2}{-1+\sqrt{5}}}x\right)}{\sqrt{10}(-1+\sqrt{5})} \\
&\quad - \sqrt{\frac{1}{2}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{-\frac{1}{2} + \frac{\sqrt{5}}{2}}\sqrt{1+x^2}\right) \\
&\quad + x \log(1+x\sqrt{1+x^2})
\end{aligned}$$

[In] Integrate[Log[1 + x*sqrt[1 + x^2]],x]

```
[Out] -2*x + ((5 + Sqrt[5])*ArcTan[Sqrt[2/(1 + Sqrt[5])]*x])/Sqrt[10*(1 + Sqrt[5]]
) + Sqrt[(1 + Sqrt[5])/2]*ArcTan[Sqrt[1/2 + Sqrt[5]/2]*Sqrt[1 + x^2]] - ((
-5 + Sqrt[5])*ArcTanh[Sqrt[2/(-1 + Sqrt[5])]*x])/Sqrt[10*(-1 + Sqrt[5])] -
Sqrt[(-1 + Sqrt[5])/2]*ArcTanh[Sqrt[-1/2 + Sqrt[5]/2]*Sqrt[1 + x^2]] + x*Lo
g[1 + x*Sqrt[1 + x^2]]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. $2(75) = 150$.

Time = 0.18 (sec) , antiderivative size = 413, normalized size of antiderivative = 4.26

method	result
parts	$x \ln(1 + x\sqrt{x^2 + 1}) - 2x + \frac{(3+\sqrt{5})\sqrt{5} \arctan\left(\frac{2x}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}} - \frac{(\sqrt{5}-3)\sqrt{5} \operatorname{arctanh}\left(\frac{2x}{\sqrt{-2+2\sqrt{5}}}\right)}{5\sqrt{-2+2\sqrt{5}}} - \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{x^2+1}}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}}$
default	$x \ln(1 + x\sqrt{x^2 + 1}) - \frac{(\sqrt{5}+1)\sqrt{5} \arctan\left(\frac{2x}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}} + \frac{(\sqrt{5}-1)\sqrt{5} \operatorname{arctanh}\left(\frac{2x}{\sqrt{-2+2\sqrt{5}}}\right)}{5\sqrt{-2+2\sqrt{5}}} - 2x + \frac{2(3+\sqrt{5})\sqrt{5} \operatorname{arctanh}\left(\frac{\sqrt{x^2+1}}{\sqrt{2+2\sqrt{5}}}\right)}{5\sqrt{2+2\sqrt{5}}}$

```
[In] int(ln(1+x*(x^2+1)^(1/2)),x,method=_RETURNVERBOSE)
```

```
[Out] x*ln(1+x*(x^2+1)^(1/2))-2*x+1/5*(3+5^(1/2))*5^(1/2)/(2+2*5^(1/2))^(1/2)*arc
tan(2*x/(2+2*5^(1/2))^(1/2))-1/5*(5^(1/2)-3)*5^(1/2)/(-2+2*5^(1/2))^(1/2)*a
rctanh(2*x/(-2+2*5^(1/2))^(1/2))-1/5*5^(1/2)/(2+5^(1/2))^(1/2)*arctanh(((x^
2+1)^(1/2)-x)/(2+5^(1/2))^(1/2))-1/5*5^(1/2)/(-2+5^(1/2))^(1/2)*arctan(((x^
2+1)^(1/2)-x)/(-2+5^(1/2))^(1/2))-1/5*(-2+5^(1/2))^(1/2)*5^(1/2)*arctanh(((
x^2+1)^(1/2)-x)/(-2+5^(1/2))^(1/2))+1/5*(2+5^(1/2))^(1/2)*5^(1/2)*arctan(((
x^2+1)^(1/2)-x)/(2+5^(1/2))^(1/2))+2/5*5^(1/2)/(2+2*5^(1/2))^(1/2)*arctan(2
*x/(2+2*5^(1/2))^(1/2))+2/5*5^(1/2)/(-2+2*5^(1/2))^(1/2)*arctanh(2*x/(-2+2*
5^(1/2))^(1/2))-1/10*(3+5^(1/2))*5^(1/2)/(2+5^(1/2))^(1/2)*arctanh(((x^2+1)
^(1/2)-x)/(2+5^(1/2))^(1/2))+1/10*(5^(1/2)-3)*5^(1/2)/(-2+5^(1/2))^(1/2)*ar
ctan(((x^2+1)^(1/2)-x)/(-2+5^(1/2))^(1/2))-1/10*(5^(1/2)-1)*5^(1/2)/(-2+5^(
1/2))^(1/2)*arctanh(((x^2+1)^(1/2)-x)/(-2+5^(1/2))^(1/2))+1/10*(5^(1/2)+1)*
5^(1/2)/(2+5^(1/2))^(1/2)*arctan(((x^2+1)^(1/2)-x)/(2+5^(1/2))^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 451 vs. 2(75) = 150.

Time = 0.26 (sec) , antiderivative size = 451, normalized size of antiderivative = 4.65

$$\begin{aligned}
\int \log(1 + x\sqrt{1+x^2}) dx = & -\frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log\left(4x^2 \right. \\
& - \sqrt{x^2+1} \left((\sqrt{5}\sqrt{2}-\sqrt{2}) \sqrt{-\sqrt{5}-1+4x} \right. \\
& \left. \left. + (\sqrt{5}\sqrt{2}x-\sqrt{2}x) \sqrt{-\sqrt{5}-1+4} \right) \right) \\
& + \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log\left(4x^2 \right. \\
& + \sqrt{x^2+1} \left((\sqrt{5}\sqrt{2}-\sqrt{2}) \sqrt{-\sqrt{5}-1-4x} \right. \\
& \left. \left. - (\sqrt{5}\sqrt{2}x-\sqrt{2}x) \sqrt{-\sqrt{5}-1+4} \right) \right) \\
& + \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log\left(4x^2 - 4\sqrt{x^2+1}x \right. \\
& \left. + (\sqrt{5}\sqrt{2}x - \sqrt{x^2+1}(\sqrt{5}\sqrt{2} + \sqrt{2}) + \sqrt{2}x) \sqrt{\sqrt{5}-1+4} \right) \\
& - \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log\left(4x^2 - 4\sqrt{x^2+1}x \right. \\
& \left. - (\sqrt{5}\sqrt{2}x - \sqrt{x^2+1}(\sqrt{5}\sqrt{2} + \sqrt{2}) + \sqrt{2}x) \sqrt{\sqrt{5}-1+4} \right) \\
& + x \log(\sqrt{x^2+1}x + 1) \\
& + \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log\left(2x + \sqrt{2} \sqrt{\sqrt{5}-1}\right) \\
& - \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log\left(2x - \sqrt{2} \sqrt{\sqrt{5}-1}\right) \\
& + \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log\left(2x + \sqrt{2} \sqrt{-\sqrt{5}-1}\right) \\
& - \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log\left(2x - \sqrt{2} \sqrt{-\sqrt{5}-1}\right) - 2x
\end{aligned}$$

[In] integrate(log(1+x*(x^2+1)^(1/2)),x, algorithm="fricas")

[Out] -1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log(4*x^2 - sqrt(x^2 + 1)*((sqrt(5)*sqrt(2) - sqrt(2))*sqrt(-sqrt(5) - 1) + 4*x) + (sqrt(5)*sqrt(2)*x - sqrt(2)*x)*sqrt(-sqrt(5) - 1) + 4) + 1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log(4*x^2 + sqrt(x^2 + 1)*((sqrt(5)*sqrt(2) - sqrt(2))*sqrt(-sqrt(5) - 1) - 4*x) - (sqrt(5)*sqrt(2)*x - sqrt(2)*x)*sqrt(-sqrt(5) - 1) + 4) - 2*x

```
(2)*x - sqrt(2)*x)*sqrt(-sqrt(5) - 1) + 4) + 1/4*sqrt(2)*sqrt(sqrt(5) - 1)*
log(4*x^2 - 4*sqrt(x^2 + 1)*x + (sqrt(5)*sqrt(2)*x - sqrt(x^2 + 1)*(sqrt(5)
*sqrt(2) + sqrt(2)) + sqrt(2)*x)*sqrt(sqrt(5) - 1) + 4) - 1/4*sqrt(2)*sqrt(
sqrt(5) - 1)*log(4*x^2 - 4*sqrt(x^2 + 1)*x - (sqrt(5)*sqrt(2)*x - sqrt(x^2
+ 1)*(sqrt(5)*sqrt(2) + sqrt(2)) + sqrt(2)*x)*sqrt(sqrt(5) - 1) + 4) + x*lo
g(sqrt(x^2 + 1)*x + 1) + 1/4*sqrt(2)*sqrt(sqrt(5) - 1)*log(2*x + sqrt(2)*sq
rt(sqrt(5) - 1)) - 1/4*sqrt(2)*sqrt(sqrt(5) - 1)*log(2*x - sqrt(2)*sqrt(sqr
t(5) - 1)) + 1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log(2*x + sqrt(2)*sqrt(-sqrt(5)
- 1)) - 1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log(2*x - sqrt(2)*sqrt(-sqrt(5) - 1
)) - 2*x
```

Sympy [F(-1)]

Timed out.

$$\int \log(1 + x\sqrt{1 + x^2}) dx = \text{Timed out}$$

```
[In] integrate(ln(1+x*(x**2+1)**(1/2)),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \log(1 + x\sqrt{1 + x^2}) dx = \int \log(\sqrt{x^2 + 1}x + 1) dx$$

```
[In] integrate(log(1+x*(x^2+1)^(1/2)),x, algorithm="maxima")
```

```
[Out] x*log(sqrt(x^2 + 1)*x + 1) - 2*x + arctan(x) + integrate((2*x^2 + 1)/(x^2 +
(x^3 + x)*sqrt(x^2 + 1) + 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 235 vs. $2(75) = 150$.

Time = 0.40 (sec) , antiderivative size = 235, normalized size of antiderivative = 2.42

$$\begin{aligned}
& \int \log \left(1 + x\sqrt{1+x^2} \right) dx \\
&= x \log \left(\sqrt{x^2+1}x + 1 \right) + \frac{1}{2} \sqrt{2\sqrt{5}+2} \arctan \left(-\frac{x - \sqrt{x^2+1} + \frac{1}{x-\sqrt{x^2+1}}}{\sqrt{2\sqrt{5}-2}} \right) \\
&+ \frac{1}{2} \sqrt{2\sqrt{5}+2} \arctan \left(\frac{x}{\sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}}} \right) \\
&- \frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(-x + \sqrt{x^2+1} + \sqrt{2\sqrt{5}+2} - \frac{1}{x - \sqrt{x^2+1}} \right) \\
&+ \frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(\left| x + \sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \right| \right) - \frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(\left| x - \sqrt{\frac{1}{2}\sqrt{5} - \frac{1}{2}} \right| \right) \\
&+ \frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(\left| -x + \sqrt{x^2+1} - \sqrt{2\sqrt{5}+2} - \frac{1}{x - \sqrt{x^2+1}} \right| \right) - 2x
\end{aligned}$$

[In] integrate(log(1+x*(x^2+1)^(1/2)),x, algorithm="giac")

[Out] x*log(sqrt(x^2 + 1)*x + 1) + 1/2*sqrt(2*sqrt(5) + 2)*arctan(-(x - sqrt(x^2 + 1) + 1/(x - sqrt(x^2 + 1)))/sqrt(2*sqrt(5) - 2)) + 1/2*sqrt(2*sqrt(5) + 2)*arctan(x/sqrt(1/2*sqrt(5) + 1/2)) - 1/4*sqrt(2*sqrt(5) - 2)*log(-x + sqrt(x^2 + 1) + sqrt(2*sqrt(5) + 2) - 1/(x - sqrt(x^2 + 1))) + 1/4*sqrt(2*sqrt(5) - 2)*log(abs(x + sqrt(1/2*sqrt(5) - 1/2))) - 1/4*sqrt(2*sqrt(5) - 2)*log(abs(x - sqrt(1/2*sqrt(5) - 1/2))) + 1/4*sqrt(2*sqrt(5) - 2)*log(abs(-x + sqrt(x^2 + 1) - sqrt(2*sqrt(5) + 2) - 1/(x - sqrt(x^2 + 1)))) - 2*x

Mupad [B] (verification not implemented)

Time = 1.50 (sec) , antiderivative size = 666, normalized size of antiderivative = 6.87

$$\begin{aligned}
& \int \log\left(1 + x\sqrt{1+x^2}\right) dx \\
&= x \ln\left(x\sqrt{x^2+1}+1\right) - 2x + \frac{\ln\left(x - \frac{\sqrt{2}\sqrt{\sqrt{5}-1}}{2}\right)\left(\frac{\sqrt{5}}{2} - \frac{5}{2}\right)}{2\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}} - \frac{\ln\left(x + \frac{\sqrt{2}\sqrt{\sqrt{5}-1}}{2}\right)\left(\frac{\sqrt{5}}{2} - \frac{5}{2}\right)}{2\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}} \\
&\quad - \frac{\ln\left(x - \frac{\sqrt{2}\sqrt{-\sqrt{5}-1}}{2}\right)\left(\frac{\sqrt{5}}{2} + \frac{5}{2}\right)}{2\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}} + \frac{\ln\left(x + \frac{\sqrt{2}\sqrt{-\sqrt{5}-1}}{2}\right)\left(\frac{\sqrt{5}}{2} + \frac{5}{2}\right)}{2\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}} \\
&\quad + \frac{\left(\ln\left(x - \frac{\sqrt{2}\sqrt{\sqrt{5}-1}}{2}\right) - \ln\left(\frac{\sqrt{2}x\sqrt{\sqrt{5}-1}}{2} + \frac{\sqrt{2}\sqrt{x^2+1}\sqrt{\sqrt{5}+1}}{2} + 1\right)\right)\left(\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 2\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)}{\left(2\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)\sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}} \\
&\quad + \frac{\left(\ln\left(x + \frac{\sqrt{2}\sqrt{\sqrt{5}-1}}{2}\right) - \ln\left(\frac{\sqrt{2}\sqrt{x^2+1}\sqrt{\sqrt{5}+1}}{2} - \frac{\sqrt{2}x\sqrt{\sqrt{5}-1}}{2} + 1\right)\right)\left(\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 2\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)}{\left(2\sqrt{\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)\sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}} \\
&\quad - \frac{\left(\ln\left(\frac{\sqrt{2}\sqrt{x^2+1}\sqrt{1-\sqrt{5}}}{2} - \frac{\sqrt{2}x\sqrt{-\sqrt{5}-1}}{2} + 1\right) - \ln\left(x + \frac{\sqrt{2}\sqrt{-\sqrt{5}-1}}{2}\right)\right)\left(\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 2\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)}{\left(2\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)\sqrt{\frac{1}{2} - \frac{\sqrt{5}}{2}}} \\
&\quad - \frac{\left(\ln\left(\frac{\sqrt{2}x\sqrt{-\sqrt{5}-1}}{2} + \frac{\sqrt{2}\sqrt{x^2+1}\sqrt{1-\sqrt{5}}}{2} + 1\right) - \ln\left(x - \frac{\sqrt{2}\sqrt{-\sqrt{5}-1}}{2}\right)\right)\left(\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 2\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)}{\left(2\sqrt{-\frac{\sqrt{5}}{2} - \frac{1}{2}} + 4\left(-\frac{\sqrt{5}}{2} - \frac{1}{2}\right)^{3/2}\right)\sqrt{\frac{1}{2} - \frac{\sqrt{5}}{2}}}
\end{aligned}$$

[In] int(log(x*(x^2 + 1)^(1/2) + 1),x)

[Out] x*log(x*(x^2 + 1)^(1/2) + 1) - 2*x + (log(x - (2^(1/2))*(5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 - 5/2)/(2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2)) - (log(x + (2^(1/2))*(5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 - 5/2)/(2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2)) - (log(x - (2^(1/2))*(-5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 + 5/2)/(2*(-5^(1/2)/2 - 1/2)^(1/2) + 4*(-5^(1/2)/2 - 1/2)^(3/2)) + (log(x + (2^(1/2))*(-5^(1/2) - 1)^(1/2))/2)*(5^(1/2)/2 + 5/2)/(2*(-5^(1/2)/2 - 1/2)^(1/2) + 4*(-5^(1/2)/2 - 1/2)^(3/2)) + ((log(x - (2^(1/2))*(5^(1/2) - 1)^(1/2))/2) - log((2^(1/2)*x*(5^(1/2) - 1)^(1/2))/2 + (2^(1/2)*(x^2 + 1)^(1/2)*(5^(1/2) + 1)^(1/2))/2 + 1))*((5^(1/2)/2 - 1/2)^(1/2) + 2*(5^(1/2)/2 - 1/2)^(3/2))/((2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2))*(5^(1/2)/2 + 1/2)^(1/2)) + ((log(x + (2^(1/2))*(5^(1/2) - 1)^(1/2))/2) - log((2^(1/2)*x*(5^(1/2) - 1)^(1/2))/2 + (2^(1/2)*(x^2 + 1)^(1/2)*(5^(1/2) + 1)^(1/2))/2 + 1))*((5^(1/2)/2 - 1/2)^(1/2) + 2*(5^(1/2)/2 - 1/2)^(3/2))/((2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(5^(1/2)/2 - 1/2)^(3/2))*(5^(1/2)/2 + 1/2)^(1/2)) + ((log(x - (2^(1/2))*(-5^(1/2) - 1)^(1/2))/2) - log((2^(1/2)*x*(-5^(1/2) - 1)^(1/2))/2 + (2^(1/2)*(x^2 + 1)^(1/2)*(-5^(1/2) - 1)^(1/2))/2 + 1))*((-5^(1/2)/2 - 1/2)^(1/2) + 2*(-5^(1/2)/2 - 1/2)^(3/2))/((-2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(-5^(1/2)/2 - 1/2)^(3/2))*(-5^(1/2)/2 - 1/2)^(1/2)) + ((log(x + (2^(1/2))*(-5^(1/2) - 1)^(1/2))/2) - log((2^(1/2)*x*(-5^(1/2) - 1)^(1/2))/2 + (2^(1/2)*(x^2 + 1)^(1/2)*(-5^(1/2) - 1)^(1/2))/2 + 1))*((-5^(1/2)/2 - 1/2)^(1/2) + 2*(-5^(1/2)/2 - 1/2)^(3/2))/((-2*(5^(1/2)/2 - 1/2)^(1/2) + 4*(-5^(1/2)/2 - 1/2)^(3/2))*(-5^(1/2)/2 - 1/2)^(1/2))

$$\begin{aligned}
& ((1/2) - 1)^{(1/2)}/2) - \log((2^{(1/2)}*(x^2 + 1)^{(1/2)}*(5^{(1/2)} + 1)^{(1/2)})/2 \\
& - (2^{(1/2)}*x*(5^{(1/2)} - 1)^{(1/2)})/2 + 1)) * ((5^{(1/2)}/2 - 1/2)^{(1/2)} + 2*(5^{(1/2)}/2 - 1/2)^{(3/2)}) / ((2*(5^{(1/2)}/2 - 1/2)^{(1/2)} + 4*(5^{(1/2)}/2 - 1/2)^{(3/2)}) * (5^{(1/2)}/2 + 1/2)^{(1/2)}) - ((\log((2^{(1/2)}*(x^2 + 1)^{(1/2)}*(1 - 5^{(1/2)})^{(1/2)})/2 - (2^{(1/2)}*x*(- 5^{(1/2)} - 1)^{(1/2)})/2 + 1) - \log(x + (2^{(1/2)}*(- 5^{(1/2)} - 1)^{(1/2)})/2)) * ((- 5^{(1/2)}/2 - 1/2)^{(1/2)} + 2*(- 5^{(1/2)}/2 - 1/2)^{(3/2)})) / ((2*(- 5^{(1/2)}/2 - 1/2)^{(1/2)} + 4*(- 5^{(1/2)}/2 - 1/2)^{(3/2)}) * (1/2 - 5^{(1/2)}/2)^{(1/2)}) - ((\log((2^{(1/2)}*x*(- 5^{(1/2)} - 1)^{(1/2)})/2 + (2^{(1/2)}*(x^2 + 1)^{(1/2)}*(1 - 5^{(1/2)})^{(1/2)})/2 + 1) - \log(x - (2^{(1/2)}*(- 5^{(1/2)} - 1)^{(1/2)})/2)) * ((- 5^{(1/2)}/2 - 1/2)^{(1/2)} + 2*(- 5^{(1/2)}/2 - 1/2)^{(3/2)})) / ((2*(- 5^{(1/2)}/2 - 1/2)^{(1/2)} + 4*(- 5^{(1/2)}/2 - 1/2)^{(3/2)}) * (1/2 - 5^{(1/2)}/2)^{(1/2)})
\end{aligned}$$

$$3.5 \quad \int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx$$

Optimal result	67
Rubi [C] (warning: unable to verify)	67
Mathematica [C] (verified)	69
Maple [C] (verified)	70
Fricas [A] (verification not implemented)	70
Sympy [F(-1)]	70
Maxima [F]	71
Giac [F]	71
Mupad [F(-1)]	71

Optimal result

Integrand size = 19, antiderivative size = 45

$$\int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx = \frac{x}{3} + \frac{1}{3} \arctan\left(\frac{\cos(x)(1+\cos^2(x))\sin(x)}{1+\cos^2(x)\sqrt{1+\cos^2(x)+\cos^4(x)}}\right)$$

[Out] 1/3*x+1/3*arctan(cos(x)*(1+cos(x)^2)*sin(x)/(1+cos(x)^2*(1+cos(x)^2+cos(x)^4)^(1/2)))

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 0.38 (sec) , antiderivative size = 289, normalized size of antiderivative = 6.42, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {6851, 1230, 1117, 1720}

$$\begin{aligned} & \int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx \\ &= \frac{\cos^2(x)\sqrt{\tan^4(x)+3\tan^2(x)+3} \arctan\left(\frac{\tan(x)}{\sqrt{\tan^4(x)+3\tan^2(x)+3}}\right)}{2\sqrt{\cos^4(x)(\tan^4(x)+3\tan^2(x)+3)}} \\ & \quad - \frac{(1+\sqrt{3})\cos^2(x)(\tan^2(x)+\sqrt{3})\sqrt{\frac{\tan^4(x)+3\tan^2(x)+3}{(\tan^2(x)+\sqrt{3})^2}} \operatorname{EllipticF}\left(2\arctan\left(\frac{\tan(x)}{\sqrt[4]{3}}\right), \frac{1}{4}(2-\sqrt{3})\right)}{4\sqrt[4]{3}\sqrt{\cos^4(x)(\tan^4(x)+3\tan^2(x)+3)}} \\ & \quad + \frac{(2+\sqrt{3})\cos^2(x)(\tan^2(x)+\sqrt{3})\sqrt{\frac{\tan^4(x)+3\tan^2(x)+3}{(\tan^2(x)+\sqrt{3})^2}} \operatorname{EllipticPi}\left(\frac{1}{6}(3-2\sqrt{3}), 2\arctan\left(\frac{\tan(x)}{\sqrt[4]{3}}\right), \frac{1}{4}(2-\sqrt{3})\right)}{4\sqrt[4]{3}\sqrt{\cos^4(x)(\tan^4(x)+3\tan^2(x)+3)}} \end{aligned}$$

[In] Int[Cos[x]^2/Sqrt[1 + Cos[x]^2 + Cos[x]^4], x]

[Out] (ArcTan[Tan[x]/Sqrt[3 + 3*Tan[x]^2 + Tan[x]^4]]*Cos[x]^2*Sqrt[3 + 3*Tan[x]^2 + Tan[x]^4])/(2*Sqrt[Cos[x]^4*(3 + 3*Tan[x]^2 + Tan[x]^4)]) - ((1 + Sqrt[3])*Cos[x]^2*EllipticF[2*ArcTan[Tan[x]/3^(1/4)], (2 - Sqrt[3])/4]*(Sqrt[3] + Tan[x]^2)*Sqrt[(3 + 3*Tan[x]^2 + Tan[x]^4)/(Sqrt[3] + Tan[x]^2)^2])/(4*3^(1/4)*Sqrt[Cos[x]^4*(3 + 3*Tan[x]^2 + Tan[x]^4)]) + ((2 + Sqrt[3])*Cos[x]^2*EllipticPi[(3 - 2*Sqrt[3])/6, 2*ArcTan[Tan[x]/3^(1/4)], (2 - Sqrt[3])/4]*(Sqrt[3] + Tan[x]^2)*Sqrt[(3 + 3*Tan[x]^2 + Tan[x]^4)/(Sqrt[3] + Tan[x]^2)^2])/(4*3^(1/4)*Sqrt[Cos[x]^4*(3 + 3*Tan[x]^2 + Tan[x]^4)])

Rule 1117

Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^2 + c*x^4])]*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1230

Int[1/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]

Rule 1720

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2)])/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4])*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/(4*a*B))], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rule 6851

Int[(u_)*((a_)*(v_)^(m_)*(w_)^(n_))^(p_), x_Symbol] := Dist[a^IntPart[p]*((a*v^m*w^n)^FracPart[p]/(v^(m*FracPart[p])*w^(n*FracPart[p]))), Int[u*v^(m*p)*w^(n*p), x], x] /; FreeQ[{a, m, n, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !FreeQ[w, x]

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{1}{(1+x^2)^2 \sqrt{\frac{3+3x^2+x^4}{(1+x^2)^2}}} dx, x, \tan(x) \right) \\
&= \frac{\left(\cos^2(x) \sqrt{3+3\tan^2(x)+\tan^4(x)} \right) \text{Subst} \left(\int \frac{1}{(1+x^2)\sqrt{3+3x^2+x^4}} dx, x, \tan(x) \right)}{\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}} \\
&= \frac{\left((-1-\sqrt{3}) \cos^2(x) \sqrt{3+3\tan^2(x)+\tan^4(x)} \right) \text{Subst} \left(\int \frac{1}{\sqrt{3+3x^2+x^4}} dx, x, \tan(x) \right)}{2\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}} \\
&\quad + \frac{\left((3+\sqrt{3}) \cos^2(x) \sqrt{3+3\tan^2(x)+\tan^4(x)} \right) \text{Subst} \left(\int \frac{1+\frac{x^2}{\sqrt{3}}}{(1+x^2)\sqrt{3+3x^2+x^4}} dx, x, \tan(x) \right)}{2\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}} \\
&= \frac{\arctan \left(\frac{\tan(x)}{\sqrt{3+3\tan^2(x)+\tan^4(x)}} \right) \cos^2(x) \sqrt{3+3\tan^2(x)+\tan^4(x)}}{2\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}} \\
&\quad - \frac{(1+\sqrt{3}) \cos^2(x) \text{EllipticF} \left(2 \arctan \left(\frac{\tan(x)}{\sqrt{3}} \right), \frac{1}{4}(2-\sqrt{3}) \right) (\sqrt{3}+\tan^2(x)) \sqrt{\frac{3+3\tan^2(x)+\tan^4(x)}{(\sqrt{3}+\tan^2(x))^2}}}{4\sqrt{3}\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}} \\
&\quad + \frac{(2+\sqrt{3}) \cos^2(x) \text{EllipticPi} \left(\frac{1}{6}(3-2\sqrt{3}), 2 \arctan \left(\frac{\tan(x)}{\sqrt{3}} \right), \frac{1}{4}(2-\sqrt{3}) \right) (\sqrt{3}+\tan^2(x)) \sqrt{\frac{3+3\tan^2(x)+\tan^4(x)}{(\sqrt{3}+\tan^2(x))^2}}}{4\sqrt{3}\sqrt{\cos^4(x) (3+3\tan^2(x)+\tan^4(x))}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3 in optimal.

Time = 0.81 (sec) , antiderivative size = 159, normalized size of antiderivative = 3.53

$$\int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx = \frac{2i \cos^2(x) \text{EllipticPi} \left(\frac{3}{2} + \frac{i\sqrt{3}}{2}, \text{I} \text{ArcSinh} \left(\sqrt{-\frac{2i}{-3i+\sqrt{3}}} \tan(x) \right), \frac{3i-\sqrt{3}}{3i+\sqrt{3}} \right) \sqrt{1-\frac{2i \tan^2(x)}{-3i+\sqrt{3}}} \sqrt{1+\frac{2i \tan^2(x)}{3i+\sqrt{3}}}}{\sqrt{-\frac{i}{-3i+\sqrt{3}}} \sqrt{15+8\cos(2x)+\cos(4x)}}$$

[In] Integrate[Cos[x]^2/Sqrt[1 + Cos[x]^2 + Cos[x]^4], x]

[Out] ((-2*I)*Cos[x]^2*EllipticPi[3/2 + (I/2)*Sqrt[3], I*ArcSinh[Sqrt[(-2*I)/(-3*I + Sqrt[3])]*Tan[x]], (3*I - Sqrt[3])/(3*I + Sqrt[3])]*Sqrt[1 - ((2*I)*Tan[x]^2)/(-3*I + Sqrt[3])]*Sqrt[1 + ((2*I)*Tan[x]^2)/(3*I + Sqrt[3])])/(Sqrt[(-I)/(-3*I + Sqrt[3])]*Sqrt[15 + 8*Cos[2*x] + Cos[4*x]])

Maple [C] (verified)

Result contains higher order function than in optimal. Order 4 vs. order 3.

Time = 2.60 (sec) , antiderivative size = 312, normalized size of antiderivative = 6.93

method	result
default	$\frac{2\sqrt{(7+4\cos(2x)+\cos^2(2x))(\sin^2(2x))} (i\sqrt{3}-3) \sqrt{\frac{(-1+i\sqrt{3})(\cos(2x)-1)}{(i\sqrt{3}-3)(1+\cos(2x))}} (1+\cos(2x))^2 \sqrt{\frac{\cos(2x)+2+i\sqrt{3}}{(i\sqrt{3}+3)(1+\cos(2x))}} \sqrt{\frac{i\sqrt{3}-\cos(2x)-2}{(i\sqrt{3}-3)(1+\cos(2x))}}}{(-1+i\sqrt{3}) \sqrt{(\cos(2x)-1)(1+\cos(2x))} (\cos(2x)+2+i\sqrt{3}) (i\sqrt{3}-\cos(2x)-2) \sin(2x) \sqrt{7+4\cos(2x)+\cos^2(2x)}}$

[In] `int(cos(x)^2/(1+cos(x)^2+cos(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-2*((7+4*\cos(2*x)+\cos(2*x)^2)*\sin(2*x)^2)^(1/2)*(I*3^(1/2)-3)*((-1+I*3^(1/2))*(\cos(2*x)-1)/(I*3^(1/2)-3)/(1+\cos(2*x)))^(1/2)*(1+\cos(2*x))^2*((\cos(2*x)+2+I*3^(1/2))/(I*3^(1/2)+3)/(1+\cos(2*x)))^(1/2)*((I*3^(1/2)-\cos(2*x)-2)/(I*3^(1/2)-3)/(1+\cos(2*x)))^(1/2)*\text{EllipticPi}(((-1+I*3^(1/2))*(\cos(2*x)-1)/(I*3^(1/2)-3)/(1+\cos(2*x)))^(1/2), (I*3^(1/2)-3)/(-1+I*3^(1/2)), ((1+I*3^(1/2))*(I*3^(1/2)-3)/(I*3^(1/2)+3)/(-1+I*3^(1/2)))^(1/2))/(-1+I*3^(1/2))/((\cos(2*x)-1)*(1+\cos(2*x))*(\cos(2*x)+2+I*3^(1/2))*(I*3^(1/2)-\cos(2*x)-2))^(1/2)/\sin(2*x)/(7+4*\cos(2*x)+\cos(2*x)^2)^(1/2)$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx = \frac{1}{6} \arctan \left(\frac{2\sqrt{\cos^4(x)+\cos^2(x)+1}\cos^3(x)\sin(x)}{2\cos^6(x)-1} \right)$$

[In] `integrate(cos(x)^2/(1+cos(x)^2+cos(x)^4)^(1/2),x, algorithm="fricas")`

[Out] `1/6*arctan(2*sqrt(cos(x)^4 + cos(x)^2 + 1)*cos(x)^3*sin(x)/(2*cos(x)^6 - 1))`

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(x)}{\sqrt{1+\cos^2(x)+\cos^4(x)}} dx = \text{Timed out}$$

[In] `integrate(cos(x)**2/(1+cos(x)**2+cos(x)**4)**(1/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^2(x)}{\sqrt{1 + \cos^2(x) + \cos^4(x)}} dx = \int \frac{\cos(x)^2}{\sqrt{\cos(x)^4 + \cos(x)^2 + 1}} dx$$

[In] integrate(cos(x)^2/(1+cos(x)^2+cos(x)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(cos(x)^2/sqrt(cos(x)^4 + cos(x)^2 + 1), x)

Giac [F]

$$\int \frac{\cos^2(x)}{\sqrt{1 + \cos^2(x) + \cos^4(x)}} dx = \int \frac{\cos(x)^2}{\sqrt{\cos(x)^4 + \cos(x)^2 + 1}} dx$$

[In] integrate(cos(x)^2/(1+cos(x)^2+cos(x)^4)^(1/2),x, algorithm="giac")

[Out] integrate(cos(x)^2/sqrt(cos(x)^4 + cos(x)^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(x)}{\sqrt{1 + \cos^2(x) + \cos^4(x)}} dx = \int \frac{\cos(x)^2}{\sqrt{\cos(x)^4 + \cos(x)^2 + 1}} dx$$

[In] int(cos(x)^2/(cos(x)^2 + cos(x)^4 + 1)^(1/2),x)

[Out] int(cos(x)^2/(cos(x)^2 + cos(x)^4 + 1)^(1/2), x)

3.6 $\int \tan(x) \sqrt{1 + \tan^4(x)} dx$

Optimal result	72
Rubi [A] (verified)	72
Mathematica [A] (verified)	74
Maple [A] (verified)	74
Fricas [B] (verification not implemented)	75
Sympy [F]	75
Maxima [F]	75
Giac [A] (verification not implemented)	76
Mupad [F(-1)]	76

Optimal result

Integrand size = 13, antiderivative size = 56

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = -\frac{1}{2} \operatorname{arcsinh}(\tan^2(x)) - \frac{\operatorname{arctanh}\left(\frac{1 - \tan^2(x)}{\sqrt{2}\sqrt{1 + \tan^4(x)}}\right)}{\sqrt{2}} + \frac{1}{2} \sqrt{1 + \tan^4(x)}$$

[Out] $-1/2*\operatorname{arcsinh}(\tan(x)^2) - 1/2*\operatorname{arctanh}(1/2*(1 - \tan(x)^2)*2^{(1/2)}/(1 + \tan(x)^4)^{(1/2)})*2^{(1/2)} + 1/2*(1 + \tan(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3751, 1262, 749, 858, 221, 739, 212}

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = -\frac{1}{2} \operatorname{arcsinh}(\tan^2(x)) - \frac{\operatorname{arctanh}\left(\frac{1 - \tan^2(x)}{\sqrt{2}\sqrt{\tan^4(x) + 1}}\right)}{\sqrt{2}} + \frac{1}{2} \sqrt{\tan^4(x) + 1}$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]*\operatorname{Sqrt}[1 + \operatorname{Tan}[x]^4], x]$

[Out] $-1/2*\operatorname{ArcSinh}[\operatorname{Tan}[x]^2] - \operatorname{ArcTanh}[(1 - \operatorname{Tan}[x]^2)/(\operatorname{Sqrt}[2]*\operatorname{Sqrt}[1 + \operatorname{Tan}[x]^4])]/\operatorname{Sqrt}[2] + \operatorname{Sqrt}[1 + \operatorname{Tan}[x]^4]/2$

Rule 212

$\operatorname{Int}[(a + (b \cdot x^2)^{-1}), x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 221


```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 739

```
Int[1/(((d_) + (e_.)*(x_))*Sqrt[(a_) + (c_.)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ
[{a, c, d, e}, x]
```

Rule 749

```
Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
(d + e*x)^(m + 1)*((a + c*x^2)^p/(e*(m + 2*p + 1))), x] + Dist[2*(p/(e*(m +
2*p + 1))), Int[(d + e*x)^m*Simp[a*e - c*d*x, x]*(a + c*x^2)^(p - 1), x],
x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && GtQ[p, 0] && Ne
Q[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] &&
IntQuadraticQ[a, 0, c, d, e, m, p, x]
```

Rule 858

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] + D
ist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d,
e, f, g, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1262

```
Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol]
:= Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
[{a, c, d, e, p, q}, x]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\text{integral} = \text{Subst} \left(\int \frac{x\sqrt{1+x^4}}{1+x^2} dx, x, \tan(x) \right)$$

$$\begin{aligned}
&= \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1+x^2}}{1+x} dx, x, \tan^2(x) \right) \\
&= \frac{1}{2} \sqrt{1+\tan^4(x)} + \frac{1}{2} \text{Subst} \left(\int \frac{1-x}{(1+x)\sqrt{1+x^2}} dx, x, \tan^2(x) \right) \\
&= \frac{1}{2} \sqrt{1+\tan^4(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \tan^2(x) \right) \\
&\quad + \text{Subst} \left(\int \frac{1}{(1+x)\sqrt{1+x^2}} dx, x, \tan^2(x) \right) \\
&= -\frac{1}{2} \operatorname{arcsinh}(\tan^2(x)) + \frac{1}{2} \sqrt{1+\tan^4(x)} - \text{Subst} \left(\int \frac{1}{2-x^2} dx, x, \frac{1-\tan^2(x)}{\sqrt{1+\tan^4(x)}} \right) \\
&= -\frac{1}{2} \operatorname{arcsinh}(\tan^2(x)) - \frac{\operatorname{arctanh}\left(\frac{1-\tan^2(x)}{\sqrt{2}\sqrt{1+\tan^4(x)}}\right)}{\sqrt{2}} + \frac{1}{2} \sqrt{1+\tan^4(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.32

$$\begin{aligned}
&\int \tan(x) \sqrt{1+\tan^4(x)} dx \\
&= \frac{\left(-2\sqrt{2}\operatorname{arcsinh}(\cos(2x))\cos^2(x) - 2\operatorname{arctanh}\left(\frac{2\sin^2(x)}{\sqrt{3+\cos(4x)}}\right)\cos^2(x) + \sqrt{3+\cos(4x)}\right)\sqrt{1+\tan^4(x)}}{2\sqrt{3+\cos(4x)}}
\end{aligned}$$

[In] Integrate[Tan[x]*Sqrt[1 + Tan[x]^4],x]

[Out] ((-2*Sqrt[2]*ArcSinh[Cos[2*x]]*Cos[x]^2 - 2*ArcTanh[(2*Sin[x]^2)/Sqrt[3 + Cos[4*x]])*Cos[x]^2 + Sqrt[3 + Cos[4*x]])*Sqrt[1 + Tan[x]^4])/(2*Sqrt[3 + Cos[4*x]])

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$\frac{\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}{2} - \frac{\operatorname{arcsinh}(\tan^2(x))}{2} - \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(-2(\tan^2(x))+2)\sqrt{2}}{4\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}\right)}{2}$	64
default	$\frac{\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}{2} - \frac{\operatorname{arcsinh}(\tan^2(x))}{2} - \frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(-2(\tan^2(x))+2)\sqrt{2}}{4\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}\right)}{2}$	64

[In] `int((1+tan(x)^4)^(1/2)*tan(x),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2} * ((1 + \tan(x)^2)^2 - 2 * \tan(x)^2)^{(1/2)} - \frac{1}{2} * \operatorname{arcsinh}(\tan(x)^2) - \frac{1}{2} * 2^{(1/2)} * \operatorname{arc} \tanh(1/4 * (-2 * \tan(x)^2 + 2) * 2^{(1/2)}) / ((1 + \tan(x)^2)^2 - 2 * \tan(x)^2)^{(1/2)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 88 vs. $2(43) = 86$.

Time = 0.31 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.57

$$\begin{aligned} & \int \tan(x) \sqrt{1 + \tan^4(x)} dx \\ &= \frac{1}{4} \sqrt{2} \log \left(\frac{3 \tan(x)^4 - 2 \tan(x)^2 + 2 \sqrt{\tan(x)^4 + 1} (\sqrt{2} \tan(x)^2 - \sqrt{2}) + 3}{\tan(x)^4 + 2 \tan(x)^2 + 1} \right) \\ &+ \frac{1}{2} \sqrt{\tan(x)^4 + 1} + \frac{1}{2} \log \left(-\tan(x)^2 + \sqrt{\tan(x)^4 + 1} \right) \end{aligned}$$

[In] `integrate((1+tan(x)^4)^(1/2)*tan(x),x, algorithm="fricas")`

[Out] $\frac{1}{4} * \sqrt{2} * \log((3 * \tan(x)^4 - 2 * \tan(x)^2 + 2 * \sqrt{\tan(x)^4 + 1}) * (\sqrt{2} * \tan(x)^2 - \sqrt{2}) + 3) / (\tan(x)^4 + 2 * \tan(x)^2 + 1) + \frac{1}{2} * \sqrt{\tan(x)^4 + 1} + \frac{1}{2} * \log(-\tan(x)^2 + \sqrt{\tan(x)^4 + 1})$

Sympy [F]

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = \int \sqrt{\tan^4(x) + 1} \tan(x) dx$$

[In] `integrate((1+tan(x)**4)**(1/2)*tan(x),x)`

[Out] `Integral(sqrt(tan(x)**4 + 1)*tan(x), x)`

Maxima [F]

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = \int \sqrt{\tan(x)^4 + 1} \tan(x) dx$$

[In] `integrate((1+tan(x)^4)^(1/2)*tan(x),x, algorithm="maxima")`

[Out] `integrate(sqrt(tan(x)^4 + 1)*tan(x), x)`

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.41

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = \frac{1}{2} \sqrt{2} \log \left(-\frac{\tan(x)^2 + \sqrt{2} - \sqrt{\tan(x)^4 + 1 + 1}}{\tan(x)^2 - \sqrt{2} - \sqrt{\tan(x)^4 + 1 + 1}} \right) + \frac{1}{2} \sqrt{\tan(x)^4 + 1} + \frac{1}{2} \log \left(-\tan(x)^2 + \sqrt{\tan(x)^4 + 1} \right)$$

```
[In] integrate((1+tan(x)^4)^(1/2)*tan(x),x, algorithm="giac")
```

```
[Out] 1/2*sqrt(2)*log(-(tan(x)^2 + sqrt(2) - sqrt(tan(x)^4 + 1) + 1)/(tan(x)^2 - sqrt(2) - sqrt(tan(x)^4 + 1) + 1)) + 1/2*sqrt(tan(x)^4 + 1) + 1/2*log(-tan(x)^2 + sqrt(tan(x)^4 + 1))
```

Mupad [F(-1)]

Timed out.

$$\int \tan(x) \sqrt{1 + \tan^4(x)} dx = \int \tan(x) \sqrt{\tan(x)^4 + 1} dx$$

```
[In] int(tan(x)*(tan(x)^4 + 1)^(1/2),x)
```

```
[Out] int(tan(x)*(tan(x)^4 + 1)^(1/2), x)
```

3.7 $\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx$

Optimal result	77
Rubi [A] (verified)	77
Mathematica [A] (verified)	78
Maple [A] (verified)	79
Fricas [B] (verification not implemented)	79
Sympy [F]	79
Maxima [B] (verification not implemented)	80
Giac [B] (verification not implemented)	80
Mupad [F(-1)]	80

Optimal result

Integrand size = 13, antiderivative size = 15

$$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx = -\frac{2}{3} \operatorname{arctanh}\left(\sqrt{1+\sec^3(x)}\right)$$

[Out] $-2/3*\operatorname{arctanh}((1+\sec(x)^3)^{(1/2)})$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {4224, 272, 65, 213}

$$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx = -\frac{2}{3} \operatorname{arctanh}\left(\sqrt{\sec^3(x)+1}\right)$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]/\operatorname{Sqrt}[1+\operatorname{Sec}[x]^3],x]$

[Out] $(-2*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+\operatorname{Sec}[x]^3]])/3$

Rule 65

$\operatorname{Int}[(a_. + (b_.)(x_))^{(m_)}*((c_.) + (d_.)(x_))^{(n_)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{n_}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 213

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4224

```
Int[((a_) + (b_)*((c_)*sec[(e_) + (f_)*(x_)])^(n_))^(p_)*tan[(e_) + (f_)*(x_)]^(m_), x_Symbol] := With[{ff = FreeFactors[Sec[e + f*x], x]}, Dist[1/f, Subst[Int[(-1 + ff^2*x^2)^(m - 1)/2]*((a + b*(c*ff*x)^n)^p/x), x], x, Sec[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2] && (GtQ[m, 0] || EqQ[n, 2] || EqQ[n, 4] || IGtQ[p, 0] || IntegersQ[2*n, p])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{1}{x\sqrt{1+x^3}} dx, x, \sec(x) \right) \\
 &= \frac{1}{3} \text{Subst} \left(\int \frac{1}{x\sqrt{1+x}} dx, x, \sec^3(x) \right) \\
 &= \frac{2}{3} \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+\sec^3(x)} \right) \\
 &= -\frac{2}{3} \operatorname{arctanh} \left(\sqrt{1+\sec^3(x)} \right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx = -\frac{2}{3} \operatorname{arctanh} \left(\sqrt{1+\sec^3(x)} \right)$$

```
[In] Integrate[Tan[x]/Sqrt[1 + Sec[x]^3], x]
```

```
[Out] (-2*ArcTanh[Sqrt[1 + Sec[x]^3]])/3
```

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.80

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}\left(\sqrt{1+\sec^3(x)}\right)}{3}$	12
default	$-\frac{2 \operatorname{arctanh}\left(\sqrt{1+\sec^3(x)}\right)}{3}$	12

[In] `int(tan(x)/(1+sec(x)^3)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-2/3*arctanh((1+sec(x)^3)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(11) = 22.

Time = 0.31 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.00

$$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx = \frac{1}{3} \log \left(2 \sqrt{\frac{\cos(x)^3 + 1}{\cos(x)^3}} \cos(x)^3 - 2 \cos(x)^3 - 1 \right)$$

[In] `integrate(tan(x)/(1+sec(x)^3)^(1/2),x, algorithm="fricas")`

[Out] `1/3*log(2*sqrt((cos(x)^3 + 1)/cos(x)^3)*cos(x)^3 - 2*cos(x)^3 - 1)`

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{1+\sec^3(x)}} dx = \int \frac{\tan(x)}{\sqrt{(\sec(x)+1)(\sec^2(x)-\sec(x)+1)}} dx$$

[In] `integrate(tan(x)/(1+sec(x)**3)**(1/2),x)`

[Out] `Integral(tan(x)/sqrt((sec(x) + 1)*(sec(x)**2 - sec(x) + 1)), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. $2(11) = 22$.

Time = 0.20 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \frac{\tan(x)}{\sqrt{1 + \sec^3(x)}} dx = -\frac{1}{3} \log \left(\sqrt{\frac{1}{\cos(x)^3} + 1} + 1 \right) + \frac{1}{3} \log \left(\sqrt{\frac{1}{\cos(x)^3} + 1} - 1 \right)$$

[In] integrate(tan(x)/(1+sec(x)^3)^(1/2),x, algorithm="maxima")

[Out] -1/3*log(sqrt(1/cos(x)^3 + 1) + 1) + 1/3*log(sqrt(1/cos(x)^3 + 1) - 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. $2(11) = 22$.

Time = 0.30 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.87

$$\int \frac{\tan(x)}{\sqrt{1 + \sec^3(x)}} dx = -\frac{1}{3} \log \left(\sqrt{\frac{1}{\cos(x)^3} + 1} + 1 \right) + \frac{1}{3} \log \left(\left| \sqrt{\frac{1}{\cos(x)^3} + 1} - 1 \right| \right)$$

[In] integrate(tan(x)/(1+sec(x)^3)^(1/2),x, algorithm="giac")

[Out] -1/3*log(sqrt(1/cos(x)^3 + 1) + 1) + 1/3*log(abs(sqrt(1/cos(x)^3 + 1) - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{1 + \sec^3(x)}} dx = \int \frac{\tan(x)}{\sqrt{\frac{1}{\cos(x)^3} + 1}} dx$$

[In] int(tan(x)/(1/cos(x)^3 + 1)^(1/2),x)

[Out] int(tan(x)/(1/cos(x)^3 + 1)^(1/2), x)

3.8 $\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx$

Optimal result	81
Rubi [A] (verified)	82
Mathematica [C] (verified)	84
Maple [B] (verified)	84
Fricas [C] (verification not implemented)	86
Sympy [F]	87
Maxima [C] (verification not implemented)	87
Giac [B] (verification not implemented)	90
Mupad [F(-1)]	91

Optimal result

Integrand size = 14, antiderivative size = 137

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx$$

$$= \operatorname{arcsinh}(1 + \tan(x)) - \sqrt{\frac{1}{2} (1 + \sqrt{5})} \arctan \left(\frac{2\sqrt{5} - (5 + \sqrt{5}) \tan(x)}{\sqrt{10} (1 + \sqrt{5}) \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right)$$

$$- \sqrt{\frac{1}{2} (-1 + \sqrt{5})} \operatorname{arctanh} \left(\frac{2\sqrt{5} + (5 - \sqrt{5}) \tan(x)}{\sqrt{10} (-1 + \sqrt{5}) \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right)$$

```
[Out] arcsinh(1+tan(x))-1/2*arctanh((2*5^(1/2)+(5-5^(1/2))*tan(x))/(-10+10*5^(1/2)
))^(1/2)/(2+2*tan(x)+tan(x)^2)^(1/2))*(-2+2*5^(1/2))^(1/2)-1/2*arctan((2*5^(
1/2)-(5+5^(1/2))*tan(x))/(10+10*5^(1/2))^(1/2)/(2+2*tan(x)+tan(x)^2)^(1/2)
)*(2+2*5^(1/2))^(1/2)
```

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {1004, 633, 221, 1050, 1044, 213, 209}

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx$$

$$= \operatorname{arcsinh}(\tan(x) + 1) - \sqrt{\frac{1}{2}(1 + \sqrt{5})} \operatorname{arctan}\left(\frac{2\sqrt{5} - (5 + \sqrt{5}) \tan(x)}{\sqrt{10(1 + \sqrt{5})} \sqrt{\tan^2(x) + 2 \tan(x) + 2}}\right)$$

$$- \sqrt{\frac{1}{2}(\sqrt{5} - 1)} \operatorname{arctanh}\left(\frac{(5 - \sqrt{5}) \tan(x) + 2\sqrt{5}}{\sqrt{10(\sqrt{5} - 1)} \sqrt{\tan^2(x) + 2 \tan(x) + 2}}\right)$$

[In] Int[Sqrt[2 + 2*Tan[x] + Tan[x]^2],x]

[Out] ArcSinh[1 + Tan[x]] - Sqrt[(1 + Sqrt[5])/2]*ArcTan[(2*Sqrt[5] - (5 + Sqrt[5])*Tan[x])/(Sqrt[10*(1 + Sqrt[5])]*Sqrt[2 + 2*Tan[x] + Tan[x]^2])] - Sqrt[(-1 + Sqrt[5])/2]*ArcTanh[(2*Sqrt[5] + (5 - Sqrt[5])*Tan[x])/(Sqrt[10*(-1 + Sqrt[5])]*Sqrt[2 + 2*Tan[x] + Tan[x]^2])]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 1004

```
Int[Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]/((d_) + (f_)*(x_)^2), x_Symbol]
  :> Dist[c/f, Int[1/Sqrt[a + b*x + c*x^2], x], x] - Dist[1/f, Int[(c*d - a*
f - b*f*x)/(Sqrt[a + b*x + c*x^2]*(d + f*x^2)), x], x] /; FreeQ[{a, b, c, d
, f}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 1044

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f
_)*(x_)^2]), x_Symbol] :> Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*
e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[
{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]
```

Rule 1050

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (
f_)*(x_)^2]), x_Symbol] :> With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist
[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*
e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Si
mp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c
*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] &&
NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{\sqrt{2+2x+x^2}}{1+x^2} dx, x, \tan(x) \right) \\
&= \text{Subst} \left(\int \frac{1}{\sqrt{2+2x+x^2}} dx, x, \tan(x) \right) - \text{Subst} \left(\int \frac{-1-2x}{(1+x^2)\sqrt{2+2x+x^2}} dx, x, \tan(x) \right) \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{4}}} dx, x, 2+2\tan(x) \right) \\
&\quad - \frac{\text{Subst} \left(\int \frac{5-\sqrt{5}-2\sqrt{5}x}{(1+x^2)\sqrt{2+2x+x^2}} dx, x, \tan(x) \right)}{2\sqrt{5}} + \frac{\text{Subst} \left(\int \frac{5+\sqrt{5}+2\sqrt{5}x}{(1+x^2)\sqrt{2+2x+x^2}} dx, x, \tan(x) \right)}{2\sqrt{5}} \\
&= \text{arcsinh}(1 + \tan(x)) \\
&\quad - \left(2(5 - \sqrt{5}) \right) \text{Subst} \left(\int \frac{1}{20(1 - \sqrt{5}) + 2x^2} dx, x, \frac{-2\sqrt{5} - (5 - \sqrt{5})\tan(x)}{\sqrt{2+2\tan(x) + \tan^2(x)}} \right) \\
&\quad - \left(2(5 + \sqrt{5}) \right) \text{Subst} \left(\int \frac{1}{20(1 + \sqrt{5}) + 2x^2} dx, x, \frac{2\sqrt{5} - (5 + \sqrt{5})\tan(x)}{\sqrt{2+2\tan(x) + \tan^2(x)}} \right)
\end{aligned}$$

$$\begin{aligned}
&= \operatorname{arcsinh}(1 + \tan(x)) - \sqrt{\frac{1}{2}(1 + \sqrt{5})} \operatorname{arctan} \left(\frac{2\sqrt{5} - (5 + \sqrt{5}) \tan(x)}{\sqrt{10(1 + \sqrt{5})} \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right) \\
&\quad - \sqrt{\frac{1}{2}(-1 + \sqrt{5})} \operatorname{arctanh} \left(\frac{2\sqrt{5} + (5 - \sqrt{5}) \tan(x)}{\sqrt{10(-1 + \sqrt{5})} \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right)
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.20 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.72

$$\begin{aligned}
&\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx \\
&= \operatorname{arcsinh}(1 + \tan(x)) + \frac{1}{2}i \left(\sqrt{1 + 2i} \operatorname{arctanh} \left(\frac{(2 + i) + (1 + i) \tan(x)}{\sqrt{1 + 2i} \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right) \right. \\
&\quad \left. - \sqrt{1 - 2i} \operatorname{arctanh} \left(\frac{(4 - 2i) + (2 - 2i) \tan(x)}{2\sqrt{1 - 2i} \sqrt{2 + 2 \tan(x) + \tan^2(x)}} \right) \right)
\end{aligned}$$

[In] Integrate[Sqrt[2 + 2*Tan[x] + Tan[x]^2], x]

[Out] ArcSinh[1 + Tan[x]] + (I/2)*(Sqrt[1 + 2*I]*ArcTanh[((2 + I) + (1 + I)*Tan[x])/(Sqrt[1 + 2*I]*Sqrt[2 + 2*Tan[x] + Tan[x]^2])]) - Sqrt[1 - 2*I]*ArcTanh[((4 - 2*I) + (2 - 2*I)*Tan[x])/(2*Sqrt[1 - 2*I]*Sqrt[2 + 2*Tan[x] + Tan[x]^2])])]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1603 vs. 2(105) = 210.

Time = 1.60 (sec) , antiderivative size = 1604, normalized size of antiderivative = 11.71

method	result	size
derivativedivides	Expression too large to display	1604
default	Expression too large to display	1604

[In] int((2+2*tan(x)+tan(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] arcsinh(tan(x)+1)-1/10*(10*(-1/2*5^(1/2)+1/2+tan(x))^2/(-1/2*5^(1/2)-1/2-tan(x))^2-2*5^(1/2)*(-1/2*5^(1/2)+1/2+tan(x))^2/(-1/2*5^(1/2)-1/2-tan(x))^2+10*2*5^(1/2))^(1/2)*5^(1/2)*(3*5^(1/2)*(-10+10*5^(1/2))^(1/2)*arctan(1/80*(-22+10*5^(1/2))^(1/2)*((5-5^(1/2))*(2*(-1/2*5^(1/2)+1/2+tan(x))^2/(-1/2*5^(1/2)-1/2-tan(x))^2+5^(1/2)+3))^(1/2)*(11*5^(1/2)*(-1/2*5^(1/2)+1/2+tan(x))^2

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.65

$$\begin{aligned}
 & \int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx = \\
 & -\frac{1}{4} \sqrt{2i-1} \log \left(\frac{\sqrt{2i-1}((9i+13) \tan(x)^2 + (7i+24) \tan(x) - 15i+20) + \sqrt{\tan(x)^2 + 2 \tan(x) + 2}}{\tan(x)^2 + 1} \right) \\
 & -\frac{1}{4} \sqrt{-2i-1} \log \left(\frac{\sqrt{-2i-1}(-(9i-13) \tan(x)^2 - (7i-24) \tan(x) + 15i+20) + \sqrt{\tan(x)^2 + 2 \tan(x) + 2}}{\tan(x)^2 + 1} \right) \\
 & +\frac{1}{4} \sqrt{-2i-1} \log \left(\frac{\sqrt{-2i-1}((9i-13) \tan(x)^2 + (7i-24) \tan(x) - 15i-20) + \sqrt{\tan(x)^2 + 2 \tan(x) + 2}}{\tan(x)^2 + 1} \right) \\
 & +\frac{1}{4} \sqrt{2i-1} \log \left(\frac{\sqrt{2i-1}(-(9i+13) \tan(x)^2 - (7i+24) \tan(x) + 15i-20) + \sqrt{\tan(x)^2 + 2 \tan(x) + 2}}{\tan(x)^2 + 1} \right) \\
 & + \log \left(-\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) - 1 \right)
 \end{aligned}$$

[In] integrate((2+2*tan(x)+tan(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/4*sqrt(2*I - 1)*log((sqrt(2*I - 1)*((9*I + 13)*tan(x)^2 + (7*I + 24)*tan(x) - 15*I + 20) + sqrt(tan(x)^2 + 2*tan(x) + 2))*((24*I - 7)*tan(x) + 7*I + 24))/(tan(x)^2 + 1) - 1/4*sqrt(-2*I - 1)*log((sqrt(-2*I - 1)*(-(9*I - 13)*tan(x)^2 - (7*I - 24)*tan(x) + 15*I + 20) + sqrt(tan(x)^2 + 2*tan(x) + 2))*(-(24*I + 7)*tan(x) - 7*I + 24))/(tan(x)^2 + 1) + 1/4*sqrt(-2*I - 1)*log((sqrt(-2*I - 1)*((9*I - 13)*tan(x)^2 + (7*I - 24)*tan(x) - 15*I - 20) + sqrt(tan(x)^2 + 2*tan(x) + 2))*(-(24*I + 7)*tan(x) - 7*I + 24))/(tan(x)^2 + 1) + 1/4*sqrt(2*I - 1)*log((sqrt(2*I - 1)*(-(9*I + 13)*tan(x)^2 - (7*I + 24)*tan(x) + 15*I - 20) + sqrt(tan(x)^2 + 2*tan(x) + 2))*((24*I - 7)*tan(x) + 7*I + 24))/(tan(x)^2 + 1) + log(-sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x) - 1)

Sympy [F]

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx = \int \sqrt{\tan^2(x) + 2 \tan(x) + 2} dx$$

[In] integrate((2+2*tan(x)+tan(x)**2)**(1/2),x)

[Out] Integral(sqrt(tan(x)**2 + 2*tan(x) + 2), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.78 (sec) , antiderivative size = 4065, normalized size of antiderivative = 29.67

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx = \text{Too large to display}$$

[In] integrate((2+2*tan(x)+tan(x)^2)^(1/2),x, algorithm="maxima")

[Out]
$$\begin{aligned} & -1/400*\sqrt{10}*(4*\sqrt{10}*(\sqrt{5}*\sqrt{2}*\sqrt{\sqrt{5} + 1} - \sqrt{5}*\sqrt{2}*\sqrt{\sqrt{5} - 1})*\arctan2(-1/2*\sqrt{2}*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5) \\ & ^{(1/4)*\sqrt{\sqrt{5} - 1}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + 1/2*\sqrt{2}*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5) \\ & ^{(1/4)*\sqrt{\sqrt{5} + 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - 2*\cos(2*x) + \sin(2*x), 1/2*\sqrt{2}*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5) \\ & ^{(1/4)*\sqrt{\sqrt{5} - 1}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + 1/2*\sqrt{2}*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5) \\ & ^{(1/4)*\sqrt{\sqrt{5} + 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + \cos(2*x) + 2*\sin(2*x) + 3) + 10*\sqrt{10}*\sqrt{2}*\sqrt{\sqrt{5} + 1}*\arctan2((6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5) \\ & ^{(1/4)*(\sqrt{2}*\sqrt{\sqrt{5} - 1}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + \sqrt{2}*\sqrt{\sqrt{5} + 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) \end{aligned}$$

$$\begin{aligned}
& + 6*\cos(2*x) + 2*\sin(4*x) + 1))) + 4, (6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos \\
& (4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(\\
& 4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5)^{(1/4)}* \\
& (\sqrt{2}*\sqrt{\sqrt{5} + 1}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2 \\
& *x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - \sqrt{2}*\sqrt{\sqrt{5} - \\
& 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\co \\
& s(2*x) + 2*\sin(4*x) + 1))) + 8) + 2*\sqrt{10}*(\sqrt{5}*\sqrt{2}*\sqrt{\sqrt{5} \\
& + 1} + \sqrt{5}*\sqrt{2}*\sqrt{\sqrt{5} - 1})*\log(\sqrt{5}*\sqrt{6*(2*\cos(2*x) - \\
& 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3 \\
& *\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24* \\
& \sin(2*x) + 5)*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4 \\
& *x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))^2 + \sqrt{5}*\sqrt{6*(2*\cos(2*x) - 4*\sin(2 \\
& *x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2 \\
& *x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x \\
&) + 5)*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + \\
& 6*\cos(2*x) + 2*\sin(4*x) + 1))^2 + (6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) \\
& + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) \\
& + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5)^{(1/4)}*((\sqrt{ \\
& t(2)*\sqrt{\sqrt{5} + 1} + 2*\sqrt{2}*\sqrt{\sqrt{5} - 1})*\cos(2*x) + (2*\sqrt{2} \\
& *\sqrt{\sqrt{5} + 1} - \sqrt{2}*\sqrt{\sqrt{5} - 1})*\sin(2*x) + 3*\sqrt{2}*\sqrt{\sqrt{5} \\
& + 1})*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4 \\
& *x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - (6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(\\
& 4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4 \\
& *x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5)^{(1/4)}* \\
& ((2*\sqrt{2}*\sqrt{\sqrt{5} + 1} - \sqrt{2}*\sqrt{\sqrt{5} - 1})*\cos(2*x) - (\sqrt{ \\
& 2}*\sqrt{\sqrt{5} + 1} + 2*\sqrt{2}*\sqrt{\sqrt{5} - 1})*\sin(2*x) - 3*\sqrt{2}*\sqrt{ \\
& \sqrt{5} - 1})*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, c \\
& os(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + 5*\cos(2*x)^2 + 5*\sin(2*x)^2 + 6*c \\
& os(2*x) + 12*\sin(2*x) + 9) + 5*\sqrt{10}*\sqrt{2}*\sqrt{\sqrt{5} - 1}*\log(4*(\sqrt{ \\
& 5}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6 \\
& *\cos(2*x) + 2*\sin(4*x) + 1))^2 + \sqrt{5}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(\\
& 4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))^2)*\sqrt{6*(\\
& 2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6 \\
& *\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\co \\
& s(2*x) + 24*\sin(2*x) + 5) + 8*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5 \\
& *\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5* \\
& \sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5)^{(1/4)}*((2*\sqrt{ \\
& 2}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\co \\
& s(2*x) + 2*\sin(4*x) + 1)) + \sqrt{2}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) \\
& + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)))*\sqrt{\sqrt{5} + \\
& 1} + (\sqrt{2}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(\\
& 4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - 2*\sqrt{2}*\sin(1/2*\arctan2(-2*\cos(4*x \\
&) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)))*\sqrt{ \\
& \sqrt{5} - 1} + 80) - 60*\sqrt{\sqrt{5} + 1}*\arctan2(-(6*(2*\cos(2*x) - 4*s \\
& \sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*si
\end{aligned}$$

$$\begin{aligned}
& n(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(\\
& 2*x) + 5)^{(1/4)}*(\sqrt{2}*\sqrt{\sqrt{5} - 1}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \\
& \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - \sqrt{2} \\
& *\sqrt{\sqrt{5} + 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \\
& \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))) - 4*\cos(2*x) + 2*\sin(2*x), (6*(2 \\
& *\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6* \\
& \cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos \\
& (2*x) + 24*\sin(2*x) + 5)^{(1/4)}*(\sqrt{2}*\sqrt{\sqrt{5} + 1}*\cos(1/2*\arctan2(- \\
& 2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) \\
& + 1)) + \sqrt{2}*\sqrt{\sqrt{5} - 1}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + \\
& 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))) + 2*\cos(2*x) + 4* \\
& \sin(2*x) + 6) - 30*\sqrt{\sqrt{5} - 1}*\log(20*\cos(2*x)^2 + 20*\sin(2*x)^2 + 4* \\
& (\sqrt{5}*\cos(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) \\
& + 6*\cos(2*x) + 2*\sin(4*x) + 1))^2 + \sqrt{5}*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))^2)*\sqrt{(\\
& 6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4 \\
& *(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12 \\
& *\cos(2*x) + 24*\sin(2*x) + 5) + 4*(6*(2*\cos(2*x) - 4*\sin(2*x) - 1)*\cos(4*x) \\
& + 5*\cos(4*x)^2 + 36*\cos(2*x)^2 + 4*(6*\cos(2*x) + 3*\sin(2*x) + 2)*\sin(4*x) + \\
& 5*\sin(4*x)^2 + 36*\sin(2*x)^2 + 12*\cos(2*x) + 24*\sin(2*x) + 5)^{(1/4)}*((\sqrt{ \\
& 2}*\cos(2*x) + 2*\sqrt{2}*\sin(2*x) + 3*\sqrt{2}))*\cos(1/2*\arctan2(-2*\cos(4*x) \\
& + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) - (2 \\
& *\sqrt{2}*\cos(2*x) - \sqrt{2}*\sin(2*x))*\sin(1/2*\arctan2(-2*\cos(4*x) + \sin(4*x) \\
&) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)))*\sqrt{\sqrt{5} \\
& + 1} + ((2*\sqrt{2}*\cos(2*x) - \sqrt{2}*\sin(2*x))*\cos(1/2*\arctan2(-2*\cos(4*x) \\
& + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1)) + (\sqrt{ \\
& 2}*\cos(2*x) + 2*\sqrt{2}*\sin(2*x) + 3*\sqrt{2}))*\sin(1/2*\arctan2(-2*\cos(4* \\
& x) + \sin(4*x) + 6*\sin(2*x) + 2, \cos(4*x) + 6*\cos(2*x) + 2*\sin(4*x) + 1))) * \\
& \sqrt{\sqrt{5} - 1}) + 24*\cos(2*x) + 48*\sin(2*x) + 36) + 20*\sqrt{10}*\log((\sqrt{ \\
& (\text{abs}(2*e^{(2*I*x)} + 2)^4 + 64*\cos(2*x)^4 + 64*\sin(2*x)^4 + 32*(\cos(2*x)*\sin(\\
& 2*x) + \cos(2*x))*\text{abs}(2*e^{(2*I*x)} + 2)^2 + 128*(\cos(2*x)^2 + 3)*\sin(2*x)^2 + \\
& 256*\sin(2*x)^3 + 128*\cos(2*x)^2 + 256*(\cos(2*x)^2 + 1)*\sin(2*x) + 64)*\cos(\\
& 1/2*\arctan2(-8*(\cos(2*x)^2 - \sin(2*x)^2 - 2*\sin(2*x) - 1)/\text{abs}(2*e^{(2*I*x)} + \\
& 2)^2, (\text{abs}(2*e^{(2*I*x)} + 2)^2 + 16*\cos(2*x)*\sin(2*x) + 16*\cos(2*x))/\text{abs}(2* \\
& e^{(2*I*x)} + 2)^2))^2 + \sqrt{(\text{abs}(2*e^{(2*I*x)} + 2)^4 + 64*\cos(2*x)^4 + 64*\sin \\
& (2*x)^4 + 32*(\cos(2*x)*\sin(2*x) + \cos(2*x))*\text{abs}(2*e^{(2*I*x)} + 2)^2 + 128*(\cos(2*x)^2 + 3)*\sin(2*x)^2 + 256*\sin(2*x)^3 + 128*\cos(2*x)^2 + 256*(\cos(2*x)^2 + 1)*\sin(2*x) + 64)*\sin(1/2*\arctan2(-8*(\cos(2*x)^2 - \sin(2*x)^2 - 2*\sin(2*x) - 1)/\text{abs}(2*e^{(2*I*x)} + 2)^2, (\text{abs}(2*e^{(2*I*x)} + 2)^2 + 16*\cos(2*x)*\sin(2*x) + 16*\cos(2*x))/\text{abs}(2*e^{(2*I*x)} + 2)^2))^2 - 4*(\text{abs}(2*e^{(2*I*x)} + 2)^4 + 64*\cos(2*x)^4 + 64*\sin(2*x)^4 + 32*(\cos(2*x)*\sin(2*x) + \cos(2*x))*\text{abs}(2*e^{(2*I*x)} + 2)^2 + 128*(\cos(2*x)^2 + 3)*\sin(2*x)^2 + 256*\sin(2*x)^3 + 128*\cos(2*x)^2 + 256*(\cos(2*x)^2 + 1)*\sin(2*x) + 64)^{(1/4)}*(\cos(2*x) + \sin(2*x) + 1)*\cos(1/2*\arctan2(-8*(\cos(2*x)^2 - \sin(2*x)^2 - 2*\sin(2*x) - 1)/\text{abs}(2*e^{(2*I*x)} + 2)^2, (\text{abs}(2*e^{(2*I*x)} + 2)^2 + 16*\cos(2*x)*\sin(2*x) + 16*\cos(2*x)
\end{aligned}$$

)/abs(2*e^(2*I*x) + 2)^2)) + 4*(abs(2*e^(2*I*x) + 2)^4 + 64*cos(2*x)^4 + 64*sin(2*x)^4 + 32*(cos(2*x)*sin(2*x) + cos(2*x))*abs(2*e^(2*I*x) + 2)^2 + 128*(cos(2*x)^2 + 3)*sin(2*x)^2 + 256*sin(2*x)^3 + 128*cos(2*x)^2 + 256*(cos(2*x)^2 + 1)*sin(2*x) + 64)^(1/4)*(cos(2*x) - sin(2*x) - 1)*sin(1/2*arctan2(-8*(cos(2*x)^2 - sin(2*x)^2 - 2*sin(2*x) - 1)/abs(2*e^(2*I*x) + 2)^2, (abs(2*e^(2*I*x) + 2)^2 + 16*cos(2*x)*sin(2*x) + 16*cos(2*x))/abs(2*e^(2*I*x) + 2)^2)) + 8*cos(2*x)^2 + 8*sin(2*x)^2 + 16*sin(2*x) + 8)/abs(2*e^(2*I*x) + 2)^2))

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 495 vs. 2(104) = 208.

Time = 0.34 (sec) , antiderivative size = 495, normalized size of antiderivative = 3.61

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx =$$

$$-\frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(256 \left(\sqrt{5} \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) + \sqrt{5} \sqrt{\sqrt{5}-2} - \sqrt{5} - 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} \right) \right.$$

$$+ 256 \left(\sqrt{5} \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) + \sqrt{5} - 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} + \sqrt{\sqrt{5}-2} + 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} \right) \left.$$

$$+ \frac{1}{4} \sqrt{2\sqrt{5}-2} \log \left(256 \left(\sqrt{5} \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) - \sqrt{5} \sqrt{\sqrt{5}-2} - \sqrt{5} - 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} \right) \right.$$

$$+ 256 \left(\sqrt{5} \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) + \sqrt{5} - 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \sqrt{\sqrt{5}-2} + 2 \sqrt{\tan(x)^2 + 2 \tan(x) + 2} \right) \left.$$

$$+ \frac{\left(\pi + 4 \arctan \left(-\frac{1}{2} \left(2\sqrt{5}\sqrt{\sqrt{5}-2} + \sqrt{5} + 4\sqrt{\sqrt{5}-2} + 3 \right) \right) \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) \right)}{4(\sqrt{5}-1)}$$

$$- \frac{\left(\pi + 4 \arctan \left(\frac{1}{2} \left(2\sqrt{5}\sqrt{\sqrt{5}-2} - \sqrt{5} + 4\sqrt{\sqrt{5}-2} - 3 \right) \right) \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) \right) - \frac{3}{2} \right)}{4(\sqrt{5}-1)}$$

$$- \log \left(\sqrt{\tan(x)^2 + 2 \tan(x) + 2} - \tan(x) - 1 \right)$$

[In] integrate((2+2*tan(x)+tan(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/4*sqrt(2*sqrt(5) - 2)*log(256*(sqrt(5)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) + sqrt(5)*sqrt(sqrt(5) - 2) - sqrt(5) - 2*sqrt(tan(x)^2 + 2*tan(x) + 2) - 2*sqrt(sqrt(5) - 2) + 2*tan(x) + 2)^2 + 256*(sqrt(5)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) + sqrt(5) - 2*sqrt(tan(x)^2 + 2*tan(x) + 2) + sqrt(tan(x)^2 + 2*tan(x) - 2)^2) + 1/4*sqrt(2*sqrt(5) - 2)*log(256*(sqrt(5)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) - sqrt(5)*sqrt(sqrt(5) - 2) - sq

```

rt(5) - 2*sqrt(tan(x)^2 + 2*tan(x) + 2) + 2*sqrt(sqrt(5) - 2) + 2*tan(x) +
2)^2 + 256*(sqrt(5)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) + sqrt(5) - 2*
sqrt(tan(x)^2 + 2*tan(x) + 2) - sqrt(sqrt(5) - 2) + 2*tan(x) - 2)^2) + 1/4*
(pi + 4*arctan(-1/2*(2*sqrt(5)*sqrt(sqrt(5) - 2) + sqrt(5) + 4*sqrt(sqrt(5)
- 2) + 3)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) + 3/2*sqrt(5)*sqrt(sqrt
(5) - 2) + 1/2*sqrt(5) + 7/2*sqrt(sqrt(5) - 2) + 3/2))*sqrt(2*sqrt(5) - 2)/
(sqrt(5) - 1) - 1/4*(pi + 4*arctan(1/2*(2*sqrt(5)*sqrt(sqrt(5) - 2) - sqrt(
5) + 4*sqrt(sqrt(5) - 2) - 3)*(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(x)) - 3/
2*sqrt(5)*sqrt(sqrt(5) - 2) + 1/2*sqrt(5) - 7/2*sqrt(sqrt(5) - 2) + 3/2))*s
qrt(2*sqrt(5) - 2)/(sqrt(5) - 1) - log(sqrt(tan(x)^2 + 2*tan(x) + 2) - tan(
x) - 1)

```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{2 + 2 \tan(x) + \tan^2(x)} dx = \int \sqrt{\tan(x)^2 + 2 \tan(x) + 2} dx$$

[In] int((2*tan(x) + tan(x)^2 + 2)^(1/2),x)

[Out] int((2*tan(x) + tan(x)^2 + 2)^(1/2), x)

3.9 $\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx$

Optimal result	92
Rubi [A] (verified)	92
Mathematica [A] (verified)	94
Maple [A] (verified)	95
Fricas [A] (verification not implemented)	95
Sympy [F]	95
Maxima [A] (verification not implemented)	96
Giac [F]	96
Mupad [B] (verification not implemented)	96

Optimal result

Integrand size = 12, antiderivative size = 41

$$\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx = \frac{1}{2} \arctan\left(\sqrt{-1 + \sec(x)}\right) - \arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \frac{1}{2} \cos(x) \sqrt{-1 + \sec(x)}$$

[Out] 1/2*arctan((-1+sec(x))^(1/2))-arctan((-1+sec(x))^(1/2))*cos(x)+1/2*cos(x)*(-1+sec(x))^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {4420, 5311, 12, 248, 44, 65, 209}

$$\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx = \frac{1}{2} \arctan\left(\sqrt{\sec(x) - 1}\right) - \cos(x) \arctan\left(\sqrt{\sec(x) - 1}\right) + \frac{1}{2} \cos(x) \sqrt{\sec(x) - 1}$$

[In] Int[ArcTan[Sqrt[-1 + Sec[x]]]*Sin[x],x]

[Out] ArcTan[Sqrt[-1 + Sec[x]]]/2 - ArcTan[Sqrt[-1 + Sec[x]]]*Cos[x] + (Cos[x]*Sqrt[-1 + Sec[x]])/2

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 248

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Subst[Int[(a + b/x^n)^p/x^2, x], x, 1/x] /; FreeQ[{a, b, p}, x] && ILtQ[n, 0]
```

Rule 4420

```
Int[(u_)*(F_)[(c_.)*((a_.) + (b_.)*(x_))], x_Symbol] := With[{d = FreeFactors[Cos[c*(a + b*x)], x]}, Dist[-d/(b*c), Subst[Int[SubstFor[1, Cos[c*(a + b*x)]]/d, u, x], x], x, Cos[c*(a + b*x)]/d, x] /; FunctionOfQ[Cos[c*(a + b*x)]]/d, u, x, True]] /; FreeQ[{a, b, c}, x] && (EqQ[F, Sin] || EqQ[F, sin])
```

Rule 5311

```
Int[ArcTan[u_], x_Symbol] := Simp[x*ArcTan[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/(1 + u^2)), x], x] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \arctan\left(\sqrt{-1 + \frac{1}{x}}\right) dx, x, \cos(x)\right)$$

$$\begin{aligned}
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \text{Subst}\left(\int -\frac{1}{2\sqrt{-1 + \frac{1}{x}}} dx, x, \cos(x)\right) \\
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{-1 + \frac{1}{x}}} dx, x, \cos(x)\right) \\
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{-1 + xx^2}} dx, x, \sec(x)\right) \\
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \frac{1}{2} \cos(x) \sqrt{-1 + \sec(x)} \\
&\quad + \frac{1}{4} \text{Subst}\left(\int \frac{1}{\sqrt{-1 + xx}} dx, x, \sec(x)\right) \\
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \frac{1}{2} \cos(x) \sqrt{-1 + \sec(x)} \\
&\quad + \frac{1}{2} \text{Subst}\left(\int \frac{1}{1 + x^2} dx, x, \sqrt{-1 + \sec(x)}\right) \\
&= \frac{1}{2} \arctan\left(\sqrt{-1 + \sec(x)}\right) - \arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) + \frac{1}{2} \cos(x) \sqrt{-1 + \sec(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.71

$$\begin{aligned}
&\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx \\
&= -\arctan\left(\sqrt{-1 + \sec(x)}\right) \cos(x) \\
&\quad + \frac{1}{2} \left(\cos(x) + \arctan\left(\frac{\tan\left(\frac{x}{2}\right)}{\sqrt{\frac{\cos(x)}{1 + \cos(x)}}}\right) \sqrt{\frac{\cos(x)}{1 + \cos(x)}} \cot\left(\frac{x}{2}\right) \right) \sqrt{-1 + \sec(x)}
\end{aligned}$$

[In] Integrate[ArcTan[Sqrt[-1 + Sec[x]]]*Sin[x],x]

[Out] -(ArcTan[Sqrt[-1 + Sec[x]]]*Cos[x]) + ((Cos[x] + ArcTan[Tan[x/2]/Sqrt[Cos[x]/(1 + Cos[x])]]*Sqrt[Cos[x]/(1 + Cos[x])]*Cot[x/2])*Sqrt[-1 + Sec[x]])/2

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{\arctan\left(\sqrt{-1+\sec(x)}\right)}{\sec(x)} + \frac{\sqrt{-1+\sec(x)}}{2\sec(x)} + \frac{\arctan\left(\sqrt{-1+\sec(x)}\right)}{2}$	36
default	$-\frac{\arctan\left(\sqrt{-1+\sec(x)}\right)}{\sec(x)} + \frac{\sqrt{-1+\sec(x)}}{2\sec(x)} + \frac{\arctan\left(\sqrt{-1+\sec(x)}\right)}{2}$	36

[In] `int(arctan((-1+sec(x))^(1/2))*sin(x),x,method=_RETURNVERBOSE)`

[Out] `-1/sec(x)*arctan((-1+sec(x))^(1/2))+1/2*(-1+sec(x))^(1/2)/sec(x)+1/2*arctan((-1+sec(x))^(1/2))`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int \arctan\left(\sqrt{-1+\sec(x)}\right) \sin(x) dx = -\frac{1}{2} (2 \cos(x) - 1) \arctan\left(\sqrt{\sec(x) - 1}\right) + \frac{1}{2} \sqrt{-\frac{\cos(x) - 1}{\cos(x)}} \cos(x)$$

[In] `integrate(arctan((-1+sec(x))^(1/2))*sin(x),x, algorithm="fricas")`

[Out] `-1/2*(2*cos(x) - 1)*arctan(sqrt(sec(x) - 1)) + 1/2*sqrt(-(cos(x) - 1)/cos(x)))*cos(x)`

Sympy [F]

$$\int \arctan\left(\sqrt{-1+\sec(x)}\right) \sin(x) dx = \int \sin(x) \operatorname{atan}\left(\sqrt{\sec(x) - 1}\right) dx$$

[In] `integrate(atan((-1+sec(x))**(1/2))*sin(x),x)`

[Out] `Integral(sin(x)*atan(sqrt(sec(x) - 1)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx = -\arctan\left(\sqrt{-\frac{\cos(x) - 1}{\cos(x)}}\right) \cos(x) - \frac{\sqrt{-\frac{\cos(x) - 1}{\cos(x)}}}{2\left(\frac{\cos(x) - 1}{\cos(x)} - 1\right)} + \frac{1}{2} \arctan\left(\sqrt{-\frac{\cos(x) - 1}{\cos(x)}}\right)$$

[In] integrate(arctan((-1+sec(x))^(1/2))*sin(x),x, algorithm="maxima")

[Out] -arctan(sqrt(-(cos(x) - 1)/cos(x)))*cos(x) - 1/2*sqrt(-(cos(x) - 1)/cos(x)) /((cos(x) - 1)/cos(x) - 1) + 1/2*arctan(sqrt(-(cos(x) - 1)/cos(x)))

Giac [F]

$$\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx = \int \arctan\left(\sqrt{\sec(x) - 1}\right) \sin(x) dx$$

[In] integrate(arctan((-1+sec(x))^(1/2))*sin(x),x, algorithm="giac")

[Out] undef

Mupad [B] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \arctan\left(\sqrt{-1 + \sec(x)}\right) \sin(x) dx = -\operatorname{atan}\left(\sqrt{\frac{1}{\cos(x)} - 1}\right) \cos(x) - \frac{\cos(x) \left(\frac{3 \operatorname{asin}\left(\sqrt{\cos(x)}\right)}{2 \cos(x)^{3/2}} - \frac{3 \sqrt{1 - \cos(x)}}{2 \cos(x)} \right) \sqrt{1 - \cos(x)}}{3 \sqrt{\frac{1}{\cos(x)} - 1}}$$

[In] int(atan((1/cos(x) - 1)^(1/2))*sin(x),x)

[Out] - atan((1/cos(x) - 1)^(1/2))*cos(x) - (cos(x)*((3*asin(cos(x)^(1/2)))/(2*cos(x)^(3/2)) - (3*(1 - cos(x))^(1/2))/(2*cos(x))))*(1 - cos(x))^(1/2))/(3*(1/cos(x) - 1)^(1/2))

3.10 $\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx$

Optimal result	97
Rubi [A] (verified)	97
Mathematica [A] (verified)	99
Maple [F]	99
Fricas [A] (verification not implemented)	99
Sympy [A] (verification not implemented)	99
Maxima [F]	100
Giac [A] (verification not implemented)	100
Mupad [F(-1)]	100

Optimal result

Integrand size = 19, antiderivative size = 44

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \frac{1}{10} e^{\arcsin(x)} \left(3x + x^3 - 3\sqrt{1-x^2} - 3x^2\sqrt{1-x^2} \right)$$

[Out] 1/10*exp(arcsin(x))*(3*x+x^3-3*(-x^2+1)^(1/2)-3*x^2*(-x^2+1)^(1/2))

Rubi [A] (verified)

Time = 0.54 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.41, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {4920, 6873, 6852, 4519, 4517}

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \frac{1}{10} x^3 e^{\arcsin(x)} - \frac{3}{10} \sqrt{1-x^2} x^2 e^{\arcsin(x)} - \frac{3}{10} \sqrt{1-x^2} e^{\arcsin(x)} + \frac{3}{10} x e^{\arcsin(x)}$$

[In] Int[(E^ArcSin[x]*x^3)/Sqrt[1 - x^2],x]

[Out] (3*E^ArcSin[x]*x)/10 + (E^ArcSin[x]*x^3)/10 - (3*E^ArcSin[x]*Sqrt[1 - x^2])/10 - (3*E^ArcSin[x]*x^2*Sqrt[1 - x^2])/10

Rule 4517

```
Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] :=
  Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; F
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Rule 4519

```
Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)]^(n_), x_Symbol]
:> Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]^n/(e^2*n^2 + b^2*c^2*Log[F]^2)), x]
+ (Dist[(n*(n - 1)*e^2)/(e^2*n^2 + b^2*c^2*Log[F]^2), Int[F^(c*(a + b*x))*Sin[d + e*x]^(n - 2), x], x]
- Simp[e*n*F^(c*(a + b*x))*Cos[d + e*x]*(Sin[d + e*x]^(n - 1)/(e^2*n^2 + b^2*c^2*Log[F]^2)), x])
/; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2*n^2 + b^2*c^2*Log[F]^2, 0] && GtQ[n, 1]
```

Rule 4920

```
Int[(u_.)*(f_)^(ArcSin[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol]
:> Dist[1/b, Subst[Int[(u /. x -> -a/b + Sin[x]/b)*f^(c*x^n)*Cos[x], x], x, ArcSin[a + b*x]], x]
/; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rule 6852

```
Int[(u_.)*((a_.)*(v_)^(m_.))^(p_), x_Symbol]
:> Dist[a^IntPart[p]*((a*v^m)^FracPart[p]/v^(m*FracPart[p])), Int[u*v^(m*p), x], x]
/; FreeQ[{a, m, p}, x] && !IntegerQ[p] && !FreeQ[v, x] && !(EqQ[a, 1] && EqQ[m, 1]) && !(EqQ[v, x] && EqQ[m, 1])
```

Rule 6873

```
Int[u_, x_Symbol]
:> With[{v = NormalizeIntegrand[u, x]}, Int[v, x] /; v != u]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{e^x \cos(x) \sin^3(x)}{\sqrt{1 - \sin^2(x)}} dx, x, \arcsin(x) \right) \\
&= \text{Subst} \left(\int \frac{e^x \cos(x) \sin^3(x)}{\sqrt{\cos^2(x)}} dx, x, \arcsin(x) \right) \\
&= 1 \text{Subst} \left(\int e^x \sin^3(x) dx, x, \arcsin(x) \right) \\
&= \frac{1}{10} e^{\arcsin(x)} x^3 - \frac{3}{10} e^{\arcsin(x)} x^2 \sqrt{1 - x^2} + \frac{3}{5} \text{Subst} \left(\int e^x \sin(x) dx, x, \arcsin(x) \right) \\
&= \frac{3}{10} e^{\arcsin(x)} x + \frac{1}{10} e^{\arcsin(x)} x^3 - \frac{3}{10} e^{\arcsin(x)} \sqrt{1 - x^2} - \frac{3}{10} e^{\arcsin(x)} x^2 \sqrt{1 - x^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.86

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = -\frac{1}{40} e^{\arcsin(x)} \left(15 \left(-x + \sqrt{1-x^2} \right) - 3 \cos(3 \arcsin(x)) + \sin(3 \arcsin(x)) \right)$$

[In] Integrate[(E^ArcSin[x]*x^3)/Sqrt[1 - x^2], x]

[Out] -1/40*(E^ArcSin[x]*(15*(-x + Sqrt[1 - x^2]) - 3*Cos[3*ArcSin[x]] + Sin[3*ArcSin[x]]))

Maple [F]

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{-x^2+1}} dx$$

[In] int(exp(arcsin(x))*x^3/(-x^2+1)^(1/2), x)

[Out] int(exp(arcsin(x))*x^3/(-x^2+1)^(1/2), x)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.64

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \frac{1}{10} \left(x^3 - 3(x^2 + 1)\sqrt{-x^2 + 1} + 3x \right) e^{\arcsin(x)}$$

[In] integrate(exp(arcsin(x))*x^3/(-x^2+1)^(1/2), x, algorithm="fricas")

[Out] 1/10*(x^3 - 3*(x^2 + 1)*sqrt(-x^2 + 1) + 3*x)*e^arcsin(x)

Sympy [A] (verification not implemented)

Time = 0.38 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.27

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \frac{x^3 e^{\arcsin(x)}}{10} - \frac{3x^2 \sqrt{1-x^2} e^{\arcsin(x)}}{10} + \frac{3x e^{\arcsin(x)}}{10} - \frac{3\sqrt{1-x^2} e^{\arcsin(x)}}{10}$$

[In] integrate(exp(asin(x))*x**3/(-x**2+1)**(1/2), x)

[Out] x**3*exp(asin(x))/10 - 3*x**2*sqrt(1 - x**2)*exp(asin(x))/10 + 3*x*exp(asin(x))/10 - 3*sqrt(1 - x**2)*exp(asin(x))/10

Maxima [F]

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \int \frac{x^3 e^{\arcsin(x)}}{\sqrt{-x^2+1}} dx$$

[In] integrate(exp(arcsin(x))*x^3/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(x^3*e^arcsin(x)/sqrt(-x^2 + 1), x)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.05

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \frac{1}{10} (x^2 - 1) x e^{\arcsin(x)} + \frac{3}{10} (-x^2 + 1)^{\frac{3}{2}} e^{\arcsin(x)} + \frac{2}{5} x e^{\arcsin(x)} - \frac{3}{5} \sqrt{-x^2 + 1} e^{\arcsin(x)}$$

[In] integrate(exp(arcsin(x))*x^3/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/10*(x^2 - 1)*x*e^arcsin(x) + 3/10*(-x^2 + 1)^(3/2)*e^arcsin(x) + 2/5*x*e^arcsin(x) - 3/5*sqrt(-x^2 + 1)*e^arcsin(x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\arcsin(x)} x^3}{\sqrt{1-x^2}} dx = \int \frac{x^3 e^{\arcsin(x)}}{\sqrt{1-x^2}} dx$$

[In] int((x^3*exp(asin(x)))/(1 - x^2)^(1/2),x)

[Out] int((x^3*exp(asin(x)))/(1 - x^2)^(1/2), x)

$$3.11 \quad \int \frac{x \log(1+x^2) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx$$

Optimal result	101
Rubi [A] (verified)	101
Mathematica [A] (verified)	103
Maple [F]	104
Fricas [A] (verification not implemented)	104
Sympy [F(-1)]	104
Maxima [F]	104
Giac [F]	105
Mupad [F(-1)]	105

Optimal result

Integrand size = 29, antiderivative size = 68

$$\int \frac{x \log(1+x^2) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx = 4x - 2 \arctan(x) - x \log(1+x^2) - 2\sqrt{1+x^2} \log(x+\sqrt{1+x^2}) + \sqrt{1+x^2} \log(1+x^2) \log(x+\sqrt{1+x^2})$$

[Out] 4*x-2*arctan(x)-x*ln(x^2+1)-2*ln(x+(x^2+1)^(1/2))*(x^2+1)^(1/2)+ln(x^2+1)*1n(x+(x^2+1)^(1/2))*(x^2+1)^(1/2)

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.276$, Rules used = {267, 2634, 8, 2637, 12, 2498, 327, 209}

$$\int \frac{x \log(1+x^2) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx = -2 \arctan(x) + x(-\log(x^2+1)) + \sqrt{x^2+1} \log(x^2+1) \log(\sqrt{x^2+1}+x) - 2\sqrt{x^2+1} \log(\sqrt{x^2+1}+x) + 4x$$

[In] Int[(x*Log[1+x^2]*Log[x+Sqrt[1+x^2]])/Sqrt[1+x^2],x]

[Out] 4*x - 2*ArcTan[x] - x*Log[1+x^2] - 2*Sqrt[1+x^2]*Log[x+Sqrt[1+x^2]] + Sqrt[1+x^2]*Log[1+x^2]*Log[x+Sqrt[1+x^2]]

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 327

```
Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2498

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)], x_Symbol] := Simp[x*Log[c*(d
+ e*x^n)^p], x] - Dist[e*n*p, Int[x^n/(d + e*x^n), x], x] /; FreeQ[{c, d,
e, n, p}, x]
```

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rule 2637

```
Int[Log[v_]*Log[w_]*(u_), x_Symbol] := With[{z = IntHide[u, x]}, Dist[Log[v
]*Log[w], z, x] + (-Int[SimplifyIntegrand[z*Log[w]*(D[v, x]/v), x], x] - In
t[SimplifyIntegrand[z*Log[v]*(D[w, x]/w), x], x]) /; InverseFunctionFreeQ[z
```

, x]] /; InverseFunctionFreeQ[v, x] && InverseFunctionFreeQ[w, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \sqrt{1+x^2} \log(1+x^2) \log(x+\sqrt{1+x^2}) \\
 &\quad - \int \log(1+x^2) dx - \int \frac{2x \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx \\
 &= -x \log(1+x^2) + \sqrt{1+x^2} \log(1+x^2) \log(x+\sqrt{1+x^2}) \\
 &\quad + 2 \int \frac{x^2}{1+x^2} dx - 2 \int \frac{x \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx \\
 &= 2x - x \log(1+x^2) - 2\sqrt{1+x^2} \log(x+\sqrt{1+x^2}) \\
 &\quad + \sqrt{1+x^2} \log(1+x^2) \log(x+\sqrt{1+x^2}) + 2 \int 1 dx - 2 \int \frac{1}{1+x^2} dx \\
 &= 4x - 2 \arctan(x) - x \log(1+x^2) - 2\sqrt{1+x^2} \log(x+\sqrt{1+x^2}) \\
 &\quad + \sqrt{1+x^2} \log(1+x^2) \log(x+\sqrt{1+x^2})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.94

$$\begin{aligned}
 \int \frac{x \log(1+x^2) \log(x+\sqrt{1+x^2})}{\sqrt{1+x^2}} dx &= 4x - 2 \arctan(x) - 2\sqrt{1+x^2} \log(x+\sqrt{1+x^2}) \\
 &\quad + \log(1+x^2) \left(-x + \sqrt{1+x^2} \log(x+\sqrt{1+x^2}) \right)
 \end{aligned}$$

[In] Integrate[(x*Log[1 + x^2]*Log[x + Sqrt[1 + x^2]])/Sqrt[1 + x^2],x]

[Out] 4*x - 2*ArcTan[x] - 2*Sqrt[1 + x^2]*Log[x + Sqrt[1 + x^2]] + Log[1 + x^2]*(
-x + Sqrt[1 + x^2]*Log[x + Sqrt[1 + x^2]])

Maple [F]

$$\int \frac{x \ln(x^2 + 1) \ln(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] `int(x*ln(x^2+1)*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x)`

[Out] `int(x*ln(x^2+1)*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.63

$$\int \frac{x \log(1 + x^2) \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \sqrt{x^2 + 1} (\log(x^2 + 1) - 2) \log(x + \sqrt{x^2 + 1}) - x \log(x^2 + 1) + 4x - 2 \arctan(x)$$

[In] `integrate(x*log(x^2+1)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="fricas")`

[Out] `sqrt(x^2 + 1)*(log(x^2 + 1) - 2)*log(x + sqrt(x^2 + 1)) - x*log(x^2 + 1) + 4*x - 2*arctan(x)`

Sympy [F(-1)]

Timed out.

$$\int \frac{x \log(1 + x^2) \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \text{Timed out}$$

[In] `integrate(x*ln(x**2+1)*ln(x+(x**2+1)**(1/2))/(x**2+1)**(1/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{x \log(1 + x^2) \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \int \frac{x \log(x^2 + 1) \log(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] `integrate(x*log(x^2+1)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="maxima")`

[Out] `-(2*x^2 - (x^2 + 1)*log(x^2 + 1) + 2)*log(x + sqrt(x^2 + 1))/sqrt(x^2 + 1) + integrate((log(x^2 + 1) - 2)/(x^2 + sqrt(x^2 + 1)*x), x) - integrate(-(2*x^2 - (x^2 + 1)*log(x^2 + 1) + 2)/(sqrt(x^2 + 1)*x), x)`

Giac [F]

$$\int \frac{x \log(1+x^2) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \int \frac{x \log(x^2+1) \log(x + \sqrt{x^2+1})}{\sqrt{x^2+1}} dx$$

[In] integrate(x*log(x^2+1)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(x*log(x^2 + 1)*log(x + sqrt(x^2 + 1))/sqrt(x^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(1+x^2) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \int \frac{x \ln(x^2+1) \ln(x + \sqrt{x^2+1})}{\sqrt{x^2+1}} dx$$

[In] int((x*log(x^2 + 1)*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2),x)

[Out] int((x*log(x^2 + 1)*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2), x)

3.12 $\int \arctan(x + \sqrt{1-x^2}) dx$

Optimal result	106
Rubi [C] (verified)	106
Mathematica [C] (verified)	111
Maple [C] (verified)	112
Fricas [A] (verification not implemented)	113
Sympy [F(-1)]	113
Maxima [F]	114
Giac [B] (verification not implemented)	114
Mupad [B] (verification not implemented)	115

Optimal result

Integrand size = 14, antiderivative size = 141

$$\int \arctan(x + \sqrt{1-x^2}) dx = -\frac{\arcsin(x)}{2} + \frac{1}{4}\sqrt{3} \arctan\left(\frac{-1 + \sqrt{3}x}{\sqrt{1-x^2}}\right) + \frac{1}{4}\sqrt{3} \arctan\left(\frac{1 + \sqrt{3}x}{\sqrt{1-x^2}}\right) - \frac{1}{4}\sqrt{3} \arctan\left(\frac{-1 + 2x^2}{\sqrt{3}}\right) + x \arctan(x + \sqrt{1-x^2}) - \frac{1}{4} \operatorname{arctanh}(x\sqrt{1-x^2}) - \frac{1}{8} \log(1-x^2+x^4)$$

```
[Out] -1/2*arcsin(x)+x*arctan(x+(-x^2+1)^(1/2))-1/4*arctanh(x*(-x^2+1)^(1/2))-1/8*ln(x^4-x^2+1)-1/4*arctan(1/3*(2*x^2-1)*3^(1/2))*3^(1/2)+1/4*arctan((-1+x*3^(1/2))/(-x^2+1)^(1/2))*3^(1/2)+1/4*arctan((1+x*3^(1/2))/(-x^2+1)^(1/2))*3^(1/2)
```

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.60 (sec) , antiderivative size = 269, normalized size of antiderivative = 1.91, number of steps used = 40, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 1.071$, Rules

used = {5311, 12, 6874, 222, 1128, 648, 632, 210, 642, 1188, 399, 385, 211, 1307, 1121}

$$\int \arctan(x + \sqrt{1-x^2}) dx = -\frac{\arcsin(x)}{2} + \frac{1}{4}\sqrt{3} \arctan\left(\frac{1-2x^2}{\sqrt{3}}\right) + \frac{1}{12}(-\sqrt{3} + 3i) \arctan\left(\frac{x}{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}\sqrt{1-x^2}}\right) + \frac{\arctan\left(\frac{x}{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}\sqrt{1-x^2}}\right)}{\sqrt{3}} - \frac{1}{12}(\sqrt{3} + 3i) \arctan\left(\frac{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}x}{\sqrt{1-x^2}}\right) + \frac{\arctan\left(\frac{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}x}{\sqrt{1-x^2}}\right)}{\sqrt{3}} + x \arctan(\sqrt{1-x^2} + x) - \frac{1}{8} \log(x^4 - x^2 + 1)$$

[In] Int[ArcTan[x + Sqrt[1 - x^2]],x]

[Out] -1/2*ArcSin[x] + (Sqrt[3]*ArcTan[(1 - 2*x^2)/Sqrt[3]])/4 + ArcTan[x/(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*Sqrt[1 - x^2])]/Sqrt[3] + ((3*I - Sqrt[3])*ArcTan[x/(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*Sqrt[1 - x^2])])/12 + ArcTan[(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*x)/Sqrt[1 - x^2]]/Sqrt[3] - ((3*I + Sqrt[3])*ArcTan[(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*x)/Sqrt[1 - x^2])])/12 + x*ArcTan[x + Sqrt[1 - x^2]] - Log[1 - x^2 + x^4]/8

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 222

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Di
st[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^
n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*
d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 632

```
Int(((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int(((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int(((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1121

```
Int[(x_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2,
Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]
```

Rule 1128

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dis
t[1/2, Subst[Int[x^((m - 1)/2)*(a + b*x + c*x^2)^p, x], x, x^2], x] /; Free
```

$Q[\{a, b, c, p\}, x] \ \&\& \ \text{IntegerQ}[(m - 1)/2]$

Rule 1188

$\text{Int}[\{(d_)+(e_)(x_)^2\}^q/\{(a_)+(b_)(x_)^2+(c_)(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{r = \text{Rt}[b^2 - 4ac, 2]\}, \text{Dist}[2(c/r), \text{Int}[(d + e x^2)^q/(b - r + 2c x^2), x], x] - \text{Dist}[2(c/r), \text{Int}[(d + e x^2)^q/(b + r + 2c x^2), x], x]] \ /; \text{FreeQ}[\{a, b, c, d, e, q\}, x] \ \&\& \ \text{NeQ}[b^2 - 4ac, 0] \ \&\& \ \text{NeQ}[c d^2 - b d e + a e^2, 0] \ \&\& \ !\text{IntegerQ}[q]$

Rule 1307

$\text{Int}[\{(f_)(x_)^m\} \{(d_)+(e_)(x_)^2\}^q/\{(a_)+(b_)(x_)^2+(c_)(x_)^4\}, x_Symbol] \rightarrow \text{Dist}[e(f^2/c), \text{Int}[(f x)^{m-2}(d + e x^2)^{q-1}, x], x] - \text{Dist}[f^2/c, \text{Int}[(f x)^{m-2}(d + e x^2)^{q-1}(\text{Simp}[a e - (c d - b e)x^2, x]/(a + b x^2 + c x^4)), x], x] \ /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{NeQ}[b^2 - 4ac, 0] \ \&\& \ !\text{IntegerQ}[q] \ \&\& \ \text{GtQ}[q, 0] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LeQ}[m, 3]$

Rule 5311

$\text{Int}[\text{ArcTan}[u_], x_Symbol] \rightarrow \text{Simp}[x \text{ArcTan}[u], x] - \text{Int}[\text{SimplifyIntegrand}[x * (D[u, x]/(1 + u^2)), x], x] \ /; \text{InverseFunctionFreeQ}[u, x]$

Rule 6874

$\text{Int}[u_, x_Symbol] \rightarrow \text{With}[\{v = \text{ExpandIntegrand}[u, x]\}, \text{Int}[v, x] \ /; \text{SumQ}[v]]$

Rubi steps

$$\begin{aligned} \text{integral} &= x \arctan\left(x + \sqrt{1 - x^2}\right) - \int \frac{x\left(1 - \frac{x}{\sqrt{1 - x^2}}\right)}{2(1 + x\sqrt{1 - x^2})} dx \\ &= x \arctan\left(x + \sqrt{1 - x^2}\right) - \frac{1}{2} \int \frac{x\left(1 - \frac{x}{\sqrt{1 - x^2}}\right)}{1 + x\sqrt{1 - x^2}} dx \\ &= x \arctan\left(x + \sqrt{1 - x^2}\right) - \frac{1}{2} \int \left(\frac{x^2}{-x + x^3 - \sqrt{1 - x^2}} + \frac{x}{1 + x\sqrt{1 - x^2}}\right) dx \\ &= x \arctan\left(x + \sqrt{1 - x^2}\right) - \frac{1}{2} \int \frac{x^2}{-x + x^3 - \sqrt{1 - x^2}} dx - \frac{1}{2} \int \frac{x}{1 + x\sqrt{1 - x^2}} dx \end{aligned}$$

$$\begin{aligned}
&= x \arctan \left(x + \sqrt{1-x^2} \right) - \frac{1}{2} \int \left(\frac{x}{1-x^2+x^4} - \frac{x^2\sqrt{1-x^2}}{1-x^2+x^4} \right) dx \\
&\quad - \frac{1}{2} \int \left(-\frac{1}{\sqrt{1-x^2}} + \frac{x^3}{1-x^2+x^4} + \frac{\sqrt{1-x^2}}{1-x^2+x^4} - \frac{x^2\sqrt{1-x^2}}{1-x^2+x^4} \right) dx \\
&= x \arctan \left(x + \sqrt{1-x^2} \right) + \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx - \frac{1}{2} \int \frac{x}{1-x^2+x^4} dx \\
&\quad - \frac{1}{2} \int \frac{x^3}{1-x^2+x^4} dx - \frac{1}{2} \int \frac{\sqrt{1-x^2}}{1-x^2+x^4} dx + 2 \left(\frac{1}{2} \int \frac{x^2\sqrt{1-x^2}}{1-x^2+x^4} dx \right) \\
&= \frac{\arcsin(x)}{2} + x \arctan \left(x + \sqrt{1-x^2} \right) \\
&\quad - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, x^2 \right) - \frac{1}{4} \text{Subst} \left(\int \frac{x}{1-x+x^2} dx, x, x^2 \right) \\
&\quad + 2 \left(-\left(\frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx \right) + \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}(1-x^2+x^4)} dx \right) \\
&\quad + \frac{i \int \frac{\sqrt{1-x^2}}{-1-i\sqrt{3}+2x^2} dx}{\sqrt{3}} - \frac{i \int \frac{\sqrt{1-x^2}}{-1+i\sqrt{3}+2x^2} dx}{\sqrt{3}} \\
&= \frac{\arcsin(x)}{2} + x \arctan \left(x + \sqrt{1-x^2} \right) - \frac{1}{8} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, x^2 \right) \\
&\quad - \frac{1}{8} \text{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, x^2 \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{-3-x^2} dx, x, -1+2x^2 \right) \\
&\quad + 2 \left(-\frac{\arcsin(x)}{2} - \frac{i \int \frac{1}{\sqrt{1-x^2}(-1-i\sqrt{3}+2x^2)} dx}{\sqrt{3}} + \frac{i \int \frac{1}{\sqrt{1-x^2}(-1+i\sqrt{3}+2x^2)} dx}{\sqrt{3}} \right) \\
&\quad + \frac{1}{6} (3-i\sqrt{3}) \int \frac{1}{\sqrt{1-x^2}(-1+i\sqrt{3}+2x^2)} dx \\
&\quad + \frac{1}{6} (3+i\sqrt{3}) \int \frac{1}{\sqrt{1-x^2}(-1-i\sqrt{3}+2x^2)} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{\arcsin(x)}{2} + \frac{\arctan\left(\frac{1-2x^2}{\sqrt{3}}\right)}{2\sqrt{3}} + x \arctan\left(x + \sqrt{1-x^2}\right) \\
&\quad - \frac{1}{8} \log(1-x^2+x^4) + \frac{1}{4} \text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2x^2\right) \\
&\quad + 2 \left(-\frac{\arcsin(x)}{2} + \frac{i \text{Subst}\left(\int \frac{1}{-1+i\sqrt{3}-(-1-i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right)}{\sqrt{3}} \right. \\
&\qquad\qquad\qquad \left. - \frac{i \text{Subst}\left(\int \frac{1}{-1-i\sqrt{3}-(-1+i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right)}{\sqrt{3}} \right) \\
&\quad + \frac{1}{6} (3-i\sqrt{3}) \text{Subst}\left(\int \frac{1}{-1+i\sqrt{3}-(-1-i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\
&\quad + \frac{1}{6} (3+i\sqrt{3}) \text{Subst}\left(\int \frac{1}{-1-i\sqrt{3}-(-1+i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\
&= \frac{\arcsin(x)}{2} + \frac{1}{4} \sqrt{3} \arctan\left(\frac{1-2x^2}{\sqrt{3}}\right) + \frac{1}{12} (3i-\sqrt{3}) \arctan\left(\frac{x}{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}}\sqrt{1-x^2}}\right) \\
&\quad - \frac{1}{12} (3i+\sqrt{3}) \arctan\left(\frac{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}}x}{\sqrt{1-x^2}}\right) \\
&\quad + 2 \left(-\frac{\arcsin(x)}{2} + \frac{\arctan\left(\frac{x}{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}}\sqrt{1-x^2}}\right)}{2\sqrt{3}} + \frac{\arctan\left(\frac{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}}x}{\sqrt{1-x^2}}\right)}{2\sqrt{3}} \right) \\
&\quad + x \arctan\left(x + \sqrt{1-x^2}\right) - \frac{1}{8} \log(1-x^2+x^4)
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.13 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.60

$$\int \arctan\left(x + \sqrt{1-x^2}\right) dx = -\arctan\left(\frac{x}{-1 + \sqrt{1-x^2}}\right) + x \arctan\left(x + \sqrt{1-x^2}\right) - \log(x) + \frac{1}{2} \log\left(-1 + \sqrt{1-x^2}\right) - \frac{1}{2} \operatorname{RootSum}\left[1 - 2\#1 + 2\#1^2 + 2\#1^3 + \#1^4\right] \frac{-\log(x) + \log\left(-1 + \sqrt{1-x^2} - x\#1\right) - \log(x)\#1 + \log\left(-1 + \sqrt{1-x^2} - x\#1\right)\#1 - 3\log(x)\#1^2 - \log(x)\#1^3 + \log\left(-1 + \sqrt{1-x^2} - x\#1\right)\#1^2 - \log(x)\#1^3}{-1 + 2\#1 + 3\#1^2 + \#1^3}$$

[In] Integrate[ArcTan[x + Sqrt[1 - x^2]],x]

[Out] -ArcTan[x/(-1 + Sqrt[1 - x^2])] + x*ArcTan[x + Sqrt[1 - x^2]] - Log[x] + Log[-1 + Sqrt[1 - x^2]]/2 - RootSum[1 - 2*#1 + 2*#1^2 + 2*#1^3 + #1^4 & , (-Log[x] + Log[-1 + Sqrt[1 - x^2] - x*#1] - Log[x]*#1 + Log[-1 + Sqrt[1 - x^2] - x*#1]*#1 - 3*Log[x]*#1^2 + 3*Log[-1 + Sqrt[1 - x^2] - x*#1]*#1^2 - Log[x]*#1^3 + Log[-1 + Sqrt[1 - x^2] - x*#1]*#1^3)/(-1 + 2*#1 + 3*#1^2 + 2*#1^3) &]/2

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.10 (sec) , antiderivative size = 471, normalized size of antiderivative = 3.34

method	result
default	$x \arctan\left(x + \sqrt{-x^2 + 1}\right) - \frac{\ln(x^4 - x^2 + 1)}{8} - \frac{\arctan\left(\frac{(2x^2 - 1)\sqrt{3}}{3}\right)\sqrt{3}}{4} - \frac{\left(\frac{i\sqrt{3}}{12} + \frac{1}{4}\right) \ln\left(\frac{(\sqrt{-x^2 + 1} - 1)^2}{x^2} + \frac{(1 + i\sqrt{3})(\sqrt{-x^2 + 1} - 1)}{x}\right)}{2}$
parts	$x \arctan\left(x + \sqrt{-x^2 + 1}\right) - \frac{\ln(x^4 - x^2 + 1)}{8} - \frac{\arctan\left(\frac{(2x^2 - 1)\sqrt{3}}{3}\right)\sqrt{3}}{4} - \frac{\left(\frac{i\sqrt{3}}{12} + \frac{1}{4}\right) \ln\left(\frac{(\sqrt{-x^2 + 1} - 1)^2}{x^2} + \frac{(1 + i\sqrt{3})(\sqrt{-x^2 + 1} - 1)}{x}\right)}{2}$

[In] int(arctan(x+(-x^2+1)^(1/2)),x,method=_RETURNVERBOSE)

[Out] x*arctan(x+(-x^2+1)^(1/2))-1/8*ln(x^4-x^2+1)-1/4*arctan(1/3*(2*x^2-1)*3^(1/2))*3^(1/2)-1/2*(1/12*I*3^(1/2)+1/4)*ln(((x^2+1)^(1/2)-1)^2/x^2+(1+I*3^(1/2)/2))*((x^2+1)^(1/2)-1)/x-1/2*(1/4-1/12*I*3^(1/2))*ln(((x^2+1)^(1/2)-1)^2/x^2+(1-I*3^(1/2)/2))*((x^2+1)^(1/2)-1)/x+1/2*(1/12*I*3^(1/2)+1/4)*ln(((x^2+1)^(1/2)-1)^2/x^2+(-1-I*3^(1/2))*((x^2+1)^(1/2)-1)/x-1)-1/12*I*3^(1/2)*ln(((x^2+1)^(1/2)-1)^2/x^2+(1+I*3^(1/2))*((x^2+1)^(1/2)-1)/x)+1/12*I*3^(1/2)*ln(((x^2+1)^(1/2)-1)^2/x^2+(1-I*3^(1/2))*((x^2+1)^(1/2)-1)/x)+arctan(((x^2+1)^(1/2)-1)/x)-1/12*I*3^(1/2)*ln(((x^2+1)^(1/2)-1)/x)

$2)-1)^{1/2}/x^2+(-1+I*3^{(1/2)})*((-x^2+1)^{(1/2)}-1)/x-1)+1/12*I*3^{(1/2)}*\ln(((x^2+1)^{(1/2)}-1)^{1/2}/x^2+(-1-I*3^{(1/2)})*((-x^2+1)^{(1/2)}-1)/x-1)$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.35

$$\begin{aligned} \int \arctan(x + \sqrt{1-x^2}) dx &= x \arctan(x + \sqrt{-x^2+1}) - \frac{1}{4} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^2-1)\right) \\ &\quad - \frac{1}{8} \sqrt{3} \arctan\left(\frac{4\sqrt{3}\sqrt{-x^2+1}x + \sqrt{3}}{3(2x^2-1)}\right) \\ &\quad - \frac{1}{8} \sqrt{3} \arctan\left(\frac{4\sqrt{3}\sqrt{-x^2+1}x - \sqrt{3}}{3(2x^2-1)}\right) \\ &\quad + \frac{1}{2} \arctan\left(\frac{\sqrt{-x^2+1}x}{x^2-1}\right) - \frac{1}{8} \log(x^4 - x^2 + 1) \\ &\quad - \frac{1}{16} \log(-x^4 + x^2 + 2\sqrt{-x^2+1}x + 1) \\ &\quad + \frac{1}{16} \log(-x^4 + x^2 - 2\sqrt{-x^2+1}x + 1) \end{aligned}$$

[In] integrate(arctan(x+(-x^2+1)^(1/2)),x, algorithm="fricas")

[Out] x*arctan(x + sqrt(-x^2 + 1)) - 1/4*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 - 1)) - 1/8*sqrt(3)*arctan(1/3*(4*sqrt(3)*sqrt(-x^2 + 1)*x + sqrt(3))/(2*x^2 - 1)) - 1/8*sqrt(3)*arctan(1/3*(4*sqrt(3)*sqrt(-x^2 + 1)*x - sqrt(3))/(2*x^2 - 1)) + 1/2*arctan(sqrt(-x^2 + 1)*x/(x^2 - 1)) - 1/8*log(x^4 - x^2 + 1) - 1/16*log(-x^4 + x^2 + 2*sqrt(-x^2 + 1)*x + 1) + 1/16*log(-x^4 + x^2 - 2*sqrt(-x^2 + 1)*x + 1)

Sympy [F(-1)]

Timed out.

$$\int \arctan(x + \sqrt{1-x^2}) dx = \text{Timed out}$$

[In] integrate(atan(x+(-x**2+1)**(1/2)),x)

[Out] Timed out

Maxima [F]

$$\int \arctan(x + \sqrt{1-x^2}) dx = \int \arctan(x + \sqrt{-x^2+1}) dx$$

[In] integrate(arctan(x+(-x^2+1)^(1/2)),x, algorithm="maxima")

[Out] x*arctan(x + sqrt(x + 1)*sqrt(-x + 1)) - integrate((x^3 + x^2*e^(1/2*log(x + 1) + 1/2*log(-x + 1)) - x)/(x^4 + (x^2 - 1)*e^(log(x + 1) + log(-x + 1)) + 2*(x^3 - x)*e^(1/2*log(x + 1) + 1/2*log(-x + 1)) - 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(110) = 220.

Time = 0.29 (sec) , antiderivative size = 364, normalized size of antiderivative = 2.58

$$\begin{aligned} & \int \arctan(x + \sqrt{1-x^2}) dx \\ &= x \arctan(x + \sqrt{-x^2+1}) - \frac{1}{4} \pi \operatorname{sgn}(x) \\ &+ \frac{1}{8} \sqrt{3} \left(\pi \operatorname{sgn}(x) + 2 \arctan \left(-\frac{\sqrt{3}x \left(\frac{\sqrt{-x^2+1}-1}{x} + \frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{3(\sqrt{-x^2+1}-1)} \right) \right) \\ &+ \frac{1}{8} \sqrt{3} \left(\pi \operatorname{sgn}(x) + 2 \arctan \left(\frac{\sqrt{3}x \left(\frac{\sqrt{-x^2+1}-1}{x} - \frac{(\sqrt{-x^2+1}-1)^2}{x^2} + 1 \right)}{3(\sqrt{-x^2+1}-1)} \right) \right) \\ &- \frac{1}{4} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3}(2x^2-1) \right) - \frac{1}{2} \arctan \left(-\frac{x \left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{2(\sqrt{-x^2+1}-1)} \right) \\ &- \frac{1}{8} \log(x^4 - x^2 + 1) + \frac{1}{8} \log \left(\left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right)^2 + \frac{2x}{\sqrt{-x^2+1}-1} \right. \\ &\quad \left. - \frac{2(\sqrt{-x^2+1}-1)}{x} + 4 \right) - \frac{1}{8} \log \left(\left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right)^2 \right. \\ &\quad \left. - \frac{2x}{\sqrt{-x^2+1}-1} + \frac{2(\sqrt{-x^2+1}-1)}{x} + 4 \right) \end{aligned}$$

[In] integrate(arctan(x+(-x^2+1)^(1/2)),x, algorithm="giac")

[Out] $x \arctan(x + \sqrt{-x^2 + 1}) - \frac{1}{4} \pi \operatorname{sgn}(x) + \frac{1}{8} \sqrt{3} (\pi \operatorname{sgn}(x) + 2 \arctan(-\frac{1}{3} \sqrt{3} x ((\sqrt{-x^2 + 1} - 1)/x + (\sqrt{-x^2 + 1} - 1)^2/x^2 - 1)/(\sqrt{-x^2 + 1} - 1))) + \frac{1}{8} \sqrt{3} (\pi \operatorname{sgn}(x) + 2 \arctan(\frac{1}{3} \sqrt{3} x ((\sqrt{-x^2 + 1} - 1)/x - (\sqrt{-x^2 + 1} - 1)^2/x^2 + 1)/(\sqrt{-x^2 + 1} - 1))) - \frac{1}{4} \sqrt{3} \arctan(\frac{1}{3} \sqrt{3} (2x^2 - 1)) - \frac{1}{2} \arctan(-\frac{1}{2} x ((\sqrt{-x^2 + 1} - 1)^2/x^2 - 1)/(\sqrt{-x^2 + 1} - 1)) - \frac{1}{8} \log(x^4 - x^2 + 1) + \frac{1}{8} \log((x/(\sqrt{-x^2 + 1} - 1) - (\sqrt{-x^2 + 1} - 1)/x)^2 + 2x/(\sqrt{-x^2 + 1} - 1) - 2(\sqrt{-x^2 + 1} - 1)/x + 4) - \frac{1}{8} \log((x/(\sqrt{-x^2 + 1} - 1) - (\sqrt{-x^2 + 1} - 1)/x)^2 - 2x/(\sqrt{-x^2 + 1} - 1) + 2(\sqrt{-x^2 + 1} - 1)/x + 4)$

Mupad [B] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 661, normalized size of antiderivative = 4.69

$$\int \arctan\left(x + \sqrt{1 - x^2}\right) dx = \text{Too large to display}$$

[In] int(atan(x + (1 - x^2)^(1/2)),x)

[Out] $x \operatorname{atan}(x + (1 - x^2)^{1/2}) - \operatorname{asin}(x)/2 + (\log(x - 3^{1/2}/2 - 1i/2) * (3^{1/2}/2 + (3^{1/2}/2 + 1i/2)^3 + 1i/2)) / (2 * 3^{1/2} - 8 * (3^{1/2}/2 + 1i/2)^3 + 2i) - (\log(x - 3^{1/2}/2 + 1i/2) * (3^{1/2}/2 + (3^{1/2}/2 - 1i/2)^3 - 1i/2)) / (8 * (3^{1/2}/2 - 1i/2)^3 - 2 * 3^{1/2} + 2i) - (\log(x + 3^{1/2}/2 - 1i/2) * (3^{1/2}/2 + (3^{1/2}/2 - 1i/2)^3 - 1i/2)) / (8 * (3^{1/2}/2 - 1i/2)^3 - 2 * 3^{1/2} + 2i) + (\log(x + 3^{1/2}/2 + 1i/2) * (3^{1/2}/2 + (3^{1/2}/2 + 1i/2)^3 + 1i/2)) / (2 * 3^{1/2} - 8 * (3^{1/2}/2 + 1i/2)^3 + 2i) + (\log(((x * (3^{1/2}/2 + 1i/2) - 1) * 1i) / (1 - (3^{1/2}/2 + 1i/2)^2)^{1/2}) - (1 - x^2)^{1/2} * 1i) / (3^{1/2}/2 - x + 1i/2) * ((3^{1/2}/2 + 1i/2)^2 + 1)) / ((1 - (3^{1/2}/2 + 1i/2)^2)^{1/2}) * (2 * 3^{1/2} - 8 * (3^{1/2}/2 + 1i/2)^3 + 2i) - (\log(((x * (3^{1/2}/2 - 1i/2) - 1) * 1i) / (1 - (3^{1/2}/2 - 1i/2)^2)^{1/2}) - (1 - x^2)^{1/2} * 1i) / (x - 3^{1/2}/2 + 1i/2) * ((3^{1/2}/2 - 1i/2)^2 + 1)) / ((1 - (3^{1/2}/2 - 1i/2)^2)^{1/2}) * (8 * (3^{1/2}/2 - 1i/2)^3 - 2 * 3^{1/2} + 2i) + (\log(((x * (3^{1/2}/2 - 1i/2) + 1) * 1i) / (1 - (3^{1/2}/2 - 1i/2)^2)^{1/2}) + (1 - x^2)^{1/2} * 1i) / (x + 3^{1/2}/2 - 1i/2) * ((3^{1/2}/2 - 1i/2)^2 + 1)) / ((1 - (3^{1/2}/2 - 1i/2)^2)^{1/2}) * (8 * (3^{1/2}/2 - 1i/2)^3 - 2 * 3^{1/2} + 2i) - (\log(((x * (3^{1/2}/2 + 1i/2) + 1) * 1i) / (1 - (3^{1/2}/2 + 1i/2)^2)^{1/2}) + (1 - x^2)^{1/2} * 1i) / (x + 3^{1/2}/2 + 1i/2) * ((3^{1/2}/2 + 1i/2)^2 + 1)) / ((1 - (3^{1/2}/2 + 1i/2)^2)^{1/2}) * (2 * 3^{1/2} - 8 * (3^{1/2}/2 + 1i/2)^3 + 2i)$

3.13 $\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx$

Optimal result	116
Rubi [C] (verified)	116
Mathematica [C] (verified)	121
Maple [F]	123
Fricas [A] (verification not implemented)	123
Sympy [F(-1)]	124
Maxima [F]	124
Giac [B] (verification not implemented)	124
Mupad [F(-1)]	126

Optimal result

Integrand size = 27, antiderivative size = 152

$$\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = -\frac{\arcsin(x)}{2} + \frac{1}{4}\sqrt{3} \arctan\left(\frac{-1 + \sqrt{3}x}{\sqrt{1-x^2}}\right) + \frac{1}{4}\sqrt{3} \arctan\left(\frac{1 + \sqrt{3}x}{\sqrt{1-x^2}}\right) - \frac{1}{4}\sqrt{3} \arctan\left(\frac{-1 + 2x^2}{\sqrt{3}}\right) - \sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) + \frac{1}{4}\operatorname{arctanh}(x\sqrt{1-x^2}) + \frac{1}{8}\log(1-x^2+x^4)$$

[Out] -1/2*arcsin(x)+1/4*arctanh(x*(-x^2+1)^(1/2))+1/8*ln(x^4-x^2+1)-1/4*arctan(1/3*(2*x^2-1)*3^(1/2))*3^(1/2)+1/4*arctan((-1+x*3^(1/2))/(-x^2+1)^(1/2))*3^(1/2)+1/4*arctan((1+x*3^(1/2))/(-x^2+1)^(1/2))*3^(1/2)-arctan(x+(-x^2+1)^(1/2))*(-x^2+1)^(1/2)

Rubi [C] (verified)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.88, number of steps used = 32, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.593$, Rules

used = {267, 5315, 12, 6874, 1121, 632, 210, 1307, 222, 1188, 385, 211, 399, 1261, 648, 642}

$$\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = -\frac{\arcsin(x)}{2} + \frac{1}{4}\sqrt{3} \arctan\left(\frac{1-2x^2}{\sqrt{3}}\right) - \frac{1}{12}(-\sqrt{3} + 3i) \arctan\left(\frac{x}{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}\sqrt{1-x^2}}\right) + \frac{\arctan\left(\frac{x}{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}\sqrt{1-x^2}}\right)}{2\sqrt{3}} + \frac{1}{12}(\sqrt{3} + 3i) \arctan\left(\frac{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}x}{\sqrt{1-x^2}}\right) + \frac{\arctan\left(\frac{\sqrt{-\frac{-\sqrt{3}+i}{\sqrt{3}+i}}x}{\sqrt{1-x^2}}\right)}{2\sqrt{3}} - \sqrt{1-x^2} \arctan(\sqrt{1-x^2} + x) + \frac{1}{8} \log(x^4 - x^2 + 1)$$

[In] Int[(x*ArcTan[x + Sqrt[1 - x^2]])/Sqrt[1 - x^2],x]

[Out] -1/2*ArcSin[x] + (Sqrt[3]*ArcTan[(1 - 2*x^2)/Sqrt[3]])/4 + ArcTan[x/(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*Sqrt[1 - x^2])]/(2*Sqrt[3]) - ((3*I - Sqrt[3])*ArcTan[x/(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*Sqrt[1 - x^2])])/12 + ArcTan[(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*x)/Sqrt[1 - x^2]]/(2*Sqrt[3]) + ((3*I + Sqrt[3])*ArcTan[(Sqrt[-((I - Sqrt[3])/(I + Sqrt[3]))]*x)/Sqrt[1 - x^2]])/12 - Sqrt[1 - x^2]*ArcTan[x + Sqrt[1 - x^2]] + Log[1 - x^2 + x^4]/8

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 267

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 399

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 1121

Int[(x_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, p}, x]

Rule 1188

```
Int[((d_) + (e_)*(x_)^2)^(q_)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol]
  := With[{r = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/r), Int[(d + e*x^2)^q/(b - r + 2*c*x^2), x], x] - Dist[2*(c/r), Int[(d + e*x^2)^q/(b + r + 2*c*x^2), x], x]] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[q]
```

Rule 1261

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol]
  := Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]
```

Rule 1307

```
Int[(((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(q_))/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol]
  := Dist[e*(f^2/c), Int[(f*x)^(m - 2)*(d + e*x^2)^(q - 1), x], x] - Dist[f^2/c, Int[(f*x)^(m - 2)*(d + e*x^2)^(q - 1)*(Simp[a*e - (c*d - b*e)*x^2, x]/(a + b*x^2 + c*x^4)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b^2 - 4*a*c, 0] && !IntegerQ[q] && GtQ[q, 0] && GtQ[m, 1] && LeQ[m, 3]
```

Rule 5315

```
Int[((a_) + ArcTan[u_]*(b_))*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[a + b*ArcTan[u], w, x] - Dist[b, Int[SimplifyIntegrand[w*(D[u, x]/(1 + u^2)), x], x], x] /; InverseFunctionFreeQ[w, x] /; FreeQ[{a, b}, x] && InverseFunctionFreeQ[u, x] && !MatchQ[v, ((c_) + (d_)*x)^(m_) /; FreeQ[{c, d, m}, x]] && FalseQ[FunctionOfLinear[v*(a + b*ArcTan[u]), x]]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\sqrt{1-x^2} \arctan\left(x + \sqrt{1-x^2}\right) - \int \frac{x - \sqrt{1-x^2}}{2(1+x\sqrt{1-x^2})} dx \\ &= -\sqrt{1-x^2} \arctan\left(x + \sqrt{1-x^2}\right) - \frac{1}{2} \int \frac{x - \sqrt{1-x^2}}{1+x\sqrt{1-x^2}} dx \\ &= -\sqrt{1-x^2} \arctan\left(x + \sqrt{1-x^2}\right) - \frac{1}{2} \int \left(\frac{x}{1+x\sqrt{1-x^2}} - \frac{\sqrt{1-x^2}}{1+x\sqrt{1-x^2}} \right) dx \end{aligned}$$

$$\begin{aligned}
&= -\sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) - \frac{1}{2} \int \frac{x}{1+x\sqrt{1-x^2}} dx + \frac{1}{2} \int \frac{\sqrt{1-x^2}}{1+x\sqrt{1-x^2}} dx \\
&= -\sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) - \frac{1}{2} \int \left(\frac{x}{1-x^2+x^4} - \frac{x^2\sqrt{1-x^2}}{1-x^2+x^4} \right) dx \\
&\quad + \frac{1}{2} \int \left(\frac{\sqrt{1-x^2}}{1-x^2+x^4} - \frac{x(1-x^2)}{1-x^2+x^4} \right) dx \\
&= -\sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) - \frac{1}{2} \int \frac{x}{1-x^2+x^4} dx \\
&\quad + \frac{1}{2} \int \frac{\sqrt{1-x^2}}{1-x^2+x^4} dx + \frac{1}{2} \int \frac{x^2\sqrt{1-x^2}}{1-x^2+x^4} dx - \frac{1}{2} \int \frac{x(1-x^2)}{1-x^2+x^4} dx \\
&= -\sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) - \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, x^2 \right) \\
&\quad - \frac{1}{4} \text{Subst} \left(\int \frac{1-x}{1-x+x^2} dx, x, x^2 \right) - \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}} dx \\
&\quad + \frac{1}{2} \int \frac{1}{\sqrt{1-x^2}(1-x^2+x^4)} dx - \frac{i \int \frac{\sqrt{1-x^2}}{-1-i\sqrt{3}+2x^2} dx}{\sqrt{3}} + \frac{i \int \frac{\sqrt{1-x^2}}{-1+i\sqrt{3}+2x^2} dx}{\sqrt{3}} \\
&= -\frac{\arcsin(x)}{2} - \sqrt{1-x^2} \arctan(x + \sqrt{1-x^2}) - \frac{1}{8} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, x^2 \right) \\
&\quad + \frac{1}{8} \text{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, x^2 \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{-3-x^2} dx, x, -1+2x^2 \right) \\
&\quad - \frac{i \int \frac{1}{\sqrt{1-x^2}(-1-i\sqrt{3}+2x^2)} dx}{\sqrt{3}} + \frac{i \int \frac{1}{\sqrt{1-x^2}(-1+i\sqrt{3}+2x^2)} dx}{\sqrt{3}} \\
&\quad - \frac{1}{6} (3-i\sqrt{3}) \int \frac{1}{\sqrt{1-x^2}(-1+i\sqrt{3}+2x^2)} dx \\
&\quad - \frac{1}{6} (3+i\sqrt{3}) \int \frac{1}{\sqrt{1-x^2}(-1-i\sqrt{3}+2x^2)} dx
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\arcsin(x)}{2} + \frac{\arctan\left(\frac{1-2x^2}{\sqrt{3}}\right)}{2\sqrt{3}} - \sqrt{1-x^2} \arctan\left(x + \sqrt{1-x^2}\right) \\
&\quad + \frac{1}{8} \log(1-x^2+x^4) + \frac{1}{4} \text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2x^2\right) \\
&\quad + \frac{i \text{Subst}\left(\int \frac{1}{-1+i\sqrt{3}-(-1-i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right)}{\sqrt{3}} \\
&\quad - \frac{i \text{Subst}\left(\int \frac{1}{-1-i\sqrt{3}-(-1+i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right)}{\sqrt{3}} \\
&\quad - \frac{1}{6} (3-i\sqrt{3}) \text{Subst}\left(\int \frac{1}{-1+i\sqrt{3}-(-1-i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\
&\quad - \frac{1}{6} (3+i\sqrt{3}) \text{Subst}\left(\int \frac{1}{-1-i\sqrt{3}-(-1+i\sqrt{3})x^2} dx, x, \frac{x}{\sqrt{1-x^2}}\right) \\
&= -\frac{\arcsin(x)}{2} + \frac{1}{4} \sqrt{3} \arctan\left(\frac{1-2x^2}{\sqrt{3}}\right) + \frac{\arctan\left(\frac{x}{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}\sqrt{1-x^2}}}\right)}{2\sqrt{3}} \\
&\quad - \frac{1}{12} (3i-\sqrt{3}) \arctan\left(\frac{x}{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}\sqrt{1-x^2}}}\right) \\
&\quad + \frac{\arctan\left(\frac{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}x}}{\sqrt{1-x^2}}\right)}{2\sqrt{3}} + \frac{1}{12} (3i+\sqrt{3}) \arctan\left(\frac{\sqrt{-\frac{i-\sqrt{3}}{i+\sqrt{3}}x}}{\sqrt{1-x^2}}\right) \\
&\quad - \sqrt{1-x^2} \arctan\left(x + \sqrt{1-x^2}\right) + \frac{1}{8} \log(1-x^2+x^4)
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.09 (sec) , antiderivative size = 2175, normalized size of antiderivative = 14.31

$$\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = \text{Result too large to show}$$

[In] Integrate[(x*ArcTan[x + Sqrt[1 - x^2]])/Sqrt[1 - x^2], x]

[Out] (-24*ArcSin[x] - 48*Sqrt[1 - x^2]*ArcTan[x + Sqrt[1 - x^2]] + (2*(-3*I + Sqrt[3])*ArcTan[(3 - I*Sqrt[3] + (-3 - I*Sqrt[3])*x^4 + 2*x*(-6*I + 2*Sqrt[3]

$$\begin{aligned}
& - I\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) - 2x^3(6I + 2\sqrt{3} + I\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) - (2I)\sqrt{3}x^2(6 + \sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}))/ (I - \sqrt{3} + (6I)(I + \sqrt{3})x - 2(-15I + \sqrt{3})x^2 + 6(1 + (3I)\sqrt{3})x^3 + (11I + 3\sqrt{3})x^4))/\sqrt{((1 - I\sqrt{3})/6) - (2(-3I + \sqrt{3})\text{ArcTan}[(3 - I\sqrt{3} + (-3 - I\sqrt{3})x^4 + 2x^3(6I + 2\sqrt{3} + I\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) + x(12I - 4\sqrt{3} + (2I)\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) - (2I)\sqrt{3}x^2(6 + \sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2})])/ (I - \sqrt{3} + (6 - (6I)\sqrt{3})x - 2(-15I + \sqrt{3})x^2 + (-6 - (18I)\sqrt{3})x^3 + (11I + 3\sqrt{3})x^4))/\sqrt{((1 - I\sqrt{3})/6) - (2(3I + \sqrt{3})\text{ArcTan}[-3 - I\sqrt{3} + (3 - I\sqrt{3})x^4 + 2x^3(-6I + 2\sqrt{3} - I\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) - 2x(6I + 2\sqrt{3} + I\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) - (2I)\sqrt{3}x^2(6 + \sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2})])/ (-I - \sqrt{3} + (-6 - (6I)\sqrt{3})x - 2(15I + \sqrt{3})x^2 + 6(1 - (3I)\sqrt{3})x^3 + (-11I + 3\sqrt{3})x^4))/\sqrt{((1 + I\sqrt{3})/6) + (2(3I + \sqrt{3})\text{ArcTan}[-3 - I\sqrt{3} + (3 - I\sqrt{3})x^4 + 2x(6I + 2\sqrt{3} + I\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) + x^3(12I - 4\sqrt{3} + (2I)\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) - (2I)\sqrt{3}x^2(6 + \sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2})])/ (-I - \sqrt{3} + (6 + (6I)\sqrt{3})x - 2(15I + \sqrt{3})x^2 + (-6 + (18I)\sqrt{3})x^3 + (-11I + 3\sqrt{3})x^4))/\sqrt{((1 + I\sqrt{3})/6) + 2\sqrt{3}(-3I + \sqrt{3})\text{Log}[I + \sqrt{3} - (2I)x^2] + 2\sqrt{3}(3I + \sqrt{3})\text{Log}[-I + \sqrt{3} + (2I)x^2] + ((3 - I\sqrt{3})\text{Log}[16(1 + \sqrt{3})x + x^2]^2))/\sqrt{((1 + I\sqrt{3})/6) + ((3 + I\sqrt{3})\text{Log}[16(1 + \sqrt{3})x + x^2]^2))/\sqrt{((1 - I\sqrt{3})/6) - (I(-3I + \sqrt{3})\text{Log}[(4 - 4\sqrt{3})x + 4x^2]^2))/\sqrt{((1 - I\sqrt{3})/6) + (I(3I + \sqrt{3})\text{Log}[(4 - 4\sqrt{3})x + 4x^2]^2))/\sqrt{((1 + I\sqrt{3})/6) - (I(-3I + \sqrt{3})\text{Log}[3I + \sqrt{3} - (-I + \sqrt{3})x^4 + (2I)\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2} + (5I)x^2(2 + \sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) + x(3 + (5I)\sqrt{3} + (3I)\sqrt{6 - (6I)\sqrt{3}}\sqrt{1 - x^2}) + Ix^3(3I + 3\sqrt{3} + \sqrt{6 - (6I)\sqrt{3}}\sqrt{1 - x^2})])/ (I - I\sqrt{3})/6) + ((3 + I\sqrt{3})\text{Log}[3I + \sqrt{3} - (-I + \sqrt{3})x^4 + (2I)\sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2} + (5I)x^2(2 + \sqrt{2 - (2I)\sqrt{3}}\sqrt{1 - x^2}) + x^3(3 - (3I)\sqrt{3} - I\sqrt{6 - (6I)\sqrt{3}}\sqrt{1 - x^2}) - Ix(-3I + 5\sqrt{3} + 3\sqrt{6 - (6I)\sqrt{3}}\sqrt{1 - x^2})])/ (I - I\sqrt{3})/6) + (I(3I + \sqrt{3})\text{Log}[-3I + \sqrt{3} - (I + \sqrt{3})x^4 - (2I)\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2} - (5I)x^2(2 + \sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) + x(3 - (5I)\sqrt{3} - (3I)\sqrt{6 + (6I)\sqrt{3}}\sqrt{1 - x^2}) - Ix^3(-3I + 3\sqrt{3} + \sqrt{6 + (6I)\sqrt{3}}\sqrt{1 - x^2})])/ (I + I\sqrt{3})/6) + ((3 - I\sqrt{3})\text{Log}[-3I + \sqrt{3} - (I + \sqrt{3})x^4 - (2I)\sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2} - (5I)x^2(2 + \sqrt{2 + (2I)\sqrt{3}}\sqrt{1 - x^2}) + x^3(3 + (3I)\sqrt{3} + I\sqrt{6 + (6I)\sqrt{3}}\sqrt{1 - x^2}) + Ix(3I + 5\sqrt{3} + 3\sqrt{6 + (6I)\sqrt{3}}\sqrt{1 - x^2})])/ (I + I\sqrt{3})/6))/48
\end{aligned}$$

Maple [F]

$$\int \frac{x \arctan(x + \sqrt{-x^2 + 1})}{\sqrt{-x^2 + 1}} dx$$

[In] int(x*arctan(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x)

[Out] int(x*arctan(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.32

$$\begin{aligned} \int \frac{x \arctan(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = & -\frac{1}{4} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3}(2x^2 - 1)\right) \\ & - \sqrt{-x^2 + 1} \arctan(x + \sqrt{-x^2 + 1}) \\ & - \frac{1}{8} \sqrt{3} \arctan\left(\frac{4\sqrt{3}\sqrt{-x^2 + 1}x + \sqrt{3}}{3(2x^2 - 1)}\right) \\ & - \frac{1}{8} \sqrt{3} \arctan\left(\frac{4\sqrt{3}\sqrt{-x^2 + 1}x - \sqrt{3}}{3(2x^2 - 1)}\right) \\ & + \frac{1}{2} \arctan\left(\frac{\sqrt{-x^2 + 1}x}{x^2 - 1}\right) + \frac{1}{8} \log(x^4 - x^2 + 1) \\ & + \frac{1}{16} \log(-x^4 + x^2 + 2\sqrt{-x^2 + 1}x + 1) \\ & - \frac{1}{16} \log(-x^4 + x^2 - 2\sqrt{-x^2 + 1}x + 1) \end{aligned}$$

[In] integrate(x*arctan(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -1/4*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 - 1)) - sqrt(-x^2 + 1)*arctan(x + sqrt(-x^2 + 1)) - 1/8*sqrt(3)*arctan(1/3*(4*sqrt(3)*sqrt(-x^2 + 1)*x + sqrt(3)))/(2*x^2 - 1) - 1/8*sqrt(3)*arctan(1/3*(4*sqrt(3)*sqrt(-x^2 + 1)*x - sqrt(3)))/(2*x^2 - 1) + 1/2*arctan(sqrt(-x^2 + 1)*x/(x^2 - 1)) + 1/8*log(x^4 - x^2 + 1) + 1/16*log(-x^4 + x^2 + 2*sqrt(-x^2 + 1)*x + 1) - 1/16*log(-x^4 + x^2 - 2*sqrt(-x^2 + 1)*x + 1)

Sympy [F(-1)]

Timed out.

$$\int \frac{x \arctan(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \text{Timed out}$$

[In] integrate(x*atan(x+(-x**2+1)**(1/2)))/(-x**2+1)**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{x \arctan(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \int \frac{x \arctan(x + \sqrt{-x^2 + 1})}{\sqrt{-x^2 + 1}} dx$$

[In] integrate(x*arctan(x+(-x^2+1)^(1/2)))/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(x + 1)*sqrt(-x + 1)*arctan(x + sqrt(x + 1)*sqrt(-x + 1)) - integrate(x/(x^2 + 2*x*e^(1/2*log(x + 1) + 1/2*log(-x + 1)) + e^(log(x + 1) + log(-x + 1)) + 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. 2(119) = 238.

Time = 0.34 (sec) , antiderivative size = 373, normalized size of antiderivative = 2.45

$$\begin{aligned}
 & \int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx \\
 &= -\frac{1}{4} \pi \operatorname{sgn}(x) + \frac{1}{8} \sqrt{3} \left(\pi \operatorname{sgn}(x) + 2 \arctan \left(-\frac{\sqrt{3}x \left(\frac{\sqrt{-x^2+1}-1}{x} + \frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{3(\sqrt{-x^2+1}-1)} \right) \right) \\
 &+ \frac{1}{8} \sqrt{3} \left(\pi \operatorname{sgn}(x) + 2 \arctan \left(\frac{\sqrt{3}x \left(\frac{\sqrt{-x^2+1}-1}{x} - \frac{(\sqrt{-x^2+1}-1)^2}{x^2} + 1 \right)}{3(\sqrt{-x^2+1}-1)} \right) \right) \\
 &- \frac{1}{4} \sqrt{3} \arctan \left(\frac{1}{3} \sqrt{3}(2x^2 - 1) \right) - \sqrt{-x^2+1} \arctan(x + \sqrt{-x^2+1}) \\
 &- \frac{1}{2} \arctan \left(-\frac{x \left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{2(\sqrt{-x^2+1}-1)} \right) + \frac{1}{8} \log(x^4 - x^2 + 1) \\
 &- \frac{1}{8} \log \left(\left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right)^2 + \frac{2x}{\sqrt{-x^2+1}-1} - \frac{2(\sqrt{-x^2+1}-1)}{x} \right. \\
 &\quad \left. + 4 \right) + \frac{1}{8} \log \left(\left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right)^2 - \frac{2x}{\sqrt{-x^2+1}-1} \right. \\
 &\quad \left. + \frac{2(\sqrt{-x^2+1}-1)}{x} + 4 \right)
 \end{aligned}$$

[In] integrate(x*arctan(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] -1/4*pi*sgn(x) + 1/8*sqrt(3)*(pi*sgn(x) + 2*arctan(-1/3*sqrt(3)*x*((sqrt(-x^2 + 1) - 1)/x + (sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))) + 1/8*sqrt(3)*(pi*sgn(x) + 2*arctan(1/3*sqrt(3)*x*((sqrt(-x^2 + 1) - 1)/x - (sqrt(-x^2 + 1) - 1)^2/x^2 + 1)/(sqrt(-x^2 + 1) - 1))) - 1/4*sqrt(3)*arctan(1/3*sqrt(3)*(2*x^2 - 1)) - sqrt(-x^2 + 1)*arctan(x + sqrt(-x^2 + 1)) - 1/2*arctan(-1/2*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1)) + 1/8*log(x^4 - x^2 + 1) - 1/8*log((x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)^2 + 2*x/(sqrt(-x^2 + 1) - 1) - 2*(sqrt(-x^2 + 1) - 1)/x + 4) + 1/8*log((x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)^2 - 2*x/(sqrt(-x^2 + 1) - 1) + 2*(sqrt(-x^2 + 1) - 1)/x + 4)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \arctan(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = \int \frac{x \operatorname{atan}(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx$$

```
[In] int((x*atan(x + (1 - x^2)^(1/2)))/(1 - x^2)^(1/2), x)
```

```
[Out] int((x*atan(x + (1 - x^2)^(1/2)))/(1 - x^2)^(1/2), x)
```

3.14 $\int \frac{\arcsin(x)}{1+\sqrt{1-x^2}} dx$

Optimal result	127
Rubi [A] (verified)	127
Mathematica [A] (verified)	130
Maple [F]	130
Fricas [A] (verification not implemented)	130
Sympy [F]	130
Maxima [F]	131
Giac [A] (verification not implemented)	131
Mupad [F(-1)]	131

Optimal result

Integrand size = 18, antiderivative size = 45

$$\int \frac{\arcsin(x)}{1+\sqrt{1-x^2}} dx = -\frac{x \arcsin(x)}{1+\sqrt{1-x^2}} + \frac{\arcsin(x)^2}{2} - \log\left(1+\sqrt{1-x^2}\right)$$

[Out] $1/2*\arcsin(x)^2-\ln(1+(-x^2+1)^{(1/2)})-x*\arcsin(x)/(1+(-x^2+1)^{(1/2)})$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.13, number of steps used = 9, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {6874, 283, 222, 4875, 4723, 272, 65, 212, 4781, 29, 4737}

$$\int \frac{\arcsin(x)}{1+\sqrt{1-x^2}} dx = \frac{\sqrt{1-x^2} \arcsin(x)}{x} + \frac{\arcsin(x)^2}{2} - \frac{\arcsin(x)}{x} - \operatorname{arctanh}\left(\sqrt{1-x^2}\right) - \log(x)$$

[In] `Int[ArcSin[x]/(1 + Sqrt[1 - x^2]),x]`

[Out] `-(ArcSin[x]/x) + (Sqrt[1 - x^2]*ArcSin[x])/x + ArcSin[x]^2/2 - ArcTanh[Sqrt[1 - x^2]] - Log[x]`

Rule 29

`Int[(x_)^(-1), x_Symbol] :> Simp[Log[x], x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +`

$d(x^{p/b})^n, x, (a + b*x)^{(1/p)}, x] /; \text{FreeQ}\{a, b, c, d, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

$\text{Int}[(a_ + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

Rule 222

$\text{Int}[1/\text{Sqrt}[(a_ + (b_.)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{GtQ}[a, 0] \&\& \text{NegQ}[b]$

Rule 272

$\text{Int}[(x_)^{(m_)}*((a_ + (b_.)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p, x}], x, x^n], x] /; \text{FreeQ}\{a, b, m, n, p, x\} \&\& \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 283

$\text{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_.)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(c*x)^{(m + 1)*((a + b*x^n)^p/(c*(m + 1))), x] - \text{Dist}[b*n*(p/(c^n*(m + 1))), \text{Int}[(c*x)^{(m + n)*(a + b*x^n)^{(p - 1)}, x], x] /; \text{FreeQ}\{a, b, c, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& \text{LtQ}[m, -1] \&\& !\text{LtQ}[(m + n*p + n + 1)/n, 0] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 4723

$\text{Int}[(a_ + \text{ArcSin}[(c_)*(x_)]*(b_))^{(n_)}*((d_)*(x_)^{(m_)}), x_Symbol] \rightarrow \text{Simp}[(d*x)^{(m + 1)*((a + b*\text{ArcSin}[c*x])^n/(d*(m + 1))), x] - \text{Dist}[b*c*(n/(d*(m + 1))), \text{Int}[(d*x)^{(m + 1)*((a + b*\text{ArcSin}[c*x])^{(n - 1)}/\text{Sqrt}[1 - c^2*x^2]), x], x] /; \text{FreeQ}\{a, b, c, d, m, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{NeQ}[m, -1]$

Rule 4737

$\text{Int}[(a_ + \text{ArcSin}[(c_)*(x_)]*(b_))^{(n_)} / \text{Sqrt}[(d_ + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(1/(b*c*(n + 1)))*\text{Simp}[\text{Sqrt}[1 - c^2*x^2]/\text{Sqrt}[d + e*x^2]]*(a + b*\text{ArcSin}[c*x])^{(n + 1)}, x] /; \text{FreeQ}\{a, b, c, d, e, n, x\} \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{NeQ}[n, -1]$

Rule 4781

$\text{Int}[(a_ + \text{ArcSin}[(c_)*(x_)]*(b_))^{(n_)}*((f_)*(x_)^{(m_)}*\text{Sqrt}[(d_ + (e_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[(f*x)^{(m + 1)*\text{Sqrt}[d + e*x^2]}*(a + b*\text{ArcS$


```
in[c*x])^n/(f*(m + 1))), x] + (-Dist[b*c*(n/(f*(m + 1)))*Simp[Sqrt[d + e*x^2]/Sqrt[1 - c^2*x^2]], Int[(f*x)^(m + 1)*(a + b*ArcSin[c*x])^(n - 1), x], x] + Dist[(c^2/(f^2*(m + 1)))*Simp[Sqrt[d + e*x^2]/Sqrt[1 - c^2*x^2]], Int[(f*x)^(m + 2)*((a + b*ArcSin[c*x])^n/Sqrt[1 - c^2*x^2]), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[c^2*d + e, 0] && GtQ[n, 0] && LtQ[m, -1]
```

Rule 4875

```
Int[((a_.) + ArcSin[(c_.)*(x_.)]*(b_.))^ (n_.)*(Px_.)*((f_.) + (g_.)*((d_.) + (e_.)*(x_.)^2)^(p_.))^(m_.), x_Symbol] := With[{u = ExpandIntegrand[Px*(f + g*(d + e*x^2)^p]^m*(a + b*ArcSin[c*x])^n, x]}, Int[u, x] /; SumQ[u] /; FreeQ[{a, b, c, d, e, f, g}, x] && PolynomialQ[Px, x] && EqQ[c^2*d + e, 0] && IGtQ[p + 1/2, 0] && IntegersQ[m, n]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{\arcsin(x)}{x^2} - \frac{\sqrt{1-x^2} \arcsin(x)}{x^2} \right) dx \\
 &= \int \frac{\arcsin(x)}{x^2} dx - \int \frac{\sqrt{1-x^2} \arcsin(x)}{x^2} dx \\
 &= -\frac{\arcsin(x)}{x} + \frac{\sqrt{1-x^2} \arcsin(x)}{x} - \int \frac{1}{x} dx + \int \frac{1}{x\sqrt{1-x^2}} dx + \int \frac{\arcsin(x)}{\sqrt{1-x^2}} dx \\
 &= -\frac{\arcsin(x)}{x} + \frac{\sqrt{1-x^2} \arcsin(x)}{x} + \frac{\arcsin(x)^2}{2} - \log(x) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1-xx}} dx, x, x^2 \right) \\
 &= -\frac{\arcsin(x)}{x} + \frac{\sqrt{1-x^2} \arcsin(x)}{x} + \frac{\arcsin(x)^2}{2} - \log(x) - \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x^2} \right) \\
 &= -\frac{\arcsin(x)}{x} + \frac{\sqrt{1-x^2} \arcsin(x)}{x} + \frac{\arcsin(x)^2}{2} - \text{arctanh}(\sqrt{1-x^2}) - \log(x)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.98

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \frac{(-1 + \sqrt{1 - x^2}) \arcsin(x)}{x} + \frac{\arcsin(x)^2}{2} - \log(1 + \sqrt{1 - x^2})$$

[In] Integrate[ArcSin[x]/(1 + Sqrt[1 - x^2]),x]

[Out] ((-1 + Sqrt[1 - x^2])*ArcSin[x])/x + ArcSin[x]^2/2 - Log[1 + Sqrt[1 - x^2]]

Maple [F]

$$\int \frac{\arcsin(x)}{1 + \sqrt{-x^2 + 1}} dx$$

[In] int(arcsin(x)/(1+(-x^2+1)^(1/2)),x)

[Out] int(arcsin(x)/(1+(-x^2+1)^(1/2)),x)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.40

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \frac{x \arcsin(x)^2 - 2x \log(x) - x \log(\sqrt{-x^2 + 1} + 1) + x \log(\sqrt{-x^2 + 1} - 1) + 2\sqrt{-x^2 + 1} \arcsin(x) - 2 \arcsin(x)}{2x}$$

[In] integrate(arcsin(x)/(1+(-x^2+1)^(1/2)),x, algorithm="fricas")

[Out] 1/2*(x*arcsin(x)^2 - 2*x*log(x) - x*log(sqrt(-x^2 + 1) + 1) + x*log(sqrt(-x^2 + 1) - 1) + 2*sqrt(-x^2 + 1)*arcsin(x) - 2*arcsin(x))/x

Sympy [F]

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \int \frac{\operatorname{asin}(x)}{\sqrt{1 - x^2} + 1} dx$$

[In] integrate(asin(x)/(1+(-x**2+1)**(1/2)),x)

[Out] Integral(asin(x)/(sqrt(1 - x**2) + 1), x)

Maxima [F]

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \int \frac{\arcsin(x)}{\sqrt{-x^2 + 1} + 1} dx$$

[In] integrate(arcsin(x)/(1+(-x^2+1)^(1/2)),x, algorithm="maxima")

[Out] integrate(arcsin(x)/(sqrt(-x^2 + 1) + 1), x)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.27

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \frac{1}{2} \arcsin(x)^2 - \frac{x \arcsin(x)}{\sqrt{-x^2 + 1} + 1} - 2 \log(2) + \log(2\sqrt{-x^2 + 1} + 2) - 2 \log(\sqrt{-x^2 + 1} + 1)$$

[In] integrate(arcsin(x)/(1+(-x^2+1)^(1/2)),x, algorithm="giac")

[Out] 1/2*arcsin(x)^2 - x*arcsin(x)/(sqrt(-x^2 + 1) + 1) - 2*log(2) + log(2*sqrt(-x^2 + 1) + 2) - 2*log(sqrt(-x^2 + 1) + 1)

Mupad [F(-1)]

Timed out.

$$\int \frac{\arcsin(x)}{1 + \sqrt{1 - x^2}} dx = \int \frac{\asin(x)}{\sqrt{1 - x^2} + 1} dx$$

[In] int(asin(x)/((1 - x^2)^(1/2) + 1),x)

[Out] int(asin(x)/((1 - x^2)^(1/2) + 1), x)

$$3.15 \quad \int \frac{\log(x + \sqrt{1+x^2})}{(1-x^2)^{3/2}} dx$$

Optimal result	132
Rubi [A] (verified)	132
Mathematica [B] (verified)	133
Maple [F]	134
Fricas [B] (verification not implemented)	134
Sympy [F(-1)]	134
Maxima [F]	134
Giac [A] (verification not implemented)	135
Mupad [F(-1)]	135

Optimal result

Integrand size = 24, antiderivative size = 34

$$\int \frac{\log(x + \sqrt{1+x^2})}{(1-x^2)^{3/2}} dx = -\frac{1}{2} \arcsin(x^2) + \frac{x \log(x + \sqrt{1+x^2})}{\sqrt{1-x^2}}$$

[Out] $-1/2*\arcsin(x^2)+x*\ln(x+(x^2+1)^{(1/2))}/(-x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {197, 2634, 281, 222}

$$\int \frac{\log(x + \sqrt{1+x^2})}{(1-x^2)^{3/2}} dx = \frac{x \log(\sqrt{x^2+1} + x)}{\sqrt{1-x^2}} - \frac{\arcsin(x^2)}{2}$$

[In] `Int[Log[x + Sqrt[1 + x^2]]/(1 - x^2)^(3/2),x]`

[Out] $-1/2*\text{ArcSin}[x^2] + (x*\text{Log}[x + \text{Sqrt}[1 + x^2]])/\text{Sqrt}[1 - x^2]$

Rule 197

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1) / a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 222

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 - x^2}} - \int \frac{x}{\sqrt{1 - x^4}} dx \\ &= \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 - x^2}} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1 - x^2}} dx, x, x^2\right) \\ &= -\frac{1}{2} \arcsin(x^2) + \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 - x^2}} \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 76 vs. 2(34) = 68.

Time = 0.07 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.24

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = \sqrt{1 - x^2} \left(\frac{\sqrt{1 + x^2} \arctan\left(\frac{\sqrt{1 - x^4}}{1 + x^2}\right)}{\sqrt{1 - x^4}} - \frac{x \log(x + \sqrt{1 + x^2})}{-1 + x^2} \right)$$

```
[In] Integrate[Log[x + Sqrt[1 + x^2]]/(1 - x^2)^(3/2), x]
```

```
[Out] Sqrt[1 - x^2]*((Sqrt[1 + x^2]*ArcTan[Sqrt[1 - x^4]/(1 + x^2)])/Sqrt[1 - x^4]
] - (x*Log[x + Sqrt[1 + x^2]])/(-1 + x^2)
```

Maple [F]

$$\int \frac{\ln(x + \sqrt{x^2 + 1})}{(-x^2 + 1)^{\frac{3}{2}}} dx$$

[In] int(ln(x+(x^2+1)^(1/2))/(-x^2+1)^(3/2),x)

[Out] int(ln(x+(x^2+1)^(1/2))/(-x^2+1)^(3/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(28) = 56.

Time = 0.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.82

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = -\frac{\sqrt{-x^2 + 1}x \log(x + \sqrt{x^2 + 1}) - (x^2 - 1) \arctan\left(\frac{\sqrt{x^2 + 1}\sqrt{-x^2 + 1} - 1}{x^2}\right)}{x^2 - 1}$$

[In] integrate(log(x+(x^2+1)^(1/2))/(-x^2+1)^(3/2),x, algorithm="fricas")

[Out] -(sqrt(-x^2 + 1)*x*log(x + sqrt(x^2 + 1)) - (x^2 - 1)*arctan((sqrt(x^2 + 1)*sqrt(-x^2 + 1) - 1)/x^2))/(x^2 - 1)

Sympy [F(-1)]

Timed out.

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = \text{Timed out}$$

[In] integrate(ln(x+(x**2+1)**(1/2))/(-x**2+1)**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = \int \frac{\log(x + \sqrt{x^2 + 1})}{(-x^2 + 1)^{\frac{3}{2}}} dx$$

[In] integrate(log(x+(x^2+1)^(1/2))/(-x^2+1)^(3/2),x, algorithm="maxima")

[Out] integrate(log(x + sqrt(x^2 + 1))/(-x^2 + 1)^(3/2), x)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = -\frac{\sqrt{-x^2 + 1} x \log(x + \sqrt{x^2 + 1})}{x^2 - 1} - \frac{1}{2} \arcsin(x^2)$$

[In] integrate(log(x+(x^2+1)^(1/2))/(-x^2+1)^(3/2),x, algorithm="giac")

[Out] -sqrt(-x^2 + 1)*x*log(x + sqrt(x^2 + 1))/(x^2 - 1) - 1/2*arcsin(x^2)

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x + \sqrt{1 + x^2})}{(1 - x^2)^{3/2}} dx = \int \frac{\ln(x + \sqrt{x^2 + 1})}{(1 - x^2)^{3/2}} dx$$

[In] int(log(x + (x^2 + 1)^(1/2))/(1 - x^2)^(3/2),x)

[Out] int(log(x + (x^2 + 1)^(1/2))/(1 - x^2)^(3/2), x)

3.16 $\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx$

Optimal result	136
Rubi [A] (verified)	136
Mathematica [A] (verified)	137
Maple [F]	137
Fricas [B] (verification not implemented)	138
Sympy [C] (verification not implemented)	138
Maxima [A] (verification not implemented)	138
Giac [A] (verification not implemented)	139
Mupad [F(-1)]	139

Optimal result

Integrand size = 12, antiderivative size = 22

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \arcsin(x)}{\sqrt{1+x^2}} - \frac{\arcsin(x^2)}{2}$$

[Out] $-1/2*\arcsin(x^2)+x*\arcsin(x)/(x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {197, 4755, 281, 222}

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \arcsin(x)}{\sqrt{x^2+1}} - \frac{\arcsin(x^2)}{2}$$

[In] `Int[ArcSin[x]/(1+x^2)^(3/2),x]`

[Out] `(x*ArcSin[x])/Sqrt[1+x^2] - ArcSin[x^2]/2`

Rule 197

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 222

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 4755

```
Int[((a_.) + ArcSin[(c_.)*(x_)])*(b_.)*((d_) + (e_.)*(x_)^2)^(p_.), x_Symbol] := With[{u = IntHide[(d + e*x^2)^p, x]}, Dist[a + b*ArcSin[c*x], u, x] - Dist[b*c, Int[SimplifyIntegrand[u/Sqrt[1 - c^2*x^2], x], x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d + e, 0] && (IGtQ[p, 0] || ILtQ[p + 1/2, 0])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x \arcsin(x)}{\sqrt{1+x^2}} - \int \frac{x}{\sqrt{1-x^2}} dx \\ &= \frac{x \arcsin(x)}{\sqrt{1+x^2}} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, x^2\right) \\ &= \frac{x \arcsin(x)}{\sqrt{1+x^2}} - \frac{\arcsin(x^2)}{2} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \arcsin(x)}{\sqrt{1+x^2}} - \frac{\arcsin(x^2)}{2}$$

```
[In] Integrate[ArcSin[x]/(1 + x^2)^(3/2), x]
```

```
[Out] (x*ArcSin[x])/Sqrt[1 + x^2] - ArcSin[x^2]/2
```

Maple [F]

$$\int \frac{\arcsin(x)}{(x^2+1)^{\frac{3}{2}}} dx$$

```
[In] int(arcsin(x)/(x^2+1)^(3/2), x)
```

```
[Out] int(arcsin(x)/(x^2+1)^(3/2), x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(18) = 36.
Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.55

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{2\sqrt{x^2+1}x \arcsin(x) + (x^2+1) \arctan\left(\frac{\sqrt{x^2+1}\sqrt{-x^2+1}x^2}{x^4-1}\right)}{2(x^2+1)}$$

[In] integrate(arcsin(x)/(x^2+1)^(3/2),x, algorithm="fricas")

[Out] 1/2*(2*sqrt(x^2 + 1)*x*arcsin(x) + (x^2 + 1)*arctan(sqrt(x^2 + 1)*sqrt(-x^2 + 1)*x^2/(x^4 - 1)))/(x^2 + 1)

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 20.25 (sec) , antiderivative size = 78, normalized size of antiderivative = 3.55

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \operatorname{asin}(x)}{\sqrt{x^2+1}} + \frac{i G_{6,6}^{6,2} \left(\begin{matrix} \frac{1}{4}, \frac{3}{4} & \frac{1}{2}, \frac{1}{2}, 1, 1 \\ 0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1, 0 \end{matrix} \middle| \frac{1}{x^4} \right)}{8\pi^{\frac{3}{2}}} - \frac{G_{6,6}^{2,6} \left(\begin{matrix} -\frac{1}{2}, -\frac{1}{4}, 0, \frac{1}{4}, \frac{1}{2}, 1 \\ -\frac{1}{4}, \frac{1}{4} & -\frac{1}{2}, 0, 0, 0 \end{matrix} \middle| \frac{e^{-2i\pi}}{x^4} \right)}{8\pi^{\frac{3}{2}}}$$

[In] integrate(asin(x)/(x**2+1)**(3/2),x)

[Out] x*asin(x)/sqrt(x**2 + 1) + I*meijerg(((1/4, 3/4), (1/2, 1/2, 1, 1)), ((0, 1/4, 1/2, 3/4, 1, 0), ()), x**(-4))/(8*pi**(3/2)) - meijerg((-1/2, -1/4, 0, 1/4, 1/2, 1), ()), ((-1/4, 1/4), (-1/2, 0, 0, 0)), exp_polar(-2*I*pi)/x**4)/(8*pi**(3/2))

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \arcsin(x)}{\sqrt{x^2+1}} - \frac{1}{2} \arcsin(x^2)$$

[In] integrate(arcsin(x)/(x^2+1)^(3/2),x, algorithm="maxima")

[Out] x*arcsin(x)/sqrt(x^2 + 1) - 1/2*arcsin(x^2)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \frac{x \arcsin(x)}{\sqrt{x^2+1}} - \frac{1}{2} \arcsin(x^2)$$

[In] integrate(arcsin(x)/(x^2+1)^(3/2),x, algorithm="giac")

[Out] x*arcsin(x)/sqrt(x^2 + 1) - 1/2*arcsin(x^2)

Mupad [F(-1)]

Timed out.

$$\int \frac{\arcsin(x)}{(1+x^2)^{3/2}} dx = \int \frac{\arcsin(x)}{(x^2+1)^{3/2}} dx$$

[In] int(asin(x)/(x^2 + 1)^(3/2),x)

[Out] int(asin(x)/(x^2 + 1)^(3/2), x)

$$3.17 \quad \int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx$$

Optimal result	140
Rubi [A] (verified)	140
Mathematica [B] (verified)	141
Maple [F]	142
Fricas [B] (verification not implemented)	142
Sympy [F]	142
Maxima [F]	142
Giac [A] (verification not implemented)	143
Mupad [F(-1)]	143

Optimal result

Integrand size = 22, antiderivative size = 32

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = -\frac{1}{2} \operatorname{arccosh}(x^2) + \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{1 + x^2}}$$

[Out] $-1/2*\operatorname{arccosh}(x^2)+x*\ln(x+(x^2-1)^{(1/2)})/(x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {197, 2634, 282, 54}

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \frac{x \log(\sqrt{x^2 - 1} + x)}{\sqrt{x^2 + 1}} - \frac{\operatorname{arccosh}(x^2)}{2}$$

[In] `Int[Log[x + Sqrt[-1 + x^2]]/(1 + x^2)^(3/2),x]`

[Out] $-1/2*\operatorname{ArcCosh}[x^2] + (x*\operatorname{Log}[x + \operatorname{Sqrt}[-1 + x^2]])/\operatorname{Sqrt}[1 + x^2]$

Rule 54

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_) + (d_.)*(x_)]), x_Symbol] :> Simp[
ArcCosh[b*(x/a)]/b, x] /; FreeQ[{a, b, c, d}, x] && EqQ[a + c, 0] && EqQ[b
- d, 0] && GtQ[a, 0]
```

Rule 197

```
Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)
/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]
```

Rule 282

```
Int[(x_)^(m_)*((a1_) + (b1_)*(x_)^(n_))^(p_)*((a2_) + (b2_)*(x_)^(n_))^(
p_), x_Symbol] := With[{k = GCD[m + 1, 2*n]}, Dist[1/k, Subst[Int[x^((m + 1)
)/k - 1)*(a1 + b1*x^(n/k))^p*(a2 + b2*x^(n/k))^p, x], x, x^k], x] /; k != 1
] /; FreeQ[{a1, b1, a2, b2, p}, x] && EqQ[a2*b1 + a1*b2, 0] && IGtQ[2*n, 0]
&& IntegerQ[m]
```

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{1 + x^2}} - \int \frac{x}{\sqrt{-1 + x^2} \sqrt{1 + x^2}} dx \\ &= \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{1 + x^2}} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{-1 + x} \sqrt{1 + x}} dx, x, x^2\right) \\ &= -\frac{1}{2} \text{arccosh}(x^2) + \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{1 + x^2}} \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 89 vs. 2(32) = 64.

Time = 0.05 (sec) , antiderivative size = 89, normalized size of antiderivative = 2.78

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \frac{4x \log(x + \sqrt{-1 + x^2}) + \frac{\sqrt{-1 + x^2}(1 + x^2) \left(\log\left(1 - \frac{x^2}{\sqrt{-1 + x^4}}\right) - \log\left(1 + \frac{x^2}{\sqrt{-1 + x^4}}\right) \right)}{\sqrt{-1 + x^4}}}{4\sqrt{1 + x^2}}$$

```
[In] Integrate[Log[x + Sqrt[-1 + x^2]]/(1 + x^2)^(3/2), x]
```

```
[Out] (4*x*Log[x + Sqrt[-1 + x^2]] + (Sqrt[-1 + x^2]*(1 + x^2)*(Log[1 - x^2/Sqrt[
-1 + x^4]] - Log[1 + x^2/Sqrt[-1 + x^4]]))/Sqrt[-1 + x^4])/(4*Sqrt[1 + x^2]
)
```

Maple [F]

$$\int \frac{\ln(x + \sqrt{x^2 - 1})}{(x^2 + 1)^{\frac{3}{2}}} dx$$

[In] `int(ln(x+(x^2-1)^(1/2))/(x^2+1)^(3/2),x)`

[Out] `int(ln(x+(x^2-1)^(1/2))/(x^2+1)^(3/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 58 vs. 2(26) = 52.

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.81

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \frac{2\sqrt{x^2 + 1}x \log(x + \sqrt{x^2 - 1}) + (x^2 + 1) \log(-x^2 + \sqrt{x^2 + 1}\sqrt{x^2 - 1})}{2(x^2 + 1)}$$

[In] `integrate(log(x+(x^2-1)^(1/2))/(x^2+1)^(3/2),x, algorithm="fricas")`

[Out] `1/2*(2*sqrt(x^2 + 1)*x*log(x + sqrt(x^2 - 1)) + (x^2 + 1)*log(-x^2 + sqrt(x^2 + 1)*sqrt(x^2 - 1)))/(x^2 + 1)`

Sympy [F]

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \int \frac{\log(x + \sqrt{x^2 - 1})}{(x^2 + 1)^{\frac{3}{2}}} dx$$

[In] `integrate(ln(x+(x**2-1)**(1/2))/(x**2+1)**(3/2),x)`

[Out] `Integral(log(x + sqrt(x**2 - 1))/(x**2 + 1)**(3/2), x)`

Maxima [F]

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \int \frac{\log(x + \sqrt{x^2 - 1})}{(x^2 + 1)^{\frac{3}{2}}} dx$$

[In] `integrate(log(x+(x^2-1)^(1/2))/(x^2+1)^(3/2),x, algorithm="maxima")`

[Out] `integrate(log(x + sqrt(x^2 - 1))/(x^2 + 1)^(3/2), x)`

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.12

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \frac{x \log(x + \sqrt{x^2 - 1})}{\sqrt{x^2 + 1}} + \frac{1}{2} \log(x^2 - \sqrt{x^4 - 1})$$

[In] integrate(log(x+(x^2-1)^(1/2))/(x^2+1)^(3/2),x, algorithm="giac")

[Out] x*log(x + sqrt(x^2 - 1))/sqrt(x^2 + 1) + 1/2*log(x^2 - sqrt(x^4 - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x + \sqrt{-1 + x^2})}{(1 + x^2)^{3/2}} dx = \int \frac{\ln(x + \sqrt{x^2 - 1})}{(x^2 + 1)^{3/2}} dx$$

[In] int(log(x + (x^2 - 1)^(1/2))/(x^2 + 1)^(3/2),x)

[Out] int(log(x + (x^2 - 1)^(1/2))/(x^2 + 1)^(3/2), x)

3.18 $\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx$

Optimal result	144
Rubi [A] (verified)	144
Mathematica [A] (verified)	145
Maple [C] (warning: unable to verify)	146
Fricas [A] (verification not implemented)	146
Sympy [A] (verification not implemented)	146
Maxima [A] (verification not implemented)	147
Giac [A] (verification not implemented)	147
Mupad [F(-1)]	147

Optimal result

Integrand size = 15, antiderivative size = 43

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \frac{\sqrt{-1+x^2}}{x} - \operatorname{arctanh}\left(\frac{x}{\sqrt{-1+x^2}}\right) + \frac{\sqrt{-1+x^2}\log(x)}{x}$$

[Out] $-\operatorname{arctanh}(x/(x^2-1)^{(1/2)})+(x^2-1)^{(1/2)}/x+\ln(x)*(x^2-1)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2373, 283, 223, 212}

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = -\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2-1}}\right) + \frac{\sqrt{x^2-1}}{x} + \frac{\sqrt{x^2-1}\log(x)}{x}$$

[In] $\operatorname{Int}[\operatorname{Log}[x]/(x^2*\operatorname{Sqrt}[-1+x^2]),x]$

[Out] $\operatorname{Sqrt}[-1+x^2]/x - \operatorname{ArcTanh}[x/\operatorname{Sqrt}[-1+x^2]] + (\operatorname{Sqrt}[-1+x^2]*\operatorname{Log}[x])/x$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] := \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+ + (b_+)(x_+)^2], x_Symbol] := \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 283

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^p/(c^(n*(m + 1))), x] - Dist[b*n*(p/(c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m + n*p + n + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2373

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*(x_)^(r_.))^(q_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^r)^(q + 1)*((a + b*Log[c*x^n])/(d*(m + 1))), x] - Dist[b*(n/(d*(m + 1))), Int[(f*x)^(m*(d + e*x^r)^(q + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, n, q, r}, x] && EqQ[m + r*(q + 1) + 1, 0] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\sqrt{-1+x^2} \log(x)}{x} - \int \frac{\sqrt{-1+x^2}}{x^2} dx \\
&= \frac{\sqrt{-1+x^2}}{x} + \frac{\sqrt{-1+x^2} \log(x)}{x} - \int \frac{1}{\sqrt{-1+x^2}} dx \\
&= \frac{\sqrt{-1+x^2}}{x} + \frac{\sqrt{-1+x^2} \log(x)}{x} - \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{-1+x^2}}\right) \\
&= \frac{\sqrt{-1+x^2}}{x} - \operatorname{arctanh}\left(\frac{x}{\sqrt{-1+x^2}}\right) + \frac{\sqrt{-1+x^2} \log(x)}{x}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2 \sqrt{-1+x^2}} dx = \frac{\sqrt{-1+x^2}}{x} + \frac{\sqrt{-1+x^2} \log(x)}{x} - \log\left(x + \sqrt{-1+x^2}\right)$$

[In] Integrate[Log[x]/(x^2*Sqrt[-1 + x^2]),x]

[Out] Sqrt[-1 + x^2]/x + (Sqrt[-1 + x^2]*Log[x])/x - Log[x + Sqrt[-1 + x^2]]

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 2.07

method	result	size
meijerg	$-\frac{\sqrt{-\operatorname{signum}(x^2-1)} \arcsin(x)}{\sqrt{\operatorname{signum}(x^2-1)}} + \frac{-\frac{\sqrt{-\operatorname{signum}(x^2-1)} \sqrt{-x^2+1}}{\sqrt{\operatorname{signum}(x^2-1)}} - \frac{\sqrt{-\operatorname{signum}(x^2-1)} \ln(x) \sqrt{-x^2+1}}{\sqrt{\operatorname{signum}(x^2-1)}}}{x}$	89

[In] `int(ln(x)/x^2/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*arcsin(x)+(-1/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*(-x^2+1)^(1/2)-1/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*ln(x)*(-x^2+1)^(1/2))/x`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \frac{x \log(-x + \sqrt{x^2-1}) + \sqrt{x^2-1}(\log(x) + 1) + x}{x}$$

[In] `integrate(log(x)/x^2/(x^2-1)^(1/2),x, algorithm="fricas")`

[Out] `(x*log(-x + sqrt(x^2 - 1)) + sqrt(x^2 - 1)*(log(x) + 1) + x)/x`

Sympy [A] (verification not implemented)

Time = 2.69 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.98

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \begin{cases} \frac{\sqrt{x^2-1}}{x} & \text{for } x > -1 \wedge x < 1 \\ \text{NaN} & \text{for } x < -1 \\ \log(x + \sqrt{x^2-1}) - \frac{\sqrt{x^2-1}}{x} & \text{for } x < 1 \\ \text{NaN} & \text{otherwise} \end{cases}$$

[In] `integrate(ln(x)/x**2/(x**2-1)**(1/2),x)`

[Out] `Piecewise((sqrt(x**2 - 1)/x, (x > -1) & (x < 1)))*log(x) - Piecewise((nan, x < -1), (log(x + sqrt(x**2 - 1)) - sqrt(x**2 - 1)/x, x < 1), (nan, True))`

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \frac{\sqrt{x^2-1} \log(x)}{x} + \frac{\sqrt{x^2-1}}{x} - \log\left(2x + 2\sqrt{x^2-1}\right)$$

[In] integrate(log(x)/x^2/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 - 1)*log(x)/x + sqrt(x^2 - 1)/x - log(2*x + 2*sqrt(x^2 - 1))

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \frac{2 \log(x)}{(x - \sqrt{x^2-1})^2 + 1} + \frac{2}{(x - \sqrt{x^2-1})^2 + 1} + \frac{1}{2} \log\left(\left(x - \sqrt{x^2-1}\right)^2\right) - \log(|x|)$$

[In] integrate(log(x)/x^2/(x^2-1)^(1/2),x, algorithm="giac")

[Out] 2*log(x)/((x - sqrt(x^2 - 1))^2 + 1) + 2/((x - sqrt(x^2 - 1))^2 + 1) + 1/2*log((x - sqrt(x^2 - 1))^2) - log(abs(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x)}{x^2\sqrt{-1+x^2}} dx = \int \frac{\ln(x)}{x^2\sqrt{x^2-1}} dx$$

[In] int(log(x)/(x^2*(x^2 - 1)^(1/2)),x)

[Out] int(log(x)/(x^2*(x^2 - 1)^(1/2)), x)

3.19 $\int \frac{\sqrt{1+x^3}}{x} dx$

Optimal result	148
Rubi [A] (verified)	148
Mathematica [A] (verified)	149
Maple [A] (verified)	150
Fricas [A] (verification not implemented)	150
Sympy [A] (verification not implemented)	150
Maxima [A] (verification not implemented)	151
Giac [A] (verification not implemented)	151
Mupad [B] (verification not implemented)	151

Optimal result

Integrand size = 13, antiderivative size = 28

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2\sqrt{1+x^3}}{3} - \frac{2}{3} \operatorname{arctanh}(\sqrt{1+x^3})$$

[Out] $-2/3*\operatorname{arctanh}((x^3+1)^{(1/2)})+2/3*(x^3+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {272, 52, 65, 213}

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2\sqrt{x^3+1}}{3} - \frac{2}{3} \operatorname{arctanh}(\sqrt{x^3+1})$$

[In] `Int[Sqrt[1 + x^3]/x, x]`

[Out] $(2*\operatorname{Sqrt}[1 + x^3])/3 - (2*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 + x^3]])/3$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 213

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3} \text{Subst} \left(\int \frac{\sqrt{1+x}}{x} dx, x, x^3 \right) \\
&= \frac{2\sqrt{1+x^3}}{3} + \frac{1}{3} \text{Subst} \left(\int \frac{1}{x\sqrt{1+x}} dx, x, x^3 \right) \\
&= \frac{2\sqrt{1+x^3}}{3} + \frac{2}{3} \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+x^3} \right) \\
&= \frac{2\sqrt{1+x^3}}{3} - \frac{2}{3} \text{arctanh}(\sqrt{1+x^3})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2\sqrt{1+x^3}}{3} - \frac{2}{3} \text{arctanh}(\sqrt{1+x^3})$$

```
[In] Integrate[Sqrt[1 + x^3]/x,x]
```

```
[Out] (2*Sqrt[1 + x^3])/3 - (2*ArcTanh[Sqrt[1 + x^3]])/3
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.75

method	result	size
default	$-\frac{2 \operatorname{arctanh}(\sqrt{x^3+1})}{3} + \frac{2\sqrt{x^3+1}}{3}$	21
elliptic	$-\frac{2 \operatorname{arctanh}(\sqrt{x^3+1})}{3} + \frac{2\sqrt{x^3+1}}{3}$	21
trager	$\frac{2\sqrt{x^3+1}}{3} - \frac{\ln\left(-\frac{x^3+2\sqrt{x^3+1}+2}{x^3}\right)}{3}$	33
pseudoelliptic	$\frac{2\sqrt{x^3+1}}{3} + \frac{\ln(\sqrt{x^3+1}-1)}{3} - \frac{\ln(1+\sqrt{x^3+1})}{3}$	35
meijerg	$-\frac{-2(2-2\ln(2)+3\ln(x))\sqrt{\pi}+4\sqrt{\pi}-4\sqrt{\pi}\sqrt{x^3+1}+4\sqrt{\pi}\ln\left(\frac{1}{2}+\frac{\sqrt{x^3+1}}{2}\right)}{6\sqrt{\pi}}$	56

```
[In] int((x^3+1)^(1/2)/x,x,method=_RETURNVERBOSE)
```

```
[Out] -2/3*arctanh((x^3+1)^(1/2))+2/3*(x^3+1)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2}{3} \sqrt{x^3+1} - \frac{1}{3} \log(\sqrt{x^3+1}+1) + \frac{1}{3} \log(\sqrt{x^3+1}-1)$$

```
[In] integrate((x^3+1)^(1/2)/x,x, algorithm="fricas")
```

```
[Out] 2/3*sqrt(x^3 + 1) - 1/3*log(sqrt(x^3 + 1) + 1) + 1/3*log(sqrt(x^3 + 1) - 1)
```

Sympy [A] (verification not implemented)

Time = 0.90 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.71

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2x^{\frac{3}{2}}}{3\sqrt{1+\frac{1}{x^3}}} - \frac{2 \operatorname{asinh}\left(\frac{1}{x^{\frac{3}{2}}}\right)}{3} + \frac{2}{3x^{\frac{3}{2}}\sqrt{1+\frac{1}{x^3}}}$$

```
[In] integrate((x**3+1)**(1/2)/x,x)
```

```
[Out] 2*x**(3/2)/(3*sqrt(1 + x**(-3))) - 2*asinh(x**(-3/2))/3 + 2/(3*x**(3/2)*sqrt(1 + x**(-3)))
```

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2}{3} \sqrt{x^3+1} - \frac{1}{3} \log(\sqrt{x^3+1}+1) + \frac{1}{3} \log(\sqrt{x^3+1}-1)$$

[In] integrate((x^3+1)^(1/2)/x,x, algorithm="maxima")

[Out] 2/3*sqrt(x^3 + 1) - 1/3*log(sqrt(x^3 + 1) + 1) + 1/3*log(sqrt(x^3 + 1) - 1)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.25

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2}{3} \sqrt{x^3+1} - \frac{1}{3} \log(\sqrt{x^3+1}+1) + \frac{1}{3} \log(|\sqrt{x^3+1}-1|)$$

[In] integrate((x^3+1)^(1/2)/x,x, algorithm="giac")

[Out] 2/3*sqrt(x^3 + 1) - 1/3*log(sqrt(x^3 + 1) + 1) + 1/3*log(abs(sqrt(x^3 + 1) - 1))

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 174, normalized size of antiderivative = 6.21

$$\int \frac{\sqrt{1+x^3}}{x} dx = \frac{2\sqrt{x^3+1}}{3} - \frac{2\left(\frac{3}{2} + \frac{\sqrt{3}1i}{2}\right) \sqrt{\frac{x-\frac{1}{2}+\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{x+1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \sqrt{\frac{\frac{1}{2}-x+\frac{\sqrt{3}1i}{2}}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}} \Pi\left(\frac{3}{2} + \frac{\sqrt{3}1i}{2}; \operatorname{asin}\left(\sqrt{\frac{x+1}{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}}\right) \middle| -\frac{\frac{3}{2}+\frac{\sqrt{3}1i}{2}}{-\frac{3}{2}+\frac{\sqrt{3}1i}{2}}\right)}{\sqrt{x^3 + \left(-\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - 1\right) x - \left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}}$$

[In] int((x^3 + 1)^(1/2)/x,x)

[Out] (2*(x^3 + 1)^(1/2))/3 - (2*((3^(1/2)*1i)/2 + 3/2)*((x + (3^(1/2)*1i)/2 - 1/2)/((3^(1/2)*1i)/2 - 3/2))^(1/2)*((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*(((3^(1/2)*1i)/2 - x + 1/2)/((3^(1/2)*1i)/2 + 3/2))^(1/2)*ellipticPi((3^(1/2)*1i)/2 + 3/2, asin(((x + 1)/((3^(1/2)*1i)/2 + 3/2))^(1/2)), -(3^(1/2)*1i)/2 + 3/2)/((3^(1/2)*1i)/2 - 3/2))/(x^3 - x*((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2) + 1) - ((3^(1/2)*1i)/2 - 1/2)*((3^(1/2)*1i)/2 + 1/2))^(1/2)

$$3.20 \quad \int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx$$

Optimal result	152
Rubi [A] (verified)	152
Mathematica [A] (verified)	153
Maple [F]	153
Fricas [A] (verification not implemented)	153
Sympy [A] (verification not implemented)	154
Maxima [A] (verification not implemented)	154
Giac [A] (verification not implemented)	154
Mupad [F(-1)]	154

Optimal result

Integrand size = 23, antiderivative size = 26

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = -x + \sqrt{-1 + x^2} \log(x + \sqrt{-1 + x^2})$$

[Out] -x+ln(x+(x^2-1)^(1/2))*(x^2-1)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {267, 2634, 8}

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = \sqrt{x^2 - 1} \log(\sqrt{x^2 - 1} + x) - x$$

[In] Int[(x*Log[x + Sqrt[-1 + x^2]])/Sqrt[-1 + x^2],x]

[Out] -x + Sqrt[-1 + x^2]*Log[x + Sqrt[-1 + x^2]]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 267

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \sqrt{-1+x^2} \log\left(x + \sqrt{-1+x^2}\right) - \int 1 dx \\ &= -x + \sqrt{-1+x^2} \log\left(x + \sqrt{-1+x^2}\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x \log\left(x + \sqrt{-1+x^2}\right)}{\sqrt{-1+x^2}} dx = -x + \sqrt{-1+x^2} \log\left(x + \sqrt{-1+x^2}\right)$$

```
[In] Integrate[(x*Log[x + Sqrt[-1 + x^2]])/Sqrt[-1 + x^2], x]
```

```
[Out] -x + Sqrt[-1 + x^2]*Log[x + Sqrt[-1 + x^2]]
```

Maple [F]

$$\int \frac{x \ln\left(x + \sqrt{x^2-1}\right)}{\sqrt{x^2-1}} dx$$

```
[In] int(x*ln(x+(x^2-1)^(1/2))/(x^2-1)^(1/2), x)
```

```
[Out] int(x*ln(x+(x^2-1)^(1/2))/(x^2-1)^(1/2), x)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log\left(x + \sqrt{-1+x^2}\right)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log\left(x + \sqrt{x^2-1}\right) - x$$

```
[In] integrate(x*log(x+(x^2-1)^(1/2))/(x^2-1)^(1/2), x, algorithm="fricas")
```

```
[Out] sqrt(x^2 - 1)*log(x + sqrt(x^2 - 1)) - x
```

Sympy [A] (verification not implemented)

Time = 3.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = -x + \sqrt{x^2 - 1} \log(x + \sqrt{x^2 - 1})$$

[In] integrate(x*ln(x+(x**2-1)**(1/2))/(x**2-1)**(1/2),x)

[Out] -x + sqrt(x**2 - 1)*log(x + sqrt(x**2 - 1))

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = \sqrt{x^2 - 1} \log(x + \sqrt{x^2 - 1}) - x$$

[In] integrate(x*log(x+(x^2-1)^(1/2))/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 - 1)*log(x + sqrt(x^2 - 1)) - x

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = \sqrt{x^2 - 1} \log(x + \sqrt{x^2 - 1}) - x$$

[In] integrate(x*log(x+(x^2-1)^(1/2))/(x^2-1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 - 1)*log(x + sqrt(x^2 - 1)) - x

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(x + \sqrt{-1 + x^2})}{\sqrt{-1 + x^2}} dx = \int \frac{x \ln(x + \sqrt{x^2 - 1})}{\sqrt{x^2 - 1}} dx$$

[In] int((x*log(x + (x^2 - 1)^(1/2)))/(x^2 - 1)^(1/2),x)

[Out] int((x*log(x + (x^2 - 1)^(1/2)))/(x^2 - 1)^(1/2), x)

3.21 $\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx$

Optimal result	155
Rubi [A] (verified)	155
Mathematica [B] (verified)	157
Maple [F]	157
Fricas [B] (verification not implemented)	157
Sympy [F]	158
Maxima [F]	158
Giac [A] (verification not implemented)	158
Mupad [F(-1)]	158

Optimal result

Integrand size = 17, antiderivative size = 38

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \frac{1}{4}x\sqrt{1+x^2} - \frac{1}{2}\sqrt{1-x^4} \arcsin(x) + \frac{\operatorname{arcsinh}(x)}{4}$$

[Out] 1/4*arcsinh(x)+1/4*x*(x^2+1)^(1/2)-1/2*arcsin(x)*(-x^4+1)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {267, 4871, 12, 26, 201, 221}

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = -\frac{1}{2}\sqrt{1-x^4} \arcsin(x) + \frac{\operatorname{arcsinh}(x)}{4} + \frac{1}{4}\sqrt{x^2+1}x$$

[In] Int[(x^3*ArcSin[x])/Sqrt[1 - x^4],x]

[Out] (x*Sqrt[1 + x^2])/4 - (Sqrt[1 - x^4]*ArcSin[x])/2 + ArcSinh[x]/4

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 26

Int[(u_.)*((a_) + (b_.)*(x_)^(n_.))^(m_.)*((c_) + (d_.)*(x_)^(j_.))^(p_.), x_Symbol] := Dist[(-b^2/d)^m, Int[u/(a - b*x^n)^m, x], x] /; FreeQ[{a, b, c, d, m, n, p}, x] && EqQ[j, 2*n] && EqQ[p, -m] && EqQ[b^2*c + a^2*d, 0] && G

tQ[a, 0] && LtQ[d, 0]

Rule 201

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 4871

```
Int[((a_.) + ArcSin[(c_.)*(x_)])*(b_.)*(u_), x_Symbol] := With[{v = IntHide
[u, x]}, Dist[a + b*ArcSin[c*x], v, x] - Dist[b*c, Int[SimplifyIntegrand[v/
Sqrt[1 - c^2*x^2], x], x], x] /; InverseFunctionFreeQ[v, x] /; FreeQ[{a, b
, c}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{2}\sqrt{1-x^4}\arcsin(x) - \int -\frac{\sqrt{1-x^4}}{2\sqrt{1-x^2}} dx \\
 &= -\frac{1}{2}\sqrt{1-x^4}\arcsin(x) + \frac{1}{2} \int \frac{\sqrt{1-x^4}}{\sqrt{1-x^2}} dx \\
 &= -\frac{1}{2}\sqrt{1-x^4}\arcsin(x) + \frac{1}{2} \int \sqrt{1+x^2} dx \\
 &= \frac{1}{4}x\sqrt{1+x^2} - \frac{1}{2}\sqrt{1-x^4}\arcsin(x) + \frac{1}{4} \int \frac{1}{\sqrt{1+x^2}} dx \\
 &= \frac{1}{4}x\sqrt{1+x^2} - \frac{1}{2}\sqrt{1-x^4}\arcsin(x) + \frac{\operatorname{arcsinh}(x)}{4}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 85 vs. $2(38) = 76$.

Time = 0.05 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.24

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \frac{1}{4} \left(\frac{x\sqrt{1-x^4}}{\sqrt{1-x^2}} - 2\sqrt{1-x^4} \arcsin(x) + \log(1-x^2) - \log\left(-x + x^3 + \sqrt{1-x^2}\sqrt{1-x^4}\right) \right)$$

[In] Integrate[(x^3*ArcSin[x])/Sqrt[1 - x^4],x]

[Out] ((x*Sqrt[1 - x^4])/Sqrt[1 - x^2] - 2*Sqrt[1 - x^4]*ArcSin[x] + Log[1 - x^2] - Log[-x + x^3 + Sqrt[1 - x^2]*Sqrt[1 - x^4]])/4

Maple [F]

$$\int \frac{x^3 \arcsin(x)}{\sqrt{-x^4+1}} dx$$

[In] int(x^3*arcsin(x)/(-x^4+1)^(1/2),x)

[Out] int(x^3*arcsin(x)/(-x^4+1)^(1/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. $2(28) = 56$.

Time = 0.26 (sec) , antiderivative size = 138, normalized size of antiderivative = 3.63

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \frac{4\sqrt{-x^4+1}(x^2-1)\arcsin(x) + 2\sqrt{-x^4+1}\sqrt{-x^2+1}x + (x^2-1)\log\left(\frac{x^3+\sqrt{-x^4+1}\sqrt{-x^2+1}-x}{x^3-x}\right) - (x^2-1)}{8(x^2-1)}$$

[In] integrate(x^3*arcsin(x)/(-x^4+1)^(1/2),x, algorithm="fricas")

[Out] -1/8*(4*sqrt(-x^4 + 1)*(x^2 - 1)*arcsin(x) + 2*sqrt(-x^4 + 1)*sqrt(-x^2 + 1)*x + (x^2 - 1)*log((x^3 + sqrt(-x^4 + 1)*sqrt(-x^2 + 1) - x)/(x^3 - x)) - (x^2 - 1)*log(-(x^3 - sqrt(-x^4 + 1)*sqrt(-x^2 + 1) - x)/(x^3 - x)))/(x^2 - 1)

Sympy [F]

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \int \frac{x^3 \operatorname{asin}(x)}{\sqrt{-(x-1)(x+1)(x^2+1)}} dx$$

[In] integrate(x**3*asin(x)/(-x**4+1)**(1/2),x)

[Out] Integral(x**3*asin(x)/sqrt(-(x - 1)*(x + 1)*(x**2 + 1)), x)

Maxima [F]

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \int \frac{x^3 \operatorname{asin}(x)}{\sqrt{-x^4+1}} dx$$

[In] integrate(x^3*arcsin(x)/(-x^4+1)^(1/2),x, algorithm="maxima")

[Out] -1/2*sqrt(x^2 + 1)*sqrt(x + 1)*sqrt(-x + 1)*arctan2(x, sqrt(x + 1)*sqrt(-x + 1)) + integrate(1/2*sqrt(x^2 + 1)/(x^2 + e^(log(x + 1) + log(-x + 1))), x)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \frac{1}{4} \sqrt{x^2+1}x - \frac{1}{2} \sqrt{-x^4+1} \arcsin(x) - \frac{1}{4} \log(-x + \sqrt{x^2+1})$$

[In] integrate(x^3*arcsin(x)/(-x^4+1)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(x^2 + 1)*x - 1/2*sqrt(-x^4 + 1)*arcsin(x) - 1/4*log(-x + sqrt(x^2 + 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3 \arcsin(x)}{\sqrt{1-x^4}} dx = \int \frac{x^3 \operatorname{asin}(x)}{\sqrt{1-x^4}} dx$$

[In] int((x^3*asin(x))/(1 - x^4)^(1/2),x)

[Out] int((x^3*asin(x))/(1 - x^4)^(1/2), x)

3.22 $\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx$

Optimal result	159
Rubi [A] (verified)	159
Mathematica [A] (verified)	161
Maple [F]	162
Fricas [B] (verification not implemented)	162
Sympy [F]	162
Maxima [F]	162
Giac [A] (verification not implemented)	163
Mupad [F(-1)]	163

Optimal result

Integrand size = 15, antiderivative size = 70

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx = -\frac{\sqrt{-1+x^4}}{2\sqrt{1-\frac{1}{x^2}x}} + \frac{1}{2}\sqrt{-1+x^4}\sec^{-1}(x) + \frac{1}{2}\operatorname{arctanh}\left(\frac{\sqrt{1-\frac{1}{x^2}x}}{\sqrt{-1+x^4}}\right)$$

[Out] $1/2*\operatorname{arctanh}(x*(1-1/x^2)^{(1/2)}/(x^4-1)^{(1/2)})+1/2*\operatorname{arcsec}(x)*(x^4-1)^{(1/2)}-1/2*(x^4-1)^{(1/2)}/x/(1-1/x^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.34, number of steps used = 7, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {267, 5354, 12, 1586, 1266, 879, 889, 209}

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx = \frac{\sqrt{1-x^2} \arctan\left(\frac{\sqrt{x^4-1}}{\sqrt{1-x^2}}\right)}{2\sqrt{1-\frac{1}{x^2}x}} + \frac{1}{2}\sqrt{x^4-1}\sec^{-1}(x) - \frac{\sqrt{x^4-1}}{2\sqrt{1-\frac{1}{x^2}x}}$$

[In] $\operatorname{Int}[(x^3*\operatorname{ArcSec}[x])/Sqrt[-1+x^4],x]$

[Out] $-1/2*Sqrt[-1+x^4]/(Sqrt[1-x^(-2)]*x) + (Sqrt[-1+x^4]*\operatorname{ArcSec}[x])/2 + (Sqrt[1-x^2]*\operatorname{ArcTan}[Sqrt[-1+x^4]/Sqrt[1-x^2]])/(2*Sqrt[1-x^(-2)]*x)$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\amp; \ !\operatorname{Match} Q[u, (b_*)(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]
```

Rule 879

```
Int[((d_) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))^(n_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-d + e*x)^m*(f + g*x)^(n + 1)*((a + c*x^2)^p/(g*(m - n - 1))), x] - Dist[c*m*((e*f + d*g)/(e^2*g*(m - n - 1))), Int[(d + e*x)^(m + 1)*(f + g*x)^n*(a + c*x^2)^(p - 1), x], x] /; FreeQ[{a, c, d, e, f, g, n}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + p, 0] && GtQ[p, 0] && NeQ[m - n - 1, 0] && !IGtQ[n, 0] && !IntegerQ[n + p] && LtQ[n + p + 2, 0] && RationalQ[n]
```

Rule 889

```
Int[Sqrt[(d_) + (e_.)*(x_)]/(((f_.) + (g_.)*(x_))*Sqrt[(a_) + (c_.)*(x_)^2]), x_Symbol] := Dist[2*e^2, Subst[Int[1/(c*(e*f + d*g) + e^2*g*x^2), x], x, Sqrt[a + c*x^2]/Sqrt[d + e*x]], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0]
```

Rule 1266

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x] && IntegerQ[(m + 1)/2]
```

Rule 1586

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^(mn_.))^(q_.)*((a_) + (c_.)*(x_)^(n2_.))^(p_.), x_Symbol] := Dist[(e^IntPart[q]*((d + e*x^mn)^FracPart[q]/(1 + d*(1/(x^mn*e))))^FracPart[q])/x^(mn*FracPart[q]), Int[x^(m + mn*q)*(1 + d*(1/(x^mn*e)))^q*(a + c*x^n2)^p, x], x] /; FreeQ[{a, c, d, e, m, mn, p, q}, x] && EqQ[n2, -2*mn] && !IntegerQ[p] && !IntegerQ[q] && PosQ[n2]
```

Rule 5354

```
Int[((a_.) + ArcSec[(c_.)*(x_)]*(b_.))*(u_), x_Symbol] := With[{v = IntHide[u, x]}, Dist[a + b*ArcSec[c*x], v, x] - Dist[b/c, Int[SimplifyIntegrand[v/
```


$(x^2 \sqrt{1 - 1/(c^2 x^2)}), x], x], x] /; \text{InverseFunctionFreeQ}[v, x]] /; \text{FreeQ}[a, b, c], x]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) - \int \frac{\sqrt{-1 + x^4}}{2 \sqrt{1 - \frac{1}{x^2} x^2}} dx \\
 &= \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) - \frac{1}{2} \int \frac{\sqrt{-1 + x^4}}{\sqrt{1 - \frac{1}{x^2} x^2}} dx \\
 &= \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) - \frac{\sqrt{1 - x^2} \int \frac{\sqrt{-1 + x^4}}{x \sqrt{1 - x^2}} dx}{2 \sqrt{1 - \frac{1}{x^2} x^2}} \\
 &= \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) - \frac{\sqrt{1 - x^2} \text{Subst}\left(\int \frac{\sqrt{-1 + x^2}}{\sqrt{1 - x x}} dx, x, x^2\right)}{4 \sqrt{1 - \frac{1}{x^2} x^2}} \\
 &= -\frac{\sqrt{-1 + x^4}}{2 \sqrt{1 - \frac{1}{x^2} x^2}} + \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) + \frac{\sqrt{1 - x^2} \text{Subst}\left(\int \frac{\sqrt{1 - x}}{x \sqrt{-1 + x^2}} dx, x, x^2\right)}{4 \sqrt{1 - \frac{1}{x^2} x^2}} \\
 &= -\frac{\sqrt{-1 + x^4}}{2 \sqrt{1 - \frac{1}{x^2} x^2}} + \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) + \frac{\sqrt{1 - x^2} \text{Subst}\left(\int \frac{1}{1 + x^2} dx, x, \frac{\sqrt{-1 + x^4}}{\sqrt{1 - x^2}}\right)}{2 \sqrt{1 - \frac{1}{x^2} x^2}} \\
 &= -\frac{\sqrt{-1 + x^4}}{2 \sqrt{1 - \frac{1}{x^2} x^2}} + \frac{1}{2} \sqrt{-1 + x^4} \sec^{-1}(x) + \frac{\sqrt{1 - x^2} \arctan\left(\frac{\sqrt{-1 + x^4}}{\sqrt{1 - x^2}}\right)}{2 \sqrt{1 - \frac{1}{x^2} x^2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.26

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1 + x^4}} dx = \frac{1}{2} \left(-\frac{\sqrt{1 - \frac{1}{x^2} x^2} \sqrt{-1 + x^4}}{-1 + x^2} + \sqrt{-1 + x^4} \sec^{-1}(x) - \log(x - x^3) \right. \\
 \left. + \log\left(1 - x^2 - \sqrt{1 - \frac{1}{x^2} x^2} \sqrt{-1 + x^4}\right) \right)$$

[In] Integrate[(x^3*ArcSec[x])/Sqrt[-1 + x^4],x]

[Out] (-((Sqrt[1 - x^(-2)]*x*Sqrt[-1 + x^4])/(-1 + x^2)) + Sqrt[-1 + x^4]*ArcSec[x] - Log[x - x^3] + Log[1 - x^2 - Sqrt[1 - x^(-2)]*x*Sqrt[-1 + x^4]])/2

Maple [F]

$$\int \frac{x^3 \operatorname{arcsec}(x)}{\sqrt{x^4 - 1}} dx$$

[In] int(x^3*arcsec(x)/(x^4-1)^(1/2),x)

[Out] int(x^3*arcsec(x)/(x^4-1)^(1/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(54) = 108.

Time = 0.26 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.57

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1 + x^4}} dx$$

$$= \frac{(x^2 - 1) \log\left(\frac{x^2 + \sqrt{x^4 - 1} \sqrt{x^2 - 1} - 1}{x^2 - 1}\right) - (x^2 - 1) \log\left(-\frac{x^2 - \sqrt{x^4 - 1} \sqrt{x^2 - 1} - 1}{x^2 - 1}\right) + 2\sqrt{x^4 - 1}((x^2 - 1) \operatorname{arcsec}(x) - \sqrt{x^4 - 1})}{4(x^2 - 1)}$$

[In] integrate(x^3*arcsec(x)/(x^4-1)^(1/2),x, algorithm="fricas")

[Out] 1/4*((x^2 - 1)*log((x^2 + sqrt(x^4 - 1)*sqrt(x^2 - 1) - 1)/(x^2 - 1)) - (x^2 - 1)*log(-(x^2 - sqrt(x^4 - 1)*sqrt(x^2 - 1) - 1)/(x^2 - 1)) + 2*sqrt(x^4 - 1)*((x^2 - 1)*arcsec(x) - sqrt(x^4 - 1)))/(x^2 - 1)

Sympy [F]

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1 + x^4}} dx = \int \frac{x^3 \operatorname{asec}(x)}{\sqrt{(x - 1)(x + 1)(x^2 + 1)}} dx$$

[In] integrate(x**3*asec(x)/(x**4-1)**(1/2),x)

[Out] Integral(x**3*asec(x)/sqrt((x - 1)*(x + 1)*(x**2 + 1)), x)

Maxima [F]

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1 + x^4}} dx = \int \frac{x^3 \operatorname{arcsec}(x)}{\sqrt{x^4 - 1}} dx$$

[In] integrate(x^3*arcsec(x)/(x^4-1)^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(x^2 + 1)*sqrt(x + 1)*sqrt(x - 1)*arctan(sqrt(x + 1)*sqrt(x - 1)) - integrate((2*(x^3*e^(3/2*log(x + 1) + 3/2*log(x - 1))) + x^3*e^(1/2*log(x + 1) + 1/2*log(x - 1)))*sqrt(x^2 + 1)*log(x) + (x^3 + x)*e^(1/2*log(x^2 + 1) + 3/2*log(x + 1) + 3/2*log(x - 1)))/((x^2 + 1)*(e^(2*log(x + 1) + 2*log(x - 1)) + e^(log(x + 1) + log(x - 1))))), x)

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.74

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx = \frac{1}{2} \sqrt{x^4-1} \arccos\left(\frac{1}{x}\right) - \frac{2\sqrt{x^2+1} - \log(\sqrt{x^2+1}+1) + \log(\sqrt{x^2+1}-1)}{4 \operatorname{sgn}(x)}$$

[In] integrate(x^3*arcsec(x)/(x^4-1)^(1/2),x, algorithm="giac")

[Out] 1/2*sqrt(x^4 - 1)*arccos(1/x) - 1/4*(2*sqrt(x^2 + 1) - log(sqrt(x^2 + 1) + 1) + log(sqrt(x^2 + 1) - 1))/sgn(x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3 \sec^{-1}(x)}{\sqrt{-1+x^4}} dx = \int \frac{x^3 \operatorname{acos}\left(\frac{1}{x}\right)}{\sqrt{x^4-1}} dx$$

[In] int((x^3*acos(1/x))/(x^4 - 1)^(1/2),x)

[Out] int((x^3*acos(1/x))/(x^4 - 1)^(1/2), x)

$$3.23 \quad \int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx$$

Optimal result	164
Rubi [A] (verified)	164
Mathematica [A] (verified)	166
Maple [F]	166
Fricas [A] (verification not implemented)	167
Sympy [F(-1)]	167
Maxima [F]	167
Giac [F]	168
Mupad [F(-1)]	168

Optimal result

Integrand size = 25, antiderivative size = 58

$$\int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = -x \arctan(x) + \frac{1}{2} \log(1+x^2) + \sqrt{1+x^2} \arctan(x) \log(x + \sqrt{1+x^2}) - \frac{1}{2} \log^2(x + \sqrt{1+x^2})$$

[Out] -x*arctan(x)+1/2*ln(x^2+1)-1/2*ln(x+(x^2+1)^(1/2))^2+arctan(x)*ln(x+(x^2+1)^(1/2))*(x^2+1)^(1/2)

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {5050, 221, 267, 2634, 8, 5320, 6818, 4930, 266}

$$\int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) \log(\sqrt{x^2+1} + x) - x \arctan(x) - \frac{1}{2} \log^2(\sqrt{x^2+1} + x) + \frac{1}{2} \log(x^2+1)$$

[In] Int[(x*ArcTan[x]*Log[x + Sqrt[1 + x^2]])/Sqrt[1 + x^2],x]

[Out] -(x*ArcTan[x]) + Log[1 + x^2]/2 + Sqrt[1 + x^2]*ArcTan[x]*Log[x + Sqrt[1 + x^2]] - Log[x + Sqrt[1 + x^2]]^2/2

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 221

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 266

`Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]`

Rule 267

`Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]`

Rule 2634

`Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]`

Rule 4930

`Int[((a_) + ArcTan[(c_)*(x_)^(n_)])*(b_)^(p_), x_Symbol] := Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])`

Rule 5050

`Int[((a_) + ArcTan[(c_)*(x_)])*(b_)^(p_)*(x)*((d_) + (e_)*(x_)^2)^(q_), x_Symbol] := Simp[(d + e*x^2)^(q + 1)*((a + b*ArcTan[c*x])^p/(2*e*(q + 1))), x] - Dist[b*(p/(2*c*(q + 1))), Int[(d + e*x^2)^q*(a + b*ArcTan[c*x])^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && EqQ[e, c^2*d] && GtQ[p, 0] && NeQ[q, -1]`

Rule 5320

`Int[ArcTan[v_]*Log[w_]*(u_), x_Symbol] := With[{z = IntHide[u, x]}, Dist[ArcTan[v]*Log[w], z, x] + (-Int[SimplifyIntegrand[z*Log[w]*(D[v, x]/(1 + v^2)), x], x] - Int[SimplifyIntegrand[z*ArcTan[v]*(D[w, x]/w), x], x]) /; InverseFunctionFreeQ[z, x] /; InverseFunctionFreeQ[v, x] && InverseFunctionFreeQ[w, x]`

Rule 6818

`Int[(u_)*(y_)^(m_.), x_Symbol] := With[{q = DerivativeDivides[y, u, x]}, Si
mp[q*(y^(m + 1)/(m + 1)), x] /; !FalseQ[q]] /; FreeQ[m, x] && NeQ[m, -1]`

Rubi steps

$$\begin{aligned} \text{integral} &= \sqrt{1+x^2} \arctan(x) \log(x + \sqrt{1+x^2}) - \int \arctan(x) dx - \int \frac{\log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx \\ &= -x \arctan(x) + \sqrt{1+x^2} \arctan(x) \log(x + \sqrt{1+x^2}) - \frac{1}{2} \log^2(x + \sqrt{1+x^2}) + \int \frac{x}{1+x^2} dx \\ &= -x \arctan(x) + \frac{1}{2} \log(1+x^2) + \sqrt{1+x^2} \arctan(x) \log(x + \sqrt{1+x^2}) - \frac{1}{2} \log^2(x \\ &\quad + \sqrt{1+x^2}) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx &= -x \arctan(x) + \frac{1}{2} \log(1+x^2) \\ &\quad + \sqrt{1+x^2} \arctan(x) \log(x + \sqrt{1+x^2}) \\ &\quad - \frac{1}{2} \log^2(x + \sqrt{1+x^2}) \end{aligned}$$

[In] `Integrate[(x*ArcTan[x]*Log[x + Sqrt[1 + x^2]])/Sqrt[1 + x^2],x]`

[Out] `-(x*ArcTan[x]) + Log[1 + x^2]/2 + Sqrt[1 + x^2]*ArcTan[x]*Log[x + Sqrt[1 + x^2]] - Log[x + Sqrt[1 + x^2]]^2/2`

Maple [F]

$$\int \frac{x \arctan(x) \ln(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] `int(x*arctan(x)*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x)`

[Out] `int(x*arctan(x)*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.83

$$\int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) \log(x + \sqrt{x^2+1}) - x \arctan(x) - \frac{1}{2} \log(x + \sqrt{x^2+1})^2 + \frac{1}{2} \log(x^2+1)$$

[In] integrate(x*arctan(x)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] sqrt(x^2 + 1)*arctan(x)*log(x + sqrt(x^2 + 1)) - x*arctan(x) - 1/2*log(x + sqrt(x^2 + 1))^2 + 1/2*log(x^2 + 1)

Sympy [F(-1)]

Timed out.

$$\int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \text{Timed out}$$

[In] integrate(x*atan(x)*ln(x+(x**2+1)**(1/2))/(x**2+1)**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{x \arctan(x) \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \int \frac{x \arctan(x) \log(x + \sqrt{x^2+1})}{\sqrt{x^2+1}} dx$$

[In] integrate(x*arctan(x)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(x*arctan(x)*log(x + sqrt(x^2 + 1))/sqrt(x^2 + 1), x)

Giac [F]

$$\int \frac{x \arctan(x) \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \int \frac{x \arctan(x) \log(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] integrate(x*arctan(x)*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="giac")

[Out] integrate(x*arctan(x)*log(x + sqrt(x^2 + 1))/sqrt(x^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \arctan(x) \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \int \frac{x \operatorname{atan}(x) \ln(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] int((x*atan(x)*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2),x)

[Out] int((x*atan(x)*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2), x)

$$3.24 \quad \int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx$$

Optimal result	169
Rubi [A] (verified)	169
Mathematica [A] (verified)	171
Maple [A] (verified)	171
Fricas [A] (verification not implemented)	171
Sympy [A] (verification not implemented)	172
Maxima [A] (verification not implemented)	172
Giac [A] (verification not implemented)	172
Mupad [B] (verification not implemented)	172

Optimal result

Integrand size = 27, antiderivative size = 55

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \sqrt{1 - x^2} - \log(1 + \sqrt{1 - x^2}) - \sqrt{1 - x^2} \log(1 + \sqrt{1 - x^2})$$

[Out] $-\ln(1+(-x^2+1)^{(1/2)})+(-x^2+1)^{(1/2)}-\ln(1+(-x^2+1)^{(1/2)})*(-x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.185$, Rules used = {267, 2634, 1605, 196, 45}

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \sqrt{1 - x^2} - \sqrt{1 - x^2} \log(\sqrt{1 - x^2} + 1) - \log(\sqrt{1 - x^2} + 1)$$

[In] `Int[(x*Log[1 + Sqrt[1 - x^2]])/Sqrt[1 - x^2],x]`

[Out] `Sqrt[1 - x^2] - Log[1 + Sqrt[1 - x^2]] - Sqrt[1 - x^2]*Log[1 + Sqrt[1 - x^2]]`

Rule 45

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 196

```
Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(1/n
- 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, p}, x] && FractionQ[n] &&
IntegerQ[1/n]
```

Rule 267

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 1605

```
Int[((a_.) + (b_.)*(Pq_)^(n_.))^(p_.)*(Qr_), x_Symbol] := With[{q = Expon[P
q, x], r = Expon[Qr, x]}, Dist[Coeff[Qr, x, r]/(q*Coeff[Pq, x, q]), Subst[Int[(a + b*x^n)^p, x], x, Pq], x] /; EqQ[r, q - 1] && EqQ[Coeff[Qr, x, r]*D[Pq, x], q*Coeff[Pq, x, q]*Qr] /; FreeQ[{a, b, n, p}, x] && PolyQ[Pq, x] && PolyQ[Qr, x]
```

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\sqrt{1-x^2} \log\left(1 + \sqrt{1-x^2}\right) - \int \frac{x}{1 + \sqrt{1-x^2}} dx \\
&= -\sqrt{1-x^2} \log\left(1 + \sqrt{1-x^2}\right) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{1 + \sqrt{x}} dx, x, 1-x^2\right) \\
&= -\sqrt{1-x^2} \log\left(1 + \sqrt{1-x^2}\right) + \text{Subst}\left(\int \frac{x}{1+x} dx, x, \sqrt{1-x^2}\right) \\
&= -\sqrt{1-x^2} \log\left(1 + \sqrt{1-x^2}\right) + \text{Subst}\left(\int \left(1 + \frac{1}{-1-x}\right) dx, x, \sqrt{1-x^2}\right) \\
&= \sqrt{1-x^2} - \log\left(1 + \sqrt{1-x^2}\right) - \sqrt{1-x^2} \log\left(1 + \sqrt{1-x^2}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.75

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \sqrt{1 - x^2} - (1 + \sqrt{1 - x^2}) \log(1 + \sqrt{1 - x^2})$$

[In] Integrate[(x*Log[1 + Sqrt[1 - x^2]])/Sqrt[1 - x^2],x]

[Out] Sqrt[1 - x^2] - (1 + Sqrt[1 - x^2])*Log[1 + Sqrt[1 - x^2]]

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.67

method	result	size
derivativedivides	$-\ln(1 + \sqrt{-x^2 + 1})(1 + \sqrt{-x^2 + 1}) + 1 + \sqrt{-x^2 + 1}$	37
default	$-\ln(1 + \sqrt{-x^2 + 1})(1 + \sqrt{-x^2 + 1}) + 1 + \sqrt{-x^2 + 1}$	37

[In] int(x*ln(1+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] -ln(1+(-x^2+1)^(1/2))*(1+(-x^2+1)^(1/2))+1+(-x^2+1)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.64

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = -(\sqrt{-x^2 + 1} + 1) \log(\sqrt{-x^2 + 1} + 1) + \sqrt{-x^2 + 1}$$

[In] integrate(x*log(1+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -(sqrt(-x^2 + 1) + 1)*log(sqrt(-x^2 + 1) + 1) + sqrt(-x^2 + 1)

Sympy [A] (verification not implemented)

Time = 2.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.56

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \sqrt{1 - x^2} - (\sqrt{1 - x^2} + 1) \log(\sqrt{1 - x^2} + 1) + 1$$

[In] integrate(x*ln(1+(-x**2+1)**(1/2))/(-x**2+1)**(1/2),x)

[Out] sqrt(1 - x**2) - (sqrt(1 - x**2) + 1)*log(sqrt(1 - x**2) + 1) + 1

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.65

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = -(\sqrt{-x^2 + 1} + 1) \log(\sqrt{-x^2 + 1} + 1) + \sqrt{-x^2 + 1} + 1$$

[In] integrate(x*log(1+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] -(sqrt(-x^2 + 1) + 1)*log(sqrt(-x^2 + 1) + 1) + sqrt(-x^2 + 1) + 1

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.65

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = -(\sqrt{-x^2 + 1} + 1) \log(\sqrt{-x^2 + 1} + 1) + \sqrt{-x^2 + 1} + 1$$

[In] integrate(x*log(1+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] -(sqrt(-x^2 + 1) + 1)*log(sqrt(-x^2 + 1) + 1) + sqrt(-x^2 + 1) + 1

Mupad [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.49

$$\int \frac{x \log(1 + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = -(\ln(\sqrt{1 - x^2} + 1) - 1) (\sqrt{1 - x^2} + 1)$$

[In] int((x*log((1 - x^2)^(1/2) + 1))/(1 - x^2)^(1/2),x)

[Out] -(log((1 - x^2)^(1/2) + 1) - 1)*((1 - x^2)^(1/2) + 1)

$$3.25 \quad \int \frac{x \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx$$

Optimal result	173
Rubi [A] (verified)	173
Mathematica [A] (verified)	174
Maple [F]	174
Fricas [A] (verification not implemented)	174
Sympy [A] (verification not implemented)	175
Maxima [A] (verification not implemented)	175
Giac [A] (verification not implemented)	175
Mupad [F(-1)]	175

Optimal result

Integrand size = 23, antiderivative size = 26

$$\int \frac{x \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = -x + \sqrt{1+x^2} \log(x + \sqrt{1+x^2})$$

[Out] $-x + \ln(x + (x^2 + 1)^{1/2}) * (x^2 + 1)^{1/2}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {267, 2634, 8}

$$\int \frac{x \log(x + \sqrt{1+x^2})}{\sqrt{1+x^2}} dx = \sqrt{x^2 + 1} \log(\sqrt{x^2 + 1} + x) - x$$

[In] $\text{Int}[(x * \text{Log}[x + \text{Sqrt}[1 + x^2]]) / \text{Sqrt}[1 + x^2], x]$

[Out] $-x + \text{Sqrt}[1 + x^2] * \text{Log}[x + \text{Sqrt}[1 + x^2]]$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 267

$\text{Int}[(x_)^{(m_)} * ((a_) + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p+1)} / (b*n*(p+1)), x] /; \text{FreeQ}\{a, b, m, n, p\}, x \ \&\& \ \text{EqQ}[m, n - 1] \ \&\& \ \text{NeQ}[p, -1]$

Rule 2634

```
Int[Log[u]*(v_), x_Symbol] :> With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \sqrt{1+x^2} \log\left(x + \sqrt{1+x^2}\right) - \int 1 dx \\ &= -x + \sqrt{1+x^2} \log\left(x + \sqrt{1+x^2}\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{x \log\left(x + \sqrt{1+x^2}\right)}{\sqrt{1+x^2}} dx = -x + \sqrt{1+x^2} \log\left(x + \sqrt{1+x^2}\right)$$

```
[In] Integrate[(x*Log[x + Sqrt[1 + x^2]])/Sqrt[1 + x^2], x]
```

```
[Out] -x + Sqrt[1 + x^2]*Log[x + Sqrt[1 + x^2]]
```

Maple [F]

$$\int \frac{x \ln\left(x + \sqrt{x^2 + 1}\right)}{\sqrt{x^2 + 1}} dx$$

```
[In] int(x*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2), x)
```

```
[Out] int(x*ln(x+(x^2+1)^(1/2))/(x^2+1)^(1/2), x)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log\left(x + \sqrt{1+x^2}\right)}{\sqrt{1+x^2}} dx = \sqrt{x^2 + 1} \log\left(x + \sqrt{x^2 + 1}\right) - x$$

```
[In] integrate(x*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2), x, algorithm="fricas")
```

```
[Out] sqrt(x^2 + 1)*log(x + sqrt(x^2 + 1)) - x
```

Sympy [A] (verification not implemented)

Time = 3.57 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = -x + \sqrt{x^2 + 1} \log(x + \sqrt{x^2 + 1})$$

[In] integrate(x*ln(x+(x**2+1)**(1/2))/(x**2+1)**(1/2),x)

[Out] -x + sqrt(x**2 + 1)*log(x + sqrt(x**2 + 1))

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \sqrt{x^2 + 1} \log(x + \sqrt{x^2 + 1}) - x$$

[In] integrate(x*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 + 1)*log(x + sqrt(x^2 + 1)) - x

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \sqrt{x^2 + 1} \log(x + \sqrt{x^2 + 1}) - x$$

[In] integrate(x*log(x+(x^2+1)^(1/2))/(x^2+1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 + 1)*log(x + sqrt(x^2 + 1)) - x

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(x + \sqrt{1 + x^2})}{\sqrt{1 + x^2}} dx = \int \frac{x \ln(x + \sqrt{x^2 + 1})}{\sqrt{x^2 + 1}} dx$$

[In] int((x*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2),x)

[Out] int((x*log(x + (x^2 + 1)^(1/2)))/(x^2 + 1)^(1/2), x)

$$3.26 \quad \int \frac{x \log(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx$$

Optimal result	176
Rubi [A] (verified)	176
Mathematica [A] (verified)	179
Maple [F]	180
Fricas [A] (verification not implemented)	180
Sympy [F]	180
Maxima [F]	181
Giac [A] (verification not implemented)	181
Mupad [F(-1)]	182

Optimal result

Integrand size = 27, antiderivative size = 78

$$\int \frac{x \log(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = \sqrt{1-x^2} + \frac{\operatorname{arctanh}(\sqrt{2}x)}{\sqrt{2}} - \frac{\operatorname{arctanh}(\sqrt{2}\sqrt{1-x^2})}{\sqrt{2}} - \sqrt{1-x^2} \log(x + \sqrt{1-x^2})$$

[Out] 1/2*arctanh(x*2^(1/2))*2^(1/2)-1/2*arctanh(2^(1/2)*(-x^2+1)^(1/2))*2^(1/2)+(-x^2+1)^(1/2)-ln(x+(-x^2+1)^(1/2))*(-x^2+1)^(1/2)

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.407$, Rules used = {267, 2634, 6874, 2132, 327, 212, 455, 52, 65, 213, 396}

$$\int \frac{x \log(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx = -\frac{\operatorname{arctanh}(\sqrt{2}\sqrt{1-x^2})}{\sqrt{2}} + \frac{\operatorname{arctanh}(\sqrt{2}x)}{\sqrt{2}} + \sqrt{1-x^2} - \sqrt{1-x^2} \log(\sqrt{1-x^2} + x)$$

[In] Int[(x*Log[x + Sqrt[1 - x^2]])/Sqrt[1 - x^2],x]

[Out] Sqrt[1 - x^2] + ArcTanh[Sqrt[2]*x]/Sqrt[2] - ArcTanh[Sqrt[2]*Sqrt[1 - x^2]]/Sqrt[2] - Sqrt[1 - x^2]*Log[x + Sqrt[1 - x^2]]

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(

$b*(m + n + 1))$, $\text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x\}$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{GtQ}[n, 0]$ && $\text{NeQ}[m + n + 1, 0]$ && $!(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] \mid\mid (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0])))$ && $!\text{ILtQ}[m + n + 2, 0]$ && $\text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] /;$ $\text{FreeQ}\{a, b, c, d\}, x\}$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{LtQ}[-1, m, 0]$ && $\text{LeQ}[-1, n, 0]$ && $\text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]]$ && $\text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

$\text{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x\}$ && $\text{NegQ}[a/b]$ && $(\text{GtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

Rule 213

$\text{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[b, 2])^{-1})*\text{ArcTanh}[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])], x] /;$ $\text{FreeQ}\{a, b\}, x\}$ && $\text{NegQ}[a/b]$ && $(\text{LtQ}[a, 0] \mid\mid \text{GtQ}[b, 0])$

Rule 267

$\text{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x_Symbol] := \text{Simp}[(a + b*x^n)^{(p + 1)}/(b*n*(p + 1)), x] /;$ $\text{FreeQ}\{a, b, m, n, p\}, x\}$ && $\text{EqQ}[m, n - 1]$ && $\text{NeQ}[p, -1]$

Rule 327

$\text{Int}[(c_.)*(x_.))^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x_Symbol] := \text{Simp}[c^{(n - 1)}*(c*x)^{(m - n + 1)}*((a + b*x^n)^{(p + 1)}/(b*(m + n*p + 1))), x] - \text{Dist}[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), \text{Int}[(c*x)^{(m - n)}*(a + b*x^n)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, p\}, x\}$ && $\text{IGtQ}[n, 0]$ && $\text{GtQ}[m, n - 1]$ && $\text{NeQ}[m + n*p + 1, 0]$ && $\text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 396

$\text{Int}[(a_. + (b_.)*(x_.)^{(n_.)})^{(p_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), x_Symbol] := \text{Simp}[d*x*((a + b*x^n)^{(p + 1)}/(b*(n*(p + 1) + 1))), x] - \text{Dist}[(a*d - b*c*(n*(p + 1) + 1))/(b*(n*(p + 1) + 1)), \text{Int}[(a + b*x^n)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n\}, x\}$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{NeQ}[n*(p + 1) + 1, 0]$

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]
```

Rule 2132

```
Int[(x_)^(m_.)/((d_.)*(x_)^(n_.) + (c_.)*Sqrt[(a_.) + (b_.)*(x_)^(p_.)]), x_Symbol] := Dist[-d, Int[x^(m + n)/(a*c^2 + (b*c^2 - d^2)*x^(2*n)), x], x] + Dist[c, Int[(x^m*Sqrt[a + b*x^(2*n)])/(a*c^2 + (b*c^2 - d^2)*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, m, n}, x] && EqQ[p, 2*n] && NeQ[b*c^2 - d^2, 0]
```

Rule 2634

```
Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\sqrt{1-x^2} \log(x + \sqrt{1-x^2}) - \int \frac{x - \sqrt{1-x^2}}{x + \sqrt{1-x^2}} dx \\
&= -\sqrt{1-x^2} \log(x + \sqrt{1-x^2}) - \int \left(\frac{x}{x + \sqrt{1-x^2}} - \frac{\sqrt{1-x^2}}{x + \sqrt{1-x^2}} \right) dx \\
&= -\sqrt{1-x^2} \log(x + \sqrt{1-x^2}) - \int \frac{x}{x + \sqrt{1-x^2}} dx + \int \frac{\sqrt{1-x^2}}{x + \sqrt{1-x^2}} dx \\
&= -\sqrt{1-x^2} \log(x + \sqrt{1-x^2}) + \int \frac{x^2}{1-2x^2} dx \\
&\quad - \int \frac{x\sqrt{1-x^2}}{1-2x^2} dx + \int \left(\frac{x\sqrt{1-x^2}}{-1+2x^2} - \frac{1-x^2}{-1+2x^2} \right) dx \\
&= -\frac{x}{2} - \sqrt{1-x^2} \log(x + \sqrt{1-x^2}) + \frac{1}{2} \int \frac{1}{1-2x^2} dx \\
&\quad - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1-x}}{1-2x} dx, x, x^2 \right) + \int \frac{x\sqrt{1-x^2}}{-1+2x^2} dx - \int \frac{1-x^2}{-1+2x^2} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{1-x^2}}{2} + \frac{\operatorname{arctanh}(\sqrt{2x})}{2\sqrt{2}} - \sqrt{1-x^2} \log(x + \sqrt{1-x^2}) \\
&\quad - \frac{1}{4} \operatorname{Subst}\left(\int \frac{1}{(1-2x)\sqrt{1-x}} dx, x, x^2\right) \\
&\quad - \frac{1}{2} \int \frac{1}{-1+2x^2} dx + \frac{1}{2} \operatorname{Subst}\left(\int \frac{\sqrt{1-x}}{-1+2x} dx, x, x^2\right) \\
&= \sqrt{1-x^2} + \frac{\operatorname{arctanh}(\sqrt{2x})}{\sqrt{2}} - \sqrt{1-x^2} \log(x + \sqrt{1-x^2}) \\
&\quad + \frac{1}{4} \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-x}(-1+2x)} dx, x, x^2\right) \\
&\quad + \frac{1}{2} \operatorname{Subst}\left(\int \frac{1}{-1+2x^2} dx, x, \sqrt{1-x^2}\right) \\
&= \sqrt{1-x^2} + \frac{\operatorname{arctanh}(\sqrt{2x})}{\sqrt{2}} - \frac{\operatorname{arctanh}(\sqrt{2}\sqrt{1-x^2})}{2\sqrt{2}} \\
&\quad - \sqrt{1-x^2} \log(x + \sqrt{1-x^2}) - \frac{1}{2} \operatorname{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \sqrt{1-x^2}\right) \\
&= \sqrt{1-x^2} + \frac{\operatorname{arctanh}(\sqrt{2x})}{\sqrt{2}} - \frac{\operatorname{arctanh}(\sqrt{2}\sqrt{1-x^2})}{\sqrt{2}} - \sqrt{1-x^2} \log(x + \sqrt{1-x^2})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.53

$$\begin{aligned}
\int \frac{x \log(x + \sqrt{1-x^2})}{\sqrt{1-x^2}} dx &= \frac{1}{4} \left(4\sqrt{1-x^2} + 2\sqrt{2} \log(\sqrt{2} + 2x) \right. \\
&\quad \left. - \sqrt{2} \log(2 - \sqrt{2}x + \sqrt{2-2x^2}) \right. \\
&\quad \left. - \sqrt{2} \log(2 + \sqrt{2}x + \sqrt{2-2x^2}) \right. \\
&\quad \left. - 4\sqrt{1-x^2} \log(x + \sqrt{1-x^2}) \right)
\end{aligned}$$

[In] Integrate[(x*Log[x + Sqrt[1 - x^2]])/Sqrt[1 - x^2], x]

[Out] (4*Sqrt[1 - x^2] + 2*Sqrt[2]*Log[Sqrt[2] + 2*x] - Sqrt[2]*Log[2 - Sqrt[2]*x + Sqrt[2 - 2*x^2]] - Sqrt[2]*Log[2 + Sqrt[2]*x + Sqrt[2 - 2*x^2]] - 4*Sqrt[1 - x^2]*Log[x + Sqrt[1 - x^2]])/4

Maple [F]

$$\int \frac{x \ln(x + \sqrt{-x^2 + 1})}{\sqrt{-x^2 + 1}} dx$$

[In] int(x*ln(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x)

[Out] int(x*ln(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.47

$$\begin{aligned} & \int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx \\ &= -\sqrt{-x^2 + 1} \log(x + \sqrt{-x^2 + 1}) \\ & \quad + \frac{1}{4} \sqrt{2} \log\left(\frac{6x^2 - 2\sqrt{2}(2x^2 - 3) + 2\sqrt{-x^2 + 1}(3\sqrt{2} - 4) - 9}{2x^2 - 1}\right) \\ & \quad + \frac{1}{4} \sqrt{2} \log\left(\frac{2x^2 + 2\sqrt{2}x + 1}{2x^2 - 1}\right) + \sqrt{-x^2 + 1} \end{aligned}$$

[In] integrate(x*log(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -sqrt(-x^2 + 1)*log(x + sqrt(-x^2 + 1)) + 1/4*sqrt(2)*log((6*x^2 - 2*sqrt(2)*(2*x^2 - 3) + 2*sqrt(-x^2 + 1)*(3*sqrt(2) - 4) - 9)/(2*x^2 - 1)) + 1/4*sqrt(2)*log((2*x^2 + 2*sqrt(2)*x + 1)/(2*x^2 - 1)) + sqrt(-x^2 + 1)

Sympy [F]

$$\int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{-(x - 1)(x + 1)}} dx$$

[In] integrate(x*ln(x+(-x**2+1)**(1/2))/(-x**2+1)**(1/2),x)

[Out] Integral(x*log(x + sqrt(1 - x**2))/sqrt(-(x - 1)*(x + 1)), x)

Maxima [F]

$$\int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \int \frac{x \log(x + \sqrt{-x^2 + 1})}{\sqrt{-x^2 + 1}} dx$$

[In] integrate(x*log(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] (x^2 - 1)*log(x + sqrt(x + 1)*sqrt(-x + 1))/(sqrt(x + 1)*sqrt(-x + 1)) - integrate((x^2 - 1)*e^(-1/2*log(x + 1) - 1/2*log(-x + 1))/x, x) - integrate(1/(x^2 + sqrt(x + 1)*x*sqrt(-x + 1)), x)

Giac [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.56

$$\int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = -\sqrt{-x^2 + 1} \log(x + \sqrt{-x^2 + 1}) - \frac{1}{4} \sqrt{2} \log \left(\frac{\left| -4\sqrt{2} + \frac{2(\sqrt{-x^2+1}-1)^2}{x^2} - 6 \right|}{\left| 4\sqrt{2} + \frac{2(\sqrt{-x^2+1}-1)^2}{x^2} - 6 \right|} \right) + \frac{1}{4} \sqrt{2} \log \left(\left| x + \frac{1}{2} \sqrt{2} \right| \right) - \frac{1}{4} \sqrt{2} \log \left(\left| x - \frac{1}{2} \sqrt{2} \right| \right) + \sqrt{-x^2 + 1}$$

[In] integrate(x*log(x+(-x^2+1)^(1/2))/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] -sqrt(-x^2 + 1)*log(x + sqrt(-x^2 + 1)) - 1/4*sqrt(2)*log(abs(-4*sqrt(2) + 2*(sqrt(-x^2 + 1) - 1)^2/x^2 - 6)/abs(4*sqrt(2) + 2*(sqrt(-x^2 + 1) - 1)^2/x^2 - 6)) + 1/4*sqrt(2)*log(abs(x + 1/2*sqrt(2))) - 1/4*sqrt(2)*log(abs(x - 1/2*sqrt(2))) + sqrt(-x^2 + 1)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx = \int \frac{x \ln(x + \sqrt{1 - x^2})}{\sqrt{1 - x^2}} dx$$

```
[In] int((x*log(x + (1 - x^2)^(1/2)))/(1 - x^2)^(1/2), x)
```

```
[Out] int((x*log(x + (1 - x^2)^(1/2)))/(1 - x^2)^(1/2), x)
```

3.27 $\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx$

Optimal result	183
Rubi [A] (verified)	183
Mathematica [A] (verified)	184
Maple [A] (verified)	184
Fricas [A] (verification not implemented)	185
Sympy [F]	185
Maxima [A] (verification not implemented)	185
Giac [B] (verification not implemented)	185
Mupad [F(-1)]	186

Optimal result

Integrand size = 17, antiderivative size = 39

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}}{x} - \arcsin(x) - \frac{\sqrt{1-x^2}\log(x)}{x}$$

[Out] $-\arcsin(x) - (-x^2+1)^{(1/2)}/x - \ln(x) * (-x^2+1)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2373, 283, 222}

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = -\arcsin(x) - \frac{\sqrt{1-x^2}}{x} - \frac{\sqrt{1-x^2}\log(x)}{x}$$

[In] `Int[Log[x]/(x^2*Sqrt[1 - x^2]),x]`

[Out] $-(\text{Sqrt}[1 - x^2]/x) - \text{ArcSin}[x] - (\text{Sqrt}[1 - x^2]*\text{Log}[x])/x$

Rule 222

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 283

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a + b*x^n)^p/(c*(m+1))), x] - Dist[b*n*(p/(c^n*(m+1))), Int[(c*x)^(m+n)*(a + b*x^n)^(p-1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && GtQ[p, 0] && LtQ[m, -1] && !ILtQ[(m+n*p+n+1)/n, 0] && IntBi`

nomialQ[a, b, c, n, m, p, x]

Rule 2373

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*
(x_)^(r_.))^(q_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^r)^(q + 1)*((a +
b*Log[c*x^n])/(d*f*(m + 1))), x] - Dist[b*(n/(d*(m + 1))), Int[(f*x)^m*(d
+ e*x^r)^(q + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, n, q, r}, x] && EqQ
[m + r*(q + 1) + 1, 0] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\sqrt{1-x^2} \log(x)}{x} + \int \frac{\sqrt{1-x^2}}{x^2} dx \\ &= -\frac{\sqrt{1-x^2}}{x} - \frac{\sqrt{1-x^2} \log(x)}{x} - \int \frac{1}{\sqrt{1-x^2}} dx \\ &= -\frac{\sqrt{1-x^2}}{x} - \arcsin(x) - \frac{\sqrt{1-x^2} \log(x)}{x} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.64

$$\int \frac{\log(x)}{x^2 \sqrt{1-x^2}} dx = -\arcsin(x) - \frac{\sqrt{1-x^2}(1+\log(x))}{x}$$

[In] Integrate[Log[x]/(x^2*Sqrt[1 - x^2]),x]

[Out] -ArcSin[x] - (Sqrt[1 - x^2]*(1 + Log[x]))/x

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

method	result	size
meijerg	$-\arcsin(x) + \frac{-\ln(x)\sqrt{-x^2+1}-\sqrt{-x^2+1}}{x}$	35

[In] int(ln(x)/x^2/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] -arcsin(x)+(-ln(x)*(-x^2+1)^(1/2)-(-x^2+1)^(1/2))/x

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = \frac{2x \arctan\left(\frac{\sqrt{-x^2+1}-1}{x}\right) - \sqrt{-x^2+1}(\log(x)+1)}{x}$$

[In] integrate(log(x)/x^2/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] (2*x*arctan((sqrt(-x^2 + 1) - 1)/x) - sqrt(-x^2 + 1)*(log(x) + 1))/x

Sympy [F]

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\log(x)}{x^2\sqrt{-(x-1)(x+1)}} dx$$

[In] integrate(ln(x)/x**2/(-x**2+1)**(1/2),x)

[Out] Integral(log(x)/(x**2*sqrt(-(x - 1)*(x + 1))), x)

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.90

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{-x^2+1}\log(x)}{x} - \frac{\sqrt{-x^2+1}}{x} - \arcsin(x)$$

[In] integrate(log(x)/x^2/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(-x^2 + 1)*log(x)/x - sqrt(-x^2 + 1)/x - arcsin(x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(35) = 70.

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.87

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = \frac{1}{2} \left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right) \log(x) + \frac{x}{2(\sqrt{-x^2+1}-1)} - \frac{\sqrt{-x^2+1}-1}{2x} - \arcsin(x)$$

[In] integrate(log(x)/x^2/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/2*(x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)*log(x) + 1/2*x/(sqrt(-x^2 + 1) - 1) - 1/2*(sqrt(-x^2 + 1) - 1)/x - arcsin(x)

Mupad **[F(-1)]**

Timed out.

$$\int \frac{\log(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\ln(x)}{x^2\sqrt{1-x^2}} dx$$

[In] int(log(x)/(x^2*(1 - x^2)^(1/2)),x)

[Out] int(log(x)/(x^2*(1 - x^2)^(1/2)), x)

3.28 $\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx$

Optimal result	187
Rubi [A] (verified)	187
Mathematica [A] (verified)	188
Maple [C] (verified)	188
Fricas [A] (verification not implemented)	188
Sympy [B] (verification not implemented)	189
Maxima [A] (verification not implemented)	189
Giac [A] (verification not implemented)	189
Mupad [F(-1)]	190

Optimal result

Integrand size = 13, antiderivative size = 17

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = -\operatorname{arcsinh}(x) + \sqrt{1+x^2} \arctan(x)$$

[Out] $-\operatorname{arcsinh}(x) + \arctan(x) \cdot (x^2 + 1)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {5050, 221}

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) - \operatorname{arcsinh}(x)$$

[In] $\text{Int}[(x \cdot \text{ArcTan}[x]) / \text{Sqrt}[1 + x^2], x]$

[Out] $-\text{ArcSinh}[x] + \text{Sqrt}[1 + x^2] \cdot \text{ArcTan}[x]$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_.) \cdot (x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2] \cdot (x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

Rule 5050

$\text{Int}[(a_.) + \text{ArcTan}[(c_.) \cdot (x_)] \cdot (b_.)]^{(p_.)} \cdot (x_.) \cdot ((d_.) + (e_.) \cdot (x_.)^2)^{(q_.)}, x_Symbol] \rightarrow \text{Simp}[(d + e \cdot x^2)^{(q+1)} \cdot ((a + b \cdot \text{ArcTan}[c \cdot x])^p / (2 \cdot e \cdot (q+1))), x] - \text{Dist}[b \cdot (p / (2 \cdot c \cdot (q+1))), \text{Int}[(d + e \cdot x^2)^q \cdot (a + b \cdot \text{ArcTan}[c \cdot x])^{(p-1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, q\}, x] \ \&\& \ \text{EqQ}[e, c^2 \cdot d] \ \&\& \ \text{GtQ}[p,$

0] && NeQ[q, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \sqrt{1+x^2} \arctan(x) - \int \frac{1}{\sqrt{1+x^2}} dx \\ &= -\operatorname{arcsinh}(x) + \sqrt{1+x^2} \arctan(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = -\operatorname{arcsinh}(x) + \sqrt{1+x^2} \arctan(x)$$

[In] Integrate[(x*ArcTan[x])/Sqrt[1 + x^2],x]

[Out] -ArcSinh[x] + Sqrt[1 + x^2]*ArcTan[x]

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 54, normalized size of antiderivative = 3.18

method	result	size
default	$\sqrt{(x-i)(x+i)} \arctan(x) + \ln\left(\frac{ix+1}{\sqrt{x^2+1}} - i\right) - \ln\left(\frac{ix+1}{\sqrt{x^2+1}} + i\right)$	54

[In] int(x*arctan(x)/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] ((x-I)*(x+I))^(1/2)*arctan(x)+ln((1+I*x)/(x^2+1)^(1/2)-I)-ln((1+I*x)/(x^2+1)^(1/2)+I)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) + \log\left(-x + \sqrt{x^2+1}\right)$$

[In] integrate(x*arctan(x)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] sqrt(x^2 + 1)*arctan(x) + log(-x + sqrt(x^2 + 1))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(14) = 28$.

Time = 0.44 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.71

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \frac{x^2 \operatorname{atan}(x)}{\sqrt{x^2+1}} - \operatorname{asinh}(x) + \frac{\operatorname{atan}(x)}{\sqrt{x^2+1}}$$

[In] integrate(x*atan(x)/(x**2+1)**(1/2),x)

[Out] x**2*atan(x)/sqrt(x**2 + 1) - asinh(x) + atan(x)/sqrt(x**2 + 1)

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) - \operatorname{arsinh}(x)$$

[In] integrate(x*arctan(x)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 + 1)*arctan(x) - arcsinh(x)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \arctan(x) + \log(-x + \sqrt{x^2+1})$$

[In] integrate(x*arctan(x)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 + 1)*arctan(x) + log(-x + sqrt(x^2 + 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{x \arctan(x)}{\sqrt{1+x^2}} dx = \int \frac{x \operatorname{atan}(x)}{\sqrt{x^2+1}} dx$$

```
[In] int((x*atan(x))/(x^2 + 1)^(1/2),x)
```

```
[Out] int((x*atan(x))/(x^2 + 1)^(1/2), x)
```

3.29 $\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx$

Optimal result	191
Rubi [A] (verified)	191
Mathematica [A] (verified)	193
Maple [F]	193
Fricas [A] (verification not implemented)	193
Sympy [F]	194
Maxima [F]	194
Giac [B] (verification not implemented)	194
Mupad [F(-1)]	195

Optimal result

Integrand size = 17, antiderivative size = 57

$$\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}\arctan(x)}{x} - \operatorname{arctanh}(\sqrt{1-x^2}) + \sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1-x^2}}{\sqrt{2}}\right)$$

[Out] $-\operatorname{arctanh}((-x^2+1)^{(1/2)})+\operatorname{arctanh}(1/2*2^{(1/2)}*(-x^2+1)^{(1/2)})*2^{(1/2)}-\operatorname{arctan}(x)*(-x^2+1)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {270, 5096, 457, 85, 65, 212}

$$\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}\arctan(x)}{x} - \operatorname{arctanh}(\sqrt{1-x^2}) + \sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{1-x^2}}{\sqrt{2}}\right)$$

[In] $\text{Int}[\text{ArcTan}[x]/(x^2*\text{Sqrt}[1-x^2]),x]$

[Out] $-((\text{Sqrt}[1-x^2]*\text{ArcTan}[x])/x) - \text{ArcTanh}[\text{Sqrt}[1-x^2]] + \text{Sqrt}[2]*\text{ArcTanh}[\text{Sqrt}[1-x^2]/\text{Sqrt}[2]]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \text{ :> With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{n_}, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 85

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(a + b*x),
x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(c + d*x), x],
x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[0, p, 1]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*
x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n,
p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 5096

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))*((f_.)*(x_))^(m_.)*((d_.) + (e_.)*(x
_)^2)^(q_.), x_Symbol] := With[{u = IntHide[(f*x)^m*(d + e*x^2)^q, x]}, Dis
t[a + b*ArcTan[c*x], u, x] - Dist[b*c, Int[SimplifyIntegrand[u/(1 + c^2*x^2
), x], x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && ((IGtQ[q, 0] && !(
ILtQ[(m - 1)/2, 0] && GtQ[m + 2*q + 3, 0])) || (IGtQ[(m + 1)/2, 0] && !(IL
tQ[q, 0] && GtQ[m + 2*q + 3, 0])) || (ILtQ[(m + 2*q + 1)/2, 0] && !ILtQ[(m
- 1)/2, 0]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\sqrt{1-x^2} \arctan(x)}{x} + \int \frac{\sqrt{1-x^2}}{x(1+x^2)} dx \\
&= -\frac{\sqrt{1-x^2} \arctan(x)}{x} + \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1-x}}{x(1+x)} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1-x^2} \arctan(x)}{x} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1-xx}} dx, x, x^2 \right) - \text{Subst} \left(\int \frac{1}{\sqrt{1-x}(1+x)} dx, x, x^2 \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{1-x^2} \arctan(x)}{x} + 2\text{Subst}\left(\int \frac{1}{2-x^2} dx, x, \sqrt{1-x^2}\right) \\
&\quad - \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-x^2}\right) \\
&= -\frac{\sqrt{1-x^2} \arctan(x)}{x} - \text{arctanh}\left(\sqrt{1-x^2}\right) + \sqrt{2}\text{arctanh}\left(\frac{\sqrt{1-x^2}}{\sqrt{2}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.35

$$\begin{aligned}
\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx &= -\frac{\sqrt{1-x^2} \arctan(x)}{x} + \log(x) - \frac{\log(1+x^2)}{\sqrt{2}} \\
&\quad + \frac{\log(3-x^2+2\sqrt{2-2x^2})}{\sqrt{2}} - \log\left(1+\sqrt{1-x^2}\right)
\end{aligned}$$

[In] Integrate[ArcTan[x]/(x^2*Sqrt[1-x^2]),x]

[Out] -((Sqrt[1-x^2]*ArcTan[x])/x) + Log[x] - Log[1+x^2]/Sqrt[2] + Log[3-x^2+2*Sqrt[2-2*x^2]]/Sqrt[2] - Log[1+Sqrt[1-x^2]]

Maple [F]

$$\int \frac{\arctan(x)}{x^2\sqrt{-x^2+1}} dx$$

[In] int(arctan(x)/x^2/(-x^2+1)^(1/2),x)

[Out] int(arctan(x)/x^2/(-x^2+1)^(1/2),x)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.42

$$\begin{aligned}
&\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx \\
&= \frac{\sqrt{2}x \log\left(\frac{x^2-2\sqrt{2}\sqrt{-x^2+1}-3}{x^2+1}\right) - x \log(\sqrt{-x^2+1}+1) + x \log(\sqrt{-x^2+1}-1) - 2\sqrt{-x^2+1} \arctan(x)}{2x}
\end{aligned}$$

[In] integrate(arctan(x)/x^2/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] 1/2*(sqrt(2)*x*log((x^2-2*sqrt(2)*sqrt(-x^2+1)-3)/(x^2+1))-x*log(sqrt(-x^2+1)+1)+x*log(sqrt(-x^2+1)-1)-2*sqrt(-x^2+1)*arctan(x))/x

Sympy [F]

$$\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\operatorname{atan}(x)}{x^2\sqrt{-(x-1)(x+1)}} dx$$

[In] integrate(atan(x)/x**2/(-x**2+1)**(1/2),x)

[Out] Integral(atan(x)/(x**2*sqrt(-(x - 1)*(x + 1))), x)

Maxima [F]

$$\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\arctan(x)}{\sqrt{-x^2+1}x^2} dx$$

[In] integrate(arctan(x)/x^2/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(arctan(x)/(sqrt(-x^2 + 1)*x^2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 104 vs. 2(48) = 96.

Time = 0.30 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.82

$$\begin{aligned} \int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx &= \frac{1}{2} \left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right) \arctan(x) \\ &\quad - \frac{1}{2} \sqrt{2} \log \left(\frac{\sqrt{2}-\sqrt{-x^2+1}}{\sqrt{2}+\sqrt{-x^2+1}} \right) \\ &\quad - \frac{1}{2} \log(\sqrt{-x^2+1}+1) + \frac{1}{2} \log(-\sqrt{-x^2+1}+1) \end{aligned}$$

[In] integrate(arctan(x)/x^2/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/2*(x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)*arctan(x) - 1/2*sqrt(2)*log((sqrt(2) - sqrt(-x^2 + 1))/(sqrt(2) + sqrt(-x^2 + 1))) - 1/2*log(sqrt(-x^2 + 1) + 1) + 1/2*log(-sqrt(-x^2 + 1) + 1)

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\operatorname{atan}(x)}{x^2\sqrt{1-x^2}} dx$$

```
[In] int(atan(x)/(x^2*(1 - x^2)^(1/2)),x)
```

```
[Out] int(atan(x)/(x^2*(1 - x^2)^(1/2)), x)
```

3.30 $\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx$

Optimal result	196
Rubi [A] (verified)	196
Mathematica [A] (verified)	198
Maple [F]	198
Fricas [A] (verification not implemented)	198
Sympy [F]	199
Maxima [C] (verification not implemented)	199
Giac [B] (verification not implemented)	200
Mupad [B] (verification not implemented)	200

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\arcsin(x) - \sqrt{1-x^2} \arctan(x) + \sqrt{2} \arctan\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)$$

[Out] -arcsin(x)+arctan(x*2^(1/2)/(-x^2+1)^(1/2))*2^(1/2)-arctan(x)*(-x^2+1)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5094, 399, 222, 385, 209}

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\arcsin(x) - \sqrt{1-x^2} \arctan(x) + \sqrt{2} \arctan\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)$$

[In] Int[(x*ArcTan[x])/Sqrt[1 - x^2],x]

[Out] -ArcSin[x] - Sqrt[1 - x^2]*ArcTan[x] + Sqrt[2]*ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 222

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 399

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]

Rule 5094

Int[((a_.) + ArcTan[(c_.)*(x_)])*(b_.)*(x_)*((d_.) + (e_.)*(x_)^2)^(q_.), x_Symbol] := Simp[(d + e*x^2)^(q + 1)*((a + b*ArcTan[c*x])/(2*e*(q + 1))), x] - Dist[b*(c/(2*e*(q + 1))), Int[(d + e*x^2)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\sqrt{1-x^2} \arctan(x) + \int \frac{\sqrt{1-x^2}}{1+x^2} dx \\
 &= -\sqrt{1-x^2} \arctan(x) + 2 \int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx - \int \frac{1}{\sqrt{1-x^2}} dx \\
 &= -\arcsin(x) - \sqrt{1-x^2} \arctan(x) + 2 \text{Subst} \left(\int \frac{1}{1+2x^2} dx, x, \frac{x}{\sqrt{1-x^2}} \right) \\
 &= -\arcsin(x) - \sqrt{1-x^2} \arctan(x) + \sqrt{2} \arctan \left(\frac{\sqrt{2}x}{\sqrt{1-x^2}} \right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\arcsin(x) - \sqrt{1-x^2} \arctan(x) + \sqrt{2} \arctan\left(\frac{\sqrt{2}x}{\sqrt{1-x^2}}\right)$$

[In] Integrate[(x*ArcTan[x])/Sqrt[1 - x^2],x]

[Out] -ArcSin[x] - Sqrt[1 - x^2]*ArcTan[x] + Sqrt[2]*ArcTan[(Sqrt[2]*x)/Sqrt[1 - x^2]]

Maple [F]

$$\int \frac{x \arctan(x)}{\sqrt{-x^2+1}} dx$$

[In] int(x*arctan(x)/(-x^2+1)^(1/2),x)

[Out] int(x*arctan(x)/(-x^2+1)^(1/2),x)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.53

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\frac{1}{2} \sqrt{2} \arctan\left(\frac{\sqrt{2}(3x^2-1)\sqrt{-x^2+1}}{4(x^3-x)}\right) - \sqrt{-x^2+1} \arctan(x) + \arctan\left(\frac{\sqrt{-x^2+1}x}{x^2-1}\right)$$

[In] integrate(x*arctan(x)/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] -1/2*sqrt(2)*arctan(1/4*sqrt(2)*(3*x^2 - 1)*sqrt(-x^2 + 1)/(x^3 - x)) - sqrt(-x^2 + 1)*arctan(x) + arctan(sqrt(-x^2 + 1)*x/(x^2 - 1))

SymPy [F]

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = \int \frac{x \operatorname{atan}(x)}{\sqrt{-(x-1)(x+1)}} dx$$

```
[In] integrate(x*atan(x)/(-x**2+1)**(1/2),x)
```

```
[Out] Integral(x*atan(x)/sqrt(-(x - 1)*(x + 1)), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 387, normalized size of antiderivative = 8.60

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\sqrt{-x^2+1} \arctan(x) + \frac{1}{2} \sqrt{2} \arctan\left(\frac{(x^4 + |ix+1|^4 - 2(x^2-1)|ix+1|^2 + 2x^2+1)^{\frac{1}{4}} \sin\left(\frac{1}{2} \arctan\left(\frac{2x}{|ix+1|^2}, -\frac{x^2-|ix+1|^2-1}{|ix+1|^2}\right)\right)}{|ix+1|}\right) + \frac{1}{2} \sqrt{2} \arctan\left(\frac{(x^4 + |ix-1|^4 - 2(x^2-1)|ix-1|^2 + 2x^2+1)^{\frac{1}{4}} \sin\left(\frac{1}{2} \arctan\left(\frac{2x}{|ix-1|^2}, -\frac{x^2-|ix-1|^2-1}{|ix-1|^2}\right)\right)}{|ix-1|}\right) - \arctan\left(x, \sqrt{-x^2+1}\right)$$

```
[In] integrate(x*arctan(x)/(-x^2+1)^(1/2),x, algorithm="maxima")
```

```
[Out] -sqrt(-x^2 + 1)*arctan(x) + 1/2*sqrt(2)*arctan2(((x^4 + abs(I*x + 1)^4 - 2*(x^2 - 1)*abs(I*x + 1)^2 + 2*x^2 + 1)^(1/4)*sin(1/2*arctan2(2*x/abs(I*x + 1)^2, -(x^2 - abs(I*x + 1)^2 - 1)/abs(I*x + 1)^2)) + x)/abs(I*x + 1), ((x^4 + abs(I*x + 1)^4 - 2*(x^2 - 1)*abs(I*x + 1)^2 + 2*x^2 + 1)^(1/4)*cos(1/2*arctan2(2*x/abs(I*x + 1)^2, -(x^2 - abs(I*x + 1)^2 - 1)/abs(I*x + 1)^2)) + 1)/abs(I*x + 1)) + 1/2*sqrt(2)*arctan2(((x^4 + abs(I*x - 1)^4 - 2*(x^2 - 1)*abs(I*x - 1)^2 + 2*x^2 + 1)^(1/4)*sin(1/2*arctan2(2*x/abs(I*x - 1)^2, -(x^2 - abs(I*x - 1)^2 - 1)/abs(I*x - 1)^2)) + x)/abs(I*x - 1), ((x^4 + abs(I*x - 1)^4 - 2*(x^2 - 1)*abs(I*x - 1)^2 + 2*x^2 + 1)^(1/4)*cos(1/2*arctan2(2*x/abs(I*x - 1)^2, -(x^2 - abs(I*x - 1)^2 - 1)/abs(I*x - 1)^2)) + 1)/abs(I*x - 1)) - arctan2(x, sqrt(-x^2 + 1))
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 108 vs. $2(37) = 74$.

Time = 0.28 (sec) , antiderivative size = 108, normalized size of antiderivative = 2.40

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = -\frac{1}{2} \pi \operatorname{sgn}(x) + \frac{1}{2} \sqrt{2} \left(\pi \operatorname{sgn}(x) + 2 \arctan \left(-\frac{\sqrt{2}x \left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{4(\sqrt{-x^2+1}-1)} \right) \right) - \sqrt{-x^2+1} \arctan(x) - \arctan \left(-\frac{x \left(\frac{(\sqrt{-x^2+1}-1)^2}{x^2} - 1 \right)}{2(\sqrt{-x^2+1}-1)} \right)$$

[In] integrate(x*arctan(x)/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] -1/2*pi*sgn(x) + 1/2*sqrt(2)*(pi*sgn(x) + 2*arctan(-1/4*sqrt(2)*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))) - sqrt(-x^2 + 1)*arctan(x) - arctan(-1/2*x*((sqrt(-x^2 + 1) - 1)^2/x^2 - 1)/(sqrt(-x^2 + 1) - 1))

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.82

$$\int \frac{x \arctan(x)}{\sqrt{1-x^2}} dx = \sqrt{2} \operatorname{atan} \left(\frac{\sqrt{2}x}{\sqrt{1-x^2}} \right) - \operatorname{atan}(x) \sqrt{1-x^2} - \operatorname{asin}(x)$$

[In] int((x*atan(x))/(1 - x^2)^(1/2),x)

[Out] 2^(1/2)*atan((2^(1/2)*x)/(1 - x^2)^(1/2)) - atan(x)*(1 - x^2)^(1/2) - asin(x)

3.31 $\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx$

Optimal result	201
Rubi [A] (verified)	201
Mathematica [A] (verified)	202
Maple [C] (verified)	203
Fricas [A] (verification not implemented)	203
Sympy [A] (verification not implemented)	203
Maxima [A] (verification not implemented)	204
Giac [B] (verification not implemented)	204
Mupad [F(-1)]	204

Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{1+x^2} \arctan(x)}{x} - \operatorname{arctanh}(\sqrt{1+x^2})$$

[Out] $-\operatorname{arctanh}((x^2+1)^{(1/2)})-\arctan(x)*(x^2+1)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5064, 272, 65, 213}

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{x^2+1} \arctan(x)}{x} - \operatorname{arctanh}(\sqrt{x^2+1})$$

[In] $\text{Int}[\text{ArcTan}[x]/(x^2*\text{Sqrt}[1+x^2]),x]$

[Out] $-((\text{Sqrt}[1+x^2]*\text{ArcTan}[x])/x) - \text{ArcTanh}[\text{Sqrt}[1+x^2]]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b)^n, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 213

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 5064

```
Int[((a_) + ArcTan[(c_)*(x_)])*(b_)^(p_)*((f_)*(x_)^(m_))*((d_) + (e_)*(x_)^2)^(q_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^2)^(q + 1)*((a + b*ArcTan[c*x])^p/(d*f*(m + 1))), x] - Dist[b*c*(p/(f*(m + 1))), Int[(f*x)^(m + 1)*(d + e*x^2)^q*(a + b*ArcTan[c*x])^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, q}, x] && EqQ[e, c^2*d] && EqQ[m + 2*q + 3, 0] && GtQ[p, 0] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sqrt{1+x^2} \arctan(x)}{x} + \int \frac{1}{x\sqrt{1+x^2}} dx \\
 &= -\frac{\sqrt{1+x^2} \arctan(x)}{x} + \frac{1}{2} \text{Subst}\left(\int \frac{1}{x\sqrt{1+x}} dx, x, x^2\right) \\
 &= -\frac{\sqrt{1+x^2} \arctan(x)}{x} + \text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+x^2}\right) \\
 &= -\frac{\sqrt{1+x^2} \arctan(x)}{x} - \text{arctanh}\left(\sqrt{1+x^2}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.14

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{1+x^2} \arctan(x)}{x} + \log(x) - \log\left(1 + \sqrt{1+x^2}\right)$$

```
[In] Integrate[ArcTan[x]/(x^2*Sqrt[1 + x^2]),x]
```

```
[Out] -((Sqrt[1 + x^2]*ArcTan[x])/x) + Log[x] - Log[1 + Sqrt[1 + x^2]]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.68 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.93

method	result	size
default	$-\frac{\sqrt{(x-i)(x+i)} \arctan(x)}{x} + \ln\left(\frac{ix+1}{\sqrt{x^2+1}} - 1\right) - \ln\left(\frac{ix+1}{\sqrt{x^2+1}} + 1\right)$	56

[In] `int(arctan(x)/x^2/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-\left((x-I)*(x+I)\right)^{(1/2)}*\arctan(x)/x+\ln\left(\frac{(1+I*x)}{(x^2+1)^{(1/2)}-1}-\ln\left(\frac{(1+I*x)}{(x^2+1)^{(1/2)}+1}\right)\right)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.62

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\frac{x \log(-x + \sqrt{x^2 + 1} + 1) - x \log(-x + \sqrt{x^2 + 1} - 1) + \sqrt{x^2 + 1} \arctan(x)}{x}$$

[In] `integrate(arctan(x)/x^2/(x^2+1)^(1/2),x, algorithm="fricas")`

[Out] $-(x*\log(-x + \text{sqrt}(x^2 + 1) + 1) - x*\log(-x + \text{sqrt}(x^2 + 1) - 1) + \text{sqrt}(x^2 + 1)*\arctan(x))/x$

Sympy [A] (verification not implemented)

Time = 3.83 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\operatorname{asinh}\left(\frac{1}{x}\right) - \frac{\sqrt{x^2+1} \operatorname{atan}(x)}{x}$$

[In] `integrate(atan(x)/x**2/(x**2+1)**(1/2),x)`

[Out] $-\operatorname{asinh}(1/x) - \text{sqrt}(x**2 + 1)*\operatorname{atan}(x)/x$

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{x^2+1}\arctan(x)}{x} - \operatorname{arsinh}\left(\frac{1}{|x|}\right)$$

[In] integrate(arctan(x)/x^2/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(x^2 + 1)*arctan(x)/x - arcsinh(1/abs(x))

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(25) = 50.

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.86

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = \frac{2\arctan(x)}{(x-\sqrt{x^2+1})^2-1} + \arctan(x) - \log\left(\left|-x+\sqrt{x^2+1}+1\right|\right) + \log\left(\left|-x+\sqrt{x^2+1}-1\right|\right)$$

[In] integrate(arctan(x)/x^2/(x^2+1)^(1/2),x, algorithm="giac")

[Out] 2*arctan(x)/((x - sqrt(x^2 + 1))^2 - 1) + arctan(x) - log(abs(-x + sqrt(x^2 + 1) + 1)) + log(abs(-x + sqrt(x^2 + 1) - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\arctan(x)}{x^2\sqrt{1+x^2}} dx = \int \frac{\operatorname{atan}(x)}{x^2\sqrt{x^2+1}} dx$$

[In] int(atan(x)/(x^2*(x^2 + 1)^(1/2)),x)

[Out] int(atan(x)/(x^2*(x^2 + 1)^(1/2)), x)

3.32 $\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx$

Optimal result	205
Rubi [A] (verified)	205
Mathematica [A] (verified)	206
Maple [A] (verified)	206
Fricas [A] (verification not implemented)	206
Sympy [F]	207
Maxima [A] (verification not implemented)	207
Giac [B] (verification not implemented)	207
Mupad [F(-1)]	207

Optimal result

Integrand size = 17, antiderivative size = 21

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2}\arcsin(x)}{x} + \log(x)$$

[Out] $\ln(x) - \arcsin(x) * (-x^2 + 1)^{(1/2)} / x$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {4771, 29}

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = \log(x) - \frac{\sqrt{1-x^2}\arcsin(x)}{x}$$

[In] $\text{Int}[\text{ArcSin}[x]/(x^2*\text{Sqrt}[1-x^2]),x]$

[Out] $-((\text{Sqrt}[1-x^2]*\text{ArcSin}[x])/x) + \text{Log}[x]$

Rule 29

$\text{Int}[(x_)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[x], x]$

Rule 4771

$\text{Int}[(a_. + \text{ArcSin}[c_.*(x_)]*(b_.))^{(n_.)}*((f_.)*(x_))^{(m_.)}*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(f*x)^{(m+1)}*(d + e*x^2)^{(p+1)}*((a + b*\text{ArcSin}[c*x])^n/(d*f*(m+1))), x] - \text{Dist}[b*c*(n/(f*(m+1)))*\text{Simp}[(d + e*x^2)^p/(1 - c^2*x^2)^p], \text{Int}[(f*x)^{(m+1)}*(1 - c^2*x^2)^{(p+1/2)}*(a + b*\text{ArcSin}[c*x])^{(n-1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x \ \&\& \ \text{EqQ}[c^2$

*d + e, 0] && GtQ[n, 0] && EqQ[m + 2*p + 3, 0] && NeQ[m, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\sqrt{1-x^2} \arcsin(x)}{x} + \int \frac{1}{x} dx \\ &= -\frac{\sqrt{1-x^2} \arcsin(x)}{x} + \log(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\arcsin(x)}{x^2 \sqrt{1-x^2}} dx = -\frac{\sqrt{1-x^2} \arcsin(x)}{x} + \log(x)$$

[In] Integrate[ArcSin[x]/(x^2*Sqrt[1 - x^2]),x]

[Out] -((Sqrt[1 - x^2]*ArcSin[x])/x) + Log[x]

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.95

method	result	size
default	$\ln(x) - \frac{\arcsin(x)\sqrt{-x^2+1}}{x}$	20

[In] int(arcsin(x)/x^2/(-x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] ln(x)-arcsin(x)*(-x^2+1)^(1/2)/x

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \frac{\arcsin(x)}{x^2 \sqrt{1-x^2}} dx = \frac{x \log(x) - \sqrt{-x^2+1} \arcsin(x)}{x}$$

[In] integrate(arcsin(x)/x^2/(-x^2+1)^(1/2),x, algorithm="fricas")

[Out] (x*log(x) - sqrt(-x^2 + 1)*arcsin(x))/x

Sympy [F]

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\operatorname{asin}(x)}{x^2\sqrt{-(x-1)(x+1)}} dx$$

[In] integrate(asin(x)/x**2/(-x**2+1)**(1/2),x)

[Out] Integral(asin(x)/(x**2*sqrt(-(x - 1)*(x + 1))), x)

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = -\frac{\sqrt{-x^2+1}\arcsin(x)}{x} + \log(x)$$

[In] integrate(arcsin(x)/x^2/(-x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(-x^2 + 1)*arcsin(x)/x + log(x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(19) = 38.

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.90

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = \frac{1}{2} \left(\frac{x}{\sqrt{-x^2+1}-1} - \frac{\sqrt{-x^2+1}-1}{x} \right) \arcsin(x) + \log(|x|)$$

[In] integrate(arcsin(x)/x^2/(-x^2+1)^(1/2),x, algorithm="giac")

[Out] 1/2*(x/(sqrt(-x^2 + 1) - 1) - (sqrt(-x^2 + 1) - 1)/x)*arcsin(x) + log(abs(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\arcsin(x)}{x^2\sqrt{1-x^2}} dx = \int \frac{\operatorname{asin}(x)}{x^2\sqrt{1-x^2}} dx$$

[In] int(asin(x)/(x^2*(1 - x^2)^(1/2)),x)

[Out] int(asin(x)/(x^2*(1 - x^2)^(1/2)), x)

3.33 $\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx$

Optimal result	208
Rubi [A] (verified)	208
Mathematica [A] (verified)	210
Maple [C] (warning: unable to verify)	210
Fricas [A] (verification not implemented)	210
Sympy [A] (verification not implemented)	211
Maxima [A] (verification not implemented)	211
Giac [A] (verification not implemented)	211
Mupad [F(-1)]	211

Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = -\sqrt{-1+x^2} + \arctan(\sqrt{-1+x^2}) + \sqrt{-1+x^2} \log(x)$$

[Out] $\arctan((x^2-1)^{(1/2)})-(x^2-1)^{(1/2)}+\ln(x)*(x^2-1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {2376, 272, 52, 65, 209}

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \arctan(\sqrt{x^2-1}) - \sqrt{x^2-1} + \sqrt{x^2-1} \log(x)$$

[In] $\text{Int}[(x*\text{Log}[x])/Sqrt[-1 + x^2], x]$

[Out] $-Sqrt[-1 + x^2] + \text{ArcTan}[Sqrt[-1 + x^2]] + Sqrt[-1 + x^2]*\text{Log}[x]$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 2376

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.)*((d_) +
(e_.)*(x_)^(r_))^(q_.), x_Symbol] := Simp[f^m*(d + e*x^r)^(q + 1)*((a + b*L
og[c*x^n])^p/(e*r*(q + 1))), x] - Dist[b*f^m*n*(p/(e*r*(q + 1))), Int[(d +
e*x^r)^(q + 1)*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d,
e, f, m, n, q, r}, x] && EqQ[m, r - 1] && IGtQ[p, 0] && (IntegerQ[m] || Gt
Q[f, 0]) && NeQ[r, n] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \sqrt{-1+x^2} \log(x) - \int \frac{\sqrt{-1+x^2}}{x} dx \\
&= \sqrt{-1+x^2} \log(x) - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{-1+x}}{x} dx, x, x^2 \right) \\
&= -\sqrt{-1+x^2} + \sqrt{-1+x^2} \log(x) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{-1+xx}} dx, x, x^2 \right) \\
&= -\sqrt{-1+x^2} + \sqrt{-1+x^2} \log(x) + \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x^2} \right) \\
&= -\sqrt{-1+x^2} + \arctan \left(\sqrt{-1+x^2} \right) + \sqrt{-1+x^2} \log(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = -\arctan\left(\frac{1}{\sqrt{-1+x^2}}\right) + \sqrt{-1+x^2}(-1 + \log(x))$$

[In] Integrate[(x*Log[x])/Sqrt[-1 + x^2],x]

[Out] -ArcTan[1/Sqrt[-1 + x^2]] + Sqrt[-1 + x^2]*(-1 + Log[x])

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.08 (sec) , antiderivative size = 119, normalized size of antiderivative = 3.50

method	result
meijerg	$-\frac{\sqrt{-\operatorname{signum}(x^2-1)}(2-2\sqrt{-x^2+1})}{4\sqrt{\operatorname{signum}(x^2-1)}} + \frac{\sqrt{-\operatorname{signum}(x^2-1)}\ln(x)(2-2\sqrt{-x^2+1})}{2\sqrt{\operatorname{signum}(x^2-1)}} + \frac{\sqrt{-\operatorname{signum}(x^2-1)}(-16+16\sqrt{-x^2+1}-32\ln(1/2+1/2\sqrt{-x^2+1}))}{32\sqrt{\operatorname{signum}(x^2-1)}}$

[In] int(x*ln(x)/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/4/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*(2-2*(-x^2+1)^(1/2))+1/2/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*ln(x)*(2-2*(-x^2+1)^(1/2))+1/32/signum(x^2-1)^(1/2)*(-signum(x^2-1))^(1/2)*(-16+16*(-x^2+1)^(1/2)-32*ln(1/2+1/2*(-x^2+1)^(1/2)))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1}(\log(x) - 1) + 2 \arctan\left(-x + \sqrt{x^2-1}\right)$$

[In] integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="fricas")

[Out] sqrt(x^2 - 1)*(log(x) - 1) + 2*arctan(-x + sqrt(x^2 - 1))

Sympy [A] (verification not implemented)

Time = 1.37 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.85

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \left\{ \sqrt{x^2-1} - \operatorname{acos}\left(\frac{1}{x}\right) \quad \text{for } x > -1 \wedge x < 1 \right.$$

[In] integrate(x*ln(x)/(x**2-1)**(1/2),x)

[Out] sqrt(x**2 - 1)*log(x) - Piecewise((sqrt(x**2 - 1) - acos(1/x), (x > -1) & (x < 1)))

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.79

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \sqrt{x^2-1} - \arcsin\left(\frac{1}{|x|}\right)$$

[In] integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 - 1)*log(x) - sqrt(x^2 - 1) - arcsin(1/abs(x))

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.82

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \log(x) - \sqrt{x^2-1} + \arctan\left(\sqrt{x^2-1}\right)$$

[In] integrate(x*log(x)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 - 1)*log(x) - sqrt(x^2 - 1) + arctan(sqrt(x^2 - 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(x)}{\sqrt{-1+x^2}} dx = \int \frac{x \ln(x)}{\sqrt{x^2-1}} dx$$

[In] int((x*log(x))/(x^2 - 1)^(1/2),x)

[Out] int((x*log(x))/(x^2 - 1)^(1/2), x)

3.34 $\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx$

Optimal result	212
Rubi [A] (verified)	212
Mathematica [A] (verified)	213
Maple [A] (verified)	213
Fricas [A] (verification not implemented)	214
Sympy [A] (verification not implemented)	214
Maxima [A] (verification not implemented)	214
Giac [A] (verification not implemented)	214
Mupad [F(-1)]	215

Optimal result

Integrand size = 15, antiderivative size = 33

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{1+x^2}}{x} + \operatorname{arcsinh}(x) - \frac{\sqrt{1+x^2}\log(x)}{x}$$

[Out] $\operatorname{arcsinh}(x) - (x^2+1)^{(1/2)}/x - \ln(x) * (x^2+1)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2373, 283, 221}

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = \operatorname{arcsinh}(x) - \frac{\sqrt{x^2+1}}{x} - \frac{\sqrt{x^2+1}\log(x)}{x}$$

[In] $\operatorname{Int}[\operatorname{Log}[x]/(x^2*\operatorname{Sqrt}[1+x^2]),x]$

[Out] $-(\operatorname{Sqrt}[1+x^2]/x) + \operatorname{ArcSinh}[x] - (\operatorname{Sqrt}[1+x^2]*\operatorname{Log}[x])/x$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_)+(b_)*(x_)^2], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 283

$\operatorname{Int}[((c_)*(x_))^{(m_)}*((a_)+(b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \operatorname{Simp}[(c*x)^{(m+1)}*((a+b*x^n)^p/(c*(m+1))), x] - \operatorname{Dist}[b*n*(p/(c^n*(m+1))), \operatorname{Int}[(c*x)^{(m+n)}*(a+b*x^n)^{(p-1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{GtQ}[p, 0] \ \&\& \operatorname{LtQ}[m, -1] \ \&\& \operatorname{!ILtQ}[(m+n*p+n+1)/n, 0] \ \&\& \operatorname{IntBi}$

nomialQ[a, b, c, n, m, p, x]

Rule 2373

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))*((f_.)*(x_)^(m_.))*((d_) + (e_.)*
(x_)^(r_.))^(q_), x_Symbol] := Simp[(f*x)^(m + 1)*(d + e*x^r)^(q + 1)*((a +
b*Log[c*x^n])/(d*f*(m + 1))), x] - Dist[b*(n/(d*(m + 1))), Int[(f*x)^m*(d
+ e*x^r)^(q + 1), x], x] /; FreeQ[{a, b, c, d, e, f, m, n, q, r}, x] && EqQ
[m + r*(q + 1) + 1, 0] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\sqrt{1+x^2} \log(x)}{x} + \int \frac{\sqrt{1+x^2}}{x^2} dx \\ &= -\frac{\sqrt{1+x^2}}{x} - \frac{\sqrt{1+x^2} \log(x)}{x} + \int \frac{1}{\sqrt{1+x^2}} dx \\ &= -\frac{\sqrt{1+x^2}}{x} + \operatorname{arcsinh}(x) - \frac{\sqrt{1+x^2} \log(x)}{x} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.64

$$\int \frac{\log(x)}{x^2 \sqrt{1+x^2}} dx = \operatorname{arcsinh}(x) - \frac{\sqrt{1+x^2}(1+\log(x))}{x}$$

```
[In] Integrate[Log[x]/(x^2*Sqrt[1 + x^2]),x]
```

```
[Out] ArcSinh[x] - (Sqrt[1 + x^2]*(1 + Log[x]))/x
```

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

method	result	size
meijerg	$\operatorname{arcsinh}(x) + \frac{-\ln(x)\sqrt{x^2+1}-\sqrt{x^2+1}}{x}$	29

```
[In] int(ln(x)/x^2/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] arcsinh(x)+(-ln(x)*(x^2+1)^(1/2)-(x^2+1)^(1/2))/x
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = -\frac{x \log(-x + \sqrt{x^2+1}) + \sqrt{x^2+1}(\log(x) + 1) + x}{x}$$

[In] integrate(log(x)/x^2/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] -(x*log(-x + sqrt(x^2 + 1)) + sqrt(x^2 + 1)*(log(x) + 1) + x)/x

Sympy [A] (verification not implemented)

Time = 2.95 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = \operatorname{asinh}(x) - \frac{\sqrt{x^2+1} \log(x)}{x} - \frac{\sqrt{x^2+1}}{x}$$

[In] integrate(ln(x)/x**2/(x**2+1)**(1/2),x)

[Out] asinh(x) - sqrt(x**2 + 1)*log(x)/x - sqrt(x**2 + 1)/x

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = -\frac{\sqrt{x^2+1} \log(x)}{x} - \frac{\sqrt{x^2+1}}{x} + \operatorname{arsinh}(x)$$

[In] integrate(log(x)/x^2/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] -sqrt(x^2 + 1)*log(x)/x - sqrt(x^2 + 1)/x + arcsinh(x)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.76

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = \frac{2 \log(x)}{(x - \sqrt{x^2+1})^2 - 1} + \frac{2}{(x - \sqrt{x^2+1})^2 - 1} - \log(-x + \sqrt{x^2+1}) + \log(|x|)$$

[In] integrate(log(x)/x^2/(x^2+1)^(1/2),x, algorithm="giac")

[Out] 2*log(x)/((x - sqrt(x^2 + 1))^2 - 1) + 2/((x - sqrt(x^2 + 1))^2 - 1) - log(-x + sqrt(x^2 + 1)) + log(abs(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\log(x)}{x^2\sqrt{1+x^2}} dx = \int \frac{\ln(x)}{x^2\sqrt{x^2+1}} dx$$

```
[In] int(log(x)/(x^2*(x^2 + 1)^(1/2)),x)
```

```
[Out] int(log(x)/(x^2*(x^2 + 1)^(1/2)), x)
```

3.35 $\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx$

Optimal result	216
Rubi [A] (verified)	216
Mathematica [A] (verified)	217
Maple [C] (warning: unable to verify)	217
Fricas [A] (verification not implemented)	217
Sympy [A] (verification not implemented)	218
Maxima [A] (verification not implemented)	218
Giac [A] (verification not implemented)	218
Mupad [F(-1)]	218

Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{-1+x^2} \sec^{-1}(x) - \frac{x \log(x)}{\sqrt{x^2}}$$

[Out] $-x*\ln(x)/(x^2)^{(1/2)}+\operatorname{arcsec}(x)*(x^2-1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {5344, 29}

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \sec^{-1}(x) - \frac{x \log(x)}{\sqrt{x^2}}$$

[In] $\operatorname{Int}[(x*\operatorname{ArcSec}[x])/Sqrt[-1+x^2],x]$

[Out] $Sqrt[-1+x^2]*\operatorname{ArcSec}[x] - (x*\operatorname{Log}[x])/Sqrt[x^2]$

Rule 29

$\operatorname{Int}[(x_)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Log}[x], x]$

Rule 5344

$\operatorname{Int}[(a_.) + \operatorname{ArcSec}[(c_.)*(x_)]*(b_.)]*(x_)*((d_.) + (e_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(d + e*x^2)^{(p+1)}*((a + b*\operatorname{ArcSec}[c*x])/ (2*e*(p+1))), x] - \operatorname{Dist}[b*c*(x/(2*e*(p+1)*Sqrt[c^2*x^2])), \operatorname{Int}[(d + e*x^2)^{(p+1)}/(x*Sqrt[c^2*x^2 - 1]), x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, p\}, x \&\& \operatorname{NeQ}[p, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \sqrt{-1+x^2} \sec^{-1}(x) - \frac{x \int \frac{1}{x} dx}{\sqrt{x^2}} \\ &= \sqrt{-1+x^2} \sec^{-1}(x) - \frac{x \log(x)}{\sqrt{x^2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.40

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \frac{(-1+x^2) \sec^{-1}(x) - \sqrt{1-\frac{1}{x^2}} x \log(x)}{\sqrt{-1+x^2}}$$

[In] Integrate[(x*ArcSec[x])/Sqrt[-1 + x^2],x]

[Out] ((-1 + x^2)*ArcSec[x] - Sqrt[1 - x^(-2)]*x*Log[x])/Sqrt[-1 + x^2]

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.39 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.36

method	result	size
default	$\text{csgn}\left(x\sqrt{1-\frac{1}{x^2}}\right) \left(\text{arcsec}(x) x\sqrt{\frac{x^2-1}{x^2}} + \ln\left(\frac{1}{x}\right)\right)$	34

[In] int(x*arcsec(x)/(x^2-1)^(1/2),x,method=_RETURNVERBOSE)

[Out] csgn(x*(1-1/x^2)^(1/2))*(arcsec(x)*x*((x^2-1)/x^2)^(1/2)+ln(1/x))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \text{arcsec}(x) - \log(x)$$

[In] integrate(x*arcsec(x)/(x^2-1)^(1/2),x, algorithm="fricas")

[Out] sqrt(x^2 - 1)*arcsec(x) - log(x)

Sympy [A] (verification not implemented)

Time = 16.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \operatorname{asec}(x) - \left\{ -\log\left(\frac{1}{x}\right) \quad \text{for } x > -1 \wedge x < 1 \right.$$

[In] integrate(x*asec(x)/(x**2-1)**(1/2),x)

[Out] sqrt(x**2 - 1)*asec(x) - Piecewise((-log(1/x), (x > -1) & (x < 1)))

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.60

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \operatorname{arcsec}(x) - \log(x)$$

[In] integrate(x*arcsec(x)/(x^2-1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 - 1)*arcsec(x) - log(x)

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \sqrt{x^2-1} \arccos\left(\frac{1}{x}\right) - \frac{\log(|x|)}{\operatorname{sgn}(x)}$$

[In] integrate(x*arcsec(x)/(x^2-1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 - 1)*arccos(1/x) - log(abs(x))/sgn(x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \sec^{-1}(x)}{\sqrt{-1+x^2}} dx = \int \frac{x \operatorname{acos}\left(\frac{1}{x}\right)}{\sqrt{x^2-1}} dx$$

[In] int((x*acos(1/x))/(x^2 - 1)^(1/2),x)

[Out] int((x*acos(1/x))/(x^2 - 1)^(1/2), x)

3.36 $\int \frac{x \log(x)}{\sqrt{1+x^2}} dx$

Optimal result	219
Rubi [A] (verified)	219
Mathematica [A] (verified)	221
Maple [A] (verified)	221
Fricas [A] (verification not implemented)	221
Sympy [A] (verification not implemented)	222
Maxima [A] (verification not implemented)	222
Giac [A] (verification not implemented)	222
Mupad [F(-1)]	223

Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = -\sqrt{1+x^2} + \operatorname{arctanh}(\sqrt{1+x^2}) + \sqrt{1+x^2} \log(x)$$

[Out] $\operatorname{arctanh}((x^2+1)^{(1/2)})-(x^2+1)^{(1/2)}+\ln(x)*(x^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {2376, 272, 52, 65, 213}

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = \operatorname{arctanh}(\sqrt{x^2+1}) - \sqrt{x^2+1} + \sqrt{x^2+1} \log(x)$$

[In] $\text{Int}[(x*\text{Log}[x])/Sqrt[1+x^2],x]$

[Out] $-\text{Sqrt}[1+x^2] + \text{ArcTanh}[Sqrt[1+x^2]] + \text{Sqrt}[1+x^2]*\text{Log}[x]$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+n+1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m+n+1))], \text{Int}[(a + b*x)^m*(c + d*x)^{(n-1)}, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m+n+1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m-n, 0]))) \ \&\& \ !\text{ILtQ}[m+n+2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 213

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 2376

```
Int[((a_.) + Log[(c_.)*(x_)^(n_.)]*(b_.))^(p_.)*((f_.)*(x_)^(m_.)*((d_) +
(e_.)*(x_)^(r_))^(q_.), x_Symbol] := Simp[f^m*(d + e*x^r)^(q + 1)*((a + b*L
og[c*x^n])^p/(e*r*(q + 1))), x] - Dist[b*f^m*n*(p/(e*r*(q + 1))), Int[(d +
e*x^r)^(q + 1)*((a + b*Log[c*x^n])^(p - 1)/x), x], x] /; FreeQ[{a, b, c, d,
e, f, m, n, q, r}, x] && EqQ[m, r - 1] && IGtQ[p, 0] && (IntegerQ[m] || Gt
Q[f, 0]) && NeQ[r, n] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \sqrt{1+x^2} \log(x) - \int \frac{\sqrt{1+x^2}}{x} dx \\
&= \sqrt{1+x^2} \log(x) - \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{1+x}}{x} dx, x, x^2 \right) \\
&= -\sqrt{1+x^2} + \sqrt{1+x^2} \log(x) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1+x}} dx, x, x^2 \right) \\
&= -\sqrt{1+x^2} + \sqrt{1+x^2} \log(x) - \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+x^2} \right) \\
&= -\sqrt{1+x^2} + \text{arctanh}(\sqrt{1+x^2}) + \sqrt{1+x^2} \log(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.18

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = -\sqrt{1+x^2} - \log(x) + \sqrt{1+x^2} \log(x) + \log\left(1 + \sqrt{1+x^2}\right)$$

[In] Integrate[(x*Log[x])/Sqrt[1 + x^2],x]

[Out] -Sqrt[1 + x^2] - Log[x] + Sqrt[1 + x^2]*Log[x] + Log[1 + Sqrt[1 + x^2]]

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.15

method	result	size
meijerg	$1 - \sqrt{x^2 + 1} + \frac{\ln(x)(-2+2\sqrt{x^2+1})}{2} + \ln\left(\frac{1}{2} + \frac{\sqrt{x^2+1}}{2}\right)$	39

[In] int(x*ln(x)/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1-(x^2+1)^(1/2)+1/2*ln(x)*(-2+2*(x^2+1)^(1/2))+ln(1/2+1/2*(x^2+1)^(1/2))

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1}(\log(x) - 1) + \log(-x + \sqrt{x^2+1} + 1) - \log(-x + \sqrt{x^2+1} - 1)$$

[In] integrate(x*log(x)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] sqrt(x^2 + 1)*(log(x) - 1) + log(-x + sqrt(x^2 + 1) + 1) - log(-x + sqrt(x^2 + 1) - 1)

Sympy [A] (verification not implemented)

Time = 2.60 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = -\frac{x}{\sqrt{1+\frac{1}{x^2}}} + \sqrt{x^2+1} \log(x) + \operatorname{asinh}\left(\frac{1}{x}\right) - \frac{1}{x\sqrt{1+\frac{1}{x^2}}}$$

[In] integrate(x*ln(x)/(x**2+1)**(1/2),x)

[Out] -x/sqrt(1 + x**(-2)) + sqrt(x**2 + 1)*log(x) + asinh(1/x) - 1/(x*sqrt(1 + x**(-2)))

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.74

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \log(x) - \sqrt{x^2+1} + \operatorname{arsinh}\left(\frac{1}{|x|}\right)$$

[In] integrate(x*log(x)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] sqrt(x^2 + 1)*log(x) - sqrt(x^2 + 1) + arcsinh(1/abs(x))

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.29

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = \sqrt{x^2+1} \log(x) - \sqrt{x^2+1} + \frac{1}{2} \log(\sqrt{x^2+1} + 1) - \frac{1}{2} \log(\sqrt{x^2+1} - 1)$$

[In] integrate(x*log(x)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] sqrt(x^2 + 1)*log(x) - sqrt(x^2 + 1) + 1/2*log(sqrt(x^2 + 1) + 1) - 1/2*log(sqrt(x^2 + 1) - 1)

Mupad [F(-1)]

Timed out.

$$\int \frac{x \log(x)}{\sqrt{1+x^2}} dx = \int \frac{x \ln(x)}{\sqrt{x^2+1}} dx$$

```
[In] int((x*log(x))/(x^2 + 1)^(1/2),x)
```

```
[Out] int((x*log(x))/(x^2 + 1)^(1/2), x)
```

3.37 $\int \frac{\sin(x)}{1+\sin^2(x)} dx$

Optimal result	224
Rubi [A] (verified)	224
Mathematica [C] (verified)	225
Maple [A] (verified)	225
Fricas [B] (verification not implemented)	226
Sympy [B] (verification not implemented)	226
Maxima [A] (verification not implemented)	226
Giac [B] (verification not implemented)	227
Mupad [B] (verification not implemented)	227

Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*\cos(x)*2^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3265, 212}

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[In] `Int[Sin[x]/(1 + Sin[x]^2), x]`

[Out] `-(ArcTanh[Cos[x]/Sqrt[2]]/Sqrt[2])`

Rule 212

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 3265

`Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, S`


```
ubst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1}{2-x^2} dx, x, \cos(x)\right) \\ &= -\frac{\operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.88

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = -\frac{i\left(\arctan\left(\frac{-i + \tan\left(\frac{x}{2}\right)}{\sqrt{2}}\right) - \arctan\left(\frac{i + \tan\left(\frac{x}{2}\right)}{\sqrt{2}}\right)\right)}{\sqrt{2}}$$

```
[In] Integrate[Sin[x]/(1 + Sin[x]^2), x]
```

```
[Out] ((-I)*(ArcTan[(-I + Tan[x/2])/Sqrt[2]] - ArcTan[(I + Tan[x/2])/Sqrt[2]]))/Sqrt[2]
```

Maple [A] (verified)

Time = 0.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{\cos(x)\sqrt{2}}{2}\right)\sqrt{2}}{2}$	14
risch	$\frac{\sqrt{2} \ln(e^{2ix} - 2\sqrt{2}e^{ix} + 1)}{4} - \frac{\sqrt{2} \ln(e^{2ix} + 2\sqrt{2}e^{ix} + 1)}{4}$	48

```
[In] int(sin(x)/(1+sin(x)^2), x, method=_RETURNVERBOSE)
```

```
[Out] -1/2*arctanh(1/2*cos(x)*2^(1/2))*2^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(13) = 26$.

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{\cos(x)^2 - 2\sqrt{2}\cos(x) + 2}{\cos(x)^2 - 2} \right)$$

[In] integrate(sin(x)/(1+sin(x)^2),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*log(-(cos(x)^2 - 2*sqrt(2)*cos(x) + 2)/(cos(x)^2 - 2))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(19) = 38$.

Time = 9.24 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.88

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = \frac{\sqrt{2} \log(\tan^2(\frac{x}{2}) - 2\sqrt{2} + 3)}{4} - \frac{\sqrt{2} \log(\tan^2(\frac{x}{2}) + 2\sqrt{2} + 3)}{4}$$

[In] integrate(sin(x)/(1+sin(x)**2),x)

[Out] sqrt(2)*log(tan(x/2)**2 - 2*sqrt(2) + 3)/4 - sqrt(2)*log(tan(x/2)**2 + 2*sqrt(2) + 3)/4

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.50

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{\sqrt{2} - \cos(x)}{\sqrt{2} + \cos(x)} \right)$$

[In] integrate(sin(x)/(1+sin(x)^2),x, algorithm="maxima")

[Out] 1/4*sqrt(2)*log(-(sqrt(2) - cos(x))/(sqrt(2) + cos(x)))

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.69

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = -\frac{1}{4} \sqrt{2} \log(\sqrt{2} + \cos(x)) + \frac{1}{4} \sqrt{2} \log(\sqrt{2} - \cos(x))$$

[In] integrate(sin(x)/(1+sin(x)^2),x, algorithm="giac")

[Out] -1/4*sqrt(2)*log(sqrt(2) + cos(x)) + 1/4*sqrt(2)*log(sqrt(2) - cos(x))

Mupad [B] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.81

$$\int \frac{\sin(x)}{1 + \sin^2(x)} dx = -\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \cos(x)}{2}\right)}{2}$$

[In] int(sin(x)/(sin(x)^2 + 1),x)

[Out] -(2^(1/2)*atanh((2^(1/2)*cos(x))/2))/2

$$3.38 \quad \int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx$$

Optimal result	228
Rubi [A] (verified)	228
Mathematica [A] (verified)	229
Maple [A] (verified)	229
Fricas [B] (verification not implemented)	230
Sympy [F]	230
Maxima [F]	230
Giac [F]	230
Mupad [F(-1)]	231

Optimal result

Integrand size = 24, antiderivative size = 23

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctanh(x*2^(1/2)/(x^4+1)^(1/2))*2^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {1713, 212}

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right)}{\sqrt{2}}$$

[In] Int[(1 + x^2)/((1 - x^2)*Sqrt[1 + x^4]),x]

[Out] ArcTanh[(Sqrt[2]*x)/Sqrt[1 + x^4]]/Sqrt[2]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 1713

Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := Dist[A, Subst[Int[1/(d + 2*a*e*x^2), x], x, x/Sqrt[a + c*x^4]

```
]], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[c*d^2 - a*e^2, 0] && EqQ[B*d + A*e, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \frac{x}{\sqrt{1+x^4}}\right) \\ &= \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}}$$

```
[In] Integrate[(1 + x^2)/((1 - x^2)*Sqrt[1 + x^4]), x]
```

```
[Out] ArcTanh[(Sqrt[2]*x)/Sqrt[1 + x^4]]/Sqrt[2]
```

Maple [A] (verified)

Time = 1.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.96

method	result	size
elliptic	$\frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{x^4+1}}{2x}\right)\sqrt{2}}{2}$	22
trager	$-\frac{\operatorname{RootOf}\left(-Z^2-2\right)\ln\left(\frac{\operatorname{RootOf}\left(-Z^2-2\right)x-\sqrt{x^4+1}}{(-1+x)(1+x)}\right)}{2}$	39
default	$\frac{\sqrt{2}\left(\operatorname{arctanh}\left(\frac{(x^2-x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)-\operatorname{arctanh}\left(\frac{(x^2+x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)\right)}{4}$	47
pseudoelliptic	$\frac{\sqrt{2}\left(\operatorname{arctanh}\left(\frac{(x^2-x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)-\operatorname{arctanh}\left(\frac{(x^2+x+1)\sqrt{2}}{\sqrt{x^4+1}}\right)\right)}{4}$	47

```
[In] int((x^2+1)/(-x^2+1)/(x^4+1)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/2*arctanh(1/2/x*2^(1/2)*(x^4+1)^(1/2))*2^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(18) = 36.

Time = 0.27 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \frac{1}{4}\sqrt{2}\log\left(\frac{x^4+2\sqrt{2}\sqrt{x^4+1}x+2x^2+1}{x^4-2x^2+1}\right)$$

[In] integrate((x^2+1)/(-x^2+1)/(x^4+1)^(1/2),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*log((x^4 + 2*sqrt(2)*sqrt(x^4 + 1)*x + 2*x^2 + 1)/(x^4 - 2*x^2 + 1))

Sympy [F]

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = -\int \frac{x^2}{x^2\sqrt{x^4+1}-\sqrt{x^4+1}} dx - \int \frac{1}{x^2\sqrt{x^4+1}-\sqrt{x^4+1}} dx$$

[In] integrate((x**2+1)/(-x**2+1)/(x**4+1)**(1/2),x)

[Out] -Integral(x**2/(x**2*sqrt(x**4 + 1) - sqrt(x**4 + 1)), x) - Integral(1/(x**2*sqrt(x**4 + 1) - sqrt(x**4 + 1)), x)

Maxima [F]

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \int -\frac{x^2+1}{\sqrt{x^4+1}(x^2-1)} dx$$

[In] integrate((x^2+1)/(-x^2+1)/(x^4+1)^(1/2),x, algorithm="maxima")

[Out] -integrate((x^2 + 1)/(sqrt(x^4 + 1)*(x^2 - 1)), x)

Giac [F]

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \int -\frac{x^2+1}{\sqrt{x^4+1}(x^2-1)} dx$$

[In] integrate((x^2+1)/(-x^2+1)/(x^4+1)^(1/2),x, algorithm="giac")

[Out] integrate(-(x^2 + 1)/(sqrt(x^4 + 1)*(x^2 - 1)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1+x^2}{(1-x^2)\sqrt{1+x^4}} dx = \int -\frac{x^2+1}{(x^2-1)\sqrt{x^4+1}} dx$$

```
[In] int(-(x^2 + 1)/((x^2 - 1)*(x^4 + 1)^(1/2)), x)
```

```
[Out] int(-(x^2 + 1)/((x^2 - 1)*(x^4 + 1)^(1/2)), x)
```

$$3.39 \quad \int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx$$

Optimal result	232
Rubi [A] (verified)	232
Mathematica [A] (verified)	233
Maple [A] (verified)	233
Fricas [A] (verification not implemented)	234
Sympy [F]	234
Maxima [F]	234
Giac [F]	234
Mupad [F(-1)]	235

Optimal result

Integrand size = 24, antiderivative size = 23

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctan(x*2^(1/2)/(x^4+1)^(1/2))*2^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {1713, 209}

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{x^4+1}}\right)}{\sqrt{2}}$$

[In] Int[(1 - x^2)/((1 + x^2)*Sqrt[1 + x^4]),x]

[Out] ArcTan[(Sqrt[2]*x)/Sqrt[1 + x^4]]/Sqrt[2]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 1713

Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := Dist[A, Subst[Int[1/(d + 2*a*e*x^2), x], x, x/Sqrt[a + c*x^4]


```
]], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[c*d^2 - a*e^2, 0] && EqQ[B*d + A*e, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{1+2x^2} dx, x, \frac{x}{\sqrt{1+x^4}}\right) \\ &= \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \frac{\arctan\left(\frac{\sqrt{2}x}{\sqrt{1+x^4}}\right)}{\sqrt{2}}$$

```
[In] Integrate[(1 - x^2)/((1 + x^2)*Sqrt[1 + x^4]), x]
```

```
[Out] ArcTan[(Sqrt[2]*x)/Sqrt[1 + x^4]]/Sqrt[2]
```

Maple [A] (verified)

Time = 1.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\arctan\left(\frac{x\sqrt{2}}{\sqrt{x^4+1}}\right)\sqrt{2}}{2}$	19
pseudoelliptic	$\frac{\arctan\left(\frac{x\sqrt{2}}{\sqrt{x^4+1}}\right)\sqrt{2}}{2}$	19
elliptic	$\frac{\arctan\left(\frac{\sqrt{2}\sqrt{x^4+1}}{2x}\right)\sqrt{2}}{2}$	22
trager	$\frac{\text{RootOf}(_Z^2+2) \ln\left(-\frac{\text{RootOf}(_Z^2+2)x-\sqrt{x^4+1}}{x^2+1}\right)}{2}$	37

```
[In] int((-x^2+1)/(x^2+1)/(x^4+1)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/2*arctan(x*2^(1/2)/(x^4+1)^(1/2))*2^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \frac{1}{2} \sqrt{2} \arctan \left(\frac{\sqrt{2}x}{\sqrt{x^4+1}} \right)$$

[In] integrate((-x^2+1)/(x^2+1)/(x^4+1)^(1/2),x, algorithm="fricas")

[Out] 1/2*sqrt(2)*arctan(sqrt(2)*x/sqrt(x^4 + 1))

Sympy [F]

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = - \int \frac{x^2}{x^2\sqrt{x^4+1} + \sqrt{x^4+1}} dx - \int \left(-\frac{1}{x^2\sqrt{x^4+1} + \sqrt{x^4+1}} \right) dx$$

[In] integrate((-x**2+1)/(x**2+1)/(x**4+1)**(1/2),x)

[Out] -Integral(x**2/(x**2*sqrt(x**4 + 1) + sqrt(x**4 + 1)), x) - Integral(-1/(x**2*sqrt(x**4 + 1) + sqrt(x**4 + 1)), x)

Maxima [F]

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \int -\frac{x^2-1}{\sqrt{x^4+1}(x^2+1)} dx$$

[In] integrate((-x^2+1)/(x^2+1)/(x^4+1)^(1/2),x, algorithm="maxima")

[Out] -integrate((x^2 - 1)/(sqrt(x^4 + 1)*(x^2 + 1)), x)

Giac [F]

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = \int -\frac{x^2-1}{\sqrt{x^4+1}(x^2+1)} dx$$

[In] integrate((-x^2+1)/(x^2+1)/(x^4+1)^(1/2),x, algorithm="giac")

[Out] integrate(-(x^2 - 1)/(sqrt(x^4 + 1)*(x^2 + 1)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1-x^2}{(1+x^2)\sqrt{1+x^4}} dx = - \int \frac{x^2-1}{(x^2+1)\sqrt{x^4+1}} dx$$

```
[In] int(-(x^2 - 1)/((x^2 + 1)*(x^4 + 1)^(1/2)), x)
```

```
[Out] -int((x^2 - 1)/((x^2 + 1)*(x^4 + 1)^(1/2)), x)
```

3.40 $\int \frac{\log(\sin(x))}{1+\sin(x)} dx$

Optimal result	236
Rubi [A] (verified)	236
Mathematica [A] (verified)	237
Maple [A] (verified)	238
Fricas [B] (verification not implemented)	238
Sympy [B] (verification not implemented)	238
Maxima [B] (verification not implemented)	239
Giac [A] (verification not implemented)	239
Mupad [B] (verification not implemented)	240

Optimal result

Integrand size = 10, antiderivative size = 22

$$\int \frac{\log(\sin(x))}{1+\sin(x)} dx = -x - \operatorname{arctanh}(\cos(x)) - \frac{\cos(x) \log(\sin(x))}{1+\sin(x)}$$

[Out] -x-arctanh(cos(x))-cos(x)*ln(sin(x))/(1+sin(x))

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {2727, 2634, 2918, 3855, 8}

$$\int \frac{\log(\sin(x))}{1+\sin(x)} dx = -\operatorname{arctanh}(\cos(x)) - x - \frac{\cos(x) \log(\sin(x))}{\sin(x) + 1}$$

[In] Int[Log[Sin[x]]/(1 + Sin[x]),x]

[Out] -x - ArcTanh[Cos[x]] - (Cos[x]*Log[Sin[x]])/(1 + Sin[x])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2634

Int[Log[u_]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x] - Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x]] /; InverseFunctionFreeQ[u, x]

Rule 2727

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c +
d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b
^2, 0]
```

Rule 2918

```
Int[((cos[(e_) + (f_)*(x_)]*(g_))^(p_)*((d_)*sin[(e_) + (f_)*(x_)])^(
n_))/((a_) + (b_)*sin[(e_) + (f_)*(x_)])], x_Symbol] := Dist[g^2/a, Int[
(g*Cos[e + f*x])^(p - 2)*(d*Sin[e + f*x])^n, x], x] - Dist[g^2/(b*d), Int[(
g*Cos[e + f*x])^(p - 2)*(d*Sin[e + f*x])^(n + 1), x], x] /; FreeQ[{a, b, d,
e, f, g, n, p}, x] && EqQ[a^2 - b^2, 0]
```

Rule 3855

```
Int[csc[(c_) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\cos(x) \log(\sin(x))}{1 + \sin(x)} + \int \frac{\cos(x) \cot(x)}{1 + \sin(x)} dx \\ &= -\frac{\cos(x) \log(\sin(x))}{1 + \sin(x)} - \int 1 dx + \int \csc(x) dx \\ &= -x - \operatorname{arctanh}(\cos(x)) - \frac{\cos(x) \log(\sin(x))}{1 + \sin(x)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.77

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = -x - 2 \log\left(\cos\left(\frac{x}{2}\right)\right) + \frac{2 \log(\sin(x)) \sin\left(\frac{x}{2}\right)}{\cos\left(\frac{x}{2}\right) + \sin\left(\frac{x}{2}\right)}$$

```
[In] Integrate[Log[Sin[x]]/(1 + Sin[x]),x]
```

```
[Out] -x - 2*Log[Cos[x/2]] + (2*Log[Sin[x]]*Sin[x/2])/(Cos[x/2] + Sin[x/2])
```

Maple [A] (verified)

Time = 0.61 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.09

method	result
parallelrisc	$\ln(\csc(x) - \cot(x)) + (-\sec(x) + \tan(x)) \ln(\sin(x)) - x$
norman	$\frac{-x - x \tan\left(\frac{x}{2}\right) + 2 \tan\left(\frac{x}{2}\right) \ln\left(\frac{2 \tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}\right)}{1 + \tan\left(\frac{x}{2}\right)} + \ln\left(1 + \tan^2\left(\frac{x}{2}\right)\right)$
risc	$\frac{2 \ln(e^{ix})}{i + e^{ix}} + \frac{-2 \ln(e^{2ix} - 1) + \ln(e^{ix} + 1) + \ln(e^{ix} - 1) - 2ix - i\pi \operatorname{csgn}(i \sin(x))^2 - i \ln(e^{ix} + 1)e^{ix} - i \ln(e^{ix} - 1)e^{ix} - i\pi \operatorname{csgn}(ie^{-ix})}{i + e^{ix}}$

[In] `int(ln(sin(x))/(sin(x)+1),x,method=_RETURNVERBOSE)`

[Out] `ln(csc(x)-cot(x))+(-sec(x)+tan(x))*ln(sin(x))-x`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 93 vs. 2(22) = 44.

Time = 0.26 (sec) , antiderivative size = 93, normalized size of antiderivative = 4.23

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = \frac{4(\cos(x) + \sin(x) + 1) \arctan\left(-\frac{\cos(x) + \sin(x) + 1}{\cos(x) - \sin(x) + 1}\right) + 4x \cos(x) + (\cos(x) + \sin(x) + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x) + \sin(x) + 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) + 2(\cos(x) - \sin(x) + 1) \log(\sin(x)) + 4x \sin(x) + 4x}{2(\cos(x) + \sin(x) + 1)}$$

[In] `integrate(log(sin(x))/(1+sin(x)),x, algorithm="fricas")`

[Out] `-1/2*(4*(cos(x) + sin(x) + 1)*arctan(-(cos(x) + sin(x) + 1)/(cos(x) - sin(x) + 1)) + 4*x*cos(x) + (cos(x) + sin(x) + 1)*log(1/2*cos(x) + 1/2) - (cos(x) + sin(x) + 1)*log(-1/2*cos(x) + 1/2) + 2*(cos(x) - sin(x) + 1)*log(sin(x)) + 4*x*sin(x) + 4*x)/(cos(x) + sin(x) + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(20) = 40.

Time = 0.66 (sec) , antiderivative size = 105, normalized size of antiderivative = 4.77

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = -\frac{x \tan\left(\frac{x}{2}\right)}{\tan\left(\frac{x}{2}\right) + 1} - \frac{x}{\tan\left(\frac{x}{2}\right) + 1} + \frac{2 \log\left(\frac{\tan\left(\frac{x}{2}\right)}{\tan^2\left(\frac{x}{2}\right) + 1}\right) \tan\left(\frac{x}{2}\right)}{\tan\left(\frac{x}{2}\right) + 1} + \frac{\log\left(\tan^2\left(\frac{x}{2}\right) + 1\right) \tan\left(\frac{x}{2}\right)}{\tan\left(\frac{x}{2}\right) + 1} + \frac{\log\left(\tan^2\left(\frac{x}{2}\right) + 1\right)}{\tan\left(\frac{x}{2}\right) + 1} + \frac{2 \log(2) \tan\left(\frac{x}{2}\right)}{\tan\left(\frac{x}{2}\right) + 1}$$

[In] integrate(ln(sin(x))/(1+sin(x)),x)

[Out] $-x \cdot \tan(x/2) / (\tan(x/2) + 1) - x / (\tan(x/2) + 1) + 2 \cdot \log(\tan(x/2) / (\tan(x/2)^2 + 1)) \cdot \tan(x/2) / (\tan(x/2) + 1) + \log(\tan(x/2)^2 + 1) \cdot \tan(x/2) / (\tan(x/2) + 1) + \log(\tan(x/2)^2 + 1) / (\tan(x/2) + 1) + 2 \cdot \log(2) \cdot \tan(x/2) / (\tan(x/2) + 1)$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. $2(22) = 44$.

Time = 0.28 (sec) , antiderivative size = 82, normalized size of antiderivative = 3.73

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = -\frac{2 \log\left(\frac{2 \sin(x)}{\left(\frac{\sin(x)^2}{(\cos(x)+1)^2 + 1\right)(\cos(x)+1)}\right)}{\frac{\sin(x)}{\cos(x)+1} + 1} - 2 \arctan\left(\frac{\sin(x)}{\cos(x) + 1}\right) + 2 \log\left(\frac{\sin(x)}{\cos(x) + 1}\right) - \log\left(\frac{\sin(x)^2}{(\cos(x) + 1)^2} + 1\right)$$

[In] integrate(log(sin(x))/(1+sin(x)),x, algorithm="maxima")

[Out] $-2 \cdot \log(2 \cdot \sin(x) / ((\sin(x)^2 / (\cos(x) + 1)^2 + 1) \cdot (\cos(x) + 1))) / (\sin(x) / (\cos(x) + 1) + 1) - 2 \cdot \arctan(\sin(x) / (\cos(x) + 1)) + 2 \cdot \log(\sin(x) / (\cos(x) + 1)) - \log(\sin(x)^2 / (\cos(x) + 1)^2 + 1)$

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.64

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = -x - \frac{2 \log(\sin(x))}{\tan\left(\frac{1}{2}x\right) + 1} - 2 \log\left(\tan\left(\frac{1}{4}x\right)^2 + 1\right) + 2 \log\left(\left|\tan\left(\frac{1}{4}x\right)\right|\right)$$

[In] integrate(log(sin(x))/(1+sin(x)),x, algorithm="giac")

[Out] $-x - 2 \cdot \log(\sin(x)) / (\tan(1/2 \cdot x) + 1) - 2 \cdot \log(\tan(1/4 \cdot x)^2 + 1) + 2 \cdot \log(\text{abs}(\tan(1/4 \cdot x)))$

Mupad [B] (verification not implemented)

Time = 0.39 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.50

$$\int \frac{\log(\sin(x))}{1 + \sin(x)} dx = -2x + \ln(2 \sin(x) - \cos(x) 2i - 2i) (-1 - i) \\ + \ln(2 \sin(x) - \cos(x) 2i + 2i) (1 - i) - \frac{2 \ln(\sin(x))}{\cos(x) + \sin(x) 1i + 1i}$$

[In] int(log(sin(x))/(sin(x) + 1),x)

[Out] log(2*sin(x) - cos(x)*2i + 2i)*(1 - 1i) - log(2*sin(x) - cos(x)*2i - 2i)*(1 + 1i) - 2*x - (2*log(sin(x)))/(cos(x) + sin(x)*1i + 1i)

3.41 $\int \log(\sin(x)) \sqrt{1 + \sin(x)} dx$

Optimal result	241
Rubi [A] (verified)	241
Mathematica [B] (verified)	243
Maple [F]	244
Fricas [B] (verification not implemented)	244
Sympy [F]	244
Maxima [F]	245
Giac [B] (verification not implemented)	245
Mupad [F(-1)]	245

Optimal result

Integrand size = 12, antiderivative size = 42

$$\int \log(\sin(x)) \sqrt{1 + \sin(x)} dx = -4 \operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{1 + \sin(x)}}\right) + \frac{4 \cos(x)}{\sqrt{1 + \sin(x)}} - \frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}}$$

[Out] $-4*\operatorname{arctanh}(\cos(x)/(1+\sin(x))^{(1/2)})+4*\cos(x)/(1+\sin(x))^{(1/2)}-2*\cos(x)*\ln(\sin(x))/(1+\sin(x))^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {2725, 2634, 12, 2953, 3060, 2852, 212}

$$\int \log(\sin(x)) \sqrt{1 + \sin(x)} dx = -4 \operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{\sin(x) + 1}}\right) + \frac{4 \cos(x)}{\sqrt{\sin(x) + 1}} - \frac{2 \cos(x) \log(\sin(x))}{\sqrt{\sin(x) + 1}}$$

[In] $\operatorname{Int}[\operatorname{Log}[\operatorname{Sin}[x]]*\operatorname{Sqrt}[1 + \operatorname{Sin}[x]],x]$

[Out] $-4*\operatorname{ArcTanh}[\operatorname{Cos}[x]/\operatorname{Sqrt}[1 + \operatorname{Sin}[x]]] + (4*\operatorname{Cos}[x])/\operatorname{Sqrt}[1 + \operatorname{Sin}[x]] - (2*\operatorname{Cos}[x]*\operatorname{Log}[\operatorname{Sin}[x]])/\operatorname{Sqrt}[1 + \operatorname{Sin}[x]]$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 2634

```
Int[Log[u]*(v_), x_Symbol] := With[{w = IntHide[v, x]}, Dist[Log[u], w, x]
- Int[SimplifyIntegrand[w*(D[u, x]/u), x], x] /; InverseFunctionFreeQ[w, x
]] /; InverseFunctionFreeQ[u, x]
```

Rule 2725

```
Int[Sqrt[(a_) + (b_.)*sin[(c_) + (d_.)*(x_)]], x_Symbol] := Simp[-2*b*(Cos
[c + d*x]/(d*Sqrt[a + b*Sin[c + d*x]])), x] /; FreeQ[{a, b, c, d}, x] && Eq
Q[a^2 - b^2, 0]
```

Rule 2852

```
Int[Sqrt[(a_) + (b_.)*sin[(e_) + (f_.)*(x_)]]/((c_) + (d_.)*sin[(e_) + (
f_.)*(x_)]), x_Symbol] := Dist[-2*(b/f), Subst[Int[1/(b*c + a*d - d*x^2), x
], x, b*(Cos[e + f*x]/Sqrt[a + b*Sin[e + f*x])], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 - b^2, 0] && NeQ[c^2 - d^2, 0]
```

Rule 2953

```
Int[cos[(e_) + (f_.)*(x_)]^2*((d_.)*sin[(e_) + (f_.)*(x_)]^(n_)*((a_) +
(b_.)*sin[(e_) + (f_.)*(x_)]^(m_)), x_Symbol] := Dist[1/b^2, Int[(d*Sin[e
+ f*x])^n*(a + b*Sin[e + f*x])^(m + 1)*(a - b*Sin[e + f*x]), x], x] /; Free
Q[{a, b, d, e, f, m, n}, x] && EqQ[a^2 - b^2, 0] && (ILtQ[m, 0] || !IGtQ[n
, 0])
```

Rule 3060

```
Int[Sqrt[(a_) + (b_.)*sin[(e_) + (f_.)*(x_)]]*((A_.) + (B_.)*sin[(e_) + (
f_.)*(x_)])*((c_) + (d_.)*sin[(e_) + (f_.)*(x_)]^(n_), x_Symbol] := Simp
[-2*b*B*Cos[e + f*x]*((c + d*Sin[e + f*x])^(n + 1)/(d*f*(2*n + 3)*Sqrt[a +
b*Sin[e + f*x]))], x] + Dist[(A*b*d*(2*n + 3) - B*(b*c - 2*a*d*(n + 1)))/(b
*d*(2*n + 3)), Int[Sqrt[a + b*Sin[e + f*x]]*(c + d*Sin[e + f*x])^n, x], x]
/; FreeQ[{a, b, c, d, e, f, A, B, n}, x] && NeQ[b*c - a*d, 0] && EqQ[a^2 -
b^2, 0] && NeQ[c^2 - d^2, 0] && !LtQ[n, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}} - \int -\frac{2 \cos(x) \cot(x)}{\sqrt{1 + \sin(x)}} dx \\
 &= -\frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}} + 2 \int \frac{\cos(x) \cot(x)}{\sqrt{1 + \sin(x)}} dx \\
 &= -\frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}} + 2 \int \csc(x)(1 - \sin(x))\sqrt{1 + \sin(x)} dx \\
 &= \frac{4 \cos(x)}{\sqrt{1 + \sin(x)}} - \frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}} + 2 \int \csc(x)\sqrt{1 + \sin(x)} dx \\
 &= \frac{4 \cos(x)}{\sqrt{1 + \sin(x)}} - \frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}} - 4 \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, \frac{\cos(x)}{\sqrt{1 + \sin(x)}} \right) \\
 &= -4 \operatorname{arctanh} \left(\frac{\cos(x)}{\sqrt{1 + \sin(x)}} \right) + \frac{4 \cos(x)}{\sqrt{1 + \sin(x)}} - \frac{2 \cos(x) \log(\sin(x))}{\sqrt{1 + \sin(x)}}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 87 vs. $2(42) = 84$.

Time = 0.07 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.07

$$\int \log(\sin(x))\sqrt{1 + \sin(x)} dx$$

$$= \frac{2(-\log(1 + \cos(\frac{x}{2}) - \sin(\frac{x}{2})) + \log(1 - \cos(\frac{x}{2}) + \sin(\frac{x}{2})) - \cos(\frac{x}{2})(-2 + \log(\sin(x))) + (-2 + \log(\sin(x)))\cos(\frac{x}{2}) + \sin(\frac{x}{2}))}{\cos(\frac{x}{2}) + \sin(\frac{x}{2})}$$

[In] Integrate[Log[Sin[x]]*Sqrt[1 + Sin[x]],x]

[Out] (2*(-Log[1 + Cos[x/2] - Sin[x/2]] + Log[1 - Cos[x/2] + Sin[x/2]] - Cos[x/2] *(-2 + Log[Sin[x]]) + (-2 + Log[Sin[x]])*Sin[x/2])*Sqrt[1 + Sin[x]])/(Cos[x/2] + Sin[x/2])

Maple [F]

$$\int \ln(\sin(x)) \sqrt{\sin(x) + 1} dx$$

[In] `int(ln(sin(x))*(sin(x)+1)^(1/2),x)`

[Out] `int(ln(sin(x))*(sin(x)+1)^(1/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(36) = 72$.

Time = 0.27 (sec) , antiderivative size = 146, normalized size of antiderivative = 3.48

$$\int \log(\sin(x)) \sqrt{1 + \sin(x)} dx =$$

$$\frac{(\cos(x) + \sin(x) + 1) \log\left(\frac{\cos(x)^2 - (\cos(x) - 1)\sin(x) + 2(\cos(x) - \sin(x) + 1)\sqrt{\sin(x) + 1} + 2\cos(x) + 1}{2(\cos(x) + \sin(x) + 1)}\right) - (\cos(x) + \sin(x) -$$

[In] `integrate(log(sin(x))*(1+sin(x))^(1/2),x, algorithm="fricas")`

[Out] `-((cos(x) + sin(x) + 1)*log(1/2*(cos(x)^2 - (cos(x) - 1)*sin(x) + 2*(cos(x) - sin(x) + 1)*sqrt(sin(x) + 1) + 2*cos(x) + 1)/(cos(x) + sin(x) + 1)) - (cos(x) + sin(x) + 1)*log(1/2*(cos(x)^2 - (cos(x) - 1)*sin(x) - 2*(cos(x) - sin(x) + 1)*sqrt(sin(x) + 1) + 2*cos(x) + 1)/(cos(x) + sin(x) + 1)) + 2*((cos(x) - sin(x) + 1)*log(sin(x)) - 2*cos(x) + 2*sin(x) - 2)*sqrt(sin(x) + 1))/(cos(x) + sin(x) + 1)`

Sympy [F]

$$\int \log(\sin(x)) \sqrt{1 + \sin(x)} dx = \int \sqrt{\sin(x) + 1} \log(\sin(x)) dx$$

[In] `integrate(ln(sin(x))*(1+sin(x))**(1/2),x)`

[Out] `Integral(sqrt(sin(x) + 1)*log(sin(x)), x)`

Maxima [F]

$$\int \log(\sin(x))\sqrt{1+\sin(x)} dx = \int \sqrt{\sin(x)+1} \log(\sin(x)) dx$$

[In] integrate(log(sin(x))*(1+sin(x))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(sin(x) + 1)*log(sin(x)), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. 2(36) = 72.

Time = 0.32 (sec) , antiderivative size = 90, normalized size of antiderivative = 2.14

$$\int \log(\sin(x))\sqrt{1+\sin(x)} dx$$

$$= \sqrt{2} \left(2 \log(\sin(x)) \operatorname{sgn}\left(\cos\left(-\frac{1}{4}\pi + \frac{1}{2}x\right)\right) \sin\left(-\frac{1}{4}\pi + \frac{1}{2}x\right) + \left(\sqrt{2} \log\left(\frac{|-2\sqrt{2} + 4 \sin(\frac{1}{4}\pi - \frac{1}{2}x)|}{|2\sqrt{2} + 4 \sin(\frac{1}{4}\pi - \frac{1}{2}x)|}\right)\right) \right)$$

[In] integrate(log(sin(x))*(1+sin(x))^(1/2),x, algorithm="giac")

[Out] sqrt(2)*(2*log(sin(x))*sgn(cos(-1/4*pi + 1/2*x))*sin(-1/4*pi + 1/2*x) + (sqrt(2)*log(abs(-2*sqrt(2) + 4*sin(1/4*pi - 1/2*x))/abs(2*sqrt(2) + 4*sin(1/4*pi - 1/2*x)))) + 4*sin(1/4*pi - 1/2*x))*sgn(cos(-1/4*pi + 1/2*x))

Mupad [F(-1)]

Timed out.

$$\int \log(\sin(x))\sqrt{1+\sin(x)} dx = \int \ln(\sin(x)) \sqrt{\sin(x)+1} dx$$

[In] int(log(sin(x))*(sin(x) + 1)^(1/2),x)

[Out] int(log(sin(x))*(sin(x) + 1)^(1/2), x)

$$3.42 \quad \int \frac{\sec(x)}{\sqrt{-1+\sec^4(x)}} dx$$

Optimal result	246
Rubi [B] (verified)	246
Mathematica [A] (verified)	248
Maple [B] (verified)	248
Fricas [B] (verification not implemented)	248
Sympy [F]	249
Maxima [F]	249
Giac [B] (verification not implemented)	249
Mupad [F(-1)]	250

Optimal result

Integrand size = 13, antiderivative size = 28

$$\int \frac{\sec(x)}{\sqrt{-1+\sec^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\cos(x)\cot(x)\sqrt{-1+\sec^4(x)}}{\sqrt{2}}\right)}{\sqrt{2}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*\cos(x)*\cot(x)*(-1+\sec(x)^4)^{(1/2)}*2^{(1/2)})*2^{(1/2)}$

Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.14 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.11, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$, Rules used = {4233, 6854, 2013, 2033, 212}

$$\int \frac{\sec(x)}{\sqrt{-1+\sec^4(x)}} dx = -\frac{\sqrt{1-\cos^4(x)}\sec^2(x)\operatorname{arctanh}\left(\frac{\sqrt{2}\sin(x)}{\sqrt{2\sin^2(x)-\sin^4(x)}}\right)}{\sqrt{2}\sqrt{\sec^4(x)-1}}$$

[In] `Int[Sec[x]/Sqrt[-1 + Sec[x]^4], x]`

[Out] $-\left(\operatorname{ArcTanh}\left[\frac{\sqrt{2}\sin(x)}{\sqrt{2\sin^2(x)-\sin^4(x)}}\right]/\sqrt{2}\sqrt{\sec^4(x)-1}\right)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 2013

Int[(u_)^(p_), x_Symbol] := Int[ExpandToSum[u, x]^p, x] /; FreeQ[p, x] && TrinomialQ[u, x] && !TrinomialMatchQ[u, x]

Rule 2033

Int[1/Sqrt[(a_)*(x_)^2 + (b_)*(x_)^(n_)], x_Symbol] := Dist[2/(2 - n), Subst[Int[1/(1 - a*x^2), x], x, x/Sqrt[a*x^2 + b*x^n], x] /; FreeQ[{a, b, n}, x] && NeQ[n, 2]

Rule 4233

Int[sec[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sec[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + b/(1 - ff^2*x^2))^(n/2)]^p/(1 - ff^2*x^2)^((m + 1)/2), x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2] && IntegerQ[n/2] && !IntegerQ[p]

Rule 6854

Int[(u_)*((a_) + (b_)*(v_)^(n_))^(p_), x_Symbol] := Dist[(a + b*v^n)^FracPart[p]/(v^(n*FracPart[p])*(b + a/v^n)^FracPart[p]), Int[u*v^(n*p)*(b + a/v^n)^p, x], x] /; FreeQ[{a, b, p}, x] && !IntegerQ[p] && ILtQ[n, 0] && BinomialQ[v, x] && !LinearQ[v, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{1}{(1-x^2) \sqrt{-1 + \frac{1}{(1-x^2)^2}}} dx, x, \sin(x) \right) \\
 &= \frac{\left(\sqrt{1 - \cos^4(x)} \sec^2(x) \right) \text{Subst} \left(\int \frac{1}{\sqrt{1-(1-x^2)^2}} dx, x, \sin(x) \right)}{\sqrt{-1 + \sec^4(x)}} \\
 &= \frac{\left(\sqrt{1 - \cos^4(x)} \sec^2(x) \right) \text{Subst} \left(\int \frac{1}{\sqrt{2x^2-x^4}} dx, x, \sin(x) \right)}{\sqrt{-1 + \sec^4(x)}} \\
 &= - \frac{\left(\sqrt{1 - \cos^4(x)} \sec^2(x) \right) \text{Subst} \left(\int \frac{1}{1-2x^2} dx, x, \frac{\sin(x)}{\sqrt{2\sin^2(x)-\sin^4(x)}} \right)}{\sqrt{-1 + \sec^4(x)}} \\
 &= - \frac{\text{arctanh} \left(\frac{\sqrt{2}\sin(x)}{\sqrt{2\sin^2(x)-\sin^4(x)}} \right) \sqrt{1 - \cos^4(x)} \sec^2(x)}{\sqrt{2}\sqrt{-1 + \sec^4(x)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.61

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{1}{2}\sqrt{4 - 2\sin^2(x)}\right) \sqrt{3 + \cos(2x)} \sec(x) \tan(x)}{2\sqrt{-1 + \sec^4(x)}}$$

[In] Integrate[Sec[x]/Sqrt[-1 + Sec[x]^4],x]

[Out] -1/2*(ArcTanh[Sqrt[4 - 2*Sin[x]^2]/2]*Sqrt[3 + Cos[2*x]]*Sec[x]*Tan[x])/Sqrt[-1 + Sec[x]^4]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(23) = 46.

Time = 0.77 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.79

method	result	size
default	$\frac{\left(\operatorname{arcsinh}\left(\frac{-1+\cos(x)}{\cos(x)+1}\right) - \operatorname{arctanh}\left(\frac{\sqrt{2}}{2\sqrt{\frac{1+\cos^2(x)}{(\cos(x)+1)^2}}}\right)\right) \sqrt{\frac{1+\cos^2(x)}{(\cos(x)+1)^2}} (\tan(x) + \sec(x) \tan(x)) \sqrt{8}}{8\sqrt{(\tan^2(x))(\sec^2(x)+1)}}$	78

[In] int(sec(x)/(-1+sec(x)^4)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/8*(arcsinh((-1+cos(x))/(cos(x)+1))-arctanh(1/2/((1+cos(x)^2)/(cos(x)+1)^2)^(1/2)*2^(1/2)))*((1+cos(x)^2)/(cos(x)+1)^2)^(1/2)/(tan(x)^2*(sec(x)^2+1))^(1/2)*(tan(x)+sec(x)*tan(x))*8^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(23) = 46.

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.93

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{2 \left(2\sqrt{2} \sqrt{-\frac{\cos(x)^4 - 1}{\cos(x)^4}} \cos(x)^2 - (\cos(x)^2 + 3) \sin(x) \right)}{(\cos(x)^2 - 1) \sin(x)} \right)$$

[In] integrate(sec(x)/(-1+sec(x)^4)^(1/2),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*log(-2*(2*sqrt(2)*sqrt(-(cos(x)^4 - 1)/cos(x)^4)*cos(x)^2 - (cos(x)^2 + 3)*sin(x))/((cos(x)^2 - 1)*sin(x)))

Sympy [F]

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = \int \frac{\sec(x)}{\sqrt{(\sec(x) - 1)(\sec(x) + 1)(\sec^2(x) + 1)}} dx$$

[In] integrate(sec(x)/(-1+sec(x)**4)**(1/2),x)

[Out] Integral(sec(x)/sqrt((sec(x) - 1)*(sec(x) + 1)*(sec(x)**2 + 1)), x)

Maxima [F]

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = \int \frac{\sec(x)}{\sqrt{\sec(x)^4 - 1}} dx$$

[In] integrate(sec(x)/(-1+sec(x)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(sec(x)/sqrt(sec(x)^4 - 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(23) = 46.

Time = 0.51 (sec) , antiderivative size = 92, normalized size of antiderivative = 3.29

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = \frac{\sqrt{2} \left(\log \left(\tan \left(\frac{1}{2} x \right)^2 - \sqrt{\tan \left(\frac{1}{2} x \right)^4 + 1} + 1 \right) - \log \left(-\tan \left(\frac{1}{2} x \right)^2 + \sqrt{\tan \left(\frac{1}{2} x \right)^4 + 1} + 1 \right) + \log \left(-\tan \left(\frac{1}{2} x \right)^2 + \sqrt{\tan \left(\frac{1}{2} x \right)^4 + 1} - 1 \right) \right)}{4 \operatorname{sgn} \left(\tan \left(\frac{1}{2} x \right)^5 + 2 \tan \left(\frac{1}{2} x \right)^3 + \tan \left(\frac{1}{2} x \right) \right)}$$

[In] integrate(sec(x)/(-1+sec(x)^4)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*(log(tan(1/2*x)^2 - sqrt(tan(1/2*x)^4 + 1) + 1) - log(-tan(1/2*x)^2 + sqrt(tan(1/2*x)^4 + 1) + 1) + log(-tan(1/2*x)^2 + sqrt(tan(1/2*x)^4 + 1) - 1))/sgn(tan(1/2*x)^5 + 2*tan(1/2*x)^3 + tan(1/2*x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(x)}{\sqrt{-1 + \sec^4(x)}} dx = \int \frac{1}{\cos(x) \sqrt{\frac{1}{\cos(x)^4} - 1}} dx$$

```
[In] int(1/(cos(x)*(1/cos(x)^4 - 1)^(1/2)),x)
```

```
[Out] int(1/(cos(x)*(1/cos(x)^4 - 1)^(1/2)), x)
```

3.43 $\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx$

Optimal result	251
Rubi [A] (verified)	251
Mathematica [A] (verified)	252
Maple [A] (verified)	253
Fricas [B] (verification not implemented)	253
Sympy [F]	254
Maxima [B] (verification not implemented)	254
Giac [A] (verification not implemented)	255
Mupad [F(-1)]	255

Optimal result

Integrand size = 13, antiderivative size = 34

$$\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{1-\tan^2(x)}{\sqrt{2}\sqrt{1+\tan^4(x)}}\right)}{2\sqrt{2}}$$

[Out] $-1/4*\operatorname{arctanh}(1/2*(1-\tan(x)^2)*2^{(1/2)}/(1+\tan(x)^4)^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3751, 1262, 739, 212}

$$\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{1-\tan^2(x)}{\sqrt{2}\sqrt{\tan^4(x)+1}}\right)}{2\sqrt{2}}$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]/\operatorname{Sqrt}[1 + \operatorname{Tan}[x]^4], x]$

[Out] $-1/2*\operatorname{ArcTanh}[(1 - \operatorname{Tan}[x]^2)/(\operatorname{Sqrt}[2]*\operatorname{Sqrt}[1 + \operatorname{Tan}[x]^4])]/\operatorname{Sqrt}[2]$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{Gt} Q[a, 0] \ || \operatorname{Lt} Q[b, 0])$

Rule 739

$\operatorname{Int}[1/(((d_+ + (e_+)(x_+))\operatorname{Sqrt}[(a_+ + (c_+)(x_+)^2]), x_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/\operatorname{Sqrt}[a + c*x^2]] /; \operatorname{FreeQ}$

[{a, c, d, e}, x]

Rule 1262

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
  :> Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
  [{a, c, d, e, p, q}, x]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
  (f_)*(x_)])^(n_))^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
  x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
  ^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
  , p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
  alQ[n]))
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{1+x^4}} dx, x, \tan(x)\right) \\
 &= \frac{1}{2} \text{Subst}\left(\int \frac{1}{(1+x)\sqrt{1+x^2}} dx, x, \tan^2(x)\right) \\
 &= -\left(\frac{1}{2} \text{Subst}\left(\int \frac{1}{2-x^2} dx, x, \frac{1-\tan^2(x)}{\sqrt{1+\tan^4(x)}}\right)\right) \\
 &= -\frac{\text{arctanh}\left(\frac{1-\tan^2(x)}{\sqrt{2}\sqrt{1+\tan^4(x)}}\right)}{2\sqrt{2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.62

$$\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx = -\frac{\sqrt{3+\cos(4x)} \log\left(\sqrt{2}\cos(2x) + \sqrt{3+\cos(4x)}\right) \sec^2(x)}{4\sqrt{2}\sqrt{1+\tan^4(x)}}$$

```
[In] Integrate[Tan[x]/Sqrt[1 + Tan[x]^4], x]
```

```
[Out] -1/4*(Sqrt[3 + Cos[4*x]]*Log[Sqrt[2]*Cos[2*x] + Sqrt[3 + Cos[4*x]])*Sec[x]^
  2)/(Sqrt[2]*Sqrt[1 + Tan[x]^4])
```

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.09

method	result	size
derivativedivides	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(-2(\tan^2(x))+2)\sqrt{2}}{4\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}\right)}{4}$	37
default	$-\frac{\sqrt{2} \operatorname{arctanh}\left(\frac{(-2(\tan^2(x))+2)\sqrt{2}}{4\sqrt{(1+\tan^2(x))^2-2(\tan^2(x))}}\right)}{4}$	37

[In] `int(tan(x)/(1+tan(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-1/4*2^{(1/2)}*\operatorname{arctanh}(1/4*(-2*\tan(x)^2+2)*2^{(1/2)}/((1+\tan(x)^2)^2-2*\tan(x)^2)^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. $2(25) = 50$.

Time = 0.26 (sec) , antiderivative size = 186, normalized size of antiderivative = 5.47

$$\int \frac{\tan(x)}{\sqrt{1+\tan^4(x)}} dx$$

$$= \frac{1}{32} \sqrt{2} \log \left(\frac{577 \tan(x)^{16} - 1912 \tan(x)^{14} + 4124 \tan(x)^{12} - 6216 \tan(x)^{10} + 7110 \tan(x)^8 - 6216 \tan(x)^6 + 4124 \tan(x)^4 - 1912 \tan(x)^2 + 577}{(51\sqrt{2}\tan(x)^{14} - 169\sqrt{2}\tan(x)^{12} + 339\sqrt{2}\tan(x)^{10} - 465\sqrt{2}\tan(x)^8 + 465\sqrt{2}\tan(x)^6 - 339\sqrt{2}\tan(x)^4 + 169\sqrt{2}\tan(x)^2 - 51\sqrt{2})\sqrt{\tan(x)^4 + 1} + 577} \right)$$

[In] `integrate(tan(x)/(1+tan(x)^4)^(1/2),x, algorithm="fricas")`

[Out] $1/32*\sqrt{2}*\log((577*\tan(x)^{16} - 1912*\tan(x)^{14} + 4124*\tan(x)^{12} - 6216*\tan(x)^{10} + 7110*\tan(x)^8 - 6216*\tan(x)^6 + 4124*\tan(x)^4 - 1912*\tan(x)^2 + 577)/(51*\sqrt{2}*\tan(x)^{14} - 169*\sqrt{2}*\tan(x)^{12} + 339*\sqrt{2}*\tan(x)^{10} - 465*\sqrt{2}*\tan(x)^8 + 465*\sqrt{2}*\tan(x)^6 - 339*\sqrt{2}*\tan(x)^4 + 169*\sqrt{2}*\tan(x)^2 - 51*\sqrt{2})*\sqrt{\tan(x)^4 + 1} + 577)/(\tan(x)^{16} + 8*\tan(x)^{14} + 28*\tan(x)^{12} + 56*\tan(x)^{10} + 70*\tan(x)^8 + 56*\tan(x)^6 + 28*\tan(x)^4 + 8*\tan(x)^2 + 1))$

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{1 + \tan^4(x)}} dx = \int \frac{\tan(x)}{\sqrt{\tan^4(x) + 1}} dx$$

```
[In] integrate(tan(x)/(1+tan(x)**4)**(1/2),x)
```

```
[Out] Integral(tan(x)/sqrt(tan(x)**4 + 1), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 565 vs. $2(25) = 50$.

Time = 0.37 (sec) , antiderivative size = 565, normalized size of antiderivative = 16.62

$$\int \frac{\tan(x)}{\sqrt{1 + \tan^4(x)}} dx = \text{Too large to display}$$

```
[In] integrate(tan(x)/(1+tan(x)^4)^(1/2),x, algorithm="maxima")
```

```
[Out] -1/16*sqrt(2)*(log(4*sqrt(2)*(6*cos(4*x) + 1)*cos(8*x) + cos(8*x)^2 + 36*cos(4*x)^2 + sin(8*x)^2 + 12*sin(8*x)*sin(4*x) + 36*sin(4*x)^2 + 12*cos(4*x) + 1)*cos(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1))^2 + 4*sqrt(2)*(6*cos(4*x) + 1)*cos(8*x) + cos(8*x)^2 + 36*cos(4*x)^2 + sin(8*x)^2 + 12*sin(8*x)*sin(4*x) + 36*sin(4*x)^2 + 12*cos(4*x) + 1)*sin(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1))^2 + 32*(2*(6*cos(4*x) + 1)*cos(8*x) + cos(8*x)^2 + 36*cos(4*x)^2 + sin(8*x)^2 + 12*sin(8*x)*sin(4*x) + 36*sin(4*x)^2 + 12*cos(4*x) + 1)^(1/4)*cos(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1)) + 64) + log(4*cos(4*x)^2 + 4*sin(4*x)^2 + 4*sqrt(2)*(6*cos(4*x) + 1)*cos(8*x) + cos(8*x)^2 + 36*cos(4*x)^2 + sin(8*x)^2 + 12*sin(8*x)*sin(4*x) + 36*sin(4*x)^2 + 12*cos(4*x) + 1)*(cos(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1))^2 + sin(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1))^2) + 8*(2*(6*cos(4*x) + 1)*cos(8*x) + cos(8*x)^2 + 36*cos(4*x)^2 + sin(8*x)^2 + 12*sin(8*x)*sin(4*x) + 36*sin(4*x)^2 + 12*cos(4*x) + 1)^(1/4)*((cos(4*x) + 3)*cos(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1)) + sin(4*x)*sin(1/2*arctan2(sin(8*x) + 6*sin(4*x), cos(8*x) + 6*cos(4*x) + 1)))) + 24*cos(4*x) + 36)
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.47

$$\int \frac{\tan(x)}{\sqrt{1 + \tan^4(x)}} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{\tan(x)^2 + \sqrt{2} - \sqrt{\tan(x)^4 + 1 + 1}}{\tan(x)^2 - \sqrt{2} - \sqrt{\tan(x)^4 + 1 + 1}} \right)$$

[In] integrate(tan(x)/(1+tan(x)^4)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*log(-(tan(x)^2 + sqrt(2) - sqrt(tan(x)^4 + 1) + 1)/(tan(x)^2 - sqrt(2) - sqrt(tan(x)^4 + 1) + 1))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(x)}{\sqrt{1 + \tan^4(x)}} dx = \int \frac{\tan(x)}{\sqrt{\tan(x)^4 + 1}} dx$$

[In] int(tan(x)/(tan(x)^4 + 1)^(1/2),x)

[Out] int(tan(x)/(tan(x)^4 + 1)^(1/2), x)

3.44 $\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx$

Optimal result	256
Rubi [A] (verified)	256
Mathematica [A] (verified)	257
Maple [B] (verified)	258
Fricas [B] (verification not implemented)	258
Sympy [F(-1)]	258
Maxima [F]	259
Giac [B] (verification not implemented)	259
Mupad [F(-1)]	259

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{3}\cos(x)(1+\sin^2(x))}{2\sqrt{1-\sin^6(x)}}\right)}{2\sqrt{3}}$$

[Out] 1/6*arctanh(1/2*cos(x)*(1+sin(x)^2)*3^(1/2)/(1-sin(x)^6)^(1/2))*3^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3295, 2021, 1918, 212}

$$\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\cos(x)(6-3\cos^2(x))}{2\sqrt{3}\sqrt{\cos^6(x)-3\cos^4(x)+3\cos^2(x)}}\right)}{2\sqrt{3}}$$

[In] Int[Sin[x]/Sqrt[1 - Sin[x]^6],x]

[Out] ArcTanh[(Cos[x]*(6 - 3*Cos[x]^2))/(2*Sqrt[3]*Sqrt[3*Cos[x]^2 - 3*Cos[x]^4 + Cos[x]^6])]/(2*Sqrt[3])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 1918


```
Int[1/Sqrt[(a_.)*(x_)^2 + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(r_.)], x_Symbol] :
> Dist[-2/(n - 2), Subst[Int[1/(4*a - x^2), x], x, x*((2*a + b*x^(n - 2))/S
qrt[a*x^2 + b*x^n + c*x^r])], x] /; FreeQ[{a, b, c, n, r}, x] && EqQ[r, 2*n
- 2] && PosQ[n - 2] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2021

```
Int[(u_)^(p_), x_Symbol] :=> Int[ExpandToSum[u, x]^p, x] /; FreeQ[p, x] && G
eneralizedTrinomialQ[u, x] && !GeneralizedTrinomialMatchQ[u, x]
```

Rule 3295

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(n_
))^p, x_Symbol] :=> With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f
, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*(1 - ff^2*x^2)^(n/2))^p, x],
x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2
] && IntegerQ[n/2]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{1}{\sqrt{1 - (1 - x^2)^3}} dx, x, \cos(x)\right) \\
 &= -\text{Subst}\left(\int \frac{1}{\sqrt{3x^2 - 3x^4 + x^6}} dx, x, \cos(x)\right) \\
 &= \text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{\cos(x)(6 - 3\cos^2(x))}{\sqrt{3\cos^2(x) - 3\cos^4(x) + \cos^6(x)}}\right) \\
 &= \frac{\text{arctanh}\left(\frac{\cos(x)(6 - 3\cos^2(x))}{2\sqrt{3}\sqrt{3\cos^2(x) - 3\cos^4(x) + \cos^6(x)}}\right)}{2\sqrt{3}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.67

$$\int \frac{\sin(x)}{\sqrt{1 - \sin^6(x)}} dx = -\frac{\text{arctanh}\left(\frac{\sqrt{\frac{3}{2}}(-3 + \cos(2x))}{\sqrt{15 - 8\cos(2x) + \cos(4x)}}\right) \cos(x) \sqrt{15 - 8\cos(2x) + \cos(4x)}}{4\sqrt{6 - 6\sin^6(x)}}$$

```
[In] Integrate[Sin[x]/Sqrt[1 - Sin[x]^6], x]
```

```
[Out] -1/4*(ArcTanh[(Sqrt[3/2]*(-3 + Cos[2*x]))/Sqrt[15 - 8*Cos[2*x] + Cos[4*x]]]
*Cos[x]*Sqrt[15 - 8*Cos[2*x] + Cos[4*x]])/Sqrt[6 - 6*Sin[x]^6]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. 2(29) = 58.

Time = 0.89 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.72

method	result	size
default	$-\frac{\cos(x)\sqrt{\cos^4(x)-3(\cos^2(x))+3}\sqrt{3}\operatorname{arctanh}\left(\frac{(\cos^2(x)-2)\sqrt{3}}{2\sqrt{\cos^4(x)-3(\cos^2(x))+3}}\right)}{6\sqrt{\cos^6(x)-3(\cos^4(x))+3(\cos^2(x))}}$	67

[In] `int(sin(x)/(1-sin(x)^6)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/6/(cos(x)^6-3*cos(x)^4+3*cos(x)^2)^(1/2)*cos(x)*(cos(x)^4-3*cos(x)^2+3)^(1/2)*3^(1/2)*arctanh(1/2*(cos(x)^2-2)*3^(1/2)/(cos(x)^4-3*cos(x)^2+3)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.62

$$\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx$$

$$= \frac{1}{12} \sqrt{3} \log \left(\frac{7 \cos(x)^5 - 24 \cos(x)^3 - 4 \sqrt{\cos(x)^6 - 3 \cos(x)^4 + 3 \cos(x)^2} (\sqrt{3} \cos(x)^2 - 2 \sqrt{3}) + 24 \cos(x)}{\cos(x)^5} \right)$$

[In] `integrate(sin(x)/(1-sin(x)^6)^(1/2),x, algorithm="fricas")`

[Out] `1/12*sqrt(3)*log((7*cos(x)^5 - 24*cos(x)^3 - 4*sqrt(cos(x)^6 - 3*cos(x)^4 + 3*cos(x)^2)*(sqrt(3)*cos(x)^2 - 2*sqrt(3)) + 24*cos(x))/cos(x)^5)`

Sympy [F(-1)]

Timed out.

$$\int \frac{\sin(x)}{\sqrt{1-\sin^6(x)}} dx = \text{Timed out}$$

[In] `integrate(sin(x)/(1-sin(x)**6)**(1/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\sin(x)}{\sqrt{1 - \sin^6(x)}} dx = \int \frac{\sin(x)}{\sqrt{-\sin(x)^6 + 1}} dx$$

[In] integrate(sin(x)/(1-sin(x)^6)^(1/2),x, algorithm="maxima")

[Out] integrate(sin(x)/sqrt(-sin(x)^6 + 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(29) = 58.

Time = 0.33 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.72

$$\int \frac{\sin(x)}{\sqrt{1 - \sin^6(x)}} dx = \frac{\sqrt{3} \log\left(\cos(x)^2 + \sqrt{3} - \sqrt{\cos(x)^4 - 3\cos(x)^2 + 3}\right) - \sqrt{3} \log\left(-\cos(x)^2 + \sqrt{3} + \sqrt{\cos(x)^4 - 3\cos(x)^2 + 3}\right)}{6 \operatorname{sgn}(\cos(x))}$$

[In] integrate(sin(x)/(1-sin(x)^6)^(1/2),x, algorithm="giac")

[Out] -1/6*(sqrt(3)*log(cos(x)^2 + sqrt(3) - sqrt(cos(x)^4 - 3*cos(x)^2 + 3)) - sqrt(3)*log(-cos(x)^2 + sqrt(3) + sqrt(cos(x)^4 - 3*cos(x)^2 + 3)))/sgn(cos(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\sin(x)}{\sqrt{1 - \sin^6(x)}} dx = \int \frac{\sin(x)}{\sqrt{1 - \sin(x)^6}} dx$$

[In] int(sin(x)/(1 - sin(x)^6)^(1/2),x)

[Out] int(sin(x)/(1 - sin(x)^6)^(1/2), x)

3.45 $\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx$

Optimal result	260
Rubi [F]	261
Mathematica [A] (warning: unable to verify)	261
Maple [F]	262
Fricas [C] (verification not implemented)	262
Sympy [F]	264
Maxima [F]	264
Giac [F]	265
Mupad [F(-1)]	265

Optimal result

Integrand size = 23, antiderivative size = 337

$$\begin{aligned}
 & \int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx \\
 &= \sqrt{2} \left(\sqrt{-1 + \sqrt{2}} \arctan \left(\frac{\sqrt{-2 + 2\sqrt{2}}(-\sqrt{2} - \sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)})}{2\sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}}} \right) \right. \\
 & \quad - \sqrt{1 + \sqrt{2}} \arctan \left(\frac{\sqrt{2 + 2\sqrt{2}}(-\sqrt{2} - \sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)})}{2\sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}}} \right) \\
 & \quad - \sqrt{1 + \sqrt{2}} \operatorname{arctanh} \left(\frac{\sqrt{-2 + 2\sqrt{2}}\sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}}}{\sqrt{2} - \sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} \right) \\
 & \quad \left. + \sqrt{-1 + \sqrt{2}} \operatorname{arctanh} \left(\frac{\sqrt{2 + 2\sqrt{2}}\sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}}}{\sqrt{2} - \sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} \right) \right) \cot(x) \sqrt{-1 + \sec(x)} \sqrt{1 + \sec(x)}
 \end{aligned}$$

```

[Out] cot(x)*2^(1/2)*(-1+sec(x))^(1/2)*(1+sec(x))^(1/2)*(arctan(1/2*(-2^(1/2)-(-1
+sec(x))^(1/2)+(1+sec(x))^(1/2))*(-2+2*2^(1/2))^(1/2)/(-(-1+sec(x))^(1/2)+(
1+sec(x))^(1/2))^(1/2))*(2^(1/2)-1)^(1/2)+arctanh((2+2*2^(1/2))^(1/2)*(-(-1
+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2)/(2^(1/2)-(-1+sec(x))^(1/2)+(1+sec(x)
)^(1/2)))*(2^(1/2)-1)^(1/2)-arctan(1/2*(-2^(1/2)-(-1+sec(x))^(1/2)+(1+sec(x)
))^(1/2))*(2+2*2^(1/2))^(1/2)/(-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2))*
(1+2^(1/2))^(1/2)-arctanh((-2+2*2^(1/2))^(1/2)*(-(-1+sec(x))^(1/2)+(1+sec(x)
))^(1/2))^(1/2)/(2^(1/2)-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2)))*(1+2^(1/2))^(
1/2))

```

Rubi [F]

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = \int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx$$

[In] Int[Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]], x]

[Out] Defer[Int][Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]], x]

Rubi steps

$$\text{integral} = \int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx$$

Mathematica [A] (warning: unable to verify)

Time = 2.85 (sec) , antiderivative size = 552, normalized size of antiderivative = 1.64

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx$$

$$= \frac{\sqrt[4]{2} \cos(x) \left(\sqrt{-1 + \sec(x)} - \sqrt{1 + \sec(x)} \right)^2 \left(2 \arctan \left(\cot \left(\frac{\pi}{8} \right) - \frac{\csc \left(\frac{\pi}{8} \right) \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}}}{\sqrt[4]{2}} \right) \right) \cos \left(\frac{\pi}{8} \right)}{1}$$

[In] Integrate[Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]], x]

[Out] (2^(1/4)*Cos[x]*(Sqrt[-1 + Sec[x]] - Sqrt[1 + Sec[x]])^2*(2*ArcTan[Cot [Pi/8] - (Csc [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]/2^(1/4)]*Cos [Pi/8] - 2*ArcTan[Cot [Pi/8] + (Csc [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]/2^(1/4)]*Cos [Pi/8] + Cos [Pi/8]*Log[2 + Sqrt[2]*(-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])] - 2*2^(3/4)*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]] *Sin [Pi/8]] - Cos [Pi/8]*Log[2 + Sqrt[2]*(-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])] + 2*2^(3/4)*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]]*Sin [Pi/8]] + 2*ArcTan[(Sec [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]/2^(1/4) - Tan [Pi/8]]*Sin [Pi/8] + 2*ArcTan[(Sec [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]/2^(1/4) + Tan [Pi/8]]*Sin [Pi/8] - Log[2 - 2*2^(3/4)*Cos [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]] + Sqrt[2]*(-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]*Sin [Pi/8] + Log[2 + 2^(1/4)*Csc [Pi/8]*Sqrt[-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]]] + Sqrt[2]*(-Sqrt[-1 + Sec[x]] + Sqrt[1 + Sec[x]])]*Sin [Pi/8]) *Sin [x])/(-1 + Cos [2*x] + 2*Cos [x]*Sqrt[-1 + Sec[x]]*Sqrt[1 + Sec[x]])

Maple [F]

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx$$

[In] int((-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2),x)

[Out] int((-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2),x)

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 337, normalized size of antiderivative = 1.00

$$\begin{aligned}
 \int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = & \frac{1}{2} \sqrt{4i + 4} \log \left(i \sqrt{4i + 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right) \\
 & - \frac{1}{2} \sqrt{4i + 4} \log \left(-i \sqrt{4i + 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right) \\
 & - \frac{1}{2} \sqrt{-4i + 4} \log \left(i \sqrt{-4i + 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right) \\
 & + \frac{1}{2} \sqrt{-4i + 4} \log \left(-i \sqrt{-4i + 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right) \\
 & - \frac{1}{2} \sqrt{4i - 4} \log \left(i \sqrt{4i - 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right) \\
 & + \frac{1}{2} \sqrt{4i - 4} \log \left(-i \sqrt{4i - 4} \right. \\
 & \left. + 2 \sqrt{\frac{\sqrt{\frac{\cos(x)+1}{\cos(x)}} (\cos(x) - \sin(x) + 1)}{\cos(x) + 1}} \right)
 \end{aligned}$$

```
[In] integrate((-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*sqrt(4*I + 4)*log(I*sqrt(4*I + 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) - 1/2*sqrt(4*I + 4)*log(-I*sqrt(4*I + 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) - 1/2*sqrt(-4*I + 4)*log(I*sqrt(-4*I + 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) + 1/2*sqrt(-4*I + 4)*log(-I*sqrt(-4*I + 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) - 1/2*sqrt(4*I - 4)*log(I*sqrt(4*I - 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) + 1/2*sqrt(4*I - 4)*log(-I*sqrt(4*I - 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) + 1/2*sqrt(-4*I - 4)*log(I*sqrt(-4*I - 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1))) - 1/2*sqrt(-4*I - 4)*log(-I*sqrt(-4*I - 4) + 2*sqrt(sqrt((cos(x) + 1)/cos(x))*(cos(x) - sin(x) + 1)/(cos(x) + 1)))
```

Sympy [F]

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = \int \sqrt{-\sqrt{\sec(x) - 1} + \sqrt{\sec(x) + 1}} dx$$

```
[In] integrate((-(-1+sec(x))**(1/2)+(1+sec(x))**(1/2))**(1/2),x)
```

```
[Out] Integral(sqrt(-sqrt(sec(x) - 1) + sqrt(sec(x) + 1)), x)
```

Maxima [F]

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = \int \sqrt{\sqrt{\sec(x) + 1} - \sqrt{\sec(x) - 1}} dx$$

```
[In] integrate((-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(sqrt(sec(x) + 1) - sqrt(sec(x) - 1)), x)
```


Giac [F]

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = \int \sqrt{\sqrt{\sec(x) + 1} - \sqrt{\sec(x) - 1}} dx$$

[In] integrate((-(-1+sec(x))^(1/2)+(1+sec(x))^(1/2))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(sqrt(sec(x) + 1) - sqrt(sec(x) - 1)), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{-\sqrt{-1 + \sec(x)} + \sqrt{1 + \sec(x)}} dx = \int \sqrt{\sqrt{\frac{1}{\cos(x)} + 1} - \sqrt{\frac{1}{\cos(x)} - 1}} dx$$

[In] int(((1/cos(x) + 1)^(1/2) - (1/cos(x) - 1)^(1/2))^(1/2),x)

[Out] int(((1/cos(x) + 1)^(1/2) - (1/cos(x) - 1)^(1/2))^(1/2), x)

3.46 $\int x \arctan(x)^2 \log(1+x^2) dx$

Optimal result	266
Rubi [A] (verified)	266
Mathematica [A] (verified)	269
Maple [A] (verified)	269
Fricas [A] (verification not implemented)	270
Sympy [A] (verification not implemented)	270
Maxima [A] (verification not implemented)	270
Giac [F]	271
Mupad [B] (verification not implemented)	271

Optimal result

Integrand size = 12, antiderivative size = 77

$$\int x \arctan(x)^2 \log(1+x^2) dx = 3x \arctan(x) - \frac{3 \arctan(x)^2}{2} - \frac{1}{2} x^2 \arctan(x)^2 - \frac{3}{2} \log(1+x^2) - x \arctan(x) \log(1+x^2) + \frac{1}{2} (1+x^2) \arctan(x)^2 \log(1+x^2) + \frac{1}{4} \log^2(1+x^2)$$

[Out] 3*x*arctan(x)-3/2*arctan(x)^2-1/2*x^2*arctan(x)^2-3/2*ln(x^2+1)-x*arctan(x)*ln(x^2+1)+1/2*(x^2+1)*arctan(x)^2*ln(x^2+1)+1/4*ln(x^2+1)^2

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {4946, 5036, 4930, 266, 5004, 5143, 5129, 2525, 2437, 2338}

$$\int x \arctan(x)^2 \log(1+x^2) dx = -\frac{1}{2} x^2 \arctan(x)^2 + \frac{1}{2} (x^2+1) \arctan(x)^2 \log(x^2+1) - x \arctan(x) \log(x^2+1) - \frac{3 \arctan(x)^2}{2} + 3x \arctan(x) + \frac{1}{4} \log^2(x^2+1) - \frac{3}{2} \log(x^2+1)$$

[In] Int[x*ArcTan[x]^2*Log[1+x^2],x]

[Out] 3*x*ArcTan[x] - (3*ArcTan[x]^2)/2 - (x^2*ArcTan[x]^2)/2 - (3*Log[1+x^2])/2 - x*ArcTan[x]*Log[1+x^2] + ((1+x^2)*ArcTan[x]^2*Log[1+x^2])/2 + Log[1+x^2]^2/4

Rule 266

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 2338

```
Int[((a_) + Log[(c_)*(x_)^(n_)]*(b_))/(x_), x_Symbol] := Simp[(a + b*Log[c*x^n])^2/(2*b*n), x] /; FreeQ[{a, b, c, n}, x]
```

Rule 2437

```
Int[((a_) + Log[(c_)*((d_) + (e_)*(x_)^(n_))]*(b_))^(p_)*((f_) + (g_)*(x_)^(q_)), x_Symbol] := Dist[1/e, Subst[Int[(f*(x/d))^q*(a + b*Log[c*x^n])^p, x], x, d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q}, x] && EqQ[e*f - d*g, 0]
```

Rule 2525

```
Int[((a_) + Log[(c_)*((d_) + (e_)*(x_)^(n_))^(p_)]*(b_))^(q_)*(x_)^(m_)*((f_) + (g_)*(x_)^(s_))^(r_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(f + g*x^(s/n))^r*(a + b*Log[c*(d + e*x)^p])^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p, q, r, s}, x] && IntegerQ[r] && IntegerQ[s/n] && IntegerQ[Simplify[(m + 1)/n]] && (GtQ[(m + 1)/n, 0] || IGtQ[q, 0])
```

Rule 4930

```
Int[((a_) + ArcTan[(c_)*(x_)^(n_)]*(b_))^(p_), x_Symbol] := Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

Rule 4946

```
Int[((a_) + ArcTan[(c_)*(x_)^(n_)]*(b_))^(p_)*(x_)^(m_), x_Symbol] := Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Dist[b*c*n*(p/(m + 1)), Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Rule 5004

```
Int[((a_) + ArcTan[(c_)*(x_)*(b_)]^(p_)/((d_) + (e_)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5036

```
Int[(((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))^ (p_.)*((f_.)*(x_.))^(m_.))/((d_.) + (e_.)*(x_)^2), x_Symbol] := Dist[f^2/e, Int[(f*x)^(m - 2)*(a + b*ArcTan[c*x])^p, x], x] - Dist[d*(f^2/e), Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

Rule 5129

```
Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))*((d_.) + Log[(f_.) + (g_.)*(x_)^2]*(e_.)), x_Symbol] := Simp[x*(d + e*Log[f + g*x^2])*(a + b*ArcTan[c*x]), x] + (-Dist[b*c, Int[x*((d + e*Log[f + g*x^2])/(1 + c^2*x^2)), x], x] - Dist[2*e*g, Int[x^2*((a + b*ArcTan[c*x])/(f + g*x^2)), x], x]) /; FreeQ[{a, b, c, d, e, f, g}, x]
```

Rule 5143

```
Int[((a_.) + ArcTan[(c_.)*(x_.)]*(b_.))^2*((d_.) + Log[(f_.) + (g_.)*(x_)^2]*(e_.))*(x_), x_Symbol] := Simp[(f + g*x^2)*(d + e*Log[f + g*x^2])*(a + b*ArcTan[c*x])^2/(2*g), x] + (-Dist[b/c, Int[(d + e*Log[f + g*x^2])*(a + b*ArcTan[c*x]), x], x] + Dist[b*c*e, Int[x^2*((a + b*ArcTan[c*x])/(1 + c^2*x^2)), x], x] - Simp[e*x^2*((a + b*ArcTan[c*x])^2/2), x]) /; FreeQ[{a, b, c, d, e, f, g}, x] && EqQ[g, c^2*f]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{1}{2}x^2 \arctan(x)^2 + \frac{1}{2}(1+x^2) \arctan(x)^2 \log(1+x^2) \\
&\quad + \int \frac{x^2 \arctan(x)}{1+x^2} dx - \int \arctan(x) \log(1+x^2) dx \\
&= -\frac{1}{2}x^2 \arctan(x)^2 - x \arctan(x) \log(1+x^2) + \frac{1}{2}(1+x^2) \arctan(x)^2 \log(1+x^2) \\
&\quad + 2 \int \frac{x^2 \arctan(x)}{1+x^2} dx + \int \arctan(x) dx - \int \frac{\arctan(x)}{1+x^2} dx + \int \frac{x \log(1+x^2)}{1+x^2} dx \\
&= x \arctan(x) - \frac{\arctan(x)^2}{2} - \frac{1}{2}x^2 \arctan(x)^2 - x \arctan(x) \log(1+x^2) \\
&\quad + \frac{1}{2}(1+x^2) \arctan(x)^2 \log(1+x^2) + \frac{1}{2} \text{Subst}\left(\int \frac{\log(1+x)}{1+x} dx, x, x^2\right) \\
&\quad + 2 \int \arctan(x) dx - 2 \int \frac{\arctan(x)}{1+x^2} dx - \int \frac{x}{1+x^2} dx
\end{aligned}$$

$$\begin{aligned}
&= 3x \arctan(x) - \frac{3 \arctan(x)^2}{2} - \frac{1}{2} x^2 \arctan(x)^2 - \frac{1}{2} \log(1+x^2) - x \arctan(x) \log(1+x^2) \\
&\quad + \frac{1}{2} (1+x^2) \arctan(x)^2 \log(1+x^2) + \frac{1}{2} \text{Subst} \left(\int \frac{\log(x)}{x} dx, x, 1+x^2 \right) - 2 \int \frac{x}{1+x^2} dx \\
&= 3x \arctan(x) - \frac{3 \arctan(x)^2}{2} - \frac{1}{2} x^2 \arctan(x)^2 - \frac{3}{2} \log(1+x^2) \\
&\quad - x \arctan(x) \log(1+x^2) + \frac{1}{2} (1+x^2) \arctan(x)^2 \log(1+x^2) + \frac{1}{4} \log^2(1+x^2)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.75

$$\begin{aligned}
\int x \arctan(x)^2 \log(1+x^2) dx = & \frac{1}{4} (-4x \arctan(x) (-3 + \log(1+x^2)) \\
& + (-6 + \log(1+x^2)) \log(1+x^2) \\
& + 2 \arctan(x)^2 (-3 - x^2 + (1+x^2) \log(1+x^2)))
\end{aligned}$$

[In] Integrate[x*ArcTan[x]^2*Log[1+x^2],x]

[Out] (-4*x*ArcTan[x]*(-3+Log[1+x^2]))+(-6+Log[1+x^2])*Log[1+x^2]+2*ArcTan[x]^2*(-3-x^2+(1+x^2)*Log[1+x^2]))/4

Maple [A] (verified)

Time = 4.50 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.01

method	result
parallelrisch	$\frac{\ln(x^2+1) \arctan(x)^2 x^2}{2} - \frac{x^2 \arctan(x)^2}{2} - x \arctan(x) \ln(x^2+1) + \frac{\arctan(x)^2 \ln(x^2+1)}{2} + 3x \arctan(x)$
default	Expression too large to display
risch	Expression too large to display

[In] int(x*arctan(x)^2*ln(x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/2*ln(x^2+1)*arctan(x)^2*x^2-1/2*x^2*arctan(x)^2-x*arctan(x)*ln(x^2+1)+1/2*arctan(x)^2*ln(x^2+1)+3*x*arctan(x)-3/2*arctan(x)^2+1/4*ln(x^2+1)^2-3/2*ln(x^2+1)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.68

$$\int x \arctan(x)^2 \log(1+x^2) dx = -\frac{1}{2} (x^2 + 3) \arctan(x)^2 + 3x \arctan(x) + \frac{1}{2} ((x^2 + 1) \arctan(x)^2 - 2x \arctan(x) - 3) \log(x^2 + 1) + \frac{1}{4} \log(x^2 + 1)^2$$

`[In] integrate(x*arctan(x)^2*log(x^2+1),x, algorithm="fricas")`

```
[Out] -1/2*(x^2 + 3)*arctan(x)^2 + 3*x*arctan(x) + 1/2*((x^2 + 1)*arctan(x)^2 - 2*x*arctan(x) - 3)*log(x^2 + 1) + 1/4*log(x^2 + 1)^2
```

Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.13

$$\int x \arctan(x)^2 \log(1+x^2) dx = \frac{x^2 \log(x^2 + 1) \operatorname{atan}^2(x)}{2} - \frac{x^2 \operatorname{atan}^2(x)}{2} - x \log(x^2 + 1) \operatorname{atan}(x) + 3x \operatorname{atan}(x) + \frac{\log(x^2 + 1)^2}{4} + \frac{\log(x^2 + 1) \operatorname{atan}^2(x)}{2} - \frac{3 \log(x^2 + 1)}{2} - \frac{3 \operatorname{atan}^2(x)}{2}$$

`[In] integrate(x*atan(x)**2*ln(x**2+1),x)`

```
[Out] x**2*log(x**2 + 1)*atan(x)**2/2 - x**2*atan(x)**2/2 - x*log(x**2 + 1)*atan(x) + 3*x*atan(x) + log(x**2 + 1)**2/4 + log(x**2 + 1)*atan(x)**2/2 - 3*log(x**2 + 1)/2 - 3*atan(x)**2/2
```

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.87

$$\int x \arctan(x)^2 \log(1+x^2) dx = -\frac{1}{2} (x^2 - (x^2 + 1) \log(x^2 + 1) + 1) \arctan(x)^2 - (x \log(x^2 + 1) - 3x + 2 \arctan(x)) \arctan(x) + \arctan(x)^2 + \frac{1}{4} \log(x^2 + 1)^2 - \frac{3}{2} \log(x^2 + 1)$$

[In] integrate(x*arctan(x)^2*log(x^2+1),x, algorithm="maxima")

[Out] $-1/2*(x^2 - (x^2 + 1)*\log(x^2 + 1) + 1)*\arctan(x)^2 - (x*\log(x^2 + 1) - 3*x + 2*\arctan(x))*\arctan(x) + \arctan(x)^2 + 1/4*\log(x^2 + 1)^2 - 3/2*\log(x^2 + 1)$

Giac [F]

$$\int x \arctan(x)^2 \log(1 + x^2) dx = \int x \arctan(x)^2 \log(x^2 + 1) dx$$

[In] integrate(x*arctan(x)^2*log(x^2+1),x, algorithm="giac")

[Out] integrate(x*arctan(x)^2*log(x^2 + 1), x)

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.01

$$\begin{aligned} \int x \arctan(x)^2 \log(1 + x^2) dx = & \frac{\ln(x^2 + 1)^2}{4} - \frac{3 \ln(x^2 + 1)}{2} \\ & - \frac{3 \operatorname{atan}(x)^2}{2} + \frac{\ln(x^2 + 1) \operatorname{atan}(x)^2}{2} \\ & + x (3 \operatorname{atan}(x) - \ln(x^2 + 1) \operatorname{atan}(x)) \\ & - x^2 \left(\frac{\operatorname{atan}(x)^2}{2} - \frac{\ln(x^2 + 1) \operatorname{atan}(x)^2}{2} \right) \end{aligned}$$

[In] int(x*log(x^2 + 1)*atan(x)^2,x)

[Out] $\log(x^2 + 1)^2/4 - (3*\log(x^2 + 1))/2 - (3*\operatorname{atan}(x)^2)/2 + (\log(x^2 + 1)*\operatorname{atan}(x)^2)/2 + x*(3*\operatorname{atan}(x) - \log(x^2 + 1)*\operatorname{atan}(x)) - x^2*(\operatorname{atan}(x)^2/2 - (\log(x^2 + 1)*\operatorname{atan}(x)^2)/2)$

3.47 $\int \arctan\left(x\sqrt{1+x^2}\right) dx$

Optimal result	272
Rubi [A] (verified)	272
Mathematica [C] (verified)	275
Maple [B] (verified)	275
Fricas [B] (verification not implemented)	276
Sympy [F]	277
Maxima [F]	277
Giac [A] (verification not implemented)	277
Mupad [B] (verification not implemented)	278

Optimal result

Integrand size = 12, antiderivative size = 120

$$\int \arctan\left(x\sqrt{1+x^2}\right) dx = x \arctan\left(x\sqrt{1+x^2}\right) + \frac{1}{2} \arctan\left(\sqrt{3} - 2\sqrt{1+x^2}\right) - \frac{1}{2} \arctan\left(\sqrt{3} + 2\sqrt{1+x^2}\right) - \frac{1}{4}\sqrt{3} \log\left(2+x^2 - \sqrt{3}\sqrt{1+x^2}\right) + \frac{1}{4}\sqrt{3} \log\left(2+x^2 + \sqrt{3}\sqrt{1+x^2}\right)$$

[Out] $-1/2*\arctan(-3^{(1/2)}+2*(x^2+1)^{(1/2)})+x*\arctan(x*(x^2+1)^{(1/2)})-1/2*\arctan(3^{(1/2)}+2*(x^2+1)^{(1/2)})-1/4*\ln(2+x^2-3^{(1/2)}*(x^2+1)^{(1/2)})*3^{(1/2)}+1/4*\ln(2+x^2+3^{(1/2)}*(x^2+1)^{(1/2)})*3^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5311, 1699, 840, 1183, 648, 632, 210, 642}

$$\int \arctan\left(x\sqrt{1+x^2}\right) dx = x \arctan\left(x\sqrt{x^2+1}\right) + \frac{1}{2} \arctan\left(\sqrt{3} - 2\sqrt{x^2+1}\right) - \frac{1}{2} \arctan\left(2\sqrt{x^2+1} + \sqrt{3}\right) - \frac{1}{4}\sqrt{3} \log\left(x^2 - \sqrt{3}\sqrt{x^2+1} + 2\right) + \frac{1}{4}\sqrt{3} \log\left(x^2 + \sqrt{3}\sqrt{x^2+1} + 2\right)$$

[In] $\text{Int}[\text{ArcTan}[x*\text{Sqrt}[1 + x^2]], x]$

[Out] $x*\text{ArcTan}[x*\text{Sqrt}[1 + x^2]] + \text{ArcTan}[\text{Sqrt}[3] - 2*\text{Sqrt}[1 + x^2]]/2 - \text{ArcTan}[\text{Sqrt}[3] + 2*\text{Sqrt}[1 + x^2]]/2 - (\text{Sqrt}[3]*\text{Log}[2 + x^2 - \text{Sqrt}[3]*\text{Sqrt}[1 + x^2]])/4 + (\text{Sqrt}[3]*\text{Log}[2 + x^2 + \text{Sqrt}[3]*\text{Sqrt}[1 + x^2]])/4$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 840

Int[((f_) + (g_)*(x_))/(Sqrt[(d_) + (e_)*(x_)]*((a_) + (b_)*(x_) + (c_)*(x_)^2)), x_Symbol] := Dist[2, Subst[Int[(e*f - d*g + g*x^2)/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0]

Rule 1183

Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]

Rule 1699

Int[(Px_)*(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[(Px /. x -> Sqrt[x])*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]

&& PolyQ[Px, x^2]

Rule 5311

Int[ArcTan[u_], x_Symbol] :> Simp[x*ArcTan[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/(1 + u^2)), x], x] /; InverseFunctionFreeQ[u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= x \arctan\left(x\sqrt{1+x^2}\right) - \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) - \frac{1}{2} \text{Subst}\left(\int \frac{1+2x}{\sqrt{1+x}(1+x+x^2)} dx, x, x^2\right) \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) - \text{Subst}\left(\int \frac{-1+2x^2}{1-x^2+x^4} dx, x, \sqrt{1+x^2}\right) \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) - \frac{\text{Subst}\left(\int \frac{-\sqrt{3}+3x}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right)}{2\sqrt{3}} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{-\sqrt{3}-3x}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right)}{2\sqrt{3}} \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) - \frac{1}{4} \text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right) \\
 &\quad - \frac{1}{4} \text{Subst}\left(\int \frac{1}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right) \\
 &\quad - \frac{1}{4} \sqrt{3} \text{Subst}\left(\int \frac{-\sqrt{3}+2x}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right) \\
 &\quad + \frac{1}{4} \sqrt{3} \text{Subst}\left(\int \frac{\sqrt{3}+2x}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2}\right) \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) - \frac{1}{4} \sqrt{3} \log\left(2+x^2-\sqrt{3}\sqrt{1+x^2}\right) \\
 &\quad + \frac{1}{4} \sqrt{3} \log\left(2+x^2+\sqrt{3}\sqrt{1+x^2}\right) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{-1-x^2} dx, x, -\sqrt{3}+2\sqrt{1+x^2}\right) \\
 &\quad + \frac{1}{2} \text{Subst}\left(\int \frac{1}{-1-x^2} dx, x, \sqrt{3}+2\sqrt{1+x^2}\right) \\
 &= x \arctan\left(x\sqrt{1+x^2}\right) + \frac{1}{2} \arctan\left(\sqrt{3}-2\sqrt{1+x^2}\right) - \frac{1}{2} \arctan\left(\sqrt{3}+2\sqrt{1+x^2}\right) \\
 &\quad - \frac{1}{4} \sqrt{3} \log\left(2+x^2-\sqrt{3}\sqrt{1+x^2}\right) + \frac{1}{4} \sqrt{3} \log\left(2+x^2+\sqrt{3}\sqrt{1+x^2}\right)
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.79

$$\int \arctan(x\sqrt{1+x^2}) dx = -\frac{1}{2}(1-i\sqrt{3}) \arctan\left(\frac{1}{2}(1-i\sqrt{3})\sqrt{1+x^2}\right) - \frac{1}{2}(1+i\sqrt{3}) \arctan\left(\frac{1}{2}(1+i\sqrt{3})\sqrt{1+x^2}\right) + x \arctan(x\sqrt{1+x^2})$$

[In] Integrate[ArcTan[x*Sqrt[1 + x^2]],x]

[Out] $-1/2*((1 - I*\text{Sqrt}[3])*\text{ArcTan}[\frac{(1 - I*\text{Sqrt}[3])*\text{Sqrt}[1 + x^2]}{2}]) - ((1 + I*\text{Sqrt}[3])*\text{ArcTan}[\frac{(1 + I*\text{Sqrt}[3])*\text{Sqrt}[1 + x^2]}{2}])/2 + x*\text{ArcTan}[x*\text{Sqrt}[1 + x^2]]$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 507 vs. $2(92) = 184$.

Time = 0.17 (sec) , antiderivative size = 508, normalized size of antiderivative = 4.23

method	result
default	$x \arctan(x\sqrt{x^2+1}) + \frac{\sqrt{2}\sqrt{\frac{2(-1+x)^2}{(-1-x)^2}+2\sqrt{3}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{2(-1+x)^2}{(-1-x)^2}+2\sqrt{3}}}{2}\right)}{3\sqrt{\frac{(-1+x)^2+1}{(-1-x)^2}+1}\left(\frac{-1+x}{-1-x}+1\right)} + \frac{\sqrt{2}\sqrt{\frac{2(1+x)^2}{(1-x)^2}+2\sqrt{3}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{2(1+x)^2}{(1-x)^2}+2\sqrt{3}}}{2}\right)}{3\sqrt{\frac{(1+x)^2+1}{(1-x)^2}+1}\left(\frac{1+x}{1-x}+1\right)}$
parts	$x \arctan(x\sqrt{x^2+1}) + \frac{\sqrt{2}\sqrt{\frac{2(-1+x)^2}{(-1-x)^2}+2\sqrt{3}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{2(-1+x)^2}{(-1-x)^2}+2\sqrt{3}}}{2}\right)}{3\sqrt{\frac{(-1+x)^2+1}{(-1-x)^2}+1}\left(\frac{-1+x}{-1-x}+1\right)} + \frac{\sqrt{2}\sqrt{\frac{2(1+x)^2}{(1-x)^2}+2\sqrt{3}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{2(1+x)^2}{(1-x)^2}+2\sqrt{3}}}{2}\right)}{3\sqrt{\frac{(1+x)^2+1}{(1-x)^2}+1}\left(\frac{1+x}{1-x}+1\right)}$

[In] int(arctan(x*(x^2+1)^(1/2)),x,method=_RETURNVERBOSE)

[Out] $x*\arctan(x*(x^2+1)^{(1/2)})+1/3*2^{(1/2)}/(((1-x)^2/(-1-x)^2+1)/((1-x)/(-1-x)+1)^{(1/2)}/((1-x)/(-1-x)+1)*(2*(-1+x)^2/(-1-x)^2+2)^{(1/2)}*3^{(1/2)}*\arctan(h(1/2*(2*(-1+x)^2/(-1-x)^2+2)^{(1/2)}*3^{(1/2)}))+1/3*2^{(1/2)}/(((1+x)^2/(1-x)^2+1)/((1+x)/(1-x)+1)^{(1/2)}/((1+x)/(1-x)+1)*(2*(1+x)^2/(1-x)^2+2)^{(1/2)}*3^{(1/2)}*\operatorname{arctanh}(1/2*(2*(1+x)^2/(1-x)^2+2)^{(1/2)}*3^{(1/2)})-1/12*2^{(1/2)}*(2*(-1+x)^2/(-1-x)^2+2)^{(1/2)}*(3^{(1/2)}*\operatorname{arctanh}(1/2*(2*(-1+x)^2/(-1-x)^2+2)^{(1/2)}*3^{(1/2)})-3*\arctan(1/((1-x)^2/(-1-x)^2+1)*(2*(-1+x)^2/(-1-x)^2+2)^{(1/2)}*(-1+x$

)/(-1-x)))/(((1+x)^2/(-1-x)^2+1)/((1+x)/(-1-x)+1)^2)^(1/2)/((1+x)/(-1-x)+1)-1/12*2^(1/2)*(2*(1+x)^2/(1-x)^2+2)^(1/2)*(3^(1/2)*arctanh(1/2*(2*(1+x)^2/(1-x)^2+2)^(1/2)*3^(1/2))-3*arctan(1/((1+x)^2/(1-x)^2+1)*(2*(1+x)^2/(1-x)^2+2)^(1/2)*(1+x)/(1-x)))/(((1+x)^2/(1-x)^2+1)/((1+x)/(1-x)+1)^2)^(1/2)/((1+x)/(1-x)+1)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 298 vs. 2(92) = 184.

Time = 0.26 (sec) , antiderivative size = 298, normalized size of antiderivative = 2.48

$$\int \arctan(x\sqrt{1+x^2}) dx = x \arctan(\sqrt{x^2+1}x) - \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{-3}+1} \log\left(4x^2 + \sqrt{2}(\sqrt{-3}x+x)\sqrt{-\sqrt{-3}+1} - \sqrt{x^2+1}\left(\sqrt{2}(\sqrt{-3}+1)\sqrt{-\sqrt{-3}+1} + 4x\right) + 4\right) + \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{-3}+1} \log\left(4x^2 - \sqrt{2}(\sqrt{-3}x+x)\sqrt{-\sqrt{-3}+1} + \sqrt{x^2+1}\left(\sqrt{2}(\sqrt{-3}+1)\sqrt{-\sqrt{-3}+1} - 4x\right) + 4\right) - \frac{1}{4} \sqrt{2} \sqrt{\sqrt{-3}+1} \log\left(4x^2 - 4\sqrt{x^2+1}x + \left(\sqrt{2}\sqrt{x^2+1}(\sqrt{-3}-1) - \sqrt{2}(\sqrt{-3}x-x)\right)\sqrt{\sqrt{-3}+1} + 4\right) + \frac{1}{4} \sqrt{2} \sqrt{\sqrt{-3}+1} \log\left(4x^2 - 4\sqrt{x^2+1}x - \left(\sqrt{2}\sqrt{x^2+1}(\sqrt{-3}-1) - \sqrt{2}(\sqrt{-3}x-x)\right)\sqrt{\sqrt{-3}+1} + 4\right)$$

[In] integrate(arctan(x*(x^2+1)^(1/2)),x, algorithm="fricas")

[Out] x*arctan(sqrt(x^2 + 1)*x) - 1/4*sqrt(2)*sqrt(-sqrt(-3) + 1)*log(4*x^2 + sqrt(2)*(sqrt(-3)*x + x)*sqrt(-sqrt(-3) + 1) - sqrt(x^2 + 1)*(sqrt(2)*(sqrt(-3) + 1)*sqrt(-sqrt(-3) + 1) + 4*x) + 4) + 1/4*sqrt(2)*sqrt(-sqrt(-3) + 1)*log(4*x^2 - sqrt(2)*(sqrt(-3)*x + x)*sqrt(-sqrt(-3) + 1) + sqrt(x^2 + 1)*(sqrt(2)*(sqrt(-3) + 1)*sqrt(-sqrt(-3) + 1) - 4*x) + 4) - 1/4*sqrt(2)*sqrt(sqrt(-3) + 1)*log(4*x^2 - 4*sqrt(x^2 + 1)*x + (sqrt(2)*sqrt(x^2 + 1)*(sqrt(-3) - 1) - sqrt(2)*(sqrt(-3)*x - x))*sqrt(sqrt(-3) + 1) + 4) + 1/4*sqrt(2)*sqrt(sqrt(-3) + 1)*log(4*x^2 - 4*sqrt(x^2 + 1)*x - (sqrt(2)*sqrt(x^2 + 1)*(sqrt(-3) - 1) - sqrt(2)*(sqrt(-3)*x - x))*sqrt(sqrt(-3) + 1) + 4)

Sympy [F]

$$\int \arctan(x\sqrt{1+x^2}) dx = \int \operatorname{atan}(x\sqrt{x^2+1}) dx$$

```
[In] integrate(atan(x*(x**2+1)**(1/2)),x)
```

```
[Out] Integral(atan(x*sqrt(x**2 + 1)), x)
```

Maxima [F]

$$\int \arctan(x\sqrt{1+x^2}) dx = \int \arctan(\sqrt{x^2+1}x) dx$$

```
[In] integrate(arctan(x*(x^2+1)^(1/2)),x, algorithm="maxima")
```

```
[Out] x*arctan(sqrt(x^2 + 1)*x) - integrate((2*x^3 + x)*sqrt(x^2 + 1)/((x^4 + x^2)
)*(x^2 + 1) + x^2 + 1), x)
```

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.77

$$\begin{aligned} \int \arctan(x\sqrt{1+x^2}) dx &= x \arctan(\sqrt{x^2+1}x) + \frac{1}{4}\sqrt{3} \log(x^2 + \sqrt{3}\sqrt{x^2+1} + 2) \\ &\quad - \frac{1}{4}\sqrt{3} \log(x^2 - \sqrt{3}\sqrt{x^2+1} + 2) \\ &\quad - \frac{1}{2} \arctan(\sqrt{3} + 2\sqrt{x^2+1}) - \frac{1}{2} \arctan(-\sqrt{3} + 2\sqrt{x^2+1}) \end{aligned}$$

```
[In] integrate(arctan(x*(x^2+1)^(1/2)),x, algorithm="giac")
```

```
[Out] x*arctan(sqrt(x^2 + 1)*x) + 1/4*sqrt(3)*log(x^2 + sqrt(3)*sqrt(x^2 + 1) + 2)
) - 1/4*sqrt(3)*log(x^2 - sqrt(3)*sqrt(x^2 + 1) + 2) - 1/2*arctan(sqrt(3) +
2*sqrt(x^2 + 1)) - 1/2*arctan(-sqrt(3) + 2*sqrt(x^2 + 1))
```

Mupad [B] (verification not implemented)

Time = 1.10 (sec) , antiderivative size = 413, normalized size of antiderivative = 3.44

$$\begin{aligned}
& \int \arctan(x\sqrt{1+x^2}) dx = x \operatorname{atan}(x\sqrt{x^2+1}) \\
& \frac{\left(\ln\left(x - \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) - \ln\left(\frac{x}{2} + \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right)\sqrt{x^2+1} + 1 + \frac{\sqrt{3}x1i}{2}\right)\right) \left(2\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{\sqrt{\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2 + 1} \left(4\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + 1 + \sqrt{3}1i\right)} \\
& \frac{\left(\ln\left(x + \frac{1}{2} - \frac{\sqrt{3}1i}{2}\right) - \ln\left(1 + \left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right)\sqrt{x^2+1} - \frac{x}{2} + \frac{\sqrt{3}x1i}{2}\right)\right) \left(2\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{\sqrt{\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2 + 1} \left(4\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 - 1 + \sqrt{3}1i\right)} \\
& \frac{\left(\ln\left(x - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - \ln\left(\frac{x}{2} + \left(\frac{\sqrt{3}}{2} - \frac{1}{2}i\right)\sqrt{x^2+1} + 1 - \frac{\sqrt{3}x1i}{2}\right)\right) \left(2\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 - \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{\sqrt{\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2 + 1} \left(4\left(-\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 - 1 + \sqrt{3}1i\right)} \\
& \frac{\left(\ln\left(x + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right) - \ln\left(1 + \left(\frac{\sqrt{3}}{2} + \frac{1}{2}i\right)\sqrt{x^2+1} - \frac{x}{2} - \frac{\sqrt{3}x1i}{2}\right)\right) \left(2\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + \frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)}{\sqrt{\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^2 + 1} \left(4\left(\frac{1}{2} + \frac{\sqrt{3}1i}{2}\right)^3 + 1 + \sqrt{3}1i\right)}
\end{aligned}$$

`[In] int(atan(x*(x^2 + 1)^(1/2)),x)`

```

[Out] x*atan(x*(x^2 + 1)^(1/2)) - ((log(x - (3^(1/2)*1i)/2 - 1/2) - log(x/2 + (3^(1/2)/2 + 1i/2)*(x^2 + 1)^(1/2) + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3 + 1/2))/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 + 1/2)^3 + 1)) - ((log(x - (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 - 1i/2)*(x^2 + 1)^(1/2) - x/2 + (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) - ((log(x + (3^(1/2)*1i)/2 - 1/2) - log(x/2 + (3^(1/2)/2 - 1i/2)*(x^2 + 1)^(1/2) - (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 - 1/2)^3 - 1/2))/((((3^(1/2)*1i)/2 - 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 - 1/2)^3 - 1)) - ((log(x + (3^(1/2)*1i)/2 + 1/2) - log((3^(1/2)/2 + 1i/2)*(x^2 + 1)^(1/2) - x/2 - (3^(1/2)*x*1i)/2 + 1))*((3^(1/2)*1i)/2 + 2*((3^(1/2)*1i)/2 + 1/2)^3 + 1/2))/((((3^(1/2)*1i)/2 + 1/2)^2 + 1)^(1/2)*(3^(1/2)*1i + 4*((3^(1/2)*1i)/2 + 1/2)^3 + 1))

```

3.48 $\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx$

Optimal result	279
Rubi [A] (warning: unable to verify)	279
Mathematica [A] (verified)	281
Maple [A] (verified)	281
Fricas [A] (verification not implemented)	281
Sympy [A] (verification not implemented)	282
Maxima [A] (verification not implemented)	282
Giac [A] (verification not implemented)	282
Mupad [B] (verification not implemented)	282

Optimal result

Integrand size = 18, antiderivative size = 31

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{2} - (1+x)\arctan(\sqrt{x} - \sqrt{1+x})$$

[Out] $-(1+x)*\arctan(x^{(1/2)}-(1+x)^{(1/2)})+1/2*x^{(1/2)}$

Rubi [A] (warning: unable to verify)

Time = 0.01 (sec), antiderivative size = 37, normalized size of antiderivative = 1.19, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5267, 8, 4930, 52, 65, 209}

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = -\frac{1}{2}x\arctan(\sqrt{x}) - \frac{\arctan(\sqrt{x})}{2} + \frac{\pi x}{4} + \frac{\sqrt{x}}{2}$$

[In] $\text{Int}[-\text{ArcTan}[\text{Sqrt}[x] - \text{Sqrt}[1 + x]], x]$

[Out] $\text{Sqrt}[x]/2 + (\text{Pi}*x)/4 - \text{ArcTan}[\text{Sqrt}[x]]/2 - (x*\text{ArcTan}[\text{Sqrt}[x]])/2$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 52

$\text{Int}[(a_.) + (b_.)*(x_)^m]*((c_.) + (d_.)*(x_)^n), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{m+1}*((c + d*x)^n/(b*(m+n+1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m+n+1))), \text{Int}[(a + b*x)^m*(c + d*x)^{n-1}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m+n+1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m-n, 0]))) \ \&\& \ !\text{ILtQ}[m+n$

+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 4930

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_.)]*(b_.))^(p_.), x_Symbol] := Simp[x*(a
+ b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p
- 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] &&
(EqQ[n, 1] || EqQ[p, 1])
```

Rule 5267

```
Int[ArcTan[(v_) + (s_.)*Sqrt[w_]]*(u_.), x_Symbol] := Dist[Pi*(s/4), Int[u,
x], x] + Dist[1/2, Int[u*ArcTan[v], x], x] /; EqQ[s^2, 1] && EqQ[w, v^2 +
1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\left(\frac{1}{2} \int \arctan(\sqrt{x}) \, dx\right) + \frac{1}{4} \pi \int 1 \, dx \\
&= \frac{\pi x}{4} - \frac{1}{2} x \arctan(\sqrt{x}) + \frac{1}{4} \int \frac{\sqrt{x}}{1+x} \, dx \\
&= \frac{\sqrt{x}}{2} + \frac{\pi x}{4} - \frac{1}{2} x \arctan(\sqrt{x}) - \frac{1}{4} \int \frac{1}{\sqrt{x}(1+x)} \, dx \\
&= \frac{\sqrt{x}}{2} + \frac{\pi x}{4} - \frac{1}{2} x \arctan(\sqrt{x}) - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x^2} \, dx, x, \sqrt{x}\right) \\
&= \frac{\sqrt{x}}{2} + \frac{\pi x}{4} - \frac{\arctan(\sqrt{x})}{2} - \frac{1}{2} x \arctan(\sqrt{x})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int -\arctan\left(\sqrt{x} - \sqrt{1+x}\right) dx = \frac{\sqrt{x}}{2} - (1+x)\arctan\left(\sqrt{x} - \sqrt{1+x}\right)$$

[In] Integrate[-ArcTan[Sqrt[x] - Sqrt[1 + x]],x]

[Out] Sqrt[x]/2 - (1 + x)*ArcTan[Sqrt[x] - Sqrt[1 + x]]

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
default	$-x \arctan\left(\sqrt{x} - \sqrt{1+x}\right) + \frac{\sqrt{x}}{2} - \frac{\arctan(\sqrt{x})}{2}$	28
parts	$-x \arctan\left(\sqrt{x} - \sqrt{1+x}\right) + \frac{\sqrt{x}}{2} - \frac{\arctan(\sqrt{x})}{2}$	28

[In] int(-arctan(x^(1/2)-(1+x)^(1/2)),x,method=_RETURNVERBOSE)

[Out] -x*arctan(x^(1/2)-(1+x)^(1/2))+1/2*x^(1/2)-1/2*arctan(x^(1/2))

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.71

$$\int -\arctan\left(\sqrt{x} - \sqrt{1+x}\right) dx = (x+1)\arctan\left(\sqrt{x+1} - \sqrt{x}\right) + \frac{1}{2}\sqrt{x}$$

[In] integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="fricas")

[Out] (x + 1)*arctan(sqrt(x + 1) - sqrt(x)) + 1/2*sqrt(x)

Sympy [A] (verification not implemented)

Time = 9.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = \frac{\sqrt{x}}{2} - x \operatorname{atan}(\sqrt{x} - \sqrt{x+1}) - \frac{\operatorname{atan}(\sqrt{x})}{2}$$

[In] integrate(-atan(x**(1/2)-(1+x)**(1/2)),x)

[Out] sqrt(x)/2 - x*atan(sqrt(x) - sqrt(x + 1)) - atan(sqrt(x))/2

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = x \arctan(\sqrt{x+1} - \sqrt{x}) + \frac{1}{2} \sqrt{x} - \frac{1}{2} \arctan(\sqrt{x})$$

[In] integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="maxima")

[Out] x*arctan(sqrt(x + 1) - sqrt(x)) + 1/2*sqrt(x) - 1/2*arctan(sqrt(x))

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = -x \arctan(-\sqrt{x+1} + \sqrt{x}) + \frac{1}{2} \sqrt{x} - \frac{1}{2} \arctan(\sqrt{x})$$

[In] integrate(-arctan(x^(1/2)-(1+x)^(1/2)),x, algorithm="giac")

[Out] -x*arctan(-sqrt(x + 1) + sqrt(x)) + 1/2*sqrt(x) - 1/2*arctan(sqrt(x))

Mupad [B] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.29

$$\int -\arctan(\sqrt{x} - \sqrt{1+x}) dx = x \operatorname{atan}(\sqrt{x+1} - \sqrt{x}) + \frac{\sqrt{x}}{2} - \frac{\ln\left(\frac{(-1+\sqrt{x}1i)^2}{x+1}\right) 1i}{4}$$

[In] int(atan((x + 1)^(1/2) - x^(1/2)),x)

[Out] x*atan((x + 1)^(1/2) - x^(1/2)) - (log((x^(1/2)*1i - 1)^2/(x + 1))*1i)/4 + x^(1/2)/2

3.49 $\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx$

Optimal result	283
Rubi [A] (verified)	283
Mathematica [A] (verified)	284
Maple [B] (verified)	285
Fricas [B] (verification not implemented)	285
Sympy [F]	286
Maxima [F]	286
Giac [A] (verification not implemented)	286
Mupad [F(-1)]	286

Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) + \arctan\left(\sqrt{1-2x^2}\right)$$

[Out] $x \arcsin(x/(-x^2+1)^{(1/2)}) + \arctan((-2x^2+1)^{(1/2)})$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {4924, 455, 65, 209}

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) + \arctan\left(\sqrt{1-2x^2}\right)$$

[In] `Int[ArcSin[x/Sqrt[1 - x^2]],x]`

[Out] `x*ArcSin[x/Sqrt[1 - x^2]] + ArcTan[Sqrt[1 - 2*x^2]]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]
```

Rule 4924

```
Int[ArcSin[u_], x_Symbol] := Simp[x*ArcSin[u], x] - Int[SimplifyIntegrand[x*(D[u, x]/Sqrt[1 - u^2]), x], x] /; InverseFunctionFreeQ[u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) - \int \frac{x}{\sqrt{1-2x^2}(1-x^2)} dx \\
 &= x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1-2x}(1-x)} dx, x, x^2\right) \\
 &= x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{\frac{1}{2} + \frac{x^2}{2}} dx, x, \sqrt{1-2x^2}\right) \\
 &= x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) + \arctan\left(\sqrt{1-2x^2}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = x \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) + \arctan\left(\sqrt{1-2x^2}\right)$$

```
[In] Integrate[ArcSin[x/Sqrt[1 - x^2]],x]
```

```
[Out] x*ArcSin[x/Sqrt[1 - x^2]] + ArcTan[Sqrt[1 - 2*x^2]]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 137 vs. $2(25) = 50$.

Time = 0.19 (sec) , antiderivative size = 138, normalized size of antiderivative = 4.76

method	result
default	$x \arcsin\left(\frac{x}{\sqrt{-x^2+1}}\right) + \frac{\sqrt{\frac{2x^2-1}{x^2-1}} \left(\sqrt{-2x^2+1} + \arctan\left(\frac{2x-1}{\sqrt{-2x^2+1}}\right) - \arctan\left(\frac{1+2x}{\sqrt{-2x^2+1}}\right) \right) \sqrt{-x^2+1}}{\sqrt{-2x^2+1} (2+\sqrt{2}) (-2+\sqrt{2})} + \frac{\sqrt{\frac{2x^2-1}{x^2-1}} \sqrt{-x^2+1}}{2}$
parts	$x \arcsin\left(\frac{x}{\sqrt{-x^2+1}}\right) + \frac{\sqrt{\frac{2x^2-1}{x^2-1}} \left(\sqrt{-2x^2+1} + \arctan\left(\frac{2x-1}{\sqrt{-2x^2+1}}\right) - \arctan\left(\frac{1+2x}{\sqrt{-2x^2+1}}\right) \right) \sqrt{-x^2+1}}{\sqrt{-2x^2+1} (2+\sqrt{2}) (-2+\sqrt{2})} + \frac{\sqrt{\frac{2x^2-1}{x^2-1}} \sqrt{-x^2+1}}{2}$

[In] `int(arcsin(1/(-x^2+1)^(1/2)*x),x,method=_RETURNVERBOSE)`

[Out] `x*arcsin(1/(-x^2+1)^(1/2)*x)+((2*x^2-1)/(x^2-1))^(1/2)*((-2*x^2+1)^(1/2)+arctan((2*x-1)/(-2*x^2+1)^(1/2))-arctan((1+2*x)/(-2*x^2+1)^(1/2)))*(-x^2+1)^(1/2)/(-2*x^2+1)^(1/2)/(2+2^(1/2))/(-2+2^(1/2))+1/2*((2*x^2-1)/(x^2-1))^(1/2)*(-x^2+1)^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(25) = 50$.

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.07

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx$$

$$= -x \arcsin\left(\frac{\sqrt{-x^2+1}x}{x^2-1}\right) + \arctan\left(\frac{x^2 + \sqrt{-x^2+1}\sqrt{\frac{2x^2-1}{x^2-1}} - 1}{x^2}\right)$$

[In] `integrate(arcsin(x/(-x^2+1)^(1/2)),x, algorithm="fricas")`

[Out] `-x*arcsin(sqrt(-x^2+1)*x/(x^2-1))+arctan((x^2+sqrt(-x^2+1)*sqrt((2*x^2-1)/(x^2-1))-1)/x^2)`

Sympy [F]

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = \int \operatorname{asin}\left(\frac{x}{\sqrt{1-x^2}}\right) dx$$

[In] integrate(asin(x/(-x**2+1)**(1/2)),x)

[Out] Integral(asin(x/sqrt(1 - x**2)), x)

Maxima [F]

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = \int \arcsin\left(\frac{x}{\sqrt{-x^2+1}}\right) dx$$

[In] integrate(arcsin(x/(-x^2+1)^(1/2)),x, algorithm="maxima")

[Out] integrate(arcsin(x/sqrt(-x^2 + 1)), x)

Giac [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = x \arcsin\left(\frac{x}{\sqrt{-x^2+1}}\right) + \frac{\arctan(\sqrt{-2x^2+1})}{\operatorname{sgn}(x^2-1)}$$

[In] integrate(arcsin(x/(-x^2+1)^(1/2)),x, algorithm="giac")

[Out] x*arcsin(x/sqrt(-x^2 + 1)) + arctan(sqrt(-2*x^2 + 1))/sgn(x^2 - 1)

Mupad [F(-1)]

Timed out.

$$\int \arcsin\left(\frac{x}{\sqrt{1-x^2}}\right) dx = \int \operatorname{asin}\left(\frac{x}{\sqrt{1-x^2}}\right) dx$$

[In] int(asin(x/(1 - x^2)^(1/2)),x)

[Out] int(asin(x/(1 - x^2)^(1/2)), x)

3.50 $\int \arctan(x\sqrt{1-x^2}) dx$

Optimal result	287
Rubi [A] (verified)	287
Mathematica [A] (verified)	289
Maple [B] (verified)	290
Fricas [B] (verification not implemented)	290
Sympy [F(-1)]	291
Maxima [F]	292
Giac [A] (verification not implemented)	292
Mupad [B] (verification not implemented)	293

Optimal result

Integrand size = 14, antiderivative size = 106

$$\int \arctan(x\sqrt{1-x^2}) dx = -\sqrt{\frac{1}{2}(1+\sqrt{5})} \arctan\left(\sqrt{\frac{1}{2}(1+\sqrt{5})}\sqrt{1-x^2}\right) \\ + x \arctan(x\sqrt{1-x^2}) \\ + \sqrt{\frac{1}{2}(-1+\sqrt{5})} \operatorname{arctanh}\left(\sqrt{\frac{1}{2}(-1+\sqrt{5})}\sqrt{1-x^2}\right)$$

[Out] x*arctan(x*(-x^2+1)^(1/2))+1/2*arctanh(1/2*(-x^2+1)^(1/2)*(-2+2*5^(1/2))^(1/2))*(-2+2*5^(1/2))^(1/2)-1/2*arctan(1/2*(-x^2+1)^(1/2)*(2+2*5^(1/2))^(1/2))* (2+2*5^(1/2))^(1/2)

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5311, 1699, 840, 1180, 210, 212}

$$\int \arctan(x\sqrt{1-x^2}) dx = -\sqrt{\frac{2}{\sqrt{5}-1}} \arctan\left(\sqrt{\frac{2}{\sqrt{5}-1}}\sqrt{1-x^2}\right) \\ + x \arctan(x\sqrt{1-x^2}) \\ + \sqrt{\frac{2}{1+\sqrt{5}}} \operatorname{arctanh}\left(\sqrt{\frac{2}{1+\sqrt{5}}}\sqrt{1-x^2}\right)$$

[In] Int[ArcTan[x*Sqrt[1 - x^2]],x]

[Out] $-(\sqrt{2/(-1 + \sqrt{5})}) \cdot \text{ArcTan}[\sqrt{2/(-1 + \sqrt{5})}] \cdot \sqrt{1 - x^2}] + x \cdot \text{ArcTan}[x \cdot \sqrt{1 - x^2}] + \sqrt{2/(1 + \sqrt{5})} \cdot \text{ArcTanh}[\sqrt{2/(1 + \sqrt{5})}] \cdot \sqrt{1 - x^2}]$

Rule 210

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1}] \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 212

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 840

$\text{Int}[(f + (g \cdot x))/(\sqrt{(d + (e \cdot x))^2} \cdot ((a + (b \cdot x) + (c \cdot x)^2))], x_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[(e \cdot f - d \cdot g + g \cdot x^2)/(c \cdot d^2 - b \cdot d \cdot e + a \cdot e^2 - (2 \cdot c \cdot d - b \cdot e) \cdot x^2 + c \cdot x^4), x], x, \sqrt{d + e \cdot x}], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, x\} \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - b \cdot d \cdot e + a \cdot e^2, 0]$

Rule 1180

$\text{Int}[(d + (e \cdot x)^2)/((a + (b \cdot x)^2 + (c \cdot x)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4 \cdot a \cdot c, 2]\}, \text{Dist}[e/2 + (2 \cdot c \cdot d - b \cdot e)/(2 \cdot q), \text{Int}[1/(b/2 - q/2 + c \cdot x^2), x], x] + \text{Dist}[e/2 - (2 \cdot c \cdot d - b \cdot e)/(2 \cdot q), \text{Int}[1/(b/2 + q/2 + c \cdot x^2), x], x]] /; \text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0] \ \&\& \ \text{NeQ}[c \cdot d^2 - a \cdot e^2, 0] \ \&\& \ \text{PosQ}[b^2 - 4 \cdot a \cdot c]$

Rule 1699

$\text{Int}[(P_x) \cdot (x) \cdot ((d + (e \cdot x)^2)^{(q \cdot x)) \cdot ((a + (b \cdot x)^2 + (c \cdot x)^4)^{(p \cdot x))}, x_Symbol] \rightarrow \text{Dist}[1/2, \text{Subst}[\text{Int}[(P_x / x \rightarrow \sqrt{x}) \cdot (d + e \cdot x)^{q \cdot (a + b \cdot x + c \cdot x^2)^p}], x, x^2], x] /; \text{FreeQ}\{a, b, c, d, e, p, q, x\} \ \&\& \ \text{PolyQ}[P_x, x^2]$

Rule 5311

$\text{Int}[\text{ArcTan}[u], x_Symbol] \rightarrow \text{Simp}[x \cdot \text{ArcTan}[u], x] - \text{Int}[\text{SimplifyIntegrand}[x \cdot (D[u, x]/(1 + u^2))], x], x] /; \text{InverseFunctionFreeQ}[u, x]$

Rubi steps

$$\begin{aligned}
\text{integral} &= x \arctan \left(x\sqrt{1-x^2} \right) - \int \frac{x(1-2x^2)}{\sqrt{1-x^2}(1+x^2-x^4)} dx \\
&= x \arctan \left(x\sqrt{1-x^2} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1-2x}{\sqrt{1-x}(1+x-x^2)} dx, x, x^2 \right) \\
&= x \arctan \left(x\sqrt{1-x^2} \right) - \text{Subst} \left(\int \frac{1-2x^2}{1+x^2-x^4} dx, x, \sqrt{1-x^2} \right) \\
&= x \arctan \left(x\sqrt{1-x^2} \right) + \text{Subst} \left(\int \frac{1}{\frac{1}{2} - \frac{\sqrt{5}}{2} - x^2} dx, x, \sqrt{1-x^2} \right) \\
&\quad + \text{Subst} \left(\int \frac{1}{\frac{1}{2} + \frac{\sqrt{5}}{2} - x^2} dx, x, \sqrt{1-x^2} \right) \\
&= -\sqrt{\frac{2}{-1+\sqrt{5}}} \arctan \left(\sqrt{\frac{2}{-1+\sqrt{5}}} \sqrt{1-x^2} \right) \\
&\quad + x \arctan \left(x\sqrt{1-x^2} \right) + \sqrt{\frac{2}{1+\sqrt{5}}} \operatorname{arctanh} \left(\sqrt{\frac{2}{1+\sqrt{5}}} \sqrt{1-x^2} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.00

$$\frac{\int \arctan \left(x\sqrt{1-x^2} \right) dx = x \arctan \left(x\sqrt{1-x^2} \right) + \frac{\sqrt{1+\sqrt{5}} \arctan \left(\sqrt{\frac{1}{2}(1+\sqrt{5})} \sqrt{1-x^2} \right) - \sqrt{-1+\sqrt{5}} \operatorname{arctanh} \left(\sqrt{\frac{1}{2}(-1+\sqrt{5})} \sqrt{1-x^2} \right)}{\sqrt{2}}}{\sqrt{2}}$$

[In] Integrate[ArcTan[x*Sqrt[1 - x^2]],x]

[Out] x*ArcTan[x*Sqrt[1 - x^2]] - (Sqrt[1 + Sqrt[5]]*ArcTan[Sqrt[(1 + Sqrt[5])/2]*Sqrt[1 - x^2]] - Sqrt[-1 + Sqrt[5]]*ArcTanh[Sqrt[(-1 + Sqrt[5])/2]*Sqrt[1 - x^2]])/Sqrt[2]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. $2(79) = 158$.

Time = 0.10 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.98

method	result
default	$x \arctan(x\sqrt{-x^2+1}) + \frac{\sqrt{5} \arctan\left(\frac{2(\sqrt{-x^2+1}-1)^2}{x^2} - 2\sqrt{5}+4\right)}{5\sqrt{-2+\sqrt{5}}} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{2(\sqrt{-x^2+1}-1)^2}{x^2} + 4+2\sqrt{5}\right)}{5\sqrt{2+\sqrt{5}}} + \frac{\left(-\frac{1}{2} + \frac{3\sqrt{5}}{10}\right)}{5\sqrt{-2+\sqrt{5}}}$
parts	$x \arctan(x\sqrt{-x^2+1}) + \frac{\sqrt{5} \arctan\left(\frac{2(\sqrt{-x^2+1}-1)^2}{x^2} - 2\sqrt{5}+4\right)}{5\sqrt{-2+\sqrt{5}}} + \frac{\sqrt{5} \operatorname{arctanh}\left(\frac{2(\sqrt{-x^2+1}-1)^2}{x^2} + 4+2\sqrt{5}\right)}{5\sqrt{2+\sqrt{5}}} + \frac{\left(-\frac{1}{2} + \frac{3\sqrt{5}}{10}\right)}{5\sqrt{-2+\sqrt{5}}}$

[In] `int(arctan(x*(-x^2+1)^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `x*arctan(x*(-x^2+1)^(1/2))+1/5*5^(1/2)/(-2+5^(1/2))^(1/2)*arctan(1/4*(2*((-x^2+1)^(1/2)-1)^2/x^2-2*5^(1/2)+4)/(-2+5^(1/2))^(1/2))+1/5*5^(1/2)/(2+5^(1/2))^(1/2)*arctanh(1/4*(2*((-x^2+1)^(1/2)-1)^2/x^2+4+2*5^(1/2))/(2+5^(1/2))^(1/2))+(-1/2+3/10*5^(1/2))/(-2+5^(1/2))^(1/2)*arctan(1/4*(2*((-x^2+1)^(1/2)-1)^2/x^2-2*5^(1/2)+4)/(-2+5^(1/2))^(1/2))+(1/2+3/10*5^(1/2))/(2+5^(1/2))^(1/2)*arctanh(1/4*(2*((-x^2+1)^(1/2)-1)^2/x^2+4+2*5^(1/2))/(2+5^(1/2))^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 205 vs. $2(79) = 158$.

Time = 0.27 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.93

$$\int \arctan \left(x\sqrt{1-x^2} \right) dx = x \arctan \left(\sqrt{-x^2+1}x \right) + \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log \left(\left(\sqrt{5}\sqrt{2} + \sqrt{2} \right) \sqrt{\sqrt{5}-1} + 4\sqrt{-x^2+1} \right) - \frac{1}{4} \sqrt{2} \sqrt{\sqrt{5}-1} \log \left(- \left(\sqrt{5}\sqrt{2} + \sqrt{2} \right) \sqrt{\sqrt{5}-1} + 4\sqrt{-x^2+1} \right) - \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log \left(\left(\sqrt{5}\sqrt{2} - \sqrt{2} \right) \sqrt{-\sqrt{5}-1} + 4\sqrt{-x^2+1} \right) + \frac{1}{4} \sqrt{2} \sqrt{-\sqrt{5}-1} \log \left(- \left(\sqrt{5}\sqrt{2} - \sqrt{2} \right) \sqrt{-\sqrt{5}-1} + 4\sqrt{-x^2+1} \right)$$

[In] integrate(arctan(x*(-x^2+1)^(1/2)),x, algorithm="fricas")

[Out] x*arctan(sqrt(-x^2 + 1)*x) + 1/4*sqrt(2)*sqrt(sqrt(5) - 1)*log((sqrt(5)*sqrt(2) + sqrt(2))*sqrt(sqrt(5) - 1) + 4*sqrt(-x^2 + 1)) - 1/4*sqrt(2)*sqrt(sqrt(5) - 1)*log(-(sqrt(5)*sqrt(2) + sqrt(2))*sqrt(sqrt(5) - 1) + 4*sqrt(-x^2 + 1)) - 1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log((sqrt(5)*sqrt(2) - sqrt(2))*sqrt(-sqrt(5) - 1) + 4*sqrt(-x^2 + 1)) + 1/4*sqrt(2)*sqrt(-sqrt(5) - 1)*log(-(sqrt(5)*sqrt(2) - sqrt(2))*sqrt(-sqrt(5) - 1) + 4*sqrt(-x^2 + 1))

Sympy [F(-1)]

Timed out.

$$\int \arctan \left(x\sqrt{1-x^2} \right) dx = \text{Timed out}$$

[In] integrate(atan(x*(-x**2+1)**(1/2)),x)

[Out] Timed out

Maxima [F]

$$\int \arctan(x\sqrt{1-x^2}) dx = \int \arctan(\sqrt{-x^2+1}x) dx$$

[In] integrate(arctan(x*(-x^2+1)^(1/2)),x, algorithm="maxima")

[Out] x*arctan(sqrt(x + 1)*x*sqrt(-x + 1)) - integrate((2*x^3 - x)*e^(1/2*log(x + 1) + 1/2*log(-x + 1))/(x^2 + (x^4 - x^2)*e^(log(x + 1) + log(-x + 1)) - 1), x)

Giac [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.05

$$\int \arctan(x\sqrt{1-x^2}) dx = x \arctan(\sqrt{-x^2+1}x) - \frac{1}{2} \sqrt{2\sqrt{5}} + 2 \arctan\left(\frac{\sqrt{-x^2+1}}{\sqrt{\frac{1}{2}\sqrt{5}-\frac{1}{2}}}\right) + \frac{1}{4} \sqrt{2\sqrt{5}} - 2 \log\left(\sqrt{-x^2+1} + \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}}\right) - \frac{1}{4} \sqrt{2\sqrt{5}} - 2 \log\left(\left|\sqrt{-x^2+1} - \sqrt{\frac{1}{2}\sqrt{5} + \frac{1}{2}}\right|\right)$$

[In] integrate(arctan(x*(-x^2+1)^(1/2)),x, algorithm="giac")

[Out] x*arctan(sqrt(-x^2 + 1)*x) - 1/2*sqrt(2*sqrt(5) + 2)*arctan(sqrt(-x^2 + 1)/sqrt(1/2*sqrt(5) - 1/2)) + 1/4*sqrt(2*sqrt(5) - 2)*log(sqrt(-x^2 + 1) + sqrt(1/2*sqrt(5) + 1/2)) - 1/4*sqrt(2*sqrt(5) - 2)*log(abs(sqrt(-x^2 + 1) - sqrt(1/2*sqrt(5) + 1/2)))

Mupad [B] (verification not implemented)

Time = 1.15 (sec) , antiderivative size = 455, normalized size of antiderivative = 4.29

$$\begin{aligned}
\int \arctan(x\sqrt{1-x^2}) dx &= x \operatorname{atan}(x\sqrt{1-x^2}) \\
&+ \frac{\ln\left(\frac{\left(x\sqrt{\frac{\sqrt{5}+1}{2}-1}\right)^{1i} - \sqrt{1-x^2} 1i}{\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}}\right)}{x - \sqrt{\frac{\sqrt{5}+1}{2}}} \left(\sqrt{\frac{\sqrt{5}+1}{2}} - 2\left(\frac{\sqrt{5}}{2} + \frac{1}{2}\right)^{3/2}\right) \\
&+ \frac{\left(2\sqrt{\frac{\sqrt{5}+1}{2}} - 4\left(\frac{\sqrt{5}}{2} + \frac{1}{2}\right)^{3/2}\right) \sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}}{\left(2\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 4\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}} \\
&\ln\left(\frac{\left(x\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}-1\right)^{1i} - \sqrt{1-x^2} 1i}{\sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}}\right) \left(\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 2\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \\
&+ \frac{\left(2\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 4\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}}{\left(2\sqrt{\frac{\sqrt{5}+1}{2}} - 4\left(\frac{\sqrt{5}}{2} + \frac{1}{2}\right)^{3/2}\right) \sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}} \\
&\ln\left(\frac{\left(x\sqrt{\frac{\sqrt{5}+1}{2}+1}\right)^{1i} + \sqrt{1-x^2} 1i}{\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}}\right) \left(\sqrt{\frac{\sqrt{5}+1}{2}} - 2\left(\frac{\sqrt{5}}{2} + \frac{1}{2}\right)^{3/2}\right) \\
&+ \frac{\left(2\sqrt{\frac{\sqrt{5}+1}{2}} - 4\left(\frac{\sqrt{5}}{2} + \frac{1}{2}\right)^{3/2}\right) \sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}}{\left(2\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 4\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}} \\
&\ln\left(\frac{\left(x\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}}+1\right)^{1i} + \sqrt{1-x^2} 1i}{\sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}}\right) \left(\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 2\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \\
&+ \frac{\left(2\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 4\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}}{\left(2\sqrt{\frac{1}{2}-\frac{\sqrt{5}}{2}} - 4\left(\frac{1}{2} - \frac{\sqrt{5}}{2}\right)^{3/2}\right) \sqrt{\frac{\sqrt{5}}{2} + \frac{1}{2}}}
\end{aligned}$$

[In] int(atan(x*(1-x^2)^(1/2)),x)

```

[Out] x*atan(x*(1-x^2)^(1/2)) + (log((((x*(5^(1/2)/2 + 1/2)^(1/2) - 1)*1i)/(1/2
- 5^(1/2)/2)^(1/2) - (1-x^2)^(1/2)*1i)/(x - (5^(1/2)/2 + 1/2)^(1/2)))*((
5^(1/2)/2 + 1/2)^(1/2) - 2*(5^(1/2)/2 + 1/2)^(3/2)))/((2*(5^(1/2)/2 + 1/2)^(
1/2) - 4*(5^(1/2)/2 + 1/2)^(3/2))*(1/2 - 5^(1/2)/2)^(1/2)) + (log((((x*(1/
2 - 5^(1/2)/2)^(1/2) - 1)*1i)/(5^(1/2)/2 + 1/2)^(1/2) - (1-x^2)^(1/2)*1i)
/(x - (1/2 - 5^(1/2)/2)^(1/2)))*((1/2 - 5^(1/2)/2)^(1/2) - 2*(1/2 - 5^(1/2)
/2)^(3/2)))/((2*(1/2 - 5^(1/2)/2)^(1/2) - 4*(1/2 - 5^(1/2)/2)^(3/2))*(5^(1/
2)/2 + 1/2)^(1/2)) + (log((((x*(5^(1/2)/2 + 1/2)^(1/2) + 1)*1i)/(1/2 - 5^(1

```

$$\begin{aligned} & /2)/2)^{(1/2)} + (1 - x^2)^{(1/2)} * 1i) / (x + (5^{(1/2)}/2 + 1/2)^{(1/2)}) * ((5^{(1/2)}/2 + 1/2)^{(1/2)} - 2 * (5^{(1/2)}/2 + 1/2)^{(3/2)}) / ((2 * (5^{(1/2)}/2 + 1/2)^{(1/2)} - 4 * (5^{(1/2)}/2 + 1/2)^{(3/2)}) * (1/2 - 5^{(1/2)}/2)^{(1/2)}) + (\log((((x * (1/2 - 5^{(1/2)}/2)^{(1/2)} + 1) * 1i) / (5^{(1/2)}/2 + 1/2)^{(1/2)} + (1 - x^2)^{(1/2)} * 1i) / (x + (1/2 - 5^{(1/2)}/2)^{(1/2)})) * ((1/2 - 5^{(1/2)}/2)^{(1/2)} - 2 * (1/2 - 5^{(1/2)}/2)^{(3/2)})) / ((2 * (1/2 - 5^{(1/2)}/2)^{(1/2)} - 4 * (1/2 - 5^{(1/2)}/2)^{(3/2)}) * (5^{(1/2)}/2 + 1/2)^{(1/2)}) \end{aligned}$$

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 295

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```



```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```
def grade_antiderivative(result,optimal):
```

```

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

```

```

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

```

```

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

```

```

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```
if str(result).find("Integral") != -1:
```

```

    grade = "F"
    grade_annotation = ""

```

```
else:
```

```
if expnType_result <= expnType_optimal:
```

```
if result.has(I):
```

```
if optimal.has(I): #both result and optimal complex
```

```
if leaf_count_result <= 2*leaf_count_optimal:
```

```

    grade = "A"
    grade_annotation = ""

```

```
else:
```

```
grade = "B"
```

```
grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
```

```
else: #result contains complex but optimal is not
```

```
grade = "C"
```

```
grade_annotation = "Result contains complex when optimal does not."
```

```
else: # result do not contain complex, this assumes optimal do not as well
```

```
if leaf_count_result <= 2*leaf_count_optimal:
```

```

    grade = "A"
    grade_annotation = ""

```

```
else:
```

```
grade = "B"
```

```
grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
```

```
else:
```

```
grade = "C"
```

```
grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)
```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #instance(expr, Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or inst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```