

# Computer Algebra Independent Integration Tests

Summer 2023 edition

1-Algebraic-functions/1.2-Trinomial-products/1.2.1-Quadratic/32-  
1.2.1.1-a+b-x+c-x<sup>2</sup>-<sup>p</sup>

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 11:22am

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
<b>2</b>	<b>detailed summary tables of results</b>	<b>21</b>
<b>3</b>	<b>Listing of integrals</b>	<b>61</b>
<b>4</b>	<b>Appendix</b>	<b>667</b>

---

---

# CHAPTER 1

---

## INTRODUCTION

1.1	Listing of CAS systems tested . . . . .	4
1.2	Results . . . . .	5
1.3	Time and leaf size Performance . . . . .	8
1.4	Performance based on number of rules Rubi used . . . . .	10
1.5	Performance based on number of steps Rubi used . . . . .	11
1.6	Solved integrals histogram based on leaf size of result . . . . .	12
1.7	Solved integrals histogram based on CPU time used . . . . .	13
1.8	Leaf size vs. CPU time used . . . . .	14
1.9	list of integrals with no known antiderivative . . . . .	15
1.10	List of integrals solved by CAS but has no known antiderivative . . . . .	15
1.11	list of integrals solved by CAS but failed verification . . . . .	15
1.12	Timing . . . . .	16
1.13	Verification . . . . .	16
1.14	Important notes about some of the results . . . . .	16
1.15	Design of the test system . . . . .	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [ 143 ]. This is test number [ 32 ].

## 1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

## 1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 ( 143 )	0.00 ( 0 )
Mathematica	100.00 ( 143 )	0.00 ( 0 )
Mupad	92.31 ( 132 )	7.69 ( 11 )
Maple	79.02 ( 113 )	20.98 ( 30 )
Fricas	79.02 ( 113 )	20.98 ( 30 )
Giac	77.62 ( 111 )	22.38 ( 32 )
Maxima	75.52 ( 108 )	24.48 ( 35 )
Sympy	70.63 ( 101 )	29.37 ( 42 )

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> <li>1. antiderivative contains a hypergeometric function and the optimal antiderivative does not.</li> <li>2. antiderivative contains a special function and the optimal antiderivative does not.</li> <li>3. antiderivative contains the imaginary unit and the optimal antiderivative does not.</li> </ol>
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	99.301	0.699	0.000	0.000
Mathematica	76.923	10.490	12.587	0.000
Maple	68.531	7.692	2.797	20.979
Maxima	67.832	4.895	2.797	24.476
Giac	55.944	18.881	2.797	22.378
Fricas	52.448	22.378	4.196	20.979
Sympy	48.252	18.881	3.497	29.371
Mupad	0.000	92.308	0.000	7.692

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Mupad	11	0.00	100.00	0.00
Fricas	30	100.00	0.00	0.00
Maple	30	100.00	0.00	0.00
Giac	32	100.00	0.00	0.00
Maxima	35	85.71	0.00	14.29
Sympy	42	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

## 1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.



System	Mean time (sec)
Rubi	0.05
Maxima	0.24
Giac	0.27
Fricas	0.32
Sympy	0.33
Mathematica	1.34
Maple	2.21
Mupad	6.26

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	39.27	0.84	33.50	0.79
Maple	41.17	0.96	31.00	0.83
Maxima	45.49	1.05	33.50	1.03
Mathematica	50.11	1.24	44.00	1.00
Giac	54.18	1.40	37.00	1.05
Fricas	63.38	1.57	43.00	1.21
Rubi	83.41	1.02	38.00	1.00
Sympy	87.38	1.83	41.00	1.06

Table 1.6: Leaf size performance for each CAS

## 1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the  $y$  axis is the percentage solved which Rubi itself needed the number of rules given the  $x$  axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

## 1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

## 1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

## 1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

## 1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

## 1.9 list of integrals with no known antiderivative

{}

## 1.10 List of integrals solved by CAS but has no known antiderivative

**Rubi** {}

**Mathematica** {}

**Maple** {}

**Maxima** {}

**Fricas** {}

**Sympy** {}

**Giac** {}

**Mupad** {}

## 1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

**Rubi** {}

**Mathematica** {}

**Maple** {65,68}

**Maxima** Verification phase not currently implemented.

**Fricas** Verification phase not currently implemented.

**Sympy** Verification phase not currently implemented.

**Giac** Verification phase not currently implemented.

**Mupad** Verification phase not currently implemented.

## 1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

## 1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

## 1.14 Important notes about some of the results

### Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.



Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

## Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

## Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at [https://ask.sagemath.org/question/57123/could-we-have-a-leaf\\_count-function-in-base-sagemath/](https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/)

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

## Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives  $\sin(x)^2/2$

## 1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



**High level overview of the CAS independent integration test build system**

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

*The following fields are present only in Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi  
June 27, 2023  
Design-vide



---

---

## CHAPTER 2

---

### DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS . . . . .	22
2.2	Detailed conclusion table per each integral for all CAS systems . . . . .	26
2.3	Detailed conclusion table specific for Rubi results . . . . .	55

## 2.1 List of integrals sorted by grade for each CAS

Rubi . . . . .	22
Mma . . . . .	22
Maple . . . . .	23
Fricas . . . . .	23
Maxima . . . . .	23
Giac . . . . .	24
Mupad . . . . .	24
Sympy . . . . .	25

### Rubi

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143 }

**B grade** { 83 }

**C grade** { }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

### Mma

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 22, 23, 24, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 116, 117, 119, 120, 121, 122, 123, 125, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143 }

**B grade** { 17, 21, 25, 26, 27, 28, 29, 83, 99, 100, 115, 118, 124, 126, 127 }

**C grade** { 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47 }

**F normal fail** { }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## Maple

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 66, 67, 69, 70, 72, 78, 79, 80, 81, 82, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 128, 129, 130, 131, 138 }

**B grade** { 25, 74, 75, 76, 77, 83, 101, 102, 125, 126, 127 }

**C grade** { 65, 68, 71, 73 }

**F normal fail** { 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

## Fricas

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 19, 20, 22, 23, 24, 26, 28, 29, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 78, 79, 81, 82, 86, 88, 89, 91, 92, 93, 94, 98, 101, 102, 103, 105, 106, 107, 108, 109, 110, 112, 113, 116, 119, 121, 123, 130, 131, 138 }

**B grade** { 17, 18, 21, 25, 27, 74, 75, 76, 77, 80, 83, 84, 85, 87, 90, 95, 96, 97, 99, 100, 104, 114, 115, 117, 118, 120, 124, 125, 126, 127, 128, 129 }

**C grade** { 70, 71, 72, 73, 111, 122 }

**F normal fail** { 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }

**F(-1) timeout fail** { }

**F(-2) exception fail** { }

## Maxima

**A grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 72, 78, 79, 80, 81, 82, 84, 85, 86, 87, 90, 91, 92, 93, 94, 97, 99, 100, 102, 103, 104, 105, 106, 107, 108, 109, 110, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 123, 124, 125, 126, 127, 128, 129, 130, 131, 138 }

**B grade** { 17, 74, 75, 76, 77, 83, 101 }

**C grade** { 71, 73, 111, 122 }

**F normal fail** { 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }

**F(-1) timeout fail { }**

**F(-2) exception fail { 88, 89, 95, 96, 98 }**

## **Giac**

**A grade { 1, 2, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 19, 20, 22, 23, 24, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 78, 79, 80, 81, 82, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 102, 103, 104, 105, 106, 107, 108, 109, 110, 112, 113, 116, 119, 121, 123, 128, 129, 130, 131, 138 }**

**B grade { 3, 4, 5, 17, 18, 21, 25, 26, 27, 28, 29, 64, 74, 75, 76, 77, 83, 101, 114, 115, 117, 118, 120, 124, 125, 126, 127 }**

**C grade { 70, 71, 72, 73 }**

**F normal fail { 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 111, 122, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }**

**F(-1) timeout fail { }**

**F(-2) exception fail { }**

## **Mupad**

**A grade { }**

**B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 138 }**

**C grade { }**

**F normal fail { }**

**F(-1) timeout fail { 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }**

**F(-2) exception fail { }**



## Sympy

**A grade** { 1, 3, 4, 5, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 21, 27, 28, 29, 49, 50, 51, 52, 55, 56, 57, 58, 59, 60, 65, 66, 68, 69, 70, 71, 72, 73, 78, 79, 80, 81, 82, 91, 92, 93, 94, 103, 104, 105, 106, 107, 108, 109, 110, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 123, 124, 125, 126, 127, 138 }

**B grade** { 2, 6, 25, 26, 53, 54, 61, 62, 63, 64, 74, 75, 76, 77, 83, 84, 85, 87, 88, 89, 90, 95, 96, 97, 98, 99, 100 }

**C grade** { 86, 101, 102, 111, 122 }

**F normal fail** { 16, 18, 19, 20, 22, 23, 24, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 67, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143 }

**F(-1) timedout fail** { }

**F(-2) exception fail** { }

## 2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as  $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$ . To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	147	147	151	128	180	258	699	130	151
N.S.	1	1.00	1.03	0.87	1.22	1.76	4.76	0.88	1.03
time (sec)	N/A	0.037	1.219	3.678	0.212	0.276	0.801	0.280	9.864

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	121	121	117	69	130	69	269	140	100
N.S.	1	1.00	0.97	0.57	1.07	0.57	2.22	1.16	0.83
time (sec)	N/A	0.020	0.202	2.131	0.265	0.298	1.244	0.292	0.293

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	86	58	103	59	162	130	80
N.S.	1	1.00	0.91	0.61	1.08	0.62	1.71	1.37	0.84
time (sec)	N/A	0.014	0.156	2.121	0.275	0.298	0.749	0.277	9.359

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	83	47	76	49	92	120	60
N.S.	1	1.00	1.20	0.68	1.10	0.71	1.33	1.74	0.87
time (sec)	N/A	0.010	0.126	2.051	0.270	0.281	0.519	0.276	0.162

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	62	31	49	39	32	110	39
N.S.	1	1.00	1.44	0.72	1.14	0.91	0.74	2.56	0.91
time (sec)	N/A	0.007	0.084	2.129	0.270	0.420	0.286	0.269	0.085

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	102	63	117	68	211	57	81
N.S.	1	1.00	1.01	0.62	1.16	0.67	2.09	0.56	0.80
time (sec)	N/A	0.018	0.170	2.429	0.275	0.384	0.711	0.270	9.299

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	92	53	90	58	129	47	63
N.S.	1	1.00	1.16	0.67	1.14	0.73	1.63	0.59	0.80
time (sec)	N/A	0.013	0.126	2.051	0.271	0.354	0.527	0.281	9.276

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	82	43	63	48	68	37	45
N.S.	1	1.00	1.44	0.75	1.11	0.84	1.19	0.65	0.79
time (sec)	N/A	0.010	0.113	2.426	0.284	0.408	0.403	0.279	0.086

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	55	28	36	38	29	27	26
N.S.	1	1.00	1.57	0.80	1.03	1.09	0.83	0.77	0.74
time (sec)	N/A	0.006	0.076	1.989	0.281	0.795	0.240	0.263	9.078

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	47	27	36	35	26	25	26
N.S.	1	1.00	1.34	0.77	1.03	1.00	0.74	0.71	0.74
time (sec)	N/A	0.006	0.074	2.206	0.278	0.274	0.233	0.267	0.048

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	55	28	36	38	29	27	26
N.S.	1	1.00	1.57	0.80	1.03	1.09	0.83	0.77	0.74
time (sec)	N/A	0.006	0.082	2.267	0.270	0.268	0.238	0.270	9.101

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	67	39	55	43	56	35	39
N.S.	1	1.00	1.31	0.76	1.08	0.84	1.10	0.69	0.76
time (sec)	N/A	0.006	0.181	2.149	0.288	0.267	0.342	0.275	9.121

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	46	32	41	32	34	33	29
N.S.	1	1.00	1.31	0.91	1.17	0.91	0.97	0.94	0.83
time (sec)	N/A	0.005	0.070	1.895	0.201	0.265	0.213	0.261	0.110

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	46	32	43	32	34	33	29
N.S.	1	1.00	1.24	0.86	1.16	0.86	0.92	0.89	0.78
time (sec)	N/A	0.005	0.074	1.863	0.187	0.267	0.217	0.272	9.139

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	50	33	43	36	32	37	29
N.S.	1	1.00	1.28	0.85	1.10	0.92	0.82	0.95	0.74
time (sec)	N/A	0.005	0.108	1.864	0.199	0.273	0.198	0.266	0.080

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	83	70	75	111	105	0	74	96
N.S.	1	1.00	0.84	0.90	1.34	1.27	0.00	0.89	1.16
time (sec)	N/A	0.012	0.200	1.921	0.192	0.276	0.000	0.278	9.223

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	51	10	21	19	8	110	19
N.S.	1	1.00	3.19	0.62	1.31	1.19	0.50	6.88	1.19
time (sec)	N/A	0.004	0.050	1.861	0.271	0.271	0.294	0.273	9.247

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	24	19	28	39	0	64	20
N.S.	1	1.00	0.92	0.73	1.08	1.50	0.00	2.46	0.77
time (sec)	N/A	0.002	0.078	1.801	0.190	0.267	0.000	0.287	0.045

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	36	39	55	63	0	74	31
N.S.	1	1.00	0.68	0.74	1.04	1.19	0.00	1.40	0.58
time (sec)	N/A	0.005	0.110	1.854	0.187	0.274	0.000	0.306	9.112

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	48	50	82	83	0	84	40
N.S.	1	1.00	0.61	0.63	1.04	1.05	0.00	1.06	0.51
time (sec)	N/A	0.008	0.146	1.904	0.188	0.283	0.000	0.289	9.231

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	46	9	8	19	8	27	8
N.S.	1	1.00	3.83	0.75	0.67	1.58	0.67	2.25	0.67
time (sec)	N/A	0.004	0.051	1.899	0.291	0.276	0.256	0.271	8.972

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	21	19	28	29	0	29	18
N.S.	1	1.00	0.95	0.86	1.27	1.32	0.00	1.32	0.82
time (sec)	N/A	0.002	0.079	1.803	0.216	0.266	0.000	0.269	0.043

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	31	31	55	46	0	39	28
N.S.	1	1.00	0.69	0.69	1.22	1.02	0.00	0.87	0.62
time (sec)	N/A	0.004	0.106	1.824	0.189	0.256	0.000	0.276	9.005

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	41	41	82	61	0	49	73
N.S.	1	1.00	0.61	0.61	1.22	0.91	0.00	0.73	1.09
time (sec)	N/A	0.008	0.129	1.829	0.189	0.398	0.000	0.276	9.082

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	57	25	21	27	54	41	42
N.S.	1	1.00	4.75	2.08	1.75	2.25	4.50	3.42	3.50
time (sec)	N/A	0.005	0.070	2.184	0.291	0.378	0.258	0.281	9.098

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	58	37	29	27	51	59	36
N.S.	1	1.00	2.42	1.54	1.21	1.12	2.12	2.46	1.50
time (sec)	N/A	0.005	0.092	2.089	0.218	0.434	0.247	0.284	9.074

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	40	7	8	18	5	25	6
N.S.	1	1.00	4.00	0.70	0.80	1.80	0.50	2.50	0.60
time (sec)	N/A	0.005	0.049	2.281	0.301	0.775	0.231	0.290	9.067

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	39	9	17	17	17	33	11
N.S.	1	1.00	2.44	0.56	1.06	1.06	1.06	2.06	0.69
time (sec)	N/A	0.003	0.039	1.899	0.204	0.257	0.238	0.280	9.374











Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	51	44	43	43	49	43	43
N.S.	1	1.00	1.00	0.86	0.84	0.84	0.96	0.84	0.84
time (sec)	N/A	0.013	0.004	2.128	0.194	0.256	0.019	0.313	0.025

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	32	31	31	32	31	31
N.S.	1	1.00	1.00	0.91	0.89	0.89	0.91	0.89	0.89
time (sec)	N/A	0.008	0.003	2.136	0.195	0.267	0.023	0.314	0.039

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	21	21	22	21	21
N.S.	1	1.00	1.00	0.88	0.84	0.84	0.88	0.84	0.84
time (sec)	N/A	0.005	0.002	2.173	0.189	0.289	0.016	0.296	0.030

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	11	10	10	8	10	10
N.S.	1	1.00	1.00	0.92	0.83	0.83	0.67	0.83	0.83
time (sec)	N/A	0.001	0.000	0.024	0.193	0.266	0.017	0.262	0.017

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	16	15	67	53	15	16
N.S.	1	1.00	1.00	0.67	0.62	2.79	2.21	0.62	0.67
time (sec)	N/A	0.003	0.005	2.069	0.277	0.300	0.063	0.278	0.055

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	45	36	35	120	78	35	33
N.S.	1	1.00	1.00	0.80	0.78	2.67	1.73	0.78	0.73
time (sec)	N/A	0.007	0.032	2.140	0.278	0.554	0.101	0.309	9.204

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	55	57	58	188	105	45	55
N.S.	1	1.00	0.89	0.92	0.94	3.03	1.69	0.73	0.89
time (sec)	N/A	0.011	0.050	2.121	0.277	0.469	0.166	0.285	9.121

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	71	59	58	146	97	63	37
N.S.	1	1.00	0.85	0.70	0.69	1.74	1.15	0.75	0.44
time (sec)	N/A	0.015	0.134	2.149	0.221	0.456	2.653	0.270	9.090

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	60	48	43	124	70	49	37
N.S.	1	1.00	0.92	0.74	0.66	1.91	1.08	0.75	0.57
time (sec)	N/A	0.010	0.093	2.109	0.205	0.756	1.583	0.292	8.969

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	48	36	28	94	41	37	35
N.S.	1	1.00	1.04	0.78	0.61	2.04	0.89	0.80	0.76
time (sec)	N/A	0.007	0.054	2.205	0.185	0.281	0.959	0.274	0.126

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	21	13	59	17	37	20
N.S.	1	1.00	1.00	0.84	0.52	2.36	0.68	1.48	0.80
time (sec)	N/A	0.005	0.008	2.371	0.199	0.272	0.509	0.273	9.004

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	14	23	17	14	14
N.S.	1	1.00	1.00	0.94	0.88	1.44	1.06	0.88	0.88
time (sec)	N/A	0.002	0.046	2.438	0.191	0.270	0.343	0.278	0.034

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	29	26	31	47	95	27	28
N.S.	1	1.00	0.74	0.67	0.79	1.21	2.44	0.69	0.72
time (sec)	N/A	0.004	0.079	2.098	0.186	0.271	0.495	0.282	8.981

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	40	37	46	69	413	41	44
N.S.	1	1.00	0.69	0.64	0.79	1.19	7.12	0.71	0.76
time (sec)	N/A	0.007	0.097	1.978	0.185	0.290	0.804	0.298	9.041

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	51	48	61	91	1265	55	61
N.S.	1	1.00	0.66	0.62	0.79	1.18	16.43	0.71	0.79
time (sec)	N/A	0.012	0.115	2.123	0.192	0.282	1.230	0.280	9.085

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	20	17	30	19	80	45	19
N.S.	1	1.00	0.87	0.74	1.30	0.83	3.48	1.96	0.83
time (sec)	N/A	0.002	0.012	2.072	0.288	0.394	0.462	0.272	0.050

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	23	23	25	16	30	9	19	26	19
N.S.	1	1.00	1.09	0.70	1.30	0.39	0.83	1.13	0.83
time (sec)	N/A	0.002	0.006	2.067	0.270	0.451	0.294	0.280	9.027

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	26	9	6	8	22	25	14
N.S.	1	1.00	0.90	0.31	0.21	0.28	0.76	0.86	0.48
time (sec)	N/A	0.003	0.007	2.253	0.279	0.443	0.315	0.273	9.091

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	20	16	9	14	0	17	21
N.S.	1	1.00	0.80	0.64	0.36	0.56	0.00	0.68	0.84
time (sec)	N/A	0.002	0.006	2.147	0.280	0.667	0.000	0.268	9.005

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	A	A	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	23	23	25	16	30	9	19	26	13
N.S.	1	1.00	1.09	0.70	1.30	0.39	0.83	1.13	0.57
time (sec)	N/A	0.002	0.014	0.172	0.272	0.257	0.422	0.277	0.114

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	26	23	6	8	22	15	14
N.S.	1	1.00	0.90	0.79	0.21	0.28	0.76	0.52	0.48
time (sec)	N/A	0.003	0.011	1.909	0.286	0.254	0.414	0.281	9.148

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	27	27	30	9	19	26	18
N.S.	1	1.00	1.17	1.17	1.30	0.39	0.83	1.13	0.78
time (sec)	N/A	0.002	0.008	1.970	0.269	0.268	0.432	0.272	9.115

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	C	C	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	28	10	6	6	24	23	15
N.S.	1	1.00	0.97	0.34	0.21	0.21	0.83	0.79	0.52
time (sec)	N/A	0.004	0.007	2.091	0.273	0.278	0.417	0.271	9.080

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	27	27	30	9	20	26	18
N.S.	1	1.00	1.17	1.17	1.30	0.39	0.87	1.13	0.78
time (sec)	N/A	0.003	0.007	2.093	0.296	0.258	0.458	0.264	0.062

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	C	C	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	28	10	6	6	24	23	15
N.S.	1	1.00	0.97	0.34	0.21	0.21	0.83	0.79	0.52
time (sec)	N/A	0.005	0.004	2.223	0.286	0.262	0.414	0.265	9.119



Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	206	273	234	233	253	334	184
N.S.	1	1.00	1.89	2.50	2.15	2.14	2.32	3.06	1.69
time (sec)	N/A	0.109	0.033	2.276	0.192	0.263	0.086	0.269	9.160

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	206	273	234	235	250	334	184
N.S.	1	1.00	1.89	2.50	2.15	2.16	2.29	3.06	1.69
time (sec)	N/A	0.105	0.047	2.232	0.194	0.450	0.089	0.269	9.165

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	198	273	234	227	253	334	176
N.S.	1	1.00	1.82	2.50	2.15	2.08	2.32	3.06	1.61
time (sec)	N/A	0.100	0.038	2.256	0.201	0.433	0.078	0.263	9.202

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	109	109	206	273	234	235	248	334	184
N.S.	1	1.00	1.89	2.50	2.15	2.16	2.28	3.06	1.69
time (sec)	N/A	0.108	0.044	2.272	0.191	0.379	0.079	0.276	9.226

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	16	22	16	16
N.S.	1	1.00	1.00	0.94	0.89	0.89	1.22	0.89	0.89
time (sec)	N/A	0.008	0.006	4.031	0.282	0.739	0.049	0.269	0.031

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	9	8	10	7	8	8
N.S.	1	1.00	1.00	0.75	0.67	0.83	0.58	0.67	0.67
time (sec)	N/A	0.006	0.018	3.558	0.284	0.264	0.081	0.258	9.296

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	34	17	27	39	39	31	15
N.S.	1	1.00	1.79	0.89	1.42	2.05	2.05	1.63	0.79
time (sec)	N/A	0.012	0.026	2.151	0.276	0.259	0.053	0.256	0.236

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	13	13	10	15	8
N.S.	1	1.00	1.00	0.92	1.00	1.00	0.77	1.15	0.62
time (sec)	N/A	0.003	0.005	2.104	0.184	0.259	0.044	0.277	0.084

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	21	14	15	15	14	17	8
N.S.	1	1.00	1.00	0.67	0.71	0.71	0.67	0.81	0.38
time (sec)	N/A	0.004	0.004	2.180	0.180	0.273	0.054	0.253	0.096

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	B	B	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	17	17	14	13	13	12	15	6
N.S.	1	2.83	2.83	2.33	2.17	2.17	2.00	2.50	1.00
time (sec)	N/A	0.003	0.005	2.043	0.199	0.265	0.050	0.264	0.079

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	38	50	76	40	23
N.S.	1	1.00	1.00	0.89	1.41	1.85	2.81	1.48	0.85
time (sec)	N/A	0.013	0.012	1.982	0.196	0.266	0.127	0.263	9.566

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	29	26	39	51	76	45	23
N.S.	1	1.00	1.07	0.96	1.44	1.89	2.81	1.67	0.85
time (sec)	N/A	0.012	0.013	2.080	0.183	0.334	0.130	0.264	9.219

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	31	28	27	41	87	27	23
N.S.	1	1.00	1.00	0.90	0.87	1.32	2.81	0.87	0.74
time (sec)	N/A	0.015	0.012	3.004	0.193	0.448	0.091	0.261	9.187

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	29	26	39	51	76	45	23
N.S.	1	1.00	1.07	0.96	1.44	1.89	2.81	1.67	0.85
time (sec)	N/A	0.015	0.015	1.988	0.202	0.401	0.126	0.271	9.236

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	35	0	113	124	34	46
N.S.	1	1.00	1.00	0.92	0.00	2.97	3.26	0.89	1.21
time (sec)	N/A	0.023	0.014	2.115	0.000	0.421	0.103	0.270	8.994

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	34	35	0	124	100	30	33
N.S.	1	1.00	0.97	1.00	0.00	3.54	2.86	0.86	0.94
time (sec)	N/A	0.023	0.011	2.313	0.000	0.306	0.122	0.257	9.062

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	41	32	49	67	102	55	28
N.S.	1	1.00	1.28	1.00	1.53	2.09	3.19	1.72	0.88
time (sec)	N/A	0.019	0.012	2.106	0.286	0.260	0.131	0.263	9.066

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	43	34	36	45	39	36	33
N.S.	1	1.00	1.00	0.79	0.84	1.05	0.91	0.84	0.77
time (sec)	N/A	0.009	0.027	2.199	0.285	0.259	0.066	0.261	0.038

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	62	37	47	68	58	51	34
N.S.	1	1.00	1.44	0.86	1.09	1.58	1.35	1.19	0.79
time (sec)	N/A	0.010	0.029	2.053	0.292	0.288	0.074	0.266	9.009

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	33	32	34	53	31	36	34
N.S.	1	1.00	0.97	0.94	1.00	1.56	0.91	1.06	1.00
time (sec)	N/A	0.006	0.013	1.978	0.189	0.332	0.061	0.267	0.090

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	42	32	34	53	32	36	34
N.S.	1	1.00	1.00	0.76	0.81	1.26	0.76	0.86	0.81
time (sec)	N/A	0.006	0.017	2.084	0.186	0.246	0.071	0.259	0.081

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	70	68	0	334	265	67	119
N.S.	1	1.00	0.99	0.96	0.00	4.70	3.73	0.94	1.68
time (sec)	N/A	0.025	0.064	2.220	0.000	0.307	0.309	0.263	0.160

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	72	86	0	317	230	75	107
N.S.	1	1.00	1.00	1.19	0.00	4.40	3.19	1.04	1.49
time (sec)	N/A	0.028	0.053	2.242	0.000	0.288	0.311	0.259	9.153

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	78	83	97	171	218	90	100
N.S.	1	1.00	1.13	1.20	1.41	2.48	3.16	1.30	1.45
time (sec)	N/A	0.025	0.071	2.603	0.289	0.283	0.292	0.266	9.118

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	62	111	0	89	212	100	110
N.S.	1	1.00	1.00	1.79	0.00	1.44	3.42	1.61	1.77
time (sec)	N/A	0.116	0.134	3.144	0.000	0.454	0.486	0.283	0.153

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	58	120	31	55	63	56	56	38
N.S.	1	1.76	3.64	0.94	1.67	1.91	1.70	1.70	1.15
time (sec)	N/A	0.022	0.118	2.302	0.191	0.426	0.140	0.274	0.154

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	58	120	31	51	59	56	54	38
N.S.	1	1.87	3.87	1.00	1.65	1.90	1.81	1.74	1.23
time (sec)	N/A	0.019	0.099	2.200	0.192	0.517	0.146	0.269	0.127

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	33	27	15	165	27	27
N.S.	1	1.00	1.00	1.94	1.59	0.88	9.71	1.59	1.59
time (sec)	N/A	0.014	0.022	3.114	0.269	0.256	0.093	0.267	9.039

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	23	39	33	21	70	33	42
N.S.	1	1.00	1.00	1.70	1.43	0.91	3.04	1.43	1.83
time (sec)	N/A	0.020	0.035	3.141	0.296	0.251	0.359	0.263	9.164

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	48	29	38	40	32	40	39
N.S.	1	1.00	1.26	0.76	1.00	1.05	0.84	1.05	1.03
time (sec)	N/A	0.008	0.080	2.773	0.297	0.246	0.294	0.261	0.076

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	49	25	38	53	26	24	23
N.S.	1	1.00	1.63	0.83	1.27	1.77	0.87	0.80	0.77
time (sec)	N/A	0.007	0.100	2.278	0.287	0.262	0.307	0.265	0.049

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	49	40	52	40	44	41	39
N.S.	1	1.00	1.00	0.82	1.06	0.82	0.90	0.84	0.80
time (sec)	N/A	0.007	0.107	2.231	0.274	0.242	0.278	0.275	0.111

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	53	35	46	58	41	53	48
N.S.	1	1.00	1.18	0.78	1.02	1.29	0.91	1.18	1.07
time (sec)	N/A	0.010	0.138	2.385	0.277	0.255	0.285	0.260	9.009

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	61	35	46	60	42	36	35
N.S.	1	1.00	1.36	0.78	1.02	1.33	0.93	0.80	0.78
time (sec)	N/A	0.012	0.198	2.216	0.274	0.262	0.292	0.270	0.051

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	56	50	58	58	53	54	48
N.S.	1	1.00	0.90	0.81	0.94	0.94	0.85	0.87	0.77
time (sec)	N/A	0.009	0.201	2.260	0.280	0.266	0.300	0.267	0.095

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	43	43	56	32	41	60	37	31	30
N.S.	1	1.00	1.30	0.74	0.95	1.40	0.86	0.72	0.70
time (sec)	N/A	0.008	0.147	2.175	0.289	0.253	0.300	0.284	8.971

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	61	50	58	58	54	54	48
N.S.	1	1.00	1.03	0.85	0.98	0.98	0.92	0.92	0.81
time (sec)	N/A	0.009	0.283	2.220	0.283	0.255	0.307	0.270	9.013

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	C	C	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	54	46	46	86	56	0	36
N.S.	1	1.00	0.92	0.78	0.78	1.46	0.95	0.00	0.61
time (sec)	N/A	0.009	0.777	3.190	0.279	0.264	0.305	0.000	0.052

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	56	50	58	58	54	54	48
N.S.	1	1.00	0.90	0.81	0.94	0.94	0.87	0.87	0.77
time (sec)	N/A	0.009	0.213	2.305	0.274	0.259	0.287	0.262	0.218

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	56	32	41	60	32	31	30
N.S.	1	1.00	1.44	0.82	1.05	1.54	0.82	0.79	0.77
time (sec)	N/A	0.006	0.142	2.272	0.281	0.273	0.302	0.266	0.050



Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	24	9	8	20	10	40	20
N.S.	1	1.00	1.71	0.64	0.57	1.43	0.71	2.86	1.43
time (sec)	N/A	0.005	0.069	2.263	0.282	0.258	0.297	0.260	9.051

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	25	7	8	33	7	24	6
N.S.	1	1.00	2.50	0.70	0.80	3.30	0.70	2.40	0.60
time (sec)	N/A	0.005	0.083	2.421	0.278	0.255	0.310	0.270	8.963

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	21	22	20	22	41	20
N.S.	1	1.00	0.96	0.84	0.88	0.80	0.88	1.64	0.80
time (sec)	N/A	0.004	0.001	2.666	0.289	0.256	0.305	0.268	9.166

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	27	15	16	38	20	53	26
N.S.	1	1.00	1.50	0.83	0.89	2.11	1.11	2.94	1.44
time (sec)	N/A	0.007	0.091	2.411	0.296	0.262	0.292	0.275	9.164

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	39	15	16	40	20	36	16
N.S.	1	1.00	2.05	0.79	0.84	2.11	1.05	1.89	0.84
time (sec)	N/A	0.007	0.110	2.370	0.308	0.248	0.289	0.268	9.298

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	30	30	28	38	32	54	26
N.S.	1	1.00	0.86	0.86	0.80	1.09	0.91	1.54	0.74
time (sec)	N/A	0.005	0.108	2.192	0.277	0.251	0.292	0.261	9.324

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	30	12	11	40	15	31	11
N.S.	1	1.00	1.76	0.71	0.65	2.35	0.88	1.82	0.65
time (sec)	N/A	0.005	0.089	2.255	0.265	0.258	0.309	0.274	9.007

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	27	30	28	37	32	54	26
N.S.	1	1.00	0.84	0.94	0.88	1.16	1.00	1.69	0.81
time (sec)	N/A	0.005	0.085	2.285	0.307	0.251	0.298	0.282	9.108

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	C	C	<b>F</b>	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	28	26	16	67	37	0	17
N.S.	1	1.00	0.85	0.79	0.48	2.03	1.12	0.00	0.52
time (sec)	N/A	0.006	0.113	2.395	0.281	0.260	0.288	0.000	9.072

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	30	30	28	38	32	54	26
N.S.	1	1.00	0.86	0.86	0.80	1.09	0.91	1.54	0.74
time (sec)	N/A	0.005	0.101	2.123	0.276	0.261	0.298	0.281	9.191

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	30	12	11	40	12	31	11
N.S.	1	1.00	2.31	0.92	0.85	3.08	0.92	2.38	0.85
time (sec)	N/A	0.003	0.088	2.228	0.279	0.252	0.294	0.261	9.094

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	44	51	16	137	80	60	40
N.S.	1	1.00	2.00	2.32	0.73	6.23	3.64	2.73	1.82
time (sec)	N/A	0.010	0.213	2.292	0.189	0.273	0.467	0.333	9.364

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	23	23	123	44	19	141	83	63	46
N.S.	1	1.00	5.35	1.91	0.83	6.13	3.61	2.74	2.00
time (sec)	N/A	0.009	0.337	2.489	0.274	0.276	0.526	0.275	9.274

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	123	44	19	143	78	61	44
N.S.	1	1.00	6.15	2.20	0.95	7.15	3.90	3.05	2.20
time (sec)	N/A	0.008	0.320	2.495	0.281	0.266	0.508	0.281	9.261

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	<b>F</b>	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	19	18	26	38	0	17	15
N.S.	1	1.00	1.00	0.95	1.37	2.00	0.00	0.89	0.79
time (sec)	N/A	0.002	0.128	2.353	0.199	0.261	0.000	0.276	8.971



Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	32	32	32	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.009	0.018	0.000	0.000	0.000	0.000	0.000	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	37	37	37	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.010	0.022	0.000	0.000	0.000	0.000	0.000	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	21	21	21	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.007	0.017	0.000	0.000	0.000	0.000	0.000	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F</b>	<b>F(-1)</b>
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	54	54	48	0	0	0	0	0	0
N.S.	1	1.00	0.89	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.007	0.096	0.000	0.000	0.000	0.000	0.000	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	16	19	20	16	32
N.S.	1	1.00	1.00	0.94	0.89	1.06	1.11	0.89	1.78
time (sec)	N/A	0.001	0.004	2.112	0.187	0.253	0.029	0.267	9.255



## 2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio  $\frac{\text{number of rules}}{\text{integrand size}}$  is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [35] had the largest ratio of [.538499999999999979]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	3	1.00	13	0.231
2	A	6	3	1.00	15	0.200
3	A	5	3	1.00	15	0.200
4	A	4	3	1.00	15	0.200
5	A	3	3	1.00	15	0.200
6	A	6	3	1.00	13	0.231
7	A	5	3	1.00	13	0.231
8	A	4	3	1.00	13	0.231
9	A	3	3	1.00	13	0.231
10	A	3	3	1.00	13	0.231
11	A	3	3	1.00	13	0.231
12	A	4	3	1.00	11	0.273
13	A	3	3	1.00	11	0.273
14	A	3	3	1.00	11	0.273
15	A	3	3	1.00	11	0.273
16	A	3	2	1.00	13	0.154
17	A	2	2	1.00	15	0.133
18	A	1	1	1.00	15	0.067
19	A	2	2	1.00	15	0.133
20	A	3	2	1.00	15	0.133
21	A	2	2	1.00	13	0.154
22	A	1	1	1.00	13	0.077
23	A	2	2	1.00	13	0.154
24	A	3	2	1.00	13	0.154

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	2	2	1.00	16	0.125
26	A	2	2	1.00	15	0.133
27	A	2	2	1.00	13	0.154
28	A	2	2	1.00	11	0.182
29	A	2	2	1.00	11	0.182
30	A	6	5	1.00	13	0.385
31	A	5	5	1.00	13	0.385
32	A	4	4	1.00	13	0.308
33	A	5	5	1.00	13	0.385
34	A	6	5	1.00	13	0.385
35	A	8	7	1.00	13	0.538
36	A	7	7	1.00	13	0.538
37	A	6	6	1.00	13	0.462
38	A	7	7	1.00	13	0.538
39	A	8	7	1.00	13	0.538
40	A	5	4	1.00	13	0.308
41	A	4	4	1.00	13	0.308
42	A	4	4	1.00	13	0.308
43	A	3	3	1.00	13	0.231
44	A	3	3	1.00	13	0.231
45	A	4	4	1.00	13	0.308
46	A	5	4	1.00	13	0.308
47	A	6	4	1.00	13	0.308
48	A	1	1	1.00	11	0.091
49	A	2	1	1.00	9	0.111
50	A	2	1	1.00	9	0.111
51	A	2	1	1.00	9	0.111
52	A	1	0	1.00	7	0.000
53	A	1	1	1.00	9	0.111
54	A	2	2	1.00	9	0.222
55	A	3	2	1.00	9	0.222
56	A	5	3	1.00	11	0.273
57	A	4	3	1.00	11	0.273
58	A	3	3	1.00	11	0.273
59	A	2	2	1.00	11	0.182

Continued on next page



Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	1	1	1.00	11	0.091
61	A	2	2	1.00	11	0.182
62	A	3	2	1.00	11	0.182
63	A	4	2	1.00	11	0.182
64	A	1	1	1.00	14	0.071
65	A	1	1	1.00	14	0.071
66	A	2	2	1.00	14	0.143
67	A	1	1	1.00	14	0.071
68	A	1	1	1.00	14	0.071
69	A	2	2	1.00	14	0.143
70	A	1	1	1.00	14	0.071
71	A	2	2	1.00	14	0.143
72	A	1	1	1.00	14	0.071
73	A	2	2	1.00	14	0.143
74	A	3	2	1.00	23	0.087
75	A	3	2	1.00	23	0.087
76	A	3	2	1.00	23	0.087
77	A	3	2	1.00	23	0.087
78	A	2	2	1.00	12	0.167
79	A	2	2	1.00	15	0.133
80	A	2	2	1.00	12	0.167
81	A	3	2	1.00	12	0.167
82	A	3	2	1.00	12	0.167
83	B	3	2	2.83	10	0.200
84	A	2	2	1.00	12	0.167
85	A	2	2	1.00	12	0.167
86	A	2	2	1.00	12	0.167
87	A	2	2	1.00	12	0.167
88	A	2	2	1.00	12	0.167
89	A	2	2	1.00	13	0.154
90	A	2	2	1.00	14	0.143
91	A	3	3	1.00	12	0.250
92	A	3	3	1.00	12	0.250
93	A	4	3	1.00	12	0.250
94	A	4	3	1.00	12	0.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
95	A	3	3	1.00	12	0.250
96	A	3	3	1.00	13	0.231
97	A	3	3	1.00	14	0.214
98	A	2	2	1.00	40	0.050
99	A	3	2	1.76	30	0.067
100	A	3	2	1.87	31	0.065
101	A	2	2	1.00	14	0.143
102	A	2	2	1.00	16	0.125
103	A	3	3	1.00	14	0.214
104	A	3	3	1.00	14	0.214
105	A	3	3	1.00	14	0.214
106	A	3	3	1.00	14	0.214
107	A	3	3	1.00	14	0.214
108	A	3	3	1.00	14	0.214
109	A	3	3	1.00	14	0.214
110	A	3	3	1.00	14	0.214
111	A	3	3	1.00	14	0.214
112	A	3	3	1.00	14	0.214
113	A	3	3	1.00	14	0.214
114	A	2	2	1.00	14	0.143
115	A	2	2	1.00	14	0.143
116	A	2	2	1.00	14	0.143
117	A	2	2	1.00	14	0.143
118	A	2	2	1.00	14	0.143
119	A	2	2	1.00	14	0.143
120	A	2	2	1.00	14	0.143
121	A	2	2	1.00	14	0.143
122	A	2	2	1.00	14	0.143
123	A	2	2	1.00	14	0.143
124	A	2	2	1.00	14	0.143
125	A	2	2	1.00	27	0.074
126	A	2	2	1.00	30	0.067
127	A	2	2	1.00	28	0.071
128	A	1	1	1.00	12	0.083
129	A	1	1	1.00	14	0.071

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
130	A	1	1	1.00	16	0.062
131	A	2	2	1.00	14	0.143
132	A	1	1	1.00	12	0.083
133	A	2	2	1.00	12	0.167
134	A	2	2	1.00	12	0.167
135	A	2	2	1.00	12	0.167
136	A	2	2	1.00	12	0.167
137	A	1	1	1.00	10	0.100
138	A	1	1	1.00	7	0.143
139	A	2	2	1.00	12	0.167
140	A	2	2	1.00	12	0.167
141	A	2	2	1.00	12	0.167
142	A	2	2	1.00	12	0.167
143	A	2	2	1.00	12	0.167



---

---

# CHAPTER 3

---

## LISTING OF INTEGRALS

3.1	$\int (bx + cx^2)^{7/2} dx$	66
3.2	$\int (3ix + 4x^2)^{7/2} dx$	73
3.3	$\int (3ix + 4x^2)^{5/2} dx$	79
3.4	$\int (3ix + 4x^2)^{3/2} dx$	84
3.5	$\int \sqrt{3ix + 4x^2} dx$	89
3.6	$\int (3x - 4x^2)^{7/2} dx$	93
3.7	$\int (3x - 4x^2)^{5/2} dx$	99
3.8	$\int (3x - 4x^2)^{3/2} dx$	104
3.9	$\int \sqrt{3x - 4x^2} dx$	109
3.10	$\int \sqrt{6x - x^2} dx$	113
3.11	$\int \sqrt{5x - 9x^2} dx$	117
3.12	$\int (x - x^2)^{3/2} dx$	121
3.13	$\int \sqrt{4x + x^2} dx$	125
3.14	$\int \sqrt{-8x + x^2} dx$	129
3.15	$\int \sqrt{-x + x^2} dx$	133
3.16	$\int \frac{1}{(bx+cx^2)^{7/2}} dx$	137
3.17	$\int \frac{1}{\sqrt{3ix+4x^2}} dx$	141
3.18	$\int \frac{1}{(3ix+4x^2)^{3/2}} dx$	145
3.19	$\int \frac{1}{(3ix+4x^2)^{5/2}} dx$	149
3.20	$\int \frac{1}{(3ix+4x^2)^{7/2}} dx$	153
3.21	$\int \frac{1}{\sqrt{3x-4x^2}} dx$	157
3.22	$\int \frac{1}{(3x-4x^2)^{3/2}} dx$	161
3.23	$\int \frac{1}{(3x-4x^2)^{5/2}} dx$	164
3.24	$\int \frac{1}{(3x-4x^2)^{7/2}} dx$	168

3.25	$\int \frac{1}{\sqrt{bx-b^2x^2}} dx$	172
3.26	$\int \frac{1}{\sqrt{bx+b^2x^2}} dx$	176
3.27	$\int \frac{1}{\sqrt{6x-x^2}} dx$	180
3.28	$\int \frac{1}{\sqrt{4x+x^2}} dx$	184
3.29	$\int \frac{1}{\sqrt{-2x+x^2}} dx$	188
3.30	$\int (bx+cx^2)^{4/3} dx$	192
3.31	$\int \sqrt[3]{bx+cx^2} dx$	198
3.32	$\int \frac{1}{(bx+cx^2)^{2/3}} dx$	204
3.33	$\int \frac{1}{(bx+cx^2)^{5/3}} dx$	209
3.34	$\int \frac{1}{(bx+cx^2)^{8/3}} dx$	215
3.35	$\int (bx+cx^2)^{5/3} dx$	221
3.36	$\int (bx+cx^2)^{2/3} dx$	231
3.37	$\int \frac{1}{\sqrt[3]{bx+cx^2}} dx$	240
3.38	$\int \frac{1}{(bx+cx^2)^{4/3}} dx$	248
3.39	$\int \frac{1}{(bx+cx^2)^{7/3}} dx$	257
3.40	$\int (bx+cx^2)^{5/4} dx$	266
3.41	$\int (bx+cx^2)^{3/4} dx$	270
3.42	$\int \sqrt[4]{bx+cx^2} dx$	274
3.43	$\int \frac{1}{\sqrt[4]{bx+cx^2}} dx$	278
3.44	$\int \frac{1}{(bx+cx^2)^{3/4}} dx$	282
3.45	$\int \frac{1}{(bx+cx^2)^{5/4}} dx$	286
3.46	$\int \frac{1}{(bx+cx^2)^{9/4}} dx$	290
3.47	$\int \frac{1}{(bx+cx^2)^{13/4}} dx$	294
3.48	$\int (bx+cx^2)^p dx$	299
3.49	$\int (a+cx^2)^4 dx$	302
3.50	$\int (a+cx^2)^3 dx$	305
3.51	$\int (a+cx^2)^2 dx$	308
3.52	$\int (a+cx^2) dx$	311
3.53	$\int \frac{1}{a+cx^2} dx$	314
3.54	$\int \frac{1}{(a+cx^2)^2} dx$	318
3.55	$\int \frac{1}{(a+cx^2)^3} dx$	322
3.56	$\int (a+cx^2)^{5/2} dx$	326
3.57	$\int (a+cx^2)^{3/2} dx$	331
3.58	$\int \sqrt{a+cx^2} dx$	335
3.59	$\int \frac{1}{\sqrt{a+cx^2}} dx$	339
3.60	$\int \frac{1}{(a+cx^2)^{3/2}} dx$	343
3.61	$\int \frac{1}{(a+cx^2)^{5/2}} dx$	346

3.62	$\int \frac{1}{(a+cx^2)^{7/2}} dx$	350
3.63	$\int \frac{1}{(a+cx^2)^{9/2}} dx$	354
3.64	$\int (4+12x+9x^2)^{3/2} dx$	359
3.65	$\int \sqrt{4+12x+9x^2} dx$	363
3.66	$\int \frac{1}{\sqrt{4+12x+9x^2}} dx$	366
3.67	$\int \frac{1}{(4+12x+9x^2)^{3/2}} dx$	370
3.68	$\int \sqrt{4-12x+9x^2} dx$	373
3.69	$\int \frac{1}{\sqrt{4-12x+9x^2}} dx$	376
3.70	$\int \sqrt{-4+12x-9x^2} dx$	380
3.71	$\int \frac{1}{\sqrt{-4+12x-9x^2}} dx$	383
3.72	$\int \sqrt{-4-12x-9x^2} dx$	387
3.73	$\int \frac{1}{\sqrt{-4-12x-9x^2}} dx$	390
3.74	$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx$	394
3.75	$\int \left( \frac{-4+b^2}{4c} + bx + cx^2 \right)^5 dx$	400
3.76	$\int \left( \frac{-9+b^2}{4c} + bx + cx^2 \right)^5 dx$	406
3.77	$\int \left( \frac{-16+b^2}{4c} + bx + cx^2 \right)^5 dx$	412
3.78	$\int \frac{1}{2+4x+3x^2} dx$	418
3.79	$\int \frac{1}{4-2\sqrt{3x+x^2}} dx$	422
3.80	$\int \frac{1}{2+4x-3x^2} dx$	425
3.81	$\int \frac{1}{2+5x+3x^2} dx$	429
3.82	$\int \frac{1}{2+5x-3x^2} dx$	432
3.83	$\int \frac{1}{3+4x+x^2} dx$	435
3.84	$\int \frac{1}{1+\pi x+2x^2} dx$	439
3.85	$\int \frac{1}{1+\pi x-2x^2} dx$	443
3.86	$\int \frac{1}{1+\pi x+3x^2} dx$	447
3.87	$\int \frac{1}{1+\pi x-3x^2} dx$	451
3.88	$\int \frac{1}{a+cx+bx^2} dx$	455
3.89	$\int \frac{1}{b+2ax+bx^2} dx$	459
3.90	$\int \frac{1}{b+2ax-bx^2} dx$	463
3.91	$\int \frac{1}{(2+4x+3x^2)^2} dx$	467
3.92	$\int \frac{1}{(2+4x-3x^2)^2} dx$	471
3.93	$\int \frac{1}{(2+5x+3x^2)^2} dx$	475
3.94	$\int \frac{1}{(2+5x-3x^2)^2} dx$	479
3.95	$\int \frac{1}{(a+cx+bx^2)^2} dx$	483
3.96	$\int \frac{1}{(b+2ax+bx^2)^2} dx$	488
3.97	$\int \frac{1}{(b+2ax-bx^2)^2} dx$	493

3.98	$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx$	498
3.99	$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx$	503
3.100	$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx$	507
3.101	$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx$	511
3.102	$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{\pi}{7}\right)} dx$	515
3.103	$\int \sqrt{5 - 6x + 9x^2} dx$	519
3.104	$\int \sqrt{3 - 4x - 4x^2} dx$	523
3.105	$\int \sqrt{-8 + 6x + 9x^2} dx$	527
3.106	$\int \sqrt{2 + 4x + 3x^2} dx$	531
3.107	$\int \sqrt{2 + 4x - 3x^2} dx$	535
3.108	$\int \sqrt{2 + 5x + 3x^2} dx$	539
3.109	$\int \sqrt{2 + 5x - 3x^2} dx$	543
3.110	$\int \sqrt{-2 + 4x + 3x^2} dx$	547
3.111	$\int \sqrt{-2 + 4x - 3x^2} dx$	551
3.112	$\int \sqrt{-2 + 5x + 3x^2} dx$	555
3.113	$\int \sqrt{-2 + 5x - 3x^2} dx$	559
3.114	$\int \frac{1}{\sqrt{5 - 6x + 9x^2}} dx$	563
3.115	$\int \frac{1}{\sqrt{3 - 4x - 4x^2}} dx$	566
3.116	$\int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx$	570
3.117	$\int \frac{1}{\sqrt{2 + 4x + 3x^2}} dx$	573
3.118	$\int \frac{1}{\sqrt{2 + 4x - 3x^2}} dx$	577
3.119	$\int \frac{1}{\sqrt{2 + 5x + 3x^2}} dx$	581
3.120	$\int \frac{1}{\sqrt{2 + 5x - 3x^2}} dx$	585
3.121	$\int \frac{1}{\sqrt{-2 + 4x + 3x^2}} dx$	588
3.122	$\int \frac{1}{\sqrt{-2 + 4x - 3x^2}} dx$	592
3.123	$\int \frac{1}{\sqrt{-2 + 5x + 3x^2}} dx$	596
3.124	$\int \frac{1}{\sqrt{-2 + 5x - 3x^2}} dx$	600
3.125	$\int \frac{1}{\sqrt{\frac{b^2 + 4c}{4c} + bx + cx^2}} dx$	604
3.126	$\int \frac{1}{\sqrt{\frac{-b^2 + 4c}{4c} + bx - cx^2}} dx$	608
3.127	$\int \frac{1}{\sqrt{\frac{-b^2 + c}{4c} + bx - cx^2}} dx$	612
3.128	$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx$	616
3.129	$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx$	619
3.130	$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx$	622
3.131	$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx$	625
3.132	$\int (a + bx + cx^2)^p dx$	629
3.133	$\int (3 + 4x + 5x^2)^p dx$	633
3.134	$\int (3 + 4x + 4x^2)^p dx$	636



3.135	$\int (3 + 4x + 3x^2)^p dx$	639
3.136	$\int (3 + 4x + 2x^2)^p dx$	642
3.137	$\int (3 + 4x + x^2)^p dx$	645
3.138	$\int (3 + 4x)^p dx$	648
3.139	$\int (3 + 4x - x^2)^p dx$	652
3.140	$\int (3 + 4x - 2x^2)^p dx$	655
3.141	$\int (3 + 4x - 3x^2)^p dx$	658
3.142	$\int (3 + 4x - 4x^2)^p dx$	661
3.143	$\int (3 + 4x - 5x^2)^p dx$	664

### 3.1 $\int (bx + cx^2)^{7/2} dx$

Optimal result . . . . .	66
Rubi [A] (verified) . . . . .	66
Mathematica [A] (verified) . . . . .	68
Maple [A] (verified) . . . . .	68
Fricas [A] (verification not implemented) . . . . .	69
Sympy [A] (verification not implemented) . . . . .	70
Maxima [A] (verification not implemented) . . . . .	71
Giac [A] (verification not implemented) . . . . .	72
Mupad [B] (verification not implemented) . . . . .	72

#### Optimal result

Integrand size = 13, antiderivative size = 147

$$\int (bx + cx^2)^{7/2} dx = -\frac{35b^6(b+2cx)\sqrt{bx+cx^2}}{16384c^4} + \frac{35b^4(b+2cx)(bx+cx^2)^{3/2}}{6144c^3} - \frac{7b^2(b+2cx)(bx+cx^2)^{5/2}}{384c^2} + \frac{(b+2cx)(bx+cx^2)^{7/2}}{16c} + \frac{35b^8 \operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{bx+cx^2}}\right)}{16384c^{9/2}}$$

[Out]  $35/6144*b^4*(2*c*x+b)*(c*x^2+b*x)^(3/2)/c^3-7/384*b^2*(2*c*x+b)*(c*x^2+b*x)^(5/2)/c^2+1/16*(2*c*x+b)*(c*x^2+b*x)^(7/2)/c+35/16384*b^8*\operatorname{arctanh}(x*c^(1/2)/(c*x^2+b*x)^(1/2))/c^(9/2)-35/16384*b^6*(2*c*x+b)*(c*x^2+b*x)^(1/2)/c^4$

#### Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 634, 212}

$$\int (bx + cx^2)^{7/2} dx = \frac{35b^8 \operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{bx+cx^2}}\right)}{16384c^{9/2}} - \frac{35b^6(b+2cx)\sqrt{bx+cx^2}}{16384c^4} + \frac{35b^4(b+2cx)(bx+cx^2)^{3/2}}{6144c^3} - \frac{7b^2(b+2cx)(bx+cx^2)^{5/2}}{384c^2} + \frac{(b+2cx)(bx+cx^2)^{7/2}}{16c}$$

[In] Int[(b\*x + c\*x^2)^(7/2), x]

[Out]  $(-35*b^6*(b+2*c*x)*\operatorname{Sqrt}[b*x+c*x^2])/(16384*c^4) + (35*b^4*(b+2*c*x)*(b*x+c*x^2)^(3/2))/(6144*c^3) - (7*b^2*(b+2*c*x)*(b*x+c*x^2)^(5/2))/(384*c^2) + ((b+2*c*x)*(b*x+c*x^2)^(7/2))/(16*c) + (35*b^8*\operatorname{ArcTanh}[\operatorname{Sqrt}[c]*x/\operatorname{Sqrt}[b*x+c*x^2]])/(16384*c^(9/2))$

Rule 212

Int[((a\_) + (b\_)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 626

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

Rule 634

Int[1/Sqrt[(b\_)\*(x\_) + (c\_)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(1 - c\*x^2), x], x, x/Sqrt[b\*x + c\*x^2]], x] /; FreeQ[{b, c}, x]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} - \frac{(7b^2) \int (bx + cx^2)^{5/2} dx}{32c} \\
&= -\frac{7b^2(b + 2cx)(bx + cx^2)^{5/2}}{384c^2} + \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} + \frac{(35b^4) \int (bx + cx^2)^{3/2} dx}{768c^2} \\
&= \frac{35b^4(b + 2cx)(bx + cx^2)^{3/2}}{6144c^3} - \frac{7b^2(b + 2cx)(bx + cx^2)^{5/2}}{384c^2} \\
&\quad + \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} - \frac{(35b^6) \int \sqrt{bx + cx^2} dx}{4096c^3} \\
&= -\frac{35b^6(b + 2cx)\sqrt{bx + cx^2}}{16384c^4} + \frac{35b^4(b + 2cx)(bx + cx^2)^{3/2}}{6144c^3} \\
&\quad - \frac{7b^2(b + 2cx)(bx + cx^2)^{5/2}}{384c^2} + \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} + \frac{(35b^8) \int \frac{1}{\sqrt{bx + cx^2}} dx}{32768c^4} \\
&= -\frac{35b^6(b + 2cx)\sqrt{bx + cx^2}}{16384c^4} + \frac{35b^4(b + 2cx)(bx + cx^2)^{3/2}}{6144c^3} \\
&\quad - \frac{7b^2(b + 2cx)(bx + cx^2)^{5/2}}{384c^2} + \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} \\
&\quad + \frac{(35b^8) \text{Subst}\left(\int \frac{1}{1 - cx^2} dx, x, \frac{x}{\sqrt{bx + cx^2}}\right)}{16384c^4} \\
&= -\frac{35b^6(b + 2cx)\sqrt{bx + cx^2}}{16384c^4} + \frac{35b^4(b + 2cx)(bx + cx^2)^{3/2}}{6144c^3} \\
&\quad - \frac{7b^2(b + 2cx)(bx + cx^2)^{5/2}}{384c^2} + \frac{(b + 2cx)(bx + cx^2)^{7/2}}{16c} + \frac{35b^8 \tanh^{-1}\left(\frac{\sqrt{cx}}{\sqrt{bx + cx^2}}\right)}{16384c^{9/2}}
\end{aligned}$$

### Mathematica [A] (verified)

Time = 1.22 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.03

$$\int (bx + cx^2)^{7/2} dx = \frac{\sqrt{x(b+cx)} \left( \sqrt{c}(-105b^7 + 70b^6cx - 56b^5c^2x^2 + 48b^4c^3x^3 + 10880b^3c^4x^4 + 25856b^2c^5x^5 + 21504b^2c^6x^6 + 6144c^7x^7) + (210b^8 \operatorname{ArcTanh}[\frac{\sqrt{c}\sqrt{x}]{-\sqrt{b} + \sqrt{b+cx}}]}(\sqrt{x}\sqrt{b+cx})) \right)}{49152c^{9/2}}$$

[In] Integrate[(b\*x + c\*x^2)^(7/2),x]

[Out] (Sqrt[x\*(b + c\*x)]\*(Sqrt[c]\*(-105\*b^7 + 70\*b^6\*c\*x - 56\*b^5\*c^2\*x^2 + 48\*b^4\*c^3\*x^3 + 10880\*b^3\*c^4\*x^4 + 25856\*b^2\*c^5\*x^5 + 21504\*b\*c^6\*x^6 + 6144\*c^7\*x^7) + (210\*b^8\*ArcTanh[(Sqrt[c]\*Sqrt[x])/(-Sqrt[b] + Sqrt[b + c\*x])])/(Sqrt[x]\*Sqrt[b + c\*x]))/(49152\*c^(9/2))

### Maple [A] (verified)

Time = 3.68 (sec) , antiderivative size = 128, normalized size of antiderivative = 0.87

method	result
risch	$-\frac{(-6144c^7x^7 - 21504bc^6x^6 - 25856b^2c^5x^5 - 10880b^3c^4x^4 - 48b^4c^3x^3 + 56b^5c^2x^2 - 70b^6cx + 105b^7)x(cx+b)}{49152c^4\sqrt{x(cx+b)}} + \frac{35b^8 \ln\left(\frac{\frac{b+cx}{2} + \sqrt{cx^2+b}}{\sqrt{c}} + \sqrt{cx^2+b}\right)}{32768c^{\frac{9}{2}}}$
default	$\frac{(2cx+b)(cx^2+bx)^{\frac{7}{2}}}{16c} - \frac{7b^2}{24c} \left( \frac{(2cx+b)(cx^2+bx)^{\frac{5}{2}}}{12c} - \frac{5b^2}{16c} \left( \frac{(2cx+b)(cx^2+bx)^{\frac{3}{2}}}{8c} - \frac{b^2 \ln\left(\frac{\frac{b+cx}{2} + \sqrt{cx^2+b}}{\sqrt{c}} + \sqrt{cx^2+b}\right)}{8c^{\frac{3}{2}}}\right) \right)$

[In] int((c\*x^2+b\*x)^(7/2),x,method=\_RETURNVERBOSE)

[Out] -1/49152\*(-6144\*c^7\*x^7-21504\*b\*c^6\*x^6-25856\*b^2\*c^5\*x^5-10880\*b^3\*c^4\*x^4-48\*b^4\*c^3\*x^3+56\*b^5\*c^2\*x^2-70\*b^6\*c\*x+105\*b^7)\*x\*(c\*x+b)/c^4/(x\*(c\*x+b))^(1/2)+35/32768\*b^8/c^(9/2)\*ln((1/2\*b+c\*x)/c^(1/2)+(c\*x^2+b\*x)^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.76

$$\int (bx + cx^2)^{7/2} dx = \frac{105 b^8 \sqrt{c} \log(2cx + b + 2\sqrt{cx^2 + bx}\sqrt{c}) + 2(6144 c^8 x^7 + 21504 bc^7 x^6 + 25856 b^2 c^6 x^5 + 10880 b^3 c^5 x^4 + 48 b^4 c^4 x^3 - 56 b^5 c^3 x^2 + 70 b^6 c^2 x - 105 b^7 c) \sqrt{cx^2 + bx}}{98304 c^5} - \frac{105 b^8 \sqrt{-c} \arctan\left(\frac{\sqrt{cx^2 + bx}\sqrt{-c}}{cx}\right) - (6144 c^8 x^7 + 21504 bc^7 x^6 + 25856 b^2 c^6 x^5 + 10880 b^3 c^5 x^4 + 48 b^4 c^4 x^3 - 56 b^5 c^3 x^2 + 70 b^6 c^2 x - 105 b^7 c) \sqrt{cx^2 + bx}}{49152 c^5}$$

[In] integrate((c\*x^2+b\*x)^(7/2),x, algorithm="fricas")

```
[Out] [1/98304*(105*b^8*sqrt(c)*log(2*c*x + b + 2*sqrt(c*x^2 + b*x)*sqrt(c)) + 2*(6144*c^8*x^7 + 21504*b*c^7*x^6 + 25856*b^2*c^6*x^5 + 10880*b^3*c^5*x^4 + 48*b^4*c^4*x^3 - 56*b^5*c^3*x^2 + 70*b^6*c^2*x - 105*b^7*c)*sqrt(c*x^2 + b*x))/c^5, -1/49152*(105*b^8*sqrt(-c)*arctan(sqrt(c*x^2 + b*x)*sqrt(-c)/(c*x)) - (6144*c^8*x^7 + 21504*b*c^7*x^6 + 25856*b^2*c^6*x^5 + 10880*b^3*c^5*x^4 + 48*b^4*c^4*x^3 - 56*b^5*c^3*x^2 + 70*b^6*c^2*x - 105*b^7*c)*sqrt(c*x^2 + b*x))/c^5]
```

## Sympy [A] (verification not implemented)

Time = 0.80 (sec) , antiderivative size = 699, normalized size of antiderivative = 4.76

$$\int (bx + cx^2)^{7/2} dx = b^3 \left\{ \begin{array}{l} \left( \begin{array}{l} \frac{\log(b+2\sqrt{c}\sqrt{bx+cx^2+2cx})}{\sqrt{c}} \text{ for } \frac{b^2}{c} \neq 0 \\ \frac{(\frac{b}{2c}+x) \log(\frac{b}{2c}+x)}{\sqrt{c}(\frac{b}{2c}+x)^2} \text{ otherwise} \end{array} \right) \\ \frac{7b^5}{256c^4} \\ \frac{2(bx)^{\frac{9}{2}}}{9b^4} \\ 0 \end{array} \right. + \sqrt{bx+cx^2} \left( -\frac{7b^4}{128c^4} + \frac{7b^3x}{192c^3} - \frac{7b^2x^2}{240c^2} + \frac{bx^3}{40c} + \frac{x^4}{5} \right) \\
 + 3b^2c \left\{ \begin{array}{l} \left( \begin{array}{l} \frac{\log(b+2\sqrt{c}\sqrt{bx+cx^2+2cx})}{\sqrt{c}} \text{ for } \frac{b^2}{c} \neq 0 \\ \frac{(\frac{b}{2c}+x) \log(\frac{b}{2c}+x)}{\sqrt{c}(\frac{b}{2c}+x)^2} \text{ otherwise} \end{array} \right) \\ \frac{21b^6}{1024c^5} \\ \frac{2(bx)^{\frac{11}{2}}}{11b^5} \\ 0 \end{array} \right. + \sqrt{bx+cx^2} \cdot \left( \frac{21b^5}{512c^5} - \frac{7b^4x}{256c^4} + \frac{7b^3x^2}{320c^3} - \frac{3b^2x^3}{160c^2} + \frac{bx^4}{60c} + \frac{x^5}{6} \right) \\
 + 3bc^2 \left\{ \begin{array}{l} \left( \begin{array}{l} \frac{\log(b+2\sqrt{c}\sqrt{bx+cx^2+2cx})}{\sqrt{c}} \text{ for } \frac{b^2}{c} \neq 0 \\ \frac{(\frac{b}{2c}+x) \log(\frac{b}{2c}+x)}{\sqrt{c}(\frac{b}{2c}+x)^2} \text{ otherwise} \end{array} \right) \\ \frac{33b^7}{2048c^6} \\ \frac{2(bx)^{\frac{13}{2}}}{13b^6} \\ 0 \end{array} \right. + \sqrt{bx+cx^2} \left( -\frac{33b^6}{1024c^6} + \frac{11b^5x}{512c^5} - \frac{11b^4x^2}{640c^4} + \frac{33b^3x^3}{2240c^3} - \frac{11b^2x^4}{840c^2} + \frac{bx^5}{140c} \right) \\
 + c^3 \left\{ \begin{array}{l} \left( \begin{array}{l} \frac{\log(b+2\sqrt{c}\sqrt{bx+cx^2+2cx})}{\sqrt{c}} \text{ for } \frac{b^2}{c} \neq 0 \\ \frac{(\frac{b}{2c}+x) \log(\frac{b}{2c}+x)}{\sqrt{c}(\frac{b}{2c}+x)^2} \text{ otherwise} \end{array} \right) \\ \frac{429b^8}{32768c^7} \\ \frac{2(bx)^{\frac{15}{2}}}{15b^7} \\ 0 \end{array} \right. + \sqrt{bx+cx^2} \cdot \left( \frac{429b^7}{16384c^7} - \frac{143b^6x}{8192c^6} + \frac{143b^5x^2}{10240c^5} - \frac{429b^4x^3}{35840c^4} + \frac{143b^3x^4}{13440c^3} - \frac{143b^2x^5}{13440c^2} + \frac{143bx^6}{13440c} \right)
 \end{array}$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(7/2),x)

```
[Out] b**3*Piecewise((7*b**5*Piecewise((log(b + 2*sqrt(c))*sqrt(b*x + c*x**2) + 2*
c*x)/sqrt(c), Ne(b**2/c, 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*
c) + x)**2), True))/(256*c**4) + sqrt(b*x + c*x**2)*(-7*b**4/(128*c**4) + 7
*b**3*x/(192*c**3) - 7*b**2*x**2/(240*c**2) + b*x**3/(40*c) + x**4/5), Ne(c
, 0)), (2*(b*x)**(9/2)/(9*b**4), Ne(b, 0)), (0, True)) + 3*b**2*c*Piecewise
((-21*b**6*Piecewise((log(b + 2*sqrt(c))*sqrt(b*x + c*x**2) + 2*c*x)/sqrt(c)
, Ne(b**2/c, 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2),
True))/(1024*c**5) + sqrt(b*x + c*x**2)*(21*b**5/(512*c**5) - 7*b**4*x/(25
6*c**4) + 7*b**3*x**2/(320*c**3) - 3*b**2*x**3/(160*c**2) + b*x**4/(60*c) +
x**5/6), Ne(c, 0)), (2*(b*x)**(11/2)/(11*b**5), Ne(b, 0)), (0, True)) + 3*
b*c**2*Piecewise((33*b**7*Piecewise((log(b + 2*sqrt(c))*sqrt(b*x + c*x**2) +
2*c*x)/sqrt(c), Ne(b**2/c, 0)), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/
(2*c) + x)**2), True))/(2048*c**6) + sqrt(b*x + c*x**2)*(-33*b**6/(1024*c**
6) + 11*b**5*x/(512*c**5) - 11*b**4*x**2/(640*c**4) + 33*b**3*x**3/(2240*c
**3) - 11*b**2*x**4/(840*c**2) + b*x**5/(84*c) + x**6/7), Ne(c, 0)), (2*(b*x
)**(13/2)/(13*b**6), Ne(b, 0)), (0, True)) + c**3*Piecewise((-429*b**8*Piec
ewise((log(b + 2*sqrt(c))*sqrt(b*x + c*x**2) + 2*c*x)/sqrt(c), Ne(b**2/c, 0)
), ((b/(2*c) + x)*log(b/(2*c) + x)/sqrt(c*(b/(2*c) + x)**2), True))/(32768*
c**7) + sqrt(b*x + c*x**2)*(429*b**7/(16384*c**7) - 143*b**6*x/(8192*c**6)
+ 143*b**5*x**2/(10240*c**5) - 429*b**4*x**3/(35840*c**4) + 143*b**3*x**4/(
13440*c**3) - 13*b**2*x**5/(1344*c**2) + b*x**6/(112*c) + x**7/8), Ne(c, 0)
), (2*(b*x)**(15/2)/(15*b**7), Ne(b, 0)), (0, True))
```

## Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.22

$$\int (bx + cx^2)^{7/2} dx = \frac{1}{8} (cx^2 + bx)^{7/2} x - \frac{35 \sqrt{cx^2 + bx} b^6 x}{8192 c^3} + \frac{35 (cx^2 + bx)^{3/2} b^4 x}{3072 c^2} - \frac{7 (cx^2 + bx)^{5/2} b^2 x}{192 c} + \frac{35 b^8 \log(2cx + b + 2\sqrt{cx^2 + bx}\sqrt{c})}{32768 c^{9/2}} - \frac{35 \sqrt{cx^2 + bx} b^7}{16384 c^4} + \frac{35 (cx^2 + bx)^{3/2} b^5}{6144 c^3} - \frac{7 (cx^2 + bx)^{5/2} b^3}{384 c^2} + \frac{(cx^2 + bx)^{7/2} b}{16 c}$$

```
[In] integrate((c*x^2+b*x)^(7/2),x, algorithm="maxima")
```

```
[Out] 1/8*(c*x^2 + b*x)^(7/2)*x - 35/8192*sqrt(c*x^2 + b*x)*b^6*x/c^3 + 35/3072*(
c*x^2 + b*x)^(3/2)*b^4*x/c^2 - 7/192*(c*x^2 + b*x)^(5/2)*b^2*x/c + 35/32768
*b^8*log(2*c*x + b + 2*sqrt(c*x^2 + b*x)*sqrt(c))/c^(9/2) - 35/16384*sqrt(c
*x^2 + b*x)*b^7/c^4 + 35/6144*(c*x^2 + b*x)^(3/2)*b^5/c^3 - 7/384*(c*x^2 +
b*x)^(5/2)*b^3/c^2 + 1/16*(c*x^2 + b*x)^(7/2)*b/c
```

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.88

$$\int (bx + cx^2)^{7/2} dx = -\frac{35 b^8 \log(|2(\sqrt{cx} - \sqrt{cx^2 + bx})\sqrt{c} + b|)}{32768 c^{\frac{9}{2}}} - \frac{1}{49152} \left( \frac{105 b^7}{c^4} - 2 \left( \frac{35 b^6}{c^3} - 4 \left( \frac{7 b^5}{c^2} - 2 \left( \frac{3 b^4}{c} + 8 (85 b^3 + 2 (101 b^2 c + 12 (2 c^3 x + 7 b c^2) x) x) x \right) x \right) x \right) x \right)$$

`[In] integrate((c*x^2+b*x)^(7/2),x, algorithm="giac")`

```
[Out] -35/32768*b^8*log(abs(2*(sqrt(c)*x - sqrt(c*x^2 + b*x))*sqrt(c) + b))/c^(9/2) - 1/49152*(105*b^7/c^4 - 2*(35*b^6/c^3 - 4*(7*b^5/c^2 - 2*(3*b^4/c + 8*(85*b^3 + 2*(101*b^2*c + 12*(2*c^3*x + 7*b*c^2)*x)*x)*x)*x)*sqrt(c*x^2 + b*x)
```

**Mupad [B] (verification not implemented)**

Time = 9.86 (sec) , antiderivative size = 151, normalized size of antiderivative = 1.03

$$\int (bx + cx^2)^{7/2} dx = \frac{(cx^2 + bx)^{7/2} \left(\frac{b}{2} + cx\right)}{8c} - \frac{7b^2 \left( \frac{(cx^2 + bx)^{5/2} \left(\frac{b}{2} + cx\right)}{6c} - \frac{5b^2 \left( \frac{(cx^2 + bx)^{3/2} \left(\frac{b}{2} + cx\right)}{4c} - \frac{3b^2 \left( \sqrt{cx^2 + bx} \left(\frac{x}{2} + \frac{b}{4c}\right) - \frac{b^2 \ln\left(\frac{\frac{b}{2} + cx}{\sqrt{c}} + \sqrt{cx^2 + bx}\right)}{8c^{3/2}} \right)}{16c} \right)}{24c} \right)}{32c}$$

`[In] int((b*x + c*x^2)^(7/2),x)`

```
[Out] ((b*x + c*x^2)^(7/2)*(b/2 + c*x))/(8*c) - (7*b^2*(((b*x + c*x^2)^(5/2)*(b/2 + c*x))/(6*c) - (5*b^2*(((b*x + c*x^2)^(3/2)*(b/2 + c*x))/(4*c) - (3*b^2*((b*x + c*x^2)^(1/2)*(x/2 + b/(4*c)) - (b^2*log((b/2 + c*x)/c^(1/2) + (b*x + c*x^2)^(1/2)))/(8*c^(3/2)))))/(16*c)))/(24*c)))/(32*c)
```



## 3.2 $\int (3ix + 4x^2)^{7/2} dx$

Optimal result . . . . .	73
Rubi [A] (verified) . . . . .	73
Mathematica [A] (verified) . . . . .	75
Maple [A] (verified) . . . . .	75
Fricas [A] (verification not implemented) . . . . .	75
Sympy [B] (verification not implemented) . . . . .	76
Maxima [A] (verification not implemented) . . . . .	77
Giac [A] (verification not implemented) . . . . .	77
Mupad [B] (verification not implemented) . . . . .	78

### Optimal result

Integrand size = 15, antiderivative size = 121

$$\int (3ix + 4x^2)^{7/2} dx = \frac{25515(3i + 8x)\sqrt{3ix + 4x^2}}{4194304} + \frac{945(3i + 8x)(3ix + 4x^2)^{3/2}}{131072} \\ + \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{229635i \arcsin\left(1 - \frac{8ix}{3}\right)}{16777216}$$

[Out] 945/131072\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(3/2)+21/2048\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(5/2)+1/64\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(7/2)-229635/16777216\*I\*arcsin(-1+8/3\*I\*x)+25515/4194304\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(1/2)

### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {626, 633, 221}

$$\int (3ix + 4x^2)^{7/2} dx = \frac{229635i \arcsin\left(1 - \frac{8ix}{3}\right)}{16777216} + \frac{1}{64}(8x + 3i)(4x^2 + 3ix)^{7/2} \\ + \frac{21(8x + 3i)(4x^2 + 3ix)^{5/2}}{2048} + \frac{945(8x + 3i)(4x^2 + 3ix)^{3/2}}{131072} + \frac{25515(8x + 3i)\sqrt{4x^2 + 3ix}}{4194304}$$

[In] Int[((3\*I)\*x + 4\*x^2)^(7/2), x]

[Out] (25515\*(3\*I + 8\*x)\*Sqrt[(3\*I)\*x + 4\*x^2])/4194304 + (945\*(3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(3/2))/131072 + (21\*(3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(5/2))/2048 + ((3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(7/2))/64 + ((229635\*I)/16777216)\*ArcSin[1 - ((8\*I)/3)\*x]

Rule 221

Int[1/Sqrt[(a\_) + (b\_)\*(x\_)^2], x\_Symbol] := Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 626

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{63}{128} \int (3ix + 4x^2)^{5/2} dx \\
&= \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{945 \int (3ix + 4x^2)^{3/2} dx}{4096} \\
&= \frac{945(3i + 8x)(3ix + 4x^2)^{3/2}}{131072} + \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} \\
&\quad + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{25515 \int \sqrt{3ix + 4x^2} dx}{262144} \\
&= \frac{25515(3i + 8x)\sqrt{3ix + 4x^2}}{4194304} + \frac{945(3i + 8x)(3ix + 4x^2)^{3/2}}{131072} \\
&\quad + \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{229635 \int \frac{1}{\sqrt{3ix + 4x^2}} dx}{8388608} \\
&= \frac{25515(3i + 8x)\sqrt{3ix + 4x^2}}{4194304} + \frac{945(3i + 8x)(3ix + 4x^2)^{3/2}}{131072} \\
&\quad + \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} \\
&\quad + \frac{76545 \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{9}}} dx, x, 3i + 8x\right)}{16777216} \\
&= \frac{25515(3i + 8x)\sqrt{3ix + 4x^2}}{4194304} + \frac{945(3i + 8x)(3ix + 4x^2)^{3/2}}{131072} \\
&\quad + \frac{21(3i + 8x)(3ix + 4x^2)^{5/2}}{2048} + \frac{1}{64}(3i + 8x)(3ix + 4x^2)^{7/2} + \frac{229635i \sin^{-1}\left(1 - \frac{8ix}{3}\right)}{16777216}
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.97

$$\int (3ix + 4x^2)^{7/2} dx = \frac{\sqrt{x}\sqrt{3i+4x}(2\sqrt{x}\sqrt{3i+4x}(76545i - 68040x - 72576ix^2 + 82944x^3 - 25067520ix^4 - 79429632x^5 + 88080384ix^6 + 33554432x^7) - 229635\sqrt{x(3i+4x)})}{8388608}$$

`[In] Integrate[((3*I)*x + 4*x^2)^(7/2),x]`

```
[Out] (Sqrt[x]*Sqrt[3*I + 4*x]*(2*Sqrt[x]*Sqrt[3*I + 4*x]*(76545*I - 68040*x - (7
2576*I)*x^2 + 82944*x^3 - (25067520*I)*x^4 - 79429632*x^5 + (88080384*I)*x^
6 + 33554432*x^7) - 229635*Log[-2*Sqrt[x] + Sqrt[3*I + 4*x]])/(8388608*Sqr
t[x*(3*I + 4*x)])
```

**Maple [A] (verified)**

Time = 2.13 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.57

method	result
risch	$\frac{(33554432x^7 + 88080384ix^6 - 79429632x^5 - 25067520ix^4 + 82944x^3 - 72576ix^2 - 68040x + 76545i)(3i+4x)x}{4194304\sqrt{x(3i+4x)}} + \frac{229635 \operatorname{arcsinh}(i + \frac{8x}{3})}{16777216}$
default	$\frac{(3i+8x)(4x^2+3ix)^{\frac{7}{2}}}{64} + \frac{21(3i+8x)(4x^2+3ix)^{\frac{5}{2}}}{2048} + \frac{945(3i+8x)(4x^2+3ix)^{\frac{3}{2}}}{131072} + \frac{25515(3i+8x)\sqrt{4x^2+3ix}}{4194304} + \frac{229635 \operatorname{arcsinh}(i + \frac{8x}{3})}{16777216}$
trager	$(21ix^6 + 8x^7 - \frac{765}{128}ix^4 - \frac{303}{16}x^5 - \frac{567}{32768}ix^2 + \frac{81}{4096}x^3 + \frac{76545}{4194304}i - \frac{8505}{524288}x)\sqrt{4x^2+3ix} + \frac{229635 \ln(4\sqrt{4x^2+3ix} + i + \frac{8x}{3})}{16777216}$

`[In] int((3*I*x+4*x^2)^(7/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/4194304*(76545*I-68040*x-72576*I*x^2+82944*x^3-25067520*I*x^4-79429632*x^
5+88080384*I*x^6+33554432*x^7)*(3*I+4*x)*x/(x*(3*I+4*x))^(1/2)+229635/16777
216*arcsinh(I+8/3*x)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.57

$$\int (3ix + 4x^2)^{7/2} dx = \frac{1}{4194304} (33554432x^7 + 88080384ix^6 - 79429632x^5 - 25067520ix^4 + 82944x^3 - 72576ix^2 - 68040x + 76545i) \sqrt{4x^2+3ix} - \frac{229635}{16777216} \log\left(-2x + \sqrt{4x^2+3ix} - \frac{3}{4}i\right) - \frac{1165671}{268435456}$$

[In] integrate((3\*I\*x+4\*x^2)^(7/2),x, algorithm="fricas")

[Out] 1/4194304\*(33554432\*x^7 + 88080384\*I\*x^6 - 79429632\*x^5 - 25067520\*I\*x^4 + 82944\*x^3 - 72576\*I\*x^2 - 68040\*x + 76545\*I)\*sqrt(4\*x^2 + 3\*I\*x) - 229635/16777216\*log(-2\*x + sqrt(4\*x^2 + 3\*I\*x) - 3/4\*I) - 1165671/268435456

## Sympy [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 269 vs. 2(107) = 214.

Time = 1.24 (sec) , antiderivative size = 269, normalized size of antiderivative = 2.22

$$\int (3ix + 4x^2)^{7/2} dx =$$

$$\begin{aligned} & -108\sqrt{4x^2 + 3ix} \left( \frac{x^5}{6} + \frac{ix^4}{80} + \frac{27x^3}{2560} - \frac{189ix^2}{20480} - \frac{567x}{65536} + \frac{5103i}{524288} \right) \\ & + 64\sqrt{4x^2 + 3ix} \left( \frac{x^7}{8} + \frac{3ix^6}{448} + \frac{39x^5}{7168} - \frac{1287ix^4}{286720} \right. \\ & \quad \left. - \frac{34749x^3}{9175040} + \frac{34749ix^2}{10485760} + \frac{104247x}{33554432} - \frac{938223i}{268435456} \right) \\ & - 27i \left( \sqrt{4x^2 + 3ix} \left( \frac{x^4}{5} + \frac{3ix^3}{160} + \frac{21x^2}{1280} - \frac{63ix}{4096} - \frac{567}{32768} \right) + \frac{1701i \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{131072} \right) \\ & + 144i \left( \sqrt{4x^2 + 3ix} \left( \frac{x^6}{7} + \frac{ix^5}{112} + \frac{33x^4}{4480} - \frac{891ix^3}{143360} - \frac{891x^2}{163840} + \frac{2673ix}{524288} + \frac{24057}{4194304} \right) \right. \\ & \quad \left. - \frac{72171i \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{16777216} \right) - \frac{16041645 \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{16777216} \end{aligned}$$

[In] integrate((3\*I\*x+4\*x\*\*2)\*\*(7/2),x)

[Out] -108\*sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*5/6 + I\*x\*\*4/80 + 27\*x\*\*3/2560 - 189\*I\*x\*\*2/20480 - 567\*x/65536 + 5103\*I/524288) + 64\*sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*7/8 + 3\*I\*x\*\*6/448 + 39\*x\*\*5/7168 - 1287\*I\*x\*\*4/286720 - 34749\*x\*\*3/9175040 + 34749\*I\*x\*\*2/10485760 + 104247\*x/33554432 - 938223\*I/268435456) - 27\*I\*(sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*4/5 + 3\*I\*x\*\*3/160 + 21\*x\*\*2/1280 - 63\*I\*x/4096 - 567/32768) + 1701\*I\*asinh(8\*x/3 + I)/131072) + 144\*I\*(sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*6/7 + I\*x\*\*5/112 + 33\*x\*\*4/4480 - 891\*I\*x\*\*3/143360 - 891\*x\*\*2/163840 + 2673\*I\*x/524288 + 24057/4194304) - 72171\*I\*asinh(8\*x/3 + I)/16777216) - 16041645\*asinh(8\*x/3 + I)/16777216

**Maxima [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.07

$$\int (3ix + 4x^2)^{7/2} dx = \frac{1}{8} (4x^2 + 3ix)^{7/2} x + \frac{3}{64} i (4x^2 + 3ix)^{7/2} \\ + \frac{21}{256} (4x^2 + 3ix)^{5/2} x + \frac{63}{2048} i (4x^2 + 3ix)^{5/2} + \frac{945}{16384} (4x^2 + 3ix)^{3/2} x \\ + \frac{2835}{131072} i (4x^2 + 3ix)^{3/2} + \frac{25515}{524288} \sqrt{4x^2 + 3ix} \\ + \frac{76545}{4194304} i \sqrt{4x^2 + 3ix} + \frac{229635}{16777216} \log(8x + 4\sqrt{4x^2 + 3ix} + 3i)$$

[In] integrate((3\*I\*x+4\*x^2)^(7/2),x, algorithm="maxima")

```
[Out] 1/8*(4*x^2 + 3*I*x)^(7/2)*x + 3/64*I*(4*x^2 + 3*I*x)^(7/2) + 21/256*(4*x^2
+ 3*I*x)^(5/2)*x + 63/2048*I*(4*x^2 + 3*I*x)^(5/2) + 945/16384*(4*x^2 + 3*I
*x)^(3/2)*x + 2835/131072*I*(4*x^2 + 3*I*x)^(3/2) + 25515/524288*sqrt(4*x^2
+ 3*I*x)*x + 76545/4194304*I*sqrt(4*x^2 + 3*I*x) + 229635/16777216*log(8*x
+ 4*sqrt(4*x^2 + 3*I*x) + 3*I)
```

**Giac [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.16

$$\int (3ix + 4x^2)^{7/2} dx = \frac{1}{8388608} (8(16(8(32(8(16(8x + 21i)x - 303)x - 765i)x + 81)x - 567i)x - 8505)x + 7654 \\ - \frac{229635}{16777216} \log\left(2\sqrt{8x^2 + 2\sqrt{16x^2 + 9}}x\left(\frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1\right) - 8x - 3i\right)$$

[In] integrate((3\*I\*x+4\*x^2)^(7/2),x, algorithm="giac")

```
[Out] 1/8388608*(8*(16*(8*(32*(8*(16*(8*x + 21*I)*x - 303)*x - 765*I)*x + 81)*x -
567*I)*x - 8505)*x + 76545*I)*sqrt(8*x^2 + 2*sqrt(16*x^2 + 9)*x)*(3*I*x/(4
*x^2 + sqrt(16*x^4 + 9*x^2)) + 1) - 229635/16777216*log(2*sqrt(8*x^2 + 2*sq
rt(16*x^2 + 9)*x)*(3*I*x/(4*x^2 + sqrt(16*x^4 + 9*x^2)) + 1) - 8*x - 3*I)
```

**Mupad [B] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.83

$$\int (3ix + 4x^2)^{7/2} dx = \frac{229635 \ln\left(x + \frac{\sqrt{x(4x+3i)}}{2} + \frac{3i}{8}\right)}{16777216} + \frac{945(4x + \frac{3i}{2})(4x^2 + x3i)^{3/2}}{65536} + \frac{21(4x + \frac{3i}{2})(4x^2 + x3i)^{5/2}}{1024} + \frac{(4x + \frac{3i}{2})(4x^2 + x3i)^{7/2}}{32} + \frac{25515\left(\frac{x}{2} + \frac{3i}{16}\right)\sqrt{4x^2 + x3i}}{262144}$$

[In] int((x\*3i + 4\*x^2)^(7/2),x)

[Out] (229635\*log(x + (x\*(4\*x + 3i))^(1/2)/2 + 3i/8))/16777216 + (945\*(4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(3/2))/65536 + (21\*(4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(5/2))/1024 + ((4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(7/2))/32 + (25515\*(x/2 + 3i/16)\*(x\*3i + 4\*x^2)^(1/2))/262144

### 3.3 $\int (3ix + 4x^2)^{5/2} dx$

Optimal result . . . . .	79
Rubi [A] (verified) . . . . .	79
Mathematica [A] (verified) . . . . .	81
Maple [A] (verified) . . . . .	81
Fricas [A] (verification not implemented) . . . . .	81
Sympy [A] (verification not implemented) . . . . .	82
Maxima [A] (verification not implemented) . . . . .	82
Giac [B] (verification not implemented) . . . . .	83
Mupad [B] (verification not implemented) . . . . .	83

#### Optimal result

Integrand size = 15, antiderivative size = 95

$$\int (3ix + 4x^2)^{5/2} dx = \frac{405(3i + 8x)\sqrt{3ix + 4x^2}}{32768} + \frac{15(3i + 8x)(3ix + 4x^2)^{3/2}}{1024} + \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{3645i \arcsin\left(1 - \frac{8ix}{3}\right)}{131072}$$

[Out] 15/1024\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(3/2)+1/48\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(5/2)-3645/131072\*I\*arcsin(-1+8/3\*I\*x)+405/32768\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {626, 633, 221}

$$\int (3ix + 4x^2)^{5/2} dx = \frac{3645i \arcsin\left(1 - \frac{8ix}{3}\right)}{131072} + \frac{1}{48}(8x + 3i)(4x^2 + 3ix)^{5/2} + \frac{15(8x + 3i)(4x^2 + 3ix)^{3/2}}{1024} + \frac{405(8x + 3i)\sqrt{4x^2 + 3ix}}{32768}$$

[In] Int[((3\*I)\*x + 4\*x^2)^(5/2), x]

[Out] (405\*(3\*I + 8\*x)\*Sqrt[(3\*I)\*x + 4\*x^2])/32768 + (15\*(3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(3/2))/1024 + ((3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(5/2))/48 + ((3645\*I)/131072)\*ArcSin[1 - ((8\*I)/3)\*x]

Rule 221

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

### Rule 626

`Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c) / (2*c*(2*p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

### Rule 633

`Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{15}{32} \int (3ix + 4x^2)^{3/2} dx \\
 &= \frac{15(3i + 8x)(3ix + 4x^2)^{3/2}}{1024} + \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{405 \int \sqrt{3ix + 4x^2} dx}{2048} \\
 &= \frac{405(3i + 8x)\sqrt{3ix + 4x^2}}{32768} + \frac{15(3i + 8x)(3ix + 4x^2)^{3/2}}{1024} \\
 &\quad + \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{3645 \int \frac{1}{\sqrt{3ix + 4x^2}} dx}{65536} \\
 &= \frac{405(3i + 8x)\sqrt{3ix + 4x^2}}{32768} + \frac{15(3i + 8x)(3ix + 4x^2)^{3/2}}{1024} \\
 &\quad + \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{1215 \text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{9}}} dx, x, 3i + 8x\right)}{131072} \\
 &= \frac{405(3i + 8x)\sqrt{3ix + 4x^2}}{32768} + \frac{15(3i + 8x)(3ix + 4x^2)^{3/2}}{1024} \\
 &\quad + \frac{1}{48}(3i + 8x)(3ix + 4x^2)^{5/2} + \frac{3645i \sin^{-1}\left(1 - \frac{8ix}{3}\right)}{131072}
 \end{aligned}$$



**Mathematica [A] (verified)**

Time = 0.16 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.91

$$\int (3ix + 4x^2)^{5/2} dx = \frac{\sqrt{x(3i+4x)} \left( 7290i - 6480x - 6912ix^2 - 497664x^3 + 983040ix^4 + 524288x^5 - \frac{10935 \log(-2\sqrt{x} + \sqrt{x\sqrt{3i+4x}})}{\sqrt{x\sqrt{3i+4x}}} \right)}{196608}$$

`[In] Integrate[((3*I)*x + 4*x^2)^(5/2),x]`

```
[Out] (Sqrt[x*(3*I + 4*x)]*(7290*I - 6480*x - (6912*I)*x^2 - 497664*x^3 + (983040
*I)*x^4 + 524288*x^5 - (10935*Log[-2*Sqrt[x] + Sqrt[3*I + 4*x]])/(Sqrt[x]*S
qrt[3*I + 4*x]))/196608
```

**Maple [A] (verified)**

Time = 2.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.61

method	result
risch	$\frac{(262144x^5 + 491520ix^4 - 248832x^3 - 3456ix^2 - 3240x + 3645i)(3i+4x)x}{98304\sqrt{x(3i+4x)}} + \frac{3645 \operatorname{arcsinh}(i + \frac{8x}{3})}{131072}$
default	$\frac{(3i+8x)(4x^2+3ix)^{5/2}}{48} + \frac{15(3i+8x)(4x^2+3ix)^{3/2}}{1024} + \frac{405(3i+8x)\sqrt{4x^2+3ix}}{32768} + \frac{3645 \operatorname{arcsinh}(i + \frac{8x}{3})}{131072}$
trager	$(5ix^4 + \frac{8}{3}x^5 - \frac{9}{256}ix^2 - \frac{81}{32}x^3 + \frac{1215}{32768}i - \frac{135}{4096}x)\sqrt{4x^2+3ix} - \frac{3645 \ln(-440x-144-165i-192i\sqrt{4x^2+3ix}+384ix+192)}{131072}$

`[In] int((3*I*x+4*x^2)^(5/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/98304*(3645*I-3240*x-3456*I*x^2-248832*x^3+491520*I*x^4+262144*x^5)*(3*I+
4*x)*x/(x*(3*I+4*x))^(1/2)+3645/131072*arcsinh(I+8/3*x)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.62

$$\int (3ix + 4x^2)^{5/2} dx = \frac{1}{98304} (262144x^5 + 491520ix^4 - 248832x^3 - 3456ix^2 - 3240x + 3645i)\sqrt{4x^2 + 3ix} - \frac{3645}{131072} \log\left(-2x + \sqrt{4x^2 + 3ix} - \frac{3}{4}i\right) - \frac{8991}{1048576}$$

`[In] integrate((3*I*x+4*x^2)^(5/2),x, algorithm="fricas")`

[Out]  $1/98304*(262144*x^5 + 491520*I*x^4 - 248832*x^3 - 3456*I*x^2 - 3240*x + 3645*I)*\sqrt{4*x^2 + 3*I*x} - 3645/131072*\log(-2*x + \sqrt{4*x^2 + 3*I*x}) - 3/4*I) - 8991/1048576$

### Sympy [A] (verification not implemented)

Time = 0.75 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.71

$$\int (3ix + 4x^2)^{5/2} dx = -9\sqrt{4x^2 + 3ix} \left( \frac{x^3}{4} + \frac{ix^2}{32} + \frac{15x}{512} - \frac{135i}{4096} \right) + 16\sqrt{4x^2 + 3ix} \left( \frac{x^5}{6} + \frac{ix^4}{80} + \frac{27x^3}{2560} - \frac{189ix^2}{20480} - \frac{567x}{65536} + \frac{5103i}{524288} \right) + 24i \left( \sqrt{4x^2 + 3ix} \left( \frac{x^4}{5} + \frac{3ix^3}{160} + \frac{21x^2}{1280} - \frac{63ix}{4096} - \frac{567}{32768} \right) + \frac{1701i \operatorname{asinh} \left( \frac{8x}{3} + i \right)}{131072} \right) + \frac{44469 \operatorname{asinh} \left( \frac{8x}{3} + i \right)}{131072}$$

[In] `integrate((3*I*x+4*x**2)**(5/2),x)`

[Out]  $-9*\sqrt{4*x**2 + 3*I*x}*(x**3/4 + I*x**2/32 + 15*x/512 - 135*I/4096) + 16*\sqrt{4*x**2 + 3*I*x}*(x**5/6 + I*x**4/80 + 27*x**3/2560 - 189*I*x**2/20480 - 567*x/65536 + 5103*I/524288) + 24*I*(\sqrt{4*x**2 + 3*I*x}*(x**4/5 + 3*I*x**3/160 + 21*x**2/1280 - 63*I*x/4096 - 567/32768) + 1701*I*\operatorname{asinh}(8*x/3 + I)/131072) + 44469*\operatorname{asinh}(8*x/3 + I)/131072$

### Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.08

$$\int (3ix + 4x^2)^{5/2} dx = \frac{1}{6} (4x^2 + 3ix)^{5/2} x + \frac{1}{16} i (4x^2 + 3ix)^{5/2} + \frac{15}{128} (4x^2 + 3ix)^{3/2} x + \frac{45}{1024} i (4x^2 + 3ix)^{3/2} + \frac{405}{4096} \sqrt{4x^2 + 3ix} x + \frac{1215}{32768} i \sqrt{4x^2 + 3ix} + \frac{3645}{131072} \log \left( 8x + 4\sqrt{4x^2 + 3ix} + 3i \right)$$

[In] `integrate((3*I*x+4*x^2)^(5/2),x, algorithm="maxima")`

[Out]  $1/6*(4*x^2 + 3*I*x)^(5/2)*x + 1/16*I*(4*x^2 + 3*I*x)^(5/2) + 15/128*(4*x^2 + 3*I*x)^(3/2)*x + 45/1024*I*(4*x^2 + 3*I*x)^(3/2) + 405/4096*\sqrt{4*x^2 + 3*I*x}*x + 1215/32768*I*\sqrt{4*x^2 + 3*I*x} + 3645/131072*\log(8*x + 4*\sqrt{4*x^2 + 3*I*x} + 3*I)$

**Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 130 vs.  $2(63) = 126$ .

Time = 0.28 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.37

$$\int (3ix + 4x^2)^{5/2} dx = \frac{1}{196608} (8(16(8(32(8x + 15i)x - 243)x - 27i)x - 405)x + 3645i)\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( -\frac{3645}{131072} \log \left( 2\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - 8x - 3i \right) \right)$$

[In] integrate((3\*I\*x+4\*x^2)^(5/2),x, algorithm="giac")

[Out] 1/196608\*(8\*(16\*(8\*(32\*(8\*x + 15\*I)\*x - 243)\*x - 27\*I)\*x - 405)\*x + 3645\*I)\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 3645/131072\*log(2\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 8\*x - 3\*I)

**Mupad [B] (verification not implemented)**

Time = 9.36 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.84

$$\int (3ix + 4x^2)^{5/2} dx = \frac{3645 \ln \left( x + \frac{\sqrt{x(4x+3i)}}{2} + \frac{3i}{8} \right)}{131072} + \frac{15(4x + \frac{3i}{2})(4x^2 + x3i)^{3/2}}{512} + \frac{(4x + \frac{3i}{2})(4x^2 + x3i)^{5/2}}{24} + \frac{405(\frac{x}{2} + \frac{3i}{16})\sqrt{4x^2 + x3i}}{2048}$$

[In] int((x\*3i + 4\*x^2)^(5/2),x)

[Out] (3645\*log(x + (x\*(4\*x + 3i))^(1/2)/2 + 3i/8))/131072 + (15\*(4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(3/2))/512 + ((4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(5/2))/24 + (405\*(x/2 + 3i/16)\*(x\*3i + 4\*x^2)^(1/2))/2048

### 3.4 $\int (3ix + 4x^2)^{3/2} dx$

Optimal result	84
Rubi [A] (verified)	84
Mathematica [A] (verified)	85
Maple [A] (verified)	86
Fricas [A] (verification not implemented)	86
Sympy [A] (verification not implemented)	86
Maxima [A] (verification not implemented)	87
Giac [B] (verification not implemented)	87
Mupad [B] (verification not implemented)	88

#### Optimal result

Integrand size = 15, antiderivative size = 69

$$\int (3ix + 4x^2)^{3/2} dx = \frac{27(3i + 8x)\sqrt{3ix + 4x^2}}{1024} + \frac{1}{32}(3i + 8x)(3ix + 4x^2)^{3/2} + \frac{243i \arcsin\left(1 - \frac{8ix}{3}\right)}{4096}$$

[Out] 1/32\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(3/2)-243/4096\*I\*arcsin(-1+8/3\*I\*x)+27/1024\*(3\*I+8\*x)\*(3\*I\*x+4\*x^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {626, 633, 221}

$$\int (3ix + 4x^2)^{3/2} dx = \frac{243i \arcsin\left(1 - \frac{8ix}{3}\right)}{4096} + \frac{1}{32}(8x + 3i)(4x^2 + 3ix)^{3/2} + \frac{27(8x + 3i)\sqrt{4x^2 + 3ix}}{1024}$$

[In] Int[((3\*I)\*x + 4\*x^2)^(3/2), x]

[Out] (27\*(3\*I + 8\*x)\*Sqrt[(3\*I)\*x + 4\*x^2])/1024 + ((3\*I + 8\*x)\*((3\*I)\*x + 4\*x^2)^(3/2))/32 + ((243\*I)/4096)\*ArcSin[1 - ((8\*I)/3)\*x]

Rule 221

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N
eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{32}(3i + 8x)(3ix + 4x^2)^{3/2} + \frac{27}{64} \int \sqrt{3ix + 4x^2} dx \\
&= \frac{27(3i + 8x)\sqrt{3ix + 4x^2}}{1024} + \frac{1}{32}(3i + 8x)(3ix + 4x^2)^{3/2} + \frac{243 \int \frac{1}{\sqrt{3ix+4x^2}} dx}{2048} \\
&= \frac{27(3i + 8x)\sqrt{3ix + 4x^2}}{1024} + \frac{1}{32}(3i + 8x)(3ix + 4x^2)^{3/2} + \frac{81 \text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{x^2}{9}}} dx, x, 3i + 8x\right)}{4096} \\
&= \frac{27(3i + 8x)\sqrt{3ix + 4x^2}}{1024} + \frac{1}{32}(3i + 8x)(3ix + 4x^2)^{3/2} + \frac{243i \sin^{-1}\left(1 - \frac{8ix}{3}\right)}{4096}
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.20

$$\int (3ix + 4x^2)^{3/2} dx = \frac{2x(-243 + 108ix - 3744x^2 + 7680ix^3 + 4096x^4) - 243\sqrt{x}\sqrt{3i + 4x} \log(-2\sqrt{x} + \sqrt{3i + 4x})}{2048\sqrt{x}(3i + 4x)}$$

```
[In] Integrate[((3*I)*x + 4*x^2)^(3/2), x]
```

```
[Out] (2*x*(-243 + (108*I)*x - 3744*x^2 + (7680*I)*x^3 + 4096*x^4) - 243*Sqrt[x]*
Sqrt[3*I + 4*x]*Log[-2*Sqrt[x] + Sqrt[3*I + 4*x]])/(2048*Sqrt[x*(3*I + 4*x)]
)
```

**Maple [A] (verified)**

Time = 2.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.68

method	result
risch	$\frac{(1024x^3+1152ix^2-72x+81i)(3i+4x)x}{1024\sqrt{x(3i+4x)}} + \frac{243 \operatorname{arcsinh}\left(i+\frac{8x}{3}\right)}{4096}$
default	$\frac{(3i+8x)(4x^2+3ix)^{\frac{3}{2}}}{32} + \frac{27(3i+8x)\sqrt{4x^2+3ix}}{1024} + \frac{243 \operatorname{arcsinh}\left(i+\frac{8x}{3}\right)}{4096}$
trager	$\left(\frac{9}{8}ix^2 + x^3 + \frac{81}{1024}i - \frac{9}{128}x\right)\sqrt{4x^2+3ix} - \frac{243 \ln\left(-440x-144-165i-192i\sqrt{4x^2+3ix}+384ix+220\sqrt{4x^2+3ix}\right)}{4096}$
pseudoelliptic	$\frac{729\left(-\frac{27 \ln\left(\frac{-2x+\sqrt{x(3i+4x)}}{x}\right)}{512} + \frac{27 \ln\left(\frac{\sqrt{x(3i+4x)+2x}}{x}\right)}{512} + \left(ix^2 + \frac{8}{9}x^3 + \frac{9}{128}i - \frac{1}{16}x\right)\sqrt{x(3i+4x)}\right)x^4}{8\left(\sqrt{x(3i+4x)}+2x\right)^4\left(2x-\sqrt{x(3i+4x)}\right)^4}$

[In] int((3\*I\*x+4\*x^2)^(3/2),x,method=\_RETURNVERBOSE)

[Out] 1/1024\*(81\*I-72\*x+1152\*I\*x^2+1024\*x^3)\*(3\*I+4\*x)\*x/(x\*(3\*I+4\*x))^(1/2)+243/4096\*arcsinh(I+8/3\*x)

**Fricas [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.71

$$\int (3ix + 4x^2)^{3/2} dx = \frac{1}{1024} (1024x^3 + 1152ix^2 - 72x + 81i)\sqrt{4x^2 + 3ix} - \frac{243}{4096} \log\left(-2x + \sqrt{4x^2 + 3ix} - \frac{3}{4}i\right) - \frac{567}{32768}$$

[In] integrate((3\*I\*x+4\*x^2)^(3/2),x, algorithm="fricas")

[Out] 1/1024\*(1024\*x^3 + 1152\*I\*x^2 - 72\*x + 81\*I)\*sqrt(4\*x^2 + 3\*I\*x) - 243/4096\*log(-2\*x + sqrt(4\*x^2 + 3\*I\*x) - 3/4\*I) - 567/32768

**Sympy [A] (verification not implemented)**

Time = 0.52 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.33

$$\int (3ix + 4x^2)^{3/2} dx = 4\sqrt{4x^2 + 3ix}\left(\frac{x^3}{4} + \frac{ix^2}{32} + \frac{15x}{512} - \frac{135i}{4096}\right) + 3i\left(\sqrt{4x^2 + 3ix}\left(\frac{x^2}{3} + \frac{ix}{16} + \frac{9}{128}\right) - \frac{27i \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{512}\right) - \frac{405 \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{4096}$$

[In] integrate((3\*I\*x+4\*x\*\*2)\*\*(3/2),x)

[Out] 4\*sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*3/4 + I\*x\*\*2/32 + 15\*x/512 - 135\*I/4096) + 3\*I\*(sqrt(4\*x\*\*2 + 3\*I\*x)\*(x\*\*2/3 + I\*x/16 + 9/128) - 27\*I\*asinh(8\*x/3 + I)/512) - 405\*asinh(8\*x/3 + I)/4096

## Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.10

$$\int (3ix + 4x^2)^{3/2} dx = \frac{1}{4} (4x^2 + 3ix)^{\frac{3}{2}} x + \frac{3}{32} i (4x^2 + 3ix)^{\frac{3}{2}} + \frac{27}{128} \sqrt{4x^2 + 3ix} x + \frac{81}{1024} i \sqrt{4x^2 + 3ix} + \frac{243}{4096} \log(8x + 4\sqrt{4x^2 + 3ix} + 3i)$$

[In] integrate((3\*I\*x+4\*x^2)^(3/2),x, algorithm="maxima")

[Out] 1/4\*(4\*x^2 + 3\*I\*x)^(3/2)\*x + 3/32\*I\*(4\*x^2 + 3\*I\*x)^(3/2) + 27/128\*sqrt(4\*x^2 + 3\*I\*x)\*x + 81/1024\*I\*sqrt(4\*x^2 + 3\*I\*x) + 243/4096\*log(8\*x + 4\*sqrt(4\*x^2 + 3\*I\*x) + 3\*I)

## Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(45) = 90.

Time = 0.28 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.74

$$\int (3ix + 4x^2)^{3/2} dx = \frac{1}{2048} (8(16(8x + 9i)x - 9)x + 81i) \sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - \frac{243}{4096} \log \left( 2\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - 8x - 3i \right)$$

[In] integrate((3\*I\*x+4\*x^2)^(3/2),x, algorithm="giac")

[Out] 1/2048\*(8\*(16\*(8\*x + 9\*I)\*x - 9)\*x + 81\*I)\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 243/4096\*log(2\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 8\*x - 3\*I)

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.87

$$\int (3ix + 4x^2)^{3/2} dx = \frac{243 \ln \left( x + \frac{\sqrt{x(4x+3i)}}{2} + \frac{3i}{8} \right)}{4096} + \frac{(4x + \frac{3i}{2})(4x^2 + x3i)^{3/2}}{16} + \frac{27 \left( \frac{x}{2} + \frac{3i}{16} \right) \sqrt{4x^2 + x3i}}{64}$$

[In] int((x\*3i + 4\*x^2)^(3/2),x)

[Out] (243\*log(x + (x\*(4\*x + 3i))^(1/2)/2 + 3i/8))/4096 + ((4\*x + 3i/2)\*(x\*3i + 4\*x^2)^(3/2))/16 + (27\*(x/2 + 3i/16)\*(x\*3i + 4\*x^2)^(1/2))/64



### 3.5 $\int \sqrt{3ix + 4x^2} dx$

Optimal result	89
Rubi [A] (verified)	89
Mathematica [A] (verified)	90
Maple [A] (verified)	90
Fricas [A] (verification not implemented)	91
Sympy [A] (verification not implemented)	91
Maxima [A] (verification not implemented)	92
Giac [B] (verification not implemented)	92
Mupad [B] (verification not implemented)	92

#### Optimal result

Integrand size = 15, antiderivative size = 43

$$\int \sqrt{3ix + 4x^2} dx = \frac{1}{16}(3i + 8x)\sqrt{3ix + 4x^2} + \frac{9}{64}i \arcsin\left(1 - \frac{8ix}{3}\right)$$

[Out]  $-9/64*I*\arcsin(-1+8/3*I*x)+1/16*(3*I+8*x)*(3*I*x+4*x^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {626, 633, 221}

$$\int \sqrt{3ix + 4x^2} dx = \frac{9}{64}i \arcsin\left(1 - \frac{8ix}{3}\right) + \frac{1}{16}\sqrt{4x^2 + 3ix}(8x + 3i)$$

[In]  $\text{Int}[\text{Sqrt}[(3*I)*x + 4*x^2], x]$

[Out]  $((3*I + 8*x)*\text{Sqrt}[(3*I)*x + 4*x^2])/16 + ((9*I)/64)*\text{ArcSin}[1 - ((8*I)/3)*x]$

#### Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

#### Rule 626

$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - \text{Dist}[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), \text{Int}[(a + b*x + c*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ N$

$eQ[b^2 - 4ac, 0] \ \&\& \ GtQ[p, 0] \ \&\& \ IntegerQ[4p]$

### Rule 633

$Int[(a_.) + (b_.)(x_) + (c_.)(x_)^2]^{(p)}, x\_Symbol] \ :> \ Dist[1/(2c*(-4*(c/(b^2 - 4ac))))^{(p)}, Subst[Int[Simp[1 - x^2/(b^2 - 4ac)], x]^{(p)}, x], x, b + 2cx], x] \ /; \ FreeQ[\{a, b, c, p\}, x] \ \&\& \ GtQ[4a - b^2/c, 0]$

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{16}(3i + 8x)\sqrt{3ix + 4x^2} + \frac{9}{32} \int \frac{1}{\sqrt{3ix + 4x^2}} dx \\ &= \frac{1}{16}(3i + 8x)\sqrt{3ix + 4x^2} + \frac{3}{64} \text{Subst} \left( \int \frac{1}{\sqrt{1 + \frac{x^2}{9}}} dx, x, 3i + 8x \right) \\ &= \frac{1}{16}(3i + 8x)\sqrt{3ix + 4x^2} + \frac{9}{64} i \sin^{-1} \left( 1 - \frac{8ix}{3} \right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\int \sqrt{3ix + 4x^2} dx = \frac{1}{32} \sqrt{x(3i + 4x)} \left( 6i + 16x - \frac{9 \log(-2\sqrt{x} + \sqrt{3i + 4x})}{\sqrt{x}\sqrt{3i + 4x}} \right)$$

[In] Integrate[Sqrt[(3\*I)\*x + 4\*x^2],x]

[Out] (Sqrt[x\*(3\*I + 4\*x)]\*(6\*I + 16\*x - (9\*Log[-2\*Sqrt[x] + Sqrt[3\*I + 4\*x]])/(Sqrt[x]\*Sqrt[3\*I + 4\*x]))) / 32

### Maple [A] (verified)

Time = 2.13 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{(3i+8x)\sqrt{4x^2+3ix}}{16} + \frac{9 \operatorname{arcsinh}(i+\frac{8x}{3})}{64}$	31
risch	$\frac{(3i+8x)(3i+4x)x}{16\sqrt{x(3i+4x)}} + \frac{9 \operatorname{arcsinh}(i+\frac{8x}{3})}{64}$	36
trager	$\left(\frac{3i}{16} + \frac{x}{2}\right) \sqrt{4x^2 + 3ix} - \frac{9 \ln(-440x - 144 - 165i - 192i\sqrt{4x^2+3ix} + 384ix + 220\sqrt{4x^2+3ix})}{64}$	64
pseudoelliptic	$-\frac{27 \left( -\frac{3 \ln\left(\frac{-2x+\sqrt{x(3i+4x)}}{x}\right)}{4} + \frac{3 \ln\left(\frac{\sqrt{x(3i+4x)+2x}}{x}\right)}{4} + \left(i+\frac{8x}{3}\right)\sqrt{x(3i+4x)} \right) x^2}{16(\sqrt{x(3i+4x)+2x})^2(2x-\sqrt{x(3i+4x)})^2}$	100

[In] `int((3*I*x+4*x^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/16*(3*I+8*x)*(3*I*x+4*x^2)^(1/2)+9/64*arcsinh(I+8/3*x)`

### Fricas [A] (verification not implemented)

none

Time = 0.42 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

$$\int \sqrt{3ix + 4x^2} dx = \frac{1}{16} \sqrt{4x^2 + 3ix} (8x + 3i) - \frac{9}{64} \log\left(-2x + \sqrt{4x^2 + 3ix} - \frac{3}{4}i\right) - \frac{9}{256}$$

[In] `integrate((3*I*x+4*x^2)^(1/2),x, algorithm="fricas")`

[Out] `1/16*sqrt(4*x^2 + 3*I*x)*(8*x + 3*I) - 9/64*log(-2*x + sqrt(4*x^2 + 3*I*x) - 3/4*I) - 9/256`

### Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

$$\int \sqrt{3ix + 4x^2} dx = \left(\frac{x}{2} + \frac{3i}{16}\right) \sqrt{4x^2 + 3ix} + \frac{9 \operatorname{asinh}\left(\frac{8x}{3} + i\right)}{64}$$

[In] `integrate((3*I*x+4*x**2)**(1/2),x)`

[Out] `(x/2 + 3*I/16)*sqrt(4*x**2 + 3*I*x) + 9*asinh(8*x/3 + I)/64`

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\int \sqrt{3ix + 4x^2} dx = \frac{1}{2} \sqrt{4x^2 + 3ix} + \frac{3}{16} i \sqrt{4x^2 + 3ix} + \frac{9}{64} \log(8x + 4\sqrt{4x^2 + 3ix} + 3i)$$

[In] integrate((3\*I\*x+4\*x^2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(4\*x^2 + 3\*I\*x)\*x + 3/16\*I\*sqrt(4\*x^2 + 3\*I\*x) + 9/64\*log(8\*x + 4\*sqrt(4\*x^2 + 3\*I\*x) + 3\*I)

**Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(27) = 54.

Time = 0.27 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.56

$$\int \sqrt{3ix + 4x^2} dx = \frac{1}{32} \sqrt{8x^2 + 2\sqrt{16x^2 + 9x}(8x + 3i)} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - \frac{9}{64} \log \left( 2\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - 8x - 3i \right)$$

[In] integrate((3\*I\*x+4\*x^2)^(1/2),x, algorithm="giac")

[Out] 1/32\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(8\*x + 3\*I)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 9/64\*log(2\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 8\*x - 3\*I)

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

$$\int \sqrt{3ix + 4x^2} dx = \frac{9 \ln \left( x + \frac{\sqrt{x(4x+3i)}}{2} + \frac{3}{8}i \right)}{64} + \left( \frac{x}{2} + \frac{3}{16}i \right) \sqrt{4x^2 + x3i}$$

[In] int((x\*3i + 4\*x^2)^(1/2),x)

[Out] (9\*log(x + (x\*(4\*x + 3i))^(1/2)/2 + 3i/8))/64 + (x/2 + 3i/16)\*(x\*3i + 4\*x^2)^(1/2)

## 3.6 $\int (3x - 4x^2)^{7/2} dx$

Optimal result . . . . .	93
Rubi [A] (verified) . . . . .	93
Mathematica [A] (verified) . . . . .	95
Maple [A] (verified) . . . . .	95
Fricas [A] (verification not implemented) . . . . .	96
Sympy [B] (verification not implemented) . . . . .	96
Maxima [A] (verification not implemented) . . . . .	97
Giac [A] (verification not implemented) . . . . .	97
Mupad [B] (verification not implemented) . . . . .	98

### Optimal result

Integrand size = 13, antiderivative size = 101

$$\int (3x - 4x^2)^{7/2} dx = -\frac{25515(3 - 8x)\sqrt{3x - 4x^2}}{4194304} - \frac{945(3 - 8x)(3x - 4x^2)^{3/2}}{131072} \\ - \frac{21(3 - 8x)(3x - 4x^2)^{5/2}}{2048} - \frac{1}{64}(3 - 8x)(3x - 4x^2)^{7/2} - \frac{229635 \arcsin\left(1 - \frac{8x}{3}\right)}{16777216}$$

[Out] -945/131072\*(3-8\*x)\*(-4\*x^2+3\*x)^(3/2)-21/2048\*(3-8\*x)\*(-4\*x^2+3\*x)^(5/2)-1/64\*(3-8\*x)\*(-4\*x^2+3\*x)^(7/2)+229635/16777216\*arcsin(-1+8/3\*x)-25515/4194304\*(3-8\*x)\*(-4\*x^2+3\*x)^(1/2)

### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int (3x - 4x^2)^{7/2} dx = -\frac{229635 \arcsin\left(1 - \frac{8x}{3}\right)}{16777216} - \frac{1}{64}(3 - 8x)(3x - 4x^2)^{7/2} \\ - \frac{21(3 - 8x)(3x - 4x^2)^{5/2}}{2048} - \frac{945(3 - 8x)(3x - 4x^2)^{3/2}}{131072} - \frac{25515(3 - 8x)\sqrt{3x - 4x^2}}{4194304}$$

[In] Int[(3\*x - 4\*x^2)^(7/2), x]

[Out] (-25515\*(3 - 8\*x)\*Sqrt[3\*x - 4\*x^2])/4194304 - (945\*(3 - 8\*x)\*(3\*x - 4\*x^2)^(3/2))/131072 - (21\*(3 - 8\*x)\*(3\*x - 4\*x^2)^(5/2))/2048 - ((3 - 8\*x)\*(3\*x - 4\*x^2)^(7/2))/64 - (229635\*ArcSin[1 - (8\*x)/3])/16777216

Rule 222

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

### Rule 626

`Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c) / (2*c*(2*p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

### Rule 633

`Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{64}(3-8x)(3x-4x^2)^{7/2} + \frac{63}{128} \int (3x-4x^2)^{5/2} dx \\
 &= -\frac{21(3-8x)(3x-4x^2)^{5/2}}{2048} - \frac{1}{64}(3-8x)(3x-4x^2)^{7/2} + \frac{945 \int (3x-4x^2)^{3/2} dx}{4096} \\
 &= -\frac{945(3-8x)(3x-4x^2)^{3/2}}{131072} - \frac{21(3-8x)(3x-4x^2)^{5/2}}{2048} \\
 &\quad - \frac{1}{64}(3-8x)(3x-4x^2)^{7/2} + \frac{25515 \int \sqrt{3x-4x^2} dx}{262144} \\
 &= -\frac{25515(3-8x)\sqrt{3x-4x^2}}{4194304} - \frac{945(3-8x)(3x-4x^2)^{3/2}}{131072} \\
 &\quad - \frac{21(3-8x)(3x-4x^2)^{5/2}}{2048} - \frac{1}{64}(3-8x)(3x-4x^2)^{7/2} + \frac{229635 \int \frac{1}{\sqrt{3x-4x^2}} dx}{8388608} \\
 &= -\frac{25515(3-8x)\sqrt{3x-4x^2}}{4194304} - \frac{945(3-8x)(3x-4x^2)^{3/2}}{131072} - \frac{21(3-8x)(3x-4x^2)^{5/2}}{2048} \\
 &\quad - \frac{1}{64}(3-8x)(3x-4x^2)^{7/2} - \frac{76545 \text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{9}}} dx, x, 3-8x\right)}{16777216} \\
 &= -\frac{25515(3-8x)\sqrt{3x-4x^2}}{4194304} - \frac{945(3-8x)(3x-4x^2)^{3/2}}{131072} \\
 &\quad - \frac{21(3-8x)(3x-4x^2)^{5/2}}{2048} - \frac{1}{64}(3-8x)(3x-4x^2)^{7/2} - \frac{229635 \sin^{-1}\left(1-\frac{8x}{3}\right)}{16777216}
 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.17 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.01

$$\int (3x - 4x^2)^{7/2} dx = \frac{\sqrt{-x(-3+4x)}(-2\sqrt{x}\sqrt{-3+4x}(76545+68040x+72576x^2+82944x^3-25067520x^4+79429632x^5-88080384x^6+33554432x^7)+229635\sqrt{x}\sqrt{-3+4x})}{8388608\sqrt{x}\sqrt{-3+4x}}$$

`[In] Integrate[(3*x - 4*x^2)^(7/2), x]`

```
[Out] (Sqrt[-(x*(-3 + 4*x))]*(-2*Sqrt[x]*Sqrt[-3 + 4*x]*(76545 + 68040*x + 72576*x^2 + 82944*x^3 - 25067520*x^4 + 79429632*x^5 - 88080384*x^6 + 33554432*x^7) + 229635*Log[-2*Sqrt[x] + Sqrt[-3 + 4*x]]))/(8388608*Sqrt[x]*Sqrt[-3 + 4*x])
```

**Maple [A] (verified)**

Time = 2.43 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.62

method	result
risch	$\frac{(33554432x^7 - 88080384x^6 + 79429632x^5 - 25067520x^4 + 82944x^3 + 72576x^2 + 68040x + 76545)x(4x-3)}{4194304\sqrt{-x(4x-3)}} + \frac{229635 \arcsin\left(-1 + \frac{8x}{3}\right)}{16777216}$
meijerg	$688905i \left( -\frac{i\sqrt{\pi}\sqrt{x}\sqrt{3}\left(\frac{33554432}{243}x^7 - \frac{29360128}{81}x^6 + \frac{26476544}{81}x^5 - \frac{2785280}{27}x^4 + \frac{1024}{3}x^3 + \frac{896}{3}x^2 + 280x + 315\right)\sqrt{-\frac{4x}{3}+1}}{1451520} + \frac{i\sqrt{\pi} \arcsin\left(\frac{2\sqrt{3}\sqrt{x}}{3}\right)}{3072} \right)$
default	$-\frac{945(3-8x)(-4x^2+3x)^{\frac{3}{2}}}{131072} - \frac{21(3-8x)(-4x^2+3x)^{\frac{5}{2}}}{2048} - \frac{(3-8x)(-4x^2+3x)^{\frac{7}{2}}}{64} + \frac{229635 \arcsin\left(-1 + \frac{8x}{3}\right)}{16777216} - \frac{25515(3-8x)\sqrt{-4x^2+3x}}{4194304}$
trager	$\left(-8x^7 + 21x^6 - \frac{303}{16}x^5 + \frac{765}{128}x^4 - \frac{81}{4096}x^3 - \frac{567}{32768}x^2 - \frac{8505}{524288}x - \frac{76545}{4194304}\right)\sqrt{-4x^2+3x} + \frac{229635 \operatorname{RootOf}(x^3 - 4x^2 + 3x - 1)}{16777216}$

`[In] int((-4*x^2+3*x)^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/4194304*(33554432*x^7-88080384*x^6+79429632*x^5-25067520*x^4+82944*x^3+72576*x^2+68040*x+76545)*x*(4*x-3)/(-x*(4*x-3))^(1/2)+229635/16777216*arcsin(-1+8/3*x)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.38 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.67

$$\int (3x - 4x^2)^{7/2} dx =$$

$$-\frac{1}{4194304} (33554432 x^7 - 88080384 x^6 + 79429632 x^5 - 25067520 x^4 + 82944 x^3 + 72576 x^2 + 68040 x + 76545) \sqrt{-4x^2 + 3x} - \frac{229635}{8388608} \arctan\left(\frac{\sqrt{-4x^2 + 3x}}{2x}\right)$$

[In] integrate((-4\*x^2+3\*x)^(7/2),x, algorithm="fricas")

[Out] -1/4194304\*(33554432\*x^7 - 88080384\*x^6 + 79429632\*x^5 - 25067520\*x^4 + 82944\*x^3 + 72576\*x^2 + 68040\*x + 76545)\*sqrt(-4\*x^2 + 3\*x) - 229635/8388608\*arctan(1/2\*sqrt(-4\*x^2 + 3\*x)/x)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. 2(90) = 180.

Time = 0.71 (sec) , antiderivative size = 211, normalized size of antiderivative = 2.09

$$\int (3x - 4x^2)^{7/2} dx = 27\sqrt{-4x^2 + 3x} \left( \frac{x^4}{5} - \frac{3x^3}{160} - \frac{21x^2}{1280} - \frac{63x}{4096} - \frac{567}{32768} \right)$$

$$- 108\sqrt{-4x^2 + 3x} \left( \frac{x^5}{6} - \frac{x^4}{80} - \frac{27x^3}{2560} - \frac{189x^2}{20480} - \frac{567x}{65536} - \frac{5103}{524288} \right)$$

$$+ 144\sqrt{-4x^2 + 3x} \left( \frac{x^6}{7} - \frac{x^5}{112} - \frac{33x^4}{4480} - \frac{891x^3}{143360} - \frac{891x^2}{163840} - \frac{2673x}{524288} - \frac{24057}{4194304} \right)$$

$$- 64\sqrt{-4x^2 + 3x} \left( \frac{x^7}{8} - \frac{3x^6}{448} - \frac{39x^5}{7168} - \frac{1287x^4}{286720} - \frac{34749x^3}{9175040} - \frac{34749x^2}{10485760} - \frac{104247x}{33554432} - \frac{938223}{268435456} \right) + \frac{229635 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{16777216}$$

[In] integrate((-4\*x\*\*2+3\*x)\*\*(7/2),x)

[Out] 27\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*4/5 - 3\*x\*\*3/160 - 21\*x\*\*2/1280 - 63\*x/4096 - 567/32768) - 108\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*5/6 - x\*\*4/80 - 27\*x\*\*3/2560 - 189\*x\*\*2/20480 - 567\*x/65536 - 5103/524288) + 144\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*6/7 - x\*\*5/112 - 33\*x\*\*4/4480 - 891\*x\*\*3/143360 - 891\*x\*\*2/163840 - 2673\*x/524288 - 24057/4194304) - 64\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*7/8 - 3\*x\*\*6/448 - 39\*x\*\*5/7168 - 1287\*x\*\*4/286720 - 34749\*x\*\*3/9175040 - 34749\*x\*\*2/10485760 - 104247\*x/33554432 - 938223/268435456) + 229635\*asin(8\*x/3 - 1)/16777216



**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.16

$$\int (3x - 4x^2)^{7/2} dx = \frac{1}{8} (-4x^2 + 3x)^{7/2} x - \frac{3}{64} (-4x^2 + 3x)^{7/2} + \frac{21}{256} (-4x^2 + 3x)^{5/2} x - \frac{63}{2048} (-4x^2 + 3x)^{5/2} + \frac{945}{16384} (-4x^2 + 3x)^{3/2} x - \frac{2835}{131072} (-4x^2 + 3x)^{3/2} + \frac{25515}{524288} \sqrt{-4x^2 + 3x} x - \frac{76545}{4194304} \sqrt{-4x^2 + 3x} - \frac{229635}{16777216} \arcsin\left(-\frac{8}{3}x + 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(7/2),x, algorithm="maxima")

```
[Out] 1/8*(-4*x^2 + 3*x)^(7/2)*x - 3/64*(-4*x^2 + 3*x)^(7/2) + 21/256*(-4*x^2 + 3*x)^(5/2)*x - 63/2048*(-4*x^2 + 3*x)^(5/2) + 945/16384*(-4*x^2 + 3*x)^(3/2)*x - 2835/131072*(-4*x^2 + 3*x)^(3/2) + 25515/524288*sqrt(-4*x^2 + 3*x)*x - 76545/4194304*sqrt(-4*x^2 + 3*x) - 229635/16777216*arcsin(-8/3*x + 1)
```

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.56

$$\int (3x - 4x^2)^{7/2} dx = -\frac{1}{4194304} (8(16(8(32(8(16(8x - 21)x + 303)x - 765)x + 81)x + 567)x + 8505)x + 76545)\sqrt{-4x^2 + 3x} + \frac{229635}{16777216} \arcsin\left(\frac{8}{3}x - 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(7/2),x, algorithm="giac")

```
[Out] -1/4194304*(8*(16*(8*(32*(8*(16*(8*x - 21)*x + 303)*x - 765)*x + 81)*x + 567)*x + 8505)*x + 76545)*sqrt(-4*x^2 + 3*x) + 229635/16777216*arcsin(8/3*x - 1)
```

**Mupad [B] (verification not implemented)**

Time = 9.30 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.80

$$\int (3x - 4x^2)^{7/2} dx = \frac{229635 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{16777216} + \frac{945 \left(4x - \frac{3}{2}\right) (3x - 4x^2)^{3/2}}{65536}$$

$$+ \frac{21 \left(4x - \frac{3}{2}\right) (3x - 4x^2)^{5/2}}{1024} + \frac{\left(4x - \frac{3}{2}\right) (3x - 4x^2)^{7/2}}{32} + \frac{25515 \left(\frac{x}{2} - \frac{3}{16}\right) \sqrt{3x - 4x^2}}{262144}$$

[In] `int((3*x - 4*x^2)^(7/2),x)`

[Out] `(229635*asin((8*x)/3 - 1))/16777216 + (945*(4*x - 3/2)*(3*x - 4*x^2)^(3/2))/65536 + (21*(4*x - 3/2)*(3*x - 4*x^2)^(5/2))/1024 + ((4*x - 3/2)*(3*x - 4*x^2)^(7/2))/32 + (25515*(x/2 - 3/16)*(3*x - 4*x^2)^(1/2))/262144`

### 3.7 $\int (3x - 4x^2)^{5/2} dx$

Optimal result . . . . .	99
Rubi [A] (verified) . . . . .	99
Mathematica [A] (verified) . . . . .	101
Maple [A] (verified) . . . . .	101
Fricas [A] (verification not implemented) . . . . .	101
Sympy [A] (verification not implemented) . . . . .	102
Maxima [A] (verification not implemented) . . . . .	102
Giac [A] (verification not implemented) . . . . .	103
Mupad [B] (verification not implemented) . . . . .	103

#### Optimal result

Integrand size = 13, antiderivative size = 79

$$\int (3x - 4x^2)^{5/2} dx = -\frac{405(3 - 8x)\sqrt{3x - 4x^2}}{32768} - \frac{15(3 - 8x)(3x - 4x^2)^{3/2}}{1024} - \frac{1}{48}(3 - 8x)(3x - 4x^2)^{5/2} - \frac{3645 \arcsin\left(1 - \frac{8x}{3}\right)}{131072}$$

[Out] -15/1024\*(3-8\*x)\*(-4\*x^2+3\*x)^(3/2)-1/48\*(3-8\*x)\*(-4\*x^2+3\*x)^(5/2)+3645/131072\*arcsin(-1+8/3\*x)-405/32768\*(3-8\*x)\*(-4\*x^2+3\*x)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int (3x - 4x^2)^{5/2} dx = -\frac{3645 \arcsin\left(1 - \frac{8x}{3}\right)}{131072} - \frac{1}{48}(3 - 8x)(3x - 4x^2)^{5/2} - \frac{15(3 - 8x)(3x - 4x^2)^{3/2}}{1024} - \frac{405(3 - 8x)\sqrt{3x - 4x^2}}{32768}$$

[In] Int[(3\*x - 4\*x^2)^(5/2), x]

[Out] (-405\*(3 - 8\*x)\*Sqrt[3\*x - 4\*x^2])/32768 - (15\*(3 - 8\*x)\*(3\*x - 4\*x^2)^(3/2))/1024 - ((3 - 8\*x)\*(3\*x - 4\*x^2)^(5/2))/48 - (3645\*ArcSin[1 - (8\*x)/3])/131072

Rule 222

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt
[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]
```

### Rule 626

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N
eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

### Rule 633

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

### Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{1}{48}(3-8x)(3x-4x^2)^{5/2} + \frac{15}{32} \int (3x-4x^2)^{3/2} dx \\
&= -\frac{15(3-8x)(3x-4x^2)^{3/2}}{1024} - \frac{1}{48}(3-8x)(3x-4x^2)^{5/2} + \frac{405 \int \sqrt{3x-4x^2} dx}{2048} \\
&= -\frac{405(3-8x)\sqrt{3x-4x^2}}{32768} - \frac{15(3-8x)(3x-4x^2)^{3/2}}{1024} \\
&\quad - \frac{1}{48}(3-8x)(3x-4x^2)^{5/2} + \frac{3645 \int \frac{1}{\sqrt{3x-4x^2}} dx}{65536} \\
&= -\frac{405(3-8x)\sqrt{3x-4x^2}}{32768} - \frac{15(3-8x)(3x-4x^2)^{3/2}}{1024} \\
&\quad - \frac{1}{48}(3-8x)(3x-4x^2)^{5/2} - \frac{1215 \text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{9}}} dx, x, 3-8x\right)}{131072} \\
&= -\frac{405(3-8x)\sqrt{3x-4x^2}}{32768} - \frac{15(3-8x)(3x-4x^2)^{3/2}}{1024} \\
&\quad - \frac{1}{48}(3-8x)(3x-4x^2)^{5/2} - \frac{3645 \sin^{-1}\left(1-\frac{8x}{3}\right)}{131072}
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.16

$$\int (3x - 4x^2)^{5/2} dx = \frac{\sqrt{-x(-3+4x)}(2\sqrt{x}\sqrt{-3+4x}(-3645-3240x-3456x^2+248832x^3-491520x^4+262144x^5) + 10935\sqrt{x}\sqrt{-3+4x}\log[-2\sqrt{x}\sqrt{-3+4x}])}{196608\sqrt{x}\sqrt{-3+4x}}$$

`[In] Integrate[(3*x - 4*x^2)^(5/2),x]`

```
[Out] (Sqrt[-(x*(-3 + 4*x))]*(2*Sqrt[x]*Sqrt[-3 + 4*x]*(-3645 - 3240*x - 3456*x^2 + 248832*x^3 - 491520*x^4 + 262144*x^5) + 10935*Log[-2*Sqrt[x] + Sqrt[-3 + 4*x]]))/(196608*Sqrt[x]*Sqrt[-3 + 4*x])
```

**Maple [A] (verified)**

Time = 2.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.67

method	result
risch	$-\frac{(262144x^5-491520x^4+248832x^3-3456x^2-3240x-3645)x(4x-3)}{98304\sqrt{-x(4x-3)}} + \frac{3645 \arcsin(-1+\frac{8x}{3})}{131072}$
default	$-\frac{15(3-8x)(-4x^2+3x)^{\frac{3}{2}}}{1024} - \frac{(3-8x)(-4x^2+3x)^{\frac{5}{2}}}{48} + \frac{3645 \arcsin(-1+\frac{8x}{3})}{131072} - \frac{405(3-8x)\sqrt{-4x^2+3x}}{32768}$
meijerg	$-\frac{10935i \left( -\frac{i\sqrt{\pi}\sqrt{x}\sqrt{3}(-\frac{1835008}{243}x^5 + \frac{1146880}{81}x^4 - 7168x^3 + \frac{896}{9}x^2 + \frac{280}{3}x + 105)\sqrt{-\frac{4x}{3}+1} + \frac{i\sqrt{\pi} \arcsin(\frac{2\sqrt{3}\sqrt{x}}{3})}{192} \right)}{1024\sqrt{\pi}}$
trager	$\left(\frac{8}{3}x^5 - 5x^4 + \frac{81}{32}x^3 - \frac{9}{256}x^2 - \frac{135}{4096}x - \frac{1215}{32768}\right)\sqrt{-4x^2+3x} - \frac{3645 \operatorname{RootOf}(\_Z^2+1) \ln(8 \operatorname{RootOf}(\_Z^2+1))}{131072}$

`[In] int((-4*x^2+3*x)^(5/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/98304*(262144*x^5-491520*x^4+248832*x^3-3456*x^2-3240*x-3645)*x*(4*x-3)/(-x*(4*x-3))^(1/2)+3645/131072*arcsin(-1+8/3*x)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.35 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.73

$$\int (3x - 4x^2)^{5/2} dx = \frac{1}{98304} (262144x^5 - 491520x^4 + 248832x^3 - 3456x^2 - 3240x - 3645)\sqrt{-4x^2 + 3x} - \frac{3645}{65536} \arctan\left(\frac{\sqrt{-4x^2 + 3x}}{2x}\right)$$

[In] integrate((-4\*x^2+3\*x)^(5/2),x, algorithm="fricas")

[Out] 1/98304\*(262144\*x^5 - 491520\*x^4 + 248832\*x^3 - 3456\*x^2 - 3240\*x - 3645)\*sqrt(-4\*x^2 + 3\*x) - 3645/65536\*arctan(1/2\*sqrt(-4\*x^2 + 3\*x)/x)

### Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.63

$$\int (3x - 4x^2)^{5/2} dx = 9\sqrt{-4x^2 + 3x}\left(\frac{x^3}{4} - \frac{x^2}{32} - \frac{15x}{512} - \frac{135}{4096}\right) - 24\sqrt{-4x^2 + 3x}\left(\frac{x^4}{5} - \frac{3x^3}{160} - \frac{21x^2}{1280} - \frac{63x}{4096} - \frac{567}{32768}\right) + 16\sqrt{-4x^2 + 3x}\left(\frac{x^5}{6} - \frac{x^4}{80} - \frac{27x^3}{2560} - \frac{189x^2}{20480} - \frac{567x}{65536} - \frac{5103}{524288}\right) + \frac{3645 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{131072}$$

[In] integrate((-4\*x\*\*2+3\*x)\*\*(5/2),x)

[Out] 9\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*3/4 - x\*\*2/32 - 15\*x/512 - 135/4096) - 24\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*4/5 - 3\*x\*\*3/160 - 21\*x\*\*2/1280 - 63\*x/4096 - 567/32768) + 16\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*5/6 - x\*\*4/80 - 27\*x\*\*3/2560 - 189\*x\*\*2/20480 - 567\*x/65536 - 5103/524288) + 3645\*asin(8\*x/3 - 1)/131072

### Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.14

$$\int (3x - 4x^2)^{5/2} dx = \frac{1}{6} (-4x^2 + 3x)^{5/2} x - \frac{1}{16} (-4x^2 + 3x)^{5/2} + \frac{15}{128} (-4x^2 + 3x)^{3/2} x - \frac{45}{1024} (-4x^2 + 3x)^{3/2} + \frac{405}{4096} \sqrt{-4x^2 + 3x} - \frac{1215}{32768} \sqrt{-4x^2 + 3x} - \frac{3645}{131072} \arcsin\left(-\frac{8}{3}x + 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(5/2),x, algorithm="maxima")

[Out] 1/6\*(-4\*x^2 + 3\*x)^(5/2)\*x - 1/16\*(-4\*x^2 + 3\*x)^(5/2) + 15/128\*(-4\*x^2 + 3\*x)^(3/2)\*x - 45/1024\*(-4\*x^2 + 3\*x)^(3/2) + 405/4096\*sqrt(-4\*x^2 + 3\*x)\*x - 1215/32768\*sqrt(-4\*x^2 + 3\*x) - 3645/131072\*arcsin(-8/3\*x + 1)

### Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.59

$$\int (3x - 4x^2)^{5/2} dx = \frac{1}{98304} (8 (16 (8 (32 (8x - 15)x + 243)x - 27)x - 405)x - 3645) \sqrt{-4x^2 + 3x} + \frac{3645}{131072} \arcsin\left(\frac{8}{3}x - 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(5/2),x, algorithm="giac")

[Out] 1/98304\*(8\*(16\*(8\*(32\*(8\*x - 15)\*x + 243)\*x - 27)\*x - 405)\*x - 3645)\*sqrt(-4\*x^2 + 3\*x) + 3645/131072\*arcsin(8/3\*x - 1)

### Mupad [B] (verification not implemented)

Time = 9.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.80

$$\int (3x - 4x^2)^{5/2} dx = \frac{3645 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{131072} + \frac{15 \left(4x - \frac{3}{2}\right) (3x - 4x^2)^{3/2}}{512} + \frac{\left(4x - \frac{3}{2}\right) (3x - 4x^2)^{5/2}}{24} + \frac{405 \left(\frac{x}{2} - \frac{3}{16}\right) \sqrt{3x - 4x^2}}{2048}$$

[In] int((3\*x - 4\*x^2)^(5/2),x)

[Out] (3645\*asin((8\*x)/3 - 1))/131072 + (15\*(4\*x - 3/2)\*(3\*x - 4\*x^2)^(3/2))/512 + ((4\*x - 3/2)\*(3\*x - 4\*x^2)^(5/2))/24 + (405\*(x/2 - 3/16)\*(3\*x - 4\*x^2)^(1/2))/2048

### 3.8 $\int (3x - 4x^2)^{3/2} dx$

Optimal result	104
Rubi [A] (verified)	104
Mathematica [A] (verified)	105
Maple [A] (verified)	106
Fricas [A] (verification not implemented)	106
Sympy [A] (verification not implemented)	107
Maxima [A] (verification not implemented)	107
Giac [A] (verification not implemented)	107
Mupad [B] (verification not implemented)	108

#### Optimal result

Integrand size = 13, antiderivative size = 57

$$\int (3x - 4x^2)^{3/2} dx = -\frac{27(3 - 8x)\sqrt{3x - 4x^2}}{1024} - \frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} - \frac{243 \arcsin\left(1 - \frac{8x}{3}\right)}{4096}$$

[Out]  $-1/32*(3-8*x)*(-4*x^2+3*x)^(3/2)+243/4096*\arcsin(-1+8/3*x)-27/1024*(3-8*x)*(-4*x^2+3*x)^(1/2)$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int (3x - 4x^2)^{3/2} dx = -\frac{243 \arcsin\left(1 - \frac{8x}{3}\right)}{4096} - \frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} - \frac{27(3 - 8x)\sqrt{3x - 4x^2}}{1024}$$

[In]  $\text{Int}[(3*x - 4*x^2)^(3/2), x]$

[Out]  $(-27*(3 - 8*x)*\text{Sqrt}[3*x - 4*x^2])/1024 - ((3 - 8*x)*(3*x - 4*x^2)^(3/2))/32 - (243*\text{ArcSin}[1 - (8*x)/3])/4096$

#### Rule 222

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{NegQ}[b]$



Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^p / (2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c) / (2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} + \frac{27}{64} \int \sqrt{3x - 4x^2} dx \\
 &= -\frac{27(3 - 8x)\sqrt{3x - 4x^2}}{1024} - \frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} + \frac{243 \int \frac{1}{\sqrt{3x - 4x^2}} dx}{2048} \\
 &= -\frac{27(3 - 8x)\sqrt{3x - 4x^2}}{1024} - \frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} - \frac{81 \text{Subst}\left(\int \frac{1}{\sqrt{1 - \frac{x^2}{9}}} dx, x, 3 - 8x\right)}{4096} \\
 &= -\frac{27(3 - 8x)\sqrt{3x - 4x^2}}{1024} - \frac{1}{32}(3 - 8x)(3x - 4x^2)^{3/2} - \frac{243 \sin^{-1}\left(1 - \frac{8x}{3}\right)}{4096}
 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.44

$$\int (3x - 4x^2)^{3/2} dx = \frac{\sqrt{-x(-3 + 4x)}(-2\sqrt{x}\sqrt{-3 + 4x}(81 + 72x - 1152x^2 + 1024x^3) + 243 \log(-2\sqrt{x} + \sqrt{-3 + 4x}))}{2048\sqrt{x}\sqrt{-3 + 4x}}$$

[In] Integrate[(3\*x - 4\*x^2)^(3/2), x]

[Out] (Sqrt[-(x\*(-3 + 4\*x))]\*(-2\*Sqrt[x]\*Sqrt[-3 + 4\*x]\*(81 + 72\*x - 1152\*x^2 + 1024\*x^3) + 243\*Log[-2\*Sqrt[x] + Sqrt[-3 + 4\*x]]))/(2048\*Sqrt[x]\*Sqrt[-3 + 4\*x])

**Maple [A] (verified)**

Time = 2.43 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.75

method	result
risch	$\frac{(1024x^3 - 1152x^2 + 72x + 81)x(4x - 3)}{1024\sqrt{-x(4x - 3)}} + \frac{243 \arcsin(-1 + \frac{8x}{3})}{4096}$
default	$-\frac{(3-8x)(-4x^2+3x)^{\frac{3}{2}}}{32} + \frac{243 \arcsin(-1 + \frac{8x}{3})}{4096} - \frac{27(3-8x)\sqrt{-4x^2+3x}}{1024}$
pseudoelliptic	$-\frac{243 \arctan\left(\frac{\sqrt{-4x^2+3x}}{2x}\right)}{2048} + \frac{(-1024x^3 + 1152x^2 - 72x - 81)\sqrt{-4x^2+3x}}{1024}$
meijerg	$-\frac{243i \left( -\frac{i\sqrt{\pi}\sqrt{x}\sqrt{3}\left(\frac{5120}{27}x^3 - \frac{640}{3}x^2 + \frac{40}{3}x + 15\right)\sqrt{-\frac{4x}{3}+1}}{360} + \frac{i\sqrt{\pi}\arcsin\left(\frac{2\sqrt{3}\sqrt{x}}{3}\right)}{16} \right)}{128\sqrt{\pi}}$
trager	$\left(-x^3 + \frac{9}{8}x^2 - \frac{9}{128}x - \frac{81}{1024}\right)\sqrt{-4x^2+3x} + \frac{243 \operatorname{RootOf}(\_Z^2+1) \ln(-8 \operatorname{RootOf}(\_Z^2+1)x + 4\sqrt{-4x^2+3x})}{4096}$

[In] int((-4\*x^2+3\*x)^(3/2),x,method=\_RETURNVERBOSE)

[Out] 1/1024\*(1024\*x^3-1152\*x^2+72\*x+81)\*x\*(4\*x-3)/(-x\*(4\*x-3))^(1/2)+243/4096\*arcsin(-1+8/3\*x)

**Fricas [A] (verification not implemented)**

none

Time = 0.41 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.84

$$\int (3x - 4x^2)^{3/2} dx = -\frac{1}{1024} (1024x^3 - 1152x^2 + 72x + 81)\sqrt{-4x^2 + 3x} - \frac{243}{2048} \arctan\left(\frac{\sqrt{-4x^2 + 3x}}{2x}\right)$$

[In] integrate((-4\*x^2+3\*x)^(3/2),x, algorithm="fricas")

[Out] -1/1024\*(1024\*x^3 - 1152\*x^2 + 72\*x + 81)\*sqrt(-4\*x^2 + 3\*x) - 243/2048\*arctan(1/2\*sqrt(-4\*x^2 + 3\*x)/x)

**Sympy [A] (verification not implemented)**

Time = 0.40 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.19

$$\int (3x - 4x^2)^{3/2} dx = 3\sqrt{-4x^2 + 3x} \left( \frac{x^2}{3} - \frac{x}{16} - \frac{9}{128} \right) - 4\sqrt{-4x^2 + 3x} \left( \frac{x^3}{4} - \frac{x^2}{32} - \frac{15x}{512} - \frac{135}{4096} \right) + \frac{243 \operatorname{asin} \left( \frac{8x}{3} - 1 \right)}{4096}$$

[In] integrate((-4\*x\*\*2+3\*x)\*\*(3/2),x)

[Out] 3\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*2/3 - x/16 - 9/128) - 4\*sqrt(-4\*x\*\*2 + 3\*x)\*(x\*\*3/4 - x\*\*2/32 - 15\*x/512 - 135/4096) + 243\*asin(8\*x/3 - 1)/4096

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.11

$$\int (3x - 4x^2)^{3/2} dx = \frac{1}{4} (-4x^2 + 3x)^{\frac{3}{2}} x - \frac{3}{32} (-4x^2 + 3x)^{\frac{3}{2}} + \frac{27}{128} \sqrt{-4x^2 + 3x} x - \frac{81}{1024} \sqrt{-4x^2 + 3x} - \frac{243}{4096} \arcsin \left( -\frac{8}{3} x + 1 \right)$$

[In] integrate((-4\*x^2+3\*x)^(3/2),x, algorithm="maxima")

[Out] 1/4\*(-4\*x^2 + 3\*x)^(3/2)\*x - 3/32\*(-4\*x^2 + 3\*x)^(3/2) + 27/128\*sqrt(-4\*x^2 + 3\*x)\*x - 81/1024\*sqrt(-4\*x^2 + 3\*x) - 243/4096\*arcsin(-8/3\*x + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.65

$$\int (3x - 4x^2)^{3/2} dx = -\frac{1}{1024} (8(16(8x - 9)x + 9)x + 81)\sqrt{-4x^2 + 3x} + \frac{243}{4096} \arcsin \left( \frac{8}{3} x - 1 \right)$$

[In] integrate((-4\*x^2+3\*x)^(3/2),x, algorithm="giac")

[Out] -1/1024\*(8\*(16\*(8\*x - 9)\*x + 9)\*x + 81)\*sqrt(-4\*x^2 + 3\*x) + 243/4096\*arcsin(8/3\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.79

$$\int (3x-4x^2)^{3/2} dx = \frac{243 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{4096} + \frac{(4x - \frac{3}{2})(3x - 4x^2)^{3/2}}{16} + \frac{27\left(\frac{x}{2} - \frac{3}{16}\right)\sqrt{3x - 4x^2}}{64}$$

[In] `int((3*x - 4*x^2)^(3/2),x)`

[Out] `(243*asin((8*x)/3 - 1))/4096 + ((4*x - 3/2)*(3*x - 4*x^2)^(3/2))/16 + (27*(x/2 - 3/16)*(3*x - 4*x^2)^(1/2))/64`

### 3.9 $\int \sqrt{3x - 4x^2} dx$

Optimal result	109
Rubi [A] (verified)	109
Mathematica [A] (verified)	110
Maple [A] (verified)	110
Fricas [A] (verification not implemented)	111
Sympy [A] (verification not implemented)	111
Maxima [A] (verification not implemented)	112
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	112

#### Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \sqrt{3x - 4x^2} dx = -\frac{1}{16}(3 - 8x)\sqrt{3x - 4x^2} - \frac{9}{64} \arcsin\left(1 - \frac{8x}{3}\right)$$

[Out] 9/64\*arcsin(-1+8/3\*x)-1/16\*(3-8\*x)\*(-4\*x^2+3\*x)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int \sqrt{3x - 4x^2} dx = -\frac{9}{64} \arcsin\left(1 - \frac{8x}{3}\right) - \frac{1}{16}\sqrt{3x - 4x^2}(3 - 8x)$$

[In] Int[Sqrt[3\*x - 4\*x^2], x]

[Out] -1/16\*((3 - 8\*x)\*Sqrt[3\*x - 4\*x^2]) - (9\*ArcSin[1 - (8\*x)/3])/64

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N

$eQ[b^2 - 4ac, 0] \ \&\& \ GtQ[p, 0] \ \&\& \ IntegerQ[4p]$

### Rule 633

$\text{Int}[(a_.) + (b_.)(x_) + (c_.)(x_)^2]^{(p_)}, x\_Symbol] \ :> \text{Dist}[1/(2c*(-4*(c/(b^2 - 4ac))))^{(p)}, \text{Subst}[\text{Int}[\text{Simp}[1 - x^2/(b^2 - 4ac)], x]^{(p)}, x], x, b + 2cx], x] \ /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ GtQ[4a - b^2/c, 0]$

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{16}(3 - 8x)\sqrt{3x - 4x^2} + \frac{9}{32} \int \frac{1}{\sqrt{3x - 4x^2}} dx \\ &= -\frac{1}{16}(3 - 8x)\sqrt{3x - 4x^2} - \frac{3}{64} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{9}}} dx, x, 3 - 8x \right) \\ &= -\frac{1}{16}(3 - 8x)\sqrt{3x - 4x^2} - \frac{9}{64} \sin^{-1} \left( 1 - \frac{8x}{3} \right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \sqrt{3x - 4x^2} dx = \frac{1}{32} \sqrt{-x(-3 + 4x)} \left( -6 + 16x + \frac{9 \log(-2\sqrt{x} + \sqrt{-3 + 4x})}{\sqrt{x}\sqrt{-3 + 4x}} \right)$$

[In] Integrate[Sqrt[3\*x - 4\*x^2], x]

[Out] (Sqrt[-(x\*(-3 + 4\*x))]\*(-6 + 16\*x + (9\*Log[-2\*Sqrt[x] + Sqrt[-3 + 4\*x]])/(Sqrt[x]\*Sqrt[-3 + 4\*x])))/32

### Maple [A] (verified)

Time = 1.99 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$\frac{9 \arcsin\left(-1 + \frac{8x}{3}\right)}{64} - \frac{(3-8x)\sqrt{-4x^2+3x}}{16}$
risch	$-\frac{(-3+8x)x(4x-3)}{16\sqrt{-x(4x-3)}} + \frac{9 \arcsin\left(-1 + \frac{8x}{3}\right)}{64}$
pseudoelliptic	$-\frac{9 \arctan\left(\frac{\sqrt{-4x^2+3x}}{2x}\right)}{32} + \frac{(-3+8x)\sqrt{-4x^2+3x}}{16}$
meijerg	$-\frac{9i \left( -\frac{i\sqrt{\pi} \sqrt{x} \sqrt{3} (3-8x) \sqrt{-\frac{4x}{3}+1}}{9} + \frac{i\sqrt{\pi} \arcsin\left(\frac{2\sqrt{3}\sqrt{x}}{3}\right)}{2} \right)}{16\sqrt{\pi}}$
trager	$\left(-\frac{3}{16} + \frac{x}{2}\right) \sqrt{-4x^2+3x} - \frac{9 \operatorname{RootOf}\left(\_Z^2+1\right) \ln\left(8 \operatorname{RootOf}\left(\_Z^2+1\right) x + 4\sqrt{-4x^2+3x} - 3 \operatorname{RootOf}\left(\_Z^2+1\right)\right)}{64}$

[In] `int((-4*x^2+3*x)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $9/64*\arcsin(-1+8/3*x)-1/16*(3-8*x)*(-4*x^2+3*x)^(1/2)$

### Fricas [A] (verification not implemented)

none

Time = 0.79 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \sqrt{3x - 4x^2} dx = \frac{1}{16} \sqrt{-4x^2 + 3x} (8x - 3) - \frac{9}{32} \arctan\left(\frac{\sqrt{-4x^2 + 3x}}{2x}\right)$$

[In] `integrate((-4*x^2+3*x)^(1/2),x, algorithm="fricas")`

[Out]  $1/16*\sqrt{-4*x^2 + 3*x}*(8*x - 3) - 9/32*\arctan(1/2*\sqrt{-4*x^2 + 3*x}/x)$

### Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \sqrt{3x - 4x^2} dx = \left(\frac{x}{2} - \frac{3}{16}\right) \sqrt{-4x^2 + 3x} + \frac{9 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{64}$$

[In] `integrate((-4*x**2+3*x)**(1/2),x)`

[Out]  $(x/2 - 3/16)*\sqrt{-4*x**2 + 3*x} + 9*\operatorname{asin}(8*x/3 - 1)/64$

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \sqrt{3x - 4x^2} dx = \frac{1}{2} \sqrt{-4x^2 + 3x} x - \frac{3}{16} \sqrt{-4x^2 + 3x} - \frac{9}{64} \arcsin\left(-\frac{8}{3}x + 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-4\*x^2 + 3\*x)\*x - 3/16\*sqrt(-4\*x^2 + 3\*x) - 9/64\*arcsin(-8/3\*x + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \sqrt{3x - 4x^2} dx = \frac{1}{16} \sqrt{-4x^2 + 3x}(8x - 3) + \frac{9}{64} \arcsin\left(\frac{8}{3}x - 1\right)$$

[In] integrate((-4\*x^2+3\*x)^(1/2),x, algorithm="giac")

[Out] 1/16\*sqrt(-4\*x^2 + 3\*x)\*(8\*x - 3) + 9/64\*arcsin(8/3\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 9.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sqrt{3x - 4x^2} dx = \frac{9 \operatorname{asin}\left(\frac{8x}{3} - 1\right)}{64} + \left(\frac{x}{2} - \frac{3}{16}\right) \sqrt{3x - 4x^2}$$

[In] int((3\*x - 4\*x^2)^(1/2),x)

[Out] (9\*asin((8\*x)/3 - 1))/64 + (x/2 - 3/16)\*(3\*x - 4\*x^2)^(1/2)



### 3.10 $\int \sqrt{6x - x^2} dx$

Optimal result	113
Rubi [A] (verified)	113
Mathematica [A] (verified)	114
Maple [A] (verified)	115
Fricas [A] (verification not implemented)	115
Sympy [A] (verification not implemented)	115
Maxima [A] (verification not implemented)	116
Giac [A] (verification not implemented)	116
Mupad [B] (verification not implemented)	116

#### Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \sqrt{6x - x^2} dx = -\frac{1}{2}(3 - x)\sqrt{6x - x^2} - \frac{9}{2} \arcsin\left(1 - \frac{x}{3}\right)$$

[Out] 9/2\*arcsin(-1+1/3\*x)-1/2\*(3-x)\*(-x^2+6\*x)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int \sqrt{6x - x^2} dx = -\frac{9}{2} \arcsin\left(1 - \frac{x}{3}\right) - \frac{1}{2}\sqrt{6x - x^2}(3 - x)$$

[In] Int[Sqrt[6\*x - x^2], x]

[Out] -1/2\*((3 - x)\*Sqrt[6\*x - x^2]) - (9\*ArcSin[1 - x/3])/2

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{2}(3-x)\sqrt{6x-x^2} + \frac{9}{2} \int \frac{1}{\sqrt{6x-x^2}} dx \\ &= -\frac{1}{2}(3-x)\sqrt{6x-x^2} - \frac{3}{4} \text{Subst} \left( \int \frac{1}{\sqrt{1-\frac{x^2}{36}}} dx, x, 6-2x \right) \\ &= -\frac{1}{2}(3-x)\sqrt{6x-x^2} - \frac{9}{2} \sin^{-1} \left( 1 - \frac{x}{3} \right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

$$\int \sqrt{6x-x^2} dx = \frac{1}{2} \sqrt{-((-6+x)x)} \left( -3+x + \frac{18 \log(\sqrt{-6+x} - \sqrt{x})}{\sqrt{-6+x}\sqrt{x}} \right)$$

[In] Integrate[Sqrt[6\*x - x^2],x]

[Out] (Sqrt[-((-6 + x)\*x)]\*(-3 + x + (18\*Log[Sqrt[-6 + x] - Sqrt[x]])/(Sqrt[-6 + x]\*Sqrt[x])))/2

**Maple [A] (verified)**

Time = 2.21 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

method	result	si
risch	$-\frac{(-3+x)x(-6+x)}{2\sqrt{-x(-6+x)}} + \frac{9 \arcsin(-1+\frac{x}{3})}{2}$	2
default	$-\frac{(6-2x)\sqrt{-x^2+6x}}{4} + \frac{9 \arcsin(-1+\frac{x}{3})}{2}$	2
pseudoelliptic	$-9 \arctan\left(\frac{\sqrt{-x(-6+x)}}{x}\right) + \frac{(-3+x)\sqrt{-x(-6+x)}}{2}$	3
meijerg	$18i \left( -\frac{i\sqrt{\pi}\sqrt{x}\sqrt{6(3-x)}\sqrt{-\frac{x}{6}+1}}{36} + \frac{i\sqrt{\pi}\arcsin\left(\frac{\sqrt{6}\sqrt{x}}{6}\right)}{2} \right)$	4
trager	$\left(-\frac{3}{2} + \frac{x}{2}\right) \sqrt{-x^2+6x} + \frac{9 \operatorname{RootOf}(-Z^2+1) \ln(-\operatorname{RootOf}(-Z^2+1)x + \sqrt{-x^2+6x} + 3 \operatorname{RootOf}(-Z^2+1))}{2}$	5

```
[In] int((-x^2+6*x)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2*(-3+x)*x*(-6+x)/(-x*(-6+x))^(1/2)+9/2*arcsin(-1+1/3*x)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \sqrt{6x-x^2} dx = \frac{1}{2} \sqrt{-x^2+6x}(x-3) - 9 \arctan\left(\frac{\sqrt{-x^2+6x}}{x}\right)$$

```
[In] integrate((-x^2+6*x)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*sqrt(-x^2 + 6*x)*(x - 3) - 9*arctan(sqrt(-x^2 + 6*x)/x)
```

**Sympy [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sqrt{6x-x^2} dx = \left(\frac{x}{2} - \frac{3}{2}\right) \sqrt{-x^2+6x} + \frac{9 \operatorname{asin}\left(\frac{x}{3} - 1\right)}{2}$$

```
[In] integrate((-x**2+6*x)**(1/2),x)
```

```
[Out] (x/2 - 3/2)*sqrt(-x**2 + 6*x) + 9*asin(x/3 - 1)/2
```

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 6x} x - \frac{3}{2} \sqrt{-x^2 + 6x} - \frac{9}{2} \arcsin\left(-\frac{1}{3}x + 1\right)$$

[In] integrate((-x^2+6\*x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-x^2 + 6\*x)\*x - 3/2\*sqrt(-x^2 + 6\*x) - 9/2\*arcsin(-1/3\*x + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \sqrt{6x - x^2} dx = \frac{1}{2} \sqrt{-x^2 + 6x}(x - 3) + \frac{9}{2} \arcsin\left(\frac{1}{3}x - 1\right)$$

[In] integrate((-x^2+6\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(-x^2 + 6\*x)\*(x - 3) + 9/2\*arcsin(1/3\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sqrt{6x - x^2} dx = \frac{9 \operatorname{asin}\left(\frac{x}{3} - 1\right)}{2} + \left(\frac{x}{2} - \frac{3}{2}\right) \sqrt{6x - x^2}$$

[In] int((6\*x - x^2)^(1/2),x)

[Out] (9\*asin(x/3 - 1))/2 + (x/2 - 3/2)\*(6\*x - x^2)^(1/2)

### 3.11 $\int \sqrt{5x - 9x^2} dx$

Optimal result	117
Rubi [A] (verified)	117
Mathematica [A] (verified)	118
Maple [A] (verified)	118
Fricas [A] (verification not implemented)	119
Sympy [A] (verification not implemented)	119
Maxima [A] (verification not implemented)	120
Giac [A] (verification not implemented)	120
Mupad [B] (verification not implemented)	120

#### Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \sqrt{5x - 9x^2} dx = -\frac{1}{36}(5 - 18x)\sqrt{5x - 9x^2} - \frac{25}{216} \arcsin\left(1 - \frac{18x}{5}\right)$$

[Out] 25/216\*arcsin(-1+18/5\*x)-1/36\*(5-18\*x)\*(-9\*x^2+5\*x)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {626, 633, 222}

$$\int \sqrt{5x - 9x^2} dx = -\frac{25}{216} \arcsin\left(1 - \frac{18x}{5}\right) - \frac{1}{36}\sqrt{5x - 9x^2}(5 - 18x)$$

[In] Int[Sqrt[5\*x - 9\*x^2],x]

[Out] -1/36\*((5 - 18\*x)\*Sqrt[5\*x - 9\*x^2]) - (25\*ArcSin[1 - (18\*x)/5])/216

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N

$eQ[b^2 - 4ac, 0] \ \&\& \ GtQ[p, 0] \ \&\& \ IntegerQ[4p]$

### Rule 633

$Int[(a_.) + (b_.)(x_) + (c_.)(x_)^2]^{(p_)}, x\_Symbol] \ :> \ Dist[1/(2c*(-4*(c/(b^2 - 4ac)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4ac)], x]^p, x], x, b + 2cx], x] \ /; \ FreeQ[\{a, b, c, p\}, x] \ \&\& \ GtQ[4a - b^2/c, 0]$

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{36}(5 - 18x)\sqrt{5x - 9x^2} + \frac{25}{72} \int \frac{1}{\sqrt{5x - 9x^2}} dx \\ &= -\frac{1}{36}(5 - 18x)\sqrt{5x - 9x^2} - \frac{5}{216} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{25}}} dx, x, 5 - 18x \right) \\ &= -\frac{1}{36}(5 - 18x)\sqrt{5x - 9x^2} - \frac{25}{216} \sin^{-1} \left( 1 - \frac{18x}{5} \right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.57

$$\int \sqrt{5x - 9x^2} dx = \frac{1}{108} \sqrt{-x(-5 + 9x)} \left( -15 + 54x + \frac{25 \log(-3\sqrt{x} + \sqrt{-5 + 9x})}{\sqrt{x}\sqrt{-5 + 9x}} \right)$$

[In] Integrate[Sqrt[5\*x - 9\*x^2], x]

[Out] (Sqrt[-(x\*(-5 + 9\*x))]\*(-15 + 54\*x + (25\*Log[-3\*Sqrt[x] + Sqrt[-5 + 9\*x]]))/(Sqrt[x]\*Sqrt[-5 + 9\*x]))/108

### Maple [A] (verified)

Time = 2.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

method	result
default	$\frac{25 \arcsin(-1 + \frac{18x}{5})}{216} - \frac{(5-18x)\sqrt{-9x^2+5x}}{36}$
risch	$-\frac{(-5+18x)x(9x-5)}{36\sqrt{-x(9x-5)}} + \frac{25 \arcsin(-1 + \frac{18x}{5})}{216}$
pseudoelliptic	$-\frac{25 \arctan\left(\frac{\sqrt{-9x^2+5x}}{3x}\right)}{108} + \frac{(-5+18x)\sqrt{-9x^2+5x}}{36}$
meijerg	$-\frac{25i \left( -\frac{i\sqrt{\pi} \sqrt{x} \sqrt{5} \left(-\frac{54x}{5} + 3\right) \sqrt{-\frac{9x}{5} + 1}}{10} + \frac{i\sqrt{\pi} \arcsin\left(\frac{3\sqrt{5}\sqrt{x}}{5}\right)}{2} \right)}{54\sqrt{\pi}}$
trager	$\left(-\frac{5}{36} + \frac{x}{2}\right) \sqrt{-9x^2 + 5x} - \frac{25 \operatorname{RootOf}(\_Z^2 + 1) \ln\left(18 \operatorname{RootOf}(\_Z^2 + 1)x - 5 \operatorname{RootOf}(\_Z^2 + 1) + 6\sqrt{-9x^2 + 5x}\right)}{216}$

[In] `int((-9*x^2+5*x)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `25/216*arcsin(-1+18/5*x)-1/36*(5-18*x)*(-9*x^2+5*x)^(1/2)`

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \sqrt{5x - 9x^2} dx = \frac{1}{36} \sqrt{-9x^2 + 5x}(18x - 5) - \frac{25}{108} \arctan\left(\frac{\sqrt{-9x^2 + 5x}}{3x}\right)$$

[In] `integrate((-9*x^2+5*x)^(1/2),x, algorithm="fricas")`

[Out] `1/36*sqrt(-9*x^2 + 5*x)*(18*x - 5) - 25/108*arctan(1/3*sqrt(-9*x^2 + 5*x)/x)`

### Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \sqrt{5x - 9x^2} dx = \left(\frac{x}{2} - \frac{5}{36}\right) \sqrt{-9x^2 + 5x} + \frac{25 \operatorname{asin}\left(\frac{18x}{5} - 1\right)}{216}$$

[In] `integrate((-9*x**2+5*x)**(1/2),x)`

[Out] `(x/2 - 5/36)*sqrt(-9*x**2 + 5*x) + 25*asin(18*x/5 - 1)/216`

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.03

$$\int \sqrt{5x - 9x^2} dx = \frac{1}{2} \sqrt{-9x^2 + 5x} x - \frac{5}{36} \sqrt{-9x^2 + 5x} - \frac{25}{216} \arcsin\left(-\frac{18}{5}x + 1\right)$$

[In] integrate((-9\*x^2+5\*x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-9\*x^2 + 5\*x)\*x - 5/36\*sqrt(-9\*x^2 + 5\*x) - 25/216\*arcsin(-18/5\*x + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.77

$$\int \sqrt{5x - 9x^2} dx = \frac{1}{36} \sqrt{-9x^2 + 5x}(18x - 5) + \frac{25}{216} \arcsin\left(\frac{18}{5}x - 1\right)$$

[In] integrate((-9\*x^2+5\*x)^(1/2),x, algorithm="giac")

[Out] 1/36\*sqrt(-9\*x^2 + 5\*x)\*(18\*x - 5) + 25/216\*arcsin(18/5\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 9.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \sqrt{5x - 9x^2} dx = \frac{25 \operatorname{asin}\left(\frac{18x}{5} - 1\right)}{216} + \left(\frac{x}{2} - \frac{5}{36}\right) \sqrt{5x - 9x^2}$$

[In] int((5\*x - 9\*x^2)^(1/2),x)

[Out] (25\*asin((18\*x)/5 - 1))/216 + (x/2 - 5/36)\*(5\*x - 9\*x^2)^(1/2)



## 3.12 $\int (x - x^2)^{3/2} dx$

Optimal result . . . . .	121
Rubi [A] (verified) . . . . .	121
Mathematica [A] (verified) . . . . .	122
Maple [A] (verified) . . . . .	122
Fricas [A] (verification not implemented) . . . . .	123
Sympy [A] (verification not implemented) . . . . .	123
Maxima [A] (verification not implemented) . . . . .	124
Giac [A] (verification not implemented) . . . . .	124
Mupad [B] (verification not implemented) . . . . .	124

### Optimal result

Integrand size = 11, antiderivative size = 51

$$\int (x - x^2)^{3/2} dx = -\frac{3}{64}(1 - 2x)\sqrt{x - x^2} - \frac{1}{8}(1 - 2x)(x - x^2)^{3/2} - \frac{3}{128}\arcsin(1 - 2x)$$

[Out] -1/8\*(1-2\*x)\*(-x^2+x)^(3/2)+3/128\*arcsin(-1+2\*x)-3/64\*(1-2\*x)\*(-x^2+x)^(1/2)

### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {626, 633, 222}

$$\int (x - x^2)^{3/2} dx = -\frac{3}{128}\arcsin(1 - 2x) - \frac{1}{8}(1 - 2x)(x - x^2)^{3/2} - \frac{3}{64}(1 - 2x)\sqrt{x - x^2}$$

[In] Int[(x - x^2)^(3/2), x]

[Out] (-3\*(1 - 2\*x)\*Sqrt[x - x^2])/64 - ((1 - 2\*x)\*(x - x^2)^(3/2))/8 - (3\*ArcSin[1 - 2\*x])/128

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*

$p + 1))$ ), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{8}(1-2x)(x-x^2)^{3/2} + \frac{3}{16} \int \sqrt{x-x^2} dx \\
 &= -\frac{3}{64}(1-2x)\sqrt{x-x^2} - \frac{1}{8}(1-2x)(x-x^2)^{3/2} + \frac{3}{128} \int \frac{1}{\sqrt{x-x^2}} dx \\
 &= -\frac{3}{64}(1-2x)\sqrt{x-x^2} - \frac{1}{8}(1-2x)(x-x^2)^{3/2} - \frac{3}{128} \text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, 1-2x\right) \\
 &= -\frac{3}{64}(1-2x)\sqrt{x-x^2} - \frac{1}{8}(1-2x)(x-x^2)^{3/2} - \frac{3}{128} \sin^{-1}(1-2x)
 \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.31

$$\int (x-x^2)^{3/2} dx = \frac{x(-3+x+26x^2-40x^3+16x^4)+6\sqrt{-1+x}\sqrt{x}\arctanh\left(\frac{\sqrt{-1+x}}{-1+\sqrt{x}}\right)}{64\sqrt{-((-1+x)x)}}$$

[In] Integrate[(x - x^2)^(3/2), x]

[Out] (x\*(-3 + x + 26\*x^2 - 40\*x^3 + 16\*x^4) + 6\*Sqrt[-1 + x]\*Sqrt[x]\*ArcTanh[Sqrt[-1 + x]/(-1 + Sqrt[x])])/(64\*Sqrt[-((-1 + x)\*x)])

### Maple [A] (verified)

Time = 2.15 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.76

method	result
risch	$\frac{(16x^3-24x^2+2x+3)(-1+x)x}{64\sqrt{-(-1+x)x}} + \frac{3\arcsin(-1+2x)}{128}$
default	$-\frac{(1-2x)(-x^2+x)^{\frac{3}{2}}}{8} + \frac{3\arcsin(-1+2x)}{128} - \frac{3(1-2x)\sqrt{-x^2+x}}{64}$
pseudoelliptic	$-\frac{3\arctan\left(\frac{\sqrt{-(-1+x)x}}{x}\right)}{64} + \frac{(-16x^3+24x^2-2x-3)\sqrt{-(-1+x)x}}{64}$
meijerg	$-\frac{3i\left(-\frac{i\sqrt{\pi}\sqrt{x}(80x^3-120x^2+10x+15)\sqrt{1-x}}{240} + \frac{i\sqrt{\pi}\arcsin(\sqrt{x})}{16}\right)}{4\sqrt{\pi}}$
trager	$\left(-\frac{1}{4}x^3 + \frac{3}{8}x^2 - \frac{1}{32}x - \frac{3}{64}\right)\sqrt{-x^2+x} - \frac{3\text{RootOf}\left(-Z^2+1\right)\ln\left(2\text{RootOf}\left(-Z^2+1\right)x+2\sqrt{-x^2+x}-\text{RootOf}\left(-Z^2+1\right)\right)}{128}$

[In] `int((-x^2+x)^(3/2),x,method=_RETURNVERBOSE)`

[Out]  $1/64*(16*x^3-24*x^2+2*x+3)*(-1+x)*x/(-(-1+x)*x)^(1/2)+3/128*\arcsin(-1+2*x)$

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (x - x^2)^{3/2} dx = -\frac{1}{64} (16x^3 - 24x^2 + 2x + 3)\sqrt{-x^2 + x} - \frac{3}{64} \arctan\left(\frac{\sqrt{-x^2 + x}}{x}\right)$$

[In] `integrate((-x^2+x)^(3/2),x, algorithm="fricas")`

[Out]  $-1/64*(16*x^3 - 24*x^2 + 2*x + 3)*\text{sqrt}(-x^2 + x) - 3/64*\arctan(\text{sqrt}(-x^2 + x)/x)$

### Sympy [A] (verification not implemented)

Time = 0.34 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.10

$$\int (x - x^2)^{3/2} dx = \sqrt{-x^2 + x} \left( \frac{x^2}{3} - \frac{x}{12} - \frac{1}{8} \right) - \sqrt{-x^2 + x} \left( \frac{x^3}{4} - \frac{x^2}{24} - \frac{5x}{96} - \frac{5}{64} \right) + \frac{3\text{asin}(2x - 1)}{128}$$

[In] `integrate((-x**2+x)**(3/2),x)`

[Out]  $\text{sqrt}(-x**2 + x)*(x**2/3 - x/12 - 1/8) - \text{sqrt}(-x**2 + x)*(x**3/4 - x**2/24 - 5*x/96 - 5/64) + 3*\text{asin}(2*x - 1)/128$

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.08

$$\int (x - x^2)^{3/2} dx = \frac{1}{4} (-x^2 + x)^{\frac{3}{2}} x - \frac{1}{8} (-x^2 + x)^{\frac{3}{2}} + \frac{3}{32} \sqrt{-x^2 + x} - \frac{3}{64} \sqrt{-x^2 + x} + \frac{3}{128} \arcsin(2x - 1)$$

[In] integrate((-x^2+x)^(3/2),x, algorithm="maxima")

[Out] 1/4\*(-x^2 + x)^(3/2)\*x - 1/8\*(-x^2 + x)^(3/2) + 3/32\*sqrt(-x^2 + x)\*x - 3/64\*sqrt(-x^2 + x) + 3/128\*arcsin(2\*x - 1)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.69

$$\int (x - x^2)^{3/2} dx = -\frac{1}{64} (2(4(2x - 3)x + 1)x + 3) \sqrt{-x^2 + x} + \frac{3}{128} \arcsin(2x - 1)$$

[In] integrate((-x^2+x)^(3/2),x, algorithm="giac")

[Out] -1/64\*(2\*(4\*(2\*x - 3)\*x + 1)\*x + 3)\*sqrt(-x^2 + x) + 3/128\*arcsin(2\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 9.12 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.76

$$\int (x - x^2)^{3/2} dx = \frac{3 \arcsin(2x - 1)}{128} + \frac{3 \sqrt{x - x^2} \left(\frac{x}{2} - \frac{1}{4}\right)}{16} + \frac{(x - x^2)^{3/2} \left(x - \frac{1}{2}\right)}{4}$$

[In] int((x - x^2)^(3/2),x)

[Out] (3\*asin(2\*x - 1))/128 + (3\*(x - x^2)^(1/2)\*(x/2 - 1/4))/16 + ((x - x^2)^(3/2)\*(x - 1/2))/4

### 3.13 $\int \sqrt{4x + x^2} dx$

Optimal result	125
Rubi [A] (verified)	125
Mathematica [A] (verified)	126
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	127
Sympy [A] (verification not implemented)	127
Maxima [A] (verification not implemented)	128
Giac [A] (verification not implemented)	128
Mupad [B] (verification not implemented)	128

#### Optimal result

Integrand size = 11, antiderivative size = 35

$$\int \sqrt{4x + x^2} dx = \frac{1}{2}(2 + x)\sqrt{4x + x^2} - 4\operatorname{arctanh}\left(\frac{x}{\sqrt{4x + x^2}}\right)$$

[Out]  $-4*\operatorname{arctanh}(x/(x^2+4*x)^{(1/2)})+1/2*(2+x)*(x^2+4*x)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {626, 634, 212}

$$\int \sqrt{4x + x^2} dx = \frac{1}{2}(x + 2)\sqrt{x^2 + 4x} - 4\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 + 4x}}\right)$$

[In] `Int[Sqrt[4*x + x^2], x]`

[Out] `((2 + x)*Sqrt[4*x + x^2])/2 - 4*ArcTanh[x/Sqrt[4*x + x^2]]`

#### Rule 212

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

#### Rule 626

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N`

`eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

#### Rule 634

`Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`

#### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}(2+x)\sqrt{4x+x^2} - 2 \int \frac{1}{\sqrt{4x+x^2}} dx \\ &= \frac{1}{2}(2+x)\sqrt{4x+x^2} - 4 \text{Subst} \left( \int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{4x+x^2}} \right) \\ &= \frac{1}{2}(2+x)\sqrt{4x+x^2} - 4 \tanh^{-1} \left( \frac{x}{\sqrt{4x+x^2}} \right) \end{aligned}$$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.31

$$\int \sqrt{4x+x^2} dx = \frac{1}{2} \sqrt{x(4+x)} \left( 2+x + \frac{8 \log(-\sqrt{x} + \sqrt{4+x})}{\sqrt{x}\sqrt{4+x}} \right)$$

[In] `Integrate[Sqrt[4*x + x^2], x]`

[Out] `(Sqrt[x*(4 + x)]*(2 + x + (8*Log[-Sqrt[x] + Sqrt[4 + x]])/(Sqrt[x]*Sqrt[4 + x])))/2`

#### Maple [A] (verified)

Time = 1.90 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
trager	$\left(\frac{x}{2} + 1\right) \sqrt{x^2 + 4x} - 2 \ln(2 + x + \sqrt{x^2 + 4x})$	32
default	$\frac{(2x+4)\sqrt{x^2+4x}}{4} - 2 \ln(2 + x + \sqrt{x^2 + 4x})$	33
risch	$\frac{(2+x)x(4+x)}{2\sqrt{x(4+x)}} - 2 \ln(2 + x + \sqrt{x^2 + 4x})$	33
meijerg	$8 \frac{\left(-\frac{\sqrt{\pi} \sqrt{x} \left(3 + \frac{3x}{2}\right) \sqrt{\frac{x}{4} + 1}}{12} + \frac{\sqrt{\pi} \operatorname{arcsinh}\left(\frac{\sqrt{x}}{2}\right)}{2}\right)}{\sqrt{\pi}}$	38
pseudoelliptic	$\frac{8x^2 \left( (2+x)\sqrt{x(4+x)} + 4 \ln\left(\frac{\sqrt{x(4+x)} - x}{x}\right) - 4 \ln\left(\frac{x + \sqrt{x(4+x)}}{x}\right) \right)}{\left(-\sqrt{x(4+x)} + x\right)^2 \left(x + \sqrt{x(4+x)}\right)^2}$	76

[In] `int((x^2+4*x)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $(1/2*x+1)*(x^2+4*x)^(1/2)-2*\ln(2+x+(x^2+4*x)^(1/2))$

### Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \sqrt{4x + x^2} dx = \frac{1}{2} \sqrt{x^2 + 4x}(x + 2) + 2 \log(-x + \sqrt{x^2 + 4x} - 2)$$

[In] `integrate((x^2+4*x)^(1/2),x, algorithm="fricas")`

[Out]  $1/2*\sqrt{x^2 + 4*x}*(x + 2) + 2*\log(-x + \sqrt{x^2 + 4*x} - 2)$

### Sympy [A] (verification not implemented)

Time = 0.21 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \sqrt{4x + x^2} dx = \left(\frac{x}{2} + 1\right) \sqrt{x^2 + 4x} - 2 \log(2x + 2\sqrt{x^2 + 4x} + 4)$$

[In] `integrate((x**2+4*x)**(1/2),x)`

[Out]  $(x/2 + 1)*\sqrt{x**2 + 4*x} - 2*\log(2*x + 2*\sqrt{x**2 + 4*x} + 4)$

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.17

$$\int \sqrt{4x + x^2} dx = \frac{1}{2} \sqrt{x^2 + 4x} + \sqrt{x^2 + 4x} - 2 \log \left( 2x + 2\sqrt{x^2 + 4x} + 4 \right)$$

[In] integrate((x^2+4\*x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(x^2 + 4\*x)\*x + sqrt(x^2 + 4\*x) - 2\*log(2\*x + 2\*sqrt(x^2 + 4\*x) + 4)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \sqrt{4x + x^2} dx = \frac{1}{2} \sqrt{x^2 + 4x}(x + 2) + 2 \log \left( \left| -x + \sqrt{x^2 + 4x} - 2 \right| \right)$$

[In] integrate((x^2+4\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(x^2 + 4\*x)\*(x + 2) + 2\*log(abs(-x + sqrt(x^2 + 4\*x) - 2))

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \sqrt{4x + x^2} dx = \sqrt{x^2 + 4x} \left( \frac{x}{2} + 1 \right) - 2 \ln \left( x + \sqrt{x(x+4)} + 2 \right)$$

[In] int((4\*x + x^2)^(1/2),x)

[Out] (4\*x + x^2)^(1/2)\*(x/2 + 1) - 2\*log(x + (x\*(x + 4))^(1/2) + 2)



### 3.14 $\int \sqrt{-8x + x^2} dx$

Optimal result	129
Rubi [A] (verified)	129
Mathematica [A] (verified)	130
Maple [A] (verified)	130
Fricas [A] (verification not implemented)	131
Sympy [A] (verification not implemented)	131
Maxima [A] (verification not implemented)	132
Giac [A] (verification not implemented)	132
Mupad [B] (verification not implemented)	132

#### Optimal result

Integrand size = 11, antiderivative size = 37

$$\int \sqrt{-8x + x^2} dx = -\frac{1}{2}(4 - x)\sqrt{-8x + x^2} - 16\operatorname{arctanh}\left(\frac{x}{\sqrt{-8x + x^2}}\right)$$

[Out]  $-16*\operatorname{arctanh}(x/(x^2-8*x)^{(1/2)})-1/2*(4-x)*(x^2-8*x)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {626, 634, 212}

$$\int \sqrt{-8x + x^2} dx = -16\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 - 8x}}\right) - \frac{1}{2}\sqrt{x^2 - 8x}(4 - x)$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[-8*x + x^2], x]$

[Out]  $-1/2*((4 - x)*\operatorname{Sqrt}[-8*x + x^2]) - 16*\operatorname{ArcTanh}[x/\operatorname{Sqrt}[-8*x + x^2]]$

#### Rule 212

$\operatorname{Int}[(a + (b \cdot x) \cdot x^{-1}), x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[-b, 2])) \cdot \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

#### Rule 626

$\operatorname{Int}[(a + (b \cdot x) + (c \cdot x)^2)^p, x\_Symbol] \rightarrow \operatorname{Simp}[(b + 2 \cdot c \cdot x) \cdot ((a + b \cdot x + c \cdot x^2)^p / (2 \cdot c \cdot (2 \cdot p + 1))), x] - \operatorname{Dist}[p \cdot ((b^2 - 4 \cdot a \cdot c) / (2 \cdot c \cdot (2 \cdot p + 1))), \operatorname{Int}[(a + b \cdot x + c \cdot x^2)^{p-1}, x], x] /;$   $\operatorname{FreeQ}\{a, b, c, x\} \ \&\& \ N$

`eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

#### Rule 634

`Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`

#### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{2}(4-x)\sqrt{-8x+x^2} - 8 \int \frac{1}{\sqrt{-8x+x^2}} dx \\ &= -\frac{1}{2}(4-x)\sqrt{-8x+x^2} - 16 \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{-8x+x^2}}\right) \\ &= -\frac{1}{2}(4-x)\sqrt{-8x+x^2} - 16 \tanh^{-1}\left(\frac{x}{\sqrt{-8x+x^2}}\right) \end{aligned}$$

#### Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.24

$$\int \sqrt{-8x+x^2} dx = \frac{1}{2} \sqrt{(-8+x)x} \left( -4+x + \frac{32 \log(\sqrt{-8+x} - \sqrt{x})}{\sqrt{-8+x}\sqrt{x}} \right)$$

[In] `Integrate[Sqrt[-8*x + x^2], x]`

[Out] `(Sqrt[(-8 + x)*x]*(-4 + x + (32*Log[Sqrt[-8 + x] - Sqrt[x]])/(Sqrt[-8 + x]*Sqrt[x]))) / 2`

#### Maple [A] (verified)

Time = 1.86 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

method	result	size
trager	$\left(\frac{x}{2} - 2\right) \sqrt{x^2 - 8x} - 8 \ln(x - 4 + \sqrt{x^2 - 8x})$	32
default	$\frac{(2x-8)\sqrt{x^2-8x}}{4} - 8 \ln(x - 4 + \sqrt{x^2 - 8x})$	33
risch	$\frac{(x-4)x(x-8)}{2\sqrt{x(x-8)}} - 8 \ln(x - 4 + \sqrt{x^2 - 8x})$	33
meijerg	$\frac{32i\sqrt{\text{signum}(x-8)} \left( -\frac{i\sqrt{\pi}\sqrt{x}\sqrt{2}\left(-\frac{3x}{4}+3\right)\sqrt{-\frac{x}{8}+1} + i\sqrt{\pi}\arcsin\left(\frac{\sqrt{2}\sqrt{x}}{4}\right)}{24} \right)}{\sqrt{\pi}\sqrt{-\text{signum}(x-8)}}$	61
pseudoelliptic	$\frac{32\left((x-4)\sqrt{x(x-8)}+16\ln\left(\frac{\sqrt{x(x-8)}-x}{x}\right)-16\ln\left(\frac{x+\sqrt{x(x-8)}}{x}\right)\right)x^2}{\left(-\sqrt{x(x-8)}+x\right)^2\left(x+\sqrt{x(x-8)}\right)^2}$	76

[In] `int((x^2-8*x)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $(1/2*x-2)*(x^2-8*x)^{(1/2)}-8*\ln(x-4+(x^2-8*x)^{(1/2)})$

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.86

$$\int \sqrt{-8x + x^2} dx = \frac{1}{2} \sqrt{x^2 - 8x}(x - 4) + 8 \log(-x + \sqrt{x^2 - 8x} + 4)$$

[In] `integrate((x^2-8*x)^(1/2),x, algorithm="fricas")`

[Out]  $1/2*\text{sqrt}(x^2 - 8*x)*(x - 4) + 8*\log(-x + \text{sqrt}(x^2 - 8*x) + 4)$

### Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int \sqrt{-8x + x^2} dx = \left(\frac{x}{2} - 2\right) \sqrt{x^2 - 8x} - 8 \log(2x + 2\sqrt{x^2 - 8x} - 8)$$

[In] `integrate((x**2-8*x)**(1/2),x)`

[Out]  $(x/2 - 2)*\text{sqrt}(x**2 - 8*x) - 8*\log(2*x + 2*\text{sqrt}(x**2 - 8*x) - 8)$

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.16

$$\int \sqrt{-8x + x^2} dx = \frac{1}{2} \sqrt{x^2 - 8x} x - 2 \sqrt{x^2 - 8x} - 8 \log(2x + 2\sqrt{x^2 - 8x} - 8)$$

[In] integrate((x^2-8\*x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(x^2 - 8\*x)\*x - 2\*sqrt(x^2 - 8\*x) - 8\*log(2\*x + 2\*sqrt(x^2 - 8\*x) - 8)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \sqrt{-8x + x^2} dx = \frac{1}{2} \sqrt{x^2 - 8x}(x - 4) + 8 \log(|-x + \sqrt{x^2 - 8x} + 4|)$$

[In] integrate((x^2-8\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(x^2 - 8\*x)\*(x - 4) + 8\*log(abs(-x + sqrt(x^2 - 8\*x) + 4))

**Mupad [B] (verification not implemented)**

Time = 9.14 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.78

$$\int \sqrt{-8x + x^2} dx = \left(\frac{x}{2} - 2\right) \sqrt{x^2 - 8x} - 8 \ln\left(x + \sqrt{x(x-8)} - 4\right)$$

[In] int((x^2 - 8\*x)^(1/2),x)

[Out] (x/2 - 2)\*(x^2 - 8\*x)^(1/2) - 8\*log(x + (x\*(x - 8))^(1/2) - 4)

### 3.15 $\int \sqrt{-x + x^2} dx$

Optimal result	133
Rubi [A] (verified)	133
Mathematica [A] (verified)	134
Maple [A] (verified)	134
Fricas [A] (verification not implemented)	135
Sympy [A] (verification not implemented)	135
Maxima [A] (verification not implemented)	136
Giac [A] (verification not implemented)	136
Mupad [B] (verification not implemented)	136

#### Optimal result

Integrand size = 11, antiderivative size = 39

$$\int \sqrt{-x + x^2} dx = -\frac{1}{4}(1 - 2x)\sqrt{-x + x^2} - \frac{1}{4}\operatorname{arctanh}\left(\frac{x}{\sqrt{-x + x^2}}\right)$$

[Out]  $-1/4*\operatorname{arctanh}(x/(x^2-x)^{(1/2)})-1/4*(1-2*x)*(x^2-x)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {626, 634, 212}

$$\int \sqrt{-x + x^2} dx = -\frac{1}{4}\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2 - x}}\right) - \frac{1}{4}\sqrt{x^2 - x}(1 - 2x)$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[-x + x^2], x]$

[Out]  $-1/4*((1 - 2*x)*\operatorname{Sqrt}[-x + x^2]) - \operatorname{ArcTanh}[x/\operatorname{Sqrt}[-x + x^2]]/4$

#### Rule 212

$\operatorname{Int}[(a + (b \cdot x + c \cdot x^2)^{-1}), x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \cdot \operatorname{Rt}[-b, 2])) * \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2] \cdot (x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

#### Rule 626

$\operatorname{Int}[(a + (b \cdot x + c \cdot x^2)^p), x\_Symbol] \rightarrow \operatorname{Simp}[(b + 2 \cdot c \cdot x) * ((a + b \cdot x + c \cdot x^2)^p / (2 \cdot c \cdot (2 \cdot p + 1))), x] - \operatorname{Dist}[p * ((b^2 - 4 \cdot a \cdot c) / (2 \cdot c \cdot (2 \cdot p + 1))), \operatorname{Int}[(a + b \cdot x + c \cdot x^2)^{p-1}, x], x] /;$   $\operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{N}$

`eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

#### Rule 634

`Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`

#### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{4}(1-2x)\sqrt{-x+x^2} - \frac{1}{8} \int \frac{1}{\sqrt{-x+x^2}} dx \\ &= -\frac{1}{4}(1-2x)\sqrt{-x+x^2} - \frac{1}{4} \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{-x+x^2}}\right) \\ &= -\frac{1}{4}(1-2x)\sqrt{-x+x^2} - \frac{1}{4} \tanh^{-1}\left(\frac{x}{\sqrt{-x+x^2}}\right) \end{aligned}$$

#### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28

$$\int \sqrt{-x+x^2} dx = \frac{1}{4} \sqrt{(-1+x)x} \left( -1+2x - \frac{2 \arctanh\left(\frac{\sqrt{-1+x}}{-1+\sqrt{x}}\right)}{\sqrt{-1+x}\sqrt{x}} \right)$$

[In] `Integrate[Sqrt[-x + x^2], x]`

[Out] `(Sqrt[(-1 + x)*x]*(-1 + 2*x - (2*ArcTanh[Sqrt[-1 + x]/(-1 + Sqrt[x])]))/(Sqrt[-1 + x]*Sqrt[x]))/4`

#### Maple [A] (verified)

Time = 1.86 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{(-1+2x)\sqrt{x^2-x}}{4} - \frac{\ln\left(x-\frac{1}{2}+\sqrt{x^2-x}\right)}{8}$	33
risch	$\frac{(-1+2x)(-1+x)x}{4\sqrt{(-1+x)x}} - \frac{\ln\left(x-\frac{1}{2}+\sqrt{x^2-x}\right)}{8}$	35
trager	$\left(-\frac{1}{4} + \frac{x}{2}\right) \sqrt{x^2-x} - \frac{\ln\left(2x-1+2\sqrt{x^2-x}\right)}{8}$	36
meijerg	$-\frac{i\sqrt{\text{signum}(-1+x)}\left(-\frac{i\sqrt{\pi}\sqrt{x}(3-6x)\sqrt{1-x}}{6} + \frac{i\sqrt{\pi}\arcsin(\sqrt{x})}{2}\right)}{2\sqrt{\pi}\sqrt{-\text{signum}(-1+x)}}$	53
pseudoelliptic	$\frac{x^2\left(4\sqrt{(-1+x)x}x + \ln\left(\frac{\sqrt{(-1+x)x-x}}{x}\right) - \ln\left(\frac{x+\sqrt{(-1+x)x}}{x}\right) - 2\sqrt{(-1+x)x}\right)}{8\left(\sqrt{(-1+x)x-x}\right)^2\left(x+\sqrt{(-1+x)x}\right)^2}$	82

[In] `int((x^2-x)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/4*(-1+2*x)*(x^2-x)^(1/2)-1/8*ln(x-1/2+(x^2-x)^(1/2))`

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int \sqrt{-x+x^2} dx = \frac{1}{4} \sqrt{x^2-x}(2x-1) + \frac{1}{8} \log\left(-2x+2\sqrt{x^2-x}+1\right)$$

[In] `integrate((x^2-x)^(1/2),x, algorithm="fricas")`

[Out] `1/4*sqrt(x^2-x)*(2*x-1)+1/8*log(-2*x+2*sqrt(x^2-x)+1)`

### Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

$$\int \sqrt{-x+x^2} dx = \left(\frac{x}{2} - \frac{1}{4}\right) \sqrt{x^2-x} - \frac{\log\left(2x+2\sqrt{x^2-x}-1\right)}{8}$$

[In] `integrate((x**2-x)**(1/2),x)`

[Out] `(x/2-1/4)*sqrt(x**2-x)-log(2*x+2*sqrt(x**2-x)-1)/8`

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \sqrt{-x + x^2} dx = \frac{1}{2} \sqrt{x^2 - x} x - \frac{1}{4} \sqrt{x^2 - x} - \frac{1}{8} \log(2x + 2\sqrt{x^2 - x} - 1)$$

[In] integrate((x^2-x)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(x^2 - x)\*x - 1/4\*sqrt(x^2 - x) - 1/8\*log(2\*x + 2\*sqrt(x^2 - x) - 1)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.95

$$\int \sqrt{-x + x^2} dx = \frac{1}{4} \sqrt{x^2 - x}(2x - 1) + \frac{1}{8} \log(|-2x + 2\sqrt{x^2 - x} + 1|)$$

[In] integrate((x^2-x)^(1/2),x, algorithm="giac")

[Out] 1/4\*sqrt(x^2 - x)\*(2\*x - 1) + 1/8\*log(abs(-2\*x + 2\*sqrt(x^2 - x) + 1))

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int \sqrt{-x + x^2} dx = \sqrt{x^2 - x} \left( \frac{x}{2} - \frac{1}{4} \right) - \frac{\ln \left( x + \sqrt{x(x-1)} - \frac{1}{2} \right)}{8}$$

[In] int((x^2 - x)^(1/2),x)

[Out] (x^2 - x)^(1/2)\*(x/2 - 1/4) - log(x + (x\*(x - 1))^(1/2) - 1/2)/8



### 3.16 $\int \frac{1}{(bx+cx^2)^{7/2}} dx$

Optimal result	137
Rubi [A] (verified)	137
Mathematica [A] (verified)	138
Maple [A] (verified)	138
Fricas [A] (verification not implemented)	139
Sympy [F]	139
Maxima [A] (verification not implemented)	140
Giac [A] (verification not implemented)	140
Mupad [B] (verification not implemented)	140

#### Optimal result

Integrand size = 13, antiderivative size = 83

$$\int \frac{1}{(bx+cx^2)^{7/2}} dx = -\frac{2(b+2cx)}{5b^2(bx+cx^2)^{5/2}} + \frac{32c(b+2cx)}{15b^4(bx+cx^2)^{3/2}} - \frac{256c^2(b+2cx)}{15b^6\sqrt{bx+cx^2}}$$

[Out]  $-2/5*(2*c*x+b)/b^2/(c*x^2+b*x)^(5/2)+32/15*c*(2*c*x+b)/b^4/(c*x^2+b*x)^(3/2)$   
 $-256/15*c^2*(2*c*x+b)/b^6/(c*x^2+b*x)^(1/2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {628, 627}

$$\int \frac{1}{(bx+cx^2)^{7/2}} dx = -\frac{256c^2(b+2cx)}{15b^6\sqrt{bx+cx^2}} + \frac{32c(b+2cx)}{15b^4(bx+cx^2)^{3/2}} - \frac{2(b+2cx)}{5b^2(bx+cx^2)^{5/2}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{-7/2}, x]$

[Out]  $(-2*(b + 2*c*x))/(5*b^2*(b*x + c*x^2)^(5/2)) + (32*c*(b + 2*c*x))/(15*b^4*(b*x + c*x^2)^(3/2)) - (256*c^2*(b + 2*c*x))/(15*b^6*\text{Sqrt}[b*x + c*x^2])$

Rule 627

$\text{Int}[(a_. + (b_.)*(x_.) + (c_.)*(x_.)^2)^{-3/2}, x\_Symbol] \text{ :> Simp}[-2*((b + 2*c*x)/((b^2 - 4*a*c)*\text{Sqrt}[a + b*x + c*x^2])), x] \text{ /; FreeQ}\{a, b, c\}, x] \&\& \text{ NeQ}[b^2 - 4*a*c, 0]$

Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{2(b + 2cx)}{5b^2 (bx + cx^2)^{5/2}} - \frac{(16c) \int \frac{1}{(bx+cx^2)^{5/2}} dx}{5b^2} \\ &= -\frac{2(b + 2cx)}{5b^2 (bx + cx^2)^{5/2}} + \frac{32c(b + 2cx)}{15b^4 (bx + cx^2)^{3/2}} + \frac{(128c^2) \int \frac{1}{(bx+cx^2)^{3/2}} dx}{15b^4} \\ &= -\frac{2(b + 2cx)}{5b^2 (bx + cx^2)^{5/2}} + \frac{32c(b + 2cx)}{15b^4 (bx + cx^2)^{3/2}} - \frac{256c^2(b + 2cx)}{15b^6 \sqrt{bx + cx^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.20 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.84

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = -\frac{2(3b^5 - 10b^4cx + 80b^3c^2x^2 + 480b^2c^3x^3 + 640bc^4x^4 + 256c^5x^5)}{15b^6(x(b + cx))^{5/2}}$$

[In] Integrate[(b\*x + c\*x^2)^(-7/2), x]

[Out] (-2\*(3\*b^5 - 10\*b^4\*c\*x + 80\*b^3\*c^2\*x^2 + 480\*b^2\*c^3\*x^3 + 640\*b\*c^4\*x^4 + 256\*c^5\*x^5))/(15\*b^6\*(x\*(b + c\*x))^(5/2))

**Maple [A] (verified)**

Time = 1.92 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.90

method	result	size
gospser	$\frac{2x(cx+b)(256c^5x^5+640bx^4c^4+480b^2c^3x^3+80x^2b^3c^2-10cxb^4+3b^5)}{15b^6(cx^2+bx)^{\frac{7}{2}}}$	75
default	$\frac{2(2cx+b)}{5b^2(cx^2+bx)^{\frac{5}{2}}} - \frac{16c\left(-\frac{2(2cx+b)}{3b^2(cx^2+bx)^{\frac{3}{2}}} + \frac{16c(2cx+b)}{3b^4\sqrt{cx^2+bx}}\right)}{5b^2}$	76
pseudoelliptic	$\frac{-\frac{512}{15}c^5x^5 - \frac{256}{3}bx^4c^4 - 64b^2c^3x^3 - \frac{32}{3}x^2b^3c^2 + \frac{4}{3}cxb^4 - \frac{2}{3}b^5}{x^2(cx+b)^2\sqrt{x(cx+b)}b^6}$	77
trager	$\frac{2(256c^5x^5+640bx^4c^4+480b^2c^3x^3+80x^2b^3c^2-10cxb^4+3b^5)\sqrt{cx^2+bx}}{15b^6(cx+b)^3x^3}$	79
risch	$\frac{2(cx+b)(128c^2x^2-19bcx+3b^2)}{15b^6x^2\sqrt{x(cx+b)}} - \frac{2c^3(128c^2x^2+275bcx+150b^2)x}{15\sqrt{x(cx+b)}(c^2x^2+2bcx+b^2)b^6}$	98

[In] `int(1/(c*x^2+b*x)^(7/2),x,method=_RETURNVERBOSE)`

[Out] 
$$-2/15*x*(c*x+b)*(256*c^5*x^5+640*b*c^4*x^4+480*b^2*c^3*x^3+80*b^3*c^2*x^2-10*b^4*c*x+3*b^5)/b^6/(c*x^2+b*x)^(7/2)$$

## Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.27

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = \frac{2(256c^5x^5 + 640bc^4x^4 + 480b^2c^3x^3 + 80b^3c^2x^2 - 10b^4cx + 3b^5)\sqrt{cx^2 + bx}}{15(b^6c^3x^6 + 3b^7c^2x^5 + 3b^8cx^4 + b^9x^3)}$$

[In] `integrate(1/(c*x^2+b*x)^(7/2),x, algorithm="fricas")`

[Out] 
$$-2/15*(256*c^5*x^5 + 640*b*c^4*x^4 + 480*b^2*c^3*x^3 + 80*b^3*c^2*x^2 - 10*b^4*c*x + 3*b^5)*\sqrt{c*x^2 + b*x}/(b^6*c^3*x^6 + 3*b^7*c^2*x^5 + 3*b^8*c*x^4 + b^9*x^3)$$

## Sympy [F]

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = \int \frac{1}{(bx + cx^2)^{\frac{7}{2}}} dx$$

[In] `integrate(1/(c*x**2+b*x)**(7/2),x)`

[Out] `Integral((b*x + c*x**2)**(-7/2), x)`

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.34

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = -\frac{4cx}{5(cx^2 + bx)^{5/2}b^2} + \frac{64c^2x}{15(cx^2 + bx)^{3/2}b^4} - \frac{512c^3x}{15\sqrt{cx^2 + bx}b^6} - \frac{2}{5(cx^2 + bx)^{5/2}b} + \frac{32c}{15(cx^2 + bx)^{3/2}b^3} - \frac{256c^2}{15\sqrt{cx^2 + bx}b^5}$$

[In] integrate(1/(c\*x^2+b\*x)^(7/2),x, algorithm="maxima")

[Out]  $-4/5*c*x/((c*x^2 + b*x)^{(5/2)*b^2}) + 64/15*c^2*x/((c*x^2 + b*x)^{(3/2)*b^4}) - 512/15*c^3*x/(sqrt(c*x^2 + b*x)*b^6) - 2/5/((c*x^2 + b*x)^{(5/2)*b}) + 32/15*c/((c*x^2 + b*x)^{(3/2)*b^3}) - 256/15*c^2/(sqrt(c*x^2 + b*x)*b^5)$

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.89

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = -\frac{2 \left( 2 \left( 8 \left( 2 \left( 4x \left( \frac{2c^5x}{b^6} + \frac{5c^4}{b^5} \right) + \frac{15c^3}{b^4} \right) x + \frac{5c^2}{b^3} \right) x - \frac{5c}{b^2} \right) x + \frac{3}{b} \right)}{15 (cx^2 + bx)^{5/2}}$$

[In] integrate(1/(c\*x^2+b\*x)^(7/2),x, algorithm="giac")

[Out]  $-2/15*(2*(8*(2*(4*x*(2*c^5*x/b^6 + 5*c^4/b^5) + 15*c^3/b^4)*x + 5*c^2/b^3)*x - 5*c/b^2)*x + 3/b)/(c*x^2 + b*x)^{(5/2)}$

**Mupad [B] (verification not implemented)**

Time = 9.22 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.16

$$\int \frac{1}{(bx + cx^2)^{7/2}} dx = \frac{6b^5 + 256bc^2(cx^2 + bx)^2 + 512c^3x(cx^2 + bx)^2 - 32b^3c(cx^2 + bx) + 12b^4cx - 64b^2c^2x(cx^2 + bx)}{15b^6(cx^2 + bx)^{5/2}}$$

[In] int(1/(b\*x + c\*x^2)^(7/2),x)

[Out]  $-(6*b^5 + 256*b*c^2*(b*x + c*x^2)^2 + 512*c^3*x*(b*x + c*x^2)^2 - 32*b^3*c*(b*x + c*x^2) + 12*b^4*c*x - 64*b^2*c^2*x*(b*x + c*x^2))/(15*b^6*(b*x + c*x^2)^{(5/2)})$

### 3.17 $\int \frac{1}{\sqrt{3ix+4x^2}} dx$

Optimal result	141
Rubi [A] (verified)	141
Mathematica [B] (verified)	142
Maple [A] (verified)	142
Fricas [B] (verification not implemented)	143
Sympy [A] (verification not implemented)	143
Maxima [B] (verification not implemented)	143
Giac [B] (verification not implemented)	144
Mupad [B] (verification not implemented)	144

#### Optimal result

Integrand size = 15, antiderivative size = 16

$$\int \frac{1}{\sqrt{3ix+4x^2}} dx = \frac{1}{2}i \arcsin\left(1 - \frac{8ix}{3}\right)$$

[Out]  $-1/2*I*\arcsin(-1+8/3*I*x)$

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {633, 221}

$$\int \frac{1}{\sqrt{3ix+4x^2}} dx = \frac{1}{2}i \arcsin\left(1 - \frac{8ix}{3}\right)$$

[In] `Int[1/Sqrt[(3*I)*x + 4*x^2],x]`

[Out] `(I/2)*ArcSin[1 - ((8*I)/3)*x]`

#### Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

#### Rule 633

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{6} \text{Subst} \left( \int \frac{1}{\sqrt{1 + \frac{x^2}{9}}} dx, x, 3i + 8x \right) \\ &= \frac{1}{2} i \sin^{-1} \left( 1 - \frac{8ix}{3} \right) \end{aligned}$$

**Mathematica [B] (verified)**

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 51 vs.  $2(16) = 32$ .

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 3.19

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = -\frac{\sqrt{x}\sqrt{3i + 4x} \log(-2\sqrt{x} + \sqrt{3i + 4x})}{\sqrt{x(3i + 4x)}}$$

[In] Integrate[1/Sqrt[(3\*I)\*x + 4\*x^2],x]

[Out] -((Sqrt[x]\*Sqrt[3\*I + 4\*x]\*Log[-2\*Sqrt[x] + Sqrt[3\*I + 4\*x]])/Sqrt[x\*(3\*I + 4\*x)])

**Maple [A] (verified)**

Time = 1.86 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.62

method	result	size
default	$\frac{\text{arcsinh}\left(i + \frac{8x}{3}\right)}{2}$	10
trager	$\frac{\ln\left(440x + 144 + 165i - 192i\sqrt{4x^2 + 3ix} - 384ix + 220\sqrt{4x^2 + 3ix}\right)}{2}$	44
pseudoelliptic	$\frac{\ln\left(\frac{\sqrt{x(3i+4x)+2x}}{x}\right)}{2} - \frac{\ln\left(\frac{-2x + \sqrt{x(3i+4x)}}{x}\right)}{2}$	44

[In] int(1/(3\*I\*x+4\*x^2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*arcsinh(I+8/3\*x)

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 19 vs.  $2(8) = 16$ .

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = -\frac{1}{2} \log \left( -2x + \sqrt{4x^2 + 3ix} - \frac{3}{4}i \right)$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(1/2),x, algorithm="fricas")

[Out] -1/2\*log(-2\*x + sqrt(4\*x^2 + 3\*I\*x) - 3/4\*I)

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = \frac{\operatorname{asinh}\left(\frac{8x}{3} + i\right)}{2}$$

[In] integrate(1/(3\*I\*x+4\*x\*\*2)\*\*(1/2),x)

[Out] asinh(8\*x/3 + I)/2

**Maxima [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 21 vs.  $2(8) = 16$ .

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.31

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = \frac{1}{2} \log \left( 8x + 4\sqrt{4x^2 + 3ix} + 3i \right)$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*log(8\*x + 4\*sqrt(4\*x^2 + 3\*I\*x) + 3\*I)

**Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 110 vs.  $2(8) = 16$ .

Time = 0.27 (sec) , antiderivative size = 110, normalized size of antiderivative = 6.88

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = \frac{1}{32} \sqrt{8x^2 + 2\sqrt{16x^2 + 9x}(8x + 3i)} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - \frac{9}{64} \log \left( 2\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}} \left( \frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1 \right) - 8x - 3i \right)$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(1/2),x, algorithm="giac")

[Out] 1/32\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(8\*x + 3\*I)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 9/64\*log(2\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1) - 8\*x - 3\*I)

**Mupad [B] (verification not implemented)**

Time = 9.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.19

$$\int \frac{1}{\sqrt{3ix + 4x^2}} dx = \frac{\ln \left( x + \frac{\sqrt{x(4x+3i)}}{2} + \frac{3i}{8} \right)}{2}$$

[In] int(1/(x\*3i + 4\*x^2)^(1/2),x)

[Out] log(x + (x\*(4\*x + 3i))^(1/2)/2 + 3i/8)/2



$$3.18 \quad \int \frac{1}{(3ix+4x^2)^{3/2}} dx$$

Optimal result . . . . .	145
Rubi [A] (verified) . . . . .	145
Mathematica [A] (verified) . . . . .	146
Maple [A] (verified) . . . . .	146
Fricas [B] (verification not implemented) . . . . .	146
Sympy [F] . . . . .	147
Maxima [A] (verification not implemented) . . . . .	147
Giac [B] (verification not implemented) . . . . .	147
Mupad [B] (verification not implemented) . . . . .	148

### Optimal result

Integrand size = 15, antiderivative size = 26

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{2(3i + 8x)}{9\sqrt{3ix + 4x^2}}$$

[Out] 2/9\*(3\*I+8\*x)/(3\*I\*x+4\*x^2)^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {627}

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{2(8x + 3i)}{9\sqrt{4x^2 + 3ix}}$$

[In] Int[((3\*I)\*x + 4\*x^2)^(-3/2), x]

[Out] (2\*(3\*I + 8\*x))/(9\*Sqrt[(3\*I)\*x + 4\*x^2])

#### Rule 627

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-3/2), x\_Symbol] :> Simp[-2\*((b + 2\*c\*x)/((b^2 - 4\*a\*c)\*Sqrt[a + b\*x + c\*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

#### Rubi steps

$$\text{integral} = \frac{2(3i + 8x)}{9\sqrt{3ix + 4x^2}}$$

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{2(3i + 8x)}{9\sqrt{x(3i + 4x)}}$$

[In] Integrate[((3\*I)\*x + 4\*x^2)^(-3/2),x]

[Out] (2\*(3\*I + 8\*x))/(9\*Sqrt[x\*(3\*I + 4\*x)])

**Maple [A] (verified)**

Time = 1.80 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.73

method	result	size
risch	$\frac{\frac{2i}{3} + \frac{16x}{9}}{\sqrt{x(3i+4x)}}$	19
pseudoelliptic	$\frac{\frac{2i}{3} + \frac{16x}{9}}{\sqrt{x(3i+4x)}}$	19
default	$\frac{\frac{2i}{3} + \frac{16x}{9}}{\sqrt{4x^2+3ix}}$	21
gosper	$\frac{2x(3i+4x)(3i+8x)}{9(4x^2+3ix)^{\frac{3}{2}}}$	28
trager	$\frac{(-\frac{14}{225} + \frac{16i}{75})(24ix+32x+12i-9)\sqrt{4x^2+3ix}}{x(12ix-16x-12i-9)}$	44

[In] int(1/(3\*I\*x+4\*x^2)^(3/2),x,method=\_RETURNVERBOSE)

[Out] 2/9\*(3\*I+8\*x)/(x\*(3\*I+4\*x))^(1/2)

**Fricas [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 39 vs.  $2(18) = 36$ .

Time = 0.27 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.50

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{2(16x^2 + \sqrt{4x^2 + 3ix}(8x + 3i) + 12ix)}{9(4x^2 + 3ix)}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(3/2),x, algorithm="fricas")

[Out] 2/9\*(16\*x^2 + sqrt(4\*x^2 + 3\*I\*x)\*(8\*x + 3\*I) + 12\*I\*x)/(4\*x^2 + 3\*I\*x)

**Sympy [F]**

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \int \frac{1}{(4x^2 + 3ix)^{\frac{3}{2}}} dx$$

[In] integrate(1/(3\*I\*x+4\*x\*\*2)\*\*(3/2),x)

[Out] Integral((4\*x\*\*2 + 3\*I\*x)\*\*(-3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{16x}{9\sqrt{4x^2 + 3ix}} + \frac{2i}{3\sqrt{4x^2 + 3ix}}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(3/2),x, algorithm="maxima")

[Out] 16/9\*x/sqrt(4\*x^2 + 3\*I\*x) + 2/3\*I/sqrt(4\*x^2 + 3\*I\*x)

**Giac [B] (verification not implemented)**

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 64 vs. 2(18) = 36.

Time = 0.29 (sec) , antiderivative size = 64, normalized size of antiderivative = 2.46

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{\sqrt{8x^2 + 2\sqrt{16x^2 + 9x}}(8x + 3i)\left(\frac{3ix}{4x^2 + \sqrt{16x^2 + 9x}} + 1\right)}{9(4x^2 + 3ix)}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(3/2),x, algorithm="giac")

[Out] 1/9\*sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(8\*x + 3\*I)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1)/(4\*x^2 + 3\*I\*x)

**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1}{(3ix + 4x^2)^{3/2}} dx = \frac{16x + 6i}{9\sqrt{4x^2 + x3i}}$$

[In] int(1/(x\*3i + 4\*x^2)^(3/2),x)

[Out] (16\*x + 6i)/(9\*(x\*3i + 4\*x^2)^(1/2))

### 3.19 $\int \frac{1}{(3ix+4x^2)^{5/2}} dx$

Optimal result . . . . .	149
Rubi [A] (verified) . . . . .	149
Mathematica [A] (verified) . . . . .	150
Maple [A] (verified) . . . . .	150
Fricas [A] (verification not implemented) . . . . .	151
Sympy [F] . . . . .	151
Maxima [A] (verification not implemented) . . . . .	151
Giac [A] (verification not implemented) . . . . .	152
Mupad [B] (verification not implemented) . . . . .	152

#### Optimal result

Integrand size = 15, antiderivative size = 53

$$\int \frac{1}{(3ix+4x^2)^{5/2}} dx = \frac{2(3i+8x)}{27(3ix+4x^2)^{3/2}} + \frac{64(3i+8x)}{243\sqrt{3ix+4x^2}}$$

[Out]  $2/27*(3*I+8*x)/(3*I*x+4*x^2)^{(3/2)}+64/243*(3*I+8*x)/(3*I*x+4*x^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {628, 627}

$$\int \frac{1}{(3ix+4x^2)^{5/2}} dx = \frac{64(8x+3i)}{243\sqrt{4x^2+3ix}} + \frac{2(8x+3i)}{27(4x^2+3ix)^{3/2}}$$

[In]  $\text{Int}[(3*I)*x + 4*x^2]^{-5/2}, x]$

[Out]  $(2*(3*I + 8*x))/(27*((3*I)*x + 4*x^2)^{(3/2)}) + (64*(3*I + 8*x))/(243*\text{Sqrt}[(3*I)*x + 4*x^2])$

#### Rule 627

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{-3/2}, x\_Symbol] \rightarrow \text{Simp}[-2*((b + 2*c*x)/((b^2 - 4*a*c)*\text{Sqrt}[a + b*x + c*x^2])), x] /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

#### Rule 628

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{p_.}, x\_Symbol] \rightarrow \text{Simp}[(b + 2*c*x)*((a + b*x + c*x^2)^{(p+1})/((p+1)*(b^2 - 4*a*c))), x] - \text{Dist}[2*c*((2*p +$

3)/((p + 1)\*(b^2 - 4\*a\*c)), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; Free Q[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2(3i + 8x)}{27(3ix + 4x^2)^{3/2}} + \frac{32}{27} \int \frac{1}{(3ix + 4x^2)^{3/2}} dx \\ &= \frac{2(3i + 8x)}{27(3ix + 4x^2)^{3/2}} + \frac{64(3i + 8x)}{243\sqrt{3ix + 4x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.68

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \frac{54i - 432x + 2304ix^2 + 2048x^3}{243(x(3i + 4x))^{3/2}}$$

[In] Integrate[((3\*I)\*x + 4\*x^2)^(-5/2),x]

[Out] (54\*I - 432\*x + (2304\*I)\*x^2 + 2048\*x^3)/(243\*(x\*(3\*I + 4\*x))^(3/2))

**Maple [A] (verified)**

Time = 1.85 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.74

method	result	size
gospers	$\frac{2x(3i+4x)(1024x^3+1152ix^2-216x+27i)}{243(4x^2+3ix)^{5/2}}$	39
risch	$\frac{\frac{2048}{243}x^3 + \frac{256}{27}ix^2 - \frac{16}{9}x + \frac{2}{9}i}{x(3i+4x)\sqrt{x(3i+4x)}}$	41
pseudoelliptic	$\frac{2048x^3+2304ix^2-432x+54i}{243x(3i+4x)\sqrt{x(3i+4x)}}$	41
default	$\frac{\frac{2i}{9} + \frac{16x}{27}}{(4x^2+3ix)^{3/2}} + \frac{\frac{64i}{81} + \frac{512x}{243}}{\sqrt{4x^2+3ix}}$	42
trager	$\frac{(\frac{88}{151875} + \frac{26i}{16875})(-76800ix^3 - 102400x^3 - 115200ix^2 + 86400x^2 + 16200ix + 21600x - 2700i + 2025)\sqrt{4x^2+3ix}}{(12ix-16x-12i-9)^2x^2}$	66

[In] int(1/(3\*I\*x+4\*x^2)^(5/2),x,method=\_RETURNVERBOSE)

[Out] 2/243\*x\*(3\*I+4\*x)\*(1152\*I\*x^2+1024\*x^3+27\*I-216\*x)/(3\*I\*x+4\*x^2)^(5/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.19

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \frac{2(2048x^4 + 3072ix^3 - 1152x^2 + (1024x^3 + 1152ix^2 - 216x + 27i)\sqrt{4x^2 + 3ix})}{243(16x^4 + 24ix^3 - 9x^2)}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(5/2),x, algorithm="fricas")

[Out] 2/243\*(2048\*x^4 + 3072\*I\*x^3 - 1152\*x^2 + (1024\*x^3 + 1152\*I\*x^2 - 216\*x + 27\*I)\*sqrt(4\*x^2 + 3\*I\*x))/(16\*x^4 + 24\*I\*x^3 - 9\*x^2)

**Sympy [F]**

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \int \frac{1}{(4x^2 + 3ix)^{5/2}} dx$$

[In] integrate(1/(3\*I\*x+4\*x\*\*2)\*\*(5/2),x)

[Out] Integral((4\*x\*\*2 + 3\*I\*x)\*\*(-5/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.04

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \frac{512x}{243\sqrt{4x^2 + 3ix}} + \frac{64i}{81\sqrt{4x^2 + 3ix}} + \frac{16x}{27(4x^2 + 3ix)^{3/2}} + \frac{2i}{9(4x^2 + 3ix)^{3/2}}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(5/2),x, algorithm="maxima")

[Out] 512/243\*x/sqrt(4\*x^2 + 3\*I\*x) + 64/81\*I/sqrt(4\*x^2 + 3\*I\*x) + 16/27\*x/(4\*x^2 + 3\*I\*x)^(3/2) + 2/9\*I/(4\*x^2 + 3\*I\*x)^(3/2)

**Giac [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.40

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \frac{(8(16(8x + 9i)x - 27)x + 27i)\sqrt{8x^2 + 2}\sqrt{16x^2 + 9x}\left(\frac{3ix}{4x^2 + \sqrt{16x^4 + 9x^2}} + 1\right)}{243(4x^2 + 3ix)^2}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(5/2),x, algorithm="giac")

[Out] 1/243\*(8\*(16\*(8\*x + 9\*I)\*x - 27)\*x + 27\*I)\*sqrt(8\*x^2 + 2)\*sqrt(16\*x^2 + 9)\*x\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) + 1)/(4\*x^2 + 3\*I\*x)^2

**Mupad [B] (verification not implemented)**

Time = 9.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.58

$$\int \frac{1}{(3ix + 4x^2)^{5/2}} dx = \frac{(16x + 6i)(128x^2 + x96i + 9)}{243(4x^2 + x3i)^{3/2}}$$

[In] int(1/(x\*3i + 4\*x^2)^(5/2),x)

[Out] ((16\*x + 6i)\*(x\*96i + 128\*x^2 + 9))/(243\*(x\*3i + 4\*x^2)^(3/2))



### 3.20 $\int \frac{1}{(3ix+4x^2)^{7/2}} dx$

Optimal result	153
Rubi [A] (verified)	153
Mathematica [A] (verified)	154
Maple [A] (verified)	154
Fricas [A] (verification not implemented)	155
Sympy [F]	155
Maxima [A] (verification not implemented)	155
Giac [A] (verification not implemented)	156
Mupad [B] (verification not implemented)	156

#### Optimal result

Integrand size = 15, antiderivative size = 79

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{2(3i + 8x)}{45(3ix + 4x^2)^{5/2}} + \frac{128(3i + 8x)}{1215(3ix + 4x^2)^{3/2}} + \frac{4096(3i + 8x)}{10935\sqrt{3ix + 4x^2}}$$

[Out] 2/45\*(3\*I+8\*x)/(3\*I\*x+4\*x^2)^(5/2)+128/1215\*(3\*I+8\*x)/(3\*I\*x+4\*x^2)^(3/2)+4096/10935\*(3\*I+8\*x)/(3\*I\*x+4\*x^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {628, 627}

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{4096(8x + 3i)}{10935\sqrt{4x^2 + 3ix}} + \frac{128(8x + 3i)}{1215(4x^2 + 3ix)^{3/2}} + \frac{2(8x + 3i)}{45(4x^2 + 3ix)^{5/2}}$$

[In] Int[((3\*I)\*x + 4\*x^2)^(-7/2), x]

[Out] (2\*(3\*I + 8\*x))/(45\*((3\*I)\*x + 4\*x^2)^(5/2)) + (128\*(3\*I + 8\*x))/(1215\*((3\*I)\*x + 4\*x^2)^(3/2)) + (4096\*(3\*I + 8\*x))/(10935\*Sqrt[(3\*I)\*x + 4\*x^2])

#### Rule 627

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-3/2), x\_Symbol] :> Simp[-2\*((b + 2\*c\*x)/((b^2 - 4\*a\*c)\*Sqrt[a + b\*x + c\*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

#### Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2(3i + 8x)}{45(3ix + 4x^2)^{5/2}} + \frac{64}{45} \int \frac{1}{(3ix + 4x^2)^{5/2}} dx \\ &= \frac{2(3i + 8x)}{45(3ix + 4x^2)^{5/2}} + \frac{128(3i + 8x)}{1215(3ix + 4x^2)^{3/2}} + \frac{2048 \int \frac{1}{(3ix+4x^2)^{3/2}} dx}{1215} \\ &= \frac{2(3i + 8x)}{45(3ix + 4x^2)^{5/2}} + \frac{128(3i + 8x)}{1215(3ix + 4x^2)^{3/2}} + \frac{4096(3i + 8x)}{10935\sqrt{3ix + 4x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.15 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.61

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{1458i - 6480x - 69120ix^2 - 552960x^3 + 983040ix^4 + 524288x^5}{10935(x(3i + 4x))^{5/2}}$$

[In] Integrate[((3\*I)\*x + 4\*x^2)^(-7/2),x]

[Out] (1458\*I - 6480\*x - (69120\*I)\*x^2 - 552960\*x^3 + (983040\*I)\*x^4 + 524288\*x^5)/(10935\*(x\*(3\*I + 4\*x))^(5/2))

**Maple [A] (verified)**

Time = 1.90 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.63

method	result
gospers	$\frac{2x(3i+4x)(262144x^5+491520ix^4-276480x^3-34560ix^2-3240x+729i)}{10935(4x^2+3ix)^{\frac{7}{2}}}$
risch	$\frac{\frac{524288}{10935}x^5 + \frac{65536}{729}ix^4 - \frac{4096}{81}x^3 - \frac{512}{81}ix^2 - \frac{16}{27}x + \frac{2}{15}i}{x^2(3i+4x)^2\sqrt{x(3i+4x)}}$
pseudoelliptic	$\frac{524288}{10935}x^5 + \frac{65536}{729}ix^4 - \frac{4096}{81}x^3 - \frac{512}{81}ix^2 - \frac{16}{27}x + \frac{2}{15}i}{x^2(3i+4x)^2\sqrt{x(3i+4x)}}$
default	$\frac{\frac{2i}{15} + \frac{16x}{45}}{(4x^2+3ix)^{\frac{5}{2}}} + \frac{\frac{128i}{405} + \frac{1024x}{1215}}{(4x^2+3ix)^{\frac{3}{2}}} + \frac{\frac{4096i}{3645} + \frac{32768x}{10935}}{\sqrt{4x^2+3ix}}$
trager	$\frac{\left(\frac{1054}{4271484375} + \frac{224i}{1423828125}\right)(1228800000ix^5 + 1638400000x^5 + 3072000000ix^4 - 2304000000x^4 - 1296000000ix^3 - 1728000000ix^3 - 1296000000ix^3 - 1728000000ix^3)}{(12ix-16x-12i-9)^3x^3}$

[In] `int(1/(3*I*x+4*x^2)^(7/2),x,method=_RETURNVERBOSE)`

[Out]  $2/10935*x*(3*I+4*x)*(491520*I*x^4+262144*x^5-34560*I*x^2-276480*x^3+729*I-3240*x)/(3*I*x+4*x^2)^(7/2)$

## Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.05

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{2(524288x^6 + 1179648ix^5 - 884736x^4 - 221184ix^3 + (262144x^5 + 491520ix^4 - 276480x^3 - 3240x + 729I)\sqrt{4x^2 + 3Ix})}{10935(64x^6 + 144ix^5 - 108x^4 - 27I^2x^3)}$$

[In] `integrate(1/(3*I*x+4*x^2)^(7/2),x, algorithm="fricas")`

[Out]  $2/10935*(524288*x^6 + 1179648*I*x^5 - 884736*x^4 - 221184*I*x^3 + (262144*x^5 + 491520*I*x^4 - 276480*x^3 - 34560*I*x^2 - 3240*x + 729*I)*\sqrt{4*x^2 + 3*I*x})/(64*x^6 + 144*I*x^5 - 108*x^4 - 27*I*x^3)$

## Sympy [F]

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \int \frac{1}{(4x^2 + 3ix)^{7/2}} dx$$

[In] `integrate(1/(3*I*x+4*x**2)**(7/2),x)`

[Out] `Integral((4*x**2 + 3*I*x)**(-7/2), x)`

## Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.04

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{32768x}{10935\sqrt{4x^2 + 3ix}} + \frac{4096i}{3645\sqrt{4x^2 + 3ix}} + \frac{1024x}{1215(4x^2 + 3ix)^{3/2}} + \frac{128i}{405(4x^2 + 3ix)^{3/2}} + \frac{16x}{45(4x^2 + 3ix)^{5/2}} + \frac{2i}{15(4x^2 + 3ix)^{5/2}}$$

[In] `integrate(1/(3*I*x+4*x^2)^(7/2),x, algorithm="maxima")`

[Out]  $32768/10935*x/\sqrt{4*x^2 + 3*I*x} + 4096/3645*I/\sqrt{4*x^2 + 3*I*x} + 1024/1215*x/(4*x^2 + 3*I*x)^(3/2) + 128/405*I/(4*x^2 + 3*I*x)^(3/2) + 16/45*x/(4*x^2 + 3*I*x)^(5/2) + 2/15*I/(4*x^2 + 3*I*x)^(5/2)$

**Giac [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.06

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = \frac{(8(32(8(16(8x + 15i)x - 135)x - 135i)x - 405)x + 729i)\sqrt{8x^2 + 2}\sqrt{16x^2 + 9x}}{10935(4x^2 + 3ix)^3}$$

[In] integrate(1/(3\*I\*x+4\*x^2)^(7/2),x, algorithm="giac")

[Out] 1/10935\*(8\*(32\*(8\*(16\*(8\*x + 15\*I)\*x - 135)\*x - 135\*I)\*x - 405)\*x + 729\*I)\*  
 sqrt(8\*x^2 + 2\*sqrt(16\*x^2 + 9)\*x)\*(3\*I\*x/(4\*x^2 + sqrt(16\*x^4 + 9\*x^2)) +  
 1)/(4\*x^2 + 3\*I\*x)^3

**Mupad [B] (verification not implemented)**

Time = 9.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.51

$$\int \frac{1}{(3ix + 4x^2)^{7/2}} dx = -\frac{-524288x^5 - x^4 983040i + 552960x^3 + x^2 69120i + 6480x - 1458i}{10935(x(4x + 3i))^{5/2}}$$

[In] int(1/(x\*3i + 4\*x^2)^(7/2),x)

[Out] -(6480\*x + x^2\*69120i + 552960\*x^3 - x^4\*983040i - 524288\*x^5 - 1458i)/(109  
 35\*(x\*(4\*x + 3i))^(5/2))

### 3.21 $\int \frac{1}{\sqrt{3x-4x^2}} dx$

Optimal result . . . . .	157
Rubi [A] (verified) . . . . .	157
Mathematica [B] (verified) . . . . .	158
Maple [A] (verified) . . . . .	158
Fricas [B] (verification not implemented) . . . . .	159
Sympy [A] (verification not implemented) . . . . .	159
Maxima [A] (verification not implemented) . . . . .	159
Giac [B] (verification not implemented) . . . . .	159
Mupad [B] (verification not implemented) . . . . .	160

#### Optimal result

Integrand size = 13, antiderivative size = 12

$$\int \frac{1}{\sqrt{3x-4x^2}} dx = -\frac{1}{2} \arcsin\left(1 - \frac{8x}{3}\right)$$

[Out] 1/2\*arcsin(-1+8/3\*x)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{3x-4x^2}} dx = -\frac{1}{2} \arcsin\left(1 - \frac{8x}{3}\right)$$

[In] Int[1/Sqrt[3\*x - 4\*x^2], x]

[Out] -1/2\*ArcSin[1 - (8\*x)/3]

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= - \left( \frac{1}{6} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{9}}} dx, x, 3 - 8x \right) \right) \\ &= -\frac{1}{2} \sin^{-1} \left( 1 - \frac{8x}{3} \right) \end{aligned}$$

**Mathematica [B] (verified)**

Leaf count is larger than twice the leaf count of optimal. 46 vs. 2(12) = 24.

Time = 0.05 (sec) , antiderivative size = 46, normalized size of antiderivative = 3.83

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = -\frac{\sqrt{x}\sqrt{-3 + 4x} \log(-2\sqrt{x} + \sqrt{-3 + 4x})}{\sqrt{-x(-3 + 4x)}}$$

[In] Integrate[1/Sqrt[3\*x - 4\*x^2],x]

[Out] -((Sqrt[x]\*Sqrt[-3 + 4\*x]\*Log[-2\*Sqrt[x] + Sqrt[-3 + 4\*x]])/Sqrt[-(x\*(-3 + 4\*x))])

**Maple [A] (verified)**

Time = 1.90 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{\arcsin\left(-1 + \frac{8x}{3}\right)}{2}$	9
meijerg	$\arcsin\left(\frac{2\sqrt{3}\sqrt{x}}{3}\right)$	10
pseudoelliptic	$-\arctan\left(\frac{\sqrt{-4x^2+3x}}{2x}\right)$	20
trager	$\frac{\text{RootOf}(\_Z^2+1) \ln\left(-8 \text{RootOf}(\_Z^2+1)x+4\sqrt{-4x^2+3x}+3 \text{RootOf}(\_Z^2+1)\right)}{2}$	41

[In] int(1/(-4\*x^2+3\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*arcsin(-1+8/3\*x)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 19 vs. 2(8) = 16.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.58

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = -\arctan\left(\frac{\sqrt{-4x^2 + 3x}}{2x}\right)$$

[In] integrate(1/(-4\*x^2+3\*x)^(1/2),x, algorithm="fricas")

[Out] -arctan(1/2\*sqrt(-4\*x^2 + 3\*x)/x)

**Sympy [A] (verification not implemented)**

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = \frac{\operatorname{asin}\left(\frac{8x}{3} - 1\right)}{2}$$

[In] integrate(1/(-4\*x\*\*2+3\*x)\*\*(1/2),x)

[Out] asin(8\*x/3 - 1)/2

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = -\frac{1}{2} \arcsin\left(-\frac{8}{3}x + 1\right)$$

[In] integrate(1/(-4\*x^2+3\*x)^(1/2),x, algorithm="maxima")

[Out] -1/2\*arcsin(-8/3\*x + 1)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(8) = 16.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.25

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = \frac{1}{16} \sqrt{-4x^2 + 3x}(8x - 3) + \frac{9}{64} \arcsin\left(\frac{8}{3}x - 1\right)$$

[In] integrate(1/(-4\*x^2+3\*x)^(1/2),x, algorithm="giac")

[Out] 1/16\*sqrt(-4\*x^2 + 3\*x)\*(8\*x - 3) + 9/64\*arcsin(8/3\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 8.97 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{\sqrt{3x - 4x^2}} dx = \frac{\operatorname{asin}\left(\frac{8x}{3} - 1\right)}{2}$$

[In] `int(1/(3*x - 4*x^2)^(1/2),x)`

[Out] `asin((8*x)/3 - 1)/2`



$$3.22 \quad \int \frac{1}{(3x-4x^2)^{3/2}} dx$$

Optimal result	161
Rubi [A] (verified)	161
Mathematica [A] (verified)	162
Maple [A] (verified)	162
Fricas [A] (verification not implemented)	162
Sympy [F]	163
Maxima [A] (verification not implemented)	163
Giac [A] (verification not implemented)	163
Mupad [B] (verification not implemented)	163

### Optimal result

Integrand size = 13, antiderivative size = 22

$$\int \frac{1}{(3x-4x^2)^{3/2}} dx = -\frac{2(3-8x)}{9\sqrt{3x-4x^2}}$$

[Out]  $-2/9*(3-8*x)/(-4*x^2+3*x)^(1/2)$

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$ , Rules used = {627}

$$\int \frac{1}{(3x-4x^2)^{3/2}} dx = -\frac{2(3-8x)}{9\sqrt{3x-4x^2}}$$

[In]  $\text{Int}[(3*x - 4*x^2)^{-3/2}, x]$

[Out]  $(-2*(3 - 8*x))/(9*\text{Sqrt}[3*x - 4*x^2])$

#### Rule 627

$\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^{-3/2}, x\_Symbol] \rightarrow \text{Simp}[-2*((b + 2*c*x)/((b^2 - 4*a*c)*\text{Sqrt}[a + b*x + c*x^2])), x] /;$   $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

#### Rubi steps

$$\text{integral} = -\frac{2(3-8x)}{9\sqrt{3x-4x^2}}$$

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.95

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = \frac{2(-3 + 8x)}{9\sqrt{-x(-3 + 4x)}}$$

[In] Integrate[(3\*x - 4\*x^2)^(-3/2),x]

[Out] (2\*(-3 + 8\*x))/(9\*Sqrt[-(x\*(-3 + 4\*x))])

**Maple [A] (verified)**

Time = 1.80 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{2(3-8x)}{9\sqrt{-4x^2+3x}}$	19
pseudoelliptic	$\frac{-\frac{2}{3} + \frac{16x}{9}}{\sqrt{-4x^2+3x}}$	19
meijerg	$-\frac{2\sqrt{3}\left(1-\frac{8x}{3}\right)}{9\sqrt{x}\sqrt{-\frac{4x}{3}+1}}$	21
gospers	$-\frac{2x(4x-3)(-3+8x)}{9(-4x^2+3x)^{\frac{3}{2}}}$	25
trager	$-\frac{2(-3+8x)\sqrt{-4x^2+3x}}{9x(4x-3)}$	29

[In] int(1/(-4\*x^2+3\*x)^(3/2),x,method=\_RETURNVERBOSE)

[Out] -2/9\*(3-8\*x)/(-4\*x^2+3\*x)^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = -\frac{2\sqrt{-4x^2 + 3x}(8x - 3)}{9(4x^2 - 3x)}$$

[In] integrate(1/(-4\*x^2+3\*x)^(3/2),x, algorithm="fricas")

[Out] -2/9\*sqrt(-4\*x^2 + 3\*x)\*(8\*x - 3)/(4\*x^2 - 3\*x)

**Sympy [F]**

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = \int \frac{1}{(-4x^2 + 3x)^{\frac{3}{2}}} dx$$

[In] integrate(1/(-4\*x\*\*2+3\*x)\*\*(3/2),x)

[Out] Integral((-4\*x\*\*2 + 3\*x)\*\*(-3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.22 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.27

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = \frac{16x}{9\sqrt{-4x^2 + 3x}} - \frac{2}{3\sqrt{-4x^2 + 3x}}$$

[In] integrate(1/(-4\*x^2+3\*x)^(3/2),x, algorithm="maxima")

[Out] 16/9\*x/sqrt(-4\*x^2 + 3\*x) - 2/3/sqrt(-4\*x^2 + 3\*x)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.32

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = -\frac{2\sqrt{-4x^2 + 3x}(8x - 3)}{9(4x^2 - 3x)}$$

[In] integrate(1/(-4\*x^2+3\*x)^(3/2),x, algorithm="giac")

[Out] -2/9\*sqrt(-4\*x^2 + 3\*x)\*(8\*x - 3)/(4\*x^2 - 3\*x)

**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int \frac{1}{(3x - 4x^2)^{3/2}} dx = \frac{16x - 6}{9\sqrt{3x - 4x^2}}$$

[In] int(1/(3\*x - 4\*x^2)^(3/2),x)

[Out] (16\*x - 6)/(9\*(3\*x - 4\*x^2)^(1/2))

### 3.23 $\int \frac{1}{(3x-4x^2)^{5/2}} dx$

Optimal result	164
Rubi [A] (verified)	164
Mathematica [A] (verified)	165
Maple [A] (verified)	165
Fricas [A] (verification not implemented)	166
Sympy [F]	166
Maxima [A] (verification not implemented)	166
Giac [A] (verification not implemented)	167
Mupad [B] (verification not implemented)	167

#### Optimal result

Integrand size = 13, antiderivative size = 45

$$\int \frac{1}{(3x-4x^2)^{5/2}} dx = -\frac{2(3-8x)}{27(3x-4x^2)^{3/2}} - \frac{64(3-8x)}{243\sqrt{3x-4x^2}}$$

[Out]  $-2/27*(3-8*x)/(-4*x^2+3*x)^(3/2)-64/243*(3-8*x)/(-4*x^2+3*x)^(1/2)$

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {628, 627}

$$\int \frac{1}{(3x-4x^2)^{5/2}} dx = -\frac{64(3-8x)}{243\sqrt{3x-4x^2}} - \frac{2(3-8x)}{27(3x-4x^2)^{3/2}}$$

[In]  $\text{Int}[(3*x - 4*x^2)^{-5/2}, x]$

[Out]  $(-2*(3 - 8*x))/(27*(3*x - 4*x^2)^(3/2)) - (64*(3 - 8*x))/(243*\text{Sqrt}[3*x - 4*x^2])$

#### Rule 627

$\text{Int}[(a_. + (b_.)*(x_) + (c_.)*(x_)^2)^{-3/2}, x\_Symbol] \rightarrow \text{Simp}[-2*((b + 2*c*x)/((b^2 - 4*a*c)*\text{Sqrt}[a + b*x + c*x^2])), x] /; \text{FreeQ}\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

#### Rule 628

$\text{Int}[(a_. + (b_.)*(x_) + (c_.)*(x_)^2)^{p_}, x\_Symbol] \rightarrow \text{Simp}[(b + 2*c*x)*((a + b*x + c*x^2)^{(p+1})/((p+1)*(b^2 - 4*a*c))), x] - \text{Dist}[2*c*((2*p + 1)*(a + b*x + c*x^2)^{p+1})/((p+1)*(b^2 - 4*a*c)), x]$

$3)/((p + 1)*(b^2 - 4*a*c))$ , Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; Free Q[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{2(3-8x)}{27(3x-4x^2)^{3/2}} + \frac{32}{27} \int \frac{1}{(3x-4x^2)^{3/2}} dx \\ &= -\frac{2(3-8x)}{27(3x-4x^2)^{3/2}} - \frac{64(3-8x)}{243\sqrt{3x-4x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

$$\int \frac{1}{(3x-4x^2)^{5/2}} dx = -\frac{54+432x-2304x^2+2048x^3}{243(-x(-3+4x))^{3/2}}$$

[In] Integrate[(3\*x - 4\*x^2)^(-5/2), x]

[Out] -1/243\*(54 + 432\*x - 2304\*x^2 + 2048\*x^3)/(-(x\*(-3 + 4\*x)))^(3/2)

**Maple [A] (verified)**

Time = 1.82 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.69

method	result	size
meijerg	$-\frac{2\sqrt{3}\left(\frac{1024}{27}x^3 - \frac{128}{3}x^2 + 8x + 1\right)}{81x^{\frac{3}{2}}\left(-\frac{4x}{3} + 1\right)^{\frac{3}{2}}}$	31
gospers	$\frac{2x(4x-3)(1024x^3-1152x^2+216x+27)}{243(-4x^2+3x)^{\frac{5}{2}}}$	35
default	$-\frac{2(3-8x)}{27(-4x^2+3x)^{\frac{3}{2}}} - \frac{64(3-8x)}{243\sqrt{-4x^2+3x}}$	38
trager	$-\frac{2(1024x^3-1152x^2+216x+27)\sqrt{-4x^2+3x}}{243(4x-3)^2x^2}$	39
pseudoelliptic	$\frac{2048x^3-2304x^2+432x+54}{\sqrt{-4x^2+3x}(972x^2-729x)}$	39

[In] int(1/(-4\*x^2+3\*x)^(5/2), x, method=\_RETURNVERBOSE)

[Out] -2/81/x^(3/2)\*3^(1/2)\*(1024/27\*x^3-128/3\*x^2+8\*x+1)/(-4/3\*x+1)^(3/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int \frac{1}{(3x - 4x^2)^{5/2}} dx = -\frac{2(1024x^3 - 1152x^2 + 216x + 27)\sqrt{-4x^2 + 3x}}{243(16x^4 - 24x^3 + 9x^2)}$$

[In] integrate(1/(-4\*x^2+3\*x)^(5/2),x, algorithm="fricas")

[Out] -2/243\*(1024\*x^3 - 1152\*x^2 + 216\*x + 27)\*sqrt(-4\*x^2 + 3\*x)/(16\*x^4 - 24\*x^3 + 9\*x^2)

**Sympy [F]**

$$\int \frac{1}{(3x - 4x^2)^{5/2}} dx = \int \frac{1}{(-4x^2 + 3x)^{5/2}} dx$$

[In] integrate(1/(-4\*x\*\*2+3\*x)\*\*(5/2),x)

[Out] Integral((-4\*x\*\*2 + 3\*x)\*\*(-5/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.22

$$\int \frac{1}{(3x - 4x^2)^{5/2}} dx = \frac{512x}{243\sqrt{-4x^2 + 3x}} - \frac{64}{81\sqrt{-4x^2 + 3x}} + \frac{16x}{27(-4x^2 + 3x)^{3/2}} - \frac{2}{9(-4x^2 + 3x)^{3/2}}$$

[In] integrate(1/(-4\*x^2+3\*x)^(5/2),x, algorithm="maxima")

[Out] 512/243\*x/sqrt(-4\*x^2 + 3\*x) - 64/81/sqrt(-4\*x^2 + 3\*x) + 16/27\*x/(-4\*x^2 + 3\*x)^(3/2) - 2/9/(-4\*x^2 + 3\*x)^(3/2)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.87

$$\int \frac{1}{(3x - 4x^2)^{5/2}} dx = -\frac{2(8(16(8x - 9)x + 27)x + 27)\sqrt{-4x^2 + 3x}}{243(4x^2 - 3x)^2}$$

[In] integrate(1/(-4\*x^2+3\*x)^(5/2),x, algorithm="giac")

[Out] -2/243\*(8\*(16\*(8\*x - 9)\*x + 27)\*x + 27)\*sqrt(-4\*x^2 + 3\*x)/(4\*x^2 - 3\*x)^2

**Mupad [B] (verification not implemented)**

Time = 9.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.62

$$\int \frac{1}{(3x - 4x^2)^{5/2}} dx = \frac{(16x - 6)(-128x^2 + 96x + 9)}{243(3x - 4x^2)^{3/2}}$$

[In] int(1/(3\*x - 4\*x^2)^(5/2),x)

[Out] ((16\*x - 6)\*(96\*x - 128\*x^2 + 9))/(243\*(3\*x - 4\*x^2)^(3/2))

### 3.24 $\int \frac{1}{(3x-4x^2)^{7/2}} dx$

Optimal result	168
Rubi [A] (verified)	168
Mathematica [A] (verified)	169
Maple [A] (verified)	169
Fricas [A] (verification not implemented)	170
Sympy [F]	170
Maxima [A] (verification not implemented)	170
Giac [A] (verification not implemented)	171
Mupad [B] (verification not implemented)	171

#### Optimal result

Integrand size = 13, antiderivative size = 67

$$\int \frac{1}{(3x-4x^2)^{7/2}} dx = -\frac{2(3-8x)}{45(3x-4x^2)^{5/2}} - \frac{128(3-8x)}{1215(3x-4x^2)^{3/2}} - \frac{4096(3-8x)}{10935\sqrt{3x-4x^2}}$$

[Out]  $-2/45*(3-8*x)/(-4*x^2+3*x)^(5/2)-128/1215*(3-8*x)/(-4*x^2+3*x)^(3/2)-4096/10935*(3-8*x)/(-4*x^2+3*x)^(1/2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {628, 627}

$$\int \frac{1}{(3x-4x^2)^{7/2}} dx = -\frac{4096(3-8x)}{10935\sqrt{3x-4x^2}} - \frac{128(3-8x)}{1215(3x-4x^2)^{3/2}} - \frac{2(3-8x)}{45(3x-4x^2)^{5/2}}$$

[In]  $\text{Int}[(3*x - 4*x^2)^{-7/2}, x]$

[Out]  $(-2*(3 - 8*x))/(45*(3*x - 4*x^2)^(5/2)) - (128*(3 - 8*x))/(1215*(3*x - 4*x^2)^(3/2)) - (4096*(3 - 8*x))/(10935*sqrt[3*x - 4*x^2])$

Rule 627

$\text{Int}[(a_. + (b_.)*(x_) + (c_.)*(x_)^2)^{-3/2}, x\_Symbol] \rightarrow \text{Simp}[-2*((b + 2*c*x)/((b^2 - 4*a*c)*sqrt[a + b*x + c*x^2])), x] /; \text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 628



```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{2(3-8x)}{45(3x-4x^2)^{5/2}} + \frac{64}{45} \int \frac{1}{(3x-4x^2)^{5/2}} dx \\ &= -\frac{2(3-8x)}{45(3x-4x^2)^{5/2}} - \frac{128(3-8x)}{1215(3x-4x^2)^{3/2}} + \frac{2048 \int \frac{1}{(3x-4x^2)^{3/2}} dx}{1215} \\ &= -\frac{2(3-8x)}{45(3x-4x^2)^{5/2}} - \frac{128(3-8x)}{1215(3x-4x^2)^{3/2}} - \frac{4096(3-8x)}{10935\sqrt{3x-4x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.61

$$\int \frac{1}{(3x-4x^2)^{7/2}} dx = \frac{2(-729-3240x-34560x^2+276480x^3-491520x^4+262144x^5)}{10935(-x(-3+4x))^{5/2}}$$

[In] Integrate[(3\*x - 4\*x^2)^(-7/2), x]

[Out] (2\*(-729 - 3240\*x - 34560\*x^2 + 276480\*x^3 - 491520\*x^4 + 262144\*x^5))/(10935\*(-x\*(-3 + 4\*x)))^(5/2)

**Maple [A] (verified)**

Time = 1.83 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.61

method	result	size
meijerg	$-\frac{2\sqrt{3} \left(-\frac{262144}{243}x^5 + \frac{163840}{81}x^4 - \frac{10240}{9}x^3 + \frac{1280}{9}x^2 + \frac{40}{3}x + 3\right)}{1215x^{\frac{5}{2}} \left(-\frac{4x}{3} + 1\right)^{\frac{5}{2}}}$	41
gospers	$-\frac{2x(4x-3)(262144x^5 - 491520x^4 + 276480x^3 - 34560x^2 - 3240x - 729)}{10935(-4x^2+3x)^{\frac{7}{2}}}$	45
trager	$-\frac{2(262144x^5 - 491520x^4 + 276480x^3 - 34560x^2 - 3240x - 729)\sqrt{-4x^2+3x}}{10935(4x-3)^3x^3}$	49
pseudoelliptic	$\frac{524288x^5 - \frac{65536}{729}x^4 + \frac{4096}{81}x^3 - \frac{512}{81}x^2 - \frac{16}{27}x - \frac{2}{15}}{(4x-3)^2x^2\sqrt{-4x^2+3x}}$	49
default	$-\frac{2(3-8x)}{45(-4x^2+3x)^{\frac{5}{2}}} - \frac{128(3-8x)}{1215(-4x^2+3x)^{\frac{3}{2}}} - \frac{4096(3-8x)}{10935\sqrt{-4x^2+3x}}$	56

[In] `int(1/(-4*x^2+3*x)^(7/2),x,method=_RETURNVERBOSE)`

[Out]  $-2/1215/x^{5/2}*3^{1/2}*(-262144/243*x^5+163840/81*x^4-10240/9*x^3+1280/9*x^2+40/3*x+3)/(-4/3*x+1)^{5/2}$

## Fricas [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.91

$$\int \frac{1}{(3x - 4x^2)^{7/2}} dx = \frac{2(262144x^5 - 491520x^4 + 276480x^3 - 34560x^2 - 3240x - 729)\sqrt{-4x^2 + 3x}}{10935(64x^6 - 144x^5 + 108x^4 - 27x^3)}$$

[In] `integrate(1/(-4*x^2+3*x)^(7/2),x, algorithm="fricas")`

[Out]  $-2/10935*(262144*x^5 - 491520*x^4 + 276480*x^3 - 34560*x^2 - 3240*x - 729)*\sqrt{-4*x^2 + 3*x}/(64*x^6 - 144*x^5 + 108*x^4 - 27*x^3)$

## Sympy [F]

$$\int \frac{1}{(3x - 4x^2)^{7/2}} dx = \int \frac{1}{(-4x^2 + 3x)^{7/2}} dx$$

[In] `integrate(1/(-4*x**2+3*x)**(7/2),x)`

[Out] `Integral((-4*x**2 + 3*x)**(-7/2), x)`

## Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.22

$$\int \frac{1}{(3x - 4x^2)^{7/2}} dx = \frac{32768x}{10935\sqrt{-4x^2 + 3x}} - \frac{4096}{3645\sqrt{-4x^2 + 3x}} + \frac{1024x}{1215(-4x^2 + 3x)^{3/2}} - \frac{128}{405(-4x^2 + 3x)^{3/2}} + \frac{16x}{45(-4x^2 + 3x)^{5/2}} - \frac{2}{15(-4x^2 + 3x)^{5/2}}$$

[In] `integrate(1/(-4*x^2+3*x)^(7/2),x, algorithm="maxima")`

[Out]  $32768/10935*x/\sqrt{-4*x^2 + 3*x} - 4096/3645/\sqrt{-4*x^2 + 3*x} + 1024/1215*x/(-4*x^2 + 3*x)^{3/2} - 128/405/(-4*x^2 + 3*x)^{3/2} + 16/45*x/(-4*x^2 + 3*x)^{5/2} - 2/15/(-4*x^2 + 3*x)^{5/2}$

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.73

$$\int \frac{1}{(3x - 4x^2)^{7/2}} dx = \frac{2(8(32(8(16(8x - 15)x + 135)x - 135)x - 405)x - 729)\sqrt{-4x^2 + 3x}}{10935(4x^2 - 3x)^3}$$

[In] integrate(1/(-4\*x^2+3\*x)^(7/2),x, algorithm="giac")

[Out] -2/10935\*(8\*(32\*(8\*(16\*(8\*x - 15)\*x + 135)\*x - 135)\*x - 405)\*x - 729)\*sqrt(-4\*x^2 + 3\*x)/(4\*x^2 - 3\*x)^3

**Mupad [B] (verification not implemented)**

Time = 9.08 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.09

$$\int \frac{1}{(3x - 4x^2)^{7/2}} dx = \frac{6480x - 9216x(3x - 4x^2) - 32768x(3x - 4x^2)^2 + 12288(3x - 4x^2)^2 - 13824x^2 + 1458}{(3x - 4x^2)^{3/2}(32805x - 43740x^2)}$$

[In] int(1/(3\*x - 4\*x^2)^(7/2),x)

[Out] -(6480\*x - 9216\*x\*(3\*x - 4\*x^2) - 32768\*x\*(3\*x - 4\*x^2)^2 + 12288\*(3\*x - 4\*x^2)^2 - 13824\*x^2 + 1458)/((3\*x - 4\*x^2)^(3/2)\*(32805\*x - 43740\*x^2))

### 3.25 $\int \frac{1}{\sqrt{bx-b^2x^2}} dx$

Optimal result	172
Rubi [A] (verified)	172
Mathematica [B] (verified)	173
Maple [B] (verified)	173
Fricas [B] (verification not implemented)	174
Sympy [B] (verification not implemented)	174
Maxima [A] (verification not implemented)	174
Giac [B] (verification not implemented)	175
Mupad [B] (verification not implemented)	175

#### Optimal result

Integrand size = 16, antiderivative size = 12

$$\int \frac{1}{\sqrt{bx-b^2x^2}} dx = -\frac{\arcsin(1-2bx)}{b}$$

[Out]  $\arcsin(2*b*x-1)/b$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{bx-b^2x^2}} dx = -\frac{\arcsin(1-2bx)}{b}$$

[In]  $\text{Int}[1/\text{Sqrt}[b*x - b^2*x^2], x]$

[Out]  $-(\text{ArcSin}[1 - 2*b*x])/b$

#### Rule 222

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /;$   $\text{FreeQ}\{a, b, x\} \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{NegQ}[b]$

#### Rule 633

$\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2]^{(p_)}, x\_Symbol] \rightarrow \text{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), \text{Subst}[\text{Int}[\text{Simp}[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b + 2*c*x], x] /;$   $\text{FreeQ}\{a, b, c, p, x\} \ \&\& \ \text{GtQ}[4*a - b^2/c, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{b^2}}} dx, x, b-2b^2x\right)}{b^2} \\ &= -\frac{\sin^{-1}(1-2bx)}{b} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 57 vs.  $2(12) = 24$ .

Time = 0.07 (sec) , antiderivative size = 57, normalized size of antiderivative = 4.75

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = -\frac{2\sqrt{x}\sqrt{-1+bx} \log\left(-\sqrt{b}\sqrt{x} + \sqrt{-1+bx}\right)}{\sqrt{b}\sqrt{-bx(-1+bx)}}$$

[In] Integrate[1/Sqrt[b\*x - b^2\*x^2],x]

[Out]  $(-2*\text{Sqrt}[x]*\text{Sqrt}[-1 + b*x]*\text{Log}[-(\text{Sqrt}[b]*\text{Sqrt}[x]) + \text{Sqrt}[-1 + b*x]])/(\text{Sqrt}[b]*\text{Sqrt}[-(b*x*(-1 + b*x))])$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 24 vs.  $2(11) = 22$ .

Time = 2.18 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.08

method	result	size
pseudoelliptic	$-\frac{2 \arctan\left(\frac{\sqrt{-bx(bx-1)}}{xb}\right)}{b}$	25
default	$\frac{\arctan\left(\frac{\sqrt{b^2}\left(x-\frac{1}{2b}\right)}{\sqrt{-b^2x^2+bx}}\right)}{\sqrt{b^2}}$	35

[In] int(1/(-b^2\*x^2+b\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out]  $-2*\arctan((-b*x*(b*x-1))^(1/2)/x/b)/b$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(11) = 22.  
 Time = 0.38 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.25

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{-b^2x^2+bx}}{bx}\right)}{b}$$

[In] integrate(1/(-b^2\*x^2+b\*x)^(1/2),x, algorithm="fricas")

[Out] -2\*arctan(sqrt(-b^2\*x^2 + b\*x)/(b\*x))/b

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. 2(8) = 16.  
 Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 4.50

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = \begin{cases} \frac{\log(-2b^2x+b+2\sqrt{-b^2}\sqrt{-b^2x^2+bx})}{\sqrt{-b^2}} & \text{for } b^2 \neq 0 \\ \frac{2\sqrt{bx}}{b} & \text{for } b \neq 0 \\ \tilde{\infty}x & \text{otherwise} \end{cases}$$

[In] integrate(1/(-b\*\*2\*x\*\*2+b\*x)\*\*(1/2),x)

[Out] Piecewise((log(-2\*b\*\*2\*x + b + 2\*sqrt(-b\*\*2)\*sqrt(-b\*\*2\*x\*\*2 + b\*x))/sqrt(-b\*\*2), Ne(b\*\*2, 0)), (2\*sqrt(b\*x)/b, Ne(b, 0)), (zoo\*x, True))

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = -\frac{\arcsin\left(\frac{-2b^2x-b}{b}\right)}{b}$$

[In] integrate(1/(-b^2\*x^2+b\*x)^(1/2),x, algorithm="maxima")

[Out] -arcsin(-(2\*b^2\*x - b)/b)/b

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(11) = 22.

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 3.42

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = \frac{1}{4} \sqrt{-b^2x^2 + bx} \left( 2x - \frac{1}{b} \right) - \frac{\arcsin(-2bx + 1) \operatorname{sgn}(b)}{8|b|}$$

[In] integrate(1/(-b^2\*x^2+b\*x)^(1/2),x, algorithm="giac")

[Out] 1/4\*sqrt(-b^2\*x^2 + b\*x)\*(2\*x - 1/b) - 1/8\*arcsin(-2\*b\*x + 1)\*sgn(b)/abs(b)

**Mupad [B] (verification not implemented)**

Time = 9.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.50

$$\int \frac{1}{\sqrt{bx - b^2x^2}} dx = \frac{\ln\left(\frac{\frac{b}{2} - b^2x}{\sqrt{-b^2}} + \sqrt{bx - b^2x^2}\right)}{\sqrt{-b^2}}$$

[In] int(1/(b\*x - b^2\*x^2)^(1/2),x)

[Out] log((b/2 - b^2\*x)/(-b^2)^(1/2) + (b\*x - b^2\*x^2)^(1/2))/(-b^2)^(1/2)

### 3.26 $\int \frac{1}{\sqrt{bx+b^2x^2}} dx$

Optimal result	176
Rubi [A] (verified)	176
Mathematica [B] (verified)	177
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	178
Sympy [B] (verification not implemented)	178
Maxima [A] (verification not implemented)	178
Giac [B] (verification not implemented)	179
Mupad [B] (verification not implemented)	179

#### Optimal result

Integrand size = 15, antiderivative size = 24

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \frac{2\operatorname{arctanh}\left(\frac{bx}{\sqrt{bx+b^2x^2}}\right)}{b}$$

[Out]  $2*\operatorname{arctanh}(b*x/(b^2*x^2+b*x)^{(1/2)})/b$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {634, 212}

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \frac{2\operatorname{arctanh}\left(\frac{bx}{\sqrt{b^2x^2+bx}}\right)}{b}$$

[In] `Int[1/Sqrt[b*x + b^2*x^2], x]`

[Out]  $(2*\operatorname{ArcTanh}[(b*x)/\operatorname{Sqrt}[b*x + b^2*x^2]])/b$

#### Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

#### Rule 634

`Int[1/Sqrt[(b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(1 - c*x^2), x], x, x/Sqrt[b*x + c*x^2]], x] /; FreeQ[{b, c}, x]`



Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{1-b^2x^2} dx, x, \frac{x}{\sqrt{bx+b^2x^2}}\right) \\ &= \frac{2 \tanh^{-1}\left(\frac{bx}{\sqrt{bx+b^2x^2}}\right)}{b} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 58 vs.  $2(24) = 48$ .

Time = 0.09 (sec) , antiderivative size = 58, normalized size of antiderivative = 2.42

$$\int \frac{1}{\sqrt{bx+b^2x^2}} dx = \frac{4\sqrt{x}\sqrt{1+bx}\text{arctanh}\left(\frac{\sqrt{b}\sqrt{x}}{-1+\sqrt{1+bx}}\right)}{\sqrt{b}\sqrt{bx(1+bx)}}$$

[In] Integrate[1/Sqrt[b\*x + b^2\*x^2],x]

[Out] (4\*Sqrt[x]\*Sqrt[1 + b\*x]\*ArcTanh[(Sqrt[b]\*Sqrt[x])/(-1 + Sqrt[1 + b\*x])])/(Sqrt[b]\*Sqrt[b\*x\*(1 + b\*x)])

### Maple [A] (verified)

Time = 2.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

method	result	size
default	$\frac{\ln\left(\frac{\frac{1}{2}b+b^2x}{\sqrt{b^2}}+\sqrt{b^2x^2+bx}\right)}{\sqrt{b^2}}$	37
pseudoelliptic	$\frac{-\ln\left(\frac{-bx+\sqrt{bx(bx+1)}}{x}\right)+\ln\left(\frac{bx+\sqrt{bx(bx+1)}}{x}\right)}{b}$	47

[In] int(1/(b^2\*x^2+b\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out] ln((1/2\*b+b^2\*x)/(b^2)^(1/2)+(b^2\*x^2+b\*x)^(1/2))/(b^2)^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.43 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = -\frac{\log(-2bx + 2\sqrt{b^2x^2 + bx} - 1)}{b}$$

[In] integrate(1/(b^2\*x^2+b\*x)^(1/2),x, algorithm="fricas")

[Out] -log(-2\*b\*x + 2\*sqrt(b^2\*x^2 + b\*x) - 1)/b

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(20) = 40.

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.12

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \begin{cases} \frac{\log(2b^2x + b + 2\sqrt{b^2x^2 + bx}\sqrt{b^2})}{\sqrt{b^2}} & \text{for } b^2 \neq 0 \\ \frac{2\sqrt{bx}}{b} & \text{for } b \neq 0 \\ \tilde{\infty}x & \text{otherwise} \end{cases}$$

[In] integrate(1/(b\*\*2\*x\*\*2+b\*x)\*\*(1/2),x)

[Out] Piecewise((log(2\*b\*\*2\*x + b + 2\*sqrt(b\*\*2\*x\*\*2 + b\*x)\*sqrt(b\*\*2))/sqrt(b\*\*2), Ne(b\*\*2, 0)), (2\*sqrt(b\*x)/b, Ne(b, 0)), (zoo\*x, True))

**Maxima [A] (verification not implemented)**

none

Time = 0.22 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \frac{\log(2b^2x + 2\sqrt{b^2x^2 + bxb} + b)}{b}$$

[In] integrate(1/(b^2\*x^2+b\*x)^(1/2),x, algorithm="maxima")

[Out] log(2\*b^2\*x + 2\*sqrt(b^2\*x^2 + b\*x)\*b + b)/b

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(22) = 44.

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.46

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \frac{1}{4} \sqrt{b^2x^2 + bx} \left( 2x + \frac{1}{b} \right) + \frac{\log(|-2(x|b| - \sqrt{b^2x^2 + bx})|b| - b|)}{8|b|}$$

[In] integrate(1/(b^2\*x^2+b\*x)^(1/2),x, algorithm="giac")

[Out] 1/4\*sqrt(b^2\*x^2 + b\*x)\*(2\*x + 1/b) + 1/8\*log(abs(-2\*(x\*abs(b) - sqrt(b^2\*x^2 + b\*x))\*abs(b) - b))/abs(b)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{bx + b^2x^2}} dx = \frac{\ln\left(\frac{x b^2 + \frac{b}{2}}{\sqrt{b^2}} + \sqrt{b^2 x^2 + b x}\right)}{\sqrt{b^2}}$$

[In] int(1/(b\*x + b^2\*x^2)^(1/2),x)

[Out] log((b/2 + b^2\*x)/(b^2)^(1/2) + (b\*x + b^2\*x^2)^(1/2))/(b^2)^(1/2)

### 3.27 $\int \frac{1}{\sqrt{6x-x^2}} dx$

Optimal result	180
Rubi [A] (verified)	180
Mathematica [B] (verified)	181
Maple [A] (verified)	181
Fricas [B] (verification not implemented)	182
Sympy [A] (verification not implemented)	182
Maxima [A] (verification not implemented)	182
Giac [B] (verification not implemented)	182
Mupad [B] (verification not implemented)	183

#### Optimal result

Integrand size = 13, antiderivative size = 10

$$\int \frac{1}{\sqrt{6x-x^2}} dx = -\arcsin\left(1 - \frac{x}{3}\right)$$

[Out] arcsin(-1+1/3\*x)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{6x-x^2}} dx = -\arcsin\left(1 - \frac{x}{3}\right)$$

[In] Int[1/Sqrt[6\*x - x^2],x]

[Out] -ArcSin[1 - x/3]

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= - \left( \frac{1}{6} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{36}}} dx, x, 6 - 2x \right) \right) \\ &= - \sin^{-1} \left( 1 - \frac{x}{3} \right) \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 40 vs.  $2(10) = 20$ .

Time = 0.05 (sec) , antiderivative size = 40, normalized size of antiderivative = 4.00

$$\int \frac{1}{\sqrt{6x - x^2}} dx = - \frac{2\sqrt{-6+x}\sqrt{x} \log(\sqrt{-6+x} - \sqrt{x})}{\sqrt{-((-6+x)x)}}$$

[In] Integrate[1/Sqrt[6\*x - x^2],x]

[Out] (-2\*Sqrt[-6 + x]\*Sqrt[x]\*Log[Sqrt[-6 + x] - Sqrt[x]])/Sqrt[-((-6 + x)\*x)]

### Maple [A] (verified)

Time = 2.28 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

method	result	size
default	$\arcsin\left(-1 + \frac{x}{3}\right)$	7
meijerg	$2 \arcsin\left(\frac{\sqrt{6}\sqrt{x}}{6}\right)$	12
pseudoelliptic	$-2 \arctan\left(\frac{\sqrt{-x(-6+x)}}{x}\right)$	16
trager	$\text{RootOf}(\_Z^2 + 1) \ln\left(-\text{RootOf}(\_Z^2 + 1) x + \sqrt{-x^2 + 6x} + 3 \text{RootOf}(\_Z^2 + 1)\right)$	38

[In] int(1/(-x^2+6\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out] arcsin(-1+1/3\*x)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 18 vs.  $2(6) = 12$ .

Time = 0.77 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.80

$$\int \frac{1}{\sqrt{6x - x^2}} dx = -2 \arctan \left( \frac{\sqrt{-x^2 + 6x}}{x} \right)$$

[In] integrate(1/(-x^2+6\*x)^(1/2),x, algorithm="fricas")

[Out] -2\*arctan(sqrt(-x^2 + 6\*x)/x)

**Sympy [A] (verification not implemented)**

Time = 0.23 (sec) , antiderivative size = 5, normalized size of antiderivative = 0.50

$$\int \frac{1}{\sqrt{6x - x^2}} dx = \operatorname{asin} \left( \frac{x}{3} - 1 \right)$$

[In] integrate(1/(-x\*\*2+6\*x)\*\*(1/2),x)

[Out] asin(x/3 - 1)

**Maxima [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{6x - x^2}} dx = -\arcsin \left( -\frac{1}{3}x + 1 \right)$$

[In] integrate(1/(-x^2+6\*x)^(1/2),x, algorithm="maxima")

[Out] -arcsin(-1/3\*x + 1)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 25 vs.  $2(6) = 12$ .

Time = 0.29 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.50

$$\int \frac{1}{\sqrt{6x - x^2}} dx = \frac{1}{2} \sqrt{-x^2 + 6x}(x - 3) + \frac{9}{2} \arcsin \left( \frac{1}{3}x - 1 \right)$$

[In] integrate(1/(-x^2+6\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(-x^2 + 6\*x)\*(x - 3) + 9/2\*arcsin(1/3\*x - 1)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{\sqrt{6x - x^2}} dx = \operatorname{asin}\left(\frac{x}{3} - 1\right)$$

[In] `int(1/(6*x - x^2)^(1/2),x)`

[Out] `asin(x/3 - 1)`

### 3.28 $\int \frac{1}{\sqrt{4x+x^2}} dx$

Optimal result	184
Rubi [A] (verified)	184
Mathematica [B] (verified)	185
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	186
Sympy [A] (verification not implemented)	186
Maxima [A] (verification not implemented)	186
Giac [B] (verification not implemented)	186
Mupad [B] (verification not implemented)	187

#### Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{\sqrt{4x+x^2}} dx = 2 \operatorname{arctanh}\left(\frac{x}{\sqrt{4x+x^2}}\right)$$

[Out] 2\*arctanh(x/(x^2+4\*x)^(1/2))

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {634, 212}

$$\int \frac{1}{\sqrt{4x+x^2}} dx = 2 \operatorname{arctanh}\left(\frac{x}{\sqrt{x^2+4x}}\right)$$

[In] Int[1/Sqrt[4\*x + x^2], x]

[Out] 2\*ArcTanh[x/Sqrt[4\*x + x^2]]

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 634

Int[1/Sqrt[(b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(1 - c\*x^2), x], x, x/Sqrt[b\*x + c\*x^2]], x] /; FreeQ[{b, c}, x]



Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{4x+x^2}}\right) \\ &= 2 \tanh^{-1}\left(\frac{x}{\sqrt{4x+x^2}}\right) \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 39 vs.  $2(16) = 32$ .

Time = 0.04 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.44

$$\int \frac{1}{\sqrt{4x+x^2}} dx = -\frac{2\sqrt{x}\sqrt{4+x} \log(-\sqrt{x} + \sqrt{4+x})}{\sqrt{x(4+x)}}$$

[In] Integrate[1/Sqrt[4\*x + x^2],x]

[Out] (-2\*Sqrt[x]\*Sqrt[4 + x]\*Log[-Sqrt[x] + Sqrt[4 + x]])/Sqrt[x\*(4 + x)]

### Maple [A] (verified)

Time = 1.90 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.56

method	result	size
meijerg	$2 \operatorname{arcsinh}\left(\frac{\sqrt{x}}{2}\right)$	9
default	$\ln(2 + x + \sqrt{x^2 + 4x})$	14
trager	$\ln(2 + x + \sqrt{x^2 + 4x})$	14
pseudoelliptic	$2 \operatorname{arctanh}\left(\frac{\sqrt{x(4+x)}}{x}\right)$	15

[In] int(1/(x^2+4\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 2\*arcsinh(1/2\*x^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{4x+x^2}} dx = -\log\left(-x + \sqrt{x^2+4x} - 2\right)$$

[In] integrate(1/(x^2+4\*x)^(1/2),x, algorithm="fricas")

[Out] -log(-x + sqrt(x^2 + 4\*x) - 2)

**Sympy [A] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{4x+x^2}} dx = \log\left(2x + 2\sqrt{x^2+4x} + 4\right)$$

[In] integrate(1/(x\*\*2+4\*x)\*\*(1/2),x)

[Out] log(2\*x + 2\*sqrt(x\*\*2 + 4\*x) + 4)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{4x+x^2}} dx = \log\left(2x + 2\sqrt{x^2+4x} + 4\right)$$

[In] integrate(1/(x^2+4\*x)^(1/2),x, algorithm="maxima")

[Out] log(2\*x + 2\*sqrt(x^2 + 4\*x) + 4)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(14) = 28.

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \frac{1}{\sqrt{4x+x^2}} dx = \frac{1}{2}\sqrt{x^2+4x}(x+2) + 2\log\left(\left|-x + \sqrt{x^2+4x} - 2\right|\right)$$

[In] integrate(1/(x^2+4\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(x^2 + 4\*x)\*(x + 2) + 2\*log(abs(-x + sqrt(x^2 + 4\*x) - 2))

**Mupad [B] (verification not implemented)**

Time = 9.37 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int \frac{1}{\sqrt{4x + x^2}} dx = \ln \left( x + \sqrt{x(x+4)} + 2 \right)$$

[In] int(1/(4\*x + x^2)^(1/2),x)

[Out] log(x + (x\*(x + 4))^(1/2) + 2)

### 3.29 $\int \frac{1}{\sqrt{-2x+x^2}} dx$

Optimal result	188
Rubi [A] (verified)	188
Mathematica [B] (verified)	189
Maple [A] (verified)	189
Fricas [A] (verification not implemented)	190
Sympy [A] (verification not implemented)	190
Maxima [A] (verification not implemented)	190
Giac [B] (verification not implemented)	190
Mupad [B] (verification not implemented)	191

#### Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{\sqrt{-2x+x^2}} dx = 2\operatorname{arctanh}\left(\frac{x}{\sqrt{-2x+x^2}}\right)$$

[Out] 2\*arctanh(x/(x^2-2\*x)^(1/2))

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {634, 212}

$$\int \frac{1}{\sqrt{-2x+x^2}} dx = 2\operatorname{arctanh}\left(\frac{x}{\sqrt{x^2-2x}}\right)$$

[In] Int[1/Sqrt[-2\*x + x^2], x]

[Out] 2\*ArcTanh[x/Sqrt[-2\*x + x^2]]

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 634

Int[1/Sqrt[(b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(1 - c\*x^2), x], x, x/Sqrt[b\*x + c\*x^2]], x] /; FreeQ[{b, c}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{x}{\sqrt{-2x+x^2}}\right) \\ &= 2 \tanh^{-1}\left(\frac{x}{\sqrt{-2x+x^2}}\right) \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 39 vs. 2(16) = 32.

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.44

$$\int \frac{1}{\sqrt{-2x+x^2}} dx = -\frac{2\sqrt{-2+x}\sqrt{x} \log(\sqrt{-2+x} - \sqrt{x})}{\sqrt{(-2+x)x}}$$

[In] Integrate[1/Sqrt[-2\*x + x^2],x]

[Out] (-2\*Sqrt[-2 + x]\*Sqrt[x]\*Log[Sqrt[-2 + x] - Sqrt[x]])/Sqrt[(-2 + x)\*x]

### Maple [A] (verified)

Time = 1.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

method	result	size
default	$\ln(-1+x+\sqrt{x^2-2x})$	14
trager	$\ln(-1+x+\sqrt{x^2-2x})$	14
pseudoelliptic	$2 \operatorname{arctanh}\left(\frac{\sqrt{x(-2+x)}}{x}\right)$	15
meijerg	$\frac{2\sqrt{-\operatorname{signum}(-2+x)} \operatorname{arcsin}\left(\frac{\sqrt{2}\sqrt{x}}{2}\right)}{\sqrt{\operatorname{signum}(-2+x)}}$	26

[In] int(1/(x^2-2\*x)^(1/2),x,method=\_RETURNVERBOSE)

[Out] ln(-1+x+(x^2-2\*x)^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{-2x + x^2}} dx = -\log\left(-x + \sqrt{x^2 - 2x + 1}\right)$$

[In] integrate(1/(x^2-2\*x)^(1/2),x, algorithm="fricas")

[Out] -log(-x + sqrt(x^2 - 2\*x) + 1)

**Sympy [A] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{-2x + x^2}} dx = \log\left(2x + 2\sqrt{x^2 - 2x} - 2\right)$$

[In] integrate(1/(x\*\*2-2\*x)\*\*(1/2),x)

[Out] log(2\*x + 2\*sqrt(x\*\*2 - 2\*x) - 2)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{-2x + x^2}} dx = \log\left(2x + 2\sqrt{x^2 - 2x} - 2\right)$$

[In] integrate(1/(x^2-2\*x)^(1/2),x, algorithm="maxima")

[Out] log(2\*x + 2\*sqrt(x^2 - 2\*x) - 2)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(14) = 28.

Time = 0.28 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.06

$$\int \frac{1}{\sqrt{-2x + x^2}} dx = \frac{1}{2} \sqrt{x^2 - 2x}(x - 1) + \frac{1}{2} \log\left(\left|-x + \sqrt{x^2 - 2x} + 1\right|\right)$$

[In] integrate(1/(x^2-2\*x)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(x^2 - 2\*x)\*(x - 1) + 1/2\*log(abs(-x + sqrt(x^2 - 2\*x) + 1))

**Mupad [B] (verification not implemented)**

Time = 9.63 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.69

$$\int \frac{1}{\sqrt{-2x + x^2}} dx = \ln \left( x + \sqrt{x(x-2)} - 1 \right)$$

[In] int(1/(x^2 - 2\*x)^(1/2),x)

[Out] log(x + (x\*(x - 2))^(1/2) - 1)

### 3.30 $\int (bx + cx^2)^{4/3} dx$

Optimal result	192
Rubi [A] (verified)	193
Mathematica [C] (verified)	195
Maple [F]	196
Fricas [F]	196
Sympy [F]	196
Maxima [F]	196
Giac [F]	197
Mupad [B] (verification not implemented)	197

#### Optimal result

Integrand size = 13, antiderivative size = 448

$$\int (bx + cx^2)^{4/3} dx = \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)(bx+cx^2)^{4/3}}{55c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{4/3}(b+2cx)(bx+cx^2)^{4/3}}{22c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{\sqrt[3]{2}3^{3/4}\sqrt{2-\sqrt{3}}b^2(bx+cx^2)^{4/3}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}{55c(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}\sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}$$

[Out]  $\frac{3}{55}(-c*x*(c*x+b)/b^2)^{(1/3)}*(2*c*x+b)*(c*x^2+b*x)^{(4/3)}/c/(-c*(c*x^2+b*x)/b^2)^{(4/3)}+3/22*(-c*x*(c*x+b)/b^2)^{(4/3)}*(2*c*x+b)*(c*x^2+b*x)^{(4/3)}/c/(-c*(c*x^2+b*x)/b^2)^{(4/3)}+1/55*2^{(1/3)}*3^{(3/4)}*b^2*(c*x^2+b*x)^{(4/3)}*(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)})*EllipticF((1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}+3^{(1/2)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)}),2*I-I*3^{(1/2)})*(1/2*6^{(1/2)}-1/2*2^{(1/2)})*((1+2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}+2*2^{(1/3)}*(-c*x*(c*x+b)/b^2)^{(2/3)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}/c/(2*c*x+b)/(-c*(c*x^2+b*x)/b^2)^{(4/3)}/((-1+2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}$



**Rubi [A] (verified)**

Time = 0.50 (sec) , antiderivative size = 448, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {636, 633, 201, 242, 225}

$$\int (bx + cx^2)^{4/3} dx = \frac{\sqrt[3]{23}^{3/4} \sqrt{2 - \sqrt{3}} b^2 (bx + cx^2)^{4/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{2 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) - \sqrt{3}}}}{55c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{\frac{1-2^{2/3}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}} + \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (b+2cx) (bx+cx^2)^{4/3}}{22c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{3 \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (b+2cx) (bx+cx^2)^{4/3}}{55c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}$$

[In] Int[(b\*x + c\*x^2)^(4/3), x]

[Out] (3\*((c\*x\*(b + c\*x))/b^2)^(1/3)\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(4/3))/(55\*c\*(-((c\*(b\*x + c\*x^2))/b^2)^(4/3)) + (3\*((c\*x\*(b + c\*x))/b^2)^(4/3)\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(4/3))/(22\*c\*(-((c\*(b\*x + c\*x^2))/b^2)^(4/3)) + (2^(1/3)\*3^(3/4)\*Sqrt[2 - Sqrt[3]]\*b^2\*(b\*x + c\*x^2)^(4/3)\*(1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3))\*Sqrt[(1 + 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3)) + 2\*2^(1/3)\*(-((c\*x\*(b + c\*x))/b^2)^(2/3))]/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3))^2]\*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3))]/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3))], -7 + 4\*Sqrt[3])]/(55\*c\*(b + 2\*c\*x)\*(-((c\*(b\*x + c\*x^2))/b^2)^(4/3)\*Sqrt[-((1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3)))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2)^(1/3))^2])])

Rule 201

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[x\*((a + b\*x^n)^p/(n\*p + 1)), x] + Dist[a\*n\*(p/(n\*p + 1)), Int[(a + b\*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2\*p] || (EqQ[n, 2] && IntegerQ[4\*p]) || (EqQ[n, 2] && IntegerQ[3\*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 225

```
Int[1/Sqrt[(a_) + (b_)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2])))*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x]] /; FreeQ[{a, b}, x
] && NegQ[a]
```

### Rule 242

```
Int[((a_) + (b_)*(x_)^2)^(-2/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x))
, Subst[Int[1/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

### Rule 633

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

### Rule 636

```
Int[((b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[(b*x + c*x^2)^p/((-
c)*((b*x + c*x^2)/b^2))^p, Int[((-c)*(x/b) - c^2*(x^2/b^2))^p, x], x] /; Fr
eeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4
```

### Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(bx + cx^2)^{4/3} \int \left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{4/3} dx}{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&= -\frac{\left(b^2(bx + cx^2)^{4/3}\right) \text{Subst}\left(\int \left(1 - \frac{b^2x^2}{c^2}\right)^{4/3} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{8 \cdot 2^{2/3} c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&= \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (b + 2cx) (bx + cx^2)^{4/3}}{22c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&= -\frac{\left(b^2(bx + cx^2)^{4/3}\right) \text{Subst}\left(\int \sqrt[3]{1 - \frac{b^2x^2}{c^2}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{11 \cdot 2^{2/3} c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)(bx+cx^2)^{4/3}}{55c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{4/3}(b+2cx)(bx+cx^2)^{4/3}}{22c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&\quad - \frac{\left(\sqrt[3]{2}b^2(bx+cx^2)^{4/3}\right) \text{Subst}\left(\int \frac{1}{(1-\frac{b^2x^2}{c^2})^{2/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{55c^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&= \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)(bx+cx^2)^{4/3}}{55c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{4/3}(b+2cx)(bx+cx^2)^{4/3}}{22c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&\quad + \frac{\left(3(bx+cx^2)^{4/3}\sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3}\sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{55\cdot 2^{2/3}\left(-\frac{c}{b}-\frac{2c^2x}{b^2}\right)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&= \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)(bx+cx^2)^{4/3}}{55c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} + \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{4/3}(b+2cx)(bx+cx^2)^{4/3}}{22c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}} \\
&\quad + \frac{\sqrt[3]{23^{3/4}}\sqrt{2-\sqrt{3}}b^2(bx+cx^2)^{4/3}\sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2}}}{55c(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}\sqrt{-1-\frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.11

$$\int (bx+cx^2)^{4/3} dx = \frac{3bx^2\sqrt[3]{x(b+cx)} \text{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{7}{3}, \frac{10}{3}, -\frac{cx}{b}\right)}{7\sqrt[3]{1+\frac{cx}{b}}}$$

[In] Integrate[(b\*x + c\*x^2)^(4/3), x]

[Out]  $(3*b*x^2*(x*(b + c*x))^{(1/3)}*Hypergeometric2F1[-4/3, 7/3, 10/3, -((c*x)/b)])/(7*(1 + (c*x)/b)^{(1/3)})$

### Maple [F]

$$\int (cx^2 + bx)^{\frac{4}{3}} dx$$

[In] `int((c*x^2+b*x)^(4/3),x)`

[Out] `int((c*x^2+b*x)^(4/3),x)`

### Fricas [F]

$$\int (bx + cx^2)^{4/3} dx = \int (cx^2 + bx)^{\frac{4}{3}} dx$$

[In] `integrate((c*x^2+b*x)^(4/3),x, algorithm="fricas")`

[Out] `integral((c*x^2 + b*x)^(4/3), x)`

### Sympy [F]

$$\int (bx + cx^2)^{4/3} dx = \int (bx + cx^2)^{\frac{4}{3}} dx$$

[In] `integrate((c*x**2+b*x)**(4/3),x)`

[Out] `Integral((b*x + c*x**2)**(4/3), x)`

### Maxima [F]

$$\int (bx + cx^2)^{4/3} dx = \int (cx^2 + bx)^{\frac{4}{3}} dx$$

[In] `integrate((c*x^2+b*x)^(4/3),x, algorithm="maxima")`

[Out] `integrate((c*x^2 + b*x)^(4/3), x)`

**Giac [F]**

$$\int (bx + cx^2)^{4/3} dx = \int (cx^2 + bx)^{\frac{4}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(4/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(4/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.08

$$\int (bx + cx^2)^{4/3} dx = \frac{3x(cx^2 + bx)^{4/3} {}_2F_1\left(-\frac{4}{3}, \frac{7}{3}; \frac{10}{3}; -\frac{cx}{b}\right)}{7\left(\frac{cx}{b} + 1\right)^{4/3}}$$

[In] int((b\*x + c\*x^2)^(4/3),x)

[Out] (3\*x\*(b\*x + c\*x^2)^(4/3)\*hypergeom([-4/3, 7/3], 10/3, -(c\*x)/b))/(7\*((c\*x)/b + 1)^(4/3))

### 3.31 $\int \sqrt[3]{bx + cx^2} dx$

Optimal result	198
Rubi [A] (verified)	199
Mathematica [C] (verified)	201
Maple [F]	202
Fricas [F]	202
Sympy [F]	202
Maxima [F]	202
Giac [F]	203
Mupad [B] (verification not implemented)	203

#### Optimal result

Integrand size = 13, antiderivative size = 387

$$\int \sqrt[3]{bx + cx^2} dx = \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)\sqrt[3]{bx+cx^2}}{10c\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}}$$

$$+ \frac{3^{3/4}\sqrt{2-\sqrt{3}}b^2\sqrt[3]{bx+cx^2}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}{5\cdot 2^{2/3}c(b+2cx)\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}\sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}$$

```
[Out] 3/10*(-c*x*(c*x+b)/b^2)^(1/3)*(2*c*x+b)*(c*x^2+b*x)^(1/3)/c/(-c*(c*x^2+b*x)
/b^2)^(1/3)+1/10*3^(3/4)*b^2*(c*x^2+b*x)^(1/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2
)^(1/3))*EllipticF((1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)*
(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)),2*I-I*3^(1/2))*(1/2*6^(1/2)-1/2*2^(1/2))*
((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(c*x+b)/b^2)^(2/3))/(1
-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)*2^(1/3)/c/(2*c*x+b)/(-c
*(c*x^2+b*x)/b^2)^(1/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))/(1-2^(2/3)*
(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 387, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {636, 633, 201, 242, 225}

$$\int \sqrt[3]{bx + cx^2} dx$$

$$= \frac{3^{3/4} \sqrt{2 - \sqrt{3}} b^{2/3} \sqrt[3]{bx + cx^2} \left( 1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right) \sqrt{\frac{2^3 \sqrt{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} \operatorname{EllipticF}}{5 \cdot 2^{2/3} c (b + 2cx) \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}}$$

$$+ \frac{3 \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (b + 2cx) \sqrt[3]{bx + cx^2}}{10c \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}}$$

[In] Int[(b\*x + c\*x^2)^(1/3), x]

[Out] (3\*((c\*x\*(b + c\*x))/b^2))^(1/3)\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(1/3)/(10\*c\*((c\*(b\*x + c\*x^2))/b^2))^(1/3) + (3^(3/4)\*Sqrt[2 - Sqrt[3]]\*b^2\*(b\*x + c\*x^2)^(1/3)\*(1 - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3)\*Sqrt[(1 + 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*((c\*x\*(b + c\*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3))^2\*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3)], -7 + 4\*Sqrt[3]]/(5\*2^(2/3)\*c\*(b + 2\*c\*x)\*((c\*(b\*x + c\*x^2))/b^2))^(1/3)\*Sqrt[-((1 - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)\*((c\*x\*(b + c\*x))/b^2))^(1/3))^2])

Rule 201

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[x\*((a + b\*x^n)^p/(n\*p + 1)), x] + Dist[a\*n\*(p/(n\*p + 1)), Int[(a + b\*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2\*p] || (EqQ[n, 2] && IntegerQ[4\*p]) || (EqQ[n, 2] && IntegerQ[3\*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 225

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2\*Sqrt[2 - Sqrt[3]]\*(s + r\*x)\*(Sqrt[(s^2 - r\*s

```
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*(s + r*x)/((1 - Sqrt[3])*s + r*x)^2]))*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

### Rule 242

```
Int[((a_) + (b_.)*(x_)^2)^(-2/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x))
, Subst[Int[1/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

### Rule 636

```
Int[((b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[(b*x + c*x^2)^p/((-
c)*((b*x + c*x^2)/b^2))^p, Int[((-c)*(x/b) - c^2*(x^2/b^2))^p, x], x] /; Fr
eeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt[3]{bx + cx^2} \int \sqrt[3]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}} dx}{\sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}} \\ &= -\frac{\left(b^2 \sqrt[3]{bx + cx^2}\right) \text{Subst}\left(\int \sqrt[3]{1 - \frac{b^2x^2}{c^2}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{2^{2/3} c^2 \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}} \\ &= \frac{3 \sqrt[3]{-\frac{cx(b + cx)}{b^2}} (b + 2cx) \sqrt[3]{bx + cx^2}}{10c \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}} - \frac{\left(b^2 \sqrt[3]{bx + cx^2}\right) \text{Subst}\left(\int \frac{1}{\left(1 - \frac{b^2x^2}{c^2}\right)^{2/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{5^{2/3} c^2 \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}} \end{aligned}$$



$$\begin{aligned}
&= \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)\sqrt[3]{bx+cx^2}}{10c\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}} \\
&\quad + \frac{\left(3\sqrt[3]{bx+cx^2}\sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3}\sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{10 \cdot 2^{2/3} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}} \\
&= \frac{3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(b+2cx)\sqrt[3]{bx+cx^2}}{10c\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}} \\
&\quad + \frac{3^{3/4}\sqrt{2-\sqrt{3}b^2}\sqrt[3]{bx+cx^2}\sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}{5 \cdot 2^{2/3}c(b+2cx)\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}\sqrt{-1-\frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.12

$$\int \sqrt[3]{bx+cx^2} dx = \frac{3x\sqrt[3]{x(b+cx)} \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{4}{3}, \frac{7}{3}, -\frac{cx}{b}\right)}{4\sqrt[3]{1+\frac{cx}{b}}}$$

[In] Integrate[(b\*x + c\*x^2)^(1/3),x]

[Out] (3\*x\*(x\*(b + c\*x))^(1/3)\*Hypergeometric2F1[-1/3, 4/3, 7/3, -((c\*x)/b)])/(4\*(1 + (c\*x)/b)^(1/3))

**Maple [F]**

$$\int (cx^2 + bx)^{\frac{1}{3}} dx$$

[In] int((c\*x^2+b\*x)^(1/3),x)

[Out] int((c\*x^2+b\*x)^(1/3),x)

**Fricas [F]**

$$\int \sqrt[3]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(1/3), x)

**Sympy [F]**

$$\int \sqrt[3]{bx + cx^2} dx = \int \sqrt[3]{bx + cx^2} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(1/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(1/3), x)

**Maxima [F]**

$$\int \sqrt[3]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(1/3), x)

**Giac [F]**

$$\int \sqrt[3]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(1/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.09

$$\int \sqrt[3]{bx + cx^2} dx = \frac{3x (cx^2 + bx)^{1/3} {}_2F_1\left(-\frac{1}{3}, \frac{4}{3}; \frac{7}{3}; -\frac{cx}{b}\right)}{4\left(\frac{cx}{b} + 1\right)^{1/3}}$$

[In] int((b\*x + c\*x^2)^(1/3),x)

[Out] (3\*x\*(b\*x + c\*x^2)^(1/3)\*hypergeom([-1/3, 4/3], 7/3, -(c\*x)/b))/(4\*((c\*x)/b + 1)^(1/3))

### 3.32 $\int \frac{1}{(bx+cx^2)^{2/3}} dx$

Optimal result	204
Rubi [A] (verified)	205
Mathematica [C] (verified)	206
Maple [F]	207
Fricas [F]	207
Sympy [F]	207
Maxima [F]	207
Giac [F]	208
Mupad [B] (verification not implemented)	208

#### Optimal result

Integrand size = 13, antiderivative size = 322

$$\int \frac{1}{(bx+cx^2)^{2/3}} dx = \frac{\sqrt[3]{2}3^{3/4}\sqrt{2-\sqrt{3}}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{c}{b^2}}\right)}}}{c(b+2cx)(bx+cx^2)^{2/3}\sqrt{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{c}{b^2}}\right)}}$$

```
[Out] 2^(1/3)*3^(3/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(2/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*EllipticF((1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)),2*I-I*3^(1/2))*(1/2*6^(1/2)-1/2*2^(1/2))*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(c*x+b)/b^2)^(2/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)/c/(2*c*x+b)/(c*x^2+b*x)^(2/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)
```

**Rubi [A] (verified)**

Time = 0.30 (sec) , antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {636, 633, 242, 225}

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \frac{\sqrt[3]{2} 3^{3/4} \sqrt{2 - \sqrt{3}} b^2 \left(-\frac{c(bx + cx^2)}{b^2}\right)^{2/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right) \sqrt{\frac{2 \sqrt[3]{2} \left(-\frac{cx(b + cx)}{b^2}\right)^{2/3} + 2^{2/3}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}{c(b + 2cx)(bx + cx^2)^{2/3} \sqrt{\left(-\frac{cx(b + cx)}{b^2}\right)^2}}$$

[In] Int[(b\*x + c\*x^2)^(-2/3), x]

[Out] (2^(1/3)\*3^(3/4)\*Sqrt[2 - Sqrt[3]]\*b^2\*(-((c\*(b\*x + c\*x^2))/b^2))^(2/3)\*(1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))\*Sqrt[(1 + 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-((c\*x\*(b + c\*x))/b^2))^(2/3))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))]^2\*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))], -7 + 4\*Sqrt[3]]/(c\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(2/3)\*Sqrt[-((1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))^(2/3))])

Rule 225

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2\*Sqrt[2 - Sqrt[3]]\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/((1 - Sqrt[3])\*s + r\*x)^2]/(3^(1/4)\*r\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])\*EllipticF[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b}, x] && NegQ[a]

Rule 242

Int[((a\_) + (b\_.)\*(x\_)^2)^(-2/3), x\_Symbol] := Dist[3\*(Sqrt[b\*x^2]/(2\*b\*x)), Subst[Int[1/Sqrt[-a + x^3], x], x, (a + b\*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2)^p, Int[(-c)\*(x/b) - c^2\*(x^2/b^2)]^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \int \frac{1}{\left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{2/3}} dx}{(bx + cx^2)^{2/3}} \\
 &= -\frac{\left(\sqrt[3]{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{2/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{c^2(bx + cx^2)^{2/3}} \\
 &= \frac{\left(3\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{2^{2/3}\left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right)(bx + cx^2)^{2/3}} \\
 &= \frac{\sqrt[3]{23}^{3/4} \sqrt{2 - \sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}}}}{c(b+2cx)(bx + cx^2)^{2/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}} \sqrt{\left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}
 \end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.13

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \frac{3x\left(1 + \frac{cx}{b}\right)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -\frac{cx}{b}\right)}{(x(b + cx))^{2/3}}$$

[In] Integrate[(b\*x + c\*x^2)^(-2/3), x]

[Out] (3\*x\*(1 + (c\*x)/b)^(2/3)\*Hypergeometric2F1[1/3, 2/3, 4/3, -(c\*x)/b])/(x\*(b + c\*x))^(2/3)

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{2}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(2/3),x)

[Out] int(1/(c\*x^2+b\*x)^(2/3),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{2}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(2/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(-2/3), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \int \frac{1}{(bx + cx^2)^{\frac{2}{3}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(2/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-2/3), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{2}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(2/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-2/3), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \int \frac{1}{(cx^2 + bx)^{2/3}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(2/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-2/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.11

$$\int \frac{1}{(bx + cx^2)^{2/3}} dx = \frac{3x \left(\frac{cx}{b} + 1\right)^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; \frac{4}{3}; -\frac{cx}{b}\right)}{(cx^2 + bx)^{2/3}}$$

[In] int(1/(b\*x + c\*x^2)^(2/3),x)

[Out] (3\*x\*((c\*x)/b + 1)^(2/3)\*hypergeom([1/3, 2/3], 4/3, -(c\*x)/b))/(b\*x + c\*x^2)^(2/3)



### 3.33 $\int \frac{1}{(bx+cx^2)^{5/3}} dx$

Optimal result	209
Rubi [A] (verified)	210
Mathematica [C] (verified)	212
Maple [F]	213
Fricas [F]	213
Sympy [F]	213
Maxima [F]	213
Giac [F]	214
Mupad [B] (verification not implemented)	214

#### Optimal result

Integrand size = 13, antiderivative size = 384

$$\int \frac{1}{(bx+cx^2)^{5/3}} dx = \frac{3(b+2cx) \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3}}{2c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{5/3}}$$

$$+ \frac{\sqrt[3]{23}^{3/4} \sqrt{2-\sqrt{3}} b^2 \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3} \left(1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}{c(b+2cx)(bx+cx^2)^{5/3} \sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}$$

```
[Out] 3/2*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(5/3)/c/(-c*x*(c*x+b)/b^2)^(2/3)/(c*x^2+
b*x)^(5/3)+2^(1/3)*3^(3/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(5/3)*(1-2^(2/3)*(-c*x*
(c*x+b)/b^2)^(1/3))*EllipticF((1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/
(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)),2*I-I*3^(1/2))*(1/2*6^(1/2)-1/
2*2^(1/2))*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(c*x+b)/b^2
)^(2/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)/c/(2*c*x+b)/
(c*x^2+b*x)^(5/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))/(1-2^(2/3)*(-c*x*(
c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)
```

**Rubi [A] (verified)**

Time = 0.33 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {636, 633, 205, 242, 225}

$$\int \frac{1}{(bx + cx^2)^{5/3}} dx = \frac{\sqrt[3]{2} 3^{3/4} \sqrt{2 - \sqrt{3}} b^2 \left(-\frac{c(bx + cx^2)}{b^2}\right)^{5/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt{2} \left(-\frac{cx(b + cx)}{b^2}\right)^{2/3} + 2^{2/3}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}{c(b + 2cx)(bx + cx^2)^{5/3} \sqrt{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}} + \frac{3(b + 2cx) \left(-\frac{c(bx + cx^2)}{b^2}\right)^{5/3}}{2c \left(-\frac{cx(b + cx)}{b^2}\right)^{2/3} (bx + cx^2)^{5/3}}$$

[In] Int[(b\*x + c\*x^2)^(-5/3), x]

[Out] (3\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(5/3)/(2\*c\*(-((c\*x\*(b + c\*x))/b^2))^(2/3)\*(b\*x + c\*x^2)^(5/3)) + (2^(1/3)\*3^(3/4)\*Sqrt[2 - Sqrt[3]]\*b^2\*(-(c\*(b\*x + c\*x^2))/b^2))^(5/3)\*(1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)\*Sqrt[(1 + 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-(c\*x\*(b + c\*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2]\*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)], -7 + 4\*Sqrt[3]]/(c\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(5/3)\*Sqrt[-((1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2])

Rule 205

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[(-x)\*((a + b\*x^n)^(p + 1)/(a\*n\*(p + 1))), x] + Dist[(n\*(p + 1) + 1)/(a\*n\*(p + 1)), Int[(a + b\*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2\*p] || (n == 2 && IntegerQ[4\*p]) || (n == 2 && IntegerQ[3\*p]) || Denominator[p + 1/n] < Denominator[p])

Rule 225

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2\*Sqrt[2 - Sqrt[3]]\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)]/((1 - Sqrt[3])\*s + r\*x)^2)/(3^(1/4)\*r\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])]\*EllipticF[ArcSin[((1 + Sqrt[3])

\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x]] /; FreeQ[{a, b}, x] && NegQ[a]

### Rule 242

Int[((a\_) + (b\_.)\*(x\_)^2)^(-2/3), x\_Symbol] := Dist[3\*(Sqrt[b\*x^2]/(2\*b\*x)), Subst[Int[1/Sqrt[-a + x^3], x], x, (a + b\*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c)], x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \int \frac{1}{\left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{5/3}} dx}{(bx+cx^2)^{5/3}} \\
 &= -\frac{\left(4\sqrt[3]{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{5/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{5/3}} \\
 &= \frac{3(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}}{2c\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}(bx+cx^2)^{5/3}} \\
 &= -\frac{\left(\sqrt[3]{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{2/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{5/3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3}}{2c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{5/3}} \\
&\quad + \frac{\left(3 \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{2^{2/3} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{5/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3}}{2c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{5/3}} \\
&\quad + \frac{\sqrt[3]{23^{3/4}} \sqrt{2 - \sqrt{3}b^2} \left(-\frac{c(b+cx^2)}{b^2}\right)^{5/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx}{b^2}}}{1-\sqrt{3}-\dots}}}{c(b+2cx) (bx+cx^2)^{5/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.12

$$\int \frac{1}{(bx+cx^2)^{5/3}} dx = -\frac{3\left(1+\frac{cx}{b}\right)^{2/3} \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{5}{3}, \frac{1}{3}, -\frac{cx}{b}\right)}{2b(x(b+cx))^{2/3}}$$

[In] Integrate[(b\*x + c\*x^2)^(-5/3),x]

[Out] (-3\*(1 + (c\*x)/b)^(2/3)\*Hypergeometric2F1[-2/3, 5/3, 1/3, -(c\*x)/b])/(2\*b\*(x\*(b + c\*x))^(2/3))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{5}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(5/3),x)

[Out] int(1/(c\*x^2+b\*x)^(5/3),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{5}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{5}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(1/3)/(c^2\*x^4 + 2\*b\*c\*x^3 + b^2\*x^2), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{5}{3}}} dx = \int \frac{1}{(bx + cx^2)^{\frac{5}{3}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(5/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-5/3), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{5}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{5}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-5/3), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{5/3}} dx = \int \frac{1}{(cx^2 + bx)^{5/3}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-5/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.09

$$\int \frac{1}{(bx + cx^2)^{5/3}} dx = -\frac{3x \left(\frac{cx}{b} + 1\right)^{5/3} {}_2F_1\left(-\frac{2}{3}, \frac{5}{3}; \frac{1}{3}; -\frac{cx}{b}\right)}{2(cx^2 + bx)^{5/3}}$$

[In] int(1/(b\*x + c\*x^2)^(5/3),x)

[Out] -(3\*x\*((c\*x)/b + 1)^(5/3)\*hypergeom([-2/3, 5/3], 1/3, -(c\*x)/b))/(2\*(b\*x + c\*x^2)^(5/3))

### 3.34 $\int \frac{1}{(bx+cx^2)^{8/3}} dx$

Optimal result	215
Rubi [A] (verified)	216
Mathematica [C] (verified)	218
Maple [F]	219
Fricas [F]	219
Sympy [F]	219
Maxima [F]	219
Giac [F]	220
Mupad [B] (verification not implemented)	220

#### Optimal result

Integrand size = 13, antiderivative size = 448

$$\int \frac{1}{(bx+cx^2)^{8/3}} dx = \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (bx+cx^2)^{8/3}} + \frac{21(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{8/3}}$$

$$+ \frac{14\sqrt[3]{23}^{3/4} \sqrt{2-\sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3} \left(1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2\sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}{5c(b+2cx)(bx+cx^2)^{8/3} \sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}$$

```
[Out] 3/5*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(8/3)/c/(-c*x*(c*x+b)/b^2)^(5/3)/(c*x^2+b*x)^(8/3)+21/5*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(8/3)/c/(-c*x*(c*x+b)/b^2)^(2/3)/(c*x^2+b*x)^(8/3)+14/5*2^(1/3)*3^(3/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(8/3)*(1-2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)*EllipticF((1-2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)),2*I-I*3^(1/2))*(1/2*6^(1/2)-1/2*2^(1/2))*((1+2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3))*(-c*x*(c*x+b)/b^2)^(2/3))/(1-2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)/c/(2*c*x+b)/(c*x^2+b*x)^(8/3)/((-1+2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3))/(1-2^(2/3))*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)
```

**Rubi [A] (verified)**

Time = 0.38 (sec) , antiderivative size = 448, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.385$ , Rules used = {636, 633, 205, 242, 225}

$$\int \frac{1}{(bx + cx^2)^{8/3}} dx = \frac{14\sqrt[3]{23}^{3/4} \sqrt{2 - \sqrt{3}} b^2 \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}}{5c(b+2cx)(bx+cx^2)^{8/3}} + \frac{21(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{8/3}} + \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (bx+cx^2)^{8/3}}$$

[In] Int[(b\*x + c\*x^2)^(-8/3), x]

[Out] (3\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(8/3)/(5\*c\*(-((c\*x\*(b + c\*x))/b^2))^(5/3)\*(b\*x + c\*x^2)^(8/3)) + (21\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(8/3)/(5\*c\*(-((c\*x\*(b + c\*x))/b^2))^(2/3)\*(b\*x + c\*x^2)^(8/3)) + (14\*2^(1/3)\*3^(3/4)\*Sqrt[2 - Sqrt[3]]\*b^2\*(-((c\*(b\*x + c\*x^2))/b^2))^(8/3)\*(1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)\*Sqrt[(1 + 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-(c\*x\*(b + c\*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2\*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)], -7 + 4\*Sqrt[3]]/(5\*c\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(8/3)\*Sqrt[-((1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2])

**Rule 205**

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[(-x)\*((a + b\*x^n)^(p + 1)/(a\*n\*(p + 1))), x] + Dist[(n\*(p + 1) + 1)/(a\*n\*(p + 1)), Int[(a + b\*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2\*p] || (n == 2 && IntegerQ[4\*p]) || (n == 2 && IntegerQ[3\*p]) || Denominator[p + 1/n] < Denominator[p])

**Rule 225**

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2\*Sqrt[2 - Sqrt[3]]\*(s + r\*x)\*(Sqrt[(s^2 - r\*s



\*x + r^2\*x^2)/((1 - Sqrt[3])\*s + r\*x)^2)/(3^(1/4)\*r\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)))\*EllipticF[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b}, x] && NegQ[a]

### Rule 242

Int[((a\_) + (b\_.)\*(x\_)^2)^(-2/3), x\_Symbol] := Dist[3\*(Sqrt[b\*x^2]/(2\*b\*x)), Subst[Int[1/Sqrt[-a + x^3], x], x, (a + b\*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c)], x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3} \int \frac{1}{\left(-\frac{cx}{b}-\frac{c^2x^2}{b^2}\right)^{8/3}} dx}{(bx+cx^2)^{8/3}} \\
 &= -\frac{\left(16\sqrt[3]{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{8/3}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{8/3}} \\
 &= \frac{3(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c\left(-\frac{cx(b+cx)}{b^2}\right)^{5/3}(bx+cx^2)^{8/3}} \\
 &\quad - \frac{\left(56\sqrt[3]{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{5/3}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{5c^2(bx+cx^2)^{8/3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (bx+cx^2)^{8/3}} + \frac{21(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{8/3}} \\
&\quad - \frac{\left(14\sqrt[3]{2}b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{2/3}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{5c^2 (bx+cx^2)^{8/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (bx+cx^2)^{8/3}} + \frac{21(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{8/3}} \\
&\quad + \frac{\left(21\sqrt[3]{2} \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{5 \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{8/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (bx+cx^2)^{8/3}} + \frac{21(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3}}{5c \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (bx+cx^2)^{8/3}} \\
&\quad + \frac{14\sqrt[3]{2}3^{3/4} \sqrt{2-\sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{8/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}}}{5c(b+2cx) (bx+cx^2)^{8/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.11

$$\int \frac{1}{(bx+cx^2)^{8/3}} dx = -\frac{3\left(1+\frac{cx}{b}\right)^{2/3} \text{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{8}{3}, -\frac{2}{3}, -\frac{cx}{b}\right)}{5b^2x(x(b+cx))^{2/3}}$$

[In] Integrate[(b\*x + c\*x^2)^(-8/3), x]

[Out] (-3\*(1 + (c\*x)/b)^(2/3)\*Hypergeometric2F1[-5/3, 8/3, -2/3, -(c\*x)/b])/(5\*b^2\*x\*(x\*(b + c\*x))^(2/3))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{8}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(8/3),x)

[Out] int(1/(c\*x^2+b\*x)^(8/3),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{8}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{8}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(8/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(1/3)/(c^3\*x^6 + 3\*b\*c^2\*x^5 + 3\*b^2\*c\*x^4 + b^3\*x^3), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{8}{3}}} dx = \int \frac{1}{(bx + cx^2)^{\frac{8}{3}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(8/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-8/3), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{8}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{8}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(8/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-8/3), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{8/3}} dx = \int \frac{1}{(cx^2 + bx)^{8/3}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(8/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-8/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.06 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.08

$$\int \frac{1}{(bx + cx^2)^{8/3}} dx = -\frac{3x \left(\frac{cx}{b} + 1\right)^{8/3} {}_2F_1\left(-\frac{5}{3}, \frac{8}{3}; -\frac{2}{3}; -\frac{cx}{b}\right)}{5 (cx^2 + bx)^{8/3}}$$

[In] int(1/(b\*x + c\*x^2)^(8/3),x)

[Out] -(3\*x\*((c\*x)/b + 1)^(8/3)\*hypergeom([-5/3, 8/3], -2/3, -(c\*x)/b))/(5\*(b\*x + c\*x^2)^(8/3))

### 3.35 $\int (bx + cx^2)^{5/3} dx$

Optimal result . . . . .	222
Rubi [A] (verified) . . . . .	223
Mathematica [C] (verified) . . . . .	228
Maple [F] . . . . .	229
Fricas [F] . . . . .	229
Sympy [F] . . . . .	229
Maxima [F] . . . . .	229
Giac [F] . . . . .	230
Mupad [B] (verification not implemented) . . . . .	230

## Optimal result

Integrand size = 13, antiderivative size = 842

$$\begin{aligned}
 \int (bx + cx^2)^{5/3} dx &= \frac{15 \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx) (bx+cx^2)^{5/3}}{364c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &+ \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b+2cx) (bx+cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &- \frac{15(b+2cx) (bx+cx^2)^{5/3}}{182\sqrt[3]{2}c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
 &- \frac{15\sqrt[4]{3}\sqrt{2+\sqrt{3}}b^2(bx+cx^2)^{5/3} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} E\left(\arcsin\left(\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}\right)\right)}{364\sqrt[3]{2}c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}} \\
 &+ \frac{5\sqrt[3]{4}b^2(bx+cx^2)^{5/3} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} \text{EllipticF}\left(\arcsin\left(\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}\right)\right)}{91\sqrt[5]{2}c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}
 \end{aligned}$$

[Out] 15/364\*(-c\*x\*(c\*x+b)/b^2)^(2/3)\*(2\*c\*x+b)\*(c\*x^2+b\*x)^(5/3)/c/(-c\*(c\*x^2+b\*x)/b^2)^(5/3)+3/26\*(-c\*x\*(c\*x+b)/b^2)^(5/3)\*(2\*c\*x+b)\*(c\*x^2+b\*x)^(5/3)/c/(-c\*(c\*x^2+b\*x)/b^2)^(5/3)-15/364\*(2\*c\*x+b)\*(c\*x^2+b\*x)^(5/3)\*2^(2/3)/c/(-c\*(c\*x^2+b\*x)/b^2)^(5/3)/(1-2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2))+5/182\*3^(3/4)\*b^2\*(c\*x^2+b\*x)^(5/3)\*(1-2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3))\*EllipticF((1-2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2)),2\*I-I\*3^(1/2))\*((1+2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3)+2\*2^(1/3)\*(-c\*x\*(c\*x+b)/b^2)^(2/3))/(1-2^(2/3)\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2))^2

$$\begin{aligned} &)^{(1/2)} * 2^{(1/6)} / c / (2 * c * x + b) / (-c * (c * x^2 + b * x) / b^2)^{(5/3)} / ((-1 + 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)}) / (1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)})^2)^{(1/2)} - 15 / 728 * 3^{(1/4)} * b^2 * (c * x^2 + b * x)^{(5/3)} * (1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)}) * \text{EllipticE}((1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} + 3^{(1/2)}) / (1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)}), 2 * I - I * 3^{(1/2)}) * ((1 + 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} + 2 * 2^{(1/3)} * (-c * x * (c * x + b) / b^2)^{(2/3)}) / (1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)})^2)^{(1/2)} * (1/2 * 6^{(1/2)} + 1/2 * 2^{(1/2)}) * 2^{(2/3)} / c / (2 * c * x + b) / (-c * (c * x^2 + b * x) / b^2)^{(5/3)} / ((-1 + 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)}) / (1 - 2^{(2/3)}) * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)})^2)^{(1/2)} \end{aligned}$$

### Rubi [A] (verified)

Time = 0.78 (sec) , antiderivative size = 842, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used

= {636, 633, 201, 241, 310, 225, 1893}

$$\int (bx + cx^2)^{5/3} dx =$$

$$\frac{15\sqrt[4]{3}\sqrt{2+\sqrt{3}}(cx^2+bx)^{5/3}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+1}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}E\left(\arcsin\left(\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1}\right)\right)}{364\sqrt[3]{2}c(b+2cx)\left(-\frac{c(cx^2+bx)}{b^2}\right)^{5/3}\sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}$$

$$+ \frac{5\sqrt[3]{3/4}(cx^2+bx)^{5/3}\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{2\sqrt[3]{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+1}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}\text{EllipticF}\left(\arcsin\left(\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1}\right)\right)}{91\sqrt[5]{2^6}c(b+2cx)\left(-\frac{c(cx^2+bx)}{b^2}\right)^{5/3}\sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}$$

$$- \frac{15(b+2cx)(cx^2+bx)^{5/3}}{182\sqrt[3]{2}c\left(-\frac{c(cx^2+bx)}{b^2}\right)^{5/3}\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)}$$

$$+ \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{5/3}(b+2cx)(cx^2+bx)^{5/3}}{26c\left(-\frac{c(cx^2+bx)}{b^2}\right)^{5/3}} + \frac{15\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}(b+2cx)(cx^2+bx)^{5/3}}{364c\left(-\frac{c(cx^2+bx)}{b^2}\right)^{5/3}}$$

[In] Int[(b\*x + c\*x^2)^(5/3), x]

[Out] (15\*(-((c\*x\*(b + c\*x))/b^2))^(2/3)\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(5/3))/(364\*c\*(-((c\*(b\*x + c\*x^2))/b^2))^(5/3)) + (3\*(-((c\*x\*(b + c\*x))/b^2))^(5/3)\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(5/3))/(26\*c\*(-((c\*(b\*x + c\*x^2))/b^2))^(5/3)) - (15\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(5/3))/(182\*2^(1/3)\*c\*(-((c\*(b\*x + c\*x^2))/b^2))^(5/3)\*(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))) - (15\*3^(1/4)\*Sqrt[2 + Sqrt[3]]\*b^2\*(b\*x + c\*x^2)^(5/3)\*(1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))\*Sqrt[(1 + 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-((c\*x\*(b + c\*x))/b^2))^(2/3))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))]^2\*EllipticE[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))], -7 + 4\*S



```

qrt[3]]/(364*2^(1/3)*c*(b + 2*c*x)*(-((c*(b*x + c*x^2))/b^2))^(5/3)*Sqrt[-
((1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*
x*(b + c*x))/b^2))^(1/3))^2)] + (5*3^(3/4)*b^2*(b*x + c*x^2)^(5/3)*(1 - 2^
(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))*Sqrt[(1 + 2^(2/3)*(-((c*x*(b + c*x))/
b^2))^(1/3) + 2*2^(1/3)*(-((c*x*(b + c*x))/b^2))^(2/3))/(1 - Sqrt[3] - 2^(2
/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2]*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2
/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x
))/b^2))^(1/3)], -7 + 4*Sqrt[3]]]/(91*2^(5/6)*c*(b + 2*c*x)*(-((c*(b*x + c
*x^2))/b^2))^(5/3)*Sqrt[-((1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1 -
Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2)]

```

#### Rule 201

```

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])

```

#### Rule 225

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2)])]*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x
] && NegQ[a]

```

#### Rule 241

```

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x))
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]

```

#### Rule 310

```

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(-1 + Sqrt[3])*(s/r), Int[1/Sqrt[a + b*x^
3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]
/; FreeQ[{a, b}, x] && NegQ[a]

```

#### Rule 633

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

```

## Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[(-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

## Rule 1893

Int[((c\_) + (d\_.)\*(x\_))/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Simplify[(1 + Sqrt[3])\*(d/c)], s = Denom[Simplify[(1 + Sqrt[3])\*(d/c)]]], Simp[2\*d\*s^3\*(Sqrt[a + b\*x^3]/(a\*r^2\*((1 - Sqrt[3])\*s + r\*x))), x] + Simp[3^(1/4)\*Sqrt[2 + Sqrt[3]]\*d\*s\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/(1 - Sqrt[3])\*s + r\*x]^2)/(r^2\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])]\*EllipticE[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b\*c^3 - 2\*(5 + 3\*Sqrt[3])\*a\*d^3, 0]

## Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(bx + cx^2)^{5/3} \int \left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{5/3} dx}{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &= -\frac{\left(b^2(bx + cx^2)^{5/3}\right) \text{Subst}\left(\int \left(1 - \frac{b^2x^2}{c^2}\right)^{5/3} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{16\sqrt[3]{2}c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &= \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b + 2cx) (bx + cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &\quad - \frac{\left(5b^2(bx + cx^2)^{5/3}\right) \text{Subst}\left(\int \left(1 - \frac{b^2x^2}{c^2}\right)^{2/3} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{104\sqrt[3]{2}c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &= \frac{15\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b + 2cx) (bx + cx^2)^{5/3}}{364c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} + \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b + 2cx) (bx + cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
 &\quad - \frac{\left(5b^2(bx + cx^2)^{5/3}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{1 - \frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{182\sqrt[3]{2}c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{15 \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{5/3}}{364c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} + \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b+2cx)(bx+cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
&\quad + \frac{\left(15(bx+cx^2)^{5/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}} \right)}{364\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
&= \frac{15 \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{5/3}}{364c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} + \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b+2cx)(bx+cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
&\quad - \frac{\left(15(bx+cx^2)^{5/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1+\sqrt{3}-x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}} \right)}{364\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
&\quad + \frac{\left(15(1+\sqrt{3})(bx+cx^2)^{5/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}} \right)}{364\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{15 \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{5/3}}{364c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} + \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{5/3} (b+2cx)(bx+cx^2)^{5/3}}{26c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3}} \\
&+ \frac{15b^2(bx+cx^2)^{5/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}}{182\sqrt[3]{2}c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
&- \frac{15\sqrt[4]{3}\sqrt{2+\sqrt{3}}b^2(bx+cx^2)^{5/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}}}}{364\sqrt[3]{2}c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}} \\
&+ \frac{5\sqrt[3]{4}b^2(bx+cx^2)^{5/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}+2\sqrt[3]{2}}{1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}}}{91\sqrt[5]{6}c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{5/3} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.06

$$\int (bx+cx^2)^{5/3} dx = \frac{3bx^2(x(b+cx))^{2/3} \text{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{8}{3}, \frac{11}{3}, -\frac{cx}{b}\right)}{8\left(1+\frac{cx}{b}\right)^{2/3}}$$

[In] Integrate[(b\*x + c\*x^2)^(5/3),x]

[Out] (3\*b\*x^2\*(x\*(b + c\*x))^(2/3)\*Hypergeometric2F1[-5/3, 8/3, 11/3, -(c\*x)/b])/ (8\*(1 + (c\*x)/b)^(2/3))

**Maple [F]**

$$\int (cx^2 + bx)^{\frac{5}{3}} dx$$

[In] int((c\*x^2+b\*x)^(5/3),x)

[Out] int((c\*x^2+b\*x)^(5/3),x)

**Fricas [F]**

$$\int (bx + cx^2)^{5/3} dx = \int (cx^2 + bx)^{\frac{5}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(5/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(5/3), x)

**Sympy [F]**

$$\int (bx + cx^2)^{5/3} dx = \int (bx + cx^2)^{\frac{5}{3}} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(5/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(5/3), x)

**Maxima [F]**

$$\int (bx + cx^2)^{5/3} dx = \int (cx^2 + bx)^{\frac{5}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(5/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(5/3), x)

**Giac [F]**

$$\int (bx + cx^2)^{5/3} dx = \int (cx^2 + bx)^{5/3} dx$$

[In] integrate((c\*x^2+b\*x)^(5/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(5/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.04

$$\int (bx + cx^2)^{5/3} dx = \frac{3x(cx^2 + bx)^{5/3} {}_2F_1\left(-\frac{5}{3}, \frac{8}{3}; \frac{11}{3}; -\frac{cx}{b}\right)}{8\left(\frac{cx}{b} + 1\right)^{5/3}}$$

[In] int((b\*x + c\*x^2)^(5/3),x)

[Out] (3\*x\*(b\*x + c\*x^2)^(5/3)\*hypergeom([-5/3, 8/3], 11/3, -(c\*x)/b))/(8\*((c\*x)/b + 1)^(5/3))

### 3.36 $\int (bx + cx^2)^{2/3} dx$

Optimal result . . . . .	232
Rubi [A] (verified) . . . . .	233
Mathematica [C] (verified) . . . . .	237
Maple [F] . . . . .	238
Fricas [F] . . . . .	238
Sympy [F] . . . . .	238
Maxima [F] . . . . .	238
Giac [F] . . . . .	239
Mupad [B] (verification not implemented) . . . . .	239

## Optimal result

Integrand size = 13, antiderivative size = 781

$$\int (bx + cx^2)^{2/3} dx = \frac{3 \left( -\frac{cx(b+cx)}{b^2} \right)^{2/3} (b+2cx) (bx+cx^2)^{2/3}}{14c \left( -\frac{c(bx+cx^2)}{b^2} \right)^{2/3}}$$


---


$$\frac{3(b+2cx) (bx+cx^2)^{2/3}}{7\sqrt[3]{2}c \left( -\frac{c(bx+cx^2)}{b^2} \right)^{2/3} \left( 1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right)}$$


---


$$3\sqrt[4]{3}\sqrt{2+\sqrt{3}}b^2(bx+cx^2)^{2/3} \left( 1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2\sqrt[3]{2} \left( -\frac{cx(b+cx)}{b^2} \right)^{2/3}}{\left( 1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right)^2}} E \left( \arcsin \left( \frac{\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}} \right) \right)$$


---


$$14\sqrt[3]{2}c(b+2cx) \left( -\frac{c(bx+cx^2)}{b^2} \right)^{2/3} \sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left( 1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right)^2}}$$


---


$$\sqrt[6]{2}3^{3/4}b^2(bx+cx^2)^{2/3} \left( 1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2\sqrt[3]{2} \left( -\frac{cx(b+cx)}{b^2} \right)^{2/3}}{\left( 1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right)^2}} \text{EllipticF} \left( \arcsin \left( \frac{\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}} \right) \right)$$


---


$$+ \frac{7c(b+2cx) \left( -\frac{c(bx+cx^2)}{b^2} \right)^{2/3}}{\sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left( 1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right)^2}}}$$

```
[Out] 3/14*(-c*x*(c*x+b)/b^2)^(2/3)*(2*c*x+b)*(c*x^2+b*x)^(2/3)/c/(-c*(c*x^2+b*x)
/b^2)^(2/3)-3/14*(2*c*x+b)*(c*x^2+b*x)^(2/3)*2^(2/3)/c/(-c*(c*x^2+b*x)/b^2)
^(2/3)/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))+1/7*2^(1/6)*3^(3/4)*b^2
*(c*x^2+b*x)^(2/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*EllipticF((1-2^(2/3)
)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(
1/2)),2*I-I*3^(1/2))*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(
c*x+b)/b^2)^(2/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)))^(1/2)/c/
(2*c*x+b)/(-c*(c*x^2+b*x)/b^2)^(2/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)
)/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)))^(1/2)-3/28*3^(1/4)*b^2*(c*
x^2+b*x)^(2/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*EllipticE((1-2^(2/3)*(-
c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2)
),2*I-I*3^(1/2))*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(c*x+
```



$$\frac{b}{b^2} \sqrt[2]{\frac{b}{b^2}} \sqrt[2]{\frac{1-2^{2/3}(-c*x*(c*x+b)/b^2)^{1/3}-3^{1/2}}{2}} \sqrt[2]{\frac{1}{2} * \left( \frac{1}{2} * 6^{1/2} + \frac{1}{2} * 2^{1/2} \right) * 2^{2/3} / c / (2*c*x+b) / (-c*(c*x^2+b*x)/b^2)^{2/3} / \left( (-1+2^{2/3}) * (-c*x*(c*x+b)/b^2)^{1/3} \right) / (1-2^{2/3} * (-c*x*(c*x+b)/b^2)^{1/3} - 3^{1/2})^2} \sqrt[2]{\frac{1}{2}}$$

### Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 781, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {636, 633, 201, 241, 310, 225, 1893}

$$\int (bx + cx^2)^{2/3} dx = \frac{\sqrt[6]{2} 3^{3/4} b^2 (bx + cx^2)^{2/3} \left( 1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right) \sqrt{\frac{2^3 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}}{7c(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}}$$

$$+ \frac{3^4 \sqrt[3]{3} \sqrt{2 + \sqrt{3}} b^2 (bx + cx^2)^{2/3} \left( 1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} \right) \sqrt{\frac{2^3 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} E\left(\arcsin\left(\frac{\sqrt{2 + \sqrt{3}} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\sqrt{3}}\right)\right)}{14^3 \sqrt[3]{2} c (b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}}$$

$$+ \frac{3 \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx) (bx+cx^2)^{2/3}}{14c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}}$$

$$- \frac{3(b+2cx) (bx+cx^2)^{2/3}}{7^3 \sqrt[3]{2} c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)}$$

[In] Int[(b\*x + c\*x^2)^(2/3), x]

```
[Out] (3*(-((c*x*(b + c*x))/b^2))^(2/3)*(b + 2*c*x)*(b*x + c*x^2)^(2/3))/(14*c*(-
((c*(b*x + c*x^2))/b^2))^(2/3)) - (3*(b + 2*c*x)*(b*x + c*x^2)^(2/3))/(7*2^
(1/3)*c*(-((c*(b*x + c*x^2))/b^2))^(2/3)*(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b
+ c*x))/b^2))^(1/3))) - (3*3^(1/4)*Sqrt[2 + Sqrt[3]]*b^2*(b*x + c*x^2)^(2/3
)*(1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))*Sqrt[(1 + 2^(2/3)*(-((c*x*(b
+ c*x))/b^2))^(1/3) + 2*2^(1/3)*(-((c*x*(b + c*x))/b^2))^(2/3))/(1 - Sqrt[
3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2]*EllipticE[ArcSin[(1 + Sqrt[
3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x
*(b + c*x))/b^2))^(1/3))], -7 + 4*Sqrt[3]])/(14*2^(1/3)*c*(b + 2*c*x)*(-((c
*(b*x + c*x^2))/b^2))^(2/3)*Sqrt[-((1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1
/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2]) + (2^(1/6)
*3^(3/4)*b^2*(b*x + c*x^2)^(2/3)*(1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3
))*Sqrt[(1 + 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3) + 2*2^(1/3)*(-((c*x*(b
+ c*x))/b^2))^(2/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))
^2]*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))
/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))], -7 + 4*Sqrt[3]])/
(7*c*(b + 2*c*x)*(-((c*(b*x + c*x^2))/b^2))^(2/3)*Sqrt[-((1 - 2^(2/3)*(-((c
*x*(b + c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(
1/3))^2])]
```

#### Rule 201

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])
```

#### Rule 225

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2])))*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x
] && NegQ[a]
```

#### Rule 241

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x))
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]
```

#### Rule 310

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[-(1 + Sqrt[3])*(s/r), Int[1/Sqrt[a + b*x^
```

3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])\*s + r\*x)/Sqrt[a + b\*x^3], x], x]]  
 /; FreeQ[{a, b}, x] && NegQ[a]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rule 1893

Int[((c\_) + (d\_.)\*(x\_))/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Simplify[(1 + Sqrt[3])\*(d/c)], s = Denom[Simplify[(1 + Sqrt[3])\*(d/c)]]], Simp[2\*d\*s^3\*(Sqrt[a + b\*x^3]/(a\*r^2\*((1 - Sqrt[3])\*s + r\*x))), x] + Simp[3^(1/4)\*Sqrt[2 + Sqrt[3]]\*d\*s\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/(1 - Sqrt[3])\*s + r\*x]^2)/(r^2\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])\*EllipticE[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b\*c^3 - 2\*(5 + 3\*Sqrt[3])\*a\*d^3, 0]

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(bx + cx^2)^{2/3} \int \left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{2/3} dx}{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
 &= -\frac{\left(b^2(bx + cx^2)^{2/3}\right) \text{Subst}\left(\int \left(1 - \frac{b^2x^2}{c^2}\right)^{2/3} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{4\sqrt[3]{2}c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
 &= \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b + 2cx) (bx + cx^2)^{2/3}}{14c \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
 &= -\frac{\left(b^2(bx + cx^2)^{2/3}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{1 - \frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{7\sqrt[3]{2}c^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{2/3}}{14c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
&\quad + \frac{\left(3(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx\left(1+\frac{cx}{b}\right)}{b}}\right)}{14\sqrt[3]{2}\left(-\frac{c}{b}-\frac{2c^2x}{b^2}\right)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
&= \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{2/3}}{14c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
&\quad - \frac{\left(3(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1+\sqrt{3}-x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx\left(1+\frac{cx}{b}\right)}{b}}\right)}{14\sqrt[3]{2}\left(-\frac{c}{b}-\frac{2c^2x}{b^2}\right)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
&\quad + \frac{\left(3(1+\sqrt{3})(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx\left(1+\frac{cx}{b}\right)}{b}}\right)}{14\sqrt[3]{2}\left(-\frac{c}{b}-\frac{2c^2x}{b^2}\right)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} (b+2cx)(bx+cx^2)^{2/3}}{14c\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3}} \\
&+ \frac{3b^2(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}}{7\sqrt[3]{2}c(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \left(1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
&- \frac{3^4\sqrt{3}\sqrt{2+\sqrt{3}}b^2(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}}}}{14\sqrt[3]{2}c(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}} \\
&+ \frac{\sqrt[6]{23}^{3/4}b^2(bx+cx^2)^{2/3} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx}{b^2}}}}} {7c(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{2/3} \sqrt{-1-\frac{4cx(b+cx)}{b^2}} \sqrt{\frac{1-\sqrt{3}-2^{2/3}\sqrt[3]{-\frac{cx}{b^2}}}{1-\sqrt{3}-2^{2/3}}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.06

$$\int (bx+cx^2)^{2/3} dx = \frac{3x(x(b+cx))^{2/3} \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{5}{3}, \frac{8}{3}, -\frac{cx}{b}\right)}{5\left(1+\frac{cx}{b}\right)^{2/3}}$$

[In] Integrate[(b\*x + c\*x^2)^(2/3), x]

[Out] (3\*x\*(x\*(b + c\*x))^(2/3)\*Hypergeometric2F1[-2/3, 5/3, 8/3, -(c\*x)/b])/ (5\*(1 + (c\*x)/b)^(2/3))

**Maple [F]**

$$\int (cx^2 + bx)^{\frac{2}{3}} dx$$

[In] int((c\*x^2+b\*x)^(2/3),x)

[Out] int((c\*x^2+b\*x)^(2/3),x)

**Fricas [F]**

$$\int (bx + cx^2)^{2/3} dx = \int (cx^2 + bx)^{\frac{2}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(2/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(2/3), x)

**Sympy [F]**

$$\int (bx + cx^2)^{2/3} dx = \int (bx + cx^2)^{\frac{2}{3}} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(2/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(2/3), x)

**Maxima [F]**

$$\int (bx + cx^2)^{2/3} dx = \int (cx^2 + bx)^{\frac{2}{3}} dx$$

[In] integrate((c\*x^2+b\*x)^(2/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(2/3), x)

**Giac [F]**

$$\int (bx + cx^2)^{2/3} dx = \int (cx^2 + bx)^{2/3} dx$$

[In] integrate((c\*x^2+b\*x)^(2/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(2/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.06 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.05

$$\int (bx + cx^2)^{2/3} dx = \frac{3x(cx^2 + bx)^{2/3} {}_2F_1\left(-\frac{2}{3}, \frac{5}{3}; \frac{8}{3}; -\frac{cx}{b}\right)}{5\left(\frac{cx}{b} + 1\right)^{2/3}}$$

[In] int((b\*x + c\*x^2)^(2/3),x)

[Out] (3\*x\*(b\*x + c\*x^2)^(2/3)\*hypergeom([-2/3, 5/3], 8/3, -(c\*x)/b))/(5\*((c\*x)/b + 1)^(2/3))

$$3.37 \quad \int \frac{1}{\sqrt[3]{bx + cx^2}} dx$$

Optimal result	240
Rubi [A] (verified)	241
Mathematica [C] (verified)	245
Maple [F]	246
Fricas [F]	246
Sympy [F]	246
Maxima [F]	246
Giac [F]	247
Mupad [B] (verification not implemented)	247

### Optimal result

Integrand size = 13, antiderivative size = 715

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = -\frac{3(b + 2cx) \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}}{\sqrt[3]{2c} \sqrt[3]{bx + cx^2} \left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)} + \frac{3^4 \sqrt{3} \sqrt{2 + \sqrt{3}} b^2 \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right) \sqrt{\frac{1 + 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}} + 2 \sqrt[3]{2} \left(-\frac{cx(b + cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}{\sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}} + \frac{2^3 \sqrt{2} c (b + 2cx) \sqrt[3]{bx + cx^2} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}{\sqrt{\frac{1 + 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}} + 2 \sqrt[3]{2} \left(-\frac{cx(b + cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}} \text{EllipticF} + \frac{c(b + 2cx) \sqrt[3]{bx + cx^2} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}{\sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right)^2}}}$$

[Out] -3/2\*(2\*c\*x+b)\*(-c\*(c\*x^2+b\*x)/b^2)^(1/3)\*2^(2/3)/c/(c\*x^2+b\*x)^(1/3)/(1-2^



$$\begin{aligned}
& (2/3)*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)}+2^{(1/6)}*3^{(3/4)}*b^2*(-c*(c*x^2+b*x) \\
& /b^2)^{(1/3)}*(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)})*EllipticF((1-2^{(2/3)}*(-c*x \\
& *(c*x+b)/b^2)^{(1/3)}+3^{(1/2)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)}),2 \\
& *I-I*3^{(1/2)})*((1+2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}+2*2^{(1/3)}*(-c*x*(c*x+b)/ \\
& b^2)^{(2/3)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}/c/(2*c*x+ \\
& b)/(c*x^2+b*x)^{(1/3)}/((-1+2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)})/(1-2^{(2/3)}*(-c \\
& x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}-3/4*3^{(1/4)}*b^2*(-c*(c*x^2+b*x)/b^2) \\
& ^{(1/3)}*(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)})*EllipticE((1-2^{(2/3)}*(-c*x*(c*x \\
& +b)/b^2)^{(1/3)}+3^{(1/2)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)}),2*I-I* \\
& 3^{(1/2)})*((1+2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}+2*2^{(1/3)}*(-c*x*(c*x+b)/b^2)^{( \\
& 2/3)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}*(1/2*6^{(1/2)}+1 \\
& /2*2^{(1/2)})*2^{(2/3)}/c/(2*c*x+b)/(c*x^2+b*x)^{(1/3)}/((-1+2^{(2/3)}*(-c*x*(c*x+b) \\
& )/b^2)^{(1/3)})/(1-2^{(2/3)}*(-c*x*(c*x+b)/b^2)^{(1/3)}-3^{(1/2)})^2)^{(1/2)}
\end{aligned}$$

### Rubi [A] (verified)

Time = 0.60 (sec) , antiderivative size = 715, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$ , Rules used

= {636, 633, 241, 310, 225, 1893}

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx$$

$$= \frac{\sqrt[6]{2} 3^{3/4} b^2 \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} \operatorname{EllipticF}\left(\dots\right)}{c(b + 2cx) \sqrt[3]{bx + cx^2} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}} - \frac{3^4 \sqrt[3]{3} \sqrt{2 + \sqrt{3}} b^2 \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} E\left(\dots\right)}{2^3 \sqrt[3]{2} c(b + 2cx) \sqrt[3]{bx + cx^2} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}}} - \frac{3(b + 2cx) \sqrt[3]{-\frac{c(bx + cx^2)}{b^2}}}{\sqrt[3]{2} c \sqrt[3]{bx + cx^2} \left(-2^{2/3} \sqrt[3]{-\frac{cx(b + cx)}{b^2}} - \sqrt{3} + 1\right)}$$

[In] Int[(b\*x + c\*x^2)^(-1/3),x]

[Out] (-3\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(1/3))/(2^(1/3)\*c\*(b\*x + c\*x^2)^(1/3)\*(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)) - (3\*3^(1/4)\*Sqrt[2 + Sqrt[3]]\*b^2\*(-(c\*(b\*x + c\*x^2))/b^2))^(1/3)\*(1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)\*Sqrt[(1 + 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-(c\*x\*(b + c\*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2]\*EllipticE[ArcSin[(1 + Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)], -7 + 4\*Sqrt[3]])/(2\*2^(1/3)\*c\*(b + 2\*c\*x)\*(b\*x + c\*x^2)^(1/3)\*Sqrt[-(1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2]] + (2^(1/6)\*3^(3/4)\*b^2\*(-(c\*(b\*x + c\*x^2))/b^2)^(1/3)\*(1 - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))\*Sqrt[(1 + 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3) + 2\*2^(1/3)\*(-(c\*x\*(b + c\*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x))/b^2))^(1/3))^2]\*EllipticF[ArcSin[(1

+ Sqrt[3] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3))  
 \*(-((c\*x\*(b + c\*x))/b^2))^(1/3)], -7 + 4\*Sqrt[3]]/(c\*(b + 2\*c\*x)\*(b\*x + c  
 \*x^2)^(1/3)\*Sqrt[-((1 - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))/(1 - Sqrt[3]  
 ] - 2^(2/3)\*(-((c\*x\*(b + c\*x))/b^2))^(1/3))^2))]

#### Rule 225

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]],  
 s = Denom[Rt[b/a, 3]]}, Simp[2\*Sqrt[2 - Sqrt[3]]\*(s + r\*x)\*(Sqrt[(s^2 - r\*s  
 \*x + r^2\*x^2)/((1 - Sqrt[3])\*s + r\*x)^2]/(3^(1/4)\*r\*Sqrt[a + b\*x^3]\*Sqrt[(-  
 s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)))]\*EllipticF[ArcSin[((1 + Sqrt[3])  
 \*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b}, x  
 ] && NegQ[a]

#### Rule 241

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1/3), x\_Symbol] := Dist[3\*(Sqrt[b\*x^2]/(2\*b\*x))  
 , Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b\*x^2)^(1/3)], x] /; FreeQ[{a, b}  
 , x]

#### Rule 310

Int[(x\_)/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Rt[b/a, 3]  
 ], s = Denom[Rt[b/a, 3]]}, Dist[(-1 + Sqrt[3])\*(s/r), Int[1/Sqrt[a + b\*x^  
 3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])\*s + r\*x)/Sqrt[a + b\*x^3], x], x]  
 /; FreeQ[{a, b}, x] && NegQ[a]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*  
 (c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b  
 + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

#### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-  
 c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; Fr  
 eeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

#### Rule 1893

Int[((c\_) + (d\_.)\*(x\_))/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = N  
 umer[Simplify[(1 + Sqrt[3])\*(d/c)]], s = Denom[Simplify[(1 + Sqrt[3])\*(d/c)  
 ]]}, Simp[2\*d\*s^3\*(Sqrt[a + b\*x^3]/(a\*r^2\*((1 - Sqrt[3])\*s + r\*x))), x] + S  
 imp[3^(1/4)\*Sqrt[2 + Sqrt[3]]\*d\*s\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/(  
 (1 - Sqrt[3])\*s + r\*x)^2]/(r^2\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - S  
 qrt[3])\*s + r\*x)^2)))]\*EllipticE[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[

3])\*(s + r\*x)], -7 + 4\*sqrt[3]], x]] /; FreeQ[{a, b, c, d}, x] && NegQ[a] &&  
EqQ[b\*c^3 - 2\*(5 + 3\*sqrt[3])\*a\*d^3, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \int \frac{1}{\sqrt[3]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}}} dx}{\sqrt[3]{bx+cx^2}} \\
 &= -\frac{\left(b^2 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{1-\frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{\sqrt[3]{2c^2} \sqrt[3]{bx+cx^2}} \\
 &= -\frac{\left(3 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \sqrt[3]{bx+cx^2}} \\
 &= -\frac{\left(3 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1+\sqrt{3}-x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \sqrt[3]{bx+cx^2}} \\
 &+ \frac{\left(3(1+\sqrt{3}) \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx(1+\frac{cx}{b})}{b}}\right)}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) \sqrt[3]{bx+cx^2}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3b^2 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}}{\sqrt[3]{2c(b+2cx)} \sqrt[3]{bx+cx^2} \left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
&\quad - \frac{3^4 \sqrt[3]{3} \sqrt{2+\sqrt{3}} b^2 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx}{b^2}}}{(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}})}}}{2^3 \sqrt[3]{2} c(b+2cx) \sqrt[3]{bx+cx^2} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}} \\
&\quad + \frac{\sqrt[6]{23^3/4} b^2 \sqrt[3]{-\frac{c(bx+cx^2)}{b^2}} \sqrt{-1-\frac{4cx}{b}-\frac{4c^2x^2}{b^2}} \left(1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}})}}}{c(b+2cx) \sqrt[3]{bx+cx^2} \sqrt{-1-\frac{4cx(b+cx)}{b^2}}} \sqrt{\frac{1}{(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}})}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.06

$$\int \frac{1}{\sqrt[3]{bx+cx^2}} dx = \frac{3x \sqrt[3]{1+\frac{cx}{b}} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{5}{3}, -\frac{cx}{b}\right)}{2 \sqrt[3]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-1/3),x]

[Out] (3\*x\*(1 + (c\*x)/b)^(1/3)\*Hypergeometric2F1[1/3, 2/3, 5/3, -((c\*x)/b)])/(2\*(x\*(b + c\*x))^(1/3))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{1}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(1/3),x)

[Out] int(1/(c\*x^2+b\*x)^(1/3),x)

**Fricas [F]**

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(-1/3), x)

**Sympy [F]**

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = \int \frac{1}{\sqrt[3]{bx + cx^2}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(1/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-1/3), x)

**Maxima [F]**

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-1/3), x)

**Giac [F]**

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-1/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.05

$$\int \frac{1}{\sqrt[3]{bx + cx^2}} dx = \frac{3x \left(\frac{cx}{b} + 1\right)^{1/3} {}_2F_1\left(\frac{1}{3}, \frac{2}{3}; \frac{5}{3}; -\frac{cx}{b}\right)}{2(cx^2 + bx)^{1/3}}$$

[In] int(1/(b\*x + c\*x^2)^(1/3),x)

[Out] (3\*x\*((c\*x)/b + 1)^(1/3)\*hypergeom([1/3, 2/3], 5/3, -(c\*x)/b))/(2\*(b\*x + c\*x^2)^(1/3))

**3.38**       $\int \frac{1}{(bx+cx^2)^{4/3}} dx$

Optimal result	249
Rubi [A] (verified)	250
Mathematica [C] (verified)	254
Maple [F]	255
Fricas [F]	255
Sympy [F]	255
Maxima [F]	255
Giac [F]	256
Mupad [B] (verification not implemented)	256



## Optimal result

Integrand size = 13, antiderivative size = 773

$$\int \frac{1}{(bx + cx^2)^{4/3}} dx = \frac{3(b + 2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx + cx^2)^{4/3}} + \frac{3 \cdot 2^{2/3} (b + 2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c (bx + cx^2)^{4/3} \left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} + \frac{3^4 \sqrt{3} \sqrt{2 + \sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} E\left(\frac{\sqrt[3]{2} c (b + 2cx) (bx + cx^2)^{4/3} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}{2^6 \sqrt{2} 3^{3/4} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2 \sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} \text{EllipticF}\left(\frac{c(b + 2cx) (bx + cx^2)^{4/3} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}\right)}{c(b + 2cx) (bx + cx^2)^{4/3} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}}{2}$$

[Out]  $3*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(4/3)/c/(-c*x*(c*x+b)/b^2)^(1/3)/(c*x^2+b*x)^(4/3)+3*2^(2/3)*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(4/3)/c/(c*x^2+b*x)^(4/3)/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))-2*2^(1/6)*3^(3/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(4/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*\text{EllipticF}\left(\frac{(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))}{(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))}, 2*I-I*3^(1/2)*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+2*2^(1/3)*(-c*x*(c*x+b)/b^2)^(2/3)))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)/c/(2*c*x+b)/(c*x^2+b*x)^(4/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)+3/2*3^(1/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(4/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*\text{EllipticE}\left(\frac{(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))}{(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))}\right)$

$), 2 * I - I * 3^{(1/2)}) * ((1 + 2^{(2/3)} * (-c * x * (c * x + b) / b^2)^{(1/3)} + 2 * 2^{(1/3)} * (-c * x * (c * x + b) / b^2)^{(2/3)}) / (1 - 2^{(2/3)} * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)})^2)^{(1/2)} * (1/2 * 6^{(1/2)} + 1/2 * 2^{(1/2)}) * 2^{(2/3)} / c / (2 * c * x + b) / (c * x^2 + b * x)^{(4/3)} / ((-1 + 2^{(2/3)} * (-c * x * (c * x + b) / b^2)^{(1/3)}) / (1 - 2^{(2/3)} * (-c * x * (c * x + b) / b^2)^{(1/3)} - 3^{(1/2)})^2)^{(1/2)}$

### Rubi [A] (verified)

Time = 0.71 (sec) , antiderivative size = 773, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {636, 633, 205, 241, 310, 225, 1893}

$$\int \frac{1}{(bx + cx^2)^{4/3}} dx =$$

$$\frac{2^6 \sqrt{2} 3^{3/4} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} \text{EllipticF}\left(\dots\right)}{c(b+2cx)(bx+cx^2)^{4/3} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)}}} + \frac{3^4 \sqrt{3} \sqrt{2 + \sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{2^3 \sqrt{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3} + 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 1}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)^2}} E\left(\dots\right)}{\sqrt[3]{2} c(b+2cx)(bx+cx^2)^{4/3} \sqrt{\frac{1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)}}} + \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{4/3}} + \frac{3 \cdot 2^{2/3} (b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c (bx+cx^2)^{4/3} \left(-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} - \sqrt{3} + 1\right)}$$

[In] Int[(b\*x + c\*x^2)^(-4/3), x]

[Out] (3\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(4/3)/(c\*(-((c\*x\*(b + c\*x))/b^2))^(1/3)\*(b\*x + c\*x^2)^(4/3)) + (3\*2^(2/3)\*(b + 2\*c\*x)\*(-(c\*(b\*x + c\*x^2))/b^2))^(4/3)/(c\*(b\*x + c\*x^2)^(4/3)\*(1 - Sqrt[3] - 2^(2/3)\*(-(c\*x\*(b + c\*x

```

)))/b^2))^(1/3))) + (3*3^(1/4)*Sqrt[2 + Sqrt[3]]*b^2*(-((c*(b*x + c*x^2))/b^
2))^(4/3)*(1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))*Sqrt[(1 + 2^(2/3)*(-
((c*x*(b + c*x))/b^2))^(1/3) + 2*2^(1/3)*(-((c*x*(b + c*x))/b^2))^(2/3))/(1
- Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2]*EllipticE[ArcSin[(1
+ Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)
*(-((c*x*(b + c*x))/b^2))^(1/3))], -7 + 4*Sqrt[3]]]/(2^(1/3)*c*(b + 2*c*x)*
(b*x + c*x^2)^(4/3)*Sqrt[-((1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))/(1
- Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2]] - (2*2^(1/6)*3^(3/
4)*b^2*(-((c*(b*x + c*x^2))/b^2))^(4/3)*(1 - 2^(2/3)*(-((c*x*(b + c*x))/b^2
))^(1/3))*Sqrt[(1 + 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3) + 2*2^(1/3)*(-((
c*x*(b + c*x))/b^2))^(2/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))
^(1/3))^2]*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))
^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))], -7 + 4*Sqr
t[3]]]/(c*(b + 2*c*x)*(b*x + c*x^2)^(4/3)*Sqrt[-((1 - 2^(2/3)*(-((c*x*(b +
c*x))/b^2))^(1/3))/(1 - Sqrt[3] - 2^(2/3)*(-((c*x*(b + c*x))/b^2))^(1/3))^2
]))

```

#### Rule 205

```

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p +
1)/(a*n*(p + 1))), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n
)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (Integ
erQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denom
inator[p + 1/n] < Denominator[p])

```

#### Rule 225

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]],
s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s
*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-
s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2)])]*EllipticF[ArcSin[((1 + Sqrt[3])
*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x
] && NegQ[a]

```

#### Rule 241

```

Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x))
, Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}
, x]

```

#### Rule 310

```

Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
], s = Denom[Rt[b/a, 3]]}, Dist[(-1 + Sqrt[3])*(s/r), Int[1/Sqrt[a + b*x^
3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])*s + r*x)/Sqrt[a + b*x^3], x], x]
/; FreeQ[{a, b}, x] && NegQ[a]

```

Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[(-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

Rule 1893

Int[((c\_) + (d\_.)\*(x\_))/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Simplify[(1 + Sqrt[3])\*(d/c)], s = Denom[Simplify[(1 + Sqrt[3])\*(d/c)]]], Simp[2\*d\*s^3\*(Sqrt[a + b\*x^3]/(a\*r^2\*((1 - Sqrt[3])\*s + r\*x))), x] + Simp[3^(1/4)\*Sqrt[2 + Sqrt[3]]\*d\*s\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/(1 - Sqrt[3])\*s + r\*x]^2)/(r^2\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])]\*EllipticE[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b\*c^3 - 2\*(5 + 3\*Sqrt[3])\*a\*d^3, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \int \frac{1}{\left(-\frac{cx}{b}-\frac{c^2x^2}{b^2}\right)^{4/3}} dx}{(bx+cx^2)^{4/3}} \\
 &= -\frac{\left(2^{2/3}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{4/3}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{4/3}} \\
 &= \frac{3(b+2cx)\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c^3\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(bx+cx^2)^{4/3}} \\
 &\quad + \frac{\left(2^{2/3}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{1-\frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{4/3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{4/3}} \\
&\quad \left(3 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&\quad \frac{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}}{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{4/3}} \\
&\quad \left(3 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1+\sqrt{3}-x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&\quad + \frac{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}}{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}} \\
&\quad \left(3(1+\sqrt{3}) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&\quad \frac{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}}{\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{4/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3}}{c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{4/3}} \\
&= \frac{3 \cdot 2^{2/3} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}{c(b+2cx) (bx+cx^2)^{4/3} \left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
&+ \frac{3^4 \sqrt{3} \sqrt{2 + \sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}{\sqrt[3]{2} c(b+2cx) (bx+cx^2)^{4/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}} \\
&- \frac{2^6 \sqrt{3} 3^{3/4} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{4/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right) \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}}{c(b+2cx) (bx+cx^2)^{4/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.06

$$\int \frac{1}{(bx+cx^2)^{4/3}} dx = -\frac{3 \sqrt[3]{1 + \frac{cx}{b}} \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{4}{3}, \frac{2}{3}, -\frac{cx}{b}\right)}{b \sqrt[3]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-4/3), x]

[Out] (-3\*(1 + (c\*x)/b)^(1/3)\*Hypergeometric2F1[-1/3, 4/3, 2/3, -(c\*x)/b])/ (b\*(x\*(b + c\*x))^(1/3))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{4}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(4/3),x)

[Out] int(1/(c\*x^2+b\*x)^(4/3),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{4}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{4}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(4/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(2/3)/(c^2\*x^4 + 2\*b\*c\*x^3 + b^2\*x^2), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{4}{3}}} dx = \int \frac{1}{(bx + cx^2)^{\frac{4}{3}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(4/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-4/3), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{4}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{4}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(4/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-4/3), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{4/3}} dx = \int \frac{1}{(cx^2 + bx)^{4/3}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(4/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-4/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.05

$$\int \frac{1}{(bx + cx^2)^{4/3}} dx = -\frac{3x \left(\frac{cx}{b} + 1\right)^{4/3} {}_2F_1\left(-\frac{1}{3}, \frac{4}{3}; \frac{2}{3}; -\frac{cx}{b}\right)}{(cx^2 + bx)^{4/3}}$$

[In] int(1/(b\*x + c\*x^2)^(4/3),x)

[Out] -(3\*x\*((c\*x)/b + 1)^(4/3)\*hypergeom([-1/3, 4/3], 2/3, -(c\*x)/b))/(b\*x + c\*x^2)^(4/3)



**3.39**       $\int \frac{1}{(bx+cx^2)^{7/3}} dx$

Optimal result	258
Rubi [A] (verified)	259
Mathematica [C] (verified)	263
Maple [F]	264
Fricas [F]	264
Sympy [F]	264
Maxima [F]	264
Giac [F]	265
Mupad [B] (verification not implemented)	265

## Optimal result

Integrand size = 13, antiderivative size = 838

$$\begin{aligned}
 \int \frac{1}{(bx + cx^2)^{7/3}} dx &= \frac{3(b + 2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx + cx^2)^{7/3}} \\
 &+ \frac{15(b + 2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{2c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx + cx^2)^{7/3}} + \frac{15(b + 2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{\sqrt[3]{2}c (bx + cx^2)^{7/3} \left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
 &+ \frac{15\sqrt[4]{3}\sqrt{2 + \sqrt{3}}b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}{\sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2\sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} E\left(\frac{\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}\right)} \\
 &+ \frac{2\sqrt[3]{2}c(b + 2cx) (bx + cx^2)^{7/3}}{\sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}} \\
 &+ \frac{5\sqrt[6]{2}3^{3/4}b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}{\sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}} + 2\sqrt[3]{2} \left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}} \text{EllipticF}\left(\frac{\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}\right)} \\
 &+ \frac{c(b + 2cx) (bx + cx^2)^{7/3}}{\sqrt{\frac{1-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)^2}}}
 \end{aligned}$$

```

[Out] 3/4*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(7/3)/c/(-c*x*(c*x+b)/b^2)^(4/3)/(c*x^2+
b*x)^(7/3)+15/2*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(7/3)/c/(-c*x*(c*x+b)/b^2)^(
1/3)/(c*x^2+b*x)^(7/3)+15/2*(2*c*x+b)*(-c*(c*x^2+b*x)/b^2)^(7/3)*2^(2/3)/c/
(c*x^2+b*x)^(7/3)/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))-5*2^(1/6)*3^(
3/4)*b^2*(-c*(c*x^2+b*x)/b^2)^(7/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*E
llipticF((1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3)*(-c*x*(c*x
+b)/b^2)^(1/3)-3^(1/2)),2*I-I*3^(1/2))*((1+2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)
+2*2^(1/3)*(-c*x*(c*x+b)/b^2)^(2/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(
1/2))^2)^(1/2)/c/(2*c*x+b)/(c*x^2+b*x)^(7/3)/((-1+2^(2/3)*(-c*x*(c*x+b)/b^
2)^(1/3))/(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)+15/4*3^(1/4
)*b^2*(-c*(c*x^2+b*x)/b^2)^(7/3)*(1-2^(2/3)*(-c*x*(c*x+b)/b^2)^(1/3))*Ellip

```

ticE((1-2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3)+3^(1/2))/(1-2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2)),2\*I-I\*3^(1/2))\*((1+2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3)+2\*2^(1/3))\*(-c\*x\*(c\*x+b)/b^2)^(2/3))/(1-2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)\*(1/2\*6^(1/2)+1/2\*2^(1/2))\*2^(2/3)/c/(2\*c\*x+b)/(c\*x^2+b\*x)^(7/3)/((-1+2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3))/(1-2^(2/3))\*(-c\*x\*(c\*x+b)/b^2)^(1/3)-3^(1/2))^2)^(1/2)

### Rubi [A] (verified)

Time = 0.72 (sec) , antiderivative size = 838, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$ , Rules used = {636, 633, 205, 241, 310, 225, 1893}

$$\int \frac{1}{(bx + cx^2)^{7/3}} dx = \frac{15\sqrt[4]{3}\sqrt{2+\sqrt{3}}b^2\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{2^3\sqrt{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}{2^3\sqrt{2}c(b+2cx)(cx^2+bx)^{7/3}\sqrt{\frac{1}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}}}{5\sqrt[6]{23^3/4}b^2\left(1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)\sqrt{\frac{2^3\sqrt{2}\left(-\frac{cx(b+cx)}{b^2}\right)^{2/3}+2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}\text{EllipticF}\left(\arcsin\left(\frac{-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1}{-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1}\right)\right)}{c(b+2cx)(cx^2+bx)^{7/3}\sqrt{\frac{1-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)^2}}}}}}}{+ \frac{15(b+2cx)\left(-\frac{c(cx^2+bx)}{b^2}\right)^{7/3}}{\sqrt[3]{2}c(cx^2+bx)^{7/3}\left(-2^{2/3}\sqrt[3]{-\frac{cx(b+cx)}{b^2}}-\sqrt{3}+1\right)}}{+ \frac{15(b+2cx)\left(-\frac{c(cx^2+bx)}{b^2}\right)^{7/3}}{2c\sqrt[3]{-\frac{cx(b+cx)}{b^2}}(cx^2+bx)^{7/3}} + \frac{3(b+2cx)\left(-\frac{c(cx^2+bx)}{b^2}\right)^{7/3}}{4c\left(-\frac{cx(b+cx)}{b^2}\right)^{4/3}(cx^2+bx)^{7/3}}}$$

[In] Int[(b\*x + c\*x^2)^(-7/3), x]

```
[Out] (3*(b + 2*c*x)*(-(c*(b*x + c*x^2))/b^2))^(7/3)/(4*c*(-((c*x*(b + c*x))/b^2))^(4/3)*(b*x + c*x^2)^(7/3)) + (15*(b + 2*c*x)*(-(c*(b*x + c*x^2))/b^2))^(7/3)/(2*c*(-((c*x*(b + c*x))/b^2))^(1/3)*(b*x + c*x^2)^(7/3)) + (15*(b + 2*c*x)*(-(c*(b*x + c*x^2))/b^2))^(7/3)/(2^(1/3)*c*(b*x + c*x^2)^(7/3)*(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)) + (15*3^(1/4)*Sqrt[2 + Sqrt[3]]*b^2*(-((c*(b*x + c*x^2))/b^2))^(7/3)*(1 - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))*Sqrt[(1 + 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3) + 2*2^(1/3)*(-(c*x*(b + c*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))^2]*EllipticE[ArcSin[(1 + Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)], -7 + 4*Sqrt[3]]/(2*2^(1/3)*c*(b + 2*c*x)*(b*x + c*x^2)^(7/3)*Sqrt[-((1 - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))^2]) - (5*2^(1/6)*3^(3/4)*b^2*(-((c*(b*x + c*x^2))/b^2))^(7/3)*(1 - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))*Sqrt[(1 + 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3) + 2*2^(1/3)*(-(c*x*(b + c*x))/b^2))^(2/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))^2]*EllipticF[ArcSin[(1 + Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)], -7 + 4*Sqrt[3]]/(c*(b + 2*c*x)*(b*x + c*x^2)^(7/3)*Sqrt[-((1 - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3)]/(1 - Sqrt[3] - 2^(2/3)*(-(c*x*(b + c*x))/b^2))^(1/3))^2])
```

#### Rule 205

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])
```

#### Rule 225

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[2*Sqrt[2 - Sqrt[3]]*(s + r*x)*(Sqrt[(s^2 - r*s*x + r^2*x^2)/((1 - Sqrt[3])*s + r*x)^2]/(3^(1/4)*r*Sqrt[a + b*x^3]*Sqrt[(-s)*((s + r*x)/((1 - Sqrt[3])*s + r*x)^2])))*EllipticF[ArcSin[((1 + Sqrt[3])*s + r*x)/((1 - Sqrt[3])*s + r*x)], -7 + 4*Sqrt[3]], x] /; FreeQ[{a, b}, x] && NegQ[a]
```

#### Rule 241

```
Int[((a_) + (b_.)*(x_)^2)^(-1/3), x_Symbol] := Dist[3*(Sqrt[b*x^2]/(2*b*x)), Subst[Int[x/Sqrt[-a + x^3], x], x, (a + b*x^2)^(1/3)], x] /; FreeQ[{a, b}, x]
```

#### Rule 310

```
Int[(x_)/Sqrt[(a_) + (b_.)*(x_)^3], x_Symbol] := With[{r = Numer[Rt[b/a, 3]
```

], s = Denom[Rt[b/a, 3]], Dist[(-(1 + Sqrt[3]))\*(s/r), Int[1/Sqrt[a + b\*x^3], x], x] + Dist[1/r, Int[((1 + Sqrt[3])\*s + r\*x)/Sqrt[a + b\*x^3], x], x] /; FreeQ[{a, b}, x] && NegQ[a]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*c/(b^2 - 4\*a\*c)))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rule 1893

Int[((c\_) + (d\_.)\*(x\_))/Sqrt[(a\_) + (b\_.)\*(x\_)^3], x\_Symbol] := With[{r = Numer[Simplify[(1 + Sqrt[3])\*(d/c)]]}, s = Denom[Simplify[(1 + Sqrt[3])\*(d/c)]]], Simp[2\*d\*s^3\*(Sqrt[a + b\*x^3]/(a\*r^2\*((1 - Sqrt[3])\*s + r\*x))), x] + Simp[3^(1/4)\*Sqrt[2 + Sqrt[3]]\*d\*s\*(s + r\*x)\*(Sqrt[(s^2 - r\*s\*x + r^2\*x^2)/(1 - Sqrt[3])\*s + r\*x]^2)/(r^2\*Sqrt[a + b\*x^3]\*Sqrt[(-s)\*((s + r\*x)/((1 - Sqrt[3])\*s + r\*x)^2)])\*EllipticE[ArcSin[((1 + Sqrt[3])\*s + r\*x)/((1 - Sqrt[3])\*s + r\*x)], -7 + 4\*Sqrt[3]], x] /; FreeQ[{a, b, c, d}, x] && NegQ[a] && EqQ[b\*c^3 - 2\*(5 + 3\*Sqrt[3])\*a\*d^3, 0]

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \int \frac{1}{\left(-\frac{cx}{b}-\frac{c^2x^2}{b^2}\right)^{7/3}} dx}{(bx+cx^2)^{7/3}} \\ &= -\frac{\left(8 \cdot 2^{2/3} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{7/3}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c^2 (bx+cx^2)^{7/3}} \\ &= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx+cx^2)^{7/3}} \\ &\quad - \frac{\left(5 \cdot 2^{2/3} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{4/3}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c^2 (bx+cx^2)^{7/3}} \end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx+cx^2)^{7/3}} + \frac{15(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{2c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{7/3}} \\
&\quad \left(5b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}\right) \text{Subst} \left( \int \frac{1}{\sqrt[3]{1-\frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2} \right) \\
&+ \frac{\quad}{\sqrt[3]{2c^2} (bx+cx^2)^{7/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx+cx^2)^{7/3}} + \frac{15(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{2c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{7/3}} \\
&\quad \left(15 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&+ \frac{\quad}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{7/3}} \\
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx+cx^2)^{7/3}} + \frac{15(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{2c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{7/3}} \\
&\quad \left(15 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1+\sqrt{3}-x}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&+ \frac{\quad}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{7/3}} \\
&\quad \left(15(1+\sqrt{3}) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}}\right) \text{Subst} \left( \int \frac{1}{\sqrt{-1+x^3}} dx, x, 2^{2/3} \sqrt[3]{-\frac{cx \left(1 + \frac{cx}{b}\right)}{b}} \right) \\
&+ \frac{\quad}{2\sqrt[3]{2} \left(-\frac{c}{b} - \frac{2c^2x}{b^2}\right) (bx+cx^2)^{7/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{4c \left(-\frac{cx(b+cx)}{b^2}\right)^{4/3} (bx+cx^2)^{7/3}} + \frac{15(b+2cx) \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3}}{2c \sqrt[3]{-\frac{cx(b+cx)}{b^2}} (bx+cx^2)^{7/3}} \\
&\quad - \frac{15b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}{\sqrt[3]{2}c(b+2cx) (bx+cx^2)^{7/3} \left(1 - \sqrt{3} - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)} \\
&\quad + \frac{15^4 \sqrt{3} \sqrt{2 + \sqrt{3}} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}{2^3 \sqrt{2} c(b+2cx) (bx+cx^2)^{7/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}} \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx}{b^2}}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx}{b^2}}\right)}}} \\
&\quad - \frac{5^6 \sqrt{2} 3^{3/4} b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{7/3} \sqrt{-1 - \frac{4cx}{b} - \frac{4c^2x^2}{b^2}} \left(1 - 2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}{c(b+2cx) (bx+cx^2)^{7/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}} \sqrt{\frac{1+2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}}} \\
&\quad - \frac{c(b+2cx) (bx+cx^2)^{7/3} \sqrt{-1 - \frac{4cx(b+cx)}{b^2}}}{\left(1-\sqrt{3}-2^{2/3} \sqrt[3]{-\frac{cx(b+cx)}{b^2}}\right)}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.06

$$\int \frac{1}{(bx+cx^2)^{7/3}} dx = -\frac{3 \sqrt[3]{1 + \frac{cx}{b}} \operatorname{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{7}{3}, -\frac{1}{3}, -\frac{cx}{b}\right)}{4b^2 x \sqrt[3]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-7/3),x]

[Out] (-3\*(1 + (c\*x)/b)^(1/3)\*Hypergeometric2F1[-4/3, 7/3, -1/3, -(c\*x)/b])/(4\*b^2\*x\*(x\*(b + c\*x))^(1/3))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{7}{3}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(7/3),x)

[Out] int(1/(c\*x^2+b\*x)^(7/3),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{7}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{7}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(7/3),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(2/3)/(c^3\*x^6 + 3\*b\*c^2\*x^5 + 3\*b^2\*c\*x^4 + b^3\*x^3), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{7}{3}}} dx = \int \frac{1}{(bx + cx^2)^{\frac{7}{3}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(7/3),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-7/3), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{7}{3}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{7}{3}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(7/3),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-7/3), x)



**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{7/3}} dx = \int \frac{1}{(cx^2 + bx)^{7/3}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(7/3),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-7/3), x)

**Mupad [B] (verification not implemented)**

Time = 9.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.04

$$\int \frac{1}{(bx + cx^2)^{7/3}} dx = -\frac{3x \left(\frac{cx}{b} + 1\right)^{7/3} {}_2F_1\left(-\frac{4}{3}, \frac{7}{3}; -\frac{1}{3}; -\frac{cx}{b}\right)}{4(cx^2 + bx)^{7/3}}$$

[In] int(1/(b\*x + c\*x^2)^(7/3),x)

[Out] -(3\*x\*((c\*x)/b + 1)^(7/3)\*hypergeom([-4/3, 7/3], -1/3, -(c\*x)/b))/(4\*(b\*x + c\*x^2)^(7/3))

### 3.40 $\int (bx + cx^2)^{5/4} dx$

Optimal result	266
Rubi [A] (verified)	266
Mathematica [C] (verified)	268
Maple [F]	268
Fricas [F]	268
Sympy [F]	269
Maxima [F]	269
Giac [F]	269
Mupad [B] (verification not implemented)	269

#### Optimal result

Integrand size = 13, antiderivative size = 119

$$\int (bx + cx^2)^{5/4} dx = -\frac{5b^2(b + 2cx)\sqrt{bx + cx^2}}{84c^2} + \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c} + \frac{5b^5\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(1 + \frac{2cx}{b}\right), 2\right)}{84\sqrt{2}c^3(bx + cx^2)^{3/4}}$$

[Out]  $-5/84*b^2*(2*c*x+b)*(c*x^2+b*x)^{(1/4)}/c^2+1/7*(2*c*x+b)*(c*x^2+b*x)^{(5/4)}/c+5/168*b^5*(-c*(c*x^2+b*x)/b^2)^{(3/4)}*(\cos(1/2*\arcsin(1+2*c*x/b))^{(1/2)})/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticF}(\sin(1/2*\arcsin(1+2*c*x/b)), 2^{(1/2)})/c^3/(c*x^2+b*x)^{(3/4)}*2^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {626, 636, 633, 238}

$$\int (bx + cx^2)^{5/4} dx = \frac{5b^5\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{2cx}{b} + 1\right), 2\right)}{84\sqrt{2}c^3(bx + cx^2)^{3/4}} - \frac{5b^2(b + 2cx)\sqrt{bx + cx^2}}{84c^2} + \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c}$$

[In] Int[(b\*x + c\*x^2)^(5/4), x]

[Out]  $(-5*b^2*(b + 2*c*x)*(b*x + c*x^2)^{(1/4)})/(84*c^2) + ((b + 2*c*x)*(b*x + c*x^2)^{(5/4)})/(7*c) + (5*b^5*(-((c*(b*x + c*x^2))/b^2))^{(3/4)}*\text{EllipticF}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/(84*\text{Sqrt}[2]*c^3*(b*x + c*x^2)^{(3/4)})$

Rule 238

Int[((a\_) + (b\_)\*(x\_)^2)^(-3/4), x\_Symbol] := Simp[(2/(a^(3/4)\*Rt[-b/a, 2]))\*EllipticF[(1/2)\*ArcSin[Rt[-b/a, 2]\*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b/a]

Rule 626

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*c/(b^2 - 4\*a\*c)))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rule 636

Int[((b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c} - \frac{(5b^2) \int \sqrt[4]{bx + cx^2} dx}{28c} \\
 &= -\frac{5b^2(b + 2cx)\sqrt[4]{bx + cx^2}}{84c^2} + \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c} + \frac{(5b^4) \int \frac{1}{(bx + cx^2)^{3/4}} dx}{336c^2} \\
 &= -\frac{5b^2(b + 2cx)\sqrt[4]{bx + cx^2}}{84c^2} + \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c} \\
 &\quad + \frac{\left(5b^4 \left(-\frac{c(bx + cx^2)}{b^2}\right)^{3/4}\right) \int \frac{1}{\left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{3/4}} dx}{336c^2 (bx + cx^2)^{3/4}} \\
 &= -\frac{5b^2(b + 2cx)\sqrt[4]{bx + cx^2}}{84c^2} + \frac{(b + 2cx)(bx + cx^2)^{5/4}}{7c} \\
 &\quad - \frac{\left(5b^6 \left(-\frac{c(bx + cx^2)}{b^2}\right)^{3/4}\right) \text{Subst}\left(\int \frac{1}{\left(1 - \frac{b^2x^2}{c^2}\right)^{3/4}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{168\sqrt{2}c^4 (bx + cx^2)^{3/4}}
 \end{aligned}$$

$$= -\frac{5b^2(b+2cx)\sqrt[4]{bx+cx^2}}{84c^2} + \frac{(b+2cx)(bx+cx^2)^{5/4}}{7c} \\ + \frac{5b^5\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} F\left(\frac{1}{2}\sin^{-1}\left(1+\frac{2cx}{b}\right)\middle|2\right)}{84\sqrt{2}c^3(bx+cx^2)^{3/4}}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.40

$$\int (bx+cx^2)^{5/4} dx = \frac{4bx^2\sqrt[4]{x(b+cx)} \operatorname{Hypergeometric2F1}\left(-\frac{5}{4}, \frac{9}{4}, \frac{13}{4}, -\frac{cx}{b}\right)}{9\sqrt[4]{1+\frac{cx}{b}}}$$

[In] Integrate[(b\*x + c\*x^2)^(5/4), x]

[Out] (4\*b\*x^2\*(x\*(b + c\*x))^(1/4)\*Hypergeometric2F1[-5/4, 9/4, 13/4, -((c\*x)/b)])/(9\*(1 + (c\*x)/b)^(1/4))

### Maple [F]

$$\int (cx^2 + bx)^{5/4} dx$$

[In] int((c\*x^2+b\*x)^(5/4), x)

[Out] int((c\*x^2+b\*x)^(5/4), x)

### Fricas [F]

$$\int (bx+cx^2)^{5/4} dx = \int (cx^2+bx)^{5/4} dx$$

[In] integrate((c\*x^2+b\*x)^(5/4), x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(5/4), x)

**Sympy [F]**

$$\int (bx + cx^2)^{5/4} dx = \int (bx + cx^2)^{\frac{5}{4}} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(5/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(5/4), x)

**Maxima [F]**

$$\int (bx + cx^2)^{5/4} dx = \int (cx^2 + bx)^{\frac{5}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(5/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(5/4), x)

**Giac [F]**

$$\int (bx + cx^2)^{5/4} dx = \int (cx^2 + bx)^{\frac{5}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(5/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(5/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.30

$$\int (bx + cx^2)^{5/4} dx = \frac{4x (cx^2 + bx)^{5/4} {}_2F_1\left(-\frac{5}{4}, \frac{9}{4}; \frac{13}{4}; -\frac{cx}{b}\right)}{9\left(\frac{cx}{b} + 1\right)^{5/4}}$$

[In] int((b\*x + c\*x^2)^(5/4),x)

[Out] (4\*x\*(b\*x + c\*x^2)^(5/4)\*hypergeom([-5/4, 9/4], 13/4, -(c\*x)/b))/(9\*((c\*x)/b + 1)^(5/4))

### 3.41 $\int (bx + cx^2)^{3/4} dx$

Optimal result	270
Rubi [A] (verified)	270
Mathematica [C] (verified)	272
Maple [F]	272
Fricas [F]	272
Sympy [F]	272
Maxima [F]	273
Giac [F]	273
Mupad [B] (verification not implemented)	273

#### Optimal result

Integrand size = 13, antiderivative size = 90

$$\int (bx + cx^2)^{3/4} dx = \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} - \frac{3b^3 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \arcsin\left(1 + \frac{2cx}{b}\right) \middle| 2\right)}{10\sqrt{2}c^2 \sqrt[4]{bx + cx^2}}$$

[Out]  $\frac{1}{5}*(2*c*x+b)*(c*x^2+b*x)^{(3/4)}/c-3/20*b^3*(-c*(c*x^2+b*x)/b^2)^{(1/4)}*(\cos(1/2*\arcsin(1+2*c*x/b))^{(1/2)}/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticE}(\sin(1/2*\arcsin(1+2*c*x/b)),2^{(1/2)}))/c^2/(c*x^2+b*x)^{(1/4)}*2^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {626, 636, 633, 234}

$$\int (bx + cx^2)^{3/4} dx = \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} - \frac{3b^3 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \arcsin\left(\frac{2cx}{b} + 1\right) \middle| 2\right)}{10\sqrt{2}c^2 \sqrt[4]{bx + cx^2}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{(3/4)}, x]$

[Out]  $((b + 2*c*x)*(b*x + c*x^2)^{(3/4)})/(5*c) - (3*b^3*(-((c*(b*x + c*x^2))/b^2))^{(1/4)}*\text{EllipticE}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/((10*\text{Sqrt}[2]*c^2*(b*x + c*x^2)^{(1/4)})$

#### Rule 234

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1/4}, x\_Symbol] \rightarrow \text{Simp}[(2/(a^{1/4})*\text{Rt}[-b/a, 2])*\text{EllipticE}[(1/2)*\text{ArcSin}[\text{Rt}[-b/a, 2]*x], 2], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{GtQ}$

[a, 0] && NegQ[b/a]

### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c)], x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2)^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} - \frac{(3b^2) \int \frac{1}{\sqrt[4]{bx + cx^2}} dx}{20c} \\
 &= \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} - \frac{\left(3b^2 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}}\right) \int \frac{1}{\sqrt[4]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}}} dx}{20c \sqrt[4]{bx + cx^2}} \\
 &= \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} + \frac{\left(3b^4 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt[4]{1 - \frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{20\sqrt{2}c^3 \sqrt[4]{bx + cx^2}} \\
 &= \frac{(b + 2cx)(bx + cx^2)^{3/4}}{5c} - \frac{3b^3 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \sin^{-1}\left(1 + \frac{2cx}{b}\right) \mid 2\right)}{10\sqrt{2}c^2 \sqrt[4]{bx + cx^2}}
 \end{aligned}$$

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.50

$$\int (bx + cx^2)^{3/4} dx = \frac{4x(x(b + cx))^{3/4} \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{7}{4}, \frac{11}{4}, -\frac{cx}{b}\right)}{7\left(1 + \frac{cx}{b}\right)^{3/4}}$$

[In] Integrate[(b\*x + c\*x^2)^(3/4), x]

[Out] (4\*x\*(x\*(b + c\*x))^(3/4)\*Hypergeometric2F1[-3/4, 7/4, 11/4, -((c\*x)/b)])/(7\*(1 + (c\*x)/b)^(3/4))

**Maple [F]**

$$\int (cx^2 + bx)^{\frac{3}{4}} dx$$

[In] int((c\*x^2+b\*x)^(3/4), x)

[Out] int((c\*x^2+b\*x)^(3/4), x)

**Fricas [F]**

$$\int (bx + cx^2)^{3/4} dx = \int (cx^2 + bx)^{\frac{3}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(3/4), x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(3/4), x)

**Sympy [F]**

$$\int (bx + cx^2)^{3/4} dx = \int (bx + cx^2)^{\frac{3}{4}} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(3/4), x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(3/4), x)



**Maxima [F]**

$$\int (bx + cx^2)^{3/4} dx = \int (cx^2 + bx)^{3/4} dx$$

[In] integrate((c\*x^2+b\*x)^(3/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(3/4), x)

**Giac [F]**

$$\int (bx + cx^2)^{3/4} dx = \int (cx^2 + bx)^{3/4} dx$$

[In] integrate((c\*x^2+b\*x)^(3/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(3/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.03 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.40

$$\int (bx + cx^2)^{3/4} dx = \frac{4x(cx^2 + bx)^{3/4} {}_2F_1\left(-\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; -\frac{cx}{b}\right)}{7\left(\frac{cx}{b} + 1\right)^{3/4}}$$

[In] int((b\*x + c\*x^2)^(3/4),x)

[Out] (4\*x\*(b\*x + c\*x^2)^(3/4)\*hypergeom([-3/4, 7/4], 11/4, -(c\*x)/b))/(7\*((c\*x)/b + 1)^(3/4))

### 3.42 $\int \sqrt[4]{bx + cx^2} dx$

Optimal result	274
Rubi [A] (verified)	274
Mathematica [C] (verified)	276
Maple [F]	276
Fricas [F]	276
Sympy [F]	276
Maxima [F]	277
Giac [F]	277
Mupad [B] (verification not implemented)	277

#### Optimal result

Integrand size = 13, antiderivative size = 90

$$\int \sqrt[4]{bx + cx^2} dx = \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} - \frac{b^3 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(1 + \frac{2cx}{b}\right), 2\right)}{3\sqrt{2}c^2 (bx + cx^2)^{3/4}}$$

[Out]  $\frac{1}{3}*(2*c*x+b)*(c*x^2+b*x)^{(1/4)}/c-1/6*b^3*(-c*(c*x^2+b*x)/b^2)^{(3/4)}*(\cos(1/2*\arcsin(1+2*c*x/b))^{2})^{(1/2)}/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticF}(\sin(1/2*\arcsin(1+2*c*x/b)), 2^{(1/2)})/c^2/(c*x^2+b*x)^{(3/4)*2^{(1/2)}}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {626, 636, 633, 238}

$$\int \sqrt[4]{bx + cx^2} dx = \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} - \frac{b^3 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{2cx}{b} + 1\right), 2\right)}{3\sqrt{2}c^2 (bx + cx^2)^{3/4}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{(1/4)}, x]$

[Out]  $((b + 2*c*x)*(b*x + c*x^2)^{(1/4)})/(3*c) - (b^3*(-((c*(b*x + c*x^2))/b^2))^{(3/4)}*\text{EllipticF}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/(3*\text{Sqrt}[2]*c^2*(b*x + c*x^2)^{(3/4)})$

#### Rule 238

$\text{Int}[(a_ + (b_.*x_)^2)^{(-3/4)}, x\_Symbol] \rightarrow \text{Simp}[(2/(a^{(3/4)}*\text{Rt}[-b/a, 2]))*\text{EllipticF}[(1/2)*\text{ArcSin}[\text{Rt}[-b/a, 2]*x], 2], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{GtQ}$

[a, 0] && NegQ[b/a]

### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c)], x]^p, x], x, b + 2\*c\*x, x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2)^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} - \frac{b^2 \int \frac{1}{(bx+cx^2)^{3/4}} dx}{12c} \\
 &= \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} - \frac{\left(b^2 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4}\right) \int \frac{1}{\left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{3/4}} dx}{12c (bx + cx^2)^{3/4}} \\
 &= \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} + \frac{\left(b^4 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4}\right) \text{Subst}\left(\int \frac{1}{\left(1 - \frac{b^2x^2}{c^2}\right)^{3/4}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{6\sqrt{2}c^3 (bx + cx^2)^{3/4}} \\
 &= \frac{(b + 2cx)\sqrt[4]{bx + cx^2}}{3c} - \frac{b^3 \left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} F\left(\frac{1}{2} \sin^{-1}\left(1 + \frac{2cx}{b}\right) \mid 2\right)}{3\sqrt{2}c^2 (bx + cx^2)^{3/4}}
 \end{aligned}$$

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.50

$$\int \sqrt[4]{bx + cx^2} dx = \frac{4x \sqrt[4]{x(b + cx)} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{5}{4}, \frac{9}{4}, -\frac{cx}{b}\right)}{5 \sqrt[4]{1 + \frac{cx}{b}}}$$

[In] Integrate[(b\*x + c\*x^2)^(1/4), x]

[Out] (4\*x\*(x\*(b + c\*x))^(1/4)\*Hypergeometric2F1[-1/4, 5/4, 9/4, -(c\*x)/b])/(5\*(1 + (c\*x)/b)^(1/4))

**Maple [F]**

$$\int (cx^2 + bx)^{\frac{1}{4}} dx$$

[In] int((c\*x^2+b\*x)^(1/4), x)

[Out] int((c\*x^2+b\*x)^(1/4), x)

**Fricas [F]**

$$\int \sqrt[4]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/4), x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(1/4), x)

**Sympy [F]**

$$\int \sqrt[4]{bx + cx^2} dx = \int \sqrt[4]{bx + cx^2} dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*(1/4), x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(1/4), x)

**Maxima [F]**

$$\int \sqrt[4]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(1/4), x)

**Giac [F]**

$$\int \sqrt[4]{bx + cx^2} dx = \int (cx^2 + bx)^{\frac{1}{4}} dx$$

[In] integrate((c\*x^2+b\*x)^(1/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(1/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.00 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.40

$$\int \sqrt[4]{bx + cx^2} dx = \frac{4x (cx^2 + bx)^{1/4} {}_2F_1\left(-\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; -\frac{cx}{b}\right)}{5\left(\frac{cx}{b} + 1\right)^{1/4}}$$

[In] int((b\*x + c\*x^2)^(1/4),x)

[Out] (4\*x\*(b\*x + c\*x^2)^(1/4)\*hypergeom([-1/4, 5/4], 9/4, -(c\*x)/b))/(5\*((c\*x)/b + 1)^(1/4))

### 3.43 $\int \frac{1}{\sqrt[4]{bx + cx^2}} dx$

Optimal result	278
Rubi [A] (verified)	278
Mathematica [C] (verified)	279
Maple [F]	280
Fricas [F]	280
Sympy [F]	280
Maxima [F]	280
Giac [F]	281
Mupad [B] (verification not implemented)	281

#### Optimal result

Integrand size = 13, antiderivative size = 58

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \frac{\sqrt{2}b^4 \sqrt{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \arcsin\left(1 + \frac{2cx}{b}\right) \middle| 2\right)}{c^4 \sqrt[4]{bx + cx^2}}$$

[Out]  $b*(-c*(c*x^2+b*x)/b^2)^(1/4)*(\cos(1/2*\arcsin(1+2*c*x/b)))^2^(1/2)/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticE}(\sin(1/2*\arcsin(1+2*c*x/b)), 2^(1/2))*2^(1/2)/c/(c*x^2+b*x)^(1/4)$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {636, 633, 234}

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \frac{\sqrt{2}b^4 \sqrt{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \arcsin\left(\frac{2cx}{b} + 1\right) \middle| 2\right)}{c^4 \sqrt[4]{bx + cx^2}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{-1/4}, x]$

[Out]  $(\text{Sqrt}[2]*b*(-((c*(b*x + c*x^2))/b^2))^(1/4)*\text{EllipticE}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/(c*(b*x + c*x^2)^(1/4))$

#### Rule 234

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1/4}, x\_Symbol] \rightarrow \text{Simp}[(2/(a^(1/4)*\text{Rt}[-b/a, 2]))*\text{EllipticE}[(1/2)*\text{ArcSin}[\text{Rt}[-b/a, 2]*x], 2], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{GtQ}$

[a, 0] && NegQ[b/a]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt[4]{-\frac{c(bx + cx^2)}{b^2}} \int \frac{1}{\sqrt[4]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}}} dx}{\sqrt[4]{bx + cx^2}} \\ &= -\frac{\left(b^2 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt[4]{1 - \frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{\sqrt{2}c^2 \sqrt[4]{bx + cx^2}} \\ &= \frac{\sqrt{2}b^4 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}} E\left(\frac{1}{2} \sin^{-1}\left(1 + \frac{2cx}{b}\right) \middle| 2\right)}{c \sqrt[4]{bx + cx^2}} \end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.78

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \frac{4x^4 \sqrt[4]{1 + \frac{cx}{b}} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{7}{4}, -\frac{cx}{b}\right)}{3 \sqrt[4]{x(b + cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-1/4),x]

[Out] (4\*x\*(1 + (c\*x)/b)^(1/4)\*Hypergeometric2F1[1/4, 3/4, 7/4, -((c\*x)/b)])/(3\*(x\*(b + c\*x))^(1/4))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{1}{4}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(1/4),x)

[Out] int(1/(c\*x^2+b\*x)^(1/4),x)

**Fricas [F]**

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/4),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(-1/4), x)

**Sympy [F]**

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \int \frac{1}{\sqrt[4]{bx + cx^2}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(1/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-1/4), x)

**Maxima [F]**

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-1/4), x)



**Giac [F]**

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{1}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(1/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-1/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.62

$$\int \frac{1}{\sqrt[4]{bx + cx^2}} dx = \frac{4x \left(\frac{cx}{b} + 1\right)^{1/4} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; \frac{7}{4}; -\frac{cx}{b}\right)}{3(cx^2 + bx)^{1/4}}$$

[In] int(1/(b\*x + c\*x^2)^(1/4),x)

[Out] (4\*x\*((c\*x)/b + 1)^(1/4)\*hypergeom([1/4, 3/4], 7/4, -(c\*x)/b))/(3\*(b\*x + c\*x^2)^(1/4))

### 3.44 $\int \frac{1}{(bx+cx^2)^{3/4}} dx$

Optimal result	282
Rubi [A] (verified)	282
Mathematica [C] (verified)	283
Maple [F]	284
Fricas [F]	284
Sympy [F]	284
Maxima [F]	284
Giac [F]	285
Mupad [B] (verification not implemented)	285

#### Optimal result

Integrand size = 13, antiderivative size = 59

$$\int \frac{1}{(bx+cx^2)^{3/4}} dx = \frac{2\sqrt{2}b\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(1 + \frac{2cx}{b}\right), 2\right)}{c(bx+cx^2)^{3/4}}$$

[Out]  $2*b*(-c*(c*x^2+b*x)/b^2)^{(3/4)}*(\cos(1/2*\arcsin(1+2*c*x/b))^2)^{(1/2)}/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticF}(\sin(1/2*\arcsin(1+2*c*x/b)), 2^{(1/2)})*2^{(1/2)}/c/(c*x^2+b*x)^{(3/4)}$

#### Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {636, 633, 238}

$$\int \frac{1}{(bx+cx^2)^{3/4}} dx = \frac{2\sqrt{2}b\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \text{EllipticF}\left(\frac{1}{2} \arcsin\left(\frac{2cx}{b} + 1\right), 2\right)}{c(bx+cx^2)^{3/4}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{-3/4}, x]$

[Out]  $(2*\text{Sqrt}[2]*b*(-((c*(b*x + c*x^2))/b^2))^{(3/4)}*\text{EllipticF}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/(c*(b*x + c*x^2)^{(3/4)})$

#### Rule 238

$\text{Int}[(a + (b \cdot x)^2)^{-3/4}, x\_Symbol] \rightarrow \text{Simp}[(2/(a^{3/4}*\text{Rt}[-b/a, 2]))*\text{EllipticF}[(1/2)*\text{ArcSin}[\text{Rt}[-b/a, 2]*x], 2], x] /;$   $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{GtQ}\{a, 0\} \ \&\& \ \text{NegQ}\{b/a\}$

Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c)], x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} \int \frac{1}{\left(-\frac{cx}{b} - \frac{c^2x^2}{b^2}\right)^{3/4}} dx}{(bx+cx^2)^{3/4}} \\ &= -\frac{\left(\sqrt{2}b^2\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4}\right) \text{Subst}\left(\int \frac{1}{\left(1-\frac{b^2x^2}{c^2}\right)^{3/4}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{c^2(bx+cx^2)^{3/4}} \\ &= \frac{2\sqrt{2}b\left(-\frac{c(bx+cx^2)}{b^2}\right)^{3/4} F\left(\frac{1}{2} \sin^{-1}\left(1 + \frac{2cx}{b}\right) \middle| 2\right)}{c(bx+cx^2)^{3/4}} \end{aligned}$$

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.73

$$\int \frac{1}{(bx+cx^2)^{3/4}} dx = \frac{4x\left(1 + \frac{cx}{b}\right)^{3/4} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{3}{4}, \frac{5}{4}, -\frac{cx}{b}\right)}{(x(b+cx))^{3/4}}$$

[In] Integrate[(b\*x + c\*x^2)^(-3/4), x]

[Out] (4\*x\*(1 + (c\*x)/b)^(3/4)\*Hypergeometric2F1[1/4, 3/4, 5/4, -(c\*x)/b])/(x\*(b + c\*x))^(3/4)

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{3}{4}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(3/4),x)

[Out] int(1/(c\*x^2+b\*x)^(3/4),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{3}{4}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{3}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(3/4),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(-3/4), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{3}{4}}} dx = \int \frac{1}{(bx + cx^2)^{\frac{3}{4}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(3/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-3/4), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{\frac{3}{4}}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{3}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(3/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-3/4), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{3/4}} dx = \int \frac{1}{(cx^2 + bx)^{3/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(3/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-3/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.61

$$\int \frac{1}{(bx + cx^2)^{3/4}} dx = \frac{4x \left(\frac{cx}{b} + 1\right)^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; \frac{5}{4}; -\frac{cx}{b}\right)}{(cx^2 + bx)^{3/4}}$$

[In] int(1/(b\*x + c\*x^2)^(3/4),x)

[Out] (4\*x\*((c\*x)/b + 1)^(3/4)\*hypergeom([1/4, 3/4], 5/4, -(c\*x)/b))/(b\*x + c\*x^2)^(3/4)

### 3.45 $\int \frac{1}{(bx+cx^2)^{5/4}} dx$

Optimal result	286
Rubi [A] (verified)	286
Mathematica [C] (verified)	288
Maple [F]	288
Fricas [F]	288
Sympy [F]	288
Maxima [F]	289
Giac [F]	289
Mupad [B] (verification not implemented)	289

#### Optimal result

Integrand size = 13, antiderivative size = 83

$$\int \frac{1}{(bx+cx^2)^{5/4}} dx = -\frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}} + \frac{4\sqrt{2}\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} E\left(\frac{1}{2}\arcsin\left(1+\frac{2cx}{b}\right)\middle|2\right)}{b^4\sqrt[4]{bx+cx^2}}$$

[Out]  $-4*(2*c*x+b)/b^2/(c*x^2+b*x)^(1/4)+4*(-c*(c*x^2+b*x)/b^2)^(1/4)*(\cos(1/2*\arcsin(1+2*c*x/b)))^(1/2)/\cos(1/2*\arcsin(1+2*c*x/b))*\text{EllipticE}(\sin(1/2*\arcsin(1+2*c*x/b)),2^(1/2))*2^(1/2)/b/(c*x^2+b*x)^(1/4)$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {628, 636, 633, 234}

$$\int \frac{1}{(bx+cx^2)^{5/4}} dx = \frac{4\sqrt{2}\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} E\left(\frac{1}{2}\arcsin\left(\frac{2cx}{b}+1\right)\middle|2\right)}{b^4\sqrt[4]{bx+cx^2}} - \frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{-5/4}, x]$

[Out]  $(-4*(b + 2*c*x))/(b^2*(b*x + c*x^2)^(1/4)) + (4*\text{Sqrt}[2]*(-((c*(b*x + c*x^2))/b^2))^(1/4)*\text{EllipticE}[\text{ArcSin}[1 + (2*c*x)/b]/2, 2])/(b*(b*x + c*x^2)^(1/4))$

#### Rule 234

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1/4}, x\_Symbol] \rightarrow \text{Simp}[(2/(a^(1/4)*\text{Rt}[-b/a, 2]))*\text{EllipticE}[(1/2)*\text{ArcSin}[\text{Rt}[-b/a, 2]*x], 2], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{GtQ}$

[a, 0] && NegQ[b/a]

### Rule 628

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p + 1) / ((p + 1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p + 3) / ((p + 1)\*(b^2 - 4\*a\*c))), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*c/(b^2 - 4\*a\*c)))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rule 636

Int[((b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p / ((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[(-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}} + \frac{(4c) \int \frac{1}{\sqrt[4]{bx+cx^2}} dx}{b^2} \\
 &= -\frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}} + \frac{\left(4c\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}}\right) \int \frac{1}{\sqrt[4]{-\frac{cx}{b}-\frac{c^2x^2}{b^2}}} dx}{b^2\sqrt[4]{bx+cx^2}} \\
 &= -\frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}} - \frac{\left(2\sqrt{2}\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-\frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b}-\frac{2c^2x}{b^2}\right)}{c\sqrt[4]{bx+cx^2}} \\
 &= -\frac{4(b+2cx)}{b^2\sqrt[4]{bx+cx^2}} + \frac{4\sqrt{2}\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} E\left(\frac{1}{2} \sin^{-1}\left(1+\frac{2cx}{b}\right) \middle| 2\right)}{b\sqrt[4]{bx+cx^2}}
 \end{aligned}$$

**Mathematica [C] (verified)**

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.54

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = -\frac{4\sqrt[4]{1 + \frac{cx}{b}} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{5}{4}, \frac{3}{4}, -\frac{cx}{b}\right)}{b\sqrt[4]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-5/4),x]

[Out] (-4\*(1 + (c\*x)/b)^(1/4)\*Hypergeometric2F1[-1/4, 5/4, 3/4, -((c\*x)/b)])/(b\*(x\*(b + c\*x))^(1/4))

**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{5/4}} dx$$

[In] int(1/(c\*x^2+b\*x)^(5/4),x)

[Out] int(1/(c\*x^2+b\*x)^(5/4),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = \int \frac{1}{(cx^2 + bx)^{5/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/4),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(3/4)/(c^2\*x^4 + 2\*b\*c\*x^3 + b^2\*x^2), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = \int \frac{1}{(bx + cx^2)^{5/4}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(5/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-5/4), x)



**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = \int \frac{1}{(cx^2 + bx)^{5/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-5/4), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = \int \frac{1}{(cx^2 + bx)^{5/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(5/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-5/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.43

$$\int \frac{1}{(bx + cx^2)^{5/4}} dx = -\frac{4x \left(\frac{cx}{b} + 1\right)^{5/4} {}_2F_1\left(-\frac{1}{4}, \frac{5}{4}; \frac{3}{4}; -\frac{cx}{b}\right)}{(cx^2 + bx)^{5/4}}$$

[In] int(1/(b\*x + c\*x^2)^(5/4),x)

[Out] -(4\*x\*((c\*x)/b + 1)^(5/4)\*hypergeom([-1/4, 5/4], 3/4, -(c\*x)/b))/(b\*x + c\*x^2)^(5/4)

$$3.46 \quad \int \frac{1}{(bx+cx^2)^{9/4}} dx$$

Optimal result	290
Rubi [A] (verified)	290
Mathematica [C] (verified)	292
Maple [F]	292
Fricas [F]	292
Sympy [F]	293
Maxima [F]	293
Giac [F]	293
Mupad [B] (verification not implemented)	293

### Optimal result

Integrand size = 13, antiderivative size = 115

$$\int \frac{1}{(bx+cx^2)^{9/4}} dx = -\frac{4(b+2cx)}{5b^2(bx+cx^2)^{5/4}} + \frac{48c(b+2cx)}{5b^4\sqrt[4]{bx+cx^2}} - \frac{48\sqrt{2}c^4\sqrt{-\frac{c(bx+cx^2)}{b^2}}E\left(\frac{1}{2}\arcsin\left(1+\frac{2cx}{b}\right)\middle|2\right)}{5b^3\sqrt[4]{bx+cx^2}}$$

[Out]  $-4/5*(2*c*x+b)/b^2/(c*x^2+b*x)^{(5/4)}+48/5*c*(2*c*x+b)/b^4/(c*x^2+b*x)^{(1/4)}$   
 $-48/5*c*(-c*(c*x^2+b*x)/b^2)^{(1/4)}*(\cos(1/2*\arcsin(1+2*c*x/b)))^2)^{(1/2)}/\cos$   
 $(1/2*\arcsin(1+2*c*x/b))*\text{EllipticE}(\sin(1/2*\arcsin(1+2*c*x/b)),2^{(1/2)})*2^{(1/2)}/b^3/(c*x^2+b*x)^{(1/4)}$

### Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {628, 636, 633, 234}

$$\int \frac{1}{(bx+cx^2)^{9/4}} dx = -\frac{48\sqrt{2}c^4\sqrt{-\frac{c(bx+cx^2)}{b^2}}E\left(\frac{1}{2}\arcsin\left(\frac{2cx}{b}+1\right)\middle|2\right)}{5b^3\sqrt[4]{bx+cx^2}} + \frac{48c(b+2cx)}{5b^4\sqrt[4]{bx+cx^2}} - \frac{4(b+2cx)}{5b^2(bx+cx^2)^{5/4}}$$

[In]  $\text{Int}[(b*x + c*x^2)^{-9/4}, x]$

[Out]  $(-4*(b + 2*c*x))/(5*b^2*(b*x + c*x^2)^{(5/4)}) + (48*c*(b + 2*c*x))/(5*b^4*(b*x + c*x^2)^{(1/4)}) - (48*sqrt[2]*c*(-((c*(b*x + c*x^2))/b^2))^{(1/4)}*EllipticE[ArcSin[1 + (2*c*x)/b]/2, 2])/(5*b^3*(b*x + c*x^2)^{(1/4)})$

#### Rule 234

Int[((a\_) + (b\_)\*(x\_)^2)^(-1/4), x\_Symbol] := Simp[(2/(a^(1/4)\*Rt[-b/a, 2]))\*EllipticE[(1/2)\*ArcSin[Rt[-b/a, 2]\*x], 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b/a]

#### Rule 628

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^(p + 1)/((p + 1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p + 3)/((p + 1)\*(b^2 - 4\*a\*c))), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

#### Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*c/(b^2 - 4\*a\*c)))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

#### Rule 636

Int[((b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[(b\*x + c\*x^2)^p/((-c)\*((b\*x + c\*x^2)/b^2))^p, Int[((-c)\*(x/b) - c^2\*(x^2/b^2))^p, x], x] /; FreeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4

#### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{4(b + 2cx)}{5b^2 (bx + cx^2)^{5/4}} - \frac{(12c) \int \frac{1}{(bx+cx^2)^{5/4}} dx}{5b^2} \\
 &= -\frac{4(b + 2cx)}{5b^2 (bx + cx^2)^{5/4}} + \frac{48c(b + 2cx)}{5b^4 \sqrt[4]{bx + cx^2}} - \frac{(48c^2) \int \frac{1}{\sqrt[4]{bx + cx^2}} dx}{5b^4} \\
 &= -\frac{4(b + 2cx)}{5b^2 (bx + cx^2)^{5/4}} + \frac{48c(b + 2cx)}{5b^4 \sqrt[4]{bx + cx^2}} - \frac{\left(48c^2 \sqrt[4]{-\frac{c(bx + cx^2)}{b^2}}\right) \int \frac{1}{\sqrt[4]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}}} dx}{5b^4 \sqrt[4]{bx + cx^2}}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{4(b+2cx)}{5b^2(bx+cx^2)^{5/4}} + \frac{48c(b+2cx)}{5b^4\sqrt[4]{bx+cx^2}} \\
&\quad + \frac{\left(24\sqrt{2}\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}}\right) \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-\frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2}\right)}{5b^2\sqrt[4]{bx+cx^2}} \\
&= -\frac{4(b+2cx)}{5b^2(bx+cx^2)^{5/4}} + \frac{48c(b+2cx)}{5b^4\sqrt[4]{bx+cx^2}} - \frac{48\sqrt{2}c\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} E\left(\frac{1}{2}\sin^{-1}\left(1+\frac{2cx}{b}\right)\middle|2\right)}{5b^3\sqrt[4]{bx+cx^2}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.43

$$\int \frac{1}{(bx+cx^2)^{9/4}} dx = -\frac{4\sqrt[4]{1+\frac{cx}{b}} \text{Hypergeometric2F1}\left(-\frac{5}{4}, \frac{9}{4}, -\frac{1}{4}, -\frac{cx}{b}\right)}{5b^2x\sqrt[4]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-9/4), x]

[Out] (-4\*(1 + (c\*x)/b)^(1/4)\*Hypergeometric2F1[-5/4, 9/4, -1/4, -((c\*x)/b)])/(5\*b^2\*x\*(x\*(b + c\*x))^(1/4))

### Maple [F]

$$\int \frac{1}{(cx^2 + bx)^{9/4}} dx$$

[In] int(1/(c\*x^2+b\*x)^(9/4), x)

[Out] int(1/(c\*x^2+b\*x)^(9/4), x)

### Fricas [F]

$$\int \frac{1}{(bx+cx^2)^{9/4}} dx = \int \frac{1}{(cx^2+bx)^{9/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(9/4), x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(3/4)/(c^3\*x^6 + 3\*b\*c^2\*x^5 + 3\*b^2\*c\*x^4 + b^3\*x^3), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{9/4}} dx = \int \frac{1}{(bx + cx^2)^{9/4}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(9/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-9/4), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{9/4}} dx = \int \frac{1}{(cx^2 + bx)^{9/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(9/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-9/4), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{9/4}} dx = \int \frac{1}{(cx^2 + bx)^{9/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(9/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-9/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.31

$$\int \frac{1}{(bx + cx^2)^{9/4}} dx = -\frac{4x \left(\frac{cx}{b} + 1\right)^{9/4} {}_2F_1\left(-\frac{5}{4}, \frac{9}{4}; -\frac{1}{4}; -\frac{cx}{b}\right)}{5(cx^2 + bx)^{9/4}}$$

[In] int(1/(b\*x + c\*x^2)^(9/4),x)

[Out] -(4\*x\*((c\*x)/b + 1)^(9/4)\*hypergeom([-5/4, 9/4], -1/4, -(c\*x)/b))/(5\*(b\*x + c\*x^2)^(9/4))

$$3.47 \quad \int \frac{1}{(bx+cx^2)^{13/4}} dx$$

Optimal result	294
Rubi [A] (verified)	294
Mathematica [C] (verified)	296
Maple [F]	297
Fricas [F]	297
Sympy [F]	297
Maxima [F]	297
Giac [F]	298
Mupad [B] (verification not implemented)	298

### Optimal result

Integrand size = 13, antiderivative size = 146

$$\int \frac{1}{(bx+cx^2)^{13/4}} dx = -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} + \frac{448\sqrt{2}c^2\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}}E\left(\frac{1}{2}\arcsin\left(1+\frac{2cx}{b}\right)\middle|2\right)}{15b^5\sqrt[4]{bx+cx^2}}$$

[Out]  $-4/9*(2*c*x+b)/b^2/(c*x^2+b*x)^(9/4)+112/45*c*(2*c*x+b)/b^4/(c*x^2+b*x)^(5/4)-448/15*c^2*(2*c*x+b)/b^6/(c*x^2+b*x)^(1/4)+448/15*c^2*(-c*(c*x^2+b*x)/b^2)^(1/4)*(cos(1/2*arcsin(1+2*c*x/b))^(1/2)/cos(1/2*arcsin(1+2*c*x/b))*EllipticE(sin(1/2*arcsin(1+2*c*x/b)),2^(1/2))*2^(1/2)/b^5/(c*x^2+b*x)^(1/4)$

### Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$ , Rules used = {628, 636, 633, 234}

$$\int \frac{1}{(bx+cx^2)^{13/4}} dx = \frac{448\sqrt{2}c^2\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}}E\left(\frac{1}{2}\arcsin\left(\frac{2cx}{b}+1\right)\middle|2\right)}{15b^5\sqrt[4]{bx+cx^2}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}}$$

[In] Int[(b\*x + c\*x^2)^(-13/4), x]

```
[Out] (-4*(b + 2*c*x))/(9*b^2*(b*x + c*x^2)^(9/4)) + (112*c*(b + 2*c*x))/(45*b^4*
(b*x + c*x^2)^(5/4)) - (448*c^2*(b + 2*c*x))/(15*b^6*(b*x + c*x^2)^(1/4)) +
(448*sqrt[2]*c^2*(-((c*(b*x + c*x^2))/b^2))^(1/4)*EllipticE[ArcSin[1 + (2*
c*x)/b]/2, 2])/(15*b^5*(b*x + c*x^2)^(1/4))
```

### Rule 234

```
Int[((a_) + (b_)*(x_)^2)^(-1/4), x_Symbol] := Simp[(2/(a^(1/4)*Rt[-b/a, 2]
))*EllipticE[(1/2)*ArcSin[Rt[-b/a, 2]*x], 2], x] /; FreeQ[{a, b}, x] && GtQ
[a, 0] && NegQ[b/a]
```

### Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

### Rule 636

```
Int[((b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[(b*x + c*x^2)^p/((-
c)*(b*x + c*x^2)/b^2))^p, Int[((-c)*(x/b) - c^2*(x^2/b^2))^p, x], x] /; Fr
eeQ[{b, c}, x] && RationalQ[p] && 3 <= Denominator[p] <= 4
```

### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} - \frac{(28c) \int \frac{1}{(bx+cx^2)^{9/4}} dx}{9b^2} \\
 &= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} + \frac{(112c^2) \int \frac{1}{(bx+cx^2)^{5/4}} dx}{15b^4} \\
 &= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} + \frac{(448c^3) \int \frac{1}{\sqrt[4]{bx+cx^2}} dx}{15b^6}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} \\
&\quad \left( 448c^3\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} \right) \int \frac{1}{\sqrt[4]{-\frac{cx}{b} - \frac{c^2x^2}{b^2}}} dx \\
&\quad + \frac{\hspace{15em}}{15b^6\sqrt[4]{bx+cx^2}} \\
&= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} \\
&\quad \left( 224\sqrt{2}c\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} \right) \text{Subst} \left( \int \frac{1}{\sqrt[4]{1 - \frac{b^2x^2}{c^2}}} dx, x, -\frac{c}{b} - \frac{2c^2x}{b^2} \right) \\
&\quad - \frac{\hspace{15em}}{15b^4\sqrt[4]{bx+cx^2}} \\
&= -\frac{4(b+2cx)}{9b^2(bx+cx^2)^{9/4}} + \frac{112c(b+2cx)}{45b^4(bx+cx^2)^{5/4}} - \frac{448c^2(b+2cx)}{15b^6\sqrt[4]{bx+cx^2}} \\
&\quad + \frac{448\sqrt{2}c^2\sqrt[4]{-\frac{c(bx+cx^2)}{b^2}} E\left(\frac{1}{2}\sin^{-1}\left(1 + \frac{2cx}{b}\right) \middle| 2\right)}{15b^5\sqrt[4]{bx+cx^2}}
\end{aligned}$$

### Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 10.02 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.34

$$\int \frac{1}{(bx+cx^2)^{13/4}} dx = -\frac{4\sqrt[4]{1 + \frac{cx}{b}} \text{Hypergeometric2F1}\left(-\frac{9}{4}, \frac{13}{4}, -\frac{5}{4}, -\frac{cx}{b}\right)}{9b^3x^2\sqrt[4]{x(b+cx)}}$$

[In] Integrate[(b\*x + c\*x^2)^(-13/4), x]

[Out] (-4\*(1 + (c\*x)/b)^(1/4)\*Hypergeometric2F1[-9/4, 13/4, -5/4, -((c\*x)/b)])/(9\*b^3\*x^2\*(x\*(b + c\*x))^(1/4))



**Maple [F]**

$$\int \frac{1}{(cx^2 + bx)^{\frac{13}{4}}} dx$$

[In] int(1/(c\*x^2+b\*x)^(13/4),x)

[Out] int(1/(c\*x^2+b\*x)^(13/4),x)

**Fricas [F]**

$$\int \frac{1}{(bx + cx^2)^{13/4}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{13}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(13/4),x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^(3/4)/(c^4\*x^8 + 4\*b\*c^3\*x^7 + 6\*b^2\*c^2\*x^6 + 4\*b^3\*c\*x^5 + b^4\*x^4), x)

**Sympy [F]**

$$\int \frac{1}{(bx + cx^2)^{13/4}} dx = \int \frac{1}{(bx + cx^2)^{\frac{13}{4}}} dx$$

[In] integrate(1/(c\*x\*\*2+b\*x)\*\*(13/4),x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*(-13/4), x)

**Maxima [F]**

$$\int \frac{1}{(bx + cx^2)^{13/4}} dx = \int \frac{1}{(cx^2 + bx)^{\frac{13}{4}}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(13/4),x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^(-13/4), x)

**Giac [F]**

$$\int \frac{1}{(bx + cx^2)^{13/4}} dx = \int \frac{1}{(cx^2 + bx)^{13/4}} dx$$

[In] integrate(1/(c\*x^2+b\*x)^(13/4),x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^(-13/4), x)

**Mupad [B] (verification not implemented)**

Time = 9.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.25

$$\int \frac{1}{(bx + cx^2)^{13/4}} dx = -\frac{4x \left(\frac{cx}{b} + 1\right)^{13/4} {}_2F_1\left(-\frac{9}{4}, \frac{13}{4}; -\frac{5}{4}; -\frac{cx}{b}\right)}{9 (cx^2 + bx)^{13/4}}$$

[In] int(1/(b\*x + c\*x^2)^(13/4),x)

[Out] -(4\*x\*((c\*x)/b + 1)^(13/4)\*hypergeom([-9/4, 13/4], -5/4, -(c\*x)/b))/(9\*(b\*x + c\*x^2)^(13/4))

### 3.48 $\int (bx + cx^2)^p dx$

Optimal result	299
Rubi [A] (verified)	299
Mathematica [A] (verified)	300
Maple [F]	300
Fricas [F]	300
Sympy [F]	300
Maxima [F]	301
Giac [F]	301
Mupad [B] (verification not implemented)	301

#### Optimal result

Integrand size = 11, antiderivative size = 55

$$\int (bx + cx^2)^p dx = -\frac{\left(-\frac{cx}{b}\right)^{-1-p} (bx + cx^2)^{1+p} \operatorname{Hypergeometric2F1}\left(-p, 1+p, 2+p, \frac{b+cx}{b}\right)}{b(1+p)}$$

[Out]  $-(c*x/b)^{-1-p}*(c*x^2+b*x)^{p+1}*hypergeom([-p, p+1], [2+p], (c*x+b)/b)/b/(p+1)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {638}

$$\int (bx + cx^2)^p dx = -\frac{\left(-\frac{cx}{b}\right)^{-p-1} (bx + cx^2)^{p+1} \operatorname{Hypergeometric2F1}\left(-p, p+1, p+2, \frac{b+cx}{b}\right)}{b(p+1)}$$

[In]  $\operatorname{Int}[(b*x + c*x^2)^p, x]$

[Out]  $-\left(\left(-\left(\frac{c*x}{b}\right)\right)^{-1-p}*(b*x + c*x^2)^{p+1}*Hypergeometric2F1[-p, 1+p, 2+p, (b+c*x)/b]\right)/(b*(1+p))$

#### Rule 638

$\operatorname{Int}[(a_. + (b_.)*(x_) + (c_.)*(x_)^2)^p, x\_Symbol] :> \operatorname{With}[\{q = \operatorname{Rt}[b^2 - 4*a*c, 2]\}, \operatorname{Simp}[(a + b*x + c*x^2)^{p+1}/(q*(p+1)*((q-b-2*c*x)/(2*q))^{p+1})]*\operatorname{Hypergeometric2F1}[-p, p+1, p+2, (b+q+2*c*x)/(2*q)], x] /;$   $\operatorname{FreeQ}\{a, b, c, p\}, x \ \&\& \ \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ !\operatorname{IntegerQ}[4*p]$

#### Rubi steps

$$\text{integral} = -\frac{\left(-\frac{cx}{b}\right)^{-1-p} (bx + cx^2)^{1+p} {}_2F_1\left(-p, 1+p; 2+p; \frac{b+cx}{b}\right)}{b(1+p)}$$

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.82

$$\int (bx + cx^2)^p dx = \frac{x(x(b + cx))^p \left(1 + \frac{cx}{b}\right)^{-p} \text{Hypergeometric2F1}\left(-p, 1 + p, 2 + p, -\frac{cx}{b}\right)}{1 + p}$$

[In] Integrate[(b\*x + c\*x^2)^p,x]

[Out] (x\*(x\*(b + c\*x))^p\*Hypergeometric2F1[-p, 1 + p, 2 + p, -(c\*x)/b])/((1 + p)\*(1 + (c\*x)/b)^p)

**Maple [F]**

$$\int (cx^2 + bx)^p dx$$

[In] int((c\*x^2+b\*x)^p,x)

[Out] int((c\*x^2+b\*x)^p,x)

**Fricas [F]**

$$\int (bx + cx^2)^p dx = \int (cx^2 + bx)^p dx$$

[In] integrate((c\*x^2+b\*x)^p,x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x)^p, x)

**Sympy [F]**

$$\int (bx + cx^2)^p dx = \int (bx + cx^2)^p dx$$

[In] integrate((c\*x\*\*2+b\*x)\*\*p,x)

[Out] Integral((b\*x + c\*x\*\*2)\*\*p, x)

**Maxima [F]**

$$\int (bx + cx^2)^p dx = \int (cx^2 + bx)^p dx$$

[In] integrate((c\*x^2+b\*x)^p,x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x)^p, x)

**Giac [F]**

$$\int (bx + cx^2)^p dx = \int (cx^2 + bx)^p dx$$

[In] integrate((c\*x^2+b\*x)^p,x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x)^p, x)

**Mupad [B] (verification not implemented)**

Time = 9.36 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.87

$$\int (bx + cx^2)^p dx = \frac{x(cx^2 + bx)^p {}_2F_1(-p, p + 1; p + 2; -\frac{cx}{b})}{(\frac{cx}{b} + 1)^p (p + 1)}$$

[In] int((b\*x + c\*x^2)^p,x)

[Out] (x\*(b\*x + c\*x^2)^p\*hypergeom([-p, p + 1], p + 2, -(c\*x)/b))/(((c\*x)/b + 1)^p\*(p + 1))

### 3.49 $\int (a + cx^2)^4 dx$

Optimal result	302
Rubi [A] (verified)	302
Mathematica [A] (verified)	303
Maple [A] (verified)	303
Fricas [A] (verification not implemented)	303
Sympy [A] (verification not implemented)	304
Maxima [A] (verification not implemented)	304
Giac [A] (verification not implemented)	304
Mupad [B] (verification not implemented)	304

#### Optimal result

Integrand size = 9, antiderivative size = 51

$$\int (a + cx^2)^4 dx = a^4x + \frac{4}{3}a^3cx^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{c^4x^9}{9}$$

[Out]  $a^4x + 4/3a^3cx^3 + 6/5a^2c^2x^5 + 4/7ac^3x^7 + 1/9c^4x^9$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {200}

$$\int (a + cx^2)^4 dx = a^4x + \frac{4}{3}a^3cx^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{c^4x^9}{9}$$

[In] Int[(a + c\*x^2)^4,x]

[Out]  $a^4x + (4a^3cx^3)/3 + (6a^2c^2x^5)/5 + (4ac^3x^7)/7 + (c^4x^9)/9$

#### Rule 200

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Int[ExpandIntegrand[(a + b\*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]

#### Rubi steps

$$\begin{aligned} \text{integral} &= \int (a^4 + 4a^3cx^2 + 6a^2c^2x^4 + 4ac^3x^6 + c^4x^8) dx \\ &= a^4x + \frac{4}{3}a^3cx^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{c^4x^9}{9} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00

$$\int (a + cx^2)^4 dx = a^4x + \frac{4}{3}a^3cx^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{c^4x^9}{9}$$

[In] Integrate[(a + c\*x^2)^4,x]

[Out] a^4\*x + (4\*a^3\*c\*x^3)/3 + (6\*a^2\*c^2\*x^5)/5 + (4\*a\*c^3\*x^7)/7 + (c^4\*x^9)/9

**Maple [A] (verified)**

Time = 2.13 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.86

method	result	size
gospers	$a^4x + \frac{4}{3}ca^3x^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{1}{9}c^4x^9$	44
default	$a^4x + \frac{4}{3}ca^3x^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{1}{9}c^4x^9$	44
norman	$a^4x + \frac{4}{3}ca^3x^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{1}{9}c^4x^9$	44
risch	$a^4x + \frac{4}{3}ca^3x^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{1}{9}c^4x^9$	44
parallelrisch	$a^4x + \frac{4}{3}ca^3x^3 + \frac{6}{5}a^2c^2x^5 + \frac{4}{7}ac^3x^7 + \frac{1}{9}c^4x^9$	44

[In] int((c\*x^2+a)^4,x,method=\_RETURNVERBOSE)

[Out] a^4\*x+4/3\*c\*a^3\*x^3+6/5\*a^2\*c^2\*x^5+4/7\*a\*c^3\*x^7+1/9\*c^4\*x^9

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^4 dx = \frac{1}{9}c^4x^9 + \frac{4}{7}ac^3x^7 + \frac{6}{5}a^2c^2x^5 + \frac{4}{3}a^3cx^3 + a^4x$$

[In] integrate((c\*x^2+a)^4,x, algorithm="fricas")

[Out] 1/9\*c^4\*x^9 + 4/7\*a\*c^3\*x^7 + 6/5\*a^2\*c^2\*x^5 + 4/3\*a^3\*c\*x^3 + a^4\*x

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.96

$$\int (a + cx^2)^4 dx = a^4x + \frac{4a^3cx^3}{3} + \frac{6a^2c^2x^5}{5} + \frac{4ac^3x^7}{7} + \frac{c^4x^9}{9}$$

[In] integrate((c\*x\*\*2+a)\*\*4,x)

[Out] a\*\*4\*x + 4\*a\*\*3\*c\*x\*\*3/3 + 6\*a\*\*2\*c\*\*2\*x\*\*5/5 + 4\*a\*c\*\*3\*x\*\*7/7 + c\*\*4\*x\*\*9/9

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^4 dx = \frac{1}{9}c^4x^9 + \frac{4}{7}ac^3x^7 + \frac{6}{5}a^2c^2x^5 + \frac{4}{3}a^3cx^3 + a^4x$$

[In] integrate((c\*x^2+a)^4,x, algorithm="maxima")

[Out] 1/9\*c^4\*x^9 + 4/7\*a\*c^3\*x^7 + 6/5\*a^2\*c^2\*x^5 + 4/3\*a^3\*c\*x^3 + a^4\*x

**Giac [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^4 dx = \frac{1}{9}c^4x^9 + \frac{4}{7}ac^3x^7 + \frac{6}{5}a^2c^2x^5 + \frac{4}{3}a^3cx^3 + a^4x$$

[In] integrate((c\*x^2+a)^4,x, algorithm="giac")

[Out] 1/9\*c^4\*x^9 + 4/7\*a\*c^3\*x^7 + 6/5\*a^2\*c^2\*x^5 + 4/3\*a^3\*c\*x^3 + a^4\*x

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^4 dx = a^4x + \frac{4a^3cx^3}{3} + \frac{6a^2c^2x^5}{5} + \frac{4ac^3x^7}{7} + \frac{c^4x^9}{9}$$

[In] int((a + c\*x^2)^4,x)

[Out] a^4\*x + (c^4\*x^9)/9 + (4\*a^3\*c\*x^3)/3 + (4\*a\*c^3\*x^7)/7 + (6\*a^2\*c^2\*x^5)/5



### 3.50 $\int (a + cx^2)^3 dx$

Optimal result	305
Rubi [A] (verified)	305
Mathematica [A] (verified)	306
Maple [A] (verified)	306
Fricas [A] (verification not implemented)	306
Sympy [A] (verification not implemented)	307
Maxima [A] (verification not implemented)	307
Giac [A] (verification not implemented)	307
Mupad [B] (verification not implemented)	307

#### Optimal result

Integrand size = 9, antiderivative size = 35

$$\int (a + cx^2)^3 dx = a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{c^3x^7}{7}$$

[Out]  $a^3x + a^2cx^3 + 3/5*a*c^2*x^5 + 1/7*c^3*x^7$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {200}

$$\int (a + cx^2)^3 dx = a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{c^3x^7}{7}$$

[In] Int[(a + c\*x^2)^3, x]

[Out]  $a^3x + a^2cx^3 + (3*a*c^2*x^5)/5 + (c^3*x^7)/7$

#### Rule 200

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] :> Int[ExpandIntegrand[(a + b\*x^n)^p, x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && IGtQ[p, 0]

#### Rubi steps

$$\begin{aligned} \text{integral} &= \int (a^3 + 3a^2cx^2 + 3ac^2x^4 + c^3x^6) dx \\ &= a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{c^3x^7}{7} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (a + cx^2)^3 dx = a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{c^3x^7}{7}$$

[In] Integrate[(a + c\*x^2)^3,x]

[Out] a^3\*x + a^2\*c\*x^3 + (3\*a\*c^2\*x^5)/5 + (c^3\*x^7)/7

**Maple [A] (verified)**

Time = 2.14 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
gospers	$a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{1}{7}c^3x^7$	32
default	$a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{1}{7}c^3x^7$	32
norman	$a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{1}{7}c^3x^7$	32
risch	$a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{1}{7}c^3x^7$	32
parallexrisch	$a^3x + a^2cx^3 + \frac{3}{5}ac^2x^5 + \frac{1}{7}c^3x^7$	32

[In] int((c\*x^2+a)^3,x,method=\_RETURNVERBOSE)

[Out] a^3\*x+a^2\*c\*x^3+3/5\*a\*c^2\*x^5+1/7\*c^3\*x^7

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int (a + cx^2)^3 dx = \frac{1}{7}c^3x^7 + \frac{3}{5}ac^2x^5 + a^2cx^3 + a^3x$$

[In] integrate((c\*x^2+a)^3,x, algorithm="fricas")

[Out] 1/7\*c^3\*x^7 + 3/5\*a\*c^2\*x^5 + a^2\*c\*x^3 + a^3\*x

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int (a + cx^2)^3 dx = a^3x + a^2cx^3 + \frac{3ac^2x^5}{5} + \frac{c^3x^7}{7}$$

[In] integrate((c\*x\*\*2+a)\*\*3,x)

[Out] a\*\*3\*x + a\*\*2\*c\*x\*\*3 + 3\*a\*c\*\*2\*x\*\*5/5 + c\*\*3\*x\*\*7/7

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int (a + cx^2)^3 dx = \frac{1}{7}c^3x^7 + \frac{3}{5}ac^2x^5 + a^2cx^3 + a^3x$$

[In] integrate((c\*x^2+a)^3,x, algorithm="maxima")

[Out] 1/7\*c^3\*x^7 + 3/5\*a\*c^2\*x^5 + a^2\*c\*x^3 + a^3\*x

**Giac [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int (a + cx^2)^3 dx = \frac{1}{7}c^3x^7 + \frac{3}{5}ac^2x^5 + a^2cx^3 + a^3x$$

[In] integrate((c\*x^2+a)^3,x, algorithm="giac")

[Out] 1/7\*c^3\*x^7 + 3/5\*a\*c^2\*x^5 + a^2\*c\*x^3 + a^3\*x

**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int (a + cx^2)^3 dx = a^3x + a^2cx^3 + \frac{3ac^2x^5}{5} + \frac{c^3x^7}{7}$$

[In] int((a + c\*x^2)^3,x)

[Out] a^3\*x + (c^3\*x^7)/7 + a^2\*c\*x^3 + (3\*a\*c^2\*x^5)/5

### 3.51 $\int (a + cx^2)^2 dx$

Optimal result . . . . .	308
Rubi [A] (verified) . . . . .	308
Mathematica [A] (verified) . . . . .	309
Maple [A] (verified) . . . . .	309
Fricas [A] (verification not implemented) . . . . .	309
Sympy [A] (verification not implemented) . . . . .	310
Maxima [A] (verification not implemented) . . . . .	310
Giac [A] (verification not implemented) . . . . .	310
Mupad [B] (verification not implemented) . . . . .	310

#### Optimal result

Integrand size = 9, antiderivative size = 25

$$\int (a + cx^2)^2 dx = a^2x + \frac{2}{3}acx^3 + \frac{c^2x^5}{5}$$

[Out]  $a^2x + 2/3acx^3 + 1/5c^2x^5$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {200}

$$\int (a + cx^2)^2 dx = a^2x + \frac{2}{3}acx^3 + \frac{c^2x^5}{5}$$

[In]  $\text{Int}[(a + c*x^2)^2, x]$

[Out]  $a^2x + (2*a*c*x^3)/3 + (c^2*x^5)/5$

#### Rule 200

$\text{Int}[(a + b*x^n)^p, x] /; \text{FreeQ}\{a, b, x\} \ \&\amp; \ \text{IGtQ}[n, 0] \ \&\amp; \ \text{IGtQ}[p, 0]$

#### Rubi steps

$$\begin{aligned} \text{integral} &= \int (a^2 + 2acx^2 + c^2x^4) dx \\ &= a^2x + \frac{2}{3}acx^3 + \frac{c^2x^5}{5} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int (a + cx^2)^2 dx = a^2x + \frac{2}{3}acx^3 + \frac{c^2x^5}{5}$$

[In] Integrate[(a + c\*x^2)^2,x]

[Out] a^2\*x + (2\*a\*c\*x^3)/3 + (c^2\*x^5)/5

**Maple [A] (verified)**

Time = 2.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
gospers	$a^2x + \frac{2}{3}acx^3 + \frac{1}{5}x^5c^2$	22
default	$a^2x + \frac{2}{3}acx^3 + \frac{1}{5}x^5c^2$	22
norman	$a^2x + \frac{2}{3}acx^3 + \frac{1}{5}x^5c^2$	22
risch	$a^2x + \frac{2}{3}acx^3 + \frac{1}{5}x^5c^2$	22
parallelrisch	$a^2x + \frac{2}{3}acx^3 + \frac{1}{5}x^5c^2$	22

[In] int((c\*x^2+a)^2,x,method=\_RETURNVERBOSE)

[Out] a^2\*x+2/3\*a\*c\*x^3+1/5\*x^5\*c^2

**Fricas [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^2 dx = \frac{1}{5}c^2x^5 + \frac{2}{3}acx^3 + a^2x$$

[In] integrate((c\*x^2+a)^2,x, algorithm="fricas")

[Out] 1/5\*c^2\*x^5 + 2/3\*a\*c\*x^3 + a^2\*x

**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int (a + cx^2)^2 dx = a^2x + \frac{2acx^3}{3} + \frac{c^2x^5}{5}$$

[In] integrate((c\*x\*\*2+a)\*\*2,x)

[Out] a\*\*2\*x + 2\*a\*c\*x\*\*3/3 + c\*\*2\*x\*\*5/5

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^2 dx = \frac{1}{5}c^2x^5 + \frac{2}{3}acx^3 + a^2x$$

[In] integrate((c\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/5\*c^2\*x^5 + 2/3\*a\*c\*x^3 + a^2\*x

**Giac [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^2 dx = \frac{1}{5}c^2x^5 + \frac{2}{3}acx^3 + a^2x$$

[In] integrate((c\*x^2+a)^2,x, algorithm="giac")

[Out] 1/5\*c^2\*x^5 + 2/3\*a\*c\*x^3 + a^2\*x

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int (a + cx^2)^2 dx = a^2x + \frac{2acx^3}{3} + \frac{c^2x^5}{5}$$

[In] int((a + c\*x^2)^2,x)

[Out] a^2\*x + (c^2\*x^5)/5 + (2\*a\*c\*x^3)/3

## 3.52 $\int (a + cx^2) dx$

Optimal result	311
Rubi [A] (verified)	311
Mathematica [A] (verified)	312
Maple [A] (verified)	312
Fricas [A] (verification not implemented)	312
Sympy [A] (verification not implemented)	313
Maxima [A] (verification not implemented)	313
Giac [A] (verification not implemented)	313
Mupad [B] (verification not implemented)	313

### Optimal result

Integrand size = 7, antiderivative size = 12

$$\int (a + cx^2) dx = ax + \frac{cx^3}{3}$$

[Out] a\*x+1/3\*c\*x^3

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 0,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$ , Rules used = {}

$$\int (a + cx^2) dx = ax + \frac{cx^3}{3}$$

[In] Int[a + c\*x^2,x]

[Out] a\*x + (c\*x^3)/3

Rubi steps

$$\text{integral} = ax + \frac{cx^3}{3}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int (a + cx^2) dx = ax + \frac{cx^3}{3}$$

[In] Integrate[a + c\*x^2,x]

[Out] a\*x + (c\*x^3)/3

**Maple [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.92

method	result	size
gospers	$ax + \frac{1}{3}cx^3$	11
default	$ax + \frac{1}{3}cx^3$	11
norman	$ax + \frac{1}{3}cx^3$	11
risch	$ax + \frac{1}{3}cx^3$	11
parallelrisch	$ax + \frac{1}{3}cx^3$	11
parts	$ax + \frac{1}{3}cx^3$	11

[In] int(c\*x^2+a,x,method=\_RETURNVERBOSE)

[Out] a\*x+1/3\*c\*x^3

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + cx^2) dx = \frac{1}{3}cx^3 + ax$$

[In] integrate(c\*x^2+a,x, algorithm="fricas")

[Out] 1/3\*c\*x^3 + a\*x



**Sympy [A] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int (a + cx^2) dx = ax + \frac{cx^3}{3}$$

[In] integrate(c\*x\*\*2+a,x)

[Out] a\*x + c\*x\*\*3/3

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + cx^2) dx = \frac{1}{3} cx^3 + ax$$

[In] integrate(c\*x^2+a,x, algorithm="maxima")

[Out] 1/3\*c\*x^3 + a\*x

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + cx^2) dx = \frac{1}{3} cx^3 + ax$$

[In] integrate(c\*x^2+a,x, algorithm="giac")

[Out] 1/3\*c\*x^3 + a\*x

**Mupad [B] (verification not implemented)**

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int (a + cx^2) dx = \frac{cx^3}{3} + ax$$

[In] int(a + c\*x^2,x)

[Out] a\*x + (c\*x^3)/3

### 3.53 $\int \frac{1}{a+cx^2} dx$

Optimal result	314
Rubi [A] (verified)	314
Mathematica [A] (verified)	315
Maple [A] (verified)	315
Fricas [A] (verification not implemented)	315
Sympy [B] (verification not implemented)	316
Maxima [A] (verification not implemented)	316
Giac [A] (verification not implemented)	316
Mupad [B] (verification not implemented)	317

#### Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \frac{1}{a+cx^2} dx = \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{c}}$$

[Out]  $\arctan(x*c^{(1/2)}/a^{(1/2)})/a^{(1/2)}/c^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.111$ , Rules used = {211}

$$\int \frac{1}{a+cx^2} dx = \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{c}}$$

[In]  $\text{Int}[(a + c*x^2)^{-1}, x]$

[Out]  $\text{ArcTan}[(\text{Sqrt}[c]*x)/\text{Sqrt}[a]]/(\text{Sqrt}[a]*\text{Sqrt}[c])$

#### Rule 211

$\text{Int}[(a_+ + (b_+)*x^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b]$

#### Rubi steps

$$\text{integral} = \frac{\tan^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{c}}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + cx^2} dx = \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{c}}$$

[In] Integrate[(a + c\*x^2)^(-1),x]

[Out] ArcTan[(Sqrt[c]\*x)/Sqrt[a]]/(Sqrt[a]\*Sqrt[c])

**Maple [A] (verified)**

Time = 2.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{\sqrt{ac}}$	16
risch	$-\frac{\ln(cx+\sqrt{-ac})}{2\sqrt{-ac}} + \frac{\ln(-cx+\sqrt{-ac})}{2\sqrt{-ac}}$	41

[In] int(1/(c\*x^2+a),x,method=\_RETURNVERBOSE)

[Out] 1/(a\*c)^(1/2)\*arctan(c\*x/(a\*c)^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.79

$$\int \frac{1}{a + cx^2} dx = \left[ -\frac{\sqrt{-ac} \log\left(\frac{cx^2 - 2\sqrt{-ac}x - a}{cx^2 + a}\right)}{2ac}, \frac{\sqrt{ac} \arctan\left(\frac{\sqrt{ac}x}{a}\right)}{ac} \right]$$

[In] integrate(1/(c\*x^2+a),x, algorithm="fricas")

[Out] [-1/2\*sqrt(-a\*c)\*log((c\*x^2 - 2\*sqrt(-a\*c)\*x - a)/(c\*x^2 + a))/(a\*c), sqrt(a\*c)\*arctan(sqrt(a\*c)\*x/a)/(a\*c)]

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(22) = 44.

Time = 0.06 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.21

$$\int \frac{1}{a + cx^2} dx = -\frac{\sqrt{-\frac{1}{ac}} \log\left(-a\sqrt{-\frac{1}{ac}} + x\right)}{2} + \frac{\sqrt{-\frac{1}{ac}} \log\left(a\sqrt{-\frac{1}{ac}} + x\right)}{2}$$

[In] integrate(1/(c\*x\*\*2+a),x)

[Out] -sqrt(-1/(a\*c))\*log(-a\*sqrt(-1/(a\*c)) + x)/2 + sqrt(-1/(a\*c))\*log(a\*sqrt(-1/(a\*c)) + x)/2

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \frac{1}{a + cx^2} dx = \frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{\sqrt{ac}}$$

[In] integrate(1/(c\*x^2+a),x, algorithm="maxima")

[Out] arctan(c\*x/sqrt(a\*c))/sqrt(a\*c)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int \frac{1}{a + cx^2} dx = \frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{\sqrt{ac}}$$

[In] integrate(1/(c\*x^2+a),x, algorithm="giac")

[Out] arctan(c\*x/sqrt(a\*c))/sqrt(a\*c)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.67

$$\int \frac{1}{a + cx^2} dx = \frac{\operatorname{atan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{c}}$$

[In] `int(1/(a + c*x^2),x)`

[Out] `atan((c^(1/2)*x)/a^(1/2))/(a^(1/2)*c^(1/2))`

### 3.54 $\int \frac{1}{(a+cx^2)^2} dx$

Optimal result	318
Rubi [A] (verified)	318
Mathematica [A] (verified)	319
Maple [A] (verified)	319
Fricas [A] (verification not implemented)	320
Sympy [B] (verification not implemented)	320
Maxima [A] (verification not implemented)	320
Giac [A] (verification not implemented)	321
Mupad [B] (verification not implemented)	321

#### Optimal result

Integrand size = 9, antiderivative size = 45

$$\int \frac{1}{(a+cx^2)^2} dx = \frac{x}{2a(a+cx^2)} + \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{c}}$$

[Out] 1/2\*x/a/(c\*x^2+a)+1/2\*arctan(x\*c^(1/2)/a^(1/2))/a^(3/2)/c^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {205, 211}

$$\int \frac{1}{(a+cx^2)^2} dx = \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{c}} + \frac{x}{2a(a+cx^2)}$$

[In] Int[(a + c\*x^2)^(-2), x]

[Out] x/(2\*a\*(a + c\*x^2)) + ArcTan[(Sqrt[c]\*x)/Sqrt[a]]/(2\*a^(3/2)\*Sqrt[c])

#### Rule 205

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])
```

#### Rule 211

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(Rt[a/b, 2]/a)\*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{2a(a+cx^2)} + \frac{\int \frac{1}{a+cx^2} dx}{2a} \\ &= \frac{x}{2a(a+cx^2)} + \frac{\tan^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{c}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a+cx^2)^2} dx = \frac{x}{2a(a+cx^2)} + \frac{\arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{c}}$$

[In] Integrate[(a + c\*x^2)^(-2),x]

[Out] x/(2\*a\*(a + c\*x^2)) + ArcTan[(Sqrt[c]\*x)/Sqrt[a]]/(2\*a^(3/2)\*Sqrt[c])

**Maple [A] (verified)**

Time = 2.14 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

method	result	size
default	$\frac{x}{2a(cx^2+a)} + \frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{2a\sqrt{ac}}$	36
risch	$\frac{x}{2a(cx^2+a)} - \frac{\ln(cx+\sqrt{-ac})}{4\sqrt{-aca}} + \frac{\ln(-cx+\sqrt{-ac})}{4\sqrt{-aca}}$	62

[In] int(1/(c\*x^2+a)^2,x,method=\_RETURNVERBOSE)

[Out] 1/2\*x/a/(c\*x^2+a)+1/2/a/(a\*c)^(1/2)\*arctan(c\*x/(a\*c)^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.55 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.67

$$\int \frac{1}{(a + cx^2)^2} dx$$

$$= \left[ \frac{2acx - (cx^2 + a)\sqrt{-ac} \log\left(\frac{cx^2 - 2\sqrt{-ac}x - a}{cx^2 + a}\right)}{4(a^2c^2x^2 + a^3c)}, \frac{acx + (cx^2 + a)\sqrt{ac} \arctan\left(\frac{\sqrt{ac}x}{a}\right)}{2(a^2c^2x^2 + a^3c)} \right]$$

[In] integrate(1/(c\*x^2+a)^2,x, algorithm="fricas")

[Out] [1/4\*(2\*a\*c\*x - (c\*x^2 + a)\*sqrt(-a\*c)\*log((c\*x^2 - 2\*sqrt(-a\*c)\*x - a)/(c\*x^2 + a)))/(a^2\*c^2\*x^2 + a^3\*c), 1/2\*(a\*c\*x + (c\*x^2 + a)\*sqrt(a\*c)\*arctan(sqrt(a\*c)\*x/a))/(a^2\*c^2\*x^2 + a^3\*c)]

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(36) = 72.

Time = 0.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.73

$$\int \frac{1}{(a + cx^2)^2} dx$$

$$= \frac{x}{2a^2 + 2acx^2} - \frac{\sqrt{-\frac{1}{a^3c}} \log\left(-a^2\sqrt{-\frac{1}{a^3c}} + x\right)}{4} + \frac{\sqrt{-\frac{1}{a^3c}} \log\left(a^2\sqrt{-\frac{1}{a^3c}} + x\right)}{4}$$

[In] integrate(1/(c\*x\*\*2+a)\*\*2,x)

[Out] x/(2\*a\*\*2 + 2\*a\*c\*x\*\*2) - sqrt(-1/(a\*\*3\*c))\*log(-a\*\*2\*sqrt(-1/(a\*\*3\*c)) + x)/4 + sqrt(-1/(a\*\*3\*c))\*log(a\*\*2\*sqrt(-1/(a\*\*3\*c)) + x)/4

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + cx^2)^2} dx = \frac{x}{2(acx^2 + a^2)} + \frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{2\sqrt{aca}}$$

[In] integrate(1/(c\*x^2+a)^2,x, algorithm="maxima")

[Out] 1/2\*x/(a\*c\*x^2 + a^2) + 1/2\*arctan(c\*x/sqrt(a\*c))/(sqrt(a\*c)\*a)



**Giac [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

$$\int \frac{1}{(a + cx^2)^2} dx = \frac{\arctan\left(\frac{cx}{\sqrt{ac}}\right)}{2\sqrt{aca}} + \frac{x}{2(cx^2 + a)a}$$

[In] integrate(1/(c\*x^2+a)^2,x, algorithm="giac")

[Out] 1/2\*arctan(c\*x/sqrt(a\*c))/(sqrt(a\*c)\*a) + 1/2\*x/((c\*x^2 + a)\*a)

**Mupad [B] (verification not implemented)**

Time = 9.20 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a + cx^2)^2} dx = \frac{x}{2a(cx^2 + a)} + \frac{\operatorname{atan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)}{2a^{3/2}\sqrt{c}}$$

[In] int(1/(a + c\*x^2)^2,x)

[Out] x/(2\*a\*(a + c\*x^2)) + atan((c^(1/2)\*x)/a^(1/2))/(2\*a^(3/2)\*c^(1/2))

### 3.55 $\int \frac{1}{(a+cx^2)^3} dx$

Optimal result	322
Rubi [A] (verified)	322
Mathematica [A] (verified)	323
Maple [A] (verified)	323
Fricas [A] (verification not implemented)	324
Sympy [A] (verification not implemented)	324
Maxima [A] (verification not implemented)	324
Giac [A] (verification not implemented)	325
Mupad [B] (verification not implemented)	325

#### Optimal result

Integrand size = 9, antiderivative size = 62

$$\int \frac{1}{(a+cx^2)^3} dx = \frac{x}{4a(a+cx^2)^2} + \frac{3x}{8a^2(a+cx^2)} + \frac{3 \arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{c}}$$

[Out] 1/4\*x/a/(c\*x^2+a)^2+3/8\*x/a^2/(c\*x^2+a)+3/8\*arctan(x\*c^(1/2)/a^(1/2))/a^(5/2)/c^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$ , Rules used = {205, 211}

$$\int \frac{1}{(a+cx^2)^3} dx = \frac{3 \arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{c}} + \frac{3x}{8a^2(a+cx^2)} + \frac{x}{4a(a+cx^2)^2}$$

[In] Int[(a + c\*x^2)^(-3),x]

[Out] x/(4\*a\*(a + c\*x^2)^2) + (3\*x)/(8\*a^2\*(a + c\*x^2)) + (3\*ArcTan[(Sqrt[c]\*x)/Sqrt[a]])/(8\*a^(5/2)\*Sqrt[c])

#### Rule 205

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[p, -1] && (IntegerQ[2*p] || (n == 2 && IntegerQ[4*p]) || (n == 2 && IntegerQ[3*p]) || Denominator[p + 1/n] < Denominator[p])
```

## Rule 211

$\text{Int}[(a_+ + (b_+)(x_+)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a) * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{PosQ}[a/b]$

## Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{4a(a+cx^2)^2} + \frac{3 \int \frac{1}{(a+cx^2)^2} dx}{4a} \\ &= \frac{x}{4a(a+cx^2)^2} + \frac{3x}{8a^2(a+cx^2)} + \frac{3 \int \frac{1}{a+cx^2} dx}{8a^2} \\ &= \frac{x}{4a(a+cx^2)^2} + \frac{3x}{8a^2(a+cx^2)} + \frac{3 \tan^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{c}} \end{aligned}$$

## Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a+cx^2)^3} dx = \frac{5ax + 3cx^3}{8a^2(a+cx^2)^2} + \frac{3 \arctan\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{c}}$$

[In] Integrate[(a + c\*x^2)^(-3),x]

[Out] (5\*a\*x + 3\*c\*x^3)/(8\*a^2\*(a + c\*x^2)^2) + (3\*ArcTan[(Sqrt[c]\*x)/Sqrt[a]])/(8\*a^(5/2)\*Sqrt[c])

## Maple [A] (verified)

Time = 2.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{x}{4a(cx^2+a)^2} + \frac{\frac{3x}{8a(cx^2+a)} + \frac{3 \arctan\left(\frac{cx}{\sqrt{ac}}\right)}{8a\sqrt{ac}}}{a}$	57
risch	$\frac{\frac{3cx^3}{8a^2} + \frac{5x}{8a}}{(cx^2+a)^2} - \frac{3 \ln(cx+\sqrt{-ac})}{16\sqrt{-ac}a^2} + \frac{3 \ln(-cx+\sqrt{-ac})}{16\sqrt{-ac}a^2}$	73

[In] int(1/(c\*x^2+a)^3,x,method=\_RETURNVERBOSE)

[Out] 1/4\*x/a/(c\*x^2+a)^2+3/4/a\*(1/2\*x/a/(c\*x^2+a)+1/2/a/(a\*c)^(1/2)\*arctan(c\*x/(a\*c)^(1/2)))

**Fricas [A] (verification not implemented)**

none

Time = 0.47 (sec) , antiderivative size = 188, normalized size of antiderivative = 3.03

$$\int \frac{1}{(a + cx^2)^3} dx = \left[ \frac{6ac^2x^3 + 10a^2cx - 3(c^2x^4 + 2acx^2 + a^2)\sqrt{-ac} \log\left(\frac{cx^2 - 2\sqrt{-ac}x - a}{cx^2 + a}\right)}{16(a^3c^3x^4 + 2a^4c^2x^2 + a^5c)}, \frac{3ac^2x^3 + 5a^2cx + 3(c^2x^4 + 2acx^2 + a^5c)}{8(a^3c^3x^4 + 2a^4c^2x^2 + a^5c)} \right]$$

```
[In] integrate(1/(c*x^2+a)^3,x, algorithm="fricas")
```

```
[Out] [1/16*(6*a*c^2*x^3 + 10*a^2*c*x - 3*(c^2*x^4 + 2*a*c*x^2 + a^2)*sqrt(-a*c)*
log((c*x^2 - 2*sqrt(-a*c)*x - a)/(c*x^2 + a)))/(a^3*c^3*x^4 + 2*a^4*c^2*x^2
+ a^5*c), 1/8*(3*a*c^2*x^3 + 5*a^2*c*x + 3*(c^2*x^4 + 2*a*c*x^2 + a^2)*sq
r t(a*c)*arctan(sqrt(a*c)*x/a))/(a^3*c^3*x^4 + 2*a^4*c^2*x^2 + a^5*c)]
```

**Sympy [A] (verification not implemented)**

Time = 0.17 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.69

$$\int \frac{1}{(a + cx^2)^3} dx = -\frac{3\sqrt{-\frac{1}{a^5c}} \log\left(-a^3\sqrt{-\frac{1}{a^5c}} + x\right)}{16} + \frac{3\sqrt{-\frac{1}{a^5c}} \log\left(a^3\sqrt{-\frac{1}{a^5c}} + x\right)}{16} + \frac{5ax + 3cx^3}{8a^4 + 16a^3cx^2 + 8a^2c^2x^4}$$

```
[In] integrate(1/(c*x**2+a)**3,x)
```

```
[Out] -3*sqrt(-1/(a**5*c))*log(-a**3*sqrt(-1/(a**5*c)) + x)/16 + 3*sqrt(-1/(a**5*
c))*log(a**3*sqrt(-1/(a**5*c)) + x)/16 + (5*a*x + 3*c*x**3)/(8*a**4 + 16*a*
*3*c*x**2 + 8*a**2*c**2*x**4)
```

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + cx^2)^3} dx = \frac{3cx^3 + 5ax}{8(a^2c^2x^4 + 2a^3cx^2 + a^4)} + \frac{3 \arctan\left(\frac{cx}{\sqrt{ac}}\right)}{8\sqrt{aca^2}}$$

```
[In] integrate(1/(c*x^2+a)^3,x, algorithm="maxima")
```

```
[Out] 1/8*(3*c*x^3 + 5*a*x)/(a^2*c^2*x^4 + 2*a^3*c*x^2 + a^4) + 3/8*arctan(c*x/sq
rt(a*c))/(sqrt(a*c)*a^2)
```

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.73

$$\int \frac{1}{(a + cx^2)^3} dx = \frac{3 \arctan\left(\frac{cx}{\sqrt{ac}}\right)}{8 \sqrt{aca^2}} + \frac{3cx^3 + 5ax}{8(cx^2 + a)^2 a^2}$$

[In] integrate(1/(c\*x^2+a)^3,x, algorithm="giac")

[Out] 3/8\*arctan(c\*x/sqrt(a\*c))/(sqrt(a\*c)\*a^2) + 1/8\*(3\*c\*x^3 + 5\*a\*x)/((c\*x^2 + a)^2\*a^2)

**Mupad [B] (verification not implemented)**

Time = 9.12 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.89

$$\int \frac{1}{(a + cx^2)^3} dx = \frac{\frac{5x}{8a} + \frac{3cx^3}{8a^2}}{a^2 + 2acx^2 + c^2x^4} + \frac{3 \operatorname{atan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)}{8a^{5/2}\sqrt{c}}$$

[In] int(1/(a + c\*x^2)^3,x)

[Out] ((5\*x)/(8\*a) + (3\*c\*x^3)/(8\*a^2))/(a^2 + c^2\*x^4 + 2\*a\*c\*x^2) + (3\*atan((c^(1/2)\*x)/a^(1/2)))/(8\*a^(5/2)\*c^(1/2))

### 3.56 $\int (a + cx^2)^{5/2} dx$

Optimal result	326
Rubi [A] (verified)	326
Mathematica [A] (verified)	327
Maple [A] (verified)	328
Fricas [A] (verification not implemented)	328
Sympy [A] (verification not implemented)	329
Maxima [A] (verification not implemented)	329
Giac [A] (verification not implemented)	329
Mupad [B] (verification not implemented)	330

#### Optimal result

Integrand size = 11, antiderivative size = 84

$$\int (a + cx^2)^{5/2} dx = \frac{5}{16}a^2x\sqrt{a + cx^2} + \frac{5}{24}ax(a + cx^2)^{3/2} + \frac{1}{6}x(a + cx^2)^{5/2} + \frac{5a^3\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{16\sqrt{c}}$$

[Out]  $5/24*a*x*(c*x^2+a)^{(3/2)}+1/6*x*(c*x^2+a)^{(5/2)}+5/16*a^3*\operatorname{arctanh}(x*c^{(1/2)}/(c*x^2+a)^{(1/2)})/c^{(1/2)}+5/16*a^2*x*(c*x^2+a)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {201, 223, 212}

$$\int (a + cx^2)^{5/2} dx = \frac{5a^3\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{16\sqrt{c}} + \frac{5}{16}a^2x\sqrt{a + cx^2} + \frac{5}{24}ax(a + cx^2)^{3/2} + \frac{1}{6}x(a + cx^2)^{5/2}$$

[In]  $\operatorname{Int}[(a + c*x^2)^{(5/2)}, x]$

[Out]  $(5*a^2*x*\operatorname{Sqrt}[a + c*x^2])/16 + (5*a*x*(a + c*x^2)^{(3/2)})/24 + (x*(a + c*x^2)^{(5/2)})/6 + (5*a^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[a + c*x^2]])/(16*\operatorname{Sqrt}[c])$

#### Rule 201

$\operatorname{Int}[(a + b*x^n)^p, x] := \operatorname{Simp}[x*(a + b*x^n)^p/(n*p + 1), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{p-1}, x], x] /;$  Free

$Q[\{a, b\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{GtQ}[p, 0] \&\& (\text{IntegerQ}[2*p] \mid\mid (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[4*p]) \mid\mid (\text{EqQ}[n, 2] \&\& \text{IntegerQ}[3*p]) \mid\mid \text{LtQ}[\text{Denominator}[p + 1/n], \text{Denominator}[p]])$

### Rule 212

$\text{Int}[(a_ + (b_ \cdot x_ )^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[-b, 2])) \cdot \text{ArcTanh}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[a, 2])], x] /;$   $\text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

### Rule 223

$\text{Int}[1/\text{Sqrt}[(a_ + (b_ \cdot x_ )^2)], x\_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(1 - b \cdot x^2)], x], x, x/\text{Sqrt}[a + b \cdot x^2] /;$   $\text{FreeQ}[\{a, b\}, x] \&\& !\text{GtQ}[a, 0]$

### Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{6}x(a + cx^2)^{5/2} + \frac{1}{6}(5a) \int (a + cx^2)^{3/2} dx \\
 &= \frac{5}{24}ax(a + cx^2)^{3/2} + \frac{1}{6}x(a + cx^2)^{5/2} + \frac{1}{8}(5a^2) \int \sqrt{a + cx^2} dx \\
 &= \frac{5}{16}a^2x\sqrt{a + cx^2} + \frac{5}{24}ax(a + cx^2)^{3/2} + \frac{1}{6}x(a + cx^2)^{5/2} + \frac{1}{16}(5a^3) \int \frac{1}{\sqrt{a + cx^2}} dx \\
 &= \frac{5}{16}a^2x\sqrt{a + cx^2} + \frac{5}{24}ax(a + cx^2)^{3/2} \\
 &\quad + \frac{1}{6}x(a + cx^2)^{5/2} + \frac{1}{16}(5a^3) \text{Subst}\left(\int \frac{1}{1 - cx^2} dx, x, \frac{x}{\sqrt{a + cx^2}}\right) \\
 &= \frac{5}{16}a^2x\sqrt{a + cx^2} + \frac{5}{24}ax(a + cx^2)^{3/2} + \frac{1}{6}x(a + cx^2)^{5/2} + \frac{5a^3 \tanh^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a + cx^2}}\right)}{16\sqrt{c}}
 \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.85

$$\int (a + cx^2)^{5/2} dx = \frac{1}{48}\sqrt{a + cx^2}(33a^2x + 26acx^3 + 8c^2x^5) - \frac{5a^3 \log(-\sqrt{cx} + \sqrt{a + cx^2})}{16\sqrt{c}}$$

[In] Integrate[(a + c\*x^2)^(5/2),x]

[Out] (Sqrt[a + c\*x^2]\*(33\*a^2\*x + 26\*a\*c\*x^3 + 8\*c^2\*x^5))/48 - (5\*a^3\*Log[-(Sqrt[c]\*x) + Sqrt[a + c\*x^2]])/(16\*Sqrt[c])

**Maple [A] (verified)**

Time = 2.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.70

method	result	size
risch	$\frac{x(8x^4c^2+26acx^2+33a^2)\sqrt{cx^2+a}}{48} + \frac{5a^3 \ln(\sqrt{cx^2+a})}{16\sqrt{c}}$	59
pseudoelliptic	$\frac{5 \operatorname{arctanh}\left(\frac{\sqrt{cx^2+a}}{x\sqrt{c}}\right)a^3}{16\sqrt{c}} + \frac{11\left(\frac{8c^{\frac{5}{2}}x^4}{33} + \frac{26ac^{\frac{3}{2}}x^2}{33} + a^2\sqrt{c}\right)x\sqrt{cx^2+a}}{16\sqrt{c}}$	67
default	$\frac{x(cx^2+a)^{\frac{5}{2}}}{6} + \frac{5a\left(\frac{x(cx^2+a)^{\frac{3}{2}}}{4} + \frac{3a\left(\frac{x\sqrt{cx^2+a}}{2} + \frac{a \ln(\sqrt{cx^2+a})}{2\sqrt{c}}\right)}{4}\right)}{6}$	68

[In] int((c\*x^2+a)^(5/2),x,method=\_RETURNVERBOSE)

[Out] 1/48\*x\*(8\*c^2\*x^4+26\*a\*c\*x^2+33\*a^2)\*(c\*x^2+a)^(1/2)+5/16\*a^3\*ln(c^(1/2)\*x+(c\*x^2+a)^(1/2))/c^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.46 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.74

$$\int (a + cx^2)^{5/2} dx = \left[ \frac{15a^3\sqrt{c} \log(-2cx^2 - 2\sqrt{cx^2+a}\sqrt{cx} - a) + 2(8c^3x^5 + 26ac^2x^3 + 33a^2cx)\sqrt{cx^2+a}}{96c}, \right. \\ \left. \frac{15a^3\sqrt{-c} \arctan\left(\frac{\sqrt{-cx}}{\sqrt{cx^2+a}}\right) - (8c^3x^5 + 26ac^2x^3 + 33a^2cx)\sqrt{cx^2+a}}{48c} \right]$$

[In] integrate((c\*x^2+a)^(5/2),x, algorithm="fricas")

[Out] [1/96\*(15\*a^3\*sqrt(c)\*log(-2\*c\*x^2 - 2\*sqrt(c\*x^2 + a)\*sqrt(c)\*x - a) + 2\*(8\*c^3\*x^5 + 26\*a\*c^2\*x^3 + 33\*a^2\*c\*x)\*sqrt(c\*x^2 + a))/c, -1/48\*(15\*a^3\*sqrt(-c)\*arctan(sqrt(-c)\*x/sqrt(c\*x^2 + a)) - (8\*c^3\*x^5 + 26\*a\*c^2\*x^3 + 33\*a^2\*c\*x)\*sqrt(c\*x^2 + a))/c]



**Sympy [A] (verification not implemented)**

Time = 2.65 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.15

$$\int (a + cx^2)^{5/2} dx = \frac{11a^{5/2}x\sqrt{1 + \frac{cx^2}{a}}}{16} + \frac{13a^{3/2}cx^3\sqrt{1 + \frac{cx^2}{a}}}{24} + \frac{\sqrt{ac^2}x^5\sqrt{1 + \frac{cx^2}{a}}}{6} + \frac{5a^3 \operatorname{asinh}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{16\sqrt{c}}$$

[In] integrate((c\*x\*\*2+a)\*\*(5/2),x)

[Out] 11\*a\*\*(5/2)\*x\*sqrt(1 + c\*x\*\*2/a)/16 + 13\*a\*\*(3/2)\*c\*x\*\*3\*sqrt(1 + c\*x\*\*2/a)/24 + sqrt(a)\*c\*\*2\*x\*\*5\*sqrt(1 + c\*x\*\*2/a)/6 + 5\*a\*\*3\*asinh(sqrt(c)\*x/sqrt(a))/(16\*sqrt(c))

**Maxima [A] (verification not implemented)**

none

Time = 0.22 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.69

$$\int (a + cx^2)^{5/2} dx = \frac{1}{6} (cx^2 + a)^{5/2} x + \frac{5}{24} (cx^2 + a)^{3/2} ax + \frac{5}{16} \sqrt{cx^2 + aa^2} x + \frac{5 a^3 \operatorname{arsinh}\left(\frac{cx}{\sqrt{ac}}\right)}{16 \sqrt{c}}$$

[In] integrate((c\*x^2+a)^(5/2),x, algorithm="maxima")

[Out] 1/6\*(c\*x^2 + a)^(5/2)\*x + 5/24\*(c\*x^2 + a)^(3/2)\*a\*x + 5/16\*sqrt(c\*x^2 + a)\*a^2\*x + 5/16\*a^3\*arcsinh(c\*x/sqrt(a\*c))/sqrt(c)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.75

$$\int (a + cx^2)^{5/2} dx = -\frac{5 a^3 \log\left(|-\sqrt{cx} + \sqrt{cx^2 + a}|\right)}{16 \sqrt{c}} + \frac{1}{48} (2(4c^2x^2 + 13ac)x^2 + 33a^2)\sqrt{cx^2 + ax}$$

[In] integrate((c\*x^2+a)^(5/2),x, algorithm="giac")

[Out] -5/16\*a^3\*log(abs(-sqrt(c)\*x + sqrt(c\*x^2 + a)))/sqrt(c) + 1/48\*(2\*(4\*c^2\*x^2 + 13\*a\*c)\*x^2 + 33\*a^2)\*sqrt(c\*x^2 + a)\*x

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.44

$$\int (a + cx^2)^{5/2} dx = \frac{x (cx^2 + a)^{5/2} {}_2F_1\left(-\frac{5}{2}, \frac{1}{2}; \frac{3}{2}; -\frac{cx^2}{a}\right)}{\left(\frac{cx^2}{a} + 1\right)^{5/2}}$$

[In] int((a + c\*x^2)^(5/2),x)

[Out] (x\*(a + c\*x^2)^(5/2)\*hypergeom([-5/2, 1/2], 3/2, -(c\*x^2)/a))/((c\*x^2)/a + 1)^(5/2)

### 3.57 $\int (a + cx^2)^{3/2} dx$

Optimal result . . . . .	331
Rubi [A] (verified) . . . . .	331
Mathematica [A] (verified) . . . . .	332
Maple [A] (verified) . . . . .	332
Fricas [A] (verification not implemented) . . . . .	333
Sympy [A] (verification not implemented) . . . . .	333
Maxima [A] (verification not implemented) . . . . .	334
Giac [A] (verification not implemented) . . . . .	334
Mupad [B] (verification not implemented) . . . . .	334

#### Optimal result

Integrand size = 11, antiderivative size = 65

$$\int (a + cx^2)^{3/2} dx = \frac{3}{8}ax\sqrt{a + cx^2} + \frac{1}{4}x(a + cx^2)^{3/2} + \frac{3a^2\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{8\sqrt{c}}$$

[Out]  $1/4*x*(c*x^2+a)^{(3/2)}+3/8*a^2*\operatorname{arctanh}(x*c^{(1/2)}/(c*x^2+a)^{(1/2)})/c^{(1/2)}+3/8*a*x*(c*x^2+a)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {201, 223, 212}

$$\int (a + cx^2)^{3/2} dx = \frac{3a^2\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{8\sqrt{c}} + \frac{3}{8}ax\sqrt{a + cx^2} + \frac{1}{4}x(a + cx^2)^{3/2}$$

[In]  $\operatorname{Int}[(a + c*x^2)^{(3/2)}, x]$

[Out]  $(3*a*x*\operatorname{Sqrt}[a + c*x^2])/8 + (x*(a + c*x^2)^{(3/2)})/4 + (3*a^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[a + c*x^2]])/(8*\operatorname{Sqrt}[c])$

#### Rule 201

```
Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{4}x(a+cx^2)^{3/2} + \frac{1}{4}(3a) \int \sqrt{a+cx^2} dx \\
&= \frac{3}{8}ax\sqrt{a+cx^2} + \frac{1}{4}x(a+cx^2)^{3/2} + \frac{1}{8}(3a^2) \int \frac{1}{\sqrt{a+cx^2}} dx \\
&= \frac{3}{8}ax\sqrt{a+cx^2} + \frac{1}{4}x(a+cx^2)^{3/2} + \frac{1}{8}(3a^2) \text{Subst}\left(\int \frac{1}{1-cx^2} dx, x, \frac{x}{\sqrt{a+cx^2}}\right) \\
&= \frac{3}{8}ax\sqrt{a+cx^2} + \frac{1}{4}x(a+cx^2)^{3/2} + \frac{3a^2 \tanh^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{8\sqrt{c}}
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.92

$$\int (a+cx^2)^{3/2} dx = \frac{1}{8}x\sqrt{a+cx^2}(5a+2cx^2) - \frac{3a^2 \log(-\sqrt{cx} + \sqrt{a+cx^2})}{8\sqrt{c}}$$

```
[In] Integrate[(a + c*x^2)^(3/2), x]
```

```
[Out] (x*Sqrt[a + c*x^2]*(5*a + 2*c*x^2))/8 - (3*a^2*Log[-(Sqrt[c]*x) + Sqrt[a +
c*x^2]])/(8*Sqrt[c])
```

**Maple [A] (verified)**

Time = 2.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.74

method	result	size
risch	$\frac{x(2cx^2+5a)\sqrt{cx^2+a}}{8} + \frac{3a^2 \ln(\sqrt{c}x + \sqrt{cx^2+a})}{8\sqrt{c}}$	48
default	$\frac{x(cx^2+a)^{\frac{3}{2}}}{4} + \frac{3a \left( \frac{x\sqrt{cx^2+a}}{2} + \frac{a \ln(\sqrt{c}x + \sqrt{cx^2+a})}{2\sqrt{c}} \right)}{4}$	52
pseudoelliptic	$\frac{2\sqrt{cx^2+a}c^{\frac{3}{2}}x^3 + 5ax\sqrt{cx^2+a}\sqrt{c} + 3 \operatorname{arctanh}\left(\frac{\sqrt{cx^2+a}}{x\sqrt{c}}\right)a^2}{8\sqrt{c}}$	62

[In] `int((c*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out]  $\frac{1}{8}x(2cx^2+5a)(cx^2+a)^{1/2} + \frac{3}{8}a^2 \ln(c^{1/2}x + (cx^2+a)^{1/2})/c^{1/2}$

### Fricas [A] (verification not implemented)

none

Time = 0.76 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.91

$$\int (a + cx^2)^{3/2} dx = \left[ \frac{3a^2\sqrt{c} \log(-2cx^2 - 2\sqrt{cx^2+a}\sqrt{cx} - a) + 2(2c^2x^3 + 5acx)\sqrt{cx^2+a}}{16c}, \right. \\ \left. - \frac{3a^2\sqrt{-c} \arctan\left(\frac{\sqrt{-cx}}{\sqrt{cx^2+a}}\right) - (2c^2x^3 + 5acx)\sqrt{cx^2+a}}{8c} \right]$$

[In] `integrate((c*x^2+a)^(3/2),x, algorithm="fricas")`

[Out]  $\frac{1}{16}(3a^2\sqrt{c}\log(-2cx^2 - 2\sqrt{cx^2+a}\sqrt{cx} - a) + 2(2c^2x^3 + 5acx)\sqrt{cx^2+a})/c, -\frac{1}{8}(3a^2\sqrt{-c}\arctan(\sqrt{-cx}/\sqrt{cx^2+a}) - (2c^2x^3 + 5acx)\sqrt{cx^2+a})/c$

### Sympy [A] (verification not implemented)

Time = 1.58 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.08

$$\int (a + cx^2)^{3/2} dx = \frac{5a^{\frac{3}{2}}x\sqrt{1 + \frac{cx^2}{a}}}{8} + \frac{\sqrt{ac}x^3\sqrt{1 + \frac{cx^2}{a}}}{4} + \frac{3a^2 \operatorname{asinh}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{8\sqrt{c}}$$

[In] `integrate((c*x**2+a)**(3/2),x)`

[Out]  $5a^{3/2}x\sqrt{1 + cx^2/a}/8 + \sqrt{a}cx^3\sqrt{1 + cx^2/a}/4 + 3a^{3/2}\operatorname{asinh}(\sqrt{c}x/\sqrt{a})/(8\sqrt{c})$

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.66

$$\int (a + cx^2)^{3/2} dx = \frac{1}{4} (cx^2 + a)^{\frac{3}{2}} x + \frac{3}{8} \sqrt{cx^2 + a} ax + \frac{3a^2 \operatorname{arsinh}\left(\frac{cx}{\sqrt{ac}}\right)}{8\sqrt{c}}$$

[In] integrate((c\*x^2+a)^(3/2),x, algorithm="maxima")

[Out] 1/4\*(c\*x^2 + a)^(3/2)\*x + 3/8\*sqrt(c\*x^2 + a)\*a\*x + 3/8\*a^2\*arcsinh(c\*x/sqrt(a\*c))/sqrt(c)

**Giac [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.75

$$\int (a + cx^2)^{3/2} dx = \frac{1}{8} (2cx^2 + 5a)\sqrt{cx^2 + a} - \frac{3a^2 \log(|-\sqrt{cx} + \sqrt{cx^2 + a}|)}{8\sqrt{c}}$$

[In] integrate((c\*x^2+a)^(3/2),x, algorithm="giac")

[Out] 1/8\*(2\*c\*x^2 + 5\*a)\*sqrt(c\*x^2 + a)\*x - 3/8\*a^2\*log(abs(-sqrt(c)\*x + sqrt(c\*x^2 + a)))/sqrt(c)

**Mupad [B] (verification not implemented)**

Time = 8.97 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.57

$$\int (a + cx^2)^{3/2} dx = \frac{x (cx^2 + a)^{3/2} {}_2F_1\left(-\frac{3}{2}, \frac{1}{2}; \frac{3}{2}; -\frac{cx^2}{a}\right)}{\left(\frac{cx^2}{a} + 1\right)^{3/2}}$$

[In] int((a + c\*x^2)^(3/2),x)

[Out] (x\*(a + c\*x^2)^(3/2)\*hypergeom([-3/2, 1/2], 3/2, -(c\*x^2)/a))/((c\*x^2)/a + 1)^(3/2)

### 3.58 $\int \sqrt{a + cx^2} dx$

Optimal result	335
Rubi [A] (verified)	335
Mathematica [A] (verified)	336
Maple [A] (verified)	336
Fricas [A] (verification not implemented)	337
Sympy [A] (verification not implemented)	337
Maxima [A] (verification not implemented)	337
Giac [A] (verification not implemented)	338
Mupad [B] (verification not implemented)	338

#### Optimal result

Integrand size = 11, antiderivative size = 46

$$\int \sqrt{a + cx^2} dx = \frac{1}{2}x\sqrt{a + cx^2} + \frac{a \operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a + cx^2}}\right)}{2\sqrt{c}}$$

[Out]  $1/2*a*\operatorname{arctanh}(x*c^{(1/2)}/(c*x^2+a)^{(1/2)})/c^{(1/2)}+1/2*x*(c*x^2+a)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$ , Rules used = {201, 223, 212}

$$\int \sqrt{a + cx^2} dx = \frac{a \operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a + cx^2}}\right)}{2\sqrt{c}} + \frac{1}{2}x\sqrt{a + cx^2}$$

[In] `Int[Sqrt[a + c*x^2], x]`

[Out] `(x*Sqrt[a + c*x^2])/2 + (a*ArcTanh[(Sqrt[c]*x)/Sqrt[a + c*x^2]])/(2*Sqrt[c])`

#### Rule 201

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[x*((a + b*x^n)^p/(n*p
+ 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; Free
Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] &&
IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n],
Denominator[p]])
```

#### Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

### Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}x\sqrt{a+cx^2} + \frac{1}{2}a \int \frac{1}{\sqrt{a+cx^2}} dx \\ &= \frac{1}{2}x\sqrt{a+cx^2} + \frac{1}{2}a \text{Subst}\left(\int \frac{1}{1-cx^2} dx, x, \frac{x}{\sqrt{a+cx^2}}\right) \\ &= \frac{1}{2}x\sqrt{a+cx^2} + \frac{a \tanh^{-1}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{2\sqrt{c}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.04

$$\int \sqrt{a+cx^2} dx = \frac{1}{2}x\sqrt{a+cx^2} - \frac{a \log(-\sqrt{cx} + \sqrt{a+cx^2})}{2\sqrt{c}}$$

```
[In] Integrate[Sqrt[a + c*x^2],x]
```

```
[Out] (x*Sqrt[a + c*x^2])/2 - (a*Log[-(Sqrt[c]*x) + Sqrt[a + c*x^2]])/(2*Sqrt[c])
```

### Maple [A] (verified)

Time = 2.20 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{x\sqrt{cx^2+a}}{2} + \frac{a \ln(\sqrt{c}x + \sqrt{cx^2+a})}{2\sqrt{c}}$	36
risch	$\frac{x\sqrt{cx^2+a}}{2} + \frac{a \ln(\sqrt{c}x + \sqrt{cx^2+a})}{2\sqrt{c}}$	36
pseudoelliptic	$\frac{\sqrt{cx^2+a}x\sqrt{c} + \arctanh\left(\frac{\sqrt{cx^2+a}}{x\sqrt{c}}\right)a}{2\sqrt{c}}$	40



[In] `int((c*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $1/2*x*(c*x^2+a)^{(1/2)}+1/2*a/c^{(1/2)}*\ln(c^{(1/2)}*x+(c*x^2+a)^{(1/2)})$

### Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 94, normalized size of antiderivative = 2.04

$$\int \sqrt{a + cx^2} dx = \left[ \frac{2\sqrt{cx^2 + acx} + a\sqrt{c} \log(-2cx^2 - 2\sqrt{cx^2 + a}\sqrt{cx} - a)}{4c}, \frac{\sqrt{cx^2 + acx} - a\sqrt{-c} \arctan\left(\frac{\sqrt{-cx}}{\sqrt{cx^2 + a}}\right)}{2c} \right]$$

[In] `integrate((c*x^2+a)^(1/2),x, algorithm="fricas")`

[Out]  $[1/4*(2*\sqrt{c*x^2 + a}*c*x + a*\sqrt{c}*\log(-2*c*x^2 - 2*\sqrt{c*x^2 + a}*\sqrt{c}*x - a))/c, 1/2*(\sqrt{c*x^2 + a}*c*x - a*\sqrt{-c}*\arctan(\sqrt{-c}*x/\sqrt{c*x^2 + a}))/c]$

### Sympy [A] (verification not implemented)

Time = 0.96 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

$$\int \sqrt{a + cx^2} dx = \frac{\sqrt{ax}\sqrt{1 + \frac{cx^2}{a}}}{2} + \frac{a \operatorname{asinh}\left(\frac{\sqrt{cx}}{\sqrt{a}}\right)}{2\sqrt{c}}$$

[In] `integrate((c*x**2+a)**(1/2),x)`

[Out]  $\sqrt{a}*x*\sqrt{1 + c*x**2/a}/2 + a*\operatorname{asinh}(\sqrt{c}*x/\sqrt{a})/(2*\sqrt{c})$

### Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.61

$$\int \sqrt{a + cx^2} dx = \frac{1}{2} \sqrt{cx^2 + ax} + \frac{a \operatorname{arsinh}\left(\frac{cx}{\sqrt{ac}}\right)}{2\sqrt{c}}$$

[In] `integrate((c*x^2+a)^(1/2),x, algorithm="maxima")`

[Out]  $1/2*\sqrt{c*x^2 + a}*x + 1/2*a*\operatorname{arcsinh}(c*x/\sqrt{a*c})/\sqrt{c}$

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int \sqrt{a + cx^2} dx = \frac{1}{2} \sqrt{cx^2 + ax} - \frac{a \log(|-\sqrt{cx} + \sqrt{cx^2 + a}|)}{2\sqrt{c}}$$

[In] integrate((c\*x^2+a)^(1/2),x, algorithm="giac")

[Out] 1/2\*sqrt(c\*x^2 + a)\*x - 1/2\*a\*log(abs(-sqrt(c)\*x + sqrt(c\*x^2 + a)))/sqrt(c)

**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

$$\int \sqrt{a + cx^2} dx = \frac{x \sqrt{cx^2 + a}}{2} + \frac{a \ln(\sqrt{cx} + \sqrt{cx^2 + a})}{2\sqrt{c}}$$

[In] int((a + c\*x^2)^(1/2),x)

[Out] (x\*(a + c\*x^2)^(1/2))/2 + (a\*log(c^(1/2)\*x + (a + c\*x^2)^(1/2)))/(2\*c^(1/2))

### 3.59 $\int \frac{1}{\sqrt{a+cx^2}} dx$

Optimal result . . . . .	339
Rubi [A] (verified) . . . . .	339
Mathematica [A] (verified) . . . . .	340
Maple [A] (verified) . . . . .	340
Fricas [A] (verification not implemented) . . . . .	340
Sympy [A] (verification not implemented) . . . . .	341
Maxima [A] (verification not implemented) . . . . .	341
Giac [A] (verification not implemented) . . . . .	341
Mupad [B] (verification not implemented) . . . . .	342

#### Optimal result

Integrand size = 11, antiderivative size = 25

$$\int \frac{1}{\sqrt{a+cx^2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{\sqrt{c}}$$

[Out]  $\operatorname{arctanh}(x*c^{(1/2)}/(c*x^2+a)^{(1/2)})/c^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {223, 212}

$$\int \frac{1}{\sqrt{a+cx^2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{cx}}{\sqrt{a+cx^2}}\right)}{\sqrt{c}}$$

[In]  $\operatorname{Int}[1/\operatorname{Sqrt}[a + c*x^2], x]$

[Out]  $\operatorname{ArcTanh}[(\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[a + c*x^2]]/\operatorname{Sqrt}[c]$

#### Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

#### Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+ + (b_+)*(x_+)^2), x\_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left( \int \frac{1}{1 - cx^2} dx, x, \frac{x}{\sqrt{a + cx^2}} \right) \\ &= \frac{\tanh^{-1} \left( \frac{\sqrt{cx}}{\sqrt{a + cx^2}} \right)}{\sqrt{c}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \frac{\text{arctanh} \left( \frac{\sqrt{cx}}{\sqrt{a + cx^2}} \right)}{\sqrt{c}}$$

[In] Integrate[1/Sqrt[a + c\*x^2],x]

[Out] ArcTanh[(Sqrt[c]\*x)/Sqrt[a + c\*x^2]]/Sqrt[c]

### Maple [A] (verified)

Time = 2.37 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

method	result	size
default	$\frac{\ln(\sqrt{c}x + \sqrt{cx^2 + a})}{\sqrt{c}}$	21
pseudoelliptic	$\frac{\text{arctanh} \left( \frac{\sqrt{cx^2 + a}}{x\sqrt{c}} \right)}{\sqrt{c}}$	22

[In] int(1/(c\*x^2+a)^(1/2),x,method=\_RETURNVERBOSE)

[Out] ln(c^(1/2)\*x+(c\*x^2+a)^(1/2))/c^(1/2)

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 2.36

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \left[ \frac{\log(-2cx^2 - 2\sqrt{cx^2 + a}\sqrt{cx} - a)}{2\sqrt{c}}, -\frac{\sqrt{-c} \arctan \left( \frac{\sqrt{-cx}}{\sqrt{cx^2 + a}} \right)}{c} \right]$$

[In] integrate(1/(c\*x^2+a)^(1/2),x, algorithm="fricas")

[Out]  $[1/2*\log(-2*c*x^2 - 2*\sqrt{c*x^2 + a}*\sqrt{c}*x - a)/\sqrt{c}, -\sqrt{-c}*\arctan(\sqrt{-c}*x/\sqrt{c*x^2 + a})/c]$

### Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \frac{\operatorname{asinh}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)}{\sqrt{c}}$$

[In] `integrate(1/(c*x**2+a)**(1/2),x)`

[Out] `asinh(sqrt(c)*x/sqrt(a))/sqrt(c)`

### Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \frac{\operatorname{arsinh}\left(\frac{cx}{\sqrt{ac}}\right)}{\sqrt{c}}$$

[In] `integrate(1/(c*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `arcsinh(c*x/sqrt(a*c))/sqrt(c)`

### Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.48

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \frac{1}{2} \sqrt{cx^2 + a} - \frac{a \log(|-\sqrt{c}x + \sqrt{cx^2 + a}|)}{2\sqrt{c}}$$

[In] `integrate(1/(c*x^2+a)^(1/2),x, algorithm="giac")`

[Out] `1/2*sqrt(c*x^2 + a)*x - 1/2*a*log(abs(-sqrt(c)*x + sqrt(c*x^2 + a)))/sqrt(c)`

**Mupad [B] (verification not implemented)**

Time = 9.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{a + cx^2}} dx = \frac{\ln(\sqrt{c}x + \sqrt{cx^2 + a})}{\sqrt{c}}$$

[In] int(1/(a + c\*x^2)^(1/2),x)

[Out] log(c^(1/2)\*x + (a + c\*x^2)^(1/2))/c^(1/2)

$$3.60 \quad \int \frac{1}{(a+cx^2)^{3/2}} dx$$

Optimal result . . . . .	343
Rubi [A] (verified) . . . . .	343
Mathematica [A] (verified) . . . . .	344
Maple [A] (verified) . . . . .	344
Fricas [A] (verification not implemented) . . . . .	344
Sympy [A] (verification not implemented) . . . . .	345
Maxima [A] (verification not implemented) . . . . .	345
Giac [A] (verification not implemented) . . . . .	345
Mupad [B] (verification not implemented) . . . . .	345

### Optimal result

Integrand size = 11, antiderivative size = 16

$$\int \frac{1}{(a+cx^2)^{3/2}} dx = \frac{x}{a\sqrt{a+cx^2}}$$

[Out] x/a/(c\*x^2+a)^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$ , Rules used = {197}

$$\int \frac{1}{(a+cx^2)^{3/2}} dx = \frac{x}{a\sqrt{a+cx^2}}$$

[In] Int[(a + c\*x^2)^(-3/2), x]

[Out] x/(a\*Sqrt[a + c\*x^2])

#### Rule 197

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] :> Simp[x\*((a + b\*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

#### Rubi steps

$$\text{integral} = \frac{x}{a\sqrt{a+cx^2}}$$

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{x}{a\sqrt{a + cx^2}}$$

[In] Integrate[(a + c\*x^2)^(-3/2),x]

[Out] x/(a\*Sqrt[a + c\*x^2])

**Maple [A] (verified)**

Time = 2.44 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

method	result	size
gospers	$\frac{x}{a\sqrt{cx^2+a}}$	15
default	$\frac{x}{a\sqrt{cx^2+a}}$	15
trager	$\frac{x}{a\sqrt{cx^2+a}}$	15
pseudoelliptic	$\frac{x}{a\sqrt{cx^2+a}}$	15

[In] int(1/(c\*x^2+a)^(3/2),x,method=\_RETURNVERBOSE)

[Out] x/a/(c\*x^2+a)^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.44

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{\sqrt{cx^2 + ax}}{acx^2 + a^2}$$

[In] integrate(1/(c\*x^2+a)^(3/2),x, algorithm="fricas")

[Out] sqrt(c\*x^2 + a)\*x/(a\*c\*x^2 + a^2)



**Sympy [A] (verification not implemented)**

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.06

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{x}{a^{3/2} \sqrt{1 + \frac{cx^2}{a}}}$$

[In] integrate(1/(c\*x\*\*2+a)\*\*(3/2),x)

[Out] x/(a\*\*(3/2)\*sqrt(1 + c\*x\*\*2/a))

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{x}{\sqrt{cx^2 + aa}}$$

[In] integrate(1/(c\*x^2+a)^(3/2),x, algorithm="maxima")

[Out] x/(sqrt(c\*x^2 + a)\*a)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{x}{\sqrt{cx^2 + aa}}$$

[In] integrate(1/(c\*x^2+a)^(3/2),x, algorithm="giac")

[Out] x/(sqrt(c\*x^2 + a)\*a)

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{1}{(a + cx^2)^{3/2}} dx = \frac{x}{a \sqrt{cx^2 + a}}$$

[In] int(1/(a + c\*x^2)^(3/2),x)

[Out] x/(a\*(a + c\*x^2)^(1/2))

### 3.61 $\int \frac{1}{(a+cx^2)^{5/2}} dx$

Optimal result	346
Rubi [A] (verified)	346
Mathematica [A] (verified)	347
Maple [A] (verified)	347
Fricas [A] (verification not implemented)	348
Sympy [B] (verification not implemented)	348
Maxima [A] (verification not implemented)	348
Giac [A] (verification not implemented)	349
Mupad [B] (verification not implemented)	349

#### Optimal result

Integrand size = 11, antiderivative size = 39

$$\int \frac{1}{(a+cx^2)^{5/2}} dx = \frac{x}{3a(a+cx^2)^{3/2}} + \frac{2x}{3a^2\sqrt{a+cx^2}}$$

[Out] 1/3\*x/a/(c\*x^2+a)^(3/2)+2/3\*x/a^2/(c\*x^2+a)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {198, 197}

$$\int \frac{1}{(a+cx^2)^{5/2}} dx = \frac{2x}{3a^2\sqrt{a+cx^2}} + \frac{x}{3a(a+cx^2)^{3/2}}$$

[In] Int[(a + c\*x^2)^(-5/2), x]

[Out] x/(3\*a\*(a + c\*x^2)^(3/2)) + (2\*x)/(3\*a^2\*Sqrt[a + c\*x^2])

#### Rule 197

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[x\*((a + b\*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

#### Rule 198

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[(-x)\*((a + b\*x^n)^(p + 1)/(a\*n\*(p + 1))), x] + Dist[(n\*(p + 1) + 1)/(a\*n\*(p + 1)), Int[(a + b\*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1],

0] && NeQ[p, -1]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{3a(a+cx^2)^{3/2}} + \frac{2 \int \frac{1}{(a+cx^2)^{3/2}} dx}{3a} \\ &= \frac{x}{3a(a+cx^2)^{3/2}} + \frac{2x}{3a^2\sqrt{a+cx^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.74

$$\int \frac{1}{(a+cx^2)^{5/2}} dx = \frac{3ax + 2cx^3}{3a^2(a+cx^2)^{3/2}}$$

[In] Integrate[(a + c\*x^2)^(-5/2), x]

[Out] (3\*a\*x + 2\*c\*x^3)/(3\*a^2\*(a + c\*x^2)^(3/2))

**Maple [A] (verified)**

Time = 2.10 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.67

method	result	size
gospers	$\frac{x(2cx^2+3a)}{3(c x^2+a)^{\frac{3}{2}} a^2}$	26
trager	$\frac{x(2cx^2+3a)}{3(c x^2+a)^{\frac{3}{2}} a^2}$	26
pseudoelliptic	$\frac{x(2cx^2+3a)}{3(c x^2+a)^{\frac{3}{2}} a^2}$	26
default	$\frac{x}{3a(c x^2+a)^{\frac{3}{2}}} + \frac{2x}{3a^2\sqrt{c x^2+a}}$	32

[In] int(1/(c\*x^2+a)^(5/2), x, method=\_RETURNVERBOSE)

[Out] 1/3\*x\*(2\*c\*x^2+3\*a)/(c\*x^2+a)^(3/2)/a^2

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.21

$$\int \frac{1}{(a + cx^2)^{5/2}} dx = \frac{(2cx^3 + 3ax)\sqrt{cx^2 + a}}{3(a^2c^2x^4 + 2a^3cx^2 + a^4)}$$

[In] integrate(1/(c\*x^2+a)^(5/2),x, algorithm="fricas")

[Out] 1/3\*(2\*c\*x^3 + 3\*a\*x)\*sqrt(c\*x^2 + a)/(a^2\*c^2\*x^4 + 2\*a^3\*c\*x^2 + a^4)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(32) = 64.

Time = 0.49 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.44

$$\int \frac{1}{(a + cx^2)^{5/2}} dx = \frac{3ax}{3a^{7/2}\sqrt{1 + \frac{cx^2}{a}} + 3a^{5/2}cx^2\sqrt{1 + \frac{cx^2}{a}}} + \frac{2cx^3}{3a^{7/2}\sqrt{1 + \frac{cx^2}{a}} + 3a^{5/2}cx^2\sqrt{1 + \frac{cx^2}{a}}}$$

[In] integrate(1/(c\*x\*\*2+a)\*\*(5/2),x)

[Out] 3\*a\*x/(3\*a\*\*(7/2)\*sqrt(1 + c\*x\*\*2/a) + 3\*a\*\*(5/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a)) + 2\*c\*x\*\*3/(3\*a\*\*(7/2)\*sqrt(1 + c\*x\*\*2/a) + 3\*a\*\*(5/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a))

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + cx^2)^{5/2}} dx = \frac{2x}{3\sqrt{cx^2 + aa^2}} + \frac{x}{3(cx^2 + a)^{3/2}a}$$

[In] integrate(1/(c\*x^2+a)^(5/2),x, algorithm="maxima")

[Out] 2/3\*x/(sqrt(c\*x^2 + a)\*a^2) + 1/3\*x/((c\*x^2 + a)^(3/2)\*a)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a + cx^2)^{5/2}} dx = \frac{x \left( \frac{2cx^2}{a^2} + \frac{3}{a} \right)}{3 (cx^2 + a)^{3/2}}$$

[In] integrate(1/(c\*x^2+a)^(5/2),x, algorithm="giac")

[Out] 1/3\*x\*(2\*c\*x^2/a^2 + 3/a)/(c\*x^2 + a)^(3/2)

**Mupad [B] (verification not implemented)**

Time = 8.98 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.72

$$\int \frac{1}{(a + cx^2)^{5/2}} dx = \frac{2x(cx^2 + a) + ax}{3a^2(cx^2 + a)^{3/2}}$$

[In] int(1/(a + c\*x^2)^(5/2),x)

[Out] (2\*x\*(a + c\*x^2) + a\*x)/(3\*a^2\*(a + c\*x^2)^(3/2))

## 3.62 $\int \frac{1}{(a+cx^2)^{7/2}} dx$

Optimal result	350
Rubi [A] (verified)	350
Mathematica [A] (verified)	351
Maple [A] (verified)	351
Fricas [A] (verification not implemented)	352
Sympy [B] (verification not implemented)	352
Maxima [A] (verification not implemented)	353
Giac [A] (verification not implemented)	353
Mupad [B] (verification not implemented)	353

### Optimal result

Integrand size = 11, antiderivative size = 58

$$\int \frac{1}{(a+cx^2)^{7/2}} dx = \frac{x}{5a(a+cx^2)^{5/2}} + \frac{4x}{15a^2(a+cx^2)^{3/2}} + \frac{8x}{15a^3\sqrt{a+cx^2}}$$

[Out] 1/5\*x/a/(c\*x^2+a)^(5/2)+4/15\*x/a^2/(c\*x^2+a)^(3/2)+8/15\*x/a^3/(c\*x^2+a)^(1/2)

### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {198, 197}

$$\int \frac{1}{(a+cx^2)^{7/2}} dx = \frac{8x}{15a^3\sqrt{a+cx^2}} + \frac{4x}{15a^2(a+cx^2)^{3/2}} + \frac{x}{5a(a+cx^2)^{5/2}}$$

[In] Int[(a + c\*x^2)^(-7/2),x]

[Out] x/(5\*a\*(a + c\*x^2)^(5/2)) + (4\*x)/(15\*a^2\*(a + c\*x^2)^(3/2)) + (8\*x)/(15\*a^3\*Sqrt[a + c\*x^2])

#### Rule 197

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[x\*((a + b\*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

#### Rule 198

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[(-x)\*((a + b\*x^n)^(p + 1)/(a\*n\*(p + 1))), x] + Dist[(n\*(p + 1) + 1)/(a\*n\*(p + 1)), Int[(a + b\*x^n

$)^{(p + 1), x], x] /; \text{FreeQ}\{a, b, n, p\}, x\} \&\& \text{ILtQ}[\text{Simplify}[1/n + p + 1], 0] \&\& \text{NeQ}[p, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{5a(a+cx^2)^{5/2}} + \frac{4 \int \frac{1}{(a+cx^2)^{5/2}} dx}{5a} \\ &= \frac{x}{5a(a+cx^2)^{5/2}} + \frac{4x}{15a^2(a+cx^2)^{3/2}} + \frac{8 \int \frac{1}{(a+cx^2)^{3/2}} dx}{15a^2} \\ &= \frac{x}{5a(a+cx^2)^{5/2}} + \frac{4x}{15a^2(a+cx^2)^{3/2}} + \frac{8x}{15a^3\sqrt{a+cx^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int \frac{1}{(a+cx^2)^{7/2}} dx = \frac{15a^2x + 20acx^3 + 8c^2x^5}{15a^3(a+cx^2)^{5/2}}$$

[In] Integrate[(a + c\*x^2)^(-7/2),x]

[Out] (15\*a^2\*x + 20\*a\*c\*x^3 + 8\*c^2\*x^5)/(15\*a^3\*(a + c\*x^2)^(5/2))

**Maple [A] (verified)**

Time = 1.98 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.64

method	result	size
gospers	$\frac{x(8x^4c^2+20acx^2+15a^2)}{15(cx^2+a)^{\frac{5}{2}}a^3}$	37
trager	$\frac{x(8x^4c^2+20acx^2+15a^2)}{15(cx^2+a)^{\frac{5}{2}}a^3}$	37
pseudoelliptic	$\frac{x(8x^4c^2+20acx^2+15a^2)}{15(cx^2+a)^{\frac{5}{2}}a^3}$	37
default	$\frac{x}{5a(cx^2+a)^{\frac{5}{2}}} + \frac{\frac{4x}{15a(cx^2+a)^{\frac{3}{2}}} + \frac{8x}{15a^2\sqrt{cx^2+a}}}{a}$	53

[In] int(1/(c\*x^2+a)^(7/2),x,method=\_RETURNVERBOSE)

[Out] 1/15\*x\*(8\*c^2\*x^4+20\*a\*c\*x^2+15\*a^2)/(c\*x^2+a)^(5/2)/a^3

**Fricas [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.19

$$\int \frac{1}{(a + cx^2)^{7/2}} dx = \frac{(8c^2x^5 + 20acx^3 + 15a^2x)\sqrt{cx^2 + a}}{15(a^3c^3x^6 + 3a^4c^2x^4 + 3a^5cx^2 + a^6)}$$

[In] integrate(1/(c\*x^2+a)^(7/2),x, algorithm="fricas")

[Out] 1/15\*(8\*c^2\*x^5 + 20\*a\*c\*x^3 + 15\*a^2\*x)\*sqrt(c\*x^2 + a)/(a^3\*c^3\*x^6 + 3\*a^4\*c^2\*x^4 + 3\*a^5\*c\*x^2 + a^6)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 413 vs. 2(51) = 102.

Time = 0.80 (sec) , antiderivative size = 413, normalized size of antiderivative = 7.12

$$\int \frac{1}{(a + cx^2)^{7/2}} dx = \frac{15a^5x}{15a^{\frac{17}{2}}\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{15}{2}}cx^2\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{13}{2}}c^2x^4\sqrt{1 + \frac{cx^2}{a}} + 15a^{\frac{11}{2}}c^3x^6\sqrt{1 + \frac{cx^2}{a}}} + \frac{35a^4cx^3}{15a^{\frac{17}{2}}\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{15}{2}}cx^2\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{13}{2}}c^2x^4\sqrt{1 + \frac{cx^2}{a}} + 15a^{\frac{11}{2}}c^3x^6\sqrt{1 + \frac{cx^2}{a}}} + \frac{28a^3c^2x^5}{15a^{\frac{17}{2}}\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{15}{2}}cx^2\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{13}{2}}c^2x^4\sqrt{1 + \frac{cx^2}{a}} + 15a^{\frac{11}{2}}c^3x^6\sqrt{1 + \frac{cx^2}{a}}} + \frac{8a^2c^3x^7}{15a^{\frac{17}{2}}\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{15}{2}}cx^2\sqrt{1 + \frac{cx^2}{a}} + 45a^{\frac{13}{2}}c^2x^4\sqrt{1 + \frac{cx^2}{a}} + 15a^{\frac{11}{2}}c^3x^6\sqrt{1 + \frac{cx^2}{a}}}$$

[In] integrate(1/(c\*x\*\*2+a)\*\*(7/2),x)

[Out] 15\*a\*\*5\*x/(15\*a\*\*(17/2)\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(15/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(13/2)\*c\*\*2\*x\*\*4\*sqrt(1 + c\*x\*\*2/a) + 15\*a\*\*(11/2)\*c\*\*3\*x\*\*6\*sqrt(1 + c\*x\*\*2/a)) + 35\*a\*\*4\*c\*x\*\*3/(15\*a\*\*(17/2)\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(15/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(13/2)\*c\*\*2\*x\*\*4\*sqrt(1 + c\*x\*\*2/a) + 15\*a\*\*(11/2)\*c\*\*3\*x\*\*6\*sqrt(1 + c\*x\*\*2/a)) + 28\*a\*\*3\*c\*\*2\*x\*\*5/(15\*a\*\*(17/2)\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(15/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(13/2)\*c\*\*2\*x\*\*4\*sqrt(1 + c\*x\*\*2/a) + 15\*a\*\*(11/2)\*c\*\*3\*x\*\*6\*sqrt(1 + c\*x\*\*2/a)) + 8\*a\*\*2\*c\*\*3\*x\*\*7/(15\*a\*\*(17/2)\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(15/2)\*c\*x\*\*2\*sqrt(1 + c\*x\*\*2/a) + 45\*a\*\*(13/2)\*c\*\*2\*x\*\*4\*sqrt(1 + c\*x\*\*2/a) + 15\*a\*\*(11/2)\*c\*\*3\*x\*\*6\*sqrt(1 + c\*x\*\*2/a))



**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + cx^2)^{7/2}} dx = \frac{8x}{15\sqrt{cx^2 + a}a^3} + \frac{4x}{15(cx^2 + a)^{3/2}a^2} + \frac{x}{5(cx^2 + a)^{5/2}a}$$

[In] integrate(1/(c\*x^2+a)^(7/2),x, algorithm="maxima")

[Out] 8/15\*x/(sqrt(c\*x^2 + a)\*a^3) + 4/15\*x/((c\*x^2 + a)^(3/2)\*a^2) + 1/5\*x/((c\*x^2 + a)^(5/2)\*a)

**Giac [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a + cx^2)^{7/2}} dx = \frac{\left(4x^2\left(\frac{2c^2x^2}{a^3} + \frac{5c}{a^2}\right) + \frac{15}{a}\right)x}{15(cx^2 + a)^{5/2}}$$

[In] integrate(1/(c\*x^2+a)^(7/2),x, algorithm="giac")

[Out] 1/15\*(4\*x^2\*(2\*c^2\*x^2/a^3 + 5\*c/a^2) + 15/a)\*x/(c\*x^2 + a)^(5/2)

**Mupad [B] (verification not implemented)**

Time = 9.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.76

$$\int \frac{1}{(a + cx^2)^{7/2}} dx = \frac{8x(cx^2 + a)^2 + 3a^2x + 4ax(cx^2 + a)}{15a^3(cx^2 + a)^{5/2}}$$

[In] int(1/(a + c\*x^2)^(7/2),x)

[Out] (8\*x\*(a + c\*x^2)^2 + 3\*a^2\*x + 4\*a\*x\*(a + c\*x^2))/(15\*a^3\*(a + c\*x^2)^(5/2))

### 3.63 $\int \frac{1}{(a+cx^2)^{9/2}} dx$

Optimal result	354
Rubi [A] (verified)	354
Mathematica [A] (verified)	355
Maple [A] (verified)	355
Fricas [A] (verification not implemented)	356
Sympy [B] (verification not implemented)	356
Maxima [A] (verification not implemented)	357
Giac [A] (verification not implemented)	358
Mupad [B] (verification not implemented)	358

#### Optimal result

Integrand size = 11, antiderivative size = 77

$$\int \frac{1}{(a+cx^2)^{9/2}} dx = \frac{x}{7a(a+cx^2)^{7/2}} + \frac{6x}{35a^2(a+cx^2)^{5/2}} + \frac{8x}{35a^3(a+cx^2)^{3/2}} + \frac{16x}{35a^4\sqrt{a+cx^2}}$$

[Out] 1/7\*x/a/(c\*x^2+a)^(7/2)+6/35\*x/a^2/(c\*x^2+a)^(5/2)+8/35\*x/a^3/(c\*x^2+a)^(3/2)+16/35\*x/a^4/(c\*x^2+a)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$ , Rules used = {198, 197}

$$\int \frac{1}{(a+cx^2)^{9/2}} dx = \frac{16x}{35a^4\sqrt{a+cx^2}} + \frac{8x}{35a^3(a+cx^2)^{3/2}} + \frac{6x}{35a^2(a+cx^2)^{5/2}} + \frac{x}{7a(a+cx^2)^{7/2}}$$

[In] Int[(a + c\*x^2)^(-9/2), x]

[Out] x/(7\*a\*(a + c\*x^2)^(7/2)) + (6\*x)/(35\*a^2\*(a + c\*x^2)^(5/2)) + (8\*x)/(35\*a^3\*(a + c\*x^2)^(3/2)) + (16\*x)/(35\*a^4\*sqrt[a + c\*x^2])

#### Rule 197

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[x\*((a + b\*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

#### Rule 198

Int[((a\_) + (b\_.)\*(x\_)^(n\_))^(p\_), x\_Symbol] := Simp[(-x)\*((a + b\*x^n)^(p + 1)/(a\*n\*(p + 1))), x] + Dist[(n\*(p + 1) + 1)/(a\*n\*(p + 1)), Int[(a + b\*x^n

$(p + 1), x], x] /; \text{FreeQ}\{a, b, n, p\}, x \ \&\& \ \text{ILtQ}[\text{Simplify}[1/n + p + 1], 0] \ \&\& \ \text{NeQ}[p, -1]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x}{7a(a+cx^2)^{7/2}} + \frac{6 \int \frac{1}{(a+cx^2)^{7/2}} dx}{7a} \\
 &= \frac{x}{7a(a+cx^2)^{7/2}} + \frac{6x}{35a^2(a+cx^2)^{5/2}} + \frac{24 \int \frac{1}{(a+cx^2)^{5/2}} dx}{35a^2} \\
 &= \frac{x}{7a(a+cx^2)^{7/2}} + \frac{6x}{35a^2(a+cx^2)^{5/2}} + \frac{8x}{35a^3(a+cx^2)^{3/2}} + \frac{16 \int \frac{1}{(a+cx^2)^{3/2}} dx}{35a^3} \\
 &= \frac{x}{7a(a+cx^2)^{7/2}} + \frac{6x}{35a^2(a+cx^2)^{5/2}} + \frac{8x}{35a^3(a+cx^2)^{3/2}} + \frac{16x}{35a^4\sqrt{a+cx^2}}
 \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.66

$$\int \frac{1}{(a+cx^2)^{9/2}} dx = \frac{35a^3x + 70a^2cx^3 + 56ac^2x^5 + 16c^3x^7}{35a^4(a+cx^2)^{7/2}}$$

[In] Integrate[(a + c\*x^2)^(-9/2), x]

[Out] (35\*a^3\*x + 70\*a^2\*c\*x^3 + 56\*a\*c^2\*x^5 + 16\*c^3\*x^7)/(35\*a^4\*(a + c\*x^2)^(7/2))

**Maple [A] (verified)**

Time = 2.12 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.62

method	result	size
gospers	$\frac{x(16c^3x^6+56a^2cx^4+70a^2cx^2+35a^3)}{35(c^2x+a)^{\frac{7}{2}}a^4}$	48
trager	$\frac{x(16c^3x^6+56a^2cx^4+70a^2cx^2+35a^3)}{35(c^2x+a)^{\frac{7}{2}}a^4}$	48
pseudoelliptic	$\frac{x(16c^3x^6+56a^2cx^4+70a^2cx^2+35a^3)}{35(c^2x+a)^{\frac{7}{2}}a^4}$	48
default	$\frac{x}{7a(c^2x+a)^{\frac{7}{2}}} + \frac{\frac{6x}{35a(c^2x+a)^{\frac{5}{2}}} + \frac{6\left(\frac{4x}{15a(c^2x+a)^{\frac{3}{2}}} + \frac{8x}{15a^2\sqrt{c^2x+a}}\right)}{7a}}{a}$	74

[In] `int(1/(c*x^2+a)^(9/2),x,method=_RETURNVERBOSE)`

[Out]  $1/35*x*(16*c^3*x^6+56*a*c^2*x^4+70*a^2*c*x^2+35*a^3)/(c*x^2+a)^(7/2)/a^4$

### Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.18

$$\int \frac{1}{(a+cx^2)^{9/2}} dx = \frac{(16c^3x^7 + 56ac^2x^5 + 70a^2cx^3 + 35a^3x)\sqrt{cx^2+a}}{35(a^4c^4x^8 + 4a^5c^3x^6 + 6a^6c^2x^4 + 4a^7cx^2 + a^8)}$$

[In] `integrate(1/(c*x^2+a)^(9/2),x, algorithm="fricas")`

[Out]  $1/35*(16*c^3*x^7 + 56*a*c^2*x^5 + 70*a^2*c*x^3 + 35*a^3*x)*\sqrt{c*x^2 + a}/(a^4*c^4*x^8 + 4*a^5*c^3*x^6 + 6*a^6*c^2*x^4 + 4*a^7*c*x^2 + a^8)$

### Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1265 vs.  $2(70) = 140$ .

Time = 1.23 (sec) , antiderivative size = 1265, normalized size of antiderivative = 16.43

$$\int \frac{1}{(a+cx^2)^{9/2}} dx = \text{Too large to display}$$

[In] `integrate(1/(c*x**2+a)**(9/2),x)`

[Out]  $35*a^{14}*x/(35*a^{37/2}*\sqrt{1 + c*x^2/a}) + 210*a^{35/2}*c*x^2*\sqrt{1 + c*x^2/a} + 525*a^{33/2}*c^2*x^4*\sqrt{1 + c*x^2/a} + 700*a^{31/2}*c^3*x^6*\sqrt{1 + c*x^2/a} + 525*a^{29/2}*c^4*x^8*\sqrt{1 + c*x^2/a} + 210*a^{27/2}*c^5*x^{10}*\sqrt{1 + c*x^2/a} + 35*a^{25/2}*c^6*x^{12}*\sqrt{1$

```

+ c*x**2/a)) + 175*a**13*c*x**3/(35*a**(37/2)*sqrt(1 + c*x**2/a) + 210*a**
35/2)*c*x**2*sqrt(1 + c*x**2/a) + 525*a**(33/2)*c**2*x**4*sqrt(1 + c*x**2/a
) + 700*a**(31/2)*c**3*x**6*sqrt(1 + c*x**2/a) + 525*a**(29/2)*c**4*x**8*sq
rt(1 + c*x**2/a) + 210*a**(27/2)*c**5*x**10*sqrt(1 + c*x**2/a) + 35*a**(25/
2)*c**6*x**12*sqrt(1 + c*x**2/a)) + 371*a**12*c**2*x**5/(35*a**(37/2)*sqrt(
1 + c*x**2/a) + 210*a**(35/2)*c*x**2*sqrt(1 + c*x**2/a) + 525*a**(33/2)*c**
2*x**4*sqrt(1 + c*x**2/a) + 700*a**(31/2)*c**3*x**6*sqrt(1 + c*x**2/a) + 52
5*a**(29/2)*c**4*x**8*sqrt(1 + c*x**2/a) + 210*a**(27/2)*c**5*x**10*sqrt(1
+ c*x**2/a) + 35*a**(25/2)*c**6*x**12*sqrt(1 + c*x**2/a)) + 429*a**11*c**3*
x**7/(35*a**(37/2)*sqrt(1 + c*x**2/a) + 210*a**(35/2)*c*x**2*sqrt(1 + c*x**
2/a) + 525*a**(33/2)*c**2*x**4*sqrt(1 + c*x**2/a) + 700*a**(31/2)*c**3*x**6
*sqrt(1 + c*x**2/a) + 525*a**(29/2)*c**4*x**8*sqrt(1 + c*x**2/a) + 210*a**(
27/2)*c**5*x**10*sqrt(1 + c*x**2/a) + 35*a**(25/2)*c**6*x**12*sqrt(1 + c*x
**2/a)) + 286*a**10*c**4*x**9/(35*a**(37/2)*sqrt(1 + c*x**2/a) + 210*a**(35/
2)*c*x**2*sqrt(1 + c*x**2/a) + 525*a**(33/2)*c**2*x**4*sqrt(1 + c*x**2/a) +
700*a**(31/2)*c**3*x**6*sqrt(1 + c*x**2/a) + 525*a**(29/2)*c**4*x**8*sqrt(
1 + c*x**2/a) + 210*a**(27/2)*c**5*x**10*sqrt(1 + c*x**2/a) + 35*a**(25/2)*
c**6*x**12*sqrt(1 + c*x**2/a)) + 104*a**9*c**5*x**11/(35*a**(37/2)*sqrt(1 +
c*x**2/a) + 210*a**(35/2)*c*x**2*sqrt(1 + c*x**2/a) + 525*a**(33/2)*c**2*x
**4*sqrt(1 + c*x**2/a) + 700*a**(31/2)*c**3*x**6*sqrt(1 + c*x**2/a) + 525*a
**(29/2)*c**4*x**8*sqrt(1 + c*x**2/a) + 210*a**(27/2)*c**5*x**10*sqrt(1 + c
*x**2/a) + 35*a**(25/2)*c**6*x**12*sqrt(1 + c*x**2/a)) + 16*a**8*c**6*x**13
/(35*a**(37/2)*sqrt(1 + c*x**2/a) + 210*a**(35/2)*c*x**2*sqrt(1 + c*x**2/a)
+ 525*a**(33/2)*c**2*x**4*sqrt(1 + c*x**2/a) + 700*a**(31/2)*c**3*x**6*sq
rt(1 + c*x**2/a) + 525*a**(29/2)*c**4*x**8*sqrt(1 + c*x**2/a) + 210*a**(27/2
)*c**5*x**10*sqrt(1 + c*x**2/a) + 35*a**(25/2)*c**6*x**12*sqrt(1 + c*x**2/a
))

```

## Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + cx^2)^{9/2}} dx = \frac{16x}{35\sqrt{cx^2 + aa^4}} + \frac{8x}{35(cx^2 + a)^{3/2}a^3} + \frac{6x}{35(cx^2 + a)^{5/2}a^2} + \frac{x}{7(cx^2 + a)^{7/2}a}$$

[In] integrate(1/(c\*x^2+a)^(9/2),x, algorithm="maxima")

[Out] 16/35\*x/(sqrt(c\*x^2 + a)\*a^4) + 8/35\*x/((c\*x^2 + a)^(3/2)\*a^3) + 6/35\*x/((c\*x^2 + a)^(5/2)\*a^2) + 1/7\*x/((c\*x^2 + a)^(7/2)\*a)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.71

$$\int \frac{1}{(a + cx^2)^{9/2}} dx = \frac{\left(2 \left(4x^2 \left(\frac{2c^3x^2}{a^4} + \frac{7c^2}{a^3}\right) + \frac{35c}{a^2}\right)x^2 + \frac{35}{a}\right)x}{35 (cx^2 + a)^{7/2}}$$

[In] integrate(1/(c\*x^2+a)^(9/2),x, algorithm="giac")

[Out] 1/35\*(2\*(4\*x^2\*(2\*c^3\*x^2/a^4 + 7\*c^2/a^3) + 35\*c/a^2)\*x^2 + 35/a)\*x/(c\*x^2 + a)^(7/2)

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a + cx^2)^{9/2}} dx = \frac{16x}{35a^4 \sqrt{cx^2 + a}} + \frac{8x}{35a^3 (cx^2 + a)^{3/2}} + \frac{6x}{35a^2 (cx^2 + a)^{5/2}} + \frac{x}{7a (cx^2 + a)^{7/2}}$$

[In] int(1/(a + c\*x^2)^(9/2),x)

[Out] (16\*x)/(35\*a^4\*(a + c\*x^2)^(1/2)) + (8\*x)/(35\*a^3\*(a + c\*x^2)^(3/2)) + (6\*x)/(35\*a^2\*(a + c\*x^2)^(5/2)) + x/(7\*a\*(a + c\*x^2)^(7/2))

### 3.64 $\int (4 + 12x + 9x^2)^{3/2} dx$

Optimal result . . . . .	359
Rubi [A] (verified) . . . . .	359
Mathematica [A] (verified) . . . . .	360
Maple [A] (verified) . . . . .	360
Fricas [A] (verification not implemented) . . . . .	360
Sympy [B] (verification not implemented) . . . . .	361
Maxima [A] (verification not implemented) . . . . .	361
Giac [B] (verification not implemented) . . . . .	361
Mupad [B] (verification not implemented) . . . . .	362

#### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{1}{12}(2 + 3x)(4 + 12x + 9x^2)^{3/2}$$

[Out] 1/12\*(2+3\*x)\*(9\*x^2+12\*x+4)^(3/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {623}

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{1}{12}(3x + 2)(9x^2 + 12x + 4)^{3/2}$$

[In] Int[(4 + 12\*x + 9\*x^2)^(3/2), x]

[Out] ((2 + 3\*x)\*(4 + 12\*x + 9\*x^2)^(3/2))/12

#### Rule 623

Int[((a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] /; FreeQ[{a, b, c, p}, x] && EqQ[b^2 - 4\*a\*c, 0] && NeQ[p, -2^(-1)]

#### Rubi steps

$$\text{integral} = \frac{1}{12}(2 + 3x)(4 + 12x + 9x^2)^{3/2}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{1}{12}(2 + 3x) ((2 + 3x)^2)^{3/2}$$

[In] Integrate[(4 + 12\*x + 9\*x^2)^(3/2),x]

[Out] ((2 + 3\*x)\*((2 + 3\*x)^2)^(3/2))/12

**Maple [A] (verified)**

Time = 2.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{(2+3x)((2+3x)^2)^{\frac{3}{2}}}{12}$	17
risch	$\frac{\sqrt{(2+3x)^2} (2+3x)^3}{12}$	19
gospers	$\frac{x(27x^3+72x^2+72x+32)((2+3x)^2)^{\frac{3}{2}}}{4(2+3x)^3}$	35

[In] int((9\*x^2+12\*x+4)^(3/2),x,method=\_RETURNVERBOSE)

[Out] 1/12\*(2+3\*x)\*((2+3\*x)^2)^(3/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.39 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{27}{4}x^4 + 18x^3 + 18x^2 + 8x$$

[In] integrate((9\*x^2+12\*x+4)^(3/2),x, algorithm="fricas")

[Out] 27/4\*x^4 + 18\*x^3 + 18\*x^2 + 8\*x



**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 80 vs.  $2(19) = 38$ .

Time = 0.46 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.48

$$\int (4 + 12x + 9x^2)^{3/2} dx = 4\left(\frac{x}{2} + \frac{1}{3}\right)\sqrt{9x^2 + 12x + 4} + 12\left(\frac{x^2}{3} + \frac{x}{9} - \frac{2}{27}\right)\sqrt{9x^2 + 12x + 4} + 9\sqrt{9x^2 + 12x + 4}\left(\frac{x^3}{4} + \frac{x^2}{18} - \frac{x}{27} + \frac{2}{81}\right)$$

[In] integrate((9\*x\*\*2+12\*x+4)\*\*(3/2),x)

[Out] 4\*(x/2 + 1/3)\*sqrt(9\*x\*\*2 + 12\*x + 4) + 12\*(x\*\*2/3 + x/9 - 2/27)\*sqrt(9\*x\*\*2 + 12\*x + 4) + 9\*sqrt(9\*x\*\*2 + 12\*x + 4)\*(x\*\*3/4 + x\*\*2/18 - x/27 + 2/81)

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{1}{4} (9x^2 + 12x + 4)^{\frac{3}{2}} x + \frac{1}{6} (9x^2 + 12x + 4)^{\frac{3}{2}}$$

[In] integrate((9\*x^2+12\*x+4)^(3/2),x, algorithm="maxima")

[Out] 1/4\*(9\*x^2 + 12\*x + 4)^(3/2)\*x + 1/6\*(9\*x^2 + 12\*x + 4)^(3/2)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 45 vs.  $2(19) = 38$ .

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.96

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{3}{4} (3x^2 + 4x)^2 \operatorname{sgn}(3x + 2) + 2(3x^2 + 4x) \operatorname{sgn}(3x + 2) + \frac{4}{3} \operatorname{sgn}(3x + 2)$$

[In] integrate((9\*x^2+12\*x+4)^(3/2),x, algorithm="giac")

[Out] 3/4\*(3\*x^2 + 4\*x)^2\*sgn(3\*x + 2) + 2\*(3\*x^2 + 4\*x)\*sgn(3\*x + 2) + 4/3\*sgn(3\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int (4 + 12x + 9x^2)^{3/2} dx = \frac{(9x + 6)(9x^2 + 12x + 4)^{3/2}}{36}$$

[In] int((12\*x + 9\*x^2 + 4)^(3/2),x)

[Out] ((9\*x + 6)\*(12\*x + 9\*x^2 + 4)^(3/2))/36

### 3.65 $\int \sqrt{4 + 12x + 9x^2} dx$

Optimal result . . . . .	363
Rubi [A] (verified) . . . . .	363
Mathematica [A] (verified) . . . . .	364
Maple [C] (warning: unable to verify) . . . . .	364
Fricas [A] (verification not implemented) . . . . .	364
Sympy [A] (verification not implemented) . . . . .	365
Maxima [A] (verification not implemented) . . . . .	365
Giac [A] (verification not implemented) . . . . .	365
Mupad [B] (verification not implemented) . . . . .	365

#### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{4 + 12x + 9x^2}$$

[Out] 1/6\*(2+3\*x)\*((2+3\*x)^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {623}

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{1}{6}(3x + 2)\sqrt{9x^2 + 12x + 4}$$

[In] Int[Sqrt[4 + 12\*x + 9\*x^2],x]

[Out] ((2 + 3\*x)\*Sqrt[4 + 12\*x + 9\*x^2])/6

#### Rule 623

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] /; FreeQ[{a, b, c, p}, x] && EqQ[b^2 - 4\*a\*c, 0] && NeQ[p, -2^(-1)]

#### Rubi steps

$$\text{integral} = \frac{1}{6}(2 + 3x)\sqrt{4 + 12x + 9x^2}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{x\sqrt{(2+3x)^2(4+3x)}}{4+6x}$$

[In] Integrate[Sqrt[4 + 12\*x + 9\*x^2], x]

[Out] (x\*Sqrt[(2 + 3\*x)^2]\*(4 + 3\*x))/(4 + 6\*x)

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 2.

Time = 2.07 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\text{csgn}(2+3x)(2+3x)^2}{6}$	16
gospers	$\frac{x(4+3x)\sqrt{(2+3x)^2}}{4+6x}$	25
risch	$\frac{3\sqrt{(2+3x)^2} x^2}{2(2+3x)} + \frac{2\sqrt{(2+3x)^2} x}{2+3x}$	42

[In] int((9\*x^2+12\*x+4)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/6\*csgn(2+3\*x)\*(2+3\*x)^2

**Fricas [A] (verification not implemented)**

none

Time = 0.45 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.39

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{3}{2} x^2 + 2x$$

[In] integrate((9\*x^2+12\*x+4)^(1/2), x, algorithm="fricas")

[Out] 3/2\*x^2 + 2\*x

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \sqrt{4 + 12x + 9x^2} dx = \left(\frac{x}{2} + \frac{1}{3}\right) \sqrt{9x^2 + 12x + 4}$$

[In] integrate((9\*x\*\*2+12\*x+4)\*\*(1/2),x)

[Out] (x/2 + 1/3)\*sqrt(9\*x\*\*2 + 12\*x + 4)

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{1}{2} \sqrt{9x^2 + 12x + 4}x + \frac{1}{3} \sqrt{9x^2 + 12x + 4}$$

[In] integrate((9\*x^2+12\*x+4)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(9\*x^2 + 12\*x + 4)\*x + 1/3\*sqrt(9\*x^2 + 12\*x + 4)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{1}{2} (3x^2 + 4x) \operatorname{sgn}(3x + 2) + \frac{2}{3} \operatorname{sgn}(3x + 2)$$

[In] integrate((9\*x^2+12\*x+4)^(1/2),x, algorithm="giac")

[Out] 1/2\*(3\*x^2 + 4\*x)\*sgn(3\*x + 2) + 2/3\*sgn(3\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 9.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \sqrt{4 + 12x + 9x^2} dx = \frac{(3x + 2) \sqrt{9x^2 + 12x + 4}}{6}$$

[In] int((12\*x + 9\*x^2 + 4)^(1/2),x)

[Out] ((3\*x + 2)\*(12\*x + 9\*x^2 + 4)^(1/2))/6

### 3.66 $\int \frac{1}{\sqrt{4+12x+9x^2}} dx$

Optimal result	366
Rubi [A] (verified)	366
Mathematica [A] (verified)	367
Maple [A] (verified)	367
Fricas [A] (verification not implemented)	368
Sympy [A] (verification not implemented)	368
Maxima [A] (verification not implemented)	368
Giac [A] (verification not implemented)	368
Mupad [B] (verification not implemented)	369

#### Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \frac{1}{\sqrt{4+12x+9x^2}} dx = \frac{(2+3x)\log(2+3x)}{3\sqrt{4+12x+9x^2}}$$

[Out] 1/3\*(2+3\*x)\*ln(2+3\*x)/((2+3\*x)^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {622, 31}

$$\int \frac{1}{\sqrt{4+12x+9x^2}} dx = \frac{(3x+2)\log(3x+2)}{3\sqrt{9x^2+12x+4}}$$

[In] Int[1/Sqrt[4 + 12\*x + 9\*x^2], x]

[Out] ((2 + 3\*x)\*Log[2 + 3\*x])/(3\*Sqrt[4 + 12\*x + 9\*x^2])

#### Rule 31

Int[((a\_) + (b\_.)\*(x\_))^(n\_), x\_Symbol] := Simp[Log[RemoveContent[a + b\*x, x]]/b, x] /; FreeQ[{a, b}, x]

#### Rule 622

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[(b/2 + c\*x)/Sqrt[a + b\*x + c\*x^2], Int[1/(b/2 + c\*x), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(6 + 9x) \int \frac{1}{6+9x} dx}{\sqrt{4 + 12x + 9x^2}} \\ &= \frac{(2 + 3x) \log(2 + 3x)}{3\sqrt{4 + 12x + 9x^2}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{(2 + 3x) \log(2 + 3x)}{3\sqrt{(2 + 3x)^2}}$$

[In] Integrate[1/Sqrt[4 + 12\*x + 9\*x^2],x]

[Out] ((2 + 3\*x)\*Log[2 + 3\*x])/(3\*Sqrt[(2 + 3\*x)^2])

### Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.31

method	result	size
meijerg	$\frac{\ln(1 + \frac{3x}{2})}{3}$	9
default	$\frac{(2+3x) \ln(2+3x)}{3\sqrt{(2+3x)^2}}$	23
risch	$\frac{\sqrt{(2+3x)^2} \ln(2+3x)}{9x+6}$	25

[In] int(1/(9\*x^2+12\*x+4)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*ln(1+3/2\*x)

**Fricas [A] (verification not implemented)**

none

Time = 0.44 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.28

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{1}{3} \log(3x + 2)$$

[In] integrate(1/(9\*x^2+12\*x+4)^(1/2),x, algorithm="fricas")

[Out] 1/3\*log(3\*x + 2)

**Sympy [A] (verification not implemented)**

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{(x + \frac{2}{3}) \log(x + \frac{2}{3})}{3\sqrt{(x + \frac{2}{3})^2}}$$

[In] integrate(1/(9\*x\*\*2+12\*x+4)\*\*(1/2),x)

[Out] (x + 2/3)\*log(x + 2/3)/(3\*sqrt((x + 2/3)\*\*2))

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{1}{3} \log\left(x + \frac{2}{3}\right)$$

[In] integrate(1/(9\*x^2+12\*x+4)^(1/2),x, algorithm="maxima")

[Out] 1/3\*log(x + 2/3)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{\log(|3x + 2| |\operatorname{sgn}(3x + 2)|)}{3 \operatorname{sgn}(3x + 2)}$$

[In] integrate(1/(9\*x^2+12\*x+4)^(1/2),x, algorithm="giac")

[Out] 1/3\*log(abs(3\*x + 2)\*abs(sgn(3\*x + 2)))/sgn(3\*x + 2)



**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.48

$$\int \frac{1}{\sqrt{4 + 12x + 9x^2}} dx = \frac{\ln(9x + 6) \operatorname{sign}(18x + 12)}{3}$$

[In] int(1/(12\*x + 9\*x^2 + 4)^(1/2),x)

[Out] (log(9\*x + 6)\*sign(18\*x + 12))/3

$$3.67 \quad \int \frac{1}{(4+12x+9x^2)^{3/2}} dx$$

Optimal result	370
Rubi [A] (verified)	370
Mathematica [A] (verified)	371
Maple [A] (verified)	371
Fricas [A] (verification not implemented)	371
Sympy [F]	372
Maxima [A] (verification not implemented)	372
Giac [A] (verification not implemented)	372
Mupad [B] (verification not implemented)	372

### Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \frac{1}{(4+12x+9x^2)^{3/2}} dx = -\frac{1}{6(2+3x)\sqrt{4+12x+9x^2}}$$

[Out] -1/6/(2+3\*x)/((2+3\*x)^2)^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {621}

$$\int \frac{1}{(4+12x+9x^2)^{3/2}} dx = -\frac{1}{6(3x+2)\sqrt{9x^2+12x+4}}$$

[In] Int[(4 + 12\*x + 9\*x^2)^(-3/2), x]

[Out] -1/6\*1/((2 + 3\*x)\*Sqrt[4 + 12\*x + 9\*x^2])

#### Rule 621

Int[((a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[2\*((a + b\*x + c\*x^2)^(p + 1)/((2\*p + 1)\*(b + 2\*c\*x))), x] /; FreeQ[{a, b, c, p}, x] && EqQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1]

#### Rubi steps

$$\text{integral} = -\frac{1}{6(2+3x)\sqrt{4+12x+9x^2}}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = -\frac{2 + 3x}{6((2 + 3x)^2)^{3/2}}$$

[In] Integrate[(4 + 12\*x + 9\*x^2)^(-3/2),x]

[Out] -1/6\*(2 + 3\*x)/((2 + 3\*x)^2)^(3/2)

**Maple [A] (verified)**

Time = 2.15 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.64

method	result	size
meijerg	$\frac{x(\frac{3x}{2}+2)}{16(1+\frac{3x}{2})^2}$	16
gosper	$-\frac{2+3x}{6((2+3x)^2)^{\frac{3}{2}}}$	17
default	$-\frac{2+3x}{6((2+3x)^2)^{\frac{3}{2}}}$	17
risch	$-\frac{\sqrt{(2+3x)^2}}{6(2+3x)^3}$	19

[In] int(1/(9\*x^2+12\*x+4)^(3/2),x,method=\_RETURNVERBOSE)

[Out] 1/16\*x\*(3/2\*x+2)/(1+3/2\*x)^2

**Fricas [A] (verification not implemented)**

none

Time = 0.67 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.56

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = -\frac{1}{6(9x^2 + 12x + 4)}$$

[In] integrate(1/(9\*x^2+12\*x+4)^(3/2),x, algorithm="fricas")

[Out] -1/6/(9\*x^2 + 12\*x + 4)

**Sympy [F]**

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = \int \frac{1}{(9x^2 + 12x + 4)^{\frac{3}{2}}} dx$$

[In] integrate(1/(9\*x\*\*2+12\*x+4)\*\*(3/2),x)

[Out] Integral((9\*x\*\*2 + 12\*x + 4)\*\*(-3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.36

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = -\frac{1}{6(3x + 2)^2}$$

[In] integrate(1/(9\*x^2+12\*x+4)^(3/2),x, algorithm="maxima")

[Out] -1/6/(3\*x + 2)^2

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = -\frac{1}{6(3x + 2)^2 \operatorname{sgn}(3x + 2)}$$

[In] integrate(1/(9\*x^2+12\*x+4)^(3/2),x, algorithm="giac")

[Out] -1/6/((3\*x + 2)^2\*sgn(3\*x + 2))

**Mupad [B] (verification not implemented)**

Time = 9.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{1}{(4 + 12x + 9x^2)^{3/2}} dx = -\frac{\sqrt{9x^2 + 12x + 4}}{6(3x + 2)^3}$$

[In] int(1/(12\*x + 9\*x^2 + 4)^(3/2),x)

[Out] -(12\*x + 9\*x^2 + 4)^(1/2)/(6\*(3\*x + 2)^3)

### 3.68 $\int \sqrt{4 - 12x + 9x^2} dx$

Optimal result	373
Rubi [A] (verified)	373
Mathematica [A] (verified)	374
Maple [C] (warning: unable to verify)	374
Fricas [A] (verification not implemented)	374
Sympy [A] (verification not implemented)	375
Maxima [A] (verification not implemented)	375
Giac [A] (verification not implemented)	375
Mupad [B] (verification not implemented)	375

#### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \sqrt{4 - 12x + 9x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{4 - 12x + 9x^2}$$

[Out]  $-1/6*(2-3*x)*((-2+3*x)^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {623}

$$\int \sqrt{4 - 12x + 9x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{9x^2 - 12x + 4}$$

[In]  $\text{Int}[\text{Sqrt}[4 - 12*x + 9*x^2], x]$

[Out]  $-1/6*((2 - 3*x)*\text{Sqrt}[4 - 12*x + 9*x^2])$

#### Rule 623

$\text{Int}[(a_ + (b_)*(x_ ) + (c_)*(x_ )^2)^{(p_ )}, x\_Symbol] :> \text{Simp}[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] /;$   $\text{FreeQ}\{a, b, c, p\}, x \ \&\& \ \text{EqQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}]$

#### Rubi steps

$$\text{integral} = -\frac{1}{6}(2 - 3x)\sqrt{4 - 12x + 9x^2}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int \sqrt{4 - 12x + 9x^2} dx = \frac{\sqrt{(2 - 3x)^2 x(-4 + 3x)}}{-4 + 6x}$$

[In] Integrate[Sqrt[4 - 12\*x + 9\*x^2], x]

[Out] (Sqrt[(2 - 3\*x)^2]\*x\*(-4 + 3\*x))/(-4 + 6\*x)

**Maple [C] (warning: unable to verify)**

Result contains higher order function than in optimal. Order 9 vs. order 2.

Time = 0.17 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\text{csgn}(-2+3x)(-2+3x)^2}{6}$	16
gospers	$\frac{x(3x-4)\sqrt{(-2+3x)^2}}{-4+6x}$	25
risch	$\frac{3\sqrt{(-2+3x)^2} x^2}{2(-2+3x)} - \frac{2\sqrt{(-2+3x)^2} x}{-2+3x}$	42

[In] int(((−2+3\*x)^2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/6\*csgn(−2+3\*x)\*(−2+3\*x)^2

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.39

$$\int \sqrt{4 - 12x + 9x^2} dx = \frac{3}{2} x^2 - 2x$$

[In] integrate(((−2+3\*x)^2)^(1/2), x, algorithm="fricas")

[Out] 3/2\*x^2 - 2\*x

**Sympy [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \sqrt{4 - 12x + 9x^2} dx = \left(\frac{x}{2} - \frac{1}{3}\right) \sqrt{9x^2 - 12x + 4}$$

[In] integrate(((−2+3\*x)\*\*2)\*\*(1/2),x)

[Out] (x/2 - 1/3)\*sqrt(9\*x\*\*2 - 12\*x + 4)

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \sqrt{4 - 12x + 9x^2} dx = \frac{1}{2} \sqrt{9x^2 - 12x + 4}x - \frac{1}{3} \sqrt{9x^2 - 12x + 4}$$

[In] integrate(((−2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(9\*x^2 - 12\*x + 4)\*x - 1/3\*sqrt(9\*x^2 - 12\*x + 4)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \sqrt{4 - 12x + 9x^2} dx = \frac{1}{2} (3x^2 - 4x) \operatorname{sgn}(3x - 2) + \frac{2}{3} \operatorname{sgn}(3x - 2)$$

[In] integrate(((−2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2\*(3\*x^2 - 4\*x)\*sgn(3\*x - 2) + 2/3\*sgn(3\*x - 2)

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.57

$$\int \sqrt{4 - 12x + 9x^2} dx = \frac{|3x - 2| (3x - 2)}{6}$$

[In] int(((3\*x - 2)^2)^(1/2),x)

[Out] (abs(3\*x - 2)\*(3\*x - 2))/6

### 3.69 $\int \frac{1}{\sqrt{4-12x+9x^2}} dx$

Optimal result	376
Rubi [A] (verified)	376
Mathematica [A] (verified)	377
Maple [A] (verified)	377
Fricas [A] (verification not implemented)	378
Sympy [A] (verification not implemented)	378
Maxima [A] (verification not implemented)	378
Giac [A] (verification not implemented)	378
Mupad [B] (verification not implemented)	379

#### Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \frac{1}{\sqrt{4-12x+9x^2}} dx = -\frac{(2-3x)\log(2-3x)}{3\sqrt{4-12x+9x^2}}$$

[Out]  $-1/3*(2-3*x)*\ln(2-3*x)/((-2+3*x)^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {622, 31}

$$\int \frac{1}{\sqrt{4-12x+9x^2}} dx = -\frac{(2-3x)\log(2-3x)}{3\sqrt{9x^2-12x+4}}$$

[In] `Int[1/Sqrt[4 - 12*x + 9*x^2], x]`

[Out]  $-1/3*((2-3*x)*\text{Log}[2-3*x])/ \text{Sqrt}[4-12*x+9*x^2]$

#### Rule 31

`Int[((a_) + (b_.)*(x_))(-1), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

#### Rule 622

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[(b/2 + c*x)/Sqrt[a + b*x + c*x^2], Int[1/(b/2 + c*x), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`



Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(-6 + 9x) \int \frac{1}{-6+9x} dx}{\sqrt{4 - 12x + 9x^2}} \\ &= -\frac{(2 - 3x) \log(2 - 3x)}{3\sqrt{4 - 12x + 9x^2}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.90

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = -\frac{(2 - 3x) \log(2 - 3x)}{3\sqrt{(2 - 3x)^2}}$$

[In] Integrate[1/Sqrt[4 - 12\*x + 9\*x^2],x]

[Out] -1/3\*((2 - 3\*x)\*Log[2 - 3\*x])/Sqrt[(2 - 3\*x)^2]

### Maple [A] (verified)

Time = 1.91 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{(-2+3x) \ln(-2+3x)}{3\sqrt{(-2+3x)^2}}$	23
risch	$\frac{\sqrt{(-2+3x)^2} \ln(-2+3x)}{-6+9x}$	25
meijerg	$-\frac{2 \ln(1-\frac{3x}{2})}{3\sqrt{(-2+3x)^2}} + \frac{x \ln(1-\frac{3x}{2})}{\sqrt{(-2+3x)^2}}$	36

[In] int(1/((-2+3\*x)^2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3/((-2+3\*x)^2)^(1/2)\*(-2+3\*x)\*ln(-2+3\*x)

**Fricas [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.28

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = \frac{1}{3} \log(3x - 2)$$

[In] integrate(1/((-2+3\*x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/3\*log(3\*x - 2)

**Sympy [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.76

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = \frac{(x - \frac{2}{3}) \log(x - \frac{2}{3})}{3\sqrt{(x - \frac{2}{3})^2}}$$

[In] integrate(1/((-2+3\*x)\*\*2)\*\*(1/2),x)

[Out] (x - 2/3)\*log(x - 2/3)/(3\*sqrt((x - 2/3)\*\*2))

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = \frac{1}{3} \log\left(x - \frac{2}{3}\right)$$

[In] integrate(1/((-2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*log(x - 2/3)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = \frac{1}{3} \log(|3x - 2|) \operatorname{sgn}(3x - 2)$$

[In] integrate(1/((-2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] 1/3\*log(abs(3\*x - 2))\*sgn(3\*x - 2)

**Mupad [B] (verification not implemented)**

Time = 9.15 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.48

$$\int \frac{1}{\sqrt{4 - 12x + 9x^2}} dx = \frac{\ln(3x - 2) \operatorname{sign}(3x - 2)}{3}$$

[In] int(1/((3\*x - 2)^2)^(1/2),x)

[Out] (log(3\*x - 2)\*sign(3\*x - 2))/3

### 3.70 $\int \sqrt{-4 + 12x - 9x^2} dx$

Optimal result	380
Rubi [A] (verified)	380
Mathematica [A] (verified)	381
Maple [A] (verified)	381
Fricas [C] (verification not implemented)	381
Sympy [A] (verification not implemented)	382
Maxima [A] (verification not implemented)	382
Giac [C] (verification not implemented)	382
Mupad [B] (verification not implemented)	382

#### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \sqrt{-4 + 12x - 9x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{-4 + 12x - 9x^2}$$

[Out]  $-1/6*(2-3*x)*(-(-2+3*x)^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {623}

$$\int \sqrt{-4 + 12x - 9x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{-9x^2 + 12x - 4}$$

[In] `Int[Sqrt[-4 + 12*x - 9*x^2], x]`

[Out]  $-1/6*((2 - 3*x)*\text{Sqrt}[-4 + 12*x - 9*x^2])$

#### Rule 623

`Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] /; FreeQ[{a, b, c, p}, x] && EqQ[b^2 - 4*a*c, 0] && NeQ[p, -2^(-1)]`

#### Rubi steps

$$\text{integral} = -\frac{1}{6}(2 - 3x)\sqrt{-4 + 12x - 9x^2}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \sqrt{-4 + 12x - 9x^2} dx = \frac{\sqrt{-(2-3x)^2}x(-4+3x)}{-4+6x}$$

[In] Integrate[Sqrt[-4 + 12\*x - 9\*x^2],x]

[Out] (Sqrt[-(2 - 3\*x)^2]\*x\*(-4 + 3\*x))/(-4 + 6\*x)

**Maple [A] (verified)**

Time = 1.97 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

method	result	size
gospers	$\frac{x(3x-4)\sqrt{-(-2+3x)^2}}{-4+6x}$	27
default	$\frac{x(3x-4)\sqrt{-(-2+3x)^2}}{-4+6x}$	27
risch	$-\frac{2\sqrt{-(-2+3x)^2}x}{-2+3x} + \frac{3\sqrt{-(-2+3x)^2}x^2}{2(-2+3x)}$	46

[In] int((-(-2+3\*x)^2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*x\*(3\*x-4)\*(-(-2+3\*x)^2)^(1/2)/(-2+3\*x)

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.39

$$\int \sqrt{-4 + 12x - 9x^2} dx = \frac{3}{2}i x^2 - 2i x$$

[In] integrate((-(-2+3\*x)^2)^(1/2),x, algorithm="fricas")

[Out] 3/2\*I\*x^2 - 2\*I\*x

**Sympy [A] (verification not implemented)**

Time = 0.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \sqrt{-4 + 12x - 9x^2} dx = \left(\frac{x}{2} - \frac{1}{3}\right) \sqrt{-9x^2 + 12x - 4}$$

[In] integrate((-(-2+3\*x)\*\*2)\*\*(1/2),x)

[Out] (x/2 - 1/3)\*sqrt(-9\*x\*\*2 + 12\*x - 4)

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \sqrt{-4 + 12x - 9x^2} dx = \frac{1}{2} \sqrt{-9x^2 + 12x - 4}x - \frac{1}{3} \sqrt{-9x^2 + 12x - 4}$$

[In] integrate((-(-2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-9\*x^2 + 12\*x - 4)\*x - 1/3\*sqrt(-9\*x^2 + 12\*x - 4)

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \sqrt{-4 + 12x - 9x^2} dx = -\frac{1}{2}i(3x^2 - 4x)\operatorname{sgn}(-3x + 2) - \frac{2}{3}i\operatorname{sgn}(-3x + 2)$$

[In] integrate((-(-2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2\*I\*(3\*x^2 - 4\*x)\*sgn(-3\*x + 2) - 2/3\*I\*sgn(-3\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 9.11 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \sqrt{-4 + 12x - 9x^2} dx = \frac{(3x - 2) \sqrt{-(3x - 2)^2}}{6}$$

[In] int((- (3\*x - 2)^2)^(1/2),x)

[Out] ((3\*x - 2)\*(-(3\*x - 2)^2)^(1/2))/6

### 3.71 $\int \frac{1}{\sqrt{-4+12x-9x^2}} dx$

Optimal result . . . . .	383
Rubi [A] (verified) . . . . .	383
Mathematica [A] (verified) . . . . .	384
Maple [C] (verified) . . . . .	384
Fricas [C] (verification not implemented) . . . . .	385
Sympy [A] (verification not implemented) . . . . .	385
Maxima [C] (verification not implemented) . . . . .	385
Giac [C] (verification not implemented) . . . . .	385
Mupad [B] (verification not implemented) . . . . .	386

#### Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \frac{1}{\sqrt{-4+12x-9x^2}} dx = -\frac{(2-3x)\log(2-3x)}{3\sqrt{-4+12x-9x^2}}$$

[Out]  $-1/3*(2-3*x)*\ln(2-3*x)/(-(-2+3*x)^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {622, 31}

$$\int \frac{1}{\sqrt{-4+12x-9x^2}} dx = -\frac{(2-3x)\log(2-3x)}{3\sqrt{-9x^2+12x-4}}$$

[In] `Int[1/Sqrt[-4 + 12*x - 9*x^2], x]`

[Out]  $-1/3*((2-3*x)*\text{Log}[2-3*x])/ \text{Sqrt}[-4+12*x-9*x^2]$

#### Rule 31

`Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

#### Rule 622

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[(b/2 + c*x)/Sqrt[a + b*x + c*x^2], Int[1/(b/2 + c*x), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(6-9x) \int \frac{1}{6-9x} dx}{\sqrt{-4+12x-9x^2}} \\ &= -\frac{(2-3x) \log(2-3x)}{3\sqrt{-4+12x-9x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{\sqrt{-4+12x-9x^2}} dx = -\frac{(2-3x) \log(2-3x)}{3\sqrt{-(2-3x)^2}}$$

[In] Integrate[1/Sqrt[-4 + 12\*x - 9\*x^2],x]

[Out] -1/3\*((2 - 3\*x)\*Log[2 - 3\*x])/Sqrt[-(2 - 3\*x)^2]

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 2.09 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.34

method	result	size
meijerg	$-\frac{i \ln(1-\frac{3x}{2})}{3}$	10
default	$\frac{(-2+3x) \ln(-2+3x)}{3\sqrt{-(-2+3x)^2}}$	25
risch	$\frac{(-2+3x) \ln(-2+3x)}{3\sqrt{-(-2+3x)^2}}$	25

[In] int(1/(-(-2+3\*x)^2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] -1/3\*I\*ln(1-3/2\*x)



**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{-4 + 12x - 9x^2}} dx = -\frac{1}{3}i \log\left(x - \frac{2}{3}\right)$$

[In] integrate(1/(-(-2+3\*x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/3\*I\*log(x - 2/3)

**Sympy [A] (verification not implemented)**

Time = 0.42 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-4 + 12x - 9x^2}} dx = \frac{(x - \frac{2}{3}) \log(x - \frac{2}{3})}{3\sqrt{-(x - \frac{2}{3})^2}}$$

[In] integrate(1/(-(-2+3\*x)\*\*2)\*\*(1/2),x)

[Out] (x - 2/3)\*log(x - 2/3)/(3\*sqrt(-(x - 2/3)\*\*2))

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{-4 + 12x - 9x^2}} dx = \frac{1}{3}i \log\left(x - \frac{2}{3}\right)$$

[In] integrate(1/(-(-2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*I\*log(x - 2/3)

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt{-4 + 12x - 9x^2}} dx = \frac{i \log((-3ix + 2i)\operatorname{sgn}(-3x + 2))}{3 \operatorname{sgn}(-3x + 2)}$$

[In] integrate(1/(-(-2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] 1/3\*I\*log((-3\*I\*x + 2\*I)\*sgn(-3\*x + 2))/sgn(-3\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 9.08 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{-4 + 12x - 9x^2}} dx = -\frac{\ln(2 - 3x) \operatorname{sign}(3x - 2) i}{3}$$

[In] int(1/(-(3\*x - 2)^2)^(1/2),x)

[Out] -(log(2 - 3\*x)\*sign(3\*x - 2)\*1i)/3

### 3.72 $\int \sqrt{-4 - 12x - 9x^2} dx$

Optimal result	387
Rubi [A] (verified)	387
Mathematica [A] (verified)	388
Maple [A] (verified)	388
Fricas [C] (verification not implemented)	388
Sympy [A] (verification not implemented)	389
Maxima [A] (verification not implemented)	389
Giac [C] (verification not implemented)	389
Mupad [B] (verification not implemented)	389

#### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{-4 - 12x - 9x^2}$$

[Out] 1/6\*(2+3\*x)\*(-(2+3\*x)^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {623}

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{1}{6}(3x + 2)\sqrt{-9x^2 - 12x - 4}$$

[In] Int[Sqrt[-4 - 12\*x - 9\*x^2],x]

[Out] ((2 + 3\*x)\*Sqrt[-4 - 12\*x - 9\*x^2])/6

#### Rule 623

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] /; FreeQ[{a, b, c, p}, x] && EqQ[b^2 - 4\*a\*c, 0] && NeQ[p, -2^(-1)]

#### Rubi steps

$$\text{integral} = \frac{1}{6}(2 + 3x)\sqrt{-4 - 12x - 9x^2}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{x\sqrt{-(2+3x)^2(4+3x)}}{4+6x}$$

[In] Integrate[Sqrt[-4 - 12\*x - 9\*x^2],x]

[Out] (x\*Sqrt[-(2 + 3\*x)^2]\*(4 + 3\*x))/(4 + 6\*x)

**Maple [A] (verified)**

Time = 2.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.17

method	result	size
gospers	$\frac{x(4+3x)\sqrt{-(2+3x)^2}}{4+6x}$	27
default	$\frac{x(4+3x)\sqrt{-(2+3x)^2}}{4+6x}$	27
risch	$\frac{2\sqrt{-(2+3x)^2}x}{2+3x} + \frac{3\sqrt{-(2+3x)^2}x^2}{2(2+3x)}$	46

[In] int((-2+3\*x)^2^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*x\*(4+3\*x)\*(-(2+3\*x)^2)^(1/2)/(2+3\*x)

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.39

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{3}{2}i x^2 + 2i x$$

[In] integrate((-2+3\*x)^2^(1/2),x, algorithm="fricas")

[Out] 3/2\*I\*x^2 + 2\*I\*x

**Sympy [A] (verification not implemented)**

Time = 0.46 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

$$\int \sqrt{-4 - 12x - 9x^2} dx = \left(\frac{x}{2} + \frac{1}{3}\right) \sqrt{-9x^2 - 12x - 4}$$

[In] integrate((-2+3\*x)\*\*2)\*\*(1/2),x

[Out] (x/2 + 1/3)\*sqrt(-9\*x\*\*2 - 12\*x - 4)

**Maxima [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{1}{2} \sqrt{-9x^2 - 12x - 4}x + \frac{1}{3} \sqrt{-9x^2 - 12x - 4}$$

[In] integrate((-2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-9\*x^2 - 12\*x - 4)\*x + 1/3\*sqrt(-9\*x^2 - 12\*x - 4)

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.13

$$\int \sqrt{-4 - 12x - 9x^2} dx = -\frac{1}{2}i(3x^2 + 4x)\operatorname{sgn}(-3x - 2) - \frac{2}{3}i\operatorname{sgn}(-3x - 2)$$

[In] integrate((-2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2\*I\*(3\*x^2 + 4\*x)\*sgn(-3\*x - 2) - 2/3\*I\*sgn(-3\*x - 2)

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

$$\int \sqrt{-4 - 12x - 9x^2} dx = \frac{(3x + 2) \sqrt{-(3x + 2)^2}}{6}$$

[In] int((-3\*x + 2)^2)^(1/2),x

[Out] ((3\*x + 2)\*(-3\*x + 2)^2)^(1/2)/6

### 3.73 $\int \frac{1}{\sqrt{-4-12x-9x^2}} dx$

Optimal result	390
Rubi [A] (verified)	390
Mathematica [A] (verified)	391
Maple [C] (verified)	391
Fricas [C] (verification not implemented)	392
Sympy [A] (verification not implemented)	392
Maxima [C] (verification not implemented)	392
Giac [C] (verification not implemented)	392
Mupad [B] (verification not implemented)	393

#### Optimal result

Integrand size = 14, antiderivative size = 29

$$\int \frac{1}{\sqrt{-4-12x-9x^2}} dx = \frac{(2+3x)\log(2+3x)}{3\sqrt{-4-12x-9x^2}}$$

[Out] 1/3\*(2+3\*x)\*ln(2+3\*x)/(-(2+3\*x)^2)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {622, 31}

$$\int \frac{1}{\sqrt{-4-12x-9x^2}} dx = \frac{(3x+2)\log(3x+2)}{3\sqrt{-9x^2-12x-4}}$$

[In] Int[1/Sqrt[-4 - 12\*x - 9\*x^2], x]

[Out] ((2 + 3\*x)\*Log[2 + 3\*x])/(3\*Sqrt[-4 - 12\*x - 9\*x^2])

#### Rule 31

Int[((a\_) + (b\_.)\*(x\_))^(n\_), x\_Symbol] := Simp[Log[RemoveContent[a + b\*x, x]]/b, x] /; FreeQ[{a, b}, x]

#### Rule 622

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[(b/2 + c\*x)/Sqrt[a + b\*x + c\*x^2], Int[1/(b/2 + c\*x), x], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= - \left( - \frac{(-6 - 9x) \int \frac{1}{-6-9x} dx}{\sqrt{-4 - 12x - 9x^2}} \right) \\ &= \frac{(2 + 3x) \log(2 + 3x)}{3\sqrt{-4 - 12x - 9x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = \frac{(2 + 3x) \log(2 + 3x)}{3\sqrt{-(2 + 3x)^2}}$$

[In] Integrate[1/Sqrt[-4 - 12\*x - 9\*x^2],x]

[Out] ((2 + 3\*x)\*Log[2 + 3\*x])/(3\*Sqrt[-(2 + 3\*x)^2])

**Maple [C] (verified)**

Result contains complex when optimal does not.

Time = 2.22 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.34

method	result	size
meijerg	$-\frac{i \ln(1 + \frac{3x}{2})}{3}$	10
default	$\frac{(2+3x) \ln(2+3x)}{3\sqrt{-(2+3x)^2}}$	25
risch	$\frac{(2+3x) \ln(2+3x)}{3\sqrt{-(2+3x)^2}}$	25

[In] int(1/(-(2+3\*x)^2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] -1/3\*I\*ln(1+3/2\*x)

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = -\frac{1}{3}i \log\left(x + \frac{2}{3}\right)$$

[In] integrate(1/(-(2+3\*x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/3\*I\*log(x + 2/3)

**Sympy [A] (verification not implemented)**

Time = 0.41 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = \frac{(x + \frac{2}{3}) \log(x + \frac{2}{3})}{3\sqrt{-(x + \frac{2}{3})^2}}$$

[In] integrate(1/(-(2+3\*x)\*\*2)\*\*(1/2),x)

[Out] (x + 2/3)\*log(x + 2/3)/(3\*sqrt(-(x + 2/3)\*\*2))

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.21

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = \frac{1}{3}i \log\left(x + \frac{2}{3}\right)$$

[In] integrate(1/(-(2+3\*x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*I\*log(x + 2/3)

**Giac [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = \frac{i \log((-3ix - 2i)\operatorname{sgn}(-3x - 2))}{3 \operatorname{sgn}(-3x - 2)}$$

[In] integrate(1/(-(2+3\*x)^2)^(1/2),x, algorithm="giac")

[Out] 1/3\*I\*log((-3\*I\*x - 2\*I)\*sgn(-3\*x - 2))/sgn(-3\*x - 2)



**Mupad [B] (verification not implemented)**

Time = 9.12 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{-4 - 12x - 9x^2}} dx = -\frac{\ln(-3x - 2) \operatorname{sign}(3x + 2) i}{3}$$

[In] int(1/(-(3\*x + 2)^2)^(1/2),x)

[Out] -(log(- 3\*x - 2)\*sign(3\*x + 2)\*1i)/3

$$3.74 \quad \int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx$$

Optimal result	394
Rubi [A] (verified)	394
Mathematica [A] (verified)	395
Maple [B] (verified)	396
Fricas [B] (verification not implemented)	396
Sympy [B] (verification not implemented)	397
Maxima [B] (verification not implemented)	397
Giac [B] (verification not implemented)	398
Mupad [B] (verification not implemented)	398

### Optimal result

Integrand size = 23, antiderivative size = 109

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{(1-b-2cx)^6}{384c^6} - \frac{5(1-b-2cx)^7}{896c^6} + \frac{5(1-b-2cx)^8}{1024c^6} - \frac{5(1-b-2cx)^9}{2304c^6} + \frac{(1-b-2cx)^{10}}{2048c^6} - \frac{(1-b-2cx)^{11}}{22528c^6}$$

[Out] 1/384\*(-2\*c\*x-b+1)^6/c^6-5/896\*(-2\*c\*x-b+1)^7/c^6+5/1024\*(-2\*c\*x-b+1)^8/c^6-5/2304\*(-2\*c\*x-b+1)^9/c^6+1/2048\*(-2\*c\*x-b+1)^10/c^6-1/22528\*(-2\*c\*x-b+1)^11/c^6

### Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {624, 45}

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx = -\frac{(-b-2cx+1)^{11}}{22528c^6} + \frac{(-b-2cx+1)^{10}}{2048c^6} - \frac{5(-b-2cx+1)^9}{2304c^6} + \frac{5(-b-2cx+1)^8}{1024c^6} - \frac{5(-b-2cx+1)^7}{896c^6} + \frac{(-b-2cx+1)^6}{384c^6}$$

[In] Int[((-1 + b^2)/(4\*c) + b\*x + c\*x^2)^5, x]

[Out] (1 - b - 2\*c\*x)^6/(384\*c^6) - (5\*(1 - b - 2\*c\*x)^7)/(896\*c^6) + (5\*(1 - b - 2\*c\*x)^8)/(1024\*c^6) - (5\*(1 - b - 2\*c\*x)^9)/(2304\*c^6) + (1 - b - 2\*c\*x)^10/(2048\*c^6) - (1 - b - 2\*c\*x)^11/(22528\*c^6)

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

### Rule 624

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[1/c^p, Int[Simp[b/2 - q/2 + c*x, x]^p*Simp[b/2 + q/2 + c
*x, x]^p, x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && PerfectSquareQ[b^2 - 4*a*c]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \left(\frac{1}{2}(-1+b) + cx\right)^5 \left(\frac{1+b}{2} + cx\right)^5 dx}{c^5} \\ &= \frac{\int \left(\left(\frac{1}{2}(-1+b) + cx\right)^5 + 5\left(\frac{1}{2}(-1+b) + cx\right)^6 + 10\left(\frac{1}{2}(-1+b) + cx\right)^7 + 10\left(\frac{1}{2}(-1+b) + cx\right)^8 + 5\left(\frac{1}{2}(-1+b) + cx\right)^9 + \left(\frac{1}{2}(-1+b) + cx\right)^{10}\right) dx}{c^5} \\ &= \frac{(1-b-2cx)^6}{384c^6} - \frac{5(1-b-2cx)^7}{896c^6} + \frac{5(1-b-2cx)^8}{1024c^6} \\ &\quad - \frac{5(1-b-2cx)^9}{2304c^6} + \frac{(1-b-2cx)^{10}}{2048c^6} - \frac{(1-b-2cx)^{11}}{22528c^6} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.89

$$\begin{aligned} \int \left(\frac{-1+b^2}{4c} + bx + cx^2\right)^5 dx &= \frac{(-1+b^2)^5 x}{1024c^5} + \frac{5b(-1+b^2)^4 x^2}{512c^4} \\ &\quad + \frac{5(-1+b^2)^3(-1+9b^2)x^3}{768c^3} + \frac{5b(-1+b^2)^2(-1+3b^2)x^4}{64c^2} \\ &\quad + \frac{(-1+b^2)(1-14b^2+21b^4)x^5}{32c} + \frac{1}{48}b(15-70b^2+63b^4)x^6 \\ &\quad + \frac{5}{56}(1-14b^2+21b^4)cx^7 + \frac{5}{8}b(-1+3b^2)c^2x^8 \\ &\quad + \frac{5}{36}(-1+9b^2)c^3x^9 + \frac{1}{2}bc^4x^{10} + \frac{c^5x^{11}}{11} \end{aligned}$$

```
[In] Integrate[((-1 + b^2)/(4*c) + b*x + c*x^2)^5, x]
```

```
[Out] ((-1 + b^2)^5*x)/(1024*c^5) + (5*b*(-1 + b^2)^4*x^2)/(512*c^4) + (5*(-1 + b
^2)^3*(-1 + 9*b^2)*x^3)/(768*c^3) + (5*b*(-1 + b^2)^2*(-1 + 3*b^2)*x^4)/(64
*c^2) + ((-1 + b^2)*(1 - 14*b^2 + 21*b^4)*x^5)/(32*c) + (b*(15 - 70*b^2 + 6
3*b^4)*x^6)/48 + (5*(1 - 14*b^2 + 21*b^4)*c*x^7)/56 + (5*b*(-1 + 3*b^2)*c^2
*x^8)/8 + (5*(-1 + 9*b^2)*c^3*x^9)/36 + (b*c^4*x^10)/2 + (c^5*x^11)/11
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 272 vs.  $2(97) = 194$ .

Time = 2.28 (sec) , antiderivative size = 273, normalized size of antiderivative = 2.50

method	result
norman	$\frac{(\frac{5}{4}b^2c^7 - \frac{5}{36}c^7)x^9 + (\frac{15}{8}b^3c^6 - \frac{5}{8}bc^6)x^8 + (\frac{15}{8}b^4c^5 - \frac{5}{4}b^2c^5 + \frac{5}{56}c^5)x^7 + (\frac{21}{16}b^5c^4 - \frac{35}{24}c^4b^3 + \frac{5}{16}bc^4)x^6 + (\frac{15}{64}b^7c^2 - \frac{35}{64}b^5c^2 + \frac{25}{64}c^2b^3 - \frac{5}{64}c^2)}{x(64512c^{10}x^{10} + 354816c^9bx^9 + 887040x^8b^2c^8 + 1330560b^3c^7x^7 + 1330560x^6b^4c^6 - 98560x^8c^8 + 931392x^5b^5c^5 - 443520bc^7x^7 + 46560x^6b^2c^8 + 1330560b^3c^7x^8 + 1330560x^7b^4c^6 - 98560x^9c^8 + 931392x^6b^5c^5 - 443520bc^7x^8 + 46560x^7b^2c^8)}$
gosper	
parallelrisch	
risch	$\frac{5bx^6}{16} - \frac{35b^3x^6}{24} - \frac{x}{1024c^5} + \frac{21b^5x^6}{16} - \frac{x^5}{32c} + \frac{15b^4cx^7}{8} + \frac{21b^6x^5}{32c} + \frac{15b^2x^5}{32c} + \frac{5c^3x^9b^2}{4} + \frac{15b^7x^4}{64c^2} - \frac{35b^5x^4}{64c^2} + \frac{5b^6x^4}{512c^2}$
default	$\frac{c^5x^{11}}{11} + \frac{bc^4x^{10}}{2} + \frac{(256(b^2-1)c^3 + 4096b^2c^3 + 4c(32(24b^2-8)c^2 + 1024b^2c^2))x^9}{9216} + \frac{(1024(b^2-1)c^2b + 4b(32(24b^2-8)c^2 + 1024b^2c^2))x^8}{9216}$

[In] int((1/4\*(b^2-1)/c+b\*x+c\*x^2)^5,x,method=\_RETURNVERBOSE)

[Out]  $((5/4*b^2*c^7 - 5/36*c^7)*x^9 + (15/8*b^3*c^6 - 5/8*b*c^6)*x^8 + (15/8*b^4*c^5 - 5/4*b^2*c^5 + 5/56*c^5)*x^7 + (21/16*b^5*c^4 - 35/24*c^4*b^3 + 5/16*b*c^4)*x^6 + (15/64*b^7*c^2 - 35/64*b^5*c^2 + 25/64*c^2*b^3 - 5/64*b*c^2)*x^4 + (21/32*c^3*b^6 - 35/32*b^4*c^3 + 15/32*b^2*c^3 - 1/32*c^3)*x^5 + (5/512*b^9 - 5/128*b^7 + 15/256*b^5 - 5/128*b^3 + 5/512*b)*x^2 + (15/256*b^8*c - 35/192*b^6*c + 25/128*b^4*c - 5/64*b^2*c + 5/768*c)*x^3 + 1/11*c^9*x^{11} + 1/2*b*c^8*x^{10} + 1/1024*(b^{10} - 5*b^8 + 10*b^6 - 10*b^4 + 5*b^2 - 1)/c^4)x/c^4$

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 233 vs.  $2(85) = 170$ .

Time = 0.26 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.14

$$\int \left( \frac{-1 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{64512c^{10}x^{11} + 354816bc^9x^{10} + 98560(9b^2 - 1)c^8x^9 + 443520(3b^3 - b)c^7x^8 + 63360(21b^4 - 14b^2 + 1)c^6x^7 + 14784(63b^5 - 70b^3 + 15b)c^5x^6 + 22176(21b^6 - 35b^4 + 15b^2 - 1)c^4x^5 + 55440(3b^7 - 7b^5 + 5b^3 - b)c^3x^4 + 4620(9b^8 - 28b^6 + 30b^4 - 12b^2 + 1)c^2x^3 + 6930(b^9 - 4b^7 + 6b^5 - 4b^3 + b)c^2x^2 + 693(b^{10} - 5b^8 + 10b^6 - 10b^4 + 5b^2 - 1)cx}{c^5}$$

[In] integrate((1/4\*(b^2-1)/c+b\*x+c\*x^2)^5,x, algorithm="fricas")

[Out]  $1/709632*(64512*c^{10}*x^{11} + 354816*b*c^9*x^{10} + 98560*(9*b^2 - 1)*c^8*x^9 + 443520*(3*b^3 - b)*c^7*x^8 + 63360*(21*b^4 - 14*b^2 + 1)*c^6*x^7 + 14784*(63*b^5 - 70*b^3 + 15*b)*c^5*x^6 + 22176*(21*b^6 - 35*b^4 + 15*b^2 - 1)*c^4*x^5 + 55440*(3*b^7 - 7*b^5 + 5*b^3 - b)*c^3*x^4 + 4620*(9*b^8 - 28*b^6 + 30*b^4 - 12*b^2 + 1)*c^2*x^3 + 6930*(b^9 - 4*b^7 + 6*b^5 - 4*b^3 + b)*c*x^2 + 693*(b^{10} - 5*b^8 + 10*b^6 - 10*b^4 + 5*b^2 - 1)*x)/c^5$

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(95) = 190.

Time = 0.09 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.32

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{bc^4x^{10}}{2} + \frac{c^5x^{11}}{11} + x^9 \cdot \left( \frac{5b^2c^3}{4} - \frac{5c^3}{36} \right) + x^8$$

$$\cdot \left( \frac{15b^3c^2}{8} - \frac{5bc^2}{8} \right) + x^7 \cdot \left( \frac{15b^4c}{8} - \frac{5b^2c}{4} + \frac{5c}{56} \right) + x^6$$

$$\cdot \left( \frac{21b^5}{16} - \frac{35b^3}{24} + \frac{5b}{16} \right) + \frac{x^5 \cdot (21b^6 - 35b^4 + 15b^2 - 1)}{32c}$$

$$+ \frac{x^4 \cdot (15b^7 - 35b^5 + 25b^3 - 5b)}{64c^2}$$

$$+ \frac{x^3 \cdot (45b^8 - 140b^6 + 150b^4 - 60b^2 + 5)}{768c^3}$$

$$+ \frac{x^2 \cdot (5b^9 - 20b^7 + 30b^5 - 20b^3 + 5b)}{512c^4}$$

$$+ \frac{x(b^{10} - 5b^8 + 10b^6 - 10b^4 + 5b^2 - 1)}{1024c^5}$$

[In] integrate((1/4\*(b\*\*2-1)/c+b\*x+c\*x\*\*2)\*\*5,x)

[Out] b\*c\*\*4\*x\*\*10/2 + c\*\*5\*x\*\*11/11 + x\*\*9\*(5\*b\*\*2\*c\*\*3/4 - 5\*c\*\*3/36) + x\*\*8\*(15\*b\*\*3\*c\*\*2/8 - 5\*b\*c\*\*2/8) + x\*\*7\*(15\*b\*\*4\*c/8 - 5\*b\*\*2\*c/4 + 5\*c/56) + x\*\*6\*(21\*b\*\*5/16 - 35\*b\*\*3/24 + 5\*b/16) + x\*\*5\*(21\*b\*\*6 - 35\*b\*\*4 + 15\*b\*\*2 - 1)/(32\*c) + x\*\*4\*(15\*b\*\*7 - 35\*b\*\*5 + 25\*b\*\*3 - 5\*b)/(64\*c\*\*2) + x\*\*3\*(45\*b\*\*8 - 140\*b\*\*6 + 150\*b\*\*4 - 60\*b\*\*2 + 5)/(768\*c\*\*3) + x\*\*2\*(5\*b\*\*9 - 20\*b\*\*7 + 30\*b\*\*5 - 20\*b\*\*3 + 5\*b)/(512\*c\*\*4) + x\*(b\*\*10 - 5\*b\*\*8 + 10\*b\*\*6 - 10\*b\*\*4 + 5\*b\*\*2 - 1)/(1024\*c\*\*5)

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(85) = 170.

Time = 0.19 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.15

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{1}{11} c^5 x^{11} + \frac{1}{2} bc^4 x^{10} + \frac{10}{9} b^2 c^3 x^9 + \frac{5}{4} b^3 c^2 x^8 + \frac{5}{7} b^4 c x^7 + \frac{1}{6} b^5 x^6$$

$$+ \frac{5(2cx^3 + 3bx^2)(b^2 - 1)^4}{1536c^4} + \frac{(6c^2x^5 + 15bcx^4 + 10b^2x^3)(b^2 - 1)^3}{192c^3}$$

$$+ \frac{(20c^3x^7 + 70bc^2x^6 + 84b^2cx^5 + 35b^3x^4)(b^2 - 1)^2}{224c^2}$$

$$+ \frac{(70c^4x^9 + 315bc^3x^8 + 540b^2c^2x^7 + 420b^3cx^6 + 126b^4x^5)(b^2 - 1)}{504c} + \frac{(b^2 - 1)^5 x}{1024c^5}$$

[In] integrate((1/4\*(b^2-1)/c+b\*x+c\*x^2)^5,x, algorithm="maxima")

[Out] 1/11\*c^5\*x^11 + 1/2\*b\*c^4\*x^10 + 10/9\*b^2\*c^3\*x^9 + 5/4\*b^3\*c^2\*x^8 + 5/7\*b^4\*c\*x^7 + 1/6\*b^5\*x^6 + 5/1536\*(2\*c\*x^3 + 3\*b\*x^2)\*(b^2 - 1)^4/c^4 + 1/192\*(6\*c^2\*x^5 + 15\*b\*c\*x^4 + 10\*b^2\*x^3)\*(b^2 - 1)^3/c^3 + 1/224\*(20\*c^3\*x^7 + 70\*b\*c^2\*x^6 + 84\*b^2\*c\*x^5 + 35\*b^3\*x^4)\*(b^2 - 1)^2/c^2 + 1/504\*(70\*c^4\*x^9 + 315\*b\*c^3\*x^8 + 540\*b^2\*c^2\*x^7 + 420\*b^3\*c\*x^6 + 126\*b^4\*x^5)\*(b^2 - 1)/c + 1/1024\*(b^2 - 1)^5\*x/c^5

## Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(85) = 170.

Time = 0.27 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.06

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{64512 c^{10} x^{11} + 354816 b c^9 x^{10} + 887040 b^2 c^8 x^9 + 1330560 b^3 c^7 x^8 + 1330560 b^4 c^6 x^7 - 98560 c^8 x^9 + 931392 b^5 c^5 x^6 - 443520 b^6 c^4 x^5 - 887040 b^7 c^3 x^4 + 166320 b^8 c^2 x^3 - 776160 b^9 c x^2 + 693 b^{10} x - 129360 b^{11} c^2 x^3 + 332640 b^{12} c^4 x^5 - 27720 b^{13} c^6 x^7 + 277200 b^{14} c^8 x^9 - 3465 b^{15} x + 138600 b^{16} c^2 x^3 - 22176 c^{14} x^5 + 41580 b^{15} c^6 x^7 - 55440 b^{16} c^8 x^9 + 6930 b^{17} c^2 x^3 - 55440 b^{18} c^4 x^5 - 27720 b^{19} c^6 x^7 - 6930 b^{20} c^8 x^9 + 4620 c^{12} x^3 + 6930 b^2 c^4 x^5 + 3465 b^4 c^6 x^7 - 693 x^9)/c^5$$

[In] integrate((1/4\*(b^2-1)/c+b\*x+c\*x^2)^5,x, algorithm="giac")

[Out] 1/709632\*(64512\*c^10\*x^11 + 354816\*b\*c^9\*x^10 + 887040\*b^2\*c^8\*x^9 + 1330560\*b^3\*c^7\*x^8 + 1330560\*b^4\*c^6\*x^7 - 98560\*c^8\*x^9 + 931392\*b^5\*c^5\*x^6 - 443520\*b^6\*c^4\*x^5 - 887040\*b^7\*c^3\*x^4 + 166320\*b^8\*c^2\*x^3 + 693\*b^10\*x - 129360\*b^11\*c^2\*x^3 + 332640\*b^12\*c^4\*x^5 - 27720\*b^13\*c^6\*x^7 + 277200\*b^14\*c^8\*x^9 - 3465\*b^15\*x + 138600\*b^16\*c^2\*x^3 - 22176\*c^14\*x^5 + 41580\*b^15\*c^6\*x^7 - 55440\*b^16\*c^8\*x^9 + 6930\*b^17\*c^2\*x^3 - 55440\*b^18\*c^4\*x^5 - 27720\*b^19\*c^6\*x^7 - 6930\*b^20\*c^8\*x^9 + 4620\*c^12\*x^3 + 6930\*b^2\*c^4\*x^5 + 3465\*b^4\*c^6\*x^7 - 693\*x^9)/c^5

## Mupad [B] (verification not implemented)

Time = 9.16 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.69

$$\int \left( \frac{-1+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{c^5 x^{11}}{11} + \frac{x (b^2 - 1)^5}{1024 c^5} + \frac{b x^6 (63 b^4 - 70 b^2 + 15)}{48} + \frac{5 c x^7 (21 b^4 - 14 b^2 + 1)}{56} + \frac{b c^4 x^{10}}{2} + \frac{5 c^3 x^9 (9 b^2 - 1)}{36} + \frac{x^5 (21 b^6 - 35 b^4 + 15 b^2 - 1)}{32 c} + \frac{5 b c^2 x^8 (3 b^2 - 1)}{8} + \frac{5 b x^2 (b^2 - 1)^4}{512 c^4} + \frac{5 x^3 (b^2 - 1)^3 (9 b^2 - 1)}{768 c^3} + \frac{5 b x^4 (b^2 - 1)^2 (3 b^2 - 1)}{64 c^2}$$

[In]  $\text{int}((b*x + c*x^2 + (b^2/4 - 1/4)/c)^5, x)$

[Out]  $(c^5*x^{11})/11 + (x*(b^2 - 1)^5)/(1024*c^5) + (b*x^6*(63*b^4 - 70*b^2 + 15))/48 + (5*c*x^7*(21*b^4 - 14*b^2 + 1))/56 + (b*c^4*x^{10})/2 + (5*c^3*x^9*(9*b^2 - 1))/36 + (x^5*(15*b^2 - 35*b^4 + 21*b^6 - 1))/(32*c) + (5*b*c^2*x^8*(3*b^2 - 1))/8 + (5*b*x^2*(b^2 - 1)^4)/(512*c^4) + (5*x^3*(b^2 - 1)^3*(9*b^2 - 1))/(768*c^3) + (5*b*x^4*(b^2 - 1)^2*(3*b^2 - 1))/(64*c^2)$

$$3.75 \quad \int \left( \frac{-4+b^2}{4c} + bx + cx^2 \right)^5 dx$$

Optimal result	400
Rubi [A] (verified)	400
Mathematica [A] (verified)	401
Maple [B] (verified)	402
Fricas [B] (verification not implemented)	402
Sympy [B] (verification not implemented)	403
Maxima [B] (verification not implemented)	403
Giac [B] (verification not implemented)	404
Mupad [B] (verification not implemented)	404

### Optimal result

Integrand size = 23, antiderivative size = 109

$$\int \left( \frac{-4+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{(2-b-2cx)^6}{12c^6} - \frac{5(2-b-2cx)^7}{56c^6} + \frac{5(2-b-2cx)^8}{128c^6} - \frac{5(2-b-2cx)^9}{576c^6} + \frac{(2-b-2cx)^{10}}{1024c^6} - \frac{(2-b-2cx)^{11}}{22528c^6}$$

[Out] 1/12\*(-2\*c\*x-b+2)^6/c^6-5/56\*(-2\*c\*x-b+2)^7/c^6+5/128\*(-2\*c\*x-b+2)^8/c^6-5/576\*(-2\*c\*x-b+2)^9/c^6+1/1024\*(-2\*c\*x-b+2)^10/c^6-1/22528\*(-2\*c\*x-b+2)^11/c^6

### Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {624, 45}

$$\int \left( \frac{-4+b^2}{4c} + bx + cx^2 \right)^5 dx = -\frac{(-b-2cx+2)^{11}}{22528c^6} + \frac{(-b-2cx+2)^{10}}{1024c^6} - \frac{5(-b-2cx+2)^9}{576c^6} + \frac{5(-b-2cx+2)^8}{128c^6} - \frac{5(-b-2cx+2)^7}{56c^6} + \frac{(-b-2cx+2)^6}{12c^6}$$

[In] Int[((-4 + b^2)/(4\*c) + b\*x + c\*x^2)^5, x]

[Out] (2 - b - 2\*c\*x)^6/(12\*c^6) - (5\*(2 - b - 2\*c\*x)^7)/(56\*c^6) + (5\*(2 - b - 2\*c\*x)^8)/(128\*c^6) - (5\*(2 - b - 2\*c\*x)^9)/(576\*c^6) + (2 - b - 2\*c\*x)^10/(1024\*c^6) - (2 - b - 2\*c\*x)^11/(22528\*c^6)

Rule 45



```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

### Rule 624

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[1/c^p, Int[Simp[b/2 - q/2 + c*x, x]^p*Simp[b/2 + q/2 + c
*x, x]^p, x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && PerfectSquareQ[b^2 - 4*a*c]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \left(\frac{1}{2}(-2+b) + cx\right)^5 \left(\frac{2+b}{2} + cx\right)^5 dx}{c^5} \\ &= \frac{\int \left(32\left(\frac{1}{2}(-2+b) + cx\right)^5 + 80\left(\frac{1}{2}(-2+b) + cx\right)^6 + 80\left(\frac{1}{2}(-2+b) + cx\right)^7 + 40\left(\frac{1}{2}(-2+b) + cx\right)^8\right)}{c^5} \\ &= \frac{(2-b-2cx)^6}{12c^6} - \frac{5(2-b-2cx)^7}{56c^6} + \frac{5(2-b-2cx)^8}{128c^6} \\ &\quad - \frac{5(2-b-2cx)^9}{576c^6} + \frac{(2-b-2cx)^{10}}{1024c^6} - \frac{(2-b-2cx)^{11}}{22528c^6} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.89

$$\begin{aligned} \int \left(\frac{-4+b^2}{4c} + bx + cx^2\right)^5 dx &= \frac{(-4+b^2)^5 x}{1024c^5} + \frac{5b(-4+b^2)^4 x^2}{512c^4} \\ &\quad + \frac{5(-4+b^2)^3(-4+9b^2)x^3}{768c^3} + \frac{5b(-4+b^2)^2(-4+3b^2)x^4}{64c^2} \\ &\quad + \frac{(-4+b^2)(16-56b^2+21b^4)x^5}{32c} \\ &\quad + \frac{1}{48}b(240-280b^2+63b^4)x^6 + \frac{5}{56}(16-56b^2+21b^4)cx^7 \\ &\quad + \frac{5}{8}b(-4+3b^2)c^2x^8 + \frac{5}{36}(-4+9b^2)c^3x^9 + \frac{1}{2}bc^4x^{10} + \frac{c^5x^{11}}{11} \end{aligned}$$

```
[In] Integrate[((-4 + b^2)/(4*c) + b*x + c*x^2)^5, x]
```

```
[Out] ((-4 + b^2)^5*x)/(1024*c^5) + (5*b*(-4 + b^2)^4*x^2)/(512*c^4) + (5*(-4 + b
^2)^3*(-4 + 9*b^2)*x^3)/(768*c^3) + (5*b*(-4 + b^2)^2*(-4 + 3*b^2)*x^4)/(64
*c^2) + ((-4 + b^2)*(16 - 56*b^2 + 21*b^4)*x^5)/(32*c) + (b*(240 - 280*b^2
+ 63*b^4)*x^6)/48 + (5*(16 - 56*b^2 + 21*b^4)*c*x^7)/56 + (5*b*(-4 + 3*b^2)
*c^2*x^8)/8 + (5*(-4 + 9*b^2)*c^3*x^9)/36 + (b*c^4*x^10)/2 + (c^5*x^11)/11
```

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(97) = 194.

Time = 2.23 (sec) , antiderivative size = 273, normalized size of antiderivative = 2.50

method	result
norman	$\frac{(\frac{5}{4}b^2c^7 - \frac{5}{9}c^7)x^9 + (\frac{15}{8}b^3c^6 - \frac{5}{2}bc^6)x^8 + (\frac{15}{8}b^4c^5 - 5b^2c^5 + \frac{10}{7}c^5)x^7 + (\frac{21}{16}b^5c^4 - \frac{35}{6}c^4b^3 + 5bc^4)x^6 + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)x^5 + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)x^4 + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)x^3 + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)x^2 + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)x + (\frac{15}{64}b^7c^2 - \frac{35}{16}b^5c^2 + \frac{25}{4}c^2b^3 - 5bc^2 + 4c^2)}$
gospers	$x(64512c^{10}x^{10} + 354816c^9bx^9 + 887040x^8b^2c^8 + 1330560b^3c^7x^7 + 1330560x^6b^4c^6 - 394240x^8c^8 + 931392x^5b^5c^5 - 1774080bc^7x^7 + 4620000c^9x^5 + 1330560b^3c^7x^7 + 1330560x^6b^4c^6 - 394240x^8c^8 + 931392x^5b^5c^5 - 1774080bc^7x^7 + 4620000c^9x^5)$
parallemrisch	$64512c^{10}x^{11} + 354816c^9bx^{10} + 887040x^9b^2c^8 + 1330560b^3c^7x^8 + 1330560x^7b^4c^6 - 394240x^9c^8 + 931392x^6b^5c^5 - 1774080bc^7x^8 + 4620000c^9x^5 + 1330560b^3c^7x^7 + 1330560x^6b^4c^6 - 394240x^8c^8 + 931392x^5b^5c^5 - 1774080bc^7x^7 + 4620000c^9x^5$
risch	$5bx^6 - \frac{35b^3x^6}{6} - \frac{x}{c^5} + \frac{21b^5x^6}{16} - \frac{2x^5}{c} + \frac{15b^4cx^7}{8} + \frac{21b^6x^5}{32c} + \frac{15b^2x^5}{2c} + \frac{5c^3x^9b^2}{4} + \frac{15b^7x^4}{64c^2} - \frac{35b^5x^4}{16c^2} + \frac{5b^9x^2}{512c^4}$
default	$\frac{c^5x^{11}}{11} + \frac{bc^4x^{10}}{2} + \frac{(256(b^2-4)c^3 + 4096b^2c^3 + 4c(32(24b^2-32)c^2 + 1024b^2c^2))x^9}{9216} + \frac{(1024(b^2-4)c^2b + 4b(32(24b^2-32)c^2 + 1024b^2c^2))x^8}{9216}$

[In] int((1/4\*(b^2-4)/c+b\*x+c\*x^2)^5,x,method=\_RETURNVERBOSE)

[Out] ((5/4\*b^2\*c^7-5/9\*c^7)\*x^9+(15/8\*b^3\*c^6-5/2\*b\*c^6)\*x^8+(15/8\*b^4\*c^5-5\*b^2\*c^5+10/7\*c^5)\*x^7+(21/16\*b^5\*c^4-35/6\*c^4\*b^3+5\*b\*c^4)\*x^6+(15/64\*b^7\*c^2-35/16\*b^5\*c^2+25/4\*c^2\*b^3-5\*b\*c^2)\*x^4+(21/32\*c^3\*b^6-35/8\*b^4\*c^3+15/2\*b^2\*c^3-2\*c^3)\*x^5+(5/512\*b^9-5/32\*b^7+15/16\*b^5-5/2\*b^3+5/2\*b)\*x^2+(15/256\*b^8\*c-35/48\*b^6\*c+25/8\*b^4\*c-5\*b^2\*c+5/3\*c)\*x^3+1/11\*c^9\*x^11+1/2\*b\*c^8\*x^10+1/1024\*(b^10-20\*b^8+160\*b^6-640\*b^4+1280\*b^2-1024)/c\*x)/c^4

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 235 vs. 2(85) = 170.

Time = 0.45 (sec) , antiderivative size = 235, normalized size of antiderivative = 2.16

$$\int \left( \frac{-4 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{64512c^{10}x^{11} + 354816bc^9x^{10} + 98560(9b^2 - 4)c^8x^9 + 443520(3b^3 - 4b)c^7x^8 + 63360(21b^4 - 56b^2 + 16)c^6x^7 + 14784(63b^5 - 280b^3 + 240b)c^5x^6 + 22176(21b^6 - 140b^4 + 240b^2 - 64)c^4x^5 + 55440(3b^7 - 28b^5 + 80b^3 - 64b)c^3x^4 + 4620(9b^8 - 112b^6 + 480b^4 - 768b^2 + 256)c^2x^3 + 6930(b^9 - 16b^7 + 96b^5 - 256b^3 + 256b)c^2x^2 + 693(b^{10} - 20b^8 + 160b^6 - 640b^4 + 1280b^2 - 1024)x}{c^5}$$

[In] integrate((1/4\*(b^2-4)/c+b\*x+c\*x^2)^5,x, algorithm="fricas")

[Out] 1/709632\*(64512\*c^10\*x^11 + 354816\*b\*c^9\*x^10 + 98560\*(9\*b^2 - 4)\*c^8\*x^9 + 443520\*(3\*b^3 - 4\*b)\*c^7\*x^8 + 63360\*(21\*b^4 - 56\*b^2 + 16)\*c^6\*x^7 + 14784\*(63\*b^5 - 280\*b^3 + 240\*b)\*c^5\*x^6 + 22176\*(21\*b^6 - 140\*b^4 + 240\*b^2 - 64)\*c^4\*x^5 + 55440\*(3\*b^7 - 28\*b^5 + 80\*b^3 - 64\*b)\*c^3\*x^4 + 4620\*(9\*b^8 - 112\*b^6 + 480\*b^4 - 768\*b^2 + 256)\*c^2\*x^3 + 6930\*(b^9 - 16\*b^7 + 96\*b^5 - 256\*b^3 + 256\*b)\*c^2\*x^2 + 693\*(b^10 - 20\*b^8 + 160\*b^6 - 640\*b^4 + 1280\*b^2 - 1024)\*x)/c^5

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 250 vs. 2(95) = 190.

Time = 0.09 (sec) , antiderivative size = 250, normalized size of antiderivative = 2.29

$$\int \left( \frac{-4 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{bc^4x^{10}}{2} + \frac{c^5x^{11}}{11} + x^9 \cdot \left( \frac{5b^2c^3}{4} - \frac{5c^3}{9} \right) + x^8$$

$$\cdot \left( \frac{15b^3c^2}{8} - \frac{5bc^2}{2} \right) + x^7 \cdot \left( \frac{15b^4c}{8} - 5b^2c + \frac{10c}{7} \right) + x^6$$

$$\cdot \left( \frac{21b^5}{16} - \frac{35b^3}{6} + 5b \right) + \frac{x^5 \cdot (21b^6 - 140b^4 + 240b^2 - 64)}{32c}$$

$$+ \frac{x^4 \cdot (15b^7 - 140b^5 + 400b^3 - 320b)}{64c^2}$$

$$+ \frac{x^3 \cdot (45b^8 - 560b^6 + 2400b^4 - 3840b^2 + 1280)}{768c^3}$$

$$+ \frac{x^2 \cdot (5b^9 - 80b^7 + 480b^5 - 1280b^3 + 1280b)}{512c^4}$$

$$+ \frac{x(b^{10} - 20b^8 + 160b^6 - 640b^4 + 1280b^2 - 1024)}{1024c^5}$$

[In] integrate((1/4\*(b\*\*2-4)/c+b\*x+c\*x\*\*2)\*\*5,x)

[Out] b\*c\*\*4\*x\*\*10/2 + c\*\*5\*x\*\*11/11 + x\*\*9\*(5\*b\*\*2\*c\*\*3/4 - 5\*c\*\*3/9) + x\*\*8\*(15\*b\*\*3\*c\*\*2/8 - 5\*b\*c\*\*2/2) + x\*\*7\*(15\*b\*\*4\*c/8 - 5\*b\*\*2\*c + 10\*c/7) + x\*\*6\*(21\*b\*\*5/16 - 35\*b\*\*3/6 + 5\*b) + x\*\*5\*(21\*b\*\*6 - 140\*b\*\*4 + 240\*b\*\*2 - 64)/(32\*c) + x\*\*4\*(15\*b\*\*7 - 140\*b\*\*5 + 400\*b\*\*3 - 320\*b)/(64\*c\*\*2) + x\*\*3\*(45\*b\*\*8 - 560\*b\*\*6 + 2400\*b\*\*4 - 3840\*b\*\*2 + 1280)/(768\*c\*\*3) + x\*\*2\*(5\*b\*\*9 - 80\*b\*\*7 + 480\*b\*\*5 - 1280\*b\*\*3 + 1280\*b)/(512\*c\*\*4) + x\*(b\*\*10 - 20\*b\*\*8 + 160\*b\*\*6 - 640\*b\*\*4 + 1280\*b\*\*2 - 1024)/(1024\*c\*\*5)

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(85) = 170.

Time = 0.19 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.15

$$\int \left( \frac{-4 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{1}{11} c^5 x^{11} + \frac{1}{2} bc^4 x^{10} + \frac{10}{9} b^2 c^3 x^9 + \frac{5}{4} b^3 c^2 x^8 + \frac{5}{7} b^4 c x^7 + \frac{1}{6} b^5 x^6$$

$$+ \frac{5(2cx^3 + 3bx^2)(b^2 - 4)^4}{1536c^4} + \frac{(6c^2x^5 + 15bcx^4 + 10b^2x^3)(b^2 - 4)^3}{192c^3}$$

$$+ \frac{(20c^3x^7 + 70bc^2x^6 + 84b^2cx^5 + 35b^3x^4)(b^2 - 4)^2}{224c^2}$$

$$+ \frac{(70c^4x^9 + 315bc^3x^8 + 540b^2c^2x^7 + 420b^3cx^6 + 126b^4x^5)(b^2 - 4)}{504c} + \frac{(b^2 - 4)^5 x}{1024c^5}$$

[In] integrate((1/4\*(b^2-4)/c+b\*x+c\*x^2)^5,x, algorithm="maxima")

[Out] 1/11\*c^5\*x^11 + 1/2\*b\*c^4\*x^10 + 10/9\*b^2\*c^3\*x^9 + 5/4\*b^3\*c^2\*x^8 + 5/7\*b^4\*c\*x^7 + 1/6\*b^5\*x^6 + 5/1536\*(2\*c\*x^3 + 3\*b\*x^2)\*(b^2 - 4)^4/c^4 + 1/192\*(6\*c^2\*x^5 + 15\*b\*c\*x^4 + 10\*b^2\*x^3)\*(b^2 - 4)^3/c^3 + 1/224\*(20\*c^3\*x^7 + 70\*b\*c^2\*x^6 + 84\*b^2\*c\*x^5 + 35\*b^3\*x^4)\*(b^2 - 4)^2/c^2 + 1/504\*(70\*c^4\*x^9 + 315\*b\*c^3\*x^8 + 540\*b^2\*c^2\*x^7 + 420\*b^3\*c\*x^6 + 126\*b^4\*x^5)\*(b^2 - 4)/c + 1/1024\*(b^2 - 4)^5\*x/c^5

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(85) = 170.

Time = 0.27 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.06

$$\int \left( \frac{-4 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{64512 c^{10} x^{11} + 354816 b c^9 x^{10} + 887040 b^2 c^8 x^9 + 1330560 b^3 c^7 x^8 + 1330560 b^4 c^6 x^7 - 394240 c^8 x^9 + 931392 c^9 x^{10} + \dots}{c^5}$$

[In] integrate((1/4\*(b^2-4)/c+b\*x+c\*x^2)^5,x, algorithm="giac")

[Out] 1/709632\*(64512\*c^10\*x^11 + 354816\*b\*c^9\*x^10 + 887040\*b^2\*c^8\*x^9 + 1330560\*b^3\*c^7\*x^8 + 1330560\*b^4\*c^6\*x^7 - 394240\*c^8\*x^9 + 931392\*b^5\*c^5\*x^6 - 1774080\*b\*c^7\*x^8 + 465696\*b^6\*c^4\*x^5 - 3548160\*b^2\*c^6\*x^7 + 166320\*b^7\*c^3\*x^4 - 4139520\*b^3\*c^5\*x^6 + 41580\*b^8\*c^2\*x^3 - 3104640\*b^4\*c^4\*x^5 + 1013760\*c^6\*x^7 + 6930\*b^9\*c\*x^2 - 1552320\*b^5\*c^3\*x^4 + 3548160\*b\*c^5\*x^6 + 693\*b^10\*x - 517440\*b^6\*c^2\*x^3 + 5322240\*b^2\*c^4\*x^5 - 110880\*b^7\*c\*x^2 + 4435200\*b^3\*c^3\*x^4 - 13860\*b^8\*x + 2217600\*b^4\*c^2\*x^3 - 1419264\*c^4\*x^5 + 665280\*b^5\*c\*x^2 - 3548160\*b\*c^3\*x^4 + 110880\*b^6\*x - 3548160\*b^2\*c^2\*x^3 - 1774080\*b^3\*c\*x^2 - 443520\*b^4\*x + 1182720\*c^2\*x^3 + 1774080\*b\*c\*x^2 + 887040\*b^2\*x - 709632\*x)/c^5

### Mupad [B] (verification not implemented)

Time = 9.17 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.69

$$\int \left( \frac{-4 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{c^5 x^{11}}{11} + \frac{x (b^2 - 4)^5}{1024 c^5} + \frac{b x^6 (63 b^4 - 280 b^2 + 240)}{48} + \frac{5 c x^7 (21 b^4 - 56 b^2 + 16)}{56} + \frac{b c^4 x^{10}}{2} + \frac{5 c^3 x^9 (9 b^2 - 4)}{36} + \frac{x^5 (21 b^6 - 140 b^4 + 240 b^2 - 64)}{32 c} + \frac{5 b c^2 x^8 (3 b^2 - 4)}{8} + \frac{5 b x^2 (b^2 - 4)^4}{512 c^4} + \frac{5 x^3 (b^2 - 4)^3 (9 b^2 - 4)}{768 c^3} + \frac{5 b x^4 (b^2 - 4)^2 (3 b^2 - 4)}{64 c^2}$$

[In] `int((b*x + c*x^2 + (b^2/4 - 1)/c)^5,x)`

[Out]  $(c^5x^{11})/11 + (x(b^2 - 4)^5)/(1024c^5) + (bx^6(63b^4 - 280b^2 + 240))/48 + (5c^3x^7(21b^4 - 56b^2 + 16))/56 + (bc^4x^{10})/2 + (5c^3x^9(9b^2 - 4))/36 + (x^5(240b^2 - 140b^4 + 21b^6 - 64))/(32c) + (5bc^2x^8(3b^2 - 4))/8 + (5bx^2(b^2 - 4)^4)/(512c^4) + (5x^3(b^2 - 4)^3(9b^2 - 4))/(768c^3) + (5bx^4(b^2 - 4)^2(3b^2 - 4))/(64c^2)$

$$3.76 \quad \int \left( \frac{-9+b^2}{4c} + bx + cx^2 \right)^5 dx$$

Optimal result . . . . .	406
Rubi [A] (verified) . . . . .	406
Mathematica [A] (verified) . . . . .	407
Maple [B] (verified) . . . . .	408
Fricas [B] (verification not implemented) . . . . .	408
Sympy [B] (verification not implemented) . . . . .	409
Maxima [B] (verification not implemented) . . . . .	410
Giac [B] (verification not implemented) . . . . .	410
Mupad [B] (verification not implemented) . . . . .	411

### Optimal result

Integrand size = 23, antiderivative size = 109

$$\int \left( \frac{-9+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{81(3-b-2cx)^6}{128c^6} - \frac{405(3-b-2cx)^7}{896c^6} + \frac{135(3-b-2cx)^8}{1024c^6} - \frac{5(3-b-2cx)^9}{256c^6} + \frac{3(3-b-2cx)^{10}}{2048c^6} - \frac{(3-b-2cx)^{11}}{22528c^6}$$

[Out] 81/128\*(-2\*c\*x-b+3)^6/c^6-405/896\*(-2\*c\*x-b+3)^7/c^6+135/1024\*(-2\*c\*x-b+3)^8/c^6-5/256\*(-2\*c\*x-b+3)^9/c^6+3/2048\*(-2\*c\*x-b+3)^10/c^6-1/22528\*(-2\*c\*x-b+3)^11/c^6

### Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {624, 45}

$$\int \left( \frac{-9+b^2}{4c} + bx + cx^2 \right)^5 dx = -\frac{(-b-2cx+3)^{11}}{22528c^6} + \frac{3(-b-2cx+3)^{10}}{2048c^6} - \frac{5(-b-2cx+3)^9}{256c^6} + \frac{135(-b-2cx+3)^8}{1024c^6} - \frac{405(-b-2cx+3)^7}{896c^6} + \frac{81(-b-2cx+3)^6}{128c^6}$$

[In] Int[((-9 + b^2)/(4\*c) + b\*x + c\*x^2)^5, x]

[Out] (81\*(3 - b - 2\*c\*x)^6)/(128\*c^6) - (405\*(3 - b - 2\*c\*x)^7)/(896\*c^6) + (135\*(3 - b - 2\*c\*x)^8)/(1024\*c^6) - (5\*(3 - b - 2\*c\*x)^9)/(256\*c^6) + (3\*(3 - b - 2\*c\*x)^10)/(2048\*c^6) - (3 - b - 2\*c\*x)^11/(22528\*c^6)

## Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

## Rule 624

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[1/c^p, Int[Simp[b/2 - q/2 + c*x, x]^p*Simp[b/2 + q/2 + c
*x, x]^p, x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && PerfectSquareQ[b^2 - 4*a*c]
```

## Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \left(\frac{1}{2}(-3+b) + cx\right)^5 \left(\frac{3+b}{2} + cx\right)^5 dx}{c^5} \\ &= \frac{\int \left(243\left(\frac{1}{2}(-3+b) + cx\right)^5 + 405\left(\frac{1}{2}(-3+b) + cx\right)^6 + 270\left(\frac{1}{2}(-3+b) + cx\right)^7 + 90\left(\frac{1}{2}(-3+b) + cx\right)^8 + 27\left(\frac{1}{2}(-3+b) + cx\right)^9 + 9\left(\frac{1}{2}(-3+b) + cx\right)^{10} + \left(\frac{1}{2}(-3+b) + cx\right)^{11}\right) dx}{c^5} \\ &= \frac{81(3-b-2cx)^6}{128c^6} - \frac{405(3-b-2cx)^7}{896c^6} + \frac{135(3-b-2cx)^8}{1024c^6} \\ &\quad - \frac{5(3-b-2cx)^9}{256c^6} + \frac{3(3-b-2cx)^{10}}{2048c^6} - \frac{(3-b-2cx)^{11}}{22528c^6} \end{aligned}$$

## Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.82

$$\begin{aligned} \int \left(\frac{-9+b^2}{4c} + bx + cx^2\right)^5 dx &= \frac{(-9+b^2)^5 x}{1024c^5} + \frac{5b(-9+b^2)^4 x^2}{512c^4} \\ &\quad + \frac{15(-9+b^2)^3(-1+b^2)x^3}{256c^3} + \frac{15b(-9+b^2)^2(-3+b^2)x^4}{64c^2} \\ &\quad + \frac{3(-9+b^2)(27-42b^2+7b^4)x^5}{32c} \\ &\quad + \frac{3}{16}b(135-70b^2+7b^4)x^6 + \frac{15}{56}(27-42b^2+7b^4)cx^7 \\ &\quad + \frac{15}{8}b(-3+b^2)c^2x^8 + \frac{5}{4}(-1+b^2)c^3x^9 + \frac{1}{2}bc^4x^{10} + \frac{c^5x^{11}}{11} \end{aligned}$$

```
[In] Integrate[((-9 + b^2)/(4*c) + b*x + c*x^2)^5, x]
```

```
[Out] ((-9 + b^2)^5*x)/(1024*c^5) + (5*b*(-9 + b^2)^4*x^2)/(512*c^4) + (15*(-9 +
b^2)^3*(-1 + b^2)*x^3)/(256*c^3) + (15*b*(-9 + b^2)^2*(-3 + b^2)*x^4)/(64*c
```

$$\begin{aligned} &^2) + (3*(-9 + b^2)*(27 - 42*b^2 + 7*b^4)*x^5)/(32*c) + (3*b*(135 - 70*b^2 \\ &+ 7*b^4)*x^6)/16 + (15*(27 - 42*b^2 + 7*b^4)*c*x^7)/56 + (15*b*(-3 + b^2)*c \\ &^2*x^8)/8 + (5*(-1 + b^2)*c^3*x^9)/4 + (b*c^4*x^10)/2 + (c^5*x^11)/11 \end{aligned}$$

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(97) = 194.

Time = 2.26 (sec) , antiderivative size = 273, normalized size of antiderivative = 2.50

method	result
norman	$\frac{(\frac{5}{4}b^2c^7 - \frac{5}{4}c^7)x^9 + (\frac{15}{8}b^3c^6 - \frac{45}{8}bc^6)x^8 + (\frac{15}{8}b^4c^5 - \frac{45}{4}b^2c^5 + \frac{405}{56}c^5)x^7 + (\frac{21}{16}b^5c^4 - \frac{105}{8}c^4b^3 + \frac{405}{16}bc^4)x^6 + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)x^5 + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)x^4 + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)x^3 + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)x^2 + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)x + (\frac{15}{64}b^7c^2 - \frac{315}{64}b^5c^2 + \frac{2025}{64}c^2b^3)}$
gospers	$x(7168c^{10}x^{10} + 39424c^9bx^9 + 98560x^8b^2c^8 + 147840b^3c^7x^7 + 147840x^6b^4c^6 - 98560x^8c^8 + 103488x^5b^5c^5 - 443520bc^7x^7 + 51744b^6c^6)$
parallelrisc	$7168c^{10}x^{11} + 39424c^9bx^{10} + 98560x^9b^2c^8 + 147840b^3c^7x^8 + 147840x^7b^4c^6 - 98560x^9c^8 + 103488x^6b^5c^5 - 443520bc^7x^8 + 51744b^6c^4c^2$
risc	$\frac{405bx^6}{16} - \frac{105b^3x^6}{8} - \frac{59049x}{1024c^5} + \frac{21b^5x^6}{16} - \frac{729x^5}{32c} + \frac{15b^4cx^7}{8} + \frac{21b^6x^5}{32c} + \frac{1215b^2x^5}{32c} + \frac{5c^3x^9b^2}{4} + \frac{15b^7x^4}{64c^2} - \frac{315b^5c^2}{64c^2}$
default	$\frac{c^5x^{11}}{11} + \frac{bc^4x^{10}}{2} + \frac{(256(b^2-9)c^3 + 4096b^2c^3 + 4c(32(24b^2-72)c^2 + 1024b^2c^2))x^9}{9216} + \frac{(1024(b^2-9)c^2b + 4b(32(24b^2-72)c^2 + 1024b^2c^2))x^8}{9216}$

[In] int((1/4\*(b^2-9)/c+b\*x+c\*x^2)^5,x,method=\_RETURNVERBOSE)

[Out] ((5/4\*b^2\*c^7-5/4\*c^7)\*x^9+(15/8\*b^3\*c^6-45/8\*b\*c^6)\*x^8+(15/8\*b^4\*c^5-45/4\*b^2\*c^5+405/56\*c^5)\*x^7+(21/16\*b^5\*c^4-105/8\*c^4\*b^3+405/16\*b\*c^4)\*x^6+(15/64\*b^7\*c^2-315/64\*b^5\*c^2+2025/64\*c^2\*b^3-3645/64\*b\*c^2)\*x^4+(21/32\*c^3\*b^6-315/32\*b^4\*c^3+1215/32\*b^2\*c^3-729/32\*c^3)\*x^5+(5/512\*b^9-45/128\*b^7+1215/256\*b^5-3645/128\*b^3+32805/512\*b)\*x^2+(15/256\*b^8\*c-105/64\*b^6\*c+2025/128\*b^4\*c-3645/64\*b^2\*c+10935/256\*c)\*x^3+1/11\*c^9\*x^11+1/2\*b\*c^8\*x^10+1/1024\*(b^10-45\*b^8+810\*b^6-7290\*b^4+32805\*b^2-59049)/c\*x)/c^4

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(85) = 170.

Time = 0.43 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.08

$$\int \left( \frac{-9 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{7168c^{10}x^{11} + 39424bc^9x^{10} + 98560(b^2-1)c^8x^9 + 147840(b^3-3b)c^7x^8 + 21120(7b^4-42b^2+27)c^6x^7 + \dots}{9216}$$

[In] integrate((1/4\*(b^2-9)/c+b\*x+c\*x^2)^5,x, algorithm="fricas")



[Out]  $\frac{1}{78848} \cdot (7168 \cdot c^{10} \cdot x^{11} + 39424 \cdot b \cdot c^9 \cdot x^{10} + 98560 \cdot (b^2 - 1) \cdot c^8 \cdot x^9 + 147840 \cdot (b^3 - 3b) \cdot c^7 \cdot x^8 + 21120 \cdot (7b^4 - 42b^2 + 27) \cdot c^6 \cdot x^7 + 14784 \cdot (7b^5 - 70b^3 + 135b) \cdot c^5 \cdot x^6 + 7392 \cdot (7b^6 - 105b^4 + 405b^2 - 243) \cdot c^4 \cdot x^5 + 18480 \cdot (b^7 - 21b^5 + 135b^3 - 243b) \cdot c^3 \cdot x^4 + 4620 \cdot (b^8 - 28b^6 + 270b^4 - 972b^2 + 729) \cdot c^2 \cdot x^3 + 770 \cdot (b^9 - 36b^7 + 486b^5 - 2916b^3 + 6561b) \cdot c \cdot x^2 + 77 \cdot (b^{10} - 45b^8 + 810b^6 - 7290b^4 + 32805b^2 - 59049) \cdot x) / c^5$

## Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(99) = 198.

Time = 0.08 (sec) , antiderivative size = 253, normalized size of antiderivative = 2.32

$$\int \left( \frac{-9 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{bc^4x^{10}}{2} + \frac{c^5x^{11}}{11} + x^9 \cdot \left( \frac{5b^2c^3}{4} - \frac{5c^3}{4} \right) + x^8 \cdot \left( \frac{15b^3c^2}{8} - \frac{45bc^2}{8} \right) + x^7 \cdot \left( \frac{15b^4c}{8} - \frac{45b^2c}{4} + \frac{405c}{56} \right) + x^6 \cdot \left( \frac{21b^5}{16} - \frac{105b^3}{8} + \frac{405b}{16} \right) + \frac{x^5 \cdot (21b^6 - 315b^4 + 1215b^2 - 729)}{32c} + \frac{x^4 \cdot (15b^7 - 315b^5 + 2025b^3 - 3645b)}{64c^2} + \frac{x^3 \cdot (15b^8 - 420b^6 + 4050b^4 - 14580b^2 + 10935)}{256c^3} + \frac{x^2 \cdot (5b^9 - 180b^7 + 2430b^5 - 14580b^3 + 32805b)}{512c^4} + \frac{x(b^{10} - 45b^8 + 810b^6 - 7290b^4 + 32805b^2 - 59049)}{1024c^5}$$

[In] `integrate((1/4*(b**2-9)/c+b*x+c*x**2)**5,x)`

[Out]  $b \cdot c^{4} \cdot x^{10} / 2 + c^{5} \cdot x^{11} / 11 + x^{9} \cdot (5 \cdot b^{2} \cdot c^{3} / 4 - 5 \cdot c^{3} / 4) + x^{8} \cdot (15 \cdot b^{3} \cdot c^{2} / 8 - 45 \cdot b \cdot c^{2} / 8) + x^{7} \cdot (15 \cdot b^{4} \cdot c / 8 - 45 \cdot b^{2} \cdot c / 4 + 405 \cdot c / 56) + x^{6} \cdot (21 \cdot b^{5} / 16 - 105 \cdot b^{3} / 8 + 405 \cdot b / 16) + x^{5} \cdot (21 \cdot b^{6} - 315 \cdot b^{4} + 1215 \cdot b^{2} - 729) / (32 \cdot c) + x^{4} \cdot (15 \cdot b^{7} - 315 \cdot b^{5} + 2025 \cdot b^{3} - 3645 \cdot b) / (64 \cdot c^{2}) + x^{3} \cdot (15 \cdot b^{8} - 420 \cdot b^{6} + 4050 \cdot b^{4} - 14580 \cdot b^{2} + 10935) / (256 \cdot c^{3}) + x^{2} \cdot (5 \cdot b^{9} - 180 \cdot b^{7} + 2430 \cdot b^{5} - 14580 \cdot b^{3} + 32805 \cdot b) / (512 \cdot c^{4}) + x \cdot (b^{10} - 45 \cdot b^{8} + 810 \cdot b^{6} - 7290 \cdot b^{4} + 32805 \cdot b^{2} - 59049) / (1024 \cdot c^{5})$

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(85) = 170.

Time = 0.20 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.15

$$\int \left( \frac{-9 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{1}{11} c^5 x^{11} + \frac{1}{2} bc^4 x^{10} + \frac{10}{9} b^2 c^3 x^9 + \frac{5}{4} b^3 c^2 x^8 + \frac{5}{7} b^4 c x^7 + \frac{1}{6} b^5 x^6$$

$$+ \frac{5(2cx^3 + 3bx^2)(b^2 - 9)^4}{1536c^4} + \frac{(6c^2x^5 + 15bcx^4 + 10b^2x^3)(b^2 - 9)^3}{192c^3}$$

$$+ \frac{(20c^3x^7 + 70bc^2x^6 + 84b^2cx^5 + 35b^3x^4)(b^2 - 9)^2}{224c^2}$$

$$+ \frac{(70c^4x^9 + 315bc^3x^8 + 540b^2c^2x^7 + 420b^3cx^6 + 126b^4x^5)(b^2 - 9)}{504c} + \frac{(b^2 - 9)^5 x}{1024c^5}$$

[In] integrate((1/4\*(b^2-9)/c+b\*x+c\*x^2)^5,x, algorithm="maxima")

[Out] 1/11\*c^5\*x^11 + 1/2\*b\*c^4\*x^10 + 10/9\*b^2\*c^3\*x^9 + 5/4\*b^3\*c^2\*x^8 + 5/7\*b^4\*c\*x^7 + 1/6\*b^5\*x^6 + 5/1536\*(2\*c\*x^3 + 3\*b\*x^2)\*(b^2 - 9)^4/c^4 + 1/192\*(6\*c^2\*x^5 + 15\*b\*c\*x^4 + 10\*b^2\*x^3)\*(b^2 - 9)^3/c^3 + 1/224\*(20\*c^3\*x^7 + 70\*b\*c^2\*x^6 + 84\*b^2\*c\*x^5 + 35\*b^3\*x^4)\*(b^2 - 9)^2/c^2 + 1/504\*(70\*c^4\*x^9 + 315\*b\*c^3\*x^8 + 540\*b^2\*c^2\*x^7 + 420\*b^3\*c\*x^6 + 126\*b^4\*x^5)\*(b^2 - 9)/c + 1/1024\*(b^2 - 9)^5\*x/c^5

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(85) = 170.

Time = 0.26 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.06

$$\int \left( \frac{-9 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{7168c^{10}x^{11} + 39424bc^9x^{10} + 98560b^2c^8x^9 + 147840b^3c^7x^8 + 147840b^4c^6x^7 - 98560c^8x^9 + 103488b^5c^5x^6 - 443520b^6c^4x^5 - 887040b^7c^3x^4 - 1034880b^8c^2x^3 - 776160b^9cx^2 + 1995840b^{10}x - 129360b^{11} - 2993760b^{12} - 27720b^{13} + 2494800b^{14} - 3465b^{15} + 1247400b^{16} - 1796256c^4x^5 + 374220b^5cx^2}{1024c^5}$$

[In] integrate((1/4\*(b^2-9)/c+b\*x+c\*x^2)^5,x, algorithm="giac")

[Out] 1/78848\*(7168\*c^10\*x^11 + 39424\*b\*c^9\*x^10 + 98560\*b^2\*c^8\*x^9 + 147840\*b^3\*c^7\*x^8 + 147840\*b^4\*c^6\*x^7 - 98560\*c^8\*x^9 + 103488\*b^5\*c^5\*x^6 - 443520\*b^6\*c^4\*x^5 + 51744\*b^6\*c^4\*x^5 - 887040\*b^2\*c^6\*x^7 + 18480\*b^7\*c^3\*x^4 - 1034880\*b^3\*c^5\*x^6 + 4620\*b^8\*c^2\*x^3 - 776160\*b^4\*c^4\*x^5 + 570240\*c^6\*x^7 + 770\*b^9\*c\*x^2 - 388080\*b^5\*c^3\*x^4 + 1995840\*b\*c^5\*x^6 + 77\*b^10\*x - 129360\*b^6\*c^2\*x^3 + 2993760\*b^2\*c^4\*x^5 - 27720\*b^7\*c\*x^2 + 2494800\*b^3\*c^3\*x^4 - 3465\*b^8\*x + 1247400\*b^4\*c^2\*x^3 - 1796256\*c^4\*x^5 + 374220\*b^5\*c\*x^2)

$$\begin{aligned} & - 4490640*b*c^3*x^4 + 62370*b^6*x - 4490640*b^2*c^2*x^3 - 2245320*b^3*c*x^2 \\ & - 561330*b^4*x + 3367980*c^2*x^3 + 5051970*b*c*x^2 + 2525985*b^2*x - 45467 \\ & 73*x)/c^5 \end{aligned}$$

### Mupad [B] (verification not implemented)

Time = 9.20 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.61

$$\begin{aligned} \int \left( \frac{-9 + b^2}{4c} + bx + cx^2 \right)^5 dx = & \frac{c^5 x^{11}}{11} + \frac{5 c^3 x^9 (b^2 - 1)}{4} + \frac{x (b^2 - 9)^5}{1024 c^5} \\ & + \frac{3 b x^6 (7 b^4 - 70 b^2 + 135)}{16} + \frac{15 c x^7 (7 b^4 - 42 b^2 + 27)}{56} \\ & + \frac{b c^4 x^{10}}{2} + \frac{3 x^5 (7 b^6 - 105 b^4 + 405 b^2 - 243)}{32 c} \\ & + \frac{15 b c^2 x^8 (b^2 - 3)}{8} + \frac{15 x^3 (b^2 - 1) (b^2 - 9)^3}{256 c^3} \\ & + \frac{5 b x^2 (b^2 - 9)^4}{512 c^4} + \frac{15 b x^4 (b^2 - 3) (b^2 - 9)^2}{64 c^2} \end{aligned}$$

[In] int((b\*x + c\*x^2 + (b^2/4 - 9/4)/c)^5,x)

[Out] (c^5\*x^11)/11 + (5\*c^3\*x^9\*(b^2 - 1))/4 + (x\*(b^2 - 9)^5)/(1024\*c^5) + (3\*b\*x^6\*(7\*b^4 - 70\*b^2 + 135))/16 + (15\*c\*x^7\*(7\*b^4 - 42\*b^2 + 27))/56 + (b\*c^4\*x^10)/2 + (3\*x^5\*(405\*b^2 - 105\*b^4 + 7\*b^6 - 243))/(32\*c) + (15\*b\*c^2\*x^8\*(b^2 - 3))/8 + (15\*x^3\*(b^2 - 1)\*(b^2 - 9)^3)/(256\*c^3) + (5\*b\*x^2\*(b^2 - 9)^4)/(512\*c^4) + (15\*b\*x^4\*(b^2 - 3)\*(b^2 - 9)^2)/(64\*c^2)

$$3.77 \quad \int \left( \frac{-16+b^2}{4c} + bx + cx^2 \right)^5 dx$$

Optimal result	412
Rubi [A] (verified)	412
Mathematica [A] (verified)	413
Maple [B] (verified)	414
Fricas [B] (verification not implemented)	415
Sympy [B] (verification not implemented)	415
Maxima [B] (verification not implemented)	416
Giac [B] (verification not implemented)	416
Mupad [B] (verification not implemented)	417

### Optimal result

Integrand size = 23, antiderivative size = 109

$$\int \left( \frac{-16+b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{8(4-b-2cx)^6}{3c^6} - \frac{10(4-b-2cx)^7}{7c^6} + \frac{5(4-b-2cx)^8}{16c^6} - \frac{5(4-b-2cx)^9}{144c^6} + \frac{(4-b-2cx)^{10}}{512c^6} - \frac{(4-b-2cx)^{11}}{22528c^6}$$

[Out]  $8/3*(-2*c*x-b+4)^6/c^6-10/7*(-2*c*x-b+4)^7/c^6+5/16*(-2*c*x-b+4)^8/c^6-5/144*(-2*c*x-b+4)^9/c^6+1/512*(-2*c*x-b+4)^{10}/c^6-1/22528*(-2*c*x-b+4)^{11}/c^6$

### Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$ , Rules used = {624, 45}

$$\int \left( \frac{-16+b^2}{4c} + bx + cx^2 \right)^5 dx = -\frac{(-b-2cx+4)^{11}}{22528c^6} + \frac{(-b-2cx+4)^{10}}{512c^6} - \frac{5(-b-2cx+4)^9}{144c^6} + \frac{5(-b-2cx+4)^8}{16c^6} - \frac{10(-b-2cx+4)^7}{7c^6} + \frac{8(-b-2cx+4)^6}{3c^6}$$

[In] Int[((-16 + b^2)/(4\*c) + b\*x + c\*x^2)^5,x]

[Out]  $(8*(4-b-2*c*x)^6)/(3*c^6) - (10*(4-b-2*c*x)^7)/(7*c^6) + (5*(4-b-2*c*x)^8)/(16*c^6) - (5*(4-b-2*c*x)^9)/(144*c^6) + (4-b-2*c*x)^{10}/(512*c^6) - (4-b-2*c*x)^{11}/(22528*c^6)$

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 624

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[1/c^p, Int[Simp[b/2 - q/2 + c*x, x]^p*Simp[b/2 + q/2 + c
*x, x]^p, x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[p,
0] && PerfectSquareQ[b^2 - 4*a*c]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \left(\frac{1}{2}(-4+b) + cx\right)^5 \left(\frac{4+b}{2} + cx\right)^5 dx}{c^5} \\ &= \frac{\int \left(1024\left(\frac{1}{2}(-4+b) + cx\right)^5 + 1280\left(\frac{1}{2}(-4+b) + cx\right)^6 + 640\left(\frac{1}{2}(-4+b) + cx\right)^7 + 160\left(\frac{1}{2}(-4+b) + cx\right)^8 + 32\left(\frac{1}{2}(-4+b) + cx\right)^9\right) dx}{c^5} \\ &= \frac{8(4-b-2cx)^6}{3c^6} - \frac{10(4-b-2cx)^7}{7c^6} + \frac{5(4-b-2cx)^8}{16c^6} \\ &\quad - \frac{5(4-b-2cx)^9}{144c^6} + \frac{(4-b-2cx)^{10}}{512c^6} - \frac{(4-b-2cx)^{11}}{22528c^6} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.89

$$\begin{aligned} \int \left(\frac{-16+b^2}{4c} + bx + cx^2\right)^5 dx &= \frac{(-16+b^2)^5 x}{1024c^5} + \frac{5b(-16+b^2)^4 x^2}{512c^4} \\ &\quad + \frac{5(-16+b^2)^3(-16+9b^2) x^3}{768c^3} \\ &\quad + \frac{5b(-16+b^2)^2(-16+3b^2) x^4}{64c^2} \\ &\quad + \frac{(-16+b^2)(256-224b^2+21b^4) x^5}{32c} \\ &\quad + \frac{1}{48}b(3840-1120b^2+63b^4) x^6 \\ &\quad + \frac{5}{56}(256-224b^2+21b^4) cx^7 + \frac{5}{8}b(-16+3b^2) c^2 x^8 \\ &\quad + \frac{5}{36}(-16+9b^2) c^3 x^9 + \frac{1}{2}bc^4 x^{10} + \frac{c^5 x^{11}}{11} \end{aligned}$$



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 235 vs. 2(85) = 170.

Time = 0.38 (sec) , antiderivative size = 235, normalized size of antiderivative = 2.16

$$\int \left( \frac{-16 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{64512 c^{10} x^{11} + 354816 bc^9 x^{10} + 98560 (9b^2 - 16)c^8 x^9 + 443520 (3b^3 - 16b)c^7 x^8 + 63360 (21b^4 - 224b^2 + 256)c^6 x^7 + 14784 (63b^5 - 1120b^3 + 3840b)c^5 x^6 + 22176 (21b^6 - 560b^4 + 3840b^2 - 4096)c^4 x^5 + 55440 (3b^7 - 112b^5 + 1280b^3 - 4096b)c^3 x^4 + 4620 (9b^8 - 448b^6 + 7680b^4 - 49152b^2 + 65536)c^2 x^3 + 6930 (b^9 - 64b^7 + 1536b^5 - 16384b^3 + 65536b)c x^2 + 693 (b^{10} - 80b^8 + 2560b^6 - 40960b^4 + 327680b^2 - 1048576)x}{c^5}$$

[In] integrate((1/4\*(b^2-16)/c+b\*x+c\*x^2)^5,x, algorithm="fricas")

[Out] 1/709632\*(64512\*c^10\*x^11 + 354816\*b\*c^9\*x^10 + 98560\*(9\*b^2 - 16)\*c^8\*x^9 + 443520\*(3\*b^3 - 16\*b)\*c^7\*x^8 + 63360\*(21\*b^4 - 224\*b^2 + 256)\*c^6\*x^7 + 14784\*(63\*b^5 - 1120\*b^3 + 3840\*b)\*c^5\*x^6 + 22176\*(21\*b^6 - 560\*b^4 + 3840\*b^2 - 4096)\*c^4\*x^5 + 55440\*(3\*b^7 - 112\*b^5 + 1280\*b^3 - 4096\*b)\*c^3\*x^4 + 4620\*(9\*b^8 - 448\*b^6 + 7680\*b^4 - 49152\*b^2 + 65536)\*c^2\*x^3 + 6930\*(b^9 - 64\*b^7 + 1536\*b^5 - 16384\*b^3 + 65536\*b)\*c\*x^2 + 693\*(b^10 - 80\*b^8 + 2560\*b^6 - 40960\*b^4 + 327680\*b^2 - 1048576)\*x)/c^5

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(97) = 194.

Time = 0.08 (sec) , antiderivative size = 248, normalized size of antiderivative = 2.28

$$\int \left( \frac{-16 + b^2}{4c} + bx + cx^2 \right)^5 dx$$

$$= \frac{bc^4 x^{10}}{2} + \frac{c^5 x^{11}}{11} + x^9 \cdot \left( \frac{5b^2 c^3}{4} - \frac{20c^3}{9} \right) + x^8 \cdot \left( \frac{15b^3 c^2}{8} - 10bc^2 \right)$$

$$+ x^7 \cdot \left( \frac{15b^4 c}{8} - 20b^2 c + \frac{160c}{7} \right) + x^6 \cdot \left( \frac{21b^5}{16} - \frac{70b^3}{3} + 80b \right)$$

$$+ \frac{x^5 \cdot (21b^6 - 560b^4 + 3840b^2 - 4096)}{32c} + \frac{x^4 \cdot (15b^7 - 560b^5 + 6400b^3 - 20480b)}{64c^2}$$

$$+ \frac{x^3 \cdot (45b^8 - 2240b^6 + 38400b^4 - 245760b^2 + 327680)}{768c^3}$$

$$+ \frac{x^2 \cdot (5b^9 - 320b^7 + 7680b^5 - 81920b^3 + 327680b)}{512c^4}$$

$$+ \frac{x(b^{10} - 80b^8 + 2560b^6 - 40960b^4 + 327680b^2 - 1048576)}{1024c^5}$$

[In] integrate((1/4\*(b\*\*2-16)/c+b\*x+c\*x\*\*2)\*\*5,x)

[Out] b\*c\*\*4\*x\*\*10/2 + c\*\*5\*x\*\*11/11 + x\*\*9\*(5\*b\*\*2\*c\*\*3/4 - 20\*c\*\*3/9) + x\*\*8\*(15\*b\*\*3\*c\*\*2/8 - 10\*b\*c\*\*2) + x\*\*7\*(15\*b\*\*4\*c/8 - 20\*b\*\*2\*c + 160\*c/7) + x\*\*

$6*(21*b^{**5}/16 - 70*b^{**3}/3 + 80*b) + x^{**5}*(21*b^{**6} - 560*b^{**4} + 3840*b^{**2} - 4096)/(32*c) + x^{**4}*(15*b^{**7} - 560*b^{**5} + 6400*b^{**3} - 20480*b)/(64*c^{**2}) + x^{**3}*(45*b^{**8} - 2240*b^{**6} + 38400*b^{**4} - 245760*b^{**2} + 327680)/(768*c^{**3}) + x^{**2}*(5*b^{**9} - 320*b^{**7} + 7680*b^{**5} - 81920*b^{**3} + 327680*b)/(512*c^{**4}) + x*(b^{**10} - 80*b^{**8} + 2560*b^{**6} - 40960*b^{**4} + 327680*b^{**2} - 1048576)/(1024*c^{**5})$

### Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(85) = 170.

Time = 0.19 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.15

$$\begin{aligned}
 & \int \left( \frac{-16 + b^2}{4c} + bx + cx^2 \right)^5 dx \\
 &= \frac{1}{11} c^5 x^{11} + \frac{1}{2} bc^4 x^{10} + \frac{10}{9} b^2 c^3 x^9 + \frac{5}{4} b^3 c^2 x^8 + \frac{5}{7} b^4 c x^7 + \frac{1}{6} b^5 x^6 \\
 &+ \frac{5(2cx^3 + 3bx^2)(b^2 - 16)^4}{1536c^4} + \frac{(6c^2x^5 + 15bcx^4 + 10b^2x^3)(b^2 - 16)^3}{192c^3} \\
 &+ \frac{(20c^3x^7 + 70bc^2x^6 + 84b^2cx^5 + 35b^3x^4)(b^2 - 16)^2}{224c^2} \\
 &+ \frac{(70c^4x^9 + 315bc^3x^8 + 540b^2c^2x^7 + 420b^3cx^6 + 126b^4x^5)(b^2 - 16)}{504c} + \frac{(b^2 - 16)^5 x}{1024c^5}
 \end{aligned}$$

[In] integrate((1/4\*(b^2-16)/c+b\*x+c\*x^2)^5,x, algorithm="maxima")

[Out] 1/11\*c^5\*x^11 + 1/2\*b\*c^4\*x^10 + 10/9\*b^2\*c^3\*x^9 + 5/4\*b^3\*c^2\*x^8 + 5/7\*b^4\*c\*x^7 + 1/6\*b^5\*x^6 + 5/1536\*(2\*c\*x^3 + 3\*b\*x^2)\*(b^2 - 16)^4/c^4 + 1/192\*(6\*c^2\*x^5 + 15\*b\*c\*x^4 + 10\*b^2\*x^3)\*(b^2 - 16)^3/c^3 + 1/224\*(20\*c^3\*x^7 + 70\*b\*c^2\*x^6 + 84\*b^2\*c\*x^5 + 35\*b^3\*x^4)\*(b^2 - 16)^2/c^2 + 1/504\*(70\*c^4\*x^9 + 315\*b\*c^3\*x^8 + 540\*b^2\*c^2\*x^7 + 420\*b^3\*c\*x^6 + 126\*b^4\*x^5)\*(b^2 - 16)/c + 1/1024\*(b^2 - 16)^5\*x/c^5

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 334 vs. 2(85) = 170.

Time = 0.28 (sec) , antiderivative size = 334, normalized size of antiderivative = 3.06

$$\begin{aligned}
 & \int \left( \frac{-16 + b^2}{4c} + bx + cx^2 \right)^5 dx \\
 &= \frac{64512c^{10}x^{11} + 354816bc^9x^{10} + 887040b^2c^8x^9 + 1330560b^3c^7x^8 + 1330560b^4c^6x^7 - 1576960c^8x^9 + 931392c^9x^8 - 1048576c^5x^6 + 1048576c^4x^5 - 1048576c^3x^4 + 1048576c^2x^3 - 1048576cx^2 + 1048576x}{1024c^5}
 \end{aligned}$$

[In] integrate((1/4\*(b^2-16)/c+b\*x+c\*x^2)^5,x, algorithm="giac")



```
[Out] 1/709632*(64512*c^10*x^11 + 354816*b*c^9*x^10 + 887040*b^2*c^8*x^9 + 133056
0*b^3*c^7*x^8 + 1330560*b^4*c^6*x^7 - 1576960*c^8*x^9 + 931392*b^5*c^5*x^6
- 7096320*b*c^7*x^8 + 465696*b^6*c^4*x^5 - 14192640*b^2*c^6*x^7 + 166320*b^
7*c^3*x^4 - 16558080*b^3*c^5*x^6 + 41580*b^8*c^2*x^3 - 12418560*b^4*c^4*x^5
+ 16220160*c^6*x^7 + 6930*b^9*c*x^2 - 6209280*b^5*c^3*x^4 + 56770560*b*c^5
*x^6 + 693*b^10*x - 2069760*b^6*c^2*x^3 + 85155840*b^2*c^4*x^5 - 443520*b^7
*c*x^2 + 70963200*b^3*c^3*x^4 - 55440*b^8*x + 35481600*b^4*c^2*x^3 - 908328
96*c^4*x^5 + 10644480*b^5*c*x^2 - 227082240*b*c^3*x^4 + 1774080*b^6*x - 227
082240*b^2*c^2*x^3 - 113541120*b^3*c*x^2 - 28385280*b^4*x + 302776320*c^2*x
^3 + 454164480*b*c*x^2 + 227082240*b^2*x - 726663168*x)/c^5
```

### Mupad [B] (verification not implemented)

Time = 9.23 (sec) , antiderivative size = 184, normalized size of antiderivative = 1.69

$$\int \left( \frac{-16 + b^2}{4c} + bx + cx^2 \right)^5 dx = \frac{c^5 x^{11}}{11} + \frac{x(b^2 - 16)^5}{1024 c^5} + \frac{b x^6 (63 b^4 - 1120 b^2 + 3840)}{48}$$

$$+ \frac{5 c x^7 (21 b^4 - 224 b^2 + 256)}{56}$$

$$+ \frac{b c^4 x^{10}}{2} + \frac{5 c^3 x^9 (9 b^2 - 16)}{36}$$

$$+ \frac{x^5 (21 b^6 - 560 b^4 + 3840 b^2 - 4096)}{32 c}$$

$$+ \frac{5 b c^2 x^8 (3 b^2 - 16)}{8} + \frac{5 b x^2 (b^2 - 16)^4}{512 c^4}$$

$$+ \frac{5 x^3 (b^2 - 16)^3 (9 b^2 - 16)}{768 c^3} + \frac{5 b x^4 (b^2 - 16)^2 (3 b^2 - 16)}{64 c^2}$$

```
[In] int((b*x + c*x^2 + (b^2/4 - 4)/c)^5,x)
```

```
[Out] (c^5*x^11)/11 + (x*(b^2 - 16)^5)/(1024*c^5) + (b*x^6*(63*b^4 - 1120*b^2 + 3
840))/48 + (5*c*x^7*(21*b^4 - 224*b^2 + 256))/56 + (b*c^4*x^10)/2 + (5*c^3*
x^9*(9*b^2 - 16))/36 + (x^5*(3840*b^2 - 560*b^4 + 21*b^6 - 4096))/(32*c) +
(5*b*c^2*x^8*(3*b^2 - 16))/8 + (5*b*x^2*(b^2 - 16)^4)/(512*c^4) + (5*x^3*(b
^2 - 16)^3*(9*b^2 - 16))/(768*c^3) + (5*b*x^4*(b^2 - 16)^2*(3*b^2 - 16))/(6
4*c^2)
```

### 3.78 $\int \frac{1}{2+4x+3x^2} dx$

Optimal result	418
Rubi [A] (verified)	418
Mathematica [A] (verified)	419
Maple [A] (verified)	419
Fricas [A] (verification not implemented)	420
Sympy [A] (verification not implemented)	420
Maxima [A] (verification not implemented)	420
Giac [A] (verification not implemented)	420
Mupad [B] (verification not implemented)	421

#### Optimal result

Integrand size = 12, antiderivative size = 18

$$\int \frac{1}{2+4x+3x^2} dx = \frac{\arctan\left(\frac{2+3x}{\sqrt{2}}\right)}{\sqrt{2}}$$

[Out] 1/2\*arctan(1/2\*(2+3\*x)\*2^(1/2))\*2^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 210}

$$\int \frac{1}{2+4x+3x^2} dx = \frac{\arctan\left(\frac{3x+2}{\sqrt{2}}\right)}{\sqrt{2}}$$

[In] Int[(2 + 4\*x + 3\*x^2)^(-1), x]

[Out] ArcTan[(2 + 3\*x)/Sqrt[2]]/Sqrt[2]

#### Rule 210

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(-(Rt[-a, 2]\*Rt[-b, 2])^(-1))\*ArcTan[Rt[-b, 2]\*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

#### Rule 632

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-1), x\_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4\*a\*c - x^2, x], x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c},

x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-8 - x^2} dx, x, 4 + 6x\right)\right) \\ &= \frac{\tan^{-1}\left(\frac{2+3x}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{\arctan\left(\frac{2+3x}{\sqrt{2}}\right)}{\sqrt{2}}$$

[In] Integrate[(2 + 4\*x + 3\*x^2)^(-1),x]

[Out] ArcTan[(2 + 3\*x)/Sqrt[2]]/Sqrt[2]

**Maple [A] (verified)**

Time = 4.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{\sqrt{2} \arctan\left(\frac{(4+6x)\sqrt{2}}{4}\right)}{2}$	17
risch	$\frac{\arctan\left(\frac{(2+3x)\sqrt{2}}{2}\right)\sqrt{2}}{2}$	17

[In] int(1/(3\*x^2+4\*x+2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*2^(1/2)\*arctan(1/4\*(4+6\*x)\*2^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.74 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (3x + 2) \right)$$

[In] integrate(1/(3\*x^2+4\*x+2),x, algorithm="fricas")

[Out] 1/2\*sqrt(2)\*arctan(1/2\*sqrt(2)\*(3\*x + 2))

**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{\sqrt{2} \operatorname{atan} \left( \frac{3\sqrt{2}x}{2} + \sqrt{2} \right)}{2}$$

[In] integrate(1/(3\*x\*\*2+4\*x+2),x)

[Out] sqrt(2)\*atan(3\*sqrt(2)\*x/2 + sqrt(2))/2

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (3x + 2) \right)$$

[In] integrate(1/(3\*x^2+4\*x+2),x, algorithm="maxima")

[Out] 1/2\*sqrt(2)\*arctan(1/2\*sqrt(2)\*(3\*x + 2))

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{1}{2} \sqrt{2} \arctan \left( \frac{1}{2} \sqrt{2} (3x + 2) \right)$$

[In] integrate(1/(3\*x^2+4\*x+2),x, algorithm="giac")

[Out] 1/2\*sqrt(2)\*arctan(1/2\*sqrt(2)\*(3\*x + 2))

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1}{2 + 4x + 3x^2} dx = \frac{\sqrt{2} \operatorname{atan}\left(\frac{\sqrt{2}(3x+2)}{2}\right)}{2}$$

[In] int(1/(4\*x + 3\*x^2 + 2),x)

[Out] (2^(1/2)\*atan((2^(1/2)\*(3\*x + 2))/2))/2

### 3.79 $\int \frac{1}{4-2\sqrt{3}x+x^2} dx$

Optimal result	422
Rubi [A] (verified)	422
Mathematica [A] (verified)	423
Maple [A] (verified)	423
Fricas [A] (verification not implemented)	423
Sympy [A] (verification not implemented)	424
Maxima [A] (verification not implemented)	424
Giac [A] (verification not implemented)	424
Mupad [B] (verification not implemented)	424

#### Optimal result

Integrand size = 15, antiderivative size = 12

$$\int \frac{1}{4-2\sqrt{3}x+x^2} dx = -\arctan(\sqrt{3}-x)$$

[Out]  $\arctan(x-3^{1/2})$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$ , Rules used = {632, 210}

$$\int \frac{1}{4-2\sqrt{3}x+x^2} dx = -\arctan(\sqrt{3}-x)$$

[In]  $\text{Int}[(4 - 2*\text{Sqrt}[3]*x + x^2)^{-1}, x]$

[Out]  $-\text{ArcTan}[\text{Sqrt}[3] - x]$

#### Rule 210

$\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 632

$\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-4-x^2} dx, x, -2\sqrt{3}+2x\right)\right) \\ &= -\tan^{-1}\left(\sqrt{3}-x\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{4-2\sqrt{3}x+x^2} dx = -\arctan\left(\sqrt{3}-x\right)$$

[In] Integrate[(4 - 2\*Sqrt[3]\*x + x^2)^(-1),x]

[Out] -ArcTan[Sqrt[3] - x]

**Maple [A] (verified)**

Time = 3.56 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.75

method	result	size
default	$\arctan(x - \sqrt{3})$	9
risch	$\arctan(x - \sqrt{3})$	9
parallelrisch	$\frac{i \ln(x+i-\sqrt{3})}{2} - \frac{i \ln(x-\sqrt{3}-i)}{2}$	28

[In] int(1/(4+x^2-2\*3^(1/2)\*x),x,method=\_RETURNVERBOSE)

[Out] arctan(x-3^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{4-2\sqrt{3}x+x^2} dx = -\arctan\left(-x+\sqrt{3}\right)$$

[In] integrate(1/(4+x^2-2\*x\*3^(1/2)),x, algorithm="fricas")

[Out] -arctan(-x + sqrt(3))

**Sympy [A] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.58

$$\int \frac{1}{4 - 2\sqrt{3}x + x^2} dx = \operatorname{atan}\left(x - \sqrt{3}\right)$$

[In] integrate(1/(4+x\*\*2-2\*x\*3\*\*(1/2)),x)

[Out] atan(x - sqrt(3))

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{4 - 2\sqrt{3}x + x^2} dx = \operatorname{arctan}\left(x - \sqrt{3}\right)$$

[In] integrate(1/(4+x^2-2\*x\*3^(1/2)),x, algorithm="maxima")

[Out] arctan(x - sqrt(3))

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{4 - 2\sqrt{3}x + x^2} dx = \operatorname{arctan}\left(x - \sqrt{3}\right)$$

[In] integrate(1/(4+x^2-2\*x\*3^(1/2)),x, algorithm="giac")

[Out] arctan(x - sqrt(3))

**Mupad [B] (verification not implemented)**

Time = 9.30 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.67

$$\int \frac{1}{4 - 2\sqrt{3}x + x^2} dx = \operatorname{atan}\left(x - \sqrt{3}\right)$$

[In] int(1/(x^2 - 2\*3^(1/2)\*x + 4),x)

[Out] atan(x - 3^(1/2))



### 3.80 $\int \frac{1}{2+4x-3x^2} dx$

Optimal result	425
Rubi [A] (verified)	425
Mathematica [A] (verified)	426
Maple [A] (verified)	426
Fricas [B] (verification not implemented)	427
Sympy [A] (verification not implemented)	427
Maxima [A] (verification not implemented)	427
Giac [A] (verification not implemented)	428
Mupad [B] (verification not implemented)	428

#### Optimal result

Integrand size = 12, antiderivative size = 19

$$\int \frac{1}{2+4x-3x^2} dx = -\frac{\operatorname{arctanh}\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{10}}$$

[Out]  $-1/10*\operatorname{arctanh}(1/10*(2-3*x)*10^{(1/2)})*10^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 212}

$$\int \frac{1}{2+4x-3x^2} dx = -\frac{\operatorname{arctanh}\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{10}}$$

[In]  $\operatorname{Int}[(2 + 4*x - 3*x^2)^{-1}, x]$

[Out]  $-(\operatorname{ArcTanh}[(2 - 3*x)/\operatorname{Sqrt}[10]]/\operatorname{Sqrt}[10])$

#### Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

#### Rule 632

$\operatorname{Int}[(a_ + (b_)*(x_ + (c_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Dist}[-2, \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /;$   $\operatorname{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{40-x^2} dx, x, 4-6x\right)\right) \\ &= -\frac{\tanh^{-1}\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{10}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.79

$$\int \frac{1}{2+4x-3x^2} dx = \frac{-\log(2+\sqrt{10}-3x) + \log(-2+\sqrt{10}+3x)}{2\sqrt{10}}$$

[In] `Integrate[(2 + 4*x - 3*x^2)^(-1), x]`

[Out] `(-Log[2 + Sqrt[10] - 3*x] + Log[-2 + Sqrt[10] + 3*x])/(2*Sqrt[10])`

**Maple [A] (verified)**

Time = 2.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\sqrt{10} \operatorname{arctanh}\left(\frac{(-4+6x)\sqrt{10}}{20}\right)}{10}$	17
risch	$\frac{\sqrt{10} \ln(3x-2+\sqrt{10})}{20} - \frac{\sqrt{10} \ln(3x-2-\sqrt{10})}{20}$	32

[In] `int(1/(-3*x^2+4*x+2), x, method=_RETURNVERBOSE)`

[Out] `1/10*10^(1/2)*arctanh(1/20*(-4+6*x)*10^(1/2))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 39 vs.  $2(16) = 32$ .

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.05

$$\int \frac{1}{2+4x-3x^2} dx = \frac{1}{20} \sqrt{10} \log \left( \frac{9x^2 + 2\sqrt{10}(3x-2) - 12x + 14}{3x^2 - 4x - 2} \right)$$

[In] integrate(1/(-3\*x^2+4\*x+2),x, algorithm="fricas")

[Out] 1/20\*sqrt(10)\*log((9\*x^2 + 2\*sqrt(10)\*(3\*x - 2) - 12\*x + 14)/(3\*x^2 - 4\*x - 2))

**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.05

$$\int \frac{1}{2+4x-3x^2} dx = \frac{\sqrt{10} \log \left( x - \frac{2}{3} + \frac{\sqrt{10}}{3} \right)}{20} - \frac{\sqrt{10} \log \left( x - \frac{\sqrt{10}}{3} - \frac{2}{3} \right)}{20}$$

[In] integrate(1/(-3\*x\*\*2+4\*x+2),x)

[Out] sqrt(10)\*log(x - 2/3 + sqrt(10)/3)/20 - sqrt(10)\*log(x - sqrt(10)/3 - 2/3)/20

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

$$\int \frac{1}{2+4x-3x^2} dx = -\frac{1}{20} \sqrt{10} \log \left( \frac{3x - \sqrt{10} - 2}{3x + \sqrt{10} - 2} \right)$$

[In] integrate(1/(-3\*x^2+4\*x+2),x, algorithm="maxima")

[Out] -1/20\*sqrt(10)\*log((3\*x - sqrt(10) - 2)/(3\*x + sqrt(10) - 2))

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.63

$$\int \frac{1}{2 + 4x - 3x^2} dx = -\frac{1}{20} \sqrt{10} \log \left( \frac{|6x - 2\sqrt{10} - 4|}{|6x + 2\sqrt{10} - 4|} \right)$$

[In] integrate(1/(-3\*x^2+4\*x+2),x, algorithm="giac")

[Out] -1/20\*sqrt(10)\*log(abs(6\*x - 2\*sqrt(10) - 4)/abs(6\*x + 2\*sqrt(10) - 4))

**Mupad [B] (verification not implemented)**

Time = 0.24 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{2 + 4x - 3x^2} dx = \frac{\sqrt{10} \operatorname{atanh}\left(\sqrt{10} \left(\frac{3x}{10} - \frac{1}{5}\right)\right)}{10}$$

[In] int(1/(4\*x - 3\*x^2 + 2),x)

[Out] (10^(1/2)\*atanh(10^(1/2)\*((3\*x)/10 - 1/5)))/10

### 3.81 $\int \frac{1}{2+5x+3x^2} dx$

Optimal result	429
Rubi [A] (verified)	429
Mathematica [A] (verified)	430
Maple [A] (verified)	430
Fricas [A] (verification not implemented)	430
Sympy [A] (verification not implemented)	431
Maxima [A] (verification not implemented)	431
Giac [A] (verification not implemented)	431
Mupad [B] (verification not implemented)	431

#### Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \frac{1}{2+5x+3x^2} dx = -\log(1+x) + \log(2+3x)$$

[Out]  $-\ln(1+x)+\ln(2+3*x)$

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {630, 31}

$$\int \frac{1}{2+5x+3x^2} dx = \log(3x+2) - \log(x+1)$$

[In]  $\text{Int}[(2+5*x+3*x^2)^{-1},x]$

[Out]  $-\text{Log}[1+x] + \text{Log}[2+3*x]$

#### Rule 31

$\text{Int}[(a_+ + (b_+)(x_+))^{-1}, x\_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}\{a, b, x\}$

#### Rule 630

$\text{Int}[(a_+ + (b_+)(x_+) + (c_+)(x_+)^2)^{-1}, x\_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 - q/2 + c*x, x], x], x] - \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 + q/2 + c*x, x], x], x] /; \text{FreeQ}\{a, b, c, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{PosQ}[b^2 - 4*a*c] \&\& \text{PerfectSquareQ}[b^2 - 4*a*c]$

Rubi steps

$$\begin{aligned} \text{integral} &= 3 \int \frac{1}{2+3x} dx - 3 \int \frac{1}{3+3x} dx \\ &= -\log(1+x) + \log(2+3x) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{2+5x+3x^2} dx = -\log(1+x) + \log(2+3x)$$

[In] Integrate[(2 + 5\*x + 3\*x^2)^(-1),x]

[Out] -Log[1 + x] + Log[2 + 3\*x]

**Maple [A] (verified)**

Time = 2.10 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
parallelsch	$-\ln(1+x) + \ln\left(\frac{2}{3} + x\right)$	12
default	$-\ln(1+x) + \ln(2+3x)$	14
norman	$-\ln(1+x) + \ln(2+3x)$	14
risch	$-\ln(1+x) + \ln(2+3x)$	14

[In] int(1/(3\*x^2+5\*x+2),x,method=\_RETURNVERBOSE)

[Out] -ln(1+x)+ln(2/3+x)

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{2+5x+3x^2} dx = \log(3x+2) - \log(x+1)$$

[In] integrate(1/(3\*x^2+5\*x+2),x, algorithm="fricas")

[Out] log(3\*x + 2) - log(x + 1)

**Sympy [A] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.77

$$\int \frac{1}{2 + 5x + 3x^2} dx = \log\left(x + \frac{2}{3}\right) - \log(x + 1)$$

[In] integrate(1/(3\*x\*\*2+5\*x+2),x)

[Out] log(x + 2/3) - log(x + 1)

**Maxima [A] (verification not implemented)**

none

Time = 0.18 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{1}{2 + 5x + 3x^2} dx = \log(3x + 2) - \log(x + 1)$$

[In] integrate(1/(3\*x^2+5\*x+2),x, algorithm="maxima")

[Out] log(3\*x + 2) - log(x + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{1}{2 + 5x + 3x^2} dx = \log(|3x + 2|) - \log(|x + 1|)$$

[In] integrate(1/(3\*x^2+5\*x+2),x, algorithm="giac")

[Out] log(abs(3\*x + 2)) - log(abs(x + 1))

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.62

$$\int \frac{1}{2 + 5x + 3x^2} dx = -2 \operatorname{atanh}(6x + 5)$$

[In] int(1/(5\*x + 3\*x^2 + 2),x)

[Out] -2\*atanh(6\*x + 5)

### 3.82 $\int \frac{1}{2+5x-3x^2} dx$

Optimal result	432
Rubi [A] (verified)	432
Mathematica [A] (verified)	433
Maple [A] (verified)	433
Fricas [A] (verification not implemented)	433
Sympy [A] (verification not implemented)	434
Maxima [A] (verification not implemented)	434
Giac [A] (verification not implemented)	434
Mupad [B] (verification not implemented)	434

#### Optimal result

Integrand size = 12, antiderivative size = 21

$$\int \frac{1}{2+5x-3x^2} dx = -\frac{1}{7} \log(2-x) + \frac{1}{7} \log(1+3x)$$

[Out] -1/7\*ln(2-x)+1/7\*ln(1+3\*x)

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {630, 31}

$$\int \frac{1}{2+5x-3x^2} dx = \frac{1}{7} \log(3x+1) - \frac{1}{7} \log(2-x)$$

[In] Int[(2 + 5\*x - 3\*x^2)^(-1), x]

[Out] -1/7\*Log[2 - x] + Log[1 + 3\*x]/7

#### Rule 31

Int[((a\_) + (b\_.)\*(x\_))^(n\_), x\_Symbol] := Simp[Log[RemoveContent[a + b\*x, x]]/b, x] /; FreeQ[{a, b}, x]

#### Rule 630

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(n\_), x\_Symbol] := With[{q = Rt[b^2 - 4\*a\*c, 2]}, Dist[c/q, Int[1/Simp[b/2 - q/2 + c\*x, x], x] - Dist[c/q, Int[1/Simp[b/2 + q/2 + c\*x, x], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && PosQ[b^2 - 4\*a\*c] && PerfectSquareQ[b^2 - 4\*a\*c]



Rubi steps

$$\begin{aligned} \text{integral} &= -\left(\frac{3}{7} \int \frac{1}{-1-3x} dx\right) + \frac{3}{7} \int \frac{1}{6-3x} dx \\ &= -\frac{1}{7} \log(2-x) + \frac{1}{7} \log(1+3x) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{1}{2+5x-3x^2} dx = -\frac{1}{7} \log(2-x) + \frac{1}{7} \log(1+3x)$$

[In] Integrate[(2 + 5\*x - 3\*x^2)^(-1),x]

[Out] -1/7\*Log[2 - x] + Log[1 + 3\*x]/7

### Maple [A] (verified)

Time = 2.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
parallelrisc	$-\frac{\ln(-2+x)}{7} + \frac{\ln(x+\frac{1}{3})}{7}$	14
default	$-\frac{\ln(-2+x)}{7} + \frac{\ln(3x+1)}{7}$	16
norman	$-\frac{\ln(-2+x)}{7} + \frac{\ln(3x+1)}{7}$	16
risc	$-\frac{\ln(-2+x)}{7} + \frac{\ln(3x+1)}{7}$	16

[In] int(1/(-3\*x^2+5\*x+2),x,method=\_RETURNVERBOSE)

[Out] -1/7\*ln(-2+x)+1/7\*ln(x+1/3)

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1}{2+5x-3x^2} dx = \frac{1}{7} \log(3x+1) - \frac{1}{7} \log(x-2)$$

[In] integrate(1/(-3\*x^2+5\*x+2),x, algorithm="fricas")

[Out] 1/7\*log(3\*x + 1) - 1/7\*log(x - 2)

**Sympy [A] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \frac{1}{2 + 5x - 3x^2} dx = -\frac{\log(x - 2)}{7} + \frac{\log(x + \frac{1}{3})}{7}$$

[In] integrate(1/(-3\*x\*\*2+5\*x+2),x)

[Out] -log(x - 2)/7 + log(x + 1/3)/7

**Maxima [A] (verification not implemented)**

none

Time = 0.18 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.71

$$\int \frac{1}{2 + 5x - 3x^2} dx = \frac{1}{7} \log(3x + 1) - \frac{1}{7} \log(x - 2)$$

[In] integrate(1/(-3\*x^2+5\*x+2),x, algorithm="maxima")

[Out] 1/7\*log(3\*x + 1) - 1/7\*log(x - 2)

**Giac [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.81

$$\int \frac{1}{2 + 5x - 3x^2} dx = \frac{1}{7} \log(|3x + 1|) - \frac{1}{7} \log(|x - 2|)$$

[In] integrate(1/(-3\*x^2+5\*x+2),x, algorithm="giac")

[Out] 1/7\*log(abs(3\*x + 1)) - 1/7\*log(abs(x - 2))

**Mupad [B] (verification not implemented)**

Time = 0.10 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.38

$$\int \frac{1}{2 + 5x - 3x^2} dx = \frac{2 \operatorname{atanh}\left(\frac{6x}{7} - \frac{5}{7}\right)}{7}$$

[In] int(1/(5\*x - 3\*x^2 + 2),x)

[Out] (2\*atanh((6\*x)/7 - 5/7))/7

### 3.83 $\int \frac{1}{3+4x+x^2} dx$

Optimal result	435
Rubi [B] (verified)	435
Mathematica [B] (verified)	436
Maple [B] (verified)	436
Fricas [B] (verification not implemented)	437
Sympy [B] (verification not implemented)	437
Maxima [B] (verification not implemented)	437
Giac [B] (verification not implemented)	438
Mupad [B] (verification not implemented)	438

#### Optimal result

Integrand size = 10, antiderivative size = 6

$$\int \frac{1}{3+4x+x^2} dx = -\operatorname{arctanh}(2+x)$$

[Out]  $-\operatorname{arctanh}(2+x)$

#### Rubi [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 17 vs.  $2(6) = 12$ .

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.83, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$ , Rules used = {630, 31}

$$\int \frac{1}{3+4x+x^2} dx = \frac{1}{2} \log(x+1) - \frac{1}{2} \log(x+3)$$

[In]  $\text{Int}[(3 + 4*x + x^2)^{-1}, x]$

[Out]  $\text{Log}[1 + x]/2 - \text{Log}[3 + x]/2$

#### Rule 31

$\text{Int}[(a + (b \cdot x)^{-1}), x\_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]/b, x] /; \text{FreeQ}\{a, b, x\}$

#### Rule 630

$\text{Int}[(a + (b \cdot x) + (c \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \text{With}\{q = \text{Rt}[b^2 - 4 \cdot a \cdot c, 2]\}, \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 - q/2 + c \cdot x, x], x], x] - \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 + q/2 + c \cdot x, x], x], x] /; \text{FreeQ}\{a, b, c, x\} \&\& \text{NeQ}[b^2$

- 4\*a\*c, 0] && PosQ[b^2 - 4\*a\*c] && PerfectSquareQ[b^2 - 4\*a\*c]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} \int \frac{1}{1+x} dx - \frac{1}{2} \int \frac{1}{3+x} dx \\ &= \frac{1}{2} \log(1+x) - \frac{1}{2} \log(3+x) \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 17 vs. 2(6) = 12.

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.83

$$\int \frac{1}{3+4x+x^2} dx = \frac{1}{2} \log(1+x) - \frac{1}{2} \log(3+x)$$

[In] Integrate[(3 + 4\*x + x^2)^(-1),x]

[Out] Log[1 + x]/2 - Log[3 + x]/2

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 13 vs. 2(6) = 12.

Time = 2.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 2.33

method	result	size
default	$-\frac{\ln(3+x)}{2} + \frac{\ln(1+x)}{2}$	14
norman	$-\frac{\ln(3+x)}{2} + \frac{\ln(1+x)}{2}$	14
risch	$-\frac{\ln(3+x)}{2} + \frac{\ln(1+x)}{2}$	14
parallelrisk	$-\frac{\ln(3+x)}{2} + \frac{\ln(1+x)}{2}$	14

[In] int(1/(x^2+4\*x+3),x,method=\_RETURNVERBOSE)

[Out] -1/2\*ln(3+x)+1/2\*ln(1+x)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 13 vs.  $2(6) = 12$ .

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 2.17

$$\int \frac{1}{3 + 4x + x^2} dx = -\frac{1}{2} \log(x + 3) + \frac{1}{2} \log(x + 1)$$

[In] integrate(1/(x^2+4\*x+3),x, algorithm="fricas")

[Out] -1/2\*log(x + 3) + 1/2\*log(x + 1)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 12 vs.  $2(5) = 10$ .

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 2.00

$$\int \frac{1}{3 + 4x + x^2} dx = \frac{\log(x + 1)}{2} - \frac{\log(x + 3)}{2}$$

[In] integrate(1/(x\*\*2+4\*x+3),x)

[Out] log(x + 1)/2 - log(x + 3)/2

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 13 vs.  $2(6) = 12$ .

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 2.17

$$\int \frac{1}{3 + 4x + x^2} dx = -\frac{1}{2} \log(x + 3) + \frac{1}{2} \log(x + 1)$$

[In] integrate(1/(x^2+4\*x+3),x, algorithm="maxima")

[Out] -1/2\*log(x + 3) + 1/2\*log(x + 1)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 15 vs.  $2(6) = 12$ .

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 2.50

$$\int \frac{1}{3 + 4x + x^2} dx = -\frac{1}{2} \log(|x + 3|) + \frac{1}{2} \log(|x + 1|)$$

[In] `integrate(1/(x^2+4*x+3),x, algorithm="giac")`

[Out] `-1/2*log(abs(x + 3)) + 1/2*log(abs(x + 1))`

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{1}{3 + 4x + x^2} dx = -\operatorname{atanh}(x + 2)$$

[In] `int(1/(4*x + x^2 + 3),x)`

[Out] `-atanh(x + 2)`

### 3.84 $\int \frac{1}{1+\pi x+2x^2} dx$

Optimal result	439
Rubi [A] (verified)	439
Mathematica [A] (verified)	440
Maple [A] (verified)	440
Fricas [B] (verification not implemented)	441
Sympy [B] (verification not implemented)	441
Maxima [A] (verification not implemented)	441
Giac [A] (verification not implemented)	442
Mupad [B] (verification not implemented)	442

#### Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{1}{1+\pi x+2x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{\pi+4x}{\sqrt{-8+\pi^2}}\right)}{\sqrt{-8+\pi^2}}$$

[Out]  $-2*\operatorname{arctanh}((\text{Pi}+4*x)/(\text{Pi}^2-8)^{(1/2)})/(\text{Pi}^2-8)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 212}

$$\int \frac{1}{1+\pi x+2x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{4x+\pi}{\sqrt{\pi^2-8}}\right)}{\sqrt{\pi^2-8}}$$

[In]  $\text{Int}[(1 + \text{Pi}*x + 2*x^2)^{-1}, x]$

[Out]  $(-2*\text{ArcTanh}[(\text{Pi} + 4*x)/\text{Sqrt}[-8 + \text{Pi}^2]])/\text{Sqrt}[-8 + \text{Pi}^2]$

#### Rule 212

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 632

$\text{Int}[(a_ + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-8 + \pi^2 - x^2} dx, x, \pi + 4x\right)\right) \\ &= -\frac{2 \tanh^{-1}\left(\frac{\pi+4x}{\sqrt{-8+\pi^2}}\right)}{\sqrt{-8 + \pi^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \pi x + 2x^2} dx = -\frac{2\text{arctanh}\left(\frac{\pi+4x}{\sqrt{-8+\pi^2}}\right)}{\sqrt{-8 + \pi^2}}$$

[In] `Integrate[(1 + Pi*x + 2*x^2)^(-1),x]`

[Out] `(-2*ArcTanh[(Pi + 4*x)/Sqrt[-8 + Pi^2]])/Sqrt[-8 + Pi^2]`

**Maple [A] (verified)**

Time = 1.98 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
default	$-\frac{2 \operatorname{arctanh}\left(\frac{\pi+4x}{\sqrt{\pi^2-8}}\right)}{\sqrt{\pi^2-8}}$	24
risch	$\frac{\ln\left(-\pi^2+\pi\sqrt{\pi^2-8}+4x\sqrt{\pi^2-8}+8\right)}{\sqrt{\pi^2-8}} - \frac{\ln\left(\pi^2+\pi\sqrt{\pi^2-8}+4x\sqrt{\pi^2-8}-8\right)}{\sqrt{\pi^2-8}}$	71

[In] `int(1/(Pi*x+2*x^2+1),x,method=_RETURNVERBOSE)`

[Out] `-2*arctanh((Pi+4*x)/(Pi^2-8)^(1/2))/(Pi^2-8)^(1/2)`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(23) = 46.

Time = 0.27 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.85

$$\int \frac{1}{1 + \pi x + 2x^2} dx = \frac{\log\left(\frac{\pi^2 + 4\pi x + 8x^2 - (\pi + 4x)\sqrt{\pi^2 - 8} - 4}{\pi x + 2x^2 + 1}\right)}{\sqrt{\pi^2 - 8}}$$

[In] integrate(1/(pi\*x+2\*x^2+1),x, algorithm="fricas")

[Out] log((pi^2 + 4\*pi\*x + 8\*x^2 - (pi + 4\*x)\*sqrt(pi^2 - 8) - 4)/(pi\*x + 2\*x^2 + 1))/sqrt(pi^2 - 8)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(26) = 52.

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.81

$$\int \frac{1}{1 + \pi x + 2x^2} dx = \frac{\log\left(x - \frac{\pi^2}{4\sqrt{-8 + \pi^2}} + \frac{\pi}{4} + \frac{2}{\sqrt{-8 + \pi^2}}\right)}{\sqrt{-8 + \pi^2}} - \frac{\log\left(x - \frac{2}{\sqrt{-8 + \pi^2}} + \frac{\pi}{4} + \frac{\pi^2}{4\sqrt{-8 + \pi^2}}\right)}{\sqrt{-8 + \pi^2}}$$

[In] integrate(1/(pi\*x+2\*x\*\*2+1),x)

[Out] log(x - pi\*\*2/(4\*sqrt(-8 + pi\*\*2)) + pi/4 + 2/sqrt(-8 + pi\*\*2))/sqrt(-8 + pi\*\*2) - log(x - 2/sqrt(-8 + pi\*\*2) + pi/4 + pi\*\*2/(4\*sqrt(-8 + pi\*\*2)))/sqrt(-8 + pi\*\*2)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.41

$$\int \frac{1}{1 + \pi x + 2x^2} dx = \frac{\log\left(\frac{\pi + 4x - \sqrt{\pi^2 - 8}}{\pi + 4x + \sqrt{\pi^2 - 8}}\right)}{\sqrt{\pi^2 - 8}}$$

[In] integrate(1/(pi\*x+2\*x^2+1),x, algorithm="maxima")

[Out] log((pi + 4\*x - sqrt(pi^2 - 8))/(pi + 4\*x + sqrt(pi^2 - 8)))/sqrt(pi^2 - 8)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.48

$$\int \frac{1}{1 + \pi x + 2x^2} dx = \frac{\log\left(\frac{|\pi + 4x - \sqrt{\pi^2 - 8}|}{\pi + 4x + \sqrt{\pi^2 - 8}}\right)}{\sqrt{\pi^2 - 8}}$$

[In] integrate(1/(pi\*x+2\*x^2+1),x, algorithm="giac")

[Out] log(abs(pi + 4\*x - sqrt(pi^2 - 8))/abs(pi + 4\*x + sqrt(pi^2 - 8)))/sqrt(pi^2 - 8)

**Mupad [B] (verification not implemented)**

Time = 9.57 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{1}{1 + \pi x + 2x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{\pi + 4x}{\sqrt{\pi^2 - 8}}\right)}{\sqrt{\pi^2 - 8}}$$

[In] int(1/(Pi\*x + 2\*x^2 + 1),x)

[Out] -(2\*atanh((Pi + 4\*x)/(Pi^2 - 8)^(1/2)))/(Pi^2 - 8)^(1/2)

### 3.85 $\int \frac{1}{1+\pi x-2x^2} dx$

Optimal result	443
Rubi [A] (verified)	443
Mathematica [A] (verified)	444
Maple [A] (verified)	444
Fricas [B] (verification not implemented)	445
Sympy [B] (verification not implemented)	445
Maxima [A] (verification not implemented)	445
Giac [A] (verification not implemented)	446
Mupad [B] (verification not implemented)	446

#### Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{1}{1+\pi x-2x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{\pi-4x}{\sqrt{8+\pi^2}}\right)}{\sqrt{8+\pi^2}}$$

[Out]  $-2*\operatorname{arctanh}((\text{Pi}-4*x)/(\text{Pi}^2+8)^{(1/2)})/(\text{Pi}^2+8)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 212}

$$\int \frac{1}{1+\pi x-2x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{\pi-4x}{\sqrt{8+\pi^2}}\right)}{\sqrt{8+\pi^2}}$$

[In]  $\text{Int}[(1 + \text{Pi}*x - 2*x^2)^{-1}, x]$

[Out]  $(-2*\text{ArcTanh}[(\text{Pi} - 4*x)/\text{Sqrt}[8 + \text{Pi}^2]])/\text{Sqrt}[8 + \text{Pi}^2]$

#### Rule 212

$\text{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 632

$\text{Int}[(a_+ + (b_+)*(x_+) + (c_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{8 + \pi^2 - x^2} dx, x, \pi - 4x\right)\right) \\ &= -\frac{2 \tanh^{-1}\left(\frac{\pi-4x}{\sqrt{8+\pi^2}}\right)}{\sqrt{8 + \pi^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{1}{1 + \pi x - 2x^2} dx = \frac{2 \operatorname{arctanh}\left(\frac{-\pi+4x}{\sqrt{8+\pi^2}}\right)}{\sqrt{8 + \pi^2}}$$

[In] `Integrate[(1 + Pi*x - 2*x^2)^(-1),x]`

[Out] `(2*ArcTanh[(-Pi + 4*x)/Sqrt[8 + Pi^2]])/Sqrt[8 + Pi^2]`

**Maple [A] (verified)**

Time = 2.08 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{2 \operatorname{arctanh}\left(\frac{-\pi+4x}{\sqrt{\pi^2+8}}\right)}{\sqrt{\pi^2+8}}$	26
risch	$\frac{\ln\left(\pi^2 - \pi\sqrt{\pi^2+8} + 4x\sqrt{\pi^2+8} + 8\right)}{\sqrt{\pi^2+8}} - \frac{\ln\left(-\pi^2 - \pi\sqrt{\pi^2+8} + 4x\sqrt{\pi^2+8} - 8\right)}{\sqrt{\pi^2+8}}$	73

[In] `int(1/(Pi*x-2*x^2+1),x,method=_RETURNVERBOSE)`

[Out] `2/(Pi^2+8)^(1/2)*arctanh((-Pi+4*x)/(Pi^2+8)^(1/2))`

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(23) = 46.

Time = 0.33 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.89

$$\int \frac{1}{1 + \pi x - 2x^2} dx = \frac{\log\left(-\frac{\pi^2 - 4\pi x + 8x^2 - (\pi - 4x)\sqrt{\pi^2 + 8} + 4}{\pi x - 2x^2 + 1}\right)}{\sqrt{\pi^2 + 8}}$$

[In] integrate(1/(pi\*x-2\*x^2+1),x, algorithm="fricas")

[Out] log(-(pi^2 - 4\*pi\*x + 8\*x^2 - (pi - 4\*x)\*sqrt(pi^2 + 8) + 4)/(pi\*x - 2\*x^2 + 1))/sqrt(pi^2 + 8)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(26) = 52.

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.81

$$\int \frac{1}{1 + \pi x - 2x^2} dx = -\frac{\log\left(x - \frac{\pi}{4} - \frac{\pi^2}{4\sqrt{8+\pi^2}} - \frac{2}{\sqrt{8+\pi^2}}\right)}{\sqrt{8+\pi^2}} + \frac{\log\left(x - \frac{\pi}{4} + \frac{2}{\sqrt{8+\pi^2}} + \frac{\pi^2}{4\sqrt{8+\pi^2}}\right)}{\sqrt{8+\pi^2}}$$

[In] integrate(1/(pi\*x-2\*x\*\*2+1),x)

[Out] -log(x - pi/4 - pi\*\*2/(4\*sqrt(8 + pi\*\*2)) - 2/sqrt(8 + pi\*\*2))/sqrt(8 + pi\*\*2) + log(x - pi/4 + 2/sqrt(8 + pi\*\*2) + pi\*\*2/(4\*sqrt(8 + pi\*\*2)))/sqrt(8 + pi\*\*2)

**Maxima [A] (verification not implemented)**

none

Time = 0.18 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.44

$$\int \frac{1}{1 + \pi x - 2x^2} dx = -\frac{\log\left(\frac{\pi - 4x + \sqrt{\pi^2 + 8}}{\pi - 4x - \sqrt{\pi^2 + 8}}\right)}{\sqrt{\pi^2 + 8}}$$

[In] integrate(1/(pi\*x-2\*x^2+1),x, algorithm="maxima")

[Out] -log((pi - 4\*x + sqrt(pi^2 + 8))/(pi - 4\*x - sqrt(pi^2 + 8)))/sqrt(pi^2 + 8)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int \frac{1}{1 + \pi x - 2x^2} dx = -\frac{\log\left(\frac{|\pi - 4x - \sqrt{\pi^2 + 8}|}{|\pi - 4x + \sqrt{\pi^2 + 8}|}\right)}{\sqrt{\pi^2 + 8}}$$

[In] integrate(1/(pi\*x-2\*x^2+1),x, algorithm="giac")

[Out] -log(abs(-pi + 4\*x - sqrt(pi^2 + 8))/abs(-pi + 4\*x + sqrt(pi^2 + 8)))/sqrt(pi^2 + 8)

**Mupad [B] (verification not implemented)**

Time = 9.22 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{1}{1 + \pi x - 2x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{\pi - 4x}{\sqrt{\pi^2 + 8}}\right)}{\sqrt{\pi^2 + 8}}$$

[In] int(1/(Pi\*x - 2\*x^2 + 1),x)

[Out] -(2\*atanh((Pi - 4\*x)/(Pi^2 + 8)^(1/2)))/(Pi^2 + 8)^(1/2)

### 3.86 $\int \frac{1}{1+\pi x+3x^2} dx$

Optimal result	447
Rubi [A] (verified)	447
Mathematica [A] (verified)	448
Maple [A] (verified)	448
Fricas [A] (verification not implemented)	449
Sympy [C] (verification not implemented)	449
Maxima [A] (verification not implemented)	449
Giac [A] (verification not implemented)	450
Mupad [B] (verification not implemented)	450

#### Optimal result

Integrand size = 12, antiderivative size = 31

$$\int \frac{1}{1+\pi x+3x^2} dx = \frac{2 \arctan\left(\frac{\pi+6x}{\sqrt{12-\pi^2}}\right)}{\sqrt{12-\pi^2}}$$

[Out]  $2*\arctan((\text{Pi}+6*x)/(-\text{Pi}^2+12)^{(1/2)})/(-\text{Pi}^2+12)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 210}

$$\int \frac{1}{1+\pi x+3x^2} dx = \frac{2 \arctan\left(\frac{6x+\pi}{\sqrt{12-\pi^2}}\right)}{\sqrt{12-\pi^2}}$$

[In]  $\text{Int}[(1 + \text{Pi}*x + 3*x^2)^{-1}, x]$

[Out]  $(2*\text{ArcTan}[(\text{Pi} + 6*x)/\text{Sqrt}[12 - \text{Pi}^2]])/\text{Sqrt}[12 - \text{Pi}^2]$

#### Rule 210

$\text{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}\{a/b\} \ \& \ (\text{LtQ}\{a, 0\} \ || \ \text{LtQ}\{b, 0\})$

#### Rule 632

$\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-12 + \pi^2 - x^2} dx, x, \pi + 6x\right)\right) \\ &= \frac{2 \tan^{-1}\left(\frac{\pi+6x}{\sqrt{12-\pi^2}}\right)}{\sqrt{12-\pi^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \pi x + 3x^2} dx = \frac{2 \arctan\left(\frac{\pi+6x}{\sqrt{12-\pi^2}}\right)}{\sqrt{12-\pi^2}}$$

[In] `Integrate[(1 + Pi*x + 3*x^2)^(-1),x]`

[Out] `(2*ArcTan[(Pi + 6*x)/Sqrt[12 - Pi^2]])/Sqrt[12 - Pi^2]`

**Maple [A] (verified)**

Time = 3.00 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.90

method	result	size
default	$\frac{2 \arctan\left(\frac{\pi+6x}{\sqrt{-\pi^2+12}}\right)}{\sqrt{-\pi^2+12}}$	28
risch	$\frac{\ln\left(-\pi^2+\pi\sqrt{\pi^2-12}+6x\sqrt{\pi^2-12}+12\right)}{\sqrt{\pi^2-12}} - \frac{\ln\left(\pi^2+\pi\sqrt{\pi^2-12}+6x\sqrt{\pi^2-12}-12\right)}{\sqrt{\pi^2-12}}$	71

[In] `int(1/(Pi*x+3*x^2+1),x,method=_RETURNVERBOSE)`

[Out] `2*arctan((Pi+6*x)/(-Pi^2+12)^(1/2))/(-Pi^2+12)^(1/2)`



**Fricas [A] (verification not implemented)**

none

Time = 0.45 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int \frac{1}{1 + \pi x + 3x^2} dx = \frac{2\sqrt{-\pi^2 + 12} \arctan\left(\frac{(\pi + 6x)\sqrt{-\pi^2 + 12}}{\pi^2 - 12}\right)}{\pi^2 - 12}$$

[In] integrate(1/(pi\*x+3\*x^2+1),x, algorithm="fricas")

[Out] 2\*sqrt(-pi^2 + 12)\*arctan((pi + 6\*x)\*sqrt(-pi^2 + 12)/(pi^2 - 12))/(pi^2 - 12)

**Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.81

$$\int \frac{1}{1 + \pi x + 3x^2} dx = -\frac{i \log\left(x + \frac{\pi}{6} - \frac{2i}{\sqrt{12 - \pi^2}} + \frac{i\pi^2}{6\sqrt{12 - \pi^2}}\right)}{\sqrt{12 - \pi^2}} + \frac{i \log\left(x + \frac{\pi}{6} - \frac{i\pi^2}{6\sqrt{12 - \pi^2}} + \frac{2i}{\sqrt{12 - \pi^2}}\right)}{\sqrt{12 - \pi^2}}$$

[In] integrate(1/(pi\*x+3\*x\*\*2+1),x)

[Out] -I\*log(x + pi/6 - 2\*I/sqrt(12 - pi\*\*2) + I\*pi\*\*2/(6\*sqrt(12 - pi\*\*2)))/sqrt(12 - pi\*\*2) + I\*log(x + pi/6 - I\*pi\*\*2/(6\*sqrt(12 - pi\*\*2)) + 2\*I/sqrt(12 - pi\*\*2))/sqrt(12 - pi\*\*2)

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{1}{1 + \pi x + 3x^2} dx = \frac{2 \arctan\left(\frac{\pi + 6x}{\sqrt{-\pi^2 + 12}}\right)}{\sqrt{-\pi^2 + 12}}$$

[In] integrate(1/(pi\*x+3\*x^2+1),x, algorithm="maxima")

[Out] 2\*arctan((pi + 6\*x)/sqrt(-pi^2 + 12))/sqrt(-pi^2 + 12)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{1}{1 + \pi x + 3x^2} dx = \frac{2 \arctan\left(\frac{\pi+6x}{\sqrt{-\pi^2+12}}\right)}{\sqrt{-\pi^2+12}}$$

[In] integrate(1/(pi\*x+3\*x^2+1),x, algorithm="giac")

[Out] 2\*arctan((pi + 6\*x)/sqrt(-pi^2 + 12))/sqrt(-pi^2 + 12)

**Mupad [B] (verification not implemented)**

Time = 9.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.74

$$\int \frac{1}{1 + \pi x + 3x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{\pi+6x}{\sqrt{\pi^2-12}}\right)}{\sqrt{\pi^2-12}}$$

[In] int(1/(Pi\*x + 3\*x^2 + 1),x)

[Out] -(2\*atanh((Pi + 6\*x)/(Pi^2 - 12)^(1/2)))/(Pi^2 - 12)^(1/2)

### 3.87 $\int \frac{1}{1+\pi x-3x^2} dx$

Optimal result	451
Rubi [A] (verified)	451
Mathematica [A] (verified)	452
Maple [A] (verified)	452
Fricas [B] (verification not implemented)	453
Sympy [B] (verification not implemented)	453
Maxima [A] (verification not implemented)	453
Giac [A] (verification not implemented)	454
Mupad [B] (verification not implemented)	454

#### Optimal result

Integrand size = 12, antiderivative size = 27

$$\int \frac{1}{1+\pi x-3x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{\pi-6x}{\sqrt{12+\pi^2}}\right)}{\sqrt{12+\pi^2}}$$

[Out]  $-2*\operatorname{arctanh}((\text{Pi}-6*x)/(\text{Pi}^2+12)^{(1/2)})/(\text{Pi}^2+12)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 212}

$$\int \frac{1}{1+\pi x-3x^2} dx = -\frac{2\operatorname{arctanh}\left(\frac{\pi-6x}{\sqrt{12+\pi^2}}\right)}{\sqrt{12+\pi^2}}$$

[In]  $\text{Int}[(1 + \text{Pi}*x - 3*x^2)^{-1}, x]$

[Out]  $(-2*\text{ArcTanh}[(\text{Pi} - 6*x)/\text{Sqrt}[12 + \text{Pi}^2]])/\text{Sqrt}[12 + \text{Pi}^2]$

#### Rule 212

$\text{Int}[(a_+ + (b_-)*(x_-)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 632

$\text{Int}[(a_+ + (b_-)*(x_-) + (c_-)*(x_-)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{12 + \pi^2 - x^2} dx, x, \pi - 6x\right)\right) \\ &= -\frac{2 \tanh^{-1}\left(\frac{\pi-6x}{\sqrt{12+\pi^2}}\right)}{\sqrt{12 + \pi^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.07

$$\int \frac{1}{1 + \pi x - 3x^2} dx = \frac{2\text{arctanh}\left(\frac{-\pi+6x}{\sqrt{12+\pi^2}}\right)}{\sqrt{12 + \pi^2}}$$

[In] Integrate[(1 + Pi\*x - 3\*x^2)^(-1),x]

[Out] (2\*ArcTanh[(-Pi + 6\*x)/Sqrt[12 + Pi^2]])/Sqrt[12 + Pi^2]

**Maple [A] (verified)**

Time = 1.99 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{2 \operatorname{arctanh}\left(\frac{-\pi+6x}{\sqrt{\pi^2+12}}\right)}{\sqrt{\pi^2+12}}$	26
risch	$\frac{\ln\left(\pi^2-\pi\sqrt{\pi^2+12}+6x\sqrt{\pi^2+12}+12\right)}{\sqrt{\pi^2+12}} - \frac{\ln\left(-\pi^2-\pi\sqrt{\pi^2+12}+6x\sqrt{\pi^2+12}-12\right)}{\sqrt{\pi^2+12}}$	73

[In] int(1/(Pi\*x-3\*x^2+1),x,method=\_RETURNVERBOSE)

[Out] 2/(Pi^2+12)^(1/2)\*arctanh((-Pi+6\*x)/(Pi^2+12)^(1/2))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 51 vs. 2(23) = 46.

Time = 0.40 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.89

$$\int \frac{1}{1 + \pi x - 3x^2} dx = \frac{\log\left(-\frac{\pi^2 - 6\pi x + 18x^2 - (\pi - 6x)\sqrt{\pi^2 + 12} + 6}{\pi x - 3x^2 + 1}\right)}{\sqrt{\pi^2 + 12}}$$

[In] integrate(1/(pi\*x-3\*x^2+1),x, algorithm="fricas")

[Out] log(-(pi^2 - 6\*pi\*x + 18\*x^2 - (pi - 6\*x)\*sqrt(pi^2 + 12) + 6)/(pi\*x - 3\*x^2 + 1))/sqrt(pi^2 + 12)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 76 vs. 2(26) = 52.

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.81

$$\int \frac{1}{1 + \pi x - 3x^2} dx = \frac{\log\left(x - \frac{\pi}{6} + \frac{\pi^2}{6\sqrt{\pi^2 + 12}} + \frac{2}{\sqrt{\pi^2 + 12}}\right)}{\sqrt{\pi^2 + 12}} - \frac{\log\left(x - \frac{\pi}{6} - \frac{2}{\sqrt{\pi^2 + 12}} - \frac{\pi^2}{6\sqrt{\pi^2 + 12}}\right)}{\sqrt{\pi^2 + 12}}$$

[In] integrate(1/(pi\*x-3\*x\*\*2+1),x)

[Out] log(x - pi/6 + pi\*\*2/(6\*sqrt(pi\*\*2 + 12)) + 2/sqrt(pi\*\*2 + 12))/sqrt(pi\*\*2 + 12) - log(x - pi/6 - 2/sqrt(pi\*\*2 + 12) - pi\*\*2/(6\*sqrt(pi\*\*2 + 12)))/sqrt(pi\*\*2 + 12)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.44

$$\int \frac{1}{1 + \pi x - 3x^2} dx = -\frac{\log\left(\frac{\pi - 6x + \sqrt{\pi^2 + 12}}{\pi - 6x - \sqrt{\pi^2 + 12}}\right)}{\sqrt{\pi^2 + 12}}$$

[In] integrate(1/(pi\*x-3\*x^2+1),x, algorithm="maxima")

[Out] -log((pi - 6\*x + sqrt(pi^2 + 12))/(pi - 6\*x - sqrt(pi^2 + 12)))/sqrt(pi^2 + 12)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.67

$$\int \frac{1}{1 + \pi x - 3x^2} dx = -\frac{\log\left(\frac{|\pi - 6x - \sqrt{\pi^2 + 12}|}{|\pi - 6x + \sqrt{\pi^2 + 12}|}\right)}{\sqrt{\pi^2 + 12}}$$

[In] integrate(1/(pi\*x-3\*x^2+1),x, algorithm="giac")

[Out] -log(abs(-pi + 6\*x - sqrt(pi^2 + 12))/abs(-pi + 6\*x + sqrt(pi^2 + 12)))/sqrt(pi^2 + 12)

**Mupad [B] (verification not implemented)**

Time = 9.24 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{1}{1 + \pi x - 3x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{\pi - 6x}{\sqrt{\pi^2 + 12}}\right)}{\sqrt{\pi^2 + 12}}$$

[In] int(1/(Pi\*x - 3\*x^2 + 1),x)

[Out] -(2\*atanh((Pi - 6\*x)/(Pi^2 + 12)^(1/2)))/(Pi^2 + 12)^(1/2)

### 3.88 $\int \frac{1}{a+cx+bx^2} dx$

Optimal result	455
Rubi [A] (verified)	455
Mathematica [A] (verified)	456
Maple [A] (verified)	456
Fricas [A] (verification not implemented)	457
Sympy [B] (verification not implemented)	457
Maxima [F(-2)]	458
Giac [A] (verification not implemented)	458
Mupad [B] (verification not implemented)	458

#### Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{1}{a+cx+bx^2} dx = \frac{2 \arctan\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$$

[Out]  $2*\arctan((2*b*x+c)/(4*a*b-c^2)^{(1/2)})/(4*a*b-c^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {632, 210}

$$\int \frac{1}{a+cx+bx^2} dx = \frac{2 \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$$

[In]  $\text{Int}[(a + c*x + b*x^2)^{-1}, x]$

[Out]  $(2*\text{ArcTan}[(c + 2*b*x)/\text{Sqrt}[4*a*b - c^2]])/\text{Sqrt}[4*a*b - c^2]$

#### Rule 210

$\text{Int}[(a + (b_*)*(x_*)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 632

$\text{Int}[(a + (b_*)*(x_*) + (c_*)*(x_*)^2)^{-1}, x\_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c\},$

x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-4ab + c^2 - x^2} dx, x, c + 2bx\right)\right) \\ &= \frac{2 \tan^{-1}\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + cx + bx^2} dx = \frac{2 \arctan\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$$

[In] Integrate[(a + c\*x + b\*x^2)^(-1),x]

[Out] (2\*ArcTan[(c + 2\*b\*x)/Sqrt[4\*a\*b - c^2]])/Sqrt[4\*a\*b - c^2]

**Maple [A] (verified)**

Time = 2.12 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{2 \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$	35
risch	$-\frac{\ln\left(2bx+\sqrt{-4ab+c^2}+c\right)}{\sqrt{-4ab+c^2}} + \frac{\ln\left(-2bx+\sqrt{-4ab+c^2}-c\right)}{\sqrt{-4ab+c^2}}$	61

[In] int(1/(b\*x^2+c\*x+a),x,method=\_RETURNVERBOSE)

[Out] 2\*arctan((2\*b\*x+c)/(4\*a\*b-c^2)^(1/2))/(4\*a\*b-c^2)^(1/2)



**Fricas [A] (verification not implemented)**

none

Time = 0.42 (sec) , antiderivative size = 113, normalized size of antiderivative = 2.97

$$\int \frac{1}{a + cx + bx^2} dx = \left[ -\frac{\sqrt{-4ab + c^2} \log\left(\frac{2b^2x^2 + 2bcx - 2ab + c^2 - \sqrt{-4ab + c^2}(2bx + c)}{bx^2 + cx + a}\right)}{4ab - c^2}, \right. \\ \left. -\frac{2 \arctan\left(-\frac{2bx + c}{\sqrt{4ab - c^2}}\right)}{\sqrt{4ab - c^2}} \right]$$

`[In] integrate(1/(b*x^2+c*x+a),x, algorithm="fricas")`

```
[Out] [-sqrt(-4*a*b + c^2)*log((2*b^2*x^2 + 2*b*c*x - 2*a*b + c^2 - sqrt(-4*a*b +
c^2)*(2*b*x + c))/(b*x^2 + c*x + a))/(4*a*b - c^2), -2*arctan(-(2*b*x + c)
/sqrt(4*a*b - c^2))/sqrt(4*a*b - c^2)]
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(32) = 64.

Time = 0.10 (sec) , antiderivative size = 124, normalized size of antiderivative = 3.26

$$\int \frac{1}{a + cx + bx^2} dx = -\sqrt{-\frac{1}{4ab - c^2}} \log\left(x + \frac{-4ab\sqrt{-\frac{1}{4ab - c^2}} + c^2\sqrt{-\frac{1}{4ab - c^2}} + c}{2b}\right) \\ + \sqrt{-\frac{1}{4ab - c^2}} \log\left(x + \frac{4ab\sqrt{-\frac{1}{4ab - c^2}} - c^2\sqrt{-\frac{1}{4ab - c^2}} + c}{2b}\right)$$

`[In] integrate(1/(b*x**2+c*x+a),x)`

```
[Out] -sqrt(-1/(4*a*b - c**2))*log(x + (-4*a*b*sqrt(-1/(4*a*b - c**2)) + c**2*sqrt(-1/(4*a*b - c**2)) + c)/(2*b)) + sqrt(-1/(4*a*b - c**2))*log(x + (4*a*b*sqrt(-1/(4*a*b - c**2)) - c**2*sqrt(-1/(4*a*b - c**2)) + c)/(2*b))
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{1}{a + cx + bx^2} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(b\*x^2+c\*x+a),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(c^2-4\*a\*b>0)', see 'assume?' for more data

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.89

$$\int \frac{1}{a + cx + bx^2} dx = \frac{2 \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$$

[In] integrate(1/(b\*x^2+c\*x+a),x, algorithm="giac")

[Out] 2\*arctan((2\*b\*x + c)/sqrt(4\*a\*b - c^2))/sqrt(4\*a\*b - c^2)

**Mupad [B] (verification not implemented)**

Time = 8.99 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.21

$$\int \frac{1}{a + cx + bx^2} dx = \frac{2 \operatorname{atan}\left(\frac{c}{\sqrt{4ab-c^2}} + \frac{2bx}{\sqrt{4ab-c^2}}\right)}{\sqrt{4ab-c^2}}$$

[In] int(1/(a + c\*x + b\*x^2),x)

[Out] (2\*atan(c/(4\*a\*b - c^2)^(1/2) + (2\*b\*x)/(4\*a\*b - c^2)^(1/2)))/(4\*a\*b - c^2)^(1/2)

### 3.89 $\int \frac{1}{b+2ax+bx^2} dx$

Optimal result	459
Rubi [A] (verified)	459
Mathematica [A] (verified)	460
Maple [A] (verified)	460
Fricas [A] (verification not implemented)	461
Sympy [B] (verification not implemented)	461
Maxima [F(-2)]	462
Giac [A] (verification not implemented)	462
Mupad [B] (verification not implemented)	462

#### Optimal result

Integrand size = 13, antiderivative size = 35

$$\int \frac{1}{b+2ax+bx^2} dx = -\frac{\operatorname{arctanh}\left(\frac{a+bx}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

[Out]  $-\operatorname{arctanh}((b*x+a)/(a^2-b^2)^{(1/2)})/(a^2-b^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$ , Rules used = {632, 212}

$$\int \frac{1}{b+2ax+bx^2} dx = -\frac{\operatorname{arctanh}\left(\frac{a+bx}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}}$$

[In]  $\operatorname{Int}[(b + 2*a*x + b*x^2)^{-1}, x]$

[Out]  $-(\operatorname{ArcTanh}[(a + b*x)/\operatorname{Sqrt}[a^2 - b^2]]/\operatorname{Sqrt}[a^2 - b^2])$

#### Rule 212

$\operatorname{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \mid\mid \operatorname{LtQ}[b, 0])$

#### Rule 632

$\operatorname{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Dist}[-2, \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \operatorname{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{4(a^2 - b^2) - x^2} dx, x, 2a + 2bx\right)\right) \\ &= -\frac{\tanh^{-1}\left(\frac{a+bx}{\sqrt{a^2-b^2}}\right)}{\sqrt{a^2-b^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int \frac{1}{b + 2ax + bx^2} dx = \frac{\arctan\left(\frac{a+bx}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$$

[In] `Integrate[(b + 2*a*x + b*x^2)^(-1),x]`

[Out] `ArcTan[(a + b*x)/Sqrt[-a^2 + b^2]]/Sqrt[-a^2 + b^2]`

**Maple [A] (verified)**

Time = 2.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{\arctan\left(\frac{2bx+2a}{2\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$	35
risch	$\frac{\ln(-bx+\sqrt{a^2-b^2}-a)}{2\sqrt{a^2-b^2}} - \frac{\ln(bx+\sqrt{a^2-b^2}+a)}{2\sqrt{a^2-b^2}}$	65

[In] `int(1/(b*x^2+2*a*x+b),x,method=_RETURNVERBOSE)`

[Out] `1/(-a^2+b^2)^(1/2)*arctan(1/2*(2*b*x+2*a)/(-a^2+b^2)^(1/2))`

**Fricas [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 124, normalized size of antiderivative = 3.54

$$\int \frac{1}{b + 2ax + bx^2} dx = \left[ \frac{\log\left(\frac{b^2x^2 + 2abx + 2a^2 - b^2 - 2\sqrt{a^2 - b^2}(bx + a)}{bx^2 + 2ax + b}\right)}{2\sqrt{a^2 - b^2}}, \right. \\ \left. - \frac{\sqrt{-a^2 + b^2} \arctan\left(-\frac{\sqrt{-a^2 + b^2}(bx + a)}{a^2 - b^2}\right)}{a^2 - b^2} \right]$$

[In] integrate(1/(b\*x^2+2\*a\*x+b),x, algorithm="fricas")

```
[Out] [1/2*log((b^2*x^2 + 2*a*b*x + 2*a^2 - b^2 - 2*sqrt(a^2 - b^2)*(b*x + a))/(b
*x^2 + 2*a*x + b))/sqrt(a^2 - b^2), -sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b
^2)*(b*x + a)/(a^2 - b^2))/(a^2 - b^2)]
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(27) = 54.

Time = 0.12 (sec) , antiderivative size = 100, normalized size of antiderivative = 2.86

$$\int \frac{1}{b + 2ax + bx^2} dx = \frac{\sqrt{\frac{1}{(a-b)(a+b)}} \log\left(x + \frac{-a^2\sqrt{\frac{1}{(a-b)(a+b)}} + a + b^2\sqrt{\frac{1}{(a-b)(a+b)}}}{b}\right)}{2} \\ - \frac{\sqrt{\frac{1}{(a-b)(a+b)}} \log\left(x + \frac{a^2\sqrt{\frac{1}{(a-b)(a+b)}} + a - b^2\sqrt{\frac{1}{(a-b)(a+b)}}}{b}\right)}{2}$$

[In] integrate(1/(b\*x\*\*2+2\*a\*x+b),x)

```
[Out] sqrt(1/((a - b)*(a + b)))*log(x + (-a**2*sqrt(1/((a - b)*(a + b))) + a + b*
*2*sqrt(1/((a - b)*(a + b))))/b)/2 - sqrt(1/((a - b)*(a + b)))*log(x + (a**
2*sqrt(1/((a - b)*(a + b))) + a - b**2*sqrt(1/((a - b)*(a + b))))/b)/2
```

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{1}{b + 2ax + bx^2} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(b\*x^2+2\*a\*x+b),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(4\*a^2-4\*b^2>0)', see 'assume?' for more details)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \frac{1}{b + 2ax + bx^2} dx = \frac{\arctan\left(\frac{bx+a}{\sqrt{-a^2+b^2}}\right)}{\sqrt{-a^2+b^2}}$$

[In] integrate(1/(b\*x^2+2\*a\*x+b),x, algorithm="giac")

[Out] arctan((b\*x + a)/sqrt(-a^2 + b^2))/sqrt(-a^2 + b^2)

**Mupad [B] (verification not implemented)**

Time = 9.06 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{1}{b + 2ax + bx^2} dx = -\frac{\operatorname{atanh}\left(\frac{a+bx}{\sqrt{a+b}\sqrt{a-b}}\right)}{\sqrt{a+b}\sqrt{a-b}}$$

[In] int(1/(b + 2\*a\*x + b\*x^2),x)

[Out] -atanh((a + b\*x)/((a + b)^(1/2)\*(a - b)^(1/2)))/((a + b)^(1/2)\*(a - b)^(1/2))

### 3.90 $\int \frac{1}{b+2ax-bx^2} dx$

Optimal result	463
Rubi [A] (verified)	463
Mathematica [A] (verified)	464
Maple [A] (verified)	464
Fricas [B] (verification not implemented)	465
Sympy [B] (verification not implemented)	465
Maxima [A] (verification not implemented)	465
Giac [A] (verification not implemented)	466
Mupad [B] (verification not implemented)	466

#### Optimal result

Integrand size = 14, antiderivative size = 32

$$\int \frac{1}{b+2ax-bx^2} dx = -\frac{\operatorname{arctanh}\left(\frac{a-bx}{\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$$

[Out]  $-\operatorname{arctanh}((-b*x+a)/(a^2+b^2)^{(1/2)})/(a^2+b^2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {632, 212}

$$\int \frac{1}{b+2ax-bx^2} dx = -\frac{\operatorname{arctanh}\left(\frac{a-bx}{\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$$

[In]  $\operatorname{Int}[(b + 2*a*x - b*x^2)^{-1}, x]$

[Out]  $-(\operatorname{ArcTanh}[(a - b*x)/\operatorname{Sqrt}[a^2 + b^2]]/\operatorname{Sqrt}[a^2 + b^2])$

#### Rule 212

$\operatorname{Int}[(a_.) + (b_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \mid\mid \operatorname{LtQ}[b, 0])$

#### Rule 632

$\operatorname{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Dist}[-2, \operatorname{Subst}[\operatorname{Int}[1/\operatorname{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \operatorname{FreeQ}\{a, b, c\},$

`x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{4(a^2 + b^2) - x^2} dx, x, 2a - 2bx\right)\right) \\ &= -\frac{\tanh^{-1}\left(\frac{a-bx}{\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.28

$$\int \frac{1}{b + 2ax - bx^2} dx = -\frac{\arctan\left(\frac{-a+bx}{\sqrt{-a^2-b^2}}\right)}{\sqrt{-a^2-b^2}}$$

[In] `Integrate[(b + 2*a*x - b*x^2)^(-1),x]`

[Out] `-(ArcTan[(-a + b*x)/Sqrt[-a^2 - b^2]]/Sqrt[-a^2 - b^2])`

**Maple [A] (verified)**

Time = 2.11 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

method	result	size
default	$-\frac{\operatorname{arctanh}\left(\frac{-2bx+2a}{2\sqrt{a^2+b^2}}\right)}{\sqrt{a^2+b^2}}$	32
risch	$\frac{\ln\left(bx+\sqrt{a^2+b^2}-a\right)}{2\sqrt{a^2+b^2}} - \frac{\ln\left(-bx+\sqrt{a^2+b^2}+a\right)}{2\sqrt{a^2+b^2}}$	57

[In] `int(1/(-b*x^2+2*a*x+b),x,method=_RETURNVERBOSE)`

[Out] `-1/(a^2+b^2)^(1/2)*arctanh(1/2*(-2*b*x+2*a)/(a^2+b^2)^(1/2))`



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 67 vs.  $2(30) = 60$ .

Time = 0.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.09

$$\int \frac{1}{b + 2ax - bx^2} dx = \frac{\log\left(\frac{b^2x^2 - 2abx + 2a^2 + b^2 + 2\sqrt{a^2 + b^2}(bx - a)}{bx^2 - 2ax - b}\right)}{2\sqrt{a^2 + b^2}}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b),x, algorithm="fricas")

[Out] 1/2\*log((b^2\*x^2 - 2\*a\*b\*x + 2\*a^2 + b^2 + 2\*sqrt(a^2 + b^2)\*(b\*x - a))/(b\*x^2 - 2\*a\*x - b))/sqrt(a^2 + b^2)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 102 vs.  $2(27) = 54$ .

Time = 0.13 (sec) , antiderivative size = 102, normalized size of antiderivative = 3.19

$$\int \frac{1}{b + 2ax - bx^2} dx = -\frac{\sqrt{\frac{1}{a^2 + b^2}} \log\left(x + \frac{-a^2 \sqrt{\frac{1}{a^2 + b^2}} - a - b^2 \sqrt{\frac{1}{a^2 + b^2}}}{b}\right)}{2} + \frac{\sqrt{\frac{1}{a^2 + b^2}} \log\left(x + \frac{a^2 \sqrt{\frac{1}{a^2 + b^2}} - a + b^2 \sqrt{\frac{1}{a^2 + b^2}}}{b}\right)}{2}$$

[In] integrate(1/(-b\*x\*\*2+2\*a\*x+b),x)

[Out] -sqrt(1/(a\*\*2 + b\*\*2))\*log(x + (-a\*\*2\*sqrt(1/(a\*\*2 + b\*\*2)) - a - b\*\*2\*sqrt(1/(a\*\*2 + b\*\*2)))/b)/2 + sqrt(1/(a\*\*2 + b\*\*2))\*log(x + (a\*\*2\*sqrt(1/(a\*\*2 + b\*\*2)) - a + b\*\*2\*sqrt(1/(a\*\*2 + b\*\*2)))/b)/2

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.53

$$\int \frac{1}{b + 2ax - bx^2} dx = -\frac{\log\left(\frac{bx - a - \sqrt{a^2 + b^2}}{bx - a + \sqrt{a^2 + b^2}}\right)}{2\sqrt{a^2 + b^2}}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b),x, algorithm="maxima")

[Out] -1/2\*log((b\*x - a - sqrt(a^2 + b^2))/(b\*x - a + sqrt(a^2 + b^2)))/sqrt(a^2 + b^2)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.72

$$\int \frac{1}{b + 2ax - bx^2} dx = -\frac{\log\left(\left|\frac{2bx - 2a - 2\sqrt{a^2 + b^2}}{2bx - 2a + 2\sqrt{a^2 + b^2}}\right|\right)}{2\sqrt{a^2 + b^2}}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b),x, algorithm="giac")

[Out] -1/2\*log(abs(2\*b\*x - 2\*a - 2\*sqrt(a^2 + b^2))/abs(2\*b\*x - 2\*a + 2\*sqrt(a^2 + b^2)))/sqrt(a^2 + b^2)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \frac{1}{b + 2ax - bx^2} dx = -\frac{\operatorname{atanh}\left(\frac{a - bx}{\sqrt{a^2 + b^2}}\right)}{\sqrt{a^2 + b^2}}$$

[In] int(1/(b + 2\*a\*x - b\*x^2),x)

[Out] -atanh((a - b\*x)/(a^2 + b^2)^(1/2))/(a^2 + b^2)^(1/2)

### 3.91 $\int \frac{1}{(2+4x+3x^2)^2} dx$

Optimal result	467
Rubi [A] (verified)	467
Mathematica [A] (verified)	468
Maple [A] (verified)	468
Fricas [A] (verification not implemented)	469
Sympy [A] (verification not implemented)	469
Maxima [A] (verification not implemented)	469
Giac [A] (verification not implemented)	470
Mupad [B] (verification not implemented)	470

#### Optimal result

Integrand size = 12, antiderivative size = 43

$$\int \frac{1}{(2+4x+3x^2)^2} dx = \frac{2+3x}{4(2+4x+3x^2)} + \frac{3 \arctan\left(\frac{2+3x}{\sqrt{2}}\right)}{4\sqrt{2}}$$

[Out] 1/4\*(2+3\*x)/(3\*x^2+4\*x+2)+3/8\*arctan(1/2\*(2+3\*x)\*2^(1/2))\*2^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {628, 632, 210}

$$\int \frac{1}{(2+4x+3x^2)^2} dx = \frac{3 \arctan\left(\frac{3x+2}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{3x+2}{4(3x^2+4x+2)}$$

[In] Int[(2 + 4\*x + 3\*x^2)^(-2), x]

[Out] (2 + 3\*x)/(4\*(2 + 4\*x + 3\*x^2)) + (3\*ArcTan[(2 + 3\*x)/Sqrt[2]])/(4\*Sqrt[2])

#### Rule 210

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(-(Rt[-a, 2]\*Rt[-b, 2])^(-1))\*ArcTan[Rt[-b, 2]\*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

#### Rule 628

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p+1) / ((p+1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p +

3)/((p + 1)\*(b^2 - 4\*a\*c)), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

### Rule 632

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-1), x\_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4\*a\*c - x^2, x], x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 + 3x}{4(2 + 4x + 3x^2)} + \frac{3}{4} \int \frac{1}{2 + 4x + 3x^2} dx \\ &= \frac{2 + 3x}{4(2 + 4x + 3x^2)} - \frac{3}{2} \text{Subst}\left(\int \frac{1}{-8 - x^2} dx, x, 4 + 6x\right) \\ &= \frac{2 + 3x}{4(2 + 4x + 3x^2)} + \frac{3 \tan^{-1}\left(\frac{2+3x}{\sqrt{2}}\right)}{4\sqrt{2}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{(2 + 4x + 3x^2)^2} dx = \frac{2 + 3x}{4(2 + 4x + 3x^2)} + \frac{3 \arctan\left(\frac{2+3x}{\sqrt{2}}\right)}{4\sqrt{2}}$$

[In] Integrate[(2 + 4\*x + 3\*x^2)^(-2), x]

[Out] (2 + 3\*x)/(4\*(2 + 4\*x + 3\*x^2)) + (3\*ArcTan[(2 + 3\*x)/Sqrt[2]])/(4\*Sqrt[2])

### Maple [A] (verified)

Time = 2.20 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

method	result	size
risch	$\frac{\frac{x}{4} + \frac{1}{6}}{x^2 + \frac{4}{3}x + \frac{2}{3}} + \frac{3 \arctan\left(\frac{(2+3x)\sqrt{2}}{2}\right)\sqrt{2}}{8}$	34
default	$\frac{4+6x}{24x^2+32x+16} + \frac{3\sqrt{2} \arctan\left(\frac{(4+6x)\sqrt{2}}{4}\right)}{8}$	37

[In] int(1/(3\*x^2+4\*x+2)^2,x,method=\_RETURNVERBOSE)

[Out]  $(1/4*x+1/6)/(x^2+4/3*x+2/3)+3/8*\arctan(1/2*(2+3*x)*2^{(1/2)})*2^{(1/2)}$

### Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.05

$$\int \frac{1}{(2+4x+3x^2)^2} dx = \frac{3\sqrt{2}(3x^2+4x+2)\arctan\left(\frac{1}{2}\sqrt{2}(3x+2)\right)+6x+4}{8(3x^2+4x+2)}$$

[In] `integrate(1/(3*x^2+4*x+2)^2,x, algorithm="fricas")`

[Out]  $1/8*(3*\sqrt{2}*(3*x^2+4*x+2)*\arctan(1/2*\sqrt{2}*(3*x+2))+6*x+4)/(3*x^2+4*x+2)$

### Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.91

$$\int \frac{1}{(2+4x+3x^2)^2} dx = \frac{3x+2}{12x^2+16x+8} + \frac{3\sqrt{2}\operatorname{atan}\left(\frac{3\sqrt{2}x}{2} + \sqrt{2}\right)}{8}$$

[In] `integrate(1/(3*x**2+4*x+2)**2,x)`

[Out]  $(3*x+2)/(12*x**2+16*x+8)+3*\sqrt{2}*\operatorname{atan}(3*\sqrt{2}*x/2+\sqrt{2})/8$

### Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

$$\int \frac{1}{(2+4x+3x^2)^2} dx = \frac{3}{8}\sqrt{2}\arctan\left(\frac{1}{2}\sqrt{2}(3x+2)\right) + \frac{3x+2}{4(3x^2+4x+2)}$$

[In] `integrate(1/(3*x^2+4*x+2)^2,x, algorithm="maxima")`

[Out]  $3/8*\sqrt{2}*\arctan(1/2*\sqrt{2}*(3*x+2))+1/4*(3*x+2)/(3*x^2+4*x+2)$

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

$$\int \frac{1}{(2 + 4x + 3x^2)^2} dx = \frac{3}{8} \sqrt{2} \arctan\left(\frac{1}{2} \sqrt{2}(3x + 2)\right) + \frac{3x + 2}{4(3x^2 + 4x + 2)}$$

[In] integrate(1/(3\*x^2+4\*x+2)^2,x, algorithm="giac")

[Out] 3/8\*sqrt(2)\*arctan(1/2\*sqrt(2)\*(3\*x + 2)) + 1/4\*(3\*x + 2)/(3\*x^2 + 4\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 0.04 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.77

$$\int \frac{1}{(2 + 4x + 3x^2)^2} dx = \frac{\frac{x}{4} + \frac{1}{6}}{x^2 + \frac{4x}{3} + \frac{2}{3}} + \frac{3\sqrt{2} \operatorname{atan}\left(\frac{3\sqrt{2}x}{2} + \sqrt{2}\right)}{8}$$

[In] int(1/(4\*x + 3\*x^2 + 2)^2,x)

[Out] (x/4 + 1/6)/((4\*x)/3 + x^2 + 2/3) + (3\*2^(1/2)\*atan((3\*2^(1/2)\*x)/2 + 2^(1/2)))/8

### 3.92 $\int \frac{1}{(2+4x-3x^2)^2} dx$

Optimal result	471
Rubi [A] (verified)	471
Mathematica [A] (verified)	472
Maple [A] (verified)	473
Fricas [A] (verification not implemented)	473
Sympy [A] (verification not implemented)	473
Maxima [A] (verification not implemented)	474
Giac [A] (verification not implemented)	474
Mupad [B] (verification not implemented)	474

#### Optimal result

Integrand size = 12, antiderivative size = 43

$$\int \frac{1}{(2+4x-3x^2)^2} dx = -\frac{2-3x}{20(2+4x-3x^2)} - \frac{3\operatorname{arctanh}\left(\frac{2-3x}{\sqrt{10}}\right)}{20\sqrt{10}}$$

[Out] 1/20\*(-2+3\*x)/(-3\*x^2+4\*x+2)-3/200\*arctanh(1/10\*(2-3\*x)\*10^(1/2))\*10^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {628, 632, 212}

$$\int \frac{1}{(2+4x-3x^2)^2} dx = -\frac{3\operatorname{arctanh}\left(\frac{2-3x}{\sqrt{10}}\right)}{20\sqrt{10}} - \frac{2-3x}{20(-3x^2+4x+2)}$$

[In] Int[(2 + 4\*x - 3\*x^2)^(-2), x]

[Out] -1/20\*(2 - 3\*x)/(2 + 4\*x - 3\*x^2) - (3\*ArcTanh[(2 - 3\*x)/Sqrt[10]])/(20\*Sqrt[10])

#### Rule 212

Int[((a\_) + (b\_)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

### Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{2-3x}{20(2+4x-3x^2)} + \frac{3}{20} \int \frac{1}{2+4x-3x^2} dx \\
 &= -\frac{2-3x}{20(2+4x-3x^2)} - \frac{3}{10} \text{Subst}\left(\int \frac{1}{40-x^2} dx, x, 4-6x\right) \\
 &= -\frac{2-3x}{20(2+4x-3x^2)} - \frac{3 \tanh^{-1}\left(\frac{2-3x}{\sqrt{10}}\right)}{20\sqrt{10}}
 \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.44

$$\begin{aligned}
 &\int \frac{1}{(2+4x-3x^2)^2} dx \\
 &= \frac{2-3x}{20(-2-4x+3x^2)} - \frac{3 \log(2+\sqrt{10}-3x)}{40\sqrt{10}} + \frac{3 \log(-2+\sqrt{10}+3x)}{40\sqrt{10}}
 \end{aligned}$$

```
[In] Integrate[(2 + 4*x - 3*x^2)^(-2), x]
```

```
[Out] (2 - 3*x)/(20*(-2 - 4*x + 3*x^2)) - (3*Log[2 + Sqrt[10] - 3*x])/(40*Sqrt[10
]) + (3*Log[-2 + Sqrt[10] + 3*x])/(40*Sqrt[10])
```



**Maple [A] (verified)**

Time = 2.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

method	result	size
default	$-\frac{-4+6x}{40(3x^2-4x-2)} + \frac{3\sqrt{10} \operatorname{arctanh}\left(\frac{(-4+6x)\sqrt{10}}{20}\right)}{200}$	37
risch	$\frac{-\frac{x}{20} + \frac{1}{30}}{x^2 - \frac{4}{3}x - \frac{2}{3}} + \frac{3\sqrt{10} \ln(3x-2+\sqrt{10})}{400} - \frac{3\sqrt{10} \ln(3x-2-\sqrt{10})}{400}$	48

[In] `int(1/(-3*x^2+4*x+2)^2,x,method=_RETURNVERBOSE)`

[Out]  $-1/40*(-4+6*x)/(3*x^2-4*x-2)+3/200*10^{(1/2)}*\operatorname{arctanh}(1/20*(-4+6*x)*10^{(1/2)})$

**Fricas [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.58

$$\int \frac{1}{(2+4x-3x^2)^2} dx = \frac{3\sqrt{10}(3x^2-4x-2) \log\left(\frac{9x^2+2\sqrt{10}(3x-2)-12x+14}{3x^2-4x-2}\right) - 60x + 40}{400(3x^2-4x-2)}$$

[In] `integrate(1/(-3*x^2+4*x+2)^2,x, algorithm="fricas")`

[Out]  $1/400*(3*\operatorname{sqrt}(10)*(3*x^2-4*x-2)*\log((9*x^2+2*\operatorname{sqrt}(10)*(3*x-2)-12*x+14)/(3*x^2-4*x-2))-60*x+40)/(3*x^2-4*x-2)$

**Sympy [A] (verification not implemented)**

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.35

$$\int \frac{1}{(2+4x-3x^2)^2} dx = \frac{2-3x}{60x^2-80x-40} + \frac{3\sqrt{10} \log\left(x - \frac{2}{3} + \frac{\sqrt{10}}{3}\right)}{400} - \frac{3\sqrt{10} \log\left(x - \frac{\sqrt{10}}{3} - \frac{2}{3}\right)}{400}$$

[In] `integrate(1/(-3*x**2+4*x+2)**2,x)`

[Out]  $(2-3*x)/(60*x**2-80*x-40)+3*\operatorname{sqrt}(10)*\log(x-2/3+\operatorname{sqrt}(10)/3)/400-3*\operatorname{sqrt}(10)*\log(x-\operatorname{sqrt}(10)/3-2/3)/400$

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.09

$$\int \frac{1}{(2+4x-3x^2)^2} dx = -\frac{3}{400} \sqrt{10} \log \left( \frac{3x - \sqrt{10} - 2}{3x + \sqrt{10} - 2} \right) - \frac{3x - 2}{20(3x^2 - 4x - 2)}$$

[In] integrate(1/(-3\*x^2+4\*x+2)^2,x, algorithm="maxima")

[Out] -3/400\*sqrt(10)\*log((3\*x - sqrt(10) - 2)/(3\*x + sqrt(10) - 2)) - 1/20\*(3\*x - 2)/(3\*x^2 - 4\*x - 2)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.19

$$\int \frac{1}{(2+4x-3x^2)^2} dx = -\frac{3}{400} \sqrt{10} \log \left( \frac{|6x - 2\sqrt{10} - 4|}{|6x + 2\sqrt{10} - 4|} \right) - \frac{3x - 2}{20(3x^2 - 4x - 2)}$$

[In] integrate(1/(-3\*x^2+4\*x+2)^2,x, algorithm="giac")

[Out] -3/400\*sqrt(10)\*log(abs(6\*x - 2\*sqrt(10) - 4)/abs(6\*x + 2\*sqrt(10) - 4)) - 1/20\*(3\*x - 2)/(3\*x^2 - 4\*x - 2)

**Mupad [B] (verification not implemented)**

Time = 9.01 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.79

$$\int \frac{1}{(2+4x-3x^2)^2} dx = \frac{3\sqrt{10} \operatorname{atanh}\left(\sqrt{10}\left(\frac{3x}{10} - \frac{1}{5}\right)\right)}{200} + \frac{\frac{x}{20} - \frac{1}{30}}{-x^2 + \frac{4x}{3} + \frac{2}{3}}$$

[In] int(1/(4\*x - 3\*x^2 + 2)^2,x)

[Out] (3\*10^(1/2)\*atanh(10^(1/2)\*((3\*x)/10 - 1/5)))/200 + (x/20 - 1/30)/((4\*x)/3 - x^2 + 2/3)

### 3.93 $\int \frac{1}{(2+5x+3x^2)^2} dx$

Optimal result	475
Rubi [A] (verified)	475
Mathematica [A] (verified)	476
Maple [A] (verified)	476
Fricas [A] (verification not implemented)	477
Sympy [A] (verification not implemented)	477
Maxima [A] (verification not implemented)	477
Giac [A] (verification not implemented)	478
Mupad [B] (verification not implemented)	478

#### Optimal result

Integrand size = 12, antiderivative size = 34

$$\int \frac{1}{(2+5x+3x^2)^2} dx = -\frac{5+6x}{2+5x+3x^2} + 6\log(1+x) - 6\log(2+3x)$$

[Out]  $(-5-6*x)/(3*x^2+5*x+2)+6*\ln(1+x)-6*\ln(2+3*x)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {628, 630, 31}

$$\int \frac{1}{(2+5x+3x^2)^2} dx = -\frac{6x+5}{3x^2+5x+2} + 6\log(x+1) - 6\log(3x+2)$$

[In]  $\text{Int}[(2+5*x+3*x^2)^{-2}, x]$

[Out]  $-((5+6*x)/(2+5*x+3*x^2))+6*\text{Log}[1+x]-6*\text{Log}[2+3*x]$

#### Rule 31

$\text{Int}[(a_+ + (b_+)(x_+))^{-1}, x\_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] \text{ ; FreeQ}\{a, b\}, x]$

#### Rule 628

$\text{Int}[(a_+ + (b_+)(x_+) + (c_+)(x_+)^2)^{(p_+)}, x\_Symbol] \rightarrow \text{Simp}[(b + 2*c*x) * ((a + b*x + c*x^2)^{(p+1}) / ((p+1)*(b^2 - 4*a*c))), x] - \text{Dist}[2*c*((2*p+3) / ((p+1)*(b^2 - 4*a*c))), \text{Int}[(a + b*x + c*x^2)^{(p+1)}, x], x] \text{ ; Free}$

$Q[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{NeQ}[p, -3/2] \ \&\& \ \text{IntegerQ}[4*p]$

### Rule 630

$\text{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^{-1}, x\_Symbol] \ :> \ \text{With}[\{q = \text{Rt}[b^2 - 4*a*c, 2]\}, \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 - q/2 + c*x, x], x], x] - \text{Dist}[c/q, \text{Int}[1/\text{Simp}[b/2 + q/2 + c*x, x], x], x]] \ /; \ \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[b^2 - 4*a*c] \ \&\& \ \text{PerfectSquareQ}[b^2 - 4*a*c]$

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{5+6x}{2+5x+3x^2} - 6 \int \frac{1}{2+5x+3x^2} dx \\ &= -\frac{5+6x}{2+5x+3x^2} - 18 \int \frac{1}{2+3x} dx + 18 \int \frac{1}{3+3x} dx \\ &= -\frac{5+6x}{2+5x+3x^2} + 6 \log(1+x) - 6 \log(2+3x) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.97

$$\int \frac{1}{(2+5x+3x^2)^2} dx = \frac{-5-6x}{2+5x+3x^2} + 6 \log(1+x) - 6 \log(2+3x)$$

[In] Integrate[(2 + 5\*x + 3\*x^2)^(-2),x]

[Out] (-5 - 6\*x)/(2 + 5\*x + 3\*x^2) + 6\*Log[1 + x] - 6\*Log[2 + 3\*x]

### Maple [A] (verified)

Time = 1.98 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.94

method	result	size
default	$-\frac{1}{1+x} + 6 \ln(1+x) - \frac{3}{2+3x} - 6 \ln(2+3x)$	32
risch	$\frac{-2x-\frac{5}{3}}{x^2+\frac{5}{3}x+\frac{2}{3}} + 6 \ln(1+x) - 6 \ln(2+3x)$	32
norman	$\frac{\frac{15}{2}x^2+\frac{13}{2}x}{3x^2+5x+2} + 6 \ln(1+x) - 6 \ln(2+3x)$	38
parallelrisch	$\frac{36 \ln(1+x)x^2 - 36 \ln(\frac{2}{3}+x)x^2 + 60 \ln(1+x)x - 60 \ln(\frac{2}{3}+x)x + 15x^2 + 24 \ln(1+x) - 24 \ln(\frac{2}{3}+x) + 13x}{6x^2+10x+4}$	68

[In] `int(1/(3*x^2+5*x+2)^2,x,method=_RETURNVERBOSE)`

[Out]  $-1/(1+x)+6*\ln(1+x)-3/(2+3*x)-6*\ln(2+3*x)$

### Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.56

$$\int \frac{1}{(2+5x+3x^2)^2} dx$$

$$= -\frac{6(3x^2+5x+2)\log(3x+2) - 6(3x^2+5x+2)\log(x+1) + 6x+5}{3x^2+5x+2}$$

[In] `integrate(1/(3*x^2+5*x+2)^2,x, algorithm="fricas")`

[Out]  $-(6*(3*x^2 + 5*x + 2)*\log(3*x + 2) - 6*(3*x^2 + 5*x + 2)*\log(x + 1) + 6*x + 5)/(3*x^2 + 5*x + 2)$

### Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.91

$$\int \frac{1}{(2+5x+3x^2)^2} dx = \frac{-6x-5}{3x^2+5x+2} - 6\log\left(x + \frac{2}{3}\right) + 6\log(x+1)$$

[In] `integrate(1/(3*x**2+5*x+2)**2,x)`

[Out]  $(-6*x - 5)/(3*x**2 + 5*x + 2) - 6*\log(x + 2/3) + 6*\log(x + 1)$

### Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{1}{(2+5x+3x^2)^2} dx = -\frac{6x+5}{3x^2+5x+2} - 6\log(3x+2) + 6\log(x+1)$$

[In] `integrate(1/(3*x^2+5*x+2)^2,x, algorithm="maxima")`

[Out]  $-(6*x + 5)/(3*x^2 + 5*x + 2) - 6*\log(3*x + 2) + 6*\log(x + 1)$

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.06

$$\int \frac{1}{(2 + 5x + 3x^2)^2} dx = -\frac{6x + 5}{3x^2 + 5x + 2} - 6 \log(|3x + 2|) + 6 \log(|x + 1|)$$

[In] integrate(1/(3\*x^2+5\*x+2)^2,x, algorithm="giac")

[Out] -(6\*x + 5)/(3\*x^2 + 5\*x + 2) - 6\*log(abs(3\*x + 2)) + 6\*log(abs(x + 1))

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00

$$\int \frac{1}{(2 + 5x + 3x^2)^2} dx = -6 \ln\left(\frac{3x + 2}{x + 1}\right) - \frac{2(3x + \frac{5}{2})}{3x^2 + 5x + 2}$$

[In] int(1/(5\*x + 3\*x^2 + 2)^2,x)

[Out] - 6\*log((3\*x + 2)/(x + 1)) - (2\*(3\*x + 5/2))/(5\*x + 3\*x^2 + 2)

### 3.94 $\int \frac{1}{(2+5x-3x^2)^2} dx$

Optimal result	479
Rubi [A] (verified)	479
Mathematica [A] (verified)	480
Maple [A] (verified)	480
Fricas [A] (verification not implemented)	481
Sympy [A] (verification not implemented)	481
Maxima [A] (verification not implemented)	482
Giac [A] (verification not implemented)	482
Mupad [B] (verification not implemented)	482

#### Optimal result

Integrand size = 12, antiderivative size = 42

$$\int \frac{1}{(2+5x-3x^2)^2} dx = -\frac{5-6x}{49(2+5x-3x^2)} - \frac{6}{343} \log(2-x) + \frac{6}{343} \log(1+3x)$$

[Out] 1/49\*(-5+6\*x)/(-3\*x^2+5\*x+2)-6/343\*ln(2-x)+6/343\*ln(1+3\*x)

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {628, 630, 31}

$$\int \frac{1}{(2+5x-3x^2)^2} dx = -\frac{5-6x}{49(-3x^2+5x+2)} - \frac{6}{343} \log(2-x) + \frac{6}{343} \log(3x+1)$$

[In] Int[(2 + 5\*x - 3\*x^2)^(-2), x]

[Out] -1/49\*(5 - 6\*x)/(2 + 5\*x - 3\*x^2) - (6\*Log[2 - x])/343 + (6\*Log[1 + 3\*x])/343

#### Rule 31

Int[((a\_) + (b\_.)\*(x\_))^(n\_), x\_Symbol] := Simp[Log[RemoveContent[a + b\*x, x]]/b, x] /; FreeQ[{a, b}, x]

#### Rule 628

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p+1) / ((p+1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p +

3)/((p + 1)\*(b^2 - 4\*a\*c))), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

### Rule 630

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-1), x\_Symbol] := With[{q = Rt[b^2 - 4\*a\*c, 2]}, Dist[c/q, Int[1/Simp[b/2 - q/2 + c\*x, x], x], x] - Dist[c/q, Int[1/Simp[b/2 + q/2 + c\*x, x], x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && PosQ[b^2 - 4\*a\*c] && PerfectSquareQ[b^2 - 4\*a\*c]

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{5-6x}{49(2+5x-3x^2)} + \frac{6}{49} \int \frac{1}{2+5x-3x^2} dx \\ &= -\frac{5-6x}{49(2+5x-3x^2)} - \frac{18}{343} \int \frac{1}{-1-3x} dx + \frac{18}{343} \int \frac{1}{6-3x} dx \\ &= -\frac{5-6x}{49(2+5x-3x^2)} - \frac{6}{343} \log(2-x) + \frac{6}{343} \log(1+3x) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{1}{(2+5x-3x^2)^2} dx = \frac{5-6x}{49(-2-5x+3x^2)} - \frac{6}{343} \log(2-x) + \frac{6}{343} \log(1+3x)$$

[In] Integrate[(2 + 5\*x - 3\*x^2)^(-2), x]

[Out] (5 - 6\*x)/(49\*(-2 - 5\*x + 3\*x^2)) - (6\*Log[2 - x])/343 + (6\*Log[1 + 3\*x])/343

### Maple [A] (verified)

Time = 2.08 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76



method	result	size
default	$-\frac{1}{49(-2+x)} - \frac{6 \ln(-2+x)}{343} - \frac{3}{49(3x+1)} + \frac{6 \ln(3x+1)}{343}$	32
risch	$\frac{-\frac{2x}{49} + \frac{5}{147}}{x^2 - \frac{5}{3}x - \frac{2}{3}} - \frac{6 \ln(-2+x)}{343} + \frac{6 \ln(3x+1)}{343}$	32
norman	$\frac{\frac{15}{98}x^2 - \frac{37}{98}x}{3x^2 - 5x - 2} - \frac{6 \ln(-2+x)}{343} + \frac{6 \ln(3x+1)}{343}$	38
parallelrisch	$-\frac{36 \ln(-2+x)x^2 - 36 \ln(x + \frac{1}{3})x^2 - 60 \ln(-2+x)x + 60 \ln(x + \frac{1}{3})x - 105x^2 - 24 \ln(-2+x) + 24 \ln(x + \frac{1}{3}) + 259x}{686(3x^2 - 5x - 2)}$	68

[In] `int(1/(-3*x^2+5*x+2)^2,x,method=_RETURNVERBOSE)`

[Out]  $-1/49/(-2+x) - 6/343 \cdot \ln(-2+x) - 3/49/(3x+1) + 6/343 \cdot \ln(3x+1)$

### Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.26

$$\int \frac{1}{(2+5x-3x^2)^2} dx = \frac{6(3x^2-5x-2) \log(3x+1) - 6(3x^2-5x-2) \log(x-2) - 42x + 35}{343(3x^2-5x-2)}$$

[In] `integrate(1/(-3*x^2+5*x+2)^2,x, algorithm="fricas")`

[Out]  $1/343 \cdot (6 \cdot (3x^2 - 5x - 2) \cdot \log(3x + 1) - 6 \cdot (3x^2 - 5x - 2) \cdot \log(x - 2) - 42x + 35) / (3x^2 - 5x - 2)$

### Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.76

$$\int \frac{1}{(2+5x-3x^2)^2} dx = \frac{5-6x}{147x^2-245x-98} - \frac{6 \log(x-2)}{343} + \frac{6 \log(x+\frac{1}{3})}{343}$$

[In] `integrate(1/(-3*x**2+5*x+2)**2,x)`

[Out]  $(5 - 6x)/(147x^2 - 245x - 98) - 6 \cdot \log(x - 2)/343 + 6 \cdot \log(x + 1/3)/343$

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int \frac{1}{(2 + 5x - 3x^2)^2} dx = -\frac{6x - 5}{49(3x^2 - 5x - 2)} + \frac{6}{343} \log(3x + 1) - \frac{6}{343} \log(x - 2)$$

[In] integrate(1/(-3\*x^2+5\*x+2)^2,x, algorithm="maxima")

[Out] -1/49\*(6\*x - 5)/(3\*x^2 - 5\*x - 2) + 6/343\*log(3\*x + 1) - 6/343\*log(x - 2)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{1}{(2 + 5x - 3x^2)^2} dx = -\frac{6x - 5}{49(3x^2 - 5x - 2)} + \frac{6}{343} \log(|3x + 1|) - \frac{6}{343} \log(|x - 2|)$$

[In] integrate(1/(-3\*x^2+5\*x+2)^2,x, algorithm="giac")

[Out] -1/49\*(6\*x - 5)/(3\*x^2 - 5\*x - 2) + 6/343\*log(abs(3\*x + 1)) - 6/343\*log(abs(x - 2))

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int \frac{1}{(2 + 5x - 3x^2)^2} dx = \frac{6 \ln\left(\frac{3x+1}{x-2}\right)}{343} + \frac{2\left(3x - \frac{5}{2}\right)}{49(-3x^2 + 5x + 2)}$$

[In] int(1/(5\*x - 3\*x^2 + 2)^2,x)

[Out] (6\*log((3\*x + 1)/(x - 2)))/343 + (2\*(3\*x - 5/2))/(49\*(5\*x - 3\*x^2 + 2))

### 3.95 $\int \frac{1}{(a+cx+bx^2)^2} dx$

Optimal result	483
Rubi [A] (verified)	483
Mathematica [A] (verified)	484
Maple [A] (verified)	485
Fricas [B] (verification not implemented)	485
Sympy [B] (verification not implemented)	486
Maxima [F(-2)]	486
Giac [A] (verification not implemented)	487
Mupad [B] (verification not implemented)	487

#### Optimal result

Integrand size = 12, antiderivative size = 71

$$\int \frac{1}{(a+cx+bx^2)^2} dx = \frac{c+2bx}{(4ab-c^2)(a+cx+bx^2)} + \frac{4b \arctan\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{(4ab-c^2)^{3/2}}$$

[Out]  $(2*b*x+c)/(4*a*b-c^2)/(b*x^2+c*x+a)+4*b*\arctan((2*b*x+c)/(4*a*b-c^2)^{(1/2)})/(4*a*b-c^2)^{(3/2)}$

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$ , Rules used = {628, 632, 210}

$$\int \frac{1}{(a+cx+bx^2)^2} dx = \frac{4b \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{(4ab-c^2)^{3/2}} + \frac{2bx+c}{(4ab-c^2)(a+bx^2+cx)}$$

[In]  $\text{Int}[(a + c*x + b*x^2)^{-2}, x]$

[Out]  $(c + 2*b*x)/((4*a*b - c^2)*(a + c*x + b*x^2)) + (4*b*\text{ArcTan}[(c + 2*b*x)/\text{Sqrt}[4*a*b - c^2]])/(4*a*b - c^2)^{(3/2)}$

#### Rule 210

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

## Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

## Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

## Rubi steps

$$\begin{aligned} \text{integral} &= \frac{c + 2bx}{(4ab - c^2)(a + cx + bx^2)} + \frac{(2b) \int \frac{1}{a+cx+bx^2} dx}{4ab - c^2} \\ &= \frac{c + 2bx}{(4ab - c^2)(a + cx + bx^2)} - \frac{(4b) \text{Subst}\left(\int \frac{1}{-4ab+c^2-x^2} dx, x, c + 2bx\right)}{4ab - c^2} \\ &= \frac{c + 2bx}{(4ab - c^2)(a + cx + bx^2)} + \frac{4b \tan^{-1}\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{(4ab - c^2)^{3/2}} \end{aligned}$$

## Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.99

$$\int \frac{1}{(a + cx + bx^2)^2} dx = \frac{c + 2bx}{(4ab - c^2)(a + x(c + bx))} + \frac{4b \arctan\left(\frac{c+2bx}{\sqrt{4ab-c^2}}\right)}{(4ab - c^2)^{3/2}}$$

```
[In] Integrate[(a + c*x + b*x^2)^(-2), x]
```

```
[Out] (c + 2*b*x)/((4*a*b - c^2)*(a + x*(c + b*x))) + (4*b*ArcTan[(c + 2*b*x)/Sqr
t[4*a*b - c^2]])/(4*a*b - c^2)^(3/2)
```

**Maple [A] (verified)**

Time = 2.22 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.96

method	result	size
default	$\frac{2bx+c}{(4ab-c^2)(bx^2+cx+a)} + \frac{4b \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{(4ab-c^2)^{\frac{3}{2}}}$	68
risch	$\frac{\frac{2bx}{4ab-c^2} + \frac{c}{4ab-c^2}}{bx^2+cx+a} + \frac{2b \ln\left(\frac{(-8ab^2+2bc^2)x+(-4ab+c^2)^{\frac{3}{2}}-4abc+c^3}{(-4ab+c^2)^{\frac{3}{2}}}\right)}{(-4ab+c^2)^{\frac{3}{2}}} - \frac{2b \ln\left(\frac{(8ab^2-2bc^2)x+(-4ab+c^2)^{\frac{3}{2}}+4abc-c^3}{(-4ab+c^2)^{\frac{3}{2}}}\right)}{(-4ab+c^2)^{\frac{3}{2}}}$	14

[In] int(1/(b\*x^2+c\*x+a)^2,x,method=\_RETURNVERBOSE)

[Out] (2\*b\*x+c)/(4\*a\*b-c^2)/(b\*x^2+c\*x+a)+4\*b\*arctan((2\*b\*x+c)/(4\*a\*b-c^2)^(1/2))/  
(4\*a\*b-c^2)^(3/2)**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(67) = 134.

Time = 0.31 (sec) , antiderivative size = 334, normalized size of antiderivative = 4.70

$$\int \frac{1}{(a+cx+bx^2)^2} dx = \frac{\left[4abc - c^3 + 2(b^2x^2 + bcx + ab)\sqrt{-4ab + c^2} \log\left(\frac{2b^2x^2 + 2bcx - 2ab + c^2 + \sqrt{-4ab + c^2}(2bx + c)}{bx^2 + cx + a}\right) + 2(4ab^2 - bc^2)x\right]}{16a^3b^2 - 8a^2bc^2 + ac^4 + (16a^2b^3 - 8ab^2c^2 + bc^4)x^2 + (16a^2b^2c - 8abc^3 + c^5)x}$$

[In] integrate(1/(b\*x^2+c\*x+a)^2,x, algorithm="fricas")

[Out] [(4\*a\*b\*c - c^3 + 2\*(b^2\*x^2 + b\*c\*x + a\*b)\*sqrt(-4\*a\*b + c^2)\*log((2\*b^2\*x^2 + 2\*b\*c\*x - 2\*a\*b + c^2 + sqrt(-4\*a\*b + c^2)\*(2\*b\*x + c))/(b\*x^2 + c\*x + a)) + 2\*(4\*a\*b^2 - b\*c^2)\*x)/(16\*a^3\*b^2 - 8\*a^2\*b\*c^2 + a\*c^4 + (16\*a^2\*b^3 - 8\*a\*b^2\*c^2 + b\*c^4)\*x^2 + (16\*a^2\*b^2\*c - 8\*a\*b\*c^3 + c^5)\*x), (4\*a\*b\*c - c^3 - 4\*(b^2\*x^2 + b\*c\*x + a\*b)\*sqrt(4\*a\*b - c^2)\*arctan(-(2\*b\*x + c)/sqrt(4\*a\*b - c^2)) + 2\*(4\*a\*b^2 - b\*c^2)\*x)/(16\*a^3\*b^2 - 8\*a^2\*b\*c^2 + a\*c^4 + (16\*a^2\*b^3 - 8\*a\*b^2\*c^2 + b\*c^4)\*x^2 + (16\*a^2\*b^2\*c - 8\*a\*b\*c^3 + c^5)\*x]]

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 265 vs. 2(60) = 120.

Time = 0.31 (sec) , antiderivative size = 265, normalized size of antiderivative = 3.73

$$\int \frac{1}{(a + cx + bx^2)^2} dx =$$

$$-2b\sqrt{-\frac{1}{(4ab - c^2)^3}} \log\left(x + \frac{-32a^2b^3\sqrt{-\frac{1}{(4ab - c^2)^3}} + 16ab^2c^2\sqrt{-\frac{1}{(4ab - c^2)^3}} - 2bc^4\sqrt{-\frac{1}{(4ab - c^2)^3}} + 2bc}{4b^2}\right)$$

$$+ 2b\sqrt{-\frac{1}{(4ab - c^2)^3}} \log\left(x + \frac{32a^2b^3\sqrt{-\frac{1}{(4ab - c^2)^3}} - 16ab^2c^2\sqrt{-\frac{1}{(4ab - c^2)^3}} + 2bc^4\sqrt{-\frac{1}{(4ab - c^2)^3}} + 2bc}{4b^2}\right)$$

$$+ \frac{2bx + c}{4a^2b - ac^2 + x^2 \cdot (4ab^2 - bc^2) + x(4abc - c^3)}$$

[In] integrate(1/(b\*x\*\*2+c\*x+a)\*\*2,x)

[Out]  $-2*b*\sqrt{-1/(4*a*b - c**2)**3}*\log(x + (-32*a**2*b**3*\sqrt{-1/(4*a*b - c**2)**3} + 16*a*b**2*c**2*\sqrt{-1/(4*a*b - c**2)**3} - 2*b*c**4*\sqrt{-1/(4*a*b - c**2)**3} + 2*b*c)/(4*b**2)) + 2*b*\sqrt{-1/(4*a*b - c**2)**3}*\log(x + (32*a**2*b**3*\sqrt{-1/(4*a*b - c**2)**3} - 16*a*b**2*c**2*\sqrt{-1/(4*a*b - c**2)**3} + 2*b*c**4*\sqrt{-1/(4*a*b - c**2)**3} + 2*b*c)/(4*b**2)) + (2*b*x + c)/(4*a**2*b - a*c**2 + x**2*(4*a*b**2 - b*c**2) + x*(4*a*b*c - c**3))$

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{1}{(a + cx + bx^2)^2} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(b\*x^2+c\*x+a)^2,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(c^2-4\*a\*b>0)', see 'assume?' for more details)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.94

$$\int \frac{1}{(a + cx + bx^2)^2} dx = \frac{4b \arctan\left(\frac{2bx+c}{\sqrt{4ab-c^2}}\right)}{(4ab-c^2)^{\frac{3}{2}}} + \frac{2bx+c}{(bx^2+cx+a)(4ab-c^2)}$$

[In] integrate(1/(b\*x^2+c\*x+a)^2,x, algorithm="giac")

[Out] 4\*b\*arctan((2\*b\*x + c)/sqrt(4\*a\*b - c^2))/(4\*a\*b - c^2)^(3/2) + (2\*b\*x + c)/((b\*x^2 + c\*x + a)\*(4\*a\*b - c^2))

**Mupad [B] (verification not implemented)**

Time = 0.16 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.68

$$\int \frac{1}{(a + cx + bx^2)^2} dx = \frac{\frac{c}{4ab-c^2} + \frac{2bx}{4ab-c^2}}{bx^2 + cx + a} - \frac{4b \operatorname{atan}\left(\frac{\left(\frac{2b(c^3-4abc)}{(4ab-c^2)^{5/2}} - \frac{4b^2x}{(4ab-c^2)^{3/2}}\right)(4ab-c^2)}{2b}\right)}{(4ab-c^2)^{3/2}}$$

[In] int(1/(a + c\*x + b\*x^2)^2,x)

[Out] (c/(4\*a\*b - c^2) + (2\*b\*x)/(4\*a\*b - c^2))/(a + c\*x + b\*x^2) - (4\*b\*atan((((2\*b\*(c^3 - 4\*a\*b\*c))/(4\*a\*b - c^2)^(5/2) - (4\*b^2\*x)/(4\*a\*b - c^2)^(3/2))\* (4\*a\*b - c^2))/(2\*b)))/(4\*a\*b - c^2)^(3/2)

### 3.96 $\int \frac{1}{(b+2ax+bx^2)^2} dx$

Optimal result	488
Rubi [A] (verified)	488
Mathematica [A] (verified)	489
Maple [A] (verified)	490
Fricas [B] (verification not implemented)	490
Sympy [B] (verification not implemented)	491
Maxima [F(-2)]	491
Giac [A] (verification not implemented)	492
Mupad [B] (verification not implemented)	492

#### Optimal result

Integrand size = 13, antiderivative size = 72

$$\int \frac{1}{(b+2ax+bx^2)^2} dx = -\frac{a+bx}{2(a^2-b^2)(b+2ax+bx^2)} + \frac{\operatorname{barctanh}\left(\frac{a+bx}{\sqrt{a^2-b^2}}\right)}{2(a^2-b^2)^{3/2}}$$

[Out] 1/2\*(-b\*x-a)/(a^2-b^2)/(b\*x^2+2\*a\*x+b)+1/2\*b\*arctanh((b\*x+a)/(a^2-b^2)^(1/2))/(a^2-b^2)^(3/2)

#### Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$ , Rules used = {628, 632, 212}

$$\int \frac{1}{(b+2ax+bx^2)^2} dx = \frac{\operatorname{barctanh}\left(\frac{a+bx}{\sqrt{a^2-b^2}}\right)}{2(a^2-b^2)^{3/2}} - \frac{a+bx}{2(a^2-b^2)(2ax+bx^2+b)}$$

[In] Int[(b + 2\*a\*x + b\*x^2)^(-2), x]

[Out] -1/2\*(a + b\*x)/((a^2 - b^2)\*(b + 2\*a\*x + b\*x^2)) + (b\*ArcTanh[(a + b\*x)/Sqrt[a^2 - b^2]])/(2\*(a^2 - b^2)^(3/2))

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])



Rule 628

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p + 1) / ((p + 1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p + 3) / ((p + 1)\*(b^2 - 4\*a\*c))), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && IntegerQ[4\*p]

Rule 632

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-1), x\_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4\*a\*c - x^2, x], x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{a + bx}{2(a^2 - b^2)(b + 2ax + bx^2)} - \frac{b \int \frac{1}{b + 2ax + bx^2} dx}{2(a^2 - b^2)} \\ &= -\frac{a + bx}{2(a^2 - b^2)(b + 2ax + bx^2)} + \frac{b \text{Subst}\left(\int \frac{1}{4(a^2 - b^2) - x^2} dx, x, 2a + 2bx\right)}{a^2 - b^2} \\ &= -\frac{a + bx}{2(a^2 - b^2)(b + 2ax + bx^2)} + \frac{b \tanh^{-1}\left(\frac{a + bx}{\sqrt{a^2 - b^2}}\right)}{2(a^2 - b^2)^{3/2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00

$$\int \frac{1}{(b + 2ax + bx^2)^2} dx = \frac{a + bx}{2(-a^2 + b^2)(b + 2ax + bx^2)} + \frac{b \arctan\left(\frac{a + bx}{\sqrt{-a^2 + b^2}}\right)}{2(-a^2 + b^2)^{3/2}}$$

[In] Integrate[(b + 2\*a\*x + b\*x^2)^(-2),x]

[Out] (a + b\*x)/(2\*(-a^2 + b^2)\*(b + 2\*a\*x + b\*x^2)) + (b\*ArcTan[(a + b\*x)/Sqrt[-a^2 + b^2]])/(2\*(-a^2 + b^2)^(3/2))

**Maple [A] (verified)**

Time = 2.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.19

method	result	size
default	$\frac{2bx+2a}{(-4a^2+4b^2)(bx^2+2ax+b)} + \frac{2b \arctan\left(\frac{2bx+2a}{2\sqrt{-a^2+b^2}}\right)}{(-4a^2+4b^2)\sqrt{-a^2+b^2}}$	86
risch	$\frac{-\frac{bx}{4(a^2-b^2)} - \frac{a}{4(a^2-b^2)}}{\frac{1}{2}bx^2+ax+\frac{1}{2}b} + \frac{b \ln\left((-a^2b+b^3)x - (a^2-b^2)^{\frac{3}{2}} - a^3+ab^2\right)}{4(a^2-b^2)^{\frac{3}{2}}} - \frac{b \ln\left((a^2b-b^3)x - (a^2-b^2)^{\frac{3}{2}} + a^3-ab^2\right)}{4(a^2-b^2)^{\frac{3}{2}}}$	150

[In] int(1/(b\*x^2+2\*a\*x+b)^2,x,method=\_RETURNVERBOSE)

[Out] (2\*b\*x+2\*a)/(-4\*a^2+4\*b^2)/(b\*x^2+2\*a\*x+b)+2\*b/(-4\*a^2+4\*b^2)/(-a^2+b^2)^(1/2)\*arctan(1/2\*(2\*b\*x+2\*a)/(-a^2+b^2)^(1/2))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(64) = 128.

Time = 0.29 (sec) , antiderivative size = 317, normalized size of antiderivative = 4.40

$$\int \frac{1}{(b+2ax+bx^2)^2} dx$$

$$= \left[ -\frac{2a^3 - 2ab^2 + (b^2x^2 + 2abx + b^2)\sqrt{a^2 - b^2} \log\left(\frac{b^2x^2 + 2abx + 2a^2 - b^2 - 2\sqrt{a^2 - b^2}(bx+a)}{bx^2 + 2ax + b}\right) + 2(a^2b - b^3)x}{4(a^4b - 2a^2b^3 + b^5 + (a^4b - 2a^2b^3 + b^5)x^2 + 2(a^5 - 2a^3b^2 + ab^4)x)}, \right.$$

$$\left. -\frac{a^3 - ab^2 - (b^2x^2 + 2abx + b^2)\sqrt{-a^2 + b^2} \arctan\left(-\frac{\sqrt{-a^2 + b^2}(bx+a)}{a^2 - b^2}\right) + (a^2b - b^3)x}{2(a^4b - 2a^2b^3 + b^5 + (a^4b - 2a^2b^3 + b^5)x^2 + 2(a^5 - 2a^3b^2 + ab^4)x)} \right]$$

[In] integrate(1/(b\*x^2+2\*a\*x+b)^2,x, algorithm="fricas")

```
[Out] [-1/4*(2*a^3 - 2*a*b^2 + (b^2*x^2 + 2*a*b*x + b^2)*sqrt(a^2 - b^2)*log((b^2*x^2 + 2*a*b*x + 2*a^2 - b^2 - 2*sqrt(a^2 - b^2)*(b*x + a))/(b*x^2 + 2*a*x + b)) + 2*(a^2*b - b^3)*x)/(a^4*b - 2*a^2*b^3 + b^5 + (a^4*b - 2*a^2*b^3 + b^5)*x^2 + 2*(a^5 - 2*a^3*b^2 + a*b^4)*x), -1/2*(a^3 - a*b^2 - (b^2*x^2 + 2*a*b*x + b^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*x + a)/(a^2 - b^2)) + (a^2*b - b^3)*x)/(a^4*b - 2*a^2*b^3 + b^5 + (a^4*b - 2*a^2*b^3 + b^5)*x^2 + 2*(a^5 - 2*a^3*b^2 + a*b^4)*x)]
```

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 230 vs. 2(58) = 116.

Time = 0.31 (sec) , antiderivative size = 230, normalized size of antiderivative = 3.19

$$\int \frac{1}{(b + 2ax + bx^2)^2} dx$$

$$= -\frac{b\sqrt{\frac{1}{(a-b)^3(a+b)^3}} \log\left(x + \frac{-a^4b\sqrt{\frac{1}{(a-b)^3(a+b)^3}} + 2a^2b^3\sqrt{\frac{1}{(a-b)^3(a+b)^3}} + ab - b^5\sqrt{\frac{1}{(a-b)^3(a+b)^3}}}{b^2}\right)}{4}$$

$$+ \frac{b\sqrt{\frac{1}{(a-b)^3(a+b)^3}} \log\left(x + \frac{a^4b\sqrt{\frac{1}{(a-b)^3(a+b)^3}} - 2a^2b^3\sqrt{\frac{1}{(a-b)^3(a+b)^3}} + ab + b^5\sqrt{\frac{1}{(a-b)^3(a+b)^3}}}{b^2}\right)}{4}$$

$$+ \frac{-a - bx}{2a^2b - 2b^3 + x^2 \cdot (2a^2b - 2b^3) + x(4a^3 - 4ab^2)}$$

[In] integrate(1/(b\*x\*\*2+2\*a\*x+b)\*\*2,x)

[Out] -b\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3))\*log(x + (-a\*\*4\*b\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)) + 2\*a\*\*2\*b\*\*3\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)) + a\*b - b\*\*5\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)))/b\*\*2)/4 + b\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3))\*log(x + (a\*\*4\*b\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)) - 2\*a\*\*2\*b\*\*3\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)) + a\*b + b\*\*5\*sqrt(1/((a - b)\*\*3\*(a + b)\*\*3)))/b\*\*2)/4 + (-a - b\*x)/(2\*a\*\*2\*b - 2\*b\*\*3 + x\*\*2\*(2\*a\*\*2\*b - 2\*b\*\*3) + x\*(4\*a\*\*3 - 4\*a\*b\*\*2))

**Maxima [F(-2)]**

Exception generated.

$$\int \frac{1}{(b + 2ax + bx^2)^2} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(b\*x^2+2\*a\*x+b)^2,x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(4\*a^2-4\*b^2>0)', see 'assume?' for more de

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.04

$$\int \frac{1}{(b + 2ax + bx^2)^2} dx = -\frac{b \arctan\left(\frac{bx+a}{\sqrt{-a^2+b^2}}\right)}{2(a^2-b^2)\sqrt{-a^2+b^2}} - \frac{bx+a}{2(bx^2+2ax+b)(a^2-b^2)}$$

[In] integrate(1/(b\*x^2+2\*a\*x+b)^2,x, algorithm="giac")

[Out] -1/2\*b\*arctan((b\*x + a)/sqrt(-a^2 + b^2))/((a^2 - b^2)\*sqrt(-a^2 + b^2)) - 1/2\*(b\*x + a)/((b\*x^2 + 2\*a\*x + b)\*(a^2 - b^2))

**Mupad [B] (verification not implemented)**

Time = 9.15 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.49

$$\int \frac{1}{(b + 2ax + bx^2)^2} dx = -\frac{\frac{a}{2(a^2-b^2)} + \frac{bx}{2(a^2-b^2)}}{bx^2 + 2ax + b} + \frac{b \operatorname{atan}\left(\frac{-a^3 \operatorname{li}-\operatorname{li} x a^2 b+a b^2 \operatorname{li}+\operatorname{li} x b^3}{(a+b)^{3/2}(a-b)^{3/2}}\right)}{2(a+b)^{3/2}(a-b)^{3/2}} \operatorname{li}$$

[In] int(1/(b + 2\*a\*x + b\*x^2)^2,x)

[Out] (b\*atan((a\*b^2\*li + b^3\*x\*li - a^3\*li - a^2\*b\*x\*li)/((a + b)^(3/2)\*(a - b)^(3/2)))\*li)/(2\*(a + b)^(3/2)\*(a - b)^(3/2)) - (a/(2\*(a^2 - b^2)) + (b\*x)/(2\*(a^2 - b^2)))/(b + 2\*a\*x + b\*x^2)

$$3.97 \quad \int \frac{1}{(b+2ax-bx^2)^2} dx$$

Optimal result . . . . .	493
Rubi [A] (verified) . . . . .	493
Mathematica [A] (verified) . . . . .	494
Maple [A] (verified) . . . . .	495
Fricas [B] (verification not implemented) . . . . .	495
Sympy [B] (verification not implemented) . . . . .	495
Maxima [A] (verification not implemented) . . . . .	496
Giac [A] (verification not implemented) . . . . .	497
Mupad [B] (verification not implemented) . . . . .	497

### Optimal result

Integrand size = 14, antiderivative size = 69

$$\int \frac{1}{(b+2ax-bx^2)^2} dx = -\frac{a-bx}{2(a^2+b^2)(b+2ax-bx^2)} - \frac{\operatorname{barctanh}\left(\frac{a-bx}{\sqrt{a^2+b^2}}\right)}{2(a^2+b^2)^{3/2}}$$

[Out]  $1/2*(b*x-a)/(a^2+b^2)/(-b*x^2+2*a*x+b)-1/2*b*\operatorname{arctanh}((-b*x+a)/(a^2+b^2)^{(1/2)))/(a^2+b^2)^{(3/2)}$

### Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {628, 632, 212}

$$\int \frac{1}{(b+2ax-bx^2)^2} dx = -\frac{\operatorname{barctanh}\left(\frac{a-bx}{\sqrt{a^2+b^2}}\right)}{2(a^2+b^2)^{3/2}} - \frac{a-bx}{2(a^2+b^2)(2ax-bx^2+b)}$$

[In]  $\operatorname{Int}[(b+2*a*x-b*x^2)^{-2}, x]$

[Out]  $-1/2*(a-b*x)/((a^2+b^2)*(b+2*a*x-b*x^2)) - (b*\operatorname{ArcTanh}[(a-b*x)/\operatorname{Sqrt}[a^2+b^2]])/(2*(a^2+b^2)^{(3/2)})$

#### Rule 212

$\operatorname{Int}[(a_0 + (b_0)*(x_0)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \operatorname{LtQ}[b, 0])$

Rule 628

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] - Dist[2*c*((2*p +
3)/((p + 1)*(b^2 - 4*a*c))), Int[(a + b*x + c*x^2)^(p + 1), x], x] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[p, -3/2] && Int
egerQ[4*p]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{a - bx}{2(a^2 + b^2)(b + 2ax - bx^2)} + \frac{b \int \frac{1}{b + 2ax - bx^2} dx}{2(a^2 + b^2)} \\
&= -\frac{a - bx}{2(a^2 + b^2)(b + 2ax - bx^2)} - \frac{b \text{Subst}\left(\int \frac{1}{4(a^2 + b^2) - x^2} dx, x, 2a - 2bx\right)}{a^2 + b^2} \\
&= -\frac{a - bx}{2(a^2 + b^2)(b + 2ax - bx^2)} - \frac{b \tanh^{-1}\left(\frac{a - bx}{\sqrt{a^2 + b^2}}\right)}{2(a^2 + b^2)^{3/2}}
\end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.13

$$\int \frac{1}{(b + 2ax - bx^2)^2} dx = \frac{-a + bx}{b + 2ax - bx^2} - \frac{b \arctan\left(\frac{-a + bx}{\sqrt{-a^2 - b^2}}\right)}{2(a^2 + b^2)}$$

```
[In] Integrate[(b + 2*a*x - b*x^2)^(-2), x]
```

```
[Out] ((-a + b*x)/(b + 2*a*x - b*x^2) - (b*ArcTan[(-a + b*x)/Sqrt[-a^2 - b^2]])/S
qrt[-a^2 - b^2])/(2*(a^2 + b^2))
```

**Maple [A] (verified)**

Time = 2.60 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.20

method	result	size
default	$\frac{-2bx+2a}{(-4a^2-4b^2)(-bx^2+2ax+b)} + \frac{2b \operatorname{arctanh}\left(\frac{-2bx+2a}{2\sqrt{a^2+b^2}}\right)}{(-4a^2-4b^2)\sqrt{a^2+b^2}}$	83
risch	$\frac{\frac{bx}{4a^2+4b^2} - \frac{a}{4(a^2+b^2)}}{-\frac{1}{2}bx^2+ax+\frac{1}{2}b} + \frac{b \ln\left((a^2b+b^3)x+(a^2+b^2)^{\frac{3}{2}}-a^3-ab^2\right)}{4(a^2+b^2)^{\frac{3}{2}}} - \frac{b \ln\left((-a^2b-b^3)x+(a^2+b^2)^{\frac{3}{2}}+a^3+ab^2\right)}{4(a^2+b^2)^{\frac{3}{2}}}$	134

[In] int(1/(-b\*x^2+2\*a\*x+b)^2,x,method=\_RETURNVERBOSE)

[Out] (-2\*b\*x+2\*a)/(-4\*a^2-4\*b^2)/(-b\*x^2+2\*a\*x+b)+2\*b/(-4\*a^2-4\*b^2)/(a^2+b^2)^(1/2)\*arctanh(1/2\*(-2\*b\*x+2\*a)/(a^2+b^2)^(1/2))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 171 vs. 2(65) = 130.

Time = 0.28 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.48

$$\int \frac{1}{(b+2ax-bx^2)^2} dx = \frac{2a^3 + 2ab^2 + (b^2x^2 - 2abx - b^2)\sqrt{a^2+b^2} \log\left(\frac{b^2x^2 - 2abx + 2a^2 + b^2 + 2\sqrt{a^2+b^2}(bx-a)}{bx^2 - 2ax - b}\right) - 2(a^2b + b^3)x}{4(a^4b + 2a^2b^3 + b^5 - (a^4b + 2a^2b^3 + b^5)x^2 + 2(a^5 + 2a^3b^2 + ab^4)x)}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b)^2,x, algorithm="fricas")

[Out] -1/4\*(2\*a^3 + 2\*a\*b^2 + (b^2\*x^2 - 2\*a\*b\*x - b^2)\*sqrt(a^2 + b^2)\*log((b^2\*x^2 - 2\*a\*b\*x + 2\*a^2 + b^2 + 2\*sqrt(a^2 + b^2)\*(b\*x - a))/(b\*x^2 - 2\*a\*x - b)) - 2\*(a^2\*b + b^3)\*x)/(a^4\*b + 2\*a^2\*b^3 + b^5 - (a^4\*b + 2\*a^2\*b^3 + b^5)\*x^2 + 2\*(a^5 + 2\*a^3\*b^2 + a\*b^4)\*x)

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 218 vs. 2(56) = 112.

Time = 0.29 (sec) , antiderivative size = 218, normalized size of antiderivative = 3.16

$$\int \frac{1}{(b + 2ax - bx^2)^2} dx = -\frac{b\sqrt{\frac{1}{(a^2+b^2)^3}} \log\left(x + \frac{-a^4b\sqrt{\frac{1}{(a^2+b^2)^3}} - 2a^2b^3\sqrt{\frac{1}{(a^2+b^2)^3}} - ab - b^5\sqrt{\frac{1}{(a^2+b^2)^3}}}{b^2}\right)}{4} + \frac{b\sqrt{\frac{1}{(a^2+b^2)^3}} \log\left(x + \frac{a^4b\sqrt{\frac{1}{(a^2+b^2)^3}} + 2a^2b^3\sqrt{\frac{1}{(a^2+b^2)^3}} - ab + b^5\sqrt{\frac{1}{(a^2+b^2)^3}}}{b^2}\right)}{4} + \frac{a - bx}{-2a^2b - 2b^3 + x^2 \cdot (2a^2b + 2b^3) + x(-4a^3 - 4ab^2)}$$

[In] integrate(1/(-b\*x\*\*2+2\*a\*x+b)\*\*2,x)

[Out] -b\*sqrt((a\*\*2 + b\*\*2)\*\*(-3))\*log(x + (-a\*\*4\*b\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)) - 2\*a\*\*2\*b\*\*3\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)) - a\*b - b\*\*5\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)))/b\*\*2)/4 + b\*sqrt((a\*\*2 + b\*\*2)\*\*(-3))\*log(x + (a\*\*4\*b\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)) + 2\*a\*\*2\*b\*\*3\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)) - a\*b + b\*\*5\*sqrt((a\*\*2 + b\*\*2)\*\*(-3)))/b\*\*2)/4 + (a - b\*x)/(-2\*a\*\*2\*b - 2\*b\*\*3 + x\*\*2\*(2\*a\*\*2\*b + 2\*b\*\*3) + x\*(-4\*a\*\*3 - 4\*a\*b\*\*2))

## Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.41

$$\int \frac{1}{(b + 2ax - bx^2)^2} dx = -\frac{b \log\left(\frac{bx-a-\sqrt{a^2+b^2}}{bx-a+\sqrt{a^2+b^2}}\right)}{4(a^2+b^2)^{\frac{3}{2}}} + \frac{bx-a}{2(a^2b+b^3-(a^2b+b^3)x^2+2(a^3+ab^2)x)}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b)^2,x, algorithm="maxima")

[Out] -1/4\*b\*log((b\*x - a - sqrt(a^2 + b^2))/(b\*x - a + sqrt(a^2 + b^2)))/(a^2 + b^2)^(3/2) + 1/2\*(b\*x - a)/(a^2\*b + b^3 - (a^2\*b + b^3)\*x^2 + 2\*(a^3 + a\*b^2)\*x)



**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.30

$$\int \frac{1}{(b + 2ax - bx^2)^2} dx = -\frac{b \log\left(\frac{|2bx - 2a - 2\sqrt{a^2 + b^2}|}{|2bx - 2a + 2\sqrt{a^2 + b^2}|}\right)}{4(a^2 + b^2)^{\frac{3}{2}}} - \frac{bx - a}{2(bx^2 - 2ax - b)(a^2 + b^2)}$$

[In] integrate(1/(-b\*x^2+2\*a\*x+b)^2,x, algorithm="giac")

[Out] -1/4\*b\*log(abs(2\*b\*x - 2\*a - 2\*sqrt(a^2 + b^2))/abs(2\*b\*x - 2\*a + 2\*sqrt(a^2 + b^2)))/(a^2 + b^2)^(3/2) - 1/2\*(b\*x - a)/((b\*x^2 - 2\*a\*x - b)\*(a^2 + b^2))

**Mupad [B] (verification not implemented)**

Time = 9.12 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.45

$$\int \frac{1}{(b + 2ax - bx^2)^2} dx = -\frac{\frac{a}{2(a^2+b^2)} - \frac{bx}{2(a^2+b^2)}}{-bx^2 + 2ax + b} + \frac{b \operatorname{atan}\left(\frac{ab^2 \operatorname{li} + a^3 \operatorname{li} - bx(a^2+b^2) \operatorname{li}}{(a^2+b^2)^{3/2}}\right) \operatorname{li}}{2(a^2 + b^2)^{3/2}}$$

[In] int(1/(b + 2\*a\*x - b\*x^2)^2,x)

[Out] (b\*atan((a\*b^2\*li + a^3\*li - b\*x\*(a^2 + b^2)\*li)/(a^2 + b^2)^(3/2))\*li)/(2\*(a^2 + b^2)^(3/2)) - (a/(2\*(a^2 + b^2)) - (b\*x)/(2\*(a^2 + b^2)))/(b + 2\*a\*x - b\*x^2)

$$3.98 \quad \int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx$$

Optimal result	498
Rubi [A] (verified)	498
Mathematica [A] (verified)	499
Maple [A] (verified)	500
Fricas [A] (verification not implemented)	500
Sympy [B] (verification not implemented)	500
Maxima [F(-2)]	501
Giac [A] (verification not implemented)	502
Mupad [B] (verification not implemented)	502

### Optimal result

Integrand size = 40, antiderivative size = 62

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx =$$

$$-\left(\frac{a}{b}\right)^{-1/n} \arctan\left(\cot\left(\frac{\pi - 2k\pi}{n}\right) - \left(\frac{a}{b}\right)^{-1/n} x \csc\left(\frac{\pi - 2k\pi}{n}\right)\right) \csc\left(\frac{\pi - 2k\pi}{n}\right)$$

[Out]  $\arctan(-\cot((-2*\text{Pi}*k+\text{Pi})/n)+x*\csc((-2*\text{Pi}*k+\text{Pi})/n)/((a/b)^{(1/n)}))*\csc((-2*\text{Pi}*k+\text{Pi})/n)/((a/b)^{(1/n)})$

### Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.050$ , Rules used = {632, 210}

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx =$$

$$-\left(\frac{a}{b}\right)^{-1/n} \csc\left(\frac{\pi - 2\pi k}{n}\right) \arctan\left(\cot\left(\frac{\pi - 2\pi k}{n}\right) - x\left(\frac{a}{b}\right)^{-1/n} \csc\left(\frac{\pi - 2\pi k}{n}\right)\right)$$

[In]  $\text{Int}[(a/b)^{(2/n)} + x^2 - 2*(a/b)^{1/n}*x*\text{Cos}[(\text{Pi} - 2*k*\text{Pi})/n]]^{(-1)}, x]$

[Out]  $-((\text{ArcTan}[\text{Cot}[(\text{Pi} - 2*k*\text{Pi})/n] - (x*\text{Csc}[(\text{Pi} - 2*k*\text{Pi})/n])/(a/b)^{1/n}])*\text{Csc}[(\text{Pi} - 2*k*\text{Pi})/n])/(a/b)^{1/n})$

Rule 210

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

### Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= - \left( 2 \text{Subst} \left( \int \frac{1}{-x^2 - 4 \left(\frac{a}{b}\right)^{2/n} \left(1 - \cos^2 \left(\frac{\pi - 2k\pi}{n}\right)\right)} dx, x, 2x \right. \right. \\ &\quad \left. \left. - 2 \left(\frac{a}{b}\right)^{\frac{1}{n}} \cos \left(\frac{\pi - 2k\pi}{n}\right) \right) \right) \\ &= - \left(\frac{a}{b}\right)^{-1/n} \tan^{-1} \left( \cot \left(\frac{\pi - 2k\pi}{n}\right) - \left(\frac{a}{b}\right)^{-1/n} x \csc \left(\frac{\pi - 2k\pi}{n}\right) \right) \csc \left(\frac{\pi - 2k\pi}{n}\right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2 \left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos \left(\frac{\pi - 2k\pi}{n}\right)} dx = \\ - \left(\frac{a}{b}\right)^{-1/n} \arctan \left( \cot \left(\frac{\pi - 2k\pi}{n}\right) - \left(\frac{a}{b}\right)^{-1/n} x \csc \left(\frac{\pi - 2k\pi}{n}\right) \right) \csc \left(\frac{\pi - 2k\pi}{n}\right) \end{aligned}$$

```
[In] Integrate[((a/b)^(2/n) + x^2 - 2*(a/b)^n^(-1)*x*Cos[(Pi - 2*k*Pi)/n])^(-1), x]
```

```
[Out] -((ArcTan[Cot[(Pi - 2*k*Pi)/n] - (x*Csc[(Pi - 2*k*Pi)/n])/(a/b)^n^(-1)]*Csc[(Pi - 2*k*Pi)/n])/(a/b)^n^(-1))
```

**Maple [A] (verified)**

Time = 3.14 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.79

method	result	size
default	$\frac{\arctan\left(\frac{2x-2\left(\frac{a}{b}\right)^{\frac{1}{n}}\cos\left(\frac{\pi(2k-1)}{n}\right)}{2\sqrt{-\left(\frac{a}{b}\right)^{\frac{2}{n}}\left(\cos^2\left(\frac{\pi(2k-1)}{n}\right)\right)+\left(\frac{a}{b}\right)^{\frac{2}{n}}}}\right)}{\sqrt{-\left(\frac{a}{b}\right)^{\frac{2}{n}}\left(\cos^2\left(\frac{\pi(2k-1)}{n}\right)\right)+\left(\frac{a}{b}\right)^{\frac{2}{n}}}}$	111
risch	Expression too large to display	1343

[In] `int(1/((a/b)^(2/n)+x^2-2*(a/b)^(1/n)*x*cos((-2*Pi*k+Pi)/n)),x,method=_RETURNVERBOSE)`

[Out]  $1/(-((a/b)^{(1/n)})^2*\cos(\text{Pi}*(2*k-1)/n)^2+(a/b)^{(2/n)})^{(1/2)}*\arctan(1/2*(2*x-2*(a/b)^{(1/n)}*\cos(\text{Pi}*(2*k-1)/n))/(-((a/b)^{(1/n)})^2*\cos(\text{Pi}*(2*k-1)/n)^2+(a/b)^{(2/n)})^{(1/2)})$

**Fricas [A] (verification not implemented)**

none

Time = 0.45 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.44

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi-2k\pi}{n}\right)} dx = -\frac{\arctan\left(\frac{\left(\frac{a}{b}\right)^{\left(\frac{1}{n}\right)} \cos\left(\frac{2\pi k - \pi}{n}\right) - x}{\left(\frac{a}{b}\right)^{\left(\frac{1}{n}\right)} \sin\left(\frac{2\pi k - \pi}{n}\right)}\right)}{\left(\frac{a}{b}\right)^{\left(\frac{1}{n}\right)} \sin\left(\frac{2\pi k - \pi}{n}\right)}$$

[In] `integrate(1/((a/b)^(2/n)+x^2-2*(a/b)^(1/n)*x*cos((-2*pi*k+pi)/n)),x, algorithm="fricas")`

[Out]  $-\arctan\left(\frac{\left(\frac{a}{b}\right)^{(1/n)}*\cos(2*\text{pi}*k/n - \text{pi}/n) - x}{\left(\frac{a}{b}\right)^{(1/n)}*\sin(2*\text{pi}*k/n - \text{pi}/n)}\right)/\left(\frac{a}{b}\right)^{(1/n)}*\sin(2*\text{pi}*k/n - \text{pi}/n)$

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(46) = 92.

Time = 0.49 (sec) , antiderivative size = 212, normalized size of antiderivative = 3.42

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi-2k\pi}{n}\right)} dx =$$

$$\frac{\sqrt{\frac{\left(\frac{a}{b}\right)^{-\frac{2}{n}}}{\cos^2\left(\frac{\pi(2k-1)}{n}\right)-1}} \log\left(x - \left(\frac{a}{b}\right)^{\frac{1}{n}} \cos\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right) - \frac{\sqrt{\frac{\left(\frac{a}{b}\right)^{-\frac{2}{n}}}{\cos^2\left(\frac{\pi(2k-1)}{n}\right)-1}} \left(-2\left(\frac{a}{b}\right)^{\frac{2}{n}} \cos^2\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right) + 2\left(\frac{a}{b}\right)^{\frac{2}{n}}\right)}{2}\right)}{2}$$

$$+ \frac{\sqrt{\frac{\left(\frac{a}{b}\right)^{-\frac{2}{n}}}{\cos^2\left(\frac{\pi(2k-1)}{n}\right)-1}} \log\left(x - \left(\frac{a}{b}\right)^{\frac{1}{n}} \cos\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right) + \frac{\sqrt{\frac{\left(\frac{a}{b}\right)^{-\frac{2}{n}}}{\cos^2\left(\frac{\pi(2k-1)}{n}\right)-1}} \left(-2\left(\frac{a}{b}\right)^{\frac{2}{n}} \cos^2\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right) + 2\left(\frac{a}{b}\right)^{\frac{2}{n}}\right)}{2}\right)}{2}$$

[In] integrate(1/((a/b)\*\*(2/n)+x\*\*2-2\*(a/b)\*\*(1/n)\*x\*cos((-2\*pi\*k+pi)/n)),x)

[Out] -sqrt(1/((a/b)\*\*(2/n)\*(cos(pi\*(2\*k - 1)/n)\*\*2 - 1)))\*log(x - (a/b)\*\*(1/n)\*cos(2\*pi\*k/n - pi/n) - sqrt(1/((a/b)\*\*(2/n)\*(cos(pi\*(2\*k - 1)/n)\*\*2 - 1)))\*(-2\*(a/b)\*\*(2/n)\*cos(2\*pi\*k/n - pi/n)\*\*2 + 2\*(a/b)\*\*(2/n))/2)/2 + sqrt(1/((a/b)\*\*(2/n)\*(cos(pi\*(2\*k - 1)/n)\*\*2 - 1)))\*log(x - (a/b)\*\*(1/n)\*cos(2\*pi\*k/n - pi/n) + sqrt(1/((a/b)\*\*(2/n)\*(cos(pi\*(2\*k - 1)/n)\*\*2 - 1)))\*(-2\*(a/b)\*\*(2/n)\*cos(2\*pi\*k/n - pi/n)\*\*2 + 2\*(a/b)\*\*(2/n))/2)/2

## Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{\frac{1}{n}} x \cos\left(\frac{\pi-2k\pi}{n}\right)} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/((a/b)^(2/n)+x^2-2\*(a/b)^(1/n)\*x\*cos((-2\*pi\*k+pi)/n)),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation \*may\* help (example of legal syntax is 'assume(1>0)', see 'assume?' for more details)Is 1

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.61

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{1/n} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx = \frac{\arctan\left(\frac{\left(\frac{a}{b}\right)^{1/n} \cos\left(-\frac{2\pi k}{n} + \frac{\pi}{n}\right) - x}{\sqrt{-\cos\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right)^2 + 1\left(\frac{a}{b}\right)^{1/n}}}\right)}{\sqrt{-\cos\left(\frac{2\pi k}{n} - \frac{\pi}{n}\right)^2 + 1\left(\frac{a}{b}\right)^{1/n}}}$$

[In] integrate(1/((a/b)^(2/n)+x^2-2\*(a/b)^(1/n)\*x\*cos((-2\*pi\*k+pi)/n)),x, algorithm="giac")

[Out] arctan(-((a/b)^(1/n)\*cos(-2\*pi\*k/n + pi/n) - x)/(sqrt(-cos(2\*pi\*k/n - pi/n)^2 + 1)\*(a/b)^(1/n)))/sqrt(-cos(2\*pi\*k/n - pi/n)^2 + 1)\*(a/b)^(1/n))

**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.77

$$\int \frac{1}{\left(\frac{a}{b}\right)^{2/n} + x^2 - 2\left(\frac{a}{b}\right)^{1/n} x \cos\left(\frac{\pi - 2k\pi}{n}\right)} dx = -\frac{\operatorname{atanh}\left(\frac{x - \cos\left(\frac{\Pi(2k-1)}{n}\right)\left(\frac{a}{b}\right)^{1/n}}{\sqrt{\cos\left(\frac{\Pi(2k-1)}{n}\right) - 1}\sqrt{\cos\left(\frac{\Pi(2k-1)}{n}\right) + 1}\left(\frac{a}{b}\right)^{1/n}}}\right)}{\sqrt{\cos\left(\frac{\Pi(2k-1)}{n}\right) - 1}\sqrt{\cos\left(\frac{\Pi(2k-1)}{n}\right) + 1}\left(\frac{a}{b}\right)^{1/n}}$$

[In] int(1/((a/b)^(2/n) + x^2 - 2\*x\*cos((Pi - 2\*Pi\*k)/n)\*(a/b)^(1/n)),x)

[Out] -atanh((x - cos((Pi\*(2\*k - 1))/n)\*(a/b)^(1/n))/((cos((Pi\*(2\*k - 1))/n) - 1)^(1/2)\*(cos((Pi\*(2\*k - 1))/n) + 1)^(1/2)\*(a/b)^(1/n)))/((cos((Pi\*(2\*k - 1))/n) - 1)^(1/2)\*(cos((Pi\*(2\*k - 1))/n) + 1)^(1/2)\*(a/b)^(1/n))

$$3.99 \quad \int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx$$

Optimal result . . . . .	503
Rubi [A] (verified) . . . . .	503
Mathematica [B] (verified) . . . . .	504
Maple [A] (verified) . . . . .	504
Fricas [B] (verification not implemented) . . . . .	505
Sympy [B] (verification not implemented) . . . . .	505
Maxima [A] (verification not implemented) . . . . .	505
Giac [A] (verification not implemented) . . . . .	506
Mupad [B] (verification not implemented) . . . . .	506

### Optimal result

Integrand size = 30, antiderivative size = 33

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = \frac{2\operatorname{arctanh}\left(\frac{-\sqrt{b^2 - 4ab^3} + 2b^2x}{b}\right)}{b}$$

[Out] 2\*arctanh((2\*b^2\*x - (-4\*a\*b^3 + b^2)^(1/2))/b)/b

### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.76, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {630, 31}

$$\begin{aligned} & \int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx \\ &= \frac{\log(-\sqrt{b^2 - 4ab^3} + 2b^2x + b)}{b} - \frac{\log(\sqrt{b^2 - 4ab^3} - 2b^2x + b)}{b} \end{aligned}$$

[In] Int[(a\*b + Sqrt[b^2 - 4\*a\*b^3]\*x - b^2\*x^2)^(-1), x]

[Out] -(Log[b + Sqrt[b^2 - 4\*a\*b^3] - 2\*b^2\*x]/b) + Log[b - Sqrt[b^2 - 4\*a\*b^3] + 2\*b^2\*x]/b

#### Rule 31

Int[((a\_) + (b\_.)\*(x\_))^(n\_), x\_Symbol] := Simp[Log[RemoveContent[a + b\*x, x]]/b, x] /; FreeQ[{a, b}, x]

#### Rule 630

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[c/q, Int[1/Simp[b/2 - q/2 + c*x, x], x], x] - Dist[c/q,
Int[1/Simp[b/2 + q/2 + c*x, x], x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2
- 4*a*c, 0] && PosQ[b^2 - 4*a*c] && PerfectSquareQ[b^2 - 4*a*c]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(b \int \frac{1}{\frac{1}{2}(-b + \sqrt{b^2 - 4ab^3}) - b^2x} dx\right) + b \int \frac{1}{\frac{1}{2}(b + \sqrt{b^2 - 4ab^3}) - b^2x} dx \\ &= -\frac{\log(b + \sqrt{b^2 - 4ab^3} - 2b^2x)}{b} + \frac{\log(b - \sqrt{b^2 - 4ab^3} + 2b^2x)}{b} \end{aligned}$$

**Mathematica [B] (verified)**

Leaf count is larger than twice the leaf count of optimal. 120 vs. 2(33) = 66.

Time = 0.12 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.64

$$\begin{aligned} &\int \frac{1}{ab + \sqrt{b^2 - 4ab^3}x - b^2x^2} dx \\ &= \frac{-2\sqrt{b^2 - 4ab^3} \arctan\left(\frac{-1+2bx}{\sqrt{-1+4ab}}\right) + 2\sqrt{b^2 - 4ab^3} \arctan\left(\frac{1+2bx}{\sqrt{-1+4ab}}\right) + b\sqrt{-1+4ab}(\log(a+x+bx^2) - \log(a+x+b*x^2) - \log[a+x*(-1+b*x)])}{2b^2\sqrt{-1+4ab}} \end{aligned}$$

```
[In] Integrate[(a*b + Sqrt[b^2 - 4*a*b^3]*x - b^2*x^2)^(-1), x]
```

```
[Out] (-2*Sqrt[b^2 - 4*a*b^3]*ArcTan[(-1 + 2*b*x)/Sqrt[-1 + 4*a*b]] + 2*Sqrt[b^2
- 4*a*b^3]*ArcTan[(1 + 2*b*x)/Sqrt[-1 + 4*a*b]] + b*Sqrt[-1 + 4*a*b]*(Log[a
+ x + b*x^2] - Log[a + x*(-1 + b*x)]))/(2*b^2*Sqrt[-1 + 4*a*b])
```

**Maple [A] (verified)**

Time = 2.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.94

method	result	size
default	$-\frac{2 \operatorname{arctanh}\left(\frac{-2b^2x + \sqrt{-b^2(4ab-1)}}{b}\right)}{b}$	31

```
[In] int(1/(a*b-b^2*x^2+x*(-4*a*b^3+b^2)^(1/2)),x,method=_RETURNVERBOSE)
```

```
[Out] -2/b*arctanh((-2*b^2*x+(-b^2*(4*a*b-1))^(1/2))/b)
```



**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(31) = 62.

Time = 0.43 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.91

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = \frac{\log\left(\frac{2b^2x+b-\sqrt{-4ab^3+b^2}}{b}\right) - \log\left(\frac{2b^2x-b-\sqrt{-4ab^3+b^2}}{b}\right)}{b}$$

[In] integrate(1/(a\*b-b^2\*x^2+x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="fricas")

[Out] (log((2\*b^2\*x + b - sqrt(-4\*a\*b^3 + b^2))/b) - log((2\*b^2\*x - b - sqrt(-4\*a\*b^3 + b^2))/b))/b

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(26) = 52.

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.70

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = -\frac{\log\left(x - \frac{1}{2b} - \frac{\sqrt{-4ab^3+b^2}}{2b^2}\right) - \log\left(x + \frac{1}{2b} - \frac{\sqrt{-4ab^3+b^2}}{2b^2}\right)}{b}$$

[In] integrate(1/(a\*b-b\*\*2\*x\*\*2+x\*(-4\*a\*b\*\*3+b\*\*2)\*\*(1/2)),x)

[Out] -(log(x - 1/(2\*b) - sqrt(-4\*a\*b\*\*3 + b\*\*2)/(2\*b\*\*2)) - log(x + 1/(2\*b) - sqrt(-4\*a\*b\*\*3 + b\*\*2)/(2\*b\*\*2)))/b

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.67

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = -\frac{\log\left(\frac{2b^2x-b-\sqrt{-4ab^3+b^2}}{2b^2x+b-\sqrt{-4ab^3+b^2}}\right)}{b}$$

[In] integrate(1/(a\*b-b^2\*x^2+x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="maxima")

[Out] -log((2\*b^2\*x - b - sqrt(-4\*a\*b^3 + b^2))/(2\*b^2\*x + b - sqrt(-4\*a\*b^3 + b^2)))/b

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.70

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3}x - b^2x^2} dx = -\frac{\log\left(\frac{|2b^2x - \sqrt{-4ab+1}|b| - |b||}{|2b^2x - \sqrt{-4ab+1}|b| + |b||}\right)}{|b|}$$

[In] integrate(1/(a\*b-b^2\*x^2+x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="giac")

[Out] -log(abs(2\*b^2\*x - sqrt(-4\*a\*b + 1)\*abs(b) - abs(b))/abs(2\*b^2\*x - sqrt(-4\*a\*b + 1)\*abs(b) + abs(b)))/abs(b)

**Mupad [B] (verification not implemented)**

Time = 0.15 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.15

$$\int \frac{1}{ab + \sqrt{b^2 - 4ab^3}x - b^2x^2} dx = -\frac{2 \operatorname{atanh}\left(\frac{\sqrt{b^2 - 4ab^3}}{\sqrt{b^2}} - \frac{2b^2x}{\sqrt{b^2}}\right)}{\sqrt{b^2}}$$

[In] int(1/(a\*b + x\*(b^2 - 4\*a\*b^3)^(1/2) - b^2\*x^2),x)

[Out] -(2\*atanh((b^2 - 4\*a\*b^3)^(1/2)/(b^2)^(1/2) - (2\*b^2\*x)/(b^2)^(1/2)))/(b^2)^(1/2)

$$3.100 \quad \int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx$$

Optimal result	507
Rubi [A] (verified)	507
Mathematica [B] (verified)	508
Maple [A] (verified)	508
Fricas [B] (verification not implemented)	509
Sympy [B] (verification not implemented)	509
Maxima [A] (verification not implemented)	509
Giac [A] (verification not implemented)	510
Mupad [B] (verification not implemented)	510

### Optimal result

Integrand size = 31, antiderivative size = 31

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{b^2 - 4ab^3 + 2b^2x}}{b}\right)}{b}$$

[Out]  $2 \operatorname{arctanh}\left(\frac{(2b^2x + (-4ab^3 + b^2)^{1/2})}{b}\right)/b$

### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.87, number of steps used = 3, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.065$ , Rules used = {630, 31}

$$\begin{aligned} & \int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx \\ &= \frac{\log(\sqrt{b^2 - 4ab^3 + 2b^2x} + b)}{b} - \frac{\log(-\sqrt{b^2 - 4ab^3} - 2b^2x + b)}{b} \end{aligned}$$

[In]  $\text{Int}[(a*b - \text{Sqrt}[b^2 - 4*a*b^3]*x - b^2*x^2)^{-1}, x]$

[Out]  $-(\text{Log}[b - \text{Sqrt}[b^2 - 4*a*b^3] - 2*b^2*x]/b) + \text{Log}[b + \text{Sqrt}[b^2 - 4*a*b^3] + 2*b^2*x]/b$

#### Rule 31

$\text{Int}[(a_) + (b_)*(x_)^{-1}, x\_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}\{a, b\}, x]$

#### Rule 630

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Dist[c/q, Int[1/Simp[b/2 - q/2 + c*x, x], x], x] - Dist[c/q,
Int[1/Simp[b/2 + q/2 + c*x, x], x], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2
- 4*a*c, 0] && PosQ[b^2 - 4*a*c] && PerfectSquareQ[b^2 - 4*a*c]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(b \int \frac{1}{\frac{1}{2}(-b - \sqrt{b^2 - 4ab^3}) - b^2x} dx\right) + b \int \frac{1}{\frac{1}{2}(b - \sqrt{b^2 - 4ab^3}) - b^2x} dx \\ &= -\frac{\log(b - \sqrt{b^2 - 4ab^3} - 2b^2x)}{b} + \frac{\log(b + \sqrt{b^2 - 4ab^3} + 2b^2x)}{b} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 120 vs. 2(31) = 62.

Time = 0.10 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.87

$$\begin{aligned} &\int \frac{1}{ab - \sqrt{b^2 - 4ab^3}x - b^2x^2} dx \\ &= \frac{2\sqrt{b^2 - 4ab^3} \arctan\left(\frac{-1+2bx}{\sqrt{-1+4ab}}\right) - 2\sqrt{b^2 - 4ab^3} \arctan\left(\frac{1+2bx}{\sqrt{-1+4ab}}\right) + b\sqrt{-1+4ab}(\log(a+x+bx^2) - \log(a+x+bx^2))}{2b^2\sqrt{-1+4ab}} \end{aligned}$$

```
[In] Integrate[(a*b - Sqrt[b^2 - 4*a*b^3]*x - b^2*x^2)^(-1), x]
```

```
[Out] (2*Sqrt[b^2 - 4*a*b^3]*ArcTan[(-1 + 2*b*x)/Sqrt[-1 + 4*a*b]] - 2*Sqrt[b^2 -
4*a*b^3]*ArcTan[(1 + 2*b*x)/Sqrt[-1 + 4*a*b]] + b*Sqrt[-1 + 4*a*b]*(Log[a
+ x + b*x^2] - Log[a + x*(-1 + b*x)]))/(2*b^2*Sqrt[-1 + 4*a*b])
```

### Maple [A] (verified)

Time = 2.20 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00

method	result	size
default	$\frac{2 \operatorname{arctanh}\left(\frac{2b^2x + \sqrt{-b^2(4ab-1)}}{b}\right)}{b}$	31

```
[In] int(1/(a*b-b^2*x^2-x*(-4*a*b^3+b^2)^(1/2)),x,method=_RETURNVERBOSE)
```

```
[Out] 2/b*arctanh((2*b^2*x+(-b^2*(4*a*b-1))^(1/2))/b)
```

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 59 vs.  $2(29) = 58$ .

Time = 0.52 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.90

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = \frac{\log\left(\frac{2b^2x+b+\sqrt{-4ab^3+b^2}}{b}\right) - \log\left(\frac{2b^2x-b+\sqrt{-4ab^3+b^2}}{b}\right)}{b}$$

[In] integrate(1/(a\*b-b^2\*x^2-x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="fricas")

[Out] (log((2\*b^2\*x + b + sqrt(-4\*a\*b^3 + b^2))/b) - log((2\*b^2\*x - b + sqrt(-4\*a\*b^3 + b^2))/b))/b

**Sympy [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 56 vs.  $2(26) = 52$ .

Time = 0.15 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.81

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = -\frac{\log\left(x - \frac{1}{2b} + \frac{\sqrt{-4ab^3+b^2}}{2b^2}\right) - \log\left(x + \frac{1}{2b} + \frac{\sqrt{-4ab^3+b^2}}{2b^2}\right)}{b}$$

[In] integrate(1/(a\*b-b\*\*2\*x\*\*2-x\*(-4\*a\*b\*\*3+b\*\*2)\*\*(1/2)),x)

[Out] -(log(x - 1/(2\*b) + sqrt(-4\*a\*b\*\*3 + b\*\*2)/(2\*b\*\*2)) - log(x + 1/(2\*b) + sqrt(-4\*a\*b\*\*3 + b\*\*2)/(2\*b\*\*2)))/b

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.65

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3x - b^2x^2}} dx = -\frac{\log\left(\frac{2b^2x-b+\sqrt{-4ab^3+b^2}}{2b^2x+b+\sqrt{-4ab^3+b^2}}\right)}{b}$$

[In] integrate(1/(a\*b-b^2\*x^2-x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="maxima")

[Out] -log((2\*b^2\*x - b + sqrt(-4\*a\*b^3 + b^2))/(2\*b^2\*x + b + sqrt(-4\*a\*b^3 + b^2)))/b

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.74

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3}x - b^2x^2} dx = -\frac{\log\left(\frac{2b^2x + \sqrt{-4ab+1}|b|-|b|}{2b^2x + \sqrt{-4ab+1}|b|+|b|}\right)}{|b|}$$

[In] integrate(1/(a\*b-b^2\*x^2-x\*(-4\*a\*b^3+b^2)^(1/2)),x, algorithm="giac")

[Out] -log(abs(2\*b^2\*x + sqrt(-4\*a\*b + 1)\*abs(b) - abs(b))/abs(2\*b^2\*x + sqrt(-4\*a\*b + 1)\*abs(b) + abs(b)))/abs(b)

**Mupad [B] (verification not implemented)**

Time = 0.13 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

$$\int \frac{1}{ab - \sqrt{b^2 - 4ab^3}x - b^2x^2} dx = \frac{2 \operatorname{atanh}\left(\frac{\sqrt{b^2 - 4ab^3}}{\sqrt{b^2}} + \frac{2b^2x}{\sqrt{b^2}}\right)}{\sqrt{b^2}}$$

[In] int(-1/(x\*(b^2 - 4\*a\*b^3)^(1/2) - a\*b + b^2\*x^2),x)

[Out] (2\*atanh((b^2 - 4\*a\*b^3)^(1/2)/(b^2)^(1/2) + (2\*b^2\*x)/(b^2)^(1/2)))/(b^2)^(1/2)

$$3.101 \quad \int \frac{1}{1+x^2+2x \cos\left(\frac{1}{7}\right)} dx$$

Optimal result	511
Rubi [A] (verified)	511
Mathematica [A] (verified)	512
Maple [B] (verified)	512
Fricas [A] (verification not implemented)	513
Sympy [C] (verification not implemented)	513
Maxima [B] (verification not implemented)	514
Giac [B] (verification not implemented)	514
Mupad [B] (verification not implemented)	514

### Optimal result

Integrand size = 14, antiderivative size = 17

$$\int \frac{1}{1+x^2+2x \cos\left(\frac{1}{7}\right)} dx = \arctan\left(\left(x + \cos\left(\frac{1}{7}\right)\right) \csc\left(\frac{1}{7}\right)\right) \csc\left(\frac{1}{7}\right)$$

[Out] arctan((x+cos(1/7))\*csc(1/7))\*csc(1/7)

### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {632, 210}

$$\int \frac{1}{1+x^2+2x \cos\left(\frac{1}{7}\right)} dx = \csc\left(\frac{1}{7}\right) \arctan\left(\csc\left(\frac{1}{7}\right) \left(x + \cos\left(\frac{1}{7}\right)\right)\right)$$

[In] Int[(1 + x^2 + 2\*x\*Cos[1/7])^(-1), x]

[Out] ArcTan[(x + Cos[1/7])\*Csc[1/7]]\*Csc[1/7]

#### Rule 210

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(-(Rt[-a, 2]\*Rt[-b, 2])^(-1))\*ArcTan[Rt[-b, 2]\*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

#### Rule 632

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-1), x\_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4\*a\*c - x^2, x], x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c},

x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-x^2 - 4\sin^2\left(\frac{1}{7}\right)} dx, x, 2x + 2\cos\left(\frac{1}{7}\right)\right)\right) \\ &= \tan^{-1}\left(\left(x + \cos\left(\frac{1}{7}\right)\right) \csc\left(\frac{1}{7}\right)\right) \csc\left(\frac{1}{7}\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx = \arctan\left(\cot\left(\frac{1}{7}\right) + x \csc\left(\frac{1}{7}\right)\right) \csc\left(\frac{1}{7}\right)$$

[In] Integrate[(1 + x^2 + 2\*x\*Cos[1/7])^(-1),x]

[Out] ArcTan[Cot[1/7] + x\*Csc[1/7]]\*Csc[1/7]

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 32 vs. 2(11) = 22.

Time = 3.11 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.94

method	result	size
default	$\frac{\arctan\left(\frac{2x+2\cos\left(\frac{1}{7}\right)}{2\sqrt{-\left(\cos^2\left(\frac{1}{7}\right)+1\right)}}\right)}{\sqrt{-\left(\cos^2\left(\frac{1}{7}\right)+1\right)}}$	33
risch	Expression too large to display	3085

[In] int(1/(1+x^2+2\*x\*cos(1/7)),x,method=\_RETURNVERBOSE)

[Out] 1/(-cos(1/7)^2+1)^(1/2)\*arctan(1/2\*(2\*x+2\*cos(1/7))/(-cos(1/7)^2+1)^(1/2))



**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx = \frac{\arctan\left(\frac{x + \cos\left(\frac{1}{7}\right)}{\sin\left(\frac{1}{7}\right)}\right)}{\sin\left(\frac{1}{7}\right)}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7)),x, algorithm="fricas")

[Out] arctan((x + cos(1/7))/sin(1/7))/sin(1/7)

**Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.09 (sec) , antiderivative size = 165, normalized size of antiderivative = 9.71

$$\begin{aligned} & \int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx \\ &= -\frac{i \log\left(x + \cos\left(\frac{1}{7}\right) - \frac{i}{\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}} + \frac{i \cos^2\left(\frac{1}{7}\right)}{\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}}\right)}{2\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}} \\ &+ \frac{i \log\left(x + \cos\left(\frac{1}{7}\right) - \frac{i \cos^2\left(\frac{1}{7}\right)}{\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}} + \frac{i}{\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}}\right)}{2\sqrt{1 - \cos\left(\frac{1}{7}\right)}\sqrt{\cos\left(\frac{1}{7}\right) + 1}} \end{aligned}$$

[In] integrate(1/(1+x\*\*2+2\*x\*cos(1/7)),x)

```
[Out] -I*log(x + cos(1/7) - I/(sqrt(1 - cos(1/7))*sqrt(cos(1/7) + 1)) + I*cos(1/7)
)**2/(sqrt(1 - cos(1/7))*sqrt(cos(1/7) + 1))/(2*sqrt(1 - cos(1/7))*sqrt(co
s(1/7) + 1)) + I*log(x + cos(1/7) - I*cos(1/7)**2/(sqrt(1 - cos(1/7))*sqrt(
cos(1/7) + 1)) + I/(sqrt(1 - cos(1/7))*sqrt(cos(1/7) + 1)))/(2*sqrt(1 - cos
(1/7))*sqrt(cos(1/7) + 1))
```

**Maxima [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx = \frac{\arctan\left(\frac{x + \cos\left(\frac{1}{7}\right)}{\sqrt{-\cos\left(\frac{1}{7}\right)^2 + 1}}\right)}{\sqrt{-\cos\left(\frac{1}{7}\right)^2 + 1}}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7)),x, algorithm="maxima")

[Out] arctan((x + cos(1/7))/sqrt(-cos(1/7)^2 + 1))/sqrt(-cos(1/7)^2 + 1)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx = \frac{\arctan\left(\frac{x + \cos\left(\frac{1}{7}\right)}{\sqrt{-\cos\left(\frac{1}{7}\right)^2 + 1}}\right)}{\sqrt{-\cos\left(\frac{1}{7}\right)^2 + 1}}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7)),x, algorithm="giac")

[Out] arctan((x + cos(1/7))/sqrt(-cos(1/7)^2 + 1))/sqrt(-cos(1/7)^2 + 1)

**Mupad [B] (verification not implemented)**

Time = 9.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.59

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{1}{7}\right)} dx = \frac{\operatorname{atan}\left(\frac{x + \cos\left(\frac{1}{7}\right)}{\sqrt{1 - \cos\left(\frac{1}{7}\right)^2}}\right)}{\sqrt{1 - \cos\left(\frac{1}{7}\right)^2}}$$

[In] int(1/(2\*x\*cos(1/7) + x^2 + 1),x)

[Out] atan((x + cos(1/7))/(1 - cos(1/7)^2)^(1/2))/(1 - cos(1/7)^2)^(1/2)

### 3.102 $\int \frac{1}{1+x^2+2x \cos(\frac{\pi}{7})} dx$

Optimal result . . . . .	515
Rubi [A] (verified) . . . . .	515
Mathematica [A] (verified) . . . . .	516
Maple [B] (verified) . . . . .	516
Fricas [A] (verification not implemented) . . . . .	517
Sympy [C] (verification not implemented) . . . . .	517
Maxima [A] (verification not implemented) . . . . .	517
Giac [A] (verification not implemented) . . . . .	518
Mupad [B] (verification not implemented) . . . . .	518

#### Optimal result

Integrand size = 16, antiderivative size = 23

$$\int \frac{1}{1+x^2+2x \cos(\frac{\pi}{7})} dx = \arctan\left(\cot\left(\frac{\pi}{7}\right) + x \csc\left(\frac{\pi}{7}\right)\right) \csc\left(\frac{\pi}{7}\right)$$

[Out] `arctan(cot(1/7*Pi)+x*csc(1/7*Pi))*csc(1/7*Pi)`

#### Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$ , Rules used = {632, 210}

$$\int \frac{1}{1+x^2+2x \cos(\frac{\pi}{7})} dx = \csc\left(\frac{\pi}{7}\right) \arctan\left(\csc\left(\frac{\pi}{7}\right) \left(x + \cos\left(\frac{\pi}{7}\right)\right)\right)$$

[In] `Int[(1 + x^2 + 2*x*Cos [Pi/7])^(-1), x]`

[Out] `ArcTan[(x + Cos [Pi/7])*Csc [Pi/7]]*Csc [Pi/7]`

#### Rule 210

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])`

#### Rule 632

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2\text{Subst}\left(\int \frac{1}{-x^2 - 4\sin^2\left(\frac{\pi}{7}\right)} dx, x, 2x + 2\cos\left(\frac{\pi}{7}\right)\right)\right) \\ &= \tan^{-1}\left(\left(x + \cos\left(\frac{\pi}{7}\right)\right) \csc\left(\frac{\pi}{7}\right)\right) \csc\left(\frac{\pi}{7}\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{\pi}{7}\right)} dx = \arctan\left(\cot\left(\frac{\pi}{7}\right) + x \csc\left(\frac{\pi}{7}\right)\right) \csc\left(\frac{\pi}{7}\right)$$

[In] Integrate[(1 + x^2 + 2\*x\*Cos[Pi/7])^(-1),x]

[Out] ArcTan[Cot[Pi/7] + x\*Csc[Pi/7]]\*Csc[Pi/7]

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(17) = 34.

Time = 3.14 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.70

method	result
default	$\frac{\arctan\left(\frac{2x+2\cos\left(\frac{\pi}{7}\right)}{2\sqrt{-(\cos^2\left(\frac{\pi}{7}\right))+1}}\right)}{\sqrt{-(\cos^2\left(\frac{\pi}{7}\right))+1}}$
norman	$\left(-\frac{4(-1)^{\frac{5}{7}}}{7} + \frac{(-1)^{\frac{4}{7}}}{7} - \frac{5(-1)^{\frac{3}{7}}}{7} + \frac{2(-1)^{\frac{2}{7}}}{7} - \frac{6(-1)^{\frac{1}{7}}}{7} + \frac{3}{7}\right) \ln\left((-1)^{\frac{5}{7}} - (-1)^{\frac{4}{7}} + (-1)^{\frac{3}{7}} - (-1)^{\frac{2}{7}} + (-1)^{\frac{1}{7}}\right)$
risch	$\frac{4\ln(x+(-1)^{\frac{1}{7}})(-1)^{\frac{5}{7}}}{7} - \frac{\ln(x+(-1)^{\frac{1}{7}})(-1)^{\frac{4}{7}}}{7} + \frac{5\ln(x+(-1)^{\frac{1}{7}})(-1)^{\frac{3}{7}}}{7} - \frac{2\ln(x+(-1)^{\frac{1}{7}})(-1)^{\frac{2}{7}}}{7} + \frac{6\ln(x+(-1)^{\frac{1}{7}})(-1)^{\frac{1}{7}}}{7}$

[In] int(1/(1+x^2+2\*x\*cos(1/7\*Pi)),x,method=\_RETURNVERBOSE)

[Out] 1/(-cos(1/7\*Pi)^2+1)^(1/2)\*arctan(1/2\*(2\*x+2\*cos(1/7\*Pi))/(-cos(1/7\*Pi)^2+1)^(1/2))

**Fricas [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int \frac{1}{1+x^2+2x\cos\left(\frac{\pi}{7}\right)} dx = \frac{\arctan\left(\frac{x+\cos\left(\frac{1}{7}\pi\right)}{\sin\left(\frac{1}{7}\pi\right)}\right)}{\sin\left(\frac{1}{7}\pi\right)}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7\*pi)),x, algorithm="fricas")

[Out] arctan((x + cos(1/7\*pi))/sin(1/7\*pi))/sin(1/7\*pi)

**Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 70, normalized size of antiderivative = 3.04

$$\int \frac{1}{1+x^2+2x\cos\left(\frac{\pi}{7}\right)} dx = -\frac{i\log\left(x+\cos\left(\frac{\pi}{7}\right)-\frac{i(2-2\cos^2\left(\frac{\pi}{7}\right))}{2\sin\left(\frac{\pi}{7}\right)}\right)}{2\sin\left(\frac{\pi}{7}\right)} + \frac{i\log\left(x+\cos\left(\frac{\pi}{7}\right)+\frac{i(2-2\cos^2\left(\frac{\pi}{7}\right))}{2\sin\left(\frac{\pi}{7}\right)}\right)}{2\sin\left(\frac{\pi}{7}\right)}$$

[In] integrate(1/(1+x\*\*2+2\*x\*cos(1/7\*pi)),x)

[Out] -I\*log(x + cos(pi/7) - I\*(2 - 2\*cos(pi/7)\*\*2)/(2\*sin(pi/7)))/(2\*sin(pi/7)) + I\*log(x + cos(pi/7) + I\*(2 - 2\*cos(pi/7)\*\*2)/(2\*sin(pi/7)))/(2\*sin(pi/7))

**Maxima [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{1}{1+x^2+2x\cos\left(\frac{\pi}{7}\right)} dx = \frac{\arctan\left(\frac{x+\cos\left(\frac{1}{7}\pi\right)}{\sqrt{-\cos\left(\frac{1}{7}\pi\right)^2+1}}\right)}{\sqrt{-\cos\left(\frac{1}{7}\pi\right)^2+1}}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7\*pi)),x, algorithm="maxima")

[Out] arctan((x + cos(1/7\*pi))/sqrt(-cos(1/7\*pi)^2 + 1))/sqrt(-cos(1/7\*pi)^2 + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{\pi}{7}\right)} dx = \frac{\arctan\left(\frac{x + \cos\left(\frac{1}{7}\pi\right)}{\sqrt{-\cos\left(\frac{1}{7}\pi\right)^2 + 1}}\right)}{\sqrt{-\cos\left(\frac{1}{7}\pi\right)^2 + 1}}$$

[In] integrate(1/(1+x^2+2\*x\*cos(1/7\*pi)),x, algorithm="giac")

[Out] arctan((x + cos(1/7\*pi))/sqrt(-cos(1/7\*pi)^2 + 1))/sqrt(-cos(1/7\*pi)^2 + 1)

**Mupad [B] (verification not implemented)**

Time = 9.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.83

$$\int \frac{1}{1 + x^2 + 2x \cos\left(\frac{\pi}{7}\right)} dx = -\frac{\operatorname{atanh}\left(\frac{x + \cos\left(\frac{\pi}{7}\right)}{\sqrt{\cos\left(\frac{\pi}{7}\right) - 1} \sqrt{\cos\left(\frac{\pi}{7}\right) + 1}}\right)}{\sqrt{\cos\left(\frac{\pi}{7}\right) - 1} \sqrt{\cos\left(\frac{\pi}{7}\right) + 1}}$$

[In] int(1/(x^2 + 2\*x\*cos(Pi/7) + 1),x)

[Out] -atanh((x + cos(Pi/7))/((cos(Pi/7) - 1)^(1/2)\*(cos(Pi/7) + 1)^(1/2)))/((cos(Pi/7) - 1)^(1/2)\*(cos(Pi/7) + 1)^(1/2)))

### 3.103 $\int \sqrt{5 - 6x + 9x^2} dx$

Optimal result	519
Rubi [A] (verified)	519
Mathematica [A] (verified)	520
Maple [A] (verified)	520
Fricas [A] (verification not implemented)	521
Sympy [A] (verification not implemented)	521
Maxima [A] (verification not implemented)	521
Giac [A] (verification not implemented)	522
Mupad [B] (verification not implemented)	522

#### Optimal result

Integrand size = 14, antiderivative size = 38

$$\int \sqrt{5 - 6x + 9x^2} dx = -\frac{1}{6}(1 - 3x)\sqrt{5 - 6x + 9x^2} + \frac{2}{3}\operatorname{arcsinh}\left(\frac{1}{2}(-1 + 3x)\right)$$

[Out] 2/3\*arcsinh(-1/2+3/2\*x)-1/6\*(1-3\*x)\*(9\*x^2-6\*x+5)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 221}

$$\int \sqrt{5 - 6x + 9x^2} dx = \frac{2}{3}\operatorname{arcsinh}\left(\frac{1}{2}(3x - 1)\right) - \frac{1}{6}(1 - 3x)\sqrt{9x^2 - 6x + 5}$$

[In] Int[Sqrt[5 - 6\*x + 9\*x^2],x]

[Out] -1/6\*((1 - 3\*x)\*Sqrt[5 - 6\*x + 9\*x^2]) + (2\*ArcSinh[(-1 + 3\*x)/2])/3

#### Rule 221

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N

`eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

### Rule 633

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{6}(1-3x)\sqrt{5-6x+9x^2} + 2 \int \frac{1}{\sqrt{5-6x+9x^2}} dx \\ &= -\frac{1}{6}(1-3x)\sqrt{5-6x+9x^2} + \frac{1}{18} \text{Subst} \left( \int \frac{1}{\sqrt{1+\frac{x^2}{144}}} dx, x, -6+18x \right) \\ &= -\frac{1}{6}(1-3x)\sqrt{5-6x+9x^2} + \frac{2}{3} \sinh^{-1} \left( \frac{1}{2}(-1+3x) \right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.26

$$\int \sqrt{5-6x+9x^2} dx = \frac{1}{6}(-1+3x)\sqrt{5-6x+9x^2} - \frac{2}{3} \log \left( 1-3x+\sqrt{5-6x+9x^2} \right)$$

[In] Integrate[Sqrt[5 - 6\*x + 9\*x^2], x]

[Out] ((-1 + 3\*x)\*Sqrt[5 - 6\*x + 9\*x^2])/6 - (2\*Log[1 - 3\*x + Sqrt[5 - 6\*x + 9\*x^2]])/3

### Maple [A] (verified)

Time = 2.77 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{(18x-6)\sqrt{9x^2-6x+5}}{36} + \frac{2 \operatorname{arcsinh}\left(-\frac{1}{2}+\frac{3x}{2}\right)}{3}$	29
risch	$\frac{\sqrt{9x^2-6x+5}(3x-1)}{6} + \frac{2 \operatorname{arcsinh}\left(-\frac{1}{2}+\frac{3x}{2}\right)}{3}$	29
trager	$\left(\frac{x}{2} - \frac{1}{6}\right) \sqrt{9x^2 - 6x + 5} + \frac{2 \ln\left(-1+3x+\sqrt{9x^2-6x+5}\right)}{3}$	40

[In] int((9\*x^2-6\*x+5)^(1/2), x, method=\_RETURNVERBOSE)



[Out]  $1/36*(18*x-6)*(9*x^2-6*x+5)^{(1/2)}+2/3*\operatorname{arcsinh}(-1/2+3/2*x)$

### Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05

$$\int \sqrt{5-6x+9x^2} dx = \frac{1}{6} \sqrt{9x^2-6x+5}(3x-1) - \frac{2}{3} \log(-3x + \sqrt{9x^2-6x+5} + 1)$$

[In] `integrate((9*x^2-6*x+5)^(1/2),x, algorithm="fricas")`

[Out]  $1/6*\operatorname{sqrt}(9*x^2-6*x+5)*(3*x-1) - 2/3*\log(-3*x + \operatorname{sqrt}(9*x^2-6*x+5) + 1)$

### Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \sqrt{5-6x+9x^2} dx = \left(\frac{x}{2} - \frac{1}{6}\right) \sqrt{9x^2-6x+5} + \frac{2 \operatorname{asinh}\left(\frac{3x}{2} - \frac{1}{2}\right)}{3}$$

[In] `integrate((9*x**2-6*x+5)**(1/2),x)`

[Out]  $(x/2 - 1/6)*\operatorname{sqrt}(9*x**2 - 6*x + 5) + 2*\operatorname{asinh}(3*x/2 - 1/2)/3$

### Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \sqrt{5-6x+9x^2} dx = \frac{1}{2} \sqrt{9x^2-6x+5}x - \frac{1}{6} \sqrt{9x^2-6x+5} + \frac{2}{3} \operatorname{arsinh}\left(\frac{3}{2}x - \frac{1}{2}\right)$$

[In] `integrate((9*x^2-6*x+5)^(1/2),x, algorithm="maxima")`

[Out]  $1/2*\operatorname{sqrt}(9*x^2-6*x+5)*x - 1/6*\operatorname{sqrt}(9*x^2-6*x+5) + 2/3*\operatorname{arcsinh}(3/2*x - 1/2)$

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.05

$$\int \sqrt{5 - 6x + 9x^2} dx = \frac{1}{6} \sqrt{9x^2 - 6x + 5}(3x - 1) - \frac{2}{3} \log(-3x + \sqrt{9x^2 - 6x + 5} + 1)$$

[In] integrate((9\*x^2-6\*x+5)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(9\*x^2 - 6\*x + 5)\*(3\*x - 1) - 2/3\*log(-3\*x + sqrt(9\*x^2 - 6\*x + 5) + 1)

**Mupad [B] (verification not implemented)**

Time = 0.08 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \sqrt{5 - 6x + 9x^2} dx = \frac{2 \ln(3x + \sqrt{9x^2 - 6x + 5} - 1)}{3} + \left(\frac{x}{2} - \frac{1}{6}\right) \sqrt{9x^2 - 6x + 5}$$

[In] int((9\*x^2 - 6\*x + 5)^(1/2),x)

[Out] (2\*log(3\*x + (9\*x^2 - 6\*x + 5)^(1/2) - 1))/3 + (x/2 - 1/6)\*(9\*x^2 - 6\*x + 5)^(1/2)

### 3.104 $\int \sqrt{3 - 4x - 4x^2} dx$

Optimal result	523
Rubi [A] (verified)	523
Mathematica [A] (verified)	524
Maple [A] (verified)	524
Fricas [B] (verification not implemented)	525
Sympy [A] (verification not implemented)	525
Maxima [A] (verification not implemented)	525
Giac [A] (verification not implemented)	526
Mupad [B] (verification not implemented)	526

#### Optimal result

Integrand size = 14, antiderivative size = 30

$$\int \sqrt{3 - 4x - 4x^2} dx = \frac{1}{4}(1 + 2x)\sqrt{3 - 4x - 4x^2} + \arcsin\left(\frac{1}{2} + x\right)$$

[Out]  $\arcsin(1/2+x)+1/4*(1+2*x)*(-4*x^2-4*x+3)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 222}

$$\int \sqrt{3 - 4x - 4x^2} dx = \arcsin\left(x + \frac{1}{2}\right) + \frac{1}{4}\sqrt{-4x^2 - 4x + 3}(2x + 1)$$

[In]  $\text{Int}[\text{Sqrt}[3 - 4*x - 4*x^2], x]$

[Out]  $((1 + 2*x)*\text{Sqrt}[3 - 4*x - 4*x^2])/4 + \text{ArcSin}[1/2 + x]$

#### Rule 222

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{NegQ}[b]$

#### Rule 626

$\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2]^{(p_)}, x\_Symbol] \rightarrow \text{Simp}[(b + 2*c*x) * ((a + b*x + c*x^2)^p / (2*c*(2*p + 1))), x] - \text{Dist}[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), \text{Int}[(a + b*x + c*x^2)^{(p-1)}, x], x] /; \text{FreeQ}\{a, b, c, x\} \ \&\& \ N$

`eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

### Rule 633

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{4}(1+2x)\sqrt{3-4x-4x^2} + 2 \int \frac{1}{\sqrt{3-4x-4x^2}} dx \\ &= \frac{1}{4}(1+2x)\sqrt{3-4x-4x^2} - \frac{1}{8} \text{Subst} \left( \int \frac{1}{\sqrt{1-\frac{x^2}{64}}} dx, x, -4-8x \right) \\ &= \frac{1}{4}(1+2x)\sqrt{3-4x-4x^2} + \sin^{-1} \left( \frac{1}{2} + x \right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.63

$$\int \sqrt{3-4x-4x^2} dx = \frac{1}{4}(1+2x)\sqrt{3-4x-4x^2} - 2 \arctan \left( \frac{\sqrt{3-4x-4x^2}}{3+2x} \right)$$

`[In] Integrate[Sqrt[3 - 4*x - 4*x^2], x]`

`[Out] ((1 + 2*x)*Sqrt[3 - 4*x - 4*x^2])/4 - 2*ArcTan[Sqrt[3 - 4*x - 4*x^2]/(3 + 2*x)]`

### Maple [A] (verified)

Time = 2.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.83

method	result
default	$-\frac{(-8x-4)\sqrt{-4x^2-4x+3}}{16} + \arcsin\left(x + \frac{1}{2}\right)$
risch	$-\frac{(4x^2+4x-3)(1+2x)}{4\sqrt{-4x^2-4x+3}} + \arcsin\left(x + \frac{1}{2}\right)$
trager	$\left(\frac{1}{4} + \frac{x}{2}\right) \sqrt{-4x^2 - 4x + 3} + \text{RootOf}(\_Z^2 + 1) \ln(-2 \text{RootOf}(\_Z^2 + 1) x - \text{RootOf}(\_Z^2 + 1)) -$

`[In] int((-4*x^2-4*x+3)^(1/2), x, method=_RETURNVERBOSE)`

[Out]  $-1/16*(-8*x-4)*(-4*x^2-4*x+3)^{(1/2)}+\arcsin(x+1/2)$

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs.  $2(24) = 48$ .

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.77

$$\int \sqrt{3 - 4x - 4x^2} dx = \frac{1}{4} \sqrt{-4x^2 - 4x + 3}(2x + 1) - \arctan\left(\frac{\sqrt{-4x^2 - 4x + 3}(2x + 1)}{4x^2 + 4x - 3}\right)$$

[In] `integrate((-4*x^2-4*x+3)^(1/2),x, algorithm="fricas")`

[Out]  $1/4*\sqrt{-4*x^2 - 4*x + 3}*(2*x + 1) - \arctan(\sqrt{-4*x^2 - 4*x + 3}*(2*x + 1)/(4*x^2 + 4*x - 3))$

### Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.87

$$\int \sqrt{3 - 4x - 4x^2} dx = \left(\frac{x}{2} + \frac{1}{4}\right) \sqrt{-4x^2 - 4x + 3} + \arcsin\left(x + \frac{1}{2}\right)$$

[In] `integrate((-4*x**2-4*x+3)**(1/2),x)`

[Out]  $(x/2 + 1/4)*\sqrt{-4*x**2 - 4*x + 3} + \arcsin(x + 1/2)$

### Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int \sqrt{3 - 4x - 4x^2} dx = \frac{1}{2} \sqrt{-4x^2 - 4x + 3}x + \frac{1}{4} \sqrt{-4x^2 - 4x + 3} - \arcsin\left(-x - \frac{1}{2}\right)$$

[In] `integrate((-4*x^2-4*x+3)^(1/2),x, algorithm="maxima")`

[Out]  $1/2*\sqrt{-4*x^2 - 4*x + 3}*x + 1/4*\sqrt{-4*x^2 - 4*x + 3} - \arcsin(-x - 1/2)$

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.80

$$\int \sqrt{3 - 4x - 4x^2} dx = \frac{1}{4} \sqrt{-4x^2 - 4x + 3}(2x + 1) + \arcsin\left(x + \frac{1}{2}\right)$$

[In] integrate((-4\*x^2-4\*x+3)^(1/2),x, algorithm="giac")

[Out] 1/4\*sqrt(-4\*x^2 - 4\*x + 3)\*(2\*x + 1) + arcsin(x + 1/2)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.77

$$\int \sqrt{3 - 4x - 4x^2} dx = \operatorname{asin}\left(x + \frac{1}{2}\right) + \left(\frac{x}{2} + \frac{1}{4}\right) \sqrt{-4x^2 - 4x + 3}$$

[In] int((3 - 4\*x^2 - 4\*x)^(1/2),x)

[Out] asin(x + 1/2) + (x/2 + 1/4)\*(3 - 4\*x^2 - 4\*x)^(1/2)

### 3.105 $\int \sqrt{-8 + 6x + 9x^2} dx$

Optimal result	527
Rubi [A] (verified)	527
Mathematica [A] (verified)	528
Maple [A] (verified)	528
Fricas [A] (verification not implemented)	529
Sympy [A] (verification not implemented)	529
Maxima [A] (verification not implemented)	529
Giac [A] (verification not implemented)	530
Mupad [B] (verification not implemented)	530

#### Optimal result

Integrand size = 14, antiderivative size = 49

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{6}(1 + 3x)\sqrt{-8 + 6x + 9x^2} - \frac{3}{2}\operatorname{arctanh}\left(\frac{1 + 3x}{\sqrt{-8 + 6x + 9x^2}}\right)$$

[Out]  $-3/2*\operatorname{arctanh}((1+3*x)/(9*x^2+6*x-8)^{(1/2)})+1/6*(1+3*x)*(9*x^2+6*x-8)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 635, 212}

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{6}(3x + 1)\sqrt{9x^2 + 6x - 8} - \frac{3}{2}\operatorname{arctanh}\left(\frac{3x + 1}{\sqrt{9x^2 + 6x - 8}}\right)$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[-8 + 6*x + 9*x^2], x]$

[Out]  $((1 + 3*x)*\operatorname{Sqrt}[-8 + 6*x + 9*x^2])/6 - (3*\operatorname{ArcTanh}[(1 + 3*x)/\operatorname{Sqrt}[-8 + 6*x + 9*x^2]])/2$

#### Rule 212

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$  FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])

#### Rule 626

$\operatorname{Int}[(a + (b \cdot x) + (c \cdot x)^2)^p, x\_Symbol] \rightarrow \operatorname{Simp}[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - \operatorname{Dist}[p*((b^2 - 4*a*c)/(2*c*(2*$

$p + 1))$ ,  $\text{Int}[(a + b*x + c*x^2)^{(p - 1)}, x], x] /;$   $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{IntegerQ}[4*p]$

### Rule 635

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_) + (c_)*(x_)^2], x\_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /;$   $\text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{6}(1 + 3x)\sqrt{-8 + 6x + 9x^2} - \frac{9}{2} \int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx \\ &= \frac{1}{6}(1 + 3x)\sqrt{-8 + 6x + 9x^2} - 9 \text{Subst}\left(\int \frac{1}{36 - x^2} dx, x, \frac{6 + 18x}{\sqrt{-8 + 6x + 9x^2}}\right) \\ &= \frac{1}{6}(1 + 3x)\sqrt{-8 + 6x + 9x^2} - \frac{3}{2} \tanh^{-1}\left(\frac{1 + 3x}{\sqrt{-8 + 6x + 9x^2}}\right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{6}(1 + 3x)\sqrt{-8 + 6x + 9x^2} - 3 \text{arctanh}\left(\frac{\sqrt{-8 + 6x + 9x^2}}{-2 + 3x}\right)$$

[In] Integrate[Sqrt[-8 + 6\*x + 9\*x^2],x]

[Out] ((1 + 3\*x)\*Sqrt[-8 + 6\*x + 9\*x^2])/6 - 3\*ArcTanh[Sqrt[-8 + 6\*x + 9\*x^2]/(-2 + 3\*x)]

### Maple [A] (verified)

Time = 2.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

method	result	size
trager	$\left(\frac{x}{2} + \frac{1}{6}\right)\sqrt{9x^2 + 6x - 8} - \frac{3 \ln(\sqrt{9x^2 + 6x - 8} + 1 + 3x)}{2}$	40
default	$\frac{(18x+6)\sqrt{9x^2+6x-8}}{36} - \frac{\ln\left(\frac{(9x+3)\sqrt{9} + \sqrt{9x^2+6x-8}}{2}\right)\sqrt{9}}{2}$	50
risch	$\frac{(3x+1)\sqrt{9x^2+6x-8}}{6} - \frac{\ln\left(\frac{(9x+3)\sqrt{9} + \sqrt{9x^2+6x-8}}{2}\right)\sqrt{9}}{2}$	50



[In] `int((9*x^2+6*x-8)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $(1/2*x+1/6)*(9*x^2+6*x-8)^(1/2)-3/2*\ln((9*x^2+6*x-8)^(1/2)+1+3*x)$

### Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.82

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{6} \sqrt{9x^2 + 6x - 8}(3x + 1) + \frac{3}{2} \log(-3x + \sqrt{9x^2 + 6x - 8} - 1)$$

[In] `integrate((9*x^2+6*x-8)^(1/2),x, algorithm="fricas")`

[Out]  $1/6*\text{sqrt}(9*x^2 + 6*x - 8)*(3*x + 1) + 3/2*\log(-3*x + \text{sqrt}(9*x^2 + 6*x - 8) - 1)$

### Sympy [A] (verification not implemented)

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int \sqrt{-8 + 6x + 9x^2} dx = \left(\frac{x}{2} + \frac{1}{6}\right) \sqrt{9x^2 + 6x - 8} - \frac{3 \log(18x + 6\sqrt{9x^2 + 6x - 8} + 6)}{2}$$

[In] `integrate((9*x**2+6*x-8)**(1/2),x)`

[Out]  $(x/2 + 1/6)*\text{sqrt}(9*x**2 + 6*x - 8) - 3*\log(18*x + 6*\text{sqrt}(9*x**2 + 6*x - 8) + 6)/2$

### Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.06

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{2} \sqrt{9x^2 + 6x - 8}x + \frac{1}{6} \sqrt{9x^2 + 6x - 8} - \frac{3}{2} \log(18x + 6\sqrt{9x^2 + 6x - 8} + 6)$$

[In] `integrate((9*x^2+6*x-8)^(1/2),x, algorithm="maxima")`

[Out]  $1/2*\text{sqrt}(9*x^2 + 6*x - 8)*x + 1/6*\text{sqrt}(9*x^2 + 6*x - 8) - 3/2*\log(18*x + 6*\text{sqrt}(9*x^2 + 6*x - 8) + 6)$

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \sqrt{-8 + 6x + 9x^2} dx = \frac{1}{6} \sqrt{9x^2 + 6x - 8}(3x + 1) + \frac{3}{2} \log \left( \left| -3x + \sqrt{9x^2 + 6x - 8} - 1 \right| \right)$$

[In] integrate((9\*x^2+6\*x-8)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(9\*x^2 + 6\*x - 8)\*(3\*x + 1) + 3/2\*log(abs(-3\*x + sqrt(9\*x^2 + 6\*x - 8) - 1))

**Mupad [B] (verification not implemented)**

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.80

$$\int \sqrt{-8 + 6x + 9x^2} dx = \left( \frac{x}{2} + \frac{1}{6} \right) \sqrt{9x^2 + 6x - 8} - \frac{3 \ln(3x + \sqrt{9x^2 + 6x - 8} + 1)}{2}$$

[In] int((6\*x + 9\*x^2 - 8)^(1/2),x)

[Out] (x/2 + 1/6)\*(6\*x + 9\*x^2 - 8)^(1/2) - (3\*log(3\*x + (6\*x + 9\*x^2 - 8)^(1/2) + 1))/2

### 3.106 $\int \sqrt{2 + 4x + 3x^2} dx$

Optimal result	531
Rubi [A] (verified)	531
Mathematica [A] (verified)	532
Maple [A] (verified)	532
Fricas [A] (verification not implemented)	533
Sympy [A] (verification not implemented)	533
Maxima [A] (verification not implemented)	534
Giac [A] (verification not implemented)	534
Mupad [B] (verification not implemented)	534

#### Optimal result

Integrand size = 14, antiderivative size = 45

$$\int \sqrt{2 + 4x + 3x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{2 + 4x + 3x^2} + \frac{\operatorname{arcsinh}\left(\frac{2+3x}{\sqrt{2}}\right)}{3\sqrt{3}}$$

[Out]  $1/9*\operatorname{arcsinh}(1/2*(2+3*x)*2^{(1/2)})*3^{(1/2)}+1/6*(2+3*x)*(3*x^2+4*x+2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 221}

$$\int \sqrt{2 + 4x + 3x^2} dx = \frac{\operatorname{arcsinh}\left(\frac{3x+2}{\sqrt{2}}\right)}{3\sqrt{3}} + \frac{1}{6}\sqrt{3x^2 + 4x + 2}(3x + 2)$$

[In] Int[Sqrt[2 + 4\*x + 3\*x^2], x]

[Out]  $((2 + 3*x)*\operatorname{Sqrt}[2 + 4*x + 3*x^2])/6 + \operatorname{ArcSinh}[(2 + 3*x)/\operatorname{Sqrt}[2]]/(3*\operatorname{Sqrt}[3])$

#### Rule 221

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p)/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*

$p + 1))$ , Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{6}(2 + 3x)\sqrt{2 + 4x + 3x^2} + \frac{1}{3} \int \frac{1}{\sqrt{2 + 4x + 3x^2}} dx \\ &= \frac{1}{6}(2 + 3x)\sqrt{2 + 4x + 3x^2} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 + \frac{x^2}{8}}} dx, x, 4 + 6x\right)}{6\sqrt{6}} \\ &= \frac{1}{6}(2 + 3x)\sqrt{2 + 4x + 3x^2} + \frac{\sinh^{-1}\left(\frac{2+3x}{\sqrt{2}}\right)}{3\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.18

$$\int \sqrt{2 + 4x + 3x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{2 + 4x + 3x^2} - \frac{\log(-2 - 3x + \sqrt{6 + 12x + 9x^2})}{3\sqrt{3}}$$

[In] Integrate[Sqrt[2 + 4\*x + 3\*x^2], x]

[Out] ((2 + 3\*x)\*Sqrt[2 + 4\*x + 3\*x^2])/6 - Log[-2 - 3\*x + Sqrt[6 + 12\*x + 9\*x^2]]/(3\*Sqrt[3])

### Maple [A] (verified)

Time = 2.38 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

method	result	size
default	$\frac{(4+6x)\sqrt{3x^2+4x+2}}{12} + \frac{\sqrt{3} \operatorname{arcsinh}\left(\frac{3\sqrt{2}\left(\frac{2}{3}+x\right)}{2}\right)}{9}$	35
risch	$\frac{(2+3x)\sqrt{3x^2+4x+2}}{6} + \frac{\sqrt{3} \operatorname{arcsinh}\left(\frac{3\sqrt{2}\left(\frac{2}{3}+x\right)}{2}\right)}{9}$	35
trager	$\left(\frac{1}{3} + \frac{x}{2}\right) \sqrt{3x^2 + 4x + 2} + \frac{\operatorname{RootOf}\left(\_Z^2 - 3\right) \ln\left(3 \operatorname{RootOf}\left(\_Z^2 - 3\right) x + 2 \operatorname{RootOf}\left(\_Z^2 - 3\right) + 3\sqrt{3x^2 + 4x + 2}\right)}{9}$	61

[In] `int((3*x^2+4*x+2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/12*(4+6*x)*(3*x^2+4*x+2)^(1/2)+1/9*3^(1/2)*arcsinh(3/2*2^(1/2)*(2/3+x))`

### Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.29

$$\int \sqrt{2+4x+3x^2} dx = \frac{1}{6} \sqrt{3x^2+4x+2}(3x+2) + \frac{1}{18} \sqrt{3} \log\left(-\sqrt{3}\sqrt{3x^2+4x+2}(3x+2) - 9x^2 - 12x - 5\right)$$

[In] `integrate((3*x^2+4*x+2)^(1/2),x, algorithm="fricas")`

[Out] `1/6*sqrt(3*x^2 + 4*x + 2)*(3*x + 2) + 1/18*sqrt(3)*log(-sqrt(3)*sqrt(3*x^2 + 4*x + 2)*(3*x + 2) - 9*x^2 - 12*x - 5)`

### Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.91

$$\int \sqrt{2+4x+3x^2} dx = \left(\frac{x}{2} + \frac{1}{3}\right) \sqrt{3x^2+4x+2} + \frac{\sqrt{3} \operatorname{asinh}\left(\frac{3\sqrt{2}\left(x+\frac{2}{3}\right)}{2}\right)}{9}$$

[In] `integrate((3*x**2+4*x+2)**(1/2),x)`

[Out] `(x/2 + 1/3)*sqrt(3*x**2 + 4*x + 2) + sqrt(3)*asinh(3*sqrt(2)*(x + 2/3)/2)/9`

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int \sqrt{2 + 4x + 3x^2} dx$$

$$= \frac{1}{2} \sqrt{3x^2 + 4x + 2} + \frac{1}{9} \sqrt{3} \operatorname{arsinh} \left( \frac{1}{2} \sqrt{2}(3x + 2) \right) + \frac{1}{3} \sqrt{3x^2 + 4x + 2}$$

[In] integrate((3\*x^2+4\*x+2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(3\*x^2 + 4\*x + 2)\*x + 1/9\*sqrt(3)\*arcsinh(1/2\*sqrt(2)\*(3\*x + 2)) + 1/3\*sqrt(3\*x^2 + 4\*x + 2)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.18

$$\int \sqrt{2 + 4x + 3x^2} dx = \frac{1}{6} \sqrt{3x^2 + 4x + 2}(3x + 2)$$

$$- \frac{1}{9} \sqrt{3} \log \left( -\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2 + 4x + 2} \right) - 2 \right)$$

[In] integrate((3\*x^2+4\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(3\*x^2 + 4\*x + 2)\*(3\*x + 2) - 1/9\*sqrt(3)\*log(-sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 4\*x + 2)) - 2)

**Mupad [B] (verification not implemented)**

Time = 9.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.07

$$\int \sqrt{2 + 4x + 3x^2} dx = \frac{\sqrt{3} \ln \left( \sqrt{3x^2 + 4x + 2} + \frac{\sqrt{3}(3x+2)}{3} \right)}{9} + \left( \frac{x}{2} + \frac{1}{3} \right) \sqrt{3x^2 + 4x + 2}$$

[In] int((4\*x + 3\*x^2 + 2)^(1/2),x)

[Out] (3^(1/2)\*log((4\*x + 3\*x^2 + 2)^(1/2) + (3^(1/2)\*(3\*x + 2))/3))/9 + (x/2 + 1/3)\*(4\*x + 3\*x^2 + 2)^(1/2)

### 3.107 $\int \sqrt{2 + 4x - 3x^2} dx$

Optimal result . . . . .	535
Rubi [A] (verified) . . . . .	535
Mathematica [A] (verified) . . . . .	536
Maple [A] (verified) . . . . .	536
Fricas [A] (verification not implemented) . . . . .	537
Sympy [A] (verification not implemented) . . . . .	537
Maxima [A] (verification not implemented) . . . . .	538
Giac [A] (verification not implemented) . . . . .	538
Mupad [B] (verification not implemented) . . . . .	538

#### Optimal result

Integrand size = 14, antiderivative size = 45

$$\int \sqrt{2 + 4x - 3x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{2 + 4x - 3x^2} - \frac{5 \arcsin\left(\frac{2-3x}{\sqrt{10}}\right)}{3\sqrt{3}}$$

[Out]  $-5/9*\arcsin(1/10*(2-3*x)*10^{(1/2)})*3^{(1/2)}-1/6*(2-3*x)*(-3*x^2+4*x+2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 222}

$$\int \sqrt{2 + 4x - 3x^2} dx = -\frac{5 \arcsin\left(\frac{2-3x}{\sqrt{10}}\right)}{3\sqrt{3}} - \frac{1}{6}\sqrt{-3x^2 + 4x + 2}(2 - 3x)$$

[In] Int[Sqrt[2 + 4\*x - 3\*x^2], x]

[Out]  $-1/6*((2 - 3*x)*\text{Sqrt}[2 + 4*x - 3*x^2]) - (5*\text{ArcSin}[(2 - 3*x)/\text{Sqrt}[10]])/(3*\text{Sqrt}[3])$

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^(p)/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*

$p + 1))$ , `Int[(a + b*x + c*x^2)^(p - 1), x], x] /;` `FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

### Rule 633

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /;` `FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{6}(2 - 3x)\sqrt{2 + 4x - 3x^2} + \frac{5}{3} \int \frac{1}{\sqrt{2 + 4x - 3x^2}} dx \\ &= -\frac{1}{6}(2 - 3x)\sqrt{2 + 4x - 3x^2} - \frac{1}{6} \sqrt{\frac{5}{6}} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{40}}} dx, x, 4 - 6x \right) \\ &= -\frac{1}{6}(2 - 3x)\sqrt{2 + 4x - 3x^2} - \frac{5 \sin^{-1} \left( \frac{2-3x}{\sqrt{10}} \right)}{3\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.36

$$\int \sqrt{2 + 4x - 3x^2} dx = \frac{1}{6}(-2 + 3x)\sqrt{2 + 4x - 3x^2} + \frac{10 \arctan \left( \frac{-2 - \sqrt{10} + 3x}{\sqrt{6 + 12x - 9x^2}} \right)}{3\sqrt{3}}$$

`[In] Integrate[Sqrt[2 + 4*x - 3*x^2], x]`

`[Out] ((-2 + 3*x)*Sqrt[2 + 4*x - 3*x^2])/6 + (10*ArcTan[(-2 - Sqrt[10] + 3*x)/Sqrt[6 + 12*x - 9*x^2]])/(3*Sqrt[3])`

### Maple [A] (verified)

Time = 2.22 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78



method	result
default	$-\frac{(4-6x)\sqrt{-3x^2+4x+2}}{12} + \frac{5\sqrt{3} \arcsin\left(\frac{3\sqrt{10}\left(-\frac{2}{3}+x\right)}{10}\right)}{9}$
risch	$-\frac{(3x^2-4x-2)(-2+3x)}{6\sqrt{-3x^2+4x+2}} + \frac{5\sqrt{3} \arcsin\left(\frac{3\sqrt{10}\left(-\frac{2}{3}+x\right)}{10}\right)}{9}$
trager	$\left(-\frac{1}{3} + \frac{x}{2}\right) \sqrt{-3x^2 + 4x + 2} - \frac{5 \operatorname{RootOf}(\_Z^2 + 3) \ln\left(3x \operatorname{RootOf}(\_Z^2 + 3) + 3\sqrt{-3x^2 + 4x + 2} - 2 \operatorname{RootOf}(\_Z^2 + 3)\right)}{9}$

[In] `int((-3*x^2+4*x+2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/12*(4-6*x)*(-3*x^2+4*x+2)^(1/2)+5/9*3^(1/2)*arcsin(3/10*10^(1/2)*(-2/3+x))`

### Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.33

$$\int \sqrt{2+4x-3x^2} dx = \frac{1}{6} \sqrt{-3x^2+4x+2}(3x-2) - \frac{5}{9} \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt{-3x^2+4x+2}(3x-2)}{3(3x^2-4x-2)}\right)$$

[In] `integrate((-3*x^2+4*x+2)^(1/2),x, algorithm="fricas")`

[Out] `1/6*sqrt(-3*x^2 + 4*x + 2)*(3*x - 2) - 5/9*sqrt(3)*arctan(1/3*sqrt(3)*sqrt(-3*x^2 + 4*x + 2)*(3*x - 2)/(3*x^2 - 4*x - 2))`

### Sympy [A] (verification not implemented)

Time = 0.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.93

$$\int \sqrt{2+4x-3x^2} dx = \left(\frac{x}{2} - \frac{1}{3}\right) \sqrt{-3x^2+4x+2} + \frac{5\sqrt{3} \operatorname{asin}\left(\frac{3\sqrt{10}\left(x-\frac{2}{3}\right)}{10}\right)}{9}$$

[In] `integrate((-3*x**2+4*x+2)**(1/2),x)`

[Out] `(x/2 - 1/3)*sqrt(-3*x**2 + 4*x + 2) + 5*sqrt(3)*asin(3*sqrt(10)*(x - 2/3)/10)/9`

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.02

$$\int \sqrt{2+4x-3x^2} dx = \frac{1}{2} \sqrt{-3x^2+4x+2}x - \frac{5}{9} \sqrt{3} \arcsin\left(-\frac{1}{10} \sqrt{10}(3x-2)\right) - \frac{1}{3} \sqrt{-3x^2+4x+2}$$

[In] integrate((-3\*x^2+4\*x+2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-3\*x^2 + 4\*x + 2)\*x - 5/9\*sqrt(3)\*arcsin(-1/10\*sqrt(10)\*(3\*x - 2)) - 1/3\*sqrt(-3\*x^2 + 4\*x + 2)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.80

$$\int \sqrt{2+4x-3x^2} dx = \frac{1}{6} \sqrt{-3x^2+4x+2}(3x-2) + \frac{5}{9} \sqrt{3} \arcsin\left(\frac{1}{10} \sqrt{10}(3x-2)\right)$$

[In] integrate((-3\*x^2+4\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(-3\*x^2 + 4\*x + 2)\*(3\*x - 2) + 5/9\*sqrt(3)\*arcsin(1/10\*sqrt(10)\*(3\*x - 2))

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.78

$$\int \sqrt{2+4x-3x^2} dx = \frac{5\sqrt{3} \operatorname{asin}\left(\frac{\sqrt{10}(3x-2)}{10}\right)}{9} + \left(\frac{x}{2} - \frac{1}{3}\right) \sqrt{-3x^2+4x+2}$$

[In] int((4\*x - 3\*x^2 + 2)^(1/2),x)

[Out] (5\*3^(1/2)\*asin((10^(1/2)\*(3\*x - 2))/10))/9 + (x/2 - 1/3)\*(4\*x - 3\*x^2 + 2)^(1/2)

### 3.108 $\int \sqrt{2 + 5x + 3x^2} dx$

Optimal result	539
Rubi [A] (verified)	539
Mathematica [A] (verified)	540
Maple [A] (verified)	540
Fricas [A] (verification not implemented)	541
Sympy [A] (verification not implemented)	541
Maxima [A] (verification not implemented)	542
Giac [A] (verification not implemented)	542
Mupad [B] (verification not implemented)	542

#### Optimal result

Integrand size = 14, antiderivative size = 62

$$\int \sqrt{2 + 5x + 3x^2} dx = \frac{1}{12}(5 + 6x)\sqrt{2 + 5x + 3x^2} - \frac{\operatorname{arctanh}\left(\frac{5+6x}{2\sqrt{3}\sqrt{2+5x+3x^2}}\right)}{24\sqrt{3}}$$

[Out]  $-1/72*\operatorname{arctanh}(1/6*(5+6*x)*3^{(1/2)}/(3*x^2+5*x+2)^{(1/2)})*3^{(1/2)}+1/12*(5+6*x)*(3*x^2+5*x+2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 635, 212}

$$\int \sqrt{2 + 5x + 3x^2} dx = \frac{1}{12}(6x + 5)\sqrt{3x^2 + 5x + 2} - \frac{\operatorname{arctanh}\left(\frac{6x+5}{2\sqrt{3}\sqrt{3x^2+5x+2}}\right)}{24\sqrt{3}}$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[2 + 5*x + 3*x^2], x]$

[Out]  $((5 + 6*x)*\operatorname{Sqrt}[2 + 5*x + 3*x^2])/12 - \operatorname{ArcTanh}[(5 + 6*x)/(2*\operatorname{Sqrt}[3]*\operatorname{Sqrt}[2 + 5*x + 3*x^2])]/(24*\operatorname{Sqrt}[3])$

#### Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

#### Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

### Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int
t[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,
b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{12}(5 + 6x)\sqrt{2 + 5x + 3x^2} - \frac{1}{24} \int \frac{1}{\sqrt{2 + 5x + 3x^2}} dx \\ &= \frac{1}{12}(5 + 6x)\sqrt{2 + 5x + 3x^2} - \frac{1}{12} \text{Subst} \left( \int \frac{1}{12 - x^2} dx, x, \frac{5 + 6x}{\sqrt{2 + 5x + 3x^2}} \right) \\ &= \frac{1}{12}(5 + 6x)\sqrt{2 + 5x + 3x^2} - \frac{\tanh^{-1} \left( \frac{5 + 6x}{2\sqrt{3}\sqrt{2 + 5x + 3x^2}} \right)}{24\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.90

$$\int \sqrt{2 + 5x + 3x^2} dx = \frac{1}{36} \left( 3(5 + 6x)\sqrt{2 + 5x + 3x^2} - \sqrt{3} \arctanh \left( \frac{\sqrt{\frac{2}{3} + \frac{5x}{3} + x^2}}{1 + x} \right) \right)$$

```
[In] Integrate[Sqrt[2 + 5*x + 3*x^2], x]
```

```
[Out] (3*(5 + 6*x)*Sqrt[2 + 5*x + 3*x^2] - Sqrt[3]*ArcTanh[Sqrt[2/3 + (5*x)/3 + x
^2]/(1 + x)]/36
```

### Maple [A] (verified)

Time = 2.26 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.81

method	result	size
default	$\frac{(5+6x)\sqrt{3x^2+5x+2}}{12} - \frac{\ln\left(\frac{\left(\frac{5}{2}+3x\right)\sqrt{3} + \sqrt{3x^2+5x+2}}{3}\right)\sqrt{3}}{72}$	50
risch	$\frac{(5+6x)\sqrt{3x^2+5x+2}}{12} - \frac{\ln\left(\frac{\left(\frac{5}{2}+3x\right)\sqrt{3} + \sqrt{3x^2+5x+2}}{3}\right)\sqrt{3}}{72}$	50
trager	$\left(\frac{5}{12} + \frac{x}{2}\right)\sqrt{3x^2+5x+2} - \frac{\text{RootOf}(\_Z^2-3)\ln\left(6\text{RootOf}(\_Z^2-3)x+5\text{RootOf}(\_Z^2-3)+6\sqrt{3x^2+5x+2}\right)}{72}$	61

```
[In] int((3*x^2+5*x+2)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/12*(5+6*x)*(3*x^2+5*x+2)^(1/2)-1/72*ln(1/3*(5/2+3*x)*3^(1/2)+(3*x^2+5*x+2)^(1/2))*3^(1/2)
```

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \sqrt{2+5x+3x^2} dx = \frac{1}{12} \sqrt{3x^2+5x+2}(6x+5) + \frac{1}{144} \sqrt{3} \log\left(-4\sqrt{3}\sqrt{3x^2+5x+2}(6x+5) + 72x^2 + 120x + 49\right)$$

```
[In] integrate((3*x^2+5*x+2)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/12*sqrt(3*x^2 + 5*x + 2)*(6*x + 5) + 1/144*sqrt(3)*log(-4*sqrt(3)*sqrt(3*x^2 + 5*x + 2)*(6*x + 5) + 72*x^2 + 120*x + 49)
```

### Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.85

$$\int \sqrt{2+5x+3x^2} dx = \left(\frac{x}{2} + \frac{5}{12}\right)\sqrt{3x^2+5x+2} - \frac{\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+5x+2} + 5)}{72}$$

```
[In] integrate((3*x**2+5*x+2)**(1/2),x)
```

```
[Out] (x/2 + 5/12)*sqrt(3*x**2 + 5*x + 2) - sqrt(3)*log(6*x + 2*sqrt(3)*sqrt(3*x**2 + 5*x + 2) + 5)/72
```

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \sqrt{2 + 5x + 3x^2} dx = \frac{1}{2} \sqrt{3x^2 + 5x + 2} - \frac{1}{72} \sqrt{3} \log \left( 2\sqrt{3}\sqrt{3x^2 + 5x + 2} + 6x + 5 \right) + \frac{5}{12} \sqrt{3x^2 + 5x + 2}$$

[In] integrate((3\*x^2+5\*x+2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(3\*x^2 + 5\*x + 2)\*x - 1/72\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 5\*x + 2) + 6\*x + 5) + 5/12\*sqrt(3\*x^2 + 5\*x + 2)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.87

$$\int \sqrt{2 + 5x + 3x^2} dx = \frac{1}{12} \sqrt{3x^2 + 5x + 2}(6x + 5) + \frac{1}{72} \sqrt{3} \log \left( \left| -2\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2 + 5x + 2} \right) - 5 \right| \right)$$

[In] integrate((3\*x^2+5\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(3\*x^2 + 5\*x + 2)\*(6\*x + 5) + 1/72\*sqrt(3)\*log(abs(-2\*sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 5\*x + 2)) - 5))

**Mupad [B] (verification not implemented)**

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.77

$$\int \sqrt{2 + 5x + 3x^2} dx = \left( \frac{x}{2} + \frac{5}{12} \right) \sqrt{3x^2 + 5x + 2} - \frac{\sqrt{3} \ln \left( \sqrt{3x^2 + 5x + 2} + \frac{\sqrt{3}(3x + \frac{5}{2})}{3} \right)}{72}$$

[In] int((5\*x + 3\*x^2 + 2)^(1/2),x)

[Out] (x/2 + 5/12)\*(5\*x + 3\*x^2 + 2)^(1/2) - (3^(1/2)\*log((5\*x + 3\*x^2 + 2)^(1/2) + (3^(1/2)\*(3\*x + 5/2))/3))/72

### 3.109 $\int \sqrt{2 + 5x - 3x^2} dx$

Optimal result	543
Rubi [A] (verified)	543
Mathematica [A] (verified)	544
Maple [A] (verified)	544
Fricas [A] (verification not implemented)	545
Sympy [A] (verification not implemented)	545
Maxima [A] (verification not implemented)	545
Giac [A] (verification not implemented)	546
Mupad [B] (verification not implemented)	546

#### Optimal result

Integrand size = 14, antiderivative size = 43

$$\int \sqrt{2 + 5x - 3x^2} dx = -\frac{1}{12}(5 - 6x)\sqrt{2 + 5x - 3x^2} - \frac{49 \arcsin\left(\frac{1}{7}(5 - 6x)\right)}{24\sqrt{3}}$$

[Out] 49/72\*arcsin(-5/7+6/7\*x)\*3^(1/2)-1/12\*(5-6\*x)\*(-3\*x^2+5\*x+2)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 222}

$$\int \sqrt{2 + 5x - 3x^2} dx = -\frac{49 \arcsin\left(\frac{1}{7}(5 - 6x)\right)}{24\sqrt{3}} - \frac{1}{12}\sqrt{-3x^2 + 5x + 2}(5 - 6x)$$

[In] Int[Sqrt[2 + 5\*x - 3\*x^2], x]

[Out] -1/12\*((5 - 6\*x)\*Sqrt[2 + 5\*x - 3\*x^2]) - (49\*ArcSin[(5 - 6\*x)/7])/(24\*Sqrt[3])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^p/(2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c)/(2\*c\*(2\*

$p + 1))$ , Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0] && GtQ[p, 0] && IntegerQ[4\*p]

### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{12}(5 - 6x)\sqrt{2 + 5x - 3x^2} + \frac{49}{24} \int \frac{1}{\sqrt{2 + 5x - 3x^2}} dx \\ &= -\frac{1}{12}(5 - 6x)\sqrt{2 + 5x - 3x^2} - \frac{7 \text{Subst}\left(\int \frac{1}{\sqrt{1 - \frac{x^2}{49}}} dx, x, 5 - 6x\right)}{24\sqrt{3}} \\ &= -\frac{1}{12}(5 - 6x)\sqrt{2 + 5x - 3x^2} - \frac{49 \sin^{-1}\left(\frac{1}{7}(5 - 6x)\right)}{24\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.30

$$\int \sqrt{2 + 5x - 3x^2} dx = \frac{1}{36} \left( 3(-5 + 6x)\sqrt{2 + 5x - 3x^2} - 49\sqrt{3} \arctan\left(\frac{\sqrt{6 + 15x - 9x^2}}{1 + 3x}\right) \right)$$

[In] Integrate[Sqrt[2 + 5\*x - 3\*x^2], x]

[Out] (3\*(-5 + 6\*x)\*Sqrt[2 + 5\*x - 3\*x^2] - 49\*Sqrt[3]\*ArcTan[Sqrt[6 + 15\*x - 9\*x^2]/(1 + 3\*x)]/36

### Maple [A] (verified)

Time = 2.18 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.74

method	result
default	$\frac{49 \arcsin\left(-\frac{5}{7} + \frac{6x}{7}\right)\sqrt{3}}{72} - \frac{(5-6x)\sqrt{-3x^2+5x+2}}{12}$
risch	$-\frac{(3x^2-5x-2)(-5+6x)}{12\sqrt{-3x^2+5x+2}} + \frac{49 \arcsin\left(-\frac{5}{7} + \frac{6x}{7}\right)\sqrt{3}}{72}$
trager	$\left(-\frac{5}{12} + \frac{x}{2}\right)\sqrt{-3x^2 + 5x + 2} + \frac{49 \text{RootOf}\left(\_Z^2 + 3\right) \ln\left(-6x \text{RootOf}\left(\_Z^2 + 3\right) + 6\sqrt{-3x^2 + 5x + 2} + 5 \text{RootOf}\left(\_Z^2 + 3\right)\right)}{72}$



[In] `int((-3*x^2+5*x+2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]  $49/72*\arcsin(-5/7+6/7*x)*3^(1/2)-1/12*(5-6*x)*(-3*x^2+5*x+2)^(1/2)$

### Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.40

$$\int \sqrt{2+5x-3x^2} dx = \frac{1}{12} \sqrt{-3x^2+5x+2}(6x-5) - \frac{49}{72} \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt{-3x^2+5x+2}(6x-5)}{6(3x^2-5x-2)}\right)$$

[In] `integrate((-3*x^2+5*x+2)^(1/2),x, algorithm="fricas")`

[Out]  $1/12*\sqrt{-3*x^2+5*x+2}*(6*x-5) - 49/72*\sqrt{3}*\arctan(1/6*\sqrt{3}*\sqrt{-3*x^2+5*x+2}*(6*x-5)/(3*x^2-5*x-2))$

### Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.86

$$\int \sqrt{2+5x-3x^2} dx = \left(\frac{x}{2} - \frac{5}{12}\right) \sqrt{-3x^2+5x+2} + \frac{49\sqrt{3} \operatorname{asin}\left(\frac{6x}{7} - \frac{5}{7}\right)}{72}$$

[In] `integrate((-3*x**2+5*x+2)**(1/2),x)`

[Out]  $(x/2 - 5/12)*\sqrt{-3*x**2+5*x+2} + 49*\sqrt{3}*\operatorname{asin}(6*x/7 - 5/7)/72$

### Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.95

$$\int \sqrt{2+5x-3x^2} dx = \frac{1}{2} \sqrt{-3x^2+5x+2} - \frac{49}{72} \sqrt{3} \arcsin\left(-\frac{6}{7}x + \frac{5}{7}\right) - \frac{5}{12} \sqrt{-3x^2+5x+2}$$

[In] `integrate((-3*x^2+5*x+2)^(1/2),x, algorithm="maxima")`

[Out]  $1/2*\sqrt{-3*x^2+5*x+2}*x - 49/72*\sqrt{3}*\arcsin(-6/7*x + 5/7) - 5/12*\sqrt{-3*x^2+5*x+2}$

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.72

$$\int \sqrt{2 + 5x - 3x^2} dx = \frac{1}{12} \sqrt{-3x^2 + 5x + 2}(6x - 5) + \frac{49}{72} \sqrt{3} \arcsin\left(\frac{6}{7}x - \frac{5}{7}\right)$$

[In] integrate((-3\*x^2+5\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(-3\*x^2 + 5\*x + 2)\*(6\*x - 5) + 49/72\*sqrt(3)\*arcsin(6/7\*x - 5/7)

**Mupad [B] (verification not implemented)**

Time = 8.97 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.70

$$\int \sqrt{2 + 5x - 3x^2} dx = \frac{49\sqrt{3} \operatorname{asin}\left(\frac{6x}{7} - \frac{5}{7}\right)}{72} + \left(\frac{x}{2} - \frac{5}{12}\right) \sqrt{-3x^2 + 5x + 2}$$

[In] int((5\*x - 3\*x^2 + 2)^(1/2),x)

[Out] (49\*3^(1/2)\*asin((6\*x)/7 - 5/7))/72 + (x/2 - 5/12)\*(5\*x - 3\*x^2 + 2)^(1/2)

### 3.110 $\int \sqrt{-2 + 4x + 3x^2} dx$

Optimal result	547
Rubi [A] (verified)	547
Mathematica [A] (verified)	548
Maple [A] (verified)	548
Fricas [A] (verification not implemented)	549
Sympy [A] (verification not implemented)	549
Maxima [A] (verification not implemented)	550
Giac [A] (verification not implemented)	550
Mupad [B] (verification not implemented)	550

#### Optimal result

Integrand size = 14, antiderivative size = 59

$$\int \sqrt{-2 + 4x + 3x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{-2 + 4x + 3x^2} - \frac{5\operatorname{arctanh}\left(\frac{2+3x}{\sqrt{3}\sqrt{-2+4x+3x^2}}\right)}{3\sqrt{3}}$$

[Out]  $-5/9*\operatorname{arctanh}(1/3*(2+3*x)*3^{(1/2)}/(3*x^2+4*x-2)^{(1/2)})*3^{(1/2)}+1/6*(2+3*x)*(3*x^2+4*x-2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 635, 212}

$$\int \sqrt{-2 + 4x + 3x^2} dx = \frac{1}{6}(3x + 2)\sqrt{3x^2 + 4x - 2} - \frac{5\operatorname{arctanh}\left(\frac{3x+2}{\sqrt{3}\sqrt{3x^2+4x-2}}\right)}{3\sqrt{3}}$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[-2 + 4*x + 3*x^2], x]$

[Out]  $((2 + 3*x)*\operatorname{Sqrt}[-2 + 4*x + 3*x^2])/6 - (5*\operatorname{ArcTanh}[(2 + 3*x)/(\operatorname{Sqrt}[3]*\operatorname{Sqrt}[-2 + 4*x + 3*x^2])])/(3*\operatorname{Sqrt}[3])$

#### Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

#### Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N
eQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

### Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int
t[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,
b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{6}(2 + 3x)\sqrt{-2 + 4x + 3x^2} - \frac{5}{3} \int \frac{1}{\sqrt{-2 + 4x + 3x^2}} dx \\ &= \frac{1}{6}(2 + 3x)\sqrt{-2 + 4x + 3x^2} - \frac{10}{3} \text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{4 + 6x}{\sqrt{-2 + 4x + 3x^2}}\right) \\ &= \frac{1}{6}(2 + 3x)\sqrt{-2 + 4x + 3x^2} - \frac{5 \tanh^{-1}\left(\frac{2+3x}{\sqrt{3}\sqrt{-2+4x+3x^2}}\right)}{3\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.03

$$\int \sqrt{-2 + 4x + 3x^2} dx = \frac{1}{6}(2 + 3x)\sqrt{-2 + 4x + 3x^2} - \frac{10 \operatorname{arctanh}\left(\frac{\sqrt{-6+12x+9x^2}}{2+\sqrt{10+3x}}\right)}{3\sqrt{3}}$$

```
[In] Integrate[Sqrt[-2 + 4*x + 3*x^2], x]
```

```
[Out] ((2 + 3*x)*Sqrt[-2 + 4*x + 3*x^2])/6 - (10*ArcTanh[Sqrt[-6 + 12*x + 9*x^2]/
(2 + Sqrt[10] + 3*x))]/(3*Sqrt[3])
```

### Maple [A] (verified)

Time = 2.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.85

method	result	size
default	$\frac{(4+6x)\sqrt{3x^2+4x-2}}{12} - \frac{5 \ln\left(\frac{(2+3x)\sqrt{3} + \sqrt{3x^2+4x-2}}{3}\right)\sqrt{3}}{9}$	50
risch	$\frac{(2+3x)\sqrt{3x^2+4x-2}}{6} - \frac{5 \ln\left(\frac{(2+3x)\sqrt{3} + \sqrt{3x^2+4x-2}}{3}\right)\sqrt{3}}{9}$	50
trager	$\left(\frac{1}{3} + \frac{x}{2}\right)\sqrt{3x^2+4x-2} - \frac{5 \operatorname{RootOf}(\_Z^2-3) \ln\left(3 \operatorname{RootOf}(\_Z^2-3)x + 2 \operatorname{RootOf}(\_Z^2-3) + 3\sqrt{3x^2+4x-2}\right)}{9}$	61

[In] `int((3*x^2+4*x-2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/12*(4+6*x)*(3*x^2+4*x-2)^(1/2)-5/9*ln(1/3*(2+3*x)*3^(1/2)+(3*x^2+4*x-2)^(1/2))*3^(1/2)`

### Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98

$$\int \sqrt{-2+4x+3x^2} dx = \frac{1}{6} \sqrt{3x^2+4x-2}(3x+2) + \frac{5}{18} \sqrt{3} \log\left(-\sqrt{3}\sqrt{3x^2+4x-2}(3x+2) + 9x^2 + 12x - 1\right)$$

[In] `integrate((3*x^2+4*x-2)^(1/2),x, algorithm="fricas")`

[Out] `1/6*sqrt(3*x^2 + 4*x - 2)*(3*x + 2) + 5/18*sqrt(3)*log(-sqrt(3)*sqrt(3*x^2 + 4*x - 2)*(3*x + 2) + 9*x^2 + 12*x - 1)`

### Sympy [A] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.92

$$\int \sqrt{-2+4x+3x^2} dx = \left(\frac{x}{2} + \frac{1}{3}\right)\sqrt{3x^2+4x-2} - \frac{5\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+4x-2} + 4)}{9}$$

[In] `integrate((3*x**2+4*x-2)**(1/2),x)`

[Out] `(x/2 + 1/3)*sqrt(3*x**2 + 4*x - 2) - 5*sqrt(3)*log(6*x + 2*sqrt(3)*sqrt(3*x**2 + 4*x - 2) + 4)/9`

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98

$$\int \sqrt{-2 + 4x + 3x^2} dx = \frac{1}{2} \sqrt{3x^2 + 4x - 2} - \frac{5}{9} \sqrt{3} \log \left( 2 \sqrt{3} \sqrt{3x^2 + 4x - 2} + 6x + 4 \right) + \frac{1}{3} \sqrt{3x^2 + 4x - 2}$$

[In] integrate((3\*x^2+4\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(3\*x^2 + 4\*x - 2)\*x - 5/9\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 4\*x - 2) + 6\*x + 4) + 1/3\*sqrt(3\*x^2 + 4\*x - 2)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.92

$$\int \sqrt{-2 + 4x + 3x^2} dx = \frac{1}{6} \sqrt{3x^2 + 4x - 2}(3x + 2) + \frac{5}{9} \sqrt{3} \log \left( \left| -\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2 + 4x - 2} \right) - 2 \right| \right)$$

[In] integrate((3\*x^2+4\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(3\*x^2 + 4\*x - 2)\*(3\*x + 2) + 5/9\*sqrt(3)\*log(abs(-sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 4\*x - 2)) - 2))

**Mupad [B] (verification not implemented)**

Time = 9.01 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.81

$$\int \sqrt{-2 + 4x + 3x^2} dx = \left( \frac{x}{2} + \frac{1}{3} \right) \sqrt{3x^2 + 4x - 2} - \frac{5\sqrt{3} \ln \left( \sqrt{3x^2 + 4x - 2} + \frac{\sqrt{3}(3x+2)}{3} \right)}{9}$$

[In] int((4\*x + 3\*x^2 - 2)^(1/2),x)

[Out] (x/2 + 1/3)\*(4\*x + 3\*x^2 - 2)^(1/2) - (5\*3^(1/2)\*log((4\*x + 3\*x^2 - 2)^(1/2) + (3^(1/2)\*(3\*x + 2))/3))/9

### 3.111 $\int \sqrt{-2 + 4x - 3x^2} dx$

Optimal result	551
Rubi [A] (verified)	551
Mathematica [A] (verified)	552
Maple [A] (verified)	552
Fricas [C] (verification not implemented)	553
Sympy [C] (verification not implemented)	553
Maxima [C] (verification not implemented)	554
Giac [F]	554
Mupad [B] (verification not implemented)	554

#### Optimal result

Integrand size = 14, antiderivative size = 59

$$\int \sqrt{-2 + 4x - 3x^2} dx = -\frac{1}{6}(2 - 3x)\sqrt{-2 + 4x - 3x^2} + \frac{\arctan\left(\frac{2-3x}{\sqrt{3}\sqrt{-2+4x-3x^2}}\right)}{3\sqrt{3}}$$

[Out]  $1/9*\arctan(1/3*(2-3*x)*3^{(1/2)/(-3*x^2+4*x-2)^{(1/2)}}*3^{(1/2)}-1/6*(2-3*x)*(-3*x^2+4*x-2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 635, 210}

$$\int \sqrt{-2 + 4x - 3x^2} dx = \frac{\arctan\left(\frac{2-3x}{\sqrt{3}\sqrt{-3x^2+4x-2}}\right)}{3\sqrt{3}} - \frac{1}{6}(2 - 3x)\sqrt{-3x^2 + 4x - 2}$$

[In] `Int[Sqrt[-2 + 4*x - 3*x^2],x]`

[Out]  $-1/6*((2 - 3*x)*\text{Sqrt}[-2 + 4*x - 3*x^2]) + \text{ArcTan}[(2 - 3*x)/(\text{Sqrt}[3]*\text{Sqrt}[-2 + 4*x - 3*x^2])]/(3*\text{Sqrt}[3])$

#### Rule 210

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

#### Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

### Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int
t[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,
b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{6}(2-3x)\sqrt{-2+4x-3x^2} - \frac{1}{3} \int \frac{1}{\sqrt{-2+4x-3x^2}} dx \\ &= -\frac{1}{6}(2-3x)\sqrt{-2+4x-3x^2} - \frac{2}{3} \text{Subst}\left(\int \frac{1}{-12-x^2} dx, x, \frac{4-6x}{\sqrt{-2+4x-3x^2}}\right) \\ &= -\frac{1}{6}(2-3x)\sqrt{-2+4x-3x^2} + \frac{\tan^{-1}\left(\frac{2-3x}{\sqrt{3}\sqrt{-2+4x-3x^2}}\right)}{3\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.78 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.92

$$\int \sqrt{-2+4x-3x^2} dx = \frac{1}{6}(-2+3x)\sqrt{-2+4x-3x^2} - \frac{\arctan\left(\frac{-2+3x}{\sqrt{-6+12x-9x^2}}\right)}{3\sqrt{3}}$$

```
[In] Integrate[Sqrt[-2 + 4*x - 3*x^2], x]
```

```
[Out] ((-2 + 3*x)*Sqrt[-2 + 4*x - 3*x^2])/6 - ArcTan[(-2 + 3*x)/Sqrt[-6 + 12*x -
9*x^2]]/(3*Sqrt[3])
```

### Maple [A] (verified)

Time = 3.19 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.78



method	result
default	$-\frac{(4-6x)\sqrt{-3x^2+4x-2}}{12} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(-\frac{2}{3}+x\right)}{\sqrt{-3x^2+4x-2}}\right)}{9}$
risch	$-\frac{(3x^2-4x+2)(-2+3x)}{6\sqrt{-3x^2+4x-2}} - \frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(-\frac{2}{3}+x\right)}{\sqrt{-3x^2+4x-2}}\right)}{9}$
trager	$\left(-\frac{1}{3} + \frac{x}{2}\right) \sqrt{-3x^2 + 4x - 2} + \frac{\text{RootOf}\left(\_Z^2+3\right) \ln\left(3x \text{RootOf}\left(\_Z^2+3\right)+3\sqrt{-3x^2+4x-2}-2 \text{RootOf}\left(\_Z^2+3\right)\right)}{9}$

[In] `int((-3*x^2+4*x-2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/12*(4-6*x)*(-3*x^2+4*x-2)^(1/2)-1/9*3^(1/2)*arctan(3^(1/2)*(-2/3+x)/(-3*x^2+4*x-2)^(1/2))`

### Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.46

$$\int \sqrt{-2+4x-3x^2} dx = \frac{1}{6} \sqrt{-3x^2+4x-2}(3x-2) - \frac{1}{18}i\sqrt{3} \log\left(-\frac{2(i\sqrt{3}\sqrt{-3x^2+4x-2}+3x-2)}{x}\right) + \frac{1}{18}i\sqrt{3} \log\left(-\frac{2(-i\sqrt{3}\sqrt{-3x^2+4x-2}+3x-2)}{x}\right)$$

[In] `integrate((-3*x^2+4*x-2)^(1/2),x, algorithm="fricas")`

[Out] `1/6*sqrt(-3*x^2 + 4*x - 2)*(3*x - 2) - 1/18*I*sqrt(3)*log(-2*(I*sqrt(3)*sqrt(-3*x^2 + 4*x - 2) + 3*x - 2)/x) + 1/18*I*sqrt(3)*log(-2*(-I*sqrt(3)*sqrt(-3*x^2 + 4*x - 2) + 3*x - 2)/x)`

### Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.95

$$\int \sqrt{-2+4x-3x^2} dx = \left(\frac{x}{2} - \frac{1}{3}\right) \sqrt{-3x^2+4x-2} + \frac{\sqrt{3}i \log(-6x + 2\sqrt{3}i\sqrt{-3x^2+4x-2} + 4)}{9}$$

[In] integrate((-3\*x\*\*2+4\*x-2)\*\*(1/2),x)

[Out] (x/2 - 1/3)\*sqrt(-3\*x\*\*2 + 4\*x - 2) + sqrt(3)\*I\*log(-6\*x + 2\*sqrt(3)\*I\*sqrt(-3\*x\*\*2 + 4\*x - 2) + 4)/9

## Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.78

$$\int \sqrt{-2 + 4x - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 4x - 2} + \frac{1}{9} i \sqrt{3} \operatorname{arsinh} \left( \frac{1}{2} \sqrt{2} (3x - 2) \right) - \frac{1}{3} \sqrt{-3x^2 + 4x - 2}$$

[In] integrate((-3\*x^2+4\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(-3\*x^2 + 4\*x - 2)\*x + 1/9\*I\*sqrt(3)\*arcsinh(1/2\*sqrt(2)\*(3\*x - 2)) - 1/3\*sqrt(-3\*x^2 + 4\*x - 2)

## Giac [F]

$$\int \sqrt{-2 + 4x - 3x^2} dx = \int \sqrt{-3x^2 + 4x - 2} dx$$

[In] integrate((-3\*x^2+4\*x-2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-3\*x^2 + 4\*x - 2), x)

## Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.61

$$\int \sqrt{-2 + 4x - 3x^2} dx = \frac{\sqrt{3} \operatorname{asin} \left( \frac{\sqrt{2} (3x-2) i i}{2} \right)}{9} + \left( \frac{x}{2} - \frac{1}{3} \right) \sqrt{-3x^2 + 4x - 2}$$

[In] int((4\*x - 3\*x^2 - 2)^(1/2),x)

[Out] (3^(1/2)\*asin((2^(1/2)\*(3\*x - 2)\*1i)/2))/9 + (x/2 - 1/3)\*(4\*x - 3\*x^2 - 2)^(1/2)

### 3.112 $\int \sqrt{-2 + 5x + 3x^2} dx$

Optimal result	555
Rubi [A] (verified)	555
Mathematica [A] (verified)	556
Maple [A] (verified)	557
Fricas [A] (verification not implemented)	557
Sympy [A] (verification not implemented)	557
Maxima [A] (verification not implemented)	558
Giac [A] (verification not implemented)	558
Mupad [B] (verification not implemented)	558

#### Optimal result

Integrand size = 14, antiderivative size = 62

$$\int \sqrt{-2 + 5x + 3x^2} dx = \frac{1}{12}(5 + 6x)\sqrt{-2 + 5x + 3x^2} - \frac{49\operatorname{arctanh}\left(\frac{5+6x}{2\sqrt{3}\sqrt{-2+5x+3x^2}}\right)}{24\sqrt{3}}$$

[Out]  $-49/72*\operatorname{arctanh}(1/6*(5+6*x)*3^{(1/2)}/(3*x^2+5*x-2)^{(1/2)})*3^{(1/2)}+1/12*(5+6*x)*(3*x^2+5*x-2)^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 635, 212}

$$\int \sqrt{-2 + 5x + 3x^2} dx = \frac{1}{12}(6x + 5)\sqrt{3x^2 + 5x - 2} - \frac{49\operatorname{arctanh}\left(\frac{6x+5}{2\sqrt{3}\sqrt{3x^2+5x-2}}\right)}{24\sqrt{3}}$$

[In]  $\operatorname{Int}[\operatorname{Sqrt}[-2 + 5*x + 3*x^2], x]$

[Out]  $((5 + 6*x)*\operatorname{Sqrt}[-2 + 5*x + 3*x^2])/12 - (49*\operatorname{ArcTanh}[(5 + 6*x)/(2*\operatorname{Sqrt}[3]*\operatorname{Sqrt}[-2 + 5*x + 3*x^2]))/(24*\operatorname{Sqrt}[3])$

#### Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$   $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt} Q[a, 0] \ || \ \operatorname{Lt} Q[b, 0])$

#### Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)
*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*
p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

### Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int
t[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,
b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

### Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{12}(5 + 6x)\sqrt{-2 + 5x + 3x^2} - \frac{49}{24} \int \frac{1}{\sqrt{-2 + 5x + 3x^2}} dx \\
&= \frac{1}{12}(5 + 6x)\sqrt{-2 + 5x + 3x^2} - \frac{49}{12} \text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{5 + 6x}{\sqrt{-2 + 5x + 3x^2}}\right) \\
&= \frac{1}{12}(5 + 6x)\sqrt{-2 + 5x + 3x^2} - \frac{49 \tanh^{-1}\left(\frac{5 + 6x}{2\sqrt{3}\sqrt{-2 + 5x + 3x^2}}\right)}{24\sqrt{3}}
\end{aligned}$$

### Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.90

$$\begin{aligned}
&\int \sqrt{-2 + 5x + 3x^2} dx \\
&= \frac{1}{36} \left( 3(5 + 6x)\sqrt{-2 + 5x + 3x^2} - 49\sqrt{3} \arctanh\left(\frac{\sqrt{-\frac{2}{3} + \frac{5x}{3} + x^2}}{2 + x}\right) \right)
\end{aligned}$$

```
[In] Integrate[Sqrt[-2 + 5*x + 3*x^2], x]
```

```
[Out] (3*(5 + 6*x)*Sqrt[-2 + 5*x + 3*x^2] - 49*Sqrt[3]*ArcTanh[Sqrt[-2/3 + (5*x)/
3 + x^2]/(2 + x)]/36
```

**Maple [A] (verified)**

Time = 2.30 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.81

method	result	
default	$\frac{(5+6x)\sqrt{3x^2+5x-2}}{12} - \frac{49 \ln\left(\frac{(\frac{5}{2}+3x)\sqrt{3}}{3} + \sqrt{3x^2+5x-2}\right)\sqrt{3}}{72}$	5
risch	$\frac{(5+6x)\sqrt{3x^2+5x-2}}{12} - \frac{49 \ln\left(\frac{(\frac{5}{2}+3x)\sqrt{3}}{3} + \sqrt{3x^2+5x-2}\right)\sqrt{3}}{72}$	5
trager	$\left(\frac{5}{12} + \frac{x}{2}\right)\sqrt{3x^2+5x-2} + \frac{49 \operatorname{RootOf}(\_Z^2-3) \ln(-6 \operatorname{RootOf}(\_Z^2-3)x - 5 \operatorname{RootOf}(\_Z^2-3) + 6\sqrt{3x^2+5x-2})}{72}$	6

```
[In] int((3*x^2+5*x-2)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/12*(5+6*x)*(3*x^2+5*x-2)^(1/2)-49/72*ln(1/3*(5/2+3*x)*3^(1/2)+(3*x^2+5*x-2)^(1/2))*3^(1/2)
```

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \sqrt{-2+5x+3x^2} dx = \frac{1}{12} \sqrt{3x^2+5x-2}(6x+5) + \frac{49}{144} \sqrt{3} \log\left(-4\sqrt{3}\sqrt{3x^2+5x-2}(6x+5) + 72x^2 + 120x + 1\right)$$

```
[In] integrate((3*x^2+5*x-2)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/12*sqrt(3*x^2 + 5*x - 2)*(6*x + 5) + 49/144*sqrt(3)*log(-4*sqrt(3)*sqrt(3*x^2 + 5*x - 2)*(6*x + 5) + 72*x^2 + 120*x + 1)
```

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.87

$$\int \sqrt{-2+5x+3x^2} dx = \left(\frac{x}{2} + \frac{5}{12}\right)\sqrt{3x^2+5x-2} - \frac{49\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+5x-2} + 5)}{72}$$

```
[In] integrate((3*x**2+5*x-2)**(1/2),x)
```

```
[Out] (x/2 + 5/12)*sqrt(3*x**2 + 5*x - 2) - 49*sqrt(3)*log(6*x + 2*sqrt(3)*sqrt(3*x**2 + 5*x - 2) + 5)/72
```

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.94

$$\int \sqrt{-2 + 5x + 3x^2} dx = \frac{1}{2} \sqrt{3x^2 + 5x - 2} - \frac{49}{72} \sqrt{3} \log \left( 2\sqrt{3}\sqrt{3x^2 + 5x - 2} + 6x + 5 \right) + \frac{5}{12} \sqrt{3x^2 + 5x - 2}$$

[In] integrate((3\*x^2+5\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/2\*sqrt(3\*x^2 + 5\*x - 2)\*x - 49/72\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 5\*x - 2) + 6\*x + 5) + 5/12\*sqrt(3\*x^2 + 5\*x - 2)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.87

$$\int \sqrt{-2 + 5x + 3x^2} dx = \frac{1}{12} \sqrt{3x^2 + 5x - 2}(6x + 5) + \frac{49}{72} \sqrt{3} \log \left( \left| -2\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2 + 5x - 2} \right) - 5 \right| \right)$$

[In] integrate((3\*x^2+5\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(3\*x^2 + 5\*x - 2)\*(6\*x + 5) + 49/72\*sqrt(3)\*log(abs(-2\*sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 5\*x - 2)) - 5))

**Mupad [B] (verification not implemented)**

Time = 0.22 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.77

$$\int \sqrt{-2 + 5x + 3x^2} dx = \left( \frac{x}{2} + \frac{5}{12} \right) \sqrt{3x^2 + 5x - 2} - \frac{49\sqrt{3} \ln \left( \sqrt{3x^2 + 5x - 2} + \frac{\sqrt{3}(3x + \frac{5}{2})}{3} \right)}{72}$$

[In] int((5\*x + 3\*x^2 - 2)^(1/2),x)

[Out] (x/2 + 5/12)\*(5\*x + 3\*x^2 - 2)^(1/2) - (49\*3^(1/2)\*log((5\*x + 3\*x^2 - 2)^(1/2) + (3^(1/2)\*(3\*x + 5/2))/3))/72

### 3.113 $\int \sqrt{-2 + 5x - 3x^2} dx$

Optimal result	559
Rubi [A] (verified)	559
Mathematica [A] (verified)	560
Maple [A] (verified)	560
Fricas [A] (verification not implemented)	561
Sympy [A] (verification not implemented)	561
Maxima [A] (verification not implemented)	561
Giac [A] (verification not implemented)	562
Mupad [B] (verification not implemented)	562

#### Optimal result

Integrand size = 14, antiderivative size = 39

$$\int \sqrt{-2 + 5x - 3x^2} dx = -\frac{1}{12}(5 - 6x)\sqrt{-2 + 5x - 3x^2} - \frac{\arcsin(5 - 6x)}{24\sqrt{3}}$$

[Out] 1/72\*arcsin(-5+6\*x)\*3^(1/2)-1/12\*(5-6\*x)\*(-3\*x^2+5\*x-2)^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$ , Rules used = {626, 633, 222}

$$\int \sqrt{-2 + 5x - 3x^2} dx = -\frac{\arcsin(5 - 6x)}{24\sqrt{3}} - \frac{1}{12}\sqrt{-3x^2 + 5x - 2}(5 - 6x)$$

[In] Int[Sqrt[-2 + 5\*x - 3\*x^2],x]

[Out] -1/12\*((5 - 6\*x)\*Sqrt[-2 + 5\*x - 3\*x^2]) - ArcSin[5 - 6\*x]/(24\*Sqrt[3])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 626

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x) \* ((a + b\*x + c\*x^2)^p / (2\*c\*(2\*p + 1))), x] - Dist[p\*((b^2 - 4\*a\*c) / (2\*c\*(2\*p + 1))), Int[(a + b\*x + c\*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && N

$eQ[b^2 - 4ac, 0] \ \&\& \ GtQ[p, 0] \ \&\& \ IntegerQ[4p]$

### Rule 633

$\text{Int}[(a_.) + (b_.)x + (c_.)x^2]^{(p)}, x\_Symbol] \rightarrow \text{Dist}[1/(2c(-4c/(b^2 - 4ac)))^p, \text{Subst}[\text{Int}[\text{Simp}[1 - x^2/(b^2 - 4ac)], x]^p, x], x, b + 2cx], x] \ /; \text{FreeQ}\{a, b, c, p\}, x] \ \&\& \ GtQ[4a - b^2/c, 0]$

### Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{1}{12}(5 - 6x)\sqrt{-2 + 5x - 3x^2} + \frac{1}{24} \int \frac{1}{\sqrt{-2 + 5x - 3x^2}} dx \\ &= -\frac{1}{12}(5 - 6x)\sqrt{-2 + 5x - 3x^2} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, 5 - 6x\right)}{24\sqrt{3}} \\ &= -\frac{1}{12}(5 - 6x)\sqrt{-2 + 5x - 3x^2} - \frac{\sin^{-1}(5 - 6x)}{24\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.44

$$\begin{aligned} &\int \sqrt{-2 + 5x - 3x^2} dx \\ &= \frac{1}{36} \left( 3(-5 + 6x)\sqrt{-2 + 5x - 3x^2} - \sqrt{3} \arctan\left(\frac{\sqrt{-6 + 15x - 9x^2}}{-2 + 3x}\right) \right) \end{aligned}$$

[In] Integrate[Sqrt[-2 + 5\*x - 3\*x^2],x]

[Out] (3\*(-5 + 6\*x)\*Sqrt[-2 + 5\*x - 3\*x^2] - Sqrt[3]\*ArcTan[Sqrt[-6 + 15\*x - 9\*x^2]/(-2 + 3\*x)]/36

### Maple [A] (verified)

Time = 2.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

method	result
default	$\frac{\arcsin(-5+6x)\sqrt{3}}{72} - \frac{(5-6x)\sqrt{-3x^2+5x-2}}{12}$
risch	$-\frac{(3x^2-5x+2)(-5+6x)}{12\sqrt{-3x^2+5x-2}} + \frac{\arcsin(-5+6x)\sqrt{3}}{72}$
trager	$\left(-\frac{5}{12} + \frac{x}{2}\right)\sqrt{-3x^2+5x-2} + \frac{\text{RootOf}(\_Z^2+3)\ln(-6x\text{RootOf}(\_Z^2+3)+5\text{RootOf}(\_Z^2+3)+6\sqrt{-3x^2+5x-2})}{72}$



[In] `int((-3*x^2+5*x-2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/72*arcsin(-5+6*x)*3^(1/2)-1/12*(5-6*x)*(-3*x^2+5*x-2)^(1/2)`

### Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.54

$$\int \sqrt{-2 + 5x - 3x^2} dx = \frac{1}{12} \sqrt{-3x^2 + 5x - 2}(6x - 5) - \frac{1}{72} \sqrt{3} \arctan \left( \frac{\sqrt{3} \sqrt{-3x^2 + 5x - 2}(6x - 5)}{6(3x^2 - 5x + 2)} \right)$$

[In] `integrate((-3*x^2+5*x-2)^(1/2),x, algorithm="fricas")`

[Out] `1/12*sqrt(-3*x^2 + 5*x - 2)*(6*x - 5) - 1/72*sqrt(3)*arctan(1/6*sqrt(3)*sqrt(-3*x^2 + 5*x - 2)*(6*x - 5)/(3*x^2 - 5*x + 2))`

### Sympy [A] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.82

$$\int \sqrt{-2 + 5x - 3x^2} dx = \left( \frac{x}{2} - \frac{5}{12} \right) \sqrt{-3x^2 + 5x - 2} + \frac{\sqrt{3} \operatorname{asin}(6x - 5)}{72}$$

[In] `integrate((-3*x**2+5*x-2)**(1/2),x)`

[Out] `(x/2 - 5/12)*sqrt(-3*x**2 + 5*x - 2) + sqrt(3)*asin(6*x - 5)/72`

### Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.05

$$\int \sqrt{-2 + 5x - 3x^2} dx = \frac{1}{2} \sqrt{-3x^2 + 5x - 2}x + \frac{1}{72} \sqrt{3} \arcsin(6x - 5) - \frac{5}{12} \sqrt{-3x^2 + 5x - 2}$$

[In] `integrate((-3*x^2+5*x-2)^(1/2),x, algorithm="maxima")`

[Out] `1/2*sqrt(-3*x^2 + 5*x - 2)*x + 1/72*sqrt(3)*arcsin(6*x - 5) - 5/12*sqrt(-3*x^2 + 5*x - 2)`

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.79

$$\int \sqrt{-2 + 5x - 3x^2} dx = \frac{1}{12} \sqrt{-3x^2 + 5x - 2}(6x - 5) + \frac{1}{72} \sqrt{3} \arcsin(6x - 5)$$

[In] integrate((-3\*x^2+5\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(-3\*x^2 + 5\*x - 2)\*(6\*x - 5) + 1/72\*sqrt(3)\*arcsin(6\*x - 5)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.77

$$\int \sqrt{-2 + 5x - 3x^2} dx = \frac{\sqrt{3} \operatorname{asin}(6x - 5)}{72} + \left( \frac{x}{2} - \frac{5}{12} \right) \sqrt{-3x^2 + 5x - 2}$$

[In] int((5\*x - 3\*x^2 - 2)^(1/2),x)

[Out] (3^(1/2)\*asin(6\*x - 5))/72 + (x/2 - 5/12)\*(5\*x - 3\*x^2 - 2)^(1/2)

### 3.114 $\int \frac{1}{\sqrt{5-6x+9x^2}} dx$

Optimal result	563
Rubi [A] (verified)	563
Mathematica [A] (verified)	564
Maple [A] (verified)	564
Fricas [B] (verification not implemented)	564
Sympy [A] (verification not implemented)	565
Maxima [A] (verification not implemented)	565
Giac [B] (verification not implemented)	565
Mupad [B] (verification not implemented)	565

#### Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{1}{3} \operatorname{arcsinh}\left(\frac{1}{2}(-1+3x)\right)$$

[Out] 1/3\*arcsinh(-1/2+3/2\*x)

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 221}

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{1}{3} \operatorname{arcsinh}\left(\frac{1}{2}(3x-1)\right)$$

[In] Int[1/Sqrt[5 - 6\*x + 9\*x^2], x]

[Out] ArcSinh[(-1 + 3\*x)/2]/3

#### Rule 221

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{36} \text{Subst} \left( \int \frac{1}{\sqrt{1 + \frac{x^2}{144}}} dx, x, -6 + 18x \right) \\ &= \frac{1}{3} \sinh^{-1} \left( \frac{1}{2}(-1 + 3x) \right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.71

$$\int \frac{1}{\sqrt{5 - 6x + 9x^2}} dx = -\frac{1}{3} \log \left( 1 - 3x + \sqrt{5 - 6x + 9x^2} \right)$$

[In] Integrate[1/Sqrt[5 - 6\*x + 9\*x^2],x]

[Out] -1/3\*Log[1 - 3\*x + Sqrt[5 - 6\*x + 9\*x^2]]

**Maple [A] (verified)**

Time = 2.26 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.64

method	result	size
default	$\frac{\text{arcsinh}\left(-\frac{1}{2} + \frac{3x}{2}\right)}{3}$	9
trager	$\frac{\ln\left(-1 + 3x + \sqrt{9x^2 - 6x + 5}\right)}{3}$	21

[In] int(1/(9\*x^2-6\*x+5)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*arcsinh(-1/2+3/2\*x)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. 2(8) = 16.

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{\sqrt{5 - 6x + 9x^2}} dx = -\frac{1}{3} \log \left( -3x + \sqrt{9x^2 - 6x + 5} + 1 \right)$$

[In] integrate(1/(9\*x^2-6\*x+5)^(1/2),x, algorithm="fricas")

[Out] -1/3\*log(-3\*x + sqrt(9\*x^2 - 6\*x + 5) + 1)

**Sympy [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{\operatorname{asinh}\left(\frac{3x}{2} - \frac{1}{2}\right)}{3}$$

[In] integrate(1/(9\*x\*\*2-6\*x+5)\*\*(1/2),x)

[Out] asinh(3\*x/2 - 1/2)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.57

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{1}{3} \operatorname{arsinh}\left(\frac{3}{2}x - \frac{1}{2}\right)$$

[In] integrate(1/(9\*x^2-6\*x+5)^(1/2),x, algorithm="maxima")

[Out] 1/3\*arcsinh(3/2\*x - 1/2)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(8) = 16.

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.86

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{1}{6} \sqrt{9x^2 - 6x + 5}(3x - 1) - \frac{2}{3} \log\left(-3x + \sqrt{9x^2 - 6x + 5} + 1\right)$$

[In] integrate(1/(9\*x^2-6\*x+5)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(9\*x^2 - 6\*x + 5)\*(3\*x - 1) - 2/3\*log(-3\*x + sqrt(9\*x^2 - 6\*x + 5) + 1)

**Mupad [B] (verification not implemented)**

Time = 9.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.43

$$\int \frac{1}{\sqrt{5-6x+9x^2}} dx = \frac{\ln\left(3x + \sqrt{9x^2 - 6x + 5} - 1\right)}{3}$$

[In] int(1/(9\*x^2 - 6\*x + 5)^(1/2),x)

[Out] log(3\*x + (9\*x^2 - 6\*x + 5)^(1/2) - 1)/3

### 3.115 $\int \frac{1}{\sqrt{3-4x-4x^2}} dx$

Optimal result	566
Rubi [A] (verified)	566
Mathematica [B] (verified)	567
Maple [A] (verified)	567
Fricas [B] (verification not implemented)	568
Sympy [A] (verification not implemented)	568
Maxima [A] (verification not implemented)	568
Giac [B] (verification not implemented)	568
Mupad [B] (verification not implemented)	569

#### Optimal result

Integrand size = 14, antiderivative size = 10

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = \frac{1}{2} \arcsin\left(\frac{1}{2} + x\right)$$

[Out] 1/2\*arcsin(1/2+x)

#### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = \frac{1}{2} \arcsin\left(x + \frac{1}{2}\right)$$

[In] Int[1/Sqrt[3 - 4\*x - 4\*x^2], x]

[Out] ArcSin[1/2 + x]/2

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= - \left( \frac{1}{16} \text{Subst} \left( \int \frac{1}{\sqrt{1 - \frac{x^2}{64}}} dx, x, -4 - 8x \right) \right) \\ &= \frac{1}{2} \sin^{-1} \left( \frac{1}{2} + x \right) \end{aligned}$$

## Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 25 vs. 2(10) = 20.

Time = 0.08 (sec) , antiderivative size = 25, normalized size of antiderivative = 2.50

$$\int \frac{1}{\sqrt{3 - 4x - 4x^2}} dx = -\arctan \left( \frac{\sqrt{3 - 4x - 4x^2}}{3 + 2x} \right)$$

[In] Integrate[1/Sqrt[3 - 4\*x - 4\*x^2],x]

[Out] -ArcTan[Sqrt[3 - 4\*x - 4\*x^2]/(3 + 2\*x)]

## Maple [A] (verified)

Time = 2.42 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

method	result	size
default	$\frac{\arcsin(x + \frac{1}{2})}{2}$	7
trager	$-\frac{\text{RootOf}(\_Z^2 + 1) \ln(2 \text{RootOf}(\_Z^2 + 1)x + \sqrt{-4x^2 - 4x + 3} + \text{RootOf}(\_Z^2 + 1))}{2}$	38

[In] int(1/(-4\*x^2-4\*x+3)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/2\*arcsin(x+1/2)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 33 vs.  $2(6) = 12$ .

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 3.30

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = -\frac{1}{2} \arctan\left(\frac{\sqrt{-4x^2-4x+3}(2x+1)}{4x^2+4x-3}\right)$$

[In] integrate(1/(-4\*x^2-4\*x+3)^(1/2),x, algorithm="fricas")

[Out] -1/2\*arctan(sqrt(-4\*x^2 - 4\*x + 3)\*(2\*x + 1)/(4\*x^2 + 4\*x - 3))

**Sympy [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = \frac{\operatorname{asin}\left(x + \frac{1}{2}\right)}{2}$$

[In] integrate(1/(-4\*x\*\*2-4\*x+3)\*\*(1/2),x)

[Out] asin(x + 1/2)/2

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = -\frac{1}{2} \arcsin\left(-x - \frac{1}{2}\right)$$

[In] integrate(1/(-4\*x^2-4\*x+3)^(1/2),x, algorithm="maxima")

[Out] -1/2\*arcsin(-x - 1/2)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 24 vs.  $2(6) = 12$ .

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 2.40

$$\int \frac{1}{\sqrt{3-4x-4x^2}} dx = \frac{1}{4} \sqrt{-4x^2-4x+3}(2x+1) + \arcsin\left(x + \frac{1}{2}\right)$$

[In] integrate(1/(-4\*x^2-4\*x+3)^(1/2),x, algorithm="giac")

[Out] 1/4\*sqrt(-4\*x^2 - 4\*x + 3)\*(2\*x + 1) + arcsin(x + 1/2)



**Mupad [B] (verification not implemented)**

Time = 8.96 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \frac{1}{\sqrt{3 - 4x - 4x^2}} dx = \frac{\operatorname{asin}\left(x + \frac{1}{2}\right)}{2}$$

[In] `int(1/(3 - 4*x^2 - 4*x)^(1/2),x)`

[Out] `asin(x + 1/2)/2`

### 3.116 $\int \frac{1}{\sqrt{-8+6x+9x^2}} dx$

Optimal result	570
Rubi [A] (verified)	570
Mathematica [A] (verified)	571
Maple [A] (verified)	571
Fricas [A] (verification not implemented)	571
Sympy [A] (verification not implemented)	572
Maxima [A] (verification not implemented)	572
Giac [A] (verification not implemented)	572
Mupad [B] (verification not implemented)	572

#### Optimal result

Integrand size = 14, antiderivative size = 25

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{1}{3} \operatorname{arctanh}\left(\frac{1+3x}{\sqrt{-8+6x+9x^2}}\right)$$

[Out] 1/3\*arctanh((1+3\*x)/(9\*x^2+6\*x-8)^(1/2))

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {635, 212}

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = \frac{1}{3} \operatorname{arctanh}\left(\frac{3x+1}{\sqrt{9x^2+6x-8}}\right)$$

[In] Int[1/Sqrt[-8 + 6\*x + 9\*x^2], x]

[Out] ArcTanh[(1 + 3\*x)/Sqrt[-8 + 6\*x + 9\*x^2]]/3

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 635

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(4\*c - x^2), x], x, (b + 2\*c\*x)/Sqrt[a + b\*x + c\*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{36-x^2} dx, x, \frac{6+18x}{\sqrt{-8+6x+9x^2}}\right) \\ &= \frac{1}{3} \tanh^{-1}\left(\frac{1+3x}{\sqrt{-8+6x+9x^2}}\right) \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = -\frac{1}{3} \log\left(-1-3x+\sqrt{-8+6x+9x^2}\right)$$

[In] Integrate[1/Sqrt[-8 + 6\*x + 9\*x^2], x]

[Out] -1/3\*Log[-1 - 3\*x + Sqrt[-8 + 6\*x + 9\*x^2]]

### Maple [A] (verified)

Time = 2.67 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

method	result	size
trager	$\frac{\ln(\sqrt{9x^2+6x-8}+1+3x)}{3}$	21
default	$\frac{\ln\left(\frac{(9x+3)\sqrt{9}+\sqrt{9x^2+6x-8}}{9}\right)\sqrt{9}}{9}$	30

[In] int(1/(9\*x^2+6\*x-8)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/3\*ln((9\*x^2+6\*x-8)^(1/2)+1+3\*x)

### Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-8+6x+9x^2}} dx = -\frac{1}{3} \log\left(-3x+\sqrt{9x^2+6x-8}-1\right)$$

[In] integrate(1/(9\*x^2+6\*x-8)^(1/2), x, algorithm="fricas")

[Out] -1/3\*log(-3\*x + sqrt(9\*x^2 + 6\*x - 8) - 1)

**Sympy [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx = \frac{\log(18x + 6\sqrt{9x^2 + 6x - 8} + 6)}{3}$$

[In] integrate(1/(9\*x\*\*2+6\*x-8)\*\*(1/2),x)

[Out] log(18\*x + 6\*sqrt(9\*x\*\*2 + 6\*x - 8) + 6)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx = \frac{1}{3} \log(18x + 6\sqrt{9x^2 + 6x - 8} + 6)$$

[In] integrate(1/(9\*x^2+6\*x-8)^(1/2),x, algorithm="maxima")

[Out] 1/3\*log(18\*x + 6\*sqrt(9\*x^2 + 6\*x - 8) + 6)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.64

$$\int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx = \frac{1}{6} \sqrt{9x^2 + 6x - 8}(3x + 1) + \frac{3}{2} \log\left(\left|-3x + \sqrt{9x^2 + 6x - 8} - 1\right|\right)$$

[In] integrate(1/(9\*x^2+6\*x-8)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(9\*x^2 + 6\*x - 8)\*(3\*x + 1) + 3/2\*log(abs(-3\*x + sqrt(9\*x^2 + 6\*x - 8) - 1))

**Mupad [B] (verification not implemented)**

Time = 9.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-8 + 6x + 9x^2}} dx = \frac{\ln(3x + \sqrt{9x^2 + 6x - 8} + 1)}{3}$$

[In] int(1/(6\*x + 9\*x^2 - 8)^(1/2),x)

[Out] log(3\*x + (6\*x + 9\*x^2 - 8)^(1/2) + 1)/3

### 3.117 $\int \frac{1}{\sqrt{2+4x+3x^2}} dx$

Optimal result . . . . .	573
Rubi [A] (verified) . . . . .	573
Mathematica [A] (verified) . . . . .	574
Maple [A] (verified) . . . . .	574
Fricas [B] (verification not implemented) . . . . .	575
Sympy [A] (verification not implemented) . . . . .	575
Maxima [A] (verification not implemented) . . . . .	575
Giac [B] (verification not implemented) . . . . .	576
Mupad [B] (verification not implemented) . . . . .	576

#### Optimal result

Integrand size = 14, antiderivative size = 18

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{\operatorname{arcsinh}\left(\frac{2+3x}{\sqrt{2}}\right)}{\sqrt{3}}$$

[Out] 1/3\*arcsinh(1/2\*(2+3\*x)\*2^(1/2))\*3^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 221}

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{\operatorname{arcsinh}\left(\frac{3x+2}{\sqrt{2}}\right)}{\sqrt{3}}$$

[In] Int[1/Sqrt[2 + 4\*x + 3\*x^2], x]

[Out] ArcSinh[(2 + 3\*x)/Sqrt[2]]/Sqrt[3]

#### Rule 221

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSinh[Rt[b, 2]\*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{x^2}{8}}} dx, x, 4+6x\right)}{2\sqrt{6}} \\ &= \frac{\sinh^{-1}\left(\frac{2+3x}{\sqrt{2}}\right)}{\sqrt{3}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.09 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.50

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = -\frac{\log(-2-3x+\sqrt{6+12x+9x^2})}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[2 + 4\*x + 3\*x^2],x]

[Out] -(Log[-2 - 3\*x + Sqrt[6 + 12\*x + 9\*x^2]]/Sqrt[3])

**Maple [A] (verified)**

Time = 2.41 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{3} \operatorname{arcsinh}\left(\frac{3\sqrt{2}\left(\frac{2}{3}+x\right)}{2}\right)}{3}$	15
trager	$\frac{\operatorname{RootOf}\left(-Z^2-3\right) \ln\left(3 \operatorname{RootOf}\left(-Z^2-3\right) x+2 \operatorname{RootOf}\left(-Z^2-3\right)+3 \sqrt{3 x^2+4 x+2}\right)}{3}$	42

[In] int(1/(3\*x^2+4\*x+2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*3^(1/2)\*arcsinh(3/2\*2^(1/2)\*(2/3+x))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 38 vs.  $2(16) = 32$ .

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.11

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{1}{6} \sqrt{3} \log \left( -\sqrt{3} \sqrt{3x^2+4x+2} (3x+2) - 9x^2 - 12x - 5 \right)$$

[In] integrate(1/(3\*x^2+4\*x+2)^(1/2),x, algorithm="fricas")

[Out] 1/6\*sqrt(3)\*log(-sqrt(3)\*sqrt(3\*x^2 + 4\*x + 2)\*(3\*x + 2) - 9\*x^2 - 12\*x - 5)

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{\sqrt{3} \operatorname{asinh} \left( \frac{3\sqrt{2}(x+\frac{2}{3})}{2} \right)}{3}$$

[In] integrate(1/(3\*x\*\*2+4\*x+2)\*\*(1/2),x)

[Out] sqrt(3)\*asinh(3\*sqrt(2)\*(x + 2/3)/2)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{1}{3} \sqrt{3} \operatorname{arsinh} \left( \frac{1}{2} \sqrt{2} (3x+2) \right)$$

[In] integrate(1/(3\*x^2+4\*x+2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*sqrt(3)\*arcsinh(1/2\*sqrt(2)\*(3\*x + 2))

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(16) = 32.

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 2.94

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{1}{6} \sqrt{3x^2+4x+2}(3x+2) - \frac{1}{9} \sqrt{3} \log\left(-\sqrt{3}\left(\sqrt{3}x - \sqrt{3x^2+4x+2}\right) - 2\right)$$

[In] integrate(1/(3\*x^2+4\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(3\*x^2 + 4\*x + 2)\*(3\*x + 2) - 1/9\*sqrt(3)\*log(-sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 4\*x + 2)) - 2)

**Mupad [B] (verification not implemented)**

Time = 9.16 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.44

$$\int \frac{1}{\sqrt{2+4x+3x^2}} dx = \frac{\sqrt{3} \ln\left(\sqrt{3}\left(x + \frac{2}{3}\right) + \sqrt{3x^2+4x+2}\right)}{3}$$

[In] int(1/(4\*x + 3\*x^2 + 2)^(1/2),x)

[Out] (3^(1/2)\*log(3^(1/2)\*(x + 2/3) + (4\*x + 3\*x^2 + 2)^(1/2)))/3



### 3.118 $\int \frac{1}{\sqrt{2+4x-3x^2}} dx$

Optimal result	577
Rubi [A] (verified)	577
Mathematica [B] (verified)	578
Maple [A] (verified)	578
Fricas [B] (verification not implemented)	579
Sympy [A] (verification not implemented)	579
Maxima [A] (verification not implemented)	579
Giac [B] (verification not implemented)	580
Mupad [B] (verification not implemented)	580

#### Optimal result

Integrand size = 14, antiderivative size = 19

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = -\frac{\arcsin\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{3}}$$

[Out]  $-1/3*\arcsin(1/10*(2-3*x)*10^{(1/2)})*3^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = -\frac{\arcsin\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{3}}$$

[In] Int[1/Sqrt[2 + 4\*x - 3\*x^2], x]

[Out] -(ArcSin[(2 - 3\*x)/Sqrt[10]]/Sqrt[3])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{40}}} dx, x, 4-6x\right)}{2\sqrt{30}} \\ &= -\frac{\sin^{-1}\left(\frac{2-3x}{\sqrt{10}}\right)}{\sqrt{3}} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 39 vs. 2(19) = 38.

Time = 0.11 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.05

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{3}x}{\sqrt{2}-\sqrt{2+4x-3x^2}}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[2 + 4\*x - 3\*x^2],x]

[Out] (-2\*ArcTan[(Sqrt[3]\*x)/(Sqrt[2] - Sqrt[2 + 4\*x - 3\*x^2])])/Sqrt[3]

### Maple [A] (verified)

Time = 2.37 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{\sqrt{3} \arcsin\left(\frac{3\sqrt{10}\left(-\frac{2}{3}+x\right)}{10}\right)}{3}$	15
trager	$-\frac{\text{RootOf}\left(\_Z^2+3\right) \ln\left(3x \text{RootOf}\left(\_Z^2+3\right)+3\sqrt{-3x^2+4x+2}-2 \text{RootOf}\left(\_Z^2+3\right)\right)}{3}$	42

[In] int(1/(-3\*x^2+4\*x+2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*3^(1/2)\*arcsin(3/10\*10^(1/2)\*(-2/3+x))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(16) = 32.

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.11

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = -\frac{1}{3} \sqrt{3} \arctan \left( \frac{\sqrt{3} \sqrt{-3x^2+4x+2}(3x-2)}{3(3x^2-4x-2)} \right)$$

[In] integrate(1/(-3\*x^2+4\*x+2)^(1/2),x, algorithm="fricas")

[Out] -1/3\*sqrt(3)\*arctan(1/3\*sqrt(3)\*sqrt(-3\*x^2 + 4\*x + 2)\*(3\*x - 2)/(3\*x^2 - 4\*x - 2))

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.05

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin} \left( \frac{3\sqrt{10}(x-\frac{2}{3})}{10} \right)}{3}$$

[In] integrate(1/(-3\*x\*\*2+4\*x+2)\*\*(1/2),x)

[Out] sqrt(3)\*asin(3\*sqrt(10)\*(x - 2/3)/10)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = -\frac{1}{3} \sqrt{3} \arcsin \left( -\frac{1}{10} \sqrt{10}(3x-2) \right)$$

[In] integrate(1/(-3\*x^2+4\*x+2)^(1/2),x, algorithm="maxima")

[Out] -1/3\*sqrt(3)\*arcsin(-1/10\*sqrt(10)\*(3\*x - 2))

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(16) = 32.

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.89

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = \frac{1}{6} \sqrt{-3x^2+4x+2}(3x-2) + \frac{5}{9} \sqrt{3} \arcsin\left(\frac{1}{10} \sqrt{10}(3x-2)\right)$$

[In] integrate(1/(-3\*x^2+4\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(-3\*x^2 + 4\*x + 2)\*(3\*x - 2) + 5/9\*sqrt(3)\*arcsin(1/10\*sqrt(10)\*(3\*x - 2))

**Mupad [B] (verification not implemented)**

Time = 9.30 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{2+4x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin}\left(\frac{\sqrt{40}(6x-4)}{40}\right)}{3}$$

[In] int(1/(4\*x - 3\*x^2 + 2)^(1/2),x)

[Out] (3^(1/2)\*asin((40^(1/2)\*(6\*x - 4))/40))/3

### 3.119 $\int \frac{1}{\sqrt{2+5x+3x^2}} dx$

Optimal result	581
Rubi [A] (verified)	581
Mathematica [A] (verified)	582
Maple [A] (verified)	582
Fricas [A] (verification not implemented)	583
Sympy [A] (verification not implemented)	583
Maxima [A] (verification not implemented)	583
Giac [A] (verification not implemented)	583
Mupad [B] (verification not implemented)	584

#### Optimal result

Integrand size = 14, antiderivative size = 35

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{5+6x}{2\sqrt{3}\sqrt{2+5x+3x^2}}\right)}{\sqrt{3}}$$

[Out] 1/3\*arctanh(1/6\*(5+6\*x)\*3^(1/2)/(3\*x^2+5\*x+2)^(1/2))\*3^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {635, 212}

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{6x+5}{2\sqrt{3}\sqrt{3x^2+5x+2}}\right)}{\sqrt{3}}$$

[In] Int[1/Sqrt[2 + 5\*x + 3\*x^2], x]

[Out] ArcTanh[(5 + 6\*x)/(2\*Sqrt[3]\*Sqrt[2 + 5\*x + 3\*x^2])]/Sqrt[3]

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 635

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(4\*c - x^2), x], x, (b + 2\*c\*x)/Sqrt[a + b\*x + c\*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{5 + 6x}{\sqrt{2 + 5x + 3x^2}}\right) \\ &= \frac{\tanh^{-1}\left(\frac{5+6x}{2\sqrt{3}\sqrt{2+5x+3x^2}}\right)}{\sqrt{3}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{2 + 5x + 3x^2}} dx = \frac{2\text{arctanh}\left(\frac{\sqrt{\frac{2}{3} + \frac{5x}{3} + x^2}}{1+x}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[2 + 5\*x + 3\*x^2], x]

[Out] (2\*ArcTanh[Sqrt[2/3 + (5\*x)/3 + x^2]/(1 + x)]/Sqrt[3]

**Maple [A] (verified)**

Time = 2.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{\ln\left(\frac{\left(\frac{5}{2}+3x\right)\sqrt{3}}{3} + \sqrt{3x^2+5x+2}\right)\sqrt{3}}{3}$	30
trager	$\frac{\text{RootOf}\left(\_Z^2-3\right)\ln\left(6\text{RootOf}\left(\_Z^2-3\right)x+5\text{RootOf}\left(\_Z^2-3\right)+6\sqrt{3x^2+5x+2}\right)}{3}$	42

[In] int(1/(3\*x^2+5\*x+2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/3\*ln(1/3\*(5/2+3\*x)\*3^(1/2)+(3\*x^2+5\*x+2)^(1/2))\*3^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{1}{6} \sqrt{3} \log \left( 4 \sqrt{3} \sqrt{3x^2+5x+2} (6x+5) + 72x^2 + 120x + 49 \right)$$

[In] integrate(1/(3\*x^2+5\*x+2)^(1/2),x, algorithm="fricas")

[Out] 1/6\*sqrt(3)\*log(4\*sqrt(3)\*sqrt(3\*x^2 + 5\*x + 2)\*(6\*x + 5) + 72\*x^2 + 120\*x + 49)

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+5x+2} + 5)}{3}$$

[In] integrate(1/(3\*x\*\*2+5\*x+2)\*\*(1/2),x)

[Out] sqrt(3)\*log(6\*x + 2\*sqrt(3)\*sqrt(3\*x\*\*2 + 5\*x + 2) + 5)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{1}{3} \sqrt{3} \log \left( 2 \sqrt{3} \sqrt{3x^2+5x+2} + 6x + 5 \right)$$

[In] integrate(1/(3\*x^2+5\*x+2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 5\*x + 2) + 6\*x + 5)

**Giac [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int \frac{1}{\sqrt{2+5x+3x^2}} dx = \frac{1}{12} \sqrt{3x^2+5x+2} (6x+5) + \frac{1}{72} \sqrt{3} \log \left( \left| -2 \sqrt{3} \left( \sqrt{3x} - \sqrt{3x^2+5x+2} \right) - 5 \right| \right)$$

[In] integrate(1/(3\*x^2+5\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(3\*x^2 + 5\*x + 2)\*(6\*x + 5) + 1/72\*sqrt(3)\*log(abs(-2\*sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 5\*x + 2)) - 5))

### Mupad [B] (verification not implemented)

Time = 9.32 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{2 + 5x + 3x^2}} dx = \frac{\sqrt{3} \ln(\sqrt{3}(x + \frac{5}{6}) + \sqrt{3x^2 + 5x + 2})}{3}$$

[In] int(1/(5\*x + 3\*x^2 + 2)^(1/2),x)

[Out] (3^(1/2)\*log(3^(1/2)\*(x + 5/6) + (5\*x + 3\*x^2 + 2)^(1/2)))/3



### 3.120 $\int \frac{1}{\sqrt{2+5x-3x^2}} dx$

Optimal result . . . . .	585
Rubi [A] (verified) . . . . .	585
Mathematica [A] (verified) . . . . .	586
Maple [A] (verified) . . . . .	586
Fricas [B] (verification not implemented) . . . . .	586
Sympy [A] (verification not implemented) . . . . .	587
Maxima [A] (verification not implemented) . . . . .	587
Giac [B] (verification not implemented) . . . . .	587
Mupad [B] (verification not implemented) . . . . .	587

#### Optimal result

Integrand size = 14, antiderivative size = 17

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = -\frac{\arcsin\left(\frac{1}{7}(5-6x)\right)}{\sqrt{3}}$$

[Out] 1/3\*arcsin(-5/7+6/7\*x)\*3^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = -\frac{\arcsin\left(\frac{1}{7}(5-6x)\right)}{\sqrt{3}}$$

[In] Int[1/Sqrt[2 + 5\*x - 3\*x^2],x]

[Out] -(ArcSin[(5 - 6\*x)/7]/Sqrt[3])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_)^2], x\_Symbol] :> Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] :> Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{49}}} dx, x, 5-6x\right)}{7\sqrt{3}} \\ &= -\frac{\sin^{-1}\left(\frac{1}{7}(5-6x)\right)}{\sqrt{3}} \end{aligned}$$

### Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.76

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{6+15x-9x^2}}{1+3x}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[2 + 5\*x - 3\*x^2],x]

[Out] (-2\*ArcTan[Sqrt[6 + 15\*x - 9\*x^2]/(1 + 3\*x)])/Sqrt[3]

### Maple [A] (verified)

Time = 2.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.71

method	result	size
default	$\frac{\arcsin\left(-\frac{5}{7}+\frac{6x}{7}\right)\sqrt{3}}{3}$	12
trager	$\frac{\text{RootOf}\left(\_Z^2+3\right)\ln\left(-6x\text{RootOf}\left(\_Z^2+3\right)+6\sqrt{-3x^2+5x+2}+5\text{RootOf}\left(\_Z^2+3\right)\right)}{3}$	42

[In] int(1/(-3\*x^2+5\*x+2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*arcsin(-5/7+6/7\*x)\*3^(1/2)

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(11) = 22.

Time = 0.26 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.35

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = -\frac{1}{3}\sqrt{3}\arctan\left(\frac{\sqrt{3}\sqrt{-3x^2+5x+2}(6x-5)}{6(3x^2-5x-2)}\right)$$

[In] integrate(1/(-3\*x^2+5\*x+2)^(1/2),x, algorithm="fricas")

[Out] -1/3\*sqrt(3)\*arctan(1/6\*sqrt(3)\*sqrt(-3\*x^2 + 5\*x + 2)\*(6\*x - 5)/(3\*x^2 - 5\*x - 2))

**Sympy [A] (verification not implemented)**

Time = 0.31 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin}\left(\frac{6x}{7} - \frac{5}{7}\right)}{3}$$

[In] integrate(1/(-3\*x\*\*2+5\*x+2)\*\*(1/2),x)

[Out] sqrt(3)\*asin(6\*x/7 - 5/7)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = -\frac{1}{3} \sqrt{3} \arcsin\left(-\frac{6}{7}x + \frac{5}{7}\right)$$

[In] integrate(1/(-3\*x^2+5\*x+2)^(1/2),x, algorithm="maxima")

[Out] -1/3\*sqrt(3)\*arcsin(-6/7\*x + 5/7)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.82

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = \frac{1}{12} \sqrt{-3x^2+5x+2}(6x-5) + \frac{49}{72} \sqrt{3} \arcsin\left(\frac{6}{7}x - \frac{5}{7}\right)$$

[In] integrate(1/(-3\*x^2+5\*x+2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(-3\*x^2 + 5\*x + 2)\*(6\*x - 5) + 49/72\*sqrt(3)\*arcsin(6/7\*x - 5/7)

**Mupad [B] (verification not implemented)**

Time = 9.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.65

$$\int \frac{1}{\sqrt{2+5x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin}\left(\frac{6x}{7} - \frac{5}{7}\right)}{3}$$

[In] int(1/(5\*x - 3\*x^2 + 2)^(1/2),x)

[Out] (3^(1/2)\*asin((6\*x)/7 - 5/7))/3

### 3.121 $\int \frac{1}{\sqrt{-2+4x+3x^2}} dx$

Optimal result	588
Rubi [A] (verified)	588
Mathematica [A] (verified)	589
Maple [A] (verified)	589
Fricas [A] (verification not implemented)	590
Sympy [A] (verification not implemented)	590
Maxima [A] (verification not implemented)	590
Giac [A] (verification not implemented)	590
Mupad [B] (verification not implemented)	591

#### Optimal result

Integrand size = 14, antiderivative size = 32

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{2+3x}{\sqrt{3}\sqrt{-2+4x+3x^2}}\right)}{\sqrt{3}}$$

[Out] 1/3\*arctanh(1/3\*(2+3\*x)\*3^(1/2)/(3\*x^2+4\*x-2)^(1/2))\*3^(1/2)

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {635, 212}

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{3x+2}{\sqrt{3}\sqrt{3x^2+4x-2}}\right)}{\sqrt{3}}$$

[In] Int[1/Sqrt[-2 + 4\*x + 3\*x^2], x]

[Out] ArcTanh[(2 + 3\*x)/(Sqrt[3]\*Sqrt[-2 + 4\*x + 3\*x^2])]/Sqrt[3]

#### Rule 212

Int[((a\_) + (b\_.)\*(x\_)^2)^(-1), x\_Symbol] := Simp[(1/(Rt[a, 2]\*Rt[-b, 2]))\*ArcTanh[Rt[-b, 2]\*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

#### Rule 635

Int[1/Sqrt[(a\_) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2], x\_Symbol] := Dist[2, Subst[Int[1/(4\*c - x^2), x], x, (b + 2\*c\*x)/Sqrt[a + b\*x + c\*x^2]], x] /; FreeQ[{a,

b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{4 + 6x}{\sqrt{-2 + 4x + 3x^2}}\right) \\ &= \frac{\tanh^{-1}\left(\frac{2+3x}{\sqrt{3}\sqrt{-2+4x+3x^2}}\right)}{\sqrt{3}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.08 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.84

$$\int \frac{1}{\sqrt{-2 + 4x + 3x^2}} dx = -\frac{\log(-2 - 3x + \sqrt{-6 + 12x + 9x^2})}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[-2 + 4\*x + 3\*x^2],x]

[Out] -(Log[-2 - 3\*x + Sqrt[-6 + 12\*x + 9\*x^2]]/Sqrt[3])

**Maple [A] (verified)**

Time = 2.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.94

method	result	size
default	$\frac{\ln\left(\frac{(2+3x)\sqrt{3} + \sqrt{3x^2+4x-2}}{3}\right)\sqrt{3}}{3}$	30
trager	$\frac{\text{RootOf}(\_Z^2-3) \ln\left(3 \text{RootOf}(\_Z^2-3)x + 2 \text{RootOf}(\_Z^2-3) + 3\sqrt{3x^2+4x-2}\right)}{3}$	42

[In] int(1/(3\*x^2+4\*x-2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*ln(1/3\*(2+3\*x)\*3^(1/2)+(3\*x^2+4\*x-2)^(1/2))\*3^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.16

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{1}{6} \sqrt{3} \log \left( \sqrt{3} \sqrt{3x^2+4x-2} (3x+2) + 9x^2 + 12x - 1 \right)$$

[In] integrate(1/(3\*x^2+4\*x-2)^(1/2),x, algorithm="fricas")

[Out] 1/6\*sqrt(3)\*log(sqrt(3)\*sqrt(3\*x^2 + 4\*x - 2)\*(3\*x + 2) + 9\*x^2 + 12\*x - 1)

**Sympy [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+4x-2} + 4)}{3}$$

[In] integrate(1/(3\*x\*\*2+4\*x-2)\*\*(1/2),x)

[Out] sqrt(3)\*log(6\*x + 2\*sqrt(3)\*sqrt(3\*x\*\*2 + 4\*x - 2) + 4)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.31 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{1}{3} \sqrt{3} \log \left( 2\sqrt{3}\sqrt{3x^2+4x-2} + 6x + 4 \right)$$

[In] integrate(1/(3\*x^2+4\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 4\*x - 2) + 6\*x + 4)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.69

$$\int \frac{1}{\sqrt{-2+4x+3x^2}} dx = \frac{1}{6} \sqrt{3x^2+4x-2} (3x+2) + \frac{5}{9} \sqrt{3} \log \left( \left| -\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2+4x-2} \right) - 2 \right| \right)$$

[In] integrate(1/(3\*x^2+4\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/6\*sqrt(3\*x^2 + 4\*x - 2)\*(3\*x + 2) + 5/9\*sqrt(3)\*log(abs(-sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 4\*x - 2)) - 2))

**Mupad [B] (verification not implemented)**

Time = 9.11 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.81

$$\int \frac{1}{\sqrt{-2 + 4x + 3x^2}} dx = \frac{\sqrt{3} \ln \left( \sqrt{3} \left( x + \frac{2}{3} \right) + \sqrt{3x^2 + 4x - 2} \right)}{3}$$

[In] int(1/(4\*x + 3\*x^2 - 2)^(1/2),x)

[Out] (3^(1/2)\*log(3^(1/2)\*(x + 2/3) + (4\*x + 3\*x^2 - 2)^(1/2)))/3

### 3.122 $\int \frac{1}{\sqrt{-2+4x-3x^2}} dx$

Optimal result	592
Rubi [A] (verified)	592
Mathematica [A] (verified)	593
Maple [A] (verified)	593
Fricas [C] (verification not implemented)	594
Sympy [C] (verification not implemented)	594
Maxima [C] (verification not implemented)	594
Giac [F]	595
Mupad [B] (verification not implemented)	595

#### Optimal result

Integrand size = 14, antiderivative size = 33

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = -\frac{\arctan\left(\frac{2-3x}{\sqrt{3}\sqrt{-2+4x-3x^2}}\right)}{\sqrt{3}}$$

[Out]  $-1/3*\arctan(1/3*(2-3*x)*3^{(1/2)/(-3*x^2+4*x-2)^{(1/2)}}*3^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {635, 210}

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = -\frac{\arctan\left(\frac{2-3x}{\sqrt{3}\sqrt{-3x^2+4x-2}}\right)}{\sqrt{3}}$$

[In]  $\text{Int}[1/\text{Sqrt}[-2 + 4*x - 3*x^2], x]$

[Out]  $-(\text{ArcTan}[(2 - 3*x)/(\text{Sqrt}[3]*\text{Sqrt}[-2 + 4*x - 3*x^2])]/\text{Sqrt}[3])$

#### Rule 210

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

#### Rule 635

$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_) + (c_)*(x_)^2)], x\_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a,$



b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{-12 - x^2} dx, x, \frac{4 - 6x}{\sqrt{-2 + 4x - 3x^2}}\right) \\ &= -\frac{\tan^{-1}\left(\frac{2-3x}{\sqrt{3}\sqrt{-2+4x-3x^2}}\right)}{\sqrt{3}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.11 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{-2 + 4x - 3x^2}} dx = -\frac{\arctan\left(\frac{2-3x}{\sqrt{-6+12x-9x^2}}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[-2 + 4\*x - 3\*x^2],x]

[Out] -(ArcTan[(2 - 3\*x)/Sqrt[-6 + 12\*x - 9\*x^2]]/Sqrt[3])

**Maple [A] (verified)**

Time = 2.40 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{\sqrt{3} \arctan\left(\frac{\sqrt{3}\left(-\frac{2}{3}+x\right)}{\sqrt{-3x^2+4x-2}}\right)}{3}$	26
trager	$\frac{\text{RootOf}\left(\_Z^2+3\right) \ln\left(-3x \text{RootOf}\left(\_Z^2+3\right)+3\sqrt{-3x^2+4x-2}+2 \text{RootOf}\left(\_Z^2+3\right)\right)}{3}$	42

[In] int(1/(-3\*x^2+4\*x-2)^(1/2),x,method=\_RETURNVERBOSE)

[Out] 1/3\*3^(1/2)\*arctan(3^(1/2)\*(-2/3+x)/(-3\*x^2+4\*x-2)^(1/2))

**Fricas [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 67, normalized size of antiderivative = 2.03

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = \frac{1}{6}i\sqrt{3}\log\left(-\frac{2(i\sqrt{3}\sqrt{-3x^2+4x-2}+3x-2)}{x}\right) - \frac{1}{6}i\sqrt{3}\log\left(-\frac{2(-i\sqrt{3}\sqrt{-3x^2+4x-2}+3x-2)}{x}\right)$$

[In] integrate(1/(-3\*x^2+4\*x-2)^(1/2),x, algorithm="fricas")

[Out] 1/6\*I\*sqrt(3)\*log(-2\*(I\*sqrt(3)\*sqrt(-3\*x^2 + 4\*x - 2) + 3\*x - 2)/x) - 1/6\*I\*sqrt(3)\*log(-2\*(-I\*sqrt(3)\*sqrt(-3\*x^2 + 4\*x - 2) + 3\*x - 2)/x)

**Sympy [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = -\frac{\sqrt{3}i\log(-6x+2\sqrt{3}i\sqrt{-3x^2+4x-2}+4)}{3}$$

[In] integrate(1/(-3\*x\*\*2+4\*x-2)\*\*(1/2),x)

[Out] -sqrt(3)\*I\*log(-6\*x + 2\*sqrt(3)\*I\*sqrt(-3\*x\*\*2 + 4\*x - 2) + 4)/3

**Maxima [C] (verification not implemented)**

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.48

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = -\frac{1}{3}i\sqrt{3}\operatorname{arsinh}\left(\frac{1}{2}\sqrt{2}(3x-2)\right)$$

[In] integrate(1/(-3\*x^2+4\*x-2)^(1/2),x, algorithm="maxima")

[Out] -1/3\*I\*sqrt(3)\*arcsinh(1/2\*sqrt(2)\*(3\*x - 2))

**Giac [F]**

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = \int \frac{1}{\sqrt{-3x^2+4x-2}} dx$$

[In] integrate(1/(-3\*x^2+4\*x-2)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(-3\*x^2 + 4\*x - 2), x)

**Mupad [B] (verification not implemented)**

Time = 9.07 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.52

$$\int \frac{1}{\sqrt{-2+4x-3x^2}} dx = -\frac{\sqrt{3} \operatorname{asin}\left(\sqrt{2}\left(\frac{3x}{2}-1\right) \operatorname{li}\right)}{3}$$

[In] int(1/(4\*x - 3\*x^2 - 2)^(1/2),x)

[Out] -(3^(1/2)\*asin(2^(1/2)\*((3\*x)/2 - 1)\*1i))/3

### 3.123 $\int \frac{1}{\sqrt{-2+5x+3x^2}} dx$

Optimal result	596
Rubi [A] (verified)	596
Mathematica [A] (verified)	597
Maple [A] (verified)	597
Fricas [A] (verification not implemented)	598
Sympy [A] (verification not implemented)	598
Maxima [A] (verification not implemented)	598
Giac [A] (verification not implemented)	598
Mupad [B] (verification not implemented)	599

#### Optimal result

Integrand size = 14, antiderivative size = 35

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{5+6x}{2\sqrt{3}\sqrt{-2+5x+3x^2}}\right)}{\sqrt{3}}$$

[Out]  $1/3*\operatorname{arctanh}(1/6*(5+6*x)*3^{(1/2)}/(3*x^2+5*x-2)^{(1/2)})*3^{(1/2)}$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {635, 212}

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{\operatorname{arctanh}\left(\frac{6x+5}{2\sqrt{3}\sqrt{3x^2+5x-2}}\right)}{\sqrt{3}}$$

[In]  $\operatorname{Int}[1/\operatorname{Sqrt}[-2 + 5*x + 3*x^2], x]$

[Out]  $\operatorname{ArcTanh}[(5 + 6*x)/(2*\operatorname{Sqrt}[3]*\operatorname{Sqrt}[-2 + 5*x + 3*x^2])]/\operatorname{Sqrt}[3]$

#### Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x\_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

#### Rule 635

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_) + (c_)*(x_)^2)], x\_Symbol] \rightarrow \operatorname{Dist}[2, \operatorname{Subst}[\operatorname{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\operatorname{Sqrt}[a + b*x + c*x^2]], x] /; \operatorname{FreeQ}\{a,$

b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2\text{Subst}\left(\int \frac{1}{12 - x^2} dx, x, \frac{5 + 6x}{\sqrt{-2 + 5x + 3x^2}}\right) \\ &= \frac{\tanh^{-1}\left(\frac{5+6x}{2\sqrt{3}\sqrt{-2+5x+3x^2}}\right)}{\sqrt{3}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{-2 + 5x + 3x^2}} dx = \frac{2\text{arctanh}\left(\frac{\sqrt{-\frac{2}{3} + \frac{5x}{3} + x^2}}{2+x}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[-2 + 5\*x + 3\*x^2], x]

[Out] (2\*ArcTanh[Sqrt[-2/3 + (5\*x)/3 + x^2]/(2 + x)]/Sqrt[3]

**Maple [A] (verified)**

Time = 2.12 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.86

method	result	size
default	$\frac{\ln\left(\frac{\left(\frac{5}{2}+3x\right)\sqrt{3}}{3} + \sqrt{3x^2+5x-2}\right)\sqrt{3}}{3}$	30
trager	$\frac{\text{RootOf}\left(\_Z^2-3\right)\ln\left(6\text{RootOf}\left(\_Z^2-3\right)x+6\sqrt{3x^2+5x-2}+5\text{RootOf}\left(\_Z^2-3\right)\right)}{3}$	42

[In] int(1/(3\*x^2+5\*x-2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/3\*ln(1/3\*(5/2+3\*x)\*3^(1/2)+(3\*x^2+5\*x-2)^(1/2))\*3^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.09

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{1}{6} \sqrt{3} \log \left( 4 \sqrt{3} \sqrt{3x^2+5x-2} (6x+5) + 72x^2 + 120x + 1 \right)$$

[In] integrate(1/(3\*x^2+5\*x-2)^(1/2),x, algorithm="fricas")

[Out] 1/6\*sqrt(3)\*log(4\*sqrt(3)\*sqrt(3\*x^2 + 5\*x - 2)\*(6\*x + 5) + 72\*x^2 + 120\*x + 1)

**Sympy [A] (verification not implemented)**

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{\sqrt{3} \log(6x + 2\sqrt{3}\sqrt{3x^2+5x-2} + 5)}{3}$$

[In] integrate(1/(3\*x\*\*2+5\*x-2)\*\*(1/2),x)

[Out] sqrt(3)\*log(6\*x + 2\*sqrt(3)\*sqrt(3\*x\*\*2 + 5\*x - 2) + 5)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.80

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{1}{3} \sqrt{3} \log \left( 2 \sqrt{3} \sqrt{3x^2+5x-2} + 6x + 5 \right)$$

[In] integrate(1/(3\*x^2+5\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*sqrt(3)\*log(2\*sqrt(3)\*sqrt(3\*x^2 + 5\*x - 2) + 6\*x + 5)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int \frac{1}{\sqrt{-2+5x+3x^2}} dx = \frac{1}{12} \sqrt{3x^2+5x-2} (6x+5) + \frac{49}{72} \sqrt{3} \log \left( \left| -2\sqrt{3} \left( \sqrt{3}x - \sqrt{3x^2+5x-2} \right) - 5 \right| \right)$$

[In] integrate(1/(3\*x^2+5\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(3\*x^2 + 5\*x - 2)\*(6\*x + 5) + 49/72\*sqrt(3)\*log(abs(-2\*sqrt(3)\*(sqrt(3)\*x - sqrt(3\*x^2 + 5\*x - 2)) - 5))

### Mupad [B] (verification not implemented)

Time = 9.19 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{-2 + 5x + 3x^2}} dx = \frac{\sqrt{3} \ln(\sqrt{3}(x + \frac{5}{6}) + \sqrt{3x^2 + 5x - 2})}{3}$$

[In] int(1/(5\*x + 3\*x^2 - 2)^(1/2),x)

[Out] (3^(1/2)\*log(3^(1/2)\*(x + 5/6) + (5\*x + 3\*x^2 - 2)^(1/2)))/3

$$3.124 \quad \int \frac{1}{\sqrt{-2+5x-3x^2}} dx$$

Optimal result	600
Rubi [A] (verified)	600
Mathematica [B] (verified)	601
Maple [A] (verified)	601
Fricas [B] (verification not implemented)	602
Sympy [A] (verification not implemented)	602
Maxima [A] (verification not implemented)	602
Giac [B] (verification not implemented)	603
Mupad [B] (verification not implemented)	603

### Optimal result

Integrand size = 14, antiderivative size = 13

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = -\frac{\arcsin(5-6x)}{\sqrt{3}}$$

[Out] 1/3\*arcsin(-5+6\*x)\*3^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = -\frac{\arcsin(5-6x)}{\sqrt{3}}$$

[In] Int[1/Sqrt[-2 + 5\*x - 3\*x^2], x]

[Out] -(ArcSin[5 - 6\*x]/Sqrt[3])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b + 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]



Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, 5-6x\right)}{\sqrt{3}} \\ &= -\frac{\sin^{-1}(5-6x)}{\sqrt{3}} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 30 vs. 2(13) = 26.

Time = 0.09 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.31

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = -\frac{2 \arctan\left(\frac{\sqrt{-6+15x-9x^2}}{-2+3x}\right)}{\sqrt{3}}$$

[In] Integrate[1/Sqrt[-2 + 5\*x - 3\*x^2], x]

[Out] (-2\*ArcTan[Sqrt[-6 + 15\*x - 9\*x^2]/(-2 + 3\*x)]/Sqrt[3])

### Maple [A] (verified)

Time = 2.23 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$\frac{\arcsin(-5+6x)\sqrt{3}}{3}$	12
trager	$\frac{\text{RootOf}(\_Z^2+3) \ln(-6x \text{RootOf}(\_Z^2+3)+5 \text{RootOf}(\_Z^2+3)+6\sqrt{-3x^2+5x-2})}{3}$	42

[In] int(1/(-3\*x^2+5\*x-2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/3\*arcsin(-5+6\*x)\*3^(1/2)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 40 vs. 2(11) = 22.

Time = 0.25 (sec) , antiderivative size = 40, normalized size of antiderivative = 3.08

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = -\frac{1}{3} \sqrt{3} \arctan \left( \frac{\sqrt{3} \sqrt{-3x^2+5x-2}(6x-5)}{6(3x^2-5x+2)} \right)$$

[In] integrate(1/(-3\*x^2+5\*x-2)^(1/2),x, algorithm="fricas")

[Out] -1/3\*sqrt(3)\*arctan(1/6\*sqrt(3)\*sqrt(-3\*x^2 + 5\*x - 2)\*(6\*x - 5)/(3\*x^2 - 5\*x + 2))

**Sympy [A] (verification not implemented)**

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin}(6x-5)}{3}$$

[In] integrate(1/(-3\*x\*\*2+5\*x-2)\*\*(1/2),x)

[Out] sqrt(3)\*asin(6\*x - 5)/3

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = \frac{1}{3} \sqrt{3} \arcsin(6x-5)$$

[In] integrate(1/(-3\*x^2+5\*x-2)^(1/2),x, algorithm="maxima")

[Out] 1/3\*sqrt(3)\*arcsin(6\*x - 5)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(11) = 22.

Time = 0.26 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.38

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = \frac{1}{12} \sqrt{-3x^2+5x-2}(6x-5) + \frac{1}{72} \sqrt{3} \arcsin(6x-5)$$

[In] integrate(1/(-3\*x^2+5\*x-2)^(1/2),x, algorithm="giac")

[Out] 1/12\*sqrt(-3\*x^2 + 5\*x - 2)\*(6\*x - 5) + 1/72\*sqrt(3)\*arcsin(6\*x - 5)

**Mupad [B] (verification not implemented)**

Time = 9.09 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{1}{\sqrt{-2+5x-3x^2}} dx = \frac{\sqrt{3} \operatorname{asin}(6x-5)}{3}$$

[In] int(1/(5\*x - 3\*x^2 - 2)^(1/2),x)

[Out] (3^(1/2)\*asin(6\*x - 5))/3

$$3.125 \quad \int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx$$

Optimal result	604
Rubi [A] (verified)	604
Mathematica [A] (verified)	605
Maple [B] (verified)	605
Fricas [B] (verification not implemented)	606
Sympy [A] (verification not implemented)	606
Maxima [A] (verification not implemented)	607
Giac [B] (verification not implemented)	607
Mupad [B] (verification not implemented)	607

### Optimal result

Integrand size = 27, antiderivative size = 22

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = \frac{\operatorname{arcsinh}\left(\frac{b+2cx}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[Out]  $\operatorname{arcsinh}(1/2*(2*c*x+b)/c^{(1/2)})/c^{(1/2)}$

### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$ , Rules used = {633, 221}

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = \frac{\operatorname{arcsinh}\left(\frac{b+2cx}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[In]  $\operatorname{Int}[1/\operatorname{Sqrt}[(b^2 + 4*c)/(4*c) + b*x + c*x^2], x]$

[Out]  $\operatorname{ArcSinh}[(b + 2*c*x)/(2*\operatorname{Sqrt}[c])]/\operatorname{Sqrt}[c]$

#### Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b, x\} \&\& \operatorname{GtQ}[a, 0] \&\& \operatorname{PosQ}[b]$

#### Rule 633

$\operatorname{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2]^{(p_)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), \operatorname{Subst}[\operatorname{Int}[\operatorname{Simp}[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b]$

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{x^2}{4c}}} dx, x, b+2cx\right)}{2c} \\ &= \frac{\sinh^{-1}\left(\frac{b+2cx}{2\sqrt{c}}\right)}{\sqrt{c}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.21 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = -\frac{\log\left(b + 2cx - \sqrt{c}\sqrt{4 + \frac{b^2}{c} + 4bx + 4cx^2}\right)}{\sqrt{c}}$$

[In] Integrate[1/Sqrt[(b^2 + 4\*c)/(4\*c) + b\*x + c\*x^2], x]

[Out] -(Log[b + 2\*c\*x - Sqrt[c]\*Sqrt[4 + b^2/c + 4\*b\*x + 4\*c\*x^2]]/Sqrt[c])

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(16) = 32.

Time = 2.29 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.32

method	result	size
default	$\frac{\ln\left(\frac{(4cx+2b)\sqrt{4} + \sqrt{\frac{b^2+4c}{c} + 4bx + 4cx^2}}{4\sqrt{c}}\right)\sqrt{4}}{2\sqrt{c}}$	51

[In] int(2/((b^2+4\*c)/c+4\*b\*x+4\*c\*x^2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/2\*ln(1/4\*(4\*c\*x+2\*b)\*4^(1/2)/c^(1/2)+((b^2+4\*c)/c+4\*b\*x+4\*c\*x^2)^(1/2))\*4^(1/2)/c^(1/2)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 71 vs.  $2(16) = 32$ .

Time = 0.27 (sec) , antiderivative size = 137, normalized size of antiderivative = 6.23

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx$$

$$= \left[ \frac{\log\left(-4c^2x^2 - 4bcx - b^2 - (2cx + b)\sqrt{c}\sqrt{\frac{4c^2x^2 + 4bcx + b^2 + 4c}{c}} - 2c\right)}{2\sqrt{c}}, \right.$$

$$\left. - \frac{\sqrt{-c} \arctan\left(\frac{(2cx+b)\sqrt{-c}\sqrt{\frac{4c^2x^2 + 4bcx + b^2 + 4c}{c}}}{4c^2x^2 + 4bcx + b^2 + 4c}\right)}{c} \right]$$

[In] integrate(2/((b^2+4\*c)/c+4\*b\*x+4\*c\*x^2)^(1/2),x, algorithm="fricas")

[Out] [1/2\*log(-4\*c^2\*x^2 - 4\*b\*c\*x - b^2 - (2\*c\*x + b)\*sqrt(c)\*sqrt((4\*c^2\*x^2 + 4\*b\*c\*x + b^2 + 4\*c)/c) - 2\*c)/sqrt(c), -sqrt(-c)\*arctan((2\*c\*x + b)\*sqrt(-c)\*sqrt((4\*c^2\*x^2 + 4\*b\*c\*x + b^2 + 4\*c)/c)/(4\*c^2\*x^2 + 4\*b\*c\*x + b^2 + 4\*c))/c]

**Sympy [A] (verification not implemented)**

Time = 0.47 (sec) , antiderivative size = 80, normalized size of antiderivative = 3.64

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = 2 \left( \begin{array}{l} \left( \frac{\log\left(4b+4\sqrt{c}\sqrt{4bx+4cx^2+\frac{b^2+4c}{c}}+8cx\right)}{2\sqrt{c}} \right) \text{ for } c \neq 0 \\ \left( \frac{\sqrt{4bx+\frac{b^2+4c}{c}}}{2b} \right) \text{ for } b \neq 0 \\ \left( \frac{x}{\sqrt{\frac{b^2+4c}{c}}} \right) \text{ otherwise} \end{array} \right)$$

[In] integrate(2/((b\*\*2+4\*c)/c+4\*b\*x+4\*c\*x\*\*2)\*\*(1/2),x)

[Out] 2\*Piecewise((log(4\*b + 4\*sqrt(c)\*sqrt(4\*b\*x + 4\*c\*x\*\*2 + (b\*\*2 + 4\*c)/c) + 8\*c\*x)/(2\*sqrt(c)), Ne(c, 0)), (sqrt(4\*b\*x + (b\*\*2 + 4\*c)/c)/(2\*b), Ne(b, 0)), (x/sqrt((b\*\*2 + 4\*c)/c), True))

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.73

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = \frac{\operatorname{arsinh}\left(\frac{2cx+b}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[In] integrate(2/((b^2+4\*c)/c+4\*b\*x+4\*c\*x^2)^(1/2),x, algorithm="maxima")

[Out] arcsinh(1/2\*(2\*c\*x + b)/sqrt(c))/sqrt(c)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(16) = 32.

Time = 0.33 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.73

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = -\frac{\log\left(\left|-bc^2 - \left(2\sqrt{c^3}x - \sqrt{4c^3x^2 + 4bc^2x + b^2c + 4c^2}\right)\sqrt{c|c}\right|\right)}{\sqrt{c}}$$

[In] integrate(2/((b^2+4\*c)/c+4\*b\*x+4\*c\*x^2)^(1/2),x, algorithm="giac")

[Out] -log(abs(-b\*c^2 - (2\*sqrt(c^3)\*x - sqrt(4\*c^3\*x^2 + 4\*b\*c^2\*x + b^2\*c + 4\*c^2))\*sqrt(c)\*abs(c)))/sqrt(c)

**Mupad [B] (verification not implemented)**

Time = 9.36 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.82

$$\int \frac{1}{\sqrt{\frac{b^2+4c}{4c} + bx + cx^2}} dx = \frac{\ln\left(\frac{b+2cx}{\sqrt{c}} + \sqrt{\frac{b^2+4c}{c} + 4bx + 4cx^2}\right)}{\sqrt{c}}$$

[In] int(2/((4\*c + b^2)/c + 4\*b\*x + 4\*c\*x^2)^(1/2),x)

[Out] log((b + 2\*c\*x)/c^(1/2) + ((4\*c + b^2)/c + 4\*b\*x + 4\*c\*x^2)^(1/2))/c^(1/2)

$$3.126 \quad \int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx$$

Optimal result	608
Rubi [A] (verified)	608
Mathematica [B] (verified)	609
Maple [B] (verified)	609
Fricas [B] (verification not implemented)	610
Sympy [A] (verification not implemented)	610
Maxima [A] (verification not implemented)	611
Giac [B] (verification not implemented)	611
Mupad [B] (verification not implemented)	611

### Optimal result

Integrand size = 30, antiderivative size = 23

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = -\frac{\arcsin\left(\frac{b-2cx}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[Out]  $-\arcsin(1/2*(-2*c*x+b)/c^{(1/2)})/c^{(1/2)}$

### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = -\frac{\arcsin\left(\frac{b-2cx}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[In]  $\text{Int}[1/\text{Sqrt}[(-b^2 + 4*c)/(4*c) + b*x - c*x^2], x]$

[Out]  $-(\text{ArcSin}[(b - 2*c*x)/(2*\text{Sqrt}[c]])/\text{Sqrt}[c])$

#### Rule 222

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x\_Symbol] \rightarrow \text{Simp}[\text{ArcSin}[\text{Rt}[-b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[-b, 2], x] /;$   $\text{FreeQ}\{a, b\}, x \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{NegQ}[b]$

#### Rule 633

$\text{Int}[(a_) + (b_)*(x_) + (c_)*(x_)^2]^{(p_)}, x\_Symbol] \rightarrow \text{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), \text{Subst}[\text{Int}[\text{Simp}[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b]$



+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{4c}}} dx, x, b-2cx\right)}{2c} \\ &= -\frac{\sin^{-1}\left(\frac{b-2cx}{2\sqrt{c}}\right)}{\sqrt{c}} \end{aligned}$$

**Mathematica [B] (verified)**

Leaf count is larger than twice the leaf count of optimal. 123 vs. 2(23) = 46.

Time = 0.34 (sec) , antiderivative size = 123, normalized size of antiderivative = 5.35

$$\begin{aligned} &\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx \\ &= 2 \left( -\frac{\arctan\left(\frac{2\sqrt{-c^2}x - \sqrt{c}\sqrt{4 - \frac{b^2}{c} + 4bx - 4cx^2}}{b}\right)}{2\sqrt{c}} \right. \\ &\quad \left. - \frac{\log\left(2c^2x^2 + c\left(-1 - bx + \sqrt{-c}\sqrt{4 - \frac{b^2}{c} + 4bx - 4cx^2}\right)\right)}{4\sqrt{-c}} \right) \end{aligned}$$

[In] Integrate[1/Sqrt[(-b^2 + 4\*c)/(4\*c) + b\*x - c\*x^2], x]

[Out] 2\*(-1/2\*ArcTan[(2\*sqrt[-c^2]\*x - sqrt[c]\*sqrt[4 - b^2/c + 4\*b\*x - 4\*c\*x^2])/b]/sqrt[c] - Log[2\*c^2\*x^2 + c\*(-1 - b\*x + sqrt[-c]\*x\*sqrt[4 - b^2/c + 4\*b\*x - 4\*c\*x^2])]/(4\*sqrt[-c])]

**Maple [B] (verified)**

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(17) = 34.

Time = 2.49 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.91

method	result	size
default	$\frac{\arctan\left(\frac{2\sqrt{c}\left(x-\frac{b}{2c}\right)}{\sqrt{-4cx^2+4bx-\frac{b^2-4c}{c}}}\right)}{\sqrt{c}}$	44

[In] `int(2/((-b^2+4*c)/c+4*b*x-4*c*x^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/c^(1/2)*arctan(2*c^(1/2)*(x-1/2/c*b)/(-4*c*x^2+4*b*x-(b^2-4*c)/c)^(1/2))`

### Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(19) = 38.

Time = 0.28 (sec) , antiderivative size = 141, normalized size of antiderivative = 6.13

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx$$

$$= \left[ \frac{\sqrt{-c} \log\left(4c^2x^2 - 4bcx + b^2 - (2cx - b)\sqrt{-c}\sqrt{\frac{-4c^2x^2 - 4bcx + b^2 - 4c}{c}} - 2c\right)}{2c}, \right.$$

$$\left. - \frac{\arctan\left(\frac{(2cx-b)\sqrt{c}\sqrt{\frac{-4c^2x^2 - 4bcx + b^2 - 4c}{c}}}{4c^2x^2 - 4bcx + b^2 - 4c}\right)}{\sqrt{c}} \right]$$

[In] `integrate(2/((-b^2+4*c)/c+4*b*x-4*c*x^2)^(1/2),x, algorithm="fricas")`

[Out] `[-1/2*sqrt(-c)*log(4*c^2*x^2 - 4*b*c*x + b^2 - (2*c*x - b)*sqrt(-c)*sqrt(-(4*c^2*x^2 - 4*b*c*x + b^2 - 4*c)/c) - 2*c)/c, -arctan((2*c*x - b)*sqrt(c)*sqrt(-(4*c^2*x^2 - 4*b*c*x + b^2 - 4*c)/c)/(4*c^2*x^2 - 4*b*c*x + b^2 - 4*c))/sqrt(c)]`

### Sympy [A] (verification not implemented)

Time = 0.53 (sec) , antiderivative size = 83, normalized size of antiderivative = 3.61

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = 2 \left( \begin{array}{ll} \frac{\log\left(4b-8cx+4\sqrt{-c}\sqrt{4bx-4cx^2+\frac{-b^2+4c}{c}}\right)}{2\sqrt{-c}} & \text{for } c \neq 0 \\ \frac{\sqrt{4bx+\frac{-b^2+4c}{c}}}{2b} & \text{for } b \neq 0 \\ \frac{x}{\sqrt{\frac{-b^2+4c}{c}}} & \text{otherwise} \end{array} \right)$$

[In] integrate(2/((-b\*\*2+4\*c)/c+4\*b\*x-4\*c\*x\*\*2)\*\*(1/2),x)

[Out] 2\*Piecewise((log(4\*b - 8\*c\*x + 4\*sqrt(-c)\*sqrt(4\*b\*x - 4\*c\*x\*\*2 + (-b\*\*2 + 4\*c)/c))/(2\*sqrt(-c)), Ne(c, 0)), (sqrt(4\*b\*x + (-b\*\*2 + 4\*c)/c)/(2\*b), Ne(b, 0)), (x/sqrt((-b\*\*2 + 4\*c)/c), True))

### Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = -\frac{\arcsin\left(-\frac{2cx-b}{2\sqrt{c}}\right)}{\sqrt{c}}$$

[In] integrate(2/((-b^2+4\*c)/c+4\*b\*x-4\*c\*x^2)^(1/2),x, algorithm="maxima")

[Out] -arcsin(-1/2\*(2\*c\*x - b)/sqrt(c))/sqrt(c)

### Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(19) = 38.

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 2.74

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = -\frac{\log(b\sqrt{-c} - (2\sqrt{-c^3}x - \sqrt{-4c^3x^2 + 4bc^2x - b^2c + 4c^2})|c|)}{\sqrt{-c}}$$

[In] integrate(2/((-b^2+4\*c)/c+4\*b\*x-4\*c\*x^2)^(1/2),x, algorithm="giac")

[Out] -log(b\*sqrt(-c)\*c - (2\*sqrt(-c^3)\*x - sqrt(-4\*c^3\*x^2 + 4\*b\*c^2\*x - b^2\*c + 4\*c^2))\*abs(c))/sqrt(-c)

### Mupad [B] (verification not implemented)

Time = 9.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{\frac{-b^2+4c}{4c} + bx - cx^2}} dx = \frac{\ln\left(\frac{b-2cx}{\sqrt{-c}} + \sqrt{4bx + \frac{4c-b^2}{c} - 4cx^2}\right)}{\sqrt{-c}}$$

[In] int(2/(4\*b\*x + (4\*c - b^2)/c - 4\*c\*x^2)^(1/2),x)

[Out] log((b - 2\*c\*x)/(-c)^(1/2) + (4\*b\*x + (4\*c - b^2)/c - 4\*c\*x^2)^(1/2))/(-c)^(1/2)

$$3.127 \quad \int \frac{1}{\sqrt{\frac{-b^2+c}{4c}+bx-cx^2}} dx$$

Optimal result	612
Rubi [A] (verified)	612
Mathematica [B] (verified)	613
Maple [B] (verified)	613
Fricas [B] (verification not implemented)	614
Sympy [A] (verification not implemented)	614
Maxima [A] (verification not implemented)	615
Giac [B] (verification not implemented)	615
Mupad [B] (verification not implemented)	615

### Optimal result

Integrand size = 28, antiderivative size = 20

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c}+bx-cx^2}} dx = -\frac{\arcsin\left(\frac{b-2cx}{\sqrt{c}}\right)}{\sqrt{c}}$$

[Out] -arcsin((-2\*c\*x+b)/c^(1/2))/c^(1/2)

### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {633, 222}

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c}+bx-cx^2}} dx = -\frac{\arcsin\left(\frac{b-2cx}{\sqrt{c}}\right)}{\sqrt{c}}$$

[In] Int[1/Sqrt[(-b^2 + c)/(4\*c) + b\*x - c\*x^2],x]

[Out] -(ArcSin[(b - 2\*c\*x)/Sqrt[c]]/Sqrt[c])

#### Rule 222

Int[1/Sqrt[(a\_) + (b\_)\*(x\_)^2], x\_Symbol] := Simp[ArcSin[Rt[-b, 2]\*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

#### Rule 633

Int[((a\_) + (b\_)\*(x\_) + (c\_)\*(x\_)^2)^(p\_), x\_Symbol] := Dist[1/(2\*c\*(-4\*(c/(b^2 - 4\*a\*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4\*a\*c), x]^p, x], x, b

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{x^2}{c}}} dx, x, b-2cx\right)}{c} \\ &= -\frac{\sin^{-1}\left(\frac{b-2cx}{\sqrt{c}}\right)}{\sqrt{c}} \end{aligned}$$

### Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 123 vs. 2(20) = 40.

Time = 0.32 (sec) , antiderivative size = 123, normalized size of antiderivative = 6.15

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx = \frac{-2\sqrt{-c} \arctan\left(\frac{\sqrt{c}(-2\sqrt{-cx} + \sqrt{1-\frac{b^2}{c} + 4bx - 4cx^2})}{b}\right) + \sqrt{c} \log\left(c\left(-1 - 4bx + 8cx^2 + 4\sqrt{-cx}\sqrt{1-\frac{b^2}{c} + 4bx - 4cx^2}\right)\right)}{2\sqrt{-c^2}}$$

[In] Integrate[1/Sqrt[(-b^2 + c)/(4\*c) + b\*x - c\*x^2], x]

[Out] -1/2\*(-2\*Sqrt[-c]\*ArcTan[(Sqrt[c]\*(-2\*Sqrt[-c]\*x + Sqrt[1 - b^2/c + 4\*b\*x - 4\*c\*x^2]))/b] + Sqrt[c]\*Log[c\*(-1 - 4\*b\*x + 8\*c\*x^2 + 4\*Sqrt[-c]\*x\*Sqrt[1 - b^2/c + 4\*b\*x - 4\*c\*x^2])])/Sqrt[-c^2]

### Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(16) = 32.

Time = 2.50 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.20

method	result	size
default	$\frac{\arctan\left(\frac{2\sqrt{c}\left(x-\frac{b}{2c}\right)}{\sqrt{-4cx^2+4bx-\frac{b^2-c}{c}}}\right)}{\sqrt{c}}$	44

[In] int(2/((-b^2+c)/c+4\*b\*x-4\*c\*x^2)^(1/2), x, method=\_RETURNVERBOSE)

[Out] 1/c^(1/2)\*arctan(2\*c^(1/2)\*(x-1/2/c\*b)/(-4\*c\*x^2+4\*b\*x-(b^2-c)/c)^(1/2))

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(19) = 38.

Time = 0.27 (sec) , antiderivative size = 143, normalized size of antiderivative = 7.15

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx$$

$$= \left[ \frac{\sqrt{-c} \log \left( 8c^2x^2 - 8bcx + 2b^2 - 2(2cx - b)\sqrt{-c} \sqrt{\frac{-4c^2x^2 - 4bcx + b^2 - c}{c}} - c \right)}{2c}, \right.$$

$$\left. - \frac{\arctan \left( \frac{(2cx - b)\sqrt{c} \sqrt{\frac{-4c^2x^2 - 4bcx + b^2 - c}{c}}}{4c^2x^2 - 4bcx + b^2 - c} \right)}{\sqrt{c}} \right]$$

[In] integrate(2/((-b^2+c)/c+4\*b\*x-4\*c\*x^2)^(1/2),x, algorithm="fricas")

[Out] [-1/2\*sqrt(-c)\*log(8\*c^2\*x^2 - 8\*b\*c\*x + 2\*b^2 - 2\*(2\*c\*x - b)\*sqrt(-c)\*sqrt(-4\*c^2\*x^2 - 4\*b\*c\*x + b^2 - c)/c) - c, -arctan((2\*c\*x - b)\*sqrt(c)\*sqrt(-4\*c^2\*x^2 - 4\*b\*c\*x + b^2 - c)/(4\*c^2\*x^2 - 4\*b\*c\*x + b^2 - c))/sqrt(c)]

**Sympy [A] (verification not implemented)**

Time = 0.51 (sec) , antiderivative size = 78, normalized size of antiderivative = 3.90

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx = 2 \left( \begin{array}{l} \frac{\log \left( 4b - 8cx + 4\sqrt{-c} \sqrt{4bx - 4cx^2 + \frac{-b^2+c}{c}} \right)}{2\sqrt{-c}} \quad \text{for } c \neq 0 \\ \frac{\sqrt{4bx + \frac{-b^2+c}{c}}}{2b} \quad \text{for } b \neq 0 \\ \frac{x}{\sqrt{\frac{-b^2+c}{c}}} \quad \text{otherwise} \end{array} \right)$$

[In] integrate(2/((-b\*\*2+c)/c+4\*b\*x-4\*c\*x\*\*2)\*\*(1/2),x)

[Out] 2\*Piecewise((log(4\*b - 8\*c\*x + 4\*sqrt(-c)\*sqrt(4\*b\*x - 4\*c\*x\*\*2 + (-b\*\*2 + c)/c))/(2\*sqrt(-c)), Ne(c, 0)), (sqrt(4\*b\*x + (-b\*\*2 + c)/c)/(2\*b), Ne(b, 0)), (x/sqrt((-b\*\*2 + c)/c), True))

**Maxima [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx = -\frac{\arcsin\left(-\frac{2cx-b}{\sqrt{c}}\right)}{\sqrt{c}}$$

[In] integrate(2/((-b^2+c)/c+4\*b\*x-4\*c\*x^2)^(1/2),x, algorithm="maxima")

[Out] -arcsin(-(2\*c\*x - b)/sqrt(c))/sqrt(c)

**Giac [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(19) = 38.

Time = 0.28 (sec) , antiderivative size = 61, normalized size of antiderivative = 3.05

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx = -\frac{\log(b\sqrt{-c} - (2\sqrt{-c^3x} - \sqrt{-4c^3x^2 + 4bc^2x - b^2c + c^2})|c|)}{\sqrt{-c}}$$

[In] integrate(2/((-b^2+c)/c+4\*b\*x-4\*c\*x^2)^(1/2),x, algorithm="giac")

[Out] -log(b\*sqrt(-c)\*c - (2\*sqrt(-c^3)\*x - sqrt(-4\*c^3\*x^2 + 4\*b\*c^2\*x - b^2\*c + c^2))\*abs(c))/sqrt(-c)

**Mupad [B] (verification not implemented)**

Time = 9.26 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.20

$$\int \frac{1}{\sqrt{\frac{-b^2+c}{4c} + bx - cx^2}} dx = \frac{\ln\left(\frac{b-2cx}{\sqrt{-c}} + \sqrt{\frac{c-b^2}{c} + 4bx - 4cx^2}\right)}{\sqrt{-c}}$$

[In] int(2/((c - b^2)/c + 4\*b\*x - 4\*c\*x^2)^(1/2),x)

[Out] log((b - 2\*c\*x)/(-c)^(1/2) + ((c - b^2)/c + 4\*b\*x - 4\*c\*x^2)^(1/2))/(-c)^(1/2)

$$3.128 \quad \int \frac{1}{(2+3x+x^2)^{3/2}} dx$$

Optimal result	616
Rubi [A] (verified)	616
Mathematica [A] (verified)	617
Maple [A] (verified)	617
Fricas [B] (verification not implemented)	617
Sympy [F]	618
Maxima [A] (verification not implemented)	618
Giac [A] (verification not implemented)	618
Mupad [B] (verification not implemented)	618

### Optimal result

Integrand size = 12, antiderivative size = 19

$$\int \frac{1}{(2+3x+x^2)^{3/2}} dx = -\frac{2(3+2x)}{\sqrt{2+3x+x^2}}$$

[Out]  $-2*(3+2*x)/(x^2+3*x+2)^{(1/2)}$

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$ , Rules used = {627}

$$\int \frac{1}{(2+3x+x^2)^{3/2}} dx = -\frac{2(2x+3)}{\sqrt{x^2+3x+2}}$$

[In]  $\text{Int}[(2+3*x+x^2)^{-3/2}, x]$

[Out]  $(-2*(3+2*x))/\text{Sqrt}[2+3*x+x^2]$

#### Rule 627

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-3/2), x_Symbol] := Simp[-2*((b + 2*c*x)/((b^2 - 4*a*c)*Sqrt[a + b*x + c*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

#### Rubi steps

$$\text{integral} = -\frac{2(3+2x)}{\sqrt{2+3x+x^2}}$$



**Mathematica [A] (verified)**

Time = 0.13 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = -\frac{2(3 + 2x)}{\sqrt{2 + 3x + x^2}}$$

[In] Integrate[(2 + 3\*x + x^2)^(-3/2),x]

[Out] (-2\*(3 + 2\*x))/Sqrt[2 + 3\*x + x^2]

**Maple [A] (verified)**

Time = 2.35 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.95

method	result	size
default	$-\frac{2(2x+3)}{\sqrt{x^2+3x+2}}$	18
trager	$-\frac{2(2x+3)}{\sqrt{x^2+3x+2}}$	18
risch	$-\frac{2(2x+3)}{\sqrt{x^2+3x+2}}$	18
gospers	$-\frac{2(2+x)(1+x)(2x+3)}{(x^2+3x+2)^{\frac{3}{2}}}$	24

[In] int(1/(x^2+3\*x+2)^(3/2),x,method=\_RETURNVERBOSE)

[Out] -2\*(2\*x+3)/(x^2+3\*x+2)^(1/2)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(17) = 34.

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.00

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = -\frac{2(2x^2 + \sqrt{x^2 + 3x + 2}(2x + 3) + 6x + 4)}{x^2 + 3x + 2}$$

[In] integrate(1/(x^2+3\*x+2)^(3/2),x, algorithm="fricas")

[Out] -2\*(2\*x^2 + sqrt(x^2 + 3\*x + 2)\*(2\*x + 3) + 6\*x + 4)/(x^2 + 3\*x + 2)

**Sympy [F]**

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = \int \frac{1}{(x^2 + 3x + 2)^{\frac{3}{2}}} dx$$

[In] integrate(1/(x\*\*2+3\*x+2)\*\*(3/2),x)

[Out] Integral((x\*\*2 + 3\*x + 2)\*\*(-3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.37

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = -\frac{4x}{\sqrt{x^2 + 3x + 2}} - \frac{6}{\sqrt{x^2 + 3x + 2}}$$

[In] integrate(1/(x^2+3\*x+2)^(3/2),x, algorithm="maxima")

[Out] -4\*x/sqrt(x^2 + 3\*x + 2) - 6/sqrt(x^2 + 3\*x + 2)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = -\frac{2(2x + 3)}{\sqrt{x^2 + 3x + 2}}$$

[In] integrate(1/(x^2+3\*x+2)^(3/2),x, algorithm="giac")

[Out] -2\*(2\*x + 3)/sqrt(x^2 + 3\*x + 2)

**Mupad [B] (verification not implemented)**

Time = 8.97 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.79

$$\int \frac{1}{(2 + 3x + x^2)^{3/2}} dx = -\frac{4\left(x + \frac{3}{2}\right)}{\sqrt{x^2 + 3x + 2}}$$

[In] int(1/(3\*x + x^2 + 2)^(3/2),x)

[Out] -(4\*(x + 3/2))/(3\*x + x^2 + 2)^(1/2)

$$3.129 \quad \int \frac{1}{(27-24x+4x^2)^{3/2}} dx$$

Optimal result	619
Rubi [A] (verified)	619
Mathematica [A] (verified)	620
Maple [A] (verified)	620
Fricas [B] (verification not implemented)	620
Sympy [F]	621
Maxima [A] (verification not implemented)	621
Giac [A] (verification not implemented)	621
Mupad [B] (verification not implemented)	621

### Optimal result

Integrand size = 14, antiderivative size = 23

$$\int \frac{1}{(27-24x+4x^2)^{3/2}} dx = \frac{3-x}{9\sqrt{27-24x+4x^2}}$$

[Out] 1/9\*(3-x)/(4\*x^2-24\*x+27)^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.071$ , Rules used = {627}

$$\int \frac{1}{(27-24x+4x^2)^{3/2}} dx = \frac{3-x}{9\sqrt{4x^2-24x+27}}$$

[In] Int[(27 - 24\*x + 4\*x^2)^(-3/2), x]

[Out] (3 - x)/(9\*sqrt[27 - 24\*x + 4\*x^2])

#### Rule 627

Int[((a\_.) + (b\_.)\*(x\_)) + (c\_.)\*(x\_)^2]^(-3/2), x\_Symbol] :> Simp[-2\*((b + 2\*c\*x)/((b^2 - 4\*a\*c)\*sqrt[a + b\*x + c\*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

#### Rubi steps

$$\text{integral} = \frac{3-x}{9\sqrt{27-24x+4x^2}}$$

**Mathematica [A] (verified)**

Time = 0.14 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = \frac{3 - x}{9\sqrt{27 - 24x + 4x^2}}$$

[In] Integrate[(27 - 24\*x + 4\*x^2)^(-3/2),x]

[Out] (3 - x)/(9\*sqrt[27 - 24\*x + 4\*x^2])

**Maple [A] (verified)**

Time = 2.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.78

method	result	size
trager	$-\frac{-3+x}{9\sqrt{4x^2-24x+27}}$	18
risch	$-\frac{-3+x}{9\sqrt{4x^2-24x+27}}$	18
default	$-\frac{8x-24}{72\sqrt{4x^2-24x+27}}$	20
gospers	$-\frac{(-3+2x)(2x-9)(-3+x)}{9(4x^2-24x+27)^{3/2}}$	28

[In] int(1/(4\*x^2-24\*x+27)^(3/2),x,method=\_RETURNVERBOSE)

[Out] -1/9\*(-3+x)/(4\*x^2-24\*x+27)^(1/2)

**Fricas [B] (verification not implemented)**

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(17) = 34.

Time = 0.28 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.78

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = -\frac{4x^2 + 2\sqrt{4x^2 - 24x + 27}(x - 3) - 24x + 27}{18(4x^2 - 24x + 27)}$$

[In] integrate(1/(4\*x^2-24\*x+27)^(3/2),x, algorithm="fricas")

[Out] -1/18\*(4\*x^2 + 2\*sqrt(4\*x^2 - 24\*x + 27)\*(x - 3) - 24\*x + 27)/(4\*x^2 - 24\*x + 27)

**Sympy [F]**

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = \int \frac{1}{(4x^2 - 24x + 27)^{\frac{3}{2}}} dx$$

[In] integrate(1/(4\*x\*\*2-24\*x+27)\*\*(3/2),x)

[Out] Integral((4\*x\*\*2 - 24\*x + 27)\*\*(-3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = -\frac{x}{9\sqrt{4x^2 - 24x + 27}} + \frac{1}{3\sqrt{4x^2 - 24x + 27}}$$

[In] integrate(1/(4\*x^2-24\*x+27)^(3/2),x, algorithm="maxima")

[Out] -1/9\*x/sqrt(4\*x^2 - 24\*x + 27) + 1/3/sqrt(4\*x^2 - 24\*x + 27)

**Giac [A] (verification not implemented)**

none

Time = 0.28 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = -\frac{x - 3}{9\sqrt{4x^2 - 24x + 27}}$$

[In] integrate(1/(4\*x^2-24\*x+27)^(3/2),x, algorithm="giac")

[Out] -1/9\*(x - 3)/sqrt(4\*x^2 - 24\*x + 27)

**Mupad [B] (verification not implemented)**

Time = 0.06 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.74

$$\int \frac{1}{(27 - 24x + 4x^2)^{3/2}} dx = -\frac{x - 3}{9\sqrt{4x^2 - 24x + 27}}$$

[In] int(1/(4\*x^2 - 24\*x + 27)^(3/2),x)

[Out] -(x - 3)/(9\*(4\*x^2 - 24\*x + 27)^(1/2))

### 3.130 $\int \frac{x}{(5-4x-x^2)^{3/2}} dx$

Optimal result	622
Rubi [A] (verified)	622
Mathematica [A] (verified)	623
Maple [A] (verified)	623
Fricas [A] (verification not implemented)	623
Sympy [F]	624
Maxima [A] (verification not implemented)	624
Giac [A] (verification not implemented)	624
Mupad [B] (verification not implemented)	624

#### Optimal result

Integrand size = 16, antiderivative size = 23

$$\int \frac{x}{(5-4x-x^2)^{3/2}} dx = \frac{5-2x}{9\sqrt{5-4x-x^2}}$$

[Out] 1/9\*(5-2\*x)/(-x^2-4\*x+5)^(1/2)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.062$ , Rules used = {650}

$$\int \frac{x}{(5-4x-x^2)^{3/2}} dx = \frac{5-2x}{9\sqrt{-x^2-4x+5}}$$

[In] Int[x/(5 - 4\*x - x^2)^(3/2), x]

[Out] (5 - 2\*x)/(9\*sqrt[5 - 4\*x - x^2])

#### Rule 650

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(3/2), x_Symbol]
:> Simp[-2*((b*d - 2*a*e + (2*c*d - b*e)*x)/((b^2 - 4*a*c)*sqrt[a + b*x
+ c*x^2])), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b
^2 - 4*a*c, 0]
```

#### Rubi steps

$$\text{integral} = \frac{5-2x}{9\sqrt{5-4x-x^2}}$$

**Mathematica [A] (verified)**

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.43

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = \frac{(-5 + 2x)\sqrt{5 - 4x - x^2}}{9(-1 + x)(5 + x)}$$

[In] Integrate[x/(5 - 4\*x - x^2)^(3/2),x]

[Out] ((-5 + 2\*x)\*Sqrt[5 - 4\*x - x^2])/(9\*(-1 + x)\*(5 + x))

**Maple [A] (verified)**

Time = 2.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.87

method	result	size
risch	$-\frac{-5+2x}{9\sqrt{-x^2-4x+5}}$	20
gosper	$\frac{(x+5)(-1+x)(-5+2x)}{9(-x^2-4x+5)^{3/2}}$	26
trager	$\frac{(-5+2x)\sqrt{-x^2-4x+5}}{9x^2+36x-45}$	30
default	$\frac{1}{\sqrt{-x^2-4x+5}} + \frac{-2x-4}{9\sqrt{-x^2-4x+5}}$	33

[In] int(x/(-x^2-4\*x+5)^(3/2),x,method=\_RETURNVERBOSE)

[Out] -1/9\*(-5+2\*x)/(-x^2-4\*x+5)^(1/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = \frac{\sqrt{-x^2 - 4x + 5}(2x - 5)}{9(x^2 + 4x - 5)}$$

[In] integrate(x/(-x^2-4\*x+5)^(3/2),x, algorithm="fricas")

[Out] 1/9\*sqrt(-x^2 - 4\*x + 5)\*(2\*x - 5)/(x^2 + 4\*x - 5)

**Sympy [F]**

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = \int \frac{x}{(-(x - 1)(x + 5))^{\frac{3}{2}}} dx$$

[In] integrate(x/(-x\*\*2-4\*x+5)\*\*(3/2),x)

[Out] Integral(x/(-(x - 1)\*(x + 5))\*\*(3/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.30

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = -\frac{2x}{9\sqrt{-x^2 - 4x + 5}} + \frac{5}{9\sqrt{-x^2 - 4x + 5}}$$

[In] integrate(x/(-x^2-4\*x+5)^(3/2),x, algorithm="maxima")

[Out] -2/9\*x/sqrt(-x^2 - 4\*x + 5) + 5/9/sqrt(-x^2 - 4\*x + 5)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.26

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = \frac{\sqrt{-x^2 - 4x + 5}(2x - 5)}{9(x^2 + 4x - 5)}$$

[In] integrate(x/(-x^2-4\*x+5)^(3/2),x, algorithm="giac")

[Out] 1/9\*sqrt(-x^2 - 4\*x + 5)\*(2\*x - 5)/(x^2 + 4\*x - 5)

**Mupad [B] (verification not implemented)**

Time = 0.05 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.83

$$\int \frac{x}{(5 - 4x - x^2)^{3/2}} dx = -\frac{2x - 5}{9\sqrt{-x^2 - 4x + 5}}$$

[In] int(x/(5 - x^2 - 4\*x)^(3/2),x)

[Out] -(2\*x - 5)/(9\*(5 - x^2 - 4\*x)^(1/2))



$$3.131 \quad \int \frac{1}{(5-4x-x^2)^{5/2}} dx$$

Optimal result	625
Rubi [A] (verified)	625
Mathematica [A] (verified)	626
Maple [A] (verified)	626
Fricas [A] (verification not implemented)	627
Sympy [F]	627
Maxima [A] (verification not implemented)	627
Giac [A] (verification not implemented)	628
Mupad [B] (verification not implemented)	628

### Optimal result

Integrand size = 14, antiderivative size = 43

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{2+x}{27(5-4x-x^2)^{3/2}} + \frac{2(2+x)}{243\sqrt{5-4x-x^2}}$$

[Out] 1/27\*(2+x)/(-x^2-4\*x+5)^(3/2)+2/243\*(2+x)/(-x^2-4\*x+5)^(1/2)

### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {628, 627}

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{2(x+2)}{243\sqrt{-x^2-4x+5}} + \frac{x+2}{27(-x^2-4x+5)^{3/2}}$$

[In] Int[(5 - 4\*x - x^2)^(-5/2), x]

[Out] (2 + x)/(27\*(5 - 4\*x - x^2)^(3/2)) + (2\*(2 + x))/(243\*sqrt[5 - 4\*x - x^2])

#### Rule 627

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(-3/2), x\_Symbol] := Simp[-2\*((b + 2\*c\*x)/((b^2 - 4\*a\*c)\*Sqrt[a + b\*x + c\*x^2])), x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4\*a\*c, 0]

#### Rule 628

Int[((a\_.) + (b\_.)\*(x\_) + (c\_.)\*(x\_)^2)^(p\_), x\_Symbol] := Simp[(b + 2\*c\*x)\*((a + b\*x + c\*x^2)^(p + 1)/((p + 1)\*(b^2 - 4\*a\*c))), x] - Dist[2\*c\*((2\*p + 3)/((p + 1)\*(b^2 - 4\*a\*c))), Int[(a + b\*x + c\*x^2)^(p + 1), x], x] /; Free

$Q[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4ac, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{NeQ}[p, -3/2] \ \&\& \ \text{Int egerQ}[4p]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2+x}{27(5-4x-x^2)^{3/2}} + \frac{2}{27} \int \frac{1}{(5-4x-x^2)^{3/2}} dx \\ &= \frac{2+x}{27(5-4x-x^2)^{3/2}} + \frac{2(2+x)}{243\sqrt{5-4x-x^2}} \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.01 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.00

$$\int \frac{1}{(5-4x-x^2)^{5/2}} dx = \frac{\sqrt{5-4x-x^2}(38+3x-12x^2-2x^3)}{243(-1+x)^2(5+x)^2}$$

[In] Integrate[(5 - 4\*x - x^2)^(-5/2), x]

[Out] (Sqrt[5 - 4\*x - x^2]\*(38 + 3\*x - 12\*x^2 - 2\*x^3))/(243\*(-1 + x)^2\*(5 + x)^2)

**Maple [A] (verified)**

Time = 2.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

method	result	size
gospers	$\frac{(x+5)(-1+x)(2x^3+12x^2-3x-38)}{243(-x^2-4x+5)^{5/2}}$	36
default	$-\frac{-2x-4}{54(-x^2-4x+5)^{3/2}} - \frac{-2x-4}{243\sqrt{-x^2-4x+5}}$	40
trager	$-\frac{(2x^3+12x^2-3x-38)\sqrt{-x^2-4x+5}}{243(x^2+4x-5)^2}$	40
risch	$\frac{2x^3+12x^2-3x-38}{243(x^2+4x-5)\sqrt{-x^2-4x+5}}$	40

[In] int(1/(-x^2-4\*x+5)^(5/2), x, method=\_RETURNVERBOSE)

[Out] 1/243\*(x+5)\*(-1+x)\*(2\*x^3+12\*x^2-3\*x-38)/(-x^2-4\*x+5)^(5/2)

**Fricas [A] (verification not implemented)**

none

Time = 0.26 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.14

$$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx = -\frac{(2x^3 + 12x^2 - 3x - 38)\sqrt{-x^2 - 4x + 5}}{243(x^4 + 8x^3 + 6x^2 - 40x + 25)}$$

[In] integrate(1/(-x^2-4\*x+5)^(5/2),x, algorithm="fricas")

[Out] -1/243\*(2\*x^3 + 12\*x^2 - 3\*x - 38)\*sqrt(-x^2 - 4\*x + 5)/(x^4 + 8\*x^3 + 6\*x^2 - 40\*x + 25)

**Sympy [F]**

$$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx = \int \frac{1}{(-x^2 - 4x + 5)^{5/2}} dx$$

[In] integrate(1/(-x\*\*2-4\*x+5)\*\*(5/2),x)

[Out] Integral((-x\*\*2 - 4\*x + 5)\*\*(-5/2), x)

**Maxima [A] (verification not implemented)**

none

Time = 0.20 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.37

$$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx = \frac{2x}{243\sqrt{-x^2 - 4x + 5}} + \frac{4}{243\sqrt{-x^2 - 4x + 5}} + \frac{x}{27(-x^2 - 4x + 5)^{3/2}} + \frac{2}{27(-x^2 - 4x + 5)^{3/2}}$$

[In] integrate(1/(-x^2-4\*x+5)^(5/2),x, algorithm="maxima")

[Out] 2/243\*x/sqrt(-x^2 - 4\*x + 5) + 4/243/sqrt(-x^2 - 4\*x + 5) + 1/27\*x/(-x^2 - 4\*x + 5)^(3/2) + 2/27/(-x^2 - 4\*x + 5)^(3/2)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.84

$$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx = -\frac{((2(x+6)x - 3)x - 38)\sqrt{-x^2 - 4x + 5}}{243(x^2 + 4x - 5)^2}$$

[In] integrate(1/(-x^2-4\*x+5)^(5/2),x, algorithm="giac")

[Out] -1/243\*((2\*(x + 6)\*x - 3)\*x - 38)\*sqrt(-x^2 - 4\*x + 5)/(x^2 + 4\*x - 5)^2

**Mupad [B] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.67

$$\int \frac{1}{(5 - 4x - x^2)^{5/2}} dx = -\frac{(4x + 8)(8x^2 + 32x - 76)}{3888(-x^2 - 4x + 5)^{3/2}}$$

[In] int(1/(5 - x^2 - 4\*x)^(5/2),x)

[Out] -((4\*x + 8)\*(32\*x + 8\*x^2 - 76))/(3888\*(5 - x^2 - 4\*x)^(3/2))

### 3.132 $\int (a + bx + cx^2)^p dx$

Optimal result	629
Rubi [A] (verified)	629
Mathematica [A] (verified)	630
Maple [F]	630
Fricas [F]	631
Sympy [F]	631
Maxima [F]	631
Giac [F]	631
Mupad [F(-1)]	632

#### Optimal result

Integrand size = 12, antiderivative size = 122

$$\int (a + bx + cx^2)^p dx = \frac{2^{1+p} \left( -\frac{b - \sqrt{b^2 - 4ac} + 2cx}{\sqrt{b^2 - 4ac}} \right)^{-1-p} (a + bx + cx^2)^{1+p} \text{Hypergeometric2F1} \left( -p, 1 + p, 2 + p, \frac{b + \sqrt{b^2 - 4ac} + 2cx}{2\sqrt{b^2 - 4ac}} \right)}{\sqrt{b^2 - 4ac}(1 + p)}$$

```
[Out] -2^(p+1)*(c*x^2+b*x+a)^(p+1)*hypergeom([-p, p+1], [2+p], 1/2*(b+2*c*x+(-4*a*c+b^2)^(1/2))/(-4*a*c+b^2)^(1/2))*((-b-2*c*x+(-4*a*c+b^2)^(1/2))/(-4*a*c+b^2)^(1/2))^(1-p)/(p+1)/(-4*a*c+b^2)^(1/2)
```

#### Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$ , Rules used = {638}

$$\int (a + bx + cx^2)^p dx = \frac{2^{p+1} \left( -\frac{\sqrt{b^2 - 4ac} + b + 2cx}{\sqrt{b^2 - 4ac}} \right)^{-p-1} (a + bx + cx^2)^{p+1} \text{Hypergeometric2F1} \left( -p, p + 1, p + 2, \frac{b + 2cx + \sqrt{b^2 - 4ac}}{2\sqrt{b^2 - 4ac}} \right)}{(p + 1)\sqrt{b^2 - 4ac}}$$

```
[In] Int[(a + b*x + c*x^2)^p, x]
```

```
[Out] -((2^(1 + p)*(-(b - Sqrt[b^2 - 4*a*c] + 2*c*x)/Sqrt[b^2 - 4*a*c]))^(-1 - p)*(a + b*x + c*x^2)^(1 + p)*Hypergeometric2F1[-p, 1 + p, 2 + p, (b + Sqrt[b^2 - 4*a*c] + 2*c*x)/(2*Sqrt[b^2 - 4*a*c])])/(Sqrt[b^2 - 4*a*c]*(1 + p))
```

## Rule 638

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Rt[b^2
- 4*a*c, 2]}, Simp[(-(a + b*x + c*x^2)^(p + 1)/(q*(p + 1)*((q - b - 2*c*x)
/(2*q))^(p + 1)))*Hypergeometric2F1[-p, p + 1, p + 2, (b + q + 2*c*x)/(2*q)
], x]] /; FreeQ[{a, b, c, p}, x] && NeQ[b^2 - 4*a*c, 0] && !IntegerQ[4*p]
```

## Rubi steps

$$\text{integral} \\ = - \frac{2^{1+p} \left( -\frac{b - \sqrt{b^2 - 4ac} + 2cx}{\sqrt{b^2 - 4ac}} \right)^{-1-p} (a + bx + cx^2)^{1+p} {}_2F_1 \left( -p, 1 + p; 2 + p; \frac{b + \sqrt{b^2 - 4ac} + 2cx}{2\sqrt{b^2 - 4ac}} \right)}{\sqrt{b^2 - 4ac}(1 + p)}$$

## Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.03

$$\int (a + bx + cx^2)^p dx \\ = \frac{2^{-1+p} (b - \sqrt{b^2 - 4ac} + 2cx) \left( \frac{b + \sqrt{b^2 - 4ac} + 2cx}{\sqrt{b^2 - 4ac}} \right)^{-p} (a + x(b + cx))^p \text{Hypergeometric2F1} \left( -p, 1 + p, 2 + p, \frac{-b + \sqrt{b^2 - 4ac} - 2cx}{2\sqrt{b^2 - 4ac}} \right)}{c(1 + p)}$$

[In] Integrate[(a + b\*x + c\*x^2)^p,x]

[Out] (2^(-1 + p)\*(b - Sqrt[b^2 - 4\*a\*c] + 2\*c\*x)\*(a + x\*(b + c\*x))^p\*Hypergeometric2F1[-p, 1 + p, 2 + p, (-b + Sqrt[b^2 - 4\*a\*c] - 2\*c\*x)/(2\*Sqrt[b^2 - 4\*a\*c])])/(c\*(1 + p)\*((b + Sqrt[b^2 - 4\*a\*c] + 2\*c\*x)/Sqrt[b^2 - 4\*a\*c])^p)

## Maple [F]

$$\int (cx^2 + bx + a)^p dx$$

[In] int((c\*x^2+b\*x+a)^p,x)

[Out] int((c\*x^2+b\*x+a)^p,x)

**Fricas [F]**

$$\int (a + bx + cx^2)^p dx = \int (cx^2 + bx + a)^p dx$$

[In] integrate((c\*x^2+b\*x+a)^p,x, algorithm="fricas")

[Out] integral((c\*x^2 + b\*x + a)^p, x)

**Sympy [F]**

$$\int (a + bx + cx^2)^p dx = \int (a + bx + cx^2)^p dx$$

[In] integrate((c\*x\*\*2+b\*x+a)\*\*p,x)

[Out] Integral((a + b\*x + c\*x\*\*2)\*\*p, x)

**Maxima [F]**

$$\int (a + bx + cx^2)^p dx = \int (cx^2 + bx + a)^p dx$$

[In] integrate((c\*x^2+b\*x+a)^p,x, algorithm="maxima")

[Out] integrate((c\*x^2 + b\*x + a)^p, x)

**Giac [F]**

$$\int (a + bx + cx^2)^p dx = \int (cx^2 + bx + a)^p dx$$

[In] integrate((c\*x^2+b\*x+a)^p,x, algorithm="giac")

[Out] integrate((c\*x^2 + b\*x + a)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (a + bx + cx^2)^p dx = \int (cx^2 + bx + a)^p dx$$

```
[In] int((a + b*x + c*x^2)^p,x)
```

```
[Out] int((a + b*x + c*x^2)^p, x)
```



### 3.133 $\int (3 + 4x + 5x^2)^p dx$

Optimal result	633
Rubi [A] (verified)	633
Mathematica [A] (verified)	634
Maple [F]	634
Fricas [F]	634
Sympy [F]	635
Maxima [F]	635
Giac [F]	635
Mupad [F(-1)]	635

#### Optimal result

Integrand size = 12, antiderivative size = 37

$$\int (3 + 4x + 5x^2)^p dx = 5^{-1-p} 11^p (2 + 5x) \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{11} (2 + 5x)^2 \right)$$

[Out]  $5^{(-1-p)} * 11^p * (2+5*x) * \operatorname{hypergeom}([1/2, -p], [3/2], -1/11*(2+5*x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x + 5x^2)^p dx = 5^{-p-1} 11^p (5x + 2) \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{11} (5x + 2)^2 \right)$$

[In]  $\operatorname{Int}[(3 + 4*x + 5*x^2)^p, x]$

[Out]  $5^{(-1 - p)} * 11^p * (2 + 5*x) * \operatorname{Hypergeometric2F1}[1/2, -p, 3/2, -1/11*(2 + 5*x)^2]$

#### Rule 251

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a^p * x * \operatorname{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /;$   $\operatorname{FreeQ}\{a, b, n, p, x\} \ \&\& \ !\operatorname{IGtQ}[p, 0] \ \&\& \ !\operatorname{IntegerQ}[1/n] \ \&\& \ !\operatorname{ILtQ}[\operatorname{Simplify}[1/n + p], 0] \ \&\& \ (\operatorname{IntegerQ}[p] \ || \ \operatorname{GtQ}[a, 0])$

#### Rule 633

$\operatorname{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^{(p)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{Simp}[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b$

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}(5^{-1-p}11^p) \text{Subst}\left(\int \left(1 + \frac{x^2}{44}\right)^p dx, x, 4 + 10x\right) \\ &= 5^{-1-p}11^p(2 + 5x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{1}{11}(2 + 5x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int (3 + 4x + 5x^2)^p dx = 5^{-1-p}11^p(2 + 5x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{11}(2 + 5x)^2\right)$$

[In] Integrate[(3 + 4\*x + 5\*x^2)^p, x]

[Out] 5^(-1 - p)\*11^p\*(2 + 5\*x)\*Hypergeometric2F1[1/2, -p, 3/2, -1/11\*(2 + 5\*x)^2]

**Maple [F]**

$$\int (5x^2 + 4x + 3)^p dx$$

[In] int((5\*x^2+4\*x+3)^p, x)

[Out] int((5\*x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3 + 4x + 5x^2)^p dx = \int (5x^2 + 4x + 3)^p dx$$

[In] integrate((5\*x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((5\*x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x + 5x^2)^p dx = \int (5x^2 + 4x + 3)^p dx$$

[In] integrate((5\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((5\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x + 5x^2)^p dx = \int (5x^2 + 4x + 3)^p dx$$

[In] integrate((5\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((5\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x + 5x^2)^p dx = \int (5x^2 + 4x + 3)^p dx$$

[In] integrate((5\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((5\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x + 5x^2)^p dx = \int (5x^2 + 4x + 3)^p dx$$

[In] int((4\*x + 5\*x^2 + 3)^p,x)

[Out] int((4\*x + 5\*x^2 + 3)^p, x)

### 3.134 $\int (3 + 4x + 4x^2)^p dx$

Optimal result	636
Rubi [A] (verified)	636
Mathematica [A] (verified)	637
Maple [F]	637
Fricas [F]	637
Sympy [F]	638
Maxima [F]	638
Giac [F]	638
Mupad [F(-1)]	638

#### Optimal result

Integrand size = 12, antiderivative size = 32

$$\int (3 + 4x + 4x^2)^p dx = 2^{-1+p}(1 + 2x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{2}(1 + 2x)^2\right)$$

[Out]  $2^{(-1+p)}*(1+2*x)*\operatorname{hypergeom}([1/2, -p], [3/2], -1/2*(1+2*x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x + 4x^2)^p dx = 2^{p-1}(2x + 1) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{2}(2x + 1)^2\right)$$

[In]  $\operatorname{Int}[(3 + 4*x + 4*x^2)^p, x]$

[Out]  $2^{(-1 + p)}*(1 + 2*x)*\operatorname{Hypergeometric2F1}[1/2, -p, 3/2, -1/2*(1 + 2*x)^2]$

#### Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F
1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p
, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] ||
GtQ[a, 0])
```

#### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c))))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= 2^{-3+p} \text{Subst} \left( \int \left( 1 + \frac{x^2}{32} \right)^p dx, x, 4 + 8x \right) \\ &= 2^{-1+p} (1 + 2x) {}_2F_1 \left( \frac{1}{2}, -p; \frac{3}{2}; -\frac{1}{2}(1 + 2x)^2 \right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int (3 + 4x + 4x^2)^p dx = 2^{-3+p} (4 + 8x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{32}(4 + 8x)^2 \right)$$

[In] Integrate[(3 + 4\*x + 4\*x^2)^p,x]

[Out] 2^(-3 + p)\*(4 + 8\*x)\*Hypergeometric2F1[1/2, -p, 3/2, -1/32\*(4 + 8\*x)^2]

**Maple [F]**

$$\int (4x^2 + 4x + 3)^p dx$$

[In] int((4\*x^2+4\*x+3)^p,x)

[Out] int((4\*x^2+4\*x+3)^p,x)

**Fricas [F]**

$$\int (3 + 4x + 4x^2)^p dx = \int (4x^2 + 4x + 3)^p dx$$

[In] integrate((4\*x^2+4\*x+3)^p,x, algorithm="fricas")

[Out] integral((4\*x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x + 4x^2)^p dx = \int (4x^2 + 4x + 3)^p dx$$

[In] integrate((4\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((4\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x + 4x^2)^p dx = \int (4x^2 + 4x + 3)^p dx$$

[In] integrate((4\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((4\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x + 4x^2)^p dx = \int (4x^2 + 4x + 3)^p dx$$

[In] integrate((4\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((4\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x + 4x^2)^p dx = \int (4x^2 + 4x + 3)^p dx$$

[In] int((4\*x + 4\*x^2 + 3)^p,x)

[Out] int((4\*x + 4\*x^2 + 3)^p, x)

### 3.135 $\int (3 + 4x + 3x^2)^p dx$

Optimal result	639
Rubi [A] (verified)	639
Mathematica [A] (verified)	640
Maple [F]	640
Fricas [F]	640
Sympy [F]	641
Maxima [F]	641
Giac [F]	641
Mupad [F(-1)]	641

#### Optimal result

Integrand size = 12, antiderivative size = 37

$$\int (3 + 4x + 3x^2)^p dx = 3^{-1-p}5^p(2 + 3x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{5}(2 + 3x)^2\right)$$

[Out]  $3^{(-1-p)}5^p(2+3x)*\operatorname{hypergeom}([1/2, -p], [3/2], -1/5*(2+3x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x + 3x^2)^p dx = 3^{-p-1}5^p(3x + 2) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{5}(3x + 2)^2\right)$$

[In]  $\operatorname{Int}[(3 + 4*x + 3*x^2)^p, x]$

[Out]  $3^{(-1 - p)}5^p(2 + 3*x)*\operatorname{Hypergeometric2F1}[1/2, -p, 3/2, -1/5*(2 + 3*x)^2]$

#### Rule 251

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a^p*x*\operatorname{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; \operatorname{FreeQ}[a, b, n, p], x \ \&\& \ !\operatorname{IGtQ}[p, 0] \ \&\& \ !\operatorname{IntegerQ}[1/n] \ \&\& \ !\operatorname{ILtQ}[\operatorname{Simplify}[1/n + p], 0] \ \&\& \ (\operatorname{IntegerQ}[p] \ || \ \operatorname{GtQ}[a, 0])$

#### Rule 633

$\operatorname{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^{(p_.)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{Simp}[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b$

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}(3^{-1-p}5^p) \text{Subst}\left(\int \left(1 + \frac{x^2}{20}\right)^p dx, x, 4 + 6x\right) \\ &= 3^{-1-p}5^p(2 + 3x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; -\frac{1}{5}(2 + 3x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int (3 + 4x + 3x^2)^p dx = 3^{-1-p}5^p(2 + 3x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, -\frac{1}{5}(2 + 3x)^2\right)$$

[In] Integrate[(3 + 4\*x + 3\*x^2)^p, x]

[Out] 3^(-1 - p)\*5^p\*(2 + 3\*x)\*Hypergeometric2F1[1/2, -p, 3/2, -1/5\*(2 + 3\*x)^2]

**Maple [F]**

$$\int (3x^2 + 4x + 3)^p dx$$

[In] int((3\*x^2+4\*x+3)^p, x)

[Out] int((3\*x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3 + 4x + 3x^2)^p dx = \int (3x^2 + 4x + 3)^p dx$$

[In] integrate((3\*x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((3\*x^2 + 4\*x + 3)^p, x)



**Sympy [F]**

$$\int (3 + 4x + 3x^2)^p dx = \int (3x^2 + 4x + 3)^p dx$$

[In] integrate((3\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((3\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x + 3x^2)^p dx = \int (3x^2 + 4x + 3)^p dx$$

[In] integrate((3\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((3\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x + 3x^2)^p dx = \int (3x^2 + 4x + 3)^p dx$$

[In] integrate((3\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((3\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x + 3x^2)^p dx = \int (3x^2 + 4x + 3)^p dx$$

[In] int((4\*x + 3\*x^2 + 3)^p,x)

[Out] int((4\*x + 3\*x^2 + 3)^p, x)

### 3.136 $\int (3 + 4x + 2x^2)^p dx$

Optimal result	642
Rubi [A] (verified)	642
Mathematica [A] (verified)	643
Maple [F]	643
Fricas [F]	643
Sympy [F]	644
Maxima [F]	644
Giac [F]	644
Mupad [F(-1)]	644

#### Optimal result

Integrand size = 12, antiderivative size = 21

$$\int (3 + 4x + 2x^2)^p dx = (1 + x) \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -2(1 + x)^2 \right)$$

[Out] (1+x)\*hypergeom([1/2, -p], [3/2], -2\*(1+x)^2)

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x + 2x^2)^p dx = (x + 1) \operatorname{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -2(x + 1)^2 \right)$$

[In] Int[(3 + 4\*x + 2\*x^2)^p, x]

[Out] (1 + x)\*Hypergeometric2F1[1/2, -p, 3/2, -2\*(1 + x)^2]

#### Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F
1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p
, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] ||
GtQ[a, 0])
```

#### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c))))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```

`+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{4} \text{Subst} \left( \int \left( 1 + \frac{x^2}{8} \right)^p dx, x, 4 + 4x \right) \\ &= (1 + x) {}_2F_1 \left( \frac{1}{2}, -p; \frac{3}{2}; -2(1 + x)^2 \right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int (3 + 4x + 2x^2)^p dx = (1 + x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, -2(1 + x)^2 \right)$$

[In] `Integrate[(3 + 4*x + 2*x^2)^p, x]`

[Out] `(1 + x)*Hypergeometric2F1[1/2, -p, 3/2, -2*(1 + x)^2]`

**Maple [F]**

$$\int (2x^2 + 4x + 3)^p dx$$

[In] `int((2*x^2+4*x+3)^p, x)`

[Out] `int((2*x^2+4*x+3)^p, x)`

**Fricas [F]**

$$\int (3 + 4x + 2x^2)^p dx = \int (2x^2 + 4x + 3)^p dx$$

[In] `integrate((2*x^2+4*x+3)^p, x, algorithm="fricas")`

[Out] `integral((2*x^2 + 4*x + 3)^p, x)`

**Sympy [F]**

$$\int (3 + 4x + 2x^2)^p dx = \int (2x^2 + 4x + 3)^p dx$$

[In] integrate((2\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((2\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x + 2x^2)^p dx = \int (2x^2 + 4x + 3)^p dx$$

[In] integrate((2\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((2\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x + 2x^2)^p dx = \int (2x^2 + 4x + 3)^p dx$$

[In] integrate((2\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((2\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x + 2x^2)^p dx = \int (2x^2 + 4x + 3)^p dx$$

[In] int((4\*x + 2\*x^2 + 3)^p,x)

[Out] int((4\*x + 2\*x^2 + 3)^p, x)

### 3.137 $\int (3 + 4x + x^2)^p dx$

Optimal result	645
Rubi [A] (verified)	645
Mathematica [A] (verified)	646
Maple [F]	646
Fricas [F]	646
Sympy [F]	647
Maxima [F]	647
Giac [F]	647
Mupad [F(-1)]	647

#### Optimal result

Integrand size = 10, antiderivative size = 54

$$\int (3 + 4x + x^2)^p dx = -\frac{2^{1+2p}(-2-2x)^{-1-p}(3+4x+x^2)^{1+p} \operatorname{Hypergeometric2F1}\left(-p, 1+p, 2+p, \frac{3+x}{2}\right)}{1+p}$$

[Out]  $-2^{(1+2*p)}*(-2-2*x)^{(-1-p)}*(x^2+4*x+3)^{(p+1)}*\operatorname{hypergeom}([-p, p+1], [2+p], 3/2+1/2*x)/(p+1)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$ , Rules used = {638}

$$\int (3 + 4x + x^2)^p dx = -\frac{2^{2p+1}(-2x-2)^{-p-1}(x^2+4x+3)^{p+1} \operatorname{Hypergeometric2F1}\left(-p, p+1, p+2, \frac{x+3}{2}\right)}{p+1}$$

[In]  $\operatorname{Int}[(3 + 4*x + x^2)^p, x]$

[Out]  $-((2^{(1+2*p)}*(-2-2*x)^{(-1-p)}*(3+4*x+x^2)^{(1+p)}*\operatorname{Hypergeometric2F1}[-p, 1+p, 2+p, (3+x)/2])/(1+p))$

#### Rule 638

$\operatorname{Int}[(a_. + (b_.)*(x_) + (c_.)*(x_)^2)^{(p_.)}, x\_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Rt}[b^2 - 4*a*c, 2]\}, \operatorname{Simp}[(-a + b*x + c*x^2)^{(p+1)}/(q*(p+1)*((q-b-2*c*x)/(2*q))^{(p+1)})]*\operatorname{Hypergeometric2F1}[-p, p+1, p+2, (b+q+2*c*x)/(2*q)]$

], x]] /; FreeQ[{a, b, c, p}, x] && NeQ[b^2 - 4\*a\*c, 0] && !IntegerQ[4\*p]

Rubi steps

$$\text{integral} = -\frac{2^{1+2p}(-2-2x)^{-1-p}(3+4x+x^2)^{1+p} {}_2F_1(-p, 1+p; 2+p; \frac{3+x}{2})}{1+p}$$

**Mathematica [A] (verified)**

Time = 0.10 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.89

$$\int (3+4x+x^2)^p dx = \frac{2^p(1+x)(3+x)^{-p}(3+4x+x^2)^p \text{Hypergeometric2F1}(-p, 1+p, 2+p, \frac{1}{2}(-1-x))}{1+p}$$

[In] Integrate[(3 + 4\*x + x^2)^p, x]

[Out] (2^p\*(1 + x)\*(3 + 4\*x + x^2)^p\*Hypergeometric2F1[-p, 1 + p, 2 + p, (-1 - x)/2])/((1 + p)\*(3 + x)^p)

**Maple [F]**

$$\int (x^2 + 4x + 3)^p dx$$

[In] int((x^2+4\*x+3)^p, x)

[Out] int((x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3+4x+x^2)^p dx = \int (x^2+4x+3)^p dx$$

[In] integrate((x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x + x^2)^p dx = \int (x^2 + 4x + 3)^p dx$$

[In] integrate((x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x + x^2)^p dx = \int (x^2 + 4x + 3)^p dx$$

[In] integrate((x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x + x^2)^p dx = \int (x^2 + 4x + 3)^p dx$$

[In] integrate((x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x + x^2)^p dx = \int (x^2 + 4x + 3)^p dx$$

[In] int((4\*x + x^2 + 3)^p,x)

[Out] int((4\*x + x^2 + 3)^p, x)

### 3.138 $\int (3 + 4x)^p dx$

Optimal result	648
Rubi [A] (verified)	648
Mathematica [A] (verified)	649
Maple [A] (verified)	649
Fricas [A] (verification not implemented)	649
Sympy [A] (verification not implemented)	650
Maxima [A] (verification not implemented)	650
Giac [A] (verification not implemented)	650
Mupad [B] (verification not implemented)	651

#### Optimal result

Integrand size = 7, antiderivative size = 18

$$\int (3 + 4x)^p dx = \frac{(3 + 4x)^{1+p}}{4(1 + p)}$$

[Out] 1/4\*(3+4\*x)^(p+1)/(p+1)

#### Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$ , Rules used = {32}

$$\int (3 + 4x)^p dx = \frac{(4x + 3)^{p+1}}{4(p + 1)}$$

[In] Int[(3 + 4\*x)^p,x]

[Out] (3 + 4\*x)^(1 + p)/(4\*(1 + p))

#### Rule 32

Int[((a\_.) + (b\_.)\*(x\_))^(m\_), x\_Symbol] := Simp[(a + b\*x)^(m + 1)/(b\*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

#### Rubi steps

$$\text{integral} = \frac{(3 + 4x)^{1+p}}{4(1 + p)}$$



**Mathematica [A] (verified)**

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (3 + 4x)^p dx = \frac{(3 + 4x)^{1+p}}{4(1 + p)}$$

[In] Integrate[(3 + 4\*x)^p,x]

[Out] (3 + 4\*x)^(1 + p)/(4\*(1 + p))

**Maple [A] (verified)**

Time = 2.11 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

method	result	size
gospers	$\frac{(4x+3)^{1+p}}{4p+4}$	17
default	$\frac{(4x+3)^{1+p}}{4p+4}$	17
meijerg	$3^p x {}_2F_1\left(1, -p; 2; -\frac{4x}{3}\right)$	17
risch	$\frac{(4x+3)(4x+3)^p}{4p+4}$	20
parallelrisch	$\frac{12(4x+3)^p x + 9(4x+3)^p}{12+12p}$	28
norman	$\frac{x e^{p \ln(4x+3)}}{1+p} + \frac{3 e^{p \ln(4x+3)}}{4(1+p)}$	34

[In] int((4\*x+3)^p,x,method=\_RETURNVERBOSE)

[Out] 1/4\*(4\*x+3)^(1+p)/(1+p)

**Fricas [A] (verification not implemented)**

none

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.06

$$\int (3 + 4x)^p dx = \frac{(4x + 3)^p (4x + 3)}{4(p + 1)}$$

[In] integrate((3+4\*x)^p,x, algorithm="fricas")

[Out] 1/4\*(4\*x + 3)^p\*(4\*x + 3)/(p + 1)

**Sympy [A] (verification not implemented)**

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (3 + 4x)^p dx = \frac{\begin{cases} \frac{(4x+3)^{p+1}}{p+1} & \text{for } p \neq -1 \\ \log(4x + 3) & \text{otherwise} \end{cases}}{4}$$

[In] integrate((3+4\*x)\*\*p,x)

[Out] Piecewise(((4\*x + 3)\*\*(p + 1)/(p + 1), Ne(p, -1)), (log(4\*x + 3), True))/4

**Maxima [A] (verification not implemented)**

none

Time = 0.19 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int (3 + 4x)^p dx = \frac{(4x + 3)^{p+1}}{4(p + 1)}$$

[In] integrate((3+4\*x)^p,x, algorithm="maxima")

[Out] 1/4\*(4\*x + 3)^(p + 1)/(p + 1)

**Giac [A] (verification not implemented)**

none

Time = 0.27 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int (3 + 4x)^p dx = \frac{(4x + 3)^{p+1}}{4(p + 1)}$$

[In] integrate((3+4\*x)^p,x, algorithm="giac")

[Out] 1/4\*(4\*x + 3)^(p + 1)/(p + 1)

**Mupad [B] (verification not implemented)**

Time = 9.25 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.78

$$\int (3 + 4x)^p dx = \begin{cases} \frac{\ln(4x+3)}{4} & \text{if } p = -1 \\ \frac{(4x+3)^{p+1}}{4(p+1)} & \text{if } p \neq -1 \end{cases}$$

[In] `int((4*x + 3)^p,x)`

[Out] `piecewise(p == -1, log(4*x + 3)/4, p ~= -1, (4*x + 3)^(p + 1)/(4*(p + 1)))`

### 3.139 $\int (3 + 4x - x^2)^p dx$

Optimal result	652
Rubi [A] (verified)	652
Mathematica [A] (verified)	653
Maple [F]	653
Fricas [F]	653
Sympy [F]	654
Maxima [F]	654
Giac [F]	654
Mupad [F(-1)]	654

#### Optimal result

Integrand size = 12, antiderivative size = 31

$$\int (3 + 4x - x^2)^p dx = -7^p(2 - x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{7}(2 - x)^2\right)$$

[Out]  $-7^p(2-x)*\operatorname{hypergeom}([1/2, -p], [3/2], 1/7*(2-x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x - x^2)^p dx = -7^p(2 - x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{7}(2 - x)^2\right)$$

[In]  $\operatorname{Int}[(3 + 4*x - x^2)^p, x]$

[Out]  $-(7^p(2 - x)*\operatorname{Hypergeometric2F1}[1/2, -p, 3/2, (2 - x)^2/7])$

#### Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F
1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p
, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] ||
GtQ[a, 0])
```

#### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c))))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(\frac{1}{2}7^p \text{Subst}\left(\int \left(1 - \frac{x^2}{28}\right)^p dx, x, 4 - 2x\right)\right) \\ &= -7^p(2 - x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; \frac{1}{7}(2 - x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int (3 + 4x - x^2)^p dx = 7^p(-2 + x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{7}(-2 + x)^2\right)$$

[In] Integrate[(3 + 4\*x - x^2)^p, x]

[Out] 7^p\*(-2 + x)\*Hypergeometric2F1[1/2, -p, 3/2, (-2 + x)^2/7]

**Maple [F]**

$$\int (-x^2 + 4x + 3)^p dx$$

[In] int((-x^2+4\*x+3)^p, x)

[Out] int((-x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3 + 4x - x^2)^p dx = \int (-x^2 + 4x + 3)^p dx$$

[In] integrate((-x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((-x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x - x^2)^p dx = \int (-x^2 + 4x + 3)^p dx$$

[In] integrate((-x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((-x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x - x^2)^p dx = \int (-x^2 + 4x + 3)^p dx$$

[In] integrate((-x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((-x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x - x^2)^p dx = \int (-x^2 + 4x + 3)^p dx$$

[In] integrate((-x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((-x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x - x^2)^p dx = \int (-x^2 + 4x + 3)^p dx$$

[In] int((4\*x - x^2 + 3)^p,x)

[Out] int((4\*x - x^2 + 3)^p, x)

### 3.140 $\int (3 + 4x - 2x^2)^p dx$

Optimal result	655
Rubi [A] (verified)	655
Mathematica [A] (verified)	656
Maple [F]	656
Fricas [F]	656
Sympy [F]	657
Maxima [F]	657
Giac [F]	657
Mupad [F(-1)]	657

#### Optimal result

Integrand size = 12, antiderivative size = 31

$$\int (3 + 4x - 2x^2)^p dx = -5^p(1-x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{2}{5}(1-x)^2\right)$$

[Out]  $-5^p(1-x)\operatorname{hypergeom}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{2}{5}(1-x)^2\right)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x - 2x^2)^p dx = -5^p(1-x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{2}{5}(1-x)^2\right)$$

[In]  $\operatorname{Int}[(3 + 4*x - 2*x^2)^p, x]$

[Out]  $-(5^p(1-x)\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, -p, \frac{3}{2}, \frac{2*(1-x)^2}{5}\right])$

#### Rule 251

$\operatorname{Int}[(a_.) + (b_.)*(x_)^{(n_)}]^{(p_)}, x\_Symbol] \rightarrow \operatorname{Simp}[a^p*x*\operatorname{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; \operatorname{FreeQ}\{a, b, n, p\}, x \ \&\& \ !\operatorname{IGtQ}[p, 0] \ \&\& \ !\operatorname{IntegerQ}[1/n] \ \&\& \ !\operatorname{ILtQ}[\operatorname{Simplify}[1/n + p], 0] \ \&\& \ (\operatorname{IntegerQ}[p] \ || \ \operatorname{GtQ}[a, 0])$

#### Rule 633

$\operatorname{Int}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^{(p_)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), \operatorname{Subst}[\operatorname{Int}[\operatorname{Simp}[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b$

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(\frac{1}{4}5^p \text{Subst}\left(\int \left(1 - \frac{x^2}{40}\right)^p dx, x, 4 - 4x\right)\right) \\ &= -5^p(1-x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; \frac{2}{5}(1-x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.84

$$\int (3 + 4x - 2x^2)^p dx = 5^p(-1 + x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{2}{5}(-1 + x)^2\right)$$

[In] Integrate[(3 + 4\*x - 2\*x^2)^p, x]

[Out] 5^p\*(-1 + x)\*Hypergeometric2F1[1/2, -p, 3/2, (2\*(-1 + x)^2)/5]

**Maple [F]**

$$\int (-2x^2 + 4x + 3)^p dx$$

[In] int((-2\*x^2+4\*x+3)^p, x)

[Out] int((-2\*x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3 + 4x - 2x^2)^p dx = \int (-2x^2 + 4x + 3)^p dx$$

[In] integrate((-2\*x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((-2\*x^2 + 4\*x + 3)^p, x)



**Sympy [F]**

$$\int (3 + 4x - 2x^2)^p dx = \int (-2x^2 + 4x + 3)^p dx$$

[In] integrate((-2\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((-2\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x - 2x^2)^p dx = \int (-2x^2 + 4x + 3)^p dx$$

[In] integrate((-2\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((-2\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x - 2x^2)^p dx = \int (-2x^2 + 4x + 3)^p dx$$

[In] integrate((-2\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((-2\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x - 2x^2)^p dx = \int (-2x^2 + 4x + 3)^p dx$$

[In] int((4\*x - 2\*x^2 + 3)^p,x)

[Out] int((4\*x - 2\*x^2 + 3)^p, x)

### 3.141 $\int (3 + 4x - 3x^2)^p dx$

Optimal result	658
Rubi [A] (verified)	658
Mathematica [A] (verified)	659
Maple [F]	659
Fricas [F]	659
Sympy [F]	660
Maxima [F]	660
Giac [F]	660
Mupad [F(-1)]	660

#### Optimal result

Integrand size = 12, antiderivative size = 38

$$\int (3 + 4x - 3x^2)^p dx = -3^{-1-p}13^p(2 - 3x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{13}(2 - 3x)^2 \right)$$

[Out]  $-3^{(-1-p)}*13^p*(2-3*x)*\text{hypergeom}([1/2, -p], [3/2], 1/13*(2-3*x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x - 3x^2)^p dx = -3^{-p-1}13^p(2 - 3x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{13}(2 - 3x)^2 \right)$$

[In]  $\text{Int}[(3 + 4*x - 3*x^2)^p, x]$

[Out]  $-(3^{(-1 - p)}*13^p*(2 - 3*x)*\text{Hypergeometric2F1}[1/2, -p, 3/2, (2 - 3*x)^2/13])$

#### Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])
```

#### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(\frac{1}{2}(3^{-1-p}13^p) \text{Subst}\left(\int\left(1-\frac{x^2}{52}\right)^p dx, x, 4-6x\right)\right) \\ &= -3^{-1-p}13^p(2-3x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; \frac{1}{13}(2-3x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.97

$$\int (3 + 4x - 3x^2)^p dx = 3^{-1-p}13^p(-2 + 3x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{13}(2 - 3x)^2\right)$$

[In] Integrate[(3 + 4\*x - 3\*x^2)^p,x]

[Out] 3^(-1 - p)\*13^p\*(-2 + 3\*x)\*Hypergeometric2F1[1/2, -p, 3/2, (2 - 3\*x)^2/13]

**Maple [F]**

$$\int (-3x^2 + 4x + 3)^p dx$$

[In] int((-3\*x^2+4\*x+3)^p,x)

[Out] int((-3\*x^2+4\*x+3)^p,x)

**Fricas [F]**

$$\int (3 + 4x - 3x^2)^p dx = \int (-3x^2 + 4x + 3)^p dx$$

[In] integrate((-3\*x^2+4\*x+3)^p,x, algorithm="fricas")

[Out] integral((-3\*x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x - 3x^2)^p dx = \int (-3x^2 + 4x + 3)^p dx$$

[In] integrate((-3\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((-3\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x - 3x^2)^p dx = \int (-3x^2 + 4x + 3)^p dx$$

[In] integrate((-3\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((-3\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x - 3x^2)^p dx = \int (-3x^2 + 4x + 3)^p dx$$

[In] integrate((-3\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((-3\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x - 3x^2)^p dx = \int (-3x^2 + 4x + 3)^p dx$$

[In] int((4\*x - 3\*x^2 + 3)^p,x)

[Out] int((4\*x - 3\*x^2 + 3)^p, x)

### 3.142 $\int (3 + 4x - 4x^2)^p dx$

Optimal result	661
Rubi [A] (verified)	661
Mathematica [A] (verified)	662
Maple [F]	662
Fricas [F]	662
Sympy [F]	663
Maxima [F]	663
Giac [F]	663
Mupad [F(-1)]	663

#### Optimal result

Integrand size = 12, antiderivative size = 35

$$\int (3 + 4x - 4x^2)^p dx = -2^{-1+2p}(1 - 2x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{4}(1 - 2x)^2\right)$$

[Out]  $-2^{(-1+2*p)}*(1-2*x)*\operatorname{hypergeom}([1/2, -p], [3/2], 1/4*(1-2*x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x - 4x^2)^p dx = -2^{2p-1}(1 - 2x) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{4}(1 - 2x)^2\right)$$

[In]  $\operatorname{Int}[(3 + 4*x - 4*x^2)^p, x]$

[Out]  $-(2^{(-1 + 2*p)}*(1 - 2*x)*\operatorname{Hypergeometric2F1}[1/2, -p, 3/2, (1 - 2*x)^2/4])$

#### Rule 251

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Simp}[a^p*x*\operatorname{Hypergeometric2F1}[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /;$   $\operatorname{FreeQ}\{a, b, n, p\}, x$  &&  $! \operatorname{IGtQ}[p, 0]$  &&  $! \operatorname{IntegerQ}[1/n]$  &&  $! \operatorname{ILtQ}[\operatorname{Simplify}[1/n + p], 0]$  &&  $(\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[a, 0])$

#### Rule 633

$\operatorname{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(p_.)}, x\_Symbol] \rightarrow \operatorname{Dist}[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^{(p)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{Simp}[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b$

+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(2^{-3+2p} \text{Subst}\left(\int \left(1 - \frac{x^2}{64}\right)^p dx, x, 4 - 8x\right)\right) \\ &= -2^{-1+2p}(1 - 2x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; \frac{1}{4}(1 - 2x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.07 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int (3 + 4x - 4x^2)^p dx = -2^{-3+2p}(4 - 8x) \text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{64}(4 - 8x)^2\right)$$

[In] Integrate[(3 + 4\*x - 4\*x^2)^p, x]

[Out] -(2^(-3 + 2\*p))\*(4 - 8\*x)\*Hypergeometric2F1[1/2, -p, 3/2, (4 - 8\*x)^2/64]

**Maple [F]**

$$\int (-4x^2 + 4x + 3)^p dx$$

[In] int((-4\*x^2+4\*x+3)^p, x)

[Out] int((-4\*x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3 + 4x - 4x^2)^p dx = \int (-4x^2 + 4x + 3)^p dx$$

[In] integrate((-4\*x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((-4\*x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x - 4x^2)^p dx = \int (-4x^2 + 4x + 3)^p dx$$

[In] integrate((-4\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((-4\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x - 4x^2)^p dx = \int (-4x^2 + 4x + 3)^p dx$$

[In] integrate((-4\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((-4\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x - 4x^2)^p dx = \int (-4x^2 + 4x + 3)^p dx$$

[In] integrate((-4\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((-4\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x - 4x^2)^p dx = \int (-4x^2 + 4x + 3)^p dx$$

[In] int((4\*x - 4\*x^2 + 3)^p,x)

[Out] int((4\*x - 4\*x^2 + 3)^p, x)

### 3.143 $\int (3 + 4x - 5x^2)^p dx$

Optimal result	664
Rubi [A] (verified)	664
Mathematica [A] (verified)	665
Maple [F]	665
Fricas [F]	665
Sympy [F]	666
Maxima [F]	666
Giac [F]	666
Mupad [F(-1)]	666

#### Optimal result

Integrand size = 12, antiderivative size = 38

$$\int (3 + 4x - 5x^2)^p dx = -5^{-1-p} 19^p (2 - 5x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{19}(2 - 5x)^2 \right)$$

[Out]  $-5^{(-1-p)} * 19^p * (2-5*x) * \text{hypergeom}([1/2, -p], [3/2], 1/19*(2-5*x)^2)$

#### Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2,  $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$ , Rules used = {633, 251}

$$\int (3 + 4x - 5x^2)^p dx = -5^{-p-1} 19^p (2 - 5x) \text{Hypergeometric2F1} \left( \frac{1}{2}, -p, \frac{3}{2}, \frac{1}{19}(2 - 5x)^2 \right)$$

[In]  $\text{Int}[(3 + 4*x - 5*x^2)^p, x]$

[Out]  $-(5^{(-1-p)} * 19^p * (2-5*x) * \text{Hypergeometric2F1}[1/2, -p, 3/2, (2-5*x)^2/19])$

#### Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] || GtQ[a, 0])
```

#### Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c))))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```



+ 2\*c\*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4\*a - b^2/c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\left(\frac{1}{2}(5^{-1-p}19^p)\text{Subst}\left(\int\left(1-\frac{x^2}{76}\right)^p dx, x, 4-10x\right)\right) \\ &= -5^{-1-p}19^p(2-5x) {}_2F_1\left(\frac{1}{2}, -p; \frac{3}{2}; \frac{1}{19}(2-5x)^2\right) \end{aligned}$$

**Mathematica [A] (verified)**

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.97

$$\int (3+4x-5x^2)^p dx = 5^{-1-p}19^p(-2+5x)\text{Hypergeometric2F1}\left(\frac{1}{2}, -p, \frac{3}{2}, \frac{1}{19}(2-5x)^2\right)$$

[In] Integrate[(3 + 4\*x - 5\*x^2)^p, x]

[Out] 5^(-1 - p)\*19^p\*(-2 + 5\*x)\*Hypergeometric2F1[1/2, -p, 3/2, (2 - 5\*x)^2/19]

**Maple [F]**

$$\int (-5x^2 + 4x + 3)^p dx$$

[In] int((-5\*x^2+4\*x+3)^p, x)

[Out] int((-5\*x^2+4\*x+3)^p, x)

**Fricas [F]**

$$\int (3+4x-5x^2)^p dx = \int (-5x^2 + 4x + 3)^p dx$$

[In] integrate((-5\*x^2+4\*x+3)^p, x, algorithm="fricas")

[Out] integral((-5\*x^2 + 4\*x + 3)^p, x)

**Sympy [F]**

$$\int (3 + 4x - 5x^2)^p dx = \int (-5x^2 + 4x + 3)^p dx$$

[In] integrate((-5\*x\*\*2+4\*x+3)\*\*p,x)

[Out] Integral((-5\*x\*\*2 + 4\*x + 3)\*\*p, x)

**Maxima [F]**

$$\int (3 + 4x - 5x^2)^p dx = \int (-5x^2 + 4x + 3)^p dx$$

[In] integrate((-5\*x^2+4\*x+3)^p,x, algorithm="maxima")

[Out] integrate((-5\*x^2 + 4\*x + 3)^p, x)

**Giac [F]**

$$\int (3 + 4x - 5x^2)^p dx = \int (-5x^2 + 4x + 3)^p dx$$

[In] integrate((-5\*x^2+4\*x+3)^p,x, algorithm="giac")

[Out] integrate((-5\*x^2 + 4\*x + 3)^p, x)

**Mupad [F(-1)]**

Timed out.

$$\int (3 + 4x - 5x^2)^p dx = \int (-5x^2 + 4x + 3)^p dx$$

[In] int((4\*x - 5\*x^2 + 3)^p,x)

[Out] int((4\*x - 5\*x^2 + 3)^p, x)

---

---

# CHAPTER 4

---

## APPENDIX

4.1 Listing of Grading functions . . . . . 667

### 4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

#### Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

## Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_coun
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```



```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`') or type(expn,'*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

## Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

## SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-t
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```



```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```