

Computer Algebra Independent Integration Tests

Summer 2023 edition

213-Goursat

Nasser M. Abbasi

October 31, 2023

Compiled on October 31, 2023 at 6:02pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	31
4	Appendix	145

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [15]. This is test number [213].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Maple	100.00 (15)	0.00 (0)
Fricas	93.33 (14)	6.67 (1)
Mathematica	73.33 (11)	26.67 (4)
Rubi	66.67 (10)	33.33 (5)
Maxima	13.33 (2)	86.67 (13)
Sympy	13.33 (2)	86.67 (13)
Mupad	6.67 (1)	93.33 (14)
Giac	0.00 (0)	100.00 (15)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Fricas	86.667	6.667	0.000	6.667
Mathematica	66.667	0.000	6.667	26.667
Rubi	20.000	0.000	46.667	33.333
Maxima	13.333	0.000	0.000	86.667
Maple	0.000	0.000	100.000	0.000
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	6.667	0.000	93.333
Sympy	0.000	0.000	13.333	86.667

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Maple	0	0.00	0.00	0.00
Fricas	1	0.00	0.00	100.00
Mathematica	4	100.00	0.00	0.00
Rubi	5	100.00	0.00	0.00
Maxima	13	100.00	0.00	0.00
Sympy	13	76.92	23.08	0.00
Mupad	14	0.00	100.00	0.00
Giac	15	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maxima	0.24
Rubi	0.31
Mupad	0.46
Fricas	0.86
Sympy	1.25
Maple	1.48
Mathematica	6.27
Giac	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	62.50	0.67	62.50	0.67
Mupad	100.00	1.09	100.00	1.09
Mathematica	103.45	0.98	107.00	1.00
Fricas	117.36	0.98	114.00	0.94
Maxima	124.00	1.33	124.00	1.33
Rubi	207.60	1.92	165.00	1.60
Maple	453.60	3.68	458.00	3.44
Giac	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for intergals that required less steps by Rubi, CAS systems had more success which indicates the intergal was not as hard to solve. As Rubi needed more steps to solve the intergal, the solved percentage decreased for most CAS systems which indicates the intergal is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {11}

Mathematica {}

Maple {1, 2, 3, 4, 5}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
```

```

Return the tree size of this expression.
"""
if expr not in SR:
    # deal with lists, tuples, vectors
    return 1 + sum(tree_size(a) for a in expr)
expr = SR(expr)
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)

```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```

try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1

```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```


integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)

```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer, 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer, Leaf size of result.
4. integer, Leaf size of the optimal antiderivative.
5. number, CPU time used to solve this integral. 0 if failed.
6. string, The integral in Latex format
7. string, The input used in CAS own syntax.
8. string, The result (antiderivative) produced by CAS in Latex format
9. string, The optimal antiderivative in Latex format.
10. integer, 0 or 1, Indicates if problem has known antiderivative or not
11. String, The result (antiderivative) in CAS own syntax.
12. String, The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String, Small string description of why the grade was given.
14. integer, 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer, Number of steps used.
16. integer, Number of rules used.
17. integer, Integrand leaf size.
18. real number, Ratio, Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String, The optimal antiderivative in Mathematica syntax

Nasir M Abbasi
June 27, 2023
Diagrams

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	29

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	23
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 4 }

B grade { }

C grade { 3, 5, 6, 7, 8, 10, 11 }

F normal fail { 9, 12, 13, 14, 15 }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 6, 7, 8, 9, 10, 13 }

B grade { }

C grade { 5 }

F normal fail { 11, 12, 14, 15 }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple**A grade** { }**B grade** { }**C grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Fricas****A grade** { 1, 2, 3, 4, 6, 7, 8, 10, 11, 12, 13, 14, 15 }**B grade** { 5 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { 9 }**Maxima****A grade** { 1, 2 }**B grade** { }**C grade** { }**F normal fail** { 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Giac****A grade** { }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 }**F(-1) timeout fail** { }**F(-2) exception fail** { }

Mupad

A grade { }

B grade { 1 }

C grade { }

F normal fail { }

F(-1) timeout fail { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 }

F(-2) exception fail { }

Sympy

A grade { }

B grade { }

C grade { 1, 2 }

F normal fail { 3, 4, 5, 7, 10, 11, 12, 13, 14, 15 }

F(-1) timeout fail { 6, 8, 9 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	C	F	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	92	95	92	113	124	87	60	0	100
N.S.	1	1.03	1.00	1.23	1.35	0.95	0.65	0.00	1.09
time (sec)	N/A	0.087	0.469	0.939	0.231	0.383	1.394	0.000	0.458

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	A	A	C	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	95	95	86	99	124	88	65	0	0
N.S.	1	1.00	0.91	1.04	1.31	0.93	0.68	0.00	0.00
time (sec)	N/A	0.047	1.152	0.669	0.240	0.397	1.100	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	107	196	107	164	0	109	0	0	0
N.S.	1	1.83	1.00	1.53	0.00	1.02	0.00	0.00	0.00
time (sec)	N/A	0.127	1.662	0.794	0.000	0.396	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	111	118	111	125	0	119	0	0	0
N.S.	1	1.06	1.00	1.13	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	0.142	0.592	0.939	0.000	0.351	0.000	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	C	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	147	239	123	133	0	125	0	0	0
N.S.	1	1.63	0.84	0.90	0.00	0.85	0.00	0.00	0.00
time (sec)	N/A	0.154	10.156	0.664	0.000	0.359	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	261	90	290	0	104	0	0	0
N.S.	1	2.90	1.00	3.22	0.00	1.16	0.00	0.00	0.00
time (sec)	N/A	0.426	0.194	0.466	0.000	1.358	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	134	85	292	0	78	0	0	0
N.S.	1	1.58	1.00	3.44	0.00	0.92	0.00	0.00	0.00
time (sec)	N/A	0.350	1.460	0.526	0.000	1.156	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	103	110	600	0	133	0	0	0
N.S.	1	0.94	1.00	5.45	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.298	1.049	5.344	0.000	1.499	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	F	F(-2)	F(-1)	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	0	119	1329	0	0	0	0	0
N.S.	1	0.00	1.00	11.17	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.000	5.407	6.459	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	A	C	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	102	298	102	607	0	106	0	0	0
N.S.	1	2.92	1.00	5.95	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.481	15.220	0.812	0.000	0.647	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	C	F	C	F	A	F	F	F(-1)
verified	N/A	No	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	125	537	0	606	0	107	0	0	0
N.S.	1	4.30	0.00	4.85	0.00	0.86	0.00	0.00	0.00
time (sec)	N/A	1.005	0.000	0.604	0.000	0.498	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	F	C	F	A	F	F	F(-1)
verified	N/A	N/A	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	166	0	0	461	0	137	0	0	0
N.S.	1	0.00	0.00	2.78	0.00	0.83	0.00	0.00	0.00
time (sec)	N/A	0.000	0.000	0.697	0.000	0.651	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	C	F	A	F	F	F(-1)
verified	N/A	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	113	0	113	458	0	130	0	0	0
N.S.	1	0.00	1.00	4.05	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	0.000	31.584	1.348	0.000	0.515	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	F	C	F	A	F	F	F(-1)
verified	N/A	N/A	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	166	0	0	682	0	146	0	0	0
N.S.	1	0.00	0.00	4.11	0.00	0.88	0.00	0.00	0.00
time (sec)	N/A	0.000	0.000	0.635	0.000	0.654	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	F	C	F	A	F	F	F(-1)
verified	N/A	N/A	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	194	0	0	845	0	174	0	0	0
N.S.	1	0.00	0.00	4.36	0.00	0.90	0.00	0.00	0.00
time (sec)	N/A	0.000	0.000	1.282	0.000	3.209	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [5] had the largest ratio of [.761900000000000022]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	7	1.03	16	0.438
2	A	8	7	1.00	18	0.389
3	C	16	15	1.83	21	0.714
4	A	16	12	1.06	29	0.414
5	C	23	16	1.63	21	0.762
6	C	14	4	2.90	26	0.154
7	C	7	3	1.58	28	0.107
8	C	7	3	0.94	30	0.100
9	F	0	0	N/A	0.000	N/A
10	C	12	10	2.92	24	0.417
11	C	14	12	4.30	27	0.444
12	F	0	0	N/A	0.000	N/A
13	F	0	0	N/A	0.000	N/A
14	F	0	0	N/A	0.000	N/A
15	F	0	0	N/A	0.000	N/A

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{1+x}{x \sqrt[3]{-1+x^3}} dx$	32
3.2	$\int \frac{-1+x^2}{x(-1+x^3)^{2/3}} dx$	38
3.3	$\int \frac{(1+x) \sqrt[3]{-1+x^3}}{(-1+x)^2 x} dx$	44
3.4	$\int \frac{(-1+x)^2 (1+x)}{x(1+x+x^2) \sqrt[3]{-1+x^3}} dx$	52
3.5	$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx$	59
3.6	$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx$	67
3.7	$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx$	73
3.8	$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx$	78
3.9	$\int \frac{(2+x+x^2) \sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx$	83
3.10	$\int \frac{1+3x}{(-1+3x) \sqrt[3]{-x+x^3}} dx$	94
3.11	$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx$	102
3.12	$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x) \sqrt[3]{-x+x^3}} dx$	112
3.13	$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx$	121
3.14	$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x) \sqrt[3]{-x+x^3}} dx$	130
3.15	$\int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx$	139

3.1 $\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx$

Optimal result	32
Rubi [A] (verified)	32
Mathematica [A] (verified)	34
Maple [C] (warning: unable to verify)	35
Fricas [A] (verification not implemented)	35
Sympy [C] (verification not implemented)	36
Maxima [A] (verification not implemented)	36
Giac [F]	37
Mupad [B] (verification not implemented)	37

Optimal result

Integrand size = 16, antiderivative size = 92

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx = -\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) - \log\left(1-x+\sqrt[3]{-1+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)$$

[Out] $-3^{(1/2)}*\arctan(3^{(1/2)}*(x^3-1)^{(1/3)/(-2+2*x+(x^3-1)^{(1/3))})-\ln(1-x+(x^3-1)^{(1/3))}+1/2*\ln(1-2*x+x^2+(-1+x)*(x^3-1)^{(1/3)}+(x^3-1)^{(2/3))}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.03, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {1858, 245, 272, 58, 632, 210, 31}

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx = \frac{\arctan\left(\frac{\sqrt[3]{x^3-1}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1-2\sqrt[3]{x^3-1}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}+1\right) - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}-x\right) + \frac{\log(x)}{2}$$

[In] $\text{Int}[(1+x)/(x*(-1+x^3)^{(1/3)}),x]$

[Out] $\text{ArcTan}[(1+(2*x)/(-1+x^3)^{(1/3)})/\text{Sqrt}[3]]/\text{Sqrt}[3] - \text{ArcTan}[(1-2*(-1+x^3)^{(1/3)})/\text{Sqrt}[3]]/\text{Sqrt}[3] + \text{Log}[x]/2 - \text{Log}[1+(-1+x^3)^{(1/3)}]/2 - \text{Log}[-x+(-1+x^3)^{(1/3)}]/2$

Rule 31

$\text{Int}[\frac{(a + b \cdot x)^{-1}}{b}, x] \text{ ; FreeQ}\{a, b, x\} \text{ := Simp}[\frac{\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]}{b}, x] \text{ ; FreeQ}\{a, b, x\}$

Rule 58

$\text{Int}[\frac{1}{((a + b \cdot x) \cdot ((c + d \cdot x)^{1/3}))}, x] \text{ := With}[\{q = \text{Rt}[-(b \cdot c - a \cdot d)/b, 3]\}, \text{Simp}[\frac{\text{Log}[\text{RemoveContent}[a + b \cdot x, x]]}{(2 \cdot b \cdot q)}, x] + (\text{Dist}[3/(2 \cdot b), \text{Subst}[\text{Int}[1/(q^2 - q \cdot x + x^2), x], x, (c + d \cdot x)^{1/3}], x] - \text{Dist}[3/(2 \cdot b \cdot q), \text{Subst}[\text{Int}[1/(q + x), x], x, (c + d \cdot x)^{1/3}], x])] \text{ ; FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NegQ}[(b \cdot c - a \cdot d)/b]$

Rule 210

$\text{Int}[\frac{(a + b \cdot x^2)^{-1}}{x}, x] \text{ := Simp}[\frac{(-\text{Rt}[-a, 2] \cdot \text{Rt}[-b, 2])^{-1} \cdot \text{ArcTan}[\text{Rt}[-b, 2] \cdot (x/\text{Rt}[-a, 2])]}{x}], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 245

$\text{Int}[\frac{(a + b \cdot x^3)^{-1/3}}{x}, x] \text{ := Simp}[\frac{\text{ArcTan}[(1 + 2 \cdot \text{Rt}[b, 3] \cdot (x/(a + b \cdot x^3)^{1/3}))]/\text{Sqrt}[3]}{(\text{Sqrt}[3] \cdot \text{Rt}[b, 3])}, x] - \text{Simp}[\frac{\text{Log}[(a + b \cdot x^3)^{1/3} - \text{Rt}[b, 3] \cdot x]}{(2 \cdot \text{Rt}[b, 3])}, x] \text{ ; FreeQ}\{a, b, x\}$

Rule 272

$\text{Int}[x^m \cdot (a + b \cdot x^n)^p, x] \text{ := Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1) \cdot (a + b \cdot x)^p}, x], x, x^n], x] \text{ ; FreeQ}\{a, b, m, n, p, x\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 632

$\text{Int}[\frac{(a + b \cdot x + c \cdot x^2)^{-1}}{x}, x] \text{ := Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4 \cdot a \cdot c - x^2, x], x], x, b + 2 \cdot c \cdot x], x] \text{ ; FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4 \cdot a \cdot c, 0]$

Rule 1858

$\text{Int}[(Pq) \cdot ((c + d \cdot x)^m \cdot (a + b \cdot x^n)^p), x] \text{ := Int}[\text{ExpandIntegrand}[(c \cdot x)^m \cdot Pq \cdot (a + b \cdot x^n)^p, x], x] \text{ ; FreeQ}\{a, b, c, m, n, p, x\} \ \&\& \ (\text{PolyQ}[Pq, x] \ || \ \text{PolyQ}[Pq, x^n]) \ \&\& \ !\text{IGtQ}[m, 0]$

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{\sqrt[3]{-1+x^3}} + \frac{1}{x\sqrt[3]{-1+x^3}} \right) dx \\
&= \int \frac{1}{\sqrt[3]{-1+x^3}} dx + \int \frac{1}{x\sqrt[3]{-1+x^3}} dx \\
&= \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log(-x + \sqrt[3]{-1+x^3}) + \frac{1}{3} \text{Subst}\left(\int \frac{1}{\sqrt[3]{-1+xx}} dx, x, x^3\right) \\
&= \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log(-x + \sqrt[3]{-1+x^3}) \\
&\quad - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3}\right) + \frac{1}{2} \text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3}\right) \\
&= \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log(1 + \sqrt[3]{-1+x^3}) \\
&\quad - \frac{1}{2} \log(-x + \sqrt[3]{-1+x^3}) - \text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1 + 2\sqrt[3]{-1+x^3}\right) \\
&= \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{-1+2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{\sqrt{3}} \\
&\quad + \frac{\log(x)}{2} - \frac{1}{2} \log(1 + \sqrt[3]{-1+x^3}) - \frac{1}{2} \log(-x + \sqrt[3]{-1+x^3})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx &= -\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) - \log(1-x+\sqrt[3]{-1+x^3}) \\
&\quad + \frac{1}{2} \log(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3})
\end{aligned}$$

[In] Integrate[(1+x)/(x*(-1+x^3)^(1/3)),x]

[Out] -(Sqrt[3]*ArcTan[(Sqrt[3]*(-1+x^3)^(1/3))/(-2+2*x+(-1+x^3)^(1/3))])
- Log[1-x+(-1+x^3)^(1/3)] + Log[1-2*x+x^2+(-1+x)*(-1+x^3)^(1/3)
+ (-1+x^3)^(2/3)]/2

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.94 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.23

method	result
meijerg	$\frac{\sqrt{3}\Gamma\left(\frac{2}{3}\right)(-\text{signum}(x^3-1))^{\frac{1}{3}}\left(\frac{2\left(-\frac{\pi\sqrt{3}}{6}-\frac{3\ln(3)}{2}+3\ln(x)+i\pi\right)\pi\sqrt{3}+2\pi\sqrt{3}x^3\text{hypergeom}\left(\left[1,1,\frac{4}{3}\right],[2,2],x^3\right)}{3\Gamma\left(\frac{2}{3}\right)}\right)}{6\pi\text{signum}(x^3-1)^{\frac{1}{3}}} + \frac{(-\text{signum}(x^3-1))^{\frac{1}{3}}x}{\text{signum}(x^3-1)}$
trager	$-\ln\left(\frac{12\text{RootOf}\left(_Z^2-_Z+1\right)^2x^2-30\text{RootOf}\left(_Z^2-_Z+1\right)^2x+99\text{RootOf}\left(_Z^2-_Z+1\right)(x^3-1)^{\frac{2}{3}}-127\text{RootOf}\left(_Z^2-_Z+1\right)(x^3-1)^{\frac{1}{3}}}{\dots}\right)$

[In] int((1+x)/x/(x^3-1)^(1/3),x,method=_RETURNVERBOSE)

[Out] 1/6/Pi*3^(1/2)*GAMMA(2/3)/signum(x^3-1)^(1/3)*(-signum(x^3-1))^(1/3)*(2/3*(-1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*Pi*3^(1/2)/GAMMA(2/3)+2/9*Pi*3^(1/2)/GAMMA(2/3)*x^3*hypergeom([1,1,4/3],[2,2],x^3))+1/signum(x^3-1)^(1/3)*(-signum(x^3-1))^(1/3)*x*hypergeom([1/3,1/3],[4/3],x^3)

Fricas [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 87, normalized size of antiderivative = 0.95

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx$$

$$= \sqrt{3} \arctan\left(\frac{4\sqrt{3}(x^3-1)^{\frac{1}{3}}(x-1) + \sqrt{3}(x^2+x+1) - 2\sqrt{3}(x^3-1)^{\frac{2}{3}}}{3(3x^2-5x+3)}\right)$$

$$- \frac{1}{2} \log\left(\frac{(x^3-1)^{\frac{1}{3}}(x-1) + x - (x^3-1)^{\frac{2}{3}}}{x}\right)$$

[In] integrate((1+x)/x/(x^3-1)^(1/3),x, algorithm="fricas")

[Out] sqrt(3)*arctan(-1/3*(4*sqrt(3)*(x^3-1)^(1/3)*(x-1) + sqrt(3)*(x^2+x+1) - 2*sqrt(3)*(x^3-1)^(2/3))/(3*x^2-5*x+3)) - 1/2*log(((x^3-1)^(1/3)*(x-1) + x - (x^3-1)^(2/3))/x)

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.39 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.65

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx = \frac{x e^{-\frac{i\pi}{3}} \Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| x^3\right)}{3\Gamma\left(\frac{4}{3}\right)} - \frac{\Gamma\left(\frac{1}{3}\right) {}_2F_1\left(\frac{1}{3}, \frac{1}{3} \middle| \frac{4}{3} \middle| \frac{e^{2i\pi}}{x^3}\right)}{3x\Gamma\left(\frac{4}{3}\right)}$$

[In] integrate((1+x)/x/(x**3-1)**(1/3),x)

[Out] x*exp(-I*pi/3)*gamma(1/3)*hyper((1/3, 1/3), (4/3,), x**3)/(3*gamma(4/3)) - gamma(1/3)*hyper((1/3, 1/3), (4/3,), exp_polar(2*I*pi)/x**3)/(3*x*gamma(4/3))

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.35

$$\begin{aligned} \int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx &= \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(2(x^3-1)^{\frac{1}{3}} - 1\right)\right) \\ &\quad - \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(\frac{2(x^3-1)^{\frac{1}{3}}}{x} + 1\right)\right) \\ &\quad + \frac{1}{6} \log\left(\left(x^3-1\right)^{\frac{2}{3}} - \left(x^3-1\right)^{\frac{1}{3}} + 1\right) - \frac{1}{3} \log\left(\left(x^3-1\right)^{\frac{1}{3}} + 1\right) \\ &\quad + \frac{1}{6} \log\left(\frac{\left(x^3-1\right)^{\frac{1}{3}}}{x} + \frac{\left(x^3-1\right)^{\frac{2}{3}}}{x^2} + 1\right) - \frac{1}{3} \log\left(\frac{\left(x^3-1\right)^{\frac{1}{3}}}{x} - 1\right) \end{aligned}$$

[In] integrate((1+x)/x/(x^3-1)^(1/3),x, algorithm="maxima")

[Out] 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(x^3 - 1)^(1/3) - 1)) - 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(x^3 - 1)^(1/3)/x + 1)) + 1/6*log((x^3 - 1)^(2/3) - (x^3 - 1)^(1/3) + 1) - 1/3*log((x^3 - 1)^(1/3) + 1) + 1/6*log((x^3 - 1)^(1/3)/x + (x^3 - 1)^(2/3)/x^2 + 1) - 1/3*log((x^3 - 1)^(1/3)/x - 1)

Giac [F]

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx = \int \frac{x+1}{(x^3-1)^{\frac{1}{3}}x} dx$$

[In] integrate((1+x)/x/(x^3-1)^(1/3),x, algorithm="giac")

[Out] integrate((x + 1)/((x^3 - 1)^(1/3)*x), x)

Mupad [B] (verification not implemented)

Time = 0.46 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.09

$$\int \frac{1+x}{x\sqrt[3]{-1+x^3}} dx = \frac{x(1-x^3)^{1/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; x^3\right)}{(x^3-1)^{1/3}} - \ln\left(9\left(-\frac{1}{6} + \frac{\sqrt{3}li}{6}\right)^2 + (x^3-1)^{1/3}\right) \left(-\frac{1}{6} + \frac{\sqrt{3}li}{6}\right) + \ln\left(9\left(\frac{1}{6} + \frac{\sqrt{3}li}{6}\right)^2 + (x^3-1)^{1/3}\right) \left(\frac{1}{6} + \frac{\sqrt{3}li}{6}\right)$$

[In] int((x + 1)/(x*(x^3 - 1)^(1/3)),x)

[Out] log(9*((3^(1/2)*1i)/6 + 1/6)^2 + (x^3 - 1)^(1/3))*((3^(1/2)*1i)/6 + 1/6) - log(9*((3^(1/2)*1i)/6 - 1/6)^2 + (x^3 - 1)^(1/3))*((3^(1/2)*1i)/6 - 1/6) - log((x^3 - 1)^(1/3) + 1)/3 + (x*(1 - x^3)^(1/3)*hypergeom([1/3, 1/3], 4/3, x^3))/(x^3 - 1)^(1/3)

3.2

$$\int \frac{-1+x^2}{x(-1+x^3)^{2/3}} dx$$

Optimal result	38
Rubi [A] (verified)	38
Mathematica [A] (verified)	40
Maple [C] (warning: unable to verify)	41
Fricas [A] (verification not implemented)	41
Sympy [C] (verification not implemented)	42
Maxima [A] (verification not implemented)	42
Giac [F]	43
Mupad [F(-1)]	43

Optimal result

Integrand size = 18, antiderivative size = 95

$$\int \frac{-1+x^2}{x(-1+x^3)^{2/3}} dx = -\frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2}\log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{1}{2}\log\left(1 + \sqrt[3]{-1+x^3}\right)$$

[Out] $-1/3*\arctan(1/3*(1+2*x/(x^3-1)^{(1/3}))*3^{(1/2)})*3^{(1/2)}+1/3*\arctan(1/3*(1-2*(x^3-1)^{(1/3}))*3^{(1/2)})*3^{(1/2)}+1/2*\ln(x)-1/2*\ln(x-(x^3-1)^{(1/3}))-1/2*\ln(1+(x^3-1)^{(1/3}))$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {1858, 272, 60, 632, 210, 31, 337}

$$\int \frac{-1+x^2}{x(-1+x^3)^{2/3}} dx = -\frac{\arctan\left(\frac{\frac{2x}{\sqrt[3]{x^3-1}}+1}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1-2\sqrt[3]{x^3-1}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2}\log\left(x - \sqrt[3]{x^3-1}\right) - \frac{1}{2}\log\left(\sqrt[3]{x^3-1} + 1\right) + \frac{\log(x)}{2}$$

[In] $\text{Int}[(-1 + x^2)/(x*(-1 + x^3)^{(2/3)}),x]$

[Out] $-(\text{ArcTan}[(1 + (2*x)/(-1 + x^3)^{(1/3)})/\text{Sqrt}[3]]/\text{Sqrt}[3]) + \text{ArcTan}[(1 - 2*(-1 + x^3)^{(1/3)})/\text{Sqrt}[3]]/\text{Sqrt}[3] + \text{Log}[x]/2 - \text{Log}[x - (-1 + x^3)^{(1/3)}]/2 - \text{Log}[1 + (-1 + x^3)^{(1/3)}]/2$

Rule 31

$\text{Int}[(a_ + (b_)*(x_))^{(-1)}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] \text{ /; FreeQ}\{a, b\}, x]$

Rule 60

$\text{Int}[1/(((a_ + (b_)*(x_))*((c_ + (d_)*(x_))^{(2/3)})), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-(b*c - a*d)/b, 3]\}, \text{Simp}[-\text{Log}[\text{RemoveContent}[a + b*x, x]]/(2*b*q^2), x] + (\text{Dist}[3/(2*b*q), \text{Subst}[\text{Int}[1/(q^2 - q*x + x^2), x], x, (c + d*x)^{(1/3)}], x] + \text{Dist}[3/(2*b*q^2), \text{Subst}[\text{Int}[1/(q + x), x], x, (c + d*x)^{(1/3)}], x]) \text{ /; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NegQ}[(b*c - a*d)/b]$

Rule 210

$\text{Int}[(a_ + (b_)*(x_)^2)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)}*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \text{ /; FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])]$

Rule 272

$\text{Int}[(x_)^{(m_)*((a_ + (b_)*(x_)^{(n_))^{(p_)}), x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n] - 1)*(a + b*x)^p}, x], x, x^n], x] \text{ /; FreeQ}\{a, b, m, n, p\}, x \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]]$

Rule 337

$\text{Int}[(x_)/((a_ + (b_)*(x_)^3)^{(2/3)}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b, 3]\}, \text{Simp}[-\text{ArcTan}[(1 + 2*q*(x/(a + b*x^3)^{(1/3)}))/\text{Sqrt}[3]]/(\text{Sqrt}[3]*q^2), x] - \text{Simp}[\text{Log}[q*x - (a + b*x^3)^{(1/3)}]/(2*q^2), x]] \text{ /; FreeQ}\{a, b\}, x]$

Rule 632

$\text{Int}[(a_ + (b_)*(x_ + (c_)*(x_)^2)^{(-1)}, x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] \text{ /; FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 1858

$\text{Int}[(Pq_)*((c_)*(x_))^{(m_)*((a_ + (b_)*(x_)^{(n_))^{(p_)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*Pq*(a + b*x^n)^p, x], x] \text{ /; FreeQ}\{a, b, c, m, n$

, p}, x] && (PolyQ[Pq, x] || PolyQ[Pq, x^n]) && !IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(-\frac{1}{x(-1+x^3)^{2/3}} + \frac{x}{(-1+x^3)^{2/3}} \right) dx \\
&= -\int \frac{1}{x(-1+x^3)^{2/3}} dx + \int \frac{x}{(-1+x^3)^{2/3}} dx \\
&= -\frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{1}{3} \text{Subst}\left(\int \frac{1}{(-1+x)^{2/3}x} dx, x, x^3\right) \\
&= -\frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) \\
&\quad - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3}\right) - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3}\right) \\
&= -\frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) \\
&\quad - \frac{1}{2} \log\left(1 + \sqrt[3]{-1+x^3}\right) + \text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2\sqrt[3]{-1+x^3}\right) \\
&= -\frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{-1+2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{\sqrt{3}} \\
&\quad + \frac{\log(x)}{2} - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{1}{2} \log\left(1 + \sqrt[3]{-1+x^3}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.15 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.91

$$\begin{aligned}
\int \frac{-1+x^2}{x(-1+x^3)^{2/3}} dx &= \frac{1}{2} \left(-2\sqrt{3} \arctan\left(\frac{\sqrt{3}(-1+x)}{-1+x+2\sqrt[3]{-1+x^3}}\right) \right. \\
&\quad \left. - 2 \log\left(1-x+\sqrt[3]{-1+x^3}\right) + \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right) \right)
\end{aligned}$$

[In] Integrate[(-1 + x^2)/(x*(-1 + x^3)^(2/3)), x]


```
[Out] (-2*Sqrt[3]*ArcTan[(Sqrt[3]*(-1 + x))/(-1 + x + 2*(-1 + x^3)^(1/3))] - 2*Log[1 - x + (-1 + x^3)^(1/3)] + Log[1 - 2*x + x^2 + (-1 + x)*(-1 + x^3)^(1/3) + (-1 + x^3)^(2/3)])/2
```

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.67 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.04

method	result
meijerg	$\frac{(-\operatorname{signum}(x^3-1))^{\frac{2}{3}} x^2 \operatorname{hypergeom}\left(\left[\frac{2}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)}{2 \operatorname{signum}(x^3-1)^{\frac{2}{3}}} \frac{(-\operatorname{signum}(x^3-1))^{\frac{2}{3}} \left(\left(\frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 3\ln(x) + i\pi \right) \Gamma\left(\frac{2}{3}\right) + \frac{2\Gamma\left(\frac{2}{3}\right) x^3 \operatorname{hypergeom}\left(\left[\frac{2}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right)}{3} \right)}{3\Gamma\left(\frac{2}{3}\right) \operatorname{signum}(x^3-1)^{\frac{2}{3}}}$
trager	$-\ln\left(\frac{-58 \operatorname{RootOf}\left(_Z^2 - _Z + 1\right)^2 x^2 - 145 \operatorname{RootOf}\left(_Z^2 - _Z + 1\right)^2 x - 99 \operatorname{RootOf}\left(_Z^2 - _Z + 1\right) (x^3 - 1)^{\frac{2}{3}} - 99 \operatorname{RootOf}\left(_Z^2 - _Z + 1\right)}{\dots}\right)$

```
[In] int((x^2-1)/x/(x^3-1)^(2/3),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2/signum(x^3-1)^(2/3)*(-signum(x^3-1))^(2/3)*x^2*hypergeom([2/3,2/3],[5/3],x^3)-1/3/GAMMA(2/3)/signum(x^3-1)^(2/3)*(-signum(x^3-1))^(2/3)*((1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*GAMMA(2/3)+2/3*GAMMA(2/3)*x^3*hypergeom([1,1,5/3],[2,2],x^3))
```

Fricas [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.93

$$\int \frac{-1 + x^2}{x(-1 + x^3)^{2/3}} dx =$$

$$-\sqrt{3} \arctan\left(\frac{4\sqrt{3}(x^3-1)^{\frac{1}{3}}(x-1) + \sqrt{3}(x^2+x+1) - 2\sqrt{3}(x^3-1)^{\frac{2}{3}}}{3(3x^2-5x+3)}\right)$$

$$-\frac{1}{2} \log\left(\frac{(x^3-1)^{\frac{1}{3}}(x-1) + x - (x^3-1)^{\frac{2}{3}}}{x}\right)$$

```
[In] integrate((x^2-1)/x/(x^3-1)^(2/3),x, algorithm="fricas")
```

```
[Out] -sqrt(3)*arctan(-1/3*(4*sqrt(3)*(x^3-1)^(1/3)*(x-1) + sqrt(3)*(x^2+x+1) - 2*sqrt(3)*(x^3-1)^(2/3))/(3*x^2-5*x+3)) - 1/2*log(((x^3-1)^(1/3)*(x-1) + x - (x^3-1)^(2/3))/x)
```

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.10 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.68

$$\int \frac{-1 + x^2}{x(-1 + x^3)^{2/3}} dx = \frac{x^2 e^{-\frac{2i\pi}{3}} \Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\frac{2}{3}, \frac{2}{3} \middle| \frac{5}{3} \middle| x^3\right)}{3\Gamma\left(\frac{5}{3}\right)} + \frac{\Gamma\left(\frac{2}{3}\right) {}_2F_1\left(\frac{2}{3}, \frac{2}{3} \middle| \frac{5}{3} \middle| \frac{e^{2i\pi}}{x^3}\right)}{3x^2\Gamma\left(\frac{5}{3}\right)}$$

[In] integrate((x**2-1)/x/(x**3-1)**(2/3),x)

[Out] x**2*exp(-2*I*pi/3)*gamma(2/3)*hyper((2/3, 2/3), (5/3,), x**3)/(3*gamma(5/3)) + gamma(2/3)*hyper((2/3, 2/3), (5/3,), exp_polar(2*I*pi)/x**3)/(3*x**2*gamma(5/3))

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.31

$$\int \frac{-1 + x^2}{x(-1 + x^3)^{2/3}} dx = -\frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(2(x^3 - 1)^{\frac{1}{3}} - 1\right)\right) + \frac{1}{3} \sqrt{3} \arctan\left(\frac{1}{3} \sqrt{3} \left(\frac{2(x^3 - 1)^{\frac{1}{3}}}{x} + 1\right)\right) + \frac{1}{6} \log\left((x^3 - 1)^{\frac{2}{3}} - (x^3 - 1)^{\frac{1}{3}} + 1\right) - \frac{1}{3} \log\left((x^3 - 1)^{\frac{1}{3}} + 1\right) + \frac{1}{6} \log\left(\frac{(x^3 - 1)^{\frac{1}{3}}}{x} + \frac{(x^3 - 1)^{\frac{2}{3}}}{x^2} + 1\right) - \frac{1}{3} \log\left(\frac{(x^3 - 1)^{\frac{1}{3}}}{x} - 1\right)$$

[In] integrate((x^2-1)/x/(x^3-1)^(2/3),x, algorithm="maxima")

[Out] -1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(x^3 - 1)^(1/3) - 1)) + 1/3*sqrt(3)*arctan(1/3*sqrt(3)*(2*(x^3 - 1)^(1/3)/x + 1)) + 1/6*log((x^3 - 1)^(2/3) - (x^3 - 1)^(1/3) + 1) - 1/3*log((x^3 - 1)^(1/3) + 1) + 1/6*log((x^3 - 1)^(1/3)/x + (x^3 - 1)^(2/3)/x^2 + 1) - 1/3*log((x^3 - 1)^(1/3)/x - 1)

Giac [F]

$$\int \frac{-1 + x^2}{x(-1 + x^3)^{2/3}} dx = \int \frac{x^2 - 1}{(x^3 - 1)^{2/3} x} dx$$

[In] integrate((x^2-1)/x/(x^3-1)^(2/3),x, algorithm="giac")

[Out] integrate((x^2 - 1)/((x^3 - 1)^(2/3)*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{-1 + x^2}{x(-1 + x^3)^{2/3}} dx = \int \frac{x^2 - 1}{x(x^3 - 1)^{2/3}} dx$$

[In] int((x^2 - 1)/(x*(x^3 - 1)^(2/3)),x)

[Out] int((x^2 - 1)/(x*(x^3 - 1)^(2/3)), x)

3.3 $\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx$

Optimal result	44
Rubi [C] (verified)	45
Mathematica [A] (verified)	49
Maple [C] (warning: unable to verify)	50
Fricas [A] (verification not implemented)	50
Sympy [F]	51
Maxima [F]	51
Giac [F]	51
Mupad [F(-1)]	51

Optimal result

Integrand size = 21, antiderivative size = 107

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx = -\frac{3\sqrt[3]{-1+x^3}}{-1+x} + \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) - \log\left(1-x+\sqrt[3]{-1+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)$$

[Out] -3*(x^3-1)^(1/3)/(-1+x)+3^(1/2)*arctan(3^(1/2)*(x^3-1)^(1/3)/(-2+2*x+(x^3-1)^(1/3)))-ln(1-x+(x^3-1)^(1/3))+1/2*ln(1-2*x+x^2+(-1+x)*(x^3-1)^(1/3)+(x^3-1)^(2/3))

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.13 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.83, number of steps used = 16, number of rules used = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {2180, 252, 251, 272, 53, 60, 632, 210, 31, 270, 267, 372, 371, 294, 337}

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx$$

$$= -\frac{\arctan\left(\frac{\sqrt[3]{x^3-1}+1}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{1-2\sqrt[3]{x^3-1}}{\sqrt{3}}\right)}{\sqrt{3}}$$

$$- \frac{3(1-x^3)^{2/3} x \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{(x^3-1)^{2/3}}$$

$$- \frac{3(1-x^3)^{2/3} x^4 \operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{4(x^3-1)^{2/3}} - \frac{3}{(x^3-1)^{2/3}}$$

$$- \frac{1}{2} \log\left(x - \sqrt[3]{x^3-1}\right) - \frac{1}{2} \log\left(\sqrt[3]{x^3-1} + 1\right) - \frac{3x^2}{(x^3-1)^{2/3}} + \frac{\log(x)}{2}$$

[In] Int[((1 + x)*(-1 + x^3)^(1/3))/((-1 + x)^2*x), x]

[Out] -3/(-1 + x^3)^(2/3) - (3*x^2)/(-1 + x^3)^(2/3) - ArcTan[(1 + (2*x)/(-1 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + ArcTan[(1 - 2*(-1 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - (3*x*(1 - x^3)^(2/3)*Hypergeometric2F1[1/3, 5/3, 4/3, x^3])/(-1 + x^3)^(2/3) - (3*x^4*(1 - x^3)^(2/3)*Hypergeometric2F1[4/3, 5/3, 7/3, x^3])/(4*(-1 + x^3)^(2/3)) + Log[x]/2 - Log[x - (-1 + x^3)^(1/3)]/2 - Log[1 + (-1 + x^3)^(1/3)]/2

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 53

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^(n + 1)/((b*c - a*d)*(m + 1)), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]

Rule 60

```
Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[
{q = Rt[-(b*c - a*d)/b, 3]}, Simp[-Log[RemoveContent[a + b*x, x]]/(2*b*q^2)
, x] + (Dist[3/(2*b*q), Subst[Int[1/(q^2 - q*x + x^2), x], x, (c + d*x)^(1/3)], x] + Dist[3/(2*b*q^2), Subst[Int[1/(q + x), x], x, (c + d*x)^(1/3)], x
])] /; FreeQ[{a, b, c, d}, x] && NegQ[(b*c - a*d)/b]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F
1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p
, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] ||
GtQ[a, 0])
```

Rule 252

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^IntPart[p]*((a + b*x
^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]), Int[(1 + b*(x^n/a))^p, x], x]
/; FreeQ[{a, b, n, p}, x] && !IGtQ[p, 0] && !IntegerQ[1/n] && !ILtQ[Sim
plify[1/n + p], 0] && !(IntegerQ[p] || GtQ[a, 0])
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)
^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n,
p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
```

, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 337

Int[(x_)/((a_) + (b_.)*(x_)^3)^(2/3), x_Symbol] :> With[{q = Rt[b, 3]}, Simp[-ArcTan[(1 + 2*q*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*q^2), x] - Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*q^2), x]] /; FreeQ[{a, b}, x]

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 372

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]), Int[(c*x)^(m*(1 + b*(x^n/a))^p], x], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && !(ILtQ[p, 0] || GtQ[a, 0])

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 2180

Int[(Px_)*(x_))^(m_.)*((c_) + (d_.)*(x_))^(q_)*((a_) + (b_.)*(x_)^3)^(p_.), x_Symbol] :> Int[ExpandIntegrand[(c^3 + d^3*x^3)^q*(a + b*x^3)^p, x^m*(Px/(c^2 - c*d*x + d^2*x^2)^q), x], x] /; FreeQ[{a, b, c, d, m, p}, x] && PolyQ[Px, x] && ILtQ[q, 0] && IntegerQ[m] && RationalQ[p] && EqQ[Denominator[p], 3]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{3}{(-1+x^3)^{5/3}} + \frac{1}{x(-1+x^3)^{5/3}} + \frac{5x}{(-1+x^3)^{5/3}} + \frac{5x^2}{(-1+x^3)^{5/3}} \right. \\
&\quad \left. + \frac{3x^3}{(-1+x^3)^{5/3}} + \frac{x^4}{(-1+x^3)^{5/3}} \right) dx \\
&= 3 \int \frac{1}{(-1+x^3)^{5/3}} dx + 3 \int \frac{x^3}{(-1+x^3)^{5/3}} dx + 5 \int \frac{x}{(-1+x^3)^{5/3}} dx \\
&\quad + 5 \int \frac{x^2}{(-1+x^3)^{5/3}} dx + \int \frac{1}{x(-1+x^3)^{5/3}} dx + \int \frac{x^4}{(-1+x^3)^{5/3}} dx \\
&= -\frac{5}{2(-1+x^3)^{2/3}} - \frac{3x^2}{(-1+x^3)^{2/3}} \\
&\quad + \frac{1}{3} \text{Subst} \left(\int \frac{1}{(-1+x)^{5/3} x} dx, x, x^3 \right) - \frac{(3(1-x^3)^{2/3}) \int \frac{1}{(1-x^3)^{5/3}} dx}{(-1+x^3)^{2/3}} - \frac{(3(1-x^3)^{2/3}) \int \frac{x^3}{(1-x^3)^{5/3}} dx}{(-1+x^3)^{2/3}} \\
&= -\frac{3}{(-1+x^3)^{2/3}} - \frac{3x^2}{(-1+x^3)^{2/3}} - \frac{\arctan \left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} \\
&\quad - \frac{3x(1-x^3)^{2/3} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3 \right)}{(-1+x^3)^{2/3}} \\
&\quad - \frac{3x^4(1-x^3)^{2/3} \text{Hypergeometric2F1} \left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3 \right)}{4(-1+x^3)^{2/3}} \\
&\quad - \frac{1}{2} \log \left(x - \sqrt[3]{-1+x^3} \right) - \frac{1}{3} \text{Subst} \left(\int \frac{1}{(-1+x)^{2/3} x} dx, x, x^3 \right) \\
&= -\frac{3}{(-1+x^3)^{2/3}} - \frac{3x^2}{(-1+x^3)^{2/3}} - \frac{\arctan \left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} \\
&\quad - \frac{3x(1-x^3)^{2/3} \text{Hypergeometric2F1} \left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3 \right)}{(-1+x^3)^{2/3}} \\
&\quad - \frac{3x^4(1-x^3)^{2/3} \text{Hypergeometric2F1} \left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3 \right)}{4(-1+x^3)^{2/3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2} \log \left(x - \sqrt[3]{-1+x^3} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3} \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3}{(-1+x^3)^{2/3}} - \frac{3x^2}{(-1+x^3)^{2/3}} - \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} \\
&\quad - \frac{3x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{(-1+x^3)^{2/3}} \\
&\quad - \frac{3x^4(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{4(-1+x^3)^{2/3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{1}{2} \log\left(1 + \sqrt[3]{-1+x^3}\right) + \operatorname{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2\sqrt[3]{-1+x^3}\right) \\
&= -\frac{3}{(-1+x^3)^{2/3}} - \frac{3x^2}{(-1+x^3)^{2/3}} - \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} \\
&\quad - \frac{\arctan\left(\frac{-1+2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{3x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{(-1+x^3)^{2/3}} \\
&\quad - \frac{3x^4(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{4(-1+x^3)^{2/3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2} \log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{1}{2} \log\left(1 + \sqrt[3]{-1+x^3}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.66 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2 x} dx &= -\frac{3\sqrt[3]{-1+x^3}}{-1+x} + \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) \\
&\quad - \log\left(1-x+\sqrt[3]{-1+x^3}\right) \\
&\quad + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)
\end{aligned}$$

[In] Integrate[((1+x)*(-1+x^3)^(1/3))/((-1+x)^2*x),x]

[Out] (-3*(-1+x^3)^(1/3))/(-1+x) + Sqrt[3]*ArcTan[(Sqrt[3]*(-1+x^3)^(1/3))/(-2+2*x+(-1+x^3)^(1/3))] - Log[1-x+(-1+x^3)^(1/3)] + Log[1-2*x+x^2+(-1+x)*(-1+x^3)^(1/3)+(-1+x^3)^(2/3)]/2

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.79 (sec) , antiderivative size = 164, normalized size of antiderivative = 1.53

method	result
risch	$-\frac{3(x^2+x+1)}{(x^3-1)^{\frac{2}{3}}} + \frac{\left(\frac{(x^3-1)^{\frac{2}{3}} (-\text{signum}(x^3-1))^{\frac{2}{3}} x^2 \text{hypergeom}\left(\left[\frac{2}{3}, \frac{2}{3}\right], \left[\frac{5}{3}\right], x^3\right) - (x^3-1)^{\frac{2}{3}} (-\text{signum}(x^3-1))^{\frac{2}{3}} \left(\left(\frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 3\ln(x) + I\pi\right) \Gamma\left(\frac{2}{3}\right)\right)}{2\left((x^3-1)^2\right)^{\frac{1}{3}} \text{signum}(x^3-1)^{\frac{2}{3}}}\right)}{(x^3-1)^{\frac{2}{3}}}$
trager	$-\frac{3(x^3-1)^{\frac{1}{3}}}{-1+x} - \ln\left(\frac{12\text{RootOf}\left(_Z^2 - _Z+1\right)^2 x^2 - 30\text{RootOf}\left(_Z^2 - _Z+1\right)^2 x - 99\text{RootOf}\left(_Z^2 - _Z+1\right)(x^3-1)^{\frac{2}{3}} + 127}{(x^3-1)^{\frac{2}{3}}}\right)$

[In] int((1+x)*(x^3-1)^(1/3)/(-1+x)^2/x,x,method=_RETURNVERBOSE)

[Out] -3*(x^2+x+1)/(x^3-1)^(2/3)+(1/2/((x^3-1)^2)^(1/3)*(x^3-1)^(2/3)/signum(x^3-1)^(2/3)*(-signum(x^3-1))^(2/3)*x^2*hypergeom([2/3,2/3],[5/3],x^3)-1/3/((x^3-1)^2)^(1/3)*(x^3-1)^(2/3)/GAMMA(2/3)/signum(x^3-1)^(2/3)*(-signum(x^3-1))^(2/3)*((1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*GAMMA(2/3)+2/3*GAMMA(2/3)*x^3*hypergeom([1,1,5/3],[2,2],x^3)))/(x^3-1)^(2/3)*((x^3-1)^2)^(1/3)

Fricas [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.02

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2 x} dx = \frac{2\sqrt{3}(x-1) \arctan\left(-\frac{4\sqrt{3}(x^3-1)^{\frac{1}{3}}(x-1)+\sqrt{3}(x^2+x+1)-2\sqrt{3}(x^3-1)^{\frac{2}{3}}}{3(3x^2-5x+3)}\right) + (x-1) \log\left(\frac{(x^3-1)^{\frac{1}{3}}(x-1)+x-(x^3-1)^{\frac{2}{3}}}{x}\right)}{2(x-1)}$$

[In] integrate((1+x)*(x^3-1)^(1/3)/(-1+x)^2/x,x, algorithm="fricas")

[Out] -1/2*(2*sqrt(3)*(x-1)*arctan(-1/3*(4*sqrt(3)*(x^3-1)^(1/3)*(x-1)+sqrt(3)*(x^2+x+1)-2*sqrt(3)*(x^3-1)^(2/3))/(3*x^2-5*x+3))+ (x-1)*log(((x^3-1)^(1/3)*(x-1)+x-(x^3-1)^(2/3))/x)+6*(x^3-1)^(1/3)/(x-1)

Sympy [F]

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx = \int \frac{\sqrt[3]{(x-1)(x^2+x+1)}(x+1)}{x(x-1)^2} dx$$

[In] integrate((1+x)*(x**3-1)**(1/3)/(-1+x)**2/x,x)

[Out] Integral(((x - 1)*(x**2 + x + 1))**(1/3)*(x + 1)/(x*(x - 1)**2), x)

Maxima [F]

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx = \int \frac{(x^3-1)^{\frac{1}{3}}(x+1)}{(x-1)^2x} dx$$

[In] integrate((1+x)*(x^3-1)^(1/3)/(-1+x)^2/x,x, algorithm="maxima")

[Out] integrate((x^3 - 1)^(1/3)*(x + 1)/((x - 1)^2*x), x)

Giac [F]

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx = \int \frac{(x^3-1)^{\frac{1}{3}}(x+1)}{(x-1)^2x} dx$$

[In] integrate((1+x)*(x^3-1)^(1/3)/(-1+x)^2/x,x, algorithm="giac")

[Out] integrate((x^3 - 1)^(1/3)*(x + 1)/((x - 1)^2*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(1+x)\sqrt[3]{-1+x^3}}{(-1+x)^2x} dx = \int \frac{(x^3-1)^{1/3}(x+1)}{x(x-1)^2} dx$$

[In] int(((x^3 - 1)^(1/3)*(x + 1))/(x*(x - 1)^2), x)

[Out] int(((x^3 - 1)^(1/3)*(x + 1))/(x*(x - 1)^2), x)

$$3.4 \quad \int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx$$

Optimal result	52
Rubi [A] (verified)	52
Mathematica [A] (verified)	56
Maple [C] (warning: unable to verify)	56
Fricas [A] (verification not implemented)	57
Sympy [F]	57
Maxima [F]	57
Giac [F]	58
Mupad [F(-1)]	58

Optimal result

Integrand size = 29, antiderivative size = 111

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = -\frac{3(-1+x^3)^{2/3}}{1+x+x^2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) - \log\left(1-x+\sqrt[3]{-1+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)$$

[Out] $-3*(x^3-1)^{(2/3)}/(x^2+x+1)-3^{(1/2)}*\arctan(3^{(1/2)}*(x^3-1)^{(1/3)}/(-2+2*x+(x^3-1)^{(1/3)}))- \ln(1-x+(x^3-1)^{(1/3)})+1/2*\ln(1-2*x+x^2+(-1+x)*(x^3-1)^{(1/3)}+(x^3-1)^{(2/3)})$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.06, number of steps used = 16, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.414$, Rules used = {2182, 21, 197, 272, 53, 58, 632, 210, 31, 267, 294, 245}

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = \frac{\arctan\left(\frac{\sqrt[3]{x^3-1}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1-2\sqrt[3]{x^3-1}}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{3x}{\sqrt[3]{x^3-1}} + \frac{3}{\sqrt[3]{x^3-1}} - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}+1\right) - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}-x\right) + \frac{\log(x)}{2}$$

```
[In] Int[((-1 + x)^2*(1 + x))/(x*(1 + x + x^2)*(-1 + x^3)^(1/3)),x]
[Out] 3/(-1 + x^3)^(1/3) - (3*x)/(-1 + x^3)^(1/3) + ArcTan[(1 + (2*x)/(-1 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] - ArcTan[(1 - 2*(-1 + x^3)^(1/3))/Sqrt[3]]/Sqrt[3] + Log[x]/2 - Log[1 + (-1 + x^3)^(1/3)]/2 - Log[-x + (-1 + x^3)^(1/3)]/2
```

Rule 21

```
Int[(u_.)*((a_) + (b_.)*(v_))^(m_.)*((c_) + (d_.)*(v_))^(n_.), x_Symbol] :=
  Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x]
  && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x,
  a + b*x])
```

Rule 31

```
Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x,
x]]/b, x] /; FreeQ[{a, b}, x]
```

Rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 58

```
Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[
{q = Rt[-(b*c - a*d)/b, 3]}, Simp[Log[RemoveContent[a + b*x, x]]/(2*b*q), x
] + (Dist[3/(2*b), Subst[Int[1/(q^2 - q*x + x^2), x], x, (c + d*x)^(1/3)],
x] - Dist[3/(2*b*q), Subst[Int[1/(q + x), x], x, (c + d*x)^(1/3)], x])] /;
FreeQ[{a, b, c, d}, x] && NegQ[(b*c - a*d)/b]
```

Rule 197

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)
/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 245

```
Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*
(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^
3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]
```

Rule 267

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)
^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] &&
NeQ[p, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 294

```
Int[((c_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2182

```
Int[(Px_)*(x_)^(m_.)*((c_) + (d_.)*(x_) + (e_.)*(x_)^2)^(q_)*((a_) + (b_.)*
(x_)^3)^(p_.), x_Symbol] := Dist[1/c^q, Int[ExpandIntegrand[(c^3 - d^3*x^3)
^q*(a + b*x^3)^p, x^m*(Px/(c - d*x)^q), x], x], x] /; FreeQ[{a, b, c, d, e,
m, p}, x] && PolyQ[Px, x] && EqQ[d^2 - c*e, 0] && ILtQ[q, 0] && IntegerQ[m
] && RationalQ[p] && EqQ[Denominator[p], 3]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(-\frac{2}{(1-x^3)\sqrt[3]{-1+x^3}} + \frac{1}{x(1-x^3)\sqrt[3]{-1+x^3}} + \frac{2x^2}{(1-x^3)\sqrt[3]{-1+x^3}} \right. \\
&\quad \left. - \frac{x^3}{(1-x^3)\sqrt[3]{-1+x^3}} \right) dx \\
&= -\left(2 \int \frac{1}{(1-x^3)\sqrt[3]{-1+x^3}} dx \right) + 2 \int \frac{x^2}{(1-x^3)\sqrt[3]{-1+x^3}} dx \\
&\quad + \int \frac{1}{x(1-x^3)\sqrt[3]{-1+x^3}} dx - \int \frac{x^3}{(1-x^3)\sqrt[3]{-1+x^3}} dx \\
&= 2 \int \frac{1}{(-1+x^3)^{4/3}} dx - 2 \int \frac{x^2}{(-1+x^3)^{4/3}} dx - \int \frac{1}{x(-1+x^3)^{4/3}} dx + \int \frac{x^3}{(-1+x^3)^{4/3}} dx \\
&= \frac{2}{\sqrt[3]{-1+x^3}} - \frac{3x}{\sqrt[3]{-1+x^3}} - \frac{1}{3} \text{Subst} \left(\int \frac{1}{(-1+x)^{4/3}x} dx, x, x^3 \right) + \int \frac{1}{\sqrt[3]{-1+x^3}} dx \\
&= \frac{3}{\sqrt[3]{-1+x^3}} - \frac{3x}{\sqrt[3]{-1+x^3}} + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} \\
&\quad - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right) + \frac{1}{3} \text{Subst} \left(\int \frac{1}{\sqrt[3]{-1+xx}} dx, x, x^3 \right) \\
&= \frac{3}{\sqrt[3]{-1+x^3}} - \frac{3x}{\sqrt[3]{-1+x^3}} + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right) \\
&\quad - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3} \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3} \right) \\
&= \frac{3}{\sqrt[3]{-1+x^3}} - \frac{3x}{\sqrt[3]{-1+x^3}} + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{\log(x)}{2} - \frac{1}{2} \log \left(1 + \sqrt[3]{-1+x^3} \right) \\
&\quad - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right) - \text{Subst} \left(\int \frac{1}{-3-x^2} dx, x, -1 + 2\sqrt[3]{-1+x^3} \right) \\
&= \frac{3}{\sqrt[3]{-1+x^3}} - \frac{3x}{\sqrt[3]{-1+x^3}} + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{\arctan \left(\frac{-1+2\sqrt[3]{-1+x^3}}{\sqrt{3}} \right)}{\sqrt{3}} \\
&\quad + \frac{\log(x)}{2} - \frac{1}{2} \log \left(1 + \sqrt[3]{-1+x^3} \right) - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.00

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = -\frac{3(-1+x^3)^{2/3}}{1+x+x^2} - \sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x+\sqrt[3]{-1+x^3}}\right) - \log\left(1-x+\sqrt[3]{-1+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)$$

[In] Integrate[((-1 + x)^2*(1 + x))/(x*(1 + x + x^2)*(-1 + x^3)^(1/3)), x]

[Out] (-3*(-1 + x^3)^(2/3))/(1 + x + x^2) - Sqrt[3]*ArcTan[(Sqrt[3]*(-1 + x^3)^(1/3))/(-2 + 2*x + (-1 + x^3)^(1/3))] - Log[1 - x + (-1 + x^3)^(1/3)] + Log[1 - 2*x + x^2 + (-1 + x)*(-1 + x^3)^(1/3) + (-1 + x^3)^(2/3)]/2

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.94 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.13

method	result
risch	$-\frac{3(-1+x)}{(x^3-1)^{\frac{1}{3}}} + \frac{\sqrt{3}\Gamma(\frac{2}{3})(-\text{signum}(x^3-1))^{\frac{1}{3}} \left(\frac{2\left(-\frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 3\ln(x) + i\pi\right)\pi\sqrt{3}}{3\Gamma(\frac{2}{3})} + \frac{2\pi\sqrt{3}x^3 \text{hypergeom}\left(\left[1, 1, \frac{4}{3}\right], [2, 2], x^3\right)}{9\Gamma(\frac{2}{3})}\right)}{6\pi \text{signum}(x^3-1)^{\frac{1}{3}}} + \frac{(-\text{signu}}{2}$
trager	Expression too large to display

[In] int((-1+x)^2*(1+x)/x/(x^2+x+1)/(x^3-1)^(1/3), x, method=_RETURNVERBOSE)

[Out] -3*(-1+x)/(x^3-1)^(1/3)+1/6/Pi*3^(1/2)*GAMMA(2/3)/signum(x^3-1)^(1/3)*(-signum(x^3-1))^(1/3)*(2/3*(-1/6*Pi*3^(1/2)-3/2*ln(3)+3*ln(x)+I*Pi)*Pi*3^(1/2)/GAMMA(2/3)+2/9*Pi*3^(1/2)/GAMMA(2/3)*x^3*hypergeom([1, 1, 4/3], [2, 2], x^3))+1/signum(x^3-1)^(1/3)*(-signum(x^3-1))^(1/3)*x*hypergeom([1/3, 1/3], [4/3], x^3)

Fricas [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.07

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx$$

$$= \frac{2\sqrt{3}(x^2+x+1)\arctan\left(-\frac{4\sqrt{3}(x^3-1)^{\frac{1}{3}}(x-1)+\sqrt{3}(x^2+x+1)-2\sqrt{3}(x^3-1)^{\frac{2}{3}}}{3(3x^2-5x+3)}\right) - (x^2+x+1)\log\left(\frac{(x^3-1)^{\frac{1}{3}}(x-1)+x-1}{x}\right)}{2(x^2+x+1)}$$

[In] integrate((-1+x)^2*(1+x)/x/(x^2+x+1)/(x^3-1)^(1/3),x, algorithm="fricas")

[Out] 1/2*(2*sqrt(3)*(x^2 + x + 1)*arctan(-1/3*(4*sqrt(3)*(x^3 - 1)^(1/3)*(x - 1) + sqrt(3)*(x^2 + x + 1) - 2*sqrt(3)*(x^3 - 1)^(2/3))/(3*x^2 - 5*x + 3)) - (x^2 + x + 1)*log(((x^3 - 1)^(1/3)*(x - 1) + x - (x^3 - 1)^(2/3))/x) - 6*(x^3 - 1)^(2/3))/(x^2 + x + 1)

Sympy [F]

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = \int \frac{(x-1)^2(x+1)}{x\sqrt[3]{(x-1)(x^2+x+1)}(x^2+x+1)} dx$$

[In] integrate((-1+x)**2*(1+x)/x/(x**2+x+1)/(x**3-1)**(1/3),x)

[Out] Integral((x - 1)**2*(x + 1)/(x*((x - 1)*(x**2 + x + 1))**(1/3)*(x**2 + x + 1)), x)

Maxima [F]

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = \int \frac{(x+1)(x-1)^2}{(x^3-1)^{\frac{1}{3}}(x^2+x+1)x} dx$$

[In] integrate((-1+x)^2*(1+x)/x/(x^2+x+1)/(x^3-1)^(1/3),x, algorithm="maxima")

[Out] integrate((x + 1)*(x - 1)^2/((x^3 - 1)^(1/3)*(x^2 + x + 1)*x), x)

Giac [F]

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = \int \frac{(x+1)(x-1)^2}{(x^3-1)^{\frac{1}{3}}(x^2+x+1)x} dx$$

[In] integrate((-1+x)^2*(1+x)/x/(x^2+x+1)/(x^3-1)^(1/3),x, algorithm="giac")

[Out] integrate((x + 1)*(x - 1)^2/((x^3 - 1)^(1/3)*(x^2 + x + 1)*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(-1+x)^2(1+x)}{x(1+x+x^2)\sqrt[3]{-1+x^3}} dx = \int \frac{(x-1)^2(x+1)}{x(x^3-1)^{1/3}(x^2+x+1)} dx$$

[In] int(((x - 1)^2*(x + 1))/(x*(x^3 - 1)^(1/3)*(x + x^2 + 1)),x)

[Out] int(((x - 1)^2*(x + 1))/(x*(x^3 - 1)^(1/3)*(x + x^2 + 1)), x)

$$3.5 \quad \int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx$$

Optimal result	59
Rubi [C] (verified)	59
Mathematica [C] (verified)	64
Maple [C] (warning: unable to verify)	65
Fricas [B] (verification not implemented)	65
Sympy [F]	66
Maxima [F]	66
Giac [F]	66
Mupad [F(-1)]	66

Optimal result

Integrand size = 21, antiderivative size = 147

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \frac{-3(-1+x^3)^{2/3} - 2\sqrt{3}(1-2x+x^2) \arctan\left(\frac{\sqrt{3}(1+x+x^2)+4\sqrt{3}(-1+x)\sqrt[3]{-1+x^3}}{3(3-5x+3x^2)}\right)}{2(1-2x+x^2)}$$

[Out] 1/2*(-2*3^(1/2)*(x^2-2*x+1)*arctan(1/3*(4*3^(1/2)*(x^3-1)^(1/3)*(-1+x)+3^(1/2)*(x^2+x+1)-2*3^(1/2)*(x^3-1)^(2/3))/(3*x^2-5*x+3))-(x^2-2*x+1)*ln(((1+x)*(x^3-1)^(1/3)+x-(x^3-1)^(2/3))/x)-3*(x^3-1)^(2/3))/(x^2-2*x+1)

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.15 (sec) , antiderivative size = 239, normalized size of antiderivative = 1.63, number of steps used = 23, number of rules used = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.762$, Rules used = {2180, 198, 197, 272, 53, 58, 632, 210, 31, 372, 371, 267, 270, 45, 294, 245}

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \frac{\arctan\left(\frac{\sqrt[3]{x^3-1}+1}{\sqrt{3}}\right)}{\sqrt{3}} - \frac{\arctan\left(\frac{1-2\sqrt[3]{x^3-1}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{9\sqrt[3]{1-x^3}x^5 \text{Hypergeometric2F1}\left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3\right)}{5\sqrt[3]{x^3-1}} + \frac{9\sqrt[3]{1-x^3}x^2 \text{Hypergeometric2F1}\left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3\right)}{2\sqrt[3]{x^3-1}} + \frac{2x}{\sqrt[3]{x^3-1}} - \frac{x}{(x^3-1)^{4/3}} - \frac{3}{\sqrt[3]{x^3-1}} - \frac{9}{2(x^3-1)^{4/3}} - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}+1\right) - \frac{1}{2} \log\left(\sqrt[3]{x^3-1}-x\right) - \frac{7x^4}{2(x^3-1)^{4/3}} + \frac{\log(x)}{2}$$

[In] Int[((1 + x)*(-1 + x^3)^(2/3))/((-1 + x)^3*x),x]

[Out]
$$-9/(2*(-1 + x^3)^{4/3}) - x/(-1 + x^3)^{4/3} - (7*x^4)/(2*(-1 + x^3)^{4/3}) - 3/(-1 + x^3)^{1/3} + (2*x)/(-1 + x^3)^{1/3} + \text{ArcTan}[(1 + (2*x)/(-1 + x^3)^{1/3})/\text{Sqrt}[3]]/\text{Sqrt}[3] - \text{ArcTan}[(1 - 2*(-1 + x^3)^{1/3})/\text{Sqrt}[3]]/\text{Sqrt}[3] + (9*x^2*(1 - x^3)^{1/3}*\text{Hypergeometric2F1}[2/3, 7/3, 5/3, x^3])/(2*(-1 + x^3)^{1/3}) + (9*x^5*(1 - x^3)^{1/3}*\text{Hypergeometric2F1}[5/3, 7/3, 8/3, x^3])/(5*(-1 + x^3)^{1/3}) + \text{Log}[x]/2 - \text{Log}[1 + (-1 + x^3)^{1/3}]/2 - \text{Log}[-x + (-1 + x^3)^{1/3}]/2$$

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LtQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 53

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && IntLinearQ[a, b, c, d, m, n, x]

Rule 58

Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(1/3)), x_Symbol] := With[{q = Rt[-(b*c - a*d)/b, 3]}, Simp[Log[RemoveContent[a + b*x, x]]/(2*b*q), x] + (Dist[3/(2*b), Subst[Int[1/(q^2 - q*x + x^2), x], x, (c + d*x)^(1/3)], x] - Dist[3/(2*b*q), Subst[Int[1/(q + x), x], x, (c + d*x)^(1/3)], x])] /; FreeQ[{a, b, c, d}, x] && NegQ[(b*c - a*d)/b]

Rule 197

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 198

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-x)*((a + b*x^n)^(p + 1)/(a*n*(p + 1))), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 245

Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]

Rule 267

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(a + b*x^n)^(p + 1)/(b*n*(p + 1)), x] /; FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]

Rule 270

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 371

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p
*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1
, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt
Q[p, 0] || GtQ[a, 0])
```

Rule 372

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^I
ntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]), Int[(c*x)^
m*(1 + b*(x^n/a))^p, x], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0]
&& !(ILtQ[p, 0] || GtQ[a, 0])
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2180

```
Int[(Px_)*(x_)^m_.*((c_) + (d_.)*(x_)^(q_))*((a_) + (b_.)*(x_)^3)^(p_.),
x_Symbol] := Int[ExpandIntegrand[(c^3 + d^3*x^3)^q*(a + b*x^3)^p, x^m*(Px/(
c^2 - c*d*x + d^2*x^2)^q), x], x] /; FreeQ[{a, b, c, d, m, p}, x] && PolyQ[
Px, x] && ILtQ[q, 0] && IntegerQ[m] && RationalQ[p] && EqQ[Denominator[p],
3]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{4}{(-1+x^3)^{7/3}} + \frac{1}{x(-1+x^3)^{7/3}} + \frac{9x}{(-1+x^3)^{7/3}} + \frac{13x^2}{(-1+x^3)^{7/3}} + \frac{13x^3}{(-1+x^3)^{7/3}} \right. \\ &\quad \left. + \frac{9x^4}{(-1+x^3)^{7/3}} + \frac{4x^5}{(-1+x^3)^{7/3}} + \frac{x^6}{(-1+x^3)^{7/3}} \right) dx \\ &= 4 \int \frac{1}{(-1+x^3)^{7/3}} dx + 4 \int \frac{x^5}{(-1+x^3)^{7/3}} dx + 9 \int \frac{x}{(-1+x^3)^{7/3}} dx \\ &\quad + 9 \int \frac{x^4}{(-1+x^3)^{7/3}} dx + 13 \int \frac{x^2}{(-1+x^3)^{7/3}} dx \\ &\quad + 13 \int \frac{x^3}{(-1+x^3)^{7/3}} dx + \int \frac{1}{x(-1+x^3)^{7/3}} dx + \int \frac{x^6}{(-1+x^3)^{7/3}} dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{13}{4(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} \\
&\quad + \frac{1}{3} \text{Subst} \left(\int \frac{1}{(-1+x)^{7/3}x} dx, x, x^3 \right) + \frac{4}{3} \text{Subst} \left(\int \frac{x}{(-1+x)^{7/3}} dx, x, x^3 \right) - 3 \int \frac{1}{(-1+x^3)^{4/3}} dx \\
&= -\frac{7}{2(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} \\
&\quad + \frac{2x}{\sqrt[3]{-1+x^3}} + \frac{9x^2 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3 \right)}{2\sqrt[3]{-1+x^3}} \\
&\quad + \frac{9x^5 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3 \right)}{5\sqrt[3]{-1+x^3}} \\
&\quad - \frac{1}{3} \text{Subst} \left(\int \frac{1}{(-1+x)^{4/3}x} dx, x, x^3 \right) + \frac{4}{3} \text{Subst} \left(\int \left(\frac{1}{(-1+x)^{7/3}} + \frac{1}{(-1+x)^{4/3}} \right) dx, x, x^3 \right) + \int \frac{1}{(-1+x^3)^{4/3}} dx \\
&= -\frac{9}{2(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} - \frac{3}{\sqrt[3]{-1+x^3}} + \frac{2x}{\sqrt[3]{-1+x^3}} \\
&\quad + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{9x^2 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3 \right)}{2\sqrt[3]{-1+x^3}} \\
&\quad + \frac{9x^5 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3 \right)}{5\sqrt[3]{-1+x^3}} \\
&\quad - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right) + \frac{1}{3} \text{Subst} \left(\int \frac{1}{\sqrt[3]{-1+xx}} dx, x, x^3 \right) \\
&= -\frac{9}{2(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} - \frac{3}{\sqrt[3]{-1+x^3}} + \frac{2x}{\sqrt[3]{-1+x^3}} \\
&\quad + \frac{\arctan \left(\frac{1 + \frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}} \right)}{\sqrt{3}} + \frac{9x^2 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3 \right)}{2\sqrt[3]{-1+x^3}} \\
&\quad + \frac{9x^5 \sqrt[3]{1-x^3} \text{Hypergeometric2F1} \left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3 \right)}{5\sqrt[3]{-1+x^3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2} \log \left(-x + \sqrt[3]{-1+x^3} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3} \right) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3} \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{9}{2(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} - \frac{3}{\sqrt[3]{-1+x^3}} + \frac{2x}{\sqrt[3]{-1+x^3}} \\
&\quad + \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{9x^2\sqrt[3]{1-x^3}\operatorname{Hypergeometric2F1}\left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3\right)}{2\sqrt[3]{-1+x^3}} \\
&\quad + \frac{9x^5\sqrt[3]{1-x^3}\operatorname{Hypergeometric2F1}\left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3\right)}{5\sqrt[3]{-1+x^3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2}\log\left(1+\sqrt[3]{-1+x^3}\right) - \frac{1}{2}\log\left(-x+\sqrt[3]{-1+x^3}\right) - \operatorname{Subst}\left(\int \frac{1}{-3-x^2} dx, x, -1+2\sqrt[3]{-1+x^3}\right) \\
&= -\frac{9}{2(-1+x^3)^{4/3}} - \frac{x}{(-1+x^3)^{4/3}} - \frac{7x^4}{2(-1+x^3)^{4/3}} - \frac{3}{\sqrt[3]{-1+x^3}} \\
&\quad + \frac{2x}{\sqrt[3]{-1+x^3}} + \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{\sqrt{3}} + \frac{\arctan\left(\frac{-1+2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{\sqrt{3}} \\
&\quad + \frac{9x^2\sqrt[3]{1-x^3}\operatorname{Hypergeometric2F1}\left(\frac{2}{3}, \frac{7}{3}, \frac{5}{3}, x^3\right)}{2\sqrt[3]{-1+x^3}} \\
&\quad + \frac{9x^5\sqrt[3]{1-x^3}\operatorname{Hypergeometric2F1}\left(\frac{5}{3}, \frac{7}{3}, \frac{8}{3}, x^3\right)}{5\sqrt[3]{-1+x^3}} + \frac{\log(x)}{2} \\
&\quad - \frac{1}{2}\log\left(1+\sqrt[3]{-1+x^3}\right) - \frac{1}{2}\log\left(-x+\sqrt[3]{-1+x^3}\right)
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 10.16 (sec) , antiderivative size = 123, normalized size of antiderivative = 0.84

$$\begin{aligned}
&\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x^3)^3 x} dx = -\frac{3(-1+x^3)^{2/3}}{2(-1+x)^2} + \frac{x\sqrt[3]{1-x^3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{3}, \frac{4}{3}, x^3\right)}{\sqrt[3]{-1+x^3}} \\
&+ \frac{1}{6}\left(-2\sqrt{3}\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right) - 2\log\left(1+\sqrt[3]{-1+x^3}\right) + \log\left(1-\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)\right)
\end{aligned}$$

[In] Integrate[((1 + x)*(-1 + x^3)^(2/3))/((-1 + x)^3*x), x]

[Out] (-3*(-1 + x^3)^(2/3))/(2*(-1 + x)^2) + (x*(1 - x^3)^(1/3)*Hypergeometric2F1[1/3, 1/3, 4/3, x^3])/(-1 + x^3)^(1/3) + (-2*sqrt[3]*ArcTan[(1 - 2*(-1 + x^3)^(1/3))/sqrt[3]] - 2*Log[1 + (-1 + x^3)^(1/3)] + Log[1 - (-1 + x^3)^(1/3) + (-1 + x^3)^(2/3)])/6

Maple [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.66 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.90

method	result
risch	$-\frac{3(x^2+x+1)}{2(-1+x)(x^3-1)^{\frac{1}{3}}} + \frac{\sqrt{3}\Gamma(\frac{2}{3})(-\text{signum}(x^3-1))^{\frac{1}{3}} \left(\frac{2\left(-\frac{\pi\sqrt{3}}{6} - \frac{3\ln(3)}{2} + 3\ln(x) + i\pi\right)\pi\sqrt{3}}{3\Gamma(\frac{2}{3})} + \frac{2\pi\sqrt{3}x^3 \text{hypergeom}\left(\left[1,1,\frac{4}{3}\right],[2,2],x^3\right)}{9\Gamma(\frac{2}{3})}\right)}{6\pi \text{signum}(x^3-1)^{\frac{1}{3}}}$
trager	$-\frac{3(x^3-1)^{\frac{2}{3}}}{2(-1+x)^2} + \ln\left(-\frac{58 \text{RootOf}(_Z^2 - _Z+1)^2 x^2 - 145 \text{RootOf}(_Z^2 - _Z+1)^2 x - 99 \text{RootOf}(_Z^2 - _Z+1)(x^3-1)^{\frac{2}{3}}}{\dots}\right)$

[In] int((1+x)*(x^3-1)^(2/3)/(-1+x)^3/x,x,method=_RETURNVERBOSE)

[Out] $-3/2*(x^2+x+1)/(-1+x)/(x^3-1)^{(1/3)}+1/6/\text{Pi}*3^{(1/2)}*\text{GAMMA}(2/3)/\text{signum}(x^3-1)^{(1/3)}*(-\text{signum}(x^3-1))^{(1/3)}*(2/3*(-1/6*\text{Pi}*3^{(1/2)}-3/2*\ln(3)+3*\ln(x)+\text{I}*\text{Pi})*\text{Pi}*3^{(1/2)}/\text{GAMMA}(2/3)+2/9*\text{Pi}*3^{(1/2)}/\text{GAMMA}(2/3)*x^3*\text{hypergeom}([1,1,4/3],[2,2],x^3))+1/\text{signum}(x^3-1)^{(1/3)}*(-\text{signum}(x^3-1))^{(1/3)}*x*\text{hypergeom}([1/3,1/3],[4/3],x^3)$

Fricas [B] (verification not implemented)

Error detected during grading. Assigning place holder grade for now.

Time = 0.36 (sec) , antiderivative size = 125, normalized size of antiderivative = 0.85

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \frac{2\sqrt{3}(x^2-2x+1) \arctan\left(-\frac{4\sqrt{3}(x^3-1)^{1/3}(x-1)+\sqrt{3}(x^2+x+1)-2\sqrt{3}(x^3-1)^{2/3}}{3(3x^2-5x+3)}\right) - (x^2 - 1) + \sqrt{3}(x^2+x+1) - 2\sqrt{3}(x^3-1)^{2/3}}{2(x^2-2x+1)}$$

[In] integrate((1+x)*(x^3-1)^(2/3)/(-1+x)^3/x,x, algorithm="fricas")

[Out] $1/2*(2*\text{sqrt}(3)*(x^2 - 2*x + 1)*\text{arctan}(-1/3*(4*\text{sqrt}(3)*(x^3 - 1)^{(1/3)}*(x - 1) + \text{sqrt}(3)*(x^2 + x + 1) - 2*\text{sqrt}(3)*(x^3 - 1)^{(2/3)})/(3*x^2 - 5*x + 3)) - (x^2 - 2*x + 1)*\log(((x^3 - 1)^{(1/3)}*(x - 1) + x - (x^3 - 1)^{(2/3)})/x) - 3*(x^3 - 1)^{(2/3)})/(x^2 - 2*x + 1)$

Sympy [F]

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \int \frac{((x-1)(x^2+x+1))^{2/3}(x+1)}{x(x-1)^3} dx$$

[In] integrate((1+x)*(x**3-1)**(2/3)/(-1+x)**3/x,x)

[Out] Integral(((x - 1)*(x**2 + x + 1))**(2/3)*(x + 1)/(x*(x - 1)**3), x)

Maxima [F]

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \int \frac{(x^3-1)^{2/3}(x+1)}{(x-1)^3 x} dx$$

[In] integrate((1+x)*(x^3-1)^(2/3)/(-1+x)^3/x,x, algorithm="maxima")

[Out] integrate((x^3 - 1)^(2/3)*(x + 1)/((x - 1)^3*x), x)

Giac [F]

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \int \frac{(x^3-1)^{2/3}(x+1)}{(x-1)^3 x} dx$$

[In] integrate((1+x)*(x^3-1)^(2/3)/(-1+x)^3/x,x, algorithm="giac")

[Out] integrate((x^3 - 1)^(2/3)*(x + 1)/((x - 1)^3*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(1+x)(-1+x^3)^{2/3}}{(-1+x)^3 x} dx = \int \frac{(x^3-1)^{2/3}(x+1)}{x(x-1)^3} dx$$

[In] int(((x^3 - 1)^(2/3)*(x + 1))/(x*(x - 1)^3),x)

[Out] int(((x^3 - 1)^(2/3)*(x + 1))/(x*(x - 1)^3), x)

$$3.6 \quad \int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx$$

Optimal result	67
Rubi [C] (verified)	68
Mathematica [A] (verified)	70
Maple [C] (verified)	70
Fricas [A] (verification not implemented)	71
Sympy [F(-1)]	71
Maxima [F]	72
Giac [F]	72
Mupad [F(-1)]	72

Optimal result

Integrand size = 26, antiderivative size = 90

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \frac{\arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2x^2+\sqrt[3]{-1+x^3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(x^2 + \sqrt[3]{-1+x^3}\right) - \frac{1}{6} \log\left(x^4 - x^2\sqrt[3]{-1+x^3} + (-1+x^3)^{2/3}\right)$$

[Out] 1/3*arctan(3^(1/2)*(x^3-1)^(1/3)/(-2*x^2+(x^3-1)^(1/3)))*3^(1/2)+1/3*ln(x^2+(x^3-1)^(1/3))-1/6*ln(x^4-x^2*(x^3-1)^(1/3)+(x^3-1)^(2/3))

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.43 (sec) , antiderivative size = 261, normalized size of antiderivative = 2.90, number of steps used = 14, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {6860, 1542, 525, 524}

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \frac{2\sqrt[3]{1-x^3}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, 1, \frac{5}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{\sqrt{5}(1+\sqrt{5})\sqrt[3]{x^3-1}} + \frac{\sqrt[3]{1-x^3}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, 1, \frac{5}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{2\sqrt{5}\sqrt[3]{x^3-1}} - \frac{2\sqrt[3]{1-x^3}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, -\frac{2x^3}{1-\sqrt{5}}, x^3\right)}{\sqrt{5}(1-\sqrt{5})\sqrt[3]{x^3-1}} - \frac{\sqrt[3]{1-x^3}x^2 \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, -\frac{2x^3}{1-\sqrt{5}}, x^3\right)}{2\sqrt{5}\sqrt[3]{x^3-1}}$$

[In] Int[(x*(-2 + x^3))/((-1 + x^3)^(1/3)*(-1 + x^3 + x^6)),x]

[Out] (x^2*(1 - x^3)^(1/3)*AppellF1[2/3, 1/3, 1, 5/3, x^3, (-2*x^3)/(1 + Sqrt[5])])/((2*Sqrt[5]*(-1 + x^3)^(1/3)) + (2*x^2*(1 - x^3)^(1/3)*AppellF1[2/3, 1/3, 1, 5/3, x^3, (-2*x^3)/(1 + Sqrt[5])])/(Sqrt[5]*(1 + Sqrt[5])*(-1 + x^3)^(1/3)) - (x^2*(1 - x^3)^(1/3)*AppellF1[2/3, 1, 1/3, 5/3, (-2*x^3)/(1 - Sqrt[5])], x^3))/(2*Sqrt[5]*(-1 + x^3)^(1/3)) - (2*x^2*(1 - x^3)^(1/3)*AppellF1[2/3, 1, 1/3, 5/3, (-2*x^3)/(1 - Sqrt[5]), x^3])/(Sqrt[5]*(1 - Sqrt[5])*(-1 + x^3)^(1/3))

Rule 524

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m + 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 525

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a)^FracPart[p])), Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 1542

Int[(((f._)*(x_)^(m_.)*((d_) + (e._)*(x_)^(n_.))^(q_))/((a_) + (c._)*(x_)^(n2_.) + (b._)*(x_)^(n_.)), x_Symbol] := Int[ExpandIntegrand[(d + e*x^n)^q, (f*x)^m/(a + b*x^n + c*x^(2*n)), x], x] /; FreeQ[{a, b, c, d, e, f, q, n}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && IGtQ[n, 0] && !IntegerQ[q] && IntegerQ[m]

Rule 6860

Int[(u_)/((a_.) + (b._)*(x_)^(n_.) + (c._)*(x_)^(n2_.)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n)), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(-\frac{2x}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} + \frac{x^4}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} \right) dx \\
&= -\left(2 \int \frac{x}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx \right) + \int \frac{x^4}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx \\
&= -\left(2 \int \left(-\frac{2x}{\sqrt{5}(-1+\sqrt{5}-2x^3)\sqrt[3]{-1+x^3}} - \frac{2x}{\sqrt{5}\sqrt[3]{-1+x^3}(1+\sqrt{5}+2x^3)} \right) dx \right) \\
&\quad + \int \left(-\frac{(-1+\sqrt{5})x}{\sqrt{5}(-1+\sqrt{5}-2x^3)\sqrt[3]{-1+x^3}} + \frac{(1+\sqrt{5})x}{\sqrt{5}\sqrt[3]{-1+x^3}(1+\sqrt{5}+2x^3)} \right) dx \\
&= \frac{4 \int \frac{x}{(-1+\sqrt{5}-2x^3)\sqrt[3]{-1+x^3}} dx}{\sqrt{5}} + \frac{4 \int \frac{x}{\sqrt[3]{-1+x^3}(1+\sqrt{5}+2x^3)} dx}{\sqrt{5}} \\
&\quad - \frac{1}{5}(5-\sqrt{5}) \int \frac{x}{(-1+\sqrt{5}-2x^3)\sqrt[3]{-1+x^3}} dx \\
&\quad + \frac{1}{5}(5+\sqrt{5}) \int \frac{x}{\sqrt[3]{-1+x^3}(1+\sqrt{5}+2x^3)} dx \\
&= \frac{(4\sqrt[3]{1-x^3}) \int \frac{x}{(-1+\sqrt{5}-2x^3)\sqrt[3]{1-x^3}} dx}{\sqrt{5}\sqrt[3]{-1+x^3}} + \frac{(4\sqrt[3]{1-x^3}) \int \frac{x}{\sqrt[3]{1-x^3}(1+\sqrt{5}+2x^3)} dx}{\sqrt{5}\sqrt[3]{-1+x^3}} \\
&\quad - \frac{((5-\sqrt{5})\sqrt[3]{1-x^3}) \int \frac{x}{(-1+\sqrt{5}-2x^3)\sqrt[3]{1-x^3}} dx}{5\sqrt[3]{-1+x^3}} \\
&\quad + \frac{((5+\sqrt{5})\sqrt[3]{1-x^3}) \int \frac{x}{\sqrt[3]{1-x^3}(1+\sqrt{5}+2x^3)} dx}{5\sqrt[3]{-1+x^3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x^2 \sqrt[3]{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, 1, \frac{5}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{2\sqrt{5} \sqrt[3]{-1+x^3}} \\
&+ \frac{2x^2 \sqrt[3]{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, \frac{1}{3}, 1, \frac{5}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{\sqrt{5}(1+\sqrt{5}) \sqrt[3]{-1+x^3}} \\
&- \frac{x^2 \sqrt[3]{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, -\frac{2x^3}{1-\sqrt{5}}, x^3\right)}{2\sqrt{5} \sqrt[3]{-1+x^3}} \\
&- \frac{2x^2 \sqrt[3]{1-x^3} \operatorname{AppellF1}\left(\frac{2}{3}, 1, \frac{1}{3}, \frac{5}{3}, -\frac{2x^3}{1-\sqrt{5}}, x^3\right)}{\sqrt{5}(1-\sqrt{5}) \sqrt[3]{-1+x^3}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \frac{\arctan\left(\frac{\sqrt{3} \sqrt[3]{-1+x^3}}{-2x^2+\sqrt[3]{-1+x^3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(x^2 + \sqrt[3]{-1+x^3}\right) - \frac{1}{6} \log\left(x^4 - x^2 \sqrt[3]{-1+x^3} + (-1+x^3)^{2/3}\right)$$

[In] Integrate[(x*(-2 + x^3))/((-1 + x^3)^(1/3)*(-1 + x^3 + x^6)),x]

[Out] ArcTan[(Sqrt[3]*(-1 + x^3)^(1/3))/(-2*x^2 + (-1 + x^3)^(1/3))]/Sqrt[3] + Log[x^2 + (-1 + x^3)^(1/3)]/3 - Log[x^4 - x^2*(-1 + x^3)^(1/3) + (-1 + x^3)^(2/3)]/6

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.47 (sec) , antiderivative size = 290, normalized size of antiderivative = 3.22

method	result
trager	$\ln\left(\frac{-9\operatorname{RootOf}(9_Z^2+3_Z+1)^2 x^6 - 9\operatorname{RootOf}(9_Z^2+3_Z+1) x^6 - 9\operatorname{RootOf}(9_Z^2+3_Z+1) (x^3-1)^{\frac{1}{3}} x^4 - 2x^6 + 3(x^3-1)^{\frac{2}{3}} x^2 + 3\operatorname{RootOf}(9_Z^2+3_Z+1)}{x^6 + x^3 - 1}\right)$
	3

[In] int(x*(x^3-2)/(x^3-1)^(1/3)/(x^6+x^3-1),x,method=_RETURNVERBOSE)

[Out] 1/3*ln(-(-9*RootOf(9*_Z^2+3*_Z+1)^2*x^6-9*RootOf(9*_Z^2+3*_Z+1)*x^6-9*RootOf(9*_Z^2+3*_Z+1)*(x^3-1)^(1/3)*x^4-2*x^6+3*(x^3-1)^(2/3)*x^2+3*RootOf(9*_Z^2+3*_Z+1))

$2+3*_Z+1)*x^3+2*x^3-3*\text{RootOf}(9*_Z^2+3*_Z+1)-2)/(x^6+x^3-1))+\text{RootOf}(9*_Z^2+3*_Z+1)*\ln((-9*\text{RootOf}(9*_Z^2+3*_Z+1)^2*x^6-3*\text{RootOf}(9*_Z^2+3*_Z+1)*x^6+6*\text{RootOf}(9*_Z^2+3*_Z+1)*(x^3-1)^{1/3})*x^4+3*\text{RootOf}(9*_Z^2+3*_Z+1)*(x^3-1)^{2/3})*x^2+(x^3-1)^{1/3}*x^4-(x^3-1)^{2/3}*x^2-3*\text{RootOf}(9*_Z^2+3*_Z+1)*x^3-x^3+3*\text{RootOf}(9*_Z^2+3*_Z+1)+1)/(x^6+x^3-1))$

Fricas [A] (verification not implemented)

none

Time = 1.36 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.16

$$\begin{aligned}
 & \int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx \\
 &= -\frac{1}{3} \sqrt{3} \arctan \left(\frac{\sqrt{3}x^6 + 2\sqrt{3}(x^3-1)^{\frac{1}{3}}x^4 + 4\sqrt{3}(x^3-1)^{\frac{2}{3}}x^2}{x^6 - 8x^3 + 8} \right) \\
 &+ \frac{1}{6} \log \left(\frac{x^6 + 3(x^3-1)^{\frac{1}{3}}x^4 + x^3 + 3(x^3-1)^{\frac{2}{3}}x^2 - 1}{x^6 + x^3 - 1} \right)
 \end{aligned}$$

[In] integrate(x*(x^3-2)/(x^3-1)^(1/3)/(x^6+x^3-1),x, algorithm="fricas")

[Out] -1/3*sqrt(3)*arctan((sqrt(3)*x^6 + 2*sqrt(3)*(x^3 - 1)^(1/3)*x^4 + 4*sqrt(3)*(x^3 - 1)^(2/3)*x^2)/(x^6 - 8*x^3 + 8)) + 1/6*log((x^6 + 3*(x^3 - 1)^(1/3)*x^4 + x^3 + 3*(x^3 - 1)^(2/3)*x^2 - 1)/(x^6 + x^3 - 1))

Sympy [F(-1)]

Timed out.

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \text{Timed out}$$

[In] integrate(x*(x**3-2)/(x**3-1)**(1/3)/(x**6+x**3-1),x)

[Out] Timed out

Maxima [F]

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \int \frac{(x^3-2)x}{(x^6+x^3-1)(x^3-1)^{\frac{1}{3}}} dx$$

[In] integrate(x*(x^3-2)/(x^3-1)^(1/3)/(x^6+x^3-1),x, algorithm="maxima")

[Out] integrate((x^3 - 2)*x/((x^6 + x^3 - 1)*(x^3 - 1)^(1/3)), x)

Giac [F]

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \int \frac{(x^3-2)x}{(x^6+x^3-1)(x^3-1)^{\frac{1}{3}}} dx$$

[In] integrate(x*(x^3-2)/(x^3-1)^(1/3)/(x^6+x^3-1),x, algorithm="giac")

[Out] integrate((x^3 - 2)*x/((x^6 + x^3 - 1)*(x^3 - 1)^(1/3)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x(-2+x^3)}{\sqrt[3]{-1+x^3}(-1+x^3+x^6)} dx = \int \frac{x(x^3-2)}{(x^3-1)^{1/3}(x^6+x^3-1)} dx$$

[In] int((x*(x^3 - 2))/((x^3 - 1)^(1/3)*(x^3 + x^6 - 1)),x)

[Out] int((x*(x^3 - 2))/((x^3 - 1)^(1/3)*(x^3 + x^6 - 1)), x)

$$3.7 \quad \int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx$$

Optimal result	73
Rubi [C] (verified)	73
Mathematica [A] (verified)	75
Maple [C] (verified)	75
Fricas [A] (verification not implemented)	76
Sympy [F]	76
Maxima [F]	76
Giac [F]	77
Mupad [F(-1)]	77

Optimal result

Integrand size = 28, antiderivative size = 85

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = -\frac{\arctan\left(\frac{\sqrt{3}x^2}{x^2-2\sqrt[3]{1+x^3}}\right)}{\sqrt{3}} - \frac{1}{3} \log\left(x^2 + \sqrt[3]{1+x^3}\right) + \frac{1}{6} \log\left(x^4 - x^2\sqrt[3]{1+x^3} + (1+x^3)^{2/3}\right)$$

[Out] $-1/3*\arctan(3^{(1/2)}*x^2/(x^2-2*(x^3+1)^{(1/3)}))*3^{(1/2)}-1/3*\ln(x^2+(x^3+1)^{(1/3)})+1/6*\ln(x^4-x^2*(x^3+1)^{(1/3)}+(x^3+1)^{(2/3)})$

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.35 (sec) , antiderivative size = 134, normalized size of antiderivative = 1.58, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {6860, 251, 440}

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = \frac{(-\sqrt{3}+i)x \operatorname{AppellF1}\left(\frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, -x^3, -\frac{2x^3}{1-i\sqrt{3}}\right)}{\sqrt{3}+i} + \frac{(\sqrt{3}+i)x \operatorname{AppellF1}\left(\frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, -x^3, -\frac{2x^3}{1+i\sqrt{3}}\right)}{-\sqrt{3}+i} + x \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -x^3\right)$$

[In] $\text{Int}[(x^3*(2+x^3))/((1+x^3)^{(2/3)}*(1+x^3+x^6)),x]$

```
[Out] ((I - Sqrt[3])*x*AppellF1[1/3, 2/3, 1, 4/3, -x^3, (-2*x^3)/(1 - I*Sqrt[3])]
)/(I + Sqrt[3]) + ((I + Sqrt[3])*x*AppellF1[1/3, 2/3, 1, 4/3, -x^3, (-2*x^3
)/(1 + I*Sqrt[3])])/(I - Sqrt[3]) + x*Hypergeometric2F1[1/3, 2/3, 4/3, -x^3
]
```

Rule 251

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p*x*Hypergeometric2F
1[-p, 1/n, 1/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, n, p}, x] && !IGtQ[p
, 0] && !IntegerQ[1/n] && !ILtQ[Simplify[1/n + p], 0] && (IntegerQ[p] ||
GtQ[a, 0])
```

Rule 440

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)
], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])
```

Rule 6860

```
Int[(u_)/((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.)), x_Symbol] := With[
{v = RationalFunctionExpand[u/(a + b*x^n + c*x^(2*n))], x}, Int[v, x] /; Su
mQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{(1+x^3)^{2/3}} - \frac{1-x^3}{(1+x^3)^{2/3}(1+x^3+x^6)} \right) dx \\
&= \int \frac{1}{(1+x^3)^{2/3}} dx - \int \frac{1-x^3}{(1+x^3)^{2/3}(1+x^3+x^6)} dx \\
&= x \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -x^3 \right) \\
&\quad - \int \left(\frac{-1-i\sqrt{3}}{(1+x^3)^{2/3}(1-i\sqrt{3}+2x^3)} + \frac{-1+i\sqrt{3}}{(1+x^3)^{2/3}(1+i\sqrt{3}+2x^3)} \right) dx \\
&= x \operatorname{Hypergeometric2F1} \left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -x^3 \right) \\
&\quad - (-1-i\sqrt{3}) \int \frac{1}{(1+x^3)^{2/3}(1-i\sqrt{3}+2x^3)} dx \\
&\quad - (-1+i\sqrt{3}) \int \frac{1}{(1+x^3)^{2/3}(1+i\sqrt{3}+2x^3)} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{(i - \sqrt{3}) x \operatorname{AppellF1}\left(\frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, -x^3, -\frac{2x^3}{1-i\sqrt{3}}\right)}{i + \sqrt{3}} \\
&+ \frac{(i + \sqrt{3}) x \operatorname{AppellF1}\left(\frac{1}{3}, \frac{2}{3}, 1, \frac{4}{3}, -x^3, -\frac{2x^3}{1+i\sqrt{3}}\right)}{i - \sqrt{3}} \\
&+ x \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, -x^3\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.46 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = -\frac{\arctan\left(\frac{\sqrt{3}x^2}{x^2-2\sqrt[3]{1+x^3}}\right)}{\sqrt{3}} - \frac{1}{3} \log\left(x^2 + \sqrt[3]{1+x^3}\right) + \frac{1}{6} \log\left(x^4 - x^2\sqrt[3]{1+x^3} + (1+x^3)^{2/3}\right)$$

[In] Integrate[(x^3*(2 + x^3))/((1 + x^3)^(2/3)*(1 + x^3 + x^6)),x]

[Out] -(ArcTan[(Sqrt[3]*x^2)/(x^2 - 2*(1 + x^3)^(1/3))]/Sqrt[3]) - Log[x^2 + (1 + x^3)^(1/3)]/3 + Log[x^4 - x^2*(1 + x^3)^(1/3) + (1 + x^3)^(2/3)]/6

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.53 (sec) , antiderivative size = 292, normalized size of antiderivative = 3.44

method	result
trager	$-\frac{\ln\left(\frac{-9\operatorname{RootOf}(9_Z^2-3_Z+1)^2x^6-3\operatorname{RootOf}(9_Z^2-3_Z+1)(x^3+1)^{\frac{1}{3}}x^4+3\operatorname{RootOf}(9_Z^2-3_Z+1)(x^3+1)^{\frac{2}{3}}x^2+2(x^3+1)^{\frac{1}{3}}}{x^6+x^3+1}\right)}{3}$

[In] int(x^3*(x^3+2)/(x^3+1)^(2/3)/(x^6+x^3+1),x,method=_RETURNVERBOSE)

[Out] -1/3*ln((-9*RootOf(9*_Z^2-3*_Z+1)^2*x^6-3*RootOf(9*_Z^2-3*_Z+1)*(x^3+1)^(1/3)*x^4+3*RootOf(9*_Z^2-3*_Z+1)*(x^3+1)^(2/3)*x^2+2*(x^3+1)^(1/3)*x^4+(x^3+1)^(2/3)*x^2+3*RootOf(9*_Z^2-3*_Z+1)*x^3+3*RootOf(9*_Z^2-3*_Z+1))/(x^6+x^3+1))+RootOf(9*_Z^2-3*_Z+1)*ln((-9*RootOf(9*_Z^2-3*_Z+1)^2*x^6+3*RootOf(9*_Z^2-3*_Z+1)*x^6+6*RootOf(9*_Z^2-3*_Z+1)*(x^3+1)^(1/3)*x^4+3*RootOf(9*_Z^2-3*_Z+1)*(x^3+1)^(2/3)*x^2-(x^3+1)^(1/3)*x^4-2*(x^3+1)^(2/3)*x^2-3*RootOf(9*_Z^2-3*_Z+1)*x^3-3*RootOf(9*_Z^2-3*_Z+1))/(x^6+x^3+1))

Fricas [A] (verification not implemented)

none

Time = 1.16 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.92

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = -\frac{1}{3}\sqrt{3}\arctan\left(-\frac{\sqrt{3}x^2-2\sqrt{3}(x^3+1)^{1/3}}{3x^2}\right) - \frac{1}{6}\log\left(\frac{x^6+3(x^3+1)^{1/3}x^4+x^3+3(x^3+1)^{2/3}x^2+1}{x^6+x^3+1}\right)$$

[In] integrate(x^3*(x^3+2)/(x^3+1)^(2/3)/(x^6+x^3+1),x, algorithm="fricas")

[Out] -1/3*sqrt(3)*arctan(-1/3*(sqrt(3)*x^2 - 2*sqrt(3)*(x^3 + 1)^(1/3))/x^2) - 1/6*log((x^6 + 3*(x^3 + 1)^(1/3)*x^4 + x^3 + 3*(x^3 + 1)^(2/3)*x^2 + 1)/(x^6 + x^3 + 1))

Sympy [F]

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = \int \frac{x^3(x^3+2)}{((x+1)(x^2-x+1))^{2/3}(x^6+x^3+1)} dx$$

[In] integrate(x**3*(x**3+2)/(x**3+1)**(2/3)/(x**6+x**3+1),x)

[Out] Integral(x**3*(x**3 + 2)/(((x + 1)*(x**2 - x + 1))**(2/3)*(x**6 + x**3 + 1)), x)

Maxima [F]

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = \int \frac{(x^3+2)x^3}{(x^6+x^3+1)(x^3+1)^{2/3}} dx$$

[In] integrate(x^3*(x^3+2)/(x^3+1)^(2/3)/(x^6+x^3+1),x, algorithm="maxima")

[Out] integrate((x^3 + 2)*x^3/((x^6 + x^3 + 1)*(x^3 + 1)^(2/3)), x)

Giac [F]

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = \int \frac{(x^3+2)x^3}{(x^6+x^3+1)(x^3+1)^{2/3}} dx$$

[In] integrate(x^3*(x^3+2)/(x^3+1)^(2/3)/(x^6+x^3+1),x, algorithm="giac")

[Out] integrate((x^3 + 2)*x^3/((x^6 + x^3 + 1)*(x^3 + 1)^(2/3)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x^3(2+x^3)}{(1+x^3)^{2/3}(1+x^3+x^6)} dx = \int \frac{x^3(x^3+2)}{(x^3+1)^{2/3}(x^6+x^3+1)} dx$$

[In] int((x^3*(x^3 + 2))/((x^3 + 1)^(2/3)*(x^3 + x^6 + 1)),x)

[Out] int((x^3*(x^3 + 2))/((x^3 + 1)^(2/3)*(x^3 + x^6 + 1)), x)

$$3.8 \quad \int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx$$

Optimal result	78
Rubi [C] (verified)	78
Mathematica [A] (verified)	80
Maple [C] (verified)	80
Fricas [A] (verification not implemented)	81
Sympy [F(-1)]	81
Maxima [F]	81
Giac [F]	82
Mupad [F(-1)]	82

Optimal result

Integrand size = 30, antiderivative size = 110

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = -\frac{\sqrt[3]{1-x^3}}{x^2} - \frac{\arctan\left(\frac{\sqrt{3}x^2}{x^2+2\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} - \frac{1}{3} \log\left(-x^2 + \sqrt[3]{1-x^3}\right) + \frac{1}{6} \log\left(x^4 + x^2\sqrt[3]{1-x^3} + (1-x^3)^{2/3}\right)$$

[Out] $-(x^3+1)^{1/3}/x^2-1/3*\arctan(3^{1/2}*x^2/(x^2+2*(x^3+1)^{1/3}))*3^{1/2}-1/3*\ln(-x^2+(x^3+1)^{1/3})+1/6*\ln(x^4+x^2*(x^3+1)^{1/3}+(x^3+1)^{2/3})$

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.30 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.94, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {6860, 371, 440}

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = -\frac{2x \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{2x^3}{1-\sqrt{5}}\right)}{1-\sqrt{5}} - \frac{2x \operatorname{AppellF1}\left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{1+\sqrt{5}} - \frac{\operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, -\frac{1}{3}, \frac{1}{3}, x^3\right)}{x^2}$$

[In] Int[((1 - x^3)^(1/3)*(-2 + x^3))/(x^3*(-1 + x^3 + x^6)), x]

[Out] $(-2*x*AppellF1[1/3, -1/3, 1, 4/3, x^3, (-2*x^3)/(1 - \text{Sqrt}[5])])/(1 - \text{Sqrt}[5]) - (2*x*AppellF1[1/3, -1/3, 1, 4/3, x^3, (-2*x^3)/(1 + \text{Sqrt}[5])])/(1 + \text{Sqrt}[5]) - \text{Hypergeometric2F1}[-2/3, -1/3, 1/3, x^3]/x^2$

Rule 371

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 440

$\text{Int}[(a_) + (b_*)*(x_)^{(n_)})^{(p_*)}*((c_) + (d_*)*(x_)^{(n_)})^{(q_)}, x_Symbol] \rightarrow \text{Simp}[a^p * c^q * x * \text{AppellF1}[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /;$ FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 6860

$\text{Int}[(u_)/((a_*) + (b_*)*(x_)^{(n_*)} + (c_*)*(x_)^{(n2_*)}), x_Symbol] \rightarrow \text{With}[\{v = \text{RationalFunctionExpand}[u/(a + b*x^n + c*x^{(2*n)})], x\}, \text{Int}[v, x] /;$ SumQ[v]] /; FreeQ[{a, b, c}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{2\sqrt[3]{1-x^3}}{x^3} + \frac{(-1-2x^3)\sqrt[3]{1-x^3}}{-1+x^3+x^6} \right) dx \\
 &= 2 \int \frac{\sqrt[3]{1-x^3}}{x^3} dx + \int \frac{(-1-2x^3)\sqrt[3]{1-x^3}}{-1+x^3+x^6} dx \\
 &= -\frac{\text{Hypergeometric2F1}\left(-\frac{2}{3}, -\frac{1}{3}, \frac{1}{3}, x^3\right)}{x^2} + \int \left(-\frac{2\sqrt[3]{1-x^3}}{1-\sqrt{5}+2x^3} - \frac{2\sqrt[3]{1-x^3}}{1+\sqrt{5}+2x^3} \right) dx \\
 &= -\frac{\text{Hypergeometric2F1}\left(-\frac{2}{3}, -\frac{1}{3}, \frac{1}{3}, x^3\right)}{x^2} - 2 \int \frac{\sqrt[3]{1-x^3}}{1-\sqrt{5}+2x^3} dx - 2 \int \frac{\sqrt[3]{1-x^3}}{1+\sqrt{5}+2x^3} dx \\
 &= -\frac{2x \text{AppellF1}\left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{2x^3}{1-\sqrt{5}}\right)}{1-\sqrt{5}} - \frac{2x \text{AppellF1}\left(\frac{1}{3}, -\frac{1}{3}, 1, \frac{4}{3}, x^3, -\frac{2x^3}{1+\sqrt{5}}\right)}{1+\sqrt{5}} \\
 &\quad - \frac{\text{Hypergeometric2F1}\left(-\frac{2}{3}, -\frac{1}{3}, \frac{1}{3}, x^3\right)}{x^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.05 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = -\frac{\sqrt[3]{1-x^3}}{x^2} - \frac{\arctan\left(\frac{\sqrt{3}x^2}{x^2+2\sqrt[3]{1-x^3}}\right)}{\sqrt{3}} - \frac{1}{3} \log\left(-x^2 + \sqrt[3]{1-x^3}\right) + \frac{1}{6} \log\left(x^4 + x^2\sqrt[3]{1-x^3} + (1-x^3)^{2/3}\right)$$

[In] Integrate[((1 - x^3)^(1/3)*(-2 + x^3))/(x^3*(-1 + x^3 + x^6)),x]

[Out] -((1 - x^3)^(1/3)/x^2) - ArcTan[(Sqrt[3]*x^2)/(x^2 + 2*(1 - x^3)^(1/3))]/Sqrt[3] - Log[-x^2 + (1 - x^3)^(1/3)]/3 + Log[x^4 + x^2*(1 - x^3)^(1/3) + (1 - x^3)^(2/3)]/6

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 5.34 (sec) , antiderivative size = 600, normalized size of antiderivative = 5.45

method	result
risch	$\frac{x^3-1}{x^2(-x^3+1)^{\frac{2}{3}}} + \ln\left(\frac{\left(9\text{RootOf}\left(9_Z^2-3_Z+1\right)^2 x^9-x^9-9\text{RootOf}\left(9_Z^2-3_Z+1\right)^2 x^6+3\text{RootOf}\left(9_Z^2-3_Z+1\right) x^6+3\left(x^6-2x^3\right)^{\frac{2}{3}}}{(-1+x)\left(x^2\right)}\right)$
trager	Expression too large to display

[In] int((-x^3+1)^(1/3)*(x^3-2)/x^3/(x^6+x^3-1),x,method=_RETURNVERBOSE)

[Out] (x^3-1)/x^2/(-x^3+1)^(2/3)+(1/3*ln((9*RootOf(9*_Z^2-3*_Z+1)^2*x^9-x^9-9*RootOf(9*_Z^2-3*_Z+1)^2*x^6+3*RootOf(9*_Z^2-3*_Z+1)*x^6+3*(x^6-2*x^3+1)^(2/3)*x^4-3*(x^6-2*x^3+1)^(1/3)*x^5+2*x^6-6*RootOf(9*_Z^2-3*_Z+1)*x^3+3*(x^6-2*x^3+1)^(1/3)*x^2-2*x^3+3*RootOf(9*_Z^2-3*_Z+1)+1)/(-1+x)/(x^2+x+1)/(x^6+x^3-1))-ln((9*RootOf(9*_Z^2-3*_Z+1)^2*x^9-x^9-9*RootOf(9*_Z^2-3*_Z+1)^2*x^6+3*RootOf(9*_Z^2-3*_Z+1)*x^6+3*(x^6-2*x^3+1)^(2/3)*x^4-3*(x^6-2*x^3+1)^(1/3)*x^5+2*x^6-6*RootOf(9*_Z^2-3*_Z+1)*x^3+3*(x^6-2*x^3+1)^(1/3)*x^2-2*x^3+3*RootOf(9*_Z^2-3*_Z+1)+1)/(-1+x)/(x^2+x+1)/(x^6+x^3-1))*RootOf(9*_Z^2-3*_Z+1)+RootOf(9*_Z^2-3*_Z+1)*ln((9*RootOf(9*_Z^2-3*_Z+1)^2*x^9-6*RootOf(9*_Z^2-3*_Z+1)*x^9-9*RootOf(9*_Z^2-3*_Z+1)^2*x^6+3*RootOf(9*_Z^2-3*_Z+1)*x^6+3*(x^6-2*x^3+1)^(2/3)*x^4-3*(x^6-2*x^3+1)^(1/3)*x^5+2*x^6+6*RootOf(9*_Z^2-3*_Z+1)*x^3+3

$$\frac{(x^6 - 2x^3 + 1)^{1/3} x^2 - 4x^3 - 3 \operatorname{RootOf}(9Z^2 - 3Z + 1) + 2}{(-1+x)(x^2+x+1)(x^6+x^3-1)} \frac{1}{(-x^3+1)^{2/3}((x^3-1)^2)^{1/3}}$$

Fricas [A] (verification not implemented)

none

Time = 1.50 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.21

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = \frac{2\sqrt{3}x^2 \arctan\left(\frac{\sqrt{3}x^6 - 2\sqrt{3}(-x^3+1)^{1/3}x^4 + 4\sqrt{3}(-x^3+1)^{2/3}x^2}{x^6 - 8x^3 + 8}\right) + x^2 \log\left(\frac{x^6 - 3(-x^3+1)^{1/3}x^4 + x^3 + 3(-x^3+1)^{2/3}x^2 - 1}{x^6 + x^3 - 1}\right) + 6}{6x^2}$$

[In] integrate((-x^3+1)^(1/3)*(x^3-2)/x^3/(x^6+x^3-1),x, algorithm="fricas")

[Out] -1/6*(2*sqrt(3)*x^2*arctan((sqrt(3)*x^6 - 2*sqrt(3)*(-x^3 + 1)^(1/3)*x^4 + 4*sqrt(3)*(-x^3 + 1)^(2/3)*x^2)/(x^6 - 8*x^3 + 8)) + x^2*log((x^6 - 3*(-x^3 + 1)^(1/3)*x^4 + x^3 + 3*(-x^3 + 1)^(2/3)*x^2 - 1)/(x^6 + x^3 - 1)) + 6*(-x^3 + 1)^(1/3))/x^2

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = \text{Timed out}$$

[In] integrate((-x**3+1)**(1/3)*(x**3-2)/x**3/(x**6+x**3-1),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = \int \frac{(x^3-2)(-x^3+1)^{1/3}}{(x^6+x^3-1)x^3} dx$$

[In] integrate((-x^3+1)^(1/3)*(x^3-2)/x^3/(x^6+x^3-1),x, algorithm="maxima")

[Out] integrate((x^3 - 2)*(-x^3 + 1)^(1/3)/((x^6 + x^3 - 1)*x^3), x)

Giac [F]

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = \int \frac{(x^3-2)(-x^3+1)^{\frac{1}{3}}}{(x^6+x^3-1)x^3} dx$$

[In] integrate((-x^3+1)^(1/3)*(x^3-2)/x^3/(x^6+x^3-1),x, algorithm="giac")

[Out] integrate((x^3 - 2)*(-x^3 + 1)^(1/3)/((x^6 + x^3 - 1)*x^3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt[3]{1-x^3}(-2+x^3)}{x^3(-1+x^3+x^6)} dx = \int \frac{(1-x^3)^{1/3}(x^3-2)}{x^3(x^6+x^3-1)} dx$$

[In] int(((1 - x^3)^(1/3)*(x^3 - 2))/(x^3*(x^3 + x^6 - 1)),x)

[Out] int(((1 - x^3)^(1/3)*(x^3 - 2))/(x^3*(x^3 + x^6 - 1)), x)

$$3.9 \quad \int \frac{(2+x+x^2) \sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx$$

Optimal result	83
Rubi [F]	83
Mathematica [A] (verified)	90
Maple [C] (verified)	90
Fricas [F(-2)]	92
Sympy [F(-1)]	92
Maxima [F]	93
Giac [F]	93
Mupad [F(-1)]	93

Optimal result

Integrand size = 39, antiderivative size = 119

$$\int \frac{(2+x+x^2) \sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \frac{\sqrt[3]{-1+x^3}}{-1+x^2} - \frac{\arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x^2+\sqrt[3]{-1+x^3}}\right)}{\sqrt{3}} + \frac{1}{3} \log\left(1-x^2+\sqrt[3]{-1+x^3}\right) - \frac{1}{6} \log\left(1-2x^2+x^4+(-1+x^2)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)$$

[Out] (x^3-1)^(1/3)/(x^2-1)-1/3*arctan(3^(1/2)*(x^3-1)^(1/3)/(-2+2*x^2+(x^3-1)^(1/3)))*3^(1/2)+1/3*ln(1-x^2+(x^3-1)^(1/3))-1/6*ln(1-2*x^2+x^4+(x^2-1)*(x^3-1)^(1/3)+(x^3-1)^(2/3))

Rubi [F]

$$\int \frac{(2+x+x^2) \sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \int \frac{(2+x+x^2) \sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx$$

[In] Int[((2 + x + x^2)*(-1 + x^3)^(1/3))/(x*(-1 + x^2)^2*(-3 - 2*x + x^2 + x^3)), x]

[Out] 1/(2*(-1 + x^3)^(2/3)) + x^2/(2*(-1 + x^3)^(2/3)) - (-1 + x^3)^(1/3)/3 - (-1 + x^3)^(1/3)/(2*(1 + x^3)) + (x*(-1 + x^3)^(1/3))/(2*(1 + x^3)) - (x^2*(-1 + x^3)^(1/3))/(2*(1 + x^3)) - (2*ArcTan[(1 - 2*(-1 + x^3)^(1/3))/Sqrt[3]])/(3*Sqrt[3]) - (x*(1 - x^3)^(2/3)*Hypergeometric2F1[1/3, 2/3, 4/3, x^3])/((

$6*(-1 + x^3)^{(2/3)} + (x*(1 - x^3)^{(2/3)}*\text{Hypergeometric2F1}[1/3, 5/3, 4/3, x^3])/(3*(-1 + x^3)^{(2/3)} + (x^4*(1 - x^3)^{(2/3)}*\text{Hypergeometric2F1}[4/3, 5/3, 7/3, x^3])/(6*(-1 + x^3)^{(2/3)} - \text{Log}[x]/3 + \text{Log}[1 + (-1 + x^3)^{(1/3)}]/3 + (2*\text{Defer}[\text{Int}] [(-1 + x^3)^{(1/3)} / (-3 - 2*x + x^2 + x^3), x]) / 3 + \text{Defer}[\text{Int}] [(x*(-1 + x^3)^{(1/3)}) / (-3 - 2*x + x^2 + x^3), x] / 3 + \text{Defer}[\text{Int}] [(x^2*(-1 + x^3)^{(1/3)}) / (-3 - 2*x + x^2 + x^3), x] / 3$

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(-\frac{\sqrt[3]{-1+x^3}}{3(-1+x)^2} + \frac{\sqrt[3]{-1+x^3}}{12(-1+x)} - \frac{2\sqrt[3]{-1+x^3}}{3x} + \frac{\sqrt[3]{-1+x^3}}{2(1+x)^2} + \frac{\sqrt[3]{-1+x^3}}{4(1+x)} \right. \\
&\quad \left. + \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{3(-3-2x+x^2+x^3)} \right) dx \\
&= \frac{1}{12} \int \frac{\sqrt[3]{-1+x^3}}{-1+x} dx + \frac{1}{4} \int \frac{\sqrt[3]{-1+x^3}}{1+x} dx - \frac{1}{3} \int \frac{\sqrt[3]{-1+x^3}}{(-1+x)^2} dx \\
&\quad + \frac{1}{3} \int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{1}{2} \int \frac{\sqrt[3]{-1+x^3}}{(1+x)^2} dx - \frac{2}{3} \int \frac{\sqrt[3]{-1+x^3}}{x} dx \\
&= \frac{1}{12} \int \left(\frac{1}{(-1+x^3)^{2/3}} + \frac{x}{(-1+x^3)^{2/3}} \right. \\
&\quad \left. + \frac{x^2}{(-1+x^3)^{2/3}} \right) dx - \frac{2}{9} \text{Subst} \left(\int \frac{\sqrt[3]{-1+x}}{x} dx, x, x^3 \right) + \frac{1}{4} \int \left(\frac{\sqrt[3]{-1+x^3}}{1+x^3} - \frac{x\sqrt[3]{-1+x^3}}{1+x^3} + \frac{x^2\sqrt[3]{-1+x^3}}{1+x^3} \right) dx \\
&= -\frac{2}{3} \sqrt[3]{-1+x^3} + \frac{1}{12} \int \frac{1}{(-1+x^3)^{2/3}} dx + \frac{1}{12} \int \frac{x}{(-1+x^3)^{2/3}} dx + \frac{1}{12} \int \frac{x^2}{(-1+x^3)^{2/3}} dx \\
&\quad + \frac{2}{9} \text{Subst} \left(\int \frac{1}{(-1+x)^{2/3}x} dx, x, x^3 \right) + \frac{1}{4} \int \frac{\sqrt[3]{-1+x^3}}{1+x^3} dx - \frac{1}{4} \int \frac{x\sqrt[3]{-1+x^3}}{1+x^3} dx + \frac{1}{4} \int \frac{x^2\sqrt[3]{-1+x^3}}{1+x^3} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{7}{12}\sqrt[3]{-1+x^3} + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad - \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{12\sqrt{3}} - \frac{\log(x)}{3} - \frac{1}{24}\log\left(x - \sqrt[3]{-1+x^3}\right) \\
&\quad + \frac{1}{12}\text{Subst}\left(\int \frac{\sqrt[3]{-1+x}}{1+x} dx, x, x^3\right) - \frac{1}{6}\int \frac{2-x^3}{(-1+x^3)^{2/3}(1+x^3)} dx \\
&\quad + \frac{1}{6}\int \frac{x(-2+3x^3)}{(-1+x^3)^{2/3}(1+x^3)} dx - \frac{1}{4}\int \frac{x}{(-1+x^3)^{2/3}} dx - \frac{1}{3}\int \frac{x}{(-1+x^3)^{2/3}} dx \\
&\quad + \frac{1}{3}\int \frac{x}{(-1+x^3)^{2/3}(1+x^3)} dx + \frac{1}{3}\int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx \\
&\quad + \frac{1}{3}\int \frac{x^2\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx - \frac{1}{3}\int \frac{-1+2x^3}{(-1+x^3)^{2/3}(1+x^3)} dx \\
&\quad + \frac{1}{3}\text{Subst}\left(\int \frac{1}{1+x} dx, x, \sqrt[3]{-1+x^3}\right) + \frac{1}{3}\text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \sqrt[3]{-1+x^3}\right) + \frac{1}{2}\int \frac{1}{(-1+x^3)^{2/3}} dx \\
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{2\sqrt{3}} \\
&\quad - \frac{5\arctan\left(\frac{1+\frac{2\sqrt[3]{2}x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{6\cdot 2^{2/3}\sqrt{3}} + \frac{x(1-x^3)^{2/3}\text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{12(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\text{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} + \frac{5\log(1+x^3)}{36\cdot 2^{2/3}} \\
&\quad + \frac{1}{4}\log\left(x - \sqrt[3]{-1+x^3}\right) - \frac{5\log\left(\sqrt[3]{2}x - \sqrt[3]{-1+x^3}\right)}{12\cdot 2^{2/3}} + \frac{1}{3}\log\left(1 + \sqrt[3]{-1+x^3}\right) + \frac{1}{6}\int \frac{1}{(-1+x^3)^{2/3}} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{\arctan\left(\frac{1+\frac{2x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{2\sqrt{3}} - \frac{5\arctan\left(\frac{1+\frac{2\sqrt[3]{2}x}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{6\cdot 2^{2/3}\sqrt{3}} \\
&\quad - \frac{2\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{12(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} + \frac{5\log(1+x^3)}{36\cdot 2^{2/3}} \\
&\quad + \frac{1}{4}\log\left(x-\sqrt[3]{-1+x^3}\right) - \frac{5\log\left(\sqrt[3]{2}x-\sqrt[3]{-1+x^3}\right)}{12\cdot 2^{2/3}} + \frac{1}{3}\log\left(1+\sqrt[3]{-1+x^3}\right) - \frac{1}{4}\int \frac{1}{(-1+x^3)^{2/3}} dx \\
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{2\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} \\
&\quad - \frac{5x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{12(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&\quad - \frac{\log\left(2^{2/3}+\frac{1-x}{\sqrt[3]{-1+x^3}}\right)}{12\cdot 2^{2/3}} - \frac{\log\left(1-\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{6\cdot 2^{2/3}} \\
&\quad + \frac{1}{3}\log\left(1+\sqrt[3]{-1+x^3}\right) + \frac{1}{8}\operatorname{Subst}\left(\int \frac{1}{2\sqrt[3]{2}-2^{2/3}x+x^2} dx, x, \frac{1-x}{\sqrt[3]{-1+x^3}}\right) + \frac{1}{3}\int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&+ \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{2 \arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} \\
&- \frac{x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{6(-1+x^3)^{2/3}} \\
&+ \frac{x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&+ \frac{x^4(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&- \frac{\log\left(2^{2/3} + \frac{1-x}{\sqrt[3]{-1+x^3}}\right)}{12 \cdot 2^{2/3}} - \frac{\log\left(1 - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{6 \cdot 2^{2/3}} \\
&+ \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(-1+x^3)^{2/3}} + \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{12 \cdot 2^{2/3}} + \frac{\log\left(2\sqrt[3]{2} + \frac{(1-x)^2}{(-1+x^3)^{2/3}} - \frac{2^{2/3}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{24 \cdot 2^{2/3}} \\
&+ \frac{1}{3} \log\left(1 + \sqrt[3]{-1+x^3}\right) + \frac{1}{4} \operatorname{Subst}\left(\int \frac{x}{4+x^3} dx, x, \frac{1-x}{\sqrt[3]{-1+x^3}}\right) + \frac{1}{3} \int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{1}{3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{\arctan\left(\frac{1-\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{4\ 2^{2/3}\sqrt{3}} - \frac{\arctan\left(\frac{1+\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{2\ 2^{2/3}\sqrt{3}} \\
&\quad - \frac{2\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{6(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&\quad - \frac{\log\left(2^{2/3} + \frac{1-x}{\sqrt[3]{-1+x^3}}\right)}{12\ 2^{2/3}} - \frac{\log\left(1 - \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{6\ 2^{2/3}} \\
&\quad + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(-1+x^3)^{2/3}} + \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{12\ 2^{2/3}} + \frac{\log\left(2\sqrt[3]{2} + \frac{(1-x)^2}{(-1+x^3)^{2/3}} - \frac{2^{2/3}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{24\ 2^{2/3}} \\
&\quad + \frac{1}{3}\log\left(1 + \sqrt[3]{-1+x^3}\right) + \frac{1}{3}\int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{1}{3}\int \frac{x^2\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{2}{3}\int \frac{\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{\arctan\left(\frac{1-\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{4\ 2^{2/3}\sqrt{3}} - \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{2\ 2^{2/3}\sqrt{3}} \\
&\quad - \frac{2\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{6(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&\quad + \frac{\log\left(1 + \frac{2^{2/3}(1-x)^2}{(-1+x^3)^{2/3}} + \frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{12\ 2^{2/3}} + \frac{\log\left(2\sqrt[3]{2} + \frac{(1-x)^2}{(-1+x^3)^{2/3}} - \frac{2^{2/3}(1-x)}{\sqrt[3]{-1+x^3}}\right)}{24\ 2^{2/3}} \\
&\quad + \frac{1}{3}\log\left(1+\sqrt[3]{-1+x^3}\right) + \frac{1}{8}\operatorname{Subst}\left(\int \frac{1}{2\sqrt[3]{2}-2^{2/3}x+x^2} dx, x, \frac{1-x}{\sqrt[3]{-1+x^3}}\right) - \frac{1}{4}\operatorname{Subst}\left(\int \frac{1}{2\sqrt[3]{2}-} \right. \\
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} + \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&\quad - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{\arctan\left(\frac{1-\frac{\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{4\ 2^{2/3}\sqrt{3}} - \frac{\arctan\left(\frac{1+\frac{2\sqrt[3]{2}(1-x)}{\sqrt[3]{-1+x^3}}}{\sqrt{3}}\right)}{2\ 2^{2/3}\sqrt{3}} \\
&\quad - \frac{2\arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} - \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{6(-1+x^3)^{2/3}} \\
&\quad + \frac{x(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&\quad + \frac{x^4(1-x^3)^{2/3}\operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&\quad + \frac{1}{3}\log\left(1+\sqrt[3]{-1+x^3}\right) + \frac{1}{3}\int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{1}{3}\int \frac{x^2\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{2}{3}\int \frac{\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2(-1+x^3)^{2/3}} + \frac{x^2}{2(-1+x^3)^{2/3}} - \frac{1}{3}\sqrt[3]{-1+x^3} - \frac{\sqrt[3]{-1+x^3}}{2(1+x^3)} \\
&+ \frac{x\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{x^2\sqrt[3]{-1+x^3}}{2(1+x^3)} - \frac{2 \arctan\left(\frac{1-2\sqrt[3]{-1+x^3}}{\sqrt{3}}\right)}{3\sqrt{3}} \\
&- \frac{x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{6(-1+x^3)^{2/3}} \\
&+ \frac{x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{5}{3}, \frac{4}{3}, x^3\right)}{3(-1+x^3)^{2/3}} \\
&+ \frac{x^4(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{4}{3}, \frac{5}{3}, \frac{7}{3}, x^3\right)}{6(-1+x^3)^{2/3}} - \frac{\log(x)}{3} \\
&+ \frac{1}{3} \log\left(1+\sqrt[3]{-1+x^3}\right) + \frac{1}{3} \int \frac{x\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{1}{3} \int \frac{x^2\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx + \frac{2}{3} \int \frac{\sqrt[3]{-1+x^3}}{-3-2x+x^2+x^3} dx
\end{aligned}$$

Mathematica [A] (verified)

Time = 5.41 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx &= \frac{\sqrt[3]{-1+x^3}}{-1+x^2} - \frac{\arctan\left(\frac{\sqrt{3}\sqrt[3]{-1+x^3}}{-2+2x^2+\sqrt[3]{-1+x^3}}\right)}{\sqrt{3}} \\
&+ \frac{1}{3} \log\left(1-x^2+\sqrt[3]{-1+x^3}\right) - \frac{1}{6} \log\left(1-2x^2+x^4+(-1+x^2)\sqrt[3]{-1+x^3}+(-1+x^3)^{2/3}\right)
\end{aligned}$$

[In] Integrate[((2 + x + x^2)*(-1 + x^3)^(1/3))/(x*(-1 + x^2)^2*(-3 - 2*x + x^2 + x^3)), x]

[Out] (-1 + x^3)^(1/3)/(-1 + x^2) - ArcTan[(Sqrt[3]*(-1 + x^3)^(1/3))/(-2 + 2*x^2 + (-1 + x^3)^(1/3))]/Sqrt[3] + Log[1 - x^2 + (-1 + x^3)^(1/3)]/3 - Log[1 - 2*x^2 + x^4 + (-1 + x^2)*(-1 + x^3)^(1/3) + (-1 + x^3)^(2/3)]/6

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 6.46 (sec) , antiderivative size = 1329, normalized size of antiderivative = 11.17

method	result	size
trager	Expression too large to display	1329
risch	Expression too large to display	1795

[In] int((x^2+x+2)*(x^3-1)^(1/3)/x/(x^2-1)^2/(x^3+x^2-2*x-3),x,method=_RETURNVER
BOSE)

[Out] (x^3-1)^(1/3)/(x^2-1)+3/2*RootOf(81*_Z^2+18*_Z+4)*ln(-(33318290353802192*x+
33719421708885792*(x^3-1)^(1/3)*x^3+33719421708885792*x^2*(x^3-1)^(1/3)+147
47439992666544*x^4-29494879985333088*x^3-33719421708885792*(x^3-1)^(1/3)-10
924029624197440*x^2+33719421708885792*(x^3-1)^(2/3)+45457696558242270*RootO
f(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x^3+45457696558242270*RootOf(81*_Z^2+18*_Z
+4)*(x^3-1)^(2/3)*x+45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)
*x^2-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x+147474399926
66544*x^5-19642219640243838*RootOf(81*_Z^2+18*_Z+4)*x^5-19642219640243838*R
ootOf(81*_Z^2+18*_Z+4)*x^4+39284439280487676*RootOf(81*_Z^2+18*_Z+4)*x^3+10
2579264381097584*RootOf(81*_Z^2+18*_Z+4)*x^2+43652605460366070*RootOf(81*_Z
^2+18*_Z+4)*x+2843018479155015*RootOf(81*_Z^2+18*_Z+4)^2*x^5-81229099404429
00*RootOf(81*_Z^2+18*_Z+4)^2+43652605460366070*RootOf(81*_Z^2+18*_Z+4)+3371
9421708885792*x*(x^3-1)^(2/3)-33719421708885792*(x^3-1)^(1/3)*x+28430184791
55015*RootOf(81*_Z^2+18*_Z+4)^2*x^4-5686036958310030*RootOf(81*_Z^2+18*_Z+4
)^2*x^3-16651965377907945*RootOf(81*_Z^2+18*_Z+4)^2*x^2-8122909940442900*Ro
otOf(81*_Z^2+18*_Z+4)^2*x+45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)
^(2/3)-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)+333182903538
02192)/(x^3+x^2-2*x-3)/x^2)-1/3*ln(-(23216580007526132*x+23617711362609732*
(x^3-1)^(1/3)*x^3+23617711362609732*x^2*(x^3-1)^(1/3)+19252773664777768*x^4
-38505547329555536*x^3-23617711362609732*(x^3-1)^(1/3)-34541740986807172*x^
2+23617711362609732*(x^3-1)^(2/3)-45457696558242270*RootOf(81*_Z^2+18*_Z+4)
*(x^3-1)^(1/3)*x^3-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(2/3)*
x-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x^2+4545769655824
2270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x+19252773664777768*x^5+20905783
408757178*RootOf(81*_Z^2+18*_Z+4)*x^5+20905783408757178*RootOf(81*_Z^2+18*_
Z+4)*x^4-41811566817514356*RootOf(81*_Z^2+18*_Z+4)*x^3-109980137882390004*R
ootOf(81*_Z^2+18*_Z+4)*x^2-47262787656118470*RootOf(81*_Z^2+18*_Z+4)*x+2843
018479155015*RootOf(81*_Z^2+18*_Z+4)^2*x^5-8122909940442900*RootOf(81*_Z^2+
18*_Z+4)^2-47262787656118470*RootOf(81*_Z^2+18*_Z+4)+23617711362609732*x*(x
^3-1)^(2/3)-23617711362609732*(x^3-1)^(1/3)*x+2843018479155015*RootOf(81*_Z
^2+18*_Z+4)^2*x^4-5686036958310030*RootOf(81*_Z^2+18*_Z+4)^2*x^3-1665196537
7907945*RootOf(81*_Z^2+18*_Z+4)^2*x^2-8122909940442900*RootOf(81*_Z^2+18*_Z
+4)^2*x-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(2/3)+45457696558
242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)+23216580007526132)/(x^3+x^2-2*
x-3)/x^2)-3/2*ln(-(23216580007526132*x+23617711362609732*(x^3-1)^(1/3)*x^3+
23617711362609732*x^2*(x^3-1)^(1/3)+19252773664777768*x^4-38505547329555536
x^3-23617711362609732(x^3-1)^(1/3)-34541740986807172*x^2+2361771136260973
2*(x^3-1)^(2/3)-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x^3
-45457696558242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(2/3)*x-4545769655824227

```

0*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(1/3)*x^2+45457696558242270*RootOf(81*_Z^
2+18*_Z+4)*(x^3-1)^(1/3)*x+19252773664777768*x^5+20905783408757178*RootOf(8
1*_Z^2+18*_Z+4)*x^5+20905783408757178*RootOf(81*_Z^2+18*_Z+4)*x^4-418115668
17514356*RootOf(81*_Z^2+18*_Z+4)*x^3-109980137882390004*RootOf(81*_Z^2+18*_
Z+4)*x^2-47262787656118470*RootOf(81*_Z^2+18*_Z+4)*x+2843018479155015*RootO
f(81*_Z^2+18*_Z+4)^2*x^5-8122909940442900*RootOf(81*_Z^2+18*_Z+4)^2-4726278
7656118470*RootOf(81*_Z^2+18*_Z+4)+23617711362609732*x*(x^3-1)^(2/3)-236177
11362609732*(x^3-1)^(1/3)*x+2843018479155015*RootOf(81*_Z^2+18*_Z+4)^2*x^4-
5686036958310030*RootOf(81*_Z^2+18*_Z+4)^2*x^3-16651965377907945*RootOf(81*
_Z^2+18*_Z+4)^2*x^2-8122909940442900*RootOf(81*_Z^2+18*_Z+4)^2*x-4545769655
8242270*RootOf(81*_Z^2+18*_Z+4)*(x^3-1)^(2/3)+45457696558242270*RootOf(81*_
Z^2+18*_Z+4)*(x^3-1)^(1/3)+23216580007526132)/(x^3+x^2-2*x-3)/x^2)*RootOf(8
1*_Z^2+18*_Z+4)

```

Fricas [F(-2)]

Exception generated.

$$\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \text{Exception raised: TypeError}$$

```

[In] integrate((x^2+x+2)*(x^3-1)^(1/3)/x/(x^2-1)^2/(x^3+x^2-2*x-3),x, algorithm=
"fricas")

```

```

[Out] Exception raised: TypeError >> Error detected within library code: integ
rate: implementation incomplete (residue poly has multiple non-linear facto
rs)

```

Sympy [F(-1)]

Timed out.

$$\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \text{Timed out}$$

```

[In] integrate((x**2+x+2)*(x**3-1)**(1/3)/x/(x**2-1)**2/(x**3+x**2-2*x-3),x)

```

```

[Out] Timed out

```

Maxima [F]

$$\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \int \frac{(x^3-1)^{\frac{1}{3}}(x^2+x+2)}{(x^3+x^2-2x-3)(x^2-1)^2x} dx$$

[In] integrate((x^2+x+2)*(x^3-1)^(1/3)/x/(x^2-1)^2/(x^3+x^2-2*x-3),x, algorithm="maxima")

[Out] integrate((x^3 - 1)^(1/3)*(x^2 + x + 2)/((x^3 + x^2 - 2*x - 3)*(x^2 - 1)^2*x), x)

Giac [F]

$$\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \int \frac{(x^3-1)^{\frac{1}{3}}(x^2+x+2)}{(x^3+x^2-2x-3)(x^2-1)^2x} dx$$

[In] integrate((x^2+x+2)*(x^3-1)^(1/3)/x/(x^2-1)^2/(x^3+x^2-2*x-3),x, algorithm="giac")

[Out] integrate((x^3 - 1)^(1/3)*(x^2 + x + 2)/((x^3 + x^2 - 2*x - 3)*(x^2 - 1)^2*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(2+x+x^2)\sqrt[3]{-1+x^3}}{x(-1+x^2)^2(-3-2x+x^2+x^3)} dx = \int -\frac{(x^3-1)^{1/3}(x^2+x+2)}{x(x^2-1)^2(-x^3-x^2+2x+3)} dx$$

[In] int(-((x^3 - 1)^(1/3)*(x + x^2 + 2))/(x*(x^2 - 1)^2*(2*x - x^2 - x^3 + 3)), x)

[Out] int(-((x^3 - 1)^(1/3)*(x + x^2 + 2))/(x*(x^2 - 1)^2*(2*x - x^2 - x^3 + 3)), x)

$$3.10 \quad \int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx$$

Optimal result	94
Rubi [C] (verified)	95
Mathematica [A] (verified)	99
Maple [C] (verified)	99
Fricas [A] (verification not implemented)	100
Sympy [F]	100
Maxima [F]	100
Giac [F]	101
Mupad [F(-1)]	101

Optimal result

Integrand size = 24, antiderivative size = 102

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx = -\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-x+x^3}}{-2+2x+\sqrt[3]{-x+x^3}}\right) - \log\left(1-x+\sqrt[3]{-x+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-x+x^3}+(-x+x^3)^{2/3}\right)$$

[Out] $-3^{(1/2)}*\arctan(3^{(1/2)}*(x^3-x)^{(1/3)/(-2+2*x+(x^3-x)^{(1/3))})-\ln(1-x+(x^3-x)^{(1/3))}+1/2*\ln(1-2*x+x^2+(-1+x)*(x^3-x)^{(1/3)}+(x^3-x)^{(2/3)})$

Rubi [C] (verified)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 0.48 (sec) , antiderivative size = 298, normalized size of antiderivative = 2.92, number of steps used = 12, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {2081, 6865, 6857, 281, 245, 1576, 476, 384, 525, 524}

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx = -\frac{18\sqrt[3]{1-x^2}x^2 \operatorname{AppellF1}\left(\frac{5}{6}, \frac{1}{3}, 1, \frac{11}{6}, x^2, 9x^2\right)}{5\sqrt[3]{x^3-x}} + \frac{\sqrt{3}\sqrt[3]{x^2-1}\sqrt[3]{x} \arctan\left(\frac{1-\frac{4x^{2/3}}{\sqrt[3]{x^2-1}}}{\sqrt{3}}\right)}{2\sqrt[3]{x^3-x}} + \frac{\sqrt{3}\sqrt[3]{x^2-1}\sqrt[3]{x} \arctan\left(\frac{\frac{2x^{2/3}}{\sqrt[3]{x^2-1}}+1}{\sqrt{3}}\right)}{2\sqrt[3]{x^3-x}} + \frac{\sqrt[3]{x^2-1}\sqrt[3]{x} \log(1-9x^2)}{4\sqrt[3]{x^3-x}} - \frac{3\sqrt[3]{x^2-1}\sqrt[3]{x} \log\left(x^{2/3}-\sqrt[3]{x^2-1}\right)}{4\sqrt[3]{x^3-x}} - \frac{3\sqrt[3]{x^2-1}\sqrt[3]{x} \log\left(2x^{2/3}+\sqrt[3]{x^2-1}\right)}{4\sqrt[3]{x^3-x}}$$

[In] Int[(1 + 3*x)/((-1 + 3*x)*(-x + x^3)^(1/3)), x]

[Out] (-18*x^2*(1 - x^2)^(1/3)*AppellF1[5/6, 1/3, 1, 11/6, x^2, 9*x^2])/(5*(-x + x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1 + x^2)^(1/3)*ArcTan[(1 - (4*x^(2/3)))/(-1 + x^2)^(1/3)]/Sqrt[3])/(2*(-x + x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1 + x^2)^(1/3)*ArcTan[(1 + (2*x^(2/3)))/(-1 + x^2)^(1/3)]/Sqrt[3])/(2*(-x + x^3)^(1/3)) + (x^(1/3)*(-1 + x^2)^(1/3)*Log[1 - 9*x^2])/(4*(-x + x^3)^(1/3)) - (3*x^(1/3)*(-1 + x^2)^(1/3)*Log[x^(2/3) - (-1 + x^2)^(1/3)])/(4*(-x + x^3)^(1/3)) - (3*x^(1/3)*(-1 + x^2)^(1/3)*Log[2*x^(2/3) + (-1 + x^2)^(1/3)])/(4*(-x + x^3)^(1/3))

Rule 245

Int[((a_) + (b_.)*(x_)^3)^(-1/3), x_Symbol] := Simp[ArcTan[(1 + 2*Rt[b, 3]*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*Rt[b, 3]), x] - Simp[Log[(a + b*x^3)^(1/3) - Rt[b, 3]*x]/(2*Rt[b, 3]), x] /; FreeQ[{a, b}, x]

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 384

```
Int[1/(((a_) + (b_.)*(x_)^3)^(1/3)*((c_) + (d_.)*(x_)^3)), x_Symbol] := Wit
h[{q = Rt[(b*c - a*d)/c, 3]}, Simp[ArcTan[(1 + (2*q*x)/(a + b*x^3)^(1/3))/S
qrt[3]]/(Sqrt[3]*c*q), x] + (-Simp[Log[q*x - (a + b*x^3)^(1/3)]/(2*c*q), x]
+ Simp[Log[c + d*x^3]/(6*c*q), x])] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c -
a*d, 0]
```

Rule 476

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_),
x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k -
1)*(a + b*x^(n/k))^p*(c + d*x^(n/k))^q, x], x, x^k], x] /; k != 1] /; FreeQ
[{a, b, c, d, p, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 524

```
Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_
))^(q_), x_Symbol] := Simp[a^p*c^q*((e*x)^(m + 1)/(e*(m + 1)))*AppellF1[(m
+ 1)/n, -p, -q, 1 + (m + 1)/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a,
b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] && NeQ[m, n
- 1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])
```

Rule 525

```
Int[((e_.)*(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_
))^(q_), x_Symbol] := Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^
n/a)^FracPart[p]), Int[(e*x)^m*(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /;
FreeQ[{a, b, c, d, e, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[m, -1] &&
NeQ[m, n - 1] && !(IntegerQ[p] || GtQ[a, 0])
```

Rule 1576

```
Int[((f_.)*(x_)^(m_.)*((d_) + (e_.)*(x_)^(n_))^(q_)*((a_) + (c_.)*(x_)^(n2
_))^(p_), x_Symbol] := Dist[(f*x)^m/x^m, Int[ExpandIntegrand[x^m*(a + c*x^(
2*n))^p, (d/(d^2 - e^2*x^(2*n)) - e*(x^n/(d^2 - e^2*x^(2*n))))^(-q), x], x]
, x] /; FreeQ[{a, c, d, e, f, m, n, p}, x] && EqQ[n2, 2*n] && !IntegerQ[p]
&& ILtQ[q, 0]
```


Rule 2081

Int[(u_)*(P_)^(p_), x_Symbol] := With[{m = MinimumMonomialExponent[P, x]}, Dist[P^FracPart[p]/(x^(m*FracPart[p]))*Distrib[1/x^m, P]^FracPart[p]], Int[u*x^(m*p)*Distrib[1/x^m, P]^p, x], x] /; FreeQ[p, x] && !IntegerQ[p] && SumQ[P] && EveryQ[BinomialQ[#1, x] & , P] && !PolyQ[P, x, 2]

Rule 6857

Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rule 6865

Int[(u_)*(x_)^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k, Subst[Int[x^(k*(m + 1) - 1)*(u /. x -> x^k), x], x, x^(1/k)], x] /; FractionQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \int \frac{1+3x}{\sqrt[3]{x(-1+3x)}\sqrt[3]{-1+x^2}} dx}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x(1+3x^3)}{(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(\frac{x}{\sqrt[3]{-1+x^6}} + \frac{2x}{(-1+3x^3)\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x}{\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
 &\quad + \frac{\left(6\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x}{(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{-1+x^3}} dx, x, x^{2/3}\right)}{2\sqrt[3]{-x+x^3}} \\
 &\quad + \frac{\left(6\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(\frac{x}{\sqrt[3]{-1+x^6(-1+9x^6)}} + \frac{3x^4}{\sqrt[3]{-1+x^6(-1+9x^6)}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
 \end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
= & \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
& + \frac{\left(6\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x}{\sqrt[3]{-1+x^6(-1+9x^6)}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
& + \frac{\left(18\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x^4}{\sqrt[3]{-1+x^6(-1+9x^6)}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
& \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
= & \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
& + \frac{\left(18\sqrt[3]{x}\sqrt[3]{1-x^2}\right) \text{Subst}\left(\int \frac{x^4}{\sqrt[3]{1-x^6(-1+9x^6)}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
& + \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{-1+x^3(-1+9x^3)}} dx, x, x^{2/3}\right)}{\sqrt[3]{-x+x^3}} \\
= & -\frac{18x^2\sqrt[3]{1-x^2} \text{AppellF1}\left(\frac{5}{6}, \frac{1}{3}, 1, \frac{11}{6}, x^2, 9x^2\right)}{5\sqrt[3]{-x+x^3}} + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1-\frac{4x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
& + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} + \frac{\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(1-9x^2)}{4\sqrt[3]{-x+x^3}} \\
& - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(2x^{2/3}+\sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 15.22 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.00

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx = -\sqrt{3} \arctan\left(\frac{\sqrt{3}\sqrt[3]{-x+x^3}}{-2+2x+\sqrt[3]{-x+x^3}}\right) - \log\left(1-x+\sqrt[3]{-x+x^3}\right) + \frac{1}{2} \log\left(1-2x+x^2+(-1+x)\sqrt[3]{-x+x^3}+(-x+x^3)^{2/3}\right)$$

[In] Integrate[(1 + 3*x)/((-1 + 3*x)*(-x + x^3)^(1/3)), x]

[Out] -(Sqrt[3]*ArcTan[(Sqrt[3]*(-x + x^3)^(1/3))/(-2 + 2*x + (-x + x^3)^(1/3))]) - Log[1 - x + (-x + x^3)^(1/3)] + Log[1 - 2*x + x^2 + (-1 + x)*(-x + x^3)^(1/3) + (-x + x^3)^(2/3)]/2

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.81 (sec) , antiderivative size = 607, normalized size of antiderivative = 5.95

method	result
trager	RootOf(_Z^2 - _Z + 1) ln $\left(\frac{-1415 \text{RootOf}(_Z^2 - _Z + 1)^2 x^2 + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{2}{3}} + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} + 3679 \text{RootOf}(_Z^2 - _Z + 1) x + 4649 \text{RootOf}(_Z^2 - _Z + 1) x^2 - 1107 (x^3 - x)^{\frac{2}{3}} - 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} - 1107 (x^3 - x)^{\frac{1}{3}} x - 1698 \text{RootOf}(_Z^2 - _Z + 1)^2 - 4786 \text{RootOf}(_Z^2 - _Z + 1) x - 2522 x^2 + 1107 (x^3 - x)^{\frac{1}{3}} + 3145 \text{RootOf}(_Z^2 - _Z + 1) + 1552 x - 1358}{(-1 + 3x)} - \ln(-1415 \text{RootOf}(_Z^2 - _Z + 1)^2 x^2 + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{2}{3}} + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} + 3679 \text{RootOf}(_Z^2 - _Z + 1) x + 4649 \text{RootOf}(_Z^2 - _Z + 1) x^2 - 1107 (x^3 - x)^{\frac{2}{3}} - 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} - 1107 (x^3 - x)^{\frac{1}{3}} x - 1698 \text{RootOf}(_Z^2 - _Z + 1)^2 + 2572 \text{RootOf}(_Z^2 - _Z + 1) x - 712 x^2 + 2127 (x^3 - x)^{\frac{1}{3}} - 251 \text{RootOf}(_Z^2 - _Z + 1) - 445 x - 89} / (-1 + 3x) \right) * \text{RootOf}(_Z^2 - _Z + 1) + \ln(-1415 \text{RootOf}(_Z^2 - _Z + 1)^2 x^2 + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{2}{3}} + 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} + 3679 \text{RootOf}(_Z^2 - _Z + 1) x + 4649 \text{RootOf}(_Z^2 - _Z + 1) x^2 - 1107 (x^3 - x)^{\frac{2}{3}} - 3234 \text{RootOf}(_Z^2 - _Z + 1) (x^3 - x)^{\frac{1}{3}} - 1107 (x^3 - x)^{\frac{1}{3}} x - 1698 \text{RootOf}(_Z^2 - _Z + 1)^2 + 2572 \text{RootOf}(_Z^2 - _Z + 1) x - 712 x^2 + 2127 (x^3 - x)^{\frac{1}{3}} - 251 \text{RootOf}(_Z^2 - _Z + 1) - 445 x - 89} / (-1 + 3x) \right)$

[In] int((1+3*x)/(-1+3*x)/(x^3-x)^(1/3), x, method=_RETURNVERBOSE)

[Out] RootOf(_Z^2 - _Z + 1) * ln((-1415*RootOf(_Z^2 - _Z + 1)^2*x^2+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)*x+3679*RootOf(_Z^2 - _Z + 1)^2*x+4649*RootOf(_Z^2 - _Z + 1)*x^2-1107*(x^3-x)^(2/3)-3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)-1107*(x^3-x)^(1/3)*x-1698*RootOf(_Z^2 - _Z + 1)^2-4786*RootOf(_Z^2 - _Z + 1)*x-2522*x^2+1107*(x^3-x)^(1/3)+3145*RootOf(_Z^2 - _Z + 1)+1552*x-1358)/(-1+3*x)) - ln((-1415*RootOf(_Z^2 - _Z + 1)^2*x^2+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)*x-3679*RootOf(_Z^2 - _Z + 1)^2*x+1819*RootOf(_Z^2 - _Z + 1)*x^2-2127*(x^3-x)^(2/3)-3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)-2127*(x^3-x)^(1/3)*x+1698*RootOf(_Z^2 - _Z + 1)^2+2572*RootOf(_Z^2 - _Z + 1)*x-712*x^2+2127*(x^3-x)^(1/3)-251*RootOf(_Z^2 - _Z + 1)-445*x-89)/(-1+3*x)) * RootOf(_Z^2 - _Z + 1) + ln((-1415*RootOf(_Z^2 - _Z + 1)^2*x^2+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)*x-3679*RootOf(_Z^2 - _Z + 1)^2*x+1819*RootOf(_Z^2 - _Z + 1)*x^2-2127*(x^3-x)^(2/3)-3234*RootOf(_Z^2 - _Z + 1)*(x^3-x)^(1/3)-2127*(x^3-x)^(1/3)*x+1698*RootOf(_Z^2 - _Z + 1)^2+2572*RootOf(_Z^2 - _Z + 1)*x-712*x^2+2127*(x^3-x)^(1/3)-251*RootOf(_Z^2 - _Z + 1)-445*x-89)/(-1+3*x))

Fricas [A] (verification not implemented)

none

Time = 0.65 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.04

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx$$

$$= \sqrt{3} \arctan \left(\frac{612314840 \sqrt{3}(x^3-x)^{\frac{1}{3}}(x-1) + \sqrt{3}(1609127381x^2 + 1235276981x + 124616800) + 2605939922\sqrt{3}(x^3-x)^{\frac{2}{3}}}{2990437623x^2 + 3108349623x - 39304000} \right)$$

$$- \frac{1}{2} \log \left(\frac{3(x^3-x)^{\frac{1}{3}}(x-1) + 3x - 3(x^3-x)^{\frac{2}{3}} - 1}{3x-1} \right)$$

```
[In] integrate((1+3*x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="fricas")
```

```
[Out] sqrt(3)*arctan((612314840*sqrt(3)*(x^3 - x)^(1/3)*(x - 1) + sqrt(3)*(1609127381*x^2 + 1235276981*x + 124616800) + 2605939922*sqrt(3)*(x^3 - x)^(2/3))/(2990437623*x^2 + 3108349623*x - 39304000)) - 1/2*log((3*(x^3 - x)^(1/3)*(x - 1) + 3*x - 3*(x^3 - x)^(2/3) - 1)/(3*x - 1))
```

Sympy [F]

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{3x+1}{\sqrt[3]{x(x-1)(x+1)}(3x-1)} dx$$

```
[In] integrate((1+3*x)/(-1+3*x)/(x**3-x)**(1/3),x)
```

```
[Out] Integral((3*x + 1)/((x*(x - 1)*(x + 1))**(1/3)*(3*x - 1)), x)
```

Maxima [F]

$$\int \frac{1+3x}{(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{3x+1}{(x^3-x)^{\frac{1}{3}}(3x-1)} dx$$

```
[In] integrate((1+3*x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="maxima")
```

```
[Out] integrate((3*x + 1)/((x^3 - x)^(1/3)*(3*x - 1)), x)
```

Giac [F]

$$\int \frac{1 + 3x}{(-1 + 3x)\sqrt[3]{-x + x^3}} dx = \int \frac{3x + 1}{(x^3 - x)^{\frac{1}{3}}(3x - 1)} dx$$

[In] integrate((1+3*x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="giac")

[Out] integrate((3*x + 1)/((x^3 - x)^(1/3)*(3*x - 1)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1 + 3x}{(-1 + 3x)\sqrt[3]{-x + x^3}} dx = \int \frac{3x + 1}{(x^3 - x)^{1/3} (3x - 1)} dx$$

[In] int((3*x + 1)/((x^3 - x)^(1/3)*(3*x - 1)),x)

[Out] int((3*x + 1)/((x^3 - x)^(1/3)*(3*x - 1)), x)

$$3.11 \quad \int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx$$

Optimal result	102
Rubi [C] (warning: unable to verify)	102
Mathematica [F]	108
Maple [C] (verified)	109
Fricas [A] (verification not implemented)	109
Sympy [F]	110
Maxima [F]	110
Giac [F]	110
Mupad [F(-1)]	111

Optimal result

Integrand size = 27, antiderivative size = 125

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx =$$

$$-\sqrt{3} \arctan \left(\frac{\sqrt{3}(124616800 + 1235276981x + 1609127381x^2) + 612314840\sqrt{3}(-1+x)\sqrt[3]{-x+x^3} + 2605939922\sqrt{3}}{-39304000 + 3108349623x + 2990437623x^2} \right)$$

$$-\frac{1}{2} \log \left(\frac{-1+3x+3(-1+x)\sqrt[3]{-x+x^3}-3(-x+x^3)^{2/3}}{-1+3x} \right)$$

```
[Out] -3^(1/2)*arctan((612314840*3^(1/2)*(x^3-x)^(1/3)*(-1+x)+3^(1/2)*(1609127381
*x^2+1235276981*x+124616800)+2605939922*3^(1/2)*(x^3-x)^(2/3))/(2990437623*
x^2+3108349623*x-39304000))-1/2*ln((3*(-1+x)*(x^3-x)^(1/3)+3*x-3*(x^3-x)^(2
/3)-1)/(-1+3*x))
```

Rubi [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 3 in optimal.

Time = 1.01 (sec) , antiderivative size = 537, normalized size of antiderivative = 4.30,
 number of steps used = 14, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules

used = {2081, 6865, 6857, 247, 231, 281, 337, 1452, 441, 440, 476, 503}

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \frac{4x(1-x^2)^{2/3} \text{AppellF1}\left(\frac{1}{6}, \frac{2}{3}, 1, \frac{7}{6}, x^2, 9x^2\right)}{(x^3-x)^{2/3}}$$

$$x(1-x^2) \left(1 - \frac{x^{2/3}}{\sqrt[3]{x^2-1}}\right) \sqrt{\frac{\frac{x^{4/3}}{(x^2-1)^{2/3}} + \frac{x^{2/3}}{\sqrt[3]{x^2-1}} + 1}{\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{x^2-1}}\right)^2}} \text{EllipticF}\left(\arccos\left(\frac{1 - \frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{x^2-1}}}{1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{x^2-1}}}\right), \frac{1}{4}(2 + \sqrt{3})\right)$$

$$2\sqrt[4]{3}(x^3-x)^{2/3} \sqrt{-\frac{x^{2/3}\left(1 - \frac{x^{2/3}}{\sqrt[3]{x^2-1}}\right)}{\sqrt[3]{x^2-1}\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{x^2-1}}\right)^2}}$$

$$\frac{\sqrt{3}x^{2/3}(x^2-1)^{2/3} \arctan\left(\frac{1 - \frac{4x^{2/3}}{\sqrt[3]{x^2-1}}}{\sqrt{3}}\right)}{2(x^3-x)^{2/3}}$$

$$\frac{\sqrt{3}x^{2/3}(x^2-1)^{2/3} \arctan\left(\frac{\frac{2x^{2/3}}{\sqrt[3]{x^2-1}} + 1}{\sqrt{3}}\right)}{2(x^3-x)^{2/3}} + \frac{x^{2/3}(x^2-1)^{2/3} \log(1-9x^2)}{4(x^3-x)^{2/3}}$$

$$\frac{3x^{2/3}(x^2-1)^{2/3} \log\left(x^{2/3} - \sqrt[3]{x^2-1}\right)}{4(x^3-x)^{2/3}} - \frac{3x^{2/3}(x^2-1)^{2/3} \log\left(2x^{2/3} + \sqrt[3]{x^2-1}\right)}{4(x^3-x)^{2/3}}$$

[In] Int[((-1 + x)*(1 + 3*x))/((-1 + 3*x)*(-x + x^3)^(2/3)),x]

[Out] (4*x*(1 - x^2)^(2/3)*AppellF1[1/6, 2/3, 1, 7/6, x^2, 9*x^2])/(-x + x^3)^(2/3) - (Sqrt[3]*x^(2/3)*(-1 + x^2)^(2/3)*ArcTan[(1 - (4*x^(2/3)))/(-1 + x^2)^(1/3)]/Sqrt[3])/(2*(-x + x^3)^(2/3)) - (Sqrt[3]*x^(2/3)*(-1 + x^2)^(2/3)*ArcTan[(1 + (2*x^(2/3)))/(-1 + x^2)^(1/3)]/Sqrt[3])/(2*(-x + x^3)^(2/3)) - (x*(1 - x^2)*(1 - x^(2/3))/(-1 + x^2)^(1/3))*Sqrt[(1 + x^(4/3))/(-1 + x^2)^(2/3) + x^(2/3)/(-1 + x^2)^(1/3)]/(1 - ((1 + Sqrt[3])*x^(2/3))/(-1 + x^2)^(1/3))^2]*EllipticF[ArcCos[(1 - ((1 - Sqrt[3])*x^(2/3))/(-1 + x^2)^(1/3))/(1 - ((1 + Sqrt[3])*x^(2/3))/(-1 + x^2)^(1/3))], (2 + Sqrt[3])/4)]/(2*3^(1/4)*(-x + x^3)^(2/3)*Sqrt[-((x^(2/3)*(1 - x^(2/3))/(-1 + x^2)^(1/3)))/((-1 + x^2)^(1/3)*(1 - ((1 + Sqrt[3])*x^(2/3))/(-1 + x^2)^(1/3))^2)]] + (x^(2/3)*(-1 + x^2)^(2/3)*Log[1 - 9*x^2])/(4*(-x + x^3)^(2/3)) - (3*x^(2/3)*(-1 + x^2)^(2/3)*Log[x^(2/3) - (-1 + x^2)^(1/3)])/(4*(-x + x^3)^(2/3)) - (3*x^(2/3)*(-1 + x^2)^(2/3)*Log[2*x^(2/3) + (-1 + x^2)^(1/3)])/(4*(-x + x^3)^(2/3))

Rule 231

Int[1/Sqrt[(a_) + (b_.)*(x_)^6], x_Symbol] := With[{r = Numer[Rt[b/a, 3]], s = Denom[Rt[b/a, 3]]}, Simp[x*(s + r*x^2)*(Sqrt[(s^2 - r*s*x^2 + r^2*x^4)/

```
(s + (1 + Sqrt[3])*r*x^2)^2]/(2*3^(1/4)*s*Sqrt[a + b*x^6]*Sqrt[r*x^2*((s +
r*x^2)/(s + (1 + Sqrt[3])*r*x^2)^2]))*EllipticF[ArcCos[(s + (1 - Sqrt[3])*
r*x^2)/(s + (1 + Sqrt[3])*r*x^2)], (2 + Sqrt[3])/4], x]] /; FreeQ[{a, b}, x
]
```

Rule 247

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[(a/(a + b*x^n))^(p + 1
/n)*(a + b*x^n)^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/
(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] &
& NeQ[p, -2^(-1)] && LtQ[Denominator[p + 1/n], Denominator[p]]
```

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 337

```
Int[(x_)/((a_) + (b_.)*(x_)^3)^(2/3), x_Symbol] := With[{q = Rt[b, 3]}, Sim
p[-ArcTan[(1 + 2*q*(x/(a + b*x^3)^(1/3)))/Sqrt[3]]/(Sqrt[3]*q^2), x] - Simp
[Log[q*x - (a + b*x^3)^(1/3)]/(2*q^2), x]] /; FreeQ[{a, b}, x]
```

Rule 440

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)
], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])
```

Rule 441

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]),
Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}
, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])
```

Rule 476

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_),
x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k -
1)*(a + b*x^(n/k))^p*(c + d*x^(n/k))^q, x], x, x^k], x] /; k != 1] /; FreeQ
```


$\{a, b, c, d, p, q\}, x \} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[n, 0] \&\& \text{IntegerQ}[m]$

Rule 503

$\text{Int}[(x_)/(((a_)+(b_)*(x_)^3)^{(2/3)}*((c_)+(d_)*(x_)^3)), x_Symbol] \text{ :>}$
 $\text{With}[\{q = \text{Rt}[(b*c - a*d)/c, 3]\}, \text{Simp}[-\text{ArcTan}[(1 + (2*q*x)/(a + b*x^3))^{(1/3)}]/\text{Sqrt}[3]]/(\text{Sqrt}[3]*c*q^2), x] + (-\text{Simp}[\text{Log}[q*x - (a + b*x^3)^{(1/3)}]/(2*c*q^2), x] + \text{Simp}[\text{Log}[c + d*x^3]/(6*c*q^2), x]) /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0]$

Rule 1452

$\text{Int}[(d_)+(e_)*(x_)^{(n_)]^{(q_)}*((a_)+(c_)*(x_)^{(n2_)]^{(p_)}, x_Symbol] \text{ :>}$
 $\text{Int}[\text{ExpandIntegrand}[(a + c*x^{(2*n)})^p, (d/(d^2 - e^2*x^{(2*n)}) - e*(x^n/(d^2 - e^2*x^{(2*n)}))^{(-q)}, x], x] /; \text{FreeQ}[\{a, c, d, e, n, p\}, x] \&\& \text{EqQ}[n, 2, 2*n] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& !\text{IntegerQ}[p] \&\& \text{ILtQ}[q, 0]$

Rule 2081

$\text{Int}[(u_)*(P_)^{(p_)}, x_Symbol] \text{ :>}$ $\text{With}[\{m = \text{MinimumMonomialExponent}[P, x]\}, \text{Dist}[P^{\text{FracPart}[p]}/(x^{(m*\text{FracPart}[p])}* \text{Distrib}[1/x^m, P]^{\text{FracPart}[p]}), \text{Int}[u*x^{(m*p)}*\text{Distrib}[1/x^m, P]^p, x], x] /; \text{FreeQ}[p, x] \&\& !\text{IntegerQ}[p] \&\& \text{SumQ}[P] \&\& \text{EveryQ}[\text{BinomialQ}[\#1, x] \& , P] \&\& !\text{PolyQ}[P, x, 2]$

Rule 6857

$\text{Int}[(u_)/((a_)+(b_)*(x_)^{(n_)}), x_Symbol] \text{ :>}$ $\text{With}[\{v = \text{RationalFunctionExpand}[u/(a + b*x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v]] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{IGtQ}[n, 0]$

Rule 6865

$\text{Int}[(u_)*(x_)^{(m_)}, x_Symbol] \text{ :>}$ $\text{With}[\{k = \text{Denominator}[m]\}, \text{Dist}[k, \text{Subst}[\text{Int}[x^{(k*(m + 1) - 1)}*(u /. x \rightarrow x^k), x], x, x^{(1/k)}], x] /; \text{FractionQ}[m]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(x^{2/3}(-1+x^2)^{2/3}\right) \int \frac{(-1+x)(1+3x)}{x^{2/3}(-1+3x)(-1+x^2)^{2/3}} dx}{(-x+x^3)^{2/3}} \\ &= \frac{\left(3x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int \frac{(-1+x^3)(1+3x^3)}{(-1+3x^3)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \end{aligned}$$

$$\begin{aligned}
&= \frac{\left(3x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\left(-\frac{1}{3(-1+x^6)^{2/3}}+\frac{x^3}{(-1+x^6)^{2/3}}-\frac{4}{3(-1+3x^3)(-1+x^6)^{2/3}}\right)dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&= -\frac{\left(x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{1}{(-1+x^6)^{2/3}}dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&\quad +\frac{\left(3x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{x^3}{(-1+x^6)^{2/3}}dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&\quad -\frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{1}{(-1+3x^3)(-1+x^6)^{2/3}}dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&= -\frac{\left(x^{2/3}\sqrt[6]{-1+x^2}\right) \text{Subst}\left(\int\frac{1}{\sqrt{1-x^6}}dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}}(-x+x^3)^{2/3}} \\
&\quad +\frac{\left(3x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{x}{(-1+x^3)^{2/3}}dx, x, x^{2/3}\right)}{2(-x+x^3)^{2/3}} \\
&\quad -\frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\left(\frac{1}{(-1+x^6)^{2/3}(-1+9x^6)}+\frac{3x^3}{(-1+x^6)^{2/3}(-1+9x^6)}\right)dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&= -\frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
&\quad -\frac{x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)\sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \text{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right), \frac{1}{4}\right)}{2\sqrt[4]{3}(-x+x^3)^{2/3}\sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
&\quad -\frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
&\quad -\frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{1}{(-1+x^6)^{2/3}(-1+9x^6)}dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&\quad -\frac{\left(12x^{2/3}(-1+x^2)^{2/3}\right) \text{Subst}\left(\int\frac{x^3}{(-1+x^6)^{2/3}(-1+9x^6)}dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
& - \frac{x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \operatorname{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right), \frac{1}{4}\right)}{2^4\sqrt{3}(-x+x^3)^{2/3} \sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
& - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
& - \frac{\left(4x^{2/3}(1-x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1-x^6)^{2/3}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(6x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{x}{(-1+x^3)^{2/3}(-1+9x^3)} dx, x, x^{2/3}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4x(1-x^2)^{2/3} \operatorname{AppellF1}\left(\frac{1}{6}, \frac{2}{3}, 1, \frac{7}{6}, x^2, 9x^2\right)}{(-x+x^3)^{2/3}} \\
&\quad - \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1-\sqrt[3]{-1+x^2}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
&\quad - \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\sqrt[3]{-1+x^2}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
&\quad - \frac{x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \operatorname{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right), \frac{1}{4}\right)}{2^4\sqrt{3}(-x+x^3)^{2/3} \sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
&\quad + \frac{x^{2/3}(-1+x^2)^{2/3} \log(1-9x^2)}{4(-x+x^3)^{2/3}} - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
&\quad - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(2x^{2/3}+\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}}
\end{aligned}$$

Mathematica [F]

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx$$

[In] Integrate[((-1 + x)*(1 + 3*x))/((-1 + 3*x)*(-x + x^3)^(2/3)), x]

[Out] Integrate[((-1 + x)*(1 + 3*x))/((-1 + 3*x)*(-x + x^3)^(2/3)), x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.60 (sec) , antiderivative size = 606, normalized size of antiderivative = 4.85

method	result
trager	$\ln \left(-\frac{1415 \operatorname{RootOf}(_Z^2 - _Z + 1)^2 x^2 + 3234 \operatorname{RootOf}(_Z^2 - _Z + 1)(x^3 - x)^{\frac{2}{3}} + 3234 \operatorname{RootOf}(_Z^2 - _Z + 1)(x^3 - x)^{\frac{1}{3}} x - 3679}{\dots} \right)$

[In] `int((-1+x)*(1+3*x)/(-1+3*x)/(x^3-x)^(2/3),x,method=_RETURNVERBOSE)`

[Out] `ln(-(1415*RootOf(_Z^2-_Z+1)^2*x^2+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)*x-3679*RootOf(_Z^2-_Z+1)^2*x+1819*RootOf(_Z^2-_Z+1)*x^2-2127*(x^3-x)^(2/3)-3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)-2127*(x^3-x)^(1/3)*x+1698*RootOf(_Z^2-_Z+1)^2+2572*RootOf(_Z^2-_Z+1)*x-712*x^2+2127*(x^3-x)^(1/3)-251*RootOf(_Z^2-_Z+1)-445*x-89)/(-1+3*x))*RootOf(_Z^2-_Z+1)-RootOf(_Z^2-_Z+1)*ln((-1415*RootOf(_Z^2-_Z+1)^2*x^2+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)*x+3679*RootOf(_Z^2-_Z+1)^2*x+4649*RootOf(_Z^2-_Z+1)*x^2-1107*(x^3-x)^(2/3)-3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)-1107*(x^3-x)^(1/3)*x-1698*RootOf(_Z^2-_Z+1)^2-4786*RootOf(_Z^2-_Z+1)*x-2522*x^2+1107*(x^3-x)^(1/3)+3145*RootOf(_Z^2-_Z+1)+1552*x-1358)/(-1+3*x))+ln((-1415*RootOf(_Z^2-_Z+1)^2*x^2+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(2/3)+3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)*x+3679*RootOf(_Z^2-_Z+1)^2*x+4649*RootOf(_Z^2-_Z+1)*x^2-1107*(x^3-x)^(2/3)-3234*RootOf(_Z^2-_Z+1)*(x^3-x)^(1/3)-1107*(x^3-x)^(1/3)*x-1698*RootOf(_Z^2-_Z+1)^2-4786*RootOf(_Z^2-_Z+1)*x-2522*x^2+1107*(x^3-x)^(1/3)+3145*RootOf(_Z^2-_Z+1)+1552*x-1358)/(-1+3*x))`

Fricas [A] (verification not implemented)

none

Time = 0.50 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.86

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx =$$

$$-\sqrt{3} \arctan \left(\frac{612314840 \sqrt{3}(x^3-x)^{1/3}(x-1) + \sqrt{3}(1609127381 x^2 + 1235276981 x + 124616800) + 26059}{2990437623 x^2 + 3108349623 x - 39304000} \right)$$

$$-\frac{1}{2} \log \left(\frac{3(x^3-x)^{1/3}(x-1) + 3x - 3(x^3-x)^{2/3} - 1}{3x-1} \right)$$

[In] `integrate((-1+x)*(1+3*x)/(-1+3*x)/(x^3-x)^(2/3),x, algorithm="fricas")`

[Out] $-\sqrt{3} \arctan\left(\frac{612314840\sqrt{3}(x^3 - x)^{1/3}(x - 1) + \sqrt{3}(1609127381x^2 + 1235276981x + 124616800) + 2605939922\sqrt{3}(x^3 - x)^{2/3}}{(2990437623x^2 + 3108349623x - 39304000)}\right) - \frac{1}{2} \log\left(\frac{3(x^3 - x)^{1/3}(x - 1) + 3x - 3(x^3 - x)^{2/3} - 1}{3x - 1}\right)$

Sympy [F]

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(x-1)(3x+1)}{(x(x-1)(x+1))^{2/3} \cdot (3x-1)} dx$$

[In] `integrate((-1+x)*(1+3*x)/(-1+3*x)/(x**3-x)**(2/3), x)`

[Out] `Integral((x - 1)*(3*x + 1)/((x*(x - 1)*(x + 1))**(2/3)*(3*x - 1)), x)`

Maxima [F]

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(3x+1)(x-1)}{(x^3-x)^{2/3}(3x-1)} dx$$

[In] `integrate((-1+x)*(1+3*x)/(-1+3*x)/(x^3-x)^(2/3), x, algorithm="maxima")`

[Out] `integrate((3*x + 1)*(x - 1)/((x^3 - x)^(2/3)*(3*x - 1)), x)`

Giac [F]

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(3x+1)(x-1)}{(x^3-x)^{2/3}(3x-1)} dx$$

[In] `integrate((-1+x)*(1+3*x)/(-1+3*x)/(x^3-x)^(2/3), x, algorithm="giac")`

[Out] `integrate((3*x + 1)*(x - 1)/((x^3 - x)^(2/3)*(3*x - 1)), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(-1+x)(1+3x)}{(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(3x+1)(x-1)}{(x^3-x)^{2/3}(3x-1)} dx$$

```
[In] int(((3*x + 1)*(x - 1))/((x^3 - x)^(2/3)*(3*x - 1)), x)
```

```
[Out] int(((3*x + 1)*(x - 1))/((x^3 - x)^(2/3)*(3*x - 1)), x)
```

$$3.12 \quad \int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx$$

Optimal result	112
Rubi [F]	112
Mathematica [F]	118
Maple [C] (verified)	118
Fricas [A] (verification not implemented)	119
Sympy [F]	119
Maxima [F]	119
Giac [F]	120
Mupad [F(-1)]	120

Optimal result

Integrand size = 37, antiderivative size = 166

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx$$

$$= \frac{-3(-x+x^3)^{2/3} + 2\sqrt{3}(1-2x+x^2) \operatorname{Arctan}\left(\frac{\sqrt{3}(124616800+1235276981x+1609127381x^2)+612314840\sqrt{3}(-1+x)\sqrt[3]{-x+x^3}}{-39304000+3108349623x+2990437623x^2}\right)}{2(1-2x+x^2)}$$

```
[Out] 1/2*(2*3^(1/2)*(x^2+x)*arctan((612314840*3^(1/2)*(x^3-x)^(1/3)*(-1+x)+3^(1/2)*(1609127381*x^2+1235276981*x+124616800)+2605939922*3^(1/2)*(x^3-x)^(2/3))/(2990437623*x^2+3108349623*x-39304000))-(x^2+x)*ln((3*(-1+x)*(x^3-x)^(1/3)+3*x-3*(x^3-x)^(2/3)-1)/(-1+3*x))-6*(x^3-x)^(2/3))/(x^2+x)
```

Rubi [F]

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx$$

```
[In] Int[((-1 + x)^2*(1 + 3*x))/(x*(1 + x)*(-1 + 3*x)*(-x + x^3)^(1/3)),x]
```

```
[Out] 3/(-x + x^3)^(1/3) + (3*(1 + Sqrt[3])*x^(2/3))/((-1 + x^2)^(1/3)*(-x + x^3)^(1/3)*(1 - ((1 + Sqrt[3])*x^(2/3))/(-1 + x^2)^(1/3))) - (18*x^2*(1 - x^2)^(1/3)*AppellF1[5/6, 1/3, 1, 11/6, x^2, 9*x^2])/(5*(-x + x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1 + x^2)^(1/3)*ArcTan[(1 - (4*x^(2/3))/(-1 + x^2)^(1/3))/Sqrt[3]])/(2*(-x + x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1 + x^2)^(1/3)*ArcTan[(1 + (2*x^(2/3))/(-1 + x^2)^(1/3))/Sqrt[3]])/(2*(-x + x^3)^(1/3)) - (3*3^(1/4))*x
```


$$\begin{aligned} &^{(2/3)}*(1-x^2)*(1-x^{(2/3)})/(-1+x^2)^{(1/3)}*\text{Sqrt}[(1+x^{(4/3)})/(-1+x^2)^{(2/3)}+x^{(2/3)}/(-1+x^2)^{(1/3)}]/(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})^2]*\text{EllipticE}[\text{ArcCos}[(1-((1-\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})/(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})]],(2+\text{Sqrt}[3])/4]/((-1+x^2)^{(1/3)}*(-x+x^3)^{(1/3)}*\text{Sqrt}[-((x^{(2/3)}*(1-x^{(2/3)})/(-1+x^2)^{(1/3)}))/((-1+x^2)^{(1/3)}*(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})^2)]])-(3^{(3/4)}*(1-\text{Sqrt}[3])*x^{(2/3)}*(1-x^2)*(1-x^{(2/3)})/(-1+x^2)^{(1/3)}*\text{Sqrt}[(1+x^{(4/3)})/(-1+x^2)^{(2/3)}+x^{(2/3)}/(-1+x^2)^{(1/3)}]/(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})^2]*\text{EllipticF}[\text{ArcCos}[(1-((1-\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})/(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})]],(2+\text{Sqrt}[3])/4]/(2*(-1+x^2)^{(1/3)}*(-x+x^3)^{(1/3)}*\text{Sqrt}[-((x^{(2/3)}*(1-x^{(2/3)})/(-1+x^2)^{(1/3)}))/((-1+x^2)^{(1/3)}*(1-((1+\text{Sqrt}[3])*x^{(2/3)})/(-1+x^2)^{(1/3)})^2)]])+(x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Log}[1-9*x^2])/((4*(-x+x^3)^{(1/3)})-(3*x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Log}[x^{(2/3)}-(-1+x^2)^{(1/3)}])/(4*(-x+x^3)^{(1/3)})-(3*x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Log}[2*x^{(2/3)}+(-1+x^2)^{(1/3)}])/(4*(-x+x^3)^{(1/3)})+(2*x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Defer}[\text{Subst}][\text{Defer}[\text{Int}][1/((1+x)*(-1+x^6)^{(1/3)}),x],x,x^{(1/3)}])/(-x+x^3)^{(1/3)}-(2*(1-I*\text{Sqrt}[3])*x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Defer}[\text{Subst}][\text{Defer}[\text{Int}][1/((-1-I*\text{Sqrt}[3]+2*x)*(-1+x^6)^{(1/3)}),x],x,x^{(1/3)}])/(-x+x^3)^{(1/3)}-(2*(1+I*\text{Sqrt}[3])*x^{(1/3)}*(-1+x^2)^{(1/3)}*\text{Defer}[\text{Subst}][\text{Defer}[\text{Int}][1/((-1+I*\text{Sqrt}[3]+2*x)*(-1+x^6)^{(1/3)}),x],x,x^{(1/3)}])/(-x+x^3)^{(1/3)} \end{aligned}$$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \int \frac{(-1+x)^2(1+3x)}{x^{4/3}(1+x)(-1+3x)\sqrt[3]{-1+x^2}} dx}{\sqrt[3]{-x+x^3}} \\ &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{(-1+x^3)^2(1+3x^3)}{x^2(1+x^3)(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\ &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(-\frac{1}{x^2\sqrt[3]{-1+x^6}} + \frac{x}{\sqrt[3]{-1+x^6}} + \frac{2}{3(1+x)\sqrt[3]{-1+x^6}} - \frac{2(1+x)}{3(1-x+x^2)\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \end{aligned}$$

$$\begin{aligned}
&= \frac{(2\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{(1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&- \frac{(2\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1+x}{(1-x+x^2)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&- \frac{(3\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{x^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&+ \frac{(3\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&+ \frac{(6\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{x}{(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&= - \frac{(3\sqrt[3]{x}) \operatorname{Subst}\left(\int \frac{1}{x^2\sqrt{1-x^6}} dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}\sqrt[6]{-1+x^2}\sqrt[3]{-x+x^3}}} \\
&+ \frac{(3\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{\sqrt[3]{-1+x^3}} dx, x, x^{2/3}\right)}{2\sqrt[3]{-x+x^3}} \\
&+ \frac{(2\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{(1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&- \frac{(2\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \left(\frac{1-i\sqrt{3}}{(-1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} + \frac{1+i\sqrt{3}}{(-1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&+ \frac{(6\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \left(\frac{x}{\sqrt[3]{-1+x^6}(-1+9x^6)} + \frac{3x^4}{\sqrt[3]{-1+x^6}(-1+9x^6)}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3}{\sqrt[3]{-x+x^3}} + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3} - \sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
&\quad + \frac{(6\sqrt[3]{x}) \operatorname{Subst}\left(\int \frac{x^4}{\sqrt{1-x^6}} dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}} \sqrt[6]{-1+x^2} \sqrt[3]{-x+x^3}} \\
&\quad + \frac{(2\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{(1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad + \frac{(6\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{x}{\sqrt[3]{-1+x^6}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad + \frac{(18\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{x^4}{\sqrt[3]{-1+x^6}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{(2(1-i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{(-1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{(2(1+i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{(-1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3}{\sqrt[3]{-x+x^3}} + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3} - \sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(18\sqrt[3]{x}\sqrt[3]{1-x^2}\right) \text{Subst}\left(\int \frac{x^4}{\sqrt[3]{1-x^6}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(3\sqrt[3]{x}\right) \text{Subst}\left(\int \frac{-1+\sqrt{3}-2x^4}{\sqrt{1-x^6}} dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}}\sqrt[6]{-1+x^2}\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(3(-1+\sqrt{3})\sqrt[3]{x}\right) \text{Subst}\left(\int \frac{1}{\sqrt{1-x^6}} dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}}\sqrt[6]{-1+x^2}\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(2\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{-1+x^3}(-1+9x^3)} dx, x, x^{2/3}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(2(1-i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(2(1+i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3}{\sqrt[3]{-x+x^3}} + \frac{3(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}\sqrt[3]{-x+x^3} \left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)} \\
&\quad - \frac{18x^2\sqrt[3]{1-x^2} \operatorname{AppellF1}\left(\frac{5}{6}, \frac{1}{3}, 1, \frac{11}{6}, x^2, 9x^2\right)}{5\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1 - \frac{4x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1 + \frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad - \frac{3^4\sqrt{3}x^{2/3}(1-x^2) \left(1 - \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1 + \frac{x^{4/3}}{(-1+x^2)^{2/3}} + \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} E\left(\arccos\left(\frac{1 - \frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right)\right)}{\frac{1}{4}} \\
&\quad - \frac{\sqrt[3]{-1+x^2}\sqrt[3]{-x+x^3} \sqrt{-\frac{x^{2/3}\left(1 - \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}}{3^{3/4}(1-\sqrt{3})x^{2/3}(1-x^2) \left(1 - \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1 + \frac{x^{4/3}}{(-1+x^2)^{2/3}} + \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \operatorname{EllipticF}\left(\arccos\left(\frac{1}{1}\right)\right)} \\
&\quad - \frac{2\sqrt[3]{-1+x^2}\sqrt[3]{-x+x^3} \sqrt{-\frac{x^{2/3}\left(1 - \frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1 - \frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}}{\frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(1-9x^2)}{4\sqrt[3]{-x+x^3}} - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(x^{2/3} - \sqrt[3]{-1+x^2})}{4\sqrt[3]{-x+x^3}}} \\
&\quad - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(2x^{2/3} + \sqrt[3]{-1+x^2})}{4\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(2\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(2(1-i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(2(1+i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
\end{aligned}$$

Mathematica [F]

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx$$

[In] Integrate[((-1 + x)^2*(1 + 3*x))/(x*(1 + x)*(-1 + 3*x)*(-x + x^3)^(1/3)),x]

[Out] Integrate[((-1 + x)^2*(1 + 3*x))/(x*(1 + x)*(-1 + 3*x)*(-x + x^3)^(1/3)), x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.70 (sec) , antiderivative size = 461, normalized size of antiderivative = 2.78

method	result
trager	$-\frac{3(x^3-x)^{\frac{2}{3}}}{(1+x)x} - \ln\left(\frac{11314080 \operatorname{RootOf}(11664_Z^2-108_Z+1)^2 x^2+349272 \operatorname{RootOf}(11664_Z^2-108_Z+1)(x^3-x)^{\frac{2}{3}}-229716}{\dots}\right)$
risch	$-\frac{3(-1+x)}{(x(x^2-1))^{\frac{1}{3}}} + \operatorname{RootOf}(_Z^2 - _Z + 1) \ln\left(\frac{-1415 \operatorname{RootOf}(_Z^2 - _Z + 1)^2 x^2+3234 \operatorname{RootOf}(_Z^2 - _Z + 1)(x^3-x)^{\frac{2}{3}}-229716}{\dots}\right)$

[In] int((-1+x)^2*(1+3*x)/x/(1+x)/(-1+3*x)/(x^3-x)^(1/3),x,method=_RETURNVERBOSE)

[Out] -3/(1+x)/x*(x^3-x)^(2/3)-ln((11314080*RootOf(11664*_Z^2-108*_Z+1)^2*x^2+349272*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(2/3)-229716*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(1/3)*x-29416608*RootOf(11664*_Z^2-108*_Z+1)^2*x-224316*RootOf(11664*_Z^2-108*_Z+1)*x^2-1107*(x^3-x)^(2/3)+229716*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(1/3)+3234*(x^3-x)^(1/3)*x+13576896*RootOf(11664*_Z^2-108*_Z+1)^2+42660*RootOf(11664*_Z^2-108*_Z+1)*x-1157*x^2-3234*(x^3-x)^(1/3)-88992*RootOf(11664*_Z^2-108*_Z+1)+712*x-623)/(-1+3*x))+108*RootOf(11664*_Z^2-108*_Z+1)*ln(-(-5190480*RootOf(11664*_Z^2-108*_Z+1)^2*x^2+349272*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(2/3)-119556*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(1/3))*x+13495248*RootOf(11664*_Z^2-108*_Z+1)^2*x-181656*RootOf(11664*_Z^2-108*_Z+1)*x^2-2127*(x^3-x)^(2/3)+119556*RootOf(11664*_Z^2-108*_Z+1)*(x^3-x)^(1/3))+3234*(x^3-x)^(1/3)*x-6228576*RootOf(11664*_Z^2-108*_Z+1)^2+224316*RootOf(11664*_Z^2-108*_Z+1)*x-1552*x^2-3234*(x^3-x)^(1/3)-135324*RootOf(11664*_Z^2-108*_Z+1)-970*x-194)/(-1+3*x))

Fricas [A] (verification not implemented)

none

Time = 0.65 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.83

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx$$

$$= \frac{2\sqrt{3}(x^2+x) \arctan\left(\frac{612314840\sqrt{3}(x^3-x)^{\frac{1}{3}}(x-1)+\sqrt{3}(1609127381x^2+1235276981x+124616800)+2605939922\sqrt{3}(x^3-x)^{\frac{2}{3}}}{2990437623x^2+3108349623x-39304000}\right) - \log((3(x^3-x)^{\frac{1}{3}}(x-1)+3x-3(x^3-x)^{\frac{2}{3}}-1)/(3x-1)) - 6(x^3-x)^{\frac{2}{3}}/(x^2+x)}{2(x^2+x)}$$

```
[In] integrate((-1+x)^2*(1+3*x)/x/(1+x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="fricas")
```

```
[Out] 1/2*(2*sqrt(3)*(x^2 + x)*arctan((612314840*sqrt(3)*(x^3 - x)^(1/3)*(x - 1) + sqrt(3)*(1609127381*x^2 + 1235276981*x + 124616800) + 2605939922*sqrt(3)*(x^3 - x)^(2/3))/(2990437623*x^2 + 3108349623*x - 39304000)) - (x^2 + x)*log((3*(x^3 - x)^(1/3)*(x - 1) + 3*x - 3*(x^3 - x)^(2/3) - 1)/(3*x - 1)) - 6*(x^3 - x)^(2/3))/(x^2 + x)
```

Sympy [F]

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(x-1)^2 \cdot (3x+1)}{x\sqrt[3]{x(x-1)(x+1)}(x+1)(3x-1)} dx$$

```
[In] integrate((-1+x)**2*(1+3*x)/x/(1+x)/(-1+3*x)/(x**3-x)**(1/3),x)
```

```
[Out] Integral((x - 1)**2*(3*x + 1)/(x*(x*(x - 1)*(x + 1))**(1/3)*(x + 1)*(3*x - 1)), x)
```

Maxima [F]

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(3x+1)(x-1)^2}{(x^3-x)^{\frac{1}{3}}(3x-1)(x+1)x} dx$$

```
[In] integrate((-1+x)^2*(1+3*x)/x/(1+x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="maxima")
```

```
[Out] integrate((3*x + 1)*(x - 1)^2/((x^3 - x)^(1/3)*(3*x - 1)*(x + 1)*x), x)
```

Giac [F]

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(3x+1)(x-1)^2}{(x^3-x)^{1/3}(3x-1)(x+1)x} dx$$

[In] integrate((-1+x)^2*(1+3*x)/x/(1+x)/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="giac")

[Out] integrate((3*x + 1)*(x - 1)^2/((x^3 - x)^(1/3)*(3*x - 1)*(x + 1)*x), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(-1+x)^2(1+3x)}{x(1+x)(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(3x+1)(x-1)^2}{x(x^3-x)^{1/3}(3x-1)(x+1)} dx$$

[In] int(((3*x + 1)*(x - 1)^2)/(x*(x^3 - x)^(1/3)*(3*x - 1)*(x + 1)),x)

[Out] int(((3*x + 1)*(x - 1)^2)/(x*(x^3 - x)^(1/3)*(3*x - 1)*(x + 1)), x)

$$3.13 \quad \int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx$$

Optimal result	121
Rubi [F]	121
Mathematica [A] (verified)	127
Maple [C] (verified)	127
Fricas [A] (verification not implemented)	128
Sympy [F]	128
Maxima [F]	128
Giac [F]	129
Mupad [F(-1)]	129

Optimal result

Integrand size = 33, antiderivative size = 113

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = -\frac{3\sqrt[3]{x(-1+x^2)}}{-1+x} + \frac{1}{2} \left(-2\sqrt{3} \arctan \left(\frac{\sqrt{3}(-1+x)}{-1+x+2\sqrt[3]{x(-1+x^2)}} \right) - 2 \log \left(1-x+\sqrt[3]{x(-1+x^2)} \right) + \log \left(1-2x+x^2+(-1+x)\sqrt[3]{x(-1+x^2)}+(x(-1+x^2))^{2/3} \right) \right)$$

[Out] $-3*(x*(x^2-1))^{(1/3)/(-1+x)-3^{(1/2)*\arctan(3^{(1/2)*(-1+x)/(-1+x+2*(x*(x^2-1))^{(1/3)})}-\ln(1-x+(x*(x^2-1))^{(1/3)})+1/2*\ln(1-2*x+x^2+(-1+x)*(x*(x^2-1))^{(1/3)}+(x*(x^2-1))^{(2/3))}$

Rubi [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx$$

[In] $\text{Int}[(x*(1+x)*(1+3*x))/((-1+x)*(-1+3*x)*(-x+x^3)^{(2/3)},x]$

[Out] $(4*x*(1-x^2)^{(2/3)*\text{AppellF1}[1/6, 2/3, 1, 7/6, x^2, 9*x^2]} / (-x+x^3)^{(2/3)} - (\text{Sqrt}[3]*x^{(2/3)*(-1+x^2)^{(2/3)*\text{ArcTan}[(1-(4*x^{(2/3)})/(-1+x^2)^{(1/3)})/\text{Sqrt}[3]]} / (2*(-x+x^3)^{(2/3)}) - (\text{Sqrt}[3]*x^{(2/3)*(-1+x^2)^{(2/3)*\text{ArcTan}[(1+(2*x^{(2/3)})/(-1+x^2)^{(1/3)})/\text{Sqrt}[3]]} / (2*(-x+x^3)^{(2/3)}) + (4*x*(1-x^2)*(1-x^{(2/3)/(-1+x^2)^{(1/3)})*\text{Sqrt}[(1+x^{(4/3)/(-1+x^2)^{(2/3)}}$

$$\begin{aligned}
& /3) + x^{2/3}/(-1 + x^2)^{1/3})/(1 - ((1 + \text{Sqrt}[3])x^{2/3})/(-1 + x^2)^{1/3}) \\
& 3))^{2/3} * \text{EllipticF}[\text{ArcCos}[(1 - ((1 - \text{Sqrt}[3])x^{2/3})/(-1 + x^2)^{1/3})/(1 - \\
& ((1 + \text{Sqrt}[3])x^{2/3})/(-1 + x^2)^{1/3})], (2 + \text{Sqrt}[3])/4)]/(3^{1/4}) * (-x \\
& + x^3)^{2/3} * \text{Sqrt}[-((x^{2/3}) * (1 - x^{2/3})/(-1 + x^2)^{1/3})/((-1 + x^2)^{1/3}) * (1 - ((1 + \text{Sqrt}[3])x^{2/3})/(-1 + x^2)^{1/3})^{2/3})] + (x^{2/3}) * (-1 + \\
& x^2)^{2/3} * \text{Log}[1 - 9x^2]/(4 * (-x + x^3)^{2/3}) - (3x^{2/3}) * (-1 + x^2)^{2/3} * \text{Log}[x^{2/3} - (-1 + x^2)^{1/3}]/(4 * (-x + x^3)^{2/3}) - (3x^{2/3}) * (-1 + \\
& x^2)^{2/3} * \text{Log}[2x^{2/3} + (-1 + x^2)^{1/3}]/(4 * (-x + x^3)^{2/3}) + (4x^{2/3}) * (-1 + x^2)^{2/3} * \text{Defer}[\text{Subst}[\text{Defer}[\text{Int}][1/((-1 + x) * (-1 + x^6)^{2/3}) \\
&], x], x, x^{1/3}]/(-x + x^3)^{2/3} - (4 * (1 - \text{I} * \text{Sqrt}[3])x^{2/3}) * (-1 + x^2)^{2/3} * \text{Defer}[\text{Subst}[\text{Defer}[\text{Int}][1/((1 - \text{I} * \text{Sqrt}[3] + 2x) * (-1 + x^6)^{2/3}) \\
&], x], x, x^{1/3}]/(-x + x^3)^{2/3} - (4 * (1 + \text{I} * \text{Sqrt}[3])x^{2/3}) * (-1 + x^2)^{2/3} * \text{Defer}[\text{Subst}[\text{Defer}[\text{Int}][1/((1 + \text{I} * \text{Sqrt}[3] + 2x) * (-1 + x^6)^{2/3}) \\
&], x, x^{1/3}]/(-x + x^3)^{2/3}
\end{aligned}$$

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(x^{2/3}(-1 + x^2)^{2/3}) \int \frac{\sqrt[3]{x(1+x)(1+3x)}}{(-1+x)(-1+3x)(-1+x^2)^{2/3}} dx}{(-x + x^3)^{2/3}} \\
&= \frac{(3x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{x^3(1+x^3)(1+3x^3)}{(-1+x^3)(-1+3x^3)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&= \frac{(3x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \left(\frac{8}{3(-1+x^6)^{2/3}} + \frac{4}{3(-1+x)(-1+x^6)^{2/3}} + \frac{x^3}{(-1+x^6)^{2/3}} - \frac{4(2+x)}{3(1+x+x^2)(-1+x^6)^{2/3}} - \frac{1}{3}\right) dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&= \frac{(3x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{x^3}{(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&+ \frac{(4x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{1}{(-1+x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&- \frac{(4x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{2+x}{(1+x+x^2)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&- \frac{(4x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{1}{(-1+3x^3)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}} \\
&+ \frac{(8x^{2/3}(-1 + x^2)^{2/3}) \text{Subst}\left(\int \frac{1}{(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x + x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
& \frac{(8x^{2/3}\sqrt[6]{-1+x^2}) \operatorname{Subst}\left(\int \frac{1}{\sqrt{1-x^6}} dx, x, \frac{\sqrt[3]{x}}{\sqrt[6]{-1+x^2}}\right)}{\sqrt{-\frac{1}{-1+x^2}}(-x+x^3)^{2/3}} \\
& + \frac{(3x^{2/3}(-1+x^2)^{2/3}) \operatorname{Subst}\left(\int \frac{x}{(-1+x^3)^{2/3}} dx, x, x^{2/3}\right)}{2(-x+x^3)^{2/3}} \\
& + \frac{(4x^{2/3}(-1+x^2)^{2/3}) \operatorname{Subst}\left(\int \frac{1}{(-1+x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{(4x^{2/3}(-1+x^2)^{2/3}) \operatorname{Subst}\left(\int \left(\frac{1-i\sqrt{3}}{(1-i\sqrt{3}+2x)(-1+x^6)^{2/3}} + \frac{1+i\sqrt{3}}{(1+i\sqrt{3}+2x)(-1+x^6)^{2/3}}\right) dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{(4x^{2/3}(-1+x^2)^{2/3}) \operatorname{Subst}\left(\int \left(\frac{1}{(-1+x^6)^{2/3}(-1+9x^6)} + \frac{3x^3}{(-1+x^6)^{2/3}(-1+9x^6)}\right) dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
& + \frac{4x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \operatorname{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right), \frac{1}{4}}\right)}{\sqrt[4]{3}(-x+x^3)^{2/3} \sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
& - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
& + \frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x^6)^{2/3}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(12x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{x^3}{(-1+x^6)^{2/3}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(4(1-i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1-i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(4(1+i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
& + \frac{4x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \operatorname{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right)\right)}{\sqrt[4]{3}(-x+x^3)^{2/3} \sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
& - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
& - \frac{\left(4x^{2/3}(1-x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1-x^6)^{2/3}(-1+9x^6)} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& + \frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(6x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{x}{(-1+x^3)^{2/3}(-1+9x^3)} dx, x, x^{2/3}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(4(1-i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1-i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
& - \frac{\left(4(1+i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4x(1-x^2)^{2/3} \operatorname{AppellF1}\left(\frac{1}{6}, \frac{2}{3}, 1, \frac{7}{6}, x^2, 9x^2\right)}{(-x+x^3)^{2/3}} \\
&\quad - \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1-\frac{4x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
&\quad - \frac{\sqrt{3}x^{2/3}(-1+x^2)^{2/3} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2(-x+x^3)^{2/3}} \\
&\quad + \frac{4x(1-x^2)\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right) \sqrt{\frac{1+\frac{x^{4/3}}{(-1+x^2)^{2/3}}+\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}}{\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}} \operatorname{EllipticF}\left(\arccos\left(\frac{1-\frac{(1-\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}{1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}}\right), \frac{1}{4}}\right)}{\sqrt[4]{3}(-x+x^3)^{2/3} \sqrt{-\frac{x^{2/3}\left(1-\frac{x^{2/3}}{\sqrt[3]{-1+x^2}}\right)}{\sqrt[3]{-1+x^2}\left(1-\frac{(1+\sqrt{3})x^{2/3}}{\sqrt[3]{-1+x^2}}\right)^2}}} \\
&\quad + \frac{x^{2/3}(-1+x^2)^{2/3} \log(1-9x^2)}{4(-x+x^3)^{2/3}} - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(x^{2/3}-\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
&\quad - \frac{3x^{2/3}(-1+x^2)^{2/3} \log\left(2x^{2/3}+\sqrt[3]{-1+x^2}\right)}{4(-x+x^3)^{2/3}} \\
&\quad + \frac{\left(4x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&\quad - \frac{\left(4(1-i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1-i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}} \\
&\quad - \frac{\left(4(1+i\sqrt{3})x^{2/3}(-1+x^2)^{2/3}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)(-1+x^6)^{2/3}} dx, x, \sqrt[3]{x}\right)}{(-x+x^3)^{2/3}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 31.58 (sec) , antiderivative size = 113, normalized size of antiderivative = 1.00

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = -\frac{3\sqrt[3]{x(-1+x^2)}}{-1+x} + \frac{1}{2} \left(-2\sqrt{3} \arctan \left(\frac{\sqrt{3}(-1+x)}{-1+x+2\sqrt[3]{x(-1+x^2)}} \right) - 2 \log \left(1-x+\sqrt[3]{x(-1+x^2)} \right) + \log \left(1-2x+x^2+(-1+x)\sqrt[3]{x(-1+x^2)}+(x(-1+x^2))^{2/3} \right) \right)$$

[In] Integrate[(x*(1+x)*(1+3*x))/((-1+x)*(-1+3*x)*(-x+x^3)^(2/3)),x]

[Out] $(-3*(x*(-1+x^2))^{1/3})/(-1+x) + (-2*\text{Sqrt}[3]*\text{ArcTan}[(\text{Sqrt}[3]*(-1+x))/(-1+x+2*(x*(-1+x^2))^{1/3})] - 2*\text{Log}[1-x+(x*(-1+x^2))^{1/3}] + \text{Log}[1-2*x+x^2+(-1+x)*(x*(-1+x^2))^{1/3}+(x*(-1+x^2))^{2/3}])/2$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.35 (sec) , antiderivative size = 458, normalized size of antiderivative = 4.05

method	result
trager	$-\frac{3(x^3-x)^{1/3}}{-1+x} - \ln \left(-\frac{-8730 \text{RootOf}(9_Z^2-3_Z+1)^2 x^2 + 9702 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{2/3} - 6381 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} + 22698 \text{RootOf}(9_Z^2-3_Z+1)^2 x - 411 \text{RootOf}(9_Z^2-3_Z+1) x^2 - 2127(x^3-x)^{2/3} + 6381 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} - 1107(x^3-x)^{1/3} x - 10476 \text{RootOf}(9_Z^2-3_Z+1)^2 - 13947 \text{RootOf}(9_Z^2-3_Z+1) x + 2264 x^2 + 1107(x^3-x)^{1/3} + 4512 \text{RootOf}(9_Z^2-3_Z+1) + 1415 x + 283}{(-1+3*x)} + 3 \text{RootOf}(9_Z^2-3_Z+1) \ln \left(\frac{4005 \text{RootOf}(9_Z^2-3_Z+1)^2 x^2 + 9702 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{2/3} - 3321 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} x - 10413 \text{RootOf}(9_Z^2-3_Z+1)^2 x - 7716 \text{RootOf}(9_Z^2-3_Z+1) x^2 - 1107(x^3-x)^{2/3} + 3321 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} - 2127(x^3-x)^{1/3} x + 4806 \text{RootOf}(9_Z^2-3_Z+1)^2 + 13173}{(-1+3*x)} \right) \right)$
risch	Expression too large to display

[In] int(x*(1+x)*(1+3*x)/(-1+x)/(-1+3*x)/(x^3-x)^(2/3),x,method=_RETURNVERBOSE)

[Out] $-3/(-1+x)*(x^3-x)^{1/3} - \ln \left(\frac{-8730 \text{RootOf}(9_Z^2-3_Z+1)^2 x^2 + 9702 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{2/3} - 6381 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} x + 22698 \text{RootOf}(9_Z^2-3_Z+1)^2 x - 411 \text{RootOf}(9_Z^2-3_Z+1) x^2 - 2127(x^3-x)^{2/3} + 6381 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} - 1107(x^3-x)^{1/3} x - 10476 \text{RootOf}(9_Z^2-3_Z+1)^2 - 13947 \text{RootOf}(9_Z^2-3_Z+1) x + 2264 x^2 + 1107(x^3-x)^{1/3} + 4512 \text{RootOf}(9_Z^2-3_Z+1) + 1415 x + 283}{(-1+3*x)} + 3 \text{RootOf}(9_Z^2-3_Z+1) \ln \left(\frac{4005 \text{RootOf}(9_Z^2-3_Z+1)^2 x^2 + 9702 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{2/3} - 3321 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} x - 10413 \text{RootOf}(9_Z^2-3_Z+1)^2 x - 7716 \text{RootOf}(9_Z^2-3_Z+1) x^2 - 1107(x^3-x)^{2/3} + 3321 \text{RootOf}(9_Z^2-3_Z+1)(x^3-x)^{1/3} - 2127(x^3-x)^{1/3} x + 4806 \text{RootOf}(9_Z^2-3_Z+1)^2 + 13173}{(-1+3*x)} \right) \right)$

*RootOf(9*_Z^2-3*_Z+1)*x+3679*x^2+2127*(x^3-x)^(1/3)-6963*RootOf(9*_Z^2-3*_Z+1)-2264*x+1981)/(-1+3*x))

Fricas [A] (verification not implemented)

none

Time = 0.52 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.15

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \frac{2\sqrt{3}(x-1) \arctan\left(\frac{612314840\sqrt{3}(x^3-x)^{1/3}(x-1)+\sqrt{3}(1609127381x^2+1235276981x+124616800)+2605939922\sqrt{3}(x^3-x)^{2/3}}{2990437623x^2+3108349623x-39304000}\right) + (x-1)}{2(x-1)}$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)/(-1+3*x)/(x^3-x)^(2/3),x, algorithm="fricas")

[Out] -1/2*(2*sqrt(3)*(x - 1)*arctan((612314840*sqrt(3)*(x^3 - x)^(1/3)*(x - 1) + sqrt(3)*(1609127381*x^2 + 1235276981*x + 124616800) + 2605939922*sqrt(3)*(x^3 - x)^(2/3))/(2990437623*x^2 + 3108349623*x - 39304000)) + (x - 1)*log((3*(x^3 - x)^(1/3)*(x - 1) + 3*x - 3*(x^3 - x)^(2/3) - 1)/(3*x - 1)) + 6*(x^3 - x)^(1/3))/(x - 1)

Sympy [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{x(x+1)(3x+1)}{(x(x-1)(x+1))^{2/3}(x-1)(3x-1)} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)/(-1+3*x)/(x**3-x)**(2/3),x)

[Out] Integral(x*(x + 1)*(3*x + 1)/((x*(x - 1)*(x + 1))**(2/3)*(x - 1)*(3*x - 1)), x)

Maxima [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(3x+1)(x+1)x}{(x^3-x)^{2/3}(3x-1)(x-1)} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)/(-1+3*x)/(x^3-x)^(2/3),x, algorithm="maxima")

[Out] integrate((3*x + 1)*(x + 1)*x/((x^3 - x)^(2/3)*(3*x - 1)*(x - 1)), x)

Giac [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{(3x+1)(x+1)x}{(x^3-x)^{2/3}(3x-1)(x-1)} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)/(-1+3*x)/(x^3-x)^(2/3),x, algorithm="giac")

[Out] integrate((3*x + 1)*(x + 1)*x/((x^3 - x)^(2/3)*(3*x - 1)*(x - 1)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x(1+x)(1+3x)}{(-1+x)(-1+3x)(-x+x^3)^{2/3}} dx = \int \frac{x(3x+1)(x+1)}{(x^3-x)^{2/3}(3x-1)(x-1)} dx$$

[In] int((x*(3*x + 1)*(x + 1))/((x^3 - x)^(2/3)*(3*x - 1)*(x - 1)),x)

[Out] int((x*(3*x + 1)*(x + 1))/((x^3 - x)^(2/3)*(3*x - 1)*(x - 1)), x)

$$3.14 \quad \int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx$$

Optimal result	130
Rubi [F]	130
Mathematica [F]	136
Maple [C] (verified)	136
Fricas [A] (verification not implemented)	137
Sympy [F]	137
Maxima [F]	137
Giac [F]	138
Mupad [F(-1)]	138

Optimal result

Integrand size = 33, antiderivative size = 166

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx$$

$$= \frac{-3(-x+x^3)^{2/3} + 2\sqrt{3}(1-2x+x^2) \arctan\left(\frac{\sqrt{3}(124616800+1235276981x+1609127381x^2)+612314840\sqrt{3}(-1+x)\sqrt[3]{-x+x^3}}{-39304000+3108349623x+2990437623x^2}\right)}{2(1-2x+x^2)}$$

[Out] 1/2*(2*3^(1/2)*(x^2-2*x+1)*arctan((612314840*3^(1/2)*(x^3-x)^(1/3)*(-1+x)+3^(1/2)*(1609127381*x^2+1235276981*x+124616800)+2605939922*3^(1/2)*(x^3-x)^(2/3))/(2990437623*x^2+3108349623*x-39304000))-(x^2-2*x+1)*ln((3*(-1+x)*(x^3-x)^(1/3)+3*x-3*(x^3-x)^(2/3)-1)/(-1+3*x))-3*(x^3-x)^(2/3))/(x^2-2*x+1)

Rubi [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx$$

[In] Int[(x*(1+x)*(1+3*x))/((-1+x)^2*(-1+3*x)*(-x+x^3)^(1/3)),x]

[Out] (-18*x^2*(1-x^2)^(1/3)*AppellF1[5/6, 1/3, 1, 11/6, x^2, 9*x^2])/(5*(-x+x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1+x^2)^(1/3)*ArcTan[(1-(4*x^(2/3)))/(-1+x^2)^(1/3)]/Sqrt[3])/(2*(-x+x^3)^(1/3)) + (Sqrt[3]*x^(1/3)*(-1+x^2)^(1/3)*ArcTan[(1+(2*x^(2/3)))/(-1+x^2)^(1/3)]/Sqrt[3])/(2*(-x+x^3)^(1/3)) + (x^(1/3)*(-1+x^2)^(1/3)*Log[1-9*x^2])/(4*(-x+x^3)^(1/3)) - (3*x^(1/3)*(-1+x^2)^(1/3)*Log[x^(2/3)-(-1+x^2)^(1/3)])/(4*(-x+x^3)^(1/3))

3)) - (3*x^(1/3)*(-1 + x^2)^(1/3)*Log[2*x^(2/3) + (-1 + x^2)^(1/3)]/(4*(-x + x^3)^(1/3)) + (16*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((-1 + I*Sqrt[3] - 2*x)^2*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(3*(-x + x^3)^(1/3)) - (((16*I)/3)*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((-1 + I*Sqrt[3] - 2*x)*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(Sqrt[3]*(-x + x^3)^(1/3)) + (4*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((-1 + x)^2*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(3*(-x + x^3)^(1/3)) + (5*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((-1 + x)*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(3*(-x + x^3)^(1/3)) - ((15 + (31*I)*Sqrt[3])*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((1 - I*Sqrt[3] + 2*x)*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(9*(-x + x^3)^(1/3)) + (16*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((1 + I*Sqrt[3] + 2*x)^2*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(3*(-x + x^3)^(1/3)) - (((16*I)/3)*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((1 + I*Sqrt[3] + 2*x)*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(Sqrt[3]*(-x + x^3)^(1/3)) - ((15 - (31*I)*Sqrt[3])*x^(1/3)*(-1 + x^2)^(1/3)*Defer[Subst][Defer[Int][1/((1 + I*Sqrt[3] + 2*x)*(-1 + x^6)^(1/3)), x], x, x^(1/3)]/(9*(-x + x^3)^(1/3))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \int \frac{x^{2/3}(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-1+x^2}} dx}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x^4(1+x^3)(1+3x^3)}{(-1+x^3)^2(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
 &= \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(\frac{4}{9(-1+x)^2\sqrt[3]{-1+x^6}} + \frac{5}{9(-1+x)\sqrt[3]{-1+x^6}} + \frac{x}{\sqrt[3]{-1+x^6}} - \frac{4}{3(1+x+x^2)^2\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\left(\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{13-5x}{(1+x+x^2)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(4\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1+x)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(5\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x}{\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&- \frac{\left(4\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(1+x+x^2)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(6\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{x}{(-1+3x^3)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&= \frac{\left(\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(\frac{-5-\frac{31i}{\sqrt{3}}}{(1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} + \frac{-5+\frac{31i}{\sqrt{3}}}{(1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(4\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1+x)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(3\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{\sqrt[3]{-1+x^3}} dx, x, x^{2/3}\right)}{2\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(5\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \frac{1}{(-1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&- \frac{\left(4\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(-\frac{4}{3(-1+i\sqrt{3}-2x)^2\sqrt[3]{-1+x^6}} + \frac{4i}{3\sqrt{3}(-1+i\sqrt{3}-2x)\sqrt[3]{-1+x^6}} - \frac{1}{3(1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}} \\
&+ \frac{\left(6\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \text{Subst}\left(\int \left(\frac{x}{\sqrt[3]{-1+x^6}(-1+9x^6)} + \frac{3x^4}{\sqrt[3]{-1+x^6}(-1+9x^6)}\right) dx, x, \sqrt[3]{x}\right)}{\sqrt[3]{-x+x^3}}
\end{aligned}$$

$$\begin{aligned}
& \sqrt{3} \sqrt[3]{x} \sqrt[3]{-1+x^2} \arctan \left(\frac{1 + \frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}} \right) \\
= & \frac{2\sqrt[3]{-x+x^3}}{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log \left(x^{2/3} - \sqrt[3]{-1+x^2} \right)} \\
- & \frac{4\sqrt[3]{-x+x^3}}{(4\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(-1+x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{3\sqrt[3]{-x+x^3}}{(5\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(-1+x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{3\sqrt[3]{-x+x^3}}{(16\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(-1+i\sqrt{3}-2x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{3\sqrt[3]{-x+x^3}}{(16\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{\sqrt[3]{-x+x^3}}{(6\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{x}{\sqrt[3]{-1+x^6}(-1+9x^6)} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{\sqrt[3]{-x+x^3}}{(18\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{x^4}{\sqrt[3]{-1+x^6}(-1+9x^6)} dx, x, \sqrt[3]{x} \right)} \\
- & \frac{3\sqrt{3}\sqrt[3]{-x+x^3}}{(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(-1+i\sqrt{3}-2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
- & \frac{3\sqrt{3}\sqrt[3]{-x+x^3}}{(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
+ & \frac{9\sqrt[3]{-x+x^3}}{((-15+31i\sqrt{3}) \sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
- & \frac{9\sqrt[3]{-x+x^3}}{((15+31i\sqrt{3}) \sqrt[3]{x}\sqrt[3]{-1+x^2}) \operatorname{Subst} \left(\int \frac{1}{(1-i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)}
\end{aligned}$$

$$\begin{aligned}
& \sqrt{3} \sqrt[3]{x} \sqrt[3]{-1+x^2} \arctan \left(\frac{1 + \frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}} \right) \\
= & \frac{2\sqrt[3]{-x+x^3}}{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(x^{2/3} - \sqrt[3]{-1+x^2})} \\
& - \frac{4\sqrt[3]{-x+x^3}}{(18\sqrt[3]{x}\sqrt[3]{1-x^2}) \text{Subst} \left(\int \frac{x^4}{\sqrt[3]{1-x^6(-1+9x^6)}} dx, x, \sqrt[3]{x} \right)} \\
& + \frac{\sqrt[3]{-x+x^3}}{(4\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(-1+x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& + \frac{3\sqrt[3]{-x+x^3}}{(5\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(-1+x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& + \frac{3\sqrt[3]{-x+x^3}}{(3\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{\sqrt[3]{-1+x^3(-1+9x^3)}} dx, x, x^{2/3} \right)} \\
& + \frac{\sqrt[3]{-x+x^3}}{(16\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(-1+i\sqrt{3}-2x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& + \frac{3\sqrt[3]{-x+x^3}}{(16\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x)^2 \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& - \frac{3\sqrt[3]{3}\sqrt[3]{-x+x^3}}{(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(-1+i\sqrt{3}-2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& - \frac{3\sqrt[3]{3}\sqrt[3]{-x+x^3}}{(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& + \frac{9\sqrt[3]{-x+x^3}}{((-15+31i\sqrt{3}) \sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(1+i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)} \\
& - \frac{9\sqrt[3]{-x+x^3}}{((15+31i\sqrt{3}) \sqrt[3]{x}\sqrt[3]{-1+x^2}) \text{Subst} \left(\int \frac{1}{(1-i\sqrt{3}+2x) \sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x} \right)}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{18x^2\sqrt[3]{1-x^2} \operatorname{AppellF1}\left(\frac{5}{6}, \frac{1}{3}, 1, \frac{11}{6}, x^2, 9x^2\right)}{5\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1-\frac{4x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\sqrt{3}\sqrt[3]{x}\sqrt[3]{-1+x^2} \arctan\left(\frac{1+\frac{2x^{2/3}}{\sqrt[3]{-1+x^2}}}{\sqrt{3}}\right)}{2\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\sqrt[3]{x}\sqrt[3]{-1+x^2} \log(1-9x^2)}{4\sqrt[3]{-x+x^3}} - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(x^{2/3} - \sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
&\quad - \frac{3\sqrt[3]{x}\sqrt[3]{-1+x^2} \log\left(2x^{2/3} + \sqrt[3]{-1+x^2}\right)}{4\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(4\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(5\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(16\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+i\sqrt{3}-2x)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left(16\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)^2\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(-1+i\sqrt{3}-2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt{3}\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left(16i\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{3\sqrt{3}\sqrt[3]{-x+x^3}} \\
&\quad + \frac{\left((-15+31i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(1+i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{9\sqrt[3]{-x+x^3}} \\
&\quad - \frac{\left((15+31i\sqrt{3})\sqrt[3]{x}\sqrt[3]{-1+x^2}\right) \operatorname{Subst}\left(\int \frac{1}{(1-i\sqrt{3}+2x)\sqrt[3]{-1+x^6}} dx, x, \sqrt[3]{x}\right)}{9\sqrt[3]{-x+x^3}}
\end{aligned}$$

Mathematica [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx$$

[In] Integrate[(x*(1+x)*(1+3*x))/((-1+x)^2*(-1+3*x)*(-x+x^3)^(1/3)),x]

[Out] Integrate[(x*(1+x)*(1+3*x))/((-1+x)^2*(-1+3*x)*(-x+x^3)^(1/3)), x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.64 (sec) , antiderivative size = 682, normalized size of antiderivative = 4.11

method	result	size
trager	Expression too large to display	682

[In] int(x*(1+x)*(1+3*x)/(-1+x)^2/(-1+3*x)/(x^3-x)^(1/3),x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned} & -3/2/(-1+x)^2*(x^3-x)^{(2/3)}-3*\ln(-(12735*\text{RootOf}(9*_Z^2-3*_Z+1)^2*x^2+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(2/3)}+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)} \\ & *x-33111*\text{RootOf}(9*_Z^2-3*_Z+1)^2*x+5457*\text{RootOf}(9*_Z^2-3*_Z+1)*x^2-2127*(x^3-x)^{(2/3)}-9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)}-2127*(x^3-x)^{(1/3)}*x+152 \\ & 82*\text{RootOf}(9*_Z^2-3*_Z+1)^2+7716*\text{RootOf}(9*_Z^2-3*_Z+1)*x-712*x^2+2127*(x^3-x)^{(1/3)}-753*\text{RootOf}(9*_Z^2-3*_Z+1)-445*x-89)/(-1+3*x))*\text{RootOf}(9*_Z^2-3*_Z+1) \\ & +3*\text{RootOf}(9*_Z^2-3*_Z+1)*\ln((-12735*\text{RootOf}(9*_Z^2-3*_Z+1)^2*x^2+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(2/3)}+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)}*x+33 \\ & 111*\text{RootOf}(9*_Z^2-3*_Z+1)^2*x+13947*\text{RootOf}(9*_Z^2-3*_Z+1)*x^2-1107*(x^3-x)^{(2/3)}-9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)}-1107*(x^3-x)^{(1/3)}*x-15282*\text{R} \\ & \text{ootOf}(9*_Z^2-3*_Z+1)^2-14358*\text{RootOf}(9*_Z^2-3*_Z+1)*x-2522*x^2+1107*(x^3-x)^{(1/3)}+9435*\text{RootOf}(9*_Z^2-3*_Z+1)+1552*x-1358)/(-1+3*x))+\ln(-(12735*\text{RootOf}(9 \\ & *_Z^2-3*_Z+1)^2*x^2+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(2/3)}+9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)}*x-33111*\text{RootOf}(9*_Z^2-3*_Z+1)^2*x+5457*\text{RootOf}(9*_ \\ & *_Z^2-3*_Z+1)*x^2-2127*(x^3-x)^{(2/3)}-9702*\text{RootOf}(9*_Z^2-3*_Z+1)*(x^3-x)^{(1/3)}-2127*(x^3-x)^{(1/3)}*x+15282*\text{RootOf}(9*_Z^2-3*_Z+1)^2+7716*\text{RootOf}(9*_Z^2-3*_ \\ & *_Z+1)*x-712*x^2+2127*(x^3-x)^{(1/3)}-753*\text{RootOf}(9*_Z^2-3*_Z+1)-445*x-89)/(-1+3*x)) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.65 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.88

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx$$

$$= \frac{2\sqrt{3}(x^2-2x+1) \arctan\left(\frac{612314840\sqrt{3}(x^3-x)^{\frac{1}{3}}(x-1)+\sqrt{3}(1609127381x^2+1235276981x+124616800)+2605939922\sqrt{3}(x^3-x)^{\frac{2}{3}}}{2990437623x^2+3108349623x-39304000}\right)}{2(x^2-2x+1)}$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)^2/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="fricas")

[Out] 1/2*(2*sqrt(3)*(x^2 - 2*x + 1)*arctan((612314840*sqrt(3)*(x^3 - x)^(1/3)*(x - 1) + sqrt(3)*(1609127381*x^2 + 1235276981*x + 124616800) + 2605939922*sqrt(3)*(x^3 - x)^(2/3))/(2990437623*x^2 + 3108349623*x - 39304000)) - (x^2 - 2*x + 1)*log((3*(x^3 - x)^(1/3)*(x - 1) + 3*x - 3*(x^3 - x)^(2/3) - 1)/(3*x - 1)) - 3*(x^3 - x)^(2/3))/(x^2 - 2*x + 1)

Sympy [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{x(x+1)(3x+1)}{\sqrt[3]{x(x-1)(x+1)}(x-1)^2 \cdot (3x-1)} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)**2/(-1+3*x)/(x**3-x)**(1/3),x)

[Out] Integral(x*(x + 1)*(3*x + 1)/((x*(x - 1)*(x + 1))**(1/3)*(x - 1)**2*(3*x - 1)), x)

Maxima [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(3x+1)(x+1)x}{(x^3-x)^{\frac{1}{3}}(3x-1)(x-1)^2} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)^2/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="maxima")

[Out] integrate((3*x + 1)*(x + 1)*x/((x^3 - x)^(1/3)*(3*x - 1)*(x - 1)^2), x)

Giac [F]

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{(3x+1)(x+1)x}{(x^3-x)^{\frac{1}{3}}(3x-1)(x-1)^2} dx$$

[In] integrate(x*(1+x)*(1+3*x)/(-1+x)^2/(-1+3*x)/(x^3-x)^(1/3),x, algorithm="giac")

[Out] integrate((3*x + 1)*(x + 1)*x/((x^3 - x)^(1/3)*(3*x - 1)*(x - 1)^2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{x(1+x)(1+3x)}{(-1+x)^2(-1+3x)\sqrt[3]{-x+x^3}} dx = \int \frac{x(3x+1)(x+1)}{(x^3-x)^{1/3}(3x-1)(x-1)^2} dx$$

[In] int((x*(3*x + 1)*(x + 1))/((x^3 - x)^(1/3)*(3*x - 1)*(x - 1)^2),x)

[Out] int((x*(3*x + 1)*(x + 1))/((x^3 - x)^(1/3)*(3*x - 1)*(x - 1)^2), x)

$$3.15 \quad \int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx$$

Optimal result	139
Rubi [F]	139
Mathematica [F]	141
Maple [C] (verified)	142
Fricas [A] (verification not implemented)	143
Sympy [F]	143
Maxima [F]	143
Giac [F]	144
Mupad [F(-1)]	144

Optimal result

Integrand size = 40, antiderivative size = 194

$$\int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx = \frac{\arctan\left(\frac{258\sqrt{3}(-1+x^3)^{2/3}(-59x^2+2x^5)-258\sqrt{3}\sqrt[3]{-1+x^3}(-24x+56x^4+13824-25667x^3+13021x^6+2197x^9)}{13824-25667x^3+13021x^6+2197x^9}\right)}{3\sqrt{3}} + \frac{1}{18} \log\left(\frac{-27+82x^3-83x^6+x^9+18(-1+x^3)^{2/3}(-5x^2+2x^5)-9\sqrt[3]{-1+x^3}(3x-13x^4+x^7)}{-27+x^3-2x^6+x^9}\right)$$

```
[Out] 1/9*3^(1/2)*arctan((258*3^(1/2)*(2*x^5-59*x^2)*(x^3-1)^(2/3)-258*3^(1/2)*(1
3*x^7+56*x^4-24*x)*(x^3-1)^(1/3)-3^(1/2)*(169*x^9+7789*x^6+7135*x^3-10368))
/(2197*x^9+13021*x^6-25667*x^3+13824))+1/18*ln((x^9-83*x^6+82*x^3+18*(2*x^5
-5*x^2)*(x^3-1)^(2/3)-9*(x^7-13*x^4+3*x)*(x^3-1)^(1/3)-27)/(x^9-2*x^6+x^3-2
7))
```

Rubi [F]

$$\int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx = \int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx$$

```
[In] Int[(3 + 8*x^3 - 3*x^6 + x^9)/((-1 + x^3)^(2/3)*(-27 + x^3 - 2*x^6 + x^9)),
x]
```

```
[Out] (x*(1 - x^3)^(2/3)*Hypergeometric2F1[1/3, 2/3, 4/3, x^3])/(-1 + x^3)^(2/3)
+ (10*Defer[Int][1/((-1 + x^3)^(2/3)*(-3 - 2*x + x^2 + x^3)), x])/9 + (7*De
fer[Int][x/((-1 + x^3)^(2/3)*(-3 - 2*x + x^2 + x^3)), x])/9 - (2*Defer[Int]
[x^2/((-1 + x^3)^(2/3)*(-3 - 2*x + x^2 + x^3)), x])/9 - (20*Defer[Int][1/((
```

$-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3 + 3*x^4 - x^5 + x^6)), x]) / 3 + (41$
 $* \text{Defer}[\text{Int}][x / ((-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3 + 3*x^4 - x^5 + x^6)), x]) / 9 - (44 * \text{Defer}[\text{Int}][x^2 / ((-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3$
 $+ 3*x^4 - x^5 + x^6)), x]) / 9 + \text{Defer}[\text{Int}][x^3 / ((-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3 + 3*x^4 - x^5 + x^6)), x] - (11 * \text{Defer}[\text{Int}][x^4 / ((-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3 + 3*x^4 - x^5 + x^6)), x]) / 9 + (2 * \text{Defer}[\text{Int}][x$
 $^5 / ((-1 + x^3)^{(2/3)} * (9 - 6*x + 7*x^2 - 4*x^3 + 3*x^4 - x^5 + x^6)), x]) / 9$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{(-1 + x^3)^{2/3}} + \frac{30 + 7x^3 - x^6}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} \right) dx \\
 &= \int \frac{1}{(-1 + x^3)^{2/3}} dx + \int \frac{30 + 7x^3 - x^6}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx \\
 &= \frac{(1 - x^3)^{2/3} \int \frac{1}{(1 - x^3)^{2/3}} dx}{(-1 + x^3)^{2/3}} \\
 &\quad + \int \left(\frac{10 + 7x - 2x^2}{9(-1 + x^3)^{2/3} (-3 - 2x + x^2 + x^3)} + \frac{-60 + 41x - 44x^2 + 9x^3 - 11x^4 + 2x^5}{9(-1 + x^3)^{2/3} (9 - 6x + 7x^2 - 4x^3 + 3x^4 - x^5 + x^6)} \right) dx \\
 &= \frac{x(1 - x^3)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{(-1 + x^3)^{2/3}} \\
 &\quad + \frac{1}{9} \int \frac{10 + 7x - 2x^2}{(-1 + x^3)^{2/3} (-3 - 2x + x^2 + x^3)} dx \\
 &\quad + \frac{1}{9} \int \frac{-60 + 41x - 44x^2 + 9x^3 - 11x^4 + 2x^5}{(-1 + x^3)^{2/3} (9 - 6x + 7x^2 - 4x^3 + 3x^4 - x^5 + x^6)} dx \\
 &= \frac{x(1 - x^3)^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{(-1 + x^3)^{2/3}} \\
 &\quad + \frac{1}{9} \int \left(\frac{10}{(-1 + x^3)^{2/3} (-3 - 2x + x^2 + x^3)} + \frac{7x}{(-1 + x^3)^{2/3} (-3 - 2x + x^2 + x^3)} - \frac{2}{(-1 + x^3)^{2/3} (-3 - 2x + x^2 + x^3)} \right) dx
 \end{aligned}$$

$$\begin{aligned}
&= \frac{x(1-x^3)^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{2}{3}, \frac{4}{3}, x^3\right)}{(-1+x^3)^{2/3}} \\
&\quad - \frac{2}{9} \int \frac{x^2}{(-1+x^3)^{2/3}(-3-2x+x^2+x^3)} dx \\
&\quad + \frac{2}{9} \int \frac{x^5}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx \\
&\quad + \frac{7}{9} \int \frac{x}{(-1+x^3)^{2/3}(-3-2x+x^2+x^3)} dx \\
&\quad + \frac{10}{9} \int \frac{1}{(-1+x^3)^{2/3}(-3-2x+x^2+x^3)} dx \\
&\quad - \frac{11}{9} \int \frac{x^4}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx \\
&\quad + \frac{41}{9} \int \frac{x}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx \\
&\quad - \frac{44}{9} \int \frac{x^2}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx \\
&\quad - \frac{20}{3} \int \frac{1}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx \\
&\quad + \int \frac{x^3}{(-1+x^3)^{2/3}(9-6x+7x^2-4x^3+3x^4-x^5+x^6)} dx
\end{aligned}$$

Mathematica [F]

$$\int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx = \int \frac{3+8x^3-3x^6+x^9}{(-1+x^3)^{2/3}(-27+x^3-2x^6+x^9)} dx$$

[In] Integrate[(3 + 8*x^3 - 3*x^6 + x^9)/((-1 + x^3)^(2/3)*(-27 + x^3 - 2*x^6 + x^9)), x]

[Out] Integrate[(3 + 8*x^3 - 3*x^6 + x^9)/((-1 + x^3)^(2/3)*(-27 + x^3 - 2*x^6 + x^9)), x]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.28 (sec) , antiderivative size = 845, normalized size of antiderivative = 4.36

method	result	size
trager	Expression too large to display	845

[In] $\text{int}((x^9-3x^6+8x^3+3)/(x^3-1)^{(2/3)}/(x^9-2x^6+x^3-27), x, \text{method}=_RETURNVE$
RBOSE)

[Out] $\frac{1}{9}\ln\left(\frac{-206955\sqrt[3]{81Z^2+9Z+1}^2x^9-1011123\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{1/3}x^7-106434\sqrt[3]{81Z^2+9Z+1}x^9+4516155\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{2/3}x^5+206955\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x^7-9271x^9-5173875\sqrt[3]{81Z^2+9Z+1}^2x^6+407187\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{2/3}x^5+35478(x^3-1)^{1/3}x^7-8088984\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{1/3}x^4-2039985\sqrt[3]{81Z^2+9Z+1}x^6-63162(x^3-1)^{2/3}x^5+2329722\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x^4+18542x^6-206955\sqrt[3]{81Z^2+9Z+1}^2x^3-2442069\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{2/3}x^2+70956(x^3-1)^{1/3}x^4+514431\sqrt[3]{81Z^2+9Z+1}x^3-419166(x^3-1)^{2/3}x^2+337041\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x+241046x^3-106434(x^3-1)^{1/3}x-620865\sqrt[3]{81Z^2+9Z+1}-250317}{(x^3+x^2-2x-3)/(x^6-x^5+3x^4-4x^3+7x^2-6x+9)}\right)+\sqrt[3]{81Z^2+9Z+1}\ln\left(\frac{-60444\sqrt[3]{81Z^2+9Z+1}^2x^9-319302\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{1/3}x^7-9271\sqrt[3]{81Z^2+9Z+1}x^9+501795\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{2/3}x^5+12483\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x^7-1511100\sqrt[3]{81Z^2+9Z+1}^2x^6+66267\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{2/3}x^5-2554416\sqrt[3]{81Z^2+9Z+1}^2(x^3-1)^{1/3}x^4-413107\sqrt[3]{81Z^2+9Z+1}x^6-5850(x^3-1)^{2/3}x^5-113004\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x^4-27813x^6-60444\sqrt[3]{81Z^2+9Z+1}^2x^3+271341\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{2/3}x^2+8322(x^3-1)^{1/3}x^4-190603\sqrt[3]{81Z^2+9Z+1}x^3-16425(x^3-1)^{2/3}x^2-106434\sqrt[3]{81Z^2+9Z+1}(x^3-1)^{1/3}x-27813x^3+4161(x^3-1)^{1/3}x+181332\sqrt[3]{81Z^2+9Z+1}+27813}{(x^3+x^2-2x-3)/(x^6-x^5+3x^4-4x^3+7x^2-6x+9)}\right)$

Fricas [A] (verification not implemented)

none

Time = 3.21 (sec) , antiderivative size = 174, normalized size of antiderivative = 0.90

$$\int \frac{3 + 8x^3 - 3x^6 + x^9}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx = \frac{1}{9} \sqrt{3} \arctan \left(\frac{258 \sqrt{3} (2x^5 - 59x^2)(x^3 - 1)^{2/3} - 258 \sqrt{3} (13x^7 - 56x^4 - 24x)(x^3 - 1)^{1/3} - \sqrt{3} (169x^9 + 7789x^6 + 7135x^3 - 10368)}{2197x^9 + 13021x^6 - 25667x^3 + 13824} \right) + \frac{1}{18} \log \left(\frac{x^9 - 83x^6 + 82x^3 + 18(2x^5 - 5x^2)(x^3 - 1)^{2/3} - 9(x^7 - 13x^4 + 3x)(x^3 - 1)^{1/3} - 27}{x^9 - 2x^6 + x^3 - 27} \right)$$

[In] integrate((x^9-3*x^6+8*x^3+3)/((x^3-1)^(2/3)*(x^9-2*x^6+x^3-27)),x, algorithm="fricas")

[Out] 1/9*sqrt(3)*arctan((258*sqrt(3)*(2*x^5 - 59*x^2)*(x^3 - 1)^(2/3) - 258*sqrt(3)*(13*x^7 + 56*x^4 - 24*x)*(x^3 - 1)^(1/3) - sqrt(3)*(169*x^9 + 7789*x^6 + 7135*x^3 - 10368))/(2197*x^9 + 13021*x^6 - 25667*x^3 + 13824)) + 1/18*log((x^9 - 83*x^6 + 82*x^3 + 18*(2*x^5 - 5*x^2)*(x^3 - 1)^(2/3) - 9*(x^7 - 13*x^4 + 3*x)*(x^3 - 1)^(1/3) - 27)/(x^9 - 2*x^6 + x^3 - 27))

Sympy [F]

$$\int \frac{3 + 8x^3 - 3x^6 + x^9}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx = \int \frac{x^9 - 3x^6 + 8x^3 + 3}{((x - 1)(x^2 + x + 1))^{2/3} (x^3 + x^2 - 2x - 3)(x^6 - x^5 + 3x^4 - 4x^3 + 7x^2 - 6x + 9)} dx$$

[In] integrate((x**9-3*x**6+8*x**3+3)/((x**3-1)**(2/3)*(x**9-2*x**6+x**3-27)),x)

[Out] Integral((x**9 - 3*x**6 + 8*x**3 + 3)/(((x - 1)*(x**2 + x + 1))**(2/3)*(x**3 + x**2 - 2*x - 3)*(x**6 - x**5 + 3*x**4 - 4*x**3 + 7*x**2 - 6*x + 9)), x)

Maxima [F]

$$\int \frac{3 + 8x^3 - 3x^6 + x^9}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx = \int \frac{x^9 - 3x^6 + 8x^3 + 3}{(x^9 - 2x^6 + x^3 - 27)(x^3 - 1)^{2/3}} dx$$

[In] integrate((x^9-3*x^6+8*x^3+3)/((x^3-1)^(2/3)*(x^9-2*x^6+x^3-27)),x, algorithm="maxima")

[Out] integrate((x^9 - 3*x^6 + 8*x^3 + 3)/((x^9 - 2*x^6 + x^3 - 27)*(x^3 - 1)^(2/3)), x)

Giac [F]

$$\int \frac{3 + 8x^3 - 3x^6 + x^9}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx = \int \frac{x^9 - 3x^6 + 8x^3 + 3}{(x^9 - 2x^6 + x^3 - 27)(x^3 - 1)^{2/3}} dx$$

[In] integrate((x^9-3*x^6+8*x^3+3)/((x^3-1)^(2/3)*(x^9-2*x^6+x^3-27)),x, algorithm="giac")

[Out] integrate((x^9 - 3*x^6 + 8*x^3 + 3)/((x^9 - 2*x^6 + x^3 - 27)*(x^3 - 1)^(2/3)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{3 + 8x^3 - 3x^6 + x^9}{(-1 + x^3)^{2/3} (-27 + x^3 - 2x^6 + x^9)} dx = \int \frac{x^9 - 3x^6 + 8x^3 + 3}{(x^3 - 1)^{2/3} (x^9 - 2x^6 + x^3 - 27)} dx$$

[In] int((8*x^3 - 3*x^6 + x^9 + 3)/((x^3 - 1)^(2/3)*(x^3 - 2*x^6 + x^9 - 27)),x)

[Out] int((8*x^3 - 3*x^6 + x^9 + 3)/((x^3 - 1)^(2/3)*(x^3 - 2*x^6 + x^9 - 27)), x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 145

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*                               Small rewrite of logic in main function to make it*)
(*                               match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)
```

```

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal},
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is"}
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<}
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

```

```

    finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],

```

```

9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions

```

```

#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022  add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

      if debug then
          print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
      fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

```

```

if ExpnType_result<=ExpnType_optimal then
  if debug then
    print("ExpnType_result<=ExpnType_optimal");
  fi;
  if is_contains_complex(result) then
    if is_contains_complex(optimal) then
      if debug then
        print("both result and optimal complex");
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        return "A"," ";
      else
        return "B",cat("Both result and optimal contain complex but leaf count of r
          convert(leaf_count_result,string)," vs. $2 (" ,
          convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
        end if
      else #result contains complex but optimal is not
        if debug then
          print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
      fi;
    else # result do not contain complex
      # this assumes optimal do not as well. No check is needed here.
      if debug then
        print("result do not contain complex, this assumes optimal do not as well")
      fi;
      if leaf_count_result<=2*leaf_count_optimal then
        if debug then
          print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
      else
        if debug then
          print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of op
          convert(leaf_count_result,string)," $ vs. $2(",
          convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
        fi;
      fi;
    else #ExpnType(result) > ExpnType(optimal)
      if debug then
        print("ExpnType(result) > ExpnType(optimal)");

```

```

    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'^^') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then

```

```

    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  else
    max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
  end if
elif type(expn,`+`) or type(expn,`*`) then
  max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif ElementaryFunctionQ(op(0,expn)) then
  max(3,ExpnType(op(1,expn)))
elif SpecialFunctionQ(op(0,expn)) then
  max(4,apply(max,map(ExpnType,[op(expn)])))
elif HypergeometricFunctionQ(op(0,expn)) then
  max(5,apply(max,map(ExpnType,[op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6,apply(max,map(ExpnType,[op(expn)])))
elif op(0,expn)='int' then
  max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)

```



```

    member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
    member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
    if nops(u)=2 then
        op(2,u)
    else
        apply(op(0,u), op(2..nops(u), u))
    end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma] [LeafCount] (u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr, Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:

```

```

    return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:

```

```

    return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
elif isinstance(expn,Pow): #type(expn,``^`)
    if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
        return expnType(expn.args[0]) #ExpnType(op(1,expn))
    elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,``+`) or type(expn,``*`)
    m1 = expnType(expn.args[0])
    m2 = expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
    return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

```

```

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""
else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    #debug=False

```

```

m = func.name() in ['exp','log','ln',
    'sin','cos','tan','cot','sec','csc',
    'arcsin','arccos','arctan','arccot','arcsec','arccsc',
    'sinh','cosh','tanh','coth','sech','csch',
    'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
    'arctan2','floor','abs'
]
if debug:
    if m:
        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
    'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
    'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
    'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
    'elliptic_pi','exp_integral_e','log_integral']

if debug:
    print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("cought exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)

```

```

    return expnType(expn.operands()[0]) #expnType(expn.args[0])
elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
    if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)
        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```
def grade_antiderivative(result,optimal):
```

```
if debug:
```

```

    print ("Enter grade_antiderivative for sagemath")
    print("Enter grade_antiderivative, result=",result)
    print("Enter grade_antiderivative, optimal=",optimal)
    print("type(anti)=",type(result))
    print("type(optimal)=",type(optimal))

```

```
leaf_count_result = tree_size(result) #leaf_count(result)
```

```
leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)
```



```

#if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```