

Computer Algebra Independent Integration Tests

Summer 2023 edition

4-Trig-functions/4.2-Cosine/82-4.2.0-a-cos-^m-b-trg-ⁿ

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 11:39am

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	97
4	Appendix	1443

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [294]. This is test number [82].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (294)	0.00 (0)
Mathematica	100.00 (294)	0.00 (0)
Fricas	67.01 (197)	32.99 (97)
Maple	66.67 (196)	33.33 (98)
Maxima	31.29 (92)	68.71 (202)
Mupad	27.21 (80)	72.79 (214)
Giac	13.27 (39)	86.73 (255)
Sympy	6.12 (18)	93.88 (276)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	97.619	0.340	2.041	0.000
Maple	39.116	24.490	3.061	33.333
Fricas	29.932	1.701	35.374	32.993
Maxima	21.088	10.204	0.000	68.707
Giac	10.544	2.721	0.000	86.735
Sympy	3.741	2.381	0.000	93.878
Mupad	0.000	27.211	0.000	72.789

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima

and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	97	93.81	0.00	6.19
Maple	98	98.98	1.02	0.00
Maxima	202	99.50	0.00	0.50
Mupad	214	0.00	100.00	0.00
Giac	255	99.61	0.39	0.00
Sympy	276	47.10	52.90	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.05
Fricas	0.20
Maxima	0.38
Mathematica	0.75
Giac	1.24
Maple	3.29
Sympy	5.12
Mupad	11.39

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	54.98	1.09	43.50	0.87
Mathematica	59.26	0.90	55.00	0.91
Sympy	60.28	1.52	46.00	1.43
Rubi	68.33	1.00	69.00	1.00
Fricas	91.11	1.45	91.00	1.21
Maple	139.54	1.99	106.00	1.66
Maxima	221.09	2.81	48.50	0.86
Giac	5516.31	76.40	34.00	0.76

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {284, 285, 286, 287, 292}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v1.0a

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	27
2.3	Detailed conclusion table specific for Rubi results	87

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	23
Maple	23
Fricas	24
Maxima	24
Giac	25
Mupad	25
Sympy	26

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 277, 278, 279, 280, 281, 282, 283, 288, 289, 290, 291, 293, 294 }

B grade { 1 }

C grade { 276, 284, 285, 286, 287, 292 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 17, 39, 40, 41, 44, 51, 52, 53, 54, 55, 56, 64, 66, 68, 78, 80, 90, 92, 102, 103, 104, 105, 115, 116, 117, 127, 128, 129, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 275, 276, 277, 278 }

B grade { 9, 10, 11, 12, 14, 15, 16, 18, 19, 20, 22, 23, 24, 42, 43, 67, 69, 70, 71, 72, 73, 74, 75, 76, 77, 79, 81, 82, 83, 84, 85, 86, 87, 88, 89, 91, 93, 94, 95, 96, 97, 98, 99, 100, 106, 107, 108, 110, 111, 112, 113, 114, 118, 119, 120, 121, 122, 123, 124, 125, 126, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 274 }

C grade { 13, 21, 45, 46, 47, 48, 49, 50, 109 }

F normal fail { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 57, 58, 59, 60, 61, 62, 63, 65, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

F(-1) timedout fail { 101 }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 43, 44, 51, 52, 53, 54, 55, 56, 64, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 261, 262, 265, 266, 267, 270 }

B grade { 42, 271, 272, 275, 276 }

C grade { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 45, 46, 47, 48, 49, 50, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 263, 264, 268, 269, 273, 274, 277, 278 }

F normal fail { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 57, 65, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

F(-1) timeout fail { }

F(-2) exception fail { 58, 59, 60, 61, 62, 63 }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 51, 52, 53, 54, 55, 56, 140, 141, 142, 143, 144, 146, 150, 151, 152, 153, 154, 156, 160, 161, 162, 163, 164, 165, 167, 171, 172, 173, 174, 175, 176, 182, 183, 184, 185, 186, 192, 193, 194, 195, 196, 261, 262, 265, 266, 267, 270, 271, 272, 275, 276 }

B grade { 42, 43, 44, 145, 147, 148, 149, 155, 157, 158, 159, 166, 168, 169, 170, 177, 178, 179, 180, 181, 187, 188, 189, 190, 191, 197, 198, 199, 200, 201 }

C grade { }

F normal fail { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 45, 46, 47, 48, 49, 50, 57, 58, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 273, 274, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

F(-1) timeout fail { }

F(-2) exception fail { 59 }

Giac

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 43, 44, 51, 52, 53, 54, 55, 56, 143, 152, 261, 262, 265, 266, 267, 270, 271, 272, 275, 276 }

B grade { 64, 140, 141, 142, 150, 151, 161, 162 }

C grade { }

F normal fail { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 45, 46, 47, 48, 49, 50, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 144, 145, 146, 147, 148, 149, 153, 154, 155, 156, 157, 158, 159, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 273, 274, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

F(-1) timeout fail { 160 }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 21, 25, 26, 27, 28, 29, 30, 37, 41, 54, 55, 56, 64, 107, 108, 109, 140, 141, 142, 143, 144, 146, 148, 150, 151, 152, 153, 154, 156, 158, 160, 161, 162, 163, 164, 165, 167, 169, 171, 172, 173, 174, 175, 176, 178, 180, 182, 183, 184, 185, 186, 188, 190, 192, 193, 194, 195, 196, 198, 200, 261, 262, 265, 270 }

C grade { }

F normal fail { }

F(-1) timeout fail { 17, 18, 19, 20, 22, 23, 24, 31, 32, 33, 34, 35, 36, 38, 39, 40, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 57, 58, 59, 60, 61, 62, 63, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 145, 147, 149, 155, 157, 159, 166, 168, 170, 177, 179, 181, 187, 189, 191, 197, 199, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 268, 269, 273, 274, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 263, 264, 266, 267, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294 }

F(-2) exception fail { }

Sympy

A grade { 1, 3, 5, 7, 41, 144, 153, 154, 175, 176, 186 }

B grade { 2, 4, 6, 8, 64, 142, 143 }

C grade { }

F normal fail { 11, 12, 13, 14, 15, 19, 20, 21, 22, 23, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 42, 43, 47, 48, 57, 59, 60, 61, 62, 63, 65, 69, 70, 71, 72, 73, 74, 75, 82, 107, 108, 109, 110, 111, 112, 113, 114, 121, 122, 123, 124, 125, 126, 135, 145, 146, 152, 177, 178, 187, 188, 189, 198, 202, 204, 205, 206, 207, 208, 209, 212, 213, 223, 224, 225, 226, 227, 228, 229, 230, 232, 233, 234, 235, 236, 237, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 253, 254, 255, 256, 257, 260, 261, 262, 263, 264, 268, 269, 271, 272, 273, 274, 275, 276, 277, 278, 280, 281, 282, 284, 285, 286, 287, 291, 292, 293 }

F(-1) timedout fail { 9, 10, 16, 17, 18, 24, 25, 31, 39, 40, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 58, 66, 67, 68, 76, 77, 78, 79, 80, 81, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 115, 116, 117, 118, 119, 120, 127, 128, 129, 130, 131, 132, 133, 134, 136, 137, 138, 139, 140, 141, 147, 148, 149, 150, 151, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 179, 180, 181, 182, 183, 184, 185, 190, 191, 192, 193, 194, 195, 196, 197, 199, 200, 201, 203, 210, 211, 214, 215, 216, 217, 218, 219, 220, 221, 222, 231, 238, 239, 252, 258, 259, 265, 266, 267, 270, 279, 283, 288, 289, 290, 294 }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	21	11	10	10	12	10	10
N.S.	1	1.00	2.10	1.10	1.00	1.00	1.20	1.00	1.00
time (sec)	N/A	0.006	0.012	0.268	0.239	0.265	0.060	0.294	14.677

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	23	19	22	22	46	18	18
N.S.	1	1.00	0.92	0.76	0.88	0.88	1.84	0.72	0.72
time (sec)	N/A	0.012	0.028	0.476	0.255	0.310	0.091	0.289	15.283

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	26	22	22	21	36	22	24
N.S.	1	1.00	1.00	0.85	0.85	0.81	1.38	0.85	0.92
time (sec)	N/A	0.014	0.010	0.911	0.254	0.262	0.124	0.293	13.520

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	33	31	33	36	95	32	31
N.S.	1	1.00	0.72	0.67	0.72	0.78	2.07	0.70	0.67
time (sec)	N/A	0.024	0.047	0.754	0.251	0.271	0.175	0.299	13.694

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	32	34	33	58	34	31
N.S.	1	1.00	1.00	0.78	0.83	0.80	1.41	0.83	0.76
time (sec)	N/A	0.016	0.016	0.996	0.240	0.286	0.238	0.296	0.054

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	43	42	48	46	139	46	42
N.S.	1	1.00	0.64	0.63	0.72	0.69	2.07	0.69	0.63
time (sec)	N/A	0.039	0.046	1.135	0.231	0.271	0.375	0.294	14.250

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	54	42	44	43	78	44	43
N.S.	1	1.00	1.00	0.78	0.81	0.80	1.44	0.81	0.80
time (sec)	N/A	0.020	0.015	1.276	0.244	0.270	0.508	0.319	13.988

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	55	55	59	56	184	60	53
N.S.	1	1.00	0.62	0.62	0.67	0.64	2.09	0.68	0.60
time (sec)	N/A	0.055	0.062	1.148	0.260	0.279	0.710	0.293	14.508

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	51	199	0	80	0	0	42
N.S.	1	1.00	0.78	3.06	0.00	1.23	0.00	0.00	0.65
time (sec)	N/A	0.034	0.124	4.632	0.000	0.096	0.000	0.000	14.502

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	40	202	0	74	0	0	42
N.S.	1	1.00	0.95	4.81	0.00	1.76	0.00	0.00	1.00
time (sec)	N/A	0.023	0.064	2.886	0.000	0.102	0.000	0.000	14.441

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	36	179	0	68	0	0	35
N.S.	1	1.00	0.86	4.26	0.00	1.62	0.00	0.00	0.83
time (sec)	N/A	0.023	0.052	2.154	0.000	0.091	0.000	0.000	14.407

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	133	0	57	0	0	15
N.S.	1	1.00	1.00	8.31	0.00	3.56	0.00	0.00	0.94
time (sec)	N/A	0.009	0.033	1.612	0.000	0.086	0.000	0.000	14.414

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	18	0	51	0	0	15
N.S.	1	1.00	1.00	1.12	0.00	3.19	0.00	0.00	0.94
time (sec)	N/A	0.012	0.038	0.246	0.000	0.090	0.000	0.000	13.594

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	182	0	93	0	0	42
N.S.	1	1.00	1.00	4.79	0.00	2.45	0.00	0.00	1.11
time (sec)	N/A	0.021	0.074	1.302	0.000	0.096	0.000	0.000	14.078

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	36	213	0	92	0	0	42
N.S.	1	1.00	0.86	5.07	0.00	2.19	0.00	0.00	1.00
time (sec)	N/A	0.023	0.072	1.456	0.000	0.087	0.000	0.000	14.248

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	59	358	0	110	0	0	42
N.S.	1	1.00	0.91	5.51	0.00	1.69	0.00	0.00	0.65
time (sec)	N/A	0.035	0.116	2.307	0.000	0.090	0.000	0.000	14.850

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	76	210	0	95	0	0	0
N.S.	1	1.00	0.78	2.14	0.00	0.97	0.00	0.00	0.00
time (sec)	N/A	0.056	0.126	4.013	0.000	0.096	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	62	213	0	91	0	0	0
N.S.	1	1.00	0.89	3.04	0.00	1.30	0.00	0.00	0.00
time (sec)	N/A	0.039	0.105	2.784	0.000	0.096	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	58	190	0	77	0	0	0
N.S.	1	1.00	0.83	2.71	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.040	0.065	2.322	0.000	0.093	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	142	0	63	0	0	0
N.S.	1	1.00	1.00	3.74	0.00	1.66	0.00	0.00	0.00
time (sec)	N/A	0.023	0.036	1.915	0.000	0.089	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	54	0	60	0	0	33
N.S.	1	1.00	1.00	1.42	0.00	1.58	0.00	0.00	0.87
time (sec)	N/A	0.021	0.042	0.388	0.000	0.082	0.000	0.000	0.157

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	50	198	0	104	0	0	0
N.S.	1	1.00	0.74	2.91	0.00	1.53	0.00	0.00	0.00
time (sec)	N/A	0.038	0.048	1.450	0.000	0.093	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	103	0	0	0
N.S.	1	1.00	0.71	3.35	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.041	0.082	1.646	0.000	0.093	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	53	53	33	32	31	40	0	34	0
N.S.	1	1.00	0.62	0.60	0.58	0.75	0.00	0.64	0.00
time (sec)	N/A	0.052	0.015	1.229	0.365	0.296	0.000	0.347	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	23	24	17	26	0	17	0
N.S.	1	1.00	0.68	0.71	0.50	0.76	0.00	0.50	0.00
time (sec)	N/A	0.036	0.008	0.783	0.388	0.301	0.000	0.319	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	15	6	15	15	9	46
N.S.	1	1.00	1.00	1.15	0.46	1.15	1.15	0.69	3.54
time (sec)	N/A	0.016	0.005	0.819	0.375	0.296	0.230	0.378	14.477

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	16	16	16	48	38	65	0	0	0
N.S.	1	1.00	1.00	3.00	2.38	4.06	0.00	0.00	0.00
time (sec)	N/A	0.018	0.004	0.905	0.385	0.284	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	26	70	304	40	0	20	0
N.S.	1	1.00	0.62	1.67	7.24	0.95	0.00	0.48	0.00
time (sec)	N/A	0.031	0.013	1.045	0.395	0.284	0.000	0.355	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	36	89	933	49	0	29	0
N.S.	1	1.00	0.59	1.46	15.30	0.80	0.00	0.48	0.00
time (sec)	N/A	0.064	0.038	1.077	0.622	0.300	0.000	0.311	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	61	117	0	92	0	0	0
N.S.	1	1.00	0.52	1.00	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.080	0.152	12.326	0.000	0.122	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	50	271	0	66	0	0	0
N.S.	1	1.00	0.75	4.04	0.00	0.99	0.00	0.00	0.00
time (sec)	N/A	0.057	0.099	6.035	0.000	0.123	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	44	44	37	92	0	59	0	0	0
N.S.	1	1.00	0.84	2.09	0.00	1.34	0.00	0.00	0.00
time (sec)	N/A	0.024	0.046	2.910	0.000	0.089	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	31	244	0	71	0	0	0
N.S.	1	1.00	0.74	5.81	0.00	1.69	0.00	0.00	0.00
time (sec)	N/A	0.025	0.039	3.470	0.000	0.101	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	44	101	0	74	0	0	0
N.S.	1	1.00	0.62	1.42	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.033	0.086	5.052	0.000	0.100	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	57	297	0	92	0	0	0
N.S.	1	1.00	0.49	2.54	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.054	0.135	8.308	0.000	0.131	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	132	132	53	56	85	68	0	57	0
N.S.	1	1.00	0.40	0.42	0.64	0.52	0.00	0.43	0.00
time (sec)	N/A	0.057	0.151	13.264	0.329	0.278	0.000	0.331	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	38	38	55	42	0	25	0
N.S.	1	1.00	0.49	0.49	0.71	0.54	0.00	0.32	0.00
time (sec)	N/A	0.034	0.097	0.821	0.352	0.291	0.000	0.295	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	25	20	22	21	0	13	0
N.S.	1	1.00	0.69	0.56	0.61	0.58	0.00	0.36	0.00
time (sec)	N/A	0.020	0.034	1.013	0.471	0.260	0.000	0.314	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	44	0	15	61	300	40
N.S.	1	1.00	1.00	1.83	0.00	0.62	2.54	12.50	1.67
time (sec)	N/A	0.022	0.047	1.199	0.000	0.264	0.470	0.565	14.224

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	74	0	0	0	0	0	0
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.049	0.062	0.000	0.000	0.000	0.000	0.000	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	123	83	234	0	98	0	0	0
N.S.	1	1.00	0.67	1.90	0.00	0.80	0.00	0.00	0.00
time (sec)	N/A	0.092	0.355	6.764	0.000	0.113	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	75	221	0	101	0	0	0
N.S.	1	1.00	0.77	2.28	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.083	0.286	4.485	0.000	0.122	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	73	208	0	88	0	0	0
N.S.	1	1.00	0.77	2.19	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	0.066	0.263	3.465	0.000	0.114	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	62	211	0	88	0	0	0
N.S.	1	1.00	0.90	3.06	0.00	1.28	0.00	0.00	0.00
time (sec)	N/A	0.045	0.163	2.844	0.000	0.122	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	61	188	0	76	0	0	0
N.S.	1	1.00	0.91	2.81	0.00	1.13	0.00	0.00	0.00
time (sec)	N/A	0.044	0.079	2.211	0.000	0.088	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	142	0	63	0	0	0
N.S.	1	1.00	1.00	3.74	0.00	1.66	0.00	0.00	0.00
time (sec)	N/A	0.019	0.039	2.045	0.000	0.091	0.000	0.000	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	142	0	57	0	0	0
N.S.	1	1.00	1.00	3.64	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.032	0.048	1.437	0.000	0.088	0.000	0.000	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	48	196	0	101	0	0	0
N.S.	1	1.00	0.76	3.11	0.00	1.60	0.00	0.00	0.00
time (sec)	N/A	0.058	0.200	1.782	0.000	0.112	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	49	239	0	100	0	0	0
N.S.	1	1.00	0.70	3.41	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.065	0.224	1.837	0.000	0.105	0.000	0.000	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	69	364	0	118	0	0	0
N.S.	1	1.00	0.73	3.83	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.094	0.375	2.546	0.000	0.095	0.000	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	69	396	0	112	0	0	0
N.S.	1	1.00	0.70	4.04	0.00	1.14	0.00	0.00	0.00
time (sec)	N/A	0.093	0.336	2.135	0.000	0.097	0.000	0.000	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	123	79	414	0	128	0	0	0
N.S.	1	1.00	0.64	3.37	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.125	0.388	3.063	0.000	0.098	0.000	0.000	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	83	236	0	102	0	0	0
N.S.	1	1.00	0.66	1.87	0.00	0.81	0.00	0.00	0.00
time (sec)	N/A	0.110	0.264	5.704	0.000	0.102	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	75	223	0	103	0	0	0
N.S.	1	1.00	0.79	2.35	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	0.075	0.266	4.483	0.000	0.102	0.000	0.000	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	73	210	0	91	0	0	0
N.S.	1	1.00	0.74	2.14	0.00	0.93	0.00	0.00	0.00
time (sec)	N/A	0.078	0.202	3.411	0.000	0.100	0.000	0.000	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	65	213	0	89	0	0	0
N.S.	1	1.00	0.97	3.18	0.00	1.33	0.00	0.00	0.00
time (sec)	N/A	0.044	0.108	2.671	0.000	0.157	0.000	0.000	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	58	190	0	77	0	0	0
N.S.	1	1.00	0.83	2.71	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.037	0.045	2.162	0.000	0.095	0.000	0.000	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	144	0	63	0	0	0
N.S.	1	1.00	1.00	3.69	0.00	1.62	0.00	0.00	0.00
time (sec)	N/A	0.031	0.014	1.798	0.000	0.097	0.000	0.000	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	57	0	0	0
N.S.	1	1.00	1.00	3.51	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.048	0.025	1.269	0.000	0.102	0.000	0.000	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	50	198	0	102	0	0	0
N.S.	1	1.00	0.76	3.00	0.00	1.55	0.00	0.00	0.00
time (sec)	N/A	0.064	0.222	1.782	0.000	0.112	0.000	0.000	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	101	0	0	0
N.S.	1	1.00	0.71	3.35	0.00	1.40	0.00	0.00	0.00
time (sec)	N/A	0.064	0.177	1.781	0.000	0.089	0.000	0.000	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	69	365	0	119	0	0	0
N.S.	1	1.00	0.70	3.72	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.088	0.267	2.375	0.000	0.105	0.000	0.000	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	69	398	0	115	0	0	0
N.S.	1	1.00	0.69	3.98	0.00	1.15	0.00	0.00	0.00
time (sec)	N/A	0.081	0.374	2.136	0.000	0.104	0.000	0.000	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	79	416	0	132	0	0	0
N.S.	1	1.00	0.63	3.30	0.00	1.05	0.00	0.00	0.00
time (sec)	N/A	0.108	0.553	2.990	0.000	0.105	0.000	0.000	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	83	236	0	108	0	0	0
N.S.	1	1.00	0.66	1.89	0.00	0.86	0.00	0.00	0.00
time (sec)	N/A	0.088	0.364	10.455	0.000	0.112	0.000	0.000	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	75	223	0	107	0	0	0
N.S.	1	1.00	0.77	2.28	0.00	1.09	0.00	0.00	0.00
time (sec)	N/A	0.071	0.256	5.656	0.000	0.105	0.000	0.000	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	76	210	0	95	0	0	0
N.S.	1	1.00	0.78	2.16	0.00	0.98	0.00	0.00	0.00
time (sec)	N/A	0.066	0.145	3.607	0.000	0.097	0.000	0.000	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	62	213	0	91	0	0	0
N.S.	1	1.00	0.89	3.04	0.00	1.30	0.00	0.00	0.00
time (sec)	N/A	0.036	0.093	2.853	0.000	0.108	0.000	0.000	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	59	190	0	79	0	0	0
N.S.	1	1.00	0.82	2.64	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.056	0.048	2.771	0.000	0.099	0.000	0.000	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	63	0	0	0
N.S.	1	1.00	1.00	3.51	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.043	0.017	6.788	0.000	0.108	0.000	0.000	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	38	144	0	57	0	0	0
N.S.	1	1.00	0.93	3.51	0.00	1.39	0.00	0.00	0.00
time (sec)	N/A	0.039	0.208	27.801	0.000	0.093	0.000	0.000	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	50	198	0	104	0	0	0
N.S.	1	1.00	0.74	2.91	0.00	1.53	0.00	0.00	0.00
time (sec)	N/A	0.060	0.177	98.968	0.000	0.089	0.000	0.000	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	103	0	0	0
N.S.	1	1.00	0.71	3.35	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.066	0.251	1.135	0.000	0.105	0.000	0.000	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	69	367	0	123	0	0	0
N.S.	1	1.00	0.69	3.67	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.084	0.431	1.673	0.000	0.114	0.000	0.000	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	69	398	0	119	0	0	0
N.S.	1	1.00	0.69	3.98	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	0.088	0.682	1.065	0.000	0.101	0.000	0.000	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F(-1)	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	128	128	79	0	0	138	0	0	0
N.S.	1	1.00	0.62	0.00	0.00	1.08	0.00	0.00	0.00
time (sec)	N/A	0.112	0.852	180.000	0.000	0.104	0.000	0.000	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	76	210	0	95	0	0	0
N.S.	1	1.00	0.78	2.14	0.00	0.97	0.00	0.00	0.00
time (sec)	N/A	0.059	0.086	3.615	0.000	0.097	0.000	0.000	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	73	233	0	101	0	0	0
N.S.	1	1.00	0.58	1.86	0.00	0.81	0.00	0.00	0.00
time (sec)	N/A	0.085	0.275	4.563	0.000	0.109	0.000	0.000	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	71	220	0	104	0	0	0
N.S.	1	1.00	0.71	2.20	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.070	0.227	3.905	0.000	0.120	0.000	0.000	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	63	207	0	91	0	0	0
N.S.	1	1.00	0.65	2.13	0.00	0.94	0.00	0.00	0.00
time (sec)	N/A	0.062	0.197	3.197	0.000	0.109	0.000	0.000	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	58	210	0	91	0	0	0
N.S.	1	1.00	0.81	2.92	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.046	0.190	2.570	0.000	0.096	0.000	0.000	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	51	187	0	79	0	0	58
N.S.	1	1.00	0.74	2.71	0.00	1.14	0.00	0.00	0.84
time (sec)	N/A	0.049	0.152	1.791	0.000	0.094	0.000	0.000	13.170

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	141	0	66	0	0	33
N.S.	1	1.00	1.00	3.44	0.00	1.61	0.00	0.00	0.80
time (sec)	N/A	0.023	0.016	1.725	0.000	0.095	0.000	0.000	0.137

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	54	0	60	0	0	33
N.S.	1	1.00	1.00	1.42	0.00	1.58	0.00	0.00	0.87
time (sec)	N/A	0.024	0.016	0.476	0.000	0.088	0.000	0.000	0.116

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	47	195	0	104	0	0	0
N.S.	1	1.00	0.72	3.00	0.00	1.60	0.00	0.00	0.00
time (sec)	N/A	0.055	0.063	1.904	0.000	0.095	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	48	238	0	103	0	0	0
N.S.	1	1.00	0.72	3.55	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.066	0.094	1.790	0.000	0.102	0.000	0.000	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	65	367	0	121	0	0	0
N.S.	1	1.00	0.67	3.78	0.00	1.25	0.00	0.00	0.00
time (sec)	N/A	0.088	0.231	2.418	0.000	0.099	0.000	0.000	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	63	267	0	115	0	0	0
N.S.	1	1.00	0.66	2.81	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.091	0.256	2.524	0.000	0.103	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	77	413	0	131	0	0	0
N.S.	1	1.00	0.62	3.30	0.00	1.05	0.00	0.00	0.00
time (sec)	N/A	0.118	0.447	3.185	0.000	0.100	0.000	0.000	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	128	76	236	0	101	0	0	0
N.S.	1	1.00	0.59	1.84	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.087	0.423	5.527	0.000	0.106	0.000	0.000	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	74	223	0	104	0	0	0
N.S.	1	1.00	0.74	2.23	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.079	0.279	4.592	0.000	0.107	0.000	0.000	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	66	210	0	91	0	0	0
N.S.	1	1.00	0.66	2.10	0.00	0.91	0.00	0.00	0.00
time (sec)	N/A	0.068	0.215	3.416	0.000	0.126	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	61	213	0	91	0	0	0
N.S.	1	1.00	0.85	2.96	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.047	0.188	2.853	0.000	0.105	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	54	190	0	79	0	0	0
N.S.	1	1.00	0.75	2.64	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.045	0.186	2.251	0.000	0.095	0.000	0.000	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	66	0	0	0
N.S.	1	1.00	1.00	3.51	0.00	1.61	0.00	0.00	0.00
time (sec)	N/A	0.027	0.018	2.027	0.000	0.097	0.000	0.000	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	60	0	0	0
N.S.	1	1.00	1.00	3.51	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.030	0.025	1.393	0.000	0.083	0.000	0.000	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	50	198	0	104	0	0	0
N.S.	1	1.00	0.74	2.91	0.00	1.53	0.00	0.00	0.00
time (sec)	N/A	0.046	0.032	1.742	0.000	0.099	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	51	241	0	103	0	0	0
N.S.	1	1.00	0.74	3.49	0.00	1.49	0.00	0.00	0.00
time (sec)	N/A	0.070	0.055	2.150	0.000	0.095	0.000	0.000	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	68	367	0	121	0	0	0
N.S.	1	1.00	0.69	3.74	0.00	1.23	0.00	0.00	0.00
time (sec)	N/A	0.132	0.133	2.653	0.000	0.098	0.000	0.000	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	66	398	0	115	0	0	0
N.S.	1	1.00	0.68	4.10	0.00	1.19	0.00	0.00	0.00
time (sec)	N/A	0.115	0.257	2.339	0.000	0.128	0.000	0.000	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	80	416	0	131	0	0	0
N.S.	1	1.00	0.63	3.30	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.168	0.346	3.118	0.000	0.107	0.000	0.000	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	128	76	236	0	101	0	0	0
N.S.	1	1.00	0.59	1.84	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.149	0.632	5.879	0.000	0.104	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	74	223	0	104	0	0	0
N.S.	1	1.00	0.74	2.23	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.116	0.515	4.423	0.000	0.104	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	66	210	0	91	0	0	0
N.S.	1	1.00	0.66	2.10	0.00	0.91	0.00	0.00	0.00
time (sec)	N/A	0.114	0.318	3.774	0.000	0.097	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	61	213	0	91	0	0	0
N.S.	1	1.00	0.85	2.96	0.00	1.26	0.00	0.00	0.00
time (sec)	N/A	0.064	0.229	2.813	0.000	0.100	0.000	0.000	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	54	190	0	79	0	0	0
N.S.	1	1.00	0.75	2.64	0.00	1.10	0.00	0.00	0.00
time (sec)	N/A	0.046	0.213	2.445	0.000	0.099	0.000	0.000	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	144	0	66	0	0	0
N.S.	1	1.00	1.00	3.51	0.00	1.61	0.00	0.00	0.00
time (sec)	N/A	0.028	0.018	1.914	0.000	0.100	0.000	0.000	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	38	144	0	60	0	0	0
N.S.	1	1.00	0.93	3.51	0.00	1.46	0.00	0.00	0.00
time (sec)	N/A	0.029	0.197	1.387	0.000	0.088	0.000	0.000	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	50	198	0	104	0	0	0
N.S.	1	1.00	0.74	2.91	0.00	1.53	0.00	0.00	0.00
time (sec)	N/A	0.047	0.040	1.742	0.000	0.089	0.000	0.000	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	51	241	0	103	0	0	0
N.S.	1	1.00	0.71	3.35	0.00	1.43	0.00	0.00	0.00
time (sec)	N/A	0.049	0.046	1.664	0.000	0.094	0.000	0.000	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	68	367	0	121	0	0	0
N.S.	1	1.00	0.70	3.78	0.00	1.25	0.00	0.00	0.00
time (sec)	N/A	0.080	0.100	2.785	0.000	0.095	0.000	0.000	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	66	398	0	115	0	0	0
N.S.	1	1.00	0.67	4.06	0.00	1.17	0.00	0.00	0.00
time (sec)	N/A	0.085	0.169	2.233	0.000	0.098	0.000	0.000	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	80	416	0	131	0	0	0
N.S.	1	1.00	0.64	3.33	0.00	1.05	0.00	0.00	0.00
time (sec)	N/A	0.113	0.394	3.171	0.000	0.108	0.000	0.000	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	68	367	0	121	0	0	0
N.S.	1	1.00	0.68	3.67	0.00	1.21	0.00	0.00	0.00
time (sec)	N/A	0.058	0.015	2.587	0.000	0.105	0.000	0.000	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	55	62	49	176	0	203	75
N.S.	1	1.00	0.56	0.63	0.50	1.80	0.00	2.07	0.77
time (sec)	N/A	0.029	0.244	3.099	0.393	0.343	0.000	2.076	14.994

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	45	40	42	39	0	71047	57
N.S.	1	1.00	0.64	0.57	0.60	0.56	0.00	1014.96	0.81
time (sec)	N/A	0.020	0.101	3.174	0.459	0.303	0.000	9.095	0.815

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	45	42	25	150	121	105	62
N.S.	1	1.00	0.71	0.67	0.40	2.38	1.92	1.67	0.98
time (sec)	N/A	0.017	0.153	3.069	0.388	0.342	32.912	1.465	14.574

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	28	60	31	44
N.S.	1	1.00	1.00	0.91	0.41	0.88	1.88	0.97	1.38
time (sec)	N/A	0.009	0.116	3.070	0.439	0.281	1.301	0.540	14.172

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	21	26	94	22	0	20
N.S.	1	1.00	1.00	0.88	1.08	3.92	0.92	0.00	0.83
time (sec)	N/A	0.003	0.030	2.269	0.337	0.295	0.615	0.000	0.109

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	40	65	113	0	0	0
N.S.	1	1.00	1.00	1.21	1.97	3.42	0.00	0.00	0.00
time (sec)	N/A	0.009	0.015	3.228	0.449	0.390	0.000	0.000	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	54	28	0	0	59
N.S.	1	1.00	1.00	0.91	1.69	0.88	0.00	0.00	1.84
time (sec)	N/A	0.012	0.024	2.915	0.436	0.267	0.000	0.000	14.109

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	52	84	661	201	0	0	0
N.S.	1	1.00	0.72	1.17	9.18	2.79	0.00	0.00	0.00
time (sec)	N/A	0.021	0.054	3.088	0.443	0.341	0.000	0.000	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	45	42	294	41	0	0	128
N.S.	1	1.00	0.64	0.60	4.20	0.59	0.00	0.00	1.83
time (sec)	N/A	0.019	0.098	2.960	0.447	0.305	0.000	0.000	16.116

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	66	103	1656	227	0	0	0
N.S.	1	1.00	0.62	0.96	15.48	2.12	0.00	0.00	0.00
time (sec)	N/A	0.041	0.126	2.977	0.483	0.317	0.000	0.000	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	55	63	53	183	0	204	76
N.S.	1	1.00	0.54	0.62	0.52	1.81	0.00	2.02	0.75
time (sec)	N/A	0.033	0.203	2.814	0.399	0.311	0.000	2.432	14.934

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	45	41	45	43	0	71048	58
N.S.	1	1.00	0.62	0.57	0.62	0.60	0.00	986.78	0.81
time (sec)	N/A	0.018	0.138	2.793	0.398	0.281	0.000	11.242	0.680

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	45	43	28	153	0	106	63
N.S.	1	1.00	0.69	0.66	0.43	2.35	0.00	1.63	0.97
time (sec)	N/A	0.017	0.147	3.043	0.395	0.391	0.000	1.198	14.497

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	30	13	29	46	0	29
N.S.	1	1.00	0.97	0.91	0.39	0.88	1.39	0.00	0.88
time (sec)	N/A	0.009	0.131	2.968	0.370	0.297	15.888	0.000	0.256

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	22	26	95	22	0	21
N.S.	1	1.00	1.00	0.88	1.04	3.80	0.88	0.00	0.84
time (sec)	N/A	0.003	0.042	2.402	0.368	0.327	9.452	0.000	13.482

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	34	34	33	41	68	114	0	0	0
N.S.	1	1.00	0.97	1.21	2.00	3.35	0.00	0.00	0.00
time (sec)	N/A	0.009	0.019	2.773	0.422	0.304	0.000	0.000	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	32	30	54	29	0	0	60
N.S.	1	1.00	0.97	0.91	1.64	0.88	0.00	0.00	1.82
time (sec)	N/A	0.015	0.024	2.845	0.403	0.282	0.000	0.000	13.914

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	52	85	691	204	0	0	0
N.S.	1	1.00	0.70	1.15	9.34	2.76	0.00	0.00	0.00
time (sec)	N/A	0.024	0.057	2.869	0.438	0.339	0.000	0.000	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	45	43	299	42	0	0	129
N.S.	1	1.00	0.62	0.60	4.15	0.58	0.00	0.00	1.79
time (sec)	N/A	0.020	0.071	2.771	0.414	0.283	0.000	0.000	15.349

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	110	110	67	104	1742	234	0	0	0
N.S.	1	1.00	0.61	0.95	15.84	2.13	0.00	0.00	0.00
time (sec)	N/A	0.039	0.103	3.033	0.481	0.391	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	57	55	77	61	0	0	73
N.S.	1	1.00	0.49	0.47	0.66	0.53	0.00	0.00	0.63
time (sec)	N/A	0.027	0.164	3.744	0.406	0.297	0.000	0.000	15.647

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	55	65	59	193	0	203	78
N.S.	1	1.00	0.51	0.61	0.55	1.80	0.00	1.90	0.73
time (sec)	N/A	0.032	0.257	2.881	0.403	0.299	0.000	2.478	15.097

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	45	43	49	47	0	71047	60
N.S.	1	1.00	0.59	0.57	0.64	0.62	0.00	934.83	0.79
time (sec)	N/A	0.020	0.152	2.898	0.430	0.276	0.000	8.371	0.669

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	45	45	32	159	0	0	40
N.S.	1	1.00	0.65	0.65	0.46	2.30	0.00	0.00	0.58
time (sec)	N/A	0.017	0.170	3.197	0.448	0.326	0.000	0.000	0.420

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	13	31	0	0	31
N.S.	1	1.00	0.91	0.91	0.37	0.89	0.00	0.00	0.89
time (sec)	N/A	0.009	0.145	2.743	0.412	0.345	0.000	0.000	14.423

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	24	24	26	97	0	0	23
N.S.	1	1.00	0.89	0.89	0.96	3.59	0.00	0.00	0.85
time (sec)	N/A	0.003	0.047	2.335	0.381	0.309	0.000	0.000	0.099

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	43	72	116	0	0	0
N.S.	1	1.00	0.92	1.19	2.00	3.22	0.00	0.00	0.00
time (sec)	N/A	0.008	0.026	2.652	0.408	0.310	0.000	0.000	0.000

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	54	31	0	0	62
N.S.	1	1.00	0.91	0.91	1.54	0.89	0.00	0.00	1.77
time (sec)	N/A	0.015	0.023	2.753	0.418	0.285	0.000	0.000	14.493

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	52	87	747	210	0	0	0
N.S.	1	1.00	0.67	1.12	9.58	2.69	0.00	0.00	0.00
time (sec)	N/A	0.024	0.065	2.814	0.399	0.313	0.000	0.000	0.000

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	45	45	311	46	0	0	131
N.S.	1	1.00	0.59	0.59	4.09	0.61	0.00	0.00	1.72
time (sec)	N/A	0.020	0.066	2.666	0.383	0.352	0.000	0.000	14.507

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	66	106	1914	244	0	0	0
N.S.	1	1.00	0.57	0.91	16.50	2.10	0.00	0.00	0.00
time (sec)	N/A	0.041	0.151	3.241	0.505	0.319	0.000	0.000	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	57	52	68	54	0	0	73
N.S.	1	1.00	0.53	0.49	0.64	0.50	0.00	0.00	0.68
time (sec)	N/A	0.027	0.118	2.912	0.412	0.277	0.000	0.000	14.695

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	55	62	49	182	0	0	78
N.S.	1	1.00	0.56	0.63	0.50	1.86	0.00	0.00	0.80
time (sec)	N/A	0.028	0.245	3.167	0.394	0.297	0.000	0.000	14.392

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	45	40	42	42	0	0	60
N.S.	1	1.00	0.64	0.57	0.60	0.60	0.00	0.00	0.86
time (sec)	N/A	0.017	0.087	2.980	0.401	0.303	0.000	0.000	0.691

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	45	42	25	157	0	0	65
N.S.	1	1.00	0.71	0.67	0.40	2.49	0.00	0.00	1.03
time (sec)	N/A	0.017	0.163	3.095	0.403	0.305	0.000	0.000	14.015

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	13	31	46	0	47
N.S.	1	1.00	1.00	0.91	0.41	0.97	1.44	0.00	1.47
time (sec)	N/A	0.009	0.123	2.837	0.378	0.284	18.278	0.000	14.623

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	21	26	97	22	0	37
N.S.	1	1.00	1.00	0.88	1.08	4.04	0.92	0.00	1.54
time (sec)	N/A	0.003	0.029	2.344	0.308	0.302	0.670	0.000	0.315

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	40	65	116	0	0	0
N.S.	1	1.00	1.00	1.21	1.97	3.52	0.00	0.00	0.00
time (sec)	N/A	0.009	0.017	2.879	0.394	0.311	0.000	0.000	0.000

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	32	29	59	31	0	0	62
N.S.	1	1.00	1.00	0.91	1.84	0.97	0.00	0.00	1.94
time (sec)	N/A	0.014	0.026	2.850	0.377	0.292	0.000	0.000	14.506

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	72	72	52	84	661	207	0	0	0
N.S.	1	1.00	0.72	1.17	9.18	2.88	0.00	0.00	0.00
time (sec)	N/A	0.025	0.040	3.315	0.414	0.298	0.000	0.000	0.000

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	45	42	294	44	0	0	131
N.S.	1	1.00	0.64	0.60	4.20	0.63	0.00	0.00	1.87
time (sec)	N/A	0.021	0.101	2.827	0.408	0.350	0.000	0.000	15.040

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	66	103	1656	233	0	0	0
N.S.	1	1.00	0.62	0.96	15.48	2.18	0.00	0.00	0.00
time (sec)	N/A	0.039	0.072	3.283	0.412	0.344	0.000	0.000	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	55	65	49	182	0	0	78
N.S.	1	1.00	0.51	0.61	0.46	1.70	0.00	0.00	0.73
time (sec)	N/A	0.034	0.411	2.847	0.413	0.329	0.000	0.000	15.224

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	45	43	42	42	0	0	60
N.S.	1	1.00	0.59	0.57	0.55	0.55	0.00	0.00	0.79
time (sec)	N/A	0.019	0.077	3.059	0.369	0.358	0.000	0.000	0.689

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	45	45	25	157	0	0	65
N.S.	1	1.00	0.65	0.65	0.36	2.28	0.00	0.00	0.94
time (sec)	N/A	0.015	0.185	2.905	0.397	0.300	0.000	0.000	14.867

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	13	31	0	0	47
N.S.	1	1.00	0.91	0.91	0.37	0.89	0.00	0.00	1.34
time (sec)	N/A	0.009	0.150	2.868	0.374	0.279	0.000	0.000	0.353

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	26	97	22	0	37
N.S.	1	1.00	1.00	0.89	0.96	3.59	0.81	0.00	1.37
time (sec)	N/A	0.003	0.036	2.253	0.325	0.411	10.108	0.000	0.296

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	43	65	116	0	0	0
N.S.	1	1.00	0.92	1.19	1.81	3.22	0.00	0.00	0.00
time (sec)	N/A	0.008	0.018	2.938	0.366	0.312	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	67	31	0	0	62
N.S.	1	1.00	0.91	0.91	1.91	0.89	0.00	0.00	1.77
time (sec)	N/A	0.014	0.019	2.985	0.394	0.298	0.000	0.000	14.559

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	52	87	670	207	0	0	0
N.S.	1	1.00	0.67	1.12	8.59	2.65	0.00	0.00	0.00
time (sec)	N/A	0.025	0.040	2.896	0.395	0.346	0.000	0.000	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	45	45	311	44	0	0	131
N.S.	1	1.00	0.59	0.59	4.09	0.58	0.00	0.00	1.72
time (sec)	N/A	0.022	0.049	3.241	0.413	0.266	0.000	0.000	16.408

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	66	106	1679	233	0	0	0
N.S.	1	1.00	0.57	0.91	14.47	2.01	0.00	0.00	0.00
time (sec)	N/A	0.039	0.055	2.937	0.418	0.332	0.000	0.000	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	58	65	49	182	0	0	78
N.S.	1	1.00	0.54	0.61	0.46	1.70	0.00	0.00	0.73
time (sec)	N/A	0.032	0.654	3.222	0.402	0.357	0.000	0.000	15.217

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	76	76	48	43	42	42	0	0	60
N.S.	1	1.00	0.63	0.57	0.55	0.55	0.00	0.00	0.79
time (sec)	N/A	0.021	0.082	2.762	0.378	0.313	0.000	0.000	0.678

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	48	45	25	157	0	0	65
N.S.	1	1.00	0.70	0.65	0.36	2.28	0.00	0.00	0.94
time (sec)	N/A	0.018	0.228	2.922	0.386	0.320	0.000	0.000	14.050

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	32	13	31	0	0	47
N.S.	1	1.00	1.00	0.91	0.37	0.89	0.00	0.00	1.34
time (sec)	N/A	0.008	0.161	2.867	0.388	0.273	0.000	0.000	13.901

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	27	24	26	97	0	0	37
N.S.	1	1.00	1.00	0.89	0.96	3.59	0.00	0.00	1.37
time (sec)	N/A	0.003	0.031	2.253	0.368	0.307	0.000	0.000	0.297

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	33	43	65	116	0	0	0
N.S.	1	1.00	0.92	1.19	1.81	3.22	0.00	0.00	0.00
time (sec)	N/A	0.009	0.029	2.756	0.384	0.333	0.000	0.000	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	32	32	67	31	0	0	87
N.S.	1	1.00	0.91	0.91	1.91	0.89	0.00	0.00	2.49
time (sec)	N/A	0.014	0.025	3.033	0.383	0.328	0.000	0.000	14.085

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.030	0.100	0.000	0.000	0.000	0.000	0.000	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	88	88	89	0	0	0	0	0	0
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.076	11.047	0.000	0.000	0.000	0.000	0.000	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	0	13	15
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.00	0.87	1.00
time (sec)	N/A	0.030	0.025	0.722	0.310	0.277	0.000	0.328	13.969

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	17	14	13	23	0	13	15
N.S.	1	1.00	1.00	0.82	0.76	1.35	0.00	0.76	0.88
time (sec)	N/A	0.031	0.030	0.602	0.217	0.280	0.000	0.337	14.252

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	53	88	0	67	0	0	0
N.S.	1	1.00	0.79	1.31	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.063	0.268	1.173	0.000	0.089	0.000	0.000	0.000

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	67	67	61	142	0	83	0	0	0
N.S.	1	1.00	0.91	2.12	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.058	0.265	1.211	0.000	0.097	0.000	0.000	0.000

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	22	0	14	16
N.S.	1	1.00	0.86	0.67	0.62	1.05	0.00	0.67	0.76
time (sec)	N/A	0.033	0.029	8.130	0.205	0.292	0.000	0.311	13.722

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	27	26	25	23	0	24	0
N.S.	1	1.00	0.82	0.79	0.76	0.70	0.00	0.73	0.00
time (sec)	N/A	0.048	0.068	0.737	0.269	0.316	0.000	0.305	0.000

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	29	26	25	33	0	26	0
N.S.	1	1.00	0.83	0.74	0.71	0.94	0.00	0.74	0.00
time (sec)	N/A	0.045	0.072	0.913	0.194	0.286	0.000	0.319	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	63	100	0	78	0	0	0
N.S.	1	1.00	0.68	1.09	0.00	0.85	0.00	0.00	0.00
time (sec)	N/A	0.134	0.306	1.222	0.000	0.093	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	63	152	0	95	0	0	0
N.S.	1	1.00	0.68	1.65	0.00	1.03	0.00	0.00	0.00
time (sec)	N/A	0.148	0.550	1.132	0.000	0.103	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	7	6	6	0	6	8
N.S.	1	1.00	1.00	0.70	0.60	0.60	0.00	0.60	0.80
time (sec)	N/A	0.019	0.011	0.715	0.209	0.244	0.000	0.301	13.425

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	47	24	41	95	0	42	0
N.S.	1	1.00	1.47	0.75	1.28	2.97	0.00	1.31	0.00
time (sec)	N/A	0.031	0.030	0.928	0.451	0.334	0.000	0.312	0.000

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	50	43	41	97	0	42	0
N.S.	1	1.00	1.61	1.39	1.32	3.13	0.00	1.35	0.00
time (sec)	N/A	0.031	0.037	1.015	0.410	0.320	0.000	0.302	0.000

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	49	123	0	82	0	0	0
N.S.	1	1.00	0.80	2.02	0.00	1.34	0.00	0.00	0.00
time (sec)	N/A	0.052	0.322	1.273	0.000	0.131	0.000	0.000	0.000

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	54	177	0	96	0	0	0
N.S.	1	1.00	0.87	2.85	0.00	1.55	0.00	0.00	0.00
time (sec)	N/A	0.055	0.282	1.365	0.000	0.100	0.000	0.000	0.000

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	74	73	65	131	0	66	0
N.S.	1	1.00	1.19	1.18	1.05	2.11	0.00	1.06	0.00
time (sec)	N/A	0.067	0.091	1.138	0.353	0.320	0.000	0.315	0.000

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	62	62	33	71	63	141	0	64	0
N.S.	1	1.00	0.53	1.15	1.02	2.27	0.00	1.03	0.00
time (sec)	N/A	0.056	0.035	1.056	0.335	0.311	0.000	0.312	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	64	168	0	98	0	0	0
N.S.	1	1.00	0.70	1.83	0.00	1.07	0.00	0.00	0.00
time (sec)	N/A	0.101	0.530	1.323	0.000	0.096	0.000	0.000	0.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	92	92	76	160	0	114	0	0	0
N.S.	1	1.00	0.83	1.74	0.00	1.24	0.00	0.00	0.00
time (sec)	N/A	0.095	0.354	1.767	0.000	0.096	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [275] had the largest ratio of [.315800000000000025]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	6	0.167
2	A	2	2	1.00	8	0.250
3	A	2	1	1.00	8	0.125
4	A	3	2	1.00	8	0.250
5	A	2	1	1.00	8	0.125
6	A	4	2	1.00	8	0.250
7	A	2	1	1.00	8	0.125
8	A	5	2	1.00	8	0.250
9	A	3	2	1.00	10	0.200
10	A	2	2	1.00	10	0.200
11	A	2	2	1.00	10	0.200
12	A	1	1	1.00	10	0.100
13	A	1	1	1.00	10	0.100
14	A	2	2	1.00	10	0.200
15	A	2	2	1.00	10	0.200
16	A	3	2	1.00	10	0.200
17	A	4	3	1.00	12	0.250
18	A	3	3	1.00	12	0.250
19	A	3	3	1.00	12	0.250
20	A	2	2	1.00	12	0.167
21	A	2	2	1.00	12	0.167
22	A	3	3	1.00	12	0.250
23	A	3	3	1.00	12	0.250
24	A	4	3	1.00	12	0.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	1	1	1.00	10	0.100
26	A	1	1	1.00	10	0.100
27	A	1	1	1.00	10	0.100
28	A	1	1	1.00	10	0.100
29	A	1	1	1.00	10	0.100
30	A	1	1	1.00	10	0.100
31	A	1	1	1.00	12	0.083
32	A	1	1	1.00	12	0.083
33	A	1	1	1.00	12	0.083
34	A	1	1	1.00	12	0.083
35	A	1	1	1.00	12	0.083
36	A	1	1	1.00	12	0.083
37	A	1	1	1.00	8	0.125
38	A	1	1	1.00	10	0.100
39	A	4	3	1.00	10	0.300
40	A	3	3	1.00	10	0.300
41	A	2	2	1.00	10	0.200
42	A	2	2	1.00	10	0.200
43	A	3	3	1.00	10	0.300
44	A	4	3	1.00	10	0.300
45	A	6	3	1.00	10	0.300
46	A	4	3	1.00	10	0.300
47	A	3	3	1.00	10	0.300
48	A	3	3	1.00	10	0.300
49	A	4	3	1.00	10	0.300
50	A	6	3	1.00	10	0.300
51	A	7	3	1.00	10	0.300
52	A	5	3	1.00	10	0.300
53	A	3	3	1.00	10	0.300
54	A	3	3	1.00	10	0.300
55	A	3	2	1.00	10	0.200
56	A	3	2	1.00	10	0.200
57	A	2	2	1.00	12	0.167
58	A	2	2	1.00	14	0.143
59	A	2	2	1.00	14	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	2	2	1.00	14	0.143
61	A	2	2	1.00	14	0.143
62	A	2	2	1.00	14	0.143
63	A	2	2	1.00	14	0.143
64	A	2	2	1.00	14	0.143
65	A	2	2	1.00	14	0.143
66	A	6	4	1.00	21	0.190
67	A	5	4	1.00	21	0.190
68	A	5	4	1.00	21	0.190
69	A	4	4	1.00	21	0.190
70	A	4	4	1.00	19	0.210
71	A	2	2	1.00	12	0.167
72	A	3	3	1.00	19	0.158
73	A	4	4	1.00	21	0.190
74	A	4	4	1.00	21	0.190
75	A	5	4	1.00	21	0.190
76	A	5	4	1.00	21	0.190
77	A	6	4	1.00	21	0.190
78	A	6	4	1.00	21	0.190
79	A	5	4	1.00	21	0.190
80	A	5	4	1.00	21	0.190
81	A	4	4	1.00	19	0.210
82	A	3	3	1.00	12	0.250
83	A	3	3	1.00	19	0.158
84	A	3	3	1.00	21	0.143
85	A	4	4	1.00	21	0.190
86	A	4	4	1.00	21	0.190
87	A	5	4	1.00	21	0.190
88	A	5	4	1.00	21	0.190
89	A	6	4	1.00	21	0.190
90	A	6	4	1.00	21	0.190
91	A	5	4	1.00	21	0.190
92	A	5	4	1.00	19	0.210
93	A	3	3	1.00	12	0.250
94	A	4	4	1.00	19	0.210

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
95	A	3	3	1.00	21	0.143
96	A	3	3	1.00	21	0.143
97	A	4	4	1.00	21	0.190
98	A	4	4	1.00	21	0.190
99	A	5	4	1.00	21	0.190
100	A	5	4	1.00	21	0.190
101	A	6	4	1.00	21	0.190
102	A	4	3	1.00	12	0.250
103	A	6	4	1.00	21	0.190
104	A	5	4	1.00	21	0.190
105	A	5	4	1.00	21	0.190
106	A	4	4	1.00	21	0.190
107	A	4	4	1.00	21	0.190
108	A	3	3	1.00	19	0.158
109	A	2	2	1.00	12	0.167
110	A	4	4	1.00	19	0.210
111	A	4	4	1.00	21	0.190
112	A	5	4	1.00	21	0.190
113	A	5	4	1.00	21	0.190
114	A	6	4	1.00	21	0.190
115	A	6	4	1.00	21	0.190
116	A	5	4	1.00	21	0.190
117	A	5	4	1.00	21	0.190
118	A	4	4	1.00	21	0.190
119	A	4	4	1.00	21	0.190
120	A	3	3	1.00	21	0.143
121	A	3	3	1.00	19	0.158
122	A	3	3	1.00	12	0.250
123	A	4	4	1.00	19	0.210
124	A	5	4	1.00	21	0.190
125	A	5	4	1.00	21	0.190
126	A	6	4	1.00	21	0.190
127	A	6	4	1.00	21	0.190
128	A	5	4	1.00	21	0.190
129	A	5	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
130	A	4	4	1.00	21	0.190
131	A	4	4	1.00	21	0.190
132	A	3	3	1.00	21	0.143
133	A	3	3	1.00	21	0.143
134	A	4	4	1.00	19	0.210
135	A	3	3	1.00	12	0.250
136	A	5	4	1.00	19	0.210
137	A	5	4	1.00	21	0.190
138	A	6	4	1.00	21	0.190
139	A	4	3	1.00	12	0.250
140	A	4	3	1.00	23	0.130
141	A	3	2	1.00	23	0.087
142	A	3	3	1.00	23	0.130
143	A	2	2	1.00	23	0.087
144	A	2	2	1.00	23	0.087
145	A	2	2	1.00	23	0.087
146	A	3	3	1.00	23	0.130
147	A	3	3	1.00	23	0.130
148	A	3	2	1.00	23	0.087
149	A	4	3	1.00	23	0.130
150	A	4	3	1.00	23	0.130
151	A	3	2	1.00	23	0.087
152	A	3	3	1.00	23	0.130
153	A	2	2	1.00	23	0.087
154	A	2	2	1.00	23	0.087
155	A	2	2	1.00	23	0.087
156	A	3	3	1.00	23	0.130
157	A	3	3	1.00	23	0.130
158	A	3	2	1.00	23	0.087
159	A	4	3	1.00	23	0.130
160	A	3	2	1.00	23	0.087
161	A	4	3	1.00	23	0.130
162	A	3	2	1.00	23	0.087
163	A	3	3	1.00	23	0.130
164	A	2	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
165	A	2	2	1.00	23	0.087
166	A	2	2	1.00	23	0.087
167	A	3	3	1.00	23	0.130
168	A	3	3	1.00	23	0.130
169	A	3	2	1.00	23	0.087
170	A	4	3	1.00	23	0.130
171	A	3	2	1.00	23	0.087
172	A	4	3	1.00	23	0.130
173	A	3	2	1.00	23	0.087
174	A	3	3	1.00	23	0.130
175	A	2	2	1.00	23	0.087
176	A	2	2	1.00	23	0.087
177	A	2	2	1.00	23	0.087
178	A	3	3	1.00	23	0.130
179	A	3	3	1.00	23	0.130
180	A	3	2	1.00	23	0.087
181	A	4	3	1.00	23	0.130
182	A	4	3	1.00	23	0.130
183	A	3	2	1.00	23	0.087
184	A	3	3	1.00	23	0.130
185	A	2	2	1.00	23	0.087
186	A	2	2	1.00	23	0.087
187	A	2	2	1.00	23	0.087
188	A	3	3	1.00	23	0.130
189	A	3	3	1.00	23	0.130
190	A	3	2	1.00	23	0.087
191	A	4	3	1.00	23	0.130
192	A	4	3	1.00	23	0.130
193	A	3	2	1.00	23	0.087
194	A	3	3	1.00	23	0.130
195	A	2	2	1.00	23	0.087
196	A	2	2	1.00	23	0.087
197	A	2	2	1.00	23	0.087
198	A	3	3	1.00	23	0.130
199	A	3	3	1.00	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
200	A	3	2	1.00	23	0.087
201	A	4	3	1.00	23	0.130
202	A	2	2	1.00	21	0.095
203	A	2	2	1.00	21	0.095
204	A	2	2	1.00	19	0.105
205	A	1	1	1.00	12	0.083
206	A	2	2	1.00	19	0.105
207	A	2	2	1.00	21	0.095
208	A	2	2	1.00	21	0.095
209	A	2	2	1.00	21	0.095
210	A	2	2	1.00	21	0.095
211	A	2	2	1.00	19	0.105
212	A	1	1	1.00	12	0.083
213	A	2	2	1.00	19	0.105
214	A	2	2	1.00	21	0.095
215	A	2	2	1.00	21	0.095
216	A	2	2	1.00	21	0.095
217	A	2	2	1.00	21	0.095
218	A	2	2	1.00	19	0.105
219	A	1	1	1.00	12	0.083
220	A	2	2	1.00	19	0.105
221	A	2	2	1.00	21	0.095
222	A	2	2	1.00	21	0.095
223	A	2	2	1.00	21	0.095
224	A	2	2	1.00	21	0.095
225	A	2	2	1.00	19	0.105
226	A	1	1	1.00	12	0.083
227	A	2	2	1.00	19	0.105
228	A	2	2	1.00	21	0.095
229	A	2	2	1.00	21	0.095
230	A	2	2	1.00	21	0.095
231	A	2	2	1.00	21	0.095
232	A	2	2	1.00	19	0.105
233	A	1	1	1.00	12	0.083
234	A	2	2	1.00	19	0.105

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
235	A	2	2	1.00	21	0.095
236	A	2	2	1.00	21	0.095
237	A	2	2	1.00	21	0.095
238	A	2	2	1.00	21	0.095
239	A	2	2	1.00	19	0.105
240	A	1	1	1.00	12	0.083
241	A	2	2	1.00	19	0.105
242	A	2	2	1.00	21	0.095
243	A	2	2	1.00	21	0.095
244	A	2	2	1.00	21	0.095
245	A	2	2	1.00	19	0.105
246	A	2	2	1.00	17	0.118
247	A	1	1	1.00	10	0.100
248	A	2	2	1.00	17	0.118
249	A	2	2	1.00	19	0.105
250	A	2	2	1.00	19	0.105
251	A	2	2	1.00	19	0.105
252	A	2	2	1.00	21	0.095
253	A	2	2	1.00	21	0.095
254	A	2	2	1.00	21	0.095
255	A	2	2	1.00	21	0.095
256	A	2	2	1.00	21	0.095
257	A	2	2	1.00	21	0.095
258	A	2	2	1.00	21	0.095
259	A	2	2	1.00	21	0.095
260	A	2	2	1.00	21	0.095
261	A	2	2	1.00	17	0.118
262	A	2	2	1.00	17	0.118
263	A	3	3	1.00	19	0.158
264	A	3	3	1.00	19	0.158
265	A	3	2	1.00	11	0.182
266	A	3	2	1.00	19	0.105
267	A	3	2	1.00	19	0.105
268	A	4	3	1.00	19	0.158
269	A	4	3	1.00	19	0.158

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
270	A	2	2	1.00	9	0.222
271	A	5	5	1.00	17	0.294
272	A	5	5	1.00	17	0.294
273	A	3	3	1.00	19	0.158
274	A	3	3	1.00	19	0.158
275	A	6	6	1.00	19	0.316
276	A	6	6	1.00	19	0.316
277	A	4	3	1.00	19	0.158
278	A	4	3	1.00	19	0.158
279	A	2	2	1.00	21	0.095
280	A	2	2	1.00	21	0.095
281	A	2	2	1.00	21	0.095
282	A	2	2	1.00	21	0.095
283	A	2	2	1.00	21	0.095
284	A	2	2	1.00	17	0.118
285	A	2	2	1.00	19	0.105
286	A	2	2	1.00	19	0.105
287	A	2	2	1.00	21	0.095
288	A	2	2	0.97	23	0.087
289	A	2	2	1.00	23	0.087
290	A	2	2	1.00	23	0.087
291	A	2	2	1.00	23	0.087
292	A	2	2	1.00	23	0.087
293	A	2	2	1.00	23	0.087
294	A	2	2	1.00	23	0.087

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \cos(a + bx) dx$	106
3.2	$\int \cos^2(a + bx) dx$	109
3.3	$\int \cos^3(a + bx) dx$	113
3.4	$\int \cos^4(a + bx) dx$	117
3.5	$\int \cos^5(a + bx) dx$	121
3.6	$\int \cos^6(a + bx) dx$	125
3.7	$\int \cos^7(a + bx) dx$	129
3.8	$\int \cos^8(a + bx) dx$	133
3.9	$\int \cos^{\frac{7}{2}}(a + bx) dx$	138
3.10	$\int \cos^{\frac{5}{2}}(a + bx) dx$	142
3.11	$\int \cos^{\frac{3}{2}}(a + bx) dx$	146
3.12	$\int \sqrt{\cos(a + bx)} dx$	150
3.13	$\int \frac{1}{\sqrt{\cos(a + bx)}} dx$	153
3.14	$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx$	156
3.15	$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx$	160
3.16	$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx$	164
3.17	$\int (c \cos(a + bx))^{\frac{7}{2}} dx$	168
3.18	$\int (c \cos(a + bx))^{\frac{5}{2}} dx$	172
3.19	$\int (c \cos(a + bx))^{\frac{3}{2}} dx$	176
3.20	$\int \sqrt{c \cos(a + bx)} dx$	180
3.21	$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx$	184
3.22	$\int \frac{1}{(c \cos(a + bx))^{\frac{3}{2}}} dx$	188
3.23	$\int \frac{1}{(c \cos(a + bx))^{\frac{5}{2}}} dx$	192
3.24	$\int \frac{1}{(c \cos(a + bx))^{\frac{7}{2}}} dx$	196
3.25	$\int \cos^{\frac{4}{3}}(a + bx) dx$	200

3.26	$\int \cos^{\frac{2}{3}}(a + bx) dx$	203
3.27	$\int \sqrt[3]{\cos(a + bx)} dx$	206
3.28	$\int \frac{1}{\sqrt[3]{\cos(a + bx)}} dx$	209
3.29	$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx$	212
3.30	$\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx$	215
3.31	$\int (c \cos(a + bx))^{4/3} dx$	218
3.32	$\int (c \cos(a + bx))^{2/3} dx$	221
3.33	$\int \sqrt[3]{c \cos(a + bx)} dx$	224
3.34	$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$	227
3.35	$\int \frac{1}{(c \cos(a+bx))^{2/3}} dx$	230
3.36	$\int \frac{1}{(c \cos(a+bx))^{4/3}} dx$	233
3.37	$\int \cos^n(a + bx) dx$	236
3.38	$\int (c \cos(a + bx))^n dx$	239
3.39	$\int (a \cos^2(x))^{5/2} dx$	242
3.40	$\int (a \cos^2(x))^{3/2} dx$	246
3.41	$\int \sqrt{a \cos^2(x)} dx$	250
3.42	$\int \frac{1}{\sqrt{a \cos^2(x)}} dx$	253
3.43	$\int \frac{1}{(a \cos^2(x))^{3/2}} dx$	257
3.44	$\int \frac{1}{(a \cos^2(x))^{5/2}} dx$	261
3.45	$\int (a \cos^3(x))^{5/2} dx$	266
3.46	$\int (a \cos^3(x))^{3/2} dx$	271
3.47	$\int \sqrt{a \cos^3(x)} dx$	275
3.48	$\int \frac{1}{\sqrt{a \cos^3(x)}} dx$	279
3.49	$\int \frac{1}{(a \cos^3(x))^{3/2}} dx$	283
3.50	$\int \frac{1}{(a \cos^3(x))^{5/2}} dx$	287
3.51	$\int (a \cos^4(x))^{5/2} dx$	292
3.52	$\int (a \cos^4(x))^{3/2} dx$	297
3.53	$\int \sqrt{a \cos^4(x)} dx$	301
3.54	$\int \frac{1}{\sqrt{a \cos^4(x)}} dx$	305
3.55	$\int \frac{1}{(a \cos^4(x))^{3/2}} dx$	309
3.56	$\int \frac{1}{(a \cos^4(x))^{5/2}} dx$	313
3.57	$\int (b \cos^m(c + dx))^n dx$	318
3.58	$\int (c \cos^m(a + bx))^{5/2} dx$	322
3.59	$\int (c \cos^m(a + bx))^{3/2} dx$	326
3.60	$\int \sqrt{c \cos^m(a + bx)} dx$	330
3.61	$\int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx$	334
3.62	$\int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx$	338

3.63	$\int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx$	342
3.64	$\int (c \cos^m(a+bx))^{\frac{1}{m}} dx$	346
3.65	$\int (a(b \cos(c+dx))^p)^n dx$	350
3.66	$\int \cos^5(c+dx) \sqrt{b \cos(c+dx)} dx$	354
3.67	$\int \cos^4(c+dx) \sqrt{b \cos(c+dx)} dx$	359
3.68	$\int \cos^3(c+dx) \sqrt{b \cos(c+dx)} dx$	363
3.69	$\int \cos^2(c+dx) \sqrt{b \cos(c+dx)} dx$	368
3.70	$\int \cos(c+dx) \sqrt{b \cos(c+dx)} dx$	372
3.71	$\int \sqrt{b \cos(c+dx)} dx$	376
3.72	$\int \sqrt{b \cos(c+dx)} \sec(c+dx) dx$	380
3.73	$\int \sqrt{b \cos(c+dx)} \sec^2(c+dx) dx$	384
3.74	$\int \sqrt{b \cos(c+dx)} \sec^3(c+dx) dx$	388
3.75	$\int \sqrt{b \cos(c+dx)} \sec^4(c+dx) dx$	392
3.76	$\int \sqrt{b \cos(c+dx)} \sec^5(c+dx) dx$	397
3.77	$\int \sqrt{b \cos(c+dx)} \sec^6(c+dx) dx$	402
3.78	$\int \cos^4(c+dx) (b \cos(c+dx))^{3/2} dx$	407
3.79	$\int \cos^3(c+dx) (b \cos(c+dx))^{3/2} dx$	412
3.80	$\int \cos^2(c+dx) (b \cos(c+dx))^{3/2} dx$	416
3.81	$\int \cos(c+dx) (b \cos(c+dx))^{3/2} dx$	420
3.82	$\int (b \cos(c+dx))^{3/2} dx$	424
3.83	$\int (b \cos(c+dx))^{3/2} \sec(c+dx) dx$	428
3.84	$\int (b \cos(c+dx))^{3/2} \sec^2(c+dx) dx$	432
3.85	$\int (b \cos(c+dx))^{3/2} \sec^3(c+dx) dx$	436
3.86	$\int (b \cos(c+dx))^{3/2} \sec^4(c+dx) dx$	440
3.87	$\int (b \cos(c+dx))^{3/2} \sec^5(c+dx) dx$	444
3.88	$\int (b \cos(c+dx))^{3/2} \sec^6(c+dx) dx$	449
3.89	$\int (b \cos(c+dx))^{3/2} \sec^7(c+dx) dx$	454
3.90	$\int \cos^3(c+dx) (b \cos(c+dx))^{5/2} dx$	459
3.91	$\int \cos^2(c+dx) (b \cos(c+dx))^{5/2} dx$	464
3.92	$\int \cos(c+dx) (b \cos(c+dx))^{5/2} dx$	468
3.93	$\int (b \cos(c+dx))^{5/2} dx$	472
3.94	$\int (b \cos(c+dx))^{5/2} \sec(c+dx) dx$	476
3.95	$\int (b \cos(c+dx))^{5/2} \sec^2(c+dx) dx$	480
3.96	$\int (b \cos(c+dx))^{5/2} \sec^3(c+dx) dx$	484
3.97	$\int (b \cos(c+dx))^{5/2} \sec^4(c+dx) dx$	488
3.98	$\int (b \cos(c+dx))^{5/2} \sec^5(c+dx) dx$	492
3.99	$\int (b \cos(c+dx))^{5/2} \sec^6(c+dx) dx$	496
3.100	$\int (b \cos(c+dx))^{5/2} \sec^7(c+dx) dx$	501
3.101	$\int (b \cos(c+dx))^{5/2} \sec^8(c+dx) dx$	506
3.102	$\int (b \cos(c+dx))^{7/2} dx$	510
3.103	$\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	514
3.104	$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	519

3.105	$\int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	524
3.106	$\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	529
3.107	$\int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	533
3.108	$\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	537
3.109	$\int \frac{1}{\sqrt{b \cos(c+dx)}} dx$	541
3.110	$\int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	545
3.111	$\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	549
3.112	$\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	553
3.113	$\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	558
3.114	$\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	563
3.115	$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	568
3.116	$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	573
3.117	$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	577
3.118	$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	581
3.119	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	585
3.120	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	589
3.121	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	593
3.122	$\int \frac{1}{(b \cos(c+dx))^{3/2}} dx$	597
3.123	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	601
3.124	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	605
3.125	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	610
3.126	$\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	615
3.127	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	620
3.128	$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	625
3.129	$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	629
3.130	$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	633
3.131	$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	637
3.132	$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	641
3.133	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	645
3.134	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	649
3.135	$\int \frac{1}{(b \cos(c+dx))^{5/2}} dx$	653
3.136	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	657

3.137	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	662
3.138	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	667
3.139	$\int \frac{1}{(b \cos(c+dx))^{7/2}} dx$	672
3.140	$\int \cos^{7/2}(c+dx) \sqrt{b \cos(c+dx)} dx$	676
3.141	$\int \cos^{5/2}(c+dx) \sqrt{b \cos(c+dx)} dx$	681
3.142	$\int \cos^{3/2}(c+dx) \sqrt{b \cos(c+dx)} dx$	732
3.143	$\int \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} dx$	736
3.144	$\int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx$	740
3.145	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{3/2}(c+dx)} dx$	744
3.146	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{5/2}(c+dx)} dx$	748
3.147	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{7/2}(c+dx)} dx$	752
3.148	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{9/2}(c+dx)} dx$	757
3.149	$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{11/2}(c+dx)} dx$	761
3.150	$\int \cos^{5/2}(c+dx) (b \cos(c+dx))^{3/2} dx$	767
3.151	$\int \cos^{3/2}(c+dx) (b \cos(c+dx))^{3/2} dx$	772
3.152	$\int \sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2} dx$	823
3.153	$\int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx$	827
3.154	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{3/2}(c+dx)} dx$	831
3.155	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{5/2}(c+dx)} dx$	835
3.156	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{7/2}(c+dx)} dx$	839
3.157	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{9/2}(c+dx)} dx$	843
3.158	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{11/2}(c+dx)} dx$	848
3.159	$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{13/2}(c+dx)} dx$	852
3.160	$\int \cos^{5/2}(c+dx) (b \cos(c+dx))^{5/2} dx$	858
3.161	$\int \cos^{3/2}(c+dx) (b \cos(c+dx))^{5/2} dx$	862
3.162	$\int \sqrt{\cos(c+dx)} (b \cos(c+dx))^{5/2} dx$	867
3.163	$\int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx$	918
3.164	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{3/2}(c+dx)} dx$	922
3.165	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{5/2}(c+dx)} dx$	925
3.166	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{7/2}(c+dx)} dx$	929
3.167	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{9/2}(c+dx)} dx$	933

3.168	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{11/2}(c+dx)} dx$	937
3.169	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx$	942
3.170	$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{15/2}(c+dx)} dx$	946
3.171	$\int \frac{\cos^{11/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	952
3.172	$\int \frac{\cos^{9/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	956
3.173	$\int \frac{\cos^{7/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	960
3.174	$\int \frac{\cos^{5/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	964
3.175	$\int \frac{\cos^{3/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$	968
3.176	$\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx$	972
3.177	$\int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} dx$	976
3.178	$\int \frac{1}{\cos^{3/2}(c+dx) \sqrt{b \cos(c+dx)}} dx$	980
3.179	$\int \frac{1}{\cos^{5/2}(c+dx) \sqrt{b \cos(c+dx)}} dx$	984
3.180	$\int \frac{1}{\cos^{7/2}(c+dx) \sqrt{b \cos(c+dx)}} dx$	989
3.181	$\int \frac{1}{\cos^{9/2}(c+dx) \sqrt{b \cos(c+dx)}} dx$	993
3.182	$\int \frac{\cos^{11/2}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	999
3.183	$\int \frac{\cos^{9/2}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	1003
3.184	$\int \frac{\cos^{7/2}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	1007
3.185	$\int \frac{\cos^{5/2}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	1011
3.186	$\int \frac{\cos^{3/2}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$	1014
3.187	$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx$	1018
3.188	$\int \frac{1}{\sqrt{\cos(c+dx)} (b \cos(c+dx))^{3/2}} dx$	1022
3.189	$\int \frac{1}{\cos^{3/2}(c+dx) (b \cos(c+dx))^{3/2}} dx$	1026
3.190	$\int \frac{1}{\cos^{5/2}(c+dx) (b \cos(c+dx))^{3/2}} dx$	1031
3.191	$\int \frac{1}{\cos^{7/2}(c+dx) (b \cos(c+dx))^{3/2}} dx$	1035
3.192	$\int \frac{\cos^{13/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1041
3.193	$\int \frac{\cos^{11/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1045
3.194	$\int \frac{\cos^{9/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1049
3.195	$\int \frac{\cos^{7/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1053
3.196	$\int \frac{\cos^{5/2}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1056

3.197	$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$	1060
3.198	$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx$	1064
3.199	$\int \frac{1}{\sqrt{\cos(c+dx)(b \cos(c+dx))^{5/2}}} dx$	1068
3.200	$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$	1073
3.201	$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$	1077
3.202	$\int \cos^m(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	1083
3.203	$\int \cos^2(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	1087
3.204	$\int \cos(c+dx) \sqrt[3]{b \cos(c+dx)} dx$	1091
3.205	$\int \sqrt[3]{b \cos(c+dx)} dx$	1095
3.206	$\int \sqrt[3]{b \cos(c+dx)} \sec(c+dx) dx$	1098
3.207	$\int \sqrt[3]{b \cos(c+dx)} \sec^2(c+dx) dx$	1102
3.208	$\int \sqrt[3]{b \cos(c+dx)} \sec^3(c+dx) dx$	1106
3.209	$\int \cos^m(c+dx) (b \cos(c+dx))^{2/3} dx$	1110
3.210	$\int \cos^2(c+dx) (b \cos(c+dx))^{2/3} dx$	1114
3.211	$\int \cos(c+dx) (b \cos(c+dx))^{2/3} dx$	1118
3.212	$\int (b \cos(c+dx))^{2/3} dx$	1122
3.213	$\int (b \cos(c+dx))^{2/3} \sec(c+dx) dx$	1125
3.214	$\int (b \cos(c+dx))^{2/3} \sec^2(c+dx) dx$	1129
3.215	$\int (b \cos(c+dx))^{2/3} \sec^3(c+dx) dx$	1133
3.216	$\int \cos^m(c+dx) (b \cos(c+dx))^{4/3} dx$	1137
3.217	$\int \cos^2(c+dx) (b \cos(c+dx))^{4/3} dx$	1141
3.218	$\int \cos(c+dx) (b \cos(c+dx))^{4/3} dx$	1145
3.219	$\int (b \cos(c+dx))^{4/3} dx$	1149
3.220	$\int (b \cos(c+dx))^{4/3} \sec(c+dx) dx$	1152
3.221	$\int (b \cos(c+dx))^{4/3} \sec^2(c+dx) dx$	1156
3.222	$\int (b \cos(c+dx))^{4/3} \sec^3(c+dx) dx$	1160
3.223	$\int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1164
3.224	$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1168
3.225	$\int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1172
3.226	$\int \frac{1}{\sqrt[3]{b \cos(c+dx)}} dx$	1176
3.227	$\int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1179
3.228	$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1183
3.229	$\int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$	1187
3.230	$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1191
3.231	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1195
3.232	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1199

3.233	$\int \frac{1}{(b \cos(c+dx))^{2/3}} dx$	1203
3.234	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1206
3.235	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1209
3.236	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx$	1212
3.237	$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1215
3.238	$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1219
3.239	$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1223
3.240	$\int \frac{1}{(b \cos(c+dx))^{4/3}} dx$	1227
3.241	$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1230
3.242	$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1233
3.243	$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx$	1236
3.244	$\int (a \cos(e+fx))^m (b \cos(e+fx))^n dx$	1239
3.245	$\int \cos^2(c+dx) (b \cos(c+dx))^n dx$	1243
3.246	$\int \cos(c+dx) (b \cos(c+dx))^n dx$	1247
3.247	$\int (b \cos(c+dx))^n dx$	1251
3.248	$\int (b \cos(c+dx))^n \sec(c+dx) dx$	1254
3.249	$\int (b \cos(c+dx))^n \sec^2(c+dx) dx$	1258
3.250	$\int (b \cos(c+dx))^n \sec^3(c+dx) dx$	1262
3.251	$\int (b \cos(c+dx))^n \sec^4(c+dx) dx$	1266
3.252	$\int \cos^{5/2}(c+dx) (b \cos(c+dx))^n dx$	1270
3.253	$\int \cos^{3/2}(c+dx) (b \cos(c+dx))^n dx$	1274
3.254	$\int \sqrt{\cos(c+dx)} (b \cos(c+dx))^n dx$	1278
3.255	$\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx$	1282
3.256	$\int \frac{(b \cos(c+dx))^n}{\cos^{3/2}(c+dx)} dx$	1286
3.257	$\int \frac{(b \cos(c+dx))^n}{\cos^{5/2}(c+dx)} dx$	1290
3.258	$\int \frac{(b \cos(c+dx))^n}{\cos^{7/2}(c+dx)} dx$	1294
3.259	$\int \frac{(b \cos(c+dx))^n}{\cos^{9/2}(c+dx)} dx$	1298
3.260	$\int (a \cos(e+fx))^m (b \sec(e+fx))^n dx$	1302
3.261	$\int \cos(a+bx) \sqrt{\csc(a+bx)} dx$	1306
3.262	$\int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1309
3.263	$\int \cos^2(a+bx) \sqrt{\csc(a+bx)} dx$	1313
3.264	$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1317
3.265	$\int \cos^3(x) \csc^{9/2}(x) dx$	1321
3.266	$\int \cos^3(a+bx) \sqrt{\csc(a+bx)} dx$	1325
3.267	$\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1329
3.268	$\int \cos^4(a+bx) \sqrt{\csc(a+bx)} dx$	1333

3.269	$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1337
3.270	$\int \cos(x) \csc^{\frac{7}{3}}(x) dx$	1341
3.271	$\int \sqrt{\csc(a+bx)} \sec(a+bx) dx$	1344
3.272	$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1349
3.273	$\int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx$	1354
3.274	$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1358
3.275	$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$	1362
3.276	$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1367
3.277	$\int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx$	1372
3.278	$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$	1376
3.279	$\int (d \cos(a+bx))^{\frac{3}{2}} \csc^p(a+bx) dx$	1380
3.280	$\int \sqrt{d \cos(a+bx)} \csc^p(a+bx) dx$	1384
3.281	$\int \frac{\csc^p(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	1388
3.282	$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{\frac{3}{2}}} dx$	1392
3.283	$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{\frac{5}{2}}} dx$	1395
3.284	$\int \cos^m(e+fx) \csc^n(e+fx) dx$	1398
3.285	$\int (a \cos(e+fx))^m \csc^n(e+fx) dx$	1402
3.286	$\int \cos^m(e+fx) (b \csc(e+fx))^n dx$	1406
3.287	$\int (a \cos(e+fx))^m (b \csc(e+fx))^n dx$	1410
3.288	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{\frac{7}{2}} dx$	1414
3.289	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{\frac{5}{2}} dx$	1418
3.290	$\int (a \cos(e+fx))^m (b \csc(e+fx))^{\frac{3}{2}} dx$	1422
3.291	$\int (a \cos(e+fx))^m \sqrt{b \csc(e+fx)} dx$	1426
3.292	$\int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx$	1430
3.293	$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{\frac{3}{2}}} dx$	1434
3.294	$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{\frac{5}{2}}} dx$	1438

3.1 $\int \cos(a + bx) dx$

Optimal result	106
Rubi [A] (verified)	106
Mathematica [B] (verified)	107
Maple [A] (verified)	107
Fricas [A] (verification not implemented)	107
Sympy [A] (verification not implemented)	108
Maxima [A] (verification not implemented)	108
Giac [A] (verification not implemented)	108
Mupad [B] (verification not implemented)	108

Optimal result

Integrand size = 6, antiderivative size = 10

$$\int \cos(a + bx) dx = \frac{\sin(a + bx)}{b}$$

[Out] $\sin(b*x+a)/b$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2717}

$$\int \cos(a + bx) dx = \frac{\sin(a + bx)}{b}$$

[In] $\text{Int}[\text{Cos}[a + b*x], x]$

[Out] $\text{Sin}[a + b*x]/b$

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rubi steps

$$\text{integral} = \frac{\sin(a + bx)}{b}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 21 vs. 2(10) = 20.

Time = 0.01 (sec) , antiderivative size = 21, normalized size of antiderivative = 2.10

$$\int \cos(a + bx) dx = \frac{\cos(bx) \sin(a)}{b} + \frac{\cos(a) \sin(bx)}{b}$$

[In] Integrate[Cos[a + b*x],x]

[Out] (Cos[b*x]*Sin[a])/b + (Cos[a]*Sin[b*x])/b

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
derivativedivides	$\frac{\sin(bx+a)}{b}$	11
default	$\frac{\sin(bx+a)}{b}$	11
risch	$\frac{\sin(bx+a)}{b}$	11
parallelrisch	$\frac{\sin(bx+a)}{b}$	11
norman	$\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}$	30
meijerg	$\frac{\cos(a) \sin(bx)}{b} - \frac{\sin(a) \sqrt{\pi} \left(\frac{1}{\sqrt{\pi}} - \frac{\cos(bx)}{\sqrt{\pi}}\right)}{b}$	35

[In] int(cos(b*x+a),x,method=_RETURNVERBOSE)

[Out] sin(b*x+a)/b

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(a + bx) dx = \frac{\sin(bx + a)}{b}$$

[In] integrate(cos(b*x+a),x, algorithm="fricas")

[Out] sin(b*x + a)/b

Sympy [A] (verification not implemented)

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \cos(a + bx) dx = \begin{cases} \frac{\sin(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a),x)

[Out] Piecewise((sin(a + b*x)/b, Ne(b, 0)), (x*cos(a), True))

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(a + bx) dx = \frac{\sin(bx + a)}{b}$$

[In] integrate(cos(b*x+a),x, algorithm="maxima")

[Out] sin(b*x + a)/b

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(a + bx) dx = \frac{\sin(bx + a)}{b}$$

[In] integrate(cos(b*x+a),x, algorithm="giac")

[Out] sin(b*x + a)/b

Mupad [B] (verification not implemented)

Time = 14.68 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(a + bx) dx = \frac{\sin(a + bx)}{b}$$

[In] int(cos(a + b*x),x)

[Out] sin(a + b*x)/b

3.2 $\int \cos^2(a + bx) dx$

Optimal result	109
Rubi [A] (verified)	109
Mathematica [A] (verified)	110
Maple [A] (verified)	110
Fricas [A] (verification not implemented)	111
Sympy [B] (verification not implemented)	111
Maxima [A] (verification not implemented)	111
Giac [A] (verification not implemented)	112
Mupad [B] (verification not implemented)	112

Optimal result

Integrand size = 8, antiderivative size = 25

$$\int \cos^2(a + bx) dx = \frac{x}{2} + \frac{\cos(a + bx) \sin(a + bx)}{2b}$$

[Out] 1/2*x+1/2*cos(b*x+a)*sin(b*x+a)/b

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\int \cos^2(a + bx) dx = \frac{\sin(a + bx) \cos(a + bx)}{2b} + \frac{x}{2}$$

[In] Int[Cos[a + b*x]^2,x]

[Out] x/2 + (Cos[a + b*x]*Sin[a + b*x])/(2*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos(a+bx)\sin(a+bx)}{2b} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} + \frac{\cos(a+bx)\sin(a+bx)}{2b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.92

$$\int \cos^2(a+bx) dx = \frac{2(a+bx) + \sin(2(a+bx))}{4b}$$

[In] Integrate[Cos[a + b*x]^2,x]

[Out] (2*(a + b*x) + Sin[2*(a + b*x)])/(4*b)

Maple [A] (verified)

Time = 0.48 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.76

method	result	size
risch	$\frac{x}{2} + \frac{\sin(2bx+2a)}{4b}$	19
parallelrisc	$\frac{2bx+\sin(2bx+2a)}{4b}$	20
derivativdivides	$\frac{\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx}{2} + \frac{a}{2}}{b}$	27
default	$\frac{\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx}{2} + \frac{a}{2}}{b}$	27
norman	$\frac{\frac{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} + x\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \frac{x}{2} - \frac{\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} + \frac{x\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^2}$	77

[In] int(cos(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/2*x+1/4/b*sin(2*b*x+2*a)

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \cos^2(a + bx) dx = \frac{bx + \cos(bx + a) \sin(bx + a)}{2b}$$

[In] integrate(cos(b*x+a)^2,x, algorithm="fricas")

[Out] 1/2*(b*x + cos(b*x + a)*sin(b*x + a))/b

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 46 vs. 2(19) = 38.

Time = 0.09 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.84

$$\int \cos^2(a + bx) dx = \begin{cases} \frac{x \sin^2(a+bx)}{2} + \frac{x \cos^2(a+bx)}{2} + \frac{\sin(a+bx) \cos(a+bx)}{2b} & \text{for } b \neq 0 \\ x \cos^2(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a)**2,x)

[Out] Piecewise((x*sin(a + b*x)**2/2 + x*cos(a + b*x)**2/2 + sin(a + b*x)*cos(a + b*x)/(2*b), Ne(b, 0)), (x*cos(a)**2, True))

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \cos^2(a + bx) dx = \frac{2bx + 2a + \sin(2bx + 2a)}{4b}$$

[In] integrate(cos(b*x+a)^2,x, algorithm="maxima")

[Out] 1/4*(2*b*x + 2*a + sin(2*b*x + 2*a))/b

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \cos^2(a + bx) dx = \frac{1}{2}x + \frac{\sin(2bx + 2a)}{4b}$$

[In] integrate(cos(b*x+a)^2,x, algorithm="giac")

[Out] 1/2*x + 1/4*sin(2*b*x + 2*a)/b

Mupad [B] (verification not implemented)

Time = 15.28 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.72

$$\int \cos^2(a + bx) dx = \frac{x}{2} + \frac{\sin(2a + 2bx)}{4b}$$

[In] int(cos(a + b*x)^2,x)

[Out] x/2 + sin(2*a + 2*b*x)/(4*b)

3.3 $\int \cos^3(a + bx) dx$

Optimal result	113
Rubi [A] (verified)	113
Mathematica [A] (verified)	114
Maple [A] (verified)	114
Fricas [A] (verification not implemented)	114
Sympy [A] (verification not implemented)	115
Maxima [A] (verification not implemented)	115
Giac [A] (verification not implemented)	115
Mupad [B] (verification not implemented)	116

Optimal result

Integrand size = 8, antiderivative size = 26

$$\int \cos^3(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[Out] $\sin(b*x+a)/b-1/3*\sin(b*x+a)^3/b$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2713}

$$\int \cos^3(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^3, x]$

[Out] $\text{Sin}[a + b*x]/b - \text{Sin}[a + b*x]^3/(3*b)$

Rule 2713

$\text{Int}[\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n - 1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ $\text{FreeQ}\{c, d, x\}$ && $\text{IGtQ}[(n - 1)/2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int (1 - x^2) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \cos^3(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[In] Integrate[Cos[a + b*x]^3,x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/(3*b)

Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

method	result	size
derivativdivides	$\frac{(2+\cos^2(bx+a)) \sin(bx+a)}{3b}$	22
default	$\frac{(2+\cos^2(bx+a)) \sin(bx+a)}{3b}$	22
parallelrisc	$\frac{9 \sin(bx+a) + \sin(3bx+3a)}{12b}$	24
risc	$\frac{3 \sin(bx+a)}{4b} + \frac{\sin(3bx+3a)}{12b}$	27

[In] int(cos(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/3/b*(2+cos(b*x+a)^2)*sin(b*x+a)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

$$\int \cos^3(a + bx) dx = \frac{(\cos(bx + a))^2 + 2}{3b} \sin(bx + a)$$

[In] integrate(cos(b*x+a)^3,x, algorithm="fricas")

[Out] 1/3*(cos(b*x + a)^2 + 2)*sin(b*x + a)/b

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.38

$$\int \cos^3(a + bx) dx = \begin{cases} \frac{2 \sin^3(a+bx)}{3b} + \frac{\sin(a+bx) \cos^2(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^3(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a)**3,x)

[Out] Piecewise((2*sin(a + b*x)**3/(3*b) + sin(a + b*x)*cos(a + b*x)**2/b, Ne(b, 0)), (x*cos(a)**3, True))

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \cos^3(a + bx) dx = -\frac{\sin(bx + a)^3 - 3 \sin(bx + a)}{3b}$$

[In] integrate(cos(b*x+a)^3,x, algorithm="maxima")

[Out] -1/3*(sin(b*x + a)^3 - 3*sin(b*x + a))/b

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \cos^3(a + bx) dx = -\frac{\sin(bx + a)^3 - 3 \sin(bx + a)}{3b}$$

[In] integrate(cos(b*x+a)^3,x, algorithm="giac")

[Out] -1/3*(sin(b*x + a)^3 - 3*sin(b*x + a))/b

Mupad [B] (verification not implemented)

Time = 13.52 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \cos^3(a + bx) dx = \frac{3 \sin(a + bx) - \sin(a + bx)^3}{3b}$$

[In] `int(cos(a + b*x)^3,x)`

[Out] `(3*sin(a + b*x) - sin(a + b*x)^3)/(3*b)`

3.4 $\int \cos^4(a + bx) dx$

Optimal result	117
Rubi [A] (verified)	117
Mathematica [A] (verified)	118
Maple [A] (verified)	118
Fricas [A] (verification not implemented)	119
Sympy [B] (verification not implemented)	119
Maxima [A] (verification not implemented)	119
Giac [A] (verification not implemented)	120
Mupad [B] (verification not implemented)	120

Optimal result

Integrand size = 8, antiderivative size = 46

$$\int \cos^4(a + bx) dx = \frac{3x}{8} + \frac{3 \cos(a + bx) \sin(a + bx)}{8b} + \frac{\cos^3(a + bx) \sin(a + bx)}{4b}$$

[Out] $3/8*x+3/8*\cos(b*x+a)*\sin(b*x+a)/b+1/4*\cos(b*x+a)^3*\sin(b*x+a)/b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\int \cos^4(a + bx) dx = \frac{\sin(a + bx) \cos^3(a + bx)}{4b} + \frac{3 \sin(a + bx) \cos(a + bx)}{8b} + \frac{3x}{8}$$

[In] Int[Cos[a + b*x]^4,x]

[Out] $(3*x)/8 + (3*\cos[a + b*x]*\sin[a + b*x])/(8*b) + (\cos[a + b*x]^3*\sin[a + b*x])/ (4*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2]

*n]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cos^3(a+bx)\sin(a+bx)}{4b} + \frac{3}{4} \int \cos^2(a+bx) dx \\
&= \frac{3\cos(a+bx)\sin(a+bx)}{8b} + \frac{\cos^3(a+bx)\sin(a+bx)}{4b} + \frac{3 \int 1 dx}{8} \\
&= \frac{3x}{8} + \frac{3\cos(a+bx)\sin(a+bx)}{8b} + \frac{\cos^3(a+bx)\sin(a+bx)}{4b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \cos^4(a+bx) dx = \frac{12(a+bx) + 8\sin(2(a+bx)) + \sin(4(a+bx))}{32b}$$

`[In] Integrate[Cos[a + b*x]^4, x]``[Out] (12*(a + b*x) + 8*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/(32*b)`**Maple [A] (verified)**

Time = 0.75 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.67

method	result
parallelrisch	$\frac{12bx + \sin(4bx+4a) + 8\sin(2bx+2a)}{32b}$
risch	$\frac{3x}{8} + \frac{\sin(4bx+4a)}{32b} + \frac{\sin(2bx+2a)}{4b}$
derivativedivides	$\frac{\left(\cos^3(bx+a) + \frac{3\cos(\frac{bx+a}{2})}{2}\right)\sin(bx+a)}{4} + \frac{3bx}{8} + \frac{3a}{8}$
default	$\frac{\left(\cos^3(bx+a) + \frac{3\cos(\frac{bx+a}{2})}{2}\right)\sin(bx+a)}{4} + \frac{3bx}{8} + \frac{3a}{8}$
norman	$\frac{\frac{3x}{8} + \frac{5\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} - \frac{3\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} - \frac{5\left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3x\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2} + \frac{9x\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4} + \frac{3x\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4}$

`[In] int(cos(b*x+a)^4, x, method=_RETURNVERBOSE)``[Out] 1/32*(12*b*x+sin(4*b*x+4*a)+8*sin(2*b*x+2*a))/b`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.78

$$\int \cos^4(a + bx) dx = \frac{3bx + (2 \cos(bx + a))^3 + 3 \cos(bx + a) \sin(bx + a)}{8b}$$

[In] integrate(cos(b*x+a)^4,x, algorithm="fricas")

[Out] 1/8*(3*b*x + (2*cos(b*x + a))^3 + 3*cos(b*x + a))*sin(b*x + a)/b

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(41) = 82.

Time = 0.17 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.07

$$\int \cos^4(a + bx) dx = \begin{cases} \frac{3x \sin^4(a+bx)}{8} + \frac{3x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{3x \cos^4(a+bx)}{8} + \frac{3 \sin^3(a+bx) \cos(a+bx)}{8b} + \frac{5 \sin(a+bx) \cos^3(a+bx)}{8b} & \text{for } b \neq 0 \\ x \cos^4(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a)**4,x)

[Out] Piecewise((3*x*sin(a + b*x)**4/8 + 3*x*sin(a + b*x)**2*cos(a + b*x)**2/4 + 3*x*cos(a + b*x)**4/8 + 3*sin(a + b*x)**3*cos(a + b*x)/(8*b) + 5*sin(a + b*x)*cos(a + b*x)**3/(8*b), Ne(b, 0)), (x*cos(a)**4, True))

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.72

$$\int \cos^4(a + bx) dx = \frac{12bx + 12a + \sin(4bx + 4a) + 8 \sin(2bx + 2a)}{32b}$$

[In] integrate(cos(b*x+a)^4,x, algorithm="maxima")

[Out] 1/32*(12*b*x + 12*a + sin(4*b*x + 4*a) + 8*sin(2*b*x + 2*a))/b

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.70

$$\int \cos^4(a + bx) dx = \frac{3}{8}x + \frac{\sin(4bx + 4a)}{32b} + \frac{\sin(2bx + 2a)}{4b}$$

[In] integrate(cos(b*x+a)^4,x, algorithm="giac")

[Out] 3/8*x + 1/32*sin(4*b*x + 4*a)/b + 1/4*sin(2*b*x + 2*a)/b

Mupad [B] (verification not implemented)

Time = 13.69 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.67

$$\int \cos^4(a + bx) dx = \frac{3x}{8} + \frac{\frac{\sin(2a+2bx)}{4} + \frac{\sin(4a+4bx)}{32}}{b}$$

[In] int(cos(a + b*x)^4,x)

[Out] (3*x)/8 + (sin(2*a + 2*b*x)/4 + sin(4*a + 4*b*x)/32)/b

3.5 $\int \cos^5(a + bx) dx$

Optimal result	121
Rubi [A] (verified)	121
Mathematica [A] (verified)	122
Maple [A] (verified)	122
Fricas [A] (verification not implemented)	122
Sympy [A] (verification not implemented)	123
Maxima [A] (verification not implemented)	123
Giac [A] (verification not implemented)	123
Mupad [B] (verification not implemented)	124

Optimal result

Integrand size = 8, antiderivative size = 41

$$\int \cos^5(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b}$$

[Out] $\sin(b*x+a)/b-2/3*\sin(b*x+a)^3/b+1/5*\sin(b*x+a)^5/b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2713}

$$\int \cos^5(a + bx) dx = \frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^5, x]$

[Out] $\text{Sin}[a + b*x]/b - (2*\text{Sin}[a + b*x]^3)/(3*b) + \text{Sin}[a + b*x]^5/(5*b)$

Rule 2713

$\text{Int}[\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n - 1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ $\text{FreeQ}\{c, d, x\}$ && $\text{IGtQ}[(n - 1)/2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \cos^5(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{2 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b}$$

[In] Integrate[Cos[a + b*x]^5,x]

[Out] Sin[a + b*x]/b - (2*Sin[a + b*x]^3)/(3*b) + Sin[a + b*x]^5/(5*b)

Maple [A] (verified)

Time = 1.00 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{5b}$	32
default	$\frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{5b}$	32
parallelrisch	$\frac{150 \sin(bx+a) + 3 \sin(5bx+5a) + 25 \sin(3bx+3a)}{240b}$	37
risch	$\frac{5 \sin(bx+a)}{8b} + \frac{\sin(5bx+5a)}{80b} + \frac{5 \sin(3bx+3a)}{48b}$	41

[In] int(cos(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/5/b*(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a)

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \cos^5(a + bx) dx = \frac{(3 \cos(bx + a)^4 + 4 \cos(bx + a)^2 + 8) \sin(bx + a)}{15b}$$

[In] integrate(cos(b*x+a)^5,x, algorithm="fricas")

[Out] 1/15*(3*cos(b*x + a)^4 + 4*cos(b*x + a)^2 + 8)*sin(b*x + a)/b

Sympy [A] (verification not implemented)

Time = 0.24 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.41

$$\int \cos^5(a + bx) dx = \begin{cases} \frac{8 \sin^5(a+bx)}{15b} + \frac{4 \sin^3(a+bx) \cos^2(a+bx)}{3b} + \frac{\sin(a+bx) \cos^4(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^5(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a)**5,x)

[Out] Piecewise((8*sin(a + b*x)**5/(15*b) + 4*sin(a + b*x)**3*cos(a + b*x)**2/(3*b) + sin(a + b*x)*cos(a + b*x)**4/b, Ne(b, 0)), (x*cos(a)**5, True))

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \cos^5(a + bx) dx = \frac{3 \sin^5(bx + a) - 10 \sin^3(bx + a) + 15 \sin(bx + a)}{15b}$$

[In] integrate(cos(b*x+a)^5,x, algorithm="maxima")

[Out] 1/15*(3*sin(b*x + a)^5 - 10*sin(b*x + a)^3 + 15*sin(b*x + a))/b

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.83

$$\int \cos^5(a + bx) dx = \frac{3 \sin^5(bx + a) - 10 \sin^3(bx + a) + 15 \sin(bx + a)}{15b}$$

[In] integrate(cos(b*x+a)^5,x, algorithm="giac")

[Out] 1/15*(3*sin(b*x + a)^5 - 10*sin(b*x + a)^3 + 15*sin(b*x + a))/b

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.76

$$\int \cos^5(a + bx) dx = \frac{\frac{\sin(a+bx)^5}{5} - \frac{2\sin(a+bx)^3}{3} + \sin(a + bx)}{b}$$

[In] int(cos(a + b*x)^5,x)

[Out] (sin(a + b*x) - (2*sin(a + b*x)^3)/3 + sin(a + b*x)^5/5)/b

3.6 $\int \cos^6(a + bx) dx$

Optimal result	125
Rubi [A] (verified)	125
Mathematica [A] (verified)	126
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	127
Sympy [B] (verification not implemented)	127
Maxima [A] (verification not implemented)	128
Giac [A] (verification not implemented)	128
Mupad [B] (verification not implemented)	128

Optimal result

Integrand size = 8, antiderivative size = 67

$$\int \cos^6(a + bx) dx = \frac{5x}{16} + \frac{5 \cos(a + bx) \sin(a + bx)}{16b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b}$$

[Out] 5/16*x+5/16*cos(b*x+a)*sin(b*x+a)/b+5/24*cos(b*x+a)^3*sin(b*x+a)/b+1/6*cos(b*x+a)^5*sin(b*x+a)/b

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\int \cos^6(a + bx) dx = \frac{\sin(a + bx) \cos^5(a + bx)}{6b} + \frac{5 \sin(a + bx) \cos^3(a + bx)}{24b} + \frac{5 \sin(a + bx) \cos(a + bx)}{16b} + \frac{5x}{16}$$

[In] Int[Cos[a + b*x]^6,x]

[Out] (5*x)/16 + (5*Cos[a + b*x]*Sin[a + b*x])/(16*b) + (5*Cos[a + b*x]^3*Sin[a + b*x])/(24*b) + (Cos[a + b*x]^5*Sin[a + b*x])/(6*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]
*((b*Sin[c + d*x])^(n - 1)/(d*n)], x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5}{6} \int \cos^4(a + bx) dx \\
 &= \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5}{8} \int \cos^2(a + bx) dx \\
 &= \frac{5 \cos(a + bx) \sin(a + bx)}{16b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b} + \frac{5 \int 1 dx}{16} \\
 &= \frac{5x}{16} + \frac{5 \cos(a + bx) \sin(a + bx)}{16b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{24b} + \frac{\cos^5(a + bx) \sin(a + bx)}{6b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.64

$$\int \cos^6(a + bx) dx = \frac{60a + 60bx + 45 \sin(2(a + bx)) + 9 \sin(4(a + bx)) + \sin(6(a + bx))}{192b}$$

```
[In] Integrate[Cos[a + b*x]^6,x]
```

```
[Out] (60*a + 60*b*x + 45*Sin[2*(a + b*x)] + 9*Sin[4*(a + b*x)] + Sin[6*(a + b*x)]
)/(192*b)
```

Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.63

method	result
parallelrisc	$\frac{60bx + \sin(6bx+6a) + 9\sin(4bx+4a) + 45\sin(2bx+2a)}{192b}$
risc	$\frac{5x}{16} + \frac{\sin(6bx+6a)}{192b} + \frac{3\sin(4bx+4a)}{64b} + \frac{15\sin(2bx+2a)}{64b}$
derivativedivides	$\frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right) \sin(bx+a)}{6} + \frac{5bx}{16} + \frac{5a}{16}$
default	$\frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right) \sin(bx+a)}{6} + \frac{5bx}{16} + \frac{5a}{16}$
norman	$\frac{5x}{16} + \frac{11 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{5 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} + \frac{15 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} - \frac{15 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{5 \left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} - \frac{11 \left(\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} + \frac{15}{(1+\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right))}$

[In] int(cos(b*x+a)^6,x,method=_RETURNVERBOSE)

[Out] 1/192*(60*b*x+sin(6*b*x+6*a)+9*sin(4*b*x+4*a)+45*sin(2*b*x+2*a))/b

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.69

$$\int \cos^6(a + bx) dx$$

$$= \frac{15bx + (8 \cos(bx + a)^5 + 10 \cos(bx + a)^3 + 15 \cos(bx + a)) \sin(bx + a)}{48b}$$

[In] integrate(cos(b*x+a)^6,x, algorithm="fricas")

[Out] 1/48*(15*b*x + (8*cos(b*x + a)^5 + 10*cos(b*x + a)^3 + 15*cos(b*x + a))*sin(b*x + a))/b

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(61) = 122.

Time = 0.37 (sec) , antiderivative size = 139, normalized size of antiderivative = 2.07

$$\int \cos^6(a + bx) dx$$

$$= \begin{cases} \frac{5x \sin^6(a+bx)}{16} + \frac{15x \sin^4(a+bx) \cos^2(a+bx)}{16} + \frac{15x \sin^2(a+bx) \cos^4(a+bx)}{16} + \frac{5x \cos^6(a+bx)}{16} + \frac{5 \sin^5(a+bx) \cos(a+bx)}{16b} + \frac{5 \sin^3(a+bx) \cos^3(a+bx)}{16b} \\ x \cos^6(a) \end{cases}$$

[In] integrate(cos(b*x+a)**6,x)

[Out] Piecewise((5*x*sin(a + b*x)**6/16 + 15*x*sin(a + b*x)**4*cos(a + b*x)**2/16 + 15*x*sin(a + b*x)**2*cos(a + b*x)**4/16 + 5*x*cos(a + b*x)**6/16 + 5*sin(a + b*x)**5*cos(a + b*x)/(16*b) + 5*sin(a + b*x)**3*cos(a + b*x)**3/(6*b) + 11*sin(a + b*x)*cos(a + b*x)**5/(16*b), Ne(b, 0)), (x*cos(a)**6, True))

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.72

$$\int \cos^6(a + bx) dx = -\frac{4 \sin(2bx + 2a)^3 - 60bx - 60a - 9 \sin(4bx + 4a) - 48 \sin(2bx + 2a)}{192b}$$

[In] integrate(cos(b*x+a)^6,x, algorithm="maxima")

[Out] -1/192*(4*sin(2*b*x + 2*a)^3 - 60*b*x - 60*a - 9*sin(4*b*x + 4*a) - 48*sin(2*b*x + 2*a))/b

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.69

$$\int \cos^6(a + bx) dx = \frac{5}{16}x + \frac{\sin(6bx + 6a)}{192b} + \frac{3 \sin(4bx + 4a)}{64b} + \frac{15 \sin(2bx + 2a)}{64b}$$

[In] integrate(cos(b*x+a)^6,x, algorithm="giac")

[Out] 5/16*x + 1/192*sin(6*b*x + 6*a)/b + 3/64*sin(4*b*x + 4*a)/b + 15/64*sin(2*b*x + 2*a)/b

Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.63

$$\int \cos^6(a + bx) dx = \frac{5x}{16} + \frac{\frac{15 \sin(2a+2bx)}{64} + \frac{3 \sin(4a+4bx)}{64} + \frac{\sin(6a+6bx)}{192}}{b}$$

[In] int(cos(a + b*x)^6,x)

[Out] (5*x)/16 + ((15*sin(2*a + 2*b*x))/64 + (3*sin(4*a + 4*b*x))/64 + sin(6*a + 6*b*x)/192)/b

3.7 $\int \cos^7(a + bx) dx$

Optimal result	129
Rubi [A] (verified)	129
Mathematica [A] (verified)	130
Maple [A] (verified)	130
Fricas [A] (verification not implemented)	130
Sympy [A] (verification not implemented)	131
Maxima [A] (verification not implemented)	131
Giac [A] (verification not implemented)	131
Mupad [B] (verification not implemented)	132

Optimal result

Integrand size = 8, antiderivative size = 54

$$\int \cos^7(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b}$$

[Out] $\sin(b*x+a)/b - \sin(b*x+a)^3/b + 3/5*\sin(b*x+a)^5/b - 1/7*\sin(b*x+a)^7/b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2713}

$$\int \cos^7(a + bx) dx = -\frac{\sin^7(a + bx)}{7b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^3(a + bx)}{b} + \frac{\sin(a + bx)}{b}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^7, x]$

[Out] $\text{Sin}[a + b*x]/b - \text{Sin}[a + b*x]^3/b + (3*\text{Sin}[a + b*x]^5)/(5*b) - \text{Sin}[a + b*x]^7/(7*b)$

Rule 2713

$\text{Int}[\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n - 1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ $\text{FreeQ}\{c, d\}, x]$
 $\&\& \text{IGtQ}[(n - 1)/2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int (1 - 3x^2 + 3x^4 - x^6) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int \cos^7(a + bx) dx = \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{b} + \frac{3 \sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b}$$

[In] Integrate[Cos[a + b*x]^7,x]

[Out] Sin[a + b*x]/b - Sin[a + b*x]^3/b + (3*Sin[a + b*x]^5)/(5*b) - Sin[a + b*x]^7/(7*b)

Maple [A] (verified)

Time = 1.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.78

method	result	size
derivativedivides	$\frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{7b}$	42
default	$\frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{7b}$	42
parallelrisc	$\frac{1225 \sin(bx+a) + 5 \sin(7bx+7a) + 49 \sin(5bx+5a) + 245 \sin(3bx+3a)}{2240b}$	48
risc	$\frac{35 \sin(bx+a)}{64b} + \frac{\sin(7bx+7a)}{448b} + \frac{7 \sin(5bx+5a)}{320b} + \frac{7 \sin(3bx+3a)}{64b}$	55

[In] int(cos(b*x+a)^7,x,method=_RETURNVERBOSE)

[Out] 1/7/b*(16/5+cos(b*x+a)^6+6/5*cos(b*x+a)^4+8/5*cos(b*x+a)^2)*sin(b*x+a)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.80

$$\int \cos^7(a + bx) dx = \frac{(5 \cos(bx + a)^6 + 6 \cos(bx + a)^4 + 8 \cos(bx + a)^2 + 16) \sin(bx + a)}{35b}$$

[In] integrate(cos(b*x+a)^7,x, algorithm="fricas")

[Out] 1/35*(5*cos(b*x + a)^6 + 6*cos(b*x + a)^4 + 8*cos(b*x + a)^2 + 16)*sin(b*x + a)/b

Sympy [A] (verification not implemented)

Time = 0.51 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

$$\int \cos^7(a + bx) dx = \begin{cases} \frac{16 \sin^7(a+bx)}{35b} + \frac{8 \sin^5(a+bx) \cos^2(a+bx)}{5b} + \frac{2 \sin^3(a+bx) \cos^4(a+bx)}{b} + \frac{\sin(a+bx) \cos^6(a+bx)}{b} & \text{for } b \neq 0 \\ x \cos^7(a) & \text{otherwise} \end{cases}$$

[In] integrate(cos(b*x+a)**7,x)

[Out] Piecewise((16*sin(a + b*x)**7/(35*b) + 8*sin(a + b*x)**5*cos(a + b*x)**2/(5*b) + 2*sin(a + b*x)**3*cos(a + b*x)**4/b + sin(a + b*x)*cos(a + b*x)**6/b, Ne(b, 0)), (x*cos(a)**7, True))

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \cos^7(a + bx) dx = -\frac{5 \sin^7(bx + a) - 21 \sin^5(bx + a) + 35 \sin^3(bx + a) - 35 \sin(bx + a)}{35 b}$$

[In] integrate(cos(b*x+a)^7,x, algorithm="maxima")

[Out] -1/35*(5*sin(b*x + a)^7 - 21*sin(b*x + a)^5 + 35*sin(b*x + a)^3 - 35*sin(b*x + a))/b

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \cos^7(a + bx) dx = -\frac{5 \sin^7(bx + a) - 21 \sin^5(bx + a) + 35 \sin^3(bx + a) - 35 \sin(bx + a)}{35 b}$$

[In] integrate(cos(b*x+a)^7,x, algorithm="giac")

[Out] -1/35*(5*sin(b*x + a)^7 - 21*sin(b*x + a)^5 + 35*sin(b*x + a)^3 - 35*sin(b*x + a))/b

Mupad [B] (verification not implemented)

Time = 13.99 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.80

$$\int \cos^7(a + bx) dx$$

$$= -\frac{\sin(a + bx) (5 \sin(a + bx)^6 - 21 \sin(a + bx)^4 + 35 \sin(a + bx)^2 - 35)}{35b}$$

[In] `int(cos(a + b*x)^7,x)`

[Out] `-(sin(a + b*x)*(35*sin(a + b*x)^2 - 21*sin(a + b*x)^4 + 5*sin(a + b*x)^6 - 35))/(35*b)`

3.8 $\int \cos^8(a + bx) dx$

Optimal result	133
Rubi [A] (verified)	133
Mathematica [A] (verified)	134
Maple [A] (verified)	135
Fricas [A] (verification not implemented)	135
Sympy [B] (verification not implemented)	136
Maxima [A] (verification not implemented)	136
Giac [A] (verification not implemented)	136
Mupad [B] (verification not implemented)	137

Optimal result

Integrand size = 8, antiderivative size = 88

$$\int \cos^8(a + bx) dx = \frac{35x}{128} + \frac{35 \cos(a + bx) \sin(a + bx)}{128b} + \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b}$$

[Out] 35/128*x+35/128*cos(b*x+a)*sin(b*x+a)/b+35/192*cos(b*x+a)^3*sin(b*x+a)/b+7/48*cos(b*x+a)^5*sin(b*x+a)/b+1/8*cos(b*x+a)^7*sin(b*x+a)/b

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\int \cos^8(a + bx) dx = \frac{\sin(a + bx) \cos^7(a + bx)}{8b} + \frac{7 \sin(a + bx) \cos^5(a + bx)}{48b} + \frac{35 \sin(a + bx) \cos^3(a + bx)}{192b} + \frac{35 \sin(a + bx) \cos(a + bx)}{128b} + \frac{35x}{128}$$

[In] Int[Cos[a + b*x]^8,x]

[Out] (35*x)/128 + (35*Cos[a + b*x]*Sin[a + b*x])/(128*b) + (35*Cos[a + b*x]^3*Sin[a + b*x])/(192*b) + (7*Cos[a + b*x]^5*Sin[a + b*x])/(48*b) + (Cos[a + b*x]^7*Sin[a + b*x])/(8*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{7}{8} \int \cos^6(a + bx) dx \\
&= \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{35}{48} \int \cos^4(a + bx) dx \\
&= \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} \\
&\quad + \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{35}{64} \int \cos^2(a + bx) dx \\
&= \frac{35 \cos(a + bx) \sin(a + bx)}{128b} + \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} \\
&\quad + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b} + \frac{35 \int 1 dx}{128} \\
&= \frac{35x}{128} + \frac{35 \cos(a + bx) \sin(a + bx)}{128b} + \frac{35 \cos^3(a + bx) \sin(a + bx)}{192b} \\
&\quad + \frac{7 \cos^5(a + bx) \sin(a + bx)}{48b} + \frac{\cos^7(a + bx) \sin(a + bx)}{8b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.62

$$\begin{aligned}
&\int \cos^8(a + bx) dx \\
&= \frac{840a + 840bx + 672 \sin(2(a + bx)) + 168 \sin(4(a + bx)) + 32 \sin(6(a + bx)) + 3 \sin(8(a + bx))}{3072b}
\end{aligned}$$

```
[In] Integrate[Cos[a + b*x]^8, x]
```

```
[Out] (840*a + 840*b*x + 672*Sin[2*(a + b*x)] + 168*Sin[4*(a + b*x)] + 32*Sin[6*(a + b*x)] + 3*Sin[8*(a + b*x)])/(3072*b)
```

Maple [A] (verified)

Time = 1.15 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.62

method	result
parallelrisc	$\frac{840bx+3\sin(8bx+8a)+32\sin(6bx+6a)+168\sin(4bx+4a)+672\sin(2bx+2a)}{3072b}$
derivativedivides	$\frac{\left(\cos^7(bx+a)+\frac{7(\cos^5(bx+a))}{6}+\frac{35(\cos^3(bx+a))}{24}+\frac{35\cos(bx+a)}{16}\right)\sin(bx+a)}{8} + \frac{35bx}{128} + \frac{35a}{128}$
default	$\frac{\left(\cos^7(bx+a)+\frac{7(\cos^5(bx+a))}{6}+\frac{35(\cos^3(bx+a))}{24}+\frac{35\cos(bx+a)}{16}\right)\sin(bx+a)}{8} + \frac{35bx}{128} + \frac{35a}{128}$
risc	$\frac{35x}{128} + \frac{\sin(8bx+8a)}{1024b} + \frac{\sin(6bx+6a)}{96b} + \frac{7\sin(4bx+4a)}{128b} + \frac{7\sin(2bx+2a)}{32b}$
norman	$\frac{35x}{128} + \frac{93\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{64b} + \frac{91\left(\tan^3\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{192b} + \frac{1799\left(\tan^5\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{192b} - \frac{1085\left(\tan^7\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{192b} + \frac{1085\left(\tan^9\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{192b} - \frac{1799\left(\tan^{11}\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{192b}$

```
[In] int(cos(b*x+a)^8,x,method=_RETURNVERBOSE)
```

```
[Out] 1/3072*(840*b*x+3*sin(8*b*x+8*a)+32*sin(6*b*x+6*a)+168*sin(4*b*x+4*a)+672*sin(2*b*x+2*a))/b
```

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.64

$$\int \cos^8(a+bx) dx$$

$$= \frac{105bx + (48\cos^7(bx+a) + 56\cos^5(bx+a) + 70\cos^3(bx+a) + 105\cos(bx+a))\sin(bx+a)}{384b}$$

```
[In] integrate(cos(b*x+a)^8,x, algorithm="fricas")
```

```
[Out] 1/384*(105*b*x + (48*cos(b*x + a)^7 + 56*cos(b*x + a)^5 + 70*cos(b*x + a)^3 + 105*cos(b*x + a))*sin(b*x + a))/b
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(82) = 164$.

Time = 0.71 (sec) , antiderivative size = 184, normalized size of antiderivative = 2.09

$$\int \cos^8(a + bx) dx$$

$$= \begin{cases} \frac{35x \sin^8(a+bx)}{128} + \frac{35x \sin^6(a+bx) \cos^2(a+bx)}{32} + \frac{105x \sin^4(a+bx) \cos^4(a+bx)}{64} + \frac{35x \sin^2(a+bx) \cos^6(a+bx)}{32} + \frac{35x \cos^8(a+bx)}{128} + \frac{35}{128} \\ x \cos^8(a) \end{cases}$$

[In] integrate(cos(b*x+a)**8,x)

[Out] Piecewise((35*x*sin(a + b*x)**8/128 + 35*x*sin(a + b*x)**6*cos(a + b*x)**2/32 + 105*x*sin(a + b*x)**4*cos(a + b*x)**4/64 + 35*x*sin(a + b*x)**2*cos(a + b*x)**6/32 + 35*x*cos(a + b*x)**8/128 + 35*sin(a + b*x)**7*cos(a + b*x)/(128*b) + 385*sin(a + b*x)**5*cos(a + b*x)**3/(384*b) + 511*sin(a + b*x)**3*cos(a + b*x)**5/(384*b) + 93*sin(a + b*x)*cos(a + b*x)**7/(128*b), Ne(b, 0)), (x*cos(a)**8, True))

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.67

$$\int \cos^8(a + bx) dx = \frac{128 \sin(2bx + 2a)^3 - 840bx - 840a - 3 \sin(8bx + 8a) - 168 \sin(4bx + 4a) - 768 \sin(2bx + 2a)}{3072b}$$

[In] integrate(cos(b*x+a)^8,x, algorithm="maxima")

[Out] -1/3072*(128*sin(2*b*x + 2*a)^3 - 840*b*x - 840*a - 3*sin(8*b*x + 8*a) - 168*sin(4*b*x + 4*a) - 768*sin(2*b*x + 2*a))/b

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.68

$$\int \cos^8(a + bx) dx = \frac{35}{128}x + \frac{\sin(8bx + 8a)}{1024b} + \frac{\sin(6bx + 6a)}{96b} + \frac{7 \sin(4bx + 4a)}{128b} + \frac{7 \sin(2bx + 2a)}{32b}$$

[In] integrate(cos(b*x+a)^8,x, algorithm="giac")

[Out] $35/128*x + 1/1024*\sin(8*b*x + 8*a)/b + 1/96*\sin(6*b*x + 6*a)/b + 7/128*\sin(4*b*x + 4*a)/b + 7/32*\sin(2*b*x + 2*a)/b$

Mupad [B] (verification not implemented)

Time = 14.51 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.60

$$\int \cos^8(a + bx) dx = \frac{35x}{128} + \frac{\frac{7 \sin(2a+2bx)}{32} + \frac{7 \sin(4a+4bx)}{128} + \frac{\sin(6a+6bx)}{96} + \frac{\sin(8a+8bx)}{1024}}{b}$$

[In] int(cos(a + b*x)^8,x)

[Out] $(35*x)/128 + ((7*\sin(2*a + 2*b*x))/32 + (7*\sin(4*a + 4*b*x))/128 + \sin(6*a + 6*b*x)/96 + \sin(8*a + 8*b*x)/1024)/b$

3.9 $\int \cos^{\frac{7}{2}}(a + bx) dx$

Optimal result	138
Rubi [A] (verified)	138
Mathematica [A] (verified)	139
Maple [B] (verified)	139
Fricas [C] (verification not implemented)	140
Sympy [F(-1)]	140
Maxima [F]	140
Giac [F]	141
Mupad [B] (verification not implemented)	141

Optimal result

Integrand size = 10, antiderivative size = 65

$$\int \cos^{\frac{7}{2}}(a + bx) dx = \frac{10 \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b} + \frac{10\sqrt{\cos(a + bx)} \sin(a + bx)}{21b} + \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b}$$

[Out] $10/21*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b+2/7*\cos(b*x+a)^{(5/2)}*\sin(b*x+a)/b+10/21*\sin(b*x+a)*\cos(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2720}

$$\int \cos^{\frac{7}{2}}(a + bx) dx = \frac{10 \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b} + \frac{2 \sin(a + bx) \cos^{\frac{5}{2}}(a + bx)}{7b} + \frac{10 \sin(a + bx) \sqrt{\cos(a + bx)}}{21b}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(7/2)}, x]$

[Out] $(10*\operatorname{EllipticF}[(a + b*x)/2, 2])/(21*b) + (10*\operatorname{Sqrt}[\operatorname{Cos}[a + b*x]]*\operatorname{Sin}[a + b*x])/(21*b) + (2*\operatorname{Cos}[a + b*x]^{(5/2)}*\operatorname{Sin}[a + b*x])/(7*b)$

Rule 2715

$\operatorname{Int}(((b_.)*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol) \rightarrow \operatorname{Simp}[(-b)*\operatorname{Cos}[c + d*x]*(b*\operatorname{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \operatorname{Dist}[b^2*((n-1)/n), \operatorname{Int}[(b*\operatorname{Sin}[$

$c + d*x]^{(n - 2), x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2 * n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} + \frac{5}{7} \int \cos^{\frac{3}{2}}(a + bx) dx \\ &= \frac{10 \sqrt{\cos(a + bx)} \sin(a + bx)}{21b} + \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} + \frac{5}{21} \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{10 \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b} + \frac{10 \sqrt{\cos(a + bx)} \sin(a + bx)}{21b} + \frac{2 \cos^{\frac{5}{2}}(a + bx) \sin(a + bx)}{7b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.78

$$\begin{aligned} &\int \cos^{\frac{7}{2}}(a + bx) dx \\ &= \frac{20 \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + \sqrt{\cos(a + bx)}(23 \sin(a + bx) + 3 \sin(3(a + bx)))}{42b} \end{aligned}$$

[In] Integrate[Cos[a + b*x]^(7/2), x]

[Out] (20*EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*(23*Sin[a + b*x] + 3*Sin[3*(a + b*x)]))/(42*b)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 198 vs. 2(81) = 162.

Time = 4.63 (sec) , antiderivative size = 199, normalized size of antiderivative = 3.06

method	result
default	$-\frac{2\sqrt{-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\left(48\left(\cos^9\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-120\left(\cos^7\left(\frac{bx}{2}+\frac{a}{2}\right)\right)+128\left(\cos^5\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-72\left(\cos^3\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)}{21\sqrt{-2\left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)+\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}}$

[In] int(cos(b*x+a)^(7/2), x, method=_RETURNVERBOSE)

```
[Out] -2/21*((-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(48*cos(1/2*
b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b
*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2
))*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*
b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(-1+2*cos(1/2*b
*x+1/2*a)^2)^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.23

$$\int \cos^{\frac{7}{2}}(a + bx) dx$$

$$= \frac{2(3 \cos^2(bx + a) + 5) \sqrt{\cos(bx + a)} \sin(bx + a) - 5i \sqrt{2} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + 5i \sqrt{2} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{21b}$$

```
[In] integrate(cos(b*x+a)^(7/2),x, algorithm="fricas")
```

```
[Out] 1/21*(2*(3*cos(b*x + a)^2 + 5)*sqrt(cos(b*x + a))*sin(b*x + a) - 5*I*sqrt(2)
)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + 5*I*sqrt(2)*w
eierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b
```

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{7}{2}}(a + bx) dx = \text{Timed out}$$

```
[In] integrate(cos(b*x+a)**(7/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^{\frac{7}{2}}(a + bx) dx = \int \cos^{\frac{7}{2}}(bx + a) dx$$

```
[In] integrate(cos(b*x+a)^(7/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(b*x + a)^(7/2), x)
```

Giac [F]

$$\int \cos^{\frac{7}{2}}(a + bx) dx = \int \cos(bx + a)^{\frac{7}{2}} dx$$

[In] integrate(cos(b*x+a)^(7/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(7/2), x)

Mupad [B] (verification not implemented)

Time = 14.50 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \cos^{\frac{7}{2}}(a + bx) dx = -\frac{2 \cos(a + bx)^{9/2} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{13}{4}; \cos(a + bx)^2\right)}{9b \sqrt{\sin(a + bx)^2}}$$

[In] int(cos(a + b*x)^(7/2),x)

[Out] -(2*cos(a + b*x)^(9/2)*sin(a + b*x)*hypergeom([1/2, 9/4], 13/4, cos(a + b*x)^2))/(9*b*(sin(a + b*x)^2)^(1/2))

3.10 $\int \cos^{\frac{5}{2}}(a + bx) dx$

Optimal result	142
Rubi [A] (verified)	142
Mathematica [A] (verified)	143
Maple [B] (verified)	143
Fricas [C] (verification not implemented)	144
Sympy [F(-1)]	144
Maxima [F]	144
Giac [F]	145
Mupad [B] (verification not implemented)	145

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \frac{6E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b} + \frac{2 \cos^{\frac{3}{2}}(a + bx) \sin(a + bx)}{5b}$$

[Out] $6/5 * (\cos(1/2*a+1/2*b*x)^2)^{(1/2)} / \cos(1/2*a+1/2*b*x) * \text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)}) / b + 2/5 * \cos(b*x+a)^{(3/2)} * \sin(b*x+a) / b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2719}

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \frac{6E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b} + \frac{2 \sin(a + bx) \cos^{\frac{3}{2}}(a + bx)}{5b}$$

[In] Int[Cos[a + b*x]^(5/2),x]

[Out] $(6 * \text{EllipticE}[(a + b*x)/2, 2]) / (5*b) + (2 * \text{Cos}[a + b*x]^{(3/2)} * \text{Sin}[a + b*x]) / (5*b)$

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \cos^{\frac{3}{2}}(a + bx) \sin(a + bx)}{5b} + \frac{3}{5} \int \sqrt{\cos(a + bx)} dx \\ &= \frac{6E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b} + \frac{2 \cos^{\frac{3}{2}}(a + bx) \sin(a + bx)}{5b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \frac{6E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sqrt{\cos(a + bx)} \sin(2(a + bx))}{5b}$$

```
[In] Integrate[Cos[a + b*x]^(5/2), x]
```

```
[Out] (6*EllipticE[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*Sin[2*(a + b*x)])/(5*b)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(62) = 124.

Time = 2.89 (sec) , antiderivative size = 202, normalized size of antiderivative = 4.81

method	result
default	$-\frac{2\sqrt{(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}(\sin^2(\frac{bx}{2}+\frac{a}{2}))}{5\sqrt{-2(\sin^4(\frac{bx}{2}+\frac{a}{2}))+\sin^2(\frac{bx}{2}+\frac{a}{2})}} \left(-8 \cos\left(\frac{bx}{2}+\frac{a}{2}\right) \left(\sin^6\left(\frac{bx}{2}+\frac{a}{2}\right) + 8 \left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right) \right) \cos\left(\frac{bx}{2}+\frac{a}{2}\right) - 2 \left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right) \right) \right) \right) \sqrt{-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2}))} \sin\left(\frac{bx}{2}+\frac{a}{2}\right)$

```
[In] int(cos(b*x+a)^(5/2), x, method=_RETURNVERBOSE)
```

```
[Out] -2/5*((-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(-8*cos(1/2*b
*x+1/2*a)*sin(1/2*b*x+1/2*a)^6+8*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-2*
sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a)-3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*s
in(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2)))/(-2*sin
(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(-1+2*cos(
1/2*b*x+1/2*a)^2)^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.76

$$\int \cos^{\frac{5}{2}}(a + bx) dx$$

$$= \frac{2 \cos(bx + a)^{\frac{3}{2}} \sin(bx + a) + 3i \sqrt{2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) - 3i \sqrt{2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)))}{5b}$$

```
[In] integrate(cos(b*x+a)^(5/2),x, algorithm="fricas")
```

```
[Out] 1/5*(2*cos(b*x + a)^(3/2)*sin(b*x + a) + 3*I*sqrt(2)*weierstrassZeta(-4, 0,
weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - 3*I*sqrt(2)*w
eierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x +
a))))/b
```

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \text{Timed out}$$

```
[In] integrate(cos(b*x+a)**(5/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \int \cos(bx + a)^{\frac{5}{2}} dx$$

```
[In] integrate(cos(b*x+a)^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(b*x + a)^(5/2), x)
```


Giac [F]

$$\int \cos^{\frac{5}{2}}(a + bx) dx = \int \cos(bx + a)^{\frac{5}{2}} dx$$

[In] integrate(cos(b*x+a)^(5/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(5/2), x)

Mupad [B] (verification not implemented)

Time = 14.44 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \cos^{\frac{5}{2}}(a + bx) dx = -\frac{2 \cos(a + bx)^{7/2} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{11}{4}; \cos(a + bx)^2\right)}{7b \sqrt{\sin(a + bx)^2}}$$

[In] int(cos(a + b*x)^(5/2),x)

[Out] -(2*cos(a + b*x)^(7/2)*sin(a + b*x)*hypergeom([1/2, 7/4], 11/4, cos(a + b*x)^2))/(7*b*(sin(a + b*x)^2)^(1/2))

3.11 $\int \cos^{\frac{3}{2}}(a + bx) dx$

Optimal result	146
Rubi [A] (verified)	146
Mathematica [A] (verified)	147
Maple [B] (verified)	147
Fricas [C] (verification not implemented)	148
Sympy [F]	148
Maxima [F]	148
Giac [F]	148
Mupad [B] (verification not implemented)	149

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b} + \frac{2\sqrt{\cos(a + bx)} \sin(a + bx)}{3b}$$

[Out] $2/3*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b+2/3*\sin(b*x+a)*\cos(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2720}

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b} + \frac{2 \sin(a + bx) \sqrt{\cos(a + bx)}}{3b}$$

[In] `Int[Cos[a + b*x]^(3/2), x]`

[Out] $(2*\operatorname{EllipticF}[(a + b*x)/2, 2])/(3*b) + (2*\operatorname{Sqrt}[\operatorname{Cos}[a + b*x]]*\operatorname{Sin}[a + b*x])/(3*b)$

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2\sqrt{\cos(a+bx)}\sin(a+bx)}{3b} + \frac{1}{3} \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\ &= \frac{2\text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b} + \frac{2\sqrt{\cos(a+bx)}\sin(a+bx)}{3b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \cos^{\frac{3}{2}}(a+bx) dx = \frac{2\left(\text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) + \sqrt{\cos(a+bx)}\sin(a+bx)\right)}{3b}$$

```
[In] Integrate[Cos[a + b*x]^(3/2), x]
```

```
[Out] (2*(EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*Sin[a + b*x]))/(3*b)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 178 vs. 2(62) = 124.

Time = 2.15 (sec) , antiderivative size = 179, normalized size of antiderivative = 4.26

method	result
default	$-\frac{2\sqrt{\left(-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{3\sqrt{-2\left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)+\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}}{b}$

```
[In] int(cos(b*x+a)^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] -2/3*((-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(4*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2)))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(-1+2*cos(1/2*b*x+1/2*a)^2)^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.62

$$\int \cos^{\frac{3}{2}}(a + bx) dx$$

$$= \frac{2 \sqrt{\cos(bx + a)} \sin(bx + a) - i \sqrt{2} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{3b}$$

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="fricas")

[Out] 1/3*(2*sqrt(cos(b*x + a))*sin(b*x + a) - I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b

Sympy [F]

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \int \cos^{\frac{3}{2}}(a + bx) dx$$

[In] integrate(cos(b*x+a)**(3/2),x)

[Out] Integral(cos(a + b*x)**(3/2), x)

Maxima [F]

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \int \cos(bx + a)^{\frac{3}{2}} dx$$

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(3/2), x)

Giac [F]

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \int \cos(bx + a)^{\frac{3}{2}} dx$$

[In] integrate(cos(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(3/2), x)

Mupad [B] (verification not implemented)

Time = 14.41 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.83

$$\int \cos^{\frac{3}{2}}(a + bx) dx = \frac{2F\left(\frac{a}{2} + \frac{bx}{2} \mid 2\right)}{3b} + \frac{2\sqrt{\cos(a + bx)} \sin(a + bx)}{3b}$$

[In] int(cos(a + b*x)^(3/2),x)

[Out] (2*ellipticF(a/2 + (b*x)/2, 2))/(3*b) + (2*cos(a + b*x)^(1/2)*sin(a + b*x))/(3*b)

3.12 $\int \sqrt{\cos(a + bx)} dx$

Optimal result	150
Rubi [A] (verified)	150
Mathematica [A] (verified)	151
Maple [B] (verified)	151
Fricas [C] (verification not implemented)	151
Sympy [F]	152
Maxima [F]	152
Giac [F]	152
Mupad [B] (verification not implemented)	152

Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \sqrt{\cos(a + bx)} dx = \frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b}$$

[Out] $2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2719}

$$\int \sqrt{\cos(a + bx)} dx = \frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b}$$

[In] Int[Sqrt[Cos[a + b*x]], x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/b

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\text{integral} = \frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \sqrt{\cos(a + bx)} dx = \frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b}$$

[In] Integrate[Sqrt[Cos[a + b*x]],x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/b

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. 2(42) = 84.

Time = 1.61 (sec) , antiderivative size = 133, normalized size of antiderivative = 8.31

method	result
default	$\frac{2\sqrt{\left(-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\sqrt{-2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)+1}E\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)+\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)\sqrt{-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}b}$
risch	$-\frac{i\sqrt{2}\sqrt{\left(e^{2i(bx+a)}+1\right)e^{-i(bx+a)}}}{b} - i\left(\frac{2\left(e^{2i(bx+a)}+1\right)}{\sqrt{\left(e^{2i(bx+a)}+1\right)e^{i(bx+a)}}} + \frac{i\sqrt{-i\left(e^{i(bx+a)}+i\right)}\sqrt{2}\sqrt{i\left(e^{i(bx+a)}-i\right)}\sqrt{ie^{i(bx+a)}}\left(-2iE\left(\sqrt{-i\left(e^{i(bx+a)}-i\right)},\sqrt{2}\right)\right)}{\sqrt{e^{3i(bx+a)}+e^{i(bx+a)}}}\right)b$

[In] int(cos(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 2*((-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(-1+2*cos(1/2*b*x+1/2*a)^2)^(1/2)/b

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 3.56

$$\int \sqrt{\cos(a + bx)} dx = \frac{i\sqrt{2}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) - i\sqrt{2}\text{weierstrassZeta}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{b}$$

[In] integrate(cos(b*x+a)^(1/2),x, algorithm="fricas")

[Out] (I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))))/b

Sympy [F]

$$\int \sqrt{\cos(a + bx)} dx = \int \sqrt{\cos(a + bx)} dx$$

[In] integrate(cos(b*x+a)**(1/2),x)

[Out] Integral(sqrt(cos(a + b*x)), x)

Maxima [F]

$$\int \sqrt{\cos(a + bx)} dx = \int \sqrt{\cos(bx + a)} dx$$

[In] integrate(cos(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(cos(b*x + a)), x)

Giac [F]

$$\int \sqrt{\cos(a + bx)} dx = \int \sqrt{\cos(bx + a)} dx$$

[In] integrate(cos(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(b*x + a)), x)

Mupad [B] (verification not implemented)

Time = 14.41 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \sqrt{\cos(a + bx)} dx = \frac{2 E\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b}$$

[In] int(cos(a + b*x)^(1/2),x)

[Out] (2*ellipticE(a/2 + (b*x)/2, 2))/b

3.13 $\int \frac{1}{\sqrt{\cos(a+bx)}} dx$

Optimal result	153
Rubi [A] (verified)	153
Mathematica [A] (verified)	154
Maple [C] (verified)	154
Fricas [C] (verification not implemented)	154
Sympy [F]	155
Maxima [F]	155
Giac [F]	155
Mupad [B] (verification not implemented)	155

Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \frac{1}{\sqrt{\cos(a+bx)}} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b}$$

[Out] $2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2720}

$$\int \frac{1}{\sqrt{\cos(a+bx)}} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b}$$

[In] `Int[1/Sqrt[Cos[a + b*x]], x]`

[Out] `(2*EllipticF[(a + b*x)/2, 2])/b`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\text{integral} = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{b}$$

[In] Integrate[1/Sqrt[Cos[a + b*x]],x]

[Out] (2*EllipticF[(a + b*x)/2, 2])/b

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{2 \operatorname{am}^{-1}\left(\frac{bx}{2} + \frac{a}{2} \sqrt{2}\right)}{b}$	18

[In] int(1/cos(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/b*InverseJacobiAM(1/2*b*x+1/2*a,2^(1/2))

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 3.19

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \frac{-i \sqrt{2} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{b}$$

[In] integrate(1/cos(b*x+a)^(1/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/b

Sympy [F]

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \int \frac{1}{\sqrt{\cos(a + bx)}} dx$$

[In] integrate(1/cos(b*x+a)**(1/2),x)

[Out] Integral(1/sqrt(cos(a + b*x)), x)

Maxima [F]

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \int \frac{1}{\sqrt{\cos(bx + a)}} dx$$

[In] integrate(1/cos(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(cos(b*x + a)), x)

Giac [F]

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \int \frac{1}{\sqrt{\cos(bx + a)}} dx$$

[In] integrate(1/cos(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(cos(b*x + a)), x)

Mupad [B] (verification not implemented)

Time = 13.59 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{1}{\sqrt{\cos(a + bx)}} dx = \frac{2F\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b}$$

[In] int(1/cos(a + b*x)^(1/2),x)

[Out] (2*ellipticF(a/2 + (b*x)/2, 2))/b

3.14 $\int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx$

Optimal result	156
Rubi [A] (verified)	156
Mathematica [A] (verified)	157
Maple [B] (verified)	157
Fricas [C] (verification not implemented)	158
Sympy [F]	158
Maxima [F]	158
Giac [F]	159
Mupad [B] (verification not implemented)	159

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx = -\frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b} + \frac{2\sin(a+bx)}{b\sqrt{\cos(a+bx)}}$$

[Out] $-2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})/b+2*\sin(b*x+a)/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2719}

$$\int \frac{1}{\cos^{\frac{3}{2}}(a+bx)} dx = \frac{2\sin(a+bx)}{b\sqrt{\cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

[In] `Int[Cos[a + b*x]^(-3/2),x]`

[Out] `(-2*EllipticE[(a + b*x)/2, 2])/b + (2*Sin[a + b*x])/(b*sqrt[Cos[a + b*x]])`

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(a + bx)}{b \sqrt{\cos(a + bx)}} - \int \sqrt{\cos(a + bx)} dx \\ &= -\frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b} + \frac{2 \sin(a + bx)}{b \sqrt{\cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx = -\frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b} + \frac{2 \sin(a + bx)}{b \sqrt{\cos(a + bx)}}$$

```
[In] Integrate[Cos[a + b*x]^(-3/2), x]
```

```
[Out] (-2*EllipticE[(a + b*x)/2, 2])/b + (2*Sin[a + b*x])/(b*Sqrt[Cos[a + b*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 181 vs. 2(62) = 124.

Time = 1.30 (sec) , antiderivative size = 182, normalized size of antiderivative = 4.79

method	result
default	$-\frac{2\left(-2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{-1 + 2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}}{b}$

```
[In] int(1/cos(b*x+a)^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] -2*(-2*cos(1/2*b*x+1/2*a)*(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*sin(1/2*b*x+1/2*a)^2+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2)))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(-1+2*cos(1/2*b*x+1/2*a)^2)^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 93, normalized size of antiderivative = 2.45

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx$$

$$= \frac{-i\sqrt{2}\cos(bx + a)\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i\sin(bx + a))) + i\sqrt{2}}$$

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*sqrt(cos(b*x + a))*sin(b*x + a))/(b*cos(b*x + a))

Sympy [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx$$

[In] integrate(1/cos(b*x+a)**(3/2),x)

[Out] Integral(cos(a + b*x)**(-3/2), x)

Maxima [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{3}{2}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-3/2), x)

Giac [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx = \int \frac{1}{\cos(bx + a)^{\frac{3}{2}}} dx$$

[In] integrate(1/cos(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-3/2), x)

Mupad [B] (verification not implemented)

Time = 14.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.11

$$\int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx = \frac{2 \sin(a + bx) {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{4}; \cos(a + bx)^2\right)}{b \sqrt{\cos(a + bx)} \sqrt{\sin(a + bx)^2}}$$

[In] int(1/cos(a + b*x)^(3/2),x)

[Out] (2*sin(a + b*x)*hypergeom([-1/4, 1/2], 3/4, cos(a + b*x)^2))/(b*cos(a + b*x)^(1/2)*(sin(a + b*x)^2)^(1/2))

3.15 $\int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx$

Optimal result	160
Rubi [A] (verified)	160
Mathematica [A] (verified)	161
Maple [B] (verified)	161
Fricas [C] (verification not implemented)	162
Sympy [F]	162
Maxima [F]	162
Giac [F]	163
Mupad [B] (verification not implemented)	163

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b} + \frac{2 \sin(a+bx)}{3b \cos^{\frac{3}{2}}(a+bx)}$$

[Out] $2/3*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b+2/3*\sin(b*x+a)/b/\cos(b*x+a)^{(3/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2720}

$$\int \frac{1}{\cos^{\frac{5}{2}}(a+bx)} dx = \frac{2 \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b} + \frac{2 \sin(a+bx)}{3b \cos^{\frac{3}{2}}(a+bx)}$$

[In] `Int[Cos[a + b*x]^(-5/2), x]`

[Out] $(2*\operatorname{EllipticF}[(a + b*x)/2, 2])/(3*b) + (2*\sin[a + b*x])/(3*b*\cos[a + b*x]^{(3/2)})$

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720


```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
]*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(a + bx)}{3b \cos^{\frac{3}{2}}(a + bx)} + \frac{1}{3} \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{2 \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b} + \frac{2 \sin(a + bx)}{3b \cos^{\frac{3}{2}}(a + bx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.86

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx = \frac{2 \left(\text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + \frac{\sin(a + bx)}{\cos^{\frac{3}{2}}(a + bx)} \right)}{3b}$$

```
[In] Integrate[Cos[a + b*x]^(-5/2), x]
```

```
[Out] (2*(EllipticF[(a + b*x)/2, 2] + Sin[a + b*x]/Cos[a + b*x]^(3/2)))/(3*b)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(62) = 124.

Time = 1.46 (sec) , antiderivative size = 213, normalized size of antiderivative = 5.07

method	result
default	$-\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right) \right) - 1} F\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right)}{3 \sqrt{-2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)} \left(-1 + 2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}$

```
[In] int(1/cos(b*x+a)^(5/2), x, method=_RETURNVERBOSE)
```

```
[Out] -2/3*(-2*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*sin(1/2*b*x+1/2*a)^2-2*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2)))*((-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/(-1+2*cos(1/2*b*x+1/2*a)^2)^(3/2)/sin(1/2*b*x+1/2*a)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.19

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx$$

$$= \frac{-i\sqrt{2}\cos(bx + a)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i\sin(bx + a)) + i\sqrt{2}\cos(bx + a)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i\sin(bx + a)) + 2\sqrt{2}\cos(bx + a)\sin(bx + a)}{3b\cos(bx + a)^2}$$

[In] integrate(1/cos(b*x+a)^(5/2),x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*sqrt(2)*cos(b*x + a)*sin(b*x + a))/(b*cos(b*x + a)^2)

Sympy [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx$$

[In] integrate(1/cos(b*x+a)**(5/2),x)

[Out] Integral(cos(a + b*x)**(-5/2), x)

Maxima [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{5}{2}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(5/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-5/2), x)

Giac [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx = \int \frac{1}{\cos(bx + a)^{\frac{5}{2}}} dx$$

[In] integrate(1/cos(b*x+a)^(5/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-5/2), x)

Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos^{\frac{5}{2}}(a + bx)} dx = \frac{2 \sin(a + bx) {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{1}{4}; \cos(a + bx)^2\right)}{3 b \cos(a + bx)^{3/2} \sqrt{\sin(a + bx)^2}}$$

[In] int(1/cos(a + b*x)^(5/2),x)

[Out] (2*sin(a + b*x)*hypergeom([-3/4, 1/2], 1/4, cos(a + b*x)^2))/(3*b*cos(a + b*x)^(3/2)*(sin(a + b*x)^2)^(1/2))

3.16 $\int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx$

Optimal result	164
Rubi [A] (verified)	164
Mathematica [A] (verified)	165
Maple [B] (verified)	165
Fricas [C] (verification not implemented)	166
Sympy [F(-1)]	166
Maxima [F]	167
Giac [F]	167
Mupad [B] (verification not implemented)	167

Optimal result

Integrand size = 10, antiderivative size = 65

$$\int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx = -\frac{6E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5b} + \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}}$$

[Out] $-6/5*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})/b+2/5*\sin(b*x+a)/b/\cos(b*x+a)^{(5/2)}+6/5*\sin(b*x+a)/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2719}

$$\int \frac{1}{\cos^{\frac{7}{2}}(a+bx)} dx = -\frac{6E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5b} + \frac{2\sin(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} + \frac{6\sin(a+bx)}{5b\sqrt{\cos(a+bx)}}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^{(-7/2)}, x]$

[Out] $(-6*\text{EllipticE}[(a + b*x)/2, 2])/(5*b) + (2*\text{Sin}[a + b*x])/(5*b*\text{Cos}[a + b*x]^{(5/2)}) + (6*\text{Sin}[a + b*x])/(5*b*\text{Sqrt}[\text{Cos}[a + b*x]])$

Rule 2716

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n+1)}/(b*d*(n+1))), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(a + bx)}{5b \cos^{\frac{5}{2}}(a + bx)} + \frac{3}{5} \int \frac{1}{\cos^{\frac{3}{2}}(a + bx)} dx \\ &= \frac{2 \sin(a + bx)}{5b \cos^{\frac{5}{2}}(a + bx)} + \frac{6 \sin(a + bx)}{5b \sqrt{\cos(a + bx)}} - \frac{3}{5} \int \sqrt{\cos(a + bx)} dx \\ &= -\frac{6E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b} + \frac{2 \sin(a + bx)}{5b \cos^{\frac{5}{2}}(a + bx)} + \frac{6 \sin(a + bx)}{5b \sqrt{\cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.91

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \frac{-6 \cos^{\frac{3}{2}}(a + bx) E\left(\frac{1}{2}(a + bx) \mid 2\right) + 3 \sin(2(a + bx)) + 2 \tan(a + bx)}{5b \cos^{\frac{3}{2}}(a + bx)}$$

[In] `Integrate[Cos[a + b*x]^(-7/2), x]`

[Out] `(-6*Cos[a + b*x]^(3/2)*EllipticE[(a + b*x)/2, 2] + 3*Sin[2*(a + b*x)] + 2*Tan[a + b*x])/(5*b*Cos[a + b*x]^(3/2))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 357 vs. 2(81) = 162.

Time = 2.31 (sec) , antiderivative size = 358, normalized size of antiderivative = 5.51

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{bx}{2} + \frac{a}{2})) + 1)(\sin^2(\frac{bx}{2} + \frac{a}{2}))}}{(24 \cos(\frac{bx}{2} + \frac{a}{2})(\sin^6(\frac{bx}{2} + \frac{a}{2})) - 12\sqrt{2(\sin^2(\frac{bx}{2} + \frac{a}{2}))} - 1) E(\cos(\frac{bx}{2} + \frac{a}{2}), \sqrt{2})} \sqrt{2}$

[In] `int(1/cos(b*x+a)^(7/2), x, method=_RETURNVERBOSE)`

[Out] `-2/5*(-(-2*cos(1/2*b*x+1/2*a)^2+1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)/sin(1/2*b*x+1/2*a)^3*(24*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^6-12*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))*(sin(1/2*b*x+1/2*a)^2)^(1/2)`

$$\frac{1}{2} \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 - 24 \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 \cos\left(\frac{1}{2}bx + \frac{1}{2}a\right) + 12 \left(2 \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - 1\right)^{\frac{1}{2}} \text{EllipticE}\left(\cos\left(\frac{1}{2}bx + \frac{1}{2}a\right), 2^{\frac{1}{2}}\right) \left(\sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2\right)^{\frac{1}{2}} \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 + 8 \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 \cos\left(\frac{1}{2}bx + \frac{1}{2}a\right) - 3 \left(\sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2\right)^{\frac{1}{2}} \left(2 \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - 1\right)^{\frac{1}{2}} \text{EllipticE}\left(\cos\left(\frac{1}{2}bx + \frac{1}{2}a\right), 2^{\frac{1}{2}}\right) \left(-2 \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + \sin\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2\right)^{\frac{1}{2}} / \left(-1 + 2 \cos\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2\right)^{\frac{1}{2}} / b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.69

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \frac{-3i \sqrt{2} \cos(bx + a)^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) + 3i}{\dots}$$

[In] integrate(1/cos(b*x+a)^(7/2),x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*cos(b*x + a)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + 3*I*sqrt(2)*cos(b*x + a)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*(3*cos(b*x + a)^2 + 1)*sqrt(cos(b*x + a))*sin(b*x + a))/(b*cos(b*x + a)^3)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \text{Timed out}$$

[In] integrate(1/cos(b*x+a)**(7/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \int \frac{1}{\cos(bx + a)^{\frac{7}{2}}} dx$$

[In] integrate(1/cos(b*x+a)^(7/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-7/2), x)

Giac [F]

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \int \frac{1}{\cos(bx + a)^{\frac{7}{2}}} dx$$

[In] integrate(1/cos(b*x+a)^(7/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-7/2), x)

Mupad [B] (verification not implemented)

Time = 14.85 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.65

$$\int \frac{1}{\cos^{\frac{7}{2}}(a + bx)} dx = \frac{2 \sin(a + bx) {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; -\frac{1}{4}; \cos(a + bx)^2\right)}{5 b \cos(a + bx)^{5/2} \sqrt{\sin(a + bx)^2}}$$

[In] int(1/cos(a + b*x)^(7/2),x)

[Out] (2*sin(a + b*x)*hypergeom([-5/4, 1/2], -1/4, cos(a + b*x)^2))/(5*b*cos(a + b*x)^(5/2)*(sin(a + b*x)^2)^(1/2))

3.17 $\int (c \cos(a + bx))^{7/2} dx$

Optimal result	168
Rubi [A] (verified)	168
Mathematica [A] (verified)	170
Maple [A] (verified)	170
Fricas [C] (verification not implemented)	170
Sympy [F(-1)]	171
Maxima [F]	171
Giac [F]	171
Mupad [F(-1)]	171

Optimal result

Integrand size = 12, antiderivative size = 98

$$\int (c \cos(a + bx))^{7/2} dx = \frac{10c^4 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b \sqrt{c \cos(a + bx)}} + \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b}$$

[Out] $2/7*c*(c*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b+10/21*c^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}+10/21*c^3*\sin(b*x+a)*(c*\cos(b*x+a))^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\int (c \cos(a + bx))^{7/2} dx = \frac{10c^4 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b \sqrt{c \cos(a + bx)}} + \frac{10c^3 \sin(a + bx) \sqrt{c \cos(a + bx)}}{21b} + \frac{2c \sin(a + bx) (c \cos(a + bx))^{5/2}}{7b}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(7/2)}, x]$

[Out] $(10*c^4*\operatorname{Sqrt}[\operatorname{Cos}[a + b*x]]*\operatorname{EllipticF}[(a + b*x)/2, 2])/(21*b*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]]) + (10*c^3*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]]*\operatorname{Sin}[a + b*x])/(21*b) + (2*c*(c*\operatorname{Cos}[a + b*x])^{(5/2)}*\operatorname{Sin}[a + b*x])/(7*b)$

Rule 2715


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} + \frac{1}{7}(5c^2) \int (c \cos(a + bx))^{3/2} dx \\
 &= \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} \\
 &\quad + \frac{1}{21}(5c^4) \int \frac{1}{\sqrt{c \cos(a + bx)}} dx \\
 &= \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b} \\
 &\quad + \frac{(5c^4 \sqrt{\cos(a + bx)}) \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{21 \sqrt{c \cos(a + bx)}} \\
 &= \frac{10c^4 \sqrt{\cos(a + bx)} \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{21b \sqrt{c \cos(a + bx)}} \\
 &\quad + \frac{10c^3 \sqrt{c \cos(a + bx)} \sin(a + bx)}{21b} + \frac{2c(c \cos(a + bx))^{5/2} \sin(a + bx)}{7b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.78

$$\int (c \cos(a + bx))^{7/2} dx = \frac{c^3 \sqrt{c \cos(a + bx)} \left(20 \operatorname{EllipticF} \left(\frac{1}{2}(a + bx), 2 \right) + \sqrt{\cos(a + bx)} (23 \sin(a + bx) + 3 \sin(3(a + bx))) \right)}{42b \sqrt{\cos(a + bx)}}$$

[In] Integrate[(c*Cos[a + b*x])^(7/2),x]

[Out] (c^3*Sqrt[c*Cos[a + b*x]]*(20*EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*(23*Sin[a + b*x] + 3*Sin[3*(a + b*x)])))/(42*b*Sqrt[Cos[a + b*x]])

Maple [A] (verified)

Time = 4.01 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.14

method	result
default	$-\frac{2\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}(\sin^2(\frac{bx}{2}+\frac{a}{2}))c^4(48(\cos^9(\frac{bx}{2}+\frac{a}{2}))-120(\cos^7(\frac{bx}{2}+\frac{a}{2}))+128(\cos^5(\frac{bx}{2}+\frac{a}{2}))-72(\cos^3(\frac{bx}{2}+\frac{a}{2})))}{21\sqrt{-c(2(\sin^4(\frac{bx}{2}+\frac{a}{2}))-(\sin^2(\frac{bx}{2}+\frac{a}{2})))}\sin(\frac{bx}{2}+\frac{a}{2})\sqrt{c(-1+2\cos^2(\frac{bx}{2}+\frac{a}{2}))}}}$

[In] int((c*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)

[Out] -2/21*(c*(-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c^4*(48*cos(1/2*b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(-1+2*cos(1/2*b*x+1/2*a)^2))^(1/2)/b

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.97

$$\int (c \cos(a + bx))^{7/2} dx = \frac{-5i \sqrt{2c^7} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + 5i \sqrt{2c^7} \operatorname{weierstrassPInverse}(\dots)}{2}$$

[In] integrate((c*cos(b*x+a))^(7/2),x, algorithm="fricas")

```
[Out] 1/21*(-5*I*sqrt(2)*c^(7/2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(
b*x + a)) + 5*I*sqrt(2)*c^(7/2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I
*sin(b*x + a)) + 2*(3*c^3*cos(b*x + a)^2 + 5*c^3)*sqrt(c*cos(b*x + a))*sin(
b*x + a))/b
```

Sympy [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{7/2} dx = \text{Timed out}$$

```
[In] integrate((c*cos(b*x+a))**(7/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int (c \cos(a + bx))^{7/2} dx = \int (c \cos(bx + a))^{\frac{7}{2}} dx$$

```
[In] integrate((c*cos(b*x+a))^(7/2),x, algorithm="maxima")
```

```
[Out] integrate((c*cos(b*x + a))^(7/2), x)
```

Giac [F]

$$\int (c \cos(a + bx))^{7/2} dx = \int (c \cos(bx + a))^{\frac{7}{2}} dx$$

```
[In] integrate((c*cos(b*x+a))^(7/2),x, algorithm="giac")
```

```
[Out] integrate((c*cos(b*x + a))^(7/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{7/2} dx = \int (c \cos(a + bx))^{7/2} dx$$

```
[In] int((c*cos(a + b*x))^(7/2),x)
```

```
[Out] int((c*cos(a + b*x))^(7/2), x)
```

3.18 $\int (c \cos(a + bx))^{5/2} dx$

Optimal result	172
Rubi [A] (verified)	172
Mathematica [A] (verified)	173
Maple [B] (verified)	173
Fricas [C] (verification not implemented)	174
Sympy [F(-1)]	174
Maxima [F]	174
Giac [F]	175
Mupad [F(-1)]	175

Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (c \cos(a + bx))^{5/2} dx = \frac{6c^2 \sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b \sqrt{\cos(a + bx)}} + \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b}$$

[Out] $2/5*c*(c*\cos(b*x+a))^(3/2)*\sin(b*x+a)/b+6/5*c^2*(\cos(1/2*a+1/2*b*x)^2)^(1/2)/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^(1/2))*(c*\cos(b*x+a))^(1/2)/b/\cos(b*x+a)^(1/2)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2719}

$$\int (c \cos(a + bx))^{5/2} dx = \frac{6c^2 E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{c \cos(a + bx)}}{5b \sqrt{\cos(a + bx)}} + \frac{2c \sin(a + bx) (c \cos(a + bx))^{3/2}}{5b}$$

[In] $\text{Int}[(c*\text{Cos}[a + b*x])^(5/2), x]$

[Out] $(6*c^2*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*c*(c*\text{Cos}[a + b*x])^(3/2)*\text{Sin}[a + b*x])/(5*b)$

Rule 2715

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^(n-1)/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^(n-2), x], x] /; \text{FreeQ}\{b, c, d, x\} \&\amp; \text{GtQ}[n, 1] \&\amp; \text{IntegerQ}[2*n]$

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} + \frac{1}{5}(3c^2) \int \sqrt{c \cos(a + bx)} dx \\ &= \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} + \frac{(3c^2 \sqrt{c \cos(a + bx)}) \int \sqrt{\cos(a + bx)} dx}{5\sqrt{\cos(a + bx)}} \\ &= \frac{6c^2 \sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b\sqrt{\cos(a + bx)}} + \frac{2c(c \cos(a + bx))^{3/2} \sin(a + bx)}{5b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.89

$$\int (c \cos(a + bx))^{5/2} dx = \frac{(c \cos(a + bx))^{5/2} \left(6E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sqrt{\cos(a + bx)} \sin(2(a + bx)) \right)}{5b \cos^{5/2}(a + bx)}$$

```
[In] Integrate[(c*Cos[a + b*x])^(5/2),x]
```

```
[Out] ((c*Cos[a + b*x])^(5/2)*(6*EllipticE[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*S
in[2*(a + b*x)]))/(5*b*Cos[a + b*x]^(5/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(86) = 172.

Time = 2.78 (sec) , antiderivative size = 213, normalized size of antiderivative = 3.04

method	result
default	$\frac{2\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}(\sin^2(\frac{bx}{2}+\frac{a}{2}))c^3(-8\cos(\frac{bx}{2}+\frac{a}{2})(\sin^6(\frac{bx}{2}+\frac{a}{2}))+8(\sin^4(\frac{bx}{2}+\frac{a}{2}))\cos(\frac{bx}{2}+\frac{a}{2}))-2(\sin^2(\frac{bx}{2}+\frac{a}{2}))}{5\sqrt{-c(2(\sin^4(\frac{bx}{2}+\frac{a}{2}))-(\sin^2(\frac{bx}{2}+\frac{a}{2})))}\sin(\frac{bx}{2}+\frac{a}{2})\sqrt{c(-1+2$

[In] `int((c*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*(c*(-1+2*\cos(1/2*b*x+1/2*a)^2)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*c^3*(-8*\cos(1/2*b*x+1/2*a)*\sin(1/2*b*x+1/2*a)^6+8*\sin(1/2*b*x+1/2*a)^4*\cos(1/2*b*x+1/2*a)-2*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a)-3*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)}))/(-c*(2*\sin(1/2*b*x+1/2*a)^4-\sin(1/2*b*x+1/2*a)^2))^{(1/2)}/\sin(1/2*b*x+1/2*a)/(c*(-1+2*\cos(1/2*b*x+1/2*a)^2))^{(1/2)}/b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int (c \cos(a + bx))^{5/2} dx = \frac{2 \sqrt{c \cos(bx + a)} c^2 \cos(bx + a) \sin(bx + a) + 3i \sqrt{2} c^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + I \sin(bx + a))) - 3i \sqrt{2} c^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - I \sin(bx + a)))}{b}$$

[In] `integrate((c*cos(b*x+a))^(5/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{c*\cos(b*x + a)}*c^2*\cos(b*x + a)*\sin(b*x + a) + 3*I*\sqrt{2}*c^{5/2}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a))) - 3*I*\sqrt{2}*c^{5/2}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a))))/b$$

Sympy [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{5/2} dx = \text{Timed out}$$

[In] `integrate((c*cos(b*x+a))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int (c \cos(a + bx))^{5/2} dx = \int (c \cos(bx + a))^{5/2} dx$$

[In] `integrate((c*cos(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] `integrate((c*cos(b*x + a))^(5/2), x)`

Giac [F]

$$\int (c \cos(a + bx))^{5/2} dx = \int (c \cos(bx + a))^{5/2} dx$$

[In] integrate((c*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{5/2} dx = \int (c \cos(a + bx))^{5/2} dx$$

[In] int((c*cos(a + b*x))^(5/2),x)

[Out] int((c*cos(a + b*x))^(5/2), x)

3.19 $\int (c \cos(a + bx))^{3/2} dx$

Optimal result	176
Rubi [A] (verified)	176
Mathematica [A] (verified)	177
Maple [B] (verified)	178
Fricas [C] (verification not implemented)	178
Sympy [F]	178
Maxima [F]	179
Giac [F]	179
Mupad [F(-1)]	179

Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (c \cos(a + bx))^{3/2} dx = \frac{2c^2 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b \sqrt{c \cos(a + bx)}} + \frac{2c \sqrt{c \cos(a + bx)} \sin(a + bx)}{3b}$$

[Out] $2/3*c^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}+2/3*c*\sin(b*x+a)*(c*\cos(b*x+a))^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\int (c \cos(a + bx))^{3/2} dx = \frac{2c^2 \sqrt{\cos(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3b \sqrt{c \cos(a + bx)}} + \frac{2c \sin(a + bx) \sqrt{c \cos(a + bx)}}{3b}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c^2*\operatorname{Sqrt}[\operatorname{Cos}[a + b*x]]*\operatorname{EllipticF}[(a + b*x)/2, 2])/(3*b*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]]) + (2*c*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]]*\operatorname{Sin}[a + b*x])/(3*b)$

Rule 2715


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2c\sqrt{c\cos(a+bx)}\sin(a+bx)}{3b} + \frac{1}{3}c^2 \int \frac{1}{\sqrt{c\cos(a+bx)}} dx \\ &= \frac{2c\sqrt{c\cos(a+bx)}\sin(a+bx)}{3b} + \frac{\left(c^2\sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{3\sqrt{c\cos(a+bx)}} \\ &= \frac{2c^2\sqrt{\cos(a+bx)}\text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3b\sqrt{c\cos(a+bx)}} + \frac{2c\sqrt{c\cos(a+bx)}\sin(a+bx)}{3b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.83

$$\int (c\cos(a+bx))^{3/2} dx = \frac{2(c\cos(a+bx))^{3/2} \left(\text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right) + \sqrt{\cos(a+bx)}\sin(a+bx) \right)}{3b\cos^{3/2}(a+bx)}$$

```
[In] Integrate[(c*cos[a + b*x])^(3/2),x]
```

```
[Out] (2*(c*cos[a + b*x])^(3/2)*(EllipticF[(a + b*x)/2, 2] + Sqrt[Cos[a + b*x]]*Sin[a + b*x]))/(3*b*cos[a + b*x]^(3/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(86) = 172$.

Time = 2.32 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.71

method	result
default	$-\frac{2\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}(\sin^2(\frac{bx}{2}+\frac{a}{2}))c^2(4(\sin^4(\frac{bx}{2}+\frac{a}{2}))\cos(\frac{bx}{2}+\frac{a}{2})-2(\sin^2(\frac{bx}{2}+\frac{a}{2}))\cos(\frac{bx}{2}+\frac{a}{2})+\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}})}{3\sqrt{-c(2(\sin^4(\frac{bx}{2}+\frac{a}{2}))-(\sin^2(\frac{bx}{2}+\frac{a}{2})))}\sin(\frac{bx}{2}+\frac{a}{2})\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}b}$

[In] `int((c*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/3*(c*(-1+2*\cos(1/2*b*x+1/2*a)^2)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*c^2*(4*\sin(1/2*b*x+1/2*a)^4*\cos(1/2*b*x+1/2*a)-2*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a)+(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*EllipticF(\cos(1/2*b*x+1/2*a),2^{(1/2)}))/(-c*(2*\sin(1/2*b*x+1/2*a)^4-\sin(1/2*b*x+1/2*a)^2))^{(1/2)}/\sin(1/2*b*x+1/2*a)/(c*(-1+2*\cos(1/2*b*x+1/2*a)^2))^{(1/2)}/b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.10

$$\int (c \cos(a + bx))^{3/2} dx = \frac{-i \sqrt{2} c^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} c^{3/2} \text{weierstrassPInverse}(\cos(bx + a) - i \sin(bx + a), -4, 0)}{3b}$$

[In] `integrate((c*cos(b*x+a))^(3/2),x, algorithm="fricas")`

[Out]
$$1/3*(-I*\sqrt{2}*c^{(3/2)}*\text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a)) + I*\sqrt{2}*c^{(3/2)}*\text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a)) + 2*\sqrt{c*\cos(b*x + a)}*c*\sin(b*x + a))/b$$

Sympy [F]

$$\int (c \cos(a + bx))^{3/2} dx = \int (c \cos(a + bx))^{\frac{3}{2}} dx$$

[In] `integrate((c*cos(b*x+a))**(3/2),x)`

[Out] `Integral((c*cos(a + b*x))**(3/2), x)`

Maxima [F]

$$\int (c \cos(a + bx))^{3/2} dx = \int (c \cos(bx + a))^{\frac{3}{2}} dx$$

[In] integrate((c*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

Giac [F]

$$\int (c \cos(a + bx))^{3/2} dx = \int (c \cos(bx + a))^{\frac{3}{2}} dx$$

[In] integrate((c*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{3/2} dx = \int (c \cos(a + bx))^{\frac{3}{2}} dx$$

[In] int((c*cos(a + b*x))^(3/2),x)

[Out] int((c*cos(a + b*x))^(3/2), x)

3.20 $\int \sqrt{c \cos(a + bx)} dx$

Optimal result	180
Rubi [A] (verified)	180
Mathematica [A] (verified)	181
Maple [B] (verified)	181
Fricas [C] (verification not implemented)	182
Sympy [F]	182
Maxima [F]	182
Giac [F]	182
Mupad [F(-1)]	183

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \sqrt{c \cos(a + bx)} dx = \frac{2\sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}}$$

[Out] $2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2721, 2719}

$$\int \sqrt{c \cos(a + bx)} dx = \frac{2E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{c \cos(a + bx)}}{b\sqrt{\cos(a + bx)}}$$

[In] Int[Sqrt[c*Cos[a + b*x]],x]

[Out] (2*Sqrt[c*Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]])

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ

[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{c \cos(a + bx)} \int \sqrt{\cos(a + bx)} dx}{\sqrt{\cos(a + bx)}} \\ &= \frac{2\sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \sqrt{c \cos(a + bx)} dx = \frac{2\sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}}$$

[In] Integrate[Sqrt[c*Cos[a + b*x]],x]

[Out] (2*Sqrt[c*Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(60) = 120.

Time = 1.92 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.74

method	result
default	$\frac{2\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}(\sin^2(\frac{bx}{2}+\frac{a}{2}))c\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\sqrt{-2(\cos^2(\frac{bx}{2}+\frac{a}{2}))+1}E(\cos(\frac{bx}{2}+\frac{a}{2}),\sqrt{2})}{\sqrt{-c(2(\sin^4(\frac{bx}{2}+\frac{a}{2}))-(\sin^2(\frac{bx}{2}+\frac{a}{2})))}\sin(\frac{bx}{2}+\frac{a}{2})\sqrt{c(-1+2(\cos^2(\frac{bx}{2}+\frac{a}{2})))}b}$
risch	$-\frac{i\sqrt{2}\sqrt{c(e^{2i(bx+a)}+1)e^{-i(bx+a)}}}{b} - i\left(\frac{2(e^{2i(bx+a)}c+c)}{c\sqrt{e^{i(bx+a)}}(e^{2i(bx+a)}c+c)} + \frac{i\sqrt{-i(e^{i(bx+a)}+i)}\sqrt{2}\sqrt{i(e^{i(bx+a)}-i)}\sqrt{ie^{i(bx+a)}}(-2iE(\sqrt{-\frac{2iE(\cos(\frac{bx}{2}+\frac{a}{2}),\sqrt{2})}{\sqrt{e^{3i(bx+a)}}c+e^{i(bx+a)}}))}}{c\sqrt{e^{i(bx+a)}}(e^{2i(bx+a)}c+c)}\right)$

[In] int((c*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] 2*(c*(-1+2*cos(1/2*b*x+1/2*a)^2)*sin(1/2*b*x+1/2*a)^2)^(1/2)*c*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(c*(-1+2*cos(1/2*b*x+1/2*a)^2)^(1/2))/b

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.66

$$\int \sqrt{c \cos(a + bx)} dx$$

$$= \frac{i \sqrt{2} \sqrt{c} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) - i \sqrt{2} \sqrt{c} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)))}{b}$$

```
[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*sqrt(c)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))))/b
```

Sympy [F]

$$\int \sqrt{c \cos(a + bx)} dx = \int \sqrt{c \cos(a + bx)} dx$$

```
[In] integrate((c*cos(b*x+a))**(1/2),x)
```

```
[Out] Integral(sqrt(c*cos(a + b*x)), x)
```

Maxima [F]

$$\int \sqrt{c \cos(a + bx)} dx = \int \sqrt{c \cos(bx + a)} dx$$

```
[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*cos(b*x + a)), x)
```

Giac [F]

$$\int \sqrt{c \cos(a + bx)} dx = \int \sqrt{c \cos(bx + a)} dx$$

```
[In] integrate((c*cos(b*x+a))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(c*cos(b*x + a)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{c \cos(a + bx)} dx = \int \sqrt{c \cos(a + bx)} dx$$

```
[In] int((c*cos(a + b*x))^(1/2),x)
```

```
[Out] int((c*cos(a + b*x))^(1/2), x)
```

3.21 $\int \frac{1}{\sqrt{c \cos(a+bx)}} dx$

Optimal result	184
Rubi [A] (verified)	184
Mathematica [A] (verified)	185
Maple [C] (verified)	185
Fricas [C] (verification not implemented)	186
Sympy [F]	186
Maxima [F]	186
Giac [F]	187
Mupad [B] (verification not implemented)	187

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{1}{\sqrt{c \cos(a+bx)}} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b\sqrt{c \cos(a+bx)}}$$

[Out] $2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(c*\cos(b*x+a))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2721, 2720}

$$\int \frac{1}{\sqrt{c \cos(a+bx)}} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b\sqrt{c \cos(a+bx)}}$$

[In] `Int[1/Sqrt[c*Cos[a + b*x]],x]`

[Out] `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[c*Cos[a + b*x]])`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ`

`[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{\sqrt{c \cos(a+bx)}} \\ &= \frac{2\sqrt{\cos(a+bx)} \text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b\sqrt{c \cos(a+bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{c \cos(a+bx)}} dx = \frac{2\sqrt{\cos(a+bx)} \text{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{b\sqrt{c \cos(a+bx)}}$$

[In] `Integrate[1/Sqrt[c*Cos[a + b*x]],x]`

[Out] `(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[c*Cos[a + b*x]])`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.39 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

method	result	size
default	$\frac{2\sqrt{-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)} \text{am}^{-1}\left(\frac{bx}{2}+\frac{a}{2} \sqrt{2}\right)}{b\sqrt{c\left(-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)}}$	54

[In] `int(1/(c*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

[Out] `2/b/(c*(-1+2*cos(1/2*b*x+1/2*a)^2))^(1/2)*(-1+2*cos(1/2*b*x+1/2*a)^2)^(1/2)
*InverseJacobiAM(1/2*b*x+1/2*a,2^(1/2))`

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.58

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx$$

$$= \frac{-i \sqrt{2} \sqrt{c} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} \sqrt{c} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))}{bc}$$

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*sqrt(c)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/(b*c)

Sympy [F]

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos(a + bx)}} dx$$

[In] integrate(1/(c*cos(b*x+a))**(1/2),x)

[Out] Integral(1/sqrt(c*cos(a + b*x)), x)

Maxima [F]

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos(bx + a)}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(c*cos(b*x + a)), x)

Giac [F]

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos(bx + a)}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(c*cos(b*x + a)), x)

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{c \cos(a + bx)}} dx = \frac{2 \sqrt{\cos(a + bx)} F\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b \sqrt{c \cos(a + bx)}}$$

[In] int(1/(c*cos(a + b*x))^(1/2),x)

[Out] (2*cos(a + b*x)^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/(b*(c*cos(a + b*x))^(1/2))

3.22 $\int \frac{1}{(c \cos(a+bx))^{3/2}} dx$

Optimal result	188
Rubi [A] (verified)	188
Mathematica [A] (verified)	189
Maple [B] (verified)	189
Fricas [C] (verification not implemented)	190
Sympy [F]	190
Maxima [F]	190
Giac [F]	191
Mupad [F(-1)]	191

Optimal result

Integrand size = 12, antiderivative size = 68

$$\int \frac{1}{(c \cos(a+bx))^{3/2}} dx = -\frac{2\sqrt{c \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{bc^2 \sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{bc \sqrt{c \cos(a+bx)}}$$

[Out] 2*sin(b*x+a)/b/c/(c*cos(b*x+a))^(1/2)-2*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*(c*cos(b*x+a))^(1/2)/b/c^2/cos(b*x+a)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$\int \frac{1}{(c \cos(a+bx))^{3/2}} dx = \frac{2 \sin(a+bx)}{bc \sqrt{c \cos(a+bx)}} - \frac{2E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{c \cos(a+bx)}}{bc^2 \sqrt{\cos(a+bx)}}$$

[In] Int[(c*cos[a + b*x])^(-3/2), x]

[Out] (-2*Sqrt[c*cos[a + b*x]]*EllipticE[(a + b*x)/2, 2])/(b*c^2*Sqrt[Cos[a + b*x]]) + (2*Sin[a + b*x])/(b*c*Sqrt[c*cos[a + b*x]])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(a + bx)}{bc\sqrt{c \cos(a + bx)}} - \frac{\int \sqrt{c \cos(a + bx)} dx}{c^2} \\ &= \frac{2 \sin(a + bx)}{bc\sqrt{c \cos(a + bx)}} - \frac{\sqrt{c \cos(a + bx)} \int \sqrt{\cos(a + bx)} dx}{c^2 \sqrt{\cos(a + bx)}} \\ &= -\frac{2\sqrt{c \cos(a + bx)}E\left(\frac{1}{2}(a + bx) \mid 2\right)}{bc^2 \sqrt{\cos(a + bx)}} + \frac{2 \sin(a + bx)}{bc\sqrt{c \cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \frac{2\left(-\sqrt{\cos(a + bx)}E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sin(a + bx)\right)}{bc\sqrt{c \cos(a + bx)}}$$

[In] Integrate[(c*Cos[a + b*x])^(-3/2),x]

[Out] (2*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/(b*c*Sqrt[c*Cos[a + b*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(88) = 176.

Time = 1.45 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.91

method	result
default	$-\frac{2\left(-2\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)c + \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)c\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} - 1\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)}{c\sqrt{-c\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{c\left(-1 + 2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}}$

```
[In] int(1/(c*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/c*(-2*cos(1/2*b*x+1/2*a)*(-2*sin(1/2*b*x+1/2*a)^4*c+sin(1/2*b*x+1/2*a)^2*c)^(1/2)*sin(1/2*b*x+1/2*a)^2+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^4*c+sin(1/2*b*x+1/2*a)^2*c)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2)))/(-c*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(c*(-1+2*cos(1/2*b*x+1/2*a)^2))^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.53

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \frac{-i \sqrt{2} \sqrt{c} \cos(bx + a) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a)))}{(c \cos(a + bx))^{3/2}}$$

```
[In] integrate(1/(c*cos(b*x+a))^(3/2),x, algorithm="fricas")
```

```
[Out] (-I*sqrt(2)*sqrt(c)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*sqrt(c)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*sqrt(c*cos(b*x + a))*sin(b*x + a))/(b*c^2*cos(b*x + a))
```

Sympy [F]

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(a + bx))^{3/2}} dx$$

```
[In] integrate(1/(c*cos(b*x+a))**(3/2),x)
```

```
[Out] Integral((c*cos(a + b*x))**(-3/2), x)
```

Maxima [F]

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(bx + a))^{3/2}} dx$$

```
[In] integrate(1/(c*cos(b*x+a))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((c*cos(b*x + a))^(3/2), x)
```

Giac [F]

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{3}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(a + bx))^{3/2}} dx$$

[In] int(1/(c*cos(a + b*x))^(3/2),x)

[Out] int(1/(c*cos(a + b*x))^(3/2), x)

3.23 $\int \frac{1}{(c \cos(a+bx))^{5/2}} dx$

Optimal result	192
Rubi [A] (verified)	192
Mathematica [A] (verified)	193
Maple [B] (verified)	193
Fricas [C] (verification not implemented)	194
Sympy [F]	194
Maxima [F]	195
Giac [F]	195
Mupad [F(-1)]	195

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(c \cos(a+bx))^{5/2}} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3bc^2\sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}}$$

[Out] $2/3*\sin(b*x+a)/b/c/(c*\cos(b*x+a))^{(3/2)}+2/3*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\operatorname{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/c^{(1/2)}/(c*\cos(b*x+a))^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2720}

$$\int \frac{1}{(c \cos(a+bx))^{5/2}} dx = \frac{2\sqrt{\cos(a+bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a+bx), 2\right)}{3bc^2\sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bc(c \cos(a+bx))^{3/2}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(-5/2)}, x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[a + b*x]]*\operatorname{EllipticF}[(a + b*x)/2, 2])/(3*b*c^2*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]]) + (2*\operatorname{Sin}[a + b*x])/(3*b*c*(c*\operatorname{Cos}[a + b*x])^{(3/2)})$

Rule 2716

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)}/(b*d*(n+1))), x] + \operatorname{Dist}[(n+2)/(b^2*(n+1)), \operatorname{Int}[(b*\operatorname{Sin}[c + d*x])^{(n+2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
]*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(a + bx)}{3bc(c \cos(a + bx))^{3/2}} + \frac{\int \frac{1}{\sqrt{c \cos(a + bx)}} dx}{3c^2} \\ &= \frac{2 \sin(a + bx)}{3bc(c \cos(a + bx))^{3/2}} + \frac{\sqrt{\cos(a + bx)} \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{3c^2 \sqrt{c \cos(a + bx)}} \\ &= \frac{2\sqrt{\cos(a + bx)} \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right)}{3bc^2 \sqrt{c \cos(a + bx)}} + \frac{2 \sin(a + bx)}{3bc(c \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \frac{2\left(\sqrt{\cos(a + bx)} \text{EllipticF}\left(\frac{1}{2}(a + bx), 2\right) + \tan(a + bx)\right)}{3bc^2 \sqrt{c \cos(a + bx)}}$$

```
[In] Integrate[(c*Cos[a + b*x])^(-5/2),x]
```

```
[Out] (2*(Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Tan[a + b*x]))/(3*b*c^2*
Sqrt[c*Cos[a + b*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(88) = 176.

Time = 1.65 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.35

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}-1F\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\cos\left(\frac{bx}{2}+\frac{a}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(bx+a)}{2}}\right)}{3c^2\sqrt{-c\left(2\left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)\left(-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)}\sin\left(\frac{bx}{2}+\frac{a}{2}\right)}$

[In] `int(1/(c*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{2}{3}\left(-2\left(\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2\right)^{1/2}\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2-1\right)^{1/2}\text{EllipticF}\left(\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right),2^{1/2}\right)\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)+\left(\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2\right)^{1/2}\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2-1\right)^{1/2}\text{EllipticF}\left(\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right),2^{1/2}\right)\right)/c^2\left(c\left(-1+2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2\right)\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{1/2}/\left(-c\left(2\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^4-\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2\right)^{1/2}/\left(-1+2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2/\sin\left(\frac{1}{2}bx+\frac{1}{2}a\right)/\left(c\left(-1+2\cos\left(\frac{1}{2}bx+\frac{1}{2}a\right)\right)^2\right)^{1/2}/b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.43

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \frac{-i \sqrt{2} \sqrt{c} \cos(bx + a)^2 \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) + i \sqrt{2} \sqrt{c} \cos(bx + a)^2 \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + 2 \sqrt{c} \cos(bx + a) \sin(bx + a)}{(b^3 c^3 \cos(bx + a)^2)}$$

[In] `integrate(1/(c*cos(b*x+a))^(5/2),x, algorithm="fricas")`

[Out]
$$\frac{1}{3}\left(-I\sqrt{2}\sqrt{c}\cos(bx+a)^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)+I\sin(bx+a))+I\sqrt{2}\sqrt{c}\cos(bx+a)^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)-I\sin(bx+a))+2\sqrt{c}\cos(bx+a)\sin(bx+a)\right)/(b^3c^3\cos(bx+a)^2)$$

Sympy [F]

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(a + bx))^{5/2}} dx$$

[In] `integrate(1/(c*cos(b*x+a))**(5/2),x)`

[Out] `Integral((c*cos(a + b*x))**(-5/2), x)`

Maxima [F]

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(-5/2), x)

Giac [F]

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{5}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(-5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(a + bx))^{\frac{5}{2}}} dx$$

[In] int(1/(c*cos(a + b*x))^(5/2),x)

[Out] int(1/(c*cos(a + b*x))^(5/2), x)

3.24 $\int \frac{1}{(c \cos(a+bx))^{7/2}} dx$

Optimal result	196
Rubi [A] (verified)	196
Mathematica [A] (verified)	197
Maple [B] (verified)	198
Fricas [C] (verification not implemented)	198
Sympy [F(-1)]	199
Maxima [F]	199
Giac [F]	199
Mupad [F(-1)]	199

Optimal result

Integrand size = 12, antiderivative size = 100

$$\int \frac{1}{(c \cos(a+bx))^{7/2}} dx = -\frac{6\sqrt{c \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bc^4 \sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{5bc(c \cos(a+bx))^{5/2}} + \frac{6 \sin(a+bx)}{5bc^3 \sqrt{c \cos(a+bx)}}$$

[Out] $2/5*\sin(b*x+a)/b/c/(c*\cos(b*x+a))^{(5/2)}+6/5*\sin(b*x+a)/b/c^3/(c*\cos(b*x+a))^{(1/2)}-6/5*(\cos(1/2*a+1/2*b*x))^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})*(c*\cos(b*x+a))^{(1/2)}/b/c^4/\cos(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$\int \frac{1}{(c \cos(a+bx))^{7/2}} dx = -\frac{6E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{c \cos(a+bx)}}{5bc^4 \sqrt{\cos(a+bx)}} + \frac{6 \sin(a+bx)}{5bc^3 \sqrt{c \cos(a+bx)}} + \frac{2 \sin(a+bx)}{5bc(c \cos(a+bx))^{5/2}}$$

[In] $\text{Int}[(c*\text{Cos}[a + b*x])^{(-7/2)}, x]$

[Out] $(-6*\text{Sqrt}[c*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*c^4*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(5*b*c*(c*\text{Cos}[a + b*x])^{(5/2)}) + (6*\text{Sin}[a + b*x])/(5*b*c^3*\text{Sqrt}[c*\text{Cos}[a + b*x]])$

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{2 \sin(a + bx)}{5bc(c \cos(a + bx))^{5/2}} + \frac{3 \int \frac{1}{(c \cos(a + bx))^{3/2}} dx}{5c^2} \\
&= \frac{2 \sin(a + bx)}{5bc(c \cos(a + bx))^{5/2}} + \frac{6 \sin(a + bx)}{5bc^3 \sqrt{c \cos(a + bx)}} - \frac{3 \int \sqrt{c \cos(a + bx)} dx}{5c^4} \\
&= \frac{2 \sin(a + bx)}{5bc(c \cos(a + bx))^{5/2}} + \frac{6 \sin(a + bx)}{5bc^3 \sqrt{c \cos(a + bx)}} - \frac{\left(3 \sqrt{c \cos(a + bx)}\right) \int \sqrt{\cos(a + bx)} dx}{5c^4 \sqrt{\cos(a + bx)}} \\
&= -\frac{6 \sqrt{c \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5bc^4 \sqrt{\cos(a + bx)}} + \frac{2 \sin(a + bx)}{5bc(c \cos(a + bx))^{5/2}} + \frac{6 \sin(a + bx)}{5bc^3 \sqrt{c \cos(a + bx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.68

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \frac{-6 \sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right) + 6 \sin(a + bx) + 2 \sec(a + bx) \tan(a + bx)}{5bc^3 \sqrt{c \cos(a + bx)}}$$

```
[In] Integrate[(c*Cos[a + b*x])^(-7/2),x]
```

```
[Out] (-6*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + 6*Sin[a + b*x] + 2*Sec[a
+ b*x]*Tan[a + b*x])/(5*b*c^3*Sqrt[c*Cos[a + b*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 365 vs. 2(112) = 224.

Time = 2.34 (sec) , antiderivative size = 366, normalized size of antiderivative = 3.66

method	result
default	$-\frac{2\sqrt{c(-1+2\left(\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right))\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{\left(24\cos\left(\frac{bx}{2}+\frac{a}{2}\right)\left(\sin^6\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-12\sqrt{2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-1}E\left(\cos\left(\frac{bx}{2}+\frac{a}{2}\right),\sqrt{2}\right)\sqrt{\frac{1}{2}}}\right)}$

[In] int(1/(c*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)

[Out]
$$-2/5*(c*(-1+2*\cos(1/2*b*x+1/2*a)^2)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/c^4/\sin(1/2*b*x+1/2*a)^3/(8*\sin(1/2*b*x+1/2*a)^6-12*\sin(1/2*b*x+1/2*a)^4+6*\sin(1/2*b*x+1/2*a)^2-1)*(24*\cos(1/2*b*x+1/2*a)*\sin(1/2*b*x+1/2*a)^6-12*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*EllipticE(\cos(1/2*b*x+1/2*a),2^{(1/2)})*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\sin(1/2*b*x+1/2*a)^4-24*\sin(1/2*b*x+1/2*a)^4*\cos(1/2*b*x+1/2*a)+12*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*EllipticE(\cos(1/2*b*x+1/2*a),2^{(1/2)})*(sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\sin(1/2*b*x+1/2*a)^2+8*\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a)-3*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*EllipticE(\cos(1/2*b*x+1/2*a),2^{(1/2)}))*(-2*\sin(1/2*b*x+1/2*a)^4*c+\sin(1/2*b*x+1/2*a)^2*c)^{(1/2)}/(c*(-1+2*\cos(1/2*b*x+1/2*a)^2))^{(1/2)}/b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.21

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \frac{-3i \sqrt{2} \sqrt{c} \cos(bx + a)^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a)))}{(c \cos(a + bx))^{7/2}}$$

[In] integrate(1/(c*cos(b*x+a))^(7/2),x, algorithm="fricas")

[Out]
$$1/5*(-3*I*\sqrt{2}*\sqrt{c}*\cos(b*x + a)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + 3*I*\sqrt{2}*\sqrt{c}*\cos(b*x + a)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a))) + 2*\sqrt{c}*\cos(b*x + a)*(3*\cos(b*x + a)^2 + 1)*\sin(b*x + a))/(b*c^4*\cos(b*x + a)^3)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \text{Timed out}$$

[In] integrate(1/(c*cos(b*x+a))**(7/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{7}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(7/2), x)

Giac [F]

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{7}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(7/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{7/2}} dx = \int \frac{1}{(c \cos(a + bx))^{7/2}} dx$$

[In] int(1/(c*cos(a + b*x))^(7/2),x)

[Out] int(1/(c*cos(a + b*x))^(7/2), x)

3.25 $\int \cos^{\frac{4}{3}}(a + bx) dx$

Optimal result	200
Rubi [A] (verified)	200
Mathematica [A] (verified)	201
Maple [F]	201
Fricas [F]	201
Sympy [F(-1)]	201
Maxima [F]	202
Giac [F]	202
Mupad [B] (verification not implemented)	202

Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \cos^{\frac{4}{3}}(a + bx) dx = -\frac{3 \cos^{\frac{7}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{7b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/7*\cos(b*x+a)^{(7/3)}*\operatorname{hypergeom}([1/2, 7/6], [13/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \cos^{\frac{4}{3}}(a + bx) dx = -\frac{3 \sin(a + bx) \cos^{\frac{7}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right)}{7b\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(4/3)}, x]$

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(7/3)}*\operatorname{Hypergeometric2F1}[1/2, 7/6, 13/6, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(7*b*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b*.)*\sin[(c*.) + (d*.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2]))*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x$
&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3 \cos^{\frac{7}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{7b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \cos^{\frac{4}{3}}(a + bx) dx$$

$$= -\frac{3 \cos^{\frac{7}{3}}(a + bx) \csc(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{7b}$$

[In] Integrate[Cos[a + b*x]^(4/3),x]

[Out] (-3*Cos[a + b*x]^(7/3)*Csc[a + b*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(7*b)

Maple [F]

$$\int \left(\cos^{\frac{4}{3}}(bx + a) \right) dx$$

[In] int(cos(b*x+a)^(4/3),x)

[Out] int(cos(b*x+a)^(4/3),x)

Fricas [F]

$$\int \cos^{\frac{4}{3}}(a + bx) dx = \int \cos(bx + a)^{\frac{4}{3}} dx$$

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(4/3), x)

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{4}{3}}(a + bx) dx = \text{Timed out}$$

[In] integrate(cos(b*x+a)**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int \cos^{\frac{4}{3}}(a + bx) dx = \int \cos(bx + a)^{\frac{4}{3}} dx$$

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(4/3), x)

Giac [F]

$$\int \cos^{\frac{4}{3}}(a + bx) dx = \int \cos(bx + a)^{\frac{4}{3}} dx$$

[In] integrate(cos(b*x+a)^(4/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(4/3), x)

Mupad [B] (verification not implemented)

Time = 13.89 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int \cos^{\frac{4}{3}}(a + bx) dx = -\frac{3 \cos(a + bx)^{7/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos(a + bx)^2\right)}{7b \sqrt{\sin(a + bx)^2}}$$

[In] int(cos(a + b*x)^(4/3),x)

[Out] -(3*cos(a + b*x)^(7/3)*sin(a + b*x)*hypergeom([1/2, 7/6], 13/6, cos(a + b*x)^2))/(7*b*(sin(a + b*x)^2)^(1/2))

3.26 $\int \cos^{\frac{2}{3}}(a + bx) dx$

Optimal result	203
Rubi [A] (verified)	203
Mathematica [A] (verified)	204
Maple [F]	204
Fricas [F]	204
Sympy [F]	204
Maxima [F]	205
Giac [F]	205
Mupad [B] (verification not implemented)	205

Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \cos^{\frac{2}{3}}(a + bx) dx = -\frac{3 \cos^{\frac{5}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{5b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/5*\cos(b*x+a)^{(5/3)}*\operatorname{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \cos^{\frac{2}{3}}(a + bx) dx = -\frac{3 \sin(a + bx) \cos^{\frac{5}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right)}{5b\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(2/3)}, x]$

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(5/3)}*\operatorname{Hypergeometric2F1}[1/2, 5/6, 11/6, \operatorname{Cos}[a + b*x]^{(2)}]*\operatorname{Sin}[a + b*x])/(5*b*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^{(2)}])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^{(2)}])]*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^{(2)}, x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x$ && $! \operatorname{IntegerQ}[2*n]$

Rubi steps

$$\text{integral} = -\frac{3 \cos^{\frac{5}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{5b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \cos^{\frac{2}{3}}(a + bx) dx$$

$$= -\frac{3 \cos^{\frac{5}{3}}(a + bx) \csc(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{5b}$$

[In] Integrate[Cos[a + b*x]^(2/3), x]

[Out] (-3*Cos[a + b*x]^(5/3)*Csc[a + b*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(5*b)

Maple [F]

$$\int \left(\cos^{\frac{2}{3}}(bx + a) \right) dx$$

[In] int(cos(b*x+a)^(2/3), x)

[Out] int(cos(b*x+a)^(2/3), x)

Fricas [F]

$$\int \cos^{\frac{2}{3}}(a + bx) dx = \int \cos(bx + a)^{\frac{2}{3}} dx$$

[In] integrate(cos(b*x+a)^(2/3), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(2/3), x)

Sympy [F]

$$\int \cos^{\frac{2}{3}}(a + bx) dx = \int \cos^{\frac{2}{3}}(a + bx) dx$$

[In] integrate(cos(b*x+a)**(2/3), x)

[Out] Integral(cos(a + b*x)**(2/3), x)

Maxima [F]

$$\int \cos^{\frac{2}{3}}(a + bx) dx = \int \cos (bx + a)^{\frac{2}{3}} dx$$

[In] integrate(cos(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(2/3), x)

Giac [F]

$$\int \cos^{\frac{2}{3}}(a + bx) dx = \int \cos (bx + a)^{\frac{2}{3}} dx$$

[In] integrate(cos(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(2/3), x)

Mupad [B] (verification not implemented)

Time = 15.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int \cos^{\frac{2}{3}}(a + bx) dx = -\frac{3 \cos(a + bx)^{5/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos(a + bx)^2\right)}{5b \sqrt{\sin(a + bx)^2}}$$

[In] int(cos(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(5/3)*sin(a + b*x)*hypergeom([1/2, 5/6], 11/6, cos(a + b*x)^2))/(5*b*(sin(a + b*x)^2)^(1/2))

3.27 $\int \sqrt[3]{\cos(a + bx)} dx$

Optimal result	206
Rubi [A] (verified)	206
Mathematica [A] (verified)	207
Maple [F]	207
Fricas [F]	207
Sympy [F]	207
Maxima [F]	208
Giac [F]	208
Mupad [B] (verification not implemented)	208

Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \sqrt[3]{\cos(a + bx)} dx = -\frac{3 \cos^{\frac{4}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4b\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/4*\cos(b*x+a)^{(4/3)}*\operatorname{hypergeom}([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b$
 $/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \sqrt[3]{\cos(a + bx)} dx = -\frac{3 \sin(a + bx) \cos^{\frac{4}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right)}{4b\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(1/3)}, x]$

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(4/3)}*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(4*b*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2, x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x$
 $\&\& \operatorname{IntegerQ}[2*n]$

Rubi steps

$$\operatorname{integral} = -\frac{3 \cos^{\frac{4}{3}}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \sqrt[3]{\cos(a+bx)} dx$$

$$= -\frac{3 \cos^{\frac{4}{3}}(a+bx) \csc(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a+bx)\right) \sqrt{\sin^2(a+bx)}}{4b}$$

[In] Integrate[Cos[a + b*x]^(1/3),x]

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(4/3)}*\operatorname{Csc}[a + b*x]*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])/(4*b)$

Maple [F]

$$\int \left(\cos^{\frac{1}{3}}(bx+a)\right) dx$$

[In] int(cos(b*x+a)^(1/3),x)

[Out] int(cos(b*x+a)^(1/3),x)

Fricas [F]

$$\int \sqrt[3]{\cos(a+bx)} dx = \int \cos(bx+a)^{\frac{1}{3}} dx$$

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(1/3), x)

Sympy [F]

$$\int \sqrt[3]{\cos(a+bx)} dx = \int \sqrt[3]{\cos(a+bx)} dx$$

[In] integrate(cos(b*x+a)**(1/3),x)

[Out] Integral(cos(a + b*x)**(1/3), x)

Maxima [F]

$$\int \sqrt[3]{\cos(a + bx)} dx = \int \cos(bx + a)^{\frac{1}{3}} dx$$

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(1/3), x)

Giac [F]

$$\int \sqrt[3]{\cos(a + bx)} dx = \int \cos(bx + a)^{\frac{1}{3}} dx$$

[In] integrate(cos(b*x+a)^(1/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(1/3), x)

Mupad [B] (verification not implemented)

Time = 15.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int \sqrt[3]{\cos(a + bx)} dx = -\frac{3 \cos(a + bx)^{4/3} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos(a + bx)^2\right)}{4b \sqrt{\sin(a + bx)^2}}$$

[In] int(cos(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(4/3)*sin(a + b*x)*hypergeom([1/2, 2/3], 5/3, cos(a + b*x)^2))/(4*b*(sin(a + b*x)^2)^(1/2))

$$3.28 \quad \int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

Optimal result	209
Rubi [A] (verified)	209
Mathematica [A] (verified)	210
Maple [F]	210
Fricas [F]	210
Sympy [F]	211
Maxima [F]	211
Giac [F]	211
Mupad [B] (verification not implemented)	211

Optimal result

Integrand size = 10, antiderivative size = 53

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

$$= -\frac{3 \cos^{\frac{2}{3}}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a+bx)\right) \sin(a+bx)}{2b\sqrt{\sin^2(a+bx)}}$$

[Out] $-3/2*\cos(b*x+a)^{(2/3)}*\operatorname{hypergeom}([1/3, 1/2], [4/3], \cos(b*x+a)^2)*\sin(b*x+a)/b$
 $/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

$$= -\frac{3 \sin(a+bx) \cos^{\frac{2}{3}}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a+bx)\right)}{2b\sqrt{\sin^2(a+bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(-1/3)}, x]$

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(2/3)}*\operatorname{Hypergeometric2F1}[1/3, 1/2, 4/3, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(2*b*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*\sin[(c_*) + (d_*)(x_*)])^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2}$

F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
 && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3 \cos^{\frac{2}{3}}(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{2b \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt[3]{\cos(a + bx)}} dx$$

$$= -\frac{3 \cos^{\frac{2}{3}}(a + bx) \csc(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{2b}$$

[In] Integrate[Cos[a + b*x]^(-1/3), x]

[Out] (-3*Cos[a + b*x]^(2/3)*Csc[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(2*b)

Maple [F]

$$\int \frac{1}{\cos(bx + a)^{\frac{1}{3}}} dx$$

[In] int(1/cos(b*x+a)^(1/3), x)

[Out] int(1/cos(b*x+a)^(1/3), x)

Fricas [F]

$$\int \frac{1}{\sqrt[3]{\cos(a + bx)}} dx = \int \frac{1}{\cos(bx + a)^{\frac{1}{3}}} dx$$

[In] integrate(1/cos(b*x+a)^(1/3), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-1/3), x)

Sympy [F]

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx = \int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

[In] integrate(1/cos(b*x+a)**(1/3),x)

[Out] Integral(cos(a + b*x)**(-1/3), x)

Maxima [F]

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx = \int \frac{1}{\cos(bx+a)^{\frac{1}{3}}} dx$$

[In] integrate(1/cos(b*x+a)^(1/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-1/3), x)

Giac [F]

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx = \int \frac{1}{\cos(bx+a)^{\frac{1}{3}}} dx$$

[In] integrate(1/cos(b*x+a)^(1/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-1/3), x)

Mupad [B] (verification not implemented)

Time = 14.95 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.79

$$\int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx = -\frac{3 \cos(a+bx)^{2/3} \sin(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos(a+bx)^2\right)}{2b \sqrt{\sin(a+bx)^2}}$$

[In] int(1/cos(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(2/3)*sin(a + b*x)*hypergeom([1/3, 1/2], 4/3, cos(a + b*x)^2))/(2*b*(sin(a + b*x)^2)^(1/2))

$$3.29 \quad \int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx$$

Optimal result	212
Rubi [A] (verified)	212
Mathematica [A] (verified)	213
Maple [F]	213
Fricas [F]	213
Sympy [F]	214
Maxima [F]	214
Giac [F]	214
Mupad [B] (verification not implemented)	214

Optimal result

Integrand size = 10, antiderivative size = 51

$$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx = -\frac{3\sqrt[3]{\cos(a+bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{b\sqrt{\sin^2(a+bx)}}$$

[Out] $-3*\cos(b*x+a)^{(1/3)}*\operatorname{hypergeom}([1/6, 1/2], [7/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx = -\frac{3 \sin(a+bx) \sqrt[3]{\cos(a+bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right)}{b\sqrt{\sin^2(a+bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^{(-2/3)}, x]$

[Out] $(-3*\operatorname{Cos}[a + b*x]^{(1/3)}*\operatorname{Hypergeometric2F1}[1/6, 1/2, 7/6, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(b*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\text{integral} = -\frac{3\sqrt[3]{\cos(a+bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{b\sqrt{\sin^2(a+bx)}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx$$

$$= -\frac{3\sqrt[3]{\cos(a+bx)} \operatorname{csc}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right) \sqrt{\sin^2(a+bx)}}{b}$$

[In] Integrate[Cos[a + b*x]^(-2/3),x]

[Out] (-3*Cos[a + b*x]^(1/3)*Csc[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/b

Maple [F]

$$\int \frac{1}{\cos^{\frac{2}{3}}(bx+a)} dx$$

[In] int(1/cos(b*x+a)^(2/3),x)

[Out] int(1/cos(b*x+a)^(2/3),x)

Fricas [F]

$$\int \frac{1}{\cos^{\frac{2}{3}}(a+bx)} dx = \int \frac{1}{\cos^{\frac{2}{3}}(bx+a)} dx$$

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-2/3), x)

Sympy [F]

$$\int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx$$

[In] integrate(1/cos(b*x+a)**(2/3),x)

[Out] Integral(cos(a + b*x)**(-2/3), x)

Maxima [F]

$$\int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{2}{3}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-2/3), x)

Giac [F]

$$\int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{2}{3}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-2/3), x)

Mupad [B] (verification not implemented)

Time = 15.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

$$\int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx = -\frac{3 \cos(a + bx)^{1/3} \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos(a + bx)^2\right)}{b \sqrt{\sin(a + bx)^2}}$$

[In] int(1/cos(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(1/3)*sin(a + b*x)*hypergeom([1/6, 1/2], 7/6, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(1/2))

3.30 $\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx$

Optimal result	215
Rubi [A] (verified)	215
Mathematica [A] (verified)	216
Maple [F]	216
Fricas [F]	216
Sympy [F]	217
Maxima [F]	217
Giac [F]	217
Mupad [B] (verification not implemented)	217

Optimal result

Integrand size = 10, antiderivative size = 51

$$\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{b \sqrt[3]{\cos(a+bx)} \sqrt{\sin^2(a+bx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sin(b*x+a)/b/cos(b*x+a)^(1/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int \frac{1}{\cos^{\frac{4}{3}}(a+bx)} dx = \frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a+bx)\right)}{b \sqrt{\sin^2(a+bx)} \sqrt[3]{\cos(a+bx)}}$$

[In] Int[Cos[a + b*x]^(-4/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*Cos[a + b*x]^(1/3)*Sqrt[Sin[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = \frac{3 \text{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{b^3 \sqrt{\cos(a + bx)} \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx$$

$$= \frac{3 \csc(a + bx) \text{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b^3 \sqrt{\cos(a + bx)}}$$

[In] Integrate[Cos[a + b*x]^(-4/3), x]

[Out] (3*Csc[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*cos[a + b*x]^(1/3))

Maple [F]

$$\int \frac{1}{\cos(bx + a)^{\frac{4}{3}}} dx$$

[In] int(1/cos(b*x+a)^(4/3), x)

[Out] int(1/cos(b*x+a)^(4/3), x)

Fricas [F]

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx = \int \frac{1}{\cos(bx + a)^{\frac{4}{3}}} dx$$

[In] integrate(1/cos(b*x+a)^(4/3), x, algorithm="fricas")

[Out] integral(cos(b*x + a)^(-4/3), x)

Sympy [F]

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx$$

[In] integrate(1/cos(b*x+a)**(4/3),x)

[Out] Integral(cos(a + b*x)**(-4/3), x)

Maxima [F]

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{4}{3}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(4/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(-4/3), x)

Giac [F]

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx = \int \frac{1}{\cos^{\frac{4}{3}}(bx + a)} dx$$

[In] integrate(1/cos(b*x+a)^(4/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(-4/3), x)

Mupad [B] (verification not implemented)

Time = 15.33 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.82

$$\int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx = \frac{3 \sin(a + bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos(a + bx)^2\right)}{b \cos(a + bx)^{1/3} \sqrt{\sin(a + bx)^2}}$$

[In] int(1/cos(a + b*x)^(4/3),x)

[Out] (3*sin(a + b*x)*hypergeom([-1/6, 1/2], 5/6, cos(a + b*x)^2))/(b*cos(a + b*x)^(1/3)*(sin(a + b*x)^2)^(1/2))

3.31 $\int (c \cos(a + bx))^{4/3} dx$

Optimal result	218
Rubi [A] (verified)	218
Mathematica [A] (verified)	219
Maple [F]	219
Fricas [F]	219
Sympy [F(-1)]	220
Maxima [F]	220
Giac [F]	220
Mupad [F(-1)]	220

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int (c \cos(a + bx))^{4/3} dx = \frac{3(c \cos(a + bx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{7bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/7*(c*\cos(b*x+a))^{(7/3)}*hypergeom([1/2, 7/6], [13/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int (c \cos(a + bx))^{4/3} dx = \frac{3 \sin(a + bx)(c \cos(a + bx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right)}{7bc\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(4/3)}, x]$

[Out] $(-3*(c*\operatorname{Cos}[a + b*x])^{(7/3)}*\operatorname{Hypergeometric2F1}[1/2, 7/6, 13/6, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(7*b*c*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2]))*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(c \cos(a + bx))^{7/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{7bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int (c \cos(a + bx))^{4/3} dx = \frac{3(c \cos(a + bx))^{4/3} \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{7b}$$

[In] Integrate[(c*Cos[a + b*x])^(4/3),x]

[Out] (-3*(c*Cos[a + b*x])^(4/3)*Cot[a + b*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(7*b)

Maple [F]

$$\int (c \cos(bx + a))^{4/3} dx$$

[In] int((c*cos(b*x+a))^(4/3),x)

[Out] int((c*cos(b*x+a))^(4/3),x)

Fricas [F]

$$\int (c \cos(a + bx))^{4/3} dx = \int (c \cos(bx + a))^{4/3} dx$$

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(1/3)*c*cos(b*x + a), x)

Sympy [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{4/3} dx = \text{Timed out}$$

[In] integrate((c*cos(b*x+a))**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int (c \cos(a + bx))^{4/3} dx = \int (c \cos(bx + a))^{\frac{4}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

Giac [F]

$$\int (c \cos(a + bx))^{4/3} dx = \int (c \cos(bx + a))^{\frac{4}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{4/3} dx = \int (c \cos(a + bx))^{4/3} dx$$

[In] int((c*cos(a + b*x))^(4/3),x)

[Out] int((c*cos(a + b*x))^(4/3), x)

3.32 $\int (c \cos(a + bx))^{2/3} dx$

Optimal result	221
Rubi [A] (verified)	221
Mathematica [A] (verified)	222
Maple [F]	222
Fricas [F]	222
Sympy [F]	223
Maxima [F]	223
Giac [F]	223
Mupad [F(-1)]	223

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int (c \cos(a + bx))^{2/3} dx = \frac{3(c \cos(a + bx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{5bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/5*(c*\cos(b*x+a))^{(5/3)}*\operatorname{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int (c \cos(a + bx))^{2/3} dx = \frac{3 \sin(a + bx)(c \cos(a + bx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right)}{5bc\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(2/3)}, x]$

[Out] $(-3*(c*\operatorname{Cos}[a + b*x])^{(5/3)}*\operatorname{Hypergeometric2F1}[1/2, 5/6, 11/6, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(5*b*c*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(c \cos(a + bx))^{5/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{5bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int (c \cos(a + bx))^{2/3} dx = \frac{3(c \cos(a + bx))^{2/3} \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{5b}$$

[In] Integrate[(c*Cos[a + b*x])^(2/3),x]

[Out] (-3*(c*Cos[a + b*x])^(2/3)*Cot[a + b*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(5*b)

Maple [F]

$$\int (c \cos(bx + a))^{2/3} dx$$

[In] int((c*cos(b*x+a))^(2/3),x)

[Out] int((c*cos(b*x+a))^(2/3),x)

Fricas [F]

$$\int (c \cos(a + bx))^{2/3} dx = \int (c \cos(bx + a))^{2/3} dx$$

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(2/3), x)

Sympy [F]

$$\int (c \cos(a + bx))^{2/3} dx = \int (c \cos(a + bx))^{\frac{2}{3}} dx$$

[In] integrate((c*cos(b*x+a))**(2/3),x)

[Out] Integral((c*cos(a + b*x))**(2/3), x)

Maxima [F]

$$\int (c \cos(a + bx))^{2/3} dx = \int (c \cos(bx + a))^{\frac{2}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

Giac [F]

$$\int (c \cos(a + bx))^{2/3} dx = \int (c \cos(bx + a))^{\frac{2}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(2/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^{2/3} dx = \int (c \cos(a + bx))^{\frac{2}{3}} dx$$

[In] int((c*cos(a + b*x))^(2/3),x)

[Out] int((c*cos(a + b*x))^(2/3), x)

3.33 $\int \sqrt[3]{c \cos(a + bx)} dx$

Optimal result	224
Rubi [A] (verified)	224
Mathematica [A] (verified)	225
Maple [F]	225
Fricas [F]	225
Sympy [F]	226
Maxima [F]	226
Giac [F]	226
Mupad [F(-1)]	226

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \sqrt[3]{c \cos(a + bx)} dx$$

$$= -\frac{3(c \cos(a + bx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/4*(c*\cos(b*x+a))^{(4/3)}*\operatorname{hypergeom}([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \sqrt[3]{c \cos(a + bx)} dx$$

$$= -\frac{3 \sin(a + bx)(c \cos(a + bx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right)}{4bc\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(1/3)}, x]$

[Out] $(-3*(c*\operatorname{Cos}[a + b*x])^{(4/3)}*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(4*b*c*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2]))*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(c \cos(a + bx))^{4/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{4bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \sqrt[3]{c \cos(a + bx)} dx$$

$$= -\frac{3\sqrt[3]{c \cos(a + bx)} \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{4b}$$

[In] Integrate[(c*Cos[a + b*x])^(1/3),x]

[Out] (-3*(c*Cos[a + b*x])^(1/3)*Cot[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(4*b)

Maple [F]

$$\int (c \cos(bx + a))^{\frac{1}{3}} dx$$

[In] int((c*cos(b*x+a))^(1/3),x)

[Out] int((c*cos(b*x+a))^(1/3),x)

Fricas [F]

$$\int \sqrt[3]{c \cos(a + bx)} dx = \int (c \cos(bx + a))^{\frac{1}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(1/3), x)

Sympy [F]

$$\int \sqrt[3]{c \cos(a + bx)} dx = \int \sqrt[3]{c \cos(a + bx)} dx$$

[In] integrate((c*cos(b*x+a))**(1/3),x)

[Out] Integral((c*cos(a + b*x))**(1/3), x)

Maxima [F]

$$\int \sqrt[3]{c \cos(a + bx)} dx = \int (c \cos(bx + a))^{\frac{1}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

Giac [F]

$$\int \sqrt[3]{c \cos(a + bx)} dx = \int (c \cos(bx + a))^{\frac{1}{3}} dx$$

[In] integrate((c*cos(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{c \cos(a + bx)} dx = \int (c \cos(a + bx))^{1/3} dx$$

[In] int((c*cos(a + b*x))^(1/3),x)

[Out] int((c*cos(a + b*x))^(1/3), x)

$$3.34 \quad \int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

Optimal result	227
Rubi [A] (verified)	227
Mathematica [A] (verified)	228
Maple [F]	228
Fricas [F]	228
Sympy [F]	229
Maxima [F]	229
Giac [F]	229
Mupad [F(-1)]	229

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

$$= -\frac{3(c \cos(a + bx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{2bc\sqrt{\sin^2(a + bx)}}$$

[Out] $-3/2*(c*\cos(b*x+a))^{(2/3)}*\operatorname{hypergeom}([1/3, 1/2], [4/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

$$= -\frac{3 \sin(a + bx)(c \cos(a + bx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right)}{2bc\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(-1/3)}, x]$

[Out] $(-3*(c*\operatorname{Cos}[a + b*x])^{(2/3)}*\operatorname{Hypergeometric2F1}[1/3, 1/2, 4/3, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(2*b*c*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2}$

F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
 && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(c \cos(a + bx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sin(a + bx)}{2bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

$$= -\frac{3 \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{2b\sqrt[3]{c \cos(a + bx)}}$$

[In] Integrate[(c*cos[a + b*x])^(-1/3),x]

[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(2*b*(c*cos[a + b*x])^(1/3))

Maple [F]

$$\int \frac{1}{(c \cos(bx + a))^{1/3}} dx$$

[In] int(1/(c*cos(b*x+a))^(1/3),x)

[Out] int(1/(c*cos(b*x+a))^(1/3),x)

Fricas [F]

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx = \int \frac{1}{(c \cos(bx + a))^{1/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(2/3)/(c*cos(b*x + a)), x)

Sympy [F]

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx = \int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx$$

[In] integrate(1/(c*cos(b*x+a))**(1/3),x)

[Out] Integral((c*cos(a + b*x))**(-1/3), x)

Maxima [F]

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{1}{3}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

Giac [F]

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx = \int \frac{1}{(c \cos(bx + a))^{\frac{1}{3}}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{c \cos(a + bx)}} dx = \int \frac{1}{(c \cos(a + bx))^{1/3}} dx$$

[In] int(1/(c*cos(a + b*x))^(1/3),x)

[Out] int(1/(c*cos(a + b*x))^(1/3), x)

3.35 $\int \frac{1}{(c \cos(a+bx))^{2/3}} dx$

Optimal result	230
Rubi [A] (verified)	230
Mathematica [A] (verified)	231
Maple [F]	231
Fricas [F]	231
Sympy [F]	232
Maxima [F]	232
Giac [F]	232
Mupad [F(-1)]	232

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(c \cos(a+bx))^{2/3}} dx = \frac{3 \sqrt[3]{c \cos(a+bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{bc \sqrt{\sin^2(a+bx)}}$$

[Out] $-3*(c*\cos(b*x+a))^{(1/3)}*\operatorname{hypergeom}([1/6, 1/2], [7/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{(c \cos(a+bx))^{2/3}} dx = \frac{3 \sin(a+bx) \sqrt[3]{c \cos(a+bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^{(-2/3)}, x]$

[Out] $(-3*(c*\operatorname{Cos}[a + b*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/6, 1/2, 7/6, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(b*c*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2}$

```
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\text{integral} = -\frac{3\sqrt[3]{c \cos(a + bx)} \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx)\right) \sin(a + bx)}{bc\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \frac{3 \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(c \cos(a + bx))^{2/3}}$$

```
[In] Integrate[(c*cos[a + b*x])^(-2/3),x]
```

```
[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[a + b*x]^2]*Sqrt[Sin[
a + b*x]^2])/(b*(c*cos[a + b*x])^(2/3))
```

Maple [F]

$$\int \frac{1}{(c \cos(bx + a))^{2/3}} dx$$

```
[In] int(1/(c*cos(b*x+a))^(2/3),x)
```

```
[Out] int(1/(c*cos(b*x+a))^(2/3),x)
```

Fricas [F]

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \int \frac{1}{(c \cos(bx + a))^{2/3}} dx$$

```
[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="fricas")
```

```
[Out] integral((c*cos(b*x + a))^(1/3)/(c*cos(b*x + a)), x)
```

Sympy [F]

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \int \frac{1}{(c \cos(a + bx))^{2/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))**(2/3),x)

[Out] Integral((c*cos(a + b*x))**(-2/3), x)

Maxima [F]

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \int \frac{1}{(c \cos(bx + a))^{2/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

Giac [F]

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \int \frac{1}{(c \cos(bx + a))^{2/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(2/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{2/3}} dx = \int \frac{1}{(c \cos(a + bx))^{2/3}} dx$$

[In] int(1/(c*cos(a + b*x))^(2/3),x)

[Out] int(1/(c*cos(a + b*x))^(2/3), x)

3.36 $\int \frac{1}{(c \cos(a+bx))^{4/3}} dx$

Optimal result	233
Rubi [A] (verified)	233
Mathematica [A] (verified)	234
Maple [F]	234
Fricas [F]	234
Sympy [F]	234
Maxima [F]	235
Giac [F]	235
Mupad [F(-1)]	235

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(c \cos(a+bx))^{4/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{bc \sqrt[3]{c \cos(a+bx)} \sqrt{\sin^2(a+bx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sin(b*x+a)/b/c/(c*cos(b*x+a))^(1/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{(c \cos(a+bx))^{4/3}} dx = \frac{3 \sin(a+bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a+bx)\right)}{bc \sqrt{\sin^2(a+bx)} \sqrt[3]{c \cos(a+bx)}}$$

[In] Int[(c*cos[a + b*x])^(-4/3),x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*c*(c*cos[a + b*x])^(1/3)*Sqrt[Sin[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a+bx)\right) \sin(a+bx)}{bc \sqrt[3]{c \cos(a+bx)} \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \frac{3 \cot(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(c \cos(a + bx))^{4/3}}$$

[In] Integrate[(c*Cos[a + b*x])^(-4/3),x]

[Out] (3*Cot[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(c*Cos[a + b*x])^(4/3))

Maple [F]

$$\int \frac{1}{(c \cos(bx + a))^{4/3}} dx$$

[In] int(1/(c*cos(b*x+a))^(4/3),x)

[Out] int(1/(c*cos(b*x+a))^(4/3),x)

Fricas [F]

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \int \frac{1}{(c \cos(bx + a))^{4/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^(2/3)/(c^2*cos(b*x + a)^2), x)

Sympy [F]

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \int \frac{1}{(c \cos(a + bx))^{4/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))**(4/3),x)

[Out] Integral((c*cos(a + b*x))**(-4/3), x)

Maxima [F]

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \int \frac{1}{(c \cos(bx + a))^{4/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

Giac [F]

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \int \frac{1}{(c \cos(bx + a))^{4/3}} dx$$

[In] integrate(1/(c*cos(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos(a + bx))^{4/3}} dx = \int \frac{1}{(c \cos(a + bx))^{4/3}} dx$$

[In] int(1/(c*cos(a + b*x))^(4/3),x)

[Out] int(1/(c*cos(a + b*x))^(4/3), x)

3.37 $\int \cos^n(a + bx) dx$

Optimal result	236
Rubi [A] (verified)	236
Mathematica [A] (verified)	237
Maple [F]	237
Fricas [F]	237
Sympy [F]	238
Maxima [F]	238
Giac [F]	238
Mupad [B] (verification not implemented)	238

Optimal result

Integrand size = 8, antiderivative size = 64

$$\int \cos^n(a + bx) dx = -\frac{\cos^{1+n}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a + bx)\right) \sin(a + bx)}{b(1+n)\sqrt{\sin^2(a + bx)}}$$

[Out] $-\cos(b*x+a)^{(1+n)}*\operatorname{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*\sin(b*x+a)/b/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2722}

$$\int \cos^n(a + bx) dx = -\frac{\sin(a + bx) \cos^{n+1}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \cos^2(a + bx)\right)}{b(n+1)\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^n, x]$

[Out] $-\left(\operatorname{Cos}[a + b*x]^{(1+n)}*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, (1+n)/2, (3+n)/2, \operatorname{Cos}[a + b*x]^2\right]*\operatorname{Sin}[a + b*x]\right)/(b*(1+n)*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, (n+1)/2, (n+3)/2, \operatorname{Sin}[c + d*x]^2, x\right] /; \operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{\cos^{1+n}(a+bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a+bx)\right) \sin(a+bx)}{b(1+n)\sqrt{\sin^2(a+bx)}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00

$$\int \cos^n(a+bx) dx = -\frac{\cos^{1+n}(a+bx) \csc(a+bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a+bx)\right) \sqrt{\sin^2(a+bx)}}{b(1+n)}$$

[In] Integrate[Cos[a + b*x]^n,x]

[Out] -(((Cos[a + b*x]^(1 + n)*Csc[a + b*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2]))/(b*(1 + n)))

Maple [F]

$$\int (\cos^n(bx+a)) dx$$

[In] int(cos(b*x+a)^n,x)

[Out] int(cos(b*x+a)^n,x)

Fricas [F]

$$\int \cos^n(a+bx) dx = \int \cos(bx+a)^n dx$$

[In] integrate(cos(b*x+a)^n,x, algorithm="fricas")

[Out] integral(cos(b*x + a)^n, x)

Sympy [F]

$$\int \cos^n(a + bx) dx = \int \cos^n(a + bx) dx$$

[In] integrate(cos(b*x+a)**n,x)

[Out] Integral(cos(a + b*x)**n, x)

Maxima [F]

$$\int \cos^n(a + bx) dx = \int \cos(bx + a)^n dx$$

[In] integrate(cos(b*x+a)^n,x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^n, x)

Giac [F]

$$\int \cos^n(a + bx) dx = \int \cos(bx + a)^n dx$$

[In] integrate(cos(b*x+a)^n,x, algorithm="giac")

[Out] integrate(cos(b*x + a)^n, x)

Mupad [B] (verification not implemented)

Time = 14.95 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.89

$$\int \cos^n(a + bx) dx = -\frac{\cos(a + bx)^{n+1} \sin(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n}{2} + \frac{1}{2}; \frac{n}{2} + \frac{3}{2}; \cos(a + bx)^2\right)}{b \sqrt{\sin(a + bx)^2} (n + 1)}$$

[In] int(cos(a + b*x)^n,x)

[Out] -(cos(a + b*x)^(n + 1)*sin(a + b*x)*hypergeom([1/2, n/2 + 1/2], n/2 + 3/2, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(1/2)*(n + 1))

3.38 $\int (c \cos(a + bx))^n dx$

Optimal result	239
Rubi [A] (verified)	239
Mathematica [A] (verified)	240
Maple [F]	240
Fricas [F]	240
Sympy [F]	241
Maxima [F]	241
Giac [F]	241
Mupad [F(-1)]	241

Optimal result

Integrand size = 10, antiderivative size = 69

$$\int (c \cos(a + bx))^n dx$$

$$= -\frac{(c \cos(a + bx))^{1+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a + bx)\right) \sin(a + bx)}{bc(1+n)\sqrt{\sin^2(a + bx)}}$$

[Out] $-(c*\cos(b*x+a))^{(1+n)}*\operatorname{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*\sin(b*x+a)/b/c/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int (c \cos(a + bx))^n dx$$

$$= -\frac{\sin(a + bx)(c \cos(a + bx))^{n+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \cos^2(a + bx)\right)}{bc(n+1)\sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x])^n, x]$

[Out] $-\left(\left(\left(c*\operatorname{Cos}[a + b*x]\right)^{(1+n)}*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \operatorname{Cos}[a + b*x]^2\right]*\operatorname{Sin}[a + b*x]\right)\right)/\left(b*c*(1+n)*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2]\right)$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(n+1)}{2}, \frac{(n+3)}{2}, \operatorname{Sin}[c + d*x]^2\right], x] /; \operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{(c \cos(a + bx))^{1+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a + bx)\right) \sin(a + bx)}{bc(1+n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.93

$$\int (c \cos(a + bx))^n dx = \frac{(c \cos(a + bx))^n \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(1+n)}$$

[In] Integrate[(c*Cos[a + b*x])^n,x]

[Out] -(((c*Cos[a + b*x])^n*Cot[a + b*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(1 + n)))

Maple [F]

$$\int (c \cos(bx + a))^n dx$$

[In] int((c*cos(b*x+a))^n,x)

[Out] int((c*cos(b*x+a))^n,x)

Fricas [F]

$$\int (c \cos(a + bx))^n dx = \int (c \cos(bx + a))^n dx$$

[In] integrate((c*cos(b*x+a))^n,x, algorithm="fricas")

[Out] integral((c*cos(b*x + a))^n, x)

Sympy [F]

$$\int (c \cos(a + bx))^n dx = \int (c \cos(a + bx))^n dx$$

[In] integrate((c*cos(b*x+a))**n,x)

[Out] Integral((c*cos(a + b*x))**n, x)

Maxima [F]

$$\int (c \cos(a + bx))^n dx = \int (c \cos(bx + a))^n dx$$

[In] integrate((c*cos(b*x+a))^n,x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a))^n, x)

Giac [F]

$$\int (c \cos(a + bx))^n dx = \int (c \cos(bx + a))^n dx$$

[In] integrate((c*cos(b*x+a))^n,x, algorithm="giac")

[Out] integrate((c*cos(b*x + a))^n, x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos(a + bx))^n dx = \int (c \cos(a + bx))^n dx$$

[In] int((c*cos(a + b*x))^n,x)

[Out] int((c*cos(a + b*x))^n, x)

3.39 $\int (a \cos^2(x))^{5/2} dx$

Optimal result	242
Rubi [A] (verified)	242
Mathematica [A] (verified)	243
Maple [A] (verified)	244
Fricas [A] (verification not implemented)	244
Sympy [F(-1)]	244
Maxima [A] (verification not implemented)	245
Giac [A] (verification not implemented)	245
Mupad [F(-1)]	245

Optimal result

Integrand size = 10, antiderivative size = 53

$$\int (a \cos^2(x))^{5/2} dx = \frac{8}{15} a^2 \sqrt{a \cos^2(x)} \tan(x) + \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x)$$

[Out] $4/15*a*(a*\cos(x)^2)^{(3/2)}*\tan(x)+1/5*(a*\cos(x)^2)^{(5/2)}*\tan(x)+8/15*a^2*(a*\cos(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3282, 3286, 2717}

$$\int (a \cos^2(x))^{5/2} dx = \frac{8}{15} a^2 \tan(x) \sqrt{a \cos^2(x)} + \frac{1}{5} \tan(x) (a \cos^2(x))^{5/2} + \frac{4}{15} a \tan(x) (a \cos^2(x))^{3/2}$$

[In] $\text{Int}[(a*\text{Cos}[x]^2)^{(5/2)}, x]$

[Out] $(8*a^2*\text{Sqrt}[a*\text{Cos}[x]^2]*\text{Tan}[x])/15 + (4*a*(a*\text{Cos}[x]^2)^{(3/2)}*\text{Tan}[x])/15 + (a*\text{Cos}[x]^2)^{(5/2)}*\text{Tan}[x])/5$

Rule 2717

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$
 $\text{FreeQ}\{c, d\}, x]$

Rule 3282

```
Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> Simp[(-Cot[e + f*x]
)]*((b*Sin[e + f*x]^2)^p/(2*f*p)), x] + Dist[b*((2*p - 1)/(2*p)), Int[(b*Si
n[e + f*x]^2)^(p - 1), x], x] /; FreeQ[{b, e, f}, x] && !IntegerQ[p] && Gt
Q[p, 1]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :> With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{5} (4a) \int (a \cos^2(x))^{3/2} dx \\
&= \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{15} (8a^2) \int \sqrt{a \cos^2(x)} dx \\
&= \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x) + \frac{1}{15} (8a^2 \sqrt{a \cos^2(x)} \sec(x)) \int \cos(x) dx \\
&= \frac{8}{15} a^2 \sqrt{a \cos^2(x)} \tan(x) + \frac{4}{15} a (a \cos^2(x))^{3/2} \tan(x) + \frac{1}{5} (a \cos^2(x))^{5/2} \tan(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.62

$$\int (a \cos^2(x))^{5/2} dx = \frac{1}{15} a^2 \sqrt{a \cos^2(x)} (15 - 10 \sin^2(x) + 3 \sin^4(x)) \tan(x)$$

[In] Integrate[(a*Cos[x]^2)^(5/2),x]

[Out] (a^2*Sqrt[a*Cos[x]^2]*(15 - 10*Sin[x]^2 + 3*Sin[x]^4)*Tan[x])/15

Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.60

method	result
default	$\frac{a^3 \cos(x) \sin(x) (3 \cos^4(x) + 4 \cos^2(x) + 8)}{15 \sqrt{a \cos^2(x)}}$
risch	$-\frac{ia^2 e^{6ix} \sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{160(e^{2ix}+1)} - \frac{5ia^2 e^{2ix} \sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{16(e^{2ix}+1)} + \frac{5ia^2 \sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{16(e^{2ix}+1)} + \frac{5ia^2 e^{-2ix} \sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{96(e^{2ix}+1)} - \frac{11ia^2}{96(e^{2ix}+1)}$

```
[In] int((a*cos(x)^2)^(5/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/15*a^3*cos(x)*sin(x)*(3*cos(x)^4+4*cos(x)^2+8)/(a*cos(x)^2)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.75

$$\int (a \cos^2(x))^{5/2} dx = \frac{(3a^2 \cos(x)^4 + 4a^2 \cos(x)^2 + 8a^2) \sqrt{a \cos(x)^2} \sin(x)}{15 \cos(x)}$$

```
[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="fricas")
```

```
[Out] 1/15*(3*a^2*cos(x)^4 + 4*a^2*cos(x)^2 + 8*a^2)*sqrt(a*cos(x)^2)*sin(x)/cos(x)
```

Sympy [F(-1)]

Timed out.

$$\int (a \cos^2(x))^{5/2} dx = \text{Timed out}$$

```
[In] integrate((a*cos(x)**2)**(5/2),x)
```

```
[Out] Timed out
```

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.58

$$\int (a \cos^2(x))^{5/2} dx = \frac{1}{240} (3 a^2 \sin(5x) + 25 a^2 \sin(3x) + 150 a^2 \sin(x)) \sqrt{a}$$

[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="maxima")

[Out] 1/240*(3*a^2*sin(5*x) + 25*a^2*sin(3*x) + 150*a^2*sin(x))*sqrt(a)

Giac [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.64

$$\int (a \cos^2(x))^{5/2} dx = \frac{1}{15} (3 a^2 \sin(x)^5 - 10 a^2 \sin(x)^3 + 15 a^2 \sin(x)) \sqrt{a} \operatorname{sgn}(\cos(x))$$

[In] integrate((a*cos(x)^2)^(5/2),x, algorithm="giac")

[Out] 1/15*(3*a^2*sin(x)^5 - 10*a^2*sin(x)^3 + 15*a^2*sin(x))*sqrt(a)*sgn(cos(x))

Mupad [F(-1)]

Timed out.

$$\int (a \cos^2(x))^{5/2} dx = \int (a \cos(x)^2)^{5/2} dx$$

[In] int((a*cos(x)^2)^(5/2),x)

[Out] int((a*cos(x)^2)^(5/2), x)

3.40 $\int (a \cos^2(x))^{3/2} dx$

Optimal result	246
Rubi [A] (verified)	246
Mathematica [A] (verified)	247
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	248
Sympy [F(-1)]	248
Maxima [A] (verification not implemented)	248
Giac [A] (verification not implemented)	248
Mupad [F(-1)]	249

Optimal result

Integrand size = 10, antiderivative size = 34

$$\int (a \cos^2(x))^{3/2} dx = \frac{2}{3}a\sqrt{a \cos^2(x)} \tan(x) + \frac{1}{3}(a \cos^2(x))^{3/2} \tan(x)$$

[Out] $1/3*(a*\cos(x)^2)^{(3/2)}*\tan(x)+2/3*a*(a*\cos(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3282, 3286, 2717}

$$\int (a \cos^2(x))^{3/2} dx = \frac{1}{3} \tan(x) (a \cos^2(x))^{3/2} + \frac{2}{3} a \tan(x) \sqrt{a \cos^2(x)}$$

[In] `Int[(a*cos[x]^2)^(3/2),x]`

[Out] `(2*a*Sqrt[a*cos[x]^2]*Tan[x])/3 + ((a*cos[x]^2)^(3/2)*Tan[x])/3`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

Rule 3282

`Int[((b_.)*sin[(e_.) + (f_.)*(x_)^2]^(p_), x_Symbol] := Simp[(-Cot[e + f*x])*((b*Sint[e + f*x]^2)^p/(2*f*p)), x] + Dist[b*((2*p - 1)/(2*p)), Int[(b*Sint[e + f*x]^2)^(p - 1), x], x] /;`
`FreeQ[{b, e, f}, x] && !IntegerQ[p] && Gt`

Q[p, 1]

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x]^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p]))], Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3}(a \cos^2(x))^{3/2} \tan(x) + \frac{1}{3}(2a) \int \sqrt{a \cos^2(x)} dx \\ &= \frac{1}{3}(a \cos^2(x))^{3/2} \tan(x) + \frac{1}{3} \left(2a \sqrt{a \cos^2(x)} \sec(x) \right) \int \cos(x) dx \\ &= \frac{2}{3} a \sqrt{a \cos^2(x)} \tan(x) + \frac{1}{3} (a \cos^2(x))^{3/2} \tan(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.68

$$\int (a \cos^2(x))^{3/2} dx = -\frac{1}{3} a \sqrt{a \cos^2(x)} (-3 + \sin^2(x)) \tan(x)$$

[In] Integrate[(a*cos[x]^2)^(3/2),x]

[Out] -1/3*(a*Sqrt[a*cos[x]^2]*(-3 + Sin[x]^2)*Tan[x])

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.71

method	result	size
default	$\frac{a^2 \cos(x) \sin(x) (\cos^2(x) + 2)}{3 \sqrt{a \cos^2(x)}}$	24
risch	$-\frac{ia e^{4ix} \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}{24(e^{2ix} + 1)} - \frac{3ia e^{2ix} \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}{8(e^{2ix} + 1)} + \frac{3ia \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}{8(e^{2ix} + 1)} + \frac{ia e^{-2ix} \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}{24 e^{2ix} + 24}$	141

[In] int((a*cos(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] 1/3*a^2*cos(x)*sin(x)*(cos(x)^2+2)/(a*cos(x)^2)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.76

$$\int (a \cos^2(x))^{3/2} dx = \frac{(a \cos(x)^2 + 2a) \sqrt{a \cos(x)^2} \sin(x)}{3 \cos(x)}$$

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="fricas")

[Out] 1/3*(a*cos(x)^2 + 2*a)*sqrt(a*cos(x)^2)*sin(x)/cos(x)

Sympy [F(-1)]

Timed out.

$$\int (a \cos^2(x))^{3/2} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**2)**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.50

$$\int (a \cos^2(x))^{3/2} dx = \frac{1}{12} (a \sin(3x) + 9a \sin(x)) \sqrt{a}$$

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="maxima")

[Out] 1/12*(a*sin(3*x) + 9*a*sin(x))*sqrt(a)

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.50

$$\int (a \cos^2(x))^{3/2} dx = -\frac{1}{3} (\sin(x)^3 - 3 \sin(x)) a^{3/2} \operatorname{sgn}(\cos(x))$$

[In] integrate((a*cos(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/3*(sin(x)^3 - 3*sin(x))*a^(3/2)*sgn(cos(x))

Mupad [F(-1)]

Timed out.

$$\int (a \cos^2(x))^{3/2} dx = \int (a \cos(x)^2)^{3/2} dx$$

```
[In] int((a*cos(x)^2)^(3/2),x)
```

```
[Out] int((a*cos(x)^2)^(3/2), x)
```

3.41 $\int \sqrt{a \cos^2(x)} dx$

Optimal result	250
Rubi [A] (verified)	250
Mathematica [A] (verified)	251
Maple [A] (verified)	251
Fricas [A] (verification not implemented)	251
Sympy [A] (verification not implemented)	252
Maxima [A] (verification not implemented)	252
Giac [A] (verification not implemented)	252
Mupad [B] (verification not implemented)	252

Optimal result

Integrand size = 10, antiderivative size = 13

$$\int \sqrt{a \cos^2(x)} dx = \sqrt{a \cos^2(x)} \tan(x)$$

[Out] (a*cos(x)^2)^(1/2)*tan(x)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3286, 2717}

$$\int \sqrt{a \cos^2(x)} dx = \tan(x) \sqrt{a \cos^2(x)}$$

[In] Int[Sqrt[a*Cos[x]^2],x]

[Out] Sqrt[a*Cos[x]^2]*Tan[x]

Rule 2717

```
Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;
FreeQ[{c, d}, x]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Ssin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
```

FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])

Rubi steps

$$\begin{aligned} \text{integral} &= \left(\sqrt{a \cos^2(x)} \sec(x) \right) \int \cos(x) dx \\ &= \sqrt{a \cos^2(x)} \tan(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{a \cos^2(x)} dx = \sqrt{a \cos^2(x)} \tan(x)$$

[In] Integrate[Sqrt[a*Cos[x]^2],x]

[Out] Sqrt[a*Cos[x]^2]*Tan[x]

Maple [A] (verified)

Time = 0.82 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{a \cos(x) \sin(x)}{\sqrt{a(\cos^2(x))}}$	15
risch	$-\frac{i\sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{2(e^{2ix}+1)} + \frac{i\sqrt{a(e^{2ix}+1)^2 e^{-2ix}}}{2e^{2ix}+2}$	67

[In] int((a*cos(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/(a*cos(x)^2)^(1/2)*a*cos(x)*sin(x)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sqrt{a \cos^2(x)} dx = \frac{\sqrt{a \cos^2(x)} \sin(x)}{\cos(x)}$$

[In] integrate((a*cos(x)^2)^(1/2),x, algorithm="fricas")

[Out] sqrt(a*cos(x)^2)*sin(x)/cos(x)

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \sqrt{a \cos^2(x)} dx = \frac{\sqrt{a \cos^2(x)} \sin(x)}{\cos(x)}$$

[In] integrate((a*cos(x)**2)**(1/2),x)

[Out] sqrt(a*cos(x)**2)*sin(x)/cos(x)

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.46

$$\int \sqrt{a \cos^2(x)} dx = \sqrt{a} \sin(x)$$

[In] integrate((a*cos(x)^2)^(1/2),x, algorithm="maxima")

[Out] sqrt(a)*sin(x)

Giac [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 9, normalized size of antiderivative = 0.69

$$\int \sqrt{a \cos^2(x)} dx = \sqrt{a} \operatorname{sgn}(\cos(x)) \sin(x)$$

[In] integrate((a*cos(x)^2)^(1/2),x, algorithm="giac")

[Out] sqrt(a)*sgn(cos(x))*sin(x)

Mupad [B] (verification not implemented)

Time = 14.48 (sec) , antiderivative size = 46, normalized size of antiderivative = 3.54

$$\int \sqrt{a \cos^2(x)} dx = \frac{\sqrt{2} \sqrt{a} \sqrt{\cos(2x) + 1} (\cos(2x) - 1 + \sin(2x) 1i)}{2 (\cos(2x) 1i - \sin(2x) + 1i)}$$

[In] int((a*cos(x)^2)^(1/2),x)

[Out] (2^(1/2)*a^(1/2)*(cos(2*x) + 1)^(1/2)*(cos(2*x) + sin(2*x)*1i - 1))/(2*(cos(2*x)*1i - sin(2*x) + 1i))

3.42 $\int \frac{1}{\sqrt{a \cos^2(x)}} dx$

Optimal result	253
Rubi [A] (verified)	253
Mathematica [A] (verified)	254
Maple [B] (verified)	254
Fricas [B] (verification not implemented)	255
Sympy [F]	255
Maxima [B] (verification not implemented)	255
Giac [F]	256
Mupad [F(-1)]	256

Optimal result

Integrand size = 10, antiderivative size = 16

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \frac{\operatorname{arctanh}(\sin(x)) \cos(x)}{\sqrt{a \cos^2(x)}}$$

[Out] $\operatorname{arctanh}(\sin(x)) \cdot \cos(x) / (a \cdot \cos(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3286, 3855}

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \frac{\cos(x) \operatorname{arctanh}(\sin(x))}{\sqrt{a \cos^2(x)}}$$

[In] `Int[1/Sqrt[a*Cos[x]^2], x]`

[Out] `(ArcTanh[Sin[x]]*Cos[x])/Sqrt[a*Cos[x]^2]`

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIn[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos(x) \int \sec(x) dx}{\sqrt{a \cos^2(x)}} \\ &= \frac{\operatorname{arctanh}(\sin(x)) \cos(x)}{\sqrt{a \cos^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \frac{\operatorname{arctanh}(\sin(x)) \cos(x)}{\sqrt{a \cos^2(x)}}$$

```
[In] Integrate[1/Sqrt[a*Cos[x]^2],x]
```

```
[Out] (ArcTanh[Sin[x]]*Cos[x])/Sqrt[a*Cos[x]^2]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. 2(14) = 28.

Time = 0.90 (sec) , antiderivative size = 48, normalized size of antiderivative = 3.00

method	result	size
default	$\frac{\cos(x) \sqrt{a(\sin^2(x))} \ln\left(\frac{2\sqrt{a} \sqrt{a(\sin^2(x)) + 2a}}{\cos(x)}\right)}{\sqrt{a} \sin(x) \sqrt{a(\cos^2(x))}}$	48
risch	$-\frac{2 \ln(e^{ix} - i) \cos(x)}{\sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}} + \frac{2 \ln(e^{ix} + i) \cos(x)}{\sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}$	64

```
[In] int(1/(a*cos(x)^2)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] cos(x)*(a*sin(x)^2)^(1/2)/a^(1/2)*ln(2*(a^(1/2)*(a*sin(x)^2)^(1/2)+a)/cos(x))
/sin(x)/(a*cos(x)^2)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(14) = 28$.

Time = 0.28 (sec) , antiderivative size = 65, normalized size of antiderivative = 4.06

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \left[-\frac{\sqrt{a \cos(x)^2} \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right)}{2a \cos(x)}, -\frac{\sqrt{-a} \arctan\left(\frac{\sqrt{a \cos(x)^2} \sqrt{-a} \sin(x)}{a \cos(x)}\right)}{a} \right]$$

[In] integrate(1/(a*cos(x)^2)^(1/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(a*cos(x)^2)*log(-(sin(x) - 1)/(sin(x) + 1))/(a*cos(x)), -sqrt(-a)*arctan(sqrt(a*cos(x)^2)*sqrt(-a)*sin(x)/(a*cos(x)))/a]

Sympy [F]

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \int \frac{1}{\sqrt{a \cos^2(x)}} dx$$

[In] integrate(1/(a*cos(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(a*cos(x)**2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 38 vs. $2(14) = 28$.

Time = 0.39 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.38

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \frac{\log(\cos(x)^2 + \sin(x)^2 + 2 \sin(x) + 1) - \log(\cos(x)^2 + \sin(x)^2 - 2 \sin(x) + 1)}{2\sqrt{a}}$$

[In] integrate(1/(a*cos(x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1))/sqrt(a)

Giac [F]

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \int \frac{1}{\sqrt{a \cos(x)^2}} dx$$

[In] integrate(1/(a*cos(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(a*cos(x)^2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a \cos^2(x)}} dx = \int \frac{1}{\sqrt{a \cos(x)^2}} dx$$

[In] int(1/(a*cos(x)^2)^(1/2),x)

[Out] int(1/(a*cos(x)^2)^(1/2), x)

3.43 $\int \frac{1}{(a \cos^2(x))^{3/2}} dx$

Optimal result	257
Rubi [A] (verified)	257
Mathematica [A] (verified)	258
Maple [B] (verified)	258
Fricas [A] (verification not implemented)	259
Sympy [F]	259
Maxima [B] (verification not implemented)	260
Giac [A] (verification not implemented)	260
Mupad [F(-1)]	260

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(x)) \cos(x)}{2a\sqrt{a \cos^2(x)}} + \frac{\tan(x)}{2a\sqrt{a \cos^2(x)}}$$

[Out] 1/2*arctanh(sin(x))*cos(x)/a/(a*cos(x)^2)^(1/2)+1/2*tan(x)/a/(a*cos(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3283, 3286, 3855}

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = \frac{\cos(x) \operatorname{arctanh}(\sin(x))}{2a\sqrt{a \cos^2(x)}} + \frac{\tan(x)}{2a\sqrt{a \cos^2(x)}}$$

[In] Int[(a*Cos[x]^2)^(-3/2), x]

[Out] (ArcTanh[Sin[x]]*Cos[x])/(2*a*Sqrt[a*Cos[x]^2]) + Tan[x]/(2*a*Sqrt[a*Cos[x]^2])

Rule 3283

```
Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> Simp[Cot[e + f*x]*
((b*Sin[e + f*x]^2)^(p + 1)/(b*f*(2*p + 1))), x] + Dist[2*((p + 1)/(b*(2*p
+ 1))), Int[(b*Sin[e + f*x]^2)^(p + 1), x], x] /; FreeQ[{b, e, f}, x] && !
IntegerQ[p] && LtQ[p, -1]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\tan(x)}{2a\sqrt{a\cos^2(x)}} + \frac{\int \frac{1}{\sqrt{a\cos^2(x)}} dx}{2a} \\ &= \frac{\tan(x)}{2a\sqrt{a\cos^2(x)}} + \frac{\cos(x) \int \sec(x) dx}{2a\sqrt{a\cos^2(x)}} \\ &= \frac{\operatorname{arctanh}(\sin(x)) \cos(x)}{2a\sqrt{a\cos^2(x)}} + \frac{\tan(x)}{2a\sqrt{a\cos^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.62

$$\int \frac{1}{(a\cos^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(x)) \cos(x) + \tan(x)}{2a\sqrt{a\cos^2(x)}}$$

```
[In] Integrate[(a*Cos[x]^2)^(-3/2), x]
```

```
[Out] (ArcTanh[Sin[x]]*Cos[x] + Tan[x])/(2*a*Sqrt[a*Cos[x]^2])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 69 vs. 2(34) = 68.

Time = 1.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.67

method	result	size
default	$\frac{\sqrt{a(\sin^2(x))} \left(\ln \left(\frac{2\sqrt{a}\sqrt{a(\sin^2(x))+2a}}{\cos(x)} \right) a(\cos^2(x)) + \sqrt{a}\sqrt{a(\sin^2(x))} \right)}{2a^{\frac{5}{2}} \cos(x) \sin(x) \sqrt{a(\cos^2(x))}}$	70
risch	$-\frac{i(e^{2ix}-1)}{a(e^{2ix}+1)\sqrt{a(e^{2ix}+1)^2e^{-2ix}}} - \frac{\ln(e^{ix}-i)\cos(x)}{a\sqrt{a(e^{2ix}+1)^2e^{-2ix}}} + \frac{\ln(e^{ix}+i)\cos(x)}{a\sqrt{a(e^{2ix}+1)^2e^{-2ix}}}$	109

[In] `int(1/(a*cos(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2} \frac{1}{a^{5/2}} \frac{1}{\cos(x)} \frac{(a \sin(x)^2)^{1/2} (\ln(2 \sqrt{a} \sqrt{a \sin^2(x) + 2a}) + \sqrt{a} \sqrt{a \sin^2(x)})}{\cos(x) a \cos^2(x) + a^{1/2} (a \sin(x)^2)^{1/2}} \frac{1}{\sin(x) (a \cos(x)^2)^{1/2}}$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.95

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = -\frac{\sqrt{a \cos^2(x)} \left(\cos^2(x) \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right) - 2 \sin(x) \right)}{4 a^2 \cos^3(x)}$$

[In] `integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="fricas")`

[Out] $-\frac{1}{4} \sqrt{a \cos^2(x)} \frac{(\cos^2(x) \log(-(\sin(x) - 1)/(\sin(x) + 1)) - 2 \sin(x))}{(a^2 \cos^3(x))}$

Sympy [F]

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = \int \frac{1}{(a \cos^2(x))^{\frac{3}{2}}} dx$$

[In] `integrate(1/(a*cos(x)**2)**(3/2),x)`

[Out] `Integral((a*cos(x)**2)**(-3/2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(34) = 68$.

Time = 0.40 (sec) , antiderivative size = 304, normalized size of antiderivative = 7.24

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = \frac{4(\sin(3x) - \sin(x)) \cos(4x) + (2(2 \cos(2x) + 1) \cos(4x) + \cos(4x)^2 + 4 \cos(2x))^2}{(a \cos^2(x))^{3/2}}$$

[In] integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="maxima")

[Out] $\frac{1}{4} \cdot (4 \cdot (\sin(3x) - \sin(x)) \cdot \cos(4x) + (2 \cdot (2 \cdot \cos(2x) + 1) \cdot \cos(4x) + \cos(4x)^2 + 4 \cdot \cos(2x)^2 + \sin(4x)^2 + 4 \cdot \sin(4x) \cdot \sin(2x) + 4 \cdot \sin(2x)^2 + 4 \cdot \cos(2x) + 1) \cdot \log(\cos(x)^2 + \sin(x)^2 + 2 \cdot \sin(x) + 1) - (2 \cdot (2 \cdot \cos(2x) + 1) \cdot \cos(4x) + \cos(4x)^2 + 4 \cdot \cos(2x)^2 + \sin(4x)^2 + 4 \cdot \sin(4x) \cdot \sin(2x) + 4 \cdot \sin(2x)^2 + 4 \cdot \cos(2x) + 1) \cdot \log(\cos(x)^2 + \sin(x)^2 - 2 \cdot \sin(x) + 1) - 4 \cdot (\cos(3x) - \cos(x)) \cdot \sin(4x) + 4 \cdot (2 \cdot \cos(2x) + 1) \cdot \sin(3x) - 8 \cdot \cos(3x) \cdot \sin(2x) + 8 \cdot \cos(x) \cdot \sin(2x) - 8 \cdot \cos(2x) \cdot \sin(x) - 4 \cdot \sin(x)) / ((a \cdot \cos(4x)^2 + 4 \cdot a \cdot \cos(2x)^2 + a \cdot \sin(4x)^2 + 4 \cdot a \cdot \sin(4x) \cdot \sin(2x) + 4 \cdot a \cdot \sin(2x)^2 + 2 \cdot (2 \cdot a \cdot \cos(2x) + a) \cdot \cos(4x) + 4 \cdot a \cdot \cos(2x) + a) \cdot \sqrt{a})$

Giac [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = -\frac{\sin(x)}{2(\sin(x)^2 - 1)a^{3/2} \operatorname{sgn}(\cos(x))}$$

[In] integrate(1/(a*cos(x)^2)^(3/2),x, algorithm="giac")

[Out] $-1/2 \cdot \sin(x) / ((\sin(x)^2 - 1) \cdot a^{3/2} \cdot \operatorname{sgn}(\cos(x)))$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^2(x))^{3/2}} dx = \int \frac{1}{(a \cos(x)^2)^{3/2}} dx$$

[In] int(1/(a*cos(x)^2)^(3/2),x)

[Out] int(1/(a*cos(x)^2)^(3/2), x)

3.44 $\int \frac{1}{(a \cos^2(x))^{5/2}} dx$

Optimal result	261
Rubi [A] (verified)	261
Mathematica [A] (verified)	262
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	263
Sympy [F(-1)]	263
Maxima [B] (verification not implemented)	264
Giac [A] (verification not implemented)	265
Mupad [F(-1)]	265

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \frac{3 \operatorname{arctanh}(\sin(x)) \cos(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}}$$

[Out] $3/8 * \operatorname{arctanh}(\sin(x)) * \cos(x) / a^2 / (a * \cos(x)^2)^{(1/2)} + 1/4 * \tan(x) / a / (a * \cos(x)^2)^{(3/2)} + 3/8 * \tan(x) / a^2 / (a * \cos(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3283, 3286, 3855}

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \frac{3 \cos(x) \operatorname{arctanh}(\sin(x))}{8a^2 \sqrt{a \cos^2(x)}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}}$$

[In] $\operatorname{Int}[(a * \operatorname{Cos}[x]^2)^{-5/2}, x]$

[Out] $(3 * \operatorname{ArcTanh}[\operatorname{Sin}[x]] * \operatorname{Cos}[x]) / (8 * a^2 * \operatorname{Sqrt}[a * \operatorname{Cos}[x]^2]) + \operatorname{Tan}[x] / (4 * a * (a * \operatorname{Cos}[x]^2)^{(3/2)}) + (3 * \operatorname{Tan}[x]) / (8 * a^2 * \operatorname{Sqrt}[a * \operatorname{Cos}[x]^2])$

Rule 3283

$\operatorname{Int}[(b * \sin[e + f * x] + (f * x)^2)^p, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cot}[e + f * x] * ((b * \sin[e + f * x]^2)^{p+1} / (b * f * (2 * p + 1))), x] + \operatorname{Dist}[2 * ((p + 1) / (b * (2 * p + 1))), \operatorname{Int}[(b * \sin[e + f * x]^2)^{p+1}, x], x] /;$ $\operatorname{FreeQ}\{b, e, f, x\} \&\& \operatorname{IntegerQ}[p] \&\& \operatorname{LtQ}[p, -1]$

Rule 3286

```
Int[(u_)*((b_)*sin[e_ + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIn[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_)*(trig_)[e + f*x])^(m_) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rule 3855

```
Int[csc[(c_.) + (d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \int \frac{1}{(a \cos^2(x))^{3/2}} dx}{4a} \\
&= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{3 \int \frac{1}{\sqrt{a \cos^2(x)}} dx}{8a^2} \\
&= \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{(3 \cos(x)) \int \sec(x) dx}{8a^2 \sqrt{a \cos^2(x)}} \\
&= \frac{3 \arctanh(\sin(x)) \cos(x)}{8a^2 \sqrt{a \cos^2(x)}} + \frac{\tan(x)}{4a (a \cos^2(x))^{3/2}} + \frac{3 \tan(x)}{8a^2 \sqrt{a \cos^2(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.59

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \frac{3 \arctanh(\sin(x)) \cos(x) + (3 + 2 \sec^2(x)) \tan(x)}{8a^2 \sqrt{a \cos^2(x)}}$$

```
[In] Integrate[(a*Cos[x]^2)^(-5/2), x]
```

```
[Out] (3*ArcTanh[Sin[x]]*Cos[x] + (3 + 2*Sec[x]^2)*Tan[x])/(8*a^2*Sqrt[a*Cos[x]^2
])
```

Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.46

method	result	size
default	$\frac{\sqrt{a(\sin^2(x))} \left(3 \ln \left(\frac{2\sqrt{a}\sqrt{a(\sin^2(x))+2a}}{\cos(x)} \right) a(\cos^4(x)) + 3\sqrt{a(\sin^2(x))} (\cos^2(x))\sqrt{a} + 2\sqrt{a}\sqrt{a(\sin^2(x))} \right)}{8a^{\frac{7}{2}} \cos(x)^3 \sin(x) \sqrt{a(\cos^2(x))}}$	89
risch	$-\frac{i(3e^{6ix} + 11e^{4ix} - 11e^{2ix} - 3)}{4a^2(e^{2ix} + 1)^3 \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}} + \frac{3 \ln(e^{ix} + i) \cos(x)}{4a^2 \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}} - \frac{3 \ln(e^{ix} - i) \cos(x)}{4a^2 \sqrt{a(e^{2ix} + 1)^2 e^{-2ix}}}$	126

[In] int(1/(a*cos(x)^2)^(5/2),x,method=_RETURNVERBOSE)

[Out] $\frac{1}{8} a^{-7/2} / \cos(x)^3 * (a * \sin(x)^2)^{1/2} * (3 * \ln(2 * (a^{1/2} * (a * \sin(x)^2)^{1/2} + a) / \cos(x)) * a * \cos(x)^4 + 3 * (a * \sin(x)^2)^{1/2} * \cos(x)^2 * a^{1/2} + 2 * a^{1/2} * (a * \sin(x)^2)^{1/2}) / \sin(x) / (a * \cos(x)^2)^{1/2}$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.80

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = -\frac{\left(3 \cos(x)^4 \log\left(-\frac{\sin(x)-1}{\sin(x)+1}\right) - 2(3 \cos(x)^2 + 2) \sin(x) \right) \sqrt{a \cos(x)^2}}{16 a^3 \cos(x)^5}$$

[In] integrate(1/(a*cos(x)^2)^(5/2),x, algorithm="fricas")

[Out] $-1/16 * (3 * \cos(x)^4 * \log(-(\sin(x) - 1) / (\sin(x) + 1)) - 2 * (3 * \cos(x)^2 + 2) * \sin(x)) * \sqrt{a * \cos(x)^2} / (a^3 * \cos(x)^5)$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \text{Timed out}$$

[In] integrate(1/(a*cos(x)**2)**(5/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 933 vs. $2(49) = 98$.

Time = 0.62 (sec) , antiderivative size = 933, normalized size of antiderivative = 15.30

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(1/(a*cos(x)^2)^(5/2),x, algorithm="maxima")

[Out] $\frac{1}{16} * (4 * (3 * \sin(7x) + 11 * \sin(5x) - 11 * \sin(3x) - 3 * \sin(x)) * \cos(8x) - 24 * (2 * \sin(6x) + 3 * \sin(4x) + 2 * \sin(2x)) * \cos(7x) + 16 * (11 * \sin(5x) - 11 * \sin(3x) - 3 * \sin(x)) * \cos(6x) - 88 * (3 * \sin(4x) + 2 * \sin(2x)) * \cos(5x) - 24 * (11 * \sin(3x) + 3 * \sin(x)) * \cos(4x) + 3 * (2 * (4 * \cos(6x) + 6 * \cos(4x) + 4 * \cos(2x) + 1) * \cos(8x) + \cos(8x)^2 + 8 * (6 * \cos(4x) + 4 * \cos(2x) + 1) * \cos(6x) + 16 * \cos(6x)^2 + 12 * (4 * \cos(2x) + 1) * \cos(4x) + 36 * \cos(4x)^2 + 16 * \cos(2x)^2 + 4 * (2 * \sin(6x) + 3 * \sin(4x) + 2 * \sin(2x)) * \sin(8x) + \sin(8x)^2 + 16 * (3 * \sin(4x) + 2 * \sin(2x)) * \sin(6x) + 16 * \sin(6x)^2 + 36 * \sin(4x)^2 + 48 * \sin(4x) * \sin(2x) + 16 * \sin(2x)^2 + 8 * \cos(2x) + 1) * \log(\cos(x)^2 + \sin(x)^2 + 2 * \sin(x) + 1) - 3 * (2 * (4 * \cos(6x) + 6 * \cos(4x) + 4 * \cos(2x) + 1) * \cos(8x) + \cos(8x)^2 + 8 * (6 * \cos(4x) + 4 * \cos(2x) + 1) * \cos(6x) + 16 * \cos(6x)^2 + 12 * (4 * \cos(2x) + 1) * \cos(4x) + 36 * \cos(4x)^2 + 16 * \cos(2x)^2 + 4 * (2 * \sin(6x) + 3 * \sin(4x) + 2 * \sin(2x)) * \sin(8x) + \sin(8x)^2 + 16 * (3 * \sin(4x) + 2 * \sin(2x)) * \sin(6x) + 16 * \sin(6x)^2 + 36 * \sin(4x)^2 + 48 * \sin(4x) * \sin(2x) + 16 * \sin(2x)^2 + 8 * \cos(2x) + 1) * \log(\cos(x)^2 + \sin(x)^2 - 2 * \sin(x) + 1) - 4 * (3 * \cos(7x) + 11 * \cos(5x) - 11 * \cos(3x) - 3 * \cos(x)) * \sin(8x) + 12 * (4 * \cos(6x) + 6 * \cos(4x) + 4 * \cos(2x) + 1) * \sin(7x) - 16 * (11 * \cos(5x) - 11 * \cos(3x) - 3 * \cos(x)) * \sin(6x) + 44 * (6 * \cos(4x) + 4 * \cos(2x) + 1) * \sin(5x) + 24 * (11 * \cos(3x) + 3 * \cos(x)) * \sin(4x) - 44 * (4 * \cos(2x) + 1) * \sin(3x) + 176 * \cos(3x) * \sin(2x) + 48 * \cos(x) * \sin(2x) - 48 * \cos(2x) * \sin(x) - 12 * \sin(x)) / ((a^2 * \cos(8x)^2 + 16 * a^2 * \cos(6x)^2 + 36 * a^2 * \cos(4x)^2 + 16 * a^2 * \cos(2x)^2 + a^2 * \sin(8x)^2 + 16 * a^2 * \sin(6x)^2 + 36 * a^2 * \sin(4x)^2 + 48 * a^2 * \sin(4x) * \sin(2x) + 16 * a^2 * \sin(2x)^2 + 8 * a^2 * \cos(2x) + a^2 + 2 * (4 * a^2 * \cos(6x) + 6 * a^2 * \cos(4x) + 4 * a^2 * \cos(2x) + a^2) * \cos(8x) + 8 * (6 * a^2 * \cos(4x) + 4 * a^2 * \cos(2x) + a^2) * \cos(6x) + 12 * (4 * a^2 * \cos(2x) + a^2) * \cos(4x) + 4 * (2 * a^2 * \sin(6x) + 3 * a^2 * \sin(4x) + 2 * a^2 * \sin(2x)) * \sin(8x) + 16 * (3 * a^2 * \sin(4x) + 2 * a^2 * \sin(2x)) * \sin(6x)) * \sqrt{a})$

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.48

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = -\frac{3 \sin(x)^3 - 5 \sin(x)}{8 (\sin(x)^2 - 1)^2 a^{5/2} \operatorname{sgn}(\cos(x))}$$

[In] integrate(1/(a*cos(x)^2)^(5/2),x, algorithm="giac")

[Out] -1/8*(3*sin(x)^3 - 5*sin(x))/((sin(x)^2 - 1)^2*a^(5/2)*sgn(cos(x)))

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^2(x))^{5/2}} dx = \int \frac{1}{(a \cos(x)^2)^{5/2}} dx$$

[In] int(1/(a*cos(x)^2)^(5/2),x)

[Out] int(1/(a*cos(x)^2)^(5/2), x)

3.45 $\int (a \cos^3(x))^{5/2} dx$

Optimal result	266
Rubi [A] (verified)	266
Mathematica [A] (verified)	268
Maple [C] (verified)	268
Fricas [C] (verification not implemented)	269
Sympy [F(-1)]	269
Maxima [F]	269
Giac [F]	269
Mupad [F(-1)]	270

Optimal result

Integrand size = 10, antiderivative size = 117

$$\int (a \cos^3(x))^{5/2} dx = \frac{26a^2 \sqrt{a \cos^3(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{77 \cos^{\frac{3}{2}}(x)} + \frac{78}{385} a^2 \cos(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{77} a^2 \sqrt{a \cos^3(x)} \tan(x)$$

[Out] $26/77*a^2*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\operatorname{EllipticF}(\sin(1/2*x), 2^{(1/2)})*(a*\cos(x)^3)^{(1/2)}/\cos(x)^{(3/2)}+78/385*a^2*\cos(x)*\sin(x)*(a*\cos(x)^3)^{(1/2)}+26/165*a^2*\cos(x)^3*\sin(x)*(a*\cos(x)^3)^{(1/2)}+2/15*a^2*\cos(x)^5*\sin(x)*(a*\cos(x)^3)^{(1/2)}+26/77*a^2*(a*\cos(x)^3)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 2720}

$$\int (a \cos^3(x))^{5/2} dx = \frac{26}{165} a^2 \sin(x) \cos^3(x) \sqrt{a \cos^3(x)} + \frac{78}{385} a^2 \sin(x) \cos(x) \sqrt{a \cos^3(x)} + \frac{26}{77} a^2 \tan(x) \sqrt{a \cos^3(x)} + \frac{26a^2 \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) \sqrt{a \cos^3(x)}}{77 \cos^{\frac{3}{2}}(x)} + \frac{2}{15} a^2 \sin(x) \cos^5(x) \sqrt{a \cos^3(x)}$$

[In] $\operatorname{Int}[(a*\cos[x]^3)^{(5/2)}, x]$

```
[Out] (26*a^2*Sqrt[a*Cos[x]^3]*EllipticF[x/2, 2])/(77*Cos[x]^(3/2)) + (78*a^2*Cos[x]*Sqrt[a*Cos[x]^3]*Sin[x])/385 + (26*a^2*Cos[x]^3*Sqrt[a*Cos[x]^3]*Sin[x])/165 + (2*a^2*Cos[x]^5*Sqrt[a*Cos[x]^3]*Sin[x])/15 + (26*a^2*Sqrt[a*Cos[x]^3]*Tan[x])/77
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] :> With[{ff = FreeFactors[SIN[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x])^n)^FracPart[p]/(SIN[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(SIN[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(a^2 \sqrt{a \cos^3(x)}\right) \int \cos^{\frac{15}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\
 &= \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{\left(13 a^2 \sqrt{a \cos^3(x)}\right) \int \cos^{\frac{11}{2}}(x) dx}{15 \cos^{\frac{3}{2}}(x)} \\
 &= \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) \\
 &\quad + \frac{\left(39 a^2 \sqrt{a \cos^3(x)}\right) \int \cos^{\frac{7}{2}}(x) dx}{55 \cos^{\frac{3}{2}}(x)} \\
 &= \frac{78}{385} a^2 \cos(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165} a^2 \cos^3(x) \sqrt{a \cos^3(x)} \sin(x) \\
 &\quad + \frac{2}{15} a^2 \cos^5(x) \sqrt{a \cos^3(x)} \sin(x) + \frac{\left(39 a^2 \sqrt{a \cos^3(x)}\right) \int \cos^{\frac{3}{2}}(x) dx}{77 \cos^{\frac{3}{2}}(x)}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{78}{385}a^2 \cos(x)\sqrt{a \cos^3(x)} \sin(x) + \frac{26}{165}a^2 \cos^3(x)\sqrt{a \cos^3(x)} \sin(x) \\
&\quad + \frac{2}{15}a^2 \cos^5(x)\sqrt{a \cos^3(x)} \sin(x) + \frac{26}{77}a^2 \sqrt{a \cos^3(x)} \tan(x) \\
&\quad + \frac{\left(13a^2 \sqrt{a \cos^3(x)}\right) \int \frac{1}{\sqrt{\cos(x)}} dx}{77 \cos^{\frac{3}{2}}(x)} \\
&= \frac{26a^2 \sqrt{a \cos^3(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{77 \cos^{\frac{3}{2}}(x)} + \frac{78}{385}a^2 \cos(x)\sqrt{a \cos^3(x)} \sin(x) \\
&\quad + \frac{26}{165}a^2 \cos^3(x)\sqrt{a \cos^3(x)} \sin(x) \\
&\quad + \frac{2}{15}a^2 \cos^5(x)\sqrt{a \cos^3(x)} \sin(x) + \frac{26}{77}a^2 \sqrt{a \cos^3(x)} \tan(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.52

$$\int (a \cos^3(x))^{5/2} dx = \frac{a(a \cos^3(x))^{3/2} \left(12480 \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) + \sqrt{\cos(x)}(15465 \sin(x) + 3657 \sin(3x) + 749 \sin(5x) + 77 \sin(7x))\right)}{36960 \cos^{\frac{9}{2}}(x)}$$

[In] Integrate[(a*Cos[x]^3)^(5/2), x]

[Out] (a*(a*Cos[x]^3)^(3/2)*(12480*EllipticF[x/2, 2] + Sqrt[Cos[x]]*(15465*Sin[x] + 3657*Sin[3*x] + 749*Sin[5*x] + 77*Sin[7*x])))/(36960*Cos[x]^(9/2))

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 12.33 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00

method	result
default	$-\frac{2\sqrt{a(\cos^3(x))}a^2\left(-77(\cos^5(x))\sin(x)-91(\cos^3(x))\sin(x)+195i\sec(x)F(i(\csc(x)-\cot(x)),i)\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}+195i(\sec^2(x))}\right)}{1155}$

[In] int((a*cos(x)^3)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/1155*(a*cos(x)^3)^(1/2)*a^2*(-77*cos(x)^5*sin(x)-91*cos(x)^3*sin(x)+195*I*sec(x)*EllipticF(I*(csc(x)-cot(x)), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+195*I*sec(x)^2*EllipticF(I*(csc(x)-cot(x)), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-117*cos(x)*sin(x)-195*tan(x))

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.79

$$\int (a \cos^3(x))^{5/2} dx = \frac{195i \sqrt{2} a^{5/2} \cos(x) \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)) - 195i \sqrt{2} a^{5/2} \cos(x)}{1}$$

[In] integrate((a*cos(x)^3)^(5/2),x, algorithm="fricas")

[Out] 1/1155*(195*I*sqrt(2)*a^(5/2)*cos(x)*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - 195*I*sqrt(2)*a^(5/2)*cos(x)*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)) + 2*(77*a^2*cos(x)^6 + 91*a^2*cos(x)^4 + 117*a^2*cos(x)^2 + 195*a^2)*sqrt(a*cos(x)^3)*sin(x))/cos(x)

Sympy [F(-1)]

Timed out.

$$\int (a \cos^3(x))^{5/2} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**3)**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int (a \cos^3(x))^{5/2} dx = \int (a \cos(x)^3)^{5/2} dx$$

[In] integrate((a*cos(x)^3)^(5/2),x, algorithm="maxima")

[Out] integrate((a*cos(x)^3)^(5/2), x)

Giac [F]

$$\int (a \cos^3(x))^{5/2} dx = \int (a \cos(x)^3)^{5/2} dx$$

[In] integrate((a*cos(x)^3)^(5/2),x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos^3(x))^{5/2} dx = \int (a \cos(x)^3)^{5/2} dx$$

```
[In] int((a*cos(x)^3)^(5/2),x)
```

```
[Out] int((a*cos(x)^3)^(5/2), x)
```

3.46 $\int (a \cos^3(x))^{3/2} dx$

Optimal result	271
Rubi [A] (verified)	271
Mathematica [A] (verified)	272
Maple [C] (verified)	273
Fricas [C] (verification not implemented)	273
Sympy [F(-1)]	274
Maxima [F]	274
Giac [F]	274
Mupad [F(-1)]	274

Optimal result

Integrand size = 10, antiderivative size = 67

$$\int (a \cos^3(x))^{3/2} dx = \frac{14a \sqrt{a \cos^3(x)} E\left(\frac{x}{2} \mid 2\right)}{15 \cos^{\frac{3}{2}}(x)} + \frac{14}{45} a \sqrt{a \cos^3(x)} \sin(x) + \frac{2}{9} a \cos^2(x) \sqrt{a \cos^3(x)} \sin(x)$$

[Out] 14/15*a*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticE(sin(1/2*x),2^(1/2))*(a*cos(x)^3)^(1/2)/cos(x)^(3/2)+14/45*a*sin(x)*(a*cos(x)^3)^(1/2)+2/9*a*cos(x)^2*sin(x)*(a*cos(x)^3)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 2719}

$$\int (a \cos^3(x))^{3/2} dx = \frac{14}{45} a \sin(x) \sqrt{a \cos^3(x)} + \frac{14a E\left(\frac{x}{2} \mid 2\right) \sqrt{a \cos^3(x)}}{15 \cos^{\frac{3}{2}}(x)} + \frac{2}{9} a \sin(x) \cos^2(x) \sqrt{a \cos^3(x)}$$

[In] Int[(a*cos[x]^3)^(3/2),x]

[Out] (14*a*Sqrt[a*cos[x]^3]*EllipticE[x/2, 2])/(15*cos[x]^(3/2)) + (14*a*Sqrt[a*cos[x]^3]*Sin[x])/45 + (2*a*cos[x]^2*Sqrt[a*cos[x]^3]*Sin[x])/9

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[

$(c + d*x)^{(n - 2)}, x, x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2 * n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \ :> \ \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3286

$\text{Int}[(u_.)*((b_.)*\sin[(e_.) + (f_.)*(x_.)]^{(n_.)})^{(p_.)}, x_Symbol] \ :> \ \text{With}[\{\text{ff} = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Dist}[(b*\text{ff}^n)^{\text{IntPart}[p]}*(b*\text{Sin}[e + f*x]^{n - \text{FracPart}[p]}/(\text{Sin}[e + f*x]/\text{ff})^{(n*\text{FracPart}[p])}), \text{Int}[\text{ActivateTrig}[u]*(\text{Sin}[e + f*x]/\text{ff})^{(n*p)}, x], x]] /; \text{FreeQ}[\{b, e, f, n, p\}, x] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{IntegerQ}[n] \ \&\& \ (\text{EqQ}[u, 1] \ || \ \text{MatchQ}[u, ((d_.)*(trig_)[e + f*x])^{(m_.)}] /; \text{FreeQ}[\{d, m\}, x] \ \&\& \ \text{MemberQ}[\{\sin, \cos, \tan, \cot, \sec, \csc\}, \text{trig}])]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(a\sqrt{a\cos^3(x)}\right) \int \cos^{\frac{9}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\ &= \frac{2}{9}a\cos^2(x)\sqrt{a\cos^3(x)}\sin(x) + \frac{\left(7a\sqrt{a\cos^3(x)}\right) \int \cos^{\frac{5}{2}}(x) dx}{9\cos^{\frac{3}{2}}(x)} \\ &= \frac{14}{45}a\sqrt{a\cos^3(x)}\sin(x) + \frac{2}{9}a\cos^2(x)\sqrt{a\cos^3(x)}\sin(x) + \frac{\left(7a\sqrt{a\cos^3(x)}\right) \int \sqrt{\cos(x)} dx}{15\cos^{\frac{3}{2}}(x)} \\ &= \frac{14a\sqrt{a\cos^3(x)}E\left(\frac{x}{2}\mid 2\right)}{15\cos^{\frac{3}{2}}(x)} + \frac{14}{45}a\sqrt{a\cos^3(x)}\sin(x) + \frac{2}{9}a\cos^2(x)\sqrt{a\cos^3(x)}\sin(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.75

$$\int (a\cos^3(x))^{3/2} dx = \frac{(a\cos^3(x))^{3/2} \left(168E\left(\frac{x}{2}\mid 2\right) + \sqrt{\cos(x)}(38\sin(2x) + 5\sin(4x))\right)}{180\cos^{\frac{9}{2}}(x)}$$

[In] Integrate[(a*Cos[x]^3)^(3/2), x]

[Out] ((a*Cos[x]^3)^(3/2)*(168*EllipticE[x/2, 2] + Sqrt[Cos[x]]*(38*Sin[2*x] + 5*Sin[4*x])))/(180*Cos[x]^(9/2))

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 6.04 (sec) , antiderivative size = 271, normalized size of antiderivative = 4.04

method	result
default	$\frac{2\sqrt{a(\cos^3(x))} a \left(5(\cos^3(x)) \sin(x) - 21iF(i(\csc(x) - \cot(x)), i) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} + 21i \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} E(i(\csc(x) - \cot(x))) \right)}{1}$

[In] `int((a*cos(x)^3)^(3/2), x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 2/45*(a*\cos(x)^3)^{(1/2)}*a/(\cos(x)+1)*(5*\cos(x)^3*\sin(x)-21*I*\text{EllipticF}(I*(\csc(x)-\cot(x)), I)*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}+21*I*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}*\text{EllipticE}(I*(\csc(x)-\cot(x)), I)+5*\cos(x)^2*\sin(x)-42*I*\sec(x)*\text{EllipticF}(I*(\csc(x)-\cot(x)), I)*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}+42*I*\sec(x)*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}*\text{EllipticE}(I*(\csc(x)-\cot(x)), I)+7*\cos(x)*\sin(x)-21*I*\sec(x)^2*\text{EllipticF}(I*(\csc(x)-\cot(x)), I)*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}+21*I*\sec(x)^2*(1/(\cos(x)+1))^{(1/2)}*(\cos(x)/(\cos(x)+1))^{(1/2)}*\text{EllipticE}(I*(\csc(x)-\cot(x)), I)+7*\sin(x)+21*\tan(x)) \end{aligned}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.99

$$\begin{aligned} & \int (a \cos^3(x))^{3/2} dx = \\ & -\frac{7}{15}i\sqrt{2}a^{3/2}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x))) \\ & + \frac{7}{15}i\sqrt{2}a^{3/2}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) - i \sin(x))) \\ & + \frac{2}{45}\sqrt{a \cos(x)^3}(5a \cos(x)^2 + 7a) \sin(x) \end{aligned}$$

[In] `integrate((a*cos(x)^3)^(3/2), x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -7/15*I*\text{sqrt}(2)*a^{(3/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) + I*\sin(x))) + 7/15*I*\text{sqrt}(2)*a^{(3/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(x) - I*\sin(x))) + 2/45*\text{sqrt}(a*\cos(x)^3)*(5*a*\cos(x)^2 + 7*a)*\sin(x) \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int (a \cos^3(x))^{3/2} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**3)**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int (a \cos^3(x))^{3/2} dx = \int (a \cos(x)^3)^{\frac{3}{2}} dx$$

[In] integrate((a*cos(x)^3)^(3/2),x, algorithm="maxima")

[Out] integrate((a*cos(x)^3)^(3/2), x)

Giac [F]

$$\int (a \cos^3(x))^{3/2} dx = \int (a \cos(x)^3)^{\frac{3}{2}} dx$$

[In] integrate((a*cos(x)^3)^(3/2),x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos^3(x))^{3/2} dx = \int (a \cos(x)^3)^{3/2} dx$$

[In] int((a*cos(x)^3)^(3/2),x)

[Out] int((a*cos(x)^3)^(3/2), x)

3.47 $\int \sqrt{a \cos^3(x)} dx$

Optimal result	275
Rubi [A] (verified)	275
Mathematica [A] (verified)	276
Maple [C] (verified)	276
Fricas [C] (verification not implemented)	277
Sympy [F]	277
Maxima [F]	278
Giac [F]	278
Mupad [F(-1)]	278

Optimal result

Integrand size = 10, antiderivative size = 44

$$\int \sqrt{a \cos^3(x)} dx = \frac{2\sqrt{a \cos^3(x)} \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{3 \cos^{\frac{3}{2}}(x)} + \frac{2}{3} \sqrt{a \cos^3(x)} \tan(x)$$

[Out] $2/3*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\operatorname{EllipticF}(\sin(1/2*x), 2^{(1/2)})*(a*\cos(x)^3)^{(1/2)}/\cos(x)^{(3/2)}+2/3*(a*\cos(x)^3)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 2720}

$$\int \sqrt{a \cos^3(x)} dx = \frac{2}{3} \tan(x) \sqrt{a \cos^3(x)} + \frac{2 \operatorname{EllipticF}\left(\frac{x}{2}, 2\right) \sqrt{a \cos^3(x)}}{3 \cos^{\frac{3}{2}}(x)}$$

[In] `Int[Sqrt[a*Cos[x]^3],x]`

[Out] $(2*\operatorname{Sqrt}[a*\operatorname{Cos}[x]^3]*\operatorname{EllipticF}[x/2, 2])/(3*\operatorname{Cos}[x]^{(3/2)}) + (2*\operatorname{Sqrt}[a*\operatorname{Cos}[x]^3]*\operatorname{Tan}[x])/3$

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
]*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p]))], Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{a \cos^3(x)} \int \cos^{\frac{3}{2}}(x) dx}{\cos^{\frac{3}{2}}(x)} \\ &= \frac{2}{3} \sqrt{a \cos^3(x)} \tan(x) + \frac{\sqrt{a \cos^3(x)} \int \frac{1}{\sqrt{\cos(x)}} dx}{3 \cos^{\frac{3}{2}}(x)} \\ &= \frac{2 \sqrt{a \cos^3(x)} \text{EllipticF}\left(\frac{x}{2}, 2\right)}{3 \cos^{\frac{3}{2}}(x)} + \frac{2}{3} \sqrt{a \cos^3(x)} \tan(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.84

$$\int \sqrt{a \cos^3(x)} dx = \frac{2 \sqrt{a \cos^3(x)} \left(\text{EllipticF}\left(\frac{x}{2}, 2\right) + \sqrt{\cos(x)} \sin(x) \right)}{3 \cos^{\frac{3}{2}}(x)}$$

```
[In] Integrate[Sqrt[a*Cos[x]^3], x]
```

```
[Out] (2*Sqrt[a*Cos[x]^3]*(EllipticF[x/2, 2] + Sqrt[Cos[x]]*Sin[x]))/(3*Cos[x]^(3/2))
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 2.91 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.09

method	result
default	$-\frac{2\sqrt{a(\cos^3(x))} \left(i \sec(x) F(i \csc(x) - \cot(x), i) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} + i (\sec^2(x)) F(i \csc(x) - \cot(x), i) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} - \tan(x) \right)}{3}$

```
[In] int((a*cos(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/3*(a*cos(x)^3)^(1/2)*(I*sec(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)+I*sec(x)^2*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)-tan(x))
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.34

$$\int \sqrt{a \cos^3(x)} dx$$

$$= \frac{i \sqrt{2} \sqrt{a} \cos(x) \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)) - i \sqrt{2} \sqrt{a} \cos(x) \operatorname{weierstrassPInverse}(-4, 0, \cos(x) - i \sin(x))}{3 \cos(x)}$$

```
[In] integrate((a*cos(x)^3)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/3*(I*sqrt(2)*sqrt(a)*cos(x)*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - I*sqrt(2)*sqrt(a)*cos(x)*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)) + 2*sqrt(a*cos(x)^3)*sin(x))/cos(x)
```

Sympy [F]

$$\int \sqrt{a \cos^3(x)} dx = \int \sqrt{a \cos^3(x)} dx$$

```
[In] integrate((a*cos(x)**3)**(1/2),x)
```

```
[Out] Integral(sqrt(a*cos(x)**3), x)
```

Maxima [F]

$$\int \sqrt{a \cos^3(x)} dx = \int \sqrt{a \cos(x)^3} dx$$

[In] integrate((a*cos(x)^3)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*cos(x)^3), x)

Giac [F]

$$\int \sqrt{a \cos^3(x)} dx = \int \sqrt{a \cos(x)^3} dx$$

[In] integrate((a*cos(x)^3)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(a*cos(x)^3), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a \cos^3(x)} dx = \int \sqrt{a \cos(x)^3} dx$$

[In] int((a*cos(x)^3)^(1/2),x)

[Out] int((a*cos(x)^3)^(1/2), x)

3.48 $\int \frac{1}{\sqrt{a \cos^3(x)}} dx$

Optimal result	279
Rubi [A] (verified)	279
Mathematica [A] (verified)	280
Maple [C] (verified)	280
Fricas [C] (verification not implemented)	281
Sympy [F]	281
Maxima [F]	282
Giac [F]	282
Mupad [F(-1)]	282

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = -\frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right)}{\sqrt{a \cos^3(x)}} + \frac{2 \cos(x) \sin(x)}{\sqrt{a \cos^3(x)}}$$

[Out] $-2*\cos(x)^{(3/2)}*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\text{EllipticE}(\sin(1/2*x), 2^{(1/2)})/(a*\cos(x)^3)^{(1/2)}+2*\cos(x)*\sin(x)/(a*\cos(x)^3)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2716, 2719}

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \frac{2 \sin(x) \cos(x)}{\sqrt{a \cos^3(x)}} - \frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right)}{\sqrt{a \cos^3(x)}}$$

[In] Int[1/Sqrt[a*Cos[x]^3], x]

[Out] $(-2*\text{Cos}[x]^{(3/2)}*\text{EllipticE}[x/2, 2])/ \text{Sqrt}[a*\text{Cos}[x]^3] + (2*\text{Cos}[x]*\text{Sin}[x])/ \text{Sqrt}[a*\text{Cos}[x]^3]$

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)^(n_)])^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p]))], Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{3}{2}}(x)} dx}{\sqrt{a \cos^3(x)}} \\ &= \frac{2 \cos(x) \sin(x)}{\sqrt{a \cos^3(x)}} - \frac{\cos^{\frac{3}{2}}(x) \int \sqrt{\cos(x)} dx}{\sqrt{a \cos^3(x)}} \\ &= -\frac{2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right)}{\sqrt{a \cos^3(x)}} + \frac{2 \cos(x) \sin(x)}{\sqrt{a \cos^3(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.74

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \frac{-2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right) + \sin(2x)}{\sqrt{a \cos^3(x)}}$$

```
[In] Integrate[1/Sqrt[a*Cos[x]^3], x]
```

```
[Out] (-2*Cos[x]^(3/2)*EllipticE[x/2, 2] + Sin[2*x])/Sqrt[a*Cos[x]^3]
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 3.47 (sec) , antiderivative size = 244, normalized size of antiderivative = 5.81

method	result
default	$\frac{2 \left(i \sqrt{\frac{\cos(x)}{\cos(x)+1}} \sqrt{\frac{1}{\cos(x)+1}} F(i(\csc(x)-\cot(x)), i)(\cos^2(x)) - i \sqrt{\frac{\cos(x)}{\cos(x)+1}} \sqrt{\frac{1}{\cos(x)+1}} E(i(\csc(x)-\cot(x)), i)(\cos^2(x)) + 2i \sqrt{\frac{\cos(x)}{\cos(x)+1}} \sqrt{\frac{1}{\cos(x)+1}} \right)}{\sqrt{a \cos^3(x)}}$


```
[In] int(1/(a*cos(x)^3)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 2*(I*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*cos(x)^2-I*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)*cos(x)^2+2*I*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*cos(x)-2*I*(cos(x)/(cos(x)+1))^(1/2)*(1/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)*cos(x)+I*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)*(1/(cos(x)+1))^(1/2)-I*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)*(1/(cos(x)+1))^(1/2)+sin(x))*cos(x)/(cos(x)+1)/(a*cos(x)^3)^(1/2)
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.69

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx$$

$$= \frac{i \sqrt{2} \sqrt{a} \cos(x)^2 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x))) - i \sqrt{2} \sqrt{a} \cos(x)^2}{a \cos(x)^2}$$

```
[In] integrate(1/(a*cos(x)^3)^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(a)*cos(x)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) + I*sin(x))) - I*sqrt(2)*sqrt(a)*cos(x)^2*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) - I*sin(x))) + 2*sqrt(a*cos(x)^3)*sin(x))/(a*cos(x)^2)
```

Sympy [F]

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \int \frac{1}{\sqrt{a \cos^3(x)}} dx$$

```
[In] integrate(1/(a*cos(x)**3)**(1/2),x)
```

```
[Out] Integral(1/sqrt(a*cos(x)**3), x)
```

Maxima [F]

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

[In] integrate(1/(a*cos(x)^3)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(a*cos(x)^3), x)

Giac [F]

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

[In] integrate(1/(a*cos(x)^3)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(a*cos(x)^3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a \cos^3(x)}} dx = \int \frac{1}{\sqrt{a \cos(x)^3}} dx$$

[In] int(1/(a*cos(x)^3)^(1/2),x)

[Out] int(1/(a*cos(x)^3)^(1/2), x)

3.49 $\int \frac{1}{(a \cos^3(x))^{3/2}} dx$

Optimal result	283
Rubi [A] (verified)	283
Mathematica [A] (verified)	284
Maple [C] (verified)	285
Fricas [C] (verification not implemented)	285
Sympy [F(-1)]	285
Maxima [F]	286
Giac [F]	286
Mupad [F(-1)]	286

Optimal result

Integrand size = 10, antiderivative size = 71

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \frac{10 \cos^{3/2}(x) \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{21a \sqrt{a \cos^3(x)}} + \frac{10 \sin(x)}{21a \sqrt{a \cos^3(x)}} + \frac{2 \sec(x) \tan(x)}{7a \sqrt{a \cos^3(x)}}$$

[Out] 10/21*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),2^(1/2))/a/(a*cos(x)^3)^(1/2)+10/21*sin(x)/a/(a*cos(x)^3)^(1/2)+2/7*sec(x)*tan(x)/a/(a*cos(x)^3)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2716, 2720}

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \frac{10 \sin(x)}{21a \sqrt{a \cos^3(x)}} + \frac{10 \cos^{3/2}(x) \operatorname{EllipticF}\left(\frac{x}{2}, 2\right)}{21a \sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec(x)}{7a \sqrt{a \cos^3(x)}}$$

[In] Int[(a*cos[x]^3)^(-3/2),x]

[Out] (10*cos[x]^(3/2)*EllipticF[x/2, 2])/(21*a*Sqrt[a*cos[x]^3]) + (10*sin[x])/(21*a*Sqrt[a*cos[x]^3]) + (2*Sec[x]*Tan[x])/(7*a*Sqrt[a*cos[x]^3])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Ssin[e + f*x]^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{9}{2}}(x)} dx}{a\sqrt{a \cos^3(x)}} \\
 &= \frac{2 \sec(x) \tan(x)}{7a\sqrt{a \cos^3(x)}} + \frac{\left(5 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\cos^{\frac{5}{2}}(x)} dx}{7a\sqrt{a \cos^3(x)}} \\
 &= \frac{10 \sin(x)}{21a\sqrt{a \cos^3(x)}} + \frac{2 \sec(x) \tan(x)}{7a\sqrt{a \cos^3(x)}} + \frac{\left(5 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\sqrt{\cos(x)}} dx}{21a\sqrt{a \cos^3(x)}} \\
 &= \frac{10 \cos^{\frac{3}{2}}(x) \text{EllipticF}\left(\frac{x}{2}, 2\right)}{21a\sqrt{a \cos^3(x)}} + \frac{10 \sin(x)}{21a\sqrt{a \cos^3(x)}} + \frac{2 \sec(x) \tan(x)}{7a\sqrt{a \cos^3(x)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.62

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \frac{2 \cos^2(x) \left(5 \cos^{\frac{5}{2}}(x) \text{EllipticF}\left(\frac{x}{2}, 2\right) + 5 \cos(x) \sin(x) + 3 \tan(x)\right)}{21 (a \cos^3(x))^{3/2}}$$

```
[In] Integrate[(a*Cos[x]^3)^(-3/2), x]
```

```
[Out] (2*Cos[x]^2*(5*Cos[x]^(5/2)*EllipticF[x/2, 2] + 5*Cos[x]*Sin[x] + 3*Tan[x])/(21*(a*Cos[x]^3)^(3/2))
```

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 5.05 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.42

method	result
default	$-\frac{2\left(5i\sqrt{\frac{\cos(x)}{\cos(x)+1}}\sqrt{\frac{1}{\cos(x)+1}}F(i(\csc(x)-\cot(x)),i)(\cos^2(x))+5i\sqrt{\frac{\cos(x)}{\cos(x)+1}}\sqrt{\frac{1}{\cos(x)+1}}F(i(\csc(x)-\cot(x)),i)\cos(x)-5\sin(x)-3\sec(x)\right)}{21a\sqrt{a(\cos^3(x))}}$

[In] `int(1/(a*cos(x)^3)^(3/2),x,method=_RETURNVERBOSE)`

[Out] `-2/21/a/(a*cos(x)^3)^(1/2)*(5*I*cos(x)^2*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)+5*I*cos(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)-5*sin(x)-3*sec(x)*tan(x))`

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.04

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \frac{5i \sqrt{2} \sqrt{a} \cos(x)^5 \text{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)) - 5i \sqrt{2} \sqrt{a} \cos(x)^5 \text{weierstrassPInverse}(-4, 0, \cos(x) - i \sin(x))}{21 a^2 \cos(x)^5}$$

[In] `integrate(1/(a*cos(x)^3)^(3/2),x, algorithm="fricas")`

[Out] `1/21*(5*I*sqrt(2)*sqrt(a)*cos(x)^5*weierstrassPInverse(-4, 0, cos(x) + I*sin(x)) - 5*I*sqrt(2)*sqrt(a)*cos(x)^5*weierstrassPInverse(-4, 0, cos(x) - I*sin(x)) + 2*sqrt(a*cos(x)^3)*(5*cos(x)^2 + 3)*sin(x))/(a^2*cos(x)^5)`

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \text{Timed out}$$

[In] `integrate(1/(a*cos(x)**3)**(3/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \int \frac{1}{(a \cos(x)^3)^{\frac{3}{2}}} dx$$

[In] integrate(1/(a*cos(x)^3)^(3/2),x, algorithm="maxima")

[Out] integrate((a*cos(x)^3)^(-3/2), x)

Giac [F]

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \int \frac{1}{(a \cos(x)^3)^{\frac{3}{2}}} dx$$

[In] integrate(1/(a*cos(x)^3)^(3/2),x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(-3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^3(x))^{3/2}} dx = \int \frac{1}{(a \cos(x)^3)^{3/2}} dx$$

[In] int(1/(a*cos(x)^3)^(3/2),x)

[Out] int(1/(a*cos(x)^3)^(3/2), x)

3.50 $\int \frac{1}{(a \cos^3(x))^{5/2}} dx$

Optimal result	287
Rubi [A] (verified)	287
Mathematica [A] (verified)	289
Maple [C] (verified)	289
Fricas [C] (verification not implemented)	290
Sympy [F(-1)]	290
Maxima [F]	290
Giac [F]	291
Mupad [F(-1)]	291

Optimal result

Integrand size = 10, antiderivative size = 117

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = -\frac{154 \cos^{3/2}(x) E\left(\frac{x}{2} \middle| 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \cos(x) \sin(x)}{195a^2 \sqrt{a \cos^3(x)}} \\ + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}}$$

[Out] $-154/195*\cos(x)^{(3/2)}*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\text{EllipticE}(\sin(1/2*x), 2^{(1/2)})/a^2/(a*\cos(x)^3)^{(1/2)}+154/195*\cos(x)*\sin(x)/a^2/(a*\cos(x)^3)^{(1/2)}+154/585*\tan(x)/a^2/(a*\cos(x)^3)^{(1/2)}+22/117*\sec(x)^2*\tan(x)/a^2/(a*\cos(x)^3)^{(1/2)}+2/13*\sec(x)^4*\tan(x)/a^2/(a*\cos(x)^3)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2716, 2719}

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \frac{154 \sin(x) \cos(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} \\ - \frac{154 \cos^{3/2}(x) E\left(\frac{x}{2} \middle| 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{2 \tan(x) \sec^4(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{22 \tan(x) \sec^2(x)}{117a^2 \sqrt{a \cos^3(x)}}$$

[In] Int[(a*cos[x]^3)^(-5/2),x]

[Out] $(-154*\cos[x]^{(3/2)}*\text{EllipticE}[x/2, 2])/((195*a^2*\text{Sqrt}[a*\cos[x]^3]) + (154*\cos[x]*\sin[x])/(195*a^2*\text{Sqrt}[a*\cos[x]^3]) + (154*\tan[x])/(585*a^2*\text{Sqrt}[a*\cos[x]$

]^3)) + (22*Sec[x]^2*Tan[x])/(117*a^2*Sqrt[a*Cos[x]^3]) + (2*Sec[x]^4*Tan[x])/(13*a^2*Sqrt[a*Cos[x]^3])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3286

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x])^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\cos^{\frac{3}{2}}(x) \int \frac{1}{\cos^{\frac{15}{2}}(x)} dx}{a^2 \sqrt{a \cos^3(x)}} \\
 &= \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(11 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\cos^{\frac{11}{2}}(x)} dx}{13a^2 \sqrt{a \cos^3(x)}} \\
 &= \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(77 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\cos^{\frac{7}{2}}(x)} dx}{117a^2 \sqrt{a \cos^3(x)}} \\
 &= \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} + \frac{\left(77 \cos^{\frac{3}{2}}(x)\right) \int \frac{1}{\cos^{\frac{3}{2}}(x)} dx}{195a^2 \sqrt{a \cos^3(x)}} \\
 &= \frac{154 \cos(x) \sin(x)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} \\
 &\quad + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}} - \frac{\left(77 \cos^{\frac{3}{2}}(x)\right) \int \sqrt{\cos(x)} dx}{195a^2 \sqrt{a \cos^3(x)}}
 \end{aligned}$$

$$= -\frac{154 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right)}{195a^2 \sqrt{a \cos^3(x)}} + \frac{154 \cos(x) \sin(x)}{195a^2 \sqrt{a \cos^3(x)}} \\ + \frac{154 \tan(x)}{585a^2 \sqrt{a \cos^3(x)}} + \frac{22 \sec^2(x) \tan(x)}{117a^2 \sqrt{a \cos^3(x)}} + \frac{2 \sec^4(x) \tan(x)}{13a^2 \sqrt{a \cos^3(x)}}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.49

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \frac{-462 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \mid 2\right) + 462 \cos(x) \sin(x) + 2(77 + 55 \sec^2(x) + 45 \sec^4(x)) \tan(x)}{585a^2 \sqrt{a \cos^3(x)}}$$

[In] Integrate[(a*Cos[x]^3)^(-5/2),x]

[Out] (-462*Cos[x]^(3/2)*EllipticE[x/2, 2] + 462*Cos[x]*Sin[x] + 2*(77 + 55*Sec[x]^2 + 45*Sec[x]^4)*Tan[x])/(585*a^2*Sqrt[a*Cos[x]^3])

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 8.31 (sec) , antiderivative size = 297, normalized size of antiderivative = 2.54

method	result
default	$-\frac{2i(231(\cos^3(x))\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}E(i(\csc(x)-\cot(x)),i)-231(\cos^3(x))\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}F(i(\csc(x)-\cot(x)),i)+462(\cos(x)\sin(x)+\tan(x)(77+55\sec^2(x)+45\sec^4(x))))}{585a^2\sqrt{a\cos^3(x)}}$

[In] int(1/(a*cos(x)^3)^(5/2),x,method=_RETURNVERBOSE)

[Out] -2/585*I/(cos(x)+1)/a^2/(a*cos(x)^3)^(1/2)*(231*cos(x)^3*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)-231*cos(x)^3*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)+462*cos(x)^2*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)-462*cos(x)^2*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)+231*cos(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(csc(x)-cot(x)),I)-231*cos(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(csc(x)-cot(x)),I)+231*I*cos(x)*sin(x)+77*I*sin(x)+77*I*tan(x)+55*I*sec(x)*tan(x)+55*I*tan(x)*sec(x)^2+45*I*tan(x)*sec(x)^3+45*I*tan(x)*sec(x)^4)

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 92, normalized size of antiderivative = 0.79

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \frac{231i \sqrt{2} \sqrt{a} \cos(x)^8 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(x) + i \sin(x)))}{(a \cos^3(x))^{5/2}}$$

[In] integrate(1/(a*cos(x)^3)^(5/2),x, algorithm="fricas")

[Out] 1/585*(231*I*sqrt(2)*sqrt(a)*cos(x)^8*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) + I*sin(x))) - 231*I*sqrt(2)*sqrt(a)*cos(x)^8*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(x) - I*sin(x))) + 2*(231*cos(x)^6 + 77*cos(x)^4 + 55*cos(x)^2 + 45)*sqrt(a*cos(x)^3)*sin(x))/(a^3*cos(x)^8)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \text{Timed out}$$

[In] integrate(1/(a*cos(x)**3)**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \int \frac{1}{(a \cos(x)^3)^{5/2}} dx$$

[In] integrate(1/(a*cos(x)^3)^(5/2),x, algorithm="maxima")

[Out] integrate((a*cos(x)^3)^(-5/2), x)

Giac [F]

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \int \frac{1}{(a \cos(x)^3)^{5/2}} dx$$

[In] integrate(1/(a*cos(x)^3)^(5/2),x, algorithm="giac")

[Out] integrate((a*cos(x)^3)^(-5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^3(x))^{5/2}} dx = \int \frac{1}{(a \cos(x)^3)^{5/2}} dx$$

[In] int(1/(a*cos(x)^3)^(5/2),x)

[Out] int(1/(a*cos(x)^3)^(5/2), x)

3.51 $\int (a \cos^4(x))^{5/2} dx$

Optimal result	292
Rubi [A] (verified)	292
Mathematica [A] (verified)	294
Maple [A] (verified)	294
Fricas [A] (verification not implemented)	295
Sympy [F(-1)]	295
Maxima [A] (verification not implemented)	295
Giac [A] (verification not implemented)	296
Mupad [F(-1)]	296

Optimal result

Integrand size = 10, antiderivative size = 132

$$\begin{aligned} \int (a \cos^4(x))^{5/2} dx &= \frac{63}{256} a^2 x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{21}{128} a^2 \cos(x) \sqrt{a \cos^4(x)} \sin(x) \\ &+ \frac{21}{160} a^2 \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{9}{80} a^2 \cos^5(x) \sqrt{a \cos^4(x)} \sin(x) \\ &+ \frac{1}{10} a^2 \cos^7(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{63}{256} a^2 \sqrt{a \cos^4(x)} \tan(x) \end{aligned}$$

[Out] $63/256*a^2*x*\sec(x)^2*(a*\cos(x)^4)^{(1/2)}+21/128*a^2*\cos(x)*\sin(x)*(a*\cos(x)^4)^{(1/2)}+21/160*a^2*\cos(x)^3*\sin(x)*(a*\cos(x)^4)^{(1/2)}+9/80*a^2*\cos(x)^5*\sin(x)*(a*\cos(x)^4)^{(1/2)}+1/10*a^2*\cos(x)^7*\sin(x)*(a*\cos(x)^4)^{(1/2)}+63/256*a^2*(a*\cos(x)^4)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 8}

$$\begin{aligned} \int (a \cos^4(x))^{5/2} dx &= \frac{21}{128} a^2 \sin(x) \cos(x) \sqrt{a \cos^4(x)} + \frac{63}{256} a^2 \tan(x) \sqrt{a \cos^4(x)} \\ &+ \frac{63}{256} a^2 x \sec^2(x) \sqrt{a \cos^4(x)} + \frac{1}{10} a^2 \sin(x) \cos^7(x) \sqrt{a \cos^4(x)} \\ &+ \frac{9}{80} a^2 \sin(x) \cos^5(x) \sqrt{a \cos^4(x)} + \frac{21}{160} a^2 \sin(x) \cos^3(x) \sqrt{a \cos^4(x)} \end{aligned}$$

[In] Int[(a*cos[x]^4)^(5/2),x]

[Out] $(63*a^2*x*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sec}[x]^2)/256 + (21*a^2*\text{Cos}[x]*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sin}[x])/128 + (21*a^2*\text{Cos}[x]^3*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sin}[x])/160 + (9*a^2*\text{Cos}[x]$

$$\begin{aligned} & ^5\sqrt{a\cos[x]^4\sin[x]}/80 + (a^2\cos[x]^7\sqrt{a\cos[x]^4\sin[x]})/10 \\ & + (63a^2\sqrt{a\cos[x]^4}\tan[x])/256 \end{aligned}$$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3286

Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[SIN[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*(b*SIN[e + f*x]^n)^FracPart[p]/(SIN[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(SIN[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned} \text{integral} &= \left(a^2\sqrt{a\cos^4(x)}\sec^2(x)\right)\int\cos^{10}(x)dx \\ &= \frac{1}{10}a^2\cos^7(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{10}\left(9a^2\sqrt{a\cos^4(x)}\sec^2(x)\right)\int\cos^8(x)dx \\ &= \frac{9}{80}a^2\cos^5(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{10}a^2\cos^7(x)\sqrt{a\cos^4(x)}\sin(x) \\ &\quad + \frac{1}{80}\left(63a^2\sqrt{a\cos^4(x)}\sec^2(x)\right)\int\cos^6(x)dx \\ &= \frac{21}{160}a^2\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{9}{80}a^2\cos^5(x)\sqrt{a\cos^4(x)}\sin(x) \\ &\quad + \frac{1}{10}a^2\cos^7(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{32}\left(21a^2\sqrt{a\cos^4(x)}\sec^2(x)\right)\int\cos^4(x)dx \\ &= \frac{21}{128}a^2\cos(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{21}{160}a^2\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) \\ &\quad + \frac{9}{80}a^2\cos^5(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{10}a^2\cos^7(x)\sqrt{a\cos^4(x)}\sin(x) \\ &\quad + \frac{1}{128}\left(63a^2\sqrt{a\cos^4(x)}\sec^2(x)\right)\int\cos^2(x)dx \end{aligned}$$

$$\begin{aligned}
&= \frac{21}{128}a^2 \cos(x)\sqrt{a \cos^4(x)} \sin(x) + \frac{21}{160}a^2 \cos^3(x)\sqrt{a \cos^4(x)} \sin(x) \\
&\quad + \frac{9}{80}a^2 \cos^5(x)\sqrt{a \cos^4(x)} \sin(x) + \frac{1}{10}a^2 \cos^7(x)\sqrt{a \cos^4(x)} \sin(x) \\
&\quad + \frac{63}{256}a^2 \sqrt{a \cos^4(x)} \tan(x) + \frac{1}{256} \left(63a^2 \sqrt{a \cos^4(x)} \sec^2(x) \right) \int 1 dx \\
&= \frac{63}{256}a^2 x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{21}{128}a^2 \cos(x)\sqrt{a \cos^4(x)} \sin(x) \\
&\quad + \frac{21}{160}a^2 \cos^3(x)\sqrt{a \cos^4(x)} \sin(x) + \frac{9}{80}a^2 \cos^5(x)\sqrt{a \cos^4(x)} \sin(x) \\
&\quad + \frac{1}{10}a^2 \cos^7(x)\sqrt{a \cos^4(x)} \sin(x) + \frac{63}{256}a^2 \sqrt{a \cos^4(x)} \tan(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.40

$$\int (a \cos^4(x))^{5/2} dx = \frac{a(a \cos^4(x))^{3/2} \sec^6(x)(2520x + 2100 \sin(2x) + 600 \sin(4x) + 150 \sin(6x) + 25 \sin(8x))}{10240}$$

[In] Integrate[(a*cos[x]^4)^(5/2),x]

[Out] (a*(a*cos[x]^4)^(3/2)*Sec[x]^6*(2520*x + 2100*Sin[2*x] + 600*Sin[4*x] + 150*Sin[6*x] + 25*Sin[8*x] + 2*Sin[10*x]))/10240

Maple [A] (verified)

Time = 13.26 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.42

method	result
default	$\frac{a^2 \sqrt{a(\cos^4(x))} (128(\cos^7(x)) \sin(x) + 144(\cos^5(x)) \sin(x) + 168(\cos^3(x)) \sin(x) + 210 \cos(x) \sin(x) + 315 \tan(x) + 315(\sec^2(x))x)}{1280}$
risch	$\frac{63a^2 e^{2ix} \sqrt{a(e^{2ix}+1)^4} e^{-4ix} x}{256(e^{2ix}+1)^2} - \frac{ia^2 e^{12ix} \sqrt{a(e^{2ix}+1)^4} e^{-4ix}}{10240(e^{2ix}+1)^2} - \frac{5ia^2 e^{10ix} \sqrt{a(e^{2ix}+1)^4} e^{-4ix}}{4096(e^{2ix}+1)^2} - \frac{105ia^2 e^{4ix} \sqrt{a(e^{2ix}+1)^4} e^{-4ix}}{1024(e^{2ix}+1)^2} +$

[In] int((a*cos(x)^4)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/1280*a^2*(a*cos(x)^4)^(1/2)*(128*cos(x)^7*sin(x)+144*cos(x)^5*sin(x)+168*cos(x)^3*sin(x)+210*cos(x)*sin(x)+315*tan(x)+315*sec(x)^2*x)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.52

$$\int (a \cos^4(x))^{5/2} dx = \frac{\sqrt{a \cos(x)^4} (315 a^2 x + (128 a^2 \cos(x)^9 + 144 a^2 \cos(x)^7 + 168 a^2 \cos(x)^5 + 210 a^2 \cos(x)^3 + 315 a^2 \cos(x)) \sin(x))}{1280 \cos(x)^2}$$

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="fricas")

[Out] 1/1280*sqrt(a*cos(x)^4)*(315*a^2*x + (128*a^2*cos(x)^9 + 144*a^2*cos(x)^7 + 168*a^2*cos(x)^5 + 210*a^2*cos(x)^3 + 315*a^2*cos(x))*sin(x))/cos(x)^2

Sympy [F(-1)]

Timed out.

$$\int (a \cos^4(x))^{5/2} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**4)**(5/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.64

$$\int (a \cos^4(x))^{5/2} dx = \frac{63}{256} a^{\frac{5}{2}} x + \frac{315 a^{\frac{5}{2}} \tan(x)^9 + 1470 a^{\frac{5}{2}} \tan(x)^7 + 2688 a^{\frac{5}{2}} \tan(x)^5 + 2370 a^{\frac{5}{2}} \tan(x)^3 + 965 a^{\frac{5}{2}} \tan(x)}{1280 (\tan(x)^{10} + 5 \tan(x)^8 + 10 \tan(x)^6 + 10 \tan(x)^4 + 5 \tan(x)^2 + 1)}$$

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="maxima")

[Out] 63/256*a^(5/2)*x + 1/1280*(315*a^(5/2)*tan(x)^9 + 1470*a^(5/2)*tan(x)^7 + 2688*a^(5/2)*tan(x)^5 + 2370*a^(5/2)*tan(x)^3 + 965*a^(5/2)*tan(x))/(tan(x)^10 + 5*tan(x)^8 + 10*tan(x)^6 + 10*tan(x)^4 + 5*tan(x)^2 + 1)

Giac [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.43

$$\int (a \cos^4(x))^{5/2} dx = \frac{1}{10240} (2520 a^2 x + 2 a^2 \sin(10x) + 25 a^2 \sin(8x) + 150 a^2 \sin(6x) + 600 a^2 \sin(4x) + \dots)$$

[In] integrate((a*cos(x)^4)^(5/2),x, algorithm="giac")

[Out] 1/10240*(2520*a^2*x + 2*a^2*sin(10*x) + 25*a^2*sin(8*x) + 150*a^2*sin(6*x) + 600*a^2*sin(4*x) + 2100*a^2*sin(2*x))*sqrt(a)

Mupad [F(-1)]

Timed out.

$$\int (a \cos^4(x))^{5/2} dx = \int (a \cos(x)^4)^{5/2} dx$$

[In] int((a*cos(x)^4)^(5/2),x)

[Out] int((a*cos(x)^4)^(5/2), x)

3.52 $\int (a \cos^4(x))^{3/2} dx$

Optimal result	297
Rubi [A] (verified)	297
Mathematica [A] (verified)	299
Maple [A] (verified)	299
Fricas [A] (verification not implemented)	299
Sympy [F(-1)]	300
Maxima [A] (verification not implemented)	300
Giac [A] (verification not implemented)	300
Mupad [F(-1)]	300

Optimal result

Integrand size = 10, antiderivative size = 78

$$\int (a \cos^4(x))^{3/2} dx = \frac{5}{16}ax \sqrt{a \cos^4(x)} \sec^2(x) + \frac{5}{24}a \cos(x) \sqrt{a \cos^4(x)} \sin(x) \\ + \frac{1}{6}a \cos^3(x) \sqrt{a \cos^4(x)} \sin(x) + \frac{5}{16}a \sqrt{a \cos^4(x)} \tan(x)$$

[Out] $5/16*a*x*\sec(x)^2*(a*\cos(x)^4)^{(1/2)}+5/24*a*\cos(x)*\sin(x)*(a*\cos(x)^4)^{(1/2)}+1/6*a*\cos(x)^3*\sin(x)*(a*\cos(x)^4)^{(1/2)}+5/16*a*(a*\cos(x)^4)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 8}

$$\int (a \cos^4(x))^{3/2} dx = \frac{5}{24}a \sin(x) \cos(x) \sqrt{a \cos^4(x)} + \frac{5}{16}a \tan(x) \sqrt{a \cos^4(x)} \\ + \frac{5}{16}ax \sec^2(x) \sqrt{a \cos^4(x)} + \frac{1}{6}a \sin(x) \cos^3(x) \sqrt{a \cos^4(x)}$$

[In] Int[(a*Cos[x]^4)^(3/2),x]

[Out] $(5*a*x*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sec}[x]^2)/16 + (5*a*\text{Cos}[x]*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sin}[x])/24 + (a*\text{Cos}[x]^3*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Sin}[x])/6 + (5*a*\text{Sqrt}[a*\text{Cos}[x]^4]*\text{Tan}[x])/16$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_)^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \left(a\sqrt{a\cos^4(x)}\sec^2(x) \right) \int \cos^6(x) dx \\
&= \frac{1}{6}a\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{6}\left(5a\sqrt{a\cos^4(x)}\sec^2(x) \right) \int \cos^4(x) dx \\
&= \frac{5}{24}a\cos(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{6}a\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) \\
&\quad + \frac{1}{8}\left(5a\sqrt{a\cos^4(x)}\sec^2(x) \right) \int \cos^2(x) dx \\
&= \frac{5}{24}a\cos(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{1}{6}a\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) \\
&\quad + \frac{5}{16}a\sqrt{a\cos^4(x)}\tan(x) + \frac{1}{16}\left(5a\sqrt{a\cos^4(x)}\sec^2(x) \right) \int 1 dx \\
&= \frac{5}{16}ax\sqrt{a\cos^4(x)}\sec^2(x) + \frac{5}{24}a\cos(x)\sqrt{a\cos^4(x)}\sin(x) \\
&\quad + \frac{1}{6}a\cos^3(x)\sqrt{a\cos^4(x)}\sin(x) + \frac{5}{16}a\sqrt{a\cos^4(x)}\tan(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.49

$$\int (a \cos^4(x))^{3/2} dx = \frac{1}{192} (a \cos^4(x))^{3/2} \sec^6(x) (60x + 45 \sin(2x) + 9 \sin(4x) + \sin(6x))$$

`[In] Integrate[(a*Cos[x]^4)^(3/2),x]``[Out] ((a*Cos[x]^4)^(3/2)*Sec[x]^6*(60*x + 45*Sin[2*x] + 9*Sin[4*x] + Sin[6*x]))/192`**Maple [A] (verified)**

Time = 0.82 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.49

method	result
default	$\frac{a \sqrt{a(\cos^4(x))} (8(\cos^3(x)) \sin(x) + 10 \cos(x) \sin(x) + 15 \tan(x) + 15(\sec^2(x))x)}{48}$
risch	$\frac{5a e^{2ix} \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{16(e^{2ix}+1)^2} - \frac{ia e^{8ix} \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{384(e^{2ix}+1)^2} - \frac{3ia e^{6ix} \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{128(e^{2ix}+1)^2} - \frac{15ia e^{4ix} \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{128(e^{2ix}+1)^2} + \frac{15ia e^{2ix} \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{128(e^{2ix}+1)^2}$

`[In] int((a*cos(x)^4)^(3/2),x,method=_RETURNVERBOSE)``[Out] 1/48*a*(a*cos(x)^4)^(1/2)*(8*cos(x)^3*sin(x)+10*cos(x)*sin(x)+15*tan(x)+15*sec(x)^2*x)`**Fricas [A] (verification not implemented)**

none

Time = 0.29 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.54

$$\int (a \cos^4(x))^{3/2} dx = \frac{\sqrt{a \cos^4(x)} (15ax + (8a \cos(x)^5 + 10a \cos(x)^3 + 15a \cos(x)) \sin(x))}{48 \cos^2(x)}$$

`[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="fricas")``[Out] 1/48*sqrt(a*cos(x)^4)*(15*a*x + (8*a*cos(x)^5 + 10*a*cos(x)^3 + 15*a*cos(x))*sin(x))/cos(x)^2`

Sympy [F(-1)]

Timed out.

$$\int (a \cos^4(x))^{3/2} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**4)**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.71

$$\int (a \cos^4(x))^{3/2} dx = \frac{5}{16} a^{\frac{3}{2}} x + \frac{15 a^{\frac{3}{2}} \tan(x)^5 + 40 a^{\frac{3}{2}} \tan(x)^3 + 33 a^{\frac{3}{2}} \tan(x)}{48 (\tan(x)^6 + 3 \tan(x)^4 + 3 \tan(x)^2 + 1)}$$

[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="maxima")

[Out] 5/16*a^(3/2)*x + 1/48*(15*a^(3/2)*tan(x)^5 + 40*a^(3/2)*tan(x)^3 + 33*a^(3/2)*tan(x))/(tan(x)^6 + 3*tan(x)^4 + 3*tan(x)^2 + 1)

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.32

$$\int (a \cos^4(x))^{3/2} dx = \frac{1}{192} a^{\frac{3}{2}} (60x + \sin(6x) + 9 \sin(4x) + 45 \sin(2x))$$

[In] integrate((a*cos(x)^4)^(3/2),x, algorithm="giac")

[Out] 1/192*a^(3/2)*(60*x + sin(6*x) + 9*sin(4*x) + 45*sin(2*x))

Mupad [F(-1)]

Timed out.

$$\int (a \cos^4(x))^{3/2} dx = \int (a \cos(x)^4)^{3/2} dx$$

[In] int((a*cos(x)^4)^(3/2),x)

[Out] int((a*cos(x)^4)^(3/2), x)

3.53 $\int \sqrt{a \cos^4(x)} dx$

Optimal result	301
Rubi [A] (verified)	301
Mathematica [A] (verified)	302
Maple [A] (verified)	302
Fricas [A] (verification not implemented)	303
Sympy [F(-1)]	303
Maxima [A] (verification not implemented)	303
Giac [A] (verification not implemented)	303
Mupad [F(-1)]	304

Optimal result

Integrand size = 10, antiderivative size = 36

$$\int \sqrt{a \cos^4(x)} dx = \frac{1}{2} x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{1}{2} \sqrt{a \cos^4(x)} \tan(x)$$

[Out] $1/2*x*\sec(x)^2*(a*\cos(x)^4)^{(1/2)}+1/2*(a*\cos(x)^4)^{(1/2)}*\tan(x)$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 2715, 8}

$$\int \sqrt{a \cos^4(x)} dx = \frac{1}{2} \tan(x) \sqrt{a \cos^4(x)} + \frac{1}{2} x \sec^2(x) \sqrt{a \cos^4(x)}$$

[In] Int[Sqrt[a*Cos[x]^4], x]

[Out] (x*Sqrt[a*Cos[x]^4]*Sec[x]^2)/2 + (Sqrt[a*Cos[x]^4]*Tan[x])/2

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \left(\sqrt{a \cos^4(x)} \sec^2(x) \right) \int \cos^2(x) dx \\ &= \frac{1}{2} \sqrt{a \cos^4(x)} \tan(x) + \frac{1}{2} \left(\sqrt{a \cos^4(x)} \sec^2(x) \right) \int 1 dx \\ &= \frac{1}{2} x \sqrt{a \cos^4(x)} \sec^2(x) + \frac{1}{2} \sqrt{a \cos^4(x)} \tan(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.69

$$\int \sqrt{a \cos^4(x)} dx = \frac{1}{2} \sqrt{a \cos^4(x)} \sec^2(x) (x + \cos(x) \sin(x))$$

[In] Integrate[Sqrt[a*Cos[x]^4], x]

[Out] (Sqrt[a*Cos[x]^4]*Sec[x]^2*(x + Cos[x]*Sin[x]))/2

Maple [A] (verified)

Time = 1.01 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

method	result	size
default	$\frac{\sqrt{a(\cos^4(x))} (\tan(x) + (\sec^2(x))x)}{2}$	20
risch	$\frac{\sqrt{a(e^{2ix}+1)^4 e^{-4ix}} e^{2ix} x}{2(e^{2ix}+1)^2} - \frac{i\sqrt{a(e^{2ix}+1)^4 e^{-4ix}} e^{4ix}}{8(e^{2ix}+1)^2} + \frac{i\sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}{8(e^{2ix}+1)^2}$	102

[In] int((a*cos(x)^4)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/2*(a*cos(x)^4)^(1/2)*(tan(x)+sec(x)^2*x)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.58

$$\int \sqrt{a \cos^4(x)} dx = \frac{\sqrt{a \cos(x)^4} (\cos(x) \sin(x) + x)}{2 \cos(x)^2}$$

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="fricas")

[Out] 1/2*sqrt(a*cos(x)^4)*(cos(x)*sin(x) + x)/cos(x)^2

Sympy [F(-1)]

Timed out.

$$\int \sqrt{a \cos^4(x)} dx = \text{Timed out}$$

[In] integrate((a*cos(x)**4)**(1/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.47 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.61

$$\int \sqrt{a \cos^4(x)} dx = \frac{1}{2} \sqrt{ax} + \frac{\sqrt{a} \tan(x)}{2 (\tan(x)^2 + 1)}$$

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(a)*x + 1/2*sqrt(a)*tan(x)/(tan(x)^2 + 1)

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.36

$$\int \sqrt{a \cos^4(x)} dx = \frac{1}{4} \sqrt{a} (2x + \sin(2x))$$

[In] integrate((a*cos(x)^4)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(a)*(2*x + sin(2*x))

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a \cos^4(x)} dx = \int \sqrt{a \cos(x)^4} dx$$

```
[In] int((a*cos(x)^4)^(1/2),x)
```

```
[Out] int((a*cos(x)^4)^(1/2), x)
```


3.54 $\int \frac{1}{\sqrt{a \cos^4(x)}} dx$

Optimal result	305
Rubi [A] (verified)	305
Mathematica [A] (verified)	306
Maple [A] (verified)	306
Fricas [A] (verification not implemented)	307
Sympy [F(-1)]	307
Maxima [A] (verification not implemented)	307
Giac [A] (verification not implemented)	307
Mupad [B] (verification not implemented)	308

Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\cos(x) \sin(x)}{\sqrt{a \cos^4(x)}}$$

[Out] $\cos(x) \sin(x) / (a \cos(x)^4)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3286, 3852, 8}

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\sin(x) \cos(x)}{\sqrt{a \cos^4(x)}}$$

[In] $\text{Int}[1/\text{Sqrt}[a \cos[x]^4], x]$

[Out] $(\cos[x] \sin[x]) / \text{Sqrt}[a \cos[x]^4]$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3286

$\text{Int}[(u_.) * ((b_.) \sin[(e_.) + (f_.) (x_)]^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\sin[e + f*x], x]\}, \text{Dist}[(b*ff^n)^{\text{IntPart}[p]} * (b \sin[e + f*x]^n)^{\text{FracPart}[p]} / (\sin[e + f*x]/ff)^{(n*\text{FracPart}[p])}], \text{Int}[\text{ActivateTrig}[u] * (\sin[e + f*x]/ff)^{(n*p)}, x], x]\} /; \text{FreeQ}\{\{b, e, f, n, p\}, x\} \&\& \text{!IntegerQ}[p] \&\& \text{IntegerQ}[n] \&\& (\text{EqQ}[u, 1] \parallel \text{MatchQ}[u, ((d_.) * (\text{trig}_)[e + f*x])^{(m_)}] /;$

```
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]]
```

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^2(x) \int \sec^2(x) dx}{\sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}(\int 1 dx, x, -\tan(x))}{\sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{\sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\cos(x) \sin(x)}{\sqrt{a \cos^4(x)}}$$

```
[In] Integrate[1/Sqrt[a*Cos[x]^4], x]
```

```
[Out] (Cos[x]*Sin[x])/Sqrt[a*Cos[x]^4]
```

Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
default	$\frac{\cos(x) \sin(x)}{\sqrt{a(\cos^4(x))}}$	14
risch	$\frac{2i(1+e^{-2ix})}{\sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}$	29

```
[In] int(1/(a*cos(x)^4)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] cos(x)*sin(x)/(a*cos(x)^4)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.20

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\sqrt{a \cos(x)^4} \sin(x)}{a \cos(x)^3}$$

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="fricas")

[Out] sqrt(a*cos(x)^4)*sin(x)/(a*cos(x)^3)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \text{Timed out}$$

[In] integrate(1/(a*cos(x)**4)**(1/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\tan(x)}{\sqrt{a}}$$

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="maxima")

[Out] tan(x)/sqrt(a)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\tan(x)}{\sqrt{a}}$$

[In] integrate(1/(a*cos(x)^4)^(1/2),x, algorithm="giac")

[Out] tan(x)/sqrt(a)

Mupad [B] (verification not implemented)

Time = 14.20 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.40

$$\int \frac{1}{\sqrt{a \cos^4(x)}} dx = \frac{\tan(x)}{\sqrt{a}}$$

[In] `int(1/(a*cos(x)^4)^(1/2),x)`

[Out] `tan(x)/a^(1/2)`

$$3.55 \quad \int \frac{1}{(a \cos^4(x))^{3/2}} dx$$

Optimal result	309
Rubi [A] (verified)	309
Mathematica [A] (verified)	310
Maple [A] (verified)	310
Fricas [A] (verification not implemented)	311
Sympy [F(-1)]	311
Maxima [A] (verification not implemented)	311
Giac [A] (verification not implemented)	311
Mupad [B] (verification not implemented)	312

Optimal result

Integrand size = 10, antiderivative size = 67

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{\cos(x) \sin(x)}{a \sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a \sqrt{a \cos^4(x)}}$$

[Out] $\cos(x) \sin(x) / a / (a \cos(x)^4)^{(1/2)} + 2/3 \sin(x)^2 \tan(x) / a / (a \cos(x)^4)^{(1/2)} + 1/5 \sin(x)^2 \tan(x)^3 / a / (a \cos(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3286, 3852}

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{\sin(x) \cos(x)}{a \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a \sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a \sqrt{a \cos^4(x)}}$$

[In] $\text{Int}[(a \cos[x]^4)^{-3/2}, x]$

[Out] $(\cos[x] \sin[x]) / (a \sqrt{a \cos[x]^4}) + (2 \sin[x]^2 \tan[x]) / (3 a \sqrt{a \cos[x]^4}) + (\sin[x]^2 \tan[x]^3) / (5 a \sqrt{a \cos[x]^4})$

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
```

```
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]]
```

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^2(x) \int \sec^6(x) dx}{a \sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}\left(\int (1 + 2x^2 + x^4) dx, x, -\tan(x)\right)}{a \sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{a \sqrt{a \cos^4(x)}} + \frac{2 \sin^2(x) \tan(x)}{3a \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^3(x)}{5a \sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.45

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{\cos(x)(8 + 6 \cos(2x) + \cos(4x)) \sin(x)}{15 (a \cos^4(x))^{3/2}}$$

```
[In] Integrate[(a*Cos[x]^4)^(-3/2), x]
```

```
[Out] (Cos[x]*(8 + 6*Cos[2*x] + Cos[4*x])*Sin[x])/(15*(a*Cos[x]^4)^(3/2))
```

Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.49

method	result	size
default	$\frac{8 \cos(x) \sin(x) + 4 \tan(x) + 3 \tan(x) (\sec^2(x))}{15a \sqrt{a(\cos^4(x))}}$	33
risch	$\frac{16i(5+11 \cos(2x)+9i \sin(2x))}{15a(e^{2ix}+1)^3 \sqrt{a(e^{2ix}+1)^4 e^{-4ix}}}$	49

```
[In] int(1/(a*cos(x)^4)^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/15/a/(a*cos(x)^4)^(1/2)*(8*cos(x)*sin(x)+4*tan(x)+3*tan(x)*sec(x)^2)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.49

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{\sqrt{a \cos^4(x)} (8 \cos^4(x) + 4 \cos^2(x) + 3) \sin(x)}{15 a^2 \cos^7(x)}$$

[In] integrate(1/(a*cos(x)^4)^(3/2),x, algorithm="fricas")

[Out] 1/15*sqrt(a*cos(x)^4)*(8*cos(x)^4 + 4*cos(x)^2 + 3)*sin(x)/(a^2*cos(x)^7)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \text{Timed out}$$

[In] integrate(1/(a*cos(x)**4)**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.33

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{3 \tan^5(x) + 10 \tan^3(x) + 15 \tan(x)}{15 a^{3/2}}$$

[In] integrate(1/(a*cos(x)^4)^(3/2),x, algorithm="maxima")

[Out] 1/15*(3*tan(x)^5 + 10*tan(x)^3 + 15*tan(x))/a^(3/2)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.33

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{3 \tan^5(x) + 10 \tan^3(x) + 15 \tan(x)}{15 a^{3/2}}$$

[In] integrate(1/(a*cos(x)^4)^(3/2),x, algorithm="giac")

[Out] 1/15*(3*tan(x)^5 + 10*tan(x)^3 + 15*tan(x))/a^(3/2)

Mupad [B] (verification not implemented)

Time = 14.81 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.54

$$\int \frac{1}{(a \cos^4(x))^{3/2}} dx = \frac{4 \sin(x)}{5 a^{3/2} \cos(x)^3} + \frac{\sin(x)}{5 a^{3/2} \cos(x)^5} - \frac{8 \sin(x)^3}{15 a^{3/2} \cos(x)^3}$$

[In] int(1/(a*cos(x)^4)^(3/2),x)

[Out] (4*sin(x))/(5*a^(3/2)*cos(x)^3) + sin(x)/(5*a^(3/2)*cos(x)^5) - (8*sin(x)^3)/(15*a^(3/2)*cos(x)^3)

3.56 $\int \frac{1}{(a \cos^4(x))^{5/2}} dx$

Optimal result	313
Rubi [A] (verified)	313
Mathematica [A] (verified)	314
Maple [A] (verified)	315
Fricas [A] (verification not implemented)	315
Sympy [F(-1)]	315
Maxima [A] (verification not implemented)	316
Giac [A] (verification not implemented)	316
Mupad [B] (verification not implemented)	316

Optimal result

Integrand size = 10, antiderivative size = 117

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{\cos(x) \sin(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}}$$

[Out] $\cos(x) \sin(x) / a^2 / (a \cos(x)^4)^{(1/2)} + 4/3 \sin(x)^2 \tan(x) / a^2 / (a \cos(x)^4)^{(1/2)} + 6/5 \sin(x)^2 \tan(x)^3 / a^2 / (a \cos(x)^4)^{(1/2)} + 4/7 \sin(x)^2 \tan(x)^5 / a^2 / (a \cos(x)^4)^{(1/2)} + 1/9 \sin(x)^2 \tan(x)^7 / a^2 / (a \cos(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3286, 3852}

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{\sin(x) \cos(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}}$$

[In] Int[(a*cos[x]^4)^(-5/2), x]

[Out] (Cos[x]*Sin[x])/(a^2*Sqrt[a*Cos[x]^4]) + (4*Sin[x]^2*Tan[x])/(3*a^2*Sqrt[a*Cos[x]^4]) + (6*Sin[x]^2*Tan[x]^3)/(5*a^2*Sqrt[a*Cos[x]^4]) + (4*Sin[x]^2*Tan[x]^5)/(7*a^2*Sqrt[a*Cos[x]^4]) + (Sin[x]^2*Tan[x]^7)/(9*a^2*Sqrt[a*Cos[x]^4])

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIN[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expa
ndIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c,
d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^2(x) \int \sec^{10}(x) dx}{a^2 \sqrt{a \cos^4(x)}} \\ &= -\frac{\cos^2(x) \text{Subst}\left(\int (1 + 4x^2 + 6x^4 + 4x^6 + x^8) dx, x, -\tan(x)\right)}{a^2 \sqrt{a \cos^4(x)}} \\ &= \frac{\cos(x) \sin(x)}{a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan(x)}{3a^2 \sqrt{a \cos^4(x)}} + \frac{6 \sin^2(x) \tan^3(x)}{5a^2 \sqrt{a \cos^4(x)}} + \frac{4 \sin^2(x) \tan^5(x)}{7a^2 \sqrt{a \cos^4(x)}} + \frac{\sin^2(x) \tan^7(x)}{9a^2 \sqrt{a \cos^4(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.40

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{(128 + 130 \cos(2x) + 46 \cos(4x) + 10 \cos(6x) + \cos(8x)) \sec^6(x) \tan(x)}{315a^2 \sqrt{a \cos^4(x)}}$$

[In] Integrate[(a*cos[x]^4)^(-5/2), x]

[Out] ((128 + 130*cos[2*x] + 46*cos[4*x] + 10*cos[6*x] + Cos[8*x])*Sec[x]^6*Tan[x])/(315*a^2*Sqrt[a*cos[x]^4])

Maple [A] (verified)

Time = 0.96 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.39

method	result	size
default	$\frac{\tan(x) (\sec^6(x) (128 \cos^8(x) + 64 \cos^6(x) + 48 \cos^4(x) + 40 \cos^2(x) + 35))}{315 a^2 \sqrt{a \cos^4(x)}}$	46
risch	$\frac{256i (126 e^{6ix} + 84 e^{4ix} + 9 + 37 \cos(2x) + 35i \sin(2x))}{315 a^2 (e^{2ix} + 1)^7 \sqrt{a (e^{2ix} + 1)^4 e^{-4ix}}}$	63

[In] `int(1/(a*cos(x)^4)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $1/315 * \tan(x) * \sec(x)^6 * (128 * \cos(x)^8 + 64 * \cos(x)^6 + 48 * \cos(x)^4 + 40 * \cos(x)^2 + 35) / a^2 / (a * \cos(x)^4)^{1/2}$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.38

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{(128 \cos(x)^8 + 64 \cos(x)^6 + 48 \cos(x)^4 + 40 \cos(x)^2 + 35) \sqrt{a \cos(x)^4} \sin(x)}{315 a^3 \cos(x)^{11}}$$

[In] `integrate(1/(a*cos(x)^4)^(5/2),x, algorithm="fricas")`

[Out] $1/315 * (128 * \cos(x)^8 + 64 * \cos(x)^6 + 48 * \cos(x)^4 + 40 * \cos(x)^2 + 35) * \sqrt{a * \cos(x)^4} * \sin(x) / (a^3 * \cos(x)^{11})$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(1/(a*cos(x)**4)**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.29

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{35 \tan(x)^9 + 180 \tan(x)^7 + 378 \tan(x)^5 + 420 \tan(x)^3 + 315 \tan(x)}{315 a^{5/2}}$$

[In] integrate(1/(a*cos(x)^4)^(5/2),x, algorithm="maxima")

[Out] 1/315*(35*tan(x)^9 + 180*tan(x)^7 + 378*tan(x)^5 + 420*tan(x)^3 + 315*tan(x))/a^(5/2)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.29

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{35 \tan(x)^9 + 180 \tan(x)^7 + 378 \tan(x)^5 + 420 \tan(x)^3 + 315 \tan(x)}{315 a^{5/2}}$$

[In] integrate(1/(a*cos(x)^4)^(5/2),x, algorithm="giac")

[Out] 1/315*(35*tan(x)^9 + 180*tan(x)^7 + 378*tan(x)^5 + 420*tan(x)^3 + 315*tan(x))/a^(5/2)

Mupad [B] (verification not implemented)

Time = 18.03 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.62

$$\int \frac{1}{(a \cos^4(x))^{5/2}} dx = \frac{e^{x 4i} \sqrt{a \left(\frac{e^{-x 1i}}{2} + \frac{e^{x 1i}}{2}\right)^4} 2048i}{5 a^3 (e^{x 2i} + 1)^5 (e^{x 2i} + 2 e^{x 4i} + e^{x 6i})} - \frac{e^{x 4i} \sqrt{a \left(\frac{e^{-x 1i}}{2} + \frac{e^{x 1i}}{2}\right)^4} 4096i}{3 a^3 (e^{x 2i} + 1)^6 (e^{x 2i} + 2 e^{x 4i} + e^{x 6i})} + \frac{e^{x 4i} \sqrt{a \left(\frac{e^{-x 1i}}{2} + \frac{e^{x 1i}}{2}\right)^4} 12288i}{7 a^3 (e^{x 2i} + 1)^7 (e^{x 2i} + 2 e^{x 4i} + e^{x 6i})} - \frac{e^{x 4i} \sqrt{a \left(\frac{e^{-x 1i}}{2} + \frac{e^{x 1i}}{2}\right)^4} 1024i}{a^3 (e^{x 2i} + 1)^8 (e^{x 2i} + 2 e^{x 4i} + e^{x 6i})} + \frac{e^{x 4i} \sqrt{a \left(\frac{e^{-x 1i}}{2} + \frac{e^{x 1i}}{2}\right)^4} 2048i}{9 a^3 (e^{x 2i} + 1)^9 (e^{x 2i} + 2 e^{x 4i} + e^{x 6i})}$$

[In] int(1/(a*cos(x)^4)^(5/2),x)

[Out] (exp(x*4i)*(a*(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*2048i)/(5*a^3*(exp(x*2i) + 1)^5*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (exp(x*4i)*(a*(exp(-x*1i)

$$\begin{aligned}
& /2 + \exp(x*1i)/2)^4)^{(1/2)}*4096i)/(3*a^3*(\exp(x*2i) + 1)^6*(\exp(x*2i) + 2*\exp(x*4i) + \exp(x*6i))) + (\exp(x*4i)*(a*(\exp(-x*1i)/2 + \exp(x*1i)/2)^4)^{(1/2)}*12288i)/(7*a^3*(\exp(x*2i) + 1)^7*(\exp(x*2i) + 2*\exp(x*4i) + \exp(x*6i))) - \\
& (\exp(x*4i)*(a*(\exp(-x*1i)/2 + \exp(x*1i)/2)^4)^{(1/2)}*1024i)/(a^3*(\exp(x*2i) + 1)^8*(\exp(x*2i) + 2*\exp(x*4i) + \exp(x*6i))) + (\exp(x*4i)*(a*(\exp(-x*1i)/2 + \exp(x*1i)/2)^4)^{(1/2)}*2048i)/(9*a^3*(\exp(x*2i) + 1)^9*(\exp(x*2i) + 2*\exp(x*4i) + \exp(x*6i)))
\end{aligned}$$

3.57 $\int (b \cos^m(c + dx))^n dx$

Optimal result	318
Rubi [A] (verified)	318
Mathematica [A] (verified)	319
Maple [F]	319
Fricas [F]	320
Sympy [F]	320
Maxima [F]	320
Giac [F]	320
Mupad [F(-1)]	321

Optimal result

Integrand size = 12, antiderivative size = 78

$$\int (b \cos^m(c + dx))^n dx = \frac{\cos(c + dx) (b \cos^m(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + mn), \frac{1}{2}(3 + mn), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + mn) \sqrt{\sin^2(c + dx)}}$$

[Out] $-\cos(d*x+c)*(b*\cos(d*x+c)^m)^n*\operatorname{hypergeom}([1/2, 1/2*m*n+1/2], [1/2*m*n+3/2], c \cos(d*x+c)^2*\sin(d*x+c)/d/(m*n+1)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3287, 2722}

$$\int (b \cos^m(c + dx))^n dx = \frac{\sin(c + dx) \cos(c + dx) (b \cos^m(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(mn + 1), \frac{1}{2}(mn + 3), \cos^2(c + dx)\right)}{d(mn + 1) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x]^m)^n, x]$

[Out] $-\left(\operatorname{Cos}[c + d*x]*(b*\operatorname{Cos}[c + d*x]^m)^n*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1 + m*n)}{2}, \frac{(3 + m*n)}{2}, \operatorname{Cos}[c + d*x]^2*\operatorname{Sin}[c + d*x]\right]/(d*(1 + m*n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])\right)$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2}$

```
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_)^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-mn}(c + dx) (b \cos^m(c + dx))^n) \int \cos^{mn}(c + dx) dx \\ &= \frac{\cos(c + dx) (b \cos^m(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + mn), \frac{1}{2}(3 + mn), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + mn) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.92

$$\int (b \cos^m(c + dx))^n dx = \frac{(b \cos^m(c + dx))^n \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + mn), \frac{1}{2}(3 + mn), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(1 + mn)}$$

```
[In] Integrate[(b*Cos[c + d*x]^m)^n,x]
```

```
[Out] -(((b*Cos[c + d*x]^m)^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (1 + m*n)/2, (3
+ m*n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1 + m*n)))
```

Maple [F]

$$\int (b(\cos^m(dx + c)))^n dx$$

```
[In] int((b*cos(d*x+c)^m)^n,x)
```

```
[Out] int((b*cos(d*x+c)^m)^n,x)
```

Fricas [F]

$$\int (b \cos^m(c + dx))^n dx = \int (b \cos(dx + c)^m)^n dx$$

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c)^m)^n, x)

Sympy [F]

$$\int (b \cos^m(c + dx))^n dx = \int (b \cos^m(c + dx))^n dx$$

[In] integrate((b*cos(d*x+c)**m)**n,x)

[Out] Integral((b*cos(c + d*x)**m)**n, x)

Maxima [F]

$$\int (b \cos^m(c + dx))^n dx = \int (b \cos(dx + c)^m)^n dx$$

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c)^m)^n, x)

Giac [F]

$$\int (b \cos^m(c + dx))^n dx = \int (b \cos(dx + c)^m)^n dx$$

[In] integrate((b*cos(d*x+c)^m)^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c)^m)^n, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos^m(c + dx))^n dx = \int (b \cos(c + dx)^m)^n dx$$

```
[In] int((b*cos(c + d*x)^m)^n,x)
```

```
[Out] int((b*cos(c + d*x)^m)^n, x)
```

3.58 $\int (c \cos^m(a + bx))^{5/2} dx$

Optimal result	322
Rubi [A] (verified)	322
Mathematica [A] (verified)	323
Maple [F]	324
Fricas [F(-2)]	324
Sympy [F(-1)]	324
Maxima [F]	324
Giac [F]	325
Mupad [F(-1)]	325

Optimal result

Integrand size = 14, antiderivative size = 89

$$\int (c \cos^m(a + bx))^{5/2} dx = \frac{2c^2 \cos^{1+2m}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 5m), \frac{1}{4}(6 + 5m), \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + 5m) \sqrt{\sin^2(a + bx)}}$$

[Out] $-2*c^2*\cos(b*x+a)^{(1+2*m)}*hypergeom([1/2, 1/2+5/4*m], [3/2+5/4*m], \cos(b*x+a)^2)*\sin(b*x+a)*(c*\cos(b*x+a)^m)^{(1/2)}/b/(2+5*m)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int (c \cos^m(a + bx))^{5/2} dx = \frac{2c^2 \sin(a + bx) \cos^{2m+1}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(5m + 2), \frac{1}{4}(5m + 6), \cos^2(a + bx)\right)}{b(5m + 2) \sqrt{\sin^2(a + bx)}}$$

[In] $\text{Int}[(c*\text{Cos}[a + b*x]^m)^{(5/2)}, x]$

[Out] $(-2*c^2*\text{Cos}[a + b*x]^{(1 + 2*m)}*\text{Sqrt}[c*\text{Cos}[a + b*x]^m]*\text{Hypergeometric2F1}[1/2, (2 + 5*m)/4, (6 + 5*m)/4, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(2 + 5*m)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \left(c^2 \cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{5m}{2}}(a + bx) dx \\ &= \frac{2c^2 \cos^{1+2m}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 5m), \frac{1}{4}(6 + 5m), \cos^2(a + bx)\right)}{b(2 + 5m) \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.83

$$\int (c \cos^m(a + bx))^{5/2} dx = \frac{2(c \cos^m(a + bx))^{5/2} \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 5m), \frac{1}{4}(6 + 5m), \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(2 + 5m)}$$

```
[In] Integrate[(c*Cos[a + b*x]^m)^(5/2),x]
```

```
[Out] (-2*(c*Cos[a + b*x]^m)^(5/2)*Cot[a + b*x]*Hypergeometric2F1[1/2, (2 + 5*m)/
4, (6 + 5*m)/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(2 + 5*m))
```

Maple [F]

$$\int (c(\cos^m(bx + a)))^{\frac{5}{2}} dx$$

[In] int((c*cos(b*x+a)^m)^(5/2),x)

[Out] int((c*cos(b*x+a)^m)^(5/2),x)

Fricas [F(-2)]

Exception generated.

$$\int (c \cos^m(a + bx))^{\frac{5}{2}} dx = \text{Exception raised: TypeError}$$

[In] integrate((c*cos(b*x+a)^m)^(5/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F(-1)]

Timed out.

$$\int (c \cos^m(a + bx))^{\frac{5}{2}} dx = \text{Timed out}$$

[In] integrate((c*cos(b*x+a)**m)**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int (c \cos^m(a + bx))^{\frac{5}{2}} dx = \int (c \cos(bx + a)^m)^{\frac{5}{2}} dx$$

[In] integrate((c*cos(b*x+a)^m)^(5/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(5/2), x)

Giac [F]

$$\int (c \cos^m(a + bx))^{5/2} dx = \int (c \cos(bx + a)^m)^{5/2} dx$$

[In] integrate((c*cos(b*x+a)^m)^(5/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos^m(a + bx))^{5/2} dx = \int (c \cos(a + bx)^m)^{5/2} dx$$

[In] int((c*cos(a + b*x)^m)^(5/2),x)

[Out] int((c*cos(a + b*x)^m)^(5/2), x)

3.59 $\int (c \cos^m(a + bx))^{3/2} dx$

Optimal result	326
Rubi [A] (verified)	326
Mathematica [A] (verified)	327
Maple [F]	328
Fricas [F(-2)]	328
Sympy [F]	328
Maxima [F(-2)]	328
Giac [F]	329
Mupad [F(-1)]	329

Optimal result

Integrand size = 14, antiderivative size = 83

$$\int (c \cos^m(a + bx))^{3/2} dx = \frac{2c \cos^{1+m}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 3m), \frac{3(2+m)}{4}, \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + 3m) \sqrt{\sin^2(a + bx)}}$$

[Out] $-2*c*\cos(b*x+a)^{(1+m)}*\operatorname{hypergeom}([1/2, 1/2+3/4*m], [3/2+3/4*m], \cos(b*x+a)^2)*\sin(b*x+a)*(c*\cos(b*x+a)^m)^{(1/2)}/b/(2+3*m)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int (c \cos^m(a + bx))^{3/2} dx = \frac{2c \sin(a + bx) \cos^{m+1}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(3m + 2), \frac{3(m+2)}{4}, \cos^2(a + bx)\right)}{b(3m + 2) \sqrt{\sin^2(a + bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a + b*x]^m)^{(3/2)}, x]$

[Out] $(-2*c*\operatorname{Cos}[a + b*x]^{(1 + m)}*\operatorname{Sqrt}[c*\operatorname{Cos}[a + b*x]^m]*\operatorname{Hypergeometric2F1}[1/2, (2 + 3*m)/4, (3*(2 + m))/4, \operatorname{Cos}[a + b*x]^2]*\operatorname{Sin}[a + b*x])/(b*(2 + 3*m)*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]^2])$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \left(c \cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{3m}{2}}(a + bx) dx \\ &= \frac{2c \cos^{1+m}(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 3m), \frac{3(2+m)}{4}, \cos^2(a + bx)\right) \sin^{\frac{3m}{2}}(a + bx)}{b(2 + 3m) \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.87

$$\begin{aligned} \int (c \cos^m(a + bx))^{3/2} dx = \\ \frac{2(c \cos^m(a + bx))^{3/2} \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 + 3m), \frac{3(2+m)}{4}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(2 + 3m)} \end{aligned}$$

```
[In] Integrate[(c*Cos[a + b*x]^m)^(3/2),x]
```

```
[Out] (-2*(c*Cos[a + b*x]^m)^(3/2)*Cot[a + b*x]*Hypergeometric2F1[1/2, (2 + 3*m)/
4, (3*(2 + m))/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(2 + 3*m))
```

Maple [F]

$$\int (c(\cos^m(bx + a)))^{\frac{3}{2}} dx$$

[In] int((c*cos(b*x+a)^m)^(3/2),x)

[Out] int((c*cos(b*x+a)^m)^(3/2),x)

Fricas [F(-2)]

Exception generated.

$$\int (c \cos^m(a + bx))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] integrate((c*cos(b*x+a)^m)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int (c \cos^m(a + bx))^{3/2} dx = \int (c \cos^m(a + bx))^{\frac{3}{2}} dx$$

[In] integrate((c*cos(b*x+a)**m)**(3/2),x)

[Out] Integral((c*cos(a + b*x)**m)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int (c \cos^m(a + bx))^{3/2} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((c*cos(b*x+a)^m)^(3/2),x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: BINDING-STACK overflow at size 10240. Stack can probably be resized.Proceed with caution.

Giac [F]

$$\int (c \cos^m(a + bx))^{3/2} dx = \int (c \cos(bx + a)^m)^{\frac{3}{2}} dx$$

[In] integrate((c*cos(b*x+a)^m)^(3/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c \cos^m(a + bx))^{3/2} dx = \int (c \cos(a + bx)^m)^{3/2} dx$$

[In] int((c*cos(a + b*x)^m)^(3/2),x)

[Out] int((c*cos(a + b*x)^m)^(3/2), x)

3.60 $\int \sqrt{c \cos^m(a + bx)} dx$

Optimal result	330
Rubi [A] (verified)	330
Mathematica [A] (verified)	331
Maple [F]	331
Fricas [F(-2)]	332
Sympy [F]	332
Maxima [F]	332
Giac [F]	332
Mupad [F(-1)]	333

Optimal result

Integrand size = 14, antiderivative size = 74

$$\int \sqrt{c \cos^m(a + bx)} dx = \frac{2 \cos(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{4}, \frac{6+m}{4}, \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + m) \sqrt{\sin^2(a + bx)}}$$

[Out] -2*cos(b*x+a)*hypergeom([1/2, 1/2+1/4*m], [3/2+1/4*m], cos(b*x+a)^2)*sin(b*x+a)*(c*cos(b*x+a)^m)^(1/2)/b/(2+m)/(sin(b*x+a)^2)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int \sqrt{c \cos^m(a + bx)} dx = \frac{2 \sin(a + bx) \cos(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{m+2}{4}, \frac{m+6}{4}, \cos^2(a + bx)\right)}{b(m + 2) \sqrt{\sin^2(a + bx)}}$$

[In] Int[Sqrt[c*Cos[a + b*x]^m], x]

[Out] (-2*Cos[a + b*x]*Sqrt[c*Cos[a + b*x]^m]*Hypergeometric2F1[1/2, (2 + m)/4, (6 + m)/4, Cos[a + b*x]^2]*Sin[a + b*x])/(b*(2 + m)*Sqrt[Sin[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2

```
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \left(\cos^{-\frac{m}{2}}(a + bx) \sqrt{c \cos^m(a + bx)} \right) \int \cos^{\frac{m}{2}}(a + bx) dx \\ &= -\frac{2 \cos(a + bx) \sqrt{c \cos^m(a + bx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{4}, \frac{6+m}{4}, \cos^2(a + bx)\right) \sin(a + bx)}{b(2 + m) \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.92

$$\begin{aligned} \int \sqrt{c \cos^m(a + bx)} dx = \\ -\frac{2 \sqrt{c \cos^m(a + bx)} \cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+m}{4}, \frac{6+m}{4}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(2 + m)} \end{aligned}$$

```
[In] Integrate[Sqrt[c*Cos[a + b*x]^m], x]
```

```
[Out] (-2*Sqrt[c*Cos[a + b*x]^m]*Cot[a + b*x]*Hypergeometric2F1[1/2, (2 + m)/4, (
6 + m)/4, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2])/(b*(2 + m))
```

Maple [F]

$$\int \sqrt{c (\cos^m(bx + a))} dx$$

```
[In] int((c*cos(b*x+a)^m)^(1/2), x)
```

```
[Out] int((c*cos(b*x+a)^m)^(1/2), x)
```

Fricas [F(-2)]

Exception generated.

$$\int \sqrt{c \cos^m(a + bx)} dx = \text{Exception raised: TypeError}$$

[In] integrate((c*cos(b*x+a)^m)^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int \sqrt{c \cos^m(a + bx)} dx = \int \sqrt{c \cos^m(a + bx)} dx$$

[In] integrate((c*cos(b*x+a)**m)**(1/2),x)

[Out] Integral(sqrt(c*cos(a + b*x)**m), x)

Maxima [F]

$$\int \sqrt{c \cos^m(a + bx)} dx = \int \sqrt{c \cos^m(bx + a)} dx$$

[In] integrate((c*cos(b*x+a)^m)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cos(b*x + a)^m), x)

Giac [F]

$$\int \sqrt{c \cos^m(a + bx)} dx = \int \sqrt{c \cos^m(bx + a)} dx$$

[In] integrate((c*cos(b*x+a)^m)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*cos(b*x + a)^m), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{c \cos^m(a + bx)} dx = \int \sqrt{c \cos(a + bx)^m} dx$$

```
[In] int((c*cos(a + b*x)^m)^(1/2),x)
```

```
[Out] int((c*cos(a + b*x)^m)^(1/2), x)
```

3.61 $\int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx$

Optimal result	334
Rubi [A] (verified)	334
Mathematica [A] (verified)	335
Maple [F]	335
Fricas [F(-2)]	336
Sympy [F]	336
Maxima [F]	336
Giac [F]	336
Mupad [F(-1)]	337

Optimal result

Integrand size = 14, antiderivative size = 80

$$\int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx$$

$$= -\frac{2 \cos(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-m}{4}, \frac{6-m}{4}, \cos^2(a+bx)\right) \sin(a+bx)}{b(2-m) \sqrt{c \cos^m(a+bx)} \sqrt{\sin^2(a+bx)}}$$

[Out] -2*cos(b*x+a)*hypergeom([1/2, 1/2-1/4*m], [3/2-1/4*m], cos(b*x+a)^2)*sin(b*x+a)/b/(2-m)/(c*cos(b*x+a)^m)^(1/2)/(sin(b*x+a)^2)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int \frac{1}{\sqrt{c \cos^m(a+bx)}} dx$$

$$= -\frac{2 \sin(a+bx) \cos(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-m}{4}, \frac{6-m}{4}, \cos^2(a+bx)\right)}{b(2-m) \sqrt{\sin^2(a+bx)} \sqrt{c \cos^m(a+bx)}}$$

[In] Int[1/Sqrt[c*Cos[a + b*x]^m], x]

[Out] (-2*Cos[a + b*x]*Hypergeometric2F1[1/2, (2 - m)/4, (6 - m)/4, Cos[a + b*x]^2]*Sin[a + b*x])/(b*(2 - m)*Sqrt[c*Cos[a + b*x]^m]*Sqrt[Sin[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2

```
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^{\frac{m}{2}}(a + bx) \int \cos^{-\frac{m}{2}}(a + bx) dx}{\sqrt{c \cos^m(a + bx)}} \\ &= -\frac{2 \cos(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-m}{4}, \frac{6-m}{4}, \cos^2(a + bx)\right) \sin(a + bx)}{b(2 - m) \sqrt{c \cos^m(a + bx)} \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.90

$$\begin{aligned} &\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx \\ &= \frac{2 \cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2-m}{4}, \frac{6-m}{4}, \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(-2 + m) \sqrt{c \cos^m(a + bx)}} \end{aligned}$$

```
[In] Integrate[1/Sqrt[c*Cos[a + b*x]^m],x]
```

```
[Out] (2*Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - m)/4, (6 - m)/4, Cos[a + b*x]^2
]*Sqrt[Sin[a + b*x]^2])/(b*(-2 + m)*Sqrt[c*Cos[a + b*x]^m])
```

Maple [F]

$$\int \frac{1}{\sqrt{c (\cos^m(bx + a))}} dx$$

```
[In] int(1/(c*cos(b*x+a)^m)^(1/2),x)
```

```
[Out] int(1/(c*cos(b*x+a)^m)^(1/2),x)
```

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx = \text{Exception raised: TypeError}$$

[In] `integrate(1/(c*cos(b*x+a)^m)^(1/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx$$

[In] `integrate(1/(c*cos(b*x+a)**m)**(1/2),x)`

[Out] `Integral(1/sqrt(c*cos(a + b*x)**m), x)`

Maxima [F]

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos^m(bx + a)}} dx$$

[In] `integrate(1/(c*cos(b*x+a)^m)^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/sqrt(c*cos(b*x + a)^m), x)`

Giac [F]

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos^m(bx + a)}} dx$$

[In] `integrate(1/(c*cos(b*x+a)^m)^(1/2),x, algorithm="giac")`

[Out] `integrate(1/sqrt(c*cos(b*x + a)^m), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{c \cos^m(a + bx)}} dx = \int \frac{1}{\sqrt{c \cos(a + bx)^m}} dx$$

```
[In] int(1/(c*cos(a + b*x)^m)^(1/2),x)
```

```
[Out] int(1/(c*cos(a + b*x)^m)^(1/2), x)
```

3.62 $\int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx$

Optimal result	338
Rubi [A] (verified)	338
Mathematica [A] (verified)	339
Maple [F]	340
Fricas [F(-2)]	340
Sympy [F]	340
Maxima [F]	340
Giac [F]	341
Mupad [F(-1)]	341

Optimal result

Integrand size = 14, antiderivative size = 89

$$\int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx = \frac{2 \cos^{1-m}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2-3m), \frac{3(2-m)}{4}, \cos^2(a+bx)\right) \sin(a+bx)}{bc(2-3m)\sqrt{c \cos^m(a+bx)}\sqrt{\sin^2(a+bx)}}$$

[Out] -2*cos(b*x+a)^(1-m)*hypergeom([1/2, 1/2-3/4*m], [3/2-3/4*m], cos(b*x+a)^2)*sin(b*x+a)/b/c/(2-3*m)/(c*cos(b*x+a)^m)^(1/2)/(sin(b*x+a)^2)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int \frac{1}{(c \cos^m(a+bx))^{3/2}} dx = \frac{2 \sin(a+bx) \cos^{1-m}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2-3m), \frac{3(2-m)}{4}, \cos^2(a+bx)\right)}{bc(2-3m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

[In] Int[(c*cos[a + b*x]^m)^(-3/2), x]

[Out] (-2*cos[a + b*x]^(1 - m)*Hypergeometric2F1[1/2, (2 - 3*m)/4, (3*(2 - m))/4, Cos[a + b*x]^2]*Sin[a + b*x])/(b*c*(2 - 3*m)*Sqrt[c*cos[a + b*x]^m]*Sqrt[Sin[a + b*x]^2])

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^{\frac{m}{2}}(a + bx) \int \cos^{-\frac{3m}{2}}(a + bx) dx}{c\sqrt{c \cos^m(a + bx)}} \\ &= -\frac{2 \cos^{1-m}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 - 3m), \frac{3(2-m)}{4}, \cos^2(a + bx)\right) \sin(a + bx)}{bc(2 - 3m)\sqrt{c \cos^m(a + bx)}\sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.81

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \frac{\cot(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 - 3m), -\frac{3}{4}(-2 + m), \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{\left(b - \frac{3bm}{2}\right) (c \cos^m(a + bx))^{3/2}}$$

```
[In] Integrate[(c*Cos[a + b*x]^m)^(-3/2), x]
```

```
[Out] -((Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - 3*m)/4, (-3*(-2 + m))/4, Cos[a
+ b*x]^2]*Sqrt[Sin[a + b*x]^2])/((b - (3*b*m)/2)*(c*Cos[a + b*x]^m)^(3/2)))
```

Maple [F]

$$\int \frac{1}{(c(\cos^m(bx+a)))^{\frac{3}{2}}} dx$$

[In] int(1/(c*cos(b*x+a)^m)^(3/2),x)

[Out] int(1/(c*cos(b*x+a)^m)^(3/2),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos^m(a + bx))^{\frac{3}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a)**m)**(3/2),x)

[Out] Integral((c*cos(a + b*x)**m)**(-3/2), x)

Maxima [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(bx + a)^m)^{\frac{3}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(-3/2), x)

Giac [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(bx + a)^m)^{3/2}} dx$$

[In] integrate(1/(c*cos(b*x+a)^m)^(3/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(-3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos^m(a + bx))^{3/2}} dx = \int \frac{1}{(c \cos(a + bx)^m)^{3/2}} dx$$

[In] int(1/(c*cos(a + b*x)^m)^(3/2),x)

[Out] int(1/(c*cos(a + b*x)^m)^(3/2), x)

3.63 $\int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx$

Optimal result	342
Rubi [A] (verified)	342
Mathematica [A] (verified)	343
Maple [F]	344
Fricas [F(-2)]	344
Sympy [F]	344
Maxima [F]	344
Giac [F]	345
Mupad [F(-1)]	345

Optimal result

Integrand size = 14, antiderivative size = 89

$$\int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx = \frac{2 \cos^{1-2m}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2-5m), \frac{1}{4}(6-5m), \cos^2(a+bx)\right) \sin(a+bx)}{bc^2(2-5m)\sqrt{c \cos^m(a+bx)}\sqrt{\sin^2(a+bx)}}$$

[Out] $-2*\cos(b*x+a)^{(1-2*m)}*\operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{4}(2-5*m)\right], \left[\frac{3}{2}-\frac{5}{4}*m\right], \cos(b*x+a)^2\right)*\sin(b*x+a)/b/c^2/(2-5*m)/(c*\cos(b*x+a)^m)^{(1/2)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int \frac{1}{(c \cos^m(a+bx))^{5/2}} dx = \frac{2 \sin(a+bx) \cos^{1-2m}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2-5m), \frac{1}{4}(6-5m), \cos^2(a+bx)\right)}{bc^2(2-5m)\sqrt{\sin^2(a+bx)}\sqrt{c \cos^m(a+bx)}}$$

[In] $\operatorname{Int}[(c*\operatorname{Cos}[a+b*x]^m)^{-5/2}, x]$

[Out] $(-2*\operatorname{Cos}[a+b*x]^{(1-2*m)}*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(2-5*m)}{4}, \frac{(6-5*m)}{4}, \operatorname{Cos}[a+b*x]^2\right]*\operatorname{Sin}[a+b*x])/(b*c^2*(2-5*m)*\operatorname{Sqrt}[c*\operatorname{Cos}[a+b*x]^m]*\operatorname{Sqrt}[\operatorname{Sin}[a+b*x]^2])$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^{\frac{m}{2}}(a + bx) \int \cos^{-\frac{5m}{2}}(a + bx) dx}{c^2 \sqrt{c \cos^m(a + bx)}} \\ &= -\frac{2 \cos^{1-2m}(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 - 5m), \frac{1}{4}(6 - 5m), \cos^2(a + bx)\right) \sin(a + bx)}{bc^2(2 - 5m) \sqrt{c \cos^m(a + bx)} \sqrt{\sin^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.83

$$\int \frac{1}{(c \cos^m(a + bx))^{5/2}} dx = -\frac{\cot(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2 - 5m), \frac{1}{4}(6 - 5m), \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{\left(b - \frac{5bm}{2}\right) (c \cos^m(a + bx))^{5/2}}$$

```
[In] Integrate[(c*Cos[a + b*x]^m)^(-5/2), x]
```

```
[Out] -((Cot[a + b*x]*Hypergeometric2F1[1/2, (2 - 5*m)/4, (6 - 5*m)/4, Cos[a + b*
x]^2]*Sqrt[Sin[a + b*x]^2])/((b - (5*b*m)/2)*(c*Cos[a + b*x]^m)^(5/2)))
```

Maple [F]

$$\int \frac{1}{(c(\cos^m(bx+a)))^{\frac{5}{2}}} dx$$

[In] int(1/(c*cos(b*x+a)^m)^(5/2),x)

[Out] int(1/(c*cos(b*x+a)^m)^(5/2),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c \cos^m(a + bx))^{\frac{5}{2}}} dx = \text{Exception raised: TypeError}$$

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (has polynomial part)

Sympy [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{\frac{5}{2}}} dx = \int \frac{1}{(c \cos^m(a + bx))^{\frac{5}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a)**m)**(5/2),x)

[Out] Integral((c*cos(a + b*x)**m)**(-5/2), x)

Maxima [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{\frac{5}{2}}} dx = \int \frac{1}{(c \cos(bx + a)^m)^{\frac{5}{2}}} dx$$

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(-5/2), x)

Giac [F]

$$\int \frac{1}{(c \cos^m(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(bx + a)^m)^{5/2}} dx$$

[In] integrate(1/(c*cos(b*x+a)^m)^(5/2),x, algorithm="giac")

[Out] integrate((c*cos(b*x + a)^m)^(-5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c \cos^m(a + bx))^{5/2}} dx = \int \frac{1}{(c \cos(a + bx)^m)^{5/2}} dx$$

[In] int(1/(c*cos(a + b*x)^m)^(5/2),x)

[Out] int(1/(c*cos(a + b*x)^m)^(5/2), x)

3.64 $\int (c \cos^m(a + bx))^{\frac{1}{m}} dx$

Optimal result	346
Rubi [A] (verified)	346
Mathematica [A] (verified)	347
Maple [A] (verified)	347
Fricas [A] (verification not implemented)	348
Sympy [B] (verification not implemented)	348
Maxima [F]	348
Giac [B] (verification not implemented)	349
Mupad [B] (verification not implemented)	349

Optimal result

Integrand size = 14, antiderivative size = 24

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \frac{(c \cos^m(a + bx))^{\frac{1}{m}} \tan(a + bx)}{b}$$

[Out] $(c \cdot \cos(b \cdot x + a))^m \wedge (1/m) \cdot \tan(b \cdot x + a) / b$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2717}

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \frac{\tan(a + bx) (c \cos^m(a + bx))^{\frac{1}{m}}}{b}$$

[In] `Int[(c*Cos[a + b*x]^m)^m^(-1), x]`

[Out] `((c*Cos[a + b*x]^m)^m^(-1)*Tan[a + b*x])/b`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /;`
`FreeQ[{c, d}, x]`

Rule 3287

`Int[(u_.)*((b_.)*((c_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.))^(p_.), x_Symbol] := Dist[b^IntPart[p]*((b*(c*Ssin[e + f*x])^n)^FracPart[p]/(c*Ssin[e + f*x])^(n*FracPart[p])), Int[ActivateTrig[u]*(c*Ssin[e + f*x])^(n*p), x], x] /;`
`FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat`

```
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \left((c \cos^m(a + bx))^{\frac{1}{m}} \sec(a + bx) \right) \int \cos(a + bx) dx \\ &= \frac{(c \cos^m(a + bx))^{\frac{1}{m}} \tan(a + bx)}{b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \frac{(c \cos^m(a + bx))^{\frac{1}{m}} \tan(a + bx)}{b}$$

```
[In] Integrate[(c*Cos[a + b*x]^m)^m^(-1), x]
```

```
[Out] ((c*Cos[a + b*x]^m)^m^(-1)*Tan[a + b*x])/b
```

Maple [A] (verified)

Time = 1.20 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.83

method	result	size
parallelrisch	$-\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right) (c(\cos^m(bx+a)))^{\frac{1}{m}}}{b \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}$	44

```
[In] int((c*cos(b*x+a)^m)^(1/m), x, method=_RETURNVERBOSE)
```

```
[Out] -2/b*tan(1/2*b*x+1/2*a)*(c*cos(b*x+a)^m)^(1/m)/(tan(1/2*b*x+1/2*a)^2-1)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.62

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \frac{c^{\frac{1}{m}} \sin(bx + a)}{b}$$

[In] integrate((c*cos(b*x+a)^m)^(1/m),x, algorithm="fricas")

[Out] c^(1/m)*sin(b*x + a)/b

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(20) = 40.

Time = 0.47 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.54

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \begin{cases} x(c \cos^m(a))^{\frac{1}{m}} & \text{for } b = 0 \\ x(0^m c)^{\frac{1}{m}} & \text{for } a = -bx + \frac{\pi}{2} \vee a = -bx + \frac{3\pi}{2} \\ \frac{(c \cos^m(a+bx))^{\frac{1}{m}} \sin(a+bx)}{b \cos(a+bx)} & \text{otherwise} \end{cases}$$

[In] integrate((c*cos(b*x+a)**m)**(1/m),x)

[Out] Piecewise((x*(c*cos(a)**m)**(1/m), Eq(b, 0)), (x*(0**m*c)**(1/m), Eq(a, -b*x + pi/2) | Eq(a, -b*x + 3*pi/2)), ((c*cos(a + b*x)**m)**(1/m)*sin(a + b*x)/(b*cos(a + b*x)), True))

Maxima [F]

$$\int (c \cos^m(a + bx))^{\frac{1}{m}} dx = \int (c \cos(bx + a)^m)^{\frac{1}{m}} dx$$

[In] integrate((c*cos(b*x+a)^m)^(1/m),x, algorithm="maxima")

[Out] integrate((c*cos(b*x + a)^m)^(1/m), x)

3.65 $\int (a(b \cos(c + dx))^p)^n dx$

Optimal result	350
Rubi [A] (verified)	350
Mathematica [A] (verified)	351
Maple [F]	351
Fricas [F]	352
Sympy [F]	352
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	353

Optimal result

Integrand size = 14, antiderivative size = 80

$$\int (a(b \cos(c + dx))^p)^n dx = \frac{\cos(c + dx) (a(b \cos(c + dx))^p)^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + np) \sqrt{\sin^2(c + dx)}}$$

[Out] `-cos(d*x+c)*(a*(b*cos(d*x+c))^p)^n*hypergeom([1/2, 1/2*n*p+1/2], [1/2*n*p+3/2], cos(d*x+c)^2)*sin(d*x+c)/d/(n*p+1)/(sin(d*x+c)^2)^(1/2)`

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3287, 2722}

$$\int (a(b \cos(c + dx))^p)^n dx = \frac{\sin(c + dx) \cos(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(np + 1), \frac{1}{2}(np + 3), \cos^2(c + dx)\right) (a(b \cos(c + dx))^p)^n}{d(np + 1) \sqrt{\sin^2(c + dx)}}$$

[In] `Int[(a*(b*Cos[c + d*x])^p)^n,x]`

[Out] `-((Cos[c + d*x]*(a*(b*Cos[c + d*x])^p)^n*Hypergeometric2F1[1/2, (1 + n*p)/2, (3 + n*p)/2, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(1 + n*p)*Sqrt[Sin[c + d*x]^2]))`

Rule 2722

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2`

```
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rule 3287

```
Int[(u_.)*((b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_)^(p_), x_Symbol] := D
ist[b^IntPart[p]*((b*(c*Sin[e + f*x])^n)^FracPart[p]/(c*Sin[e + f*x])^(n*Fr
acPart[p])), Int[ActivateTrig[u]*(c*Sin[e + f*x])^(n*p), x], x] /; FreeQ[{b
, c, e, f, n, p}, x] && !IntegerQ[p] && !IntegerQ[n] && (EqQ[u, 1] || Mat
chQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin,
cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned} \text{integral} &= ((b \cos(c + dx))^{-np} (a(b \cos(c + dx))^p)^n) \int (b \cos(c + dx))^{np} dx \\ &= \frac{\cos(c + dx) (a(b \cos(c + dx))^p)^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + np) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.92

$$\int (a(b \cos(c + dx))^p)^n dx = \frac{(a(b \cos(c + dx))^p)^n \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + np), \frac{1}{2}(3 + np), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(1 + np)}$$

```
[In] Integrate[(a*(b*Cos[c + d*x])^p)^n,x]
```

```
[Out] -(((a*(b*Cos[c + d*x])^p)^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (1 + n*p)/2
, (3 + n*p)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1 + n*p)))
```

Maple [F]

$$\int (a(\cos(dx + c)b)^p)^n dx$$

```
[In] int((a*(cos(d*x+c)*b)^p)^n,x)
```

```
[Out] int((a*(cos(d*x+c)*b)^p)^n,x)
```

Fricas [F]

$$\int (a(b \cos(c + dx))^p)^n dx = \int ((b \cos(dx + c))^p a)^n dx$$

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="fricas")

[Out] integral(((b*cos(d*x + c))^p*a)^n, x)

Sympy [F]

$$\int (a(b \cos(c + dx))^p)^n dx = \int (a(b \cos(c + dx))^p)^n dx$$

[In] integrate((a*(b*cos(d*x+c))**p)**n,x)

[Out] Integral((a*(b*cos(c + d*x))**p)**n, x)

Maxima [F]

$$\int (a(b \cos(c + dx))^p)^n dx = \int ((b \cos(dx + c))^p a)^n dx$$

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="maxima")

[Out] integrate(((b*cos(d*x + c))^p*a)^n, x)

Giac [F]

$$\int (a(b \cos(c + dx))^p)^n dx = \int ((b \cos(dx + c))^p a)^n dx$$

[In] integrate((a*(b*cos(d*x+c))^p)^n,x, algorithm="giac")

[Out] integrate(((b*cos(d*x + c))^p*a)^n, x)

Mupad [F(-1)]

Timed out.

$$\int (a(b \cos(c + dx))^p)^n dx = \int (a(b \cos(c + dx))^p)^n dx$$

```
[In] int((a*(b*cos(c + d*x))^p)^n,x)
```

```
[Out] int((a*(b*cos(c + d*x))^p)^n, x)
```

3.66 $\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	354
Rubi [A] (verified)	354
Mathematica [A] (verified)	356
Maple [A] (verified)	357
Fricas [C] (verification not implemented)	357
Sympy [F(-1)]	357
Maxima [F]	358
Giac [F]	358
Mupad [F(-1)]	358

Optimal result

Integrand size = 21, antiderivative size = 123

$$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{30b \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{30 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^2d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d}$$

```
[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^2/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^4/d+30/77*b*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d
```

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used

= {16, 2715, 2721, 2720}

$$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2 \sin(c + dx) (b \cos(c + dx))^{9/2}}{11b^4d} + \frac{18 \sin(c + dx) (b \cos(c + dx))^{5/2}}{77b^2d} + \frac{30 \sin(c + dx) \sqrt{b \cos(c + dx)}}{77d} + \frac{30b \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}}$$

[In] Int[Cos[c + d*x]^5*Sqrt[b*Cos[c + d*x]],x]

[Out] (30*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^2*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^4*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\text{integral} = \frac{\int (b \cos(c + dx))^{11/2} dx}{b^5}$$

$$\begin{aligned}
&= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b^3} \\
&= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^2d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d} + \frac{45 \int (b \cos(c + dx))^{3/2} dx}{77b} \\
&= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^2d} \\
&\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d} + \frac{1}{77}(15b) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^2d} \\
&\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d} + \frac{\left(15b\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{77\sqrt{b \cos(c + dx)}} \\
&= \frac{30b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d\sqrt{b \cos(c + dx)}} + \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} \\
&\quad + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^2d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^4d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.67

$$\begin{aligned}
&\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx \\
&= \frac{\sqrt{b \cos(c + dx)} \left(240 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)} (290 \sin(c + dx) + 57 \sin(3(c + dx))) + 7 \sin(5(c + dx)) \right)}{616d\sqrt{\cos(c + dx)}}
\end{aligned}$$

[In] Integrate[Cos[c + d*x]^5*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)])))/(616*d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 6.76 (sec) , antiderivative size = 234, normalized size of antiderivative = 1.90

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1084\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-370\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{77\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}$

[In] int(cos(d*x+c)^5*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

```
[Out] -2/77*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(448*cos(1/2*d*x+1/2*c)^13-1568*cos(1/2*d*x+1/2*c)^11+2384*cos(1/2*d*x+1/2*c)^9-2040*cos(1/2*d*x+1/2*c)^7+1084*cos(1/2*d*x+1/2*c)^5-370*cos(1/2*d*x+1/2*c)^3+15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+62*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 0.80

$$\int \cos^5(c+dx)\sqrt{b\cos(c+dx)}dx$$

$$= \frac{2(7\cos(dx+c)^4+9\cos(dx+c)^2+15)\sqrt{b\cos(dx+c)}\sin(dx+c)-15i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c))+I\sin(dx+c)+15I\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c))-I\sin(dx+c))}{77d}$$

[In] integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

```
[Out] 1/77*(2*(7*cos(d*x+c)^4+9*cos(d*x+c)^2+15)*sqrt(b*cos(d*x+c))*sin(d*x+c)-15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4,0,cos(d*x+c))+I*sin(d*x+c)+15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4,0,cos(d*x+c))-I*sin(d*x+c))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^5(c+dx)\sqrt{b\cos(c+dx)}dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**5*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^5 dx$$

[In] integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^5, x)

Giac [F]

$$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^5 dx$$

[In] integrate(cos(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^5, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^5(c + dx) \sqrt{b \cos(c + dx)} dx = \int \cos(c + dx)^5 \sqrt{b \cos(c + dx)} dx$$

[In] int(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2), x)

3.67 $\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	359
Rubi [A] (verified)	359
Mathematica [A] (verified)	361
Maple [B] (verified)	361
Fricas [C] (verification not implemented)	361
Sympy [F(-1)]	362
Maxima [F]	362
Giac [F]	362
Mupad [F(-1)]	362

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3d}$$

[Out] $14/45*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/b/d+2/9*(b*\cos(d*x+c))^{(7/2)}*\sin(d*x+c)/b^3/d+14/15*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2 \sin(c + dx) (b \cos(c + dx))^{7/2}}{9b^3d} + \frac{14 \sin(c + dx) (b \cos(c + dx))^{3/2}}{45bd} + \frac{14 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^4*\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] (14*sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*d*sqrt[Cos[c + d*x]]) + (14*(b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b*d) + (2*(b*cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^4} \\
 &= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b^2} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3d} + \frac{7}{15} \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3d} \\
 &\quad + \frac{\left(7\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)}} \\
 &= \frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} \\
 &\quad + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45bd} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{\sqrt{b \cos(c + dx)} \left(168 E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)} (38 \sin(2(c + dx)) + 5 \sin(4(c + dx))) \right)}{180 d \sqrt{\cos(c + dx)}}$$

[In] Integrate[Cos[c + d*x]^4*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Sqrt[Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(109) = 218.

Time = 4.48 (sec) , antiderivative size = 221, normalized size of antiderivative = 2.28

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+256\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-128\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] int(cos(d*x+c)^4*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2/45*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-24*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.04

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{2 \left(5 \cos(dx + c)^3 + 7 \cos(dx + c) \right) \sqrt{b \cos(dx + c)} \sin(dx + c) + 21i \sqrt{2} \sqrt{b} \text{weierstrassZeta}(-4, 0, \text{weiers}}{}$$

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

```
[Out] 1/45*(2*(5*cos(d*x + c)^3 + 7*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x +
c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0,
cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4,
0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**4*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^4 dx$$

```
[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^4, x)
```

Giac [F]

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^4 dx$$

```
[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int \cos^4(c + dx) \sqrt{b \cos(c + dx)} dx = \int \cos(c + dx)^4 \sqrt{b \cos(c + dx)} dx$$

```
[In] int(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2), x)
```

3.68 $\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	363
Rubi [A] (verified)	363
Mathematica [A] (verified)	365
Maple [A] (verified)	365
Fricas [C] (verification not implemented)	366
Sympy [F(-1)]	366
Maxima [F]	366
Giac [F]	367
Mupad [F(-1)]	367

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}} + \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^2d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^2/d+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2 \sin(c + dx) (b \cos(c + dx))^{5/2}}{7b^2d} + \frac{10 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^3*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]], x]$

[Out] (10*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(21*d*Sqrt[b*Cos[c + d*x]]) + (10*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(21*d) + (2*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(7*b^2*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^3} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^2d} + \frac{5 \int (b \cos(c + dx))^{3/2} dx}{7b} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^2d} \\
 &\quad + \frac{1}{21}(5b) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^2d} \\
 &\quad + \frac{(5b\sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

$$= \frac{10b\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b\cos(c+dx)}} + \frac{10\sqrt{b\cos(c+dx)} \sin(c+dx)}{21d} + \frac{2(b\cos(c+dx))^{5/2} \sin(c+dx)}{7b^2d}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.77

$$\int \cos^3(c+dx) \sqrt{b\cos(c+dx)} dx$$

$$= \frac{\sqrt{b\cos(c+dx)} \left(20 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sqrt{\cos(c+dx)} (23 \sin(c+dx) + 3 \sin(3(c+dx))) \right)}{42d\sqrt{\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^3*Sqrt[b*Cos[c + d*x]], x]

[Out] (Sqrt[b*Cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.46 (sec) , antiderivative size = 208, normalized size of antiderivative = 2.19

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}b\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}$

[In] int(cos(d*x+c)^3*(cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)

[Out]
$$-2/21*\left(\left(2*\cos(1/2*d*x+1/2*c)\right)^2-1\right)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b*(48*\cos(1/2*d*x+1/2*c)^9-120*\cos(1/2*d*x+1/2*c)^7+128*\cos(1/2*d*x+1/2*c)^5-72*\cos(1/2*d*x+1/2*c)^3+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})+16*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.93

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{2 \sqrt{b \cos(dx + c)} (3 \cos(dx + c)^2 + 5) \sin(dx + c) - 5i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))}{21 d}$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/21*(2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 5)*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^3 dx$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^3, x)
```

Giac [F]

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^3 dx$$

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^3(c + dx) \sqrt{b \cos(c + dx)} dx = \int \cos(c + dx)^3 \sqrt{b \cos(c + dx)} dx$$

[In] int(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2), x)

3.69 $\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	368
Rubi [A] (verified)	368
Mathematica [A] (verified)	369
Maple [B] (verified)	370
Fricas [C] (verification not implemented)	370
Sympy [F]	371
Maxima [F]	371
Giac [F]	371
Mupad [F(-1)]	371

Optimal result

Integrand size = 21, antiderivative size = 69

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd}$$

[Out] $2/5*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/b/d+6/5*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2 \sin(c + dx) (b \cos(c + dx))^{3/2}}{5bd} + \frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[In] `Int[Cos[c + d*x]^2*Sqrt[b*Cos[c + d*x]],x]`

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(5*b*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n-1)/(d*n)), x] + Dist[b^2*((n-1)/n), Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b^2} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd} + \frac{3}{5} \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
 &= \frac{6\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5bd}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.90

$$\begin{aligned}
 &\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx \\
 &= \frac{\sqrt{b \cos(c + dx)} \left(6E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)} \sin(2(c + dx))\right)}{5d\sqrt{\cos(c + dx)}}
 \end{aligned}$$

`[In] Integrate[Cos[c + d*x]^2*Sqrt[b*Cos[c + d*x]],x]`

`[Out] (Sqrt[b*Cos[c + d*x]]*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*d*Sqrt[Cos[c + d*x]])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. $2(85) = 170$.

Time = 2.84 (sec) , antiderivative size = 211, normalized size of antiderivative = 3.06

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)}b\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}$

[In] `int(cos(d*x+c)^2*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.28

$$\int \cos^2(c+dx)\sqrt{b\cos(c+dx)}dx$$

$$= \frac{2\sqrt{b\cos(dx+c)}\cos(dx+c)\sin(dx+c)+3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+\sin(dx+c))) - 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-\sin(dx+c)))}{d}$$

[In] `integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{b*\cos(dx+c)}*\cos(dx+c)*\sin(dx+c)+3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+\sin(dx+c)))) - 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-\sin(dx+c))))/d$$

Sympy [F]

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(c + dx)} \cos^2(c + dx) dx$$

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*cos(c + d*x)**2, x)

Maxima [F]

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^2, x)

Giac [F]

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*cos(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx) \sqrt{b \cos(c + dx)} dx = \int \cos(c + dx)^2 \sqrt{b \cos(c + dx)} dx$$

[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2), x)

3.70 $\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	372
Rubi [A] (verified)	372
Mathematica [A] (verified)	373
Maple [B] (verified)	374
Fricas [C] (verification not implemented)	374
Sympy [F]	375
Maxima [F]	375
Giac [F]	375
Mupad [F(-1)]	375

Optimal result

Integrand size = 19, antiderivative size = 67

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2b \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}$$

[Out] 2/3*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+2/3*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d} + \frac{2b \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

[In] Int[Cos[c + d*x]*Sqrt[b*Cos[c + d*x]],x]

[Out] (2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*d*Sqrt[b*Cos[c + d*x]]) + (2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{3/2} dx}{b} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{1}{3}b \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{(b\sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2b\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.91

$$\begin{aligned}
 &\int \cos(c + dx)\sqrt{b \cos(c + dx)} dx \\
 &= \frac{2(b \cos(c + dx))^{3/2} \left(\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)} \sin(c + dx) \right)}{3bd \cos^{3/2}(c + dx)}
 \end{aligned}$$

```
[In] Integrate[Cos[c + d*x]*Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (2*(b*Cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*b*d*Cos[c + d*x]^(3/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. $2(83) = 166$.

Time = 2.21 (sec) , antiderivative size = 188, normalized size of antiderivative = 2.81

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\right)}}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$

```
[In] int(cos(d*x+c)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.13

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{3d}$$

```
[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/3*(-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*sin(d*x + c))/d
```

Sympy [F]

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(c + dx)} \cos(c + dx) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))**(1/2),x)`

[Out] `Integral(sqrt(b*cos(c + d*x))*cos(c + d*x), x)`

Maxima [F]

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*cos(d*x + c))*cos(d*x + c), x)`

Giac [F]

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(b*cos(d*x + c))*cos(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx) \sqrt{b \cos(c + dx)} dx = \int \cos(c + dx) \sqrt{b \cos(c + dx)} dx$$

[In] `int(cos(c + d*x)*(b*cos(c + d*x))^(1/2),x)`

[Out] `int(cos(c + d*x)*(b*cos(c + d*x))^(1/2), x)`

3.71 $\int \sqrt{b \cos(c + dx)} dx$

Optimal result	376
Rubi [A] (verified)	376
Mathematica [A] (verified)	377
Maple [B] (verified)	377
Fricas [C] (verification not implemented)	378
Sympy [F]	378
Maxima [F]	378
Giac [F]	378
Mupad [F(-1)]	379

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \sqrt{b \cos(c + dx)} dx = \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2721, 2719}

$$\int \sqrt{b \cos(c + dx)} dx = \frac{2E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[In] Int[Sqrt[b*Cos[c + d*x]],x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ

[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \sqrt{b \cos(c + dx)} dx = \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]],x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(60) = 120.

Time = 2.04 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.74

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)bd}}$
risch	$-\frac{i\sqrt{2}\sqrt{(e^{2i(dx+c)}+1)be^{-i(dx+c)}}}{d} - \frac{i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)\left(-2iE\left(\sqrt{-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}}}\right)\right)}{d}$

[In] int((cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] 2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.66

$$\int \sqrt{b \cos(c + dx)} dx$$

$$= \frac{i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

```
[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d
```

Sympy [F]

$$\int \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(c + dx)} dx$$

```
[In] integrate((b*cos(d*x+c))**(1/2),x)
```

```
[Out] Integral(sqrt(b*cos(c + d*x)), x)
```

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} dx$$

```
[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c)), x)
```

Giac [F]

$$\int \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(dx + c)} dx$$

```
[In] integrate((b*cos(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*cos(d*x + c)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} dx = \int \sqrt{b \cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^(1/2),x)
```

```
[Out] int((b*cos(c + d*x))^(1/2), x)
```

3.72 $\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$

Optimal result	380
Rubi [A] (verified)	380
Mathematica [A] (verified)	381
Maple [B] (verified)	381
Fricas [C] (verification not implemented)	382
Sympy [F]	382
Maxima [F]	382
Giac [F]	382
Mupad [F(-1)]	383

Optimal result

Integrand size = 19, antiderivative size = 39

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d\sqrt{b \cos(c + dx)}}$$

[Out] 2*b*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2721, 2720}

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d\sqrt{b \cos(c + dx)}}$$

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x],x]

[Out] (2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])ⁿ/Sin[c + d*x]ⁿ, Int[Sin[c + d*x]ⁿ, x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \frac{2b \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x],x]

[Out] (2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(61) = 122.

Time = 1.44 (sec) , antiderivative size = 142, normalized size of antiderivative = 3.64

method	result	size
default	$\frac{2 \sqrt{\left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{\left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) b d}}$	142

[In] int(sec(d*x+c)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.46

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

```
[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] (-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/d
```

Sympy [F]

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \int \sqrt{b \cos(c + dx)} \sec(c + dx) dx$$

```
[In] integrate(sec(d*x+c)*(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x), x)
```

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c) dx$$

```
[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c), x)
```

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c) dx$$

```
[In] integrate(sec(d*x+c)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c), x)
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x), x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x), x)
```

3.73 $\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$

Optimal result	384
Rubi [A] (verified)	384
Mathematica [A] (verified)	385
Maple [B] (verified)	386
Fricas [C] (verification not implemented)	386
Sympy [F]	387
Maxima [F]	387
Giac [F]	387
Mupad [F(-1)]	387

Optimal result

Integrand size = 21, antiderivative size = 63

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = -\frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}$$

[Out] $2*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[In] `Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2,x]`

[Out] `(-2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]) + (2*b*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]])`

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.76

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \frac{2b \left(-\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2,x]
```

```
[Out] (2*b*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*S
qrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 195 vs. $2(83) = 166$.

Time = 1.78 (sec) , antiderivative size = 196, normalized size of antiderivative = 3.11

method	result
default	$\frac{2b\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)}}$

[In] `int(sec(d*x+c)^2*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2*b*(-2*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}}}{d}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.60

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} \sqrt{b} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 \sqrt{b \cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))}$$

[In] `integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$\frac{(-I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*\sqrt{b*\cos(d*x + c)}*\sin(d*x + c)}{(d*\cos(d*x + c))}$$

Sympy [F]

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx$$

[In] integrate(sec(d*x+c)**2*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x)**2, x)

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^2 dx$$

[In] integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^2, x)

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^2 dx$$

[In] integrate(sec(d*x+c)^2*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^2(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^2} dx$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^2, x)

3.74 $\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$

Optimal result	388
Rubi [A] (verified)	388
Mathematica [A] (verified)	389
Maple [B] (verified)	390
Fricas [C] (verification not implemented)	390
Sympy [F]	391
Maxima [F]	391
Giac [F]	391
Mupad [F(-1)]	391

Optimal result

Integrand size = 21, antiderivative size = 70

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}$$

[Out] $2/3*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+2/3*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sec}[c + d*x]^3, x]$

[Out] $(2*b*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^2*\operatorname{Sin}[c + d*x])/(3*d*(b*\operatorname{Cos}[c + d*x])^{(3/2)})$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3}b \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\left(b\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2b\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.70

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \frac{2b\left(\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx)\right)}{3d\sqrt{b \cos(c + dx)}}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^3,x]

[Out] (2*b*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(86) = 172$.

Time = 1.84 (sec) , antiderivative size = 239, normalized size of antiderivative = 3.41

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\sin\left(\frac{dx}{2}\right)}$

[In] `int(sec(d*x+c)^3*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{2}{3}\left(-2\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)^{\frac{1}{2}}\text{EllipticF}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)+\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)\right)^2\right)^{\frac{1}{2}}\left(2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)^{\frac{1}{2}}\text{EllipticF}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)\right)*b*\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}/\left(-b*\left(2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4-\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)\right)^{\frac{1}{2}}/\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)/\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)/\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b\right)^{\frac{1}{2}}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.43

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 \sqrt{b \cos(dx + c)} \sin(dx + c)}{3 d \cos(dx + c)^2}$$

[In] `integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$\frac{1}{3}\left(-I\sqrt{2}\sqrt{b}\cos(dx + c)^2\text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I\sin(dx + c)) + I\sqrt{2}\sqrt{b}\cos(dx + c)^2\text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I\sin(dx + c)) + 2\sqrt{b\cos(dx + c)}\sin(dx + c)\right)/(d\cos(dx + c)^2)$$

Sympy [F]

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx$$

[In] integrate(sec(d*x+c)**3*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x)**3, x)

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^3 dx$$

[In] integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^3, x)

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^3 dx$$

[In] integrate(sec(d*x+c)^3*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^3(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^3} dx$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^3, x)

3.75 $\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$

Optimal result	392
Rubi [A] (verified)	392
Mathematica [A] (verified)	394
Maple [B] (verified)	394
Fricas [C] (verification not implemented)	395
Sympy [F]	395
Maxima [F]	395
Giac [F]	396
Mupad [F(-1)]	396

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = -\frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}$$

[Out] $2/5*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+6/5*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-6/5*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[In] `Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^4,x]`

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^(5/2)) + (6*b*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^4 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5} (3b^2) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{3}{5} \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
 &= -\frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.73

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$$

$$= \frac{2\sqrt{b \cos(c + dx)} \sec^2(c + dx) \left(-3 \cos^{\frac{3}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + \frac{3}{2} \sin(2(c + dx)) + \tan(c + dx) \right)}{5d}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^4,x]

[Out] (2*Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^2*(-3*Cos[c + d*x]^(3/2)*EllipticE[(c + d*x)/2, 2] + (3*Sin[2*(c + d*x)]/2 + Tan[c + d*x]))/(5*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 363 vs. 2(107) = 214.

Time = 2.55 (sec) , antiderivative size = 364, normalized size of antiderivative = 3.83

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))} \left(24 \cos(\frac{dx}{2} + \frac{c}{2}) (\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12 \sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} E(\cos(\frac{dx}{2} + \frac{c}{2})) \right)}{5d}$

[In] int(sec(d*x+c)^4*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-2/5*(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)/\sin(1/2*d*x+1/2*c)^3/(8*\sin(1/2*d*x+1/2*c)^6-12*\sin(1/2*d*x+1/2*c)^4+6*\sin(1/2*d*x+1/2*c)^2-1)*(24*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-12*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(\cos(1/2*d*x+1/2*c),2^(1/2))*\sin(1/2*d*x+1/2*c)^4-24*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+12*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(\cos(1/2*d*x+1/2*c),2^(1/2))*\sin(1/2*d*x+1/2*c)^2+8*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(\cos(1/2*d*x+1/2*c),2^(1/2))*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^(1/2)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.24

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$$

$$= \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)))}{1}$$

[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 1)*sin(d*x + c))/(d*cos(d*x + c)^3)

Sympy [F]

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = \int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx$$

[In] integrate(sec(d*x+c)**4*(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))*sec(c + d*x)**4, x)

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^4 dx$$

[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^4, x)

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^4 dx$$

[In] integrate(sec(d*x+c)^4*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^4, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^4(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^4} dx$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^4, x)

3.76 $\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx$

Optimal result	397
Rubi [A] (verified)	397
Mathematica [A] (verified)	399
Maple [B] (verified)	399
Fricas [C] (verification not implemented)	400
Sympy [F(-1)]	400
Maxima [F]	400
Giac [F]	401
Mupad [F(-1)]	401

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}$$

[Out] $2/7*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sec}[c + d*x]^5, x]$

[Out] $(10*b*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^4*\operatorname{Sin}[c + d*x])/(7*d*(b*\operatorname{Cos}[c + d*x])^{(7/2)}) + (10*b^2*\operatorname{Sin}[c + d*x])/(21*d*(b*\operatorname{Cos}[c + d*x])^{(3/2)})$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^5 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{1}{7}(5b^3) \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{1}{21}(5b) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{\left(5b\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21\sqrt{b \cos(c + dx)}} \\
 &= \frac{10b\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^2 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.34 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx$$

$$= \frac{\sqrt{b \cos(c + dx)} \sec^3(c + dx) \left(10 \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21d}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^5,x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^3*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 395 vs. 2(110) = 220.

Time = 2.14 (sec) , antiderivative size = 396, normalized size of antiderivative = 4.04

method	result
default	$-\frac{2\left(-40\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-40\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+60\sqrt{\frac{1}{2}-\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}\right)}{21d}$

[In] int(sec(d*x+c)^5*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^6-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*b*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx$$

$$= \frac{-5i \sqrt{2} \sqrt{b} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} \sqrt{b} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 \sqrt{b} \cos(dx + c) (5 \cos^2(dx + c) + 3) \sin(dx + c)}{21 d \cos(dx + c)^4}$$

[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*(5*cos(d*x + c)^2 + 3)*sin(d*x + c))/(d*cos(d*x + c)^4)

Sympy [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \text{Timed out}$$

[In] integrate(sec(d*x+c)**5*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^5 dx$$

[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^5, x)

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^5 dx$$

[In] integrate(sec(d*x+c)^5*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^5, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^5(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^5} dx$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^5,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^5, x)

3.77 $\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx$

Optimal result	402
Rubi [A] (verified)	402
Mathematica [A] (verified)	404
Maple [B] (verified)	404
Fricas [C] (verification not implemented)	405
Sympy [F(-1)]	405
Maxima [F]	405
Giac [F]	406
Mupad [F(-1)]	406

Optimal result

Integrand size = 21, antiderivative size = 123

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = -\frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}}$$

[Out] $2/9*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[In] Int[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^6,x]

[Out] $(-14\sqrt{b\cos[c + dx]}\text{EllipticE}[(c + dx)/2, 2]) / (15d\sqrt{\cos[c + dx]}) + (2b^5\sin[c + dx]) / (9d(b\cos[c + dx])^{9/2}) + (14b^3\sin[c + dx]) / (45d(b\cos[c + dx])^{5/2}) + (14b\sin[c + dx]) / (15d\sqrt{b\cos[c + dx]})$

Rule 16

$\text{Int}[(u_.)*(v_)^{(m_.)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \&\& \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_.)\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\cos[c + dx]*((b*\sin[c + dx])^{(n+1)} / (b*d*(n+1))), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\sin[c + dx])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d, x\} \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\sqrt{\sin[(c_.) + (d_.)*(x_)]}, x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \pi/2 + dx), 2], x] /; \text{FreeQ}\{c, d, x\}$

Rule 2721

$\text{Int}[(b_.)\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\sin[c + dx])^n / \sin[c + dx]^n, \text{Int}[\sin[c + dx]^n, x], x] /; \text{FreeQ}\{b, c, d, x\} \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^6 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9}(7b^4) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{1}{15}(7b^2) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7}{15} \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} \\
 &\quad + \frac{14b \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{(7\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)}}
 \end{aligned}$$

$$= -\frac{14\sqrt{b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15d\sqrt{\cos(c+dx)}} + \frac{2b^5\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}}$$

$$+ \frac{14b^3\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}} + \frac{14b\sin(c+dx)}{15d\sqrt{b\cos(c+dx)}}$$

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.64

$$\int \sqrt{b\cos(c+dx)}\sec^6(c+dx)dx$$

$$= \frac{\sqrt{b\cos(c+dx)}\sec^5(c+dx)\left(-336\cos^{\frac{9}{2}}(c+dx)E\left(\frac{1}{2}(c+dx)\middle|2\right) + 150\sin(c+dx) + 91\sin(3(c+dx)) + 21\sin(5(c+dx))\right)}{360d}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^6,x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sec[c + d*x]^5*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)])/(360*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 413 vs. 2(131) = 262.

Time = 3.06 (sec) , antiderivative size = 414, normalized size of antiderivative = 3.37

method	result
default	$-\frac{2\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1\right)}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b\left(-\frac{\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{144b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)-\frac{1}{2}\right)^5}-\frac{7\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}{180b\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)-\frac{1}{2}\right)^5}\right)}{360d}$

[In] int(sec(d*x+c)^6*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b*(-1/144*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/180*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^3-14/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)+7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)

$)^2)^{1/2} * (\text{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{1/2}) - \text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^{1/2})) / \sin(1/2*d*x+1/2*c) / ((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{1/2} / d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.04

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx$$

$$= \frac{-21i \sqrt{2} \sqrt{b} \cos(dx + c)^5 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)))}{d}$$

[In] integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*cos(d*x + c)^4 + 7*cos(d*x + c)^2 + 5)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)

Sympy [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = \text{Timed out}$$

[In] integrate(sec(d*x+c)**6*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^6 dx$$

[In] integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^6, x)

Giac [F]

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = \int \sqrt{b \cos(dx + c)} \sec(dx + c)^6 dx$$

[In] integrate(sec(d*x+c)^6*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))*sec(d*x + c)^6, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{b \cos(c + dx)} \sec^6(c + dx) dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^6} dx$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^6,x)

[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^6, x)

3.78 $\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal result	407
Rubi [A] (verified)	407
Mathematica [A] (verified)	409
Maple [A] (verified)	409
Fricas [C] (verification not implemented)	410
Sympy [F(-1)]	410
Maxima [F]	410
Giac [F]	411
Mupad [F(-1)]	411

Optimal result

Integrand size = 21, antiderivative size = 126

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{30b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{30b \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77bd} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^3d}$$

[Out] $18/77*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b/d+2/11*(b*\cos(d*x+c))^{(9/2)}*\sin(d*x+c)/b^3/d+30/77*b^2*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+30/77*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^3d} + \frac{30b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77bd} + \frac{30b \sin(c + dx) \sqrt{b \cos(c + dx)}}{77d}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^4*(b*\operatorname{Cos}[c + d*x])^{(3/2)}, x]$

[Out] (30*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^4} \\
 &= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^3d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b^2} \\
 &= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77bd} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^3d} + \frac{45}{77} \int (b \cos(c + dx))^{3/2} dx \\
 &= \frac{30b\sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77bd} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^3d} + \frac{1}{77} (15b^2) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{30b\sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77bd} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^3d} + \frac{\left(15b^2 \sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{77\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

$$= \frac{30b^2 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77d \sqrt{b \cos(c+dx)}} + \frac{30b \sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} \\ + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77bd} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^3d}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.66

$$\int \cos^4(c+dx)(b \cos(c+dx))^{3/2} dx = \frac{(b \cos(c+dx))^{3/2} \left(240 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sqrt{\cos(c+dx)}(290 \sin(c+dx) + 57 \sin(3(c+dx))) \right)}{616d \cos^{3/2}(c+dx)}$$

[In] Integrate[Cos[c + d*x]^4*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)])))/(616*d*Cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 5.70 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.87

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1084\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-370\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15\left(\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\right)^{1/2}\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^{1/2}\operatorname{EllipticF}\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right), 2\right)^{1/2}+62\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}{77\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\right)}/\sin\left(\frac{dx}{2}+\frac{c}{2}\right)/\left(\left(2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-1\right)b\right)^{1/2}/d}$

[In] int(cos(d*x+c)^4*(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/77*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(448*cos(1/2*d*x+1/2*c)^13-1568*cos(1/2*d*x+1/2*c)^11+2384*cos(1/2*d*x+1/2*c)^9-2040*cos(1/2*d*x+1/2*c)^7+1084*cos(1/2*d*x+1/2*c)^5-370*cos(1/2*d*x+1/2*c)^3+15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2)^(1/2)+62*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.81

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{-15i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 15i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{b^2 \cos^2(dx + c) \sin^2(dx + c)}$$

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/77*(-15*I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 15*I*sqrt(2)*b^(3/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(7*b*cos(d*x + c)^4 + 9*b*cos(d*x + c)^2 + 15*b)*sqrt(b*cos(d*x + c))*sin(d*x + c))/d

Sympy [F(-1)]

Timed out.

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**4*(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{3/2} \cos^4(dx + c) dx$$

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^4, x)

Giac [F]

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

[In] integrate(cos(d*x+c)^4*(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^4, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^4(c + dx)(b \cos(c + dx))^{3/2} dx = \int \cos(c + dx)^4 (b \cos(c + dx))^{3/2} dx$$

[In] int(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2), x)

3.79 $\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal result	412
Rubi [A] (verified)	412
Mathematica [A] (verified)	414
Maple [B] (verified)	414
Fricas [C] (verification not implemented)	414
Sympy [F(-1)]	415
Maxima [F]	415
Giac [F]	415
Mupad [F(-1)]	415

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{14b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d}$$

[Out] $14/45*(b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/d+2/9*(b*\cos(d*x+c))^{7/2}*\sin(d*x+c)/b^2/d+14/15*b*(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{1/2})*(b*\cos(d*x+c))^{1/2}/d/\cos(d*x+c)^{1/2}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9b^2d} + \frac{14 \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d} + \frac{14bE\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^3*(b*\text{Cos}[c + d*x])^{3/2}, x]$

[Out] $(14*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (14*(b*\text{Cos}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(45*d) + (2*(b*\text{Cos}[c + d*x])^{7/2}*\text{Sin}[c + d*x])/(9*b^2*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^3} \\
 &= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} \\
 &\quad + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} + \frac{1}{15}(7b) \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d} \\
 &\quad + \frac{\left(7b\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)}} \\
 &= \frac{14b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} \\
 &\quad + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.79

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(b \cos(c + dx))^{3/2} \left(168E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)}(38 \sin(2(c + dx)) + 5 \sin(4(c + dx))) \right)}{180d \cos^{\frac{3}{2}}(c + dx)}$$

[In] Integrate[Cos[c + d*x]^3*(b*Cos[c + d*x])^(3/2),x]

[Out] ((b*Cos[c + d*x])^(3/2)*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Cos[c + d*x]^(3/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 222 vs. 2(107) = 214.

Time = 4.48 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.35

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}\right)}$

[In] int(cos(d*x+c)^3*(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/45*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2))*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-4*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.08

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{21i \sqrt{2} b^{\frac{3}{2}} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 21}{1}$$

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

```
[Out] 1/45*(21*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(5*b*cos(d*x + c)^3 + 7*b*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(3/2), x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2), x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^3, x)
```

Giac [F]

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(3/2), x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(b \cos(c + dx))^{3/2} dx = \int \cos(c + dx)^3 (b \cos(c + dx))^{3/2} dx$$

```
[In] int(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2), x)
```

```
[Out] int(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2), x)
```

3.80 $\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal result	416
Rubi [A] (verified)	416
Mathematica [A] (verified)	418
Maple [A] (verified)	418
Fricas [C] (verification not implemented)	418
Sympy [F(-1)]	419
Maxima [F]	419
Giac [F]	419
Mupad [F(-1)]	419

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b/d+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{5/2}}{7bd} + \frac{10b \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2*(b*\operatorname{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(10*b^2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (10*b*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(21*d) + (2*(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sin}[c + d*x])/(7*b*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^2} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} + \frac{5}{7} \int (b \cos(c + dx))^{3/2} dx \\
 &= \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} \\
 &\quad + \frac{1}{21} (5b^2) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd} \\
 &\quad + \frac{\left(5b^2 \sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21\sqrt{b \cos(c + dx)}} \\
 &= \frac{10b^2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{10b\sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7bd}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.74

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(b \cos(c + dx))^{3/2} \left(20 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)}(23 \sin(c + dx) + 3 \sin(3(c + dx))) \right)}{42d \cos^{\frac{3}{2}}(c + dx)}$$

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(3/2),x]

[Out] ((b*Cos[c + d*x])^(3/2)*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 3.41 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.14

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^2\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.93

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{-5i\sqrt{2}b^{\frac{3}{2}}\operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i\sqrt{2}b^{\frac{3}{2}}\operatorname{weierstrassPInverse}(\dots)}{21}$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{21}(-5I\sqrt{2}b^{3/2}\text{weierstrassPInverse}(-4, 0, \cos(dx + c)) + I\sin(dx + c)) + 5I\sqrt{2}b^{3/2}\text{weierstrassPInverse}(-4, 0, \cos(dx + c)) - I\sin(dx + c) + 2(3b\cos(dx + c)^2 + 5b)\sqrt{b\cos(dx + c)}\sin(dx + c))/d$

Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(3/2), x)`

[Out] Timed out

Maxima [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

[In] `integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^2, x)`

Giac [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

[In] `integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(3/2), x, algorithm="giac")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{3/2} dx = \int \cos(c + dx)^2 (b \cos(c + dx))^{3/2} dx$$

[In] `int(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2), x)`

[Out] `int(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2), x)`

3.81 $\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal result	420
Rubi [A] (verified)	420
Mathematica [A] (verified)	421
Maple [B] (verified)	422
Fricas [C] (verification not implemented)	422
Sympy [F(-1)]	423
Maxima [F]	423
Giac [F]	423
Mupad [F(-1)]	423

Optimal result

Integrand size = 19, antiderivative size = 67

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{6b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d}$$

[Out] $2/5*(b*\cos(d*x+c))^(3/2)*\sin(d*x+c)/d+6/5*b*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2715, 2721, 2719}

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{2 \sin(c + dx)(b \cos(c + dx))^{3/2}}{5d} + \frac{6bE\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]*(b*\text{Cos}[c + d*x])^(3/2),x]$

[Out] $(6*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^(3/2)*\text{Sin}[c + d*x])/(5*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{1}{5}(3b) \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{(3b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
 &= \frac{6b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.97

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(b \cos(c + dx))^{5/2} \left(6E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)} \sin(2(c + dx)) \right)}{5bd \cos^{5/2}(c + dx)}$$

`[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(3/2), x]`

`[Out] ((b*Cos[c + d*x])^(5/2)*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*b*d*Cos[c + d*x]^(5/2))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. $2(83) = 166$.

Time = 2.67 (sec) , antiderivative size = 213, normalized size of antiderivative = 3.18

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}b^2\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)$

[In] `int(cos(d*x+c)*(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/(2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.16 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.33

$$\int \cos(c+dx)(b\cos(c+dx))^{3/2} dx = \frac{2\sqrt{b\cos(dx+c)}b\cos(dx+c)\sin(dx+c)+3i\sqrt{2}b^{3/2}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+I*\sin(dx+c)))}{5\sqrt{2}b^{3/2}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+I*\sin(dx+c)))}-3I\sqrt{2}b^{3/2}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-I*\sin(dx+c)))}/d$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{b*\cos(dx+c)}*b*\cos(dx+c)*\sin(dx+c)+3*I*\sqrt{2}*b^{3/2}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+I*\sin(dx+c))))-3*I*\sqrt{2}*b^{3/2}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)-I*\sin(dx+c))))/d$$

Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c), x)

Giac [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*cos(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{3/2} dx = \int \cos(c + dx) (b \cos(c + dx))^{3/2} dx$$

[In] int(cos(c + d*x)*(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(3/2), x)

3.82 $\int (b \cos(c + dx))^{3/2} dx$

Optimal result	424
Rubi [A] (verified)	424
Mathematica [A] (verified)	425
Maple [B] (verified)	426
Fricas [C] (verification not implemented)	426
Sympy [F]	426
Maxima [F]	427
Giac [F]	427
Mupad [F(-1)]	427

Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (b \cos(c + dx))^{3/2} dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}$$

[Out] $2/3*b^2*(\cos(1/2*d*x+1/2*c))^2^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\int (b \cos(c + dx))^{3/2} dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(2*b^2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/((3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(3*d))$

Rule 2715


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} + \frac{1}{3}b^2 \int \frac{1}{\sqrt{b\cos(c+dx)}} dx \\ &= \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} + \frac{\left(b^2\sqrt{\cos(c+dx)}\right) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3\sqrt{b\cos(c+dx)}} \\ &= \frac{2b^2\sqrt{\cos(c+dx)}\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d\sqrt{b\cos(c+dx)}} + \frac{2b\sqrt{b\cos(c+dx)}\sin(c+dx)}{3d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.83

$$\int (b\cos(c+dx))^{3/2} dx = \frac{2(b\cos(c+dx))^{3/2} \left(\text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sqrt{\cos(c+dx)}\sin(c+dx) \right)}{3d\cos^{3/2}(c+dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(3/2),x]
```

```
[Out] (2*(b*Cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d*Cos[c + d*x]^(3/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(86) = 172$.

Time = 2.16 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.71

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}b^2\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$

[In] `int((cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/3*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(4*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticF}(\cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2)^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.10

$$\int (b \cos(c + dx))^{3/2} dx = \frac{-i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(\cos(dx + c) - i \sin(dx + c), -4, 0)}{3d}$$

[In] `integrate((b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$1/3*(-I*\sqrt{2}*b^{3/2}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*b^{3/2}*\text{weierstrassPInverse}(\cos(d*x + c) - I*\sin(d*x + c), -4, 0) + 2*\sqrt{b*\cos(d*x + c)}*b*\sin(d*x + c))/d$$

Sympy [F]

$$\int (b \cos(c + dx))^{3/2} dx = \int (b \cos(c + dx))^{3/2} dx$$

[In] `integrate((b*cos(d*x+c))**(3/2),x)`

[Out] `Integral((b*cos(c + d*x))**(3/2), x)`

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} dx$$

[In] integrate((b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int (b \cos(c + dx))^{3/2} dx = \int (b \cos(dx + c))^{\frac{3}{2}} dx$$

[In] integrate((b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} dx = \int (b \cos(c + dx))^{\frac{3}{2}} dx$$

[In] int((b*cos(c + d*x))^(3/2),x)

[Out] int((b*cos(c + d*x))^(3/2), x)

3.83 $\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx$

Optimal result	428
Rubi [A] (verified)	428
Mathematica [A] (verified)	429
Maple [B] (verified)	429
Fricas [C] (verification not implemented)	430
Sympy [F(-1)]	430
Maxima [F]	430
Giac [F]	431
Mupad [F(-1)]	431

Optimal result

Integrand size = 19, antiderivative size = 39

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}}$$

[Out] $2*b*(\cos(1/2*d*x+1/2*c))^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2})*(b*\cos(d*x+c))^{1/2}/d/\cos(d*x+c)^{1/2}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2721, 2719}

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \frac{2bE\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{3/2}*\text{Sec}[c + d*x], x]$

[Out] $(2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*)^{(n_*)}), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \&\& \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])ⁿ/Sin[c + d*x]ⁿ, Int[Sin[c + d*x]ⁿ, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{(b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x], x]

[Out] (2*b*Sqrt[b*Cos[c + d*x])*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(61) = 122.

Time = 1.80 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.69

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} b^2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1} E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right) b d}$
risch	$-\frac{i\sqrt{2}b\sqrt{(e^{2i(dx+c)}+1)be^{-i(dx+c)}}}{d} - i\left(\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}(-2iE(\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}))}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)$

[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c), x, method=_RETURNVERBOSE)

[Out] $2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.62

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \frac{i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i \sqrt{2} b^{3/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)))}{d}$$

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="fricas")`

[Out] $(I*\sqrt{2}*b^{(3/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - I*\sqrt{2}*b^{(3/2)}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/d$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c),x)`

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c) dx$$

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c), x)`

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x),x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x), x)

3.84 $\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx$

Optimal result	432
Rubi [A] (verified)	432
Mathematica [A] (verified)	433
Maple [B] (verified)	433
Fricas [C] (verification not implemented)	434
Sympy [F(-1)]	434
Maxima [F]	434
Giac [F]	435
Mupad [F(-1)]	435

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2720}

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[In] `Int[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^2,x]`

[Out] `(2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2720


```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
]*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b^2 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \frac{2b^2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^2,x]
```

```
[Out] (2*b^2*Sqrt[Cos[c + d*x])*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]
])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 1.27 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result	size
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^2\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)bd}}$	144

```
[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^2,x,method=_RETURNVERBOSE)
```

[Out] $-2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c), 2^{(1/2)})/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \frac{-i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{3/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="fricas")`

[Out] $(-I*\sqrt{2}*b^{(3/2)}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*b^{(3/2)}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)))/d$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**2,x)`

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^2 dx$$

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^2, x)`

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^2} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^2, x)

3.85 $\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx$

Optimal result	436
Rubi [A] (verified)	436
Mathematica [A] (verified)	437
Maple [B] (verified)	438
Fricas [C] (verification not implemented)	438
Sympy [F(-1)]	439
Maxima [F]	439
Giac [F]	439
Mupad [F(-1)]	439

Optimal result

Integrand size = 21, antiderivative size = 66

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = -\frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}$$

[Out] $2*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{2bE\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^3,x]$

[Out] $(-2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - b \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{(b\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2b\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.76

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \frac{2b^2 \left(-\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^3,x]
```

```
[Out] (2*b^2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d
*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(86) = 172$.

Time = 1.78 (sec) , antiderivative size = 198, normalized size of antiderivative = 3.00

method	result
default	$\frac{2b^2 \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b + b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)}{\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{\left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$

[In] `int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^3,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2*b^2*(-2*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)/d}}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.55

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \frac{-i \sqrt{2} b^{3/2} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} b^{3/2} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 \sqrt{b \cos(dx + c)} * b * \sin(dx + c) / (d * \cos(dx + c))}{1}$$

[In] `integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="fricas")`

[Out]
$$(-I*\sqrt{2}*b^{(3/2)*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + I*\sqrt{2}*b^{(3/2)*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*\sqrt{b*\cos(d*x + c)}*b*\sin(d*x + c))/(d*\cos(d*x + c))$$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**3,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^3, x)

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^3} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^3, x)

3.86 $\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx$

Optimal result	440
Rubi [A] (verified)	440
Mathematica [A] (verified)	441
Maple [B] (verified)	442
Fricas [C] (verification not implemented)	442
Sympy [F(-1)]	443
Maxima [F]	443
Giac [F]	443
Mupad [F(-1)]	443

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}$$

[Out] $2/3*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(3/2)+2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c),2^(1/2))*\cos(d*x+c)^(1/2)/d/(b*\cos(d*x+c))^(1/2)$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^(3/2)*\operatorname{Sec}[c + d*x]^4,x]$

[Out] $(2*b^2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/((3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^3*\operatorname{Sin}[c + d*x])/(3*d*(b*\operatorname{Cos}[c + d*x])^(3/2))$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^4 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} b^2 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\left(b^2 \sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2b^2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\begin{aligned}
 &\int (b \cos(c + dx))^{3/2} \sec^4(c \\
 &+ dx) dx = \frac{2b^2 \left(\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx) \right)}{3d\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

[In] Integrate[(b*cos[c + d*x])^(3/2)*Sec[c + d*x]^4,x]

[Out] (2*b^2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(88) = 176.

Time = 1.78 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.35

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^4,x,method=_RETURNVERBOSE)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*b^2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.40

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \frac{-i \sqrt{2} b^{3/2} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{3/2} \cos(dx + c)}{3 d \cos(dx + c)}$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2)*b^(3/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(3/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*b*sin(d*x + c))/(d*cos(d*x + c)^2)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**4,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^4 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^4, x)

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^4 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^4,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^4, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^4(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^4} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^4, x)

3.87 $\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx$

Optimal result	444
Rubi [A] (verified)	444
Mathematica [A] (verified)	446
Maple [B] (verified)	446
Fricas [C] (verification not implemented)	447
Sympy [F(-1)]	447
Maxima [F]	447
Giac [F]	448
Mupad [F(-1)]	448

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = -\frac{6b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}$$

[Out] $2/5*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \frac{2b^4 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^2 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{6bE\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{5d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sec}[c + d*x]^5,x]$

[Out] $(-6*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*b^2*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c+d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c-Pi/2+d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c+d*x])^n/Sin[c+d*x]^n, Int[Sin[c+d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^5 \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
 &= \frac{2b^4 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{1}{5} (3b^3) \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
 &= \frac{2b^4 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6b^2 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}} - \frac{1}{5} (3b) \int \sqrt{b \cos(c+dx)} dx \\
 &= \frac{2b^4 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6b^2 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}} - \frac{(3b\sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{5\sqrt{\cos(c+dx)}} \\
 &= -\frac{6b\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d\sqrt{\cos(c+dx)}} + \frac{2b^4 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6b^2 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.70

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \frac{(b \cos(c + dx))^{3/2} \sec^4(c + dx) \left(-12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10d}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^5,x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(110) = 220.

Time = 2.38 (sec) , antiderivative size = 365, normalized size of antiderivative = 3.72

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}b(24\cos(\frac{dx}{2} + \frac{c}{2})(\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}})E(\cos(\frac{dx}{2} + \frac{c}{2}))}{10d}$

[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^5,x,method=_RETURNVERBOSE)

[Out]
$$-2/5*(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b/\sin(1/2*d*x+1/2*c)^3/(8*\sin(1/2*d*x+1/2*c)^6-12*\sin(1/2*d*x+1/2*c)^4+6*\sin(1/2*d*x+1/2*c)^2-1)*(24*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-12*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^4-24*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+12*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2+8*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.21

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \frac{-3i \sqrt{2} b^{3/2} \cos(dx + c)^3 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)))}{(d \cos(dx + c))^3}$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*b^(3/2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*b^(3/2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*b*cos(d*x + c)^2 + b)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**5,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^5 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^5, x)

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^5 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^5,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^5, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^5(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^5} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^5,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^5, x)

3.88 $\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx$

Optimal result	449
Rubi [A] (verified)	449
Mathematica [A] (verified)	451
Maple [B] (verified)	451
Fricas [C] (verification not implemented)	452
Sympy [F(-1)]	452
Maxima [F]	452
Giac [F]	453
Mupad [F(-1)]	453

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}$$

[Out] $2/7*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^2 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(3/2)}*\operatorname{Sec}[c + d*x]^6, x]$

[Out] $(10*b^2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^5*\operatorname{Sin}[c + d*x])/(7*d*(b*\operatorname{Cos}[c + d*x])^{(7/2)}) + (10*b^3*\operatorname{Sin}[c + d*x])/(21*d*(b*\operatorname{Cos}[c + d*x])^{(3/2)})$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^6 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{1}{7} (5b^4) \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{1}{21} (5b^2) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{(5b^2 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
 &= \frac{10b^2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^3 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.69

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \frac{(b \cos(c + dx))^{3/2} \sec^4(c + dx) \left(10 \cos^{\frac{5}{2}}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21d}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^6,x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^4*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(112) = 224.

Time = 2.14 (sec) , antiderivative size = 398, normalized size of antiderivative = 3.98

method	result
default	$-\frac{2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} F \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 40 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) + 60 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \right)}{\dots}$

[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^6,x,method=_RETURNVERBOSE)

[Out]
$$-2/21*(-40*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^6-40*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+60*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^4+40*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-30*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2-16*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+5*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^{2-1})^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)}))*b^2*((2*\cos(1/2*d*x+1/2*c)^{2-1})*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^{2-1})^3/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^{2-1})*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.15

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \frac{-5i \sqrt{2} b^{3/2} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} b^{3/2} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2(5b \cos(dx + c)^2 + 3b \sqrt{b \cos(dx + c)} \sin(dx + c)) / (d \cos(dx + c)^4)}{1}$$

```
[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="fricas")
```

```
[Out] 1/21*(-5*I*sqrt(2)*b^(3/2)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*b^(3/2)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(5*b*cos(d*x + c)^2 + 3*b*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^4)
```

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \text{Timed out}$$

```
[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**6,x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^6 dx$$

```
[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^6, x)
```

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^6 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^6,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^6, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^6(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^6} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^6,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^6, x)

3.89 $\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx$

Optimal result	454
Rubi [A] (verified)	454
Mathematica [A] (verified)	456
Maple [B] (verified)	456
Fricas [C] (verification not implemented)	457
Sympy [F(-1)]	457
Maxima [F]	457
Giac [F]	458
Mupad [F(-1)]	458

Optimal result

Integrand size = 21, antiderivative size = 126

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = -\frac{14b\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} + \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}}$$

[Out] $2/9*b^6*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*b*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \frac{2b^6 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^2 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{14bE\left(\frac{1}{2}(c + dx) \mid 2\right)\sqrt{b \cos(c + dx)}}{15d\sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(3/2)*\text{Sec}[c + d*x]^7,x]$

[Out] $(-14*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/((15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^6*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b^4*\text{Sin}[c + d*x])/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*b^2*\text{Sin}[c + d*x])/(15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c+d*x]*((b*\text{Sin}[c+d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c+d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c+d*x])^n/\text{Sin}[c+d*x]^n, \text{Int}[\text{Sin}[c+d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^7 \int \frac{1}{(b \cos(c+dx))^{11/2}} dx \\
 &= \frac{2b^6 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{1}{9}(7b^5) \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
 &= \frac{2b^6 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^4 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{1}{15}(7b^3) \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
 &= \frac{2b^6 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^4 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} \\
 &\quad + \frac{14b^2 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{1}{15}(7b) \int \sqrt{b \cos(c+dx)} dx \\
 &= \frac{2b^6 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^4 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} \\
 &\quad + \frac{14b^2 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{(7b\sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{15\sqrt{\cos(c+dx)}} \\
 &= -\frac{14b\sqrt{b \cos(c+dx)}E(\frac{1}{2}(c+dx)|2)}{15d\sqrt{\cos(c+dx)}} + \frac{2b^6 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} \\
 &\quad + \frac{14b^4 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14b^2 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.63

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \frac{(b \cos(c + dx))^{3/2} \sec^6(c + dx) \left(-336 \cos^{\frac{9}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) + 150 \sin(c + dx) + 91 \sin(3(c + dx)) \right) + 21 \sin(5(c + dx))}{360d}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^7,x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sec[c + d*x]^6*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)]))/(360*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. 2(134) = 268.

Time = 2.99 (sec) , antiderivative size = 416, normalized size of antiderivative = 3.30

method	result
default	$- \frac{2 \sqrt{-\left(-2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right) b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b^2 \left(- \frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{144 b \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^5} - \frac{7 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{180 b \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^5}}{\dots}$

[In] int((cos(d*x+c)*b)^(3/2)*sec(d*x+c)^7,x,method=_RETURNVERBOSE)

[Out] -2*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2*(-1/144*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/180*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^3-14/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)+7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 132, normalized size of antiderivative = 1.05

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \frac{-21i \sqrt{2} b^{3/2} \cos(dx + c)^5 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 21i \sqrt{2} b^{3/2} \cos(dx + c)^5 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2(21b^2 \cos(dx + c)^4 + 7b^2 \cos(dx + c)^2 + 5b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^5}$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*b^(3/2)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*b^(3/2)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*b*cos(d*x + c)^4 + 7*b*cos(d*x + c)^2 + 5*b)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)*sec(d*x+c)**7,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \int (b \cos(dx + c))^{3/2} \sec(dx + c)^7 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^7, x)

Giac [F]

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \int (b \cos(dx + c))^{\frac{3}{2}} \sec(dx + c)^7 dx$$

[In] integrate((b*cos(d*x+c))^(3/2)*sec(d*x+c)^7,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)*sec(d*x + c)^7, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{3/2} \sec^7(c + dx) dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^7} dx$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^7,x)

[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^7, x)

3.90 $\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal result	459
Rubi [A] (verified)	459
Mathematica [A] (verified)	461
Maple [A] (verified)	461
Fricas [C] (verification not implemented)	462
Sympy [F(-1)]	462
Maxima [F]	462
Giac [F]	463
Mupad [F(-1)]	463

Optimal result

Integrand size = 21, antiderivative size = 125

$$\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{30b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{30b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d}$$

[Out] $18/77*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/d+2/11*(b*\cos(d*x+c))^{(9/2)}*\sin(d*x+c)/b^2/d+30/77*b^3*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+30/77*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{30b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{77d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{9/2}}{11b^2d} + \frac{30b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{77d} + \frac{18 \sin(c + dx)(b \cos(c + dx))^{5/2}}{77d}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^3*(b*\operatorname{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(30*b^3*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2])/(77*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (30*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(77*d) + (18*(b*\text{Cos}[c + d*x])^{5/2}*\text{Sin}[c + d*x])/(77*d) + (2*(b*\text{Cos}[c + d*x])^{9/2}*\text{Sin}[c + d*x])/(11*b^2*d)$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*)^{(n_*)}), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

$\text{Int}[(b_*)*\text{sin}[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\text{sin}[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d, x\}$

Rule 2721

$\text{Int}[(b_*)*\text{sin}[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^3} \\ &= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b} \\ &= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d} \\ &\quad + \frac{1}{77}(45b) \int (b \cos(c + dx))^{3/2} dx \\ &= \frac{30b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{77d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77d} \\ &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^2d} + \frac{1}{77}(15b^3) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \end{aligned}$$

$$\begin{aligned}
&= \frac{30b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77d} \\
&\quad + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^2d} + \frac{\left(15b^3 \sqrt{\cos(c+dx)}\right) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{77\sqrt{b \cos(c+dx)}} \\
&= \frac{30b^3 \sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77d\sqrt{b \cos(c+dx)}} + \frac{30b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{77d} \\
&\quad + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77d} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^2d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.36 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.66

$$\int \cos^3(c+dx)(b \cos(c+dx))^{5/2} dx = \frac{(b \cos(c+dx))^{5/2} \left(240 \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + \sqrt{\cos(c+dx)}(290 \sin(c+dx) + 57 \sin(3(c+dx)))\right)}{616d \cos^{\frac{5}{2}}(c+dx)}$$

[In] Integrate[Cos[c + d*x]^3*(b*Cos[c + d*x])^(5/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(240*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(290*Sin[c + d*x] + 57*Sin[3*(c + d*x)] + 7*Sin[5*(c + d*x)]))/616*d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 10.46 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.89

method	result
default	$ -\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}b^3\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1084\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-370\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15\left(\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\right)^{1/2}\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^{1/2}\operatorname{EllipticF}\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right), 2\right)^{1/2}+62\cos\left(\frac{dx}{2}+\frac{c}{2}\right)}{77\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\right)}} $

[In] int(cos(d*x+c)^3*(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/77*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(448*cos(1/2*d*x+1/2*c)^13-1568*cos(1/2*d*x+1/2*c)^11+2384*cos(1/2*d*x+1/2*c)^9-2040*cos(1/2*d*x+1/2*c)^7+1084*cos(1/2*d*x+1/2*c)^5-370*cos(1/2*d*x+1/2*c)^3+15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2)^(1/2)+62*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.86

$$\int \cos^3(c + dx)(b \cos(c + dx))^5 dx = \frac{-15i \sqrt{2} b^{\frac{5}{2}} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 15i \sqrt{2} b^{\frac{5}{2}} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

```
[Out] 1/77*(-15*I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 15*I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(7*b^2*cos(d*x + c)^4 + 9*b^2*cos(d*x + c)^2 + 15*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(b \cos(c + dx))^5 dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**3*(b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^3(c + dx)(b \cos(c + dx))^5 dx = \int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^3 dx$$

```
[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^3, x)
```

Giac [F]

$$\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^3 dx$$

[In] integrate(cos(d*x+c)^3*(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^3(c + dx)(b \cos(c + dx))^{5/2} dx = \int \cos(c + dx)^3 (b \cos(c + dx))^{5/2} dx$$

[In] int(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2), x)

3.91 $\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal result	464
Rubi [A] (verified)	464
Mathematica [A] (verified)	466
Maple [B] (verified)	466
Fricas [C] (verification not implemented)	466
Sympy [F(-1)]	467
Maxima [F]	467
Giac [F]	467
Mupad [F(-1)]	467

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{14b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d \sqrt{\cos(c + dx)}} + \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd}$$

[Out] $14/45*b*(b*\cos(d*x+c))^{3/2}*\sin(d*x+c)/d+2/9*(b*\cos(d*x+c))^{7/2}*\sin(d*x+c)/b/d+14/15*b^2*(\cos(1/2*d*x+1/2*c)^2)^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2})*(b*\cos(d*x+c))^{1/2}/d/\cos(d*x+c)^{1/2}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{14b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{15d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{7/2}}{9bd} + \frac{14b \sin(c + dx)(b \cos(c + dx))^{3/2}}{45d}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^2*(b*\text{Cos}[c + d*x])^{5/2}, x]$

[Out] $(14*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (14*b*(b*\text{Cos}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(45*d) + (2*(b*\text{Cos}[c + d*x])^{7/2}*\text{Sin}[c + d*x])/(9*b*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^2} \\
 &= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} + \frac{7}{9} \int (b \cos(c + dx))^{5/2} dx \\
 &= \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} \\
 &\quad + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} + \frac{1}{15} (7b^2) \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd} \\
 &\quad + \frac{\left(7b^2 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)}} \\
 &= \frac{14b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d\sqrt{\cos(c + dx)}} \\
 &\quad + \frac{14b(b \cos(c + dx))^{3/2} \sin(c + dx)}{45d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9bd}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{(b \cos(c + dx))^{5/2} \left(168E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)}(38 \sin(2(c + dx)) + 5 \sin(4(c + dx))) \right)}{180d \cos^{5/2}(c + dx)}$$

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(5/2),x]

[Out] ((b*Cos[c + d*x])^(5/2)*(168*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)])))/(180*d*Cos[c + d*x]^(5/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 222 vs. 2(110) = 220.

Time = 5.66 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.28

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^3\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}\right)}$

[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] -2/45*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))-4*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.09

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{21i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - 21}{1}$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

```
[Out] 1/45*(21*I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(5*b^2*cos(d*x + c)^3 + 7*b^2*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c))/d
```

Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(5/2), x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

```
[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2), x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^2, x)
```

Giac [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

```
[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(5/2), x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{5/2} dx = \int \cos(c + dx)^2 (b \cos(c + dx))^{5/2} dx$$

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2), x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2), x)
```

3.92 $\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal result	468
Rubi [A] (verified)	468
Mathematica [A] (verified)	470
Maple [A] (verified)	470
Fricas [C] (verification not implemented)	470
Sympy [F(-1)]	471
Maxima [F]	471
Giac [F]	471
Mupad [F(-1)]	471

Optimal result

Integrand size = 19, antiderivative size = 97

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{10b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/d+10/21*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2715, 2721, 2720}

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{10b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{2 \sin(c + dx)(b \cos(c + dx))^{5/2}}{7d}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]*(b*\operatorname{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(10*b^3*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (10*b^2*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(21*d) + (2*(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sin}[c + d*x])/(7*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{7}(5b) \int (b \cos(c + dx))^{3/2} dx \\
 &= \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} \\
 &\quad + \frac{1}{21}(5b^3) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} \\
 &\quad + \frac{(5b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
 &= \frac{10b^3 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{10b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.78

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \cos(c + dx)} \left(20 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)}(23 \sin(c + dx) + 3 \sin(3(c + dx))) \right)}{42d \sqrt{\cos(c + dx)}}$$

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(5/2),x]

[Out] (b^2*Sqrt[b*Cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.61 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.16

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^3\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)}}$

[In] int(cos(d*x+c)*(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.98

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{-5i \sqrt{2} b^{5/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} b^{5/2} \operatorname{weierstrassPInverse}(\dots)}{2}$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] $1/21*(-5*I*\sqrt{2}*b^{(5/2)}*weierstrassPInverse(-4, 0, \cos(dx + c) + I*\sin(dx + c)) + 5*I*\sqrt{2}*b^{(5/2)}*weierstrassPInverse(-4, 0, \cos(dx + c) - I*\sin(dx + c)) + 2*(3*b^2*\cos(dx + c)^2 + 5*b^2)*\sqrt{b*\cos(dx + c)}*\sin(dx + c))/d$

Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{5/2} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c), x)`

Giac [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{5/2} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(5/2),x, algorithm="giac")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*cos(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{5/2} dx = \int \cos(c + dx) (b \cos(c + dx))^{5/2} dx$$

[In] `int(cos(c + d*x)*(b*cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)*(b*cos(c + d*x))^(5/2), x)`

3.93 $\int (b \cos(c + dx))^{5/2} dx$

Optimal result	472
Rubi [A] (verified)	472
Mathematica [A] (verified)	473
Maple [B] (verified)	473
Fricas [C] (verification not implemented)	474
Sympy [F(-1)]	474
Maxima [F]	474
Giac [F]	475
Mupad [F(-1)]	475

Optimal result

Integrand size = 12, antiderivative size = 70

$$\int (b \cos(c+dx))^{5/2} dx = \frac{6b^2 \sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5d \sqrt{\cos(c+dx)}} + \frac{2b(b \cos(c+dx))^{3/2} \sin(c+dx)}{5d}$$

[Out] $2/5*b*(b*\cos(d*x+c))^(3/2)*\sin(d*x+c)/d+6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2719}

$$\int (b \cos(c+dx))^{5/2} dx = \frac{6b^2 E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5d \sqrt{\cos(c+dx)}} + \frac{2b \sin(c+dx)(b \cos(c+dx))^{3/2}}{5d}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(5/2), x]$

[Out] $(6*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b*(b*\text{Cos}[c + d*x])^(3/2)*\text{Sin}[c + d*x])/(5*d)$

Rule 2715

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^(n-1)/(d*n)), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^(n-2), x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{1}{5}(3b^2) \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} + \frac{(3b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\ &= \frac{6b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b(b \cos(c + dx))^{3/2} \sin(c + dx)}{5d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.89

$$\int (b \cos(c + dx))^{5/2} dx = \frac{(b \cos(c + dx))^{5/2} \left(6E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sqrt{\cos(c + dx)} \sin(2(c + dx)) \right)}{5d \cos^{5/2}(c + dx)}$$

[In] `Integrate[(b*Cos[c + d*x])^(5/2),x]`

[Out] `((b*Cos[c + d*x])^(5/2)*(6*EllipticE[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[2*(c + d*x)]))/(5*d*Cos[c + d*x]^(5/2))`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(86) = 172.

Time = 2.85 (sec) , antiderivative size = 213, normalized size of antiderivative = 3.04

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}} b^3 \left(-8 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \left(\sin^6\left(\frac{dx}{2} + \frac{c}{2}\right) \right) + 8 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \cos\left(\frac{dx}{2} + \frac{c}{2}\right) - 2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right) \right) \right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}$

```
[In] int((cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2/5*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(-8*cos(
1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+8*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*
c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)
*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2)))/(-
b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/(
(2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.30

$$\int (b \cos(c + dx))^{5/2} dx = \frac{2 \sqrt{b \cos(dx + c)} b^2 \cos(dx + c) \sin(dx + c) + 3i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + I \sin(dx + c))) - 3i \sqrt{2} b^{5/2} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - I \sin(dx + c)))}{d}$$

```
[In] integrate((b*cos(d*x+c))^(5/2),x, algorithm="fricas")
```

```
[Out] 1/5*(2*sqrt(b*cos(d*x + c))*b^2*cos(d*x + c)*sin(d*x + c) + 3*I*sqrt(2)*b^(
5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin
(d*x + c))) - 3*I*sqrt(2)*b^(5/2)*weierstrassZeta(-4, 0, weierstrassPInvers
e(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/d
```

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} dx = \text{Timed out}$$

```
[In] integrate((b*cos(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{5/2} dx$$

```
[In] integrate((b*cos(d*x+c))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(5/2), x)
```

Giac [F]

$$\int (b \cos(c + dx))^{5/2} dx = \int (b \cos(dx + c))^{5/2} dx$$

[In] integrate((b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} dx = \int (b \cos(c + dx))^{5/2} dx$$

[In] int((b*cos(c + d*x))^(5/2),x)

[Out] int((b*cos(c + d*x))^(5/2), x)

3.94 $\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx$

Optimal result	476
Rubi [A] (verified)	476
Mathematica [A] (verified)	477
Maple [B] (verified)	478
Fricas [C] (verification not implemented)	478
Sympy [F(-1)]	479
Maxima [F]	479
Giac [F]	479
Mupad [F(-1)]	479

Optimal result

Integrand size = 19, antiderivative size = 72

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}$$

[Out] $2/3*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2715, 2721, 2720}

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3d}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sec}[c + d*x], x]$

[Out] $(2*b^3*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/((3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^2*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(3*d))$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b \int (b \cos(c + dx))^{3/2} dx \\
 &= \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{1}{3} b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d} + \frac{(b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2b^3 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.82

$$\begin{aligned}
 &\int (b \cos(c + dx))^{5/2} \sec(c \\
 &+ dx) dx = \frac{2b(b \cos(c + dx))^{3/2} \left(\text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)} \sin(c + dx) \right)}{3d \cos^{3/2}(c + dx)}
 \end{aligned}$$

[In] Integrate[(b*cos[c + d*x])^(5/2)*Sec[c + d*x],x]

[Out] (2*b*(b*cos[c + d*x])^(3/2)*(EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*Sin[c + d*x]))/(3*d*cos[c + d*x]^(3/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(88) = 176.

Time = 2.77 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.64

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^3\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)\sqrt{2}}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)}bd}$

[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c),x,method=_RETURNVERBOSE)

[Out] -2/3*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(4*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.10

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \frac{-i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{3d}$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c),x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(5/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*b^2*sin(d*x + c))/d

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c), x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c), x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c), x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c), x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x), x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x), x)

3.95 $\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx$

Optimal result	480
Rubi [A] (verified)	480
Mathematica [A] (verified)	481
Maple [B] (verified)	481
Fricas [C] (verification not implemented)	482
Sympy [F(-1)]	482
Maxima [F]	483
Giac [F]	483
Mupad [F(-1)]	483

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)}}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2719}

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] `Int[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^2,x]`

[Out] `(2*b^2*Sqrt[b*Cos[c + d*x])*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]])`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2719


```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \sqrt{b \cos(c + dx)} dx \\ &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^2,x]
```

```
[Out] (2*b^2*Sqrt[b*Cos[c + d*x])*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]
])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 6.79 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result
default	$\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b^3\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}bd}$
risch	$-\frac{i\sqrt{2}b^2\sqrt{(e^{2i(dx+c)}+1)be^{-i(dx+c)}}}{d} - i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}} + \frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)\left(-2iE\left(\sqrt{-\frac{1}{2}\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1},\sqrt{2}\right)\right)$

[In] `int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^2,x,method=_RETURNVERBOSE)`

[Out] $2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*b^3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*\cos(1/2*d*x+1/2*c)^2+1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.54

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \frac{i\sqrt{2}b^{5/2}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) - i\sqrt{2}b^{5/2}\text{weierstrassZeta}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

[In] `integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="fricas")`

[Out] $(I*\text{sqrt}(2)*b^{5/2}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) - I*\text{sqrt}(2)*b^{5/2}*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))))/d$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**2,x)`

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^2, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^2} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^2,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^2, x)

3.96 $\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx$

Optimal result	484
Rubi [A] (verified)	484
Mathematica [A] (verified)	485
Maple [B] (verified)	485
Fricas [C] (verification not implemented)	486
Sympy [F(-1)]	486
Maxima [F]	486
Giac [F]	487
Mupad [F(-1)]	487

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[Out] $2*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2720}

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[In] `Int[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^3,x]`

[Out] `(2*b^3*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
]*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{(b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{2b^3 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \frac{2(b \cos(c + dx))^{5/2} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{d \cos^{5/2}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^3,x]
```

```
[Out] (2*(b*Cos[c + d*x])^(5/2)*EllipticF[(c + d*x)/2, 2])/(d*Cos[c + d*x]^(5/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 27.80 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result	size
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^3\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}F\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)bd}$	144

```
[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^3,x,method=_RETURNVERBOSE)
```

[Out] $-2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*b^3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.39

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \frac{-i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{5/2} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{d}$$

[In] `integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="fricas")`

[Out] $(-I*\sqrt{2}*b^{(5/2)}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*b^{(5/2)}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)))/d$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**3,x)`

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^3 dx$$

[In] `integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^3, x)`

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^3} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^3,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^3, x)

3.97 $\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx$

Optimal result	488
Rubi [A] (verified)	488
Mathematica [A] (verified)	489
Maple [B] (verified)	490
Fricas [C] (verification not implemented)	490
Sympy [F(-1)]	491
Maxima [F]	491
Giac [F]	491
Mupad [F(-1)]	491

Optimal result

Integrand size = 21, antiderivative size = 68

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = -\frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d \sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{d \sqrt{b \cos(c + dx)}}$$

[Out] $2*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \frac{2b^3 \sin(c + dx)}{d \sqrt{b \cos(c + dx)}} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(5/2)}*\text{Sec}[c + d*x]^4,x]$

[Out] $(-2*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)^{(m_*)}*(v_*)^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{b, n, x\} \ \&\amp; \ \text{IntegerQ}[m]$

Rule 2716


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b^4 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - b^2 \int \sqrt{b \cos(c + dx)} dx \\
&= \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}} - \frac{(b^2 \sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)}} \\
&= -\frac{2b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{d\sqrt{\cos(c + dx)}} + \frac{2b^3 \sin(c + dx)}{d\sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \frac{2b^3 \left(-\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^4,x]
```

```
[Out] (2*b^3*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d
*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(88) = 176$.

Time = 98.97 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.91

method	result
default	$\frac{2b^3 \left(-2 \cos\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) b + b \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \sqrt{-2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} \right)}{\sqrt{-b \left(2 \left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right) \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{\left(2 \left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$

[In] `int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^4,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2b^3(-2\cos(1/2dx+1/2c)*(-2\sin(1/2dx+1/2c)^4+b\sin(1/2dx+1/2c)^2)^{(1/2)}\sin(1/2dx+1/2c)^2+(\sin(1/2dx+1/2c)^2)^{(1/2)}*(2\sin(1/2dx+1/2c)^2-1)^{(1/2)}*(-2\sin(1/2dx+1/2c)^4+b\sin(1/2dx+1/2c)^2)^{(1/2)}*EllipticE(\cos(1/2dx+1/2c),2^{(1/2)})/(-b*(2\sin(1/2dx+1/2c)^4-\sin(1/2dx+1/2c)^2))^{(1/2)}/\sin(1/2dx+1/2c)/((2\cos(1/2dx+1/2c)^2-1)*b)^{(1/2)}/d}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.53

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \frac{-i \sqrt{2} b^{5/2} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + i \sqrt{2} b^{5/2} \cos(dx + c) \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 \sqrt{b \cos(dx + c)} b^2 \sin(dx + c) / (d \cos(dx + c))}{1}$$

[In] `integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="fricas")`

[Out]
$$(-I*\sqrt{2}*b^{(5/2)}*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + I*\sqrt{2}*b^{(5/2)}*\cos(d*x + c)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*\sqrt{b*\cos(d*x + c)}*b^2*\sin(d*x + c)/(d*\cos(d*x + c))$$

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**4,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^4 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^4, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^4 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^4,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^4, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^4(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^4} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^4,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^4, x)

3.98 $\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx$

Optimal result	492
Rubi [A] (verified)	492
Mathematica [A] (verified)	493
Maple [B] (verified)	494
Fricas [C] (verification not implemented)	494
Sympy [F(-1)]	495
Maxima [F]	495
Giac [F]	495
Mupad [F(-1)]	495

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}$$

[Out] $2/3*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^(3/2)+2/3*b^3*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c),2^(1/2))*\cos(d*x+c)^(1/2)/d/(b*\cos(d*x+c))^(1/2)$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d \sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^(5/2)*\operatorname{Sec}[c + d*x]^5,x]$

[Out] $(2*b^3*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^4*\operatorname{Sin}[c + d*x])/(3*d*(b*\operatorname{Cos}[c + d*x])^(3/2))$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^5 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} b^3 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\left(b^3 \sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b^4 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\begin{aligned}
 &\int (b \cos(c + dx))^{5/2} \sec^5(c \\
 &+ dx) dx = \frac{2b^3 \left(\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx) \right)}{3d\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

[In] Integrate[(b*cos[c + d*x])^(5/2)*Sec[c + d*x]^5,x]

[Out] (2*b^3*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt[b*cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(88) = 176.

Time = 1.14 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.35

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} F \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}{3 \sqrt{-b \left(2 \left(\sin^4 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) \right)} \left(2 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \right)}$$

[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^5,x)

[Out] -2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*b^3*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.43

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \frac{-i \sqrt{2} b^{5/2} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} b^{5/2} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 \sqrt{b \cos(dx + c)} b^2 \sin(dx + c)}{3 d \cos(dx + c)}$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2)*b^(5/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*b^(5/2)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*b^2*sin(d*x + c))/(d*cos(d*x + c)^2)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**5,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^5 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^5, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^5 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^5,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^5, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^5(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^5} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^5,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^5, x)

3.99 $\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx$

Optimal result	496
Rubi [A] (verified)	496
Mathematica [A] (verified)	498
Maple [B] (verified)	498
Fricas [C] (verification not implemented)	499
Sympy [F(-1)]	499
Maxima [F]	499
Giac [F]	500
Mupad [F(-1)]	500

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = -\frac{6b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d \sqrt{\cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d \sqrt{b \cos(c + dx)}}$$

[Out] $2/5*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+6/5*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d \sqrt{b \cos(c + dx)}} - \frac{6b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(5/2)*\text{Sec}[c + d*x]^6,x]$

[Out] $(-6*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x])*E(\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^5*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^(5/2)) + (6*b^3*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^6 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5}(3b^4) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{1}{5}(3b^2) \int \sqrt{b \cos(c + dx)} dx \\
 &= \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{\left(3b^2 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)}} \\
 &= -\frac{6b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5d\sqrt{\cos(c + dx)}} + \frac{2b^5 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6b^3 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.69

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \frac{(b \cos(c + dx))^{5/2} \sec^5(c + dx) \left(-12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 7 \sin(c + dx) + 3 \sin(3(c + dx)) \right)}{10d}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^6,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^5*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(112) = 224.

Time = 1.67 (sec) , antiderivative size = 367, normalized size of antiderivative = 3.67

$$2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1) b (\sin^2(\frac{dx}{2} + \frac{c}{2}))} b^2 \left(24 \cos(\frac{dx}{2} + \frac{c}{2}) (\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12\sqrt{2} (\sin^2(\frac{dx}{2} + \frac{c}{2})) \right)$$

[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^6,x)

[Out] -2/5*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^2/sin(1/2*d*x+1/2*c)^3/(8*sin(1/2*d*x+1/2*c)^6-12*sin(1/2*d*x+1/2*c)^4+6*sin(1/2*d*x+1/2*c)^2-1)*(24*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6-12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4-24*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)+12*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2+8*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)-3*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))*(-2*sin(1/2*d*x+1/2*c)^4*b+b*sin(1/2*d*x+1/2*c)^2)^(1/2)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.23

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \frac{-3i \sqrt{2} b^{5/2} \cos(dx + c)^3 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 3i \sqrt{2} b^{5/2} \cos(dx + c)^3 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2(3b^2 \cos(dx + c)^2 + b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^3}$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*b^(5/2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*b^(5/2)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(3*b^2*cos(d*x + c)^2 + b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^3)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**6,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^6 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^6, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^6 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^6,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^6, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^6(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^6} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^6,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^6, x)

3.100 $\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx$

Optimal result	501
Rubi [A] (verified)	501
Mathematica [A] (verified)	503
Maple [B] (verified)	503
Fricas [C] (verification not implemented)	504
Sympy [F(-1)]	504
Maxima [F]	504
Giac [F]	505
Mupad [F(-1)]	505

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \frac{10b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}$$

[Out] $2/7*b^6*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{10b^3 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sec}[c + d*x]^7, x]$

[Out] $(10*b^3*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b^6*\operatorname{Sin}[c + d*x])/(7*d*(b*\operatorname{Cos}[c + d*x])^{(7/2)}) + (10*b^4*\operatorname{Sin}[c + d*x])/(21*d*(b*\operatorname{Cos}[c + d*x])^{(3/2)})$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^7 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
 &= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{1}{7}(5b^5) \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{1}{21}(5b^3) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}} + \frac{(5b^3 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
 &= \frac{10b^3 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{2b^6 \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10b^4 \sin(c + dx)}{21d(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.68 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.69

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \frac{(b \cos(c + dx))^{5/2} \sec^5(c + dx) \left(10 \cos^{5/2}(c + dx) \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 5 \sin(2(c + dx)) + 6 \tan(c + dx) \right)}{21d}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^7,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^5*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(112) = 224.

Time = 1.06 (sec) , antiderivative size = 398, normalized size of antiderivative = 3.98

$$2 \left(-40 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 1} F \left(\cos \left(\frac{dx}{2} + \frac{c}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{dx}{2} + \frac{c}{2} \right) \right) - 40 \cos \left(\frac{dx}{2} + \frac{c}{2} \right) \left(\sin \left(\frac{dx}{2} + \frac{c}{2} \right) \right)^5 \right)$$

[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^7,x)

[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^6-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^4+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))*b^3*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.19

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \frac{-5i \sqrt{2} b^{5/2} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} b^{5/2} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2(5b^2 \cos(dx + c)^2 + 3b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{(d \cos(dx + c))^4}$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="fricas")

[Out] 1/21*(-5*I*sqrt(2)*b^(5/2)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*b^(5/2)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(5*b^2*cos(d*x + c)^2 + 3*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^4)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**7,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^7 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^7, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^7 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^7,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^7, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^7(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^7} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^7,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^7, x)

3.101 $\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx$

Optimal result	506
Rubi [A] (verified)	506
Mathematica [A] (verified)	508
Maple [F(-1)]	508
Fricas [C] (verification not implemented)	508
Sympy [F(-1)]	509
Maxima [F]	509
Giac [F]	509
Mupad [F(-1)]	509

Optimal result

Integrand size = 21, antiderivative size = 128

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = -\frac{14b^2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15d \sqrt{\cos(c + dx)}} + \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d \sqrt{b \cos(c + dx)}}$$

[Out] $2/9*b^7*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^5*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \frac{2b^7 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^5 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14b^3 \sin(c + dx)}{15d \sqrt{b \cos(c + dx)}} - \frac{14b^2 E\left(\frac{1}{2}(c + dx) \mid 2\right) \sqrt{b \cos(c + dx)}}{15d \sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^(5/2)*\text{Sec}[c + d*x]^8,x]$

[Out] $(-14*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/ (15*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^7*\text{Sin}[c + d*x])/ (9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b^5*\text{Sin}[c + d*x])/ (45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*b^3*\text{Sin}[c + d*x])/ (15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c+d*x]*((b*\text{Sin}[c+d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c+d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c+d*x])^n/\text{Sin}[c+d*x]^n, \text{Int}[\text{Sin}[c+d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^8 \int \frac{1}{(b \cos(c+dx))^{11/2}} dx \\
 &= \frac{2b^7 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{1}{9}(7b^6) \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
 &= \frac{2b^7 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^5 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{1}{15}(7b^4) \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
 &= \frac{2b^7 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^5 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} \\
 &\quad + \frac{14b^3 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{1}{15}(7b^2) \int \sqrt{b \cos(c+dx)} dx \\
 &= \frac{2b^7 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^5 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} \\
 &\quad + \frac{14b^3 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{(7b^2 \sqrt{b \cos(c+dx)}) \int \sqrt{\cos(c+dx)} dx}{15\sqrt{\cos(c+dx)}} \\
 &= -\frac{14b^2 \sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15d\sqrt{\cos(c+dx)}} + \frac{2b^7 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} \\
 &\quad + \frac{14b^5 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14b^3 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.85 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.62

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \frac{(b \cos(c + dx))^{5/2} \sec^7(c + dx) \left(-336 \cos^{\frac{9}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \mid 2\right) + 150 \sin(c + dx) + 91 \sin(3(c + dx)) \right) + 21 \sin(5(c + dx))}{360d}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^8,x]

[Out] ((b*Cos[c + d*x])^(5/2)*Sec[c + d*x]^7*(-336*Cos[c + d*x]^(9/2)*EllipticE[(c + d*x)/2, 2] + 150*Sin[c + d*x] + 91*Sin[3*(c + d*x)] + 21*Sin[5*(c + d*x)]))/(360*d)

Maple [F(-1)]

Timed out.

hanged

[In] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^8,x)

[Out] int((cos(d*x+c)*b)^(5/2)*sec(d*x+c)^8,x)

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 138, normalized size of antiderivative = 1.08

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \frac{-21i \sqrt{2} b^{\frac{5}{2}} \cos(dx + c)^5 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + 21i \sqrt{2} b^{\frac{5}{2}} \cos(dx + c)^5 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2*(21*b^2*\cos(d*x + c)^4 + 7*b^2*\cos(d*x + c)^2 + 5*b^2)*\sqrt{b*\cos(d*x + c)}*\sin(d*x + c)}{(d*\cos(d*x + c))^5}$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*b^(5/2)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*b^(5/2)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*b^2*cos(d*x + c)^4 + 7*b^2*cos(d*x + c)^2 + 5*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^5)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)*sec(d*x+c)**8,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^8 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^8, x)

Giac [F]

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \int (b \cos(dx + c))^{5/2} \sec(dx + c)^8 dx$$

[In] integrate((b*cos(d*x+c))^(5/2)*sec(d*x+c)^8,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)*sec(d*x + c)^8, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{5/2} \sec^8(c + dx) dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^8} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^8,x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^8, x)

3.102 $\int (b \cos(c + dx))^{7/2} dx$

Optimal result	510
Rubi [A] (verified)	510
Mathematica [A] (verified)	512
Maple [A] (verified)	512
Fricas [C] (verification not implemented)	512
Sympy [F(-1)]	513
Maxima [F]	513
Giac [F]	513
Mupad [F(-1)]	513

Optimal result

Integrand size = 12, antiderivative size = 98

$$\int (b \cos(c + dx))^{7/2} dx = \frac{10b^4 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}$$

[Out] $2/7*b*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/d+10/21*b^4*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*b^3*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\int (b \cos(c + dx))^{7/2} dx = \frac{10b^4 \sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} + \frac{10b^3 \sin(c + dx) \sqrt{b \cos(c + dx)}}{21d} + \frac{2b \sin(c + dx) (b \cos(c + dx))^{5/2}}{7d}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(7/2)}, x]$

[Out] $(10*b^4*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (10*b^3*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(21*d) + (2*b*(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sin}[c + d*x])/(7*d)$

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} + \frac{1}{7}(5b^2) \int (b \cos(c + dx))^{3/2} dx \\
&= \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} \\
&\quad + \frac{1}{21}(5b^4) \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d} \\
&\quad + \frac{(5b^4 \sqrt{\cos(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21 \sqrt{b \cos(c + dx)}} \\
&= \frac{10b^4 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d \sqrt{b \cos(c + dx)}} \\
&\quad + \frac{10b^3 \sqrt{b \cos(c + dx)} \sin(c + dx)}{21d} + \frac{2b(b \cos(c + dx))^{5/2} \sin(c + dx)}{7d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.78

$$\int (b \cos(c + dx))^{7/2} dx = \frac{b^3 \sqrt{b \cos(c + dx)} \left(20 \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sqrt{\cos(c + dx)} (23 \sin(c + dx) + 3 \sin(3(c + dx))) \right)}{42d \sqrt{\cos(c + dx)}}$$

[In] Integrate[(b*Cos[c + d*x])^(7/2),x]

[Out] (b^3*Sqrt[b*Cos[c + d*x]]*(20*EllipticF[(c + d*x)/2, 2] + Sqrt[Cos[c + d*x]]*(23*Sin[c + d*x] + 3*Sin[3*(c + d*x)])))/(42*d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.62 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.14

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)b^4\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 72\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$

[In] int((cos(d*x+c)*b)^(7/2),x,method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*b^4*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.97

$$\int (b \cos(c + dx))^{7/2} dx = \frac{-5i \sqrt{2} b^{7/2} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} b^{7/2} \operatorname{weierstrassPInverse}(\dots)}{2}$$

[In] integrate((b*cos(d*x+c))^(7/2),x, algorithm="fricas")


```
[Out] 1/21*(-5*I*sqrt(2)*b^(7/2)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*b^(7/2)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*(3*b^3*cos(d*x + c)^2 + 5*b^3)*sqrt(b*cos(d*x + c))*sin(d*x + c))/d
```

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{7/2} dx = \text{Timed out}$$

```
[In] integrate((b*cos(d*x+c))**(7/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int (b \cos(c + dx))^{7/2} dx = \int (b \cos(dx + c))^{\frac{7}{2}} dx$$

```
[In] integrate((b*cos(d*x+c))^(7/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^(7/2), x)
```

Giac [F]

$$\int (b \cos(c + dx))^{7/2} dx = \int (b \cos(dx + c))^{\frac{7}{2}} dx$$

```
[In] integrate((b*cos(d*x+c))^(7/2),x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^(7/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{7/2} dx = \int (b \cos(c + dx))^{7/2} dx$$

```
[In] int((b*cos(c + d*x))^(7/2),x)
```

```
[Out] int((b*cos(c + d*x))^(7/2), x)
```

3.103 $\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	514
Rubi [A] (verified)	514
Mathematica [A] (verified)	516
Maple [A] (verified)	516
Fricas [C] (verification not implemented)	517
Sympy [F(-1)]	517
Maxima [F]	517
Giac [F]	518
Mupad [F(-1)]	518

Optimal result

Integrand size = 21, antiderivative size = 125

$$\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77d\sqrt{b \cos(c+dx)}} + \frac{30\sqrt{b \cos(c+dx)} \sin(c+dx)}{77bd} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d}$$

[Out] $18/77*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^3/d+2/11*(b*\cos(d*x+c))^{(9/2)}*\sin(d*x+c)/b^5/d+30/77*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+30/77*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b/d$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^6(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^5d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^3d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77bd} + \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77d\sqrt{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^6/Sqrt[b*Cos[c + d*x]],x]
 [Out] (30*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*d*Sqrt[b*Cos[c + d*x]]) + (30*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^3*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^5*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Ssin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Ssin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Ssin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^6} \\
 &= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^5d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b^4} \\
 &= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^3d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^5d} + \frac{45 \int (b \cos(c + dx))^{3/2} dx}{77b^2} \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77bd} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^3d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^5d} + \frac{15}{77} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77bd} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^3d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^5d} + \frac{\left(15\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{77\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

$$= \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77d\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77bd} \\ + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^3d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^5d}$$

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.58

$$\int \frac{\cos^6(c+dx)}{\sqrt{b\cos(c+dx)}} dx \\ = \frac{480\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 347 \sin(2(c+dx)) + 64 \sin(4(c+dx)) + 7 \sin(6(c+dx))}{1232d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^6/Sqrt[b*Cos[c + d*x]], x]

[Out] (480*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*d*sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 4.56 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.86

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1084\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-370\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{77\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}$

[In] int(cos(d*x+c)^6/(cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)

[Out]
$$-2/77*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(448*\cos(1/2*d*x+1/2*c)^13-1568*\cos(1/2*d*x+1/2*c)^11+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*\cos(1/2*d*x+1/2*c)^2+1)^(1/2)*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^(1/2))+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.81

$$\int \frac{\cos^6(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{2(7 \cos(dx + c)^4 + 9 \cos(dx + c)^2 + 15) \sqrt{b \cos(dx + c)} \sin(dx + c) - 15i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c)) + 15i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c)) - I \sin(dx + c)}{77 b}$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/77*(2*(7*cos(d*x + c)^4 + 9*cos(d*x + c)^2 + 15)*sqrt(b*cos(d*x + c))*sin(d*x + c) - 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c)) + I*sin(d*x + c) + 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c)) - I*sin(d*x + c))/b*d

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^6(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^6}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^6/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\cos^6(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^6}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^6}{\sqrt{b \cos(c + dx)}} dx$$

[In] int(cos(c + d*x)^6/(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^6/(b*cos(c + d*x))^(1/2), x)

3.104 $\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	519
Rubi [A] (verified)	519
Mathematica [A] (verified)	521
Maple [A] (verified)	521
Fricas [C] (verification not implemented)	521
Sympy [F(-1)]	522
Maxima [F]	522
Giac [F]	522
Mupad [F(-1)]	523

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{14\sqrt{b \cos(c+dx)}E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15bd\sqrt{\cos(c+dx)}} + \frac{14(b \cos(c+dx))^{3/2} \sin(c+dx)}{45b^2d} + \frac{2(b \cos(c+dx))^{7/2} \sin(c+dx)}{9b^4d}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^4/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^4d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^2d} + \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^5/Sqrt[b*Cos[c + d*x]],x]

[Out] (14*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b*d*Sqrt[Cos[c + d*x]]) + (14*(b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^2*d) + (2*(b*Cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^4*d)

Rule 16

```
Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2715

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c+d*x]*((b*Sin[c+d*x])^(n-1)/(d*n)), x] + Dist[b^2*((n-1)/n), Int[(b*Sin[c+d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c-Pi/2+d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c+d*x])^n/Sin[c+d*x]^n, Int[Sin[c+d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^5} \\
&= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^4d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b^3} \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^2d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^4d} + \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b} \\
&= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^2d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^4d} \\
&\quad + \frac{\left(7\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15b\sqrt{\cos(c + dx)}} \\
&= \frac{14\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15bd\sqrt{\cos(c + dx)}} \\
&\quad + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^2d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^4d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.71

$$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \frac{168\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + \cos(c+dx)(38\sin(2(c+dx)) + 5\sin(4(c+dx)))}{180d\sqrt{b \cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^5/Sqrt[b*Cos[c + d*x]],x]

[Out] (168*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.90 (sec) , antiderivative size = 220, normalized size of antiderivative = 2.20

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}\right)}$

[In] int(cos(d*x+c)^5/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-2/45*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(160*\cos(1/2*d*x+1/2*c)^11-480*\cos(1/2*d*x+1/2*c)^9+616*\cos(1/2*d*x+1/2*c)^7-432*\cos(1/2*d*x+1/2*c)^5+160*\cos(1/2*d*x+1/2*c)^3-21*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*\cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(\cos(1/2*d*x+1/2*c),2^(1/2))-24*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2)^(1/2))/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.12 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04

$$\int \frac{\cos^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \frac{2(5 \cos(dx+c)^3 + 7 \cos(dx+c))\sqrt{b \cos(dx+c)} \sin(dx+c) + 21i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weiers})}{1}$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

```
[Out] 1/45*(2*(5*cos(d*x + c)^3 + 7*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x +
c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0,
cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4,
0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b*d)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \frac{\cos^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^5}{\sqrt{b \cos(dx + c)}} dx$$

```
[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(d*x + c)^5/sqrt(b*cos(d*x + c)), x)
```

Giac [F]

$$\int \frac{\cos^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^5}{\sqrt{b \cos(dx + c)}} dx$$

```
[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(cos(d*x + c)^5/sqrt(b*cos(d*x + c)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^5}{\sqrt{b \cos(c + dx)}} dx$$

```
[In] int(cos(c + d*x)^5/(b*cos(c + d*x))^(1/2), x)
```

```
[Out] int(cos(c + d*x)^5/(b*cos(c + d*x))^(1/2), x)
```

3.105 $\int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	524
Rubi [A] (verified)	524
Mathematica [A] (verified)	526
Maple [A] (verified)	526
Fricas [C] (verification not implemented)	526
Sympy [F(-1)]	527
Maxima [F]	527
Giac [F]	527
Mupad [F(-1)]	528

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b \cos(c+dx)}} + \frac{10\sqrt{b \cos(c+dx)} \sin(c+dx)}{21bd} + \frac{2(b \cos(c+dx))^{5/2} \sin(c+dx)}{7b^3d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^3/d+10/21*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b/d$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^3d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21bd} + \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^4/\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]], x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (10*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(21*b*d) + (2*(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sin}[c + d*x])/(7*b^3*d)$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c+d*x]*(b*\sin[c+d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\sin[c+d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\sin[c+d*x])^n/\sin[c+d*x]^n, \text{Int}[\sin[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^4} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^3d} + \frac{5 \int (b \cos(c + dx))^{3/2} dx}{7b^2} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21bd} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^3d} + \frac{5}{21} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21bd} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^3d} \\
 &\quad + \frac{\left(5\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21\sqrt{b \cos(c + dx)}} \\
 &= \frac{10\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21d\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21bd} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.65

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{40\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 26 \sin(2(c + dx)) + 3 \sin(4(c + dx))}{84d\sqrt{b \cos(c + dx)}}$$

[In] Integrate[Cos[c + d*x]^4/Sqrt[b*Cos[c + d*x]],x]

[Out] (40*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*d*sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.20 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.13

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\left(48\left(\cos^9\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 120\left(\cos^7\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 128\left(\cos^5\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 72\left(\cos^3\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 5\right)}{21\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)} \sin\left(\frac{dx}{2} + \frac{c}{2}\right) \sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1}}$

[In] int(cos(d*x+c)^4/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.94

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{2\sqrt{b \cos(dx + c)}(3 \cos(dx + c)^2 + 5) \sin(dx + c) - 5i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i)}{21bd}$$

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

```
[Out] 1/21*(2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 5)*sin(d*x + c) - 5*I*sqrt
(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I
*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))
/(b*d)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)}{\sqrt{b} \cos(c + dx)} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \frac{\cos^4(c + dx)}{\sqrt{b} \cos(c + dx)} dx = \int \frac{\cos(dx + c)^4}{\sqrt{b} \cos(dx + c)} dx$$

```
[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(d*x + c)^4/sqrt(b*cos(d*x + c)), x)
```

Giac [F]

$$\int \frac{\cos^4(c + dx)}{\sqrt{b} \cos(c + dx)} dx = \int \frac{\cos(dx + c)^4}{\sqrt{b} \cos(dx + c)} dx$$

```
[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(cos(d*x + c)^4/sqrt(b*cos(d*x + c)), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^4}{\sqrt{b \cos(c + dx)}} dx$$

```
[In] int(cos(c + d*x)^4/(b*cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^4/(b*cos(c + d*x))^(1/2), x)
```


3.106 $\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	529
Rubi [A] (verified)	529
Mathematica [A] (verified)	530
Maple [B] (verified)	531
Fricas [C] (verification not implemented)	531
Sympy [F(-1)]	532
Maxima [F]	532
Giac [F]	532
Mupad [F(-1)]	532

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5bd\sqrt{\cos(c+dx)}} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^2d}$$

[Out] $2/5*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/b^2/d+6/5*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^2d} + \frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

[In] `Int[Cos[c + d*x]^3/Sqrt[b*Cos[c + d*x]],x]`

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(5*b^2*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b^3} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^2d} + \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^2d} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b\sqrt{\cos(c + dx)}} \\
 &= \frac{6\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5bd\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{6\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos(c + dx) \sin(2(c + dx))}{5d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^3/Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(88) = 176.

Time = 2.57 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.92

method	result
default	$-\frac{2\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{5\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] `int(cos(d*x+c)^3/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.26

$$\int \frac{\cos^3(c+dx)}{\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{2\sqrt{b\cos(dx+c)}\cos(dx+c)\sin(dx+c) + 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c) - i\sin(dx+c)))}{(b*d)}$$

[In] `integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{b*\cos(d*x+c)}*\cos(d*x+c)*\sin(d*x+c) + 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c) + I*\sin(d*x+c))) - 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c) - I*\sin(d*x+c))))/(b*d)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^3}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^3}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^3}{\sqrt{b \cos(c + dx)}} dx$$

[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(1/2), x)

$$3.107 \quad \int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	533
Rubi [A] (verified)	533
Mathematica [A] (verified)	534
Maple [B] (verified)	535
Fricas [C] (verification not implemented)	535
Sympy [F]	535
Maxima [F]	536
Giac [F]	536
Mupad [B] (verification not implemented)	536

Optimal result

Integrand size = 21, antiderivative size = 69

$$\begin{aligned} & \int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx \\ &= \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d\sqrt{b \cos(c+dx)}} + \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3bd} \end{aligned}$$

[Out] $2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}+2/3*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b/d$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\begin{aligned} & \int \frac{\cos^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx \\ &= \frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3bd} + \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d\sqrt{b \cos(c+dx)}} \end{aligned}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^2/\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]], x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(3*b*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{3/2} dx}{b^2} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3bd} + \frac{1}{3} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3bd} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
 &= \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3bd}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx))}{3d\sqrt{b \cos(c + dx)}}$$

`[In] Integrate[Cos[c + d*x]^2/Sqrt[b*Cos[c + d*x]],x]`

`[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*d*Sqrt[b*Cos[c + d*x]])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(85) = 170.

Time = 1.79 (sec) , antiderivative size = 187, normalized size of antiderivative = 2.71

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$

[In] `int(cos(d*x+c)^2/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/3*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(4*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(\cos(1/2*d*x+1/2*c),2^(1/2)))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.14

$$\int \frac{\cos^2(c+dx)}{\sqrt{b}\cos(c+dx)} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))}{3bd}$$

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out]
$$1/3*(-I*\sqrt{2}*\sqrt{b}*\text{weierstrassPInverse}(-4,0,\cos(d*x+c))+I*\sin(d*x+c))+I*\sqrt{2}*\sqrt{b}*\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*\sqrt{b*\cos(d*x+c)}*\sin(d*x+c))/(b*d)$$

Sympy [F]

$$\int \frac{\cos^2(c+dx)}{\sqrt{b}\cos(c+dx)} dx = \int \frac{\cos^2(c+dx)}{\sqrt{b}\cos(c+dx)} dx$$

[In] `integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(1/2),x)`

[Out] `Integral(cos(c+d*x)**2/sqrt(b*cos(c+d*x)),x)`

Maxima [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 13.17 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.84

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2 \sqrt{\cos(c + dx)} F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{3d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{3bd}$$

[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/2),x)

[Out] (2*cos(c + d*x)^(1/2)*ellipticF(c/2 + (d*x)/2, 2))/(3*d*(b*cos(c + d*x))^(1/2)) + (2*sin(c + d*x)*(b*cos(c + d*x))^(1/2))/(3*b*d)

3.108 $\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	537
Rubi [A] (verified)	537
Mathematica [A] (verified)	538
Maple [B] (verified)	538
Fricas [C] (verification not implemented)	539
Sympy [F]	539
Maxima [F]	539
Giac [F]	540
Mupad [B] (verification not implemented)	540

Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{bd\sqrt{\cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c))^{1/2}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{1/2})*(b*\cos(d*x+c))^{1/2}/b/d/\cos(d*x+c)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2721, 2719}

$$\int \frac{\cos(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

[In] `Int[Cos[c + d*x]/Sqrt[b*Cos[c + d*x]],x]`

[Out] `(2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]])`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \sqrt{b \cos(c + dx)} dx}{b} \\ &= \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{b \sqrt{\cos(c + dx)}} \\ &= \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{bd \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{bd \sqrt{\cos(c + dx)}}$$

[In] `Integrate[Cos[c + d*x]/Sqrt[b*Cos[c + d*x]],x]`

[Out] `(2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(63) = 126.

Time = 1.72 (sec) , antiderivative size = 141, normalized size of antiderivative = 3.44

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1}E\left(\cos\left(\frac{dx}{2} + \frac{c}{2}\right), \sqrt{2}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)bd}$
risch	$-\frac{i\left(e^{2i(dx+c)} + 1\right)\sqrt{2}e^{-i(dx+c)}}{d\sqrt{\left(e^{2i(dx+c)} + 1\right)be^{-i(dx+c)}}} - \frac{i\left(\frac{2\left(b e^{2i(dx+c)} + b\right)}{b\sqrt{e^{i(dx+c)}\left(b e^{2i(dx+c)} + b\right)}} + \frac{i\sqrt{-i\left(e^{i(dx+c)} + i\right)}\sqrt{2}\sqrt{i\left(e^{i(dx+c)} - i\right)}\sqrt{ie^{i(dx+c)}}\left(-2iE\left(\sqrt{-i\left(e^{i(dx+c)} + i\right)}\right)\right)}{\sqrt{b e^{3i(dx+c)} + b e^{i(dx+c)}}}\right)}{d\sqrt{\left(e^{2i(dx+c)} + 1\right)be^{-i(dx+c)}}}$

[In] `int(cos(d*x+c)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c),2^(1/2))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.61

$$\int \frac{\cos(c+dx)}{\sqrt{b\cos(c+dx)}} dx = \frac{i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c))) - i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c)))}{bd}$$

```
[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b*d)
```

Sympy [F]

$$\int \frac{\cos(c+dx)}{\sqrt{b\cos(c+dx)}} dx = \int \frac{\cos(c+dx)}{\sqrt{b\cos(c+dx)}} dx$$

```
[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Integral(cos(c + d*x)/sqrt(b*cos(c + d*x)), x)
```

Maxima [F]

$$\int \frac{\cos(c+dx)}{\sqrt{b\cos(c+dx)}} dx = \int \frac{\cos(dx+c)}{\sqrt{b\cos(dx+c)}} dx$$

```
[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(d*x + c)/sqrt(b*cos(d*x + c)), x)
```

Giac [F]

$$\int \frac{\cos(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.80

$$\int \frac{\cos(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2 \sqrt{\cos(c + dx)} E\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[In] int(cos(c + d*x)/(b*cos(c + d*x))^(1/2),x)

[Out] (2*cos(c + d*x)^(1/2)*ellipticE(c/2 + (d*x)/2, 2))/(d*(b*cos(c + d*x))^(1/2))

3.109 $\int \frac{1}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	541
Rubi [A] (verified)	541
Mathematica [A] (verified)	542
Maple [C] (verified)	542
Fricas [C] (verification not implemented)	543
Sympy [F]	543
Maxima [F]	543
Giac [F]	544
Mupad [B] (verification not implemented)	544

Optimal result

Integrand size = 12, antiderivative size = 38

$$\int \frac{1}{\sqrt{b \cos(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d\sqrt{b \cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2721, 2720}

$$\int \frac{1}{\sqrt{b \cos(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d\sqrt{b \cos(c+dx)}}$$

[In] `Int[1/Sqrt[b*Cos[c + d*x]],x]`

[Out] `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ`

`[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{b \cos(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d\sqrt{b \cos(c+dx)}}$$

[In] `Integrate[1/Sqrt[b*Cos[c + d*x]],x]`

[Out] `(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(d*Sqrt[b*Cos[c + d*x]])`

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.48 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.42

method	result	size
default	$\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1} \operatorname{am}^{-1}\left(\frac{dx}{2} + \frac{c}{2} \sqrt{2}\right)}{d\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)b}}$	54

[In] `int(1/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `2/d/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)*(2*cos(1/2*d*x+1/2*c)^2-1)^(1/2)*InverseJacobiAM(1/2*d*x+1/2*c,2^(1/2))`

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.58

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{-i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{bd}$$

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b*d)

Sympy [F]

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(1/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(1/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{b \cos(c + dx)}} dx = \frac{2 \sqrt{\cos(c + dx)} F\left(\frac{c}{2} + \frac{dx}{2} \mid 2\right)}{d \sqrt{b \cos(c + dx)}}$$

[In] int(1/(b*cos(c + d*x))^(1/2),x)

[Out] (2*cos(c + d*x)^(1/2)*ellipticF(c/2 + (d*x)/2, 2))/(d*(b*cos(c + d*x))^(1/2))

$$3.110 \quad \int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	545
Rubi [A] (verified)	545
Mathematica [A] (verified)	546
Maple [B] (verified)	547
Fricas [C] (verification not implemented)	547
Sympy [F]	548
Maxima [F]	548
Giac [F]	548
Mupad [F(-1)]	548

Optimal result

Integrand size = 19, antiderivative size = 65

$$\int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx = -\frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd\sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{d\sqrt{b \cos(c+dx)}}$$

[Out] 2*sin(d*x+c)/d/(b*cos(d*x+c))^(1/2)-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 \sin(c+dx)}{d\sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{bd\sqrt{\cos(c+dx)}}$$

[In] Int[Sec[c + d*x]/Sqrt[b*Cos[c + d*x]],x]

[Out] (-2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]) + (2*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
&= \frac{2 \sin(c + dx)}{d \sqrt{b \cos(c + dx)}} - \frac{\int \sqrt{b \cos(c + dx)} dx}{b} \\
&= \frac{2 \sin(c + dx)}{d \sqrt{b \cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{b \sqrt{\cos(c + dx)}} \\
&= -\frac{2 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{bd \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)}{d \sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.72

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2 \left(-\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx) \right)}{d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Sec[c + d*x]/Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(d*Sqr
t[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(85) = 170.

Time = 1.90 (sec) , antiderivative size = 195, normalized size of antiderivative = 3.00

method	result
default	$\frac{2\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)}{\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)}}$

[In] `int(sec(d*x+c)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2*(-2*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c))^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)*(-2*\sin(1/2*d*x+1/2*c))^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})}/(-b*(2*\sin(1/2*d*x+1/2*c))^4-\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}}{d}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.60

$$\int \frac{\sec(c+dx)}{\sqrt{b\cos(c+dx)}} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))) + \dots}{\dots}$$

[In] `integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2),x,algorithm="fricas")`

[Out]
$$(-I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))) + I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))) + 2*\sqrt{b*\cos(d*x+c)}*\sin(d*x+c))/(b*d*\cos(d*x+c))$$

Sympy [F]

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx) \sqrt{b \cos(c + dx)}} dx$$

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/2)), x)

3.111 $\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	549
Rubi [A] (verified)	549
Mathematica [A] (verified)	550
Maple [B] (verified)	551
Fricas [C] (verification not implemented)	551
Sympy [F]	552
Maxima [F]	552
Giac [F]	552
Mupad [F(-1)]	552

Optimal result

Integrand size = 21, antiderivative size = 67

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d\sqrt{b \cos(c+dx)}} + \frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}}$$

[Out] $2/3*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+2/3*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \frac{\sec^2(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2b \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3d\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c + d*x]^2/\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]], x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(3*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (2*b*\operatorname{Sin}[c + d*x])/(3*d*(b*\operatorname{Cos}[c + d*x])^{(3/2)})$

Rule 16

$\operatorname{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n, x\} \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2b \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{1}{3} \int \frac{1}{\sqrt{b \cos(c + dx)}} dx \\
&= \frac{2b \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3\sqrt{b \cos(c + dx)}} \\
&= \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3d\sqrt{b \cos(c + dx)}} + \frac{2b \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.72

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{2\left(\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx)\right)}{3d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Sec[c + d*x]^2/Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d*Sqrt
[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(83) = 166.

Time = 1.79 (sec) , antiderivative size = 238, normalized size of antiderivative = 3.55

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-2/3*(-2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)}))*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.54

$$\int \frac{\sec^2(c+dx)}{\sqrt{b}\cos(c+dx)} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c)) + 2\sqrt{b}\cos(dx+c)\sin(dx+c)}{3bd\cos(dx+c)^2}$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out]
$$1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)^2*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x+c) + I*\sin(d*x+c)) + I*\sqrt{2}*\sqrt{b}*\cos(d*x+c)^2*\operatorname{weierstrassPInverse}(-4, 0, \cos(d*x+c) - I*\sin(d*x+c)) + 2*\sqrt{b}*\cos(d*x+c)*\sin(d*x+c))/(\sqrt{b*d}\cos(d*x+c)^2)$$

Sympy [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)**2/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^2 \sqrt{b \cos(c + dx)}} dx$$

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/2)), x)

3.112 $\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	553
Rubi [A] (verified)	553
Mathematica [A] (verified)	555
Maple [B] (verified)	555
Fricas [C] (verification not implemented)	556
Sympy [F]	556
Maxima [F]	556
Giac [F]	557
Mupad [F(-1)]	557

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx = -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5bd\sqrt{\cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}}$$

[Out] $2/5*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2b^2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5d\sqrt{b \cos(c+dx)}} - \frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5bd\sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]^3/\text{Sqrt}[b*\text{Cos}[c + d*x]],x]$

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2716

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{1}{5}(3b) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b} \\
 &= \frac{2b^2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}} - \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b\sqrt{\cos(c + dx)}} \\
 &= -\frac{6\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5bd\sqrt{\cos(c + dx)}} + \frac{2b^2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5d\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.67

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{-6\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \middle| 2\right) + 6\sin(c + dx) + 2\sec(c + dx)\tan(c + dx)}{5d\sqrt{b \cos(c + dx)}}$$

[In] Integrate[Sec[c + d*x]^3/Sqrt[b*Cos[c + d*x]],x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(109) = 218.

Time = 2.42 (sec) , antiderivative size = 367, normalized size of antiderivative = 3.78

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{(24\cos(\frac{dx}{2} + \frac{c}{2})(\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}} E(\cos(\frac{dx+c}{2}))$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-2/5*(-(-2*\cos(1/2*d*x+1/2*c)^2+1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)/b/\sin(1/2*d*x+1/2*c)^3/(8*\sin(1/2*d*x+1/2*c)^6-12*\sin(1/2*d*x+1/2*c)^4+6*\sin(1/2*d*x+1/2*c)^2-1)*(24*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6-12*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^(1/2))*\sin(1/2*d*x+1/2*c)^4-24*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)+12*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^(1/2))*\sin(1/2*d*x+1/2*c)^2+8*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c), 2^(1/2)))*(-2*\sin(1/2*d*x+1/2*c)^4*b+b*\sin(1/2*d*x+1/2*c)^2)^(1/2)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.25

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^3 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c))) + \dots}{\dots}$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 1)*sin(d*x + c))/(b*d*cos(d*x + c)^3)

Sympy [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)**3/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^3 \sqrt{b \cos(c + dx)}} dx$$

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/2)), x)

3.113 $\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	558
Rubi [A] (verified)	558
Mathematica [A] (verified)	560
Maple [B] (verified)	560
Fricas [C] (verification not implemented)	561
Sympy [F]	561
Maxima [F]	561
Giac [F]	562
Mupad [F(-1)]	562

Optimal result

Integrand size = 21, antiderivative size = 95

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b \cos(c+dx)}} + \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}$$

[Out] $2/7*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c+d*x]^4/\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]], x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(21*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*b^3*\operatorname{Sin}[c+d*x])/(7*d*(b*\operatorname{Cos}[c+d*x])^{(7/2)}) + (10*b*\operatorname{Sin}[c+d*x])/(21*d*(b*\operatorname{Cos}[c+d*x])^{(3/2)})$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c+d*x]*((b*\sin[c+d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\sin[c+d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\sin[c+d*x])^n/\sin[c+d*x]^n, \text{Int}[\sin[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^4 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
 &= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{1}{7}(5b^2) \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
 &= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{5}{21} \int \frac{1}{\sqrt{b \cos(c+dx)}} dx \\
 &= \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21\sqrt{b \cos(c+dx)}} \\
 &= \frac{10\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21d\sqrt{b \cos(c+dx)}} + \frac{2b^3 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10b \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.66

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{10\sqrt{\cos(c + dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + 2(5 + 3 \sec^2(c + dx)) \tan(c + dx)}{21d\sqrt{b \cos(c + dx)}}$$

[In] Integrate[Sec[c + d*x]^4/Sqrt[b*Cos[c + d*x]],x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(107) = 214.

Time = 2.52 (sec) , antiderivative size = 267, normalized size of antiderivative = 2.81

method	result
default	$-\frac{\sqrt{-\left(-2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + 1\right)b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}{\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)bd}} \left(-\frac{\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{28b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^4} - \frac{5\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}}{21b\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right) - \frac{1}{2}\right)^4} \right)$

[In] int(sec(d*x+c)^4/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1/28*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^4-5/21*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^2+10/21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.21

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{-5i \sqrt{2} \sqrt{b} \cos(dx + c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + 5i \sqrt{2} \sqrt{b} \cos(dx + c)^4}{21 b d c}$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*(5*cos(d*x + c)^2 + 3)*sin(d*x + c))/(b*d*cos(d*x + c)^4)

Sympy [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**4/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)**4/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^4}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^4}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^4(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^4 \sqrt{b \cos(c + dx)}} dx$$

[In] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(1/2)), x)

3.114 $\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	563
Rubi [A] (verified)	563
Mathematica [A] (verified)	565
Maple [B] (verified)	565
Fricas [C] (verification not implemented)	566
Sympy [F]	566
Maxima [F]	566
Giac [F]	567
Mupad [F(-1)]	567

Optimal result

Integrand size = 21, antiderivative size = 125

$$\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx = -\frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15bd\sqrt{\cos(c+dx)}} + \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}}$$

$$+ \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}}$$

[Out] $2/9*b^4*\sin(d*x+c)/d/(b*\cos(d*x+c))^(9/2)+14/45*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^(5/2)+14/15*\sin(d*x+c)/d/(b*\cos(d*x+c))^(1/2)-14/15*(\cos(1/2*d*x+1/2*c))^2)^(1/2)/\cos(1/2*d*x+1/2*c)*EllipticE(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/b/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec^5(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2b^4 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b^2 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}}$$

$$+ \frac{14 \sin(c+dx)}{15d\sqrt{b \cos(c+dx)}} - \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15bd\sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]^5/\text{Sqrt}[b*\text{Cos}[c + d*x]],x]$

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^(9/2)) + (14*b^2*\text{Sin}[c +$

$d*x]/(45*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (14*\text{Sin}[c + d*x])/(15*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x] \&\& \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\text{sin}[(c_.) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n+1)}/(b*d*(n+1))), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\text{sin}[(c_.) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^{n+1}/\text{Sin}[c + d*x]^{n+1}, \text{Int}[\text{Sin}[c + d*x]^{n+1}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^5 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9}(7b^3) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{1}{15}(7b) \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b} \\
 &= \frac{2b^4 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} \\
 &\quad + \frac{14 \sin(c + dx)}{15d\sqrt{b \cos(c + dx)}} - \frac{\left(7\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15b\sqrt{\cos(c + dx)}}
 \end{aligned}$$

$$= -\frac{14\sqrt{b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15bd\sqrt{\cos(c+dx)}} + \frac{2b^4\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}}$$

$$+ \frac{14b^2\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}} + \frac{14\sin(c+dx)}{15d\sqrt{b\cos(c+dx)}}$$

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.62

$$\int \frac{\sec^5(c+dx)}{\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{-42\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + 42\sin(c+dx) + 2\sec(c+dx)(7+5\sec^2(c+dx))\tan(c+dx)}{45d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^5/Sqrt[b*Cos[c + d*x]], x]

[Out] (-42*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 412 vs. 2(133) = 266.

Time = 3.18 (sec) , antiderivative size = 413, normalized size of antiderivative = 3.30

method	result
default	$-\frac{\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{72b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5} - \frac{7\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{90b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5}\right)}$

[In] int(sec(d*x+c)^5/(cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)

[Out] -(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)*(-1/72*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/90*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^3-28/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)+14/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-14/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.05

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{-21i \sqrt{2} \sqrt{b} \cos(dx + c)^5 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)))}{1}$$

[In] integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*cos(d*x + c)^4 + 7*cos(d*x + c)^2 + 5)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^5)

Sympy [F]

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**5/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(sec(c + d*x)**5/sqrt(b*cos(c + d*x)), x)

Maxima [F]

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^5}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^5/sqrt(b*cos(d*x + c)), x)

Giac [F]

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^5}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(sec(d*x+c)^5/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^5/sqrt(b*cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^5 \sqrt{b \cos(c + dx)}} dx$$

[In] int(1/(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^5*(b*cos(c + d*x))^(1/2)), x)

3.115 $\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx$

Optimal result	568
Rubi [A] (verified)	568
Mathematica [A] (verified)	570
Maple [A] (verified)	570
Fricas [C] (verification not implemented)	571
Sympy [F(-1)]	571
Maxima [F]	571
Giac [F]	571
Mupad [F(-1)]	572

Optimal result

Integrand size = 21, antiderivative size = 128

$$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77bd\sqrt{b \cos(c+dx)}} + \frac{30\sqrt{b \cos(c+dx)} \sin(c+dx)}{77b^2d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^4/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^6/d+30/77*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^2/d

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^6d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^4d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^2d} + \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77bd\sqrt{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^7/(b*Cos[c + d*x])^(3/2), x]

[Out] (30*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*b*d*sqrt[b*Cos[c + d*x]]) + (30*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b^2*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^4*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^6*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^7} \\
 &= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^6 d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b^5} \\
 &= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^4 d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^6 d} + \frac{45 \int (b \cos(c + dx))^{3/2} dx}{77b^3} \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77b^2 d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^4 d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^6 d} + \frac{15 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{77b} \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77b^2 d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^4 d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^6 d} + \frac{\left(15\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{77b\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

$$= \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77bd\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^2d} \\ + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^4d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^6d}$$

Mathematica [A] (verified)

Time = 0.42 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.59

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{480\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 347 \sin(2(c+dx)) + 64 \sin(4(c+dx))}{1232bd\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^7/(b*Cos[c + d*x])^(3/2), x]

[Out] (480*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*b*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 5.53 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.84

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1084\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-370\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+15\left(\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\right)^2\right)^{1/2}\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+1\right)^{1/2}\operatorname{EllipticF}\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right), 2\right)^{1/2}+62\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{77b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}$

[In] int(cos(d*x+c)^7/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/77*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(448*cos(1/2*d*x+1/2*c)^13-1568*cos(1/2*d*x+1/2*c)^11+2384*cos(1/2*d*x+1/2*c)^9-2040*cos(1/2*d*x+1/2*c)^7+1084*cos(1/2*d*x+1/2*c)^5-370*cos(1/2*d*x+1/2*c)^3+15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2)^(1/2)+62*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.79

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{2(7\cos(dx+c)^4 + 9\cos(dx+c)^2 + 15)\sqrt{b\cos(dx+c)}\sin(dx+c) - 15i\sqrt{2}\sqrt{b\cos(dx+c)}}{(b\cos(c+dx))^{3/2}}$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/77*(2*(7*cos(d*x + c)^4 + 9*cos(d*x + c)^2 + 15)*sqrt(b*cos(d*x + c))*sin(d*x + c) - 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**7/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(dx+c)^7}{(b\cos(dx+c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \int \frac{\cos(dx+c)^7}{(b\cos(dx+c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^7(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^7}{(b \cos(c + dx))^{3/2}} dx$$

```
[In] int(cos(c + d*x)^7/(b*cos(c + d*x))^(3/2), x)
```

```
[Out] int(cos(c + d*x)^7/(b*cos(c + d*x))^(3/2), x)
```

$$3.116 \quad \int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	573
Rubi [A] (verified)	573
Mathematica [A] (verified)	575
Maple [A] (verified)	575
Fricas [C] (verification not implemented)	575
Sympy [F(-1)]	576
Maxima [F]	576
Giac [F]	576
Mupad [F(-1)]	576

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{14(b \cos(c+dx))^{3/2} \sin(c+dx)}{45b^3 d} + \frac{2(b \cos(c+dx))^{7/2} \sin(c+dx)}{9b^5 d}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^3/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^5/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^2/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^5 d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^3 d} + \frac{14 E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15b^2 d \sqrt{\cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^6/(b*Cos[c + d*x])^(3/2),x]

[Out] (14*sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b^2*d*sqrt[Cos[c + d*x]]) + (14*(b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^3*d) + (2*(b*cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^5*d)

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)*((b_)*(v_))^{(n_)}}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c+d*x]*(b*\text{Sin}[c+d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c+d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c+d*x])^{n-1}/\text{Sin}[c+d*x]^n, \text{Int}[\text{Sin}[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^6} \\
 &= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^5 d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b^4} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^3 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^5 d} + \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b^2} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^3 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^5 d} \\
 &\quad + \frac{\left(7\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15b^2 \sqrt{\cos(c + dx)}} \\
 &= \frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15b^2 d \sqrt{\cos(c + dx)}} \\
 &\quad + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^3 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^5 d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.74

$$\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{168\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + \cos(c+dx)(38\sin(2(c+dx)) + 5\sin(4(c+dx)))}{180bd\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^6/(b*Cos[c + d*x])^(3/2), x]

[Out] (168*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*b*d*sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 4.59 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.23

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] int(cos(d*x+c)^6/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/45*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2)^(1/2))-24*cos(1/2*d*x+1/2*c)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04

$$\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{2(5\cos(dx+c)^3 + 7\cos(dx+c))\sqrt{b\cos(dx+c)}\sin(dx+c) + 21i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4, 0, \cos(dx+c) + i\sin(dx+c)) - 21i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{b^2d}$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] 1/45*(2*(5*cos(d*x + c)^3 + 7*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b^2*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^6}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^6}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^6}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(cos(c + d*x)^6/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^6/(b*cos(c + d*x))^(3/2), x)

$$3.117 \quad \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	577
Rubi [A] (verified)	577
Mathematica [A] (verified)	579
Maple [A] (verified)	579
Fricas [C] (verification not implemented)	579
Sympy [F(-1)]	580
Maxima [F]	580
Giac [F]	580
Mupad [F(-1)]	580

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21bd\sqrt{b \cos(c+dx)}} + \frac{10\sqrt{b \cos(c+dx)} \sin(c+dx)}{21b^2d} + \frac{2(b \cos(c+dx))^{5/2} \sin(c+dx)}{7b^4d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^4/d+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^4d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^2d} + \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^5/(b*\operatorname{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(21*b*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]) + (10*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]]*\operatorname{Sin}[c + d*x])/(21*b^2*d) + (2*(b*\operatorname{Cos}[c + d*x])^{(5/2)}*\operatorname{Sin}[c + d*x])/(7*b^4*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^5} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^4d} + \frac{5 \int (b \cos(c + dx))^{3/2} dx}{7b^3} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^2d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^4d} + \frac{5 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{21b} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^2d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^4d} \\
 &\quad + \frac{\left(5\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21b\sqrt{b \cos(c + dx)}} \\
 &= \frac{10\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21bd\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^2d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^4d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.66

$$\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{40\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 26\sin(2(c+dx)) + 3\sin(4(c+dx))}{84bd\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^5/(b*Cos[c + d*x])^(3/2), x]

[Out] (40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*b*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.42 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.10

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] int(cos(d*x+c)^5/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.91

$$\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{2\sqrt{b\cos(dx+c)}(3\cos(dx+c)^2+5)\sin(dx+c)-5i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}}{(b\cos(c+dx))^{3/2}}$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] 1/21*(2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 5)*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^5}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^5}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^5}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(cos(c + d*x)^5/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^5/(b*cos(c + d*x))^(3/2), x)

$$3.118 \quad \int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	581
Rubi [A] (verified)	581
Mathematica [A] (verified)	582
Maple [B] (verified)	583
Fricas [C] (verification not implemented)	583
Sympy [F(-1)]	583
Maxima [F]	584
Giac [F]	584
Mupad [F(-1)]	584

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5b^2 d \sqrt{\cos(c+dx)}} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^3 d}$$

[Out] $2/5*(b*\cos(d*x+c))^(3/2)*\sin(d*x+c)/b^3/d+6/5*(\cos(1/2*d*x+1/2*c))^2^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))*(b*\cos(d*x+c))^(1/2)/b^2/d/\cos(d*x+c)^(1/2)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^3 d} + \frac{6 E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5b^2 d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^4/(b*\text{Cos}[c + d*x])^(3/2), x]$

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^(3/2)*\text{Sin}[c + d*x])/(5*b^3*d)$

Rule 16

$\text{Int}[(u_*)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^(m+n), x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b^4} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^3d} + \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b^2} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^3d} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b^2 \sqrt{\cos(c + dx)}} \\
 &= \frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5b^2d \sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.85

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{6\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos(c + dx) \sin(2(c + dx))}{5bd \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^4/(b*Cos[c + d*x])^(3/2),x]
```

```
[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*b*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(88) = 176.

Time = 2.85 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.96

method	result
default	$-\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{5b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] `int(cos(d*x+c)^4/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)}))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.26

$$\int \frac{\cos^4(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{2\sqrt{b\cos(dx+c)}\cos(dx+c)\sin(dx+c) + 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)) + i\sin(dx+c)) - 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c) - i\sin(dx+c)))}{(b^2*d)}$$

[In] `integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{b*\cos(d*x+c)}*\cos(d*x+c)*\sin(d*x+c) + 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c) + I*\sin(d*x+c))) - 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c) - I*\sin(d*x+c))))/(b^2*d)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^4}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^4}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^4}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(cos(c + d*x)^4/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^4/(b*cos(c + d*x))^(3/2), x)

$$3.119 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	585
Rubi [A] (verified)	585
Mathematica [A] (verified)	586
Maple [B] (verified)	587
Fricas [C] (verification not implemented)	587
Sympy [F(-1)]	587
Maxima [F]	588
Giac [F]	588
Mupad [F(-1)]	588

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3bd\sqrt{b \cos(c+dx)}} + \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^2d}$$

[Out] $2/3*(\cos(1/2*d*x+1/2*c))^{1/2}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{1/2})*\cos(d*x+c)^{1/2}/b/d/(b*\cos(d*x+c))^{1/2}+2/3*\sin(d*x+c)*(b*\cos(d*x+c))^{1/2}/b^2/d$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^2d} + \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^3/(b*\operatorname{Cos}[c+d*x])^{3/2}, x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(3*b*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(3*b^2*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{3/2} dx}{b^3} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^2d} + \frac{\int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{3b} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^2d} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b\sqrt{b \cos(c + dx)}} \\
 &= \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3bd\sqrt{b \cos(c + dx)}} + \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx))}{3bd\sqrt{b \cos(c + dx)}}$$

`[In] Integrate[Cos[c + d*x]^3/(b*Cos[c + d*x])^(3/2), x]`

`[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*b*d*Sqrt[b*Cos[c + d*x]])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(88) = 176.

Time = 2.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.64

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2}\right)}{3b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$

[In] `int(cos(d*x+c)^3/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{2}{3} * \left((2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * b * \sin(1/2 * d * x + 1/2 * c) ^ 2 \right) ^ {1/2} / b * \left(4 * \sin(1/2 * d * x + 1/2 * c) ^ 4 * \cos(1/2 * d * x + 1/2 * c) - 2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 * \cos(1/2 * d * x + 1/2 * c) + \left(\sin(1/2 * d * x + 1/2 * c) ^ 2 \right) ^ {1/2} * \left(2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1 \right) ^ {1/2} * \text{EllipticF}(\cos(1/2 * d * x + 1/2 * c), 2 ^ {1/2}) \right) / \left(-b * \left(2 * \sin(1/2 * d * x + 1/2 * c) ^ 4 - \sin(1/2 * d * x + 1/2 * c) ^ 2 \right) ^ {1/2} / \sin(1/2 * d * x + 1/2 * c) / \left((2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * b \right) ^ {1/2} / d \right)$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.10

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{3b^2 d}$$

[In] `integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$\frac{1}{3} * \left(-I * \sqrt{2} * \sqrt{b} * \text{weierstrassPInverse}(-4, 0, \cos(d * x + c) + I * \sin(d * x + c)) + I * \sqrt{2} * \sqrt{b} * \text{weierstrassPInverse}(-4, 0, \cos(d * x + c) - I * \sin(d * x + c)) + 2 * \sqrt{b * \cos(d * x + c)} * \sin(d * x + c) \right) / (b^2 * d)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^3}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(3/2), x)

$$3.120 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	589
Rubi [A] (verified)	589
Mathematica [A] (verified)	590
Maple [B] (verified)	590
Fricas [C] (verification not implemented)	591
Sympy [F(-1)]	591
Maxima [F]	592
Giac [F]	592
Mupad [F(-1)]	592

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{b^2 d \sqrt{\cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2719}

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^2/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(2*\text{Sqrt}[b*\text{Cos}[c + d*x])*E[\text{EllipticE}[(c + d*x)/2, 2]]/(b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)^{(v_*)^{(m_*)}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] := \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \sqrt{b \cos(c + dx)} dx}{b^2} \\ &= \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{b^2 \sqrt{\cos(c + dx)}} \\ &= \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{b^2 d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{b^2 d \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(3/2),x]
```

```
[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]
])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 2.03 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}$
risch	$-\frac{i(e^{2i(dx+c)}+1)\sqrt{2}e^{-i(dx+c)}}{db\sqrt{(e^{2i(dx+c)}+1)be^{-i(dx+c)}}}-i\left(-\frac{2(b e^{2i(dx+c)}+b)}{b\sqrt{e^{i(dx+c)}(b e^{2i(dx+c)}+b)}}+\frac{i\sqrt{-i(e^{i(dx+c)}+i)}\sqrt{2}\sqrt{i(e^{i(dx+c)}-i)}\sqrt{ie^{i(dx+c)}}(-2iE(\sqrt{-i(e^{i(dx+c)}-i)}\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}})}))}{\sqrt{b e^{3i(dx+c)}+b e^{i(dx+c)}}}\right)db\sqrt{(e^{2i(dx+c)}+1)}$

[In] `int(cos(d*x+c)^2/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^(1/2)*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*\cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(\cos(1/2*d*x+1/2*c),2^(1/2))/b/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.61

$$\int \frac{\cos^2(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)))}{(b\cos(c+dx))^{3/2}}$$

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out] $(I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))-I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))))/(b^2*d)$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(3/2), x)

$$3.121 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	593
Rubi [A] (verified)	593
Mathematica [A] (verified)	594
Maple [B] (verified)	594
Fricas [C] (verification not implemented)	595
Sympy [F]	595
Maxima [F]	595
Giac [F]	595
Mupad [F(-1)]	596

Optimal result

Integrand size = 19, antiderivative size = 41

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd\sqrt{b \cos(c+dx)}}$$

[Out] 2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 2721, 2720}

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd\sqrt{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(3/2),x]

[Out] (2*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(b*d*sqrt[b*Cos[c + d*x]])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{b} \\ &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b \sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{bd \sqrt{b \cos(c+dx)}}$$

```
[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(3/2), x]
```

```
[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(b*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 1.39 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result	size
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$	144

```
[In] int(cos(d*x+c)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] -2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(sin(1/2*d*x
+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c
)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))/sin(
1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.08 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{-i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))}{b^2 d}$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^2*d)

Sympy [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(cos(c + d*x)/(b*cos(c + d*x))**(3/2), x)

Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{3/2}} dx$$

```
[In] int(cos(c + d*x)/(b*cos(c + d*x))^(3/2), x)
```

```
[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(3/2), x)
```

3.122 $\int \frac{1}{(b \cos(c+dx))^{3/2}} dx$

Optimal result	597
Rubi [A] (verified)	597
Mathematica [A] (verified)	598
Maple [B] (verified)	598
Fricas [C] (verification not implemented)	599
Sympy [F]	599
Maxima [F]	599
Giac [F]	600
Mupad [F(-1)]	600

Optimal result

Integrand size = 12, antiderivative size = 68

$$\int \frac{1}{(b \cos(c+dx))^{3/2}} dx = -\frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{bd \sqrt{b \cos(c+dx)}}$$

[Out] 2*sin(d*x+c)/b/d/(b*cos(d*x+c))^(1/2)-2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^2/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$\int \frac{1}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)}{bd \sqrt{b \cos(c+dx)}} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^2 d \sqrt{\cos(c+dx)}}$$

[In] Int[(b*Cos[c + d*x])^(-3/2),x]

[Out] (-2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]]) + (2*Sin[c + d*x])/(b*d*Sqrt[b*Cos[c + d*x]])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(c + dx)}{bd\sqrt{b \cos(c + dx)}} - \frac{\int \sqrt{b \cos(c + dx)} dx}{b^2} \\ &= \frac{2 \sin(c + dx)}{bd\sqrt{b \cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{b^2 \sqrt{\cos(c + dx)}} \\ &= -\frac{2\sqrt{b \cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right)}{b^2 d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)}{bd\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \frac{2\left(-\sqrt{\cos(c + dx)}E\left(\frac{1}{2}(c + dx) \mid 2\right) + \sin(c + dx)\right)}{bd\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(-3/2),x]
```

```
[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(b*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(88) = 176.

Time = 1.74 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.91

method	result
default	$-\frac{2\left(-2\cos\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}b + b\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) + \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)} - 1\sqrt{-2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right)}\right)}{b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - \left(\sin^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2} + \frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2} + \frac{c}{2}\right)\right) - 1\right)}}$

[In] `int(1/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2/b*(-2*\cos(1/2*d*x+1/2*c)*(-2*\sin(1/2*d*x+1/2*c)^4+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*(-2*\sin(1/2*d*x+1/2*c)^4+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)})}{(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.53

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \frac{-i \sqrt{2} \sqrt{b} \cos(dx + c) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c)))}{(b \cos(c + dx))^{3/2}}$$

[In] `integrate(1/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$(-I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c))) + I*\sqrt{2}*\sqrt{b}*\cos(d*x + c)*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c))) + 2*\sqrt{b*\cos(d*x + c)}*\sin(d*x + c))/(b^2*d*\cos(d*x + c))$$

Sympy [F]

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(c + dx))^{\frac{3}{2}}} dx$$

[In] `integrate(1/(b*cos(d*x+c))**(3/2),x)`

[Out] `Integral((b*cos(c + d*x))**(-3/2), x)`

Maxima [F]

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] `integrate(1/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(3/2), x)`

Giac [F]

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(-3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(c + dx))^{3/2}} dx$$

[In] int(1/(b*cos(c + d*x))^(3/2),x)

[Out] int(1/(b*cos(c + d*x))^(3/2), x)

3.123 $\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx$

Optimal result	601
Rubi [A] (verified)	601
Mathematica [A] (verified)	602
Maple [B] (verified)	603
Fricas [C] (verification not implemented)	603
Sympy [F]	603
Maxima [F]	604
Giac [F]	604
Mupad [F(-1)]	604

Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3bd\sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}}$$

[Out] 2/3*sin(d*x+c)/d/(b*cos(d*x+c))^(3/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2716, 2721, 2720}

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2 \sin(c+dx)}{3d(b \cos(c+dx))^{3/2}} + \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3bd\sqrt{b \cos(c+dx)}}$$

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(3*b*d*Sqrt[b*Cos[c + d*x]]) + (2*Sin[c + d*x])/(3*d*(b*Cos[c + d*x])^(3/2))

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
&= \frac{2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{3b} \\
&= \frac{2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b \sqrt{b \cos(c + dx)}} \\
&= \frac{2 \sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3bd \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx)}{3d(b \cos(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.74

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{2 \left(\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx) \right)}{3bd \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(3/2), x]
```

```
[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b*d*Sq
rt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(85) = 170.

Time = 2.15 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.49

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3b\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] `int(sec(d*x+c)/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/3*(-2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)})*\sin(1/2*d*x+1/2*c)^2-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticF(\cos(1/2*d*x+1/2*c),2^{(1/2)}))/b*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/(2*\cos(1/2*d*x+1/2*c)^2-1)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.49

$$\int \frac{\sec(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\cos(dx+c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c)) + 2*\sqrt{b*\cos(dx+c)}*\sin(dx+c)}{(b^2*d*\cos(dx+c))^2}$$

[In] `integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out]
$$1/3*(-I*\sqrt{2}*\sqrt{b}*\cos(dx+c)^2*\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + I*\sin(dx+c)) + I*\sqrt{2}*\sqrt{b}*\cos(dx+c)^2*\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - I*\sin(dx+c)) + 2*\sqrt{b*\cos(dx+c)}*\sin(dx+c))/(b^2*d*\cos(dx+c)^2)$$

Sympy [F]

$$\int \frac{\sec(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \int \frac{\sec(c+dx)}{(b\cos(c+dx))^{\frac{3}{2}}} dx$$

[In] `integrate(sec(d*x+c)/(b*cos(d*x+c))**(3/2),x)`

[Out] `Integral(sec(c + d*x)/(b*cos(c + d*x))**(3/2), x)`

Maxima [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{3/2}} dx$$

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(3/2)), x)

$$3.124 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	605
Rubi [A] (verified)	605
Mathematica [A] (verified)	607
Maple [B] (verified)	607
Fricas [C] (verification not implemented)	608
Sympy [F]	608
Maxima [F]	608
Giac [F]	609
Mupad [F(-1)]	609

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx = -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5b^2 d \sqrt{\cos(c+dx)}} + \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd \sqrt{b \cos(c+dx)}}$$

[Out] $2/5*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{5b^2 d \sqrt{\cos(c+dx)}} + \frac{6 \sin(c+dx)}{5bd \sqrt{b \cos(c+dx)}} + \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]^2/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

```
Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2716

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c+d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c-Pi/2+d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c+d*x])^n/Sin[c+d*x]^n, Int[Sin[c+d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^2 \int \frac{1}{(b \cos(c+dx))^{7/2}} dx \\
 &= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{3}{5} \int \frac{1}{(b \cos(c+dx))^{3/2}} dx \\
 &= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}} - \frac{3 \int \sqrt{b \cos(c+dx)} dx}{5b^2} \\
 &= \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}} - \frac{\left(3\sqrt{b \cos(c+dx)}\right) \int \sqrt{\cos(c+dx)} dx}{5b^2 \sqrt{\cos(c+dx)}} \\
 &= -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5b^2 d \sqrt{\cos(c+dx)}} + \frac{2b \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5bd\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.69

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{-6\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + 6\sin(c+dx) + 2\sec(c+dx)\tan(c+dx)}{5bd\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(3/2), x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*b*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(110) = 220.

Time = 2.65 (sec) , antiderivative size = 367, normalized size of antiderivative = 3.74

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{(24\cos(\frac{dx}{2} + \frac{c}{2})(\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}} E(\cos(\frac{dx}{2} + \frac{c}{2}))$

[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out]
$$-\frac{2}{5} * (-(-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) * b * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) / b ^ 2 / \sin(1/2 * d * x + 1/2 * c) ^ 3 / (8 * \sin(1/2 * d * x + 1/2 * c) ^ 6 - 12 * \sin(1/2 * d * x + 1/2 * c) ^ 4 + 6 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) * (24 * \cos(1/2 * d * x + 1/2 * c) * \sin(1/2 * d * x + 1/2 * c) ^ 6 - 12 * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2)) * \sin(1/2 * d * x + 1/2 * c) ^ 4 - 24 * \sin(1/2 * d * x + 1/2 * c) ^ 4 * \cos(1/2 * d * x + 1/2 * c) + 12 * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2)) * \sin(1/2 * d * x + 1/2 * c) ^ 2 + 8 * \sin(1/2 * d * x + 1/2 * c) ^ 2 * \cos(1/2 * d * x + 1/2 * c) - 3 * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2))) * (-2 * \sin(1/2 * d * x + 1/2 * c) ^ 4 * b + b * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) / ((2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * b) ^ (1/2) / d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.23

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^3 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c)))}{(b \cos(c + dx))^{3/2}}$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 1)*sin(d*x + c))/(b^2*d*cos(d*x + c)^3)

Sympy [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(3/2), x)

Maxima [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{3/2}} dx$$

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(3/2)), x)

3.125 $\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx$

Optimal result	610
Rubi [A] (verified)	610
Mathematica [A] (verified)	612
Maple [B] (verified)	612
Fricas [C] (verification not implemented)	613
Sympy [F]	613
Maxima [F]	613
Giac [F]	613
Mupad [F(-1)]	614

Optimal result

Integrand size = 21, antiderivative size = 97

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21bd\sqrt{b \cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}$$

[Out] $2/7*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(3/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec), antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21bd\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c+d*x]^3/(b*\operatorname{Cos}[c+d*x])^{(3/2)}, x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(21*b*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*b^2*\operatorname{Sin}[c+d*x])/(7*d*(b*\operatorname{Cos}[c+d*x])^{(7/2)}) + (10*\operatorname{Sin}[c+d*x])/(21*d*(b*\operatorname{Cos}[c+d*x])^{(3/2)})$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c+d*x]*((b*\sin[c+d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\sin[c+d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\sin[c+d*x])^n/\sin[c+d*x]^n, \text{Int}[\sin[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^3 \int \frac{1}{(b \cos(c+dx))^{9/2}} dx \\
 &= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{1}{7}(5b) \int \frac{1}{(b \cos(c+dx))^{5/2}} dx \\
 &= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{5 \int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{21b} \\
 &= \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}} + \frac{(5\sqrt{\cos(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b\sqrt{b \cos(c+dx)}} \\
 &= \frac{10\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21bd\sqrt{b \cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21d(b \cos(c+dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.68

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 2(5 + 3\sec^2(c+dx)) \tan(c+dx)}{21bd\sqrt{b \cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(3/2), x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*b*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(109) = 218.

Time = 2.34 (sec) , antiderivative size = 398, normalized size of antiderivative = 4.10

method	result
default	$-\frac{2\left(-40\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-40\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+60\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{\dots}$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^6-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^4+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^2-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2)))/b*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.19

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{-5i \sqrt{2} \sqrt{b} \cos(dx+c)^4 \operatorname{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))}{(b \cos(c+dx))^{3/2}}$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*(5*cos(d*x + c)^2 + 3)*sin(d*x + c))/(b^2*d*cos(d*x + c)^4)

Sympy [F]

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(3/2), x)

Maxima [F]

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \int \frac{\sec(dx+c)^3}{(b \cos(dx+c))^{\frac{3}{2}}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{3/2}} dx$$

```
[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2)),x)
```

```
[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(3/2)), x)
```

3.126 $\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx$

Optimal result	615
Rubi [A] (verified)	615
Mathematica [A] (verified)	617
Maple [B] (verified)	617
Fricas [C] (verification not implemented)	618
Sympy [F]	618
Maxima [F]	618
Giac [F]	619
Mupad [F(-1)]	619

Optimal result

Integrand size = 21, antiderivative size = 126

$$\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx = -\frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd \sqrt{b \cos(c+dx)}}$$

[Out] $2/9*b^3*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(9/2)}+14/45*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+14/15*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}-14/15*(\cos(1/2*d*x+1/2*c))^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^2/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec^4(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{2b^3 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} - \frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15b^2 d \sqrt{\cos(c+dx)}} + \frac{14b \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15bd \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]^4/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^{(9/2)}) + (14*b*\text{Sin}[c +$

$d*x]]/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*\text{Sin}[c + d*x])/(15*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^(m+n), x], x] /; \text{FreeQ}\{b, n\}, x] \&\& \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\text{sin}[(c_.) + (d_)*(x_)]^(n_), x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^(n+1)/(b*d*(n+1))), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^(n+2), x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\text{sin}[(c_.) + (d_)*(x_)]^(n_), x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^4 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9}(7b^2) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{7}{15} \int \frac{1}{(b \cos(c + dx))^{3/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14 \sin(c + dx)}{15bd\sqrt{b \cos(c + dx)}} - \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b^2} \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14b \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} \\
 &\quad + \frac{14 \sin(c + dx)}{15bd\sqrt{b \cos(c + dx)}} - \frac{(7\sqrt{b \cos(c + dx)}) \int \sqrt{\cos(c + dx)} dx}{15b^2 \sqrt{\cos(c + dx)}}
 \end{aligned}$$

$$= -\frac{14\sqrt{b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15b^2d\sqrt{\cos(c+dx)}} + \frac{2b^3\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}}$$

$$+ \frac{14b\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}} + \frac{14\sin(c+dx)}{15bd\sqrt{b\cos(c+dx)}}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.63

$$\int \frac{\sec^4(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{-42\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + 42\sin(c+dx) + 2\sec(c+dx)(7+5\sec^2(c+dx))}{45bd\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^4/(b*Cos[c + d*x])^(3/2), x]

[Out] (-42*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*b*d*sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. 2(134) = 268.

Time = 3.12 (sec) , antiderivative size = 416, normalized size of antiderivative = 3.30

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{\left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{144b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5} - \frac{7\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{180b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5}\right)}$

[In] int(sec(d*x+c)^4/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b*(-1/144*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/180*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^3-14/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)+7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.04

$$\int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{-21i \sqrt{2} \sqrt{b} \cos(dx + c)^5 \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(dx + c)))}{(b \cos(c + dx))^{3/2}}$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*cos(d*x + c)^4 + 7*cos(d*x + c)^2 + 5)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(b^2*d*cos(d*x + c)^5)

Sympy [F]

$$\int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx$$

[In] integrate(sec(d*x+c)**4/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sec(c + d*x)**4/(b*cos(c + d*x))**(3/2), x)

Maxima [F]

$$\int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^4}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

Giac [F]

$$\int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sec(dx + c)^4}{(b \cos(dx + c))^{3/2}} dx$$

[In] integrate(sec(d*x+c)^4/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^4(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^4 (b \cos(c + dx))^{3/2}} dx$$

[In] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^4*(b*cos(c + d*x))^(3/2)), x)

3.127 $\int \frac{\cos^8(c+dx)}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	620
Rubi [A] (verified)	620
Mathematica [A] (verified)	622
Maple [A] (verified)	622
Fricas [C] (verification not implemented)	623
Sympy [F(-1)]	623
Maxima [F]	623
Giac [F]	623
Mupad [F(-1)]	624

Optimal result

Integrand size = 21, antiderivative size = 128

$$\int \frac{\cos^8(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77b^2d\sqrt{b \cos(c+dx)}} + \frac{30\sqrt{b \cos(c+dx)} \sin(c+dx)}{77b^3d} + \frac{18(b \cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d} + \frac{2(b \cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d}$$

[Out] 18/77*(b*cos(d*x+c))^(5/2)*sin(d*x+c)/b^5/d+2/11*(b*cos(d*x+c))^(9/2)*sin(d*x+c)/b^7/d+30/77*(cos(1/2*d*x+1/2*c))^2^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c), 2^(1/2))*cos(d*x+c)^(1/2)/b^2/d/(b*cos(d*x+c))^(1/2)+30/77*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/b^3/d

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^8(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{9/2}}{11b^7d} + \frac{18 \sin(c+dx)(b \cos(c+dx))^{5/2}}{77b^5d} + \frac{30 \sin(c+dx)\sqrt{b \cos(c+dx)}}{77b^3d} + \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77b^2d\sqrt{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^8/(b*Cos[c + d*x])^(5/2), x]

[Out] (30*sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2])/(77*b^2*d*sqrt[b*Cos[c + d*x]]) + (30*sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(77*b^3*d) + (18*(b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(77*b^5*d) + (2*(b*Cos[c + d*x])^(9/2)*Sin[c + d*x])/(11*b^7*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{11/2} dx}{b^8} \\
 &= \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^7d} + \frac{9 \int (b \cos(c + dx))^{7/2} dx}{11b^6} \\
 &= \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^5d} + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^7d} + \frac{45 \int (b \cos(c + dx))^{3/2} dx}{77b^4} \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77b^3d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^5d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^7d} + \frac{15 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{77b^2} \\
 &= \frac{30\sqrt{b \cos(c + dx)} \sin(c + dx)}{77b^3d} + \frac{18(b \cos(c + dx))^{5/2} \sin(c + dx)}{77b^5d} \\
 &\quad + \frac{2(b \cos(c + dx))^{9/2} \sin(c + dx)}{11b^7d} + \frac{\left(15\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{77b^2\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

$$= \frac{30\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{77b^2d\sqrt{b\cos(c+dx)}} + \frac{30\sqrt{b\cos(c+dx)} \sin(c+dx)}{77b^3d} \\ + \frac{18(b\cos(c+dx))^{5/2} \sin(c+dx)}{77b^5d} + \frac{2(b\cos(c+dx))^{9/2} \sin(c+dx)}{11b^7d}$$

Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.59

$$\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{480\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 347 \sin(2(c+dx)) + 64 \sin(4(c+dx))}{1232b^2d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^8/(b*Cos[c + d*x])^(5/2), x]

[Out] (480*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 347*Sin[2*(c + d*x)] + 64*Sin[4*(c + d*x)] + 7*Sin[6*(c + d*x)])/(1232*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 5.88 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.84

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}{\left(448\left(\cos^{13}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1568\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+2384\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2040\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1568\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-448\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+64\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{77b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}}$

[In] int(cos(d*x+c)^8/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out]
$$-2/77*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(448*\cos(1/2*d*x+1/2*c)^{13}-1568*\cos(1/2*d*x+1/2*c)^{11}+2384*\cos(1/2*d*x+1/2*c)^9-2040*\cos(1/2*d*x+1/2*c)^7+1084*\cos(1/2*d*x+1/2*c)^5-370*\cos(1/2*d*x+1/2*c)^3+15*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}*\operatorname{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})+62*\cos(1/2*d*x+1/2*c))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.79

$$\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{2(7\cos(dx+c)^4 + 9\cos(dx+c)^2 + 15)\sqrt{b\cos(dx+c)}\sin(dx+c) - 15i\sqrt{2}\sqrt{b\cos(dx+c)}}{(b\cos(c+dx))^{5/2}}$$

[In] integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/77*(2*(7*cos(d*x + c)^4 + 9*cos(d*x + c)^2 + 15)*sqrt(b*cos(d*x + c))*sin(d*x + c) - 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 15*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^3*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**8/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(dx+c)^8}{(b\cos(dx+c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^8/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^8(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(dx+c)^8}{(b\cos(dx+c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^8/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^8/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^8(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^8}{(b \cos(c + dx))^{5/2}} dx$$

```
[In] int(cos(c + d*x)^8/(b*cos(c + d*x))^(5/2), x)
```

```
[Out] int(cos(c + d*x)^8/(b*cos(c + d*x))^(5/2), x)
```


$$3.128 \quad \int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	625
Rubi [A] (verified)	625
Mathematica [A] (verified)	627
Maple [A] (verified)	627
Fricas [C] (verification not implemented)	627
Sympy [F(-1)]	628
Maxima [F]	628
Giac [F]	628
Mupad [F(-1)]	628

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{14\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15b^3 d \sqrt{\cos(c+dx)}} + \frac{14(b \cos(c+dx))^{3/2} \sin(c+dx)}{45b^4 d} + \frac{2(b \cos(c+dx))^{7/2} \sin(c+dx)}{9b^6 d}$$

[Out] 14/45*(b*cos(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+2/9*(b*cos(d*x+c))^(7/2)*sin(d*x+c)/b^6/d+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^3/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^7(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{7/2}}{9b^6 d} + \frac{14 \sin(c+dx)(b \cos(c+dx))^{3/2}}{45b^4 d} + \frac{14 E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15b^3 d \sqrt{\cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^7/(b*Cos[c + d*x])^(5/2),x]

[Out] (14*sqrt[b*cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(15*b^3*d*sqrt[Cos[c + d*x]]) + (14*(b*cos[c + d*x])^(3/2)*Sin[c + d*x])/(45*b^4*d) + (2*(b*cos[c + d*x])^(7/2)*Sin[c + d*x])/(9*b^6*d)

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c+d*x]*(b*\text{Sin}[c+d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c+d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c+d*x])^{n-1}/\text{Sin}[c+d*x]^n, \text{Int}[\text{Sin}[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{9/2} dx}{b^7} \\
 &= \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^6 d} + \frac{7 \int (b \cos(c + dx))^{5/2} dx}{9b^5} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^4 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^6 d} + \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b^3} \\
 &= \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^4 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^6 d} \\
 &\quad + \frac{\left(7\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15b^3 \sqrt{\cos(c + dx)}} \\
 &= \frac{14\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{15b^3 d \sqrt{\cos(c + dx)}} \\
 &\quad + \frac{14(b \cos(c + dx))^{3/2} \sin(c + dx)}{45b^4 d} + \frac{2(b \cos(c + dx))^{7/2} \sin(c + dx)}{9b^6 d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.52 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.74

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{168\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + \cos(c+dx)(38\sin(2(c+dx)) + 5\sin(4(c+dx)))}{180b^2d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^7/(b*Cos[c + d*x])^(5/2), x]

[Out] (168*sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*(38*Sin[2*(c + d*x)] + 5*Sin[4*(c + d*x)]))/(180*b^2*d*sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 4.42 (sec) , antiderivative size = 223, normalized size of antiderivative = 2.23

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(160\left(\cos^{11}\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-480\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+616\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-432\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{45b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] int(cos(d*x+c)^7/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/45*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b^2*(160*cos(1/2*d*x+1/2*c)^11-480*cos(1/2*d*x+1/2*c)^9+616*cos(1/2*d*x+1/2*c)^7-432*cos(1/2*d*x+1/2*c)^5+160*cos(1/2*d*x+1/2*c)^3-21*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticE(cos(1/2*d*x+1/2*c), 2)^(1/2))-24*cos(1/2*d*x+1/2*c)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2)^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.04

$$\int \frac{\cos^7(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{2(5\cos(dx+c)^3 + 7\cos(dx+c))\sqrt{b\cos(dx+c)}\sin(dx+c) + 21i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4, 0, \cos(dx+c) + i\sin(dx+c)) - 21i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{(b^3d)}$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] 1/45*(2*(5*cos(d*x + c)^3 + 7*cos(d*x + c))*sqrt(b*cos(d*x + c))*sin(d*x + c) + 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) - 21*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))))/(b^3*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^7(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**7/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^7(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^7}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^7(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^7}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^7/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^7/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^7(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^7}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^7/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^7/(b*cos(c + d*x))^(5/2), x)

$$3.129 \quad \int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	629
Rubi [A] (verified)	629
Mathematica [A] (verified)	631
Maple [A] (verified)	631
Fricas [C] (verification not implemented)	631
Sympy [F(-1)]	632
Maxima [F]	632
Giac [F]	632
Mupad [F(-1)]	632

Optimal result

Integrand size = 21, antiderivative size = 100

$$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21b^2d\sqrt{b \cos(c+dx)}} + \frac{10\sqrt{b \cos(c+dx)} \sin(c+dx)}{21b^3d} + \frac{2(b \cos(c+dx))^{5/2} \sin(c+dx)}{7b^5d}$$

[Out] $2/7*(b*\cos(d*x+c))^{(5/2)}*\sin(d*x+c)/b^5/d+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}+10/21*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^6(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{5/2}}{7b^5d} + \frac{10 \sin(c+dx)\sqrt{b \cos(c+dx)}}{21b^3d} + \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21b^2d\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^6/(b*\operatorname{Cos}[c+d*x])^{(5/2)}, x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(21*b^2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (10*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(21*b^3*d) + (2*(b*\operatorname{Cos}[c+d*x])^{(5/2)}*\operatorname{Sin}[c+d*x])/(7*b^5*d)$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] := \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] := \text{Simp}[(-b)*\text{Cos}[c+d*x]*(b*\sin[c+d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\sin[c+d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_)+(d_)*(x_)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c-\text{Pi}/2+d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] := \text{Dist}[(b*\sin[c+d*x])^{n-1}/\sin[c+d*x]^n, \text{Int}[\sin[c+d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{7/2} dx}{b^6} \\
 &= \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^5d} + \frac{5 \int (b \cos(c + dx))^{3/2} dx}{7b^4} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^3d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^5d} + \frac{5 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{21b^2} \\
 &= \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^3d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^5d} \\
 &\quad + \frac{\left(5\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21b^2\sqrt{b \cos(c + dx)}} \\
 &= \frac{10\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21b^2d\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{10\sqrt{b \cos(c + dx)} \sin(c + dx)}{21b^3d} + \frac{2(b \cos(c + dx))^{5/2} \sin(c + dx)}{7b^5d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.66

$$\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{40\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 26\sin(2(c+dx)) + 3\sin(4(c+dx))}{84b^2d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^6/(b*Cos[c + d*x])^(5/2), x]

[Out] (40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)])/(84*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.77 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.10

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(48\left(\cos^9\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-120\left(\cos^7\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+128\left(\cos^5\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-72\left(\cos^3\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{21b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] int(cos(d*x+c)^6/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/21*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b^2*(48*cos(1/2*d*x+1/2*c)^9-120*cos(1/2*d*x+1/2*c)^7+128*cos(1/2*d*x+1/2*c)^5-72*cos(1/2*d*x+1/2*c)^3+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))+16*cos(1/2*d*x+1/2*c))/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.91

$$\int \frac{\cos^6(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{2\sqrt{b\cos(dx+c)}(3\cos(dx+c)^2+5)\sin(dx+c)-5i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}}{(b\cos(c+dx))^{5/2}}$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] 1/21*(2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 5)*sin(d*x + c) - 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)))/(b^3*d)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**6/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^6}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^6}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^6/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^6/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^6(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^6}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^6/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^6/(b*cos(c + d*x))^(5/2), x)

$$3.130 \quad \int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	633
Rubi [A] (verified)	633
Mathematica [A] (verified)	634
Maple [B] (verified)	635
Fricas [C] (verification not implemented)	635
Sympy [F(-1)]	635
Maxima [F]	636
Giac [F]	636
Mupad [F(-1)]	636

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5b^3 d \sqrt{\cos(c+dx)}} + \frac{2(b \cos(c+dx))^{3/2} \sin(c+dx)}{5b^4 d}$$

[Out] $2/5*(b*\cos(d*x+c))^{(3/2)}*\sin(d*x+c)/b^4/d+6/5*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2719}

$$\int \frac{\cos^5(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx)(b \cos(c+dx))^{3/2}}{5b^4 d} + \frac{6 E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5b^3 d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^5/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*(b*\text{Cos}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(5*b^4*d)$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{5/2} dx}{b^5} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^4d} + \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b^3} \\
 &= \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^4d} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b^3 \sqrt{\cos(c + dx)}} \\
 &= \frac{6\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5b^3d\sqrt{\cos(c + dx)}} + \frac{2(b \cos(c + dx))^{3/2} \sin(c + dx)}{5b^4d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.85

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{6\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + \cos(c + dx) \sin(2(c + dx))}{5b^2d\sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^5/(b*Cos[c + d*x])^(5/2), x]
```

```
[Out] (6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]*Sin[2*(c + d*x)])/(5*b^2*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(88) = 176.

Time = 2.81 (sec) , antiderivative size = 213, normalized size of antiderivative = 2.96

method	result
default	$-\frac{2\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(-8\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+8\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}{5b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] `int(cos(d*x+c)^5/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(-8*\cos(1/2*d*x+1/2*c)*\sin(1/2*d*x+1/2*c)^6+8*\sin(1/2*d*x+1/2*c)^4*\cos(1/2*d*x+1/2*c)-2*\sin(1/2*d*x+1/2*c)^2*\cos(1/2*d*x+1/2*c)-3*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)}*EllipticE(\cos(1/2*d*x+1/2*c),2^{(1/2)}))/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.26

$$\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{2\sqrt{b\cos(dx+c)}\cos(dx+c)\sin(dx+c) + 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)) + i\sin(dx+c)) - 3i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c) - i\sin(dx+c)))}{(b^3*d)}$$

[In] `integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out]
$$1/5*(2*\sqrt{b*\cos(d*x+c)}*\cos(d*x+c)*\sin(d*x+c) + 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)) + I*\sin(d*x+c))) - 3*I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c) - I*\sin(d*x+c))))/(b^3*d)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^5(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**5/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^5}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^5}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^5/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^5/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^5(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^5}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^5/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^5/(b*cos(c + d*x))^(5/2), x)

$$3.131 \quad \int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	637
Rubi [A] (verified)	637
Mathematica [A] (verified)	638
Maple [B] (verified)	639
Fricas [C] (verification not implemented)	639
Sympy [F(-1)]	639
Maxima [F]	640
Giac [F]	640
Mupad [F(-1)]	640

Optimal result

Integrand size = 21, antiderivative size = 72

$$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2\sqrt{b \cos(c+dx)} \sin(c+dx)}{3b^3 d}$$

[Out] $2/3*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}+2/3*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2715, 2721, 2720}

$$\int \frac{\cos^4(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{3b^3 d} + \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^4/(b*\operatorname{Cos}[c+d*x])^{(5/2)}, x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(3*b^2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(3*b^3*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n-1)/(d*n)), x] + Dist[b^2*((n-1)/n), Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\int (b \cos(c + dx))^{3/2} dx}{b^4} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^3 d} + \frac{\int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{3b^2} \\
 &= \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^3 d} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2 \sqrt{b \cos(c + dx)}} \\
 &= \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3b^2 d \sqrt{b \cos(c + dx)}} + \frac{2\sqrt{b \cos(c + dx)} \sin(c + dx)}{3b^3 d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.75

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \sin(2(c + dx))}{3b^2 d \sqrt{b \cos(c + dx)}}$$

`[In] Integrate[Cos[c + d*x]^4/(b*Cos[c + d*x])^(5/2), x]`

`[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)])/(3*b^2*d*Sqrt[b*Cos[c + d*x]])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(88) = 176.

Time = 2.44 (sec) , antiderivative size = 190, normalized size of antiderivative = 2.64

method	result
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(4\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2}\right)}{3b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b}$

[In] `int(cos(d*x+c)^4/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{2}{3}\frac{\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\sqrt{b}}{\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b^2*\left(4\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)-2\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)+\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}*\left(2*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)^{\frac{1}{2}}*\text{EllipticF}\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)\right)}{\left(-b*\left(2*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4-\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)\right)^{\frac{1}{2}}/\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)/\left(\left(2*\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b\right)^{\frac{1}{2}}/d}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.10

$$\int \frac{\cos^4(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c))+i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(dx+c)-i\sin(dx+c))}{3b^2\sqrt{b}\cos(dx+c)}$$

[In] `integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2),x,algorithm="fricas")`

[Out]
$$\frac{1}{3}\frac{\left(-I*\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c))+I*\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))+2*\sqrt{b*\cos(d*x+c)}*\sin(d*x+c)\right)}{b^2\sqrt{b}\cos(dx+c)}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^4(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**4/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^4}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^4}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^4/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^4}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^4/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^4/(b*cos(c + d*x))^(5/2), x)

$$3.132 \quad \int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	641
Rubi [A] (verified)	641
Mathematica [A] (verified)	642
Maple [B] (verified)	642
Fricas [C] (verification not implemented)	643
Sympy [F(-1)]	643
Maxima [F]	644
Giac [F]	644
Mupad [F(-1)]	644

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{b^3 d \sqrt{\cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2719}

$$\int \frac{\cos^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^3/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(2*\text{Sqrt}[b*\text{Cos}[c + d*x])*E[\text{EllipticE}[(c + d*x)/2, 2]]/(b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)^{(v_*)^{(m_*)}*(b_*)^{(v_*)^{(n_*)}}, x_Symbol] := \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \sqrt{b \cos(c + dx)} dx}{b^3} \\ &= \frac{\sqrt{b \cos(c + dx)} \int \sqrt{\cos(c + dx)} dx}{b^3 \sqrt{\cos(c + dx)}} \\ &= \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{b^3 d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{2\sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{b^3 d \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^3/(b*Cos[c + d*x])^(5/2),x]
```

```
[Out] (2*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(b^3*d*Sqrt[Cos[c + d*x]
])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 1.91 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result
default	$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}E\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}$
risch	$-\frac{i\left(e^{2i(dx+c)}+1\right)\sqrt{2}e^{-i(dx+c)}}{db^2\sqrt{\left(e^{2i(dx+c)}+1\right)be^{-i(dx+c)}}}-\frac{i\left(-\frac{2\left(be^{2i(dx+c)}+b\right)}{b\sqrt{e^{i(dx+c)}\left(be^{2i(dx+c)}+b\right)}}+\frac{i\sqrt{-i\left(e^{i(dx+c)}+i\right)}\sqrt{2}\sqrt{i\left(e^{i(dx+c)}-i\right)}\sqrt{ie^{i(dx+c)}}\left(-2iE\left(\sqrt{-\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)\right)}{\sqrt{be^{3i(dx+c)}+be^{i(dx+c)}}}\right)}{db^2\sqrt{\left(e^{2i(dx+c)}+1\right)be^{-i(dx+c)}}}$

[In] `int(cos(d*x+c)^3/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $2\left(\left(2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}*\left(\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)^{\frac{1}{2}}*\left(-2\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2+1\right)^{\frac{1}{2}}*EllipticE\left(\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right),2^{\frac{1}{2}}\right)/b^2/\left(-b*\left(2*\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^4-\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2\right)\right)^{\frac{1}{2}}/\sin\left(\frac{1}{2}d*x+\frac{1}{2}c\right)/\left(\left(2*\cos\left(\frac{1}{2}d*x+\frac{1}{2}c\right)^2-1\right)*b\right)^{\frac{1}{2}}/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.61

$$\int \frac{\cos^3(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{i\sqrt{2}\sqrt{b}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)+i\sin(dx+c)))}{(b\cos(c+dx))^{5/2}}$$

[In] `integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out] $(I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))-I*\sqrt{2}*\sqrt{b}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c))))/(b^3*d)$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**3/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^3}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^3}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^3/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^3/(b*cos(c + d*x))^(5/2), x)

3.133 $\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	645
Rubi [A] (verified)	645
Mathematica [A] (verified)	646
Maple [B] (verified)	646
Fricas [C] (verification not implemented)	647
Sympy [F(-1)]	647
Maxima [F]	647
Giac [F]	648
Mupad [F(-1)]	648

Optimal result

Integrand size = 21, antiderivative size = 41

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2)^{(1/2)}*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 2721, 2720}

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2/(b*\operatorname{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*\operatorname{EllipticF}[(c + d*x)/2, 2])/(b^2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n, x\} \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{b^2} \\ &= \frac{\sqrt{\cos(c+dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{2\sqrt{\cos(c+dx)} \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.93

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \cos^{5/2}(c+dx) \text{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{d(b \cos(c+dx))^{5/2}}$$

```
[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]
```

```
[Out] (2*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2])/(d*(b*Cos[c + d*x])^(5/2))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. 2(63) = 126.

Time = 1.39 (sec) , antiderivative size = 144, normalized size of antiderivative = 3.51

method	result	size
default	$-\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{-2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+1}F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)bd}}$	144

```
[In] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)
```

[Out] $-2*((2*\cos(1/2*d*x+1/2*c)^2-1)*b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)}/b^2*(\sin(1/2*d*x+1/2*c)^2)^{(1/2)}*(-2*\cos(1/2*d*x+1/2*c)^2+1)^{(1/2)}/(-b*(2*\sin(1/2*d*x+1/2*c)^4-\sin(1/2*d*x+1/2*c)^2))^{(1/2)}*\text{EllipticF}(\cos(1/2*d*x+1/2*c), 2^{(1/2)})/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)}/d$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.46

$$\int \frac{\cos^2(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i\sin(dx+c)) + i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4, 0, \cos(dx+c) - i\sin(dx+c))}{b^3d}$$

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")`

[Out] $(-I*\sqrt{2}*\sqrt{b}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) + I*\sin(d*x + c)) + I*\sqrt{2}*\sqrt{b}*\text{weierstrassPInverse}(-4, 0, \cos(d*x + c) - I*\sin(d*x + c)))/(b^3*d)$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(5/2), x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^2(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \int \frac{\cos(dx+c)^2}{(b\cos(dx+c))^{\frac{5}{2}}} dx$$

[In] `integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)`

Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(5/2), x)

3.134 $\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	649
Rubi [A] (verified)	649
Mathematica [A] (verified)	650
Maple [B] (verified)	651
Fricas [C] (verification not implemented)	651
Sympy [F(-1)]	651
Maxima [F]	652
Giac [F]	652
Mupad [F(-1)]	652

Optimal result

Integrand size = 19, antiderivative size = 68

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx = -\frac{2\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^3 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $2*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{2 E\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \cos(c+dx)}}{b^3 d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(-2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2716

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\int \frac{1}{(b \cos(c+dx))^{3/2}} dx}{b} \\
&= \frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\int \sqrt{b \cos(c+dx)} dx}{b^3} \\
&= \frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sqrt{b \cos(c+dx)} \int \sqrt{\cos(c+dx)} dx}{b^3 \sqrt{\cos(c+dx)}} \\
&= -\frac{2 \sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{b^3 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.74

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \left(-\sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right) + \sin(c+dx) \right)}{b^2 d \sqrt{b \cos(c+dx)}}$$

```
[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(5/2),x]
```

```
[Out] (2*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/(b^2*d
*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(88) = 176.

Time = 1.74 (sec) , antiderivative size = 198, normalized size of antiderivative = 2.91

method	result
default	$\frac{2\left(-2\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)b+b\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}\sqrt{-2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}\right)}{b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\sin\left(\frac{dx}{2}+\frac{c}{2}\right)\sqrt{2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1}}$

[In] `int(cos(d*x+c)/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{-2/b^2(-2\cos(1/2*d*x+1/2*c))*(-2\sin(1/2*d*x+1/2*c))^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\sin(1/2*d*x+1/2*c)^2+(\sin(1/2*d*x+1/2*c)^2)^{(1/2)*(2*\sin(1/2*d*x+1/2*c)^2-1)^{(1/2)*(-2*\sin(1/2*d*x+1/2*c))^4*b+b*\sin(1/2*d*x+1/2*c)^2)^{(1/2)*\text{EllipticE}(\cos(1/2*d*x+1/2*c),2^{(1/2)})/(-b*(2*\sin(1/2*d*x+1/2*c))^4-\sin(1/2*d*x+1/2*c)^2)^{(1/2)/\sin(1/2*d*x+1/2*c)/((2*\cos(1/2*d*x+1/2*c)^2-1)*b)^{(1/2)/d}}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.53

$$\int \frac{\cos(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{-i\sqrt{2}\sqrt{b}\cos(dx+c)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(dx+c)))}{(b\cos(c+dx))^{5/2}}$$

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out]
$$(-I*\sqrt{2)*\sqrt{b)*\cos(d*x+c)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)+I*\sin(d*x+c)))+I*\sqrt{2)*\sqrt{b)*\cos(d*x+c)*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(d*x+c)-I*\sin(d*x+c)))+2*\sqrt{b*\cos(d*x+c)*\sin(d*x+c)}/(b^3*d*\cos(d*x+c))$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(cos(c + d*x)/(b*cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(5/2), x)

3.135 $\int \frac{1}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	653
Rubi [A] (verified)	653
Mathematica [A] (verified)	654
Maple [B] (verified)	654
Fricas [C] (verification not implemented)	655
Sympy [F]	655
Maxima [F]	656
Giac [F]	656
Mupad [F(-1)]	656

Optimal result

Integrand size = 12, antiderivative size = 72

$$\int \frac{1}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}}$$

[Out] $2/3*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(3/2)}+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2720}

$$\int \frac{1}{(b \cos(c+dx))^{5/2}} dx = \frac{2\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{3b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{3bd(b \cos(c+dx))^{3/2}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c+d*x])^{(-5/2)}, x]$

[Out] $(2*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(3*b^2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*\operatorname{Sin}[c+d*x])/(3*b*d*(b*\operatorname{Cos}[c+d*x])^{(3/2)})$

Rule 2716

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c+d*x]*((b*\operatorname{Sin}[c+d*x])^{(n+1)}/(b*d*(n+1))), x] + \operatorname{Dist}[(n+2)/(b^2*(n+1)), \operatorname{Int}[(b*\operatorname{Sin}[c+d*x])^{(n+2)}, x], x] /; \operatorname{FreeQ}\{b, c, d\}, x] \&\& \operatorname{LtQ}[n, -1] \&\& \operatorname{IntegerQ}[2*n]$

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \sin(c + dx)}{3bd(b \cos(c + dx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b \cos(c+dx)}} dx}{3b^2} \\ &= \frac{2 \sin(c + dx)}{3bd(b \cos(c + dx))^{3/2}} + \frac{\sqrt{\cos(c + dx)} \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^2 \sqrt{b \cos(c + dx)}} \\ &= \frac{2\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{3b^2 d \sqrt{b \cos(c + dx)}} + \frac{2 \sin(c + dx)}{3bd(b \cos(c + dx))^{3/2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.71

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \frac{2\left(\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right) + \tan(c + dx)\right)}{3b^2 d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(-5/2),x]
```

```
[Out] (2*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b^2*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(88) = 176.

Time = 1.66 (sec) , antiderivative size = 241, normalized size of antiderivative = 3.35

method	result
default	$-\frac{2\left(-2\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\cos\left(\frac{dx}{2}+\frac{c}{2}\right)+\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{3b^2\sqrt{-b\left(2\left(\sin^4\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)\right)}\left(2\left(\cos^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-1\right)\sin\left(\frac{dx}{2}+\frac{c}{2}\right)}$

[In] `int(1/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`

[Out] `-2/3*(-2*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2))*sin(1/2*d*x+1/2*c)^2-2*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c),2^(1/2)))/b^2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d`

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.43

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \frac{-i \sqrt{2} \sqrt{b} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) + i \sin(dx + c)) + i \sqrt{2} \sqrt{b} \cos(dx + c)^2 \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c)) + 2 \sqrt{b \cos(dx + c)} \sin(dx + c)}{(b^3 d \cos(dx + c))^2}$$

[In] `integrate(1/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out] `1/3*(-I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + I*sqrt(2)*sqrt(b)*cos(d*x + c)^2*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^2)`

Sympy [F]

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(c + dx))^{\frac{5}{2}}} dx$$

[In] `integrate(1/(b*cos(d*x+c))**(5/2),x)`

[Out] `Integral((b*cos(c + d*x))**(-5/2), x)`

Maxima [F]

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(-5/2), x)

Giac [F]

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(-5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{(b \cos(c + dx))^{5/2}} dx$$

[In] int(1/(b*cos(c + d*x))^(5/2),x)

[Out] int(1/(b*cos(c + d*x))^(5/2), x)

3.136 $\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	657
Rubi [A] (verified)	657
Mathematica [A] (verified)	659
Maple [B] (verified)	659
Fricas [C] (verification not implemented)	660
Sympy [F(-1)]	660
Maxima [F]	660
Giac [F]	661
Mupad [F(-1)]	661

Optimal result

Integrand size = 19, antiderivative size = 97

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx = -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5b^3 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $2/5*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+6/5*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}-6/5*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.211$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{5/2}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5b^3 d \sqrt{\cos(c+dx)}} + \frac{6 \sin(c+dx)}{5b^2 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{5d(b \cos(c+dx))^{5/2}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(-6*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(5*b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*\text{Sin}[c + d*x])/(5*d*(b*\text{Cos}[c + d*x])^{(5/2)}) + (6*\text{Sin}[c + d*x])/(5*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

```
Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2716

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c+d*x]*((b*Sin[c+d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c+d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c+d*x])^n/Sin[c+d*x]^n, Int[Sin[c+d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
&= \frac{2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{3 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx}{5b} \\
&= \frac{2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^2 d \sqrt{b \cos(c + dx)}} - \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b^3} \\
&= \frac{2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^2 d \sqrt{b \cos(c + dx)}} - \frac{\left(3 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b^3 \sqrt{\cos(c + dx)}} \\
&= -\frac{6 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5b^3 d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)}{5d(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^2 d \sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.70

$$\int \frac{\sec(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{-6\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + 6\sin(c+dx) + 2\sec(c+dx)\tan(c+dx)}{5b^2d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(5/2), x]

[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c + d*x]*Tan[c + d*x])/(5*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. 2(109) = 218.

Time = 2.78 (sec) , antiderivative size = 367, normalized size of antiderivative = 3.78

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{(24\cos(\frac{dx}{2} + \frac{c}{2})(\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12\sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1)\sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}}} E(\cos(\frac{dx}{2} + \frac{c}{2}))$

[In] int(sec(d*x+c)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out]
$$-\frac{2}{5} * (-(-2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 + 1) * b * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) / b ^ 3 / \sin(1/2 * d * x + 1/2 * c) ^ 3 / (8 * \sin(1/2 * d * x + 1/2 * c) ^ 6 - 12 * \sin(1/2 * d * x + 1/2 * c) ^ 4 + 6 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) * (24 * \cos(1/2 * d * x + 1/2 * c) * \sin(1/2 * d * x + 1/2 * c) ^ 6 - 12 * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2)) * \sin(1/2 * d * x + 1/2 * c) ^ 4 - 24 * \sin(1/2 * d * x + 1/2 * c) ^ 4 * \cos(1/2 * d * x + 1/2 * c) + 12 * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2)) * \sin(1/2 * d * x + 1/2 * c) ^ 2 + 8 * \sin(1/2 * d * x + 1/2 * c) ^ 2 * \cos(1/2 * d * x + 1/2 * c) - 3 * (\sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) * (2 * \sin(1/2 * d * x + 1/2 * c) ^ 2 - 1) ^ (1/2) * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2 ^ (1/2))) * (-2 * \sin(1/2 * d * x + 1/2 * c) ^ 4 * b + b * \sin(1/2 * d * x + 1/2 * c) ^ 2) ^ (1/2) / ((2 * \cos(1/2 * d * x + 1/2 * c) ^ 2 - 1) * b) ^ (1/2) / d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.25

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c)))}{(b \cos(c + dx))^{5/2}}$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/5*(-3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 3*I*sqrt(2)*sqrt(b)*cos(d*x + c)^3*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 1)*sin(d*x + c))/(b^3*d*cos(d*x + c)^3)

Sympy [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{5/2}} dx$$

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(5/2)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(5/2)), x)

$$3.137 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	662
Rubi [A] (verified)	662
Mathematica [A] (verified)	664
Maple [B] (verified)	664
Fricas [C] (verification not implemented)	665
Sympy [F(-1)]	665
Maxima [F]	665
Giac [F]	665
Mupad [F(-1)]	666

Optimal result

Integrand size = 21, antiderivative size = 98

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21b^2d\sqrt{b \cos(c+dx)}} + \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}} + \frac{10 \sin(c+dx)}{21bd(b \cos(c+dx))^{3/2}}$$

[Out] $2/7*b*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(7/2)}+10/21*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(3/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\operatorname{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2720}

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right)}{21b^2d\sqrt{b \cos(c+dx)}} + \frac{10 \sin(c+dx)}{21bd(b \cos(c+dx))^{3/2}} + \frac{2b \sin(c+dx)}{7d(b \cos(c+dx))^{7/2}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c+d*x]^2/(b*\operatorname{Cos}[c+d*x])^{(5/2)}, x]$

[Out] $(10*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{EllipticF}[(c+d*x)/2, 2])/(21*b^2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + (2*b*\operatorname{Sin}[c+d*x])/(7*d*(b*\operatorname{Cos}[c+d*x])^{(7/2)}) + (10*\operatorname{Sin}[c+d*x])/(21*b*d*(b*\operatorname{Cos}[c+d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2716

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1))), x] + Dist[(n+2)/(b^2*(n+1)), Int[(b*Sin[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{9/2}} dx \\
 &= \frac{2b \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{5}{7} \int \frac{1}{(b \cos(c + dx))^{5/2}} dx \\
 &= \frac{2b \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10 \sin(c + dx)}{21bd(b \cos(c + dx))^{3/2}} + \frac{5 \int \frac{1}{\sqrt{b \cos(c + dx)}} dx}{21b^2} \\
 &= \frac{2b \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10 \sin(c + dx)}{21bd(b \cos(c + dx))^{3/2}} + \frac{\left(5\sqrt{\cos(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21b^2 \sqrt{b \cos(c + dx)}} \\
 &= \frac{10\sqrt{\cos(c + dx)} \text{EllipticF}\left(\frac{1}{2}(c + dx), 2\right)}{21b^2 d \sqrt{b \cos(c + dx)}} + \frac{2b \sin(c + dx)}{7d(b \cos(c + dx))^{7/2}} + \frac{10 \sin(c + dx)}{21bd(b \cos(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.67

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{10\sqrt{\cos(c+dx)} \operatorname{EllipticF}\left(\frac{1}{2}(c+dx), 2\right) + 2(5+3\sec^2(c+dx)) \tan(c+dx)}{21b^2 d \sqrt{b \cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(5/2), x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 2*(5 + 3*Sec[c + d*x]^2)*Tan[c + d*x])/(21*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(110) = 220.

Time = 2.23 (sec) , antiderivative size = 398, normalized size of antiderivative = 4.06

method	result
default	$-\frac{2\left(-40\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\sqrt{2\left(\sin^2\left(\frac{dx}{2}+\frac{c}{2}\right)\right)}-1F\left(\cos\left(\frac{dx}{2}+\frac{c}{2}\right),\sqrt{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)-40\cos\left(\frac{dx}{2}+\frac{c}{2}\right)\left(\sin^6\left(\frac{dx}{2}+\frac{c}{2}\right)\right)+60\sqrt{\frac{1}{2}-\frac{\cos(dx+c)}{2}}\right)}{\dots}$

[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/21*(-40*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^6-40*cos(1/2*d*x+1/2*c)*sin(1/2*d*x+1/2*c)^6+60*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^4+40*sin(1/2*d*x+1/2*c)^4*cos(1/2*d*x+1/2*c)-30*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))*sin(1/2*d*x+1/2*c)^2-16*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)+5*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(2*sin(1/2*d*x+1/2*c)^2-1)^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2)))/b^2*((2*cos(1/2*d*x+1/2*c)^2-1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(2*cos(1/2*d*x+1/2*c)^2-1)^3/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.17

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{-5i \sqrt{2} \sqrt{b} \cos(dx+c)^4 \text{weierstrassPInverse}(-4, 0, \cos(dx+c) + i \sin(dx+c))}{(b \cos(c+dx))^{5/2}}$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/21*(-5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c)) + 5*I*sqrt(2)*sqrt(b)*cos(d*x + c)^4*weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c)) + 2*sqrt(b*cos(d*x + c))*(5*cos(d*x + c)^2 + 3)*sin(d*x + c))/(b^3*d*cos(d*x + c)^4)

Sympy [F(-1)]

Timed out.

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \int \frac{\sec(dx+c)^2}{(b \cos(dx+c))^{\frac{5}{2}}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{5/2}} dx$$

```
[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2)),x)
```

```
[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(5/2)), x)
```

3.138 $\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx$

Optimal result	667
Rubi [A] (verified)	667
Mathematica [A] (verified)	669
Maple [B] (verified)	669
Fricas [C] (verification not implemented)	670
Sympy [F(-1)]	670
Maxima [F]	670
Giac [F]	671
Mupad [F(-1)]	671

Optimal result

Integrand size = 21, antiderivative size = 125

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx = -\frac{14\sqrt{b \cos(c+dx)}E\left(\frac{1}{2}(c+dx) \mid 2\right)}{15b^3d\sqrt{\cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}} + \frac{14 \sin(c+dx)}{15b^2d\sqrt{b \cos(c+dx)}}$$

[Out] $2/9*b^2*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(9/2)}+14/45*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/2)}+14/15*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}-14/15*(\cos(1/2*d*x+1/2*c))^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*(b*\cos(d*x+c))^{(1/2)}/b^3/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 2716, 2721, 2719}

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{5/2}} dx = -\frac{14E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{15b^3d\sqrt{\cos(c+dx)}} + \frac{2b^2 \sin(c+dx)}{9d(b \cos(c+dx))^{9/2}} + \frac{14 \sin(c+dx)}{15b^2d\sqrt{b \cos(c+dx)}} + \frac{14 \sin(c+dx)}{45d(b \cos(c+dx))^{5/2}}$$

[In] $\text{Int}[\text{Sec}[c + d*x]^3/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(-14*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{EllipticE}[(c + d*x)/2, 2])/(15*b^3*d*\text{Sqrt}[\text{Cos}[c + d*x]]) + (2*b^2*\text{Sin}[c + d*x])/(9*d*(b*\text{Cos}[c + d*x])^{(9/2)}) + (14*\text{Sin}[c + d$

$*x]/(45*d*(b*\text{Cos}[c + d*x])^(5/2)) + (14*\text{Sin}[c + d*x])/(15*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] := \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x\} \&\& \text{IntegerQ}[m]$

Rule 2716

$\text{Int}[(b_)*\text{sin}[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] := \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n+1)}/(b*d*(n+1))), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x\} \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_)+(d_)*(x_)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x\}$

Rule 2721

$\text{Int}[(b_)*\text{sin}[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] := \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x\} \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{11/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{1}{9}(7b) \int \frac{1}{(b \cos(c + dx))^{7/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{7 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx}{15b} \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} + \frac{14 \sin(c + dx)}{15b^2 d \sqrt{b \cos(c + dx)}} - \frac{7 \int \sqrt{b \cos(c + dx)} dx}{15b^3} \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \cos(c + dx))^{9/2}} + \frac{14 \sin(c + dx)}{45d(b \cos(c + dx))^{5/2}} \\
 &\quad + \frac{14 \sin(c + dx)}{15b^2 d \sqrt{b \cos(c + dx)}} - \frac{\left(7 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{15b^3 \sqrt{\cos(c + dx)}}
 \end{aligned}$$

$$= -\frac{14\sqrt{b\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15b^3d\sqrt{\cos(c+dx)}} + \frac{2b^2\sin(c+dx)}{9d(b\cos(c+dx))^{9/2}}$$

$$+ \frac{14\sin(c+dx)}{45d(b\cos(c+dx))^{5/2}} + \frac{14\sin(c+dx)}{15b^2d\sqrt{b\cos(c+dx)}}$$

Mathematica [A] (verified)

Time = 0.39 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.64

$$\int \frac{\sec^3(c+dx)}{(b\cos(c+dx))^{5/2}} dx = \frac{-42\sqrt{\cos(c+dx)}E\left(\frac{1}{2}(c+dx)\middle|2\right) + 42\sin(c+dx) + 2\sec(c+dx)(7+5\sec^2(c+dx))}{45b^2d\sqrt{b\cos(c+dx)}}$$

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(5/2), x]

[Out] (-42*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 42*Sin[c + d*x] + 2*Sec[c + d*x]*(7 + 5*Sec[c + d*x]^2)*Tan[c + d*x])/(45*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 415 vs. 2(133) = 266.

Time = 3.17 (sec) , antiderivative size = 416, normalized size of antiderivative = 3.33

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2})) + 1)b(\sin^2(\frac{dx}{2} + \frac{c}{2}))}}{144b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5} \left(-\frac{\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{144b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5} - \frac{7\cos(\frac{dx}{2} + \frac{c}{2})\sqrt{-b(2(\sin^4(\frac{dx}{2} + \frac{c}{2})) - (\sin^2(\frac{dx}{2} + \frac{c}{2})))}}{180b(\cos^2(\frac{dx}{2} + \frac{c}{2}) - \frac{1}{2})^5} \right)$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2*(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)/b^2*(-1/144*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^5-7/180*cos(1/2*d*x+1/2*c)/b*(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)/(cos(1/2*d*x+1/2*c)^2-1/2)^3-14/15*sin(1/2*d*x+1/2*c)^2*cos(1/2*d*x+1/2*c)/(-(-2*cos(1/2*d*x+1/2*c)^2+1)*b*sin(1/2*d*x+1/2*c)^2)^(1/2)+7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-7/15*(sin(1/2*d*x+1/2*c)^2)^(1/2)*(-2*cos(1/2*d*x+1/2*c)^2+1)^(1/2)/(-b*(2*sin(1/2*d*x+1/2*c)^4-sin(1/2*d*x+1/2*c)^2))^(1/2)*(EllipticF(cos(1/2*d*x+1/2*c), 2^(1/2))-EllipticE(cos(1/2*d*x+1/2*c), 2^(1/2)))/sin(1/2*d*x+1/2*c)/((2*cos(1/2*d*x+1/2*c)^2-1)*b)^(1/2)/d

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.11 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.05

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{-21i \sqrt{2} \sqrt{b} \cos(dx + c)^5 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c))) + 21i \sqrt{2} \sqrt{b} \cos(dx + c)^5 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c) - i \sin(dx + c))) + 2 * (21 \cos(dx + c)^4 + 7 \cos(dx + c)^2 + 5) \sqrt{b \cos(dx + c)} \sin(dx + c)}{(b^3 d \cos(dx + c))^5}$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/45*(-21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) + I*sin(d*x + c))) + 21*I*sqrt(2)*sqrt(b)*cos(d*x + c)^5*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(d*x + c) - I*sin(d*x + c))) + 2*(21*cos(d*x + c)^4 + 7*cos(d*x + c)^2 + 5)*sqrt(b*cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^5)

Sympy [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

Giac [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{5/2}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{5/2}} dx$$

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(5/2)), x)

$$3.139 \quad \int \frac{1}{(b \cos(c+dx))^{7/2}} dx$$

Optimal result	672
Rubi [A] (verified)	672
Mathematica [A] (verified)	673
Maple [B] (verified)	674
Fricas [C] (verification not implemented)	674
Sympy [F(-1)]	675
Maxima [F]	675
Giac [F]	675
Mupad [F(-1)]	675

Optimal result

Integrand size = 12, antiderivative size = 100

$$\int \frac{1}{(b \cos(c+dx))^{7/2}} dx = -\frac{6\sqrt{b \cos(c+dx)} E\left(\frac{1}{2}(c+dx) \mid 2\right)}{5b^4 d \sqrt{\cos(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \cos(c+dx))^{5/2}} + \frac{6 \sin(c+dx)}{5b^3 d \sqrt{b \cos(c+dx)}}$$

[Out] 2/5*sin(d*x+c)/b/d/(b*cos(d*x+c))^(5/2)+6/5*sin(d*x+c)/b^3/d/(b*cos(d*x+c))^(1/2)-6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))*(b*cos(d*x+c))^(1/2)/b^4/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$\int \frac{1}{(b \cos(c+dx))^{7/2}} dx = -\frac{6E\left(\frac{1}{2}(c+dx) \mid 2\right) \sqrt{b \cos(c+dx)}}{5b^4 d \sqrt{\cos(c+dx)}} + \frac{6 \sin(c+dx)}{5b^3 d \sqrt{b \cos(c+dx)}} + \frac{2 \sin(c+dx)}{5bd(b \cos(c+dx))^{5/2}}$$

[In] Int[(b*Cos[c + d*x])^(-7/2),x]

[Out] (-6*Sqrt[b*Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2])/(5*b^4*d*Sqrt[Cos[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Cos[c + d*x])^(5/2)) + (6*Sin[c + d*x])/(5*b^3*d*Sqrt[b*Cos[c + d*x]])

Rule 2716


```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), In
t[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] &&
IntegerQ[2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{2 \sin(c + dx)}{5bd(b \cos(c + dx))^{5/2}} + \frac{3 \int \frac{1}{(b \cos(c + dx))^{3/2}} dx}{5b^2} \\
&= \frac{2 \sin(c + dx)}{5bd(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^3 d \sqrt{b \cos(c + dx)}} - \frac{3 \int \sqrt{b \cos(c + dx)} dx}{5b^4} \\
&= \frac{2 \sin(c + dx)}{5bd(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^3 d \sqrt{b \cos(c + dx)}} - \frac{\left(3 \sqrt{b \cos(c + dx)}\right) \int \sqrt{\cos(c + dx)} dx}{5b^4 \sqrt{\cos(c + dx)}} \\
&= -\frac{6 \sqrt{b \cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right)}{5b^4 d \sqrt{\cos(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \cos(c + dx))^{5/2}} + \frac{6 \sin(c + dx)}{5b^3 d \sqrt{b \cos(c + dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.68

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \frac{-6 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \mid 2\right) + 6 \sin(c + dx) + 2 \sec(c + dx) \tan(c + dx)}{5b^3 d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^(-7/2),x]
```

```
[Out] (-6*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + 6*Sin[c + d*x] + 2*Sec[c
+ d*x]*Tan[c + d*x])/(5*b^3*d*Sqrt[b*Cos[c + d*x]])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 366 vs. $2(112) = 224$.

Time = 2.59 (sec) , antiderivative size = 367, normalized size of antiderivative = 3.67

method	result
default	$-\frac{2\sqrt{-(-2(\cos^2(\frac{dx}{2} + \frac{c}{2}))) + 1} b(\sin^2(\frac{dx}{2} + \frac{c}{2})) \left(24 \cos(\frac{dx}{2} + \frac{c}{2}) (\sin^6(\frac{dx}{2} + \frac{c}{2})) - 12 \sqrt{2(\sin^2(\frac{dx}{2} + \frac{c}{2}))} - 1 \sqrt{\frac{1}{2} - \frac{\cos(dx+c)}{2}} E(\cos(\frac{dx}{2} + \frac{c}{2})) \right)}{\dots}$

[In] `int(1/(cos(d*x+c)*b)^(7/2),x,method=_RETURNVERBOSE)`

[Out]
$$-2/5 * (-(-2 * \cos(1/2 * d * x + 1/2 * c)^2 + 1) * b * \sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} / b^4 / \sin(1/2 * d * x + 1/2 * c)^3 / (8 * \sin(1/2 * d * x + 1/2 * c)^6 - 12 * \sin(1/2 * d * x + 1/2 * c)^4 + 6 * \sin(1/2 * d * x + 1/2 * c)^2 - 1) * (24 * \cos(1/2 * d * x + 1/2 * c) * \sin(1/2 * d * x + 1/2 * c)^6 - 12 * (2 * \sin(1/2 * d * x + 1/2 * c)^2 - 1)^{(1/2)} * (\sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2^{(1/2)}) * \sin(1/2 * d * x + 1/2 * c)^4 - 24 * \sin(1/2 * d * x + 1/2 * c)^4 * \cos(1/2 * d * x + 1/2 * c) + 12 * (2 * \sin(1/2 * d * x + 1/2 * c)^2 - 1)^{(1/2)} * (\sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2^{(1/2)}) * \sin(1/2 * d * x + 1/2 * c)^2 + 8 * \sin(1/2 * d * x + 1/2 * c)^2 * \cos(1/2 * d * x + 1/2 * c) - 3 * (\sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} * (2 * \sin(1/2 * d * x + 1/2 * c)^2 - 1)^{(1/2)} * \text{EllipticE}(\cos(1/2 * d * x + 1/2 * c), 2^{(1/2)}) * (-2 * \sin(1/2 * d * x + 1/2 * c)^4 * b + b * \sin(1/2 * d * x + 1/2 * c)^2)^{(1/2)} / ((2 * \cos(1/2 * d * x + 1/2 * c)^2 - 1) * b)^{(1/2)} / d$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.21

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \frac{-3i \sqrt{2} \sqrt{b} \cos(dx + c)^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c)))}{\dots}$$

[In] `integrate(1/(b*cos(d*x+c))^(7/2),x, algorithm="fricas")`

[Out]
$$1/5 * (-3 * I * \sqrt{2} * \sqrt{b} * \cos(dx + c)^3 * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c))) + I * \sin(dx + c)) + 3 * I * \sqrt{2} * \sqrt{b} * \cos(dx + c)^3 * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(dx + c))) - I * \sin(dx + c)) + 2 * \sqrt{b} * \cos(dx + c) * (3 * \cos(dx + c)^2 + 1) * \sin(dx + c) / (b^4 * d * \cos(dx + c)^3)$$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \text{Timed out}$$

[In] integrate(1/(b*cos(d*x+c))**(7/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \int \frac{1}{(b \cos(dx + c))^{7/2}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

Giac [F]

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \int \frac{1}{(b \cos(dx + c))^{7/2}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(7/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{7/2}} dx = \int \frac{1}{(b \cos(c + dx))^{7/2}} dx$$

[In] int(1/(b*cos(c + d*x))^(7/2),x)

[Out] int(1/(b*cos(c + d*x))^(7/2), x)

3.140 $\int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	676
Rubi [A] (verified)	676
Mathematica [A] (verified)	678
Maple [A] (verified)	678
Fricas [A] (verification not implemented)	678
Sympy [F(-1)]	679
Maxima [A] (verification not implemented)	679
Giac [B] (verification not implemented)	679
Mupad [B] (verification not implemented)	680

Optimal result

Integrand size = 23, antiderivative size = 98

$$\int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{3x \sqrt{b \cos(c + dx)}}{8 \sqrt{\cos(c + dx)}} + \frac{3 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d}$$

[Out] $1/4*\cos(d*x+c)^{(5/2)}*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d+3/8*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+3/8*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{3x \sqrt{b \cos(c + dx)}}{8 \sqrt{\cos(c + dx)}} + \frac{\sin(c + dx) \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)}}{4d} + \frac{3 \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}{8d}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(3*x*\sqrt{b*\cos[c + d*x]})/(8*\sqrt{\cos[c + d*x]}) + (3*\sqrt{\cos[c + d*x]}*\sqrt{b*\cos[c + d*x]}*\sin[c + d*x])/(8*d) + (\cos[c + d*x]^{5/2}*\sqrt{b*\cos[c + d*x]}*\sin[c + d*x])/(4*d)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \cos^4(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
 &= \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \cos^2(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\
 &= \frac{3\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} \\
 &\quad + \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int 1 dx}{8\sqrt{\cos(c + dx)}} \\
 &= \frac{3x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{3\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} \\
 &\quad + \frac{\cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.56

$$\int \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx$$

$$= \frac{\sqrt{b \cos(c+dx)} (12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32d \sqrt{\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

method	result
default	$\frac{\sqrt{\cos(dx+c)b} (2 \sin(dx+c) (\cos^3(dx+c) + 3 \cos(dx+c) \sin(dx+c) + 3dx + 3c))}{8d \sqrt{\cos(dx+c)}}$
risch	$\frac{3 \sqrt{\cos(dx+c)b} (\sqrt{\cos(dx+c)}) e^{i(dx+c)} x}{4(e^{2i(dx+c)} + 1)} - \frac{i \sqrt{\cos(dx+c)b} (\sqrt{\cos(dx+c)}) e^{5i(dx+c)}}{32(e^{2i(dx+c)} + 1)d} + \frac{i \sqrt{\cos(dx+c)b} (\sqrt{\cos(dx+c)}) e^{-i(dx+c)}}{4(e^{2i(dx+c)} + 1)d} - \frac{7i \sqrt{\cos(dx+c)b} (\sqrt{\cos(dx+c)}) e^{-3i(dx+c)}}{32(e^{2i(dx+c)} + 1)d}$

[In] int(cos(d*x+c)^(7/2)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/8/d*(cos(d*x+c)*b)^(1/2)*(2*sin(d*x+c)*cos(d*x+c)^3+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.80

$$\int \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx$$

$$= \left[\frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) + 3 \sqrt{-b} \log(2b \cos(dx+c)^2 - 2 \sqrt{b \cos(dx+c)})}{16d} \right]$$

[In] integrate(cos(d*x+c)^(7/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/16*(2*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 3)*sqrt(cos(d*x + c))*sin(d*x + c) + 3*sqrt(-b)*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt

$(-b)\sqrt{\cos(dx + c)}\sin(dx + c) - b)/d, 1/8(\sqrt{b\cos(dx + c)}(2\cos(dx + c)^2 + 3)\sqrt{\cos(dx + c)}\sin(dx + c) + 3\sqrt{b}\arctan(\sqrt{b\cos(dx + c)}\sin(dx + c)/(\sqrt{b}\cos(dx + c)^{3/2}))))/d]$

Sympy [F(-1)]

Timed out.

$$\int \cos^{7/2}(c + dx)\sqrt{b\cos(c + dx)} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(7/2)*(b*cos(d*x+c))**(1/2), x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.50

$$\int \cos^{7/2}(c + dx)\sqrt{b\cos(c + dx)} dx$$

$$= \frac{(12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c))))\sqrt{b}}{32 d}$$

[In] integrate(cos(d*x+c)^(7/2)*(b*cos(d*x+c))^(1/2), x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(80) = 160.

Time = 2.08 (sec) , antiderivative size = 203, normalized size of antiderivative = 2.07

$$\int \cos^{7/2}(c + dx)\sqrt{b\cos(c + dx)} dx$$

$$= \frac{3\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 12\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 - 10\sqrt{b} \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 18\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 6\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 3\sqrt{b}dx + 10\sqrt{b}\tan(\frac{1}{2}dx + \frac{1}{2}c)}{8(d \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 4d \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 + d)}$$

[In] integrate(cos(d*x+c)^(7/2)*(b*cos(d*x+c))^(1/2), x, algorithm="giac")

[Out] 1/8*(3*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^8 + 12*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^6 - 10*sqrt(b)*tan(1/2*d*x + 1/2*c)^7 + 18*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^4 + 6*sqrt(b)*tan(1/2*d*x + 1/2*c)^5 + 12*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^2 - 6*sqrt(b)*tan(1/2*d*x + 1/2*c)^3 + 3*sqrt(b)*d*x + 10*sqrt(b)*tan(1/2*d*x + 1/2*c))/(d*tan(1/2*d*x + 1/2*c)^8 + 4*d*tan(1/2*d*x + 1/2*c)^6 + 6*d*tan(1/2*d*x + 1/2*c)^4 + 4*d*tan(1/2*d*x + 1/2*c)^2 + d)

Mupad [B] (verification not implemented)

Time = 14.99 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.77

$$\int \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24dx \cos(c + dx))}{32d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x) + 1))

3.141 $\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	681
Rubi [A] (verified)	681
Mathematica [A] (verified)	682
Maple [A] (verified)	682
Fricas [A] (verification not implemented)	683
Sympy [F(-1)]	683
Maxima [A] (verification not implemented)	683
Giac [B] (verification not implemented)	684
Mupad [B] (verification not implemented)	731

Optimal result

Integrand size = 23, antiderivative size = 70

$$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx = \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}} - \frac{\sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \sqrt{\cos(c+dx)}}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}-1/3*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)} dx = \frac{\sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} - \frac{\sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) - (\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] := \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2713

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \cos^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\sqrt{b \cos(c + dx)} \text{Subst}(\int (1 - x^2) dx, x, -\sin(c + dx))}{d\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{\sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} (5 + \cos(2(c + dx))) \sin(c + dx)}{6d\sqrt{\cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (Sqrt[b*Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Sqrt[Cos[c + d*x]])
```

Maple [A] (verified)

Time = 3.17 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.57

method	result
default	$\frac{(2 + \cos^2(dx+c)) \sin(dx+c) \sqrt{\cos(dx+c)b}}{3d\sqrt{\cos(dx+c)}}$
risch	$-\frac{i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{4i(dx+c)}}{12(e^{2i(dx+c)}+1)d} - \frac{3i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{2i(dx+c)}}{4(e^{2i(dx+c)}+1)d} + \frac{3i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})}{4(e^{2i(dx+c)}+1)d} + \frac{i\sqrt{\cos(dx+c)b}}{4(e^{2i(dx+c)}+1)d}$

```
[In] int(cos(d*x+c)^(5/2)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.56

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3 d \sqrt{\cos(dx + c)}}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.46 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b} (\sin(3 dx + 3 c) + 9 \sin(\frac{1}{3} \arctan(\sin(3 dx + 3 c), \cos(3 dx + 3 c))))}{12 d}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/12*sqrt(b)*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71047 vs. 2(60) = 120.

Time = 9.09 (sec) , antiderivative size = 71047, normalized size of antiderivative = 1014.96

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \text{Too large to display}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] -1/96*(3*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c)^2 - 3*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6 - 24*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^6*tan(c) - 24*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c) + 9*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^4*tan(c)^2 - 18*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^6*tan(c)^2 - 48*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^6*tan(c)^2 + 9*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c)^2 - 18*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c)^2 - 9*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^4 + 18*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^6 + 48*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^6 - 9*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6 + 18*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6 - 72*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^4*tan(c) - 72*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^4*tan(c) + 24*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)*tan(1/3*c)^6*tan(c) + 144*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^6*tan(c) - 72*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^6*tan(c) + 144*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^6*tan(c) - 72*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c) + 24*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^6*tan(c) + 9*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^2*tan(c)^2 - 54*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^6*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^4*tan(c)^2 - 144*sqrt(b)*d*x^4*tan(1/2*d*x + 1/2*c

$$\begin{aligned}
&)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 \tan(c)^2 + \\
&27*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x \\
&+ 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 - 54*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \\
&* \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 + 3*s \\
&qrt(b)*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan \\
&(c)^2 + 48*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
&-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan(c)^2 - 54*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/ \\
&2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 \\
&+ 108*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2 \\
&*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 144*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2 \\
&*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 \\
&+ 48*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x \\
&+ 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 9*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 * \\
&\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 54*s \\
&qrt(b)*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1 \\
&/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 3*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
&-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 9*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1 \\
&/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 54*\sqrt{b} \\
& *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^2 \tan(1/3*c)^4 + 144*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x \\
&+ 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 27*\sqrt{b}*d*x^4 \tan(1/2* \\
&d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \\
&+ 54*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d \\
&*x + 1/2*c)^4 \tan(1/3*c)^4 - 3*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2 \\
&*d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(\\
&1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4 \tan \\
&(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3 \\
&*c)^6 - 108*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan \\
&(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 144*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^ \\
&3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 - 48*\sqrt{b}* \\
&d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan \\
&(1/3*c)^6 - 9*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \\
&* \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2* \\
&c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 3*\sqrt{b} \\
&)*d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 72*\sqrt{b} \\
& *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^3 \tan(1/3*c)^2 \tan(c) - 72*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/ \\
&2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 \tan(c) + 72*\sqrt{b}*d \\
&*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan \\
&(1/3*c)^4 \tan(c) + 432*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
&1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c) - 216*\sqrt{b}*d*x^4 \tan \\
&(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3 \\
&*c)^4 \tan(c) + 432*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c \\
&)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 \tan(c) - 216*\sqrt{b}*d*x^4 \tan(1/2 \\
&*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^6*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 144* \\
& \sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 3*\sqrt{b}*d*x^ \\
& 4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 54 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - \\
& 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^3*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 2 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c \\
&)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 324*\sqrt{b}*d*x \\
& ^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 27*\sqrt{b} \\
& (b)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 96*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*\sqrt{b} \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 18 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 - \\
& 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 \\
& - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*\sqrt{b}*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan \\
& (1/3*c)^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-
\end{aligned}$$

$$\begin{aligned}
& 1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 9*\sqrt{b}*d \\
& *x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b} \\
& (b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& n(1/3*c)^6 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*\sqrt{b}*d*x^4*\tan \\
& n(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) \\
& + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{ \\
& rt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^3*\tan(1/3*c)^2*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan \\
& n(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b} \\
& (b)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)*\tan(1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 1296*\sqrt{b}*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6* \\
& c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^ \\
& 4*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4 \\
& *\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c \\
&) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3* \\
& c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c) \\
& ^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan \\
& (1/3*c)^6*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c) - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^3*\tan(1/3*c)^6*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& *x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 72*\sqrt{b}*d*x^
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)
\end{aligned}$$

$$\begin{aligned}
&)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 + \\
&192 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^2 \tan(1/3*c)^5 \tan(c)^2 - 1440 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2* \\
&d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 \tan(c)^2 + 1728 \sqrt{b} \\
&* \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(\\
&1/3*c)^5 \tan(c)^2 - 384 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \\
&^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 \tan(c)^2 + 9 \sqrt{b} * d*x^4 \tan(1/2* \\
&d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 54 \sqrt{b} * d* \\
&x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 3 \\
&* \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan(c)^2 + 144 \sqrt{b} * d \\
&* x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) * \tan \\
&(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + \\
&1/6*c)^4 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan(c)^2 - 24 \sqrt{b} * \tan(1/2*d \\
&* x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan \\
&(c)^2 - 18 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan \\
&(1/3*c)^6 \tan(c)^2 + 324 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + \\
&1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 54 \sqrt{b} * d*x^4 * \\
&\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192 * \\
&\sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c) \\
&^2 \tan(1/3*c)^6 \tan(c)^2 - 288 \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
&1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} * d*x^4 * \\
&\tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 144 * \\
&\sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/ \\
&2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 72 \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d* \\
&x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} * \tan \\
&(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3* \\
&c)^6 \tan(c)^2 - 18 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2* \\
&c)^4 \tan(1/3*c)^6 \tan(c)^2 + 9 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/ \\
&2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 320 \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 \\
&* \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 432 \\
&* \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c \\
&)^4 \tan(1/3*c)^6 \tan(c)^2 - 384 \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x \\
&+ 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} * \tan(1 \\
&/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \\
&* \tan(c)^2 + 18 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 * \\
&\tan(-1/2*d*x + 1/2*c)^2 + 48 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d \\
&* x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 - 9 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c \\
&)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 + 18 \sqrt{b} * d*x^4 \tan(1 \\
&/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 - 9 \sqrt{b} (b \\
&)* d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^2 - 144 \sqrt{b} \\
& * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c) * \tan(1/3*c)^2 + 162 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + \\
&1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 - 324 \sqrt{b} * d*x^4 \tan(1/2*d \\
&* x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + \\
&432 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d
\end{aligned}$$

$$\begin{aligned}
& *x + 1/2*c)^3*\tan(1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*ta \\
& n(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 27*\sqrt{b}*d*x^ \\
& 4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 54*\sqrt{b}*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 - 432*\sqrt{ \\
& b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)*\tan(1/3*c)^4 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 972*sq \\
& rt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c) \\
& ^4 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^3*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 27*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1440*\sqrt{b}*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^4 + 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1440*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + \\
& 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^5 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 3*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^6 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c) \\
& ^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^6 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*ta \\
& n(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6 - 288*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan
\end{aligned}$$

$$\begin{aligned}
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9*\sqrt{b} \\
&)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 640*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 96*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
& 54\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^4\tan(-1/2dx \\
& x + 1/2c)^2 - 108\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c \\
&)^6\tan(-1/2dx + 1/2c)^2 + 144\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^3\tan(\\
& 1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)^3 - 48\sqrt{b}d^4x^4\tan(1/2dx + \\
& 1/2c)\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^3 - 9\sqrt{b}d^4x^4\tan \\
& n(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^4 + 54\sqrt{b} \\
& rt(b)d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/ \\
& 2c)^4 - 3\sqrt{b}d^4x^4\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^4 + 9 \\
& 6\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2 \\
& c)^4\tan(1/3c) - 27\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6 \\
& c)^4\tan(1/3c)^2 + 54\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + \\
& 1/6c)^6\tan(1/3c)^2 - 432\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^3\tan(1/2dx \\
& x + 1/6c)^4\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 144\sqrt{b}d^4x^4\tan(1/2 \\
& *dx + 1/2c)\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 1 \\
& 62\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^2\tan(-1/2dx \\
& + 1/2c)^2\tan(1/3c)^2 - 972\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2 \\
& *dx + 1/6c)^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 54\sqrt{b}d^4x^4\tan \\
& (1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 432\sqrt{b}d^4x^ \\
& 4\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^3\tan \\
& (1/3c)^2 - 432\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^4\tan \\
& an(-1/2dx + 1/2c)^3\tan(1/3c)^2 + 72\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan \\
& (1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^3\tan(1/3c)^2 - 9\sqrt{b}d^4x^4 \\
& tan(1/2dx + 1/2c)^4\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 + 162\sqrt{b}d \\
& *x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^4 \\
& tan(1/3c)^2 - 27\sqrt{b}d^4x^4\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c \\
&)^4\tan(1/3c)^2 + 1440\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c) \\
& ^5\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 - 1008\sqrt{b}d^4x^4\tan(1/2dx + 1/2c) \\
& ^3\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 - 640\sqrt{b} \\
&)d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^2\tan \\
& (1/3c)^3 + 4800\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^4\tan(\\
& -1/2dx + 1/2c)^4\tan(1/3c)^3 - 5760\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^3\tan(\\
& 1/2dx + 1/6c)^5\tan(-1/2dx + 1/2c)^4\tan(1/3c)^3 + 1280\sqrt{b}d^4x^4 \\
& tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)^4\tan(1/3c \\
&)^3 - 27\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^2\tan(1/ \\
& 3c)^4 + 162\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^4\tan \\
& n(1/3c)^4 - 9\sqrt{b}d^4x^4\tan(1/2dx + 1/6c)^6\tan(1/3c)^4 - 432\sqrt{b} \\
& (b)d^4x^4\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2 \\
& c)\tan(1/3c)^4 + 432\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6 \\
& c)^4\tan(-1/2dx + 1/2c)\tan(1/3c)^4 + 72\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4 \\
& *tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)\tan(1/3c)^4 + 54\sqrt{b}d^4x^ \\
& ^4\tan(1/2dx + 1/2c)^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^4 - 972\sqrt{b} \\
& (b)d^4x^4\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c) \\
& ^2\tan(1/3c)^4 + 162\sqrt{b}d^4x^4\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1 \\
& /2c)^2\tan(1/3c)^4 - 2880\sqrt{b}d^4x^4\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/ \\
& 6c)^5\tan(-1/2dx + 1/2c)^2\tan(1/3c)^4 + 864\sqrt{b}d^4x^4\tan(1/2dx + 1/2
\end{aligned}$$

$$\begin{aligned}
& *c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 144*\sqrt{b} \\
& t(b)*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 43 \\
& 2*\sqrt{b}*d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + \\
& 1/2*c)^3 \tan(1/3*c)^4 + 216*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/ \\
& 6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 144*\sqrt{b} \tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 54*\sqrt{b} \\
& (b)*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 27* \\
& \sqrt{b}*d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + \\
& 4800*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + \\
& 1/2*c)^4 \tan(1/3*c)^4 - 11664*\sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
& 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 5760*\sqrt{b} \tan(1/2*d*x + \\
& 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 1008 \\
& *\sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^ \\
& 4 \tan(1/3*c)^4 + 96*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan \\
& an(1/3*c)^5 - 2880*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan \\
& (-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 + 3456*\sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan \\
& n(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 - 768*\sqrt{b} \tan \\
& (1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3* \\
& c)^5 + 1440*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2* \\
& d*x + 1/2*c)^4 \tan(1/3*c)^5 - 5760*\sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d \\
& *x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 + 5760*\sqrt{b} \tan(1/2*d \\
& *x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 - \\
& 1728*\sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/ \\
& 2*c)^4 \tan(1/3*c)^5 + 96*\sqrt{b} \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2* \\
& c)^4 \tan(1/3*c)^5 - 3*\sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 5 \\
& 4*\sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 \\
& - 9*\sqrt{b} *d*x^4 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 + 96*\sqrt{b} \tan(1/2* \\
& d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(1/3*c)^6 - 144*\sqrt{b} \tan(1/2*d* \\
& x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48*\sqrt{b} *d*x^4 \tan(1/2 \\
& *d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 + 144*\sqrt{b} *d*x^4 \tan(\\
& 1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \\
& + 72*\sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1 \\
& /2*c) \tan(1/3*c)^6 + 48*\sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \\
& ^6 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 - 108*\sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 54*\sqrt{b} *d*x^4 \tan(1/2*d*x + \\
& 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 640*\sqrt{b} \tan(1/2*d*x + \\
& 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 864* \\
& \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) \\
& ^2 \tan(1/3*c)^6 - 768*\sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \\
& * \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 96*\sqrt{b} \tan(1/2*d*x + 1/2*c) \tan \\
& (1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 48*\sqrt{b} *d*x^4 \\
& * \tan(1/2*d*x + 1/2*c) \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 + 72*\sqrt{b} \tan \\
& (1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3* \\
& c)^6 + 144*\sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d \\
& *x + 1/2*c)^3 \tan(1/3*c)^6 + 24*\sqrt{b} \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/3*c)^6 - 3*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) - 1296 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) - 216 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 216 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) - 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^6 * \tan(c) - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^6 * \tan(c) - 24 * \sqrt{b} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) + 24 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 9 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(c)^2 - 18 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(c)^2 + 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 324 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 18 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 24 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 3 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 9 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 192 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) * \tan(c)^2 - 1440 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 + 1728 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 - 384 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(
\end{aligned}$$

$$\begin{aligned}
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 27*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c \\
&)^2*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c \\
&)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*s \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)* \\
& \tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 1 \\
& 62*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2 \\
& *c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3* \\
& c)^2*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(\\
& c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*t \\
& \tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*t \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 \\
& - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 4800*s \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*t \\
& \tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^5*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^5*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*
\end{aligned}$$

$$\begin{aligned}
& c)^6 \tan(c)^2 - 320 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 432 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 384 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 18 \sqrt{b} *d*x^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c) \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 2016 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 2560 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 2016 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 384 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 432 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 320 \sqrt{b} \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 9 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 + 18 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 - 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) + 48 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) + 54 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 - 324 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 + 18 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 + 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 - 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 + 24 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 - 3 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^4 + 54 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 - 9 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 - 96 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 + 192 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c) - 1440 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c) + 1728 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^5 \tan(-1
\end{aligned}$$

$$\begin{aligned}
& /2*d*x + 1/2*c)^4*\tan(1/3*c) - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 27*\sqrt{b}*d*x^4*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 162*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 - 9*\sqrt{b}*d*x^4*t \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 432*\sqrt{b}* \\
& d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan \\
& (1/3*c)^2 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 162*\sqrt{b}* \\
& d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 \\
& *\tan(1/3*c)^2 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9936* \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5 \\
& *\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 + 9600*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^3 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2 \\
& *d*x + 1/2*c)^2*\tan(1/3*c)^3 + 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 4800*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 \\
& + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^3 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + 5760*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 320* \\
& \sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 9*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 - 27*\sqrt{b}*d*x^4*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^5*\tan(1/3*c)^4 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& -1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/
\end{aligned}$$

$$\begin{aligned}
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 \\
& + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 23328*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 4 \\
& 32*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^3*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c \\
&)^3*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1 \\
& 440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(1/3*c)^4 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9936*\sqrt{ \\
& t(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (1/3*c)^5 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(\\
& 1/3*c)^5 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/ \\
& 3*c)^5 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
& ^5 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^5 + 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^5 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^5 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/ \\
& 6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + 5760*\sqrt{ \\
& t(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^5 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^5 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 9*\sqrt{b} \\
& (b)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 - 320*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan \\
& (-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 +
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c) - 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
&)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 \\
& + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 1440*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 \\
& - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 324*s \\
& qrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*ta \\
& n(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2* \\
& an(1/3*c)^2*\tan(c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/ \\
& 6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*ta \\
& n(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2 \\
& *c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + \\
& 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(- \\
& 1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 7 \\
& 2*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c \\
&)^2 + 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440* \\
& sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 19200*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4* \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
&)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 + 128 \\
& 0*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c) \\
& ^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*t \\
& an(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400 \\
& *\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x \\
& + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 640*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 320*\sqrt{ \\
& b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - \\
& 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^3*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^3*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3 \\
& *c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c \\
&)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c) \\
& ^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& *\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - \\
& 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^4*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4*\tan(c)^2 - 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 \\
& - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4* \\
& tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2 \\
& *\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + \\
& 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{ \\
& b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 1440*\sqrt{b}*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 5760*s \\
& qrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2
\end{aligned}$$

$$\begin{aligned}
& + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^5\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^5\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^5\tan(c)^2 - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 3456\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 - 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 - 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 + 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 + 3\sqrt{b}*d*x^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(1/3*c)^6\tan(c)^2 - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^6\tan(c)^2 + 1280\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^6\tan(c)^2 - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^6\tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 768\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 + 864\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 640\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 48\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^6\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^6\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^6\tan(c)^2 - 9\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2 + 54\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4 - 3\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^6 - 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c) + 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c) + 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c) + 18\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2 - 324\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2 + 54\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2 - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^5\tan(-1/2*d*x + 1/2*c)^2 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c)^2 + 48\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^3 - 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3 + 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3 + 18*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3 \\
& *\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4 - 144*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 96*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c) - 2880*\sqrt{b}*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c) + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 1440*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 5760*s \\
& qrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c) + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1 \\
& /2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) + 96*\sqrt{b}*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 9*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*t \\
& an(1/3*c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)* \\
& tan(1/3*c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^ \\
& 2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 144*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^ \\
& 2 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*t \\
& an(1/3*c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 19872*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 11520*\sqrt{b}* \\
& tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 432*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - \\
& 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - \\
& 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440 \\
& *sqrt(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 4800 \\
& *sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 5760* \\
& sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3 + 1280*s \\
& qrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 - 9600*sqr \\
& rt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 \\
& *tan(1/3*c)^3 + 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3 \\
& *tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*sqrt(\\
& b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^3 - 640*sqrt(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^3 + 320*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^3 - 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^3 + 19200*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 19200*sqrt(b)*\tan(1/2 \\
& *d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 4800*sqrt(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 54*sqrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 27*sqrt(b)*d*x^4*ta \\
& n(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 - 11664*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4 - 1008*sqrt(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(1/3*c)^4 + 144*sqrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 72* \\
& sqrt(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 54*sqrt \\
& (b)*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*sqrt(b)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872 \\
& *sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^4 - 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c \\
&)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872*sqrt(b)*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*sqrt(\\
& b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*sqrt(b) \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 432*sqrt(b)* \\
& tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1 \\
& /3*c)^4 + 216*sqrt(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4 - 1008*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^4 - 11664*sqrt(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 4800*sqrt(b)*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 1440*sqrt(b)*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 5760*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5 + 5760*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5 - 1728*sqrt(b)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^5 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c) \\
& ^6*\tan(1/3*c)^5 - 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c) \\
&)^2*\tan(1/3*c)^5 + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c) \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^5 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^5 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^5 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^5 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^5 - 3*\sqrt{b}*d*x^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 1280*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(1/3*c)^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 864*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 6 \\
& 40*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 48 \\
& *\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*s \\
& qrt(b)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*sq \\
& rt(b)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*\sqrt{b}*d* \\
& x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 72*\sqrt{b}*d*x \\
& ^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*\sqrt{b}*d*x^4* \\
& tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) + \\
& 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - \\
& 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 24 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*sq \\
& rt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4* \\
& tan(c) + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) \\
&) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) \\
&)^2*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& 1/3*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/3*c)^2 * \tan(c) - 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x \\
& + 1/2*c) * \tan(1/3*c)^2 * \tan(c) + 216 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan \\
& (-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c) - 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + \\
& 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^2 * \tan(c) - 72 * \sqrt{b} * d*x^4 * \tan(- \\
& 1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^2 * \tan(c) + 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \\
& \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2 \\
& *c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + \\
& 216 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan \\
& (c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) - 216 * \sqrt{b} \\
& (b) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^4 * \tan(c) - 432 * \\
& \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^4 * \tan(c) - \\
& 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^4 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan \\
& (-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) \\
& ^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 216 * \sqrt{b} * \tan(1/2*d*x + \\
& 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24 * \sqrt{b} * \tan(1/2*d \\
& *x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan \\
& (1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) - 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \\
& \tan(1/3*c)^6 * \tan(c) + 24 * \sqrt{b} * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c \\
&) + 3 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan \\
& (1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(c)^2 + 9 * \sqrt{b} * d*x^4 * \tan(1/ \\
& 2*d*x + 1/6*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + \\
& 1/6*c)^5 * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c) \\
& ^6 * \tan(c)^2 + 48 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c) \\
& * \tan(c)^2 - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan \\
& (-1/2*d*x + 1/2*c) * \tan(c)^2 - 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d \\
& *x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2 \\
& *c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 + 108 * \sqrt{b} * d \\
& *x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 54 * \sqrt{b} * d \\
& *x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 640 * \sqrt{b} * \\
& \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c \\
&)^2 - 2016 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d \\
& *x + 1/2*c)^2 * \tan(c)^2 + 768 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1 \\
& /6*c)^5 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 288 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) \\
& * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 48 * \sqrt{b} * d*x^4 \\
& * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 72 * \sqrt{b} * \tan(1/2 \\
& *d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 1 \\
& 44 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2 \\
& *c)^3 * \tan(c)^2 - 24 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \\
& \tan(c)^2 + 3 * \sqrt{b} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan \\
& (1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 \\
& + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + \\
& 1/2*c)^4 * \tan(c)^2 - 1280 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c \\
&)^3 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) * \tan \\
& (1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d \\
& *x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 1440 * \sqrt{b} * \tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 54*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 640*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 96*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)
\end{aligned}$$

$$\begin{aligned}
&)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(-1/2 dx + 1/2 c) \tan(1/3 c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c) \\
& \tan(1/3 c)^6 \tan(c)^2 - 288 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2 dx + 1/6 c) \tan(-1/2 dx \\
& x + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(-1/2 dx + 1/2 c)^3 \tan(1/3 c)^6 \tan(c)^2 - 3 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^4 + 54 \sqrt{b} dx \\
& x^4 \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^2 - 9 \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^4 - 96 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^5 \\
& - 48 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^6 - 48 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^3 \tan(-1/2 dx + 1/2 c) + 144 \sqrt{b} dx^4 \tan(1/2 \\
& dx + 1/2 c) \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c) + 72 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c) + 48 \sqrt{b} \\
& b) \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c) - 108 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^2 \tan(-1/2 dx + 1/2 c)^2 + 54 \sqrt{b} \\
& dx^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^2 + 640 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx + 1/2 c)^2 - 2016 \sqrt{b} \\
& b) \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 + 768 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/ \\
& 2 c)^2 - 288 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^2 - 48 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c) \tan(-1/2 dx + 1/2 c) \\
& ^3 + 72 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^3 + 144 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^4 \tan(\\
& -1/2 dx + 1/2 c)^3 + 24 \sqrt{b} \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^3 - 3 \sqrt{b} dx^4 \tan(-1/2 dx + 1/2 c)^4 - 96 \sqrt{b} \tan(1/2 dx + 1 \\
& /2 c)^4 \tan(1/2 dx + 1/6 c) \tan(-1/2 dx + 1/2 c)^4 + 432 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^4 - 1280 \sqrt{b} \\
&) \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx + 1/2 c)^4 + 432 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c \\
&)^4 - 96 \sqrt{b} \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^4 - 1440 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^4 \tan(1/3 c) + 1728 \sqrt{b} \\
& \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^5 \tan(1/3 c) - 384 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(1/3 c) + 2880 \sqrt{b} \tan(1/ \\
& 2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) - 11520 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx \\
& + 1/2 c)^2 \tan(1/3 c) + 11520 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) - 3456 \sqrt{b} \tan(1/2 dx + 1/ \\
& 2 c) \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) + 192 \sqrt{b} \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) - 96 \sqrt{b} \tan \\
& n(1/2 dx + 1/2 c)^4 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) + 1728 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c) \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) \\
& - 5760 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) + 5760 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c \\
& c)^3 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) - 1440 \sqrt{b} \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) + 54 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c \\
& ^2 \tan(1/3 c)^2 - 27 \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^2 \tan(1/3 c)^2 -
\end{aligned}$$

$$\begin{aligned}
& 4800\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^2 + 9 \\
& 936\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^2 - 57 \\
& 60\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^2 + 432 \\
& \sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^2 + 144\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^2 + 216\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan \\
& \tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan \\
& (-1/2*d*x + 1/2*c)\tan(1/3*c)^2 + 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(-1/ \\
& 2*d*x + 1/2*c)\tan(1/3*c)^2 + 54\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/3*c)^2 + 2880\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(-1/ \\
& 2*d*x + 1/2*c)^2\tan(1/3*c)^2 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/ \\
& 2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^2 + 38400\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^2 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x \\
& + 1/2*c)^2\tan(1/3*c)^2 + 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(-1/2*d*x \\
& + 1/2*c)^2\tan(1/3*c)^2 + 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + \\
& 1/2*c)^3\tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1 \\
& /6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^2 + 216\sqrt{b}\tan(1/2*d*x + 1/ \\
& 6*c)^4\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2 - 5760\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2 + 9936\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan \\
& (1/3*c)^2 - 4800\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan \\
& (1/3*c)^2 - 4800\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(\\
& 1/3*c)^3 + 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^3\tan(\\
& 1/3*c)^3 - 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(\\
& 1/3*c)^3 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan(1/3 \\
& *c)^3 - 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^3 + 640\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 - 11520\sqrt{b}\tan(\\
& 1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^3 + 38400\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d* \\
& x + 1/2*c)^2\tan(1/3*c)^3 - 38400\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x \\
& + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 + 9600\sqrt{b}\tan(1/2*d*x \\
& + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 - 1280\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 + 5760\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 - 4800\sqrt{b} \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 - 9\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + \\
& 1/6*c)\tan(1/3*c)^4 + 9936\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6 \\
& *c)^2\tan(1/3*c)^4 - 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6 \\
& *c)^3\tan(1/3*c)^4 + 9936\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c) \\
& ^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^4 + 72\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^4 + 432\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1 \\
& /3*c)^4 + 216\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*
\end{aligned}$$

$$\begin{aligned}
& c)^4 - 2016\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3* \\
& c)^4 + 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d \\
& *x + 1/2*c)^2\tan(1/3*c)^4 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x \\
& + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^4 + 9600\sqrt{b}\tan(1/2*d*x \\
& + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^4 + 144\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^4 + 216\sqrt{b}\tan(1/2*d*x + \\
& 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^4 + 432\sqrt{b}\tan(1/2*d*x + \\
& 1/2*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/ \\
& 6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c) \\
& ^4\tan(1/3*c)^5 + 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)* \\
& \tan(1/3*c)^5 - 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan \\
& (1/3*c)^5 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(\\
& 1/3*c)^5 - 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^5 + 768\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5 - 3456\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^5 + 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^5 - 96\sqrt{b}\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5 - 48\sqrt{b}\tan(1/2*d \\
& *x + 1/2*c)^3\tan(1/3*c)^6 - 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x \\
& + 1/6*c)\tan(1/3*c)^6 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6 \\
& *c)^2\tan(1/3*c)^6 - 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^6 + 48\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6 + 72\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6 - 288\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6 + 192\sqrt{b}\tan \\
& (1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6 + 24\sqrt{b}\tan(-1/ \\
& 2*d*x + 1/2*c)^3\tan(1/3*c)^6 - 24\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan \\
& (c) + 72\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(c) - \\
& 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^4\tan(c) - 48\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^6\tan(c) - 144\sqrt{b}*d*x^4 \\
& *tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c) + 72\sqrt{b}*d*x^4\tan \\
& (1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c) - 144\sqrt{b}*d*x^4\tan(1/ \\
& 2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^2\tan(c) - 24\sqrt{b}*d*x^4\tan(-1/2*d \\
& *x + 1/2*c)^3\tan(c) + 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2 \\
& *c)^4\tan(c) + 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan \\
& (-1/2*d*x + 1/2*c)^4\tan(c) + 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d \\
& *x + 1/2*c)^4\tan(c) + 72\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/3*c)^2\tan \\
& (c) - 216\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c) \\
&)^2\tan(c) - 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(\\
& 1/3*c)^2\tan(c) - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^2\tan(c) + 7 \\
& 2\sqrt{b}*d*x^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^2\tan(c) + 144\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2\tan(c) + 216\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2\tan(c) - 72\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/3*c)^4\tan(c) - 432\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^4\tan(c) - 216\sqrt{b}\tan(1 \\
& /2*d*x + 1/6*c)^4\tan(1/3*c)^4\tan(c) + 72\sqrt{b}\tan(-1/2*d*x + 1/2*c)^4\sqrt{b} \\
& \tan(1/3*c)^4\tan(c) - 48\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)
\end{aligned}$$

$$\begin{aligned}
& - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^6\tan(c) - 18\sqrt{b}*d*x^4 \\
& * \tan(1/2*d*x + 1/2*c)^2\tan(c)^2 + 9\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^2\tan \\
& \tan(c)^2 + 320\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^3\tan(c)^2 \\
& - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^4\tan(c)^2 + 3 \\
& 84\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^5\tan(c)^2 - 144\sqrt{b} \\
& \tan(b)\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^6\tan(c)^2 - 48\sqrt{b}*d*x^4 \\
& \tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)\tan(c)^2 - 72\sqrt{b}\tan(1/2* \\
& d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c)^2 - 144* \\
& \sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c) \\
& *\tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c)\tan(c)^2 \\
& - 18\sqrt{b}*d*x^4\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 192\sqrt{b}\tan(1/2 \\
& *d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 + 864 \\
& *\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c \\
&)^2\tan(c)^2 - 2560\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan \\
& \tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 + 864\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2* \\
& d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 192\sqrt{b}\tan(1/2*d*x + \\
& 1/6*c)^5\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/2*c \\
&)^4\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 72\sqrt{b}\tan(1/2 \\
& *d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 - 1008\sqrt{b}* \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 \\
& + 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 + 1 \\
& 440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)\tan(c) \\
& ^2 - 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)* \\
& \tan(c)^2 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(1 \\
& /3*c)\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan \\
& \tan(1/3*c)\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)\tan(c)^2 \\
& - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)\tan \\
& (c)^2 + 3456\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d \\
& *x + 1/2*c)^2\tan(1/3*c)\tan(c)^2 - 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)\tan(c)^2 + 11520\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/3*c)\tan(c)^2 - 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2 \\
& *c)^2\tan(1/3*c)\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x \\
& + 1/2*c)^4\tan(1/3*c)\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2 \\
& *d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)\tan(c)^2 + 1440\sqrt{b}*tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)\tan(c)^2 + 9\sqrt{b} \\
&)*d*x^4\tan(1/3*c)^2\tan(c)^2 + 1440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2 \\
& *d*x + 1/6*c)\tan(1/3*c)^2\tan(c)^2 - 11664\sqrt{b}\tan(1/2*d*x + 1/2*c)^3 \\
& \tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^2\tan(c)^2 + 19200\sqrt{b}\tan(1/2*d*x + \\
& 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^2\tan(c)^2 - 11664\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^2\tan(c)^2 + 1440\sqrt{b} \\
& *\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^2\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 320*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 - 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 + 18*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 - 9*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6 + 48*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) + 18*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^2 - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 - 2560*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2 + 24*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3 + 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3 + 72*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 - 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4 + 384*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4 - 1008*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 320*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) - 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c) + 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c) - 1728*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c) + 96*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c) - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 3456*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 2880*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 384*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 1728*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 9*\text{sqrt}(b)*d*x^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^2 - 11664*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 19200*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2 - 11664*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2 + 72*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 432*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 19872*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 9600*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 1008*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 320*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3 + 19200*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3 - 19200*\text{sqrt}(b)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 320*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/3*c)^6*\tan(c) + 3*\sqrt{b}*d*x^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 288*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 640*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
& c)^3 \tan(1/3*c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) \\
&)*\tan(1/3*c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(1/3*c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2 + 144*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\sqrt{b}*\tan \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 - 2016*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 72*\sqrt{b}*\tan(-1/2* \\
& d*x + 1/2*c)^3*\tan(1/3*c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^3 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3 - \\
& 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3 + 640*\sqrt{b}*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^4 - \\
& 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 96*\sqrt{b}*\tan(1/3*c)^5 - 24*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(c) + 24*\sqrt{b}*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) \\
& - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*\tan(\\
& 1/3*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) \\
&)*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192* \\
& \sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 24*\sqrt{b}*\tan \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)*\tan(c)^2 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan \\
& (1/3*c)*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)*\tan(c)^2 \\
& - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 1008*\sqrt{b}*\tan \\
& \tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c \\
&)*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan \\
& (c)^2 + 320*\sqrt{b}*\tan(1/3*c)^3*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c \\
&)^3 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) - 1008*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /6*c)^3 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) + 72*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) - 96*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2 - 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2 \\
& *d*x + 1/2*c)^2 + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3 + 384*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)*\tan(1/3*c) + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) - 192*s \\
& \sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^2 + 72*\sqrt{b} \\
&)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 320*\sqrt{b}*\tan(1/3*c)^3 - 48*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(c) \\
& - 72*\sqrt{b}*\tan(1/3*c)^2*\tan(c) - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 24*\sqrt{b}*\tan(-1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)*\tan(c)^2 - 96*\sqrt{b}*\tan(1/3*c)*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c) + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c) \\
& - 96*\sqrt{b}*\tan(1/3*c) - 24*\sqrt{b}*\tan(c))/(d*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 2* \\
& d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \\
& *\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 \\
& + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 6*d* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^4*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3* \\
& c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*t \\
& an(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 6* \\
& d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + \\
& 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*t \\
& an(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + \\
& 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*t \\
& an(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4*t
\end{aligned}$$

$$\begin{aligned}
& \text{an}(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 12*d* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 \\
& *\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + \\
& 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 9*d*t \\
& \text{an}(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*t \\
& \text{an}(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 3*d \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 3*d*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 12*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 2*d \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + d*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c \\
&)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 \\
& + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2* \\
& \tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(\\
& c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6
\end{aligned}$$

$$\begin{aligned}
& *c)^4 \tan(1/3*c)^4 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6* \\
& c)^6 \tan(1/3*c)^4 \tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6* \\
& c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c)^2 \\
& + 6*d*\tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c)^2 \\
& + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^ \\
& 4 \tan(1/3*c)^4 \tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^ \\
& 4 \tan(1/3*c)^4 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \\
& * \tan(1/3*c)^6 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 * \\
& \tan(1/3*c)^6 \tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan(c)^2 + 2* \\
& d*\tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 12 \\
& *d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 * \tan \\
& (1/3*c)^6 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 * \tan \\
& (1/3*c)^6 \tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 * \tan \\
& (1/3*c)^6 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 * \tan \\
& (1/3*c)^6 \tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan \\
& (-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 * \tan \\
& (-1/2*d*x + 1/2*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 * \tan \\
& (-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 * \\
& \tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2* \\
& d*x + 1/2*c)^2 \tan(1/3*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6 \\
& *c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4 \tan \\
& (1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 18*d*\tan(1/2*d*x \\
& + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 3 \\
& *d*\tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 9*d*\tan(1/ \\
& 2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^4 \tan \\
& (1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 36*d*\tan(1/2*d* \\
& x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + \\
& 6*d*\tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 3*d*\tan(1 \\
& /2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + \\
& 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 9*d \\
& * \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 3*d*\tan(1/2* \\
& d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2* \\
& c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/6*c)^6 \tan(1/3 \\
& *c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 1 \\
& 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 * \tan \\
& (1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c) \\
& ^6 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 3*d* \\
& \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + d*\tan(1/2*d*x \\
& + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4 * \tan \\
& (1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 4*d*\tan(1/2*d*x + 1 \\
& /2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 3*d*\tan(1 \\
& /2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(c)^2 +
\end{aligned}$$

$$\begin{aligned}
& 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*d* \\
& \tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*t \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 18*d*t \\
& \tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^2*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1 \\
& /2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1/ \\
& 2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + d*\tan(1/2* \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& *d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c \\
&)^6*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^6*\tan(c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2* \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2* \\
& *d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)^4 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1 \\
& /3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 + \\
& 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& *d*x + 1/2*c)^2*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2 \\
& *c)^2*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3 \\
& *c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 3 \\
& *d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/6*c)^4* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2* \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^4 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*
\end{aligned}$$

$$\begin{aligned}
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c) \\
& ^4*\tan(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^6 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + \\
& d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c \\
&)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2 \\
& *c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6 \\
& *c)^4*\tan(1/3*c)^4*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1 \\
& /2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*t \\
& an(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x \\
& + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4 + 2*d*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (-1/2*d*x + 1/2*c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 6 \\
& *d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + \\
& 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 9*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 + \\
& 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 36*d*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 6 \\
& *d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*t \\
& an(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 12*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/6*c)^2 \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3 \\
& *c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c)^
\end{aligned}$$

$$\begin{aligned}
& 2*\tan(1/3*c)^6 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 2*d*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 6*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan \\
& (c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + d*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^4 + d*\tan(1/2*d*x + 1/6*c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan \\
& (-1/2*d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/ \\
& 6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + \\
& 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 6*d*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + d*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c \\
&)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + \\
& 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 6*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/3*c)^4*\tan(\\
& c)^2 + d*\tan(1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 + d*\tan \\
& (-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 6*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + \\
& 3*d*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 + 2*d* \\
& \tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/3*c)^2 + d*\tan(c)^2 + d
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.82 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.81

$$\int \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12d (\cos(2c + 2dx) + 1)}$$

[In] `int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2),x)`

[Out] `(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*d*(cos(2*c + 2*d*x) + 1))`

3.142 $\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx$

Optimal result	732
Rubi [A] (verified)	732
Mathematica [A] (verified)	733
Maple [A] (verified)	733
Fricas [A] (verification not implemented)	734
Sympy [B] (verification not implemented)	734
Maxima [A] (verification not implemented)	735
Giac [B] (verification not implemented)	735
Mupad [B] (verification not implemented)	735

Optimal result

Integrand size = 23, antiderivative size = 63

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{x \sqrt{b \cos(c + dx)}}{2 \sqrt{\cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d}$$

[Out] $1/2*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+1/2*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{x \sqrt{b \cos(c + dx)}}{2 \sqrt{\cos(c + dx)}} + \frac{\sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}{2d}$$

[In] `Int[Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]],x]`

[Out] $(x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(2*\text{Sqrt}[\text{Cos}[c + d*x]]) + (\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(2*d)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)
)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \cos^2(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} + \frac{\sqrt{b \cos(c + dx)} \int 1 dx}{2\sqrt{\cos(c + dx)}} \\ &= \frac{x \sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} (2(c + dx) + \sin(2(c + dx)))}{4d \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (Sqrt[b*Cos[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[Cos[c + d
*x]])
```

Maple [A] (verified)

Time = 3.07 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{\sqrt{\cos(dx+c)b}(\cos(dx+c)\sin(dx+c)+dx+c)}{2d\sqrt{\cos(dx+c)}}$	42
risch	$\frac{\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{i(dx+c)}x}{e^{2i(dx+c)}+1} - \frac{i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{3i(dx+c)}}{4(e^{2i(dx+c)}+1)d} + \frac{i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{-i(dx+c)}}{4(e^{2i(dx+c)}+1)d}$	136

[In] `int(cos(d*x+c)^(3/2)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $1/2/d*(\cos(d*x+c)*b)^(1/2)*(\cos(d*x+c)*\sin(d*x+c)+d*x+c)/\cos(d*x+c)^(1/2)$

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 150, normalized size of antiderivative = 2.38

$$\int \cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}dx$$

$$= \left[\frac{2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) + \sqrt{-b}\log\left(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{\cos(dx+c)}\right)}{4d} \right]$$

[In] `integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] $[1/4*(2*\sqrt{b*\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c) + \sqrt{-b}*\log(2*b*\cos(d*x+c)^2 - 2*\sqrt{b*\cos(d*x+c)}*\sqrt{-b}*\sqrt{\cos(d*x+c)}*\sin(d*x+c) - b))/d, 1/2*(\sqrt{b*\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c) + \sqrt{b}*\arctan(\sqrt{b*\cos(d*x+c)}*\sin(d*x+c)/(\sqrt{b}*\cos(d*x+c)^(3/2)))/d]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(56) = 112$.

Time = 32.91 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.92

$$\int \cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}dx$$

$$= \begin{cases} x\sqrt{b\cos(c)}\cos^{\frac{3}{2}}(c) & \text{for } d = 0 \\ 0 & \text{for } c = -dx + \frac{\pi}{2} \vee c = -c \\ \frac{x\sqrt{b\cos(c+dx)}\sin^2(c+dx)}{2\sqrt{\cos(c+dx)}} + \frac{x\sqrt{b\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)}{2} + \frac{\sqrt{b\cos(c+dx)}\sin(c+dx)\sqrt{\cos(c+dx)}}{2d} & \text{otherwise} \end{cases}$$

[In] `integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(1/2),x)`

[Out] Piecewise((x*sqrt(b*cos(c))*cos(c)**(3/2), Eq(d, 0)), (0, Eq(c, -d*x + pi/2) | Eq(c, -d*x + 3*pi/2)), (x*sqrt(b*cos(c + d*x))*sin(c + d*x)**2/(2*sqrt(cos(c + d*x))) + x*sqrt(b*cos(c + d*x))*cos(c + d*x)**(3/2)/2 + sqrt(b*cos(c + d*x))*sin(c + d*x)*sqrt(cos(c + d*x))/(2*d), True))

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.40

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{(2 dx + 2 c + \sin(2 dx + 2 c)) \sqrt{b}}{4 d}$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))*sqrt(b)/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(51) = 102.

Time = 1.47 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.67

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b} dx \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 2 \sqrt{b} dx \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 2 \sqrt{b} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + \sqrt{b} dx + 2 \sqrt{b} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{2 \left(d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 2 d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + d \right)}$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] 1/2*(sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^4 + 2*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^2 - 2*sqrt(b)*tan(1/2*d*x + 1/2*c)^3 + sqrt(b)*d*x + 2*sqrt(b)*tan(1/2*d*x + 1/2*c))/(d*tan(1/2*d*x + 1/2*c)^4 + 2*d*tan(1/2*d*x + 1/2*c)^2 + d)

Mupad [B] (verification not implemented)

Time = 14.57 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.98

$$\int \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{\cos(c + d x)} \sqrt{b \cos(c + d x)} (\sin(c + d x) + \sin(3 c + 3 d x) + 4 d x \cos(c + d x))}{4 d (\cos(2 c + 2 d x) + 1)}$$

[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*d*(cos(2*c + 2*d*x) + 1))

3.143 $\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx$

Optimal result	736
Rubi [A] (verified)	736
Mathematica [A] (verified)	737
Maple [A] (verified)	737
Fricas [A] (verification not implemented)	737
Sympy [B] (verification not implemented)	738
Maxima [A] (verification not implemented)	738
Giac [A] (verification not implemented)	738
Mupad [B] (verification not implemented)	739

Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx = \frac{\sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] `Int[Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]],x]`

[Out] `(Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2) * b^(n - 1/2) * (Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \cos(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}}$$

[In] Integrate[Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)\sqrt{\cos(dx+c)b}}{d\sqrt{\cos(dx+c)}}$	29
risch	$-\frac{i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{2i(dx+c)}}{(e^{2i(dx+c)}+1)d} + \frac{i\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})}{(e^{2i(dx+c)}+1)d}$	85

[In] int(cos(d*x+c)^(1/2)*(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{d \sqrt{\cos(dx + c)}}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(29) = 58$.

Time = 1.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.88

$$\int \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} dx = \begin{cases} x \sqrt{b \cos(c)} \sqrt{\cos(c)} & \text{for } d = 0 \\ 0 & \text{for } c = -dx + \frac{\pi}{2} \vee c = -dx + \frac{3\pi}{2} \\ \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}} & \text{otherwise} \end{cases}$$

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(1/2),x)

[Out] Piecewise((x*sqrt(b*cos(c))*sqrt(cos(c)), Eq(d, 0)), (0, Eq(c, -d*x + pi/2) | Eq(c, -d*x + 3*pi/2)), (sqrt(b*cos(c + d*x))*sin(c + d*x)/(d*sqrt(cos(c + d*x))), True))

Maxima [A] (verification not implemented)

none

Time = 0.44 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.41

$$\int \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} dx = \frac{\sqrt{b} \sin(dx+c)}{d}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] sqrt(b)*sin(d*x + c)/d

Giac [A] (verification not implemented)

none

Time = 0.54 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

$$\int \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} dx = \frac{2 \sqrt{b} \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{d \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + d}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] 2*sqrt(b)*tan(1/2*d*x + 1/2*c)/(d*tan(1/2*d*x + 1/2*c)^2 + d)

Mupad [B] (verification not implemented)

Time = 14.17 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.38

$$\int \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} dx = \frac{\sqrt{\cos(c + dx)} \sin(2c + 2dx) \sqrt{b \cos(c + dx)}}{d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(d*(cos(2*c + 2*d*x) + 1))

$$3.144 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx$$

Optimal result	740
Rubi [A] (verified)	740
Mathematica [A] (verified)	741
Maple [A] (verified)	741
Fricas [A] (verification not implemented)	741
Sympy [A] (verification not implemented)	742
Maxima [A] (verification not implemented)	742
Giac [F]	742
Mupad [B] (verification not implemented)	743

Optimal result

Integrand size = 23, antiderivative size = 24

$$\int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx = \frac{x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] $x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}} dx = \frac{x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[In] `Int[Sqrt[b*Cos[c + d*x]]/Sqrt[Cos[c + d*x]],x]`

[Out] `(x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int 1 dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx = \frac{x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Sqrt[Cos[c + d*x]],x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

Maple [A] (verified)

Time = 2.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
risch	$\frac{x \sqrt{\cos(dx+c)b}}{\sqrt{\cos(dx+c)}}$	21
default	$\frac{\sqrt{\cos(dx+c)b} (dx+c)}{d \sqrt{\cos(dx+c)}}$	28

[In] int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] x*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 94, normalized size of antiderivative = 3.92

$$\begin{aligned} &\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx \\ &= \left[\frac{\sqrt{-b} \log \left(2 b \cos(dx + c)^2 - 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)} \sin(dx + c) - b \right)}{2 d}, \frac{\sqrt{b} \arctan \left(\frac{\sqrt{b \cos(dx + c)}}{\sqrt{b \cos(dx + c)}} \right)}{d} \right] \end{aligned}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/d, sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]

Sympy [A] (verification not implemented)

Time = 0.61 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx = \frac{x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(1/2),x)

[Out] x*sqrt(b*cos(c + d*x))/sqrt(cos(c + d*x))

Maxima [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx = \frac{2 \sqrt{b} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/sqrt(cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

$$\int \frac{\sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} dx = \frac{x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(1/2),x)

[Out] (x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

$$3.145 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal result	744
Rubi [A] (verified)	744
Mathematica [A] (verified)	745
Maple [A] (verified)	745
Fricas [A] (verification not implemented)	745
Sympy [F]	746
Maxima [B] (verification not implemented)	746
Giac [F]	746
Mupad [F(-1)]	747

Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[Out] $\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3855}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]/\operatorname{Cos}[c+d*x]^{(3/2)}, x]$

[Out] $(\operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]]*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]])/(d*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]])$

Rule 17

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\operatorname{Sqrt}[b*v]/\operatorname{Sqrt}[a*v]), \operatorname{Int}[u*v^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, m\}, x$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{IGtQ}[n+1/2, 0]$ && $\operatorname{IntegerQ}[m+n]$

Rule 3855

$\operatorname{Int}[\operatorname{csc}[(c_*)+(d_*)*(x_*)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c+d*x]]/d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sec(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx = \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 3.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

method	result	size
default	$-\frac{2\sqrt{\cos(dx+c)}b \operatorname{arctanh}(\cot(dx+c)-\csc(dx+c))}{d\sqrt{\cos(dx+c)}}$	40
risch	$-\frac{\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}-i)}{\sqrt{\cos(dx+c)}d} + \frac{\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}+i)}{\sqrt{\cos(dx+c)}d}$	73

[In] int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/d*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)*arctanh(cot(d*x+c)-csc(d*x+c))

Fricas [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 113, normalized size of antiderivative = 3.42

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx = \left[\frac{\sqrt{b} \log \left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3} \right)}{2d}, \right. \\ \left. - \frac{\sqrt{-b} \operatorname{arctan} \left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b} \sin(dx+c)}{b\sqrt{\cos(dx+c)}} \right)}{d} \right]$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] [1/2*sqrt(b)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/d, -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/d]

Sympy [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx$$

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(3/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))/cos(c + d*x)**(3/2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(29) = 58.

Time = 0.45 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.97

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{b}(\log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - \log(\cos(dx + c)^2 + \sin(dx + c)^2 - 2 \sin(dx + c) + 1))}{2d}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] 1/2*sqrt(b)*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/d

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{3}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{3/2}} dx$$

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(3/2), x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(3/2), x)
```

$$3.146 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal result	748
Rubi [A] (verified)	748
Mathematica [A] (verified)	749
Maple [A] (verified)	749
Fricas [A] (verification not implemented)	750
Sympy [F]	750
Maxima [A] (verification not implemented)	750
Giac [F]	750
Mupad [B] (verification not implemented)	751

Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(3/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3852, 8}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[In] `Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(5/2),x]`

[Out] `(Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sec^2(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\sqrt{b \cos(c + dx)} \text{Subst}(\int 1 dx, x, -\tan(c + dx))}{d \sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)}$$

```
[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(5/2), x]
```

```
[Out] (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))
```

Maple [A] (verified)

Time = 2.92 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)\sqrt{\cos(dx+c)b}}{d \cos(dx+c)^{\frac{3}{2}}}$	29
risch	$\frac{2i\sqrt{\cos(dx+c)b}}{\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)}$	38

```
[In] int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(5/2), x, method=_RETURNVERBOSE)
```

```
[Out] sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(3/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.88

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)} \sin(dx + c)}{d \cos(dx + c)^{\frac{3}{2}}}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

Sympy [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx$$

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(5/2),x)

[Out] Integral(sqrt(b*cos(c + d*x))/cos(c + d*x)**(5/2), x)

Maxima [A] (verification not implemented)

none

Time = 0.44 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.69

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{2 \sqrt{b} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c))^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1} d$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c))^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{5}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(5/2), x)

Mupad [B] (verification not implemented)

Time = 14.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.84

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + 1i)}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

[In] `int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(5/2),x)`

[Out] `((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))`

$$3.147 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx$$

Optimal result	752
Rubi [A] (verified)	752
Mathematica [A] (verified)	753
Maple [A] (verified)	753
Fricas [A] (verification not implemented)	754
Sympy [F(-1)]	754
Maxima [B] (verification not implemented)	755
Giac [F]	755
Mupad [F(-1)]	756

Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{\frac{5}{2}}(c+dx)}$$

[Out] 1/2*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+1/2*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{7}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{\sin(c+dx)\sqrt{b \cos(c+dx)}}{2d \cos^{\frac{5}{2}}(c+dx)}$$

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(7/2),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(2*d*Sqrt[Cos[c + d*x]]) + (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(2*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3853


```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] & & IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sec^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{2d \cos^{\frac{5}{2}}(c + dx)} + \frac{\sqrt{b \cos(c + dx)} \int \sec(c + dx) dx}{2\sqrt{\cos(c + dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{2d \sqrt{\cos(c + dx)}} + \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{2d \cos^{\frac{5}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.72

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} (\operatorname{arctanh}(\sin(c + dx)) \cos^2(c + dx) + \sin(c + dx))}{2d \cos^{\frac{5}{2}}(c + dx)}$$

```
[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(7/2), x]
```

```
[Out] (Sqrt[b*Cos[c + d*x]]*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/
(2*d*Cos[c + d*x]^(5/2)))
```

Maple [A] (verified)

Time = 3.09 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{(-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c)) \sqrt{\cos(dx+c)b}}{2d \cos(dx+c)^{\frac{5}{2}}}$	84
risch	$-\frac{i \sqrt{\cos(dx+c)b} (e^{3i(dx+c)} - e^{i(dx+c)})}{\sqrt{\cos(dx+c)} d (e^{2i(dx+c)} + 1)^2} + \frac{\sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)} + i)}{2\sqrt{\cos(dx+c)} d} - \frac{\sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)} - i)}{2\sqrt{\cos(dx+c)} d}$	132

[In] `int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(7/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}d(-\cos(dx+c)^2 \ln(-\cot(dx+c) + \csc(dx+c) - 1) + \cos(dx+c)^2 \ln(-\cot(dx+c) + \csc(dx+c) + 1) + \sin(dx+c)) \cdot (\cos(dx+c) \cdot b)^{1/2} / \cos(dx+c)^{5/2}$

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.79

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx$$

$$= \left[\frac{\sqrt{b \cos(dx + c)}^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b \cos(dx+c)}\sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b \cos(dx + c)}\sqrt{\cos(dx + c)}}{4d \cos(dx + c)^3} - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b} \sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right) \cos(dx + c)^3 - \sqrt{b \cos(dx + c)}\sqrt{\cos(dx + c)} \sin(dx + c)}{2d \cos(dx + c)^3} \right]$$

[In] `integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2),x, algorithm="fricas")`

[Out] $\left[\frac{1}{4} \cdot (\sqrt{b} \cdot \cos(dx + c))^3 \cdot \log(-(\sqrt{b} \cdot \cos(dx + c))^3 - 2 \cdot \sqrt{b} \cdot \cos(dx + c) \cdot \sqrt{\cos(dx + c)} \cdot \sin(dx + c) - 2 \cdot b \cdot \cos(dx + c)) / \cos(dx + c)^3 + 2 \cdot \sqrt{b} \cdot \cos(dx + c) \cdot \sqrt{\cos(dx + c)} \cdot \sin(dx + c) / (d \cdot \cos(dx + c)^3), -\frac{1}{2} \cdot (\sqrt{-b} \cdot \arctan(\sqrt{b \cos(dx + c)} \cdot \sqrt{-b} \cdot \sin(dx + c) / (b \cdot \sqrt{\cos(dx + c)})) \cdot \cos(dx + c)^3 - \sqrt{b \cos(dx + c)} \cdot \sqrt{\cos(dx + c)} \cdot \sin(dx + c)) / (d \cdot \cos(dx + c)^3) \right]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(7/2),x)`

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 661 vs. $2(60) = 120$.

Time = 0.44 (sec) , antiderivative size = 661, normalized size of antiderivative = 9.18

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx = \frac{\left(4(\sin(4dx + 4c) + 2\sin(2dx + 2c))\cos\left(\frac{3}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) - 4(\sin(4dx + 4c) + 2\sin(2dx + 2c))\cos\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) + 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) + 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{3}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right)\right)\sqrt{b}}{\left(2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1\right)^2 + \sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right)^2 + 2\sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) + 1 + \left(2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1\right)\log\left(\cos\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right), \cos(2dx + 2c)\right)^2 + \sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right)^2 - 2\sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) + 1 - 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{3}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right) + 4(\cos(4dx + 4c) + 2\cos(2dx + 2c) + 1)\sin\left(\frac{1}{2}\arctan(\sin(2dx + 2c), \cos(2dx + 2c))\right)\right)\sqrt{b}}{\left(2(2\cos(2dx + 2c) + 1)\cos(4dx + 4c) + \cos(4dx + 4c)^2 + 4\cos(2dx + 2c)^2 + \sin(4dx + 4c)^2 + 4\sin(4dx + 4c)\sin(2dx + 2c) + 4\sin(2dx + 2c)^2 + 4\cos(2dx + 2c) + 1\right)d}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out]
$$\begin{aligned} & -1/4*(4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) + (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) - 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))*\sqrt{b}/((2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*d) \end{aligned}$$

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\cos^{\frac{7}{2}}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(7/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{7/2}} dx$$

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(7/2), x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(7/2), x)
```

$$3.148 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{9}{2}}(c+dx)} dx$$

Optimal result	757
Rubi [A] (verified)	757
Mathematica [A] (verified)	758
Maple [A] (verified)	758
Fricas [A] (verification not implemented)	759
Sympy [F(-1)]	759
Maxima [B] (verification not implemented)	759
Giac [F]	760
Mupad [B] (verification not implemented)	760

Optimal result

Integrand size = 23, antiderivative size = 70

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} + \frac{\sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{\frac{7}{2}}(c+dx)}$$

[Out] $\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(3/2)}+1/3*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(7/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3852}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{\sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{\sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[In] $\text{Int}[\text{Sqrt}[b*\text{Cos}[c + d*x]]/\text{Cos}[c + d*x]^{(9/2)}, x]$

[Out] $(\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Cos}[c + d*x]^{(3/2)}) + (\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Cos}[c + d*x]^{(7/2)})$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sec^4(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\sqrt{b \cos(c + dx)} \text{Subst}\left(\int (1 + x^2) dx, x, -\tan(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)} + \frac{\sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \cos^{\frac{7}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(c + dx)} \left(\tan(c + dx) + \frac{1}{3} \tan^3(c + dx) \right)}{d \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(9/2), x]
```

```
[Out] (Sqrt[b*Cos[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[Cos[c + d*x]])
```

Maple [A] (verified)

Time = 2.96 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

method	result	size
default	$\frac{(2(\cos^2(dx+c))+1)\sqrt{\cos(dx+c)b} \sin(dx+c)}{3d \cos(dx+c)^{\frac{7}{2}}}$	42
risch	$\frac{4i\sqrt{\cos(dx+c)b}(3e^{2i(dx+c)}+1)}{3\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)^3}$	51

```
[In] int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(9/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(7/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.59

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)}(2 \cos(dx + c)^2 + 1) \sin(dx + c)}{3 d \cos(dx + c)^{\frac{7}{2}}}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(9/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(60) = 120.

Time = 0.45 (sec) , antiderivative size = 294, normalized size of antiderivative = 4.20

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{4((3 \cos(2 dx + 2 c) + 1) \sin(6 dx + 6 c) + 3(3 \cos(2 dx + 2 c) + 1) \sin(4 dx + 4 c) - 3 \cos(6 dx + 6 c) \sin(2 dx + 2 c) - 9 \cos(4 dx + 4 c) \sin(2 dx + 2 c)) \sqrt{b}}{3(2(3 \cos(4 dx + 4 c) + 3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + \cos(6 dx + 6 c)^2 + 6(3 \cos(2 dx + 2 c) + 1) \cos(6 dx + 6 c) + 9 \cos(4 dx + 4 c)^2 + 9 \cos(2 dx + 2 c)^2 + 6(\sin(4 dx + 4 c) + \sin(2 dx + 2 c)) \sin(6 dx + 6 c) + \sin(6 dx + 6 c)^2 + 9 \sin(4 dx + 4 c)^2 + 18 \sin(4 dx + 4 c) \sin(2 dx + 2 c) + 9 \sin(2 dx + 2 c)^2 + 6 \cos(2 dx + 2 c) + 1) d}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="maxima")

```
[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)
*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)
)*sin(2*d*x + 2*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) +
1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4
*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x +
4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*
x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 +
6*cos(2*d*x + 2*c) + 1)*d)
```

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\cos(dx + c)^{\frac{9}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(9/2), x)

Mupad [B] (verification not implemented)

Time = 16.12 (sec) , antiderivative size = 128, normalized size of antiderivative = 1.83

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{9}{2}}(c + dx)} dx$$

$$= \frac{2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + \sin(6c + 6dx) + 10i)}{3d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 10)}$$

[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(9/2),x)

[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.149 \quad \int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx$$

Optimal result	761
Rubi [A] (verified)	761
Mathematica [A] (verified)	763
Maple [A] (verified)	763
Fricas [A] (verification not implemented)	763
Sympy [F(-1)]	764
Maxima [B] (verification not implemented)	764
Giac [F]	765
Mupad [F(-1)]	766

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{3 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{\sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3 \sqrt{b \cos(c+dx)} \sin(c+dx)}{8d \cos^{\frac{5}{2}}(c+dx)}$$

[Out] 1/4*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(9/2)+3/8*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+3/8*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{\sqrt{b \cos(c+dx)}}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{3 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{3 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{\frac{5}{2}}(c+dx)} + \frac{\sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{\frac{9}{2}}(c+dx)}$$

[In] Int[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(11/2),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(8*d*Sqrt[Cos[c + d*x]]) + (Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(4*d*Cos[c + d*x]^(9/2)) + (3*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(8*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{b \cos(c + dx)} \int \sec^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
 &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sec^3(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\
 &= \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{3\sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{\frac{5}{2}}(c + dx)} \\
 &\quad + \frac{\left(3\sqrt{b \cos(c + dx)}\right) \int \sec(c + dx) dx}{8\sqrt{\cos(c + dx)}} \\
 &= \frac{3\text{arctanh}(\sin(c + dx))\sqrt{b \cos(c + dx)}}{8d\sqrt{\cos(c + dx)}} \\
 &\quad + \frac{\sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{3\sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{\frac{5}{2}}(c + dx)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$= \frac{\sqrt{b \cos(c + dx)}(3 \operatorname{arctanh}(\sin(c + dx)) \cos^4(c + dx) + (2 + 3 \cos^2(c + dx)) \sin(c + dx))}{8d \cos^{\frac{9}{2}}(c + dx)}$$

[In] Integrate[Sqrt[b*Cos[c + d*x]]/Cos[c + d*x]^(11/2),x]

[Out] (Sqrt[b*Cos[c + d*x]]*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(9/2))

Maple [A] (verified)

Time = 2.98 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.96

method	result
default	$\frac{(-3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+3(\cos^2(dx+c)) \sin(dx+c)+2 \sin(dx+c)) \sqrt{b \cos(dx+c)}}{8d \cos(dx+c)^{\frac{9}{2}}}$
risch	$-\frac{i\sqrt{\cos(dx+c)b}(3e^{7i(dx+c)}+11e^{5i(dx+c)}-11e^{3i(dx+c)}-3e^{i(dx+c)})}{4\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)^4} + \frac{3\sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)}+i)}{8\sqrt{\cos(dx+c)}d} - \frac{3\sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)}-i)}{8\sqrt{\cos(dx+c)}d}$

[In] int((cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(11/2),x,method=_RETURNVERBOSE)

[Out] 1/8/d*(-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)+3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(9/2)

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 227, normalized size of antiderivative = 2.12

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx$$

$$= \frac{\left[3\sqrt{b} \cos(dx+c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b \cos(dx+c)}(3 \cos(dx+c)^2 + 2) \sqrt{\cos(dx+c)} \right]}{16d \cos(dx+c)^5}$$

$$- \frac{3\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b \sin(dx+c)}}{b\sqrt{\cos(dx+c)}}\right) \cos(dx+c)^5 - \sqrt{b \cos(dx+c)}(3 \cos(dx+c)^2 + 2) \sqrt{\cos(dx+c)}}{8d \cos(dx+c)^5}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(d*cos(d*x + c)^5)]

Sympy [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(1/2)/cos(d*x+c)**(11/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1656 vs. 2(89) = 178.

Time = 0.48 (sec) , antiderivative size = 1656, normalized size of antiderivative = 15.48

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx = \text{Too large to display}$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2),x, algorithm="maxima")

[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)

```

*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*s
in(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + 3
*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(
8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2
*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(
2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8
*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin
(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*
x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)
*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arcta
n2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) +
6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*
c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(
4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x
+ 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*
x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c)
+ 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x +
2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4
*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos
(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x
+ 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))
*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d
*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2
+ 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d
*x + 2*c) + 1)*d)

```

Giac [F]

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(dx + c)}}{\cos^{\frac{11}{2}}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(1/2)/cos(d*x+c)^(11/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cos(d*x + c))/cos(d*x + c)^(11/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{b \cos(c + dx)}}{\cos^{\frac{11}{2}}(c + dx)} dx = \int \frac{\sqrt{b \cos(c + dx)}}{\cos(c + dx)^{11/2}} dx$$

```
[In] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(11/2),x)
```

```
[Out] int((b*cos(c + d*x))^(1/2)/cos(c + d*x)^(11/2), x)
```

3.150 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx$

Optimal result	767
Rubi [A] (verified)	767
Mathematica [A] (verified)	768
Maple [A] (verified)	769
Fricas [A] (verification not implemented)	769
Sympy [F(-1)]	769
Maxima [A] (verification not implemented)	770
Giac [B] (verification not implemented)	770
Mupad [B] (verification not implemented)	770

Optimal result

Integrand size = 23, antiderivative size = 101

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{3bx \sqrt{b \cos(c + dx)}}{8 \sqrt{\cos(c + dx)}} + \frac{3b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d} + \frac{b \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d}$$

[Out] $1/4*b*\cos(d*x+c)^{(5/2)}*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d+3/8*b*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+3/8*b*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{3bx \sqrt{b \cos(c + dx)}}{8 \sqrt{\cos(c + dx)}} + \frac{b \sin(c + dx) \cos^{\frac{5}{2}}(c + dx) \sqrt{b \cos(c + dx)}}{4d} + \frac{3b \sin(c + dx) \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}{8d}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)}*(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(3*b*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(8*\text{Sqrt}[\text{Cos}[c + d*x]]) + (3*b*\text{Sqrt}[\text{Cos}[c + d*x]])*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]/(8*d) + (b*\text{Cos}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(4*d)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \cos^4(c+dx) dx}{\sqrt{\cos(c+dx)}} \\
 &= \frac{b\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}\sin(c+dx)}{4d} + \frac{\left(3b\sqrt{b\cos(c+dx)}\right) \int \cos^2(c+dx) dx}{4\sqrt{\cos(c+dx)}} \\
 &= \frac{3b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} \\
 &\quad + \frac{b\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}\sin(c+dx)}{4d} + \frac{\left(3b\sqrt{b\cos(c+dx)}\right) \int 1 dx}{8\sqrt{\cos(c+dx)}} \\
 &= \frac{3bx\sqrt{b\cos(c+dx)}}{8\sqrt{\cos(c+dx)}} + \frac{3b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} \\
 &\quad + \frac{b\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}\sin(c+dx)}{4d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.54

$$\begin{aligned}
 &\int \cos^{\frac{5}{2}}(c+dx)(b\cos(c \\
 &\quad + dx))^{\frac{3}{2}} dx = \frac{(b\cos(c+dx))^{\frac{3}{2}}(12(c+dx) + 8\sin(2(c+dx)) + \sin(4(c+dx)))}{32d\cos^{\frac{3}{2}}(c+dx)}
 \end{aligned}$$

`[In] Integrate[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2), x]`

`[Out] ((b*Cos[c + d*x])^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*d*Cos[c + d*x]^(3/2))`

Maple [A] (verified)

Time = 2.81 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.62

method	result
default	$\frac{b\sqrt{\cos(dx+c)}(2\sin(dx+c)(\cos^3(dx+c))+3\cos(dx+c)\sin(dx+c)+3dx+3c)}{8d\sqrt{\cos(dx+c)}}$
risch	$\frac{3b\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})e^{i(dx+c)}x}{4(e^{2i(dx+c)}+1)} - \frac{ib\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})e^{5i(dx+c)}}{32(e^{2i(dx+c)}+1)d} + \frac{ib\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})e^{-i(dx+c)}}{4(e^{2i(dx+c)}+1)d}$

[In] `int(cos(d*x+c)^(5/2)*(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`[Out]
$$\frac{1}{8}d*b*(\cos(d*x+c)*b)^{(1/2)}*(2*\sin(d*x+c)*\cos(d*x+c)^3+3*\cos(d*x+c)*\sin(d*x+c)+3*d*x+3*c)/\cos(d*x+c)^{(1/2)}$$
Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.81

$$\int \cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{3}{2}} dx = \left[\frac{2(2b\cos(dx+c)^2+3b)\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c)+3\sqrt{-bb}\log(2b\cos(dx+c))}{16d} \right]$$

[In] `integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(3/2),x,algorithm="fricas")`[Out]
$$\frac{1}{16}*(2*(2*b*\cos(d*x+c)^2+3*b)*\sqrt{b*\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)+3*\sqrt{-b}*b*\log(2*b*\cos(d*x+c)^2-2*\sqrt{b*\cos(d*x+c)}*\sqrt{-b}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-b))/d, \frac{1}{8}*((2*b*\cos(d*x+c)^2+3*b)*\sqrt{b*\cos(d*x+c)}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)+3*b^{(3/2)}*\arctan(\sqrt{b*\cos(d*x+c)}*\sin(d*x+c)/(\sqrt{b*\cos(d*x+c)}^{(3/2)})))/d]$$
Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{3}{2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.52

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(12(dx + c)b + b \sin(4dx + 4c) + 8b \sin(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c)))) \sqrt{b}}{32d}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/32*(12*(d*x + c)*b + b*sin(4*d*x + 4*c) + 8*b*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(83) = 166.

Time = 2.43 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.02

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(3\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 12\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 - 10\sqrt{b} \tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 18\sqrt{b}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^5 - 6\sqrt{b} \tan(\frac{1}{2}dx + \frac{1}{2}c)^3 + 3\sqrt{b}dx + 10\sqrt{b} \tan(\frac{1}{2}dx + \frac{1}{2}c)) * b / (d \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 4d \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 + 6d \tan(\frac{1}{2}dx + \frac{1}{2}c)^4 + 4d \tan(\frac{1}{2}dx + \frac{1}{2}c)^2 + d)}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] 1/8*(3*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^8 + 12*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^6 - 10*sqrt(b)*tan(1/2*d*x + 1/2*c)^7 + 18*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^4 + 6*sqrt(b)*tan(1/2*d*x + 1/2*c)^5 + 12*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^2 - 6*sqrt(b)*tan(1/2*d*x + 1/2*c)^3 + 3*sqrt(b)*d*x + 10*sqrt(b)*tan(1/2*d*x + 1/2*c))*b/(d*tan(1/2*d*x + 1/2*c)^8 + 4*d*tan(1/2*d*x + 1/2*c)^6 + 6*d*tan(1/2*d*x + 1/2*c)^4 + 4*d*tan(1/2*d*x + 1/2*c)^2 + d)

Mupad [B] (verification not implemented)

Time = 14.93 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.75

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24 dx)}{32d (\cos(2c + 2dx) + 1)}$$

```
[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(3/2),x)
```

```
[Out] (b*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c +  
3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x) +  
1))
```

3.151 $\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{3}{2}} dx$

Optimal result	772
Rubi [A] (verified)	772
Mathematica [A] (verified)	773
Maple [A] (verified)	773
Fricas [A] (verification not implemented)	774
Sympy [F(-1)]	774
Maxima [A] (verification not implemented)	774
Giac [B] (verification not implemented)	775
Mupad [B] (verification not implemented)	822

Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{3}{2}} dx = \frac{b\sqrt{b \cos(c + dx)} \sin(c + dx)}{d\sqrt{\cos(c + dx)}} - \frac{b\sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d\sqrt{\cos(c + dx)}}$$

[Out] $b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}-1/3*b*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{3}{2}} dx = \frac{b \sin(c + dx) \sqrt{b \cos(c + dx)}}{d\sqrt{\cos(c + dx)}} - \frac{b \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d\sqrt{\cos(c + dx)}}$$

[In] Int[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2),x]

[Out] $(b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) - (b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2713

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \cos^3(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{\left(b\sqrt{b\cos(c+dx)}\right) \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{d\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}} - \frac{b\sqrt{b\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{\frac{3}{2}} dx = \frac{(b\cos(c+dx))^{\frac{3}{2}}(5+\cos(2(c+dx)))\sin(c+dx)}{6d\cos^{\frac{3}{2}}(c+dx)}$$

[In] Integrate[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2), x]

[Out] ((b*Cos[c + d*x])^(3/2)*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 2.79 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.57

method	result
default	$\frac{b(2+\cos^2(dx+c))\sqrt{\cos(dx+c)}b\sin(dx+c)}{3d\sqrt{\cos(dx+c)}}$
risch	$-\frac{ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})e^{4i(dx+c)}}{12(e^{2i(dx+c)}+1)d} - \frac{3ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})e^{2i(dx+c)}}{4(e^{2i(dx+c)}+1)d} + \frac{3ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})}{4(e^{2i(dx+c)}+1)d} + \frac{ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})}{4(e^{2i(dx+c)}+1)d}$

[In] int(cos(d*x+c)^(3/2)*(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/3/d*b*(2+cos(d*x+c)^2)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.60

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(b \cos(dx + c)^2 + 2b) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3 d \sqrt{\cos(dx + c)}}$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/3*(b*cos(d*x + c)^2 + 2*b)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{3/2} dx = \frac{(b \sin(3 dx + 3 c) + 9 b \sin(\frac{1}{3} \arctan(\sin(3 dx + 3 c), \cos(3 dx + 3 c)))) \sqrt{b}}{12 d}$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/12*(b*sin(3*d*x + 3*c) + 9*b*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71048 vs. 2(62) = 124.

Time = 11.24 (sec) , antiderivative size = 71048, normalized size of antiderivative = 986.78

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{3}{2}} dx = \text{Too large to display}$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out]
$$\begin{aligned} & -1/96*(3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \end{aligned}$$

$$\begin{aligned}
&)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 \tan(c)^2 + \\
& 27 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x \\
& + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 - 54 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \\
& * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 + 3 * s \\
& qrt(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan \\
& (c)^2 + 48 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
& -1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan(c)^2 - 54 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/ \\
& 2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^6 \tan(c)^2 \\
& + 108 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2 \\
& * d*x + 1/2*c)^2 * \tan(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2 \\
& * c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 \\
& + 48 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x \\
& + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 9 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 * \\
& \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 54 * s \\
& qrt(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1 \\
& /2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 3 * sqrt(b) * d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
& -1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 9 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1 \\
& /2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 54 * sq \\
& rt(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
& 2*c)^2 * \tan(1/3*c)^4 + 144 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x \\
& + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 27 \sqrt{b} * d*x^4 \tan(1/2* \\
& d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \\
& + 54 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d \\
& * x + 1/2*c)^4 \tan(1/3*c)^4 - 3 * sqrt(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2 \\
& * d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(\\
& 1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 + 54 \sqrt{b} * d*x^4 * ta \\
& n(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3 \\
& * c)^6 - 108 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 \tan \\
& (-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^6 + 144 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^ \\
& 3 * \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 - 48 \sqrt{b} * \\
& d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 * t \\
& an(1/3*c)^6 - 9 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \\
& * \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 54 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2* \\
& c)^2 * \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 3 * sqrt(b \\
&) * d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 72 * sq \\
& rt(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
& 2*c)^3 * \tan(1/3*c)^2 * \tan(c) - 72 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 * \tan(1/ \\
& 2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 * \tan(c) + 72 \sqrt{b} * d \\
& * x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * ta \\
& n(1/3*c)^4 * \tan(c) + 432 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + \\
& 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) - 216 \sqrt{b} * d*x^4 * ta \\
& n(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3 \\
& * c)^4 * \tan(c) + 432 \sqrt{b} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c \\
&)^6 \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) - 216 \sqrt{b} * d*x^4 \tan(1/2 \\
& * d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^6*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 144* \\
& \sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 3*\sqrt{b}*d*x^4 \\
& *4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 54 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - \\
& 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^3*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 2 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c \\
&)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 324*\sqrt{b}*d*x \\
& ^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 27*\sqrt{b} \\
& (b)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 96*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*\sqrt{ \\
& b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 18 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 - \\
& 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 \\
& - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*\sqrt{b}*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan \\
& (1/3*c)^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c
\end{aligned}$$

$$\begin{aligned}
&)^3 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^4 \tan(c)^2 + \\
&192 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/ \\
&2 c)^2 \tan(1/3 c)^5 \tan(c)^2 - 1440 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 * \\
&dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^5 \tan(c)^2 + 1728 \sqrt{b} \\
&* \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^4 \tan(\\
&1/3 c)^5 \tan(c)^2 - 384 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c) \\
&^6 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^5 \tan(c)^2 + 9 \sqrt{b} dx^4 \tan(1/2 * \\
&dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^2 \tan(1/3 c)^6 \tan(c)^2 - 54 \sqrt{b} dx \\
&x^4 \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^4 \tan(1/3 c)^6 \tan(c)^2 + 3 \\
&* \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^6 \tan(1/3 c)^6 \tan(c)^2 + 144 \sqrt{b} dx \\
&* x^4 \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c) * ta \\
&n(1/3 c)^6 \tan(c)^2 - 144 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c) \tan(1/2 dx + \\
&1/6 c)^4 \tan(-1/2 dx + 1/2 c) \tan(1/3 c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(1/2 dx \\
&* dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c) \tan(1/3 c)^6 \tan \\
&(c)^2 - 18 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^4 \tan(-1/2 dx + 1/2 c)^2 \tan \\
&(1/3 c)^6 \tan(c)^2 + 324 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + \\
&1/6 c)^2 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 - 54 \sqrt{b} dx^4 \\
&\tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 + 192 * \\
&\sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c) \\
&^2 \tan(1/3 c)^6 \tan(c)^2 - 288 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + \\
&1/6 c)^6 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 - 48 \sqrt{b} dx^4 \\
&\tan(1/2 dx + 1/2 c)^3 \tan(-1/2 dx + 1/2 c)^3 \tan(1/3 c)^6 \tan(c)^2 + 144 * \\
&\sqrt{b} dx^4 \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/ \\
&2 c)^3 \tan(1/3 c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx \\
&x + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^3 \tan(1/3 c)^6 \tan(c)^2 - 48 \sqrt{b} \tan \\
&(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^3 \tan(1/3 * \\
&c)^6 \tan(c)^2 - 18 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^2 \tan(-1/2 dx + 1/2 * \\
&c)^4 \tan(1/3 c)^6 \tan(c)^2 + 9 \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/ \\
&2 dx + 1/2 c)^4 \tan(1/3 c)^6 \tan(c)^2 - 320 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \\
&* \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^6 \tan(c)^2 + 432 \\
&* \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c) \\
&)^4 \tan(1/3 c)^6 \tan(c)^2 - 384 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx \\
&+ 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1 \\
&/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c)^6 \\
&* \tan(c)^2 + 18 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^6 * \\
&\tan(-1/2 dx + 1/2 c)^2 + 48 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx \\
&* dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^3 - 9 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c) \\
&)^4 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^4 + 18 \sqrt{b} dx^4 \tan(1 \\
&/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^4 - 9 \sqrt{b} (b \\
&)* dx^4 \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^6 \tan(1/3 c)^2 - 144 * \sqrt{ \\
&rt(b) dx^4 \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/ \\
&2 c) \tan(1/3 c)^2 + 162 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + \\
&1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^2 - 324 \sqrt{b} dx^4 \tan(1/2 dx \\
&* dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^2 + \\
&432 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx
\end{aligned}$$

$$\begin{aligned}
& *x + 1/2*c)^3*\tan(1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2* \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*t \\
& an(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 27*\sqrt{b}*d*x^ \\
& 4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 54*\sqrt{b}*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 - 432*\sqrt{b} \\
& (b)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)*\tan(1/3*c)^4 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 972*\sqrt{ \\
& rt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c) \\
& ^4 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^3*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 27*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1440*\sqrt{b}*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^4 + 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1440*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + \\
& 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^5 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 3*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^6 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c) \\
& ^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^6 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 54*\sqrt{b}*d*x^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6 - 288*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan
\end{aligned}$$

$$\begin{aligned}
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2 \\
& *c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9*\sqrt{b} \\
&)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^ \\
& 2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6 \\
& *c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2* \\
& \tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 640*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 4 \\
& 800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 1280*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)^3*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b} \\
&)*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1 \\
& /6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2* \\
& c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
&)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + \\
& 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x \\
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) \\
& ^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) \\
& ^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^ \\
& 3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan \\
& (c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 96*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
& \text{rt}(b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 - \\
& 2880 \cdot \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(-1/2*d*x + \\
& 1/2*c)^2 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 + 3456 \cdot \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^3 \cdot \tan(1/ \\
& 2*d*x + 1/6*c)^5 \cdot \tan(-1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 - 768 \cdot \text{sqrt}(b) \\
&) \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1/2*c)^2 \cdot \tan \\
& (1/3*c)^5 \cdot \tan(c)^2 + 1440 \cdot \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6* \\
& c)^2 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 - 5760 \cdot \text{sqrt}(b) \cdot \tan(1/2*d \\
& *x + 1/2*c)^3 \cdot \tan(1/2*d*x + 1/6*c)^3 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^5 \cdot \tan \\
& an(c)^2 + 5760 \cdot \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(-1 \\
& /2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 - 1728 \cdot \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c) \\
& * \tan(1/2*d*x + 1/6*c)^5 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^5 \cdot \tan(c)^2 + 96 * \\
& \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^5 \cdot \tan(c)^ \\
& 2 + 3 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 54 * \text{sqrt}(\\
& b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^ \\
& 2 + 9 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 + 96 * \text{sqrt}(\\
& b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^5 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 14 \\
& 4 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^3 \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(1/3*c)^6 \cdot \tan(c) \\
& ^2 + 48 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^3 \cdot \tan(-1/2*d*x + 1/2*c) \cdot \tan(1/3 * \\
& c)^6 \cdot \tan(c)^2 - 144 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c) \cdot \tan(1/2*d*x + 1/6*c) \\
& ^2 \cdot \tan(-1/2*d*x + 1/2*c) \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 72 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1 \\
& /2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(-1/2*d*x + 1/2*c) \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 \\
& - 48 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1 \\
& /2*c) \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 + 108 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(\\
& -1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 54 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + \\
& 1/6*c)^2 \cdot \tan(-1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 640 * \text{sqrt}(b) \cdot \tan(1/ \\
& 2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^3 \cdot \tan(-1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^ \\
& 6 \cdot \tan(c)^2 + 864 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^3 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(\\
& -1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 768 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c) \\
&)^2 \cdot \tan(1/2*d*x + 1/6*c)^5 \cdot \tan(-1/2*d*x + 1/2*c)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - \\
& 96 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c) \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1/2*c) \\
&)^2 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 + 48 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c) \cdot \tan(-1/2 * \\
& d*x + 1/2*c)^3 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 72 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan \\
& n(1/2*d*x + 1/6*c)^2 \cdot \tan(-1/2*d*x + 1/2*c)^3 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 144 * \text{sq} \\
& rt(b) \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^4 \cdot \tan(-1/2*d*x + 1/2*c)^3 \\
& * \tan(1/3*c)^6 \cdot \tan(c)^2 - 24 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1 \\
& /2*c)^3 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 + 3 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan \\
& (1/3*c)^6 \cdot \tan(c)^2 + 96 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c) \\
& * \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 1008 * \text{sqrt}(b) \cdot \tan(1/2*d*x + \\
& 1/2*c)^3 \cdot \tan(1/2*d*x + 1/6*c)^2 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c) \\
&)^2 + 1280 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c)^2 \cdot \tan(1/2*d*x + 1/6*c)^3 \cdot \tan(-1/2*d \\
& *x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - 1008 * \text{sqrt}(b) \cdot \tan(1/2*d*x + 1/2*c) \cdot \tan \\
& (1/2*d*x + 1/6*c)^4 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 + 96 * \text{sqrt} \\
& (b) \cdot \tan(1/2*d*x + 1/6*c)^5 \cdot \tan(-1/2*d*x + 1/2*c)^4 \cdot \tan(1/3*c)^6 \cdot \tan(c)^2 - \\
& 3 * \text{sqrt}(b) \cdot d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^4 \cdot \tan(1/2*d*x + 1/6*c)^6 - 48 * \text{sqrt}(b) * \\
& d*x^4 \cdot \tan(1/2*d*x + 1/2*c)^3 \cdot \tan(1/2*d*x + 1/6*c)^6 \cdot \tan(-1/2*d*x + 1/2*c) +
\end{aligned}$$

$$\begin{aligned}
& *c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 144 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 43 \\
& 2 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + \\
& 1/2*c)^3 \tan(1/3*c)^4 + 216 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/ \\
& 6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 144 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 54 * \text{sqrt} \\
& (b) * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 27 * \\
& \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + \\
& 4800 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + \\
& 1/2*c)^4 \tan(1/3*c)^4 - 11664 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
& 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 5760 * \text{sqrt}(b) * \tan(1/2*d*x + \\
& 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 1008 \\
& * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^ \\
& 4 \tan(1/3*c)^4 + 96 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan \\
& (1/3*c)^5 - 2880 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan \\
& (-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 + 3456 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^3 \tan \\
& (1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 - 768 * \text{sqrt}(b) * \tan \\
& (1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3* \\
& c)^5 + 1440 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2* \\
& d*x + 1/2*c)^4 \tan(1/3*c)^5 - 5760 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d \\
& *x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 + 5760 * \text{sqrt}(b) * \tan(1/2*d \\
& *x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 - \\
& 1728 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/ \\
& 2*c)^4 \tan(1/3*c)^5 + 96 * \text{sqrt}(b) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2* \\
& c)^4 \tan(1/3*c)^5 - 3 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 5 \\
& 4 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 \\
& - 9 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 + 96 * \text{sqrt}(b) * \tan(1/2* \\
& d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(1/3*c)^6 - 144 * \text{sqrt}(b) * \tan(1/2*d* \\
& x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48 * \text{sqrt}(b) * d*x^4 \tan(1/2 \\
& *d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 + 144 * \text{sqrt}(b) * d*x^4 \tan(\\
& 1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \\
& + 72 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1 \\
& /2*c) \tan(1/3*c)^6 + 48 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \\
& ^6 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 - 108 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 54 * \text{sqrt}(b) * d*x^4 \tan(1/2*d*x + \\
& 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 640 * \text{sqrt}(b) * \tan(1/2*d*x + \\
& 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 864 * \\
& \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) \\
& ^2 \tan(1/3*c)^6 - 768 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \\
& * \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 96 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c) \tan \\
& (1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 48 * \text{sqrt}(b) * d*x^4 \\
& * \tan(1/2*d*x + 1/2*c) \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 + 72 * \text{sqrt}(b) * \tan \\
& (1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3* \\
& c)^6 + 144 * \text{sqrt}(b) * \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d \\
& *x + 1/2*c)^3 \tan(1/3*c)^6 + 24 * \text{sqrt}(b) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/3*c)^6 - 3*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) \\
& - 1296 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) \\
& - 216 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) \\
& + 216 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) \\
& + 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) \\
& - 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^6 * \tan(c) - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) \\
& + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^6 * \tan(c) - 24 * \sqrt{b} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) \\
& + 24 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) \\
& + 9 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(c)^2 - 18 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(c)^2 + 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 324 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 18 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 24 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 3 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 9 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 192 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) * \tan(c)^2 - 1440 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 + 1728 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 - 384 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(
\end{aligned}$$

$$\begin{aligned}
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 27*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c \\
&)^2*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c \\
&)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*s \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)* \\
& \tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c \\
&)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 1 \\
& 62*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2 \\
& *c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3* \\
& c)^2*\tan(c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(\\
& c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*t \\
& \tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*t \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 \\
& - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 4800*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*t \\
& \tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 144*\sqrt{b}*d*x^4* \\
& \tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432* \\
& \sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/ \\
& 2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x \\
& + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 \\
& *\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^4*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4 \\
& *\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 233 \\
& 28*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2016*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^4*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2 \\
& *c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3 \\
& *c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3* \\
& \tan(1/3*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(- \\
& 1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c \\
&)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - \\
& 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2* \\
& d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b} \\
& *\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 144 \\
& 0*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c) \\
& ^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^ \\
& 5*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& 1/3*c)^5*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan \\
& (c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c \\
&)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 19 \\
& 2*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c \\
&)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 5*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(- \\
& 1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c \\
&)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + \\
& 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^5*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^6*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*
\end{aligned}$$

$$\begin{aligned}
& c)^6 \tan(c)^2 - 320 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 432 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 384 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 \tan(c)^2 - 18 \sqrt{b} *d*x^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c) \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 2016 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 2560 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 2016 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 144 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 384 \sqrt{b} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 432 \sqrt{b} \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 320 \sqrt{b} \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 9 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 + 18 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 - 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) + 48 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) + 54 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 - 324 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 + 18 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 + 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 - 144 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 + 24 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 - 3 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^4 + 54 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 - 9 \sqrt{b} *d*x^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 - 96 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 - 48 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 + 192 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c) - 1440 \sqrt{b} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c) + 1728 \sqrt{b} \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^5 \tan(-1
\end{aligned}$$

$$\begin{aligned}
& /2*d*x + 1/2*c)^4*\tan(1/3*c) - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 27*\sqrt{b}*d*x^4*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 162*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 - 9*\sqrt{b}*d*x^4*t \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 432*\sqrt{b}* \\
& d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan \\
& (1/3*c)^2 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 972*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 162*\sqrt{b}* \\
& d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*\sqrt{ \\
& rt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\sqrt{ \\
& rt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^ \\
& 3*\tan(1/3*c)^2 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9936* \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 + 9600*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^3 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2 \\
& *d*x + 1/2*c)^2*\tan(1/3*c)^3 + 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 4800*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 \\
& + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^3 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + 5760*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 320* \\
& \sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 9*\sqrt{ \\
& t{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 - 27*\sqrt{b}*d*x^4*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^5*\tan(1/3*c)^4 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& -1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/
\end{aligned}$$

$$\begin{aligned}
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 \\
& + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 23328*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 144*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 4 \\
& 32*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^3*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c \\
&)^3*\tan(1/3*c)^4 - 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1 \\
& 440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(1/3*c)^4 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9936*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (1/3*c)^5 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(\\
& 1/3*c)^5 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/ \\
& 3*c)^5 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
& ^5 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^5 + 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^5 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^5 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/ \\
& 6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + 5760*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^5 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^5 + 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 9*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 - 320*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan \\
& (-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 +
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c) - 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 3*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 18*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 \\
& + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 1440*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 \\
& - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^2*\tan(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 324*s \\
& qrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*ta \\
& n(c)^2 - 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2* \\
& an(1/3*c)^2*\tan(c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/ \\
& 6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*ta \\
& n(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2 \\
& *c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + \\
& 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(- \\
& 1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 7 \\
& 2*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c \\
&)^2 + 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440* \\
& sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 19200*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4* \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
&)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 + 128 \\
& 0*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c) \\
& ^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*t \\
& an(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400 \\
& *\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x \\
& + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 640*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 320*\sqrt{ \\
& b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - \\
& 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^3*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^3*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3 \\
& *c)^4*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c \\
&)^2 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c) \\
& ^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& *\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - \\
& 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^4*\tan(c)^2 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4*\tan(c)^2 - 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 \\
& - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4* \\
& tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2 \\
& *\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + \\
& 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 1008*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{ \\
& b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 1440*\sqrt{b}*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 5760*s \\
& qrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2
\end{aligned}$$

$$\begin{aligned}
& + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^5\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^5\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^5\tan(c)^2 - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 3456\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 - 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 - 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 + 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5\tan(c)^2 + 3\sqrt{b}*d*x^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(1/3*c)^6\tan(c)^2 - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^6\tan(c)^2 + 1280\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^6\tan(c)^2 - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^6\tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6\tan(c)^2 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 768\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 + 864\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 640\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)^2 - 48\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^6\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^6\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^6\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^6\tan(c)^2 - 9\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2 + 54\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4 - 3\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^6 - 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c) + 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c) + 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c) + 18\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2 - 324\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2 + 54\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2 - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^5\tan(-1/2*d*x + 1/2*c)^2 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c)^2 + 48\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^3 - 144\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3 + 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3 + 18*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4 - 9*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 320*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3 \\
& *\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4 - 144*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 96*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c) - 2880*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c) + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 1440*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 5760*\sqrt{b} \\
& *\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c) + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1 \\
& /2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) + 96*\sqrt{b}*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 9*\sqrt{b}*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 - 27*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan \\
& (1/3*c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)* \\
& \tan(1/3*c)^2 + 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^ \\
& 2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 144*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^ \\
& 2 - 324*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^2 + 162*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 19872*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 11520*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^2 - 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 432*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - \\
& 9*\sqrt{b}*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - \\
& 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 11664*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440 \\
& *sqrt(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 4800 \\
& *sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 5760* \\
& sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3 + 1280*s \\
& qrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 - 9600*sq \\
& rt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^3 + 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3 \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*sqrt(\\
& b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^3 - 640*sqrt(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^3 + 320*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^3 - 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^3 + 19200*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 19200*sqrt(b)*\tan(1/2 \\
& *d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 4800*sqrt(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 54*sqrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 27*sqrt(b)*d*x^4*ta \\
& n(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 - 11664*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4 - 1008*sqrt(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(1/3*c)^4 + 144*sqrt(b)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*sqrt(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 72* \\
& sqrt(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 54*sqrt \\
& (b)*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*sqrt(b)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872 \\
& *sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^4 - 38400*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c \\
&)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872*sqrt(b)*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*sqrt(\\
& b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*sqrt(b) \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 432*sqrt(b)* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1 \\
& /3*c)^4 + 216*sqrt(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4 - 1008*sqrt(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 5760*sqrt(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^4 - 11664*sqrt(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 4800*sqrt(b)*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 1440*sqrt(b)*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 5760*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5 + 5760*sqrt(b)*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5 - 1728*sqrt(b)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^5 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^5 - 192*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 3*\sqrt{b}*d*x^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 640*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 432*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) + 144*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 24*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 216*\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/3*c)^2 * \tan(c) - 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x \\
& x + 1/2*c) * \tan(1/3*c)^2 * \tan(c) + 216 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan \\
& \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c) - 432 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + \\
& 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^2 * \tan(c) - 72 * \sqrt{b} * d*x^4 * \tan(- \\
& 1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^2 * \tan(c) + 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \\
& \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2 \\
& *c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + \\
& 216 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan \\
& (c) + 72 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) - 216 * \sqrt{b} (\\
& b) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^4 * \tan(c) - 432 * \\
& \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^4 * \tan(c) - \\
& 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^4 * \tan(c) + 72 * \sqrt{b} * d*x^4 * \tan \\
& \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) \\
& ^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 216 * \sqrt{b} * \tan(1/2*d*x + \\
& 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24 * \sqrt{b} * \tan(1/2*d \\
& *x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) - 144 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(\\
& 1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) - 72 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^4 * \\
& \tan(1/3*c)^6 * \tan(c) + 24 * \sqrt{b} * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c \\
&) + 3 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54 * \sqrt{b} * d*x^4 * \tan(\\
& 1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(c)^2 + 9 * \sqrt{b} * d*x^4 * \tan(1/ \\
& 2*d*x + 1/6*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + \\
& 1/6*c)^5 * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c) \\
& ^6 * \tan(c)^2 + 48 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c) \\
& * \tan(c)^2 - 144 * \sqrt{b} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan \\
& \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 72 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d \\
& *x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48 * \sqrt{b} * \tan(1/2*d*x + 1/2 \\
& *c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 + 108 * \sqrt{b} * d \\
& *x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 54 * \sqrt{b} * d \\
& *x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 640 * \sqrt{b} * \\
& \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c \\
&)^2 - 2016 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d \\
& *x + 1/2*c)^2 * \tan(c)^2 + 768 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1 \\
& /6*c)^5 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 288 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) \\
& * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 48 * \sqrt{b} * d*x^4 \\
& * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 72 * \sqrt{b} * \tan(1/2 \\
& *d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 1 \\
& 44 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2 \\
& *c)^3 * \tan(c)^2 - 24 * \sqrt{b} * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \\
& \tan(c)^2 + 3 * \sqrt{b} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan \\
& \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 \\
& + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + \\
& 1/2*c)^4 * \tan(c)^2 - 1280 * \sqrt{b} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c \\
&)^3 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 432 * \sqrt{b} * \tan(1/2*d*x + 1/2*c) * \tan \\
& (1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96 * \sqrt{b} * \tan(1/2*d \\
& *x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 1440 * \sqrt{b} * \tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 + 2880*\sqrt{b} \\
& * \tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6 \\
& *c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan \\
& (c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4 \\
& *\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - \\
& 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x \\
& + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 1440*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 54*\sqrt{b} \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*\sqrt{b}*d*x^4*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c \\
&)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 + 432*\sqrt{b} \\
& * \tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 - 144 \\
& *\sqrt{b}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(\\
& c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b} \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 54*sq \\
& rt(b)*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^2*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c \\
&)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan \\
& (c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3 \\
& *c)^2*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan \\
& (c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^2*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 640*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*\sqrt{b}*d*x^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 23328*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2 - 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 96*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 320*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c
\end{aligned}$$

$$\begin{aligned}
&)^6 \tan(c)^2 - 48 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(-1/2 dx + 1/2 c) \tan(1/3 c)^6 \tan(c)^2 - 72 \sqrt{b} \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c) \\
& * \tan(1/3 c)^6 \tan(c)^2 - 288 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 + 192 \sqrt{b} \tan(1/2 dx + 1/6 c) \tan(-1/2 dx \\
& x + 1/2 c)^2 \tan(1/3 c)^6 \tan(c)^2 - 24 \sqrt{b} \tan(-1/2 dx + 1/2 c)^3 \tan(1/3 c)^6 \tan(c)^2 - 3 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^4 + 54 \sqrt{b} dx \\
& x^4 \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^2 - 9 \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^4 - 96 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^5 \\
& - 48 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^6 - 48 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^3 \tan(-1/2 dx + 1/2 c) + 144 \sqrt{b} dx^4 \tan(1/2 \\
& dx + 1/2 c) \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c) + 72 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c) + 48 \sqrt{b} \\
& (b) \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c) - 108 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c)^2 \tan(-1/2 dx + 1/2 c)^2 + 54 \sqrt{b} \\
& dx^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^2 + 640 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx + 1/2 c)^2 - 2016 \sqrt{b} \\
& (b) \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 + 768 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/ \\
& 2 c)^2 - 288 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx \\
& * x + 1/2 c)^2 - 48 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 c) \tan(-1/2 dx + 1/2 c) \\
& ^3 + 72 \sqrt{b} \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx \\
& + 1/2 c)^3 + 144 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^4 \tan(\\
& -1/2 dx + 1/2 c)^3 + 24 \sqrt{b} \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 \\
& c)^3 - 3 \sqrt{b} dx^4 \tan(-1/2 dx + 1/2 c)^4 - 96 \sqrt{b} \tan(1/2 dx + 1 \\
& /2 c)^4 \tan(1/2 dx + 1/6 c) \tan(-1/2 dx + 1/2 c)^4 + 432 \sqrt{b} \tan(1/2 \\
& dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^4 - 1280 \sqrt{b} \\
&) \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx + 1/2 c)^4 + 4 \\
& 32 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 c)^4 \tan(-1/2 dx + 1/2 c \\
&)^4 - 96 \sqrt{b} \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^4 - 1440 \sqrt{b} \\
& (b) \tan(1/2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^4 \tan(1/3 c) + 1728 \sqrt{b} \\
& * \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^5 \tan(1/3 c) - 384 \sqrt{b} \tan \\
& (1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^6 \tan(1/3 c) + 2880 \sqrt{b} \tan(1/ \\
& 2 dx + 1/2 c)^4 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) \\
& - 11520 \sqrt{b} \tan(1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c)^3 \tan(-1/2 dx \\
& + 1/2 c)^2 \tan(1/3 c) + 11520 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + \\
& 1/6 c)^4 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) - 3456 \sqrt{b} \tan(1/2 dx + 1/ \\
& 2 c) \tan(1/2 dx + 1/6 c)^5 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) + 192 \sqrt{b} \\
&) \tan(1/2 dx + 1/6 c)^6 \tan(-1/2 dx + 1/2 c)^2 \tan(1/3 c) - 96 \sqrt{b} \tan \\
& n(1/2 dx + 1/2 c)^4 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) + 1728 \sqrt{b} \tan(\\
& 1/2 dx + 1/2 c)^3 \tan(1/2 dx + 1/6 c) \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) \\
& - 5760 \sqrt{b} \tan(1/2 dx + 1/2 c)^2 \tan(1/2 dx + 1/6 c)^2 \tan(-1/2 dx + \\
& 1/2 c)^4 \tan(1/3 c) + 5760 \sqrt{b} \tan(1/2 dx + 1/2 c) \tan(1/2 dx + 1/6 \\
& c)^3 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) - 1440 \sqrt{b} \tan(1/2 dx + 1/6 c) \\
& ^4 \tan(-1/2 dx + 1/2 c)^4 \tan(1/3 c) + 54 \sqrt{b} dx^4 \tan(1/2 dx + 1/2 \\
& c)^2 \tan(1/3 c)^2 - 27 \sqrt{b} dx^4 \tan(1/2 dx + 1/6 c)^2 \tan(1/3 c)^2 -
\end{aligned}$$

$$\begin{aligned}
& 4800\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^2 + 9 \\
& 936\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^2 - 57 \\
& 60\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^2 + 432 \\
& \sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^2 + 144\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^2 + 216\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan \\
& \tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan \\
& (-1/2*d*x + 1/2*c)\tan(1/3*c)^2 + 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(-1/ \\
& 2*d*x + 1/2*c)\tan(1/3*c)^2 + 54\sqrt{b}*d*x^4\tan(-1/2*d*x + 1/2*c)^2\tan(\\
& 1/3*c)^2 + 2880\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(-1/ \\
& 2*d*x + 1/2*c)^2\tan(1/3*c)^2 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/ \\
& 2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^2 + 38400\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^2 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x \\
& + 1/2*c)^2\tan(1/3*c)^2 + 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(-1/2*d*x \\
& + 1/2*c)^2\tan(1/3*c)^2 + 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + \\
& 1/2*c)^3\tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1 \\
& /6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^2 + 216\sqrt{b}\tan(1/2*d*x + 1/ \\
& 6*c)^4\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^2 + 432\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2 - 5760\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2 + 9936\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan \\
& (1/3*c)^2 - 4800\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan \\
& (1/3*c)^2 - 4800\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(\\
& 1/3*c)^3 + 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^3\tan(\\
& 1/3*c)^3 - 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(\\
& 1/3*c)^3 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan(1/3 \\
& *c)^3 - 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^3 + 640\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 - 11520\sqrt{b}\tan(\\
& 1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^ \\
& 3 + 38400\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d* \\
& x + 1/2*c)^2\tan(1/3*c)^3 - 38400\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x \\
& + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 + 9600\sqrt{b}\tan(1/2*d*x \\
& + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^3 - 1280\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 + 5760\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 - 4800\sqrt{b} \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^3 - 9\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + \\
& 1/6*c)\tan(1/3*c)^4 + 9936\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6 \\
& *c)^2\tan(1/3*c)^4 - 19200\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6 \\
& *c)^3\tan(1/3*c)^4 + 9936\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c) \\
& ^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^4 + 72\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^4 + 432\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1 \\
& /3*c)^4 + 216\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*
\end{aligned}$$

$$\begin{aligned}
& c)^4 - 2016\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3* \\
& c)^4 + 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d \\
& *x + 1/2*c)^2\tan(1/3*c)^4 - 23328\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x \\
& + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^4 + 9600\sqrt{b}\tan(1/2*d*x \\
& + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^4 + 144\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^4 + 216\sqrt{b}\tan(1/2*d*x + \\
& 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(1/3*c)^4 + 432\sqrt{b}\tan(1/2*d*x + \\
& 1/2*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 1440\sqrt{b}\tan(1/2*d*x + 1/ \\
& 6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^4 - 96\sqrt{b}\tan(1/2*d*x + 1/2*c) \\
& ^4\tan(1/3*c)^5 + 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)* \\
& \tan(1/3*c)^5 - 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan \\
& (1/3*c)^5 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(\\
& 1/3*c)^5 - 1440\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^5 + 768\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^5 - 3456\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^5 + 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c) \\
& ^5 - 96\sqrt{b}\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^5 - 48\sqrt{b}\tan(1/2*d \\
& *x + 1/2*c)^3\tan(1/3*c)^6 - 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x \\
& + 1/6*c)\tan(1/3*c)^6 + 432\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6 \\
& *c)^2\tan(1/3*c)^6 - 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^6 + 48*s \\
& qrt(b)\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6 + 72\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^6 - 288\sqrt{b}* \\
& \tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6 + 192\sqrt{b}\tan \\
& (1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)^6 + 24\sqrt{b}\tan(-1/ \\
& 2*d*x + 1/2*c)^3\tan(1/3*c)^6 - 24\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)^3\tan \\
& (c) + 72\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(c) - \\
& 72\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^4\tan(c) - 48\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^6\tan(c) - 144\sqrt{b}*d*x^4 \\
& *tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c) + 72\sqrt{b}*d*x^4\tan \\
& (1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c) - 144\sqrt{b}*d*x^4\tan(1/ \\
& 2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)^2\tan(c) - 24\sqrt{b}*d*x^4\tan(-1/2*d \\
& *x + 1/2*c)^3\tan(c) + 24\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2 \\
& *c)^4\tan(c) + 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan \\
& (-1/2*d*x + 1/2*c)^4\tan(c) + 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d \\
& *x + 1/2*c)^4\tan(c) + 72\sqrt{b}*d*x^4\tan(1/2*d*x + 1/2*c)\tan(1/3*c)^2\tan \\
& (c) - 216\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c) \\
&)^2\tan(c) - 432\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(\\
& 1/3*c)^2\tan(c) - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)^2\tan(c) + 7 \\
& 2\sqrt{b}*d*x^4\tan(-1/2*d*x + 1/2*c)\tan(1/3*c)^2\tan(c) + 144\sqrt{b}\tan \\
& (1/2*d*x + 1/2*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2\tan(c) + 216\sqrt{b} \\
& (b)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)^2\tan(c) - 72* \\
& sqrt(b)\tan(1/2*d*x + 1/2*c)^4\tan(1/3*c)^4\tan(c) - 432\sqrt{b}\tan(1/2*d* \\
& x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^4\tan(c) - 216\sqrt{b}\tan(1 \\
& /2*d*x + 1/6*c)^4\tan(1/3*c)^4\tan(c) + 72\sqrt{b}\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^4\tan(c) - 48\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/3*c)^6\tan(c)
\end{aligned}$$

$$\begin{aligned}
& - 72\sqrt{b}\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^6\tan(c) - 18\sqrt{b}*d*x^4 \\
& * \tan(1/2*d*x + 1/2*c)^2\tan(c)^2 + 9\sqrt{b}*d*x^4\tan(1/2*d*x + 1/6*c)^2\tan \\
& \tan(c)^2 + 320\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^3\tan(c)^2 \\
& - 1008\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^4\tan(c)^2 + 3 \\
& 84\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^5\tan(c)^2 - 144\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^6\tan(c)^2 - 48\sqrt{b}*d*x^4 \\
& * \tan(1/2*d*x + 1/2*c)\tan(-1/2*d*x + 1/2*c)\tan(c)^2 - 72\sqrt{b}\tan(1/2* \\
& d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)\tan(c)^2 - 144* \\
& \sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c) \\
& * \tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(-1/2*d*x + 1/2*c)\tan(c)^2 \\
& - 18\sqrt{b}*d*x^4\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 192\sqrt{b}\tan(1/2 \\
& *d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 + 864 \\
& * \sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c \\
&)^2\tan(c)^2 - 2560\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan \\
& \tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 + 864\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2* \\
& d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 192\sqrt{b}\tan(1/2*d*x + \\
& 1/6*c)^5\tan(-1/2*d*x + 1/2*c)^2\tan(c)^2 - 24\sqrt{b}\tan(1/2*d*x + 1/2*c \\
&)^4\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 72\sqrt{b}\tan(1/2 \\
& *d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2*c)^3\tan(c)^2 - 144\sqrt{b}\tan(1/2*d*x \\
& + 1/2*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2 \\
& *c)^2\tan(1/2*d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 - 1008\sqrt{b}* \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 \\
& + 320\sqrt{b}\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^4\tan(c)^2 + 1 \\
& 440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)\tan(c) \\
& ^2 - 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)* \\
& \tan(c)^2 + 5760\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(1/2*d*x + 1/6*c)^4\tan(1 \\
& /3*c)\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^5\tan \\
& \tan(1/3*c)\tan(c)^2 + 96\sqrt{b}\tan(1/2*d*x + 1/6*c)^6\tan(1/3*c)\tan(c)^2 \\
& - 192\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)\tan \\
& (c)^2 + 3456\sqrt{b}\tan(1/2*d*x + 1/2*c)^3\tan(1/2*d*x + 1/6*c)\tan(-1/2*d \\
& *x + 1/2*c)^2\tan(1/3*c)\tan(c)^2 - 11520\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^2\tan(1/3*c)\tan(c)^2 + 11520\sqrt{b} \\
& \tan(1/2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^3\tan(-1/2*d*x + 1/2*c)^2\tan \\
& \tan(1/3*c)\tan(c)^2 - 2880\sqrt{b}\tan(1/2*d*x + 1/6*c)^4\tan(-1/2*d*x + 1/2 \\
& *c)^2\tan(1/3*c)\tan(c)^2 + 384\sqrt{b}\tan(1/2*d*x + 1/2*c)^2\tan(-1/2*d*x \\
& + 1/2*c)^4\tan(1/3*c)\tan(c)^2 - 1728\sqrt{b}\tan(1/2*d*x + 1/2*c)\tan(1/2 \\
& *d*x + 1/6*c)\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)\tan(c)^2 + 1440\sqrt{b}\tan \\
& \tan(1/2*d*x + 1/6*c)^2\tan(-1/2*d*x + 1/2*c)^4\tan(1/3*c)\tan(c)^2 + 9\sqrt{b} \\
&)*d*x^4\tan(1/3*c)^2\tan(c)^2 + 1440\sqrt{b}\tan(1/2*d*x + 1/2*c)^4\tan(1/2 \\
& *d*x + 1/6*c)\tan(1/3*c)^2\tan(c)^2 - 11664\sqrt{b}\tan(1/2*d*x + 1/2*c)^3 \\
& \tan(1/2*d*x + 1/6*c)^2\tan(1/3*c)^2\tan(c)^2 + 19200\sqrt{b}\tan(1/2*d*x + \\
& 1/2*c)^2\tan(1/2*d*x + 1/6*c)^3\tan(1/3*c)^2\tan(c)^2 - 11664\sqrt{b}\tan(1 \\
& /2*d*x + 1/2*c)\tan(1/2*d*x + 1/6*c)^4\tan(1/3*c)^2\tan(c)^2 + 1440\sqrt{b} \\
& * \tan(1/2*d*x + 1/6*c)^5\tan(1/3*c)^2\tan(c)^2 - 72\sqrt{b}\tan(1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*\sqrt{b}*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(\\
& c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^2*\tan(c)^2 - 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6 \\
& *c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*\sqrt{b}*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan \\
& (c)^2 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3 \\
& *c)^2*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3 \\
& *\tan(1/3*c)^2*\tan(c)^2 - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)*\tan(1/3*c)^3*\tan(c)^2 + 19200*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\t \\
& an(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 19200*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 + 4800*\sqrt{b}*\tan(1/2*d \\
& *x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - \\
& 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\t \\
& an(c)^2 + 320*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 1008* \\
& \sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 5760*\sqrt{b}*\tan(1/2 \\
& *d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 - 11664*\sqrt{b}*\t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 4800*\sq \\
& rt(b)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*\sqrt{b}*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 + 864*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880 \\
& *\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 \\
& - 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 384*\sqrt{b}*\t \\
& an(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/ \\
& 2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x \\
& + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^5*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 \\
& + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 - 24*\sqrt{b}*\tan(-1 \\
& /2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 + 18*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)^2 - 9*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^3 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 2*d*x + 1/6*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 5 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6 + 48*\sqrt{b}*\tan \\
& x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c) + 72*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) + 144*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) + 24*\sqrt{b}*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) + 18*\sqrt{b}*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^2 - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 - 2560*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2 + 24*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3 + 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3 + 72*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 - 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4 + 384*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4 - 1008*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 320*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) - 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c) + 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c) - 1728*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c) + 96*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c) - 192*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 3456*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 2880*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 384*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 1728*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 9*\text{sqrt}(b)*d*x^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^2 - 11664*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 19200*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2 - 11664*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2 + 72*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 432*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 864*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 11520*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 19872*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 9600*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 1008*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440*\text{sqrt}(b)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 320*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 5760*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3 + 19200*\text{sqrt}(b)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3 - 19200*\text{sqrt}(b)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 2560*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 9600*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 320*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 - 11664*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 + 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 864*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 24*\sqrt{b}*\tan(1/3*c)^6*\tan(c) + 3*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(c)^2 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^5*\tan(c)^2 - 24*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 288*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 768*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 2016*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 640*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*\sqrt{b}
\end{aligned}$$

$$\begin{aligned}
& c)^3 \tan(1/3*c)^2 - 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) \\
&)*\tan(1/3*c)^2 + 9936*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*t \\
& \tan(1/3*c)^2 - 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2 + 144*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*\sqrt{b}*t \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 - 2016*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*\sqrt{b}*\tan(1 \\
& /2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 72*\sqrt{b}*\tan(-1/2* \\
& d*x + 1/2*c)^3*\tan(1/3*c)^2 - 1280*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^3 + 5760*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3 - \\
& 4800*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3 + 640*\sqrt{b}*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3 + 432*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^4 - \\
& 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 + 72*\sqrt{b}*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 96*\sqrt{b}*\tan(1/3*c)^5 - 24*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^4*\tan(c) + 24*\sqrt{b}*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) \\
& - 216*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 72*\sqrt{b}*\tan(\\
& 1/3*c)^4*\tan(c) - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 1008*\sqrt{b}*\tan(1/ \\
& 2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) \\
&)*\tan(c)^2 - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192* \\
& \sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 24*\sqrt{b}* \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)*\tan(c)^2 - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan \\
& (1/3*c)*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)*\tan(c)^2 \\
& - 192*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 1008*\sqrt{b}*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c) \\
&)*\tan(1/3*c)^2*\tan(c)^2 - 72*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan \\
& (c)^2 + 320*\sqrt{b}*\tan(1/3*c)^3*\tan(c)^2 - 144*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
&)^3 + 384*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) - 1008*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2 + 320*\sqrt{b}*\tan(1/2*d*x + 1 \\
& /6*c)^3 + 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) + 72*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) - 96*\sqrt{b}*\tan(1/2*d*x + \\
& 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2 - 192*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2 \\
& *d*x + 1/2*c)^2 + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c)^3 + 384*\sqrt{b}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 1728*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)*\tan(1/3*c) + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) - 192*s \\
& \sqrt{b}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 1008*\sqrt{b}*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/3*c)^2 + 1440*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^2 + 72*\sqrt{b} \\
&)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 320*\sqrt{b}*\tan(1/3*c)^3 - 48*\sqrt{b} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(c) - 72*\sqrt{b}*\tan(1/2*d*x + 1/6*c)^2*\tan(c) \\
& - 72*\sqrt{b}*\tan(1/3*c)^2*\tan(c) - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 24*\sqrt{b}*\tan(-1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)*\tan(c)^2 - 96*\sqrt{b}*\tan(1/3*c)*\tan(c)^2 - 48*\sqrt{b}*\tan(1/2*d*x + 1/2*c) \\
& - 96*\sqrt{b}*\tan(1/2*d*x + 1/6*c) + 24*\sqrt{b}*\tan(-1/2*d*x + 1/2*c) \\
& - 96*\sqrt{b}*\tan(1/3*c) - 24*\sqrt{b}*\tan(c))*b/(d*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c \\
&)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(1/3*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c \\
&)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + \\
& 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*t \\
& \tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c) \\
& ^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 6* \\
& d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/ \\
& 3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 \\
& *\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1 \\
& /6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*t \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + \\
& 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*t \\
& \tan(1/3*c)^4 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 \\
& + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1 \\
& /6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6 \\
& *\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4
\end{aligned}$$

$$\begin{aligned}
& /6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/ \\
& 6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^ \\
& 2 + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^ \\
& 2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^ \\
& 2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c)^2 + \\
& 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + \\
& 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*t \\
& \tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4* \\
& \tan(-1/2*d*x + 1/2*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6 \\
& *\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*t \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 18*d*\tan(1/2*d \\
& *x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + \\
& 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 36*d*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 \\
& + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 3*d*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9 \\
& *d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 3*d*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(1 \\
& /3*c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + \\
& 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3* \\
& d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 4*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2
\end{aligned}$$

$$\begin{aligned}
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c
\end{aligned}$$

$$\begin{aligned}
&)^2 \tan(1/3*c)^6 + 2*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6 \tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 \tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^2 \tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^4 \tan(c)^2 + 6*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c)^2 + d*\tan(1/3*c)^6 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 + d*\tan(1/2*d*x + 1/6*c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^4 + 6*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + d*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c)^4 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4 \tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4 \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 6*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 \tan(c)^2 + 3*d*\tan(1/3*c)^4 \tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4 + 4*d*\tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^2 + 6*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + 3*d*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2 \tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 \tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2 \tan(c)^2 + 3*d*\tan(1/3*c)^2 \tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/3*c)^2 + d*\tan(c)^2 + d)
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.81

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{3}{2}} dx = \frac{b \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12d (\cos(2c + 2dx) + 1)}$$

[In] `int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2),x)`

[Out] `(b*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*d*(cos(2*c + 2*d*x) + 1))`

3.152 $\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{3/2} dx$

Optimal result	823
Rubi [A] (verified)	823
Mathematica [A] (verified)	824
Maple [A] (verified)	825
Fricas [A] (verification not implemented)	825
Sympy [F]	825
Maxima [A] (verification not implemented)	826
Giac [A] (verification not implemented)	826
Mupad [B] (verification not implemented)	826

Optimal result

Integrand size = 23, antiderivative size = 65

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{3/2} dx = \frac{bx\sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{b\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}\sin(c + dx)}{2d}$$

[Out] $1/2*b*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+1/2*b*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{3/2} dx = \frac{bx\sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{b \sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{2d}$$

[In] $\text{Int}[\text{Sqrt}[\text{Cos}[c + d*x]]*(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(b*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(2*\text{Sqrt}[\text{Cos}[c + d*x]]) + (b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(2*d)$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

```
Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m + 1/2)
)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \cos^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{2d} + \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int 1 dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{bx\sqrt{b\cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{2d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.69

$$\int \sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2} dx = \frac{(b\cos(c+dx))^{3/2}(2(c+dx) + \sin(2(c+dx)))}{4d\cos^{3/2}(c+dx)}$$

```
[In] Integrate[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2), x]
```

```
[Out] ((b*Cos[c + d*x])^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Cos[c + d*x]
^(3/2))
```


Maple [A] (verified)

Time = 3.04 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.66

method	result
default	$\frac{b\sqrt{\cos(dx+c)}b(\cos(dx+c)\sin(dx+c)+dx+c)}{2d\sqrt{\cos(dx+c)}}$
risch	$\frac{b\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})e^{i(dx+c)}x}{e^{2i(dx+c)}+1} - \frac{ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})e^{3i(dx+c)}}{4(e^{2i(dx+c)}+1)d} + \frac{ib\sqrt{\cos(dx+c)}b(\sqrt{\cos(dx+c)})e^{-i(dx+c)}}{4(e^{2i(dx+c)}+1)d}$

[In] `int(cos(d*x+c)^(1/2)*(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`[Out] `1/2/d*b*(cos(d*x+c)*b)^(1/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/cos(d*x+c)^(1/2)`**Fricas [A] (verification not implemented)**

none

Time = 0.39 (sec) , antiderivative size = 153, normalized size of antiderivative = 2.35

$$\int \sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2} dx = \left[\frac{2\sqrt{b\cos(dx+c)}b\sqrt{\cos(dx+c)}\sin(dx+c) + \sqrt{-bb}\log\left(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)}\right)}{4d} \right]$$

[In] `integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`

[Out] `[1/4*(2*sqrt(b*cos(d*x + c))*b*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(-b)*b*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/d, 1/2*(sqrt(b*cos(d*x + c))*b*sqrt(cos(d*x + c))*sin(d*x + c) + b^(3/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]`

Sympy [F]

$$\int \sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2} dx = \int (b\cos(c+dx))^{\frac{3}{2}} \sqrt{\cos(c+dx)} dx$$

[In] `integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(3/2),x)`[Out] `Integral((b*cos(c + d*x))**(3/2)*sqrt(cos(c + d*x)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.43

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2} dx = \frac{(2(dx+c)b + b \sin(2dx+2c))\sqrt{b}}{4d}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/4*(2*(d*x + c)*b + b*sin(2*d*x + 2*c))*sqrt(b)/d

Giac [A] (verification not implemented)

none

Time = 1.20 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.63

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2} dx = \frac{\left(\sqrt{b}dx \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 + 2\sqrt{b}dx \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 - 2\sqrt{b} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 + \sqrt{b}dx + 2\sqrt{b} \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)\right)}{2\left(d \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^4 + 2d \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^2 + d\right)}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] 1/2*(sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^4 + 2*sqrt(b)*d*x*tan(1/2*d*x + 1/2*c)^2 - 2*sqrt(b)*tan(1/2*d*x + 1/2*c)^3 + sqrt(b)*d*x + 2*sqrt(b)*tan(1/2*d*x + 1/2*c))*b/(d*tan(1/2*d*x + 1/2*c)^4 + 2*d*tan(1/2*d*x + 1/2*c)^2 + d)

Mupad [B] (verification not implemented)

Time = 14.50 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.97

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2} dx = \frac{b \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (\sin(c+dx) + \sin(3c+3dx) + 4dx \cos(c+dx))}{4d (\cos(2c+2dx) + 1)}$$

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(3/2),x)

[Out] (b*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*d*(cos(2*c + 2*d*x) + 1))

$$3.153 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx$$

Optimal result	827
Rubi [A] (verified)	827
Mathematica [A] (verified)	828
Maple [A] (verified)	828
Fricas [A] (verification not implemented)	828
Sympy [A] (verification not implemented)	829
Maxima [A] (verification not implemented)	829
Giac [F]	829
Mupad [B] (verification not implemented)	830

Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx = \frac{b\sqrt{b \cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}}$$

[Out] $b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx = \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}/\text{Sqrt}[\text{Cos}[c + d*x]], x]$

[Out] $(b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2717

$\text{Int}[\sin[\text{Pi}/2 + (c_*) + (d_*)*(x_*)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \cos(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{d\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{(b\cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx = \frac{(b\cos(c+dx))^{3/2} \sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Sqrt[Cos[c + d*x]],x]

[Out] ((b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 2.97 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b \sin(dx+c) \sqrt{\cos(dx+c)} b}{d \sqrt{\cos(dx+c)}}$	30
risch	$\frac{b \sin(dx+c) \sqrt{\cos(dx+c)} b}{d \sqrt{\cos(dx+c)}}$	30

[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] b*sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{(b\cos(c+dx))^{3/2}}{\sqrt{\cos(c+dx)}} dx = \frac{\sqrt{b\cos(dx+c)} b \sin(dx+c)}{d\sqrt{\cos(dx+c)}}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [A] (verification not implemented)

Time = 15.89 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.39

$$\int \frac{(b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx = \begin{cases} \frac{(b \cos(c + dx))^{\frac{3}{2}} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)} & \text{for } d \neq 0 \\ \frac{x(b \cos(c))^{\frac{3}{2}}}{\sqrt{\cos(c)}} & \text{otherwise} \end{cases}$$

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(1/2),x)

[Out] Piecewise(((b*cos(c + d*x))**(3/2)*sin(c + d*x)/(d*cos(c + d*x)**(3/2)), Ne(d, 0)), (x*(b*cos(c))**(3/2)/sqrt(cos(c)), True))

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.39

$$\int \frac{(b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx = \frac{b^{\frac{3}{2}} \sin(dx + c)}{d}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] b^(3/2)*sin(d*x + c)/d

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/sqrt(cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{(b \cos(c + dx))^{3/2}}{\sqrt{\cos(c + dx)}} dx = \frac{b \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] `int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(1/2),x)`

[Out] `(b*sin(c + d*x)*(b*cos(c + d*x))^(1/2))/(d*cos(c + d*x)^(1/2))`

$$3.154 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal result	831
Rubi [A] (verified)	831
Mathematica [A] (verified)	832
Maple [A] (verified)	832
Fricas [A] (verification not implemented)	832
Sympy [A] (verification not implemented)	833
Maxima [A] (verification not implemented)	833
Giac [F]	833
Mupad [B] (verification not implemented)	834

Optimal result

Integrand size = 23, antiderivative size = 25

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{bx \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] $b*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{bx \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[(b*\text{Cos}[c+d*x])^{(3/2)}/\text{Cos}[c+d*x]^{(3/2)},x]$

[Out] $(b*x*\text{Sqrt}[b*\text{Cos}[c+d*x]])/\text{Sqrt}[\text{Cos}[c+d*x]]$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)*((b_)*(v_))^{(n_)}], x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int 1 dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{bx\sqrt{b\cos(c+dx)}}{\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{3/2}(c+dx)} dx = \frac{bx\sqrt{b\cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(3/2), x]

[Out] (b*x*Sqrt[b*Cos[c + d*x]])/Sqrt[Cos[c + d*x]]

Maple [A] (verified)

Time = 2.40 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

method	result	size
risch	$\frac{bx\sqrt{\cos(dx+c)b}}{\sqrt{\cos(dx+c)}}$	22
default	$\frac{b\sqrt{\cos(dx+c)b}(dx+c)}{d\sqrt{\cos(dx+c)}}$	29

[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(3/2), x, method=_RETURNVERBOSE)

[Out] b*x*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 95, normalized size of antiderivative = 3.80

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{3/2}(c+dx)} dx = \left[\frac{\sqrt{-bb} \log\left(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{\cos(dx+c)}\sin(dx+c) - \dots\right)}{2d} \right]$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2), x, algorithm="fricas")

[Out] $[1/2\sqrt{-b}b\log(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)})\sqrt{-b}\sqrt{\cos(dx+c)}\sin(dx+c) - b]/d, b^{3/2}\arctan(\sqrt{b\cos(dx+c)})\sin(dx+c)/(\sqrt{b}\cos(dx+c)^{3/2})/d]$

Sympy [A] (verification not implemented)

Time = 9.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.88

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)} dx = \frac{x(b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)}$$

[In] `integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(3/2),x)`

[Out] `x*(b*cos(c + d*x))**(3/2)/cos(c + d*x)**(3/2)`

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.04

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)} dx = \frac{2b^{3/2} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

[In] `integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")`

[Out] `2*b^(3/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d`

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{3/2}}{\cos(dx + c)^{3/2}} dx$$

[In] `integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(3/2),x, algorithm="giac")`

[Out] `integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 13.48 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.84

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{3/2}(c + dx)} dx = \frac{bx \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(3/2),x)

[Out] (b*x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

$$3.155 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal result	835
Rubi [A] (verified)	835
Mathematica [A] (verified)	836
Maple [A] (verified)	836
Fricas [A] (verification not implemented)	836
Sympy [F(-1)]	837
Maxima [B] (verification not implemented)	837
Giac [F]	837
Mupad [F(-1)]	838

Optimal result

Integrand size = 23, antiderivative size = 34

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[Out] b*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3855}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{d\sqrt{\cos(c+dx)}}$$

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(5/2),x]

[Out] (b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3855

Int[csc[(c_)+(d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{\text{barctanh}(\sin(c+dx))\sqrt{b\cos(c+dx)}}{d\sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.97

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{5/2}(c+dx)} dx = \frac{\text{arctanh}(\sin(c+dx))(b\cos(c+dx))^{3/2}}{d\cos^{3/2}(c+dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*(b*Cos[c + d*x])^(3/2))/(d*Cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 2.77 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.21

method	result	size
default	$-\frac{2\sqrt{\cos(dx+c)}b\text{arctanh}(\cot(dx+c)-\csc(dx+c))}{d\sqrt{\cos(dx+c)}}$	41
risch	$-\frac{b\sqrt{\cos(dx+c)}b\ln(e^{i(dx+c)}-i)}{\sqrt{\cos(dx+c)}d} + \frac{b\sqrt{\cos(dx+c)}b\ln(e^{i(dx+c)}+i)}{\sqrt{\cos(dx+c)}d}$	75

[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(5/2), x, method=_RETURNVERBOSE)

[Out] -2/d*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)*b*arctanh(cot(d*x+c)-csc(d*x+c))

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 114, normalized size of antiderivative = 3.35

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{5/2}(c+dx)} dx = \left[\frac{b^{3/2} \log\left(-\frac{b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right)}{2d}, \right. \\ \left. -\frac{\sqrt{-bb}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)}{d} \right]$$

```
[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="fricas")
[Out] [1/2*b^(3/2)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(c
os(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3/d, -sqrt(-b)*
b*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))
/d]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

```
[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(5/2),x)
```

```
[Out] Timed out
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. 2(30) = 60.

Time = 0.42 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.00

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{5/2}(c + dx)} dx = \frac{(b \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1)) \sqrt{b}}{2d}$$

```
[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")
```

```
[Out] 1/2*(b*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b*log(co
s(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d
```

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{5/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{3/2}}{\cos^{5/2}(dx + c)} dx$$

```
[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(5/2),x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(5/2), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{5/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^{5/2}} dx$$

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(5/2), x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(5/2), x)
```

$$3.156 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{7}{2}}(c+dx)} dx$$

Optimal result	839
Rubi [A] (verified)	839
Mathematica [A] (verified)	840
Maple [A] (verified)	840
Fricas [A] (verification not implemented)	841
Sympy [F(-1)]	841
Maxima [A] (verification not implemented)	841
Giac [F]	842
Mupad [B] (verification not implemented)	842

Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{7}{2}}(c+dx)} dx = \frac{b \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $b \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(3/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3852, 8}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{7}{2}}(c+dx)} dx = \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[In] $\text{Int}[(b \cos[c + d*x])^{(3/2)} / \cos[c + d*x]^{(7/2)}, x]$

[Out] $(b \sqrt{b \cos[c + d*x]} * \sin[c + d*x]) / (d \cos[c + d*x]^{(3/2)})$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_))^{(m_)} * ((b_.) * (v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)} * b^{(n - 1/2)} * (\text{Sqrt}[b*v] / \text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec^2(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{\left(b\sqrt{b\cos(c+dx)}\right) \text{Subst}\left(\int 1 dx, x, -\tan(c+dx)\right)}{d\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.97

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{7/2}(c+dx)} dx = \frac{(b\cos(c+dx))^{3/2} \sin(c+dx)}{d\cos^{5/2}(c+dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(7/2), x]
```

```
[Out] ((b*Cos[c + d*x])^(3/2)*Sin[c + d*x])/(d*Cos[c + d*x]^(5/2))
```

Maple [A] (verified)

Time = 2.84 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b \sin(dx+c) \sqrt{\cos(dx+c)} b}{d \cos(dx+c)^{\frac{3}{2}}}$	30
risch	$\frac{2ib \sqrt{\cos(dx+c)} b}{\sqrt{\cos(dx+c)} d(e^{2i(dx+c)}+1)}$	39

```
[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(7/2), x, method=_RETURNVERBOSE)
```

```
[Out] b*sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(3/2)
```


Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{7/2}(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)} b \sin(dx + c)}{d \cos(dx + c)^{3/2}}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(7/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.64

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{7/2}(c + dx)} dx = \frac{2 b^{3/2} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c))^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1} d$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out] 2*b^(3/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{7/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{3/2}}{\cos^{7/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(7/2), x)

Mupad [B] (verification not implemented)

Time = 13.91 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.82

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{7/2}(c + dx)} dx = \frac{b \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + 1i)}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(7/2),x)

[Out] (b*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

$$3.157 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx$$

Optimal result	843
Rubi [A] (verified)	843
Mathematica [A] (verified)	844
Maple [A] (verified)	844
Fricas [A] (verification not implemented)	845
Sympy [F(-1)]	845
Maxima [B] (verification not implemented)	846
Giac [F]	846
Mupad [F(-1)]	847

Optimal result

Integrand size = 23, antiderivative size = 74

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{b\sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{5/2}(c+dx)}$$

[Out] 1/2*b*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+1/2*b*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^2(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{b \cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{b \sin(c+dx)\sqrt{b \cos(c+dx)}}{2d \cos^{5/2}(c+dx)}$$

[In] Int[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(9/2),x]

[Out] (b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(2*d*Sqrt[Cos[c + d*x]]) + (b*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(2*d*Cos[c + d*x]^(5/2))

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]
*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec^3(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)} + \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec(c+dx) dx}{2\sqrt{\cos(c+dx)}} \\ &= \frac{\text{barctanh}(\sin(c+dx))\sqrt{b\cos(c+dx)}}{2d\sqrt{\cos(c+dx)}} + \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{2d\cos^{\frac{5}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.70

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{9/2}(c+dx)} dx = \frac{(b\cos(c+dx))^{3/2} (\text{arctanh}(\sin(c+dx)) \cos^2(c+dx) + \sin(c+dx))}{2d\cos^{7/2}(c+dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(9/2), x]
```

```
[Out] ((b*Cos[c + d*x])^(3/2)*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x
]))/(2*d*Cos[c + d*x]^(7/2))
```

Maple [A] (verified)

Time = 2.87 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{b(-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c))\sqrt{\cos(dx+c)}b}{2d\cos(dx+c)^{\frac{5}{2}}}$	85
risch	$-\frac{ib\sqrt{\cos(dx+c)}b(e^{3i(dx+c)}-e^{i(dx+c)})}{\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)^2} - \frac{b\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}-i)}{2\sqrt{\cos(dx+c)}d} + \frac{b\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}+i)}{2\sqrt{\cos(dx+c)}d}$	135

[In] `int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(9/2),x,method=_RETURNVERBOSE)`

[Out] $1/2/d*b*(-\cos(d*x+c)^2*\ln(-\cot(d*x+c)+\csc(d*x+c)-1)+\cos(d*x+c)^2*\ln(-\cot(d*x+c)+\csc(d*x+c)+1)+\sin(d*x+c))*(\cos(d*x+c)*b)^(1/2)/\cos(d*x+c)^(5/2)$

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.76

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{\frac{9}{2}}(c + dx)} dx = \left[\frac{b^{\frac{3}{2}} \cos(dx + c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + \sqrt{-bb} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b \sin(dx+c)}}{b \sqrt{\cos(dx+c)}}\right) \cos(dx + c)^3 - \sqrt{b \cos(dx + c)} b \sqrt{\cos(dx + c)} \sin(dx + c)}{4 d \cos(dx + c)^3} \right]$$

[In] `integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2),x, algorithm="fricas")`

[Out] $[1/4*(b^{(3/2)*\cos(d*x + c)^3*\log(-(b*\cos(d*x + c)^3 - 2*\sqrt{b*\cos(d*x + c)}*\sqrt{b*\cos(d*x + c)}*\sin(d*x + c) - 2*b*\cos(d*x + c)))/\cos(d*x + c)^3) + 2*\sqrt{b*\cos(d*x + c)}*b*\sqrt{\cos(d*x + c)}*\sin(d*x + c))/(d*\cos(d*x + c)^3), -1/2*(\sqrt{-b}*b*\arctan(\sqrt{b*\cos(d*x + c)}*\sqrt{-b*\sin(d*x + c)})/(b*\sqrt{\cos(d*x + c)}))*\cos(d*x + c)^3 - \sqrt{b*\cos(d*x + c)}*b*\sqrt{\cos(d*x + c)}*\sin(d*x + c))/(d*\cos(d*x + c)^3)]$

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(9/2),x)`

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 691 vs. $2(62) = 124$.

Time = 0.44 (sec) , antiderivative size = 691, normalized size of antiderivative = 9.34

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{9/2}(c + dx)} dx = \text{Too large to display}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2),x, algorithm="maxima")

[Out] $-1/4*(4*(b*\sin(4*d*x + 4*c) + 2*b*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 4*(b*\sin(4*d*x + 4*c) + 2*b*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - (b*\cos(4*d*x + 4*c))^2 + 4*b*\cos(2*d*x + 2*c)^2 + b*\sin(4*d*x + 4*c)^2 + 4*b*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*b*\sin(2*d*x + 2*c)^2 + 2*(2*b*\cos(2*d*x + 2*c) + b)*\cos(4*d*x + 4*c) + 4*b*\cos(2*d*x + 2*c) + b)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) + (b*\cos(4*d*x + 4*c))^2 + 4*b*\cos(2*d*x + 2*c)^2 + b*\sin(4*d*x + 4*c)^2 + 4*b*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*b*\sin(2*d*x + 2*c)^2 + 2*(2*b*\cos(2*d*x + 2*c) + b)*\cos(4*d*x + 4*c) + 4*b*\cos(2*d*x + 2*c) + b)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) - 4*(b*\cos(4*d*x + 4*c) + 2*b*\cos(2*d*x + 2*c) + b)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 4*(b*\cos(4*d*x + 4*c) + 2*b*\cos(2*d*x + 2*c) + b)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))*\sqrt{b}/((2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c))^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*d$

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{9/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{3/2}}{\cos^{9/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(9/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{9/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^{9/2}} dx$$

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(9/2), x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(9/2), x)
```

$$3.158 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx$$

Optimal result	848
Rubi [A] (verified)	848
Mathematica [A] (verified)	849
Maple [A] (verified)	849
Fricas [A] (verification not implemented)	850
Sympy [F(-1)]	850
Maxima [B] (verification not implemented)	850
Giac [F]	851
Mupad [B] (verification not implemented)	851

Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{b\sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)} + \frac{b\sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{\frac{7}{2}}(c+dx)}$$

[Out] $b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(3/2)}+1/3*b*\sin(d*x+c)^3*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(7/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3852}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{b \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{\frac{7}{2}}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(3/2)}/\text{Cos}[c + d*x]^{(11/2)}, x]$

[Out] $(b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/ (d*\text{Cos}[c + d*x]^{(3/2)}) + (b*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Cos}[c + d*x]^{(7/2)})$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3852


```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec^4(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= -\frac{\left(b\sqrt{b\cos(c+dx)}\right) \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d\sqrt{\cos(c+dx)}} \\ &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{d\cos^{\frac{3}{2}}(c+dx)} + \frac{b\sqrt{b\cos(c+dx)} \sin^3(c+dx)}{3d\cos^{\frac{7}{2}}(c+dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.62

$$\int \frac{(b\cos(c+dx))^{3/2}}{\cos^{\frac{11}{2}}(c+dx)} dx = \frac{(b\cos(c+dx))^{3/2} \left(\tan(c+dx) + \frac{1}{3}\tan^3(c+dx)\right)}{d\cos^{\frac{3}{2}}(c+dx)}$$

[In] Integrate[(b*cos[c + d*x])^(3/2)/Cos[c + d*x]^(11/2), x]

[Out] ((b*cos[c + d*x])^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*cos[c + d*x]^(3/2))

Maple [A] (verified)

Time = 2.77 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.60

method	result	size
default	$\frac{b(2(\cos^2(dx+c))+1)\sqrt{\cos(dx+c)}b\sin(dx+c)}{3d\cos(dx+c)^{\frac{7}{2}}}$	43
risch	$\frac{4ib\sqrt{\cos(dx+c)}b(3e^{2i(dx+c)}+1)}{3\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)^3}$	52

[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(11/2), x, method=_RETURNVERBOSE)

[Out] 1/3/d*b*(2*cos(d*x+c)^2+1)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.58

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{11/2}(c + dx)} dx = \frac{(2b \cos(dx + c)^2 + b) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \cos(dx + c)^{7/2}}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2),x, algorithm="fricas")

[Out] 1/3*(2*b*cos(d*x + c)^2 + b)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(3/2)/cos(d*x+c)**(11/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(62) = 124.

Time = 0.41 (sec) , antiderivative size = 299, normalized size of antiderivative = 4.15

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{11/2}(c + dx)} dx =$$

$$\frac{4(3b \cos(6dx + 6c) + 3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1) * d}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1)}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2),x, algorithm="maxima")

[Out] -4/3*(3*b*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b*cos(2*d*x + 2*c) + b)*sin(6*d*x + 6*c) - 3*(3*b*cos(2*d*x + 2*c) + b)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{\frac{11}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{\frac{3}{2}}}{\cos^{\frac{11}{2}}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(11/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(11/2), x)

Mupad [B] (verification not implemented)

Time = 15.35 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.79

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{\frac{11}{2}}(c + dx)} dx = \frac{2b \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx) 1i + 9 \sin(2c + 2dx) + 6 \sin(4c + 4dx) + \sin(6c + 6dx) + 10i)}{3d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 10)}$$

[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(11/2),x)

[Out] (2*b*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.159 \quad \int \frac{(b \cos(c+dx))^{3/2}}{\cos^{13/2}(c+dx)} dx$$

Optimal result	852
Rubi [A] (verified)	852
Mathematica [A] (verified)	854
Maple [A] (verified)	854
Fricas [A] (verification not implemented)	854
Sympy [F(-1)]	855
Maxima [B] (verification not implemented)	855
Giac [F]	856
Mupad [F(-1)]	857

Optimal result

Integrand size = 23, antiderivative size = 110

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{13/2}(c+dx)} dx = \frac{3b \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{b \sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{9/2}(c+dx)} + \frac{3b \sqrt{b \cos(c+dx)} \sin(c+dx)}{8d \cos^{5/2}(c+dx)}$$

[Out] $1/4*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(9/2)}+3/8*b*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(5/2)}+3/8*b*\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{(b \cos(c+dx))^{3/2}}{\cos^{13/2}(c+dx)} dx = \frac{3b \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{3b \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{5/2}(c+dx)} + \frac{b \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{9/2}(c+dx)}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c+d*x])^{(3/2)}/\operatorname{Cos}[c+d*x]^{(13/2)},x]$

[Out] $(3*b*\operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]]*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]])/(8*d*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]) + (b*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(4*d*\operatorname{Cos}[c+d*x]^{(9/2)}) + (3*b*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(8*d*\operatorname{Cos}[c+d*x]^{(5/2)})$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(b\sqrt{b\cos(c+dx)}\right) \int \sec^5(c+dx) dx}{\sqrt{\cos(c+dx)}} \\
 &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{\left(3b\sqrt{b\cos(c+dx)}\right) \int \sec^3(c+dx) dx}{4\sqrt{\cos(c+dx)}} \\
 &= \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b\sqrt{b\cos(c+dx)} \sin(c+dx)}{8d \cos^{\frac{5}{2}}(c+dx)} \\
 &\quad + \frac{\left(3b\sqrt{b\cos(c+dx)}\right) \int \sec(c+dx) dx}{8\sqrt{\cos(c+dx)}} \\
 &= \frac{3b \operatorname{arctanh}(\sin(c+dx)) \sqrt{b\cos(c+dx)}}{8d\sqrt{\cos(c+dx)}} \\
 &\quad + \frac{b\sqrt{b\cos(c+dx)} \sin(c+dx)}{4d \cos^{\frac{9}{2}}(c+dx)} + \frac{3b\sqrt{b\cos(c+dx)} \sin(c+dx)}{8d \cos^{\frac{5}{2}}(c+dx)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.61

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{13/2}(c + dx)} dx = \frac{b\sqrt{b \cos(c + dx)}(3 \operatorname{arctanh}(\sin(c + dx)) \cos^4(c + dx) + (2 + 3 \cos^2(c + dx)) \sin(c + dx))}{8d \cos^{9/2}(c + dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(3/2)/Cos[c + d*x]^(13/2),x]

[Out] (b*Sqrt[b*Cos[c + d*x]]*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(9/2))

Maple [A] (verified)

Time = 3.03 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.95

method	result
default	$\frac{b(-3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+3(\cos^2(dx+c)) \sin(dx+c)+2 \sin(dx+c))}{8d \cos(dx+c)^{9/2}}$
risch	$-\frac{ib\sqrt{\cos(dx+c)}b(3e^{7i(dx+c)}+11e^{5i(dx+c)}-11e^{3i(dx+c)}-3e^{i(dx+c)})}{4\sqrt{\cos(dx+c)}d(e^{2i(dx+c)}+1)^4} + \frac{3b\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}+i)}{8\sqrt{\cos(dx+c)}d} - \frac{3b\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}-i)}{8\sqrt{\cos(dx+c)}d}$

[In] int((cos(d*x+c)*b)^(3/2)/cos(d*x+c)^(13/2),x,method=_RETURNVERBOSE)

[Out] 1/8/d*b*(-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)+3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(9/2)

Fricas [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.13

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{13/2}(c + dx)} dx = \frac{\left[3b^{3/2} \cos(dx + c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 3\sqrt{-bb} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b \sin(dx+c)}}{b\sqrt{\cos(dx+c)}}\right) \cos(dx + c)^5 - (3b \cos(dx + c)^2 + 2b) \sqrt{b \cos(dx + c)} \sqrt{\cos(dx + c)} \right]}{8d \cos(dx + c)^5}$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2),x, algorithm="fricas")

[Out] [1/16*(3*b^(3/2)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x +


```

2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + 3*
(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2 +
16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2 +
36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*si
n(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos
(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d
*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x +
4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4*
c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2*
b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 12*
(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos
(2*d*x + 2*c) + b)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4
4*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*c
os(2*d*x + 2*c) + b)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) +
44*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b
*cos(2*d*x + 2*c) + b)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))
+ 12*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4
*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*
c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*c
os(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2
*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x +
2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*s
in(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x
+ 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 +
48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x
+ 2*c) + 1)*d)

```

Giac [F]

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{13/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{3/2}}{\cos^{13/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(3/2)/cos(d*x+c)^(13/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(3/2)/cos(d*x + c)^(13/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{3/2}}{\cos^{13/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{3/2}}{\cos(c + dx)^{13/2}} dx$$

```
[In] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(13/2), x)
```

```
[Out] int((b*cos(c + d*x))^(3/2)/cos(c + d*x)^(13/2), x)
```

3.160 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx$

Optimal result	858
Rubi [A] (verified)	858
Mathematica [A] (verified)	859
Maple [A] (verified)	860
Fricas [A] (verification not implemented)	860
Sympy [F(-1)]	860
Maxima [A] (verification not implemented)	861
Giac [F(-1)]	861
Mupad [B] (verification not implemented)	861

Optimal result

Integrand size = 23, antiderivative size = 116

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} + \frac{b^2 \sqrt{b \cos(c + dx)} \sin^5(c + dx)}{5d \sqrt{\cos(c + dx)}}$$

[Out] $b^2 \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)} - 2/3 * b^2 \sin(d*x+c)^3 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)} + 1/5 * b^2 \sin(d*x+c)^5 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{b^2 \sin^5(c + dx) \sqrt{b \cos(c + dx)}}{5d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}} + \frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)} * (b * \text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(b^2 * \text{Sqrt}[b * \text{Cos}[c + d*x]] * \text{Sin}[c + d*x]) / (d * \text{Sqrt}[\text{Cos}[c + d*x]]) - (2 * b^2 * \text{Sqrt}[b * \text{Cos}[c + d*x]] * \text{Sin}[c + d*x]^3) / (3 * d * \text{Sqrt}[\text{Cos}[c + d*x]]) + (b^2 * \text{Sqrt}[b * \text{Cos}[c + d*x]] * \text{Sin}[c + d*x]^5) / (5 * d * \text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2713

`Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \cos^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
 &= -\frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\
 &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{2b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} \\
 &\quad + \frac{b^2 \sqrt{b \cos(c + dx)} \sin^5(c + dx)}{5d \sqrt{\cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.49

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{(b \cos(c + dx))^{\frac{5}{2}} \sin(c + dx) (15 - 10 \sin^2(c + dx) + 3 \sin^4(c + dx))}{15d \cos^{\frac{5}{2}}(c + dx)}$$

`[In] Integrate[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2), x]`

`[Out] ((b*Cos[c + d*x])^(5/2)*Sin[c + d*x]*(15 - 10*Sin[c + d*x]^2 + 3*Sin[c + d*x]^4))/(15*d*Cos[c + d*x]^(5/2))`

Maple [A] (verified)

Time = 3.74 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.47

method	result
default	$\frac{b^2(3(\cos^4(dx+c))+4(\cos^2(dx+c))+8)\sqrt{\cos(dx+c)b}\sin(dx+c)}{15d\sqrt{\cos(dx+c)}}$
risch	$-\frac{ib^2\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{6i(dx+c)}}{80(e^{2i(dx+c)}+1)d} - \frac{5ib^2\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})e^{2i(dx+c)}}{8(e^{2i(dx+c)}+1)d} + \frac{5ib^2\sqrt{\cos(dx+c)b}(\sqrt{\cos(dx+c)})}{8(e^{2i(dx+c)}+1)d} + \dots$

[In] `int(cos(d*x+c)^(5/2)*(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`[Out] `1/15/d*b^2*(3*cos(d*x+c)^4+4*cos(d*x+c)^2+8)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(1/2)`**Fricas [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.53

$$\int \cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}} dx = \frac{(3b^2\cos(dx+c)^4 + 4b^2\cos(dx+c)^2 + 8b^2)\sqrt{b\cos(dx+c)}\sin(dx+c)}{15d\sqrt{\cos(dx+c)}}$$

[In] `integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x,algorithm="fricas")`[Out] `1/15*(3*b^2*cos(d*x+c)^4 + 4*b^2*cos(d*x+c)^2 + 8*b^2)*sqrt(b*cos(d*x+c))*sin(d*x+c)/(d*sqrt(cos(d*x+c)))`**Sympy [F(-1)]**

Timed out.

$$\int \cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.66

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{(3b^2 \sin(5dx + 5c) + 25b^2 \sin(\frac{3}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))) + 150b^2 \sin(\frac{1}{5} \arctan(\sin(5dx + 5c), \cos(5dx + 5c))))}{240d}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/240*(3*b^2*sin(5*d*x + 5*c) + 25*b^2*sin(3/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))) + 150*b^2*sin(1/5*arctan2(sin(5*d*x + 5*c), cos(5*d*x + 5*c))))*sqrt(b)/d

Giac [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] Timed out

Mupad [B] (verification not implemented)

Time = 15.65 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.63

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (175 \sin(2c + 2dx) + 28 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{240d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2),x)

[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(175*sin(2*c + 2*d*x) + 28*sin(4*c + 4*d*x) + 3*sin(6*c + 6*d*x)))/(240*d*(cos(2*c + 2*d*x) + 1))

3.161 $\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx$

Optimal result	862
Rubi [A] (verified)	862
Mathematica [A] (verified)	864
Maple [A] (verified)	864
Fricas [A] (verification not implemented)	864
Sympy [F(-1)]	865
Maxima [A] (verification not implemented)	865
Giac [B] (verification not implemented)	865
Mupad [B] (verification not implemented)	866

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{3b^2x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{3b^2\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}\sin(c + dx)}{8d} + \frac{b^2 \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}\sin(c + dx)}{4d}$$

[Out] $1/4*b^2*\cos(d*x+c)^{(5/2)}*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d+3/8*b^2*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+3/8*b^2*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{3b^2x\sqrt{b \cos(c + dx)}}{8\sqrt{\cos(c + dx)}} + \frac{b^2 \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)\sqrt{b \cos(c + dx)}}{4d} + \frac{3b^2 \sin(c + dx)\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}}{8d}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(3/2)}*(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(3b^2x\sqrt{b\cos[c+dx]})/(8\sqrt{\cos[c+dx]}) + (3b^2\sqrt{\cos[c+dx]})\sqrt{b\cos[c+dx]}\sin[c+dx]/(8d) + (b^2\cos[c+dx]^{5/2})\sqrt{b\cos[c+dx]}\sin[c+dx]/(4d)$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\sqrt{b*v}/\sqrt{a*v}), \text{Int}[u*v^{(m+n)}, x], x] \text{ /; FreeQ}[\{a, b, m\}, x] \&\& \text{!IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2715

$\text{Int}[(b_)*\sin[(c_)+(d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\cos[c+dx]*((b*\sin[c+dx])^{(n-1)})/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\sin[c+dx])^{(n-2)}, x], x] \text{ /; FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2\sqrt{b\cos(c+dx)}\right) \int \cos^4(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \cos^{\frac{5}{2}}(c+dx) \sqrt{b\cos(c+dx)} \sin(c+dx)}{4d} + \frac{\left(3b^2\sqrt{b\cos(c+dx)}\right) \int \cos^2(c+dx) dx}{4\sqrt{\cos(c+dx)}} \\ &= \frac{3b^2\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} \\ &\quad + \frac{b^2 \cos^{\frac{5}{2}}(c+dx) \sqrt{b\cos(c+dx)} \sin(c+dx)}{4d} + \frac{\left(3b^2\sqrt{b\cos(c+dx)}\right) \int 1 dx}{8\sqrt{\cos(c+dx)}} \\ &= \frac{3b^2x\sqrt{b\cos(c+dx)}}{8\sqrt{\cos(c+dx)}} + \frac{3b^2\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}\sin(c+dx)}{8d} \\ &\quad + \frac{b^2 \cos^{\frac{5}{2}}(c+dx) \sqrt{b\cos(c+dx)} \sin(c+dx)}{4d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{(b \cos(c + dx))^{\frac{5}{2}}(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))}{32d \cos^{\frac{5}{2}}(c + dx)}$$

[In] Integrate[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2),x]

[Out] ((b*Cos[c + d*x])^(5/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 2.88 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.61

method	result
default	$\frac{b^2 \sqrt{\cos(dx+c)} b (2 \sin(dx+c) (\cos^3(dx+c)) + 3 \cos(dx+c) \sin(dx+c) + 3dx + 3c)}{8d \sqrt{\cos(dx+c)}}$
risch	$\frac{3b^2 \sqrt{\cos(dx+c)} b (\sqrt{\cos(dx+c)}) e^{i(dx+c)} x}{4(e^{2i(dx+c)} + 1)} - \frac{ib^2 \sqrt{\cos(dx+c)} b (\sqrt{\cos(dx+c)}) e^{5i(dx+c)}}{32(e^{2i(dx+c)} + 1)d} + \frac{ib^2 \sqrt{\cos(dx+c)} b (\sqrt{\cos(dx+c)}) e^{-i(dx+c)}}{4(e^{2i(dx+c)} + 1)d}$

[In] int(cos(d*x+c)^(3/2)*(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/8/d*b^2*(cos(d*x+c)*b)^(1/2)*(2*sin(d*x+c)*cos(d*x+c)^3+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 193, normalized size of antiderivative = 1.80

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \left[\frac{3 \sqrt{-bb^2} \log \left(2 b \cos(dx + c)^2 - 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)} \sin(dx + c) - b \right) + 2 (2 b \cos(dx + c)^2 - 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)} \sin(dx + c) - b)}{16 d} \right]$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(-b)*b^2*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b) + 2*(2*b^2*cos(d*x + c)^2 + 3*b^2)


```
*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/d, 1/8*(3*b^(5/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2))) + (2*b^2*cos(d*x + c)^2 + 3*b^2)*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/d]
```

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**(5/2), x)
```

```
[Out] Timed out
```

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.55

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{(12(dx + c)b^2 + b^2 \sin(4dx + 4c) + 8b^2 \sin(\frac{1}{2} \arctan(\sin(4dx + 4c), \cos(4dx + 4c))))\sqrt{b}}{32d}$$

```
[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(5/2), x, algorithm="maxima")
```

```
[Out] 1/32*(12*(d*x + c)*b^2 + b^2*sin(4*d*x + 4*c) + 8*b^2*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(89) = 178.

Time = 2.48 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.90

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{5/2} dx = \frac{3b^{\frac{5}{2}}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 12b^{\frac{5}{2}}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^6 - 10b^{\frac{5}{2}}\tan(\frac{1}{2}dx + \frac{1}{2}c)^7 + 18b^{\frac{5}{2}}dx \tan(\frac{1}{2}dx + \frac{1}{2}c)^5}{8(d \tan(\frac{1}{2}dx + \frac{1}{2}c)^8 + 4d \tan(\frac{1}{2}dx + \frac{1}{2}c)^6)}$$

```
[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^(5/2), x, algorithm="giac")
```

```
[Out] 1/8*(3*b^(5/2)*d*x*tan(1/2*d*x + 1/2*c)^8 + 12*b^(5/2)*d*x*tan(1/2*d*x + 1/2*c)^6 - 10*b^(5/2)*tan(1/2*d*x + 1/2*c)^7 + 18*b^(5/2)*d*x*tan(1/2*d*x + 1/2*c)^5)
```

$$\begin{aligned} & /2*c)^4 + 6*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^5 + 12*b^{(5/2)}*d*x*\tan(1/2*d*x + 1 \\ & /2*c)^2 - 6*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3 + 3*b^{(5/2)}*d*x + 10*b^{(5/2)}*\tan \\ & (1/2*d*x + 1/2*c))/(d*\tan(1/2*d*x + 1/2*c)^8 + 4*d*\tan(1/2*d*x + 1/2*c)^6 + \\ & 6*d*\tan(1/2*d*x + 1/2*c)^4 + 4*d*\tan(1/2*d*x + 1/2*c)^2 + d) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 15.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.73

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}} dx = \frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx) + 24d \cos(c + dx))}{32d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(5/2),x)

[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*d*(cos(2*c + 2*d*x) + 1))

3.162 $\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{5/2} dx$

Optimal result	867
Rubi [A] (verified)	867
Mathematica [A] (verified)	868
Maple [A] (verified)	868
Fricas [A] (verification not implemented)	869
Sympy [F(-1)]	869
Maxima [A] (verification not implemented)	869
Giac [B] (verification not implemented)	870
Mupad [B] (verification not implemented)	917

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{5/2} dx = \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}}$$

[Out] $b^2 \sin(dx+c) (b \cos(dx+c))^{1/2} / d \cos(dx+c)^{1/2} - 1/3 b^2 \sin(dx+c)^3 (b \cos(dx+c))^{1/2} / d \cos(dx+c)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{5/2} dx = \frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sin^3(c + dx) \sqrt{b \cos(c + dx)}}{3d \sqrt{\cos(c + dx)}}$$

[In] $\text{Int}[\text{Sqrt}[\text{Cos}[c + d*x]]*(b*\text{Cos}[c + d*x])^{5/2}, x]$

[Out] $(b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]) - (b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m)}*((b_*)*(v_))^{(n)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /;$ FreeQ[{a, b

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2713

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \cos^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \sqrt{\cos(c + dx)}} - \frac{b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2} dx = \frac{(b \cos(c + dx))^{5/2} (5 + \cos(2(c + dx))) \sin(c + dx)}{6d \cos^{5/2}(c + dx)}$$

[In] Integrate[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2),x]

[Out] ((b*Cos[c + d*x])^(5/2)*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 2.90 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.57

method	result
default	$\frac{b^2(2+\cos^2(dx+c))\sqrt{\cos(dx+c)}\sin(dx+c)}{3d\sqrt{\cos(dx+c)}}$
risch	$-\frac{ib^2\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})e^{4i(dx+c)}}{12(e^{2i(dx+c)}+1)d} - \frac{3ib^2\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})e^{2i(dx+c)}}{4(e^{2i(dx+c)}+1)d} + \frac{3ib^2\sqrt{\cos(dx+c)}(\sqrt{\cos(dx+c)})}{4(e^{2i(dx+c)}+1)d} + ib^2\sqrt{\cos(dx+c)}$

[In] int(cos(d*x+c)^(1/2)*(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/3/d*b^2*(2+cos(d*x+c)^2)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.62

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2} dx = \frac{(b^2 \cos(dx+c)^2 + 2b^2) \sqrt{b \cos(dx+c)} \sin(dx+c)}{3d \sqrt{\cos(dx+c)}}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*(b^2*cos(d*x + c)^2 + 2*b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.43 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.64

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2} dx = \frac{(b^2 \sin(3dx+3c) + 9b^2 \sin(\frac{1}{3} \arctan(\sin(3dx+3c), \cos(3dx+3c)))) \sqrt{b}}{12d}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/12*(b^2*sin(3*d*x + 3*c) + 9*b^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 71047 vs. 2(66) = 132.

Time = 8.37 (sec) , antiderivative size = 71047, normalized size of antiderivative = 934.83

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^{5/2} dx = \text{Too large to display}$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out]
$$\begin{aligned} & -1/96*(3*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 3*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 48*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 48*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) + 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c) - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) + 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c \end{aligned}$$

$$\begin{aligned}
&)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 \tan(c)^2 + \\
&27*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x \\
&+ 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 - 54*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \\
&* \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 + 3*b \\
&^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan \\
&(c)^2 + 48*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
&-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan(c)^2 - 54*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/ \\
&2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 \\
&+ 108*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2 \\
&*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 144*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2 \\
&*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 \\
&+ 48*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x \\
&+ 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 9*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 * \\
&\tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 54*b \\
&^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1 \\
&/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 3*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(\\
&-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 9*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1 \\
&/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 + 54*b^{(\\
&5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^2 \tan(1/3*c)^4 + 144*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x \\
&+ 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 27*b^{(5/2)}*d*x^4 \tan(1/2* \\
&d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \\
&+ 54*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d \\
&*x + 1/2*c)^4 \tan(1/3*c)^4 - 3*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2 \\
&*d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(\\
&1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 + 54*b^{(5/2)}*d*x^4 \tan \\
&(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3 \\
&*c)^6 - 108*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan \\
&(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 144*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^ \\
&3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 - 48*b^{(5/2)}* \\
&d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan \\
&(1/3*c)^6 - 9*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \\
&* \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 54*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2* \\
&c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 3*b^{(5/2)} \\
&)*d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 - 72*b^{(\\
&5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^3 \tan(1/3*c)^2 \tan(c) - 72*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/ \\
&2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^2 \tan(c) + 72*b^{(5/2)}*d \\
&*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan \\
&(1/3*c)^4 \tan(c) + 432*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
&1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 \tan(c) - 216*b^{(5/2)}*d*x^4 \tan \\
&(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3 \\
&*c)^4 \tan(c) + 432*b^{(5/2)}*d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c \\
&)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 \tan(c) - 216*b^{(5/2)}*d*x^4 \tan(1/2 \\
&*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^6*\tan(c)^2 - 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 144* \\
& b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 3*b^(5/2)*d*x^ \\
& 4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 54 \\
& *b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*ta \\
& n(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b^(5/2)*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - \\
& 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 3*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 54*b^(5/2)*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144 \\
& *b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^3*\tan(1/3*c)^2 - 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d* \\
& x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 54*b^(5/2)*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 2 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c \\
&)^4 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*ta \\
& n(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 324*b^(5/2)*d*x \\
& ^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*ta \\
& n(1/3*c)^4 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 27*b^(5/ \\
& 2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 96*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*b^(\\
& 5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 18 \\
& *b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 - \\
& 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^6 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 54*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 \\
& - 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6 + 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*b^(5/2)*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*ta \\
& n(1/3*c)^6 + 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-
\end{aligned}$$

$$\begin{aligned}
& 1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 3*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 9*b^(5/2)*d \\
& *x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*b^(5/ \\
& 2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& n(1/3*c)^6 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*b^(5/2)*d*x^4*\tan \\
& n(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) \\
& + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 216*b^(\\
& (5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^3*\tan(1/3*c)^2*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2) \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan \\
& n(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 216*b^(5/ \\
& 2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)*\tan(1/3*c)^4*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 1296*b^(5/2)*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^4*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6* \\
& c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^ \\
& 4*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4 \\
& *\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) - 216*b^(5/2)*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c \\
&) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3* \\
& c)^6*\tan(c) + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(1/3*c)^6*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) - 432*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c) \\
& ^6*\tan(c) + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan \\
& (1/3*c)^6*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c) - 432*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6*\tan(c) - 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^3*\tan(1/3*c)^6*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& *x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 72*b^(5/2)*d*x^
\end{aligned}$$

$$\begin{aligned}
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c) - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) - 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 162*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 324*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 27*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 162*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 27*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 162*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 27*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 432*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)
\end{aligned}$$

$$\begin{aligned}
&)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 \tan(c)^2 + \\
&192*b^{(5/2)} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c)^2 \tan(1/3*c)^5 \tan(c)^2 - 1440*b^{(5/2)} \tan(1/2*d*x + 1/2*c)^4 \tan(1/2* \\
&d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 \tan(c)^2 + 1728*b^{(5/2)} \\
&* \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(\\
&1/3*c)^5 \tan(c)^2 - 384*b^{(5/2)} \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c) \\
&^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 \tan(c)^2 + 9*b^{(5/2)} * d*x^4 \tan(1/2* \\
&d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 54*b^{(5/2)} * d* \\
&x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 3 \\
&* b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 \tan(c)^2 + 144*b^{(5/2)} * d \\
&>* x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) * ta \\
&n(1/3*c)^6 \tan(c)^2 - 144*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + \\
&1/6*c)^4 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan(c)^2 - 24*b^{(5/2)} * \tan(1/2*d \\
&>* x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 \tan \\
&(c)^2 - 18*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan \\
&(1/3*c)^6 \tan(c)^2 + 324*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + \\
&1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 54*b^{(5/2)} * d*x^4 * \\
&\tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 + 192* \\
&b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c) \\
&^2 \tan(1/3*c)^6 \tan(c)^2 - 288*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + \\
&1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 \tan(c)^2 - 48*b^{(5/2)} * d*x^4 * \\
&\tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 + 144* \\
&b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/ \\
&2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 72*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d* \\
&x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 \tan(c)^2 - 48*b^{(5/2)} * \tan \\
&(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3* \\
&c)^6 \tan(c)^2 - 18*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2* \\
&c)^4 \tan(1/3*c)^6 \tan(c)^2 + 9*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/ \\
&2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 320*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 \\
&* \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 + 432 \\
&* b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c \\
&)^4 \tan(1/3*c)^6 \tan(c)^2 - 384*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x \\
&+ 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \tan(c)^2 - 48*b^{(5/2)} * \tan(1 \\
&/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 \\
&* \tan(c)^2 + 18*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 * \\
&\tan(-1/2*d*x + 1/2*c)^2 + 48*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d \\
&>* x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 - 9*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c \\
&)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 + 18*b^{(5/2)} * d*x^4 \tan(1 \\
&/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 - 9*b^{(5/2)} \\
&)* d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^2 - 144*b^{(5/2)} \\
&)* d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/ \\
&2*c) * \tan(1/3*c)^2 + 162*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + \\
&1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 - 324*b^{(5/2)} * d*x^4 \tan(1/2*d \\
&>* x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^2 + \\
&432*b^{(5/2)} * d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d
\end{aligned}$$

$$\begin{aligned}
& *x + 1/2*c)^3*\tan(1/3*c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 27*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*ta \\
& n(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 27*b^(5/2)*d*x^ \\
& 4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 54*b^(5/2)*d \\
& *x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 - 432*b^(5/ \\
& 2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)*\tan(1/3*c)^4 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 972*b^(\\
& (5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4 + 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 - 432*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c) \\
& ^4 + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^3*\tan(1/3*c)^4 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 27*b^(5/2)*d*x^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1440*b^(5/2)*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^4 + 192*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1440*b^(5/2)*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + \\
& 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^5 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 54*b^(5/2)*d*x^4*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 3*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^6 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*b^(5/2)*d*x^4*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c) \\
& ^6 + 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)*\tan(1/3*c)^6 + 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^6 - 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 54*b^(5/2)*d*x^4*ta \\
& n(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*b^(5/2)*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6 - 288*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)^6 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan
\end{aligned}$$

$$\begin{aligned}
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*
\end{aligned}$$

$$\begin{aligned}
& c)^3 \tan(1/3c)^4 \tan(c) - 216b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/6c)^4 \tan(-1/ \\
& 2dx + 1/2c)^3 \tan(1/3c)^4 \tan(c) - 72b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c \\
&)^3 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^4 \tan(c) + 216b^{(5/2)} d^4 x^4 \tan(1/2 \\
& dx + 1/2c) \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^4 \tan \\
& (c) + 216b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2 \\
& dx + 1/2c)^4 \tan(1/3c)^4 \tan(c) + 144b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan \\
& (1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^4 \tan(c) - 72b^{(5/2)} \\
&) d^4 x^4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(1/3c)^6 \tan(c) + \\
& 72b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^4 \tan(1/3c)^6 \tan \\
& (c) - 24b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(1/3c \\
&)^6 \tan(c) + 24b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c) \tan \\
& (1/3c)^6 \tan(c) - 432b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx \\
& + 1/6c)^2 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c) + 72b^{(5/2)} d^4 x^4 \tan \\
& (1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c) + 144b^{(5/2)} \\
&) d^4 x^4 \tan(1/2dx + 1/2c)^3 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c) - \\
& 432b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^2 \tan(-1/2dx \\
& + 1/2c)^2 \tan(1/3c)^6 \tan(c) + 144b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^2 \tan \\
& (-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c) - 72b^{(5/2)} d^4 x^4 \tan(1/2dx \\
& + 1/6c)^2 \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c) + 24b^{(5/2)} d^4 x^4 \tan \\
& (1/2dx + 1/2c) \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c) + 72b^{(5/2)} \\
&) \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 \tan \\
& (1/3c)^6 \tan(c) + 144b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \\
& \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c) + 24b^{(5/2)} \tan(1/2dx + 1/ \\
& 6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c) + 3b^{(5/2)} d^4 x^4 \tan(1/ \\
& 2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(c)^2 + 48b^{(5/2)} d^4 x^4 \tan(1/2 \\
& dx + 1/2c)^3 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) \tan(c)^2 - 54b^{(5/2)} \\
&) d^4 x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + \\
& 1/2c)^2 \tan(c)^2 + 108b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + \\
& 1/6c)^6 \tan(-1/2dx + 1/2c)^2 \tan(c)^2 - 144b^{(5/2)} d^4 x^4 \tan(1/2dx + \\
& 1/2c)^3 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^3 \tan(c)^2 + 48b^{(5/ \\
& 2)} d^4 x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) \\
&)^3 \tan(c)^2 + 9b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \\
& \tan(-1/2dx + 1/2c)^4 \tan(c)^2 - 54b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^2 \\
& \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^4 \tan(c)^2 + 3b^{(5/2)} d^4 x^4 \tan \\
& (1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(c)^2 + 96b^{(5/2)} \tan(1/ \\
& 2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) \tan \\
& (c)^2 + 27b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan \\
& (1/3c)^2 \tan(c)^2 - 54b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx \\
& + 1/6c)^6 \tan(1/3c)^2 \tan(c)^2 + 432b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^3 \\
& \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) \tan(1/3c)^2 \tan(c)^2 - 144b^{(5/2)} \\
&) d^4 x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2 \\
& c) \tan(1/3c)^2 \tan(c)^2 - 162b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/2c)^4 \tan(1/ \\
& 2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^2 \tan(c)^2 + 972b^{(5/2)} \\
&) d^4 x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) \\
&)^2 \tan(1/3c)^2 \tan(c)^2 - 54b^{(5/2)} d^4 x^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 640*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 1280*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 972*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 96*b^
\end{aligned}$$

$$\begin{aligned}
& (5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^5*\tan(c)^2 - \\
& 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 768*b^(5/2) \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan \\
& (1/3*c)^5*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d \\
& *x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*t \\
& an(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c) \\
& *\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 96* \\
& b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^ \\
& 2 + 3*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 54*b^(5/ \\
& 2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^ \\
& 2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b^(5/ \\
& 2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6*\tan(c)^2 - 14 \\
& 4*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c) \\
& ^2 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3* \\
& c)^6*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 \\
& - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)*\tan(1/3*c)^6*\tan(c)^2 + 108*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 640*b^(5/2)*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 6*\tan(c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 768*b^(5/2)*\tan(1/2*d*x + 1/2*c) \\
&)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 - \\
& 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) \\
&)^2*\tan(1/3*c)^6*\tan(c)^2 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2* \\
& d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*ta \\
& n(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 144*b^ \\
& (5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 \\
& *\tan(1/3*c)^6*\tan(c)^2 - 24*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^3*\tan(1/3*c)^6*\tan(c)^2 + 3*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^6*\tan(c)^2 + 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c) \\
& *\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) \\
&)^2 + 1280*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b^(5 \\
& /2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - \\
& 3*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6 - 48*b^(5/2)* \\
& d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) +
\end{aligned}$$

$$\begin{aligned}
& 54b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx \\
& x + 1/2c)^2 - 108b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c \\
&)^6 \tan(-1/2dx + 1/2c)^2 + 144b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^3 \tan(\\
& 1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^3 - 48b^{(5/2)}d^4x^4 \tan(1/2dx + \\
& 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^3 - 9b^{(5/2)}d^4x^4 \tan \\
& n(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 + 54b^{(5/2)} \\
& (5/2)d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/ \\
& 2c)^4 - 3b^{(5/2)}d^4x^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 + 9 \\
& 6b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c \\
& c)^4 \tan(1/3c) - 27b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6 \\
& c)^4 \tan(1/3c)^2 + 54b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + \\
& 1/6c)^6 \tan(1/3c)^2 - 432b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx \\
& x + 1/6c)^4 \tan(-1/2dx + 1/2c) \tan(1/3c)^2 + 144b^{(5/2)}d^4x^4 \tan(1/2 \\
& *dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) \tan(1/3c)^2 + 1 \\
& 62b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx \\
& + 1/2c)^2 \tan(1/3c)^2 - 972b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2 \\
& *dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^2 + 54b^{(5/2)}d^4x^4 \tan \\
& (1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^2 + 432b^{(5/2)}d^4x^ \\
& 4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^3 \tan \\
& (1/3c)^2 - 432b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^4 \tan \\
& an(-1/2dx + 1/2c)^3 \tan(1/3c)^2 + 72b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan \\
& (1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^2 - 9b^{(5/2)}d^4x^4 \\
& \tan(1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^2 + 162b^{(5/2)}d \\
& *x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 \\
& \tan(1/3c)^2 - 27b^{(5/2)}d^4x^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c \\
&)^4 \tan(1/3c)^2 + 1440b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c) \\
& ^5 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^2 - 1008b^{(5/2)} \tan(1/2dx + 1/2c) \\
& ^3 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^2 - 640b^{(5/2)} \\
&) \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 \tan \\
& (1/3c)^3 + 4800b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(\\
& -1/2dx + 1/2c)^4 \tan(1/3c)^3 - 5760b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(\\
& 1/2dx + 1/6c)^5 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^3 + 1280b^{(5/2)} \tan(\\
& 1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 \tan(1/3c \\
&)^3 - 27b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(1/ \\
& 3c)^4 + 162b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan \\
& n(1/3c)^4 - 9b^{(5/2)}d^4x^4 \tan(1/2dx + 1/6c)^6 \tan(1/3c)^4 - 432b^{(5/2)} \\
& (5/2)d^4x^4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c \\
& c) \tan(1/3c)^4 + 432b^{(5/2)}d^4x^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c \\
& c)^4 \tan(-1/2dx + 1/2c) \tan(1/3c)^4 + 72b^{(5/2)} \tan(1/2dx + 1/2c)^4 \\
& * \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) \tan(1/3c)^4 + 54b^{(5/2)}d^4x^ \\
& ^4 \tan(1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^4 - 972b^{(5/2)} \\
&)d^4x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c) \\
& ^2 \tan(1/3c)^4 + 162b^{(5/2)}d^4x^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1 \\
& /2c)^2 \tan(1/3c)^4 - 2880b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/ \\
& 6c)^5 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^4 + 864b^{(5/2)} \tan(1/2dx + 1/2
\end{aligned}$$

$$\begin{aligned}
& *c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^4 + 144*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 - 43 \\
& 2*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 216*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 144*b^(5/2) \tan(1/2*d*x + 1/2 \\
& *c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^4 + 54*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 27* \\
& b^(5/2) *d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 4800*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 11664*b^(5/2) \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 5760*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 - 1008 \\
& *b^(5/2) \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^4 + 96*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^5 - 2880*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 + 3456*b^(5/2) \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 - 768*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^5 + 1440*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 - 5760*b^(5/2) \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 + 5760*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 - 1728*b^(5/2) \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 + 96*b^(5/2) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^4 \tan(1/3*c)^5 - 3*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^4 \tan(1/3*c)^6 + 54*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^2 \tan(1/3*c)^6 - 9*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/6*c)^4 \tan(1/3*c)^6 + 96*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^5 \tan(1/3*c)^6 - 144*b^(5/2) \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^6 \tan(1/3*c)^6 - 48*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^3 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 + 144*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 + 72*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 + 48*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c) \tan(1/3*c)^6 - 108*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 54*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 640*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^3 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 + 864*b^(5/2) \tan(1/2*d*x + 1/2*c)^3 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 768*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^5 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 96*b^(5/2) \tan(1/2*d*x + 1/2*c) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x + 1/2*c)^2 \tan(1/3*c)^6 - 48*b^(5/2) *d*x^4 \tan(1/2*d*x + 1/2*c) \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 + 72*b^(5/2) \tan(1/2*d*x + 1/2*c)^4 \tan(1/2*d*x + 1/6*c)^2 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 + 144*b^(5/2) \tan(1/2*d*x + 1/2*c)^2 \tan(1/2*d*x + 1/6*c)^4 \tan(-1/2*d*x + 1/2*c)^3 \tan(1/3*c)^6 + 24*b^(5/2) \tan(1/2*d*x + 1/6*c)^6 \tan(-1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/3*c)^6 - 3*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 1280*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) + 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 1296*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 432*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) - 1296*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^4 * \tan(c) + 432*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) - 216*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^4 * \tan(c) + 72*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 216*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 432*b^(5/2) * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 72*b^(5/2) * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) + 72*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) - 72*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^6 * \tan(c) - 48*b^(5/2) * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^6 * \tan(c) - 144*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) + 72*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^6 * \tan(c) - 144*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^6 * \tan(c) - 24*b^(5/2) * d*x^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^6 * \tan(c) + 24*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 144*b^(5/2) * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 72*b^(5/2) * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 9*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(c)^2 - 18*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(c)^2 + 144*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 54*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 324*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 18*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 144*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 144*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 24*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 3*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 9*b^(5/2) * d*x^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 48*b^(5/2) * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 192*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) * \tan(c)^2 - 1440*b^(5/2) * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 + 1728*b^(5/2) * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan(c)^2 - 384*b^(5/2) * \tan(1/2*d*x + 1/2*c)^2 * \tan(
\end{aligned}$$

$$\begin{aligned}
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 27*b^(5/2) \\
& *d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c \\
&)^2*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 \\
& + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c \\
&)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*b \\
& ^{(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)* \\
& \tan(1/3*c)^2*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 972*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 1 \\
& 62*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2 \\
& *c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3* \\
& c)^2*\tan(c)^2 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(\\
& c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*b^(5/2) \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^2*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d \\
& *x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2* \\
& \tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 9600*b^(5/2)*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 \\
& - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 2560*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 4800*b^ \\
& (5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^3*\tan(c)^2 + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*b^(5/2)*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^3*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5* \\
& \tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 320*b^(5/2)*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 + 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 23328*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^5*\tan(c)^2 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^5*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 192*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*
\end{aligned}$$

$$\begin{aligned}
& c)^6 \tan(c)^2 - 320b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^3 \tan(1/3c)^6 \tan(c)^2 + 432b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^4 \tan(1/3c)^6 \tan(c)^2 - 384b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^5 \tan(1/3c)^6 \tan(c)^2 - 48b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(1/3c)^6 \tan(c)^2 - 48b^{(5/2)} dx^4 \tan(1/2dx + 1/2c) \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c)^2 - 72b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c)^2 - 144b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c)^2 - 24b^{(5/2)} \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c)^2 - 18b^{(5/2)} dx^4 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 + 192b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c) \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 - 2016b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 + 2560b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^3 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 - 2016b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 + 192b^{(5/2)} \tan(1/2dx + 1/6c)^5 \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 - 24b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c)^2 - 144b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c)^2 - 72b^{(5/2)} \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c)^2 - 48b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c)^2 - 384b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c) \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c)^2 + 432b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c)^2 - 320b^{(5/2)} \tan(1/2dx + 1/6c)^3 \tan(-1/2dx + 1/2c)^4 \tan(1/3c)^6 \tan(c)^2 - 9b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 + 18b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^6 - 144b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) + 48b^{(5/2)} dx^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) + 54b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^2 - 324b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^2 + 18b^{(5/2)} dx^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 + 144b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^3 - 144b^{(5/2)} dx^4 \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^3 + 24b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^3 - 3b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c)^4 + 54b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 - 9b^{(5/2)} dx^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^4 - 96b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^5 \tan(-1/2dx + 1/2c)^4 - 48b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^4 + 192b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 \tan(1/3c) - 1440b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) + 1728b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^5 \tan(-1
\end{aligned}$$

$$\begin{aligned}
& /2*d*x + 1/2*c)^4*\tan(1/3*c) - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 27*b^(5/2)*d*x^4*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 162*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 - 9*b^(5/2)*d*x^4*t \\
& an(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& ^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 432*b^(5/2)* \\
& d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan \\
& (1/3*c)^2 + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)*\tan(1/3*c)^2 + 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 972*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 162*b^(5/2)* \\
& d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*b^ \\
& (5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^2 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*b \\
& ^5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^ \\
& 3*\tan(1/3*c)^2 + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c) \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 4800*b^(5/2)*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9936* \\
& b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*t \\
& an(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 320*b^(5/2)*ta \\
& n(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 + 9600*b^(5/2)*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^3 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2 \\
& *d*x + 1/2*c)^2*\tan(1/3*c)^3 + 2560*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 4800*b^(5/2)*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 \\
& + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^3 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + 5760*b^(5/2)*\tan(1/2*d*x \\
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 320* \\
& b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 9*b^(\\
& 5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 - 27*b^(5/2)*d*x^4*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^4 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^5*\tan(1/3*c)^4 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2* \\
& d*x + 1/6*c)^6*\tan(1/3*c)^4 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(\\
& -1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan \\
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*b^(5/2)*\tan(1/
\end{aligned}$$

$$\begin{aligned}
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 \\
& + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9600*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 23328*b^(5/2)*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 144*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 216*b^(5/2)*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 4 \\
& 32*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^3*\tan(1/3*c)^4 + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c \\
&)^3*\tan(1/3*c)^4 - 9*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1 \\
& 440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(1/3*c)^4 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9936*b^(\\
& 5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^4 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^4 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan \\
& (1/3*c)^5 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(\\
& 1/3*c)^5 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/ \\
& 3*c)^5 + 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*b^(5/2)*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
& ^5 - 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^5 + 192*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^5 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^5 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/ \\
& 6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 + 5760*b^(\\
& 5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^5 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*ta \\
& n(1/3*c)^5 + 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 9*b^(5/ \\
& 2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 - 320*b^(5/2)*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 + 432*b^(5/2)*\tan(1/2*d*x + 1/2* \\
& c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)*ta \\
& n(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*t \\
& an(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 144*b^(5/2)*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 +
\end{aligned}$$

$$\begin{aligned}
& 24*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 18*b^{(5/2)}*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 2016*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 2560*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 2016*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 24*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 432*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 320*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(c) + 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 216*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) + 216*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) - 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) - 1296*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 216*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) - 1296*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) + 432*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) - 216*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2)
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 72*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c) - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c) - 24*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c) + 24*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c) + 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c) + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 3*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 96*b^(5/2)*\tan(1/2*d*x + 1/2*c) \\
&)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 - 2880*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^ \\
& 2 + 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 768*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 1440*b^(5/2) \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(\\
& 1/3*c)*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^ \\
& 3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 \\
& - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^2*\tan(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^ \\
& 4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1 \\
& /2*c)^3*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*d*x^4*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 324*b \\
& ^ (5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*ta \\
& n(c)^2 - 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*ta \\
& n(1/3*c)^2*\tan(c)^2 - 9600*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/ \\
& 6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*b^(5/2)*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 2*\tan(c)^2 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*ta \\
& n(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2 \\
& *c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + \\
& 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& *\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(- \\
& 1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c) \\
& ^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 7 \\
& 2*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c \\
&)^2 + 9*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440* \\
& b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 19200*b^(5/2)*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3* \\
& c)^2*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*ta \\
& n(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1 \\
& /6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 4800*b^(5/2)*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*b^(5/2)
\end{aligned}$$

$$\begin{aligned}
&)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 + 128 \\
& 0*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c) \\
& ^2 - 9600*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*t \\
& an(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400 \\
& *b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x \\
& + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 640*b^(5/2)*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 320*b^(5 \\
& /2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - \\
& 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^3*\tan(c)^2 + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 19200*b^(5/2 \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^3*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c \\
&)^4*\tan(1/3*c)^3*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3 \\
& *c)^4*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c \\
&)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c) \\
& ^4*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(1/3*c)^4*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/ \\
& 6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c) \\
& *\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - \\
& 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1 \\
& /2*c)*\tan(1/3*c)^4*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^4*\tan(c)^2 - 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*b^(5/2)*\tan(1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 4*\tan(c)^2 - 38400*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 19872*b^(5/2)*\tan(1/2*d*x + 1 \\
& /2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 \\
& - 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4* \\
& tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2 \\
& *\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + \\
& 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 1008*b^(5/2)*\tan(1 \\
& /2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 5760*b^(5 \\
& /2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c \\
&)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 1440*b^(5/2)*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 5760*b \\
& ^{(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2
\end{aligned}$$

$$\begin{aligned}
& + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5* \\
& \tan(c)^2 - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3* \\
& c)^5*\tan(c)^2 + 96*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^5*\tan(c)^2 - 1 \\
& 92*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(\\
& c)^2 + 3456*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 11520*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*t \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 11520 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^5*\tan(c)^2 - 2880*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(- \\
& 1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c \\
&)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1440 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c) \\
& ^2 + 3*b^{(5/2)}*d*x^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b^{(5/2)}*\tan(1/2*d*x + 1/2*c \\
&)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 - 1008*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 1280*b^{(5/2)}*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 1008*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b \\
& ^{(5/2)}*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^6*\tan(c)^2 - 24*b^{(5/2)}*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 - 144*b^{(5/2)}*\tan(\\
& 1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^ \\
& 6*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/ \\
& 3*c)^6*\tan(c)^2 - 96*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^6*\tan(c)^2 - 768*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 864*b^{(5/2)}*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan \\
& (c)^2 - 640*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^6*\tan(c)^2 - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*t \\
& \tan(1/3*c)^6*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2 \\
& *c)^3*\tan(1/3*c)^6*\tan(c)^2 - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 96*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(-1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 - 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^2 + 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^4 - 3*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^6 - 144*b^{(5/2)}*d* \\
& x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) + 1 \\
& 44*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + \\
& 1/2*c) + 24*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2 \\
& *d*x + 1/2*c) + 18*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2* \\
& c)^2 - 324*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(\\
& -1/2*d*x + 1/2*c)^2 + 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2 - 192*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*\tan(\\
& -1/2*d*x + 1/2*c)^2 - 96*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)^2 + 48*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(- \\
& 1/2*d*x + 1/2*c)^3 - 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^3 + 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2
\end{aligned}$$

$$\begin{aligned}
& *d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3 + 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2 \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3 + 18*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d \\
& *x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3 \\
& *\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 384*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4 - 144*b^(5/2)*\tan(\\
& 1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 96*b^(5/2) \\
&)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c) - 2880*b^(5/2)*t \\
& an(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c) + 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 768*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) + 1440*b^(5/2)*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 5760*b \\
& ^{(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/3*c) + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1 \\
& /2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) + 96*b^(5/2)*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c) - 9*b^(5/2)*d*x^4*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^2 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 - 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(1/3*c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^5*t \\
& an(1/3*c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (1/3*c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)* \\
& \tan(1/3*c)^2 + 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^ \\
& 2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 144*b^(5/2)*\tan(\\
& 1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^ \\
& 2 - 324*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^2 + 162*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*t \\
& an(1/3*c)^2 - 9600*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 19872*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*t \\
& an(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 - 11520*b^(5/2)* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1 \\
& /2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 + 432*b^(5/2)*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 \\
& + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2 - \\
& 9*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440*b^(5/2)*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - \\
& 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^2 + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 - 11664*b^(5/2)*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 1440 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 4800 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 5760* \\
& b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3 + 1280*b \\
& ^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3 - 9600*b^{(5/2)} \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 \\
& *\tan(1/3*c)^3 + 38400*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3 \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 38400*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^3 - 640*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^3 + 320*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^3 - 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^3 + 19200*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 19200*b^{(5/2)}*\tan(1/2 \\
& *d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 4800*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 + \\
& 54*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 27*b^{(5/2)}*d*x^4*ta \\
& n(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 - 11664*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4 - 1008*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(1/3*c)^4 + 144*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(- \\
& 1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 432*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 72* \\
& b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 54*b^{(5 \\
& /2)}*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c \\
&)^2*\tan(1/3*c)^4 - 38400*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c \\
&)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 19872*b^{(5/2)}*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*b^{(5/2)} \\
& *\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 432*b^{(5/2)}* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1 \\
& /3*c)^4 + 216*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/ \\
& 3*c)^4 - 1008*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^4 + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^4 - 11664*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d* \\
& x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 4800*b^{(5/2)}*\tan(1/2*d* \\
& x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 1440*b^{(5/2)}*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 5760*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5 + 5760*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5 - 1728*b^{(5/2)}*\tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^5 + 96*b^(5/2)*\tan(1/2*d*x + 1/6*c) \\
& ^6*\tan(1/3*c)^5 - 192*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c) \\
&)^2*\tan(1/3*c)^5 + 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c) \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 + 11520*b^(5/ \\
& 2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^5 - 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(\\
& 1/3*c)^5 + 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^5 - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^5 + 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(1/3*c)^5 - 3*b^(5/2)*d*x^4*\tan(1/3*c)^6 + 96*b^(5/2)*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 - 1008*b^(5/2)*\tan(1/2 \\
& *d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 1280*b^(5/2)*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 - 1008*b^(5/2)*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 96*b^(5/2)*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(1/3*c)^6 + 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)*\tan(1/3*c)^6 + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(- \\
& 1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 768*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 864*b^(5/2)*\tan(1/2*d \\
& *x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 - 6 \\
& 40*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 48 \\
& *b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 + 72*b \\
& ^5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 144*b^ \\
& (5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 96*b^(5/2) \\
&)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 - 72*b^(5/2)*d* \\
& x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c) + 72*b^(5/2)*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*b^(5/2)*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 432*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 72*b^(5/2)*d*x \\
& ^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 144*b^(5/2)*d*x^4* \\
& tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 432*b^(5/2)*d*x^4*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) + \\
& 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) - \\
& 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c) + 24 \\
& *b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^(\\
& 5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4* \\
& tan(c) + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(c) + 24*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(c) - 72*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c) \\
&) + 216*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) \\
&)^2*\tan(c) - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& 1/3*c)^2*\tan(c) - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6
\end{aligned}$$

$$\begin{aligned}
& * \tan(1/3*c)^2 * \tan(c) - 432*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c) + 216*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c) - 432*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^2 * \tan(c) - 72*b^{(5/2)} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^2 * \tan(c) + 72*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + 432*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + 216*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^2 * \tan(c) + 72*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) - 216*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^4 * \tan(c) - 432*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^4 * \tan(c) - 72*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^6 * \tan(1/3*c)^4 * \tan(c) + 72*b^{(5/2)} * d*x^4 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c) + 144*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) + 216*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^4 * \tan(c) - 24*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) - 144*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^6 * \tan(c) - 72*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c)^6 * \tan(c) + 24*b^{(5/2)} * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c)^6 * \tan(c) + 3*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 54*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(c)^2 + 9*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(c)^2 - 96*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^5 * \tan(c)^2 - 48*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^6 * \tan(c)^2 + 48*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 144*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 72*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 - 48*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c) * \tan(c)^2 + 108*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 54*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 640*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 2016*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 768*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 - 288*b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(c)^2 + 48*b^{(5/2)} * d*x^4 * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 72*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 144*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 - 24*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^6 * \tan(-1/2*d*x + 1/2*c)^3 * \tan(c)^2 + 3*b^{(5/2)} * d*x^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 432*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 1280*b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^3 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 + 432*b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 96*b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^5 * \tan(-1/2*d*x + 1/2*c)^4 * \tan(c)^2 - 1440*b^{(5/2)} * \tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)*\tan(c)^2 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 192*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 27*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 4800*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 - 144*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 54*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 23328*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 38400*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 23328*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 - 4800*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^3*\tan(c)^2 - 320*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^3*\tan(c)^2 + 640*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 38400*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 38400*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 9600*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 1280*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 + 9*b^(5/2)*d*x^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 9936*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^4*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 23328*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9600*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^5*\tan(c)^2 - 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5*\tan(c)^2 + 768*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 3456*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 + 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 96*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5*\tan(c)^2 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^6*\tan(c)^2 - 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 - 320*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)
\end{aligned}$$

$$\begin{aligned}
&)^6 \tan(c)^2 - 48b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(-1/2dx + 1/2c) \tan(1/3c)^6 \tan(c)^2 - 72b^{(5/2)} \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c) \\
& \tan(1/3c)^6 \tan(c)^2 - 288b^{(5/2)} \tan(1/2dx + 1/2c) \tan(-1/2dx + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 + 192b^{(5/2)} \tan(1/2dx + 1/6c) \tan(-1/2dx \\
& x + 1/2c)^2 \tan(1/3c)^6 \tan(c)^2 - 24b^{(5/2)} \tan(-1/2dx + 1/2c)^3 \tan(1/3c)^6 \tan(c)^2 - 3b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^4 + 54b^{(5/2)} dx \\
& x^4 \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 - 9b^{(5/2)} dx^4 \tan(1/2dx + 1/6c)^4 - 96b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^5 \\
& - 48b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^6 - 48b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^3 \tan(-1/2dx + 1/2c) + 144b^{(5/2)} dx^4 \tan(1/2 \\
& dx + 1/2c) \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c) + 72b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c) + 48b^{(5/2)} \\
& \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c) - 108b^{(5/2)} dx^4 \tan(1/2dx + 1/2c)^2 \tan(-1/2dx + 1/2c)^2 + 54b^{(5/2)} \\
& dx^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^2 + 640b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^3 \tan(-1/2dx + 1/2c)^2 - 2016b^{(5/2)} \\
& \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^2 + 768b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^5 \tan(-1/2dx + 1/ \\
& 2c)^2 - 288b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 - 48b^{(5/2)} dx^4 \tan(1/2dx + 1/2c) \tan(-1/2dx + 1/2c) \\
& ^3 + 72b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^3 + 144b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(\\
& -1/2dx + 1/2c)^3 + 24b^{(5/2)} \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^3 - 3b^{(5/2)} dx^4 \tan(-1/2dx + 1/2c)^4 - 96b^{(5/2)} \tan(1/2dx + 1 \\
& /2c)^4 \tan(1/2dx + 1/6c) \tan(-1/2dx + 1/2c)^4 + 432b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 - 1280b^{(5/2)} \\
& \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^3 \tan(-1/2dx + 1/2c)^4 + 432b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c \\
&)^4 - 96b^{(5/2)} \tan(1/2dx + 1/6c)^5 \tan(-1/2dx + 1/2c)^4 - 1440b^{(5/2)} \tan(1/2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^4 \tan(1/3c) + 1728b^{(5/2)} \\
& \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^5 \tan(1/3c) - 384b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^6 \tan(1/3c) + 2880b^{(5/2)} \tan(1/ \\
& 2dx + 1/2c)^4 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^2 \tan(1/3c) - 11520b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c)^3 \tan(-1/2dx \\
& + 1/2c)^2 \tan(1/3c) + 11520b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^2 \tan(1/3c) - 3456b^{(5/2)} \tan(1/2dx + 1/ \\
& 2c) \tan(1/2dx + 1/6c)^5 \tan(-1/2dx + 1/2c)^2 \tan(1/3c) + 192b^{(5/2)} \tan(1/2dx + 1/6c)^6 \tan(-1/2dx + 1/2c)^2 \tan(1/3c) - 96b^{(5/2)} \tan \\
& (1/2dx + 1/2c)^4 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) + 1728b^{(5/2)} \tan(1/2dx + 1/2c)^3 \tan(1/2dx + 1/6c) \tan(-1/2dx + 1/2c)^4 \tan(1/3c) \\
& - 5760b^{(5/2)} \tan(1/2dx + 1/2c)^2 \tan(1/2dx + 1/6c)^2 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) + 5760b^{(5/2)} \tan(1/2dx + 1/2c) \tan(1/2dx + 1/6c \\
& c)^3 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) - 1440b^{(5/2)} \tan(1/2dx + 1/6c)^4 \tan(-1/2dx + 1/2c)^4 \tan(1/3c) + 54b^{(5/2)} dx^4 \tan(1/2dx + 1/2c \\
&)^2 \tan(1/3c)^2 - 27b^{(5/2)} dx^4 \tan(1/2dx + 1/6c)^2 \tan(1/3c)^2 -
\end{aligned}$$

$$\begin{aligned}
& 4800b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^3\tan(1/3c)^2 + 9936b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^4\tan(1/3c)^2 - 5760b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^5\tan(1/3c)^2 + 432b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^6\tan(1/3c)^2 + 144b^{(5/2)}dx^4\tan(1/2dx + 1/2c)\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 216b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 432b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 72b^{(5/2)}\tan(1/2dx + 1/6c)^6\tan(-1/2dx + 1/2c)\tan(1/3c)^2 + 54b^{(5/2)}dx^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 2880b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 - 23328b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 38400b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^3\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 - 23328b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 2880b^{(5/2)}\tan(1/2dx + 1/6c)^5\tan(-1/2dx + 1/2c)^2\tan(1/3c)^2 + 72b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(-1/2dx + 1/2c)^3\tan(1/3c)^2 + 432b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^3\tan(1/3c)^2 + 216b^{(5/2)}\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)^3\tan(1/3c)^2 + 432b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 - 5760b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 + 9936b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 - 4800b^{(5/2)}\tan(1/2dx + 1/6c)^3\tan(-1/2dx + 1/2c)^4\tan(1/3c)^2 - 4800b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)^2\tan(1/3c)^3 + 19200b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^3\tan(1/3c)^3 - 19200b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^4\tan(1/3c)^3 + 5760b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^5\tan(1/3c)^3 - 320b^{(5/2)}\tan(1/2dx + 1/6c)^6\tan(1/3c)^3 + 640b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^3 - 11520b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)\tan(-1/2dx + 1/2c)^2\tan(1/3c)^3 + 38400b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^2\tan(1/3c)^3 - 38400b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^3\tan(-1/2dx + 1/2c)^2\tan(1/3c)^3 + 9600b^{(5/2)}\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)^2\tan(1/3c)^3 - 1280b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(-1/2dx + 1/2c)^4\tan(1/3c)^3 + 5760b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)\tan(-1/2dx + 1/2c)^4\tan(1/3c)^3 - 4800b^{(5/2)}\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)^4\tan(1/3c)^3 - 9b^{(5/2)}dx^4\tan(1/3c)^4 - 1440b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(1/2dx + 1/6c)\tan(1/3c)^4 + 9936b^{(5/2)}\tan(1/2dx + 1/2c)^3\tan(1/2dx + 1/6c)^2\tan(1/3c)^4 - 19200b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^3\tan(1/3c)^4 + 9936b^{(5/2)}\tan(1/2dx + 1/2c)\tan(1/2dx + 1/6c)^4\tan(1/3c)^4 - 1440b^{(5/2)}\tan(1/2dx + 1/6c)^5\tan(1/3c)^4 + 72b^{(5/2)}\tan(1/2dx + 1/2c)^4\tan(-1/2dx + 1/2c)\tan(1/3c)^4 + 432b^{(5/2)}\tan(1/2dx + 1/2c)^2\tan(1/2dx + 1/6c)^2\tan(-1/2dx + 1/2c)\tan(1/3c)^4 + 216b^{(5/2)}\tan(1/2dx + 1/6c)^4\tan(-1/2dx + 1/2c)\tan(1/3c)^4
\end{aligned}$$

$$\begin{aligned}
& c)^4 - 2016*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^4 + 11520*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)^4 - 23328*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x \\
& + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9600*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 144*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 216*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 432*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 1440*b^{(5/2)}*\tan(1/2*d*x + 1/ \\
& 6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 - 96*b^{(5/2)}*\tan(1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^5 + 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)* \\
& \tan(1/3*c)^5 - 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*t \\
& an(1/3*c)^5 + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(\\
& 1/3*c)^5 - 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^5 + 768*b^{(5/2)}*t \\
& an(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 3456*b^{(5/2)}*t \\
& an(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
& ^5 + 2880*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
& ^5 - 96*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^5 - 48*b^{(5/2)}*\tan(1/2*d \\
& *x + 1/2*c)^3*\tan(1/3*c)^6 - 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)*\tan(1/3*c)^6 + 432*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6 \\
& *c)^2*\tan(1/3*c)^6 - 320*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^6 + 48*b \\
& ^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 72*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 - 288*b^{(5/2)}* \\
& \tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 192*b^{(5/2)}*\tan \\
& (1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 24*b^{(5/2)}*\tan(-1/ \\
& 2*d*x + 1/2*c)^3*\tan(1/3*c)^6 - 24*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)^3*\tan \\
& (c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(c) - \\
& 72*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 48*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(c) - 144*b^{(5/2)}*d*x^4 \\
& *\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan \\
& (1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c) - 144*b^{(5/2)}*d*x^4*\tan(1/ \\
& 2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c) - 24*b^{(5/2)}*d*x^4*\tan(-1/2*d \\
& *x + 1/2*c)^3*\tan(c) + 24*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(c) + 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*t \\
& an(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d \\
& *x + 1/2*c)^4*\tan(c) + 72*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^2*t \\
& an(c) - 216*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c \\
&)^2*\tan(c) - 432*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(\\
& 1/3*c)^2*\tan(c) - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c) + 7 \\
& 2*b^{(5/2)}*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c) + 144*b^{(5/2)}*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) + 216*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 72* \\
& b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c) - 432*b^{(5/2)}*\tan(1/2*d* \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 216*b^{(5/2)}*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c) + 72*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^4*\tan(c) - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)
\end{aligned}$$

$$\begin{aligned}
& - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c) - 18*b^{(5/2)}*d*x^4 \\
& * \tan(1/2*d*x + 1/2*c)^2*\tan(c)^2 + 9*b^{(5/2)}*d*x^4*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(c)^2 + 320*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3*\tan(c)^2 \\
& - 1008*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 3 \\
& 84*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5*\tan(c)^2 - 144*b^{(5/2)} \\
& * \tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 - 48*b^{(5/2)}*d*x^4 \\
& * \tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144* \\
& b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& * \tan(c)^2 - 24*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 \\
& - 18*b^{(5/2)}*d*x^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192*b^{(5/2)}*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 864 \\
& *b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) \\
&)^2*\tan(c)^2 - 2560*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 864*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2* \\
& d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/6*c)^5*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 24*b^{(5/2)}*\tan(1/2*d*x + 1/2*c) \\
&)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2 \\
& *d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 144*b^{(5/2)}*\tan(1/2*d*x \\
& + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 384*b^{(5/2)}*\tan(1/2*d*x + 1/2 \\
& *c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 1008*b^{(5/2)}* \\
& \tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 \\
& + 320*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 1 \\
& 440*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)*\tan(c) \\
& ^2 - 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)* \\
& \tan(c)^2 + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1 \\
& /3*c)*\tan(c)^2 - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^5* \\
& \tan(1/3*c)*\tan(c)^2 + 96*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)*\tan(c)^2 \\
& - 192*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan \\
& (c)^2 + 3456*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d \\
& *x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 11520*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 + 11520*b^{(5/2)} \\
& * \tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)*\tan(c)^2 - 2880*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2 \\
& *c)^2*\tan(1/3*c)*\tan(c)^2 + 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2 \\
& *d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1440*b^{(5/2)}* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 9*b^{(5/2)} \\
&)*d*x^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)*\tan(1/3*c)^2*\tan(c)^2 - 11664*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3* \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19200*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 11664*b^{(5/2)}*\tan(1 \\
& /2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^{(5/2)} \\
& * \tan(1/2*d*x + 1/6*c)^5*\tan(1/3*c)^2*\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 432*b^(5/2)*\tan(1/2*d*x \\
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 19872*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 9600*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^2*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3*\tan(c)^2 + 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 19200*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 2560*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 - 9600*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3*\tan(c)^2 + 320*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3*\tan(c)^2 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4*\tan(c)^2 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4*\tan(c)^2 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 - 72*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4*\tan(c)^2 + 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5*\tan(c)^2 + 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 192*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 + 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6*\tan(c)^2 - 24*b^(5/2)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6*\tan(c)^2 + 18*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)^2 - 9*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/6*c)^2 + 320*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^3 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^4 + 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^5 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^6 + 48*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c) + 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) + 24*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) + 18*b^(5/2)*d*x^4*\tan(-1/2*d*x
\end{aligned}$$

$$\begin{aligned}
& + 1/2*c)^2 - 192*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^2 + 864*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^2 - 2560*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^3*tan(-1/2*d*x + 1/2*c)^2 + 864*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^2 - 192*b^(5/2)*tan(1/2*d*x + 1/6*c)^5*tan(-1/2*d*x + 1/2*c)^2 + 24*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(-1/2*d*x + 1/2*c)^3 + 144*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^3 + 72*b^(5/2)*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^3 - 144*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(-1/2*d*x + 1/2*c)^4 + 384*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^4 - 1008*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^4 + 320*b^(5/2)*tan(1/2*d*x + 1/6*c)^3*tan(-1/2*d*x + 1/2*c)^4 + 1440*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)^2*tan(1/3*c) - 5760*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^3*tan(1/3*c) + 5760*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^4*tan(1/3*c) - 1728*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^5*tan(1/3*c) + 96*b^(5/2)*tan(1/2*d*x + 1/6*c)^6*tan(1/3*c) - 192*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c) + 3456*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c) - 11520*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c) + 11520*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^3*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c) - 2880*b^(5/2)*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c) + 384*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c) - 1728*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c) + 1440*b^(5/2)*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c) - 9*b^(5/2)*d*x^4*tan(1/3*c)^2 + 1440*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(1/2*d*x + 1/6*c)*tan(1/3*c)^2 - 11664*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)^2*tan(1/3*c)^2 + 19200*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^3*tan(1/3*c)^2 - 11664*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^4*tan(1/3*c)^2 + 1440*b^(5/2)*tan(1/2*d*x + 1/6*c)^5*tan(1/3*c)^2 + 72*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(-1/2*d*x + 1/2*c)*tan(1/3*c)^2 + 432*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)*tan(1/3*c)^2 + 216*b^(5/2)*tan(1/2*d*x + 1/6*c)^4*tan(-1/2*d*x + 1/2*c)*tan(1/3*c)^2 + 864*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^2 - 11520*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^2 + 19872*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^2 - 9600*b^(5/2)*tan(1/2*d*x + 1/6*c)^3*tan(-1/2*d*x + 1/2*c)^2*tan(1/3*c)^2 + 144*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^2 + 216*b^(5/2)*tan(1/2*d*x + 1/6*c)^2*tan(-1/2*d*x + 1/2*c)^3*tan(1/3*c)^2 - 1008*b^(5/2)*tan(1/2*d*x + 1/2*c)*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^2 + 1440*b^(5/2)*tan(1/2*d*x + 1/6*c)*tan(-1/2*d*x + 1/2*c)^4*tan(1/3*c)^2 + 320*b^(5/2)*tan(1/2*d*x + 1/2*c)^4*tan(1/3*c)^3 - 5760*b^(5/2)*tan(1/2*d*x + 1/2*c)^3*tan(1/2*d*x + 1/6*c)*tan(1/3*c)^3 + 19200*b^(5/2)*tan(1/2*d*x + 1/2*c)^2*tan(1/2*d*x + 1/6*c)^2*tan(1/3*c)^3 - 19200*b^(5/2)*tan(1/2*d*x +
\end{aligned}$$

$$\begin{aligned}
& 1/2*c)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^3 + 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^3 - 2560*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 11520*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 - 9600*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^3 + 320*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^3 - 1008*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 5760*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 - 11664*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 4800*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^4 + 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^4 + 864*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 - 2880*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 72*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^3*\tan(1/3*c)^4 + 384*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 1728*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^5 + 1440*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^5 - 192*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^5 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^6 + 24*b^(5/2)*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^6 + 24*b^(5/2)*d*x^4*\tan(1/2*d*x + 1/2*c)*\tan(c) - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c) - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(c) - 24*b^(5/2)*\tan(1/2*d*x + 1/6*c)^6*\tan(c) + 24*b^(5/2)*d*x^4*\tan(-1/2*d*x + 1/2*c)*\tan(c) + 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) + 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c) - 72*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c) + 72*b^(5/2)*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c) - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c) - 216*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c) - 24*b^(5/2)*\tan(1/3*c)^6*\tan(c) + 3*b^(5/2)*d*x^4*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 - 1280*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^3*\tan(c)^2 + 432*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)^5*\tan(c)^2 - 24*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 144*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)*\tan(c)^2 - 288*b^(5/2)*\tan(1/2*d*x + 1/2*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 768*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 2016*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 640*b^(5/2)*\tan(1/2*d*x + 1/6*c)^3*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 72*b^(5/2)*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 - 48*b^(5/2)*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 - 96*b^(5/2)*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)*\tan(c)^2 + 1728*b^(5/2)
\end{aligned}$$

$$\begin{aligned}
& /2) * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c) * \tan(1/3*c) * \tan(c)^2 - 5760 * \\
& b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c) * \tan(c)^2 + \\
& 5760 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^3 * \tan(1/3*c) * \tan(c) \\
& ^2 - 1440 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c) * \tan(c)^2 + 768 * b^{(5/2)} * \\
& \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) * \tan(c)^2 - 3456 * b \\
& ^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan \\
& (1/3*c) * \tan(c)^2 + 2880 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2 * \\
& c)^2 * \tan(1/3*c) * \tan(c)^2 - 96 * b^{(5/2)} * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) * \tan \\
& (c)^2 + 432 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^2 * \tan(c)^2 - 5760 * b^{(5/2)} \\
& * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c) * \tan(1/3*c)^2 * \tan(c)^2 + 9 \\
& 936 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^2 * \tan(c) \\
& ^2 - 4800 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^3 * \tan(1/3*c)^2 * \tan(c)^2 - 144 * b^{(5/2)} \\
&) * \tan(1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c)^2 - 216 * \\
& b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^2 * \tan(c)^2 \\
& - 2016 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^2 * \tan \\
& (c)^2 + 2880 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3 * \\
& c)^2 * \tan(c)^2 - 72 * b^{(5/2)} * \tan(-1/2*d*x + 1/2*c)^3 * \tan(1/3*c)^2 * \tan(c)^2 - \\
& 1280 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/3*c)^3 * \tan(c)^2 + 5760 * b^{(5/2)} * \tan \\
& (1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c) * \tan(1/3*c)^3 * \tan(c)^2 - 4800 * b^{(5/2)} \\
&) * \tan(1/2*d*x + 1/6*c)^2 * \tan(1/3*c)^3 * \tan(c)^2 + 640 * b^{(5/2)} * \tan(-1/2*d*x + \\
& 1/2*c)^2 * \tan(1/3*c)^3 * \tan(c)^2 + 432 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/3 * \\
& c)^4 * \tan(c)^2 - 1440 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c) * \tan(1/3*c)^4 * \tan(c)^2 - 7 \\
& 2 * b^{(5/2)} * \tan(-1/2*d*x + 1/2*c) * \tan(1/3*c)^4 * \tan(c)^2 - 96 * b^{(5/2)} * \tan(1/3 * \\
& c)^5 * \tan(c)^2 - 3 * b^{(5/2)} * d*x^4 - 96 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(1/2 \\
& * d*x + 1/6*c) + 432 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1/6*c)^2 - \\
& 1280 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^3 + 432 * b^{(5/2)} * \tan \\
& (1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^4 - 96 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c \\
&)^5 + 24 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^4 * \tan(-1/2*d*x + 1/2*c) + 144 * b^{(5/2)} \\
& * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c) + 72 * b \\
& ^{(5/2)} * \tan(1/2*d*x + 1/6*c)^4 * \tan(-1/2*d*x + 1/2*c) - 288 * b^{(5/2)} * \tan(1/2*d \\
& * x + 1/2*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 + 768 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \\
& \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^2 - 2016 * b^{(5/2)} * \tan(1/2*d*x + 1 \\
& /2*c) * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 + 640 * b^{(5/2)} * \tan(1/2 * \\
& d*x + 1/6*c)^3 * \tan(-1/2*d*x + 1/2*c)^2 + 48 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \\
& \tan(-1/2*d*x + 1/2*c)^3 + 72 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + \\
& 1/2*c)^3 - 48 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(-1/2*d*x + 1/2*c)^4 - 96 * b^{(5/2)} \\
& * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^4 - 96 * b^{(5/2)} * \tan(1/2*d*x \\
& + 1/2*c)^4 * \tan(1/3*c) + 1728 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^3 * \tan(1/2*d*x + 1 \\
& /6*c) * \tan(1/3*c) - 5760 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c)^2 * \tan(1/2*d*x + 1/6*c) \\
& ^2 * \tan(1/3*c) + 5760 * b^{(5/2)} * \tan(1/2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c)^3 * \tan \\
& (1/3*c) - 1440 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^4 * \tan(1/3*c) + 768 * b^{(5/2)} * \tan \\
& (1/2*d*x + 1/2*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) - 3456 * b^{(5/2)} * \tan(1 \\
& /2*d*x + 1/2*c) * \tan(1/2*d*x + 1/6*c) * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) + 2 \\
& 880 * b^{(5/2)} * \tan(1/2*d*x + 1/6*c)^2 * \tan(-1/2*d*x + 1/2*c)^2 * \tan(1/3*c) - 96 * \\
& b^{(5/2)} * \tan(-1/2*d*x + 1/2*c)^4 * \tan(1/3*c) + 432 * b^{(5/2)} * \tan(1/2*d*x + 1/2 *
\end{aligned}$$

$$\begin{aligned}
& c)^3 \tan(1/3*c)^2 - 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) \\
&)*\tan(1/3*c)^2 + 9936*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan \\
& \tan(1/3*c)^2 - 4800*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^3*\tan(1/3*c)^2 + 144*b^{(5/2)} \\
&)*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 216*b^{(5/2)}*\tan \\
& \tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 - 2016*b^{(5/2)}*\tan \\
& (1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 2880*b^{(5/2)}*\tan(1 \\
& /2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 72*b^{(5/2)}*\tan(-1/2* \\
& d*x + 1/2*c)^3*\tan(1/3*c)^2 - 1280*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c) \\
&)^3 + 5760*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^3 - \\
& 4800*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^3 + 640*b^{(5/2)}*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^3 + 432*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/3*c)^4 - \\
& 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^4 + 72*b^{(5/2)}*\tan(-1/2*d*x + \\
& 1/2*c)*\tan(1/3*c)^4 - 96*b^{(5/2)}*\tan(1/3*c)^5 - 24*b^{(5/2)}*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(c) - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2* \\
& \tan(c) - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^4*\tan(c) + 24*b^{(5/2)}*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(c) - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c) \\
& - 216*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c) - 72*b^{(5/2)}*\tan(\\
& 1/3*c)^4*\tan(c) - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*b^{(5/2)} \\
& *\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 1008*b^{(5/2)}*\tan(1/ \\
& 2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + 320*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/6*c)^3*\tan(c)^2 - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) \\
& *\tan(c)^2 - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 192* \\
& b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 - 24*b^{(5/2)}* \\
& \tan(-1/2*d*x + 1/2*c)^3*\tan(c)^2 + 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)*\tan(c)^2 - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)*\tan \\
& (1/3*c)*\tan(c)^2 + 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)*\tan(c)^2 \\
& - 192*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)*\tan(c)^2 - 1008*b^{(5/2)}*\tan \\
& (1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan(c)^2 + 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c) \\
&)*\tan(1/3*c)^2*\tan(c)^2 - 72*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2*\tan \\
& (c)^2 + 320*b^{(5/2)}*\tan(1/3*c)^3*\tan(c)^2 - 144*b^{(5/2)}*\tan(1/2*d*x + 1/2*c) \\
&)^3 + 384*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c) - 1008*b^{(5/2)} \\
&)*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + 1/6*c)^2 + 320*b^{(5/2)}*\tan(1/2*d*x + 1 \\
& /6*c)^3 + 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c) + 72*b^{(5 \\
& /2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c) - 96*b^{(5/2)}*\tan(1/2*d*x + \\
& 1/2*c)*\tan(-1/2*d*x + 1/2*c)^2 - 192*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(-1/2 \\
& *d*x + 1/2*c)^2 + 24*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c)^3 + 384*b^{(5/2)}*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c) - 1728*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(1/2*d*x + \\
& 1/6*c)*\tan(1/3*c) + 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c) - 192*b \\
& ^{(5/2)}*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c) - 1008*b^{(5/2)}*\tan(1/2*d*x + 1/2* \\
& c)*\tan(1/3*c)^2 + 1440*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(1/3*c)^2 + 72*b^{(5/ \\
& 2)}*\tan(-1/2*d*x + 1/2*c)*\tan(1/3*c)^2 + 320*b^{(5/2)}*\tan(1/3*c)^3 - 48*b^{(5/ \\
& 2)}*\tan(1/2*d*x + 1/2*c)^2*\tan(c) - 72*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)^2*\tan(c) \\
& - 72*b^{(5/2)}*\tan(1/3*c)^2*\tan(c) - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c)*\tan(c)^ \\
& 2 - 96*b^{(5/2)}*\tan(1/2*d*x + 1/6*c)*\tan(c)^2 - 24*b^{(5/2)}*\tan(-1/2*d*x + 1/
\end{aligned}$$

$$\begin{aligned}
& 2*c)*\tan(c)^2 - 96*b^{(5/2)}*\tan(1/3*c)*\tan(c)^2 - 48*b^{(5/2)}*\tan(1/2*d*x + 1/2*c) \\
& /2*c) - 96*b^{(5/2)}*\tan(1/2*d*x + 1/6*c) + 24*b^{(5/2)}*\tan(-1/2*d*x + 1/2*c) \\
& - 96*b^{(5/2)}*\tan(1/3*c) - 24*b^{(5/2)}*\tan(c))/(d*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& 1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/ \\
& 2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^ \\
& 6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^ \\
& 4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 2* \\
& d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(\\
& -1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + \\
& 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4 \\
& *\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2* \\
& d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 \\
& + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 6*d* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1 \\
& /3*c)^4*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3* \\
& c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d \\
& *x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d \\
& *x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6* \\
& \tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan \\
& (1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 6* \\
& d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^4 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + \\
& 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c) \\
& ^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/6 \\
& *c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + \\
& 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6* \\
& \tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4* \\
& \tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4* \\
\end{aligned}$$

$$\begin{aligned}
& \text{an}(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 12*d* \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1 \\
& /3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1 \\
& /2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 \\
& *\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + \\
& 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 9*d*t \\
& \text{an}(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\text{ta} \\
& \text{n}(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c \\
&)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 3*d \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4 + 3*d*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 12*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 2*d \\
& *\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + d*\tan(1/2*d* \\
& x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c \\
&)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 2*d*\tan(1/2*d*x + \\
& 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(\\
& 1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 \\
& + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4 \\
& *\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2* \\
& \tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x \\
& + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(\\
& c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/ \\
& 2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6
\end{aligned}$$

$$\begin{aligned}
& 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4* \\
& \tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 9*d* \\
& \tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*t \\
& \tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 18*d*t \\
& \tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^2*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1 \\
& /2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1 \\
& /2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(1/ \\
& 2*d*x + 1/6*c)^6*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(1 \\
& /2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + d*\tan(1/2 \\
& *d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c \\
&)^6*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c \\
&)^6*\tan(c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6*\tan(c)^2 + d*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2* \\
& d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2 \\
& *d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1 \\
& /2*d*x + 1/2*c)^4 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4*\tan(1 \\
& /3*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 + \\
& 18*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2* \\
& \tan(1/3*c)^2 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + 1/2 \\
& *c)^2*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan \\
& (1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c) \\
& ^4*\tan(1/3*c)^2 + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3 \\
& *c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 3 \\
& *d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 36*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/6*c)^4* \\
& \tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2* \\
& d*x + 1/2*c)^4*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2 \\
& *c)^4*\tan(1/3*c)^4 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 6*d*\tan(1/2*d*
\end{aligned}$$

$$\begin{aligned}
& x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c) \\
& ^4*\tan(1/3*c)^6 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/ \\
& 3*c)^6 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + \\
& d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2 \\
& *d*x + 1/6*c)^4*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^ \\
& 6*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x \\
& + 1/2*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*t \\
& an(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c \\
&)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2* \\
& c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 36*d*\tan(1/2*d*x + 1/2 \\
& *c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6 \\
& *c)^4*\tan(1/3*c)^4*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/ \\
& 2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1 \\
& /2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^4*t \\
& an(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2*d*x \\
& + 1/6*c)^2*\tan(1/3*c)^6*\tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^ \\
& 6*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^4 + 2*d*\tan(1/ \\
& 2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^6 + 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 + 2*d*\tan(1/2*d*x + 1/6*c)^6*\tan \\
& (-1/2*d*x + 1/2*c)^2 + d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 6 \\
& *d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + \\
& 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^4 + 9*d*\tan(1/2*d*x + 1/2* \\
& c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(\\
& 1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^6*\tan(1/3*c)^2 + \\
& 6*d*\tan(1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 36*d*\tan \\
& (1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3* \\
& c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 6 \\
& *d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 9*d*\tan(1/ \\
& 2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 3*d*\tan(1/2*d*x + 1 \\
& /2*c)^4*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*t \\
& an(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^4 + 12*d*\tan(1/2*d*x + \\
& 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 18*d*\tan(1/2*d*x + 1/6*c)^2 \\
& *\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3 \\
& *c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x + 1/6*c)^
\end{aligned}$$

$$\begin{aligned}
& 2*\tan(1/3*c)^6 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^6 + 3*d*\tan(1/2*d*x \\
& + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan \\
& (1/2*d*x + 1/6*c)^4*\tan(c)^2 + d*\tan(1/2*d*x + 1/6*c)^6*\tan(c)^2 + 2*d*\tan \\
& (1/2*d*x + 1/2*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1 \\
& /2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 6*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2* \\
& c)^2*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)^4*\tan(c)^2 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 \\
& + 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 9*d*\tan(1/2*d*x + 1/6*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 12*d*\tan(1/2*d*x + 1/ \\
& 2*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 18*d*\tan(1/2*d*x + 1 \\
& /6*c)^2*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(-1/2*d*x + \\
& 1/2*c)^4*\tan(1/3*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4*\tan \\
& (c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4*\tan(c)^2 + 6*d*\tan(-1/2*d* \\
& x + 1/2*c)^2*\tan(1/3*c)^4*\tan(c)^2 + d*\tan(1/3*c)^6*\tan(c)^2 + 3*d*\tan(1/2* \\
& d*x + 1/2*c)^4*\tan(1/2*d*x + 1/6*c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2* \\
& d*x + 1/6*c)^4 + d*\tan(1/2*d*x + 1/6*c)^6 + 2*d*\tan(1/2*d*x + 1/2*c)^4*\tan(\\
& -1/2*d*x + 1/2*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan \\
& (-1/2*d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^4*\tan(-1/2*d*x + 1/2*c)^2 \\
& + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/ \\
& 6*c)^2*\tan(-1/2*d*x + 1/2*c)^4 + 3*d*\tan(1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + \\
& 18*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^4*\tan(1/3*c)^2 + 12*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2*d*x + \\
& 1/2*c)^2*\tan(1/3*c)^2 + 18*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^2 + 3*d*\tan(-1/2*d*x + 1/2*c)^4*\tan(1/3*c)^2 + 6*d*\tan(1/2*d*x \\
& + 1/2*c)^2*\tan(1/3*c)^4 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^4 + 6*d*\tan \\
& (-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^4 + d*\tan(1/3*c)^6 + d*\tan(1/2*d*x + 1/2*c \\
&)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + 1/6*c)^2*\tan(c)^2 + \\
& 3*d*\tan(1/2*d*x + 1/6*c)^4*\tan(c)^2 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2* \\
& d*x + 1/2*c)^2*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^ \\
& 2*\tan(c)^2 + d*\tan(-1/2*d*x + 1/2*c)^4*\tan(c)^2 + 6*d*\tan(1/2*d*x + 1/2*c)^ \\
& 2*\tan(1/3*c)^2*\tan(c)^2 + 9*d*\tan(1/2*d*x + 1/6*c)^2*\tan(1/3*c)^2*\tan(c)^2 \\
& + 6*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2*\tan(c)^2 + 3*d*\tan(1/3*c)^4*\tan(\\
& c)^2 + d*\tan(1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/2*d*x + \\
& 1/6*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^4 + 4*d*\tan(1/2*d*x + 1/2*c)^2*\tan(-1/2 \\
& *d*x + 1/2*c)^2 + 6*d*\tan(1/2*d*x + 1/6*c)^2*\tan(-1/2*d*x + 1/2*c)^2 + d*\tan \\
& (-1/2*d*x + 1/2*c)^4 + 6*d*\tan(1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + 9*d*\tan(1 \\
& /2*d*x + 1/6*c)^2*\tan(1/3*c)^2 + 6*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(1/3*c)^2 + \\
& 3*d*\tan(1/3*c)^4 + 2*d*\tan(1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(1/2*d*x + \\
& 1/6*c)^2*\tan(c)^2 + 2*d*\tan(-1/2*d*x + 1/2*c)^2*\tan(c)^2 + 3*d*\tan(1/3*c)^ \\
& 2*\tan(c)^2 + 2*d*\tan(1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/2*d*x + 1/6*c)^2 + 2*d* \\
& \tan(-1/2*d*x + 1/2*c)^2 + 3*d*\tan(1/3*c)^2 + d*\tan(c)^2 + d
\end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.79

$$\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2} dx = \frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2),x)

[Out] (b^2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*d*(cos(2*c + 2*d*x) + 1))

3.163 $\int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx$

Optimal result	918
Rubi [A] (verified)	918
Mathematica [A] (verified)	919
Maple [A] (verified)	919
Fricas [A] (verification not implemented)	920
Sympy [F(-1)]	920
Maxima [A] (verification not implemented)	920
Giac [F]	921
Mupad [B] (verification not implemented)	921

Optimal result

Integrand size = 23, antiderivative size = 69

$$\int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx = \frac{b^2 x \sqrt{b \cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b^2 \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d}$$

[Out] $1/2*b^2*x*(b*\cos(d*x+c))^{(1/2)}/\cos(d*x+c)^{(1/2)}+1/2*b^2*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}*(b*\cos(d*x+c))^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\sqrt{\cos(c+dx)}} dx = \frac{b^2 x \sqrt{b \cos(c+dx)}}{2\sqrt{\cos(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}{2d}$$

[In] $\text{Int}[(b*\text{Cos}[c + d*x])^{(5/2)}/\text{Sqrt}[\text{Cos}[c + d*x]],x]$

[Out] $(b^2*x*\text{Sqrt}[b*\text{Cos}[c + d*x]])/(2*\text{Sqrt}[\text{Cos}[c + d*x]]) + (b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(2*d)$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b$

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \cos^2(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} + \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int 1 dx}{2\sqrt{\cos(c + dx)}} \\ &= \frac{b^2 x \sqrt{b \cos(c + dx)}}{2\sqrt{\cos(c + dx)}} + \frac{b^2 \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \frac{(b \cos(c + dx))^{5/2} (2(c + dx) + \sin(2(c + dx)))}{4d \cos^{5/2}(c + dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Sqrt[Cos[c + d*x]],x]

[Out] ((b*Cos[c + d*x])^(5/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 3.20 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{b^2 \sqrt{\cos(dx+c)} b (\cos(dx+c) \sin(dx+c) + dx+c)}{2d \sqrt{\cos(dx+c)}}$	45
risch	$\frac{b^2 x \sqrt{\cos(dx+c)} b}{2 \sqrt{\cos(dx+c)}} + \frac{b^2 \sqrt{\cos(dx+c)} b \sin(2dx+2c)}{4 \sqrt{\cos(dx+c)} d}$	61

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] $1/2/d*b^2*(\cos(d*x+c)*b)^{(1/2)}*(\cos(d*x+c)*\sin(d*x+c)+d*x+c)/\cos(d*x+c)^{(1/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.30

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \left[\frac{2 \sqrt{b \cos(dx + c)} b^2 \sqrt{\cos(dx + c)} \sin(dx + c) + \sqrt{-bb^2} \log(2b \cos(dx + c)^2 - 2}{4d} \right]$$

[In] `integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="fricas")`

[Out] `[1/4*(2*sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(-b)*b^2*log(2*b*cos(d*x + c)^2 - 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/d, 1/2*(sqrt(b*cos(d*x + c))*b^2*sqrt(cos(d*x + c))*sin(d*x + c) + b^(5/2)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/d]`

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(1/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.45 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.46

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \frac{(2(dx + c)b^2 + b^2 \sin(2dx + 2c))\sqrt{b}}{4d}$$

[In] `integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="maxima")`

[Out] `1/4*(2*(d*x + c)*b^2 + b^2*sin(2*d*x + 2*c))*sqrt(b)/d`

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/sqrt(cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.58

$$\int \frac{(b \cos(c + dx))^{5/2}}{\sqrt{\cos(c + dx)}} dx = \frac{b^2 \sqrt{b \cos(c + dx)} (\sin(2c + 2dx) + 2dx)}{4d \sqrt{\cos(c + dx)}}$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(1/2),x)

[Out] (b^2*(b*cos(c + d*x))^(1/2)*(sin(2*c + 2*d*x) + 2*d*x))/(4*d*cos(c + d*x)^(1/2))

$$3.164 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal result	922
Rubi [A] (verified)	922
Mathematica [A] (verified)	923
Maple [A] (verified)	923
Fricas [A] (verification not implemented)	923
Sympy [F(-1)]	924
Maxima [A] (verification not implemented)	924
Giac [F]	924
Mupad [B] (verification not implemented)	924

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}}$$

[Out] $b^2 \sin(dx+c) (b \cos(dx+c))^{1/2} / d \cos(dx+c)^{1/2}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{3}{2}}(c+dx)} dx = \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

[In] `Int[(b*cos[c + d*x])^(5/2)/Cos[c + d*x]^(3/2),x]`

[Out] `(b^2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Cos[c + d*x]])`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m + 1/2) * b^(n - 1/2) * (Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2717

`Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c+dx)}\right) \int \cos(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{3/2}(c+dx)} dx = \frac{(b \cos(c+dx))^{5/2} \sin(c+dx)}{d \cos^{5/2}(c+dx)}$$

[In] Integrate[(b*cos[c + d*x])^(5/2)/Cos[c + d*x]^(3/2),x]

[Out] ((b*cos[c + d*x])^(5/2)*Sin[c + d*x])/(d*cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 2.74 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b^2 \sin(dx+c) \sqrt{\cos(dx+c)b}}{d \sqrt{\cos(dx+c)}}$	32
risch	$\frac{b^2 \sin(dx+c) \sqrt{\cos(dx+c)b}}{d \sqrt{\cos(dx+c)}}$	32

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(3/2),x,method=_RETURNVERBOSE)

[Out] b^2*sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{3/2}(c+dx)} dx = \frac{\sqrt{b \cos(dx+c)} b^2 \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b^2*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{3/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{3/2}(c + dx)} dx = \frac{b^{5/2} \sin(dx + c)}{d}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] b^(5/2)*sin(d*x + c)/d

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{3/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos^{3/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(3/2), x)

Mupad [B] (verification not implemented)

Time = 14.42 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{3/2}(c + dx)} dx = \frac{b^2 \sin(c + dx) \sqrt{b \cos(c + dx)}}{d \sqrt{\cos(c + dx)}}$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(3/2),x)

[Out] (b^2*sin(c + d*x)*(b*cos(c + d*x))^(1/2))/(d*cos(c + d*x)^(1/2))

$$3.165 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal result	925
Rubi [A] (verified)	925
Mathematica [A] (verified)	926
Maple [A] (verified)	926
Fricas [A] (verification not implemented)	926
Sympy [F(-1)]	927
Maxima [A] (verification not implemented)	927
Giac [F]	927
Mupad [B] (verification not implemented)	928

Optimal result

Integrand size = 23, antiderivative size = 27

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{b^2 x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[Out] $b^2 x (b \cos(dx+c))^{1/2} / \cos(dx+c)^{1/2}$

Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{5}{2}}(c+dx)} dx = \frac{b^2 x \sqrt{b \cos(c+dx)}}{\sqrt{\cos(c+dx)}}$$

[In] $\text{Int}[(b \cos[c + d*x])^{5/2} / \cos[c + d*x]^{5/2}, x]$

[Out] $(b^2 x \sqrt{b \cos[c + d*x]}) / \sqrt{\cos[c + d*x]}$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{!IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int 1 dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{b^2 x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \frac{x(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(5/2), x]

[Out] (x*(b*Cos[c + d*x])^(5/2))/Cos[c + d*x]^(5/2)

Maple [A] (verified)

Time = 2.34 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
risch	$\frac{b^2 x \sqrt{\cos(dx+c)} b}{\sqrt{\cos(dx+c)}}$	24
default	$\frac{b^2 \sqrt{\cos(dx+c)} b (dx+c)}{d \sqrt{\cos(dx+c)}}$	31

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(5/2), x, method=_RETURNVERBOSE)

[Out] b^2*x*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.59

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \left[\frac{\sqrt{-bb^2} \log \left(2 b \cos(dx + c)^2 - 2 \sqrt{b \cos(dx + c)} \sqrt{-b} \sqrt{\cos(dx + c)} \sin(dx + c) \right)}{2 d} \right]$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2), x, algorithm="fricas")

[Out] $[1/2\sqrt{-b}b^2\log(2b\cos(dx+c)^2 - 2\sqrt{b\cos(dx+c)})\sqrt{-b}\sqrt{\cos(dx+c)}\sin(dx+c) - b)/d, b^{5/2}\arctan(\sqrt{b\cos(dx+c)}\sin(dx+c)/(\sqrt{b}\cos(dx+c)^{3/2}))/d]$

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \text{Timed out}$$

[In] `integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \frac{2b^{5/2} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

[In] `integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x, algorithm="maxima")`

[Out] $2*b^{5/2}*\arctan(\sin(d*x + c)/(\cos(d*x + c) + 1))/d$

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos(dx + c)^{5/2}} dx$$

[In] `integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(5/2),x, algorithm="giac")`

[Out] `integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(5/2), x)`

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.85

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{5/2}(c + dx)} dx = \frac{b^2 x \sqrt{b \cos(c + dx)}}{\sqrt{\cos(c + dx)}}$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(5/2),x)

[Out] (b^2*x*(b*cos(c + d*x))^(1/2))/cos(c + d*x)^(1/2)

$$3.166 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx$$

Optimal result	929
Rubi [A] (verified)	929
Mathematica [A] (verified)	930
Maple [A] (verified)	930
Fricas [A] (verification not implemented)	931
Sympy [F(-1)]	931
Maxima [B] (verification not implemented)	931
Giac [F]	932
Mupad [F(-1)]	932

Optimal result

Integrand size = 23, antiderivative size = 36

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx = \frac{b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

[Out] b^2*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3855}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^2(c+dx)} dx = \frac{b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}}$$

[In] Int[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(7/2), x]

[Out] (b^2*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(d*Sqrt[Cos[c + d*x]])

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3855

Int[csc[(c_)+(d_)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c+dx)}\right) \int \sec(c+dx) dx}{\sqrt{\cos(c+dx)}} \\ &= \frac{b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{d \sqrt{\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{7/2}(c+dx)} dx = \frac{\operatorname{arctanh}(\sin(c+dx))(b \cos(c+dx))^{5/2}}{d \cos^{5/2}(c+dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(7/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*(b*Cos[c + d*x])^(5/2))/(d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 2.65 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.19

method	result	size
default	$-\frac{2\sqrt{\cos(dx+c)}b^2 \operatorname{arctanh}(\cot(dx+c)-\csc(dx+c))}{d\sqrt{\cos(dx+c)}}$	43
risch	$\frac{b^2\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}+i)}{\sqrt{\cos(dx+c)}d} - \frac{b^2\sqrt{\cos(dx+c)}b \ln(e^{i(dx+c)}-i)}{\sqrt{\cos(dx+c)}d}$	79

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(7/2), x, method=_RETURNVERBOSE)

[Out] -2/d*(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(1/2)*b^2*arctanh(cot(d*x+c)-csc(d*x+c))

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.22

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{7/2}(c + dx)} dx = \left[\frac{b^{5/2} \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right)}{2d}, \right. \\ \left. -\frac{\sqrt{-bb^2} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b \sin(dx+c)}}{b\sqrt{\cos(dx+c)}}\right)}{d} \right]$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="fricas")

```
[Out] [1/2*b^(5/2)*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(c
os(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3)/d, -sqrt(-b)*
b^2*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c)
)))/d]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{7/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(7/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(32) = 64.

Time = 0.41 (sec) , antiderivative size = 72, normalized size of antiderivative = 2.00

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{7/2}(c + dx)} dx = \frac{(b^2 \log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - b^2 \log(\cos(dx + c)^2)}{2d}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="maxima")

```
[Out] 1/2*(b^2*log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - b^2*lo
g(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))*sqrt(b)/d
```

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{7/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos(dx + c)^{7/2}} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(7/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{7/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^{7/2}} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(7/2),x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(7/2), x)

$$3.167 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{9}{2}}(c+dx)} dx$$

Optimal result	933
Rubi [A] (verified)	933
Mathematica [A] (verified)	934
Maple [A] (verified)	934
Fricas [A] (verification not implemented)	935
Sympy [F(-1)]	935
Maxima [A] (verification not implemented)	935
Giac [F]	936
Mupad [B] (verification not implemented)	936

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{\frac{3}{2}}(c+dx)}$$

[Out] $b^2 \sin(dx+c) (b \cos(dx+c))^{1/2} / d / \cos(dx+c)^{3/2}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3852, 8}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{\frac{9}{2}}(c+dx)} dx = \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{\frac{3}{2}}(c+dx)}$$

[In] $\text{Int}[(b \cos[c + dx])^{5/2} / \cos[c + dx]^{9/2}, x]$

[Out] $(b^2 \sqrt{b \cos[c + dx]} \sin[c + dx]) / (d \cos[c + dx]^{3/2})$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_))^{(m_)} * ((b_.) * (v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)} * b^{(n - 1/2)} * (\text{Sqrt}[b*v] / \text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sec^2(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \text{Subst}\left(\int 1 dx, x, -\tan(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \frac{(b \cos(c + dx))^{5/2} \sin(c + dx)}{d \cos^{7/2}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(9/2), x]
```

```
[Out] ((b*Cos[c + d*x])^(5/2)*Sin[c + d*x])/(d*Cos[c + d*x]^(7/2))
```

Maple [A] (verified)

Time = 2.75 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{b^2 \sin(dx+c) \sqrt{\cos(dx+c)b}}{d \cos(dx+c)^{\frac{3}{2}}}$	32
risch	$\frac{2ib^2 \sqrt{\cos(dx+c)b}}{\sqrt{\cos(dx+c)} d(e^{2i(dx+c)}+1)}$	41

```
[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(9/2), x, method=_RETURNVERBOSE)
```

```
[Out] b^2*sin(d*x+c)*(cos(d*x+c)*b)^(1/2)/d/cos(d*x+c)^(3/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \frac{\sqrt{b \cos(dx + c)} b^2 \sin(dx + c)}{d \cos(dx + c)^{3/2}}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*b^2*sin(d*x + c)/(d*cos(d*x + c)^(3/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(9/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.42 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.54

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \frac{2 b^{5/2} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c))^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1} d$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="maxima")

[Out] 2*b^(5/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c))^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos^{9/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(9/2), x)

Mupad [B] (verification not implemented)

Time = 14.49 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.77

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{9/2}(c + dx)} dx = \frac{b^2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + 1)}{d \sqrt{\cos(c + dx)} (\cos(2c + 2dx) + 1)}$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(9/2),x)

[Out] (b^2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/
(d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

$$3.168 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{11/2}(c+dx)} dx$$

Optimal result	937
Rubi [A] (verified)	937
Mathematica [A] (verified)	938
Maple [A] (verified)	939
Fricas [A] (verification not implemented)	939
Sympy [F(-1)]	940
Maxima [B] (verification not implemented)	940
Giac [F]	941
Mupad [F(-1)]	941

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{11/2}(c+dx)} dx = \frac{b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{2d \sqrt{\cos(c+dx)}} + \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{2d \cos^{5/2}(c+dx)}$$

[Out] 1/2*b^2*sin(d*x+c)*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(5/2)+1/2*b^2*arctanh(sin(d*x+c))*(b*cos(d*x+c))^(1/2)/d/cos(d*x+c)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{11/2}(c+dx)} dx = \frac{b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{2d \sqrt{\cos(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{2d \cos^{5/2}(c+dx)}$$

[In] Int[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(11/2), x]

[Out] (b^2*ArcTanh[Sin[c + d*x]]*Sqrt[b*Cos[c + d*x]])/(2*d*Sqrt[Cos[c + d*x]]) + (b^2*Sqrt[b*Cos[c + d*x]]*Sin[c + d*x])/(2*d*Cos[c + d*x]^(5/2))

Rule 17

```
Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[a^(m + 1/2)
)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sec^3(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d \cos^{\frac{5}{2}}(c + dx)} + \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sec(c + dx) dx}{2\sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{2d \sqrt{\cos(c + dx)}} + \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{2d \cos^{\frac{5}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.67

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{\frac{11}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^{5/2} (\operatorname{arctanh}(\sin(c + dx)) \cos^2(c + dx) + \sin(c + dx))}{2d \cos^{\frac{9}{2}}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(11/2), x]
```

```
[Out] ((b*Cos[c + d*x])^(5/2)*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x
]))/(2*d*Cos[c + d*x]^(9/2))
```

Maple [A] (verified)

Time = 2.81 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{b^2(-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c)) \sqrt{\cos(dx+c)b}}{2d \cos(dx+c)^{\frac{5}{2}}}$	87
risch	$-\frac{ib^2 \sqrt{\cos(dx+c)b} (e^{3i(dx+c)} - e^{i(dx+c)})}{\sqrt{\cos(dx+c)} d (e^{2i(dx+c)} + 1)^2} - \frac{b^2 \sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)} - i)}{2\sqrt{\cos(dx+c)} d} + \frac{b^2 \sqrt{\cos(dx+c)b} \ln(e^{i(dx+c)} + i)}{2\sqrt{\cos(dx+c)} d}$	14

[In] `int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(11/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}d^2 b^2 (-\cos(dx+c)^2 \ln(-\cot(dx+c)+\csc(dx+c)-1) + \cos(dx+c)^2 \ln(-\cot(dx+c)+\csc(dx+c)+1) + \sin(dx+c)) (\cos(dx+c)b)^{1/2} / \cos(dx+c)^{5/2}$

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 210, normalized size of antiderivative = 2.69

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx = \left[\frac{b^{5/2} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + \sqrt{-bb^2} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b \sin(dx+c)}}{b \sqrt{\cos(dx+c)}}\right) \cos(dx+c)^3 - \sqrt{b \cos(dx+c)} b^2 \sqrt{\cos(dx+c)} \sin(dx+c)}{4d \cos(dx+c)^3} \right]$$

[In] `integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2),x, algorithm="fricas")`

[Out] $\left[\frac{1}{4} b^{5/2} \cos(dx+c)^3 \log(-b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)) / \cos(dx+c)^3 + 2 \sqrt{b \cos(dx+c)} b^2 \sqrt{\cos(dx+c)} \sin(dx+c) / (d \cos(dx+c)^3), -1/2 (\sqrt{-b} b^2 \arctan(\sqrt{b \cos(dx+c)} \sqrt{-b \sin(dx+c)} / (b \sqrt{\cos(dx+c)}))) \cos(dx+c)^3 - \sqrt{b \cos(dx+c)} b^2 \sqrt{\cos(dx+c)} \sin(dx+c) / (d \cos(dx+c)^3) \right]$

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(11/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 747 vs. 2(66) = 132.

Time = 0.40 (sec) , antiderivative size = 747, normalized size of antiderivative = 9.58

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx = \text{Too large to display}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2),x, algorithm="maxima")

```
[Out] -1/4*(4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos^{11/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(11/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(11/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos^{11/2}(c + dx)} dx$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(11/2),x)

[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(11/2), x)

$$3.169 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx$$

Optimal result	942
Rubi [A] (verified)	942
Mathematica [A] (verified)	943
Maple [A] (verified)	943
Fricas [A] (verification not implemented)	944
Sympy [F(-1)]	944
Maxima [B] (verification not implemented)	944
Giac [F]	945
Mupad [B] (verification not implemented)	945

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx = \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{d \cos^{3/2}(c+dx)} + \frac{b^2 \sqrt{b \cos(c+dx)} \sin^3(c+dx)}{3d \cos^{7/2}(c+dx)}$$

[Out] $b^2 \sin(d*x+c) * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(3/2)} + 1/3 * b^2 \sin(d*x+c)^3 * (b \cos(d*x+c))^{(1/2)} / d / \cos(d*x+c)^{(7/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3852}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{13/2}(c+dx)} dx = \frac{b^2 \sin^3(c+dx) \sqrt{b \cos(c+dx)}}{3d \cos^{7/2}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{d \cos^{3/2}(c+dx)}$$

[In] Int[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(13/2),x]

[Out] $(b^2 \sqrt{b \cos[c + d*x]} \sin[c + d*x]) / (d \cos[c + d*x]^{(3/2)}) + (b^2 \sqrt{b \cos[c + d*x]} \sin^3[c + d*x]) / (3 * d \cos[c + d*x]^{(7/2)})$

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2) * b^(n - 1/2) * (Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sec^4(c + dx) dx}{\sqrt{\cos(c + dx)}} \\ &= -\frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \text{Subst}\left(\int (1 + x^2) dx, x, -\tan(c + dx)\right)}{d \sqrt{\cos(c + dx)}} \\ &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{d \cos^{\frac{3}{2}}(c + dx)} + \frac{b^2 \sqrt{b \cos(c + dx)} \sin^3(c + dx)}{3d \cos^{\frac{7}{2}}(c + dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{\frac{13}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^{5/2} (\tan(c + dx) + \frac{1}{3} \tan^3(c + dx))}{d \cos^{\frac{5}{2}}(c + dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(13/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 2.67 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

method	result	size
default	$\frac{b^2 (2(\cos^2(dx+c))+1) \sqrt{\cos(dx+c)} b \sin(dx+c)}{3d \cos(dx+c)^{\frac{7}{2}}}$	45
risch	$\frac{4ib^2 \sqrt{\cos(dx+c)} b (3e^{2i(dx+c)}+1)}{3\sqrt{\cos(dx+c)} d(e^{2i(dx+c)}+1)^3}$	54

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(13/2), x, method=_RETURNVERBOSE)

[Out] 1/3/d*b^2*(2*cos(d*x+c)^2+1)*(cos(d*x+c)*b)^(1/2)*sin(d*x+c)/cos(d*x+c)^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.61

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{13/2}(c + dx)} dx = \frac{(2b^2 \cos(dx + c)^2 + b^2) \sqrt{b \cos(dx + c)} \sin(dx + c)}{3d \cos(dx + c)^{7/2}}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2),x, algorithm="fricas")

[Out] 1/3*(2*b^2*cos(d*x + c)^2 + b^2)*sqrt(b*cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(7/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{13/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(5/2)/cos(d*x+c)**(13/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(66) = 132.

Time = 0.38 (sec) , antiderivative size = 311, normalized size of antiderivative = 4.09

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{13/2}(c + dx)} dx =$$

$$\frac{4(3b^2 \cos(6dx + 6c) + 3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1)d}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1)d}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2),x, algorithm="maxima")

[Out] -4/3*(3*b^2*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b^2*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b^2*cos(2*d*x + 2*c) + b^2)*sin(6*d*x + 6*c) - 3*(3*b^2*cos(2*d*x + 2*c) + b^2)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{13/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos^{13/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(13/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(13/2), x)

Mupad [B] (verification not implemented)

Time = 14.51 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.72

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{13/2}(c + dx)} dx = \frac{2 b^2 \sqrt{b \cos(c + dx)} (\cos(2c + 2dx) 15i + \cos(4c + 4dx) 6i + \cos(6c + 6dx))}{3 d \sqrt{\cos(c + dx)} (15 \cos(2c + 2dx) + 6 \cos(4c + 4dx) + \cos(6c + 6dx) + 10)}$$

[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(13/2),x)

[Out] (2*b^2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.170 \quad \int \frac{(b \cos(c+dx))^{5/2}}{\cos^{15/2}(c+dx)} dx$$

Optimal result	946
Rubi [A] (verified)	946
Mathematica [A] (verified)	948
Maple [A] (verified)	948
Fricas [A] (verification not implemented)	948
Sympy [F(-1)]	949
Maxima [B] (verification not implemented)	949
Giac [F]	950
Mupad [F(-1)]	951

Optimal result

Integrand size = 23, antiderivative size = 116

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{15/2}(c+dx)} dx = \frac{3b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{4d \cos^{9/2}(c+dx)} + \frac{3b^2 \sqrt{b \cos(c+dx)} \sin(c+dx)}{8d \cos^{5/2}(c+dx)}$$

[Out] $1/4*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(9/2)}+3/8*b^2*\sin(d*x+c)*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(5/2)}+3/8*b^2*\operatorname{arctanh}(\sin(d*x+c))*(b*\cos(d*x+c))^{(1/2)}/d/\cos(d*x+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3853, 3855}

$$\int \frac{(b \cos(c+dx))^{5/2}}{\cos^{15/2}(c+dx)} dx = \frac{3b^2 \operatorname{arctanh}(\sin(c+dx)) \sqrt{b \cos(c+dx)}}{8d \sqrt{\cos(c+dx)}} + \frac{3b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{8d \cos^{5/2}(c+dx)} + \frac{b^2 \sin(c+dx) \sqrt{b \cos(c+dx)}}{4d \cos^{9/2}(c+dx)}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c+d*x])^{(5/2)}/\operatorname{Cos}[c+d*x]^{(15/2)},x]$

[Out] $(3*b^2*\operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]]*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]])/(8*d*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]) + (b^2*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(4*d*\operatorname{Cos}[c+d*x]^{(9/2)}) + (3*b^2*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]*\operatorname{Sin}[c+d*x])/(8*d*\operatorname{Cos}[c+d*x]^{(5/2)})$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\left(b^2 \sqrt{b \cos(c + dx)}\right) \int \sec^5(c + dx) dx}{\sqrt{\cos(c + dx)}} \\
 &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{\left(3b^2 \sqrt{b \cos(c + dx)}\right) \int \sec^3(c + dx) dx}{4\sqrt{\cos(c + dx)}} \\
 &= \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{3b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{\frac{5}{2}}(c + dx)} \\
 &\quad + \frac{\left(3b^2 \sqrt{b \cos(c + dx)}\right) \int \sec(c + dx) dx}{8\sqrt{\cos(c + dx)}} \\
 &= \frac{3b^2 \operatorname{arctanh}(\sin(c + dx)) \sqrt{b \cos(c + dx)}}{8d \sqrt{\cos(c + dx)}} \\
 &\quad + \frac{b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{4d \cos^{\frac{9}{2}}(c + dx)} + \frac{3b^2 \sqrt{b \cos(c + dx)} \sin(c + dx)}{8d \cos^{\frac{5}{2}}(c + dx)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.57

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \frac{(b \cos(c + dx))^{5/2} (3 \operatorname{arctanh}(\sin(c + dx)) \cos^4(c + dx) + (2 + 3 \cos^2(c + dx)) \sin(c + dx))}{8d \cos^{13/2}(c + dx)}$$

[In] Integrate[(b*Cos[c + d*x])^(5/2)/Cos[c + d*x]^(15/2), x]

[Out] ((b*Cos[c + d*x])^(5/2)*(3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x]))/(8*d*Cos[c + d*x]^(13/2))

Maple [A] (verified)

Time = 3.24 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.91

method	result
default	$-\frac{(3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)-3(\cos^4(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)-3(\cos^2(dx+c)) \sin(dx+c)-2 \sin(dx+c))}{8d \cos(dx+c)^{9/2}}$
risch	$-\frac{ib^2 \sqrt{\cos(dx+c)} b (3e^{7i(dx+c)}+11e^{5i(dx+c)}-11e^{3i(dx+c)}-3e^{i(dx+c)})}{4\sqrt{\cos(dx+c)} d(e^{2i(dx+c)}+1)^4} + \frac{3b^2 \sqrt{\cos(dx+c)} b \ln(e^{i(dx+c)}+i)}{8\sqrt{\cos(dx+c)} d} - \frac{3b^2 \sqrt{\cos(dx+c)} b \ln(e^{i(dx+c)}-i)}{8\sqrt{\cos(dx+c)} d}$

[In] int((cos(d*x+c)*b)^(5/2)/cos(d*x+c)^(15/2), x, method=_RETURNVERBOSE)

[Out] -1/8/d*(3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)-3*cos(d*x+c)^2*sin(d*x+c)-2*sin(d*x+c))*(cos(d*x+c)*b)^(1/2)*b^2/cos(d*x+c)^(9/2)

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 244, normalized size of antiderivative = 2.10

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \frac{3b^{5/2} \cos(dx+c)^5 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)} \sqrt{b \cos(dx+c)} \sin(dx+c) - 2b \cos(dx+c)}{\cos(dx+c)^3}\right) + 3\sqrt{-bb^2} \arctan\left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b \sin(dx+c)}}{b \sqrt{\cos(dx+c)}}\right) \cos(dx+c)^5 - (3b^2 \cos(dx+c)^2 + 2b^2) \sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)}}{8d \cos(dx+c)^5}$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2), x, algorithm="fricas")

[Out] [1/16*(3*b^(5/2)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x +

$c)^3 + 2*(3*b^2*\cos(dx + c)^2 + 2*b^2)*\sqrt{b*\cos(dx + c)}*\sqrt{\cos(dx + c)}*\sin(dx + c))/(d*\cos(dx + c)^5)$, $-1/8*(3*\sqrt{-b}*b^2*\arctan(\sqrt{b*\cos(dx + c)}*\sqrt{-b}*\sin(dx + c)/(b*\sqrt{\cos(dx + c)})))*\cos(dx + c)^5 - (3*b^2*\cos(dx + c)^2 + 2*b^2)*\sqrt{b*\cos(dx + c)}*\sqrt{\cos(dx + c)}*\sin(dx + c))/(d*\cos(dx + c)^5]$

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(dx+c))**(5/2)/cos(dx+c)**(15/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1914 vs. 2(98) = 196.

Time = 0.51 (sec) , antiderivative size = 1914, normalized size of antiderivative = 16.50

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \text{Too large to display}$$

[In] integrate((b*cos(dx+c))^(5/2)/cos(dx+c)^(15/2),x, algorithm="maxima")

[Out] $-1/16*(12*(b^2*\sin(8*dx + 8*c) + 4*b^2*\sin(6*dx + 6*c) + 6*b^2*\sin(4*dx + 4*c) + 4*b^2*\sin(2*dx + 2*c))*\cos(7/2*\arctan2(\sin(2*dx + 2*c), \cos(2*dx + 2*c))) + 44*(b^2*\sin(8*dx + 8*c) + 4*b^2*\sin(6*dx + 6*c) + 6*b^2*\sin(4*dx + 4*c) + 4*b^2*\sin(2*dx + 2*c))*\cos(5/2*\arctan2(\sin(2*dx + 2*c), \cos(2*dx + 2*c))) - 44*(b^2*\sin(8*dx + 8*c) + 4*b^2*\sin(6*dx + 6*c) + 6*b^2*\sin(4*dx + 4*c) + 4*b^2*\sin(2*dx + 2*c))*\cos(3/2*\arctan2(\sin(2*dx + 2*c), \cos(2*dx + 2*c))) - 12*(b^2*\sin(8*dx + 8*c) + 4*b^2*\sin(6*dx + 6*c) + 6*b^2*\sin(4*dx + 4*c) + 4*b^2*\sin(2*dx + 2*c))*\cos(1/2*\arctan2(\sin(2*dx + 2*c), \cos(2*dx + 2*c))) - 3*(b^2*\cos(8*dx + 8*c)^2 + 16*b^2*\cos(6*dx + 6*c)^2 + 36*b^2*\cos(4*dx + 4*c)^2 + 16*b^2*\cos(2*dx + 2*c)^2 + b^2*\sin(8*dx + 8*c)^2 + 16*b^2*\sin(6*dx + 6*c)^2 + 36*b^2*\sin(4*dx + 4*c)^2 + 4*8*b^2*\sin(4*dx + 4*c)*\sin(2*dx + 2*c) + 16*b^2*\sin(2*dx + 2*c)^2 + 8*b^2*\cos(2*dx + 2*c) + b^2 + 2*(4*b^2*\cos(6*dx + 6*c) + 6*b^2*\cos(4*dx + 4*c) + 4*b^2*\cos(2*dx + 2*c) + b^2)*\cos(8*dx + 8*c) + 8*(6*b^2*\cos(4*dx + 4*c) + 4*b^2*\cos(2*dx + 2*c) + b^2)*\cos(6*dx + 6*c) + 12*(4*b^2*\cos(2*dx + 2*c) + b^2)*\cos(4*dx + 4*c) + 4*(2*b^2*\sin(6*dx + 6*c) + 3*b^2*\sin(4*dx + 4*c) + 2*b^2*\sin(2*dx + 2*c))*\sin(8*dx + 8*c) + 16*(3*b^2*\sin(4*dx + 4*c) + 2*b^2*\sin(2*dx + 2*c))*\sin(6*dx + 6*c))*\log(\cos(1/2*\arctan2(\sin(2$

```

*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2
*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) +
1) + 3*(b^2*cos(8*d*x + 8*c)^2 + 16*b^2*cos(6*d*x + 6*c)^2 + 36*b^2*cos(4*d
*x + 4*c)^2 + 16*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(8*d*x + 8*c)^2 + 16*b^2*s
in(6*d*x + 6*c)^2 + 36*b^2*sin(4*d*x + 4*c)^2 + 48*b^2*sin(4*d*x + 4*c)*sin
(2*d*x + 2*c) + 16*b^2*sin(2*d*x + 2*c)^2 + 8*b^2*cos(2*d*x + 2*c) + b^2 +
2*(4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c)
+ b^2)*cos(8*d*x + 8*c) + 8*(6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*
c) + b^2)*cos(6*d*x + 6*c) + 12*(4*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x +
4*c) + 4*(2*b^2*sin(6*d*x + 6*c) + 3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x
+ 2*c))*sin(8*d*x + 8*c) + 16*(3*b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x +
2*c))*sin(6*d*x + 6*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 12*(b^2*cos(8*d*x + 8
*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2
*c) + b^2)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b^2*c
os(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*c) + 4*b^2*c
os(2*d*x + 2*c) + b^2)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))
+ 44*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*cos(4*d*x + 4*
c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*
d*x + 2*c))) + 12*(b^2*cos(8*d*x + 8*c) + 4*b^2*cos(6*d*x + 6*c) + 6*b^2*co
s(4*d*x + 4*c) + 4*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4
*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*
cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x
+ 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*
c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) +
2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*
x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 3
6*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x
+ 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*d)

```

Giac [F]

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \int \frac{(b \cos(dx + c))^{5/2}}{\cos^{15/2}(dx + c)} dx$$

[In] integrate((b*cos(d*x+c))^(5/2)/cos(d*x+c)^(15/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(5/2)/cos(d*x + c)^(15/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^{5/2}}{\cos^{15/2}(c + dx)} dx = \int \frac{(b \cos(c + dx))^{5/2}}{\cos(c + dx)^{15/2}} dx$$

```
[In] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(15/2), x)
```

```
[Out] int((b*cos(c + d*x))^(5/2)/cos(c + d*x)^(15/2), x)
```

$$3.171 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	952
Rubi [A] (verified)	952
Mathematica [A] (verified)	953
Maple [A] (verified)	953
Fricas [A] (verification not implemented)	954
Sympy [F(-1)]	954
Maxima [A] (verification not implemented)	954
Giac [F]	955
Mupad [B] (verification not implemented)	955

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b \cos(c+dx)}} - \frac{2\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \sin^5(c+dx)}{5d\sqrt{b \cos(c+dx)}}$$

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)-2/3*sin(d*x+c)^3*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)+1/5*sin(d*x+c)^5*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sin^5(c+dx)\sqrt{\cos(c+dx)}}{5d\sqrt{b \cos(c+dx)}} - \frac{2\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^(11/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]]) - (2*Sqrt[Cos[c + d*x]]*Sin[c + d*x]^3)/(3*d*Sqrt[b*Cos[c + d*x]]) + (Sqrt[Cos[c + d*x]]*Sin[c + d*x]^5)/(5*d*Sqrt[b*Cos[c + d*x]])

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x] \&\& !\text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2713

$\text{Int}[\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1-x^2)^{((n-1)/2)}, x], x], x, \text{Cos}[c+d*x]], x] /; \text{FreeQ}\{c, d\}, x] \&\& \text{IGtQ}[(n-1)/2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos^5(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}(\int (1-2x^2+x^4) dx, x, -\sin(c+dx))}{d\sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b \cos(c+dx)}} - \frac{2\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \sin^5(c+dx)}{5d\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.53

$$\int \frac{\cos^{11/2}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx) (15 - 10 \sin^2(c+dx) + 3 \sin^4(c+dx))}{15d\sqrt{b \cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(11/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x]*(15 - 10*Sin[c + d*x]^2 + 3*Sin[c + d*x]^4))/(15*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.91 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.49

method	result	size
default	$\frac{(3(\cos^4(dx+c))+4(\cos^2(dx+c))+8) \sin(dx+c)(\sqrt{\cos(dx+c)})}{15d\sqrt{\cos(dx+c)b}}$	52
risch	$\frac{5 \sin(dx+c)(\sqrt{\cos(dx+c)})}{8d\sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(5dx+5c)}{80\sqrt{\cos(dx+c)b}d} + \frac{5(\sqrt{\cos(dx+c)}) \sin(3dx+3c)}{48\sqrt{\cos(dx+c)b}d}$	95

[In] `int(cos(d*x+c)^(11/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{15} \frac{1}{d} \frac{(3 \cos(dx+c)^4 + 4 \cos(dx+c)^2 + 8) \sin(dx+c) \cos(dx+c)^{1/2}}{(\cos(dx+c)*b)^{1/2}}$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.50

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{(3 \cos(dx+c)^4 + 4 \cos(dx+c)^2 + 8) \sqrt{b \cos(dx+c)} \sin(dx+c)}{15 b d \sqrt{\cos(dx+c)}}$$

[In] `integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{15} \frac{(3 \cos(dx+c)^4 + 4 \cos(dx+c)^2 + 8) \sqrt{b \cos(dx+c)} \sin(dx+c)}{(b*d*\sqrt{\cos(dx+c)})}$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.64

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{3 \sin(5 dx + 5 c) + 25 \sin\left(\frac{3}{5} \arctan\left(\frac{\sin(5 dx + 5 c)}{\cos(5 dx + 5 c)}\right)\right) + 150 \sin\left(\frac{1}{5} \arctan\left(\frac{\sin(5 dx + 5 c)}{\cos(5 dx + 5 c)}\right)\right)}{240 \sqrt{bd}}$$

[In] `integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] $\frac{1}{240} \frac{(3 \sin(5 dx + 5 c) + 25 \sin(\frac{3}{5} \arctan2(\sin(5 dx + 5 c), \cos(5 dx + 5 c))) + 150 \sin(\frac{1}{5} \arctan2(\sin(5 dx + 5 c), \cos(5 dx + 5 c))))}{(\sqrt{b*d})}$

Giac [F]

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^{\frac{11}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(11/2)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 14.70 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.68

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx$$

$$= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (175 \sin(2c + 2dx) + 28 \sin(4c + 4dx) + 3 \sin(6c + 6dx))}{240bd (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(175*sin(2*c + 2*d*x) + 28*sin(4*c + 4*d*x) + 3*sin(6*c + 6*d*x)))/(240*b*d*(cos(2*c + 2*d*x) + 1))

3.172 $\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$

Optimal result	956
Rubi [A] (verified)	956
Mathematica [A] (verified)	957
Maple [A] (verified)	958
Fricas [A] (verification not implemented)	958
Sympy [F(-1)]	958
Maxima [A] (verification not implemented)	959
Giac [F]	959
Mupad [B] (verification not implemented)	959

Optimal result

Integrand size = 23, antiderivative size = 98

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b \cos(c+dx)}} + \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8d\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b \cos(c+dx)}}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4d\sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{8d\sqrt{b \cos(c+dx)}}$$

[In] `Int[Cos[c + d*x]^(9/2)/Sqrt[b*Cos[c + d*x]],x]`

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 17

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)
)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos^4(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\
 &= \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b \cos(c+dx)}} + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int \cos^2(c+dx) dx}{4\sqrt{b \cos(c+dx)}} \\
 &= \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8d\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b \cos(c+dx)}} + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int 1 dx}{8\sqrt{b \cos(c+dx)}} \\
 &= \frac{3x\sqrt{\cos(c+dx)}}{8\sqrt{b \cos(c+dx)}} + \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8d\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4d\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.56

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)}(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32d\sqrt{b \cos(c+dx)}}$$

```
[In] Integrate[Cos[c + d*x]^(9/2)/Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (Sqrt[Cos[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))
/(32*d*Sqrt[b*Cos[c + d*x]])
```

Maple [A] (verified)

Time = 3.17 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.63

method	result	size
default	$\frac{(2 \sin(dx+c)(\cos^3(dx+c))+3 \cos(dx+c) \sin(dx+c)+3dx+3c) (\sqrt{\cos(dx+c)})}{8d\sqrt{\cos(dx+c)b}}$	62
risch	$\frac{3x(\sqrt{\cos(dx+c)})}{8\sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(4dx+4c)}{32\sqrt{\cos(dx+c)b}d} + \frac{(\sqrt{\cos(dx+c)}) \sin(2dx+2c)}{4\sqrt{\cos(dx+c)b}d}$	87

[In] `int(cos(d*x+c)^(9/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{8}d*(2*\sin(d*x+c)*\cos(d*x+c)^3+3*\cos(d*x+c)*\sin(d*x+c)+3*d*x+3*c)*\cos(d*x+c)^(1/2)/(\cos(d*x+c)*b)^(1/2)$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.86

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b} \log(2b \cos(dx+c)^2 + 2 \sqrt{b \cos(dx+c)})}{16bd}$$

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")`

[Out] $[1/16*(2*\sqrt{b*\cos(d*x+c)}*(2*\cos(d*x+c)^2+3)*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-3*\sqrt{-b}*\log(2*b*\cos(d*x+c)^2+2*\sqrt{b*\cos(d*x+c)})*\sqrt{-b}*\sqrt{\cos(d*x+c)}*\sin(d*x+c)-b)]/(b*d), 1/8*(\sqrt{b*\cos(d*x+c)}*(2*\cos(d*x+c)^2+3)*\sqrt{\cos(d*x+c)}*\sin(d*x+c)+3*\sqrt{b}*\arctan(\sqrt{b*\cos(d*x+c)}*\sin(d*x+c)/(\sqrt{b}*\cos(d*x+c)^(3/2))))/(b*d)]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(1/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.50

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \frac{12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c))\right)}{32 \sqrt{bd}}$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(sqrt(b)*d)

Giac [F]

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \int \frac{\cos(dx+c)^{\frac{9}{2}}}{\sqrt{b \cos(dx+c)}} dx$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(9/2)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 14.39 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.80

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \frac{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (8 \sin(c+dx) + 9 \sin(3c+3dx) + \sin(5c+5dx) + 24 dx \cos(c+dx))}{32 b d (\cos(2c+2dx) + 1)}$$

[In] int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*b*d*(cos(2*c + 2*d*x) + 1))

$$3.173 \quad \int \frac{\cos^{\frac{7}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	960
Rubi [A] (verified)	960
Mathematica [A] (verified)	961
Maple [A] (verified)	961
Fricas [A] (verification not implemented)	962
Sympy [F(-1)]	962
Maxima [A] (verification not implemented)	962
Giac [F]	963
Mupad [B] (verification not implemented)	963

Optimal result

Integrand size = 23, antiderivative size = 70

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d\sqrt{b \cos(c+dx)}} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3d\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3d\sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}/\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) - (\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 2713


```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^3(c + dx) dx}{\sqrt{b \cos(c + dx)}} \\ &= -\frac{\sqrt{\cos(c + dx)} \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{b \cos(c + dx)}} \\ &= \frac{\sqrt{\cos(c + dx)} \sin(c + dx)}{d \sqrt{b \cos(c + dx)}} - \frac{\sqrt{\cos(c + dx)} \sin^3(c + dx)}{3d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sqrt{\cos(c + dx)}(5 + \cos(2(c + dx))) \sin(c + dx)}{6d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^(7/2)/Sqrt[b*Cos[c + d*x]],x]
```

```
[Out] (Sqrt[Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*Sqrt[b*Cos[c + d*x]])
```

Maple [A] (verified)

Time = 2.98 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.57

method	result	size
default	$\frac{(2 + \cos^2(dx+c)) \sin(dx+c) (\sqrt{\cos(dx+c)})}{3d \sqrt{\cos(dx+c)b}}$	40
risch	$\frac{3 \sin(dx+c) (\sqrt{\cos(dx+c)})}{4d \sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(3dx+3c)}{12 \sqrt{\cos(dx+c)b} d}$	63

```
[In] int(cos(d*x+c)^(7/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3bd\sqrt{\cos(dx + c)}}$$

```
[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan\left(\frac{\sin(3dx + 3c)}{\cos(3dx + 3c)}\right)\right)}{12\sqrt{bd}}$$

```
[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(sqrt(b)*d)
```

Giac [F]

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^{\frac{7}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.86

$$\begin{aligned} & \int \frac{\cos^{\frac{7}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx \\ &= \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12bd(\cos(2c + 2dx) + 1)} \end{aligned}$$

[In] int(cos(c + d*x)^(7/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*b*d*(cos(2*c + 2*d*x) + 1))

$$3.174 \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	964
Rubi [A] (verified)	964
Mathematica [A] (verified)	965
Maple [A] (verified)	965
Fricas [A] (verification not implemented)	966
Sympy [F(-1)]	966
Maxima [A] (verification not implemented)	966
Giac [F]	967
Mupad [B] (verification not implemented)	967

Optimal result

Integrand size = 23, antiderivative size = 63

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{x \sqrt{\cos(c+dx)}}{2\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{x \sqrt{\cos(c+dx)}}{2\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2d\sqrt{b \cos(c+dx)}}$$

[In] `Int[Cos[c + d*x]^(5/2)/Sqrt[b*Cos[c + d*x]],x]`

[Out] $(x*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b`

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^2(c + dx) dx}{\sqrt{b \cos(c + dx)}} \\ &= \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2d\sqrt{b \cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \int 1 dx}{2\sqrt{b \cos(c + dx)}} \\ &= \frac{x\sqrt{\cos(c + dx)}}{2\sqrt{b \cos(c + dx)}} + \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2d\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.71

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sqrt{\cos(c + dx)}(2(c + dx) + \sin(2(c + dx)))}{4d\sqrt{b \cos(c + dx)}}$$

[In] Integrate[Cos[c + d*x]^(5/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.10 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.67

method	result	size
default	$\frac{(\cos(dx+c) \sin(dx+c)+dx+c)(\sqrt{\cos(dx+c)})}{2d\sqrt{\cos(dx+c)b}}$	42
risch	$\frac{x(\sqrt{\cos(dx+c)})}{2\sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(2dx+2c)}{4\sqrt{\cos(dx+c)b}d}$	55

[In] int(cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.49

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

$$= \left[\frac{2 \sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)} \sin(dx+c) - \sqrt{-b} \log \left(2 b \cos(dx+c)^2 + 2 \sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \right)}{4 b d} \right]$$

```
[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
```

```
[Out] [1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2))))/(b*d)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \text{Timed out}$$

```
[In] integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.40

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{2 dx + 2 c + \sin(2 dx + 2 c)}{4 \sqrt{b} d}$$

```
[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(sqrt(b)*d)
```

Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \int \frac{\cos(dx+c)^{\frac{5}{2}}}{\sqrt{b \cos(dx+c)}} dx$$

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(5/2)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 14.02 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int \frac{\cos^{\frac{5}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx \\ &= \frac{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (\sin(c+dx) + \sin(3c+3dx) + 4dx \cos(c+dx))}{4bd (\cos(2c+2dx) + 1)} \end{aligned}$$

[In] int(cos(c + d*x)^(5/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*b*d*(cos(2*c + 2*d*x) + 1))

$$3.175 \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	968
Rubi [A] (verified)	968
Mathematica [A] (verified)	969
Maple [A] (verified)	969
Fricas [A] (verification not implemented)	969
Sympy [A] (verification not implemented)	970
Maxima [A] (verification not implemented)	970
Giac [F]	970
Mupad [B] (verification not implemented)	971

Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sin(c+dx) \sqrt{\cos(c+dx)}}{d \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(3/2)}/\text{Sqrt}[b*\text{Cos}[c + d*x]], x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(\text{d}*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rule 2717

$\text{Int}[\sin[\text{Pi}/2 + (c_*) + (d_*)*(x_*)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{d \sqrt{b \cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(3/2)/Sqrt[b*Cos[c + d*x]],x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.84 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{d\sqrt{\cos(dx+c)b}}$	29
risch	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{d\sqrt{\cos(dx+c)b}}$	29

[In] int(cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/d/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{bd \sqrt{\cos(dx+c)}}$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

Sympy [A] (verification not implemented)

Time = 18.28 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.44

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \begin{cases} \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{d\sqrt{b \cos(c+dx)}} & \text{for } d \neq 0 \\ \frac{x \cos^{\frac{3}{2}}(c)}{\sqrt{b \cos(c)}} & \text{otherwise} \end{cases}$$

[In] integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Piecewise((sin(c + d*x)*sqrt(cos(c + d*x))/(d*sqrt(b*cos(c + d*x))), Ne(d, 0)), (x*cos(c)**(3/2)/sqrt(b*cos(c)), True))

Maxima [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.41

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sin(dx + c)}{\sqrt{bd}}$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(sqrt(b)*d)

Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^{\frac{3}{2}}}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(3/2)/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 14.62 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.47

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{\sqrt{b \cos(c + dx)}} dx = \frac{\sqrt{\cos(c + dx)} \sin(2c + 2dx) \sqrt{b \cos(c + dx)}}{bd (\cos(2c + 2dx) + 1)}$$

[In] `int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(1/2),x)`

[Out] `(cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(b*d*(cos(2*c + 2*d*x) + 1))`

$$3.176 \quad \int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx$$

Optimal result	972
Rubi [A] (verified)	972
Mathematica [A] (verified)	973
Maple [A] (verified)	973
Fricas [A] (verification not implemented)	973
Sympy [A] (verification not implemented)	974
Maxima [A] (verification not implemented)	974
Giac [F]	974
Mupad [B] (verification not implemented)	975

Optimal result

Integrand size = 23, antiderivative size = 24

$$\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx = \frac{x \sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

[Out] $x \cos(dx+c)^{(1/2)} / (b \cos(dx+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx = \frac{x \sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

[In] `Int[Sqrt[Cos[c + d*x]]/Sqrt[b*Cos[c + d*x]],x]`

[Out] `(x*Sqrt[Cos[c + d*x]])/Sqrt[b*Cos[c + d*x]]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{x \sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx = \frac{x \sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}}$$

[In] Integrate[Sqrt[Cos[c + d*x]]/Sqrt[b*Cos[c + d*x]],x]

[Out] (x*Sqrt[Cos[c + d*x]])/Sqrt[b*Cos[c + d*x]]

Maple [A] (verified)

Time = 2.34 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.88

method	result	size
risch	$\frac{x(\sqrt{\cos(dx+c)})}{\sqrt{\cos(dx+c)b}}$	21
default	$\frac{(\sqrt{\cos(dx+c)})(dx+c)}{d\sqrt{\cos(dx+c)b}}$	28

[In] int(cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] x*cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 97, normalized size of antiderivative = 4.04

$$\begin{aligned} &\int \frac{\sqrt{\cos(c+dx)}}{\sqrt{b \cos(c+dx)}} dx \\ &= \left[-\frac{\sqrt{-b} \log \left(2b \cos(dx+c)^2 + 2\sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) - b \right)}{2bd}, \frac{\arctan \left(\frac{\sqrt{b \cos(dx+c)}}{\sqrt{b \cos(dx+c)}} \right)}{\sqrt{bd}} \right] \end{aligned}$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b)/(b*d), arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(sqrt(b)*d)]

Sympy [A] (verification not implemented)

Time = 0.67 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\cos(c + dx)}}{\sqrt{b \cos(c + dx)}} dx = \frac{x \sqrt{\cos(c + dx)}}{\sqrt{b \cos(c + dx)}}$$

[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(1/2),x)

[Out] x*sqrt(cos(c + d*x))/sqrt(b*cos(c + d*x))

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.08

$$\int \frac{\sqrt{\cos(c + dx)}}{\sqrt{b \cos(c + dx)}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{\sqrt{bd}}$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(sqrt(b)*d)

Giac [F]

$$\int \frac{\sqrt{\cos(c + dx)}}{\sqrt{b \cos(c + dx)}} dx = \int \frac{\sqrt{\cos(dx + c)}}{\sqrt{b \cos(dx + c)}} dx$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/sqrt(b*cos(d*x + c)), x)

Mupad [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.54

$$\int \frac{\sqrt{\cos(c + dx)}}{\sqrt{b \cos(c + dx)}} dx = \frac{2x \cos(c + dx)^{3/2} \sqrt{b \cos(c + dx)}}{b (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(1/2),x)

[Out] (2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b*(cos(2*c + 2*d*x) + 1))

$$3.177 \quad \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx$$

Optimal result	976
Rubi [A] (verified)	976
Mathematica [A] (verified)	977
Maple [A] (verified)	977
Fricas [A] (verification not implemented)	977
Sympy [F]	978
Maxima [B] (verification not implemented)	978
Giac [F]	979
Mupad [F(-1)]	979

Optimal result

Integrand size = 23, antiderivative size = 33

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{d\sqrt{b\cos(c+dx)}}$$

[Out] $\operatorname{arctanh}(\sin(d*x+c))*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3855}

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)}\operatorname{arctanh}(\sin(c+dx))}{d\sqrt{b\cos(c+dx)}}$$

[In] $\operatorname{Int}[1/(\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]),x]$

[Out] $(\operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]]*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]])/(d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]])$

Rule 18

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \operatorname{Dist}[a^{(m-1/2)}*b^{(n+1/2)}*(\operatorname{Sqrt}[a*v]/\operatorname{Sqrt}[b*v]), \operatorname{Int}[u*v^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, m\}, x$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{ILtQ}[n-1/2, 0]$ && $\operatorname{IntegerQ}[m+n]$

Rule 3855

$\operatorname{Int}[\operatorname{csc}[(c_*)+(d_*)*(x_)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c+d*x]]/d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{d \sqrt{b \cos(c+dx)}}$$

[In] Integrate[1/(Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.88 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.21

method	result	size
default	$-\frac{2(\sqrt{\cos(dx+c)} \operatorname{arctanh}(\cot(dx+c)-\csc(dx+c)))}{d \sqrt{\cos(dx+c)} b}$	40
risch	$\frac{(\sqrt{\cos(dx+c)} \ln(e^{i(dx+c)}+i))}{\sqrt{\cos(dx+c)} b d} - \frac{(\sqrt{\cos(dx+c)} \ln(e^{i(dx+c)}-i))}{\sqrt{\cos(dx+c)} b d}$	73

[In] int(1/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] -2/d*cos(d*x+c)^(1/2)*arctanh(cot(d*x+c)-csc(d*x+c))/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.52

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx$$

$$= \left[\frac{\log\left(-\frac{b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right)}{2\sqrt{bd}}, \right.$$

$$\left. -\frac{\sqrt{-b}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)}{bd} \right]$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3/(sqrt(b)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b*d)]

Sympy [F]

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(c+dx)}\sqrt{\cos(c+dx)}} dx$$

[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(1/(sqrt(b*cos(c + d*x))*sqrt(cos(c + d*x))), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(29) = 58.

Time = 0.39 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.97

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c))}{2\sqrt{bd}}$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(sqrt(b)*d)

Giac [F]

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}} dx$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} dx$$

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)), x)

$$3.178 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

Optimal result	980
Rubi [A] (verified)	980
Mathematica [A] (verified)	981
Maple [A] (verified)	981
Fricas [A] (verification not implemented)	982
Sympy [F]	982
Maxima [B] (verification not implemented)	982
Giac [F]	983
Mupad [B] (verification not implemented)	983

Optimal result

Integrand size = 23, antiderivative size = 32

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

[Out] $\sin(d*x+c)/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3852, 8}

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

[In] $\text{Int}[1/(\text{Cos}[c + d*x]^{(3/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]]),x]$

[Out] $\text{Sin}[c + d*x]/(d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m - 1/2)}*b^{(n + 1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{ILtQ}[n - 1/2, 0] \&\& \text{IntegerQ}[m + n]$

Rule 3852

`Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \sec^2(c + dx) dx}{\sqrt{b \cos(c + dx)}} \\ &= -\frac{\sqrt{\cos(c + dx)} \text{Subst}(\int 1 dx, x, -\tan(c + dx))}{d \sqrt{b \cos(c + dx)}} \\ &= \frac{\sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00

$$\int \frac{1}{\cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}} dx = \frac{\sin(c + dx)}{d \sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)}}$$

[In] `Integrate[1/(Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]]),x]`

[Out] `Sin[c + d*x]/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]])`

Maple [A] (verified)

Time = 2.85 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)}{d \sqrt{\cos(dx+c)} \sqrt{\cos(dx+c)b}}$	29
risch	$\frac{ie^{-i(dx+c)}}{\sqrt{\cos(dx+c)b} \sqrt{\cos(dx+c)} d}$	34

[In] `int(1/cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `sin(d*x+c)/d/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)`

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.97

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sqrt{b\cos(dx+c)}\sin(dx+c)}{bd\cos(dx+c)^{\frac{3}{2}}}$$

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b*d*cos(d*x + c)^(3/2))

Sympy [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(c+dx)}\cos^{\frac{3}{2}}(c+dx)} dx$$

[In] integrate(1/cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Integral(1/(sqrt(b*cos(c + d*x))*cos(c + d*x)**(3/2)), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(28) = 56.

Time = 0.38 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.84

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{2\sqrt{b}\sin(2dx+2c)}{(b\cos(2dx+2c)^2 + b\sin(2dx+2c)^2 + 2b\cos(2dx+2c) + b)d}$$

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((b*cos(2*d*x + 2*c)^2 + b*sin(2*d*x + 2*c)^2 + 2*b*cos(2*d*x + 2*c) + b)*d)

Giac [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(dx+c)}\cos(dx+c)^{\frac{3}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(3/2)), x)

Mupad [B] (verification not implemented)

Time = 14.51 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.94

$$\begin{aligned} & \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx \\ &= \frac{\sqrt{b\cos(c+dx)}(\cos(2c+2dx) \operatorname{li} + \sin(2c+2dx) + \operatorname{li})}{bd\sqrt{\cos(c+dx)}(\cos(2c+2dx) + 1)} \end{aligned}$$

[In] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] ((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*li + sin(2*c + 2*d*x) + li))/(b*d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

$$3.179 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx$$

Optimal result	984
Rubi [A] (verified)	984
Mathematica [A] (verified)	985
Maple [A] (verified)	986
Fricas [A] (verification not implemented)	986
Sympy [F(-1)]	987
Maxima [B] (verification not implemented)	987
Giac [F]	988
Mupad [F(-1)]	988

Optimal result

Integrand size = 23, antiderivative size = 72

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{2d \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/d/\cos(d*x+c)^{(3/2)}/(b*\cos(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\cos(d*x+c)^{(1/2)}/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{2d \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[1/(\operatorname{Cos}[c+d*x])^{(5/2)}*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]],x]$

[Out] $(\operatorname{ArcTanh}[\operatorname{Sin}[c+d*x]]*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]])/(2*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]]) + \operatorname{Sin}[c+d*x]/(2*d*\operatorname{Cos}[c+d*x]^{(3/2)}*\operatorname{Sqrt}[b*\operatorname{Cos}[c+d*x]])$

Rule 18


```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m - 1/2)
)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{2\sqrt{b \cos(c+dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{2d \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.72

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \cos^2(c+dx) + \sin(c+dx)}{2d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

```
[In] Integrate[1/(Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]]),x]
```

```
[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/(2*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])
```

Maple [A] (verified)

Time = 3.32 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.17

method	result	size
default	$\frac{-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c)}{2d\sqrt{\cos(dx+c)b} \cos(dx+c)^{\frac{3}{2}}}$	84
risch	$-\frac{i(e^{2i(dx+c)}-1)}{2\sqrt{\cos(dx+c)b} \sqrt{\cos(dx+c)} (e^{2i(dx+c)}+1)d} - \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}-i)}{2\sqrt{\cos(dx+c)b}d} + \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}+i)}{2\sqrt{\cos(dx+c)b}d}$	122

[In] int(1/cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2/d*(-cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)-1)+cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)+1)+sin(d*x+c))/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(3/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.88

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

$$= \left[\frac{\sqrt{b}\cos(dx+c)^3 \log\left(-\frac{b\cos(dx+c)^3-2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c)-2b\cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}}{4bd\cos(dx+c)^3} \right. \\ \left. - \frac{\sqrt{-b}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)\cos(dx+c)^3 - \sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c)}{2bd\cos(dx+c)^3} \right]$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^3)]

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \text{Timed out}$$

[In] integrate(1/cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 661 vs. 2(60) = 120.

Time = 0.41 (sec) , antiderivative size = 661, normalized size of antiderivative = 9.18

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx =$$

$$\frac{4(\sin(4dx+4c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) - 4(\sin(4dx+2c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) + 4(\sin(4dx+4c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right) - 4(\sin(4dx+2c) + 2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right)}{\sqrt{b}d}$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

```
[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*sqrt(b)*d)
```

Giac [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(dx+c)}\cos(dx+c)^{\frac{5}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(5/2)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\cos(c+dx)^{5/2}\sqrt{b\cos(c+dx)}} dx$$

[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(1/2)), x)

$$3.180 \quad \int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

Optimal result	989
Rubi [A] (verified)	989
Mathematica [A] (verified)	990
Maple [A] (verified)	990
Fricas [A] (verification not implemented)	991
Sympy [F(-1)]	991
Maxima [B] (verification not implemented)	991
Giac [F]	992
Mupad [B] (verification not implemented)	992

Optimal result

Integrand size = 23, antiderivative size = 70

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} + \frac{\sin^3(c+dx)}{3d\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

[Out] $1/3*\sin(d*x+c)^3/d/\cos(d*x+c)^{(5/2)}/(b*\cos(d*x+c))^{(1/2)}+\sin(d*x+c)/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3852}

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sin^3(c+dx)}{3d\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}}$$

[In] $\text{Int}[1/(\text{Cos}[c + d*x]^{(7/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]]), x]$

[Out] $\text{Sin}[c + d*x]/(d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + \text{Sin}[c + d*x]^3/(3*d*\text{Cos}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m-1/2)}*b^{(n+1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b

, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d\sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{d\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3d \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.64

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b \cos(c+dx)}} dx = \frac{\sqrt{\cos(c+dx)}(\tan(c+dx) + \frac{1}{3}\tan^3(c+dx))}{d\sqrt{b \cos(c+dx)}}$$

[In] Integrate[1/(Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] (Sqrt[Cos[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.83 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.60

method	result	size
default	$\frac{(2(\cos^2(dx+c))+1)\sin(dx+c)}{3d\sqrt{\cos(dx+c)}b\cos(dx+c)^{\frac{5}{2}}}$	42
risch	$\frac{2i(4\cos(dx+c)+2i\sin(dx+c))}{3\sqrt{\cos(dx+c)}b\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)^2d}$	56

[In] int(1/cos(d*x+c)^(7/2)/(cos(d*x+c)*b)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*sin(d*x+c)/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(5/2)

Fricas [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.63

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{\sqrt{b\cos(dx+c)}(2\cos(dx+c)^2+1)\sin(dx+c)}{3bd\cos(dx+c)^{\frac{7}{2}}}$$

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(b*d*cos(d*x + c)^(7/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \text{Timed out}$$

[In] integrate(1/cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(1/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 294 vs. 2(60) = 120.

Time = 0.41 (sec) , antiderivative size = 294, normalized size of antiderivative = 4.20

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{4((3\cos(2dx+2c)+1)\sin(6dx+6c) + 3(3\cos(2dx+2c)+1)\sin(4dx+4c) - 3\cos(6dx+6c)\sin(2dx+2c) - 9\cos(4dx+4c)\sin(2dx+2c))}{3(2(3\cos(4dx+4c)+3\cos(2dx+2c)+1)\cos(6dx+6c) + \cos(6dx+6c)^2 + 6(3\cos(2dx+2c)+1)\cos(6dx+6c) + 9\cos(4dx+4c)^2 + 9\cos(2dx+2c)^2 + 6(\sin(4dx+4c) + \sin(2dx+2c))\sin(6dx+6c) + \sin(6dx+6c)^2 + 9\sin(4dx+4c)^2 + 18\sin(4dx+4c)\sin(2dx+2c) + 9\sin(2dx+2c)^2 + 6\cos(2dx+2c)+1)\sqrt{b}d}$$

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")

```
[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)
*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)
*sin(2*d*x + 2*c))/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6
*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4
*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + s
in(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)
^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*
d*x + 2*c) + 1)*sqrt(b)*d)
```

Giac [F]

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(dx+c)}\cos(dx+c)^{\frac{7}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(7/2)), x)

Mupad [B] (verification not implemented)

Time = 15.04 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.87

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{2\sqrt{b\cos(c+dx)}(\cos(2c+2dx)15i + \cos(4c+4dx)6i + \cos(6c+6dx)1i + 9\sin(2c+2dx) + 6\sin(4c+4dx) + \sin(6c+6dx) + 10i)}{3bd\sqrt{\cos(c+dx)}(15\cos(2c+2dx) + 6\cos(4c+4dx) + \cos(6c+6dx) + 10)}$$

[In] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(1/2)),x)

[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.181 \quad \int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

Optimal result	993
Rubi [A] (verified)	993
Mathematica [A] (verified)	995
Maple [A] (verified)	995
Fricas [A] (verification not implemented)	995
Sympy [F(-1)]	996
Maxima [B] (verification not implemented)	996
Giac [F]	997
Mupad [F(-1)]	998

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{3\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{8d\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{4d\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{3\sin(c+dx)}{8d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/d/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2)+3/8*sin(d*x+c)/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+3/8*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \frac{3\sqrt{\cos(c+dx)}\operatorname{arctanh}(\sin(c+dx))}{8d\sqrt{b\cos(c+dx)}} + \frac{3\sin(c+dx)}{8d\cos^{\frac{3}{2}}(c+dx)\sqrt{b\cos(c+dx)}} + \frac{\sin(c+dx)}{4d\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(9/2)*Sqrt[b*Cos[c + d*x]]),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(8*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(4*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]) + (3*Sin[c + d*x])/(8*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \sec^5(c + dx) dx}{\sqrt{b \cos(c + dx)}} \\
 &= \frac{\sin(c + dx)}{4d \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)}} + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int \sec^3(c + dx) dx}{4\sqrt{b \cos(c + dx)}} \\
 &= \frac{\sin(c + dx)}{4d \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)}} + \frac{3 \sin(c + dx)}{8d \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int \sec(c + dx) dx}{8\sqrt{b \cos(c + dx)}} \\
 &= \frac{3 \arctanh(\sin(c + dx)) \sqrt{\cos(c + dx)}}{8d \sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{\sin(c + dx)}{4d \cos^{\frac{7}{2}}(c + dx) \sqrt{b \cos(c + dx)}} + \frac{3 \sin(c + dx)}{8d \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.62

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{3\operatorname{arctanh}(\sin(c+dx))\cos^4(c+dx) + (2+3\cos^2(c+dx))\sin(c+dx)}{8d\cos^{\frac{7}{2}}(c+dx)\sqrt{b\cos(c+dx)}}$$

[In] Integrate[1/(Cos[c + d*x]^(9/2)*Sqrt[b*Cos[c + d*x]]), x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/ (8*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 3.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.96

method	result
default	$\frac{-3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)-1)+3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)+1)+3(\cos^2(dx+c))\sin(dx+c)+2\sin(dx+c)}{8d\sqrt{\cos(dx+c)b}\cos(dx+c)^{\frac{7}{2}}}$
risch	$-\frac{i(3e^{6i(dx+c)}+11e^{4i(dx+c)}-11e^{2i(dx+c)}-3)}{8\sqrt{\cos(dx+c)b}\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)^3d} + \frac{3(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}+i)}{8\sqrt{\cos(dx+c)b}d} - \frac{3(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}-i)}{8\sqrt{\cos(dx+c)b}d}$

[In] int(1/cos(d*x+c)^(9/2)/(cos(d*x+c)*b)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/8/d*(-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)+3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.18

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx$$

$$= \frac{\left[3\sqrt{b}\cos(dx+c)^5 \log\left(\frac{-b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right) + 2\sqrt{b\cos(dx+c)}(3\cos(dx+c)^2 + 2) \right]}{16bd\cos(dx+c)^5}$$

$$- \frac{3\sqrt{-b}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)\cos(dx+c)^5 - \sqrt{b\cos(dx+c)}(3\cos(dx+c)^2 + 2)\sqrt{\cos(dx+c)}}{8bd\cos(dx+c)^5}$$

```
[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="fricas")
[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^5), -1/8*(3*sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^5 - sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*sin(d*x + c))/(b*d*cos(d*x + c)^5)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{9}{2}}(c + dx)\sqrt{b \cos(c + dx)}} dx = \text{Timed out}$$

```
[In] integrate(1/cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1656 vs. 2(89) = 178.

Time = 0.41 (sec) , antiderivative size = 1656, normalized size of antiderivative = 15.48

$$\int \frac{1}{\cos^{\frac{9}{2}}(c + dx)\sqrt{b \cos(c + dx)}} dx = \text{Too large to display}$$

```
[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="maxima")
```

```
[Out] -1/16*(12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(sin(8*d*x + 8*c) + 4*sin(6*d*x + 6*c) + 6*sin(4*d*x + 4*c) + 4*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*s
```

```

in(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3
*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(
8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2
*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(
2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8
*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin
(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*
x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)
*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan
2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c)))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) +
6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*
c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(
4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x
+ 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*
x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c)
+ 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c)))))/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1
)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d
*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x +
2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2
+ 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d
*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c
))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*si
n(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c
) + 1)*sqrt(b)*d)

```

Giac [F]

$$\int \frac{1}{\cos^{\frac{9}{2}}(c+dx)\sqrt{b\cos(c+dx)}} dx = \int \frac{1}{\sqrt{b\cos(dx+c)}\cos(dx+c)^{\frac{9}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*cos(d*x + c))*cos(d*x + c)^(9/2)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{9}{2}}(c + dx) \sqrt{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^{9/2} \sqrt{b \cos(c + dx)}} dx$$

```
[In] int(1/(cos(c + d*x)^(9/2)*(b*cos(c + d*x))^(1/2)), x)
```

```
[Out] int(1/(cos(c + d*x)^(9/2)*(b*cos(c + d*x))^(1/2)), x)
```

$$3.182 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	999
Rubi [A] (verified)	999
Mathematica [A] (verified)	1000
Maple [A] (verified)	1001
Fricas [A] (verification not implemented)	1001
Sympy [F(-1)]	1001
Maxima [A] (verification not implemented)	1002
Giac [F]	1002
Mupad [B] (verification not implemented)	1002

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{3x \sqrt{\cos(c+dx)}}{8b \sqrt{b \cos(c+dx)}} + \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8bd \sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4bd \sqrt{b \cos(c+dx)}}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/b/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{3x \sqrt{\cos(c+dx)}}{8b \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4bd \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{8bd \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(11/2)}/(b*\text{Cos}[c + d*x]^{(3/2)}), x]$

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

```
Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]
```

Rule 17

```
Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m + 1/2)
)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]
```

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[
c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2
*n]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos^4(c+dx) dx}{b\sqrt{b \cos(c+dx)}} \\
 &= \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4bd\sqrt{b \cos(c+dx)}} + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int \cos^2(c+dx) dx}{4b\sqrt{b \cos(c+dx)}} \\
 &= \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8bd\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4bd\sqrt{b \cos(c+dx)}} + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int 1 dx}{8b\sqrt{b \cos(c+dx)}} \\
 &= \frac{3x\sqrt{\cos(c+dx)}}{8b\sqrt{b \cos(c+dx)}} + \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8bd\sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4bd\sqrt{b \cos(c+dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.41 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.51

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{\cos^{\frac{3}{2}}(c+dx)(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))}{32d(b \cos(c+dx))^{3/2}}$$

```
[In] Integrate[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(3/2), x]
```

```
[Out] (Cos[c + d*x]^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])
/(32*d*(b*Cos[c + d*x])^(3/2))
```


Maple [A] (verified)

Time = 2.85 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.61

method	result	size
default	$\frac{(\sqrt{\cos(dx+c)})(2\sin(dx+c)(\cos^3(dx+c))+3\cos(dx+c)\sin(dx+c)+3dx+3c)}{8db\sqrt{\cos(dx+c)}b}$	65
risch	$\frac{3x(\sqrt{\cos(dx+c)})}{8b\sqrt{\cos(dx+c)}b} + \frac{(\sqrt{\cos(dx+c)})\sin(4dx+4c)}{32b\sqrt{\cos(dx+c)}bd} + \frac{(\sqrt{\cos(dx+c)})\sin(2dx+2c)}{4b\sqrt{\cos(dx+c)}bd}$	96

[In] `int(cos(d*x+c)^(11/2)/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)`[Out]
$$\frac{1}{8} \frac{d \cos(d*x+c)^{(1/2)} (2 \sin(d*x+c) \cos(d*x+c)^3 + 3 \cos(d*x+c) \sin(d*x+c) + 3 d*x + 3c)}{b (\cos(d*x+c) \cdot b)^{(1/2)}}$$
Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.70

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \left[\frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b} \log \left(\frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b}}{16 b^2 d} \right)}{16 b^2 d} \right]$$

[In] `integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")`[Out]
$$\left[\frac{1}{16} (2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b} \log(2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b})) / (b^2 d), \frac{1}{8} (\sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) + 3 \sqrt{b} \arctan(\sqrt{b \cos(dx+c)} \sin(dx+c) / (\sqrt{b} \cos(dx+c)^{(3/2)})) / (b^2 d) \right]$$
Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(3/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.46

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c))\right)}{32 b^{\frac{3}{2}} d}$$

```
[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")
```

```
[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(3/2)*d)
```

Giac [F]

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^{\frac{11}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

```
[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")
```

```
[Out] integrate(cos(d*x + c)^(11/2)/(b*cos(d*x + c))^(3/2), x)
```

Mupad [B] (verification not implemented)

Time = 15.22 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.73

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx))}{32 b^2 d (\cos(2c + 2dx) + 1)}$$

```
[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(3/2),x)
```

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*b^2*d*(cos(2*c + 2*d*x) + 1))
```

$$3.183 \quad \int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Optimal result	1003
Rubi [A] (verified)	1003
Mathematica [A] (verified)	1004
Maple [A] (verified)	1004
Fricas [A] (verification not implemented)	1005
Sympy [F(-1)]	1005
Maxima [A] (verification not implemented)	1005
Giac [F]	1006
Mupad [B] (verification not implemented)	1006

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3bd\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx = \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\cos(c+dx)}}{3bd\sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(9/2)}/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) - (\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2713

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^3(c + dx) dx}{b\sqrt{b \cos(c + dx)}} \\ &= -\frac{\sqrt{\cos(c + dx)} \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{bd\sqrt{b \cos(c + dx)}} \\ &= \frac{\sqrt{\cos(c + dx)} \sin(c + dx)}{bd\sqrt{b \cos(c + dx)}} - \frac{\sqrt{\cos(c + dx)} \sin^3(c + dx)}{3bd\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \frac{\cos^{\frac{3}{2}}(c + dx)(5 + \cos(2(c + dx))) \sin(c + dx)}{6d(b \cos(c + dx))^{\frac{3}{2}}}$$

```
[In] Integrate[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(3/2), x]
```

```
[Out] (Cos[c + d*x]^(3/2)*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*d*(b*Cos[c + d*x])^(3/2))
```

Maple [A] (verified)

Time = 3.06 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.57

method	result	size
default	$\frac{(2 + \cos^2(dx+c)) \sin(dx+c) (\sqrt{\cos(dx+c)})}{3db \sqrt{\cos(dx+c)b}}$	43
risch	$\frac{3 \sin(dx+c) (\sqrt{\cos(dx+c)})}{4bd \sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(3dx+3c)}{12b \sqrt{\cos(dx+c)b d}}$	69

```
[In] int(cos(d*x+c)^(9/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/b/(cos(d*x+c)*b)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.36 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\sqrt{b \cos(dx + c)} (\cos(dx + c)^2 + 2) \sin(dx + c)}{3 b^2 d \sqrt{\cos(dx + c)}}$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\sin(3 dx + 3 c) + 9 \sin\left(\frac{1}{3} \arctan\left(\frac{\sin(3 dx + 3 c)}{\cos(3 dx + 3 c)}\right)\right)}{12 b^{\frac{3}{2}} d}$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(3/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^{\frac{9}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(9/2)/(b*cos(d*x + c))^(3/2), x)

Mupad [B] (verification not implemented)

Time = 0.69 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.79

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12b^2 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(3/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*b^2*d*(cos(2*c + 2*d*x) + 1))

$$3.184 \quad \int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx$$

Optimal result	1007
Rubi [A] (verified)	1007
Mathematica [A] (verified)	1008
Maple [A] (verified)	1008
Fricas [A] (verification not implemented)	1009
Sympy [F(-1)]	1009
Maxima [A] (verification not implemented)	1009
Giac [F]	1010
Mupad [B] (verification not implemented)	1010

Optimal result

Integrand size = 23, antiderivative size = 69

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx = \frac{x \sqrt{\cos(c+dx)}}{2b \sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2bd \sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/b/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{3}{2}}} dx = \frac{x \sqrt{\cos(c+dx)}}{2b \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2bd \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(x*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*b*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(2*b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] := \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] := \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b$

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^2(c + dx) dx}{b\sqrt{b \cos(c + dx)}} \\ &= \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2bd\sqrt{b \cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \int 1 dx}{2b\sqrt{b \cos(c + dx)}} \\ &= \frac{x\sqrt{\cos(c + dx)}}{2b\sqrt{b \cos(c + dx)}} + \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2bd\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\cos^{\frac{3}{2}}(c + dx)(2(c + dx) + \sin(2(c + dx)))}{4d(b \cos(c + dx))^{3/2}}$$

[In] Integrate[Cos[c + d*x]^(7/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (Cos[c + d*x]^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*(b*Cos[c + d*x])^(3/2))

Maple [A] (verified)

Time = 2.90 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{(\sqrt{\cos(dx+c)})(\cos(dx+c) \sin(dx+c)+dx+c)}{2db\sqrt{\cos(dx+c)b}}$	45
risch	$\frac{x(\sqrt{\cos(dx+c)})}{2b\sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(2dx+2c)}{4b\sqrt{\cos(dx+c)b}d}$	61

[In] int(cos(d*x+c)^(7/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/2/d*cos(d*x+c)^(1/2)*(cos(d*x+c)*sin(d*x+c)+d*x+c)/b/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.28

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{3}{2}}} dx = \left[\frac{2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) - \sqrt{-b}\log\left(2b\cos(dx+c)^2 + 2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) - b\right)}{4b^2d} \right]$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

```
[Out] [1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^2*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^2*d)]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{3}{2}}} dx = \frac{2dx + 2c + \sin(2dx + 2c)}{4b^{\frac{3}{2}}d}$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(3/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^{\frac{7}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/(b*cos(d*x + c))^(3/2), x)

Mupad [B] (verification not implemented)

Time = 14.87 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.94

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx)) + 4dx \cos(c + dx)}{4b^2 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(7/2)/(b*cos(c + d*x))^(3/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*b^2*d*(cos(2*c + 2*d*x) + 1))

$$3.185 \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1011
Rubi [A] (verified)	1011
Mathematica [A] (verified)	1012
Maple [A] (verified)	1012
Fricas [A] (verification not implemented)	1012
Sympy [F(-1)]	1013
Maxima [A] (verification not implemented)	1013
Giac [F]	1013
Mupad [B] (verification not implemented)	1013

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{bd\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{\sin(c+dx)\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)}/(b*\text{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m)}*((b_*)*(v_))^{(n)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2717

$\text{Int}[\sin[\text{Pi}/2 + (c_*) + (d_*)*(x)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{b\sqrt{b\cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{bd\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{d(b\cos(c+dx))^{3/2}}$$

[In] Integrate[Cos[c + d*x]^(5/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (Cos[c + d*x]^(3/2)*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(3/2))

Maple [A] (verified)

Time = 2.87 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{bd\sqrt{\cos(dx+c)b}}$	32
risch	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{bd\sqrt{\cos(dx+c)b}}$	32

[In] int(cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/b/d/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{\sqrt{b\cos(dx+c)} \sin(dx+c)}{b^2 d \sqrt{\cos(dx+c)}}$$

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(3/2), x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \frac{\sin(dx + c)}{b^{\frac{3}{2}}d}$$

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2), x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(3/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \int \frac{\cos(dx + c)^{\frac{5}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(5/2)/(b*cos(d*x + c))^(3/2), x)

Mupad [B] (verification not implemented)

Time = 0.35 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \frac{\sqrt{\cos(c + dx)} \sin(2c + 2dx) \sqrt{b \cos(c + dx)}}{b^2 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(5/2)/(b*cos(c + d*x))^(3/2), x)

[Out] (cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(b^2*d*(cos(2*c + 2*d*x) + 1))

$$3.186 \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1014
Rubi [A] (verified)	1014
Mathematica [A] (verified)	1015
Maple [A] (verified)	1015
Fricas [A] (verification not implemented)	1015
Sympy [A] (verification not implemented)	1016
Maxima [A] (verification not implemented)	1016
Giac [F]	1016
Mupad [B] (verification not implemented)	1017

Optimal result

Integrand size = 23, antiderivative size = 27

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{x \sqrt{\cos(c+dx)}}{b \sqrt{b \cos(c+dx)}}$$

[Out] $x \cos(dx+c)^{(1/2)} / b (b \cos(dx+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{3/2}} dx = \frac{x \sqrt{\cos(c+dx)}}{b \sqrt{b \cos(c+dx)}}$$

[In] `Int[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(3/2),x]`

[Out] `(x*Sqrt[Cos[c + d*x]])/(b*Sqrt[b*Cos[c + d*x]])`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{b\sqrt{b\cos(c+dx)}} \\ &= \frac{x\sqrt{\cos(c+dx)}}{b\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \frac{x\sqrt{b\cos(c+dx)}}{b^2\sqrt{\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(3/2), x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/(b^2*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
risch	$\frac{x(\sqrt{\cos(dx+c)})}{b\sqrt{\cos(dx+c)b}}$	24
default	$\frac{(\sqrt{\cos(dx+c)})(dx+c)}{db\sqrt{\cos(dx+c)b}}$	31

[In] int(cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] x*cos(d*x+c)^(1/2)/b/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.59

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b\cos(c+dx))^{3/2}} dx = \left[-\frac{\sqrt{-b} \log\left(2b\cos(dx+c)^2 + 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{\cos(dx+c)}\sin(dx+c)\right)}{2b^2d} \right]$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2), x, algorithm="fricas")

[Out] $[-1/2*\sqrt{-b}*\log(2*b*\cos(dx + c)^2 + 2*\sqrt{b*\cos(dx + c)}*\sqrt{-b}*\sqrt{\cos(dx + c)}*\sin(dx + c) - b)/(b^2*d), \arctan(\sqrt{b*\cos(dx + c)}*\sin(dx + c)/(\sqrt{b*\cos(dx + c)}^{3/2}))/b^{3/2}*d]$

Sympy [A] (verification not implemented)

Time = 10.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{x \cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}}$$

[In] `integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(3/2),x)`

[Out] `x*cos(c + d*x)**(3/2)/(b*cos(c + d*x))**(3/2)`

Maxima [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{3}{2}}d}$$

[In] `integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/b^(3/2)*d`

Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\cos(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] `integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)^(3/2)/(b*cos(d*x + c))^(3/2), x)`

Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.37

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{3}{2}}} dx = \frac{2x \cos(c + dx)^{\frac{3}{2}} \sqrt{b \cos(c + dx)}}{b^2 (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(3/2),x)

[Out] (2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b^2*(cos(2*c + 2*d*x) + 1)
)

$$3.187 \quad \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1018
Rubi [A] (verified)	1018
Mathematica [A] (verified)	1019
Maple [A] (verified)	1019
Fricas [A] (verification not implemented)	1019
Sympy [F]	1020
Maxima [B] (verification not implemented)	1020
Giac [F]	1020
Mupad [F(-1)]	1021

Optimal result

Integrand size = 23, antiderivative size = 36

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{bd\sqrt{b \cos(c+dx)}}$$

[Out] $\operatorname{arctanh}(\sin(dx+c))\cos(dx+c)^{(1/2)}/b/d/(b\cos(dx+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3855}

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx = \frac{\sqrt{\cos(c+dx)}\operatorname{arctanh}(\sin(c+dx))}{bd\sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]/(b*\operatorname{Cos}[c + d*x])^{(3/2)}, x]$

[Out] $(\operatorname{ArcTanh}[\operatorname{Sin}[c + d*x]]*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]])/(b*d*\operatorname{Sqrt}[b*\operatorname{Cos}[c + d*x]])$

Rule 17

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \operatorname{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\operatorname{Sqrt}[b*v]/\operatorname{Sqrt}[a*v]), \operatorname{Int}[u*v^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, m\}, x \&\& \operatorname{IntegerQ}[m] \&\& \operatorname{IGtQ}[n+1/2, 0] \&\& \operatorname{IntegerQ}[m+n]$

Rule 3855

$\operatorname{Int}[\operatorname{csc}[(c_*) + (d_*)*(x_)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]]/d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{b\sqrt{b\cos(c+dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{bd\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{\sqrt{\cos(c+dx)}}{(b\cos(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\cos^{3/2}(c+dx)}{d(b\cos(c+dx))^{3/2}}$$

[In] Integrate[Sqrt[Cos[c + d*x]]/(b*Cos[c + d*x])^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^(3/2))/(d*(b*Cos[c + d*x])^(3/2))

Maple [A] (verified)

Time = 2.94 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.19

method	result	size
default	$-\frac{2(\sqrt{\cos(dx+c)})\operatorname{arctanh}(\cot(dx+c)-\csc(dx+c))}{d\sqrt{\cos(dx+c)}bb}$	43
risch	$-\frac{(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}-i)}{b\sqrt{\cos(dx+c)}bd} + \frac{(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}+i)}{b\sqrt{\cos(dx+c)}bd}$	79

[In] int(cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] -2/d*cos(d*x+c)^(1/2)*arctanh(cot(d*x+c)-csc(d*x+c))/(cos(d*x+c)*b)^(1/2)/b

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.22

$$\int \frac{\sqrt{\cos(c+dx)}}{(b\cos(c+dx))^{3/2}} dx = \left[\frac{\log\left(-\frac{b\cos(dx+c)^3 - 2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c) - 2b\cos(dx+c)}{\cos(dx+c)^3}\right)}{2b^{3/2}d}, \right. \\ \left. -\frac{\sqrt{-b}\operatorname{arctan}\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)}{b^2d} \right]$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3/(b^(3/2)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b^2*d)]

Sympy [F]

$$\int \frac{\sqrt{\cos(c + dx)}}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sqrt{\cos(c + dx)}}{(b \cos(c + dx))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(sqrt(cos(c + d*x))/(b*cos(c + d*x))**(3/2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.37 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.81

$$\int \frac{\sqrt{\cos(c + dx)}}{(b \cos(c + dx))^{3/2}} dx = \frac{\log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - \log(\cos(dx + c)^2 + \sin(dx + c)^2 - 2 \sin(dx + c) + 1)}{2 b^{\frac{3}{2}} d}$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(3/2)*d)

Giac [F]

$$\int \frac{\sqrt{\cos(c + dx)}}{(b \cos(c + dx))^{3/2}} dx = \int \frac{\sqrt{\cos(dx + c)}}{(b \cos(dx + c))^{\frac{3}{2}}} dx$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/(b*cos(d*x + c))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx = \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{3/2}} dx$$

```
[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(3/2), x)
```

```
[Out] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(3/2), x)
```

$$3.188 \quad \int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1022
Rubi [A] (verified)	1022
Mathematica [A] (verified)	1023
Maple [A] (verified)	1023
Fricas [A] (verification not implemented)	1024
Sympy [F]	1024
Maxima [B] (verification not implemented)	1024
Giac [F]	1024
Mupad [B] (verification not implemented)	1025

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2}} dx = \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)/b/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3852, 8}

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{3/2}} dx = \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[1/(\text{Sqrt}[\text{Cos}[c + d*x]]*(b*\text{Cos}[c + d*x])^{(3/2)}),x]$

[Out] $\text{Sin}[c + d*x]/(b*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m - 1/2)}*b^{(n + 1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \ \&\& \ \text{IntegerQ}[m] \ \&\& \ \text{ILtQ}[n - 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \sec^2(c + dx) dx}{b\sqrt{b \cos(c + dx)}} \\ &= -\frac{\sqrt{\cos(c + dx)} \text{Subst}(\int 1 dx, x, -\tan(c + dx))}{bd\sqrt{b \cos(c + dx)}} \\ &= \frac{\sin(c + dx)}{bd\sqrt{\cos(c + dx)}\sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{1}{\sqrt{\cos(c + dx)}(b \cos(c + dx))^{3/2}} dx = \frac{\sqrt{\cos(c + dx)} \sin(c + dx)}{d(b \cos(c + dx))^{3/2}}$$

```
[In] Integrate[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(3/2)),x]
```

```
[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(3/2))
```

Maple [A] (verified)

Time = 2.98 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)}{bd\sqrt{\cos(dx+c)}\sqrt{\cos(dx+c)b}}$	32
risch	$\frac{ie^{-i(dx+c)}}{b\sqrt{\cos(dx+c)b}\sqrt{\cos(dx+c)}d}$	37

```
[In] int(1/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] sin(d*x+c)/b/d/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2}} dx = \frac{\sqrt{b\cos(dx+c)}\sin(dx+c)}{b^2 d \cos(dx+c)^{\frac{3}{2}}}$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^2*d*cos(d*x + c)^(3/2))

Sympy [F]

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2}} dx = \int \frac{1}{(b\cos(c+dx))^{\frac{3}{2}}\sqrt{\cos(c+dx)}} dx$$

[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Integral(1/((b*cos(c + d*x))**(3/2)*sqrt(cos(c + d*x))), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(31) = 62.

Time = 0.39 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.91

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2}} dx = \frac{2\sqrt{b}\sin(2dx+2c)}{(b^2\cos(2dx+2c)^2 + b^2\sin(2dx+2c)^2 + 2b^2\cos(2dx+2c) + b^2)}$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((b^2*cos(2*d*x + 2*c)^2 + b^2*sin(2*d*x + 2*c)^2 + 2*b^2*cos(2*d*x + 2*c) + b^2)*d)

Giac [F]

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2}} dx = \int \frac{1}{(b\cos(dx+c))^{\frac{3}{2}}\sqrt{\cos(dx+c)}} dx$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*sqrt(cos(d*x + c))), x)

Mupad [B] (verification not implemented)

Time = 14.56 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.77

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{3/2}} dx = \frac{\sqrt{b\cos(c+dx)}(\cos(2c+2dx)1i + \sin(2c+2dx) + 1i)}{b^2 d \sqrt{\cos(c+dx)}(\cos(2c+2dx) + 1)}$$

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(3/2)),x)

[Out] ((b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(b^2*d*cos(c + d*x)^(1/2)*(cos(2*c + 2*d*x) + 1))

$$3.189 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1026
Rubi [A] (verified)	1026
Mathematica [A] (verified)	1027
Maple [A] (verified)	1028
Fricas [A] (verification not implemented)	1028
Sympy [F]	1029
Maxima [B] (verification not implemented)	1029
Giac [F]	1030
Mupad [F(-1)]	1030

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2bd\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}}$$

[Out] 1/2*sin(d*x+c)/b/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+1/2*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{\sqrt{\cos(c+dx)}\operatorname{arctanh}(\sin(c+dx))}{2bd\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2bd \cos^{\frac{3}{2}}(c+dx)\sqrt{b \cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]])/(2*b*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(2*b*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[a^(m - 1/2)
)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^3(c+dx) dx}{b\sqrt{b}\cos(c+dx)} \\ &= \frac{\sin(c+dx)}{2bd\cos^{\frac{3}{2}}(c+dx)\sqrt{b}\cos(c+dx)} + \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{2b\sqrt{b}\cos(c+dx)} \\ &= \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2bd\sqrt{b}\cos(c+dx)} + \frac{\sin(c+dx)}{2bd\cos^{\frac{3}{2}}(c+dx)\sqrt{b}\cos(c+dx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.67

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{\frac{3}{2}}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\cos^2(c+dx) + \sin(c+dx)}{2d\sqrt{\cos(c+dx)}(b\cos(c+dx))^{\frac{3}{2}}}$$

```
[In] Integrate[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(3/2)),x]
```

```
[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x])/(2*d*Sqrt[Cos[c + d*x]
])* (b*Cos[c + d*x])^(3/2)
```

Maple [A] (verified)

Time = 2.90 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c)}{2db\sqrt{\cos(dx+c)}b \cos(dx+c)^{\frac{3}{2}}}$	87
risch	$-\frac{i(e^{2i(dx+c)}-1)}{2b\sqrt{\cos(dx+c)}b\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)d} + \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}+i)}{2b\sqrt{\cos(dx+c)}bd} - \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}-i)}{2b\sqrt{\cos(dx+c)}bd}$	131

```
[In] int(1/cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2/d*(-cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)-1)+cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)+1)+sin(d*x+c))/b/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(3/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.65

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{\frac{3}{2}}} dx = \left[\frac{\sqrt{b} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^3 - 2\sqrt{b \cos(dx+c)}\sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^3}\right)}{4b^2d \cos(dx+c)^3} - \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{b \cos(dx+c)}\sqrt{-b} \sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right) \cos(dx+c)^3 - \sqrt{b \cos(dx+c)} \sqrt{\cos(dx+c)} \sin(dx+c)}{2b^2d \cos(dx+c)^3} \right]$$

```
[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")
```

```
[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^2*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b^2*d*cos(d*x + c)^3)]
```


Giac [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \int \frac{1}{(b\cos(dx+c))^{\frac{3}{2}}\cos(dx+c)^{\frac{3}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(3/2)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \int \frac{1}{\cos(c+dx)^{3/2}(b\cos(c+dx))^{3/2}} dx$$

[In] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(3/2)), x)

$$3.190 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1031
Rubi [A] (verified)	1031
Mathematica [A] (verified)	1032
Maple [A] (verified)	1032
Fricas [A] (verification not implemented)	1033
Sympy [F(-1)]	1033
Maxima [B] (verification not implemented)	1033
Giac [F]	1034
Mupad [B] (verification not implemented)	1034

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3bd \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}}$$

[Out] $1/3*\sin(d*x+c)^3/b/d/\cos(d*x+c)^{(5/2)/(b*\cos(d*x+c))^{(1/2)}+\sin(d*x+c)/b/d/c \cos(d*x+c)^{(1/2)/(b*\cos(d*x+c))^{(1/2)}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3852}

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{\sin^3(c+dx)}{3bd \cos^{\frac{5}{2}}(c+dx)\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b \cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] Sin[c + d*x]/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]^3/(3*b*d*Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b

, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{b\sqrt{b\cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{bd\sqrt{b\cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{bd\sqrt{\cos(c+dx)}\sqrt{b\cos(c+dx)}} + \frac{\sin^3(c+dx)}{3bd\cos^{\frac{5}{2}}(c+dx)\sqrt{b\cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{\cos^{\frac{3}{2}}(c+dx) (\tan(c+dx) + \frac{1}{3}\tan^3(c+dx))}{d(b\cos(c+dx))^{3/2}}$$

[In] Integrate[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2)), x]

[Out] (Cos[c + d*x]^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*(b*Cos[c + d*x])^(3/2))

Maple [A] (verified)

Time = 3.24 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

method	result	size
default	$\frac{(2(\cos^2(dx+c))+1)\sin(dx+c)}{3db\sqrt{\cos(dx+c)}b\cos(dx+c)^{\frac{5}{2}}}$	45
risch	$\frac{2i(4\cos(dx+c)+2i\sin(dx+c))}{3b\sqrt{\cos(dx+c)}b\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)^2d}$	59

[In] int(1/cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*sin(d*x+c)/b/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(5/2)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.58

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{\sqrt{b\cos(dx+c)}(2\cos(dx+c)^2+1)\sin(dx+c)}{3b^2d\cos(dx+c)^{\frac{7}{2}}}$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(2*cos(d*x + c)^2 + 1)*sin(d*x + c)/(b^2*d*cos(d*x + c)^(7/2))

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(1/cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(3/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(66) = 132.

Time = 0.41 (sec) , antiderivative size = 311, normalized size of antiderivative = 4.09

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{1}{3(b\cos(6dx+6c)^2+9b\cos(4dx+4c)^2+9b\cos(2dx+2c)^2+b}$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="maxima")

```
[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)
*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)
*sin(2*d*x + 2*c))/((b*cos(6*d*x + 6*c)^2 + 9*b*cos(4*d*x + 4*c)^2 + 9*b*cos(2*d*x + 2*c)^2 + b*sin(6*d*x + 6*c)^2 + 9*b*sin(4*d*x + 4*c)^2 + 18*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*b*sin(2*d*x + 2*c)^2 + 2*(3*b*cos(4*d*x + 4*c) + 3*b*cos(2*d*x + 2*c) + b)*cos(6*d*x + 6*c) + 6*(3*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 6*b*cos(2*d*x + 2*c) + 6*(b*sin(4*d*x + 4*c) + b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b)*sqrt(b)*d
```

Giac [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \int \frac{1}{(b\cos(dx+c))^{\frac{3}{2}}\cos(dx+c)^{\frac{5}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(5/2)), x)

Mupad [B] (verification not implemented)

Time = 16.41 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.72

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{2\sqrt{b\cos(c+dx)}(\cos(2c+2dx)15i + \cos(4c+4dx)6i + \cos(6c+6dx)1i + 9\sin(2c+2dx) + 6\sin(4c+4dx) + \sin(6c+6dx) + 10i)}{3b^2d\sqrt{\cos(c+dx)}(15\cos(2c+2dx) + 6\cos(4c+4dx) + \cos(6c+6dx) + 10)}$$

[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(3/2)),x)

[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b^2*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.191 \quad \int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx$$

Optimal result	1035
Rubi [A] (verified)	1035
Mathematica [A] (verified)	1037
Maple [A] (verified)	1037
Fricas [A] (verification not implemented)	1037
Sympy [F(-1)]	1038
Maxima [B] (verification not implemented)	1038
Giac [F]	1039
Mupad [F(-1)]	1040

Optimal result

Integrand size = 23, antiderivative size = 116

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{3 \operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{8bd \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx)}{8bd \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/b/d/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2)+3/8*sin(d*x+c)/b/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+3/8*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b \cos(c+dx))^{3/2}} dx = \frac{3 \sqrt{\cos(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{8bd \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx)}{8bd \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4bd \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]]/(8*b*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(4*b*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]) + (3*Sin[c + d*x])/(8*b*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[a^{(m - 1/2)}*b^{(n + 1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& !\text{IntegerQ}[m] \&\& \text{ILtQ}[n - 1/2, 0] \&\& \text{IntegerQ}[m + n]$

Rule 3853

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n - 1)}/(d*(n - 1)), x] + \text{Dist}[b^2*((n - 2)/(n - 1)), \text{Int}[(b*\text{Csc}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \& \& \text{IntegerQ}[2*n]$

Rule 3855

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \sec^5(c + dx) dx}{b\sqrt{b \cos(c + dx)}} \\
 &= \frac{\sin(c + dx)}{4bd \cos^{\frac{7}{2}}(c + dx)\sqrt{b \cos(c + dx)}} + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int \sec^3(c + dx) dx}{4b\sqrt{b \cos(c + dx)}} \\
 &= \frac{\sin(c + dx)}{4bd \cos^{\frac{7}{2}}(c + dx)\sqrt{b \cos(c + dx)}} + \frac{3 \sin(c + dx)}{8bd \cos^{\frac{3}{2}}(c + dx)\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int \sec(c + dx) dx}{8b\sqrt{b \cos(c + dx)}} \\
 &= \frac{3\text{arctanh}(\sin(c + dx))\sqrt{\cos(c + dx)}}{8bd\sqrt{b \cos(c + dx)}} \\
 &\quad + \frac{\sin(c + dx)}{4bd \cos^{\frac{7}{2}}(c + dx)\sqrt{b \cos(c + dx)}} + \frac{3 \sin(c + dx)}{8bd \cos^{\frac{3}{2}}(c + dx)\sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.57

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{3\arctanh(\sin(c+dx))\cos^4(c+dx) + (2+3\cos^2(c+dx))\sin(c+dx)}{8d\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{3/2}}$$

[In] Integrate[1/(Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^(3/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/ (8*d*Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(3/2))

Maple [A] (verified)

Time = 2.94 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.91

method	result
default	$-\frac{3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)-1)-3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)+1)-3(\cos^2(dx+c))\sin(dx+c)-2\sin(dx+c)}{8db\sqrt{\cos(dx+c)}b\cos(dx+c)^{\frac{7}{2}}}$
risch	$-\frac{i(3e^{6i(dx+c)}+11e^{4i(dx+c)}-11e^{2i(dx+c)}-3)}{8b\sqrt{\cos(dx+c)}b\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)^3d} + \frac{3(\sqrt{\cos(dx+c)}\ln(e^{i(dx+c)}+i))}{8b\sqrt{\cos(dx+c)}bd} - \frac{3(\sqrt{\cos(dx+c)}\ln(e^{i(dx+c)}-i))}{8b\sqrt{\cos(dx+c)}bd}$

[In] int(1/cos(d*x+c)^(7/2)/(cos(d*x+c)*b)^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/8/d*(3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)-3*cos(d*x+c)^2*sin(d*x+c)-2*sin(d*x+c))/b/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.01

$$\int \frac{1}{\cos^{\frac{7}{2}}(c+dx)(b\cos(c+dx))^{3/2}} dx = \frac{\left[\frac{3\sqrt{b}\cos(dx+c)^5 \log\left(-\frac{b\cos(dx+c)^3-2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c)}{\cos(dx+c)^3}\right)}{16} + \frac{3\sqrt{-b}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)\cos(dx+c)^5 - \sqrt{b\cos(dx+c)}(3\cos(dx+c)^2+2)\sqrt{\cos(dx+c)}}{8b^2d\cos(dx+c)^5} \right]}{16}$$

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*si

$n(dx + c)/(b^2 d \cos(dx + c)^5)$, $-1/8*(3*\sqrt{-b}*\arctan(\sqrt{b*\cos(dx + c)})*\sqrt{-b}*\sin(dx + c)/(b*\sqrt{\cos(dx + c)}))*\cos(dx + c)^5 - \sqrt{b*\cos(dx + c)}*(3*\cos(dx + c)^2 + 2)*\sqrt{\cos(dx + c)}*\sin(dx + c)/(b^2*d*\cos(dx + c)^5]$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^{3/2}} dx = \text{Timed out}$$

[In] integrate(1/cos(dx+c)**(7/2)/(b*cos(dx+c))**(3/2), x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1679 vs. 2(98) = 196.

Time = 0.42 (sec) , antiderivative size = 1679, normalized size of antiderivative = 14.47

$$\int \frac{1}{\cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(1/cos(dx+c)^(7/2)/(b*cos(dx+c))^(3/2), x, algorithm="maxima")

[Out] $-1/16*(12*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(7/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) + 44*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(5/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 44*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(3/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 12*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 3*(2*(4*\cos(6dx + 6c) + 6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos(8dx + 8c) + \cos(8dx + 8c)^2 + 8*(6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos(6dx + 6c) + 16*\cos(6dx + 6c)^2 + 12*(4*\cos(2dx + 2c) + 1)*\cos(4dx + 4c) + 36*\cos(4dx + 4c)^2 + 16*\cos(2dx + 2c)^2 + 4*(2*\sin(6dx + 6c) + 3*\sin(4dx + 4c) + 2*\sin(2dx + 2c))*\sin(8dx + 8c) + \sin(8dx + 8c)^2 + 16*(3*\sin(4dx + 4c) + 2*\sin(2dx + 2c))*\sin(6dx + 6c) + 16*\sin(6dx + 6c)^2 + 36*\sin(4dx + 4c)^2 + 48*\sin(4dx + 4c)*\sin(2dx + 2c) + 16*\sin(2dx + 2c)^2 + 8*\cos(2dx + 2c) + 1)*\log(\cos(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c)))^2 + \sin(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c)))^2 + 2*\sin(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) + 1) + 3*(2*(4*\cos(6dx + 6c) + 6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos($

$8*d*x + 8*c) + \cos(8*d*x + 8*c)^2 + 8*(6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\cos(6*d*x + 6*c) + 16*\cos(6*d*x + 6*c)^2 + 12*(4*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + 36*\cos(4*d*x + 4*c)^2 + 16*\cos(2*d*x + 2*c)^2 + 4*(2*\sin(6*d*x + 6*c) + 3*\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\sin(8*d*x + 8*c) + \sin(8*d*x + 8*c)^2 + 16*(3*\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\sin(6*d*x + 6*c) + 16*\sin(6*d*x + 6*c)^2 + 36*\sin(4*d*x + 4*c)^2 + 48*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 16*\sin(2*d*x + 2*c)^2 + 8*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))) + 1) - 12*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(7/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 44*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(5/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 44*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 12*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))))/((b*\cos(8*d*x + 8*c)^2 + 16*b*\cos(6*d*x + 6*c)^2 + 36*b*\cos(4*d*x + 4*c)^2 + 16*b*\cos(2*d*x + 2*c)^2 + b*\sin(8*d*x + 8*c)^2 + 16*b*\sin(6*d*x + 6*c)^2 + 36*b*\sin(4*d*x + 4*c)^2 + 48*b*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 16*b*\sin(2*d*x + 2*c)^2 + 2*(4*b*\cos(6*d*x + 6*c) + 6*b*\cos(4*d*x + 4*c) + 4*b*\cos(2*d*x + 2*c) + b)*\cos(8*d*x + 8*c) + 8*(6*b*\cos(4*d*x + 4*c) + 4*b*\cos(2*d*x + 2*c) + b)*\cos(6*d*x + 6*c) + 12*(4*b*\cos(2*d*x + 2*c) + b)*\cos(4*d*x + 4*c) + 8*b*\cos(2*d*x + 2*c) + 4*(2*b*\sin(6*d*x + 6*c) + 3*b*\sin(4*d*x + 4*c) + 2*b*\sin(2*d*x + 2*c))*\sin(8*d*x + 8*c) + 16*(3*b*\sin(4*d*x + 4*c) + 2*b*\sin(2*d*x + 2*c))*\sin(6*d*x + 6*c) + b)*\sqrt{b}*d)$

Giac [F]

$$\int \frac{1}{\cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{3}{2}} \cos(dx + c)^{\frac{7}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(3/2)*cos(d*x + c)^(7/2)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^{3/2}} dx = \int \frac{1}{\cos(c + dx)^{7/2} (b \cos(c + dx))^{3/2}} dx$$

```
[In] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(3/2)), x)
```

```
[Out] int(1/(cos(c + d*x)^(7/2)*(b*cos(c + d*x))^(3/2)), x)
```


$$3.192 \quad \int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1041
Rubi [A] (verified)	1041
Mathematica [A] (verified)	1042
Maple [A] (verified)	1043
Fricas [A] (verification not implemented)	1043
Sympy [F(-1)]	1043
Maxima [A] (verification not implemented)	1044
Giac [F]	1044
Mupad [B] (verification not implemented)	1044

Optimal result

Integrand size = 23, antiderivative size = 107

$$\int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{3x \sqrt{\cos(c+dx)}}{8b^2 \sqrt{b \cos(c+dx)}} + \frac{3 \cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{8b^2 d \sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{7}{2}}(c+dx) \sin(c+dx)}{4b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $3/8*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+1/4*\cos(d*x+c)^{(7/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+3/8*x*\cos(d*x+c)^{(1/2)}/b^2/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{3x \sqrt{\cos(c+dx)}}{8b^2 \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{7}{2}}(c+dx)}{4b^2 d \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{8b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(13/2)}/(b*\text{Cos}[c + d*x]^{(5/2)}, x]$

[Out] $(3*x*\text{Sqrt}[\text{Cos}[c + d*x]])/(8*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (3*\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(8*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(7/2)}*\text{Sin}[c + d*x])/(4*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[a^(m + 1/2)*b^(n - 1/2)*(Sqrt[b*v]/Sqrt[a*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 2715

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^4(c + dx) dx}{b^2 \sqrt{b \cos(c + dx)}} \\
 &= \frac{\cos^{\frac{7}{2}}(c + dx) \sin(c + dx)}{4b^2 d \sqrt{b \cos(c + dx)}} + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int \cos^2(c + dx) dx}{4b^2 \sqrt{b \cos(c + dx)}} \\
 &= \frac{3 \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{8b^2 d \sqrt{b \cos(c + dx)}} + \frac{\cos^{\frac{7}{2}}(c + dx) \sin(c + dx)}{4b^2 d \sqrt{b \cos(c + dx)}} + \frac{\left(3\sqrt{\cos(c + dx)}\right) \int 1 dx}{8b^2 \sqrt{b \cos(c + dx)}} \\
 &= \frac{3x \sqrt{\cos(c + dx)}}{8b^2 \sqrt{b \cos(c + dx)}} + \frac{3 \cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{8b^2 d \sqrt{b \cos(c + dx)}} + \frac{\cos^{\frac{7}{2}}(c + dx) \sin(c + dx)}{4b^2 d \sqrt{b \cos(c + dx)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.54

$$\int \frac{\cos^{\frac{13}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)}(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))}{32b^2 d \sqrt{b \cos(c + dx)}}$$

`[In] Integrate[Cos[c + d*x]^(13/2)/(b*Cos[c + d*x])^(5/2), x]`

`[Out] (Sqrt[Cos[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)]))/(32*b^2*d*Sqrt[b*Cos[c + d*x]])`

Maple [A] (verified)

Time = 3.22 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.61

method	result	size
default	$\frac{(\sqrt{\cos(dx+c)})(2\sin(dx+c)(\cos^3(dx+c))+3\cos(dx+c)\sin(dx+c)+3dx+3c)}{8db^2\sqrt{\cos(dx+c)b}}$	65
risch	$\frac{3x(\sqrt{\cos(dx+c)})}{8b^2\sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)})\sin(4dx+4c)}{32b^2\sqrt{\cos(dx+c)b}d} + \frac{(\sqrt{\cos(dx+c)})\sin(2dx+2c)}{4b^2\sqrt{\cos(dx+c)b}d}$	96

[In] `int(cos(d*x+c)^(13/2)/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)`[Out]
$$\frac{1}{8} \frac{d \cos(dx+c)^{1/2} (2 \sin(dx+c) \cos(dx+c)^3 + 3 \cos(dx+c) \sin(dx+c) + 3 dx + 3c)}{b^2 (\cos(dx+c) b)^{1/2}}$$
Fricas [A] (verification not implemented)

none

Time = 0.36 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.70

$$\int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \left[\frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b} \log \left(\frac{2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b}}{16 b^3 d} \right)}{16 b^3 d} \right]$$

[In] `integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`[Out]
$$\left[\frac{1}{16} (2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b} \log(2 \sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) - 3 \sqrt{-b})) / (b^3 d), \frac{1}{8} (\sqrt{b \cos(dx+c)} (2 \cos(dx+c)^2 + 3) \sqrt{\cos(dx+c)} \sin(dx+c) + 3 \sqrt{b} \arctan(\sqrt{b \cos(dx+c)} \sin(dx+c) / (\sqrt{b} \cos(dx+c)^{3/2})) / (b^3 d)) \right]$$
Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{13}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(13/2)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.46

$$\int \frac{\cos^{\frac{13}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin\left(\frac{1}{2} \arctan(\sin(4 dx + 4 c), \cos(4 dx + 4 c))\right)}{32 b^{\frac{5}{2}} d}$$

```
[In] integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")
```

```
[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(5/2)*d)
```

Giac [F]

$$\int \frac{\cos^{\frac{13}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^{\frac{13}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

```
[In] integrate(cos(d*x+c)^(13/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")
```

```
[Out] integrate(cos(d*x + c)^(13/2)/(b*cos(d*x + c))^(5/2), x)
```

Mupad [B] (verification not implemented)

Time = 15.22 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.73

$$\int \frac{\cos^{\frac{13}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (8 \sin(c + dx) + 9 \sin(3c + 3dx) + \sin(5c + 5dx))}{32 b^3 d (\cos(2c + 2dx) + 1)}$$

```
[In] int(cos(c + d*x)^(13/2)/(b*cos(c + d*x))^(5/2),x)
```

```
[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(8*sin(c + d*x) + 9*sin(3*c + 3*d*x) + sin(5*c + 5*d*x) + 24*d*x*cos(c + d*x)))/(32*b^3*d*(cos(2*c + 2*d*x) + 1))
```

$$3.193 \quad \int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1045
Rubi [A] (verified)	1045
Mathematica [A] (verified)	1046
Maple [A] (verified)	1046
Fricas [A] (verification not implemented)	1047
Sympy [F(-1)]	1047
Maxima [A] (verification not implemented)	1047
Giac [F]	1048
Mupad [B] (verification not implemented)	1048

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sqrt{\cos(c+dx)} \sin^3(c+dx)}{3b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2713}

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sin(c+dx) \sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}} - \frac{\sin^3(c+dx) \sqrt{\cos(c+dx)}}{3b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(11/2)}/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) - (\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[a^{(m+1/2)}*b^{(n-1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2713

```
Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x]
&& IGtQ[(n - 1)/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^3(c + dx) dx}{b^2 \sqrt{b \cos(c + dx)}} \\ &= -\frac{\sqrt{\cos(c + dx)} \text{Subst}(\int (1 - x^2) dx, x, -\sin(c + dx))}{b^2 d \sqrt{b \cos(c + dx)}} \\ &= \frac{\sqrt{\cos(c + dx)} \sin(c + dx)}{b^2 d \sqrt{b \cos(c + dx)}} - \frac{\sqrt{\cos(c + dx)} \sin^3(c + dx)}{3b^2 d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.63

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{5}{2}}} dx = \frac{\sqrt{\cos(c + dx)}(5 + \cos(2(c + dx))) \sin(c + dx)}{6b^2 d \sqrt{b \cos(c + dx)}}$$

```
[In] Integrate[Cos[c + d*x]^(11/2)/(b*Cos[c + d*x])^(5/2), x]
```

```
[Out] (Sqrt[Cos[c + d*x]]*(5 + Cos[2*(c + d*x)])*Sin[c + d*x])/(6*b^2*d*Sqrt[b*Cos[c + d*x]])
```

Maple [A] (verified)

Time = 2.76 (sec) , antiderivative size = 43, normalized size of antiderivative = 0.57

method	result	size
default	$\frac{(2 + \cos^2(dx + c)) \sin(dx + c) (\sqrt{\cos(dx + c)})}{3d b^2 \sqrt{\cos(dx + c)} b}$	43
risch	$\frac{3 \sin(dx + c) (\sqrt{\cos(dx + c)})}{4b^2 d \sqrt{\cos(dx + c)} b} + \frac{(\sqrt{\cos(dx + c)}) \sin(3dx + 3c)}{12b^2 \sqrt{\cos(dx + c)} b d}$	69

```
[In] int(cos(d*x+c)^(11/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/3/d*(2+cos(d*x+c)^2)*sin(d*x+c)*cos(d*x+c)^(1/2)/b^2/(cos(d*x+c)*b)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{\sqrt{b\cos(dx+c)}(\cos(dx+c)^2+2)\sin(dx+c)}{3b^3d\sqrt{\cos(dx+c)}}$$

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*sqrt(b*cos(d*x + c))*(cos(d*x + c)^2 + 2)*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(11/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.55

$$\int \frac{\cos^{\frac{11}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{\sin(3dx+3c) + 9\sin\left(\frac{1}{3}\arctan\left(\frac{\sin(3dx+3c)}{\cos(3dx+3c)}\right)\right)}{12b^{\frac{5}{2}}d}$$

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(5/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^{\frac{11}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^(11/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(11/2)/(b*cos(d*x + c))^(5/2), x)

Mupad [B] (verification not implemented)

Time = 0.68 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.79

$$\int \frac{\cos^{\frac{11}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (10 \sin(2c + 2dx) + \sin(4c + 4dx))}{12b^3 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(11/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(10*sin(2*c + 2*d*x) + sin(4*c + 4*d*x)))/(12*b^3*d*(cos(2*c + 2*d*x) + 1))

$$3.194 \quad \int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1049
Rubi [A] (verified)	1049
Mathematica [A] (verified)	1050
Maple [A] (verified)	1050
Fricas [A] (verification not implemented)	1051
Sympy [F(-1)]	1051
Maxima [A] (verification not implemented)	1051
Giac [F]	1052
Mupad [B] (verification not implemented)	1052

Optimal result

Integrand size = 23, antiderivative size = 69

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{x \sqrt{\cos(c+dx)}}{2b^2 \sqrt{b \cos(c+dx)}} + \frac{\cos^{\frac{3}{2}}(c+dx) \sin(c+dx)}{2b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\cos(d*x+c)^{(3/2)}*\sin(d*x+c)/b^2/d/(b*\cos(d*x+c))^{(1/2)}+1/2*x*\cos(d*x+c)^{(1/2)}/b^2/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2715, 8}

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{x \sqrt{\cos(c+dx)}}{2b^2 \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx) \cos^{\frac{3}{2}}(c+dx)}{2b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(9/2)}/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(x*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*b^2*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + (\text{Cos}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(2*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \text{ :> Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] \text{ /; FreeQ}\{[a, b$

, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2715

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \cos^2(c + dx) dx}{b^2 \sqrt{b \cos(c + dx)}} \\ &= \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2b^2 d \sqrt{b \cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \int 1 dx}{2b^2 \sqrt{b \cos(c + dx)}} \\ &= \frac{x \sqrt{\cos(c + dx)}}{2b^2 \sqrt{b \cos(c + dx)}} + \frac{\cos^{\frac{3}{2}}(c + dx) \sin(c + dx)}{2b^2 d \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.70

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)}(2(c + dx) + \sin(2(c + dx)))}{4b^2 d \sqrt{b \cos(c + dx)}}$$

[In] Integrate[Cos[c + d*x]^(9/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.92 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.65

method	result	size
default	$\frac{(\sqrt{\cos(dx+c)})(\cos(dx+c) \sin(dx+c)+dx+c)}{2d b^2 \sqrt{\cos(dx+c)b}}$	45
risch	$\frac{x(\sqrt{\cos(dx+c)})}{2b^2 \sqrt{\cos(dx+c)b}} + \frac{(\sqrt{\cos(dx+c)}) \sin(2dx+2c)}{4b^2 \sqrt{\cos(dx+c)b} d}$	61

[In] int(cos(d*x+c)^(9/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] $1/2/d*\cos(d*x+c)^{(1/2)}*(\cos(d*x+c)*\sin(d*x+c)+d*x+c)/b^2/(\cos(d*x+c)*b)^{(1/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 157, normalized size of antiderivative = 2.28

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \left[\frac{2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) - \sqrt{-b}\log\left(2b\cos(dx+c)^2 + 2\sqrt{b\cos(dx+c)}\sqrt{\cos(dx+c)}\sin(dx+c) - b\right)}{4b^3d} \right]$$

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out] `[1/4*(2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - sqrt(-b)*log(2*b*cos(d*x + c)^2 + 2*sqrt(b*cos(d*x + c))*sqrt(-b)*sqrt(cos(d*x + c))*sin(d*x + c) - b))/(b^3*d), 1/2*(sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) + sqrt(b)*arctan(sqrt(b*cos(d*x + c))*sin(d*x + c)/(sqrt(b)*cos(d*x + c)^(3/2)))/(b^3*d)]`

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(9/2)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.36

$$\int \frac{\cos^{\frac{9}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{2dx + 2c + \sin(2dx + 2c)}{4b^{\frac{5}{2}}d}$$

[In] `integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(5/2)*d)`

Giac [F]

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^{\frac{9}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^(9/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(9/2)/(b*cos(d*x + c))^(5/2), x)

Mupad [B] (verification not implemented)

Time = 14.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.94

$$\int \frac{\cos^{\frac{9}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)} \sqrt{b \cos(c + dx)} (\sin(c + dx) + \sin(3c + 3dx) + 4dx \cos(c + dx))}{4b^3 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(9/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(sin(c + d*x) + sin(3*c + 3*d*x) + 4*d*x*cos(c + d*x)))/(4*b^3*d*(cos(2*c + 2*d*x) + 1))

$$3.195 \quad \int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1053
Rubi [A] (verified)	1053
Mathematica [A] (verified)	1054
Maple [A] (verified)	1054
Fricas [A] (verification not implemented)	1054
Sympy [F(-1)]	1055
Maxima [A] (verification not implemented)	1055
Giac [F]	1055
Mupad [B] (verification not implemented)	1055

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2717}

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sin(c+dx) \sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(7/2)}/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_*))^{(m)}*((b_*)*(v_*))^{(n)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x \ \&\amp; \ !\text{IntegerQ}[m] \ \&\amp; \ \text{IGtQ}[n + 1/2, 0] \ \&\amp; \ \text{IntegerQ}[m + n]$

Rule 2717

$\text{Int}[\sin[\text{Pi}/2 + (c_*) + (d_*)*(x_*)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \cos(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{\cos(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(7/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (Sqrt[Cos[c + d*x]]*Sin[c + d*x])/(b^2*d*Sqrt[b*Cos[c + d*x]])

Maple [A] (verified)

Time = 2.87 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{b^2 d \sqrt{\cos(dx+c)b}}$	32
risch	$\frac{\sin(dx+c)(\sqrt{\cos(dx+c)})}{b^2 d \sqrt{\cos(dx+c)b}}$	32

[In] int(cos(d*x+c)^(7/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] sin(d*x+c)*cos(d*x+c)^(1/2)/b^2/d/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{\cos^{\frac{7}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{b^3 d \sqrt{\cos(dx+c)}}$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(7/2)/(b*cos(d*x+c))**(5/2), x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.37

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sin(dx + c)}{b^{\frac{5}{2}} d}$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2), x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(5/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^{\frac{7}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(7/2)/(b*cos(d*x + c))^(5/2), x)

Mupad [B] (verification not implemented)

Time = 13.90 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.34

$$\int \frac{\cos^{\frac{7}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{\cos(c + dx)} \sin(2c + 2dx) \sqrt{b \cos(c + dx)}}{b^3 d (\cos(2c + 2dx) + 1)}$$

[In] int(cos(c + d*x)^(7/2)/(b*cos(c + d*x))^(5/2), x)

[Out] (cos(c + d*x)^(1/2)*sin(2*c + 2*d*x)*(b*cos(c + d*x))^(1/2))/(b^3*d*(cos(2*c + 2*d*x) + 1))

$$3.196 \quad \int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Optimal result	1056
Rubi [A] (verified)	1056
Mathematica [A] (verified)	1057
Maple [A] (verified)	1057
Fricas [A] (verification not implemented)	1057
Sympy [F(-1)]	1058
Maxima [A] (verification not implemented)	1058
Giac [F]	1058
Mupad [B] (verification not implemented)	1059

Optimal result

Integrand size = 23, antiderivative size = 27

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{x \sqrt{\cos(c+dx)}}{b^2 \sqrt{b \cos(c+dx)}}$$

[Out] $x \cos(dx+c)^{(1/2)} / b^2 / (b \cos(dx+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{x \sqrt{\cos(c+dx)}}{b^2 \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^{(5/2)} / (b * \text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(x * \text{Sqrt}[\text{Cos}[c + d*x]]) / (b^2 * \text{Sqrt}[b * \text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_)*((a_)*(v_))^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)} * b^{(n - 1/2)} * (\text{Sqrt}[b*v] / \text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n + 1/2, 0] \&\& \text{IntegerQ}[m + n]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int 1 dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{x \sqrt{\cos(c+dx)}}{b^2 \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.00

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{x \sqrt{b \cos(c+dx)}}{b^3 \sqrt{\cos(c+dx)}}$$

[In] Integrate[Cos[c + d*x]^(5/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (x*Sqrt[b*Cos[c + d*x]])/(b^3*Sqrt[Cos[c + d*x]])

Maple [A] (verified)

Time = 2.25 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.89

method	result	size
risch	$\frac{x(\sqrt{\cos(dx+c)})}{b^2 \sqrt{\cos(dx+c)b}}$	24
default	$\frac{(\sqrt{\cos(dx+c)})(dx+c)}{d b^2 \sqrt{\cos(dx+c)b}}$	31

[In] int(cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] x*cos(d*x+c)^(1/2)/b^2/(cos(d*x+c)*b)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.59

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \left[-\frac{\sqrt{-b} \log \left(2 b \cos(dx+c)^2 + 2 \sqrt{b \cos(dx+c)} \sqrt{-b} \sqrt{\cos(dx+c)} \sin(dx+c) \right)}{2 b^3 d} \right]$$

[In] integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2), x, algorithm="fricas")

[Out] $[-1/2\sqrt{-b}\log(2b\cos(dx+c)^2 + 2\sqrt{b\cos(dx+c)}\sqrt{-b}\sqrt{t(\cos(dx+c))\sin(dx+c) - b})/(b^3d), \arctan(\sqrt{b\cos(dx+c)}\sin(dx+c)/(\sqrt{b}\cos(dx+c)^{3/2}))]/(b^{5/2}d)]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**(5/2)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{5}{2}}d}$$

[In] `integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] $2*\arctan(\sin(dx+c)/(\cos(dx+c)+1))/(b^{5/2}*d)$

Giac [F]

$$\int \frac{\cos^{\frac{5}{2}}(c+dx)}{(b\cos(c+dx))^{\frac{5}{2}}} dx = \int \frac{\cos(dx+c)^{\frac{5}{2}}}{(b\cos(dx+c))^{\frac{5}{2}}} dx$$

[In] `integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")`

[Out] `integrate(cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x)`

Mupad [B] (verification not implemented)

Time = 0.30 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.37

$$\int \frac{\cos^{\frac{5}{2}}(c + dx)}{(b \cos(c + dx))^{\frac{5}{2}}} dx = \frac{2x \cos(c + dx)^{\frac{3}{2}} \sqrt{b \cos(c + dx)}}{b^3 (\cos(2c + 2dx) + 1)}$$

[In] `int(cos(c + d*x)^(5/2)/(b*cos(c + d*x))^(5/2),x)`

[Out] `(2*x*cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(1/2))/(b^3*(cos(2*c + 2*d*x) + 1))`

$$3.197 \quad \int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

Optimal result	1060
Rubi [A] (verified)	1060
Mathematica [A] (verified)	1061
Maple [A] (verified)	1061
Fricas [A] (verification not implemented)	1061
Sympy [F(-1)]	1062
Maxima [B] (verification not implemented)	1062
Giac [F]	1062
Mupad [F(-1)]	1063

Optimal result

Integrand size = 23, antiderivative size = 36

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}}$$

[Out] $\operatorname{arctanh}(\sin(dx+c)) \cdot \cos(dx+c)^{(1/2)} / b^2/d / (b \cdot \cos(dx+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3855}

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{\frac{5}{2}}} dx = \frac{\sqrt{\cos(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{b^2 d \sqrt{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^{(3/2)} / (b * \operatorname{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $(\operatorname{ArcTanh}[\operatorname{Sin}[c + d*x]] * \operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]) / (b^2 * d * \operatorname{Sqrt}[b * \operatorname{Cos}[c + d*x]])$

Rule 17

$\operatorname{Int}[(u_.) * ((a_.) * (v_))^{(m_)} * ((b_.) * (v_))^{(n_)}, x_Symbol] \rightarrow \operatorname{Dist}[a^{(m + 1/2)} * b^{(n - 1/2)} * (\operatorname{Sqrt}[b*v] / \operatorname{Sqrt}[a*v]), \operatorname{Int}[u*v^{(m + n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, m\}, x$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{IGtQ}[n + 1/2, 0]$ && $\operatorname{IntegerQ}[m + n]$

Rule 3855

$\operatorname{Int}[\operatorname{csc}[(c_.) + (d_.) * (x_)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]] / d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{b^2 d \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.92

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx)) \cos^{\frac{5}{2}}(c+dx)}{d(b \cos(c+dx))^{5/2}}$$

[In] Integrate[Cos[c + d*x]^(3/2)/(b*Cos[c + d*x])^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^(5/2))/(d*(b*Cos[c + d*x])^(5/2))

Maple [A] (verified)

Time = 2.76 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.19

method	result	size
default	$-\frac{2(\sqrt{\cos(dx+c)} \operatorname{arctanh}(\cot(dx+c)-\csc(dx+c)))}{d \sqrt{\cos(dx+c)} b b^2}$	43
risch	$-\frac{(\sqrt{\cos(dx+c)} \ln(e^{i(dx+c)}-i))}{b^2 \sqrt{\cos(dx+c)} b d} + \frac{(\sqrt{\cos(dx+c)} \ln(e^{i(dx+c)}+i))}{b^2 \sqrt{\cos(dx+c)} b d}$	79

[In] int(cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)

[Out] $-2/d \cos(d*x+c)^{(1/2)} * \operatorname{arctanh}(\cot(d*x+c) - \csc(d*x+c)) / (\cos(d*x+c) * b)^{(1/2)} / b^2$

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 116, normalized size of antiderivative = 3.22

$$\int \frac{\cos^{\frac{3}{2}}(c+dx)}{(b \cos(c+dx))^{5/2}} dx = \left[\frac{\log \left(-\frac{b \cos(dx+c)^3 - 2 \sqrt{b \cos(dx+c)} \sqrt{b} \sqrt{\cos(dx+c)} \sin(dx+c) - 2 b \cos(dx+c)}{\cos(dx+c)^3} \right)}{2 b^{\frac{5}{2}} d}, \right. \\ \left. - \frac{\sqrt{-b} \arctan \left(\frac{\sqrt{b \cos(dx+c)} \sqrt{-b} \sin(dx+c)}{b \sqrt{\cos(dx+c)}} \right)}{b^3 d} \right]$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3/(b^(5/2)*d), -sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))/(b^3*d)]

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(32) = 64.

Time = 0.38 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.81

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \frac{\log(\cos(dx + c)^2 + \sin(dx + c)^2 + 2 \sin(dx + c) + 1) - \log(\cos(dx + c)^2 + \sin(dx + c)^2)}{2 b^{\frac{5}{2}} d}$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/2*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/(b^(5/2)*d)

Giac [F]

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(dx + c)^{\frac{3}{2}}}{(b \cos(dx + c))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^(3/2)/(b*cos(d*x + c))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^{\frac{3}{2}}(c + dx)}{(b \cos(c + dx))^{5/2}} dx = \int \frac{\cos(c + dx)^{3/2}}{(b \cos(c + dx))^{5/2}} dx$$

```
[In] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(5/2), x)
```

```
[Out] int(cos(c + d*x)^(3/2)/(b*cos(c + d*x))^(5/2), x)
```

$$3.198 \quad \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1064
Rubi [A] (verified)	1064
Mathematica [A] (verified)	1065
Maple [A] (verified)	1065
Fricas [A] (verification not implemented)	1066
Sympy [F]	1066
Maxima [B] (verification not implemented)	1066
Giac [F]	1067
Mupad [B] (verification not implemented)	1067

Optimal result

Integrand size = 23, antiderivative size = 35

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[Out] $\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3852, 8}

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[\text{Sqrt}[\text{Cos}[c + d*x]]/(b*\text{Cos}[c + d*x])^{(5/2)}, x]$

[Out] $\text{Sin}[c + d*x]/(b^2*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)*((b_.)*(v_))^{(n_)}], x_Symbol] \rightarrow \text{Dist}[a^{(m + 1/2)}*b^{(n - 1/2)}*(\text{Sqrt}[b*v]/\text{Sqrt}[a*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& !\text{IntegerQ}[m] \&\& \text{IGtQ}[n + 1/2, 0] \&\& \text{IntegerQ}[m + n]$

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^2(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{b^2 d \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{\cos^{3/2}(c+dx) \sin(c+dx)}{d(b \cos(c+dx))^{5/2}}$$

```
[In] Integrate[Sqrt[Cos[c + d*x]]/(b*Cos[c + d*x])^(5/2), x]
```

```
[Out] (Cos[c + d*x]^(3/2)*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(5/2))
```

Maple [A] (verified)

Time = 3.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
default	$\frac{\sin(dx+c)}{b^2 d \sqrt{\cos(dx+c)} \sqrt{\cos(dx+c)b}}$	32
risch	$\frac{2i(\sqrt{\cos(dx+c)})}{b^2 \sqrt{\cos(dx+c)b} d(e^{2i(dx+c)}+1)}$	41

```
[In] int(cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(5/2), x, method=_RETURNVERBOSE)
```

```
[Out] sin(d*x+c)/b^2/d/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{b \cos(dx+c)} \sin(dx+c)}{b^3 d \cos(dx+c)^{\frac{3}{2}}}$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] sqrt(b*cos(d*x + c))*sin(d*x + c)/(b^3*d*cos(d*x + c)^(3/2))

Sympy [F]

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{\frac{5}{2}}} dx$$

[In] integrate(cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Integral(sqrt(cos(c + d*x))/(b*cos(c + d*x))**(5/2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 67 vs. 2(31) = 62.

Time = 0.38 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.91

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sqrt{b} \sin(2 dx + 2 c)}{(b^3 \cos(2 dx + 2 c))^2 + b^3 \sin(2 dx + 2 c)^2 + 2 b^3 \cos(2 dx + 2 c) + b^3} d$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*sin(2*d*x + 2*c)/((b^3*cos(2*d*x + 2*c)^2 + b^3*sin(2*d*x + 2*c)^2 + 2*b^3*cos(2*d*x + 2*c) + b^3)*d)

Giac [F]

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \int \frac{\sqrt{\cos(dx+c)}}{(b \cos(dx+c))^{5/2}} dx$$

[In] integrate(cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(cos(d*x + c))/(b*cos(d*x + c))^(5/2), x)

Mupad [B] (verification not implemented)

Time = 14.09 (sec) , antiderivative size = 87, normalized size of antiderivative = 2.49

$$\int \frac{\sqrt{\cos(c+dx)}}{(b \cos(c+dx))^{5/2}} dx = \frac{2 \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)} (\sin(c+dx) + \sin(3c+3dx) + \cos(c+dx))}{b^3 d (4 \cos(2c+2dx) + \cos(4c+4dx) + 3)}$$

[In] int(cos(c + d*x)^(1/2)/(b*cos(c + d*x))^(5/2),x)

[Out] (2*cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(1/2)*(cos(c + d*x)*3i + sin(c + d*x) + cos(3*c + 3*d*x)*1i + sin(3*c + 3*d*x)))/(b^3*d*(4*cos(2*c + 2*d*x) + cos(4*c + 4*d*x) + 3))

$$3.199 \quad \int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1068
Rubi [A] (verified)	1068
Mathematica [A] (verified)	1069
Maple [A] (verified)	1070
Fricas [A] (verification not implemented)	1070
Sympy [F(-1)]	1071
Maxima [B] (verification not implemented)	1071
Giac [F]	1072
Mupad [F(-1)]	1072

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2}} dx = \frac{\operatorname{arctanh}(\sin(c+dx))\sqrt{\cos(c+dx)}}{2b^2d\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2b^2d \cos^{3/2}(c+dx)\sqrt{b \cos(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(3/2)}/(b*\cos(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\cos(d*x+c)^{(1/2)}/b^2/d/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b \cos(c+dx))^{5/2}} dx = \frac{\sqrt{\cos(c+dx)}\operatorname{arctanh}(\sin(c+dx))}{2b^2d\sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{2b^2d \cos^{3/2}(c+dx)\sqrt{b \cos(c+dx)}}$$

[In] $\text{Int}[1/(\text{Sqrt}[\text{Cos}[c + d*x]]*(b*\text{Cos}[c + d*x])^{(5/2)}), x]$

[Out] $(\text{ArcTanh}[\text{Sin}[c + d*x]]*\text{Sqrt}[\text{Cos}[c + d*x]])/(2*b^2*d*\text{Sqrt}[b*\text{Cos}[c + d*x]]) + \text{Sin}[c + d*x]/(2*b^2*d*\text{Cos}[c + d*x]^{(3/2)}*\text{Sqrt}[b*\text{Cos}[c + d*x]])$

Rule 18

$\text{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a^{(m - 1/2)}*b^{(n + 1/2)}*(\text{Sqrt}[a*v]/\text{Sqrt}[b*v]), \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b$

, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c + dx)} \int \sec^3(c + dx) dx}{b^2 \sqrt{b \cos(c + dx)}} \\ &= \frac{\sin(c + dx)}{2b^2 d \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}} + \frac{\sqrt{\cos(c + dx)} \int \sec(c + dx) dx}{2b^2 \sqrt{b \cos(c + dx)}} \\ &= \frac{\operatorname{arctanh}(\sin(c + dx)) \sqrt{\cos(c + dx)}}{2b^2 d \sqrt{b \cos(c + dx)}} + \frac{\sin(c + dx)}{2b^2 d \cos^{\frac{3}{2}}(c + dx) \sqrt{b \cos(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.71

$$\int \frac{1}{\sqrt{\cos(c + dx)} (b \cos(c + dx))^{5/2}} dx = \frac{\sqrt{b \cos(c + dx)} (\operatorname{arctanh}(\sin(c + dx)) \cos^2(c + dx) + \sin(c + dx))}{2b^3 d \cos^{\frac{5}{2}}(c + dx)}$$

[In] Integrate[1/(Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^(5/2)),x]

[Out] (Sqrt[b*Cos[c + d*x]]*(ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^2 + Sin[c + d*x]))/(2*b^3*d*Cos[c + d*x]^(5/2))

Maple [A] (verified)

Time = 3.19 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.12

method	result	size
default	$\frac{-(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)-1)+(\cos^2(dx+c)) \ln(-\cot(dx+c)+\csc(dx+c)+1)+\sin(dx+c)}{2db^2\sqrt{\cos(dx+c)b}\cos(dx+c)^{\frac{3}{2}}}$	87
risch	$-\frac{i(e^{2i(dx+c)}-1)}{2b^2\sqrt{\cos(dx+c)b}\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)d} + \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}+i)}{2b^2\sqrt{\cos(dx+c)b}d} - \frac{(\sqrt{\cos(dx+c)}) \ln(e^{i(dx+c)}-i)}{2b^2\sqrt{\cos(dx+c)b}d}$	131

[In] int(1/cos(d*x+c)^(1/2)/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/2/d*(-cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)-1)+cos(d*x+c)^2*ln(-cot(d*x+c)+csc(d*x+c)+1)+sin(d*x+c))/b^2/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(3/2)

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 207, normalized size of antiderivative = 2.65

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx = \left[\frac{\sqrt{b}\cos(dx+c)^3 \log\left(-\frac{b\cos(dx+c)^3-2\sqrt{b}\cos(dx+c)\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c)-2}{\cos(dx+c)^3}\right)}{4b^3d\cos(dx+c)^3} - \frac{\sqrt{-b}\arctan\left(\frac{\sqrt{b}\cos(dx+c)\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)\cos(dx+c)^3 - \sqrt{b}\cos(dx+c)\sqrt{\cos(dx+c)}\sin(dx+c)}{2b^3d\cos(dx+c)^3} \right]$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^3*d*cos(d*x + c)^3), -1/2*(sqrt(-b)*arctan(sqrt(b*cos(d*x + c))*sqrt(-b)*sin(d*x + c)/(b*sqrt(cos(d*x + c))))*cos(d*x + c)^3 - sqrt(b*cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c))/(b^3*d*cos(d*x + c)^3)]

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(1/cos(d*x+c)**(1/2)/(b*cos(d*x+c))**(5/2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 688 vs. 2(66) = 132.

Time = 0.45 (sec) , antiderivative size = 688, normalized size of antiderivative = 8.82

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")

```
[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))/(b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*sqrt(b)*d
```

Giac [F]

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx = \int \frac{1}{(b\cos(dx+c))^{\frac{5}{2}}\sqrt{\cos(dx+c)}} dx$$

[In] integrate(1/cos(d*x+c)^(1/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*sqrt(cos(d*x + c))), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx = \int \frac{1}{\sqrt{\cos(c+dx)}(b\cos(c+dx))^{5/2}} dx$$

[In] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2)),x)

[Out] int(1/(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^(5/2)), x)

$$3.200 \quad \int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1073
Rubi [A] (verified)	1073
Mathematica [A] (verified)	1074
Maple [A] (verified)	1074
Fricas [A] (verification not implemented)	1075
Sympy [F(-1)]	1075
Maxima [B] (verification not implemented)	1075
Giac [F]	1076
Mupad [B] (verification not implemented)	1076

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx = \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3b^2 d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[Out] $1/3*\sin(d*x+c)^3/b^2/d/\cos(d*x+c)^{(5/2)}/(b*\cos(d*x+c))^{(1/2)}+\sin(d*x+c)/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\cos(d*x+c))^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 3852}

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx = \frac{\sin^3(c+dx)}{3b^2 d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] Sin[c + d*x]/(b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]^3/(3*b^2*d*Cos[c + d*x]^(5/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[a^(m - 1/2)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b

, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^4(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\ &= -\frac{\sqrt{\cos(c+dx)} \text{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{b^2 d \sqrt{b \cos(c+dx)}} \\ &= \frac{\sin(c+dx)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \cos(c+dx)}} + \frac{\sin^3(c+dx)}{3b^2 d \cos^{\frac{5}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx = \frac{\cos^{\frac{5}{2}}(c+dx) \left(\tan(c+dx) + \frac{1}{3} \tan^3(c+dx)\right)}{d(b \cos(c+dx))^{5/2}}$$

[In] Integrate[1/(Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] (Cos[c + d*x]^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*(b*Cos[c + d*x])^(5/2))

Maple [A] (verified)

Time = 2.88 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.59

method	result	size
default	$\frac{(2(\cos^2(dx+c))+1) \sin(dx+c)}{3db^2 \sqrt{\cos(dx+c)b} \cos(dx+c)^{\frac{5}{2}}}$	45
risch	$\frac{2i(4 \cos(dx+c)+2i \sin(dx+c))}{3b^2 \sqrt{\cos(dx+c)b} \sqrt{\cos(dx+c)} (e^{2i(dx+c)}+1)^2 d}$	59

[In] int(1/cos(d*x+c)^(3/2)/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] $1/3/d*(2*\cos(d*x+c)^2+1)*\sin(d*x+c)/b^2/(\cos(d*x+c)*b)^{(1/2)}/\cos(d*x+c)^{(5/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.58

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{5/2}} dx = \frac{\sqrt{b\cos(dx+c)}(2\cos(dx+c)^2+1)\sin(dx+c)}{3b^3d\cos(dx+c)^{\frac{7}{2}}}$$

[In] `integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")`

[Out] $1/3*\text{sqrt}(b*\cos(d*x + c))*(2*\cos(d*x + c)^2 + 1)*\sin(d*x + c)/(b^3*d*\cos(d*x + c)^{(7/2)})$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{5/2}} dx = \text{Timed out}$$

[In] `integrate(1/cos(d*x+c)**(3/2)/(b*cos(d*x+c))**(5/2),x)`

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 343 vs. $2(66) = 132$.

Time = 0.39 (sec) , antiderivative size = 343, normalized size of antiderivative = 4.51

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{5/2}} dx = \frac{1}{3(b^2\cos(6dx+6c)^2+9b^2\cos(4dx+4c)^2+9b^2\cos(2dx+2c)^2 -$$

[In] `integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] $4/3*((3*\cos(2*d*x + 2*c) + 1)*\sin(6*d*x + 6*c) + 3*(3*\cos(2*d*x + 2*c) + 1)*\sin(4*d*x + 4*c) - 3*\cos(6*d*x + 6*c)*\sin(2*d*x + 2*c) - 9*\cos(4*d*x + 4*c)*\sin(2*d*x + 2*c))/((b^2*\cos(6*d*x + 6*c)^2 + 9*b^2*\cos(4*d*x + 4*c)^2 + 9*b^2*\cos(2*d*x + 2*c)^2 + b^2*\sin(6*d*x + 6*c)^2 + 9*b^2*\sin(4*d*x + 4*c)^2 + 18*b^2*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 9*b^2*\sin(2*d*x + 2*c)^2 + 6*b^2*\cos(2*d*x + 2*c) + b^2 + 2*(3*b^2*\cos(4*d*x + 4*c) + 3*b^2*\cos(2*d*x + 2*c) + b^2)*\cos(6*d*x + 6*c) + 6*(3*b^2*\cos(2*d*x + 2*c) + b^2)*\cos(4*d*x + 4*c) + 6*(b^2*\sin(4*d*x + 4*c) + b^2*\sin(2*d*x + 2*c))*\sin(6*d*x + 6*c))*\text{sqrt}(b)*d$

Giac [F]

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{5/2}} dx = \int \frac{1}{(b\cos(dx+c))^{\frac{5}{2}} \cos(dx+c)^{\frac{3}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*cos(d*x + c)^(3/2)), x)

Mupad [B] (verification not implemented)

Time = 15.13 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.72

$$\int \frac{1}{\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{5/2}} dx = \frac{2\sqrt{b\cos(c+dx)}(\cos(2c+2dx)15i + \cos(4c+4dx)6i + \cos(6c+6dx)1i + 9\sin(2c+2dx) + 6\sin(4c+4dx) + \sin(6c+6dx) + 10i)}{3b^3d\sqrt{\cos(c+dx)}(15\cos(2c+2dx) + 6\cos(4c+4dx) + \cos(6c+6dx) + 10)}$$

[In] int(1/(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^(5/2)),x)

[Out] (2*(b*cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*15i + cos(4*c + 4*d*x)*6i + cos(6*c + 6*d*x)*1i + 9*sin(2*c + 2*d*x) + 6*sin(4*c + 4*d*x) + sin(6*c + 6*d*x) + 10i))/(3*b^3*d*cos(c + d*x)^(1/2)*(15*cos(2*c + 2*d*x) + 6*cos(4*c + 4*d*x) + cos(6*c + 6*d*x) + 10))

$$3.201 \quad \int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx$$

Optimal result	1077
Rubi [A] (verified)	1077
Mathematica [A] (verified)	1079
Maple [A] (verified)	1079
Fricas [A] (verification not implemented)	1079
Sympy [F(-1)]	1080
Maxima [B] (verification not implemented)	1080
Giac [F]	1081
Mupad [F(-1)]	1082

Optimal result

Integrand size = 23, antiderivative size = 116

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx = \frac{3 \operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{8b^2 d \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/b^2/d/cos(d*x+c)^(7/2)/(b*cos(d*x+c))^(1/2)+3/8*sin(d*x+c)/b^2/d/cos(d*x+c)^(3/2)/(b*cos(d*x+c))^(1/2)+3/8*arctanh(sin(d*x+c))*cos(d*x+c)^(1/2)/b^2/d/(b*cos(d*x+c))^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 3853, 3855}

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b \cos(c+dx))^{5/2}} dx = \frac{3 \sqrt{\cos(c+dx)} \operatorname{arctanh}(\sin(c+dx))}{8b^2 d \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}}$$

[In] Int[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[Cos[c + d*x]]/(8*b^2*d*Sqrt[b*Cos[c + d*x]]) + Sin[c + d*x]/(4*b^2*d*Cos[c + d*x]^(7/2)*Sqrt[b*Cos[c + d*x]]) + (3*Sin[c + d*x])/(8*b^2*d*Cos[c + d*x]^(3/2)*Sqrt[b*Cos[c + d*x]])

Rule 18

```
Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[a^(m - 1/2)
)*b^(n + 1/2)*(Sqrt[a*v]/Sqrt[b*v]), Int[u*v^(m + n), x], x] /; FreeQ[{a, b
, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]
```

Rule 3853

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*
x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)),
Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &
& IntegerQ[2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\sqrt{\cos(c+dx)} \int \sec^5(c+dx) dx}{b^2 \sqrt{b \cos(c+dx)}} \\
&= \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int \sec^3(c+dx) dx}{4b^2 \sqrt{b \cos(c+dx)}} \\
&= \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \\
&\quad + \frac{\left(3\sqrt{\cos(c+dx)}\right) \int \sec(c+dx) dx}{8b^2 \sqrt{b \cos(c+dx)}} \\
&= \frac{3 \operatorname{arctanh}(\sin(c+dx)) \sqrt{\cos(c+dx)}}{8b^2 d \sqrt{b \cos(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \cos^{\frac{7}{2}}(c+dx) \sqrt{b \cos(c+dx)}} \\
&\quad + \frac{3 \sin(c+dx)}{8b^2 d \cos^{\frac{3}{2}}(c+dx) \sqrt{b \cos(c+dx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 66, normalized size of antiderivative = 0.57

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{3\operatorname{arctanh}(\sin(c+dx))\cos^4(c+dx) + (2+3\cos^2(c+dx))\sin(c+dx)}{8d\cos^{\frac{3}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}}}$$

[In] Integrate[1/(Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^(5/2)),x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Cos[c + d*x]^4 + (2 + 3*Cos[c + d*x]^2)*Sin[c + d*x])/ (8*d*Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^(5/2))

Maple [A] (verified)

Time = 3.39 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.91

method	result
default	$\frac{-3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)-1)+3(\cos^4(dx+c))\ln(-\cot(dx+c)+\csc(dx+c)+1)+3(\cos^2(dx+c))\sin(dx+c)+2\sin(dx+c)}{8db^2\sqrt{\cos(dx+c)}b\cos(dx+c)^{\frac{7}{2}}}$
risch	$-\frac{i(3e^{6i(dx+c)}+11e^{4i(dx+c)}-11e^{2i(dx+c)}-3)}{8b^2\sqrt{\cos(dx+c)}b\sqrt{\cos(dx+c)}(e^{2i(dx+c)}+1)^3d} - \frac{3(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}-i)}{8b^2\sqrt{\cos(dx+c)}bd} + \frac{3(\sqrt{\cos(dx+c)})\ln(e^{i(dx+c)}+i)}{8b^2\sqrt{\cos(dx+c)}bd}$

[In] int(1/cos(d*x+c)^(5/2)/(cos(d*x+c)*b)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/8/d*(-3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)-1)+3*cos(d*x+c)^4*ln(-cot(d*x+c)+csc(d*x+c)+1)+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))/b^2/(cos(d*x+c)*b)^(1/2)/cos(d*x+c)^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.42 (sec) , antiderivative size = 233, normalized size of antiderivative = 2.01

$$\int \frac{1}{\cos^{\frac{5}{2}}(c+dx)(b\cos(c+dx))^{\frac{5}{2}}} dx = \frac{\left[\frac{3\sqrt{b}\cos(dx+c)^5 \log\left(-\frac{b\cos(dx+c)^3-2\sqrt{b\cos(dx+c)}\sqrt{b}\sqrt{\cos(dx+c)}\sin(dx+c)}{\cos(dx+c)^3}\right)}{16} + \frac{3\sqrt{-b}\arctan\left(\frac{\sqrt{b\cos(dx+c)}\sqrt{-b}\sin(dx+c)}{b\sqrt{\cos(dx+c)}}\right)\cos(dx+c)^5 - \sqrt{b\cos(dx+c)}(3\cos(dx+c)^2+2)\sqrt{\cos(dx+c)}}{8b^3d\cos(dx+c)^5} \right]}{16}$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^5*log(-(b*cos(d*x + c))^3 - 2*sqrt(b*cos(d*x + c))*sqrt(b)*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b*cos(d*x + c))/cos(d*x + c)^3) + 2*sqrt(b*cos(d*x + c))*(3*cos(d*x + c)^2 + 2)*sqrt(cos(d*x + c))*si

$n(dx + c)/(b^3 d \cos(dx + c)^5)$, $-1/8*(3*\sqrt{-b}*\arctan(\sqrt{b*\cos(dx + c)})*\sqrt{-b}*\sin(dx + c)/(b*\sqrt{\cos(dx + c)}))*\cos(dx + c)^5 - \sqrt{b*\cos(dx + c)}*(3*\cos(dx + c)^2 + 2)*\sqrt{\cos(dx + c)}*\sin(dx + c)/(b^3*d*\cos(dx + c)^5]$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Timed out}$$

[In] integrate(1/cos(dx+c)**(5/2)/(b*cos(dx+c))**(5/2), x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1729 vs. 2(98) = 196.

Time = 0.44 (sec) , antiderivative size = 1729, normalized size of antiderivative = 14.91

$$\int \frac{1}{\cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}}} dx = \text{Too large to display}$$

[In] integrate(1/cos(dx+c)^(5/2)/(b*cos(dx+c))^(5/2), x, algorithm="maxima")

[Out] $-1/16*(12*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(7/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) + 44*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(5/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 44*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(3/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 12*(\sin(8dx + 8c) + 4*\sin(6dx + 6c) + 6*\sin(4dx + 4c) + 4*\sin(2dx + 2c))*\cos(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) - 3*(2*(4*\cos(6dx + 6c) + 6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos(8dx + 8c) + \cos(8dx + 8c)^2 + 8*(6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos(6dx + 6c) + 16*\cos(6dx + 6c)^2 + 12*(4*\cos(2dx + 2c) + 1)*\cos(4dx + 4c) + 36*\cos(4dx + 4c)^2 + 16*\cos(2dx + 2c)^2 + 4*(2*\sin(6dx + 6c) + 3*\sin(4dx + 4c) + 2*\sin(2dx + 2c))*\sin(8dx + 8c) + \sin(8dx + 8c)^2 + 16*(3*\sin(4dx + 4c) + 2*\sin(2dx + 2c))*\sin(6dx + 6c) + 16*\sin(6dx + 6c)^2 + 36*\sin(4dx + 4c)^2 + 48*\sin(4dx + 4c)*\sin(2dx + 2c) + 16*\sin(2dx + 2c)^2 + 8*\cos(2dx + 2c) + 1)*\log(\cos(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c)))^2 + \sin(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c)))^2 + 2*\sin(1/2*\arctan2(\sin(2dx + 2c), \cos(2dx + 2c))) + 1) + 3*(2*(4*\cos(6dx + 6c) + 6*\cos(4dx + 4c) + 4*\cos(2dx + 2c) + 1)*\cos($

$8*d*x + 8*c) + \cos(8*d*x + 8*c)^2 + 8*(6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\cos(6*d*x + 6*c) + 16*\cos(6*d*x + 6*c)^2 + 12*(4*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + 36*\cos(4*d*x + 4*c)^2 + 16*\cos(2*d*x + 2*c)^2 + 4*(2*\sin(6*d*x + 6*c) + 3*\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\sin(8*d*x + 8*c) + \sin(8*d*x + 8*c)^2 + 16*(3*\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\sin(6*d*x + 6*c) + 16*\sin(6*d*x + 6*c)^2 + 36*\sin(4*d*x + 4*c)^2 + 48*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 16*\sin(2*d*x + 2*c)^2 + 8*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))) + 1) - 12*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(7/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 44*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(5/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 44*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 12*(\cos(8*d*x + 8*c) + 4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))/((b^2*\cos(8*d*x + 8*c)^2 + 16*b^2*\cos(6*d*x + 6*c)^2 + 36*b^2*\cos(4*d*x + 4*c)^2 + 16*b^2*\cos(2*d*x + 2*c)^2 + b^2*\sin(8*d*x + 8*c)^2 + 16*b^2*\sin(6*d*x + 6*c)^2 + 36*b^2*\sin(4*d*x + 4*c)^2 + 48*b^2*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 16*b^2*\sin(2*d*x + 2*c)^2 + 8*b^2*\cos(2*d*x + 2*c) + b^2 + 2*(4*b^2*\cos(6*d*x + 6*c) + 6*b^2*\cos(4*d*x + 4*c) + 4*b^2*\cos(2*d*x + 2*c) + b^2)*\cos(8*d*x + 8*c) + 8*(6*b^2*\cos(4*d*x + 4*c) + 4*b^2*\cos(2*d*x + 2*c) + b^2)*\cos(6*d*x + 6*c) + 12*(4*b^2*\cos(2*d*x + 2*c) + b^2)*\cos(4*d*x + 4*c) + 4*(2*b^2*\sin(6*d*x + 6*c) + 3*b^2*\sin(4*d*x + 4*c) + 2*b^2*\sin(2*d*x + 2*c))*\sin(8*d*x + 8*c) + 16*(3*b^2*\sin(4*d*x + 4*c) + 2*b^2*\sin(2*d*x + 2*c))*\sin(6*d*x + 6*c))*\sqrt{b}*d$

Giac [F]

$$\int \frac{1}{\cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{5}{2}} \cos(dx + c)^{\frac{5}{2}}} dx$$

[In] integrate(1/cos(d*x+c)^(5/2)/(b*cos(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*cos(d*x + c))^(5/2)*cos(d*x + c)^(5/2)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^{\frac{5}{2}}} dx = \int \frac{1}{\cos(c + dx)^{\frac{5}{2}}(b \cos(c + dx))^{\frac{5}{2}}} dx$$

```
[In] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2)), x)
```

```
[Out] int(1/(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^(5/2)), x)
```

3.202 $\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal result	1083
Rubi [A] (verified)	1083
Mathematica [A] (verified)	1084
Maple [F]	1085
Fricas [F]	1085
Sympy [F]	1085
Maxima [F]	1085
Giac [F]	1086
Mupad [F(-1)]	1086

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \frac{3 \cos^{1+m}(c + dx) \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(4 + 3m), \frac{1}{6}(10 + 3m), \cos^2(c + dx)\right) \sin(c + dx)}{d(4 + 3m) \sqrt{\sin^2(c + dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*(b*\cos(d*x+c))^{(1/3)}*\operatorname{hypergeom}([1/2, 2/3+1/2*m], [5/3+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(4+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+1}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m + 4), \frac{1}{6}(3m + 10), \cos^2(c + dx)\right)}{d(3m + 4) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^m*(b*\operatorname{Cos}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*\operatorname{Cos}[c + d*x]^{(1 + m)}*(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/2, (4 + 3*m)/6, (10 + 3*m)/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(4 + 3*m)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt[3]{b \cos(c + dx)} \int \cos^{\frac{1}{3}+m}(c + dx) dx}{\sqrt[3]{\cos(c + dx)}} \\ &= \frac{3 \cos^{1+m}(c + dx) \sqrt[3]{b \cos(c + dx)} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(4 + 3m), \frac{1}{6}(10 + 3m), \cos^2(c + dx)\right)}{d(4 + 3m) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \frac{\cos^{1+m}(c + dx) \sqrt[3]{b \cos(c + dx)} \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{4}{3} + m\right), \frac{1}{2}\left(\frac{10}{3} + m\right), \cos^2(c + dx)\right)}{d\left(\frac{4}{3} + m\right)}$$

```
[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(1/3), x]
```

```
[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(1/3)*Csc[c + d*x]*Hypergeometric2
F1[1/2, (4/3 + m)/2, (10/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d
*(4/3 + m))
```

Maple [F]

$$\int (\cos^m(dx + c)) (\cos(dx + c)b)^{\frac{1}{3}} dx$$

[In] `int(cos(d*x+c)^m*(cos(d*x+c)*b)^(1/3),x)`

[Out] `int(cos(d*x+c)^m*(cos(d*x+c)*b)^(1/3),x)`

Fricas [F]

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m dx$$

[In] `integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)`

Sympy [F]

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int \sqrt[3]{b \cos(c + dx)} \cos^m(c + dx) dx$$

[In] `integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(1/3),x)`

[Out] `Integral((b*cos(c + d*x))**(1/3)*cos(c + d*x)**m, x)`

Maxima [F]

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m dx$$

[In] `integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)`

Giac [F]

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^m dx$$

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^m(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int \cos(c + dx)^m (b \cos(c + dx))^{1/3} dx$$

[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(1/3),x)

[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(1/3), x)

3.203 $\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal result	1087
Rubi [A] (verified)	1087
Mathematica [A] (verified)	1088
Maple [F]	1088
Fricas [F]	1089
Sympy [F(-1)]	1089
Maxima [F]	1089
Giac [F]	1089
Mupad [F(-1)]	1090

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3(b \cos(c + dx))^{10/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10b^3 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/10*(b*\cos(d*x+c))^{(10/3)}*\operatorname{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3 \sin(c + dx) (b \cos(c + dx))^{10/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right)}{10b^3 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2*(b*\operatorname{Cos}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(10/3)}*\operatorname{Hypergeometric2F1}[1/2, 5/3, 8/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(10*b^3*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)}*(b_*)^{(v_*)}*(v_*)^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}[\{b, n\}, x] \ \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{7/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{10/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \frac{3 \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{10d}$$

```
[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(1/3), x]
```

```
[Out] (-3*Cos[c + d*x]^2*(b*Cos[c + d*x])^(1/3)*Cot[c + d*x]*Hypergeometric2F1[1/
2, 5/3, 8/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(10*d)
```

Maple [F]

$$\int (\cos^2(dx + c)) (\cos(dx + c) b)^{\frac{1}{3}} dx$$

```
[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(1/3), x)
```

```
[Out] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(1/3), x)
```


Fricas [F]

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(1/3),x)

[Out] Timed out

Maxima [F]

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

Giac [F]

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int \cos(c + dx)^2 (b \cos(c + dx))^{1/3} dx$$

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3),x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3), x)
```

3.204 $\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx$

Optimal result	1091
Rubi [A] (verified)	1091
Mathematica [A] (verified)	1092
Maple [F]	1092
Fricas [F]	1093
Sympy [F]	1093
Maxima [F]	1093
Giac [F]	1093
Mupad [F(-1)]	1094

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3(b \cos(c + dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7b^2 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\operatorname{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3 \sin(c + dx) (b \cos(c + dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right)}{7b^2 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]*(b*\operatorname{Cos}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(7/3)}*\operatorname{Hypergeometric2F1}[1/2, 7/6, 13/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(7*b^2*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)}*(b_*)^{(v_*)}*(v_*)^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{4/3} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{7/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \\ -\frac{3(b \cos(c + dx))^{4/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{7bd} \end{aligned}$$

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(1/3),x]

[Out] (-3*(b*Cos[c + d*x])^(4/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, C
os[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(7*b*d)

Maple [F]

$$\int \cos(dx + c) (\cos(dx + c) b)^{\frac{1}{3}} dx$$

[In] int(cos(d*x+c)*(cos(d*x+c)*b)^(1/3),x)

[Out] int(cos(d*x+c)*(cos(d*x+c)*b)^(1/3),x)

Fricas [F]

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

Sympy [F]

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int \sqrt[3]{b \cos(c + dx)} \cos(c + dx) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(1/3)*cos(c + d*x), x)

Maxima [F]

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

Giac [F]

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*cos(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx) \sqrt[3]{b \cos(c + dx)} dx = \int \cos(c + dx) (b \cos(c + dx))^{1/3} dx$$

```
[In] int(cos(c + d*x)*(b*cos(c + d*x))^(1/3),x)
```

```
[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(1/3), x)
```

3.205 $\int \sqrt[3]{b \cos(c + dx)} dx$

Optimal result	1095
Rubi [A] (verified)	1095
Mathematica [A] (verified)	1096
Maple [F]	1096
Fricas [F]	1096
Sympy [F]	1097
Maxima [F]	1097
Giac [F]	1097
Mupad [F(-1)]	1097

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3(b \cos(c + dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{(4/3)}*\operatorname{hypergeom}([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3 \sin(c + dx)(b \cos(c + dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right)}{4bd\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(4/3)}*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(4*b*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(b \cos(c + dx))^{4/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \sqrt[3]{b \cos(c + dx)} dx$$

$$= -\frac{3\sqrt[3]{b \cos(c + dx)} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{4d}$$

[In] Integrate[(b*Cos[c + d*x])^(1/3),x]

[Out] (-3*(b*Cos[c + d*x])^(1/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(4*d)

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{1}{3}} dx$$

[In] int((cos(d*x+c)*b)^(1/3),x)

[Out] int((cos(d*x+c)*b)^(1/3),x)

Fricas [F]

$$\int \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3), x)

Sympy [F]

$$\int \sqrt[3]{b \cos(c + dx)} dx = \int \sqrt[3]{b \cos(c + dx)} dx$$

[In] integrate((b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(dx + c))^{\frac{1}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{b \cos(c + dx)} dx = \int (b \cos(c + dx))^{1/3} dx$$

[In] int((b*cos(c + d*x))^(1/3),x)

[Out] int((b*cos(c + d*x))^(1/3), x)

3.206 $\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$

Optimal result	1098
Rubi [A] (verified)	1098
Mathematica [A] (verified)	1099
Maple [F]	1099
Fricas [F]	1100
Sympy [F]	1100
Maxima [F]	1100
Giac [F]	1100
Mupad [F(-1)]	1101

Optimal result

Integrand size = 19, antiderivative size = 53

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$$

$$= -\frac{3 \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3*(b*\cos(d*x+c))^{(1/3)}*\operatorname{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$$

$$= -\frac{3 \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Sec}[c + d*x], x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/6, 1/2, 7/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)^{(m_*)}}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{2/3}} dx \\ &= -\frac{3\sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\begin{aligned} &\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx \\ &= -\frac{3b \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(b \cos(c + dx))^{2/3}} \end{aligned}$$

[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x],x]

[Out] (-3*b*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(2/3))

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{1}{3}} \sec(dx + c) dx$$

[In] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c),x)

[Out] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c),x)

Fricas [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

Sympy [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx = \int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx$$

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c),x)

[Out] Integral((b*cos(c + d*x))**(1/3)*sec(c + d*x), x)

Maxima [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

Giac [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{b \cos(c + dx)} \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^{1/3}}{\cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x),x)
```

```
[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x), x)
```

3.207 $\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$

Optimal result	1102
Rubi [A] (verified)	1102
Mathematica [A] (verified)	1103
Maple [F]	1103
Fricas [F]	1104
Sympy [F]	1104
Maxima [F]	1104
Giac [F]	1104
Mupad [F(-1)]	1105

Optimal result

Integrand size = 21, antiderivative size = 56

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$$

$$= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}}$$

[Out] $3/2*b*\operatorname{hypergeom}([-1/3, 1/2], [2/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{2/3}/(\sin(d*x+c)^2)^{1/2}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$$

$$= \frac{3b \sin(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}(b \cos(c + dx))^{2/3}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{1/3}*\operatorname{Sec}[c + d*x]^2,x]$

[Out] $(3*b*\operatorname{Hypergeometric2F1}[-1/3, 1/2, 2/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/ (2*d*(b*\operatorname{Cos}[c + d*x])^{2/3}*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)^{(m_*)}*(b_*)^{(v_*)^{(n_*)}}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{5/3}} dx \\ &= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

$$\begin{aligned} &\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx \\ &= \frac{3b \csc(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2d(b \cos(c + dx))^{2/3}} \end{aligned}$$

```
[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^2,x]
```

```
[Out] (3*b*Csc[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(2/3))
```

Maple [F]

$$\int (\cos(dx + c)b)^{\frac{1}{3}} (\sec^2(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c)^2,x)
```

```
[Out] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c)^2,x)
```

Fricas [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec^2(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

Sympy [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx = \int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx$$

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c)**2,x)

[Out] Integral((b*cos(c + d*x))**(1/3)*sec(c + d*x)**2, x)

Maxima [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec^2(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

Giac [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec^2(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{b \cos(c + dx)} \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^{1/3}}{\cos(c + dx)^2} dx$$

```
[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^2,x)
```

```
[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^2, x)
```

3.208 $\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx$

Optimal result	1106
Rubi [A] (verified)	1106
Mathematica [A] (verified)	1107
Maple [F]	1107
Fricas [F]	1108
Sympy [F]	1108
Maxima [F]	1108
Giac [F]	1108
Mupad [F(-1)]	1109

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx$$

$$= \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \cos(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}}$$

[Out] $3/5*b^2*\operatorname{hypergeom}([-5/6, 1/2], [1/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(5/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx$$

$$= \frac{3b^2 \sin(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right)}{5d\sqrt{\sin^2(c + dx)}(b \cos(c + dx))^{5/3}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Sec}[c + d*x]^3, x]$

[Out] $(3*b^2*\operatorname{Hypergeometric2F1}[-5/6, 1/2, 1/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(5*d*(b*\operatorname{Cos}[c + d*x])^{(5/3)}*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{8/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \cos(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\begin{aligned} &\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx \\ &= \frac{3 \sqrt[3]{b \cos(c + dx)} \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right) \sec^2(c + dx) \sqrt{\sin^2(c + dx)}}{5d} \end{aligned}$$

[In] Integrate[(b*Cos[c + d*x])^(1/3)*Sec[c + d*x]^3,x]

[Out] (3*(b*Cos[c + d*x])^(1/3)*Csc[c + d*x]*Hypergeometric2F1[-5/6, 1/2, 1/6, Co
s[c + d*x]^2]*Sec[c + d*x]^2*Sqrt[Sin[c + d*x]^2])/(5*d)

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{1}{3}} (\sec^3(dx + c)) dx$$

[In] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c)^3,x)

[Out] int((cos(d*x+c)*b)^(1/3)*sec(d*x+c)^3,x)

Fricas [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

Sympy [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx = \int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx$$

[In] integrate((b*cos(d*x+c))**(1/3)*sec(d*x+c)**3,x)

[Out] Integral((b*cos(c + d*x))**(1/3)*sec(c + d*x)**3, x)

Maxima [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

Giac [F]

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{\frac{1}{3}} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(1/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt[3]{b \cos(c + dx)} \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^{1/3}}{\cos(c + dx)^3} dx$$

```
[In] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^3,x)
```

```
[Out] int((b*cos(c + d*x))^(1/3)/cos(c + d*x)^3, x)
```

3.209 $\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal result	1110
Rubi [A] (verified)	1110
Mathematica [A] (verified)	1111
Maple [F]	1111
Fricas [F]	1112
Sympy [F]	1112
Maxima [F]	1112
Giac [F]	1112
Mupad [F(-1)]	1113

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3 \cos^{1+m}(c + dx)(b \cos(c + dx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(5 + 3m), \frac{1}{6}(11 + 3m), \cos^2(c + dx)\right) \sin(c + dx)}{d(5 + 3m) \sqrt{\sin^2(c + dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*(b*\cos(d*x+c))^{(2/3)}*\operatorname{hypergeom}\left(\left[\frac{1}{2}, 5/6+1/2*m\right], \left[\frac{11}{6}+1/2*m\right], \cos(d*x+c)^2\right)*\sin(d*x+c)/d/(5+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} \cos^{m+1}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m + 5), \frac{1}{6}(3m + 11), \cos^2(c + dx)\right)}{d(3m + 5) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^m*(b*\operatorname{Cos}[c + d*x])^{(2/3)}, x]$

[Out] $(-3*\operatorname{Cos}[c + d*x]^{(1 + m)}*(b*\operatorname{Cos}[c + d*x])^{(2/3)}*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, (5 + 3*m)/6, (11 + 3*m)/6, \operatorname{Cos}[c + d*x]^2\right]*\operatorname{Sin}[c + d*x])/(d*(5 + 3*m)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[b^{\operatorname{IntPart}[n]}*((b*v)^{\operatorname{FracPart}[n]}/(a^{\operatorname{IntPart}[n]}*(a*v)^{\operatorname{FracPart}[n]})), \operatorname{Int}[u*(a*v)^{(m + n}$

), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(b \cos(c + dx))^{2/3} \int \cos^{2/3+m}(c + dx) dx}{\cos^{2/3}(c + dx)} \\ &= \frac{3 \cos^{1+m}(c + dx) (b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(5 + 3m), \frac{1}{6}(11 + 3m), \cos^2(c + dx)\right)}{d(5 + 3m) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \cos^m(c + dx) (b \cos(c + dx))^{2/3} dx = \frac{\cos^{1+m}(c + dx) (b \cos(c + dx))^{2/3} \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{5}{3} + m\right), \frac{1}{2}\left(\frac{11}{3} + m\right), \cos^2(c + dx)\right)}{d\left(\frac{5}{3} + m\right)}$$

[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(2/3), x]

[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(2/3)*Csc[c + d*x]*Hypergeometric2F1[1/2, (5/3 + m)/2, (11/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(5/3 + m))

Maple [F]

$$\int (\cos^m(dx + c)) (\cos(dx + c)b)^{2/3} dx$$

[In] int(cos(d*x+c)^m*(cos(d*x+c)*b)^(2/3), x)

[Out] int(cos(d*x+c)^m*(cos(d*x+c)*b)^(2/3), x)

Fricas [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m dx$$

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

Sympy [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(c + dx))^{\frac{2}{3}} \cos^m(c + dx) dx$$

[In] integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(2/3),x)

[Out] Integral((b*cos(c + d*x))**(2/3)*cos(c + d*x)**m, x)

Maxima [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m dx$$

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

Giac [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^m dx$$

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^m(c + dx)(b \cos(c + dx))^{2/3} dx = \int \cos(c + dx)^m (b \cos(c + dx))^{2/3} dx$$

```
[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(2/3), x)
```

```
[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(2/3), x)
```

3.210 $\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal result	1114
Rubi [A] (verified)	1114
Mathematica [A] (verified)	1115
Maple [F]	1115
Fricas [F]	1116
Sympy [F(-1)]	1116
Maxima [F]	1116
Giac [F]	1116
Mupad [F(-1)]	1117

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3(b \cos(c + dx))^{11/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{17}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{11b^3 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/11*(b*\cos(d*x+c))^{(11/3)}*hypergeom([1/2, 11/6], [17/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{11/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{17}{6}, \cos^2(c + dx)\right)}{11b^3 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2*(b*\operatorname{Cos}[c + d*x])^{(2/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(11/3)}*\operatorname{Hypergeometric2F1}[1/2, 11/6, 17/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/((11*b^3*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)^{(m_*)}}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{8/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{11/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{17}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{11b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\begin{aligned} \int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \\ \frac{3 \cos^2(c + dx)(b \cos(c + dx))^{2/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{11}{6}, \frac{17}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{11d} \end{aligned}$$

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(2/3),x]

[Out] (-3*Cos[c + d*x]^2*(b*Cos[c + d*x])^(2/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 11/6, 17/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(11*d)

Maple [F]

$$\int (\cos^2(dx + c)) (\cos(dx + c) b)^{\frac{2}{3}} dx$$

[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(2/3),x)

[Out] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(2/3),x)

Fricas [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(2/3),x)

[Out] Timed out

Maxima [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

Giac [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{2/3} dx = \int \cos(c + dx)^2 (b \cos(c + dx))^{2/3} dx$$

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3), x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3), x)
```

3.211 $\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx$

Optimal result	1118
Rubi [A] (verified)	1118
Mathematica [A] (verified)	1119
Maple [F]	1119
Fricas [F]	1120
Sympy [F(-1)]	1120
Maxima [F]	1120
Giac [F]	1120
Mupad [F(-1)]	1121

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3(b \cos(c + dx))^{8/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{8b^2 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/8*(b*\cos(d*x+c))^{(8/3)}*\operatorname{hypergeom}([1/2, 4/3], [7/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{8/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right)}{8b^2 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]*(b*\operatorname{Cos}[c + d*x])^{(2/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(8/3)}*\operatorname{Hypergeometric2F1}[1/2, 4/3, 7/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(8*b^2*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)}*(b_*)^{(v_*)}*(n_*), x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x] \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{5/3} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{8/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{8b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \\ \frac{3(b \cos(c + dx))^{5/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{8bd} \end{aligned}$$

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(2/3),x]

[Out] (-3*(b*Cos[c + d*x])^(5/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 4/3, 7/3, Co
s[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(8*b*d)

Maple [F]

$$\int \cos(dx + c) (\cos(dx + c) b)^{\frac{2}{3}} dx$$

[In] int(cos(d*x+c)*(cos(d*x+c)*b)^(2/3),x)

[Out] int(cos(d*x+c)*(cos(d*x+c)*b)^(2/3),x)

Fricas [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**(2/3),x)

[Out] Timed out

Maxima [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

Giac [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*cos(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{2/3} dx = \int \cos(c + dx) (b \cos(c + dx))^{2/3} dx$$

```
[In] int(cos(c + d*x)*(b*cos(c + d*x))^(2/3), x)
```

```
[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(2/3), x)
```

3.212 $\int (b \cos(c + dx))^{2/3} dx$

Optimal result	1122
Rubi [A] (verified)	1122
Mathematica [A] (verified)	1123
Maple [F]	1123
Fricas [F]	1123
Sympy [F]	1124
Maxima [F]	1124
Giac [F]	1124
Mupad [F(-1)]	1124

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int (b \cos(c + dx))^{2/3} dx = \frac{3(b \cos(c + dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/5*(b*\cos(d*x+c))^{(5/3)*\operatorname{hypergeom}([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int (b \cos(c + dx))^{2/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right)}{5bd\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(2/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(5/3)*\operatorname{Hypergeometric2F1}[1/2, 5/6, 11/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(5*b*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 2722

$\operatorname{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2]))*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(b \cos(c + dx))^{5/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int (b \cos(c + dx))^{2/3} dx = \frac{3(b \cos(c + dx))^{2/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{5d}$$

[In] Integrate[(b*Cos[c + d*x])^(2/3),x]

[Out] (-3*(b*Cos[c + d*x])^(2/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(5*d)

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{2}{3}} dx$$

[In] int((cos(d*x+c)*b)^(2/3),x)

[Out] int((cos(d*x+c)*b)^(2/3),x)

Fricas [F]

$$\int (b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3), x)

Sympy [F]

$$\int (b \cos(c + dx))^{2/3} dx = \int (b \cos(c + dx))^{\frac{2}{3}} dx$$

[In] integrate((b*cos(d*x+c))**(2/3),x)

[Out] Integral((b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int (b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int (b \cos(c + dx))^{2/3} dx = \int (b \cos(dx + c))^{\frac{2}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} dx = \int (b \cos(c + dx))^{\frac{2}{3}} dx$$

[In] int((b*cos(c + d*x))^(2/3),x)

[Out] int((b*cos(c + d*x))^(2/3), x)

3.213 $\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx$

Optimal result	1125
Rubi [A] (verified)	1125
Mathematica [A] (verified)	1126
Maple [F]	1126
Fricas [F]	1127
Sympy [F]	1127
Maxima [F]	1127
Giac [F]	1127
Mupad [F(-1)]	1128

Optimal result

Integrand size = 19, antiderivative size = 55

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \frac{3(b \cos(c + dx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/2*(b*\cos(d*x+c))^{(2/3)}*\operatorname{hypergeom}([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \frac{3 \sin(c + dx) (b \cos(c + dx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(2/3)}*\operatorname{Sec}[c + d*x], x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(2/3)}*\operatorname{Hypergeometric2F1}[1/3, 1/2, 4/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(2*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)^{(m_*)}}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}[\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= -\frac{3(b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

$$\begin{aligned} \int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \\ -\frac{3b \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2d \sqrt[3]{b \cos(c + dx)}} \end{aligned}$$

[In] Integrate[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x],x]

[Out] (-3*b*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sqrt[Si
n[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(1/3))

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{2}{3}} \sec(dx + c) dx$$

[In] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c),x)

[Out] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c),x)

Fricas [F]

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

Sympy [F]

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \int (b \cos(c + dx))^{2/3} \sec(c + dx) dx$$

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c),x)

[Out] Integral((b*cos(c + d*x))**(2/3)*sec(c + d*x), x)

Maxima [F]

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

Giac [F]

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^{2/3}}{\cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x),x)
```

```
[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x), x)
```


3.214 $\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx$

Optimal result	1129
Rubi [A] (verified)	1129
Mathematica [A] (verified)	1130
Maple [F]	1130
Fricas [F]	1131
Sympy [F(-1)]	1131
Maxima [F]	1131
Giac [F]	1131
Mupad [F(-1)]	1132

Optimal result

Integrand size = 21, antiderivative size = 54

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt[3]{b \cos(c + dx)} \sqrt{\sin^2(c + dx)}}$$

[Out] 3*b*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \frac{3b \sin(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)} \sqrt[3]{b \cos(c + dx)}}$$

[In] Int[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^2,x]

[Out] (3*b*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{4/3}} dx \\ &= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt[3]{b \cos(c + dx)} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

$$\begin{aligned} &\int (b \cos(c + dx))^{2/3} \sec^2(c \\ &+ dx) dx = \frac{3b \operatorname{csc}(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d \sqrt[3]{b \cos(c + dx)}} \end{aligned}$$

```
[In] Integrate[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^2,x]
```

```
[Out] (3*b*Csc[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(1/3))
```

Maple [F]

$$\int (\cos(dx + c)b)^{\frac{2}{3}} (\sec^2(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c)^2,x)
```

```
[Out] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c)^2,x)
```

Fricas [F]

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c)**2,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

Giac [F]

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^{2/3}}{\cos(c + dx)^2} dx$$

```
[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^2,x)
```

```
[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^2, x)
```

3.215 $\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx$

Optimal result	1133
Rubi [A] (verified)	1133
Mathematica [A] (verified)	1134
Maple [F]	1134
Fricas [F]	1135
Sympy [F(-1)]	1135
Maxima [F]	1135
Giac [F]	1135
Mupad [F(-1)]	1136

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \cos(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}}$$

[Out] $3/4*b^2*\operatorname{hypergeom}([-2/3, 1/2], [1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{\wedge}(4/3)/(\sin(d*x+c)^2)^{\wedge}(1/2)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \frac{3b^2 \sin(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)}(b \cos(c + dx))^{4/3}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{\wedge}(2/3)*\operatorname{Sec}[c + d*x]^{\wedge}3, x]$

[Out] $(3*b^2*\operatorname{Hypergeometric2F1}[-2/3, 1/2, 1/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(4*d*(b*\operatorname{Cos}[c + d*x])^{\wedge}(4/3)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{\wedge}(v_*)^{\wedge}(m_*)*((b_*)(v_*))^{\wedge}(n_*), x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{\wedge}(m+n), x], x] /; \operatorname{FreeQ}[\{b, n\}, x] \ \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{7/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \cos(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \frac{3(b \cos(c + dx))^{2/3} \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sec^2(c + dx) \sqrt{\sin^2(c + dx)}}{4d}$$

```
[In] Integrate[(b*Cos[c + d*x])^(2/3)*Sec[c + d*x]^3,x]
```

```
[Out] (3*(b*Cos[c + d*x])^(2/3)*Csc[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Co
s[c + d*x]^2]*Sec[c + d*x]^2*Sqrt[Sin[c + d*x]^2])/(4*d)
```

Maple [F]

$$\int (\cos(dx + c)b)^{\frac{2}{3}} (\sec^3(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c)^3,x)
```

```
[Out] int((cos(d*x+c)*b)^(2/3)*sec(d*x+c)^3,x)
```

Fricas [F]

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(2/3)*sec(d*x+c)**3,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

Giac [F]

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{2/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(2/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{2/3} \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^{2/3}}{\cos(c + dx)^3} dx$$

```
[In] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^3,x)
```

```
[Out] int((b*cos(c + d*x))^(2/3)/cos(c + d*x)^3, x)
```


3.216 $\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx$

Optimal result	1137
Rubi [A] (verified)	1137
Mathematica [A] (verified)	1138
Maple [F]	1139
Fricas [F]	1139
Sympy [F(-1)]	1139
Maxima [F]	1139
Giac [F]	1140
Mupad [F(-1)]	1140

Optimal result

Integrand size = 21, antiderivative size = 83

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3b \cos^{2+m}(c + dx) \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(7 + 3m), \frac{1}{6}(13 + 3m), \cos^2(c + dx)\right) \sin(c + dx)}{d(7 + 3m)\sqrt{\sin^2(c + dx)}}$$

[Out] $-3*b*\cos(d*x+c)^{(2+m)}*(b*\cos(d*x+c))^{(1/3)}*\operatorname{hypergeom}([1/2, 7/6+1/2*m], [13/6+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(7+3*m)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \cos^{m+2}(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m + 7), \frac{1}{6}(3m + 13), \cos^2(c + dx)\right)}{d(3m + 7)\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^m*(b*\operatorname{Cos}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*b*\operatorname{Cos}[c + d*x]^{(2 + m)}*(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/2, (7 + 3*m)/6, (13 + 3*m)/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(7 + 3*m)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\left(b\sqrt[3]{b\cos(c+dx)}\right) \int \cos^{\frac{4}{3}+m}(c+dx) dx}{\sqrt[3]{\cos(c+dx)}} \\ &= \frac{3b\cos^{2+m}(c+dx)\sqrt[3]{b\cos(c+dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(7+3m), \frac{1}{6}(13+3m), \cos^2(c+dx)\right)}{d(7+3m)\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.99

$$\begin{aligned} \int \cos^m(c+dx)(b\cos(c+dx))^{4/3} dx = \\ \frac{\cos^{1+m}(c+dx)(b\cos(c+dx))^{4/3} \operatorname{csc}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{7}{3}+m\right), \frac{1}{2}\left(\frac{13}{3}+m\right), \cos^2(c+dx)\right)}{d\left(\frac{7}{3}+m\right)} \end{aligned}$$

```
[In] Integrate[Cos[c + d*x]^m*(b*Cos[c + d*x])^(4/3), x]
```

```
[Out] -((Cos[c + d*x]^(1 + m)*(b*Cos[c + d*x])^(4/3)*Csc[c + d*x]*Hypergeometric2
F1[1/2, (7/3 + m)/2, (13/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d
*(7/3 + m))
```

Maple [F]

$$\int (\cos^m(dx + c)) (\cos(dx + c)b)^{\frac{4}{3}} dx$$

[In] `int(cos(d*x+c)^m*(cos(d*x+c)*b)^(4/3),x)`

[Out] `int(cos(d*x+c)^m*(cos(d*x+c)*b)^(4/3),x)`

Fricas [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^m dx$$

[In] `integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^m*cos(d*x + c), x)`

Sympy [F(-1)]

Timed out.

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)**m*(b*cos(d*x+c))**(4/3),x)`

[Out] Timed out

Maxima [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^m dx$$

[In] `integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^m, x)`

Giac [F]

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{4/3} \cos(dx + c)^m dx$$

[In] integrate(cos(d*x+c)^m*(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^m, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^m(c + dx)(b \cos(c + dx))^{4/3} dx = \int \cos(c + dx)^m (b \cos(c + dx))^{4/3} dx$$

[In] int(cos(c + d*x)^m*(b*cos(c + d*x))^(4/3),x)

[Out] int(cos(c + d*x)^m*(b*cos(c + d*x))^(4/3), x)

3.217 $\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx$

Optimal result	.1141
Rubi [A] (verified)	.1141
Mathematica [A] (verified)	.1142
Maple [F]	.1142
Fricas [F]	.1143
Sympy [F(-1)]	.1143
Maxima [F]	.1143
Giac [F]	.1143
Mupad [F(-1)]	.1144

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3(b \cos(c + dx))^{13/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{13b^3 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/13*(b*\cos(d*x+c))^{(13/3)}*\operatorname{hypergeom}([1/2, 13/6], [19/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^{3/d}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{13/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c + dx)\right)}{13b^3 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2*(b*\operatorname{Cos}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(13/3)}*\operatorname{Hypergeometric2F1}[1/2, 13/6, 19/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(13*b^{3*d}*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)^{(m_*)}}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}[\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{10/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{13/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{13b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\begin{aligned} \int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \\ \frac{3 \cos^2(c + dx)(b \cos(c + dx))^{4/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{13}{6}, \frac{19}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{13d} \end{aligned}$$

[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^(4/3),x]

[Out] (-3*Cos[c + d*x]^2*(b*Cos[c + d*x])^(4/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 13/6, 19/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(13*d)

Maple [F]

$$\int (\cos^2(dx + c)) (\cos(dx + c) b)^{\frac{4}{3}} dx$$

[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(4/3),x)

[Out] int(cos(d*x+c)^2*(cos(d*x+c)*b)^(4/3),x)

Fricas [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^3, x)

Sympy [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^2, x)

Giac [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^{4/3} dx = \int \cos(c + dx)^2 (b \cos(c + dx))^{4/3} dx$$

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3),x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3), x)
```


3.218 $\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx$

Optimal result	1145
Rubi [A] (verified)	1145
Mathematica [A] (verified)	1146
Maple [F]	1146
Fricas [F]	1147
Sympy [F(-1)]	1147
Maxima [F]	1147
Giac [F]	1147
Mupad [F(-1)]	1148

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3(b \cos(c + dx))^{10/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10b^2 d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/10*(b*\cos(d*x+c))^{(10/3)}*\operatorname{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{10/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right)}{10b^2 d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]*(b*\operatorname{Cos}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(10/3)}*\operatorname{Hypergeometric2F1}[1/2, 5/3, 8/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(10*b^2*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_.)*(v_)^{(m_.)}*((b_.)*(v_))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{7/3} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{10/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \\ \frac{3(b \cos(c + dx))^{7/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{10bd} \end{aligned}$$

[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^(4/3),x]

[Out] (-3*(b*Cos[c + d*x])^(7/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 5/3, 8/3, Co
s[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(10*b*d)

Maple [F]

$$\int \cos(dx + c) (\cos(dx + c) b)^{\frac{4}{3}} dx$$

[In] int(cos(d*x+c)*(cos(d*x+c)*b)^(4/3),x)

[Out] int(cos(d*x+c)*(cos(d*x+c)*b)^(4/3),x)

Fricas [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{4/3} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="fricas")`

[Out] `integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)^2, x)`

Sympy [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \text{Timed out}$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))**(4/3),x)`

[Out] Timed out

Maxima [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{4/3} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c), x)`

Giac [F]

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{4/3} \cos(dx + c) dx$$

[In] `integrate(cos(d*x+c)*(b*cos(d*x+c))^(4/3),x, algorithm="giac")`

[Out] `integrate((b*cos(d*x + c))^(4/3)*cos(d*x + c), x)`

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^{4/3} dx = \int \cos(c + dx) (b \cos(c + dx))^{4/3} dx$$

```
[In] int(cos(c + d*x)*(b*cos(c + d*x))^(4/3), x)
```

```
[Out] int(cos(c + d*x)*(b*cos(c + d*x))^(4/3), x)
```

3.219 $\int (b \cos(c + dx))^{4/3} dx$

Optimal result	1149
Rubi [A] (verified)	1149
Mathematica [A] (verified)	1150
Maple [F]	1150
Fricas [F]	1150
Sympy [F(-1)]	1151
Maxima [F]	1151
Giac [F]	1151
Mupad [F(-1)]	1151

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int (b \cos(c + dx))^{4/3} dx = \frac{3(b \cos(c + dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7bd\sqrt{\sin^2(c + dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\operatorname{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int (b \cos(c + dx))^{4/3} dx = \frac{3 \sin(c + dx)(b \cos(c + dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right)}{7bd\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(7/3)}*\operatorname{Hypergeometric2F1}[1/2, 7/6, 13/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(7*b*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(b \cos(c + dx))^{7/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int (b \cos(c + dx))^{4/3} dx = \frac{3(b \cos(c + dx))^{4/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{7d}$$

[In] Integrate[(b*Cos[c + d*x])^(4/3),x]

[Out] (-3*(b*Cos[c + d*x])^(4/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(7*d)

Maple [F]

$$\int (\cos(dx + c) b)^{4/3} dx$$

[In] int((cos(d*x+c)*b)^(4/3),x)

[Out] int((cos(d*x+c)*b)^(4/3),x)

Fricas [F]

$$\int (b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{4/3} dx$$

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c), x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int (b \cos(c + dx))^{4/3} dx = \int (b \cos(dx + c))^{\frac{4}{3}} dx$$

[In] integrate((b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} dx = \int (b \cos(c + dx))^{4/3} dx$$

[In] int((b*cos(c + d*x))^(4/3),x)

[Out] int((b*cos(c + d*x))^(4/3), x)

3.220 $\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx$

Optimal result	1152
Rubi [A] (verified)	1152
Mathematica [A] (verified)	1153
Maple [F]	1153
Fricas [F]	1154
Sympy [F(-1)]	1154
Maxima [F]	1154
Giac [F]	1154
Mupad [F(-1)]	1155

Optimal result

Integrand size = 19, antiderivative size = 55

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \frac{3(b \cos(c + dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{(4/3)}*\operatorname{hypergeom}([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \frac{3 \sin(c + dx) (b \cos(c + dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right)}{4d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(4/3)}*\operatorname{Sec}[c + d*x], x]$

[Out] $(-3*(b*\operatorname{Cos}[c + d*x])^{(4/3)}*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(4*d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \sqrt[3]{b \cos(c + dx)} dx \\ &= -\frac{3(b \cos(c + dx))^{4/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

$$\begin{aligned} \int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \\ -\frac{3b \sqrt[3]{b \cos(c + dx)} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{4d} \end{aligned}$$

[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x],x]

[Out] (-3*b*(b*Cos[c + d*x])^(1/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(4*d)

Maple [F]

$$\int (\cos(dx + c) b)^{\frac{4}{3}} \sec(dx + c) dx$$

[In] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c),x)

[Out] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c),x)

Fricas [F]

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c), x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c),x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c), x)

Giac [F]

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^{4/3}}{\cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x),x)
```

```
[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x), x)
```

3.221 $\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx$

Optimal result	1156
Rubi [A] (verified)	1156
Mathematica [A] (verified)	1157
Maple [F]	1157
Fricas [F]	1158
Sympy [F(-1)]	1158
Maxima [F]	1158
Giac [F]	1158
Mupad [F(-1)]	1159

Optimal result

Integrand size = 21, antiderivative size = 54

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \frac{3b \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}}$$

[Out] $-3*b*(b*\cos(d*x+c))^{(1/3)}*\operatorname{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \frac{3b \sin(c + dx) \sqrt[3]{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{(4/3)}*\operatorname{Sec}[c + d*x]^2, x]$

[Out] $(-3*b*(b*\operatorname{Cos}[c + d*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/6, 1/2, 7/6, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{2/3}} dx \\ &= -\frac{3b^3 \sqrt{b \cos(c + dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.04

$$\begin{aligned} \int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \\ \frac{3b^2 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(b \cos(c + dx))^{2/3}} \end{aligned}$$

[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^2,x]

[Out] (-3*b^2*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(2/3))

Maple [F]

$$\int (\cos(dx + c)b)^{\frac{4}{3}} (\sec^2(dx + c)) dx$$

[In] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c)^2,x)

[Out] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c)^2,x)

Fricas [F]

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c)^2, x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c)**2,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^2, x)

Giac [F]

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^2 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^{4/3}}{\cos(c + dx)^2} dx$$

```
[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^2,x)
```

```
[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^2, x)
```

3.222 $\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx$

Optimal result	1160
Rubi [A] (verified)	1160
Mathematica [A] (verified)	1161
Maple [F]	1161
Fricas [F]	1162
Sympy [F(-1)]	1162
Maxima [F]	1162
Giac [F]	1162
Mupad [F(-1)]	1163

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}}$$

[Out] $\frac{3/2*b^2*\operatorname{hypergeom}([-1/3, 1/2], [2/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{2/3}/(\sin(d*x+c)^2)^{1/2}}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \frac{3b^2 \sin(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}(b \cos(c + dx))^{2/3}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^{4/3}*\operatorname{Sec}[c + d*x]^3,x]$

[Out] $(3*b^2*\operatorname{Hypergeometric2F1}[-1/3, 1/2, 2/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(2*d*(b*\operatorname{Cos}[c + d*x])^{2/3}*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\amp; \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{5/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} &\int (b \cos(c + dx))^{4/3} \sec^3(c \\ &+ dx) dx = \frac{3b^2 \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2d(b \cos(c + dx))^{2/3}} \end{aligned}$$

```
[In] Integrate[(b*Cos[c + d*x])^(4/3)*Sec[c + d*x]^3,x]
```

```
[Out] (3*b^2*Csc[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sqrt[
Sin[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(2/3))
```

Maple [F]

$$\int (\cos(dx + c)b)^{\frac{4}{3}} (\sec^3(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c)^3,x)
```

```
[Out] int((cos(d*x+c)*b)^(4/3)*sec(d*x+c)^3,x)
```

Fricas [F]

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c)^3, x)

Sympy [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**(4/3)*sec(d*x+c)**3,x)

[Out] Timed out

Maxima [F]

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^3, x)

Giac [F]

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \int (b \cos(dx + c))^{4/3} \sec(dx + c)^3 dx$$

[In] integrate((b*cos(d*x+c))^(4/3)*sec(d*x+c)^3,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3)*sec(d*x + c)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^{4/3} \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^{4/3}}{\cos(c + dx)^3} dx$$

```
[In] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^3,x)
```

```
[Out] int((b*cos(c + d*x))^(4/3)/cos(c + d*x)^3, x)
```

$$3.223 \quad \int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1164
Rubi [A] (verified)	1164
Mathematica [A] (verified)	1165
Maple [F]	1165
Fricas [F]	1166
Sympy [F]	1166
Maxima [F]	1166
Giac [F]	1166
Mupad [F(-1)]	1167

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3 \cos^{1+m}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(2+3m), \frac{1}{6}(8+3m), \cos^2(c+dx)\right) \sin(c+dx)}{d(2+3m) \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*\operatorname{hypergeom}([1/2, 1/3+1/2*m], [4/3+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(2+3*m)/(b*\cos(d*x+c))^{(1/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{\cos^m(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3 \sin(c+dx) \cos^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m+2), \frac{1}{6}(3m+8), \cos^2(c+dx)\right)}{d(3m+2) \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^m/(b*\operatorname{Cos}[c+d*x])^{(1/3)}, x]$

[Out] $(-3*\operatorname{Cos}[c+d*x]^{(1+m)}*\operatorname{Hypergeometric2F1}[1/2, (2+3*m)/6, (8+3*m)/6, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(d*(2+3*m)*(b*\operatorname{Cos}[c+d*x])^{(1/3)}*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt[3]{\cos(c+dx)} \int \cos^{-\frac{1}{3}+m}(c+dx) dx}{\sqrt[3]{b} \cos(c+dx)} \\ &= -\frac{3 \cos^{1+m}(c+dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(2+3m), \frac{1}{6}(8+3m), \cos^2(c+dx)\right) \sin(c+dx)}{d(2+3m) \sqrt[3]{b} \cos(c+dx) \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \frac{\cos^m(c+dx)}{\sqrt[3]{b} \cos(c+dx)} dx = -\frac{\cos^{1+m}(c+dx) \csc(c+dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{2}{3}+m\right), \frac{1}{2}\left(\frac{8}{3}+m\right), \cos^2(c+dx)\right) \sqrt{\sin^2(c+dx)}}{d\left(\frac{2}{3}+m\right) \sqrt[3]{b} \cos(c+dx)}$$

```
[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(1/3), x]
```

```
[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (2/3 + m)/2, (8
/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(2/3 + m)*(b*Cos[c + d*
x])^(1/3))
```

Maple [F]

$$\int \frac{\cos^m(dx+c)}{(\cos(dx+c)b)^{\frac{1}{3}}} dx$$

```
[In] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(1/3), x)
```

```
[Out] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(1/3), x)
```

Fricas [F]

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^m(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{1/3}} dx$$

```
[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(1/3), x)
```

```
[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(1/3), x)
```

$$3.224 \quad \int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1168
Rubi [A] (verified)	1168
Mathematica [A] (verified)	1169
Maple [F]	1169
Fricas [F]	1170
Sympy [F]	1170
Maxima [F]	1170
Giac [F]	1170
Mupad [F(-1)]	1171

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

$$= -\frac{3(b \cos(c+dx))^{8/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{8b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/8*(b*\cos(d*x+c))^{(8/3)}*\operatorname{hypergeom}([1/2, 4/3], [7/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

$$= -\frac{3 \sin(c+dx)(b \cos(c+dx))^{8/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c+dx)\right)}{8b^3 d \sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^2/(b*\operatorname{Cos}[c+d*x])^{(1/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c+d*x])^{(8/3)}*\operatorname{Hypergeometric2F1}[1/2, 4/3, 7/3, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(8*b^3*d*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{5/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{8/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{8b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\begin{aligned} &\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= -\frac{3 \cos^2(c + dx) \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{4}{3}, \frac{7}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{8d \sqrt[3]{b \cos(c + dx)}} \end{aligned}$$

[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(1/3), x]

[Out] (-3*Cos[c + d*x]^2*Cot[c + d*x]*Hypergeometric2F1[1/2, 4/3, 7/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(8*d*(b*Cos[c + d*x])^(1/3))

Maple [F]

$$\int \frac{\cos^2(dx + c)}{(\cos(dx + c)b)^{1/3}} dx$$

[In] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(1/3), x)

[Out] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(1/3), x)

Fricas [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)/b, x)

Sympy [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(cos(c + d*x)**2/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{1/3}} dx$$

```
[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/3), x)
```

```
[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(1/3), x)
```

$$3.225 \quad \int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1172
Rubi [A] (verified)	1172
Mathematica [A] (verified)	1173
Maple [F]	1173
Fricas [F]	1174
Sympy [F]	1174
Maxima [F]	1174
Giac [F]	1174
Mupad [F(-1)]	1175

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

$$= -\frac{3(b \cos(c+dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{5b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] -3/5*(b*cos(d*x+c))^(5/3)*hypergeom([1/2, 5/6],[11/6],cos(d*x+c)^2)*sin(d*x+c)/b^2/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\cos(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

$$= -\frac{3 \sin(c+dx) (b \cos(c+dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right)}{5b^2 d \sqrt{\sin^2(c+dx)}}$$

[In] Int[Cos[c + d*x]/(b*Cos[c + d*x])^(1/3),x]

[Out] (-3*(b*Cos[c + d*x])^(5/3)*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sin[c + d*x])/(5*b^2*d*Sqrt[Sin[c + d*x]^2])

Rule 16

```
Int[(u_.)*(v_)^(m_.)*((b_.)*(v_)^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{2/3} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{5/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = -\frac{3(b \cos(c + dx))^{2/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{5bd}$$

```
[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(1/3), x]
```

```
[Out] (-3*(b*Cos[c + d*x])^(2/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(5*b*d)
```

Maple [F]

$$\int \frac{\cos(dx + c)}{(\cos(dx + c)b)^{1/3}} dx$$

```
[In] int(cos(d*x+c)/(cos(d*x+c)*b)^(1/3), x)
```

```
[Out] int(cos(d*x+c)/(cos(d*x+c)*b)^(1/3), x)
```

Fricas [F]

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/b, x)

Sympy [F]

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(cos(c + d*x)/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{1/3}} dx$$

```
[In] int(cos(c + d*x)/(b*cos(c + d*x))^(1/3), x)
```

```
[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(1/3), x)
```

$$3.226 \quad \int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

Optimal result	1176
Rubi [A] (verified)	1176
Mathematica [A] (verified)	1177
Maple [F]	1177
Fricas [F]	1177
Sympy [F]	1178
Maxima [F]	1178
Giac [F]	1178
Mupad [F(-1)]	1178

Optimal result

Integrand size = 12, antiderivative size = 58

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

$$= -\frac{3(b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2bd\sqrt{\sin^2(c + dx)}}$$

[Out] -3/2*(b*cos(d*x+c))^(2/3)*hypergeom([1/3, 1/2],[4/3],cos(d*x+c)^2)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

$$= -\frac{3 \sin(c + dx)(b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right)}{2bd\sqrt{\sin^2(c + dx)}}$$

[In] Int[(b*Cos[c + d*x])^(-1/3),x]

[Out] (-3*(b*Cos[c + d*x])^(2/3)*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sin[c + d*x])/(2*b*d*Sqrt[Sin[c + d*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2

F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x]
 && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3(b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.95

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

$$= -\frac{3 \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2d\sqrt[3]{b \cos(c + dx)}}$$

[In] Integrate[(b*Cos[c + d*x])^(-1/3),x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(1/3))

Maple [F]

$$\int \frac{1}{(\cos(dx + c)b)^{1/3}} dx$$

[In] int(1/(cos(d*x+c)*b)^(1/3),x)

[Out] int(1/(cos(d*x+c)*b)^(1/3),x)

Fricas [F]

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c))^{1/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(1/(b*cos(d*x+c))**(1/3),x)

[Out] Integral((b*cos(c + d*x))**(-1/3), x)

Maxima [F]

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{(b \cos(c + dx))^{1/3}} dx$$

[In] int(1/(b*cos(c + d*x))^(1/3),x)

[Out] int(1/(b*cos(c + d*x))^(1/3), x)

$$3.227 \quad \int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1179
Rubi [A] (verified)	1179
Mathematica [A] (verified)	1180
Maple [F]	1180
Fricas [F]	1181
Sympy [F]	1181
Maxima [F]	1181
Giac [F]	1181
Mupad [F(-1)]	1182

Optimal result

Integrand size = 19, antiderivative size = 53

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{d \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\sec(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{4/3}} dx \\ &= \frac{3 \text{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{d \sqrt[3]{b \cos(c + dx)} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.02

$$\begin{aligned} &\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= \frac{3b \cot(c + dx) \text{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(b \cos(c + dx))^{4/3}} \end{aligned}$$

```
[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(1/3),x]
```

```
[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[Si
n[c + d*x]^2])/(d*(b*Cos[c + d*x])^(4/3))
```

Maple [F]

$$\int \frac{\sec(dx + c)}{(\cos(dx + c)b)^{1/3}} dx$$

```
[In] int(sec(d*x+c)/(cos(d*x+c)*b)^(1/3),x)
```

```
[Out] int(sec(d*x+c)/(cos(d*x+c)*b)^(1/3),x)
```

Fricas [F]

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{1/3}} dx$$

```
[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/3)),x)
```

```
[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(1/3)), x)
```

$$3.228 \quad \int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1183
Rubi [A] (verified)	1183
Mathematica [A] (verified)	1184
Maple [F]	1184
Fricas [F]	1185
Sympy [F]	1185
Maxima [F]	1185
Giac [F]	1185
Mupad [F(-1)]	1186

Optimal result

Integrand size = 21, antiderivative size = 56

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{4d(b \cos(c+dx))^{4/3} \sqrt{\sin^2(c+dx)}}$$

[Out] $3/4*b*\operatorname{hypergeom}([-2/3, 1/2], [1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(4/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3b \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{4/3}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c+d*x]^2/(b*\operatorname{Cos}[c+d*x])^{(1/3)}, x]$

[Out] $(3*b*\operatorname{Hypergeometric2F1}[-2/3, 1/2, 1/3, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(4*d*(b*\operatorname{Cos}[c+d*x])^{(4/3)}*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{7/3}} dx \\ &= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \cos(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.04

$$\begin{aligned} &\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= \frac{3b^2 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{4d(b \cos(c + dx))^{7/3}} \end{aligned}$$

[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(1/3),x]

[Out] (3*b^2*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(4*d*(b*Cos[c + d*x])^(7/3))

Maple [F]

$$\int \frac{\sec^2(dx + c)}{(\cos(dx + c)b)^{\frac{1}{3}}} dx$$

[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(1/3),x)

[Out] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(1/3),x)

Fricas [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{1/3}} dx$$

```
[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3)),x)
```

```
[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(1/3)), x)
```

$$3.229 \quad \int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx$$

Optimal result	1187
Rubi [A] (verified)	1187
Mathematica [A] (verified)	1188
Maple [F]	1188
Fricas [F]	1189
Sympy [F]	1189
Maxima [F]	1189
Giac [F]	1189
Mupad [F(-1)]	1190

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{7d(b \cos(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/7*b^2*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(7/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^3(c+dx)}{\sqrt[3]{b \cos(c+dx)}} dx = \frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(1/3), x]

[Out] (3*b^2*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Cos[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{10/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7d(b \cos(c + dx))^{7/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} &\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx \\ &= \frac{3b^2 \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{7d(b \cos(c + dx))^{7/3}} \end{aligned}$$

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(1/3),x]

[Out] (3*b^2*Csc[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(7*d*(b*Cos[c + d*x])^(7/3))

Maple [F]

$$\int \frac{\sec^3(dx + c)}{(\cos(dx + c) b)^{1/3}} dx$$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(1/3),x)

[Out] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(1/3),x)

Fricas [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(1/3), x)

Maxima [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(1/3), x)

Giac [F]

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{\frac{1}{3}}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(1/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{\sqrt[3]{b \cos(c + dx)}} dx = \int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{1/3}} dx$$

```
[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/3)),x)
```

```
[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(1/3)), x)
```

$$3.230 \quad \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal result	1191
Rubi [A] (verified)	1191
Mathematica [A] (verified)	1192
Maple [F]	1192
Fricas [F]	1193
Sympy [F]	1193
Maxima [F]	1193
Giac [F]	1193
Mupad [F(-1)]	1194

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \cos^{1+m}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(1+3m), \frac{1}{6}(7+3m), \cos^2(c+dx)\right) \sin(c+dx)}{d(1+3m)(b \cos(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}}$$

[Out] $-3*\cos(d*x+c)^{(1+m)}*\operatorname{hypergeom}([1/2, 1/6+1/2*m], [7/6+1/2*m], \cos(d*x+c)^2)*\sin(d*x+c)/d/(1+3*m)/(b*\cos(d*x+c))^{(2/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \sin(c+dx) \cos^{m+1}(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m+1), \frac{1}{6}(3m+7), \cos^2(c+dx)\right)}{d(3m+1) \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{2/3}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^m/(b*\operatorname{Cos}[c+d*x])^{(2/3)}, x]$

[Out] $(-3*\operatorname{Cos}[c+d*x]^{(1+m)}*\operatorname{Hypergeometric2F1}[1/2, (1+3*m)/6, (7+3*m)/6, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/d*(1+3*m)*(b*\operatorname{Cos}[c+d*x])^{(2/3)}*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*(b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m +
n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\cos^{\frac{2}{3}}(c + dx) \int \cos^{-\frac{2}{3}+m}(c + dx) dx}{(b \cos(c + dx))^{2/3}} \\ &= -\frac{3 \cos^{1+m}(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(1 + 3m), \frac{1}{6}(7 + 3m), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + 3m)(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \frac{\cos^{1+m}(c + dx) \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{1}{3} + m\right), \frac{1}{2}\left(\frac{7}{3} + m\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(\frac{1}{3} + m\right)(b \cos(c + dx))^{2/3}}$$

```
[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(2/3),x]
```

```
[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (1/3 + m)/2, (7
/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1/3 + m)*(b*Cos[c + d*
x])^(2/3))
```

Maple [F]

$$\int \frac{\cos^m(dx + c)}{(\cos(dx + c)b)^{\frac{2}{3}}} dx$$

```
[In] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(2/3),x)
```

```
[Out] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(2/3),x)
```


Fricas [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)^m/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{2/3}} dx$$

```
[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(2/3),x)
```

```
[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(2/3), x)
```

$$3.231 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal result	1195
Rubi [A] (verified)	1195
Mathematica [A] (verified)	1196
Maple [F]	1196
Fricas [F]	1197
Sympy [F(-1)]	1197
Maxima [F]	1197
Giac [F]	1197
Mupad [F(-1)]	1198

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3(b \cos(c+dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{7b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/7*(b*\cos(d*x+c))^{(7/3)}*\operatorname{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \sin(c+dx)(b \cos(c+dx))^{7/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c+dx)\right)}{7b^3 d \sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]^2/(b*\operatorname{Cos}[c+d*x])^{(2/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c+d*x])^{(7/3)}*\operatorname{Hypergeometric2F1}[1/2, 7/6, 13/6, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(7*b^3*d*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 16

```
Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)
^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{4/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{7/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.09

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \frac{3 \cos^2(c + dx) \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{7}{6}, \frac{13}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{7d(b \cos(c + dx))^{2/3}}$$

```
[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]
```

```
[Out] (-3*Cos[c + d*x]^2*Cot[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[c + d
*x]^2]*Sqrt[Sin[c + d*x]^2])/(7*d*(b*Cos[c + d*x])^(2/3))
```

Maple [F]

$$\int \frac{\cos^2(dx + c)}{(\cos(dx + c)b)^{2/3}} dx$$

```
[In] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(2/3), x)
```

```
[Out] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(2/3), x)
```

Fricas [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*cos(d*x + c)/b, x)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(2/3),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{2/3}} dx$$

```
[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(2/3),x)
```

```
[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(2/3), x)
```

$$3.232 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal result	1199
Rubi [A] (verified)	1199
Mathematica [A] (verified)	1200
Maple [F]	1200
Fricas [F]	1201
Sympy [F]	1201
Maxima [F]	1201
Giac [F]	1201
Mupad [F(-1)]	1202

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3(b \cos(c+dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{4b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/4*(b*\cos(d*x+c))^{4/3}*hypergeom([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{1/2}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \sin(c+dx)(b \cos(c+dx))^{4/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c+dx)\right)}{4b^2 d \sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]/(b*\operatorname{Cos}[c+d*x])^{2/3}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c+d*x])^{4/3}*\operatorname{Hypergeometric2F1}[1/2, 2/3, 5/3, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(4*b^2*d*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 16

```
Int[(u_.)*(v_)^(m_.)*((b_.)*(v_.))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \sqrt[3]{b \cos(c + dx)} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{4/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = -\frac{3 \sqrt[3]{b \cos(c + dx)} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2}{3}, \frac{5}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{4bd}$$

```
[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(2/3), x]
```

```
[Out] (-3*(b*Cos[c + d*x])^(1/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(4*b*d)
```

Maple [F]

$$\int \frac{\cos(dx + c)}{(\cos(dx + c)b)^{2/3}} dx$$

```
[In] int(cos(d*x+c)/(cos(d*x+c)*b)^(2/3), x)
```

```
[Out] int(cos(d*x+c)/(cos(d*x+c)*b)^(2/3), x)
```


Fricas [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)/b, x)

Sympy [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(cos(c + d*x)/(b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

```
[In] int(cos(c + d*x)/(b*cos(c + d*x))^(2/3), x)
```

```
[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(2/3), x)
```

$$3.233 \quad \int \frac{1}{(b \cos(c+dx))^{2/3}} dx$$

Optimal result	1203
Rubi [A] (verified)	1203
Mathematica [A] (verified)	1204
Maple [F]	1204
Fricas [F]	1204
Sympy [F]	1205
Maxima [F]	1205
Giac [F]	1205
Mupad [F(-1)]	1205

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(b \cos(c+dx))^{2/3}} dx = \frac{3\sqrt[3]{b \cos(c+dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{bd\sqrt{\sin^2(c+dx)}}$$

[Out] $-3*(b*\cos(d*x+c))^{(1/3)}*\operatorname{hypergeom}([1/6, 1/2], [7/6], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \sin(c+dx) \sqrt[3]{b \cos(c+dx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c+dx)\right)}{bd\sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c+d*x])^{(-2/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c+d*x])^{(1/3)}*\operatorname{Hypergeometric2F1}[1/6, 1/2, 7/6, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(b*d*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 2722

$\operatorname{Int}[(b_*)*\sin((c_*) + (d_*)(x_))]^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c+d*x]*((b*\operatorname{Sin}[c+d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]^2])]*\operatorname{Hypergeometric2}$

F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
 && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{3\sqrt[3]{b \cos(c + dx)} \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{bd\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = -\frac{3 \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(b \cos(c + dx))^{2/3}}$$

[In] Integrate[(b*Cos[c + d*x])^(-2/3),x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(2/3))

Maple [F]

$$\int \frac{1}{(\cos(dx + c)b)^{2/3}} dx$$

[In] int(1/(cos(d*x+c)*b)^(2/3),x)

[Out] int(1/(cos(d*x+c)*b)^(2/3),x)

Fricas [F]

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))**(2/3),x)

[Out] Integral((b*cos(c + d*x))**(-2/3), x)

Maxima [F]

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{(b \cos(c + dx))^{2/3}} dx$$

[In] int(1/(b*cos(c + d*x))^(2/3),x)

[Out] int(1/(b*cos(c + d*x))^(2/3), x)

3.234 $\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx$

Optimal result	1206
Rubi [A] (verified)	1206
Mathematica [A] (verified)	1207
Maple [F]	1207
Fricas [F]	1207
Sympy [F]	1208
Maxima [F]	1208
Giac [F]	1208
Mupad [F(-1)]	1208

Optimal result

Integrand size = 19, antiderivative size = 55

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{2d(b \cos(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/2*hypergeom([-1/3, 1/2], [2/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c+dx)\right)}{2d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{2/3}}$$

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sin[c + d*x])/(2*d*(b*Cos[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{5/3}} dx \\ &= \frac{3 \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2d(b \cos(c + dx))^{2/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \frac{3b \cot(c + dx) \text{Hypergeometric2F1}\left(-\frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2d(b \cos(c + dx))^{5/3}}$$

```
[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(2/3), x]
```

```
[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*Sqrt[Si
n[c + d*x]^2])/(2*d*(b*Cos[c + d*x])^(5/3))
```

Maple [F]

$$\int \frac{\sec(dx + c)}{(\cos(dx + c)b)^{2/3}} dx$$

```
[In] int(sec(d*x+c)/(cos(d*x+c)*b)^(2/3), x)
```

```
[Out] int(sec(d*x+c)/(cos(d*x+c)*b)^(2/3), x)
```

Fricas [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

```
[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3), x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)/(b*cos(d*x + c)), x)
```

Sympy [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{2/3}} dx$$

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(2/3)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(2/3)), x)

$$3.235 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx$$

Optimal result	1209
Rubi [A] (verified)	1209
Mathematica [A] (verified)	1210
Maple [F]	1210
Fricas [F]	1210
Sympy [F]	1211
Maxima [F]	1211
Giac [F]	1211
Mupad [F(-1)]	1211

Optimal result

Integrand size = 21, antiderivative size = 56

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{5d(b \cos(c+dx))^{5/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/5*b*hypergeom([-5/6, 1/2], [1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(5/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3b \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c+dx)\right)}{5d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{5/3}}$$

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*b*Hypergeometric2F1[-5/6, 1/2, 1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(5*d*(b*Cos[c + d*x])^(5/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{8/3}} dx \\ &= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5d(b \cos(c + dx))^{5/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.04

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \frac{3b^2 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{6}, \frac{1}{2}, \frac{1}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{5d(b \cos(c + dx))^{8/3}}$$

```
[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(2/3), x]
```

```
[Out] (3*b^2*Cot[c + d*x]*Hypergeometric2F1[-5/6, 1/2, 1/6, Cos[c + d*x]^2]*Sqrt[
Sin[c + d*x]^2])/(5*d*(b*Cos[c + d*x])^(8/3))
```

Maple [F]

$$\int \frac{\sec^2(dx + c)}{(\cos(dx + c)b)^{2/3}} dx$$

```
[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(2/3), x)
```

```
[Out] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(2/3), x)
```

Fricas [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

```
[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3), x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^2/(b*cos(d*x + c)), x)
```

Sympy [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(2/3), x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(2/3), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{2/3}} dx$$

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3)), x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(2/3)), x)

3.236 $\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx$

Optimal result	1212
Rubi [A] (verified)	1212
Mathematica [A] (verified)	1213
Maple [F]	1213
Fricas [F]	1213
Sympy [F]	1214
Maxima [F]	1214
Giac [F]	1214
Mupad [F(-1)]	1214

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{1}{2}, -\frac{1}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{8d(b \cos(c+dx))^{8/3} \sqrt{\sin^2(c+dx)}}$$

[Out] $3/8*b^2*\operatorname{hypergeom}([-4/3, 1/2], [-1/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\cos(d*x+c))^{(8/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{2/3}} dx = \frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{1}{2}, -\frac{1}{3}, \cos^2(c+dx)\right)}{8d\sqrt{\sin^2(c+dx)}(b \cos(c+dx))^{8/3}}$$

[In] $\operatorname{Int}[\operatorname{Sec}[c + d*x]^3/(b*\operatorname{Cos}[c + d*x])^{(2/3)}, x]$

[Out] $(3*b^2*\operatorname{Hypergeometric2F1}[-4/3, 1/2, -1/3, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(8*d*(b*\operatorname{Cos}[c + d*x])^{(8/3)}*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)^{(v_*)}*((b_*)^{(v_*)})^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n, x\} \ \&\amp; \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{11/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{1}{2}, -\frac{1}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{8d(b \cos(c + dx))^{8/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \frac{3b^2 \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{4}{3}, \frac{1}{2}, -\frac{1}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{8d(b \cos(c + dx))^{8/3}}$$

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(2/3), x]

[Out] (3*b^2*Csc[c + d*x]*Hypergeometric2F1[-4/3, 1/2, -1/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(8*d*(b*Cos[c + d*x])^(8/3))

Maple [F]

$$\int \frac{\sec^3(dx + c)}{(\cos(dx + c)b)^{2/3}} dx$$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(2/3), x)

[Out] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(2/3), x)

Fricas [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(1/3)*sec(d*x + c)^3/(b*cos(d*x + c)), x)

Sympy [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(2/3),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(2/3), x)

Maxima [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(2/3), x)

Giac [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{2/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(2/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{2/3}} dx = \int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{2/3}} dx$$

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(2/3)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(2/3)), x)

$$3.237 \quad \int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal result	1215
Rubi [A] (verified)	1215
Mathematica [A] (verified)	1216
Maple [F]	1216
Fricas [F]	1217
Sympy [F]	1217
Maxima [F]	1217
Giac [F]	1217
Mupad [F(-1)]	1218

Optimal result

Integrand size = 21, antiderivative size = 83

$$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \cos^m(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-1+3m), \frac{1}{6}(5+3m), \cos^2(c+dx)\right) \sin(c+dx)}{bd(1-3m) \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

[Out] 3*cos(d*x+c)^m*hypergeom([1/2, -1/6+1/2*m], [5/6+1/2*m], cos(d*x+c)^2)*sin(d*x+c)/b/d/(1-3*m)/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \sin(c+dx) \cos^m(c+dx) \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(3m-1), \frac{1}{6}(3m+5), \cos^2(c+dx)\right)}{bd(1-3m) \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[In] Int[Cos[c + d*x]^m/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*Cos[c + d*x]^m*Hypergeometric2F1[1/2, (-1 + 3*m)/6, (5 + 3*m)/6, Cos[c + d*x]^2]*Sin[c + d*x])/ (b*d*(1 - 3*m)*(b*Cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt[3]{\cos(c+dx)} \int \cos^{-\frac{4}{3}+m}(c+dx) dx}{b^3 \sqrt[3]{b \cos(c+dx)}} \\ &= \frac{3 \cos^m(c+dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{6}(-1+3m), \frac{1}{6}(5+3m), \cos^2(c+dx)\right) \sin(c+dx)}{bd(1-3m) \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.99

$$\int \frac{\cos^m(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{\cos^{1+m}(c+dx) \csc(c+dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{1}{3}+m\right), \frac{1}{2}\left(\frac{5}{3}+m\right), \cos^2(c+dx)\right) \sqrt{\sin^2(c+dx)}}{d\left(-\frac{1}{3}+m\right) (b \cos(c+dx))^{4/3}}$$

```
[In] Integrate[Cos[c + d*x]^m/(b*Cos[c + d*x])^(4/3), x]
```

```
[Out] -((Cos[c + d*x]^(1 + m)*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/3 + m)/2, (
5/3 + m)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-1/3 + m)*(b*Cos[c +
d*x])^(4/3))
```

Maple [F]

$$\int \frac{\cos^m(dx+c)}{(\cos(dx+c)b)^{\frac{4}{3}}} dx$$

```
[In] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(4/3), x)
```

```
[Out] int(cos(d*x+c)^m/(cos(d*x+c)*b)^(4/3), x)
```


Fricas [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*cos(d*x + c)^m/(b^2*cos(d*x + c)^2), x)

Sympy [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

[In] integrate(cos(d*x+c)**m/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(cos(c + d*x)**m/(b*cos(c + d*x))**(4/3), x)

Maxima [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^m}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^m/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^m/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^m(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(c + dx)^m}{(b \cos(c + dx))^{4/3}} dx$$

```
[In] int(cos(c + d*x)^m/(b*cos(c + d*x))^(4/3),x)
```

```
[Out] int(cos(c + d*x)^m/(b*cos(c + d*x))^(4/3), x)
```

$$3.238 \quad \int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal result	1219
Rubi [A] (verified)	1219
Mathematica [A] (verified)	1220
Maple [F]	1220
Fricas [F]	1221
Sympy [F(-1)]	1221
Maxima [F]	1221
Giac [F]	1221
Mupad [F(-1)]	1222

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3(b \cos(c+dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{5b^3 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/5*(b*\cos(d*x+c))^{5/3}*hypergeom([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/b^3/d/(\sin(d*x+c)^2)^{1/2}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\cos^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \sin(c+dx)(b \cos(c+dx))^{5/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c+dx)\right)}{5b^3 d \sqrt{\sin^2(c+dx)}}$$

[In] $\text{Int}[\text{Cos}[c + d*x]^2/(b*\text{Cos}[c + d*x])^{4/3}, x]$

[Out] $(-3*(b*\text{Cos}[c + d*x])^{5/3}*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*b^3*d*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

```
Int[(u_)*(v_)^(m_)*((b_)*(v_)^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]
```

Rule 2722

```
Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n+1)/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{2/3} dx}{b^2} \\ &= -\frac{3(b \cos(c + dx))^{5/3} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{5b^3 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \\ -\frac{3(b \cos(c + dx))^{2/3} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{5b^2 d} \end{aligned}$$

```
[In] Integrate[Cos[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]
```

```
[Out] (-3*(b*Cos[c + d*x])^(2/3)*Cot[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(5*b^2*d)
```

Maple [F]

$$\int \frac{\cos^2(dx + c)}{(\cos(dx + c)b)^{4/3}} dx$$

```
[In] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(4/3), x)
```

```
[Out] int(cos(d*x+c)^2/(cos(d*x+c)*b)^(4/3), x)
```

Fricas [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/b^2, x)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**2/(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)^2/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(c + dx)^2}{(b \cos(c + dx))^{4/3}} dx$$

```
[In] int(cos(c + d*x)^2/(b*cos(c + d*x))^(4/3), x)
```

```
[Out] int(cos(c + d*x)^2/(b*cos(c + d*x))^(4/3), x)
```

$$3.239 \quad \int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal result	1223
Rubi [A] (verified)	1223
Mathematica [A] (verified)	1224
Maple [F]	1224
Fricas [F]	1225
Sympy [F(-1)]	1225
Maxima [F]	1225
Giac [F]	1225
Mupad [F(-1)]	1226

Optimal result

Integrand size = 19, antiderivative size = 58

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3(b \cos(c+dx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{2b^2 d \sqrt{\sin^2(c+dx)}}$$

[Out] $-3/2*(b*\cos(d*x+c))^{(2/3)}*\operatorname{hypergeom}([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/b^2/d/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\cos(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \sin(c+dx) (b \cos(c+dx))^{2/3} \operatorname{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c+dx)\right)}{2b^2 d \sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c+d*x]/(b*\operatorname{Cos}[c+d*x])^{(4/3)}, x]$

[Out] $(-3*(b*\operatorname{Cos}[c+d*x])^{(2/3)}*\operatorname{Hypergeometric2F1}[1/3, 1/2, 4/3, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/(2*b^2*d*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2722

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt[3]{b \cos(c + dx)}} dx}{b} \\ &= -\frac{3(b \cos(c + dx))^{2/3} \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{2b^2 d \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\begin{aligned} \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \\ -\frac{3 \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{3}, \frac{1}{2}, \frac{4}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{2bd \sqrt[3]{b \cos(c + dx)}} \end{aligned}$$

[In] Integrate[Cos[c + d*x]/(b*Cos[c + d*x])^(4/3),x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(2*b*d*(b*Cos[c + d*x])^(1/3))

Maple [F]

$$\int \frac{\cos(dx + c)}{(\cos(dx + c)b)^{4/3}} dx$$

[In] int(cos(d*x+c)/(cos(d*x+c)*b)^(4/3),x)

[Out] int(cos(d*x+c)/(cos(d*x+c)*b)^(4/3),x)

Fricas [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/(b^2*cos(d*x + c)), x)

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))**(4/3),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(cos(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\cos(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

```
[In] int(cos(c + d*x)/(b*cos(c + d*x))^(4/3), x)
```

```
[Out] int(cos(c + d*x)/(b*cos(c + d*x))^(4/3), x)
```

$$3.240 \quad \int \frac{1}{(b \cos(c+dx))^{4/3}} dx$$

Optimal result	1227
Rubi [A] (verified)	1227
Mathematica [A] (verified)	1228
Maple [F]	1228
Fricas [F]	1228
Sympy [F]	1228
Maxima [F]	1229
Giac [F]	1229
Mupad [F(-1)]	1229

Optimal result

Integrand size = 12, antiderivative size = 56

$$\int \frac{1}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{bd \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*sin(d*x+c)/b/d/(b*cos(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\int \frac{1}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \cos(c+dx)}}$$

[In] Int[(b*cos[c + d*x])^(-4/3),x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sin[c + d*x])/(b*d*(b*cos[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\text{integral} = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{bd \sqrt[3]{b \cos(c+dx)} \sqrt{\sin^2(c+dx)}}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.95

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \frac{3 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{6}, \frac{1}{2}, \frac{5}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(b \cos(c + dx))^{4/3}}$$

[In] Integrate[(b*Cos[c + d*x])^(-4/3),x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(b*Cos[c + d*x])^(4/3))

Maple [F]

$$\int \frac{1}{(\cos(dx + c)b)^{\frac{4}{3}}} dx$$

[In] int(1/(cos(d*x+c)*b)^(4/3),x)

[Out] int(1/(cos(d*x+c)*b)^(4/3),x)

Fricas [F]

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c))^{\frac{4}{3}}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)/(b^2*cos(d*x + c)^2), x)

Sympy [F]

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(c + dx))^{\frac{4}{3}}} dx$$

[In] integrate(1/(b*cos(d*x+c))**(4/3),x)

[Out] Integral((b*cos(c + d*x))**(-4/3), x)

Maxima [F]

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(1/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{(b \cos(c + dx))^{4/3}} dx$$

[In] int(1/(b*cos(c + d*x))^(4/3),x)

[Out] int(1/(b*cos(c + d*x))^(4/3), x)

3.241 $\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx$

Optimal result	1230
Rubi [A] (verified)	1230
Mathematica [A] (verified)	.1231
Maple [F]	.1231
Fricas [F]	.1231
Sympy [F]	1232
Maxima [F]	1232
Giac [F]	1232
Mupad [F(-1)]	1232

Optimal result

Integrand size = 19, antiderivative size = 55

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{4d(b \cos(c+dx))^{4/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(4/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \frac{\sec(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{4/3}}$$

[In] Int[Sec[c + d*x]/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*Sin[c + d*x])/(4*d*(b*Cos[c + d*x])^(4/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{1}{(b \cos(c + dx))^{7/3}} dx \\ &= \frac{3 \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{4d(b \cos(c + dx))^{4/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \frac{3b \cot(c + dx) \text{Hypergeometric2F1}\left(-\frac{2}{3}, \frac{1}{2}, \frac{1}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{4d(b \cos(c + dx))^{7/3}}$$

```
[In] Integrate[Sec[c + d*x]/(b*Cos[c + d*x])^(4/3), x]
```

```
[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*Sqrt[Si
n[c + d*x]^2])/(4*d*(b*Cos[c + d*x])^(7/3))
```

Maple [F]

$$\int \frac{\sec(dx + c)}{(\cos(dx + c)b)^{4/3}} dx$$

```
[In] int(sec(d*x+c)/(cos(d*x+c)*b)^(4/3), x)
```

```
[Out] int(sec(d*x+c)/(cos(d*x+c)*b)^(4/3), x)
```

Fricas [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

```
[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3), x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)/(b^2*cos(d*x + c)^2), x)
```

Sympy [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(sec(c + d*x)/(b*cos(c + d*x))**(4/3), x)

Maxima [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{\cos(c + dx) (b \cos(c + dx))^{4/3}} dx$$

[In] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(4/3)),x)

[Out] int(1/(cos(c + d*x)*(b*cos(c + d*x))^(4/3)), x)

$$3.242 \quad \int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx$$

Optimal result	1233
Rubi [A] (verified)	1233
Mathematica [A] (verified)	1234
Maple [F]	1234
Fricas [F]	1234
Sympy [F]	1235
Maxima [F]	1235
Giac [F]	1235
Mupad [F(-1)]	1235

Optimal result

Integrand size = 21, antiderivative size = 56

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c+dx)\right) \sin(c+dx)}{7d(b \cos(c+dx))^{7/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/7*b*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(7/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^2(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3b \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{7/3}}$$

[In] Int[Sec[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sin[c + d*x])/(7*d*(b*Cos[c + d*x])^(7/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int \frac{1}{(b \cos(c + dx))^{10/3}} dx \\ &= \frac{3b \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c + dx)\right) \sin(c + dx)}{7d(b \cos(c + dx))^{7/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.04

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \frac{3b^2 \cot(c + dx) \operatorname{Hypergeometric2F1}\left(-\frac{7}{6}, \frac{1}{2}, -\frac{1}{6}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{7d(b \cos(c + dx))^{10/3}}$$

```
[In] Integrate[Sec[c + d*x]^2/(b*Cos[c + d*x])^(4/3), x]
```

```
[Out] (3*b^2*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*Sqrt
[Sin[c + d*x]^2])/(7*d*(b*Cos[c + d*x])^(10/3))
```

Maple [F]

$$\int \frac{\sec^2(dx + c)}{(\cos(dx + c)b)^{4/3}} dx$$

```
[In] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(4/3), x)
```

```
[Out] int(sec(d*x+c)^2/(cos(d*x+c)*b)^(4/3), x)
```

Fricas [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

```
[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3), x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^2/(b^2*cos(d*x + c)^2), x)
```

Sympy [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

[In] integrate(sec(d*x+c)**2/(b*cos(d*x+c))**(4/3), x)

[Out] Integral(sec(c + d*x)**2/(b*cos(c + d*x))**(4/3), x)

Maxima [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^2}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)^2/(b*cos(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{\cos(c + dx)^2 (b \cos(c + dx))^{4/3}} dx$$

[In] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3)), x)

[Out] int(1/(cos(c + d*x)^2*(b*cos(c + d*x))^(4/3)), x)

3.243 $\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx$

Optimal result	1236
Rubi [A] (verified)	1236
Mathematica [A] (verified)	1237
Maple [F]	1237
Fricas [F]	1237
Sympy [F]	1238
Maxima [F]	1238
Giac [F]	1238
Mupad [F(-1)]	1238

Optimal result

Integrand size = 21, antiderivative size = 58

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3b^2 \operatorname{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{1}{2}, -\frac{2}{3}, \cos^2(c+dx)\right) \sin(c+dx)}{10d(b \cos(c+dx))^{10/3} \sqrt{\sin^2(c+dx)}}$$

[Out] 3/10*b^2*hypergeom([-5/3, 1/2], [-2/3], cos(d*x+c)^2)*sin(d*x+c)/d/(b*cos(d*x+c))^(10/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {16, 2722}

$$\int \frac{\sec^3(c+dx)}{(b \cos(c+dx))^{4/3}} dx = \frac{3b^2 \sin(c+dx) \operatorname{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{1}{2}, -\frac{2}{3}, \cos^2(c+dx)\right)}{10d \sqrt{\sin^2(c+dx)} (b \cos(c+dx))^{10/3}}$$

[In] Int[Sec[c + d*x]^3/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b^2*Hypergeometric2F1[-5/3, 1/2, -2/3, Cos[c + d*x]^2]*Sin[c + d*x])/(10*d*(b*Cos[c + d*x])^(10/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int \frac{1}{(b \cos(c + dx))^{13/3}} dx \\ &= \frac{3b^2 \text{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{1}{2}, -\frac{2}{3}, \cos^2(c + dx)\right) \sin(c + dx)}{10d(b \cos(c + dx))^{10/3} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \frac{3b^2 \csc(c + dx) \text{Hypergeometric2F1}\left(-\frac{5}{3}, \frac{1}{2}, -\frac{2}{3}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{10d(b \cos(c + dx))^{10/3}}$$

[In] Integrate[Sec[c + d*x]^3/(b*Cos[c + d*x])^(4/3), x]

[Out] (3*b^2*Csc[c + d*x]*Hypergeometric2F1[-5/3, 1/2, -2/3, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(10*d*(b*Cos[c + d*x])^(10/3))

Maple [F]

$$\int \frac{\sec^3(dx + c)}{(\cos(dx + c)b)^{4/3}} dx$$

[In] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(4/3), x)

[Out] int(sec(d*x+c)^3/(cos(d*x+c)*b)^(4/3), x)

Fricas [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^(2/3)*sec(d*x + c)^3/(b^2*cos(d*x + c)^2), x)

Sympy [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx$$

[In] integrate(sec(d*x+c)**3/(b*cos(d*x+c))**(4/3),x)

[Out] Integral(sec(c + d*x)**3/(b*cos(c + d*x))**(4/3), x)

Maxima [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(4/3), x)

Giac [F]

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{\sec(dx + c)^3}{(b \cos(dx + c))^{4/3}} dx$$

[In] integrate(sec(d*x+c)^3/(b*cos(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*cos(d*x + c))^(4/3), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(c + dx)}{(b \cos(c + dx))^{4/3}} dx = \int \frac{1}{\cos(c + dx)^3 (b \cos(c + dx))^{4/3}} dx$$

[In] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(4/3)),x)

[Out] int(1/(cos(c + d*x)^3*(b*cos(c + d*x))^(4/3)), x)

3.244 $\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx$

Optimal result	1239
Rubi [A] (verified)	1239
Mathematica [A] (verified)	1240
Maple [F]	1240
Fricas [F]	1241
Sympy [F]	1241
Maxima [F]	1241
Giac [F]	1241
Mupad [F(-1)]	1242

Optimal result

Integrand size = 21, antiderivative size = 82

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \frac{(a \cos(e + fx))^{1+m} (b \cos(e + fx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), \cos^2(e + fx)\right)}{af(1 + m + n)\sqrt{\sin^2(e + fx)}}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*(b*\cos(f*x+e))^n*\operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}+1/2*m+1/2*n\right], \left[\frac{3}{2}+1/2*m+1/2*n\right], \cos(f*x+e)^2*\sin(f*x+e)/a/f/(1+m+n)/(\sin(f*x+e)^2)^{(1/2)}\right)$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \frac{\sin(e + fx)(a \cos(e + fx))^{m+1} (b \cos(e + fx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(m + n + 1), \frac{1}{2}(m + n + 3), \cos^2(e + fx)\right)}{af(m + n + 1)\sqrt{\sin^2(e + fx)}}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e + f*x])^m*(b*\operatorname{Cos}[e + f*x])^n, x]$

[Out] $-\left(\left(\left(a*\operatorname{Cos}[e + f*x]\right)^{(1 + m)}*(b*\operatorname{Cos}[e + f*x])^n*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1 + m + n)}{2}, \frac{(3 + m + n)}{2}, \operatorname{Cos}[e + f*x]^2*\operatorname{Sin}[e + f*x]\right]/(a*f*(1 + m + n)*\operatorname{Sqrt}[\operatorname{Sin}[e + f*x]^2])\right)\right)$

Rule 20

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[b^{\operatorname{IntPart}[n]}*((b*v)^{\operatorname{FracPart}[n]}/(a^{\operatorname{IntPart}[n]}*(a*v)^{\operatorname{FracPart}[n]})), \operatorname{Int}[u*(a*v)^{(m + n)}$

), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \text{integral} &= ((a \cos(e + fx))^{-n} (b \cos(e + fx))^n) \int (a \cos(e + fx))^{m+n} dx \\ &= \frac{(a \cos(e + fx))^{1+m} (b \cos(e + fx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), \cos^2(e + fx)\right)}{af(1 + m + n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.94

$$\begin{aligned} \int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \\ \frac{(a \cos(e + fx))^m (b \cos(e + fx))^n \cot(e + fx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + m + n), \frac{1}{2}(3 + m + n), \cos^2(e + fx)\right)}{f(1 + m + n)} \end{aligned}$$

[In] Integrate[(a*Cos[e + f*x])^m*(b*Cos[e + f*x])^n,x]

[Out] -(((a*Cos[e + f*x])^m*(b*Cos[e + f*x])^n*Cot[e + f*x]*Hypergeometric2F1[1/2, (1 + m + n)/2, (3 + m + n)/2, Cos[e + f*x]^2]*Sqrt[Sin[e + f*x]^2])/(f*(1 + m + n)))

Maple [F]

$$\int (\cos(fx + e) a)^m (b \cos(fx + e))^n dx$$

[In] int((cos(f*x+e)*a)^m*(b*cos(f*x+e))^n,x)

[Out] int((cos(f*x+e)*a)^m*(b*cos(f*x+e))^n,x)

Fricas [F]

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cos(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

Sympy [F]

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \int (a \cos(e + fx))^m (b \cos(e + fx))^n dx$$

[In] integrate((a*cos(f*x+e))**m*(b*cos(f*x+e))**n,x)

[Out] Integral((a*cos(e + f*x))**m*(b*cos(e + f*x))**n, x)

Maxima [F]

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cos(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

Giac [F]

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \int (a \cos(fx + e))^m (b \cos(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*cos(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*cos(f*x + e))^n, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \cos(e + fx))^n dx = \int (a \cos(e + fx))^m (b \cos(e + fx))^n dx$$

```
[In] int((a*cos(e + f*x))^m*(b*cos(e + f*x))^n,x)
```

```
[Out] int((a*cos(e + f*x))^m*(b*cos(e + f*x))^n, x)
```

3.245 $\int \cos^2(c + dx)(b \cos(c + dx))^n dx$

Optimal result	1243
Rubi [A] (verified)	1243
Mathematica [A] (verified)	1244
Maple [F]	1244
Fricas [F]	1245
Sympy [F]	1245
Maxima [F]	1245
Giac [F]	1245
Mupad [F(-1)]	1246

Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx$$

$$= -\frac{(b \cos(c + dx))^{3+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+n}{2}, \frac{5+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{b^3 d(3+n) \sqrt{\sin^2(c + dx)}}$$

[Out] $-(b \cos(d*x+c))^{(3+n)} \operatorname{hypergeom}\left(\left[\frac{1}{2}, 3/2+1/2*n\right], \left[5/2+1/2*n\right], \cos(d*x+c)^2\right) * \sin(d*x+c) / b^3 d / (3+n) / (\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx$$

$$= -\frac{\sin(c + dx)(b \cos(c + dx))^{n+3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n+3}{2}, \frac{n+5}{2}, \cos^2(c + dx)\right)}{b^3 d(n+3) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^2 * (b * \operatorname{Cos}[c + d*x])^n, x]$

[Out] $-(((b * \operatorname{Cos}[c + d*x])^{(3 + n)} * \operatorname{Hypergeometric2F1}[1/2, (3 + n)/2, (5 + n)/2, \operatorname{Cos}[c + d*x]^2] * \operatorname{Sin}[c + d*x]) / (b^3 * d * (3 + n) * \operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2]))$

Rule 16

$\operatorname{Int}[(u_.) * (v_.)^{(m_.)} * ((b_.) * (v_))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u * (b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}[\{b, n\}, x] \ \&\& \ \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{2+n} dx}{b^2} \\ &= -\frac{(b \cos(c + dx))^{3+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+n}{2}, \frac{5+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{b^3 d(3+n) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.04

$$\int \cos^2(c + dx) (b \cos(c + dx))^n dx = \frac{\cos^2(c + dx) (b \cos(c + dx))^n \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{3+n}{2}, \frac{5+n}{2}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(3+n)}$$

```
[In] Integrate[Cos[c + d*x]^2*(b*Cos[c + d*x])^n,x]
```

```
[Out] -((Cos[c + d*x]^2*(b*Cos[c + d*x])^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (3
+ n)/2, (5 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(3 + n)))
```

Maple [F]

$$\int (\cos^2(dx + c)) (\cos(dx + c) b)^n dx$$

```
[In] int(cos(d*x+c)^2*(cos(d*x+c)*b)^n,x)
```

```
[Out] int(cos(d*x+c)^2*(cos(d*x+c)*b)^n,x)
```

Fricas [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

Sympy [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(c + dx))^n \cos^2(c + dx) dx$$

[In] integrate(cos(d*x+c)**2*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*cos(c + d*x)**2, x)

Maxima [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

Giac [F]

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^2 dx$$

[In] integrate(cos(d*x+c)^2*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(c + dx)(b \cos(c + dx))^n dx = \int \cos(c + dx)^2 (b \cos(c + dx))^n dx$$

```
[In] int(cos(c + d*x)^2*(b*cos(c + d*x))^n,x)
```

```
[Out] int(cos(c + d*x)^2*(b*cos(c + d*x))^n, x)
```

3.246 $\int \cos(c + dx)(b \cos(c + dx))^n dx$

Optimal result	1247
Rubi [A] (verified)	1247
Mathematica [A] (verified)	1248
Maple [F]	1248
Fricas [F]	1249
Sympy [F]	1249
Maxima [F]	1249
Giac [F]	1249
Mupad [F(-1)]	1250

Optimal result

Integrand size = 17, antiderivative size = 69

$$\int \cos(c + dx)(b \cos(c + dx))^n dx$$

$$= -\frac{(b \cos(c + dx))^{2+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+n}{2}, \frac{4+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{b^2 d(2+n) \sqrt{\sin^2(c + dx)}}$$

[Out] $-(b \cos(d*x+c))^{(2+n)} \operatorname{hypergeom}\left(\left[\frac{1}{2}, 1+1/2*n\right], [2+1/2*n], \cos(d*x+c)^2\right) \sin(d*x+c) / b^2 d / (2+n) / (\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {16, 2722}

$$\int \cos(c + dx)(b \cos(c + dx))^n dx$$

$$= -\frac{\sin(c + dx)(b \cos(c + dx))^{n+2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n+2}{2}, \frac{n+4}{2}, \cos^2(c + dx)\right)}{b^2 d(n+2) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x] * (b * \operatorname{Cos}[c + d*x])^n, x]$

[Out] $-\left(\left(b * \operatorname{Cos}[c + d*x]\right)^{(2+n)} \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(2+n)}{2}, \frac{(4+n)}{2}, \operatorname{Cos}[c + d*x]^2\right] * \operatorname{Sin}[c + d*x]\right) / \left(b^2 * d * (2+n) * \operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2]\right)$

Rule 16

$\operatorname{Int}[(u_.) * (v_.)^{(m_.)} * ((b_.) * (v_))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/b^m, \operatorname{Int}[u * (b*v)^{(m+n)}, x], x] /; \operatorname{FreeQ}\{b, n\}, x \ \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (b \cos(c + dx))^{1+n} dx}{b} \\ &= -\frac{(b \cos(c + dx))^{2+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+n}{2}, \frac{4+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{b^2 d(2+n) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \frac{\cos(c + dx)(b \cos(c + dx))^n \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{2+n}{2}, \frac{4+n}{2}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(2+n)}$$

```
[In] Integrate[Cos[c + d*x]*(b*Cos[c + d*x])^n,x]
```

```
[Out] -((Cos[c + d*x]*(b*Cos[c + d*x])^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (2 +
n)/2, (4 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(2 + n)))
```

Maple [F]

$$\int \cos(dx + c) (\cos(dx + c) b)^n dx$$

```
[In] int(cos(d*x+c)*(cos(d*x+c)*b)^n,x)
```

```
[Out] int(cos(d*x+c)*(cos(d*x+c)*b)^n,x)
```


Fricas [F]

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c), x)

Sympy [F]

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(c + dx))^n \cos(c + dx) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*cos(c + d*x), x)

Maxima [F]

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c), x)

Giac [F]

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c) dx$$

[In] integrate(cos(d*x+c)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int \cos(c + dx)(b \cos(c + dx))^n dx = \int \cos(c + dx) (b \cos(c + dx))^n dx$$

```
[In] int(cos(c + d*x)*(b*cos(c + d*x))^n,x)
```

```
[Out] int(cos(c + d*x)*(b*cos(c + d*x))^n, x)
```

3.247 $\int (b \cos(c + dx))^n dx$

Optimal result	1251
Rubi [A] (verified)	1251
Mathematica [A] (verified)	1252
Maple [F]	1252
Fricas [F]	1252
Sympy [F]	1253
Maxima [F]	1253
Giac [F]	1253
Mupad [F(-1)]	1253

Optimal result

Integrand size = 10, antiderivative size = 69

$$\int (b \cos(c + dx))^n dx$$

$$= -\frac{(b \cos(c + dx))^{1+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{bd(1+n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-(b*\cos(d*x+c))^{(1+n)}*\operatorname{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/b/d/(1+n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\int (b \cos(c + dx))^n dx$$

$$= -\frac{\sin(c + dx)(b \cos(c + dx))^{n+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n+1}{2}, \frac{n+3}{2}, \cos^2(c + dx)\right)}{bd(n+1)\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^n, x]$

[Out] $-\left(\left((b*\operatorname{Cos}[c + d*x])^{(1+n)}*\operatorname{Hypergeometric2F1}[1/2, (1+n)/2, (3+n)/2, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x]\right)/(b*d*(1+n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])\right)$

Rule 2722

$\operatorname{Int}[(b_*)*\sin[(c_*) + (d_*)(x_*)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Cos}[c + d*x]*((b*\operatorname{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)*\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]^2])]*\operatorname{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \operatorname{Sin}[c + d*x]^2], x] /;$ $\operatorname{FreeQ}\{b, c, d, n\}, x]$

&& !IntegerQ[2*n]

Rubi steps

$$\text{integral} = -\frac{(b \cos(c + dx))^{1+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{bd(1+n)\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.93

$$\int (b \cos(c + dx))^n dx = \frac{(b \cos(c + dx))^n \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1+n}{2}, \frac{3+n}{2}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(1+n)}$$

[In] Integrate[(b*Cos[c + d*x])^n,x]

[Out] -(((b*Cos[c + d*x])^n*Cot[c + d*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1 + n)))

Maple [F]

$$\int (\cos(dx + c) b)^n dx$$

[In] int((cos(d*x+c)*b)^n,x)

[Out] int((cos(d*x+c)*b)^n,x)

Fricas [F]

$$\int (b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n dx$$

[In] integrate((b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n, x)

Sympy [F]

$$\int (b \cos(c + dx))^n dx = \int (b \cos(c + dx))^n dx$$

[In] integrate((b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n, x)

Maxima [F]

$$\int (b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n dx$$

[In] integrate((b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n, x)

Giac [F]

$$\int (b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n dx$$

[In] integrate((b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n, x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^n dx = \int (b \cos(c + dx))^n dx$$

[In] int((b*cos(c + d*x))^n,x)

[Out] int((b*cos(c + d*x))^n, x)

3.248 $\int (b \cos(c + dx))^n \sec(c + dx) dx$

Optimal result	1254
Rubi [A] (verified)	1254
Mathematica [A] (verified)	1255
Maple [F]	1255
Fricas [F]	1256
Sympy [F]	1256
Maxima [F]	1256
Giac [F]	1256
Mupad [F(-1)]	1257

Optimal result

Integrand size = 17, antiderivative size = 60

$$\int (b \cos(c + dx))^n \sec(c + dx) dx$$

$$= -\frac{(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{dn \sqrt{\sin^2(c + dx)}}$$

[Out] $-(b \cos(dx+c))^n \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}n\right], \left[1+\frac{1}{2}n\right], \cos(dx+c)^2\right) \sin(dx+c)$
 $/d/n/(\sin(dx+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^n \sec(c + dx) dx$$

$$= -\frac{\sin(c + dx) (b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{n+2}{2}, \cos^2(c + dx)\right)}{dn \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b \operatorname{Cos}[c + d*x])^n \operatorname{Sec}[c + d*x], x]$

[Out] $-\left(\left(b \operatorname{Cos}[c + d*x]\right)^n \operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{n}{2}, \frac{(2 + n)}{2}, \operatorname{Cos}[c + d*x]^2\right] \operatorname{Sin}[c + d*x]\right) / \left(d * n * \operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2]\right)$

Rule 16

$\operatorname{Int}[(u_*) \cdot (v_*)^{(m_*)} \cdot ((b_*) \cdot (v_*))^{(n_*)}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u \cdot (b \cdot v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b \int (b \cos(c + dx))^{-1+n} dx \\ &= -\frac{(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{dn \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.05

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \frac{b(b \cos(c + dx))^{-1+n} \cot(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n}{2}, \frac{2+n}{2}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{dn}$$

[In] Integrate[(b*Cos[c + d*x])^n*Sec[c + d*x],x]

[Out] -((b*(b*Cos[c + d*x])^(-1 + n)*Cot[c + d*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*n))

Maple [F]

$$\int (\cos(dx + c) b)^n \sec(dx + c) dx$$

[In] int((cos(d*x+c)*b)^n*sec(d*x+c),x)

[Out] int((cos(d*x+c)*b)^n*sec(d*x+c),x)

Fricas [F]

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sec(d*x + c), x)

Sympy [F]

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \int (b \cos(c + dx))^n \sec(c + dx) dx$$

[In] integrate((b*cos(d*x+c))**n*sec(d*x+c),x)

[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x), x)

Maxima [F]

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c), x)

Giac [F]

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c) dx$$

[In] integrate((b*cos(d*x+c))^n*sec(d*x+c),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c), x)

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^n \sec(c + dx) dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x),x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x), x)
```

3.249 $\int (b \cos(c + dx))^n \sec^2(c + dx) dx$

Optimal result	1258
Rubi [A] (verified)	1258
Mathematica [A] (verified)	1259
Maple [F]	1259
Fricas [F]	1260
Sympy [F]	1260
Maxima [F]	1260
Giac [F]	1260
Mupad [F(-1)]	1261

Optimal result

Integrand size = 19, antiderivative size = 68

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx$$

$$= \frac{b(b \cos(c + dx))^{-1+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-1 + n), \frac{1+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{d(1 - n) \sqrt{\sin^2(c + dx)}}$$

[Out] b*(b*cos(d*x+c))⁽⁻¹⁺ⁿ⁾*hypergeom([1/2, -1/2+1/2*n], [1/2+1/2*n], cos(d*x+c)²)*sin(d*x+c)/d/(1-n)/(sin(d*x+c)²)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx$$

$$= \frac{b \sin(c + dx) (b \cos(c + dx))^{n-1} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n-1}{2}, \frac{n+1}{2}, \cos^2(c + dx)\right)}{d(1 - n) \sqrt{\sin^2(c + dx)}}$$

[In] Int[(b*Cos[c + d*x])ⁿ*Sec[c + d*x]²,x]

[Out] (b*(b*Cos[c + d*x])^(-1 + n)*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Cos[c + d*x]²]*Sin[c + d*x])/(d*(1 - n)*Sqrt[Sin[c + d*x]²])

Rule 16

Int[(u_.)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^2 \int (b \cos(c + dx))^{-2+n} dx \\ &= \frac{b(b \cos(c + dx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-1 + n), \frac{1+n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{d(1 - n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.99

$$\begin{aligned} \int (b \cos(c + dx))^n \sec^2(c + dx) dx = \\ \frac{b(b \cos(c + dx))^{-1+n} \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-1 + n), \frac{1+n}{2}, \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d(-1 + n)} \end{aligned}$$

[In] Integrate[(b*cos[c + d*x])^n*Sec[c + d*x]^2,x]

[Out] -((b*(b*cos[c + d*x])^(-1 + n)*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-1 + n)))

Maple [F]

$$\int (\cos(dx + c) b)^n (\sec^2(dx + c)) dx$$

[In] int((cos(d*x+c)*b)^n*sec(d*x+c)^2,x)

[Out] int((cos(d*x+c)*b)^n*sec(d*x+c)^2,x)

Fricas [F]

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^2 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^2, x)
```

Sympy [F]

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx = \int (b \cos(c + dx))^n \sec^2(c + dx) dx$$

```
[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**2,x)
```

```
[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x)**2, x)
```

Maxima [F]

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^2 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^2, x)
```

Giac [F]

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^2 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^2,x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^n \sec^2(c + dx) dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^2} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^2,x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^2, x)
```

3.250 $\int (b \cos(c + dx))^n \sec^3(c + dx) dx$

Optimal result	1262
Rubi [A] (verified)	1262
Mathematica [A] (verified)	1263
Maple [F]	1263
Fricas [F]	1264
Sympy [F]	1264
Maxima [F]	1264
Giac [F]	1264
Mupad [F(-1)]	1265

Optimal result

Integrand size = 19, antiderivative size = 68

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx$$

$$= \frac{b^2 (b \cos(c + dx))^{-2+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2 + n), \frac{n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{d(2 - n) \sqrt{\sin^2(c + dx)}}$$

[Out] $b^2*(b*\cos(d*x+c))^{(-2+n)}*\operatorname{hypergeom}([1/2, -1+1/2*n], [1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(2-n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx$$

$$= \frac{b^2 \sin(c + dx) (b \cos(c + dx))^{n-2} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n-2}{2}, \frac{n}{2}, \cos^2(c + dx)\right)}{d(2 - n) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^n*\operatorname{Sec}[c + d*x]^3,x]$

[Out] $(b^2*(b*\operatorname{Cos}[c + d*x])^{(-2 + n)}*\operatorname{Hypergeometric2F1}[1/2, (-2 + n)/2, n/2, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/d*(2 - n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= b^3 \int (b \cos(c + dx))^{-3+n} dx \\ &= \frac{b^2 (b \cos(c + dx))^{-2+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2 + n), \frac{n}{2}, \cos^2(c + dx)\right) \sin(c + dx)}{d(2 - n) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.03

$$\begin{aligned} \int (b \cos(c + dx))^n \sec^3(c + dx) dx = \\ \frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-2 + n), \frac{n}{2}, \cos^2(c + dx)\right) \sec^2(c + dx) \sqrt{\sin^2(c + dx)}}{d(-2 + n)} \end{aligned}$$

```
[In] Integrate[(b*Cos[c + d*x])^n*Sec[c + d*x]^3,x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-2 + n)/2, n/2,
Cos[c + d*x]^2]*Sec[c + d*x]^2*Sqrt[Sin[c + d*x]^2])/(d*(-2 + n)))
```

Maple [F]

$$\int (\cos(dx + c) b)^n (\sec^3(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^n*sec(d*x+c)^3,x)
```

```
[Out] int((cos(d*x+c)*b)^n*sec(d*x+c)^3,x)
```

Fricas [F]

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^3 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^3, x)
```

Sympy [F]

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx = \int (b \cos(c + dx))^n \sec^3(c + dx) dx$$

```
[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**3,x)
```

```
[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x)**3, x)
```

Maxima [F]

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^3 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^3, x)
```

Giac [F]

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^3 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^3,x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^3, x)
```


Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^n \sec^3(c + dx) dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^3} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^3,x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^3, x)
```

3.251 $\int (b \cos(c + dx))^n \sec^4(c + dx) dx$

Optimal result	1266
Rubi [A] (verified)	1266
Mathematica [A] (verified)	1267
Maple [F]	1267
Fricas [F]	1268
Sympy [F]	1268
Maxima [F]	1268
Giac [F]	1268
Mupad [F(-1)]	1269

Optimal result

Integrand size = 19, antiderivative size = 70

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx$$

$$= \frac{b^3 (b \cos(c + dx))^{-3+n} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-3 + n), \frac{1}{2}(-1 + n), \cos^2(c + dx)\right) \sin(c + dx)}{d(3 - n) \sqrt{\sin^2(c + dx)}}$$

[Out] $b^3*(b*\cos(d*x+c))^{(-3+n)}*\operatorname{hypergeom}([1/2, -3/2+1/2*n], [-1/2+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(3-n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {16, 2722}

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx$$

$$= \frac{b^3 \sin(c + dx) (b \cos(c + dx))^{n-3} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{n-3}{2}, \frac{n-1}{2}, \cos^2(c + dx)\right)}{d(3 - n) \sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c + d*x])^n*\operatorname{Sec}[c + d*x]^4,x]$

[Out] $(b^3*(b*\operatorname{Cos}[c + d*x])^{(-3 + n)}*\operatorname{Hypergeometric2F1}[1/2, (-3 + n)/2, (-1 + n)/2, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(3 - n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 16

$\operatorname{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] := \operatorname{Dist}[1/b^m, \operatorname{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{b, n\}, x \&\& \operatorname{IntegerQ}[m]$

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\text{integral} = b^4 \int (b \cos(c + dx))^{-4+n} dx$$

$$= \frac{b^3 (b \cos(c + dx))^{-3+n} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-3 + n), \frac{1}{2}(-1 + n), \cos^2(c + dx)\right) \sin(c + dx)}{d(3 - n)\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.03

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx =$$

$$\frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(-3 + n), \frac{1}{2}(-1 + n), \cos^2(c + dx)\right) \sec^3(c + dx)}{d(-3 + n)}$$

```
[In] Integrate[(b*Cos[c + d*x])^n*Sec[c + d*x]^4,x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-3 + n)/2, (-1 +
n)/2, Cos[c + d*x]^2]*Sec[c + d*x]^3*Sqrt[Sin[c + d*x]^2])/(d*(-3 + n)))
```

Maple [F]

$$\int (\cos(dx + c) b)^n (\sec^4(dx + c)) dx$$

```
[In] int((cos(d*x+c)*b)^n*sec(d*x+c)^4,x)
```

```
[Out] int((cos(d*x+c)*b)^n*sec(d*x+c)^4,x)
```

Fricas [F]

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^4 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="fricas")
```

```
[Out] integral((b*cos(d*x + c))^n*sec(d*x + c)^4, x)
```

Sympy [F]

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx = \int (b \cos(c + dx))^n \sec^4(c + dx) dx$$

```
[In] integrate((b*cos(d*x+c))**n*sec(d*x+c)**4,x)
```

```
[Out] Integral((b*cos(c + d*x))**n*sec(c + d*x)**4, x)
```

Maxima [F]

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^4 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="maxima")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^4, x)
```

Giac [F]

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx = \int (b \cos(dx + c))^n \sec(dx + c)^4 dx$$

```
[In] integrate((b*cos(d*x+c))^n*sec(d*x+c)^4,x, algorithm="giac")
```

```
[Out] integrate((b*cos(d*x + c))^n*sec(d*x + c)^4, x)
```

Mupad [F(-1)]

Timed out.

$$\int (b \cos(c + dx))^n \sec^4(c + dx) dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^4} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^4,x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^4, x)
```

3.252 $\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx$

Optimal result	1270
Rubi [A] (verified)	1270
Mathematica [A] (verified)	1271
Maple [F]	1271
Fricas [F]	1272
Sympy [F(-1)]	1272
Maxima [F]	1272
Giac [F]	1272
Mupad [F(-1)]	1273

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{2 \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(7 + 2n), \frac{1}{4}(11 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(7 + 2n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(7/2)}*(b*\cos(d*x+c))^n*\operatorname{hypergeom}([1/2, 7/4+1/2*n], [11/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(7+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{2 \sin(c + dx) \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n + 7), \frac{1}{4}(2n + 11), \cos^2(c + dx)\right)}{d(2n + 7)\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^{(5/2)}*(b*\operatorname{Cos}[c + d*x])^n, x]$

[Out] $(-2*\operatorname{Cos}[c + d*x]^{(7/2)}*(b*\operatorname{Cos}[c + d*x])^n*\operatorname{Hypergeometric2F1}[1/2, (7 + 2*n)/4, (11 + 2*n)/4, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(7 + 2*n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{\frac{5}{2}+n}(c + dx) dx \\ &= \frac{2 \cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(7 + 2n), \frac{1}{4}(11 + 2n), \cos^2(c + dx)\right) \sin}{d(7 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{\cos^{\frac{7}{2}}(c + dx)(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{7}{2} + n\right), \frac{1}{2}\left(\frac{11}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(\frac{7}{2} + n\right)}$$

```
[In] Integrate[Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^n,x]
```

```
[Out] -((Cos[c + d*x]^(7/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2
, (7/2 + n)/2, (11/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(7/2
+ n)))
```

Maple [F]

$$\int \left(\cos^{\frac{5}{2}}(dx + c) \right) (\cos(dx + c) b)^n dx$$

```
[In] int(cos(d*x+c)^(5/2)*(cos(d*x+c)*b)^n,x)
```

```
[Out] int(cos(d*x+c)^(5/2)*(cos(d*x+c)*b)^n,x)
```

Fricas [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}} dx$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

Sympy [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \text{Timed out}$$

[In] integrate(cos(d*x+c)**(5/2)*(b*cos(d*x+c))**n,x)

[Out] Timed out

Maxima [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}} dx$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

Giac [F]

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{5}{2}} dx$$

[In] integrate(cos(d*x+c)^(5/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n dx = \int \cos(c + dx)^{5/2} (b \cos(c + dx))^n dx$$

```
[In] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^n, x)
```

```
[Out] int(cos(c + d*x)^(5/2)*(b*cos(c + d*x))^n, x)
```

3.253 $\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx$

Optimal result	1274
Rubi [A] (verified)	1274
Mathematica [A] (verified)	1275
Maple [F]	1275
Fricas [F]	1276
Sympy [F]	1276
Maxima [F]	1276
Giac [F]	1276
Mupad [F(-1)]	1277

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{2 \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(5 + 2n), \frac{1}{4}(9 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(5 + 2n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(5/2)}*(b*\cos(d*x+c))^n*\operatorname{hypergeom}([1/2, 5/4+1/2*n], [9/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(5+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{2 \sin(c + dx) \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n + 5), \frac{1}{4}(2n + 9), \cos^2(c + dx)\right)}{d(2n + 5)\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Cos}[c + d*x]^{(3/2)}*(b*\operatorname{Cos}[c + d*x])^n, x]$

[Out] $(-2*\operatorname{Cos}[c + d*x]^{(5/2)}*(b*\operatorname{Cos}[c + d*x])^n*\operatorname{Hypergeometric2F1}[1/2, (5 + 2*n)/4, (9 + 2*n)/4, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(5 + 2*n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{\frac{3}{2}+n}(c + dx) dx \\ &= \frac{2 \cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(5 + 2n), \frac{1}{4}(9 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(5 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \frac{\cos^{\frac{5}{2}}(c + dx)(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{5}{2} + n\right), \frac{1}{2}\left(\frac{9}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(\frac{5}{2} + n\right)}$$

```
[In] Integrate[Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^n,x]
```

```
[Out] -((Cos[c + d*x]^(5/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2
, (5/2 + n)/2, (9/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(5/2 +
n)))
```

Maple [F]

$$\int \left(\cos^{\frac{3}{2}}(dx + c) \right) (\cos(dx + c) b)^n dx$$

```
[In] int(cos(d*x+c)^(3/2)*(cos(d*x+c)*b)^n,x)
```

```
[Out] int(cos(d*x+c)^(3/2)*(cos(d*x+c)*b)^n,x)
```

Fricas [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}} dx$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

Sympy [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(c + dx))^n \cos^{\frac{3}{2}}(c + dx) dx$$

[In] integrate(cos(d*x+c)**(3/2)*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*cos(c + d*x)**(3/2), x)

Maxima [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}} dx$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

Giac [F]

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \int (b \cos(dx + c))^n \cos(dx + c)^{\frac{3}{2}} dx$$

[In] integrate(cos(d*x+c)^(3/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*cos(d*x + c)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n dx = \int \cos(c + dx)^{3/2} (b \cos(c + dx))^n dx$$

```
[In] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^n, x)
```

```
[Out] int(cos(c + d*x)^(3/2)*(b*cos(c + d*x))^n, x)
```

3.254 $\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^n dx$

Optimal result	1278
Rubi [A] (verified)	1278
Mathematica [A] (verified)	1279
Maple [F]	1279
Fricas [F]	1280
Sympy [F]	1280
Maxima [F]	1280
Giac [F]	1280
Mupad [F(-1)]	1281

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^n dx = \frac{2 \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(3 + 2n), \frac{1}{4}(7 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(3 + 2n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-2*\cos(d*x+c)^{(3/2)}*(b*\cos(d*x+c))^n*\operatorname{hypergeom}([1/2, 3/4+1/2*n], [7/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)/d/(3+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^n dx = \frac{2 \sin(c + dx) \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n + 3), \frac{1}{4}(2n + 7), \cos^2(c + dx)\right)}{d(2n + 3)\sqrt{\sin^2(c + dx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[\operatorname{Cos}[c + d*x]]*(b*\operatorname{Cos}[c + d*x])^n, x]$

[Out] $(-2*\operatorname{Cos}[c + d*x]^{(3/2)}*(b*\operatorname{Cos}[c + d*x])^n*\operatorname{Hypergeometric2F1}[1/2, (3 + 2*n)/4, (7 + 2*n)/4, \operatorname{Cos}[c + d*x]^2]*\operatorname{Sin}[c + d*x])/(d*(3 + 2*n)*\operatorname{Sqrt}[\operatorname{Sin}[c + d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{\frac{1}{2}+n}(c + dx) dx \\ &= \frac{2 \cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(3 + 2n), \frac{1}{4}(7 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(3 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \sqrt{\cos(c + dx)}(b \cos(c + dx))^n dx = \frac{\cos^{\frac{3}{2}}(c + dx)(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{3}{2} + n\right), \frac{1}{2}\left(\frac{7}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\cos(c + dx)}}{d\left(\frac{3}{2} + n\right)}$$

```
[In] Integrate[Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^n,x]
```

```
[Out] -((Cos[c + d*x]^(3/2)*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2
, (3/2 + n)/2, (7/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(3/2 +
n)))
```

Maple [F]

$$\int (\sqrt{\cos(dx + c)}) (\cos(dx + c)b)^n dx$$

```
[In] int(cos(d*x+c)^(1/2)*(cos(d*x+c)*b)^n,x)
```

```
[Out] int(cos(d*x+c)^(1/2)*(cos(d*x+c)*b)^n,x)
```

Fricas [F]

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^n dx = \int (b \cos(dx+c))^n \sqrt{\cos(dx+c)} dx$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

Sympy [F]

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^n dx = \int (b \cos(c+dx))^n \sqrt{\cos(c+dx)} dx$$

[In] integrate(cos(d*x+c)**(1/2)*(b*cos(d*x+c))**n,x)

[Out] Integral((b*cos(c + d*x))**n*sqrt(cos(c + d*x)), x)

Maxima [F]

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^n dx = \int (b \cos(dx+c))^n \sqrt{\cos(dx+c)} dx$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

Giac [F]

$$\int \sqrt{\cos(c+dx)}(b \cos(c+dx))^n dx = \int (b \cos(dx+c))^n \sqrt{\cos(dx+c)} dx$$

[In] integrate(cos(d*x+c)^(1/2)*(b*cos(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n*sqrt(cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx = \int \sqrt{\cos(c + dx)} (b \cos(c + dx))^n dx$$

```
[In] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^n,x)
```

```
[Out] int(cos(c + d*x)^(1/2)*(b*cos(c + d*x))^n, x)
```

3.255 $\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx$

Optimal result	1282
Rubi [A] (verified)	1282
Mathematica [A] (verified)	1283
Maple [F]	1283
Fricas [F]	1284
Sympy [F]	1284
Maxima [F]	1284
Giac [F]	1284
Mupad [F(-1)]	1285

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx = \frac{2\sqrt{\cos(c+dx)}(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(1+2n), \frac{1}{4}(5+2n), \cos^2(c+dx)\right) \sin(c+dx)}{d(1+2n)\sqrt{\sin^2(c+dx)}}$$

[Out] $-2*(b*\cos(d*x+c))^n*\operatorname{hypergeom}([1/2, 1/4+1/2*n], [5/4+1/2*n], \cos(d*x+c)^2)*\sin(d*x+c)*\cos(d*x+c)^{(1/2)}/d/(1+2*n)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{(b \cos(c+dx))^n}{\sqrt{\cos(c+dx)}} dx = \frac{2 \sin(c+dx) \sqrt{\cos(c+dx)} (b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n+1), \frac{1}{4}(2n+5), \cos^2(c+dx)\right)}{d(2n+1)\sqrt{\sin^2(c+dx)}}$$

[In] $\operatorname{Int}[(b*\operatorname{Cos}[c+d*x])^n/\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]],x]$

[Out] $(-2*\operatorname{Sqrt}[\operatorname{Cos}[c+d*x]]*(b*\operatorname{Cos}[c+d*x])^n*\operatorname{Hypergeometric2F1}[1/2, (1+2*n)/4, (5+2*n)/4, \operatorname{Cos}[c+d*x]^2]*\operatorname{Sin}[c+d*x])/d*(1+2*n)*\operatorname{Sqrt}[\operatorname{Sin}[c+d*x]^2])$

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{-\frac{1}{2}+n}(c + dx) dx \\ &= \frac{2\sqrt{\cos(c + dx)}(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(1 + 2n), \frac{1}{4}(5 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 + 2n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \frac{\sqrt{\cos(c + dx)}(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(\frac{1}{2} + n\right), \frac{1}{2}\left(\frac{5}{2} + n\right), \cos^2(c + dx)\right)}{d\left(\frac{1}{2} + n\right)}$$

```
[In] Integrate[(b*Cos[c + d*x])^n/Sqrt[Cos[c + d*x]], x]
```

```
[Out] -((Sqrt[Cos[c + d*x]]*(b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2
, (1/2 + n)/2, (5/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(1/2 +
n)))
```

Maple [F]

$$\int \frac{(\cos(dx + c)b)^n}{\sqrt{\cos(dx + c)}} dx$$

```
[In] int((cos(d*x+c)*b)^n/cos(d*x+c)^(1/2), x)
```

```
[Out] int((cos(d*x+c)*b)^n/cos(d*x+c)^(1/2), x)
```

Fricas [F]

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

Sympy [F]

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx$$

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(1/2),x)

[Out] Integral((b*cos(c + d*x))**n/sqrt(cos(c + d*x)), x)

Maxima [F]

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

Giac [F]

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(dx + c))^n}{\sqrt{\cos(dx + c)}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/sqrt(cos(d*x + c)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx = \int \frac{(b \cos(c + dx))^n}{\sqrt{\cos(c + dx)}} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(1/2), x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(1/2), x)
```

$$3.256 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{3}{2}}(c+dx)} dx$$

Optimal result	1286
Rubi [A] (verified)	1286
Mathematica [A] (verified)	1287
Maple [F]	1287
Fricas [F]	1288
Sympy [F]	1288
Maxima [F]	1288
Giac [F]	1288
Mupad [F(-1)]	1289

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{2(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-1+2n), \frac{1}{4}(3+2n), \cos^2(c+dx)\right) \sin(c+dx)}{d(1-2n)\sqrt{\cos(c+dx)}\sqrt{\sin^2(c+dx)}}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -1/4+1/2*n], [3/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(1-2*n)/cos(d*x+c)^(1/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{3}{2}}(c+dx)} dx$$

$$= \frac{2 \sin(c+dx)(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n-1), \frac{1}{4}(2n+3), \cos^2(c+dx)\right)}{d(1-2n)\sqrt{\sin^2(c+dx)}\sqrt{\cos(c+dx)}}$$

[In] Int[(b*Cos[c + d*x])^n/Cos[c + d*x]^(3/2),x]

[Out] (2*(b*Cos[c + d*x])^n*Hypergeometric2F1[1/2, (-1 + 2*n)/4, (3 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(1 - 2*n)*Sqrt[Cos[c + d*x]]*Sqrt[Sin[c + d*x]^2])

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{-\frac{3}{2}+n}(c + dx) dx \\ &= \frac{2(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-1 + 2n), \frac{1}{4}(3 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(1 - 2n) \sqrt{\cos(c + dx)} \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{1}{2} + n\right), \frac{1}{2}\left(\frac{3}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(-\frac{1}{2} + n\right) \sqrt{\cos(c + dx)}}$$

```
[In] Integrate[(b*Cos[c + d*x])^n/Cos[c + d*x]^(3/2), x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/2 + n)/2, (3/
2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-1/2 + n)*Sqrt[Cos[c +
d*x]]))
```

Maple [F]

$$\int \frac{(\cos(dx + c)b)^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

```
[In] int((cos(d*x+c)*b)^n/cos(d*x+c)^(3/2), x)
```

```
[Out] int((cos(d*x+c)*b)^n/cos(d*x+c)^(3/2), x)
```

Fricas [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

Sympy [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx$$

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(3/2),x)

[Out] Integral((b*cos(c + d*x))**n/cos(c + d*x)**(3/2), x)

Maxima [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

Giac [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{3}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{3}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{3/2}} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(3/2), x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(3/2), x)
```

$$3.257 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{5}{2}}(c+dx)} dx$$

Optimal result	1290
Rubi [A] (verified)	1290
Mathematica [A] (verified)	1291
Maple [F]	1291
Fricas [F]	1292
Sympy [F]	1292
Maxima [F]	1292
Giac [F]	1292
Mupad [F(-1)]	1293

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{2(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-3+2n), \frac{1}{4}(1+2n), \cos^2(c+dx)\right) \sin(c+dx)}{d(3-2n) \cos^{\frac{3}{2}}(c+dx) \sqrt{\sin^2(c+dx)}}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -3/4+1/2*n], [1/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(3-2*n)/cos(d*x+c)^(3/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{5}{2}}(c+dx)} dx$$

$$= \frac{2 \sin(c+dx) (b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n-3), \frac{1}{4}(2n+1), \cos^2(c+dx)\right)}{d(3-2n) \sqrt{\sin^2(c+dx)} \cos^{\frac{3}{2}}(c+dx)}$$

[In] Int[(b*Cos[c + d*x])^n/Cos[c + d*x]^(5/2),x]

[Out] (2*(b*Cos[c + d*x])^n*Hypergeometric2F1[1/2, (-3 + 2*n)/4, (1 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(3 - 2*n)*Cos[c + d*x]^(3/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{-\frac{5}{2}+n}(c + dx) dx \\ &= \frac{2(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-3 + 2n), \frac{1}{4}(1 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(3 - 2n) \cos^{\frac{3}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{3}{2} + n\right), \frac{1}{2}\left(\frac{1}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(-\frac{3}{2} + n\right) \cos^{\frac{3}{2}}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^n/Cos[c + d*x]^(5/2), x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-3/2 + n)/2, (1/
2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-3/2 + n)*Cos[c + d*x]^
(3/2)))
```

Maple [F]

$$\int \frac{(\cos(dx + c)b)^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

```
[In] int((cos(d*x+c)*b)^n/cos(d*x+c)^(5/2), x)
```

```
[Out] int((cos(d*x+c)*b)^n/cos(d*x+c)^(5/2), x)
```

Fricas [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

Sympy [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx$$

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(5/2),x)

[Out] Integral((b*cos(c + d*x))**n/cos(c + d*x)**(5/2), x)

Maxima [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

Giac [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{5}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{5}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{5/2}} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(5/2), x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(5/2), x)
```

$$3.258 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{7}{2}}(c+dx)} dx$$

Optimal result	1294
Rubi [A] (verified)	1294
Mathematica [A] (verified)	1295
Maple [F]	1295
Fricas [F]	1296
Sympy [F(-1)]	1296
Maxima [F]	1296
Giac [F]	1296
Mupad [F(-1)]	1297

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-5+2n), \frac{1}{4}(-1+2n), \cos^2(c+dx)\right) \sin(c+dx)}{d(5-2n) \cos^{\frac{5}{2}}(c+dx) \sqrt{\sin^2(c+dx)}}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -5/4+1/2*n], [-1/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(5-2*n)/cos(d*x+c)^(5/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{7}{2}}(c+dx)} dx$$

$$= \frac{2 \sin(c+dx) (b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n-5), \frac{1}{4}(2n-1), \cos^2(c+dx)\right)}{d(5-2n) \sqrt{\sin^2(c+dx)} \cos^{\frac{5}{2}}(c+dx)}$$

[In] Int[(b*cos[c + d*x])^n/Cos[c + d*x]^(7/2),x]

[Out] (2*(b*cos[c + d*x])^n*Hypergeometric2F1[1/2, (-5 + 2*n)/4, (-1 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(5 - 2*n)*Cos[c + d*x]^(5/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{-\frac{7}{2}+n}(c + dx) dx \\ &= \frac{2(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-5 + 2n), \frac{1}{4}(-1 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(5 - 2n) \cos^{\frac{5}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{5}{2} + n\right), \frac{1}{2}\left(-\frac{1}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(-\frac{5}{2} + n\right) \cos^{\frac{5}{2}}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^n/Cos[c + d*x]^(7/2), x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-5/2 + n)/2, (-1
/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-5/2 + n)*Cos[c + d*x]
^(5/2)))
```

Maple [F]

$$\int \frac{(\cos(dx + c)b)^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

```
[In] int((cos(d*x+c)*b)^n/cos(d*x+c)^(7/2), x)
```

```
[Out] int((cos(d*x+c)*b)^n/cos(d*x+c)^(7/2), x)
```

Fricas [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(7/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

Giac [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{7}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(7/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{7}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{7/2}} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(7/2), x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(7/2), x)
```

$$3.259 \quad \int \frac{(b \cos(c+dx))^n}{\cos^{\frac{9}{2}}(c+dx)} dx$$

Optimal result	1298
Rubi [A] (verified)	1298
Mathematica [A] (verified)	1299
Maple [F]	1299
Fricas [F]	1300
Sympy [F(-1)]	1300
Maxima [F]	1300
Giac [F]	1300
Mupad [F(-1)]	1301

Optimal result

Integrand size = 21, antiderivative size = 80

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2(b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-7+2n), \frac{1}{4}(-3+2n), \cos^2(c+dx)\right) \sin(c+dx)}{d(7-2n) \cos^{\frac{7}{2}}(c+dx) \sqrt{\sin^2(c+dx)}}$$

[Out] 2*(b*cos(d*x+c))^n*hypergeom([1/2, -7/4+1/2*n], [-3/4+1/2*n], cos(d*x+c)^2)*sin(d*x+c)/d/(7-2*n)/cos(d*x+c)^(7/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\int \frac{(b \cos(c+dx))^n}{\cos^{\frac{9}{2}}(c+dx)} dx$$

$$= \frac{2 \sin(c+dx) (b \cos(c+dx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(2n-7), \frac{1}{4}(2n-3), \cos^2(c+dx)\right)}{d(7-2n) \sqrt{\sin^2(c+dx)} \cos^{\frac{7}{2}}(c+dx)}$$

[In] Int[(b*cos[c + d*x])^n/Cos[c + d*x]^(9/2),x]

[Out] (2*(b*cos[c + d*x])^n*Hypergeometric2F1[1/2, (-7 + 2*n)/4, (-3 + 2*n)/4, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(7 - 2*n)*Cos[c + d*x]^(7/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

```
Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[
n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n
), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]
```

Rule 2722

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((
b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2
F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\cos^{-n}(c + dx)(b \cos(c + dx))^n) \int \cos^{-\frac{9}{2}+n}(c + dx) dx \\ &= \frac{2(b \cos(c + dx))^n \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{4}(-7 + 2n), \frac{1}{4}(-3 + 2n), \cos^2(c + dx)\right) \sin(c + dx)}{d(7 - 2n) \cos^{\frac{7}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \frac{(b \cos(c + dx))^n \csc(c + dx) \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}\left(-\frac{7}{2} + n\right), \frac{1}{2}\left(-\frac{3}{2} + n\right), \cos^2(c + dx)\right) \sqrt{\sin^2(c + dx)}}{d\left(-\frac{7}{2} + n\right) \cos^{\frac{7}{2}}(c + dx)}$$

```
[In] Integrate[(b*Cos[c + d*x])^n/Cos[c + d*x]^(9/2), x]
```

```
[Out] -(((b*Cos[c + d*x])^n*Csc[c + d*x]*Hypergeometric2F1[1/2, (-7/2 + n)/2, (-3
/2 + n)/2, Cos[c + d*x]^2]*Sqrt[Sin[c + d*x]^2])/(d*(-7/2 + n)*Cos[c + d*x]
^(7/2)))
```

Maple [F]

$$\int \frac{(\cos(dx + c)b)^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

```
[In] int((cos(d*x+c)*b)^n/cos(d*x+c)^(9/2), x)
```

```
[Out] int((cos(d*x+c)*b)^n/cos(d*x+c)^(9/2), x)
```

Fricas [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2),x, algorithm="fricas")

[Out] integral((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

Sympy [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \text{Timed out}$$

[In] integrate((b*cos(d*x+c))**n/cos(d*x+c)**(9/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2),x, algorithm="maxima")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

Giac [F]

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \int \frac{(b \cos(dx + c))^n}{\cos(dx + c)^{\frac{9}{2}}} dx$$

[In] integrate((b*cos(d*x+c))^n/cos(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate((b*cos(d*x + c))^n/cos(d*x + c)^(9/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(b \cos(c + dx))^n}{\cos^{\frac{9}{2}}(c + dx)} dx = \int \frac{(b \cos(c + dx))^n}{\cos(c + dx)^{9/2}} dx$$

```
[In] int((b*cos(c + d*x))^n/cos(c + d*x)^(9/2), x)
```

```
[Out] int((b*cos(c + d*x))^n/cos(c + d*x)^(9/2), x)
```

3.260 $\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx$

Optimal result	1302
Rubi [A] (verified)	1302
Mathematica [A] (verified)	1303
Maple [F]	1303
Fricas [F]	1304
Sympy [F]	1304
Maxima [F]	1304
Giac [F]	1304
Mupad [F(-1)]	1305

Optimal result

Integrand size = 21, antiderivative size = 88

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \frac{(a \cos(e + fx))^{1+m} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1+m-n), \frac{1}{2}(3+m-n), \cos^2(e + fx)\right) (b \sec(e + fx))^n}{af(1+m-n)\sqrt{\sin^2(e + fx)}}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*\operatorname{hypergeom}([1/2, 1/2+1/2*m-1/2*n], [3/2+1/2*m-1/2*n], \cos(f*x+e)^2)*(b*\sec(f*x+e))^n*\sin(f*x+e)/a/f/(1+m-n)/(\sin(f*x+e)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2668, 2722}

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \frac{\sin(e + fx)(a \cos(e + fx))^{m+1} (b \sec(e + fx))^n \operatorname{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(m-n+1), \frac{1}{2}(m-n+3), \cos^2(e + fx)\right)}{af(m-n+1)\sqrt{\sin^2(e + fx)}}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e + f*x])^m*(b*\operatorname{Sec}[e + f*x])^n, x]$

[Out] $-\left(\left(a*\operatorname{Cos}[e + f*x]\right)^{(1+m)}*\operatorname{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1+m-n)}{2}, \frac{(3+m-n)}{2}, \operatorname{Cos}[e + f*x]^2\right]*(b*\operatorname{Sec}[e + f*x])^n*\operatorname{Sin}[e + f*x]\right)/\left(a*f*(1+m-n)*\operatorname{Sqrt}[\operatorname{Sin}[e + f*x]^2]\right)$

Rule 2668

$\operatorname{Int}[(\operatorname{csc}[e_.] + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \operatorname{Dist}[(a*b)^{\operatorname{IntPart}[n]}*(a*\sin[e + f*x])^{\operatorname{FracPart}[n]}*(b*\operatorname{Csc}$

$[e + f*x]^{\text{FracPart}[n]}$, $\text{Int}[(a*\text{Sin}[e + f*x])^{(m - n)}, x], x] /;$ $\text{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& \text{!IntegerQ}[m] \&\& \text{!IntegerQ}[n]$

Rule 2722

$\text{Int}[(b_*)*\text{sin}[(c_*) + (d_*)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]))*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2], x] /;$ $\text{FreeQ}\{b, c, d, n\}, x\} \&\& \text{!IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned} \text{integral} &= ((a \cos(e + fx))^n (b \sec(e + fx))^n) \int (a \cos(e + fx))^{m-n} dx \\ &= \frac{(a \cos(e + fx))^{1+m} \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + m - n), \frac{1}{2}(3 + m - n), \cos^2(e + fx)\right) (b \sec(e + fx))^n}{af(1 + m - n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 11.05 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.01

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \frac{\cos(e + fx)(a \cos(e + fx))^m \text{Hypergeometric2F1}\left(\frac{1}{2}, \frac{1}{2}(1 + m - n), \frac{1}{2}(3 + m - n), \cos^2(e + fx)\right) (b \sec(e + fx))^n}{f(1 + m - n)\sqrt{\sin^2(e + fx)}}$$

[In] $\text{Integrate}[(a*\text{Cos}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n,x]$

[Out] $-((\text{Cos}[e + f*x]*(a*\text{Cos}[e + f*x])^m*\text{Hypergeometric2F1}[1/2, (1 + m - n)/2, (3 + m - n)/2, \text{Cos}[e + f*x]^2]*(b*\text{Sec}[e + f*x])^n*\text{Sin}[e + f*x])/(f*(1 + m - n)*\text{Sqrt}[\text{Sin}[e + f*x]^2]))$

Maple [F]

$$\int (\cos(fx + e)a)^m (b \sec(fx + e))^n dx$$

[In] $\text{int}((\cos(f*x+e)*a)^m*(b*\sec(f*x+e))^n,x)$

[Out] $\text{int}((\cos(f*x+e)*a)^m*(b*\sec(f*x+e))^n,x)$

Fricas [F]

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

Sympy [F]

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \int (a \cos(e + fx))^m (b \sec(e + fx))^n dx$$

[In] integrate((a*cos(f*x+e))**m*(b*sec(f*x+e))**n,x)

[Out] Integral((a*cos(e + f*x))**m*(b*sec(e + f*x))**n, x)

Maxima [F]

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

Giac [F]

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \int (a \cos(fx + e))^m (b \sec(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*sec(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*sec(f*x + e))^n, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \sec(e + fx))^n dx = \int (a \cos(e + fx))^m \left(\frac{b}{\cos(e + fx)} \right)^n dx$$

```
[In] int((a*cos(e + f*x))^m*(b/cos(e + f*x))^n,x)
```

```
[Out] int((a*cos(e + f*x))^m*(b/cos(e + f*x))^n, x)
```

3.261 $\int \cos(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal result	1306
Rubi [A] (verified)	1306
Mathematica [A] (verified)	1307
Maple [A] (verified)	1307
Fricas [A] (verification not implemented)	1307
Sympy [F]	1308
Maxima [A] (verification not implemented)	1308
Giac [A] (verification not implemented)	1308
Mupad [B] (verification not implemented)	1308

Optimal result

Integrand size = 17, antiderivative size = 15

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2}{b\sqrt{\csc(a + bx)}}$$

[Out] 2/b/csc(b*x+a)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2701, 30}

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2}{b\sqrt{\csc(a + bx)}}$$

[In] Int[Cos[a + b*x]*Sqrt[Csc[a + b*x]],x]

[Out] 2/(b*Sqrt[Csc[a + b*x]])

Rule 30

Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x^{3/2}} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{2}{b\sqrt{\csc(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cos(a + bx)\sqrt{\csc(a + bx)} dx = \frac{2}{b\sqrt{\csc(a + bx)}}$$

[In] Integrate[Cos[a + b*x]*Sqrt[Csc[a + b*x]],x]

[Out] 2/(b*Sqrt[Csc[a + b*x]])

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

method	result	size
derivativedivides	$\frac{2}{b\sqrt{\csc(bx+a)}}$	14
default	$\frac{2}{b\sqrt{\csc(bx+a)}}$	14
risch	$\frac{2\sqrt{2}\sqrt{\frac{ie^{i(bx+a)}}{e^{2i(bx+a)}-1}}\sin(bx+a)}{b}$	42

[In] int(cos(b*x+a)*csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/b/csc(b*x+a)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cos(a + bx)\sqrt{\csc(a + bx)} dx = \frac{2\sqrt{\sin(bx + a)}}{b}$$

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] 2*sqrt(sin(b*x + a))/b

Sympy [F]

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos(a + bx) \sqrt{\csc(a + bx)} dx$$

[In] integrate(cos(b*x+a)*csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)*sqrt(csc(a + b*x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2 \sqrt{\sin(bx + a)}}{b}$$

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(sin(b*x + a))/b

Giac [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2 \sqrt{\sin(bx + a)}}{b}$$

[In] integrate(cos(b*x+a)*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] 2*sqrt(sin(b*x + a))/b

Mupad [B] (verification not implemented)

Time = 13.97 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \cos(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2}{b \sqrt{\frac{1}{\sin(a+bx)}}}$$

[In] int(cos(a + b*x)*(1/sin(a + b*x))^(1/2),x)

[Out] 2/(b*(1/sin(a + b*x))^(1/2))

3.262 $\int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1309
Rubi [A] (verified)	1309
Mathematica [A] (verified)	1310
Maple [A] (verified)	1310
Fricas [A] (verification not implemented)	1311
Sympy [F]	1311
Maxima [A] (verification not implemented)	1311
Giac [A] (verification not implemented)	1311
Mupad [B] (verification not implemented)	1312

Optimal result

Integrand size = 17, antiderivative size = 17

$$\int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

[Out] 2/3/b/csc(b*x+a)^(3/2)

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2701, 30}

$$\int \frac{\cos(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

[In] Int[Cos[a + b*x]/Sqrt[Csc[a + b*x]],x]

[Out] 2/(3*b*Csc[a + b*x]^(3/2))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2701

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n

+ 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x^{5/2}} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{2}{3b \csc^{3/2}(a + bx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2}{3b \csc^{3/2}(a + bx)}$$

[In] Integrate[Cos[a + b*x]/Sqrt[Csc[a + b*x]],x]

[Out] 2/(3*b*Csc[a + b*x]^(3/2))

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.82

method	result	size
derivativedivides	$\frac{2}{3b \csc(bx+a)^{3/2}}$	14
default	$\frac{2}{3b \csc(bx+a)^{3/2}}$	14

[In] int(cos(b*x+a)/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/3/b/csc(b*x+a)^(3/2)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.35

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = -\frac{2(\cos(bx + a)^2 - 1)}{3b\sqrt{\sin(bx + a)}}$$

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] -2/3*(cos(b*x + a)^2 - 1)/(b*sqrt(sin(b*x + a)))

Sympy [F]

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

[In] integrate(cos(b*x+a)/csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)/sqrt(csc(a + b*x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.22 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2 \sin(bx + a)^{\frac{3}{2}}}{3b}$$

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/3*sin(b*x + a)^(3/2)/b

Giac [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.76

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2 \sin(bx + a)^{\frac{3}{2}}}{3b}$$

[In] integrate(cos(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] 2/3*sin(b*x + a)^(3/2)/b

Mupad [B] (verification not implemented)

Time = 14.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{\cos(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2}{3b \left(\frac{1}{\sin(a+bx)} \right)^{3/2}}$$

[In] int(cos(a + b*x)/(1/sin(a + b*x))^(1/2),x)

[Out] 2/(3*b*(1/sin(a + b*x))^(3/2))

3.263 $\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal result	1313
Rubi [A] (verified)	1313
Mathematica [A] (verified)	1314
Maple [A] (verified)	1315
Fricas [C] (verification not implemented)	1315
Sympy [F]	1315
Maxima [F]	1316
Giac [F]	1316
Mupad [F(-1)]	1316

Optimal result

Integrand size = 19, antiderivative size = 67

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx$$

$$= \frac{2 \cos(a + bx)}{3b \sqrt{\csc(a + bx)}} + \frac{4 \sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right), 2\right) \sqrt{\sin(a + bx)}}{3b}$$

[Out] 2/3*cos(b*x+a)/b/csc(b*x+a)^(1/2)-4/3*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2708, 3856, 2720}

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx$$

$$= \frac{2 \cos(a + bx)}{3b \sqrt{\csc(a + bx)}} + \frac{4 \sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right), 2\right)}{3b}$$

[In] Int[Cos[a + b*x]^2*Sqrt[Csc[a + b*x]],x]

[Out] (2*Cos[a + b*x])/(3*b*Sqrt[Csc[a + b*x]]) + (4*Sqrt[Csc[a + b*x]]*EllipticF[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(3*b)

Rule 2708

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{2}{3} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{1}{3} \left(2\sqrt{\csc(a + bx)}\sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{2 \cos(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{4\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a - \frac{\pi}{2} + bx), 2\right) \sqrt{\sin(a + bx)}}{3b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.27 (sec) , antiderivative size = 53, normalized size of antiderivative = 0.79

$$\begin{aligned} &\int \cos^2(a + bx)\sqrt{\csc(a + bx)} dx \\ &= \frac{\sqrt{\csc(a + bx)} \left(-4 \operatorname{EllipticF}\left(\frac{1}{4}(-2a + \pi - 2bx), 2\right) \sqrt{\sin(a + bx)} + \sin(2(a + bx)) \right)}{3b} \end{aligned}$$

```
[In] Integrate[Cos[a + b*x]^2*Sqrt[Csc[a + b*x]],x]
```

```
[Out] (Sqrt[Csc[a + b*x]]*(-4*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + Sin[2*(a + b*x)])/(3*b)
```

Maple [A] (verified)

Time = 1.17 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.31

method	result	size
default	$\frac{2\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}F\left(\sqrt{\sin(bx+a)+1},\frac{\sqrt{2}}{2}\right)+\frac{2(\cos^2(bx+a)\sin(bx+a))}{3}}{\cos(bx+a)\sqrt{\sin(bx+a)}b}$	88

[In] `int(cos(b*x+a)^2*csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $(2/3*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticF}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})+2/3*\cos(b*x+a)^2*\sin(b*x+a))/\cos(b*x+a)/\sin(b*x+a)^{(1/2)}/b$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00

$$\int \cos^2(a+bx)\sqrt{\csc(a+bx)} dx = \frac{2\left(\cos(bx+a)\sqrt{\sin(bx+a)} - i\sqrt{2i}\text{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a)) + i\sqrt{-2i}\text{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a))\right)}{3b}$$

[In] `integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2),x, algorithm="fricas")`

[Out] $2/3*(\cos(b*x+a)*\text{sqrt}(\sin(b*x+a)) - I*\text{sqrt}(2*I)*\text{weierstrassPInverse}(4,0,\cos(b*x+a)+I*\sin(b*x+a)) + I*\text{sqrt}(-2*I)*\text{weierstrassPInverse}(4,0,\cos(b*x+a)-I*\sin(b*x+a)))/b$

Sympy [F]

$$\int \cos^2(a+bx)\sqrt{\csc(a+bx)} dx = \int \cos^2(a+bx)\sqrt{\csc(a+bx)} dx$$

[In] `integrate(cos(b*x+a)**2*csc(b*x+a)**(1/2),x)`

[Out] `Integral(cos(a+b*x)**2*sqrt(csc(a+b*x)),x)`

Maxima [F]

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos(bx + a)^2 \sqrt{\csc(bx + a)} dx$$

[In] integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^2*sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos(bx + a)^2 \sqrt{\csc(bx + a)} dx$$

[In] integrate(cos(b*x+a)^2*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^2*sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos(a + bx)^2 \sqrt{\frac{1}{\sin(a + bx)}} dx$$

[In] int(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2), x)

3.264 $\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1317
Rubi [A] (verified)	1317
Mathematica [A] (verified)	1318
Maple [A] (verified)	1319
Fricas [C] (verification not implemented)	1319
Sympy [F]	1319
Maxima [F]	1320
Giac [F]	1320
Mupad [F(-1)]	1320

Optimal result

Integrand size = 19, antiderivative size = 67

$$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{4\sqrt{\csc(a+bx)}E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{5b}$$

[Out] 2/5*cos(b*x+a)/b/csc(b*x+a)^(3/2)-4/5*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2708, 3856, 2719}

$$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2 \cos(a+bx)}{5b \csc^{\frac{3}{2}}(a+bx)} + \frac{4\sqrt{\sin(a+bx)}\sqrt{\csc(a+bx)}E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{5b}$$

[In] Int[Cos[a + b*x]^2/Sqrt[Csc[a + b*x]],x]

[Out] (2*Cos[a + b*x])/(5*b*Csc[a + b*x]^(3/2)) + (4*Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(5*b)

Rule 2708

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1]

] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \cos(a + bx)}{5b \csc^{\frac{3}{2}}(a + bx)} + \frac{2}{5} \int \frac{1}{\sqrt{\csc(a + bx)}} dx \\ &= \frac{2 \cos(a + bx)}{5b \csc^{\frac{3}{2}}(a + bx)} + \frac{1}{5} \left(2\sqrt{\csc(a + bx)}\sqrt{\sin(a + bx)} \right) \int \sqrt{\sin(a + bx)} dx \\ &= \frac{2 \cos(a + bx)}{5b \csc^{\frac{3}{2}}(a + bx)} + \frac{4\sqrt{\csc(a + bx)}E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{5b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.91

$$\begin{aligned} &\int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx \\ &= -\frac{2\sqrt{\csc(a + bx)}\left(2E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sqrt{\sin(a + bx)} - \cos(a + bx) \sin^2(a + bx)\right)}{5b} \end{aligned}$$

[In] Integrate[Cos[a + b*x]^2/Sqrt[Csc[a + b*x]],x]

[Out] (-2*Sqrt[Csc[a + b*x]]*(2*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] - Cos[a + b*x]*Sin[a + b*x]^2))/(5*b)

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.12

method	result
default	$\frac{-\frac{2(\sin^4(bx+a))}{5} + \frac{2(\sin^2(bx+a))}{5} - \frac{4\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}E\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) + \frac{2\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)}}{\cos(bx+a)\sqrt{\sin(bx+a)}}}{b}$

[In] int(cos(b*x+a)^2/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] $(-2/5*\sin(b*x+a)^4+2/5*\sin(b*x+a)^2-4/5*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticE}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)})+2/5*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticF}((\sin(b*x+a)+1)^{(1/2)},1/2*2^{(1/2)}))/\cos(b*x+a)/\sin(b*x+a)^{(1/2)}/b$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.24

$$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2\left(\sqrt{2i}\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a))) + \sqrt{-2i}\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a)))\right)}{5b}$$

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] $2/5*(\text{sqrt}(2*I)*\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(b*x+a)+I*\sin(b*x+a))) + \text{sqrt}(-2*I)*\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(b*x+a)-I*\sin(b*x+a)))) - (\cos(b*x+a)^3 - \cos(b*x+a))/\text{sqrt}(\sin(b*x+a)))/b$

Sympy [F]

$$\int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \int \frac{\cos^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

[In] integrate(cos(b*x+a)**2/csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)**2/sqrt(csc(a + b*x)), x)

Maxima [F]

$$\int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(bx + a)^2}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^2/sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(bx + a)^2}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(cos(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^2/sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(a + bx)^2}{\sqrt{\frac{1}{\sin(a + bx)}}} dx$$

[In] int(cos(a + b*x)^2/(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^2/(1/sin(a + b*x))^(1/2), x)

3.265 $\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx$

Optimal result	1321
Rubi [A] (verified)	1321
Mathematica [A] (verified)	1322
Maple [A] (verified)	1322
Fricas [A] (verification not implemented)	1323
Sympy [F(-1)]	1323
Maxima [A] (verification not implemented)	1323
Giac [A] (verification not implemented)	1323
Mupad [B] (verification not implemented)	1324

Optimal result

Integrand size = 11, antiderivative size = 21

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x)$$

[Out] $2/3*\csc(x)^{(3/2)}-2/7*\csc(x)^{(7/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2701, 14}

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x)$$

[In] $\text{Int}[\text{Cos}[x]^3*\text{Csc}[x]^{(9/2)}, x]$

[Out] $(2*\text{Csc}[x]^{(3/2)})/3 - (2*\text{Csc}[x]^{(7/2)})/7$

Rule 14

$\text{Int}[(u_*)*((c_*)*(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2701

$\text{Int}[(\csc[(e_*) + (f_*)*(x_)]*(a_))^{(m_*)}*\sec[(e_*) + (f_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(f*a^n)^{-1}, \text{Subst}[\text{Int}[x^{(m+n-1)} / (-1 + x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Csc}[e + f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n

+ 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \sqrt{x}(-1+x^2) dx, x, \csc(x)\right) \\ &= -\text{Subst}\left(\int (-\sqrt{x}+x^{5/2}) dx, x, \csc(x)\right) \\ &= \frac{2}{3} \csc^{\frac{3}{2}}(x) - \frac{2}{7} \csc^{\frac{7}{2}}(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.86

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2}{21} \csc^{\frac{3}{2}}(x) (7 - 3 \csc^2(x))$$

[In] Integrate[Cos[x]^3*Csc[x]^(9/2),x]

[Out] (2*Csc[x]^(3/2)*(7 - 3*Csc[x]^2))/21

Maple [A] (verified)

Time = 8.13 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

method	result	size
default	$-\frac{2}{7 \sin(x)^{\frac{7}{2}}} + \frac{2}{3 \sin(x)^{\frac{3}{2}}}$	14

[In] int(cos(x)^3*csc(x)^(9/2),x,method=_RETURNVERBOSE)

[Out] -2/7/sin(x)^(7/2)+2/3/sin(x)^(3/2)

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2(7 \cos(x)^2 - 4)}{21(\cos(x)^2 - 1) \sin(x)^{\frac{3}{2}}}$$

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="fricas")

[Out] 2/21*(7*cos(x)^2 - 4)/((cos(x)^2 - 1)*sin(x)^(3/2))

Sympy [F(-1)]

Timed out.

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \text{Timed out}$$

[In] integrate(cos(x)**3*csc(x)**(9/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.62

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2}{3 \sin(x)^{\frac{3}{2}}} - \frac{2}{7 \sin(x)^{\frac{7}{2}}}$$

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="maxima")

[Out] 2/3/sin(x)^(3/2) - 2/7/sin(x)^(7/2)

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.67

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2(7 \sin(x)^2 - 3)}{21 \sin(x)^{\frac{7}{2}}}$$

[In] integrate(cos(x)^3*csc(x)^(9/2),x, algorithm="giac")

[Out] 2/21*(7*sin(x)^2 - 3)/sin(x)^(7/2)

Mupad [B] (verification not implemented)

Time = 13.72 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.76

$$\int \cos^3(x) \csc^{\frac{9}{2}}(x) dx = \frac{2(7\sin(x)^2 - 3) \left(\frac{1}{\sin(x)}\right)^{7/2}}{21}$$

[In] int(cos(x)^3*(1/sin(x))^(9/2),x)

[Out] (2*(7*sin(x)^2 - 3)*(1/sin(x))^(7/2))/21

3.266 $\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal result	1325
Rubi [A] (verified)	1325
Mathematica [A] (verified)	1326
Maple [A] (verified)	1326
Fricas [A] (verification not implemented)	1327
Sympy [F(-1)]	1327
Maxima [A] (verification not implemented)	1327
Giac [A] (verification not implemented)	1327
Mupad [F(-1)]	1328

Optimal result

Integrand size = 19, antiderivative size = 33

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = -\frac{2}{5b \csc^{\frac{5}{2}}(a + bx)} + \frac{2}{b \sqrt{\csc(a + bx)}}$$

[Out] $-2/5/b/\csc(b*x+a)^{(5/2)}+2/b/\csc(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2701, 14}

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2}{b \sqrt{\csc(a + bx)}} - \frac{2}{5b \csc^{\frac{5}{2}}(a + bx)}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^3*\text{Sqrt}[\text{Csc}[a + b*x]],x]$

[Out] $-2/(5*b*\text{Csc}[a + b*x]^{(5/2)}) + 2/(b*\text{Sqrt}[\text{Csc}[a + b*x]])$

Rule 14

$\text{Int}[(u_*)*((c_*)*(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2701

$\text{Int}[(\csc[(e_*) + (f_)*(x_)]*(a_))^{(m_*)}\text{sec}[(e_*) + (f_)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(f*a^n)^{-1}, \text{Subst}[\text{Int}[x^{(m+n-1)} / (-1 + x^2/a^2)^{(n+1)/2}], x], x, a*\text{Csc}[e + f*x], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n

+ 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{-1+x^2}{x^{7/2}} dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{1}{x^{7/2}} + \frac{1}{x^{3/2}}\right) dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{2}{5b \csc^{5/2}(a+bx)} + \frac{2}{b\sqrt{\csc(a+bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int \cos^3(a+bx)\sqrt{\csc(a+bx)} dx = \frac{9 + \cos(2(a+bx))}{5b\sqrt{\csc(a+bx)}}$$

[In] Integrate[Cos[a + b*x]^3*Sqrt[Csc[a + b*x]], x]

[Out] (9 + Cos[2*(a + b*x)])/(5*b*Sqrt[Csc[a + b*x]])

Maple [A] (verified)

Time = 0.74 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.79

method	result	size
default	$-\frac{2\left(\sin^{\frac{5}{2}}(bx+a)\right)}{5} + 2\left(\sqrt{\sin(bx+a)}\right)$	26

[In] int(cos(b*x+a)^3*csc(b*x+a)^(1/2), x, method=_RETURNVERBOSE)

[Out] (-2/5*sin(b*x+a)^(5/2)+2*sin(b*x+a)^(1/2))/b

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2 (\cos(bx + a)^2 + 4) \sqrt{\sin(bx + a)}}{5b}$$

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] 2/5*(cos(b*x + a)^2 + 4)*sqrt(sin(b*x + a))/b

Sympy [F(-1)]

Timed out.

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = \text{Timed out}$$

[In] integrate(cos(b*x+a)**3*csc(b*x+a)**(1/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = \frac{2 \left(\frac{5}{\sin(bx+a)^2} - 1 \right) \sin(bx + a)^{\frac{5}{2}}}{5b}$$

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/5*(5/sin(b*x + a)^2 - 1)*sin(b*x + a)^(5/2)/b

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.73

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = -\frac{2 \left(\sin(bx + a)^{\frac{5}{2}} - 5 \sqrt{\sin(bx + a)} \right)}{5b}$$

[In] integrate(cos(b*x+a)^3*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] -2/5*(sin(b*x + a)^(5/2) - 5*sqrt(sin(b*x + a)))/b

Mupad [F(-1)]

Timed out.

$$\int \cos^3(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos(a + bx)^3 \sqrt{\frac{1}{\sin(a + bx)}} dx$$

```
[In] int(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2),x)
```

```
[Out] int(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2), x)
```


3.267 $\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1329
Rubi [A] (verified)	1329
Mathematica [A] (verified)	1330
Maple [A] (verified)	1330
Fricas [A] (verification not implemented)	1331
Sympy [F(-1)]	1331
Maxima [A] (verification not implemented)	1331
Giac [A] (verification not implemented)	1331
Mupad [F(-1)]	1332

Optimal result

Integrand size = 19, antiderivative size = 35

$$\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = -\frac{2}{7b \csc^{\frac{7}{2}}(a+bx)} + \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)}$$

[Out] $-2/7/b/\csc(b*x+a)^{(7/2)}+2/3/b/\csc(b*x+a)^{(3/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2701, 14}

$$\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{2}{7b \csc^{\frac{7}{2}}(a+bx)}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^3/\text{Sqrt}[\text{Csc}[a + b*x]], x]$

[Out] $-2/(7*b*\text{Csc}[a + b*x]^{(7/2)}) + 2/(3*b*\text{Csc}[a + b*x]^{(3/2)})$

Rule 14

$\text{Int}[(u_*)*((c_*)*(x_))^{(m_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2701

$\text{Int}[(\csc[(e_*) + (f_*)*(x_)]*(a_))^{(m_*)}*\sec[(e_*) + (f_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(f*a^n)^{-1}, \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{(n+1)}$

1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{-1+x^2}{x^{9/2}} dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{1}{x^{9/2}} + \frac{1}{x^{5/2}}\right) dx, x, \csc(a+bx)\right)}{b} \\ &= -\frac{2}{7b \csc^{7/2}(a+bx)} + \frac{2}{3b \csc^{3/2}(a+bx)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.83

$$\int \frac{\cos^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2(-3 + 7 \csc^2(a+bx))}{21b \csc^{7/2}(a+bx)}$$

[In] Integrate[Cos[a + b*x]^3/Sqrt[Csc[a + b*x]],x]

[Out] (2*(-3 + 7*Csc[a + b*x]^2))/(21*b*Csc[a + b*x]^(7/2))

Maple [A] (verified)

Time = 0.91 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

method	result	size
default	$-\frac{2\left(\sin^{7/2}(bx+a)\right)}{7} + \frac{2\left(\sin^{3/2}(bx+a)\right)}{3}$	26

[In] int(cos(b*x+a)^3/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] (-2/7*sin(b*x+a)^(7/2)+2/3*sin(b*x+a)^(3/2))/b

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.94

$$\int \frac{\cos^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = -\frac{2(3 \cos(bx + a)^4 + \cos(bx + a)^2 - 4)}{21 b \sqrt{\sin(bx + a)}}$$

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] -2/21*(3*cos(b*x + a)^4 + cos(b*x + a)^2 - 4)/(b*sqrt(sin(b*x + a)))

Sympy [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \text{Timed out}$$

[In] integrate(cos(b*x+a)**3/csc(b*x+a)**(1/2),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.71

$$\int \frac{\cos^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2 \left(\frac{7}{\sin(bx+a)^2} - 3 \right) \sin(bx + a)^{\frac{7}{2}}}{21 b}$$

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 2/21*(7/sin(b*x + a)^2 - 3)*sin(b*x + a)^(7/2)/b

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.74

$$\int \frac{\cos^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = -\frac{2 \left(3 \sin(bx + a)^{\frac{7}{2}} - 7 \sin(bx + a)^{\frac{3}{2}} \right)}{21 b}$$

[In] integrate(cos(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] -2/21*(3*sin(b*x + a)^(7/2) - 7*sin(b*x + a)^(3/2))/b

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(a + bx)^3}{\sqrt{\frac{1}{\sin(a + bx)}}} dx$$

```
[In] int(cos(a + b*x)^3/(1/sin(a + b*x))^(1/2),x)
```

```
[Out] int(cos(a + b*x)^3/(1/sin(a + b*x))^(1/2), x)
```

3.268 $\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$

Optimal result	1333
Rubi [A] (verified)	1333
Mathematica [A] (verified)	1335
Maple [A] (verified)	1335
Fricas [C] (verification not implemented)	1335
Sympy [F]	1336
Maxima [F]	1336
Giac [F]	1336
Mupad [F(-1)]	1336

Optimal result

Integrand size = 19, antiderivative size = 92

$$\begin{aligned} & \int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx \\ &= \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} \\ &+ \frac{8 \sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a - \frac{\pi}{2} + bx), 2\right) \sqrt{\sin(a + bx)}}{7b} \end{aligned}$$

[Out] $4/7*\cos(b*x+a)/b/\csc(b*x+a)^{(1/2)}+2/7*\cos(b*x+a)^3/b/\csc(b*x+a)^{(1/2)}-8/7*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\operatorname{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\csc(b*x+a)^{(1/2)}*\sin(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2708, 3856, 2720}

$$\begin{aligned} & \int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx \\ &= \frac{2 \cos^3(a + bx)}{7b \sqrt{\csc(a + bx)}} + \frac{4 \cos(a + bx)}{7b \sqrt{\csc(a + bx)}} \\ &+ \frac{8 \sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx - \frac{\pi}{2}), 2\right)}{7b} \end{aligned}$$

[In] $\operatorname{Int}[\cos[a + b*x]^4*\sqrt{\csc[a + b*x]}, x]$

[Out] $(4\cos[a + bx])/(7b\sqrt{\csc[a + bx]}) + (2\cos[a + bx]^3)/(7b\sqrt{\csc[a + bx]}) + (8\sqrt{\csc[a + bx]}*\text{EllipticF}[(a - \pi/2 + bx)/2, 2]*\sqrt{\sin[a + bx]})/(7b)$

Rule 2708

$\text{Int}[(\csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*\sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\csc[e + f*x])^{m-1}*((b*\sec[e + f*x])^{n+1})/(b*f*(m+n)), x] + \text{Dist}[(n+1)/(b^2*(m+n)), \text{Int}[(a*\csc[e + f*x])^m*(b*\sec[e + f*x])^{n+2}, x], x] /;$ FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

$\text{Int}[1/\sqrt{\sin[(c_.) + (d_.)*(x_)]}, x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \pi/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\sin[c + d*x]^n, \text{Int}[1/\sin[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{2 \cos^3(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{6}{7} \int \cos^2(a + bx) \sqrt{\csc(a + bx)} dx \\ &= \frac{4 \cos(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{4}{7} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{4 \cos(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{1}{7} \left(4\sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{4 \cos(a + bx)}{7b\sqrt{\csc(a + bx)}} + \frac{2 \cos^3(a + bx)}{7b\sqrt{\csc(a + bx)}} \\ &\quad + \frac{8\sqrt{\csc(a + bx)} \text{EllipticF}\left(\frac{1}{2}(a - \frac{\pi}{2} + bx), 2\right) \sqrt{\sin(a + bx)}}{7b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.31 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.68

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$$

$$= \frac{\sqrt{\csc(a + bx)} \left(-32 \operatorname{EllipticF} \left(\frac{1}{4}(-2a + \pi - 2bx), 2 \right) \sqrt{\sin(a + bx)} + 10 \sin(2(a + bx)) + \sin(4(a + bx)) \right)}{28b}$$

[In] Integrate[Cos[a + b*x]^4*Sqrt[Csc[a + b*x]],x]

[Out] (Sqrt[Csc[a + b*x]]*(-32*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + 10*Sin[2*(a + b*x)] + Sin[4*(a + b*x)]))/(28*b)

Maple [A] (verified)

Time = 1.22 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.09

method	result	size
default	$\frac{4\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}F\left(\sqrt{\sin(bx+a)+1},\frac{\sqrt{2}}{2}\right) + \frac{2(\sin^5(bx+a))}{7} - \frac{8(\sin^3(bx+a))}{7} + \frac{6\sin(bx+a)}{7}}{\cos(bx+a)\sqrt{\sin(bx+a)}b}$	100

[In] int(cos(b*x+a)^4*csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] (4/7*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))+2/7*sin(b*x+a)^5-8/7*sin(b*x+a)^3+6/7*sin(b*x+a))/cos(b*x+a)/sin(b*x+a)^(1/2)/b

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.09 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.85

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$$

$$= \frac{2 \left((\cos(bx + a))^3 + 2 \cos(bx + a) \right) \sqrt{\sin(bx + a)} - 2i \sqrt{2i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx + a) + i \sin(bx + a))}{7b}$$

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] 2/7*((cos(b*x + a)^3 + 2*cos(b*x + a))*sqrt(sin(b*x + a)) - 2*I*sqrt(2*I)*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + 2*I*sqrt(-2*I)*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)))/b

Sympy [F]

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx$$

[In] integrate(cos(b*x+a)**4*csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)**4*sqrt(csc(a + b*x)), x)

Maxima [F]

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos^4(bx + a) \sqrt{\csc(bx + a)} dx$$

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^4*sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos^4(bx + a) \sqrt{\csc(bx + a)} dx$$

[In] integrate(cos(b*x+a)^4*csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^4*sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \cos^4(a + bx) \sqrt{\csc(a + bx)} dx = \int \cos^4(a + bx) \sqrt{\frac{1}{\sin(a + bx)}} dx$$

[In] int(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2), x)

3.269 $\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1337
Rubi [A] (verified)	1337
Mathematica [A] (verified)	1338
Maple [A] (verified)	1339
Fricas [C] (verification not implemented)	1339
Sympy [F]	1339
Maxima [F]	1340
Giac [F]	1340
Mupad [F(-1)]	1340

Optimal result

Integrand size = 19, antiderivative size = 92

$$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{4 \cos(a+bx)}{15b \csc^{\frac{3}{2}}(a+bx)} + \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{8 \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{15b}$$

[Out] $4/15*\cos(b*x+a)/b/\csc(b*x+a)^{(3/2)}+2/9*\cos(b*x+a)^3/b/\csc(b*x+a)^{(3/2)}-8/15*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x),2^{(1/2)})*\csc(b*x+a)^{(1/2)}*\sin(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2708, 3856, 2719}

$$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2 \cos^3(a+bx)}{9b \csc^{\frac{3}{2}}(a+bx)} + \frac{4 \cos(a+bx)}{15b \csc^{\frac{3}{2}}(a+bx)} + \frac{8 \sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{15b}$$

[In] $\text{Int}[\text{Cos}[a + b*x]^4/\text{Sqrt}[\text{Csc}[a + b*x]], x]$

[Out] $(4*\text{Cos}[a + b*x])/(15*b*\text{Csc}[a + b*x]^{(3/2)}) + (2*\text{Cos}[a + b*x]^3)/(9*b*\text{Csc}[a + b*x]^{(3/2)}) + (8*\text{Sqrt}[\text{Csc}[a + b*x]]*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[\text{Sin}[a + b*x]])/(15*b)$

Rule 2708

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(b*f*(m + n))), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{2 \cos^3(a + bx)}{9b \csc^{\frac{3}{2}}(a + bx)} + \frac{2}{3} \int \frac{\cos^2(a + bx)}{\sqrt{\csc(a + bx)}} dx \\
 &= \frac{4 \cos(a + bx)}{15b \csc^{\frac{3}{2}}(a + bx)} + \frac{2 \cos^3(a + bx)}{9b \csc^{\frac{3}{2}}(a + bx)} + \frac{4}{15} \int \frac{1}{\sqrt{\csc(a + bx)}} dx \\
 &= \frac{4 \cos(a + bx)}{15b \csc^{\frac{3}{2}}(a + bx)} + \frac{2 \cos^3(a + bx)}{9b \csc^{\frac{3}{2}}(a + bx)} + \frac{1}{15} \left(4 \sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \sqrt{\sin(a + bx)} dx \\
 &= \frac{4 \cos(a + bx)}{15b \csc^{\frac{3}{2}}(a + bx)} + \frac{2 \cos^3(a + bx)}{9b \csc^{\frac{3}{2}}(a + bx)} + \frac{8 \sqrt{\csc(a + bx)} E\left(\frac{1}{2}(a - \frac{\pi}{2} + bx) \mid 2\right) \sqrt{\sin(a + bx)}}{15b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.55 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.68

$$\int \frac{\cos^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{39 \cos(a + bx) + 5 \cos(3(a + bx)) - \frac{48 E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right)}{\sin^{\frac{3}{2}}(a + bx)}}{90b \csc^{\frac{3}{2}}(a + bx)}$$

```
[In] Integrate[Cos[a + b*x]^4/Sqrt[Csc[a + b*x]], x]
```

```
[Out] (39*Cos[a + b*x] + 5*Cos[3*(a + b*x)] - (48*EllipticE[(-2*a + Pi - 2*b*x)/4, 2])/Sin[a + b*x]^(3/2))/(90*b*Csc[a + b*x]^(3/2))
```

Maple [A] (verified)

Time = 1.13 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.65

method	result
default	$\frac{-\frac{2(\cos^6(bx+a))}{9} - \frac{8\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)} E\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{15} + \frac{4\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}}{15}}{\cos(bx+a)\sqrt{\sin(bx+a)}b}$

[In] int(cos(b*x+a)^4/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] $(-2/9*\cos(b*x+a)^6 - 8/15*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticE}((\sin(b*x+a)+1)^{(1/2)}, 1/2*2^{(1/2)}) + 4/15*(\sin(b*x+a)+1)^{(1/2)}*(-2*\sin(b*x+a)+2)^{(1/2)}*(-\sin(b*x+a))^{(1/2)}*\text{EllipticF}((\sin(b*x+a)+1)^{(1/2)}, 1/2*2^{(1/2)}) - 2/45*\cos(b*x+a)^4 + 4/15*\cos(b*x+a)^2)/\cos(b*x+a)/\sin(b*x+a)^{(1/2)}/b$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.03

$$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

$$= \frac{2 \left(6\sqrt{2}i \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a))) + 6\sqrt{-2i} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a))) \right) + 6\sqrt{-2i} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)))}{45b}$$

[In] integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] $2/45*(6*\text{sqrt}(2*I)*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x+a) + I*\sin(b*x+a))) + 6*\text{sqrt}(-2*I)*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x+a) - I*\sin(b*x+a))) - (5*\cos(b*x+a)^5 + \cos(b*x+a)^3 - 6*\cos(b*x+a))/\text{sqrt}(\sin(b*x+a)))/b$

Sympy [F]

$$\int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \int \frac{\cos^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

[In] integrate(cos(b*x+a)**4/csc(b*x+a)**(1/2),x)

[Out] Integral(cos(a + b*x)**4/sqrt(csc(a + b*x)), x)

Maxima [F]

$$\int \frac{\cos^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(bx + a)^4}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^4/sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \frac{\cos^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(bx + a)^4}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(cos(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^4/sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cos^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\cos(a + bx)^4}{\sqrt{\frac{1}{\sin(a + bx)}}} dx$$

[In] int(cos(a + b*x)^4/(1/sin(a + b*x))^(1/2),x)

[Out] int(cos(a + b*x)^4/(1/sin(a + b*x))^(1/2), x)

3.270 $\int \cos(x) \csc^{\frac{7}{3}}(x) dx$

Optimal result	1341
Rubi [A] (verified)	1341
Mathematica [A] (verified)	1342
Maple [A] (verified)	1342
Fricas [A] (verification not implemented)	1342
Sympy [F(-1)]	1343
Maxima [A] (verification not implemented)	1343
Giac [A] (verification not implemented)	1343
Mupad [B] (verification not implemented)	1343

Optimal result

Integrand size = 9, antiderivative size = 10

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

[Out] $-3/4*\csc(x)^{(4/3)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2701, 30}

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

[In] `Int[Cos[x]*Csc[x]^(7/3),x]`

[Out] `(-3*Csc[x]^(4/3))/4`

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2701

`Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \sqrt[3]{x} dx, x, \csc(x)\right) \\ &= -\frac{3}{4} \csc^{\frac{4}{3}}(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4} \csc^{\frac{4}{3}}(x)$$

[In] Integrate[Cos[x]*Csc[x]^(7/3),x]

[Out] (-3*Csc[x]^(4/3))/4

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.70

method	result	size
derivativedivides	$-\frac{3(\csc^{\frac{4}{3}}(x))}{4}$	7
default	$-\frac{3(\csc^{\frac{4}{3}}(x))}{4}$	7

[In] int(cos(x)*csc(x)^(7/3),x,method=_RETURNVERBOSE)

[Out] -3/4*csc(x)^(4/3)

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

[In] integrate(cos(x)*csc(x)^(7/3),x, algorithm="fricas")

[Out] -3/4/sin(x)^(4/3)

Sympy [F(-1)]

Timed out.

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = \text{Timed out}$$

[In] integrate(cos(x)*csc(x)**(7/3),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

[In] integrate(cos(x)*csc(x)^(7/3),x, algorithm="maxima")

[Out] -3/4/sin(x)^(4/3)

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 6, normalized size of antiderivative = 0.60

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3}{4 \sin(x)^{\frac{4}{3}}}$$

[In] integrate(cos(x)*csc(x)^(7/3),x, algorithm="giac")

[Out] -3/4/sin(x)^(4/3)

Mupad [B] (verification not implemented)

Time = 13.43 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \cos(x) \csc^{\frac{7}{3}}(x) dx = -\frac{3 \left(\frac{1}{\sin(x)}\right)^{\frac{4}{3}}}{4}$$

[In] int(cos(x)*(1/sin(x))^(7/3),x)

[Out] -(3*(1/sin(x))^(4/3))/4

3.271 $\int \sqrt{\csc(a + bx)} \sec(a + bx) dx$

Optimal result	1344
Rubi [A] (verified)	1344
Mathematica [A] (verified)	1346
Maple [A] (verified)	1346
Fricas [B] (verification not implemented)	1346
Sympy [F]	1347
Maxima [A] (verification not implemented)	1347
Giac [A] (verification not implemented)	1347
Mupad [F(-1)]	1348

Optimal result

Integrand size = 17, antiderivative size = 32

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx = -\frac{\arctan\left(\sqrt{\csc(a + bx)}\right)}{b} + \frac{\operatorname{arctanh}\left(\sqrt{\csc(a + bx)}\right)}{b}$$

[Out] $-\arctan(\csc(b*x+a)^{(1/2)})/b + \operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2701, 335, 304, 209, 212}

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx = \frac{\operatorname{arctanh}\left(\sqrt{\csc(a + bx)}\right)}{b} - \frac{\arctan\left(\sqrt{\csc(a + bx)}\right)}{b}$$

[In] $\text{Int}[\text{Sqrt}[\text{Csc}[a + b*x]]*\text{Sec}[a + b*x], x]$

[Out] $-(\text{ArcTan}[\text{Sqrt}[\text{Csc}[a + b*x]])/b) + \text{ArcTanh}[\text{Sqrt}[\text{Csc}[a + b*x]]]/b$

Rule 209

$\text{Int}[\left((a_) + (b_.)*(x_)^2\right)^{-1}, x_Symbol] \rightarrow \text{Simp}[\left(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2])\right)*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 212

$\text{Int}[\left((a_) + (b_.)*(x_)^2\right)^{-1}, x_Symbol] \rightarrow \text{Simp}[\left(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2])\right)*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{Gt}$

Q[a, 0] || LtQ[b, 0])

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2701

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_), x_S
ymbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n +
1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n
+ 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \csc(a+bx)\right)}{b} \\
 &= -\frac{2\text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
 &= -\frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\text{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.47

$$\int \sqrt{\csc(a+bx)} \sec(a+bx) dx$$

$$= \frac{\left(\arctan\left(\sqrt{\sin(a+bx)}\right) + \operatorname{arctanh}\left(\sqrt{\sin(a+bx)}\right) \right) \sqrt{\csc(a+bx)} \sqrt{\sin(a+bx)}}{b}$$

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x],x]

[Out] ((ArcTan[Sqrt[Sin[a + b*x]]] + ArcTanh[Sqrt[Sin[a + b*x]]])*Sqrt[Csc[a + b*x]]*Sqrt[Sin[a + b*x]])/b

Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{\operatorname{arctanh}(\sqrt{\sin(bx+a)}) + \arctan(\sqrt{\sin(bx+a)})}{b}$	24

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a),x,method=_RETURNVERBOSE)

[Out] (arctanh(sin(b*x+a)^(1/2))+arctan(sin(b*x+a)^(1/2)))/b

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(28) = 56.

Time = 0.33 (sec) , antiderivative size = 95, normalized size of antiderivative = 2.97

$$\int \sqrt{\csc(a+bx)} \sec(a+bx) dx$$

$$= \frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) + \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6\sin(bx+a)-2}{\cos(bx+a)^2 + 2\sin(bx+a)-2}\right)}{4b}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a),x, algorithm="fricas")

[Out] 1/4*(2*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a))) + log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)))/b

Sympy [F]

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx = \int \sqrt{\csc(a + bx)} \sec(a + bx) dx$$

[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a),x)

[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x), x)

Maxima [A] (verification not implemented)

none

Time = 0.45 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.28

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx$$

$$= -\frac{2 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) - \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) + \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{2b}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a),x, algorithm="maxima")

[Out] -1/2*(2*arctan(1/sqrt(sin(b*x + a))) - log(1/sqrt(sin(b*x + a)) + 1) + log(1/sqrt(sin(b*x + a)) - 1))/b

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.31

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx$$

$$= \frac{2 \arctan\left(\sqrt{\sin(bx+a)}\right) + \log\left(\sqrt{\sin(bx+a)} + 1\right) - \log\left(\left|\sqrt{\sin(bx+a)} - 1\right|\right)}{2b}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a),x, algorithm="giac")

[Out] 1/2*(2*arctan(sqrt(sin(b*x + a))) + log(sqrt(sin(b*x + a)) + 1) - log(abs(sqrt(sin(b*x + a)) - 1)))/b

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\csc(a + bx)} \sec(a + bx) dx = \int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a + bx)} dx$$

```
[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x),x)
```

```
[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x), x)
```

$$3.272 \quad \int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

Optimal result	1349
Rubi [A] (verified)	1349
Mathematica [A] (verified)	1351
Maple [A] (verified)	1351
Fricas [B] (verification not implemented)	1351
Sympy [F]	1352
Maxima [A] (verification not implemented)	1352
Giac [A] (verification not implemented)	1352
Mupad [F(-1)]	1353

Optimal result

Integrand size = 17, antiderivative size = 31

$$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\operatorname{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{b}$$

[Out] $\arctan(\csc(b*x+a)^{(1/2)})/b + \operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {2701, 335, 218, 212, 209}

$$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\operatorname{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{b}$$

[In] $\text{Int}[\text{Sec}[a + b*x]/\text{Sqrt}[\text{Csc}[a + b*x]], x]$

[Out] $\text{ArcTan}[\text{Sqrt}[\text{Csc}[a + b*x]]]/b + \text{ArcTanh}[\text{Sqrt}[\text{Csc}[a + b*x]]]/b$

Rule 209

$\text{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2701

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_S
ymbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n +
1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n
+ 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x(-1+x^2)}} dx, x, \csc(a+bx)\right)}{b} \\
&= -\frac{2\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{b} \\
&= \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{b} + \frac{\text{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.61

$$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

$$= -\frac{\left(\arctan\left(\sqrt{\sin(a+bx)}\right) - \operatorname{arctanh}\left(\sqrt{\sin(a+bx)}\right)\right) \sqrt{\csc(a+bx)} \sqrt{\sin(a+bx)}}{b}$$

[In] Integrate[Sec[a + b*x]/Sqrt[Csc[a + b*x]],x]

[Out] -(((ArcTan[Sqrt[Sin[a + b*x]]] - ArcTanh[Sqrt[Sin[a + b*x]])]*Sqrt[Csc[a + b*x]]*Sqrt[Sin[a + b*x]])/b)

Maple [A] (verified)

Time = 1.02 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.39

method	result	size
default	$\frac{-\frac{\ln(\sqrt{\sin(bx+a)}-1)}{2} + \frac{\ln(1+\sqrt{\sin(bx+a)})}{2} - \arctan(\sqrt{\sin(bx+a)})}{b}$	43

[In] int(sec(b*x+a)/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] (-1/2*ln(sin(b*x+a)^(1/2)-1)+1/2*ln(1+sin(b*x+a)^(1/2))-arctan(sin(b*x+a)^(1/2)))/b

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(27) = 54.

Time = 0.32 (sec) , antiderivative size = 97, normalized size of antiderivative = 3.13

$$\int \frac{\sec(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

$$= -\frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) - \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6\sin(bx+a)-2}{\cos(bx+a)^2 + 2\sin(bx+a)-2}\right)}{4b}$$

[In] integrate(sec(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out] -1/4*(2*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a))) - log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)))/b

Sympy [F]

$$\int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

[In] integrate(sec(b*x+a)/csc(b*x+a)**(1/2),x)

[Out] Integral(sec(a + b*x)/sqrt(csc(a + b*x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.32

$$\int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) + \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) - \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{2b}$$

[In] integrate(sec(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] 1/2*(2*arctan(1/sqrt(sin(b*x + a))) + log(1/sqrt(sin(b*x + a)) + 1) - log(1/sqrt(sin(b*x + a)) - 1))/b

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.35

$$\int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{2 \arctan\left(\sqrt{\sin(bx + a)}\right) - \log\left(\sqrt{\sin(bx + a)} + 1\right) + \log\left(\left|\sqrt{\sin(bx + a)} - 1\right|\right)}{2b}$$

[In] integrate(sec(b*x+a)/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] -1/2*(2*arctan(sqrt(sin(b*x + a))) - log(sqrt(sin(b*x + a)) + 1) + log(abs(sqrt(sin(b*x + a)) - 1)))/b

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{1}{\cos(a + bx) \sqrt{\frac{1}{\sin(a + bx)}}} dx$$

```
[In] int(1/(cos(a + b*x)*(1/sin(a + b*x))^(1/2)),x)
```

```
[Out] int(1/(cos(a + b*x)*(1/sin(a + b*x))^(1/2)), x)
```

3.273 $\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx$

Optimal result	1354
Rubi [A] (verified)	1354
Mathematica [A] (verified)	1355
Maple [A] (verified)	1356
Fricas [C] (verification not implemented)	1356
Sympy [F]	1356
Maxima [F]	1357
Giac [F]	1357
Mupad [F(-1)]	1357

Optimal result

Integrand size = 19, antiderivative size = 61

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx$$

$$= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right), 2\right) \sqrt{\sin(a + bx)}}{b}$$

[Out] $\sec(b*x+a)/b/\csc(b*x+a)^{(1/2)} - (\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\operatorname{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\csc(b*x+a)^{(1/2)}*\sin(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2706, 3856, 2720}

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx$$

$$= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\sin(a + bx)} \sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right), 2\right)}{b}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]*\operatorname{Sec}[a + b*x]^2, x]$

[Out] $\operatorname{Sec}[a + b*x]/(b*\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]) + (\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]*\operatorname{EllipticF}[(a - \operatorname{Pi}/2 + b*x)/2, 2]*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]])/b$

Rule 2706

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Dist[b^2*((m + n - 2)/(n - 1)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{1}{2} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{1}{2} \left(\sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{\sec(a + bx)}{b\sqrt{\csc(a + bx)}} + \frac{\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right), 2\right) \sqrt{\sin(a + bx)}}{b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.32 (sec) , antiderivative size = 49, normalized size of antiderivative = 0.80

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx = \frac{\sec(a + bx) + \frac{\operatorname{EllipticF}\left(\frac{1}{4}(2a - \pi + 2bx), 2\right)}{\sqrt{\sin(a + bx)}}}{b\sqrt{\csc(a + bx)}}$$

```
[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^2,x]
```

```
[Out] (Sec[a + b*x] + EllipticF[(2*a - Pi + 2*b*x)/4, 2]/Sqrt[Sin[a + b*x]])/(b*Sqrt[Csc[a + b*x]])
```

Maple [A] (verified)

Time = 1.27 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.02

method	result	size
default	$\frac{\sqrt{\cos^2(bx+a)} \sin(bx+a) \left(\sqrt{\sin(bx+a)+1} \sqrt{-2 \sin(bx+a)+2} \sqrt{-\sin(bx+a)} F\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) + 2 \sin(bx+a) \right)}{2 \sqrt{-\sin(bx+a)} (\sin(bx+a)-1) (\sin(bx+a)+1) \cos(bx+a) \sqrt{\sin(bx+a)} b}$	123

```
[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*(cos(b*x+a)^2*sin(b*x+a))^(1/2)*((sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))+2*sin(b*x+a))/(-sin(b*x+a)*(sin(b*x+a)-1)*(sin(b*x+a)+1))^(1/2)/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.13 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.34

$$\int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx = \frac{-i\sqrt{2i} \cos(bx+a) \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + i\sqrt{-2i} \cos(bx+a) \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a))}{2b \cos(bx+a)}$$

```
[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] 1/2*(-I*sqrt(2*I)*cos(b*x + a)*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(-2*I)*cos(b*x + a)*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*sqrt(sin(b*x + a)))/(b*cos(b*x + a))
```

Sympy [F]

$$\int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx = \int \sqrt{\csc(a+bx)} \sec^2(a+bx) dx$$

```
[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**2,x)
```

```
[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x)**2, x)
```

Maxima [F]

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx = \int \sqrt{\csc(bx + a)} \sec(bx + a)^2 dx$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^2, x)

Giac [F]

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx = \int \sqrt{\csc(bx + a)} \sec(bx + a)^2 dx$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^2,x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx = \int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a + bx)^2} dx$$

[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^2,x)

[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^2, x)

3.274 $\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1358
Rubi [A] (verified)	1358
Mathematica [A] (verified)	1359
Maple [B] (verified)	1360
Fricas [C] (verification not implemented)	1360
Sympy [F]	1360
Maxima [F]	1361
Giac [F]	1361
Mupad [F(-1)]	1361

Optimal result

Integrand size = 19, antiderivative size = 62

$$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{b}$$

[Out] sec(b*x+a)/b/csc(b*x+a)^(3/2)+(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2706, 3856, 2719}

$$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\sec(a+bx)}{b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

[In] Int[Sec[a + b*x]^2/Sqrt[Csc[a + b*x]],x]

[Out] Sec[a + b*x]/(b*Csc[a + b*x]^(3/2)) - (Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/b

Rule 2706

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Dist[b^2*((m + n - 2)/(n - 1)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1

] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sec(a + bx)}{b \csc^{\frac{3}{2}}(a + bx)} - \frac{1}{2} \int \frac{1}{\sqrt{\csc(a + bx)}} dx \\ &= \frac{\sec(a + bx)}{b \csc^{\frac{3}{2}}(a + bx)} - \frac{1}{2} \left(\sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \sqrt{\sin(a + bx)} dx \\ &= \frac{\sec(a + bx)}{b \csc^{\frac{3}{2}}(a + bx)} - \frac{\sqrt{\csc(a + bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.28 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.87

$$\begin{aligned} &\int \frac{\sec^2(a + bx)}{\sqrt{\csc(a + bx)}} dx \\ &= \frac{\sqrt{\csc(a + bx)} \left(E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sqrt{\sin(a + bx)} + \sin(a + bx) \tan(a + bx) \right)}{b} \end{aligned}$$

[In] Integrate[Sec[a + b*x]^2/Sqrt[Csc[a + b*x]], x]

[Out] (Sqrt[Csc[a + b*x]]*(EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + Sin[a + b*x]*Tan[a + b*x]))/b

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 176 vs. $2(85) = 170$.

Time = 1.36 (sec) , antiderivative size = 177, normalized size of antiderivative = 2.85

method	result
default	$\frac{\sqrt{(\cos^2(bx+a)) \sin(bx+a)} \left(2\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} E\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) - \sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)} \right)}{2\sqrt{-\sin(bx+a)(\sin(bx+a)-1)(\sin(bx+a)+1)} \cos(bx+a) \sqrt{\sin(bx+a)}}$

[In] `int(sec(b*x+a)^2/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2}(\cos(bx+a)^2 \sin(bx+a))^{1/2} (2(\sin(bx+a)+1)^{1/2} (-2\sin(bx+a)+2)^{1/2} (-\sin(bx+a))^{1/2} \text{EllipticE}(\sin(bx+a)+1, 1/2 \cdot 2^{1/2}) - (\sin(bx+a)+1)^{1/2} (-2\sin(bx+a)+2)^{1/2} (-\sin(bx+a))^{1/2} \text{EllipticF}(\sin(bx+a)+1, 1/2 \cdot 2^{1/2}) - 2\cos(bx+a)^2) / (-\sin(bx+a) (\sin(bx+a)-1) (\sin(bx+a)+1))^{1/2} / \cos(bx+a) / \sin(bx+a)^{1/2} / b$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.55

$$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\sqrt{2i} \cos(bx+a) \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a))) + \sqrt{-2i} \cos(bx+a) \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)))}{2b \cos(bx+a)}$$

[In] `integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="fricas")`

[Out]
$$-1/2(\sqrt{2I} \cos(bx+a) \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + I \sin(bx+a))) + \sqrt{-2I} \cos(bx+a) \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - I \sin(bx+a)))) + 2(\cos(bx+a)^2 - 1) / \sqrt{\sin(bx+a)} / (b \cos(bx+a))$$

Sympy [F]

$$\int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx = \int \frac{\sec^2(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

[In] `integrate(sec(b*x+a)**2/csc(b*x+a)**(1/2),x)`

[Out] `Integral(sec(a + b*x)**2/sqrt(csc(a + b*x)), x)`

Maxima [F]

$$\int \frac{\sec^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec(bx + a)^2}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^2/sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \frac{\sec^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec(bx + a)^2}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(sec(b*x+a)^2/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^2/sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^2(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{1}{\cos(a + bx)^2 \sqrt{\frac{1}{\sin(a + bx)}}} dx$$

[In] int(1/(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2)),x)

[Out] int(1/(cos(a + b*x)^2*(1/sin(a + b*x))^(1/2)), x)

3.275 $\int \sqrt{\csc(a + bx)} \sec^3(a + bx) dx$

Optimal result	1362
Rubi [A] (verified)	1362
Mathematica [A] (verified)	1364
Maple [A] (verified)	1364
Fricas [B] (verification not implemented)	1365
Sympy [F]	1365
Maxima [A] (verification not implemented)	1365
Giac [A] (verification not implemented)	1366
Mupad [F(-1)]	1366

Optimal result

Integrand size = 19, antiderivative size = 62

$$\int \sqrt{\csc(a + bx)} \sec^3(a + bx) dx = -\frac{3 \arctan\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{3 \operatorname{arctanh}\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{\sec^2(a + bx)}{2b\sqrt{\csc(a + bx)}}$$

[Out] $-3/4*\arctan(\csc(b*x+a)^{(1/2)})/b+3/4*\operatorname{arctanh}(\csc(b*x+a)^{(1/2)})/b+1/2*\sec(b*x+a)^2/b/\csc(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2701, 294, 335, 304, 209, 212}

$$\int \sqrt{\csc(a + bx)} \sec^3(a + bx) dx = -\frac{3 \arctan\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{3 \operatorname{arctanh}\left(\sqrt{\csc(a + bx)}\right)}{4b} + \frac{\sec^2(a + bx)}{2b\sqrt{\csc(a + bx)}}$$

[In] $\text{Int}[\text{Sqrt}[\text{Csc}[a + b*x]]*\text{Sec}[a + b*x]^3,x]$

[Out] $(-3*\text{ArcTan}[\text{Sqrt}[\text{Csc}[a + b*x]]])/(4*b) + (3*\text{ArcTanh}[\text{Sqrt}[\text{Csc}[a + b*x]]])/(4*b) + \text{Sec}[a + b*x]^2/(2*b*\text{Sqrt}[\text{Csc}[a + b*x]])$

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 304

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^{5/2}}{(-1+x^2)^2} dx, x, \csc(a+bx)\right)}{b} \\ &= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} - \frac{3\text{Subst}\left(\int \frac{\sqrt{x}}{-1+x^2} dx, x, \csc(a+bx)\right)}{4b} \end{aligned}$$

$$\begin{aligned}
&= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} - \frac{3\text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{2b} \\
&= \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}} + \frac{3\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} \\
&\quad - \frac{3\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} \\
&= -\frac{3\arctan\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{3\text{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\sec^2(a+bx)}{2b\sqrt{\csc(a+bx)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.19

$$\begin{aligned}
&\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx \\
&= \frac{\sqrt{\csc(a+bx)} \left(3\arctan\left(\sqrt{\sin(a+bx)}\right) + 3\text{arctanh}\left(\sqrt{\sin(a+bx)}\right) + 2\sec^2(a+bx)\sqrt{\sin(a+bx)} \right) \sqrt{\sin(a+bx)}}{4b}
\end{aligned}$$

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^3,x]

[Out] (Sqrt[Csc[a + b*x]]*(3*ArcTan[Sqrt[Sin[a + b*x]]] + 3*ArcTanh[Sqrt[Sin[a + b*x]]] + 2*Sec[a + b*x]^2*Sqrt[Sin[a + b*x]])*Sqrt[Sin[a + b*x]])/(4*b)

Maple [A] (verified)

Time = 1.14 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.18

method	result	size
default	$\frac{-(-3\ln(1+\sqrt{\sin(bx+a)})+3\ln(\sqrt{\sin(bx+a)}-1)-6\arctan(\sqrt{\sin(bx+a)}))(\cos^2(bx+a))+4(\sqrt{\sin(bx+a)})}{8\cos(bx+a)^2b}$	73

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/8*(-(-3*ln(1+sin(b*x+a)^(1/2))+3*ln(sin(b*x+a)^(1/2)-1)-6*arctan(sin(b*x+a)^(1/2)))*cos(b*x+a)^2+4*sin(b*x+a)^(1/2))/cos(b*x+a)^2/b

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 131 vs. 2(50) = 100.

Time = 0.32 (sec) , antiderivative size = 131, normalized size of antiderivative = 2.11

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$$

$$= \frac{6 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) \cos(bx+a)^2 + 3 \cos(bx+a)^2 \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6 \sin(bx+a) - 2}{\cos(bx+a)^2 + 2 \sin(bx+a) - 2}\right)}{16 b \cos(bx+a)^2}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="fricas")

[Out] 1/16*(6*arctan(1/2*(sin(b*x + a) - 1)/sqrt(sin(b*x + a)))*cos(b*x + a)^2 + 3*cos(b*x + a)^2*log((cos(b*x + a)^2 + 4*(cos(b*x + a)^2 - sin(b*x + a) - 1)/sqrt(sin(b*x + a)) - 6*sin(b*x + a) - 2)/(cos(b*x + a)^2 + 2*sin(b*x + a) - 2)) + 8*sqrt(sin(b*x + a)))/(b*cos(b*x + a)^2)

Sympy [F]

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx = \int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$$

[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**3,x)

[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x)**3, x)

Maxima [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.05

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx$$

$$= \frac{\frac{4}{\left(\frac{1}{\sin(bx+a)^2} - 1\right) \sin(bx+a)^{\frac{3}{2}}} - 6 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) + 3 \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) - 3 \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{8 b}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="maxima")

[Out] 1/8*(4/((1/sin(b*x + a)^2 - 1)*sin(b*x + a)^(3/2)) - 6*arctan(1/sqrt(sin(b*x + a))) + 3*log(1/sqrt(sin(b*x + a)) + 1) - 3*log(1/sqrt(sin(b*x + a)) - 1))/b

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.06

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx = \frac{\frac{4\sqrt{\sin(bx+a)}}{\sin(bx+a)^2-1} - 6 \arctan\left(\sqrt{\sin(bx+a)}\right) - 3 \log\left(\sqrt{\sin(bx+a)}+1\right) + 3 \log\left(\left|\sqrt{\sin(bx+a)}-1\right|\right)}{8b}$$

```
[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^3,x, algorithm="giac")
```

```
[Out] -1/8*(4*sqrt(sin(b*x + a))/(sin(b*x + a)^2 - 1) - 6*arctan(sqrt(sin(b*x + a))) - 3*log(sqrt(sin(b*x + a)) + 1) + 3*log(abs(sqrt(sin(b*x + a)) - 1)))/b
```

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\csc(a+bx)} \sec^3(a+bx) dx = \int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a+bx)^3} dx$$

```
[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^3,x)
```

```
[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^3, x)
```

3.276 $\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1367
Rubi [A] (verified)	1367
Mathematica [C] (verified)	1369
Maple [A] (verified)	1369
Fricas [B] (verification not implemented)	1369
Sympy [F]	1370
Maxima [A] (verification not implemented)	1370
Giac [A] (verification not implemented)	1370
Mupad [F(-1)]	1371

Optimal result

Integrand size = 19, antiderivative size = 62

$$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\operatorname{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)}$$

[Out] 1/4*arctan(csc(b*x+a)^(1/2))/b+1/4*arctanh(csc(b*x+a)^(1/2))/b+1/2*sec(b*x+a)^2/b/csc(b*x+a)^(3/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2701, 294, 335, 218, 212, 209}

$$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\operatorname{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)}$$

[In] Int[Sec[a + b*x]^3/Sqrt[Csc[a + b*x]],x]

[Out] ArcTan[Sqrt[Csc[a + b*x]]]/(4*b) + ArcTanh[Sqrt[Csc[a + b*x]]]/(4*b) + Sec[a + b*x]^2/(2*b*Csc[a + b*x]^(3/2))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 335

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^{3/2}}{(-1+x^2)^2} dx, x, \csc(a+bx)\right)}{b} \\ &= \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x(-1+x^2)}} dx, x, \csc(a+bx)\right)}{4b} \\ &= \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \sqrt{\csc(a+bx)}\right)}{2b} \end{aligned}$$

$$\begin{aligned}
&= \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{\csc(a+bx)}\right)}{4b} \\
&= \frac{\arctan\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\text{arctanh}\left(\sqrt{\csc(a+bx)}\right)}{4b} + \frac{\sec^2(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.53

$$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2 \text{Hypergeometric2F1}\left(\frac{3}{4}, 2, \frac{7}{4}, \sin^2(a+bx)\right)}{3b \csc^{\frac{3}{2}}(a+bx)}$$

[In] Integrate[Sec[a + b*x]^3/Sqrt[Csc[a + b*x]],x]

[Out] (2*Hypergeometric2F1[3/4, 2, 7/4, Sin[a + b*x]^2])/(3*b*Csc[a + b*x]^(3/2))

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.15

method	result	size
default	$\frac{-(-\ln(1+\sqrt{\sin(bx+a)})+\ln(\sqrt{\sin(bx+a)}-1)+2\arctan(\sqrt{\sin(bx+a)}))(\cos^2(bx+a))+4(\sin^{\frac{3}{2}}(bx+a))}{8\cos(bx+a)^2b}$	71

[In] int(sec(b*x+a)^3/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/8*(-(-ln(1+sin(b*x+a)^(1/2))+ln(sin(b*x+a)^(1/2)-1)+2*arctan(sin(b*x+a)^(1/2))) * cos(b*x+a)^2+4*sin(b*x+a)^(3/2))/cos(b*x+a)^2/b

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(50) = 100.

Time = 0.31 (sec) , antiderivative size = 141, normalized size of antiderivative = 2.27

$$\int \frac{\sec^3(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{2 \arctan\left(\frac{\sin(bx+a)-1}{2\sqrt{\sin(bx+a)}}\right) \cos(bx+a)^2 - \cos(bx+a)^2 \log\left(\frac{\cos(bx+a)^2 + \frac{4(\cos(bx+a)^2 - \sin(bx+a)-1)}{\sqrt{\sin(bx+a)}} - 6\sin(bx+a)-2}{\cos(bx+a)^2 + 2\sin(bx+a)-2}\right)}{16b \cos(bx+a)^2}$$

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="fricas")

[Out]
$$-1/16*(2*\arctan(1/2*(\sin(b*x + a) - 1)/\sqrt{\sin(b*x + a)}))*\cos(b*x + a)^2 - \cos(b*x + a)^2*\log((\cos(b*x + a)^2 + 4*(\cos(b*x + a)^2 - \sin(b*x + a) - 1)/\sqrt{\sin(b*x + a)}) - 6*\sin(b*x + a) - 2)/(\cos(b*x + a)^2 + 2*\sin(b*x + a) - 2)) + 8*(\cos(b*x + a)^2 - 1)/\sqrt{\sin(b*x + a)}}/(b*\cos(b*x + a)^2)$$

Sympy [F]

$$\int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx$$

[In] integrate(sec(b*x+a)**3/csc(b*x+a)**(1/2),x)

[Out] Integral(sec(a + b*x)**3/sqrt(csc(a + b*x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.02

$$\int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{\frac{4}{(\frac{1}{\sin(bx+a)^2} - 1)\sqrt{\sin(bx+a)}} + 2 \arctan\left(\frac{1}{\sqrt{\sin(bx+a)}}\right) + \log\left(\frac{1}{\sqrt{\sin(bx+a)}} + 1\right) - \log\left(\frac{1}{\sqrt{\sin(bx+a)}} - 1\right)}{8b}$$

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out]
$$1/8*(4/((1/\sin(b*x + a)^2 - 1)*\sqrt{\sin(b*x + a)})) + 2*\arctan(1/\sqrt{\sin(b*x + a)}) + \log(1/\sqrt{\sin(b*x + a)} + 1) - \log(1/\sqrt{\sin(b*x + a)} - 1))/b$$

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.03

$$\int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{\frac{4 \sin(bx+a)^{\frac{3}{2}}}{\sin(bx+a)^2 - 1} + 2 \arctan\left(\sqrt{\sin(bx+a)}\right) - \log\left(\sqrt{\sin(bx+a)} + 1\right) + \log\left(\left|\sqrt{\sin(bx+a)} - 1\right|\right)}{8b}$$

[In] integrate(sec(b*x+a)^3/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out]
$$-1/8*(4*\sin(b*x + a)^(3/2)/(\sin(b*x + a)^2 - 1) + 2*\arctan(\sqrt{\sin(b*x + a)}) - \log(\sqrt{\sin(b*x + a)} + 1) + \log(\text{abs}(\sqrt{\sin(b*x + a)} - 1)))/b$$

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^3(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{1}{\cos(a + bx)^3 \sqrt{\frac{1}{\sin(a + bx)}}} dx$$

```
[In] int(1/(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2)), x)
```

```
[Out] int(1/(cos(a + b*x)^3*(1/sin(a + b*x))^(1/2)), x)
```

3.277 $\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx$

Optimal result	1372
Rubi [A] (verified)	1372
Mathematica [A] (verified)	1374
Maple [A] (verified)	1374
Fricas [C] (verification not implemented)	1374
Sympy [F]	1375
Maxima [F]	1375
Giac [F]	1375
Mupad [F(-1)]	1375

Optimal result

Integrand size = 19, antiderivative size = 92

$$\begin{aligned} & \int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx \\ &= \frac{5 \sec(a + bx)}{6b\sqrt{\csc(a + bx)}} + \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} \\ & \quad + \frac{5\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a - \frac{\pi}{2} + bx), 2\right) \sqrt{\sin(a + bx)}}{6b} \end{aligned}$$

[Out] $5/6*\sec(b*x+a)/b/\csc(b*x+a)^{(1/2)}+1/3*\sec(b*x+a)^3/b/\csc(b*x+a)^{(1/2)}-5/6*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\operatorname{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\csc(b*x+a)^{(1/2)}*\sin(b*x+a)^{(1/2)}/b$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2706, 3856, 2720}

$$\begin{aligned} & \int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx \\ &= \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{5 \sec(a + bx)}{6b\sqrt{\csc(a + bx)}} \\ & \quad + \frac{5\sqrt{\sin(a + bx)}\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a + bx - \frac{\pi}{2}), 2\right)}{6b} \end{aligned}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]*\operatorname{Sec}[a + b*x]^4, x]$

[Out] $(5 \operatorname{Sec}[a + b*x]) / (6*b*\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]) + \operatorname{Sec}[a + b*x]^3 / (3*b*\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]) + (5*\operatorname{Sqrt}[\operatorname{Csc}[a + b*x]]*\operatorname{EllipticF}[(a - \pi/2 + b*x)/2, 2]*\operatorname{Sqrt}[\operatorname{Sin}[a + b*x]]) / (6*b)$

Rule 2706

$\operatorname{Int}[(\operatorname{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\operatorname{sec}[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[a*b*(a*\operatorname{Csc}[e + f*x])^{(m-1)}*((b*\operatorname{Sec}[e + f*x])^{(n-1)})/(f*(n-1)), x] + \operatorname{Dist}[b^2*((m+n-2)/(n-1)), \operatorname{Int}[(a*\operatorname{Csc}[e + f*x])^m*(b*\operatorname{Sec}[e + f*x])^{(n-2)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, m\}, x \&\& \operatorname{GtQ}[n, 1] \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 2720

$\operatorname{Int}[1/\operatorname{Sqrt}[\operatorname{sin}[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \operatorname{Simp}[(2/d)*\operatorname{EllipticF}[(1/2)*(c - \pi/2 + d*x), 2], x] /;$ $\operatorname{FreeQ}\{c, d\}, x]$

Rule 3856

$\operatorname{Int}[(\operatorname{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Dist}[(b*\operatorname{Csc}[c + d*x])^n*\operatorname{Sin}[c + d*x]^n, \operatorname{Int}[1/\operatorname{Sin}[c + d*x]^n, x], x] /;$ $\operatorname{FreeQ}\{b, c, d\}, x \&\& \operatorname{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{5}{6} \int \sqrt{\csc(a + bx)} \sec^2(a + bx) dx \\ &= \frac{5 \sec(a + bx)}{6b\sqrt{\csc(a + bx)}} + \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{5}{12} \int \sqrt{\csc(a + bx)} dx \\ &= \frac{5 \sec(a + bx)}{6b\sqrt{\csc(a + bx)}} + \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} + \frac{1}{12} \left(5\sqrt{\csc(a + bx)}\sqrt{\sin(a + bx)} \right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{5 \sec(a + bx)}{6b\sqrt{\csc(a + bx)}} + \frac{\sec^3(a + bx)}{3b\sqrt{\csc(a + bx)}} \\ &\quad + \frac{5\sqrt{\csc(a + bx)} \operatorname{EllipticF}\left(\frac{1}{2}(a - \frac{\pi}{2} + bx), 2\right) \sqrt{\sin(a + bx)}}{6b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.53 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.70

$$\int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx$$

$$= \frac{\sqrt{\csc(a+bx)} \left(-5 \operatorname{EllipticF} \left(\frac{1}{4}(-2a + \pi - 2bx), 2 \right) \sqrt{\sin(a+bx)} + (5 + 2 \sec^2(a+bx)) \tan(a+bx) \right)}{6b}$$

[In] Integrate[Sqrt[Csc[a + b*x]]*Sec[a + b*x]^4,x]

[Out] (Sqrt[Csc[a + b*x]]*(-5*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]] + (5 + 2*Sec[a + b*x]^2)*Tan[a + b*x]))/(6*b)

Maple [A] (verified)

Time = 1.32 (sec) , antiderivative size = 168, normalized size of antiderivative = 1.83

method	result
default	$-\frac{\sqrt{(\cos^2(bx+a)) \sin(bx+a)} \left(5\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} F\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) (\cos^2(bx+a))+10(\cos^2(bx+a)) \right)}{12(\sin(bx+a)+1)(\sin(bx+a)-1)\sqrt{-\sin(bx+a)(\sin(bx+a)-1)(\sin(bx+a)+1)} \cos(bx+a)\sqrt{\sin(bx+a)}}$

[In] int(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] -1/12*(cos(b*x+a)^2*sin(b*x+a))^(1/2)*(5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))*cos(b*x+a)^2+10*cos(b*x+a)^2*sin(b*x+a)+4*sin(b*x+a))/(sin(b*x+a)+1)/(sin(b*x+a)-1)/(-sin(b*x+a)*(sin(b*x+a)-1)*(sin(b*x+a)+1))^(1/2)/cos(b*x+a)/sin(b*x+a)^(1/2)/b

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.07

$$\int \sqrt{\csc(a+bx)} \sec^4(a+bx) dx$$

$$= \frac{-5i\sqrt{2i} \cos(bx+a)^3 \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + 5i\sqrt{-2i} \cos(bx+a)^3 \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a))}{12b \cos(bx+a)^3}$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="fricas")

[Out] 1/12*(-5*I*sqrt(2*I)*cos(b*x + a)^3*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + 5*I*sqrt(-2*I)*cos(b*x + a)^3*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*(5*cos(b*x + a)^2 + 2)*sqrt(sin(b*x + a)))/(b*cos(b*x + a)^3)

Sympy [F]

$$\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx = \int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx$$

[In] integrate(csc(b*x+a)**(1/2)*sec(b*x+a)**4,x)

[Out] Integral(sqrt(csc(a + b*x))*sec(a + b*x)**4, x)

Maxima [F]

$$\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx = \int \sqrt{\csc(bx + a)} \sec(bx + a)^4 dx$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="maxima")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^4, x)

Giac [F]

$$\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx = \int \sqrt{\csc(bx + a)} \sec(bx + a)^4 dx$$

[In] integrate(csc(b*x+a)^(1/2)*sec(b*x+a)^4,x, algorithm="giac")

[Out] integrate(sqrt(csc(b*x + a))*sec(b*x + a)^4, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{\csc(a + bx)} \sec^4(a + bx) dx = \int \frac{\sqrt{\frac{1}{\sin(a+bx)}}}{\cos(a + bx)^4} dx$$

[In] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^4,x)

[Out] int((1/sin(a + b*x))^(1/2)/cos(a + b*x)^4, x)

3.278 $\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$

Optimal result	1376
Rubi [A] (verified)	1376
Mathematica [A] (verified)	1377
Maple [A] (verified)	1378
Fricas [C] (verification not implemented)	1378
Sympy [F]	1378
Maxima [F]	1379
Giac [F]	1379
Mupad [F(-1)]	1379

Optimal result

Integrand size = 19, antiderivative size = 92

$$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\sec(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} + \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{2b}$$

[Out] 1/2*sec(b*x+a)/b/csc(b*x+a)^(3/2)+1/3*sec(b*x+a)^3/b/csc(b*x+a)^(3/2)+1/2*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))*csc(b*x+a)^(1/2)*sin(b*x+a)^(1/2)/b

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {2706, 3856, 2719}

$$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \frac{\sec^3(a+bx)}{3b \csc^{\frac{3}{2}}(a+bx)} + \frac{\sec(a+bx)}{2b \csc^{\frac{3}{2}}(a+bx)} - \frac{\sqrt{\sin(a+bx)} \sqrt{\csc(a+bx)} E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{2b}$$

[In] Int[Sec[a + b*x]^4/Sqrt[Csc[a + b*x]],x]

[Out] Sec[a + b*x]/(2*b*Csc[a + b*x]^(3/2)) + Sec[a + b*x]^3/(3*b*Csc[a + b*x]^(3/2)) - (Sqrt[Csc[a + b*x]]*EllipticE[(a - Pi/2 + b*x)/2, 2]*Sqrt[Sin[a + b*x]])/(2*b)

Rule 2706


```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Dist[b^2*((m + n - 2)/(n - 1)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\sec^3(a + bx)}{3b \csc^{\frac{3}{2}}(a + bx)} + \frac{1}{2} \int \frac{\sec^2(a + bx)}{\sqrt{\csc(a + bx)}} dx \\
 &= \frac{\sec(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} + \frac{\sec^3(a + bx)}{3b \csc^{\frac{3}{2}}(a + bx)} - \frac{1}{4} \int \frac{1}{\sqrt{\csc(a + bx)}} dx \\
 &= \frac{\sec(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} + \frac{\sec^3(a + bx)}{3b \csc^{\frac{3}{2}}(a + bx)} - \frac{1}{4} \left(\sqrt{\csc(a + bx)} \sqrt{\sin(a + bx)} \right) \int \sqrt{\sin(a + bx)} dx \\
 &= \frac{\sec(a + bx)}{2b \csc^{\frac{3}{2}}(a + bx)} + \frac{\sec^3(a + bx)}{3b \csc^{\frac{3}{2}}(a + bx)} - \frac{\sqrt{\csc(a + bx)} E\left(\frac{1}{2}(a - \frac{\pi}{2} + bx) \mid 2\right) \sqrt{\sin(a + bx)}}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.83

$$\int \frac{\sec^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \frac{\cos(a + bx) \sqrt{\csc(a + bx)} \left(-3 + \sec^2(a + bx) + 2 \sec^4(a + bx) + 3 E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sec(a + bx) \right)}{6b}$$

```
[In] Integrate[Sec[a + b*x]^4/Sqrt[Csc[a + b*x]], x]
```

```
[Out] (Cos[a + b*x]*Sqrt[Csc[a + b*x]]*(-3 + Sec[a + b*x]^2 + 2*Sec[a + b*x]^4 + 3*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sec[a + b*x]*Sqrt[Sin[a + b*x]]))/(6*b)
```

Maple [A] (verified)

Time = 1.77 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.74

method	result
default	$\frac{6\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}E\left(\sqrt{\sin(bx+a)+1},\frac{\sqrt{2}}{2}\right)(\cos^2(bx+a))-3\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}}{12\cos(bx+a)^3\sqrt{\sin(bx+a)}b}$

[In] `int(sec(b*x+a)^4/csc(b*x+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{12} \frac{\cos(bx+a)^3 \sin(bx+a)^{(1/2)} (6(\sin(bx+a)+1)^{(1/2)} (-2\sin(bx+a)+2)^{(1/2)} (-\sin(bx+a))^{(1/2)} \text{EllipticE}((\sin(bx+a)+1)^{(1/2)}, 1/2\sqrt{2}) \cos(bx+a)^2 - 3(\sin(bx+a)+1)^{(1/2)} (-2\sin(bx+a)+2)^{(1/2)} (-\sin(bx+a))^{(1/2)} \text{EllipticF}((\sin(bx+a)+1)^{(1/2)}, 1/2\sqrt{2}) \cos(bx+a)^2 - 6\cos(bx+a)^4 + 2\cos(bx+a)^2 + 4)}{b}$$

Fricas [C] (verification not implemented)

Result contains higher order function than in optimal. Order 9 vs. order 4.

Time = 0.10 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.24

$$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx =$$

$$\frac{3\sqrt{2}i \cos(bx+a)^3 \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a))) + 3\sqrt{-2}i \cos(bx+a)^3 \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)))}{(b \cos(bx+a))^3}$$

[In] `integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="fricas")`

[Out]
$$-1/12 * (3 * \sqrt{2} * I * \cos(bx+a)^3 * \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + I * \sin(bx+a))) + 3 * \sqrt{-2} * I * \cos(bx+a)^3 * \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - I * \sin(bx+a)))) + 2 * (3 * \cos(bx+a)^4 - \cos(bx+a)^2 - 2) / \sqrt{\sin(bx+a)} / (b * \cos(bx+a))^3$$

Sympy [F]

$$\int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx = \int \frac{\sec^4(a+bx)}{\sqrt{\csc(a+bx)}} dx$$

[In] `integrate(sec(b*x+a)**4/csc(b*x+a)**(1/2),x)`

[Out] `Integral(sec(a + b*x)**4/sqrt(csc(a + b*x)), x)`

Maxima [F]

$$\int \frac{\sec^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec(bx + a)^4}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^4/sqrt(csc(b*x + a)), x)

Giac [F]

$$\int \frac{\sec^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{\sec(bx + a)^4}{\sqrt{\csc(bx + a)}} dx$$

[In] integrate(sec(b*x+a)^4/csc(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^4/sqrt(csc(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^4(a + bx)}{\sqrt{\csc(a + bx)}} dx = \int \frac{1}{\cos(a + bx)^4 \sqrt{\frac{1}{\sin(a + bx)}}} dx$$

[In] int(1/(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2)),x)

[Out] int(1/(cos(a + b*x)^4*(1/sin(a + b*x))^(1/2)), x)

3.279 $\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx$

Optimal result	1380
Rubi [A] (verified)	1380
Mathematica [A] (verified)	1381
Maple [F]	1381
Fricas [F]	1382
Sympy [F(-1)]	1382
Maxima [F]	1382
Giac [F]	1382
Mupad [F(-1)]	1383

Optimal result

Integrand size = 21, antiderivative size = 76

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \frac{d \sqrt{d \cos(a + bx)} \csc^{-1+p}(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p) \sqrt[4]{\cos^2(a + bx)}}$$

[Out] d*csc(b*x+a)^(-1+p)*hypergeom([-1/4, 1/2-1/2*p], [3/2-1/2*p], sin(b*x+a)^2)*(d*cos(b*x+a))^(1/2)/b/(1-p)/(cos(b*x+a)^2)^(1/4)

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \frac{d \sqrt{d \cos(a + bx)} \csc^{p-1}(a + bx) \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p) \sqrt[4]{\cos^2(a + bx)}}$$

[In] Int[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^p,x]

[Out] (d*Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[-1/4, (1 - p)/2, (3 - p)/2, Sin[a + b*x]^2])/(b*(1 - p)*(Cos[a + b*x]^2)^(1/4))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac

Part[(n - 1)/2]]*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FractionPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \text{integral} &= (\csc^p(a + bx) \sin^p(a + bx)) \int (d \cos(a + bx))^{3/2} \sin^{-p}(a + bx) dx \\ &= \frac{d \sqrt{d \cos(a + bx)} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p) \sqrt[4]{\cos^2(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 32.63 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.38

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \frac{2(d \cos(a + bx))^{5/2} \csc^{-1+p}(a + bx) (9 \text{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1}{2}(-1 + p), \frac{9}{4}, \cos^2(a + bx)\right) + 5 \cos^2(a + bx))}{45bd}$$

[In] Integrate[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^p,x]

[Out] (-2*(d*Cos[a + b*x])^(5/2)*Csc[a + b*x]^(-1 + p)*(9*Hypergeometric2F1[5/4, (-1 + p)/2, 9/4, Cos[a + b*x]^2] + 5*Cos[a + b*x]^2*Hypergeometric2F1[9/4, (1 + p)/2, 13/4, Cos[a + b*x]^2])*(Sin[a + b*x]^2)^((-1 + p)/2))/(45*b*d)

Maple [F]

$$\int (d \cos(bx + a))^{3/2} (\csc^p(bx + a)) dx$$

[In] int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x)

[Out] int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x)

Fricas [F]

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \int (d \cos(bx + a))^{\frac{3}{2}} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*d*csc(b*x + a)^p*cos(b*x + a), x)

Sympy [F(-1)]

Timed out.

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \text{Timed out}$$

[In] integrate((d*cos(b*x+a))**(3/2)*csc(b*x+a)**p,x)

[Out] Timed out

Maxima [F]

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \int (d \cos(bx + a))^{\frac{3}{2}} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^p, x)

Giac [F]

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \int (d \cos(bx + a))^{\frac{3}{2}} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^p,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^p, x)

Mupad [F(-1)]

Timed out.

$$\int (d \cos(a + bx))^{3/2} \csc^p(a + bx) dx = \int (d \cos(a + bx))^{3/2} \left(\frac{1}{\sin(a + bx)} \right)^p dx$$

```
[In] int((d*cos(a + b*x))^(3/2)*(1/sin(a + b*x))^p,x)
```

```
[Out] int((d*cos(a + b*x))^(3/2)*(1/sin(a + b*x))^p, x)
```

3.280 $\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$

Optimal result	1384
Rubi [A] (verified)	1384
Mathematica [A] (verified)	1385
Maple [F]	1385
Fricas [F]	1386
Sympy [F]	1386
Maxima [F]	1386
Giac [F]	1386
Mupad [F(-1)]	1387

Optimal result

Integrand size = 21, antiderivative size = 76

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$$

$$= \frac{d^4 \sqrt{\cos^2(a + bx)} \csc^{-1+p}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}}$$

[Out] $d*(\cos(b*x+a)^2)^{(1/4)}*\csc(b*x+a)^{-1+p}*\operatorname{hypergeom}([1/4, 1/2-1/2*p], [3/2-1/2*p], \sin(b*x+a)^2)/b/(1-p)/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$$

$$= \frac{d^4 \sqrt{\cos^2(a + bx)} \csc^{p-1}(a + bx) \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}}$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]*\operatorname{Csc}[a + b*x]^p, x]$

[Out] $(d*(\operatorname{Cos}[a + b*x]^2)^{(1/4)}*\operatorname{Csc}[a + b*x]^{-1 + p}*\operatorname{Hypergeometric2F1}[1/4, (1 - p)/2, (3 - p)/2, \operatorname{Sin}[a + b*x]^2])/(b*(1 - p)*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])$

Rule 2657

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b^{(2*\operatorname{IntPart}[(n - 1)/2] + 1)}*(b*\operatorname{Cos}[e + f*x])^{(2*\operatorname{Frac$

Part[(n - 1)/2]]*(a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \text{integral} &= (\csc^p(a + bx) \sin^p(a + bx)) \int \sqrt{d \cos(a + bx)} \sin^{-p}(a + bx) dx \\ &= \frac{d^4 \sqrt{\cos^2(a + bx)} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p)\sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 10.47 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.92

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \frac{2(d \cos(a + bx))^{3/2} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1+p}{2}, \frac{7}{4}, \cos^2(a + bx)\right) \sin^2(a + bx)^{\frac{1}{2}(-1+p)}}{3bd}$$

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^p,x]

[Out] (-2*(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[3/4, (1 + p)/2, 7/4, Cos[a + b*x]^2]*(Sin[a + b*x]^2)^((-1 + p)/2))/(3*b*d)

Maple [F]

$$\int \sqrt{d \cos(bx + a)} (\csc^p(bx + a)) dx$$

[In] int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x)

[Out] int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x)

Fricas [F]

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \int \sqrt{d \cos(bx + a)} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)

Sympy [F]

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx$$

[In] integrate((d*cos(b*x+a))**(1/2)*csc(b*x+a)**p,x)

[Out] Integral(sqrt(d*cos(a + b*x))*csc(a + b*x)**p, x)

Maxima [F]

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \int \sqrt{d \cos(bx + a)} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)

Giac [F]

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \int \sqrt{d \cos(bx + a)} \csc(bx + a)^p dx$$

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^p,x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^p, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{d \cos(a + bx)} \csc^p(a + bx) dx = \int \sqrt{d \cos(a + bx)} \left(\frac{1}{\sin(a + bx)} \right)^p dx$$

```
[In] int((d*cos(a + b*x))^(1/2)*(1/sin(a + b*x))^p,x)
```

```
[Out] int((d*cos(a + b*x))^(1/2)*(1/sin(a + b*x))^p, x)
```

$$3.281 \quad \int \frac{\csc^p(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal result	1388
Rubi [A] (verified)	1388
Mathematica [A] (verified)	1389
Maple [F]	1389
Fricas [F]	1390
Sympy [F]	1390
Maxima [F]	1390
Giac [F]	1390
Mupad [F(-1)]	1391

Optimal result

Integrand size = 21, antiderivative size = 76

$$\int \frac{\csc^p(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

$$= \frac{d \cos^2(a+bx)^{3/4} \csc^{-1+p}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{b(1-p)(d \cos(a+bx))^{3/2}}$$

[Out] $d*(\cos(b*x+a)^2)^{(3/4)}*\csc(b*x+a)^{-1+p}*\operatorname{hypergeom}([3/4, 1/2-1/2*p], [3/2-1/2*p], \sin(b*x+a)^2)/b/(1-p)/(d*\cos(b*x+a))^{(3/2)}$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int \frac{\csc^p(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

$$= \frac{d \cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{b(1-p)(d \cos(a+bx))^{3/2}}$$

[In] `Int[Csc[a + b*x]^p/Sqrt[d*Cos[a + b*x]], x]`

[Out] $(d*(\cos[a + b*x]^2)^{(3/4)}*\csc[a + b*x]^{-1 + p}*\operatorname{Hypergeometric2F1}[3/4, (1 - p)/2, (3 - p)/2, \sin[a + b*x]^2])/(b*(1 - p)*(d*\cos[a + b*x])^{(3/2)})$

Rule 2657

`Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac`

Part[(n - 1)/2]]*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \text{integral} &= (\csc^p(a + bx) \sin^p(a + bx)) \int \frac{\sin^{-p}(a + bx)}{\sqrt{d \cos(a + bx)}} dx \\ &= \frac{d \cos^2(a + bx)^{3/4} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{b(1-p)(d \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 10.54 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.89

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \frac{2\sqrt{d \cos(a + bx)} \csc^{1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1+p}{2}, \frac{5}{4}, \cos^2(a + bx)\right) \sin^2(a + bx)^{\frac{1+p}{2}}}{bd}$$

[In] Integrate[Csc[a + b*x]^p/Sqrt[d*Cos[a + b*x]], x]

[Out] (-2*Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^(1 + p)*Hypergeometric2F1[1/4, (1 + p)/2, 5/4, Cos[a + b*x]^2]*(Sin[a + b*x]^2)^((1 + p)/2))/(b*d)

Maple [F]

$$\int \frac{\csc^p(bx + a)}{\sqrt{d \cos(bx + a)}} dx$$

[In] int(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2), x)

[Out] int(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2), x)

Fricas [F]

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \int \frac{\csc^p(bx + a)}{\sqrt{d \cos(bx + a)}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d*cos(b*x + a)), x)

Sympy [F]

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx$$

[In] integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(1/2),x)

[Out] Integral(csc(a + b*x)**p/sqrt(d*cos(a + b*x)), x)

Maxima [F]

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \int \frac{\csc^p(bx + a)}{\sqrt{d \cos(bx + a)}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(csc(b*x + a)^p/sqrt(d*cos(b*x + a)), x)

Giac [F]

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \int \frac{\csc^p(bx + a)}{\sqrt{d \cos(bx + a)}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)^p/sqrt(d*cos(b*x + a)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\csc^p(a + bx)}{\sqrt{d \cos(a + bx)}} dx = \int \frac{\left(\frac{1}{\sin(a + bx)}\right)^p}{\sqrt{d \cos(a + bx)}} dx$$

```
[In] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(1/2), x)
```

```
[Out] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(1/2), x)
```

$$3.282 \quad \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal result	1392
Rubi [A] (verified)	1392
Mathematica [A] (verified)	1393
Maple [F]	1393
Fricas [F]	1393
Sympy [F]	1394
Maxima [F]	1394
Giac [F]	1394
Mupad [F(-1)]	1394

Optimal result

Integrand size = 21, antiderivative size = 78

$$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx = \frac{\sqrt[4]{\cos^2(a+bx)} \csc^{-1+p}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{bd(1-p)\sqrt{d \cos(a+bx)}}$$

[Out] $(\cos(b*x+a)^2)^{(1/4)} * \csc(b*x+a)^{-1+p} * \operatorname{hypergeom}([5/4, 1/2-1/2*p], [3/2-1/2*p], \sin(b*x+a)^2) / b/d/(1-p)/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A] (verified)

Time = 0.20 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{3/2}} dx = \frac{\sqrt[4]{\cos^2(a+bx)} \csc^{p-1}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{bd(1-p)\sqrt{d \cos(a+bx)}}$$

[In] $\operatorname{Int}[\operatorname{Csc}[a + b*x]^p / (d*\operatorname{Cos}[a + b*x])^{(3/2)}, x]$

[Out] $((\operatorname{Cos}[a + b*x]^2)^{(1/4)} * \operatorname{Csc}[a + b*x]^{-1+p} * \operatorname{Hypergeometric2F1}[5/4, (1-p)/2, (3-p)/2, \operatorname{Sin}[a + b*x]^2]) / (b*d*(1-p)*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])$

Rule 2657

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)} * ((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b^{(2*\operatorname{IntPart}[(n-1)/2] + 1)} * (b*\operatorname{Cos}[e + f*x])^{(2*\operatorname{FracPart}[(n-1)/2])} * ((a*\operatorname{Sin}[e + f*x])^{(m+1)}) / (a*f*(m+1)*(\operatorname{Cos}[e + f*x]^2)^{\operatorname{FracPart}[(n-1)/2]}) * \operatorname{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \operatorname{Sin}[e + f*x]^2], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, m, n\}, x]$

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\csc^p(a + bx) \sin^p(a + bx)) \int \frac{\sin^{-p}(a + bx)}{(d \cos(a + bx))^{3/2}} dx \\ &= \frac{\sqrt[4]{\cos^2(a + bx)} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{bd(1-p)\sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 10.64 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.87

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \frac{2 \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1+p}{2}, \frac{3}{4}, \cos^2(a + bx)\right) \sin^2(a + bx)^{\frac{1}{2}(-1+p)}}{bd\sqrt{d \cos(a + bx)}}$$

[In] Integrate[Csc[a + b*x]^p/(d*Cos[a + b*x])^(3/2), x]

[Out] (2*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[-1/4, (1 + p)/2, 3/4, Cos[a + b*x]^2]*(Sin[a + b*x]^2)^((-1 + p)/2))/(b*d*Sqrt[d*Cos[a + b*x]])

Maple [F]

$$\int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

[In] int(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x)

[Out] int(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x)

Fricas [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d^2*cos(b*x + a)^2), x)

Sympy [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

[In] integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(3/2),x)

[Out] Integral(csc(a + b*x)**p/(d*cos(a + b*x))**(3/2), x)

Maxima [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(3/2), x)

Giac [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{3/2}} dx = \int \frac{\left(\frac{1}{\sin(a + bx)}\right)^p}{(d \cos(a + bx))^{3/2}} dx$$

[In] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(3/2),x)

[Out] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(3/2), x)

$$3.283 \quad \int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal result	1395
Rubi [A] (verified)	1395
Mathematica [A] (verified)	1396
Maple [F]	1396
Fricas [F]	1396
Sympy [F(-1)]	1397
Maxima [F]	1397
Giac [F]	1397
Mupad [F(-1)]	1397

Optimal result

Integrand size = 21, antiderivative size = 78

$$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx = \frac{\cos^2(a+bx)^{3/4} \csc^{-1+p}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{7}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{bd(1-p)(d \cos(a+bx))^{3/2}}$$

[Out] $(\cos(b*x+a)^2)^{(3/4)} * \csc(b*x+a)^{-1+p} * \operatorname{hypergeom}([7/4, 1/2-1/2*p], [3/2-1/2*p], \sin(b*x+a)^2) / b/d/(1-p)/(d*\cos(b*x+a))^{(3/2)}$

Rubi [A] (verified)

Time = 0.19 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int \frac{\csc^p(a+bx)}{(d \cos(a+bx))^{5/2}} dx = \frac{\cos^2(a+bx)^{3/4} \csc^{p-1}(a+bx) \operatorname{Hypergeometric2F1}\left(\frac{7}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a+bx)\right)}{bd(1-p)(d \cos(a+bx))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Csc}[a + b*x]^p / (d*\operatorname{Cos}[a + b*x])^{(5/2)}, x]$

[Out] $((\operatorname{Cos}[a + b*x]^2)^{(3/4)} * \operatorname{Csc}[a + b*x]^{-1+p} * \operatorname{Hypergeometric2F1}[7/4, (1-p)/2, (3-p)/2, \operatorname{Sin}[a + b*x]^2]) / (b*d*(1-p)*(d*\operatorname{Cos}[a + b*x])^{(3/2)})$

Rule 2657

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] \rightarrow \operatorname{Simp}[b^{(2*\operatorname{IntPart}[(n-1)/2] + 1)}*(b*\operatorname{Cos}[e + f*x])^{(2*\operatorname{FracPart}[(n-1)/2])}*((a*\operatorname{Sin}[e + f*x])^{(m+1)})/(a*f^{(m+1)}*(\operatorname{Cos}[e + f*x]^2)^{\operatorname{FracPart}[(n-1)/2]})]*\operatorname{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \operatorname{Sin}[e + f*x]^2], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, m, n\}, x]$

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (\csc^p(a + bx) \sin^p(a + bx)) \int \frac{\sin^{-p}(a + bx)}{(d \cos(a + bx))^{5/2}} dx \\ &= \frac{\cos^2(a + bx)^{3/4} \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(\frac{7}{4}, \frac{1-p}{2}, \frac{3-p}{2}, \sin^2(a + bx)\right)}{bd(1-p)(d \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 10.66 (sec) , antiderivative size = 70, normalized size of antiderivative = 0.90

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \frac{2 \csc^{-1+p}(a + bx) \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1+p}{2}, \frac{1}{4}, \cos^2(a + bx)\right) \sin^2(a + bx)^{\frac{1}{2}(-1+p)}}{3bd(d \cos(a + bx))^{3/2}}$$

```
[In] Integrate[Csc[a + b*x]^p/(d*Cos[a + b*x])^(5/2), x]
```

```
[Out] (2*Csc[a + b*x]^(-1 + p)*Hypergeometric2F1[-3/4, (1 + p)/2, 1/4, Cos[a + b*x]^2]*(Sin[a + b*x]^2)^((-1 + p)/2))/(3*b*d*(d*Cos[a + b*x])^(3/2))
```

Maple [F]

$$\int \frac{\csc^p(bx + a)}{(d \cos(bx + a))^{5/2}} dx$$

```
[In] int(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x)
```

```
[Out] int(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x)
```

Fricas [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{5/2}} dx$$

```
[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(d*cos(b*x + a))*csc(b*x + a)^p/(d^3*cos(b*x + a)^3), x)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \text{Timed out}$$

[In] integrate(csc(b*x+a)**p/(d*cos(b*x+a))**(5/2), x)

[Out] Timed out

Maxima [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="maxima")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(5/2), x)

Giac [F]

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \int \frac{\csc(bx + a)^p}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

[In] integrate(csc(b*x+a)^p/(d*cos(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate(csc(b*x + a)^p/(d*cos(b*x + a))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\csc^p(a + bx)}{(d \cos(a + bx))^{5/2}} dx = \int \frac{\left(\frac{1}{\sin(a+bx)}\right)^p}{(d \cos(a + bx))^{5/2}} dx$$

[In] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(5/2), x)

[Out] int((1/sin(a + b*x))^p/(d*cos(a + b*x))^(5/2), x)

3.284 $\int \cos^m(e + fx) \csc^n(e + fx) dx$

Optimal result	1398
Rubi [A] (verified)	1398
Mathematica [C] (warning: unable to verify)	1399
Maple [F]	1400
Fricas [F]	1400
Sympy [F]	1400
Maxima [F]	1400
Giac [F]	1401
Mupad [F(-1)]	1401

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \cos^m(e + fx) \csc^n(e + fx) dx$$

$$= \frac{\cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] $\cos(f*x+e)^{-1+m}*(\cos(f*x+e)^2)^{(1/2-1/2*m)}*\csc(f*x+e)^{-1+n}*\operatorname{hypergeom}([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], \sin(f*x+e)^2)/f/(1-n)$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2667, 2657}

$$\int \cos^m(e + fx) \csc^n(e + fx) dx$$

$$= \frac{\cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[In] $\operatorname{Int}[\operatorname{Cos}[e + f*x]^m*\operatorname{Csc}[e + f*x]^n,x]$

[Out] $(\operatorname{Cos}[e + f*x]^{-1+m}*(\operatorname{Cos}[e + f*x]^2)^{((1-m)/2)}*\operatorname{Csc}[e + f*x]^{-1+n}*\operatorname{Hypergeometric2F1}[(1-m)/2, (1-n)/2, (3-n)/2, \operatorname{Sin}[e + f*x]^2])/f*(1-n)$

Rule 2657

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b^{(2*\operatorname{IntPart}[(n-1)/2] + 1)}*(b*\operatorname{Cos}[e + f*x])^{(2*\operatorname{Frac}$

```
Part[(n - 1)/2]]*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\text{integral} = (\csc^n(e + fx) \sin^n(e + fx)) \int \cos^m(e + fx) \sin^{-n}(e + fx) dx$$

$$= \frac{\cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 2.39 (sec) , antiderivative size = 312, normalized size of antiderivative = 3.67

$$\int \cos^m(e + fx) \csc^n(e + fx) dx =$$

$$\frac{2(-3 + n) \text{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)}{f(-1 + n) \left((-3 + n) \text{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}$$

```
[In] Integrate[Cos[e + f*x]^m*Csc[e + f*x]^n,x]
```

```
[Out] (-2*(-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, Tan[(e + f*x)/2]
^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*Cos[e + f*x]^m*Csc[e + f*x]^n*S
in[(e + f*x)/2])/(f*(-1 + n)*((-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3
/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 + 2*(
m*AppellF1[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan
[(e + f*x)/2]^2] + (1 + m - n)*AppellF1[3/2 - n/2, -m, 2 + m - n, 5/2 - n/2
, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2))
```

Maple [F]

$$\int (\cos^m (fx + e)) (\csc^n (fx + e)) dx$$

[In] `int(cos(f*x+e)^m*csc(f*x+e)^n,x)`

[Out] `int(cos(f*x+e)^m*csc(f*x+e)^n,x)`

Fricas [F]

$$\int \cos^m(e + fx) \csc^n(e + fx) dx = \int \cos (fx + e)^m \csc (fx + e)^n dx$$

[In] `integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="fricas")`

[Out] `integral(cos(f*x + e)^m*csc(f*x + e)^n, x)`

Sympy [F]

$$\int \cos^m(e + fx) \csc^n(e + fx) dx = \int \cos^m (e + fx) \csc^n (e + fx) dx$$

[In] `integrate(cos(f*x+e)**m*csc(f*x+e)**n,x)`

[Out] `Integral(cos(e + f*x)**m*csc(e + f*x)**n, x)`

Maxima [F]

$$\int \cos^m(e + fx) \csc^n(e + fx) dx = \int \cos (fx + e)^m \csc (fx + e)^n dx$$

[In] `integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="maxima")`

[Out] `integrate(cos(f*x + e)^m*csc(f*x + e)^n, x)`

Giac [F]

$$\int \cos^m(e + fx) \csc^n(e + fx) dx = \int \cos(fx + e)^m \csc(fx + e)^n dx$$

[In] integrate(cos(f*x+e)^m*csc(f*x+e)^n,x, algorithm="giac")

[Out] integrate(cos(f*x + e)^m*csc(f*x + e)^n, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^m(e + fx) \csc^n(e + fx) dx = \int \cos(e + fx)^m \left(\frac{1}{\sin(e + fx)} \right)^n dx$$

[In] int(cos(e + f*x)^m*(1/sin(e + f*x))^n,x)

[Out] int(cos(e + f*x)^m*(1/sin(e + f*x))^n, x)

3.285 $\int (a \cos(e + fx))^m \csc^n(e + fx) dx$

Optimal result	1402
Rubi [A] (verified)	1402
Mathematica [C] (warning: unable to verify)	1403
Maple [F]	1404
Fricas [F]	1404
Sympy [F]	1404
Maxima [F]	1404
Giac [F]	1405
Mupad [F(-1)]	1405

Optimal result

Integrand size = 19, antiderivative size = 88

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx$$

$$= \frac{a(a \cos(e + fx))^{-1+m} \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) \operatorname{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] a*(a*cos(f*x+e))^(-1+m)*(cos(f*x+e)²)^(1/2-1/2*m)*csc(f*x+e)⁽⁻¹⁺ⁿ⁾*hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], sin(f*x+e)²)/f/(1-n)

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2667, 2657}

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx$$

$$= \frac{a \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{n-1}(e + fx) (a \cos(e + fx))^{m-1} \operatorname{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[In] Int[(a*cos[e + f*x])^m*Csc[e + f*x]ⁿ,x]

[Out] (a*(a*cos[e + f*x])^(-1 + m)*(cos[e + f*x]²)^{((1 - m)/2)}*Csc[e + f*x]^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²)]/(f*(1 - n))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[b^{(2*IntPart[(n - 1)/2] + 1)}*(b*cos[e + f*x])^{(2*Frac}

```
Part[(n - 1)/2]]*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\text{integral} = (\csc^n(e + fx) \sin^n(e + fx)) \int (a \cos(e + fx))^m \sin^{-n}(e + fx) dx$$

$$= \frac{a(a \cos(e + fx))^{-1+m} \cos^2(e + fx)^{\frac{1-m}{2}} \csc^{-1+n}(e + fx) \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.00 (sec) , antiderivative size = 314, normalized size of antiderivative = 3.57

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx =$$

$$\frac{2(-3 + n) \text{AppellF1}\left(\frac{1}{2}, -\frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)}{f(-1 + n) \left((-3 + n) \text{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}$$

```
[In] Integrate[(a*Cos[e + f*x])^m*Csc[e + f*x]^n,x]
```

```
[Out] (-2*(-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, Tan[(e + f*x)/2]
^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*(a*Cos[e + f*x])^m*Csc[e + f*x]
^n*Sin[(e + f*x)/2])/(f*(-1 + n)*((-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m -
n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 +
2*(m*AppellF1[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2,
-Tan[(e + f*x)/2]^2] + (1 + m - n)*AppellF1[3/2 - n/2, -m, 2 + m - n, 5/2 -
n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2)
```

Maple [F]

$$\int (\cos(fx + e) a)^m (\csc^n(fx + e)) dx$$

[In] int((cos(f*x+e)*a)^m*csc(f*x+e)^n,x)

[Out] int((cos(f*x+e)*a)^m*csc(f*x+e)^n,x)

Fricas [F]

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \int (a \cos(fx + e))^m \csc^n(fx + e) dx$$

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="fricas")

[Out] integral((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

Sympy [F]

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \int (a \cos(e + fx))^m \csc^n(e + fx) dx$$

[In] integrate((a*cos(f*x+e))**m*csc(f*x+e)**n,x)

[Out] Integral((a*cos(e + f*x))**m*csc(e + f*x)**n, x)

Maxima [F]

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \int (a \cos(fx + e))^m \csc^n(fx + e) dx$$

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

Giac [F]

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \int (a \cos(fx + e))^m \csc(fx + e)^n dx$$

[In] integrate((a*cos(f*x+e))^m*csc(f*x+e)^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*csc(f*x + e)^n, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m \csc^n(e + fx) dx = \int (a \cos(e + fx))^m \left(\frac{1}{\sin(e + fx)} \right)^n dx$$

[In] int((a*cos(e + f*x))^m*(1/sin(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(1/sin(e + f*x))^n, x)

3.286 $\int \cos^m(e + fx)(b \csc(e + fx))^n dx$

Optimal result	1406
Rubi [A] (verified)	1406
Mathematica [C] (warning: unable to verify)	1407
Maple [F]	1408
Fricas [F]	1408
Sympy [F]	1408
Maxima [F]	1408
Giac [F]	1409
Mupad [F(-1)]	1409

Optimal result

Integrand size = 19, antiderivative size = 88

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx$$

$$= \frac{b \cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*cos(f*x+e)^(-1+m)*(cos(f*x+e)^2)^(1/2-1/2*m)*(b*csc(f*x+e))^(-1+n)*hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2667, 2657}

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx$$

$$= \frac{b \cos^{m-1}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[In] Int[Cos[e + f*x]^m*(b*Csc[e + f*x])^n,x]

[Out] (b*Cos[e + f*x]^(-1 + m)*(Cos[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(1 - n))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*Frac

```
Part[(n - 1)/2]]*(a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^Fr
acPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[
e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]
```

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\text{integral} = (b^2(b \csc(e + fx))^{-1+n}(b \sin(e + fx))^{-1+n}) \int \cos^m(e + fx)(b \sin(e + fx))^{-n} dx$$

$$= \frac{b \cos^{-1+m}(e + fx) \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 1.01 (sec) , antiderivative size = 314, normalized size of antiderivative = 3.57

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx =$$

$$\frac{2(-3 + n) \text{AppellF1}\left(\frac{1}{2}, -\frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)}{f(-1 + n) \left((-3 + n) \text{AppellF1}\left(\frac{1}{2} - \frac{n}{2}, -m, 1 + m - n, \frac{3}{2} - \frac{n}{2}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}$$

```
[In] Integrate[Cos[e + f*x]^m*(b*Csc[e + f*x])^n,x]
```

```
[Out] (-2*(-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m - n, 3/2 - n/2, Tan[(e + f*x)/2]
^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*Cos[e + f*x]^m*(b*Csc[e + f*x])
^n*Sin[(e + f*x)/2])/(f*(-1 + n)*((-3 + n)*AppellF1[1/2 - n/2, -m, 1 + m -
n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 +
2*(m*AppellF1[3/2 - n/2, 1 - m, 1 + m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2,
-Tan[(e + f*x)/2]^2] + (1 + m - n)*AppellF1[3/2 - n/2, -m, 2 + m - n, 5/2 -
n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2)
```

Maple [F]

$$\int (\cos^m(fx + e)) (b \csc(fx + e))^n dx$$

[In] `int(cos(f*x+e)^m*(b*csc(f*x+e))^n,x)`

[Out] `int(cos(f*x+e)^m*(b*csc(f*x+e))^n,x)`

Fricas [F]

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^m dx$$

[In] `integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="fricas")`

[Out] `integral((b*csc(f*x + e))^n*cos(f*x + e)^m, x)`

Sympy [F]

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(e + fx))^n \cos^m(e + fx) dx$$

[In] `integrate(cos(f*x+e)**m*(b*csc(f*x+e))**n,x)`

[Out] `Integral((b*csc(e + f*x))**n*cos(e + f*x)**m, x)`

Maxima [F]

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^m dx$$

[In] `integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*cos(f*x + e)^m, x)`

Giac [F]

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx = \int (b \csc(fx + e))^n \cos(fx + e)^m dx$$

[In] integrate(cos(f*x+e)^m*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^m, x)

Mupad [F(-1)]

Timed out.

$$\int \cos^m(e + fx)(b \csc(e + fx))^n dx = \int \cos(e + fx)^m \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

[In] int(cos(e + f*x)^m*(b/sin(e + f*x))^n,x)

[Out] int(cos(e + f*x)^m*(b/sin(e + f*x))^n, x)

3.287 $\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$

Optimal result	1410
Rubi [A] (verified)	1410
Mathematica [C] (warning: unable to verify)	1411
Maple [F]	1412
Fricas [F]	1412
Sympy [F]	1412
Maxima [F]	1412
Giac [F]	1413
Mupad [F(-1)]	1413

Optimal result

Integrand size = 21, antiderivative size = 91

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$$

$$= \frac{ab(a \cos(e + fx))^{-1+m} \cos^2(e + fx)^{\frac{1-m}{2}} (b \csc(e + fx))^{-1+n} \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] a*b*(a*cos(f*x+e))^(-1+m)*(cos(f*x+e)²)^(1/2-1/2*m)*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾*
hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], sin(f*x+e)²)/f/(1-n)

Rubi [A] (verified)

Time = 0.19 (sec), antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2657}

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$$

$$= \frac{ab \cos^2(e + fx)^{\frac{1-m}{2}} (a \cos(e + fx))^{m-1} (b \csc(e + fx))^{n-1} \text{Hypergeometric2F1}\left(\frac{1-m}{2}, \frac{1-n}{2}, \frac{3-n}{2}, \sin^2(e + fx)\right)}{f(1-n)}$$

[In] Int[(a*cos[e + f*x])^m*(b*Csc[e + f*x])ⁿ,x]

[Out] (a*b*(a*cos[e + f*x])^(-1 + m)*(cos[e + f*x]²)^{((1 - m)/2)}*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²]/(f*(1 - n))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b^{(2*IntPart[(n - 1)/2] + 1)}*(b*cos[e + f*x])^{(2*Frac}

Maple [F]

$$\int (\cos(fx + e) a)^m (b \csc(fx + e))^n dx$$

```
[In] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^n,x)
```

```
[Out] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^n,x)
```

Fricas [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = \int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

```
[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="fricas")
```

```
[Out] integral((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)
```

Sympy [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = \int (a \cos(e + fx))^m (b \csc(e + fx))^n dx$$

```
[In] integrate((a*cos(f*x+e))**m*(b*csc(f*x+e))**n,x)
```

```
[Out] Integral((a*cos(e + f*x))**m*(b*csc(e + f*x))**n, x)
```

Maxima [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = \int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

```
[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="maxima")
```

```
[Out] integrate((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)
```

Giac [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = \int (a \cos(fx + e))^m (b \csc(fx + e))^n dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m*(b*csc(f*x + e))^n, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^n dx = \int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^n,x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^n, x)

3.288 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx$

Optimal result	1414
Rubi [A] (verified)	1414
Mathematica [A] (verified)	1415
Maple [F]	1416
Fricas [F]	1416
Sympy [F(-1)]	1416
Maxima [F]	1416
Giac [F]	1417
Mupad [F(-1)]	1417

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx = \frac{b^3 (a \cos(e + fx))^{1+m} \sqrt{b \csc(e + fx)} \operatorname{Hypergeometric2F1}\left(\frac{9}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sqrt[4]{\sin^2(e + fx)}}{af(1+m)}$$

[Out] $-b^3*(a*\cos(f*x+e))^{(1+m)}*\operatorname{hypergeom}([9/4, 1/2+1/2*m], [3/2+1/2*m], \cos(f*x+e)^2)*(\sin(f*x+e)^2)^{(1/4)}*(b*\csc(f*x+e))^{(1/2)}/a/f/(1+m)$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2656}

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx = \frac{b \sin^2(e + fx)^{5/4} (b \csc(e + fx))^{5/2} (a \cos(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{9}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e + fx)\right)}{af(m+1)}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e + f*x])^m*(b*\operatorname{Csc}[e + f*x])^{(7/2)}, x]$

[Out] $-((b*(a*\operatorname{Cos}[e + f*x])^{(1+m)}*(b*\operatorname{Csc}[e + f*x])^{(5/2)}*\operatorname{Hypergeometric2F1}[9/4, (1+m)/2, (3+m)/2, \operatorname{Cos}[e + f*x]^2]*(\operatorname{Sin}[e + f*x]^2)^{(5/4)})/(a*f*(1+m))$

Rule 2656

```
Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]
```

Rule 2667

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \text{integral} &= (b^2(b \csc(e + fx))^{5/2}(b \sin(e + fx))^{5/2}) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{7/2}} dx \\ &= \frac{b(a \cos(e + fx))^{1+m}(b \csc(e + fx))^{5/2} \text{Hypergeometric2F1}\left(\frac{9}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)}{af(1+m)} \end{aligned}$$

Mathematica [A] (verified)

Time = 16.30 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.21

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx = \frac{2ab(a \cos(e + fx))^{-1+m} (-\cot^2(e + fx))^{\frac{1-m}{2}} (b \csc(e + fx))^{5/2} \text{Hypergeometric2F1}\left(\frac{1}{4}(7 - 2m), \frac{1-m}{2}, \frac{11-2m}{4}, \csc^2(e + fx)\right)}{f(-7 + 2m)}$$

```
[In] Integrate[(a*Cos[e + f*x])^m*(b*Csc[e + f*x])^(7/2),x]
```

```
[Out] (2*a*b*(a*Cos[e + f*x])^(-1 + m)*(-Cot[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(5/2)*Hypergeometric2F1[(7 - 2*m)/4, (1 - m)/2, (11 - 2*m)/4, Csc[e + f*x]^2])/(f*(-7 + 2*m))
```

Maple [F]

$$\int (\cos (fx + e) a)^m (b \csc (fx + e))^{\frac{7}{2}} dx$$

[In] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(7/2),x)

[Out] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(7/2),x)

Fricas [F]

$$\int (a \cos (e + fx))^m (b \csc (e + fx))^{7/2} dx = \int (b \csc (fx + e))^{\frac{7}{2}} (a \cos (fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b^3*csc(f*x + e)^3, x)

Sympy [F(-1)]

Timed out.

$$\int (a \cos (e + fx))^m (b \csc (e + fx))^{7/2} dx = \text{Timed out}$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x)

[Out] Timed out

Maxima [F]

$$\int (a \cos (e + fx))^m (b \csc (e + fx))^{7/2} dx = \int (b \csc (fx + e))^{\frac{7}{2}} (a \cos (fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^(7/2)*(a*cos(f*x + e))^m, x)

Giac [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx = \int (b \csc(fx + e))^{7/2} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(7/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^(7/2)*(a*cos(f*x + e))^m, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{7/2} dx = \int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^{7/2} dx$$

[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(7/2),x)

[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(7/2), x)

3.289 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx$

Optimal result	1418
Rubi [A] (verified)	1418
Mathematica [A] (verified)	1419
Maple [F]	1419
Fricas [F]	1420
Sympy [F(-1)]	1420
Maxima [F]	1420
Giac [F]	1420
Mupad [F(-1)]	1421

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \frac{b(a \cos(e + fx))^{1+m} (b \csc(e + fx))^{3/2} \operatorname{Hypergeometric2F1}\left(\frac{7}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)^{3/4}}{af(1+m)}$$

[Out] -b*(a*cos(f*x+e))^(1+m)*(b*csc(f*x+e))^(3/2)*hypergeom([7/4, 1/2+1/2*m], [3/2+1/2*m], cos(f*x+e)^2)*(sin(f*x+e)^2)^(3/4)/a/f/(1+m)

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2656}

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \frac{b \sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{7}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e + fx)\right)}{af(m+1)}$$

[In] Int[(a*cos[e + f*x])^m*(b*csc[e + f*x])^(5/2),x]

[Out] -((b*(a*cos[e + f*x])^(1 + m)*(b*csc[e + f*x])^(3/2)*Hypergeometric2F1[7/4, (1 + m)/2, (3 + m)/2, Cos[e + f*x]^2]*(Sin[e + f*x]^2)^(3/4))/(a*f*(1 + m))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sine[e + f*x])^(2*F

```

racPart[(n - 1)/2])*((a*cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)
^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, C
os[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

```

Rule 2667

```

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*sin[e + f*x])^m/(b*cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= (b^2(b \csc(e + fx))^{3/2}(b \sin(e + fx))^{3/2}) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{5/2}} dx \\
&= \frac{b(a \cos(e + fx))^{1+m}(b \csc(e + fx))^{3/2} \text{Hypergeometric2F1}\left(\frac{7}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)}{af(1+m)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 10.76 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.24

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \frac{2ab(a \cos(e + fx))^{-1+m} (-\cot^2(e + fx))^{\frac{1-m}{2}} (b \csc(e + fx))^{3/2} \text{Hypergeometric2F1}\left(\frac{1}{4}(5 - 2m), \frac{1-m}{2}, \frac{9-2m}{4}, \csc^2(e + fx)\right)}{f(-5 + 2m)}$$

```
[In] Integrate[(a*cos[e + f*x])^m*(b*Csc[e + f*x])^(5/2),x]
```

```
[Out] (2*a*b*(a*cos[e + f*x])^(-1 + m)*(-Cot[e + f*x]^2)^((1 - m)/2)*(b*Csc[e + f*x])^(3/2)*Hypergeometric2F1[(5 - 2*m)/4, (1 - m)/2, (9 - 2*m)/4, Csc[e + f*x]^2])/(f*(-5 + 2*m))
```

Maple [F]

$$\int (\cos(fx + e) a)^m (b \csc(fx + e))^{\frac{5}{2}} dx$$

```
[In] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(5/2),x)
```

```
[Out] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(5/2),x)
```

Fricas [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \int (b \csc(fx + e))^{5/2} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b^2*csc(f*x + e)^2, x)

Sympy [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \text{Timed out}$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x)

[Out] Timed out

Maxima [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \int (b \csc(fx + e))^{5/2} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^(5/2)*(a*cos(f*x + e))^m, x)

Giac [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \int (b \csc(fx + e))^{5/2} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^(5/2)*(a*cos(f*x + e))^m, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{5/2} dx = \int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^{5/2} dx$$

```
[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(5/2),x)
```

```
[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(5/2), x)
```

3.290 $\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx$

Optimal result	1422
Rubi [A] (verified)	1422
Mathematica [A] (verified)	1423
Maple [F]	1423
Fricas [F]	1424
Sympy [F(-1)]	1424
Maxima [F]	1424
Giac [F]	1424
Mupad [F(-1)]	1425

Optimal result

Integrand size = 23, antiderivative size = 76

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \frac{b(a \cos(e + fx))^{1+m} \sqrt{b \csc(e + fx)} \operatorname{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sqrt[4]{\sin^2(e + fx)}}{af(1+m)}$$

[Out] $-b*(a*\cos(f*x+e))^{(1+m)}*\operatorname{hypergeom}\left(\left[\frac{5}{4}, 1/2+1/2*m\right], \left[\frac{3}{2}+1/2*m\right], \cos(f*x+e)^2\right)*(\sin(f*x+e)^2)^{(1/4)}*(b*\csc(f*x+e))^{(1/2)}/a/f/(1+m)$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2656}

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \frac{b^4 \sqrt{\sin^2(e + fx)} \sqrt{b \csc(e + fx)} (a \cos(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{5}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e + fx)\right)}{af(m+1)}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e + f*x])^m*(b*\operatorname{Csc}[e + f*x])^{(3/2)}, x]$

[Out] $-((b*(a*\operatorname{Cos}[e + f*x])^{(1+m)}*\operatorname{Sqrt}[b*\operatorname{Csc}[e + f*x]]*\operatorname{Hypergeometric2F1}[5/4, (1+m)/2, (3+m)/2, \operatorname{Cos}[e + f*x]^2]*(\operatorname{Sin}[e + f*x]^2)^{(1/4)})/(a*f*(1+m))$

Rule 2656

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}), x_Symbol] \rightarrow \operatorname{Simp}[(-b^{(2*\operatorname{IntPart}[(n-1)/2] + 1)}*(b*\operatorname{Sin}[e + f*x])^{(2*F$

```

racPart[(n - 1)/2])*((a*cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)
^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, C
os[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

```

Rule 2667

```

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[b^2*(b*cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1)
, Int[(a*sin[e + f*x])^m/(b*cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m
, n}, x] && !IntegerQ[m] && !IntegerQ[n]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \left(b^2 \sqrt{b \csc(e + fx)} \sqrt{b \sin(e + fx)} \right) \int \frac{(a \cos(e + fx))^m}{(b \sin(e + fx))^{3/2}} dx \\
&= \\
&= \frac{b(a \cos(e + fx))^{1+m} \sqrt{b \csc(e + fx)} \text{Hypergeometric2F1}\left(\frac{5}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sqrt[4]{\sin^2(e + fx)}}{af(1+m)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 9.29 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.24

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \frac{2ab(a \cos(e + fx))^{-1+m} (-\cot^2(e + fx))^{\frac{1-m}{2}} \sqrt{b \csc(e + fx)} \text{Hypergeometric2F1}\left(\frac{1}{4}(3 - 2m), \frac{1-m}{2}, \frac{7-2m}{4}, \csc^2(e + fx)\right)}{f(-3 + 2m)}$$

```
[In] Integrate[(a*cos[e + f*x])^m*(b*Csc[e + f*x])^(3/2),x]
```

```
[Out] (2*a*b*(a*cos[e + f*x])^(-1 + m)*(-Cot[e + f*x]^2)^((1 - m)/2)*Sqrt[b*Csc[e + f*x]]*Hypergeometric2F1[(3 - 2*m)/4, (1 - m)/2, (7 - 2*m)/4, Csc[e + f*x]^2])/(f*(-3 + 2*m))
```

Maple [F]

$$\int (\cos(fx + e) a)^m (b \csc(fx + e))^{\frac{3}{2}} dx$$

```
[In] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(3/2),x)
```

```
[Out] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(3/2),x)
```

Fricas [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \int (b \csc(fx + e))^{\frac{3}{2}} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m*b*csc(f*x + e), x)

Sympy [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \text{Timed out}$$

[In] integrate((a*cos(f*x+e))**m*(b*csc(f*x+e))**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \int (b \csc(fx + e))^{\frac{3}{2}} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^(3/2)*(a*cos(f*x + e))^m, x)

Giac [F]

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \int (b \csc(fx + e))^{\frac{3}{2}} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^(3/2)*(a*cos(f*x + e))^m, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + fx))^m (b \csc(e + fx))^{3/2} dx = \int (a \cos(e + fx))^m \left(\frac{b}{\sin(e + fx)} \right)^{3/2} dx$$

```
[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(3/2),x)
```

```
[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(3/2), x)
```

3.291 $\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx$

Optimal result	1426
Rubi [A] (verified)	1426
Mathematica [A] (verified)	1427
Maple [F]	1427
Fricas [F]	1428
Sympy [F]	1428
Maxima [F]	1428
Giac [F]	1428
Mupad [F(-1)]	1429

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \frac{(a \cos(e + fx))^{1+m} (b \csc(e + fx))^{3/2} \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)^{3/4}}{abf(1+m)}$$

[Out] $-(a \cos(fx+e))^{(1+m)} (b \csc(fx+e))^{(3/2)} \operatorname{hypergeom}\left(\left[\frac{3}{4}, \frac{1}{2} + \frac{1}{2}m\right], \left[\frac{3}{2} + \frac{1}{2}m\right], \cos(fx+e)^2\right) (\sin(fx+e)^2)^{(3/4)} / a/b/f/(1+m)$

Rubi [A] (verified)

Time = 0.10 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2656}

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \frac{\sin^2(e + fx)^{3/4} (b \csc(e + fx))^{3/2} (a \cos(e + fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{3}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e + fx)\right)}{abf(m+1)}$$

[In] $\operatorname{Int}[(a \cos[e + fx])^m \sqrt{b \csc[e + fx]}], x]$

[Out] $-\left(\left(a \cos[e + fx]\right)^{(1+m)} (b \csc[e + fx])^{(3/2)} \operatorname{Hypergeometric2F1}\left[\frac{3}{4}, (1+m)/2, (3+m)/2, \cos[e + fx]^2\right] (\sin[e + fx]^2)^{(3/4)}\right) / (a*b*f*(1+m))$

Rule 2656

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.))^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \operatorname{Simp}[(-b^{(2*\operatorname{IntPart}[(n-1)/2] + 1)}*(b*\sin[e + fx])^{(2*F$

```

racPart[(n - 1)/2])*((a*cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)
^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, C
os[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

```

Rule 2666

```

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[(1/b^2)*(b*cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n
+ 1), Int[(a*sin[e + f*x])^m/(b*cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e,
f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{((b \csc(e + fx))^{3/2} (b \sin(e + fx))^{3/2}) \int \frac{(a \cos(e + fx))^m}{\sqrt{b \sin(e + fx)}} dx}{b^2} \\
&= \frac{(a \cos(e + fx))^{1+m} (b \csc(e + fx))^{3/2} \text{Hypergeometric2F1}\left(\frac{3}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sin^2(e + fx)}{abf(1+m)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 8.90 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.23

$$\begin{aligned}
&\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx \\
&= \frac{2(a \cos(e + fx))^m (-\cot^2(e + fx))^{\frac{1-m}{2}} \sqrt{b \csc(e + fx)} \text{Hypergeometric2F1}\left(\frac{1}{4}(1 - 2m), \frac{1-m}{2}, \frac{1}{4}(5 - 2m), \cos^2(e + fx)\right)}{f(-1 + 2m)}
\end{aligned}$$

```
[In] Integrate[(a*cos[e + f*x])^m*Sqrt[b*Csc[e + f*x]],x]
```

```
[Out] (2*(a*cos[e + f*x])^m*(-Cot[e + f*x]^2)^((1 - m)/2)*Sqrt[b*Csc[e + f*x]]*Hypergeometric2F1[(1 - 2*m)/4, (1 - m)/2, (5 - 2*m)/4, Csc[e + f*x]^2]*Tan[e + f*x])/(f*(-1 + 2*m))
```

Maple [F]

$$\int (\cos(fx + e) a)^m \sqrt{b \csc(fx + e)} dx$$

```
[In] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(1/2),x)
```

```
[Out] int((cos(f*x+e)*a)^m*(b*csc(f*x+e))^(1/2),x)
```

Fricas [F]

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \int \sqrt{b \csc(fx + e)} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

Sympy [F]

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx$$

[In] integrate((a*cos(f*x+e))**m*(b*csc(f*x+e))**(1/2),x)

[Out] Integral((a*cos(e + f*x))**m*sqrt(b*csc(e + f*x)), x)

Maxima [F]

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \int \sqrt{b \csc(fx + e)} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

Giac [F]

$$\int (a \cos(e + fx))^m \sqrt{b \csc(e + fx)} dx = \int \sqrt{b \csc(fx + e)} (a \cos(fx + e))^m dx$$

[In] integrate((a*cos(f*x+e))^m*(b*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m, x)

Mupad [F(-1)]

Timed out.

$$\int (a \cos(e + f x))^m \sqrt{b \csc(e + f x)} dx = \int (a \cos(e + f x))^m \sqrt{\frac{b}{\sin(e + f x)}} dx$$

```
[In] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(1/2),x)
```

```
[Out] int((a*cos(e + f*x))^m*(b/sin(e + f*x))^(1/2), x)
```

3.292 $\int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx$

Optimal result	1430
Rubi [A] (verified)	1430
Mathematica [C] (warning: unable to verify)	1431
Maple [F]	1432
Fricas [F]	1432
Sympy [F]	1432
Maxima [F]	1432
Giac [F]	1433
Mupad [F(-1)]	1433

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx = \frac{(a \cos(e+fx))^{1+m} \sqrt{b \csc(e+fx)} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right) \sqrt[4]{\sin^2(e+fx)}}{abf(1+m)}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*\operatorname{hypergeom}([1/4, 1/2+1/2*m], [3/2+1/2*m], \cos(f*x+e)^2)*(\sin(f*x+e)^2)^{(1/4)}*(b*\csc(f*x+e))^{(1/2)}/a/b/f/(1+m)$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2656}

$$\int \frac{(a \cos(e+fx))^m}{\sqrt{b \csc(e+fx)}} dx = \frac{\sqrt[4]{\sin^2(e+fx)} \sqrt{b \csc(e+fx)} (a \cos(e+fx))^{m+1} \operatorname{Hypergeometric2F1}\left(\frac{1}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{abf(m+1)}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e+f*x])^m/\operatorname{Sqrt}[b*\operatorname{Csc}[e+f*x]],x]$

[Out] $-\left(\left(a*\operatorname{Cos}[e+f*x]\right)^{(1+m)}*\operatorname{Sqrt}[b*\operatorname{Csc}[e+f*x]]*\operatorname{Hypergeometric2F1}\left[\frac{1}{4}, \left(1+m\right)/2, \left(3+m\right)/2, \operatorname{Cos}[e+f*x]^2\right]*\left(\operatorname{Sin}[e+f*x]^2\right)^{(1/4)}\right)/\left(a*b*f*(1+m)\right)$

Rule 2656

```
Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*SIN[e + f*x])^(2*FracPart[(n - 1)/2])*((a*cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(SIN[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]
```

Rule 2666

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[(1/b^2)*(b*cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*SIN[e + f*x])^m/(b*cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]
```

Rubi steps

$$\text{integral} = \frac{\left(\sqrt{b \csc(e + fx)} \sqrt{b \sin(e + fx)}\right) \int (a \cos(e + fx))^m \sqrt{b \sin(e + fx)} dx}{b^2}$$

$$= \frac{(a \cos(e + fx))^{1+m} \sqrt{b \csc(e + fx)} \text{Hypergeometric2F1}\left(\frac{1}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right) \sqrt[4]{\sin^2(e + fx)}}{abf(1+m)}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

Time = 28.33 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.88

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx$$

$$= \frac{3f(b \csc(e + fx))^{3/2} \left(7 \text{AppellF1}\left(\frac{3}{4}, -m, \frac{3}{2} + m, \frac{7}{4}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 2(2m \text{AppellF1}\left(\frac{3}{4}, -m, \frac{3}{2} + m, \frac{7}{4}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}{14b \text{AppellF1}\left(\frac{3}{4}, -m, \frac{3}{2} + m, \frac{7}{4}, \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)}$$

```
[In] Integrate[(a*cos[e + f*x])^m/Sqrt[b*Csc[e + f*x]],x]
```

```
[Out] (14*b*AppellF1[3/4, -m, 3/2 + m, 7/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(a*cos[e + f*x])^m)/(3*f*(b*Csc[e + f*x])^(3/2)*(7*AppellF1[3/4, -m, 3/2 + m, 7/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*(2*m*AppellF1[7/4, 1 - m, 3/2 + m, 11/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + (3 + 2*m)*AppellF1[7/4, -m, 5/2 + m, 11/4, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]))*Tan[(e + f*x)/2]^2)
```

Maple [F]

$$\int \frac{(\cos(fx + e) a)^m}{\sqrt{b \csc(fx + e)}} dx$$

[In] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(1/2),x)

[Out] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(1/2),x)

Fricas [F]

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx = \int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b*csc(f*x + e)), x)

Sympy [F]

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx = \int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

[In] integrate((a*cos(f*x+e))**m/(b*csc(f*x+e))**(1/2),x)

[Out] Integral((a*cos(e + f*x))**m/sqrt(b*csc(e + f*x)), x)

Maxima [F]

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx = \int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/sqrt(b*csc(f*x + e)), x)

Giac [F]

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx = \int \frac{(a \cos(fx + e))^m}{\sqrt{b \csc(fx + e)}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/sqrt(b*csc(f*x + e)), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(a \cos(e + fx))^m}{\sqrt{b \csc(e + fx)}} dx = \int \frac{(a \cos(e + fx))^m}{\sqrt{\frac{b}{\sin(e+fx)}}} dx$$

[In] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(1/2),x)

[Out] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(1/2), x)

3.293 $\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx$

Optimal result	1434
Rubi [A] (verified)	1434
Mathematica [A] (verified)	1435
Maple [F]	1435
Fricas [F]	1436
Sympy [F]	1436
Maxima [F]	1436
Giac [F]	1436
Mupad [F(-1)]	1437

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx = \frac{(a \cos(e+fx))^{1+m} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right)}{abf(1+m)\sqrt{b \csc(e+fx)}\sqrt[4]{\sin^2(e+fx)}}$$

[Out] $-(a*\cos(f*x+e))^{(1+m)}*\operatorname{hypergeom}([-1/4, 1/2+1/2*m], [3/2+1/2*m], \cos(f*x+e)^2)/a/b/f/(1+m)/(\sin(f*x+e)^2)^{(1/4)}/(b*\csc(f*x+e))^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2656}

$$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{3/2}} dx = \frac{(a \cos(e+fx))^{m+1} \operatorname{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{abf(m+1)\sqrt[4]{\sin^2(e+fx)}\sqrt{b \csc(e+fx)}}$$

[In] $\operatorname{Int}[(a*\operatorname{Cos}[e+f*x])^m/(b*\operatorname{Csc}[e+f*x])^{(3/2)}, x]$

[Out] $-\left(\left(\left(a*\operatorname{Cos}[e+f*x]\right)^{(1+m)}*\operatorname{Hypergeometric2F1}\left[-1/4, (1+m)/2, (3+m)/2, \operatorname{Cos}[e+f*x]^2\right]\right)/(a*b*f*(1+m)*\operatorname{Sqrt}[b*\operatorname{Csc}[e+f*x]]*(\operatorname{Sin}[e+f*x]^2)^{(1/4)})$

Rule 2656

```
Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]
```

Rule 2666

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[(1/b^2)*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (a \cos(e + fx))^m (b \sin(e + fx))^{3/2} dx}{b^2 \sqrt{b \csc(e + fx)} \sqrt{b \sin(e + fx)}} \\ &= -\frac{(a \cos(e + fx))^{1+m} \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{abf(1+m) \sqrt{b \csc(e + fx)} \sqrt[4]{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 11.13 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.49

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \frac{2a(a \cos(e + fx))^{-1+m} \cos(2(e + fx)) (-\cot^2(e + fx))^{\frac{1-m}{2}} \text{Hypergeometric2F1}\left(-\frac{1}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{bf(3+2m) \sqrt{b \csc(e + fx)} (-2 + \csc^2(e + fx))}$$

```
[In] Integrate[(a*Cos[e + f*x])^m/(b*Csc[e + f*x])^(3/2), x]
```

```
[Out] (2*a*(a*Cos[e + f*x])^(-1 + m)*Cos[2*(e + f*x)]*(-Cot[e + f*x]^2)^((1 - m)/2)*Hypergeometric2F1[(-3 - 2*m)/4, (1 - m)/2, (1 - 2*m)/4, Csc[e + f*x]^2])/(b*f*(3 + 2*m)*Sqrt[b*Csc[e + f*x]]*(-2 + Csc[e + f*x]^2))
```

Maple [F]

$$\int \frac{(\cos(fx + e) a)^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

```
[In] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(3/2), x)
```

```
[Out] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(3/2), x)
```

Fricas [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b^2*csc(f*x + e)^2), x)

Sympy [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{\frac{3}{2}}} dx$$

[In] integrate((a*cos(f*x+e))**m/(b*csc(f*x+e))**(3/2),x)

[Out] Integral((a*cos(e + f*x))**m/(b*csc(e + f*x))**(3/2), x)

Maxima [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(3/2), x)

Giac [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{3/2}} dx = \int \frac{(a \cos(e + fx))^m}{\left(\frac{b}{\sin(e + fx)}\right)^{3/2}} dx$$

```
[In] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(3/2),x)
```

```
[Out] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(3/2), x)
```

$$3.294 \quad \int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx$$

Optimal result	1438
Rubi [A] (verified)	1438
Mathematica [A] (verified)	1439
Maple [F]	1439
Fricas [F]	1440
Sympy [F(-1)]	1440
Maxima [F]	1440
Giac [F]	1440
Mupad [F(-1)]	1441

Optimal result

Integrand size = 23, antiderivative size = 78

$$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx = \frac{(a \cos(e+fx))^{1+m} \sqrt{b \csc(e+fx)} \operatorname{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e+fx)\right) \sqrt[4]{\sin^2(e+fx)}}{ab^3 f(1+m)}$$

[Out] -(a*cos(f*x+e))^(1+m)*hypergeom([-3/4, 1/2+1/2*m], [3/2+1/2*m], cos(f*x+e)^2) *(sin(f*x+e)^2)^(1/4)*(b*csc(f*x+e))^(1/2)/a/b^3/f/(1+m)

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2656}

$$\int \frac{(a \cos(e+fx))^m}{(b \csc(e+fx))^{5/2}} dx = \frac{(a \cos(e+fx))^{m+1} \operatorname{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{m+1}{2}, \frac{m+3}{2}, \cos^2(e+fx)\right)}{abf(m+1) \sin^2(e+fx)^{3/4} (b \csc(e+fx))^{3/2}}$$

[In] Int[(a*cos[e + f*x])^m/(b*Csc[e + f*x])^(5/2),x]

[Out] -(((a*cos[e + f*x])^(1 + m)*Hypergeometric2F1[-3/4, (1 + m)/2, (3 + m)/2, C os[e + f*x]^2])/(a*b*f*(1 + m)*(b*Csc[e + f*x])^(3/2)*(Sin[e + f*x]^2)^(3/4)))

Rule 2656

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x])^(2*FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]
```

Rule 2666

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(1/b^2)*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int (a \cos(e + fx))^m (b \sin(e + fx))^{5/2} dx}{b^2 (b \csc(e + fx))^{3/2} (b \sin(e + fx))^{3/2}} \\ &= -\frac{(a \cos(e + fx))^{1+m} \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{abf(1+m)(b \csc(e + fx))^{3/2} \sin^2(e + fx)^{3/4}} \end{aligned}$$

Mathematica [A] (verified)

Time = 11.38 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.60

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \frac{2(a \cos(e + fx))^m (1 + 2 \cos(2(e + fx))) (-\cot^2(e + fx))^{\frac{1-m}{2}} \text{Hypergeometric2F1}\left(-\frac{3}{4}, \frac{1+m}{2}, \frac{3+m}{2}, \cos^2(e + fx)\right)}{b^2 f (5 + 2m) \sqrt{b \csc(e + fx)} (-4 + \dots)}$$

```
[In] Integrate[(a*Cos[e + f*x])^m/(b*Csc[e + f*x])^(5/2), x]
```

```
[Out] (2*(a*Cos[e + f*x])^m*(1 + 2*Cos[2*(e + f*x)])*(-Cot[e + f*x]^2)^((1 - m)/2)*Hypergeometric2F1[(-5 - 2*m)/4, (1 - m)/2, (-1 - 2*m)/4, Csc[e + f*x]^2]*Tan[e + f*x]/(b^2*f*(5 + 2*m)*Sqrt[b*Csc[e + f*x]]*(-4 + 3*Csc[e + f*x]^2))
```

Maple [F]

$$\int \frac{(\cos(fx + e)a)^m}{(b \csc(fx + e))^{\frac{5}{2}}} dx$$

```
[In] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(5/2), x)
```

```
[Out] int((cos(f*x+e)*a)^m/(b*csc(f*x+e))^(5/2), x)
```

Fricas [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{5/2}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(b*csc(f*x + e))*(a*cos(f*x + e))^m/(b^3*csc(f*x + e)^3), x)

Sympy [F(-1)]

Timed out.

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \text{Timed out}$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{5/2}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="maxima")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(5/2), x)

Giac [F]

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \int \frac{(a \cos(fx + e))^m}{(b \csc(fx + e))^{5/2}} dx$$

[In] integrate((a*cos(f*x+e))^m/(b*csc(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((a*cos(f*x + e))^m/(b*csc(f*x + e))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(a \cos(e + fx))^m}{(b \csc(e + fx))^{5/2}} dx = \int \frac{(a \cos(e + fx))^m}{\left(\frac{b}{\sin(e+fx)}\right)^{5/2}} dx$$

```
[In] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(5/2),x)
```

```
[Out] int((a*cos(e + f*x))^m/(b/sin(e + f*x))^(5/2), x)
```

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 1443

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
  fi;
else # result do not contain complex
  # this assumes optimal do not as well. No check is needed here.
  if debug then
    print("result do not contain complex, this assumes optimal do not as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A"," ";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
  fi;
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C",cat("Result contains higher order function than in optimal. Order ",
                convert(ExpnType_result,string)," vs. order ",
                convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

```



```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```



```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```