

Computer Algebra Independent Integration Tests

Summer 2023 edition

4-Trig-functions/4.2-Cosine/87-4.2.13-d+e-x-^m-cos-a+b-x+c-x²-
ⁿ

Nasser M. Abbasi

September 5, 2023 Compiled on September 5, 2023 at 4:43pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	35
4	Appendix	209

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [34]. This is test number [87].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (34)	0.00 (0)
Mathematica	100.00 (34)	0.00 (0)
Fricas	100.00 (34)	0.00 (0)
Maple	94.12 (32)	5.88 (2)
Giac	94.12 (32)	5.88 (2)
Maxima	94.12 (32)	5.88 (2)
Sympy	52.94 (18)	47.06 (16)
Mupad	44.12 (15)	55.88 (19)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	70.588	0.000	0.000	29.412
Mathematica	70.588	0.000	0.000	29.412
Fricas	70.588	0.000	0.000	29.412
Maple	64.706	0.000	0.000	35.294
Sympy	17.647	5.882	0.000	76.471
Giac	0.000	0.000	64.706	35.294
Mupad	0.000	14.706	0.000	85.294
Maxima	0.000	0.000	64.706	35.294

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	2	100.00	0.00	0.00
Giac	2	100.00	0.00	0.00
Maxima	2	100.00	0.00	0.00
Sympy	16	100.00	0.00	0.00
Mupad	19	0.00	100.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.10
Fricas	0.28
Giac	0.41
Maxima	0.56
Maple	0.84
Mathematica	1.89
Sympy	2.79
Mupad	10.14

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	32.87	1.01	19.00	1.01
Sympy	46.06	1.18	18.00	0.91
Mathematica	83.97	0.95	86.50	0.96
Fricas	89.09	0.95	94.50	0.99
Maple	95.19	0.90	68.00	0.85
Rubi	101.38	1.00	98.50	1.00
Giac	112.19	1.15	98.50	1.11
Maxima	509.28	3.35	125.00	2.46

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

[{4, 9, 14, 15, 19, 23, 27, 28, 31, 34}](#)

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi [{}](#)

Mathematica [{}](#)

Maple [{}](#)

Maxima [{}](#)

Fricas [{}](#)

Sympy [{}](#)

Giac [{}](#)

Mupad [{}](#)

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi [{}](#)

Mathematica [{}](#)

Maple [{}](#)

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```

x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)

```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```

try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1

```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```

integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)

```

Which gives $\sin(x)^{2/2}$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	33

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	23
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }
B grade { }
C grade { }
F normal fail { }
F(-1) timeout fail { }
F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }
B grade { }
C grade { }
F normal fail { }
F(-1) timeout fail { }
F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

B grade { }

C grade { }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 5, 6, 7, 8, 10, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { }

B grade { }

C grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { }

B grade { }

C grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 20, 21, 22, 24, 25, 26, 29, 30, 32, 33 }

F normal fail { 5, 10 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 3,8,11,12,13 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1,2,5,6,7,10,16,17,18,20,21,22,24,25,26,29,30,32,33 }

F(-2) exception fail { }

Sympy

A grade { 3,8,13,18,22,26 }

B grade { 12,25 }

C grade { }

F normal fail { 1,2,5,6,7,10,11,16,17,20,21,24,29,30,32,33 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	249	249	160	204	1570	173	0	227	0
N.S.	1	1.00	0.64	0.82	6.31	0.69	0.00	0.91	0.00
time (sec)	N/A	0.345	0.471	1.853	1.070	0.320	0.000	0.338	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	123	123	112	99	580	119	0	183	0
N.S.	1	1.00	0.91	0.80	4.72	0.97	0.00	1.49	0.00
time (sec)	N/A	0.074	0.197	0.844	0.630	0.272	0.000	0.333	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	85	81	112	103	88	135	99
N.S.	1	1.00	0.87	0.83	1.14	1.05	0.90	1.38	1.01
time (sec)	N/A	0.031	0.115	0.357	0.239	0.302	0.218	0.333	0.080

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	15	15	17	15	17	17	14	17	17
N.S.	1	1.00	1.13	1.00	1.13	1.13	0.93	1.13	1.13
time (sec)	N/A	0.013	2.150	0.227	0.361	0.262	0.383	0.340	13.744

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	111	111	110	0	0	115	0	0	0
N.S.	1	1.00	0.99	0.00	0.00	1.04	0.00	0.00	0.00
time (sec)	N/A	0.144	1.719	0.000	0.000	0.287	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	251	251	164	224	1570	178	0	229	0
N.S.	1	1.00	0.65	0.89	6.25	0.71	0.00	0.91	0.00
time (sec)	N/A	0.326	0.458	1.755	1.104	0.286	0.000	0.325	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	124	124	116	110	580	125	0	185	0
N.S.	1	1.00	0.94	0.89	4.68	1.01	0.00	1.49	0.00
time (sec)	N/A	0.060	0.206	0.855	0.618	0.293	0.000	0.373	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	99	99	88	88	112	106	94	137	105
N.S.	1	1.00	0.89	0.89	1.13	1.07	0.95	1.38	1.06
time (sec)	N/A	0.032	0.118	0.332	0.246	0.275	0.226	0.363	0.054

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	20	20	14	18	18
N.S.	1	1.00	1.12	1.00	1.25	1.25	0.88	1.12	1.12
time (sec)	N/A	0.017	2.046	0.251	0.366	0.266	0.398	0.338	14.376

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	111	111	114	0	0	123	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	1.11	0.00	0.00	0.00
time (sec)	N/A	0.156	2.419	0.000	0.000	0.274	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	67	59	160	59	0	75	64
N.S.	1	1.00	0.82	0.72	1.95	0.72	0.00	0.91	0.78
time (sec)	N/A	0.055	0.095	1.862	0.440	0.260	0.000	0.433	0.116

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	B	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	39	30	126	32	155	65	32
N.S.	1	1.00	0.95	0.73	3.07	0.78	3.78	1.59	0.78
time (sec)	N/A	0.022	0.033	1.584	0.401	0.266	0.661	0.473	14.528

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	24	24	24	20	70	22	29	39	19
N.S.	1	1.00	1.00	0.83	2.92	0.92	1.21	1.62	0.79
time (sec)	N/A	0.007	0.019	0.662	0.320	0.274	0.200	0.398	0.043

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	12	13	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	0.92	1.00	1.00
time (sec)	N/A	0.010	9.941	0.183	0.349	0.268	0.354	0.406	13.770

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	13	13	15	11	13	13	14	13	13
N.S.	1	1.00	1.15	0.85	1.00	1.00	1.08	1.00	1.00
time (sec)	N/A	0.043	10.619	0.208	0.360	0.281	0.326	0.407	14.017

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	248	170	191	1617	178	0	214	0
N.S.	1	1.00	0.69	0.77	6.52	0.72	0.00	0.86	0.00
time (sec)	N/A	0.333	0.454	0.684	1.061	0.345	0.000	0.462	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	116	95	610	124	0	172	0
N.S.	1	1.00	0.92	0.75	4.84	0.98	0.00	1.37	0.00
time (sec)	N/A	0.075	0.173	0.722	0.635	0.290	0.000	0.438	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	97	72	124	93	83	122	0
N.S.	1	1.00	0.97	0.72	1.24	0.93	0.83	1.22	0.00
time (sec)	N/A	0.060	0.070	0.590	0.325	0.270	0.466	0.444	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	17	17	19	17	28	19	15	19	19
N.S.	1	1.00	1.12	1.00	1.65	1.12	0.88	1.12	1.12
time (sec)	N/A	0.034	3.959	0.428	0.399	0.270	0.798	0.485	13.750

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	248	248	175	199	1617	187	0	216	0
N.S.	1	1.00	0.71	0.80	6.52	0.75	0.00	0.87	0.00
time (sec)	N/A	0.258	0.399	0.663	1.055	0.269	0.000	0.418	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	122	99	611	133	0	174	0
N.S.	1	1.00	0.97	0.79	4.85	1.06	0.00	1.38	0.00
time (sec)	N/A	0.081	0.173	0.666	0.699	0.287	0.000	0.431	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	100	76	124	96	88	124	0
N.S.	1	1.00	1.00	0.76	1.24	0.96	0.88	1.24	0.00
time (sec)	N/A	0.057	0.072	0.682	0.330	0.285	0.452	0.438	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	28	22	15	20	20
N.S.	1	1.00	1.11	1.00	1.56	1.22	0.83	1.11	1.11
time (sec)	N/A	0.032	4.132	0.393	0.382	0.270	0.838	0.500	13.403

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	77	64	171	57	0	64	0
N.S.	1	1.00	0.91	0.75	2.01	0.67	0.00	0.75	0.00
time (sec)	N/A	0.070	0.104	1.428	0.430	0.275	0.000	0.503	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	B	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	46	46	41	35	136	37	121	54	0
N.S.	1	1.00	0.89	0.76	2.96	0.80	2.63	1.17	0.00
time (sec)	N/A	0.035	0.043	1.313	0.382	0.273	0.919	0.460	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	A	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	27	27	26	20	34	19	22	26	0
N.S.	1	1.00	0.96	0.74	1.26	0.70	0.81	0.96	0.00
time (sec)	N/A	0.017	0.011	1.264	0.304	0.266	0.374	0.593	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	24	15	14	15	15
N.S.	1	1.00	1.13	0.87	1.60	1.00	0.93	1.00	1.00
time (sec)	N/A	0.029	7.145	0.363	0.358	0.265	0.662	0.533	13.385

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	15	15	17	13	26	15	15	15	15
N.S.	1	1.00	1.13	0.87	1.73	1.00	1.00	1.00	1.00
time (sec)	N/A	0.058	7.053	0.401	0.339	0.273	0.594	0.405	13.308

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	285	285	187	396	2271	230	0	289	0
N.S.	1	1.00	0.66	1.39	7.97	0.81	0.00	1.01	0.00
time (sec)	N/A	0.310	0.788	2.376	1.336	0.269	0.000	0.347	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	140	140	129	182	695	136	0	199	0
N.S.	1	1.00	0.92	1.30	4.96	0.97	0.00	1.42	0.00
time (sec)	N/A	0.055	0.440	0.752	0.593	0.287	0.000	0.333	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	19	19	21	19	21	21	17	21	21
N.S.	1	1.00	1.11	1.00	1.11	1.11	0.89	1.11	1.11
time (sec)	N/A	0.014	3.237	0.429	0.385	0.269	0.345	0.351	13.456

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	291	291	215	378	2360	257	0	292	0
N.S.	1	1.00	0.74	1.30	8.11	0.88	0.00	1.00	0.00
time (sec)	N/A	0.370	0.699	1.391	1.382	0.303	0.000	0.371	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	A	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	150	139	170	737	149	0	195	0
N.S.	1	1.00	0.93	1.13	4.91	0.99	0.00	1.30	0.00
time (sec)	N/A	0.112	0.326	0.776	0.649	0.280	0.000	0.353	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	21	21	23	21	120	23	19	23	23
N.S.	1	1.00	1.10	1.00	5.71	1.10	0.90	1.10	1.10
time (sec)	N/A	0.045	4.466	0.544	0.382	0.279	41.990	0.396	14.124

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [24] had the largest ratio of [.46670000000000004]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	8	6	1.00	15	0.400
2	A	4	4	1.00	13	0.308
3	A	3	3	1.00	11	0.273
4	N/A	0	0	1.00	15	0.000
5	A	5	4	1.00	32	0.125
6	A	8	6	1.00	16	0.375
7	A	4	4	1.00	14	0.286
8	A	3	3	1.00	12	0.250
9	N/A	0	0	1.00	16	0.000
10	A	5	4	1.00	34	0.118
11	A	6	6	1.00	13	0.462
12	A	3	3	1.00	11	0.273
13	A	2	2	1.00	9	0.222
14	N/A	0	0	1.00	13	0.000
15	N/A	0	0	1.00	13	0.000
16	A	10	7	1.00	17	0.412
17	A	6	5	1.00	15	0.333
18	A	5	4	1.00	13	0.308
19	N/A	0	0	1.00	17	0.000
20	A	10	7	1.00	18	0.389
21	A	6	5	1.00	16	0.312
22	A	5	4	1.00	14	0.286
23	N/A	0	0	1.00	18	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
24	A	8	7	1.00	15	0.467
25	A	5	4	1.00	13	0.308
26	A	4	3	1.00	11	0.273
27	N/A	0	0	1.00	15	0.000
28	N/A	0	0	1.00	15	0.000
29	A	8	6	1.00	19	0.316
30	A	4	4	1.00	17	0.235
31	N/A	0	0	1.00	19	0.000
32	A	10	7	1.00	21	0.333
33	A	6	5	1.00	19	0.263
34	N/A	0	0	1.00	21	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^2 \cos(a + bx + cx^2) dx$	37
3.2	$\int x \cos(a + bx + cx^2) dx$	44
3.3	$\int \cos(a + bx + cx^2) dx$	50
3.4	$\int \frac{\cos(a+bx+cx^2)}{x} dx$	55
3.5	$\int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx$	58
3.6	$\int x^2 \cos(a + bx - cx^2) dx$	63
3.7	$\int x \cos(a + bx - cx^2) dx$	70
3.8	$\int \cos(a + bx - cx^2) dx$	76
3.9	$\int \frac{\cos(a+bx-cx^2)}{x} dx$	81
3.10	$\int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx$	84
3.11	$\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx$	89
3.12	$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx$	94
3.13	$\int \cos\left(\frac{1}{4} + x + x^2\right) dx$	99
3.14	$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$	103
3.15	$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$	106
3.16	$\int x^2 \cos^2(a + bx + cx^2) dx$	110
3.17	$\int x \cos^2(a + bx + cx^2) dx$	118
3.18	$\int \cos^2(a + bx + cx^2) dx$	124
3.19	$\int \frac{\cos^2(a+bx+cx^2)}{x} dx$	129
3.20	$\int x^2 \cos^2(a + bx - cx^2) dx$	132
3.21	$\int x \cos^2(a + bx - cx^2) dx$	140
3.22	$\int \cos^2(a + bx - cx^2) dx$	146
3.23	$\int \frac{\cos^2(a+bx-cx^2)}{x} dx$	151
3.24	$\int x^2 \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	154
3.25	$\int x \cos^2\left(\frac{1}{4} + x + x^2\right) dx$	159

3.26	$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$	164
3.27	$\int \frac{\cos^2(\frac{1}{4}+x+x^2)}{x} dx$	168
3.28	$\int \frac{\cos^2(\frac{1}{4}+x+x^2)}{x^2} dx$	171
3.29	$\int (d+ex)^2 \cos(a+bx+cx^2) dx$	175
3.30	$\int (d+ex) \cos(a+bx+cx^2) dx$	183
3.31	$\int \frac{\cos(a+bx+cx^2)}{d+ex} dx$	189
3.32	$\int (d+ex)^2 \cos^2(a+bx+cx^2) dx$	192
3.33	$\int (d+ex) \cos^2(a+bx+cx^2) dx$	200
3.34	$\int \frac{\cos^2(a+bx+cx^2)}{d+ex} dx$	206

3.1 $\int x^2 \cos(a + bx + cx^2) dx$

Optimal result	37
Rubi [A] (verified)	38
Mathematica [A] (verified)	40
Maple [A] (verified)	40
Fricas [A] (verification not implemented)	41
Sympy [F]	41
Maxima [C] (verification not implemented)	41
Giac [C] (verification not implemented)	43
Mupad [F(-1)]	43

Optimal result

Integrand size = 15, antiderivative size = 249

$$\begin{aligned} \int x^2 \cos(a + bx + cx^2) dx = & \frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & - \frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c} \end{aligned}$$

```
[Out] -1/4*b*sin(c*x^2+b*x+a)/c^2+1/2*x*sin(c*x^2+b*x+a)/c+1/8*b^2*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(5/2)-1/4*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(3/2)-1/4*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(3/2)-1/8*b^2*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, number of rules / integrand size = 0.400, Rules used = {3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int x^2 \cos(a + bx + cx^2) dx = & -\frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & + \frac{\sqrt{\frac{\pi}{2}} b^2 \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} b^2 \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c} \end{aligned}$$

[In] `Int[x^2*Cos[a + b*x + c*x^2], x]`

[Out] `(b^2*.Sqrt[Pi/2]*Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/(4*c^(5/2)) - (Sqrt[Pi/2]*Cos[a - b^2/(4*c)]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])/(2*c^(3/2)) - (Sqrt[Pi/2]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a - b^2/(4*c)]/(2*c^(3/2)) - (b^2*Sqrt[Pi/2]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])])*Sin[a - b^2/(4*c)]/(4*c^(5/2)) - (b*Sin[a + b*x + c*x^2])/(4*c^2) + (x*Sin[a + b*x + c*x^2])/(2*c)`

Rule 3432

```
Int[ Sin[(d_.)*(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ Cos[(d_.)*(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3528

```
Int[ Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[ Cos[(b^2 - 4*a*c)/(4*c)], Int[ Sin[(b + 2*c*x)^2/(4*c)], x], x] - Dist[ Sin[(b^2 - 4*a*c)/(4*c)], Int[ Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3529

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_))^(m_), x_Sym-
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[
(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - D-
ist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x])
/; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x \sin(a + bx + cx^2)}{2c} - \frac{\int \sin(a + bx + cx^2) dx}{2c} - \frac{b \int x \cos(a + bx + cx^2) dx}{2c} \\
&= -\frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c} + \frac{b^2 \int \cos(a + bx + cx^2) dx}{4c^2} \\
&\quad - \frac{\cos\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} - \frac{\sin\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\
&= -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\
&\quad - \frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c} \\
&\quad + \frac{\left(b^2 \cos\left(a - \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx}{4c^2} - \frac{\left(b^2 \sin\left(a - \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{4c^2} \\
&= \frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} - \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\
&\quad - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\
&\quad - \frac{b \sin(a + bx + cx^2)}{4c^2} + \frac{x \sin(a + bx + cx^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.64

$$\int x^2 \cos(a + bx + cx^2) dx \\ = \frac{-\sqrt{2\pi} \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a - \frac{b^2}{4c}\right) + b^2 \sin\left(a - \frac{b^2}{4c}\right)\right) + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a - \frac{b^2}{4c}\right) - 2c \sin\left(a - \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

[In] `Integrate[x^2*Cos[a + b*x + c*x^2], x]`

[Out] $\left(-\frac{(-\operatorname{FresnelS}\left(\frac{(b+2c)x}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a - \frac{b^2}{4c}\right) + b^2 \sin\left(a - \frac{b^2}{4c}\right)\right) + \operatorname{FresnelC}\left(\frac{(b+2c)x}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a - \frac{b^2}{4c}\right) - 2c \sin\left(a - \frac{b^2}{4c}\right)\right))}{8c^{5/2}} \right)$

Maple [A] (verified)

Time = 1.85 (sec) , antiderivative size = 204, normalized size of antiderivative = 0.82

method	result
default	$\frac{x \sin(cx^2 + bx + a)}{2c} - \frac{b \left(\frac{\sin(cx^2 + bx + a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2 - ac}{4c}\right) C\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2 - ac}{4c}\right) S\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2 - ac}{4c}\right) C\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2 - ac}{4c}\right) S\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}}$
risch	$\frac{b^2 \sqrt{\pi} e^{-\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}} - \frac{i\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} - \frac{b^2 \sqrt{\pi} e^{\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} - \frac{i\sqrt{\pi} e^{\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2 \cos\left(\frac{b^2 - ac}{4c}\right) C\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi}x^2 \sin\left(\frac{b^2 - ac}{4c}\right) S\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\sqrt{2}\pi^{\frac{3}{2}} \left(\cos\left(\frac{4ac-b^2}{4c}\right) C\left(\frac{\sqrt{2}\sqrt{c}x + \frac{\sqrt{2}b}{2\sqrt{\pi}\sqrt{c}}}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac-b^2}{4c}\right) S\left(\frac{\sqrt{2}\sqrt{c}x + \frac{\sqrt{2}b}{2\sqrt{\pi}\sqrt{c}}}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}}$

[In] `int(x^2*cos(c*x^2+b*x+a), x, method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2}x \sin(cx^2 + bx + a)/c - \frac{1}{2}b/c * (\frac{1}{2} \sin(cx^2 + bx + a)/c - \frac{1}{4}b/c^{(3/2)} * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} * (\cos((1/4)b^2 - a*c)/c) * \operatorname{FresnelC}(2^{(1/2)}/\operatorname{Pi}^{(1/2)}/c^{(1/2)} * (c*x + 1/2*b)) + \sin((1/4)b^2 - a*c)/c * \operatorname{FresnelS}(2^{(1/2)}/\operatorname{Pi}^{(1/2)}/c^{(1/2)} * (c*x + 1/2*b))) - \frac{1}{4}c^{(3/2)} * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} * (\cos((1/4)b^2 - a*c)/c) * \operatorname{FresnelS}(2^{(1/2)}/\operatorname{Pi}^{(1/2)}/c^{(1/2)})$$

$/2)/c^{(1/2)*(c*x+1/2*b))}-\sin((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)*(c*x+1/2*b)})$

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.69

$$\int x^2 \cos(a + bx + cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \cos\left(-\frac{b^2 - 4ac}{4c}\right) - 2\pi c \sin\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} \text{C}\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2} \left(\pi b^2 \sin\left(-\frac{b^2 - 4ac}{4c}\right) + 2\pi c \cos\left(-\frac{b^2 - 4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}}}{8c^3}$$

```
[In] integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="fricas")
[Out] 1/8*(sqrt(2)*(pi*b^2*cos(-1/4*(b^2 - 4*a*c)/c) - 2*pi*c*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) - sqrt(2)*(pi*b^2*sin(-1/4*(b^2 - 4*a*c)/c) + 2*pi*c*cos(-1/4*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x + b)*sqrt(c/pi)/c) + 2*(2*c^2*x - b*c)*sin(c*x^2 + b*x + a))/c^3
```

Sympy [F]

$$\int x^2 \cos(a + bx + cx^2) dx = \int x^2 \cos(a + bx + cx^2) dx$$

```
[In] integrate(x**2*cos(c*x**2+b*x+a),x)
[Out] Integral(x**2*cos(a + b*x + c*x**2), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.07 (sec) , antiderivative size = 1570, normalized size of antiderivative = 6.31

$$\int x^2 \cos(a + bx + cx^2) dx = \text{Too large to display}$$

```
[In] integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="maxima")
[Out] 1/32*(8*(((-(I - 1)*sqrt(2)*sqrt(pi))*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1)) + 2*(2*c^2*x - b*c)*sin(c*x^2 + b*x + a))/c^3
```

$$\begin{aligned}
& 4*I*b*c*x + I*b^2)/c)) - 1)*b^2*c^3 + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(\\
& 4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I* \\
& c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/4*(b^2 - 4*a*c)/c) + ((-(I + 1) \\
&)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) \\
& + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
&)/c)) - 1))*b^2*c^3 + 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I* \\
& b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c* \\
& x + I*b^2)/c))*c^4)*sin(-1/4*(b^2 - 4*a*c)/c))*x^3 + 12*((-(I - 1)*sqrt(2) \\
&)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1) \\
&)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) \\
& - 1))*b^3*c^2 + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + \\
& I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
&)/c))*b*c^3)*cos(-1/4*(b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(\\
& 1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(\\
& pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 4* \\
& (-I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1) \\
&)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*s \\
& in(-1/4*(b^2 - 4*a*c)/c))*x^2 + 8*(b*c^2*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x \\
& + I*b^2)/c) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^ \\
& 2 - 4*a*c)/c) - b*c^2*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + e^(-1/ \\
& 4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/4*(b^2 - 4*a*c)/c))*((4*c^2* \\
& x^2 + 4*b*c*x + b^2)/c)^(3/2) + 6*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(\\
& ((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(er \\
& f(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 4*((I + 1)* \\
& sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(\\
& 2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*cos(-1/4* \\
& (b^2 - 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + \\
& 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(- \\
& 4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 4*(-(I - 1)*sqrt(2)*gamma(3 \\
& /2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, - \\
& 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*sin(-1/4*(b^2 - 4*a*c)/c) \\
&))*x - (((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + \\
& I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4* \\
& I*b*c*x + I*b^2)/c)) - 1))*b^5 - 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2* \\
& x^2 + 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + \\
& 4*I*b*c*x + I*b^2)/c))*b^3*c)*cos(-1/4*(b^2 - 4*a*c)/c) - (((I + 1)*sqrt(\\
& 2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I \\
& - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) \\
& - 1))*b^5 - 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 + 4*I*b*c*x + \\
& I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
&)/c))*b^3*c)*sin(-1/4*(b^2 - 4*a*c)/c))/(c^4*((4*c^2*x^2 + 4*b*c*x + b^2)/c) \\
&)^(3/2))
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.34 (sec) , antiderivative size = 227, normalized size of antiderivative = 0.91

$$\int x^2 \cos(a + bx + cx^2) dx =$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2+2ic)\operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4ic}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(-2ix-\frac{ib}{c}\right) + 2ib\right)e^{(icx^2+ibx+ia)}$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2-2ic)\operatorname{erf}\left(\frac{1}{4}i\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4ic}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(2ix+\frac{ib}{c}\right) - 2ib\right)e^{(-icx^2-ibx-ia)}$$

$$16c^2$$

[In] `integrate(x^2*cos(c*x^2+b*x+a),x, algorithm="giac")`

[Out] $-1/16*(-I*\sqrt{2}*\sqrt{\pi}*(b^2 + 2*I*c)*\operatorname{erf}(-1/4*I*\sqrt{2}*(2*x + b/c)*(I*c/abs(c) + 1)*\sqrt{abs(c)})*e^{(-1/4*(I*b^2 - 4*I*a*c)/c)}/((I*c/abs(c) + 1)*\sqrt{abs(c)}) - 2*(c*(-2*I*x - I*b/c) + 2*I*b)*e^{(I*c*x^2 + I*b*x + I*a)}/c^2 - 1/16*(I*\sqrt{2}*\sqrt{\pi}*(b^2 - 2*I*c)*\operatorname{erf}(1/4*I*\sqrt{2}*(2*x + b/c)*(-I*c/abs(c) + 1)*\sqrt{abs(c)})*e^{(-1/4*(-I*b^2 + 4*I*a*c)/c)}/((-I*c/abs(c) + 1)*\sqrt{abs(c)}) - 2*(c*(2*I*x + I*b/c) - 2*I*b)*e^{(-I*c*x^2 - I*b*x - I*a)}/c^2$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + bx + cx^2) dx = \int x^2 \cos(cx^2 + bx + a) dx$$

[In] `int(x^2*cos(a + b*x + c*x^2),x)`

[Out] `int(x^2*cos(a + b*x + c*x^2), x)`

3.2 $\int x \cos(a + bx + cx^2) dx$

Optimal result	44
Rubi [A] (verified)	44
Mathematica [A] (verified)	46
Maple [A] (verified)	46
Fricas [A] (verification not implemented)	47
Sympy [F]	47
Maxima [C] (verification not implemented)	47
Giac [C] (verification not implemented)	48
Mupad [F(-1)]	49

Optimal result

Integrand size = 13, antiderivative size = 123

$$\begin{aligned} \int x \cos(a + bx + cx^2) dx = & -\frac{b\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & + \frac{b\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} + \frac{\sin(a + bx + cx^2)}{2c} \end{aligned}$$

[Out] $1/2*\sin(c*x^2+b*x+a)/c-1/4*b*\cos(a-1/4*b^2/c)*\text{FresnelC}(1/2*(2*c*x+b)/c^{(1/2)}*2^{(1/2)}/\text{Pi}^{(1/2)})*2^{(1/2)}*\text{Pi}^{(1/2)}/c^{(3/2)}+1/4*b*\text{FresnelS}(1/2*(2*c*x+b)/c^{(1/2)}*2^{(1/2)}/\text{Pi}^{(1/2)})*\sin(a-1/4*b^2/c)*2^{(1/2)}*\text{Pi}^{(1/2)}/c^{(3/2)}$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3543, 3529, 3433, 3432}

$$\begin{aligned} \int x \cos(a + bx + cx^2) dx = & -\frac{\sqrt{\frac{\pi}{2}} b \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & + \frac{\sqrt{\frac{\pi}{2}} b \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} + \frac{\sin(a + bx + cx^2)}{2c} \end{aligned}$$

[In] $\text{Int}[x*\cos[a + b*x + c*x^2], x]$

[Out] $-1/2*(b*\text{Sqrt}[\text{Pi}/2]*\cos[a - b^2/(4*c)]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])/c^{(3/2)} + (b*\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*\text{sin}[a - b^2/(4*c)])/(2*c^{(3/2)}) + \text{Sin}[a + b*x + c*x^2]/(2*c)$

Rule 3432

```
Int[ $\sin[(d_.) * ((e_.) + (f_.) * (x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ ) * FresnelS[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot e + f \cdot x$ ], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\cos[(d_.) * ((e_.) + (f_.) * (x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ ) * FresnelC[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot e + f \cdot x$ ], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[ $\cos[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2]$ , x_Symbol] :> Dist[ $\cos[(b^2 - 4a*c)/(4*c)]$ , Int[ $\cos[(b + 2c*x)^2/(4*c)]$ , x], x] + Dist[ $\sin[(b^2 - 4a*c)/(4*c)]$ , Int[ $\sin[(b + 2c*x)^2/(4*c)]$ , x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4a*c, 0]
```

Rule 3543

```
Int[ $\cos[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2] * ((d_.) + (e_.) * (x_))$ , x_Symbol] :> Simp[e * (Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[ $\cos[a + b*x + c*x^2]$ , x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\sin(a + bx + cx^2)}{2c} - \frac{b \int \cos(a + bx + cx^2) dx}{2c} \\
&= \frac{\sin(a + bx + cx^2)}{2c} - \frac{\left(b \cos\left(a - \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\
&\quad + \frac{\left(b \sin\left(a - \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\
&= -\frac{b \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\
&\quad + \frac{b \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} + \frac{\sin(a + bx + cx^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.91

$$\int x \cos(a + bx + cx^2) dx \\ = \frac{-b\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) + 2\sqrt{c} \sin(a + x(b + cx))}{4c^{3/2}}$$

[In] `Integrate[x*Cos[a + b*x + c*x^2], x]`

[Out] $\left(-\frac{(b \sqrt{2\pi}) \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{3/2}} + \frac{b \sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{3/2}} + \frac{2\sqrt{c} \sin(a + x(b + cx))}{4c^{3/2}} \right)$

Maple [A] (verified)

Time = 0.84 (sec) , antiderivative size = 99, normalized size of antiderivative = 0.80

method	result
default	$\frac{\sin(cx^2 + bx + a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{\frac{b^2}{4} - ac}{c}\right) \text{C}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{\frac{b^2}{4} - ac}{c}\right) \text{S}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}}$
risch	$-\frac{b\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{4c}} \text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} + \frac{b\sqrt{\pi} e^{\frac{i(4ac-b^2)}{4c}} \text{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} + \frac{\sin(cx^2 + bx + a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi} x \cos\left(\frac{\frac{b^2}{4} - ac}{c}\right) \text{C}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi} x \sin\left(\frac{\frac{b^2}{4} - ac}{c}\right) \text{S}\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right)}{2\sqrt{c}} - \frac{\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{\frac{b^2}{4} - ac}{c}\right) \sqrt{2}\sqrt{\pi} \text{C}\left(\frac{\sqrt{2}\sqrt{c}x + \frac{b}{2}}{\sqrt{\pi}}\right) + \sin\left(\frac{\frac{b^2}{4} - ac}{c}\right) \sqrt{2}\sqrt{\pi} \text{S}\left(\frac{\sqrt{2}\sqrt{c}x + \frac{b}{2}}{\sqrt{\pi}}\right) \right)}{2\sqrt{c}}$

[In] `int(x*cos(c*x^2+b*x+a), x, method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2} \sin(cx^2 + bx + a) / c - \frac{1}{4} b / c^{(3/2)} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4)b^2 - a*c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2*b)) + \sin((1/4)b^2 - a*c) / c * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2*b))$$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.97

$$\int x \cos(a + bx + cx^2) dx = -\frac{\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right) - 2c \sin(cx^2 + b^2/4c)}{4c^2}$$

[In] `integrate(x*cos(c*x^2+b*x+a),x, algorithm="fricas")`

[Out]
$$-1/4 * (\sqrt{2} * \pi * b * \sqrt{c/\pi}) * \cos(-1/4 * (b^2 - 4*a*c)/c) * \text{fresnel_cos}(1/2 * \sqrt{2} * (2*c*x + b) * \sqrt{c/\pi}/c) - \sqrt{2} * \pi * b * \sqrt{c/\pi} * \text{fresnel_sin}(1/2 * \sqrt{2} * (2*c*x + b) * \sqrt{c/\pi}/c) * \sin(-1/4 * (b^2 - 4*a*c)/c) - 2*c * \sin(c*x^2 + b*x + a)/c^2$$

Sympy [F]

$$\int x \cos(a + bx + cx^2) dx = \int x \cos(a + bx + cx^2) dx$$

[In] `integrate(x*cos(c*x**2+b*x+a),x)`

[Out] `Integral(x*cos(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.63 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.72

$$\int x \cos(a + bx + cx^2) dx = \frac{\left((i-1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf}\left(\frac{1}{2} \sqrt{\frac{4 i c^2 x^2+4 i b c x+i b^2}{c}}\right)-1\right)-(i+1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf}\left(\frac{1}{2} \sqrt{\frac{-4 i c^2 x^2+4 i b c x+i b^2}{c}}\right)-1\right)\right) b^2}{b^2}$$

[In] `integrate(x*cos(c*x^2+b*x+a),x, algorithm="maxima")`

[Out]
$$1/16 * ((I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}\left(1/2 * \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}\right) - 1) - (I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}\left(1/2 * \sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}\right) - 1)) * b^2 * \cos(-1/4 * (b^2 - 4*a*c)/c) + ((I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}\left(1/2 * \sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}\right) - 1) - (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}\left(1/2 * \sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}\right) - 1)) * b^2 * \sin(-1/4 * (b^2 - 4*a*c)/c)$$

$$\begin{aligned}
& \text{qrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}((4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1) - (\text{I} - 1) * \\
& \text{sqrt}(2) * \text{sqrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}(-(4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1)) * \\
& \text{b}^2 * \sin(-1/4 * (\text{b}^2 - 4 * \text{a} * \text{c}) / \text{c}) - 2 * ((-(\text{I} - 1) * \text{sqrt}(2) * \text{sqrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}((4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1) + (\text{I} + 1) * \text{sqrt}(2) * \text{sqrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}(-(4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1)) * \text{b} * \text{c} * \cos(-1/4 * (\text{b}^2 - \\
& 4 * \text{a} * \text{c}) / \text{c}) + (-(\text{I} + 1) * \text{sqrt}(2) * \text{sqrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}((4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1) + (\text{I} - 1) * \text{sqrt}(2) * \text{sqrt}(\text{pi}) * (\text{erf}(1/2 * \text{sqrt}(-(4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})) - 1)) * \text{b} * \text{c} * \sin(-1/4 * (\text{b}^2 - 4 * \text{a} * \text{c}) / \text{c})) * \text{x} - 4 * (\text{c} * (\text{I} * \text{e}^{(1/4 * (4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})} - \text{I} * \text{e}^{(-1/4 * (4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})}) * \cos(-1/4 * (\text{b}^2 - 4 * \text{a} * \text{c}) / \text{c}) - \text{c} * (\text{e}^{(1/4 * (4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})} + \text{e}^{(-1/4 * (4 * \text{I} * \text{c}^2 * \text{x}^2 + 4 * \text{I} * \text{b} * \text{c} * \text{x} + \text{I} * \text{b}^2) / \text{c})}) * \sin(-1/4 * (\text{b}^2 - 4 * \text{a} * \text{c}) / \text{c})) * \text{sqrt}((4 * \text{c}^2 * \text{x}^2 + 4 * \text{b} * \text{c} * \text{x} + \text{b}^2) / \text{c})) / (\text{c}^2 * \text{sqrt}((4 * \text{c}^2 * \text{x}^2 + 4 * \text{b} * \text{c} * \text{x} + \text{b}^2) / \text{c}))
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.49

$$\begin{aligned}
& \int x \cos(a + bx + cx^2) dx \\
& = -\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{4} i \sqrt{2} \left(2x + \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{i b^2 - 4 i a c}{4 c}\right)}}{\left(\frac{i c}{|c|} + 1\right) \sqrt{|c|}} + 2 i e^{(i c x^2 + i b x + i a)} \\
& - \frac{8 c}{-\frac{i \sqrt{2} \sqrt{\pi} b \operatorname{erf}\left(\frac{1}{4} i \sqrt{2} \left(2x + \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}\right) e^{\left(-\frac{-i b^2 + 4 i a c}{4 c}\right)}}{\left(-\frac{i c}{|c|} + 1\right) \sqrt{|c|}} - 2 i e^{(-i c x^2 - i b x - i a)}}
\end{aligned}$$

```

[In] integrate(x*cos(c*x^2+b*x+a),x, algorithm="giac")
[Out] -1/8*(I*sqrt(2)*sqrt(pi)*b*erf(-1/4*I*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*
sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c)))
+ 2*I*e^(I*c*x^2 + I*b*x + I*a))/c - 1/8*(-I*sqrt(2)*sqrt(pi)*b*erf(1/4*I*s
qrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*
c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*e^(-I*c*x^2 - I*b*x - I*a))/c

```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + bx + cx^2) dx = \int x \cos(cx^2 + bx + a) dx$$

[In] `int(x*cos(a + b*x + c*x^2),x)`

[Out] `int(x*cos(a + b*x + c*x^2), x)`

3.3 $\int \cos(a + bx + cx^2) dx$

Optimal result	50
Rubi [A] (verified)	50
Mathematica [A] (verified)	51
Maple [A] (verified)	52
Fricas [A] (verification not implemented)	52
Sympy [A] (verification not implemented)	52
Maxima [C] (verification not implemented)	53
Giac [C] (verification not implemented)	53
Mupad [B] (verification not implemented)	54

Optimal result

Integrand size = 11, antiderivative size = 98

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{\sqrt{c}}$$

[Out] $\frac{1}{2} \cos(a - \frac{b^2}{4c}) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3529, 3433, 3432}

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}$$

[In] $\text{Int}[\cos[a + b*x + c*x^2], x]$

[Out] $\frac{(\sqrt{\frac{\pi}{2}} \cos(a - \frac{b^2}{4c}) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right))}{\sqrt{c}} - \frac{(\sqrt{\frac{\pi}{2}} \sin(a - \frac{b^2}{4c}) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right))}{\sqrt{c}}$

Rule 3432

```
Int[ $\sin[(d_.) \cdot ((e_.) + (f_.) \cdot (x_))^2]$ , x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\cos[(d_.) \cdot ((e_.) + (f_.) \cdot (x_))^2]$ , x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[ $\cos[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2]$ , x_Symbol] :> Dist[ $\cos[(b^2 - 4*a*c)/(4*c)]$ , Int[ $\cos[(b + 2*c*x)^2/(4*c)]$ , x], x] + Dist[ $\sin[(b^2 - 4*a*c)/(4*c)]$ , Int[ $\sin[(b + 2*c*x)^2/(4*c)]$ , x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \cos\left(a - \frac{b^2}{4c}\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) dx - \sin\left(a - \frac{b^2}{4c}\right) \int \sin\left(\frac{(b + 2cx)^2}{4c}\right) dx \\ &= \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{\sqrt{c}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 85, normalized size of antiderivative = 0.87

$$\begin{aligned} &\int \cos(a + bx + cx^2) dx \\ &= \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \right)}{\sqrt{c}} \end{aligned}$$

[In] `Integrate[Cos[a + b*x + c*x^2], x]`

[Out] `(Sqrt[Pi/2]*(Cos[a - b^2/(4*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] - FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a - b^2/(4*c)]))/Sqrt[c]`

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \left(\cos\left(\frac{b^2}{4} - \frac{ac}{c}\right) C\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi} \sqrt{c}}\right) + \sin\left(\frac{b^2}{4} - \frac{ac}{c}\right) S\left(\frac{\sqrt{2}(cx + \frac{b}{2})}{\sqrt{\pi} \sqrt{c}}\right) \right)}{2\sqrt{c}}$	81
risch	$\frac{\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac-b^2)}{4c}} \operatorname{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}}$	99

[In] `int(cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2} \cdot 2^{(1/2)} \cdot \pi^{(1/2)} \cdot c^{(1/2)} \cdot (\cos((1/4 \cdot b^2 - a \cdot c)/c) \cdot \operatorname{FresnelC}(2^{(1/2)}/\pi^{(1/2)})/c^{(1/2)} \cdot (c \cdot x + 1/2 \cdot b)) + \sin((1/4 \cdot b^2 - a \cdot c)/c) \cdot \operatorname{FresnelS}(2^{(1/2)}/\pi^{(1/2)})/c^{(1/2)} \cdot (c \cdot x + 1/2 \cdot b))$$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.05

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\pi \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2-4ac}{4c}\right)}{2c}$$

[In] `integrate(cos(c*x^2+b*x+a),x, algorithm="fricas")`

[Out]
$$\frac{1}{2} \cdot (\sqrt{2} \cdot \pi \cdot \sqrt{c/\pi}) \cdot \cos(-1/4 \cdot (b^2 - 4 \cdot a \cdot c)/c) \cdot \operatorname{fresnel_cos}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x + b) \cdot \sqrt{c/\pi}/c) - \sqrt{2} \cdot \pi \cdot \sqrt{c/\pi} \cdot \operatorname{fresnel_sin}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x + b) \cdot \sqrt{c/\pi}/c) \cdot \sin(-1/4 \cdot (b^2 - 4 \cdot a \cdot c)/c))/c$$

Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.90

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(-\sin\left(a - \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(a - \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b+2cx)}{2\sqrt{\pi}\sqrt{c}}\right) \right) \sqrt{\frac{1}{c}}}{2}$$

[In] `integrate(cos(c*x**2+b*x+a),x)`

```
[Out] sqrt(2)*sqrt(pi)*(-sin(a - b**2/(4*c))*fresnels(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))) + cos(a - b**2/(4*c))*fresnelc(sqrt(2)*(b + 2*c*x)/(2*sqrt(pi)*sqrt(c))))*sqrt(1/c)/2
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.24 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.14

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left((i-1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) + \left((i+1) \cos\left(-\frac{b^2-4ac}{4c}\right) + (i-1) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{-2icx+ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

```
[In] integrate(cos(c*x^2+b*x+a),x, algorithm="maxima")
```

```
[Out] -1/8*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + ((I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))/sqrt(c)
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.38

$$\int \cos(a + bx + cx^2) dx = \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2-4icac}{4c}\right)}}{4\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{1}{4}i\sqrt{2}(2x + \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2+4icac}{4c}\right)}}{4\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}}$$

```
[In] integrate(cos(c*x^2+b*x+a),x, algorithm="giac")
```

```
[Out] 1/4*I*sqrt(2)*sqrt(pi)*erf(-1/4*I*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 1/4*I*sqrt(2)*sqrt(pi)*erf(1/4*I*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c)))
```

Mupad [B] (verification not implemented)

Time = 0.08 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.01

$$\int \cos(a + bx + cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} \cos\left(\frac{4ac - b^2}{4c}\right) C\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2} - \frac{\sqrt{2} \sqrt{\pi} \sin\left(\frac{4ac - b^2}{4c}\right) S\left(\frac{\sqrt{2}\left(\frac{b}{2} + cx\right)\sqrt{\frac{1}{c}}}{\sqrt{\pi}}\right) \sqrt{\frac{1}{c}}}{2}$$

[In] int(cos(a + b*x + c*x^2),x)

[Out] $(2^{(1/2)}*\pi^{(1/2)}*\cos((4*a*c - b^2)/(4*c))*\text{fresnelc}((2^{(1/2)}*(b/2 + c*x)*(1/c)^{(1/2)})/\pi^{(1/2)})*(1/c)^{(1/2)})/2 - (2^{(1/2)}*\pi^{(1/2)}*\sin((4*a*c - b^2)/(4*c))*\text{fresnels}((2^{(1/2)}*(b/2 + c*x)*(1/c)^{(1/2)})/\pi^{(1/2)})*(1/c)^{(1/2)})/2$

3.4 $\int \frac{\cos(a+bx+cx^2)}{x} dx$

Optimal result	55
Rubi [N/A]	55
Mathematica [N/A]	56
Maple [N/A] (verified)	56
Fricas [N/A]	56
Sympy [N/A]	56
Maxima [N/A]	57
Giac [N/A]	57
Mupad [N/A]	57

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \text{Int}\left(\frac{\cos(a + bx + cx^2)}{x}, x\right)$$

[Out] Unintegrable($\cos(c*x^2+b*x+a)/x$, x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec), antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

[In] Int[$\cos[a + b*x + c*x^2]/x$, x]

[Out] Defer[Int][$\cos[a + b*x + c*x^2]/x$, x]

Rubi steps

$$\text{integral} = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 2.15 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

[In] `Integrate[Cos[a + b*x + c*x^2]/x, x]`

[Out] `Integrate[Cos[a + b*x + c*x^2]/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.23 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)}{x} dx$$

[In] `int(cos(c*x^2+b*x+a)/x,x)`

[Out] `int(cos(c*x^2+b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 + b*x + a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(a + bx + cx^2)}{x} dx$$

[In] `integrate(cos(c*x**2+b*x+a)/x,x)`

[Out] `Integral(cos(a + b*x + c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="maxima")`

[Out] `integrate(cos(c*x^2 + b*x + a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(cos(c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 13.74 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)}{x} dx$$

[In] `int(cos(a + b*x + c*x^2)/x,x)`

[Out] `int(cos(a + b*x + c*x^2)/x, x)`

3.5 $\int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx$

Optimal result	58
Rubi [A] (verified)	58
Mathematica [A] (verified)	60
Maple [F]	60
Fricas [A] (verification not implemented)	60
Sympy [F]	61
Maxima [F]	61
Giac [F]	61
Mupad [F(-1)]	62

Optimal result

Integrand size = 32, antiderivative size = 111

$$\begin{aligned} & \int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx \\ &= -\frac{\cos(a+bx+cx^2)}{x} - \sqrt{c}\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \end{aligned}$$

[Out] $-\cos(c*x^2+b*x+a)/x - \cos(a-1/4*b^2/c)*\text{FresnelS}(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*c^(1/2)*2^(1/2)*Pi^(1/2) - \text{FresnelC}(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*\sin(a-1/4*b^2/c)*c^(1/2)*2^(1/2)*Pi^(1/2)$

Rubi [A] (verified)

Time = 0.14 (sec), antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3547, 3528, 3432, 3433}

$$\begin{aligned} & \int \left(\frac{\cos(a+bx+cx^2)}{x^2} + \frac{b \sin(a+bx+cx^2)}{x} \right) dx \\ &= -\sqrt{2\pi}\sqrt{c} \sin\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{2\pi}\sqrt{c} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \frac{\cos(a+bx+cx^2)}{x} \end{aligned}$$

[In] $\text{Int}[\text{Cos}[a + b*x + c*x^2]/x^2 + (b*\text{Sin}[a + b*x + c*x^2])/x, x]$

[Out] $-(\cos[a + bx + cx^2]/x) - \sqrt{c} \operatorname{Sqrt}[2\pi] \cos[a - b^2/(4c)] \operatorname{FresnelS}[(b + 2cx)/(\operatorname{Sqrt}[c] \operatorname{Sqrt}[2\pi])] - \sqrt{c} \operatorname{Sqrt}[2\pi] \operatorname{FresnelC}[(b + 2cx)/(\operatorname{Sqrt}[c] \operatorname{Sqrt}[2\pi])] \sin[a - b^2/(4c)]$

Rule 3432

$\operatorname{Int}[\sin[(d_.) \cdot (e_.) + (f_.) \cdot (x_.)^2], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2]/(f \cdot \operatorname{Rt}[d, 2])) \operatorname{FresnelS}[\operatorname{Sqrt}[2/\pi] \cdot \operatorname{Rt}[d, 2] \cdot (e + f \cdot x)], x] /; \operatorname{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\operatorname{Int}[\cos[(d_.) \cdot (e_.) + (f_.) \cdot (x_.)^2], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Sqrt}[\pi/2]/(f \cdot \operatorname{Rt}[d, 2])) \operatorname{FresnelC}[\operatorname{Sqrt}[2/\pi] \cdot \operatorname{Rt}[d, 2] \cdot (e + f \cdot x)], x] /; \operatorname{FreeQ}[\{d, e, f\}, x]$

Rule 3528

$\operatorname{Int}[\sin[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2], x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[\cos[(b^2 - 4a \cdot c)/(4c)], \operatorname{Int}[\sin[(b + 2cx)^2/(4c)], x], x] - \operatorname{Dist}[\sin[(b^2 - 4a \cdot c)/(4c)], \operatorname{Int}[\cos[(b + 2cx)^2/(4c)], x], x] /; \operatorname{FreeQ}[\{a, b, c\}, x] \& \operatorname{NeQ}[b^2 - 4a \cdot c, 0]$

Rule 3547

$\operatorname{Int}[\cos[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2] \cdot ((d_.) + (e_.) \cdot (x_.)^{(m)}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(d + e \cdot x)^{(m+1)} \cdot (\cos[a + bx + cx^2]/(e \cdot (m+1))), x] + (\operatorname{Dist}[(b \cdot e - 2 \cdot c \cdot d)/(e^{2(m+1)}), \operatorname{Int}[(d + e \cdot x)^{(m+1)} \cdot \sin[a + bx + cx^2], x], x] + \operatorname{Dist}[2 \cdot (c/(e^{2(m+1)})), \operatorname{Int}[(d + e \cdot x)^{(m+2)} \cdot \sin[a + bx + cx^2], x], x]) /; \operatorname{FreeQ}[\{a, b, c, d, e\}, x] \& \operatorname{NeQ}[b \cdot e - 2 \cdot c \cdot d, 0] \& \operatorname{LtQ}[m, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{\sin(a + bx + cx^2)}{x} dx + \int \frac{\cos(a + bx + cx^2)}{x^2} dx \\ &= -\frac{\cos(a + bx + cx^2)}{x} - (2c) \int \sin(a + bx + cx^2) dx \\ &= -\frac{\cos(a + bx + cx^2)}{x} - \left(2c \cos\left(a - \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b + 2cx)^2}{4c}\right) dx \\ &\quad - \left(2c \sin\left(a - \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b + 2cx)^2}{4c}\right) dx \\ &= -\frac{\cos(a + bx + cx^2)}{x} - \sqrt{c} \sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b + 2cx}{\sqrt{c} \sqrt{2\pi}}\right) \\ &\quad - \sqrt{c} \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{b + 2cx}{\sqrt{c} \sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 1.72 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.99

$$\int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx =$$

$$-\frac{\cos(a + x(b + cx)) + \sqrt{c}\sqrt{2\pi}x \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{c}\sqrt{2\pi}x \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{x}$$

[In] Integrate[Cos[a + b*x + c*x^2]/x^2 + (b*Sin[a + b*x + c*x^2])/x, x]

[Out] $-\left(\left(\cos\left(a+x(b+c x)\right)+\sqrt{c} \sqrt{2 \pi } x \cos \left(a-\frac{b^2}{4 c}\right)\right) \text{FresnelS}\left(\frac{(b+2 c x) (\sqrt{c} \sqrt{2 \pi })}{\sqrt{c} \sqrt{2 \pi }}\right)+\sqrt{c} \sqrt{2 \pi } x \text{FresnelC}\left(\frac{(b+2 c x) (\sqrt{c} \sqrt{2 \pi })}{\sqrt{c} \sqrt{2 \pi }}\right) \sin \left(a-\frac{b^2}{4 c}\right)\right) / x$

Maple [F]

$$\int \left(\frac{\cos(cx^2 + bx + a)}{x^2} + \frac{b \sin(cx^2 + bx + a)}{x} \right) dx$$

[In] int(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x, x)

[Out] int(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x, x)

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.04

$$\int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx =$$

$$-\frac{\sqrt{2} \pi x \sqrt{\frac{c}{\pi}} \cos \left(-\frac{b^2-4 a c}{4 c}\right) S\left(\frac{\sqrt{2} (2 c x+b) \sqrt{\frac{c}{\pi }}}{2 c}\right)+\sqrt{2} \pi x \sqrt{\frac{c}{\pi }} C\left(\frac{\sqrt{2} (2 c x+b) \sqrt{\frac{c}{\pi }}}{2 c}\right) \sin \left(-\frac{b^2-4 a c}{4 c}\right)+\cos \left(c x^2+b x+a\right)}{x}$$

[In] integrate(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x, x, algorithm="fricas")

[Out] $-\left(\sqrt{2} \pi x \sqrt{\frac{c}{\pi }} \cos \left(-\frac{b^2-4 a c}{4 c}\right) S\left(\frac{\sqrt{2} (2 c x+b) \sqrt{\frac{c}{\pi }}}{2 c}\right)+\sqrt{2} \pi x \sqrt{\frac{c}{\pi }} C\left(\frac{\sqrt{2} (2 c x+b) \sqrt{\frac{c}{\pi }}}{2 c}\right) \sin \left(-\frac{b^2-4 a c}{4 c}\right)+\cos \left(c x^2+b x+a\right)\right) / x$

Sympy [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{bx \sin(a + bx + cx^2) + \cos(a + bx + cx^2)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(c*x**2+b*x+a)/x**2+b*sin(c*x**2+b*x+a)/x,x)`

[Out] `Integral((b*x*sin(a + b*x + c*x**2) + cos(a + b*x + c*x**2))/x**2, x)`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(cx^2 + bx + a)}{x} + \frac{\cos(cx^2 + bx + a)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x,x, algorithm="maxima")`

[Out] `integrate(b*sin(c*x^2 + b*x + a)/x + cos(c*x^2 + b*x + a)/x^2, x)`

Giac [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(cx^2 + bx + a)}{x} + \frac{\cos(cx^2 + bx + a)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(c*x^2+b*x+a)/x^2+b*sin(c*x^2+b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(b*sin(c*x^2 + b*x + a)/x + cos(c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(\frac{\cos(a + bx + cx^2)}{x^2} + \frac{b \sin(a + bx + cx^2)}{x} \right) dx \\ &= \int \frac{\cos(cx^2 + bx + a)}{x^2} + \frac{b \sin(cx^2 + bx + a)}{x} dx \end{aligned}$$

[In] `int(cos(a + b*x + c*x^2)/x^2 + (b*sin(a + b*x + c*x^2))/x, x)`

[Out] `int(cos(a + b*x + c*x^2)/x^2 + (b*sin(a + b*x + c*x^2))/x, x)`

3.6 $\int x^2 \cos(a + bx - cx^2) dx$

Optimal result	63
Rubi [A] (verified)	64
Mathematica [A] (verified)	66
Maple [A] (verified)	66
Fricas [A] (verification not implemented)	67
Sympy [F]	67
Maxima [C] (verification not implemented)	67
Giac [C] (verification not implemented)	69
Mupad [F(-1)]	69

Optimal result

Integrand size = 16, antiderivative size = 251

$$\begin{aligned} \int x^2 \cos(a + bx - cx^2) dx = & -\frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & - \frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c} \end{aligned}$$

```
[Out] -1/4*b*sin(-c*x^2+b*x+a)/c^2-1/2*x*sin(-c*x^2+b*x+a)/c-1/8*b^2*cos(a+1/4*b^2/c)*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(5/2)+1/4*cos(a+1/4*b^2/c)*FresnelS(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(3/2)-1/4*FresnelC(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(3/2)-1/8*b^2*FresnelS(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a+1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.375, Rules used = {3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int x^2 \cos(a + bx - cx^2) dx = & -\frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} b^2 \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} b^2 \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c} \end{aligned}$$

[In] $\text{Int}[x^2 \cos[a + bx - cx^2], x]$

[Out]
$$\begin{aligned} & -1/4*(b^2*\text{Sqrt}[\text{Pi}/2]*\cos[a + b^2/(4*c)]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}]])]/c^{(5/2)} + (\text{Sqrt}[\text{Pi}/2]*\cos[a + b^2/(4*c)]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqr} t[c]*\text{Sqrt}[2*\text{Pi}])])/({2*c}^{(3/2)}) - (\text{Sqrt}[\text{Pi}/2]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*\text{Sin}[a + b^2/(4*c)])/({2*c}^{(3/2)}) - (b^2*\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*\text{Sin}[a + b^2/(4*c)])/(4*c^{(5/2)}) - (b*\text{Sin}[a + b*x - c*x^2])/(4*c^2) - (x*\text{Sin}[a + b*x - c*x^2])/(2*c) \end{aligned}$$

Rule 3432

$\text{Int}[\text{Sin}[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\text{Cos}[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\text{Sqrt}[2/\text{Pi}]*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3528

$\text{Int}[\text{Sin}[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_{\text{Symbol}}] \Rightarrow \text{Dist}[\text{Cos}[(b^2 - 4*a*c)/(4*c)], \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Dist}[\text{Sin}[(b^2 - 4*a*c)/(4*c)], \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 3529

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_))^(m_), x_Sym-
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[
(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - D-
ist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x])
/; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x \sin(a + bx - cx^2)}{2c} + \frac{\int \sin(a + bx - cx^2) dx}{2c} + \frac{b \int x \cos(a + bx - cx^2) dx}{2c} \\
&= -\frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c} + \frac{b^2 \int \cos(a + bx - cx^2) dx}{4c^2} \\
&\quad - \frac{\cos\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} + \frac{\sin\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} \\
&= \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} \\
&\quad - \frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c} \\
&\quad + \frac{\left(b^2 \cos\left(a + \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx}{4c^2} + \frac{\left(b^2 \sin\left(a + \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx}{4c^2} \\
&= -\frac{b^2 \sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} + \frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\
&\quad - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} - \frac{b^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{4c^{5/2}} \\
&\quad - \frac{b \sin(a + bx - cx^2)}{4c^2} - \frac{x \sin(a + bx - cx^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.65

$$\int x^2 \cos(a + bx - cx^2) dx \\ = \frac{-\sqrt{2\pi} \operatorname{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a + \frac{b^2}{4c}\right) - b^2 \sin\left(a + \frac{b^2}{4c}\right)\right) + \sqrt{2\pi} \operatorname{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a + \frac{b^2}{4c}\right) + 2c \sin\left(a + \frac{b^2}{4c}\right)\right)}{8c^{5/2}}$$

[In] `Integrate[x^2*Cos[a + b*x - c*x^2], x]`

[Out] $\left(-\frac{\operatorname{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2c \cos\left(a + \frac{b^2}{4c}\right) - b^2 \sin\left(a + \frac{b^2}{4c}\right)\right) + \operatorname{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(b^2 \cos\left(a + \frac{b^2}{4c}\right) + 2c \sin\left(a + \frac{b^2}{4c}\right)\right)}{8c^{5/2}} \right)$

Maple [A] (verified)

Time = 1.76 (sec) , antiderivative size = 224, normalized size of antiderivative = 0.89

method	result
default	$-\frac{x \sin(-cx^2+bx+a)}{2c} + \frac{b \left(-\frac{\sin(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{4c}\right) C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) - \sin\left(\frac{b^2+ac}{4c}\right) S\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) \right)}{4c\sqrt{-c}} \right)}{2c}$
risch	$\frac{b^2\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\operatorname{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{16c^2\sqrt{-ic}} + \frac{i\sqrt{\pi}e^{-\frac{i(4ac+b^2)}{4c}}\operatorname{erf}\left(\sqrt{-ic}x+\frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} - \frac{b^2\sqrt{\pi}e^{\frac{i(4ac+b^2)}{4c}}\operatorname{erf}\left(-\sqrt{ic}x+\frac{ib}{2\sqrt{ic}}\right)}{16c^2\sqrt{ic}} + \frac{i\sqrt{2}\pi^{\frac{3}{2}}}{\sqrt{2}} \left(\cos\left(\frac{4ac+b^2}{4c}\right) C\left(-\frac{\sqrt{2}cx}{\sqrt{\pi}\sqrt{-c}}+\frac{b}{2}\right) - \sin\left(\frac{4ac+b^2}{4c}\right) S\left(-\frac{\sqrt{2}cx}{\sqrt{\pi}\sqrt{-c}}+\frac{b}{2}\right) \right)$
parts	$\frac{\sqrt{2}\sqrt{\pi}x^2 \cos\left(\frac{b^2+ac}{4c}\right) C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}x^2 \sin\left(\frac{b^2+ac}{4c}\right) S\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} +$

[In] `int(x^2*cos(-c*x^2+b*x+a), x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -\frac{1}{2}x \sin(-cx^2+b*x+a)/c + \frac{1}{2}b/c * (-\frac{1}{2} \sin(-cx^2+b*x+a)/c + \frac{1}{4}b/c * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} / (-c)^{(1/2)} * (\cos((1/4b^2+a*c)/c) * \operatorname{FresnelC}(2^{(1/2)} / \operatorname{Pi}^{(1/2)} / (-c)^{(1/2)} * (-c*x+1/2*b)) - \sin((1/4b^2+a*c)/c) * \operatorname{FresnelS}(2^{(1/2)} / \operatorname{Pi}^{(1/2)} / (-c)^{(1/2)} * (-c*x+1/2*b))) + \frac{1}{4}c * 2^{(1/2)} * \operatorname{Pi}^{(1/2)} / (-c)^{(1/2)} * (\cos((1/4b^2+a*c)/c) * \operatorname{FresnelC}(2^{(1/2)} / \operatorname{Pi}^{(1/2)} / (-c)^{(1/2)} * (-c*x+1/2*b))) \end{aligned}$$

$$\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b))+\sin((1/4*b^2+a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/(-c)^{(1/2)}*(-c*x+1/2*b)))$$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.71

$$\int x^2 \cos(a + bx - cx^2) dx = \frac{\sqrt{2} \left(\pi b^2 \cos\left(\frac{b^2+4ac}{4c}\right) + 2\pi c \sin\left(\frac{b^2+4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2} \left(\pi b^2 \sin\left(\frac{b^2+4ac}{4c}\right) - 2\pi c \cos\left(\frac{b^2+4ac}{4c}\right) \right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{8c^3}$$

```
[In] integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="fricas")
[Out] 1/8*(sqrt(2)*(pi*b^2*cos(1/4*(b^2 + 4*a*c)/c) + 2*pi*c*sin(1/4*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_cos(1/2*sqrt(2)*(2*c*x - b)*sqrt(c/pi)/c) + sqrt(2)*(pi*b^2*sin(1/4*(b^2 + 4*a*c)/c) - 2*pi*c*cos(1/4*(b^2 + 4*a*c)/c))*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x - b)*sqrt(c/pi)/c) + 2*(2*c^2*x + b*c)*sin(c*x^2 - b*x - a))/c^3
```

Sympy [F]

$$\int x^2 \cos(a + bx - cx^2) dx = \int x^2 \cos(a + bx - cx^2) dx$$

```
[In] integrate(x**2*cos(-c*x**2+b*x+a),x)
```

[Out] $\text{Integral}(x^{**2} \cos(a + b*x - c*x^{**2}), x)$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.10 (sec), antiderivative size = 1570, normalized size of antiderivative = 6.25

$$\int x^2 \cos(a + bx - cx^2) dx = \text{Too large to display}$$

```
[In] integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="maxima")
```

```
[Out] 1/32*(8*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))
```

$$\begin{aligned}
& 4*I*b*c*x + I*b^2)/c)) - 1)*b^2*c^3 + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(\\
& 4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I* \\
& c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*cos(1/4*(b^2 + 4*a*c)/c) + (((I + 1)* \\
& sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - \\
& (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1)) *b^2*c^3 + 4*((I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b* \\
& c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x \\
& + I*b^2)/c))*c^4)*sin(1/4*(b^2 + 4*a*c)/c))*x^3 + 12*((((I - 1)*sqrt(2)*sqrt(\\
& pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b \\
& ^3*c^2 + 4*(-(I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) *b*c^3)*cos(1/4*(b^2 + 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)*sin(1/4*(b^2 + 4*a*c)/c))*x^2 + 8*(b*c^2*(-I*e^(1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) *cos(1/4*(b^2 + 4*a*c)/c))*((4*c^2*x^2 - 4*b*c*x + b^2)/c)^3/2 + 6*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 4*((I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*cos(1/4*(b^2 + 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 4*((I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*sin(1/4*(b^2 + 4*a*c)/c))*x - (((I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^5 - 4*(-(I + 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^3*c)*cos(1/4*(b^2 + 4*a*c)/c) - (((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^5 - 4*(-(I - 1)*sqrt(2)*gamma(3/2, 1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^3*c)*sin(1/4*(b^2 + 4*a*c)/c))/(c^4*((4*c^2*x^2 - 4*b*c*x + b^2)/c)^3/2)
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 229, normalized size of antiderivative = 0.91

$$\int x^2 \cos(a + bx - cx^2) dx =$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2+2ic)\operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}\left(2x-\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2+4ic}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(-2ix+\frac{ib}{c}\right) - 2ib\right)e^{(icx^2-ibx-ia)}$$

$$-\frac{i\sqrt{2}\sqrt{\pi}(b^2-2ic)\operatorname{erf}\left(\frac{1}{4}i\sqrt{2}\left(2x-\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2-4ic}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2\left(c\left(2ix-\frac{ib}{c}\right) + 2ib\right)e^{(-icx^2+ibx+ia)}$$

$$16c^2$$

```
[In] integrate(x^2*cos(-c*x^2+b*x+a),x, algorithm="giac")
[Out] -1/16*(-I*sqrt(2)*sqrt(pi)*(b^2 + 2*I*c)*erf(-1/4*I*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 2*(c*(-2*I*x + I*b/c) - 2*I*b)*e^(I*c*x^2 - I*b*x - I*a))/c^2 - 1/16*(I*sqrt(2)*sqrt(pi)*(b^2 - 2*I*c)*erf(1/4*I*sqrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*(c*(2*I*x - I*b/c) + 2*I*b)*e^(-I*c*x^2 + I*b*x + I*a))/c^2
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos(a + bx - cx^2) dx = \int x^2 \cos(-cx^2 + bx + a) dx$$

```
[In] int(x^2*cos(a + b*x - c*x^2),x)
[Out] int(x^2*cos(a + b*x - c*x^2), x)
```

3.7 $\int x \cos(a + bx - cx^2) dx$

Optimal result	70
Rubi [A] (verified)	70
Mathematica [A] (verified)	72
Maple [A] (verified)	72
Fricas [A] (verification not implemented)	73
Sympy [F]	73
Maxima [C] (verification not implemented)	73
Giac [C] (verification not implemented)	74
Mupad [F(-1)]	75

Optimal result

Integrand size = 14, antiderivative size = 124

$$\begin{aligned} \int x \cos(a + bx - cx^2) dx = & -\frac{b\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{b\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} - \frac{\sin(a + bx - cx^2)}{2c} \end{aligned}$$

[Out] $-1/2*\sin(-c*x^2+b*x+a)/c-1/4*b*cos(a+1/4*b^2/c)*\text{FresnelC}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2))*2^(1/2)*\text{Pi}^(1/2)/c^(3/2)-1/4*b*\text{FresnelS}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2))*\sin(a+1/4*b^2/c)*2^(1/2)*\text{Pi}^(1/2)/c^(3/2)$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.286, Rules used = {3543, 3529, 3433, 3432}

$$\begin{aligned} \int x \cos(a + bx - cx^2) dx = & -\frac{\sqrt{\frac{\pi}{2}} b \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{\sqrt{\frac{\pi}{2}} b \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{\sin(a + bx - cx^2)}{2c} \end{aligned}$$

[In] $\text{Int}[x*\cos[a + b*x - c*x^2], x]$

[Out] $-1/2*(b*\text{Sqrt}[\text{Pi}/2]*\cos[a + b^2/(4*c)]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])/c^(3/2) - (b*\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])*s\text{in}[a + b^2/(4*c)]/(2*c^(3/2)) - \text{Sin}[a + b*x - c*x^2]/(2*c)$

Rule 3432

```
Int[ $\sin[(d_.) * ((e_.) + (f_.) * (x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ ) * FresnelS[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot e + f \cdot x$ ], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\cos[(d_.) * ((e_.) + (f_.) * (x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ ) * FresnelC[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot e + f \cdot x$ ], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[ $\cos[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2]$ , x_Symbol] :> Dist[ $\cos[(b^2 - 4a*c)/(4*c)]$ , Int[ $\cos[(b + 2*c*x)^2/(4*c)]$ , x], x] + Dist[ $\sin[(b^2 - 4a*c)/(4*c)]$ , Int[ $\sin[(b + 2*c*x)^2/(4*c)]$ , x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[ $\cos[(a_.) + (b_.) * (x_.) + (c_.) * (x_.)^2] * ((d_.) + (e_.) * (x_))$ , x_Symbol] :> Simp[e * (Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[ $\cos[a + b*x + c*x^2]$ , x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\sin(a + bx - cx^2)}{2c} + \frac{b \int \cos(a + bx - cx^2) dx}{2c} \\
&= -\frac{\sin(a + bx - cx^2)}{2c} + \frac{\left(b \cos\left(a + \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} \\
&\quad + \frac{\left(b \sin\left(a + \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b-2cx)^2}{4c}\right) dx}{2c} \\
&= -\frac{b \sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\
&\quad - \frac{b \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{2c^{3/2}} - \frac{\sin(a + bx - cx^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.21 (sec) , antiderivative size = 116, normalized size of antiderivative = 0.94

$$\int x \cos(a + bx - cx^2) dx \\ = \frac{b\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + b\sqrt{2\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) - 2\sqrt{c} \sin(a + x(b - cx))}{4c^{3/2}}$$

[In] `Integrate[x*Cos[a + b*x - c*x^2], x]`

[Out] $(b*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a + b^2/(4*c)]*\text{FresnelC}[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*\text{Pi}])] + b*\text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*\text{Pi}]])*\text{Sin}[a + b^2/(4*c)] - 2*\text{Sqrt}[c]*\text{Sin}[a + x*(b - c*x)])/(4*c^{(3/2)})$

Maple [A] (verified)

Time = 0.86 (sec) , antiderivative size = 110, normalized size of antiderivative = 0.89

method	result
default	$-\frac{\sin(-cx^2+bx+a)}{2c} + \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2+ac}{c}\right) C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) - \sin\left(\frac{b^2+ac}{c}\right) S\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right) \right)}{4c\sqrt{-c}}$
risch	$\frac{b\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{4c}} \operatorname{erf}\left(\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{8c\sqrt{-ic}} - \frac{b\sqrt{\pi} e^{\frac{i(4ac+b^2)}{4c}} \operatorname{erf}\left(-\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{8c\sqrt{ic}} - \frac{\sin(-cx^2+bx+a)}{2c}$
parts	$\frac{\sqrt{2}\sqrt{\pi}x \cos\left(\frac{b^2+ac}{c}\right) C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}x \sin\left(\frac{b^2+ac}{c}\right) S\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}} - \frac{\sqrt{2}\sqrt{\pi}x \cos\left(\frac{b^2+ac}{c}\right) \sqrt{2}\sqrt{\pi}\sqrt{-c} C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{-c}}\right)}{2\sqrt{-c}}$

[In] `int(x*cos(-c*x^2+b*x+a), x, method=_RETURNVERBOSE)`

[Out] $-1/2*\sin(-c*x^2+b*x+a)/c+1/4*b/c*2^(1/2)*\text{Pi}^(1/2)/(-c)^(1/2)*(\cos((1/4*b^2+a*c)/c)*\text{FresnelC}(2^(1/2)/\text{Pi}^(1/2)/(-c)^(1/2)*(-c*x+1/2*b))-\sin((1/4*b^2+a*c)/c)*\text{FresnelS}(2^(1/2)/\text{Pi}^(1/2)/(-c)^(1/2)*(-c*x+1/2*b)))$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.01

$$\int x \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi b \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) + 2c \sin(cx^2 - bx - a)}{4c^2}$$

[In] `integrate(x*cos(-c*x^2+b*x+a),x, algorithm="fricas")`

[Out] `1/4*(sqrt(2)*pi*b*sqrt(c/pi)*cos(1/4*(b^2 + 4*a*c)/c)*fresnel_cos(1/2*sqrt(2)*(2*c*x - b)*sqrt(c/pi)/c) + sqrt(2)*pi*b*sqrt(c/pi)*fresnel_sin(1/2*sqrt(2)*(2*c*x - b)*sqrt(c/pi)/c)*sin(1/4*(b^2 + 4*a*c)/c) + 2*c*sin(c*x^2 - bx - a))/c^2`

Sympy [F]

$$\int x \cos(a + bx - cx^2) dx = \int x \cos(a + bx - cx^2) dx$$

[In] `integrate(x*cos(-c*x**2+b*x+a),x)`

[Out] `Integral(x*cos(a + b*x - c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 580, normalized size of antiderivative = 4.68

$$\int x \cos(a + bx - cx^2) dx = -\frac{\left(-(i-1) \sqrt{2} \sqrt{\pi } \left(\text{erf}\left(\frac{1}{2} \sqrt{\frac{4 i c^2 x^2-4 i b c x+i b^2}{c}}\right)-1\right)+(i+1) \sqrt{2} \sqrt{\pi } \left(\text{erf}\left(\frac{1}{2} \sqrt{-\frac{4 i c^2 x^2-4 i b c x+i b^2}{c}}\right)-1\right)\right)}{c}$$

[In] `integrate(x*cos(-c*x^2+b*x+a),x, algorithm="maxima")`

[Out] `-1/16*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(1/4*(b^2 + 4*a*c)/c) + ((I + 1)*sqrt(2)*`

```

sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)
*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1)
)*b^2*sin(1/4*(b^2 + 4*a*c)/c) - 2*(-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqr
t((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(er
f(1/2*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(1/4*(b^2 +
4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 - 4*I*b*c*
x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 -
4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(1/4*(b^2 + 4*a*c)/c))*x - 4*(c*(-I*e
^(1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + I*e^(-1/4*(4*I*c^2*x^2 - 4*I*b
*c*x + I*b^2)/c))*cos(1/4*(b^2 + 4*a*c)/c) - c*(e^(1/4*(4*I*c^2*x^2 - 4*I*b
*c*x + I*b^2)/c) + e^(-1/4*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*sin(1/4*(b
^2 + 4*a*c)/c))*sqrt((4*c^2*x^2 - 4*b*c*x + b^2)/c)/(c^2*sqrt((4*c^2*x^2 -
4*b*c*x + b^2)/c))

```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.49

$$\begin{aligned}
& \int x \cos(a + bx - cx^2) dx \\
&= -\frac{\frac{i\sqrt{2}\sqrt{\pi}b \operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}\left(2x - \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2+4iac}{4c}\right)}}{\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} + 2ie^{(icx^2 - ibx - ia)}}{8c} \\
&\quad - \frac{\frac{i\sqrt{2}\sqrt{\pi}b \operatorname{erf}\left(\frac{1}{4}i\sqrt{2}\left(2x - \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2-4iac}{4c}\right)}}{\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - 2ie^{(-icx^2 + ibx + ia)}}{8c}
\end{aligned}$$

```

[In] integrate(x*cos(-c*x^2+b*x+a),x, algorithm="giac")
[Out] -1/8*(-I*sqrt(2)*sqrt(pi)*b*erf(-1/4*I*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)
*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c)))
+ 2*I*e^(I*c*x^2 - I*b*x - I*a))/c - 1/8*(I*sqrt(2)*sqrt(pi)*b*erf(1/4*I*s
qrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*e^(-I*c*x^2 + I*b*x + I*a))/c

```

Mupad [F(-1)]

Timed out.

$$\int x \cos(a + bx - cx^2) dx = \int x \cos(-cx^2 + bx + a) dx$$

[In] `int(x*cos(a + b*x - c*x^2),x)`

[Out] `int(x*cos(a + b*x - c*x^2), x)`

3.8 $\int \cos(a + bx - cx^2) dx$

Optimal result	76
Rubi [A] (verified)	76
Mathematica [A] (verified)	77
Maple [A] (verified)	78
Fricas [A] (verification not implemented)	78
Sympy [A] (verification not implemented)	78
Maxima [C] (verification not implemented)	79
Giac [C] (verification not implemented)	79
Mupad [B] (verification not implemented)	80

Optimal result

Integrand size = 12, antiderivative size = 99

$$\int \cos(a + bx - cx^2) dx = -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\ -\frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{\sqrt{c}}$$

[Out] $-1/2*\cos(a+1/4*b^2/c)*\text{FresnelC}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2))*2^(1/2)*\text{Pi}^(1/2)/c^(1/2)-1/2*\text{FresnelS}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2)) * \sin(a+1/4*b^2/c)*2^(1/2)*\text{Pi}^(1/2)/c^(1/2)$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3529, 3433, 3432}

$$\int \cos(a + bx - cx^2) dx = -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} \\ -\frac{\sqrt{\frac{\pi}{2}} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}}$$

[In] $\text{Int}[\cos[a + b*x - c*x^2], x]$

[Out] $-((\text{Sqrt}[\text{Pi}/2]*\cos[a + b^2/(4*c)]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])])/\text{Sqrt}[c]) - (\text{Sqrt}[\text{Pi}/2]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*)\sin[a + b^2/(4*c)])/\text{Sqrt}[c]$

Rule 3432

```
Int[ $\text{Sin}[(d_{\cdot})*((e_{\cdot}) + (f_{\cdot})*(x_{\cdot}))^2]$ , x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\text{Cos}[(d_{\cdot})*((e_{\cdot}) + (f_{\cdot})*(x_{\cdot}))^2]$ , x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[ $\text{Cos}[(a_{\cdot}) + (b_{\cdot})*(x_{\cdot}) + (c_{\cdot})*(x_{\cdot})^2]$ , x_Symbol] :> Dist[ $\text{Cos}[(b^2 - 4*a*c)/(4*c)]$ , Int[ $\text{Cos}[(b + 2*c*x)^2/(4*c)]$ , x], x] + Dist[ $\text{Sin}[(b^2 - 4*a*c)/(4*c)]$ , Int[ $\text{Sin}[(b + 2*c*x)^2/(4*c)]$ , x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \cos\left(a + \frac{b^2}{4c}\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) dx + \sin\left(a + \frac{b^2}{4c}\right) \int \sin\left(\frac{(b - 2cx)^2}{4c}\right) dx \\ &= -\frac{\sqrt{\frac{\pi}{2}} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{\sqrt{c}} - \frac{\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)}{\sqrt{c}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.12 (sec), antiderivative size = 88, normalized size of antiderivative = 0.89

$$\begin{aligned} &\int \cos(a + bx - cx^2) dx \\ &= \frac{\sqrt{\frac{\pi}{2}} \left(\cos\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \right)}{\sqrt{c}} \end{aligned}$$

[In] `Integrate[Cos[a + b*x - c*x^2], x]`

[Out] `(Sqrt[Pi/2]*(Cos[a + b^2/(4*c)]*FresnelC[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])] + FresnelS[(-b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*Sin[a + b^2/(4*c)]))/Sqrt[c]`

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.89

method	result	size
default	$\frac{\sqrt{2} \sqrt{\pi} \left(\cos\left(\frac{\frac{b^2}{4}+ac}{c}\right) C\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi} \sqrt{-c}}\right) - \sin\left(\frac{\frac{b^2}{4}+ac}{c}\right) S\left(\frac{\sqrt{2}(-cx+\frac{b}{2})}{\sqrt{\pi} \sqrt{-c}}\right) \right)}{2\sqrt{-c}}$	88
risch	$\frac{\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{4c}} \operatorname{erf}\left(\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)}{4\sqrt{-ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac+b^2)}{4c}} \operatorname{erf}\left(-\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)}{4\sqrt{ic}}$	95

[In] `int(cos(-c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2} \cdot 2^{(1/2)} \cdot \pi^{(1/2)} / (-c)^{(1/2)} \cdot (\cos((1/4 \cdot b^2 + a \cdot c) / c) * \text{FresnelC}(2^{(1/2)} / \pi^{(1/2)} / (-c)^{(1/2)} * (-c \cdot x + 1/2 \cdot b)) - \sin((1/4 \cdot b^2 + a \cdot c) / c) * \text{FresnelS}(2^{(1/2)} / \pi^{(1/2)} / (-c)^{(1/2)} * (-c \cdot x + 1/2 \cdot b)))$$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.07

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2}\pi\sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right)}{2c}$$

[In] `integrate(cos(-c*x^2+b*x+a),x, algorithm="fricas")`

[Out]
$$\frac{1}{2} \cdot (\sqrt{2} \cdot \pi \cdot \sqrt{c/\pi} \cdot \cos(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c) \cdot \text{fresnel_cos}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) + \sqrt{2} \cdot \pi \cdot \sqrt{c/\pi} \cdot \text{fresnel_sin}(1/2 \cdot \sqrt{2} \cdot (2 \cdot c \cdot x - b) \cdot \sqrt{c/\pi} / c) \cdot \sin(1/4 \cdot (b^2 + 4 \cdot a \cdot c) / c)) / c$$

Sympy [A] (verification not implemented)

Time = 0.23 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.95

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\sqrt{\pi}\sqrt{-\frac{1}{c}} \left(-\sin\left(a + \frac{b^2}{4c}\right) S\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(a + \frac{b^2}{4c}\right) C\left(\frac{\sqrt{2}(b-2cx)}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{2}$$

[In] `integrate(cos(-c*x**2+b*x+a),x)`

```
[Out] sqrt(2)*sqrt(pi)*sqrt(-1/c)*(-sin(a + b**2/(4*c))*fresnels(sqrt(2)*(b - 2*c*x)/(2*sqrt(pi)*sqrt(-c))) + cos(a + b**2/(4*c))*fresnelc(sqrt(2)*(b - 2*c*x)/(2*sqrt(pi)*sqrt(-c)))))/2
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.13

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2}\sqrt{\pi} \left(\left(-(i-1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i+1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx-ib}{2\sqrt{ic}}\right) + \left(-(i+1) \cos\left(\frac{b^2+4ac}{4c}\right) + (i-1) \sin\left(\frac{b^2+4ac}{4c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}}$$

```
[In] integrate(cos(-c*x^2+b*x+a),x, algorithm="maxima")
[Out] 1/8*sqrt(2)*sqrt(pi)*((-I - 1)*cos(1/4*(b^2 + 4*a*c)/c) + (I + 1)*sin(1/4*(b^2 + 4*a*c)/c))*erf(1/2*(2*I*c*x - I*b)/sqrt(I*c)) + (-I + 1)*cos(1/4*(b^2 + 4*a*c)/c) + (I - 1)*sin(1/4*(b^2 + 4*a*c)/c))*erf(1/2*(2*I*c*x - I*b)/sqrt(-I*c))/sqrt(c)
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.38

$$\int \cos(a + bx - cx^2) dx = \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}(2x - \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{ib^2+4ic}{4c}\right)}}{4\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - \frac{i\sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{1}{4}i\sqrt{2}(2x - \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right) e^{\left(-\frac{-ib^2-4ic}{4c}\right)}}{4\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}}$$

```
[In] integrate(cos(-c*x^2+b*x+a),x, algorithm="giac")
[Out] 1/4*I*sqrt(2)*sqrt(pi)*erf(-1/4*I*sqrt(2)*(2*x - b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 + 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 1/4*I*sqrt(2)*sqrt(pi)*erf(1/4*I*sqrt(2)*(2*x - b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 - 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c)))
```

Mupad [B] (verification not implemented)

Time = 0.05 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.06

$$\int \cos(a + bx - cx^2) dx = \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \cos\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2} - \frac{\sqrt{2} \sqrt{\pi} S\left(\frac{\sqrt{2}\left(\frac{b}{2}-cx\right)\sqrt{-\frac{1}{c}}}{\sqrt{\pi}}\right) \sin\left(\frac{b^2+4ac}{4c}\right) \sqrt{-\frac{1}{c}}}{2}$$

[In] int(cos(a + b*x - c*x^2),x)

[Out] $(2^{(1/2)}*\pi^{(1/2)}*\text{fresnelc}((2^{(1/2)}*(b/2 - c*x)*(-1/c)^{(1/2)})/\pi^{(1/2)})*\cos((4*a*c + b^2)/(4*c))*(-1/c)^{(1/2)})/2 - (2^{(1/2)}*\pi^{(1/2)}*\text{fresnels}((2^{(1/2)}*(b/2 - c*x)*(-1/c)^{(1/2)})/\pi^{(1/2)})*\sin((4*a*c + b^2)/(4*c))*(-1/c)^{(1/2)})/2$

3.9 $\int \frac{\cos(a+bx-cx^2)}{x} dx$

Optimal result	81
Rubi [N/A]	81
Mathematica [N/A]	82
Maple [N/A] (verified)	82
Fricas [N/A]	82
Sympy [N/A]	82
Maxima [N/A]	83
Giac [N/A]	83
Mupad [N/A]	83

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \text{Int}\left(\frac{\cos(a + bx - cx^2)}{x}, x\right)$$

[Out] Unintegrable($\cos(-c*x^2+b*x+a)/x$, x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

[In] Int[$\cos[a + b*x - c*x^2]/x$, x]

[Out] Defer[Int][$\cos[a + b*x - c*x^2]/x$, x]

Rubi steps

$$\text{integral} = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 2.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

[In] `Integrate[Cos[a + b*x - c*x^2]/x, x]`

[Out] `Integrate[Cos[a + b*x - c*x^2]/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.25 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

[In] `int(cos(-c*x^2+b*x+a)/x, x)`

[Out] `int(cos(-c*x^2+b*x+a)/x, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(-c*x^2+b*x+a)/x, x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 - b*x - a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(a + bx - cx^2)}{x} dx$$

[In] `integrate(cos(-c*x**2+b*x+a)/x, x)`

[Out] `Integral(cos(a + b*x - c*x**2)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.25

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(-c*x^2+b*x+a)/x,x, algorithm="maxima")`

[Out] `integrate(cos(c*x^2 - b*x - a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

[In] `integrate(cos(-c*x^2+b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(cos(-c*x^2 + b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 14.38 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\cos(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)}{x} dx$$

[In] `int(cos(a + b*x - c*x^2)/x,x)`

[Out] `int(cos(a + b*x - c*x^2)/x, x)`

3.10 $\int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx$

Optimal result	84
Rubi [A] (verified)	84
Mathematica [A] (verified)	86
Maple [F]	86
Fricas [A] (verification not implemented)	86
Sympy [F]	87
Maxima [F]	87
Giac [F]	87
Mupad [F(-1)]	88

Optimal result

Integrand size = 34, antiderivative size = 111

$$\begin{aligned} & \int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx \\ &= -\frac{\cos(a+bx-cx^2)}{x} + \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad - \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \end{aligned}$$

[Out] $-\cos(-c*x^2+b*x+a)/x+\cos(a+1/4*b^2/c)*\text{FresnelS}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2))*c^(1/2)*2^(1/2)*\text{Pi}^(1/2)-\text{FresnelC}(1/2*(-2*c*x+b)/c^(1/2)*2^(1/2)/\text{Pi}^(1/2))*\sin(a+1/4*b^2/c)*c^(1/2)*2^(1/2)*\text{Pi}^(1/2)$

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {3547, 3528, 3432, 3433}

$$\begin{aligned} & \int \left(\frac{\cos(a+bx-cx^2)}{x^2} + \frac{b \sin(a+bx-cx^2)}{x} \right) dx \\ &= -\sqrt{2\pi}\sqrt{c} \sin\left(a + \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &\quad + \sqrt{2\pi}\sqrt{c} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{2\pi}}\right) - \frac{\cos(a+bx-cx^2)}{x} \end{aligned}$$

[In] $\text{Int}[\text{Cos}[a + b*x - c*x^2]/x^2 + (b*\text{Sin}[a + b*x - c*x^2])/x, x]$

[Out] $-(\cos[a + bx - cx^2]/x) + \sqrt{c} \cdot \sqrt{2\pi} \cdot \cos[a + b^2/(4c)] \cdot \text{FresnelS}[(b - 2cx)/(\sqrt{c} \cdot \sqrt{2\pi})] - \sqrt{c} \cdot \sqrt{2\pi} \cdot \text{FresnelC}[(b - 2cx)/(\sqrt{c} \cdot \sqrt{2\pi})] \cdot \sin[a + b^2/(4c)]$

Rule 3432

$\text{Int}[\sin[(d_.) \cdot ((e_.) + (f_.) \cdot (x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f \cdot \text{Rt}[d, 2]) \cdot \text{FresnelS}[\sqrt{2/\pi} \cdot \text{Rt}[d, 2] \cdot (e + f \cdot x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\cos[(d_.) \cdot ((e_.) + (f_.) \cdot (x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f \cdot \text{Rt}[d, 2]) \cdot \text{FresnelC}[\sqrt{2/\pi} \cdot \text{Rt}[d, 2] \cdot (e + f \cdot x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3528

$\text{Int}[\sin[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2], x_{\text{Symbol}}] \Rightarrow \text{Dist}[\cos[(b^2 - 4a \cdot c)/(4c)], \text{Int}[\sin[(b + 2cx)^2/(4c)], x], x] - \text{Dist}[\sin[(b^2 - 4a \cdot c)/(4c)], \text{Int}[\cos[(b + 2cx)^2/(4c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \& \text{NeQ}[b^2 - 4a \cdot c, 0]$

Rule 3547

$\text{Int}[\cos[(a_.) + (b_.) \cdot (x_.) + (c_.) \cdot (x_.)^2] \cdot ((d_.) + (e_.) \cdot (x_.)^{(m)}), x_{\text{Symbol}}] \Rightarrow \text{Simp}[(d + e \cdot x)^{(m+1)} \cdot (\cos[a + bx + cx^2]/(e \cdot (m+1))), x] + (\text{Dist}[(b \cdot e - 2 \cdot c \cdot d)/(e^{2(m+1)}), \text{Int}[(d + e \cdot x)^{(m+1)} \cdot \sin[a + bx + cx^2], x], x] + \text{Dist}[2 \cdot (c/(e^{2(m+1)})), \text{Int}[(d + e \cdot x)^{(m+2)} \cdot \sin[a + bx + cx^2], x], x]) /; \text{FreeQ}[\{a, b, c, d, e\}, x] \& \text{NeQ}[b \cdot e - 2 \cdot c \cdot d, 0] \& \text{LtQ}[m, -1]$

Rubi steps

$$\begin{aligned} \text{integral} &= b \int \frac{\sin(a + bx - cx^2)}{x} dx + \int \frac{\cos(a + bx - cx^2)}{x^2} dx \\ &= -\frac{\cos(a + bx - cx^2)}{x} + (2c) \int \sin(a + bx - cx^2) dx \\ &= -\frac{\cos(a + bx - cx^2)}{x} - \left(2c \cos\left(a + \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b - 2cx)^2}{4c}\right) dx \\ &\quad + \left(2c \sin\left(a + \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b - 2cx)^2}{4c}\right) dx \\ &= -\frac{\cos(a + bx - cx^2)}{x} + \sqrt{c} \sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b - 2cx}{\sqrt{c} \sqrt{2\pi}}\right) \\ &\quad - \sqrt{c} \sqrt{2\pi} \text{FresnelC}\left(\frac{b - 2cx}{\sqrt{c} \sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 2.42 (sec) , antiderivative size = 114, normalized size of antiderivative = 1.03

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= -\frac{\cos(a + x(b - cx))}{x} - \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) \\ &+ \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right) \end{aligned}$$

[In] `Integrate[Cos[a + b*x - c*x^2]/x^2 + (b*Sin[a + b*x - c*x^2])/x, x]`

[Out] $-\frac{\cos(a + x(b - cx))}{x} - \sqrt{c}\sqrt{2\pi} \cos\left(a + \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) + \sqrt{c}\sqrt{2\pi} \text{FresnelC}\left(\frac{-b + 2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a + \frac{b^2}{4c}\right)$

Maple [F]

$$\int \left(\frac{\cos(-cx^2 + bx + a)}{x^2} + \frac{b \sin(-cx^2 + bx + a)}{x} \right) dx$$

[In] `int(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x, x)`

[Out] `int(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x, x)`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.11

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx = \\ & -\frac{\sqrt{2}\pi x \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) + \cos(cx^2 - bx - a)}{x} \end{aligned}$$

[In] `integrate(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x, x, algorithm="fricas")`

[Out] $-\frac{(\sqrt{2})\pi x \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{4c}\right) S\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi x \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx-b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(\frac{b^2+4ac}{4c}\right) + \cos(cx^2 - bx - a)}{x}$

Sympy [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{bx \sin(a + bx - cx^2) + \cos(a + bx - cx^2)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(-c*x**2+b*x+a)/x**2+b*sin(-c*x**2+b*x+a)/x,x)`

[Out] `Integral((b*x*sin(a + b*x - c*x**2) + cos(a + b*x - c*x**2))/x**2, x)`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(-cx^2 + bx + a)}{x} + \frac{\cos(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x,x, algorithm="maxima")`

[Out] `integrate(-b*sin(c*x^2 - b*x - a)/x + cos(c*x^2 - b*x - a)/x^2, x)`

Giac [F]

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{b \sin(-cx^2 + bx + a)}{x} + \frac{\cos(-cx^2 + bx + a)}{x^2} dx \end{aligned}$$

[In] `integrate(cos(-c*x^2+b*x+a)/x^2+b*sin(-c*x^2+b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(b*sin(-c*x^2 + b*x + a)/x + cos(-c*x^2 + b*x + a)/x^2, x)`

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \left(\frac{\cos(a + bx - cx^2)}{x^2} + \frac{b \sin(a + bx - cx^2)}{x} \right) dx \\ &= \int \frac{\cos(-cx^2 + bx + a)}{x^2} + \frac{b \sin(-cx^2 + bx + a)}{x} dx \end{aligned}$$

[In] `int(cos(a + b*x - c*x^2)/x^2 + (b*sin(a + b*x - c*x^2))/x, x)`

[Out] `int(cos(a + b*x - c*x^2)/x^2 + (b*sin(a + b*x - c*x^2))/x, x)`

3.11 $\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	89
Rubi [A] (verified)	89
Mathematica [A] (verified)	91
Maple [A] (verified)	91
Fricas [A] (verification not implemented)	92
Sympy [F]	92
Maxima [C] (verification not implemented)	92
Giac [C] (verification not implemented)	93
Mupad [B] (verification not implemented)	93

Optimal result

Integrand size = 13, antiderivative size = 82

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{2\pi}} \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{2\pi}} \right) \\ - \frac{1}{4} \sin \left(\frac{1}{4} + x + x^2 \right) + \frac{1}{2} x \sin \left(\frac{1}{4} + x + x^2 \right)$$

[Out] $-1/4*\sin(1/4+x+x^2)+1/2*x*\sin(1/4+x+x^2)+1/8*\operatorname{FresnelC}(1/2*(1+2*x)*2^{(1/2)}/\operatorname{Pi}^{(1/2)})*2^{(1/2)}*\operatorname{Pi}^{(1/2)}-1/4*\operatorname{FresnelS}(1/2*(1+2*x)*2^{(1/2)}/\operatorname{Pi}^{(1/2)})*2^{(1/2)}*\operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.462$, Rules used = {3545, 3543, 3527, 3433, 3526, 3432}

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{2\pi}} \right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{2\pi}} \right) \\ + \frac{1}{2} x \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{4} \sin \left(x^2 + x + \frac{1}{4} \right)$$

[In] $\operatorname{Int}[x^2 \cos[1/4 + x + x^2], x]$

[Out] $(\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[2*\operatorname{Pi}]])/4 - (\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelS}[(1 + 2*x)/\operatorname{Sqrt}[2*\operatorname{Pi}]])/2 - \operatorname{Sin}[1/4 + x + x^2]/4 + (x*\operatorname{Sin}[1/4 + x + x^2])/2$

Rule 3432

```
Int[ $\sin[(d_*)((e_*) + (f_*)(x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ )/(f*Rt[d, 2])*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\cos[(d_*)((e_*) + (f_*)(x_))^2]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ )/(f*Rt[d, 2])*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3526

```
Int[ $\sin[(a_*) + (b_*)(x_*) + (c_*)(x_*)^2]$ , x_Symbol] :> Int[ $\sin[(b + 2*c*x)^2/(4*c)]$ , x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3527

```
Int[ $\cos[(a_*) + (b_*)(x_*) + (c_*)(x_*)^2]$ , x_Symbol] :> Int[ $\cos[(b + 2*c*x)^2/(4*c)]$ , x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[ $\cos[(a_*) + (b_*)(x_*) + (c_*)(x_*)^2*((d_*) + (e_*)(x_*))$ , x_Symbol]
:> Simp[e*( $\sin[a + b*x + c*x^2]/(2*c)$ ), x] + Dist[(2*c*d - b*e)/(2*c), Int[ $\cos[a + b*x + c*x^2]$ , x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[ $\cos[(a_*) + (b_*)(x_*) + (c_*)(x_*)^2*((d_*) + (e_*)(x_*)^{(m)})$ , x_Symbol]
:> Simp[e*( $(d + e*x)^{(m-1)}*\sin[a + b*x + c*x^2]/(2*c)$ ), x] + (-Dist[(b*e - 2*c*d)/(2*c), Int[( $d + e*x)^{(m-1)}*\cos[a + b*x + c*x^2]$ , x], x] - Dist[e^2*((m-1)/(2*c)), Int[( $d + e*x)^{(m-2)}*\sin[a + b*x + c*x^2]$ , x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2}x\sin\left(\frac{1}{4} + x + x^2\right) - \frac{1}{2}\int x\cos\left(\frac{1}{4} + x + x^2\right) dx - \frac{1}{2}\int \sin\left(\frac{1}{4} + x + x^2\right) dx \\ &= -\frac{1}{4}\sin\left(\frac{1}{4} + x + x^2\right) + \frac{1}{2}x\sin\left(\frac{1}{4} + x + x^2\right) \\ &\quad + \frac{1}{4}\int \cos\left(\frac{1}{4} + x + x^2\right) dx - \frac{1}{2}\int \sin\left(\frac{1}{4}(1+2x)^2\right) dx \\ &= -\frac{1}{2}\sqrt{\frac{\pi}{2}}\text{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) - \frac{1}{4}\sin\left(\frac{1}{4} + x + x^2\right) \\ &\quad + \frac{1}{2}x\sin\left(\frac{1}{4} + x + x^2\right) + \frac{1}{4}\int \cos\left(\frac{1}{4}(1+2x)^2\right) dx \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{4} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) \\
&\quad - \frac{1}{4} \sin\left(\frac{1}{4} + x + x^2\right) + \frac{1}{2} x \sin\left(\frac{1}{4} + x + x^2\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec), antiderivative size = 67, normalized size of antiderivative = 0.82

$$\begin{aligned}
\int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx &= \frac{1}{8} \left(\sqrt{2\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) \right. \\
&\quad \left. - 2 \left(\sqrt{2\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) + (1-2x) \sin\left(\frac{1}{4} + x + x^2\right) \right) \right)
\end{aligned}$$

[In] `Integrate[x^2*Cos[1/4 + x + x^2], x]`

[Out] `(Sqrt[2*Pi]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]] - 2*(Sqrt[2*Pi]*FresnelS[(1 + 2*x)/Sqrt[2*Pi]] + (1 - 2*x)*Sin[1/4 + x + x^2]))/8`

Maple [A] (verified)

Time = 1.86 (sec), antiderivative size = 59, normalized size of antiderivative = 0.72

method	result
default	$\frac{x \sin\left(\frac{1}{4}+x+x^2\right)}{2}-\frac{\sin\left(\frac{1}{4}+x+x^2\right)}{4}+\frac{\sqrt{2} \sqrt{\pi } \operatorname{C}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi }}\right)}{8}-\frac{\sqrt{2} \sqrt{\pi } \operatorname{S}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi }}\right)}{4}$
risch	$-\frac{\sqrt{\pi } (-1)^{\frac{3}{4}} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{16}-\frac{(-1)^{\frac{1}{4}} \sqrt{\pi } \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8}+\frac{\sqrt{\pi } \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{16 \sqrt{-i}}+\frac{i \sqrt{\pi } \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{8 \sqrt{-i}}$
parts	$\frac{\sqrt{2} \sqrt{\pi } \operatorname{C}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi }}\right) x^2}{2}-\frac{\sqrt{2} \pi ^{\frac{3}{2}} \left(\frac{\operatorname{C}\left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right) \left(\left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right)^2 \sqrt{\pi }-\left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right) \sqrt{2}\right)}{\sqrt{\pi }}-\frac{\left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right) \sin \left(\frac{\pi \left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right)^2}{2}\right)}{\sqrt{\pi }}}{4}$

[In] `int(x^2*cos(1/4+x+x^2), x, method=_RETURNVERBOSE)`

[Out] `1/2*x*sin(1/4+x+x^2)-1/4*sin(1/4+x+x^2)+1/8*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))-1/4*2^(1/2)*Pi^(1/2)*FresnelS(2^(1/2)/Pi^(1/2)*(x+1/2))`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.72

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{8} \sqrt{2} \sqrt{\pi} C \left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}} \right) - \frac{1}{4} \sqrt{2} \sqrt{\pi} S \left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}} \right) + \frac{1}{4} (2x-1) \sin \left(x^2 + x + \frac{1}{4} \right)$$

```
[In] integrate(x^2*cos(1/4+x+x^2),x, algorithm="fricas")
[Out] 1/8*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) - 1/4*sqrt(2)*sqrt(pi)*fresnel_sin(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) + 1/4*(2*x - 1)*sin(x^2 + x + 1/4)
```

Sympy [F]

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \cos \left(x^2 + x + \frac{1}{4} \right) dx$$

```
[In] integrate(x**2*cos(1/4+x+x**2),x)
[Out] Integral(x**2*cos(x**2 + x + 1/4), x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.95

$$\int x^2 \cos \left(\frac{1}{4} + x + x^2 \right) dx = \frac{-16x \left(-i e^{(ix^2+ix+\frac{1}{4}i)} + i e^{(-ix^2-ix-\frac{1}{4}i)} \right) + \sqrt{4x^2+4x+1} \left((i-1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{ix^2+ix+\frac{1}{4}i} \right) - 1 \right) - (I+1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{-ix^2-ix-\frac{1}{4}i} \right) - 1 \right) - (4I+4) \sqrt{2} \gamma(3/2, ix^2+ix+1/4) + (4I-4) \sqrt{2} \gamma(3/2, -ix^2-ix-1/4) \right)}{(2x+1)}$$

```
[In] integrate(x^2*cos(1/4+x+x^2),x, algorithm="maxima")
[Out] -1/32*(16*x*(-I*e^(Ix^2 + I*x + 1/4*I) + I*e^(-Ix^2 - I*x - 1/4*I)) + sqrt(4*x^2 + 4*x + 1)*((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(I*x^2 + I*x + 1/4*I)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(-Ix^2 - I*x - 1/4*I)) - 1) - (4*I + 4)*sqrt(2)*gamma(3/2, I*x^2 + I*x + 1/4*I) + (4*I - 4)*sqrt(2)*gamma(3/2, -Ix^2 - I*x - 1/4*I)) - 8*I*e^(Ix^2 + I*x + 1/4*I) + 8*I*e^(-Ix^2 - I*x - 1/4*I))/(2*x + 1)
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.91

$$\begin{aligned} \int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = & -\left(\frac{3}{32}i - \frac{1}{32}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2x+1)\right) \\ & + \left(\frac{3}{32}i + \frac{1}{32}\right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2x+1)\right) \\ & + \frac{1}{8} (-2ix + i)e^{(ix^2 + ix + \frac{1}{4}i)} + \frac{1}{8} (2ix - i)e^{(-ix^2 - ix - \frac{1}{4}i)} \end{aligned}$$

```
[In] integrate(x^2*cos(1/4+x+x^2),x, algorithm="giac")
[Out] -(3/32*I - 1/32)*sqrt(2)*sqrt(pi)*erf((1/4*I - 1/4)*sqrt(2)*(2*x + 1)) + (3/32*I + 1/32)*sqrt(2)*sqrt(pi)*erf(-(1/4*I + 1/4)*sqrt(2)*(2*x + 1)) + 1/8*(-2*I*x + I)*e^(I*x^2 + I*x + 1/4*I) + 1/8*(2*I*x - I)*e^(-I*x^2 - I*x - 1/4*I)
```

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.78

$$\begin{aligned} \int x^2 \cos\left(\frac{1}{4} + x + x^2\right) dx = & \frac{x \sin(x^2 + x + \frac{1}{4})}{2} - \frac{\sin(x^2 + x + \frac{1}{4})}{4} \\ & + \frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{8} - \frac{\sqrt{2}\sqrt{\pi} S\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{4} \end{aligned}$$

```
[In] int(x^2*cos(x + x^2 + 1/4),x)
[Out] (x*sin(x + x^2 + 1/4))/2 - sin(x + x^2 + 1/4)/4 + (2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(2*x + 1))/(2*pi^(1/2)))/8 - (2^(1/2)*pi^(1/2)*fresnels((2^(1/2)*(2*x + 1))/(2*pi^(1/2))))/4
```

3.12 $\int x \cos\left(\frac{1}{4} + x + x^2\right) dx$

Optimal result	94
Rubi [A] (verified)	94
Mathematica [A] (verified)	95
Maple [A] (verified)	95
Fricas [A] (verification not implemented)	96
Sympy [B] (verification not implemented)	96
Maxima [C] (verification not implemented)	97
Giac [C] (verification not implemented)	97
Mupad [B] (verification not implemented)	98

Optimal result

Integrand size = 11, antiderivative size = 41

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) + \frac{1}{2} \sin\left(\frac{1}{4} + x + x^2\right)$$

[Out] $\frac{1}{2} \sin(1/4+x+x^2) - \frac{1}{4} \operatorname{FresnelC}(1/2*(1+2*x)*2^{(1/2)}/\operatorname{Pi}^{(1/2)}) * 2^{(1/2)} * \operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3543, 3527, 3433}

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{2} \sin\left(x^2 + x + \frac{1}{4}\right) - \frac{1}{2} \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{2x+1}{\sqrt{2\pi}}\right)$$

[In] $\operatorname{Int}[x \cos[1/4 + x + x^2], x]$

[Out] $-\frac{1}{2} (\operatorname{Sqrt}[\operatorname{Pi}/2] \operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[2*\operatorname{Pi}]]) + \operatorname{Sin}[1/4 + x + x^2]/2$

Rule 3433

```
Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3527

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_.)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_) + (b_)*(x_) + (c_)*(x_)^2]*((d_) + (e_)*(x_)), x_Symbol]
:= Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} \sin\left(\frac{1}{4} + x + x^2\right) - \frac{1}{2} \int \cos\left(\frac{1}{4} + x + x^2\right) dx \\ &= \frac{1}{2} \sin\left(\frac{1}{4} + x + x^2\right) - \frac{1}{2} \int \cos\left(\frac{1}{4}(1+2x)^2\right) dx \\ &= -\frac{1}{2} \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) + \frac{1}{2} \sin\left(\frac{1}{4} + x + x^2\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.95

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{4} \left(-\sqrt{2\pi} \text{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right) + 2 \sin\left(\frac{1}{4} + x + x^2\right) \right)$$

[In] `Integrate[x*Cos[1/4 + x + x^2], x]`

[Out] `(-(Sqrt[2*Pi])*FresnelC[(1 + 2*x)/Sqrt[2*Pi]]) + 2*Sin[1/4 + x + x^2])/4`

Maple [A] (verified)

Time = 1.58 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

method	result	size
default	$\frac{\sin\left(\frac{1}{4}+x+x^2\right)}{2}-\frac{\sqrt{2} \sqrt{\pi } \text{C}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi }}\right)}{4}$	30
risch	$\frac{\sqrt{\pi } (-1)^{\frac{3}{4}} \text{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{8}-\frac{\sqrt{\pi } \text{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{8 \sqrt{-i}}+\frac{\sin \left(\frac{(1+2 x)^2}{4}\right)}{2}$	58
parts	$\frac{\sqrt{2} \sqrt{\pi } \text{C}\left(\frac{\sqrt{2} \left(x+\frac{1}{2}\right)}{\sqrt{\pi }}\right) x}{2}-\frac{\pi \left(\text{C}\left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right) \left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right)-\frac{\sin \left(\frac{\pi \left(\frac{\sqrt{2} x}{\sqrt{\pi }}+\frac{\sqrt{2}}{2 \sqrt{\pi }}\right)^2}{2}\right)}{\pi }\right)}{2}$	90

[In] `int(x*cos(1/4+x+x^2),x,method=_RETURNVERBOSE)`
[Out] `1/2*sin(1/4+x+x^2)-1/4*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = -\frac{1}{4} \sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right) + \frac{1}{2} \sin\left(x^2 + x + \frac{1}{4}\right)$$

[In] `integrate(x*cos(1/4+x+x^2),x, algorithm="fricas")`
[Out] `-1/4*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi)) + 1/2*sin(x^2 + x + 1/4)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(39) = 78$.

Time = 0.66 (sec) , antiderivative size = 155, normalized size of antiderivative = 3.78

$$\begin{aligned} \int x \cos\left(\frac{1}{4} + x + x^2\right) dx &= -\frac{\sqrt{2}\sqrt{\pi}xC\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{1}{4}\right)}{8\Gamma\left(\frac{5}{4}\right)} + \frac{\sqrt{2}\sqrt{\pi}xC\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)}{2} \\ &\quad + \frac{\sin\left((x + \frac{1}{2})^2\right)\Gamma\left(\frac{1}{4}\right)}{8\Gamma\left(\frac{5}{4}\right)} - \frac{\sqrt{2}\sqrt{\pi}C\left(\frac{\sqrt{2}x}{\sqrt{\pi}} + \frac{\sqrt{2}}{2\sqrt{\pi}}\right)\Gamma\left(\frac{1}{4}\right)}{16\Gamma\left(\frac{5}{4}\right)} \end{aligned}$$

[In] `integrate(x*cos(1/4+x+x**2),x)`
[Out] `-sqrt(2)*sqrt(pi)*x*fresnelc(sqrt(2)*x/sqrt(pi) + sqrt(2)/(2*sqrt(pi)))*gamma(1/4)/(8*gamma(5/4)) + sqrt(2)*sqrt(pi)*x*fresnelc(sqrt(2)*x/sqrt(pi) + sqrt(2)/(2*sqrt(pi)))/2 + sin((x + 1/2)**2)*gamma(1/4)/(8*gamma(5/4)) - sqrt(2)*sqrt(pi)*fresnelc(sqrt(2)*x/sqrt(pi) + sqrt(2)/(2*sqrt(pi)))*gamma(1/4)/(16*gamma(5/4))`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 126, normalized size of antiderivative = 3.07

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx \\ = \frac{8x \left(-i e^{(ix^2+ix+\frac{1}{4}i)} + i e^{(-ix^2-ix-\frac{1}{4}i)}\right) + \sqrt{4x^2+4x+1} \left((i-1) \sqrt{2}\sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{ix^2+ix+\frac{1}{4}i}\right) - 1\right) - (i+1) \sqrt{2}\sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{-ix^2-ix-\frac{1}{4}i}\right) - 1\right)\right)}{16(2x+1)}$$

[In] `integrate(x*cos(1/4+x+x^2),x, algorithm="maxima")`

[Out] $\frac{1}{16} \left(8x \left(-i e^{(Ix^2 + Ix + 1/4I)} + i e^{(-Ix^2 - Ix - 1/4I)} \right) + \sqrt{4x^2 + 4x + 1} \left((I - 1) \sqrt{2}\sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{Ix^2 + Ix + 1/4I}\right) - 1 \right) - (I + 1) \sqrt{2}\sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{-Ix^2 - Ix - 1/4I}\right) - 1 \right) \right) \right) / (2x + 1)$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.47 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = \left(\frac{1}{16}i + \frac{1}{16} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right) \sqrt{2}(2x+1)\right) \\ - \left(\frac{1}{16}i - \frac{1}{16} \right) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right) \sqrt{2}(2x+1)\right) \\ - \frac{1}{4}i e^{(ix^2+ix+\frac{1}{4}i)} + \frac{1}{4}i e^{(-ix^2-ix-\frac{1}{4}i)}$$

[In] `integrate(x*cos(1/4+x+x^2),x, algorithm="giac")`

[Out] $\frac{(1/16I + 1/16) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left((1/4I - 1/4) \sqrt{2}(2x+1)\right) - (1/16I - 1/16) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(-(1/4I + 1/4) \sqrt{2}(2x+1)\right) - 1/4I * e^{(Ix^2 + Ix + 1/4I)} + 1/4I * e^{(-Ix^2 - Ix - 1/4I)}}{2x+1}$

Mupad [B] (verification not implemented)

Time = 14.53 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.78

$$\int x \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sin(x^2 + x + \frac{1}{4})}{2} - \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{4}$$

[In] `int(x*cos(x + x^2 + 1/4),x)`

[Out] `sin(x + x^2 + 1/4)/2 - (2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(2*x + 1))/(2*pi^(1/2))))/4`

3.13 $\int \cos\left(\frac{1}{4} + x + x^2\right) dx$

Optimal result	99
Rubi [A] (verified)	99
Mathematica [A] (verified)	100
Maple [A] (verified)	100
Fricas [A] (verification not implemented)	100
Sympy [A] (verification not implemented)	101
Maxima [C] (verification not implemented)	101
Giac [C] (verification not implemented)	101
Mupad [B] (verification not implemented)	102

Optimal result

Integrand size = 9, antiderivative size = 24

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

[Out] $1/2*\operatorname{FresnelC}(1/2*(1+2*x)*2^{(1/2)}/\operatorname{Pi}^{(1/2)})*2^{(1/2)}*\operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3527, 3433}

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \operatorname{FresnelC}\left(\frac{2x+1}{\sqrt{2\pi}}\right)$$

[In] $\operatorname{Int}[\operatorname{Cos}[1/4 + x + x^2], x]$

[Out] $\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[2*\operatorname{Pi}]]$

Rule 3433

```
Int[Cos[(d_.)*(e_.) + (f_.)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3527

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}\text{integral} &= \int \cos\left(\frac{1}{4}(1+2x)^2\right) dx \\ &= \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right)\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.00

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{1+2x}{\sqrt{2\pi}}\right)$$

[In] `Integrate[Cos[1/4 + x + x^2], x]`
 [Out] `Sqrt[Pi/2]*FresnelC[(1 + 2*x)/Sqrt[2*Pi]]`

Maple [A] (verified)

Time = 0.66 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{2}$	20
risch	$-\frac{\sqrt{\pi}(-1)^{\frac{3}{4}} \operatorname{erf}\left((-1)^{\frac{1}{4}} x+\frac{(-1)^{\frac{1}{4}}}{2}\right)}{4}+\frac{\sqrt{\pi} \operatorname{erf}\left(\sqrt{-i} x-\frac{i}{2 \sqrt{-i}}\right)}{4 \sqrt{-i}}$	46

[In] `int(cos(1/4+x+x^2),x,method=_RETURNVERBOSE)`
 [Out] `1/2*2^(1/2)*Pi^(1/2)*FresnelC(2^(1/2)/Pi^(1/2)*(x+1/2))`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.92

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{2} \sqrt{2}\sqrt{\pi} C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)$$

[In] `integrate(cos(1/4+x+x^2),x, algorithm="fricas")`
 [Out] `1/2*sqrt(2)*sqrt(pi)*fresnel_cos(1/2*sqrt(2)*(2*x + 1)/sqrt(pi))`

Sympy [A] (verification not implemented)

Time = 0.20 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.21

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2}\sqrt{\pi}C\left(\frac{\sqrt{2}(2x+1)}{2\sqrt{\pi}}\right)}{2}$$

[In] `integrate(cos(1/4+x+x**2),x)`

[Out] `sqrt(2)*sqrt(pi)*fresnelc(sqrt(2)*(2*x + 1)/(2*sqrt(pi)))/2`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 70, normalized size of antiderivative = 2.92

$$\begin{aligned} \int \cos\left(\frac{1}{4} + x + x^2\right) dx = \\ -\frac{1}{16}\sqrt{\pi}\left((i-1)\sqrt{2}\operatorname{erf}\left(-\frac{1}{2}(-1)^{\frac{3}{4}}(2ix+i)\right) + (i-1)\sqrt{2}\operatorname{erf}\left(-\left(\frac{1}{4}i-\frac{1}{4}\right)\sqrt{2}(2ix+i)\right) - (i+1)\right. \end{aligned}$$

[In] `integrate(cos(1/4+x+x^2),x, algorithm="maxima")`

[Out] `-1/16*sqrt(pi)*((I - 1)*sqrt(2)*erf(-1/2*(-1)^(3/4)*(2*I*x + I)) + (I - 1)*sqrt(2)*erf(-(1/4*I - 1/4)*sqrt(2)*(2*I*x + I)) - (I + 1)*sqrt(2)*erf(-(1/4*I + 1/4)*sqrt(2)*(2*I*x + I)) + (I + 1)*sqrt(2)*erf(1/2*(2*I*x + I)/sqrt(-I)))`

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.40 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.62

$$\begin{aligned} \int \cos\left(\frac{1}{4} + x + x^2\right) dx = -\left(\frac{1}{8}i + \frac{1}{8}\right)\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(\left(\frac{1}{4}i - \frac{1}{4}\right)\sqrt{2}(2x+1)\right) \\ + \left(\frac{1}{8}i - \frac{1}{8}\right)\sqrt{2}\sqrt{\pi}\operatorname{erf}\left(-\left(\frac{1}{4}i + \frac{1}{4}\right)\sqrt{2}(2x+1)\right) \end{aligned}$$

[In] `integrate(cos(1/4+x+x^2),x, algorithm="giac")`

[Out] `-(1/8*I + 1/8)*sqrt(2)*sqrt(pi)*erf((1/4*I - 1/4)*sqrt(2)*(2*x + 1)) + (1/8*I - 1/8)*sqrt(2)*sqrt(pi)*erf(-(1/4*I + 1/4)*sqrt(2)*(2*x + 1))`

Mupad [B] (verification not implemented)

Time = 0.04 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.79

$$\int \cos\left(\frac{1}{4} + x + x^2\right) dx = \frac{\sqrt{2} \sqrt{\pi} C\left(\frac{\sqrt{2}(x+\frac{1}{2})}{\sqrt{\pi}}\right)}{2}$$

[In] `int(cos(x + x^2 + 1/4),x)`

[Out] `(2^(1/2)*pi^(1/2)*fresnelc((2^(1/2)*(x + 1/2))/pi^(1/2)))/2`

3.14 $\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	103
Rubi [N/A]	103
Mathematica [N/A]	104
Maple [N/A] (verified)	104
Fricas [N/A]	104
Sympy [N/A]	104
Maxima [N/A]	105
Giac [N/A]	105
Mupad [N/A]	105

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \text{Int}\left(\frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x}, x\right)$$

[Out] Unintegrable(cos(1/4+x+x^2)/x,x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec), antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] Int[Cos[1/4 + x + x^2]/x,x]

[Out] Defer[Int][Cos[1/4 + x + x^2]/x, x]

Rubi steps

$$\text{integral} = \int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 9.94 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] `Integrate[Cos[1/4 + x + x^2]/x, x]`

[Out] `Integrate[Cos[1/4 + x + x^2]/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.18 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] `int(cos(1/4+x+x^2)/x, x)`

[Out] `int(cos(1/4+x+x^2)/x, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)/x, x, algorithm="fricas")`

[Out] `integral(cos(x^2 + x + 1/4)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `integrate(cos(1/4+x+x**2)/x, x)`

[Out] `Integral(cos(x**2 + x + 1/4)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)/x,x, algorithm="maxima")`

[Out] `integrate(cos(x^2 + x + 1/4)/x, x)`

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)/x,x, algorithm="giac")`

[Out] `integrate(cos(x^2 + x + 1/4)/x, x)`

Mupad [N/A]

Not integrable

Time = 13.77 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `int(cos(x + x^2 + 1/4)/x,x)`

[Out] `int(cos(x + x^2 + 1/4)/x, x)`

3.15 $\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	106
Rubi [N/A]	106
Mathematica [N/A]	107
Maple [N/A] (verified)	107
Fricas [N/A]	107
Sympy [N/A]	108
Maxima [N/A]	108
Giac [N/A]	108
Mupad [N/A]	109

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx = -\frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x} - \sqrt{2\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) \\ - \operatorname{Int}\left(\frac{\sin\left(\frac{1}{4}+x+x^2\right)}{x}, x\right)$$

[Out] $-\cos(1/4+x+x^2)/x - \operatorname{FresnelS}(1/2*(1+2*x)*2^(1/2)/\operatorname{Pi}^{(1/2)})*2^(1/2)*\operatorname{Pi}^{(1/2)} - \text{nintegrable}(\sin(1/4+x+x^2)/x, x)$

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx = \int \frac{\cos\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$$

[In] $\operatorname{Int}[\operatorname{Cos}[1/4 + x + x^2]/x^2, x]$

[Out] $-(\operatorname{Cos}[1/4 + x + x^2]/x) - \operatorname{Sqrt}[2*\operatorname{Pi}]*\operatorname{FresnelS}[(1 + 2*x)/\operatorname{Sqrt}[2*\operatorname{Pi}]] - \operatorname{Defer}[\operatorname{Int}[\operatorname{Sin}[1/4 + x + x^2]/x, x]]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\cos(\frac{1}{4} + x + x^2)}{x} - 2 \int \sin\left(\frac{1}{4} + x + x^2\right) dx - \int \frac{\sin(\frac{1}{4} + x + x^2)}{x} dx \\
 &= -\frac{\cos(\frac{1}{4} + x + x^2)}{x} - 2 \int \sin\left(\frac{1}{4}(1+2x)^2\right) dx - \int \frac{\sin(\frac{1}{4} + x + x^2)}{x} dx \\
 &= -\frac{\cos(\frac{1}{4} + x + x^2)}{x} - \sqrt{2\pi} \text{FresnelS}\left(\frac{1+2x}{\sqrt{2\pi}}\right) - \int \frac{\sin(\frac{1}{4} + x + x^2)}{x} dx
 \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 10.62 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

$$\int \frac{\cos(\frac{1}{4} + x + x^2)}{x^2} dx = \int \frac{\cos(\frac{1}{4} + x + x^2)}{x^2} dx$$

[In] `Integrate[Cos[1/4 + x + x^2]/x^2, x]`

[Out] `Integrate[Cos[1/4 + x + x^2]/x^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.21 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \frac{\cos(\frac{1}{4} + x + x^2)}{x^2} dx$$

[In] `int(cos(1/4+x+x^2)/x^2, x)`

[Out] `int(cos(1/4+x+x^2)/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos(\frac{1}{4} + x + x^2)}{x^2} dx = \int \frac{\cos(x^2 + x + \frac{1}{4})}{x^2} dx$$

[In] `integrate(cos(1/4+x+x^2)/x^2, x, algorithm="fricas")`

[Out] `integral(cos(x^2 + x + 1/4)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.08

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

```
[In] integrate(cos(1/4+x+x**2)/x**2,x)
[Out] Integral(cos(x**2 + x + 1/4)/x**2, x)
```

Maxima [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

```
[In] integrate(cos(1/4+x+x^2)/x^2,x, algorithm="maxima")
[Out] integrate(cos(x^2 + x + 1/4)/x^2, x)
```

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

```
[In] integrate(cos(1/4+x+x^2)/x^2,x, algorithm="giac")
[Out] integrate(cos(x^2 + x + 1/4)/x^2, x)
```

Mupad [N/A]

Not integrable

Time = 14.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\cos\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

[In] int(cos(x + x^2 + 1/4)/x^2, x)

[Out] int(cos(x + x^2 + 1/4)/x^2, x)

3.16 $\int x^2 \cos^2(a + bx + cx^2) dx$

Optimal result	110
Rubi [A] (verified)	111
Mathematica [A] (verified)	113
Maple [A] (verified)	113
Fricas [A] (verification not implemented)	114
Sympy [F]	114
Maxima [C] (verification not implemented)	115
Giac [C] (verification not implemented)	116
Mupad [F(-1)]	117

Optimal result

Integrand size = 17, antiderivative size = 248

$$\begin{aligned} \int x^2 \cos^2(a + bx + cx^2) dx = & \frac{x^3}{6} + \frac{b^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

```
[Out] 1/6*x^3-1/16*b*sin(2*c*x^2+2*b*x+2*a)/c^2+1/8*x*sin(2*c*x^2+2*b*x+2*a)/c+1/
16*b^2*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/c^(5/2)-
1/16*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/c^(3/2)-
1/16*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)*Pi^(1/2)/c^(3/2)-
1/16*b^2*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.33 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.412, Rules used = {3549, 3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int x^2 \cos^2(a + bx + cx^2) dx = & -\frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & + \frac{\sqrt{\pi}b^2 \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & - \frac{\sqrt{\pi}b^2 \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} + \frac{x^3}{6} \end{aligned}$$

[In] $\text{Int}[x^2 \cos[a + b*x + c*x^2]^2, x]$

[Out] $x^3/6 + (b^2 * \text{Sqrt}[\text{Pi}] * \cos[2*a - b^2/(2*c)] * \text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(5/2)}) - (\text{Sqrt}[\text{Pi}] * \cos[2*a - b^2/(2*c)] * \text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])])/(16*c^{(3/2)}) - (\text{Sqrt}[\text{Pi}] * \text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])] * \text{Sin}[2*a - b^2/(2*c)])/(16*c^{(3/2)}) - (b^2 * \text{Sqrt}[\text{Pi}] * \text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])] * \text{Sin}[2*a - b^2/(2*c)])/(16*c^{(5/2)}) - (b * \text{Sin}[2*a + 2*b*x + 2*c*x^2])/(16*c^2) + (x * \text{Sin}[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Rule 3432

$\text{Int}[\text{Sin}[(d_.) * ((e_.) + (f_.) * (x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f * \text{Rt}[d, 2])) * \text{FresnelS}[\text{Sqrt}[2/\text{Pi}] * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\text{Cos}[(d_.) * ((e_.) + (f_.) * (x_))^2], x_{\text{Symbol}}] \Rightarrow \text{Simp}[(\text{Sqrt}[\text{Pi}/2]/(f * \text{Rt}[d, 2])) * \text{FresnelC}[\text{Sqrt}[2/\text{Pi}] * \text{Rt}[d, 2] * (e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3528

$\text{Int}[\text{Sin}[(a_.) + (b_.) * (x_) + (c_.) * (x_)^2], x_{\text{Symbol}}] \Rightarrow \text{Dist}[\text{Cos}[(b^2 - 4*a*c)/(4*c)], \text{Int}[\text{Sin}[(b + 2*c*x)^2/(4*c)], x], x] - \text{Dist}[\text{Sin}[(b^2 - 4*a*c)/(4*c)], \text{Int}[\text{Cos}[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 3529

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]*((d_.) + (e_ .)*(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]*((d_.) + (e_ .)*(x_)^(m_)), x_Sym
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[
(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - D
ist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x]
) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rule 3549

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]^(n_)*((d_.) + (e_ .)*(x_)^(m_)), x_Symbol]
:> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{x^2}{2} + \frac{1}{2}x^2 \cos(2a + 2bx + 2cx^2) \right) dx \\
&= \frac{x^3}{6} + \frac{1}{2} \int x^2 \cos(2a + 2bx + 2cx^2) dx \\
&= \frac{x^3}{6} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} - \frac{\int \sin(2a + 2bx + 2cx^2) dx}{8c} - \frac{b \int x \cos(2a + 2bx + 2cx^2) dx}{4c} \\
&= \frac{x^3}{6} - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} \\
&\quad + \frac{b^2 \int \cos(2a + 2bx + 2cx^2) dx}{8c^2} - \frac{\cos\left(2a - \frac{b^2}{2c}\right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c} \\
&\quad - \frac{\sin\left(2a - \frac{b^2}{2c}\right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x^3}{6} - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\
&\quad - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c} \\
&\quad + \frac{\left(b^2 \cos\left(2a - \frac{b^2}{2c}\right)\right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c^2} \\
&\quad - \frac{\left(b^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c^2} \\
&= \frac{x^3}{6} + \frac{b^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} - \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\
&\quad - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\
&\quad - \frac{b \sin(2a + 2bx + 2cx^2)}{16c^2} + \frac{x \sin(2a + 2bx + 2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.45 (sec), antiderivative size = 170, normalized size of antiderivative = 0.69

$$\begin{aligned}
&\int x^2 \cos^2(a + bx + cx^2) dx \\
&= \frac{-3\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a - \frac{b^2}{2c}\right) + b^2 \sin\left(2a - \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a - \frac{b^2}{2c}\right)\right)}{48c^{5/2}}
\end{aligned}$$

[In] Integrate[x^2*Cos[a + b*x + c*x^2]^2, x]

[Out]
$$\begin{aligned}
&(-3*\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])*(c*\text{Cos}[2*a - b^2/(2*c)]) \\
&+ b^2*\text{Sin}[2*a - b^2/(2*c)] + 3*\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*(b^2*\text{Cos}[2*a - b^2/(2*c)] - c*\text{Sin}[2*a - b^2/(2*c)]) + \text{Sqrt}[c]*(8*c^2*x^3 - 3*(b - 2*c*x)*\text{Sin}[2*(a + x*(b + c*x))]))/(48*c^{(5/2)})
\end{aligned}$$

Maple [A] (verified)

Time = 0.68 (sec), antiderivative size = 191, normalized size of antiderivative = 0.77

method	result
default	$\frac{x^3}{6} + \frac{x \sin(2cx^2+2bx+2a)}{8c} - \frac{b \left(\frac{\sin(2cx^2+2bx+2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^2} \right)}{4c} - \frac{\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^2}$
risch	$\frac{x^3}{6} + \frac{b^2 \sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2} \sqrt{ic} x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c^2\sqrt{ic}} - \frac{i\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2} \sqrt{ic} x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{64c\sqrt{ic}} - \frac{b^2 \sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \operatorname{erf}\left(-\sqrt{2} \sqrt{ic} x - \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c^2\sqrt{-2ic}}$

[In] `int(x^2*cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{6}x^3 + \frac{1}{8}x \sin(2cx^2+2bx+2a) - \frac{b}{4c} \left(\frac{\sin(2cx^2+2bx+2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^2} \right) - \frac{\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^2}$$

Fricas [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 178, normalized size of antiderivative = 0.72

$$\int x^2 \cos^2(a + bx + cx^2) dx = \frac{8c^3x^3 + 6(2c^2x - bc)\cos(cx^2 + bx + a)\sin(cx^2 + bx + a) + 3\left(\pi b^2 \cos\left(-\frac{b^2-4ac}{2c}\right) - \pi c \sin\left(-\frac{b^2-4ac}{2c}\right)\right)\sqrt{c}}{48c^3}$$

[In] `integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out]
$$\frac{1}{48}(8c^3x^3 + 6(2c^2x - bc)\cos(cx^2 + bx + a)\sin(cx^2 + bx + a) + 3(\pi b^2 \cos(-\frac{b^2-4ac}{2c}) - \pi c \sin(-\frac{b^2-4ac}{2c}))\sqrt{c})$$

Sympy [F]

$$\int x^2 \cos^2(a + bx + cx^2) dx = \int x^2 \cos^2(a + bx + cx^2) dx$$

[In] `integrate(x**2*cos(c*x**2+b*x+a)**2,x)`

[Out] `Integral(x**2*cos(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.06 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

[In] `integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")`

[Out]

```

-1/384*sqrt(2)*(24*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*sin(-1/2*(b^2 - 4*a*c)/c))*x^3 + 36*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*sin(-1/2*(b^2 - 4*a*c)/c))*x^2 - 4*sqrt(2)*(8*c^4*x^3 - 3*b*c^2*(-I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) - 3*b*c^2*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*(4*c^2*x^2 + 4*b*c*x + b^2/c)^(3/2) + 18*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*sin(-1/2*(b^2 - 4*a*c)/c))*x + 3*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*sin(-1/2*(b^2 - 4*a*c)/c)) + 3*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^2*c^2)*cos(-1/2*(b^2 - 4*a*c)/c))

```

$$\begin{aligned}
& /2)*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I + 1)*\sqrt(2)*\sqrt(pi)*(erf(sqrt(1/2)*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1))*b^5 + \\
& 2*(-(I + 1)*\sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*\sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^3*c) \\
& *cos(-1/2*(b^2 - 4*a*c)/c) + 3*((I + 1)*\sqrt(2)*\sqrt(pi)*(erf(sqrt(1/2)*\sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*\sqrt(2)*\sqrt(pi)*(erf(sqrt(1/2)*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1))*b^5 + 2*((I - 1)*\sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*\sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^3*c)*sin(-1/2*(b^2 - 4*a*c)/c))/(c^4*((4*c^2*x^2 + 4*b*c*x + b^2)/c)^(3/2))
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 214, normalized size of antiderivative = 0.86

$$\begin{aligned}
& \int x^2 \cos^2(a + bx + cx^2) dx \\
& = \frac{1}{6} x^3 \\
& - \frac{(c(2i x + \frac{ib}{c}) - 2i b)e^{(2icx^2+2ibx+2ia)} - \frac{i\sqrt{\pi}(b^2+ic)\operatorname{erf}\left(-\frac{1}{2}i\sqrt{c}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{ib^2+4iac}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|}+1\right)}}{32c^2} \\
& - \frac{(c(-2i x - \frac{ib}{c}) + 2i b)e^{(-2icx^2-2ibx-2ia)} + \frac{i\sqrt{\pi}(b^2-ic)\operatorname{erf}\left(\frac{1}{2}i\sqrt{c}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\right)e^{\left(-\frac{-ib^2+4iac}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|}+1\right)}}{32c^2}
\end{aligned}$$

```
[In] integrate(x^2*cos(c*x^2+b*x+a)^2,x, algorithm="giac")
[Out] 1/6*x^3 - 1/32*((c*(2*I*x + I*b/c) - 2*I*b)*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a)
- I*sqrt(pi)*(b^2 + I*c)*erf(-1/2*I*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1))/c^2 - 1/32*((c*(-2*I*x - I*b/c) + 2*I*b)*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a) + I*sqrt(pi)*(b^2 - I*c)*erf(1/2*I*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1))/c^2
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2(a + bx + cx^2) dx = \int x^2 \cos(cx^2 + bx + a)^2 dx$$

[In] int(x^2*cos(a + b*x + c*x^2)^2,x)

[Out] int(x^2*cos(a + b*x + c*x^2)^2, x)

3.17 $\int x \cos^2(a + bx + cx^2) dx$

Optimal result	118
Rubi [A] (verified)	118
Mathematica [A] (verified)	120
Maple [A] (verified)	120
Fricas [A] (verification not implemented)	121
Sympy [F]	121
Maxima [C] (verification not implemented)	121
Giac [C] (verification not implemented)	122
Mupad [F(-1)]	123

Optimal result

Integrand size = 15, antiderivative size = 126

$$\begin{aligned} \int x \cos^2(a + bx + cx^2) dx = & \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & + \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} + \frac{\sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

[Out] $1/4*x^2 + 1/8*\sin(2*c*x^2 + 2*b*x + 2*a)/c - 1/8*b*\cos(2*a - 1/2*b^2/c)*\text{FresnelC}((2*c*x + b)/c^{(1/2)}/\text{Pi}^{(1/2)})*\text{Pi}^{(1/2)}/c^{(3/2)} + 1/8*b*\text{FresnelS}((2*c*x + b)/c^{(1/2)}/\text{Pi}^{(1/2)})*\sin(2*a - 1/2*b^2/c)*\text{Pi}^{(1/2)}/c^{(3/2)}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3549, 3543, 3529, 3433, 3432}

$$\begin{aligned} \int x \cos^2(a + bx + cx^2) dx = & -\frac{\sqrt{\pi}b \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & + \frac{\sqrt{\pi}b \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & + \frac{\sin(2a + 2bx + 2cx^2)}{8c} + \frac{x^2}{4} \end{aligned}$$

[In] Int[x*Cos[a + b*x + c*x^2]^2, x]

[Out] $x^{2/4} - (b\sqrt{\pi})\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]/(8c^{3/2}) + (b\sqrt{\pi})\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)]/(8c^{3/2}) + \sin[2a + 2bx + 2cx^2]/(8c)$

Rule 3432

$\text{Int}[\sin[(d_.)\cdot(e_.) + (f_.)\cdot(x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f\sqrt{d}), x] \cdot \text{FresnelS}[\sqrt{2/\pi}\sqrt{d}\cdot(e + f\cdot x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\cos[(d_.)\cdot(e_.) + (f_.)\cdot(x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f\sqrt{d}), x] \cdot \text{FresnelC}[\sqrt{2/\pi}\sqrt{d}\cdot(e + f\cdot x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3529

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Dist}[\cos[b^2 - 4a*c]/(4c), \text{Int}[\cos[(b + 2cx)^2/(4c)], x], x] + \text{Dist}[\sin[b^2 - 4a*c]/(4c), \text{Int}[\sin[(b + 2cx)^2/(4c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4a*c, 0]$

Rule 3543

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2]\cdot((d_.) + (e_.)\cdot(x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[e\cdot(\sin[a + bx + cx^2]/(2c)), x] + \text{Dist}[(2c*d - b*e)/(2c), \text{Int}[\cos[a + bx + cx^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2c*d - b*e, 0]$

Rule 3549

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2]^n\cdot((d_.) + (e_.)\cdot(x_.)^m), x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \cos[a + bx + cx^2]^n], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{IGtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{x}{2} + \frac{1}{2}x \cos(2a + 2bx + 2cx^2) \right) dx \\ &= \frac{x^2}{4} + \frac{1}{2} \int x \cos(2a + 2bx + 2cx^2) dx \\ &= \frac{x^2}{4} + \frac{\sin(2a + 2bx + 2cx^2)}{8c} - \frac{b \int \cos(2a + 2bx + 2cx^2) dx}{4c} \\ &= \frac{x^2}{4} + \frac{\sin(2a + 2bx + 2cx^2)}{8c} - \frac{\left(b \cos\left(2a - \frac{b^2}{2c}\right) \right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{4c} \\ &\quad + \frac{\left(b \sin\left(2a - \frac{b^2}{2c}\right) \right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{4c} \end{aligned}$$

$$\begin{aligned}
&= \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\
&\quad + \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} + \frac{\sin(2a + 2bx + 2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec), antiderivative size = 116, normalized size of antiderivative = 0.92

$$\begin{aligned}
&\int x \cos^2(a + bx + cx^2) dx \\
&= \frac{-b\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + b\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 + \sin(2(a + x(b + cx^2))))}{8c^{3/2}}
\end{aligned}$$

[In] `Integrate[x*Cos[a + b*x + c*x^2]^2, x]`

[Out] $\frac{(-b\sqrt{\pi} \cos(2a - \frac{b^2}{2c}) \text{FresnelC}((b + 2c*x)/(\sqrt{c}\sqrt{\pi})) + b\sqrt{\pi} \text{FresnelS}((b + 2c*x)/(\sqrt{c}\sqrt{\pi})) \sin(2a - \frac{b^2}{2c}) + \sqrt{c}(2cx^2 + \sin(2(a + x(b + cx^2))))}{8c^{3/2}}$

Maple [A] (verified)

Time = 0.72 (sec), antiderivative size = 95, normalized size of antiderivative = 0.75

method	result	size
default	$\frac{x^2}{4} + \frac{\sin(2cx^2 + 2bx + 2a)}{8c} - \frac{b\sqrt{\pi} (\cos(\frac{-4ac+b^2}{2c}) \text{C}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}) + \sin(\frac{-4ac+b^2}{2c}) \text{S}(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}))}{8c^{3/2}}$	95
risch	$\frac{x^2}{4} - \frac{b\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \text{erf}(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{32c\sqrt{ic}} + \frac{b\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \text{erf}(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{16c\sqrt{-2ic}} + \frac{\sin(2cx^2 + 2bx + 2a)}{8c}$	141

[In] `int(x*cos(c*x^2+b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{4}x^2 + \frac{1}{8}\sin(2cx^2 + 2bx + 2a)/c - \frac{1}{8}b/c^{(3/2)}\text{Pi}^{(1/2)}(\cos(1/2*(-4a*c + b^2)/c)\text{FresnelC}((2c*x + b)/c^{(1/2)}/\text{Pi}^{(1/2)}) + \sin(1/2*(-4a*c + b^2)/c)\text{FresnelS}((2c*x + b)/c^{(1/2)}/\text{Pi}^{(1/2)}))$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 124, normalized size of antiderivative = 0.98

$$\int x \cos^2(a + bx + cx^2) dx = -\frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2 - 4ac}{2c}\right) - 2c^2x^2 - 2c \cos(cx^2 + bx + a)}{8c^2}$$

[In] `integrate(x*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out] $-1/8*(\pi*b*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - \pi*b*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) - 2*c^2*x^2 - 2*c*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2$

Sympy [F]

$$\int x \cos^2(a + bx + cx^2) dx = \int x \cos^2(a + bx + cx^2) dx$$

[In] `integrate(x*cos(c*x**2+b*x+a)**2,x)`

[Out] `Integral(x*cos(a + b*x + c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.64 (sec) , antiderivative size = 610, normalized size of antiderivative = 4.84

$$\int x \cos^2(a + bx + cx^2) dx = \frac{\sqrt{2} \left(\left((i-1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{\frac{1}{2}} \sqrt{\frac{4ic^2x^2+4ibcx+ib^2}{c}} \right) - 1 \right) - (i+1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{\frac{1}{2}} \sqrt{-\frac{4ic^2x^2+4ibcx+ib^2}{c}} \right) - 1 \right) \right)}{8c^2}$$

[In] `integrate(x*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")`

[Out] $\frac{1}{64}\sqrt{2}(((I - 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2}\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{1/2}\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1))*b^2*\cos(-1/2*(b^2 - 4*a*c)/c)$

$$\begin{aligned}
& + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) + c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) + e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.37

$$\begin{aligned}
& \int x \cos^2(a + bx + cx^2) dx \\
& = \frac{1}{4} x^2 - \frac{\frac{i \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} i \sqrt{c} \left(2x + \frac{b}{c}\right) \left(\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{i b^2 - 4 i a c}{2 c}\right)}}{\sqrt{c} \left(\frac{i c}{|c|} + 1\right)} + i e^{(2 i c x^2 + 2 i b x + 2 i a)}}{16 c} \\
& - \frac{\frac{i \sqrt{\pi} b \operatorname{erf}\left(\frac{1}{2} i \sqrt{c} \left(2x + \frac{b}{c}\right) \left(-\frac{i c}{|c|} + 1\right)\right) e^{\left(-\frac{-i b^2 + 4 i a c}{2 c}\right)}}{\sqrt{c} \left(-\frac{i c}{|c|} + 1\right)} - i e^{(-2 i c x^2 - 2 i b x - 2 i a)}}{16 c}
\end{aligned}$$

[In] integrate(x*cos(c*x^2+b*x+a)^2,x, algorithm="giac")

[Out]
$$\begin{aligned}
& 1/4*x^2 - 1/16*(I*sqrt(pi)*b*erf(-1/2*I*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1)))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) + I*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a))/c - 1/16*(-I*sqrt(pi)*b*erf(1/2*I*sqrt(c)*(2*x + b/c)*(-I*c/abs(c) + 1)))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - I*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a))/c
\end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int x \cos^2(a + bx + cx^2) dx = \int x \cos(cx^2 + bx + a)^2 dx$$

[In] int(x*cos(a + b*x + c*x^2)^2, x)

[Out] int(x*cos(a + b*x + c*x^2)^2, x)

3.18 $\int \cos^2(a + bx + cx^2) dx$

Optimal result	124
Rubi [A] (verified)	124
Mathematica [A] (verified)	125
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	126
Sympy [A] (verification not implemented)	127
Maxima [C] (verification not implemented)	127
Giac [C] (verification not implemented)	127
Mupad [F(-1)]	128

Optimal result

Integrand size = 13, antiderivative size = 100

$$\int \cos^2(a + bx + cx^2) dx = \frac{x}{2} + \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

[Out] $\frac{1}{2}x + \frac{1}{4}\cos(2a - \frac{b^2}{2c})\text{FresnelC}((2c*x+b)/c^{1/2}/\text{Pi}^{1/2})*\text{Pi}^{1/2}/c^{1/2} - \frac{1}{4}\text{FresnelS}((2c*x+b)/c^{1/2}/\text{Pi}^{1/2})*\sin(2a - \frac{b^2}{2c})*\text{Pi}^{1/2}/c^{1/2}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3531, 3529, 3433, 3432}

$$\int \cos^2(a + bx + cx^2) dx = \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2}$$

[In] Int[Cos[a + b*x + c*x^2]^2, x]

[Out] $\frac{x}{2} + (\text{Sqrt}[\text{Pi}]*\cos[2a - b^2/(2c)]*\text{FresnelC}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]/(4*\text{Sqrt}[c]) - (\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b + 2c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\sin[2a - b^2/(2c)])/(4*\text{Sqrt}[c])$

Rule 3432

```
Int[Sin[(d_)*(e_) + (f_)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[Cos[(d_)*(e_) + (f_)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[Cos[(a_)*(x_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sim[(b^2 - 4*a*c)/(4*c)], Int[Sim[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3531

```
Int[Cos[(a_)*(x_) + (b_)*(x_) + (c_)*(x_)^2]^n, x_Symbol] :> Int[ExpandTri gReduce[Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2} + \frac{1}{2} \cos(2a + 2bx + 2cx^2) \right) dx \\ &= \frac{x}{2} + \frac{1}{2} \int \cos(2a + 2bx + 2cx^2) dx \\ &= \frac{x}{2} + \frac{1}{2} \cos\left(2a - \frac{b^2}{2c}\right) \int \cos\left(\frac{(2b + 4cx)^2}{8c}\right) dx - \frac{1}{2} \sin\left(2a - \frac{b^2}{2c}\right) \int \sin\left(\frac{(2b + 4cx)^2}{8c}\right) dx \\ &= \frac{x}{2} + \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec), antiderivative size = 97, normalized size of antiderivative = 0.97

$$\begin{aligned} &\int \cos^2(a + bx + cx^2) dx \\ &= \frac{2\sqrt{c}x + \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

[In] `Integrate[Cos[a + b*x + c*x^2]^2, x]`

[Out] $(2\sqrt{c}x + \sqrt{\pi}\cos[2a - b^2/(2c)]\text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] - \sqrt{\pi}\text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a - b^2/(2c)])/(4\sqrt{c})$

Maple [A] (verified)

Time = 0.59 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} (\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right))}{4\sqrt{c}}$	72
risch	$\frac{x}{2} + \frac{\sqrt{\pi} e^{-\frac{i(4ac-b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{16\sqrt{ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac-b^2)}{2c}} \operatorname{erf}\left(-\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{8\sqrt{-2ic}}$	111

[In] `int(cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}x + \frac{1}{4}\pi^{1/2}/c^{1/2}(\cos(1/2*(-4*a*c+b^2)/c)\text{FresnelC}((2*c*x+b)/c^{1/2}/\pi^{1/2}) + \sin(1/2*(-4*a*c+b^2)/c)\text{FresnelS}((2*c*x+b)/c^{1/2}/\pi^{1/2}))$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.93

$$\int \cos^2(a + bx + cx^2) dx \\ = \frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2-4ac}{2c}\right) C\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx+b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2-4ac}{2c}\right) + 2cx}{4c}$$

[In] `integrate(cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out] $\frac{1}{4}(\pi*\sqrt{c/\pi}*\cos(-1/2*(b^2 - 4*a*c)/c)*\text{fresnel_cos}((2*c*x + b)*\sqrt{c/\pi}/c) - \pi*\sqrt{c/\pi}*\text{fresnel_sin}((2*c*x + b)*\sqrt{c/\pi}/c)*\sin(-1/2*(b^2 - 4*a*c)/c) + 2*c*x)/c$

Sympy [A] (verification not implemented)

Time = 0.47 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.83

$$\int \cos^2(a + bx + cx^2) dx \\ = \frac{x}{2} + \frac{\sqrt{\pi} \left(-\sin\left(2a - \frac{b^2}{2c}\right) S\left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}}\right) + \cos\left(2a - \frac{b^2}{2c}\right) C\left(\frac{2b+4cx}{2\sqrt{\pi}\sqrt{c}}\right) \right) \sqrt{\frac{1}{c}}}{4}$$

[In] `integrate(cos(c*x**2+b*x+a)**2,x)`

[Out] $x/2 + \sqrt{\pi} * (-\sin(2a - b^2/(2*c)) * \text{fresnels}((2*b + 4*c*x)/(2*sqrt(pi)*sqrt(c))) + \cos(2a - b^2/(2*c)) * \text{fresnelc}((2*b + 4*c*x)/(2*sqrt(pi)*sqrt(c)))) * \sqrt{1/c}) / 4$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx + cx^2) dx = \\ -\frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(\left((i-1) \cos\left(-\frac{b^2-4ac}{2c}\right) + (i+1) \sin\left(-\frac{b^2-4ac}{2c}\right) \right) \operatorname{erf}\left(\frac{2icx+ib}{\sqrt{2ic}}\right) + \left((i+1) \cos\left(-\frac{b^2-4ac}{2c}\right) + (i-1) \sin\left(-\frac{b^2-4ac}{2c}\right) \right) \operatorname{erfc}\left(\frac{2icx+ib}{\sqrt{2ic}}\right) \right)}{32c^2}$$

[In] `integrate(cos(c*x^2+b*x+a)^2,x, algorithm="maxima")`

[Out] $-1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*c os(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erfc((2*I*c*x + I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)/c^2$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.22

$$\int \cos^2(a + bx + cx^2) dx = \frac{1}{2}x + \frac{i\sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2}i\sqrt{c}(2x + \frac{b}{c})\left(\frac{ic}{|c|} + 1\right)\right) e^{-\frac{ib^2-4iac}{2c}}}{8\sqrt{c}\left(\frac{ic}{|c|} + 1\right)} \\ - \frac{i\sqrt{\pi} \operatorname{erf}\left(\frac{1}{2}i\sqrt{c}(2x + \frac{b}{c})\left(-\frac{ic}{|c|} + 1\right)\right) e^{-\frac{-ib^2+4iac}{2c}}}{8\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)}$$

[In] `integrate(cos(c*x^2+b*x+a)^2,x, algorithm="giac")`
[Out] $\frac{1}{2}x + \frac{1}{8}\sqrt{\pi}\operatorname{erf}\left(\frac{-\sqrt{c}(2x + b/c)}{\sqrt{4a + b^2}}\right) + \frac{1}{8}\sqrt{\pi}\operatorname{erf}\left(\frac{\sqrt{c}(2x + b/c)}{\sqrt{4a + b^2}}\right)$

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 dx$$

[In] `int(cos(a + b*x + c*x^2)^2,x)`
[Out] `int(cos(a + b*x + c*x^2)^2, x)`

3.19 $\int \frac{\cos^2(a+bx+cx^2)}{x} dx$

Optimal result	129
Rubi [N/A]	129
Mathematica [N/A]	130
Maple [N/A] (verified)	130
Fricas [N/A]	130
Sympy [N/A]	130
Maxima [N/A]	131
Giac [N/A]	131
Mupad [N/A]	131

Optimal result

Integrand size = 17, antiderivative size = 17

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{x}, x\right)$$

[Out] $1/2*\ln(x)+1/2*\text{Unintegrable}(\cos(2*c*x^2+2*b*x+2*a)/x,x)$

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos^2(a + bx + cx^2)}{x} dx$$

[In] $\text{Int}[\text{Cos}[a + b*x + c*x^2]^2/x, x]$

[Out] $\text{Log}[x]/2 + \text{Defer}[\text{Int}][\text{Cos}[2*a + 2*b*x + 2*c*x^2]/x, x]/2$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2x} + \frac{\cos(2a + 2bx + 2cx^2)}{2x} \right) dx \\ &= \frac{\log(x)}{2} + \frac{1}{2} \int \frac{\cos(2a + 2bx + 2cx^2)}{x} dx \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 3.96 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos^2(a + bx + cx^2)}{x} dx$$

[In] `Integrate[Cos[a + b*x + c*x^2]^2/x, x]`

[Out] `Integrate[Cos[a + b*x + c*x^2]^2/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.43 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(cx^2 + bx + a)}{x} dx$$

[In] `int(cos(c*x^2+b*x+a)^2/x,x)`

[Out] `int(cos(c*x^2+b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 + b*x + a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.80 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos^2(a + bx + cx^2)}{x} dx$$

[In] `integrate(cos(c*x**2+b*x+a)**2/x,x)`

[Out] `Integral(cos(a + b*x + c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.65

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="maxima")`

[Out] `1/2*integrate(cos(2*c*x^2 + 2*b*x + 2*a)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/x,x, algorithm="giac")`

[Out] `integrate(cos(c*x^2 + b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 13.75 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{\cos^2(a + bx + cx^2)}{x} dx = \int \frac{\cos(cx^2 + bx + a)^2}{x} dx$$

[In] `int(cos(a + b*x + c*x^2)^2/x,x)`

[Out] `int(cos(a + b*x + c*x^2)^2/x, x)`

3.20 $\int x^2 \cos^2(a + bx - cx^2) dx$

Optimal result	132
Rubi [A] (verified)	133
Mathematica [A] (verified)	135
Maple [A] (verified)	135
Fricas [A] (verification not implemented)	136
Sympy [F]	137
Maxima [C] (verification not implemented)	137
Giac [C] (verification not implemented)	138
Mupad [F(-1)]	139

Optimal result

Integrand size = 18, antiderivative size = 248

$$\begin{aligned} \int x^2 \cos^2(a + bx - cx^2) dx = & \frac{x^3}{6} - \frac{b^2 \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & + \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{3/2}} \\ & - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{5/2}} \\ & - \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

```
[Out] 1/6*x^3-1/16*b*sin(-2*c*x^2+2*b*x+2*a)/c^2-1/8*x*sin(-2*c*x^2+2*b*x+2*a)/c-
1/16*b^2*cos(2*a+1/2*b^2/c)*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/
c^(5/2)+1/16*cos(2*a+1/2*b^2/c)*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1
/2)/c^(3/2)-1/16*FresnelC((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*b^2/c)*P
i^(1/2)/c^(3/2)-1/16*b^2*FresnelS((-2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a+1/2*
b^2/c)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.389, Rules used = {3549, 3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int x^2 \cos^2(a + bx - cx^2) dx = & -\frac{\sqrt{\pi} \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & -\frac{\sqrt{\pi}b^2 \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & -\frac{\sqrt{\pi}b^2 \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ & +\frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ & -\frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} + \frac{x^3}{6} \end{aligned}$$

[In] `Int[x^2*Cos[a + b*x - c*x^2]^2,x]`

[Out] $x^{3/2} - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(5/2)}) + (\text{Sqrt}[\text{Pi}]*\text{Cos}[2*a + b^{2/(2*c)}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}]])]/(16*c^{(3/2)}) - (\text{Sqrt}[\text{Pi}]*\text{FresnelC}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(3/2)}) - (b^{2*}\text{Sqrt}[\text{Pi}]*\text{FresnelS}[(b - 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[\text{Pi}])]*)\text{Sin}[2*a + b^{2/(2*c)}]/(16*c^{(5/2)}) - (b*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(16*c^2) - (x*\text{Sin}[2*a + 2*b*x - 2*c*x^2])/(8*c)$

Rule 3432

`Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simplify[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Rule 3433

`Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simplify[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Rule 3528

`Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] - Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && Neq[b^2 - 4*a*c, 0]`

Rule 3529

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]*((d_.) + (e_ .)*(x_)), x_Symbol]
:> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]*((d_.) + (e_ .)*(x_)^(m_)), x_Sym
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[
(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - D
ist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x]
) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rule 3549

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]^(n_)*((d_.) + (e_ .)*(x_)^(m_)), x_Symbol]
:> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{x^2}{2} + \frac{1}{2}x^2 \cos(2a + 2bx - 2cx^2) \right) dx \\
&= \frac{x^3}{6} + \frac{1}{2} \int x^2 \cos(2a + 2bx - 2cx^2) dx \\
&= \frac{x^3}{6} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} + \frac{\int \sin(2a + 2bx - 2cx^2) dx}{8c} + \frac{b \int x \cos(2a + 2bx - 2cx^2) dx}{4c} \\
&= \frac{x^3}{6} - \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} \\
&\quad + \frac{b^2 \int \cos(2a + 2bx - 2cx^2) dx}{8c^2} - \frac{\cos\left(2a + \frac{b^2}{2c}\right) \int \sin\left(\frac{(2b-4cx)^2}{8c}\right) dx}{8c} \\
&\quad + \frac{\sin\left(2a + \frac{b^2}{2c}\right) \int \cos\left(\frac{(2b-4cx)^2}{8c}\right) dx}{8c}
\end{aligned}$$

$$\begin{aligned}
&= \frac{x^3}{6} + \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{3/2}} \\
&\quad - \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c} \\
&\quad + \frac{\left(b^2 \cos\left(2a + \frac{b^2}{2c}\right)\right) \int \cos\left(\frac{(2b-4cx)^2}{8c}\right) dx}{8c^2} \\
&\quad + \frac{\left(b^2 \sin\left(2a + \frac{b^2}{2c}\right)\right) \int \sin\left(\frac{(2b-4cx)^2}{8c}\right) dx}{8c^2} \\
&= \frac{x^3}{6} - \frac{b^2 \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} + \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\
&\quad - \frac{\sqrt{\pi} \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{3/2}} - \frac{b^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{16c^{5/2}} \\
&\quad - \frac{b \sin(2a + 2bx - 2cx^2)}{16c^2} - \frac{x \sin(2a + 2bx - 2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.40 (sec), antiderivative size = 175, normalized size of antiderivative = 0.71

$$\begin{aligned}
&\int x^2 \cos^2(a + bx - cx^2) dx \\
&= \frac{-3\sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(c \cos\left(2a + \frac{b^2}{2c}\right) - b^2 \sin\left(2a + \frac{b^2}{2c}\right)\right) + 3\sqrt{\pi} \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(b^2 \cos\left(2a + \frac{b^2}{2c}\right) - c \sin\left(2a + \frac{b^2}{2c}\right)\right)}{48c^{5/2}}
\end{aligned}$$

[In] Integrate[x^2*Cos[a + b*x - c*x^2]^2, x]

[Out] $\frac{(-3\sqrt{\pi} \text{FresnelS}\left[\frac{(-b + 2c)x}{\sqrt{c}\sqrt{\pi}}\right]) * (c \cos[2a + \frac{b^2}{2c}] - b^2 \sin[2a + \frac{b^2}{2c}]) + 3\sqrt{\pi} \text{FresnelC}\left[\frac{(-b + 2c)x}{\sqrt{c}\sqrt{\pi}}\right]) * (b^2 \cos[2a + \frac{b^2}{2c}] - c \sin[2a + \frac{b^2}{2c}])}{48c^{5/2}}$

Maple [A] (verified)

Time = 0.66 (sec), antiderivative size = 199, normalized size of antiderivative = 0.80

method	result
default	$\frac{x^3}{6} - \frac{x \sin(-2cx^2 + 2bx + 2a)}{8c} + \frac{b \left(-\frac{\sin(-2cx^2 + 2bx + 2a)}{4c} + \frac{b\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) C\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) S\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^2} \right)}{4c} - \frac{\sqrt{\pi} \left(\cos\left(\frac{4ac+b^2}{2c}\right) C\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{4ac+b^2}{2c}\right) S\left(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^2}$
risch	$\frac{x^3}{6} + \frac{b^2 \sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \operatorname{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c^2 \sqrt{-2ic}} + \frac{i\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \operatorname{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{32c \sqrt{-2ic}} - \frac{b^2 \sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(-\sqrt{2}\sqrt{ic}x - \frac{ib}{\sqrt{2}\sqrt{ic}}\right)}{64c^2 \sqrt{ic}}$

[In] `int(x^2*cos(-c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/6*x^3 - 1/8*x*\sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c*(-1/4*\sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/4*b/c^{(3/2)}*Pi^{(1/2)}*(\cos(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^{(1/2)}/c^{(1/2)}*(2*c*x - b)) + \sin(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^{(1/2)}/c^{(1/2)}*(2*c*x - b))) - 1/16/c^{(3/2)}*Pi^{(1/2)}*(\cos(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^{(1/2)}/c^{(1/2)}*(2*c*x - b)) - \sin(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^{(1/2)}/c^{(1/2)}*(2*c*x - b))) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec), antiderivative size = 187, normalized size of antiderivative = 0.75

$$\begin{aligned} & \int x^2 \cos^2(a + bx - cx^2) dx \\ & = \frac{8c^3x^3 + 6(2c^2x + bc)\cos(cx^2 - bx - a)\sin(cx^2 - bx - a) + 3\left(\pi b^2 \cos\left(\frac{b^2 + 4ac}{2c}\right) + \pi c \sin\left(\frac{b^2 + 4ac}{2c}\right)\right)\sqrt{\frac{c}{\pi}} C}{48c^3} \end{aligned}$$

[In] `integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/48*(8*c^3*x^3 + 6*(2*c^2*x + b*c)*\cos(c*x^2 - b*x - a)*\sin(c*x^2 - b*x - a) + 3*(\pi*b^2*\cos(1/2*(b^2 + 4*a*c)/c) + \pi*c*\sin(1/2*(b^2 + 4*a*c)/c))*\sqrt{c/\pi}*\operatorname{fresnel_cos}((2*c*x - b)*\sqrt{c/\pi}/c) + 3*(\pi*b^2*\sin(1/2*(b^2 + 4*a*c)/c) - \pi*c*\cos(1/2*(b^2 + 4*a*c)/c))*\sqrt{c/\pi}*\operatorname{fresnel_sin}((2*c*x - b)*\sqrt{c/\pi}/c))/c^3 \end{aligned}$$

Sympy [F]

$$\int x^2 \cos^2(a + bx - cx^2) dx = \int x^2 \cos^2(a + bx - cx^2) dx$$

```
[In] integrate(x**2*cos(-c*x**2+b*x+a)**2,x)
```

[Out] $\text{Integral}(x^{**2} \cos(a + b*x - c*x^{**2})^{**2}, x)$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.05 (sec) , antiderivative size = 1617, normalized size of antiderivative = 6.52

$$\int x^2 \cos^2(a + bx - cx^2) dx = \text{Too large to display}$$

```
[In] integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")
```

```
[Out] -1/384*sqrt(2)*(24*((((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*cos(1/2*(b^2 + 4*a*c)/c) + ((-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*sin(1/2*(b^2 + 4*a*c)/c))*x^3 + 36*(((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(1/2*(b^2 + 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b*c^3)*sin(1/2*(b^2 + 4*a*c)/c))*x^2 - 4*sqrt(2)*(8*c^4*x^3 - 3*b*c^2*(I*e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - I*e^(-1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*cos(1/2*(b^2 + 4*a*c)/c) - 3*b*c^2*(e^(1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + e^(-1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*sin(1/2*(b^2 + 4*a*c)/c))*(4*c^2*x^2 - 4*b*c*x + b^2)/c)^(3/2) + 18*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*sin(1/2*(b^2 + 4*a*c)/c)))*((4*c^2*x^2 - 4*b*c*x + b^2)/c)^(3/2) + 18*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*c^4)*cos(1/2*(b^2 + 4*a*c)/c)))*((4*c^2*x^2 - 4*b*c*x + b^2)/c)^(3/2)
```

$$\begin{aligned}
& 2*\sqrt{(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c}) - 1) - (I + 1)*\sqrt{2}*\sqrt{p} \\
& i)*(\operatorname{erf}(\sqrt{1/2})*\sqrt{-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c}) - 1))*b^4*c + \\
& 2*(-(I + 1)*\sqrt{2}*\operatorname{gamma}(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + \\
& (I - 1)*\sqrt{2}*\operatorname{gamma}(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^2*c^2 \\
& ^2)*\cos(1/2*(b^2 + 4*a*c)/c) + ((-(I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1}) + (I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1})*b^4*c + 2*(\\
& -(I - 1)*\sqrt{2}*\operatorname{gamma}(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) + (I + 1)*\sqrt{2}*\operatorname{gamma}(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^2*c^2)* \\
& \sin(1/2*(b^2 + 4*a*c)/c)*x + 3*((-(I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1}) + (I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1})*b^5 + 2*((\\
& I + 1)*\sqrt{2}*\operatorname{gamma}(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I - 1)*\sqrt{2}*\operatorname{gamma}(3/2, -1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c))*b^3*c)*\cos(\\
& 1/2*(b^2 + 4*a*c)/c) + 3*((I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1}) - (I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1})*b^5 + 2*((I - 1)*\sqrt{2}*\operatorname{gamma}(3/2, 1/2*(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c) - (I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(\sqrt{1/2})*\sqrt{((4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)) - 1})*b^3*c)*\sin(1/2*(b^2 + 4*a*c)/c))/(c^4*((4*c^2*x^2 - 4*b*c*x + b^2)/c)^{3/2}))
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.87

$$\begin{aligned}
& \int x^2 \cos^2(a + bx - cx^2) dx \\
& = \frac{1}{6} x^3 - \frac{(c(2ix - \frac{ib}{c}) + 2ib)e^{(2icx^2 - 2ibx - 2ia)} - \frac{i\sqrt{\pi}(b^2 + ic)\operatorname{erf}\left(-\frac{1}{2}i\sqrt{c}\left(2x - \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 + 4icac}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{32c^2} \\
& - \frac{(c(-2ix + \frac{ib}{c}) - 2ib)e^{(-2icx^2 + 2ibx + 2ia)} + \frac{i\sqrt{\pi}(b^2 - ic)\operatorname{erf}\left(\frac{1}{2}i\sqrt{c}\left(2x - \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 - 4icac}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)}}{32c^2}
\end{aligned}$$

[In] `integrate(x^2*cos(-c*x^2+b*x+a)^2,x, algorithm="giac")`

[Out]

$$\begin{aligned}
& 1/6*x^3 - 1/32*((c*(2*I*x - I*b/c) + 2*I*b)*e^{(2*I*c*x^2 - 2*I*b*x - 2*I*a)} \\
& - I*\sqrt{\pi}*(b^2 + I*c)*\operatorname{erf}(-1/2*I*\sqrt{c}*(2*x - b/c)*(I*c/abs(c) + 1))* \\
& e^{(-1/2*(I*b^2 + 4*I*a*c)/c)}/(\sqrt{c}*(I*c/abs(c) + 1))/c^2 - 1/32*((c*(-2 \\
& *I*x + I*b/c) - 2*I*b)*e^{(-2*I*c*x^2 + 2*I*b*x + 2*I*a)} + I*\sqrt{\pi}*(b^2 - \\
& I*c)*\operatorname{erf}(1/2*I*\sqrt{c}*(2*x - b/c)*(-I*c/abs(c) + 1))*e^{(-1/2*(-I*b^2 - 4*I*a*c)/c)}/(\sqrt{c}*(-I*c/abs(c) + 1))/c^2
\end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2(a + bx - cx^2) dx = \int x^2 \cos(-cx^2 + bx + a)^2 dx$$

[In] int(x^2*cos(a + b*x - c*x^2)^2,x)

[Out] int(x^2*cos(a + b*x - c*x^2)^2, x)

3.21 $\int x \cos^2(a + bx - cx^2) dx$

Optimal result	140
Rubi [A] (verified)	140
Mathematica [A] (verified)	142
Maple [A] (verified)	142
Fricas [A] (verification not implemented)	143
Sympy [F]	143
Maxima [C] (verification not implemented)	143
Giac [C] (verification not implemented)	144
Mupad [F(-1)]	145

Optimal result

Integrand size = 16, antiderivative size = 126

$$\begin{aligned} \int x \cos^2(a + bx - cx^2) dx = & \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & - \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{8c^{3/2}} \\ & - \frac{\sin(2a + 2bx - 2cx^2)}{8c} \end{aligned}$$

[Out] $1/4*x^2 - 1/8*\sin(-2*c*x^2 + 2*b*x + 2*a)/c - 1/8*b*\cos(2*a + 1/2*b^2/c)*\text{FresnelC}((-2*c*x + b)/c^{1/2}/\text{Pi}^{1/2})*\text{Pi}^{1/2}/c^{3/2} - 1/8*b*\text{FresnelS}((-2*c*x + b)/c^{1/2}/\text{Pi}^{1/2})*\sin(2*a + 1/2*b^2/c)*\text{Pi}^{1/2}/c^{3/2}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3549, 3543, 3529, 3433, 3432}

$$\begin{aligned} \int x \cos^2(a + bx - cx^2) dx = & -\frac{\sqrt{\pi}b \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & - \frac{\sqrt{\pi}b \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & - \frac{\sin(2a + 2bx - 2cx^2)}{8c} + \frac{x^2}{4} \end{aligned}$$

[In] $\text{Int}[x*\text{Cos}[a + b*x - c*x^2]^2, x]$

[Out] $x^{2/4} - (b\sqrt{\pi})\cos[2a + b^2/(2c)]\text{FresnelC}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})]/(8c^{3/2}) - (b\sqrt{\pi})\text{FresnelS}[(b - 2cx)/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)]/(8c^{3/2}) - \sin[2a + 2bx - 2cx^2]/(8c)$

Rule 3432

$\text{Int}[\sin[(d_.)\cdot(e_.) + (f_.)\cdot(x_.)^2], x] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f\sqrt{d}), \text{FresnelS}[\sqrt{2/\pi}\sqrt{d}\cdot(e + f\cdot x)], x] /; \text{FreeQ}[d, e, f, x]$

Rule 3433

$\text{Int}[\cos[(d_.)\cdot(e_.) + (f_.)\cdot(x_.)^2], x] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f\sqrt{d}), \text{FresnelC}[\sqrt{2/\pi}\sqrt{d}\cdot(e + f\cdot x)], x] /; \text{FreeQ}[d, e, f, x]$

Rule 3529

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2], x] \rightarrow \text{Dist}[\cos[b^2 - 4a\cdot c]/(4c), \text{Int}[\cos[(b + 2cx)^2/(4c)], x], x] + \text{Dist}[\sin[b^2 - 4a\cdot c]/(4c), \text{Int}[\sin[(b + 2cx)^2/(4c)], x], x] /; \text{FreeQ}[a, b, c, x] \&& \text{NeQ}[b^2 - 4a\cdot c, 0]$

Rule 3543

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2]\cdot((d_.) + (e_.)\cdot(x_)), x] \rightarrow \text{Simp}[e\cdot(\sin[a + bx + cx^2]/(2c)), x] + \text{Dist}[(2c\cdot d - b\cdot e)/(2c), \text{Int}[\cos[a + bx + cx^2], x], x] /; \text{FreeQ}[a, b, c, d, e, x] \&& \text{NeQ}[2c\cdot d - b\cdot e, 0]$

Rule 3549

$\text{Int}[\cos[(a_.) + (b_.)\cdot(x_.) + (c_.)\cdot(x_.)^2]\cdot(n_*)\cdot((d_.) + (e_.)\cdot(x_.)^m), x] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e\cdot x)^m, \cos[a + bx + cx^2]^n], x] /; \text{FreeQ}[a, b, c, d, e, m, x] \&& \text{IGtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{x}{2} + \frac{1}{2}x \cos(2a + 2bx - 2cx^2) \right) dx \\ &= \frac{x^2}{4} + \frac{1}{2} \int x \cos(2a + 2bx - 2cx^2) dx \\ &= \frac{x^2}{4} - \frac{\sin(2a + 2bx - 2cx^2)}{8c} + \frac{b \int \cos(2a + 2bx - 2cx^2) dx}{4c} \\ &= \frac{x^2}{4} - \frac{\sin(2a + 2bx - 2cx^2)}{8c} + \frac{\left(b \cos\left(2a + \frac{b^2}{2c}\right) \right) \int \cos\left(\frac{(2b-4cx)^2}{8c}\right) dx}{4c} \\ &\quad + \frac{\left(b \sin\left(2a + \frac{b^2}{2c}\right) \right) \int \sin\left(\frac{(2b-4cx)^2}{8c}\right) dx}{4c} \end{aligned}$$

$$\begin{aligned}
&= \frac{x^2}{4} - \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\
&\quad - \frac{b\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{8c^{3/2}} - \frac{\sin(2a + 2bx - 2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec), antiderivative size = 122, normalized size of antiderivative = 0.97

$$\begin{aligned}
&\int x \cos^2(a + bx - cx^2) dx \\
&= \frac{b\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + b\sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right) + \sqrt{c}(2cx^2 - \sin(2(a + x(b - cx^2))))}{8c^{3/2}}
\end{aligned}$$

[In] `Integrate[x*Cos[a + b*x - c*x^2]^2, x]`

[Out] `(b*.Sqrt[Pi]*Cos[2*a + b^2/(2*c)]*FresnelC[(-b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])] + b*.Sqrt[Pi]*FresnelS[(-b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a + b^2/(2*c)] + Sqrt[c]*(2*c*x^2 - Sin[2*(a + x*(b - c*x))]))/(8*c^(3/2))`

Maple [A] (verified)

Time = 0.67 (sec), antiderivative size = 99, normalized size of antiderivative = 0.79

method	result	size
default	$\frac{x^2}{4} - \frac{\sin(-2cx^2 + 2bx + 2a)}{8c} + \frac{b\sqrt{\pi} (\cos(\frac{4ac+b^2}{2c}) C(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}) + \sin(\frac{4ac+b^2}{2c}) S(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}))}{8c^{3/2}}$	99
risch	$\frac{x^2}{4} + \frac{b\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \operatorname{erf}\left(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}}\right)}{16c\sqrt{-2ic}} - \frac{b\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \operatorname{erf}\left(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)}{32c\sqrt{ic}} - \frac{\sin(-2cx^2 + 2bx + 2a)}{8c}$	137

[In] `int(x*cos(-c*x^2+b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out] `1/4*x^2 - 1/8*sin(-2*c*x^2 + 2*b*x + 2*a)/c + 1/8*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(4*a*c + b^2)/c)*FresnelC(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)) + sin(1/2*(4*a*c + b^2)/c)*FresnelS(1/Pi^(1/2)/c^(1/2)*(2*c*x - b)))`

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.06

$$\int x \cos^2(a + bx - cx^2) dx = \frac{\pi b \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi b \sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) + 2c^2x^2 + 2c \cos(cx^2 - bx - a)}{8c^2}$$

[In] `integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out] $\frac{1}{8} \cdot \frac{\pi b \sqrt{c/\pi} \cos(1/2*(b^2 + 4*a*c)/c) \cdot \text{fresnel_cos}(2*c*x - b) \cdot \sqrt{c/\pi}/c + \pi b \sqrt{c/\pi} \cdot \text{fresnel_sin}(2*c*x - b) \cdot \sqrt{c/\pi}/c \cdot \sin(1/2*(b^2 + 4*a*c)/c) + 2*c^2*x^2 + 2*c \cdot \cos(c*x^2 - b*x - a) \cdot \sin(c*x^2 - b*x - a)}{c^2}$

Sympy [F]

$$\int x \cos^2(a + bx - cx^2) dx = \int x \cos^2(a + bx - cx^2) dx$$

[In] `integrate(x*cos(-c*x**2+b*x+a)**2,x)`

[Out] `Integral(x*cos(a + b*x - c*x**2)**2, x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.70 (sec) , antiderivative size = 611, normalized size of antiderivative = 4.85

$$\int x \cos^2(a + bx - cx^2) dx = \frac{\sqrt{2} \left(\left((i-1) \sqrt{2} \sqrt{\pi} \left(\text{erf} \left(\sqrt{\frac{1}{2}} \sqrt{\frac{4ic^2x^2 - 4ibcx + ib^2}{c}} \right) - 1 \right) - (i+1) \sqrt{2} \sqrt{\pi} \left(\text{erf} \left(\sqrt{\frac{1}{2}} \sqrt{\frac{-4ic^2x^2 - 4ibcx + ib^2}{c}} \right) - 1 \right) \right)}{64}$$

[In] `integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")`

[Out] $\frac{1}{64} \cdot \frac{\sqrt{2} \cdot (((I-1)*\sqrt{2}*\sqrt{\pi}) * (\text{erf}(\sqrt{1/2})*\sqrt{(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c}) - 1) - (I+1)*\sqrt{2}*\sqrt{\pi} * (\text{erf}(\sqrt{1/2})*\sqrt{(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)}) - 1) * b^2 * \cos(1/2*(b^2 + 4*a*c)/c) +$

$$\begin{aligned}
& - (I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c}) - 1) + (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)} - 1)) * b^2 * \sin(1/2 * (b^2 + 4*a*c)/c) - 2 * (((I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c})) - 1) - (I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)} - 1)) * b*c*cos(1/2 * (b^2 + 4*a*c)/c) + (-(I + 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c})) - 1) + (I - 1) * \sqrt{2} * \sqrt{\pi} * (\operatorname{erf}(\sqrt{1/2}) * \sqrt{(-(4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)} - 1)) * b*c*sin(1/2 * (b^2 + 4*a*c)/c)) * x + 2 * \sqrt{2} * (4*c^2*x^2 - c * (I * e^{(1/2 * (4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)} - I * e^{-(-1/2 * (4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)}) * \cos(1/2 * (b^2 + 4*a*c)/c) - c * (e^{(1/2 * (4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)} + e^{-(-1/2 * (4*I*c^2*x^2 - 4*I*b*c*x + I*b^2)/c)}) * \sin(1/2 * (b^2 + 4*a*c)/c)) * \sqrt{(4*c^2*x^2 - 4*b*c*x + b^2)/c}) / (c^2 * \sqrt{(4*c^2*x^2 - 4*b*c*x + b^2)/c})
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.38

$$\int x \cos^2(a + bx - cx^2) dx$$

$$= \frac{1}{4} x^2 - \frac{-\frac{i \sqrt{\pi} b \operatorname{erf}\left(-\frac{1}{2} i \sqrt{c} \left(2x - \frac{b}{c}\right) \left(\frac{ic}{|c|} + 1\right)\right)}{\sqrt{c} \left(\frac{ic}{|c|} + 1\right)} e^{\left(-\frac{ib^2 + 4icac}{2c}\right)}}{16c} + i e^{(2icx^2 - 2ibx - 2ia)}$$

$$- \frac{i \sqrt{\pi} b \operatorname{erf}\left(\frac{1}{2} i \sqrt{c} \left(2x - \frac{b}{c}\right) \left(-\frac{ic}{|c|} + 1\right)\right)}{\sqrt{c} \left(-\frac{ic}{|c|} + 1\right)} e^{\left(-\frac{-ib^2 - 4icac}{2c}\right)} - i e^{(-2icx^2 + 2ibx + 2ia)}$$

$$16c$$

```
[In] integrate(x*cos(-c*x^2+b*x+a)^2,x, algorithm="giac")
```

```
[Out] 1/4*x^2 - 1/16*(-I*sqrt(pi)*b*erf(-1/2*I*sqrt(c)*(2*x - b/c)*(I*c/abs(c) + 1))*e^(-1/2*(I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)) + I*e^(2*I*c*x^2 - 2*I*b*x - 2*I*a)/c - 1/16*(I*sqrt(pi)*b*erf(1/2*I*sqrt(c)*(2*x - b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)) - I*e^(-2*I*c*x^2 + 2*I*b*x + 2*I*a))/c
```

Mupad [F(-1)]

Timed out.

$$\int x \cos^2(a + bx - cx^2) dx = \int x \cos(-cx^2 + bx + a)^2 dx$$

[In] int(x*cos(a + b*x - c*x^2)^2, x)

[Out] int(x*cos(a + b*x - c*x^2)^2, x)

3.22 $\int \cos^2(a + bx - cx^2) dx$

Optimal result	146
Rubi [A] (verified)	146
Mathematica [A] (verified)	147
Maple [A] (verified)	148
Fricas [A] (verification not implemented)	148
Sympy [A] (verification not implemented)	149
Maxima [C] (verification not implemented)	149
Giac [C] (verification not implemented)	149
Mupad [F(-1)]	150

Optimal result

Integrand size = 14, antiderivative size = 100

$$\int \cos^2(a + bx - cx^2) dx = \frac{x}{2} - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}}$$

[Out] $\frac{1}{2}x - \frac{1}{4}\cos(2a + \frac{b^2}{2c})\text{FresnelC}((-\frac{2cx+b}{\sqrt{c}})/\sqrt{\pi})\text{Pi}^{(1/2)} + \frac{1}{4}\text{FresnelS}((-\frac{2cx+b}{\sqrt{c}})/\sqrt{\pi})\text{Pi}^{(1/2)}\sin(2a + \frac{b^2}{2c})\text{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {3531, 3529, 3433, 3432}

$$\int \cos^2(a + bx - cx^2) dx = -\frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} \\ - \frac{\sqrt{\pi} \sin\left(2a + \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} + \frac{x}{2}$$

[In] Int[Cos[a + b*x - c*x^2]^2, x]

[Out] $\frac{x}{2} - \frac{(\text{Sqrt}[\text{Pi}] \cos(2a + \frac{b^2}{2c}) \text{FresnelC}((b - 2cx)/(\text{Sqrt}[c]\text{Sqrt}[\text{Pi}]))/(4\text{Sqrt}[c]) - (\text{Sqrt}[\text{Pi}] \text{FresnelS}((b - 2cx)/(\text{Sqrt}[c]\text{Sqrt}[\text{Pi}]))\text{Sin}(2a + \frac{b^2}{2c}))/(4\text{Sqrt}[c])}{4}$

Rule 3432

```
Int[Sin[(d_)*(e_) + (f_)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelS[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[Cos[(d_)*(e_) + (f_)*(x_)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3529

```
Int[Cos[(a_)*(x_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sim[(b^2 - 4*a*c)/(4*c)], Int[Sim[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3531

```
Int[Cos[(a_)*(x_) + (b_)*(x_) + (c_)*(x_)^2]^n, x_Symbol] :> Int[ExpandTri gReduce[Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2} + \frac{1}{2} \cos(2a + 2bx - 2cx^2) \right) dx \\ &= \frac{x}{2} + \frac{1}{2} \int \cos(2a + 2bx - 2cx^2) dx \\ &= \frac{x}{2} + \frac{1}{2} \cos\left(2a + \frac{b^2}{2c}\right) \int \cos\left(\frac{(2b - 4cx)^2}{8c}\right) dx + \frac{1}{2} \sin\left(2a + \frac{b^2}{2c}\right) \int \sin\left(\frac{(2b - 4cx)^2}{8c}\right) dx \\ &= \frac{x}{2} - \frac{\sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right)}{4\sqrt{c}} - \frac{\sqrt{\pi} \text{FresnelS}\left(\frac{b-2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec), antiderivative size = 100, normalized size of antiderivative = 1.00

$$\begin{aligned} &\int \cos^2(a + bx - cx^2) dx \\ &= \frac{2\sqrt{c}x + \sqrt{\pi} \cos\left(2a + \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) + \sqrt{\pi} \text{FresnelS}\left(\frac{-b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a + \frac{b^2}{2c}\right)}{4\sqrt{c}} \end{aligned}$$

[In] `Integrate[Cos[a + b*x - c*x^2]^2, x]`

[Out] $(2\sqrt{c}x + \sqrt{\pi}\cos[2a + b^2/(2c)]\text{FresnelC}[-b + 2cx]/(\sqrt{c}\sqrt{\pi}) + \sqrt{\pi}\text{FresnelS}[-b + 2cx]/(\sqrt{c}\sqrt{\pi})]\sin[2a + b^2/(2c)])/(4\sqrt{c})$

Maple [A] (verified)

Time = 0.68 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x}{2} + \frac{\sqrt{\pi} (\cos(\frac{4ac+b^2}{2c}) C(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}) + \sin(\frac{4ac+b^2}{2c}) S(\frac{2cx-b}{\sqrt{\pi}\sqrt{c}}))}{4\sqrt{c}}$	76
risch	$\frac{x}{2} + \frac{\sqrt{\pi} e^{-\frac{i(4ac+b^2)}{2c}} \operatorname{erf}(\sqrt{-2ic}x + \frac{ib}{\sqrt{-2ic}})}{8\sqrt{-2ic}} - \frac{\sqrt{\pi} e^{\frac{i(4ac+b^2)}{2c}} \sqrt{2} \operatorname{erf}(-\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}})}{16\sqrt{ic}}$	107

[In] `int(cos(-c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $1/2*x + 1/4*\pi^{(1/2)}/c^{(1/2)}*(\cos(1/2*(4*a*c+b^2)/c)*\text{FresnelC}(1/\pi^{(1/2)}/c^{(1/2)}*(2*c*x-b)) + \sin(1/2*(4*a*c+b^2)/c)*\text{FresnelS}(1/\pi^{(1/2)}/c^{(1/2)}*(2*c*x-b)))$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.96

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{\pi\sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2+4ac}{2c}\right) C\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) + \pi\sqrt{\frac{c}{\pi}} S\left(\frac{(2cx-b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(\frac{b^2+4ac}{2c}\right) + 2cx}{4c}$$

[In] `integrate(cos(-c*x^2+b*x+a)^2,x, algorithm="fricas")`

[Out] $1/4*(\pi*\sqrt{c/\pi}*\cos(1/2*(b^2 + 4*a*c)/c)*\text{fresnel_cos}((2*c*x - b)*\sqrt{c/\pi}/c) + \pi*\sqrt{c/\pi}*\text{fresnel_sin}((2*c*x - b)*\sqrt{c/\pi}/c)*\sin(1/2*(b^2 + 4*a*c)/c) + 2*c*x)/c$

Sympy [A] (verification not implemented)

Time = 0.45 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.88

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{x}{2} + \frac{\sqrt{\pi} \sqrt{-\frac{1}{c}} \left(-\sin\left(2a + \frac{b^2}{2c}\right) S\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) + \cos\left(2a + \frac{b^2}{2c}\right) C\left(\frac{2b-4cx}{2\sqrt{\pi}\sqrt{-c}}\right) \right)}{4}$$

[In] `integrate(cos(-c*x**2+b*x+a)**2,x)`

[Out] $x/2 + \sqrt{\pi} \sqrt{-1/c} (-\sin(2a + b^2/(2*c)) * \text{fresnels}((2*b - 4*c*x)/(2 * \sqrt{\pi} * \sqrt{-c})) + \cos(2*a + b^2/(2*c)) * \text{fresnelc}((2*b - 4*c*x)/(2 * \sqrt{\pi} * \sqrt{-c}))) / 4$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx - cx^2) dx \\ = \frac{4^{1/4} \sqrt{2} \sqrt{\pi} \left(\left(-(i-1) \cos\left(\frac{b^2+4ac}{2c}\right) + (i+1) \sin\left(\frac{b^2+4ac}{2c}\right) \right) \operatorname{erf}\left(\frac{2icx-i b}{\sqrt{2} i c}\right) + \left(-(i+1) \cos\left(\frac{b^2+4ac}{2c}\right) + (i-1) \sin\left(\frac{b^2+4ac}{2c}\right) \right) \operatorname{erf}\left(\frac{2icx+i b}{\sqrt{2} i c}\right) \right)}{32 c^2}$$

[In] `integrate(cos(-c*x^2+b*x+a)^2,x, algorithm="maxima")`

[Out] $\frac{1}{32} (4^{(1/4)} \sqrt{2} \sqrt{\pi} ((-(I-1) \cos(1/2*(b^2 + 4*a*c)/c) + (I+1) \sin(1/2*(b^2 + 4*a*c)/c)) * \operatorname{erf}((2*I*c*x - I*b)/\sqrt{2*I*c}) + (-(I+1) \cos(1/2*(b^2 + 4*a*c)/c) + (I-1) \sin(1/2*(b^2 + 4*a*c)/c)) * \operatorname{erf}((2*I*c*x - I*b)/\sqrt{-2*I*c})) * c^{(3/2)} + 16*c^2*x)/c^2$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.44 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.24

$$\int \cos^2(a + bx - cx^2) dx = \frac{1}{2} x + \frac{i \sqrt{\pi} \operatorname{erf}\left(-\frac{1}{2} i \sqrt{c} (2x - \frac{b}{c}) \left(\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{ib^2+4icac}{2c}\right)}}{8 \sqrt{c} \left(\frac{ic}{|c|} + 1\right)} \\ - \frac{i \sqrt{\pi} \operatorname{erf}\left(\frac{1}{2} i \sqrt{c} (2x - \frac{b}{c}) \left(-\frac{ic}{|c|} + 1\right)\right) e^{\left(-\frac{-ib^2-4icac}{2c}\right)}}{8 \sqrt{c} \left(-\frac{ic}{|c|} + 1\right)}$$

```
[In] integrate(cos(-c*x^2+b*x+a)^2,x, algorithm="giac")
[Out] 
$$\frac{1}{2}x + \frac{1}{8}\sqrt{\pi}\operatorname{erf}\left(\frac{-\sqrt{c}(2x - b/c)}{\sqrt{4a + b^2}}\right) + C$$

```

Mupad [F(-1)]

Timed out.

$$\int \cos^2(a + bx - cx^2) dx = \int \cos(-cx^2 + bx + a)^2 dx$$

```
[In] int(cos(a + b*x - c*x^2)^2,x)
[Out] int(cos(a + b*x - c*x^2)^2, x)
```

3.23 $\int \frac{\cos^2(a+bx-cx^2)}{x} dx$

Optimal result	151
Rubi [N/A]	151
Mathematica [N/A]	152
Maple [N/A] (verified)	152
Fricas [N/A]	152
Sympy [N/A]	152
Maxima [N/A]	153
Giac [N/A]	153
Mupad [N/A]	153

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx - 2cx^2)}{x}, x\right)$$

[Out] $1/2*\ln(x)+1/2*\text{Unintegrable}(\cos(-2*c*x^2+2*b*x+2*a)/x,x)$

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos^2(a + bx - cx^2)}{x} dx$$

[In] $\text{Int}[\text{Cos}[a + b*x - c*x^2]^2/x, x]$

[Out] $\text{Log}[x]/2 + \text{Defer}[\text{Int}][\text{Cos}[2*a + 2*b*x - 2*c*x^2]/x, x]/2$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2x} + \frac{\cos(2a + 2bx - 2cx^2)}{2x} \right) dx \\ &= \frac{\log(x)}{2} + \frac{1}{2} \int \frac{\cos(2a + 2bx - 2cx^2)}{x} dx \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 4.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos^2(a + bx - cx^2)}{x} dx$$

[In] `Integrate[Cos[a + b*x - c*x^2]^2/x, x]`

[Out] `Integrate[Cos[a + b*x - c*x^2]^2/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.39 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(-cx^2 + bx + a)}{x} dx$$

[In] `int(cos(-c*x^2+b*x+a)^2/x, x)`

[Out] `int(cos(-c*x^2+b*x+a)^2/x, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

[In] `integrate(cos(-c*x^2+b*x+a)^2/x, x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 - b*x - a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.84 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos^2(a + bx - cx^2)}{x} dx$$

[In] `integrate(cos(-c*x**2+b*x+a)**2/x, x)`

[Out] `Integral(cos(a + b*x - c*x**2)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.56

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

```
[In] integrate(cos(-c*x^2+b*x+a)^2/x,x, algorithm="maxima")
[Out] 1/2*integrate(cos(2*c*x^2 - 2*b*x - 2*a)/x, x) + 1/2*log(x)
```

Giac [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

```
[In] integrate(cos(-c*x^2+b*x+a)^2/x,x, algorithm="giac")
[Out] integrate(cos(-c*x^2 + b*x + a)^2/x, x)
```

Mupad [N/A]

Not integrable

Time = 13.40 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{\cos^2(a + bx - cx^2)}{x} dx = \int \frac{\cos(-cx^2 + bx + a)^2}{x} dx$$

```
[In] int(cos(a + b*x - c*x^2)^2/x,x)
[Out] int(cos(a + b*x - c*x^2)^2/x, x)
```

3.24 $\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	154
Rubi [A] (verified)	154
Mathematica [A] (verified)	156
Maple [A] (verified)	156
Fricas [A] (verification not implemented)	157
Sympy [F]	157
Maxima [C] (verification not implemented)	157
Giac [C] (verification not implemented)	158
Mupad [F(-1)]	158

Optimal result

Integrand size = 15, antiderivative size = 85

$$\begin{aligned} \int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{x^3}{6} + \frac{1}{16}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS} \left(\frac{1+2x}{\sqrt{\pi}} \right) \\ & - \frac{1}{16} \sin \left(\frac{1}{2} + 2x + 2x^2 \right) + \frac{1}{8}x \sin \left(\frac{1}{2} + 2x + 2x^2 \right) \end{aligned}$$

[Out] $1/6*x^3 - 1/16*\sin(1/2+2*x+2*x^2) + 1/8*x*\sin(1/2+2*x+2*x^2) + 1/16*\operatorname{FresnelC}((1+2*x)/\operatorname{Pi}^{(1/2)})*\operatorname{Pi}^{(1/2)} - 1/16*\operatorname{FresnelS}((1+2*x)/\operatorname{Pi}^{(1/2)})*\operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3549, 3545, 3543, 3527, 3433, 3526, 3432}

$$\begin{aligned} \int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{1}{16}\sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS} \left(\frac{2x+1}{\sqrt{\pi}} \right) \\ & + \frac{x^3}{6} + \frac{1}{8}x \sin \left(2x^2 + 2x + \frac{1}{2} \right) - \frac{1}{16} \sin \left(2x^2 + 2x + \frac{1}{2} \right) \end{aligned}$$

[In] $\operatorname{Int}[x^2 \cos[1/4 + x + x^2]^2, x]$

[Out] $x^{3/6} + (\operatorname{Sqrt}[\operatorname{Pi}] * \operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[\operatorname{Pi}]])/16 - (\operatorname{Sqrt}[\operatorname{Pi}] * \operatorname{FresnelS}[(1 + 2*x)/\operatorname{Sqrt}[\operatorname{Pi}]])/16 - \operatorname{Sin}[1/2 + 2*x + 2*x^2]/16 + (x * \operatorname{Sin}[1/2 + 2*x + 2*x^2])/8$

Rule 3432

```
Int[ $\sin[(d_*) \cdot ((e_*) + (f_*) \cdot (x_*)^2)]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ )] * FresnelS[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot \text{FresnelS}[e + f \cdot x]$ , x] /; FreeQ[{d, e, f}, x]
```

Rule 3433

```
Int[ $\cos[(d_*) \cdot ((e_*) + (f_*) \cdot (x_*)^2)]$ , x_Symbol] :> Simp[( $\sqrt{\pi}/2$ ) / ( $f \cdot \sqrt{d}$ )] * FresnelC[ $\sqrt{2/\pi} \cdot \sqrt{d} \cdot \text{FresnelC}[e + f \cdot x]$ , x] /; FreeQ[{d, e, f}, x]
```

Rule 3526

```
Int[ $\sin[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2]$ , x_Symbol] :> Int[ $\sin[(b + 2c*x)^2 / (4*c)]$ , x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3527

```
Int[ $\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2]$ , x_Symbol] :> Int[ $\cos[(b + 2c*x)^2 / (4*c)]$ , x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[ $\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2] \cdot ((d_*) + (e_*) \cdot (x_*))$ , x_Symbol] :> Simp[e * (Sin[a + b*x + c*x^2] / (2*c)), x] + Dist[(2*c*d - b*e) / (2*c), Int[ $\cos[a + b*x + c*x^2]$ , x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[ $\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2] \cdot ((d_*) + (e_*) \cdot (x_*)^{(m)})$ , x_Symbol] :> Simp[e * (d + e*x)^(m - 1) * (Sin[a + b*x + c*x^2] / (2*c)), x] + (-Dist[(b*e - 2*c*d) / (2*c), Int[(d + e*x)^(m - 1) * Cos[a + b*x + c*x^2], x], x] - Dist[e^2 * ((m - 1) / (2*c)), Int[(d + e*x)^(m - 2) * Sin[a + b*x + c*x^2], x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rule 3549

```
Int[ $\cos[(a_*) + (b_*) \cdot (x_*) + (c_*) \cdot (x_*)^2]^{(n)} \cdot ((d_*) + (e_*) \cdot (x_*)^{(m)})$ , x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{x^2}{2} + \frac{1}{2}x^2 \cos\left(\frac{1}{2} + 2x + 2x^2\right) \right) dx \\ &= \frac{x^3}{6} + \frac{1}{2} \int x^2 \cos\left(\frac{1}{2} + 2x + 2x^2\right) dx \end{aligned}$$

$$\begin{aligned}
&= \frac{x^3}{6} + \frac{1}{8}x \sin\left(\frac{1}{2} + 2x + 2x^2\right) - \frac{1}{8} \int \sin\left(\frac{1}{2} + 2x + 2x^2\right) dx - \frac{1}{4} \int x \cos\left(\frac{1}{2} + 2x + 2x^2\right) dx \\
&\quad + 2x^2 \Big) dx \\
&= \frac{x^3}{6} - \frac{1}{16} \sin\left(\frac{1}{2} + 2x + 2x^2\right) + \frac{1}{8}x \sin\left(\frac{1}{2} + 2x + 2x^2\right) \\
&\quad + \frac{1}{8} \int \cos\left(\frac{1}{2} + 2x + 2x^2\right) dx - \frac{1}{8} \int \sin\left(\frac{1}{8}(2+4x)^2\right) dx \\
&= \frac{x^3}{6} - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) - \frac{1}{16} \sin\left(\frac{1}{2} + 2x + 2x^2\right) \\
&\quad + \frac{1}{8}x \sin\left(\frac{1}{2} + 2x + 2x^2\right) + \frac{1}{8} \int \cos\left(\frac{1}{8}(2+4x)^2\right) dx \\
&= \frac{x^3}{6} + \frac{1}{16}\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right) - \frac{1}{16}\sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) \\
&\quad - \frac{1}{16} \sin\left(\frac{1}{2} + 2x + 2x^2\right) + \frac{1}{8}x \sin\left(\frac{1}{2} + 2x + 2x^2\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.91

$$\int x^2 \cos^2\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{48} \left(8x^3 + 3\sqrt{\pi} \operatorname{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right) - 3\sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) \right. \\
\left. - 3 \sin\left(\frac{1}{2}(1+2x)^2\right) + 6x \sin\left(\frac{1}{2}(1+2x)^2\right) \right)$$

[In] `Integrate[x^2*Cos[1/4 + x + x^2]^2, x]`

[Out] `(8*x^3 + 3*.Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]] - 3*.Sqrt[Pi]*FresnelS[(1 + 2*x)/Sqrt[Pi]] - 3*Sin[(1 + 2*x)^2/2] + 6*x*Sin[(1 + 2*x)^2/2])/48`

Maple [A] (verified)

Time = 1.43 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.75

method	result
default	$\frac{x^3}{6} - \frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{16} + \frac{x \sin\left(\frac{1}{2} + 2x + 2x^2\right)}{8} + \frac{\operatorname{C}\left(\frac{1+2x}{\sqrt{\pi}}\right)\sqrt{\pi}}{16} - \frac{\operatorname{S}\left(\frac{1+2x}{\sqrt{\pi}}\right)\sqrt{\pi}}{16}$
risch	$\frac{x^3}{6} - \frac{\sqrt{\pi}\sqrt{2}(-1)^{\frac{3}{4}}\operatorname{erf}\left(\sqrt{2}(-1)^{\frac{1}{4}}x + \frac{\sqrt{2}(-1)^{\frac{1}{4}}}{2}\right)}{64} - \frac{(-1)^{\frac{1}{4}}\sqrt{\pi}\sqrt{2}\operatorname{erf}\left(\sqrt{2}(-1)^{\frac{1}{4}}x + \frac{\sqrt{2}(-1)^{\frac{1}{4}}}{2}\right)}{64} + \frac{\sqrt{\pi}\operatorname{erf}\left(\sqrt{-2i}x - \frac{i}{\sqrt{-2i}}\right)}{32\sqrt{-2i}}$

```
[In] int(x^2*cos(1/4+x+x^2)^2,x,method=_RETURNVERBOSE)
```

[Out] $\frac{1}{6}x^3 - \frac{1}{16}\sin(1/2 + 2x + 2x^2) + \frac{1}{8}x\sin(1/2 + 2x + 2x^2) + \frac{1}{16}\text{FresnelC}((1+2x)/\sqrt{\pi})\sqrt{\pi} - \frac{1}{16}\text{FresnelS}((1+2x)/\sqrt{\pi})\sqrt{\pi}$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 57, normalized size of antiderivative = 0.67

$$\begin{aligned} \int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = & \frac{1}{6} x^3 + \frac{1}{8} (2x - 1) \cos \left(x^2 + x + \frac{1}{4} \right) \sin \left(x^2 + x + \frac{1}{4} \right) \\ & + \frac{1}{16} \sqrt{\pi} C \left(\frac{2x + 1}{\sqrt{\pi}} \right) - \frac{1}{16} \sqrt{\pi} S \left(\frac{2x + 1}{\sqrt{\pi}} \right) \end{aligned}$$

```
[In] integrate(x^2*cos(1/4+x+x^2)^2,x, algorithm="fricas")
```

```
[Out] 1/6*x^3 + 1/8*(2*x - 1)*cos(x^2 + x + 1/4)*sin(x^2 + x + 1/4) + 1/16*sqrt(pi)*fresnel_cos((2*x + 1)/sqrt(pi)) - 1/16*sqrt(pi)*fresnel_sin((2*x + 1)/sqrt(pi))
```

Sympy [F]

$$\int x^2 \cos^2\left(\frac{1}{4} + x + x^2\right) dx = \int x^2 \cos^2\left(x^2 + x + \frac{1}{4}\right) dx$$

[In] `integrate(x**2*cos(1/4+x+x**2)**2,x)`

[Out] $\text{Integral}(x^{**2} \cos(x^{**2} + x + 1/4)^{**2}, x)$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.43 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.01

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx \\ \equiv \frac{128 x^4 + 64 x^3 - 48 x \left(-i e^{(2i x^2 + 2i x + \frac{1}{2} i)} + i e^{(-2i x^2 - 2i x - \frac{1}{2} i)} \right) - 3 \sqrt{8 x^2 + 8 x + 2} \left((i - 1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf} \left(\sqrt{2} \left(x + \frac{1}{4} \right) \right) + \operatorname{erfc} \left(\sqrt{2} \left(x + \frac{1}{4} \right) \right) \right) \right)}{16 x^4 + 8 x^3 + 2 x^2 + 2 x + 1}$$

```
[In] integrate(x^2*cos(1/4+x+x^2)^2,x, algorithm="maxima")
```

[Out] $\frac{1}{384} \left(128x^4 + 64x^3 - 48x(-I e^{(2Ix^2 + 2Ix + 1/2I)} + I e^{-(-2Ix^2 - 2Ix - 1/2I)}) - 3\sqrt{8x^2 + 8x + 2}((I - 1)\sqrt{2}\sqrt{\pi}) \right.$
 $\left. \operatorname{erf}(\sqrt{2Ix^2 + 2Ix + 1/2I}) - 1) - (I + 1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{-2Ix^2 - 2Ix - 1/2I}) - 1) - (2I + 2)\sqrt{2}\gamma(3/2, 2Ix^2 + 2Ix + 1/2I) + (2I - 2)\sqrt{2}\gamma(3/2, -2Ix^2 - 2Ix - 1/2I)) + 24Ie^{(2Ix^2 + 2Ix + 1/2I)} - 24Ie^{-(-2Ix^2 - 2Ix - 1/2I)}) / (2x + 1)$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.50 (sec), antiderivative size = 64, normalized size of antiderivative = 0.75

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{6} x^3 - \frac{1}{32} (2ix - i) e^{(2ix^2 + 2ix + \frac{1}{2}i)} - \frac{1}{32} (-2ix + i) e^{(-2ix^2 - 2ix - \frac{1}{2}i)} \\ - \frac{1}{32} i \sqrt{\pi} \operatorname{erf} \left((i - 1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ + \frac{1}{32} i \sqrt{\pi} \operatorname{erf} \left(-(i + 1)x - \frac{1}{2}i - \frac{1}{2} \right)$$

[In] `integrate(x^2*cos(1/4+x+x^2)^2, x, algorithm="giac")`

[Out] $\frac{1}{6}x^3 - \frac{1}{32}(2Ix - I)e^{(2Ix^2 + 2Ix + 1/2I)} - \frac{1}{32}(-2Ix + I)e^{(-2Ix^2 - 2Ix - 1/2I)} - \frac{1}{32}I\sqrt{\pi}\operatorname{erf}((I - 1)x + 1/2I - 1/2) + \frac{1}{32}I\sqrt{\pi}\operatorname{erf}(-(I + 1)x - 1/2I - 1/2)$

Mupad [F(-1)]

Timed out.

$$\int x^2 \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x^2 \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

[In] `int(x^2*cos(x + x^2 + 1/4)^2, x)`

[Out] `int(x^2*cos(x + x^2 + 1/4)^2, x)`

3.25 $\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	159
Rubi [A] (verified)	159
Mathematica [A] (verified)	160
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	161
Sympy [B] (verification not implemented)	161
Maxima [C] (verification not implemented)	162
Giac [C] (verification not implemented)	162
Mupad [F(-1)]	163

Optimal result

Integrand size = 13, antiderivative size = 46

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x^2}{4} - \frac{1}{8} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) + \frac{1}{8} \sin \left(\frac{1}{2} + 2x + 2x^2 \right)$$

[Out] $1/4*x^2 + 1/8*\sin(1/2 + 2*x + 2*x^2) - 1/8*\operatorname{FresnelC}((1+2*x)/\operatorname{Pi}^{(1/2)})*\operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3549, 3543, 3527, 3433}

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{1}{8} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x^2}{4} + \frac{1}{8} \sin \left(2x^2 + 2x + \frac{1}{2} \right)$$

[In] $\operatorname{Int}[x*\operatorname{Cos}[1/4 + x + x^2]^2, x]$

[Out] $x^2/4 - (\operatorname{Sqrt}[\operatorname{Pi}]*\operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[\operatorname{Pi}]])/8 + \operatorname{Sin}[1/2 + 2*x + 2*x^2]/8$

Rule 3433

```
Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3527

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.)*(x_)), x_Symbol]
 :> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[
 Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3549

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]^n*((d_.) + (e_.)*(x_))^(m_), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x],
 x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{x}{2} + \frac{1}{2}x \cos\left(\frac{1}{2} + 2x + 2x^2\right) \right) dx \\ &= \frac{x^2}{4} + \frac{1}{2} \int x \cos\left(\frac{1}{2} + 2x + 2x^2\right) dx \\ &= \frac{x^2}{4} + \frac{1}{8} \sin\left(\frac{1}{2} + 2x + 2x^2\right) - \frac{1}{4} \int \cos\left(\frac{1}{2} + 2x + 2x^2\right) dx \\ &= \frac{x^2}{4} + \frac{1}{8} \sin\left(\frac{1}{2} + 2x + 2x^2\right) - \frac{1}{4} \int \cos\left(\frac{1}{8}(2+4x)^2\right) dx \\ &= \frac{x^2}{4} - \frac{1}{8}\sqrt{\pi} \text{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right) + \frac{1}{8} \sin\left(\frac{1}{2} + 2x + 2x^2\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.89

$$\int x \cos^2\left(\frac{1}{4} + x + x^2\right) dx = \frac{1}{8} \left(2x^2 - \sqrt{\pi} \text{FresnelC}\left(\frac{1+2x}{\sqrt{\pi}}\right) + \sin\left(\frac{1}{2}(1+2x)^2\right) \right)$$

[In] `Integrate[x*Cos[1/4 + x + x^2]^2, x]`

[Out] `(2*x^2 - Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]] + Sin[(1 + 2*x)^2/2])/8`

Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.76

method	result	size
default	$\frac{x^2}{4} + \frac{\sin(\frac{1}{2}+2x+2x^2)}{8} - \frac{C\left(\frac{1+2x}{\sqrt{\pi}}\right)\sqrt{\pi}}{8}$	35
risch	$\frac{x^2}{4} + \frac{\sqrt{\pi}\sqrt{2}(-1)^{\frac{3}{4}}\operatorname{erf}\left(\sqrt{2}(-1)^{\frac{1}{4}}x+\frac{\sqrt{2}(-1)^{\frac{1}{4}}}{2}\right)}{32} - \frac{\sqrt{\pi}\operatorname{erf}\left(\sqrt{-2i}x-\frac{i}{\sqrt{-2i}}\right)}{16\sqrt{-2i}} + \frac{\sin\left(\frac{(1+2x)^2}{2}\right)}{8}$	72

[In] `int(x*cos(1/4+x+x^2)^2,x,method=_RETURNVERBOSE)`

[Out] `1/4*x^2+1/8*sin(1/2+2*x+2*x^2)-1/8*FresnelC((1+2*x)/Pi^(1/2))*Pi^(1/2)`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 37, normalized size of antiderivative = 0.80

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} x^2 + \frac{1}{4} \cos \left(x^2 + x + \frac{1}{4} \right) \sin \left(x^2 + x + \frac{1}{4} \right) - \frac{1}{8} \sqrt{\pi} C \left(\frac{2x+1}{\sqrt{\pi}} \right)$$

[In] `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="fricas")`

[Out] `1/4*x^2 + 1/4*cos(x^2 + x + 1/4)*sin(x^2 + x + 1/4) - 1/8*sqrt(pi)*fresnel_cos((2*x + 1)/sqrt(pi))`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(37) = 74$.

Time = 0.92 (sec) , antiderivative size = 121, normalized size of antiderivative = 2.63

$$\begin{aligned} \int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx &= \frac{x^2}{4} - \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{16 \Gamma \left(\frac{5}{4} \right)} + \frac{\sqrt{\pi} x C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right)}{4} \\ &\quad + \frac{\sin \left(2 \left(x + \frac{1}{2} \right)^2 \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)} - \frac{\sqrt{\pi} C \left(\frac{2x}{\sqrt{\pi}} + \frac{1}{\sqrt{\pi}} \right) \Gamma \left(\frac{1}{4} \right)}{32 \Gamma \left(\frac{5}{4} \right)} \end{aligned}$$

[In] `integrate(x*cos(1/4+x+x**2)**2,x)`

[Out] `x**2/4 - sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(16*gamma(5/4)) + sqrt(pi)*x*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))/4 + sin(2*(x + 1/2)**2)*gamma(1/4)/(32*gamma(5/4)) - sqrt(pi)*fresnelc(2*x/sqrt(pi) + 1/sqrt(pi))*gamma(1/4)/(32*gamma(5/4))`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.38 (sec) , antiderivative size = 136, normalized size of antiderivative = 2.96

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{32x^3 + 16x^2 + 8x(-i e^{(2ix^2+2ix+\frac{1}{2}i)} + i e^{(-2ix^2-2ix-\frac{1}{2}i)}) + \sqrt{8x^2+8x+2}((i-1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{2ix^2+2ix+\frac{1}{2}}) - \operatorname{erf}(\sqrt{2ix^2+2ix+\frac{1}{2}} - \sqrt{2}))}{64(2)}$$

[In] `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="maxima")`

[Out] $\frac{1}{64} (32x^3 + 16x^2 + 8x(-I e^{(2Ix^2+2Ix+\frac{1}{2}I)} + I e^{(-2Ix^2-2Ix-\frac{1}{2}I)}) + \sqrt{8x^2+8x+2}((I-1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{2Ix^2+2Ix+\frac{1}{2}}) - \operatorname{erf}(\sqrt{2Ix^2+2Ix+\frac{1}{2}} - \sqrt{2})) - (I+1)\sqrt{2}\sqrt{\pi}(\operatorname{erf}(\sqrt{2Ix^2+2Ix+\frac{1}{2}}) - \operatorname{erf}(\sqrt{2Ix^2+2Ix+\frac{1}{2}} - \sqrt{2})) - 4I e^{(2Ix^2+2Ix+\frac{1}{2}I)} + 4I e^{(-2Ix^2-2Ix-\frac{1}{2}I)})/(2x+1)$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.46 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.17

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4}x^2 + \left(\frac{1}{32}i + \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) - \left(\frac{1}{32}i - \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) - \frac{1}{16}i e^{(2ix^2+2ix+\frac{1}{2}i)} + \frac{1}{16}i e^{(-2ix^2-2ix-\frac{1}{2}i)}$$

[In] `integrate(x*cos(1/4+x+x^2)^2,x, algorithm="giac")`

[Out] $\frac{1}{4}x^2 + \left(\frac{1}{32}I + \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left((I-1)x + \frac{1}{2}I - \frac{1}{2} \right) - \left(\frac{1}{32}I - \frac{1}{32} \right) \sqrt{\pi} \operatorname{erf} \left(-(I+1)x - \frac{1}{2}I - \frac{1}{2} \right) - \frac{1}{16}I e^{(2Ix^2+2Ix+\frac{1}{2}I)} + \frac{1}{16}I e^{(-2Ix^2-2Ix-\frac{1}{2}I)}$

Mupad [F(-1)]

Timed out.

$$\int x \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int x \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

[In] `int(x*cos(x + x^2 + 1/4)^2,x)`

[Out] `int(x*cos(x + x^2 + 1/4)^2, x)`

3.26 $\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx$

Optimal result	164
Rubi [A] (verified)	164
Mathematica [A] (verified)	165
Maple [A] (verified)	165
Fricas [A] (verification not implemented)	166
Sympy [A] (verification not implemented)	166
Maxima [C] (verification not implemented)	166
Giac [C] (verification not implemented)	167
Mupad [F(-1)]	167

Optimal result

Integrand size = 11, antiderivative size = 27

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x}{2} + \frac{1}{4} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right)$$

[Out] $1/2*x+1/4*\operatorname{FresnelC}((1+2*x)/\operatorname{Pi}^{(1/2)})*\operatorname{Pi}^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3531, 3527, 3433}

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\pi} \operatorname{FresnelC} \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{x}{2}$$

[In] $\operatorname{Int}[\operatorname{Cos}[1/4 + x + x^2]^2, x]$

[Out] $x/2 + (\operatorname{Sqrt}[\operatorname{Pi}] * \operatorname{FresnelC}[(1 + 2*x)/\operatorname{Sqrt}[\operatorname{Pi}]])/4$

Rule 3433

```
Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simp[(Sqrt[Pi/2]/(f*Rt[d, 2]))*FresnelC[Sqrt[2/Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]
```

Rule 3527

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_.)^2], x_Symbol] :> Int[Cos[(b + 2*c*x)^2/(4*c)], x] /; FreeQ[{a, b, c}, x] && EqQ[b^2 - 4*a*c, 0]
```

Rule 3531

```
Int[Cos[(a_.) + (b_ .)*(x_) + (c_ .)*(x_)^2]^n_, x_Symbol] :> Int[ExpandTri
gReduce[Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2} + \frac{1}{2} \cos \left(\frac{1}{2} + 2x + 2x^2 \right) \right) dx \\ &= \frac{x}{2} + \frac{1}{2} \int \cos \left(\frac{1}{2} + 2x + 2x^2 \right) dx \\ &= \frac{x}{2} + \frac{1}{2} \int \cos \left(\frac{1}{8}(2+4x)^2 \right) dx \\ &= \frac{x}{2} + \frac{1}{4}\sqrt{\pi} \text{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \left(2x + \sqrt{\pi} \text{FresnelC} \left(\frac{1+2x}{\sqrt{\pi}} \right) \right)$$

```
[In] Integrate[Cos[1/4 + x + x^2]^2, x]
[Out] (2*x + Sqrt[Pi]*FresnelC[(1 + 2*x)/Sqrt[Pi]])/4
```

Maple [A] (verified)

Time = 1.26 (sec), antiderivative size = 20, normalized size of antiderivative = 0.74

method	result	size
default	$\frac{x}{2} + \frac{\text{C}\left(\frac{1+2x}{\sqrt{\pi}}\right)\sqrt{\pi}}{4}$	20
risch	$\frac{x}{2} - \frac{\sqrt{\pi}\sqrt{2}(-1)^{\frac{3}{4}}\text{erf}\left(\sqrt{2}(-1)^{\frac{1}{4}}x + \frac{\sqrt{2}(-1)^{\frac{1}{4}}}{2}\right)}{16} + \frac{\sqrt{\pi}\text{erf}\left(\sqrt{-2i}x - \frac{i}{\sqrt{-2i}}\right)}{8\sqrt{-2i}}$	58

```
[In] int(cos(1/4+x+x^2)^2, x, method=_RETURNVERBOSE)
[Out] 1/2*x+1/4*FresnelC((1+2*x)/Pi^(1/2))*Pi^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.70

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{1}{4} \sqrt{\pi} C \left(\frac{2x+1}{\sqrt{\pi}} \right) + \frac{1}{2} x$$

[In] integrate(cos(1/4+x+x^2)^2,x, algorithm="fricas")

[Out] $\frac{1}{4} \sqrt{\pi} \operatorname{fresnel_cos}((2x + 1)/\sqrt{\pi}) + \frac{1}{2} x$

Sympy [A] (verification not implemented)

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.81

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \frac{x}{2} + \frac{\sqrt{\pi} C \left(\frac{4x+2}{2\sqrt{\pi}} \right)}{4}$$

[In] integrate(cos(1/4+x+x**2)**2,x)

[Out] $\frac{x}{2} + \sqrt{\pi} \operatorname{fresnelc}((4x + 2)/(2\sqrt{\pi}))/4$

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.26

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = -\frac{1}{16} \sqrt{\pi} \left((i-1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{2i}} \right) + (i+1) \operatorname{erf} \left(\frac{2ix+i}{\sqrt{-2i}} \right) \right) + \frac{1}{2} x$$

[In] integrate(cos(1/4+x+x^2)^2,x, algorithm="maxima")

[Out] $-\frac{1}{16} \sqrt{\pi} ((i-1) \operatorname{erf}((2ix+i)/\sqrt{2i}) + (i+1) \operatorname{erf}((2ix+i)/\sqrt{-2i})) + \frac{1}{2} x$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.96

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = - \left(\frac{1}{16}i + \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left((i-1)x + \frac{1}{2}i - \frac{1}{2} \right) \\ + \left(\frac{1}{16}i - \frac{1}{16} \right) \sqrt{\pi} \operatorname{erf} \left(-(i+1)x - \frac{1}{2}i - \frac{1}{2} \right) + \frac{1}{2}x$$

[In] `integrate(cos(1/4+x+x^2)^2,x, algorithm="giac")`

[Out] `-(1/16*I + 1/16)*sqrt(pi)*erf((I - 1)*x + 1/2*I - 1/2) + (1/16*I - 1/16)*sqrt(pi)*erf(-(I + 1)*x - 1/2*I - 1/2) + 1/2*x`

Mupad [F(-1)]

Timed out.

$$\int \cos^2 \left(\frac{1}{4} + x + x^2 \right) dx = \int \cos \left(x^2 + x + \frac{1}{4} \right)^2 dx$$

[In] `int(cos(x + x^2 + 1/4)^2,x)`

[Out] `int(cos(x + x^2 + 1/4)^2, x)`

3.27 $\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x} dx$

Optimal result	168
Rubi [N/A]	168
Mathematica [N/A]	169
Maple [N/A] (verified)	169
Fricas [N/A]	169
Sympy [N/A]	169
Maxima [N/A]	170
Giac [N/A]	170
Mupad [N/A]	170

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \frac{\log(x)}{2} + \frac{1}{2} \text{Int}\left(\frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

[Out] $1/2*\ln(x)+1/2*\text{Unintegrable}(\cos(1/2+2*x+2*x^2)/x,x)$

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] $\text{Int}[\cos[1/4 + x + x^2]^2/x, x]$

[Out] $\text{Log}[x]/2 + \text{Defer}[\text{Int}][\cos[1/2 + 2*x + 2*x^2]/x, x]/2$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2x} + \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} \right) dx \\ &= \frac{\log(x)}{2} + \frac{1}{2} \int \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{x} dx \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 7.14 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] `Integrate[Cos[1/4 + x + x^2]^2/x, x]`

[Out] `Integrate[Cos[1/4 + x + x^2]^2/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.36 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx$$

[In] `int(cos(1/4+x+x^2)^2/x, x)`

[Out] `int(cos(1/4+x+x^2)^2/x, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x, x, algorithm="fricas")`

[Out] `integral(cos(x^2 + x + 1/4)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.66 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.93

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos^2\left(x^2 + x + \frac{1}{4}\right)}{x} dx$$

[In] `integrate(cos(1/4+x+x**2)**2/x, x)`

[Out] `Integral(cos(x**2 + x + 1/4)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.60

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x,x, algorithm="maxima")`

[Out] `1/2*integrate(cos(2*x^2 + 2*x + 1/2)/x, x) + 1/2*log(x)`

Giac [N/A]

Not integrable

Time = 0.53 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x,x, algorithm="giac")`

[Out] `integrate(cos(x^2 + x + 1/4)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 13.38 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x} dx = \int \frac{\cos\left(x^2 + x + \frac{1}{4}\right)^2}{x} dx$$

[In] `int(cos(x + x^2 + 1/4)^2/x,x)`

[Out] `int(cos(x + x^2 + 1/4)^2/x, x)`

3.28 $\int \frac{\cos^2\left(\frac{1}{4}+x+x^2\right)}{x^2} dx$

Optimal result	171
Rubi [N/A]	171
Mathematica [N/A]	172
Maple [N/A] (verified)	172
Fricas [N/A]	173
Sympy [N/A]	173
Maxima [N/A]	173
Giac [N/A]	174
Mupad [N/A]	174

Optimal result

Integrand size = 15, antiderivative size = 15

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = -\frac{1}{2x} - \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} - \sqrt{\pi} \operatorname{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) \\ - \operatorname{Int}\left(\frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x}, x\right)$$

[Out] $-1/2/x - 1/2*\cos(1/2+2*x+2*x^2)/x - \operatorname{FresnelS}((1+2*x)/\operatorname{Pi}^{(1/2)})*\operatorname{Pi}^{(1/2)} - \operatorname{Unintegrible}(\sin(1/2+2*x+2*x^2)/x, x)$

Rubi [N/A]

Not integrable

Time = 0.06 (sec), antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

[In] $\operatorname{Int}[\operatorname{Cos}[1/4 + x + x^2]^2/x^2, x]$

[Out] $-1/2*1/x - \operatorname{Cos}[1/2 + 2*x + 2*x^2]/(2*x) - \operatorname{Sqrt}[\operatorname{Pi}]*\operatorname{FresnelS}[(1 + 2*x)/\operatorname{Sqrt}[\operatorname{Pi}]] - \operatorname{Defer}[\operatorname{Int}][\operatorname{Sin}[1/2 + 2*x + 2*x^2]/x, x]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{2x^2} + \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x^2} \right) dx \\
 &= -\frac{1}{2x} + \frac{1}{2} \int \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{x^2} dx \\
 &= -\frac{1}{2x} - \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} - 2 \int \sin\left(\frac{1}{2} + 2x + 2x^2\right) dx - \int \frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x} dx \\
 &= -\frac{1}{2x} - \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} - 2 \int \sin\left(\frac{1}{8}(2+4x)^2\right) dx - \int \frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x} dx \\
 &= -\frac{1}{2x} - \frac{\cos\left(\frac{1}{2} + 2x + 2x^2\right)}{2x} - \sqrt{\pi} \text{FresnelS}\left(\frac{1+2x}{\sqrt{\pi}}\right) - \int \frac{\sin\left(\frac{1}{2} + 2x + 2x^2\right)}{x} dx
 \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 7.05 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

[In] `Integrate[Cos[1/4 + x + x^2]^2/x^2, x]`

[Out] `Integrate[Cos[1/4 + x + x^2]^2/x^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.40 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.87

$$\int \frac{\cos^2\left(\frac{1}{4} + x + x^2\right)}{x^2} dx$$

[In] `int(cos(1/4+x+x^2)^2/x^2, x)`

[Out] `int(cos(1/4+x+x^2)^2/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x^2,x, algorithm="fricas")`

[Out] `integral(cos(x^2 + x + 1/4)^2/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.59 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos^2 \left(x^2 + x + \frac{1}{4}\right)}{x^2} dx$$

[In] `integrate(cos(1/4+x+x**2)**2/x**2,x)`

[Out] `Integral(cos(x**2 + x + 1/4)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.73

$$\int \frac{\cos^2 \left(\frac{1}{4} + x + x^2\right)}{x^2} dx = \int \frac{\cos \left(x^2 + x + \frac{1}{4}\right)^2}{x^2} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x^2,x, algorithm="maxima")`

[Out] `1/2*(x*integrate(cos(2*x^2 + 2*x + 1/2)/x^2, x) - 1)/x`

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(\frac{1}{4} + x + x^2)}{x^2} dx = \int \frac{\cos(x^2 + x + \frac{1}{4})^2}{x^2} dx$$

[In] `integrate(cos(1/4+x+x^2)^2/x^2,x, algorithm="giac")`

[Out] `integrate(cos(x^2 + x + 1/4)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 13.31 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(\frac{1}{4} + x + x^2)}{x^2} dx = \int \frac{\cos(x^2 + x + \frac{1}{4})^2}{x^2} dx$$

[In] `int(cos(x + x^2 + 1/4)^2/x^2,x)`

[Out] `int(cos(x + x^2 + 1/4)^2/x^2, x)`

3.29 $\int (d + ex)^2 \cos(a + bx + cx^2) dx$

Optimal result	175
Rubi [A] (verified)	176
Mathematica [A] (verified)	178
Maple [A] (verified)	179
Fricas [A] (verification not implemented)	179
Sympy [F]	180
Maxima [C] (verification not implemented)	180
Giac [C] (verification not implemented)	182
Mupad [F(-1)]	182

Optimal result

Integrand size = 19, antiderivative size = 285

$$\begin{aligned} \int (d + ex)^2 \cos(a + bx + cx^2) dx = & \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ & - \frac{e^2 \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ & - \frac{e^2 \sqrt{\frac{\pi}{2}} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ & - \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\ & + \frac{e(2cd - be) \sin(a + bx + cx^2)}{4c^2} \\ & + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \end{aligned}$$

```
[Out] 1/4*e*(-b*e+2*c*d)*sin(c*x^2+b*x+a)/c^2+1/2*e*(e*x+d)*sin(c*x^2+b*x+a)/c+1/
8*(-b*e+2*c*d)^2*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi
^(1/2))*2^(1/2)*Pi^(1/2)/c^(5/2)-1/4*e^2*cos(a-1/4*b^2/c)*FresnelS(1/2*(2*c
*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(3/2)-1/4*e^2*FresnelC(1
/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^
(3/2)-1/8*(-b*e+2*c*d)^2*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*s
in(a-1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.316, Rules used = {3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int (d + ex)^2 \cos(a + bx + cx^2) dx &= \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) (2cd - be)^2 \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) (2cd - be)^2 \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} e^2 \sin\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} e^2 \cos\left(a - \frac{b^2}{4c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad + \frac{e(2cd - be) \sin(a + bx + cx^2)}{4c^2} \\ &\quad + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \end{aligned}$$

[In] `Int[(d + e*x)^2*Cos[a + b*x + c*x^2], x]`

[Out] $((2*c*d - b*e)^2*\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{Cos}[a - b^2/(4*c)]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]]/(4*c^{5/2}) - (e^{2*\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{Cos}[a - b^2/(4*c)]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]}]/(2*c^{3/2}) - (e^{2*\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]})*\operatorname{Sin}[a - b^2/(4*c)]/(2*c^{3/2}) - ((2*c*d - b*e)^2*\operatorname{Sqrt}[\operatorname{Pi}/2]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[2*\operatorname{Pi}])]])*\operatorname{Sin}[a - b^2/(4*c)]/(4*c^{5/2}) + (e*(2*c*d - b*e))*\operatorname{Sin}[a + b*x + c*x^2]/(4*c^2) + (e*(d + e*x))*\operatorname{Sin}[a + b*x + c*x^2]/(2*c)$

Rule 3432

`Int[$\operatorname{Sin}[(d_{_})*(e_{_} + f_{_})*(x_{_})^2]$, x_Symbol] :> Simp[($\operatorname{Sqrt}[\operatorname{Pi}/2]/(f*\operatorname{Rt}[d, 2])$)* $\operatorname{FresnelS}[\operatorname{Sqrt}[2/\operatorname{Pi}]*\operatorname{Rt}[d, 2]*(e + f*x)]$, x] /; FreeQ[{d, e, f}, x]`

Rule 3433

`Int[$\operatorname{Cos}[(d_{_})*(e_{_} + f_{_})*(x_{_})^2]$, x_Symbol] :> Simp[($\operatorname{Sqrt}[\operatorname{Pi}/2]/(f*\operatorname{Rt}[d, 2])$)* $\operatorname{FresnelC}[\operatorname{Sqrt}[2/\operatorname{Pi}]*\operatorname{Rt}[d, 2]*(e + f*x)]$, x] /; FreeQ[{d, e, f}, x]`

Rule 3528

`Int[$\operatorname{Sin}[(a_{_}) + (b_{_})*(x_{_}) + (c_{_})*(x_{_})^2]$, x_Symbol] :> Dist[$\operatorname{Cos}[(b^2 - 4*a*c)/(4*c)]$, Int[$\operatorname{Sin}[(b + 2*c*x)^2/(4*c)]$, x], x] - Dist[$\operatorname{Sin}[(b^2 - 4*a*c)/(4*c)]$, Int[$\operatorname{Cos}[(b + 2*c*x)^2/(4*c)]$, x], x] /; FreeQ[{a, b, c}, x] && NeQ[`

$b^2 - 4*a*c, 0]$

Rule 3529

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]*((d_.) + (e_.*)(x_)), x_Symbol]
: > Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_))^(m_), x_Sym
bol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[
(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - D
ist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x]
) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} - \frac{e^2 \int \sin(a + bx + cx^2) dx}{2c} \\
&\quad - \frac{(-2cd + be) \int (d + ex) \cos(a + bx + cx^2) dx}{2c} \\
&= \frac{e(2cd - be) \sin(a + bx + cx^2)}{4c^2} + \frac{e(d + ex) \sin(a + bx + cx^2)}{2c} \\
&\quad + \frac{(2cd - be)^2 \int \cos(a + bx + cx^2) dx}{4c^2} - \frac{\left(e^2 \cos\left(a - \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\
&\quad - \frac{\left(e^2 \sin\left(a - \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{e^2 \sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c}\right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{e^2 \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin \left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\
&\quad + \frac{e(2cd - be) \sin (a + bx + cx^2)}{4c^2} + \frac{e(d + ex) \sin (a + bx + cx^2)}{2c} \\
&\quad + \frac{\left((2cd - be)^2 \cos \left(a - \frac{b^2}{4c}\right)\right) \int \cos \left(\frac{(b+2cx)^2}{4c}\right) dx}{4c^2} \\
&\quad - \frac{\left((2cd - be)^2 \sin \left(a - \frac{b^2}{4c}\right)\right) \int \sin \left(\frac{(b+2cx)^2}{4c}\right) dx}{4c^2} \\
&= \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c}\right) \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{4c^{5/2}} \\
&\quad - \frac{e^2 \sqrt{\frac{\pi}{2}} \cos \left(a - \frac{b^2}{4c}\right) \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} - \frac{e^2 \sqrt{\frac{\pi}{2}} \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin \left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\
&\quad - \frac{(2cd - be)^2 \sqrt{\frac{\pi}{2}} \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin \left(a - \frac{b^2}{4c}\right)}{4c^{5/2}} \\
&\quad + \frac{e(2cd - be) \sin (a + bx + cx^2)}{4c^2} + \frac{e(d + ex) \sin (a + bx + cx^2)}{2c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.79 (sec), antiderivative size = 187, normalized size of antiderivative = 0.66

$$\begin{aligned}
&\int (d + ex)^2 \cos (a + bx + cx^2) dx \\
&= \frac{\sqrt{2\pi} \text{FresnelC} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left((-2cd + be)^2 \cos \left(a - \frac{b^2}{4c}\right) - 2ce^2 \sin \left(a - \frac{b^2}{4c}\right)\right) - \sqrt{2\pi} \text{FresnelS} \left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \left(2ce^2 \cos \left(a - \frac{b^2}{4c}\right) + (-2cd + be)^2 \sin \left(a - \frac{b^2}{4c}\right)\right)}{8c^{5/2}}
\end{aligned}$$

```

[In] Integrate[(d + e*x)^2*Cos[a + b*x + c*x^2], x]
[Out] (Sqrt[2*Pi]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*((-2*c*d + b*e)^2*Cos[a - b^2/(4*c)] - 2*c*e^2*Sin[a - b^2/(4*c)]) - Sqrt[2*Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[2*Pi])]*(2*c*e^2*Cos[a - b^2/(4*c)] + (-2*c*d + b*e)^2*Sin[a - b^2/(4*c)]) + 2*Sqrt[c]*e*(4*c*d - b*e + 2*c*e*x)*Sin[a + x*(b + c*x)])/(8*c^(5/2))

```

Maple [A] (verified)

Time = 2.38 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.39

method	result
default	$\frac{e^2 b \left(\frac{\sin(cx^2+bx+a)}{2c} - \frac{b\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2-ac}{4c}\right) C\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2-ac}{4c}\right) S\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{\frac{3}{2}}} \right)}{2c} - \frac{e^2 \sqrt{2} \cdot \text{...}}{2c}$
risch	$\frac{\operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}d^2 e^{-\frac{i(4ac-b^2)}{4c}}}{4\sqrt{ic}} + \frac{e^2 \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}b^2 e^{-\frac{i(4ac-b^2)}{4c}}}{16\sqrt{ic}c^2} - \frac{ie^2 \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}}{8\sqrt{ic}c} - \frac{de^2 \operatorname{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}e^{-\frac{i(4ac-b^2)}{4c}}}{16\sqrt{ic}c^2}$
parts	Expression too large to display

[In] `int((e*x+d)^2*cos(c*x^2+b*x+a),x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned} & 1/2 * e^2 / c * x * \sin(c*x^2 + b*x + a) - 1/2 * e^2 * b / c * (1/2 * \sin(c*x^2 + b*x + a) / c - 1/4 * b / c^{(3/2)} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a*c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b)) + \sin((1/4 * b^2 - a*c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b))) - 1/4 * e^2 / c^{(3/2)} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a*c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b)) - \sin((1/4 * b^2 - a*c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b))) + d * e / c * \sin(c*x^2 + b*x + a) - 1/2 * d * e * b / c^{(3/2)} * 2^{(1/2)} * \text{Pi}^{(1/2)} * (\cos((1/4 * b^2 - a*c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b)) + \sin((1/4 * b^2 - a*c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b))) + 1/2 * 2^{(1/2)} * \text{Pi}^{(1/2)} / c^{(1/2)} * d^2 * (\cos((1/4 * b^2 - a*c) / c) * \text{FresnelC}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b)) + \sin((1/4 * b^2 - a*c) / c) * \text{FresnelS}(2^{(1/2)} / \text{Pi}^{(1/2)} / c^{(1/2)} * (c*x + 1/2 * b))) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 230, normalized size of antiderivative = 0.81

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx =$$

$$-\frac{\sqrt{2} \left(2 \pi c e^2 \sin\left(-\frac{b^2-4ac}{4c}\right) - \pi (4c^2 d^2 - 4bcde + b^2 e^2) \cos\left(-\frac{b^2-4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) + \sqrt{2} \left(2 \pi c e^2 \cos\left(-\frac{b^2-4ac}{4c}\right) - \pi (4c^2 d^2 - 4bcde + b^2 e^2) \sin\left(-\frac{b^2-4ac}{4c}\right) \right) \sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right)}{2c}$$

[In] `integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="fricas")`

[Out]

$$\begin{aligned} & -1/8 * (\sqrt{2} * (2 * \pi * c * e^2 * \sin(-1/4 * (b^2 - 4*a*c) / c) - \pi * (4*c^2 * d^2 - 4*b*c*d*e + b^2 * e^2) * \cos(-1/4 * (b^2 - 4*a*c) / c) * \sqrt{c / \pi} * \text{fresnel_cos}(1/2 * \sqrt{c / \pi} * (2*c*x + b) * \sqrt{c / \pi} / c) + \sqrt{2} * (2 * \pi * c * e^2 * \cos(-1/4 * (b^2 - 4*a*c) / c) - \pi * (4*c^2 * d^2 - 4*b*c*d*e + b^2 * e^2) * \sin(-1/4 * (b^2 - 4*a*c) / c) * \sqrt{c / \pi} * \text{fresnel_sin}(1/2 * \sqrt{c / \pi} * (2*c*x + b) * \sqrt{c / \pi} / c))) \end{aligned}$$

$$\pi)*\text{fresnel_sin}(1/2*\sqrt{2}*(2*c*x + b)*\sqrt{c/\pi})/c) - 2*(2*c^2*e^{2*x} + 4*c^2*d*e - b*c*e^2)*\sin(c*x^2 + b*x + a))/c^3$$

Sympy [F]

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \int (d + ex)^2 \cos(a + bx + cx^2) dx$$

[In] `integrate((e*x+d)**2*cos(c*x**2+b*x+a),x)`
[Out] `Integral((d + e*x)**2*cos(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.34 (sec) , antiderivative size = 2271, normalized size of antiderivative = 7.97

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \text{Too large to display}$$

[In] `integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="maxima")`
[Out]
$$\begin{aligned} & -1/8*\sqrt{2}*\sqrt{\pi}*((I - 1)*\cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*\sin(-1/4*(b^2 - 4*a*c)/c)) * \operatorname{erf}(1/2*(2*I*c*x + I*b)/\sqrt{I*c}) + ((I + 1)*\cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*\sin(-1/4*(b^2 - 4*a*c)/c)) * \operatorname{erf}(1/2*(2*I*c*x + I*b)/\sqrt{-I*c})) * d^2/\sqrt{c} + 1/8*((I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1)) * b^2*\cos(-1/4*(b^2 - 4*a*c)/c) + ((I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) - (I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1)) * b^2*\sin(-1/4*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) + (I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1)) * b*c*\cos(-1/4*(b^2 - 4*a*c)/c) + (-(I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1)) * b*c*\sin(-1/4*(b^2 - 4*a*c)/c)) * x - 4*(c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) * \cos(-1/4*(b^2 - 4*a*c)/c) - c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) * \sin(-1/4*(b^2 - 4*a*c)/c)) * \sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c)) + 1/32*(8*((-(I - 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1) + (I + 1)*\sqrt{2}*\sqrt{\pi})*(\operatorname{erf}(1/2*\sqrt{-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c}) - 1))$$

$$\begin{aligned}
& - 1)) * b^2 * c^3 + 4 * ((I + 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * \\
& I * b * c * x + I * b^2) / c) - (I - 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * \\
& c * x + I * b^2) / c)) * c^4) * \cos(-1/4 * (b^2 - 4 * a * c) / c) + ((-(I + 1) * \sqrt{2} * \sqrt{\pi}) * \\
& (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) + (I - 1) * \sqrt{2} * \\
& \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^2 * \\
& c^3 + 4 * (- (I - 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / \\
& c) + (I + 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * \\
& c^4) * \sin(-1/4 * (b^2 - 4 * a * c) / c) * x^3 + 12 * (((-(I - 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \\
& \sqrt{(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) + (I + 1) * \sqrt{2} * \sqrt{\pi}) * \\
& (\text{erf}(1/2 * \sqrt{-(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^3 * c^2 + 4 * \\
& ((I + 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c) - (I - \\
& 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * b * c^3) * \\
& \cos(-1/4 * (b^2 - 4 * a * c) / c) + ((-(I + 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * \\
& x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) + (I - 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{ \\
& -(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^3 * c^2 + 4 * (- (I - 1) * \sqrt{2} * \\
& \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c) + (I + 1) * \sqrt{2} * \text{gamma}(3/2, \\
& -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * b * c^3) * \sin(-1/4 * (b^2 - 4 * \\
& a * c) / c) * x^2 + 8 * (b * c^2 * (I * e^{(1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)} - I * \\
& e^{(-1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)}) * \cos(-1/4 * (b^2 - 4 * a * c) / c) - \\
& b * c^2 * (e^{(1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)} + e^{(-1/4 * (4 * I * c^2 * x^2 + \\
& 4 * I * b * c * x + I * b^2) / c)}) * \sin(-1/4 * (b^2 - 4 * a * c) / c) * ((4 * c^2 * x^2 + 4 * b * c * x + \\
& b^2) / c)^{(3/2)} + 6 * (((-(I - 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 + \\
& 4 * I * b * c * x + I * b^2) / c}) - 1) + (I + 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{ \\
& -(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^4 * c + 4 * ((I + 1) * \sqrt{2} * \text{gamma}(3/2, \\
& 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c) - (I - 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * \\
& (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * b^2 * c^2) * \cos(-1/4 * (b^2 - 4 * a * c) / c) \\
& + ((-(I + 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) + (I - 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{ \\
& -(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^4 * c + 4 * (- (I - 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * \\
& (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c) + (I + 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * \\
& x^2 + 4 * I * b * c * x + I * b^2) / c)) * b^2 * c^2 * \sin(-1/4 * (b^2 - 4 * a * c) / c) * x - (((I - 1) * \\
& \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) - (I + 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{ \\
& -(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^5 - 4 * ((I + 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + \\
& I * b^2) / c) - (I - 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * b^3 * c) * \\
& \cos(-1/4 * (b^2 - 4 * a * c) / c) - (((I + 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1) - (I - 1) * \sqrt{2} * \sqrt{\pi}) * (\text{erf}(1/2 * \sqrt{ \\
& -(4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c}) - 1)) * b^5 - 4 * ((I - 1) * \sqrt{2} * \text{gamma}(3/2, 1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c) + (I + 1) * \sqrt{2} * \text{gamma}(3/2, -1/4 * (4 * I * c^2 * x^2 + 4 * I * b * c * x + I * b^2) / c)) * b^3 * c) * \sin \\
& (-1/4 * (b^2 - 4 * a * c) / c) * e^{(2 * (4 * c^2 * x^2 + 4 * b * c * x + b^2) / c)^{(3/2)}})
\end{aligned}$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.01

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx =$$

$$-\frac{\sqrt{2}\sqrt{\pi}(-4ic^2d^2+4ibcde-i b^2e^2+2ce^2)\operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}\left(2x+\frac{b}{c}\right)\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2-4iac}{4c}\right)}}{\left(\frac{ic}{|c|}+1\right)\sqrt{|c|}} + 2i\left(ce^2\left(2x+\frac{b}{c}\right)+4cde-2be^2\right)e^{\left(-\frac{ib^2+4iac}{4c}\right)}$$

$$-\frac{\sqrt{2}\sqrt{\pi}(4ic^2d^2-4ibcde+i b^2e^2+2ce^2)\operatorname{erf}\left(\frac{1}{4}i\sqrt{2}\left(2x+\frac{b}{c}\right)\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2+4iac}{4c}\right)}}{\left(-\frac{ic}{|c|}+1\right)\sqrt{|c|}} - 2i\left(ce^2\left(2x+\frac{b}{c}\right)+4cde-2be^2\right)e^{\left(-\frac{-ib^2-4iac}{4c}\right)}$$

$$16c^2$$

[In] `integrate((e*x+d)^2*cos(c*x^2+b*x+a),x, algorithm="giac")`

[Out] $-1/16*(\sqrt{2}*\sqrt{\pi})*(-4*I*c^2*d^2 + 4*I*b*c*d*e - I*b^2*e^2 + 2*c*e^2)*\operatorname{erf}\left(-1/4*I*\sqrt{2}*(2*x + b/c)*(I*c/abs(c) + 1)*\sqrt{abs(c)}\right)*e^{(-1/4*(I*b^2 - 4*I*a*c)/c)} / ((I*c/abs(c) + 1)*\sqrt{abs(c)}) + 2*I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^{(I*c*x^2 + I*b*x + I*a)}/c^2 - 1/16*(\sqrt{2}*\sqrt{\pi})*(4*I*c^2*d^2 - 4*I*b*c*d*e + I*b^2*e^2 + 2*c*e^2)*\operatorname{erf}\left(1/4*I*\sqrt{2}*(2*x + b/c)*(-I*c/abs(c) + 1)*\sqrt{abs(c)}\right)*e^{(-1/4*(-I*b^2 + 4*I*a*c)/c)} / ((-I*c/abs(c) + 1)*\sqrt{abs(c)}) - 2*I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^{(-I*c*x^2 - I*b*x - I*a)}/c^2$

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \cos(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a) (d + ex)^2 dx$$

[In] `int(cos(a + b*x + c*x^2)*(d + e*x)^2,x)`

[Out] `int(cos(a + b*x + c*x^2)*(d + e*x)^2, x)`

3.30 $\int (d + ex) \cos(a + bx + cx^2) dx$

Optimal result	183
Rubi [A] (verified)	183
Mathematica [A] (verified)	185
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	186
Sympy [F]	186
Maxima [C] (verification not implemented)	186
Giac [C] (verification not implemented)	187
Mupad [F(-1)]	188

Optimal result

Integrand size = 17, antiderivative size = 140

$$\begin{aligned} \int (d + ex) \cos(a + bx + cx^2) dx &= \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{(2cd - be)\sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} \\ &\quad + \frac{e \sin(a + bx + cx^2)}{2c} \end{aligned}$$

```
[Out] 1/2*e*sin(c*x^2+b*x+a)/c+1/4*(-b*e+2*c*d)*cos(a-1/4*b^2/c)*FresnelC(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*2^(1/2)*Pi^(1/2)/c^(3/2)-1/4*(-b*e+2*c*d)*FresnelS(1/2*(2*c*x+b)/c^(1/2)*2^(1/2)/Pi^(1/2))*sin(a-1/4*b^2/c)*2^(1/2)*Pi^(1/2)/c^(3/2)
```

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3543, 3529, 3433, 3432}

$$\begin{aligned} \int (d + ex) \cos(a + bx + cx^2) dx &= \frac{\sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) (2cd - be) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{\sqrt{\frac{\pi}{2}} \sin\left(a - \frac{b^2}{4c}\right) (2cd - be) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad + \frac{e \sin(a + bx + cx^2)}{2c} \end{aligned}$$

[In] $\text{Int}[(d + e*x)*\cos[a + b*x + c*x^2], x]$

[Out] $((2*c*d - b*e)*\sqrt{\pi/2}*\cos[a - b^2/(4*c)]*\text{FresnelC}[(b + 2*c*x)/(\sqrt{c}*\sqrt{2*\pi})])/(2*c^{(3/2)}) - ((2*c*d - b*e)*\sqrt{\pi/2}*\text{FresnelS}[(b + 2*c*x)/(\sqrt{c}*\sqrt{2*\pi})])*\sin[a - b^2/(4*c)]/(2*c^{(3/2)}) + (e*\sin[a + b*x + c*x^2])/(2*c)$

Rule 3432

$\text{Int}[\sin[(d_*)*(e_*) + (f_*)*(x_*)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi/2}/(f*\text{Rt}[d, 2]))*\text{FresnelS}[\sqrt{2/\pi}*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\cos[(d_*)*(e_*) + (f_*)*(x_*)^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi/2}/(f*\text{Rt}[d, 2]))*\text{FresnelC}[\sqrt{2/\pi}*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3529

$\text{Int}[\cos[(a_*) + (b_*)*(x_*) + (c_*)*(x_*)^2], x_{\text{Symbol}}] \rightarrow \text{Dist}[\cos[(b^2 - 4*a*c)/(4*c)], \text{Int}[\cos[(b + 2*c*x)^2/(4*c)], x], x] + \text{Dist}[\sin[(b^2 - 4*a*c)/(4*c)], \text{Int}[\sin[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 3543

$\text{Int}[\cos[(a_*) + (b_*)*(x_*) + (c_*)*(x_*)^2]*((d_*) + (e_*)*(x_*)), x_{\text{Symbol}}] \rightarrow \text{Simp}[e*(\sin[a + b*x + c*x^2]/(2*c)), x] + \text{Dist}[(2*c*d - b*e)/(2*c), \text{Int}[\cos[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{e \sin(a + bx + cx^2)}{2c} + \frac{(2cd - be) \int \cos(a + bx + cx^2) dx}{2c} \\ &= \frac{e \sin(a + bx + cx^2)}{2c} + \frac{\left((2cd - be) \cos\left(a - \frac{b^2}{4c}\right)\right) \int \cos\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\ &\quad - \frac{\left((2cd - be) \sin\left(a - \frac{b^2}{4c}\right)\right) \int \sin\left(\frac{(b+2cx)^2}{4c}\right) dx}{2c} \\ &= \frac{(2cd - be) \sqrt{\frac{\pi}{2}} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right)}{2c^{3/2}} \\ &\quad - \frac{(2cd - be) \sqrt{\frac{\pi}{2}} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right)}{2c^{3/2}} + \frac{e \sin(a + bx + cx^2)}{2c} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.44 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.92

$$\int (d + ex) \cos(a + bx + cx^2) dx \\ = \frac{(2cd - be)\sqrt{2\pi} \cos\left(a - \frac{b^2}{4c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) - (2cd - be)\sqrt{2\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{2\pi}}\right) \sin\left(a - \frac{b^2}{4c}\right) + 2\sqrt{c}e \sin(a + bx + cx^2)}{4c^{3/2}}$$

[In] Integrate[(d + e*x)*Cos[a + b*x + c*x^2], x]

[Out] $((2*c*d - b*e)*\text{Sqrt}[2*\text{Pi}]*\text{Cos}[a - b^2/(4*c)]*\text{FresnelC}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])] - (2*c*d - b*e)*\text{Sqrt}[2*\text{Pi}]*\text{FresnelS}[(b + 2*c*x)/(\text{Sqrt}[c]*\text{Sqrt}[2*\text{Pi}])]*)\text{Sin}[a - b^2/(4*c)] + 2*\text{Sqrt}[c]*e*\text{Sin}[a + x*(b + c*x)])/(4*c^{(3/2)})$

Maple [A] (verified)

Time = 0.75 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.30

method	result
default	$\frac{e \sin(cx^2 + bx + a)}{2c} - \frac{eb\sqrt{2}\sqrt{\pi} \left(\cos\left(\frac{b^2-ac}{c}\right) C\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2-ac}{c}\right) S\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}} + \frac{\sqrt{2}\sqrt{\pi} d \left(\cos\left(\frac{b^2-ac}{c}\right) C\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) + \sin\left(\frac{b^2-ac}{c}\right) S\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) \right)}{4c^{3/2}}$
risch	$\frac{\text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{4c}}}{4\sqrt{ic}} - \frac{e\text{erf}\left(\sqrt{ic}x + \frac{ib}{2\sqrt{ic}}\right)\sqrt{\pi}be^{-\frac{i(4ac-b^2)}{4c}}}{8\sqrt{ic}c} - \frac{\text{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi}de^{\frac{i(4ac-b^2)}{4c}}}{4\sqrt{-ic}} + \frac{e\text{erf}\left(-\sqrt{-ic}x + \frac{ib}{2\sqrt{-ic}}\right)\sqrt{\pi}be^{\frac{i(4ac-b^2)}{4c}}}{8\sqrt{-ic}c}$
parts	$\frac{\sqrt{2}\sqrt{\pi} \cos\left(\frac{b^2-ac}{c}\right) C\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi} \sin\left(\frac{b^2-ac}{c}\right) S\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) ex}{2\sqrt{c}} + \frac{\sqrt{2}\sqrt{\pi} \cos\left(\frac{b^2-ac}{c}\right) C\left(\frac{\sqrt{2}(cx+\frac{b}{2})}{\sqrt{\pi}\sqrt{c}}\right) d}{2\sqrt{c}}$

[In] int((e*x+d)*cos(c*x^2+b*x+a), x, method=_RETURNVERBOSE)

[Out] $1/2*e*\sin(c*x^2+b*x+a)/c - 1/4*e*b/c^{(3/2)}*2^{(1/2)}*\text{Pi}^{(1/2)}*(\cos((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)) + \sin((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b))) + 1/2*2^{(1/2)}*\text{Pi}^{(1/2)}/c^{(1/2)}*d*(\cos((1/4*b^2-a*c)/c)*\text{FresnelC}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)) + \sin((1/4*b^2-a*c)/c)*\text{FresnelS}(2^{(1/2)}/\text{Pi}^{(1/2)}/c^{(1/2)}*(c*x+1/2*b)))$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 136, normalized size of antiderivative = 0.97

$$\int (d + ex) \cos(a + bx + cx^2) dx = \frac{\sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4ac}{4c}\right) C\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) - \sqrt{2}\pi(2cd - be)\sqrt{\frac{c}{\pi}} S\left(\frac{\sqrt{2}(2cx+b)\sqrt{\frac{c}{\pi}}}{2c}\right) \sin\left(-\frac{b^2 - 4ac}{4c}\right)}{4c^2}$$

[In] `integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="fricas")`

[Out] $\frac{1}{4} \left(\sqrt{2} \pi (2 c d - b e) \sqrt{\frac{c}{\pi}} \cos\left(\frac{b^2 - 4 a c}{4 c}\right) \text{fresnel_cos}\left(\frac{1}{2} \sqrt{2} (2 c x + b) \sqrt{\frac{c}{\pi}}\right) - \sqrt{2} \pi (2 c d - b e) \sqrt{\frac{c}{\pi}} \text{fresnel_sin}\left(\frac{1}{2} \sqrt{2} (2 c x + b) \sqrt{\frac{c}{\pi}}\right) \right) \text{sqrt}\left(\frac{c}{\pi}\right) + 2 c e \sin(c x^2 + b x + a)) / c^2$

Sympy [F]

$$\int (d + ex) \cos(a + bx + cx^2) dx = \int (d + ex) \cos(a + bx + cx^2) dx$$

[In] `integrate((e*x+d)*cos(c*x**2+b*x+a),x)`

[Out] `Integral((d + e*x)*cos(a + b*x + c*x**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.59 (sec) , antiderivative size = 695, normalized size of antiderivative = 4.96

$$\begin{aligned} \int (d + ex) \cos(a + bx + cx^2) dx = & \\ & - \frac{\sqrt{2}\sqrt{\pi} \left(((i-1) \cos\left(-\frac{b^2 - 4ac}{4c}\right) + (i+1) \sin\left(-\frac{b^2 - 4ac}{4c}\right)) \operatorname{erf}\left(\frac{2icx+ib}{2\sqrt{ic}}\right) + ((i+1) \cos\left(-\frac{b^2 - 4ac}{4c}\right) + (i-1) \sin\left(-\frac{b^2 - 4ac}{4c}\right)) \operatorname{erf}\left(\frac{2icx-ib}{2\sqrt{ic}}\right) \right)}{8\sqrt{c}} \\ & + \frac{\left(((i-1) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{1}{2}\sqrt{\frac{4ic^2x^2+4ibcx+ib^2}{c}}\right) - 1) - (i+1) \sqrt{2}\sqrt{\pi} \operatorname{erf}\left(\frac{1}{2}\sqrt{-\frac{4ic^2x^2+4ibcx+ib^2}{c}}\right) - 1 \right)}{8\sqrt{c}} \end{aligned}$$

[In] `integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="maxima")`

```
[Out] -1/8*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(I*c)) + ((I + 1)*cos(-1/4*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/4*(b^2 - 4*a*c)/c))*erf(1/2*(2*I*c*x + I*b)/sqrt(-I*c)))*d/sqrt(c) + 1/16*((((I - 1)*sqrt(2)*sqrt(pi))*(erf(1/2*sqrt(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/4*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/4*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/4*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(1/2*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/4*(b^2 - 4*a*c)/c))*x - 4*(c*(I*e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - I*e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/4*(b^2 - 4*a*c)/c) - c*(e^(1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + e^(-1/4*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/4*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))/e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 199, normalized size of antiderivative = 1.42

$$\begin{aligned} & \int (d + ex) \cos(a + bx + cx^2) dx \\ &= \frac{\frac{\sqrt{2}\sqrt{\pi}(2icd - ibe)\operatorname{erf}\left(-\frac{1}{4}i\sqrt{2}\left(2x + \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{ib^2 - 4iac}{4c}\right)}}{\left(\frac{ic}{|c|} + 1\right)\sqrt{|c|}} - 2ie^{(icx^2 + ibx + ia)}}{8c} \\ &+ \frac{\frac{\sqrt{2}\sqrt{\pi}(-2icd + ibe)\operatorname{erf}\left(\frac{1}{4}i\sqrt{2}\left(2x + \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}\right)e^{\left(-\frac{-ib^2 + 4iac}{4c}\right)}}{\left(-\frac{ic}{|c|} + 1\right)\sqrt{|c|}} + 2ie^{(-icx^2 - ibx - ia)}}{8c} \end{aligned}$$

```
[In] integrate((e*x+d)*cos(c*x^2+b*x+a),x, algorithm="giac")
```

```
[Out] 1/8*(sqrt(2)*sqrt(pi)*(2*I*c*d - I*b*e)*erf(-1/4*I*sqrt(2)*(2*x + b/c)*(I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(I*b^2 - 4*I*a*c)/c)/((I*c/abs(c) + 1)*sqrt(abs(c))) - 2*I*e*e^(I*c*x^2 + I*b*x + I*a)/c + 1/8*(sqrt(2)*sqrt(pi)*(-2*I*c*d + I*b*e)*erf(1/4*I*sqrt(2)*(2*x + b/c)*(-I*c/abs(c) + 1)*sqrt(abs(c)))*e^(-1/4*(-I*b^2 + 4*I*a*c)/c)/((-I*c/abs(c) + 1)*sqrt(abs(c))) + 2*I*e*e^(-I*c*x^2 - I*b*x - I*a)/c
```

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \cos(a + bx + cx^2) \, dx = \int \cos(cx^2 + bx + a) (d + ex) \, dx$$

[In] `int(cos(a + b*x + c*x^2)*(d + e*x),x)`

[Out] `int(cos(a + b*x + c*x^2)*(d + e*x), x)`

3.31 $\int \frac{\cos(a+bx+cx^2)}{d+ex} dx$

Optimal result	189
Rubi [N/A]	189
Mathematica [N/A]	190
Maple [N/A] (verified)	190
Fricas [N/A]	190
Sympy [N/A]	190
Maxima [N/A]	191
Giac [N/A]	191
Mupad [N/A]	191

Optimal result

Integrand size = 19, antiderivative size = 19

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \text{Int}\left(\frac{\cos(a + bx + cx^2)}{d + ex}, x\right)$$

[Out] Unintegrable($\cos(c*x^2+b*x+a)/(e*x+d)$, x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec), antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

[In] Int[$\cos(a + b*x + c*x^2)/(d + e*x)$, x]

[Out] Defer[Int][$\cos(a + b*x + c*x^2)/(d + e*x)$, x]

Rubi steps

$$\text{integral} = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

Mathematica [N/A]

Not integrable

Time = 3.24 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

[In] `Integrate[Cos[a + b*x + c*x^2]/(d + e*x), x]`

[Out] `Integrate[Cos[a + b*x + c*x^2]/(d + e*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.43 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.00

$$\int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

[In] `int(cos(c*x^2+b*x+a)/(e*x+d), x)`

[Out] `int(cos(c*x^2+b*x+a)/(e*x+d), x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/(e*x+d), x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.89

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(a + bx + cx^2)}{d + ex} dx$$

[In] `integrate(cos(c*x**2+b*x+a)/(e*x+d), x)`

[Out] `Integral(cos(a + b*x + c*x**2)/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/(e*x+d),x, algorithm="maxima")`

[Out] `integrate(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)/(e*x+d),x, algorithm="giac")`

[Out] `integrate(cos(c*x^2 + b*x + a)/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 13.46 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\cos(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)}{d + ex} dx$$

[In] `int(cos(a + b*x + c*x^2)/(d + e*x),x)`

[Out] `int(cos(a + b*x + c*x^2)/(d + e*x), x)`

3.32 $\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$

Optimal result	192
Rubi [A] (verified)	193
Mathematica [A] (verified)	195
Maple [A] (verified)	196
Fricas [A] (verification not implemented)	196
Sympy [F]	197
Maxima [C] (verification not implemented)	197
Giac [C] (verification not implemented)	199
Mupad [F(-1)]	199

Optimal result

Integrand size = 21, antiderivative size = 291

$$\begin{aligned} \int (d + ex)^2 \cos^2(a + bx + cx^2) dx &= \frac{(d + ex)^3}{6e} \\ &+ \frac{(2cd - be)^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ &- \frac{e^2 \sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ &- \frac{e^2 \sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{3/2}} \\ &- \frac{(2cd - be)^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{16c^{5/2}} \\ &+ \frac{e(2cd - be) \sin(2a + 2bx + 2cx^2)}{16c^2} \\ &+ \frac{e(d + ex) \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

```
[Out] 1/6*(e*x+d)^3/e+1/16*e*(-b*e+2*c*d)*sin(2*c*x^2+2*b*x+2*a)/c^2+1/8*e*(e*x+d)*sin(2*c*x^2+2*b*x+2*a)/c+1/16*(-b*e+2*c*d)^2*cos(2*a-1/2*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/c^(5/2)-1/16*e^2*cos(2*a-1/2*b^2/c)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/c^(3/2)-1/16*e^2*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)*Pi^(1/2)/c^(3/2)-1/16*(-b*e+2*c*d)^2*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)*Pi^(1/2)/c^(5/2)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 291, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3549, 3545, 3543, 3529, 3433, 3432, 3528}

$$\begin{aligned} \int (d + ex)^2 \cos^2(a + bx + cx^2) dx &= \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd - be)^2 \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ &\quad - \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd - be)^2 \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\ &\quad - \frac{\sqrt{\pi} e^2 \sin\left(2a - \frac{b^2}{2c}\right) \operatorname{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ &\quad - \frac{\sqrt{\pi} e^2 \cos\left(2a - \frac{b^2}{2c}\right) \operatorname{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\ &\quad + \frac{e(2cd - be) \sin(2a + 2bx + 2cx^2)}{16c^2} \\ &\quad + \frac{e(d + ex) \sin(2a + 2bx + 2cx^2)}{8c} + \frac{(d + ex)^3}{6e} \end{aligned}$$

[In] `Int[(d + e*x)^2*Cos[a + b*x + c*x^2]^2, x]`

[Out] $(d + e*x)^3/(6*e) + ((2*c*d - b*e)^2*\operatorname{Sqrt}[\Pi]*\operatorname{Cos}[2*a - b^2/(2*c)]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[\Pi])])/(16*c^{(5/2)}) - (e^{2*}\operatorname{Sqrt}[\Pi]*\operatorname{Cos}[2*a - b^2/(2*c)]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[\Pi])])/(16*c^{(3/2)}) - (e^{2*}\operatorname{Sqrt}[\Pi]*\operatorname{FresnelC}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[\Pi])]*\operatorname{Sin}[2*a - b^2/(2*c)])/(16*c^{(3/2)}) - ((2*c*d - b*e)^2*\operatorname{Sqrt}[\Pi]*\operatorname{FresnelS}[(b + 2*c*x)/(\operatorname{Sqrt}[c]*\operatorname{Sqrt}[\Pi])]*\operatorname{Sin}[2*a - b^2/(2*c)])/(16*c^{(5/2)}) + (e*(2*c*d - b*e)*\operatorname{Sin}[2*a + 2*b*x + 2*c*x^2])/(16*c^2) + (e*(d + e*x)*\operatorname{Sin}[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Rule 3432

`Int[Sin[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simplify[(Sqrt[\Pi/2]/(f*Rt[d, 2]))*\operatorname{FresnelS}[\operatorname{Sqrt}[2/\Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Rule 3433

`Int[Cos[(d_.)*(e_.) + (f_.)*(x_.)^2], x_Symbol] :> Simplify[(Sqrt[\Pi/2]/(f*Rt[d, 2]))*\operatorname{FresnelC}[\operatorname{Sqrt}[2/\Pi]*Rt[d, 2]*(e + f*x)], x] /; FreeQ[{d, e, f}, x]`

Rule 3528

`Int[Sin[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] - Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && Neq[`

$b^2 - 4*a*c, 0]$

Rule 3529

```
Int[Cos[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] :> Dist[Cos[(b^2 - 4*a*c)/(4*c)], Int[Cos[(b + 2*c*x)^2/(4*c)], x], x] + Dist[Sin[(b^2 - 4*a*c)/(4*c)], Int[Sin[(b + 2*c*x)^2/(4*c)], x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 3543

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_)), x_Symbol] :> Simp[e*(Sin[a + b*x + c*x^2]/(2*c)), x] + Dist[(2*c*d - b*e)/(2*c), Int[Cos[a + b*x + c*x^2], x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0]
```

Rule 3545

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]*((d_.) + (e_.*)(x_)^(m_)), x_Symbol] :> Simp[e*(d + e*x)^(m - 1)*(Sin[a + b*x + c*x^2]/(2*c)), x] + (-Dist[(b*e - 2*c*d)/(2*c), Int[(d + e*x)^(m - 1)*Cos[a + b*x + c*x^2], x], x] - Dist[e^2*((m - 1)/(2*c)), Int[(d + e*x)^(m - 2)*Sin[a + b*x + c*x^2], x], x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[b*e - 2*c*d, 0] && GtQ[m, 1]
```

Rule 3549

```
Int[Cos[(a_.) + (b_.*)(x_) + (c_.*)(x_)^2]^(n_)*((d_.) + (e_.*)(x_)^(m_)), x_Symbol] :> Int[ExpandTrigReduce[(d + e*x)^m, Cos[a + b*x + c*x^2]^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && IGtQ[n, 1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2}(d + ex)^2 + \frac{1}{2}(d + ex)^2 \cos(2a + 2bx + 2cx^2) \right) dx \\ &= \frac{(d + ex)^3}{6e} + \frac{1}{2} \int (d + ex)^2 \cos(2a + 2bx + 2cx^2) dx \\ &= \frac{(d + ex)^3}{6e} + \frac{e(d + ex) \sin(2a + 2bx + 2cx^2)}{8c} - \frac{e^2 \int \sin(2a + 2bx + 2cx^2) dx}{8c} \\ &\quad + \frac{(2cd - be) \int (d + ex) \cos(2a + 2bx + 2cx^2) dx}{4c} \end{aligned}$$

$$\begin{aligned}
&= \frac{(d+ex)^3}{6e} + \frac{e(2cd-be) \sin(2a+2bx+2cx^2)}{16c^2} \\
&\quad + \frac{e(d+ex) \sin(2a+2bx+2cx^2)}{8c} + \frac{(2cd-be)^2 \int \cos(2a+2bx+2cx^2) dx}{8c^2} \\
&\quad - \frac{\left(e^2 \cos\left(2a-\frac{b^2}{2c}\right)\right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c} \\
&\quad - \frac{\left(e^2 \sin\left(2a-\frac{b^2}{2c}\right)\right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c} \\
&= \frac{(d+ex)^3}{6e} - \frac{e^2 \sqrt{\pi} \cos\left(2a-\frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} \\
&\quad - \frac{e^2 \sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a-\frac{b^2}{2c}\right)}{16c^{3/2}} \\
&\quad + \frac{e(2cd-be) \sin(2a+2bx+2cx^2)}{16c^2} + \frac{e(d+ex) \sin(2a+2bx+2cx^2)}{8c} \\
&\quad + \frac{\left((2cd-be)^2 \cos\left(2a-\frac{b^2}{2c}\right)\right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c^2} \\
&\quad - \frac{\left((2cd-be)^2 \sin\left(2a-\frac{b^2}{2c}\right)\right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{8c^2} \\
&= \frac{(d+ex)^3}{6e} + \frac{(2cd-be)^2 \sqrt{\pi} \cos\left(2a-\frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{5/2}} \\
&\quad - \frac{e^2 \sqrt{\pi} \cos\left(2a-\frac{b^2}{2c}\right) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{16c^{3/2}} - \frac{e^2 \sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a-\frac{b^2}{2c}\right)}{16c^{3/2}} \\
&\quad - \frac{(2cd-be)^2 \sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a-\frac{b^2}{2c}\right)}{16c^{5/2}} \\
&\quad + \frac{e(2cd-be) \sin(2a+2bx+2cx^2)}{16c^2} + \frac{e(d+ex) \sin(2a+2bx+2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.70 (sec), antiderivative size = 215, normalized size of antiderivative = 0.74

$$\begin{aligned}
&\int (d+ex)^2 \cos^2(a+bx+cx^2) dx \\
&= \frac{3\sqrt{\pi} \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left((-2cd+be)^2 \cos\left(2a-\frac{b^2}{2c}\right) - ce^2 \sin\left(2a-\frac{b^2}{2c}\right)\right) - 3\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \left(ce^2 \cos\left(2a-\frac{b^2}{2c}\right) + (-2cd+be)^2 \sin\left(2a-\frac{b^2}{2c}\right)\right)}{16c^2}
\end{aligned}$$

[In] Integrate[(d + e*x)^2*Cos[a + b*x + c*x^2]^2, x]

[Out]
$$\frac{(3\sqrt{\pi} * \text{FresnelC}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] * ((-2cd + be)^2 * \cos[2a - b^2/(2c)] - c e^{2\sqrt{\pi}} \sin[2a - b^2/(2c)]) - 3\sqrt{\pi} * \text{FresnelS}[(b + 2cx)/(\sqrt{c}\sqrt{\pi})] * (c e^{2\sqrt{\pi}} \cos[2a - b^2/(2c)] + (-2cd + be)^2 * \sin[2a - b^2/(2c)]) + \sqrt{c} * (8c^2 x * (3d^2 + 3de*x + e^2*x^2) + 3e * (4cd - be + 2c*x) * \sin[2(a + x*(b + cx))]))}{(48c^{5/2})}$$

Maple [A] (verified)

Time = 1.39 (sec), antiderivative size = 378, normalized size of antiderivative = 1.30

method	result
default	$\frac{e^2 x \sin(2cx^2 + 2bx + 2a)}{8c} - \frac{e^2 b \left(\frac{\sin(2cx^2 + 2bx + 2a)}{4c} - \frac{b\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c^{3/2}} \right)}{4c} - \frac{e^2 \sqrt{\pi} \left(\cos\left(\frac{-4ac-b^2}{2c}\right) C\left(\frac{2cx-b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac-b^2}{2c}\right) S\left(\frac{2cx-b}{\sqrt{c}\sqrt{\pi}}\right) \right)}{4c}$
risch	$\frac{dex^2}{2} + \frac{d^2x}{2} + \frac{e^2x^3}{6} + \frac{\operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}d^2e^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} + \frac{e^2\operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}b^2e^{-\frac{i(4ac-b^2)}{2c}}}{64\sqrt{ic}c^2} - \frac{ie^2\operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}b^2e^{-\frac{i(4ac-b^2)}{2c}}}{64\sqrt{ic}c^2}$

[In] `int((e*x+d)^2*cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/8 * e^2 * c * x * \sin(2cx^2 + 2bx + 2a) - 1/4 * e^2 * b / c * (1/4 * \sin(2cx^2 + 2bx + 2a) / c - 1/4 * b / c^{3/2} * \Pi^{1/2} * (\cos(1/2 * (-4ac + b^2) / c) * \text{FresnelC}((2cx + b) / c^{1/2}) / \Pi^{1/2}) + \sin(1/2 * (-4ac + b^2) / c) * \text{FresnelS}((2cx + b) / c^{1/2}) / \Pi^{1/2})) - \\ & 1/16 * e^2 / c^{3/2} * \Pi^{1/2} * (\cos(1/2 * (-4ac + b^2) / c) * \text{FresnelS}((2cx + b) / c^{1/2}) / \Pi^{1/2}) - \sin(1/2 * (-4ac + b^2) / c) * \text{FresnelC}((2cx + b) / c^{1/2}) / \Pi^{1/2}) + \\ & 1/4 * d * e / c * \sin(2cx^2 + 2bx + 2a) - 1/4 * d * e * b / c^{3/2} * \Pi^{1/2} * (\cos(1/2 * (-4ac + b^2) / c) * \text{FresnelC}((2cx + b) / c^{1/2}) / \Pi^{1/2}) + \sin(1/2 * (-4ac + b^2) / c) * \text{FresnelS}((2cx + b) / c^{1/2}) / \Pi^{1/2})) + 1/4 * \Pi^{1/2} / c^{1/2} * d^2 * (\cos(1/2 * (-4ac + b^2) / c) * \text{FresnelC}((2cx + b) / c^{1/2}) / \Pi^{1/2}) + \sin(1/2 * (-4ac + b^2) / c) * \text{FresnelS}((2cx + b) / c^{1/2}) / \Pi^{1/2})) + 1/2 * d * e * x^2 + 1/2 * d^2 * x + 1/6 * e^2 * x^3 \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec), antiderivative size = 257, normalized size of antiderivative = 0.88

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx$$

$$= \frac{8c^3e^2x^3 + 24c^3dex^2 + 24c^3d^2x + 6(2c^2e^2x + 4c^2de - bce^2)\cos(cx^2 + bx + a)\sin(cx^2 + bx + a) - 3(\pi ce^2x^2 + 6c^2dex^2 + 6c^2d^2x + 2(2c^2e^2x + 4c^2de - bce^2)\cos(cx^2 + bx + a)\sin(cx^2 + bx + a))}{16c^2}$$

[In] `integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")`

```
[Out] 1/48*(8*c^3*e^2*x^3 + 24*c^3*d*x^2 + 24*c^3*d^2*x + 6*(2*c^2*e^2*x + 4*c^2*d*e - b*c*e^2)*cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a) - 3*(pi*c*e^2*sin(-1/2*(b^2 - 4*a*c)/c) - pi*(4*c^2*d^2 - 4*b*c*d*e + b^2*e^2)*cos(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_cos((2*c*x + b)*sqrt(c/pi)/c) - 3*(pi*c*e^2*cos(-1/2*(b^2 - 4*a*c)/c) + pi*(4*c^2*d^2 - 4*b*c*d*e + b^2*e^2)*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt(c/pi)*fresnel_sin((2*c*x + b)*sqrt(c/pi)/c))/c^3
```

Sympy [F]

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \int (d + ex)^2 \cos^2(a + bx + cx^2) dx$$

```
[In] integrate((e*x+d)**2*cos(c*x**2+b*x+a)**2, x)
```

```
[Out] Integral((d + e*x)**2*cos(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.38 (sec), antiderivative size = 2360, normalized size of antiderivative = 8.11

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \text{Too large to display}$$

```
[In] integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2, x, algorithm="maxima")
```

```
[Out] -1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)*d^2/c^2 + 1/32*sqrt(2)*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*((-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c))*x + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*co
```

$$\begin{aligned}
& s(-1/2*(b^2 - 4*a*c)/c) + c*(e^{(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)} + \\
& e^{-(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}*\sin(-1/2*(b^2 - 4*a*c)/c))*sq \\
& rt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*d*e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2) \\
& /c)) - 1/384*sqrt(2)*(24*((((I - 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi))*(erf(s \\
& qrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*(-(I \\
& + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1) \\
& *sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*cos(-1/ \\
& 2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*c^3 + 2*((I - 1)*s \\
& qrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2) \\
& *gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*c^4)*sin(-1/2*(b^2 \\
& - 4*a*c)/c))*x^3 + 36*((((I - 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*(-(I + 1) \\
& *sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) + (I - 1)*s \\
& qrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*cos(-1/2*(b^2 \\
& - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^3*c^2 + 2*((I - 1)*s \\
& qrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*sqrt(2) \\
& *gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b*c^3)*sin(-1/2*(b^2 \\
& - 4*a*c)/c))*x^2 - 4*sqrt(2)*(8*c^4*x^3 - 3*b*c^2*(-I*e^{(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)}) + I*e^{(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)})*co \\
& s(-1/2*(b^2 - 4*a*c)/c) - 3*b*c^2*(e^{(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c)} + e^{-(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)})*sin(-1/2*(b^2 - 4*a*c)/c) \\
& *((4*c^2*x^2 + 4*b*c*x + b^2)/c)^{(3/2)} + 18*((((I - 1)*sqrt(2)*sqrt(pi))*(\\
& erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*s \\
& qrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c))*b^2*c^2)*cos(-1/2*(b^2 - 4*a*c)/c) + (((I + 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2) \\
& *sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^4*c + 2*((I - 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c) - (I + 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c))*b^2*c^2)*sin(-1/2*(b^2 - 4*a*c)/c))*x + 3*((((I - 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2) \\
& *sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^5 + 2*(-(I + 1)*sqrt(2)*gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c) + (I - 1)*sqrt(2)*gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2) \\
& /c))*b^3*c)*cos(-1/2*(b^2 - 4*a*c)/c) + 3*((((I + 1)*sqrt(2)*sqrt(pi))*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi) \\
& *(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^5 +
\end{aligned}$$

$$2*((I - 1)*\sqrt{2}*\gamma(3/2, 1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c) - (I + 1)*\sqrt{2}*\gamma(3/2, -1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*b^3*c)*\sin(-1/2*(b^2 - 4*a*c)/c))*e^2/(c^4*((4*c^2*x^2 + 4*b*c*x + b^2)/c)^(3/2))$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.37 (sec), antiderivative size = 292, normalized size of antiderivative = 1.00

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \frac{1}{6} e^2 x^3 + \frac{1}{2} d e x^2 + \frac{1}{2} d^2 x \\ - \frac{i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(2icx^2 + 2ibx + 2ia)} + \frac{\sqrt{\pi}(-4ic^2d^2 + 4ibcde - ib^2e^2 + ce^2)\operatorname{erf}\left(-\frac{1}{2}i\sqrt{c}(2x + \frac{b}{c})(\frac{ic}{|c|} + 1)\right)e^{\left(-\frac{ic}{|c|}(2x + \frac{b}{c})^2\right)}}{\sqrt{c}(\frac{ic}{|c|} + 1)}}{32c^2} \\ - \frac{-i (ce^2(2x + \frac{b}{c}) + 4cde - 2be^2)e^{(-2icx^2 - 2ibx - 2ia)} + \frac{\sqrt{\pi}(4ic^2d^2 - 4ibcde + ib^2e^2 + ce^2)\operatorname{erf}\left(\frac{1}{2}i\sqrt{c}(2x + \frac{b}{c})(-\frac{ic}{|c|} + 1)\right)e^{\left(-\frac{ic}{|c|}(2x + \frac{b}{c})^2\right)}}{\sqrt{c}(-\frac{ic}{|c|} + 1)}}{32c^2}$$

[In] integrate((e*x+d)^2*cos(c*x^2+b*x+a)^2,x, algorithm="giac")

[Out] $\frac{1}{6}e^2x^3 + \frac{1}{2}d^2x^2 + \frac{1}{2}dex^2 - \frac{1}{32}(I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^{(2*I*c*x^2 + 2*I*b*x + 2*I*a)} + \sqrt{\pi}*(-4*I*c^2*d^2 + 4*I*b*c*d*e - I*b^2*e^2 + c*e^2)*\operatorname{erf}\left(-\frac{1}{2}\sqrt{c}(2*x + b/c)(I*c/abs(c) + 1)\right)*e^{(-1/2*(I*b^2 - 4*I*a*c)/c)} / (\sqrt{c}*(I*c/abs(c) + 1))) / c^2 - \frac{1}{32}(-I*(c*e^2*(2*x + b/c) + 4*c*d*e - 2*b*e^2)*e^{(-2*I*c*x^2 - 2*I*b*x - 2*I*a)} + \sqrt{\pi}*(4*I*c^2*d^2 - 4*I*b*c*d*e + I*b^2*e^2 + c*e^2)*\operatorname{erf}\left(\frac{1}{2}\sqrt{c}(2*x + b/c)(-I*c/abs(c) + 1)\right)*e^{(-1/2*(-I*b^2 + 4*I*a*c)/c)} / (\sqrt{c}*(-I*c/abs(c) + 1))) / c^2$

Mupad [F(-1)]

Timed out.

$$\int (d + ex)^2 \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 (d + ex)^2 dx$$

[In] int(cos(a + b*x + c*x^2)^2*(d + e*x)^2,x)

[Out] int(cos(a + b*x + c*x^2)^2*(d + e*x)^2, x)

$$\mathbf{3.33} \quad \int (d + ex) \cos^2(a + bx + cx^2) dx$$

Optimal result	200
Rubi [A] (verified)	200
Mathematica [A] (verified)	202
Maple [A] (verified)	202
Fricas [A] (verification not implemented)	203
Sympy [F]	203
Maxima [C] (verification not implemented)	203
Giac [C] (verification not implemented)	204
Mupad [F(-1)]	205

Optimal result

Integrand size = 19, antiderivative size = 150

$$\begin{aligned} \int (d+ex) \cos^2(a+bx+cx^2) dx = & \frac{(d+ex)^2}{4e} + \frac{(2cd-be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & - \frac{(2cd-be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} \\ & + \frac{e \sin(2a + 2bx + 2cx^2)}{8c} \end{aligned}$$

```
[Out] 1/4*(e*x+d)^2/e+1/8*e*sin(2*c*x^2+2*b*x+2*a)/c+1/8*(-b*e+2*c*d)*cos(2*a-1/2
*b^2/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))*Pi^(1/2)/c^(3/2)-1/8*(-b*e+2*c
*d)*FresnelS((2*c*x+b)/c^(1/2)/Pi^(1/2))*sin(2*a-1/2*b^2/c)*Pi^(1/2)/c^(3/2)
)
```

Rubi [A] (verified)

Time = 0.11 (sec), antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.263, Rules used = {3549, 3543, 3529, 3433, 3432}

$$\begin{aligned} \int (d+ex) \cos^2(a+bx+cx^2) dx = & \frac{\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) (2cd-be) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & - \frac{\sqrt{\pi} \sin\left(2a - \frac{b^2}{2c}\right) (2cd-be) \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\ & + \frac{e \sin(2a + 2bx + 2cx^2)}{8c} + \frac{(d+ex)^2}{4e} \end{aligned}$$

[In] $\text{Int}[(d + e*x)*\cos[a + b*x + c*x^2]^2, x]$

[Out] $(d + e*x)^2/(4*e) + ((2*c*d - b*e)*\sqrt{\pi}*\cos[2*a - b^2/(2*c)]*\text{FresnelC}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})])/(8*c^{(3/2)}) - ((2*c*d - b*e)*\sqrt{\pi}*\text{FresnelS}[(b + 2*c*x)/(\sqrt{c}*\sqrt{\pi})]*\sin[2*a - b^2/(2*c)])/(8*c^{(3/2)}) + (e*\sin[2*a + 2*b*x + 2*c*x^2])/(8*c)$

Rule 3432

$\text{Int}[\sin[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f*\text{Rt}[d, 2])*\text{FresnelS}[\sqrt{2/\pi}*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3433

$\text{Int}[\cos[(d_.)*((e_.) + (f_.)*(x_))^2], x_{\text{Symbol}}] \rightarrow \text{Simp}[(\sqrt{\pi}/2)/(f*\text{Rt}[d, 2])*\text{FresnelC}[\sqrt{2/\pi}*\text{Rt}[d, 2]*(e + f*x)], x] /; \text{FreeQ}[\{d, e, f\}, x]$

Rule 3529

$\text{Int}[\cos[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2], x_{\text{Symbol}}] \rightarrow \text{Dist}[\cos[b^2 - 4*a*c]/(4*c)], \text{Int}[\cos[(b + 2*c*x)^2/(4*c)], x], x] + \text{Dist}[\sin[(b^2 - 4*a*c)/(4*c)], \text{Int}[\sin[(b + 2*c*x)^2/(4*c)], x], x] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 3543

$\text{Int}[\cos[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]*((d_.) + (e_.)*(x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[e*(\sin[a + b*x + c*x^2]/(2*c)), x] + \text{Dist}[(2*c*d - b*e)/(2*c), \text{Int}[\cos[a + b*x + c*x^2], x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{NeQ}[2*c*d - b*e, 0]$

Rule 3549

$\text{Int}[\cos[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{(n_*)}*((d_.) + (e_.)*(x_))^{(m_*)}, x_{\text{Symbol}}] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(d + e*x)^m, \cos[a + b*x + c*x^2]^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{IGtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2}(d + ex) + \frac{1}{2}(d + ex)\cos(2a + 2bx + 2cx^2) \right) dx \\ &= \frac{(d + ex)^2}{4e} + \frac{1}{2} \int (d + ex)\cos(2a + 2bx + 2cx^2) dx \\ &= \frac{(d + ex)^2}{4e} + \frac{e \sin(2a + 2bx + 2cx^2)}{8c} + \frac{(4cd - 2be) \int \cos(2a + 2bx + 2cx^2) dx}{8c} \end{aligned}$$

$$\begin{aligned}
&= \frac{(d+ex)^2}{4e} + \frac{e \sin(2a + 2bx + 2cx^2)}{8c} \\
&\quad + \frac{\left((2cd-be)\cos\left(2a - \frac{b^2}{2c}\right)\right) \int \cos\left(\frac{(2b+4cx)^2}{8c}\right) dx}{4c} \\
&\quad - \frac{\left((2cd-be)\sin\left(2a - \frac{b^2}{2c}\right)\right) \int \sin\left(\frac{(2b+4cx)^2}{8c}\right) dx}{4c} \\
&= \frac{(d+ex)^2}{4e} + \frac{(2cd-be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}} \\
&\quad - \frac{(2cd-be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right)}{8c^{3/2}} + \frac{e \sin(2a + 2bx + 2cx^2)}{8c}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.93

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \frac{(2cd - be)\sqrt{\pi} \cos\left(2a - \frac{b^2}{2c}\right) \text{FresnelC}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) - (2cd - be)\sqrt{\pi} \text{FresnelS}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right) \sin\left(2a - \frac{b^2}{2c}\right) + \sqrt{c}(2cx(2d - b^2) + 4e^2x^2 + 4ex^3 + 2dx^2 + 2dx + 2d) \text{FresnelI}\left(\frac{b+2cx}{\sqrt{c}\sqrt{\pi}}\right)}{8c^{3/2}}$$

[In] `Integrate[(d + e*x)*Cos[a + b*x + c*x^2]^2,x]`

```
[Out] ((2*c*d - b*e)*Sqrt[Pi]*Cos[2*a - b^2/(2*c)]*FresnelC[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])] - (2*c*d - b*e)*Sqrt[Pi]*FresnelS[(b + 2*c*x)/(Sqrt[c]*Sqrt[Pi])]*Sin[2*a - b^2/(2*c)] + Sqrt[c]*(2*c*x*(2*d + e*x) + e*Sin[2*(a + x*(b + c*x))]))/(8*c^(3/2))
```

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.13

method	result
default	$\frac{e \sin(2cx^2 + 2bx + 2a)}{8c} - \frac{eb\sqrt{\pi} \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{8c^{\frac{3}{2}}} + \frac{\sqrt{\pi} d \left(\cos\left(\frac{-4ac+b^2}{2c}\right) C\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right) + \sin\left(\frac{-4ac+b^2}{2c}\right) S\left(\frac{2cx+b}{\sqrt{c}\sqrt{\pi}}\right)\right)}{4\sqrt{c}}$
risch	$\frac{e x^2}{4} + \frac{dx}{2} + \frac{\operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}de^{-\frac{i(4ac-b^2)}{2c}}}{16\sqrt{ic}} - \frac{e\operatorname{erf}\left(\sqrt{2}\sqrt{ic}x + \frac{ib\sqrt{2}}{2\sqrt{ic}}\right)\sqrt{2}\sqrt{\pi}be^{-\frac{i(4ac-b^2)}{2c}}}{32\sqrt{ic}c} - \frac{\operatorname{erf}\left(-\sqrt{-2ic}x + \frac{i}{\sqrt{-2ic}}\right)}{8\sqrt{ic}}$

```
[In] int((e*x+d)*cos(c*x^2+b*x+a)^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/8*e*sin(2*c*x^2+2*b*x+2*a)/c-1/8*e*b/c^(3/2)*Pi^(1/2)*(cos(1/2*(-4*a*c+b^2)/c)*FresnelC((2*c*x+b)/c^(1/2)/Pi^(1/2))+sin(1/2*(-4*a*c+b^2)/c)*FresnelS
```

$$\left(\frac{((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)})+1/4*\text{Pi}^{(1/2)}/c^{(1/2)}*d*(\cos(1/2*(-4*a*c+b^2)/c)*\text{FresnelC}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)})+\sin(1/2*(-4*a*c+b^2)/c)*\text{FresnelS}((2*c*x+b)/c^{(1/2)}/\text{Pi}^{(1/2)}))+1/2*d*x+1/4*e*x^2}{\text{None}}$$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 149, normalized size of antiderivative = 0.99

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \frac{2 c^2 e x^2 + \pi (2 cd - be) \sqrt{\frac{c}{\pi}} \cos\left(-\frac{b^2 - 4 ac}{2 c}\right) C\left(\frac{(2 cx + b)\sqrt{\frac{c}{\pi}}}{c}\right) - \pi (2 cd - be) \sqrt{\frac{c}{\pi}} S\left(\frac{(2 cx + b)\sqrt{\frac{c}{\pi}}}{c}\right) \sin\left(-\frac{b^2 - 4 ac}{2 c}\right)}{8 c^2}$$

```
[In] integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="fricas")
[Out] 1/8*(2*c^2*e*x^2 + pi*(2*c*d - b*e)*sqrt(c/pi)*cos(-1/2*(b^2 - 4*a*c)/c)*fr
esnel_cos((2*c*x + b)*sqrt(c/pi)/c) - pi*(2*c*d - b*e)*sqrt(c/pi)*fresnel_s
in((2*c*x + b)*sqrt(c/pi)/c)*sin(-1/2*(b^2 - 4*a*c)/c) + 4*c^2*d*x + 2*c*e*
cos(c*x^2 + b*x + a)*sin(c*x^2 + b*x + a))/c^2
```

Sympy [F]

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \int (d + ex) \cos^2(a + bx + cx^2) dx$$

```
[In] integrate((e*x+d)*cos(c*x**2+b*x+a)**2,x)
[Out] Integral((d + e*x)*cos(a + b*x + c*x**2)**2, x)
```

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.65 (sec) , antiderivative size = 737, normalized size of antiderivative = 4.91

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \frac{- \left(4^{\frac{1}{4}} \sqrt{2} \sqrt{\pi} \left(\left((i - 1) \cos\left(-\frac{b^2 - 4 ac}{2 c}\right) + (i + 1) \sin\left(-\frac{b^2 - 4 ac}{2 c}\right)\right) \operatorname{erf}\left(\frac{2 i c x + i b}{\sqrt{2} i c}\right) + \left((i + 1) \cos\left(-\frac{b^2 - 4 ac}{2 c}\right) - (i - 1) \sin\left(-\frac{b^2 - 4 ac}{2 c}\right)\right) \operatorname{erf}\left(\frac{2 i c x - i b}{\sqrt{2} i c}\right)\right) + \sqrt{2} \left(\left((i - 1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{\frac{1}{2}} \sqrt{\frac{4 i c^2 x^2 + 4 i b c x + i b^2}{c}}\right) - 1\right) - (i + 1) \sqrt{2} \sqrt{\pi} \left(\operatorname{erf}\left(\sqrt{\frac{1}{2}} \sqrt{-\frac{4 i c^2 x^2 + 4 i b c x + i b^2}{c}}\right) - 1\right)\right)}{32 c^2}$$

```
[In] integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="maxima")
[Out] -1/32*(4^(1/4)*sqrt(2)*sqrt(pi)*(((I - 1)*cos(-1/2*(b^2 - 4*a*c)/c) + (I + 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(2*I*c)) + ((I + 1)*c*cos(-1/2*(b^2 - 4*a*c)/c) + (I - 1)*sin(-1/2*(b^2 - 4*a*c)/c))*erf((2*I*c*x + I*b)/sqrt(-2*I*c)))*c^(3/2) - 16*c^2*x)*d/c^2 + 1/64*sqrt(2)*(((I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*cos(-1/2*(b^2 - 4*a*c)/c) + ((I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) - (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b^2*sin(-1/2*(b^2 - 4*a*c)/c) - 2*(-(I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*cos(-1/2*(b^2 - 4*a*c)/c) + (-(I + 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt((4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1) + (I - 1)*sqrt(2)*sqrt(pi)*(erf(sqrt(1/2)*sqrt(-(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - 1))*b*c*sin(-1/2*(b^2 - 4*a*c)/c)*x + 2*sqrt(2)*(4*c^2*x^2 - c*(I*e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c)) - I*e^(-1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*cos(-1/2*(b^2 - 4*a*c)/c) + c*(e^(1/2*(4*I*c^2*x^2 + 4*I*b*c*x + I*b^2)/c))*sin(-1/2*(b^2 - 4*a*c)/c))*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))*e/(c^2*sqrt((4*c^2*x^2 + 4*b*c*x + b^2)/c))
```

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.35 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.30

$$\int (d + ex) \cos^2(a + bx + cx^2) dx$$

$$= \frac{1}{4} ex^2 + \frac{1}{2} dx - \frac{i ee^{(2i cx^2 + 2i bx + 2i a)} - \frac{\sqrt{\pi}(2i cd - i be) \operatorname{erf}\left(-\frac{1}{2}i\sqrt{c}\left(2x + \frac{b}{c}\right)\left(\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{ib^2 - 4ia}{2c}\right)}}{\sqrt{c}\left(\frac{ic}{|c|} + 1\right)}}{16c}$$

$$- \frac{-i ee^{(-2i cx^2 - 2i bx - 2i a)} - \frac{\sqrt{\pi}(-2i cd + i be) \operatorname{erf}\left(\frac{1}{2}i\sqrt{c}\left(2x + \frac{b}{c}\right)\left(-\frac{ic}{|c|} + 1\right)\right)e^{\left(-\frac{-ib^2 + 4ia}{2c}\right)}}{\sqrt{c}\left(-\frac{ic}{|c|} + 1\right)}}{16c}$$

```
[In] integrate((e*x+d)*cos(c*x^2+b*x+a)^2,x, algorithm="giac")
[Out] 1/4*e*x^2 + 1/2*d*x - 1/16*(I*e*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a) - sqrt(pi)*(2*I*c*d - I*b*e)*erf(-1/2*I*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)))/c - 1/16*(-I*e*e^(-2*I*c*x^2 - 2*I*b*x - 2*I*a) - sqrt(pi)*(-2*I*c*d + I*b*e)*erf(1/2*I*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))/c) + 1/16*(I*e*e^(2*I*c*x^2 + 2*I*b*x + 2*I*a) - sqrt(pi)*(2*I*c*d - I*b*e)*erf(-1/2*I*sqrt(c)*(2*x + b/c)*(I*c/abs(c) + 1))*e^(-1/2*(I*b^2 - 4*I*a*c)/c)/(sqrt(c)*(I*c/abs(c) + 1)))/c
```

+ b/c)*(-I*c/abs(c) + 1))*e^(-1/2*(-I*b^2 + 4*I*a*c)/c)/(sqrt(c)*(-I*c/abs(c) + 1)))/c

Mupad [F(-1)]

Timed out.

$$\int (d + ex) \cos^2(a + bx + cx^2) dx = \int \cos(cx^2 + bx + a)^2 (d + ex) dx$$

[In] int(cos(a + b*x + c*x^2)^2*(d + e*x),x)

[Out] int(cos(a + b*x + c*x^2)^2*(d + e*x), x)

3.34 $\int \frac{\cos^2(a+bx+cx^2)}{d+ex} dx$

Optimal result	206
Rubi [N/A]	206
Mathematica [N/A]	207
Maple [N/A] (verified)	207
Fricas [N/A]	207
Sympy [N/A]	207
Maxima [N/A]	208
Giac [N/A]	208
Mupad [N/A]	208

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \frac{\log(d + ex)}{2e} + \frac{1}{2} \text{Int}\left(\frac{\cos(2a + 2bx + 2cx^2)}{d + ex}, x\right)$$

[Out] $1/2*\ln(e*x+d)/e+1/2*\text{Unintegrable}(\cos(2*c*x^2+2*b*x+2*a)/(e*x+d),x)$

Rubi [N/A]

Not integrable

Time = 0.05 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx$$

[In] $\text{Int}[\cos[a + b*x + c*x^2]^2/(d + e*x), x]$

[Out] $\text{Log}[d + e*x]/(2*e) + \text{Defer}[\text{Int}][\cos[2*a + 2*b*x + 2*c*x^2]/(d + e*x), x]/2$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{2(d + ex)} + \frac{\cos(2a + 2bx + 2cx^2)}{2(d + ex)} \right) dx \\ &= \frac{\log(d + ex)}{2e} + \frac{1}{2} \int \frac{\cos(2a + 2bx + 2cx^2)}{d + ex} dx \end{aligned}$$

Mathematica [N/A]

Not integrable

Time = 4.47 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx$$

[In] `Integrate[Cos[a + b*x + c*x^2]^2/(d + e*x), x]`

[Out] `Integrate[Cos[a + b*x + c*x^2]^2/(d + e*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.54 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int \frac{\cos^2(cx^2 + bx + a)}{ex + d} dx$$

[In] `int(cos(c*x^2+b*x+a)^2/(e*x+d), x)`

[Out] `int(cos(c*x^2+b*x+a)^2/(e*x+d), x)`

Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/(e*x+d), x, algorithm="fricas")`

[Out] `integral(cos(c*x^2 + b*x + a)^2/(e*x + d), x)`

Sympy [N/A]

Not integrable

Time = 41.99 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx$$

[In] `integrate(cos(c*x**2+b*x+a)**2/(e*x+d), x)`

[Out] `Integral(cos(a + b*x + c*x**2)**2/(d + e*x), x)`

Maxima [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 120, normalized size of antiderivative = 5.71

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="maxima")`

[Out] `-1/2*(2*e*integrate(-1/4*(cos(2*c*x^2 + 2*b*x)*cos(2*a) - sin(2*c*x^2 + 2*b*x)*sin(2*a))/((cos(2*a)^2 + sin(2*a)^2)*e*x + (cos(2*a)^2 + sin(2*a)^2)*d), x) - 2*e*integrate(1/4*cos(2*c*x^2 + 2*b*x + 2*a)/(e*x + d), x) - log(e*x + d))/e`

Giac [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{ex + d} dx$$

[In] `integrate(cos(c*x^2+b*x+a)^2/(e*x+d),x, algorithm="giac")`

[Out] `integrate(cos(c*x^2 + b*x + a)^2/(e*x + d), x)`

Mupad [N/A]

Not integrable

Time = 14.12 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int \frac{\cos^2(a + bx + cx^2)}{d + ex} dx = \int \frac{\cos(cx^2 + bx + a)^2}{d + ex} dx$$

[In] `int(cos(a + b*x + c*x^2)^2/(d + e*x),x)`

[Out] `int(cos(a + b*x + c*x^2)^2/(d + e*x), x)`

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	209
--	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                                         Small rewrite of logic in main function to make it*)
(*                                         match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal}
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A",""}
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count is different."}
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)
    finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
  ]
]
,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>ToString[Order[result]]},
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];
finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn] === Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]] === Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
              1,
              Max[ExpnType[expn[[1]]], 2]],
            Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
        If[Head[expn] === Plus || Head[expn] === Times,
          Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
              If[HypergeometricFunctionQ[Head[expn]],
                Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
                If[AppellFunctionQ[Head[expn]],
                  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
                  If[Head[expn] === RootSum,
                    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
                    If[Head[expn] === Integrate || Head[expn] === Int,
                      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
                      9]]]]]]]]]
]

ElementaryFunctionQ[func_] :=
  MemberQ[{  

    Exp, Log,  

    Sin, Cos, Tan, Cot, Sec, Csc,  

    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
  }]

```

```

Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]

```

```

SpecialFunctionQ[func_] :=
MemberQ[{{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func}]

```

```

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (",

```

```

        convert(leaf_count_optimal,string), " ) = ",convert(2*leaf_
    end if
else #result contains complex but optimal is not
if debug then
    print("result contains complex but optimal is not");
fi;
return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
# this assumes optimal do not as well. No check is needed here.
if debug then
    print("result do not contain complex, this assumes optimal do not as well")
fi;
if leaf_count_result<=2*leaf_count_optimal then
if debug then
    print("leaf_count_result<=2*leaf_count_optimal");
fi;
return "A"," ";
else
if debug then
    print("leaf_count_result>2*leaf_count_optimal");
fi;
return "B",cat("Leaf count of result is larger than twice the leaf count of op-
    convert(leaf_count_result,string)," vs. $2(", 
    convert(leaf_count_optimal,string),")=",convert(2*leaf_count_
fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
if debug then
    print("ExpnType(result) > ExpnType(optimal)");
fi;
return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:
```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+``') or type(expn,'`*``') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except:
        return False
```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0]))  #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow):  #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0])  #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0]))  #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+`) or type(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0]))  #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1)  #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sageMath")
    #print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count(result))-str(leaf_count(optimal))
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType(result))-str(ExpnType(optimal))

```

```
#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#          Albert Rich to use with Sagemath. This is used to
#          grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#          'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#          issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:  #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False
```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than optimal"
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```