

Computer Algebra Independent Integration Tests

Summer 2023 edition

4-Trig-functions/4.3-Tangent/99-4.3.10-c+d-x-^m-a+b-tan-^n

Nasser M. Abbasi

September 5, 2023 Compiled on September 5, 2023 at 9:17pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	41
4	Appendix	433

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [63]. This is test number [99].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Mathematica	100.00 (63)	0.00 (0)
Fricas	100.00 (63)	0.00 (0)
Rubi	98.41 (62)	1.59 (1)
Maple	92.06 (58)	7.94 (5)
Maxima	77.78 (49)	22.22 (14)
Giac	55.56 (35)	44.44 (28)
Mupad	50.79 (32)	49.21 (31)
Sympy	44.44 (28)	55.56 (35)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	69.841	0.000	0.000	30.159
Mathematica	58.730	12.698	0.000	28.571
Maple	44.444	19.048	0.000	36.508
Fricas	39.683	31.746	0.000	28.571
Giac	19.048	7.937	0.000	73.016
Sympy	15.873	0.000	0.000	84.127
Maxima	12.698	36.508	0.000	50.794
Mupad	0.000	22.222	0.000	77.778

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Rubi	1	100.00	0.00	0.00
Maple	5	100.00	0.00	0.00
Maxima	14	35.71	0.00	64.29
Giac	28	100.00	0.00	0.00
Mupad	31	0.00	100.00	0.00
Sympy	35	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Fricas	0.25
Rubi	0.41
Maple	0.64
Sympy	1.09
Maxima	1.89
Giac	3.52
Mupad	3.64
Mathematica	5.07

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	91.44	1.30	22.00	1.11
Sympy	143.93	1.25	19.00	1.00
Rubi	172.63	1.00	126.50	1.00
Fricas	275.24	1.77	130.00	1.58
Giac	295.46	1.58	23.00	1.10
Mathematica	300.63	1.46	171.00	1.17
Maple	433.60	1.73	118.00	1.00
Maxima	910.49	15.56	424.00	4.78

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

$\{4, 5, 9, 10, 14, 15, 34, 35, 42, 43, 47, 48, 52, 53, 57, 58, 62, 63\}$

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {44, 45, 49, 50, 59}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```

x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)

```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```

try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1

```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```

integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)

```

Which gives $\sin(x)^{2/2}$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	38

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

B grade { }

C grade { }

F normal fail { 17 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 6, 8, 11, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 40, 41, 46, 51, 54, 55, 56, 60 }

B grade { 7, 39, 44, 45, 49, 50, 59, 61 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 41, 46, 51 }

B grade { 39, 40, 44, 45, 49, 50, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { 16, 17, 36, 37, 38 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 8, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 36, 37, 38, 46, 50, 51 }

B grade { 1, 2, 3, 6, 7, 11, 12, 13, 39, 40, 41, 44, 45, 49, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 21, 22, 23, 27, 28, 32, 33, 41 }

B grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

C grade { }

F normal fail { 16, 17, 36, 37, 38 }

F(-1) timeout fail { }

F(-2) exception fail { 18, 19, 20, 24, 25, 26, 29, 30, 31 }

Giac

A grade { 18, 19, 20, 21, 24, 25, 26, 27, 29, 30, 31, 32 }

B grade { 8, 22, 23, 28, 33 }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 3, 8, 16, 17, 18, 19, 20, 24, 25, 26, 29, 30, 31, 41 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 6, 7, 11, 12, 13, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

F(-2) exception fail { }

Sympy

A grade { 8, 18, 19, 20, 24, 25, 26, 29, 30, 31 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 21, 22, 23, 27, 28, 32, 33, 36, 37, 38, 39, 40, 41, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 61 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	106	106	106	125	243	286	0	0	0
N.S.	1	1.00	1.00	1.18	2.29	2.70	0.00	0.00	0.00
time (sec)	N/A	0.183	0.013	0.490	0.600	0.280	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	77	77	77	103	163	200	0	0	0
N.S.	1	1.00	1.00	1.34	2.12	2.60	0.00	0.00	0.00
time (sec)	N/A	0.166	0.009	0.429	0.682	0.259	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	54	78	92	122	0	0	129
N.S.	1	1.00	1.00	1.44	1.70	2.26	0.00	0.00	2.39
time (sec)	N/A	0.095	0.006	0.438	0.397	0.242	0.000	0.000	3.346

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	8	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	0.80	1.20	1.20
time (sec)	N/A	0.018	1.555	0.240	0.446	0.253	0.315	0.340	3.237

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	12	12	10	12	12
N.S.	1	1.00	1.20	1.00	1.20	1.20	1.00	1.20	1.20
time (sec)	N/A	0.018	2.420	0.205	0.452	0.237	0.281	0.337	3.065

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	115	133	639	225	0	0	0
N.S.	1	1.00	1.17	1.36	6.52	2.30	0.00	0.00	0.00
time (sec)	N/A	0.198	0.686	0.632	0.397	0.256	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	73	73	160	108	257	144	0	0	0
N.S.	1	1.00	2.19	1.48	3.52	1.97	0.00	0.00	0.00
time (sec)	N/A	0.146	5.051	0.550	0.631	0.247	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	30	30	43	34	214	38	41	162	35
N.S.	1	1.00	1.43	1.13	7.13	1.27	1.37	5.40	1.17
time (sec)	N/A	0.031	0.146	0.459	0.524	0.251	0.216	0.604	0.163

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	243	14	10	14	14
N.S.	1	1.00	1.17	1.00	20.25	1.17	0.83	1.17	1.17
time (sec)	N/A	0.034	3.216	0.339	0.956	0.242	0.309	0.633	3.566

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	250	14	12	14	14
N.S.	1	1.00	1.17	1.00	20.83	1.17	1.00	1.17	1.17
time (sec)	N/A	0.035	2.442	0.311	0.447	0.237	0.287	0.671	2.928

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	205	370	251	1205	344	0	0	0
N.S.	1	1.00	1.80	1.22	5.88	1.68	0.00	0.00	0.00
time (sec)	N/A	0.358	6.536	0.355	0.471	0.274	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	128	128	172	180	736	240	0	0	0
N.S.	1	1.00	1.34	1.41	5.75	1.88	0.00	0.00	0.00
time (sec)	N/A	0.223	2.281	0.299	0.659	0.261	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	171	122	386	146	0	0	0
N.S.	1	1.00	1.90	1.36	4.29	1.62	0.00	0.00	0.00
time (sec)	N/A	0.136	4.577	0.271	0.783	0.257	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	534	14	10	14	14
N.S.	1	1.00	1.17	1.00	44.50	1.17	0.83	1.17	1.17
time (sec)	N/A	0.032	4.854	0.174	0.913	0.242	0.334	0.921	3.176

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	536	14	12	14	14
N.S.	1	1.00	1.17	1.00	44.67	1.17	1.00	1.17	1.17
time (sec)	N/A	0.035	2.584	0.188	0.713	0.242	0.284	0.962	2.826

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	18	0	0	16	0	0	50
N.S.	1	1.00	1.00	0.00	0.00	0.89	0.00	0.00	2.78
time (sec)	N/A	3.196	1.811	0.000	0.000	0.262	0.000	0.000	3.610

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	F	A	F	F	A	F	F	B
verified	N/A	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	17	0	17	0	0	15	0	0	45
N.S.	1	0.00	1.00	0.00	0.00	0.88	0.00	0.00	2.65
time (sec)	N/A	0.000	0.916	0.000	0.000	0.250	0.000	0.000	5.453

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	189	189	278	170	0	164	258	188	423
N.S.	1	1.00	1.47	0.90	0.00	0.87	1.37	0.99	2.24
time (sec)	N/A	0.217	0.802	0.674	0.000	0.255	0.263	0.382	4.118

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	137	178	108	0	105	165	119	241
N.S.	1	1.00	1.30	0.79	0.00	0.77	1.20	0.87	1.76
time (sec)	N/A	0.137	0.493	0.520	0.000	0.251	0.190	0.359	3.706

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	96	50	0	55	92	62	105
N.S.	1	1.00	1.14	0.60	0.00	0.65	1.10	0.74	1.25
time (sec)	N/A	0.059	0.741	0.471	0.000	0.243	0.148	0.353	3.446

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	161	161	166	65	112	51	0	138	0
N.S.	1	1.00	1.03	0.40	0.70	0.32	0.00	0.86	0.00
time (sec)	N/A	0.316	0.610	0.601	0.282	0.252	0.000	0.371	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	168	168	224	96	119	83	0	1013	0
N.S.	1	1.00	1.33	0.57	0.71	0.49	0.00	6.03	0.00
time (sec)	N/A	0.280	0.767	0.508	0.585	0.259	0.000	2.299	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	227	227	285	216	158	130	0	532	0
N.S.	1	1.00	1.26	0.95	0.70	0.57	0.00	2.34	0.00
time (sec)	N/A	0.376	1.000	0.569	0.595	0.250	0.000	0.442	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	270	473	283	0	267	665	368	289
N.S.	1	1.00	1.75	1.05	0.00	0.99	2.46	1.36	1.07
time (sec)	N/A	0.334	2.541	0.746	0.000	0.251	0.415	0.434	4.301

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	202	202	282	173	0	163	418	217	183
N.S.	1	1.00	1.40	0.86	0.00	0.81	2.07	1.07	0.91
time (sec)	N/A	0.235	1.612	0.780	0.000	0.245	0.329	0.449	3.651

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	151	151	130	82	0	79	226	101	103
N.S.	1	1.00	0.86	0.54	0.00	0.52	1.50	0.67	0.68
time (sec)	N/A	0.185	1.070	0.801	0.000	0.248	0.258	0.416	3.130

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	305	305	211	114	194	84	0	404	0
N.S.	1	1.00	0.69	0.37	0.64	0.28	0.00	1.32	0.00
time (sec)	N/A	0.941	0.848	0.871	0.304	0.244	0.000	0.417	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	436	436	467	175	211	142	0	1967	0
N.S.	1	1.00	1.07	0.40	0.48	0.33	0.00	4.51	0.00
time (sec)	N/A	0.828	1.672	0.892	0.480	0.278	0.000	11.298	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	396	396	667	396	0	370	945	548	411
N.S.	1	1.00	1.68	1.00	0.00	0.93	2.39	1.38	1.04
time (sec)	N/A	0.466	3.764	1.209	0.000	0.245	0.538	0.580	4.249

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	294	294	405	238	0	221	588	315	263
N.S.	1	1.00	1.38	0.81	0.00	0.75	2.00	1.07	0.89
time (sec)	N/A	0.339	2.736	1.082	0.000	0.248	0.504	0.619	3.949

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	209	205	114	0	105	311	142	146
N.S.	1	1.00	0.98	0.55	0.00	0.50	1.49	0.68	0.70
time (sec)	N/A	0.282	1.516	0.884	0.000	0.246	0.343	0.559	3.940

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	449	449	336	163	276	117	0	810	0
N.S.	1	1.00	0.75	0.36	0.61	0.26	0.00	1.80	0.00
time (sec)	N/A	2.598	1.137	0.934	0.315	0.255	0.000	0.584	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	712	712	833	254	297	197	0	2915	0
N.S.	1	1.00	1.17	0.36	0.42	0.28	0.00	4.09	0.00
time (sec)	N/A	2.118	2.853	1.063	0.431	0.247	0.000	30.478	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	23	23	25	21	520	115	51	23	24
N.S.	1	1.00	1.09	0.91	22.61	5.00	2.22	1.00	1.04
time (sec)	N/A	0.058	42.842	0.192	1.833	0.250	5.317	0.641	3.372

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	21	21	23	19	79	34	29	21	22
N.S.	1	1.00	1.10	0.90	3.76	1.62	1.38	1.00	1.05
time (sec)	N/A	0.033	9.865	0.163	0.500	0.242	2.400	0.357	3.337

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	98	98	133	0	0	85	0	0	0
N.S.	1	1.00	1.36	0.00	0.00	0.87	0.00	0.00	0.00
time (sec)	N/A	0.147	1.771	0.000	0.000	0.085	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	171	171	192	0	0	144	0	0	0
N.S.	1	1.00	1.12	0.00	0.00	0.84	0.00	0.00	0.00
time (sec)	N/A	0.229	4.489	0.000	0.000	0.087	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	251	251	269	0	0	198	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	0.79	0.00	0.00	0.00
time (sec)	N/A	0.291	13.029	0.000	0.000	0.087	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	152	152	342	500	672	500	0	0	0
N.S.	1	1.00	2.25	3.29	4.42	3.29	0.00	0.00	0.00
time (sec)	N/A	0.292	0.146	0.641	0.715	0.274	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	115	115	191	314	373	317	0	0	0
N.S.	1	1.00	1.66	2.73	3.24	2.76	0.00	0.00	0.00
time (sec)	N/A	0.242	0.080	0.583	0.520	0.261	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	87	143	130	160	0	0	161
N.S.	1	1.00	1.04	1.70	1.55	1.90	0.00	0.00	1.92
time (sec)	N/A	0.141	0.013	0.547	0.377	0.260	0.000	0.000	3.597

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	86	20	15	20	20
N.S.	1	1.00	1.11	1.00	4.78	1.11	0.83	1.11	1.11
time (sec)	N/A	0.033	1.914	0.217	0.440	0.243	0.663	0.389	3.534

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	142	31	17	20	20
N.S.	1	1.00	1.11	1.00	7.89	1.72	0.94	1.11	1.11
time (sec)	N/A	0.030	5.096	0.220	0.463	0.242	2.494	4.371	3.676

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	300	300	1337	952	2525	759	0	0	0
N.S.	1	1.00	4.46	3.17	8.42	2.53	0.00	0.00	0.00
time (sec)	N/A	0.595	7.326	1.603	1.726	0.276	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	229	229	649	575	1263	450	0	0	0
N.S.	1	1.00	2.83	2.51	5.52	1.97	0.00	0.00	0.00
time (sec)	N/A	0.438	6.833	1.389	0.912	0.264	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	136	136	200	238	527	219	0	0	0
N.S.	1	1.00	1.47	1.75	3.88	1.61	0.00	0.00	0.00
time (sec)	N/A	0.197	7.410	1.308	0.409	0.258	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	491	36	17	22	22
N.S.	1	1.00	1.10	1.00	24.55	1.80	0.85	1.10	1.10
time (sec)	N/A	0.061	19.689	0.366	0.827	0.252	1.091	0.565	4.172

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	609	47	19	22	22
N.S.	1	1.00	1.10	1.00	30.45	2.35	0.95	1.10	1.10
time (sec)	N/A	0.052	14.033	0.388	1.683	0.254	2.107	15.426	3.644

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	612	612	2594	1930	6861	1177	0	0	0
N.S.	1	1.00	4.24	3.15	11.21	1.92	0.00	0.00	0.00
time (sec)	N/A	1.118	7.568	1.210	10.883	0.268	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	A	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	436	436	1846	1138	3407	688	0	0	0
N.S.	1	1.00	4.23	2.61	7.81	1.58	0.00	0.00	0.00
time (sec)	N/A	0.726	7.164	1.058	1.972	0.290	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	277	277	493	1319	327	0	0	0
N.S.	1	1.00	1.00	1.78	4.76	1.18	0.00	0.00	0.00
time (sec)	N/A	0.368	8.323	0.915	0.670	0.266	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1935	52	17	22	22
N.S.	1	1.00	1.10	1.00	96.75	2.60	0.85	1.10	1.10
time (sec)	N/A	0.058	14.685	0.621	3.085	0.251	1.330	1.075	3.911

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	2194	63	19	22	22
N.S.	1	1.00	1.10	1.00	109.70	3.15	0.95	1.10	1.10
time (sec)	N/A	0.060	17.927	0.760	9.084	0.253	2.439	18.056	4.929

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	243	243	297	1468	983	1183	0	0	0
N.S.	1	1.00	1.22	6.04	4.05	4.87	0.00	0.00	0.00
time (sec)	N/A	0.393	1.676	0.780	0.568	0.282	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	181	236	940	715	834	0	0	0
N.S.	1	1.00	1.30	5.19	3.95	4.61	0.00	0.00	0.00
time (sec)	N/A	0.323	1.426	0.693	0.553	0.272	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	177	462	399	527	0	0	0
N.S.	1	1.00	1.42	3.70	3.19	4.22	0.00	0.00	0.00
time (sec)	N/A	0.182	1.560	0.637	0.603	0.272	0.000	0.000	0.000

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	279	27	17	22	22
N.S.	1	1.00	1.10	1.00	13.95	1.35	0.85	1.10	1.10
time (sec)	N/A	0.071	2.414	0.424	0.890	0.239	1.022	0.417	3.963

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	424	51	19	22	22
N.S.	1	1.00	1.10	1.00	21.20	2.55	0.95	1.10	1.10
time (sec)	N/A	0.069	4.558	0.317	1.726	0.252	1.759	4.307	4.091

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	848	848	1724	5959	4631	2537	0	0	0
N.S.	1	1.00	2.03	7.03	5.46	2.99	0.00	0.00	0.00
time (sec)	N/A	2.306	9.637	0.967	2.409	0.327	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	654	654	761	3901	2530	1576	0	0	0
N.S.	1	1.00	1.16	5.96	3.87	2.41	0.00	0.00	0.00
time (sec)	N/A	1.599	7.529	0.969	1.023	0.286	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	214	214	745	2001	1156	843	0	0	0
N.S.	1	1.00	3.48	9.35	5.40	3.94	0.00	0.00	0.00
time (sec)	N/A	0.302	7.889	0.862	0.705	0.268	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1498	55	19	22	22
N.S.	1	1.00	1.10	1.00	74.90	2.75	0.95	1.10	1.10
time (sec)	N/A	0.067	13.937	0.550	8.731	0.252	1.670	0.541	4.538

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1977	96	20	22	22
N.S.	1	1.00	1.10	1.00	98.85	4.80	1.00	1.10	1.10
time (sec)	N/A	0.065	15.452	0.652	27.223	0.248	3.181	22.558	3.988

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [11] had the largest ratio of [.83330000000000041]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	6	1.00	10	0.600
2	A	5	5	1.00	10	0.500
3	A	4	4	1.00	8	0.500
4	N/A	0	0	1.00	10	0.000
5	N/A	0	0	1.00	10	0.000
6	A	7	7	1.00	12	0.583
7	A	6	6	1.00	12	0.500
8	A	3	3	1.00	10	0.300
9	N/A	0	0	1.00	12	0.000
10	N/A	0	0	1.00	12	0.000
11	A	13	10	1.00	12	0.833
12	A	9	8	1.00	12	0.667
13	A	7	7	1.00	10	0.700
14	N/A	0	0	1.00	12	0.000
15	N/A	0	0	1.00	12	0.000
16	A	76	12	1.00	45	0.267
17	F	0	0	N/A	0.000	N/A
18	A	5	3	1.00	23	0.130
19	A	4	3	1.00	23	0.130
20	A	3	3	1.00	21	0.143
21	A	7	4	1.00	23	0.174
22	A	7	4	1.00	23	0.174
23	A	8	5	1.00	23	0.217

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
24	A	10	3	1.00	23	0.130
25	A	8	3	1.00	23	0.130
26	A	7	3	1.00	21	0.143
27	A	21	5	1.00	23	0.217
28	A	24	7	1.00	23	0.304
29	A	14	3	1.00	23	0.130
30	A	11	3	1.00	23	0.130
31	A	11	3	1.00	21	0.143
32	A	53	7	1.00	23	0.304
33	A	60	9	1.00	23	0.391
34	N/A	0	0	1.00	23	0.000
35	N/A	0	0	1.00	21	0.000
36	A	2	2	1.00	23	0.087
37	A	4	2	1.00	23	0.087
38	A	5	2	1.00	23	0.087
39	A	8	7	1.00	18	0.389
40	A	7	6	1.00	18	0.333
41	A	6	5	1.00	16	0.312
42	N/A	0	0	1.00	18	0.000
43	N/A	0	0	1.00	18	0.000
44	A	15	9	1.00	20	0.450
45	A	13	10	1.00	20	0.500
46	A	9	7	1.00	18	0.389
47	N/A	0	0	1.00	20	0.000
48	N/A	0	0	1.00	20	0.000
49	A	28	11	1.00	20	0.550
50	A	22	11	1.00	20	0.550
51	A	16	9	1.00	18	0.500
52	N/A	0	0	1.00	20	0.000
53	N/A	0	0	1.00	20	0.000
54	A	6	6	1.00	20	0.300
55	A	5	5	1.00	20	0.250
56	A	4	4	1.00	18	0.222
57	N/A	0	0	1.00	20	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
58	N/A	0	0	1.00	20	0.000
59	A	21	9	1.00	20	0.450
60	A	18	10	1.00	20	0.500
61	A	5	5	1.00	18	0.278
62	N/A	0	0	1.00	20	0.000
63	N/A	0	0	1.00	20	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \tan(a + bx) dx$	44
3.2	$\int x^2 \tan(a + bx) dx$	49
3.3	$\int x \tan(a + bx) dx$	54
3.4	$\int \frac{\tan(a+bx)}{x} dx$	58
3.5	$\int \frac{\tan(a+bx)}{x^2} dx$	61
3.6	$\int x^3 \tan^2(a + bx) dx$	64
3.7	$\int x^2 \tan^2(a + bx) dx$	70
3.8	$\int x \tan^2(a + bx) dx$	75
3.9	$\int \frac{\tan^2(a+bx)}{x} dx$	79
3.10	$\int \frac{\tan^2(a+bx)}{x^2} dx$	82
3.11	$\int x^3 \tan^3(a + bx) dx$	85
3.12	$\int x^2 \tan^3(a + bx) dx$	93
3.13	$\int x \tan^3(a + bx) dx$	99
3.14	$\int \frac{\tan^3(a+bx)}{x} dx$	104
3.15	$\int \frac{\tan^3(a+bx)}{x^2} dx$	108
3.16	$\int \left(\frac{x^2}{\tan^2(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$	112
3.17	$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx$	121
3.18	$\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx$	124
3.19	$\int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx$	130
3.20	$\int \frac{c+dx}{a+ia\tan(e+fx)} dx$	135
3.21	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx$	139
3.22	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx$	144
3.23	$\int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx$	150

3.24	$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx$	156
3.25	$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx$	163
3.26	$\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx$	168
3.27	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx$	173
3.28	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx$	180
3.29	$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx$	189
3.30	$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$	197
3.31	$\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx$	204
3.32	$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx$	210
3.33	$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx$	219
3.34	$\int (c+dx)^m (a+ia\tan(e+fx))^2 dx$	233
3.35	$\int (c+dx)^m (a+ia\tan(e+fx)) dx$	237
3.36	$\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx$	240
3.37	$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx$	244
3.38	$\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx$	248
3.39	$\int (c+dx)^3 (a+b\tan(e+fx)) dx$	253
3.40	$\int (c+dx)^2 (a+b\tan(e+fx)) dx$	260
3.41	$\int (c+dx) (a+b\tan(e+fx)) dx$	266
3.42	$\int \frac{a+b\tan(e+fx)}{c+dx} dx$	271
3.43	$\int \frac{a+b\tan(e+fx)}{(c+dx)^2} dx$	274
3.44	$\int (c+dx)^3 (a+b\tan(e+fx))^2 dx$	277
3.45	$\int (c+dx)^2 (a+b\tan(e+fx))^2 dx$	288
3.46	$\int (c+dx) (a+b\tan(e+fx))^2 dx$	297
3.47	$\int \frac{(a+b\tan(e+fx))^2}{c+dx} dx$	303
3.48	$\int \frac{(a+b\tan(e+fx))^2}{(c+dx)^2} dx$	306
3.49	$\int (c+dx)^3 (a+b\tan(e+fx))^3 dx$	310
3.50	$\int (c+dx)^2 (a+b\tan(e+fx))^3 dx$	328
3.51	$\int (c+dx) (a+b\tan(e+fx))^3 dx$	342
3.52	$\int \frac{(a+b\tan(e+fx))^3}{c+dx} dx$	350
3.53	$\int \frac{(a+b\tan(e+fx))^3}{(c+dx)^2} dx$	354
3.54	$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$	359
3.55	$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$	367
3.56	$\int \frac{c+dx}{a+b\tan(e+fx)} dx$	374
3.57	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$	379
3.58	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$	382
3.59	$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$	386
3.60	$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$	404
3.61	$\int \frac{c+dx}{(a+b\tan(e+fx))^2} dx$	417

3.62	$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$	425
3.63	$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$	429

3.1 $\int x^3 \tan(a + bx) dx$

Optimal result	44
Rubi [A] (verified)	44
Mathematica [A] (verified)	46
Maple [A] (verified)	47
Fricas [B] (verification not implemented)	47
Sympy [F]	47
Maxima [B] (verification not implemented)	48
Giac [F]	48
Mupad [F(-1)]	48

Optimal result

Integrand size = 10, antiderivative size = 106

$$\int x^3 \tan(a + bx) dx = \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ - \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4}$$

[Out] $1/4*I*x^4-x^3*ln(1+exp(2*I*(b*x+a)))/b+3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a)))$
 $/b^2-3/2*x*polylog(3,-exp(2*I*(b*x+a)))/b^3-3/4*I*polylog(4,-exp(2*I*(b*x+a)))/b^4$

Rubi [A] (verified)

Time = 0.18 (sec), antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3800, 2221, 2611, 6744, 2320, 6724}

$$\int x^3 \tan(a + bx) dx = -\frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} - \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\ + \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix^4}{4}$$

[In] Int[x^3*Tan[a + b*x], x]

[Out] $(I/4)*x^4 - (x^3*Log[1 + E^((2*I)*(a + b*x))])/b + (((3*I)/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2 - (3*x*PolyLog[3, -E^((2*I)*(a + b*x))])/(2*b^3) - (((3*I)/4)*PolyLog[4, -E^((2*I)*(a + b*x))])/b^4$

Rule 2221

```

Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^m_.)/
((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^n_.), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2320

```

Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 2611

```

Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^n_.)]*((f_.) + (g_.)
*(x_.))^m_.], x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]

```

Rule 3800

```

Int[((c_.) + (d_.)*(x_.))^m_*tan[(e_.) + (f_.)*(x_.)], x_Symbol] :> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]

```

Rule 6724

```

Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_.))^p_.]/((d_.) + (e_.)*(x_.)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]

```

Rule 6744

```

Int[((e_.) + (f_.)*(x_.))^m_*PolyLog[n_, (d_.)*(F_)^((c_.)*(a_.) + (b_.
)*(x_.)))^p_.], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{ix^4}{4} - 2i \int \frac{e^{2i(a+bx)}x^3}{1 + e^{2i(a+bx)}} dx \\
&= \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3 \int x^2 \log(1 + e^{2i(a+bx)}) dx}{b} \\
&= \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad - \frac{(3i) \int x \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b^2} \\
&= \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad - \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} + \frac{3 \int \text{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b^3} \\
&= \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad - \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{(3i) \text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2i(a+bx)}\right)}{4b^4} \\
&= \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad - \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \text{PolyLog}(4, -e^{2i(a+bx)})}{4b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 106, normalized size of antiderivative = 1.00

$$\int x^3 \tan(a + bx) dx = \frac{ix^4}{4} - \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3ix^2 \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
- \frac{3x \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3i \text{PolyLog}(4, -e^{2i(a+bx)})}{4b^4}$$

[In] `Integrate[x^3*Tan[a + b*x], x]`

[Out] `(I/4)*x^4 - (x^3*Log[1 + E^((2*I)*(a + b*x))])/b + (((3*I)/2)*x^2*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2 - (3*x*PolyLog[3, -E^((2*I)*(a + b*x))])/(2*b^3) - (((3*I)/4)*PolyLog[4, -E^((2*I)*(a + b*x))])/b^4`

Maple [A] (verified)

Time = 0.49 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.18

method	result
risch	$\frac{ix^4}{4} - \frac{3x \operatorname{Li}_3(-e^{2i(bx+a)})}{2b^3} + \frac{2ia^3x}{b^3} + \frac{3ix^2 \operatorname{Li}_2(-e^{2i(bx+a)})}{2b^2} + \frac{3ia^4}{2b^4} - \frac{3i \operatorname{Li}_4(-e^{2i(bx+a)})}{4b^4} - \frac{2a^3 \ln(e^{i(bx+a)})}{b^4} - \frac{x^3 \ln(e^{2i(bx+a)})}{b^4}$

[In] `int(x^3*tan(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $1/4*I*x^4-3/2*x*polylog(3,-exp(2*I*(b*x+a)))/b^3+2*I/b^3*a^3*x+3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a)))/b^2+3/2*I/b^4*a^4-3/4*I*polylog(4,-exp(2*I*(b*x+a)))/b^4-2/b^4*a^3*ln(exp(I*(b*x+a)))-x^3*ln(exp(2*I*(b*x+a))+1)/b$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs. $2(83) = 166$.

Time = 0.28 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.70

$$\int x^3 \tan(a + bx) dx = \frac{-4 b^3 x^3 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 4 b^3 x^3 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 6i b^2 x^2 \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 6i b^2 x^2 \operatorname{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{b^4}$$

[In] `integrate(x^3*tan(b*x+a),x, algorithm="fricas")`

[Out] $-1/8*(4*b^3*x^3*log(-2*(I*tan(b*x+a)-1)/(tan(b*x+a)^2+1))+4*b^3*x^3*log(-2*(-I*tan(b*x+a)-1)/(tan(b*x+a)^2+1))+6*I*b^2*x^2*dilog(2*(I*tan(b*x+a)-1)/(tan(b*x+a)^2+1)+1)-6*I*b^2*x^2*dilog(2*(-I*tan(b*x+a)-1)/(tan(b*x+a)^2+1)+1)+6*b*x*polylog(3,(tan(b*x+a)^2+2*I*tan(b*x+a)-1)/(tan(b*x+a)^2+1))+6*b*x*polylog(3,(tan(b*x+a)^2-2*I*tan(b*x+a)-1)/(tan(b*x+a)^2+1))-3*I*polylog(4,(tan(b*x+a)^2+2*I*tan(b*x+a)-1)/(tan(b*x+a)^2+1))+3*I*polylog(4,(tan(b*x+a)^2-2*I*tan(b*x+a)-1)/(tan(b*x+a)^2+1)))/b^4$

Sympy [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + bx) dx$$

[In] `integrate(x**3*tan(b*x+a),x)`

[Out] `Integral(x**3*tan(a + b*x), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 243 vs. $2(83) = 166$.

Time = 0.60 (sec) , antiderivative size = 243, normalized size of antiderivative = 2.29

$$\int x^3 \tan(a + bx) dx =$$

$$-\frac{-3i(bx + a)^4 + 12i(bx + a)^3a - 18i(bx + a)^2a^2 + 12a^3 \log(\sec(bx + a)) - 4(-4i(bx + a)^3 + 9i(bx + a)^2a)}{b^4}$$

[In] `integrate(x^3*tan(b*x+a),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -\frac{1}{12}(-3I*(bx + a)^4 + 12I*(bx + a)^3a - 18I*(bx + a)^2a^2 + 12a^3 \log(\sec(bx + a)) - 4(-4I*(bx + a)^3 + 9I*(bx + a)^2a)*\arctan2(\sin(2bx + 2a), \cos(2bx + 2a) + 1) - 6(4I*(bx + a)^2 - 6I*(bx + a)*a + 3I*a^2)*\text{dilog}(-e^{(2I*bx + 2I*a)}) + 2(4*(bx + a)^3 - 9*(bx + a)^2a + 9*(bx + a)*a^2)*\log(\cos(2bx + 2a)^2 + \sin(2bx + 2a)^2 + 2*\cos(2bx + 2a) + 1) + 6(4bx + a)*\text{polylog}(3, -e^{(2I*bx + 2I*a)}) + 12I*\text{polylog}(4, -e^{(2I*bx + 2I*a)}))/b^4 \end{aligned}$$

Giac [F]

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(bx + a) dx$$

[In] `integrate(x^3*tan(b*x+a),x, algorithm="giac")`

[Out] `integrate(x^3*tan(b*x + a), x)`

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan(a + bx) dx = \int x^3 \tan(a + b x) dx$$

[In] `int(x^3*tan(a + b*x),x)`

[Out] `int(x^3*tan(a + b*x), x)`

3.2 $\int x^2 \tan(a + bx) dx$

Optimal result	49
Rubi [A] (verified)	49
Mathematica [A] (verified)	51
Maple [A] (verified)	51
Fricas [B] (verification not implemented)	51
Sympy [F]	52
Maxima [B] (verification not implemented)	52
Giac [F]	53
Mupad [F(-1)]	53

Optimal result

Integrand size = 10, antiderivative size = 77

$$\begin{aligned} \int x^2 \tan(a + bx) dx = & \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} \\ & + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \end{aligned}$$

[Out] $\frac{1}{3}I*x^3 - x^2 \ln(1 + \exp(2*I*(b*x+a))) / b + I*x*\operatorname{polylog}(2, -\exp(2*I*(b*x+a))) / b^2 - \frac{1}{2}\operatorname{polylog}(3, -\exp(2*I*(b*x+a))) / b^3$

Rubi [A] (verified)

Time = 0.17 (sec), antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3800, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int x^2 \tan(a + bx) dx = & -\frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} \\ & - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix^3}{3} \end{aligned}$$

[In] $\operatorname{Int}[x^2 \tan(a + b*x), x]$

[Out] $(I/3)*x^3 - (x^2 \operatorname{Log}[1 + E^{((2*I)*(a + b*x))}]) / b + (I*x*\operatorname{PolyLog}[2, -E^{((2*I)*(a + b*x))}]) / b^2 - \operatorname{PolyLog}[3, -E^{((2*I)*(a + b*x))}] / (2*b^3)$

Rule 2221

$\operatorname{Int}[(((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_*)}) / ((a_) + (b_*)*((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}$

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_.))^m_, x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_.))^m_*tan[(e_.) + (f_.)*(x_.)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^p]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{ix^3}{3} - 2i \int \frac{e^{2i(a+bx)}x^2}{1 + e^{2i(a+bx)}} dx \\
 &= \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{2 \int x \log(1 + e^{2i(a+bx)}) dx}{b} \\
 &= \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{i \int \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b^2} \\
 &= \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^2} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(a+bx)}\right)}{2b^3}
 \end{aligned}$$

$$= \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.00

$$\int x^2 \tan(a + bx) dx = \frac{ix^3}{3} - \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3}$$

[In] `Integrate[x^2*Tan[a + b*x], x]`

[Out] $\frac{(I/3)*x^3 - (x^2*\log[1 + E^{((2*I)*(a + b*x))}])/b + (I*x*\operatorname{PolyLog}[2, -E^{((2*I)*(a + b*x))}])/b^2 - \operatorname{PolyLog}[3, -E^{((2*I)*(a + b*x))}]}{(2*b^3)}$

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.34

method	result	size
risch	$\frac{ix^3}{3} - \frac{2ia^2x}{b^2} - \frac{4ia^3}{3b^3} - \frac{x^2 \ln(e^{2i(bx+a)}+1)}{b} + \frac{ix \operatorname{Li}_2(-e^{2i(bx+a)})}{b^2} - \frac{\operatorname{Li}_3(-e^{2i(bx+a)})}{2b^3} + \frac{2a^2 \ln(e^{i(bx+a)})}{b^3}$	103

[In] `int(x^2*tan(b*x+a), x, method=_RETURNVERBOSE)`

[Out] $\frac{1/3*I*x^3-2*I/b^2*a^2*x-4/3*I/b^3*a^3-x^2*ln(exp(2*I*(b*x+a))+1)/b+I*x*polylog(2,-exp(2*I*(b*x+a)))/b^2-1/2*polylog(3,-exp(2*I*(b*x+a)))/b^3+2/b^3*a^2*ln(exp(I*(b*x+a)))}{4b^3}$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(62) = 124$.

Time = 0.26 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.60

$$\int x^2 \tan(a + bx) dx = \frac{-2b^2 x^2 \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2b^2 x^2 \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2i b x \operatorname{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - 2i b x \operatorname{Li}_3\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right)}{4b^3}$$

[In] `integrate(x^2*tan(b*x+a), x, algorithm="fricas")`

```
[Out] -1/4*(2*b^2*x^2*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b^2*x^2*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*I*b*x*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*I*b*x*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^3
```

Sympy [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + bx) dx$$

```
[In] integrate(x**2*tan(b*x+a),x)
```

```
[Out] Integral(x**2*tan(a + b*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(62) = 124$.

Time = 0.68 (sec) , antiderivative size = 163, normalized size of antiderivative = 2.12

$$\int x^2 \tan(a + bx) dx = \frac{-2i(bx + a)^3 + 6i(bx + a)^2a - 6i b x \text{Li}_2(-e^{(2ibx+2ia)}) - 6a^2 \log(\sec(bx + a)) - 6(-i(bx + a)^2 + 2i(bx + a))}{b^3}$$

```
[In] integrate(x^2*tan(b*x+a),x, algorithm="maxima")
```

```
[Out] -1/6*(-2*I*(b*x + a)^3 + 6*I*(b*x + a)^2*a - 6*I*b*x*dilog(-e^(2*I*b*x + 2*I*a)) - 6*a^2*log(sec(b*x + a)) - 6*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^2 - 2*(b*x + a)*a)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 3*polylog(3, -e^(2*I*b*x + 2*I*a)))/b^3
```

Giac [F]

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(bx + a) dx$$

[In] integrate(x^2*tan(b*x+a),x, algorithm="giac")

[Out] integrate(x^2*tan(b*x + a), x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan(a + bx) dx = \int x^2 \tan(a + b x) dx$$

[In] int(x^2*tan(a + b*x),x)

[Out] int(x^2*tan(a + b*x), x)

3.3 $\int x \tan(a + bx) dx$

Optimal result	54
Rubi [A] (verified)	54
Mathematica [A] (verified)	55
Maple [A] (verified)	56
Fricas [B] (verification not implemented)	56
Sympy [F]	56
Maxima [B] (verification not implemented)	57
Giac [F]	57
Mupad [B] (verification not implemented)	57

Optimal result

Integrand size = 8, antiderivative size = 54

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

[Out] $1/2*I*x^2 - x*\ln(1+\exp(2*I*(b*x+a)))/b + 1/2*I*\operatorname{polylog}(2, -\exp(2*I*(b*x+a)))/b^2$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3800, 2221, 2317, 2438}

$$\int x \tan(a + bx) dx = \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{ix^2}{2}$$

[In] $\operatorname{Int}[x \tan(a + b*x), x]$

[Out] $(I/2)*x^2 - (x*\operatorname{Log}[1 + E^{((2*I)*(a + b*x))}])/b + ((I/2)*\operatorname{PolyLog}[2, -E^{((2*I)*(a + b*x))}])/b^2$

Rule 2221

```
Int[((F_)^(g_)*(e_) + (f_)*(x_)))^(n_)*(c_) + (d_)*(x_)^(m_))/((a_) + (b_)*(F_)^(g_)*(e_) + (f_)*(x_)))^(n_), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*(c_.) + (d_)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[((c_.)*(d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[((c_.) + (d_)*(x_)^m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^m/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{ix^2}{2} - 2i \int \frac{e^{2i(a+bx)}x}{1 + e^{2i(a+bx)}} dx \\ &= \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{\int \log(1 + e^{2i(a+bx)}) dx}{b} \\ &= \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(a+bx)}\right)}{2b^2} \\ &= \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 54, normalized size of antiderivative = 1.00

$$\int x \tan(a + bx) dx = \frac{ix^2}{2} - \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{i \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2}$$

[In] `Integrate[x*Tan[a + b*x], x]`

[Out] `(I/2)*x^2 - (x*Log[1 + E^((2*I)*(a + b*x))])/b + ((I/2)*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2`

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.44

method	result
risch	$\frac{ix^2}{2} + \frac{2iax}{b} + \frac{ia^2}{b^2} - \frac{x \ln(e^{2i(bx+a)}+1)}{b} + \frac{i \text{Li}_2(-e^{2i(bx+a)})}{2b^2} - \frac{2a \ln(e^{i(bx+a)})}{b^2}$
parts	$\frac{\ln(1+\tan^2(bx+a))x}{2b} - \frac{i \left(\ln(\tan(bx+a)-i) \ln(1+\tan^2(bx+a)) - \text{dilog}\left(-\frac{i(\tan(bx+a)+i)}{2}\right) - \ln(\tan(bx+a)-i) \ln\left(-\frac{i(\tan(bx+a)+i)}{2}\right) - \ln(\tan(bx+a)-i) \ln\left(\frac{i(\tan(bx+a)+i)}{2}\right) \right)}{2}$

[In] `int(x*tan(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}I*x^2 + 2*I/b*a*x + I/b^2*a^2 - x*\ln(\exp(2*I*(b*x+a))+1)/b + 1/2*I*\text{polylog}(2,-e^{2*I*(b*x+a)})/b^2 - 2/b^2*a*\ln(\exp(I*(b*x+a)))$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 122 vs. $2(41) = 82$.

Time = 0.24 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.26

$$\int x \tan(a + bx) dx = -\frac{2 b x \log\left(-\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + 2 b x \log\left(-\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1}\right) + i \text{Li}_2\left(\frac{2(i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right) - i \text{Li}_2\left(\frac{2(-i \tan(bx+a)-1)}{\tan(bx+a)^2+1} + 1\right)}{4 b^2}$$

[In] `integrate(x*tan(b*x+a),x, algorithm="fricas")`

[Out] $\frac{-1/4*(2*b*x*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b*x*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + I*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - I*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1))/b^2}{b^2}$

Sympy [F]

$$\int x \tan(a + bx) dx = \int x \tan(a + bx) dx$$

[In] `integrate(x*tan(b*x+a),x)`

[Out] `Integral(x*tan(a + b*x), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 92 vs. $2(41) = 82$.

Time = 0.40 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.70

$$\int x \tan(a + bx) dx = -\frac{-i b^2 x^2 + 2i b x \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) + b x \log(\cos(2bx + 2a)^2 + \sin(2bx + 2a)^2)}{2 b^2}$$

```
[In] integrate(x*tan(b*x+a),x, algorithm="maxima")
[Out] -1/2*(-I*b^2*x^2 + 2*I*b*x*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1)
+ b*x*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1)
- I*dilog(-e^(2*I*b*x + 2*I*a)))/b^2
```

Giac [F]

$$\int x \tan(a + bx) dx = \int x \tan(bx + a) dx$$

```
[In] integrate(x*tan(b*x+a),x, algorithm="giac")
[Out] integrate(x*tan(b*x + a), x)
```

Mupad [B] (verification not implemented)

Time = 3.35 (sec) , antiderivative size = 129, normalized size of antiderivative = 2.39

$$\int x \tan(a + bx) dx = -\frac{\pi \ln(\cos(bx)) + \text{polylog}(2, -e^{-a2i} e^{-bx2i})}{2 b^2} 1i - \frac{\pi \ln(e^{-a2i} e^{-bx2i} + 1) + 2a \ln(e^{-a2i} e^{-bx2i} + 1) - \pi}{2 b^2}$$

```
[In] int(x*tan(a + b*x),x)
[Out] -(polylog(2, -exp(-a*2i)*exp(-b*x*2i))*1i - pi*log(exp(b*x*2i) + 1) - pi*log(exp(-a*2i)*exp(-b*x*2i) + 1) + 2*a*log(exp(-a*2i)*exp(-b*x*2i) + 1) + pi*log(cos(b*x)) + b^2*x^2*1i - log(cos(a + b*x))*(2*a - pi) + 2*b*x*log(exp(-a*2i)*exp(-b*x*2i) + 1) + a*b*x*2i)/(2*b^2)
```

3.4 $\int \frac{\tan(a+bx)}{x} dx$

Optimal result	58
Rubi [N/A]	58
Mathematica [N/A]	59
Maple [N/A] (verified)	59
Fricas [N/A]	59
Sympy [N/A]	59
Maxima [N/A]	60
Giac [N/A]	60
Mupad [N/A]	60

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a + bx)}{x} dx = \text{Int}\left(\frac{\tan(a + bx)}{x}, x\right)$$

[Out] Unintegrable(tan(b*x+a)/x,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + bx)}{x} dx$$

[In] Int[Tan[a + b*x]/x,x]

[Out] Defer[Int][Tan[a + b*x]/x, x]

Rubi steps

$$\text{integral} = \int \frac{\tan(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 1.55 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + bx)}{x} dx$$

[In] Integrate[Tan[a + b*x]/x, x]

[Out] Integrate[Tan[a + b*x]/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.24 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx + a)}{x} dx$$

[In] int(tan(b*x+a)/x, x)

[Out] int(tan(b*x+a)/x, x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

[In] integrate(tan(b*x+a)/x, x, algorithm="fricas")

[Out] integral(tan(b*x + a)/x, x)

Sympy [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + bx)}{x} dx$$

[In] integrate(tan(b*x+a)/x, x)

[Out] Integral(tan(a + b*x)/x, x)

Maxima [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

[In] `integrate(tan(b*x+a)/x,x, algorithm="maxima")`

[Out] `integrate(tan(b*x + a)/x, x)`

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(bx + a)}{x} dx$$

[In] `integrate(tan(b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 3.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x} dx = \int \frac{\tan(a + b x)}{x} dx$$

[In] `int(tan(a + b*x)/x,x)`

[Out] `int(tan(a + b*x)/x, x)`

3.5 $\int \frac{\tan(a+bx)}{x^2} dx$

Optimal result	61
Rubi [N/A]	61
Mathematica [N/A]	62
Maple [N/A] (verified)	62
Fricas [N/A]	62
Sympy [N/A]	62
Maxima [N/A]	63
Giac [N/A]	63
Mupad [N/A]	63

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\tan(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable(tan(b*x+a)/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)}{x^2} dx$$

[In] Int[Tan[a + b*x]/x^2, x]

[Out] Defer[Int][Tan[a + b*x]/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tan(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 2.42 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)}{x^2} dx$$

[In] `Integrate[Tan[a + b*x]/x^2, x]`

[Out] `Integrate[Tan[a + b*x]/x^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.20 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(bx + a)}{x^2} dx$$

[In] `int(tan(b*x+a)/x^2, x)`

[Out] `int(tan(b*x+a)/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

[In] `integrate(tan(b*x+a)/x^2, x, algorithm="fricas")`

[Out] `integral(tan(b*x + a)/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)}{x^2} dx$$

[In] `integrate(tan(b*x+a)/x**2, x)`

[Out] `Integral(tan(a + b*x)/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

[In] `integrate(tan(b*x+a)/x^2,x, algorithm="maxima")`

[Out] `integrate(tan(b*x + a)/x^2, x)`

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)}{x^2} dx$$

[In] `integrate(tan(b*x+a)/x^2,x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 3.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\tan(a + bx)}{x^2} dx = \int \frac{\tan(a + b x)}{x^2} dx$$

[In] `int(tan(a + b*x)/x^2,x)`

[Out] `int(tan(a + b*x)/x^2, x)`

3.6 $\int x^3 \tan^2(a + bx) dx$

Optimal result	64
Rubi [A] (verified)	64
Mathematica [A] (verified)	66
Maple [A] (verified)	67
Fricas [B] (verification not implemented)	67
Sympy [F]	67
Maxima [B] (verification not implemented)	68
Giac [F]	68
Mupad [F(-1)]	69

Optimal result

Integrand size = 12, antiderivative size = 98

$$\begin{aligned} \int x^3 \tan^2(a + bx) dx = & -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\ & + \frac{3 \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^4} + \frac{x^3 \tan(a + bx)}{b} \end{aligned}$$

[Out] $-\text{I}*\text{x}^3/\text{b}-1/4*\text{x}^4+3*\text{x}^2*\ln(1+\exp(2*\text{I}*(\text{b}*\text{x}+\text{a})))/\text{b}^2-3*\text{I}*\text{x}*\operatorname{polylog}(2, -\exp(2*\text{I}*(\text{b}*\text{x}+\text{a})))/\text{b}^3+3/2*\operatorname{polylog}(3, -\exp(2*\text{I}*(\text{b}*\text{x}+\text{a})))/\text{b}^4+\text{x}^3\tan(\text{b}*\text{x}+\text{a})/\text{b}$

Rubi [A] (verified)

Time = 0.20 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3801, 3800, 2221, 2611, 2320, 6724, 30}

$$\begin{aligned} \int x^3 \tan^2(a + bx) dx = & \frac{3 \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^4} - \frac{3ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\ & + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} + \frac{x^3 \tan(a + bx)}{b} - \frac{ix^3}{b} - \frac{x^4}{4} \end{aligned}$$

[In] $\text{Int}[\text{x}^3 \tan[a + b*x]^2, \text{x}]$

[Out] $((-\text{I})*\text{x}^3)/\text{b}-\text{x}^4/4+(3*\text{x}^2*\log[1+\text{E}^{((2*\text{I})*(a+b*x))}])/\text{b}^2-((3*\text{I})*\text{x}*\operatorname{PolyLog}[2, -\text{E}^{((2*\text{I})*(a+b*x))}])/\text{b}^3+(3*\operatorname{PolyLog}[3, -\text{E}^{((2*\text{I})*(a+b*x))}])/((2*\text{b}^4)+(\text{x}^3 \tan[a+b*x])/\text{b})$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_)^(m_.))/
((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x))
))^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]])]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_)^(m_.))*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d}
```

, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^3 \tan(a + bx)}{b} - \frac{3 \int x^2 \tan(a + bx) dx}{b} - \int x^3 dx \\
 &= -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{x^3 \tan(a + bx)}{b} + \frac{(6i) \int \frac{e^{2i(a+bx)} x^2}{1+e^{2i(a+bx)}} dx}{b} \\
 &= -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} + \frac{x^3 \tan(a + bx)}{b} - \frac{6 \int x \log(1 + e^{2i(a+bx)}) dx}{b^2} \\
 &= -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\
 &\quad + \frac{x^3 \tan(a + bx)}{b} + \frac{(3i) \int \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b^3} \\
 &= -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\
 &\quad + \frac{x^3 \tan(a + bx)}{b} + \frac{3 \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(a+bx)}\right)}{2b^4} \\
 &= -\frac{ix^3}{b} - \frac{x^4}{4} + \frac{3x^2 \log(1 + e^{2i(a+bx)})}{b^2} - \frac{3ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^3} \\
 &\quad + \frac{3 \text{PolyLog}(3, -e^{2i(a+bx)})}{2b^4} + \frac{x^3 \tan(a + bx)}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.69 (sec), antiderivative size = 115, normalized size of antiderivative = 1.17

$$\begin{aligned}
 \int x^3 \tan^2(a + bx) dx &= -\frac{x^4}{4} \\
 &\quad + \frac{2b^2 x^2 \left(\frac{2ibx}{1+e^{2ia}} + 3 \log(1 + e^{-2i(a+bx)}) \right) + 6ibx \text{PolyLog}(2, -e^{-2i(a+bx)}) + 3 \text{PolyLog}(3, -e^{-2i(a+bx)})}{2b^4} \\
 &\quad + \frac{x^3 \sec(a) \sec(a + bx) \sin(bx)}{b}
 \end{aligned}$$

[In] Integrate[x^3*Tan[a + b*x]^2, x]

[Out] $-1/4*x^4 + (2*b^2*x^2*((2*I)*b*x)/(1 + E^((2*I)*a)) + 3*Log[1 + E^((-2*I)*(a + b*x))]) + (6*I)*b*x*PolyLog[2, -E^((-2*I)*(a + b*x))] + 3*PolyLog[3, -E^((-2*I)*(a + b*x))]/(2*b^4) + (x^3*Sec[a]*Sec[a + b*x]*Sin[b*x])/b$

Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.36

method	result
risch	$-\frac{x^4}{4} + \frac{2ix^3}{b(e^{2i(bx+a)}+1)} - \frac{2ix^3}{b} + \frac{6ia^2x}{b^3} + \frac{4ia^3}{b^4} + \frac{3x^2 \ln(e^{2i(bx+a)}+1)}{b^2} - \frac{3ix \text{Li}_2(-e^{2i(bx+a)})}{b^3} + \frac{3 \text{Li}_3(-e^{2i(bx+a)})}{2b^4} -$

```
[In] int(x^3*tan(b*x+a)^2,x,method=_RETURNVERBOSE)
[Out] -1/4*x^4+2*I*x^3/b/(exp(2*I*(b*x+a))+1)-2*I/b*x^3+6*I/b^3*a^2*x+4*I/b^4*a^3
+3*x^2*ln(exp(2*I*(b*x+a))+1)/b^2-3*I*x*polylog(2,-exp(2*I*(b*x+a)))/b^3+3/
2*polylog(3,-exp(2*I*(b*x+a)))/b^4-6/b^4*a^2*ln(exp(I*(b*x+a)))
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 225 vs. 2(83) = 166.

Time = 0.26 (sec) , antiderivative size = 225, normalized size of antiderivative = 2.30

$$\int x^3 \tan^2(a + bx) dx =$$

$$-\frac{b^4 x^4 - 4 b^3 x^3 \tan(bx + a) - 6 b^2 x^2 \log\left(-\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) - 6 b^2 x^2 \log\left(-\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) - 6 i b x \text{Li}_2\left(\frac{-i \tan(bx + a)}{\tan(bx + a)^2 + 1}\right) + 6 i b x \text{Li}_2\left(\frac{i \tan(bx + a)}{\tan(bx + a)^2 + 1}\right) + 6 b^2 x^2 \text{dilog}(2 * (\text{I} * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1) + 1) + 6 b^2 x^2 \text{dilog}(2 * (-\text{I} * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1) + 1) - 3 * \text{polylog}(3, (\tan(bx + a)^2 + 2 * \text{I} * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1)) - 3 * \text{polylog}(3, (\tan(bx + a)^2 - 2 * \text{I} * \tan(bx + a) - 1) / (\tan(bx + a)^2 + 1)))}{b^4}$$

```
[In] integrate(x^3*tan(b*x+a)^2,x, algorithm="fricas")
[Out] -1/4*(b^4*x^4 - 4*b^3*x^3*tan(b*x + a) - 6*b^2*x^2*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 6*b^2*x^2*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 6*I*b*x*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + 6*I*b*x*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 3*polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 3*polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^4
```

Sympy [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan^2(a + bx) dx$$

```
[In] integrate(x**3*tan(b*x+a)**2,x)
[Out] Integral(x**3*tan(a + b*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 639 vs. $2(83) = 166$.

Time = 0.40 (sec) , antiderivative size = 639, normalized size of antiderivative = 6.52

$$\int x^3 \tan^2(a + bx) dx \\ = \frac{2(bx + a - \tan(bx + a))a^3 - \frac{3((bx+a)^2 \cos(2bx+2a)^2 + (bx+a)^2 \sin(2bx+2a)^2 + 2(bx+a)^2 \cos(2bx+2a) + (bx+a)^2 - (\cos(2bx+2a)^2 + \cos(2bx+2a)^2)}{\cos(2bx+2a)^2 + \cos(2bx+2a)^2}}{}$$

```
[In] integrate(x^3*tan(b*x+a)^2,x, algorithm="maxima")
[Out] 1/2*(2*(b*x + a - tan(b*x + a))*a^3 - 3*((b*x + a)^2*cos(2*b*x + 2*a)^2 + (b*x + a)^2*sin(2*b*x + 2*a)^2 + 2*(b*x + a)^2*cos(2*b*x + 2*a) + (b*x + a)^2 - (cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 4*(b*x + a)*sin(2*b*x + 2*a))*a^2/(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 2*(I*(b*x + a)^4 - 4*I*(b*x + a)^3*a + 12*((b*x + a)^2 - 2*(b*x + a)*a + ((b*x + a)^2 - 2*(b*x + a)*a)*cos(2*b*x + 2*a) - (-I*(b*x + a)^2 + 2*I*(b*x + a)*a)*sin(2*b*x + 2*a))*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + (I*(b*x + a)^4 - 4*(b*x + a)^3*(I*a + 2) + 24*(b*x + a)^2*a)*cos(2*b*x + 2*a) - 12*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*dilog(-e^(2*I*b*x + 2*I*a)) - 6*(I*(b*x + a)^2 - 2*I*(b*x + a)*a + (I*(b*x + a)^2 - 2*I*(b*x + a)*a)*cos(2*b*x + 2*a) - ((b*x + a)^2 - 2*(b*x + a)*a)*sin(2*b*x + 2*a))*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 6*(I*cos(2*b*x + 2*a) - sin(2*b*x + 2*a) + I)*polylog(3, -e^(2*I*b*x + 2*I*a)) - ((b*x + a)^4 - 4*(b*x + a)^3*(a - 2*I) - 24*I*(b*x + a)^2*a)*sin(2*b*x + 2*a))/(-4*I*cos(2*b*x + 2*a) + 4*sin(2*b*x + 2*a) - 4*I))/b^4
```

Giac [F]

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(bx + a)^2 dx$$

```
[In] integrate(x^3*tan(b*x+a)^2,x, algorithm="giac")
[Out] integrate(x^3*tan(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan^2(a + bx) dx = \int x^3 \tan(a + bx)^2 dx$$

[In] int(x^3*tan(a + b*x)^2, x)

[Out] int(x^3*tan(a + b*x)^2, x)

3.7 $\int x^2 \tan^2(a + bx) dx$

Optimal result	70
Rubi [A] (verified)	70
Mathematica [B] (verified)	72
Maple [A] (verified)	72
Fricas [B] (verification not implemented)	73
Sympy [F]	73
Maxima [B] (verification not implemented)	73
Giac [F]	74
Mupad [F(-1)]	74

Optimal result

Integrand size = 12, antiderivative size = 73

$$\begin{aligned} \int x^2 \tan^2(a + bx) dx = & -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} \\ & - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{x^2 \tan(a + bx)}{b} \end{aligned}$$

[Out] $-\text{I}*\text{x}^2/\text{b}-1/3*\text{x}^3+2*\text{x}*\ln(1+\exp(2*\text{I}*(\text{b}*\text{x}+\text{a})))/\text{b}^2-\text{I}*\operatorname{polylog}(2,-\exp(2*\text{I}*(\text{b}*\text{x}+\text{a})))/\text{b}^3+\text{x}^2\tan(\text{b}*\text{x}+\text{a})/\text{b}$

Rubi [A] (verified)

Time = 0.15 (sec), antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3801, 3800, 2221, 2317, 2438, 30}

$$\begin{aligned} \int x^2 \tan^2(a + bx) dx = & -\frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} \\ & + \frac{x^2 \tan(a + bx)}{b} - \frac{ix^2}{b} - \frac{x^3}{3} \end{aligned}$$

[In] $\text{Int}[\text{x}^2 \tan[a + b*x]^2, \text{x}]$

[Out] $((-\text{I})*\text{x}^2)/\text{b}-\text{x}^3/3+(2*\text{x}*\operatorname{Log}[1+\text{E}^{((2*\text{I})*(a+b*x))}])/\text{b}^2-(\text{I}*\operatorname{PolyLog}[2,-\text{E}^{((2*\text{I})*(a+b*x))}])/\text{b}^3+(\text{x}^2 \tan[a + b*x])/b$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] := Simplify[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2221

```
Int[((F_)((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*(F_)((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*(F_)((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x^2 \tan(a + bx)}{b} - \frac{2 \int x \tan(a + bx) dx}{b} - \int x^2 dx \\
 &= -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{x^2 \tan(a + bx)}{b} + \frac{(4i) \int \frac{e^{2i(a+bx)} x}{1+e^{2i(a+bx)}} dx}{b} \\
 &= -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} + \frac{x^2 \tan(a + bx)}{b} - \frac{2 \int \log(1 + e^{2i(a+bx)}) dx}{b^2} \\
 &= -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} + \frac{x^2 \tan(a + bx)}{b} + \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(a+bx)}\right)}{b^3}
 \end{aligned}$$

$$= -\frac{ix^2}{b} - \frac{x^3}{3} + \frac{2x \log(1 + e^{2i(a+bx)})}{b^2} - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^3} + \frac{x^2 \tan(a + bx)}{b}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 160 vs. $2(73) = 146$.

Time = 5.05 (sec), antiderivative size = 160, normalized size of antiderivative = 2.19

$$\int x^2 \tan^2(a + bx) dx = -\frac{x^3}{3} + \frac{x^2 \sec(a) \sec(a + bx) \sin(bx)}{b} + \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))}) - \pi \log(1 - e^{2i(bx - \arctan(\cot(a)))})}{b^3}$$

[In] `Integrate[x^2*Tan[a + b*x]^2, x]`

[Out]
$$-\frac{1}{3}x^3 + (x^2 \sec(a) \sec(a + bx) \sin(bx))/b + (I*b*x*(Pi + 2*ArcTan[Cot[a]]) + Pi*Log[1 + E^((-2*I)*b*x)] + 2*(b*x - ArcTan[Cot[a]])*Log[1 - E^((2*I)*(b*x - ArcTan[Cot[a]]))] - Pi*Log[Cos[b*x]] + 2*ArcTan[Cot[a]]*Log[Sin[b*x - ArcTan[Cot[a]]]] - I*PolyLog[2, E^((2*I)*(b*x - ArcTan[Cot[a]]))]) + (b^2*x^2*Sqrt[Csc[a]^2]*Tan[a])/E^(I*ArcTan[Cot[a]]))/b^3$$

Maple [A] (verified)

Time = 0.55 (sec), antiderivative size = 108, normalized size of antiderivative = 1.48

method	result	size
risch	$-\frac{x^3}{3} + \frac{2ix^2}{b(e^{2i(bx+a)}+1)} - \frac{2ix^2}{b} - \frac{4iax}{b^2} - \frac{2ia^2}{b^3} + \frac{2x \ln(e^{2i(bx+a)}+1)}{b^2} - \frac{i \operatorname{Li}_2(-e^{2i(bx+a)})}{b^3} + \frac{4a \ln(e^{i(bx+a)})}{b^3}$	108

[In] `int(x^2*tan(b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out]
$$-\frac{1}{3}x^3 + 2*I*x^2/b/(\exp(2*I*(b*x+a))+1) - 2*I/b*x^2 - 4*I/b^2*a*x - 2*I/b^3*a^2*x^2 + x*x*ln(\exp(2*I*(b*x+a))+1)/b^2 - I*polylog(2, -\exp(2*I*(b*x+a)))/b^3 + 4/b^3*a*ln(\exp(I*(b*x+a)))$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 144 vs. $2(62) = 124$.

Time = 0.25 (sec) , antiderivative size = 144, normalized size of antiderivative = 1.97

$$\int x^2 \tan^2(a + bx) dx = \frac{2 b^3 x^3 - 6 b^2 x^2 \tan(bx + a) - 6 b x \log\left(-\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) - 6 b x \log\left(-\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) - 3i \text{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) - 3i \text{Li}_2\left(\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right)}{6 b^3}$$

```
[In] integrate(x^2*tan(b*x+a)^2,x, algorithm="fricas")
[Out] -1/6*(2*b^3*x^3 - 6*b^2*x^2*tan(b*x + a) - 6*b*x*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 6*b*x*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) - 3*I*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) + 3*I*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1))/b^3
```

Sympy [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan^2(a + bx) dx$$

```
[In] integrate(x**2*tan(b*x+a)**2,x)
[Out] Integral(x**2*tan(a + b*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 257 vs. $2(62) = 124$.

Time = 0.63 (sec) , antiderivative size = 257, normalized size of antiderivative = 3.52

$$\int x^2 \tan^2(a + bx) dx = \frac{i b^3 x^3 + 6 (bx \cos(2bx + 2a) + i bx \sin(2bx + 2a) + bx) \arctan(\sin(2bx + 2a), \cos(2bx + 2a) + 1) + (-3b^2 x^2 \cos(2bx + 2a) - 6b^2 x \sin(2bx + 2a) - 3b^2 \cos(2bx + 2a) - 6b^2 \sin(2bx + 2a) - 3b^2) \operatorname{dilog}(-e^{(2*I*b*x + 2*I*a)}) - 3(I*b*x*cos(2*b*x + 2*a) - b*x*sin(2*b*x + 2*a) + I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - (b^3*x^3 + 6*I*b^2*x^2)*sin(2*b*x + 2*a))/(-3*I*b^3*cos(2*b*x + 2*a) + 3*b^3*sin(2*b*x + 2*a) - 3*I*b^3)}$$

```
[In] integrate(x^2*tan(b*x+a)^2,x, algorithm="maxima")
[Out] (I*b^3*x^3 + 6*(b*x*cos(2*b*x + 2*a) + I*b*x*sin(2*b*x + 2*a) + b*x)*arctan(2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + (I*b^3*x^3 - 6*b^2*x^2)*cos(2*b*x + 2*a) - 3*(cos(2*b*x + 2*a) + I*sin(2*b*x + 2*a) + 1)*dilog(-e^(2*I*b*x + 2*I*a)) - 3*(I*b*x*cos(2*b*x + 2*a) - b*x*sin(2*b*x + 2*a) + I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - (b^3*x^3 + 6*I*b^2*x^2)*sin(2*b*x + 2*a))/(-3*I*b^3*cos(2*b*x + 2*a) + 3*b^3*sin(2*b*x + 2*a) - 3*I*b^3))
```

Giac [F]

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(bx + a)^2 dx$$

[In] integrate(x^2*tan(b*x+a)^2,x, algorithm="giac")
 [Out] integrate(x^2*tan(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan^2(a + bx) dx = \int x^2 \tan(a + b x)^2 dx$$

[In] int(x^2*tan(a + b*x)^2,x)
 [Out] int(x^2*tan(a + b*x)^2, x)

3.8 $\int x \tan^2(a + bx) dx$

Optimal result	75
Rubi [A] (verified)	75
Mathematica [A] (verified)	76
Maple [A] (verified)	76
Fricas [A] (verification not implemented)	77
Sympy [A] (verification not implemented)	77
Maxima [B] (verification not implemented)	77
Giac [B] (verification not implemented)	78
Mupad [B] (verification not implemented)	78

Optimal result

Integrand size = 10, antiderivative size = 30

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b}$$

[Out] $-1/2*x^2 + \ln(\cos(b*x+a))/b^2 + x*tan(b*x+a)/b$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3801, 3556, 30}

$$\int x \tan^2(a + bx) dx = \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b} - \frac{x^2}{2}$$

[In] $\text{Int}[x*\text{Tan}[a + b*x]^2, x]$

[Out] $-1/2*x^2 + \text{Log}[\text{Cos}[a + b*x]]/b^2 + (x*\text{Tan}[a + b*x])/b$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 3556

```
Int[tan[(c_.) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3801

```
Int[((c_) + (d_)*(x_))^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}\text{integral} &= \frac{x \tan(a + bx)}{b} - \frac{\int \tan(a + bx) dx}{b} - \int x dx \\ &= -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \tan(a + bx)}{b}\end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.43

$$\int x \tan^2(a + bx) dx = -\frac{x^2}{2} + \frac{\log(\cos(a + bx))}{b^2} + \frac{x \sec(a) \sec(a + bx) \sin(bx)}{b} + \frac{x \tan(a)}{b}$$

[In] `Integrate[x*Tan[a + b*x]^2, x]`

[Out]
$$-\frac{1}{2}x^2 + \frac{\log(\cos(a + bx))}{b^2} + \frac{(x \sec(a) \sec(a + bx) \sin(bx))}{b} + \frac{(x \tan(a))}{b}$$

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.13

method	result	size
norman	$\frac{x \tan(bx+a)}{b} - \frac{x^2}{2} - \frac{\ln(1+\tan^2(bx+a))}{2b^2}$	34
parallelrisch	$-\frac{x^2 b^2 - 2x \tan(bx+a) b + \ln(1+\tan^2(bx+a))}{2b^2}$	35
default	$-\frac{x^2}{2} + \frac{(bx+a) \tan(bx+a) + \ln(\cos(bx+a)) - a \tan(bx+a)}{b^2}$	40
risch	$-\frac{x^2}{2} - \frac{2ix}{b} - \frac{2ia}{b^2} + \frac{2ix}{b(e^{2i(bx+a)}+1)} + \frac{\ln(e^{2i(bx+a)}+1)}{b^2}$	57

[In] `int(x*tan(b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out]
$$x \tan(bx+a)/b - 1/2x^2 - 1/2/b^2 \ln(1+\tan(bx+a)^2)$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.27

$$\int x \tan^2(a + bx) dx = -\frac{b^2 x^2 - 2 b x \tan(bx + a) - \log\left(\frac{1}{\tan(bx+a)^2+1}\right)}{2 b^2}$$

[In] `integrate(x*tan(b*x+a)^2,x, algorithm="fricas")`

[Out] `-1/2*(b^2*x^2 - 2*b*x*tan(b*x + a) - log(1/(tan(b*x + a)^2 + 1)))/b^2`

Sympy [A] (verification not implemented)

Time = 0.22 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.37

$$\int x \tan^2(a + bx) dx = \begin{cases} -\frac{x^2}{2} + \frac{x \tan(a+bx)}{b} - \frac{\log(\tan^2(a+bx)+1)}{2b^2} & \text{for } b \neq 0 \\ \frac{x^2 \tan^2(a)}{2} & \text{otherwise} \end{cases}$$

[In] `integrate(x*tan(b*x+a)**2,x)`

[Out] `Piecewise((-x**2/2 + x*tan(a + b*x)/b - log(tan(a + b*x)**2 + 1)/(2*b**2), Ne(b, 0)), (x**2*tan(a)**2/2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(28) = 56.

Time = 0.52 (sec) , antiderivative size = 214, normalized size of antiderivative = 7.13

$$\int x \tan^2(a + bx) dx = \frac{2 (bx + a - \tan(bx + a))a - \frac{(bx+a)^2 \cos(2bx+2a)^2 + (bx+a)^2 \sin(2bx+2a)^2 + 2(bx+a)^2 \cos(2bx+2a) + (bx+a)^2 - (\cos(2bx+2a)^2 + \sin(2bx+2a)^2)}{\cos(2bx+2a)^2 + \sin(2bx+2a)^2}}{2 b^2}$$

[In] `integrate(x*tan(b*x+a)^2,x, algorithm="maxima")`

[Out] `1/2*(2*(b*x + a - tan(b*x + a))*a - ((b*x + a)^2*cos(2*b*x + 2*a)^2 + (b*x + a)^2*sin(2*b*x + 2*a)^2 + 2*(b*x + a)^2*cos(2*b*x + 2*a) + (b*x + a)^2 - (cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 4*(b*x + a)*sin(2*b*x + 2*a))/(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1))/b^2`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 162 vs. $2(28) = 56$.

Time = 0.60 (sec) , antiderivative size = 162, normalized size of antiderivative = 5.40

$$\int x \tan^2(a + bx) dx = -\frac{b^2 x^2 \tan(bx) \tan(a) - b^2 x^2 + 2bx \tan(bx) + 2bx \tan(a) - \log\left(\frac{4(\tan(bx)^2 \tan(a)^2 - 2\tan(bx) \tan(a) + 1)}{\tan(bx)^2 \tan(a)^2 + \tan(bx)^2 + \tan(a)^2 + 1}\right) \tan(bx) + 2(b^2 \tan(bx) \tan(a) - b^2)}{2(b^2 \tan(bx) \tan(a) - b^2)}$$

[In] `integrate(x*tan(b*x+a)^2,x, algorithm="giac")`

[Out]
$$\begin{aligned} & -1/2*(b^2*x^2*tan(b*x)*tan(a) - b^2*x^2 + 2*b*x*tan(b*x) + 2*b*x*tan(a) - \log(4*(\tan(b*x)^2*\tan(a)^2 - 2*\tan(b*x)*\tan(a) + 1)/(\tan(b*x)^2*\tan(a)^2 - 2*\tan(b*x)*\tan(a) + 1))) * \tan(b*x) + \\ & \log(4*(\tan(b*x)^2*\tan(a)^2 - 2*\tan(b*x)*\tan(a) + 1)/(\tan(b*x)^2*\tan(a)^2 + \tan(b*x)^2 + \tan(a)^2 + 1)))/(b^2*tan(b*x)*tan(a) - b^2) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 0.16 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.17

$$\int x \tan^2(a + bx) dx = -\frac{\ln(\tan(a+bx)^2+1)}{2} - \frac{bx \tan(a+bx)}{b^2} - \frac{x^2}{2}$$

[In] `int(x*tan(a + b*x)^2,x)`

[Out]
$$-\log(\tan(a + b*x)^2 + 1)/2 - b*x*tan(a + b*x))/b^2 - x^2/2$$

3.9 $\int \frac{\tan^2(a+bx)}{x} dx$

Optimal result	79
Rubi [N/A]	79
Mathematica [N/A]	80
Maple [N/A] (verified)	80
Fricas [N/A]	80
Sympy [N/A]	80
Maxima [N/A]	81
Giac [N/A]	81
Mupad [N/A]	81

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a + bx)}{x} dx = \text{Int}\left(\frac{\tan^2(a + bx)}{x}, x\right)$$

[Out] Unintegrable(tan(b*x+a)^2/x,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

[In] Int[Tan[a + b*x]^2/x, x]

[Out] Defer[Int][Tan[a + b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\tan^2(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 3.22 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

[In] `Integrate[Tan[a + b*x]^2/x, x]`

[Out] `Integrate[Tan[a + b*x]^2/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.34 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(bx + a)}{x} dx$$

[In] `int(tan(b*x+a)^2/x,x)`

[Out] `int(tan(b*x+a)^2/x,x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(bx + a)}{x} dx$$

[In] `integrate(tan(b*x+a)^2/x,x, algorithm="fricas")`

[Out] `integral(tan(b*x + a)^2/x, x)`

Sympy [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan^2(a + bx)}{x} dx$$

[In] `integrate(tan(b*x+a)**2/x,x)`

[Out] `Integral(tan(a + b*x)**2/x, x)`

Maxima [N/A]

Not integrable

Time = 0.96 (sec) , antiderivative size = 243, normalized size of antiderivative = 20.25

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

[In] `integrate(tan(b*x+a)^2/x,x, algorithm="maxima")`

[Out] `-(b*x*cos(2*b*x + 2*a)^2*log(x) + b*x*log(x)*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a)*log(x) + b*x*log(x) - 2*(b^2*x*cos(2*b*x + 2*a)^2 + b^2*x*sin(2*b*x + 2*a)^2 + 2*b^2*x*cos(2*b*x + 2*a) + b^2*x)*integrate(sin(2*b*x + 2*a)/(b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2), x) - 2*sin(2*b*x + 2*a))/(b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + b*x)`

Giac [N/A]

Not integrable

Time = 0.63 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(bx + a)^2}{x} dx$$

[In] `integrate(tan(b*x+a)^2/x,x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 3.57 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x} dx = \int \frac{\tan(a + b x)^2}{x} dx$$

[In] `int(tan(a + b*x)^2/x,x)`

[Out] `int(tan(a + b*x)^2/x, x)`

3.10 $\int \frac{\tan^2(a+bx)}{x^2} dx$

Optimal result	82
Rubi [N/A]	82
Mathematica [N/A]	83
Maple [N/A] (verified)	83
Fricas [N/A]	83
Sympy [N/A]	83
Maxima [N/A]	84
Giac [N/A]	84
Mupad [N/A]	84

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan^2(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable(tan(b*x+a)^2/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

[In] Int[Tan[a + b*x]^2/x^2,x]

[Out] Defer[Int][Tan[a + b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tan^2(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 2.44 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

[In] `Integrate[Tan[a + b*x]^2/x^2, x]`

[Out] `Integrate[Tan[a + b*x]^2/x^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.31 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(bx + a)}{x^2} dx$$

[In] `int(tan(b*x+a)^2/x^2, x)`

[Out] `int(tan(b*x+a)^2/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(bx + a)}{x^2} dx$$

[In] `integrate(tan(b*x+a)^2/x^2, x, algorithm="fricas")`

[Out] `integral(tan(b*x + a)^2/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan^2(a + bx)}{x^2} dx$$

[In] `integrate(tan(b*x+a)**2/x**2, x)`

[Out] `Integral(tan(a + b*x)**2/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 250, normalized size of antiderivative = 20.83

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

[In] `integrate(tan(b*x+a)^2/x^2,x, algorithm="maxima")`

[Out] `(b*x*cos(2*b*x + 2*a)^2 + b*x*sin(2*b*x + 2*a)^2 + 2*b*x*cos(2*b*x + 2*a) + b*x + 4*(b^2*x^2*cos(2*b*x + 2*a)^2 + b^2*x^2*sin(2*b*x + 2*a)^2 + 2*b^2*x^2*cos(2*b*x + 2*a) + b^2*x^2)*integrate(sin(2*b*x + 2*a)/(b^2*x^3*cos(2*b*x + 2*a)^2 + b^2*x^3*sin(2*b*x + 2*a)^2 + 2*b^2*x^3*cos(2*b*x + 2*a) + b^2*x^3), x) + 2*sin(2*b*x + 2*a))/(b*x^2*cos(2*b*x + 2*a)^2 + b*x^2*sin(2*b*x + 2*a)^2 + 2*b*x^2*cos(2*b*x + 2*a) + b*x^2)`

Giac [N/A]

Not integrable

Time = 0.67 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^2}{x^2} dx$$

[In] `integrate(tan(b*x+a)^2/x^2,x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 2.93 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(a + bx)}{x^2} dx = \int \frac{\tan(a + b x)^2}{x^2} dx$$

[In] `int(tan(a + b*x)^2/x^2,x)`

[Out] `int(tan(a + b*x)^2/x^2, x)`

3.11 $\int x^3 \tan^3(a + bx) dx$

Optimal result	85
Rubi [A] (verified)	85
Mathematica [A] (verified)	89
Maple [A] (verified)	89
Fricas [B] (verification not implemented)	90
Sympy [F]	90
Maxima [B] (verification not implemented)	90
Giac [F]	91
Mupad [F(-1)]	92

Optimal result

Integrand size = 12, antiderivative size = 205

$$\begin{aligned} \int x^3 \tan^3(a + bx) dx = & \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} \\ & + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} \\ & - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\ & + \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} - \frac{3x^2 \tan(a + bx)}{2b^2} + \frac{x^3 \tan^2(a + bx)}{2b} \end{aligned}$$

[Out] $\frac{3/2*I*x^2/b^2+1/2*x^3/b-1/4*I*x^4-3*x*ln(1+exp(2*I*(b*x+a)))/b^3+x^3*ln(1+exp(2*I*(b*x+a)))/b+3/2*I*polylog(2,-exp(2*I*(b*x+a)))/b^4-3/2*I*x^2*polylog(2,-exp(2*I*(b*x+a)))/b^2+3/2*x*polylog(3,-exp(2*I*(b*x+a)))/b^3+3/4*I*polylog(4,-exp(2*I*(b*x+a)))/b^4-3/2*x^2*tan(b*x+a)/b^2+1/2*x^3*tan(b*x+a)^2/b}{b}$

Rubi [A] (verified)

Time = 0.36 (sec), antiderivative size = 205, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.833, Rules used = {3801, 3800, 2221, 2317, 2438, 30, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int x^3 \tan^3(a + bx) dx = & \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} + \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} \\ & + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} \\ & - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{3x^2 \tan(a + bx)}{2b^2} \\ & + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{x^3 \tan^2(a + bx)}{2b} + \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} \end{aligned}$$

[In] $\text{Int}[x^3 \tan[a + b*x]^3, x]$

[Out] $\frac{((3\pi)/2)x^2}{b^2} + \frac{x^3}{(2b)} - \frac{(\pi/4)x^4}{(1 + E^{((2\pi)(a + b*x))})} - \frac{(3x \log[1 + E^{((2\pi)(a + b*x))}])}{b^3} + \frac{(x^3 \log[1 + E^{((2\pi)(a + b*x))}])}{b^4} + \frac{((3\pi)/2) \text{PolyLog}[2, -E^{((2\pi)(a + b*x))}]}{b^2} - \frac{((3\pi)/2)x^2 \text{PolyLog}[2, -E^{((2\pi)(a + b*x))}]}{(2b^3)} + \frac{((3\pi)/4) \text{PolyLog}[4, -E^{((2\pi)(a + b*x))}]}{b^4} - \frac{(3x^2 \tan[a + b*x])}{(2b^2)} + \frac{(x^3 \tan[a + b*x]^2)}{(2b)}$

Rule 30

$\text{Int}[(x^m), x_{\text{Symbol}}] \rightarrow \text{Simp}[x^{(m+1)/(m+1)}, x] /; \text{FreeQ}[m, x] \& \& \text{NeQ}[m, -1]$

Rule 2221

$\text{Int}[((F(g_*(e_*(c_*(d_*(x_))))^n_* + f_*(x_))))^m_* + ((a_*) + (b_*)^m_* \text{Log}[F]) * \text{Log}[1 + b*((F(g*(e + f*x)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{m-1} * \text{Log}[1 + b*((F(g*(e + f*x)))^n/a)], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \& \& \text{IGtQ}[m, 0]$

Rule 2317

$\text{Int}[\text{Log}[(a_*) + (b_*)^m_* \text{Log}[F]], x_{\text{Symbol}}] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F(e*(c + d*x)))^n], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \& \& \text{GtQ}[a, 0]$

Rule 2320

$\text{Int}[u, x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \& \& \text{!MatchQ}[u, (w_*)^m_* ((a_*)^n_* (v_*)^m_*)^n_*] /; \text{FreeQ}[\{a, m, n\}, x] \& \& \text{IntegerQ}[m*n] \& \& \text{!MatchQ}[u, E^((c_*)^n_* ((a_*) + (b_*)^n_*)^m_*)^n_*] /; \text{FreeQ}[\{a, b, c\}, x] \& \& \text{InverseFunctionQ}[F[x]]]$

Rule 2438

$\text{Int}[\text{Log}[(c_*)^m_* ((d_*) + (e_*)^n_* (x_*)^m_*)^n_*]/(x_), x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \& \& \text{EqQ}[c*d, 1]$

Rule 2611

$\text{Int}[\text{Log}[1 + (e_*)^m_* ((F(g_*(c_*(a_*) + (b_*)^n_* (x_*)^m_*))^n_*)^m_* + (f_*)^m_* (g_*)^n_* (x_*)^m_*)^n_*], x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(f + g*x)^m) * \text{PolyLog}[2, (-e)*(F(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F]), x] + \text{Dist}[g*(m/(b*c*n*\text{Log}[F])), \text{Int}[(f + g*x)^{m-1} * \text{PolyLog}[2, (-e)*(F(c*(a + b*x)))^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e,$

```
f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x]; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^n_, x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]); FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_ + (b_)*(x_))^p]/((d_)*(e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x]; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*((F_)^((c_)*(a_ + (b_)*(x_))))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x]; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^3 \tan^2(a + bx)}{2b} - \frac{3 \int x^2 \tan^2(a + bx) dx}{2b} - \int x^3 \tan(a + bx) dx \\
&= -\frac{ix^4}{4} - \frac{3x^2 \tan(a + bx)}{2b^2} + \frac{x^3 \tan^2(a + bx)}{2b} \\
&\quad + 2i \int \frac{e^{2i(a+bx)} x^3}{1 + e^{2i(a+bx)}} dx + \frac{3 \int x \tan(a + bx) dx}{b^2} + \frac{3 \int x^2 dx}{2b} \\
&= \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} - \frac{3x^2 \tan(a + bx)}{2b^2} \\
&\quad + \frac{x^3 \tan^2(a + bx)}{2b} - \frac{(6i) \int \frac{e^{2i(a+bx)} x}{1 + e^{2i(a+bx)}} dx}{b^2} - \frac{3 \int x^2 \log(1 + e^{2i(a+bx)}) dx}{b}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} \\
&\quad - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{3x^2 \tan(a+bx)}{2b^2} + \frac{x^3 \tan^2(a+bx)}{2b} \\
&\quad + \frac{3 \int \log(1 + e^{2i(a+bx)}) dx}{b^3} + \frac{(3i) \int x \operatorname{PolyLog}(2, -e^{2i(a+bx)}) dx}{b^2} \\
&= \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} \\
&\quad + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3x^2 \tan(a+bx)}{2b^2} + \frac{x^3 \tan^2(a+bx)}{2b} \\
&\quad - \frac{(3i) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(a+bx)}\right)}{2b^4} - \frac{3 \int \operatorname{PolyLog}(3, -e^{2i(a+bx)}) dx}{2b^3} \\
&= \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} \\
&\quad + \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\
&\quad + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{3x^2 \tan(a+bx)}{2b^2} \\
&\quad + \frac{x^3 \tan^2(a+bx)}{2b} + \frac{(3i) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(3, -x)}{x} dx, x, e^{2i(a+bx)}\right)}{4b^4} \\
&= \frac{3ix^2}{2b^2} + \frac{x^3}{2b} - \frac{ix^4}{4} - \frac{3x \log(1 + e^{2i(a+bx)})}{b^3} \\
&\quad + \frac{x^3 \log(1 + e^{2i(a+bx)})}{b} + \frac{3i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^4} \\
&\quad - \frac{3ix^2 \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} + \frac{3x \operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\
&\quad + \frac{3i \operatorname{PolyLog}(4, -e^{2i(a+bx)})}{4b^4} - \frac{3x^2 \tan(a+bx)}{2b^2} + \frac{x^3 \tan^2(a+bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 6.54 (sec) , antiderivative size = 370, normalized size of antiderivative = 1.80

$$\int x^3 \tan^3(a + bx) dx$$

$$= \frac{i e^{ia} (2 b^4 e^{-2 i a} x^4 - 4 i b^3 (1 + e^{-2 i a}) x^3 \log(1 + e^{-2 i (a + b x)}) + 6 b^2 (1 + e^{-2 i a}) x^2 \text{PolyLog}(2, -e^{-2 i (a + b x)}) - 6 i b}{8 b^4}$$

$$+ \frac{x^3 \sec^2(a + bx)}{2b}$$

$$- \frac{3 \csc(a) \left(b^2 e^{-i \arctan(\cot(a))} x^2 - \frac{\cot(a) (ibx(-\pi - 2 \arctan(\cot(a))) - \pi \log(1 + e^{-2 i b x}) - 2(bx - \arctan(\cot(a))) \log(1 - e^{2 i (bx - \arctan(\cot(a)))})}{2b^4 \sqrt{\csc^2(a) (\cos^2(a) + \sin^2(a))}}\right)}{2b^2}$$

$$- \frac{3 x^2 \sec(a) \sec(a + bx) \sin(bx)}{2b^2} - \frac{1}{4} x^4 \tan(a)$$

[In] `Integrate[x^3*Tan[a + b*x]^3, x]`

[Out]
$$\begin{aligned} & ((I/8)*E^{-(I*a)*((2*b^4*x^4)/E^{((2*I)*a})} - (4*I)*b^3*(1 + E^{((-2*I)*a)})*x^3* \\ & \log[1 + E^{((-2*I)*(a + b*x))}] + 6*b^2*(1 + E^{((-2*I)*a)})*x^2*\text{PolyLog}[2, -E^{((-2*I)*(a + b*x))}] - (6*I)*b*(1 + E^{((-2*I)*a)})*x*\text{PolyLog}[3, -E^{((-2*I)*(a + b*x))}] - 3*(1 + E^{((-2*I)*a)})*\text{PolyLog}[4, -E^{((-2*I)*(a + b*x))}]*\text{Sec}[a]) \\ & /b^4 + (x^3*\text{Sec}[a + b*x]^2)/(2*b) - (3*\text{Csc}[a]*((b^2*x^2)/E^{(I*\text{ArcTan}[\text{Cot}[a]]}) - (\text{Cot}[a]*(I*b*x*(-Pi - 2*\text{ArcTan}[\text{Cot}[a]])) - \text{Pi}*\text{Log}[1 + E^{((-2*I)*b*x)}] - 2*(b*x - \text{ArcTan}[\text{Cot}[a]])*\text{Log}[1 - E^{((2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]))}]) + \text{Pi}*\text{Log}[\text{Cos}[b*x]] - 2*\text{ArcTan}[\text{Cot}[a]]*\text{Log}[\text{Sin}[b*x - \text{ArcTan}[\text{Cot}[a]]]] + I*\text{PolyLog}[2, E^{((2*I)*(b*x - \text{ArcTan}[\text{Cot}[a]))}]]))/\text{Sqrt}[1 + \text{Cot}[a]^2]*\text{Sec}[a]/(2*b^4*\text{Sqrt}[\text{Csc}[a]^2*(\text{Cos}[a]^2 + \text{Sin}[a]^2)]) - (3*x^2*\text{Sec}[a]*\text{Sec}[a + b*x]*\text{Sin}[b*x])/(2*b^2) - (x^4*\text{Tan}[a])/4 \end{aligned}$$

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 251, normalized size of antiderivative = 1.22

method	result
risch	$\frac{3i \text{Li}_2(-e^{2i(bx+a)})}{2b^4} + \frac{x^2(2bx e^{2i(bx+a)} - 3ie^{2i(bx+a)} - 3i)}{b^2(e^{2i(bx+a)} + 1)^2} + \frac{3ix^2}{b^2} - \frac{ix^4}{4} + \frac{3i \text{Li}_4(-e^{2i(bx+a)})}{4b^4} - \frac{3ia^4}{2b^4} - \frac{3ix^2 \text{Li}_2(-e^{2i(bx+a)})}{2b^2}$

[In] `int(x^3*tan(b*x+a)^3, x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 3/2*I*\text{polylog}(2, -\exp(2*I*(b*x+a)))/b^4 + x^2*(2*b*x*\exp(2*I*(b*x+a)) - 3*I*\exp(2*I*(b*x+a)) - 3*I)/b^2 + (\exp(2*I*(b*x+a)) + 1)^{-2 + 3*I}/b^2*x^2 - 1/4*I*x^4 + 3/4*I*\text{polylog}(4, -\exp(2*I*(b*x+a)))/b^4 - 3/2*I/b^4*a^4 - 3/2*I*x^2*\text{polylog}(2, -\exp(2*I*(b*x+a)))/b^2 + 6*I/b^3*a*x - 2*I/b^3*a^3*x - 6/b^4*a*\ln(\exp(I*(b*x+a))) + 2/b^4*a^3*\ln(\exp(I*(b*x+a))) + 3*I/b^4*a^2*x^3*\ln(\exp(2*I*(b*x+a)) + 1)/b + 3/2*x*\text{polylog}(3, -\exp(2*I*(b*x+a)))/b^3 - 3*x*\ln(\exp(2*I*(b*x+a)) + 1)/b^3 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 344 vs. $2(163) = 326$.

Time = 0.27 (sec) , antiderivative size = 344, normalized size of antiderivative = 1.68

$$\int x^3 \tan^3(a + bx) dx = \frac{4 b^3 x^3 \tan(bx + a)^2 + 4 b^3 x^3 - 12 b^2 x^2 \tan(bx + a) + 6 b x \text{polylog}\left(3, \frac{\tan(bx+a)^2+2i \tan(bx+a)-1}{\tan(bx+a)^2+1}\right) + 6 b x \text{polylog}(4, \frac{\tan(bx+a)^2+2i \tan(bx+a)-1}{\tan(bx+a)^2+1})}{b^4}$$

```
[In] integrate(x^3*tan(b*x+a)^3,x, algorithm="fricas")
[Out] 1/8*(4*b^3*x^3*tan(b*x + a)^2 + 4*b^3*x^3 - 12*b^2*x^2*tan(b*x + a) + 6*b*x
*polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) +
6*b*x*polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) -
6*(-I*b^2*x^2 + I)*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) -
6*(I*b^2*x^2 - I)*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) +
4*(b^3*x^3 - 3*b*x)*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) +
4*(b^3*x^3 - 3*b*x)*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) -
3*I*polylog(4, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) +
3*I*polylog(4, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^4
```

Sympy [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan^3(a + bx) dx$$

```
[In] integrate(x**3*tan(b*x+a)**3,x)
[Out] Integral(x**3*tan(a + b*x)**3, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1205 vs. $2(163) = 326$.

Time = 0.47 (sec) , antiderivative size = 1205, normalized size of antiderivative = 5.88

$$\int x^3 \tan^3(a + bx) dx = \text{Too large to display}$$

```
[In] integrate(x^3*tan(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] 1/2*(a^3*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) - 2*(3*(b*x + a)^4 - 12*(b*x + a)^3*a + 18*(b*x + a)^2*a^2 + 36*a^2 - 4*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*cos(2*b*x + 2*a) - (-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a) - 9*I*a)*sin(4*b*x + 4*a) - 2*(-4*I*(b*x + a)^3 + 9*I*(b*x + a)^2*a + 9*(-I*a^2 + I)*(b*x + a) - 9*I*a)*sin(2*b*x + 2*a) + 9*a)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 3*((b*x + a)^4 - 4*(b*x + a)^3*a + 6*(a^2 - 2)*(b*x + a)^2 + 24*(b*x + a)*a)*cos(4*b*x + 4*a) + 6*((b*x + a)^4 - 4*(b*x + a)^3*(a - I) + 6*(a^2 - 2*I*a - 1)*(b*x + a)^2 + 12*(I*a^2 + a)*(b*x + a) + 6*a^2)*cos(2*b*x + 2*a) + 6*(4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 + (4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 - 3)*cos(4*b*x + 4*a) + 2*(4*(b*x + a)^2 - 6*(b*x + a)*a + 3*a^2 - 3)*cos(2*b*x + 2*a) + (4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2 - 3*I)*sin(4*b*x + 4*a) + 2*(4*I*(b*x + a)^2 - 6*I*(b*x + a)*a + 3*I*a^2 - 3*I)*sin(2*b*x + 2*a) - 3)*dilog(-e^(2*I*b*x + 2*I*a)) + 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + (4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*I*a)*cos(4*b*x + 4*a) + 2*(4*I*(b*x + a)^3 - 9*I*(b*x + a)^2*a + 9*(I*a^2 - I)*(b*x + a) + 9*I*a)*cos(2*b*x + 2*a) - (4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*sin(4*b*x + 4*a) - 2*(4*(b*x + a)^3 - 9*(b*x + a)^2*a + 9*(a^2 - 1)*(b*x + a) + 9*a)*sin(2*b*x + 2*a) + 9*I*a)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) - 12*(cos(4*b*x + 4*a) + 2*cos(2*b*x + 2*a) + I*sin(4*b*x + 4*a) + 2*I*sin(2*b*x + 2*a) + 1)*polylog(4, -e^(2*I*b*x + 2*I*a)) + 6*(4*I*b*x + (4*I*b*x + I*a)*cos(4*b*x + 4*a) + 2*(4*I*b*x + I*a)*cos(2*b*x + 2*a) - (4*b*x + a)*sin(4*b*x + 4*a) - 2*(4*b*x + a)*sin(2*b*x + 2*a) + I*a)*polylog(3, -e^(2*I*b*x + 2*I*a)) + 3*(I*(b*x + a)^4 - 4*I*(b*x + a)^3*a + 6*(I*a^2 - 2*I)*(b*x + a)^2 + 24*I*(b*x + a)*a)*sin(4*b*x + 4*a) + 6*(I*(b*x + a)^4 + 4*(b*x + a)^3*(-I*a - 1) + 6*(I*a^2 + 2*a - I)*(b*x + a)^2 - 12*(a^2 - I*a)*(b*x + a) + 6*I*a^2)*sin(2*b*x + 2*a))/(-12*I*cos(4*b*x + 4*a) - 24*I*cos(2*b*x + 2*a) + 12*sin(4*b*x + 4*a) + 24*sin(2*b*x + 2*a) - 12*I))/b^4
```

Giac [F]

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(bx + a)^3 dx$$

```
[In] integrate(x^3*tan(b*x+a)^3,x, algorithm="giac")
```

```
[Out] integrate(x^3*tan(b*x + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \tan^3(a + bx) dx = \int x^3 \tan(a + bx)^3 dx$$

[In] int(x^3*tan(a + b*x)^3, x)

[Out] int(x^3*tan(a + b*x)^3, x)

3.12 $\int x^2 \tan^3(a + bx) dx$

Optimal result	93
Rubi [A] (verified)	93
Mathematica [A] (verified)	96
Maple [A] (verified)	96
Fricas [B] (verification not implemented)	96
Sympy [F]	97
Maxima [B] (verification not implemented)	97
Giac [F]	98
Mupad [F(-1)]	98

Optimal result

Integrand size = 12, antiderivative size = 128

$$\begin{aligned} \int x^2 \tan^3(a + bx) dx = & \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} \\ & - \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} + \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} \\ & - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} \end{aligned}$$

[Out] $1/2*x^2/b - 1/3*I*x^3 + x^2*\ln(1+\exp(2*I*(b*x+a)))/b - \ln(\cos(b*x+a))/b^3 - I*x*\operatorname{polylog}(2, -\exp(2*I*(b*x+a)))/b^2 + 1/2*\operatorname{polylog}(3, -\exp(2*I*(b*x+a)))/b^3 - x*\tan(b*x+a)/b^2 + 1/2*x^2*\tan(b*x+a)^2/b$

Rubi [A] (verified)

Time = 0.22 (sec), antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.667, Rules used = {3801, 3556, 30, 3800, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int x^2 \tan^3(a + bx) dx = & \frac{\operatorname{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{\log(\cos(a + bx))}{b^3} \\ & - \frac{ix \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{b^2} - \frac{x \tan(a + bx)}{b^2} \\ & + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} + \frac{x^2 \tan^2(a + bx)}{2b} + \frac{x^2}{2b} - \frac{ix^3}{3} \end{aligned}$$

[In] $\operatorname{Int}[x^2 \tan[a + b*x]^3, x]$

[Out] $x^{2/(2*b)} - (I/3)*x^3 + (x^{2*\text{Log}[1 + E^{((2*I)*(a + b*x))}]} / b - \text{Log}[\cos[a + b*x]] / b^3 - (I*x*\text{PolyLog}[2, -E^{((2*I)*(a + b*x))}] / b^2 + \text{PolyLog}[3, -E^{((2*I)*(a + b*x))}] / (2*b^3) - (x*\text{Tan}[a + b*x]) / b^2 + (x^{2*\text{Tan}[a + b*x]^2}) / (2*b)$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*(c_.) + (d_.)*(x_))^(m_.)) / ((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m / (b*f*g*n*Log[F])) * Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m / (b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1) * Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*(F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n] / (b*c*n*Log[F])), x] + Dist[g*(m / (b*c*n*Log[F])), Int[(f + g*x)^(m - 1) * PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1) / (d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m * (E^(2*I*(e + f*x))) / (1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```

Int[((c_) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symb
o1] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

```

Rule 6724

```

Int[PolyLog[n_, (c_)*(a_ + b*x)^p]/((d_*) + (e_)*(x_)), x_Symb
ybol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
, e, n, p}, x] && EqQ[b*d, a*e]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x^2 \tan^2(a + bx)}{2b} - \frac{\int x \tan^2(a + bx) dx}{b} - \int x^2 \tan(a + bx) dx \\
&= -\frac{ix^3}{3} - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} + 2i \int \frac{e^{2i(a+bx)} x^2}{1 + e^{2i(a+bx)}} dx + \frac{\int \tan(a + bx) dx}{b^2} \\
&\quad + \frac{\int x dx}{b} \\
&= \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} \\
&\quad - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} - \frac{2 \int x \log(1 + e^{2i(a+bx)}) dx}{b} \\
&= \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} - \frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^2} \\
&\quad - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} + \frac{i \int \text{PolyLog}(2, -e^{2i(a+bx)}) dx}{b^2} \\
&= \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} - \frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^2} \\
&\quad - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b} + \frac{\text{Subst}\left(\int \frac{\text{PolyLog}(2, -e^{2i(a+bx)})}{x} dx, x, e^{2i(a+bx)}\right)}{2b^3} \\
&= \frac{x^2}{2b} - \frac{ix^3}{3} + \frac{x^2 \log(1 + e^{2i(a+bx)})}{b} - \frac{\log(\cos(a + bx))}{b^3} - \frac{ix \text{PolyLog}(2, -e^{2i(a+bx)})}{b^2} \\
&\quad + \frac{\text{PolyLog}(3, -e^{2i(a+bx)})}{2b^3} - \frac{x \tan(a + bx)}{b^2} + \frac{x^2 \tan^2(a + bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.28 (sec) , antiderivative size = 172, normalized size of antiderivative = 1.34

$$\int x^2 \tan^3(a + bx) dx \\ = \frac{e^{-ia} (2b^2 x^2 (2ibx + 3(1 + e^{2ia}) \log(1 + e^{-2i(a+bx)})) + 6ib(1 + e^{2ia}) x \text{PolyLog}(2, -e^{-2i(a+bx)}) + 3(1 + e^{2ia}) \text{PolyLog}(3, -e^{-2i(a+bx)}))}{12b^3}$$

[In] `Integrate[x^2*Tan[a + b*x]^3, x]`

[Out] $\frac{((2*b^2*x^2*((2*I)*b*x + 3*(1 + E^((2*I)*a))*Log[1 + E^((-2*I)*(a + b*x))]) + (6*I)*b*(1 + E^((2*I)*a))*x*PolyLog[2, -E^((-2*I)*(a + b*x))]) + 3*(1 + E^((2*I)*a))*PolyLog[3, -E^((-2*I)*(a + b*x))])*Sec[a])/E^(I*a) + 6*b^2*x^2*Sec[a + b*x]^2 - 12*b*x*Sec[a]*Sec[a + b*x]*Sin[b*x] - 4*b^3*x^3*Tan[a] - 12*(Log[Cos[a + b*x]] + b*x*Tan[a]))/(12*b^3)$

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 180, normalized size of antiderivative = 1.41

method	result
risch	$-\frac{ix^3}{3} + \frac{2x(bx e^{2i(bx+a)} - ie^{2i(bx+a)} - i)}{b^2(e^{2i(bx+a)} + 1)^2} - \frac{2a^2 \ln(e^{i(bx+a)})}{b^3} + \frac{4ia^3}{3b^3} + \frac{2ia^2x}{b^2} + \frac{x^2 \ln(e^{2i(bx+a)} + 1)}{b} - \frac{ix \text{Li}_2(-e^{2i(bx+a)})}{b^2} +$

[In] `int(x^2*tan(b*x+a)^3, x, method=_RETURNVERBOSE)`

[Out]
$$\frac{-1/3*I*x^3+2*x*(b*x*exp(2*I*(b*x+a))-I*exp(2*I*(b*x+a))-I)/b^2/(exp(2*I*(b*x+a))+1)^2-2/b^3*a^2*ln(exp(I*(b*x+a)))+4/3*I/b^3*a^3+2*I/b^2*a^2*x+x^2*ln(exp(2*I*(b*x+a))+1)/b-I*x*polylog(2, -exp(2*I*(b*x+a)))/b^2+1/2*polylog(3, -exp(2*I*(b*x+a)))/b^3-1/b^3*ln(exp(2*I*(b*x+a))+1)+2/b^3*ln(exp(I*(b*x+a)))}{b^3}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(109) = 218.

Time = 0.26 (sec) , antiderivative size = 240, normalized size of antiderivative = 1.88

$$\int x^2 \tan^3(a + bx) dx \\ = \frac{2 b^2 x^2 \tan(bx + a)^2 + 2 b^2 x^2 + 2 i b x \text{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1} + 1\right) - 2 i b x \text{Li}_2\left(\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1} + 1\right) - 4 b x \tan(bx + a)^2}{12 b^3}$$

[In] `integrate(x^2*tan(b*x+a)^3, x, algorithm="fricas")`

```
[Out] 1/4*(2*b^2*x^2*tan(b*x + a)^2 + 2*b^2*x^2 + 2*I*b*x*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*I*b*x*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 4*b*x*tan(b*x + a) + 2*(b^2*x^2 - 1)*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*(b^2*x^2 - 1)*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + polylog(3, (tan(b*x + a)^2 + 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + polylog(3, (tan(b*x + a)^2 - 2*I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)))/b^3
```

Sympy [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan^3(a + bx) dx$$

```
[In] integrate(x**2*tan(b*x+a)**3,x)
[Out] Integral(x**2*tan(a + b*x)**3, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 736 vs. 2(109) = 218.

Time = 0.66 (sec), antiderivative size = 736, normalized size of antiderivative = 5.75

$$\int x^2 \tan^3(a + bx) dx = \text{Too large to display}$$

```
[In] integrate(x^2*tan(b*x+a)^3,x, algorithm="maxima")
[Out] -1/2*(a^2*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1)) + 2*(2*(b*x + a)^3 - 6*(b*x + a)^2*a - 6*((b*x + a)^2 - 2*(b*x + a)*a + ((b*x + a)^2 - 2*(b*x + a)*a - 1)*cos(4*b*x + 4*a) + 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*cos(2*b*x + 2*a) - (-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(4*b*x + 4*a) - 2*(-I*(b*x + a)^2 + 2*I*(b*x + a)*a + I)*sin(2*b*x + 2*a) - 1)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) + 2*((b*x + a)^3 - 3*(b*x + a)^2*a - 6*b*x - 6*a)*cos(4*b*x + 4*a) + 4*((b*x + a)^3 - 3*(b*x + a)^2*a - (b*x + a)*(-2*I*a - 1) - 3*a)*cos(2*b*x + 2*a) + 6*(b*x*cos(4*b*x + 4*a) + 2*b*x*cos(2*b*x + 2*a) + I*b*x*sin(4*b*x + 4*a) + 2*I*b*x*sin(2*b*x + 2*a) + b*x)*dilog(-e^(2*I*b*x + 2*I*a)) + 3*(I*(b*x + a)^2 - 2*I*(b*x + a)*a + (I*(b*x + a)^2 - 2*I*(b*x + a)*a - I)*cos(4*b*x + 4*a) + 2*(I*(b*x + a)^2 - 2*I*(b*x + a)*a - I)*cos(2*b*x + 2*a) - ((b*x + a)^2 - 2*(b*x + a)*a - 1)*sin(4*b*x + 4*a) - 2*((b*x + a)^2 - 2*(b*x + a)*a - 1)*sin(2*b*x + 2*a) - I)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 3*(I*cos(4*b*x + 4*a) + 2*I*cos(2*b*x + 2*a) - sin(4*b*x + 4*a) - 2*sin(2*b*x + 2*a))
```

```
2*a) + I)*polylog(3, -e^(2*I*b*x + 2*I*a)) + 2*(I*(b*x + a)^3 - 3*I*(b*x + a)^2*a - 6*I*b*x - 6*I*a)*sin(4*b*x + 4*a) + 4*(I*(b*x + a)^3 + 3*(b*x + a)^2*(-I*a - 1) + 3*(b*x + a)*(2*a - I) - 3*I*a)*sin(2*b*x + 2*a) - 12*a)/(-6*I*cos(4*b*x + 4*a) - 12*I*cos(2*b*x + 2*a) + 6*sin(4*b*x + 4*a) + 12*sin(2*b*x + 2*a) - 6*I))/b^3
```

Giac [F]

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(bx + a)^3 dx$$

```
[In] integrate(x^2*tan(b*x+a)^3,x, algorithm="giac")
[Out] integrate(x^2*tan(b*x + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \tan^3(a + bx) dx = \int x^2 \tan(a + b x)^3 dx$$

```
[In] int(x^2*tan(a + b*x)^3,x)
[Out] int(x^2*tan(a + b*x)^3, x)
```

3.13 $\int x \tan^3(a + bx) dx$

Optimal result	99
Rubi [A] (verified)	99
Mathematica [A] (verified)	101
Maple [A] (verified)	101
Fricas [B] (verification not implemented)	102
Sympy [F]	102
Maxima [B] (verification not implemented)	102
Giac [F]	103
Mupad [F(-1)]	103

Optimal result

Integrand size = 10, antiderivative size = 90

$$\begin{aligned} \int x \tan^3(a + bx) dx = & \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} \\ & - \frac{\tan(a + bx)}{2b^2} + \frac{x \tan^2(a + bx)}{2b} \end{aligned}$$

[Out] $1/2*x/b - 1/2*I*x^2 + x*ln(1 + exp(2*I*(b*x+a)))/b - 1/2*I*polylog(2, -exp(2*I*(b*x+a)))/b^2 - 1/2*tan(b*x+a)/b^2 + 1/2*x*tan(b*x+a)^2/b$

Rubi [A] (verified)

Time = 0.14 (sec), antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3801, 3554, 8, 3800, 2221, 2317, 2438}

$$\begin{aligned} \int x \tan^3(a + bx) dx = & -\frac{i \operatorname{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{\tan(a + bx)}{2b^2} \\ & + \frac{x \log(1 + e^{2i(a+bx)})}{b} + \frac{x \tan^2(a + bx)}{2b} + \frac{x}{2b} - \frac{ix^2}{2} \end{aligned}$$

[In] $\operatorname{Int}[x*Tan[a + b*x]^3, x]$

[Out] $x/(2*b) - (I/2)*x^2 + (x*Log[1 + E^((2*I)*(a + b*x))])/b - ((I/2)*PolyLog[2, -E^((2*I)*(a + b*x))])/b^2 - Tan[a + b*x]/(2*b^2) + (x*Tan[a + b*x]^2)/(2*b)$

Rule 8

$\operatorname{Int}[a_, x_Symbol] := \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_)*(x_))))^(n_.)*((c_.) + (d_)*(x_))^(m_.))/((a_.) + (b_)*(F_)^((g_.)*(e_.) + (f_)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_.) + (b_)*(F_)^((e_.)*(c_.) + (d_)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3554

```
Int[((b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3800

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x \tan^2(a + bx)}{2b} - \frac{\int \tan^2(a + bx) dx}{2b} - \int x \tan(a + bx) dx \\ &= -\frac{ix^2}{2} - \frac{\tan(a + bx)}{2b^2} + \frac{x \tan^2(a + bx)}{2b} + 2i \int \frac{e^{2i(a+bx)}x}{1 + e^{2i(a+bx)}} dx + \frac{\int 1 dx}{2b} \end{aligned}$$

$$\begin{aligned}
&= \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{\tan(a+bx)}{2b^2} + \frac{x \tan^2(a+bx)}{2b} - \frac{\int \log(1 + e^{2i(a+bx)}) dx}{b} \\
&= \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{\tan(a+bx)}{2b^2} \\
&\quad + \frac{x \tan^2(a+bx)}{2b} + \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(a+bx)}\right)}{2b^2} \\
&= \frac{x}{2b} - \frac{ix^2}{2} + \frac{x \log(1 + e^{2i(a+bx)})}{b} - \frac{i \text{PolyLog}(2, -e^{2i(a+bx)})}{2b^2} - \frac{\tan(a+bx)}{2b^2} + \frac{x \tan^2(a+bx)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 4.58 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.90

$$\begin{aligned}
&\int x \tan^3(a+bx) dx \\
&= \frac{ibx(\pi + 2 \arctan(\cot(a))) + \pi \log(1 + e^{-2ibx}) + 2(bx - \arctan(\cot(a))) \log(1 - e^{2i(bx - \arctan(\cot(a)))}) - \pi \log(1 - e^{2i(bx - \arctan(\cot(a)))})}{b^2}
\end{aligned}$$

[In] Integrate[x*Tan[a + b*x]^3, x]

[Out] $(I*b*x*(Pi + 2*ArcTan[Cot[a]]) + Pi*Log[1 + E^{(-2*I)*b*x}] + 2*(b*x - ArcTan[Cot[a]])*Log[1 - E^{((2*I)*(b*x - ArcTan[Cot[a]]))}] - Pi*Log[Cos[b*x]] + 2*ArcTan[Cot[a]]*Log[Sin[b*x - ArcTan[Cot[a]]]] - I*PolyLog[2, E^{((2*I)*(b*x - ArcTan[Cot[a]]))}] + b*x*Sec[a + b*x]^2 - Sec[a]*Sec[a + b*x]*Sin[b*x] - b^2*x^2*Tan[a] + (b^2*x^2*.Sqrt[Csc[a]^2]*Tan[a])/E^{(I*ArcTan[Cot[a]])})/(2*b^2)$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.36

method	result	size
risch	$-\frac{ix^2}{2} + \frac{2bx e^{2i(bx+a)} - ie^{2i(bx+a)} - i}{b^2(e^{2i(bx+a)} + 1)^2} - \frac{2i a x}{b} - \frac{ia^2}{b^2} + \frac{x \ln(e^{2i(bx+a)} + 1)}{b} - \frac{i \text{Li}_2(-e^{2i(bx+a)})}{2b^2} + \frac{2a \ln(e^{i(bx+a)})}{b^2}$	122

[In] int(x*tan(b*x+a)^3, x, method=_RETURNVERBOSE)

[Out] $-1/2*I*x^2 + (2*b*x*exp(2*I*(b*x+a)) - I*exp(2*I*(b*x+a)) - I)/b^2/(exp(2*I*(b*x+a)) + 1)^2 - 2*I/b*a*x - I/b^2*a^2*x*ln(exp(2*I*(b*x+a)) + 1)/b - 1/2*I*polylog(2, -exp(2*I*(b*x+a)))/b^2 + 2/b^2*a*ln(exp(I*(b*x+a)))$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 146 vs. $2(71) = 142$.

Time = 0.26 (sec) , antiderivative size = 146, normalized size of antiderivative = 1.62

$$\int x \tan^3(a + bx) dx = \frac{2 b x \tan(bx + a)^2 + 2 b x \log\left(-\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x \log\left(-\frac{2(-i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right) + 2 b x + i \operatorname{Li}_2\left(\frac{2(i \tan(bx + a) - 1)}{\tan(bx + a)^2 + 1}\right)}{4 b^2}$$

```
[In] integrate(x*tan(b*x+a)^3,x, algorithm="fricas")
```

```
[Out] 1/4*(2*b*x*tan(b*x + a)^2 + 2*b*x*log(-2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b*x*log(-2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1)) + 2*b*x + I*dilog(2*(I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - I*dilog(2*(-I*tan(b*x + a) - 1)/(tan(b*x + a)^2 + 1) + 1) - 2*tan(b*x + a))/b^2
```

Sympy [F]

$$\int x \tan^3(a + bx) dx = \int x \tan^3 (a + bx) dx$$

```
[In] integrate(x*tan(b*x+a)**3,x)
```

[Out] $\text{Integral}(x \cdot \tan(a + b \cdot x)^3, x)$

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 386 vs. $2(71) = 142$.

Time = 0.78 (sec) , antiderivative size = 386, normalized size of antiderivative = 4.29

$$\int x \tan^3(a + bx) dx = -\frac{b^2 x^2 \cos(4bx + 4a) + i b^2 x^2 \sin(4bx + 4a) + b^2 x^2 - 2(bx \cos(4bx + 4a) + 2bx \cos(2bx + 2a) + i bx \sin(2bx + 2a))}{16b^3}$$

```
[In] integrate(x*tan(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] -(b^2*x^2*cos(4*b*x + 4*a) + I*b^2*x^2*sin(4*b*x + 4*a) + b^2*x^2 - 2*(b*x*cos(4*b*x + 4*a) + 2*b*x*cos(2*b*x + 2*a) + I*b*x*sin(4*b*x + 4*a) + 2*I*b*x*sin(2*b*x + 2*a) + b*x)*arctan2(sin(2*b*x + 2*a), cos(2*b*x + 2*a) + 1) +
```

$$\begin{aligned}
& 2*(b^2*x^2 + 2*I*b*x + 1)*cos(2*b*x + 2*a) + (cos(4*b*x + 4*a) + 2*cos(2*b*x + 2*a) + I*sin(4*b*x + 4*a) + 2*I*sin(2*b*x + 2*a) + 1)*dilog(-e^{(2*I*b*x + 2*I*a)}) - (-I*b*x*cos(4*b*x + 4*a) - 2*I*b*x*cos(2*b*x + 2*a) + b*x*sin(4*b*x + 4*a) + 2*b*x*sin(2*b*x + 2*a) - I*b*x)*log(cos(2*b*x + 2*a)^2 + sin(2*b*x + 2*a)^2 + 2*cos(2*b*x + 2*a) + 1) + 2*(I*b^2*x^2 - 2*b*x + I)*sin(2*b*x + 2*a) + 2)/(-2*I*b^2*cos(4*b*x + 4*a) - 4*I*b^2*cos(2*b*x + 2*a) + 2*b^2*sin(4*b*x + 4*a) + 4*b^2*sin(2*b*x + 2*a) - 2*I*b^2)
\end{aligned}$$

Giac [F]

$$\int x \tan^3(a + bx) dx = \int x \tan(bx + a)^3 dx$$

[In] `integrate(x*tan(b*x+a)^3,x, algorithm="giac")`

[Out] `integrate(x*tan(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x \tan^3(a + bx) dx = \int x \tan(a + b x)^3 dx$$

[In] `int(x*tan(a + b*x)^3,x)`

[Out] `int(x*tan(a + b*x)^3, x)`

3.14 $\int \frac{\tan^3(a+bx)}{x} dx$

Optimal result	104
Rubi [N/A]	104
Mathematica [N/A]	105
Maple [N/A] (verified)	105
Fricas [N/A]	105
Sympy [N/A]	105
Maxima [N/A]	106
Giac [N/A]	106
Mupad [N/A]	107

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a + bx)}{x} dx = \text{Int}\left(\frac{\tan^3(a + bx)}{x}, x\right)$$

[Out] Unintegrable(tan(b*x+a)^3/x,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

[In] Int[Tan[a + b*x]^3/x,x]

[Out] Defer[Int][Tan[a + b*x]^3/x, x]

Rubi steps

$$\text{integral} = \int \frac{\tan^3(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 4.85 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

[In] `Integrate[Tan[a + b*x]^3/x, x]`

[Out] `Integrate[Tan[a + b*x]^3/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.17 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(bx + a)}{x} dx$$

[In] `int(tan(b*x+a)^3/x,x)`

[Out] `int(tan(b*x+a)^3/x,x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(bx + a)}{x} dx$$

[In] `integrate(tan(b*x+a)^3/x,x, algorithm="fricas")`

[Out] `integral(tan(b*x + a)^3/x, x)`

Sympy [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan^3(a + bx)}{x} dx$$

[In] `integrate(tan(b*x+a)**3/x,x)`

[Out] `Integral(tan(a + b*x)**3/x, x)`

Maxima [N/A]

Not integrable

Time = 0.91 (sec) , antiderivative size = 534, normalized size of antiderivative = 44.50

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

[In] `integrate(tan(b*x+a)^3/x, x, algorithm="maxima")`

[Out]
$$(4*b*x*\cos(2*b*x + 2*a)^2 + 4*b*x*\sin(2*b*x + 2*a)^2 + 2*b*x*\cos(2*b*x + 2*a) + (2*b*x*\cos(2*b*x + 2*a) - \sin(2*b*x + 2*a))*\cos(4*b*x + 4*a) - (b^2*x^2*\cos(4*b*x + 4*a)^2 + 4*b^2*x^2*\cos(2*b*x + 2*a)^2 + b^2*x^2*\sin(4*b*x + 4*a)^2 + 4*b^2*x^2*\sin(4*b*x + 4*a)*\sin(2*b*x + 2*a) + 4*b^2*x^2*\sin(2*b*x + 2*a)^2 + 4*b^2*x^2*\cos(2*b*x + 2*a) + b^2*x^2)*\cos(4*b*x + 4*a))*\integrate(2*(b^2*x^2 - 1)*\sin(2*b*x + 2*a)/(b^2*x^3*\cos(2*b*x + 2*a)^2 + b^2*x^3*\sin(2*b*x + 2*a)^2 + 2*b^2*x^3*\cos(2*b*x + 2*a) + b^2*x^3), x) + (2*b*x*\sin(2*b*x + 2*a) + \cos(2*b*x + 2*a) + 1)*\sin(4*b*x + 4*a) + \sin(2*b*x + 2*a))/(b^2*x^2*\cos(4*b*x + 4*a)^2 + 4*b^2*x^2*\cos(2*b*x + 2*a)^2 + b^2*x^2*\sin(4*b*x + 4*a)^2 + 4*b^2*x^2*\sin(4*b*x + 4*a)*\sin(2*b*x + 2*a) + 4*b^2*x^2*\sin(2*b*x + 2*a)^2 + 4*b^2*x^2*\cos(2*b*x + 2*a) + b^2*x^2 + 2*(2*b^2*x^2*\cos(2*b*x + 2*a) + b^2*x^2)*\cos(4*b*x + 4*a))$$

Giac [N/A]

Not integrable

Time = 0.92 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(bx + a)^3}{x} dx$$

[In] `integrate(tan(b*x+a)^3/x, x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)^3/x, x)`

Mupad [N/A]

Not integrable

Time = 3.18 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x} dx = \int \frac{\tan(a + bx)^3}{x} dx$$

[In] int(tan(a + b*x)^3/x, x)

[Out] int(tan(a + b*x)^3/x, x)

3.15 $\int \frac{\tan^3(a+bx)}{x^2} dx$

Optimal result	108
Rubi [N/A]	108
Mathematica [N/A]	109
Maple [N/A] (verified)	109
Fricas [N/A]	109
Sympy [N/A]	109
Maxima [N/A]	110
Giac [N/A]	110
Mupad [N/A]	111

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \text{Int}\left(\frac{\tan^3(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable(tan(b*x+a)^3/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

[In] Int[Tan[a + b*x]^3/x^2,x]

[Out] Defer[Int][Tan[a + b*x]^3/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tan^3(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 2.58 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

[In] `Integrate[Tan[a + b*x]^3/x^2, x]`

[Out] `Integrate[Tan[a + b*x]^3/x^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.19 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(bx + a)}{x^2} dx$$

[In] `int(tan(b*x+a)^3/x^2, x)`

[Out] `int(tan(b*x+a)^3/x^2, x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(bx + a)}{x^2} dx$$

[In] `integrate(tan(b*x+a)^3/x^2, x, algorithm="fricas")`

[Out] `integral(tan(b*x + a)^3/x^2, x)`

Sympy [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan^3(a + bx)}{x^2} dx$$

[In] `integrate(tan(b*x+a)**3/x**2, x)`

[Out] `Integral(tan(a + b*x)**3/x**2, x)`

Maxima [N/A]

Not integrable

Time = 0.71 (sec) , antiderivative size = 536, normalized size of antiderivative = 44.67

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

[In] `integrate(tan(b*x+a)^3/x^2,x, algorithm="maxima")`

[Out]
$$(4*b*x*\cos(2*b*x + 2*a)^2 + 4*b*x*\sin(2*b*x + 2*a)^2 + 2*b*x*\cos(2*b*x + 2*a) + 2*(b*x*\cos(2*b*x + 2*a) - \sin(2*b*x + 2*a))*\cos(4*b*x + 4*a) - (b^2*x^3*\cos(4*b*x + 4*a)^2 + 4*b^2*x^3*\cos(2*b*x + 2*a)^2 + b^2*x^3*\sin(4*b*x + 4*a)^2 + 4*b^2*x^3*\sin(4*b*x + 4*a)*\sin(2*b*x + 2*a) + 4*b^2*x^3*\sin(2*b*x + 2*a)^2 + 4*b^2*x^3*\cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*\cos(2*b*x + 2*a) + b^2*x^3)*\cos(4*b*x + 4*a))*\text{integrate}(2*(b^2*x^2 - 3)*\sin(2*b*x + 2*a)/(b^2*x^4*\cos(2*b*x + 2*a)^2 + b^2*x^4*\sin(2*b*x + 2*a)^2 + 2*b^2*x^4*\cos(2*b*x + 2*a) + b^2*x^4), x) + 2*(b*x*\sin(2*b*x + 2*a) + \cos(2*b*x + 2*a) + 1)*\sin(4*b*x + 4*a) + 2*\sin(2*b*x + 2*a))/(b^2*x^3*\cos(4*b*x + 4*a)^2 + 4*b^2*x^3*\cos(2*b*x + 2*a)^2 + b^2*x^3*\sin(4*b*x + 4*a)^2 + 4*b^2*x^3*\sin(4*b*x + 4*a)*\sin(2*b*x + 2*a) + 4*b^2*x^3*\sin(2*b*x + 2*a)^2 + 4*b^2*x^3*\cos(2*b*x + 2*a) + b^2*x^3 + 2*(2*b^2*x^3*\cos(2*b*x + 2*a) + b^2*x^3)*\cos(4*b*x + 4*a))$$

Giac [N/A]

Not integrable

Time = 0.96 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(bx + a)^3}{x^2} dx$$

[In] `integrate(tan(b*x+a)^3/x^2,x, algorithm="giac")`

[Out] `integrate(tan(b*x + a)^3/x^2, x)`

Mupad [N/A]

Not integrable

Time = 2.83 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\tan^3(a + bx)}{x^2} dx = \int \frac{\tan(a + bx)^3}{x^2} dx$$

[In] int(tan(a + b*x)^3/x^2,x)

[Out] int(tan(a + b*x)^3/x^2, x)

3.16 $\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx$

Optimal result	112
Rubi [A] (verified)	112
Mathematica [A] (verified)	119
Maple [F]	119
Fricas [A] (verification not implemented)	119
Sympy [F]	119
Maxima [F]	120
Giac [F]	120
Mupad [B] (verification not implemented)	120

Optimal result

Integrand size = 45, antiderivative size = 18

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

[Out] $-2*x^2/b/\tan(b*x+a)^{(1/2)}$

Rubi [A] (verified)

Time = 3.20 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 76, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3802, 3819, 209, 281, 6857, 6139, 6057, 2449, 2352, 2497, 6131, 6055}

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a+bx)}}$$

[In] $\text{Int}[x^2/\tan[a + b*x]^{(3/2)} - (4*x)/(b*\text{Sqrt}[\tan[a + b*x]]) + x^2*\text{Sqrt}[\tan[a + b*x]], x]$

[Out] $(-2*x^2)/(b*\text{Sqrt}[\tan[a + b*x]])$

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a,
, 0] || GtQ[b, 0])
```

Rule 281

```
Int[(x_)^(m_)*(a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 2352

```
Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_)/((d_) + (e_)*(x_))] / ((f_) + (g_)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] :> With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]
```

Rule 3802

```
Int[((c_) + (d_)*(x_))^(m_)*(b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(c + d*x)^m*((b*Tan[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + (-Dist[d*(m/(b*f*(n + 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n + 1), x], x] - Dist[1/b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n + 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && LtQ[n, -1] && GtQ[m, 0]
```

Rule 3819

```
Int[((c_) + (d_)*(x_))/Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)]], x_Symbol] :> Simp[(-I)*((c + d*x)/(f*Rt[a - I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x] + (Dist[I*(d/(f*Rt[a - I*b, 2])), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*(d/(f*Rt[a + I*b, 2])), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*((c + d*x)/(f*Rt[a + I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 6055

```
Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_)/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2,
```

0]

Rule 6057

```
Int[((a_.) + ArcTanh[(c_)*(x_)]*(b_.))/((d_) + (e_)*(x_)), x_Symbol] :> S
imp[(-(a + b*ArcTanh[c*x]))*(Log[2/(1 + c*x)]/e), x] + Dist[b*(c/e), Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x)/(c*d + e)*(1 + c*x))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x)/(c*d + e)*(1 + c*x))]/e), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Rule 6131

```
Int[((((a_.) + ArcTanh[(c_)*(x_)]*(b_.))^p_)*(x_))/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/(c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Rule 6139

```
Int[((((a_.) + ArcTanh[(c_)*(x_)]*(b_.))*(x_)^m_))/((d_) + (e_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0])
```

Rule 6857

```
Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{4 \int \frac{x}{\sqrt{\tan(a+bx)}} dx}{b} + \int \frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} dx + \int x^2 \sqrt{\tan(a+bx)} dx \\ &= \frac{4(-1)^{3/4} x \operatorname{arctanh}\left(\sqrt[4]{-1} \sqrt{\tan(a+bx)}\right)}{b^2} + \frac{4 \sqrt[4]{-1} x \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(a+bx)}\right)}{b^2} \\ &\quad - \frac{2x^2}{b \sqrt{\tan(a+bx)}} - \frac{(4 \sqrt[4]{-1}) \int \operatorname{arctanh}\left((-1)^{3/4} \sqrt{\tan(a+bx)}\right) dx}{b^2} \\ &\quad - \frac{(4(-1)^{3/4}) \int \operatorname{arctanh}\left(\sqrt[4]{-1} \sqrt{\tan(a+bx)}\right) dx}{b^2} + \frac{4 \int \frac{x}{\sqrt{\tan(a+bx)}} dx}{b} \end{aligned}$$

$$\begin{aligned}
&= -\frac{2x^2}{b\sqrt{\tan(a+bx)}} - \frac{(4\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}((-1)^{3/4}\sqrt{x})}{1+x^2} dx, x, \tan(a+bx)\right)}{b^3} \\
&\quad - \frac{(4(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}(\sqrt[4]{-1}\sqrt{x})}{1+x^2} dx, x, \tan(a+bx)\right)}{b^3} \\
&\quad + \frac{(4\sqrt[4]{-1}) \int \operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(a+bx)}\right) dx}{b^2} \\
&\quad + \frac{(4(-1)^{3/4}) \int \operatorname{arctanh}\left(\sqrt[4]{-1}\sqrt{\tan(a+bx)}\right) dx}{b^2} \\
&= -\frac{2x^2}{b\sqrt{\tan(a+bx)}} + \frac{(4\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}((-1)^{3/4}\sqrt{x})}{1+x^2} dx, x, \tan(a+bx)\right)}{b^3} \\
&\quad - \frac{(8\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}((-1)^{3/4}x)}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}(\sqrt[4]{-1}\sqrt{x})}{1+x^2} dx, x, \tan(a+bx)\right)}{b^3} \\
&\quad - \frac{(8(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}(\sqrt[4]{-1}x)}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&= -\frac{2x^2}{b\sqrt{\tan(a+bx)}} + \frac{(8\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}((-1)^{3/4}x)}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(8\sqrt[4]{-1}) \operatorname{Subst}\left(\int \left(-\frac{i x \operatorname{arctanh}((-1)^{3/4}x)}{2(-i+x^2)} + \frac{i x \operatorname{arctanh}((-1)^{3/4}x)}{2(i+x^2)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(8(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}(\sqrt[4]{-1}x)}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(8(-1)^{3/4}) \operatorname{Subst}\left(\int \left(-\frac{i x \operatorname{arctanh}(\sqrt[4]{-1}x)}{2(-i+x^2)} + \frac{i x \operatorname{arctanh}(\sqrt[4]{-1}x)}{2(i+x^2)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2x^2}{b\sqrt{\tan(a+bx)}} - \frac{(4\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{-i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4\sqrt[4]{-1}) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(8\sqrt[4]{-1}) \operatorname{Subst}\left(\int \left(-\frac{ix \operatorname{arctanh}((-1)^{3/4}x)}{2(-i+x^2)} + \frac{ix \operatorname{arctanh}((-1)^{3/4}x)}{2(i+x^2)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}((-1)^{3/4}x)}{-i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4(-1)^{3/4}) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}((-1)^{3/4}x)}{i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(8(-1)^{3/4}) \operatorname{Subst}\left(\int \left(-\frac{ix \operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2(-i+x^2)} + \frac{ix \operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2(i+x^2)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2\sqrt[4]{-1}\operatorname{arctanh}\left(\sqrt[4]{-1}\sqrt{\tan(a+bx)}\right)^2}{b^3} + \frac{2(-1)^{3/4}\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(a+bx)}\right)^2}{b^3} \\
&\quad - \frac{2x^2}{b\sqrt{\tan(a+bx)}} - \frac{(4i)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{1-\sqrt[4]{-1}x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4i)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{1-(-1)^{3/4}x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4\sqrt[4]{-1})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{-i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4\sqrt[4]{-1})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4\sqrt[4]{-1})\operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2\left(\sqrt[4]{-1}-x\right)} + \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2\left(\sqrt[4]{-1}+x\right)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4(-1)^{3/4})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left((-1)^{3/4}x\right)}{-i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4(-1)^{3/4})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left((-1)^{3/4}x\right)}{i+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4(-1)^{3/4})\operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{2(-(-1)^{3/4}-x)} + \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{2(-(-1)^{3/4}+x)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{4\sqrt[4]{-1}\operatorname{arctanh}\left(\sqrt[4]{-1}\sqrt{\tan(a+bx)}\right)\log\left(\frac{2}{1-\sqrt[4]{-1}\sqrt{\tan(a+bx)}}\right)}{b^3} \\
&\quad - \frac{4(-1)^{3/4}\operatorname{arctanh}\left((-1)^{3/4}\sqrt{\tan(a+bx)}\right)\log\left(\frac{2}{1-(-1)^{3/4}\sqrt{\tan(a+bx)}}\right)}{b^3} \\
&\quad - \frac{2x^2}{b\sqrt{\tan(a+bx)}} + \frac{(4i)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{1-\sqrt[4]{-1}x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4i)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{1-(-1)^{3/4}x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4i)\operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-\sqrt[4]{-1}x}\right)}{1-ix^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(4i)\operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-(-1)^{3/4}x}\right)}{1+ix^2} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(2\sqrt[4]{-1})\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{\sqrt[4]{-1-x}} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(2\sqrt[4]{-1})\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{\sqrt[4]{-1+x}} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4\sqrt[4]{-1})\operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2(\sqrt[4]{-1}-x)} + \frac{\operatorname{arctanh}\left(\sqrt[4]{-1}x\right)}{2(\sqrt[4]{-1}+x)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(2(-1)^{3/4})\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{-(-1)^{3/4}-x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad - \frac{(2(-1)^{3/4})\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{-(-1)^{3/4}+x} dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&\quad + \frac{(4(-1)^{3/4})\operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{2(-(-1)^{3/4}-x)} + \frac{\operatorname{arctanh}\left((-1)^{3/4}x\right)}{2(-(-1)^{3/4}+x)}\right) dx, x, \sqrt{\tan(a+bx)}\right)}{b^3} \\
&= \text{Too large to display}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.81 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(a + bx)}}$$

[In] `Integrate[x^2/Tan[a + b*x]^(3/2) - (4*x)/(b*.Sqrt[Tan[a + b*x]]) + x^2*Sqrt[Tan[a + b*x]], x]`

[Out] `(-2*x^2)/(b*.Sqrt[Tan[a + b*x]])`

Maple [F]

$$\int \left(-\frac{4x}{b\sqrt{\tan(bx + a)}} + x^2\left(\sqrt{\tan}(bx + a)\right) + \frac{x^2}{\tan(bx + a)^{\frac{3}{2}}} \right) dx$$

[In] `int(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2), x)`

[Out] `int(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2), x)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.89

$$\int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx = -\frac{2x^2}{b\sqrt{\tan(bx + a)}}$$

[In] `integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2), x, algorithm="fricas")`

[Out] `-2*x^2/(b*sqrt(tan(b*x + a)))`

Sympy [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a + bx)} - \frac{4x}{b\sqrt{\tan(a + bx)}} + x^2\sqrt{\tan(a + bx)} \right) dx \\ &= \frac{\int \left(-\frac{4x}{\sqrt{\tan(a+bx)}} \right) dx + \int \frac{bx^2}{\tan^{\frac{3}{2}}(a+bx)} dx + \int bx^2\sqrt{\tan(a+bx)} dx}{b} \end{aligned}$$

[In] `integrate(-4*x/b/tan(b*x+a)**(1/2)+x**2*tan(b*x+a)**(1/2)+x**2/tan(b*x+a)**(3/2), x)`

[Out] `(Integral(-4*x/sqrt(tan(a + b*x)), x) + Integral(b*x**2/tan(a + b*x)**(3/2), x) + Integral(b*x**2*sqrt(tan(a + b*x)), x))/b`

Maxima [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx \\ &= \int x^2\sqrt{\tan(bx+a)} + \frac{x^2}{\tan(bx+a)^{\frac{3}{2}}} - \frac{4x}{b\sqrt{\tan(bx+a)}} dx \end{aligned}$$

[In] integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2), x, algorithm="maxima")

[Out] integrate(x^2*sqrt(tan(b*x + a)) + x^2/tan(b*x + a)^(3/2) - 4*x/(b*sqrt(tan(b*x + a))), x)

Giac [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx \\ &= \int x^2\sqrt{\tan(bx+a)} + \frac{x^2}{\tan(bx+a)^{\frac{3}{2}}} - \frac{4x}{b\sqrt{\tan(bx+a)}} dx \end{aligned}$$

[In] integrate(-4*x/b/tan(b*x+a)^(1/2)+x^2*tan(b*x+a)^(1/2)+x^2/tan(b*x+a)^(3/2), x, algorithm="giac")

[Out] integrate(x^2*sqrt(tan(b*x + a)) + x^2/tan(b*x + a)^(3/2) - 4*x/(b*sqrt(tan(b*x + a))), x)

Mupad [B] (verification not implemented)

Time = 3.61 (sec) , antiderivative size = 50, normalized size of antiderivative = 2.78

$$\begin{aligned} & \int \left(\frac{x^2}{\tan^{\frac{3}{2}}(a+bx)} - \frac{4x}{b\sqrt{\tan(a+bx)}} + x^2\sqrt{\tan(a+bx)} \right) dx \\ &= -\frac{x^2 \sin(2a + 2bx) \sqrt{\frac{\sin(2a + 2bx)}{\cos(2a + 2bx) + 1}}}{b \sin(a + bx)^2} \end{aligned}$$

[In] int(x^2*tan(a + b*x)^(1/2) + x^2/tan(a + b*x)^(3/2) - (4*x)/(b*tan(a + b*x)^(1/2)), x)

[Out] -(x^2*sin(2*a + 2*b*x)*(sin(2*a + 2*b*x)/(cos(2*a + 2*b*x) + 1))^(1/2))/(b*sin(a + b*x)^2)

$$3.17 \quad \int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx$$

Optimal result	121
Rubi [F]	121
Mathematica [A] (verified)	122
Maple [F]	122
Fricas [A] (verification not implemented)	122
Sympy [F]	122
Maxima [F]	123
Giac [F]	123
Mupad [B] (verification not implemented)	123

Optimal result

Integrand size = 49, antiderivative size = 17

$$\int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx = \frac{x\sqrt{\tan(a+bx^2)}}{b}$$

[Out] $x*\tan(b*x^2+a)^{(1/2)}/b$

Rubi [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx \\ &= \int \left(\frac{x^2}{\sqrt{\tan(a+bx^2)}} + \frac{\sqrt{\tan(a+bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a+bx^2) \right) dx \end{aligned}$$

[In] $\text{Int}[x^2/\text{Sqrt}[\text{Tan}[a+b*x^2]] + \text{Sqrt}[\text{Tan}[a+b*x^2]]/b + x^2*\text{Tan}[a+b*x^2]^{(3/2)}, x]$

[Out] $\text{Defer}[\text{Int}][x^2/\text{Sqrt}[\text{Tan}[a+b*x^2]], x] + \text{Defer}[\text{Int}][\text{Sqrt}[\text{Tan}[a+b*x^2]], x]/b + \text{Defer}[\text{Int}][x^2*\text{Tan}[a+b*x^2]^{(3/2)}, x]$

Rubi steps

$$\text{integral} = \frac{\int \sqrt{\tan(a+bx^2)} dx}{b} + \int \frac{x^2}{\sqrt{\tan(a+bx^2)}} dx + \int x^2 \tan^{\frac{3}{2}}(a+bx^2) dx$$

Mathematica [A] (verified)

Time = 0.92 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{x \sqrt{\tan(a + bx^2)}}{b}$$

[In] Integrate[x^2/Sqrt[Tan[a + b*x^2]] + Sqrt[Tan[a + b*x^2]]/b + x^2*Tan[a + b*x^2]^(3/2), x]

[Out] (x*Sqrt[Tan[a + b*x^2]])/b

Maple [F]

$$\int \left(\frac{x^2}{\sqrt{\tan(x^2b + a)}} + \frac{\sqrt{\tan}(x^2b + a)}{b} + x^2 \left(\tan^{\frac{3}{2}}(x^2b + a) \right) \right) dx$$

[In] int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2), x)

[Out] int(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2), x)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.88

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{x \sqrt{\tan(bx^2 + a)}}{b}$$

[In] integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2), x, algorithm="fricas")

[Out] x*sqrt(tan(b*x^2 + a))/b

Sympy [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx \\ &= \frac{\int \frac{bx^2}{\sqrt{\tan(a+bx^2)}} dx + \int bx^2 \tan^{\frac{3}{2}}(a + bx^2) dx + \int \sqrt{\tan(a + bx^2)} dx}{b} \end{aligned}$$

[In] integrate(x**2/tan(b*x**2+a)**(1/2)+tan(b*x**2+a)**(1/2)/b+x**2*tan(b*x**2+a)**(3/2), x)

[Out] (Integral(b*x**2/sqrt(tan(a + b*x**2)), x) + Integral(b*x**2*tan(a + b*x**2)**(3/2), x) + Integral(sqrt(tan(a + b*x**2)), x))/b

Maxima [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx \\ &= \int x^2 \tan(bx^2 + a)^{\frac{3}{2}} + \frac{x^2}{\sqrt{\tan(bx^2 + a)}} + \frac{\sqrt{\tan(bx^2 + a)}}{b} dx \end{aligned}$$

[In] `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="maxima")`

[Out] `integrate(x^2*tan(b*x^2 + a)^(3/2) + x^2/sqrt(tan(b*x^2 + a)) + sqrt(tan(b*x^2 + a))/b, x)`

Giac [F]

$$\begin{aligned} & \int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx \\ &= \int x^2 \tan(bx^2 + a)^{\frac{3}{2}} + \frac{x^2}{\sqrt{\tan(bx^2 + a)}} + \frac{\sqrt{\tan(bx^2 + a)}}{b} dx \end{aligned}$$

[In] `integrate(x^2/tan(b*x^2+a)^(1/2)+tan(b*x^2+a)^(1/2)/b+x^2*tan(b*x^2+a)^(3/2),x, algorithm="giac")`

[Out] `integrate(x^2*tan(b*x^2 + a)^(3/2) + x^2/sqrt(tan(b*x^2 + a)) + sqrt(tan(b*x^2 + a))/b, x)`

Mupad [B] (verification not implemented)

Time = 5.45 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

$$\int \left(\frac{x^2}{\sqrt{\tan(a + bx^2)}} + \frac{\sqrt{\tan(a + bx^2)}}{b} + x^2 \tan^{\frac{3}{2}}(a + bx^2) \right) dx = \frac{x \sqrt{-\frac{e^{2i b x^2 + a 2i} 1i - i}{e^{2i b x^2 + a 2i} + 1}}}{b}$$

[In] `int(tan(a + b*x^2)^(1/2)/b + x^2/tan(a + b*x^2)^(1/2) + x^2*tan(a + b*x^2)^(3/2),x)`

[Out] `(x*(-(exp(a*2i + b*x^2*2i)*1i - 1i)/(exp(a*2i + b*x^2*2i) + 1))^(1/2))/b`

3.18 $\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx$

Optimal result	124
Rubi [A] (verified)	124
Mathematica [A] (verified)	126
Maple [A] (verified)	126
Fricas [A] (verification not implemented)	127
Sympy [A] (verification not implemented)	127
Maxima [F(-2)]	128
Giac [A] (verification not implemented)	128
Mupad [B] (verification not implemented)	128

Optimal result

Integrand size = 23, antiderivative size = 189

$$\begin{aligned} \int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx = & \frac{3id^3x}{8af^3} - \frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} \\ & - \frac{3d^3}{8f^4(a+ia\tan(e+fx))} - \frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} \\ & + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} \end{aligned}$$

[Out] $3/8*I*d^3*x/a/f^3-3/8*d*(d*x+c)^2/a/f^2-1/4*I*(d*x+c)^3/a/f+1/8*(d*x+c)^4/a/d-3/8*d^3/f^4/(a+I*a*tan(f*x+e))-3/4*I*d^2*(d*x+c)/f^3/(a+I*a*tan(f*x+e))+3/4*d*(d*x+c)^2/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^3/f/(a+I*a*tan(f*x+e))$

Rubi [A] (verified)

Time = 0.22 (sec), antiderivative size = 189, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3804, 3560, 8}

$$\begin{aligned} \int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx = & -\frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} \\ & + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} - \frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} \\ & + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+ia\tan(e+fx))} + \frac{3id^3x}{8af^3} \end{aligned}$$

[In] Int[(c + d*x)^3/(a + I*a*Tan[e + f*x]), x]

```
[Out] (((3*I)/8)*d^3*x)/(a*f^3) - (3*d*(c + d*x)^2)/(8*a*f^2) - ((I/4)*(c + d*x)^3)/(a*f) + (c + d*x)^4/(8*a*d) - (3*d^3)/(8*f^4*(a + I*a*Tan[e + f*x])) - ((3*I)/4)*d^2*(c + d*x))/(f^3*(a + I*a*Tan[e + f*x])) + (3*d*(c + d*x)^2)/(4*f^2*(a + I*a*Tan[e + f*x])) + ((I/2)*(c + d*x)^3)/(f*(a + I*a*Tan[e + f*x]))
```

Rule 8

```
Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]
```

Rule 3560

```
Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n)/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^4}{8ad} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} - \frac{(3id)\int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx}{2f} \\
&= -\frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} \\
&\quad + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} - \frac{(3d^2)\int \frac{c+dx}{a+ia\tan(e+fx)} dx}{2f^2} \\
&= -\frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} \\
&\quad + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} + \frac{(3id^3)\int \frac{1}{a+ia\tan(e+fx)} dx}{4f^3} \\
&= -\frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+ia\tan(e+fx))} \\
&\quad - \frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} \\
&\quad + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))} + \frac{(3id^3)\int 1 dx}{8af^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3id^3x}{8af^3} - \frac{3d(c+dx)^2}{8af^2} - \frac{i(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+ia\tan(e+fx))} \\
&\quad - \frac{3id^2(c+dx)}{4f^3(a+ia\tan(e+fx))} + \frac{3d(c+dx)^2}{4f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^3}{2f(a+ia\tan(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.47

$$\begin{aligned}
&\int \frac{(c+dx)^3}{a+ia\tan(e+fx)} dx \\
&= \frac{\sec(e+fx)(\cos(fx)+i\sin(fx))((4ic^3f^3+6c^2df^2(1+2ifx)+6cd^2f(-i+2fx+2if^2x^2)+d^3(-3-6ifx)))}{}
\end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x]), x]

[Out] $\left(\frac{d^3x^4}{8a} + \frac{d^2cx^3}{2a} + \frac{3dc^2x^2}{4a} + \frac{c^3x}{2a} + \frac{c^4}{8ad} + \frac{i(4d^3x^3f^3+12cd^2f^3x^2-6id^3f^2x^2+12c^2df^3x-12icd^2f^2x+4c^3f^3-6ic^2df^2-6d^3fx)}{16af^4}\right)$

Maple [A] (verified)

Time = 0.67 (sec) , antiderivative size = 170, normalized size of antiderivative = 0.90

method	result
risch	$\frac{d^3x^4}{8a} + \frac{d^2cx^3}{2a} + \frac{3dc^2x^2}{4a} + \frac{c^3x}{2a} + \frac{c^4}{8ad} + \frac{i(4d^3x^3f^3+12cd^2f^3x^2-6id^3f^2x^2+12c^2df^3x-12icd^2f^2x+4c^3f^3-6ic^2df^2-6d^3fx)}{16af^4}$
default	Expression too large to display

[In] int((d*x+c)^3/(a+I*a*tan(f*x+e)), x, method=_RETURNVERBOSE)

[Out] $\frac{1}{8}a^2d^3x^4 + \frac{1}{2}a^2d^2c^2x^3 + \frac{3}{4}a^2cd^2x^2 + \frac{c^3x}{2a} + \frac{c^4}{8ad} + \frac{i(4d^3x^3f^3+12cd^2f^3x^2-6id^3f^2x^2+12c^2df^3x-12icd^2f^2x+4c^3f^3-6ic^2df^2-6d^3fx)}{16af^4}$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.87

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ = \frac{(4i d^3 f^3 x^3 + 4i c^3 f^3 + 6 c^2 d f^2 - 6i c d^2 f - 3 d^3 - 6 (-2i c d^2 f^3 - d^3 f^2)x^2 - 6 (-2i c^2 d f^3 - 2 c d^2 f^2 + i d^3 f^4))}{16 a f^4}$$

```
[In] integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
[Out] 1/16*(4*I*d^3*f^3*x^3 + 4*I*c^3*f^3 + 6*c^2*d*f^2 - 6*I*c*d^2*f - 3*d^3 - 6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 - 6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3*f)*x + 2*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x)*e^(-2*I*f*x - 2*I*e)/(a*f^4)
```

Sympy [A] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.37

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ = \begin{cases} \frac{(4ic^3f^3 + 12ic^2df^3x + 6c^2df^2 + 12icd^2f^3x^2 + 12cd^2f^2x - 6icd^2f + 4id^3f^3x^3 + 6d^3f^2x^2 - 6id^3fx - 3d^3)e^{-2ie}e^{-2ifx}}{16af^4} & \text{for } af^4e^{2ie} \neq 0 \\ \frac{c^3xe^{-2ie}}{2a} + \frac{3c^2dx^2e^{-2ie}}{4a} + \frac{cd^2x^3e^{-2ie}}{2a} + \frac{d^3x^4e^{-2ie}}{8a} \\ + \frac{c^3x}{2a} + \frac{3c^2dx^2}{4a} + \frac{cd^2x^3}{2a} + \frac{d^3x^4}{8a} & \text{otherwise} \end{cases}$$

```
[In] integrate((d*x+c)**3/(a+I*a*tan(f*x+e)),x)
[Out] Piecewise(((4*I*c**3*f**3 + 12*I*c**2*d*f**3*x + 6*c**2*d*f**2 + 12*I*c*d**2*f**3*x**2 + 12*c*d**2*f**2*x - 6*I*c*d**2*f + 4*I*d**3*f**3*x**3 + 6*d**3*f**2*x**2 - 6*I*d**3*f*x - 3*d**3)*exp(-2*I*e)*exp(-2*I*f*x)/(16*a*f**4),
Ne(a*f**4*exp(2*I*e), 0)), (c**3*x*exp(-2*I*e)/(2*a) + 3*c**2*d*x**2*exp(-2*I*e)/(4*a) + c*d**2*x**3*exp(-2*I*e)/(2*a) + d**3*x**4*exp(-2*I*e)/(8*a),
True)) + c**3*x/(2*a) + 3*c**2*d*x**2/(4*a) + c*d**2*x**3/(2*a) + d**3*x**4/(8*a)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 188, normalized size of antiderivative = 0.99

$$\begin{aligned} & \int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ &= \frac{(2 d^3 f^4 x^4 e^{(2i f x + 2i e)} + 8 c d^2 f^4 x^3 e^{(2i f x + 2i e)} + 12 c^2 d f^4 x^2 e^{(2i f x + 2i e)} + 4 i d^3 f^3 x^3 + 8 c^3 f^4 x e^{(2i f x + 2i e)} + 12 i c d^2 f^3 x^2 e^{(2i f x + 2i e)})}{16 a f^4} \end{aligned}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

[Out] $\frac{1}{16} (2 d^3 f^4 x^4 e^{(2 I f x + 2 I e)} + 8 c d^2 f^4 x^3 e^{(2 I f x + 2 I e)} + 12 c^2 d f^4 x^2 e^{(2 I f x + 2 I e)} + 4 I d^3 f^3 x^3 + 8 c^3 f^4 x e^{(2 I f x + 2 I e)} + 12 I c d^2 f^3 x^2 + 12 I c^2 d f^4 x^3 + 6 d^3 f^2 x^2 - 6 I d^3 f x - 6 I c d^2 f - 3 d^3) e^{(-2 I f x - 2 I e)} / (a f^4)$

Mupad [B] (verification not implemented)

Time = 4.12 (sec) , antiderivative size = 423, normalized size of antiderivative = 2.24

$$\begin{aligned} & \int \frac{(c + dx)^3}{a + ia \tan(e + fx)} dx \\ &= \frac{8 c^3 f^4 x - 3 d^3 \cos(2 e + 2 f x) + 4 c^3 f^3 \sin(2 e + 2 f x) + 2 d^3 f^4 x^4 + 6 c^2 d f^2 \cos(2 e + 2 f x) + 12 c^2 d^2 f^3 x^3}{\dots} \end{aligned}$$

[In] `int((c + d*x)^3/(a + a*tan(e + f*x)*1i),x)`

[Out] $(d^3 \sin(2 e + 2 f x) * 3i - 3 d^3 \cos(2 e + 2 f x) + 8 c^3 f^4 x + c^3 f^3 \cos(2 e + 2 f x) * 4i + 4 c^3 f^3 \sin(2 e + 2 f x) + 2 d^3 f^4 x^4 + 6 c^2 d f^2 \cos(2 e + 2 f x) - c^2 d f^2 \sin(2 e + 2 f x) * 6i + 12 c^2 d f^4 x^2 + 8 c^2 d^2 f^3 x^3) / (a^2 f^4)$

$$\begin{aligned} & c*d^2*f^4*x^3 + 6*d^3*f^2*x^2*\cos(2*e + 2*f*x) + d^3*f^3*x^3*\cos(2*e + 2*f*x)*4i - d^3*f^2*x^2*\sin(2*e + 2*f*x)*6i + 4*d^3*f^3*x^3*\sin(2*e + 2*f*x) - \\ & c*d^2*f*\cos(2*e + 2*f*x)*6i - 6*c*d^2*f*\sin(2*e + 2*f*x) - d^3*f*x*\cos(2*e + 2*f*x)*6i - 6*d^3*f*x*\sin(2*e + 2*f*x) + 12*c*d^2*f^2*x*\cos(2*e + 2*f*x) \\ & + c^2*d*f^3*x*\cos(2*e + 2*f*x)*12i - c*d^2*f^2*x*\sin(2*e + 2*f*x)*12i + 12*c^2*d*f^3*x*\sin(2*e + 2*f*x) + c*d^2*f^3*x^2*\cos(2*e + 2*f*x)*12i + 12*c*d^2*f^3*x^2*\sin(2*e + 2*f*x)) / (16*a*f^4) \end{aligned}$$

3.19 $\int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx$

Optimal result	130
Rubi [A] (verified)	130
Mathematica [A] (verified)	132
Maple [A] (verified)	132
Fricas [A] (verification not implemented)	132
Sympy [A] (verification not implemented)	133
Maxima [F(-2)]	133
Giac [A] (verification not implemented)	133
Mupad [B] (verification not implemented)	134

Optimal result

Integrand size = 23, antiderivative size = 137

$$\begin{aligned} \int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx = & -\frac{d^2x}{4af^2} - \frac{i(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{id^2}{4f^3(a+ia\tan(e+fx))} \\ & + \frac{d(c+dx)}{2f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))} \end{aligned}$$

[Out] $-1/4*d^2*x/a/f^2-1/4*I*(d*x+c)^2/a/f+1/6*(d*x+c)^3/a/d-1/4*I*d^2/f^3/(a+I*a*tan(f*x+e))+1/2*d*(d*x+c)/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)^2/f/(a+I*a*tan(f*x+e))$

Rubi [A] (verified)

Time = 0.14 (sec), antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3804, 3560, 8}

$$\begin{aligned} \int \frac{(c+dx)^2}{a+ia\tan(e+fx)} dx = & \frac{d(c+dx)}{2f^2(a+ia\tan(e+fx))} + \frac{i(c+dx)^2}{2f(a+ia\tan(e+fx))} \\ & - \frac{i(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{id^2}{4f^3(a+ia\tan(e+fx))} - \frac{d^2x}{4af^2} \end{aligned}$$

[In] $\text{Int}[(c+d*x)^2/(a+I*a*Tan[e+f*x]),x]$

[Out] $-1/4*(d^2*x)/(a*f^2) - ((I/4)*(c+d*x)^2)/(a*f) + (c+d*x)^3/(6*a*d) - ((I/4)*d^2)/(f^3*(a+I*a*Tan[e+f*x])) + (d*(c+d*x))/(2*f^2*(a+I*a*Tan[e+f*x])) + ((I/2)*(c+d*x)^2)/(f*(a+I*a*Tan[e+f*x]))$

Rule 8

```
Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]
```

Rule 3560

```
Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_.) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(c + dx)^3}{6ad} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} - \frac{(id) \int \frac{c + dx}{a + ia \tan(e + fx)} dx}{f} \\
 &= -\frac{i(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} + \frac{d(c + dx)}{2f^2(a + ia \tan(e + fx))} \\
 &\quad + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} - \frac{d^2 \int \frac{1}{a + ia \tan(e + fx)} dx}{2f^2} \\
 &= -\frac{i(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} - \frac{id^2}{4f^3(a + ia \tan(e + fx))} \\
 &\quad + \frac{d(c + dx)}{2f^2(a + ia \tan(e + fx))} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))} - \frac{d^2 \int 1 dx}{4af^2} \\
 &= -\frac{d^2 x}{4af^2} - \frac{i(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} - \frac{id^2}{4f^3(a + ia \tan(e + fx))} \\
 &\quad + \frac{d(c + dx)}{2f^2(a + ia \tan(e + fx))} + \frac{i(c + dx)^2}{2f(a + ia \tan(e + fx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.49 (sec) , antiderivative size = 178, normalized size of antiderivative = 1.30

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx \\ = \sec(e + fx)(\cos(fx) + i \sin(fx)) ((d + (1 + i)cf + (1 + i)dfx)((1 + i)cf + d(-i + (1 + i)fx)) \cos(2fx)(co$$

[In] Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x]), x]

[Out] $(\sec(e + fx)(\cos(fx) + i \sin(fx))((d + (1 + i)cf + (1 + i)dfx)((1 + i)cf + d(-i + (1 + i)fx)) \cos(2fx)(co$
 $x^*(3c^2 + 3c*d*x + d^2*x^2)*(Cos[e] + I*Sin[e])))/3 - I*(d + (1 + i)cf + (1 + i)dfx)*((1 + i)cf + d*(-i + (1 + i)fx)) * (\cos(e) - I*\sin(e))*Si$
 $n[2*f*x])/ (8*f^3*(a + I*a*Tan[e + f*x]))$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 108, normalized size of antiderivative = 0.79

method	result
risch	$\frac{d^2x^3}{6a} + \frac{dx^2}{2a} + \frac{c^2x}{2a} + \frac{c^3}{6ad} + \frac{i(2d^2x^2f^2 + 4cd f^2x - 2id^2fx + 2c^2f^2 - 2icdf - d^2)e^{-2i(fx+e)}}{8af^3}$
default	$\frac{ic^2(\cos^2(fx+e))}{2} - \frac{icde(\cos^2(fx+e))}{f} - \frac{2icd\left(-\frac{(fx+e)(\cos^2(fx+e))}{2} + \frac{\sin(fx+e)\cos(fx+e)}{4} + \frac{fx}{4} + \frac{e}{4}\right)}{f} + \frac{id^2e^2(\cos^2(fx+e))}{2f^2} + \frac{2id^2e\left(-\frac{(fx+e)(\cos^2(fx+e))}{2} + \frac{\sin(fx+e)\cos(fx+e)}{4} + \frac{fx}{4} + \frac{e}{4}\right)}{f}$

[In] int((d*x+c)^2/(a+I*a*tan(f*x+e)), x, method=_RETURNVERBOSE)

[Out] $1/6/a*d^2*x^3 + 1/2/a*d*c*x^2 + 1/2/a*c^2*x + 1/6/a/d*c^3 + 1/8*I*(2*d^2*x^2*f^2 - 2*I*d^2*f*x + 4*c*d*f^2*x - 2*I*c*d*f + 2*c^2*f^2 - d^2)/a/f^3*\exp(-2*I*(f*x+e))$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.77

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx \\ = \frac{(6i d^2 f^2 x^2 + 6i c^2 f^2 + 6 cdf - 3i d^2 - 6 (-2i cdf^2 - d^2 f)x + 4 (d^2 f^3 x^3 + 3 cdf^3 x^2 + 3 c^2 f^3 x)e^{(2i f x + 2i e)})e^{(-2i f x - 2i e)}}{24 a f^3}$$

[In] integrate((d*x+c)^2/(a+I*a*tan(f*x+e)), x, algorithm="fricas")

[Out] $\frac{1}{24} \left(6I^2 d^2 f^2 x^2 + 6I^2 c^2 f^2 x^2 + 6c^2 d^2 f - 3I^2 d^2 - 6(-2I^2 c^2 d^2 f^2 - d^2 f^2)x + 4(d^2 f^2 x^3 + 3c^2 d^2 f^2 x^2 + 3c^2 f^2 x^3)e^{(2Ifx + 2Ie)} - e^{(-2Ifx - 2Ie)}(a^2 f^3) \right)$

Sympy [A] (verification not implemented)

Time = 0.19 (sec), antiderivative size = 165, normalized size of antiderivative = 1.20

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \begin{cases} \frac{(2ic^2 f^2 + 4icdf^2 x + 2cdf + 2id^2 f^2 x^2 + 2d^2 f x - id^2)e^{-2ie} e^{-2ifx}}{8af^3} & \text{for } af^3 e^{2ie} \neq 0 \\ \frac{c^2 x e^{-2ie}}{2a} + \frac{cdx^2 e^{-2ie}}{2a} + \frac{d^2 x^3 e^{-2ie}}{6a} & \text{otherwise} \\ + \frac{c^2 x}{2a} + \frac{cdx^2}{2a} + \frac{d^2 x^3}{6a} \end{cases}$$

[In] `integrate((d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

[Out] `Piecewise(((2*I*c**2*f**2 + 4*I*c*d*f**2*x + 2*c*d*f + 2*I*d**2*f**2*x**2 + 2*d**2*f*x - I*d**2)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**3), Ne(a*f**3*exp(2*I*e), 0)), (c**2*x*exp(-2*I*e)/(2*a) + c*d*x**2*exp(-2*I*e)/(2*a) + d**2*x**3*exp(-2*I*e)/(6*a), True)) + c**2*x/(2*a) + c*d*x**2/(2*a) + d**2*x**3/(6*a)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

[Out] `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

none

Time = 0.36 (sec), antiderivative size = 119, normalized size of antiderivative = 0.87

$$\int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx = \frac{(4d^2 f^3 x^3 e^{(2ifx + 2ie)} + 12cdf^3 x^2 e^{(2ifx + 2ie)} + 12c^2 f^3 x e^{(2ifx + 2ie)} + 6i d^2 f^2 x^2 + 12i cdf^2 x + 6i c^2 f^2 + 6d^2 f^2)}{24af^3}$$

[In] `integrate((d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

[Out] $\frac{1}{24} \cdot (4 \cdot d^2 \cdot f^3 \cdot x^3 \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 12 \cdot c \cdot d \cdot f^3 \cdot x^2 \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 12 \cdot c^2 \cdot f^3 \cdot x^3 \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + 6 \cdot I \cdot d^2 \cdot f^2 \cdot x^2 + 12 \cdot I \cdot c \cdot d \cdot f^2 \cdot x + 6 \cdot I \cdot c^2 \cdot f^2 + 6 \cdot d^2 \cdot f \cdot x + 6 \cdot c \cdot d \cdot f - 3 \cdot I \cdot d^2) \cdot e^{(-2 \cdot I \cdot f \cdot x - 2 \cdot I \cdot e)} / (a \cdot f^3)$

Mupad [B] (verification not implemented)

Time = 3.71 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.76

$$\begin{aligned} & \int \frac{(c + dx)^2}{a + ia \tan(e + fx)} dx \\ &= \frac{12 c^2 f^3 x - 3 d^2 \sin(2e + 2fx) + 6 c^2 f^2 \sin(2e + 2fx) + 4 d^2 f^3 x^3 + 6 cd f \cos(2e + 2fx) + 6 d^2 f^2 x^2}{\dots} \end{aligned}$$

[In] `int((c + d*x)^2/(a + a*tan(e + f*x)*1i),x)`

[Out] $\frac{(12 \cdot c^2 \cdot f^3 \cdot x - 3 \cdot d^2 \cdot \sin(2 \cdot e + 2 \cdot f \cdot x) - d^2 \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) \cdot 3i + c^2 \cdot f^2 \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) \cdot 6i + 6 \cdot c^2 \cdot f^2 \cdot \sin(2 \cdot e + 2 \cdot f \cdot x) + 4 \cdot d^2 \cdot f^3 \cdot x^3 + 6 \cdot c \cdot d \cdot f \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) - c \cdot d \cdot f \cdot \sin(2 \cdot e + 2 \cdot f \cdot x) \cdot 6i + d^2 \cdot f^2 \cdot x^2 \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) \cdot 6i + 6 \cdot d^2 \cdot f^2 \cdot x^2 \cdot \sin(2 \cdot e + 2 \cdot f \cdot x) + 12 \cdot c \cdot d \cdot f^3 \cdot x^2 + 6 \cdot d^2 \cdot f \cdot x \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) - d^2 \cdot f \cdot x \cdot \sin(2 \cdot e + 2 \cdot f \cdot x) \cdot 6i + c \cdot d \cdot f^2 \cdot x \cdot \cos(2 \cdot e + 2 \cdot f \cdot x) \cdot 12i + 12 \cdot c \cdot d \cdot f^2 \cdot x \cdot \sin(2 \cdot e + 2 \cdot f \cdot x))}{(24 \cdot a \cdot f^3)}$

3.20 $\int \frac{c+dx}{a+ia\tan(e+fx)} dx$

Optimal result	135
Rubi [A] (verified)	135
Mathematica [A] (verified)	136
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	137
Sympy [A] (verification not implemented)	137
Maxima [F(-2)]	138
Giac [A] (verification not implemented)	138
Mupad [B] (verification not implemented)	138

Optimal result

Integrand size = 21, antiderivative size = 84

$$\begin{aligned} \int \frac{c+dx}{a+ia\tan(e+fx)} dx = & -\frac{idx}{4af} + \frac{(c+dx)^2}{4ad} + \frac{d}{4f^2(a+ia\tan(e+fx))} \\ & + \frac{i(c+dx)}{2f(a+ia\tan(e+fx))} \end{aligned}$$

[Out] $-1/4*I*d*x/a/f+1/4*(d*x+c)^2/a/d+1/4*d/f^2/(a+I*a*tan(f*x+e))+1/2*I*(d*x+c)/f/(a+I*a*tan(f*x+e))$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3804, 3560, 8}

$$\begin{aligned} \int \frac{c+dx}{a+ia\tan(e+fx)} dx = & \frac{i(c+dx)}{2f(a+ia\tan(e+fx))} + \frac{(c+dx)^2}{4ad} \\ & + \frac{d}{4f^2(a+ia\tan(e+fx))} - \frac{idx}{4af} \end{aligned}$$

[In] $\text{Int}[(c+d*x)/(a+I*a*Tan[e+f*x]),x]$

[Out] $((-1/4*I)*d*x)/(a*f) + (c+d*x)^2/(4*a*d) + d/(4*f^2*(a+I*a*Tan[e+f*x])) + ((I/2)*(c+d*x))/(f*(a+I*a*Tan[e+f*x]))$

Rule 8

$\text{Int}[a_, x_Symbol] :> \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3560

```
Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a +
b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_.) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)),
Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(c + dx)^2}{4ad} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} - \frac{(id) \int \frac{1}{a + ia \tan(e + fx)} dx}{2f} \\ &= \frac{(c + dx)^2}{4ad} + \frac{d}{4f^2(a + ia \tan(e + fx))} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} - \frac{(id) \int 1 dx}{4af} \\ &= -\frac{idx}{4af} + \frac{(c + dx)^2}{4ad} + \frac{d}{4f^2(a + ia \tan(e + fx))} + \frac{i(c + dx)}{2f(a + ia \tan(e + fx))} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.74 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.14

$$\begin{aligned} &\int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ &= \frac{-i(2cf(i + 2fx) + d(1 + 2ifx + 2f^2x^2)) + (2cf(-i + 2fx) + d(-1 - 2ifx + 2f^2x^2)) \tan(e + fx)}{8af^2(-i + \tan(e + fx))} \end{aligned}$$

[In] `Integrate[(c + d*x)/(a + I*a*Tan[e + f*x]), x]`

[Out] `((-I)*(2*c*f*(I + 2*f*x) + d*(1 + (2*I)*f*x + 2*f^2*x^2)) + (2*c*f*(-I + 2*f*x) + d*(-1 - (2*I)*f*x + 2*f^2*x^2))*Tan[e + f*x])/ (8*a*f^2*(-I + Tan[e + f*x]))`

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 50, normalized size of antiderivative = 0.60

method	result
risch	$\frac{dx^2}{4a} + \frac{cx}{2a} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8af^2}$
norman	$\frac{dx^2}{4a} + \frac{2icf+d}{4af^2} + \frac{dx^2(\tan^2(fx+e))}{4a} + \frac{(2cf-id)\tan(fx+e)}{4af^2} + \frac{(2cf+id)x}{4af} + \frac{dx\tan(fx+e)}{2af} + \frac{(2cf-id)x(\tan^2(fx+e))}{4fa}$ $1+\tan^2(fx+e)$
default	$\frac{ic(\cos^2(fx+e))}{2} - \frac{ide(\cos^2(fx+e))}{2f} - \frac{id\left(-\frac{(fx+e)(\cos^2(fx+e))}{2} + \frac{\sin(fx+e)\cos(fx+e)}{4} + \frac{fx}{4} + \frac{e}{4}\right)}{f} + c\left(\frac{\sin(fx+e)\cos(fx+e)}{2} + \frac{fx}{2} + \frac{e}{2}\right) - \frac{de(\sin(fx+e)\cos(fx+e))}{af}$

[In] `int((d*x+c)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4}/a*d*x^2+1/2/a*c*x+1/8*I*(2*d*f*x-I*d+2*c*f)/a/f^2*\exp(-2*I*(f*x+e))$

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 55, normalized size of antiderivative = 0.65

$$\int \frac{c+dx}{a+ia\tan(e+fx)} dx = \frac{(2i dfx + 2i cf + 2(df^2 x^2 + 2cf^2 x)e^{(2ifx+2ie)} + d)e^{(-2ifx-2ie)}}{8af^2}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

[Out] $\frac{1}{8}*(2*I*d*f*x + 2*I*c*f + 2*(d*f^2*x^2 + 2*c*f^2*x)*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-2*I*f*x - 2*I*e)}/(a*f^2)$

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.10

$$\int \frac{c+dx}{a+ia\tan(e+fx)} dx = \begin{cases} \frac{(2icf+2idfx+d)e^{-2ie}e^{-2ifx}}{8af^2} & \text{for } af^2 e^{2ie} \neq 0 \\ \frac{cxe^{-2ie}}{2a} + \frac{dx^2 e^{-2ie}}{4a} & \text{otherwise} \end{cases} + \frac{cx}{2a} + \frac{dx^2}{4a}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x)`

[Out] `Piecewise(((2*I*c*f + 2*I*d*f*x + d)*exp(-2*I*e)*exp(-2*I*f*x)/(8*a*f**2), Ne(a*f**2*exp(2*I*e), 0)), (c*x*exp(-2*I*e)/(2*a) + d*x**2*exp(-2*I*e)/(4*a), True)) + c*x/(2*a) + d*x**2/(4*a)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{a + ia \tan(e + fx)} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.74

$$\begin{aligned} & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ &= \frac{(2 df^2 x^2 e^{(2i f x + 2i e)} + 4 c f^2 x e^{(2i f x + 2i e)} + 2i d f x + 2i c f + d) e^{(-2i f x - 2i e)}}{8 a f^2} \end{aligned}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

[Out] $\frac{1}{8} (2 d f^2 x^2 e^{(2 I f x + 2 I e)} + 4 c f^2 x e^{(2 I f x + 2 I e)} + 2 I d f x + 2 I c f + d) e^{(-2 I f x - 2 I e)} / (a f^2)$

Mupad [B] (verification not implemented)

Time = 3.45 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.25

$$\begin{aligned} & \int \frac{c + dx}{a + ia \tan(e + fx)} dx \\ &= \frac{d \cos(2e + 2fx) + 2df^2x^2 + 2cf \sin(2e + 2fx) + 4cf^2x + 2dfx \sin(2e + 2fx) - d \sin(2e + 2fx)}{8af^2} \end{aligned}$$

[In] `int((c + d*x)/(a + a*tan(e + f*x)*1i),x)`

[Out] $(d \cos(2e + 2fx) - d \sin(2e + 2fx) * 1i + 2d f^2 x^2 + c f \cos(2e + 2fx) * 2i + 2c f \sin(2e + 2fx) + 4c f^2 x + d f x \cos(2e + 2fx) * 2i + 2d f x \sin(2e + 2fx)) / (8a f^2)$

3.21 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx$

Optimal result	139
Rubi [A] (verified)	139
Mathematica [A] (verified)	141
Maple [A] (verified)	141
Fricas [A] (verification not implemented)	141
Sympy [F]	142
Maxima [A] (verification not implemented)	142
Giac [A] (verification not implemented)	142
Mupad [F(-1)]	143

Optimal result

Integrand size = 23, antiderivative size = 161

$$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx = \frac{\cos\left(2e - \frac{2cf}{d}\right) \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{2ad} + \frac{\log(c+dx)}{2ad} \\ - \frac{i \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{2ad} \\ - \frac{i \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{2ad} \\ - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{2ad}$$

[Out] $1/2*\text{Ci}(2*c*f/d+2*f*x)*\cos(-2*e+2*c*f/d)/a/d+1/2*\ln(d*x+c)/a/d-1/2*I*\cos(-2*e+2*c*f/d)*\text{Si}(2*c*f/d+2*f*x)/a/d+1/2*I*\text{Ci}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a/d+1/2*\text{Si}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a/d$

Rubi [A] (verified)

Time = 0.32 (sec), antiderivative size = 161, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {3807, 3384, 3380, 3383}

$$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))} dx = -\frac{i \text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{2ad} \\ + \frac{\text{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{2ad} \\ - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{2ad} \\ - \frac{i \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(2xf + \frac{2cf}{d}\right)}{2ad} + \frac{\log(c+dx)}{2ad}$$

[In] $\text{Int}[1/((c + d*x)*(a + I*a*Tan[e + f*x])), x]$
[Out] $(\text{Cos}[2e - (2*c*f)/d]*\text{CosIntegral}[(2*c*f)/d + 2*f*x])/(2*a*d) + \text{Log}[c + d*x]/(2*a*d) - ((I/2)*\text{CosIntegral}[(2*c*f)/d + 2*f*x]*\text{Sin}[2e - (2*c*f)/d])/(a*d) - ((I/2)*\text{Cos}[2e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/(a*d) - (\text{Si}[2e - (2*c*f)/d]*\text{SinIntegral}[(2*c*f)/d + 2*f*x])/(2*a*d)$

Rule 3380

$\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{SinIntegral}[e + f*x]/d, x] /; \text{FreeQ}[\{c, d, e, f\}, x] \& \text{EqQ}[d*e - c*f, 0]$

Rule 3383

$\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{CosIntegral}[e - \pi/2 + f*x]/d, x] /; \text{FreeQ}[\{c, d, e, f\}, x] \& \text{EqQ}[d*(e - \pi/2) - c*f, 0]$

Rule 3384

$\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \text{Dist}[\text{Cos}[(d*e - c*f)/d], \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \text{Dist}[\text{Sin}[(d*e - c*f)/d], \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \& \text{NeQ}[d*e - c*f, 0]$

Rule 3807

$\text{Int}[1/(((c_.) + (d_.)*(x_))*(a_.) + (b_.)*\tan[(e_.) + (f_.)*(x_)]), x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{Log}[c + d*x]/(2*a*d), x] + (\text{Dist}[1/(2*a), \text{Int}[\text{Cos}[2e + 2*f*x]/(c + d*x), x], x] + \text{Dist}[1/(2*b), \text{Int}[\text{Sin}[2e + 2*f*x]/(c + d*x), x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \& \text{EqQ}[a^2 + b^2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\log(c + dx)}{2ad} - \frac{i \int \frac{\sin(2e + 2fx)}{c + dx} dx}{2a} + \frac{\int \frac{\cos(2e + 2fx)}{c + dx} dx}{2a} \\ &= \frac{\log(c + dx)}{2ad} - \frac{(i \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} + \frac{\cos(2e - \frac{2cf}{d}) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} \\ &\quad - \frac{(i \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} - \frac{\sin(2e - \frac{2cf}{d}) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} \\ &= \frac{\cos(2e - \frac{2cf}{d}) \text{CosIntegral}(\frac{2cf}{d} + 2fx)}{2ad} + \frac{\log(c + dx)}{2ad} \\ &\quad - \frac{i \text{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2ad} \\ &\quad - \frac{i \cos(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{2ad} - \frac{\sin(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{2ad} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.03

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \frac{\sec(e + fx) (-i \cos(f(\frac{c}{d} + x)) + \sin(f(\frac{c}{d} + x))) \left(\text{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) (\cos(e - \frac{cf}{d}) - i \sin(e - \frac{cf}{d}))\right)}{2ad(-i + \tan(e + fx))}$$

```
[In] Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])),x]
[Out] (Sec[e + f*x]*((-I)*Cos[f*(c/d + x)] + Sin[f*(c/d + x)])*(CosIntegral[(2*f*(c + d*x))/d]*(Cos[e - (c*f)/d] - I*Sin[e - (c*f)/d]) + Log[f*(c + d*x)]*(Cos[e - (c*f)/d] + I*Sin[e - (c*f)/d]) + ((-I)*Cos[e - (c*f)/d] - Sin[e - (c*f)/d])*SinIntegral[(2*f*(c + d*x))/d]))/(2*a*d*(-I + Tan[e + f*x]))
```

Maple [A] (verified)

Time = 0.60 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.40

method	result
risch	$\frac{\ln(dx+c)}{2ad} - \frac{e^{\frac{2i(cf-de)}{d}} \text{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{2ad}$
default	$= \frac{if\left(\frac{2 \text{Si}(2fx+2e+\frac{2cf-2de}{d}) \cos(\frac{2cf-2de}{d}) - 2 \text{Ci}(2fx+2e+\frac{2cf-2de}{d}) \sin(\frac{2cf-2de}{d})}{d}\right)}{4} + \frac{f\left(\frac{2 \text{Si}(2fx+2e+\frac{2cf-2de}{d}) \sin(\frac{2cf-2de}{d}) + 2 \text{Ci}(2fx+2e+\frac{2cf-2de}{d}) \cos(\frac{2cf-2de}{d})}{d}\right)}{4af}$

```
[In] int(1/(d*x+c)/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
[Out] 1/2*ln(d*x+c)/a/d-1/2/a/d*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.32

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \frac{\text{Ei}\left(-\frac{2(i dfx+icf)}{d}\right) e^{\left(-\frac{2(i de-icf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{2 ad}$$

```
[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
[Out] 1/2*(Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + log((d*x + c)/d))/(a*d)
```

Sympy [F]

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = -\frac{i \int \frac{1}{ctan(e+fx)-ic+dx\tan(e+fx)-idx} dx}{a}$$

[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x)

[Out] $-I*Integral(1/(c*tan(e + f*x) - I*c + d*x*tan(e + f*x) - I*d*x), x)/a$

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 112, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \\ -\frac{f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + i f E_1\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) - f \log((J))}{2 adf}$$

[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="maxima")

[Out] $-1/2*(f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f))/(a*d*f)$

Giac [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.86

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \\ \frac{\cos\left(\frac{2cf}{d}\right) \text{Ci}\left(-\frac{2(df+cf)}{d}\right) + \cos(2e) \log(dx + c) + i \log(dx + c) \sin(2e) + i \text{Ci}\left(-\frac{2(df+cf)}{d}\right) \sin\left(\frac{2cf}{d}\right) - i \text{Si}\left(-\frac{2(df+cf)}{d}\right)}{2(ad \cos(2e) + i ad \sin(2e))}$$

[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e)),x, algorithm="giac")

[Out] $1/2*(\cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + \cos(2*e)*log(d*x + c) + I*log(d*x + c)*sin(2*e) + I*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - I*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d))/(a*d*cos(2*e) + I*a*d*sin(2*e))$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))} dx = \int \frac{1}{(a + a \tan(e + fx) 1i) (c + dx)} dx$$

[In] `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)),x)`

[Out] `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)), x)`

3.22 $\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx$

Optimal result	144
Rubi [A] (verified)	144
Mathematica [A] (verified)	146
Maple [A] (verified)	147
Fricas [A] (verification not implemented)	147
Sympy [F]	147
Maxima [A] (verification not implemented)	148
Giac [B] (verification not implemented)	148
Mupad [F(-1)]	149

Optimal result

Integrand size = 23, antiderivative size = 168

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx = & -\frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^2} \\ & - \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^2} \\ & - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^2} \\ & + \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^2} \\ & - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

```
[Out] -I*f*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a/d^2-f*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^2+f*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^2-I*f*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^2-1/d/(d*x+c)/(a+I*a*tan(f*x+e))
```

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.174, Rules used

$$= \{3805, 3384, 3380, 3383\}$$

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx = & -\frac{f \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ & - \frac{if \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ & + \frac{if \sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{ad^2} \\ & - \frac{f \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{ad^2} \\ & - \frac{1}{d(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

[In] $\operatorname{Int}[1/((c+d*x)^2*(a+I*a*Tan[e+f*x])), x]$

[Out] $\frac{(-I)*f*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2) - (f*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sin}[2*e - (2*c*f)/d])/(a*d^2) - (f*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2) + (I*f*\operatorname{Sin}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2) - 1/(d*(c+d*x)*(a+I*a*Tan[e+f*x]))}{}$

Rule 3380

$\operatorname{Int}[\sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x]/d, x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \&& \operatorname{EqQ}[d*e - c*f, 0]$

Rule 3383

$\operatorname{Int}[\sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[\operatorname{CosIntegral}[e - \pi/2 + f*x]/d, x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \&& \operatorname{EqQ}[d*(e - \pi/2) - c*f, 0]$

Rule 3384

$\operatorname{Int}[\sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_{\text{Symbol}}] \Rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \&& \operatorname{NeQ}[d*e - c*f, 0]$

Rule 3805

$\operatorname{Int}[1/(((c_.) + (d_.)*(x_))^2*((a_) + (b_.)*\operatorname{tan}[(e_.) + (f_.)*(x_)])), x_{\text{Symbol}}] \Rightarrow -\operatorname{Simp}[(d*(c + d*x)*(a + b*\operatorname{Tan}[e + f*x]))^{(-1)}, x] + (-\operatorname{Dist}[f/(a*d), \operatorname{Int}[\operatorname{Sin}[2*e + 2*f*x]/(c + d*x), x], x] + \operatorname{Dist}[f/(b*d), \operatorname{Int}[\operatorname{Cos}[2*e + 2*f*x]/(c + d*x), x], x]) /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \operatorname{EqQ}[a^2 + b^2, 0]$

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{1}{d(c+dx)(a+ia\tan(e+fx))} - \frac{(if) \int \frac{\cos(2e+2fx)}{c+dx} dx}{ad} - \frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{ad} \\
&= -\frac{1}{d(c+dx)(a+ia\tan(e+fx))} - \frac{(if \cos(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{ad} \\
&\quad - \frac{(f \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{ad} \\
&\quad + \frac{(if \sin(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{ad} - \frac{(f \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{ad} \\
&= -\frac{if \cos(2e - \frac{2cf}{d}) \text{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^2} \\
&\quad - \frac{f \text{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^2} - \frac{f \cos(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^2} \\
&\quad + \frac{if \sin(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^2} - \frac{1}{d(c+dx)(a+ia\tan(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.77 (sec), antiderivative size = 224, normalized size of antiderivative = 1.33

$$\begin{aligned}
&\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx \\
&= \frac{\sec(e+fx) (\cos(\frac{cf}{d}) + i \sin(\frac{cf}{d})) \left(d(i \cos(e+f(-\frac{c}{d}+x)) + i \cos(e+f(\frac{c}{d}+x)) - \sin(e+f(-\frac{c}{d}+x)) \right. \\
&\quad \left. + I \cos(e+f(c/d+x)) - \sin(e+f(-(c/d)+x)) + \sin(e+f(c/d+x)) \right) - 2*f*(c+d*x)*\text{CosIntegral}[(2*f*(c+d*x))/d]*(\cos[e - (f*(c+d*x))/d] - \\
&\quad I*\sin[e - (f*(c+d*x))/d]) + 2*f*(c+d*x)*(I*\cos[e - (f*(c+d*x))/d] + \\
&\quad \sin[e - (f*(c+d*x))/d])* \text{SinIntegral}[(2*f*(c+d*x))/d])/(2*a*d^2*(c+d*x)*(-I + \tan[e+f*x]))
\end{aligned}$$

```

[In] Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])),x]
[Out] (Sec[e + f*x]*(Cos[(c*f)/d] + I*Sin[(c*f)/d])*(d*(I*Cos[e + f*(-(c/d) + x)] +
+ I*Cos[e + f*(c/d + x)] - Sin[e + f*(-(c/d) + x)] + Sin[e + f*(c/d + x)]) -
- 2*f*(c + d*x)*CosIntegral[(2*f*(c + d*x))/d]*(Cos[e - (f*(c + d*x))/d] -
I*Sin[e - (f*(c + d*x))/d]) + 2*f*(c + d*x)*(I*Cos[e - (f*(c + d*x))/d] +
Sin[e - (f*(c + d*x))/d])*SinIntegral[(2*f*(c + d*x))/d]))/(2*a*d^2*(c + d*x)*(-I + Tan[e + f*x]))

```

Maple [A] (verified)

Time = 0.51 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.57

method	result
risch	$-\frac{1}{2d(dx+c)a} - \frac{f e^{-2i(fx+e)}}{2a(df x+cf)d} + \frac{if e^{\frac{2i(cf-de)}{d}} \text{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{ad^2}$
default	$-\frac{if^2 \left(-\frac{2 \sin(2fx+2e)}{(cf-de+d(fx+e))d} + \frac{4 \text{Si}\left(2fx+2e+\frac{2cf-2de}{d}\right) \sin\left(\frac{2cf-2de}{d}\right)}{d} + \frac{4 \text{Ci}\left(2fx+2e+\frac{2cf-2de}{d}\right) \cos\left(\frac{2cf-2de}{d}\right)}{d} \right)}{4} + \frac{f^2 \left(-\frac{2 \cos(2fx+2e)}{(cf-de+d(fx+e))d} \right)}{af}$

[In] `int(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{1}{2}/d/(d*x+c)/a - \frac{1}{2}/a*f*exp(-2*I*(f*x+e))/(d*f*x+c*f)/d + I/a*f/d^2*exp(2*I*(c*f-d*e)/d)*Ei(1, 2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)$$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 83, normalized size of antiderivative = 0.49

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx \\ = -\frac{\left(\left(2(i df x + i cf) \text{Ei}\left(-\frac{2(i df x + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + d \right) e^{(2i fx + 2i e)} + d \right) e^{(-2i fx - 2i e)}}{2(ad^3 x + acd^2)}$$

[In] `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

[Out]
$$-\frac{1}{2}*((2*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^{-(-2*(I*d*e - I*c*f)/d)} + d)*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-2*I*f*x - 2*I*e)/(a*d^3*x + a*c*d^2)}$$

Sympy [F]

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx \\ = -\frac{i \int \frac{1}{c^2 \tan(e + fx) - ic^2 + 2cdx \tan(e + fx) - 2icdx + d^2 x^2 \tan(e + fx) - id^2 x^2} dx}{a}$$

[In] `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e)),x)`

[Out]
$$-I*\text{Integral}(1/(c**2*tan(e + f*x) - I*c**2 + 2*c*d*x*tan(e + f*x) - 2*I*c*d*x + d**2*x**2*tan(e + f*x) - I*d**2*x**2), x)/a$$

Maxima [A] (verification not implemented)

none

Time = 0.58 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.71

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx =$$

$$-\frac{f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + i f^2 E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + f^2}{2((fx+e)ad^2 - ad^2e + acdf)f}$$

```
[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="maxima")
[Out] -1/2*(f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^2)/(((f*x + e)*a*d^2 - a*d^2*e + a*c*d*f)*f)
```

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1013 vs. 2(161) = 322.

Time = 2.30 (sec) , antiderivative size = 1013, normalized size of antiderivative = 6.03

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx = \text{Too large to display}$$

```
[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e)),x, algorithm="giac")
[Out] 1/2*(-2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) + 2*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*sin(-2*(d*e - c*f)/d) - 2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*d*e*f^2*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*c*f^3*cos(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*sin(-2*(d*e - c*f)/d)*sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) +
```

$f) - d*e + c*f)/d) + 2*I*d*e*f^2*\sin(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*I*c*f^3*\sin(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - d*f^2*\cos(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - I*d*f^2*\sin(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - d*f^2*d^2/((d*x + c)*a*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a*d^5*e + a*c*d^4*f)*f)$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))} dx = \int \frac{1}{(a + a \tan(e + fx) 1i) (c + dx)^2} dx$$

[In] `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2),x)`

[Out] `int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^2), x)`

3.23 $\int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx$

Optimal result	150
Rubi [A] (verified)	151
Mathematica [A] (verified)	153
Maple [A] (verified)	153
Fricas [A] (verification not implemented)	154
Sympy [F]	154
Maxima [A] (verification not implemented)	154
Giac [B] (verification not implemented)	155
Mupad [F(-1)]	155

Optimal result

Integrand size = 23, antiderivative size = 227

$$\begin{aligned} \int \frac{1}{(c+dx)^3(a+ia\tan(e+fx))} dx = & -\frac{if}{2ad^2(c+dx)} \\ & - \frac{f^2 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & + \frac{if^2 \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^3} \\ & + \frac{if^2 \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & + \frac{f^2 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} \\ & + \frac{if}{d^2(c+dx)(a+ia\tan(e+fx))} \end{aligned}$$

```
[Out] -1/2*I*f/a/d^2/(d*x+c)-f^2*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a/d^3+I*f^2*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a/d^3-I*f^2*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^3-f^2*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a/d^3-1/2/d/(d*x+c)^2/(a+I*a*tan(f*x+e))+I*f/d^2/(d*x+c)/(a+I*a*tan(f*x+e))
```

Rubi [A] (verified)

Time = 0.38 (sec) , antiderivative size = 227, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {3806, 3805, 3384, 3380, 3383}

$$\begin{aligned} \int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx = & \frac{if^2 \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \sin(2e - \frac{2cf}{d})}{ad^3} \\ & - \frac{f^2 \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \cos(2e - \frac{2cf}{d})}{ad^3} \\ & + \frac{f^2 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{ad^3} \\ & + \frac{if^2 \cos(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{ad^3} \\ & + \frac{if}{d^2(c + dx)(a + ia \tan(e + fx))} - \frac{if}{2ad^2(c + dx)} \\ & - \frac{1}{2d(c + dx)^2(a + ia \tan(e + fx))} \end{aligned}$$

[In] $\operatorname{Int}[1/((c + d*x)^3(a + I*a*\operatorname{Tan}[e + f*x])), x]$

[Out] $\frac{((-1/2*I)*f)/(a*d^2*(c + d*x)) - (f^2*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/(a*d^3) + (I*f^2*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])* \operatorname{Sin}[2*e - (2*c*f)/d]/(a*d^3) + (I*f^2*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a*d^3) + (f^2*\operatorname{Sin}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/ (a*d^3) - 1/(2*d*(c + d*x)^2*(a + I*a*\operatorname{Tan}[e + f*x])) + (I*f)/(d^2*(c + d*x)*(a + I*a*\operatorname{Tan}[e + f*x]))}{2d(c + dx)^2(a + ia \tan(e + fx))}$

Rule 3380

$\operatorname{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.*)(x_.)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[\operatorname{SinIntegral}[e + f*x]/d, x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \& \operatorname{EqQ}[d*e - c*f, 0]$

Rule 3383

$\operatorname{Int}[\sin[(e_.) + (f_.*)(x_.)]/((c_.) + (d_.*)(x_.)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[\operatorname{CosIntegral}[e - \pi/2 + f*x]/d, x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \& \operatorname{EqQ}[d*(e - \pi/2) - c*f, 0]$

Rule 3384

$\operatorname{Int}[\sin[(e_.) + (f_.*)(x_.)]/((c_.) + (d_.*)(x_.)), x_{\text{Symbol}}] \Rightarrow \operatorname{Dist}[\operatorname{Cos}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \operatorname{Dist}[\operatorname{Sin}[(d*e - c*f)/d], \operatorname{Int}[\operatorname{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \operatorname{FreeQ}[\{c, d, e, f\}, x] \& \operatorname{NeQ}[d*e - c*f, 0]$

Rule 3805

```
Int[1/(((c_) + (d_)*(x_))^2*((a_) + (b_)*tan[(e_) + (f_)*(x_)])), x_Symbol] :> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Dist[f/(a*d)], Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Dist[f/(b*d)], Int[Cos[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]
```

Rule 3806

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] :> Simp[f*((c + d*x)^(m + 2)/(b*d^2*(m + 1)*(m + 2))), x] + (Dist[2*b*(f/(a*d*(m + 1))), Int[(c + d*x)^(m + 1)/(a + b*Tan[e + f*x]), x], x] + Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + b*Tan[e + f*x])), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[m, -1] && NeQ[m, -2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} - \frac{(if)\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))} dx}{d} \\
&= -\frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} \\
&\quad + \frac{if}{d^2(c+dx)(a+ia\tan(e+fx))} + \frac{(if^2)\int \frac{\sin(2e+2fx)}{c+dx} dx}{ad^2} - \frac{f^2\int \frac{\cos(2e+2fx)}{c+dx} dx}{ad^2} \\
&= -\frac{if}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} + \frac{if}{d^2(c+dx)(a+ia\tan(e+fx))} \\
&\quad + \frac{(if^2 \cos(2e - \frac{2cf}{d}))\int \frac{\sin(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} - \frac{(f^2 \cos(2e - \frac{2cf}{d}))\int \frac{\cos(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&\quad + \frac{(if^2 \sin(2e - \frac{2cf}{d}))\int \frac{\cos(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} + \frac{(f^2 \sin(2e - \frac{2cf}{d}))\int \frac{\sin(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&= -\frac{if}{2ad^2(c+dx)} - \frac{f^2 \cos(2e - \frac{2cf}{d}) \text{CosIntegral}(\frac{2cf}{d} + 2fx)}{ad^3} \\
&\quad + \frac{if^2 \text{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{ad^3} \\
&\quad + \frac{if^2 \cos(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^3} + \frac{f^2 \sin(2e - \frac{2cf}{d}) \text{Si}(\frac{2cf}{d} + 2fx)}{ad^3} \\
&\quad - \frac{1}{2d(c+dx)^2(a+ia\tan(e+fx))} + \frac{if}{d^2(c+dx)(a+ia\tan(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.00 (sec) , antiderivative size = 285, normalized size of antiderivative = 1.26

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ = \frac{\sec(e + fx) (\cos(\frac{cf}{d}) + i \sin(\frac{cf}{d})) \left(d(id \cos(e + f(-\frac{c}{d} + x)) + (id + 2cf + 2dfx) \cos(e + f(\frac{c}{d} + x))) -$$

```
[In] Integrate[1/((c + d*x)^3*(a + I*a*Tan[e + f*x])),x]
[Out] (Sec[e + f*x]*(Cos[(c*f)/d] + I*Sin[(c*f)/d])*(d*(I*d*Cos[e + f*(-(c/d) + x)]) + (I*d + 2*c*f + 2*d*f*x)*Cos[e + f*(c/d + x)] - d*Sin[e + f*(-(c/d) + x)] + d*Sin[e + f*(c/d + x)] - (2*I)*c*f*Sin[e + f*(c/d + x)] - (2*I)*d*f*x*Sin[e + f*(c/d + x)] + 4*f^2*(c + d*x)^2*CosIntegral[(2*f*(c + d*x))/d]*(I*Cos[e - (f*(c + d*x))/d] + Sin[e - (f*(c + d*x))/d]) + 4*f^2*(c + d*x)^2*(Cos[e - (f*(c + d*x))/d] - I*Sin[e - (f*(c + d*x))/d])*SinIntegral[(2*f*(c + d*x))/d]))/(4*a*d^3*(c + d*x)^2*(-I + Tan[e + f*x]))
```

Maple [A] (verified)

Time = 0.57 (sec) , antiderivative size = 216, normalized size of antiderivative = 0.95

method	result
risch	$-\frac{1}{4ad(dx+c)^2} + \frac{if^3e^{-2i(fx+e)}x}{2ad(d^2x^2f^2+2cd^2x^2+c^2f^2)} - \frac{f^2e^{-2i(fx+e)}}{4ad(d^2x^2f^2+2cd^2x^2+c^2f^2)} + \frac{if^3e^{-2i(fx+e)}c}{2a^2(d^2x^2f^2+2cd^2x^2+c^2f^2)} + \frac{f^2e^{2i(cf-d)}}{4a^2(d^2x^2f^2+2cd^2x^2+c^2f^2)}$ $if^3 \left(-\frac{\sin(2fx+2e)}{(cf-de+d(fx+e))^2d} + \frac{-2\cos(2fx+2e)}{(cf-de+d(fx+e))d} - \frac{2 \left(\frac{2 \operatorname{Si}(2fx+2e+\frac{2cf-2de}{d}) \cos(\frac{2cf-2de}{d})}{d} - \frac{2 \operatorname{Ci}(2fx+2e+\frac{2cf-2de}{d}) \sin(\frac{2cf-2de}{d})}{d} \right)}{d} \right)$
default	$-\frac{1}{4}$

```
[In] int(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x,method=_RETURNVERBOSE)
[Out] -1/4/a/d/(d*x+c)^2+1/2*I/a*f^3*exp(-2*I*(f*x+e))/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2*x)+c^2*f^2*x-1/4/a*f^2*exp(-2*I*(f*x+e))/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)+1/2*I/a*f^3*exp(-2*I*(f*x+e))/d^2/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*c+1/a*f^2/d^3*exp(2*I*(c*f-d*e))/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e))/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.57

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ = \frac{\left(2i d^2 f x + 2i c d f - d^2 - \left(4(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2) \text{Ei}\left(-\frac{2(i d f x + i c f)}{d}\right) e^{\left(-\frac{2(i d e - i c f)}{d}\right)} + d^2 \right) e^{(2i f x + 2i e)} \right)}{4(ad^5 x^2 + 2acd^4 x + ac^2 d^3)}$$

```
[In] integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="fricas")
[Out] 1/4*(2*I*d^2*f*x + 2*I*c*d*f - d^2 - (4*(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + d^2)*e^(2*I*f*x + 2*I*e))*e^(-2*I*f*x - 2*I*e)/(a*d^5*x^2 + 2*a*c*d^4*x + a*c^2*d^3)
```

Sympy [F]

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx \\ = -\frac{i \int \frac{1}{c^3 \tan(e+fx) - ic^3 + 3c^2 dx \tan(e+fx) - 3ic^2 dx + 3cd^2 x^2 \tan(e+fx) - 3icd^2 x^2 + d^3 x^3 \tan(e+fx) - id^3 x^3} dx}{a}$$

```
[In] integrate(1/(d*x+c)**3/(a+I*a*tan(f*x+e)),x)
[Out] -I*Integral(1/(c**3*tan(e + f*x) - I*c**3 + 3*c**2*d*x*tan(e + f*x) - 3*I*c**2*d*x + 3*c*d**2*x**2*tan(e + f*x) - 3*I*c*d**2*x**2 + d**3*x**3*tan(e + f*x) - I*d**3*x**3), x)/a
```

Maxima [A] (verification not implemented)

none

Time = 0.60 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.70

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx = \\ -\frac{2 f^3 \cos\left(-\frac{2(de-cf)}{d}\right) E_3\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + 2i f^3 E_3\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) \sin\left(-\frac{2(de-cf)}{d}\right) + f^3}{4((fx+e)^2 ad^3 + ad^3 e^2 - 2acd^2 ef + ac^2 df^2 - 2(ad^3 e - acd^2 f)(fx+e))f}$$

```
[In] integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="maxima")
[Out] -1/4*(2*f^3*cos(-2*(d*e - c*f)/d)*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^3*exp_integral_e(3, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + f^3)/(((f*x + e)^2*a*d^3 + a*d^3*e^2 - 2*a*c*d^2*e*f + a*c^2*d*f^2 - 2*(a*d^3*e - a*c*d^2*f)*(f*x + e))*f)
```

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(212) = 424$.

Time = 0.44 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.34

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx =$$

$$-\frac{4 d^2 f^2 x^2 \cos\left(\frac{2 c f}{d}\right) \text{Ci}\left(-\frac{2 (d f x+c f)}{d}\right)+4 i d^2 f^2 x^2 \text{Ci}\left(-\frac{2 (d f x+c f)}{d}\right) \sin\left(\frac{2 c f}{d}\right)-4 i d^2 f^2 x^2 \cos\left(\frac{2 c f}{d}\right) \text{Si}\left(\frac{2 (d f x+c f)}{d}\right)}{ }$$

```
[In] integrate(1/(d*x+c)^3/(a+I*a*tan(f*x+e)),x, algorithm="giac")
[Out] -1/4*(4*d^2*f^2*x^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 4*I*d^2*f^2*x^2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 4*I*d^2*f^2*x^2*cos_integral(2*(d*f*x + c*f)/d) + 4*d^2*f^2*x^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 8*c*d*f^2*x*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 8*I*c*d*f^2*x*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 8*I*c*d*f^2*x*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 8*c*d*f^2*x*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 4*c^2*f^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + 4*I*c^2*f^2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 4*I*c^2*f^2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 4*c^2*f^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) - 2*I*d^2*f*x*cos(2*f*x) - 2*d^2*f*x*sin(2*f*x) - 2*I*c*d*f*cos(2*f*x) - 2*c*d*f*sin(2*f*x) + d^2*cos(2*f*x) + d^2*cos(2*e) - I*d^2*sin(2*f*x) + I*d^2*sin(2*e))/(a*d^5*x^2*cos(2*e) + I*a*d^5*x^2*sin(2*e) + 2*a*c*d^4*x*cos(2*e) + 2*I*a*c*d^4*x*sin(2*e) + a*c^2*d^3*cos(2*e) + I*a*c^2*d^3*sin(2*e))
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^3(a + ia \tan(e + fx))} dx = \int \frac{1}{(a + a \tan(e + fx) 1i) (c + dx)^3} dx$$

```
[In] int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3),x)
```

```
[Out] int(1/((a + a*tan(e + f*x)*1i)*(c + d*x)^3), x)
```

3.24 $\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx$

Optimal result	156
Rubi [A] (verified)	156
Mathematica [A] (verified)	158
Maple [A] (verified)	159
Fricas [A] (verification not implemented)	159
Sympy [A] (verification not implemented)	160
Maxima [F(-2)]	160
Giac [A] (verification not implemented)	161
Mupad [B] (verification not implemented)	161

Optimal result

Integrand size = 23, antiderivative size = 270

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx = & -\frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4} - \frac{3id^2e^{-2ie-2ifx}(c+dx)}{8a^2f^3} \\ & - \frac{3id^2e^{-4ie-4ifx}(c+dx)}{128a^2f^3} + \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2f^2} \\ & + \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} \\ & + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} + \frac{(c+dx)^4}{16a^2d} \end{aligned}$$

```
[Out] -3/16*d^3*exp(-2*I*e-2*I*f*x)/a^2/f^4-3/512*d^3*exp(-4*I*e-4*I*f*x)/a^2/f^4
-3/8*I*d^2*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^3-3/128*I*d^2*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^2/f^3+3/8*d*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^2/f^2
+1/16*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^2/f^2+1/16*(d*x+c)^4/a^2/d
```

Rubi [A] (verified)

Time = 0.33 (sec), antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used

= {3810, 2207, 2225}

$$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx = -\frac{3id^2(c+dx)e^{-2ie-2ifx}}{8a^2f^3} - \frac{3id^2(c+dx)e^{-4ie-4ifx}}{128a^2f^3} + \frac{3d(c+dx)^2e^{-2ie-2ifx}}{8a^2f^2} + \frac{3d(c+dx)^2e^{-4ie-4ifx}}{64a^2f^2} + \frac{i(c+dx)^3e^{-2ie-2ifx}}{4a^2f} + \frac{i(c+dx)^3e^{-4ie-4ifx}}{16a^2f} + \frac{(c+dx)^4}{16a^2d} - \frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4}$$

[In] $\text{Int}[(c+d*x)^3/(a+I*a*\text{Tan}[e+f*x])^2, x]$

[Out] $(-3*d^3*E^((-2*I)*e - (2*I)*f*x))/(16*a^2*f^4) - (3*d^3*E^((-4*I)*e - (4*I)*f*x))/(512*a^2*f^4) - (((3*I)/8)*d^2*E^((-2*I)*e - (2*I)*f*x)*(c+d*x))/(a^2*f^3) - (((3*I)/128)*d^2*E^((-4*I)*e - (4*I)*f*x)*(c+d*x))/(a^2*f^3) + (3*d*E^((-2*I)*e - (2*I)*f*x)*(c+d*x)^2)/(8*a^2*f^2) + (3*d*E^((-4*I)*e - (4*I)*f*x)*(c+d*x)^2)/(64*a^2*f^2) + ((I/4)*E^((-2*I)*e - (2*I)*f*x)*(c+d*x)^3)/(a^2*f) + ((I/16)*E^((-4*I)*e - (4*I)*f*x)*(c+d*x)^3)/(a^2*f) + (c+d*x)^4/(16*a^2*d)$

Rule 2207

$\text{Int}[((b_)*(F_)^((g_)*(e_) + (f_)*(x_)))^{(n_)}*((c_)*(d_)*(x_))^{(m_)}, x_{\text{Symbol}}] := \text{Simp}[(c+d*x)^m*((b*F^((g*(e+f*x))))^n/(f*g*n*\text{Log}[F])), x] - \text{Dist}[d*(m/(f*g*n*\text{Log}[F])), \text{Int}[(c+d*x)^{(m-1)}*(b*F^((g*(e+f*x))))^n, x], x] /; \text{FreeQ}[\{F, b, c, d, e, f, g, n\}, x] \&& \text{GtQ}[m, 0] \&& \text{IntegerQ}[2*m] \&& \text{!TrueQ}[\$UseGamma]$

Rule 2225

$\text{Int}[(F_)^((c_)*(a_) + (b_)*(x_)))^{(n_)}, x_{\text{Symbol}}] := \text{Simp}[(F^(c*(a+b*x)))^n/(b*c*n*\text{Log}[F]), x] /; \text{FreeQ}[\{F, a, b, c, n\}, x]$

Rule 3810

$\text{Int}[(c_*(d_)*(x_))^{(m_)}*((a_)*(b_)*\text{tan}[(e_)*(f_)*(x_)])^{(n_)}, x_{\text{Symbol}}] := \text{Int}[\text{ExpandIntegrand}[(c+d*x)^m, (1/(2*a) + E^(2*(a/b)*(e+f*x))/(2*a))^{(-n)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, 0]$

Rubi steps

$$\text{integral} = \int \left(\frac{(c+dx)^3}{4a^2} + \frac{e^{-2ie-2ifx}(c+dx)^3}{2a^2} + \frac{e^{-4ie-4ifx}(c+dx)^3}{4a^2} \right) dx$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{16a^2d} + \frac{\int e^{-4ie-4ifx}(c+dx)^3 dx}{4a^2} + \frac{\int e^{-2ie-2ifx}(c+dx)^3 dx}{2a^2} \\
&= \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} + \frac{(c+dx)^4}{16a^2d} \\
&\quad - \frac{(3id)\int e^{-4ie-4ifx}(c+dx)^2 dx}{16a^2f} - \frac{(3id)\int e^{-2ie-2ifx}(c+dx)^2 dx}{4a^2f} \\
&= \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2f^2} + \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} \\
&\quad + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} + \frac{(c+dx)^4}{16a^2d} - \frac{(3d^2)\int e^{-4ie-4ifx}(c+dx) dx}{32a^2f^2} \\
&\quad - \frac{(3d^2)\int e^{-2ie-2ifx}(c+dx) dx}{4a^2f^2} \\
&= -\frac{3id^2e^{-2ie-2ifx}(c+dx)}{8a^2f^3} - \frac{3id^2e^{-4ie-4ifx}(c+dx)}{128a^2f^3} + \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2f^2} \\
&\quad + \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} \\
&\quad + \frac{(c+dx)^4}{16a^2d} + \frac{(3id^3)\int e^{-4ie-4ifx} dx}{128a^2f^3} + \frac{(3id^3)\int e^{-2ie-2ifx} dx}{8a^2f^3} \\
&= -\frac{3d^3e^{-2ie-2ifx}}{16a^2f^4} - \frac{3d^3e^{-4ie-4ifx}}{512a^2f^4} - \frac{3id^2e^{-2ie-2ifx}(c+dx)}{8a^2f^3} \\
&\quad - \frac{3id^2e^{-4ie-4ifx}(c+dx)}{128a^2f^3} + \frac{3de^{-2ie-2ifx}(c+dx)^2}{8a^2f^2} + \frac{3de^{-4ie-4ifx}(c+dx)^2}{64a^2f^2} \\
&\quad + \frac{ie^{-2ie-2ifx}(c+dx)^3}{4a^2f} + \frac{ie^{-4ie-4ifx}(c+dx)^3}{16a^2f} + \frac{(c+dx)^4}{16a^2d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.54 (sec), antiderivative size = 473, normalized size of antiderivative = 1.75

$$\begin{aligned}
&\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx \\
&= \frac{\sec^2(e+fx)(\cos(fx)+i\sin(fx))^2 ((4ic^3f^3+6c^2df^2(1+2ifx)+6cd^2f(-i+2fx+2if^2x^2)+d^3(-3-6ifx^2))}{(a+ia\tan(e+fx))^2}
\end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^2, x]

[Out] $\left(\frac{1}{2} \operatorname{Sec}[e+f x]^2 (\operatorname{Cos}[f x]+\operatorname{I} \operatorname{Sin}[f x])^2 ((4 \operatorname{I}) c^3 f^3+6 c^2 d f^2 (1+2 i f x)+6 c d^2 f (-i+2 f x+2 i f^2 x^2)+d^3 (-3-(6 \operatorname{I}) f x)+6 f^2 x^2+(4 \operatorname{I}) f^3 x^3)) \operatorname{Cos}[2 f x]+((32 \operatorname{I}) c^3 f^3+24 c^2 d f^2 (1+(4 \operatorname{I}) f x)+12 c d^2 f (-i+4 f x+(8 \operatorname{I}) f^2 x^2)+d^3 (-3-(12 \operatorname{I}) f x)+24 f^2 x^2+(32 \operatorname{I}) f^3 x^3)) \operatorname{Cos}[4 f x] (\operatorname{Cos}[2 e]-\operatorname{I} \operatorname{Sin}[2 e]))\right)$

$$32 + f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cos[2*e] + I*Sin[2*e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(-I + 2*f*x) + 6*c*d^2*f*(-1 - (2*I)*f*x + 2*f^2*x^2) + d^3*(3*I - 6*f*x - (6*I)*f^2*x^2 + 4*f^3*x^3))*Sin[2*f*x] + ((3*2*c^3*f^3 + 24*c^2*d*f^2*(-I + 4*f*x) + 12*c*d^2*f*(-1 - (4*I)*f*x + 8*f^2*x^2) + d^3*(3*I - 12*f*x - (24*I)*f^2*x^2 + 32*f^3*x^3))*(Cos[2*e] - I*Sin[2*e])*Sin[4*f*x])/32))/((16*f^4*(a + I*a*Tan[e + f*x]))^2)$$

Maple [A] (verified)

Time = 0.75 (sec), antiderivative size = 283, normalized size of antiderivative = 1.05

method	result
risch	$\frac{d^3x^4}{16a^2} + \frac{d^2cx^3}{4a^2} + \frac{3dc^2x^2}{8a^2} + \frac{c^3x}{4a^2} + \frac{c^4}{16a^2d} + \frac{i(4d^3x^3f^3 + 12cd^2f^3x^2 - 6id^3f^2x^2 + 12c^2df^3x - 12icd^2f^2x + 4c^3f^3 - 6ic^2df^2 - 6ic^3f^4)}{16a^2f^4}$

[In] `int((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{16}/a^2*d^3*x^4 + \frac{1}{4}/a^2*d^2*c*x^3 + \frac{3}{8}/a^2*d*c^2*x^2 + \frac{1}{4}/a^2*c^3*x + \frac{1}{16}/a^2/d^4 + \frac{1}{16*I}*(4*d^3*x^3*f^3 - 6*I*d^3*x^2*f^2*x^2 + 12*c*d^2*f^3*x^2 - 12*I*c*d^2*f^2*x^3 + 12*c^2*d*f^3*x^3 - 6*I*c^2*d*f^2*x^2 + 4*c^3*f^3*x^3 - 6*d^3*x^3*f*x + 3*I*d^3*x^3 - 6*c*d^2*f^2)/a^2/f^4*\exp(-2*I*(f*x+e)) + \frac{1}{512*I}*(32*d^3*x^3*f^3 - 24*I*d^3*x^2*f^2*x^2 + 96*c*d^2*f^3*x^2 - 48*I*c*d^2*f^2*x + 96*c^2*d*f^3*x^2 - 24*I*c^2*d*f^2*x^3 - 32*c^3*f^3*x^3 - 12*d^3*f*x^3 - 3*I*d^3*x^3 - 12*c*d^2*f^2)/a^2/f^4*\exp(-4*I*(f*x+e))$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec), antiderivative size = 267, normalized size of antiderivative = 0.99

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(32i d^3 f^3 x^3 + 32i c^3 f^3 + 24 c^2 d f^2 - 12i c d^2 f - 3 d^3 - 24 (-4i c d^2 f^3 - d^3 f^2) x^2 - 12 (-8i c^2 d f^3 - 4 c d^2 f^2))}{(a^2 + 2ia \tan(e + fx))^3}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

[Out] $\frac{1}{512}*(32*I*d^3*x^3*f^3 + 32*I*c^3*x^3*f^3 + 24*c^2*d*f^2 - 12*I*c*d^2*f^2 - 3*d^3 - 24*(-4*I*c*d^2*f^3 - d^3*f^2)*x^2 - 12*(-8*I*c^2*d*f^3 - 4*c*d^2*f^2 + I*d^3*f)*x + 32*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x)*e^(4*I*f*x + 4*I*e) - 32*(-4*I*d^3*x^3*f^3 - 4*I*c^3*x^3*f^3 - 6*c^2*d*f^2 + 6*I*c*d^2*f^2 + 3*d^3 + 6*(-2*I*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(-2*I*c^2*d*f^3 - 2*c*d^2*f^2 + I*d^3*f)*x)*e^(2*I*f*x + 2*I*e)*e^(-4*I*f*x - 4*I*e)/(a^2*f^4)$

Sympy [A] (verification not implemented)

Time = 0.42 (sec) , antiderivative size = 665, normalized size of antiderivative = 2.46

$$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx$$

$$= \begin{cases} \frac{((512ia^2c^3f^7e^{2ie}+1536ia^2c^2df^7xe^{2ie}+384a^2c^2df^6e^{2ie}+1536ia^2cd^2f^7x^2e^{2ie}+768a^2cd^2f^6xe^{2ie}-192ia^2cd^2f^5e^{2ie}+512ia^2d^3f^7x^3e^{2ie}+384a^2d^2e^{2ie})}{(16a^2)} + \frac{x^3\cdot(2cd^2e^{2ie}+cd^2)e^{-4ie}}{4a^2} + \frac{x^2\cdot(6c^2de^{2ie}+3c^2d)e^{-4ie}}{8a^2} + \frac{x(2c^3e^{2ie}+c^3)e^{-4ie}}{4a^2} \\ + \frac{c^3x}{4a^2} + \frac{3c^2dx^2}{8a^2} + \frac{cd^2x^3}{4a^2} + \frac{d^3x^4}{16a^2} \end{cases}$$

[In] `integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**2,x)`

[Out] `Piecewise(((512*I*a**2*c**3*f**7*exp(2*I*e) + 1536*I*a**2*c**2*d*f**7*x*exp(2*I*e) + 384*a**2*c**2*d*f**6*exp(2*I*e) + 1536*I*a**2*c*d**2*f**7*x**2*exp(2*I*e) + 768*a**2*c*d**2*f**6*x*exp(2*I*e) - 192*I*a**2*c*d**2*f**5*exp(2*I*e) + 512*I*a**2*d**3*f**7*x**3*exp(2*I*e) + 384*a**2*d**3*f**6*x**2*exp(2*I*e) - 192*I*a**2*d**3*f**5*x*exp(2*I*e) - 48*a**2*d**3*f**4*exp(2*I*e))*exp(-4*I*f*x) + (2048*I*a**2*c**3*f**7*exp(4*I*e) + 6144*I*a**2*c**2*d*f**7*x*exp(4*I*e) + 3072*a**2*c**2*d*f**6*exp(4*I*e) + 6144*I*a**2*c*d**2*f**7*x**2*exp(4*I*e) + 6144*a**2*c*d**2*f**6*x*exp(4*I*e) - 3072*I*a**2*c*d**2*f**5*exp(4*I*e) + 2048*I*a**2*d**3*f**7*x**3*exp(4*I*e) + 3072*a**2*d**3*f**6*x**2*exp(4*I*e) - 3072*I*a**2*d**3*f**5*x*exp(4*I*e) - 1536*a**2*d**3*f**4*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(8192*a**4*f**8*exp(6*I*e), 0)), (x**4*(2*d**3*exp(2*I*e) + d**3)*exp(-4*I*e)/(16*a**2) + x**3*(2*c*d**2*exp(2*I*e) + c*d**2)*exp(-4*I*e)/(4*a**2) + x**2*(6*c**2*d*exp(2*I*e) + 3*c**2*d)*exp(-4*I*e)/(8*a**2) + x*(2*c**3*exp(2*I*e) + c**3)*exp(-4*I*e)/(4*a**2), True)) + c**3*x/(4*a**2) + 3*c**2*d*x**2/(8*a**2) + c*d**2*x**3/(4*a**2) + d**3*x**4/(16*a**2)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^2} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

[Out] `Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.`

Giac [A] (verification not implemented)

none

Time = 0.43 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.36

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(32 d^3 f^4 x^4 e^{(4i f x + 4i e)} + 128 c d^2 f^4 x^3 e^{(4i f x + 4i e)} + 192 c^2 d f^4 x^2 e^{(4i f x + 4i e)} + 128 i d^3 f^3 x^3 e^{(2i f x + 2i e)} + 32 i d^3 f^3 x^2 e^{(2i f x + 2i e)})}{(a + ia \tan(e + fx))^2}$$

```
[In] integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
[Out] 1/512*(32*d^3*f^4*x^4*e^(4*I*f*x + 4*I*e) + 128*c*d^2*f^4*x^3*e^(4*I*f*x + 4*I*e) + 192*c^2*d*f^4*x^2*e^(4*I*f*x + 4*I*e) + 128*I*d^3*f^3*x^3*e^(2*I*f*x + 2*I*e) + 32*I*d^3*f^3*x^3 + 128*c^3*f^4*x*e^(4*I*f*x + 4*I*e) + 384*I*c*d^2*f^3*x^2*e^(2*I*f*x + 2*I*e) + 96*I*c*d^2*f^3*x^2 + 384*I*c^2*d*f^3*x*x^2*e^(2*I*f*x + 2*I*e) + 192*d^3*f^2*x^2*e^(2*I*f*x + 2*I*e) + 96*I*c^2*d*f^3*x + 24*d^3*f^2*x^2 + 128*I*c^3*f^3*x^3*e^(2*I*f*x + 2*I*e) + 384*c*d^2*f^2*x*x^2*e^(2*I*f*x + 2*I*e) + 32*I*c^3*f^3 + 48*c*d^2*f^2*x + 192*c^2*d*f^2*x^2*e^(2*I*f*x + 2*I*e) - 192*I*d^3*f*x*x^2*e^(2*I*f*x + 2*I*e) + 24*c^2*d*f^2*x^2 - 12*I*d^3*f*x^3 - 192*I*c*d^2*f*x^2*e^(2*I*f*x + 2*I*e) - 12*I*c*d^2*f*x^2 - 96*d^3*x^3*e^(2*I*f*x + 2*I*e) - 3*d^3*x^3)*e^(-4*I*f*x - 4*I*e)/(a^2*f^4)
```

Mupad [B] (verification not implemented)

Time = 4.30 (sec) , antiderivative size = 289, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^2} dx = e^{-e 2i - f x 2i} \left(\frac{(4 c^3 f^3 - c^2 d f^2 6i - 6 c d^2 f + d^3 3i) 1i}{16 a^2 f^4} + \frac{d^3 x^3 1i}{4 a^2 f} \right. \\ - \frac{d x (-2 c^2 f^2 + c d f 2i + d^2) 3i}{8 a^2 f^3} - \frac{d^2 x^2 (-2 c f + d 1i) 3i}{8 a^2 f^2} \\ \left. + e^{-e 4i - f x 4i} \left(\frac{(32 c^3 f^3 - c^2 d f^2 24i - 12 c d^2 f + d^3 3i) 1i}{512 a^2 f^4} \right. \right. \\ + \frac{d^3 x^3 1i}{16 a^2 f} - \frac{d x (-8 c^2 f^2 + c d f 4i + d^2) 3i}{128 a^2 f^3} \\ \left. \left. - \frac{d^2 x^2 (-4 c f + d 1i) 3i}{64 a^2 f^2} \right) + \frac{c^3 x}{4 a^2} + \frac{d^3 x^4}{16 a^2} + \frac{3 c^2 d x^2}{8 a^2} + \frac{c d^2 x^3}{4 a^2} \right)$$

```
[In] int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^2,x)
```

```
[Out] exp(- e*2i - f*x*2i)*(((d^3*3i + 4*c^3*f^3 - c^2*d*f^2*6i - 6*c*d^2*f)*1i)/(16*a^2*f^4) + (d^3*x^3*1i)/(4*a^2*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f*2i)*3i)/(8*a^2*f^3) - (d^2*x^2*(d*1i - 2*c*f)*3i)/(8*a^2*f^2)) + exp(- e*4i - f*x*4i)*(((d^3*3i + 32*c^3*f^3 - c^2*d*f^2*24i - 12*c*d^2*f)*1i)/(512*a^2*f^4)
```

$$\begin{aligned} & + (d^3*x^3*1i)/(16*a^2*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)*3i)/(128*a^2*f^3) \\ & - (d^2*x^2*(d*1i - 4*c*f)*3i)/(64*a^2*f^2)) + (c^3*x)/(4*a^2) + (d^3*x^4)/(16*a^2) \\ & + (3*c^2*d*x^2)/(8*a^2) + (c*d^2*x^3)/(4*a^2) \end{aligned}$$

3.25 $\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx$

Optimal result	163
Rubi [A] (verified)	163
Mathematica [A] (verified)	165
Maple [A] (verified)	165
Fricas [A] (verification not implemented)	165
Sympy [A] (verification not implemented)	166
Maxima [F(-2)]	166
Giac [A] (verification not implemented)	167
Mupad [B] (verification not implemented)	167

Optimal result

Integrand size = 23, antiderivative size = 202

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx = & -\frac{id^2 e^{-2ie-2ifx}}{8a^2 f^3} - \frac{id^2 e^{-4ie-4ifx}}{128a^2 f^3} + \frac{de^{-2ie-2ifx}(c+dx)}{4a^2 f^2} \\ & + \frac{de^{-4ie-4ifx}(c+dx)}{32a^2 f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2 f} \\ & + \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2 f} + \frac{(c+dx)^3}{12a^2 d} \end{aligned}$$

[Out] $-1/8*I*d^2*exp(-2*I*e-2*I*f*x)/a^2/f^3-1/128*I*d^2*exp(-4*I*e-4*I*f*x)/a^2/f^3+1/4*d*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^2/f^2+1/32*d*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^2/f^2+1/4*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^2/f+1/16*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^2/f+1/12*(d*x+c)^3/a^2/d$

Rubi [A] (verified)

Time = 0.24 (sec), antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used = {3810, 2207, 2225}

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx = & \frac{d(c+dx)e^{-2ie-2ifx}}{4a^2 f^2} + \frac{d(c+dx)e^{-4ie-4ifx}}{32a^2 f^2} \\ & + \frac{i(c+dx)^2 e^{-2ie-2ifx}}{4a^2 f} + \frac{i(c+dx)^2 e^{-4ie-4ifx}}{16a^2 f} \\ & + \frac{(c+dx)^3}{12a^2 d} - \frac{id^2 e^{-2ie-2ifx}}{8a^2 f^3} - \frac{id^2 e^{-4ie-4ifx}}{128a^2 f^3} \end{aligned}$$

[In] $\text{Int}[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2, x]$

[Out] $\frac{((-1/8)*I)*d^2*E^((-2*I)*e - (2*I)*f*x))}{(a^2*f^3)} - \frac{((I/128)*d^2*E^((-4*I)*e - (4*I)*f*x))}{(a^2*f^3)} + \frac{(d*E^((-2*I)*e - (2*I)*f*x)*(c + d*x))}{(4*a^2*f^2)} + \frac{(d*E^((-4*I)*e - (4*I)*f*x)*(c + d*x))}{(32*a^2*f^2)} + \frac{((I/4)*E^((-2*I)*e - (2*I)*f*x)*(c + d*x)^2)}{(a^2*f)} + \frac{((I/16)*E^((-4*I)*e - (4*I)*f*x)*(c + d*x)^2)}{(a^2*f)} + \frac{(c + d*x)^3}{(12*a^2*d)}$

Rule 2207

```
Int[((b_)*(F_)^((g_)*(e_)+(f_)*(x_)))^(n_)*((c_)+(d_)*(x_))^m_, x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_)*(a_)+(b_)*(x_)))^n_, x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```
Int[((c_)+(d_)*(x_))^m*((a_)+(b_)*tan[(e_)+(f_)*(x_)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^2}{4a^2} + \frac{e^{-2ie-2ifx}(c+dx)^2}{2a^2} + \frac{e^{-4ie-4ifx}(c+dx)^2}{4a^2} \right) dx \\
&= \frac{(c+dx)^3}{12a^2d} + \frac{\int e^{-4ie-4ifx}(c+dx)^2 dx}{4a^2} + \frac{\int e^{-2ie-2ifx}(c+dx)^2 dx}{2a^2} \\
&= \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2f} + \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2f} + \frac{(c+dx)^3}{12a^2d} \\
&\quad - \frac{(id)\int e^{-4ie-4ifx}(c+dx) dx}{8a^2f} - \frac{(id)\int e^{-2ie-2ifx}(c+dx) dx}{2a^2f} \\
&= \frac{de^{-2ie-2ifx}(c+dx)}{4a^2f^2} + \frac{de^{-4ie-4ifx}(c+dx)}{32a^2f^2} + \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2f} \\
&\quad + \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2f} + \frac{(c+dx)^3}{12a^2d} - \frac{d^2 \int e^{-4ie-4ifx} dx}{32a^2f^2} - \frac{d^2 \int e^{-2ie-2ifx} dx}{4a^2f^2} \\
&= -\frac{id^2e^{-2ie-2ifx}}{8a^2f^3} - \frac{id^2e^{-4ie-4ifx}}{128a^2f^3} + \frac{de^{-2ie-2ifx}(c+dx)}{4a^2f^2} + \frac{de^{-4ie-4ifx}(c+dx)}{32a^2f^2} \\
&\quad + \frac{ie^{-2ie-2ifx}(c+dx)^2}{4a^2f} + \frac{ie^{-4ie-4ifx}(c+dx)^2}{16a^2f} + \frac{(c+dx)^3}{12a^2d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.61 (sec) , antiderivative size = 282, normalized size of antiderivative = 1.40

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ = \frac{\sec^2(e + fx)(\cos(fx) + i \sin(fx))^2 ((d + (1 + i)cf + (1 + i)dfx)((1 + i)cf + d(-i + (1 + i)fx)) \cos(2fx) \\ - (1 + i)cd \sin(2fx)))}{128 a^2 f^3}$$

[In] Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^2, x]

[Out] $\left(\sec^2(e + fx)\right)^2 \left(\cos^2(fx) + \sin^2(fx)\right)^2 \left((d + (1 + i)c f + (1 + i)d f x) \left((1 + i)c f + d(-i + (1 + i)f x)\right) \cos(2f x) - (1 + i)c d \sin(2f x)\right)/128 a^2 f^3$

Maple [A] (verified)

Time = 0.78 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.86

method	result
risch	$\frac{d^2 x^3}{12 a^2} + \frac{d c x^2}{4 a^2} + \frac{c^2 x}{4 a^2} + \frac{c^3}{12 a^2 d} + \frac{i (2 d^2 x^2 f^2 + 4 c d f^2 x - 2 i d^2 f x + 2 c^2 f^2 - 2 i c d f - d^2) e^{-2i(fx+e)}}{8 a^2 f^3} + \frac{i (8 d^2 x^2 f^2 + 16 c d f^2 x - 4 i d^2 f x - 16 c^2 f^2 - 8 c d f) e^{-2i(fx+e)}}{128 a^2 f^3}$

[In] int((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{12} a^2 d^2 x^3 + \frac{1}{4} a^2 d^2 c x^2 + \frac{1}{4} a^2 c^2 x + \frac{1}{12} a^2 d c^3 + \frac{1}{8} i (2 d^2 x^2 f^2 + 4 c d f^2 x - 2 i d^2 f x + 2 c^2 f^2 - 2 i c d f - d^2) e^{-2i(fx+e)} + \frac{1}{128} i (8 d^2 x^2 f^2 - 4 i d^2 f x + 16 c d f^2 x - 4 i c d f + 8 c^2 f^2 - d^2) e^{-2i(fx+e)}$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.81

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(24i d^2 f^2 x^2 + 24i c^2 f^2 + 12 c d f - 3i d^2 - 12 (-4i c d f^2 - d^2 f)x + 32 (d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x)e^{(4i f x + 2i c d f^2)})}{384 a^2 f^3}$$

[In] `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")`

[Out] $\frac{1}{384} \left(24 I d^2 f^2 x^2 + 24 I c^2 f^2 + 12 c d f - 3 I d^2 - 12 (-4 I c d^2 f^2 - d^2 f^2) x + 32 (d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x) e^{(4 I f x + 4 I e)} - 48 (-2 I d^2 f^2 x^2 - 2 I c^2 f^2 - 2 c d f + I d^2 + 2 (-2 I c d f^2 - d^2 f^2) x) e^{(2 I f x + 2 I e)} \right) e^{-(-4 I f x - 4 I e)} / (a^2 f^3)$

Sympy [A] (verification not implemented)

Time = 0.33 (sec), antiderivative size = 418, normalized size of antiderivative = 2.07

$$\begin{aligned} & \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx \\ &= \left\{ \frac{((64ia^2c^2f^5e^{2ie}+128ia^2cdf^5xe^{2ie}+32a^2cdf^4e^{2ie}+64ia^2d^2f^5x^2e^{2ie}+32a^2d^2f^4xe^{2ie}-8ia^2d^2f^3e^{2ie})e^{-4ifx}+(256ia^2c^2f^5e^{4ie}+512ia^2cdf^5xe^{4ie})e^{-4ifx})}{1024a^4f^6} \right. \\ & \quad \left. + \frac{x^3 \cdot (2d^2e^{2ie}+d^2)e^{-4ie}}{12a^2} + \frac{x^2 \cdot (2cde^{2ie}+cd)e^{-4ie}}{4a^2} + \frac{x(2c^2e^{2ie}+c^2)e^{-4ie}}{4a^2} \right. \\ & \quad \left. + \frac{c^2x}{4a^2} + \frac{cdx^2}{4a^2} + \frac{d^2x^3}{12a^2} \right) \end{aligned}$$

[In] `integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**2,x)`

[Out] Piecewise(((64*I*a**2*c**2*f**5*exp(2*I*e) + 128*I*a**2*c*d*f**5*x*exp(2*I*e) + 32*a**2*c*d*f**4*exp(2*I*e) + 64*I*a**2*d**2*f**5*x**2*exp(2*I*e) + 32*a**2*d**2*f**4*x*exp(2*I*e) - 8*I*a**2*d**2*f**3*exp(2*I*e))*exp(-4*I*f*x) + (256*I*a**2*c**2*f**5*exp(4*I*e) + 512*I*a**2*c*d*f**5*x*exp(4*I*e) + 256*a**2*c*d*f**4*exp(4*I*e) + 256*I*a**2*d**2*f**5*x**2*exp(4*I*e) + 256*a**2*d**2*f**4*x*exp(4*I*e) - 128*I*a**2*d**2*f**3*exp(4*I*e))*exp(-2*I*f*x))*exp(-6*I*e)/(1024*a**4*f**6), Ne(a**4*f**6*exp(6*I*e), 0)), (x**3*(2*d**2*exp(2*I*e) + d**2)*exp(-4*I*e)/(12*a**2) + x**2*(2*c*d*exp(2*I*e) + c*d)*exp(-4*I*e)/(4*a**2) + x*(2*c**2*exp(2*I*e) + c**2)*exp(-4*I*e)/(4*a**2), True) + c**2*x/(4*a**2) + c*d*x**2/(4*a**2) + d**2*x**3/(12*a**2)

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^2} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.45 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(32 d^2 f^3 x^3 e^{(4i f x + 4i e)} + 96 c d f^3 x^2 e^{(4i f x + 4i e)} + 96 c^2 f^3 x e^{(4i f x + 4i e)} + 96 i d^2 f^2 x^2 e^{(2i f x + 2i e)} + 24 i d^2 f^2 x^2 + 1)}{1}$$

```
[In] integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
[Out] 1/384*(32*d^2*f^3*x^3*e^(4*I*f*x + 4*I*e) + 96*c*d*f^3*x^2*e^(4*I*f*x + 4*I*e) + 96*c^2*f^3*x*e^(4*I*f*x + 4*I*e) + 96*I*d^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 24*I*d^2*f^2*x^2 + 192*I*c*d*f^2*x*x*e^(2*I*f*x + 2*I*e) + 48*I*c*d*f^2*x + 96*I*c^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 96*d^2*f*x*x*e^(2*I*f*x + 2*I*e) + 24*I*c^2*f^2*x^2 + 12*d^2*f*x + 96*c*d*f*x*e^(2*I*f*x + 2*I*e) + 12*c*d*f - 48*I*d^2*x^2*e^(2*I*f*x + 2*I*e) - 3*I*d^2)*e^(-4*I*f*x - 4*I*e)/(a^2*f^3)
```

Mupad [B] (verification not implemented)

Time = 3.65 (sec) , antiderivative size = 183, normalized size of antiderivative = 0.91

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^2} dx = \frac{c^2 x}{4 a^2} - e^{-e^{4i-fx} 4i} \left(\frac{(-8 c^2 f^2 + c d f 4i + d^2) 1i}{128 a^2 f^3} - \frac{d^2 x^2 1i}{16 a^2 f} \right. \\ \left. + \frac{d x (-4 c f + d 1i) 1i}{32 a^2 f^2} \right) - e^{-e^{2i-fx} 2i} \left(\frac{(-2 c^2 f^2 + c d f 2i + d^2) 1i}{8 a^2 f^3} - \frac{d^2 x^2 1i}{4 a^2 f} \right. \\ \left. + \frac{d x (-2 c f + d 1i) 1i}{4 a^2 f^2} \right) + \frac{d^2 x^3}{12 a^2} + \frac{c d x^2}{4 a^2}$$

```
[In] int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^2,x)
[Out] (c^2*x)/(4*a^2) - exp(- e*4i - f*x*4i)*((d^2 - 8*c^2*f^2 + c*d*f*4i)*1i)/(128*a^2*f^3) - (d^2*x^2*1i)/(16*a^2*f) + (d*x*(d*1i - 4*c*f)*1i)/(32*a^2*f^2) - exp(- e*2i - f*x*2i)*((d^2 - 2*c^2*f^2 + c*d*f*2i)*1i)/(8*a^2*f^3) - (d^2*x^2*1i)/(4*a^2*f) + (d*x*(d*1i - 2*c*f)*1i)/(4*a^2*f^2) + (d^2*x^3)/(12*a^2) + (c*d*x^2)/(4*a^2)
```

3.26 $\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx$

Optimal result	168
Rubi [A] (verified)	168
Mathematica [A] (verified)	170
Maple [A] (verified)	170
Fricas [A] (verification not implemented)	170
Sympy [A] (verification not implemented)	171
Maxima [F(-2)]	171
Giac [A] (verification not implemented)	172
Mupad [B] (verification not implemented)	172

Optimal result

Integrand size = 21, antiderivative size = 151

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx = & -\frac{3idx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} \\ & + \frac{d}{16f^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} \\ & + \frac{3d}{16f^2(a^2+ia^2\tan(e+fx))} + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} \end{aligned}$$

[Out]
$$\begin{aligned} -3/16*I*d*x/a^2/f-1/8*d*x^2/a^2+1/4*x*(d*x+c)/a^2+1/16*d/f^2/(a+I*a*tan(f*x+e))^2+1/4*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^2+3/16*d/f^2/(a^2+I*a^2*tan(f*x+e)) \\ +1/4*I*(d*x+c)/f/(a^2+I*a^2*tan(f*x+e)) \end{aligned}$$

Rubi [A] (verified)

Time = 0.18 (sec), antiderivative size = 151, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.143, Rules used = {3560, 8, 3811}

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx = & \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} + \frac{x(c+dx)}{4a^2} \\ & + \frac{3d}{16f^2(a^2+ia^2\tan(e+fx))} - \frac{3idx}{16a^2f} - \frac{dx^2}{8a^2} \\ & + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} + \frac{d}{16f^2(a+ia\tan(e+fx))^2} \end{aligned}$$

[In] Int[(c + d*x)/(a + I*a*Tan[e + f*x])^2, x]

```
[Out] (((-3*I)/16)*d*x)/(a^2*f) - (d*x^2)/(8*a^2) + (x*(c + d*x))/(4*a^2) + d/(16*f^2*(a + I*a*Tan[e + f*x])^2) + ((I/4)*(c + d*x))/(f*(a + I*a*Tan[e + f*x])^2) + (3*d)/(16*f^2*(a^2 + I*a^2*Tan[e + f*x])) + ((I/4)*(c + d*x))/(f*(a^2 + I*a^2*Tan[e + f*x]))
```

Rule 8

```
Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]
```

Rule 3560

```
Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n)/(2*b*d*n), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3811

```
Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Dist[(c + d*x)^m, u, x] - Dist[d*m, Int[Dist[(c + d*x)^(m - 1), u, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{x(c+dx)}{4a^2} + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} \\
&\quad - d \int \left(\frac{x}{4a^2} + \frac{i}{4f(a+ia\tan(e+fx))^2} + \frac{i}{4f(a^2+ia^2\tan(e+fx))} \right) dx \\
&= -\frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} \\
&\quad - \frac{(id) \int \frac{1}{(a+ia\tan(e+fx))^2} dx}{4f} - \frac{(id) \int \frac{1}{a^2+ia^2\tan(e+fx)} dx}{4f} \\
&= -\frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} + \frac{d}{16f^2(a+ia\tan(e+fx))^2} \\
&\quad + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} + \frac{d}{8f^2(a^2+ia^2\tan(e+fx))} \\
&\quad + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} - \frac{(id) \int 1 dx}{8a^2f} - \frac{(id) \int \frac{1}{a+ia\tan(e+fx)} dx}{8af} \\
&= -\frac{idx}{8a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} + \frac{d}{16f^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} \\
&\quad + \frac{3d}{16f^2(a^2+ia^2\tan(e+fx))} + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))} - \frac{(id) \int 1 dx}{16a^2f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3idx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} + \frac{d}{16f^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{4f(a+ia\tan(e+fx))^2} \\
&\quad + \frac{3d}{16f^2(a^2+ia^2\tan(e+fx))} + \frac{i(c+dx)}{4f(a^2+ia^2\tan(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.86

$$\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx = \frac{-\sec^2(e+fx)(8(d+2icf+2idfx)+(4cf(i+4fx)+d(1+4ifx+8f^2x^2))\cos(2(e+fx))+(4cf(1+4ifx+8f^2x^2))\sin(2(e+fx))}{64a^2f^2(-i+\tan(e+fx))^2}$$

[In] `Integrate[(c + d*x)/(a + I*a*Tan[e + f*x])^2, x]`

[Out]
$$\begin{aligned}
&-1/64*(\text{Sec}[e+f*x]^2*(8*(d+(2*I)*c*f+(2*I)*d*f*x)+(4*c*f*(I+4*f*x) \\
&+d*(1+(4*I)*f*x+8*f^2*x^2))*\text{Cos}[2*(e+f*x)]+(4*c*f*(1+(4*I)*f*x) \\
&+d*(-I+4*f*x+(8*I)*f^2*x^2))*\text{Sin}[2*(e+f*x)]))/((a^2*f^2*(-I+\text{Tan}[e \\
&+f*x]))^2)
\end{aligned}$$

Maple [A] (verified)

Time = 0.80 (sec) , antiderivative size = 82, normalized size of antiderivative = 0.54

method	result	size
risch	$\frac{dx^2}{8a^2} + \frac{cx}{4a^2} + \frac{i(2dfx+2cf-id)e^{-2i(fx+e)}}{8a^2f^2} + \frac{i(4dfx+4cf-id)e^{-4i(fx+e)}}{64a^2f^2}$	82

[In] `int((d*x+c)/(a+I*a*tan(f*x+e))^2, x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned}
&1/8*d*x^2/a^2+1/4/a^2*c*x+1/8*I*(2*d*f*x-I*d+2*c*f)/a^2/f^2*\exp(-2*I*(f*x+e)) \\
&+1/64*I*(4*d*f*x-I*d+4*c*f)/a^2/f^2*\exp(-4*I*(f*x+e))
\end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 0.52

$$\begin{aligned}
&\int \frac{c+dx}{(a+ia\tan(e+fx))^2} dx \\
&= \frac{(4i dfx + 4i cf + 8(df^2x^2 + 2cf^2x)e^{(4ifx+4ie)} - 8(-2i dfx - 2i cf - d)e^{(2ifx+2ie)} + d)e^{(-4ifx-4ie)}}{64a^2f^2}
\end{aligned}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e))^2, x, algorithm="fricas")`

[Out] $\frac{1}{64} \cdot (4 \cdot I \cdot d \cdot f \cdot x + 4 \cdot I \cdot c \cdot f + 8 \cdot (d \cdot f^2 \cdot x^2 + 2 \cdot c \cdot f^2 \cdot x) \cdot e^{(4 \cdot I \cdot f \cdot x + 4 \cdot I \cdot e)} - 8 \cdot (-2 \cdot I \cdot d \cdot f \cdot x - 2 \cdot I \cdot c \cdot f - d) \cdot e^{(2 \cdot I \cdot f \cdot x + 2 \cdot I \cdot e)} + d) \cdot e^{(-4 \cdot I \cdot f \cdot x - 4 \cdot I \cdot e)}) / (a^2 \cdot f^2)$

Sympy [A] (verification not implemented)

Time = 0.26 (sec), antiderivative size = 226, normalized size of antiderivative = 1.50

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx$$

$$= \begin{cases} \frac{((32ia^2cf^3e^{2ie} + 32ia^2df^3xe^{2ie} + 8a^2df^2e^{2ie})e^{-4ifx} + (128ia^2cf^3e^{4ie} + 128ia^2df^3xe^{4ie} + 64a^2df^2e^{4ie})e^{-2ifx})e^{-6ie}}{512a^4f^4} & \text{for } a^4f^4e^{6ie} \neq 0 \\ \frac{x^2 \cdot (2de^{2ie} + d)e^{-4ie}}{8a^2} + \frac{x(2ce^{2ie} + c)e^{-4ie}}{4a^2} & \text{otherwise} \\ + \frac{cx}{4a^2} + \frac{dx^2}{8a^2} \end{cases}$$

[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))**2,x)

[Out] Piecewise(((32*I*a**2*c*f**3*exp(2*I*e) + 32*I*a**2*d*f**3*x*exp(2*I*e) + 8*a**2*d*f**2*exp(2*I*e))*exp(-4*I*f*x) + (128*I*a**2*c*f**3*exp(4*I*e) + 128*I*a**2*d*f**3*x*exp(4*I*e) + 64*a**2*d*f**2*exp(4*I*e))*exp(-2*I*f*x))*e^(-6*I*e) / (512*a**4*f**4), Ne(a**4*f**4*exp(6*I*e), 0)), (x**2*(2*d*exp(2*I*e) + d)*exp(-4*I*e) / (8*a**2) + x*(2*c*exp(2*I*e) + c)*exp(-4*I*e) / (4*a**2), True)) + c*x / (4*a**2) + d*x**2 / (8*a**2)

Maxima [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))**2,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.42 (sec) , antiderivative size = 101, normalized size of antiderivative = 0.67

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx \\ = \frac{(8df^2x^2e^{(4ifx+4ie)} + 16cf^2xe^{(4ifx+4ie)} + 16idfxe^{(2ifx+2ie)} + 4idfx + 16icfe^{(2ifx+2ie)} + 4icf + 8de^{(2ifx+2ie)})}{64a^2f^2}$$

[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")

[Out] $\frac{1}{64} (8d^2f^2x^2e^{(4Ifx+4Ie)} + 16cf^2x^2e^{(4Ifx+4Ie)} + 16Idfxe^{(2Ifx+2Ie)} + 4Idfx + 16icfe^{(2Ifx+2Ie)} + 4icf + 8d^2e^{(2Ifx+2Ie)}) / (a^2f^2)$

Mupad [B] (verification not implemented)

Time = 3.13 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.68

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^2} dx = \frac{dx^2}{8a^2} - e^{-e^{4i-fx}4i} \left(\frac{(-4cf + d1i)1i}{64a^2f^2} - \frac{dx1i}{16a^2f} \right) \\ - e^{-e^{2i-fx}2i} \left(\frac{(-2cf + d1i)1i}{8a^2f^2} - \frac{dx1i}{4a^2f} \right) + \frac{cx}{4a^2}$$

[In] int((c + d*x)/(a + a*tan(e + f*x)*1i)^2,x)

[Out] $(d*x^2)/(8*a^2) - \exp(-e*4i - f*x*4i)*(((d*1i - 4*c*f)*1i)/(64*a^2*f^2) - (d*x*1i)/(16*a^2*f)) - \exp(-e*2i - f*x*2i)*(((d*1i - 2*c*f)*1i)/(8*a^2*f^2) - (d*x*1i)/(4*a^2*f)) + (c*x)/(4*a^2)$

3.27 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx$

Optimal result	173
Rubi [A] (verified)	174
Mathematica [A] (verified)	176
Maple [A] (verified)	177
Fricas [A] (verification not implemented)	177
Sympy [F]	178
Maxima [A] (verification not implemented)	178
Giac [A] (verification not implemented)	178
Mupad [F(-1)]	179

Optimal result

Integrand size = 23, antiderivative size = 305

$$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^2} dx = \frac{\cos\left(2e - \frac{2cf}{d}\right) \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\ + \frac{\cos\left(4e - \frac{4cf}{d}\right) \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} \\ + \frac{\log(c+dx)}{4a^2d} \\ - \frac{i \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right) \sin\left(4e - \frac{4cf}{d}\right)}{4a^2d} \\ - \frac{i \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{2a^2d} \\ - \frac{i \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\ - \frac{\sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\ - \frac{i \cos\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} \\ - \frac{\sin\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d}$$

```
[Out] 1/4*Ci(4*c*f/d+4*f*x)*cos(-4*e+4*c*f/d)/a^2/d+1/2*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a^2/d+1/4*ln(d*x+c)/a^2/d-1/2*I*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^2/d-1/4*I*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^2/d+1/4*I*Ci(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d+1/4*Si(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d+1/2*I*Ci(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d+1/2*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d
```

Rubi [A] (verified)

Time = 0.94 (sec) , antiderivative size = 305, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.217$, Rules used = {3809, 3384, 3380, 3383, 3393}

$$\begin{aligned} \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx = & -\frac{i \operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \sin\left(2e - \frac{2cf}{d}\right)}{2a^2 d} \\ & - \frac{i \operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \sin\left(4e - \frac{4cf}{d}\right)}{4a^2 d} \\ & + \frac{\operatorname{CosIntegral}\left(2xf + \frac{2cf}{d}\right) \cos\left(2e - \frac{2cf}{d}\right)}{2a^2 d} \\ & + \frac{\operatorname{CosIntegral}\left(4xf + \frac{4cf}{d}\right) \cos\left(4e - \frac{4cf}{d}\right)}{4a^2 d} \\ & - \frac{\sin\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2 d} \\ & - \frac{\sin\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2 d} \\ & - \frac{i \cos\left(2e - \frac{2cf}{d}\right) \operatorname{Si}\left(2xf + \frac{2cf}{d}\right)}{2a^2 d} \\ & - \frac{i \cos\left(4e - \frac{4cf}{d}\right) \operatorname{Si}\left(4xf + \frac{4cf}{d}\right)}{4a^2 d} + \frac{\log(c + dx)}{4a^2 d} \end{aligned}$$

[In] $\operatorname{Int}[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2), x]$

[Out] $(\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/(2*a^2*d) + (\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x])/(4*a^2*d) + \operatorname{Log}[c + d*x]/(4*a^2*d) - ((I/4)*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x]*\operatorname{Sin}[4*e - (4*c*f)/d])/(a^2*d) - ((I/2)*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sin}[2*e - (2*c*f)/d])/(a^2*d) - ((I/2)*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(a^2*d) - (\operatorname{Sin}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/(2*a^2*d) - ((I/4)*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(a^2*d) - (\operatorname{Sin}[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/(4*a^2*d)$

Rule 3380

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

Rule 3383

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> In[t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)} + \frac{\cos(2e+2fx)}{2a^2(c+dx)} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)} - \frac{i \sin(2e+2fx)}{2a^2(c+dx)} \right. \\
&\quad \left. - \frac{\sin^2(2e+2fx)}{4a^2(c+dx)} - \frac{i \sin(4e+4fx)}{4a^2(c+dx)} \right) dx \\
&= \frac{\log(c+dx)}{4a^2d} - \frac{i \int \frac{\sin(4e+4fx)}{c+dx} dx}{4a^2} - \frac{i \int \frac{\sin(2e+2fx)}{c+dx} dx}{2a^2} \\
&\quad + \frac{\int \frac{\cos^2(2e+2fx)}{c+dx} dx}{4a^2} - \frac{\int \frac{\sin^2(2e+2fx)}{c+dx} dx}{4a^2} + \frac{\int \frac{\cos(2e+2fx)}{c+dx} dx}{2a^2} \\
&= \frac{\log(c+dx)}{4a^2d} - \frac{\int \left(\frac{1}{2(c+dx)} - \frac{\cos(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} + \frac{\int \left(\frac{1}{2(c+dx)} + \frac{\cos(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} \\
&\quad - \frac{(i \cos(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c+dx} dx}{4a^2} - \frac{(i \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} \\
&\quad + \frac{\cos(2e - \frac{2cf}{d}) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} - \frac{(i \sin(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c+dx} dx}{4a^2} \\
&\quad - \frac{(i \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} - \frac{\sin(2e - \frac{2cf}{d}) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2a^2d} - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} + 2 \int \frac{\cos(4e + 4fx)}{c + dx} dx \\
&= \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2a^2d} - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} \\
&\quad + 2 \left(\frac{\cos(4e - \frac{4cf}{d}) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c + dx} dx}{8a^2} - \frac{\sin(4e - \frac{4cf}{d}) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c + dx} dx}{8a^2} \right) \\
&= \frac{\cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{4a^2d} \\
&\quad - \frac{i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{2a^2d} - \frac{i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{4a^2d} \\
&\quad + 2 \left(\frac{\cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{8a^2d} - \frac{\sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^2d} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.85 (sec), antiderivative size = 211, normalized size of antiderivative = 0.69

$$\begin{aligned}
&\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\
&= (\cos(2e - \frac{2cf}{d}) - i \sin(2e - \frac{2cf}{d})) \left(2 \operatorname{CosIntegral}\left(\frac{2f(c+dx)}{d}\right) + \cos(2e - \frac{2cf}{d}) \log(f(c + dx)) + \operatorname{CosIntegral}\left(\frac{4f(c+dx)}{d}\right) \right)
\end{aligned}$$

[In] Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^2), x]

```
[Out] ((Cos[2*e - (2*c*f)/d] - I*Sin[2*e - (2*c*f)/d])*(2*CosIntegral[(2*f*(c + d*x))/d] + Cos[2*e - (2*c*f)/d]*Log[f*(c + d*x)] + CosIntegral[(4*f*(c + d*x))/d]*(Cos[2*e - (2*c*f)/d] - I*Sin[2*e - (2*c*f)/d]) + I*Log[f*(c + d*x)]*Sin[2*e - (2*c*f)/d] - (2*I)*SinIntegral[(2*f*(c + d*x))/d] - I*Cos[2*e - (2*c*f)/d]*SinIntegral[(4*f*(c + d*x))/d] - Sin[2*e - (2*c*f)/d]*SinIntegral[(4*f*(c + d*x))/d]))/(4*a^2*d)
```

Maple [A] (verified)

Time = 0.87 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.37

method	result	size
risch	$\frac{\ln(dx+c)}{4a^2d} - \frac{e^{\frac{4i(cf-de)}{d}} \text{Ei}_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{4a^2d} - \frac{e^{\frac{2i(cf-de)}{d}} \text{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{2a^2d}$	114

```
[In] int(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/4*ln(d*x+c)/a^2/d-1/4/a^2/d*exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-1/2/a^2/d*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 0.28

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx \\ = \frac{2 \text{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + \text{Ei}\left(-\frac{4(ifx + cf)}{d}\right) e^{\left(-\frac{4(de - cf)}{d}\right)} + \log\left(\frac{dx+c}{d}\right)}{4 a^2 d}$$

```
[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] 1/4*(2*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + Ei(-4*(I*d*f*x + I*c*f)/d)*e^(-4*(I*d*e - I*c*f)/d) + log((d*x + c)/d))/(a^2*d)
```

Sympy [F]

$$\int \frac{1}{(c+dx)(a+i a \tan(e+fx))^2} dx \\ = -\frac{\int \frac{1}{c \tan^2(e+fx)-2 i c \tan(e+fx)-c+dx \tan^2(e+fx)-2 i d x \tan(e+fx)-d x} dx}{a^2}$$

[In] `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**2,x)`

[Out] `-Integral(1/(c*tan(e + f*x)**2 - 2*I*c*tan(e + f*x) - c + d*x*tan(e + f*x)*
*2 - 2*I*d*x*tan(e + f*x) - d*x), x)/a**2`

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 194, normalized size of antiderivative = 0.64

$$\int \frac{1}{(c+dx)(a+i a \tan(e+fx))^2} dx = \\ -\frac{2 f \cos\left(-\frac{2 (d e-c f)}{d}\right) E_1\left(-\frac{2 (-i (f x+e) d+i d e-i c f)}{d}\right)+f \cos\left(-\frac{4 (d e-c f)}{d}\right) E_1\left(-\frac{4 (-i (f x+e) d+i d e-i c f)}{d}\right)+2 i f E_1\left(-\frac{4 (-i (f x+e) d+i d e-i c f)}{d}\right)}{(d e-c f)^2}$$

[In] `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**2,x, algorithm="maxima")`

[Out] `-1/4*(2*f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f*cos(-4*(d*e - c*f)/d)*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + I*f*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f))/(a^2*d*f)`

Giac [A] (verification not implemented)

none

Time = 0.42 (sec) , antiderivative size = 404, normalized size of antiderivative = 1.32

$$\int \frac{1}{(c+dx)(a+i a \tan(e+fx))^2} dx \\ = \frac{2 \cos(2 e) \cos\left(\frac{2 c f}{d}\right) \text{Ci}\left(-\frac{2 (d f x+c f)}{d}\right)+\cos(2 e)^2 \log(d x+c)+2 i \cos\left(\frac{2 c f}{d}\right) \text{Ci}\left(-\frac{2 (d f x+c f)}{d}\right) \sin(2 e)+2 i c \sin(2 e) \text{Si}\left(-\frac{2 (d f x+c f)}{d}\right)}{(d e-c f)^2}$$

[In] `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**2,x, algorithm="giac")`

```
[Out] 1/4*(2*cos(2*e)*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + cos(2*e)^2*log(d*x + c) + 2*I*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e) + 2*I*cos(2*e)*log(d*x + c)*sin(2*e) - log(d*x + c)*sin(2*e)^2 + 2*I*cos(2*e)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 2*I*cos(2*e)*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 2*cos(2*c*f/d)*sin(2*e)*sin_integral(2*(d*f*x + c*f)/d) + 2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 2*I*sin(2*e)*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + cos(4*c*f/d)*cos_integral(-4*(d*f*x + c*f)/d) + I*cos_integral(-4*(d*f*x + c*f)/d)*sin(4*c*f/d) - I*cos(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) + sin(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d))/(a^2*d*cos(2*e)^2 + 2*I*a^2*d*cos(2*e)*sin(2*e) - a^2*d*sin(2*e)^2)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^2} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^2 (c + dx)} dx$$

```
[In] int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)),x)
```

```
[Out] int(1/((a + a*tan(e + f*x)*1i)^2*(c + d*x)), x)
```

3.28 $\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx$

Optimal result	180
Rubi [A] (verified)	181
Mathematica [A] (verified)	184
Maple [A] (verified)	185
Fricas [A] (verification not implemented)	185
Sympy [F]	186
Maxima [A] (verification not implemented)	186
Giac [B] (verification not implemented)	186
Mupad [F(-1)]	188

Optimal result

Integrand size = 23, antiderivative size = 436

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx = & -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} \\ & - \frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & - \frac{if \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\ & - \frac{f \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{a^2d^2} \\ & - \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{a^2d^2} \\ & + \frac{i \sin(2e + 2fx)}{2a^2d(c+dx)} + \frac{\sin^2(2e + 2fx)}{4a^2d(c+dx)} \\ & + \frac{i \sin(4e + 4fx)}{4a^2d(c+dx)} - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & + \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & - \frac{f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\ & + \frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2} \end{aligned}$$

```
[Out] -1/4/a^2/d/(d*x+c)-I*f*Ci(4*c*f/d+4*f*x)*cos(-4*e+4*c*f/d)/a^2/d^2-I*f*Ci(2*c*f/d+2*f*x)*cos(-2*e+2*c*f/d)/a^2/d^2-1/2*cos(2*f*x+2*e)/a^2/d/(d*x+c)-1/4*cos(2*f*x+2*e)^2/a^2/d/(d*x+c)-f*cos(-2*e+2*c*f/d)*Si(2*c*f/d+2*f*x)/a^2/d^2-f*cos(-4*e+4*c*f/d)*Si(4*c*f/d+4*f*x)/a^2/d^2+f*Ci(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d^2-I*f*Si(4*c*f/d+4*f*x)*sin(-4*e+4*c*f/d)/a^2/d^2+f*Ci(2*
```

$$*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d^2-I*f*Si(2*c*f/d+2*f*x)*sin(-2*e+2*c*f/d)/a^2/d^2+1/2*I*sin(2*f*x+2*e)/a^2/d/(d*x+c)+1/4*sin(2*f*x+2*e)^2/a^2/d/(d*x+c)+1/4*I*sin(4*f*x+4*e)/a^2/d/(d*x+c)$$

Rubi [A] (verified)

Time = 0.83 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.304, Rules used = {3809, 3378, 3384, 3380, 3383, 3394, 12}

$$\begin{aligned} \int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx = & -\frac{f \operatorname{CosIntegral}(4xf + \frac{4cf}{d}) \sin(4e - \frac{4cf}{d})}{a^2 d^2} \\ & -\frac{f \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \sin(2e - \frac{2cf}{d})}{a^2 d^2} \\ & -\frac{if \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \cos(2e - \frac{2cf}{d})}{a^2 d^2} \\ & -\frac{if \operatorname{CosIntegral}(4xf + \frac{4cf}{d}) \cos(4e - \frac{4cf}{d})}{a^2 d^2} \\ & +\frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{a^2 d^2} \\ & +\frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(4xf + \frac{4cf}{d})}{a^2 d^2} \\ & -\frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{a^2 d^2} \\ & -\frac{f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(4xf + \frac{4cf}{d})}{a^2 d^2} + \frac{\sin^2(2e + 2fx)}{4a^2 d(c + dx)} \\ & +\frac{i \sin(2e + 2fx)}{2a^2 d(c + dx)} + \frac{i \sin(4e + 4fx)}{4a^2 d(c + dx)} \\ & -\frac{\cos^2(2e + 2fx)}{4a^2 d(c + dx)} - \frac{\cos(2e + 2fx)}{2a^2 d(c + dx)} - \frac{1}{4a^2 d(c + dx)} \end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2), x]

[Out]
$$\begin{aligned} & -1/4*1/(a^2*d*(c + d*x)) - \operatorname{Cos}[2*e + 2*f*x]/(2*a^2*d*(c + d*x)) - \operatorname{Cos}[2*e + 2*f*x]^2/(4*a^2*d*(c + d*x)) - (I*f*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/({a^2*d^2}) - (I*f*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x])/({a^2*d^2}) - (f*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x]*\operatorname{Sin}[4*e - (4*c*f)/d])/({a^2*d^2}) - (f*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sin}[2*e - (2*c*f)/d])/({a^2*d^2}) + ((I/2)*\operatorname{Sin}[2*e + 2*f*x])/({a^2*d*(c + d*x)}) + \operatorname{Sin}[2*e + 2*f*x]^2/(4*a^2*d*(c + d*x)) + ((I/4)*\operatorname{Sin}[4*e + 4*f*x])/({a^2*d*(c + d*x)}) - (f*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/({a^2*d^2}) + (I*f*\operatorname{Sin}[2*e - (2*c*f)/d]*\operatorname{SinIntegral}[(2*c*f)/d + 2*f*x])/({a^2*d^2}) - (f*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/({a^2*d^2}) + (I*f*\operatorname{Sin}[4*e - (4*c*f)/d]*\operatorname{SinIntegral}[(4*c*f)/d + 4*f*x])/({a^2*d^2}) \end{aligned}$$

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 3378

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[((c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]]
```

Rule 3380

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

Rule 3383

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3394

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1))), Int[ExpandTrigReduce[((c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n - 1), x], x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] && LtQ[m, -1]]
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)^2} + \frac{\cos(2e+2fx)}{2a^2(c+dx)^2} + \frac{\cos^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{i \sin(2e+2fx)}{2a^2(c+dx)^2} \right. \\
&\quad \left. - \frac{\sin^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{i \sin(4e+4fx)}{4a^2(c+dx)^2} \right) dx \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{i \int \frac{\sin(4e+4fx)}{(c+dx)^2} dx}{4a^2} - \frac{i \int \frac{\sin(2e+2fx)}{(c+dx)^2} dx}{2a^2} \\
&\quad + \frac{\int \frac{\cos^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} - \frac{\int \frac{\sin^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} + \frac{\int \frac{\cos(2e+2fx)}{(c+dx)^2} dx}{2a^2} \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} + \frac{i \sin(2e+2fx)}{2a^2d(c+dx)} \\
&\quad + \frac{\sin^2(2e+2fx)}{4a^2d(c+dx)} + \frac{i \sin(4e+4fx)}{4a^2d(c+dx)} - \frac{(if) \int \frac{\cos(2e+2fx)}{c+dx} dx}{a^2d} - \frac{(if) \int \frac{\cos(4e+4fx)}{c+dx} dx}{a^2d} \\
&\quad - \frac{f \int \frac{\sin(2e+2fx)}{c+dx} dx}{a^2d} + \frac{f \int -\frac{\sin(4e+4fx)}{2(c+dx)} dx}{a^2d} - \frac{f \int \frac{\sin(4e+4fx)}{2(c+dx)} dx}{a^2d} \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} + \frac{i \sin(2e+2fx)}{2a^2d(c+dx)} \\
&\quad + \frac{\sin^2(2e+2fx)}{4a^2d(c+dx)} + \frac{i \sin(4e+4fx)}{4a^2d(c+dx)} - 2 \frac{f \int \frac{\sin(4e+4fx)}{c+dx} dx}{2a^2d} \\
&\quad - \frac{(if \cos(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} - \frac{(if \cos(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} \\
&\quad - \frac{(f \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} + \frac{(if \sin(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} \\
&\quad + \frac{(if \sin(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} - \frac{(f \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad - \frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad - \frac{if \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{a^2d^2} + \frac{i \sin(2e + 2fx)}{2a^2d(c+dx)} \\
&\quad + \frac{\sin^2(2e + 2fx)}{4a^2d(c+dx)} + \frac{i \sin(4e + 4fx)}{4a^2d(c+dx)} - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad + \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} + \frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - 2 \left(\frac{(f \cos(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^2d} + \frac{(f \sin(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^2d} \right) \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cos(2e+2fx)}{2a^2d(c+dx)} - \frac{\cos^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad - \frac{if \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad - \frac{if \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - \frac{f \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{a^2d^2} + \frac{i \sin(2e + 2fx)}{2a^2d(c+dx)} \\
&\quad + \frac{\sin^2(2e + 2fx)}{4a^2d(c+dx)} + \frac{i \sin(4e + 4fx)}{4a^2d(c+dx)} - \frac{f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad + \frac{if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{a^2d^2} + \frac{if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - 2 \left(\frac{f \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{2a^2d^2} + \frac{f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{2a^2d^2} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.67 (sec), antiderivative size = 467, normalized size of antiderivative = 1.07

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx =$$

$$\frac{(\cos(2(e+f(-\frac{c}{d}+x))) - i \sin(2(e+f(-\frac{c}{d}+x)))) \left(2d \cos(\frac{2cf}{d}) + d \cos(2(e+f(-\frac{c}{d}+x))) + d \cos(2(e+f(-\frac{c}{d}+x))) \sin(\frac{2cf}{d}) - 2 \sin(2(e+f(-\frac{c}{d}+x))) \sin(\frac{2cf}{d})\right)}{a^2 d^2 (c+dx)^2 (a+ia\tan(e+fx))^2}$$

[In] Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^2), x]

```
[Out] -1/4*((Cos[2*(e + f*(-(c/d) + x))] - I*Sin[2*(e + f*(-(c/d) + x))])*(2*d*Cos[(2*c*f)/d] + d*Cos[2*(e + f*(-(c/d) + x))] + d*Cos[2*(e + f*(c/d + x))]) - (2*I)*d*Sin[(2*c*f)/d] + (4*I)*f*(c + d*x)*CosIntegral[(2*f*(c + d*x))/d]*(Cos[2*f*x] + I*Sin[2*f*x]) + I*d*Sin[2*(e + f*(-(c/d) + x))] - I*d*Sin[2*(e + f*(c/d + x))] + 4*f*(c + d*x)*CosIntegral[(4*f*(c + d*x))/d]*(I*Cos[2*e - (2*f*(c + d*x))/d] + Sin[2*e - (2*f*(c + d*x))/d]) + 4*c*f*Cos[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 4*d*f*x*Cos[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (4*I)*c*f*Sin[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (4*I)*d*f*x*Sin[2*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 4*c*f*Cos[2*e - (2*f*(c + d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] + 4*d*f*x*Cos[2*e - (2*f*(c + d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] - (4*I)*c*f*Sin[2*e - (2*f*(c + d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] - (4*I)*d*f*x*Sin[2*e - (2*f*(c + d*x))/d]*SinIntegral[(4*f*(c + d*x))/d]))/(a^2*d^2*(c + d*x))
```

Maple [A] (verified)

Time = 0.89 (sec), antiderivative size = 175, normalized size of antiderivative = 0.40

method	result
risch	$-\frac{1}{4a^2d(dx+c)} - \frac{fe^{-4i(fx+e)}}{4a^2(df+cf)d} + \frac{ife^{\frac{4i(cf-de)}{d}}\text{Ei}_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{a^2d^2} - \frac{fe^{-2i(fx+e)}}{2a^2(df+cf)d} + \frac{ife^{\frac{2i(cf-de)}{d}}\text{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{a^2d^2}$

```
[In] int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
[Out] -1/4/a^2/d/(d*x+c)-1/4/a^2*f*exp(-4*I*(f*x+e))/(d*f*x+c*f)/d+I/a^2*f/d^2*exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-1/2/a^2*f*exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+I/a^2*f/d^2*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec), antiderivative size = 142, normalized size of antiderivative = 0.33

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx = \\ - \frac{\left(\left(4(i dfx + i cf) \text{Ei} \left(-\frac{2(i dfx + i cf)}{d} \right) e^{\left(-\frac{2(i de - i cf)}{d} \right)} + 4(i dfx + i cf) \text{Ei} \left(-\frac{4(i dfx + i cf)}{d} \right) e^{\left(-\frac{4(i de - i cf)}{d} \right)} + d \right) \right)}{4(a^2 d^3 x + a^2 c d^2)}$$

```
[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="fricas")
[Out] -1/4*((4*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^(-2*(I*d*e - I*c*f)/d) + 4*(I*d*f*x + I*c*f)*Ei(-4*(I*d*f*x + I*c*f)/d)*e^(-4*(I*d*e - I*c*f)/d) + d)*e^(4*I*f*x + 4*I*e) + 2*d*e^(2*I*f*x + 2*I*e) + d)*e^(-4*I*f*x - 4*I*e)/(a^2*d^3*x + a^2*c*d^2)
```

Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx = -\frac{\int \frac{1}{c^2\tan^2(e+fx)-2ic^2\tan(e+fx)-c^2+2cdx\tan^2(e+fx)-4icdx\tan(e+fx)-2cdx+d^2x^2\tan^2(e+fx)-2id^2x^2\tan(e+fx)-d^2x^2} dx}{a^2}$$

[In] `integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**2,x)`

[Out] `-Integral(1/(c**2*tan(e + f*x)**2 - 2*I*c**2*tan(e + f*x) - c**2 + 2*c*d*x*tan(e + f*x)**2 - 4*I*c*d*x*tan(e + f*x) - 2*c*d*x + d**2*x**2*tan(e + f*x)**2 - 2*I*d**2*x**2*tan(e + f*x) - d**2*x**2), x)/a**2`

Maxima [A] (verification not implemented)

none

Time = 0.48 (sec) , antiderivative size = 211, normalized size of antiderivative = 0.48

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx = -\frac{2f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) + 2if^2}{4((fx+e)a^2d^2-a^2)}$$

[In] `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

[Out] `-1/4*(2*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 2*I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + I*f^2*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + f^2)/(((f*x + e)*a^2*d^2 - a^2*d^2 - a^2*d^2*e + a^2*c*d*f)*f)`

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1967 vs. $2(410) = 820$.

Time = 11.30 (sec) , antiderivative size = 1967, normalized size of antiderivative = 4.51

$$\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^2} dx = \text{Too large to display}$$

[In] `integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

$d^5 e + a^2 c d^4 f) * f)$

Mupad [**F(-1)**]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^2} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^2 (c + dx)^2} dx$$

[In] $\text{int}(1/((a + a*\tan(e + f*x)*1i)^2*(c + d*x)^2), x)$

[Out] $\text{int}(1/((a + a*\tan(e + f*x)*1i)^2*(c + d*x)^2), x)$

3.29 $\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx$

Optimal result	189
Rubi [A] (verified)	190
Mathematica [A] (verified)	192
Maple [A] (verified)	193
Fricas [A] (verification not implemented)	193
Sympy [A] (verification not implemented)	194
Maxima [F(-2)]	195
Giac [A] (verification not implemented)	195
Mupad [B] (verification not implemented)	196

Optimal result

Integrand size = 23, antiderivative size = 396

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx = & -\frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4} \\ & - \frac{9id^2e^{-2ie-2ifx}(c+dx)}{32a^3f^3} - \frac{9id^2e^{-4ie-4ifx}(c+dx)}{256a^3f^3} \\ & - \frac{id^2e^{-6ie-6ifx}(c+dx)}{288a^3f^3} + \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} \\ & + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} \\ & + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} \\ & + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d} \end{aligned}$$

```
[Out] -9/64*d^3*exp(-2*I*e-2*I*f*x)/a^3/f^4-9/1024*d^3*exp(-4*I*e-4*I*f*x)/a^3/f^4-1/1728*d^3*exp(-6*I*e-6*I*f*x)/a^3/f^4-9/32*I*d^2*exp(-2*I*e-2*I*f*x)*(d*x+c)/a^3/f^3-9/256*I*d^2*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^3-1/288*I*d^2*exp(-6*I*e-6*I*f*x)*(d*x+c)/a^3/f^3+9/32*d*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f^2+9/128*d*exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f^2+1/96*d*exp(-6*I*e-6*I*f*x)*(d*x+c)^2/a^3/f^2+3/16*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^3/a^3/f^3+3/32*I*exp(-4*I*e-4*I*f*x)*(d*x+c)^3/a^3/f^3+48*I*exp(-6*I*e-6*I*f*x)*(d*x+c)^3/a^3/f^3+32*(d*x+c)^4/a^3/d
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {3810, 2207, 2225}

$$\int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx = -\frac{9id^2(c+dx)e^{-2ie-2ifx}}{32a^3f^3} - \frac{9id^2(c+dx)e^{-4ie-4ifx}}{256a^3f^3} \\ - \frac{id^2(c+dx)e^{-6ie-6ifx}}{288a^3f^3} + \frac{9d(c+dx)^2e^{-2ie-2ifx}}{32a^3f^2} \\ + \frac{9d(c+dx)^2e^{-4ie-4ifx}}{128a^3f^2} + \frac{d(c+dx)^2e^{-6ie-6ifx}}{96a^3f^2} \\ + \frac{3i(c+dx)^3e^{-2ie-2ifx}}{16a^3f} + \frac{3i(c+dx)^3e^{-4ie-4ifx}}{32a^3f} \\ + \frac{i(c+dx)^3e^{-6ie-6ifx}}{48a^3f} + \frac{(c+dx)^4}{32a^3d} \\ - \frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4}$$

[In] $\text{Int}[(c+d*x)^3/(a+I*a*\text{Tan}[e+f*x])^3, x]$

[Out] $(-9*d^3*E^((-2*I)*e - (2*I)*f*x))/(64*a^3*f^4) - (9*d^3*E^((-4*I)*e - (4*I)*f*x))/(1024*a^3*f^4) - (d^3*E^((-6*I)*e - (6*I)*f*x))/(1728*a^3*f^4) - (((9*I)/32)*d^2*E^((-2*I)*e - (2*I)*f*x)*(c+d*x))/(a^3*f^3) - (((9*I)/256)*d^2*E^((-4*I)*e - (4*I)*f*x)*(c+d*x))/(a^3*f^3) - ((I/288)*d^2*E^((-6*I)*e - (6*I)*f*x)*(c+d*x))/(a^3*f^3) + (9*d*E^((-2*I)*e - (2*I)*f*x)*(c+d*x)^2)/(32*a^3*f^2) + (9*d*E^((-4*I)*e - (4*I)*f*x)*(c+d*x)^2)/(128*a^3*f^2) + (d*E^((-6*I)*e - (6*I)*f*x)*(c+d*x)^2)/(96*a^3*f^2) + (((3*I)/16)*E^((-2*I)*e - (2*I)*f*x)*(c+d*x)^3)/(a^3*f) + (((3*I)/32)*E^((-4*I)*e - (4*I)*f*x)*(c+d*x)^3)/(a^3*f) + ((I/48)*E^((-6*I)*e - (6*I)*f*x)*(c+d*x)^3)/(a^3*f) + (c+d*x)^4/(32*a^3*d)$

Rule 2207

```
Int[((b_)*(F_)^((g_.)*((e_.)+(f_)*(x_))))^(n_.)*((c_.)+(d_)*(x_))^(m_), x_Symbol] :> Simp[(c+d*x)^m*((b*F^(g*(e+f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c+d*x)^(m-1)*(b*F^(g*(e+f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_.)*((a_.)+(b_)*(x_))))^(n_.), x_Symbol] :> Simp[(F^(c*(a+b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^3}{8a^3} + \frac{3e^{-2ie-2ifx}(c+dx)^3}{8a^3} + \frac{3e^{-4ie-4ifx}(c+dx)^3}{8a^3} \right. \\
&\quad \left. + \frac{e^{-6ie-6ifx}(c+dx)^3}{8a^3} \right) dx \\
&= \frac{(c+dx)^4}{32a^3d} + \frac{\int e^{-6ie-6ifx}(c+dx)^3 dx}{8a^3} + \frac{3 \int e^{-2ie-2ifx}(c+dx)^3 dx}{8a^3} + \frac{3 \int e^{-4ie-4ifx}(c+dx)^3 dx}{8a^3} \\
&= \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} \\
&\quad + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d} - \frac{(id) \int e^{-6ie-6ifx}(c+dx)^2 dx}{16a^3f} \\
&\quad - \frac{(9id) \int e^{-4ie-4ifx}(c+dx)^2 dx}{32a^3f} - \frac{(9id) \int e^{-2ie-2ifx}(c+dx)^2 dx}{16a^3f} \\
&= \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} \\
&\quad + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} \\
&\quad + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d} - \frac{d^2 \int e^{-6ie-6ifx}(c+dx) dx}{48a^3f^2} \\
&\quad - \frac{(9d^2) \int e^{-4ie-4ifx}(c+dx) dx}{64a^3f^2} - \frac{(9d^2) \int e^{-2ie-2ifx}(c+dx) dx}{16a^3f^2} \\
&= -\frac{9id^2e^{-2ie-2ifx}(c+dx)}{32a^3f^3} - \frac{9id^2e^{-4ie-4ifx}(c+dx)}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}(c+dx)}{288a^3f^3} \\
&\quad + \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} \\
&\quad + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d} \\
&\quad + \frac{(id^3) \int e^{-6ie-6ifx} dx}{288a^3f^3} + \frac{(9id^3) \int e^{-4ie-4ifx} dx}{256a^3f^3} + \frac{(9id^3) \int e^{-2ie-2ifx} dx}{32a^3f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{9d^3e^{-2ie-2ifx}}{64a^3f^4} - \frac{9d^3e^{-4ie-4ifx}}{1024a^3f^4} - \frac{d^3e^{-6ie-6ifx}}{1728a^3f^4} - \frac{9id^2e^{-2ie-2ifx}(c+dx)}{32a^3f^3} \\
&\quad - \frac{9id^2e^{-4ie-4ifx}(c+dx)}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}(c+dx)}{288a^3f^3} + \frac{9de^{-2ie-2ifx}(c+dx)^2}{32a^3f^2} \\
&\quad + \frac{9de^{-4ie-4ifx}(c+dx)^2}{128a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)^2}{96a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^3}{16a^3f} \\
&\quad + \frac{3ie^{-4ie-4ifx}(c+dx)^3}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^3}{48a^3f} + \frac{(c+dx)^4}{32a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.76 (sec) , antiderivative size = 667, normalized size of antiderivative = 1.68

$$\begin{aligned}
&\int \frac{(c+dx)^3}{(a+i a \tan(e+fx))^3} dx \\
&= \frac{i \sec^3(e+fx) (243(32ic^3f^3 + 8c^2df^2(5 + 12ifx) + 4cd^2f(-9i + 20fx + 24if^2x^2) + d^3(-17 - 36ifx + 40j
\end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + I*a*Tan[e + f*x])^3, x]

[Out] ((I/27648)*Sec[e + f*x]^3*(243*((32*I)*c^3*f^3 + 8*c^2*d*f^2*(5 + (12*I)*f*x) + 4*c*d^2*f*(-9*I + 20*f*x + (24*I)*f^2*x^2) + d^3*(-17 - (36*I)*f*x + 40*f^2*x^2 + (32*I)*f^3*x^3))*Cos[e + f*x] + 16*(36*c^3*f^3*(I + 6*f*x) + 18*c^2*d*f^2*(1 + (6*I)*f*x + 18*f^2*x^2) + 6*c*d^2*f*(-I + 6*f*x + (18*I)*f^2*x^2 + 36*f^3*x^3) + d^3*(-1 - (6*I)*f*x + 18*f^2*x^2 + (36*I)*f^3*x^3 + 54*f^4*x^4))*Cos[3*(e + f*x)] - (3645*I)*d^3*Sin[e + f*x] + 6804*c*d^2*f*Sin[e + f*x] + (5832*I)*c^2*d*f^2*Sin[e + f*x] - 2592*c^3*f^3*Sin[e + f*x] + 6804*d^3*f*x*Sin[e + f*x] + (11664*I)*c*d^2*f^2*x*Sin[e + f*x] - 7776*c^2*d*f^3*x*Sin[e + f*x] + (5832*I)*d^3*f^2*x^2*Sin[e + f*x] - 7776*c*d^2*f^3*x^2*Sin[e + f*x] - 2592*d^3*f^3*x^3*Sin[e + f*x] + (16*I)*d^3*Sin[3*(e + f*x)] - 96*c*d^2*f*Sin[3*(e + f*x)] - (288*I)*c^2*d*f^2*Sin[3*(e + f*x)] + 576*c^3*f^3*Sin[3*(e + f*x)] - 96*d^3*f*x*Sin[3*(e + f*x)] - (576*I)*c*d^2*f^2*x*Sin[3*(e + f*x)] + 1728*c^2*d*f^3*x*Sin[3*(e + f*x)] + (3456*I)*c^3*f^4*x*Sin[3*(e + f*x)] - (288*I)*d^3*f^2*x^2*Sin[3*(e + f*x)] + 1728*c*d^2*f^3*x^2*Sin[3*(e + f*x)] + (5184*I)*c^2*d*f^4*x^2*Sin[3*(e + f*x)] + 576*d^3*f^3*x^3*Sin[3*(e + f*x)] + (3456*I)*c*d^2*f^4*x^3*Sin[3*(e + f*x)] + (864*I)*d^3*f^4*x^4*Sin[3*(e + f*x)]))/((a^3*f^4*(-I + Tan[e + f*x]))^3)

Maple [A] (verified)

Time = 1.21 (sec) , antiderivative size = 396, normalized size of antiderivative = 1.00

method	result
risch	$\frac{d^3x^4}{32a^3} + \frac{d^2cx^3}{8a^3} + \frac{3dc^2x^2}{16a^3} + \frac{c^3x}{8a^3} + \frac{c^4}{32a^3d} + \frac{3i(4d^3x^3f^3+12cd^2f^3x^2-6id^3f^2x^2+12c^2df^3x-12icd^2f^2x+4c^3f^3-6ic^2df^2-6cd^2f^3)}{64a^3f^4}$

[In] `int((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/32/a^3*d^3*x^4 + 1/8/a^3*d^2*c*x^3 + 3/16/a^3*d*c^2*x^2 + 1/8/a^3*c^3*x + 1/32/a^3/d*c^4 + 3/64*I*(4*d^3*x^3*f^3 - 6*I*d^3*x^2*f^2*x^2 + 12*c*d^2*f^3*x^2 - 12*I*c*d^2*f^2*x^2 + 12*c^2*d*f^3*x^3 - 6*I*c^2*d*f^2*x^3 + 3*I*d^3*x^3*f*x + 3*I*d^3*x^3*f^2*x^2 + 96*c*d^2*f^2*x^2 + 32*d^3*x^3*f^3 - 24*I*d^3*x^2*f^2*x^2 + 32*c^3*x^3*f^3 - 12*d^3*x^3*f*x^2 + 48*I*c*d^2*f^2*x^3 + 96*c^2*d*f^3*x^2 - 24*I*c^2*d*f^2*x^2 + 32*c^3*x^3*f^3 - 12*d^3*x^3*f*x^2 + 108*c*d^2*f^2*x^3 - 36*I*c*d^2*f^2*x^2 + 108*c^2*d*f^3*x^3 - 18*I*c^2*d*f^2*x^2 + 36*c^3*x^3*f^3 - 6*d^3*x^3*f*x + I*d^3*x^3*f^2*x^2)/a^3/f^4 + exp(-6*I*(f*x+e)) + 1/1728*I*(36*d^3*x^3*f^3 - 18*I*d^3*x^3*f^2*x^2 + 108*c*d^2*f^3*x^2 - 36*I*c*d^2*f^2*x^3 + 108*c^2*d*f^3*x^3 - 18*I*c^2*d*f^2*x^2 + 36*c^3*x^3*f^3 - 6*d^3*x^3*f*x + I*d^3*x^3*f^2*x^2)/a^3/f^4 + exp(-6*I*(f*x+e))) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 370, normalized size of antiderivative = 0.93

$$\begin{aligned} & \int \frac{(c+dx)^3}{(a+ia\tan(e+fx))^3} dx \\ &= \frac{(576i d^3 f^3 x^3 + 576i c^3 f^3 + 288 c^2 d f^2 - 96 i c d^2 f - 16 d^3 - 288 (-6i c d^2 f^3 - d^3 f^2) x^2 - 96 (-18i c^2 d f^3 - 6i c^3 f^2) x^3 + 288 (-6i c * d^2 f^3 - d^3 f^2) x^4)}{(a+ia\tan(e+fx))^3} \end{aligned}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/27648*(576*I*d^3*x^3*f^3 + 576*I*c^3*x^3*f^3 + 288*c^2*d*f^2 - 96*I*c*d^2*f^2 - 16*d^3 - 288*(-6*I*c*d^2*f^3 - d^3*f^2)*x^2 - 96*(-18*I*c^2*d*f^3 - 6*c*d^2*f^2 + I*d^3*f)*x + 864*(d^3*x^4*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*x^4*f*x + 6*I*e) - 1296*(-4*I*d^3*x^3*f^3*x^3 - 4*I*c^3*x^3*f^3 - 6*c^2*d*f^2*x^2 + 6*I*c*d^2*f^2*x + 3*d^3*x^3 + 6*(-2*I*c*d^2*f^3*x^3 - d^3*f^2*x^2)*x^2 + 6*(-2*I*c^2*d*f^3*x^3 - 2*c*d^2*f^2*x^2 + I*d^3*f*x)*x)*e^(4*I*f*x + 4*I*e) - 81*(-32*I*d^3*x^3*f^3*x^3 - 32*I*c^3*x^3*f^3*x^3 - 24*c^2*d*f^2*x^2 + 12*I*c*d^2*f^2*x^2 + 3*d^3*x^3 + 24*(-4*I*c*d^2*f^3*x^3 - d^3*f^2*x^2)*x^2 + 12*(-8*I*c^2*d*f^3*x^3 - 4*c*d^2*f^2*x^2 + I*d^3*f*x)*x)*e^(2*I*f*x + 2*I*e)*e^(-6*I*f*x - 6*I*e)/(a^3*f^4) \end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 945, normalized size of antiderivative = 2.39

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx$$

$$= \begin{cases} \frac{((2359296ia^6c^3f^{11}e^{6ie} + 7077888ia^6c^2df^{11}xe^{6ie} + 1179648a^6c^2df^{10}e^{6ie} + 7077888ia^6cd^2f^{11}x^2e^{6ie} + 2359296a^6cd^2f^{10}xe^{6ie} - 393216ia^6cd^2f^9e^{6ie})e^{-6ie}}{32a^3} + \frac{x^3 \cdot (3cd^2e^{4ie} + 3cd^2e^{2ie} + cd^2)e^{-6ie}}{8a^3} + \frac{x^2 \cdot (9c^2de^{4ie} + 9c^2de^{2ie} + 3c^2d)e^{-6ie}}{16a^3} + \frac{x(3c^3e^{4ie} + 3c^3e^{2ie} + c^3)e^{-6ie}}{8a^3} \\ + \frac{c^3x}{8a^3} + \frac{3c^2dx^2}{16a^3} + \frac{cd^2x^3}{8a^3} + \frac{d^3x^4}{32a^3} \end{cases}$$

[In] `integrate((d*x+c)**3/(a+I*a*tan(f*x+e))**3,x)`

[Out] Piecewise(((2359296*I*a**6*c**3*f**11*exp(6*I*e) + 7077888*I*a**6*c**2*d*f**11*x*exp(6*I*e) + 1179648*a**6*c**2*d*f**10*exp(6*I*e) + 7077888*I*a**6*c*d**2*f**10*x*exp(6*I*e) - 393216*I*a**6*c*d**2*f**9*exp(6*I*e) + 2359296*a**6*c*d**2*f**10*x*exp(6*I*e) - 393216*I*a**6*d**3*f**11*x**3*exp(6*I*e) + 1179648*a**6*d**3*f**10*x**2*exp(6*I*e) - 65536*a**6*d**3*f**8*exp(6*I*e))*exp(-6*I*f*x) + (10616832*I*a**6*c**3*f**11*exp(8*I*e) + 31850496*I*a**6*c**2*d*f**11*x*exp(8*I*e) + 7962624*a**6*c**2*d*f**10*exp(8*I*e) + 31850496*I*a**6*c*d**2*f**11*x**2*exp(8*I*e) + 15925248*a**6*c*d**2*f**10*x*exp(8*I*e) - 3981312*I*a**6*c*d**2*f**9*exp(8*I*e) + 10616832*I*a**6*d**3*f**11*x**3*exp(8*I*e) + 7962624*a**6*d**3*f**10*x**2*exp(8*I*e) - 3981312*I*a**6*d**3*f**9*x*exp(8*I*e) - 995328*a**6*d**3*f**8*exp(8*I*e))*exp(-4*I*f*x) + (21233664*I*a**6*c**3*f**11*exp(10*I*e) + 63700992*I*a**6*c**2*d*f**11*x*exp(10*I*e) + 31850496*a**6*c**2*d*f**10*exp(10*I*e) + 63700992*I*a**6*c*d**2*f**11*x**2*exp(10*I*e) + 63700992*a**6*c*d**2*f**10*x*exp(10*I*e) - 31850496*I*a**6*c*d**2*f**9*exp(10*I*e) + 21233664*I*a**6*d**3*f**11*x**3*exp(10*I*e) + 31850496*a**6*d**3*f**10*x**2*exp(10*I*e) - 31850496*I*a**6*d**3*f**9*x*exp(10*I*e) - 15925248*a**6*d**3*f**8*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(113246208*a**9*f**12), Ne(a**9*f**12*exp(12*I*e), 0)), (x**4*(3*d**3*exp(4*I*e) + 3*d**3*exp(2*I*e) + d**3)*exp(-6*I*e)/(32*a**3) + x**3*(3*c*d**2*exp(4*I*e) + 3*c*d**2*exp(2*I*e) + c*d**2)*exp(-6*I*e)/(8*a**3) + x**2*(9*c**2*d*exp(4*I*e) + 9*c**2*d*exp(2*I*e) + 3*c**2*d)*exp(-6*I*e)/(16*a**3) + x*(3*c**3*exp(4*I*e) + 3*c**3*exp(2*I*e) + c**3)*exp(-6*I*e)/(8*a**3), True)) + c**3*x/(8*a**3) + 3*c**2*d*x**2/(16*a**3) + c*d**2*x**3/(8*a**3) + d**3*x**4/(32*a**3)

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.58 (sec) , antiderivative size = 548, normalized size of antiderivative = 1.38

$$\begin{aligned} & \int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx \\ &= \frac{(864 d^3 f^4 x^4 e^{(6i f x + 6i e)} + 3456 c d^2 f^4 x^3 e^{(6i f x + 6i e)} + 5184 c^2 d f^4 x^2 e^{(6i f x + 6i e)} + 5184 i d^3 f^3 x^3 e^{(4i f x + 4i e)} + 2592 i c^2 d^2 f^3 x^2 e^{(4i f x + 4i e)} + 15552 i c d^2 f^2 x e^{(4i f x + 4i e)} + 7776 i c^3 d f x e^{(4i f x + 4i e)} + 1728 i c^2 d^2 f x^2 e^{(4i f x + 4i e)} + 5184 i c^3 f^3 x^3 e^{(4i f x + 4i e)} + 1944 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} + 3888 i c d^2 f^2 x^2 e^{(4i f x + 4i e)} + 576 i c^3 d^2 f x^3 e^{(4i f x + 4i e)} + 7776 i c^2 d^2 f^2 x^3 e^{(4i f x + 4i e)} - 7776 i c d^3 f x^2 e^{(4i f x + 4i e)} - 1944 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} - 972 i c d^3 f x^3 e^{(4i f x + 4i e)} - 288 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} - 96 i c d^3 f x^4 e^{(4i f x + 4i e)} - 3888 i c^3 d^2 f x^3 e^{(4i f x + 4i e)} - 243 i c^2 d^3 f x^4 e^{(4i f x + 4i e)} - 16 d^3) e^{(-6i f x - 6i e)})}{(a^3 f^4)}$$

[In] `integrate((d*x+c)^3/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")`

[Out] $\frac{1}{27648} (864 d^3 f^4 x^4 e^{(6i f x + 6i e)} + 3456 c d^2 f^4 x^3 e^{(6i f x + 6i e)} + 5184 c^2 d f^4 x^2 e^{(6i f x + 6i e)} + 5184 i d^3 f^3 x^3 e^{(4i f x + 4i e)} + 2592 i c^2 d^2 f^3 x^2 e^{(4i f x + 4i e)} + 15552 i c d^2 f^2 x e^{(4i f x + 4i e)} + 7776 i c^3 d f x e^{(4i f x + 4i e)} + 1728 i c^2 d^2 f x^2 e^{(4i f x + 4i e)} + 5184 i c^3 f^3 x^3 e^{(4i f x + 4i e)} + 1944 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} + 3888 i c d^2 f^2 x^2 e^{(4i f x + 4i e)} + 576 i c^3 d^2 f x^3 e^{(4i f x + 4i e)} + 7776 i c^2 d^2 f^2 x^3 e^{(4i f x + 4i e)} - 7776 i c d^3 f x^2 e^{(4i f x + 4i e)} - 1944 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} - 972 i c d^3 f x^3 e^{(4i f x + 4i e)} - 288 i c^2 d^2 f^2 x^2 e^{(4i f x + 4i e)} - 96 i c d^3 f x^4 e^{(4i f x + 4i e)} - 3888 i c^3 d^2 f x^3 e^{(4i f x + 4i e)} - 243 i c^2 d^3 f x^4 e^{(4i f x + 4i e)} - 16 d^3) e^{(-6i f x - 6i e)}) / (a^3 f^4)$

Mupad [B] (verification not implemented)

Time = 4.25 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.04

$$\int \frac{(c + dx)^3}{(a + ia \tan(e + fx))^3} dx = e^{-e^{2i-fx}2i} \left(\frac{(12c^3f^3 - c^2d^2f^218i - 18cd^2f + d^39i)1i}{64a^3f^4} \right. \\ \left. + \frac{d^3x^33i}{16a^3f} - \frac{dx(-2c^2f^2 + cd^2f2i + d^2)9i}{32a^3f^3} \right. \\ \left. - \frac{d^2x^2(-2cf + d1i)9i}{32a^3f^2} \right) \\ + e^{-e^{4i-fx}4i} \left(\frac{(96c^3f^3 - c^2d^2f^272i - 36cd^2f + d^39i)1i}{1024a^3f^4} \right. \\ \left. + \frac{d^3x^33i}{32a^3f} - \frac{dx(-8c^2f^2 + cd^2f4i + d^2)9i}{256a^3f^3} \right. \\ \left. - \frac{d^2x^2(-4cf + d1i)9i}{128a^3f^2} \right) \\ + e^{-e^{6i-fx}6i} \left(\frac{(36c^3f^3 - c^2d^2f^218i - 6cd^2f + d^31i)1i}{1728a^3f^4} \right. \\ \left. + \frac{d^3x^31i}{48a^3f} - \frac{dx(-18c^2f^2 + cd^2f6i + d^2)1i}{288a^3f^3} \right. \\ \left. - \frac{d^2x^2(-6cf + d1i)1i}{96a^3f^2} \right) + \frac{c^3x}{8a^3} + \frac{d^3x^4}{32a^3} + \frac{3c^2dx^2}{16a^3} + \frac{cd^2x^3}{8a^3}$$

[In] `int((c + d*x)^3/(a + a*tan(e + f*x)*1i)^3, x)`

[Out] `exp(- e*2i - f*x*2i)*(((d^3*9i + 12*c^3*f^3 - c^2*d*f^2*18i - 18*c*d^2*f)*1i)/(64*a^3*f^4) + (d^3*x^3*3i)/(16*a^3*f) - (d*x*(d^2 - 2*c^2*f^2 + c*d*f*2i)*9i)/(32*a^3*f^3) - (d^2*x^2*(d*1i - 2*c*f)*9i)/(32*a^3*f^2)) + exp(- e*4i - f*x*4i)*(((d^3*9i + 96*c^3*f^3 - c^2*d*f^2*72i - 36*c*d^2*f)*1i)/(1024*a^3*f^4) + (d^3*x^3*3i)/(32*a^3*f) - (d*x*(d^2 - 8*c^2*f^2 + c*d*f*4i)*9i)/(256*a^3*f^3) - (d^2*x^2*(d*1i - 4*c*f)*9i)/(128*a^3*f^2)) + exp(- e*6i - f*x*6i)*(((d^3*1i + 36*c^3*f^3 - c^2*d*f^2*18i - 6*c*d^2*f)*1i)/(1728*a^3*f^4) + (d^3*x^3*1i)/(48*a^3*f) - (d*x*(d^2 - 18*c^2*f^2 + c*d*f*6i)*1i)/(288*a^3*f^3) - (d^2*x^2*(d*1i - 6*c*f)*1i)/(96*a^3*f^2)) + (c^3*x)/(8*a^3) + (d^3*x^4)/(32*a^3) + (3*c^2*d*x^2)/(16*a^3) + (c*d^2*x^3)/(8*a^3)`

3.30 $\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$

Optimal result	197
Rubi [A] (verified)	197
Mathematica [A] (verified)	199
Maple [A] (verified)	200
Fricas [A] (verification not implemented)	200
Sympy [A] (verification not implemented)	201
Maxima [F(-2)]	201
Giac [A] (verification not implemented)	202
Mupad [B] (verification not implemented)	202

Optimal result

Integrand size = 23, antiderivative size = 294

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx = & -\frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3} \\ & + \frac{3de^{-2ie-2ifx}(c+dx)}{16a^3f^2} + \frac{3de^{-4ie-4ifx}(c+dx)}{64a^3f^2} \\ & + \frac{de^{-6ie-6ifx}(c+dx)}{144a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} \\ & + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} + \frac{(c+dx)^3}{24a^3d} \end{aligned}$$

```
[Out] -3/32*I*d^2*exp(-2*I*e-2*I*f*x)/a^3/f^3-3/256*I*d^2*exp(-4*I*e-4*I*f*x)/a^3
/f^3-1/864*I*d^2*exp(-6*I*e-6*I*f*x)/a^3/f^3+3/16*d*exp(-2*I*e-2*I*f*x)*(d*
x+c)/a^3/f^2+3/64*d*exp(-4*I*e-4*I*f*x)*(d*x+c)/a^3/f^2+1/144*d*exp(-6*I*e-
6*I*f*x)*(d*x+c)/a^3/f^2+3/16*I*exp(-2*I*e-2*I*f*x)*(d*x+c)^2/a^3/f+3/32*I*
exp(-4*I*e-4*I*f*x)*(d*x+c)^2/a^3/f+1/48*I*exp(-6*I*e-6*I*f*x)*(d*x+c)^2/a^
3/f+1/24*(d*x+c)^3/a^3/d
```

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.130, Rules used

$$= \{3810, 2207, 2225\}$$

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx = & \frac{3d(c+dx)e^{-2ie-2ifx}}{16a^3f^2} + \frac{3d(c+dx)e^{-4ie-4ifx}}{64a^3f^2} \\ & + \frac{d(c+dx)e^{-6ie-6ifx}}{144a^3f^2} + \frac{3i(c+dx)^2e^{-2ie-2ifx}}{16a^3f} \\ & + \frac{3i(c+dx)^2e^{-4ie-4ifx}}{32a^3f} + \frac{i(c+dx)^2e^{-6ie-6ifx}}{48a^3f} + \frac{(c+dx)^3}{24a^3d} \\ & - \frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3} \end{aligned}$$

[In] $\text{Int}[(c+d*x)^2/(a+I*a*\text{Tan}[e+f*x])^3, x]$

[Out] $\left(\left((-3*I)/32\right)*d^2*E^{\left((-2*I)*e-(2*I)*f*x\right)}/(a^3*f^3)-\left((3*I)/256\right)*d^2*E^{\left((-4*I)*e-(4*I)*f*x\right)}/(a^3*f^3)-\left((I/864)*d^2*E^{\left((-6*I)*e-(6*I)*f*x\right)}/(a^3*f^3)+\left(3*d*E^{\left((-2*I)*e-(2*I)*f*x\right)*(c+d*x)}\right)/(16*a^3*f^2)+\left(3*d*E^{\left((-4*I)*e-(4*I)*f*x\right)*(c+d*x)}\right)/(64*a^3*f^2)+\left(d*E^{\left((-6*I)*e-(6*I)*f*x\right)*(c+d*x)}\right)/(144*a^3*f^2)+\left((3*I)/16\right)*E^{\left((-2*I)*e-(2*I)*f*x\right)*(c+d*x)^2}/(a^3*f)+\left((3*I)/32\right)*E^{\left((-4*I)*e-(4*I)*f*x\right)*(c+d*x)^2}/(a^3*f)+\left(I/48\right)*E^{\left((-6*I)*e-(6*I)*f*x\right)*(c+d*x)^2}/(a^3*f)+(c+d*x)^3/(24*a^3*d)\right)$

Rule 2207

```
Int[((b_)*(F_))^((g_)*(e_)+(f_)*(x_)))^(n_)*((c_)+(d_)*(x_))^(m_),
x_Symbol] :> Simp[(c+d*x)^m*((b*F^(g*(e+f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c+d*x)^(m-1)*(b*F^(g*(e+f*x)))^n, x], x]; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_)*(a_)+(b_)*(x_)))^(n_), x_Symbol] :> Simp[(F^(c*(a+b*x)))^n/(b*c*n*Log[F]), x]; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```
Int[((c_)+(d_)*(x_))^(m_)*((a_)+(b_)*tan[(e_)+(f_)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c+d*x)^m, (1/(2*a)+E^(2*(a/b))*(e+f*x))/(2*a)^(-n)], x]; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2+b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^2}{8a^3} + \frac{3e^{-2ie-2ifx}(c+dx)^2}{8a^3} + \frac{3e^{-4ie-4ifx}(c+dx)^2}{8a^3} \right. \\
&\quad \left. + \frac{e^{-6ie-6ifx}(c+dx)^2}{8a^3} \right) dx \\
&= \frac{(c+dx)^3}{24a^3d} + \frac{\int e^{-6ie-6ifx}(c+dx)^2 dx}{8a^3} + \frac{3 \int e^{-2ie-2ifx}(c+dx)^2 dx}{8a^3} + \frac{3 \int e^{-4ie-4ifx}(c+dx)^2 dx}{8a^3} \\
&= \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} \\
&\quad + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} + \frac{(c+dx)^3}{24a^3d} - \frac{(id) \int e^{-6ie-6ifx}(c+dx) dx}{24a^3f} \\
&\quad - \frac{(3id) \int e^{-4ie-4ifx}(c+dx) dx}{16a^3f} - \frac{(3id) \int e^{-2ie-2ifx}(c+dx) dx}{8a^3f} \\
&= \frac{3de^{-2ie-2ifx}(c+dx)}{16a^3f^2} + \frac{3de^{-4ie-4ifx}(c+dx)}{64a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)}{144a^3f^2} \\
&\quad + \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} \\
&\quad + \frac{(c+dx)^3}{24a^3d} - \frac{d^2 \int e^{-6ie-6ifx} dx}{144a^3f^2} - \frac{(3d^2) \int e^{-4ie-4ifx} dx}{64a^3f^2} - \frac{(3d^2) \int e^{-2ie-2ifx} dx}{16a^3f^2} \\
&= -\frac{3id^2e^{-2ie-2ifx}}{32a^3f^3} - \frac{3id^2e^{-4ie-4ifx}}{256a^3f^3} - \frac{id^2e^{-6ie-6ifx}}{864a^3f^3} + \frac{3de^{-2ie-2ifx}(c+dx)}{16a^3f^2} \\
&\quad + \frac{3de^{-4ie-4ifx}(c+dx)}{64a^3f^2} + \frac{de^{-6ie-6ifx}(c+dx)}{144a^3f^2} + \frac{3ie^{-2ie-2ifx}(c+dx)^2}{16a^3f} \\
&\quad + \frac{3ie^{-4ie-4ifx}(c+dx)^2}{32a^3f} + \frac{ie^{-6ie-6ifx}(c+dx)^2}{48a^3f} + \frac{(c+dx)^3}{24a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.74 (sec), antiderivative size = 405, normalized size of antiderivative = 1.38

$$\begin{aligned}
&\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx \\
&= \frac{i \sec^3(e+fx) (81(24ic^2f^2 + 4cdf(5 + 12ifx) + d^2(-9i + 20fx + 24if^2x^2)) \cos(e+fx) + 8(18c^2f^2(i+6ifx)))}{(a+ia\tan(e+fx))^3}
\end{aligned}$$

[In] `Integrate[(c + d*x)^2/(a + I*a*Tan[e + f*x])^3, x]`

[Out] `((I/6912)*Sec[e + f*x]^3*(81*((24*I)*c^2*f^2 + 4*c*d*f*(5 + (12*I)*f*x) + d^2*(-9*I + 20*f*x + (24*I)*f^2*x^2))*Cos[e + f*x] + 8*(18*c^2*f^2*(I + 6*f*x) + 6*c*d*f*(1 + (6*I)*f*x + 18*f^2*x^2) + d^2*(-I + 6*f*x + (18*I)*f^2*x^2))`

$$\begin{aligned}
& 2 + 36*f^3*x^3)*Cos[3*(e + f*x)] + 567*d^2*Sin[e + f*x] + (972*I)*c*d*f*Sin[e + f*x] - 648*c^2*f^2*Sin[e + f*x] + (972*I)*d^2*f*x*Sin[e + f*x] - 1296*c*d*f^2*x*Sin[e + f*x] - 648*d^2*f^2*x^2*Sin[e + f*x] - 8*d^2*Sin[3*(e + f*x)] - (48*I)*c*d*f*Sin[3*(e + f*x)] + 144*c^2*f^2*Sin[3*(e + f*x)] - (48*I)*d^2*f*x*Sin[3*(e + f*x)] + 288*c*d*f^2*x*Sin[3*(e + f*x)] + (864*I)*c^2*f^3*x*Sin[3*(e + f*x)] + 144*d^2*f^2*x^2*Sin[3*(e + f*x)] + (864*I)*c*d*f^3*x^2*Sin[3*(e + f*x)] + (288*I)*d^2*f^3*x^3*Sin[3*(e + f*x)])/(a^3*f^3*(-I + Tan[e + f*x]))^3
\end{aligned}$$

Maple [A] (verified)

Time = 1.08 (sec) , antiderivative size = 238, normalized size of antiderivative = 0.81

method	result
risch	$\frac{d^2x^3}{24a^3} + \frac{dcx^2}{8a^3} + \frac{c^2x}{8a^3} + \frac{c^3}{24a^3d} + \frac{3i(2d^2x^2f^2+4cd f^2x-2id^2fx+2c^2f^2-2icdf-d^2)e^{-2i(fx+e)}}{32a^3f^3} + \frac{3i(8d^2x^2f^2+16cd f^2x-4id^2fx-2icdf-d^2)e^{-2i(fx+e)}}{256a^3f^3}$

[In] `int((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`
[Out]
$$\begin{aligned}
& 1/24/a^3*d^2*x^3+1/8/a^3*d*c*x^2+1/8/a^3*c^2*x+1/24/a^3/d*c^3+3/32*I*(2*d^2 \\
& *x^2*f^2-2*I*d^2*f*x+4*c*d*f^2*x-2*I*c*d*f+2*c^2*f^2-d^2)/a^3/f^3*exp(-2*I* \\
& (f*x+e))+3/256*I*(8*d^2*x^2*f^2-4*I*d^2*f*x+16*c*d*f^2*x-4*I*c*d*f+8*c^2*f^2 \\
& -2*d^2)/a^3/f^3*exp(-4*I*(f*x+e))+1/864*I*(18*d^2*x^2*f^2-6*I*d^2*f*x+36*c*d \\
& *f^2*x-6*I*c*d*f+18*c^2*f^2-d^2)/a^3/f^3*exp(-6*I*(f*x+e))
\end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 221, normalized size of antiderivative = 0.75

$$\begin{aligned}
& \int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx \\
& = \frac{(144i d^2 f^2 x^2 + 144i c^2 f^2 + 48 cdf - 8i d^2 - 48 (-6i cdf^2 - d^2 f)x + 288 (d^2 f^3 x^3 + 3 cdf^3 x^2 + 3 c^2 f^3 x)) e^{(6i f x + 6i e)}}{(a + ia \tan(e + fx))^3}
\end{aligned}$$

[In] `integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`
[Out]
$$\begin{aligned}
& 1/6912*(144*I*d^2*f^2*x^2 + 144*I*c^2*f^2 + 48*c*d*f - 8*I*d^2 - 48*(-6*I*c \\
& *d*f^2 - d^2*f)*x + 288*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x)*e^(6*I* \\
& f*x + 6*I*e) - 648*(-2*I*d^2*f^2*x^2 - 2*I*c^2*f^2 - 2*c*d*f + I*d^2 + 2*(- \\
& 2*I*c*d*f^2 - d^2*f)*x)*e^(4*I*f*x + 4*I*e) - 81*(-8*I*d^2*f^2*x^2 - 8*I*c \\
& *f^2 - 4*c*d*f + I*d^2 + 4*(-4*I*c*d*f^2 - d^2*f)*x)*e^(2*I*f*x + 2*I*e))* \\
& e^(-6*I*f*x - 6*I*e)/(a^3*f^3)
\end{aligned}$$

Sympy [A] (verification not implemented)

Time = 0.50 (sec) , antiderivative size = 588, normalized size of antiderivative = 2.00

$$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx$$

$$= \left\{ \begin{array}{l} \frac{((147456ia^6c^2f^8e^{6ie} + 294912ia^6cdf^8xe^{6ie} + 49152a^6cdf^7e^{6ie} + 147456ia^6d^2f^8x^2e^{6ie} + 49152a^6d^2f^7xe^{6ie} - 8192ia^6d^2f^6e^{6ie})e^{-6ifx} + (663552*)}{24a^3} \\ + \frac{x^3 \cdot (3d^2e^{4ie} + 3d^2e^{2ie} + d^2)e^{-6ie}}{8a^3} + \frac{x^2 \cdot (3cde^{4ie} + 3cde^{2ie} + cd)e^{-6ie}}{8a^3} + \frac{x(3c^2e^{4ie} + 3c^2e^{2ie} + c^2)e^{-6ie}}{8a^3} \\ + \frac{c^2x}{8a^3} + \frac{cdx^2}{8a^3} + \frac{d^2x^3}{24a^3} \end{array} \right.$$

[In] integrate((d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)

[Out] Piecewise(((147456*I*a**6*c**2*f**8*exp(6*I*e) + 294912*I*a**6*c*d*f**8*x*exp(6*I*e) + 49152*a**6*c*d*f**7*exp(6*I*e) + 147456*I*a**6*d**2*f**8*x**2*exp(6*I*e) + 49152*a**6*d**2*f**7*x*exp(6*I*e) - 8192*I*a**6*d**2*f**6*exp(6*I*e))*exp(-6*I*f*x) + (663552*I*a**6*c**2*f**8*exp(8*I*e) + 1327104*I*a**6*c*d*f**8*x*exp(8*I*e) + 331776*a**6*c*d*f**7*exp(8*I*e) + 663552*I*a**6*d**2*f**8*x**2*exp(8*I*e) + 331776*a**6*d**2*f**7*x*exp(8*I*e) - 82944*I*a**6*d**2*f**6*exp(8*I*e))*exp(-4*I*f*x) + (1327104*I*a**6*c**2*f**8*exp(10*I*e) + 2654208*I*a**6*c*d*f**8*x*exp(10*I*e) + 1327104*a**6*c*d*f**7*exp(10*I*e) + 1327104*I*a**6*d**2*f**8*x**2*exp(10*I*e) + 1327104*a**6*d**2*f**7*x*exp(10*I*e) - 663552*I*a**6*d**2*f**6*exp(10*I*e))*exp(-2*I*f*x))*exp(-12*I*e)/(7077888*a**9*f**9), Ne(a**9*f**9*exp(12*I*e), 0)), (x**3*(3*d**2*exp(4*I*e) + 3*d**2*exp(2*I*e) + d**2)*exp(-6*I*e)/(24*a**3) + x**2*(3*c*d*exp(4*I*e) + 3*c*d*exp(2*I*e) + c*d)*exp(-6*I*e)/(8*a**3) + x*(3*c**2*exp(4*I*e) + 3*c**2*exp(2*I*e) + c**2)*exp(-6*I*e)/(8*a**3), True)) + c**2*x/(8*a**3) + c*d*x**2/(8*a**3) + d**2*x**3/(24*a**3)

Maxima [F(-2)]

Exception generated.

$$\int \frac{(c+dx)^2}{(a+ia\tan(e+fx))^3} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.62 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.07

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx$$

$$= \frac{(288 d^2 f^3 x^3 e^{(6i f x + 6i e)} + 864 c d f^3 x^2 e^{(6i f x + 6i e)} + 864 c^2 f^3 x e^{(6i f x + 6i e)} + 1296 i d^2 f^2 x^2 e^{(4i f x + 4i e)} + 648 i d^2 f^2 x e^{(4i f x + 4i e)})}{(a + ia \tan(e + fx))^3}$$

```
[In] integrate((d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
[Out] 1/6912*(288*d^2*f^3*x^3*e^(6*I*f*x + 6*I*e) + 864*c*d*f^3*x^2*e^(6*I*f*x + 6*I*e) + 864*c^2*f^3*x*e^(6*I*f*x + 6*I*e) + 1296*I*d^2*f^2*x^2*e^(4*I*f*x + 4*I*e) + 648*I*d^2*f^2*x^2*e^(2*I*f*x + 2*I*e) + 144*I*d^2*f^2*x^2 + 2592*I*c*d*f^2*x*e^(4*I*f*x + 4*I*e) + 1296*I*c*d*f^2*x*e^(2*I*f*x + 2*I*e) + 288*I*c*d*f^2*x + 1296*I*c^2*f^2*x*e^(4*I*f*x + 4*I*e) + 1296*d^2*f*x*x*e^(4*I*f*x + 4*I*e) + 648*I*c^2*f^2*x*e^(2*I*f*x + 2*I*e) + 324*d^2*f*x*x*e^(2*I*f*x + 2*I*e) + 144*I*c^2*f^2 + 48*d^2*f*x + 1296*c*d*f*e^(4*I*f*x + 4*I*e) + 324*c*d*f*e^(2*I*f*x + 2*I*e) + 48*c*d*f - 648*I*d^2*x*e^(4*I*f*x + 4*I*e) - 81*I*d^2*x*e^(2*I*f*x + 2*I*e) - 8*I*d^2*x*e^(-6*I*f*x - 6*I*e)/(a^3*f^3)
```

Mupad [B] (verification not implemented)

Time = 3.95 (sec) , antiderivative size = 263, normalized size of antiderivative = 0.89

$$\int \frac{(c + dx)^2}{(a + ia \tan(e + fx))^3} dx = \frac{c^2 x}{8 a^3} - e^{-e 2i - f x 2i} \left(\frac{(-6 c^2 f^2 + c d f 6i + 3 d^2) 1i}{32 a^3 f^3} - \frac{d^2 x^2 3i}{16 a^3 f} \right.$$

$$\left. + \frac{d x (-2 c f + d 1i) 3i}{16 a^3 f^2} \right)$$

$$- e^{-e 4i - f x 4i} \left(\frac{(-24 c^2 f^2 + c d f 12i + 3 d^2) 1i}{256 a^3 f^3} - \frac{d^2 x^2 3i}{32 a^3 f} \right.$$

$$\left. + \frac{d x (-4 c f + d 1i) 3i}{64 a^3 f^2} \right)$$

$$- e^{-e 6i - f x 6i} \left(\frac{(-18 c^2 f^2 + c d f 6i + d^2) 1i}{864 a^3 f^3} - \frac{d^2 x^2 1i}{48 a^3 f} \right.$$

$$\left. + \frac{d x (-6 c f + d 1i) 1i}{144 a^3 f^2} \right) + \frac{d^2 x^3}{24 a^3} + \frac{c d x^2}{8 a^3}$$

```
[In] int((c + d*x)^2/(a + a*tan(e + f*x)*1i)^3,x)
```

```
[Out] (c^2*x)/(8*a^3) - exp(- e*2i - f*x*2i)*((3*d^2 - 6*c^2*f^2 + c*d*f*6i)*1i)/(32*a^3*f^3) - (d^2*x^2*3i)/(16*a^3*f) + (d*x*(d*1i - 2*c*f)*3i)/(16*a^3*f)
```

$$\begin{aligned} & \sim 2)) - \exp(-e*4i - f*x*4i)*(((3*d^2 - 24*c^2*f^2 + c*d*f*12i)*1i)/(256*a^3 \\ & *f^3) - (d^2*x^2*3i)/(32*a^3*f) + (d*x*(d*1i - 4*c*f)*3i)/(64*a^3*f^2)) - e \\ & \exp(-e*6i - f*x*6i)*(((d^2 - 18*c^2*f^2 + c*d*f*6i)*1i)/(864*a^3*f^3) - (d^2*x^2*1i)/(48*a^3*f) + (d*x*(d*1i - 6*c*f)*1i)/(144*a^3*f^2)) + (d^2*x^3)/(24*a^3) + (c*d*x^2)/(8*a^3) \end{aligned}$$

3.31 $\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx$

Optimal result	204
Rubi [A] (verified)	204
Mathematica [A] (verified)	207
Maple [A] (verified)	207
Fricas [A] (verification not implemented)	207
Sympy [A] (verification not implemented)	208
Maxima [F(-2)]	208
Giac [A] (verification not implemented)	209
Mupad [B] (verification not implemented)	209

Optimal result

Integrand size = 21, antiderivative size = 209

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx = & -\frac{11idx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} \\ & + \frac{d}{36f^2(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\ & + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\ & + \frac{11d}{96f^2(a^3+ia^3\tan(e+fx))} + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} \end{aligned}$$

[Out] $-11/96*I*d*x/a^3/f-1/16*d*x^2/a^3+1/8*x*(d*x+c)/a^3+1/36*d/f^2/(a+I*a*tan(f*x+e))^3+1/6*I*(d*x+c)/f/(a+I*a*tan(f*x+e))^3+5/96*d/a/f^2/(a+I*a*tan(f*x+e))^2+1/8*I*(d*x+c)/a/f/(a+I*a*tan(f*x+e))^2+11/96*d/f^2/(a^3+I*a^3*tan(f*x+e))+1/8*I*(d*x+c)/f/(a^3+I*a^3*tan(f*x+e))$

Rubi [A] (verified)

Time = 0.28 (sec), antiderivative size = 209, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.143, Rules used

$$= \{3560, 8, 3811\}$$

$$\begin{aligned} \int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx = & \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} + \frac{x(c+dx)}{8a^3} \\ & + \frac{11d}{96f^2(a^3+ia^3\tan(e+fx))} - \frac{11idx}{96a^3f} - \frac{dx^2}{16a^3} \\ & + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\ & + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} \end{aligned}$$

[In] `Int[(c + d*x)/(a + I*a*Tan[e + f*x])^3, x]`

[Out] $\left(\frac{(-11I)}{96}d^2x\right) - \frac{(d^2x^2)(16a^3)}{(a^3+ia^3\tan(e+fx))^3} + \frac{(x(c+d*x))(8a^3)}{(a^3+ia^3\tan(e+fx))^3} + \frac{d(36f^2(a^3+ia^3\tan(e+fx))^2)}{(a^3+ia^3\tan(e+fx))^3} + \frac{(I/6)(c+d*x)(f(a+I*a*Tan[e+f*x]))}{(a^3+ia^3\tan(e+fx))^3} + \frac{(5d)(96a^2f^2(a^3+ia^3\tan(e+fx))^2)}{(a^3+ia^3\tan(e+fx))^3} + \frac{(I/8)(c+d*x)(a*f^2(a+I*a*Tan[e+f*x]))}{(a^3+ia^3\tan(e+fx))^3} + \frac{(I/8)(c+d*x)(f^2(a^3+I*a^3*Tan[e+f*x]))}{(a^3+ia^3\tan(e+fx))^3}$

Rule 8

`Int[a_, x_Symbol] :> Simplify[a*x, x] /; FreeQ[a, x]`

Rule 3560

`Int[((a_) + (b_)*tan((c_) + (d_)*(x_)))^(n_), x_Symbol] :> Simplify[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]`

Rule 3811

`Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan((e_) + (f_)*(x_)))^(n_), x_Symbol] :> With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Dist[(c + d*x)^m, u, x] - Dist[d*m, Int[Dist[(c + d*x)^(m - 1), u, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]]`

Rubi steps

$$\begin{aligned} \text{integral} = & \frac{x(c+dx)}{8a^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\ & + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} - d \int \left(\frac{x}{8a^3} + \frac{i}{6f(a+ia\tan(e+fx))^3} \right. \\ & \quad \left. + \frac{i}{8af(a+ia\tan(e+fx))^2} + \frac{i}{8f(a^3+ia^3\tan(e+fx))} \right) dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\
&\quad + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} - \frac{(id)\int \frac{1}{a^3+ia^3\tan(e+fx)} dx}{8f} \\
&\quad - \frac{(id)\int \frac{1}{(a+ia\tan(e+fx))^3} dx}{6f} - \frac{(id)\int \frac{1}{(a+ia\tan(e+fx))^2} dx}{8af} \\
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\
&\quad + \frac{d}{32af^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\
&\quad + \frac{d}{16f^2(a^3+ia^3\tan(e+fx))} + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} \\
&\quad - \frac{(id)\int 1 dx}{16a^3f} - \frac{(id)\int \frac{1}{a+ia\tan(e+fx)} dx}{16a^2f} - \frac{(id)\int \frac{1}{(a+ia\tan(e+fx))^2} dx}{12af} \\
&= -\frac{idx}{16a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} \\
&\quad + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} \\
&\quad + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} + \frac{3d}{32f^2(a^3+ia^3\tan(e+fx))} \\
&\quad + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} - \frac{(id)\int 1 dx}{32a^3f} - \frac{(id)\int \frac{1}{a+ia\tan(e+fx)} dx}{24a^2f} \\
&= -\frac{3idx}{32a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\
&\quad + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\
&\quad + \frac{11d}{96f^2(a^3+ia^3\tan(e+fx))} + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))} - \frac{(id)\int 1 dx}{48a^3f} \\
&= -\frac{11idx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} + \frac{d}{36f^2(a+ia\tan(e+fx))^3} + \frac{i(c+dx)}{6f(a+ia\tan(e+fx))^3} \\
&\quad + \frac{5d}{96af^2(a+ia\tan(e+fx))^2} + \frac{i(c+dx)}{8af(a+ia\tan(e+fx))^2} \\
&\quad + \frac{11d}{96f^2(a^3+ia^3\tan(e+fx))} + \frac{i(c+dx)}{8f(a^3+ia^3\tan(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.52 (sec) , antiderivative size = 205, normalized size of antiderivative = 0.98

$$\int \frac{c + dx}{(a + i a \tan(e + fx))^3} dx \\ = \frac{i \sec^3(e + fx) (27(12icf + d(5 + 12ifx)) \cos(e + fx) + 4(6cf(i + 6fx) + d(1 + 6ifx + 18f^2x^2)) \cos(3(e + fx)))}{(a^3 f^2 (-I + \tan(e + fx)))^3}$$

[In] `Integrate[(c + d*x)/(a + I*a*Tan[e + f*x])^3, x]`

[Out]
$$\begin{aligned} & ((I/1152)*\text{Sec}[e + f*x]^3*(27*((12*I)*c*f + d*(5 + (12*I)*f*x))*\text{Cos}[e + f*x] \\ & + 4*(6*c*f*(I + 6*f*x) + d*(1 + (6*I)*f*x + 18*f^2*x^2))*\text{Cos}[3*(e + f*x)] \\ & + (81*I)*d*\text{Sin}[e + f*x] - 108*c*f*\text{Sin}[e + f*x] - 108*d*f*x*\text{Sin}[e + f*x] - (\\ & 4*I)*d*\text{Sin}[3*(e + f*x)] + 24*c*f*\text{Sin}[3*(e + f*x)] + 24*d*f*x*\text{Sin}[3*(e + f*x)] \\ & + (144*I)*c*f^2*x*\text{Sin}[3*(e + f*x)] + (72*I)*d*f^2*x^2*\text{Sin}[3*(e + f*x)]) \\ & /(a^3 f^2 (-I + \tan(e + f*x)))^3) \end{aligned}$$

Maple [A] (verified)

Time = 0.88 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.55

method	result	size
risch	$\frac{dx^2}{16a^3} + \frac{cx}{8a^3} + \frac{3i(2dfx+2cf-id)e^{-2i(fx+e)}}{32a^3f^2} + \frac{3i(4dfx+4cf-id)e^{-4i(fx+e)}}{128a^3f^2} + \frac{i(6dfx+6cf-id)e^{-6i(fx+e)}}{288a^3f^2}$	114

[In] `int((d*x+c)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/16*d*x^2/a^3+1/8*a^3*c*x+3/32*I*(2*d*f*x-I*d+2*c*f)/a^3/f^2*exp(-2*I*(f*x+e))+3/128*I*(4*d*f*x-I*d+4*c*f)/a^3/f^2*exp(-4*I*(f*x+e))+1/288*I*(6*d*f*x-I*d+6*c*f)/a^3/f^2*exp(-6*I*(f*x+e)) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.50

$$\int \frac{c + dx}{(a + i a \tan(e + fx))^3} dx \\ = \frac{(24i dfx + 24i cf + 72(df^2 x^2 + 2cf^2 x)e^{(6ifx+6ie)} - 108(-2i dfx - 2i cf - d)e^{(4ifx+4ie)} - 27(-4i dfx - 1152a^3f^2)e^{(2ifx+2ie)})}{1152a^3f^2}$$

[In] `integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

[Out] $\frac{1}{1152} \left(24I^*d^*f*x + 24I^*c^*f + 72(d*f^2*x^2 + 2*c*f^2*x)*e^{(6I^*f*x + 6I^*e)} - 108(-2I^*d^*f*x - 2I^*c^*f - d)*e^{(4I^*f*x + 4I^*e)} - 27(-4I^*d^*f*x - 4I^*c^*f - d)*e^{(2I^*f*x + 2I^*e)} + 4d)*e^{(-6I^*f*x - 6I^*e)} \right) / (a^3*f^2)$

Sympy [A] (verification not implemented)

Time = 0.34 (sec), antiderivative size = 311, normalized size of antiderivative = 1.49

$$\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx$$

$$= \begin{cases} \frac{((24576ia^6cf^5e^{6ie}+24576ia^6df^5xe^{6ie}+4096a^6df^4e^{6ie})e^{-6ifx}+(110592ia^6cf^5e^{8ie}+110592ia^6df^5xe^{8ie}+27648a^6df^4e^{8ie})e^{-4ifx}+(221184ia^6cf^5xe^{10ie}+221184ia^6df^5xe^{10ie}+110592a^6df^4e^{10ie})e^{-2ifx}+(1179648a^9f^6)e^{-12ifx})}{1179648a^9f^6} \\ \frac{x^2\cdot(3de^{4ie}+3de^{2ie}+d)e^{-6ie}}{16a^3} + \frac{x(3ce^{4ie}+3ce^{2ie}+c)e^{-6ie}}{8a^3} \\ + \frac{cx}{8a^3} + \frac{dx^2}{16a^3} \end{cases}$$

[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))**3,x)

[Out] Piecewise(((24576*I*a**6*c*f**5*exp(6*I^*e) + 24576*I*a**6*d*f**5*x*exp(6*I^*e) + 4096*a**6*d*f**4*exp(6*I^*e))*exp(-6*I^*f*x) + (110592*I*a**6*c*f**5*exp(8*I^*e) + 110592*I*a**6*d*f**5*x*exp(8*I^*e) + 27648*a**6*d*f**4*exp(8*I^*e))*exp(-4*I^*f*x) + (221184*I*a**6*c*f**5*exp(10*I^*e) + 221184*I*a**6*d*f**5*x*exp(10*I^*e))*exp(-2*I^*f*x))*exp(-12*I^*e)/(1179648*a**9*f**6), Ne(a**9*f**6*exp(12*I^*e), 0)), (x**2*(3*d*exp(4*I^*e) + 3*d*exp(2*I^*e) + d)*exp(-6*I^*e)/(16*a**3) + x*(3*c*exp(4*I^*e) + 3*c*exp(2*I^*e) + c)*exp(-6*I^*e)/(8*a**3), True)) + c*x/(8*a**3) + d*x**2/(16*a**3)

Maxima [F(-2)]

Exception generated.

$$\int \frac{c+dx}{(a+ia\tan(e+fx))^3} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))**3,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: expt: undefined: 0 to a negative exponent.

Giac [A] (verification not implemented)

none

Time = 0.56 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.68

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{(72 df^2 x^2 e^{(6i f x + 6i e)} + 144 c f^2 x e^{(6i f x + 6i e)} + 216 i d f x e^{(4i f x + 4i e)} + 108 i d f x e^{(2i f x + 2i e)} + 24 i d f x + 216 i c f e^{(2i f x + 2i e)})}{1152 a^3 f^2}$$

```
[In] integrate((d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
[Out] 1/1152*(72*d*f^2*x^2*e^(6*I*f*x + 6*I*e) + 144*c*f^2*x*e^(6*I*f*x + 6*I*e)
+ 216*I*d*f*x*e^(4*I*f*x + 4*I*e) + 108*I*d*f*x*e^(2*I*f*x + 2*I*e) + 24*I*
d*f*x + 216*I*c*f*e^(4*I*f*x + 4*I*e) + 108*I*c*f*e^(2*I*f*x + 2*I*e) + 24*
I*c*f + 108*d*e^(4*I*f*x + 4*I*e) + 27*d*e^(2*I*f*x + 2*I*e) + 4*d)*e^(-6*I
*f*x - 6*I*e)/(a^3*f^2)
```

Mupad [B] (verification not implemented)

Time = 3.94 (sec) , antiderivative size = 146, normalized size of antiderivative = 0.70

$$\int \frac{c + dx}{(a + ia \tan(e + fx))^3} dx = \frac{d x^2}{16 a^3} - e^{-e^{4i-fx} 4i} \left(\frac{(-12 c f + d 3i) 1i}{128 a^3 f^2} - \frac{d x 3i}{32 a^3 f} \right) - e^{-e^{6i-fx} 6i} \left(\frac{(-6 c f + d 1i) 1i}{288 a^3 f^2} - \frac{d x 1i}{48 a^3 f} \right) - e^{-e^{2i-fx} 2i} \left(\frac{(-6 c f + d 3i) 1i}{32 a^3 f^2} - \frac{d x 3i}{16 a^3 f} \right) + \frac{c x}{8 a^3}$$

```
[In] int((c + d*x)/(a + a*tan(e + f*x)*1i)^3,x)
[Out] (d*x^2)/(16*a^3) - exp(- e*4i - f*x*4i)*(((d*3i - 12*c*f)*1i)/(128*a^3*f^2)
- (d*x*3i)/(32*a^3*f)) - exp(- e*6i - f*x*6i)*(((d*1i - 6*c*f)*1i)/(288*a^
3*f^2) - (d*x*1i)/(48*a^3*f)) - exp(- e*2i - f*x*2i)*(((d*3i - 6*c*f)*1i)/(
32*a^3*f^2) - (d*x*3i)/(16*a^3*f)) + (c*x)/(8*a^3)
```

3.32 $\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx$

Optimal result	210
Rubi [A] (verified)	211
Mathematica [A] (verified)	215
Maple [A] (verified)	216
Fricas [A] (verification not implemented)	216
Sympy [F]	216
Maxima [A] (verification not implemented)	217
Giac [A] (verification not implemented)	217
Mupad [F(-1)]	218

Optimal result

Integrand size = 23, antiderivative size = 449

$$\begin{aligned} \int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx = & \frac{3 \cos\left(2e - \frac{2cf}{d}\right) \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ & + \frac{3 \cos\left(4e - \frac{4cf}{d}\right) \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ & + \frac{\cos\left(6e - \frac{6cf}{d}\right) \text{CosIntegral}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} \\ & + \frac{\log(c+dx)}{8a^3d} \\ & - \frac{i \text{CosIntegral}\left(\frac{6cf}{d} + 6fx\right) \sin\left(6e - \frac{6cf}{d}\right)}{8a^3d} \\ & - \frac{3i \text{CosIntegral}\left(\frac{4cf}{d} + 4fx\right) \sin\left(4e - \frac{4cf}{d}\right)}{8a^3d} \\ & - \frac{3i \text{CosIntegral}\left(\frac{2cf}{d} + 2fx\right) \sin\left(2e - \frac{2cf}{d}\right)}{8a^3d} \\ & - \frac{3i \cos\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ & - \frac{3 \sin\left(2e - \frac{2cf}{d}\right) \text{Si}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\ & - \frac{3i \cos\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ & - \frac{3 \sin\left(4e - \frac{4cf}{d}\right) \text{Si}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\ & - \frac{i \cos\left(6e - \frac{6cf}{d}\right) \text{Si}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} \\ & - \frac{\sin\left(6e - \frac{6cf}{d}\right) \text{Si}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} \end{aligned}$$

[Out]
$$\begin{aligned} & 1/8*\text{Ci}(6*c*f/d+6*f*x)*\cos(-6*e+6*c*f/d)/a^3/d+3/8*\text{Ci}(4*c*f/d+4*f*x)*\cos(-4*e+4*c*f/d)/a^3/d+3/8*\text{Ci}(2*c*f/d+2*f*x)*\cos(-2*e+2*c*f/d)/a^3/d+1/8*\ln(d*x+c)/a^3/d-3/8*I*\cos(-2*e+2*c*f/d)*\text{Si}(2*c*f/d+2*f*x)/a^3/d-3/8*I*\cos(-4*e+4*c*f/d)*\text{Si}(4*c*f/d+4*f*x)/a^3/d-1/8*I*\cos(-6*e+6*c*f/d)*\text{Si}(6*c*f/d+6*f*x)/a^3/d+1/8*I*\text{Ci}(6*c*f/d+6*f*x)*\sin(-6*e+6*c*f/d)/a^3/d+1/8*\text{Si}(6*c*f/d+6*f*x)*\sin(-6*e+6*c*f/d)/a^3/d+3/8*I*\text{Ci}(4*c*f/d+4*f*x)*\sin(-4*e+4*c*f/d)/a^3/d+3/8*\text{Si}(4*c*f/d+4*f*x)*\sin(-4*e+4*c*f/d)/a^3/d+3/8*I*\text{Ci}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a^3/d+3/8*\text{Si}(2*c*f/d+2*f*x)*\sin(-2*e+2*c*f/d)/a^3/d \end{aligned}$$

Rubi [A] (verified)

Time = 2.60 (sec), antiderivative size = 449, normalized size of antiderivative = 1.00, number of steps used = 53, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.304, Rules used = {3809, 3384, 3380, 3383, 3393, 4491, 4513}

$$\begin{aligned} \int \frac{1}{(c+dx)(a+i a \tan(e+fx))^3} dx = & -\frac{3 i \operatorname{CosIntegral}\left(2 x f+\frac{2 c f}{d}\right) \sin \left(2 e-\frac{2 c f}{d}\right)}{8 a^3 d} \\ & -\frac{i \operatorname{CosIntegral}\left(6 x f+\frac{6 c f}{d}\right) \sin \left(6 e-\frac{6 c f}{d}\right)}{8 a^3 d} \\ & -\frac{3 i \operatorname{CosIntegral}\left(4 x f+\frac{4 c f}{d}\right) \sin \left(4 e-\frac{4 c f}{d}\right)}{8 a^3 d} \\ & +\frac{3 \operatorname{CosIntegral}\left(2 x f+\frac{2 c f}{d}\right) \cos \left(2 e-\frac{2 c f}{d}\right)}{8 a^3 d} \\ & +\frac{3 \operatorname{CosIntegral}\left(4 x f+\frac{4 c f}{d}\right) \cos \left(4 e-\frac{4 c f}{d}\right)}{8 a^3 d} \\ & +\frac{\operatorname{CosIntegral}\left(6 x f+\frac{6 c f}{d}\right) \cos \left(6 e-\frac{6 c f}{d}\right)}{8 a^3 d} \\ & -\frac{3 \sin \left(2 e-\frac{2 c f}{d}\right) \operatorname{Si}\left(2 x f+\frac{2 c f}{d}\right)}{8 a^3 d} \\ & -\frac{3 \sin \left(4 e-\frac{4 c f}{d}\right) \operatorname{Si}\left(4 x f+\frac{4 c f}{d}\right)}{8 a^3 d} \\ & -\frac{\sin \left(6 e-\frac{6 c f}{d}\right) \operatorname{Si}\left(6 x f+\frac{6 c f}{d}\right)}{8 a^3 d} \\ & -\frac{3 i \cos \left(2 e-\frac{2 c f}{d}\right) \operatorname{Si}\left(2 x f+\frac{2 c f}{d}\right)}{8 a^3 d} \\ & -\frac{3 i \cos \left(4 e-\frac{4 c f}{d}\right) \operatorname{Si}\left(4 x f+\frac{4 c f}{d}\right)}{8 a^3 d} \\ & -\frac{i \cos \left(6 e-\frac{6 c f}{d}\right) \operatorname{Si}\left(6 x f+\frac{6 c f}{d}\right)}{8 a^3 d}+\frac{\log (c+d x)}{8 a^3 d} \end{aligned}$$

[In] $\text{Int}[1/((c+d*x)*(a+I*a*Tan[e+f*x])^3),x]$

[Out]
$$(3*\text{Cos}[2*e - (2*c*f)/d]*\text{CosIntegral}[(2*c*f)/d + 2*f*x])/({8*a^3*d}) + (3*\text{Cos}[4*e - (4*c*f)/d]*\text{CosIntegral}[(4*c*f)/d + 4*f*x])/({8*a^3*d}) + (\text{Cos}[6*e - (6*c*f)/d]*\text{CosIntegral}[(6*c*f)/d + 6*f*x])/({8*a^3*d})$$

```
c*f)/d]*CosIntegral[(6*c*f)/d + 6*f*x])/(8*a^3*d) + Log[c + d*x]/(8*a^3*d)
- ((I/8)*CosIntegral[(6*c*f)/d + 6*f*x]*Sin[6*e - (6*c*f)/d])/(a^3*d) - (((3*I)/8)*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d])/(a^3*d) - (((3*I)/8)*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/(a^3*d) - (((3*I)/8)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^3*d) - (3*Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(8*a^3*d) - (((3*I)/8)*Cos[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^3*d) - (3*Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(8*a^3*d) - ((I/8)*Cos[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(a^3*d) - (Sin[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(8*a^3*d)
```

Rule 3380

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[SinIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]
```

Rule 3383

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_.))^m_*sin[(e_.) + (f_.)*(x_.)]^n, x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_.))^m_*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rule 4491

```
Int[Cos[(a_.) + (b_.)*(x_.)]^p_*((c_.) + (d_.)*(x_.))^m_*Sin[(a_.) + (b_.)*(x_.)]^n, x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[a + b*x]^n*Cos[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IG
```

$tQ[p, 0]$

Rule 4513

```
Int[((e_) + (f_)*(x_))^(m_)*Sin[(a_) + (b_)*(x_)]^(p_)*Sin[(c_) + (d_)*(x_)]^(q_), x_Symbol] :> Int[ExpandTrigReduce[(e + f*x)^m, Sin[a + b*x]^p*Sin[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{8a^3(c+dx)} + \frac{3\cos(2e+2fx)}{8a^3(c+dx)} + \frac{3\cos^2(2e+2fx)}{8a^3(c+dx)} + \frac{\cos^3(2e+2fx)}{8a^3(c+dx)} \right. \\
 &\quad - \frac{3i\sin(2e+2fx)}{8a^3(c+dx)} - \frac{3i\cos^2(2e+2fx)\sin(2e+2fx)}{8a^3(c+dx)} - \frac{3\sin^2(2e+2fx)}{8a^3(c+dx)} \\
 &\quad \left. + \frac{i\sin^3(2e+2fx)}{8a^3(c+dx)} - \frac{3i\sin(4e+4fx)}{8a^3(c+dx)} - \frac{3\sin(2e+2fx)\sin(4e+4fx)}{16a^3(c+dx)} \right) dx \\
 &= \frac{\log(c+dx)}{8a^3d} + \frac{i \int \frac{\sin^3(2e+2fx)}{c+dx} dx}{8a^3} - \frac{(3i) \int \frac{\sin(2e+2fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{(3i) \int \frac{\cos^2(2e+2fx)\sin(2e+2fx)}{c+dx} dx}{8a^3} - \frac{(3i) \int \frac{\sin(4e+4fx)}{c+dx} dx}{8a^3} \\
 &\quad + \frac{\int \frac{\cos^3(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sin(2e+2fx)\sin(4e+4fx)}{c+dx} dx}{16a^3} \\
 &\quad + \frac{3 \int \frac{\cos(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\cos^2(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sin^2(2e+2fx)}{c+dx} dx}{8a^3} \\
 &= \frac{\log(c+dx)}{8a^3d} + \frac{i \int \left(\frac{3\sin(2e+2fx)}{4(c+dx)} - \frac{\sin(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} \\
 &\quad - \frac{(3i) \int \left(\frac{\sin(2e+2fx)}{4(c+dx)} + \frac{\sin(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} + \frac{\int \left(\frac{3\cos(2e+2fx)}{4(c+dx)} + \frac{\cos(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} \\
 &\quad - \frac{3 \int \left(\frac{\cos(2e+2fx)}{2(c+dx)} - \frac{\cos(6e+6fx)}{2(c+dx)} \right) dx}{16a^3} - \frac{3 \int \left(\frac{1}{2(c+dx)} - \frac{\cos(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} \\
 &\quad + \frac{3 \int \left(\frac{1}{2(c+dx)} + \frac{\cos(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} - \frac{(3i \cos(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{(3i \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} \\
 &\quad + \frac{(3 \cos(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} - \frac{(3i \sin(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{(3i \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} - \frac{(3 \sin(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{8a^3d} \\
&\quad - \frac{3i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^3d} - \frac{i \int \frac{\sin(6e+6fx)}{c+dx} dx}{32a^3} - \frac{(3i) \int \frac{\sin(6e+6fx)}{c+dx} dx}{32a^3} \\
&\quad + \frac{\int \frac{\cos(6e+6fx)}{c+dx} dx}{32a^3} + \frac{3 \int \frac{\cos(6e+6fx)}{c+dx} dx}{32a^3} + 2 \frac{3 \int \frac{\cos(4e+4fx)}{c+dx} dx}{16a^3} \\
&= \frac{3 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{8a^3d} - \frac{3i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^3d} \\
&\quad - \frac{(i \cos(6e - \frac{6cf}{d})) \int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} - \frac{(3i \cos(6e - \frac{6cf}{d})) \int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} \\
&\quad + \frac{\cos(6e - \frac{6cf}{d}) \int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} + \frac{(3 \cos(6e - \frac{6cf}{d})) \int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} \\
&\quad - \frac{(i \sin(6e - \frac{6cf}{d})) \int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} - \frac{(3i \sin(6e - \frac{6cf}{d})) \int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} \\
&\quad - \frac{\sin(6e - \frac{6cf}{d}) \int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} - \frac{(3 \sin(6e - \frac{6cf}{d})) \int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{32a^3} \\
&\quad + 2 \left(\frac{(3 \cos(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d}+4fx)}{c+dx} dx}{16a^3} - \frac{(3 \sin(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d}+4fx)}{c+dx} dx}{16a^3} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{3 \cos(2e - \frac{2cf}{d}) \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{\cos(6e - \frac{6cf}{d}) \operatorname{CosIntegral}(\frac{6cf}{d} + 6fx)}{8a^3d} \\
&\quad + \frac{\log(c + dx)}{8a^3d} - \frac{i \operatorname{CosIntegral}(\frac{6cf}{d} + 6fx) \sin(6e - \frac{6cf}{d})}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx) \sin(4e - \frac{4cf}{d})}{8a^3d} \\
&\quad - \frac{3i \operatorname{CosIntegral}(\frac{2cf}{d} + 2fx) \sin(2e - \frac{2cf}{d})}{8a^3d} - \frac{3i \cos(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3 \sin(2e - \frac{2cf}{d}) \operatorname{Si}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3i \cos(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{8a^3d} \\
&\quad + 2 \left(\frac{3 \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{16a^3d} - \frac{3 \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{16a^3d} \right) \\
&\quad - \frac{i \cos(6e - \frac{6cf}{d}) \operatorname{Si}(\frac{6cf}{d} + 6fx)}{8a^3d} - \frac{\sin(6e - \frac{6cf}{d}) \operatorname{Si}(\frac{6cf}{d} + 6fx)}{8a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.14 (sec), antiderivative size = 336, normalized size of antiderivative = 0.75

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx$$

$$\frac{\sec^3(e + fx)(\cos(fx) + i \sin(fx))^3 \left(\cos(3e) \log(f(c + dx)) + i \log(f(c + dx)) \sin(3e) + (\cos(e - \frac{4cf}{d}) - \right.}{= \left. \frac{3 \cos(4e - \frac{4cf}{d}) \operatorname{CosIntegral}(\frac{4cf}{d} + 4fx)}{16a^3d} - \frac{3 \sin(4e - \frac{4cf}{d}) \operatorname{Si}(\frac{4cf}{d} + 4fx)}{16a^3d} \right)$$

```

[In] Integrate[1/((c + d*x)*(a + I*a*Tan[e + f*x])^3), x]
[Out] (Sec[e + f*x]^3*(Cos[f*x] + I*Sin[f*x])^3*(Cos[3*e]*Log[f*(c + d*x)] + I*Log[f*(c + d*x)]*Sin[3*e] + (Cos[e - (4*c*f)/d] - I*Sin[e - (4*c*f)/d])*(
 3*CosIntegral[(4*f*(c + d*x))/d] + Cos[2*e - (2*c*f)/d]*CosIntegral[(6*f*(c + d*x))/d] + 3*CosIntegral[(2*f*(c + d*x))/d]*(Cos[2*e - (2*c*f)/d] + I*Sin[2*e - (2*c*f)/d]) - I*CosIntegral[(6*f*(c + d*x))/d]*Sin[2*e - (2*c*f)/d] - (3*I)*Cos[2*e - (2*c*f)/d]*SinIntegral[(2*f*(c + d*x))/d] + 3*Sin[2*e - (2*c*f)/d]*SinIntegral[(2*f*(c + d*x))/d] - (3*I)*SinIntegral[(4*f*(c + d*x))/d] - I*Cos[2*e - (2*c*f)/d]*SinIntegral[(6*f*(c + d*x))/d] - Sin[2*e - (2*c*f)/d]*SinIntegral[(6*f*(c + d*x))/d]))/(8*d*(a + I*a*Tan[e + f*x])^3)

```

Maple [A] (verified)

Time = 0.93 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.36

method	result
risch	$\frac{\ln(dx+c)}{8a^3d} - \frac{e^{\frac{6i(cf-de)}{d}} \text{Ei}_1\left(6ifx+6ie+\frac{6i(cf-de)}{d}\right)}{8a^3d} - \frac{3e^{\frac{4i(cf-de)}{d}} \text{Ei}_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{8a^3d} - \frac{3e^{\frac{2i(cf-de)}{d}} \text{Ei}_1\left(2ifx+2ie+\frac{2i(cf-de)}{d}\right)}{8a^3d}$

[In] `int(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & \frac{1}{8} \ln(d*x+c)/a^3/d - \frac{1}{8} a^3/d * \exp(6*I*(c*f-d*e)/d) * \text{Ei}(1, 6*I*f*x + 6*I*e + 6*I*(c*f-d*e)/d) \\ & - \frac{3}{8} a^3/d * \exp(4*I*(c*f-d*e)/d) * \text{Ei}(1, 4*I*f*x + 4*I*e + 4*I*(c*f-d*e)/d) \\ & - \frac{3}{8} a^3/d * \exp(2*I*(c*f-d*e)/d) * \text{Ei}(1, 2*I*f*x + 2*I*e + 2*I*(c*f-d*e)/d) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 117, normalized size of antiderivative = 0.26

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ &= \frac{3 \text{Ei}\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + 3 \text{Ei}\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + \text{Ei}\left(-\frac{6(i dfx + i cf)}{d}\right) e^{\left(-\frac{6(i de - i cf)}{d}\right)} + \log((d*x + c)/d))}{8 a^3 d} \end{aligned}$$

[In] `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & \frac{1}{8} (3 \text{Ei}(-2*(I*d*f*x + I*c*f)/d) * e^{-(-2*(I*d*e - I*c*f)/d)} + 3 \text{Ei}(-4*(I*d*f*x + I*c*f)/d) * e^{-(-4*(I*d*e - I*c*f)/d)} + \text{Ei}(-6*(I*d*f*x + I*c*f)/d) * e^{-(-6*(I*d*e - I*c*f)/d)} + \log((d*x + c)/d)) / (a^3*d) \end{aligned}$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx \\ &= \frac{i \int \frac{1}{c \tan^3(e + fx) - 3ic \tan^2(e + fx) - 3c \tan(e + fx) + ic + dx \tan^3(e + fx) - 3idx \tan^2(e + fx) - 3dx \tan(e + fx) + idx} dx}{a^3} \end{aligned}$$

[In] `integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))**3,x)`

[Out]
$$\begin{aligned} & I * \text{Integral}(1/(c*\tan(e + f*x)**3 - 3*I*c*\tan(e + f*x)**2 - 3*c*\tan(e + f*x) + I*c + d*x*\tan(e + f*x)**3 - 3*I*d*x*\tan(e + f*x)**2 - 3*d*x*\tan(e + f*x) + I*d*x), x)/a**3 \end{aligned}$$

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 276, normalized size of antiderivative = 0.61

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx =$$

$$3 f \cos\left(-\frac{2(de-cf)}{d}\right) E_1\left(-\frac{2(-i(fx+e)d+i(de-i)cf)}{d}\right) + 3 f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+i(de-i)cf)}{d}\right) + f \cos\left(-\frac{4(de-cf)}{d}\right) E_1\left(-\frac{4(-i(fx+e)d+i(de-i)cf)}{d}\right)$$

```
[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] -1/8*(3*f*cos(-2*(d*e - c*f)/d)*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*f*cos(-4*(d*e - c*f)/d)*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f*cos(-6*(d*e - c*f)/d)*exp_integral_e(1, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*I*f*exp_integral_e(1, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + 3*I*f*exp_integral_e(1, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + I*f*exp_integral_e(1, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-6*(d*e - c*f)/d) - f*log((f*x + e)*d - d*e + c*f))/(a^3*d*f)
```

Giac [A] (verification not implemented)

none

Time = 0.58 (sec) , antiderivative size = 810, normalized size of antiderivative = 1.80

$$\int \frac{1}{(c+dx)(a+ia\tan(e+fx))^3} dx = \text{Too large to display}$$

```
[In] integrate(1/(d*x+c)/(a+I*a*tan(f*x+e))^(3,x), algorithm="giac")
```

```
[Out] 1/8*(3*cos(2*e)^2*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d) + cos(2*e)^3*log(d*x + c) + 6*I*cos(2*e)*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e) + 3*I*cos(2*e)^2*log(d*x + c)*sin(2*e) - 3*cos(2*c*f/d)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)^2 - 3*cos(2*e)*log(d*x + c)*sin(2*e)^2 - I*log(d*x + c)*sin(2*e)^3 + 3*I*cos(2*e)^2*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*c*f/d) - 6*cos(2*e)*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)*sin(2*c*f/d) - 3*I*cos_integral(-2*(d*f*x + c*f)/d)*sin(2*e)^2*sin(2*c*f/d) - 3*I*cos(2*e)^2*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 6*cos(2*e)*cos(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 3*I*cos(2*c*f/d)*sin(2*e)^2*sin_integral(2*(d*f*x + c*f)/d) + 3*cos(2*e)^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 6*I*cos(2*e)*sin(2*e)*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) - 3*sin(2*e)^2*sin(2*c*f/d)*sin_integral(2*(d*f*x + c*f)/d) + 3*cos(2*e)*cos(4*c*f/d)*cos_integral(-4*(d*f*x + c*f)/d) + 3*I*cos(4*c*f/d)
```

```

)*cos_integral(-4*(d*f*x + c*f)/d)*sin(2*e) + 3*I*cos(2*e)*cos_integral(-4*
(d*f*x + c*f)/d)*sin(4*c*f/d) - 3*cos_integral(-4*(d*f*x + c*f)/d)*sin(2*e)
*sin(4*c*f/d) - 3*I*cos(2*e)*cos(4*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) +
3*cos(4*c*f/d)*sin(2*e)*sin_integral(4*(d*f*x + c*f)/d) + 3*cos(2*e)*sin(4
*c*f/d)*sin_integral(4*(d*f*x + c*f)/d) + 3*I*sin(2*e)*sin(4*c*f/d)*sin_int
egral(4*(d*f*x + c*f)/d) + cos(6*c*f/d)*cos_integral(-6*(d*f*x + c*f)/d) +
I*cos_integral(-6*(d*f*x + c*f)/d)*sin(6*c*f/d) - I*cos(6*c*f/d)*sin_integr
al(6*(d*f*x + c*f)/d) + sin(6*c*f/d)*sin_integral(6*(d*f*x + c*f)/d))/(a^3*
d*cos(2*e)^3 + 3*I*a^3*d*cos(2*e)^2*sin(2*e) - 3*a^3*d*cos(2*e)*sin(2*e)^2
- I*a^3*d*sin(2*e)^3)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + ia \tan(e + fx))^3} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^3 (c + dx)} dx$$

[In] `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)),x)`
[Out] `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)), x)`

3.33 $\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx$

Optimal result	220
Rubi [A] (verified)	221
Mathematica [A] (verified)	228
Maple [A] (verified)	229
Fricas [A] (verification not implemented)	229
Sympy [F]	230
Maxima [A] (verification not implemented)	230
Giac [B] (verification not implemented)	231
Mupad [F(-1)]	232

Optimal result

Integrand size = 23, antiderivative size = 712

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = & -\frac{1}{8a^3d(c+dx)} - \frac{9\cos(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} \\
 & - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} - \frac{3\cos(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3if\cos(2e-\frac{2cf}{d})\text{CosIntegral}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & - \frac{3if\cos(4e-\frac{4cf}{d})\text{CosIntegral}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & - \frac{3if\cos(6e-\frac{6cf}{d})\text{CosIntegral}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{6cf}{d}+6fx)\sin(6e-\frac{6cf}{d})}{4a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{4cf}{d}+4fx)\sin(4e-\frac{4cf}{d})}{2a^3d^2} \\
 & - \frac{3f\text{CosIntegral}(\frac{2cf}{d}+2fx)\sin(2e-\frac{2cf}{d})}{4a^3d^2} \\
 & + \frac{15i\sin(2e+2fx)}{32a^3d(c+dx)} \\
 & + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} \\
 & + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i\sin(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3f\cos(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & + \frac{3if\sin(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
 & - \frac{3f\cos(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & + \frac{3if\sin(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
 & - \frac{3f\cos(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
 & + \frac{3if\sin(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2}
 \end{aligned}$$

```
[Out] -1/8/a^3/d/(d*x+c)-1/8*I*sin(2*f*x+2*e)^3/a^3/d/(d*x+c)-3/4*I*f*Ci(6*c*f/d+
6*f*x)*cos(-6*e+6*c*f/d)/a^3/d^2+3/8*I*sin(4*f*x+4*e)/a^3/d/(d*x+c)-9/32*co
s(2*f*x+2*e)/a^3/d/(d*x+c)-3/8*cos(2*f*x+2*e)^2/a^3/d/(d*x+c)-1/8*cos(2*f*x+
2*e)^3/a^3/d/(d*x+c)-3/32*cos(6*f*x+6*e)/a^3/d/(d*x+c)-3/4*f*cos(-2*e+2*c*
```

$$\begin{aligned}
& f/d) * \text{Si}(2*c*f/d + 2*f*x)/a^3/d^2 - 3/2*f*\cos(-4*e + 4*c*f/d)*\text{Si}(4*c*f/d + 4*f*x)/a^3/d^2 - 3/4*f*\cos(-6*e + 6*c*f/d)*\text{Si}(6*c*f/d + 6*f*x)/a^3/d^2 + 3/4*f*\text{Ci}(6*c*f/d + 6*f*x)*\sin(-6*e + 6*c*f/d)/a^3/d^2 - 3/2*I*f*\text{Si}(4*c*f/d + 4*f*x)*\sin(-4*e + 4*c*f/d)/a^3/d^2 + 3/2*f*\text{Ci}(4*c*f/d + 4*f*x)*\sin(-4*e + 4*c*f/d)/a^3/d^2 - 3/4*I*f*\text{Si}(6*c*f/d + 6*f*x)*\sin(-6*e + 6*c*f/d)/a^3/d^2 + 3/4*f*\text{Ci}(2*c*f/d + 2*f*x)*\sin(-2*e + 2*c*f/d)/a^3/d^2 - 3/4*I*f*\text{Ci}(2*c*f/d + 2*f*x)*\cos(-2*e + 2*c*f/d)/a^3/d^2 + 3/32*I*\sin(6*f*x + 6*e)/a^3/d/(d*x + c) + 3/8*\sin(2*f*x + 2*e)^2/a^3/d/(d*x + c) + 15/32*I*\sin(2*f*x + 2*e)/a^3/d/(d*x + c) - 3/2*I*f*\text{Ci}(4*c*f/d + 4*f*x)*\cos(-4*e + 4*c*f/d)/a^3/d^2 - 3/4*I*f*\text{Si}(2*c*f/d + 2*f*x)*\sin(-2*e + 2*c*f/d)/a^3/d^2
\end{aligned}$$

Rubi [A] (verified)

Time = 2.12 (sec), antiderivative size = 712, normalized size of antiderivative = 1.00, number of steps used = 60, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$, Rules used

$$= \{3809, 3378, 3384, 3380, 3383, 3394, 12, 4491, 4513\}$$

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = & -\frac{3f \operatorname{CosIntegral}(6xf + \frac{6cf}{d}) \sin(6e - \frac{6cf}{d})}{4a^3 d^2} \\ & -\frac{3f \operatorname{CosIntegral}(4xf + \frac{4cf}{d}) \sin(4e - \frac{4cf}{d})}{2a^3 d^2} \\ & -\frac{3f \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \sin(2e - \frac{2cf}{d})}{4a^3 d^2} \\ & -\frac{3if \operatorname{CosIntegral}(2xf + \frac{2cf}{d}) \cos(2e - \frac{2cf}{d})}{4a^3 d^2} \\ & -\frac{3if \operatorname{CosIntegral}(4xf + \frac{4cf}{d}) \cos(4e - \frac{4cf}{d})}{2a^3 d^2} \\ & -\frac{3if \operatorname{CosIntegral}(6xf + \frac{6cf}{d}) \cos(6e - \frac{6cf}{d})}{4a^3 d^2} \\ & +\frac{3if \sin(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\ & +\frac{3if \sin(4e - \frac{4cf}{d}) \operatorname{Si}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\ & +\frac{3if \sin(6e - \frac{6cf}{d}) \operatorname{Si}(6xf + \frac{6cf}{d})}{4a^3 d^2} \\ & -\frac{3f \cos(2e - \frac{2cf}{d}) \operatorname{Si}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\ & -\frac{3f \cos(4e - \frac{4cf}{d}) \operatorname{Si}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\ & -\frac{3f \cos(6e - \frac{6cf}{d}) \operatorname{Si}(6xf + \frac{6cf}{d})}{4a^3 d^2} - \frac{i \sin^3(2e + 2fx)}{8a^3 d(c+dx)} \\ & +\frac{3 \sin^2(2e + 2fx)}{8a^3 d(c+dx)} + \frac{15i \sin(2e + 2fx)}{32a^3 d(c+dx)} \\ & +\frac{3i \sin(4e + 4fx)}{8a^3 d(c+dx)} + \frac{3i \sin(6e + 6fx)}{32a^3 d(c+dx)} \\ & -\frac{\cos^3(2e + 2fx)}{8a^3 d(c+dx)} - \frac{3 \cos^2(2e + 2fx)}{8a^3 d(c+dx)} \\ & -\frac{9 \cos(2e + 2fx)}{32a^3 d(c+dx)} - \frac{3 \cos(6e + 6fx)}{32a^3 d(c+dx)} - \frac{1}{8a^3 d(c+dx)} \end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3),x]

[Out]
$$\begin{aligned} & -1/8*1/(a^3*d*(c + d*x)) - (9*\operatorname{Cos}[2*e + 2*f*x])/({32*a^3*d*(c + d*x)}) - (3*\operatorname{Cos}[2*e + 2*f*x]^2)/({8*a^3*d*(c + d*x)}) - \operatorname{Cos}[2*e + 2*f*x]^3/({8*a^3*d*(c + d*x)}) - (3*\operatorname{Cos}[6*e + 6*f*x])/({32*a^3*d*(c + d*x)}) - (((3*I)/4)*f*\operatorname{Cos}[2*e - (2*c*f)/d]*\operatorname{CosIntegral}[(2*c*f)/d + 2*f*x])/({a^3*d^2}) - (((3*I)/2)*f*\operatorname{Cos}[4*e - (4*c*f)/d]*\operatorname{CosIntegral}[(4*c*f)/d + 4*f*x])/({a^3*d^2}) - (((3*I)/4)*f*\operatorname{Cos}[6*e - (6*c*f)/d]*\operatorname{CosIntegral}[(6*c*f)/d + 6*f*x])/({a^3*d^2}) - (3*f*\operatorname{CosIntegral}[(3*c*f)/d + 3*f*x])/({a^3*d^2}) \end{aligned}$$

```

1[((6*c*f)/d + 6*f*x]*Sin[6*e - (6*c*f)/d])/(4*a^3*d^2) - (3*f*CosIntegral[(4*c*f)/d + 4*f*x]*Sin[4*e - (4*c*f)/d])/(2*a^3*d^2) - (3*f*CosIntegral[(2*c*f)/d + 2*f*x]*Sin[2*e - (2*c*f)/d])/(4*a^3*d^2) + (((15*I)/32)*Sin[2*e + 2*f*x])/(a^3*d*(c + d*x)) + (3*Sin[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) - ((I/8)*Sin[2*e + 2*f*x]^3)/(a^3*d*(c + d*x)) + (((3*I)/8)*Sin[4*e + 4*f*x])/(a^3*d*(c + d*x)) + (((3*I)/32)*Sin[6*e + 6*f*x])/(a^3*d*(c + d*x)) - (3*f*CosIntegral[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(4*a^3*d^2) + (((3*I)/4)*f*Sin[2*e - (2*c*f)/d]*SinIntegral[(2*c*f)/d + 2*f*x])/(a^3*d^2) - (3*f*CosIntegral[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(2*a^3*d^2) + (((3*I)/2)*f*Sin[4*e - (4*c*f)/d]*SinIntegral[(4*c*f)/d + 4*f*x])/(a^3*d^2) - (3*f*CosIntegral[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(4*a^3*d^2) + (((3*I)/4)*f*Sin[6*e - (6*c*f)/d]*SinIntegral[(6*c*f)/d + 6*f*x])/(a^3*d^2)

```

Rule 12

```

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

```

Rule 3378

```

Int[((c_.) + (d_.)*(x_.))^(m_)*sin[(e_.) + (f_.)*(x_.)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]

```

Rule 3380

```

Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[SiIntegral[e + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*e - c*f, 0]

```

Rule 3383

```

Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CosIntegral[e - Pi/2 + f*x]/d, x] /; FreeQ[{c, d, e, f}, x] && EqQ[d*(e - Pi/2) - c*f, 0]

```

Rule 3384

```

Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[SiIntegral[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[SiIntegral[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

```

Rule 3394

```

Int[((c_.) + (d_.)*(x_.))^(m_)*sin[(e_.) + (f_.)*(x_.)]^(n_), x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1)))

```

```
))), Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n - 1), x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] && LtQ[m, -1]
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rule 4491

```
Int[Cos[(a_.) + (b_.)*(x_)]^(p_)*((c_.) + (d_.)*(x_))^(m_.)*Sin[(a_.) + (b_.)*(x_)]^(n_.), x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[a + b*x]^n*Cos[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

Rule 4513

```
Int[((e_.) + (f_.)*(x_))^(m_)*Sin[(a_.) + (b_.)*(x_)]^(p_)*Sin[(c_.) + (d_.)*(x_)]^(q_), x_Symbol] :> Int[ExpandTrigReduce[(e + f*x)^m, Sin[a + b*x]^p*Sin[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{8a^3(c+dx)^2} + \frac{3 \cos(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \cos^2(2e+2fx)}{8a^3(c+dx)^2} + \frac{\cos^3(2e+2fx)}{8a^3(c+dx)^2} \right. \\
&\quad - \frac{3i \sin(2e+2fx)}{8a^3(c+dx)^2} - \frac{3i \cos^2(2e+2fx) \sin(2e+2fx)}{8a^3(c+dx)^2} - \frac{3 \sin^2(2e+2fx)}{8a^3(c+dx)^2} \\
&\quad \left. + \frac{i \sin^3(2e+2fx)}{8a^3(c+dx)^2} - \frac{3i \sin(4e+4fx)}{8a^3(c+dx)^2} - \frac{3 \sin(2e+2fx) \sin(4e+4fx)}{16a^3(c+dx)^2} \right) dx \\
&= -\frac{1}{8a^3d(c+dx)} + \frac{i \int \frac{\sin^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{(3i) \int \frac{\sin(2e+2fx)}{(c+dx)^2} dx}{8a^3} \\
&\quad - \frac{(3i) \int \frac{\cos^2(2e+2fx) \sin(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{(3i) \int \frac{\sin(4e+4fx)}{(c+dx)^2} dx}{8a^3} \\
&\quad + \frac{\int \frac{\cos^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{3 \int \frac{\sin(2e+2fx) \sin(4e+4fx)}{(c+dx)^2} dx}{16a^3} \\
&\quad + \frac{3 \int \frac{\cos(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\cos^2(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{3 \int \frac{\sin^2(2e+2fx)}{(c+dx)^2} dx}{8a^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{3\cos(2e+2fx)}{8a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3i\sin(2e+2fx)}{8a^3d(c+dx)} + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} - \frac{(3i) \int \left(\frac{\sin(2e+2fx)}{4(c+dx)^2} + \frac{\sin(6e+6fx)}{4(c+dx)^2} \right) dx}{8a^3} \\
&\quad - \frac{3 \int \left(\frac{\cos(2e+2fx)}{2(c+dx)^2} - \frac{\cos(6e+6fx)}{2(c+dx)^2} \right) dx}{16a^3} - \frac{(3if) \int \frac{\cos(2e+2fx)}{c+dx} dx}{4a^3d} \\
&\quad + \frac{(3if) \int \left(\frac{\cos(2e+2fx)}{4(c+dx)} - \frac{\cos(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} - \frac{(3if) \int \frac{\cos(4e+4fx)}{c+dx} dx}{2a^3d} \\
&\quad - \frac{(3f) \int \frac{\sin(2e+2fx)}{c+dx} dx}{4a^3d} + \frac{(3f) \int \left(-\frac{\sin(2e+2fx)}{4(c+dx)} - \frac{\sin(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} \\
&\quad + \frac{(3f) \int -\frac{\sin(4e+4fx)}{2(c+dx)} dx}{2a^3d} - \frac{(3f) \int \frac{\sin(4e+4fx)}{2(c+dx)} dx}{2a^3d} \\
&= -\frac{1}{8a^3d(c+dx)} - \frac{3\cos(2e+2fx)}{8a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3i\sin(2e+2fx)}{8a^3d(c+dx)} + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} \\
&\quad - \frac{(3i) \int \frac{\sin(2e+2fx)}{(c+dx)^2} dx}{32a^3} - \frac{(3i) \int \frac{\sin(6e+6fx)}{(c+dx)^2} dx}{32a^3} - \frac{3 \int \frac{\cos(2e+2fx)}{(c+dx)^2} dx}{32a^3} + \frac{3 \int \frac{\cos(6e+6fx)}{(c+dx)^2} dx}{32a^3} \\
&\quad + \frac{(3if) \int \frac{\cos(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(3if) \int \frac{\cos(6e+6fx)}{c+dx} dx}{16a^3d} - \frac{(3f) \int \frac{\sin(2e+2fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f) \int \frac{\sin(6e+6fx)}{c+dx} dx}{16a^3d} - 2 \frac{(3f) \int \frac{\sin(4e+4fx)}{c+dx} dx}{4a^3d} - \frac{(3if \cos(4e - \frac{4cf}{d})) \int \frac{\cos(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} \\
&\quad - \frac{(3if \cos(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} - \frac{(3f \cos(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} \\
&\quad + \frac{(3if \sin(4e - \frac{4cf}{d})) \int \frac{\sin(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} + \frac{(3if \sin(2e - \frac{2cf}{d})) \int \frac{\sin(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} \\
&\quad - \frac{(3f \sin(2e - \frac{2cf}{d})) \int \frac{\cos(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9\cos(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3\cos(6e+6fx)}{32a^3d(c+dx)} - \frac{3if\cos(2e-\frac{2cf}{d})\text{CosIntegral}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad - \frac{3if\cos(4e-\frac{4cf}{d})\text{CosIntegral}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad - \frac{3f\text{CosIntegral}(\frac{2cf}{d}+2fx)\sin(2e-\frac{2cf}{d})}{4a^3d^2} + \frac{15i\sin(2e+2fx)}{32a^3d(c+dx)} \\
&\quad + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i\sin(6e+6fx)}{32a^3d(c+dx)} \\
&\quad - \frac{3f\cos(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} + \frac{3if\sin(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad + \frac{3if\sin(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} - \frac{(3if)\int \frac{\cos(2e+2fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(9if)\int \frac{\cos(6e+6fx)}{c+dx} dx}{16a^3d} + \frac{(3f)\int \frac{\sin(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(9f)\int \frac{\sin(6e+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3if\cos(6e-\frac{6cf}{d}))\int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} - \frac{(3f\cos(6e-\frac{6cf}{d}))\int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3if\cos(2e-\frac{2cf}{d}))\int \frac{\cos(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f\cos(2e-\frac{2cf}{d}))\int \frac{\sin(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3if\sin(6e-\frac{6cf}{d}))\int \frac{\sin(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} - \frac{(3f\sin(6e-\frac{6cf}{d}))\int \frac{\cos(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - 2 \left(\frac{(3f\cos(4e-\frac{4cf}{d}))\int \frac{\sin(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} + \frac{(3f\sin(4e-\frac{4cf}{d}))\int \frac{\cos(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} \right) \\
&\quad - \frac{(3if\sin(2e-\frac{2cf}{d}))\int \frac{\sin(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f\sin(2e-\frac{2cf}{d}))\int \frac{\cos(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9\cos(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3\cos(6e+6fx)}{32a^3d(c+dx)} - \frac{9if\cos(2e-\frac{2cf}{d})\text{CosIntegral}(\frac{2cf}{d}+2fx)}{16a^3d^2} \\
&\quad - \frac{3if\cos(4e-\frac{4cf}{d})\text{CosIntegral}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad - \frac{3if\cos(6e-\frac{6cf}{d})\text{CosIntegral}(\frac{6cf}{d}+6fx)}{16a^3d^2} \\
&\quad - \frac{3f\text{CosIntegral}(\frac{6cf}{d}+6fx)\sin(6e-\frac{6cf}{d})}{16a^3d^2} \\
&\quad - \frac{15f\text{CosIntegral}(\frac{2cf}{d}+2fx)\sin(2e-\frac{2cf}{d})}{16a^3d^2} + \frac{15i\sin(2e+2fx)}{32a^3d(c+dx)} \\
&\quad + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i\sin(6e+6fx)}{32a^3d(c+dx)} \\
&\quad - \frac{15f\cos(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{16a^3d^2} + \frac{9if\sin(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{16a^3d^2} \\
&\quad + \frac{3if\sin(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} - 2\left(\frac{3f\text{CosIntegral}(\frac{4cf}{d}+4fx)\sin(4e-\frac{4cf}{d})}{4a^3d^2}\right. \\
&\quad \quad \quad \left.+ \frac{3f\cos(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{4a^3d^2}\right) \\
&\quad - \frac{3f\cos(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{16a^3d^2} + \frac{3if\sin(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{16a^3d^2} \\
&\quad - \frac{(9if\cos(6e-\frac{6cf}{d}))\int\frac{\cos(\frac{6cf}{d}+6fx)}{c+dx}dx}{16a^3d} - \frac{(9f\cos(6e-\frac{6cf}{d}))\int\frac{\sin(\frac{6cf}{d}+6fx)}{c+dx}dx}{16a^3d} \\
&\quad - \frac{(3if\cos(2e-\frac{2cf}{d}))\int\frac{\cos(\frac{2cf}{d}+2fx)}{c+dx}dx}{16a^3d} + \frac{(3f\cos(2e-\frac{2cf}{d}))\int\frac{\sin(\frac{2cf}{d}+2fx)}{c+dx}dx}{16a^3d} \\
&\quad + \frac{(9if\sin(6e-\frac{6cf}{d}))\int\frac{\sin(\frac{6cf}{d}+6fx)}{c+dx}dx}{16a^3d} - \frac{(9f\sin(6e-\frac{6cf}{d}))\int\frac{\cos(\frac{6cf}{d}+6fx)}{c+dx}dx}{16a^3d} \\
&\quad + \frac{(3if\sin(2e-\frac{2cf}{d}))\int\frac{\sin(\frac{2cf}{d}+2fx)}{c+dx}dx}{16a^3d} + \frac{(3f\sin(2e-\frac{2cf}{d}))\int\frac{\cos(\frac{2cf}{d}+2fx)}{c+dx}dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9\cos(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cos^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cos^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3\cos(6e+6fx)}{32a^3d(c+dx)} - \frac{3if\cos(2e-\frac{2cf}{d})\text{CosIntegral}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad - \frac{3if\cos(4e-\frac{4cf}{d})\text{CosIntegral}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad - \frac{3if\cos(6e-\frac{6cf}{d})\text{CosIntegral}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
&\quad - \frac{3f\text{CosIntegral}(\frac{6cf}{d}+6fx)\sin(6e-\frac{6cf}{d})}{4a^3d^2} \\
&\quad - \frac{3f\text{CosIntegral}(\frac{2cf}{d}+2fx)\sin(2e-\frac{2cf}{d})}{4a^3d^2} + \frac{15i\sin(2e+2fx)}{32a^3d(c+dx)} \\
&\quad + \frac{3\sin^2(2e+2fx)}{8a^3d(c+dx)} - \frac{i\sin^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3i\sin(4e+4fx)}{8a^3d(c+dx)} + \frac{3i\sin(6e+6fx)}{32a^3d(c+dx)} \\
&\quad - \frac{3f\cos(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} + \frac{3if\sin(2e-\frac{2cf}{d})\text{Si}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad + \frac{3if\sin(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{2a^3d^2} - 2\left(\frac{3f\text{CosIntegral}(\frac{4cf}{d}+4fx)\sin(4e-\frac{4cf}{d})}{4a^3d^2}\right. \\
&\quad \left.+ \frac{3f\cos(4e-\frac{4cf}{d})\text{Si}(\frac{4cf}{d}+4fx)}{4a^3d^2}\right) \\
&\quad - \frac{3f\cos(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2} + \frac{3if\sin(6e-\frac{6cf}{d})\text{Si}(\frac{6cf}{d}+6fx)}{4a^3d^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.85 (sec), antiderivative size = 833, normalized size of antiderivative = 1.17

$$\begin{aligned}
&\int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx \\
&= \frac{\sec^3(e+fx)(-i\cos(\frac{3cf}{d})+\sin(\frac{3cf}{d}))\left(3d\cos(e+f(-\frac{3c}{d}+x))+d\cos(3(e+f(-\frac{c}{d}+x)))+d\cos(3(e+f(-\frac{c}{d}+x)))\right)}{(c+dx)^2(a+ia\tan(e+fx))^3}
\end{aligned}$$

```

[In] Integrate[1/((c + d*x)^2*(a + I*a*Tan[e + f*x])^3), x]
[Out] (Sec[e + f*x]^3*(-I)*Cos[(3*c*f)/d] + Sin[(3*c*f)/d])*(3*d*Cos[e + f*((-3*c)/d + x)] + d*Cos[3*(e + f*(-(c/d) + x))] + d*Cos[3*(e + f*(c/d + x))] + 3*d*Cos[e + f*((3*c)/d + x)] + (6*I)*c*f*Cos[3*e - (3*f*(c + d*x))/d]*CosIntegral[(6*f*(c + d*x))/d] + (6*I)*d*f*x*Cos[3*e - (3*f*(c + d*x))/d]*CosIntegral[(6*f*(c + d*x))/d] + (6*I)*f*(c + d*x)*CosIntegral[(2*f*(c + d*x))/d]*(Cos[e - (c*f)/d + 3*f*x] + I*Sin[e - (c*f)/d + 3*f*x]) + (3*I)*d*Sin[e + f*((-3*c)/d + x)] + I*d*Sin[3*(e + f*(-(c/d) + x))] - I*d*Sin[3*(e + f*(c/d + x))])

```

$$\begin{aligned}
& + x)) - (3*I)*d*Sin[e + f*((3*c)/d + x)] + 6*c*f*CosIntegral[(6*f*(c + d*x))/d]*Sin[3*e - (3*f*(c + d*x))/d] + 6*d*f*x*CosIntegral[(6*f*(c + d*x))/d]*Sin[3*e - (3*f*(c + d*x))/d] + 12*f*(c + d*x)*CosIntegral[(4*f*(c + d*x))/d]*(I*Cos[e - (f*(c + 3*d*x))/d] + Sin[e - (f*(c + 3*d*x))/d]) + 6*c*f*Cos[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 6*d*f*x*Cos[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (6*I)*c*f*Sin[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + (6*I)*d*f*x*Sin[e - (c*f)/d + 3*f*x]*SinIntegral[(2*f*(c + d*x))/d] + 12*c*f*Cos[e - (f*(c + 3*d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] + 12*d*f*x*Cos[e - (f*(c + 3*d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] - (12*I)*c*f*Sin[e - (f*(c + 3*d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] - (12*I)*d*f*x*Sin[e - (f*(c + 3*d*x))/d]*SinIntegral[(4*f*(c + d*x))/d] + 6*c*f*Cos[3*e - (3*f*(c + d*x))/d]*SinIntegral[(6*f*(c + d*x))/d] + 6*d*f*x*Cos[3*e - (3*f*(c + d*x))/d]*SinIntegral[(6*f*(c + d*x))/d] - (6*I)*c*f*Sin[3*e - (3*f*(c + d*x))/d]*SinIntegral[(6*f*(c + d*x))/d] - (6*I)*d*f*x*Sin[3*e - (3*f*(c + d*x))/d]*SinIntegral[(6*f*(c + d*x))/d]))/(8*a^3*d^2*(c + d*x)*(-I + Tan[e + f*x])^3)
\end{aligned}$$

Maple [A] (verified)

Time = 1.06 (sec) , antiderivative size = 254, normalized size of antiderivative = 0.36

method	result
risch	$ -\frac{1}{8a^3d(dx+c)} - \frac{fe^{-6i(fx+e)}}{8a^3(df+cf)d} + \frac{\frac{3if e^{\frac{6i(cf-de)}{d}} Ei_1\left(6ifx+6ie+\frac{6i(cf-de)}{d}\right)}{4a^3d^2} - \frac{3fe^{-4i(fx+e)}}{8a^3(df+cf)d} + \frac{\frac{3if e^{\frac{4i(cf-de)}{d}} Ei_1\left(4ifx+4ie+\frac{4i(cf-de)}{d}\right)}{2a^3d^2}} $

```

[In] int(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x,method=_RETURNVERBOSE)
[Out] -1/8/a^3/d/(d*x+c)-1/8/a^3*f*exp(-6*I*(f*x+e))/(d*f*x+c*f)/d+3/4*I/a^3*f/d^2*exp(6*I*(c*f-d*e)/d)*Ei(1,6*I*f*x+6*I*e+6*I*(c*f-d*e)/d)-3/8/a^3*f*exp(-4*I*(f*x+e))/(d*f*x+c*f)/d+3/2*I/a^3*f/d^2*exp(4*I*(c*f-d*e)/d)*Ei(1,4*I*f*x+4*I*e+4*I*(c*f-d*e)/d)-3/8/a^3*f*exp(-2*I*(f*x+e))/(d*f*x+c*f)/d+3/4*I/a^3*f/d^2*exp(2*I*(c*f-d*e)/d)*Ei(1,2*I*f*x+2*I*e+2*I*(c*f-d*e)/d)

```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 197, normalized size of antiderivative = 0.28

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx = \\
-\frac{\left(\left(6(i dfx + i cf) Ei\left(-\frac{2(i dfx + i cf)}{d}\right) e^{\left(-\frac{2(i de - i cf)}{d}\right)} + 12(i dfx + i cf) Ei\left(-\frac{4(i dfx + i cf)}{d}\right) e^{\left(-\frac{4(i de - i cf)}{d}\right)} + 6(i dfx + i cf) Ei\left(-\frac{6(i dfx + i cf)}{d}\right) e^{\left(-\frac{6(i de - i cf)}{d}\right)} \right)}{8(a^3d^2)^2}$$

```
[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")
```

[Out]
$$\begin{aligned} & -1/8*((6*(I*d*f*x + I*c*f)*Ei(-2*(I*d*f*x + I*c*f)/d)*e^{(-2*(I*d*e - I*c*f)/d)} + 12*(I*d*f*x + I*c*f)*Ei(-4*(I*d*f*x + I*c*f)/d)*e^{(-4*(I*d*e - I*c*f)/d)} + 6*(I*d*f*x + I*c*f)*Ei(-6*(I*d*f*x + I*c*f)/d)*e^{(-6*(I*d*e - I*c*f)/d)} + d)*e^{(6*I*f*x + 6*I*e)} + 3*d*e^{(4*I*f*x + 4*I*e)} + 3*d*e^{(2*I*f*x + 2*I*e)} + d)*e^{(-6*I*f*x - 6*I*e)}/(a^3*d^3*x + a^3*c*d^2) \end{aligned}$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx \\ = & \frac{i \int \frac{1}{c^2\tan^3(e+fx)-3c^2\tan^2(e+fx)-3c^2\tan(e+fx)+ic^2+2cdx\tan^3(e+fx)-6icdx\tan^2(e+fx)-6cdx\tan(e+fx)+2icdx+d^2x^2\tan^3(e+fx)-d^2x^2\tan(e+fx)} dx}{a^3} \end{aligned}$$

[In] integrate(1/(d*x+c)**2/(a+I*a*tan(f*x+e))**3,x)

[Out]
$$\begin{aligned} & I*Integral(1/(c**2*tan(e + f*x)**3 - 3*I*c**2*tan(e + f*x)**2 - 3*c**2*tan(e + f*x) + I*c**2 + 2*c*d*x*tan(e + f*x)**3 - 6*I*c*d*x*tan(e + f*x)**2 - 6*c*d*x*tan(e + f*x) + 2*I*c*d*x + d**2*x**2*tan(e + f*x)**3 - 3*I*d**2*x**2*tan(e + f*x)**2 - 3*d**2*x**2*tan(e + f*x) + I*d**2*x**2), x)/a**3 \end{aligned}$$

Maxima [A] (verification not implemented)

none

Time = 0.43 (sec) , antiderivative size = 297, normalized size of antiderivative = 0.42

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+ia\tan(e+fx))^3} dx = \\ & \frac{-3 f^2 \cos\left(-\frac{2(de-cf)}{d}\right) E_2\left(-\frac{2(-i(fx+e)d+i(de-icf))}{d}\right) + 3 f^2 \cos\left(-\frac{4(de-cf)}{d}\right) E_2\left(-\frac{4(-i(fx+e)d+i(de-icf))}{d}\right) + f^2 \cos\left(-\frac{6(de-cf)}{d}\right) E_2\left(-\frac{6(-i(fx+e)d+i(de-icf))}{d}\right)}{(a+ia\tan(e+fx))^3} \end{aligned}$$

[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")

[Out]
$$\begin{aligned} & -1/8*(3*f^2*cos(-2*(d*e - c*f)/d)*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*f^2*cos(-4*(d*e - c*f)/d)*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + f^2*cos(-6*(d*e - c*f)/d)*exp_integral_e(2, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d) + 3*I*f^2*exp_integral_e(2, -2*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-2*(d*e - c*f)/d) + 3*I*f^2*exp_integral_e(2, -4*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-4*(d*e - c*f)/d) + I*f^2*exp_integral_e(2, -6*(-I*(f*x + e)*d + I*d*e - I*c*f)/d)*sin(-6*(d*e - c*f)/d) + f^2)/(((f*x + e)*a^3*d^2 - a^3*d^2*e + a^3*c*d*f)*f) \end{aligned}$$

Giac [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2915 vs. $2(648) = 1296$.

Time = 30.48 (sec) , antiderivative size = 2915, normalized size of antiderivative = 4.09

$$\int \frac{1}{(c + dx)^2(a + i a \tan(e + fx))^3} dx = \text{Too large to display}$$

```
[In] integrate(1/(d*x+c)^2/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")
[Out] 1/8*(-6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos(-6*(d*e - c*f)/d)*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 12*I*d*e*f^2*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*c*f^3*cos(-4*(d*e - c*f)/d)*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos(-2*(d*e - c*f)/d)*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*d*e*f^2*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*cos_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 12*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*d*e*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*sin(-4*(d*e - c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*d*e*f^2*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*c*f^3*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(-6*(d*e - c*f)/d) - 6*d*e*f^2*cos_integral(-6*(d*e - c*f)/d) + 6*c*f^3*cos_integral(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*sin(-6*(d*e - c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(-6*(d*e - c*f)/d) - 6*d*e*f^2*cos_integral(-6*(d*e - c*f)/d) + 6*d*e*f^2*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*sin(-6*(d*e - c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*cos_integral(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*d*e*f^2*cos_integral(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*d*e*f^2*cos_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*sin(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)
```

$$\begin{aligned}
& -6*c*f^3*\cos(-6*(d*e - c*f)/d)*\sin_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*\sin(-6*(d*e - c*f)/d)*\sin_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*\sin(-6*(d*e - c*f)/d)*\sin_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*\sin(-6*(d*e - c*f)/d)*\sin_integral(6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*\cos(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 12*d*e*f^2*\cos(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*c*f^3*\cos(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*\sin(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 12*I*d*e*f^2*\sin(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 12*I*c*f^3*\sin(-4*(d*e - c*f)/d)*\sin_integral(4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*\cos(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*d*e*f^2*\cos(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*c*f^3*\cos(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*\sin(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 6*I*d*e*f^2*\sin(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 6*I*c*f^3*\sin(-2*(d*e - c*f)/d)*\sin_integral(2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 3*d*f^2*\cos(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - 3*d*f^2*\cos(-4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - d*f^2*\cos(-6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - 3*I*d*f^2*\sin(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - 3*I*d*f^2*\sin(-4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - I*d*f^2*\sin(-6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - d*f^2*d^2/(((d*x + c)*a^3*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a^3*d^5*e + a^3*c*d^4*f)*f)
\end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + ia \tan(e + fx))^3} dx = \int \frac{1}{(a + a \tan(e + fx) 1i)^3 (c + dx)^2} dx$$

[In] `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2),x)`

[Out] `int(1/((a + a*tan(e + f*x)*1i)^3*(c + d*x)^2), x)`

3.34 $\int (c + dx)^m (a + ia \tan(e + fx))^2 dx$

Optimal result	233
Rubi [N/A]	233
Mathematica [N/A]	234
Maple [N/A] (verified)	234
Fricas [N/A]	234
Sympy [N/A]	235
Maxima [N/A]	235
Giac [N/A]	236
Mupad [N/A]	236

Optimal result

Integrand size = 23, antiderivative size = 23

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx))^2, x)$$

[Out] Unintegrable((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec), antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

[In] Int[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2,x]

[Out] Defer[Int][(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

Mathematica [N/A]

Not integrable

Time = 42.84 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.09

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (c + dx)^m (a + ia \tan(e + fx))^2 dx$$

[In] `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]`

[Out] `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x])^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.19 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.91

$$\int (dx + c)^m (a + ia \tan(fx + e))^2 dx$$

[In] `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2, x)`

[Out] `int((d*x+c)^m*(a+I*a*tan(f*x+e))^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 5.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (i a \tan(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2, x, algorithm="fricas")`

[Out] `(-2*I*(d*x + c)^m*a^2 + (f*e^(2*I*f*x + 2*I*e) + f)*integral(-2*(-I*a^2*d*m - 2*(a^2*d*f*x + a^2*c*f)*e^(2*I*f*x + 2*I*e))*(d*x + c)^m/(d*f*x + c*f + (d*f*x + c*f)*e^(2*I*f*x + 2*I*e)), x))/(f*e^(2*I*f*x + 2*I*e) + f)`

Sympy [N/A]

Not integrable

Time = 5.32 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.22

$$\begin{aligned} \int (c + dx)^m (a + ia \tan(e + fx))^2 dx &= -a^2 \left(\int (c + dx)^m \tan^2(e + fx) dx \right. \\ &\quad + \int (-2i(c + dx)^m \tan(e + fx)) dx \\ &\quad \left. + \int (-(c + dx)^m) dx \right) \end{aligned}$$

[In] `integrate((d*x+c)**m*(a+I*a*tan(f*x+e))**2,x)`

[Out] `-a**2*(Integral((c + d*x)**m*tan(e + f*x)**2, x) + Integral(-2*I*(c + d*x)*m*tan(e + f*x), x) + Integral(-(c + d*x)**m, x))`

Maxima [N/A]

Not integrable

Time = 1.83 (sec) , antiderivative size = 520, normalized size of antiderivative = 22.61

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (ia \tan(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")`

[Out] `(d*x + c)^(m + 1)*a^2/(d*(m + 1)) + integrate((3*(d*x + c)^m*a^2*cos(4*f*x + 4*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e)^2 + 3*(d*x + c)^m*a^2*sin(4*f*x + 4*e)^2 + 4*(d*x + c)^m*a^2*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) - 4*(d*x + c)^m*a^2*sin(2*f*x + 2*e)^2 - 4*(d*x + c)^m*a^2*cos(2*f*x + 2*e) - (d*x + c)^m*a^2 + 2*(2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*cos(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x) + I*integrate(e(-4*(2*(d*x + c)^m*a^2*cos(4*f*x + 4*e)*sin(2*f*x + 2*e) - (2*(d*x + c)^m*a^2*cos(2*f*x + 2*e) + (d*x + c)^m*a^2)*sin(4*f*x + 4*e))/(2*(2*cos(2*f*x + 2*e) + 1)*cos(4*f*x + 4*e) + cos(4*f*x + 4*e)^2 + 4*cos(2*f*x + 2*e)^2 + sin(4*f*x + 4*e)^2 + 4*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*sin(2*f*x + 2*e)^2 + 4*cos(2*f*x + 2*e) + 1), x)`

Giac [N/A]

Not integrable

Time = 0.64 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (i a \tan(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((I*a*tan(f*x + e) + a)^2*(d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 3.37 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.04

$$\int (c + dx)^m (a + ia \tan(e + fx))^2 dx = \int (a + a \tan(e + fx) \cdot 1i)^2 (c + dx)^m dx$$

[In] `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m,x)`

[Out] `int((a + a*tan(e + f*x)*1i)^2*(c + d*x)^m, x)`

3.35 $\int (c + dx)^m (a + ia \tan(e + fx)) dx$

Optimal result	237
Rubi [N/A]	237
Mathematica [N/A]	238
Maple [N/A] (verified)	238
Fricas [N/A]	238
Sympy [N/A]	238
Maxima [N/A]	239
Giac [N/A]	239
Mupad [N/A]	239

Optimal result

Integrand size = 21, antiderivative size = 21

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \text{Int}((c + dx)^m (a + ia \tan(e + fx)), x)$$

[Out] Unintegrable((d*x+c)^m*(a+I*a*tan(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (c + dx)^m (a + ia \tan(e + fx)) dx$$

[In] Int[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]

[Out] Defer[Int][(c + d*x)^m*(a + I*a*Tan[e + f*x]), x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + ia \tan(e + fx)) dx$$

Mathematica [N/A]

Not integrable

Time = 9.86 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (c + dx)^m (a + ia \tan(e + fx)) dx$$

[In] `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]),x]`

[Out] `Integrate[(c + d*x)^m*(a + I*a*Tan[e + f*x]), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.90

$$\int (dx + c)^m (a + ia \tan(fx + e)) dx$$

[In] `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

[Out] `int((d*x+c)^m*(a+I*a*tan(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.62

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="fricas")`

[Out] `integral(2*(d*x + c)^m*a*e^(2*I*f*x + 2*I*e)/(e^(2*I*f*x + 2*I*e) + 1), x)`

Sympy [N/A]

Not integrable

Time = 2.40 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.38

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = ia \left(\int (-i(c + dx)^m) dx + \int (c + dx)^m \tan(e + fx) dx \right)$$

[In] `integrate((d*x+c)**m*(a+I*a*tan(f*x+e)),x)`

[Out] `I*a*(Integral(-I*(c + d*x)**m, x) + Integral((c + d*x)**m*tan(e + f*x), x))`

Maxima [N/A]

Not integrable

Time = 0.50 (sec) , antiderivative size = 79, normalized size of antiderivative = 3.76

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

[Out] `2*I*a*integrate((d*x + c)^m*sin(2*f*x + 2*e)/(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1), x) + (d*x + c)^(m + 1)*a/(d*(m + 1))`

Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.00

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (ia \tan(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+I*a*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate((I*a*tan(f*x + e) + a)*(d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 3.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.05

$$\int (c + dx)^m (a + ia \tan(e + fx)) dx = \int (a + a \tan(e + fx) \cdot 1i) (c + dx)^m dx$$

[In] `int((a + a*tan(e + f*x)*1i)*(c + d*x)^m,x)`

[Out] `int((a + a*tan(e + f*x)*1i)*(c + d*x)^m, x)`

3.36 $\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx$

Optimal result	240
Rubi [A] (verified)	240
Mathematica [A] (verified)	241
Maple [F]	242
Fricas [A] (verification not implemented)	242
Sympy [F]	242
Maxima [F]	243
Giac [F]	243
Mupad [F(-1)]	243

Optimal result

Integrand size = 23, antiderivative size = 98

$$\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx = \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{af}$$

[Out] $\frac{1/2*(d*x+c)^(1+m)/a/d/(1+m)+I*2^(-2-m)*(d*x+c)^m*\text{GAMMA}(1+m, 2*I*f*(d*x+c)/d)}{a/\exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)}$

Rubi [A] (verified)

Time = 0.15 (sec), antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {3808, 2212}

$$\int \frac{(c+dx)^m}{a+ia\tan(e+fx)} dx = \frac{(c+dx)^{m+1}}{2ad(m+1)} + \frac{i2^{-m-2}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2if(c+dx)}{d}\right)}{af}$$

[In] $\text{Int}[(c + d*x)^m/(a + I*a*Tan[e + f*x]), x]$

[Out] $(c + d*x)^(1 + m)/(2*a*d*(1 + m)) + (I*2^(-2 - m)*(c + d*x)^m*\text{Gamma}[1 + m, ((2*I)*f*(c + d*x))/d])/((a*E^((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)^m))$

Rule 2212

```

Int[(F_)^((g_ .)*(e_ .) + (f_ .)*(x_ ))*((c_ .) + (d_ .)*(x_ ))^(m_ ), x_Symbol]
:> Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d)
)^((IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]

```

Rule 3808

```

Int[((c_ .) + (d_ .)*(x_ ))^(m_ )/((a_ .) + (b_ .)*tan[(e_ .) + (f_ .)*(x_ )]), x_Sym
bol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + Dist[1/(2*a), Int[(c +
d*x)^m*E^(2*(a/b)*(e + f*x)), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
EqQ[a^2 + b^2, 0] && !IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c + dx)^{1+m}}{2ad(1+m)} + \frac{\int e^{-2i(e+fx)}(c + dx)^m dx}{2a} \\
&= \frac{(c + dx)^{1+m}}{2ad(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c + dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{af}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.77 (sec), antiderivative size = 133, normalized size of antiderivative = 1.36

$$\begin{aligned}
&\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx \\
&= \frac{e^{-ie}(c + dx)^m \left(\frac{2e^{2ie}f(c+dx)}{d(1+m)} + i2^{-m}e^{\frac{2icf}{d}} \left(\frac{if(c+dx)}{d} \right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right) \right) \sec(e + fx)(\cos(fx) + i \sin(fx))}{4f(a + ia \tan(e + fx))}
\end{aligned}$$

[In] `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x]), x]`

[Out] `((c + d*x)^m*((2*E^((2*I)*e)*f*(c + d*x))/(d*(1 + m)) + (I*E^(((2*I)*c*f)/d)*Gamma[1 + m, ((2*I)*f*(c + d*x))/d])/(2^m*((I*f*(c + d*x))/d)^m))*Sec[e + f*x]*(Cos[f*x] + I*Sin[f*x]))/(4*E^(I*e)*f*(a + I*a*Tan[e + f*x]))`

Maple [F]

$$\int \frac{(dx + c)^m}{a + ia \tan(fx + e)} dx$$

[In] `int((d*x+c)^m/(a+I*a*tan(f*x+e)),x)`
[Out] `int((d*x+c)^m/(a+I*a*tan(f*x+e)),x)`

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.87

$$\begin{aligned} & \int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx \\ &= \frac{(i dm + i d)e^{\left(-\frac{dm \log(\frac{2if}{d}) + 2ie - 2icf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-idfx - icf)}{d}\right) + 2(df x + cf)(dx + c)^m}{4(adfm + adf)} \end{aligned}$$

[In] `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="fricas")`
[Out] `1/4*((I*d*m + I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(m + 1, -2*(-I*d*f*x - I*c*f)/d) + 2*(d*f*x + c*f)*(d*x + c)^m)/(a*d*f*m + a*d*f)`

Sympy [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = -\frac{i \int \frac{(c+dx)^m}{\tan(e+fx)-i} dx}{a}$$

[In] `integrate((d*x+c)**m/(a+I*a*tan(f*x+e)),x)`
[Out] `-I*Integral((c + d*x)**m/(\tan(e + f*x) - I), x)/a`

Maxima [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

[In] `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="maxima")`

[Out] `1/2*((d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*i integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c)))/(a*d*m + a*d)`

Giac [F]

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(dx + c)^m}{i a \tan(fx + e) + a} dx$$

[In] `integrate((d*x+c)^m/(a+I*a*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate((d*x + c)^m/(I*a*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{a + ia \tan(e + fx)} dx = \int \frac{(c + dx)^m}{a + a \tan(e + fx) \text{ li}} dx$$

[In] `int((c + d*x)^m/(a + a*tan(e + f*x)*1i),x)`

[Out] `int((c + d*x)^m/(a + a*tan(e + f*x)*1i), x)`

3.37 $\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx$

Optimal result	244
Rubi [A] (verified)	244
Mathematica [A] (verified)	245
Maple [F]	246
Fricas [A] (verification not implemented)	246
Sympy [F]	246
Maxima [F]	247
Giac [F]	247
Mupad [F(-1)]	247

Optimal result

Integrand size = 23, antiderivative size = 171

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^2f} \\ &+ \frac{i4^{-2-m}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^2f} \end{aligned}$$

[Out] $1/4*(d*x+c)^(1+m)/a^2/d/(1+m)+I*2^(-2-m)*(d*x+c)^m*\text{GAMMA}(1+m, 2*I*f*(d*x+c)/d)/a^2/\exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*4^(-2-m)*(d*x+c)^m*\text{GAMMA}(1+m, 4*I*f*(d*x+c)/d)/a^2/\exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)$

Rubi [A] (verified)

Time = 0.23 (sec), antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.087, Rules used = {3810, 2212}

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ &= \frac{i2^{-m-2}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2if(c+dx)}{d}\right)}{a^2f} \\ &+ \frac{i4^{-m-2}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{4if(c+dx)}{d}\right)}{a^2f} + \frac{(c+dx)^{m+1}}{4a^2d(m+1)} \end{aligned}$$

[In] $\text{Int}[(c + d*x)^m / (a + I*a*\text{Tan}[e + f*x])^2, x]$

[Out] $(c + d*x)^{1+m} / (4*a^2*d*(1+m)) + (I*2^{-2-m}*(c + d*x)^m * \text{Gamma}[1+m, ((2*I)*f*(c + d*x))/d]) / (a^{2*E^{((2*I)*(e - (c*f)/d))}} * f*((I*f*(c + d*x))/d)^m) + (I*4^{-2-m}*(c + d*x)^m * \text{Gamma}[1+m, ((4*I)*f*(c + d*x))/d]) / (a^{2*E^{((4*I)*(e - (c*f)/d))}} * f*((I*f*(c + d*x))/d)^m)$

Rule 2212

```
Int[(F_)^((g_)*(e_) + (f_)*(x_))*((c_) + (d_)*(x_))^m, x_Symbol]
: > Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d))
)^((IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d)^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

Rule 3810

```
Int[((c_) + (d_)*(x_))^m*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^n, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{-n}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^m}{4a^2} + \frac{e^{-2ie-2ifx}(c+dx)^m}{2a^2} + \frac{e^{-4ie-4ifx}(c+dx)^m}{4a^2} \right) dx \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{\int e^{-4ie-4ifx}(c+dx)^m dx}{4a^2} + \frac{\int e^{-2ie-2ifx}(c+dx)^m dx}{2a^2} \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{i2^{-2-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^2f} \\ &\quad + \frac{i4^{-2-m}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^2f} \end{aligned}$$

Mathematica [A] (verified)

Time = 4.49 (sec), antiderivative size = 192, normalized size of antiderivative = 1.12

$$\begin{aligned} &\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^2} dx \\ &= \frac{(c+dx)^m \left(\frac{4e^{2ie}f(c+dx)}{d(1+m)} + i2^{2-m}e^{\frac{2icf}{d}} \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2if(c+dx)}{d}\right) + i4^{-m}e^{-2ie+\frac{4icf}{d}} \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4if(c+dx)}{d}\right) \right)}{16f(a+ia\tan(e+fx))^2} \end{aligned}$$

[In] $\text{Integrate}[(c + d*x)^m/(a + I*a*\tan[e + f*x])^2, x]$

[Out] $((c + d*x)^m * ((4*E^((2*I)*e)*f*(c + d*x))/(d*(1 + m)) + (I*2^(2 - m)*E^(((2*I)*c*f)/d)*\text{Gamma}[1 + m, ((2*I)*f*(c + d*x))/d])/((I*f*(c + d*x))/d)^m + (I*E^((-2*I)*e + ((4*I)*c*f)/d)*\text{Gamma}[1 + m, ((4*I)*f*(c + d*x))/d])/4^m * ((I*f*(c + d*x))/d)^m) * \text{Sec}[e + f*x]^2 * (\cos[f*x] + I*\sin[f*x])^2)/(16*f*(a + I*a*\tan[e + f*x])^2)$

Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^2} dx$$

[In] $\text{int}((d*x+c)^m/(a+I*a*tan(f*x+e))^2, x)$

[Out] $\text{int}((d*x+c)^m/(a+I*a*tan(f*x+e))^2, x)$

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec), antiderivative size = 144, normalized size of antiderivative = 0.84

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \frac{-4(-i dm - i d)e^{\left(-\frac{dm \log(\frac{2if}{d}) + 2i de - 2i cf}{d}\right)} \Gamma(m + 1, -\frac{2(-i dfx - i cf)}{d}) - (i dm + i d)e^{\left(-\frac{dm \log(\frac{4if}{d}) + 4i de - 4i cf}{d}\right)} \Gamma(m + 1, -\frac{2(-i dfx - i cf)}{d})}{16(a^2 dfm + a^2 df)}$$

[In] $\text{integrate}((d*x+c)^m/(a+I*a*tan(f*x+e))^2, x, \text{algorithm}=\text{fricas})$

[Out] $-1/16 * (4*(-I*d*m - I*d)*e^{-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d}*\text{gamma}(m + 1, -2*(-I*d*f*x - I*c*f)/d) - (I*d*m + I*d)*e^{-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d}*\text{gamma}(m + 1, -4*(-I*d*f*x - I*c*f)/d) - 4*(d*f*x + c*f)*(d*x + c)^m)/(a^2*d*f*m + a^2*d*f)$

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = -\frac{\int \frac{(c + dx)^m}{\tan^2(e + fx) - 2i \tan(e + fx) - 1} dx}{a^2}$$

[In] $\text{integrate}((d*x+c)**m/(a+I*a*tan(f*x+e))**2, x)$

[Out] $-\text{Integral}((c + d*x)**m/(\tan(e + f*x)**2 - 2*I*tan(e + f*x) - 1), x)/a**2$

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^2} dx$$

```
[In] integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="maxima")
[Out] 1/4*((d*m + d)*integrate((d*x + c)^m*cos(4*f*x + 4*e), x) + 2*(d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*integrate((d*x + c)^m*sin(4*f*x + 4*e), x) + 2*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c)))/(a^2*d*m + a^2*d)
```

Giac [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^2} dx$$

```
[In] integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^2,x, algorithm="giac")
[Out] integrate((d*x + c)^m/(I*a*tan(f*x + e) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^2} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + f x) 1i)^2} dx$$

```
[In] int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^2,x)
[Out] int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^2, x)
```

3.38 $\int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx$

Optimal result	248
Rubi [A] (verified)	248
Mathematica [A] (verified)	250
Maple [F]	250
Fricas [A] (verification not implemented)	251
Sympy [F]	251
Maxima [F]	251
Giac [F]	252
Mupad [F(-1)]	252

Optimal result

Integrand size = 23, antiderivative size = 251

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\ &= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3i2^{-4-m}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^3f} \\ &+ \frac{3i2^{-5-2m}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^3f} \\ &+ \frac{i2^{-4-m}3^{-1-m}e^{-6i\left(e-\frac{cf}{d}\right)}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{6if(c+dx)}{d}\right)}{a^3f} \end{aligned}$$

[Out] $1/8*(d*x+c)^(1+m)/a^3/d/(1+m)+3*I*2^(-4-m)*(d*x+c)^m*\text{GAMMA}(1+m, 2*I*f*(d*x+c)/d)/a^3/\exp(2*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+3*I*2^(-5-2*m)*(d*x+c)^m*\text{GAMMA}(1+m, 4*I*f*(d*x+c)/d)/a^3/\exp(4*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)+I*2^(-4-m)*3^(-1-m)*(d*x+c)^m*\text{GAMMA}(1+m, 6*I*f*(d*x+c)/d)/a^3/\exp(6*I*(e-c*f/d))/f/((I*f*(d*x+c)/d)^m)$

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 251, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used

= {3810, 2212}

$$\begin{aligned}
 & \int \frac{(c+dx)^m}{(a+ia\tan(e+fx))^3} dx \\
 = & \frac{3i2^{-m-4}e^{-2i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2if(c+dx)}{d}\right)}{a^3 f} \\
 + & \frac{3i2^{-2m-5}e^{-4i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{4if(c+dx)}{d}\right)}{a^3 f} \\
 + & \frac{i2^{-m-4}3^{-m-1}e^{-6i\left(e-\frac{cf}{d}\right)}(c+dx)^m \left(\frac{if(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{6if(c+dx)}{d}\right)}{a^3 f} + \frac{(c+dx)^{m+1}}{8a^3 d(m+1)}
 \end{aligned}$$

[In] Int[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3, x]

[Out] $(c + d*x)^{(1 + m)}/(8*a^3*d*(1 + m)) + ((3*I)*2^{(-4 - m)}*(c + d*x)^m*\text{Gamma}[1 + m, ((2*I)*f*(c + d*x))/d])/((a^3*E^{((2*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)})^m) + ((3*I)*2^{(-5 - 2*m)}*(c + d*x)^m*\text{Gamma}[1 + m, ((4*I)*f*(c + d*x))/d])/((a^3*E^{((4*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)})^m) + (I*2^{(-4 - m)}*3^{(-1 - m)}*(c + d*x)^m*\text{Gamma}[1 + m, ((6*I)*f*(c + d*x))/d])/((a^3*E^{((6*I)*(e - (c*f)/d))*f*((I*f*(c + d*x))/d)})^m)$

Rule 2212

```

Int[(F_)^((g_.)*((e_.) + (f_.*(x_)))*((c_.) + (d_.*(x_))^m), x_Symbol]
:> Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*((-f)*g*(Log[F]/d))^
(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d)^FracPart[m]))*\Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]

```

Rule 3810

```

Int[((c_.) + (d_.*(x_))^m)*((a_.) + (b_.*tan[(e_.) + (f_.*(x_))])^n), x_Symbol]
:> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{(c+dx)^m}{8a^3} + \frac{3e^{-2ie-2ifx}(c+dx)^m}{8a^3} + \frac{3e^{-4ie-4ifx}(c+dx)^m}{8a^3} \right. \\
 &\quad \left. + \frac{e^{-6ie-6ifx}(c+dx)^m}{8a^3} \right) dx \\
 &= \frac{(c+dx)^{1+m}}{8a^3 d(1+m)} + \frac{\int e^{-6ie-6ifx}(c+dx)^m dx}{8a^3} \\
 &\quad + \frac{3 \int e^{-2ie-2ifx}(c+dx)^m dx}{8a^3} + \frac{3 \int e^{-4ie-4ifx}(c+dx)^m dx}{8a^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3i2^{-4-m}e^{-2i(e-\frac{cf}{d})}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{2if(c+dx)}{d}\right)}{a^3f} \\
&\quad + \frac{3i2^{-5-2m}e^{-4i(e-\frac{cf}{d})}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{4if(c+dx)}{d}\right)}{a^3f} \\
&\quad + \frac{i2^{-4-m}3^{-1-m}e^{-6i(e-\frac{cf}{d})}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\Gamma\left(1+m, \frac{6if(c+dx)}{d}\right)}{a^3f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 13.03 (sec), antiderivative size = 269, normalized size of antiderivative = 1.07

$$\begin{aligned}
&\int \frac{(c+dx)^m}{(a + ia \tan(e + fx))^3} dx \\
&= 2^{-5-2m}3^{-1-m}e^{-3ie}(c+dx)^m\left(\frac{if(c+dx)}{d}\right)^{-m}\left(12^{1+m}e^{6ie}f(c+dx)\left(\frac{if(c+dx)}{d}\right)^m + i2^{1+m}3^{2+m}de^{2i(2e+\frac{cf}{d})}(1+m)\right)
\end{aligned}$$

[In] `Integrate[(c + d*x)^m/(a + I*a*Tan[e + f*x])^3, x]`

[Out] `(2^(-5 - 2*m)*3^(-1 - m)*(c + d*x)^m*(12^(1 + m)*E^((6*I)*e)*f*(c + d*x)*((I*f*(c + d*x))/d)^m + I*2^(1 + m)*3^(2 + m)*d*E^((2*I)*(2*e + (c*f)/d))*(1 + m)*Gamma[1 + m, ((2*I)*f*(c + d*x))/d] + I*3^(2 + m)*d*E^((2*I)*e + ((4*I)*c*f)/d)*(1 + m)*Gamma[1 + m, ((4*I)*f*(c + d*x))/d] + I*2^(1 + m)*d*E^(((6*I)*c*f)/d)*(1 + m)*Gamma[1 + m, ((6*I)*f*(c + d*x))/d])*Sec[e + f*x]^3*(Cos[f*x] + I*Sin[f*x])^3)/(d*E^((3*I)*e)*f*(1 + m)*((I*f*(c + d*x))/d)^m*(a + I*a*Tan[e + f*x])^3)`

Maple [F]

$$\int \frac{(dx + c)^m}{(a + ia \tan(fx + e))^3} dx$$

[In] `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3, x)`

[Out] `int((d*x+c)^m/(a+I*a*tan(f*x+e))^3, x)`

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec) , antiderivative size = 198, normalized size of antiderivative = 0.79

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx =$$

$$-\frac{18(-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{2if}{d}\right) + 2i de - 2i cf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-i dfx - i cf)}{d}\right) + 9(-i dm - i d)e^{\left(-\frac{dm \log\left(\frac{4if}{d}\right) + 4i de - 4i cf}{d}\right)} \Gamma\left(m + 1, -\frac{2(-i dfx - i cf)}{d}\right)}{a^3}$$

```
[In] integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="fricas")
[Out] -1/96*(18*(-I*d*m - I*d)*e^(-(d*m*log(2*I*f/d) + 2*I*d*e - 2*I*c*f)/d)*gamma(a(m + 1, -2*(-I*d*f*x - I*c*f)/d) + 9*(-I*d*m - I*d)*e^(-(d*m*log(4*I*f/d) + 4*I*d*e - 4*I*c*f)/d)*gamma(m + 1, -4*(-I*d*f*x - I*c*f)/d) + 2*(-I*d*m - I*d)*e^(-(d*m*log(6*I*f/d) + 6*I*d*e - 6*I*c*f)/d)*gamma(m + 1, -6*(-I*d*f*x - I*c*f)/d) - 12*(d*f*x + c*f)*(d*x + c)^m)/(a^3*d*f*m + a^3*d*f)
```

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \frac{i \int \frac{(c + dx)^m}{\tan^3(e + fx) - 3i \tan^2(e + fx) - 3 \tan(e + fx) + i} dx}{a^3}$$

```
[In] integrate((d*x+c)**m/(a+I*a*tan(f*x+e))**3,x)
[Out] I*Integral((c + d*x)**m/(tan(e + f*x)**3 - 3*I*tan(e + f*x)**2 - 3*tan(e + f*x) + I), x)/a**3
```

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^3} dx$$

```
[In] integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="maxima")
[Out] 1/8*((d*m + d)*integrate((d*x + c)^m*cos(6*f*x + 6*e), x) + 3*(d*m + d)*integrate((d*x + c)^m*cos(4*f*x + 4*e), x) + 3*(d*m + d)*integrate((d*x + c)^m*cos(2*f*x + 2*e), x) - (I*d*m + I*d)*integrate((d*x + c)^m*sin(6*f*x + 6*e), x) + 3*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(4*f*x + 4*e), x) + 3*(-I*d*m - I*d)*integrate((d*x + c)^m*sin(2*f*x + 2*e), x) + e^(m*log(d*x + c) + log(d*x + c)))/(a^3*d*m + a^3*d)
```

Giac [F]

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(dx + c)^m}{(i a \tan(fx + e) + a)^3} dx$$

[In] integrate((d*x+c)^m/(a+I*a*tan(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^m/(I*a*tan(f*x + e) + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + ia \tan(e + fx))^3} dx = \int \frac{(c + dx)^m}{(a + a \tan(e + fx) \text{ li})^3} dx$$

[In] int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^3,x)

[Out] int((c + d*x)^m/(a + a*tan(e + f*x)*1i)^3, x)

$$\mathbf{3.39} \quad \int (c + dx)^3 (a + b \tan(e + fx)) dx$$

Optimal result	253
Rubi [A] (verified)	253
Mathematica [B] (verified)	256
Maple [B] (verified)	257
Fricas [B] (verification not implemented)	257
Sympy [F]	258
Maxima [B] (verification not implemented)	258
Giac [F]	259
Mupad [F(-1)]	259

Optimal result

Integrand size = 18, antiderivative size = 152

$$\begin{aligned} \int (c + dx)^3 (a + b \tan(e + fx)) dx = & \frac{a(c + dx)^4}{4d} + \frac{ib(c + dx)^4}{4d} - \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ibd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{3bd^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\ & - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \end{aligned}$$

```
[Out] 1/4*a*(d*x+c)^4/d+1/4*I*b*(d*x+c)^4/d-b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f+
3/2*I*b*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2-3/2*b*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3-3/4*I*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4
```

Rubi [A] (verified)

Time = 0.29 (sec), antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3803, 3800, 2221, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int (c + dx)^3 (a + b \tan(e + fx)) dx = & \frac{a(c + dx)^4}{4d} - \frac{3bd^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\ & + \frac{3ibd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} + \frac{ib(c + dx)^4}{4d} \\ & - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \end{aligned}$$

[In] $\text{Int}[(c + d*x)^3*(a + b*\tan[e + f*x]), x]$

[Out] $(a*(c + d*x)^4)/(4*d) + ((I/4)*b*(c + d*x)^4)/d - (b*(c + d*x)^3*\log[1 + E^((2*I)*(e + f*x))])/f + (((3*I)/2)*b*d*(c + d*x)^2*\text{PolyLog}[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*d^2*(c + d*x)*\text{PolyLog}[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*\text{PolyLog}[4, -E^((2*I)*(e + f*x))])/f^4$

Rule 2221

$\text{Int}[(((F_)^((g_)*(e_)+(f_)*(x_))))^{(n_)}*((c_)+(d_)*(x_))^{(m_)})/((a_)+(b_)*(F_)^((g_)*(e_)+(f_)*(x_))))^{(n_)}, x]$ Symbol :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

$\text{Int}[u_, x]$ Symbol :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*(a_)+(b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

$\text{Int}[\log[1 + (e_)*(F_)^((c_)*(a_)+(b_)*(x_)))^{(n_)}]*((f_)+(g_)*(x_))^{(m_)}, x]$ Symbol :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3800

$\text{Int}[((c_)+(d_)*(x_))^{(m_)}*\tan[(e_)+(f_)*(x_)], x]$ Symbol :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

Rule 3803

$\text{Int}[((c_)+(d_)*(x_))^{(m_)}*((a_)+(b_)*\tan[(e_)+(f_)*(x_)])^{(n_)}, x]$ Symbol :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 6724

$\text{Int}[\text{PolyLog}[n_, (c_)*(a_)+(b_)*(x_))^{(p_)}]/((d_)+(e_)*(x_)), x]$ Symbol :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d}

```
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c+dx)^3 + b(c+dx)^3 \tan(e+fx)) \, dx \\
&= \frac{a(c+dx)^4}{4d} + b \int (c+dx)^3 \tan(e+fx) \, dx \\
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - (2ib) \int \frac{e^{2i(e+fx)}(c+dx)^3}{1+e^{2i(e+fx)}} \, dx \\
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - \frac{b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{(3bd) \int (c+dx)^2 \log(1+e^{2i(e+fx)}) \, dx}{f} \\
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - \frac{b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ibd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{(3ibd^2) \int (c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)}) \, dx}{f^2} \\
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - \frac{b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ibd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{3bd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{(3bd^3) \int \operatorname{PolyLog}(3, -e^{2i(e+fx)}) \, dx}{2f^3} \\
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - \frac{b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ibd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3bd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad - \frac{(3ibd^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(3, -x)}{x} \, dx, x, e^{2i(e+fx)}\right)}{4f^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{a(c+dx)^4}{4d} + \frac{ib(c+dx)^4}{4d} - \frac{b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ibd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{3bd^2(c+dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
\end{aligned}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 342 vs. $2(152) = 304$.

Time = 0.15 (sec), antiderivative size = 342, normalized size of antiderivative = 2.25

$$\begin{aligned}
\int (c+dx)^3(a+b\tan(e+fx)) dx &= ac^3x + \frac{3}{2}ac^2dx^2 + \frac{3}{2}ibc^2dx^2 + acd^2x^3 + ibcd^2x^3 \\
&\quad + \frac{1}{4}ad^3x^4 + \frac{1}{4}ibd^3x^4 - \frac{3bc^2dx \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{3bcd^2x^2 \log(1+e^{2i(e+fx)})}{f} - \frac{bd^3x^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{bc^3 \log(\cos(e+fx))}{f} + \frac{3ibc^2d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad + \frac{3ibcd^2x \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad + \frac{3ibd^3x^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{3bcd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad - \frac{3bd^3x \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad - \frac{3ibd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
\end{aligned}$$

[In] `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x]), x]`

[Out] `a*c^3*x + (3*a*c^2*d*x^2)/2 + ((3*I)/2)*b*c^2*d*x^2 + a*c*d^2*x^3 + I*b*c*d^2*x^3 + (a*d^3*x^4)/4 + (I/4)*b*d^3*x^4 - (3*b*c^2*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (3*b*c*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^3*x^3*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^3*Log[Cos[e + f*x]])/f + (((3*I)/2)*b*c^2*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + ((3*I)*b*c*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (((3*I)/2)*b*d^3*x^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (3*b*c*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (3*b*d^3*x*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((3*I)/4)*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 499 vs. $2(131) = 262$.

Time = 0.64 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

method	result
risch	$-\frac{ib c^4}{4d} - ib c^3 x + \frac{id^3 b x^4}{4} - \frac{b c^3 \ln(e^{2i(fx+e)}+1)}{f} + \frac{2b c^3 \ln(e^{i(fx+e)})}{f} - \frac{b d^3 \ln(e^{2i(fx+e)}+1)x^3}{f} - \frac{3b d^3 \text{Li}_3(-e^{2i(fx+e)})}{2f^3}$

[In] `int((d*x+c)^3*(a+b*tan(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 3*I/f^2*b*d^2*c*\text{polylog}(2, -\exp(2*I*(f*x+e)))*x - 6*I/f^2*b*d^2*c*e^2*x + 6*I/f^2 \\ & b*d*c^2*e*x + I*d^2*b*c*x^3 - 1/4*I/d*b*c^4 + d^2*a*c*x^3 + 3/2*d*a*c^2*x^2 + a*c^3*x \\ & - I*b*c^3*x + 1/4*I*d^3*b*x^4 - 1/f*b*c^3*\ln(\exp(2*I*(f*x+e)) + 1) + 2/f*b*c^3*\ln(\exp(I*(f*x+e))) - 3/f*b*d^2*c*\ln(\exp(2*I*(f*x+e)) + 1)*x^2 - 3/f*b*d*c^2*\ln(\exp(2*I*(f*x+e)) + 1)*x^3 + 3/2*I/f^2*b*d^3*\text{polylog}(2, -\exp(2*I*(f*x+e)))*x^2 + 3*I/f^2*b*d^2*c^2*x^2 - 4*I/f^3*b*d^2*c*e^3 + 2*I/f^3*b*d^3*x^3 + 3/2*I/f^2*b*d*c^2*\text{polylog}(2, -\exp(2*I*(f*x+e))) + 1/4*d^3*a*x^4 + 1/4/d*a*c^4 + 6/f^3*b*c^2*d^2*c*\ln(\exp(I*(f*x+e))) - 6/f^2*b*c*d*c^2*\ln(\exp(I*(f*x+e))) - 1/f*b*d^3*\ln(\exp(2*I*(f*x+e)) + 1)*x^3 - 3/2/f^3*b*d^3*c*\text{polylog}(3, -\exp(2*I*(f*x+e)))*x^2 - 2/f^4*b*c^3*d^3*\ln(\exp(I*(f*x+e))) - 3/2/f^3*b*d^2*c*\text{polylog}(3, -\exp(2*I*(f*x+e))) + 3/2*I/f^4*b*d^3*c^4 + 3/2*I*d*b*c^2*x^2 - 3/4*I*b*d^3*c*\text{polylog}(4, -\exp(2*I*(f*x+e)))/f^4 \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 500 vs. $2(127) = 254$.

Time = 0.27 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.29

$$\begin{aligned} & \int (c + dx)^3(a + b \tan(e + fx)) dx \\ & = \frac{2 ad^3 f^4 x^4 + 8 acd^2 f^4 x^3 + 12 ac^2 df^4 x^2 + 8 ac^3 f^4 x + 3i bd^3 \text{polylog}\left(4, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3i bd^3 \text{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right)}{f^4} \end{aligned}$$

[In] `integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/8*(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x \\ & *x + 3*I*b*d^3*\text{polylog}(4, (\tan(f*x + e))^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 3*I*b*d^3*\text{polylog}(4, (\tan(f*x + e))^2 - 2*I*\tan(f*x + e) - 1) \\ & /(\tan(f*x + e)^2 + 1)) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x + I*b*c^2*d*f^2)*\text{dilog}(2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x - I*b*c^2*d*f^2)*\text{dilog}(2*(-I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1) + 1) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*f^3*x) \end{aligned}$$

$$\begin{aligned} & d*f^3*x + b*c^3*f^3)*\log(-2*(I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 4* \\ & (b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*\log(-2*(- \\ & I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(b*d^3*f*x + b*c*d^2*f)*\text{polyl} \\ & \text{og}(3, (\tan(f*x + e)^2 + 2*I*\tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(b* \\ & d^3*f*x + b*c*d^2*f)*\text{polylog}(3, (\tan(f*x + e)^2 - 2*I*\tan(f*x + e) - 1)/(\ta \\ & n(f*x + e)^2 + 1)))/f^4 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^3(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx)^3 dx$$

```
[In] integrate((d*x+c)**3*(a+b*tan(f*x+e)),x)
[Out] Integral((a + b*tan(e + f*x))*(c + d*x)**3, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 672 vs. $2(127) = 254$.

Time = 0.72 (sec), antiderivative size = 672, normalized size of antiderivative = 4.42

$$\begin{aligned} & \int (c + dx)^3(a + b \tan(e + fx)) dx \\ & = \frac{12(fx + e)ac^3 + \frac{3(fx+e)^4ad^3}{f^3} - \frac{12(fx+e)^3ad^3e}{f^3} + \frac{18(fx+e)^2ad^3e^2}{f^3} - \frac{12(fx+e)ad^3e^3}{f^3} + \frac{12(fx+e)^3acd^2}{f^2} - \frac{36(fx+e)^2acd^2e}{f^2} + \dots}{\dots} \end{aligned}$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="maxima")
[Out] 1/12*(12*(f*x + e)*a*c^3 + 3*(f*x + e)^4*a*d^3/f^3 - 12*(f*x + e)^3*a*d^3*e/f^3 + 18*(f*x + e)^2*a*d^3*e^2/f^3 - 12*(f*x + e)*a*d^3*e^3/f^3 + 12*(f*x + e)^3*a*c*d^2/f^2 - 36*(f*x + e)^2*a*c*d^2*e/f^2 + 36*(f*x + e)*a*c*d^2*e^2/f^2 + 18*(f*x + e)^2*a*c^2*d/f - 36*(f*x + e)*a*c^2*d*e/f + 12*b*c^3*log(sec(f*x + e)) - 12*b*d^3*e^3*log(sec(f*x + e))/f^3 + 36*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*b*c^2*d*e*log(sec(f*x + e))/f - (-3*I*(f*x + e)^4*b*d^3 + 12*I*b*d^3*polylog(4, -e^(2*I*f*x + 2*I*e)) - 12*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e)^3 - 18*(I*b*d^3*e^2 - 2*I*b*c*d^2*e*f + I*b*c^2*d*f^2)*(f*x + e)^2 - 4*(-4*I*(f*x + e)^3*b*d^3 + 9*(I*b*d^3*e - I*b*c*d^2*f)*(f*x + e)^2 + 9*(-I*b*d^3*e^2 + 2*I*b*c*d^2*e*f - I*b*c^2*d*f^2)*(f*x + e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*d^2*e*f + 3*I*b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog(-e^(2*I*f*x + 2*I*e)) + 2*(4*(f*x + e)^3*b*d^3 - 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2*b*c*d^2*e*f + b*c^2*d*f^2)*(f*x + e)))
```

$$+ e)) * \log(\cos(2fx + 2e)^2 + \sin(2fx + 2e)^2 + 2\cos(2fx + 2e) + 1) + 6*(4*(fx + e)*b*d^3 - 3*b*d^3*e + 3*b*c*d^2*f)*\text{polylog}(3, -e^{(2*I*fx + 2*I*e)})/f^3)/f$$

Giac [F]

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (dx + c)^3 (b \tan(fx + e) + a) dx$$

[In] `integrate((d*x+c)^3*(a+b*tan(f*x+e)),x, algorithm="giac")`
[Out] `integrate((d*x + c)^3*(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^3 dx$$

[In] `int((a + b*tan(e + f*x))*(c + d*x)^3,x)`
[Out] `int((a + b*tan(e + f*x))*(c + d*x)^3, x)`

3.40 $\int (c + dx)^2(a + b \tan(e + fx)) dx$

Optimal result	260
Rubi [A] (verified)	260
Mathematica [A] (verified)	262
Maple [B] (verified)	263
Fricas [B] (verification not implemented)	263
Sympy [F]	264
Maxima [B] (verification not implemented)	264
Giac [F]	265
Mupad [F(-1)]	265

Optimal result

Integrand size = 18, antiderivative size = 115

$$\begin{aligned} \int (c + dx)^2(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^3}{3d} + \frac{ib(c + dx)^3}{3d} - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ibd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \end{aligned}$$

[Out] $1/3*a*(d*x+c)^3/d+1/3*I*b*(d*x+c)^3/d-b*(d*x+c)^2*\ln(1+\exp(2*I*(f*x+e)))/f+$
 $I*b*d*(d*x+c)*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))/f^2-1/2*b*d^2*\operatorname{polylog}(3, -\exp(2*I*(f*x+e)))/f^3$

Rubi [A] (verified)

Time = 0.24 (sec), antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3803, 3800, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int (c + dx)^2(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^3}{3d} + \frac{ibd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ib(c + dx)^3}{3d} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2*(a + b*Tan[e + f*x]), x]$

```
[Out] (a*(c + d*x)^3)/(3*d) + ((I/3)*b*(c + d*x)^3)/d - (b*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f + (I*b*d*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)
```

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*(c_.) + (d_.)*(x_)^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]])]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.)), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_)^(m_.))*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c+dx)^2 + b(c+dx)^2 \tan(e+fx)) \, dx \\
&= \frac{a(c+dx)^3}{3d} + b \int (c+dx)^2 \tan(e+fx) \, dx \\
&= \frac{a(c+dx)^3}{3d} + \frac{ib(c+dx)^3}{3d} - (2ib) \int \frac{e^{2i(e+fx)}(c+dx)^2}{1+e^{2i(e+fx)}} \, dx \\
&= \frac{a(c+dx)^3}{3d} + \frac{ib(c+dx)^3}{3d} - \frac{b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{(2bd) \int (c+dx) \log(1+e^{2i(e+fx)}) \, dx}{f} \\
&= \frac{a(c+dx)^3}{3d} + \frac{ib(c+dx)^3}{3d} - \frac{b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{ibd(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{(ibd^2) \int \text{PolyLog}(2, -e^{2i(e+fx)}) \, dx}{f^2} \\
&= \frac{a(c+dx)^3}{3d} + \frac{ib(c+dx)^3}{3d} - \frac{b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{ibd(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{(bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} \, dx, x, e^{2i(e+fx)}\right)}{2f^3} \\
&= \frac{a(c+dx)^3}{3d} + \frac{ib(c+dx)^3}{3d} - \frac{b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{ibd(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{bd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec), antiderivative size = 191, normalized size of antiderivative = 1.66

$$\begin{aligned}
\int (c+dx)^2(a+b \tan(e+fx)) \, dx &= ac^2x + acdx^2 + ibcdx^2 + \frac{1}{3}ad^2x^3 + \frac{1}{3}ibd^2x^3 \\
&\quad - \frac{2bcdx \log(1+e^{2i(e+fx)})}{f} - \frac{bd^2x^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{bc^2 \log(\cos(e+fx))}{f} + \frac{ibcd \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad + \frac{ibd^2x \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{bd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3}
\end{aligned}$$

[In] `Integrate[(c + d*x)^2*(a + b*Tan[e + f*x]), x]`

[Out] $a*c^2*x + a*c*d*x^2 + I*b*c*d*x^2 + (a*d^2*x^3)/3 + (I/3)*b*d^2*x^3 - (2*b*c*d*x*Log[1 + E^((2*I)*(e + f*x))])/f - (b*d^2*x^2*Log[1 + E^((2*I)*(e + f*x))])/f - (b*c^2*Log[Cos[e + f*x]]/f + (I*b*c*d*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (I*b*d^2*x*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (b*d^2*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3)$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 313 vs. $2(101) = 202$.

Time = 0.58 (sec), antiderivative size = 314, normalized size of antiderivative = 2.73

method	result
risch	$\frac{id^2 b x^3}{3} - ib c^2 x + \frac{d^2 a x^3}{3} + \frac{4ibdcex}{f} + dac x^2 + \frac{2ibdce^2}{f^2} + a c^2 x + \frac{a c^3}{3d} - \frac{b c^2 \ln(e^{2i(fx+e)}+1)}{f} - \frac{2bdcln(e^{2i(fx+e)})}{f}$

[In] `int((d*x+c)^2*(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)`

[Out] $1/3*I*d^2*b*x^3 - I*b*c^2*x^2 + 1/3*d^2*a*x^3 + 4*I/f*b*d*c*e*x + d*a*c*x^2 + 2*I/f^2*b*d*c^2 + a*c^2*x^2 + 1/3/d*a*c^3 - 1/f*b*c^2*ln(exp(2*I*(f*x+e))+1) - 2/f*b*d*c*ln(exp(2*I*(f*x+e))+1)*x + I*d*b*c*x^2 + 2/f^3*b*d^2*x^2*ln(exp(I*(f*x+e))) - 2*I/f^2*b*d^2*x^2 - 1/f*b*d^2*ln(exp(2*I*(f*x+e))+1)*x^2 - 1/2*b*d^2*polylog(3, -exp(2*I*(f*x+e)))/f^3 + I/f^2*b*d*c*polylog(2, -exp(2*I*(f*x+e)))+I/f^2*b*d^2*polylog(2, -exp(2*I*(f*x+e)))*x - 1/3*I/d*b*c^3 - 4/3*I/f^3*b*d^2*x^3 - 4/f^2*b*c*d*e*ln(exp(I*(f*x+e)))+2/f*b*c^2*ln(exp(I*(f*x+e)))$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(98) = 196$.

Time = 0.26 (sec), antiderivative size = 317, normalized size of antiderivative = 2.76

$$\int (c + dx)^2(a + b \tan(e + fx)) dx = \frac{4 ad^2 f^3 x^3 + 12 acdf^3 x^2 + 12 ac^2 f^3 x - 3 bd^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3 bd^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 - 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right)}{12}$$

[In] `integrate((d*x+c)^2*(a+b*tan(f*x+e)), x, algorithm="fricas")`

[Out] $1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x - 3*b*d^2*polylog(3, (\tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 3*b*d^2*polylog(3, (\tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) + 3*b*c^2*f*x^2 + 6*a*c*d*f*x + 3*a*c^2*f*x - 3*b*d^2*polylog(3, (\tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) + 3*b*d^2*polylog(3, (\tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1)) + 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(\tan(f*x + e)^2 + 1))$

$$\begin{aligned} & 1) + 1) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x + b*c^2*f^2)*log(-2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x + b*c^2*f^2)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)))/f^3 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^2(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx)^2 dx$$

```
[In] integrate((d*x+c)**2*(a+b*tan(f*x+e)),x)
[Out] Integral((a + b*tan(e + f*x))*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 373 vs. $2(98) = 196$.

Time = 0.52 (sec), antiderivative size = 373, normalized size of antiderivative = 3.24

$$\begin{aligned} & \int (c + dx)^2(a + b \tan(e + fx)) dx \\ &= \frac{6(fx + e)ac^2 + \frac{2(fx+e)^3ad^2}{f^2} - \frac{6(fx+e)^2ad^2e}{f^2} + \frac{6(fx+e)ad^2e^2}{f^2} + \frac{6(fx+e)^2acd}{f} - \frac{12(fx+e)acde}{f} + 6bc^2 \log(\sec(fx + e))}{f^2} \end{aligned}$$

```
[In] integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="maxima")
[Out] 1/6*(6*(f*x + e)*a*c^2 + 2*(f*x + e)^3*a*d^2/f^2 - 6*(f*x + e)^2*a*d^2*e/f^2
2 + 6*(f*x + e)*a*d^2*e^2/f^2 + 6*(f*x + e)^2*a*c*d/f - 12*(f*x + e)*a*c*d*f
e/f + 6*b*c^2*log(sec(f*x + e)) + 6*b*d^2*e^2*log(sec(f*x + e))/f^2 - 12*b*
c*d*e*log(sec(f*x + e))/f - (-2*I*(f*x + e)^3*b*d^2 + 3*b*d^2*polylog(3, -e
^(2*I*f*x + 2*I*e)) - 6*(-I*b*d^2*e + I*b*c*d*f)*(f*x + e)^2 - 6*(-I*(f*x +
e)^2*b*d^2 + 2*(I*b*d^2*e - I*b*c*d*f)*(f*x + e))*arctan2(sin(2*f*x + 2*e),
cos(2*f*x + 2*e) + 1) - 6*(I*(f*x + e)*b*d^2 - I*b*d^2*e + I*b*c*d*f)*dil
og(-e^(2*I*f*x + 2*I*e)) + 3*((f*x + e)^2*b*d^2 - 2*(b*d^2*e - b*c*d*f)*(f
*x + e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) +
1))/f^2)/f
```

Giac [F]

$$\int (c + dx)^2(a + b \tan(e + fx)) dx = \int (dx + c)^2(b \tan(fx + e) + a) dx$$

[In] `integrate((d*x+c)^2*(a+b*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate((d*x + c)^2*(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx)) (c + dx)^2 dx$$

[In] `int((a + b*tan(e + f*x))*(c + d*x)^2,x)`

[Out] `int((a + b*tan(e + f*x))*(c + d*x)^2, x)`

3.41 $\int (c + dx)(a + b \tan(e + fx)) dx$

Optimal result	266
Rubi [A] (verified)	266
Mathematica [A] (verified)	268
Maple [A] (verified)	268
Fricas [B] (verification not implemented)	268
Sympy [F]	269
Maxima [A] (verification not implemented)	269
Giac [F]	269
Mupad [B] (verification not implemented)	270

Optimal result

Integrand size = 16, antiderivative size = 84

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} - \frac{b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \end{aligned}$$

[Out] $\frac{1}{2}a(d*x+c)^2/d + \frac{1}{2}I*b*(d*x+c)^2/d - b*(d*x+c)*\ln(1+\exp(2*I*(f*x+e)))/f + \frac{1}{2}I*b*d*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))/f^2$

Rubi [A] (verified)

Time = 0.14 (sec), antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.312, Rules used = {3803, 3800, 2221, 2317, 2438}

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx)) dx = & \frac{a(c + dx)^2}{2d} - \frac{b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ib(c + dx)^2}{2d} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Tan}[e + f*x]), x]$

[Out] $(a*(c + d*x)^2)/(2*d) + ((I/2)*b*(c + d*x)^2)/d - (b*(c + d*x)*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f + ((I/2)*b*d*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2$

Rule 2221

$\operatorname{Int}[((F_.)^((g_.)*(e_.) + (f_.)*(x_.)))^((n_.)*(c_.) + (d_.)*(x_.)^(m_.)))/((a_) + (b_.)*(F_.)^((g_.)*(e_.) + (f_.)*(x_.)))^((n_.))), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}$

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*(F_)^((e_)*(c_) + (d_)*(x_)))^n_), x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*(d_) + (e_)*(x_)^n_)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[((c_) + (d_)*(x_))^m_*tan[(e_) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_) + (d_)*(x_))^m_*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c + dx) + b(c + dx)\tan(e + fx)) dx \\
&= \frac{a(c + dx)^2}{2d} + b \int (c + dx)\tan(e + fx) dx \\
&= \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} - (2ib) \int \frac{e^{2i(e+fx)}(c + dx)}{1 + e^{2i(e+fx)}} dx \\
&= \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} - \frac{b(c + dx)\log(1 + e^{2i(e+fx)})}{f} + \frac{(bd) \int \log(1 + e^{2i(e+fx)}) dx}{f} \\
&= \frac{a(c + dx)^2}{2d} + \frac{ib(c + dx)^2}{2d} - \frac{b(c + dx)\log(1 + e^{2i(e+fx)})}{f} \\
&\quad - \frac{(ibd)\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^2}
\end{aligned}$$

$$= \frac{a(c+dx)^2}{2d} + \frac{ib(c+dx)^2}{2d} - \frac{b(c+dx) \log(1+e^{2i(e+fx)})}{f} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 87, normalized size of antiderivative = 1.04

$$\int (c+dx)(a+b\tan(e+fx)) dx = acx + \frac{1}{2}adx^2 + \frac{1}{2}ibdx^2 - \frac{bdx \log(1+e^{2i(e+fx)})}{f} - \frac{bc \log(\cos(e+fx))}{f} + \frac{ibd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2}$$

[In] `Integrate[(c + d*x)*(a + b*Tan[e + f*x]), x]`

[Out] $a*c*x + (a*d*x^2)/2 + (I/2)*b*d*x^2 - (b*d*x*\ln[1 + E^(2*I)*(e + f*x)])/f - (b*c*\ln[\cos(e + f*x)])/f + ((I/2)*b*d*\operatorname{PolyLog}[2, -E^(2*I)*(e + f*x)])/f^2$

Maple [A] (verified)

Time = 0.55 (sec), antiderivative size = 143, normalized size of antiderivative = 1.70

method	result
risch	$\frac{ibd x^2}{2} + \frac{ad x^2}{2} - ibcx + acx - \frac{bc \ln(e^{2i(fx+e)}+1)}{f} + \frac{2bc \ln(e^{i(fx+e)})}{f} + \frac{2ibdex}{f} + \frac{ibde^2}{f^2} - \frac{bd \ln(e^{2i(fx+e)}+1)x}{f} + \frac{ibd e \ln(e^{2i(fx+e)}+1)}{f^3}$

[In] `int((d*x+c)*(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)`

[Out] $1/2*I*b*d*x^2+1/2*a*d*x^2-I*b*c*x+a*c*x-1/f*b*c*\ln(\exp(2*I*(f*x+e))+1)+2/f*b*c*\ln(\exp(I*(f*x+e)))+2*I/f*b*d*e*x+I/f^2*b*d*e^2-1/f*b*d*\ln(\exp(2*I*(f*x+e))+1)*x+1/2*I*b*d*\operatorname{polylog}(2, -\exp(2*I*(f*x+e)))/f^2-2/f^2*b*d*e*\ln(\exp(I*(f*x+e)))$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(69) = 138$.

Time = 0.26 (sec), antiderivative size = 160, normalized size of antiderivative = 1.90

$$\int (c+dx)(a+b\tan(e+fx)) dx = \frac{2 adf^2 x^2 + 4 acf^2 x - i bd \operatorname{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i bd \operatorname{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) - 2(bdfx + bcf) \log\left(-\frac{2 i \tan(fx+e)}{\tan(fx+e)^2+1}\right)}{4f^2}$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="fricas")`

[Out]
$$\frac{1}{4} \left(2a^2 f^2 x^2 + 4a c f^2 x - b d \operatorname{dilog}(2(\tan(f x) + e) - 1) \right) / (\tan(f x) + e)^2 + \frac{1}{4} + b d \operatorname{dilog}(2(-\tan(f x) + e) - 1) / (\tan(f x) + e)^2 + 1 - 2(b d f x + b c f) \log(-2(\tan(f x) + e) - 1) / (\tan(f x) + e)^2 - 2(b d f x + b c f) \log(-2(-\tan(f x) + e) - 1) / (\tan(f x) + e)^2 + 1 \right) / f^2$$

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (a + b \tan(e + fx))(c + dx) dx$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e)),x)`

[Out] `Integral((a + b*tan(e + f*x))*(c + d*x), x)`

Maxima [A] (verification not implemented)

none

Time = 0.38 (sec) , antiderivative size = 130, normalized size of antiderivative = 1.55

$$\int (c + dx)(a + b \tan(e + fx)) dx = \frac{(a + i b) d f^2 x^2 + 2(a + i b) c f^2 x + i b d \operatorname{Li}_2(-e^{(2 i f x + 2 i e)}) + 2(-i b d f x - i b c f) \arctan(\sin(2 f x + 2 e), \cos(2 f x + 2 e))}{2 f^2}$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="maxima")`

[Out]
$$\frac{1}{2} \left((a + I b) d f^2 x^2 + 2(a + I b) c f^2 x + b d \operatorname{dilog}(-e^{(2 I f x + 2 I e)}) + 2(-I b d f x - I b c f) \arctan(\sin(2 f x + 2 e), \cos(2 f x + 2 e)) + 1 - (b d f x + b c f) \log(\cos(2 f x + 2 e)^2 + \sin(2 f x + 2 e)^2 + 2 \cos(2 f x + 2 e) + 1) \right) / f^2$$

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx)) dx = \int (dx + c)(b \tan(f x + e) + a) dx$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate((d*x + c)*(b*tan(f*x + e) + a), x)`

Mupad [B] (verification not implemented)

Time = 3.60 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.92

$$\int (c + dx)(a + b \tan(e + fx)) dx = \frac{ax(2c + dx)}{2} - \frac{bd(\pi \ln(\cos(fx)) - \pi \ln(e^{fx^2i} + 1) + \text{polylog}(2, -e^{-e^2i} e^{-fx^2i}) 1i - \pi \ln(e^{-e^2i} e^{-fx^2i} + 1) + 2e \ln(e^{fx^2i} + 1)))}{2f^2} + \frac{bc \ln(\tan(e + fx)^2 + 1)}{2f}$$

[In] `int((a + b*tan(e + f*x))*(c + d*x),x)`

[Out] `(a*x*(2*c + d*x))/2 - (b*d*(polylog(2, -exp(-e*2i)*exp(-f*x*2i))*1i - pi*log(exp(f*x*2i) + 1) - log(cos(e + f*x))*(2*e - pi) - pi*log(exp(-e*2i)*exp(-f*x*2i) + 1) + 2*e*log(exp(-e*2i)*exp(-f*x*2i) + 1) + pi*log(cos(f*x)) + f^2*x^2*1i + 2*f*x*log(exp(-e*2i)*exp(-f*x*2i) + 1) + e*f*x*2i))/(2*f^2) + (b*c*log(tan(e + f*x)^2 + 1))/(2*f)`

3.42 $\int \frac{a+b \tan(e+fx)}{c+dx} dx$

Optimal result	271
Rubi [N/A]	271
Mathematica [N/A]	272
Maple [N/A] (verified)	272
Fricas [N/A]	272
Sympy [N/A]	272
Maxima [N/A]	273
Giac [N/A]	273
Mupad [N/A]	273

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

[In] Int[(a + b*Tan[e + f*x])/(c + d*x),x]

[Out] Defer[Int][(a + b*Tan[e + f*x])/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 1.91 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

[In] `Integrate[(a + b*Tan[e + f*x])/(c + d*x), x]`

[Out] `Integrate[(a + b*Tan[e + f*x])/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{dx + c} dx$$

[In] `int((a+b*tan(f*x+e))/(d*x+c), x)`

[Out] `int((a+b*tan(f*x+e))/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b*tan(f*x + e) + a)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 0.66 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + fx)}{c + dx} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c), x)`

[Out] `Integral((a + b*tan(e + f*x))/(c + d*x), x)`

Maxima [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 86, normalized size of antiderivative = 4.78

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="maxima")`

[Out] `(2*b*d*integrate(sin(2*f*x + 2*e)/((d*x + c)*cos(2*f*x + 2*e)^2 + (d*x + c)*sin(2*f*x + 2*e)^2 + d*x + 2*(d*x + c)*cos(2*f*x + 2*e) + c), x) + a*log(d*x + c))/d`

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{b \tan(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c),x, algorithm="giac")`

[Out] `integrate((b*tan(f*x + e) + a)/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 3.53 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{c + dx} dx = \int \frac{a + b \tan(e + f x)}{c + d x} dx$$

[In] `int((a + b*tan(e + f*x))/(c + d*x),x)`

[Out] `int((a + b*tan(e + f*x))/(c + d*x), x)`

3.43 $\int \frac{a+b \tan(e+fx)}{(c+dx)^2} dx$

Optimal result	274
Rubi [N/A]	274
Mathematica [N/A]	275
Maple [N/A] (verified)	275
Fricas [N/A]	275
Sympy [N/A]	275
Maxima [N/A]	276
Giac [N/A]	276
Mupad [N/A]	276

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \text{Int}\left(\frac{a + b \tan(e + fx)}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

[In] Int[(a + b*Tan[e + f*x])/((c + d*x)^2),x]

[Out] Defer[Int][(a + b*Tan[e + f*x])/((c + d*x)^2), x]

Rubi steps

$$\text{integral} = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 5.10 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

```
[In] Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2, x]
[Out] Integrate[(a + b*Tan[e + f*x])/(c + d*x)^2, x]
```

Maple [N/A] (verified)

Not integrable

Time = 0.22 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tan(fx + e)}{(dx + c)^2} dx$$

```
[In] int((a+b*tan(f*x+e))/(d*x+c)^2, x)
[Out] int((a+b*tan(f*x+e))/(d*x+c)^2, x)
```

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))/(d*x+c)^2, x, algorithm="fricas")
[Out] integral((b*tan(f*x + e) + a)/(d^2*x^2 + 2*c*d*x + c^2), x)
```

Sympy [N/A]

Not integrable

Time = 2.49 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))/(d*x+c)**2, x)
[Out] Integral((a + b*tan(e + f*x))/(c + d*x)**2, x)
```

Maxima [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 142, normalized size of antiderivative = 7.89

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="maxima")`

[Out] `(2*(b*d^2*x + b*c*d)*integrate(sin(2*f*x + 2*e)/(d^2*x^2 + 2*c*d*x + (d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e))^2 + (d^2*x^2 + 2*c*d*x + c^2)*sin(2*f*x + 2*e)^2 + c^2 + 2*(d^2*x^2 + 2*c*d*x + c^2)*cos(2*f*x + 2*e)), x) - a)/(d^2*x + c*d)`

Giac [N/A]

Not integrable

Time = 4.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{b \tan(fx + e) + a}{(dx + c)^2} dx$$

[In] `integrate((a+b*tan(f*x+e))/(d*x+c)^2,x, algorithm="giac")`

[Out] `integrate((b*tan(f*x + e) + a)/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 3.68 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tan(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tan(e + f x)}{(c + d x)^2} dx$$

[In] `int((a + b*tan(e + f*x))/(c + d*x)^2,x)`

[Out] `int((a + b*tan(e + f*x))/(c + d*x)^2, x)`

3.44 $\int (c + dx)^3 (a + b \tan(e + fx))^2 dx$

Optimal result	277
Rubi [A] (verified)	278
Mathematica [B] (warning: unable to verify)	282
Maple [B] (verified)	283
Fricas [B] (verification not implemented)	284
Sympy [F]	285
Maxima [B] (verification not implemented)	285
Giac [F]	286
Mupad [F(-1)]	287

Optimal result

Integrand size = 20, antiderivative size = 300

$$\begin{aligned} \int (c + dx)^3 (a + b \tan(e + fx))^2 dx = & -\frac{ib^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} + \frac{iab(c + dx)^4}{2d} \\ & - \frac{b^2(c + dx)^4}{4d} + \frac{3b^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\ & - \frac{2ab(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\ & - \frac{3ib^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & + \frac{3iabd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & + \frac{3b^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\ & - \frac{3abd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & - \frac{3iabd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} \\ & + \frac{b^2(c + dx)^3 \tan(e + fx)}{f} \end{aligned}$$

```
[Out] -I*b^2*(d*x+c)^3/f+1/4*a^2*(d*x+c)^4/d+1/2*I*a*b*(d*x+c)^4/d-1/4*b^2*(d*x+c)^4/d+3*b^2*d*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f-3*I*b^2*d^2*(d*x+c)*polylog(2, -exp(2*I*(f*x+e)))/f^3+3*I*a*b*d*(d*x+c)^2*polylog(2, -exp(2*I*(f*x+e)))/f^2+3/2*b^2*d^3*polylog(3, -exp(2*I*(f*x+e)))/f^4-3*a*b*d^2*(d*x+c)*polylog(3, -exp(2*I*(f*x+e)))/f^3-3/2*I*a*b*d^3*polylog(4, -exp(2*I*(f*x+e)))/f^4+b^2*(d*x+c)^3*tan(f*x+e)/f
```

Rubi [A] (verified)

Time = 0.59 (sec) , antiderivative size = 300, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used = {3803, 3800, 2221, 2611, 6744, 2320, 6724, 3801, 32}

$$\begin{aligned} \int (c + dx)^3 (a + b \tan(e + fx))^2 dx = & \frac{a^2(c + dx)^4}{4d} - \frac{3abd^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & + \frac{3iabd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{2ab(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{iab(c + dx)^4}{2d} - \frac{3iabd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} \\ & - \frac{3ib^2d^2(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & + \frac{3b^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\ & + \frac{b^2(c + dx)^3 \tan(e + fx)}{f} - \frac{ib^2(c + dx)^3}{f} \\ & - \frac{b^2(c + dx)^4}{4d} + \frac{3b^2d^3 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \end{aligned}$$

[In] `Int[(c + d*x)^3*(a + b*Tan[e + f*x])^2, x]`

[Out] $\frac{((-I)*b^2*(c + d*x)^3)/f + (a^2*(c + d*x)^4)/(4*d) + ((I/2)*a*b*(c + d*x)^4)/(d) - (b^2*(c + d*x)^4)/(4*d) + (3*b^2*d*(c + d*x)^2*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f^2 - (2*a*b*(c + d*x)^3*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f - ((3*I)*b^2*d^2*(c + d*x)*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^3 + ((3*I)*a*b*d*(c + d*x)^2*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 + (3*b^2*d^3*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/f^4 - (3*a*b*d^2*(c + d*x)*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/f^3 - (((3*I)/2)*a*b*d^3*\operatorname{PolyLog}[4, -E^{((2*I)*(e + f*x))}])/f^4 + (b^2*(c + d*x)^3*\operatorname{Tan}[e + f*x])/f}$

Rule 32

```
Int[((a_) + (b_)*(x_))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^((g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*\operatorname{Log}[F])), Int[(c + d*x)^(m - 1)*\operatorname{Log}[1 + b*((F^((g*(e + f*x))))^n/a)], x]]
```

```
)n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_.) + (b_)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*((a_.) + (b_)*(x_))^(p_.)]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```

Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.*((a_.) + (b_.
)*(x_))))^p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a +
b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx)^3 + 2ab(c+dx)^3 \tan(e+fx) + b^2(c+dx)^3 \tan^2(e+fx)) \, dx \\
&= \frac{a^2(c+dx)^4}{4d} + (2ab) \int (c+dx)^3 \tan(e+fx) \, dx + b^2 \int (c+dx)^3 \tan^2(e+fx) \, dx \\
&= \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad - (4iab) \int \frac{e^{2i(e+fx)}(c+dx)^3}{1+e^{2i(e+fx)}} \, dx - b^2 \int (c+dx)^3 \, dx \\
&\quad - \frac{(3b^2d) \int (c+dx)^2 \tan(e+fx) \, dx}{f} \\
&= -\frac{ib^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} - \frac{b^2(c+dx)^4}{4d} \\
&\quad - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad + \frac{(6abd) \int (c+dx)^2 \log(1+e^{2i(e+fx)}) \, dx}{f} + \frac{(6ib^2d) \int \frac{e^{2i(e+fx)}(c+dx)^2}{1+e^{2i(e+fx)}} \, dx}{f} \\
&= -\frac{ib^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} - \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3iabd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad - \frac{(6iabd^2) \int (c+dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)}) \, dx}{f^2} \\
&\quad - \frac{(6b^2d^2) \int (c+dx) \log(1+e^{2i(e+fx)}) \, dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{ib^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} - \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{3ib^2d^2(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{3iabd(c+dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3abd^2(c+dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad + \frac{(3abd^3) \int \text{PolyLog}(3, -e^{2i(e+fx)}) dx}{f^3} + \frac{(3ib^2d^3) \int \text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^3} \\
&= -\frac{ib^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} - \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{3ib^2d^2(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{3iabd(c+dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3abd^2(c+dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad - \frac{(3iabd^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^4} \\
&\quad + \frac{(3b^2d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^4} \\
&= -\frac{ib^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} + \frac{iab(c+dx)^4}{2d} - \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{3ib^2d^2(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{3iabd(c+dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad + \frac{3b^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} - \frac{3abd^2(c+dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\
&\quad - \frac{3iabd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{2f^4} + \frac{b^2(c+dx)^3 \tan(e+fx)}{f}
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1337 vs. $2(300) = 600$.

Time = 7.33 (sec) , antiderivative size = 1337, normalized size of antiderivative = 4.46

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\
 &= \frac{i b^2 d^3 e^{-ie} (2 f^2 x^2 (2 f x - 3 i (1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6 (1 + e^{2ie}) f x \text{PolyLog}(2, -e^{-2i(e+fx)}) - 3 i (1 + e^{2ie}) f^2 x^2 \log(1 + e^{-2i(e+fx)}))}{4 f^4} \\
 & - \frac{i a b c d^2 e^{-ie} (2 f^2 x^2 (2 f x - 3 i (1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6 (1 + e^{2ie}) f x \text{PolyLog}(2, -e^{-2i(e+fx)}) - 3 i (1 + e^{2ie}) f^2 x^2 \log(1 + e^{-2i(e+fx)}))}{2 f^3} \\
 & - \frac{i a b d^3 e^{ie} (2 e^{-2ie} f^4 x^4 - 4 i (1 + e^{-2ie}) f^3 x^3 \log(1 + e^{-2i(e+fx)}) + 6 (1 + e^{-2ie}) f^2 x^2 \text{PolyLog}(2, -e^{-2i(e+fx)}))}{4 f^4} \\
 & + \frac{3 b^2 c^2 d \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f^2 (\cos^2(e) + \sin^2(e))} \\
 & - \frac{2 a b c^3 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f (\cos^2(e) + \sin^2(e))} \\
 & + \frac{3 b^2 c d^2 \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))}))}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \\
 & - \frac{3 a b c^2 d \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))}))}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \\
 & + \frac{\sec(e) \sec(e + f x) (4 a^2 c^3 f x \cos(f x) - 4 b^2 c^3 f x \cos(f x) + 6 a^2 c^2 d f x^2 \cos(f x) - 6 b^2 c^2 d f x^2 \cos(f x) + 4 a^2 c^2 d^2 f x^3 \cos(f x))}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}}
 \end{aligned}$$

[In] `Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^2,x]`

[Out]
$$\begin{aligned}
 & ((I/4)*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) \\
 & - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^4) - ((I/2)*a*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) \\
 & - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) - ((I/4)*a*b*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e)) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))] + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))] - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e])/f^4 + (3*b^2*c^2*d*Sec[e]*(Cos[e]*Log[Cos[e]]*Cos[f*x] - Sin[e]*Sin[f*x]) + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (2*a*b*c^3*Sec[e]*(Cos[e]*Log[Cos[e]]*Cos[f*x] - Sin[e]*Sin[f*x]) + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (3*b^2*c*d^2*2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)]))
 \end{aligned}$$

$$\begin{aligned}
& *x]) - 2*(f*x - \text{ArcTan}[\text{Cot}[e]])*\text{Log}[1 - E^((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]]))] + \\
& \text{Pi}*\text{Log}[\text{Cos}[f*x] - 2*\text{ArcTan}[\text{Cot}[e]]*\text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + I*\text{PolyLog}[2, E^((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]])))]/\text{Sqrt}[1 + \text{Cot}[e]^2]*\text{Sec}[e]/(f^3*\text{Sqrt}[\text{Csc}[e]^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)]) - (3*a*b*c^2*d*\text{Csc}[e]*((f^2*x^2)/E^((I*\text{ArcTan}[\text{Cot}[e]])) - (\text{Cot}[e]*(I*f*x*(-\text{Pi} - 2*\text{ArcTan}[\text{Cot}[e]])) - \text{Pi}*\text{Log}[1 + E^((-2*I)*f*x)] - 2*(f*x - \text{ArcTan}[\text{Cot}[e]])*\text{Log}[1 - E^((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]]))]) + \text{Pi}*\text{Log}[\text{Cos}[f*x] - 2*\text{ArcTan}[\text{Cot}[e]]*\text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + I*\text{PolyLog}[2, E^((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]])))]/\text{Sqrt}[1 + \text{Cot}[e]^2]*\text{Sec}[e]/(f^2*\text{Sqrt}[\text{Csc}[e]^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)]) + (\text{Sec}[e]*\text{Sec}[e + f*x]*(4*a^2*c^3*f*x*\text{Cos}[f*x] - 4*b^2*c^3*f*x*\text{Cos}[f*x] + 6*a^2*c^2*d*f*x^2*\text{Cos}[f*x] - 6*b^2*c^2*d*f*x^2*\text{Cos}[f*x] + 4*a^2*c*d^2*f*x^3*\text{Cos}[f*x] - 4*b^2*c*d^2*f*x^3*\text{Cos}[f*x] + a^2*d^3*f*x^4*\text{Cos}[f*x] - b^2*d^3*f*x^4*\text{Cos}[f*x] + 4*a^2*c^3*f*x*\text{Cos}[2*e + f*x] - 4*b^2*c^3*f*x*\text{Cos}[2*e + f*x] + 6*a^2*c^2*d*f*x^2*\text{Cos}[2*e + f*x] - 6*b^2*c^2*d*f*x^2*\text{Cos}[2*e + f*x] + 4*a^2*c*d^2*f*x^3*\text{Cos}[2*e + f*x] - 4*b^2*c*d^2*f*x^3*\text{Cos}[2*e + f*x] + a^2*d^3*f*x^4*\text{Cos}[2*e + f*x] - b^2*d^3*f*x^4*\text{Cos}[2*e + f*x] + 8*b^2*c^3*\text{Sin}[f*x] + 24*b^2*c^2*d*x*\text{Sin}[f*x] - 8*a*b*c^3*f*x*\text{Sin}[f*x] + 24*b^2*c*d^2*x^2*\text{Sin}[f*x] - 12*a*b*c^2*d*f*x^2*\text{Sin}[f*x] + 8*b^2*d^3*x^3*\text{Sin}[f*x] - 8*a*b*c*d^2*f*x^3*\text{Sin}[f*x] - 2*a*b*d^3*f*x^4*\text{Sin}[f*x] + 8*a*b*c^3*f*x*\text{Sin}[2*e + f*x] + 12*a*b*c^2*d*f*x^2*\text{Sin}[2*e + f*x] + 8*a*b*c*d^2*f*x^3*\text{Sin}[2*e + f*x] + 2*a*b*d^3*f*x^4*\text{Sin}[2*e + f*x]))/(8*f)
\end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 951 vs. 2(271) = 542.

Time = 1.60 (sec), antiderivative size = 952, normalized size of antiderivative = 3.17

method	result	size
risch	Expression too large to display	952

```

[In] int((d*x+c)^3*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] -12*I/f^2*b*e^2*d^2*c*a*x+6*I/f^2*b*a*c*d^2*polylog(2,-exp(2*I*(f*x+e)))*x+
12*I/f*b*d*c^2*a*e*x+2*I*b^2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/f/(exp(2*I*(f*x+e))+1)+3/f^2*b^2*c^2*d*ln(exp(2*I*(f*x+e))+1)-6/f^2*b^2*c^2*d*ln(exp(I*(f*x+e)))+3/f^2*b^2*d^3*ln(exp(2*I*(f*x+e))+1)*x^2-2/f*b*a*c^3*ln(exp(2*I*(f*x+e))+1)+4/f*b*a*c^3*ln(exp(I*(f*x+e)))-6/f^4*b^2*e^2*d^3*ln(exp(I*(f*x+e)))-2*I/f*b^2*d^3*x^3+4*I/f^4*b^2*d^3*e^3+1/2*I*d^3*a*b*x^4-3/2*d*b^2*c^2*x^2+1/4*d^3*a^2*x^4+1/4*d*a^2*c^4-1/4*d^3*b^2*x^4-b^2*c^3*x-1/4/d*b^2*c^4-2*I*a*b*c^3*x-1/2*I/d*a*b*c^4+d^2*a^2*c*x^3+3/2*d*a^2*c^2*x^2+a^2*c^3*x-d^2*b^2*c*x^3-3/2*I*a*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-4/f^4*b*e^3*a*d^3*ln(exp(I*(f*x+e)))-3/f^3*b*d^3*a*polylog(3,-exp(2*I*(f*x+e)))*x+6/f^2*b^2*d^2*c*ln(exp(2*I*(f*x+e))+1)*x-2/f*b*d^3*a*ln(exp(2*I*(f*x+e))+1)*x^3-3/f^3*b*a*c*d^2*polylog(3,-exp(2*I*(f*x+e)))+12/f^3*b^2*e*c*d^2*ln(exp(I*(f*x+e)))

```

$$\begin{aligned} &) - 3*I/f^3*b^2*d^3*polylog(2, -exp(2*I*(f*x+e)))*x + 3*I/f^4*b*e^4*a*d^3 - 6*I/f \\ & *b^2*d^2*c*x^2 - 6*I/f^3*b^2*d^2*c*e^2 - 3*I/f^3*b^2*d^2*c*polylog(2, -exp(2*I*(f*x+e))) + 6*I/f^3*b^2*d^3*e^2*x + 2*I*d^2*a*b*c*x^3 + 3*I*d*a*b*c^2*x^2 + 3/2*b^2*d^3*polylog(3, -exp(2*I*(f*x+e)))/f^4 - 12/f^2*b*e*a*c^2*d*ln(exp(I*(f*x+e))) - 6/f*b*d*c^2*a*ln(exp(2*I*(f*x+e))+1)*x - 6/f*b*a*c*d^2*ln(exp(2*I*(f*x+e))+1)*x^2 + 12/f^3*b*e^2*a*c*d^2*ln(exp(I*(f*x+e)))+6*I/f^2*b*d*c^2*a*e^2 + 3*I/f^2*b*d^3*a*polylog(2, -exp(2*I*(f*x+e)))+3*I/f^2*b*d^3*a*polylog(2, -exp(2*I*(f*x+e)))*x^2 + 4*I/f^3*b*e^3*a*d^3*x - 8*I/f^3*b*e^3*d^2*c*a - 12*I/f^2*b^2*d^2*c^2*e*x \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 759 vs. 2(264) = 528.

Time = 0.28 (sec) , antiderivative size = 759, normalized size of antiderivative = 2.53

$$\begin{aligned} & \int (c + dx)^3 (a + b \tan(e + fx))^2 dx \\ & = \frac{(a^2 - b^2)d^3 f^4 x^4 + 4(a^2 - b^2)cd^2 f^4 x^3 + 6(a^2 - b^2)c^2 df^4 x^2 + 4(a^2 - b^2)c^3 f^4 x + 3i abd^3 \text{polylog}\left(4, \frac{\tan(fx+e)}{\tan(fx+e)}\right)}{(a^2 - b^2)d^3 f^4 x^4 + 4(a^2 - b^2)cd^2 f^4 x^3 + 6(a^2 - b^2)c^2 df^4 x^2 + 4(a^2 - b^2)c^3 f^4 x + 3i abd^3 \text{polylog}\left(4, \frac{\tan(fx+e)}{\tan(fx+e)}\right)} \end{aligned}$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="fricas")
[Out] 1/4*((a^2 - b^2)*d^3*f^4*x^4 + 4*(a^2 - b^2)*c*d^2*f^4*x^3 + 6*(a^2 - b^2)*c^2*d*f^4*x^2 + 4*(a^2 - b^2)*c^3*f^4*x + 3*I*a*b*d^3*polylog(4, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*a*b*d^3*polylog(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*(I*a*b*d^3*f^4*x^2 + I*a*b*c^2*d*f^2 - I*b^2*c*d^2*f + I*(2*a*b*c*d^2*f^2 - b^2*d^3*f)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-I*a*b*d^3*f^2*x^2 - I*a*b*c^2*d*f^2 + I*b^2*c*d^2*f - I*(2*a*b*c*d^2*f^2 - b^2*d^3*f)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 - 3*b^2*c^2*d^2*f^2 + 3*(2*a*b*c*d^2*f^3 - b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 - b^2*c*d^2*f^2)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 - 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3 - b^2*d^3*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f - b^2*d^3)*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 4*(b^2*d^3*f^3*x^3 + 3*b^2*c*d^2*f^3*x^2 + 3*b^2*c^2*d*f^3*x + b^2*c^3*f^4*x)/f^4
```

Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^3 dx$$

```
[In] integrate((d*x+c)**3*(a+b*tan(f*x+e))**2,x)
[Out] Integral((a + b*tan(e + f*x))**2*(c + d*x)**3, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2525 vs. $2(264) = 528$.

Time = 1.73 (sec), antiderivative size = 2525, normalized size of antiderivative = 8.42

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="maxima")
[Out] 1/4*(4*(f*x + e)*a^2*c^3 + (f*x + e)^4*a^2*d^3/f^3 - 4*(f*x + e)^3*a^2*d^3*f/e/f^3 + 6*(f*x + e)^2*a^2*d^3*e^2/f^3 - 4*(f*x + e)*a^2*d^3*e^3/f^3 + 4*(f*x + e)^3*a^2*c*d^2/f^2 - 12*(f*x + e)^2*a^2*c*d^2*e/f^2 + 12*(f*x + e)*a^2*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^2*c^2*d/f - 12*(f*x + e)*a^2*c^2*d*e/f + 8*a*b*c^3*log(sec(f*x + e)) - 8*a*b*d^3*e^3*log(sec(f*x + e))/f^3 + 24*a*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 24*a*b*c^2*d*e*log(sec(f*x + e))/f + 4*(3*(2*a*b + I*b^2)*(f*x + e)^4*d^3 - 24*b^2*d^3*e^3 + 72*b^2*c*d^2*e^2*f - 72*b^2*c^2*d*e*f^2 + 24*b^2*c^3*f^3 - 12*((2*a*b + I*b^2)*d^3*e - (2*a*b + I*b^2)*c*d^2*f)*(f*x + e)^3 + 18*((2*a*b + I*b^2)*d^3*e^2 - 2*(2*a*b + I*b^2)*c*d^2*e*f + (2*a*b + I*b^2)*c^2*d*f^2)*(f*x + e)^2 + 12*(-I*b^2*d^3*e^3 + 3*I*b^2*c*d^2*e^2*f - 3*I*b^2*c^2*d*e*f^2 + I*b^2*c^3*f^3)*(f*x + e) - 4*(8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2*c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a*b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f*x + e) + (8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e*f - 9*b^2*c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 + 18*(a*b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f)*(f*x + e))*cos(2*f*x + 2*e) - (-8*I*(f*x + e)^3*a*b*d^3 + 9*I*b^2*d^3*e^2 - 18*I*b^2*c*d^2*e*f + 9*I*b^2*c^2*d*f^2 + 9*(2*I*a*b*d^3*e - 2*I*a*b*c*d^2*f + I*b^2*d^3)*(f*x + e)^2 + 18*(-I*a*b*d^3*e^2 - I*a*b*c^2*d*f^2 - I*b^2*d^3*e + (2*I*a*b*c*d^2*e + I*b^2*c*d^2)*f)*(f*x + e))*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + 3*((2*a*b + I*b^2)*(f*x + e)^4*d^3 - 4*(2*b^2*d^3 + (2*a*b + I*b^2)*d^3*e - (2*a*b + I*b^2)*c*d^2*f)*(f*x + e)^3 + 6*(4*b^2*d^3*e + (2*a*b + I*b^2)*d^3*e^2 + (2*a*b + I*b^2)*c^2*d*f^2 - 2*(2*b^2*c*d^2*
```

$$\begin{aligned}
& 2 + (2*a*b + I*b^2)*c*d^2*e*f)*(f*x + e)^2 + 4*(-I*b^2*d^3*e^3 + I*b^2*c^3 \\
& *f^3 - 6*b^2*d^3*e^2 + 3*(-I*b^2*c^2*d*e - 2*b^2*c^2*d)*f^2 + 3*(I*b^2*c*d^2 \\
& *e^2 + 4*b^2*c*d^2*e)*f)*(f*x + e))*\cos(2*f*x + 2*e) + 12*(4*(f*x + e)^2*a \\
& *b*d^3 + 3*a*b*d^3*e^2 + 3*a*b*c^2*d*f^2 + 3*b^2*d^3*e - 3*(2*a*b*d^3*e - 2 \\
& *a*b*c*d^2*f + b^2*d^3)*(f*x + e) - 3*(2*a*b*c*d^2*e + b^2*c*d^2)*f + (4*(f \\
& *x + e)^2*a*b*d^3 + 3*a*b*d^3*e^2 + 3*a*b*c^2*d*f^2 + 3*b^2*d^3*e - 3*(2*a* \\
& b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e) - 3*(2*a*b*c*d^2*e + b^2*c*d^2 \\
&)*f)*\cos(2*f*x + 2*e) + (4*I*(f*x + e)^2*a*b*d^3 + 3*I*a*b*d^3*e^2 + 3*I*a* \\
& b*c^2*d*f^2 + 3*I*b^2*d^3*e + 3*(-2*I*a*b*d^3*e + 2*I*a*b*c*d^2*f - I*b^2*d \\
& ^3)*(f*x + e) + 3*(-2*I*a*b*c*d^2*e - I*b^2*c*d^2)*f)*\sin(2*f*x + 2*e))*\text{dil} \\
& \text{og}(-e^2*(2*I*f*x + 2*I*e)) + 2*(8*I*(f*x + e)^3*a*b*d^3 - 9*I*b^2*d^3*e^2 + 1 \\
& 8*I*b^2*c*d^2*e*f - 9*I*b^2*c^2*d*f^2 + 9*(-2*I*a*b*d^3*e + 2*I*a*b*c*d^2*f \\
& - I*b^2*d^3)*(f*x + e)^2 + 18*(I*a*b*d^3*e^2 + I*a*b*c^2*d*f^2 + I*b^2*d^3 \\
& *e + (-2*I*a*b*c*d^2*e - I*b^2*c*d^2)*f)*(f*x + e) + (8*I*(f*x + e)^3*a*b*d \\
& ^3 - 9*I*b^2*d^3*e^2 + 18*I*b^2*c*d^2*e*f - 9*I*b^2*c^2*d*f^2 + 9*(-2*I*a*b* \\
& d^3*e + 2*I*a*b*c*d^2*f - I*b^2*d^3)*(f*x + e)^2 + 18*(I*a*b*d^3*e^2 + I*a* \\
& b*c^2*d*f^2 + I*b^2*d^3*e + (-2*I*a*b*c*d^2*e - I*b^2*c*d^2)*f)*(f*x + e)) \\
& *\cos(2*f*x + 2*e) - (8*(f*x + e)^3*a*b*d^3 - 9*b^2*d^3*e^2 + 18*b^2*c*d^2*e \\
& *f - 9*b^2*c^2*d*f^2 - 9*(2*a*b*d^3*e - 2*a*b*c*d^2*f + b^2*d^3)*(f*x + e)^2 \\
& + 18*(a*b*d^3*e^2 + a*b*c^2*d*f^2 + b^2*d^3*e - (2*a*b*c*d^2*e + b^2*c*d^2)*f) \\
& *(f*x + e))*\sin(2*f*x + 2*e))*\log(\cos(2*f*x + 2*e)^2 + \sin(2*f*x + 2*e) \\
&)^2 + 2*\cos(2*f*x + 2*e) + 1) - 24*(a*b*d^3*\cos(2*f*x + 2*e) + I*a*b*d^3*\sin \\
& (2*f*x + 2*e) + a*b*d^3)*\text{polylog}(4, -e^2*(2*I*f*x + 2*I*e)) + 6*(8*I*(f*x + \\
& e)*a*b*d^3 - 6*I*a*b*d^3*e + 6*I*a*b*c*d^2*f - 3*I*b^2*d^3 + (8*I*(f*x + e) \\
& *a*b*d^3 - 6*I*a*b*d^3*e + 6*I*a*b*c*d^2*f - 3*I*b^2*d^3)*\cos(2*f*x + 2*e) \\
& - (8*(f*x + e)*a*b*d^3 - 6*a*b*d^3*e + 6*a*b*c*d^2*f - 3*b^2*d^3)*\sin(2*f*x \\
& + 2*e))*\text{polylog}(3, -e^2*(2*I*f*x + 2*I*e)) + 3*((2*I*a*b - b^2)*(f*x + e)^4 \\
& *d^3 + 4*(-2*I*b^2*d^3 + (-2*I*a*b + b^2)*d^3*e + (2*I*a*b - b^2)*c*d^2*f)*(\\
& f*x + e)^3 + 6*(4*I*b^2*d^3*e + (2*I*a*b - b^2)*d^3*e^2 + (2*I*a*b - b^2)*c \\
& ^2*d*f^2 + 2*(-2*I*b^2*c*d^2 + (-2*I*a*b + b^2)*c*d^2*f)*f)*(f*x + e)^2 + 4 \\
& *(b^2*d^3*e^3 - b^2*c^3*f^3 - 6*I*b^2*d^3*e^2 + 3*(b^2*c^2*d*e - 2*I*b^2*c \\
& ^2*d)*f^2 - 3*(b^2*c*d^2*e^2 - 4*I*b^2*c*d^2*f)*f)*(f*x + e))*\sin(2*f*x + 2* \\
& e))/(-12*I*f^3*\cos(2*f*x + 2*e) + 12*f^3*\sin(2*f*x + 2*e) - 12*I*f^3))/f
\end{aligned}$$

Giac [F]

$$\int (c + dx)^3 (a + b \tan(f*x + e))^2 dx = \int (dx + c)^3 (b \tan(f*x + e) + a)^2 dx$$

[In] `integrate((d*x+c)^3*(a+b*tan(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)^3*(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + f x))^2 (c + d x)^3 dx$$

[In] `int((a + b*tan(e + f*x))^2*(c + d*x)^3,x)`

[Out] `int((a + b*tan(e + f*x))^2*(c + d*x)^3, x)`

3.45 $\int (c + dx)^2 (a + b \tan(e + fx))^2 dx$

Optimal result	288
Rubi [A] (verified)	289
Mathematica [B] (warning: unable to verify)	292
Maple [B] (verified)	294
Fricas [B] (verification not implemented)	294
Sympy [F]	295
Maxima [B] (verification not implemented)	295
Giac [F]	296
Mupad [F(-1)]	296

Optimal result

Integrand size = 20, antiderivative size = 229

$$\begin{aligned} \int (c + dx)^2 (a + b \tan(e + fx))^2 dx = & -\frac{ib^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} + \frac{2iab(c + dx)^3}{3d} \\ & - \frac{b^2(c + dx)^3}{3d} + \frac{2b^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\ & - \frac{2ab(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & - \frac{ib^2d^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & + \frac{2iabd(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{abd^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & + \frac{b^2(c + dx)^2 \tan(e + fx)}{f} \end{aligned}$$

```
[Out] -I*b^2*(d*x+c)^2/f+1/3*a^2*(d*x+c)^3/d+2/3*I*a*b*(d*x+c)^3/d-1/3*b^2*(d*x+c)^3/d+2*b^2*d*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^2-2*a*b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f-I*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3+2*I*a*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-a*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+b^2*(d*x+c)^2*tan(f*x+e)/f
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {3803, 3800, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32}

$$\begin{aligned} \int (c + dx)^2 (a + b \tan(e + fx))^2 dx = & \frac{a^2(c + dx)^3}{3d} + \frac{2iabd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{2ab(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{2iab(c + dx)^3}{3d} - \frac{abd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{f^3} \\ & + \frac{2b^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\ & + \frac{b^2(c + dx)^2 \tan(e + fx)}{f} - \frac{ib^2(c + dx)^2}{f} \\ & - \frac{b^2(c + dx)^3}{3d} - \frac{ib^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2 * (a + b*\operatorname{Tan}[e + f*x])^2, x]$

[Out] $\frac{((-I)*b^2*(c + d*x)^2)/f + (a^2*(c + d*x)^3)/(3*d) + (((2*I)/3)*a*b*(c + d*x)^3)/d - (b^2*(c + d*x)^3)/(3*d) + (2*b^2*d*(c + d*x)*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f^2 - (2*a*b*(c + d*x)^2*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f - (I*b^2*d^2*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^3 + ((2*I)*a*b*d*(c + d*x)*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - (a*b*d^2*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/f^3 + (b^2*(c + d*x)^2*\operatorname{Tan}[e + f*x])/f}{}$

Rule 32

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}, x \text{ Symbol}] \Rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /; \operatorname{FreeQ}[\{a, b, m\}, x] \&& \operatorname{NeQ}[m, -1]$

Rule 2221

$\operatorname{Int}[(((F_.)^{(g_.)}*((e_.) + (f_.)*(x_.)))^{(n_.)}*((c_.) + (d_.)*(x_.)^{(m_.)}))/((a_) + (b_.)*((F_.)^{(g_.)}*((e_.) + (f_.)*(x_.)))^{(n_.)}), x \text{ Symbol}] \Rightarrow \operatorname{Simp}[((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], x] - \operatorname{Dist}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^{(m - 1)}*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^{n/a})], x], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \operatorname{IGtQ}[m, 0]$

Rule 2317

$\operatorname{Int}[\operatorname{Log}[(a_) + (b_.)*((F_.)^{(e_.)}*((c_.) + (d_.)*(x_.)))^{(n_.)}], x \text{ Symbol}] \Rightarrow \operatorname{Dist}[1/(d*e*n*\operatorname{Log}[F]), \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[a + b*x]/x, x], x, (F^(e*(c + d*x)))]$

```
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_.) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.))*((f_.) + (g_.)
*(x_)^m_.], x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^
(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_.) + (d_.)*(x_))^m_*tan[(e_.) + (f_.)*(x_)], x_Symbol] :> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^m_*((b_.)*tan[(e_.) + (f_.)*(x_)])^n_, x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^m_*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^n_, x_Symb
ol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx)^2 + 2ab(c+dx)^2 \tan(e+fx) + b^2(c+dx)^2 \tan^2(e+fx)) \, dx \\
&= \frac{a^2(c+dx)^3}{3d} + (2ab) \int (c+dx)^2 \tan(e+fx) \, dx + b^2 \int (c+dx)^2 \tan^2(e+fx) \, dx \\
&= \frac{a^2(c+dx)^3}{3d} + \frac{2iab(c+dx)^3}{3d} + \frac{b^2(c+dx)^2 \tan(e+fx)}{f} \\
&\quad - (4iab) \int \frac{e^{2i(e+fx)}(c+dx)^2}{1+e^{2i(e+fx)}} \, dx - b^2 \int (c+dx)^2 \, dx \\
&\quad - \frac{(2b^2d) \int (c+dx) \tan(e+fx) \, dx}{f} \\
&= -\frac{ib^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} + \frac{2iab(c+dx)^3}{3d} - \frac{b^2(c+dx)^3}{3d} \\
&\quad - \frac{2ab(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \frac{b^2(c+dx)^2 \tan(e+fx)}{f} \\
&\quad + \frac{(4abd) \int (c+dx) \log(1+e^{2i(e+fx)}) \, dx}{f} + \frac{(4ib^2d) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} \, dx}{f} \\
&= -\frac{ib^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} + \frac{2iab(c+dx)^3}{3d} - \frac{b^2(c+dx)^3}{3d} \\
&\quad + \frac{2b^2d(c+dx) \log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{2iabd(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{b^2(c+dx)^2 \tan(e+fx)}{f} \\
&\quad - \frac{(2iabd^2) \int \text{PolyLog}(2, -e^{2i(e+fx)}) \, dx}{f^2} - \frac{(2b^2d^2) \int \log(1+e^{2i(e+fx)}) \, dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{ib^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} + \frac{2iab(c+dx)^3}{3d} - \frac{b^2(c+dx)^3}{3d} \\
&\quad + \frac{2b^2d(c+dx)\log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^2\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{2iabd(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{b^2(c+dx)^2\tan(e+fx)}{f} \\
&\quad - \frac{(abd^2)\text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{f^3} \\
&\quad + \frac{(ib^2d^2)\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{f^3} \\
&= -\frac{ib^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} + \frac{2iab(c+dx)^3}{3d} - \frac{b^2(c+dx)^3}{3d} \\
&\quad + \frac{2b^2d(c+dx)\log(1+e^{2i(e+fx)})}{f^2} - \frac{2ab(c+dx)^2\log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{ib^2d^2\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{2iabd(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{abd^2\text{PolyLog}(3, -e^{2i(e+fx)})}{f^3} + \frac{b^2(c+dx)^2\tan(e+fx)}{f}
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 649 vs. $2(229) = 458$.

Time = 6.83 (sec) , antiderivative size = 649, normalized size of antiderivative = 2.83

$$\begin{aligned}
 & \int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \\
 & - \frac{iabd^2 e^{-ie} (2f^2 x^2 (2fx - 3i(1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6(1 + e^{2ie}) fx \text{PolyLog}(2, -e^{-2i(e+fx)}) - 3i)}{6f^3} \\
 & + \frac{1}{3} x (3c^2 + 3cdx + d^2 x^2) \sec(e) (a^2 \cos(e) - b^2 \cos(e) + 2ab \sin(e)) \\
 & + \frac{2b^2 cd \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f^2 (\cos^2(e) + \sin^2(e))} \\
 & - \frac{2abc^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + fx \sin(e))}{f (\cos^2(e) + \sin^2(e))} \\
 & + \frac{b^2 d^2 \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e)))}))}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{2abcd \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (ifx(-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2ifx}) - 2(fx - \arctan(\cot(e))) \log(1 - e^{2i(fx - \arctan(\cot(e)))}))}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)} \\
 & + \frac{\sec(e) \sec(e + fx) (b^2 c^2 \sin(fx) + 2b^2 cdx \sin(fx) + b^2 d^2 x^2 \sin(fx))}{f}
 \end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^2, x]

[Out]
$$\begin{aligned}
 & ((-1/6*I)*a*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))])*Sec[e])/(E^(I*e)*f^3) + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Sec[e]*(a^2*Cos[e] - b^2*Cos[e] + 2*a*b*Sin[e]))/3 + (2*b^2*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (2*a*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^2*d^2*2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))])/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^3*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) - (2*a*b*c*d*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))])/Sqrt[1 + Cot[e]^2])*Sec[e])/(f^2*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) + (Sec[e]*Sec[e + f*x]*(b^2*c^2*Sin[f*x] + 2*b^2*c*d*x*Sin[f*x] + b^2*d^2*x^2*Sin[f*x]))/f
 \end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 574 vs. $2(209) = 418$.

Time = 1.39 (sec) , antiderivative size = 575, normalized size of antiderivative = 2.51

method	result
risch	$\frac{8ibdcaex}{f} + \frac{d^2a^2x^3}{3} + \frac{a^2c^3}{3d} - \frac{d^2b^2x^3}{3} - b^2c^2x - \frac{b^2c^3}{3d} + \frac{4b^2e d^2 \ln(e^{i(fx+e)})}{f^3} + da^2cx^2 + a^2c^2x - db^2cx^2 - \frac{2b^2c^3}{3d}$

```
[In] int((d*x+c)^2*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
[Out] 8*I/f*b*d*c*a*e*x+1/3*d^2*a^2*x^3+1/3/d*a^2*c^3-1/3*d^2*b^2*x^3-b^2*c^2*x-1
/3/d*b^2*c^3+4/f^3*b^2*c*d^2*ln(exp(I*(f*x+e)))-2/f*b*a*c^2*ln(exp(2*I*(f*x
+e))+1)+d*a^2*c*x^2+a^2*c^2*x-d*b^2*c*x^2-2*I*a*b*c^2*x-2/3*I/d*a*b*c^3+2*I
*b^2*(d^2*x^2+2*c*d*x+c^2)/f/(exp(2*I*(f*x+e))+1)+4/f*b*a*c^2*ln(exp(I*(f*x
+e)))+2/f^2*b^2*c*d*ln(exp(2*I*(f*x+e))+1)-4/f^2*b^2*c*d*ln(exp(I*(f*x+e)))
+2/f^2*b^2*d^2*ln(exp(2*I*(f*x+e))+1)*x-2*I/f*b^2*d^2*x^2-2*I/f^3*b^2*d^2*e
^2+2/3*I*d^2*a*b*x^3-a*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+2*I*d*a*b*c*x
^2-8/f^2*b*e*a*c*d*ln(exp(I*(f*x+e)))-4/f*b*d*c*a*ln(exp(2*I*(f*x+e))+1)*x-
4*I/f^2*b*a*d^2*x^2+2*I/f^2*b*a*d^2*polylog(2,-exp(2*I*(f*x+e)))*x+2*I/f^
2*b*d*c*a*polylog(2,-exp(2*I*(f*x+e)))+4*I/f^2*b*d*c*a*e^2-I*b^2*d^2*polylo
g(2,-exp(2*I*(f*x+e)))/f^3+4/f^3*b*e^2*a*d^2*ln(exp(I*(f*x+e)))-2/f*b*a*d^2
*ln(exp(2*I*(f*x+e))+1)*x^2-8/3*I/f^3*b*a*d^2*e^3-4*I/f^2*b^2*d^2*e*x
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 450 vs. $2(203) = 406$.

Time = 0.26 (sec) , antiderivative size = 450, normalized size of antiderivative = 1.97

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx \\ = 2(a^2 - b^2)d^2 f^3 x^3 + 6(a^2 - b^2)c d f^3 x^2 + 6(a^2 - b^2)c^2 f^3 x - 3abd^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3a$$

```
[In] integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="fricas")
[Out] 1/6*(2*(a^2 - b^2)*d^2*f^3*x^3 + 6*(a^2 - b^2)*c*d*f^3*x^2 + 6*(a^2 - b^2)*
c^2*f^3*x - 3*a*b*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(t
an(f*x + e)^2 + 1)) - 3*a*b*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x +
e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(2*I*a*b*d^2*f*x + 2*I*a*b*c*d*f - I*b^2*
d^2)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 3*(-2*I*a*b*d
^2*f*x - 2*I*a*b*c*d*f + I*b^2*d^2)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x
```

$$\begin{aligned}
& + e)^2 + 1) + 1) - 6*(a*b*d^2*f^2*x^2 + a*b*c^2*f^2 - b^2*c*d*f + (2*a*b*c*d*f^2 - b^2*d^2*f)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6 \\
& *(a*b*d^2*f^2*x^2 + a*b*c^2*f^2 - b^2*c*d*f + (2*a*b*c*d*f^2 - b^2*d^2*f)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^2*d^2*f^2*x^2 + 2*b^2*c*d*f^2*x + b^2*c^2*f^2)*tan(f*x + e))/f^3
\end{aligned}$$

Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx)^2 dx$$

```
[In] integrate((d*x+c)**2*(a+b*tan(f*x+e))**2,x)
[Out] Integral((a + b*tan(e + f*x))**2*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1263 vs. $2(203) = 406$.

Time = 0.91 (sec), antiderivative size = 1263, normalized size of antiderivative = 5.52

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="maxima")
[Out] 1/3*(3*(f*x + e)*a^2*c^2 + (f*x + e)^3*a^2*d^2/f^2 - 3*(f*x + e)^2*a^2*d^2*c^2*e/f^2 + 3*(f*x + e)*a^2*d^2*c^2*e^2/f^2 + 3*(f*x + e)^2*a^2*c*d/f - 6*(f*x + e)*a^2*c*d*e/f + 6*a*b*c^2*log(sec(f*x + e)) + 6*a*b*d^2*e^2*log(sec(f*x + e))/f^2 - 12*a*b*c*d*e*log(sec(f*x + e))/f + 3*((2*a*b + I*b^2)*(f*x + e)^3*d^2 + 6*b^2*d^2*e^2 - 12*b^2*c*d*e*f + 6*b^2*c^2*f^2 - 3*((2*a*b + I*b^2)*d^2*2*e - (2*a*b + I*b^2)*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*2*e^2 - 2*I*b^2*c*d*e*f + I*b^2*c^2*f^2)*(f*x + e) - 6*((f*x + e)^2*a*b*d^2 + b^2*d^2*2*e - b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*(f*x + e) + ((f*x + e)^2*a*b*d^2 + b^2*d^2*2*e - b^2*c*d*f - (2*a*b*d^2*e - 2*a*b*c*d*f + b^2*d^2)*(f*x + e))*cos(2*f*x + 2*e) - (-I*(f*x + e)^2*a*b*d^2 - I*b^2*d^2*2*e + I*b^2*c*d*f + (2*I*a*b*d^2*e - 2*I*a*b*c*d*f + I*b^2*d^2)*(f*x + e))*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2)*(f*x + e)^3*d^2 - 3*(2*b^2*d^2 + (2*a*b + I*b^2)*d^2*2*e - (2*a*b + I*b^2)*c*d*f)*(f*x + e)^2 + 3*(I*b^2*d^2*2*e^2 + I*b^2*c^2*f^2 + 4*b^2*d^2*2*e + 2*(-I*b^2*c*d*e - 2*b^2*c*d*f)*(f*x + e))*cos(2*f*x + 2*e) + 3*(2*(f*x + e)*a*b*d^2 - 2*a*b*d^2*2*e + 2*a*b*c*d*f - b^2*d^2 + (2*(f*x + e)*a*b*d^2 - 2*a*b*d^2*2*e + 2*a*b*c*d*f - b^2*d^2)*cos(2*f*x + 2*e) + (2*I*(f*x + e)*a*b*d^2 - 2*I*a*b*d^2*c^2*e))
```

$$\begin{aligned}
& + 2*I*a*b*c*d*f - I*b^2*d^2)*sin(2*f*x + 2*e))*dilog(-e^(2*I*f*x + 2*I*e)) \\
& + 3*(I*(f*x + e)^2*a*b*d^2 + I*b^2*d^2*e - I*b^2*c*d*f + (-2*I*a*b*d^2*e + \\
& 2*I*a*b*c*d*f - I*b^2*d^2)*(f*x + e) + (I*(f*x + e)^2*a*b*d^2 + I*b^2*d^2*e - \\
& I*b^2*c*d*f + (-2*I*a*b*d^2*e + 2*I*a*b*c*d*f - I*b^2*d^2)*(f*x + e))*c \\
& os(2*f*x + 2*e) - ((f*x + e)^2*a*b*d^2 + b^2*d^2*e - b^2*c*d*f - (2*a*b*d^2 \\
& *e - 2*a*b*c*d*f + b^2*d^2)*(f*x + e))*sin(2*f*x + 2*e))*log(cos(2*f*x + 2*e)^2 + \\
& sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) + 3*(I*a*b*d^2*cos(2*f*x + 2*e) - \\
& a*b*d^2*sin(2*f*x + 2*e) + I*a*b*d^2)*polylog(3, -e^(2*I*f*x + 2*I*e)) - \\
& ((-2*I*a*b + b^2)*(f*x + e)^3*d^2 - 3*(-2*I*b^2*d^2 + (-2*I*a*b + \\
& b^2)*d^2*e + (2*I*a*b - b^2)*c*d*f)*(f*x + e)^2 + 3*(b^2*d^2*e^2 + b^2*c^2*f^2 - \\
& 4*I*b^2*d^2*e - 2*(b^2*c*d*e - 2*I*b^2*c*d)*f)*(f*x + e))*sin(2*f*x + 2*e)) / \\
& (-3*I*f^2*cos(2*f*x + 2*e) + 3*f^2*sin(2*f*x + 2*e) - 3*I*f^2)/f
\end{aligned}$$

Giac [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (dx + c)^2 (b \tan(fx + e) + a)^2 dx$$

[In] integrate((d*x+c)^2*(a+b*tan(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tan(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + f x))^2 (c + d x)^2 dx$$

[In] int((a + b*tan(e + f*x))^2*(c + d*x)^2,x)

[Out] int((a + b*tan(e + f*x))^2*(c + d*x)^2, x)

3.46 $\int (c + dx)(a + b \tan(e + fx))^2 dx$

Optimal result	297
Rubi [A] (verified)	297
Mathematica [A] (verified)	299
Maple [A] (verified)	300
Fricas [A] (verification not implemented)	300
Sympy [F]	301
Maxima [B] (verification not implemented)	301
Giac [F]	302
Mupad [F(-1)]	302

Optimal result

Integrand size = 18, antiderivative size = 136

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^2 dx = & -b^2 cx - \frac{1}{2} b^2 dx^2 + \frac{a^2(c + dx)^2}{2d} + \frac{iab(c + dx)^2}{d} \\ & - \frac{2ab(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^2 d \log(\cos(e + fx))}{f^2} + \frac{iabd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & + \frac{b^2(c + dx) \tan(e + fx)}{f} \end{aligned}$$

[Out] $-b^2 c x - 1/2 b^2 d x^2 + 1/2 a^2 (d x + c)^2 d + I a b (d x + c)^2 / d - 2 a b (d x + c) \ln(1 + \exp(2 I (f x + e))) / f + b^2 d \ln(\cos(f x + e)) / f^2 + I a b d \operatorname{polylog}(2, -\exp(2 I (f x + e))) / f^2 + b^2 (d x + c) \tan(f x + e) / f$

Rubi [A] (verified)

Time = 0.20 (sec), antiderivative size = 136, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3803, 3800, 2221, 2317, 2438, 3801, 3556}

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^2 dx = & \frac{a^2(c + dx)^2}{2d} - \frac{2ab(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{iab(c + dx)^2}{d} + \frac{iabd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & + \frac{b^2(c + dx) \tan(e + fx)}{f} - b^2 cx \\ & + \frac{b^2 d \log(\cos(e + fx))}{f^2} - \frac{1}{2} b^2 dx^2 \end{aligned}$$

[In] $\text{Int}[(c + d*x)*(a + b*\text{Tan}[e + f*x])^2, x]$

[Out] $-\frac{(b^2*c*x)}{2} - \frac{(b^2*d*x^2)}{2} + \frac{(a^2*(c + d*x)^2)}{(2*d)} + \frac{(I*a*b*(c + d*x)^2)}{d} - \frac{(2*a*b*(c + d*x)*\text{Log}[1 + E^{(2*I)*(e + f*x)}])}{f} + \frac{(b^2*d*\text{Log}[\text{Cos}[e + f*x]])}{f^2} + \frac{(I*a*b*d*\text{PolyLog}[2, -E^{(2*I)*(e + f*x)}])}{f^2} + \frac{(b^2*(c + d*x)*\text{Tan}[e + f*x])}{f}$

Rule 2221

$\text{Int}[((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_*)})/((a_*) + (b_*)*((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[(c + d*x)^m/(b*f*g*n*\text{Log}[F])*(\text{Log}[1 + b*(F^((g*(e + f*x)))^n/a)]), x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{m-1}*\text{Log}[1 + b*(F^((g*(e + f*x)))^n/a)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

Rule 2317

$\text{Int}[\text{Log}[(a_*) + (b_*)*((F_*)^((e_*)*(c_*) + (d_*)*(x_))))^{(n_*)}], x_{\text{Symbol}}] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^((e*(c + d*x)))^n)], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$

Rule 2438

$\text{Int}[\text{Log}[(c_*)*((d_*) + (e_*)*(x_))^{(n_*)})/(x_*)], x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \&& \text{EqQ}[c*d, 1]$

Rule 3556

$\text{Int}[\tan[(c_*) + (d_*)*(x_*)], x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x]], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3800

$\text{Int}[((c_*) + (d_*)*(x_))^{(m_*)}*\tan[(e_*) + (f_*)*(x_*)], x_{\text{Symbol}}] \rightarrow \text{Simp}[I*((c + d*x)^(m+1)/(d*(m+1))), x] - \text{Dist}[2*I, \text{Int}[(c + d*x)^m*(E^{(2*I)*(e + f*x)})/(1 + E^{(2*I)*(e + f*x)})), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{IGtQ}[m, 0]$

Rule 3801

$\text{Int}[((c_*) + (d_*)*(x_))^{(m_*)}*((b_*)*\tan[(e_*) + (f_*)*(x_)])^{(n_*)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[b*(c + d*x)^m*((b*\text{Tan}[e + f*x])^{(n-1)}/(f*(n-1))), x] + (-\text{Dist}[b*d*(m/(f*(n-1))), \text{Int}[(c + d*x)^{m-1}*(b*\text{Tan}[e + f*x])^{(n-1)}, x], x] - \text{Dist}[b^2, \text{Int}[(c + d*x)^m*(b*\text{Tan}[e + f*x])^{(n-2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

Rule 3803

```
Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx) + 2ab(c+dx)\tan(e+fx) + b^2(c+dx)\tan^2(e+fx)) \, dx \\
&= \frac{a^2(c+dx)^2}{2d} + (2ab) \int (c+dx)\tan(e+fx) \, dx + b^2 \int (c+dx)\tan^2(e+fx) \, dx \\
&= \frac{a^2(c+dx)^2}{2d} + \frac{iab(c+dx)^2}{d} + \frac{b^2(c+dx)\tan(e+fx)}{f} \\
&\quad - (4iab) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} \, dx - b^2 \int (c+dx) \, dx - \frac{(b^2d) \int \tan(e+fx) \, dx}{f} \\
&= -b^2cx - \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} + \frac{iab(c+dx)^2}{d} - \frac{2ab(c+dx)\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\cos(e+fx))}{f^2} + \frac{b^2(c+dx)\tan(e+fx)}{f} + \frac{(2abd) \int \log(1+e^{2i(e+fx)}) \, dx}{f} \\
&= -b^2cx - \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} + \frac{iab(c+dx)^2}{d} - \frac{2ab(c+dx)\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\cos(e+fx))}{f^2} + \frac{b^2(c+dx)\tan(e+fx)}{f} - \frac{(iabd)\text{Subst}\left(\int \frac{\log(1+x)}{x} \, dx, x, e^{2i(e+fx)}\right)}{f^2} \\
&= -b^2cx - \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} + \frac{iab(c+dx)^2}{d} - \frac{2ab(c+dx)\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\cos(e+fx))}{f^2} + \frac{iabd \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} + \frac{b^2(c+dx)\tan(e+fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.41 (sec), antiderivative size = 200, normalized size of antiderivative = 1.47

$$\begin{aligned}
&\int (c+dx)(a+b\tan(e+fx))^2 \, dx \\
&= \frac{\cos(e+fx) (\cos(e+fx) ((-((e+fx)(-2iabd(e+fx)+a^2(de-2cf-dfx))+b^2(-de+2cf+dfx))) -
\end{aligned}$$

[In] `Integrate[(c + d*x)*(a + b*Tan[e + f*x])^2, x]`

[Out] `(Cos[e + f*x]*(Cos[e + f*x]*(-(e + f*x)*((-2*I)*a*b*d*(e + f*x) + a^2*(d*e - 2*c*f - d*f*x) + b^2*(-(d*e) + 2*c*f + d*f*x))) - 4*a*b*d*(e + f*x)*Log[`

$$1 + E^{\left((2*I)*(e + f*x)\right)} + 2*b*(b*d + 2*a*d*e - 2*a*c*f)*\text{Log}[\text{Cos}[e + f*x]] \\ + (2*I)*a*b*d*\text{Cos}[e + f*x]*\text{PolyLog}[2, -E^{\left((2*I)*(e + f*x)\right)}] + 2*b^2*f*(c + d*x)*\text{Sin}[e + f*x]*(a + b*\text{Tan}[e + f*x])^2)/(2*f^2*(a*\text{Cos}[e + f*x] + b*\text{Sin}[e + f*x])^2)$$

Maple [A] (verified)

Time = 1.31 (sec) , antiderivative size = 238, normalized size of antiderivative = 1.75

method	result
risch	$-\frac{b^2 dx^2}{2} + \frac{4ibadex}{f} + \frac{a^2 dx^2}{2} - b^2 cx - 2iabcx + a^2 cx + \frac{2ibade^2}{f^2} + \frac{b^2 d \ln(e^{2i(fx+e)}+1)}{f^2} - \frac{2b^2 d \ln(e^{i(fx+e)})}{f^2} - \frac{2ba^2 d \ln(e^{i(fx+e)})}{f^2}$

```
[In] int((d*x+c)*(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
[Out] -1/2*b^2*d*x^2+4*I/f*b*a*d*e*x+1/2*a^2*d*x^2-b^2*c*x-2*I*a*b*c*x+a^2*c*x+2*I/f^2*b*a*d*e^2+1/f^2*b^2*d*ln(exp(2*I*(f*x+e))+1)-2/f^2*b^2*d*ln(exp(I*(f*x+e)))-2/f*b*a*c*ln(exp(2*I*(f*x+e))+1)+4/f*b*a*c*ln(exp(I*(f*x+e)))-4/f^2*b*e*a*d*ln(exp(I*(f*x+e)))+2*I*b^2*(d*x+c)/f/(exp(2*I*(f*x+e))+1)+I*a*b*d*polylog(2,-exp(2*I*(f*x+e)))/f^2+I*a*b*d*x^2-2/f*b*a*d*ln(exp(2*I*(f*x+e))+1)*x
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 219, normalized size of antiderivative = 1.61

$$\int (c + dx)(a + b \tan(e + fx))^2 dx \\ = \frac{(a^2 - b^2)df^2x^2 + 2(a^2 - b^2)cf^2x - i abd\text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1}+1\right) + i abd\text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1}+1\right) - (2abd)}$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="fricas")
[Out] 1/2*((a^2 - b^2)*d*f^2*x^2 + 2*(a^2 - b^2)*c*f^2*x - I*a*b*d*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + I*a*b*d*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - (2*a*b*d*f*x + 2*a*b*c*f - b^2*d)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - (2*a*b*d*f*x + 2*a*b*c*f - b^2*d)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(b^2*d*f*x + b^2*c*f)*tan(f*x + e))/f^2
```

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx) dx$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))**2,x)
[Out] Integral((a + b*tan(e + f*x))**2*(c + d*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs. $2(123) = 246$.

Time = 0.41 (sec) , antiderivative size = 527, normalized size of antiderivative = 3.88

$$\begin{aligned} & \int (c + dx)(a + b \tan(e + fx))^2 dx \\ &= \frac{2(fx + e)a^2c + \frac{(fx+e)^2a^2d}{f} - \frac{2(fx+e)a^2de}{f} + 4abc \log(\sec(fx + e)) - \frac{4abde \log(\sec(fx+e))}{f} + \frac{2((2ab+ib^2)(fx+e)^2d-fc^2)}{f}}{f} \end{aligned}$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))**2,x, algorithm="maxima")
[Out] 1/2*(2*(f*x + e)*a^2*c + (f*x + e)^2*a^2*d/f - 2*(f*x + e)*a^2*d*e/f + 4*a*b*c*log(sec(f*x + e)) - 4*a*b*d*e*log(sec(f*x + e))/f + 2*((2*a*b + I*b^2)*(f*x + e)^2*d - 4*b^2*d*e + 4*b^2*c*f + 2*(-I*b^2*d*e + I*b^2*c*f)*(f*x + e)) - 2*(2*(f*x + e)*a*b*d - b^2*d + (2*(f*x + e)*a*b*d - b^2*d)*cos(2*f*x + 2*e) - (-2*I*(f*x + e)*a*b*d + I*b^2*d)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + ((2*a*b + I*b^2)*(f*x + e)^2*d + 2*(-I*b^2*d*e + I*b^2*c*f - 2*b^2*d)*(f*x + e))*cos(2*f*x + 2*e) + 2*(a*b*d*cos(2*f*x + 2*e) + I*a*b*d*sin(2*f*x + 2*e) + a*b*d)*dilog(-e^(2*I*f*x + 2*I*e)) - (-2*I*(f*x + e)*a*b*d + I*b^2*d + (-2*I*(f*x + e)*a*b*d + I*b^2*d)*cos(2*f*x + 2*e) + (2*(f*x + e)*a*b*d - b^2*d)*sin(2*f*x + 2*e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) - ((-2*I*a*b + b^2)*(f*x + e)^2*d - 2*(b^2*d*e - b^2*c*f - 2*I*b^2*d)*(f*x + e))*sin(2*f*x + 2*e)) / (-2*I*f*cos(2*f*x + 2*e) + 2*f*sin(2*f*x + 2*e) - 2*I*f)/f
```

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (dx + c)(b \tan(fx + e) + a)^2 dx$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((d*x + c)*(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^2 dx = \int (a + b \tan(e + fx))^2 (c + dx) dx$$

[In] `int((a + b*tan(e + f*x))^2*(c + d*x),x)`

[Out] `int((a + b*tan(e + f*x))^2*(c + d*x), x)`

3.47 $\int \frac{(a+b\tan(e+fx))^2}{c+dx} dx$

Optimal result	303
Rubi [N/A]	303
Mathematica [N/A]	304
Maple [N/A] (verified)	304
Fricas [N/A]	304
Sympy [N/A]	304
Maxima [N/A]	305
Giac [N/A]	305
Mupad [N/A]	305

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))^2/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

[In] Int[(a + b*Tan[e + f*x])^2/(c + d*x),x]

[Out] Defer[Int][(a + b*Tan[e + f*x])^2/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 19.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

[In] `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

[Out] `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.37 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{dx + c} dx$$

[In] `int((a+b*tan(f*x+e))^2/(d*x+c), x)`

[Out] `int((a+b*tan(f*x+e))^2/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))^2/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b^2*tan(f*x + e))^2 + 2*a*b*tan(f*x + e) + a^2)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 1.09 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^2}{c + dx} dx$$

[In] `integrate((a+b*tan(f*x+e))**2/(d*x+c), x)`

[Out] `Integral((a + b*tan(e + f*x))**2/(c + d*x), x)`

Maxima [N/A]

Not integrable

Time = 0.83 (sec) , antiderivative size = 491, normalized size of antiderivative = 24.55

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="maxima")`

[Out] $((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)^2*log(d*x + c) + 2*b^2*d*sin(2*f*x + 2*e) + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*log(d*x + c)*sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)*log(d*x + c) + (d^2*f*x + c*d*f + (d^2*f*x + c*d*f)*cos(2*f*x + 2*e)^2 + (d^2*f*x + c*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*cos(2*f*x + 2*e))*integrate(2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)*sin(2*f*x + 2*e)/(d^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2 + 2*c*d*f*x + c^2*f)*cos(2*f*x + 2*e)^2 + (d^2*f*x^2 + 2*c*d*f*x + c^2*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x^2 + 2*c*d*f*x + c^2*f)*cos(2*f*x + 2*e)), x) + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*log(d*x + c))/(d^2*f*x + c*d*f + (d^2*f*x + c*d*f)*cos(2*f*x + 2*e)^2 + (d^2*f*x + c*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^2*f*x + c*d*f)*cos(2*f*x + 2*e))$

Giac [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))^2/(d*x+c),x, algorithm="giac")`

[Out] `integrate((b*tan(f*x + e) + a)^2/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 4.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tan(e + f x))^2}{c + d x} dx$$

[In] `int((a + b*tan(e + f*x))^2/(c + d*x),x)`

[Out] `int((a + b*tan(e + f*x))^2/(c + d*x), x)`

3.48 $\int \frac{(a+b \tan(e+fx))^2}{(c+dx)^2} dx$

Optimal result	306
Rubi [N/A]	306
Mathematica [N/A]	307
Maple [N/A] (verified)	307
Fricas [N/A]	307
Sympy [N/A]	308
Maxima [N/A]	308
Giac [N/A]	309
Mupad [N/A]	309

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^2}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))^2/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.05 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

[In] Int[(a + b*Tan[e + f*x])^2/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 14.03 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

[In] `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]`

[Out] `Integrate[(a + b*Tan[e + f*x])^2/(c + d*x)^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^2}{(dx + c)^2} dx$$

[In] `int((a+b*tan(f*x+e))^2/(d*x+c)^2, x)`

[Out] `int((a+b*tan(f*x+e))^2/(d*x+c)^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2, x, algorithm="fricas")`

[Out] `integral((b^2*tan(f*x + e)^2 + 2*a*b*tan(f*x + e) + a^2)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.11 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))**2/(d*x+c)**2,x)
[Out] Integral((a + b*tan(e + f*x))**2/(c + d*x)**2, x)
```

Maxima [N/A]

Not integrable

Time = 1.68 (sec) , antiderivative size = 609, normalized size of antiderivative = 30.45

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="maxima")
[Out] -((a^2 - b^2)*d*f*x - 2*b^2*d*sin(2*f*x + 2*e) + (a^2 - b^2)*c*f + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e)^2 + ((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*sin(2*f*x + 2*e)^2 + 2*((a^2 - b^2)*d*f*x + (a^2 - b^2)*c*f)*cos(2*f*x + 2*e) - (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)) *integrate(4*(a*b*d*f*x + a*b*c*f + b^2*d)*sin(2*f*x + 2*e)/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f)*cos(2*f*x + 2*e)), x))/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e)^2 + (d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f)*cos(2*f*x + 2*e))
```

Giac [N/A]

Not integrable

Time = 15.43 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] `integrate((a+b*tan(f*x+e))^2/(d*x+c)^2,x, algorithm="giac")`

[Out] `integrate((b*tan(f*x + e) + a)^2/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 3.64 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + f x))^2}{(c + d x)^2} dx$$

[In] `int((a + b*tan(e + f*x))^2/(c + d*x)^2,x)`

[Out] `int((a + b*tan(e + f*x))^2/(c + d*x)^2, x)`

$$\mathbf{3.49} \quad \int (c + dx)^3 (a + b \tan(e + fx))^3 \, dx$$

Optimal result	311
Rubi [A] (verified)	312
Mathematica [B] (warning: unable to verify)	319
Maple [B] (verified)	320
Fricas [B] (verification not implemented)	322
Sympy [F]	323
Maxima [B] (verification not implemented)	323
Giac [F]	326
Mupad [F(-1)]	327

Optimal result

Integrand size = 20, antiderivative size = 612

$$\begin{aligned}
\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = & \frac{3ib^3d(c + dx)^2}{2f^2} - \frac{3iab^2(c + dx)^3}{f} + \frac{b^3(c + dx)^3}{2f} \\
& + \frac{a^3(c + dx)^4}{4d} + \frac{3ia^2b(c + dx)^4}{4d} - \frac{3ab^2(c + dx)^4}{4d} \\
& - \frac{ib^3(c + dx)^4}{4d} - \frac{3b^3d^2(c + dx) \log(1 + e^{2i(e+fx)})}{f^3} \\
& + \frac{9ab^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\
& - \frac{3a^2b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
& + \frac{b^3(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
& + \frac{3ib^3d^3 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} \\
& - \frac{9iab^2d^2(c + dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
& + \frac{9ia^2bd(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
& - \frac{3ib^3d(c + dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
& + \frac{9ab^2d^3 \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\
& - \frac{9a^2bd^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
& + \frac{3b^3d^2(c + dx) \text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
& - \frac{9ia^2bd^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
& + \frac{3ib^3d^3 \text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
& - \frac{3b^3d(c + dx)^2 \tan(e + fx)}{2f^2} \\
& + \frac{3ab^2(c + dx)^3 \tan(e + fx)}{f} \\
& + \frac{b^3(c + dx)^3 \tan^2(e + fx)}{2f}
\end{aligned}$$

```
[Out] -9*I*a*b^2*d^2*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^3-1/4*I*b^3*(d*x+c)^4
/d+1/2*b^3*(d*x+c)^3/f+1/4*a^3*(d*x+c)^4/d+9/2*I*a^2*b*d*(d*x+c)^2*polylog(
2,-exp(2*I*(f*x+e)))/f^2-3/4*a*b^2*(d*x+c)^4/d+3/2*I*b^3*d^3*polylog(2,-exp
(2*I*(f*x+e)))/f^4-3*b^3*d^2*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^3+9*a*b^2*d*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f^2-3*a^2*b*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))
/f+b^3*(d*x+c)^3*ln(1+exp(2*I*(f*x+e)))/f-3*I*a*b^2*(d*x+c)^3/f-9/4*I*a^2*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+3/2*I*b^3*d*(d*x+c)^2/f^2+3/4*I*a^2*b*(d*x+c)^4/d+9/2*a*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^4-9/2*a^2*b*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3+3/2*b^3*d^2*(d*x+c)*polylog(3,-exp(2*I*(f*x+e)))/f^3-3/2*I*b^3*d*(d*x+c)^2*polylog(2,-exp(2*I*(f*x+e)))/f^2+3/4*I*b^3*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-3/2*b^3*d*(d*x+c)^2*tan(f*x+e)
/f^2+3*a*b^2*(d*x+c)^3*tan(f*x+e)/f+1/2*b^3*(d*x+c)^3*tan(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 1.12 (sec) , antiderivative size = 612, normalized size of antiderivative = 1.00, number of steps used = 28, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules

used = {3803, 3800, 2221, 2611, 6744, 2320, 6724, 3801, 32, 2317, 2438}

$$\begin{aligned}
\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = & \frac{a^3(c + dx)^4}{4d} - \frac{9a^2bd^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
& + \frac{9ia^2bd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
& - \frac{3a^2b(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
& + \frac{3ia^2b(c + dx)^4}{4d} - \frac{9ia^2bd^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
& - \frac{9iab^2d^2(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
& + \frac{9ab^2d(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f^2} \\
& + \frac{3ab^2(c + dx)^3 \tan(e + fx)}{f} - \frac{3iab^2(c + dx)^3}{f} \\
& - \frac{3ab^2(c + dx)^4}{4d} + \frac{9ab^2d^3 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} \\
& + \frac{3b^3d^2(c + dx) \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
& - \frac{3b^3d^2(c + dx) \log(1 + e^{2i(e+fx)})}{f^3} \\
& - \frac{3ib^3d(c + dx)^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
& - \frac{3b^3d(c + dx)^2 \tan(e + fx)}{2f^2} \\
& + \frac{b^3(c + dx)^3 \log(1 + e^{2i(e+fx)})}{f} \\
& + \frac{b^3(c + dx)^3 \tan^2(e + fx)}{2f} + \frac{3ib^3d(c + dx)^2}{2f^2} \\
& + \frac{b^3(c + dx)^3}{2f} - \frac{ib^3(c + dx)^4}{4d} \\
& + \frac{3ib^3d^3 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} \\
& + \frac{3ib^3d^3 \operatorname{PolyLog}(4, -e^{2i(e+fx)})}{4f^4}
\end{aligned}$$

[In] Int[(c + d*x)^3*(a + b*Tan[e + f*x])^3, x]

[Out] (((3*I)/2)*b^3*d*(c + d*x)^2)/f^2 - ((3*I)*a*b^2*(c + d*x)^3)/f + (b^3*(c + d*x)^3)/(2*f) + (a^3*(c + d*x)^4)/(4*d) + (((3*I)/4)*a^2*b*(c + d*x)^4)/d

$$\begin{aligned}
& - \frac{(3*a*b^2*(c + d*x)^4)/(4*d) - ((I/4)*b^3*(c + d*x)^4)/d - (3*b^3*d^2*(c + d*x)*Log[1 + E^((2*I)*(e + f*x))])/f^3 + (9*a*b^2*d*(c + d*x)^2*Log[1 + E^((2*I)*(e + f*x))])/f^2 - (3*a^2*b*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f + (b^3*(c + d*x)^3*Log[1 + E^((2*I)*(e + f*x))])/f^4 - ((9*I)*a*b^2*d^2*(c + d*x)*PolyLog[2, -E^((2*I)*(e + f*x))])/f^3 + (((9*I)/2)*a^2*b*d*(c + d*x)^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 - (((3*I)/2)*b^3*d*(c + d*x)^2*PolyLog[2, -E^((2*I)*(e + f*x))])/f^2 + (9*a*b^2*d^3*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^4) - (9*a^2*b*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) + (3*b^3*d^2*(c + d*x)*PolyLog[3, -E^((2*I)*(e + f*x))])/(2*f^3) - (((9*I)/4)*a^2*b*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4 + (((3*I)/4)*b^3*d^3*PolyLog[4, -E^((2*I)*(e + f*x))])/f^4 - (3*b^3*d*(c + d*x)^2*Tan[e + f*x])/(2*f^2) + (3*a*b^2*(c + d*x)^3*Tan[e + f*x])/f + (b^3*(c + d*x)^3*Tan[e + f*x]^2)/(2*f)
\end{aligned}$$
Rule 32

```
Int[((a_) + (b_)*(x_))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((((F_)^((g_)*(e_) + (f_)*(x_))))^(n_)*(c_)*(d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*(d_ + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_) + (b_)*(x_)))^(n_.)]*((f_) + (g_)*
(x_))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m -
1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] :> Simp[I*
((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e +
f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 3801

```
Int[((c_) + (d_)*(x_))^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_.),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_) + (b_)*(x_))^p]/((d_*) + (e_)*(x_)), x_Sy
mbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*(F_)^((c_)*(a_) + (b_)*
(x_)))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a +
b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m -
1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = \int (a^3(c+dx)^3 + 3a^2b(c+dx)^3 \tan(e+fx) + 3ab^2(c+dx)^3 \tan^2(e+fx) + b^3(c+dx)^3 \tan^3(e+fx)) dx$$

$$\begin{aligned}
&= \frac{a^3(c+dx)^4}{4d} + (3a^2b) \int (c+dx)^3 \tan(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx)^3 \tan^2(e+fx) dx + b^3 \int (c+dx)^3 \tan^3(e+fx) dx \\
&= \frac{a^3(c+dx)^4}{4d} + \frac{3ia^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^3 \tan^2(e+fx)}{2f} - (6ia^2b) \int \frac{e^{2i(e+fx)}(c+dx)^3}{1+e^{2i(e+fx)}} dx \\
&\quad - (3ab^2) \int (c+dx)^3 dx - b^3 \int (c+dx)^3 \tan(e+fx) dx \\
&\quad - \frac{(9ab^2d) \int (c+dx)^2 \tan(e+fx) dx}{f} - \frac{(3b^3d) \int (c+dx)^2 \tan^2(e+fx) dx}{2f} \\
&= -\frac{3iab^2(c+dx)^3}{f} + \frac{a^3(c+dx)^4}{4d} + \frac{3ia^2b(c+dx)^4}{4d} - \frac{3ab^2(c+dx)^4}{4d} \\
&\quad - \frac{ib^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{3b^3d(c+dx)^2 \tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx)^3 \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^3 \tan^2(e+fx)}{2f} + (2ib^3) \int \frac{e^{2i(e+fx)}(c+dx)^3}{1+e^{2i(e+fx)}} dx \\
&\quad + \frac{(3b^3d^2) \int (c+dx) \tan(e+fx) dx}{f^2} + \frac{(9a^2bd) \int (c+dx)^2 \log(1+e^{2i(e+fx)}) dx}{f} \\
&\quad + \frac{(18iab^2d) \int \frac{e^{2i(e+fx)}(c+dx)^2}{1+e^{2i(e+fx)}} dx}{f} + \frac{(3b^3d) \int (c+dx)^2 dx}{2f} \\
&= \frac{3ib^3d(c+dx)^2}{2f^2} - \frac{3iab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} + \frac{3ia^2b(c+dx)^4}{4d} \\
&\quad - \frac{3ab^2(c+dx)^4}{4d} - \frac{ib^3(c+dx)^4}{4d} + \frac{9ab^2d(c+dx)^2 \log(1+e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2b(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^3 \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{9ia^2bd(c+dx)^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3b^3d(c+dx)^2 \tan(e+fx)}{2f^2} \\
&\quad + \frac{3ab^2(c+dx)^3 \tan(e+fx)}{f} + \frac{b^3(c+dx)^3 \tan^2(e+fx)}{2f} \\
&\quad - \frac{(9ia^2bd^2) \int (c+dx) \text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^2} \\
&\quad - \frac{(18ab^2d^2) \int (c+dx) \log(1+e^{2i(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6ib^3d^2) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} dx}{f^2} - \frac{(3b^3d) \int (c+dx)^2 \log(1+e^{2i(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3ib^3d(c+dx)^2}{2f^2} - \frac{3iab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} + \frac{3ia^2b(c+dx)^4}{4d} \\
&\quad - \frac{3ab^2(c+dx)^4}{4d} - \frac{ib^3(c+dx)^4}{4d} - \frac{3b^3d^2(c+dx)\log(1+e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ab^2d(c+dx)^2\log(1+e^{2i(e+fx)})}{f^2} - \frac{3a^2b(c+dx)^3\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3\log(1+e^{2i(e+fx)})}{f} - \frac{9iab^2d^2(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ia^2bd(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3ib^3d(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{3b^3d(c+dx)^2\tan(e+fx)}{2f^2} \\
&\quad + \frac{3ab^2(c+dx)^3\tan(e+fx)}{f} + \frac{b^3(c+dx)^3\tan^2(e+fx)}{2f} \\
&\quad + \frac{(9a^2bd^3)\int \text{PolyLog}(3, -e^{2i(e+fx)}) dx}{2f^3} + \frac{(9iab^2d^3)\int \text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^3} \\
&\quad + \frac{(3b^3d^3)\int \log(1+e^{2i(e+fx)}) dx}{f^3} + \frac{(3ib^3d^2)\int (c+dx)\text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^2} \\
&= \frac{3ib^3d(c+dx)^2}{2f^2} - \frac{3iab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} + \frac{3ia^2b(c+dx)^4}{4d} \\
&\quad - \frac{3ab^2(c+dx)^4}{4d} - \frac{ib^3(c+dx)^4}{4d} - \frac{3b^3d^2(c+dx)\log(1+e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ab^2d(c+dx)^2\log(1+e^{2i(e+fx)})}{f^2} - \frac{3a^2b(c+dx)^3\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3\log(1+e^{2i(e+fx)})}{f} - \frac{9iab^2d^2(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ia^2bd(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3ib^3d(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{3b^3d^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad - \frac{3b^3d(c+dx)^2\tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx)^3\tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^3\tan^2(e+fx)}{2f} - \frac{(9ia^2bd^3)\text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{4f^4} \\
&\quad + \frac{(9ab^2d^3)\text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^4} \\
&\quad - \frac{(3ib^3d^3)\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^4} - \frac{(3b^3d^3)\int \text{PolyLog}(3, -e^{2i(e+fx)}) dx}{2f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3ib^3d(c+dx)^2}{2f^2} - \frac{3iab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad + \frac{3ia^2b(c+dx)^4}{4d} - \frac{3ab^2(c+dx)^4}{4d} - \frac{ib^3(c+dx)^4}{4d} \\
&\quad - \frac{3b^3d^2(c+dx)\log(1+e^{2i(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2\log(1+e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2b(c+dx)^3\log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^3\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ib^3d^3\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} - \frac{9iab^2d^2(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ia^2bd(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3ib^3d(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad + \frac{9ab^2d^3\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad + \frac{3b^3d^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{9ia^2bd^3\text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
&\quad - \frac{3b^3d(c+dx)^2\tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx)^3\tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^3\tan^2(e+fx)}{2f} + \frac{(3ib^3d^3)\text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{4f^4} \\
&= \frac{3ib^3d(c+dx)^2}{2f^2} - \frac{3iab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad + \frac{3ia^2b(c+dx)^4}{4d} - \frac{3ab^2(c+dx)^4}{4d} - \frac{ib^3(c+dx)^4}{4d} \\
&\quad - \frac{3b^3d^2(c+dx)\log(1+e^{2i(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2\log(1+e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2b(c+dx)^3\log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^3\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{3ib^3d^3\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^4} - \frac{9iab^2d^2(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{9ia^2bd(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{3ib^3d(c+dx)^2\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad + \frac{9ab^2d^3\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^4} - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad + \frac{3b^3d^2(c+dx)\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} - \frac{9ia^2bd^3\text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} \\
&\quad + \frac{3ib^3d^3\text{PolyLog}(4, -e^{2i(e+fx)})}{4f^4} - \frac{3b^3d(c+dx)^2\tan(e+fx)}{2f^2} \\
&\quad + \frac{3ab^2(c+dx)^3\tan(e+fx)}{f} + \frac{b^3(c+dx)^3\tan^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 2594 vs. $2(612) = 1224$.

Time = 7.57 (sec) , antiderivative size = 2594, normalized size of antiderivative = 4.24

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Result too large to show}$$

```
[In] Integrate[(c + d*x)^3*(a + b*Tan[e + f*x])^3,x]
[Out] (((3*I)/4)*a*b^2*d^3*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/((E^(I*e)*f^4) - (((3*I)/4)*a^2*b*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) + ((I/4)*b^3*c*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/(E^(I*e)*f^3) - (((3*I)/8)*a^2*b*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((-2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e]/f^4 + ((I/8)*b^3*d^3*E^(I*e)*((2*f^4*x^4)/E^((2*I)*e) - (4*I)*(1 + E^((-2*I)*e))*f^3*x^3*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((-2*I)*e))*f^2*x^2*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (6*I)*(1 + E^((-2*I)*e))*f*x*PolyLog[3, -E^((-2*I)*(e + f*x))] - 3*(1 + E^((-2*I)*e))*PolyLog[4, -E^((-2*I)*(e + f*x))]*Sec[e]/f^4 + ((b^3*c^3 + 3*b^3*c^2*d*x + 3*b^3*c*d^2*x^2 + b^3*d^3*x^3)*Sec[e + f*x]^2)/(2*f) - (3*b^3*c*d^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^3*(Cos[e]^2 + Sin[e]^2)) + (9*a*b^2*c^2*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (3*a^2*b*c^3*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^3*c^3*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) - (3*b^3*d^3*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))]))/Sqrt[1 + Cot[e]^2]*Sec[e]/(2*f^4*Sqrt[Csc[e]^2*(Cos[e]^2 + Sin[e]^2)]) + (9*a*b^2*c*d^2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)] - 2*(f*x - ArcTan[Cot[e]])*Log[1 - E^((2*I)*(f*x - ArcTan[Cot[e]]))]) + Pi*Log[Cos[f*x]] - 2*ArcTan[Cot[e]]*Log[Sin[f*x - ArcTan[Cot[e]]]] + I*PolyLog[2, E^((2*I)*(f*x - ArcTan[Cot[e]]))])
```

$$\begin{aligned}
& *I*(f*x - \text{ArcTan}[\text{Cot}[e]])))/\text{Sqrt}[1 + \text{Cot}[e]^2])*\text{Sec}[e])/(f^3*\text{Sqrt}[\text{Csc}[e] \\
& ^{-2}*(\text{Cos}[e]^2 + \text{Sin}[e]^2]) - (9*a^2*b*c^2*d*\text{Csc}[e]*((f^2*x^2)/E^(\text{I}*\text{ArcTan}[\text{C} \\
& \text{ot}[e]])) - (\text{Cot}[e]*(\text{I}*\text{f}*\text{x}*(-\text{Pi} - 2*\text{ArcTan}[\text{Cot}[e]])) - \text{Pi}*\text{Log}[1 + E^((-2*\text{I})*f*x)] \\
& - 2*(f*x - \text{ArcTan}[\text{Cot}[e]])))*\text{Log}[1 - E^((2*\text{I})*(f*x - \text{ArcTan}[\text{Cot}[e]]))] + \\
& \text{Pi}*\text{Log}[\text{Cos}[f*x]] - 2*\text{ArcTan}[\text{Cot}[e]]*\text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + \text{I}*\text{Poly} \\
& \text{Log}[2, E^((2*\text{I})*(f*x - \text{ArcTan}[\text{Cot}[e]])))]))/\text{Sqrt}[1 + \text{Cot}[e]^2])*\text{Sec}[e])/(2*f \\
& ^{-2}*\text{Sqrt}[\text{Csc}[e]^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)]) + (3*b^3*c^2*d*\text{Csc}[e]*((f^2*x^2)/E \\
& ^(\text{I}*\text{ArcTan}[\text{Cot}[e]])) - (\text{Cot}[e]*(\text{I}*\text{f}*\text{x}*(-\text{Pi} - 2*\text{ArcTan}[\text{Cot}[e]])) - \text{Pi}*\text{Log}[1 + \\
& E^((-2*\text{I})*f*x)] - 2*(f*x - \text{ArcTan}[\text{Cot}[e]])))*\text{Log}[1 - E^((2*\text{I})*(f*x - \text{ArcTan}[\text{C} \\
& \text{ot}[e]]))] + \text{Pi}*\text{Log}[\text{Cos}[f*x]] - 2*\text{ArcTan}[\text{Cot}[e]]*\text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]] \\
&] + \text{I}*\text{PolyLog}[2, E^((2*\text{I})*(f*x - \text{ArcTan}[\text{Cot}[e]])))]))/\text{Sqrt}[1 + \text{Cot}[e]^2)* \\
& \text{Sec}[e])/(2*f^2*\text{Sqrt}[\text{Csc}[e]^2*(\text{Cos}[e]^2 + \text{Sin}[e]^2)]) + (3*x^2*(a^3*c^2*d + \\
& (3*I)*a^2*b*c^2*d - 3*a*b^2*c^2*d - I*b^3*c^2*d + a^3*c^2*d*\text{Cos}[2*e] - (3*I) \\
&)*a^2*b*c^2*d*\text{Cos}[2*e] - 3*a*b^2*c^2*d*\text{Cos}[2*e] + I*b^3*c^2*d*\text{Cos}[2*e] + I* \\
& a^3*c^2*d*\text{Sin}[2*e] + 3*a^2*b*c^2*d*\text{Sin}[2*e] - (3*I)*a*b^2*c^2*d*\text{Sin}[2*e] - \\
& b^3*c^2*d*\text{Sin}[2*e]))/(2*(1 + \text{Cos}[2*e] + \text{I}*\text{Sin}[2*e])) + (x^3*(a^3*c*d^2 + (3 \\
& *I)*a^2*b*c*d^2 - 3*a*b^2*c*d^2 - I*b^3*c*d^2 + a^3*c*d^2*\text{Cos}[2*e] - (3*I)* \\
& a^2*b*c*d^2*\text{Cos}[2*e] - 3*a*b^2*c*d^2*\text{Cos}[2*e] + I*b^3*c*d^2*\text{Cos}[2*e] + I*a^ \\
& 3*c*d^2*\text{Sin}[2*e] + 3*a^2*b*c*d^2*\text{Sin}[2*e] - (3*I)*a*b^2*c*d^2*\text{Sin}[2*e] - b^ \\
& 3*c*d^2*\text{Sin}[2*e]))/(1 + \text{Cos}[2*e] + \text{I}*\text{Sin}[2*e]) + (x^4*(a^3*d^3 + (3*I)*a^2* \\
& b*d^3 - 3*a*b^2*d^3 - I*b^3*d^3 + a^3*d^3*\text{Cos}[2*e] - (3*I)*a^2*b*d^3*\text{Cos}[2* \\
& e] - 3*a*b^2*d^3*\text{Cos}[2*e] + I*b^3*d^3*\text{Cos}[2*e] + I*a^3*d^3*\text{Sin}[2*e] + 3*a^2* \\
& *b*d^3*\text{Sin}[2*e] - (3*I)*a*b^2*d^3*\text{Sin}[2*e] - b^3*d^3*\text{Sin}[2*e]))/(4*(1 + \text{Cos} \\
& [2*e] + \text{I}*\text{Sin}[2*e])) + x*(a^3*c^3 - 3*a*b^2*c^3 + ((3*I)*a^2*b*c^3)/(1 + \text{Co} \\
& s[2*e] + \text{I}*\text{Sin}[2*e]) + ((-3*I)*a^2*b*c^3*\text{Cos}[2*e] + 3*a^2*b*c^3*\text{Sin}[2*e])/ \\
& ((1 + \text{Cos}[2*e] + \text{I}*\text{Sin}[2*e]) + ((2*I)*b^3*c^3*\text{Cos}[2*e] - 2*b^3*c^3*\text{Sin}[2*e])/ \\
& ((1 + \text{Cos}[2*e] + \text{I}*\text{Sin}[2*e])*((1 - \text{Cos}[2*e] + \text{Cos}[4*e] - \text{I}*\text{Sin}[2*e] + \text{I}*\text{Sin}[\\
& 4*e])) + ((-2*I)*b^3*c^3*\text{Cos}[4*e] + 2*b^3*c^3*\text{Sin}[4*e])/((1 + \text{Cos}[2*e] + \text{I}*\text{Sin}[2* \\
& e])*((1 - \text{Cos}[2*e] + \text{Cos}[4*e] - \text{I}*\text{Sin}[2*e] + \text{I}*\text{Sin}[4*e])) - (I*b^3*c^3 \\
&)/(1 + \text{Cos}[6*e] + \text{I}*\text{Sin}[6*e]) + (I*b^3*c^3*\text{Cos}[6*e] - b^3*c^3*\text{Sin}[6*e])/(1 \\
& + \text{Cos}[6*e] + \text{I}*\text{Sin}[6*e]) + (3*\text{Sec}[e]*\text{Sec}[e + f*x]*(-b^3*c^2*d*\text{Sin}[f*x]) + \\
& 2*a*b^2*c^3*f*\text{Sin}[f*x] - 2*b^3*c*d^2*x*\text{Sin}[f*x] + 6*a*b^2*c^2*d*f*x*\text{Sin}[f* \\
& x] - b^3*d^3*x^2*\text{Sin}[f*x] + 6*a*b^2*c*d^2*f*x^2*\text{Sin}[f*x] + 2*a*b^2*d^3*f*x^ \\
& 3*\text{Sin}[f*x]))/(2*f^2)
\end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1929 vs. 2(544) = 1088.

Time = 1.21 (sec), antiderivative size = 1930, normalized size of antiderivative = 3.15

method	result	size
risch	Expression too large to display	1930

[In] `int((d*x+c)^3*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned}
& -1/4*I*b^3*d^3*x^4+d^2*a^3*c*x^3+3/2*d*a^3*c^2*x^2+a^3*c^3*x-3/4*d^3*a*b^2*x^4-3*a*b^2*c^3*x-3/4/d*a*b^2*c^4+I*b^3*c^3*x+1/4*I/d*b^3*c^4+1/4*d^3*a^3*x^4+1/4/d*c^4*a^3-36*I/f^2*b^2*d^2*c*a*e*x+3/4*I*b^3*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4+1/f*b^3*c^3*ln(exp(2*I*(f*x+e))+1)-2/f*b^3*c^3*ln(exp(I*(f*x+e)))+b^2*(-6*I*c*d^2*x*b-6*I*b*c*d^2*x*exp(2*I*(f*x+e))+2*b*d^3*f*x^3*exp(2*I*(f*x+e))+18*I*a*c*d^2*f*x^2-3*I*c^2*d*b-3*I*d^3*x^2*b+6*b*c*d^2*f*x^2*exp(2*I*(f*x+e))-3*I*b*d^3*x^2*exp(2*I*(f*x+e))+18*I*a*c*d^2*f*x^2*exp(2*I*(f*x+e))+6*I*a*d^3*f*x^3*exp(2*I*(f*x+e))+6*b*c^2*d*f*x*exp(2*I*(f*x+e))-3*I*b*c^2*d*exp(2*I*(f*x+e))+6*I*a*c^3*f*exp(2*I*(f*x+e))+18*I*a*c^2*d*f*x+2*b*c^3*f*exp(2*I*(f*x+e))+18*I*a*c^2*d*f*x*exp(2*I*(f*x+e))+6*I*a*d^3*f*x^3+6*I*a*c^3*f)/f^2/(exp(2*I*(f*x+e))+1)^2+3/4*I*d^3*a^2*b*x^4-3/2*I*b^3*d*c^2*x^2-18*I/f^2*b*a^2*c*d^2*x+9*I/f^2*b*a^2*c*d^2*polylog(2,-exp(2*I*(f*x+e)))*x+18*I/f*b*d*c^2*a^2*e*x-6/f^4*b^3*e*d^3*ln(exp(I*(f*x+e)))+3*I/f^2*b^3*d^3*x^2+3*I/f^4*b^3*d^3*e^2-3/2*I/f^4*b^3*d^3*e^4-9/4*I*a^2*b*d^3*polylog(4,-exp(2*I*(f*x+e)))/f^4-I*d^2*b^3*c*x^3-3*d^2*a*b^2*c*x^3-9/2*d*a*b^2*c^2*x^2-3*I*a^2*b*c^3*x-3/4*I/d*a^2*b*c^4+3/2*I*b^3*d^3*polylog(2,-exp(2*I*(f*x+e)))/f^4-3/f^3*b^3*c*d^2*ln(exp(2*I*(f*x+e))+1)+6/f^3*b^3*c*d^2*ln(exp(I*(f*x+e)))+1/f*b^3*d^3*ln(exp(2*I*(f*x+e))+1)*x^3+3/2/f^3*b^3*c*d^2*polylog(3,-exp(2*I*(f*x+e)))-3/f*b*a^2*c^3*ln(exp(2*I*(f*x+e))+1)+6/f*b*a^2*c^3*ln(exp(I*(f*x+e)))-3/f^3*b^3*d^3*ln(exp(2*I*(f*x+e))+1)*x+3/2/f^3*b^3*d^3*polylog(3,-exp(2*I*(f*x+e)))*x+2/f^4*b^3*e^3*d^3*ln(exp(I*(f*x+e)))+9/2*a*b^2*d^3*polylog(3,-exp(2*I*(f*x+e)))/f^4-3*I/f^2*b^3*d*c^2*e^2-3/2*I/f^2*b^3*d*c^2*polylog(2,-exp(2*I*(f*x+e)))-2*I/f^3*b^3*d^3*e^3*x+6*I/f^3*b^3*d^3*e*x-6*I/f*b^3*d*c^2*e*x+9/2*I/f^2*b*d^3*a^2*polylog(2,-exp(2*I*(f*x+e)))*x^2+6*I/f^3*b*e^3*a^2*d^3*x-18*I/f*b^2*d^2*c*a*x^2+9*I/f^2*b*d*c^2*a^2*e^2+6*I/f^2*b^3*c*d^2*e^2*x-3*I/f^2*b^3*c*d^2*polylog(2,-exp(2*I*(f*x+e)))*x+18*I/f^3*b^2*a*d^3*e^2*x-9*I/f^3*b^2*a*d^3*polylog(2,-exp(2*I*(f*x+e)))*x+3*I*d^2*a^2*b*c*x^3+9/2*I*d*a^2*b*c^2*x^2+6/f^2*b^3*e*d*c^2*ln(exp(I*(f*x+e)))+9/f^2*b^2*a*c^2*d*ln(exp(2*I*(f*x+e))+1)-18/f^2*b^2*a*c^2*d*ln(exp(I*(f*x+e)))-18/f^4*b^2*e^2*a*d^3*ln(exp(I*(f*x+e)))-3/f*b*d^3*a^2*ln(exp(2*I*(f*x+e))+1)*x^3-9/2/f^3*b*a^2*c*d^2*polylog(3,-exp(2*I*(f*x+e)))-9/2/f^3*b*d^3*a^2*polylog(3,-exp(2*I*(f*x+e)))*x-6/f^4*b*e^3*a^2*d^3*ln(exp(I*(f*x+e)))-6/f^3*b^3*e^2*c*d^2*ln(exp(I*(f*x+e)))+3/f*b^3*d*c^2*ln(exp(2*I*(f*x+e))+1)*x+9/f^2*b^2*a*d^3*ln(exp(2*I*(f*x+e))+1)*x^2+3/f*b^3*c*d^2*ln(exp(2*I*(f*x+e))+1)*x^2+4*I/f^3*b^3*c*d^2*e^3+9/2*I/f^4*b*e^4*a^2*d^3-3/2*I/f^2*b^3*d^3*polylog(2,-exp(2*I*(f*x+e)))*x^2-6*I/f*b^2*a*d^3*x^3+12*I/f^4*b^2*a*d^3*e^3+36/f^3*b^2*e*a*c*d^2*ln(exp(I*(f*x+e)))+18/f^3*b*e^2*a^2*c*d^2*ln(exp(I*(f*x+e)))-9/f*b*a^2*c*d^2*ln(exp(2*I*(f*x+e))+1)*x^2-9/f*b*d*c^2*a^2*ln(exp(2*I*(f*x+e))+1)*x+18/f^2*b^2*d^2*c*a*ln(exp(2*I*(f*x+e))+1)*x-18/f^2*b*e*a^2*c^2*d*ln(exp(I*(f*x+e)))-12*I/f^3*b*a^2*c*d^2*e^3-18*I/f^3*b^2*d^2*c*a*e^2-9*I/f^3*b^2*d^2*c*a*polylog(2,-exp(2*I*(f*x+e)))+9/2*I/f^2*b*d*c^2*a^2*polylog(2,-exp(2*I*(f*x+e)))
\end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1177 vs. $2(530) = 1060$.

Time = 0.27 (sec) , antiderivative size = 1177, normalized size of antiderivative = 1.92

$$\int (c + dx)^3 (a + b \tan(f*x + e))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="fricas")
[Out] 1/8*(2*(a^3 - 3*a*b^2)*d^3*f^4*x^4 + 3*I*(3*a^2*b - b^3)*d^3*polylog(4, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*I*(3*a^2*b - b^3)*d^3*polylog(4, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 4*(b^3*d^3*f^3 + 2*(a^3 - 3*a*b^2)*c*d^2*f^4)*x^3 + 12*(b^3*c*d^2*f^3 + (a^3 - 3*a*b^2)*c^2*d*f^4)*x^2 + 4*(b^3*d^3*f^3*x^3 + 3*b^3*c*d^2*f^3*x^2 + 3*b^3*c^2*d*f^3*x + b^3*c^3*f^3)*tan(f*x + e)^2 + 4*(3*b^3*c^2*d*f^3 + 2*(a^3 - 3*a*b^2)*c^3*f^4)*x - 6*(I*(3*a^2*b - b^3)*d^3*f^2*x^2 - 6*I*a*b^2*c*d^2*f + I*b^3*d^3 + I*(3*a^2*b - b^3)*c^2*d*f^2 - 2*I*(3*a*b^2*d^3*f - (3*a^2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*(-I*(3*a^2*b - b^3)*d^3*f^2*x^2 + 6*I*a*b^2*c*d^2*f - I*b^3*d^3 - I*(3*a^2*b - b^3)*c^2*d*f^2 + 2*I*(3*a*b^2*d^3*f - (3*a^2*b - b^3)*c*d^2*f^2)*x)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 4*((3*a^2*b - b^3)*d^3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f + (3*a^2*b - b^3)*c^3*f^3 - 3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2 - b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 4*((3*a^2*b - b^3)*d^3*f^3*x^3 - 9*a*b^2*c^2*d*f^2 + 3*b^3*c*d^2*f^2 + (3*a^2*b - b^3)*c^3*f^3 - 3*(3*a*b^2*d^3*f^2 - (3*a^2*b - b^3)*c*d^2*f^3)*x^2 - 3*(6*a*b^2*c*d^2*f^2 - b^3*d^3*f - (3*a^2*b - b^3)*c^2*d*f^3)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(3*a*b^2*d^3 - (3*a^2*b - b^3)*d^3*f*x - (3*a^2*b - b^3)*c*d^2*f)*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(3*a*b^2*d^3 - (3*a^2*b - b^3)*d^3*f*x - (3*a^2*b - b^3)*c*d^2*f)*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 12*(2*a*b^2*d^3*f^3*x^3 + 2*a*b^2*c^3*f^3 - b^3*c^2*d*f^2 + (6*a*b^2*c*d^2*f^3 - b^3*d^3*f^2)*x^2 + 2*(3*a*b^2*c^2*d*f^3 - b^3*c*d^2*f^2)*x)*tan(f*x + e))/f^4
```

Sympy [F]

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 \, dx = \int (a + b \tan(e + fx))^3 (c + dx)^3 \, dx$$

```
[In] integrate((d*x+c)**3*(a+b*tan(f*x+e))**3,x)
```

[Out] $\text{Integral}((a + b \tan(e + f x))^3 (c + d x)^3, x)$

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 6861 vs. $2(530) = 1060$.

Time = 10.88 (sec) , antiderivative size = 6861, normalized size of antiderivative = 11.21

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/4*(4*(f*x + e)*a^3*c^3 + (f*x + e)^4*a^3*d^3/f^3 - 4*(f*x + e)^3*a^3*d^3*f^3
e/f^3 + 6*(f*x + e)^2*a^3*d^3*e^2/f^3 - 4*(f*x + e)*a^3*d^3*e^3/f^3 + 4*(f*x + e)^3*a^3*c*d^2/f^2 - 12*(f*x + e)^2*a^3*c*d^2*e/f^2 + 12*(f*x + e)*a^3*c*d^2*e^2/f^2 + 6*(f*x + e)^2*a^3*c^2*d/f - 12*(f*x + e)*a^3*c^2*d*e/f + 12*a^2*b*c^3*log(sec(f*x + e)) - 12*a^2*b*d^3*e^3*log(sec(f*x + e))/f^3 + 36*a^2*b*c*d^2*e^2*log(sec(f*x + e))/f^2 - 36*a^2*b*c^2*d*e*log(sec(f*x + e))/f - 4*(72*a*b^2*d^3*e^3 - 72*a*b^2*c^3*f^3 - 3*(3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^4*d^3 + 36*b^3*d^3*e^2 + 12*((3*a^2*b + 3*I*a*b^2 - b^3)*d^3*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*f)*(f*x + e)^3 - 18*((3*a^2*b + 3*I*a*b^2 - b^3)*d^3*e^2 - 2*(3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*e*f + (3*a^2*b + 3*I*a*b^2 - b^3)*c^2*d*f^2)*(f*x + e)^2 + 36*(6*a*b^2*c^2*d^2*e + b^3*c^2*d)*f^2 + 12*((3*I*a*b^2 - b^3)*d^3*e^3 + 3*(-3*I*a*b^2 + b^3)*c*d^2*e^2*f + 3*(3*I*a*b^2 - b^3)*c^2*d*e*f^2 + (-3*I*a*b^2 + b^3)*c^3*f^3)*(f*x + e) - 72*(3*a*b^2*c*d^2*e^2 + b^3*c*d^2*e)*f + 4*(3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*d^2*e)*f)*(f*x + e) - 9*(b^3*c*d^2*e^2 - 6*a*b^2*c*d^2*e - b^3*c*d^2)*f + (3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*c*d^2*e)*f)*(f*x + e) - 9*(b^3*c*d^2*f^2)^2)
```

$$\begin{aligned}
& e^2 - 6*a*b^2*c*d^2*e - b^3*c*d^2)*f)*cos(4*f*x + 4*e) + 2*(3*b^3*d^3*e^3 - \\
& 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b \\
& ^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f \\
&)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^ \\
& 3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c \\
& d^2 + (3*a^2*b - b^3)*c*d^2*e)*f)*(f*x + e) - 9*(b^3*c*d^2*e^2 - 6*a*b^2*c \\
& d^2*e - b^3*c*d^2)*f)*cos(2*f*x + 2*e) + (3*I*b^3*d^3*e^3 - 3*I*b^3*c^3*f^3 \\
& - 27*I*a*b^2*d^3*e^2 + 4*(3*I*a^2*b - I*b^3)*(f*x + e)^3*d^3 - 9*I*b^3*d^3 \\
& *e + 9*(-3*I*a*b^2*d^3 + (-3*I*a^2*b + I*b^3)*d^3*e + (3*I*a^2*b - I*b^3)*c \\
& *d^2*f)*(f*x + e)^2 + 9*(I*b^3*c^2*d*e - 3*I*a*b^2*c^2*d)*f^2 + 9*(6*I*a*b \\
& 2*d^3*e + I*b^3*d^3 + (3*I*a^2*b - I*b^3)*d^3*e^2 + (3*I*a^2*b - I*b^3)*c^2 \\
& *d*f^2 + 2*(-3*I*a*b^2*c*d^2 + (-3*I*a^2*b + I*b^3)*c*d^2*e)*f)*(f*x + e) + \\
& 9*(-I*b^3*c*d^2*e^2 + 6*I*a*b^2*c*d^2*e + I*b^3*c*d^2)*f)*sin(4*f*x + 4*e) \\
& + 2*(3*I*b^3*d^3*e^3 - 3*I*b^3*c^3*f^3 - 27*I*a*b^2*d^3*e^2 + 4*(3*I*a^2*b \\
& - I*b^3)*(f*x + e)^3*d^3 - 9*I*b^3*d^3*e + 9*(-3*I*a*b^2*d^3 + (-3*I*a^2*b \\
& + I*b^3)*d^3*e + (3*I*a^2*b - I*b^3)*c*d^2*f)*(f*x + e)^2 + 9*(I*b^3*c^2*d \\
& *e - 3*I*a*b^2*c^2*d)*f^2 + 9*(6*I*a*b^2*d^3*e + I*b^3*d^3 + (3*I*a^2*b - I \\
& *b^3)*d^3*e^2 + (3*I*a^2*b - I*b^3)*c^2*d*f^2 + 2*(-3*I*a*b^2*c*d^2 + (-3*I \\
& *a^2*b + I*b^3)*c*d^2*e)*f)*(f*x + e) + 9*(-I*b^3*c*d^2*e^2 + 6*I*a*b^2*c*d \\
& ^2*f)*(f*x + e)^3 + 6*(12*a*b^2*d^3*e + 2*b^3*d^3 + (3*a^2*b + 3*I*a*b^2 - b \\
& ^3)*d^3*e^2 + (3*a^2*b + 3*I*a*b^2 - b^3)*c^2*d*f^2 - 2*(6*a*b^2*c*d^2 + (3 \\
& *a^2*b + 3*I*a*b^2 - b^3)*c*d^2*e)*f^2 - 3*(12*a*b^2*c*d^2*e + 2*b^3*c \\
& *d^2 - (-3*I*a*b^2 + b^3)*c*d^2*e^2)*f)*(f*x + e))*cos(4*f*x + 4*e) - 6*((\\
& 3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^4*d^3 - 6*b^3*d^3*e^2 - 4*(3*a*b^2 - I \\
& *b^3)*d^3*e^3 + 4*(3*a*b^2 - I*b^3)*c^3*f^3 - 4*((3*a^2*b + 3*I*a*b^2 - b^3) \\
&)*d^3*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d^2*f + (3*a*b^2 + I*b^3)*d^3)*(f*x \\
& + e)^3 + 6*(b^3*d^3 + (3*a^2*b + 3*I*a*b^2 - b^3)*d^3*e^2 + (3*a^2*b + 3*I \\
& *a*b^2 - b^3)*c^2*d*f^2 + 2*(3*a*b^2 + I*b^3)*d^3*e - 2*((3*a^2*b + 3*I*a*b \\
& ^2 - b^3)*c*d^2*e + (3*a*b^2 + I*b^3)*c*d^2)*f)*(f*x + e)^2 - 6*(b^3*c^2*d \\
& + 2*(3*a*b^2 - I*b^3)*c^2*d*e)*f^2 - 4*(3*b^3*d^3*e + (3*I*a*b^2 - b^3)*d^3 \\
& *e^3 + (-3*I*a*b^2 + b^3)*c^3*f^3 + 3*(3*a*b^2 + I*b^3)*d^3*e^2 + 3*((3*I*a \\
& *b^2 - b^3)*c^2*d*e + (3*a*b^2 + I*b^3)*c^2*d)*f^2 - 3*(b^3*c*d^2 - (-3*I*a \\
& *b^2 + b^3)*c*d^2*e^2 + 2*(3*a*b^2 + I*b^3)*c*d^2*e)*f)*(f*x + e) + 12*(b^3 \\
& *c*d^2*e + (3*a*b^2 - I*b^3)*c*d^2*e^2)*f)*cos(2*f*x + 2*e) - 6*(18*a*b^2*d \\
& ^3*e + 4*(3*a^2*b - b^3)*(f*x + e)^2*d^3 + 3*b^3*d^3 + 3*(3*a^2*b - b^3)*d^3 \\
& *e^2 + 3*(3*a^2*b - b^3)*c^2*d*f^2 - 6*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3* \\
& e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e) - 6*(3*a*b^2*c*d^2 + (3*a^2*b - b^3) \\
& *c*d^2*e)*f + (18*a*b^2*d^3*e + 4*(3*a^2*b - b^3)*(f*x + e)^2*d^3 + 3*b^3*d \\
& ^3 + 3*(3*a^2*b - b^3)*d^3*e^2 + 3*(3*a^2*b - b^3)*c^2*d*f^2 - 6*(3*a*b^2*d \\
& ^3 + (3*a^2*b - b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e) - 6*(3*a*b^2
\end{aligned}$$

$$\begin{aligned}
& 2*c*d^2 + (3*a^2*b - b^3)*c*d^2*f)*cos(4*f*x + 4*e) + 2*(18*a*b^2*d^3*e \\
& + 4*(3*a^2*b - b^3)*(f*x + e)^2*d^3 + 3*b^3*d^3 + 3*(3*a^2*b - b^3)*d^3*e^2 \\
& + 3*(3*a^2*b - b^3)*c^2*d*f^2 - 6*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - \\
& (3*a^2*b - b^3)*c*d^2*f)*(f*x + e) - 6*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*c*d^2 \\
& 2*f)*cos(2*f*x + 2*e) - (-18*I*a*b^2*d^3*e + 4*(-3*I*a^2*b + I*b^3)*(f*x \\
& + e)^2*d^3 - 3*I*b^3*d^3 + 3*(-3*I*a^2*b + I*b^3)*d^3*e^2 + 3*(-3*I*a^2*b \\
& + I*b^3)*c^2*d*f^2 + 6*(3*I*a*b^2*d^3 + (3*I*a^2*b - I*b^3)*d^3*e + (-3*I*a \\
& ^2*b + I*b^3)*c*d^2*f)*(f*x + e) + 6*(3*I*a*b^2*c*d^2 + (3*I*a^2*b - I*b^3) \\
& *c*d^2*e)*f)*sin(4*f*x + 4*e) - 2*(-18*I*a*b^2*d^3*e + 4*(-3*I*a^2*b + I*b^3) \\
& *(f*x + e)^2*d^3 - 3*I*b^3*d^3 + 3*(-3*I*a^2*b + I*b^3)*d^3*e^2 + 3*(-3*I \\
& *a^2*b + I*b^3)*c^2*d*f^2 + 6*(3*I*a*b^2*d^3 + (3*I*a^2*b - I*b^3)*d^3*e + \\
& (-3*I*a^2*b + I*b^3)*c*d^2*f)*(f*x + e) + 6*(3*I*a*b^2*c*d^2 + (3*I*a^2*b - \\
& I*b^3)*c*d^2*e)*f)*sin(2*f*x + 2*e))*dilog(-e^(2*I*f*x + 2*I*e)) + 2*(-3*I \\
& *b^3*d^3*e^3 + 3*I*b^3*c^3*f^3 + 27*I*a*b^2*d^3*e^2 + 4*(-3*I*a^2*b + I*b^3) \\
&)*(f*x + e)^3*d^3 + 9*I*b^3*d^3*e + 9*(3*I*a*b^2*d^3 + (3*I*a^2*b - I*b^3)* \\
& d^3*e + (-3*I*a^2*b + I*b^3)*c*d^2*f)*(f*x + e)^2 + 9*(-I*b^3*c^2*d*e + 3*I \\
& *a*b^2*c^2*d)*f^2 + 9*(-6*I*a*b^2*d^3*e - I*b^3*d^3 + (-3*I*a^2*b + I*b^3)* \\
& d^3*e^2 + (-3*I*a^2*b + I*b^3)*c^2*d*f^2 + 2*(3*I*a*b^2*c*d^2 + (3*I*a^2*b \\
& - I*b^3)*c*d^2*e)*f)*(f*x + e) + 9*(I*b^3*c*d^2*e^2 - 6*I*a*b^2*c*d^2*e - \\
& I*b^3*c*d^2)*f + (-3*I*b^3*d^3*e^3 + 3*I*b^3*c^3*f^3 + 27*I*a*b^2*d^3*e^2 + \\
& 4*(-3*I*a^2*b + I*b^3)*(f*x + e)^3*d^3 + 9*I*b^3*d^3*e + 9*(3*I*a*b^2*d^3 + \\
& (3*I*a^2*b - I*b^3)*d^3*e + (-3*I*a^2*b + I*b^3)*c*d^2*f)*(f*x + e)^2 + 9* \\
& (-I*b^3*c^2*d*e + 3*I*a*b^2*c^2*d)*f^2 + 9*(-6*I*a*b^2*d^3*e - I*b^3*d^3 + \\
& (-3*I*a^2*b + I*b^3)*d^3*e^2 + (-3*I*a^2*b + I*b^3)*c^2*d*f^2 + 2*(3*I*a*b^ \\
& 2*c*d^2 + (3*I*a^2*b - I*b^3)*c*d^2*e)*f)*(f*x + e) + 9*(I*b^3*c*d^2*e^2 - \\
& 6*I*a*b^2*c*d^2*e - I*b^3*c*d^2)*f)*cos(4*f*x + 4*e) + 2*(-3*I*b^3*d^3*e^3 \\
& + 3*I*b^3*c^3*f^3 + 27*I*a*b^2*d^3*e^2 + 4*(-3*I*a^2*b + I*b^3)*(f*x + e)^3 \\
& *d^3 + 9*I*b^3*d^3*e + 9*(3*I*a*b^2*d^3 + (3*I*a^2*b - I*b^3)*d^3*e + (-3*I \\
& *a^2*b + I*b^3)*c*d^2*f)*(f*x + e)^2 + 9*(-I*b^3*c^2*d*e + 3*I*a*b^2*c^2*d) \\
& *f^2 + 9*(-6*I*a*b^2*d^3*e - I*b^3*d^3 + (-3*I*a^2*b + I*b^3)*d^3*e^2 + (-3 \\
& *I*a^2*b + I*b^3)*c^2*d*f^2 + 2*(3*I*a*b^2*c*d^2 + (3*I*a^2*b - I*b^3)*c*d^ \\
& 2*e)*f)*(f*x + e) + 9*(I*b^3*c*d^2*e^2 - 6*I*a*b^2*c*d^2*e - I*b^3*c*d^2)*f \\
&)*cos(2*f*x + 2*e) + (3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4* \\
& (3*a^2*b - b^3)*(f*x + e)^3*d^3 - 9*b^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - \\
& b^3)*d^3*e - (3*a^2*b - b^3)*c*d^2*f)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b \\
& ^2*c^2*d)*f^2 + 9*(6*a*b^2*d^3*e + b^3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a \\
& ^2*b - b^3)*c^2*d*f^2 - 2*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*c*d^2*e)*f)*(f*x \\
& + e) - 9*(b^3*c*d^2*e^2 - 6*a*b^2*c*d^2*e - b^3*c*d^2)*f)*sin(4*f*x + 4*e) \\
& + 2*(3*b^3*d^3*e^3 - 3*b^3*c^3*f^3 - 27*a*b^2*d^3*e^2 + 4*(3*a^2*b - b^3)* \\
& (f*x + e)^3*d^3 - 9*b^3*d^3*e - 9*(3*a*b^2*d^3 + (3*a^2*b - b^3)*d^3*e - (3 \\
& *a^2*b - b^3)*c*d^2*f)*(f*x + e)^2 + 9*(b^3*c^2*d*e - 3*a*b^2*c^2*d)*f^2 + \\
& 9*(6*a*b^2*d^3*e + b^3*d^3 + (3*a^2*b - b^3)*d^3*e^2 + (3*a^2*b - b^3)*c^2* \\
& d*f^2 - 2*(3*a*b^2*c*d^2 + (3*a^2*b - b^3)*c*d^2*e)*f)*(f*x + e) - 9*(b^3*c \\
& *d^2*e^2 - 6*a*b^2*c*d^2*e - b^3*c*d^2)*f)*sin(2*f*x + 2*e))*log(cos(2*f*x \\
& + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) + 12*((3*a^2*b - b^
\end{aligned}$$

$$\begin{aligned}
& 3*d^3*\cos(4*f*x + 4*e) + 2*(3*a^2*b - b^3)*d^3*\cos(2*f*x + 2*e) + (3*I*a^2*b - I*b^3)*d^3*\sin(4*f*x + 4*e) + 2*(3*I*a^2*b - I*b^3)*d^3*\sin(2*f*x + 2*e) + (3*a^2*b - b^3)*d^3)*\text{polylog}(4, -e^{(2*I*f*x + 2*I*e)}) + 6*(9*I*a*b^2*d^3 + 4*(-3*I*a^2*b + I*b^3)*(f*x + e)*d^3 + 3*(3*I*a^2*b - I*b^3)*d^3*e + 3*(-3*I*a^2*b + I*b^3)*c*d^2*f + (9*I*a*b^2*d^3 + 4*(-3*I*a^2*b + I*b^3)*(f*x + e)*d^3 + 3*(3*I*a^2*b - I*b^3)*d^3*e + 3*(-3*I*a^2*b + I*b^3)*c*d^2*f)*\cos(4*f*x + 4*e) + 2*(9*I*a*b^2*d^3 + 4*(-3*I*a^2*b + I*b^3)*(f*x + e)*d^3 + 3*(3*I*a^2*b - I*b^3)*d^3*e + 3*(-3*I*a^2*b + I*b^3)*c*d^2*f)*\cos(2*f*x + 2*e) - (9*a*b^2*d^3 - 4*(3*a^2*b - b^3)*(f*x + e)*d^3 + 3*(3*a^2*b - b^3)*d^3*e - 3*(3*a^2*b - b^3)*c*d^2*f)*\sin(4*f*x + 4*e) - 2*(9*a*b^2*d^3 - 4*(3*a^2*b - b^3)*(f*x + e)*d^3 + 3*(3*a^2*b - b^3)*d^3*e - 3*(3*a^2*b - b^3)*c*d^2*f)*\sin(2*f*x + 2*e))*\text{polylog}(3, -e^{(2*I*f*x + 2*I*e)}) + 3*((-3*I*a^2*b + 3*a*b^2 + I*b^3)*(f*x + e)^4*d^3 + 4*(6*I*a*b^2*d^3 + (3*I*a^2*b - 3*a*b^2 - I*b^3)*d^3*e + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c*d^2*f)*(f*x + e)^3 + 6*(-12*I*a*b^2*d^3*e - 2*I*b^3*d^3 + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*d^3*e^2 + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c^2*d*f^2 + 2*(6*I*a*b^2*c*d^2 + (3*I*a^2*b - 3*a*b^2 - I*b^3)*c*d^2*e - (3*a*b^2 + I*b^3)*d^3*e^3 + (3*a*b^2 + I*b^3)*c^3*f^3 + 3*(6*I*a*b^2*c^2*d - (3*a*b^2 + I*b^3)*c^2*d*e)*f^2 + 3*(-12*I*a*b^2*c*d^2*e - 2*I*b^3*c*d^2 + (3*a*b^2 + I*b^3)*c*d^2*e^2)*f)*(f*x + e)^2 + 4*(18*I*a*b^2*d^3*e^2 + 6*I*b^3*d^3*e - (3*a*b^2 + I*b^3)*d^3*e^3 + (3*a*b^2 + I*b^3)*c^3*f^3 + 3*(6*I*a*b^2*c^2*d - (3*a*b^2 + I*b^3)*c^2*d*e)*f^2 + 3*(-12*I*a*b^2*c*d^2*e - 2*I*b^3*c*d^2 + (3*a*b^2 + I*b^3)*c*d^2*e^2)*f)*(f*x + e)))*\sin(4*f*x + 4*e) + 6*((-3*I*a^2*b + 3*a*b^2 + I*b^3)*(f*x + e)^4*d^3 + 6*I*b^3*d^3*e^2 + 4*(3*I*a*b^2 + b^3)*d^3*e^3 + 4*(-3*I*a*b^2 - b^3)*c^3*f^3 + 4*((3*I*a^2*b - 3*a*b^2 - I*b^3)*d^3*e + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c*d^2*f + (3*I*a*b^2 - b^3)*d^3)*(f*x + e)^3 + 6*(-I*b^3*d^3 + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*d^3)*e^2 + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*d^3 + 2*((3*I*a^2*b - 3*a*b^2 - I*b^3)*c*d^2*e + (3*I*a*b^2 - b^3)*c*d^2*f)*(f*x + e)^2 + 6*(I*b^3*c^2*d + 2*(3*I*a*b^2 + b^3)*c^2*d*e)*f^2 + 4*(3*I*b^3*d^3*e - (3*a*b^2 + I*b^3)*d^3*e^3 + (3*a*b^2 + I*b^3)*c^3*f^3 + 3*(3*I*a*b^2 - b^3)*d^3*e^2 - 3*((3*a*b^2 + I*b^3)*c^2*d*e - (3*I*a*b^2 - b^3)*c^2*d)f^2 + 3*(-I*b^3*c*d^2 + (3*a*b^2 + I*b^3)*c*d^2*e^2 + 2*(-3*I*a*b^2 + b^3)*d^3 + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c^2*d*f^2 + 2*(6*I*a*b^2*c*d^2 - (3*a*b^2 + I*b^3)*c*d^2*e)*f)*(f*x + e) + 12*(-I*b^3*c*d^2*e + (-3*I*a*b^2 - b^3)*c*d^2*e^2)*f)*\sin(2*f*x + 2*e))/(-12*I*f^3*\cos(4*f*x + 4*e) - 24*I*f^3*\cos(2*f*x + 2*e) + 12*f^3*\sin(4*f*x + 4*e) + 24*f^3*\sin(2*f*x + 2*e) - 12*I*f^3))/f
\end{aligned}$$

Giac [F]

$$\int (c + dx)^3(a + b \tan(e + fx))^3 dx = \int (dx + c)^3(b \tan(fx + e) + a)^3 dx$$

```
[In] integrate((d*x+c)^3*(a+b*tan(f*x+e))^3,x, algorithm="giac")
[Out] integrate((d*x + c)^3*(b*tan(f*x + e) + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + f x))^3 (c + d x)^3 dx$$

[In] `int((a + b*tan(e + f*x))^3*(c + d*x)^3,x)`

[Out] `int((a + b*tan(e + f*x))^3*(c + d*x)^3, x)`

$$\mathbf{3.50} \quad \int (c + dx)^2 (a + b \tan(e + fx))^3 \, dx$$

Optimal result	329
Rubi [A] (verified)	330
Mathematica [B] (warning: unable to verify)	335
Maple [B] (verified)	337
Fricas [A] (verification not implemented)	338
Sympy [F]	339
Maxima [B] (verification not implemented)	339
Giac [F]	341
Mupad [F(-1)]	341

Optimal result

Integrand size = 20, antiderivative size = 436

$$\begin{aligned}
 \int (c + dx)^2 (a + b \tan(e + fx))^3 dx = & \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3iab^2(c + dx)^2}{f} + \frac{a^3(c + dx)^3}{3d} \\
 & + \frac{ia^2 b(c + dx)^3}{d} - \frac{ab^2(c + dx)^3}{d} - \frac{ib^3(c + dx)^3}{3d} \\
 & + \frac{6ab^2 d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2 b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & + \frac{b^3(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\
 & - \frac{b^3 d^2 \log(\cos(e + fx))}{f^3} \\
 & - \frac{3iab^2 d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
 & + \frac{3ia^2 bd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{ib^3 d(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
 & - \frac{3a^2 bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & + \frac{b^3 d^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
 & - \frac{b^3 d(c + dx) \tan(e + fx)}{f^2} \\
 & + \frac{3ab^2(c + dx)^2 \tan(e + fx)}{f} \\
 & + \frac{b^3(c + dx)^2 \tan^2(e + fx)}{2f}
 \end{aligned}$$

```
[Out] b^3*c*d*x/f+1/2*b^3*d^2*x^2/f-3*I*a*b^2*(d*x+c)^2/f+1/3*a^3*(d*x+c)^3/d-I*b^3*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-a*b^2*(d*x+c)^3/d+I*a^2*b*(d*x+c)^3/d+6*a*b^2*d*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f^2-3*a^2*b*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f+b^3*(d*x+c)^2*ln(1+exp(2*I*(f*x+e)))/f-b^3*d^2*ln(cos(f*x+e))/f^3+3*I*a^2*b*d*(d*x+c)*polylog(2,-exp(2*I*(f*x+e)))/f^2-3*I*a*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3-1/3*I*b^3*(d*x+c)^3/d-3/2*a^2*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+1/2*b^3*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3-b^3*d*(d*x+c)*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)^2*tan(f*x+e)/f+1/2*b^3*(d*x+c)^2*tan(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules used = {3803, 3800, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32, 3556}

$$\begin{aligned} \int (c + dx)^2 (a + b \tan(e + fx))^3 dx = & \frac{a^3(c + dx)^3}{3d} + \frac{3ia^2bd(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{3a^2b(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{ia^2b(c + dx)^3}{d} - \frac{3a^2bd^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\ & + \frac{6ab^2d(c + dx) \log(1 + e^{2i(e+fx)})}{f^2} \\ & + \frac{3ab^2(c + dx)^2 \tan(e + fx)}{f} - \frac{3iab^2(c + dx)^2}{f} \\ & - \frac{ab^2(c + dx)^3}{d} - \frac{3iab^2d^2 \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\ & - \frac{ib^3d(c + dx) \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\ & - \frac{b^3d(c + dx) \tan(e + fx)}{f^2} \\ & + \frac{b^3(c + dx)^2 \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^3(c + dx)^2 \tan^2(e + fx)}{2f} + \frac{b^3cdx}{f} \\ & - \frac{ib^3(c + dx)^3}{3d} + \frac{b^3d^2 \operatorname{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\ & - \frac{b^3d^2 \log(\cos(e + fx))}{f^3} + \frac{b^3d^2x^2}{2f} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2*(a + b*\operatorname{Tan}[e + f*x])^3, x]$

[Out] $(b^{3*c*d*x}/f + (b^{3*d^2*x^2}/(2*f) - ((3*I)*a*b^{2*(c + d*x)^2}/f + (a^{3*(c + d*x)^3}/(3*d) + (I*a^{2*b*(c + d*x)^3}/d - (a*b^{2*(c + d*x)^3})/d - ((I/3)*b^{3*(c + d*x)^3}/d + (6*a*b^{2*d*(c + d*x)}*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f^2 - (3*a^{2*b*(c + d*x)^2}*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (b^{3*(c + d*x)^2}*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f - (b^{3*d^2*2*\operatorname{Log}[\operatorname{Cos}[e + f*x]]}/f^3 - ((3*I)*a*b^{2*d^2*2*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^3 + ((3*I)*a^{2*b*d*(c + d*x)}*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - (I*b^{3*d*(c + d*x)}*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - (3*a^{2*b*d^2*2*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/(2*f^3) + (b^{3*d^2*2*\operatorname{PolyLog}[3, -E^{((2*I)*(e + f*x))}])/(2*f^3) - (b^{3*d*(c + d*x)}))$

$$)*\text{Tan}[e + f*x])/f^2 + (3*a*b^2*(c + d*x)^2*\text{Tan}[e + f*x])/f + (b^3*(c + d*x)^2*\text{Tan}[e + f*x]^2)/(2*f)$$
Rule 32

$$\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /; \text{FreeQ}[\{a, b, m\}, x] \&& \text{NeQ}[m, -1]$$
Rule 2221

$$\text{Int}[((F_.)^{(g_.)*(e_.) + (f_.)*(x_.)})^{(n_.)}*((c_.) + (d_.)*(x_.)^{(m_.)})/((a_.) + (b_.)*(F_.)^{(g_.)*(e_.) + (f_.)*(x_.)})^{(n_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{(m - 1)}*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$$
Rule 2317

$$\text{Int}[\text{Log}[(a_.) + (b_.)*(F_.)^{(e_.)*(c_.) + (d_.)*(x_.)})^{(n_.)}], x_{\text{Symbol}}] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$$
Rule 2320

$$\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /; \text{FunctionOfExponentialQ}[u, x] \&& \text{!MatchQ}[u, (w_)*(a_)*(v_)^{(n_.)}^{(m_.)} /; \text{FreeQ}[\{a, m, n\}, x] \&& \text{IntegerQ}[m*n]] \&& \text{!MatchQ}[u, E^{(c_.)*(a_.) + (b_.)*x})*(F_)[v_] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{InverseFunctionQ}[F[x]]]$$
Rule 2438

$$\text{Int}[\text{Log}[(c_.)*(d_.) + (e_.)*(x_.)^{(n_.)})]/(x_), x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \&& \text{EqQ}[c*d, 1]$$
Rule 2611

$$\text{Int}[\text{Log}[1 + (e_.)*(F_.)^{(c_.)*(a_.) + (b_.)*(x_.)})^{(n_.)}*((f_.) + (g_.)*(x_.)^{(m_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*\text{Log}[F])), x] + \text{Dist}[g*(m/(b*c*n*\text{Log}[F])), \text{Int}[(f + g*x)^{(m - 1)}*\text{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&& \text{GtQ}[m, 0]$$
Rule 3556

$$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$$

Rule 3800

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_)*(x_))^(m_)*((a_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*((a_.) + (b_)*x)^p]/((d_.) + (e_)*x), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^3(c + dx)^2 + 3a^2b(c + dx)^2 \tan(e + fx) + 3ab^2(c + dx)^2 \tan^2(e + fx) \\
&\quad + b^3(c + dx)^2 \tan^3(e + fx)) \, dx \\
&= \frac{a^3(c + dx)^3}{3d} + (3a^2b) \int (c + dx)^2 \tan(e + fx) \, dx \\
&\quad + (3ab^2) \int (c + dx)^2 \tan^2(e + fx) \, dx + b^3 \int (c + dx)^2 \tan^3(e + fx) \, dx \\
&= \frac{a^3(c + dx)^3}{3d} + \frac{ia^2b(c + dx)^3}{d} + \frac{3ab^2(c + dx)^2 \tan(e + fx)}{f} \\
&\quad + \frac{b^3(c + dx)^2 \tan^2(e + fx)}{2f} - (6ia^2b) \int \frac{e^{2i(e+fx)}(c + dx)^2}{1 + e^{2i(e+fx)}} \, dx \\
&\quad - (3ab^2) \int (c + dx)^2 \, dx - b^3 \int (c + dx)^2 \tan(e + fx) \, dx \\
&\quad - \frac{(6ab^2d) \int (c + dx) \tan(e + fx) \, dx}{f} - \frac{(b^3d) \int (c + dx) \tan^2(e + fx) \, dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3iab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} + \frac{ia^2b(c+dx)^3}{d} - \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{ib^3(c+dx)^3}{3d} - \frac{3a^2b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{b^3d(c+dx) \tan(e+fx)}{f^2} + \frac{3ab^2(c+dx)^2 \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^2 \tan^2(e+fx)}{2f} + (2ib^3) \int \frac{e^{2i(e+fx)}(c+dx)^2}{1+e^{2i(e+fx)}} dx \\
&\quad + \frac{(b^3d^2) \int \tan(e+fx) dx}{f^2} + \frac{(6a^2bd) \int (c+dx) \log(1+e^{2i(e+fx)}) dx}{f} \\
&\quad + \frac{(12iab^2d) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} dx}{f} + \frac{(b^3d) \int (c+dx) dx}{f} \\
&= \frac{b^3cdx}{f} + \frac{b^3d^2x^2}{2f} - \frac{3iab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} + \frac{ia^2b(c+dx)^3}{d} \\
&\quad - \frac{ab^2(c+dx)^3}{d} - \frac{ib^3(c+dx)^3}{3d} + \frac{6ab^2d(c+dx) \log(1+e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2b(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^2 \log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{b^3d^2 \log(\cos(e+fx))}{f^3} + \frac{3ia^2bd(c+dx) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{b^3d(c+dx) \tan(e+fx)}{f^2} + \frac{3ab^2(c+dx)^2 \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx)^2 \tan^2(e+fx)}{2f} - \frac{(3ia^2bd^2) \int \text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6ab^2d^2) \int \log(1+e^{2i(e+fx)}) dx}{f^2} - \frac{(2b^3d) \int (c+dx) \log(1+e^{2i(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c d x}{f} + \frac{b^3 d^2 x^2}{2 f} - \frac{3 i a b^2 (c + d x)^2}{f} + \frac{a^3 (c + d x)^3}{3 d} + \frac{i a^2 b (c + d x)^3}{d} \\
&\quad - \frac{a b^2 (c + d x)^3}{d} - \frac{i b^3 (c + d x)^3}{3 d} + \frac{6 a b^2 d (c + d x) \log(1 + e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3 a^2 b (c + d x)^2 \log(1 + e^{2i(e+fx)})}{f} + \frac{b^3 (c + d x)^2 \log(1 + e^{2i(e+fx)})}{f} \\
&\quad - \frac{b^3 d^2 \log(\cos(e + f x))}{f^3} + \frac{3 i a^2 b d (c + d x) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{i b^3 d (c + d x) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{b^3 d (c + d x) \tan(e + f x)}{f^2} \\
&\quad + \frac{3 a b^2 (c + d x)^2 \tan(e + f x)}{f} + \frac{b^3 (c + d x)^2 \tan^2(e + f x)}{2 f} \\
&\quad - \frac{(3 a^2 b d^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{2 f^3} \\
&\quad + \frac{(3 i a b^2 d^2) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{f^3} + \frac{(i b^3 d^2) \int \text{PolyLog}(2, -e^{2i(e+fx)}) dx}{f^2} \\
&= \frac{b^3 c d x}{f} + \frac{b^3 d^2 x^2}{2 f} - \frac{3 i a b^2 (c + d x)^2}{f} + \frac{a^3 (c + d x)^3}{3 d} \\
&\quad + \frac{i a^2 b (c + d x)^3}{d} - \frac{a b^2 (c + d x)^3}{d} - \frac{i b^3 (c + d x)^3}{3 d} \\
&\quad + \frac{6 a b^2 d (c + d x) \log(1 + e^{2i(e+fx)})}{f^2} - \frac{3 a^2 b (c + d x)^2 \log(1 + e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3 (c + d x)^2 \log(1 + e^{2i(e+fx)})}{f} - \frac{b^3 d^2 \log(\cos(e + f x))}{f^3} \\
&\quad - \frac{3 i a b^2 d^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} + \frac{3 i a^2 b d (c + d x) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{i b^3 d (c + d x) \text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{3 a^2 b d^2 \text{PolyLog}(3, -e^{2i(e+fx)})}{2 f^3} \\
&\quad - \frac{b^3 d (c + d x) \tan(e + f x)}{f^2} + \frac{3 a b^2 (c + d x)^2 \tan(e + f x)}{f} \\
&\quad + \frac{b^3 (c + d x)^2 \tan^2(e + f x)}{2 f} + \frac{(b^3 d^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2i(e+fx)}\right)}{2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3iab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} + \frac{ia^2b(c+dx)^3}{d} \\
&\quad - \frac{ab^2(c+dx)^3}{d} - \frac{ib^3(c+dx)^3}{3d} + \frac{6ab^2d(c+dx)\log(1+e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2b(c+dx)^2\log(1+e^{2i(e+fx)})}{f} + \frac{b^3(c+dx)^2\log(1+e^{2i(e+fx)})}{f} \\
&\quad - \frac{b^3 d^2 \log(\cos(e+fx))}{f^3} - \frac{3iab^2 d^2 \text{PolyLog}(2, -e^{2i(e+fx)})}{f^3} \\
&\quad + \frac{3ia^2bd(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} - \frac{ib^3d(c+dx)\text{PolyLog}(2, -e^{2i(e+fx)})}{f^2} \\
&\quad - \frac{3a^2bd^2\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} + \frac{b^3d^2\text{PolyLog}(3, -e^{2i(e+fx)})}{2f^3} \\
&\quad - \frac{b^3d(c+dx)\tan(e+fx)}{f^2} + \frac{3ab^2(c+dx)^2\tan(e+fx)}{f} + \frac{b^3(c+dx)^2\tan^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1846 vs. $2(436) = 872$.

Time = 7.16 (sec) , antiderivative size = 1846, normalized size of antiderivative = 4.23

$$\begin{aligned}
& \int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \\
& - \frac{i a^2 b d^2 e^{-ie} (2 f^2 x^2 (2 f x - 3 i (1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6 (1 + e^{2ie}) f x \text{PolyLog}(2, -e^{-2i(e+fx)}) - 3 i (1 + e^{2ie}) f^3 \log(1 + e^{-2i(e+fx)}))}{4 f^3} \\
& + \frac{i b^3 d^2 e^{-ie} (2 f^2 x^2 (2 f x - 3 i (1 + e^{2ie}) \log(1 + e^{-2i(e+fx)})) + 6 (1 + e^{2ie}) f x \text{PolyLog}(2, -e^{-2i(e+fx)}) - 3 i (1 + e^{2ie}) f^3 \log(1 + e^{-2i(e+fx)}))}{12 f^3} \\
& - \frac{b^3 d^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f^3 (\cos^2(e) + \sin^2(e))} \\
& + \frac{6 a b^2 c d \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f^2 (\cos^2(e) + \sin^2(e))} \\
& - \frac{3 a^2 b c^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f (\cos^2(e) + \sin^2(e))} \\
& + \frac{b^3 c^2 \sec(e) (\cos(e) \log(\cos(e) \cos(fx) - \sin(e) \sin(fx)) + f x \sin(e))}{f (\cos^2(e) + \sin^2(e))} \\
& + \frac{3 a b^2 d^2 \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))})}{f^3 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{3 a^2 b c d \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))})}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)} \\
& + \frac{b^3 c d \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))})}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)}{3 a^2 b c d \csc(e) \left(e^{-i \arctan(\cot(e))} f^2 x^2 - \frac{\cot(e) (i f x (-\pi - 2 \arctan(\cot(e))) - \pi \log(1 + e^{-2i f x}) - 2 (f x - \arctan(\cot(e))) \log(1 - e^{2i(f x - \arctan(\cot(e)))})}{f^2 \sqrt{\csc^2(e) (\cos^2(e) + \sin^2(e))}} \right)} \\
& + \frac{\sec(e) \sec^2(e + f x) (6 b^3 c^2 f \cos(e) + 12 b^3 c d f x \cos(e) + 6 a^3 c^2 f^2 x \cos(e) - 18 a b^2 c^2 f^2 x \cos(e) + 6 b^3 d^2 f x^2 \cos(e))}{\sin(e) \sin(f x) (6 b^3 c^2 f \cos(e) + 12 b^3 c d f x \cos(e) + 6 a^3 c^2 f^2 x \cos(e) - 18 a b^2 c^2 f^2 x \cos(e) + 6 b^3 d^2 f x^2 \cos(e))}
\end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Tan[e + f*x])^3,x]

[Out]
$$\begin{aligned}
& ((-1/4*I)*a^2*b*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))] - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e])/((E^(I*e)*f^3) + ((I/12)*b^3*d^2*(2*f^2*x^2*(2*f*x - (3*I)*(1 + E^((2*I)*e)))*Log[1 + E^((-2*I)*(e + f*x))]) + 6*(1 + E^((2*I)*e))*f*x*PolyLog[2, -E^((-2*I)*(e + f*x))]) - (3*I)*(1 + E^((2*I)*e))*PolyLog[3, -E^((-2*I)*(e + f*x))]*Sec[e]/((E^(I*e)*f^3) - (b^3*d^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^3*(Cos[e]^2 + Sin[e]^2)) + (6*a*b^2*c*d*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f^2*(Cos[e]^2 + Sin[e]^2)) - (3*a^2*b*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (b^3*c^2*Sec[e]*(Cos[e]*Log[Cos[e]*Cos[f*x] - Sin[e]*Sin[f*x]] + f*x*Sin[e]))/(f*(Cos[e]^2 + Sin[e]^2)) + (3*a*b^2*d^2*2*Csc[e]*((f^2*x^2)/E^(I*ArcTan[Cot[e]])) - (Cot[e]*(I*f*x*(-Pi - 2*ArcTan[Cot[e]])) - Pi*Log[1 + E^((-2*I)*f*x)]) - 2*(f*x - ArcT
\end{aligned}$$

$$\begin{aligned}
& \text{an}[\text{Cot}[e]] * \text{Log}[1 - E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e])))}}] + \text{Pi} * \text{Log}[\text{Cos}[f*x]] - \\
& 2 * \text{ArcTan}[\text{Cot}[e]] * \text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + I * \text{PolyLog}[2, E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]))}})] / \text{Sqrt}[1 + \text{Cot}[e]^2] * \text{Sec}[e] / (f^3 * \text{Sqrt}[\text{Csc}[e]^2 * (\text{Cos}[e]^2 + \text{Sin}[e]^2)]) - (3*a^2 * b * c * d * \text{Csc}[e] * ((f^2 * x^2) / E^{\text{(I*ArcTan}[\text{Cot}[e])}}) - \\
& (\text{Cot}[e] * (I * f * x * (-\text{Pi} - 2 * \text{ArcTan}[\text{Cot}[e]])) - \text{Pi} * \text{Log}[1 + E^{\text{((-2*I)*f*x)}]}] - 2 * (f*x - \text{ArcTan}[\text{Cot}[e]]) * \text{Log}[1 - E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]))}})] + \text{Pi} * \text{Log}[\text{Cos}[f*x]] - 2 * \text{ArcTan}[\text{Cot}[e]] * \text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + I * \text{PolyLog}[2, E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]))}})] / \text{Sqrt}[1 + \text{Cot}[e]^2] * \text{Sec}[e] / (f^2 * \text{Sqrt}[\text{Csc}[e]^2 * (\text{Cos}[e]^2 + \text{Sin}[e]^2)]) + (b^3 * c * d * \text{Csc}[e] * ((f^2 * x^2) / E^{\text{(I*ArcTan}[\text{Cot}[e])}}) - (\text{Cot}[e] * (I * f * x * (-\text{Pi} - 2 * \text{ArcTan}[\text{Cot}[e]])) - \text{Pi} * \text{Log}[1 + E^{\text{((-2*I)*f*x)}]}] - 2 * (f*x - \text{ArcTan}[\text{Cot}[e]]) * \text{Log}[1 - E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]))}})] + \text{Pi} * \text{Log}[\text{Cos}[f*x]] - 2 * \text{ArcTan}[\text{Cot}[e]] * \text{Log}[\text{Sin}[f*x - \text{ArcTan}[\text{Cot}[e]]]] + I * \text{PolyLog}[2, E^{\text{((2*I)*(f*x - \text{ArcTan}[\text{Cot}[e]))}})] / \text{Sqrt}[1 + \text{Cot}[e]^2] * \text{Sec}[e] / (f^2 * \text{Sqrt}[\text{Csc}[e]^2 * (\text{Cos}[e]^2 + \text{Sin}[e]^2)]) + (\text{Sec}[e] * \text{Sec}[e + f*x]^2 * (6 * b^3 * c^2 * f * \text{Cos}[e] + 12 * b^3 * c * d * f * x * \text{Cos}[e] + 6 * a^3 * c^2 * f^2 * x^2 * \text{Cos}[e] - 18 * a * b^2 * c^2 * f^2 * x * \text{Cos}[e] + 6 * b^3 * d^2 * f^2 * x^2 * \text{Cos}[e] + 6 * a^3 * c * d * f^2 * x^2 * \text{Cos}[e] - 18 * a * b^2 * c * d * f^2 * x^2 * \text{Cos}[e] + 2 * a^3 * d^2 * f^2 * x^2 * \text{Cos}[e] - 6 * a * b^2 * d^2 * f^2 * x^2 * \text{Cos}[e] + 3 * a^3 * c^2 * f^2 * x * \text{Cos}[e + 2 * f*x] - 9 * a * b^2 * c^2 * f^2 * x * \text{Cos}[e + 2 * f*x] + 3 * a^3 * c * d * f^2 * x^2 * \text{Cos}[e + 2 * f*x] - 9 * a * b^2 * c * d * f^2 * x^2 * \text{Cos}[e + 2 * f*x] + a^3 * d^2 * f^2 * x^2 * \text{Cos}[e + 2 * f*x] - 3 * a * b^2 * d^2 * f^2 * x^2 * \text{Cos}[e + 2 * f*x] + 3 * a^3 * c^2 * f^2 * x^2 * \text{Cos}[3 * e + 2 * f*x] - 9 * a * b^2 * c^2 * f^2 * x * \text{Cos}[3 * e + 2 * f*x] + 3 * a^3 * c * d * f^2 * x^2 * \text{Cos}[3 * e + 2 * f*x] - 9 * a * b^2 * c * d * f^2 * x^2 * \text{Cos}[3 * e + 2 * f*x] + a^3 * d^2 * f^2 * x^2 * \text{Cos}[3 * e + 2 * f*x] - 3 * a * b^2 * d^2 * f^2 * x^2 * \text{Cos}[3 * e + 2 * f*x] + 6 * b^3 * c * d * \text{Sin}[e] - 18 * a * b^2 * c^2 * f * \text{Sin}[e] + 6 * b^3 * d^2 * x * \text{Sin}[e] - 36 * a * b^2 * c * d * f * x * \text{Sin}[e] + 18 * a^2 * b * c^2 * f^2 * x * \text{Sin}[e] - 6 * b^3 * c^2 * f^2 * x * \text{Sin}[e] - 18 * a * b^2 * d^2 * f^2 * x^2 * \text{Sin}[e] + 18 * a^2 * b * c * d * f^2 * x^2 * \text{Sin}[e] - 6 * b^3 * c * d * f^2 * x^2 * \text{Sin}[e] + 6 * a^2 * b * d^2 * f^2 * x^2 * \text{Sin}[e] - 2 * b^3 * d^2 * f^2 * x^2 * \text{Sin}[e] - 6 * b^3 * c * d * \text{Sin}[e + 2 * f*x] + 18 * a * b^2 * c^2 * f * \text{Sin}[e + 2 * f*x] - 6 * b^3 * d^2 * x * \text{Sin}[e + 2 * f*x] + 36 * a * b^2 * c * d * f * x * \text{Sin}[e + 2 * f*x] - 9 * a^2 * b * c^2 * f^2 * x * \text{Sin}[e + 2 * f*x] + 3 * b^3 * c^2 * f^2 * x * \text{Sin}[e + 2 * f*x] + 18 * a * b^2 * d^2 * f * x^2 * \text{Sin}[e + 2 * f*x] - 9 * a^2 * b * c * d * f^2 * x^2 * \text{Sin}[e + 2 * f*x] + 3 * b^3 * c * d * f^2 * x^2 * \text{Sin}[e + 2 * f*x] - 3 * a^2 * b * d^2 * f^2 * x^2 * \text{Sin}[e + 2 * f*x] + b^3 * d^2 * f^2 * x^2 * \text{Sin}[e + 2 * f*x] + 9 * a^2 * b * c^2 * f^2 * x * \text{Sin}[3 * e + 2 * f*x] - 3 * b^3 * c^2 * f^2 * x * \text{Sin}[3 * e + 2 * f*x] + 9 * a^2 * b * c * d * f^2 * x^2 * \text{Sin}[3 * e + 2 * f*x] - 3 * b^3 * c * d * f^2 * x^2 * \text{Sin}[3 * e + 2 * f*x] + 3 * a^2 * b * d^2 * f^2 * x^2 * \text{Sin}[3 * e + 2 * f*x] - b^3 * d^2 * f^2 * x^2 * \text{Sin}[3 * e + 2 * f*x])) / (12 * f^2)
\end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1137 vs. $2(402) = 804$.

Time = 1.06 (sec), antiderivative size = 1138, normalized size of antiderivative = 2.61

method	result	size
risch	Expression too large to display	1138

[In] `int((d*x+c)^2*(a+b*tan(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned} & 3*I*d*a^2*b*c*x^2+12*I/f*b*a^2*c*d*e*x+6/f^2*b^2*a*c*d*ln(exp(2*I*(f*x+e))+1)-12/f^2*b^2*a*c*d*ln(exp(I*(f*x+e)))+4/f^2*b^3*c*d*ln(exp(I*(f*x+e)))+6 \\ & /f^2*b^2*a*d^2*ln(exp(2*I*(f*x+e))+1)*x+2/f*b^3*c*d*ln(exp(2*I*(f*x+e))+1)*x+6/f^3*b*e^2*d^2*a^2*ln(exp(I*(f*x+e)))-3/f*b*d^2*a^2*ln(exp(2*I*(f*x+e))+1)*x^2+12/f^3*b^2*c*a*d^2*ln(exp(I*(f*x+e)))-6*I/f^3*b^2*a*d^2*e^2-I/f^2*b^3*d^2*polylog(2,-exp(2*I*(f*x+e)))*x-2*I/f^2*b^3*c*d*e^2-I/f^2*b^3*c*d*polylog(2,-exp(2*I*(f*x+e)))+2*I/f^2*b^3*d^2*e^2*x-6*I/f*b^2*a*d^2*x^2-4*I/f^3*b*d^2*a^2*e^3+1/f*b^3*d^2*ln(exp(2*I*(f*x+e))+1)*x^2-3/f*b*a^2*c^2*ln(exp(2*I*(f*x+e))+1)+6/f*b*a^2*c^2*ln(exp(I*(f*x+e)))-2/f^3*b^3*c^2*d^2*ln(exp(I*(f*x+e)))+4/3*I/f^3*b^3*d^2*c^3-I*d*b^3*c*x^2-3*d*a*b^2*c*x^2+I*d^2*a^2*b*x^3-3*I*a^2*b*c^2*x-I/d*a^2*b*c^3+d*a^3*c*x^2+a^3*c^2*x-d^2*a*b^2*x^3-3*a*b^2*c^2*x-1/d*a*b^2*c^3+I*b^3*c^2*x+1/3*I/d*b^3*c^3+1/3*d^2*a^3*x^3+1/3/d*c^3*a^3-1/3*I*b^3*d^2*x^3+1/f*b^3*c^2*ln(exp(2*I*(f*x+e))+1)-2/f*b^3*c^2*ln(exp(I*(f*x+e)))-1/f^3*b^3*d^2*ln(exp(2*I*(f*x+e))+1)+2/f^3*b^3*d^2*ln(exp(I*(f*x+e)))+2*b^2*(3*I*a*d^2*f*x^2*exp(2*I*(f*x+e))+6*I*a*c*d*f*x*exp(2*I*(f*x+e))+b*d^2*f*x^2*exp(2*I*(f*x+e))+3*I*a*c^2*f*exp(2*I*(f*x+e))+3*I*a*d^2*f*x^2-I*b*d^2*x*exp(2*I*(f*x+e))+2*b*c*d*f*x*exp(2*I*(f*x+e))+6*I*a*c*d*f*x-I*b*c*d*exp(2*I*(f*x+e))+b*c^2*f*exp(2*I*(f*x+e))+3*I*a*c^2*f-I*d^2*x*b-I*c*d*b)/f^2/(exp(2*I*(f*x+e))+1)^2-3*I*a*b^2*d^2*polylog(2,-exp(2*I*(f*x+e)))/f^3-3/2*a^2*b*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3+1/2*b^3*d^2*polylog(3,-exp(2*I*(f*x+e)))/f^3-12/f^2*b*e*a^2*c*d*ln(exp(I*(f*x+e)))-6/f*b*a^2*c*d*ln(exp(2*I*(f*x+e))+1)*x+6*I/f^2*b*a^2*c*d*e^2-12*I/f^2*b^2*a*d^2*e*x+3*I/f^2*b*d^2*a^2*polylog(2,-exp(2*I*(f*x+e)))*x+3*I/f^2*b*a^2*c*d*polylog(2,-exp(2*I*(f*x+e)))-6*I/f^2*b*d^2*a^2*e^2*x-4*I/f*b^3*c*d*e*x \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 688, normalized size of antiderivative = 1.58

$$\begin{aligned} & \int (c + dx)^2(a + b \tan(e + fx))^3 dx \\ & = \frac{4 (a^3 - 3 ab^2) d^2 f^3 x^3 - 3 (3 a^2 b - b^3) d^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 + 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right) - 3 (3 a^2 b - b^3) d^2 \text{polylog}\left(3, \frac{\tan(fx+e)^2 - 2i \tan(fx+e) - 1}{\tan(fx+e)^2 + 1}\right)}{12*(4*(a^3 - 3*a*b^2)*d^2*f^3*x^3 - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^3*d^2*f^2 + 2*(a^3 - 3*a*b^2)*c*d*f^3)*x^2 + 6*(b^3*d^2*f^2*x^2 + 2*b^3*c*d*f^2*x + b^3*c^2*f^2)*tan(f*x + e)^2 + 12*(b^3*c*d*f^2 + (a^3 - 3*a*b^2)*d^2*f^3)*x + 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)))}) \end{aligned}$$

[In] `integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="fricas")`

[Out]

$$\begin{aligned} & 1/12*(4*(a^3 - 3*a*b^2)*d^2*f^3*x^3 - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 6*(b^3*d^2*f^2 + 2*(a^3 - 3*a*b^2)*c*d*f^3)*x^2 + 6*(b^3*d^2*f^2*x^2 + 2*b^3*c*d*f^2*x + b^3*c^2*f^2)*tan(f*x + e)^2 + 12*(b^3*c*d*f^2 + (a^3 - 3*a*b^2)*d^2*f^3)*x + 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 + 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 3*(3*a^2*b - b^3)*d^2*polylog(3, (tan(f*x + e)^2 - 2*I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1))) \end{aligned}$$

$$\begin{aligned}
& 3*a*b^2*c^2*f^3*x - 6*(-3*I*a*b^2*d^2 + I*(3*a^2*b - b^3)*d^2*f*x + I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) \\
& - 6*(3*I*a*b^2*d^2 - I*(3*a^2*b - b^3)*d^2*f*x - I*(3*a^2*b - b^3)*c*d*f)*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) - 6*((3*a^2*b - b^3)*d^2*f^2*x^2 - 6*a*b^2*c*d*f + b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 - 2*(3*a*b^2*d^2*f - (3*a^2*b - b^3)*c*d*f^2)*x)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 12*(3*a*b^2*d^2*f^2*x^2 + 3*a*b^2*c^2*f^2 - b^3*c*d*f + (6*a*b^2*c*d*f^2 - b^3*d^2*f)*x)*tan(f*x + e))/f^3
\end{aligned}$$

Sympy [F]

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx)^2 dx$$

```
[In] integrate((d*x+c)**2*(a+b*tan(f*x+e))**3,x)
[Out] Integral((a + b*tan(e + f*x))**3*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3407 vs. $2(393) = 786$.

Time = 1.97 (sec), antiderivative size = 3407, normalized size of antiderivative = 7.81

$$\int (c + dx)^2 (a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="maxima")
[Out] 1/3*(3*(f*x + e)*a^3*c^2 + (f*x + e)^3*a^3*d^2/f^2 - 3*(f*x + e)^2*a^3*d^2*e/f^2 + 3*(f*x + e)*a^3*d^2*e^2/f^2 + 3*(f*x + e)^2*a^3*c*d/f - 6*(f*x + e)*a^3*c*d*e/f + 9*a^2*b*c^2*log(sec(f*x + e)) + 9*a^2*b*d^2*e^2*log(sec(f*x + e))/f^2 - 18*a^2*b*c*d*e*log(sec(f*x + e))/f + 3*(36*a*b^2*d^2*e^2 + 36*a*b^2*c^2*f^2 + 2*(3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^3*d^2 + 12*b^3*d^2*e - 6*((3*a^2*b + 3*I*a*b^2 - b^3)*d^2*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d*f)*(f*x + e)^2 - 6*((-3*I*a*b^2 + b^3)*d^2*e^2 + 2*(3*I*a*b^2 - b^3)*c*d*e*f + (-3*I*a*b^2 + b^3)*c^2*f^2)*(f*x + e) - 12*(6*a*b^2*c*d*e + b^3*c*d)*f + 6*(b^3*d^2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f + (b^3*d^2*e^2 + b^3*c^2*f^2
```

$$\begin{aligned}
& 2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(4*f*x + 4*e) + 2*(b^3*d^2*2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*cos(2*f*x + 2*e) - (-I*b^3*d^2*2*e^2 - I*b^3*c^2*f^2 + 6*I*a*b^2*d^2*e + (3*I*a^2*b - I*b^3)*(f*x + e)^2*d^2 + I*b^3*d^2 + 2*(-3*I*a*b^2*d^2 + (-3*I*a^2*b + I*b^3)*d^2*e + (3*I*a^2*b - I*b^3)*c*d*f)*(f*x + e) + 2*(I*b^3*c*d*e - 3*I*a*b^2*c*d)*f)*sin(4*f*x + 4*e) - 2*(-I*b^3*d^2*2*e^2 - I*b^3*c^2*f^2 + 6*I*a*b^2*d^2*e + (3*I*a^2*b - I*b^3)*(f*x + e)^2*d^2 + I*b^3*d^2 + 2*(-3*I*a*b^2*d^2 + (-3*I*a^2*b + I*b^3)*d^2*e + (3*I*a^2*b - I*b^3)*c*d*f)*(f*x + e) + 2*(I*b^3*c*d*e - 3*I*a*b^2*c*d)*f)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) + 2*((3*a^2*b + 3*I*a*b^2 - b^3)*d^2*2*e - (3*a^2*b + 3*I*a*b^2 - b^3)*c*d*f)*(f*x + e)^2 + 3*(12*a*b^2*d^2*2*e + 2*b^3*d^2 - (-3*I*a*b^2 + b^3)*d^2*2*e^2 - (-3*I*a*b^2 + b^3)*c^2*f^2 - 2*(6*a*b^2*c*d + (3*I*a*b^2 - b^3)*c*d*e)*f)*(f*x + e))*cos(4*f*x + 4*e) + 4*((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^3*d^2 + 3*b^3*d^2*2*e + 3*(3*a*b^2 - I*b^3)*d^2*2*e^2 + 3*(3*a*b^2 - I*b^3)*c^2*f^2 - 3*((3*a^2*b + 3*I*a*b^2 - b^3)*c*d*f + (3*a*b^2 + I*b^3)*d^2)*(f*x + e)^2 + 3*(b^3*d^2 - (-3*I*a*b^2 + b^3)*d^2*2*e^2 - (-3*I*a*b^2 + b^3)*c^2*f^2 + 2*(3*a*b^2 + I*b^3)*d^2*2*e - 2*((3*I*a*b^2 - b^3)*c*d*e + (3*a*b^2 + I*b^3)*c*d)*f)*(f*x + e) - 3*(b^3*c*d + 2*(3*a*b^2 - I*b^3)*c*d*e)*f)*cos(2*f*x + 2*e) - 6*(3*a*b^2*d^2 - (3*a^2*b - b^3)*(f*x + e)*d^2 + (3*a^2*b - b^3)*d^2*2*e - (3*a^2*b - b^3)*c*d*f + (3*a*b^2*d^2 - (3*a^2*b - b^3)*(f*x + e)*d^2 + (3*a^2*b - b^3)*c*d*f)*cos(4*f*x + 4*e) + 2*(3*a*b^2*d^2 - (3*a^2*b - b^3)*(f*x + e)*d^2 + (3*a^2*b - b^3)*c*d*f - 2*(3*I*a*b^2*d^2 - (3*a^2*b - b^3)*(f*x + e)*d^2 + (3*a^2*b - b^3)*c*d*f)*(f*x + e) + 2*(-I*b^3*c*d*e + 3*I*a*b^2*c*d)*f)*cos(2*f*x + 2*e) + (-3*I*a^2*b + I*b^3)*(f*x + e)*d^2 + (3*I*a^2*b - I*b^3)*d^2*2*e + (-3*I*a^2*b + I*b^3)*c*d*f)*sin(4*f*x + 4*e) + 2*(3*I*a^2*b^2*d^2 - (-3*I*a^2*b + I*b^3)*(f*x + e)*d^2 + (3*I*a^2*b - I*b^3)*d^2*2*e + (-3*I*a^2*b + I*b^3)*c*d*f)*sin(2*f*x + 2*e))*dilog(-e^(2*I*f*x + 2*I*e)) - 3*(I*b^3*d^2*2*e^2 + I*b^3*c^2*f^2 - 6*I*a*b^2*d^2*2*e + (-3*I*a^2*b + I*b^3)*(f*x + e)^2*d^2 - I*b^3*d^2 + 2*(3*I*a*b^2*d^2 - (3*I*a^2*b - I*b^3)*d^2*2*e + (-3*I*a^2*b + I*b^3)*c*d*f)*(f*x + e) + 2*(-I*b^3*c*d*e + 3*I*a*b^2*c*d)*f + (I*b^3*d^2*2*e^2 + I*b^3*c^2*f^2 - 6*I*a*b^2*d^2*2*e + (-3*I*a^2*b + I*b^3)*(f*x + e)^2*d^2 - I*b^3*d^2 + 2*(3*I*a*b^2*d^2 - (3*I*a^2*b - I*b^3)*d^2*2*e + (-3*I*a^2*b + I*b^3)*c*d*f)*(f*x + e) + 2*(-I*b^3*c*d*e + 3*I*a*b^2*c*d)*f)*cos(4*f*x + 4*e) + 2*(I*b^3*d^2*2*e^2 + I*b^3*c^2*f^2 - 6*I*a*b^2*d^2*2*e + (-3*I*a^2*b + I*b^3)*(f*x + e)^2*d^2 - I*b^3*d^2 + 2*(3*I*a*b^2*d^2 - (3*I*a^2*b - I*b^3)*d^2*2*e + (-3*I*a^2*b + I*b^3)*c*d*f)*(f*x + e) + 2*(-I*b^3*c*d*e + 3*I*a*b^2*c*d)*f)*cos(2*f*x + 2*e) - (b^3*d^2*2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 - (3*a^2*b - b^3)*d^2*2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*sin(4*f*x + 4*e) - 2*(b^3*d^2*2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 - (3*a^2*b - b^3)*d^2*2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*sin(4*f*x + 4*e) - 2*(b^3*d^2*2*e^2 + b^3*c^2*f^2 - 6*a*b^2*d^2*2*e - (3*a^2*b - b^3)*(f*x + e)^2*d^2 - b^3*d^2 + 2*(3*a*b^2*d^2 - (3*a^2*b - b^3)*d^2*2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)
\end{aligned}$$

$$\begin{aligned}
& \sim 2 + (3*a^2*b - b^3)*d^2*e - (3*a^2*b - b^3)*c*d*f)*(f*x + e) - 2*(b^3*c*d*e - 3*a*b^2*c*d)*f)*\sin(2*f*x + 2*e))*\log(\cos(2*f*x + 2*e)^2 + \sin(2*f*x + 2*e)^2 + 2*\cos(2*f*x + 2*e) + 1) - 3*((-3*I*a^2*b + I*b^3)*d^2*\cos(4*f*x + 4*e) + 2*(-3*I*a^2*b + I*b^3)*d^2*\cos(2*f*x + 2*e) + (3*a^2*b - b^3)*d^2*\sin(4*f*x + 4*e) + 2*(3*a^2*b - b^3)*d^2*\sin(2*f*x + 2*e) + (-3*I*a^2*b + I*b^3)*d^2)*\text{polylog}(3, -e^(2*I*f*x + 2*I*e)) - 2*((-3*I*a^2*b + 3*a*b^2 + I*b^3)*(f*x + e)^3*d^2 + 3*(6*I*a*b^2*d^2 + (3*I*a^2*b - 3*a*b^2 - I*b^3)*d^2*e + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c*d*f)*(f*x + e)^2 + 3*(-12*I*a*b^2*d^2*e - 2*I*b^3*d^2 + (3*a*b^2 + I*b^3)*c^2*f^2 + 2*(6*I*a*b^2*c*d - (3*a*b^2 + I*b^3)*c*d*e)*f)*(f*x + e))*\sin(4*f*x + 4*e) - 4*((-3*I*a^2*b + 3*a*b^2 + I*b^3)*(f*x + e)^3*d^2 - 3*I*b^3*d^2*e + 3*(-3*I*a*b^2 - b^3)*d^2*e^2 + 3*(-3*I*a*b^2 - b^3)*c^2*f^2 + 3*((3*I*a^2*b - 3*a*b^2 - I*b^3)*d^2*e + (-3*I*a^2*b + 3*a*b^2 + I*b^3)*c*d*f + (3*I*a*b^2 - b^3)*d^2)*(f*x + e)^2 + 3*(-I*b^3*d^2 + (3*a*b^2 + I*b^3)*d^2*e^2 + (3*a*b^2 + I*b^3)*c^2*f^2 + 2*(-3*I*a*b^2 + b^3)*d^2*e - 2*((3*a*b^2 + I*b^3)*c*d*e - (3*I*a*b^2 - b^3)*c*d*f)*(f*x + e) + 3*(I*b^3*c*d + 2*(3*I*a*b^2 + b^3)*c*d*e)*f)*\sin(2*f*x + 2*e))/(-6*I*f^2*\cos(4*f*x + 4*e) - 12*I*f^2*\cos(2*f*x + 2*e) + 6*f^2*\sin(4*f*x + 4*e) + 12*f^2*\sin(2*f*x + 2*e) - 6*I*f^2))/f
\end{aligned}$$

Giac [F]

$$\int (c + dx)^2(a + b\tan(e + fx))^3 dx = \int (dx + c)^2(b\tan(fx + e) + a)^3 dx$$

[In] integrate((d*x+c)^2*(a+b*tan(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tan(f*x + e) + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2(a + b\tan(e + fx))^3 dx = \int (a + b\tan(e + fx))^3(c + dx)^2 dx$$

[In] int((a + b*tan(e + f*x))^3*(c + d*x)^2,x)

[Out] int((a + b*tan(e + f*x))^3*(c + d*x)^2, x)

3.51 $\int (c + dx)(a + b \tan(e + fx))^3 dx$

Optimal result	342
Rubi [A] (verified)	343
Mathematica [A] (verified)	346
Maple [A] (verified)	346
Fricas [A] (verification not implemented)	347
Sympy [F]	347
Maxima [B] (verification not implemented)	348
Giac [F]	349
Mupad [F(-1)]	349

Optimal result

Integrand size = 18, antiderivative size = 277

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^3 dx = & -3ab^2cx + \frac{b^3dx}{2f} - \frac{3}{2}ab^2dx^2 + \frac{a^3(c + dx)^2}{2d} \\ & + \frac{3ia^2b(c + dx)^2}{2d} - \frac{ib^3(c + dx)^2}{2d} \\ & - \frac{3a^2b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^3(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ab^2d \log(\cos(e + fx))}{f^2} \\ & + \frac{3ia^2bd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & - \frac{ib^3d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d \tan(e + fx)}{2f^2} \\ & + \frac{3ab^2(c + dx) \tan(e + fx)}{f} + \frac{b^3(c + dx) \tan^2(e + fx)}{2f} \end{aligned}$$

```
[Out] -3*a*b^2*c*x+1/2*b^3*d*x/f-3/2*a*b^2*d*x^2+1/2*a^3*(d*x+c)^2/d+3/2*I*a^2*b*(d*x+c)^2/d-1/2*I*b^3*(d*x+c)^2/d-3*a^2*b*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f+b^3*(d*x+c)*ln(1+exp(2*I*(f*x+e)))/f+3*a*b^2*d*ln(cos(f*x+e))/f^2+3/2*I*a^2*b*d*polylog(2, -exp(2*I*(f*x+e)))/f^2-1/2*I*b^3*d*polylog(2, -exp(2*I*(f*x+e)))/f^2-1/2*b^3*d*tan(f*x+e)/f^2+3*a*b^2*(d*x+c)*tan(f*x+e)/f+1/2*b^3*(d*x+c)*tan(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3803, 3800, 2221, 2317, 2438, 3801, 3556, 3554, 8}

$$\begin{aligned} \int (c + dx)(a + b \tan(e + fx))^3 dx = & \frac{a^3(c + dx)^2}{2d} - \frac{3a^2b(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{3ia^2b(c + dx)^2}{2d} + \frac{3ia^2bd \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\ & + \frac{3ab^2(c + dx) \tan(e + fx)}{f} - 3ab^2cx \\ & + \frac{3ab^2d \log(\cos(e + fx))}{f^2} - \frac{3}{2}ab^2dx^2 \\ & + \frac{b^3(c + dx) \log(1 + e^{2i(e+fx)})}{f} \\ & + \frac{b^3(c + dx) \tan^2(e + fx)}{2f} - \frac{ib^3(c + dx)^2}{2d} \\ & - \frac{ib^3d \operatorname{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d \tan(e + fx)}{2f^2} + \frac{b^3dx}{2f} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Tan}[e + f*x])^3, x]$

[Out]
$$\begin{aligned} & -3*a*b^2*c*x + (b^3*d*x)/(2*f) - (3*a*b^2*d*x^2)/2 + (a^3*(c + d*x)^2)/(2*d) \\ & + (((3*I)/2)*a^2*b*(c + d*x)^2)/d - ((I/2)*b^3*(c + d*x)^2)/d - (3*a^2*b*(c + d*x)*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (b^3*(c + d*x)*\operatorname{Log}[1 + E^{((2*I)*(e + f*x))}])/f + (3*a*b^2*d*\operatorname{Log}[\operatorname{Cos}[e + f*x]])/f^2 + (((3*I)/2)*a^2*b*d*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - ((I/2)*b^3*d*\operatorname{PolyLog}[2, -E^{((2*I)*(e + f*x))}])/f^2 - (b^3*d*\operatorname{Tan}[e + f*x])/(2*f^2) + (3*a*b^2*(c + d*x)*\operatorname{Tan}[e + f*x])/f + (b^3*(c + d*x)*\operatorname{Tan}[e + f*x]^2)/(2*f) \end{aligned}$$

Rule 8

$\operatorname{Int}[a_, x_{\text{Symbol}}] :> \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 2221

$\operatorname{Int}[(((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_*)})/((a_) + (b_*)*((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}), x_{\text{Symbol}}] :> \operatorname{Simp}[((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - \operatorname{Distr}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^(m - 1)*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \operatorname{IGtQ}[m, 0]$

Rule 2317

```

Int[Log[(a_) + (b_)*(F_)^((e_.)*(c_.) + (d_)*(x_)))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2438

```

Int[Log[(c_)*(d_ + (e_)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 3554

```

Int[((b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

```

Rule 3556

```

Int[tan[(c_.) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

```

Rule 3800

```

Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x))/(1 + E^(2*I*(e + f*x)))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

```

Rule 3801

```

Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

```

Rule 3803

```

Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

```

Rubi steps

integral =
$$\int (a^3(c + dx) + 3a^2b(c + dx)\tan(e + fx) + 3ab^2(c + dx)\tan^2(e + fx) + b^3(c + dx)\tan^3(e + fx)) \, dx$$

$$\begin{aligned}
&= \frac{a^3(c+dx)^2}{2d} + (3a^2b) \int (c+dx) \tan(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx) \tan^2(e+fx) dx + b^3 \int (c+dx) \tan^3(e+fx) dx \\
&= \frac{a^3(c+dx)^2}{2d} + \frac{3ia^2b(c+dx)^2}{2d} + \frac{3ab^2(c+dx) \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx) \tan^2(e+fx)}{2f} - (6ia^2b) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} dx - (3ab^2) \int (c+dx) dx \\
&\quad - b^3 \int (c+dx) \tan(e+fx) dx - \frac{(3ab^2d) \int \tan(e+fx) dx}{f} - \frac{(b^3d) \int \tan^2(e+fx) dx}{2f} \\
&= -3ab^2cx - \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} + \frac{3ia^2b(c+dx)^2}{2d} - \frac{ib^3(c+dx)^2}{2d} \\
&\quad - \frac{3a^2b(c+dx) \log(1+e^{2i(e+fx)})}{f} + \frac{3ab^2d \log(\cos(e+fx))}{f^2} \\
&\quad - \frac{b^3d \tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx) \tan(e+fx)}{f} + \frac{b^3(c+dx) \tan^2(e+fx)}{2f} \\
&\quad + (2ib^3) \int \frac{e^{2i(e+fx)}(c+dx)}{1+e^{2i(e+fx)}} dx + \frac{(3a^2bd) \int \log(1+e^{2i(e+fx)}) dx}{f} + \frac{(b^3d) \int 1 dx}{2f} \\
&= -3ab^2cx + \frac{b^3dx}{2f} - \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} + \frac{3ia^2b(c+dx)^2}{2d} \\
&\quad - \frac{ib^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx) \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx) \log(1+e^{2i(e+fx)})}{f} + \frac{3ab^2d \log(\cos(e+fx))}{f^2} \\
&\quad - \frac{b^3d \tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx) \tan(e+fx)}{f} + \frac{b^3(c+dx) \tan^2(e+fx)}{2f} \\
&\quad - \frac{(3ia^2bd) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^2} - \frac{(b^3d) \int \log(1+e^{2i(e+fx)}) dx}{f} \\
&= -3ab^2cx + \frac{b^3dx}{2f} - \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} + \frac{3ia^2b(c+dx)^2}{2d} \\
&\quad - \frac{ib^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx) \log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx) \log(1+e^{2i(e+fx)})}{f} + \frac{3ab^2d \log(\cos(e+fx))}{f^2} \\
&\quad + \frac{3ia^2bd \text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{b^3d \tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx) \tan(e+fx)}{f} \\
&\quad + \frac{b^3(c+dx) \tan^2(e+fx)}{2f} + \frac{(ib^3d) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i(e+fx)}\right)}{2f^2}
\end{aligned}$$

$$\begin{aligned}
&= -3ab^2cx + \frac{b^3dx}{2f} - \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} + \frac{3ia^2b(c+dx)^2}{2d} \\
&\quad - \frac{ib^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)\log(1+e^{2i(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)\log(1+e^{2i(e+fx)})}{f} + \frac{3ab^2d\log(\cos(e+fx))}{f^2} \\
&\quad + \frac{3ia^2bd\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} - \frac{ib^3d\text{PolyLog}(2, -e^{2i(e+fx)})}{2f^2} \\
&\quad - \frac{b^3d\tan(e+fx)}{2f^2} + \frac{3ab^2(c+dx)\tan(e+fx)}{f} + \frac{b^3(c+dx)\tan^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 8.32 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.00

$$\begin{aligned}
&\int (c+dx)(a+b\tan(e+fx))^3 dx \\
&= \cos(e+fx) (\cos^2(e+fx) ((-((e+fx)(-3ia^2bd(e+fx)+ib^3d(e+fx)+3ab^2(-de+2cf+dfx)+a^3(-
\end{aligned}$$

[In] `Integrate[(c + d*x)*(a + b*Tan[e + f*x])^3, x]`

[Out] `(Cos[e + f*x]*(Cos[e + f*x]^2*(-((e + f*x)*((-3*I)*a^2*b*d*(e + f*x) + I*b^3*d*(e + f*x) + 3*a*b^2*(-(d*e) + 2*c*f + d*f*x) + a^3*(-2*c*f + d*(e - f*x)))) + 2*b*(-3*a^2 + b^2)*d*(e + f*x)*Log[1 + E^((2*I)*(e + f*x))] + 2*b*(3*a*b*d + 3*a^2*(d*e - c*f) + b^2*(-(d*e) + c*f))*Log[Cos[e + f*x]]) - I*b*(-3*a^2 + b^2)*d*Cos[e + f*x]^2*PolyLog[2, -E^((2*I)*(e + f*x))] + (b^2*(2*b*f*(c + d*x) + (-b*d) + 6*a*f*(c + d*x))*Sin[2*(e + f*x)]))/2)*(a + b*Tan[e + f*x])^3)/(2*f^2*(a*Cos[e + f*x] + b*Sin[e + f*x])^3)`

Maple [A] (verified)

Time = 0.92 (sec) , antiderivative size = 493, normalized size of antiderivative = 1.78

method	result
risch	$-3ia^2bcx + \frac{b^3d\ln(e^{2i(fx+e)}+1)x}{f} + \frac{3b^2ad\ln(e^{2i(fx+e)}+1)}{f^2} - \frac{6b^2ad\ln(e^{i(fx+e)})}{f^2} - \frac{3ba^2c\ln(e^{2i(fx+e)}+1)}{f} + \frac{6ba^2c\ln(e^{i(fx+e)})}{f}$

[In] `int((d*x+c)*(a+b*tan(f*x+e))^3, x, method=_RETURNVERBOSE)`

[Out] `1/2*a^3*d*x^2+a^3*c*x+I*b^3*c*x-3*I*a^2*b*c*x+1/f*b^3*c*ln(exp(2*I*(f*x+e))+1)-2/f*b^3*c*ln(exp(I*(f*x+e)))-1/2*I*b^3*d*x^2+1/f*b^3*d*ln(exp(2*I*(f*x+e))+1)*x+3/f^2*b^2*a*d*ln(exp(2*I*(f*x+e))+1)-6/f^2*b^2*a*d*ln(exp(I*(f*x+e)))-3/f*b*a^2*c*ln(exp(2*I*(f*x+e))+1)+6/f*b*a^2*c*ln(exp(I*(f*x+e)))+2/f^2`

$$\begin{aligned} & *b^3 * e * d * \ln(\exp(I * (f * x + e))) - I / f^2 * b^3 * d * e^2 + 3/2 * I * a^2 * b * d * x^2 + b^2 * (6 * I * a * d * \\ & f * x * \exp(2 * I * (f * x + e)) + 6 * I * a * c * f * \exp(2 * I * (f * x + e)) + 2 * b * d * f * x * \exp(2 * I * (f * x + e)) + \\ & 6 * I * a * d * f * x - I * b * d * \exp(2 * I * (f * x + e)) + 2 * b * c * f * \exp(2 * I * (f * x + e)) + 6 * I * a * c * f - I * b * d \\ &) / f^2 / (\exp(2 * I * (f * x + e)) + 1)^2 + 3/2 * I * a^2 * b * d * \text{polylog}(2, -\exp(2 * I * (f * x + e))) / f^2 \\ & - 1/2 * I * b^3 * d * \text{polylog}(2, -\exp(2 * I * (f * x + e))) / f^2 - 3 * a * b^2 * c * x - 3/2 * a * b^2 * d * x^2 - 6 \\ & / f^2 * b * e * a^2 * d * \ln(\exp(I * (f * x + e))) - 3 / f * b * a^2 * d * \ln(\exp(2 * I * (f * x + e)) + 1) * x - 2 * I / \\ & f * b^3 * d * e * x + 3 * I / f^2 * b * a^2 * d * e^2 + 6 * I / f * b * a^2 * d * e * x \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec), antiderivative size = 327, normalized size of antiderivative = 1.18

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \frac{2(a^3 - 3ab^2)df^2x^2 - i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) + i(3a^2b - b^3)d\text{Li}_2\left(\frac{2(-i \tan(fx+e)-1)}{\tan(fx+e)^2+1} + 1\right) +}{}$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="fricas")
[Out] 1/4*(2*(a^3 - 3*a*b^2)*d*f^2*x^2 - I*(3*a^2*b - b^3)*d*dilog(2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + I*(3*a^2*b - b^3)*d*dilog(2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1) + 1) + 2*(b^3*d*f*x + b^3*c*f)*tan(f*x + e)^2 + 2*(b^3*d*f + 2*(a^3 - 3*a*b^2)*c*f^2)*x + 2*(3*a*b^2*d - (3*a^2*b - b^3)*d*f*x - (3*a^2*b - b^3)*c*f)*log(-2*(I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(3*a*b^2*d - (3*a^2*b - b^3)*d*f*x - (3*a^2*b - b^3)*c*f)*log(-2*(-I*tan(f*x + e) - 1)/(tan(f*x + e)^2 + 1)) + 2*(6*a*b^2*d*f*x + 6*a*b^2*c*f - b^3*d)*tan(f*x + e))/f^2
```

Sympy [F]

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))**3,x)
[Out] Integral((a + b*tan(e + f*x))**3*(c + d*x), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1319 vs. $2(241) = 482$.

Time = 0.67 (sec) , antiderivative size = 1319, normalized size of antiderivative = 4.76

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="maxima")
[Out] 1/2*(2*(f*x + e)*a^3*c + (f*x + e)^2*a^3*d/f - 2*(f*x + e)*a^3*d*e/f + 6*a^2*b*c*log(sec(f*x + e)) - 6*a^2*b*d*e*log(sec(f*x + e))/f - 2*(12*a*b^2*d*e - 12*a*b^2*c*f - (3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d + 2*b^3*d + 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d + (b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) + 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*cos(2*f*x + 2*e) + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(4*f*x + 4*e) + 2*(I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*sin(2*f*x + 2*e))*arctan2(sin(2*f*x + 2*e), cos(2*f*x + 2*e) + 1) - ((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - 2*(6*a*b^2*d + (3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f)*(f*x + e))*cos(4*f*x + 4*e) - 2*((3*a^2*b + 3*I*a*b^2 - b^3)*(f*x + e)^2*d - b^3*d - 2*(3*a*b^2 - I*b^3)*d*e + 2*(3*a*b^2 - I*b^3)*c*f - 2*((3*I*a*b^2 - b^3)*d*e + (-3*I*a*b^2 + b^3)*c*f + (3*a*b^2 + I*b^3)*d)*(f*x + e))*cos(2*f*x + 2*e) - ((3*a^2*b - b^3)*d*cos(4*f*x + 4*e) + 2*(3*a^2*b - b^3)*d*cos(2*f*x + 2*e) + (3*I*a^2*b - I*b^3)*d*sin(4*f*x + 4*e) - 2*(-3*I*a^2*b + I*b^3)*d*sin(2*f*x + 2*e) + (3*a^2*b - b^3)*d) *dilog(-e^(2*I*f*x + 2*I*e)) - (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d + (I*b^3*d*e - I*b^3*c*f - 3*I*a*b^2*d + (3*I*a^2*b - I*b^3)*(f*x + e)*d)*cos(4*f*x + 4*e) - 2*(-I*b^3*d*e + I*b^3*c*f + 3*I*a*b^2*d + (-3*I*a^2*b + I*b^3)*(f*x + e)*d)*cos(2*f*x + 2*e) - (b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*sin(4*f*x + 4*e) - 2*(b^3*d*e - b^3*c*f - 3*a*b^2*d + (3*a^2*b - b^3)*(f*x + e)*d)*sin(2*f*x + 2*e))*log(cos(2*f*x + 2*e)^2 + sin(2*f*x + 2*e)^2 + 2*cos(2*f*x + 2*e) + 1) - ((3*I*a^2*b - 3*a*b^2 - I*b^3)*(f*x + e)^2*d - 2*(6*I*a*b^2*d - (3*a*b^2 + I*b^3)*d*e + (3*a*b^2 + I*b^3)*c*f)*(f*x + e))*sin(4*f*x + 4*e) + 2*((-3*I*a^2*b + 3*a*b^2 + I*b^3)*(f*x + e)^2*d + I*b^3*d + 2*(3*I*a*b^2 + b^3)*d*e + 2*(-3*I*a*b^2 - b^3)*c*f - 2*((3*a*b^2 + I*b^3)*d*e - (3*a*b^2 + I*b^3)*c*f - (3*I*a*b^2 - b^3)*d)*(f*x + e))*sin(2*f*x + 2*e)/(-2*I*f*cos(4*f*x + 4*e) - 4*I*f*cos(2*f*x + 2*e) + 2*f*sin(4*f*x + 4*e) + 4*f*sin(2*f*x + 2*e) - 2*I*f)/f
```

Giac [F]

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (dx + c)(b \tan(fx + e) + a)^3 dx$$

[In] `integrate((d*x+c)*(a+b*tan(f*x+e))^3,x, algorithm="giac")`

[Out] `integrate((d*x + c)*(b*tan(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tan(e + fx))^3 dx = \int (a + b \tan(e + fx))^3 (c + dx) dx$$

[In] `int((a + b*tan(e + f*x))^3*(c + d*x),x)`

[Out] `int((a + b*tan(e + f*x))^3*(c + d*x), x)`

3.52 $\int \frac{(a+b\tan(e+fx))^3}{c+dx} dx$

Optimal result	350
Rubi [N/A]	350
Mathematica [N/A]	351
Maple [N/A] (verified)	351
Fricas [N/A]	351
Sympy [N/A]	352
Maxima [N/A]	352
Giac [N/A]	353
Mupad [N/A]	353

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^3}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))^3/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

[In] Int[(a + b*Tan[e + f*x])^3/(c + d*x),x]

[Out] Defer[Int][(a + b*Tan[e + f*x])^3/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 14.69 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

[In] `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

[Out] `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.62 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{dx + c} dx$$

[In] `int((a+b*tan(f*x+e))^3/(d*x+c), x)`

[Out] `int((a+b*tan(f*x+e))^3/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

[In] `integrate((a+b*tan(f*x+e))^3/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b^3*tan(f*x + e)^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 1.33 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tan(e + fx))^3}{c + dx} dx$$

```
[In] integrate((a+b*tan(f*x+e))**3/(d*x+c),x)
[Out] Integral((a + b*tan(e + f*x))**3/(c + d*x), x)
```

Maxima [N/A]

Not integrable

Time = 3.08 (sec) , antiderivative size = 1935, normalized size of antiderivative = 96.75

$$\int \frac{(a + b \tan(e + fx))^3}{c + dx} dx = \int \frac{(b \tan(fx + e) + a)^3}{dx + c} dx$$

```
[In] integrate((a+b*tan(f*x+e))**3/(d*x+c),x, algorithm="maxima")
[Out] (((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*cos(4*f*x + 4*e)^2*log(d*x + c) + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)*sin(4*f*x + 4*e)^2 + 4*(b^3*d^2*f*x + b^3*c*d*f + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*cos(2*f*x + 2*e)^2 + 4*(b^3*d^2*f*x + b^3*c*d*f + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c))*sin(2*f*x + 2*e)^2 + (2*(b^3*d^2*f*x + b^3*c*d*f + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)))*cos(2*f*x + 2*e) + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)*cos(2*f*x + 2*e) + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)*sin(2*f*x + 2*e) - (6*a*b^2*d^2*f*x + 6*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))*cos(4*f*x + 4*e) + 2*(b^3*d^2*f*x + b^3*c*d*f + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*log(d*x + c)))*cos(2*f*x + 2*e) - (d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(2*f*x + 2*e)^2 + (d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(4*f*x + 4*e)^2 + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*sin(2*f*x + 2*e)^2 + 2*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(2*f*x + 2*e))*cos(4*f*x + 4*e) + 4*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*cos(2*f*x + 2*e)*cos(4*f*x + 4*e) - 2*((3*a^2*b - b^3)*d^2*f^2*x^2 + 3*a*b^2*c*d*f)*integrate(-2*((3*a^2*b - b^3)*d^2*f^2*x^2 + 3*a*b^2*c*d*f), x)
```

$$\begin{aligned}
& + b^3 d^2 + (3 a^2 b - b^3) c^2 f^2 + (3 a b^2 d^2 f + 2 (3 a^2 b - b^3) c \\
& * d f^2) * x) * \sin(2 f x + 2 e) / (d^3 f^2 x^3 + 3 c d^2 f^2 x^2 + 3 c^2 d f^2 x \\
& + c^3 f^2 + (d^3 f^2 x^3 + 3 c d^2 f^2 x^2 + 3 c^2 d f^2 x + c^3 f^2) * \cos(2 \\
& * f x + 2 e)^2 + (d^3 f^2 x^3 + 3 c d^2 f^2 x^2 + 3 c^2 d f^2 x + c^3 f^2) * \sin \\
& (2 f x + 2 e)^2 + 2 (d^3 f^2 x^3 + 3 c d^2 f^2 x^2 + 3 c^2 d f^2 x + c^3 f^2) * \cos \\
& (2 f x + 2 e)), x) + ((a^3 - 3 a b^2) d^2 f^2 x^2 + 2 (a^3 - 3 a b^2) \\
& * c d f^2 x + (a^3 - 3 a b^2) c^2 f^2) * \log(d x + c) + (6 a b^2 d^2 f^2 x + 6 \\
& * a b^2 c d f + b^3 d^2 + (6 a b^2 d^2 f^2 x + 6 a b^2 c d f + b^3 d^2) * \cos \\
& (2 f x + 2 e) + 2 (b^3 d^2 f^2 x + b^3 c d f + 2 ((a^3 - 3 a b^2) d^2 f^2 x^2 + \\
& 2 (a^3 - 3 a b^2) c d f^2 x + (a^3 - 3 a b^2) c^2 f^2) * \log(d x + c)) * \sin \\
& (2 f x + 2 e)) * \sin(4 f x + 4 e) + (6 a b^2 d^2 f^2 x + 6 a b^2 c d f + b^3 d^2) * \\
& \sin(2 f x + 2 e)) / (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d f^2 + (d^3 f^2 x^2 + \\
& 2 c d^2 f^2 x + c^2 d f^2) * \cos(4 f x + 4 e)^2 + 4 (d^3 f^2 x^2 + 2 c d^2 f^2 \\
& x^2 + c^2 d f^2) * \cos(2 f x + 2 e)^2 + (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d \\
& * f^2) * \sin(4 f x + 4 e)^2 + 4 (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d f^2) * \sin \\
& (4 f x + 4 e) * \sin(2 f x + 2 e) + 4 (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d f^2) * \\
& \sin(2 f x + 2 e)^2 + 2 (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d f^2) * \cos(2 f x + 2 e)) * \cos \\
& (4 f x + 4 e) + 4 (d^3 f^2 x^2 + 2 c d^2 f^2 x + c^2 d f^2) * \cos(2 f x + 2 e))
\end{aligned}$$

Giac [N/A]

Not integrable

Time = 1.08 (sec), antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^3}{c + d x} dx = \int \frac{(b \tan(f x + e) + a)^3}{d x + c} dx$$

[In] integrate((a+b*tan(f*x+e))^3/(d*x+c),x, algorithm="giac")

[Out] integrate((b*tan(f*x + e) + a)^3/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 3.91 (sec), antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^3}{c + d x} dx = \int \frac{(a + b \tan(e + f x))^3}{c + d x} dx$$

[In] int((a + b*tan(e + f*x))^3/(c + d*x),x)

[Out] int((a + b*tan(e + f*x))^3/(c + d*x), x)

3.53 $\int \frac{(a+b \tan(e+fx))^3}{(c+dx)^2} dx$

Optimal result	354
Rubi [N/A]	354
Mathematica [N/A]	355
Maple [N/A] (verified)	355
Fricas [N/A]	355
Sympy [N/A]	356
Maxima [N/A]	356
Giac [N/A]	357
Mupad [N/A]	358

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tan(e + fx))^3}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tan(f*x+e))^3/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

[In] Int[(a + b*Tan[e + f*x])^3/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 17.93 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

[In] `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]`

[Out] `Integrate[(a + b*Tan[e + f*x])^3/(c + d*x)^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tan(fx + e))^3}{(dx + c)^2} dx$$

[In] `int((a+b*tan(f*x+e))^3/(d*x+c)^2, x)`

[Out] `int((a+b*tan(f*x+e))^3/(d*x+c)^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] `integrate((a+b*tan(f*x+e))^3/(d*x+c)^2, x, algorithm="fricas")`

[Out] `integral((b^3*tan(f*x + e)^3 + 3*a*b^2*tan(f*x + e)^2 + 3*a^2*b*tan(f*x + e) + a^3)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.44 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))**3/(d*x+c)**2,x)
[Out] Integral((a + b*tan(e + f*x))**3/(c + d*x)**2, x)
```

Maxima [N/A]

Not integrable

Time = 9.08 (sec) , antiderivative size = 2194, normalized size of antiderivative = 109.70

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

```
[In] integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="maxima")
[Out] -((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*cos(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e)^2 + ((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2)*sin(4*f*x + 4*e)^2 + 4*((a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + (a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 2*(a^3 - 3*a*b^2)*c*d*f^2)*x)*sin(2*f*x + 2*e)^2 + 2*((a^3 - 3*a*b^2)*d^2*f^2*x^2 + 2*(a^3 - 3*a*b^2)*c*d*f^2*x + (a^3 - 3*a*b^2)*c^2*f^2 + (2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (3*a*b^2*d^2*f^2*x + 3*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))*cos(4*f*x + 4*e) + 2*(2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*cos(2*f*x + 2*e) + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2 + d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(2*f*x + 2*e)^2 + 2*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)^2 + 2*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)
```

```

x + 2*e))*cos(4*f*x + 4*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + c^3*d*f^2)*cos(2*f*x + 2*e))*integrate(-2*((3*a^2*b - b^3)*d^2*f^2*x^2 + 6*a*b^2*c*d*f + 3*b^3*d^2 + (3*a^2*b - b^3)*c^2*f^2 + 2*(3*a*b^2*d^2*f^2 + (3*a^2*b - b^3)*c*d*f^2)*x)*sin(2*f*x + 2*e)/(d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2 + (d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2)*cos(2*f*x + 2*e))^2 + (d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2)*sin(2*f*x + 2*e)^2 + 2*(d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2)*cos(2*f*x + 2*e)), x) - 2*(3*a*b^2*d^2*f*x + 3*a*b^2*c*d*f + b^3*d^2 + (3*a*b^2*d^2*f*x + 3*a*b^2*c*d*f + b^3*d^2)*cos(2*f*x + 2*e) - (2*(a^3 - 3*a*b^2)*d^2*f^2*x^2 - b^3*c*d*f + 2*(a^3 - 3*a*b^2)*c^2*f^2 - (b^3*d^2*f - 4*(a^3 - 3*a*b^2)*c*d*f^2)*x)*sin(2*f*x + 2*e))*sin(4*f*x + 4*e) - 2*(3*a*b^2*d^2*f*x + 3*a*b^2*c*d*f + b^3*d^2)*sin(2*f*x + 2*e))/(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + c^3*d*f^2)*cos(4*f*x + 4*e)^2 + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)^2 + (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(4*f*x + 4*e)*sin(2*f*x + 2*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*sin(2*f*x + 2*e)^2 + 2*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e))*cos(4*f*x + 4*e) + 4*(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2)*cos(2*f*x + 2*e)))

```

Giac [N/A]

Not integrable

Time = 18.06 (sec), antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tan(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] integrate((a+b*tan(f*x+e))^3/(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*tan(f*x + e) + a)^3/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 4.93 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tan(e + f x))^3}{(c + d x)^2} dx = \int \frac{(a + b \tan(e + f x))^3}{(c + d x)^2} dx$$

[In] int((a + b*tan(e + f*x))^3/(c + d*x)^2,x)

[Out] int((a + b*tan(e + f*x))^3/(c + d*x)^2, x)

3.54 $\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$

Optimal result	359
Rubi [A] (verified)	360
Mathematica [A] (verified)	363
Maple [B] (verified)	363
Fricas [B] (verification not implemented)	364
Sympy [F]	365
Maxima [B] (verification not implemented)	365
Giac [F]	366
Mupad [F(-1)]	366

Optimal result

Integrand size = 20, antiderivative size = 243

$$\begin{aligned} \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx &= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\ &\quad - \frac{3ibd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2} \\ &\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3} \\ &\quad + \frac{3ibd^3 \operatorname{PolyLog}\left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{4(a^2+b^2)f^4} \end{aligned}$$

```
[Out] 1/4*(d*x+c)^4/(a+I*b)/d+b*(d*x+c)^3*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-3/2*I*b*d*(d*x+c)^2*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+3/2*b*d^2*(d*x+c)*polylog(3,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3+3/4*I*b*d^3*polylog(4,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^4
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 243, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.300, Rules used = {3813, 2221, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx &= \frac{3bd^2(c+dx)\text{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f^3(a^2+b^2)} \\ &\quad - \frac{3ibd(c+dx)^2\text{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f^2(a^2+b^2)} \\ &\quad + \frac{b(c+dx)^3\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f(a^2+b^2)} \\ &\quad + \frac{3ibd^3\text{PolyLog}\left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{4f^4(a^2+b^2)} + \frac{(c+dx)^4}{4d(a+ib)} \end{aligned}$$

[In] `Int[(c + d*x)^3/(a + b*Tan[e + f*x]), x]`

[Out] $(c + d*x)^4/(4*(a + I*b)*d) + (b*(c + d*x)^3*\text{Log}[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (((3*I)/2)*b*d*(c + d*x)^2*\text{PolyLog}[2, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/((a^2 + b^2)*f^2) + (3*b*d^2*(c + d*x)*\text{PolyLog}[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/(2*(a^2 + b^2)*f^3) + (((3*I)/4)*b*d^3*\text{PolyLog}[4, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)])/((a^2 + b^2)*f^4)$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*(c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_.))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*F^(c*(a +
```

```
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3813

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^p_]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^p_], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^4}{4(a+ib)d} + (2ib) \int \frac{e^{2i(e+fx)}(c+dx)^3}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx \\
&= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\
&\quad - \frac{(3bd) \int (c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) dx}{(a^2+b^2)f} \\
&= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\
&\quad - \frac{3ibd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2} \\
&\quad + \frac{(3ibd^2) \int (c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) dx}{(a^2+b^2)f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} \\
&\quad - \frac{3ibd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^3} \\
&\quad - \frac{(3bd^3) \int \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{2(a^2+b^2)f^3} \\
&= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} \\
&\quad - \frac{3ibd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^3} \\
&\quad + \frac{(3ibd^3) \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)x}{(a+ib)^2} \right)}{x} dx, x, e^{2i(e+fx)} \right)}{4(a^2+b^2)f^4} \\
&= \frac{(c+dx)^4}{4(a+ib)d} + \frac{b(c+dx)^3 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} \\
&\quad - \frac{3ibd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^3} + \frac{3ibd^3 \operatorname{PolyLog} \left(4, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{4(a^2+b^2)f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.68 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.22

$$\begin{aligned} & \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx \\ &= \frac{1}{4}b \left(-\frac{2(c+dx)^4}{(ia+b)d(-ib(-1+e^{2ie})+a(1+e^{2ie}))} + \frac{4(c+dx)^3 \log\left(1+\frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2+b^2)f} \right. \\ & \quad \left. + \frac{3d\left(2if^2(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right) + d\left(2f(c+dx) \operatorname{PolyLog}\left(3, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right) - id \operatorname{PolyLog}\left(4, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)\right)}{(a^2+b^2)f^4} \right) \\ & \quad + \frac{x(4c^3 + 6c^2dx + 4cd^2x^2 + d^3x^3) \cos(e)}{4(a \cos(e) + b \sin(e))} \end{aligned}$$

```
[In] Integrate[(c + d*x)^3/(a + b*Tan[e + f*x]), x]
[Out] (b*((-2*(c + d*x)^4)/((I*a + b)*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))) + (4*(c + d*x)^3*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f) + (3*d*((2*I)*f^2*(c + d*x)^2*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]) + d*(2*f*(c + d*x)*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]) - I*d*PolyLog[4, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x))))]))/((a^2 + b^2)*f^4))/4 + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cos[e])/((4*(a*Cos[e] + b*Sin[e])))
```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1467 vs. 2(220) = 440.

Time = 0.78 (sec) , antiderivative size = 1468, normalized size of antiderivative = 6.04

method	result	size
risch	Expression too large to display	1468

```
[In] int((d*x+c)^3/(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)
[Out] 3*I/f^3/(I*a+b)*b/(-I*b-a)*e^2*c*d^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) - 3*I/f/(I*a+b)*b/(-I*b-a)*d*c^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x - 3*I/f^2/(I*a+b)*b/(-I*b-a)*d*c^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*e - 3*I/f^2/(I*a+b)*b*e*d*c^2/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))) - I*b-a - 6*I/f^3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(exp(I*(f*x+e)))+3*I/f^3/(I*a+b)*b*e^2*c*d^2/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a) - 1/2/(I*a+b)*b/(-I*b-a)*d^3*x^4 - 3/2/f^4/(I*a+b)*b/(-I*b-a)*d^3*e^4 + 3/4/f^4/(I*a+b)*b/(-I*b-a)*d^3*polylog(4, (a-I*b)*exp(2*I*(f*x+e))/(-I*b-a)) - 3/(I*a+b)
```

$$\begin{aligned}
& *b/(-I*b-a)*d*c^2*x^2-2/(I*a+b)*b/(-I*b-a)*d^2*c*x^3-3*I/f/(I*a+b)*b/(-I*b-a)*d^2*c*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x^2+6*I/f^2/(I*a+b)*b*e*d*c^2/(a+I*b)*ln(exp(I*(f*x+e)))-1/4/d/(I*b-a)*c^4-d^2/(I*b-a)*c*x^3-3/2*d/(I*b-a)*c^2*x^2-1/4*d^3/(I*b-a)*x^4-1/(I*b-a)*c^3*x^6/f^2/(I*a+b)*b/(-I*b-a)*e^2*c*d^2*x^3/f^2/(I*a+b)*b/(-I*b-a)*d^2*c*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x-6/f/(I*a+b)*b/(-I*b-a)*d*c^2*e*x-I/f/(I*a+b)*b/(-I*b-a)*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x^3-I/f^4/(I*a+b)*b/(-I*b-a)*e^3*d^3*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-3/2*I/f^3/(I*a+b)*b/(-I*b-a)*d^3*polylog(3,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x-3/2*I/f^3/(I*a+b)*b/(-I*b-a)*d^2*c*polylog(3,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))+2*I/f^4/(I*a+b)*b*e^3*d^3/(a+I*b)*ln(exp(I*(f*x+e)))-I/f^4/(I*a+b)*b*e^3*d^3/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)-2/f^3/(I*a+b)*b/(-I*b-a)*d^3*e^3*x-3/f^2/(I*a+b)*b/(-I*b-a)*d*c^2*e^2-3/2*f^2/(I*a+b)*b/(-I*b-a)*d*c^2*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))+4/f^3/(I*a+b)*b/(-I*b-a)*e^3*c*d^2-3/2*f^2/(I*a+b)*b/(-I*b-a)*d^3*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x^2+I/f/(I*a+b)*b*c^3/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)-2*I/f/(I*a+b)*b*c^3/(a+I*b)*ln(exp(I*(f*x+e)))
\end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1183 vs. $2(212) = 424$.

Time = 0.28 (sec) , antiderivative size = 1183, normalized size of antiderivative = 4.87

$$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="fricas")
[Out] 1/8*(2*a*d^3*f^4*x^4 + 8*a*c*d^2*f^4*x^3 + 12*a*c^2*d*f^4*x^2 + 8*a*c^3*f^4*x - 3*I*b*d^3*polylog(4, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e))^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 3*I*b*d^3*polylog(4, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 6*(-I*b*d^3*f^2*x^2 - 2*I*b*c*d^2*f^2*x^2 - I*b*c^2*d*f^2*x^2 - I*b*c^2*d*f^2*x^2)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2 + 1)) - 6*(I*b*d^3*f^2*x^2 + 2*I*b*c*d^2*f^2*x^2 + I*b*c^2*d*f^2*x^2)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2 + 1)) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I
```

$$\begin{aligned}
& *a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + \\
& a^2 + b^2)) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3) \\
& *log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/ \\
& (tan(f*x + e)^2 + 1)) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - \\
& b*c^3*f^3)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/ \\
& (tan(f*x + e)^2 + 1)) + 6*(b*d^3*f*x + b*c*d^2*f)*polylog(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/ \\
& ((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^3*f*x + b*c*d^2*f)*polylog(3, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/ \\
& ((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)))/((a^2 + b^2)*f^4)
\end{aligned}$$

Sympy [F]

$$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx = \int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx$$

[In] `integrate((d*x+c)**3/(a+b*tan(f*x+e)),x)`
[Out] `Integral((c + d*x)**3/(a + b*tan(e + f*x)), x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 983 vs. $2(212) = 424$.

Time = 0.57 (sec), antiderivative size = 983, normalized size of antiderivative = 4.05

$$\int \frac{(c+dx)^3}{a+b\tan(e+fx)} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="maxima")`
[Out]
$$\begin{aligned}
& -1/12*(18*c^2*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e) + a)/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 6*(2*(f*x + e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(tan(f*x + e)^2 + 1)/(a^2 + b^2))*c^3 - (3*(f*x + e)^4*(a - I*b)*d^3 + 12*I*b*d^3*polylog(4, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b))) - 12*((a - I*b)*d^3*(a - I*b)*c^3 - (a - I*b)*c*d^2*f)*(f*x + e)^3 + 18*((a - I*b)*d^3*c^2 - 2*(a - I*b)*c*d^2*f + (a - I*b)*c^2*d*f^2)*(f*x + e)^2 - 12*((a - I*b)*d^3*c^3 - 3*(a - I*b)*c*d^2*e^2*f)*(f*x + e) - 12*(I*b*d^3*c^3 - 3*I*b*c*d^2*e^2*f)*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) - 4*(4*I*(f*x + e)^3*b*d^3 + 9*(-I*b*d^3*c^2 + I*b*c*d^2*f)*(f*x + e)^2 + 9*(I*b*d^3*c^2 - 2*I*b*c*d^2*f + I*b*c^2*d*f^2)*(f*x + e))
\end{aligned}$$

```

arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2)
, (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2
+ b^2)) - 6*(4*I*(f*x + e)^2*b*d^3 + 3*I*b*d^3*e^2 - 6*I*b*c*d^2*e*f + 3*I*
b*c^2*d*f^2 + 6*(-I*b*d^3*e + I*b*c*d^2*f)*(f*x + e))*dilog((I*a + b)*e^(2*
I*f*x + 2*I*e)/(-I*a + b)) - 6*(b*d^3*e^3 - 3*b*c*d^2*e^2*f)*log((a^2 + b^2)
)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e
)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*(4*(f*x + e)^3*b*d^3
- 9*(b*d^3*e - b*c*d^2*f)*(f*x + e)^2 + 9*(b*d^3*e^2 - 2*b*c*d^2*e*f + b*c^
2*d*f^2)*(f*x + e))*log(((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x +
2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x
+ 2*e))/(a^2 + b^2)) + 6*(4*(f*x + e)*b*d^3 - 3*b*d^3*e + 3*b*c*d^2*f)*po
lylog(3, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)))/((a^2 + b^2)*f^3))/f

```

Giac [F]

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^3}{b \tan(fx + e) + a} dx$$

```

[In] integrate((d*x+c)^3/(a+b*tan(f*x+e)),x, algorithm="giac")
[Out] integrate((d*x + c)^3/(b*tan(f*x + e) + a), x)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tan(e + fx)} dx$$

```

[In] int((c + d*x)^3/(a + b*tan(e + f*x)),x)
[Out] int((c + d*x)^3/(a + b*tan(e + f*x)), x)

```

3.55 $\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$

Optimal result	367
Rubi [A] (verified)	368
Mathematica [A] (verified)	370
Maple [B] (verified)	370
Fricas [B] (verification not implemented)	371
Sympy [F]	372
Maxima [B] (verification not implemented)	372
Giac [F]	373
Mupad [F(-1)]	373

Optimal result

Integrand size = 20, antiderivative size = 181

$$\begin{aligned} \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx = & \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\ & - \frac{ibd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f^2} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^3} \end{aligned}$$

[Out] $1/3*(d*x+c)^3/(a+I*b)/d+b*(d*x+c)^2*\ln(1+(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-I*b*d*(d*x+c)*\operatorname{polylog}(2, -(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2+1/2*b*d^2*\operatorname{polylog}(3, -(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^3$

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.250, Rules used = {3813, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx = & -\frac{ibd(c+dx)\operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f^2(a^2+b^2)} \\ & + \frac{b(c+dx)^2 \log\left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f(a^2+b^2)} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f^3(a^2+b^2)} + \frac{(c+dx)^3}{3d(a+ib)} \end{aligned}$$

[In] `Int[(c + d*x)^2/(a + b*Tan[e + f*x]), x]`

[Out] $(c + d*x)^3/(3*(a + I*b)*d) + (b*(c + d*x)^2*\operatorname{Log}[1 + ((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f) - (I*b*d*(c + d*x)*\operatorname{PolyLog}[2, -((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2])/((a^2 + b^2)*f^2) + (b*d^2*\operatorname{PolyLog}[3, -(((a^2 + b^2)*E^((2*I)*(e + f*x)))/(a + I*b)^2)]/(2*(a^2 + b^2)*f^3))$

Rule 2221

```
Int[((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simplify[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3813

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Sy
mbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int
[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*
I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2,
0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_ + (b_)*(x_))^p]/((d_)*(e_)*(x_)), x_Sy
mbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^3}{3(a+ib)d} + (2ib) \int \frac{e^{2i(e+fx)}(c+dx)^2}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx \\
&= \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\
&\quad - \frac{(2bd) \int (c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) dx}{(a^2+b^2)f} \\
&= \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\
&\quad - \frac{ibd(c+dx) \text{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f^2} \\
&\quad + \frac{(ibd^2) \int \text{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) dx}{(a^2+b^2)f^2} \\
&= \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} \\
&\quad - \frac{ibd(c+dx) \text{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f^2} \\
&\quad + \frac{(bd^2) \text{Subst} \left(\int \frac{\text{PolyLog} \left(2, -\frac{(a^2+b^2)x}{(a+ib)^2}\right)}{x} dx, x, e^{2i(e+fx)} \right)}{2(a^2+b^2)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^3}{3(a+ib)d} + \frac{b(c+dx)^2 \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} \\
&\quad - \frac{i b d (c+dx) \operatorname{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f^2} + \frac{b d^2 \operatorname{PolyLog} \left(3, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.43 (sec), antiderivative size = 236, normalized size of antiderivative = 1.30

$$\begin{aligned}
&\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx \\
&= \frac{1}{6}b \left(-\frac{4(c+dx)^3}{(ia+b)d(-ib(-1+e^{2ie})+a(1+e^{2ie}))} + \frac{6(c+dx)^2 \log \left(1 + \frac{(a+ib)e^{-2i(e+fx)}}{a-ib} \right)}{(a^2+b^2)f} \right. \\
&\quad \left. + \frac{3d \left(2if(c+dx) \operatorname{PolyLog} \left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib} \right) + d \operatorname{PolyLog} \left(3, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib} \right) \right)}{(a^2+b^2)f^3} \right) \\
&\quad + \frac{x(3c^2+3cdx+d^2x^2)\cos(e)}{3(a\cos(e)+b\sin(e))}
\end{aligned}$$

[In] `Integrate[(c + d*x)^2/(a + b*Tan[e + f*x]), x]`

[Out] `(b*(-4*(c + d*x)^3)/((I*a + b)*d*((-I)*b*(-1 + E^((2*I)*e))) + a*(1 + E^((2*I)*e))) + (6*(c + d*x)^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]/((a^2 + b^2)*f) + (3*d*((2*I)*f*(c + d*x)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] + d*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^3))/6 + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[e])/(3*(a*Cos[e] + b*Sin[e])))`

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 939 vs. $2(166) = 332$.

Time = 0.69 (sec), antiderivative size = 940, normalized size of antiderivative = 5.19

method	result
risch	$-\frac{d^2x^3}{3(ib-a)} - \frac{dcx^2}{ib-a} - \frac{c^2x}{ib-a} - \frac{c^3}{3d(ib-a)} + \frac{4be^3d^2}{3f^3(ia+b)(-ib-a)} - \frac{2ibcd \ln \left(1 - \frac{(-ib+a)e^{2i(fx+e)}}{-ib-a} \right) x}{f(ia+b)(-ib-a)} + \frac{2be^2d^2x}{f^2(ia+b)(-ib-a)} - \frac{b^2e^4d^4}{f^4(ia+b)(-ib-a)}$

[In] `int((d*x+c)^2/(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)`

```
[Out] -1/3*d^2/(I*b-a)*x^3-d/(I*b-a)*c*x^2-1/(I*b-a)*c^2*x-1/3/d/(I*b-a)*c^3+4/3/
f^3/(I*a+b)*b/(-I*b-a)*e^3*d^2-2*I/f/(I*a+b)*b/(-I*b-a)*c*d*ln(1-(a-I*b)*ex
p(2*I*(f*x+e))/(-I*b-a))*x+2/f^2/(I*a+b)*b/(-I*b-a)*e^2*d^2*x-1/f^2/(I*a+b)
*b/(-I*b-a)*d^2*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x-2*I/f^2/(I*a
+b)*b/(-I*b-a)*c*d*ln(1-(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*e-2*I/f^3/(I*a+b)
)*b*d^2*e^2/(a+I*b)*ln(exp(I*(f*x+e)))-2/(I*a+b)*b/(-I*b-a)*c*d*x^2-4/f/(I*
a+b)*b/(-I*b-a)*c*d*e*x-2/f^2/(I*a+b)*b/(-I*b-a)*c*d*e^2-1/f^2/(I*a+b)*b/(-
I*b-a)*c*d*polylog(2,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-1/2*I/f^3/(I*a+b)*b
/(-I*b-a)*d^2*polylog(3,(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))-2*I/f^2/(I*a+b)*
b*c*d*e/(a+I*b)*ln(I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)-2*I/f/(I*
a+b)*b*c^2/(a+I*b)*ln(exp(I*(f*x+e)))+I/f/(I*a+b)*b*c^2/(a+I*b)*ln(I*exp(2*
I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)+4*I/f^2/(I*a+b)*b*c*d*e/(a+I*b)*ln(e
xp(I*(f*x+e)))+I/f^3/(I*a+b)*b/(-I*b-a)*e^2*d^2*ln(1-(a-I*b)*exp(2*I*(f*x+e))
)/(-I*b-a))-2/3/(I*a+b)*b/(-I*b-a)*d^2*x^3-I/f/(I*a+b)*b/(-I*b-a)*d^2*ln(1-
(a-I*b)*exp(2*I*(f*x+e))/(-I*b-a))*x^2+I/f^3/(I*a+b)*b*d^2*e^2/(a+I*b)*ln(
I*exp(2*I*(f*x+e))*b-a*exp(2*I*(f*x+e))-I*b-a)
```

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 834 vs. $2(160) = 320$.

Time = 0.27 (sec), antiderivative size = 834, normalized size of antiderivative = 4.61

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

$$= \frac{4 ad^2 f^3 x^3 + 12 acdf^3 x^2 + 12 ac^2 f^3 x + 3 bd^2 \text{polylog}\left(3, \frac{(a^2 + 2i ab - b^2) \tan(fx+e)^2 - a^2 - 2i ab + b^2 - 2 (-i a^2 + 2 ab + i b^2) \tan(fx+e)^2}{(a^2 + b^2) \tan(fx+e)^2 + a^2 + b^2}\right)}{a^2 + b^2}$$

```
[In] integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="fricas")

[Out] 1/12*(4*a*d^2*f^3*x^3 + 12*a*c*d*f^3*x^2 + 12*a*c^2*f^3*x + 3*b*d^2*polylog
(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2
+ 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) +
3*b*d^2*polylog(3, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b +
b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2
+ a^2 + b^2)) - 6*(-I*b*d^2*f*x - I*b*c*d*f)*dilog(2*((I*a*b - b^2)*tan(f*x
+ e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*
tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*b*d^2*f*x + I*b*c*d*f)*dilog(2*((-I
*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x
+ e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 6*(b*d^2*f^2*x^2 + 2
*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-2*((I*a*b - b^2)*tan(f*x + e)^
2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*
x + e)^2 + a^2 + b^2)) + 6*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b
```

```

*c*d*e*f)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2
*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^
2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + 6*(b*d^2*
e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*
a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)))/((a^2 + b^2)*f^3
)

```

Sympy [F]

$$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx = \int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx$$

```
[In] integrate((d*x+c)**2/(a+b*tan(f*x+e)),x)
[Out] Integral((c + d*x)**2/(a + b*tan(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 715 vs. $2(160) = 320$.

Time = 0.55 (sec) , antiderivative size = 715, normalized size of antiderivative = 3.95

$$\int \frac{(c+dx)^2}{a+b\tan(e+fx)} dx =$$

$$6 cde \left(\frac{2(fx+e)a}{(a^2+b^2)f} + \frac{2b\log(b\tan(fx+e)+a)}{(a^2+b^2)f} - \frac{b\log(\tan(fx+e)^2+1)}{(a^2+b^2)f} \right) - 3 \left(\frac{2(fx+e)a}{a^2+b^2} + \frac{2b\log(b\tan(fx+e)+a)}{a^2+b^2} - \frac{b\log(\tan(fx+e)^2+1)}{a^2+b^2} \right)$$

```
[In] integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")
[Out] -1/6*(6*c*d*e*(2*(f*x + e)*a/((a^2 + b^2)*f) + 2*b*log(b*tan(f*x + e) + a)/((a^2 + b^2)*f) - b*log(tan(f*x + e)^2 + 1)/((a^2 + b^2)*f)) - 3*(2*(f*x + e)*a/(a^2 + b^2) + 2*b*log(b*tan(f*x + e) + a)/(a^2 + b^2) - b*log(tan(f*x + e)^2 + 1)/(a^2 + b^2))*c^2 - (2*(f*x + e)^3*(a - I*b)*d^2 + 6*(f*x + e)*(a - I*b)*d^2*e^2 + 6*I*b*d^2*e^2*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) + 3*b*d^2*e^2*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 3*b*d^2*polylog(3, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*((a - I*b)*d^2*e - (a - I*b)*c*d*f)*(f*x + e)^2 - 6*(I*(f*x + e)^2*b*d^2 + 2*(-I*b*d^2*e + I*b*c*d*f)*(f*x + e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e)^2 + 2*a*b*sin(2*f*x + 2*e)^2 + a^2*b^2 - 2*a*b*cos(2*f*x + 2*e)*sin(2*f*x + 2*e))/((a^2 - b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 - b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)), (a^2 - b^2)*cos(2*f*x + 2*e) + 2*a*b*sin(2*f*x + 2*e) + (a^2 - b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 3*b*d^2*polylog(3, (I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) - 6*((a - I*b)*d^2*e - (a - I*b)*c*d*f)*(f*x + e)^2 - 6*(I*(f*x + e)^2*b*d^2 + 2*(-I*b*d^2*e + I*b*c*d*f)*(f*x + e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e)^2 + 2*a*b*sin(2*f*x + 2*e)^2 + a^2*b^2 - 2*a*b*cos(2*f*x + 2*e)*sin(2*f*x + 2*e))/((a^2 - b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 - b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)), (a^2 - b^2)*cos(2*f*x + 2*e) + 2*a*b*sin(2*f*x + 2*e) + (a^2 - b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e))
```

$$\frac{e))}{(a^2 + b^2)}, \frac{(2*a*b*\sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*\cos(2*f*x + 2*e))}{(a^2 + b^2)} - 6*(I*(f*x + e)*b*d^2 - I*b*d^2*e + I*b*c*d*f)*dilo g((I*a + b)*e^{(2*I*f*x + 2*I*e)/(-I*a + b)}) + 3*((f*x + e)^2*b*d^2 - 2*(b*d^2*e - b*c*d*f)*(f*x + e))*\log((a^2 + b^2)*\cos(2*f*x + 2*e)^2 + 4*a*b*\sin(2*f*x + 2*e) + (a^2 + b^2)*\sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*\cos(2*f*x + 2*e))/(a^2 + b^2))/((a^2 + b^2)*f^2))/f$$

Giac [F]

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(dx + c)^2}{b \tan(fx + e) + a} dx$$

[In] integrate((d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")
[Out] integrate((d*x + c)^2/(b*tan(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tan(e + fx)} dx$$

[In] int((c + d*x)^2/(a + b*tan(e + f*x)),x)
[Out] int((c + d*x)^2/(a + b*tan(e + f*x)), x)

3.56 $\int \frac{c+dx}{a+b\tan(e+fx)} dx$

Optimal result	374
Rubi [A] (verified)	374
Mathematica [A] (verified)	376
Maple [B] (verified)	376
Fricas [B] (verification not implemented)	377
Sympy [F]	377
Maxima [B] (verification not implemented)	378
Giac [F]	378
Mupad [F(-1)]	378

Optimal result

Integrand size = 18, antiderivative size = 125

$$\int \frac{c+dx}{a+b\tan(e+fx)} dx = \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)f} - \frac{ibd\operatorname{PolyLog}\left(2,-\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2(a^2+b^2)f^2}$$

[Out] $1/2*(d*x+c)^2/(a+I*b)/d+b*(d*x+c)*\ln(1+(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f-1/2*I*b*d*\operatorname{polylog}(2,-(a^2+b^2)*\exp(2*I*(f*x+e))/(a+I*b)^2)/(a^2+b^2)/f^2$

Rubi [A] (verified)

Time = 0.18 (sec), antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.222, Rules used = {3813, 2221, 2317, 2438}

$$\int \frac{c+dx}{a+b\tan(e+fx)} dx = \frac{b(c+dx)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f(a^2+b^2)} - \frac{ibd\operatorname{PolyLog}\left(2,-\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{2f^2(a^2+b^2)} + \frac{(c+dx)^2}{2d(a+ib)}$$

[In] $\operatorname{Int}[(c+d*x)/(a+b*\operatorname{Tan}[e+f*x]),x]$

[Out] $(c+d*x)^2/(2*(a+I*b)*d) + (b*(c+d*x)*\operatorname{Log}[1+((a^2+b^2)*E^{(2*I)*(e+f*x)}))/(a+I*b)^2])/((a^2+b^2)*f) - ((I/2)*b*d*\operatorname{PolyLog}[2,-(((a^2+b^2)*E^{(2*I)*(e+f*x)}))/(a+I*b)^2])/((a^2+b^2)*f^2)$

Rule 2221

```
Int[((F_)((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*(F_)((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*(F_)((e_.)*(c_.) + (d_.)*(x_.)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_) + (e_.)*(x_.)^n_]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3813

```
Int[((c_.) + (d_.)*(x_.))^m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^2}{2(a+ib)d} + (2ib) \int \frac{e^{2i(e+fx)}(c+dx)}{(a+ib)^2 + (a^2+b^2)e^{2i(e+fx)}} dx \\
&= \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} - \frac{(bd) \int \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right) dx}{(a^2+b^2)f} \\
&= \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} \\
&\quad + \frac{(ibd) \text{Subst} \left(\int \frac{\log \left(1 + \frac{(a^2+b^2)x}{(a+ib)^2} \right)}{x} dx, x, e^{2i(e+fx)} \right)}{2(a^2+b^2)f^2} \\
&= \frac{(c+dx)^2}{2(a+ib)d} + \frac{b(c+dx) \log \left(1 + \frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{(a^2+b^2)f} - \frac{ibd \text{PolyLog} \left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2} \right)}{2(a^2+b^2)f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.56 (sec) , antiderivative size = 177, normalized size of antiderivative = 1.42

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \frac{b(c + dx)^2}{(a - ib)d(b - be^{2ie} - ia(1 + e^{2ie}))} \\ + \frac{b(c + dx) \log\left(1 + \frac{(a+ib)e^{-2i(e+fx)}}{a-ib}\right)}{(a^2 + b^2)f} \\ + \frac{ibd \operatorname{PolyLog}\left(2, \frac{(-a-ib)e^{-2i(e+fx)}}{a-ib}\right)}{2(a^2 + b^2)f^2} + \frac{x(2c + dx) \cos(e)}{2(a \cos(e) + b \sin(e))}$$

[In] `Integrate[(c + d*x)/(a + b*Tan[e + f*x]), x]`

[Out]
$$\frac{(b*(c + d*x)^2)/((a - I*b)*d*(b - b*E^{(2*I)*e} - I*a*(1 + E^{(2*I)*e}))) + (b*(c + d*x)*Log[1 + (a + I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a^2 + b^2)*f) + ((I/2)*b*d*PolyLog[2, (-a - I*b)/((a - I*b)*E^{(2*I)*(e + f*x)})])/((a^2 + b^2)*f^2) + (x*(2*c + d*x)*Cos[e])/(2*(a*Cos[e] + b*Sin[e]))}{(a^2 + b^2)*f^2}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(113) = 226.

Time = 0.64 (sec) , antiderivative size = 462, normalized size of antiderivative = 3.70

method	result
risch	$-\frac{dx^2}{2(ib-a)} - \frac{cx}{ib-a} + \frac{bc \ln(ie^{2i(fx+e)}b-a e^{2i(fx+e)}-ib-a)}{f(-ib+a)(ib+a)} - \frac{2bc \ln(e^{i(fx+e)})}{f(-ib+a)(ib+a)} - \frac{bd \ln\left(1 - \frac{(-ib+a)e^{2i(fx+e)}}{-ib-a}\right)x}{f(-ib+a)(-ib-a)} + \frac{ibd x^2}{(-ib+a)(-ib-a)}$

[In] `int((d*x+c)/(a+b*tan(f*x+e)), x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/2/(I*b-a)*d*x^2-1/(I*b-a)*c*x+1/f*b/(a-I*b)*c/(a+I*b)*\ln(I*\exp(2*I*(f*x+e)))*b-a*\exp(2*I*(f*x+e))-I*b-a-2/f*b/(a-I*b)*c/(a+I*b)*\ln(\exp(I*(f*x+e)))-1/f*b/(a-I*b)/(-I*b-a)*d*\ln(1-(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))*x+I*b/(a-I*b)/(-I*b-a)*d*x^2-1/f^2*b/(a-I*b)/(-I*b-a)*d*\ln(1-(a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))/(-I*b-a)*e+2*I/f*b/(a-I*b)/(-I*b-a)*d*e*x+I/f^2*b/(a-I*b)/(-I*b-a)*d*e^2+1/2*I/f^2*b/(a-I*b)/(-I*b-a)*d*polylog(2, (a-I*b)*\exp(2*I*(f*x+e))/(-I*b-a))-1/f^2*b/(a-I*b)*d*e/(a+I*b)*\ln(I*\exp(2*I*(f*x+e)))*b-a*\exp(2*I*(f*x+e))-I*b-a)+2/f^2*b/(a-I*b)*d*e/(a+I*b)*\ln(\exp(I*(f*x+e))) \end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 527 vs. $2(108) = 216$.

Time = 0.27 (sec) , antiderivative size = 527, normalized size of antiderivative = 4.22

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \frac{2 adf^2 x^2 + 4 acf^2 x + i bd \text{Li}_2\left(\frac{2((i ab - b^2) \tan(fx+e)^2 - a^2 - i ab + (i a^2 - 2 ab - i b^2) \tan(fx+e))}{(a^2 + b^2) \tan(fx+e)^2 + a^2 + b^2}\right) + 1}{2 adf^2 x^2 + 4 acf^2 x + i bd \text{Li}_2\left(\frac{2((-i ab - b^2) \tan(fx+e)^2 - a^2 - i ab + (i a^2 - 2 ab - i b^2) \tan(fx+e))}{(a^2 + b^2) \tan(fx+e)^2 + a^2 + b^2}\right) + 1}$$

```
[In] integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fricas")
[Out] 1/4*(2*a*d*f^2*x^2 + 4*a*c*f^2*x + I*b*d*dilog(2*((I*a*b - b^2)*tan(f*x + e))^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - I*b*d*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(b*d*f*x + b*d*e)*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(b*d*f*x + b*d*e)*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 2*(b*d*e - b*c*f)*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 2*(b*d*e - b*c*f)*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)))/((a^2 + b^2)*f^2)
```

Sympy [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + dx}{a + b \tan(e + fx)} dx$$

```
[In] integrate((d*x+c)/(a+b*tan(f*x+e)),x)
[Out] Integral((c + d*x)/(a + b*tan(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 399 vs. $2(108) = 216$.

Time = 0.60 (sec) , antiderivative size = 399, normalized size of antiderivative = 3.19

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \frac{(a - i b) df^2 x^2 + 2 (a - i b) c f^2 x - 2 i b d f x \arctan\left(\frac{2 a b \cos(2 f x + 2 e) - (a^2 - b^2) \sin(2 f x + 2 e)}{a^2 + b^2}, \frac{2 a b \sin(2 f x + 2 e) + a^2 + b^2 + (a^2 - b^2) \cos(2 f x + 2 e)}{a^2 + b^2}\right)}{a^2 + b^2}$$

[In] `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

[Out] $\frac{1}{2} ((a - I b) d f^2 x^2 + 2 (a - I b) c f^2 x - 2 I b d f x \operatorname{arctan2}(2 a b \cos(2 f x + 2 e) - (a^2 - b^2) \sin(2 f x + 2 e), (2 a b \sin(2 f x + 2 e) + a^2 + b^2 + (a^2 - b^2) \cos(2 f x + 2 e)) / (a^2 + b^2)) + b d f x \log((a^2 + b^2) \cos(2 f x + 2 e))^2 + 4 a b \sin(2 f x + 2 e) + (a^2 + b^2) \sin(2 f x + 2 e)^2 + a^2 + b^2 + 2 (a^2 - b^2) \cos(2 f x + 2 e)) / (a^2 + b^2) + 2 I b c f \operatorname{arctan2}(-b \cos(2 f x + 2 e) + a \sin(2 f x + 2 e) + b, a \cos(2 f x + 2 e) + b \sin(2 f x + 2 e) + a) + b c f \log((a^2 + b^2) \cos(2 f x + 2 e))^2 + 4 a b \sin(2 f x + 2 e) + (a^2 + b^2) \sin(2 f x + 2 e)^2 + a^2 + b^2 + 2 (a^2 - b^2) \cos(2 f x + 2 e)) - I b d \operatorname{dilog}(I a + b) e^{(2 I f x + 2 I e) / (-I a + b)}) / ((a^2 + b^2) f^2)$

Giac [F]

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{dx + c}{b \tan(f x + e) + a} dx$$

[In] `integrate((d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate((d*x + c)/(b*tan(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{a + b \tan(e + fx)} dx = \int \frac{c + d x}{a + b \tan(e + f x)} dx$$

[In] `int((c + d*x)/(a + b*tan(e + f*x)),x)`

[Out] `int((c + d*x)/(a + b*tan(e + f*x)), x)`

3.57 $\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$

Optimal result	379
Rubi [N/A]	379
Mathematica [N/A]	380
Maple [N/A] (verified)	380
Fricas [N/A]	380
Sympy [N/A]	380
Maxima [N/A]	381
Giac [N/A]	381
Mupad [N/A]	381

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\tan(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*tan(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.07 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx = \int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$$

[In] Int[1/((c + d*x)*(a + b*Tan[e + f*x])),x]

[Out] Defer[Int][1/((c + d*x)*(a + b*Tan[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b\tan(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 2.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx$$

[In] `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])),x]`

[Out] `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.42 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tan(fx + e))} dx$$

[In] `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

[Out] `int(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.35

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="fricas")`

[Out] `integral(1/(a*d*x + a*c + (b*d*x + b*c)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x)`

[Out] `Integral(1/((a + b*tan(e + f*x))*(c + d*x)), x)`

Maxima [N/A]

Not integrable

Time = 0.89 (sec) , antiderivative size = 279, normalized size of antiderivative = 13.95

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & (2*(a^2*b + b^3)*d*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d*x + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d*x + (a^4 + 2*a^2*b^2 + b^4)*c)*sin(2*f*x + 2*e)^2 + (a^4 + 2*a^2*b^2 + b^4)*c + 2*((a^4 - b^4)*d*x + (a^4 - b^4)*c)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d*x + (a^3*b + a*b^3)*c)*sin(2*f*x + 2*e)), x) + a*log(d*x + c))/((a^2 + b^2)*d) \end{aligned}$$

Giac [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate(1/((d*x + c)*(b*tan(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 3.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + f x)) (c + d x)} dx$$

[In] `int(1/((a + b*tan(e + f*x))*(c + d*x)),x)`

[Out] `int(1/((a + b*tan(e + f*x))*(c + d*x)), x)`

3.58 $\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$

Optimal result	382
Rubi [N/A]	382
Mathematica [N/A]	383
Maple [N/A] (verified)	383
Fricas [N/A]	383
Sympy [N/A]	384
Maxima [N/A]	384
Giac [N/A]	384
Mupad [N/A]	385

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\tan(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*tan(f*x+e)), x)

Rubi [N/A]

Not integrable

Time = 0.07 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

[In] Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 4.56 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx$$

[In] `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]`

[Out] `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx+c)^2(a+b\tan(fx+e))} dx$$

[In] `int(1/(d*x+c)^2/(a+b*tan(f*x+e)), x)`

[Out] `int(1/(d*x+c)^2/(a+b*tan(f*x+e)), x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)), x, algorithm="fricas")`

[Out] `integral(1/(a*d^2*x^2 + 2*a*c*d*x + a*c^2 + (b*d^2*x^2 + 2*b*c*d*x + b*c^2)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx))(c + dx)^2} dx$$

[In] `integrate(1/(d*x+c)**2/(a+b*tan(f*x+e)),x)`

[Out] `Integral(1/((a + b*tan(e + f*x))*(c + d*x)**2), x)`

Maxima [N/A]

Not integrable

Time = 1.73 (sec) , antiderivative size = 424, normalized size of antiderivative = 21.20

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="maxima")`

[Out] `(2*((a^2*b + b^3)*d^2*x + (a^2*b + b^3)*c*d)*integrate((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*cos(2*f*x + 2*e)^2 + ((a^4 + 2*a^2*b^2 + b^4)*d^2*x^2 + 2*(a^4 + 2*a^2*b^2 + b^4)*c*d*x + (a^4 + 2*a^2*b^2 + b^4)*c^2)*sin(2*f*x + 2*e)^2 + 2*((a^4 - b^4)*d^2*x^2 + 2*(a^4 - b^4)*c*d*x + (a^4 - b^4)*c^2)*cos(2*f*x + 2*e) + 4*((a^3*b + a*b^3)*d^2*x^2 + 2*(a^3*b + a*b^3)*c*d*x + (a^3*b + a*b^3)*c^2)*sin(2*f*x + 2*e)), x) - a)/((a^2 + b^2)*d^2*x + (a^2 + b^2)*c*d)`

Giac [N/A]

Not integrable

Time = 4.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e)),x, algorithm="giac")`

[Out] `integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 4.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))} dx = \int \frac{1}{(a + b \tan(e + fx)) (c + dx)^2} dx$$

[In] `int(1/((a + b*tan(e + f*x))*(c + d*x)^2),x)`

[Out] `int(1/((a + b*tan(e + f*x))*(c + d*x)^2), x)`

3.59 $\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$

Optimal result	387
Rubi [A] (verified)	388
Mathematica [B] (warning: unable to verify)	396
Maple [B] (verified)	398
Fricas [B] (verification not implemented)	398
Sympy [F]	400
Maxima [B] (verification not implemented)	400
Giac [F]	403
Mupad [F(-1)]	403

Optimal result

Integrand size = 20, antiderivative size = 848

$$\begin{aligned}
\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx = & -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2 (ia-b+(ia+b)e^{2ie+2ifx}) f} \\
& + \frac{(c+dx)^4}{4(a-ib)^2 d} + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} \\
& + \frac{3b^2d(c+dx)^2 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
& + \frac{2b(c+dx)^3 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2(a+ib)f} \\
& - \frac{2ib^2(c+dx)^3 \log \left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f} \\
& - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3} \\
& + \frac{3bd(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(ia-b)(a-ib)^2 f^2} \\
& - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog} \left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^2} \\
& + \frac{3b^2d^3 \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4} \\
& + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a-ib)^2(a+ib)f^3} \\
& - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog} \left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{(a^2+b^2)^2 f^3} \\
& - \frac{3bd^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(ia-b)(a-ib)^2 f^4} \\
& + \frac{3b^2d^3 \operatorname{PolyLog} \left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib} \right)}{2(a^2+b^2)^2 f^4}
\end{aligned}$$

```
[Out] -3*I*b^2*d^2*(d*x+c)*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+2*b^2*(d*x+c)^3/(a+I*b)/(I*a+b)^2/(I*a-b+(I*a+b)*exp(2*I*e+2*I*f*x))/f+1/4*(d*x+c)^4/(a-I*b)^2/d+b*(d*x+c)^4/(I*a-b)/(a-I*b)^2/d-b^2*(d*x+c)^4/(a^2+b^2)^2/d+3*b^2*d*(d*x+c)^2*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/
```

$$\begin{aligned}
& (a^2+b^2)^2/f^2+2*b*(d*x+c)^3*\ln(1+(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f-2*I*b^2*(d*x+c)^3*\ln(1+(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/ \\
& (a^2+b^2)^2/f-3*I*b^2*d^2*(d*x+c)*\text{polylog}(3,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+3*b*d*(d*x+c)^2*\text{polylog}(2,-(a-I*b)*\exp(2*I*e+2*I*f*x) \\
& /(a+I*b))/(I*a-b)/(a-I*b)^2/f^2-3*b^2*d*(d*x+c)^2*\text{polylog}(2,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+3/2*b^2*d^3*\text{polylog}(3,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b)) \\
& /(a^2+b^2)^2/f^4+3*b*d^2*(d*x+c)*\text{polylog}(3,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/f^4-2*I*b^2*(d*x+c)^3/(a^2+b^2) \\
& ^2/f-3/2*b*d^3*\text{polylog}(4,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^4+3/2*b^2*d^3*\text{polylog}(4,-(a-I*b)*\exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^4
\end{aligned}$$

Rubi [A] (verified)

Time = 2.31 (sec), antiderivative size = 848, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.450, Rules used

$$= \{3815, 2216, 2215, 2221, 2611, 6744, 2320, 6724, 2222\}$$

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx &= \frac{b(c+dx)^4}{(ia-b)(a-ib)^2d} + \frac{(c+dx)^4}{4(a-ib)^2d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2d} \\ &\quad + \frac{2b \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right) (c+dx)^3}{(a-ib)^2(a+ib)f} \\ &\quad - \frac{2ib^2 \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right) (c+dx)^3}{(a^2+b^2)^2f} \\ &\quad + \frac{2b^2(c+dx)^3}{(a+ib)(ia+b)^2(iia+(ia+b)e^{2ie+2ifx}-b)f} \\ &\quad - \frac{2ib^2(c+dx)^3}{(a^2+b^2)^2f} + \frac{3b^2d \log\left(\frac{e^{2ie+2ifx}(a-ib)}{a+ib} + 1\right) (c+dx)^2}{(a^2+b^2)^2f^2} \\ &\quad + \frac{3bd \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right) (c+dx)^2}{(ia-b)(a-ib)^2f^2} \\ &\quad - \frac{3b^2d \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right) (c+dx)^2}{(a^2+b^2)^2f^2} \\ &\quad - \frac{3ib^2d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right) (c+dx)}{(a^2+b^2)^2f^3} \\ &\quad + \frac{3bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right) (c+dx)}{(a-ib)^2(a+ib)f^3} \\ &\quad - \frac{3ib^2d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right) (c+dx)}{(a^2+b^2)^2f^3} \\ &\quad + \frac{3b^2d^3 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2f^4} \\ &\quad - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(ia-b)(a-ib)^2f^4} \\ &\quad + \frac{3b^2d^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2f^4} \end{aligned}$$

[In] Int[(c + d*x)^3/(a + b*Tan[e + f*x])^2, x]

[Out] $\frac{((-2*I)*b^2*(c + d*x)^3)/((a^2 + b^2)^2*f) + (2*b^2*(c + d*x)^3)/((a + I*b)*(I*a + b)^2*(I*a - b + (I*a + b)*E^{((2*I)*e + (2*I)*f*x))*f) + (c + d*x)^4)/(4*(a - I*b)^2*d) + (b*(c + d*x)^4)/((I*a - b)*(a - I*b)^2*d) - (b^2*(c + d*x)^4)/((a^2 + b^2)^2*d) + (3*b^2*d*(c + d*x)^2*\operatorname{Log}[1 + ((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})])}{(a^2 + b^2)^2*f^4}$

$$\begin{aligned} & (*e + (2*I)*f*x)/(a + I*b)])/((a^2 + b^2)^2*f^2) + (2*b*(c + d*x)^3*\text{Log}[1 \\ & + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])/((a - I*b)^2*(a + I*b)*f) \\ & - ((2*I)*b^2*(c + d*x)^3*\text{Log}[1 + ((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])/((a^2 + b^2)^2*f) - ((3*I)*b^2*d^2*(c + d*x)*\text{PolyLog}[2, -(((a - I*b) \\ & *E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^3) + (3*b*d*(c + d*x)^2*\text{PolyLog}[2, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((I*a - b)*(a - I*b)^2*f^2) - (3*b^2*d*(c + d*x)^2*\text{PolyLog}[2, -(((a - I*b)*E^((2*I) \\ & *e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^2) + (3*b^2*d^3*\text{PolyLog}[3, -((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b)])]/(2*(a^2 + b^2)^2*f^4) + (3*b*d^2*(c + d*x)*\text{PolyLog}[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a - I*b)^2*(a + I*b)*f^3) - ((3*I)*b^2*d^2*(c + d*x)*\text{PolyLog}[3, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/((a^2 + b^2)^2*f^3) - (3*b*d^3*\text{PolyLog}[4, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/(2*(I*a - b)*(a - I*b)^2*f^4) + (3*b^2*d^3*\text{PolyLog}[4, -(((a - I*b)*E^((2*I)*e + (2*I)*f*x))/(a + I*b))]/(2*(a^2 + b^2)^2*f^4) \end{aligned}$$
Rule 2215

$$\text{Int}[(c_+ + d_-)*(x_-)^m/((a_- + b_-)*(F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n), x] \rightarrow \text{Simp}[(c + d*x)^{m+1}/(a*d*(m+1)), x] - \text{Dist}[b/a, \text{Int}[(c + d*x)^m*((F^((g*(e + f*x))))^n/(a + b*(F^((g*(e + f*x))))^n)), x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$$
Rule 2216

$$\text{Int}[(a_- + b_-)*(F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n*(c_- + d_-)*(x_-)^p, x] \rightarrow \text{Dist}[1/a, \text{Int}[(c + d*x)^m*(a + b*(F^((g*(e + f*x))))^n)^{p+1}, x], x] - \text{Dist}[b/a, \text{Int}[(c + d*x)^m*(F^((g*(e + f*x))))^n*(a + b*(F^((g*(e + f*x))))^n)^p, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{ILtQ}[p, 0] \&& \text{IGtQ}[m, 0]$$
Rule 2221

$$\text{Int}[((F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n*(c_- + d_-)*(x_-)^m/((a_- + b_-)*(F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n), x] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^((g*(e + f*x))))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{(m-1)}*\text{Log}[1 + b*((F^((g*(e + f*x))))^n/a)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$$
Rule 2222

$$\text{Int}[(F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n*(a_- + b_-)*(F_-)^((g_-)*(e_-) + (f_-)*(x_-))^p/((a_- + b_-)*(F_-)^((g_-)*(e_-) + (f_-)*(x_-)))^n), x] \rightarrow \text{Simp}[(c + d*x)^m*((a + b*(F^((g*(e + f*x))))^n)^{p+1})/(b*f*g*n*(p+1)*\text{Log}[F])), x] - \text{Dist}[d*(m/(b*f*g*n*(p+1)*\text{Log}[F])), \text{Int}[(c + d*x)^{(m-1)}*(a + b*(F^((g*(e + f*x))))^n)^{p+1}, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, m\}, x]$$

, n, p}, x] && NeQ[p, -1]

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*(a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3815

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(p_.)], x_Symbol] :> Simp[((e + f*x)^m)*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = \int \left(\frac{(c + dx)^3}{(a - ib)^2} - \frac{4b^2(c + dx)^3}{(ia + b)^2 \left(ia \left(1 + \frac{ib}{a} \right) + ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} \right)^2} \right. \\ \left. + \frac{4b(c + dx)^3}{(a - ib)^2 \left(ia \left(1 + \frac{ib}{a} \right) + ia \left(1 - \frac{ib}{a} \right) e^{2ie+2ifx} \right)} \right) dx$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a-ib)^2d} + \frac{(4b) \int \frac{(c+dx)^3}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{(a-ib)^2} - \frac{(4b^2) \int \frac{(c+dx)^3}{(ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx})^2} dx}{(ia+b)^2} \\
&= \frac{(c+dx)^4}{4(a-ib)^2d} + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2d} + \frac{(4b^2) \int \frac{(c+dx)^3}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{(ia-b)(a-ib)^2} \\
&\quad - \frac{(4b) \int \frac{e^{2ie+2ifx}(c+dx)^3}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{a^2+b^2} - \frac{(4b^2) \int \frac{e^{2ie+2ifx}(c+dx)^3}{(ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx})^2} dx}{a^2+b^2} \\
&= -\frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^4}{4(a-ib)^2d} + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2d} \\
&\quad - \frac{b^2(c+dx)^4}{(a^2+b^2)^2d} + \frac{2b(c+dx)^3 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{(4b^2) \int \frac{e^{2ie+2ifx}(c+dx)^3}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{(a+ib)^2(ia+b)} \\
&\quad - \frac{(6bd) \int (c+dx)^2 \log\left(1 + \frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a-ib)^2(a+ib)f} + \frac{(6b^2d) \int \frac{(c+dx)^2}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{(a-ib)^2(a+ib)f} \\
&= -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2f} - \frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^4}{4(a-ib)^2d} \\
&\quad + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2d} + \frac{2b(c+dx)^3 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\
&\quad - \frac{2ib^2(c+dx)^3 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2f} + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2f^2} \\
&\quad - \frac{(6bd^2) \int (c+dx) \operatorname{PolyLog}\left(2, -\frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(ia-b)(a-ib)^2f^2} \\
&\quad - \frac{(6b^2d) \int \frac{e^{2ie+2ifx}(c+dx)^2}{ia(1+\frac{ib}{a})+ia(1-\frac{ib}{a})e^{2ie+2ifx}} dx}{(a-ib)(a+ib)^2f} \\
&\quad + \frac{(6ib^2d) \int (c+dx)^2 \log\left(1 + \frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^4}{4(a-ib)^2 d} \\
&\quad + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} + \frac{3b^2d(c+dx)^2 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{2b(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{2ib^2(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{(3bd^3) \int \operatorname{PolyLog}\left(3, -\frac{\left(1-\frac{ib}{a}\right)e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{(6b^2d^2) \int (c+dx) \log\left(1+\frac{\left(1-\frac{ib}{a}\right)e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{(6b^2d^2) \int (c+dx) \operatorname{PolyLog}\left(2, -\frac{\left(1-\frac{ib}{a}\right)e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^4}{4(a-ib)^2 d} \\
&\quad + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} + \frac{3b^2d(c+dx)^2 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{2b(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{2ib^2(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
&\quad - \frac{(3bd^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, -\frac{(a-ib)x}{a+ib}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{2(ia-b)(a-ib)^2 f^4} \\
&\quad + \frac{(3ib^2d^3) \int \operatorname{PolyLog}\left(2, -\frac{\left(1-\frac{ib}{a}\right)e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^3} \\
&\quad + \frac{(3ib^2d^3) \int \operatorname{PolyLog}\left(3, -\frac{\left(1-\frac{ib}{a}\right)e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^4}{4(a-ib)^2 d} \\
&\quad + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} + \frac{3b^2d(c+dx)^2 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{2b(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{2ib^2(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(ia-b)(a-ib)^2 f^4} \\
&\quad + \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, -\frac{(a-ib)x}{a+ib}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{2(a^2+b^2)^2 f^4} \\
&\quad + \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, -\frac{(a-ib)x}{a+ib}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{2(a^2+b^2)^2 f^4}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^3}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^3}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx}) f} + \frac{(c+dx)^4}{4(a-ib)^2 d} \\
&\quad + \frac{b(c+dx)^4}{(ia-b)(a-ib)^2 d} - \frac{b^2(c+dx)^4}{(a^2+b^2)^2 d} + \frac{3b^2d(c+dx)^2 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{2b(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{2ib^2(c+dx)^3 \log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} - \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{3b^2d^3 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2 f^4} + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{3ib^2d^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\
&\quad - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(ia-b)(a-ib)^2 f^4} + \frac{3b^2d^3 \operatorname{PolyLog}\left(4, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{2(a^2+b^2)^2 f^4}
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1724 vs. $2(848) = 1696$.

Time = 9.64 (sec) , antiderivative size = 1724, normalized size of antiderivative = 2.03

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx \\
 &= b \left(-\frac{4c^2(-ib(-1+e^{2ie})+a(1+e^{2ie}))(3bd+2acf)x}{a^2+b^2} + \frac{4b(c+dx)^3}{a-ib} + \frac{2af(c+dx)^4}{(a-ib)d} + \frac{12cd(-ib(-1+e^{2ie})+a(1+e^{2ie}))(bd+acf)x \log\left(1+\frac{a}{a+ib}\right)}{(a+ib)(ia+b)f} \right. \\
 & \quad + \frac{3x^2(ac^2d - ibc^2d + ac^2d \cos(2e) + ibc^2d \cos(2e) + iac^2d \sin(2e) - bc^2d \sin(2e))}{2(a-ib)(a+ib)(a+ib+a \cos(2e) - ib \cos(2e) + ia \sin(2e) + b \sin(2e))} \\
 & \quad + \frac{x^3(acd^2 - ibcd^2 + acd^2 \cos(2e) + ibcd^2 \cos(2e) + iacd^2 \sin(2e) - bcd^2 \sin(2e))}{(a-ib)(a+ib)(a+ib+a \cos(2e) - ib \cos(2e) + ia \sin(2e) + b \sin(2e))} \\
 & \quad + \frac{x^4(ad^3 - ibd^3 + ad^3 \cos(2e) + ibd^3 \cos(2e) + iad^3 \sin(2e) - bd^3 \sin(2e))}{4(a-ib)(a+ib)(a+ib+a \cos(2e) - ib \cos(2e) + ia \sin(2e) + b \sin(2e))} \\
 & \quad + x \left(\frac{c^3}{a^2 + 2iab - b^2 + a^2 \cos(4e) - 2iab \cos(4e) - b^2 \cos(4e) + ia^2 \sin(4e) + 2ab \sin(4e) - ib^2 \sin(4e)} \right. \\
 & \quad \quad \left. (-a - ib + a \cos(2e) - ib \cos(2e) + ia \sin(2e) + b \sin(2e)) (-4ia) \right. \\
 & \quad + \frac{(a-ib)(a+ib)(a+ib+a \cos(2e) - ib \cos(2e) + ia \sin(2e) + b \sin(2e)) (a^2 + 2iab - b^2 + a^2 \cos(4e) - c^3 \cos(4e) + ic^3 \sin(4e))}{(a^2 + 2iab - b^2 + a^2 \cos(4e) - 2iab \cos(4e) - b^2 \cos(4e) + ia^2 \sin(4e) + 2ab \sin(4e) - ib^2 \sin(4e))} \\
 & \quad + \frac{b^2c^3 \sin(fx) + 3b^2c^2dx \sin(fx) + 3b^2cd^2x^2 \sin(fx) + b^2d^3x^3 \sin(fx)}{(a-ib)(a+ib)f(a \cos(e) + b \sin(e))(a \cos(e+fx) + b \sin(e+fx))}
 \end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + b*Tan[e + f*x])^2, x]

[Out]

```

(b*((-4*c^2*(-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(3*b*d + 2*a*c*f)*x)/(a^2 + b^2) + (4*b*(c + d*x)^3)/(a - I*b) + (2*a*f*(c + d*x)^4)/((a - I*b)*d) + (12*c*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)*f) + (6*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*x^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)*f) + (4*a*d^3*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*x^3*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)) + (2*c^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(3*b*d + 2*a*c*f)*Log[a + I*b + (a - I*b)*E^((2*I)*(e + f*x))])/((a + I*b)*(I*a + b)*f) + (6*c*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^2) + (3*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^3) + (3*a*d^3*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(2*f^2*x^2*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - (2*I)*f*x*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - PolyLog[4, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))

```

$$\begin{aligned}
& \frac{2*I)*(e + f*x))])))/((a^2 + b^2)*f^3)))/(2*(a - I*b)*(a + I*b)*(b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e)))*f) + (3*x^2*(a*c^2*d - I*b*c^2*d + a*c^2*d*\cos[2*e] + I*b*c^2*d*\cos[2*e] + I*a*c^2*d*\sin[2*e] - b*c^2*d*\sin[2*e])))/(2*(a - I*b)*(a + I*b)*(a + I*b + a*\cos[2*e] - I*b*\cos[2*e] + I*a*\sin[2*e] + b*\sin[2*e])) + (x^3*(a*c*d^2 - I*b*c*d^2 + a*c*d^2*\cos[2*e] + I*b*c*d^2*\cos[2*e] + I*a*c*d^2*\sin[2*e] - b*c*d^2*\sin[2*e])))/((a - I*b)*(a + I*b)*(a + I*b + a*\cos[2*e] - I*b*\cos[2*e] + I*a*\sin[2*e] + b*\sin[2*e])) + (x^4*(a*d^3 - I*b*d^3 + a*d^3*\cos[2*e] + I*b*d^3*\cos[2*e] + I*a*d^3*\sin[2*e] - b*d^3*\sin[2*e]))/(4*(a - I*b)*(a + I*b)*(a + I*b + a*\cos[2*e] - I*b*\cos[2*e] + I*a*\sin[2*e] + b*\sin[2*e])) + x*(c^3/(a^2 + (2*I)*a*b - b^2 + a^2*\cos[4*e] - (2*I)*a*b*\cos[4*e] - b^2*\cos[4*e] + I*a^2*\sin[4*e] + 2*a*b*\sin[4*e] - I*b^2*\sin[4*e]) + ((-a - I*b + a*\cos[2*e] - I*b*\cos[2*e] + I*a*\sin[2*e] + b*\sin[2*e]))*(-(-4*I)*a*b*c^3*\cos[2*e] + 4*a*b*c^3*\sin[2*e])))/((a - I*b)*(a + I*b)*(a + I*b + a*\cos[2*e] - I*b*\cos[2*e] + I*a*\sin[2*e] + b*\sin[2*e]))*(a^2 + (2*I)*a*b - b^2 + a^2*\cos[4*e] - (2*I)*a*b*\cos[4*e] - b^2*\cos[4*e] + I*a^2*\sin[4*e] + 2*a*b*\sin[4*e] - I*b^2*\sin[4*e])) + (c^3*\cos[4*e] + I*c^3*\sin[4*e])/((a^2 + (2*I)*a*b - b^2 + a^2*\cos[4*e] - (2*I)*a*b*\cos[4*e] - b^2*\cos[4*e] + I*a^2*\sin[4*e] + 2*a*b*\sin[4*e] - I*b^2*\sin[4*e])) + (b^2*c^3*\sin[f*x] + 3*b^2*c^2*d*x*\sin[f*x] + 3*b^2*c*d^2*x^2*\sin[f*x] + b^2*d^3*x^3*\sin[f*x]))/((a - I*b)*(a + I*b)*f*(a*\cos[e] + b*\sin[e])*(a*\cos[e + f*x] + b*\sin[e + f*x]))
\end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 5958 vs. $2(763) = 1526$.

Time = 0.97 (sec), antiderivative size = 5959, normalized size of antiderivative = 7.03

method	result	size
risch	Expression too large to display	5959

[In] `int((d*x+c)^3/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] result too large to display

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2537 vs. $2(695) = 1390$.

Time = 0.33 (sec), antiderivative size = 2537, normalized size of antiderivative = 2.99

$$\int \frac{(c + dx)^3}{(a + b\tan(e + fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="fricas")`

[Out] $\frac{1}{4}((a^3 - a*b^2)*d^3*f^4*x^4 - 4*b^3*c^3*f^3 - 4*(b^3*d^3*f^3 - (a^3 - a*b^2)*c*d^2*f^4)*x^3 - 6*(2*b^3*c*d^2*f^3 - (a^3 - a*b^2)*c^2*d*f^4)*x^2 - 4*(3*b^3*c^2*d*f^3 - (a^3 - a*b^2)*c^3*f^4)*x - 6*(-I*a^2*b*d^3*f^2*x^2 - I*a^2*b*c^2*d*f^2 - I*a*b^2*c*d^2*f^2 - I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)*x + (-I*a*b^2*d^3*f^2*x^2 - I*a*b^2*c^2*d*f^2 - I*b^3*c*d^2*f^2 - I*(2*a*b^2*c*d^2*f^2 + a*b^2*d^3*f)*x + (-I*a*b^2*d^3*f^2*x^2 - I*a*b^2*c^2*d*f^2 - I*b^3*c*d^2*f^2 - I*(2*a*b^2*c*d^2*f^2 + a*b^2*d^3*f)*x + (-I*a*b^2*c*d^2*f^2 + I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)*x + (I*a*b^2*d^3*f^2*x^2 + I*a*b^2*c^2*d*f^2 + I*a*b^2*c*d^2*f^2 + I*a*b^2*c^2*d*f^2 + I*b^3*c*d^2*f^2 + I*(2*a*b^2*c*d^2*f^2 + b^3*d^3*f)*x)*tan(f*x + e))*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 6*(I*a^2*b*d^3*f^2*x^2 + I*a^2*b*c^2*d*f^2 + I*a*b^2*c*d^2*f^2 + I*(2*a^2*b*c*d^2*f^2 + a*b^2*d^3*f)*x + (I*a*b^2*d^3*f^2*x^2 + I*a*b^2*c^2*d*f^2 + I*b^3*c*d^2*f^2 + I*(2*a*b^2*c*d^2*f^2 + b^3*d^3*f)*x)*tan(f*x + e))*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(2*a^2*b*d^3*f^3*x^3 + 2*a^2*b*d^3*f^3 + 6*a^2*b*c^2*d*e*f^2 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c*d^2*f^3 + a*b^2*d^3*f^2)*x^2 - 6*(a^2*b*c*d^2*e^2 - a*b^2*c*d^2*e)*f + 6*(a^2*b*c^2*d*f^3 + a*b^2*c*d^2*f^2)*x + (2*a*b^2*d^3*f^3*x^3 + 2*a*b^2*d^3*e^3 + 6*a*b^2*c^2*d*f^2 - 3*b^3*d^3*e^2 + 3*(2*a*b^2*c*d^2*f^3 + b^3*d^3*f^2)*x^2 - 6*(a*b^2*c*d^2*e^2 - b^3*c*d^2*e)*f + 6*(a*b^2*c*d^2*f^3 + b^3*c*d^2*f^2)*x)*tan(f*x + e))*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(2*a^2*b*d^3*f^3*x^3 + 2*a^2*b*d^3*f^3 + 6*a^2*b*c^2*d*f^2 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c*d^2*f^3 + a*b^2*d^3*f^2)*x^2 - 6*(a^2*b*c*d^2*e^2 - a*b^2*c*d^2*e)*f + 6*(a^2*b*c^2*d*f^3 + a*b^2*c*d^2*f^2)*x + (2*a*b^2*d^3*f^3*x^3 + 2*a*b^2*d^3*e^3 + 6*a*b^2*c^2*d*f^2 - 3*b^3*d^3*e^2 + 3*(2*a*b^2*c*d^2*f^3 + b^3*d^3*f^2)*x^2 - 6*(a*b^2*c*d^2*e^2 - b^3*c*d^2*e)*f + 6*(a*b^2*c^2*d*f^3 + b^3*c*d^2*f^2)*x)*tan(f*x + e))*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 2*(2*a^2*b*d^3*f^3 - 2*a^2*b*c^3*f^3 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c^2*d*f^3 - a*b^2*c^2*d*f^2 - 6*(a^2*b*c*d^2*e^2 - a*b^2*c*d^2*e)*f + (2*a*b^2*d^3*f^3 - 2*a*b^2*c^3*f^3 - 3*b^3*d^3*e^2 + 3*(2*a*b^2*c^2*d*f^3 - b^3*c^2*d)*f^2 - 6*(a*b^2*c*d^2*e^2 - b^3*c*d^2*e)*f)*tan(f*x + e))*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 2*(2*a^2*b*d^3*f^3 - 2*a^2*b*c^3*f^3 - 3*a*b^2*d^3*e^2 + 3*(2*a^2*b*c^2*d*f^3 - b^3*c^2*d)*f^2 - 6*(a*b^2*c*d^2*e^2 - b^3*c*d^2*e)*f)*tan(f*x + e))*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - 3*(I*a*b^2*d^3*tan(f*x + e) + I*a^2*b*d^3)*polylog(4, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*(-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - 3*(-I*a*b^2*d^3*tan(f*x + e) - I*a^2*b*d^3)*polylog(4, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 3*(2*a^2*b*d^3*f*x + 2*a^2*b*c*d^2*f + a*b^2*d^3 + (2*a*b^2*d^3*f*x + 2*a*b^2*c*d^2*f + b^3*d^3)*tan(f*x + e))*polylog(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 - 2*I*a*b + b^2 - 2*$

$$\begin{aligned}
& (-I*a^2 + 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) \\
& + 3*(2*a^2*b*d^3*f*x + 2*a^2*b*c*d^2*f + a*b^2*d^3 + (2*a*b^2*d^3*f*x \\
& + 2*a*b^2*c*d^2*f + b^3*d^3)*tan(f*x + e))*polylog(3, ((a^2 - 2*I*a*b - b^2)*tan(f*x + e)^2 - a^2 + 2*I*a*b + b^2 - 2*(I*a^2 + 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + ((a^2*b - b^3)*d^3*f^4*x^4 + 4*a*b^2*c^3*f^3 + 4*(a*b^2*d^3*f^3 + (a^2*b - b^3)*c*d^2*f^4)*x^3 + 6*(2*a*b^2*c*d^2*f^3 + (a^2*b - b^3)*c^2*d*f^4)*x^2 + 4*(3*a*b^2*c^2*d*f^3 + (a^2*b - b^3)*c^3*f^4)*x)*tan(f*x + e))/((a^4*b + 2*a^2*b^3 + b^5)*f^4*tan(f*x + e) + (a^5 + 2*a^3*b^2 + a*b^4)*f^4)
\end{aligned}$$

Sympy [F]

$$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx = \int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx$$

[In] `integrate((d*x+c)**3/(a+b*tan(f*x+e))**2,x)`
[Out] `Integral((c + d*x)**3/(a + b*tan(e + f*x))**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4631 vs. $2(695) = 1390$.

Time = 2.41 (sec), antiderivative size = 4631, normalized size of antiderivative = 5.46

$$\int \frac{(c+dx)^3}{(a+b\tan(e+fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="maxima")`
[Out]
$$\begin{aligned}
& -1/12*(36*c^2*d*e*(2*a*b*log(b*tan(f*x + e) + a)/((a^4 + 2*a^2*b^2 + b^4)*f) \\
& - a*b*log(tan(f*x + e)^2 + 1)/((a^4 + 2*a^2*b^2 + b^4)*f) - b/((a^2*b + b^3)*f*tan(f*x + e) + (a^3 + a*b^2)*f) + (a^2 - b^2)*(f*x + e)/((a^4 + 2*a^2*b^2 + b^4)*f)) - 12*(2*a*b*log(b*tan(f*x + e) + a)/(a^4 + 2*a^2*b^2 + b^4) \\
& - a*b*log(tan(f*x + e)^2 + 1)/(a^4 + 2*a^2*b^2 + b^4) + (a^2 - b^2)*(f*x + e)/(a^4 + 2*a^2*b^2 + b^4) - b/(a^3 + a*b^2 + (a^2*b + b^3)*tan(f*x + e))) \\
& *c^3 - (3*(a^3 - I*a^2*b + a*b^2 - I*b^3)*(f*x + e)^4*d^3 - 24*(I*a*b^2 - b^3)*d^3*e^3 - 72*(-I*a*b^2 + b^3)*c*d^2*e^2*f - 12*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e - (a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*f)*(f*x + e)^3 + 18*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^2 - 2*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*e*f + (a^3 - I*a^2*b + a*b^2 - I*b^3)*c^2*d*f^2)*(f*x + e)^2 - 12*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d^3*e^3 - 3*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*d^2*e^2*f)*(f*x + e) - 12*(2*(I*a^2*b - a*b^2)*d^3*e^3 + 3*(-I*a*b^2 + b^3)*c*d^2*e^2*f)*(f*x + e))
\end{aligned}$$

$$\begin{aligned}
& \sim 3)*d^3*e^2 + 3*(-I*a*b^2 + b^3)*c^2*d*f^2 + 6*((-I*a^2*b + a*b^2)*c*d^2*e^2 \\
& + (I*a*b^2 - b^3)*c*d^2*e)*f + (2*(I*a^2*b + a*b^2)*d^3*e^3 + 3*(-I*a*b^2 \\
& - b^3)*d^3*e^2 + 3*(-I*a*b^2 - b^3)*c^2*d*f^2 + 6*((-I*a^2*b - a*b^2)*c*d^2 \\
& *e^2 + (I*a*b^2 + b^3)*c*d^2*e)*f)*cos(2*f*x + 2*e) - (2*(a^2*b - I*a*b^2) \\
& *d^3*e^3 - 3*(a*b^2 - I*b^3)*c^2*d*f^2 - 6*((a^2*b - I*a*b^2)*c*d^2*e^2 \\
& - (a*b^2 - I*b^3)*c*d^2*e)*f)*sin(2*f*x + 2*e))*arctan2(-b*cos(2*f*x + 2*e) \\
& + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b* \\
& sin(2*f*x + 2*e) + a) - 4*(8*(I*a^2*b - a*b^2)*(f*x + e)^3*d^3 + 9*(2*(-I*a \\
& ^2*b + a*b^2)*d^3*e + 2*(I*a^2*b - a*b^2)*c*d^2*f + (I*a*b^2 - b^3)*d^3)*(f \\
& *x + e)^2 + 18*((I*a^2*b - a*b^2)*d^3*e^2 + (I*a^2*b - a*b^2)*c^2*d*f^2 + \\
& (-I*a*b^2 + b^3)*d^3*e + (2*(-I*a^2*b + a*b^2)*c*d^2*e + (I*a*b^2 - b^3)*c*d \\
& ^2)*f)*(f*x + e) + (8*(I*a^2*b + a*b^2)*(f*x + e)^3*d^3 + 9*(2*(-I*a^2*b - \\
& a*b^2)*d^3*e + 2*(I*a^2*b + a*b^2)*c*d^2*f + (I*a*b^2 + b^3)*d^3)*(f*x + e) \\
& ^2 + 18*((I*a^2*b + a*b^2)*d^3*e^2 + (I*a^2*b + a*b^2)*c^2*d*f^2 + (-I*a*b^ \\
& 2 - b^3)*d^3*e + (2*(-I*a^2*b - a*b^2)*c*d^2*e + (I*a*b^2 + b^3)*c*d^2)*f) \\
& *(f*x + e))*cos(2*f*x + 2*e) - (8*(a^2*b - I*a*b^2)*(f*x + e)^3*d^3 - 9*(2*(\\
& a^2*b - I*a*b^2)*d^3*e - 2*(a^2*b - I*a*b^2)*c*d^2*f - (a*b^2 - I*b^3)*d^3) \\
& *(f*x + e)^2 + 18*((a^2*b - I*a*b^2)*d^3*e^2 + (a^2*b - I*a*b^2)*c^2*d*f^2 \\
& - (a*b^2 - I*b^3)*d^3*e - (2*(a^2*b - I*a*b^2)*c*d^2*e - (a*b^2 - I*b^3)*c \\
& d^2)*f)*(f*x + e))*sin(2*f*x + 2*e))*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 \\
& - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 \\
& + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + 3*((a^3 - 3*I*a^2*b - 3*a*b^ \\
& 2 + I*b^3)*(f*x + e)^4*d^3 - 4*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d^3*e - \\
& (a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*c*d^2*f + 2*(I*a*b^2 + b^3)*d^3)*(f*x \\
& + e)^3 + 6*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d^3*e^2 + (a^3 - 3*I*a^2*b \\
& - 3*a*b^2 + I*b^3)*c^2*d*f^2 - 4*(-I*a*b^2 - b^3)*d^3*e - 2*((a^3 - 3*I*a^2 \\
& *b - 3*a*b^2 + I*b^3)*c*d^2*f + 2*(I*a*b^2 + b^3)*c*d^2)*f)*(f*x + e)^2 - 4 \\
& *((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d^3*e^3 + 6*(I*a*b^2 + b^3)*d^3*e^2 + \\
& 6*(I*a*b^2 + b^3)*c^2*d*f^2 - 3*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*c*d^2 \\
& *e^2 - 4*(-I*a*b^2 - b^3)*c*d^2)*f)*(f*x + e))*cos(2*f*x + 2*e) - 12*(4*(\\
& I*a^2*b - a*b^2)*(f*x + e)^2*d^3 + 3*(I*a^2*b - a*b^2)*d^3*e^2 + 3*(I*a^2*b \\
& - a*b^2)*c^2*d*f^2 + 3*(-I*a*b^2 + b^3)*d^3*e + 3*(2*(-I*a^2*b + a*b^2)*d \\
& 3*e + 2*(I*a^2*b - a*b^2)*c*d^2*f + (I*a*b^2 - b^3)*d^3)*(f*x + e) + 3*(2*(\\
& -I*a^2*b + a*b^2)*c*d^2*e + (I*a*b^2 - b^3)*c*d^2)*f + (4*(I*a^2*b + a*b^2) \\
& *(f*x + e)^2*d^3 + 3*(I*a^2*b + a*b^2)*d^3*e^2 + 3*(I*a^2*b + a*b^2)*c^2*d \\
& f^2 + 3*(-I*a*b^2 - b^3)*d^3*e + 3*(2*(-I*a^2*b - a*b^2)*d^3*e + 2*(I*a^2*b \\
& + a*b^2)*c*d^2*f + (I*a*b^2 + b^3)*d^3)*(f*x + e) + 3*(2*(-I*a^2*b - a*b^2) \\
&)*c*d^2*e + (I*a*b^2 + b^3)*c*d^2)*f)*cos(2*f*x + 2*e) - (4*(a^2*b - I*a*b^ \\
& 2)*(f*x + e)^2*d^3 + 3*(a^2*b - I*a*b^2)*d^3*e^2 + 3*(a^2*b - I*a*b^2)*c^2 \\
& d*f^2 - 3*(a*b^2 - I*b^3)*d^3*e - 3*(2*(a^2*b - I*a*b^2)*d^3*e - 2*(a^2*b - \\
& I*a*b^2)*c*d^2*f - (a*b^2 - I*b^3)*d^3)*(f*x + e) - 3*(2*(a^2*b - I*a*b^2) \\
&)*c*d^2*e - (a*b^2 - I*b^3)*c*d^2)*f)*sin(2*f*x + 2*e))*dilog((I*a + b)*e^(2 \\
& *I*f*x + 2*I*e)/(-I*a + b)) - 6*(2*(a^2*b + I*a*b^2)*d^3*e^3 - 3*(a*b^2 + I \\
& *b^3)*d^3*e^2 - 3*(a*b^2 + I*b^3)*c^2*d*f^2 - 6*((a^2*b + I*a*b^2)*c*d^2*e^ \\
& 2 - (a*b^2 + I*b^3)*c*d^2)*f + (2*(a^2*b - I*a*b^2)*d^3*e^3 - 3*(a*b^2 -
\end{aligned}$$

$$\begin{aligned}
& I*b^3)*d^3*e^2 - 3*(a*b^2 - I*b^3)*c^2*d*f^2 - 6*((a^2*b - I*a*b^2)*c*d^2*e \\
& ^2 - (a*b^2 - I*b^3)*c*d^2*e)*f)*cos(2*f*x + 2*e) + (2*(I*a^2*b + a*b^2)*d^ \\
& 3*e^3 + 3*(-I*a*b^2 - b^3)*d^3*e^2 + 3*(-I*a*b^2 - b^3)*c^2*d*f^2 + 6*((-I* \\
& a^2*b - a*b^2)*c*d^2*e^2 + (I*a*b^2 + b^3)*c*d^2*e)*f)*sin(2*f*x + 2*e))*lo \\
& g((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin \\
& (2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*(8*(a^2*b \\
& + I*a*b^2)*(f*x + e)^3*d^3 - 9*(2*(a^2*b + I*a*b^2)*d^3*e - 2*(a^2*b + I*a \\
& *b^2)*c*d^2*f - (a*b^2 + I*b^3)*d^3)*(f*x + e)^2 + 18*((a^2*b + I*a*b^2)*d^ \\
& 3*e^2 + (a^2*b + I*a*b^2)*c^2*d*f^2 - (a*b^2 + I*b^3)*d^3*e - (2*(a^2*b + I \\
& *a*b^2)*c*d^2*e - (a*b^2 + I*b^3)*c*d^2)*f)*(f*x + e) + (8*(a^2*b - I*a*b^2) \\
&)*(f*x + e)^3*d^3 - 9*(2*(a^2*b - I*a*b^2)*d^3*e - 2*(a^2*b - I*a*b^2)*c*d^ \\
& 2*f - (a*b^2 - I*b^3)*d^3)*(f*x + e)^2 + 18*((a^2*b - I*a*b^2)*d^3*e^2 + (a \\
& ^2*b - I*a*b^2)*c^2*d*f^2 - (a*b^2 - I*b^3)*d^3*e - (2*(a^2*b - I*a*b^2)*c* \\
& d^2*e - (a*b^2 - I*b^3)*c*d^2)*f)*(f*x + e))*cos(2*f*x + 2*e) - (8*(-I*a^2*b \\
& - a*b^2)*(f*x + e)^3*d^3 + 9*(2*(I*a^2*b + a*b^2)*d^3*e + 2*(-I*a^2*b - a \\
& *b^2)*c*d^2*f + (-I*a*b^2 - b^3)*d^3)*(f*x + e)^2 + 18*((-I*a^2*b - a*b^2)* \\
& d^3*e^2 + (-I*a^2*b - a*b^2)*c^2*d*f^2 + (I*a*b^2 + b^3)*d^3*e + (2*(I*a^2*b \\
& + a*b^2)*c*d^2*e + (-I*a*b^2 - b^3)*c*d^2)*f)*(f*x + e))*sin(2*f*x + 2*e) \\
&)*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2) \\
&)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b \\
& ^2) - 24*((-I*a^2*b - a*b^2)*d^3*cos(2*f*x + 2*e) + (a^2*b - I*a*b^2)*d^3* \\
& sin(2*f*x + 2*e) + (-I*a^2*b + a*b^2)*d^3)*polylog(4, (I*a + b)*e^(2*I*f*x \\
& + 2*I*e)/(-I*a + b)) + 6*(8*(a^2*b + I*a*b^2)*(f*x + e)*d^3 - 6*(a^2*b + I* \\
& a*b^2)*d^3*e + 6*(a^2*b + I*a*b^2)*c*d^2*f + 3*(a*b^2 + I*b^3)*d^3 + (8*(a^ \\
& 2*b - I*a*b^2)*(f*x + e)*d^3 - 6*(a^2*b - I*a*b^2)*d^3*e + 6*(a^2*b - I*a*b \\
& ^2)*c*d^2*f + 3*(a*b^2 - I*b^3)*d^3)*cos(2*f*x + 2*e) - (8*(-I*a^2*b - a*b^ \\
& 2)*(f*x + e)*d^3 + 6*(I*a^2*b + a*b^2)*d^3*e + 6*(-I*a^2*b - a*b^2)*c*d^2*f \\
& + 3*(-I*a*b^2 - b^3)*d^3)*sin(2*f*x + 2*e))*polylog(3, (I*a + b)*e^(2*I*f*x \\
& + 2*I*e)/(-I*a + b)) - 3*((-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3)*(f*x + e)^ \\
& 4*d^3 + 4*((I*a^3 + 3*a^2*b - 3*I*a*b^2 - b^3)*d^3*e + (-I*a^3 - 3*a^2*b + \\
& 3*I*a*b^2 + b^3)*c*d^2*f - 2*(a*b^2 - I*b^3)*d^3)*(f*x + e)^3 + 6*((-I*a^3 \\
& - 3*a^2*b + 3*I*a*b^2 + b^3)*d^3*e^2 + (-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3) \\
& *c^2*d*f^2 + 4*(a*b^2 - I*b^3)*d^3*e + 2*((I*a^3 + 3*a^2*b - 3*I*a*b^2 - b^ \\
& 3)*c*d^2*f^2 - 2*(a*b^2 - I*b^3)*c*d^2)*f)*(f*x + e)^2 + 4*((I*a^3 + 3*a^2*b \\
& - 3*I*a*b^2 - b^3)*d^3*e^3 - 6*(a*b^2 - I*b^3)*d^3*e^2 - 6*(a*b^2 - I*b^3)* \\
& c^2*d*f^2 + 3*((-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3)*c*d^2*f^2 + 4*(a*b^2 - \\
& I*b^3)*c*d^2*f)*(f*x + e))*sin(2*f*x + 2*e))/((a^5 - I*a^4*b + 2*a^3*b^2 \\
& - 2*I*a^2*b^3 + a*b^4 - I*b^5)*f^3*cos(2*f*x + 2*e) - (-I*a^5 - a^4*b - 2* \\
& I*a^3*b^2 - 2*a^2*b^3 - I*a*b^4 - b^5)*f^3*sin(2*f*x + 2*e) + (a^5 + I*a^4*b \\
& + 2*a^3*b^2 + 2*I*a^2*b^3 + a*b^4 + I*b^5)*f^3)/f
\end{aligned}$$

Giac [F]

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(dx + c)^3}{(b \tan(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)^3/(a+b*tan(f*x+e))^2,x, algorithm="giac")
[Out] integrate((d*x + c)^3/(b*tan(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b \tan(e + fx))^2} dx$$

[In] int((c + d*x)^3/(a + b*tan(e + f*x))^2,x)
[Out] int((c + d*x)^3/(a + b*tan(e + f*x))^2, x)

3.60 $\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$

Optimal result	404
Rubi [A] (verified)	405
Mathematica [A] (verified)	410
Maple [B] (verified)	411
Fricas [B] (verification not implemented)	413
Sympy [F]	414
Maxima [B] (verification not implemented)	414
Giac [F]	415
Mupad [F(-1)]	416

Optimal result

Integrand size = 20, antiderivative size = 654

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = & -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a+ib)(ia+b)^2 (ia-b+(ia+b)e^{2ie+2ifx}) f} \\ & + \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} \\ & + \frac{2b^2d(c+dx)\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\ & + \frac{2b(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\ & - \frac{2ib^2(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\ & - \frac{ib^2d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \\ & + \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\ & - \frac{2b^2d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\ & - \frac{ib^2d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} \end{aligned}$$

[Out] $-2*I*b^2*(d*x+c)^2/(a^2+b^2)^2/f+2*b^2*(d*x+c)^2/(a+I*b)/(I*a+b)^2/(I*a-b+(I*a+b)*exp(2*I*e+2*I*f*x))/f+1/3*(d*x+c)^3/(a-I*b)^2/d+4/3*b*(d*x+c)^3/(I*a-b)/(a-I*b)^2/d-4/3*b^2*(d*x+c)^3/(a^2+b^2)^2/d+2*b^2*d*(d*x+c)*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+2*b*(d*x+c)^2*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f-2*I*b^2*(d*x+c)^2*ln(1+(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f-I*b^2*d^2*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3+2*b*d*(d*x+c)*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(I*a-b)/(a-I*b)^2/f^2-2*b^2*d*(d*x+c)*polylog(2,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^2+2*b*d^2*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a-I*b)^2/(a+I*b)/f^3-I*b^2*d^2*polylog(3,-(a-I*b)*exp(2*I*e+2*I*f*x)/(a+I*b))/(a^2+b^2)^2/f^3$

Rubi [A] (verified)

Time = 1.60 (sec) , antiderivative size = 654, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules used = {3815, 2216, 2215, 2221, 2611, 2320, 6724, 2222, 2317, 2438}

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = & -\frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2 (a^2+b^2)^2} \\ & + \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2 (a^2+b^2)^2} \\ & - \frac{2ib^2 (c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f (a^2+b^2)^2} - \frac{2ib^2 (c+dx)^2}{f (a^2+b^2)^2} \\ & - \frac{4b^2 (c+dx)^3}{3d (a^2+b^2)^2} - \frac{ib^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3 (a^2+b^2)^2} \\ & - \frac{ib^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3 (a^2+b^2)^2} \\ & + \frac{2b^2 (c+dx)^2}{f(a+ib)(b+ia)^2 ((b+ia)e^{2ie+2ifx} + ia - b)} \\ & + \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^2 (-b+ia)(a-ib)^2} \\ & + \frac{2b(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f(a-ib)^2 (a+ib)} + \frac{4b(c+dx)^3}{3d(-b+ia)(a-ib)^2} \\ & + \frac{(c+dx)^3}{3d(a-ib)^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{f^3 (a-ib)^2 (a+ib)} \end{aligned}$$

[In] $\operatorname{Int}[(c+d*x)^2/(a+b*\operatorname{Tan}[e+f*x])^2, x]$

[Out] $\frac{((-2*I)*b^2*(c + d*x)^2)/((a^2 + b^2)^2*f) + (2*b^2*(c + d*x)^2)/((a + I*b)*(I*a + b)^2*(I*a - b + (I*a + b)*E^{((2*I)*e + (2*I)*f*x))*f) + (c + d*x)^3/(3*(a - I*b)^2*d) + (4*b*(c + d*x)^3)/(3*(I*a - b)*(a - I*b)^2*d) - (4*b^2*(c + d*x)^3)/(3*(a^2 + b^2)^2*d) + (2*b^2*d*(c + d*x)*Log[1 + ((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])/((a^2 + b^2)^2*f^2) + (2*b*(c + d*x)^2*Log[1 + ((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])/((a - I*b)^2*(a + I*b)*f) - ((2*I)*b^2*(c + d*x)^2*Log[1 + ((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])/((a^2 + b^2)^2*f) - (I*b^2*d^2*PolyLog[2, -((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])]/((a^2 + b^2)^2*f^3) + (2*b*d*(c + d*x)*PolyLog[2, -((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])]/((I*a - b)*(a - I*b)^2*f^2) - (2*b^2*d*(c + d*x)*PolyLog[2, -((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])]/((a^2 + b^2)^2*f^2) + (b*d^2*PolyLog[3, -((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])]/((a - I*b)^2*(a + I*b)*f^3) - (I*b^2*d^2*PolyLog[3, -((a - I*b)*E^{((2*I)*e + (2*I)*f*x)})/(a + I*b)])]/((a^2 + b^2)^2*f^3)$

Rule 2215

```
Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Dist[b/a, Int[(c + d*x)^m*((F^(g*(e + f*x)))^n/(a + b*(F^(g*(e + f*x)))^n)), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2216

```
Int[((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.))^p*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] :> Dist[1/a, Int[(c + d*x)^m*(a + b*(F^(g*(e + f*x)))^n)^(p + 1), x], x] - Dist[b/a, Int[(c + d*x)^m*(F^(g*(e + f*x)))^n*(a + b*(F^(g*(e + f*x)))^n)^p, x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && ILtQ[p, 0] && IGtQ[m, 0]
```

Rule 2221

```
Int[((((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2222

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.)*((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_.))))^(n_.))^p*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((a + b*(F^(g*(e + f*x)))^n)^p + 1)/(b*f*g*n*(p + 1)*Log[F]), x] - Dist[d*(m/(b*f*g*n*(p + 1)*Log[F))), Int[(c + d*x)^(m - 1)*(a + b*(F^(g*(e + f*x)))^n)^p + 1, x], x] /; FreeQ[{F, a, b, c, d, e, f, g, m}
```

```
, n, p}, x] && NeQ[p, -1]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*(c_.) + (d_)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x],
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_)*x))*(F_[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[((d_) + (e_)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2,
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^m_, x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m -
1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3815

```
Int[((c_.) + (d_)*(x_)^m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 +
b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^(-n), x], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_)*(x_))^p_]/((d_.) + (e_)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^2}{(a-ib)^2} - \frac{4b^2(c+dx)^2}{(ia+b)^2 (ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx})^2} \right. \\
&\quad \left. + \frac{4b(c+dx)^2}{(a-ib)^2 (ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx})} \right) dx \\
&= \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{(4b) \int \frac{(c+dx)^2}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{(a-ib)^2} - \frac{(4b^2) \int \frac{(c+dx)^2}{(ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx})^2} dx}{(ia+b)^2} \\
&= \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(i a - b)(a-ib)^2 d} + \frac{(4b^2) \int \frac{(c+dx)^2}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{(ia-b)(a-ib)^2} \\
&\quad - \frac{(4b) \int \frac{e^{2ie+2ifx}(c+dx)^2}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{a^2 + b^2} - \frac{(4b^2) \int \frac{e^{2ie+2ifx}(c+dx)^2}{(ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx})^2} dx}{a^2 + b^2} \\
&= -\frac{2b^2(c+dx)^2}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(i a - b)(a-ib)^2 d} \\
&\quad - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} + \frac{2b(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{(4b^2) \int \frac{e^{2ie+2ifx}(c+dx)^2}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{(a+ib)^2(ia+b)} \\
&\quad - \frac{(4bd) \int (c+dx) \log\left(1 + \frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a-ib)^2(a+ib)f} + \frac{(4b^2d) \int \frac{c+dx}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{(a-ib)^2(a+ib)f} \\
&= -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^2}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^3}{3(a-ib)^2 d} \\
&\quad + \frac{4b(c+dx)^3}{3(i a - b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} + \frac{2b(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\
&\quad - \frac{2ib^2(c+dx)^2 \log\left(1 + \frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} + \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{(2bd^2) \int \text{PolyLog}\left(2, -\frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(ia-b)(a-ib)^2 f^2} - \frac{(4b^2d) \int \frac{e^{2ie+2ifx}(c+dx)}{ia(1+\frac{ib}{a}) + ia(1-\frac{ib}{a}) e^{2ie+2ifx}} dx}{(a-ib)(a+ib)^2 f} \\
&\quad + \frac{(4ib^2d) \int (c+dx) \log\left(1 + \frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^2}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} \\
&\quad + \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} \\
&\quad + \frac{2b^2d(c+dx)\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} + \frac{2b(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\
&\quad - \frac{2ib^2(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} + \frac{2bd(c+dx)\text{PolyLog}\left(2,-\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{2b^2d(c+dx)\text{PolyLog}\left(2,-\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{(bd^2)\text{Subst}\left(\int \frac{\text{PolyLog}\left(2,-\frac{(a-ib)x}{a+ib}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad - \frac{(2b^2d^2)\int \log\left(1+\frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{(2b^2d^2)\int \text{PolyLog}\left(2,-\frac{(1-\frac{ib}{a})e^{2ie+2ifx}}{1+\frac{ib}{a}}\right) dx}{(a^2+b^2)^2 f^2} \\
&= -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^2}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} \\
&\quad + \frac{(c+dx)^3}{3(a-ib)^2 d} + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} \\
&\quad + \frac{2b^2d(c+dx)\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} + \frac{2b(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} \\
&\quad - \frac{2ib^2(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} + \frac{2bd(c+dx)\text{PolyLog}\left(2,-\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{2b^2d(c+dx)\text{PolyLog}\left(2,-\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} + \frac{bd^2\text{PolyLog}\left(3,-\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} \\
&\quad + \frac{(ib^2d^2)\text{Subst}\left(\int \frac{\log\left(1+\frac{(1-\frac{ib}{a})x}{1+\frac{ib}{a}}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{(a^2+b^2)^2 f^3} \\
&\quad - \frac{(ib^2d^2)\text{Subst}\left(\int \frac{\text{PolyLog}\left(2,-\frac{(a-ib)x}{a+ib}\right)}{x} dx, x, e^{2ie+2ifx}\right)}{(a^2+b^2)^2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2ib^2(c+dx)^2}{(a^2+b^2)^2 f} - \frac{2b^2(c+dx)^2}{(a-ib)^2(a+ib)(ia-b+(ia+b)e^{2ie+2ifx})f} + \frac{(c+dx)^3}{3(a-ib)^2 d} \\
&\quad + \frac{4b(c+dx)^3}{3(ia-b)(a-ib)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2+b^2)^2 d} + \frac{2b^2d(c+dx)\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{2b(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f} - \frac{2ib^2(c+dx)^2\log\left(1+\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f} \\
&\quad - \frac{ib^2d^2 \text{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3} + \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(ia-b)(a-ib)^2 f^2} \\
&\quad - \frac{2b^2d(c+dx) \text{PolyLog}\left(2, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^2} \\
&\quad + \frac{bd^2 \text{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a-ib)^2(a+ib)f^3} - \frac{ib^2d^2 \text{PolyLog}\left(3, -\frac{(a-ib)e^{2ie+2ifx}}{a+ib}\right)}{(a^2+b^2)^2 f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.53 (sec), antiderivative size = 761, normalized size of antiderivative = 1.16

$$\begin{aligned}
&\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx \\
&= \frac{2b \left(\frac{12c(bd+acf)x}{a-ib} - \frac{12c(-ib(-1+e^{2ie})+a(1+e^{2ie}))}{a^2+b^2}(bd+acf)x + \frac{6d(bd+2acf)x^2}{a-ib} + \frac{4ad^2fx^3}{a-ib} + \frac{6d(-ib(-1+e^{2ie})+a(1+e^{2ie}))(bd+2acf)x \log\left(1+\frac{(a+ib)e^{-2ie}}{a-ib}\right)}{(a+ib)(ia+b)f} \right)}{(a-ib)^2(a+ib)f^3}
\end{aligned}$$

[In] Integrate[(c + d*x)^2/(a + b*Tan[e + f*x])^2, x]

[Out] $((2*b*((12*c*(b*d + a*c*f)*x)/(a - I*b) - (12*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*x)/(a^2 + b^2) + (6*d*(b*d + 2*a*c*f)*x^2)/(a - I*b) + (4*a*d^2*f*x^3)/(a - I*b) + (6*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*x*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a + I*b)*(I*a + b)*f) + (6*a*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*x^2*Log[1 + (a + I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a + I*b)*(I*a + b)) + (6*c*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + a*c*f)*Log[a + I*b + (a - I*b)*E^((2*I)*(e + f*x))])/((a + I*b)*(I*a + b)*f) + (3*d*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(b*d + 2*a*c*f)*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))])/((a^2 + b^2)*f^2) + (3*a*d^2*((-I)*b*(-1 + E^((2*I)*e)) + a*(1 + E^((2*I)*e)))*(2*f*x*PolyLog[2, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))] - I*PolyLog[3, (-a - I*b)/((a - I*b)*E^((2*I)*(e + f*x)))]))/((a^2 + b^2)*f^2))/(b - b*E^((2*I)*e) - I*a*(1 + E^((2*I)*e)) + ((a^2 - b^2)*f*x*(3*c^2 + 3*c$

$$\begin{aligned} & *d*x + d^2*x^2)*Cos[f*x] + (a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cos[\\ & 2*e + f*x] + 2*b*(3*b*(c + d*x)^2 + a*f*x*(3*c^2 + 3*c*d*x + d^2*x^2))*Sin[\\ & f*x])/((a*Cos[e] + b*Sin[e])*(a*Cos[e + f*x] + b*Sin[e + f*x]))/(6*(a^2 + \\ & b^2)*f) \end{aligned}$$

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3900 vs. $2(588) = 1176$.

Time = 0.97 (sec), antiderivative size = 3901, normalized size of antiderivative = 5.96

method	result	size
risch	Expression too large to display	3901

```
[In] int((d*x+c)^2/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
[Out] 2/(I*a+b)^2/f^2/(b-I*a)*b^2*e*d*a*c/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))
+b^2*exp(4*I*(f*x+e))+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2
)+I/(I*a+b)^2/f^3/(b-I*a)*b^2*e*d^2/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))
+b^2*exp(4*I*(f*x+e))+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2
)*a+4/(I*a+b)^2/f^2/(b-I*a)*b*e*d*a^2*c/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*
I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-4/(I*a+b)^2/f^2/(b-I
*a)*b*e*d*a^2*c/(a+I*b)/(I*b-a)*arctan(1/b*a)+2/(I*a+b)^2/f^2/(b-I*a)*b^2*c
*d/(a+I*b)/(I*b-a)*arctan(1/b*a)*a-2/(I*a+b)^2/f^2/(b-I*a)*b^2*c*d/(a+I*b)/
(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/
2/b*a)*a-2/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a^2*d^2/(a+I*b)/(I*b-a)*arctan(1/2/a
*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)-4*I/(I*a+b)^2
/f^3/(b-I*a)*b*e^2*a*d^2/(a+I*b)*ln(exp(I*(f*x+e)))+2*I/(I*a+b)^2/f/(b-I*a
)*b^2*a*c^2/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*
exp(2*I*(f*x+e))+1/2/b*a)-2*I/(I*a+b)^2/f/(b-I*a)*b^2*a*c^2/(a+I*b)/(I*b-a)
*arctan(1/b*a)-2*I/(I*a+b)^2/f^3/(b-I*a)*b/(a+I*b)*e^2*a*d^2*ln(1-(I*b-a)*e
xp(2*I*(f*x+e))/(a+I*b))-2*I/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^2/(a+I*b)/(I*b-a)
)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)
+2*I/(I*a+b)^2/f^3/(b-I*a)*b^3*e*d^2/(a+I*b)/(I*b-a)*arctan(1/b*a)+2*I/(I*a
+b)^2/f/(b-I*a)*b/(a+I*b)*d^2*a*ln(1-(I*b-a)*exp(2*I*(f*x+e))/(a+I*b))*x^2-
2*I/(I*a+b)^2/f^2/(b-I*a)*b^3*c*d/(a+I*b)/(I*b-a)*arctan(1/b*a)-I/(I*a+b)^2
/f/(b-I*a)*b*a^2*c^2/(a+I*b)/(I*b-a)*ln(a^2*exp(4*I*(f*x+e))+b^2*exp(4*I*(f
*x+e))+2*a^2*exp(2*I*(f*x+e))-2*b^2*exp(2*I*(f*x+e))+a^2+b^2)+2*I/(I*a+b)^2
/f^2/(b-I*a)*b^3*c*d/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x+e))*b-1/2*b/
a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)+2/(I*a+b)^2/f^3/(b-I*a)*b*e^2*a^2*d^2/(a
+I*b)/(I*b-a)*arctan(1/b*a)-1/(I*a+b)^2/f^3/(b-I*a)*b^2*e^2*a*d^2/(a+I*b)/
(I*b-a)*ln(a^2*exp(4*I*(f*x+e))+b^2*exp(4*I*(f*x+e))+2*a^2*exp(2*I*(f*x+e))
-2*b^2*exp(2*I*(f*x+e))+a^2+b^2)+8/(I*a+b)^2/f/(b-I*a)*b/(a+I*b)*d*a*c*e*x+
2/(I*a+b)^2/f^3/(b-I*a)*b^2*e*d^2/(a+I*b)/(I*b-a)*arctan(1/2/a*exp(2*I*(f*x
+e))*b-1/2*b/a+1/2/b*a*exp(2*I*(f*x+e))+1/2/b*a)*a-1/3*d^2/(2*I*a*b-a^2+b^2
```

$$\begin{aligned}
&) * x^3 - 1 / (2 * I * a * b - a^2 + b^2) * c^2 * x - 1 / 3 / d / (2 * I * a * b - a^2 + b^2) * c * x^2 + 2 * I / (I * a + b)^2 / f^2 / (b - I * a) * b * e * d * a^2 * c / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 - 4 * I / (I * a + b)^2 / f^2 / (b - I * a) * b^2 * e * d * a * c / (a + I * b) / (I * b - a) * \arctan(1 / 2 / a * \exp(2 * I * (f * x + e))) * b - 1 / 2 * b / a + 1 / 2 / b * a * \exp(2 * I * (f * x + e)) + 1 / 2 / b * a + 4 * I / (I * a + b)^2 / f^2 / (b - I * a) * b^2 * e * d * a * c / (a + I * b) / (I * b - a) * \arctan(1 / b * a) - 8 / 3 / (I * a + b)^2 / f^3 / (b - I * a) * b / (a + I * b) * e^3 * a * d^2 + 4 / (I * a + b)^2 / f^2 / (b - I * a) * b^2 / (a + I * b) * d^2 * e * x + 4 / (I * a + b)^2 / (b - I * a) * b / (a + I * b) * d * a * c * x^2 + 4 * I / (I * a + b)^2 / f^3 / (b - I * a) * b^2 * e * d^2 - 2 / (a + I * b) * \ln(\exp(I * (f * x + e))) - 2 / (I * a + b)^2 / f^3 / (b - I * a) * b^2 * e * d^2 / (a + I * b) / (I * b - a) * \arctan(1 / b * a) * a + 2 / (I * a + b)^2 / f^2 / (b - I * a) * b / (a + I * b) * d * a * c * \text{polylog}(2, (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) + 2 / (I * a + b)^2 / f^2 / (b - I * a) * b / (a + I * b) * d^2 * a * \text{polylog}(2, (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) * x - 1 / (I * a + b)^2 / f^2 / (b - I * a) * b^3 * c * d / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 - 2 / (I * a + b)^2 / f / (b - I * a) * b * a^2 * c^2 / (a + I * b) / (I * b - a) * \arctan(1 / 2 / a * \exp(2 * I * (f * x + e))) * b - 1 / 2 * b / a + 1 / 2 / b * a * \exp(2 * I * (f * x + e)) + 1 / 2 / b * a + 2 / (I * a + b)^2 / f / (b - I * a) * b * a^2 * c^2 / (a + I * b) / (I * b - a) * \arctan(1 / b * a) - 4 / (I * a + b)^2 / f^2 / (b - I * a) * b / (a + I * b) * e^2 * a * d^2 * x + I / (I * a + b)^2 / f^3 / (b - I * a) * b / (a + I * b) * d^2 * a * \text{polylog}(3, (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) - 1 / (I * a + b)^2 / f / (b - I * a) * b^2 * a * c^2 / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 + 1 / (I * a + b)^2 / f^3 / (b - I * a) * b^3 * e * d^2 / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 - 4 * I / (I * a + b)^2 / f^2 / (b - I * a) * b^2 * c * d / (a + I * b) * \ln(\exp(I * (f * x + e))) - 4 * I / (I * a + b)^2 / f / (b - I * a) * b * a * c^2 / (a + I * b) * \ln(\exp(I * (f * x + e))) + 2 * I / (I * a + b)^2 / f^2 / (b - I * a) * b^2 / (a + I * b) * d^2 * \ln(1 - (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) * x + 2 * I / (I * a + b)^2 / f^3 / (b - I * a) * b^2 / (a + I * b) * d^2 * \ln(1 - (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) * e + 4 / (I * a + b)^2 / f^2 / (b - I * a) * b / (a + I * b) * d * a * c * e^2 - I / (I * a + b)^2 / f^2 / (b - I * a) * b * a^2 * c^2 / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 - I / (I * a + b)^2 / f^3 / (b - I * a) * b * e^2 * a^2 * d^2 / (a + I * b) / (I * b - a) * \ln(a^2 * \exp(4 * I * (f * x + e))) + b^2 * \exp(4 * I * (f * x + e)) + 2 * a^2 * \exp(2 * I * (f * x + e)) - 2 * b^2 * \exp(2 * I * (f * x + e)) + a^2 + b^2 + 4 * I / (I * a + b)^2 / f / (b - I * a) * b / (a + I * b) * d * a * c * \ln(1 - (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) * x + 2 * I / (I * a + b)^2 / f^3 / (b - I * a) * b^2 * e^2 * a * d^2 / (a + I * b) / (I * b - a) * \arctan(1 / 2 / a * \exp(2 * I * (f * x + e))) * b - 1 / 2 * b / a + 1 / 2 / b * a * \exp(2 * I * (f * x + e)) + 1 / 2 / b * a - 2 * I / (I * a + b)^2 / f^3 / (b - I * a) * b^2 * e^2 * a * d^2 / (a + I * b) / (I * b - a) * \arctan(1 / b * a) - 2 * I * b^2 * (d^2 * x^2 + 2 * c * d * x + c^2) / (b - I * a) / f / (I * a + b)^2 / (b * \exp(2 * I * (f * x + e))) + I * a * \exp(2 * I * (f * x + e)) - b + I * a + 4 / 3 / (I * a + b)^2 / (b - I * a) * b / (a + I * b) * d^2 * a * x^3 + 2 / (I * a + b)^2 / f / (b - I * a) * b^2 / (a + I * b) * d^2 * x^2 + 2 / (I * a + b)^2 / f^3 / (b - I * a) * b^2 / (a + I * b) * d^2 * \text{polylog}(2, (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) + 8 * I / (I * a + b)^2 / f^2 / (b - I * a) * b * e * d * a * c / (a + I * b) * \ln(\exp(I * (f * x + e))) + 4 * I / (I * a + b)^2 / f^2 / (b - I * a) * b / (a + I * b) * d * a * c * \ln(1 - (I * b - a) * \exp(2 * I * (f * x + e)) / (a + I * b)) * e
\end{aligned}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1576 vs. $2(534) = 1068$.

Time = 0.29 (sec) , antiderivative size = 1576, normalized size of antiderivative = 2.41

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
[Out] 1/6*(2*(a^3 - a*b^2)*d^2*f^3*x^3 - 6*b^3*c^2*f^2 - 6*(b^3*d^2*f^2 - (a^3 - a*b^2)*c*d*f^3)*x^2 - 6*(2*b^3*c*d*f^2 - (a^3 - a*b^2)*c^2*f^3)*x - 3*(-2*I*a^2*b*d^2*f*x - 2*I*a^2*b*c*d*f - I*a*b^2*d^2 + (-2*I*a*b^2*d^2*f*x - 2*I*a*b^2*c*d*f - I*b^3*d^2)*tan(f*x + e))*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) - 3*(2*I*a^2*b*d^2*f*x + 2*I*a^2*b*c*d*f + I*a*b^2*d^2 + (2*I*a*b^2*d^2*f*x + 2*I*a*b^2*c*d*f + I*b^3*d^2)*tan(f*x + e))*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 6*(a^2*b*d^2*f^2*x^2 - a^2*b*d^2*e^2 + 2*a^2*b*c*d*e*f + a*b^2*d^2*e + (2*a^2*b*c*d*f^2 + a*b^2*d^2*f)*x + (a*b^2*d^2*f^2*x^2 - a*b^2*d^2*e^2 + 2*a*b^2*c*d*e*f + b^3*d^2*e + (2*a*b^2*c*d*f^2 + b^3*d^2*f)*x)*tan(f*x + e))*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(a^2*b*d^2*f^2*x^2 - a^2*b*d^2*e^2 + 2*a^2*b*c*d*f^2 + a*b^2*d^2*f*x + (a*b^2*d^2*f^2*x^2 - a*b^2*d^2*e^2 + 2*a*b^2*c*d*e*f + b^3*d^2*e + (2*a*b^2*c*d*f^2 + b^3*d^2*f)*x)*tan(f*x + e))*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 6*(a^2*b*d^2*e^2 + a^2*b*c^2*f^2 - a*b^2*d^2*e - (2*a^2*b*c*d*e - a*b^2*c*d)*f + (a*b^2*d^2*e^2 + a*b^2*c^2*f^2 - b^3*d^2*e - (2*a*b^2*c*d*e - b^3*c*d)*f)*tan(f*x + e))*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + 6*(a^2*b*d^2*e^2 + a^2*b*c^2*f^2 - a*b^2*d^2*e - (2*a^2*b*c*d*e - a*b^2*c*d)*f + (a*b^2*d^2*e^2 + a*b^2*c^2*f^2 - b^3*d^2*e - (2*a*b^2*c*d*e - b^3*c*d)*f)*tan(f*x + e))*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + 3*(a*b^2*d^2*tan(f*x + e) + a^2*b*d^2)*polylog(3, ((a^2 + 2*I*a*b - b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*((a^2 - b^3)*d^2*f^3*x^3 + 3*a*b^2*c^2*f^2 + 3*(a*b^2*d^2*f^2 + (a^2*b - b^3)*c*d*f^3)*x^2 + 3*(2*a*b^2*c*d*f^2 + (a^2*b - b^3)*c^2*f^3)*x)*tan(f*x + e)) /((a^4*b + 2*a^2*b^3 + b^5)*f^3*tan(f*x + e) + (a^5 + 2*a^3*b^2 + a*b^4)*f^3)
```

Sympy [F]

$$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = \int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx$$

```
[In] integrate((d*x+c)**2/(a+b*tan(f*x+e))**2,x)
[Out] Integral((c + d*x)**2/(a + b*tan(e + f*x))**2, x)
```

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2530 vs. $2(534) = 1068$.

Time = 1.02 (sec) , antiderivative size = 2530, normalized size of antiderivative = 3.87

$$\int \frac{(c+dx)^2}{(a+b\tan(e+fx))^2} dx = \text{Too large to display}$$

$$\begin{aligned}
& -3)*d^2*e - (a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*c*d*f + 2*(I*a*b^2 + b^3)*d \\
& - 2)*(f*x + e)^2 + 3*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d^2*e^2 - 4*(-I*a*b^2 - b^3)*d^2*e - 4*(I*a*b^2 + b^3)*c*d*f)*(f*x + e))*\cos(2*f*x + 2*e) - 3 \\
& *(2*(I*a^2*b - a*b^2)*(f*x + e)*d^2 + 2*(-I*a^2*b + a*b^2)*d^2*e + 2*(I*a^2*b - a*b^2)*c*d*f + (I*a*b^2 - b^3)*d^2 + (2*(I*a^2*b + a*b^2)*(f*x + e)*d^2 \\
& + 2*(-I*a^2*b - a*b^2)*d^2*e + 2*(I*a^2*b + a*b^2)*c*d*f + (I*a*b^2 + b^3)*d^2)*\cos(2*f*x + 2*e) - (2*(a^2*b - I*a*b^2)*(f*x + e)*d^2 - 2*(a^2*b - I*a*b^2)*d^2*e + 2*(a^2*b - I*a*b^2)*c*d*f + (a*b^2 - I*b^3)*d^2)*\sin(2*f*x + 2*e))*\operatorname{dilog}((I*a + b)*e^{(2*I*f*x + 2*I*e)/(-I*a + b)}) + 3*((a^2*b + I*a*b^2)*d^2*e^2 - (a*b^2 + I*b^3)*d^2*e + (a*b^2 + I*b^3)*c*d*f + ((a^2*b - I*a*b^2)*d^2*e^2 - (a*b^2 - I*b^3)*d^2*e + (a*b^2 - I*b^3)*c*d*f)*\cos(2*f*x + 2*e) - ((-I*a^2*b - a*b^2)*d^2*e^2 + (I*a*b^2 + b^3)*d^2*e + (-I*a*b^2 - b^3)*c*d*f)*\sin(2*f*x + 2*e))*\log((a^2 + b^2)*\cos(2*f*x + 2*e)^2 + 4*a*b*\sin(2*f*x + 2*e) + (a^2 + b^2)*\sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*c \\
& \os(2*f*x + 2*e)) + 3*((a^2*b + I*a*b^2)*(f*x + e)^2*d^2 - (2*(a^2*b + I*a*b^2)*d^2*e - 2*(a^2*b + I*a*b^2)*c*d*f - (a*b^2 + I*b^3)*d^2)*(f*x + e) + ((a^2*b - I*a*b^2)*(f*x + e)^2*d^2 - (2*(a^2*b - I*a*b^2)*d^2*e - 2*(a^2*b - I*a*b^2)*c*d*f - (a*b^2 - I*b^3)*d^2)*(f*x + e))*\cos(2*f*x + 2*e) - ((-I*a^2*b - a*b^2)*(f*x + e)^2*d^2 + (2*(I*a^2*b + a*b^2)*d^2*e + 2*(-I*a^2*b - a*b^2)*c*d*f + (-I*a*b^2 - b^3)*d^2)*(f*x + e))*\sin(2*f*x + 2*e))*\log(((a^2 + b^2)*\cos(2*f*x + 2*e)^2 + 4*a*b*\sin(2*f*x + 2*e) + (a^2 + b^2)*\sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*\cos(2*f*x + 2*e))/(a^2 + b^2)) + 3*((a^2*b - I*a*b^2)*d^2*2*\cos(2*f*x + 2*e) - (-I*a^2*b - a*b^2)*d^2*\sin(2*f*x + 2*e) + (a^2*b + I*a*b^2)*d^2)*\operatorname{polylog}(3, (I*a + b)*e^{(2*I*f*x + 2*I*e)/(-I*a + b)}) + ((I*a^3 + 3*a^2*b - 3*I*a*b^2 - b^3)*(f*x + e)^3*d^2 - 3*((I*a^3 + 3*a^2*b - 3*I*a*b^2 - b^3)*d^2*e + (-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3)*c*d*f - 2*(a*b^2 - I*b^3)*d^2)*(f*x + e)^2 - 3*((-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3)*d^2*e^2 + 4*(a*b^2 - I*b^3)*d^2*e - 4*(a*b^2 - I*b^3)*c*d*f)*(f*x + e))*\sin(2*f*x + 2*e))/((a^5 - I*a^4*b + 2*a^3*b^2 - 2*I*a^2*b^3 + a*b^4 - I*b^5)*f^2*\cos(2*f*x + 2*e) - (-I*a^5 - a^4*b - 2*I*a^3*b^2 - 2*a^2*b^3 - I*a*b^4 - b^5)*f^2*\sin(2*f*x + 2*e) + (a^5 + I*a^4*b + 2*a^3*b^2 + 2*I*a^2*b^3 + a*b^4 + I*b^5)*f^2)/f
\end{aligned}$$

Giac [F]

$$\int \frac{(c + dx)^2}{(a + b\tan(e + fx))^2} dx = \int \frac{(dx + c)^2}{(b\tan(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((d*x + c)^2/(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{(a + b \tan(e + fx))^2} dx = \int \frac{(c + d x)^2}{(a + b \tan (e + f x))^2} dx$$

[In] `int((c + d*x)^2/(a + b*tan(e + f*x))^2,x)`

[Out] `int((c + d*x)^2/(a + b*tan(e + f*x))^2, x)`

3.61 $\int \frac{c+dx}{(a+b\tan(e+fx))^2} dx$

Optimal result	417
Rubi [A] (verified)	417
Mathematica [B] (verified)	420
Maple [B] (verified)	421
Fricas [B] (verification not implemented)	422
Sympy [F]	423
Maxima [B] (verification not implemented)	423
Giac [F]	424
Mupad [F(-1)]	424

Optimal result

Integrand size = 18, antiderivative size = 214

$$\begin{aligned} \int \frac{c+dx}{(a+b\tan(e+fx))^2} dx = & -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adf)^2}{4a(a+ib)(a^2+b^2)df^2} \\ & + \frac{b(bd+2acf+2adf)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\ & - \frac{iabd\operatorname{PolyLog}\left(2,-\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\ & - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} \end{aligned}$$

```
[Out] -1/2*(d*x+c)^2/(a^2+b^2)/d+1/4*(2*a*d*f*x+2*a*c*f+b*d)^2/a/(a+I*b)/(a^2+b^2)
/d/f^2+b*(2*a*d*f*x+2*a*c*f+b*d)*ln(1+(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2
)/(a^2+b^2)^2/f^2-I*a*b*d*polylog(2,-(a^2+b^2)*exp(2*I*(f*x+e))/(a+I*b)^2)/
(a^2+b^2)^2/f^2-b*(d*x+c)/(a^2+b^2)/f/(a+b*tan(f*x+e))
```

Rubi [A] (verified)

Time = 0.30 (sec), antiderivative size = 214, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.278, Rules used

$$= \{3814, 3813, 2221, 2317, 2438\}$$

$$\begin{aligned} \int \frac{c+dx}{(a+b\tan(e+fx))^2} dx &= \frac{b(2acf+2adf x+bd)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f^2(a^2+b^2)^2} \\ &\quad - \frac{b(c+dx)}{f(a^2+b^2)(a+b\tan(e+fx))} + \frac{(2acf+2adf x+bd)^2}{4adf^2(a+ib)(a^2+b^2)} \\ &\quad - \frac{(c+dx)^2}{2d(a^2+b^2)} - \frac{iabd \operatorname{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{f^2(a^2+b^2)^2} \end{aligned}$$

[In] $\operatorname{Int}[(c+d*x)/(a+b*\operatorname{Tan}[e+f*x])^2, x]$

[Out]
$$\begin{aligned} &-1/2*(c+d*x)^2/((a^2+b^2)*d) + (b*d+2*a*c*f+2*a*d*f*x)^2/(4*a*(a+I*b)*(a^2+b^2)*d*f^2) + (b*(b*d+2*a*c*f+2*a*d*f*x)*\operatorname{Log}[1+((a^2+b^2)*E^((2*I)*(e+f*x)))/(a+I*b)^2])/((a^2+b^2)^2*f^2) - (I*a*b*d*\operatorname{PolyLog}[2, -(((a^2+b^2)*E^((2*I)*(e+f*x)))/(a+I*b)^2)])/((a^2+b^2)^2*f^2) \\ &- (b*(c+d*x))/((a^2+b^2)*f*(a+b*\operatorname{Tan}[e+f*x])) \end{aligned}$$

Rule 2221

$\operatorname{Int}[(((F_)((g_)*(e_)+(f_)*(x_)))^(n_)*((c_)+(d_)*(x_))^(m_))/((a_)+(b_)*(F_)((g_)*(e_)+(f_)*(x_)))^(n_)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[((c+d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1+b*((F^((g*(e+f*x)))^n)/a)], x] - \operatorname{Dist}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c+d*x)^(m-1)*\operatorname{Log}[1+b*((F^((g*(e+f*x)))^n)/a)], x]] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \operatorname{IGtQ}[m, 0]$

Rule 2317

$\operatorname{Int}[\operatorname{Log}[(a_)+(b_)*(F_)((e_)*(c_)+(d_)*(x_)))^(n_)], x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[1/(d*e*n*\operatorname{Log}[F]), \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[a+b*x]/x, x], x, (F^((e*(c+d*x)))^n), x]] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \operatorname{GtQ}[a, 0]$

Rule 2438

$\operatorname{Int}[\operatorname{Log}[(c_)*(d_)+(e_)*(x_)^(n_)]/(x_), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[-\operatorname{PolyLog}[2, (-c)*e*x^n]/n, x] /; \operatorname{FreeQ}[\{c, d, e, n\}, x] \&& \operatorname{EqQ}[c*d, 1]$

Rule 3813

$\operatorname{Int}[(c_)+(d_)*(x_)^(m_)/((a_)+(b_)*\operatorname{tan}[(e_)+(f_)*(x_)]), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}[(c+d*x)^(m+1)/(d*(m+1)*(a+I*b)), x] + \operatorname{Dist}[2*I*b, \operatorname{Int}[(c+d*x)^m*(E^{\operatorname{Simp}[2*I*(e+f*x), x]}/((a+I*b)^2+(a^2+b^2)*E^{\operatorname{Simp}[2*I*(e+f*x), x]}), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \operatorname{NeQ}[a^2+b^2, 0] \&& \operatorname{IGtQ}[m, 0]$

Rule 3814

```

Int[((c_) + (d_)*(x_))/((a_) + (b_)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol
] :> Simp[-(c + d*x)^2/(2*d*(a^2 + b^2)), x] + (Dist[1/(f*(a^2 + b^2)), Int
[(b*d + 2*a*c*f + 2*a*d*f*x)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((c + d*
x)/(f*(a^2 + b^2)*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f},
x] && NeQ[a^2 + b^2, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c+dx)^2}{2(a^2+b^2)d} - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} + \frac{\int \frac{bd+2acf+2adf}{a+b\tan(e+fx)} dx}{(a^2+b^2)f} \\
&= -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adf)^2}{4a(a+ib)(a^2+b^2)df^2} \\
&\quad - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} + \frac{(2ib)\int \frac{e^{2i(e+fx)}(bd+2acf+2adf)}{(a+ib)^2+(a^2+b^2)e^{2i(e+fx)}} dx}{(a^2+b^2)f} \\
&= -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adf)^2}{4a(a+ib)(a^2+b^2)df^2} + \frac{b(bd+2acf+2adf)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\
&\quad - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} - \frac{(2abd)\int \log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right) dx}{(a^2+b^2)^2f} \\
&= -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adf)^2}{4a(a+ib)(a^2+b^2)df^2} + \frac{b(bd+2acf+2adf)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\
&\quad - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))} + \frac{(iabd)\text{Subst}\left(\int \frac{\log\left(1+\frac{(a^2+b^2)x}{(a+ib)^2}\right)}{x} dx, x, e^{2i(e+fx)}\right)}{(a^2+b^2)^2f^2} \\
&= -\frac{(c+dx)^2}{2(a^2+b^2)d} + \frac{(bd+2acf+2adf)^2}{4a(a+ib)(a^2+b^2)df^2} \\
&\quad + \frac{b(bd+2acf+2adf)\log\left(1+\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} \\
&\quad - \frac{iabd \text{PolyLog}\left(2, -\frac{(a^2+b^2)e^{2i(e+fx)}}{(a+ib)^2}\right)}{(a^2+b^2)^2f^2} - \frac{b(c+dx)}{(a^2+b^2)f(a+b\tan(e+fx))}
\end{aligned}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 745 vs. $2(214) = 428$.

Time = 7.89 (sec) , antiderivative size = 745, normalized size of antiderivative = 3.48

$$\begin{aligned}
 & \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx \\
 = & \frac{(e + fx)(-2de + 2cf + d(e + fx)) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{2(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2} \\
 & + \frac{b^2d(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{a(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2} \\
 & - \frac{2bde(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f^2(a + b \tan(e + fx))^2} \\
 & + \frac{2bc(-b(e + fx) + a \log(a \cos(e + fx) + b \sin(e + fx))) \sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx))^2}{(a - ib)(a + ib)(a^2 + b^2)f(a + b \tan(e + fx))^2} \\
 & - \frac{d \left(e^{i \arctan(\frac{a}{b})} (e + fx)^2 + \frac{a(i(e+fx)(-\pi+2\arctan(\frac{a}{b}))-\pi \log(1+e^{-2i(e+fx)})-2(e+fx+\arctan(\frac{a}{b})) \log(1-e^{2i(e+fx+\arctan(\frac{a}{b}))})}{\sqrt{1+\frac{a^2}{b^2}}}\right)}{(a - ib)(a + ib)} \\
 & + \frac{\sec^2(e + fx)(a \cos(e + fx) + b \sin(e + fx)) (-b^2de \sin(e + fx) + b^2cf \sin(e + fx) + b^2d(e + fx) \sin(e + fx))}{a(a - ib)(a + ib)f^2(a + b \tan(e + fx))^2}
 \end{aligned}$$

[In] `Integrate[(c + d*x)/(a + b*Tan[e + f*x])^2, x]`

[Out]

```

((e + f*x)*(-2*d*e + 2*c*f + d*(e + f*x))*Sec[e + f*x]^2*(a*Cos[e + f*x] +
b*Sin[e + f*x])^2)/(2*(a - I*b)*(a + I*b)*f^2*(a + b*Tan[e + f*x])^2) + (b^
2*d*(-(b*(e + f*x)) + a*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])*Sec[e + f*x]^
2*(a*Cos[e + f*x] + b*Sin[e + f*x])^2)/(a*(a - I*b)*(a + I*b)*(a^2 + b^2)*f
^2*(a + b*Tan[e + f*x])^2) - (2*b*d*e*(-(b*(e + f*x)) + a*Log[a*Cos[e + f*x]
+ b*Sin[e + f*x]])*Sec[e + f*x]^2*(a*Cos[e + f*x] + b*Sin[e + f*x])^2)/((a
- I*b)*(a + I*b)*(a^2 + b^2)*f^2*(a + b*Tan[e + f*x])^2) + (2*b*c*(-(b*(e
+ f*x)) + a*Log[a*Cos[e + f*x] + b*Sin[e + f*x]])*Sec[e + f*x]^2*(a*Cos[e
+ f*x] + b*Sin[e + f*x])^2)/((a - I*b)*(a + I*b)*(a^2 + b^2)*f*(a + b*Tan[e
+ f*x])^2) - (d*(E^(I*ArcTan[a/b]))*(e + f*x)^2 + (a*(I*(e + f*x)*(-Pi + 2*
ArcTan[a/b]) - Pi*Log[1 + E^((-2*I)*(e + f*x))]) - 2*(e + f*x + ArcTan[a/b])
*Log[1 - E^((2*I)*(e + f*x + ArcTan[a/b]))]) + Pi*Log[Cos[e + f*x]] + 2*ArcT
an[a/b]*Log[Sin[e + f*x + ArcTan[a/b]]] + I*PolyLog[2, E^((2*I)*(e + f*x +
ArcTan[a/b]))])/((a - I*b)*(a + I*b)*Sqrt[(a^2 + b^2)/b^2])*f^2*(a + b*Tan[
e + f*x])^2) + ((a - I*b)*(a + I*b)*Sqrt[(a^2 + b^2)/b^2])*f^2*(a + b*Tan[e
+ f*x])^2 + (Sec[e + f*x]^2*(a*Cos[e + f*x] + b*Sin[e + f*x])*(-(b^2*d*e
*Sin[e + f*x]) + b^2*c*f*Sin[e + f*x] + b^2*d*(e + f*x)*Sin[e + f*x]))/(a*(a
- I*b)*(a + I*b)*f^2*(a + b*Tan[e + f*x])^2)

```

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2000 vs. $2(202) = 404$.

Time = 0.86 (sec) , antiderivative size = 2001, normalized size of antiderivative = 9.35

method	result	size
risch	Expression too large to display	2001

```
[In] int((d*x+c)/(a+b*tan(f*x+e))^2,x,method=_RETURNVERBOSE)
```

$$\frac{p(2*I*(f*x+e))*b - 1/2*b/a + 1/2/b*a*exp(2*I*(f*x+e)) + 1/2/b*a + 2*I/(I*a+b)^2/f^2}{2/(b-I*a)*b^2*a*d*e/(a+I*b)/(I*b-a)*arctan(1/b*a)}$$

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 843 vs. $2(197) = 394$.

Time = 0.27 (sec) , antiderivative size = 843, normalized size of antiderivative = 3.94

$$\int \frac{c + dx}{(a + b\tan(e + fx))^2} dx \\ = \frac{(a^3 - ab^2)df^2x^2 - 2b^3cf - 2(b^3df - (a^3 - ab^2)cf^2)x + (iab^2d\tan(fx + e) + ia^2bd)\text{Li}_2\left(\frac{2((iab - b^2)\tan(fx + e)}{(a^2 + b^2)}\right)}{(a^2 + b^2)^2}$$

```
[In] integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="fricas")
[Out] 1/2*((a^3 - a*b^2)*d*f^2*x^2 - 2*b^3*c*f - 2*(b^3*d*f - (a^3 - a*b^2)*c*f^2)*x + (I*a*b^2*d*tan(f*x + e) + I*a^2*b*d)*dilog(2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + (-I*a*b^2*d*tan(f*x + e) - I*a^2*b*d)*dilog(2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2) + 1) + 2*(a^2*b*d*f*x + a^2*b*d*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((I*a*b - b^2)*tan(f*x + e)^2 - a^2 - I*a*b + (I*a^2 - 2*a*b - I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) + 2*(a^2*b*d*f*x + a^2*b*d*e + (a*b^2*d*f*x + a*b^2*d*e)*tan(f*x + e))*log(-2*((-I*a*b - b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (-I*a^2 - 2*a*b + I*b^2)*tan(f*x + e))/((a^2 + b^2)*tan(f*x + e)^2 + a^2 + b^2)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (2*a*b^2*d*e - 2*a*b^2*c*f - b^3*d)*tan(f*x + e))*log(((I*a*b + b^2)*tan(f*x + e)^2 - a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) - (2*a^2*b*d*e - 2*a^2*b*c*f - a*b^2*d + (2*a*b^2*d*e - 2*a*b^2*c*f - b^3*d)*tan(f*x + e))*log(((I*a*b - b^2)*tan(f*x + e)^2 + a^2 + I*a*b + (I*a^2 + I*b^2)*tan(f*x + e))/(tan(f*x + e)^2 + 1)) + ((a^2*b - b^3)*d*f^2*x^2 + 2*a*b^2*c*f + 2*(a*b^2*d*f + (a^2*b - b^3)*c*f^2)*x)*tan(f*x + e)/((a^4*b + 2*a^2*b^3 + b^5)*f^2*tan(f*x + e) + (a^5 + 2*a^3*b^2 + a*b^4)*f^2)
```

Sympy [F]

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tan(e + fx))^2} dx$$

[In] `integrate((d*x+c)/(a+b*tan(f*x+e))**2,x)`
[Out] `Integral((c + d*x)/(a + b*tan(e + f*x))**2, x)`

Maxima [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1156 vs. $2(197) = 394$.

Time = 0.71 (sec), antiderivative size = 1156, normalized size of antiderivative = 5.40

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)/(a+b*tan(f*x+e))**2,x, algorithm="maxima")`
[Out]
$$\begin{aligned} & 1/2*((a^3 - I*a^2*b + a*b^2 - I*b^3)*d*f^2*x^2 + 2*(a^3 - I*a^2*b + a*b^2 - I*b^3)*c*f^2*x - 4*(-I*a*b^2 + b^3)*c*f - 2*(2*(-I*a^2*b + a*b^2)*c*f + (-I*a*b^2 + b^3)*d + (2*(-I*a^2*b - a*b^2)*c*f + (-I*a*b^2 - b^3)*d)*cos(2*f*x + 2*e) + (2*(a^2*b - I*a*b^2)*c*f + (a*b^2 - I*b^3)*d)*sin(2*f*x + 2*e))*arctan2(-b*cos(2*f*x + 2*e) + a*sin(2*f*x + 2*e) + b, a*cos(2*f*x + 2*e) + b*sin(2*f*x + 2*e) + a) - 4*((I*a^2*b + a*b^2)*d*f*x*cos(2*f*x + 2*e) - (a^2*b - I*a*b^2)*d*f*x*sin(2*f*x + 2*e) + (I*a^2*b - a*b^2)*d*f*x)*arctan2((2*a*b*cos(2*f*x + 2*e) - (a^2 - b^2)*sin(2*f*x + 2*e))/(a^2 + b^2), (2*a*b*sin(2*f*x + 2*e) + a^2 + b^2 + (a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + ((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*d*f^2*x^2 + 2*((a^3 - 3*I*a^2*b - 3*a*b^2 + I*b^3)*c*f^2 - 2*(I*a*b^2 + b^3)*d*f*x)*cos(2*f*x + 2*e) - 2*((I*a^2*b + a*b^2)*d*cos(2*f*x + 2*e) - (a^2*b - I*a*b^2)*d*sin(2*f*x + 2*e) + (I*a^2*b - a*b^2)*d*dilog((I*a + b)*e^(2*I*f*x + 2*I*e)/(-I*a + b)) + (2*(a^2*b + I*a*b^2)*c*f + (a*b^2 + I*b^3)*d + (2*(a^2*b - I*a*b^2)*c*f + (a*b^2 - I*b^3)*d)*cos(2*f*x + 2*e) - (2*(-I*a^2*b - a*b^2)*c*f - (I*a*b^2 + b^3)*d)*sin(2*f*x + 2*e))*log((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e)) + 2*((a^2*b - I*a*b^2)*d*f*x*cos(2*f*x + 2*e) - (-I*a^2*b - a*b^2)*d*f*x*sin(2*f*x + 2*e) + (a^2*b + I*a*b^2)*d*f*x)*log(((a^2 + b^2)*cos(2*f*x + 2*e)^2 + 4*a*b*sin(2*f*x + 2*e) + (a^2 + b^2)*sin(2*f*x + 2*e)^2 + a^2 + b^2 + 2*(a^2 - b^2)*cos(2*f*x + 2*e))/(a^2 + b^2)) + ((I*a^3 + 3*a^2*b - 3*I*a*b^2 - b^3)*d*f^2*x^2 - 2*((-I*a^3 - 3*a^2*b + 3*I*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 - I*b^3)*d*f*x)*sin(2*f*x + 2*e))/((a^5 - I*a^4*b + 2*a^3*b^2 - 2*I*a^2*b^3 + a*b^4 - I*b^5)*f^2*cos(2*f*x + 2*e) - (-I*a^5 - a^4*b - 2*I*a^3*b^2 - 2*a^2*b^3 - I*a*b^4 - b^5)*f^2*sin(2*f*x + 2*e) + (a^5 + I*a^4*b + 2*a^3*b^2 + 2*I*a^2*b^3 + a*b^4 + I*b^5)*f^2) \end{aligned}$$

Giac [F]

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{dx + c}{(b \tan(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)/(b*tan(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{(a + b \tan(e + fx))^2} dx = \int \frac{c + d x}{(a + b \tan (e + f x))^2} dx$$

[In] `int((c + d*x)/(a + b*tan(e + f*x))^2,x)`
[Out] `int((c + d*x)/(a + b*tan(e + f*x))^2, x)`

3.62 $\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$

Optimal result	425
Rubi [N/A]	425
Mathematica [N/A]	426
Maple [N/A] (verified)	426
Fricas [N/A]	426
Sympy [N/A]	427
Maxima [N/A]	427
Giac [N/A]	428
Mupad [N/A]	428

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\tan(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*tan(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.07 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

[In] Int[1/((c + d*x)*(a + b*Tan[e + f*x])^2),x]

[Out] Defer[Int][1/((c + d*x)*(a + b*Tan[e + f*x])^2), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 13.94 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx$$

[In] `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2), x]`

[Out] `Integrate[1/((c + d*x)*(a + b*Tan[e + f*x])^2), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.55 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tan(fx + e))^2} dx$$

[In] `int(1/(d*x+c)/(a+b*tan(f*x+e))^2, x)`

[Out] `int(1/(d*x+c)/(a+b*tan(f*x+e))^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \frac{1}{(c + dx)(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \tan(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2, x, algorithm="fricas")`

[Out] `integral(1/(a^2*d*x + a^2*c + (b^2*d*x + b^2*c)*tan(f*x + e)^2 + 2*(a*b*d*x + a*b*c)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.67 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(a+b\tan(e+fx))^2(c+dx)} dx$$

```
[In] integrate(1/(d*x+c)/(a+b*tan(f*x+e))**2,x)
```

[Out] $\text{Integral}(1/((a + b\tan(e + f*x))^{**2}*(c + d*x)), x)$

Maxima [N/A]

Not integrable

Time = 8.73 (sec) , antiderivative size = 1498, normalized size of antiderivative = 74.90

$$\int \frac{1}{(c+dx)(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)(b\tan(fx+e)+a)^2} dx$$

```
[In] integrate(1/(d*x+c)/(a+b*tan(f*x+e))^(2),x, algorithm="maxima")
```

```
[Out] (((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*cos(2*f*x + 2*e)^2*log(d*x + c) + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*log(d*x + c)*sin(2*f*x + 2*e)^2 - 2*(2*a*b^3*d - ((a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*log(d*x + c))*cos(2*f*x + 2*e) + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c*d*f)*sin(2*f*x + 2*e))*integrate(2*(2*(2*a^2*b^2*d*f*x + 2*a^2*b^2*c*f - a*b^3*d)*cos(2*f*x + 2*e) - (2*(a^3*b - a*b^3)*d*f*x + 2*(a^3*b - a*b^3)*c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e)))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^2*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c^2*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^2*f*x^2 + 2*(a^5*b + 2*a^3*b^3 + a*b^5)*c*d*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c^2*f)
```

$$\begin{aligned} & \text{^5)*c^2*f)*sin(2*f*x + 2*e)), x) + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*log(d*x + c) + 2*((a^2*b^2 - b^4)*d + 2*((a^3*b - a*b^3)*c*f)*log(d*x + c))*sin(2*f*x + 2*e))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c*d*f)*sin(2*f*x + 2*e)) \end{aligned}$$

Giac [N/A]

Not integrable

Time = 0.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b\tan(e + fx))^2} dx = \int \frac{1}{(dx + c)(b\tan(fx + e) + a)^2} dx$$

```
[In] integrate(1/(d*x+c)/(a+b*tan(f*x+e))^2,x, algorithm="giac")
[Out] integrate(1/((d*x + c)*(b*tan(f*x + e) + a)^2), x)
```

Mupad [N/A]

Not integrable

Time = 4.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b\tan(e + fx))^2} dx = \int \frac{1}{(a + b\tan(e + fx))^2 (c + dx)} dx$$

```
[In] int(1/((a + b*tan(e + f*x))^2*(c + d*x)),x)
[Out] int(1/((a + b*tan(e + f*x))^2*(c + d*x)), x)
```

3.63 $\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$

Optimal result	429
Rubi [N/A]	429
Mathematica [N/A]	430
Maple [N/A] (verified)	430
Fricas [N/A]	430
Sympy [N/A]	431
Maxima [N/A]	431
Giac [N/A]	432
Mupad [N/A]	432

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\tan(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.06 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$$

[In] Int[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2),x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 15.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx$$

[In] `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]`

[Out] `Integrate[1/((c + d*x)^2*(a + b*Tan[e + f*x])^2), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.65 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2 (a + b \tan(fx + e))^2} dx$$

[In] `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2, x)`

[Out] `int(1/(d*x+c)^2/(a+b*tan(f*x+e))^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 96, normalized size of antiderivative = 4.80

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2, x, algorithm="fricas")`

[Out] `integral(1/(a^2*d^2*x^2 + 2*a^2*c*d*x + a^2*c^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*tan(f*x + e)^2 + 2*(a*b*d^2*x^2 + 2*a*b*c*d*x + a*b*c^2)*tan(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 3.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(a+b\tan(e+fx))^2(c+dx)^2} dx$$

```
[In] integrate(1/(d*x+c)**2/(a+b*tan(f*x+e))**2,x)
```

[Out] $\text{Integral}(1/((a + b\tan(e + f*x))^{2*(c + d*x)}), x)$

Maxima [N/A]

Not integrable

Time = 27.22 (sec) , antiderivative size = 1977, normalized size of antiderivative = 98.85

$$\int \frac{1}{(c+dx)^2(a+b\tan(e+fx))^2} dx = \int \frac{1}{(dx+c)^2(b\tan(fx+e)+a)^2} dx$$

```
[In] integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="maxima")
```

```
[Out] -((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*cos(2*f*x + 2*e)^2 + ((a^4 - b^4)*d*f*x + (a^4 - b^4)*c*f)*sin(2*f*x + 2*e)^2 + 2*(2*a*b^3*d + (a^4 - 2*a^2*b^2 + b^4)*d*f*x + (a^4 - 2*a^2*b^2 + b^4)*c*f)*cos(2*f*x + 2*e) - ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^2 + 2*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f)*sin(2*f*x + 2*e)^2 + 2*((a^6 + a^4*b^2 - a^2*b^4 - b^6)*d^3*f*x^2 + 2*(a^6 + a^4*b^2 - a^2*b^4 - b^6)*c*d^2*f*x + (a^6 + a^4*b^2 - a^2*b^4 - b^6)*c^2*d*f)*cos(2*f*x + 2*e) + 4*((a^5*b + 2*a^3*b^3 + a*b^5)*d^3*f*x^2 + 2*(a^5*b + 2*a^3*b^3 + a*b^5)*c*d^2*f*x + (a^5*b + 2*a^3*b^3 + a*b^5)*c^2*d*f)*sin(2*f*x + 2*e))*integrate(4*(2*(a^2*b^2*d*f*x + a^2*b^2*c*f - a*b^3*d)*cos(2*f*x + 2*e) - ((a^3*b - a*b^3)*d*f*x + (a^3*b - a*b^3)*c*f - (a^2*b^2 - b^4)*d)*sin(2*f*x + 2*e))/((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^3*f + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^3*f)*cos(2*f*x + 2*e)^2 + ((a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*d^3*f*x^3 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c*d^2*f*x^2 + 3*(a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^2*d*f*x + (a^6 + 3*a^4*b^2 + 3*a^2*b^4 + b^6)*c^3*f)*sin(2*f*x + 2*e)^2)
```

$$\begin{aligned}
& \hat{4} * b^2 + 3 * a^2 * b^4 + b^6) * c * d^2 * f * x^2 + 3 * (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) \\
& * c^2 * d * f * x + (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * c^3 * f * \sin(2 * f * x + 2 * e)^2 \\
& + 2 * ((a^6 + a^4 * b^2 - a^2 * b^4 - b^6) * d^3 * f * x^3 + 3 * (a^6 + a^4 * b^2 - a^2 * b^4 \\
& - b^6) * c * d^2 * f * x^2 + 3 * (a^6 + a^4 * b^2 - a^2 * b^4 - b^6) * c^2 * d * f * x + (a^6 \\
& + a^4 * b^2 - a^2 * b^4 - b^6) * c^3 * f * \cos(2 * f * x + 2 * e) + 4 * ((a^5 * b + 2 * a^3 * b^3 \\
& + a * b^5) * d^3 * f * x^3 + 3 * (a^5 * b + 2 * a^3 * b^3 + a * b^5) * c * d^2 * f * x^2 + 3 * (a^5 * b \\
& + 2 * a^3 * b^3 + a * b^5) * c^2 * d * f * x + (a^5 * b + 2 * a^3 * b^3 + a * b^5) * c^3 * f * \sin(2 * f * \\
& x + 2 * e)), x) + 2 * (2 * (a^3 * b - a * b^3) * d * f * x + 2 * (a^3 * b - a * b^3) * c * f - (a^2 * b \\
& ^2 - b^4) * d * \sin(2 * f * x + 2 * e)) / ((a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * d^3 * f * x \\
& ^2 + 2 * (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * c * d^2 * f * x + (a^6 + 3 * a^4 * b^2 + 3 \\
& * a^2 * b^4 + b^6) * c^2 * d * f + ((a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * d^3 * f * x^2 + \\
& 2 * (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * c * d^2 * f * x + (a^6 + 3 * a^4 * b^2 + 3 * a^2 * \\
& b^4 + b^6) * c^2 * d * f * \cos(2 * f * x + 2 * e)^2 + ((a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) \\
& * d^3 * f * x^2 + 2 * (a^6 + 3 * a^4 * b^2 + 3 * a^2 * b^4 + b^6) * c * d^2 * f * x + (a^6 + 3 * a \\
& ^4 * b^2 + 3 * a^2 * b^4 + b^6) * c^2 * d * f * \sin(2 * f * x + 2 * e)^2 + 2 * ((a^6 + a^4 * b^2 - \\
& a^2 * b^4 - b^6) * d^3 * f * x^2 + 2 * (a^6 + a^4 * b^2 - a^2 * b^4 - b^6) * c * d^2 * f * x + (a^6 + a^4 * b^2 \\
& - a^2 * b^4 - b^6) * c^2 * d * f * \cos(2 * f * x + 2 * e) + 4 * ((a^5 * b + 2 * a^3 * b^3 + a * b^5) * d^3 * f * x^2 + \\
& 2 * (a^5 * b + 2 * a^3 * b^3 + a * b^5) * c * d^2 * f * x + (a^5 * b + 2 * a^3 * b^3 + a * b^5) * c^2 * d * f * \sin(2 * f * x + 2 * e))
\end{aligned}$$

Giac [N/A]

Not integrable

Time = 22.56 (sec), antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tan(fx + e) + a)^2} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tan(f*x+e))^2,x, algorithm="giac")

[Out] integrate(1/((d*x + c)^2*(b*tan(f*x + e) + a)^2), x)

Mupad [N/A]

Not integrable

Time = 3.99 (sec), antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tan(e + fx))^2} dx = \int \frac{1}{(a + b \tan(e + fx))^2(c + dx)^2} dx$$

[In] int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2),x)

[Out] int(1/((a + b*tan(e + f*x))^2*(c + d*x)^2), x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	433
--------------------------------------------	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                                         Small rewrite of logic in main function to make it*)
(*                                         match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal}
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A",""}
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count is different."}
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)
    finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
  ]
]
,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>ToString[Order[result]]},
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];
finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn] === Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]] === Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
              1,
              Max[ExpnType[expn[[1]]], 2]],
            Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
        If[Head[expn] === Plus || Head[expn] === Times,
          Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
              If[HypergeometricFunctionQ[Head[expn]],
                Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
                If[AppellFunctionQ[Head[expn]],
                  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
                  If[Head[expn] === RootSum,
                    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
                    If[Head[expn] === Integrate || Head[expn] === Int,
                      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
                      9]]]]]]]]]
]

ElementaryFunctionQ[func_] :=
  MemberQ[{  

    Exp, Log,  

    Sin, Cos, Tan, Cot, Sec, Csc,  

    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
  }]

```

```

Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]

```

```

SpecialFunctionQ[func_] :=
MemberQ[{{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func}]

```

```

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (",

```

```

        convert(leaf_count_optimal,string), " ) = ",convert(2*leaf_
    end if
else #result contains complex but optimal is not
if debug then
    print("result contains complex but optimal is not");
fi;
return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
# this assumes optimal do not as well. No check is needed here.
if debug then
    print("result do not contain complex, this assumes optimal do not as well")
fi;
if leaf_count_result<=2*leaf_count_optimal then
if debug then
    print("leaf_count_result<=2*leaf_count_optimal");
fi;
return "A"," ";
else
if debug then
    print("leaf_count_result>2*leaf_count_optimal");
fi;
return "B",cat("Leaf count of result is larger than twice the leaf count of op-
    convert(leaf_count_result,string)," vs. $2(", 
    convert(leaf_count_optimal,string),")=",convert(2*leaf_count_
fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
if debug then
    print("ExpnType(result) > ExpnType(optimal)");
fi;
return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:
```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+``') or type(expn,'`*``') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except:
        return False
```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+`) or type(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sageMath")
    #print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count(result))-str(leaf_count(optimal))
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType(result))-str(ExpnType(optimal))

```

```
#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#          Albert Rich to use with Sagemath. This is used to
#          grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#          'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#          issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:  #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False
```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than optimal."
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```