

Computer Algebra Independent Integration Tests

Summer 2023 edition

4-Trig-functions/4.4-Cotangent/113-4.4.7-d-trig- \hat{m} -a+b-c-cot- \hat{n} -
 \hat{p}

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 4:29pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	43
4	Appendix	407

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [64]. This is test number [113].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (64)	0.00 (0)
Mathematica	100.00 (64)	0.00 (0)
Fricas	100.00 (64)	0.00 (0)
Maple	98.44 (63)	1.56 (1)
Giac	84.38 (54)	15.62 (10)
Mupad	60.94 (39)	39.06 (25)
Maxima	35.94 (23)	64.06 (41)
Sympy	17.19 (11)	82.81 (53)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Maple	64.062	32.812	1.562	1.562
Mathematica	62.500	12.500	25.000	0.000
Maxima	25.000	7.812	3.125	64.062
Giac	23.438	53.125	7.812	15.625
Fricas	17.188	76.562	6.250	0.000
Sympy	12.500	4.688	0.000	82.812
Mupad	0.000	60.938	0.000	39.062

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	1	100.00	0.00	0.00
Giac	10	0.00	10.00	90.00
Mupad	25	0.00	100.00	0.00
Maxima	41	56.10	0.00	43.90
Sympy	53	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.12
Fricas	0.34
Maxima	0.38
Maple	0.42
Giac	0.75
Mathematica	1.23
Sympy	6.49
Mupad	12.79

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	70.42	1.00	60.50	1.00
Maxima	142.13	4.02	52.00	1.21
Mathematica	158.78	1.85	72.00	1.03
Maple	161.00	2.05	84.00	1.23
Giac	246.44	2.91	164.50	2.30
Mupad	311.87	3.12	47.00	1.00
Fricas	454.30	5.52	337.00	5.31
Sympy	1072.82	8.82	88.00	1.45

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {25, 35, 36, 37, 40, 43, 48, 53, 55, 58}

Maple {24, 30, 52, 53, 58}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v1.0a

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	39

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 3, 4, 5, 6, 7, 10, 11, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 26, 27, 28, 29, 31, 32, 33, 39, 40, 41, 42, 43, 44, 46, 47, 49, 60, 61, 62, 63, 64 }

B grade { 8, 9, 12, 30, 34, 38, 45, 50 }

C grade { 2, 24, 25, 35, 36, 37, 48, 51, 52, 53, 54, 55, 56, 57, 58, 59 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple

A grade { 1, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 51, 54, 55, 56, 59, 60, 62 }

B grade { 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 47, 48, 52, 53, 58, 61, 63, 64 }

C grade { 2 }

F normal fail { 57 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 5, 14, 18, 21, 24, 25, 29, 30, 47, 48, 59 }

B grade { 2, 3, 4, 6, 7, 8, 9, 10, 15, 16, 17, 19, 20, 22, 23, 26, 27, 28, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 60, 61, 62, 63, 64 }

C grade { 1, 11, 12, 13 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 1, 2, 3, 4, 5, 6, 7, 10, 12, 13, 14, 15, 16, 17, 18, 59 }

B grade { 8, 9, 11, 40, 43 }

C grade { 39, 42 }

F normal fail { 21, 22, 24, 25, 27, 29, 30, 31, 32, 38, 41, 45, 47, 48, 52, 53, 57, 58, 60, 61, 62, 63, 64 }

F(-1) timeout fail { }

F(-2) exception fail { 19, 20, 23, 26, 28, 33, 34, 35, 36, 37, 44, 46, 49, 50, 51, 54, 55, 56 }

Giac

A grade { 1, 2, 5, 6, 7, 8, 9, 14, 15, 16, 17, 18, 59, 62, 63 }

B grade { 3, 4, 10, 20, 21, 23, 24, 25, 30, 34, 35, 36, 37, 38, 41, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 60, 61, 64 }

C grade { 11, 12, 13, 39, 40 }

F normal fail { }

F(-1) timeout fail { 42 }

F(-2) exception fail { 19, 22, 26, 27, 28, 29, 31, 32, 33 }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 26, 28, 29, 34, 38, 39, 40, 42, 43, 44, 46, 47, 49, 51, 52, 54, 56, 57, 59 }

C grade { }

F normal fail { }

F(-1) timeout fail { 15, 22, 23, 24, 25, 27, 30, 31, 32, 33, 35, 36, 37, 41, 45, 48, 50, 53, 55, 58, 60, 61, 62, 63, 64 }

F(-2) exception fail { }

Sympy

A grade { 2, 3, 4, 10, 16, 51, 56, 59 }

B grade { 5, 6, 7 }

C grade { }

F normal fail { 1, 8, 9, 11, 12, 13, 14, 15, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 57, 58, 60, 61, 62, 63, 64 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	233	233	148	160	179	465	0	248	828
N.S.	1	1.00	0.64	0.69	0.77	2.00	0.00	1.06	3.55
time (sec)	N/A	0.370	0.905	2.079	0.322	0.272	0.000	0.473	13.647

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	34	29	23	48	22	40	20
N.S.	1	1.00	1.70	1.45	1.15	2.40	1.10	2.00	1.00
time (sec)	N/A	0.016	0.016	0.023	0.321	0.306	0.065	0.274	13.255

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	71	48	63	127	68	114	45
N.S.	1	1.00	1.51	1.02	1.34	2.70	1.45	2.43	0.96
time (sec)	N/A	0.046	1.259	0.054	0.299	0.271	0.096	0.320	0.121

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	111	77	112	253	126	229	76
N.S.	1	1.00	1.42	0.99	1.44	3.24	1.62	2.94	0.97
time (sec)	N/A	0.059	2.954	0.091	0.305	0.265	0.134	0.378	12.915

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	49	56	48	252	238	65	41
N.S.	1	1.00	1.00	1.14	0.98	5.14	4.86	1.33	0.84
time (sec)	N/A	0.086	0.070	0.076	0.323	0.284	0.740	0.305	0.118

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	97	97	90	99	115	534	2125	123	119
N.S.	1	1.00	0.93	1.02	1.19	5.51	21.91	1.27	1.23
time (sec)	N/A	0.125	1.021	0.118	0.324	0.305	9.066	0.316	12.401

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	150	150	138	148	228	1068	8964	206	4866
N.S.	1	1.00	0.92	0.99	1.52	7.12	59.76	1.37	32.44
time (sec)	N/A	0.194	0.352	0.210	0.324	0.328	48.202	0.398	16.089

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	22	22	51	19	300	91	0	32	18
N.S.	1	1.00	2.32	0.86	13.64	4.14	0.00	1.45	0.82
time (sec)	N/A	0.022	0.134	0.098	0.373	0.270	0.000	0.265	13.184

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	5	5	28	6	35	53	0	10	5
N.S.	1	1.00	5.60	1.20	7.00	10.60	0.00	2.00	1.00
time (sec)	N/A	0.020	0.033	0.033	0.360	0.293	0.000	0.273	12.971

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	12	12	12	13	10	21	14	28	12
N.S.	1	1.00	1.00	1.08	0.83	1.75	1.17	2.33	1.00
time (sec)	N/A	0.023	0.030	0.018	0.314	0.260	0.187	0.280	13.109

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	C	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	48	32	284	73	0	34	31
N.S.	1	1.00	1.37	0.91	8.11	2.09	0.00	0.97	0.89
time (sec)	N/A	0.030	0.115	0.039	0.390	0.271	0.000	0.268	13.017

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	C	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	30	15	17	19	0	11	14
N.S.	1	1.00	2.14	1.07	1.21	1.36	0.00	0.79	1.00
time (sec)	N/A	0.026	0.039	0.042	0.338	0.276	0.000	0.280	12.874

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	14	14	14	15	12	14	0	28	13
N.S.	1	1.00	1.00	1.07	0.86	1.00	0.00	2.00	0.93
time (sec)	N/A	0.025	0.030	0.032	0.315	0.255	0.000	0.265	13.203

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	19	29	24	27	0	25	17
N.S.	1	1.00	0.68	1.04	0.86	0.96	0.00	0.89	0.61
time (sec)	N/A	0.121	0.030	0.079	0.237	0.263	0.000	0.278	13.096

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	32	38	27	77	0	49	0
N.S.	1	1.00	1.03	1.23	0.87	2.48	0.00	1.58	0.00
time (sec)	N/A	0.118	0.152	0.044	0.373	0.276	0.000	0.291	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	8	27	12	11	10
N.S.	1	1.00	1.00	1.10	0.80	2.70	1.20	1.10	1.00
time (sec)	N/A	0.061	0.032	0.037	0.232	0.274	0.541	0.264	12.978

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	19	39	52	78	0	12	20
N.S.	1	1.00	0.53	1.08	1.44	2.17	0.00	0.33	0.56
time (sec)	N/A	0.097	0.027	0.419	0.346	0.278	0.000	0.272	13.185

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	19	24	18	35	0	25	34
N.S.	1	1.00	0.66	0.83	0.62	1.21	0.00	0.86	1.17
time (sec)	N/A	0.121	0.141	0.242	0.319	0.271	0.000	0.266	13.354

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	66	66	65	84	0	330	0	0	66
N.S.	1	1.00	0.98	1.27	0.00	5.00	0.00	0.00	1.00
time (sec)	N/A	0.141	0.203	0.163	0.000	0.342	0.000	0.000	15.109

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	48	48	48	71	0	248	0	95	53
N.S.	1	1.00	1.00	1.48	0.00	5.17	0.00	1.98	1.10
time (sec)	N/A	0.087	0.034	0.032	0.000	0.343	0.000	0.340	13.957

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	60	194	0	351	0	187	69
N.S.	1	1.00	1.00	3.23	0.00	5.85	0.00	3.12	1.15
time (sec)	N/A	0.114	0.034	7.154	0.000	0.287	0.000	0.325	13.781

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	140	174	0	768	0	0	0
N.S.	1	1.00	1.57	1.96	0.00	8.63	0.00	0.00	0.00
time (sec)	N/A	0.146	4.414	0.056	0.000	0.312	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	78	137	0	515	0	210	0
N.S.	1	1.00	1.20	2.11	0.00	7.92	0.00	3.23	0.00
time (sec)	N/A	0.053	0.173	0.046	0.000	0.284	0.000	0.498	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	51	51	44	312	0	193	0	239	0
N.S.	1	1.00	0.86	6.12	0.00	3.78	0.00	4.69	0.00
time (sec)	N/A	0.103	0.102	0.870	0.000	0.310	0.000	0.294	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	174	495	0	239	0	476	0
N.S.	1	1.00	2.05	5.82	0.00	2.81	0.00	5.60	0.00
time (sec)	N/A	0.168	1.763	1.508	0.000	0.329	0.000	0.329	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	91	150	0	486	0	0	120
N.S.	1	1.00	1.03	1.70	0.00	5.52	0.00	0.00	1.36
time (sec)	N/A	0.164	0.607	0.038	0.000	0.353	0.000	0.000	24.407

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	127	253	286	0	1134	0	0	0
N.S.	1	1.00	1.99	2.25	0.00	8.93	0.00	0.00	0.00
time (sec)	N/A	0.268	1.316	0.031	0.000	0.335	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	63	136	0	330	0	0	70
N.S.	1	1.00	0.91	1.97	0.00	4.78	0.00	0.00	1.01
time (sec)	N/A	0.108	0.188	0.024	0.000	0.333	0.000	0.000	16.991

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	F(-2)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	75	341	0	565	0	0	506
N.S.	1	1.00	1.00	4.55	0.00	7.53	0.00	0.00	6.75
time (sec)	N/A	0.145	0.087	1.279	0.000	0.790	0.000	0.000	0.260

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	80	80	222	642	0	543	0	625	0
N.S.	1	1.00	2.78	8.02	0.00	6.79	0.00	7.81	0.00
time (sec)	N/A	0.139	0.815	0.821	0.000	0.761	0.000	18.701	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	171	169	462	0	1520	0	0	0
N.S.	1	1.00	0.99	2.70	0.00	8.89	0.00	0.00	0.00
time (sec)	N/A	0.231	0.885	0.185	0.000	0.323	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	143	298	0	1071	0	0	0
N.S.	1	1.00	1.13	2.37	0.00	8.50	0.00	0.00	0.00
time (sec)	N/A	0.118	0.499	0.035	0.000	0.321	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	107	170	0	703	0	0	0
N.S.	1	1.00	1.23	1.95	0.00	8.08	0.00	0.00	0.00
time (sec)	N/A	0.070	0.077	0.048	0.000	0.302	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	111	68	0	239	0	97	41
N.S.	1	1.00	2.36	1.45	0.00	5.09	0.00	2.06	0.87
time (sec)	N/A	0.047	0.482	0.111	0.000	0.315	0.000	1.323	13.726

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	85	85	231	102	0	526	0	300	0
N.S.	1	1.00	2.72	1.20	0.00	6.19	0.00	3.53	0.00
time (sec)	N/A	0.085	3.719	0.046	0.000	0.336	0.000	0.846	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	135	135	367	162	0	898	0	1160	0
N.S.	1	1.00	2.72	1.20	0.00	6.65	0.00	8.59	0.00
time (sec)	N/A	0.141	8.138	0.039	0.000	0.354	0.000	1.136	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	190	190	2553	253	0	1452	0	3249	0
N.S.	1	1.00	13.44	1.33	0.00	7.64	0.00	17.10	0.00
time (sec)	N/A	0.270	15.969	0.042	0.000	0.420	0.000	2.089	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	123	51	0	110	0	257	104
N.S.	1	1.00	2.28	0.94	0.00	2.04	0.00	4.76	1.93
time (sec)	N/A	0.060	0.505	0.090	0.000	0.310	0.000	0.330	14.224

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	32	32	62	34	507	68	0	170	88
N.S.	1	1.00	1.94	1.06	15.84	2.12	0.00	5.31	2.75
time (sec)	N/A	0.029	0.104	0.078	0.848	0.276	0.000	0.285	14.192

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	C	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	28	28	26	31	90	56	0	34	85
N.S.	1	1.00	0.93	1.11	3.21	2.00	0.00	1.21	3.04
time (sec)	N/A	0.026	0.056	0.081	0.481	0.306	0.000	0.270	13.177

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	121	48	0	170	0	179	0
N.S.	1	1.00	1.98	0.79	0.00	2.79	0.00	2.93	0.00
time (sec)	N/A	0.056	0.255	0.041	0.000	0.282	0.000	0.520	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F	F(-1)	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	60	35	941	123	0	0	34
N.S.	1	1.00	1.43	0.83	22.40	2.93	0.00	0.00	0.81
time (sec)	N/A	0.035	0.090	0.030	0.692	0.294	0.000	0.000	13.918

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	B	B
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	48	21	143	60	0	45	20
N.S.	1	1.00	1.85	0.81	5.50	2.31	0.00	1.73	0.77
time (sec)	N/A	0.020	0.073	0.063	0.530	0.301	0.000	0.301	13.907

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	52	52	52	44	0	284	0	96	44
N.S.	1	1.00	1.00	0.85	0.00	5.46	0.00	1.85	0.85
time (sec)	N/A	0.127	0.191	0.037	0.000	0.352	0.000	0.364	14.609

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	64	64	158	80	0	588	0	229	0
N.S.	1	1.00	2.47	1.25	0.00	9.19	0.00	3.58	0.00
time (sec)	N/A	0.125	0.353	0.035	0.000	0.318	0.000	0.544	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	33	29	0	127	0	61	27
N.S.	1	1.00	1.00	0.88	0.00	3.85	0.00	1.85	0.82
time (sec)	N/A	0.077	0.027	0.078	0.000	0.285	0.000	0.328	13.513

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	60	149	0	419	0	203	93
N.S.	1	1.00	1.00	2.48	0.00	6.98	0.00	3.38	1.55
time (sec)	N/A	0.133	0.056	0.802	0.000	0.321	0.000	0.372	13.099

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	134	283	0	229	0	216	0
N.S.	1	1.00	2.48	5.24	0.00	4.24	0.00	4.00	0.00
time (sec)	N/A	0.110	0.852	0.914	0.000	0.330	0.000	0.337	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	59	68	0	385	0	109	52
N.S.	1	1.00	1.00	1.15	0.00	6.53	0.00	1.85	0.88
time (sec)	N/A	0.154	0.259	0.057	0.000	0.310	0.000	0.320	14.397

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	137	99	0	388	0	159	0
N.S.	1	1.00	2.32	1.68	0.00	6.58	0.00	2.69	0.00
time (sec)	N/A	0.108	0.806	0.058	0.000	0.368	0.000	0.339	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	44	56	0	344	88	105	47
N.S.	1	1.00	0.80	1.02	0.00	6.25	1.60	1.91	0.85
time (sec)	N/A	0.099	0.052	0.038	0.000	0.309	4.816	0.305	14.409

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	B
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	84	84	75	517	0	863	0	295	1451
N.S.	1	1.00	0.89	6.15	0.00	10.27	0.00	3.51	17.27
time (sec)	N/A	0.158	0.065	1.522	0.000	0.369	0.000	0.354	0.479

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	92	92	674	631	0	393	0	359	0
N.S.	1	1.00	7.33	6.86	0.00	4.27	0.00	3.90	0.00
time (sec)	N/A	0.196	6.947	2.931	0.000	0.354	0.000	0.328	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	69	88	0	698	0	219	88
N.S.	1	1.00	0.84	1.07	0.00	8.51	0.00	2.67	1.07
time (sec)	N/A	0.166	0.110	0.062	0.000	0.309	0.000	0.294	16.211

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	F	B	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	197	161	0	720	0	281	0
N.S.	1	1.00	2.10	1.71	0.00	7.66	0.00	2.99	0.00
time (sec)	N/A	0.151	6.340	0.060	0.000	0.386	0.000	0.340	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F(-2)	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	47	75	0	627	110	215	82
N.S.	1	1.00	0.60	0.96	0.00	8.04	1.41	2.76	1.05
time (sec)	N/A	0.115	0.056	0.046	0.000	0.337	7.393	0.320	17.273

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	B	F	B	B
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	118	118	78	0	0	1531	0	483	2817
N.S.	1	1.00	0.66	0.00	0.00	12.97	0.00	4.09	23.87
time (sec)	N/A	0.224	0.074	0.000	0.000	0.488	0.000	0.383	13.682

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	B	F(-1)
verified	N/A	Yes	No	No	TBD	TBD	TBD	TBD	TBD
size	141	141	1450	934	0	647	0	537	0
N.S.	1	1.00	10.28	6.62	0.00	4.59	0.00	3.81	0.00
time (sec)	N/A	0.296	8.162	1.563	0.000	0.343	0.000	0.355	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	57	31	33	24	34	34	37
N.S.	1	1.00	1.54	0.84	0.89	0.65	0.92	0.92	1.00
time (sec)	N/A	0.076	0.047	0.135	0.294	0.289	0.124	0.264	13.795

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	86	139	0	1063	0	204	0
N.S.	1	1.00	0.96	1.54	0.00	11.81	0.00	2.27	0.00
time (sec)	N/A	0.170	0.187	0.242	0.000	0.451	0.000	0.303	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	126	126	167	245	0	1486	0	445	0
N.S.	1	1.00	1.33	1.94	0.00	11.79	0.00	3.53	0.00
time (sec)	N/A	0.241	4.810	0.071	0.000	0.483	0.000	0.539	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	41	65	0	264	0	58	0
N.S.	1	1.00	1.00	1.59	0.00	6.44	0.00	1.41	0.00
time (sec)	N/A	0.078	0.022	0.202	0.000	0.385	0.000	0.320	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	73	248	0	670	0	111	0
N.S.	1	1.00	0.99	3.35	0.00	9.05	0.00	1.50	0.00
time (sec)	N/A	0.128	0.332	1.025	0.000	0.451	0.000	0.290	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	114	586	0	1365	0	276	0
N.S.	1	1.00	0.97	5.01	0.00	11.67	0.00	2.36	0.00
time (sec)	N/A	0.224	0.831	0.135	0.000	0.517	0.000	0.313	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [59] had the largest ratio of [.750000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	15	12	1.00	25	0.480
2	A	3	2	1.00	12	0.167
3	A	4	3	1.00	14	0.214
4	A	4	3	1.00	14	0.214
5	A	3	3	1.00	14	0.214
6	A	5	5	1.00	14	0.357
7	A	6	6	1.00	14	0.429
8	A	4	4	1.00	10	0.400
9	A	3	3	1.00	10	0.300
10	A	3	3	1.00	10	0.300
11	A	5	5	1.00	12	0.417
12	A	4	4	1.00	12	0.333
13	A	3	3	1.00	12	0.250
14	A	4	3	1.00	17	0.176
15	A	5	5	1.00	17	0.294
16	A	3	3	1.00	15	0.200
17	A	5	5	1.00	15	0.333
18	A	5	4	1.00	17	0.235
19	A	6	6	1.00	17	0.353
20	A	5	5	1.00	15	0.333
21	A	7	5	1.00	15	0.333
22	A	7	7	1.00	17	0.412
23	A	6	6	1.00	12	0.500
24	A	5	5	1.00	17	0.294

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	6	6	1.00	17	0.353
26	A	7	6	1.00	17	0.353
27	A	8	8	1.00	17	0.471
28	A	6	5	1.00	15	0.333
29	A	8	6	1.00	15	0.400
30	A	7	7	1.00	17	0.412
31	A	8	8	1.00	16	0.500
32	A	7	7	1.00	16	0.438
33	A	6	6	1.00	16	0.375
34	A	3	3	1.00	16	0.188
35	A	4	4	1.00	16	0.250
36	A	6	6	1.00	16	0.375
37	A	7	6	1.00	16	0.375
38	A	6	6	1.00	12	0.500
39	A	5	5	1.00	12	0.417
40	A	3	3	1.00	12	0.250
41	A	7	6	1.00	10	0.600
42	A	6	5	1.00	10	0.500
43	A	3	3	1.00	10	0.300
44	A	5	5	1.00	17	0.294
45	A	6	6	1.00	17	0.353
46	A	4	4	1.00	15	0.267
47	A	7	5	1.00	15	0.333
48	A	5	5	1.00	17	0.294
49	A	5	5	1.00	17	0.294
50	A	4	4	1.00	17	0.235
51	A	5	5	1.00	15	0.333
52	A	8	6	1.00	15	0.400
53	A	6	6	1.00	17	0.353
54	A	6	6	1.00	17	0.353
55	A	6	6	1.00	17	0.353
56	A	6	5	1.00	15	0.333
57	A	9	7	1.00	15	0.467
58	A	7	7	1.00	17	0.412
59	A	7	6	1.00	8	0.750

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	8	7	1.00	15	0.467
61	A	9	8	1.00	15	0.533
62	A	4	4	1.00	15	0.267
63	A	6	6	1.00	15	0.400
64	A	7	7	1.00	15	0.467

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{A+C \cot^2(c+dx)}{\sqrt{b \tan(c+dx)}} dx \dots\dots\dots$	46
3.2	$\int (a + b \cot^2(c + dx)) dx \dots\dots\dots$	55
3.3	$\int (a + b \cot^2(c + dx))^2 dx \dots\dots\dots$	59
3.4	$\int (a + b \cot^2(c + dx))^3 dx \dots\dots\dots$	64
3.5	$\int \frac{1}{a+b \cot^2(c+dx)} dx \dots\dots\dots$	69
3.6	$\int \frac{1}{(a+b \cot^2(c+dx))^2} dx \dots\dots\dots$	74
3.7	$\int \frac{1}{(a+b \cot^2(c+dx))^3} dx \dots\dots\dots$	80
3.8	$\int (1 + \cot^2(x))^{3/2} dx \dots\dots\dots$	93
3.9	$\int \sqrt{1 + \cot^2(x)} dx \dots\dots\dots$	97
3.10	$\int \frac{1}{\sqrt{1+\cot^2(x)}} dx \dots\dots\dots$	101
3.11	$\int (-1 - \cot^2(x))^{3/2} dx \dots\dots\dots$	105
3.12	$\int \sqrt{-1 - \cot^2(x)} dx \dots\dots\dots$	110
3.13	$\int \frac{1}{\sqrt{-1-\cot^2(x)}} dx \dots\dots\dots$	114
3.14	$\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx \dots\dots\dots$	118
3.15	$\int \frac{\cot^2(x)}{\sqrt{a+a \cot^2(x)}} dx \dots\dots\dots$	122
3.16	$\int \frac{\cot(x)}{\sqrt{a+a \cot^2(x)}} dx \dots\dots\dots$	126
3.17	$\int \frac{\tan(x)}{\sqrt{a+a \cot^2(x)}} dx \dots\dots\dots$	130
3.18	$\int \frac{\tan^2(x)}{\sqrt{a+a \cot^2(x)}} dx \dots\dots\dots$	135
3.19	$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx \dots\dots\dots$	139
3.20	$\int \cot(x) \sqrt{a + b \cot^2(x)} dx \dots\dots\dots$	144
3.21	$\int \sqrt{a + b \cot^2(x)} \tan(x) dx \dots\dots\dots$	149
3.22	$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx \dots\dots\dots$	155
3.23	$\int \sqrt{a + b \cot^2(x)} dx \dots\dots\dots$	161

3.24	$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx$	167
3.25	$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx$	172
3.26	$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx$	179
3.27	$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx$	185
3.28	$\int \cot(x) (a + b \cot^2(x))^{3/2} dx$	191
3.29	$\int (a + b \cot^2(x))^{3/2} \tan(x) dx$	196
3.30	$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx$	203
3.31	$\int (a + b \cot^2(c + dx))^{5/2} dx$	210
3.32	$\int (a + b \cot^2(c + dx))^{3/2} dx$	217
3.33	$\int \sqrt{a + b \cot^2(c + dx)} dx$	223
3.34	$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx$	228
3.35	$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx$	233
3.36	$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx$	238
3.37	$\int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx$	245
3.38	$\int (1 - \cot^2(x))^{3/2} dx$	255
3.39	$\int \sqrt{1 - \cot^2(x)} dx$	260
3.40	$\int \frac{1}{\sqrt{1 - \cot^2(x)}} dx$	265
3.41	$\int (-1 + \cot^2(x))^{3/2} dx$	269
3.42	$\int \sqrt{-1 + \cot^2(x)} dx$	274
3.43	$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx$	279
3.44	$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx$	283
3.45	$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx$	288
3.46	$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx$	293
3.47	$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx$	297
3.48	$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx$	303
3.49	$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx$	308
3.50	$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx$	313
3.51	$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx$	318
3.52	$\int \frac{\tan(x)}{(a + b \cot^2(x))^{3/2}} dx$	323
3.53	$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx$	330
3.54	$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx$	336
3.55	$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{5/2}} dx$	342
3.56	$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx$	348
3.57	$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx$	354

3.58	$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{5/2}} dx$	362
3.59	$\int \frac{1}{1+\cot^3(x)} dx$	370
3.60	$\int \cot(x) \sqrt{a + b \cot^4(x)} dx$	375
3.61	$\int \cot(x) (a + b \cot^4(x))^{3/2} dx$	381
3.62	$\int \frac{\cot(x)}{\sqrt{a+b \cot^4(x)}} dx$	388
3.63	$\int \frac{\cot(x)}{(a+b \cot^4(x))^{3/2}} dx$	393
3.64	$\int \frac{\cot(x)}{(a+b \cot^4(x))^{5/2}} dx$	399

3.1 $\int \frac{A+C \cot^2(c+dx)}{\sqrt{b \tan(c+dx)}} dx$

Optimal result	46
Rubi [A] (verified)	47
Mathematica [A] (verified)	50
Maple [A] (verified)	51
Fricas [C] (verification not implemented)	51
Sympy [F]	52
Maxima [A] (verification not implemented)	52
Giac [A] (verification not implemented)	53
Mupad [B] (verification not implemented)	54

Optimal result

Integrand size = 25, antiderivative size = 233

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx = -\frac{(A - C) \arctan\left(1 - \frac{\sqrt{2}\sqrt{b \tan(c + dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} + \frac{(A - C) \arctan\left(1 + \frac{\sqrt{2}\sqrt{b \tan(c + dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} - \frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) - \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} + \frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) + \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} - \frac{2bC}{3d(b \tan(c + dx))^{3/2}}$$

```
[Out] -1/2*(A-C)*arctan(1-2^(1/2)*(b*tan(d*x+c))^(1/2)/b^(1/2))/d*2^(1/2)/b^(1/2)
+1/2*(A-C)*arctan(1+2^(1/2)*(b*tan(d*x+c))^(1/2)/b^(1/2))/d*2^(1/2)/b^(1/2)
-1/4*(A-C)*ln(b^(1/2)-2^(1/2)*(b*tan(d*x+c))^(1/2)+b^(1/2)*tan(d*x+c))/d*2^(1/2)/b^(1/2)
+1/4*(A-C)*ln(b^(1/2)+2^(1/2)*(b*tan(d*x+c))^(1/2)+b^(1/2)*tan(d*x+c))/d*2^(1/2)/b^(1/2)
-2/3*b*C/d/(b*tan(d*x+c))^(3/2)
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 233, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.480$, Rules used = {3754, 3710, 12, 16, 3557, 335, 217, 1179, 642, 1176, 631, 210}

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx = -\frac{(A - C) \arctan\left(1 - \frac{\sqrt{2}\sqrt{b \tan(c + dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} + \frac{(A - C) \arctan\left(\frac{\sqrt{2}\sqrt{b \tan(c + dx)}}{\sqrt{b}} + 1\right)}{\sqrt{2}\sqrt{bd}} - \frac{(A - C) \log\left(\sqrt{b} \tan(c + dx) - \sqrt{2}\sqrt{b \tan(c + dx)} + \sqrt{b}\right)}{2\sqrt{2}\sqrt{bd}} + \frac{(A - C) \log\left(\sqrt{b} \tan(c + dx) + \sqrt{2}\sqrt{b \tan(c + dx)} + \sqrt{b}\right)}{2\sqrt{2}\sqrt{bd}} - \frac{2bC}{3d(b \tan(c + dx))^{3/2}}$$

[In] Int[(A + C*Cot[c + d*x]^2)/Sqrt[b*Tan[c + d*x]],x]

[Out] -(((A - C)*ArcTan[1 - (Sqrt[2]*Sqrt[b*Tan[c + d*x]])/Sqrt[b]])/(Sqrt[2]*Sqrt[b]*d)) + ((A - C)*ArcTan[1 + (Sqrt[2]*Sqrt[b*Tan[c + d*x]])/Sqrt[b]])/(Sqrt[2]*Sqrt[b]*d) - ((A - C)*Log[Sqrt[b] + Sqrt[b]*Tan[c + d*x] - Sqrt[2]*Sqrt[b*Tan[c + d*x]])/(2*Sqrt[2]*Sqrt[b]*d) + ((A - C)*Log[Sqrt[b] + Sqrt[b]*Tan[c + d*x] + Sqrt[2]*Sqrt[b*Tan[c + d*x]])/(2*Sqrt[2]*Sqrt[b]*d) - (2*b*C)/(3*d*(b*Tan[c + d*x])^(3/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```


Rule 3710

Int[((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((A_.) + (C_.)*tan[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[(A*b^2 + a^2*C)*((a + b*Tan[e + f*x])^(m + 1)/(b*f*(m + 1)*(a^2 + b^2))), x] + Dist[1/(a^2 + b^2), Int[(a + b*Tan[e + f*x])^(m + 1)*Simp[a*(A - C) - (A*b - b*C)*Tan[e + f*x], x], x], x] /; FreeQ[{a, b, e, f, A, C}, x] && NeQ[A*b^2 + a^2*C, 0] && LtQ[m, -1] && NeQ[a^2 + b^2, 0]

Rule 3754

Int[(cot[(e_.) + (f_.)*(x_)]*(d_.))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^p, x_Symbol] := Dist[d^(n*p), Int[(d*Cot[e + f*x])^(m - n*p)*(b + a*Cot[e + f*x])^n]^p, x], x] /; FreeQ[{a, b, d, e, f, m, n, p}, x] && !IntegerQ[m] && IntegerQ[n, p]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^2 \int \frac{C + A \tan^2(c + dx)}{(b \tan(c + dx))^{5/2}} dx \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \int \frac{b(A - C) \tan(c + dx)}{(b \tan(c + dx))^{3/2}} dx \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + (b(A - C)) \int \frac{\tan(c + dx)}{(b \tan(c + dx))^{3/2}} dx \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + (A - C) \int \frac{1}{\sqrt{b \tan(c + dx)}} dx \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \frac{(b(A - C)) \text{Subst}\left(\int \frac{1}{\sqrt{x(b^2 + x^2)}} dx, x, b \tan(c + dx)\right)}{d} \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \frac{(2b(A - C)) \text{Subst}\left(\int \frac{1}{b^2 + x^4} dx, x, \sqrt{b \tan(c + dx)}\right)}{d} \\
 &= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \frac{(A - C) \text{Subst}\left(\int \frac{b - x^2}{b^2 + x^4} dx, x, \sqrt{b \tan(c + dx)}\right)}{d} \\
 &\quad + \frac{(A - C) \text{Subst}\left(\int \frac{b + x^2}{b^2 + x^4} dx, x, \sqrt{b \tan(c + dx)}\right)}{d}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \frac{(A - C) \text{Subst}\left(\int \frac{1}{b - \sqrt{2}\sqrt{bx+x^2}} dx, x, \sqrt{b \tan(c + dx)}\right)}{2d} \\
&+ \frac{(A - C) \text{Subst}\left(\int \frac{1}{b + \sqrt{2}\sqrt{bx+x^2}} dx, x, \sqrt{b \tan(c + dx)}\right)}{2d} \\
&- \frac{(A - C) \text{Subst}\left(\int \frac{\sqrt{2}\sqrt{b+2x}}{-b - \sqrt{2}\sqrt{bx-x^2}} dx, x, \sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&- \frac{(A - C) \text{Subst}\left(\int \frac{\sqrt{2}\sqrt{b-2x}}{-b + \sqrt{2}\sqrt{bx-x^2}} dx, x, \sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&= -\frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) - \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&+ \frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) + \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&- \frac{2bC}{3d(b \tan(c + dx))^{3/2}} + \frac{(A - C) \text{Subst}\left(\int \frac{1}{-1-x^2} dx, x, 1 - \frac{\sqrt{2}\sqrt{b \tan(c+dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} \\
&- \frac{(A - C) \text{Subst}\left(\int \frac{1}{-1-x^2} dx, x, 1 + \frac{\sqrt{2}\sqrt{b \tan(c+dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} \\
&= -\frac{(A - C) \arctan\left(1 - \frac{\sqrt{2}\sqrt{b \tan(c+dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} + \frac{(A - C) \arctan\left(1 + \frac{\sqrt{2}\sqrt{b \tan(c+dx)}}{\sqrt{b}}\right)}{\sqrt{2}\sqrt{bd}} \\
&- \frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) - \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&+ \frac{(A - C) \log\left(\sqrt{b} + \sqrt{b} \tan(c + dx) + \sqrt{2}\sqrt{b \tan(c + dx)}\right)}{2\sqrt{2}\sqrt{bd}} \\
&- \frac{2bC}{3d(b \tan(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.91 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.64

$$\begin{aligned}
&\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx \\
&= \frac{-8C \cot(c + dx) - 3\sqrt{2}(A - C) \left(2 \arctan\left(1 - \sqrt{2}\sqrt{\tan(c + dx)}\right) - 2 \arctan\left(1 + \sqrt{2}\sqrt{\tan(c + dx)}\right)\right) + 1}{12d\sqrt{b \tan(c + dx)}}
\end{aligned}$$

[In] Integrate[(A + C*Cot[c + d*x]^2)/Sqrt[b*Tan[c + d*x]],x]

[Out] (-8*C*Cot[c + d*x] - 3*Sqrt[2]*(A - C)*(2*ArcTan[1 - Sqrt[2]*Sqrt[Tan[c + d*x]]) - 2*ArcTan[1 + Sqrt[2]*Sqrt[Tan[c + d*x]]) + Log[1 - Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]] - Log[1 + Sqrt[2]*Sqrt[Tan[c + d*x]] + Tan[c + d*x]])*Sqrt[Tan[c + d*x]])/(12*d*Sqrt[b*Tan[c + d*x]])

Maple [A] (verified)

Time = 2.08 (sec) , antiderivative size = 160, normalized size of antiderivative = 0.69

method	result
derivativedivides	$2b \left(-\frac{C}{3(b \tan(dx+c))^{\frac{3}{2}}} + \frac{(A-C)(b^2)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{b \tan(dx+c) + (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}}{b \tan(dx+c) - (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{b \tan(dx+c)} + 1}{(b^2)^{\frac{1}{4}}} \right) \right)}{8b^2} \right) \frac{1}{d}$
default	$2b \left(-\frac{C}{3(b \tan(dx+c))^{\frac{3}{2}}} + \frac{(A-C)(b^2)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{b \tan(dx+c) + (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}}{b \tan(dx+c) - (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{b \tan(dx+c)} + 1}{(b^2)^{\frac{1}{4}}} \right) \right)}{8b^2} \right) \frac{1}{d}$
parts	$\frac{A(b^2)^{\frac{1}{4}} \sqrt{2} \left(\ln \left(\frac{b \tan(dx+c) + (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}}{b \tan(dx+c) - (b^2)^{\frac{1}{4}} \sqrt{b \tan(dx+c)} \sqrt{2} + \sqrt{b^2}} \right) + 2 \arctan \left(\frac{\sqrt{2} \sqrt{b \tan(dx+c)} + 1}{(b^2)^{\frac{1}{4}}} \right) - 2 \arctan \left(-\frac{\sqrt{2} \sqrt{b \tan(dx+c)}}{(b^2)^{\frac{1}{4}}} \right) \right)}{4db}$

[In] int((A+C*cot(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/d*b*(-1/3*C/(b*tan(d*x+c))^(3/2)+1/8*(A-C)*(b^2)^(1/4)/b^2*2^(1/2)*(ln((b*tan(d*x+c)+(b^2)^(1/4)*(b*tan(d*x+c))^(1/2)*2^(1/2)+(b^2)^(1/2)))/(b*tan(d*x+c)-(b^2)^(1/4)*(b*tan(d*x+c))^(1/2)*2^(1/2)+(b^2)^(1/2)))+2*arctan(2^(1/2)/(b^2)^(1/4)*(b*tan(d*x+c))^(1/2)+1)-2*arctan(-2^(1/2)/(b^2)^(1/4)*(b*tan(d*x+c))^(1/2)+1)))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 465, normalized size of antiderivative = 2.00

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx =$$

$$\frac{3bd \left(-\frac{A^4 - 4A^3C + 6A^2C^2 - 4AC^3 + C^4}{b^2d^4} \right)^{\frac{1}{4}} \log \left(bd \left(-\frac{A^4 - 4A^3C + 6A^2C^2 - 4AC^3 + C^4}{b^2d^4} \right)^{\frac{1}{4}} - \sqrt{b \tan(dx + c)}(A - C) \right)}{1}$$

[In] integrate((A+C*cot(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x, algorithm="fricas")

[Out]
$$-1/6*(3*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4)*\log(b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4) - \sqrt{b*\tan(d*x + c)}*(A - C))*\tan(d*x + c)^2 + 3*I*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4)*\log(I*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4) - \sqrt{b*\tan(d*x + c)}*(A - C))*\tan(d*x + c)^2 - 3*I*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4)*\log(-I*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4) - \sqrt{b*\tan(d*x + c)}*(A - C))*\tan(d*x + c)^2 - 3*b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4)*\log(-b*d*(-(A^4 - 4*A^3*C + 6*A^2*C^2 - 4*A*C^3 + C^4)/(b^2*d^4))^(1/4) - \sqrt{b*\tan(d*x + c)}*(A - C))*\tan(d*x + c)^2 + 4*\sqrt{b*\tan(d*x + c)}*C)/(b*d*\tan(d*x + c)^2)$$

Sympy [F]

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx = \int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx$$

[In] integrate((A+C*cot(d*x+c)**2)/(b*tan(d*x+c))**(1/2),x)

[Out] Integral((A + C*cot(c + d*x)**2)/sqrt(b*tan(c + d*x)), x)

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 179, normalized size of antiderivative = 0.77

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx = \frac{3 \left(2 \sqrt{2} \sqrt{b} \arctan \left(\frac{\sqrt{2}(\sqrt{2}\sqrt{b}+2\sqrt{b \tan(dx+c)})}{2\sqrt{b}} \right) + 2 \sqrt{2} \sqrt{b} \arctan \left(-\frac{\sqrt{2}(\sqrt{2}\sqrt{b}-2\sqrt{b \tan(dx+c)})}{2\sqrt{b}} \right) + \sqrt{2} \sqrt{b} \log(b \tan(dx+c)) \right)}{12}$$

[In] integrate((A+C*cot(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x, algorithm="maxima")

[Out]
$$1/12*(3*(2*\sqrt{2}*\sqrt{b}*\arctan(1/2*\sqrt{2}*(\sqrt{2}*\sqrt{b} + 2*\sqrt{b*\tan(d*x + c)}))/\sqrt{b}) + 2*\sqrt{2}*\sqrt{b}*\arctan(-1/2*\sqrt{2}*(\sqrt{2}*\sqrt{b} - 2*\sqrt{b*\tan(d*x + c)}))/\sqrt{b}) + \sqrt{2}*\sqrt{b}*\log(b*\tan(d*x + c)) + \sqrt{2}*\sqrt{b*\tan(d*x + c)}*\sqrt{b} + b) - \sqrt{2}*\sqrt{b}*\log(b*\tan(d*x + c) - \sqrt{2}*\sqrt{b*\tan(d*x + c)}*\sqrt{b} + b))*(A - C) - 8*C*b^2/(b*d*\tan(d*x + c))^(3/2))/(b*d)$$

Giac [A] (verification not implemented)

none

Time = 0.47 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.06

$$\begin{aligned}
& \int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx \\
&= \frac{\sqrt{2} \left(A \sqrt{|b|} - C \sqrt{|b|} \right) \arctan \left(\frac{\sqrt{2} \left(\sqrt{2} \sqrt{|b|} + 2 \sqrt{b \tan(dx+c)} \right)}{2 \sqrt{|b|}} \right)}{2bd} \\
&+ \frac{\sqrt{2} \left(A \sqrt{|b|} - C \sqrt{|b|} \right) \arctan \left(-\frac{\sqrt{2} \left(\sqrt{2} \sqrt{|b|} - 2 \sqrt{b \tan(dx+c)} \right)}{2 \sqrt{|b|}} \right)}{2bd} \\
&+ \frac{\sqrt{2} \left(A \sqrt{|b|} - C \sqrt{|b|} \right) \log \left(b \tan(dx+c) + \sqrt{2} \sqrt{b \tan(dx+c)} \sqrt{|b|} + |b| \right)}{4bd} \\
&- \frac{\sqrt{2} \left(A \sqrt{|b|} - C \sqrt{|b|} \right) \log \left(b \tan(dx+c) - \sqrt{2} \sqrt{b \tan(dx+c)} \sqrt{|b|} + |b| \right)}{4bd} \\
&- \frac{2C}{3 \sqrt{b \tan(dx+c)} d \tan(dx+c)}
\end{aligned}$$

```
[In] integrate((A+C*cot(d*x+c)^2)/(b*tan(d*x+c))^(1/2),x, algorithm="giac")
```

```
[Out] 1/2*sqrt(2)*(A*sqrt(abs(b)) - C*sqrt(abs(b)))*arctan(1/2*sqrt(2)*(sqrt(2)*sqrt(abs(b)) + 2*sqrt(b*tan(d*x + c)))/sqrt(abs(b)))/(b*d) + 1/2*sqrt(2)*(A*sqrt(abs(b)) - C*sqrt(abs(b)))*arctan(-1/2*sqrt(2)*(sqrt(2)*sqrt(abs(b)) - 2*sqrt(b*tan(d*x + c)))/sqrt(abs(b)))/(b*d) + 1/4*sqrt(2)*(A*sqrt(abs(b)) - C*sqrt(abs(b)))*log(b*tan(d*x + c) + sqrt(2)*sqrt(b*tan(d*x + c))*sqrt(abs(b) + abs(b)))/(b*d) - 1/4*sqrt(2)*(A*sqrt(abs(b)) - C*sqrt(abs(b)))*log(b*tan(d*x + c) - sqrt(2)*sqrt(b*tan(d*x + c))*sqrt(abs(b) + abs(b)))/(b*d) - 2/3*C/(sqrt(b*tan(d*x + c))*d*tan(d*x + c))
```

Mupad [B] (verification not implemented)

Time = 13.65 (sec) , antiderivative size = 828, normalized size of antiderivative = 3.55

$$\int \frac{A + C \cot^2(c + dx)}{\sqrt{b \tan(c + dx)}} dx = -\frac{2 C b}{3 d (b \tan(c + dx))^{3/2}}$$

$$(-1)^{1/4} \operatorname{atan} \left(\frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d} \right) + \frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d}$$

$$+ \frac{(-1)^{1/4} \operatorname{atan} \left(\frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d} + \frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d}$$

$$+ \frac{(-1)^{1/4} \operatorname{atan} \left(\frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d} + \frac{(-1)^{1/4} (A-C) \left(\sqrt{b \tan(c+dx)} (16 A^2 b^2 d^3 - 32 A C b^2 d^3 + 16 C^2 b^2 d^3) - \frac{(-1)^{1/4} (A-C) (32 A b^3 d^4 - 32 C b^3 d^4)}{2 \sqrt{b} d} \right)}{2 \sqrt{b} d}$$

```
[In] int((A + C*cot(c + d*x)^2)/(b*tan(c + d*x))^(1/2),x)
```

```
[Out] ((-1)^(1/4)*atan((((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) - ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4)))/(2*b^(1/2)*d))*1i)/(2*b^(1/2)*d) + ((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) + ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4)))/(2*b^(1/2)*d))*1i)/(2*b^(1/2)*d))/((((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) - ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4)))/(2*b^(1/2)*d)))/(2*b^(1/2)*d) - ((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) + ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4)))/(2*b^(1/2)*d)))/(2*b^(1/2)*d)))*(A - C)*1i)/(b^(1/2)*d) - (2*C*b)/(3*d*(b*tan(c + d*x))^(3/2)) + ((-1)^(1/4)*atan((((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) - ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4))*1i)/(2*b^(1/2)*d)))/(2*b^(1/2)*d) + ((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) + ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4))*1i)/(2*b^(1/2)*d)))/(2*b^(1/2)*d))/((((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) - ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4))*1i)/(2*b^(1/2)*d)))/(2*b^(1/2)*d) - ((-1)^(1/4)*(A - C)*((b*tan(c + d*x))^(1/2)*(16*A^2*b^2*d^3 + 16*C^2*b^2*d^3 - 32*A*C*b^2*d^3) + ((-1)^(1/4)*(A - C)*(32*A*b^3*d^4 - 32*C*b^3*d^4))*1i)/(2*b^(1/2)*d)))/(2*b^(1/2)*d)))*(A - C)/(b^(1/2)*d)
```

3.2 $\int (a + b \cot^2(c + dx)) dx$

Optimal result	55
Rubi [A] (verified)	55
Mathematica [C] (verified)	56
Maple [C] (verified)	56
Fricas [B] (verification not implemented)	57
Sympy [A] (verification not implemented)	57
Maxima [A] (verification not implemented)	57
Giac [A] (verification not implemented)	58
Mupad [B] (verification not implemented)	58

Optimal result

Integrand size = 12, antiderivative size = 20

$$\int (a + b \cot^2(c + dx)) dx = ax - bx - \frac{b \cot(c + dx)}{d}$$

[Out] a*x-b*x-b*cot(d*x+c)/d

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3554, 8}

$$\int (a + b \cot^2(c + dx)) dx = ax - \frac{b \cot(c + dx)}{d} - bx$$

[In] Int[a + b*Cot[c + d*x]^2,x]

[Out] a*x - b*x - (b*Cot[c + d*x])/d

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned}
\text{integral} &= ax + b \int \cot^2(c + dx) dx \\
&= ax - \frac{b \cot(c + dx)}{d} - b \int 1 dx \\
&= ax - bx - \frac{b \cot(c + dx)}{d}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.02 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.70

$$\int (a + b \cot^2(c + dx)) dx = ax - \frac{b \cot(c + dx) \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, -\tan^2(c + dx)\right)}{d}$$

[In] Integrate[a + b*Cot[c + d*x]^2,x]

[Out] a*x - (b*Cot[c + d*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[c + d*x]^2])/d

Maple [C] (verified)

Result contains complex when optimal does not.

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.45

method	result	size
risch	$ax - bx - \frac{2ib}{d(e^{2i(dx+c)} - 1)}$	29
norman	$\frac{(a-b)x \tan(dx+c) - \frac{b}{d}}{\tan(dx+c)}$	30
parallelrisc	$\frac{b(-\tan(dx+c)xd-1)}{d \tan(dx+c)} + ax$	30
default	$ax + \frac{b(-\cot(dx+c) + \frac{\pi}{2} - \text{arccot}(\cot(dx+c)))}{d}$	31
parts	$ax + \frac{b(-\cot(dx+c) + \frac{\pi}{2} - \text{arccot}(\cot(dx+c)))}{d}$	31
derivativedivides	$\frac{-b \cot(dx+c) + (-a+b)(\frac{\pi}{2} - \text{arccot}(\cot(dx+c)))}{d}$	34

[In] int(a+b*cot(d*x+c)^2,x,method=_RETURNVERBOSE)

[Out] a*x-b*x-2*I*b/d/(exp(2*I*(d*x+c))-1)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 48 vs. 2(20) = 40.

Time = 0.31 (sec) , antiderivative size = 48, normalized size of antiderivative = 2.40

$$\int (a + b \cot^2(c + dx)) dx = \frac{(a - b)dx \sin(2dx + 2c) - b \cos(2dx + 2c) - b}{d \sin(2dx + 2c)}$$

[In] integrate(a+b*cot(d*x+c)^2,x, algorithm="fricas")

[Out] ((a - b)*d*x*sin(2*d*x + 2*c) - b*cos(2*d*x + 2*c) - b)/(d*sin(2*d*x + 2*c))

Sympy [A] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (a + b \cot^2(c + dx)) dx = ax + b \begin{cases} -x - \frac{\cot(c+dx)}{d} & \text{for } d \neq 0 \\ x \cot^2(c) & \text{otherwise} \end{cases}$$

[In] integrate(a+b*cot(d*x+c)**2,x)

[Out] a*x + b*Piecewise((-x - cot(c + d*x)/d, Ne(d, 0)), (x*cot(c)**2, True))

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.15

$$\int (a + b \cot^2(c + dx)) dx = ax - \frac{\left(dx + c + \frac{1}{\tan(dx+c)}\right)b}{d}$$

[In] integrate(a+b*cot(d*x+c)^2,x, algorithm="maxima")

[Out] a*x - (d*x + c + 1/tan(d*x + c))*b/d

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.00

$$\int (a + b \cot^2(c + dx)) dx = ax - \frac{\left(2 dx + 2 c + \frac{1}{\tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)} - \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)\right) b}{2 d}$$

[In] integrate(a+b*cot(d*x+c)^2,x, algorithm="giac")

[Out] a*x - 1/2*(2*d*x + 2*c + 1/tan(1/2*d*x + 1/2*c) - tan(1/2*d*x + 1/2*c))*b/d

Mupad [B] (verification not implemented)

Time = 13.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (a + b \cot^2(c + dx)) dx = x(a - b) - \frac{b \cot(c + dx)}{d}$$

[In] int(a + b*cot(c + d*x)^2,x)

[Out] x*(a - b) - (b*cot(c + d*x))/d

3.3 $\int (a + b \cot^2(c + dx))^2 dx$

Optimal result	59
Rubi [A] (verified)	59
Mathematica [A] (verified)	60
Maple [A] (verified)	61
Fricas [B] (verification not implemented)	61
Sympy [A] (verification not implemented)	62
Maxima [A] (verification not implemented)	62
Giac [B] (verification not implemented)	62
Mupad [B] (verification not implemented)	63

Optimal result

Integrand size = 14, antiderivative size = 47

$$\int (a + b \cot^2(c + dx))^2 dx = (a - b)^2 x - \frac{(2a - b)b \cot(c + dx)}{d} - \frac{b^2 \cot^3(c + dx)}{3d}$$

[Out] (a-b)^2*x-(2*a-b)*b*cot(d*x+c)/d-1/3*b^2*cot(d*x+c)^3/d

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 209}

$$\int (a + b \cot^2(c + dx))^2 dx = -\frac{b(2a - b) \cot(c + dx)}{d} + x(a - b)^2 - \frac{b^2 \cot^3(c + dx)}{3d}$$

[In] Int[(a + b*Cot[c + d*x]^2)^2,x]

[Out] (a - b)^2*x - ((2*a - b)*b*Cot[c + d*x])/d - (b^2*Cot[c + d*x]^3)/(3*d)

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 398

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0]

0] && GeQ[p, -q]

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{(a+bx^2)^2}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int \left((2a-b)b + b^2x^2 + \frac{(a-b)^2}{1+x^2}\right) dx, x, \cot(c+dx)\right)}{d} \\
 &= -\frac{(2a-b)b \cot(c+dx)}{d} - \frac{b^2 \cot^3(c+dx)}{3d} - \frac{(a-b)^2 \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\
 &= (a-b)^2 x - \frac{(2a-b)b \cot(c+dx)}{d} - \frac{b^2 \cot^3(c+dx)}{3d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.26 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.51

$$\int (a + b \cot^2(c + dx))^2 dx = \frac{\cot(c + dx) \left(b(6a - 3b + b \cot^2(c + dx)) + 3(a - b)^2 \operatorname{arctanh}\left(\sqrt{-\tan^2(c + dx)}\right) \sqrt{-\tan^2(c + dx)} \right)}{3d}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^2,x]

[Out] -1/3*(Cot[c + d*x]*(b*(6*a - 3*b + b*Cot[c + d*x]^2) + 3*(a - b)^2*ArcTanh[Sqrt[-Tan[c + d*x]^2]]*Sqrt[-Tan[c + d*x]^2]))/d

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.02

method	result	size
parallelrisc	$\frac{-b^2 \cot(dx+c)^3 + (-6ab+3b^2) \cot(dx+c) + 3dx(a-b)^2}{3d}$	48
norman	$\frac{(a^2-2ab+b^2)x \tan(dx+c)^3 - \frac{b^2}{3d} - \frac{b(2a-b) \tan(dx+c)^2}{d}}{\tan(dx+c)^3}$	61
derivativedivides	$\frac{-\frac{b^2 \cot(dx+c)^3}{3} - 2 \cot(dx+c)ab + \cot(dx+c)b^2 + (-a^2+2ab-b^2) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right)}{d}$	68
default	$\frac{-\frac{b^2 \cot(dx+c)^3}{3} - 2 \cot(dx+c)ab + \cot(dx+c)b^2 + (-a^2+2ab-b^2) \left(\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))\right)}{d}$	68
parts	$a^2x + \frac{b^2 \left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) - \frac{\pi}{2} + \operatorname{arccot}(\cot(dx+c)) \right)}{d} + \frac{2ab \left(-\cot(dx+c) + \frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right)}{d}$	69
risc	$a^2x - 2abx + b^2x + \frac{4ib(-3ae^{4i(dx+c)} + 3be^{4i(dx+c)} + 6e^{2i(dx+c)}a - 3e^{2i(dx+c)}b - 3a + 2b)}{3d(e^{2i(dx+c)} - 1)^3}$	92

[In] `int((a+b*cot(d*x+c))^2,x,method=_RETURNVERBOSE)`

[Out] $1/3*(-b^2*\cot(d*x+c)^3+(-6*a*b+3*b^2)*\cot(d*x+c)+3*d*x*(a-b)^2)/d$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. $2(45) = 90$.

Time = 0.27 (sec) , antiderivative size = 127, normalized size of antiderivative = 2.70

$$\int (a + b \cot^2(c + dx))^2 dx$$

$$= \frac{2b^2 \cos(2dx + 2c) - 2(3ab - 2b^2) \cos(2dx + 2c)^2 + 6ab - 2b^2 + 3((a^2 - 2ab + b^2)dx \cos(2dx + 2c))}{3(d \cos(2dx + 2c) - d) \sin(2dx + 2c)}$$

[In] `integrate((a+b*cot(d*x+c))^2,x, algorithm="fricas")`

[Out] $1/3*(2*b^2*\cos(2*d*x + 2*c) - 2*(3*a*b - 2*b^2)*\cos(2*d*x + 2*c)^2 + 6*a*b - 2*b^2 + 3*((a^2 - 2*a*b + b^2)*d*x*\cos(2*d*x + 2*c) - (a^2 - 2*a*b + b^2)*d*x*\sin(2*d*x + 2*c))/((d*\cos(2*d*x + 2*c) - d)*\sin(2*d*x + 2*c))$

Sympy [A] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.45

$$\int (a + b \cot^2(c + dx))^2 dx$$

$$= \begin{cases} a^2x - 2abx - \frac{2ab \cot(c+dx)}{d} + b^2x - \frac{b^2 \cot^3(c+dx)}{3d} + \frac{b^2 \cot(c+dx)}{d} & \text{for } d \neq 0 \\ x(a + b \cot^2(c))^2 & \text{otherwise} \end{cases}$$

[In] integrate((a+b*cot(d*x+c)**2)**2,x)

[Out] Piecewise((a**2*x - 2*a*b*x - 2*a*b*cot(c + d*x)/d + b**2*x - b**2*cot(c + d*x)**3/(3*d) + b**2*cot(c + d*x)/d, Ne(d, 0)), (x*(a + b*cot(c)**2)**2, True))

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.34

$$\int (a + b \cot^2(c + dx))^2 dx = a^2x - \frac{2 \left(dx + c + \frac{1}{\tan(dx+c)} \right) ab}{d} + \frac{\left(3 dx + 3c + \frac{3 \tan(dx+c)^2 - 1}{\tan(dx+c)^3} \right) b^2}{3d}$$

[In] integrate((a+b*cot(d*x+c)^2)^2,x, algorithm="maxima")

[Out] a^2*x - 2*(d*x + c + 1/tan(d*x + c))*a*b/d + 1/3*(3*d*x + 3*c + (3*tan(d*x + c)^2 - 1)/tan(d*x + c)^3)*b^2/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. 2(45) = 90.

Time = 0.32 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.43

$$\int (a + b \cot^2(c + dx))^2 dx$$

$$= \frac{b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^3 + 24 ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) - 15 b^2 \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right) + 24 (a^2 - 2 ab + b^2)(dx + c) - \frac{24 ab \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)}{24 d}}{24 d}$$

[In] integrate((a+b*cot(d*x+c)^2)^2,x, algorithm="giac")

[Out] 1/24*(b^2*tan(1/2*d*x + 1/2*c)^3 + 24*a*b*tan(1/2*d*x + 1/2*c) - 15*b^2*tan(1/2*d*x + 1/2*c) + 24*(a^2 - 2*a*b + b^2)*(d*x + c) - (24*a*b*tan(1/2*d*x + 1/2*c)^2 - 15*b^2*tan(1/2*d*x + 1/2*c)^2 + b^2)/tan(1/2*d*x + 1/2*c)^3)/d

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int (a + b \cot^2(c + dx))^2 dx = x(a - b)^2 - \frac{b^2 \cot(c + dx)^3}{3d} - \frac{b \cot(c + dx) (2a - b)}{d}$$

[In] int((a + b*cot(c + d*x)^2)^2,x)

[Out] x*(a - b)^2 - (b^2*cot(c + d*x)^3)/(3*d) - (b*cot(c + d*x)*(2*a - b))/d

3.4 $\int (a + b \cot^2(c + dx))^3 dx$

Optimal result	64
Rubi [A] (verified)	64
Mathematica [A] (verified)	65
Maple [A] (verified)	66
Fricas [B] (verification not implemented)	66
Sympy [A] (verification not implemented)	67
Maxima [A] (verification not implemented)	67
Giac [B] (verification not implemented)	68
Mupad [B] (verification not implemented)	68

Optimal result

Integrand size = 14, antiderivative size = 78

$$\int (a + b \cot^2(c + dx))^3 dx = (a - b)^3 x - \frac{b(3a^2 - 3ab + b^2) \cot(c + dx)}{d} - \frac{(3a - b)b^2 \cot^3(c + dx)}{3d} - \frac{b^3 \cot^5(c + dx)}{5d}$$

[Out] (a-b)^3*x-b*(3*a^2-3*a*b+b^2)*cot(d*x+c)/d-1/3*(3*a-b)*b^2*cot(d*x+c)^3/d-1/5*b^3*cot(d*x+c)^5/d

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3742, 398, 209}

$$\int (a + b \cot^2(c + dx))^3 dx = -\frac{b(3a^2 - 3ab + b^2) \cot(c + dx)}{d} - \frac{b^2(3a - b) \cot^3(c + dx)}{3d} + x(a - b)^3 - \frac{b^3 \cot^5(c + dx)}{5d}$$

[In] Int[(a + b*Cot[c + d*x]^2)^3,x]

[Out] (a - b)^3*x - (b*(3*a^2 - 3*a*b + b^2)*Cot[c + d*x])/d - ((3*a - b)*b^2*Cot[c + d*x]^3)/(3*d) - (b^3*Cot[c + d*x]^5)/(5*d)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{(a+bx^2)^3}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\ &= -\frac{\text{Subst}\left(\int \left(b(3a^2 - 3ab + b^2) + (3a - b)b^2x^2 + b^3x^4 + \frac{(a-b)^3}{1+x^2}\right) dx, x, \cot(c+dx)\right)}{d} \\ &= -\frac{b(3a^2 - 3ab + b^2) \cot(c+dx)}{d} - \frac{(3a - b)b^2 \cot^3(c+dx)}{3d} \\ &\quad - \frac{b^3 \cot^5(c+dx)}{5d} - \frac{(a-b)^3 \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\ &= (a-b)^3 x - \frac{b(3a^2 - 3ab + b^2) \cot(c+dx)}{d} - \frac{(3a - b)b^2 \cot^3(c+dx)}{3d} - \frac{b^3 \cot^5(c+dx)}{5d} \end{aligned}$$

Mathematica [A] (verified)

Time = 2.95 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.42

$$\int (a + b \cot^2(c + dx))^3 dx = \frac{\cot^5(c + dx) \left(\frac{15(a-b)^3 \arctanh\left(\frac{\sqrt{-\tan^2(c+dx)}}{-\tan^2(c+dx)}\right) \tan^8(c+dx)}{(-\tan^2(c+dx))^{3/2}} + b(3b^2 + 5(3a-b)b \tan^2(c+dx) + 15(3a^2 - 3ab + b^2)) \right)}{15d}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^3,x]

[Out] -1/15*(Cot[c + d*x]^5*((15*(a - b)^3*ArcTanh[Sqrt[-Tan[c + d*x]^2]]*Tan[c + d*x]^8)/(-Tan[c + d*x]^2)^(3/2) + b*(3*b^2 + 5*(3*a - b)*b*Tan[c + d*x]^2 + 15*(3*a^2 - 3*a*b + b^2)*Tan[c + d*x]^4))/d

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.99

method	result
parallelrisc	$\frac{-3b^3 \cot(dx+c)^5 + 5(-3ab^2+b^3) \cot(dx+c)^3 + 15(-3a^2b+3ab^2-b^3) \cot(dx+c) + 15dx(a-b)^3}{15d}$
norman	$\frac{(a^3-3a^2b+3ab^2-b^3)x \tan(dx+c)^5 - \frac{b^3}{5d} - \frac{b(3a^2-3ab+b^2) \tan(dx+c)^4}{d} - \frac{b^2(3a-b) \tan(dx+c)^2}{3d}}{\tan(dx+c)^5}$
derivativedivides	$\frac{-\frac{b^3 \cot(dx+c)^5}{5} - ab^2 \cot(dx+c)^3 + \frac{b^3 \cot(dx+c)^3}{3} - 3a^2b \cot(dx+c) + 3 \cot(dx+c)ab^2 - \cot(dx+c)b^3 + (-a^3+3a^2b-3ab^2+b^3)}{d}$
default	$\frac{-\frac{b^3 \cot(dx+c)^5}{5} - ab^2 \cot(dx+c)^3 + \frac{b^3 \cot(dx+c)^3}{3} - 3a^2b \cot(dx+c) + 3 \cot(dx+c)ab^2 - \cot(dx+c)b^3 + (-a^3+3a^2b-3ab^2+b^3)}{d}$
parts	$a^3x + \frac{b^3 \left(-\frac{\cot(dx+c)^5}{5} + \frac{\cot(dx+c)^3}{3} - \cot(dx+c) + \frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c)) \right)}{d} + \frac{3ab^2 \left(-\frac{\cot(dx+c)^3}{3} + \cot(dx+c) - \frac{\pi}{2} + \operatorname{arccot}(\cot(dx+c)) \right)}{d}$
risc	$a^3x - 3a^2bx + 3ab^2x - b^3x - \frac{2ib(45a^2e^{8i(dx+c)} - 90ab e^{8i(dx+c)} + 45b^2e^{8i(dx+c)} - 180a^2e^{6i(dx+c)} + 270ab e^{6i(dx+c)} - 180a^2e^{4i(dx+c)} + 270ab e^{4i(dx+c)} - 180a^2e^{2i(dx+c)} + 270ab e^{2i(dx+c)} - 180a^2 + 270ab - 180b^2)}{d}$

```
[In] int((a+b*cot(d*x+c)^2)^3,x,method=_RETURNVERBOSE)
```

```
[Out] 1/15*(-3*b^3*cot(d*x+c)^5+5*(-3*a*b^2+b^3)*cot(d*x+c)^3+15*(-3*a^2*b+3*a*b^2-b^3)*cot(d*x+c)+15*d*x*(a-b)^3)/d
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(74) = 148.

Time = 0.26 (sec) , antiderivative size = 253, normalized size of antiderivative = 3.24

$$\int (a + b \cot^2(c + dx))^3 dx = \frac{(45a^2b - 60ab^2 + 23b^3) \cos(2dx + 2c)^3 + 45a^2b - 30ab^2 + 13b^3 - (45a^2b - 30ab^2 + b^3) \cos(2dx + 2c)^2 + (45a^2b - 60ab^2 + 11b^3) \cos(2dx + 2c) - 15((a^3 - 3a^2b + 3ab^2 - b^3) d^2 x \cos(2dx + 2c)^2 - 2(a^3 - 3a^2b + 3ab^2 - b^3) d^2 x \cos(2dx + 2c) + (a^3 - 3a^2b + 3ab^2 - b^3) d^2 x \sin(2dx + 2c))}{(d \cos(2dx + 2c))^2 - 2d \cos(2dx + 2c) + d} \sin(2dx + 2c)$$

```
[In] integrate((a+b*cot(d*x+c)^2)^3,x, algorithm="fricas")
```

```
[Out] -1/15*((45*a^2*b - 60*a*b^2 + 23*b^3)*cos(2*d*x + 2*c)^3 + 45*a^2*b - 30*a*b^2 + 13*b^3 - (45*a^2*b - 30*a*b^2 + b^3)*cos(2*d*x + 2*c)^2 - (45*a^2*b - 60*a*b^2 + 11*b^3)*cos(2*d*x + 2*c) - 15*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*cos(2*d*x + 2*c)^2 - 2*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*cos(2*d*x + 2*c) + (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*x*sin(2*d*x + 2*c))/((d*cos(2*d*x + 2*c))^2 - 2*d*cos(2*d*x + 2*c) + d)*sin(2*d*x + 2*c)
```

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.62

$$\int (a + b \cot^2(c + dx))^3 dx$$

$$= \begin{cases} a^3 x - 3a^2 b x - \frac{3a^2 b \cot(c+dx)}{d} + 3ab^2 x - \frac{ab^2 \cot^3(c+dx)}{d} + \frac{3ab^2 \cot(c+dx)}{d} - b^3 x - \frac{b^3 \cot^5(c+dx)}{5d} + \frac{b^3 \cot^3(c+dx)}{3d} - b^3 \cot(c+dx) \\ x(a + b \cot^2(c))^3 \end{cases}$$

[In] integrate((a+b*cot(d*x+c)**2)**3,x)

[Out] Piecewise((a**3*x - 3*a**2*b*x - 3*a**2*b*cot(c + d*x)/d + 3*a*b**2*x - a*b**2*cot(c + d*x)**3/d + 3*a*b**2*cot(c + d*x)/d - b**3*x - b**3*cot(c + d*x)**5/(5*d) + b**3*cot(c + d*x)**3/(3*d) - b**3*cot(c + d*x)/d, Ne(d, 0)), (x*(a + b*cot(c)**2)**3, True))

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.44

$$\int (a + b \cot^2(c + dx))^3 dx = a^3 x - \frac{3 \left(dx + c + \frac{1}{\tan(dx+c)} \right) a^2 b}{d}$$

$$+ \frac{\left(3 dx + 3 c + \frac{3 \tan(dx+c)^2 - 1}{\tan(dx+c)^3} \right) a b^2}{d}$$

$$- \frac{\left(15 dx + 15 c + \frac{15 \tan(dx+c)^4 - 5 \tan(dx+c)^2 + 3}{\tan(dx+c)^5} \right) b^3}{15 d}$$

[In] integrate((a+b*cot(d*x+c)^2)^3,x, algorithm="maxima")

[Out] a^3*x - 3*(d*x + c + 1/tan(d*x + c))*a^2*b/d + (3*d*x + 3*c + (3*tan(d*x + c)^2 - 1)/tan(d*x + c)^3)*a*b^2/d - 1/15*(15*d*x + 15*c + (15*tan(d*x + c)^4 - 5*tan(d*x + c)^2 + 3)/tan(d*x + c)^5)*b^3/d

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(74) = 148.

Time = 0.38 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.94

$$\int (a + b \cot^2(c + dx))^3 dx$$

$$= \frac{3b^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^5 + 60ab^2 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right)^3 - 35b^3 \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) + 720a^2b \tan\left(\frac{1}{2}dx + \frac{1}{2}c\right) - 900a^2}{d}$$

[In] integrate((a+b*cot(d*x+c)^2)^3,x, algorithm="giac")

[Out] 1/480*(3*b^3*tan(1/2*d*x + 1/2*c)^5 + 60*a*b^2*tan(1/2*d*x + 1/2*c)^3 - 35*b^3*tan(1/2*d*x + 1/2*c) + 720*a^2*b*tan(1/2*d*x + 1/2*c) - 900*a*b^2*tan(1/2*d*x + 1/2*c) + 330*b^3*tan(1/2*d*x + 1/2*c) + 480*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*(d*x + c) - (720*a^2*b*tan(1/2*d*x + 1/2*c)^4 - 900*a*b^2*tan(1/2*d*x + 1/2*c)^4 + 330*b^3*tan(1/2*d*x + 1/2*c)^4 + 60*a*b^2*tan(1/2*d*x + 1/2*c)^2 - 35*b^3*tan(1/2*d*x + 1/2*c)^2 + 3*b^3)/tan(1/2*d*x + 1/2*c)^5)/d

Mupad [B] (verification not implemented)

Time = 12.92 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.97

$$\int (a + b \cot^2(c + dx))^3 dx = x(a - b)^3 - \frac{b^3 \cot(c + dx)^5}{5d} - \frac{\cot(c + dx)^3 (3ab^2 - b^3)}{3d} - \frac{b \cot(c + dx) (3a^2 - 3ab + b^2)}{d}$$

[In] int((a + b*cot(c + d*x)^2)^3,x)

[Out] x*(a - b)^3 - (b^3*cot(c + d*x)^5)/(5*d) - (cot(c + d*x)^3*(3*a*b^2 - b^3))/(3*d) - (b*cot(c + d*x)*(3*a^2 - 3*a*b + b^2))/d

3.5 $\int \frac{1}{a+b \cot^2(c+dx)} dx$

Optimal result	69
Rubi [A] (verified)	69
Mathematica [A] (verified)	70
Maple [A] (verified)	70
Fricas [A] (verification not implemented)	71
Sympy [B] (verification not implemented)	72
Maxima [A] (verification not implemented)	72
Giac [A] (verification not implemented)	73
Mupad [B] (verification not implemented)	73

Optimal result

Integrand size = 14, antiderivative size = 49

$$\int \frac{1}{a+b \cot^2(c+dx)} dx = \frac{x}{a-b} + \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{\sqrt{a}(a-b)d}$$

[Out] $x/(a-b)+\arctan(\cot(d*x+c)*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/(a-b)/d/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {3741, 3756, 211}

$$\int \frac{1}{a+b \cot^2(c+dx)} dx = \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{\sqrt{ad}(a-b)} + \frac{x}{a-b}$$

[In] $\text{Int}[(a + b*\text{Cot}[c + d*x]^2)^{-1}, x]$

[Out] $x/(a - b) + (\text{Sqrt}[b]*\text{ArcTan}[(\text{Sqrt}[b]*\text{Cot}[c + d*x])/ \text{Sqrt}[a]])/(\text{Sqrt}[a]*(a - b)*d)$

Rule 211

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

Rule 3741

$\text{Int}[(a_ + (b_)*\tan[(e_ + (f_)*(x_)^2])^{-1}, x_Symbol] \rightarrow \text{Simp}[x/(a - b), x] - \text{Dist}[b/(a - b), \text{Int}[\text{Sec}[e + f*x]^2/(a + b*\text{Tan}[e + f*x]^2), x], x]$

```
;/ FreeQ[{a, b, e, f}, x] && NeQ[a, b]
```

Rule 3756

```
Int[sec[(e_.) + (f_.)*(x_.)]^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_.)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/(c^(m - 1)*f), Subst[Int[(c^2 + ff^2*x^2)^(m/2 - 1)*(a + b*(ff*x)^n)^p, x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[m/2] && (IntegersQ[n, p] || IGtQ[m, 0] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{a-b} - \frac{b \int \frac{\csc^2(c+dx)}{a+b \cot^2(c+dx)} dx}{a-b} \\ &= \frac{x}{a-b} + \frac{b \text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cot(c+dx)\right)}{(a-b)d} \\ &= \frac{x}{a-b} + \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{\sqrt{a}(a-b)d} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00

$$\int \frac{1}{a+b \cot^2(c+dx)} dx = \frac{\arctan(\tan(c+dx)) - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a} \tan(c+dx)}{\sqrt{b}}\right)}{\sqrt{a}}}{ad-bd}$$

```
[In] Integrate[(a + b*Cot[c + d*x]^2)^(-1), x]
```

```
[Out] (ArcTan[Tan[c + d*x]] - (Sqrt[b]*ArcTan[(Sqrt[a]*Tan[c + d*x])/Sqrt[b]])/Sqrt[a])/(a*d - b*d)
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$\frac{-\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{a-b} + \frac{b \arctan\left(\frac{b \cot(dx+c)}{\sqrt{ab}}\right)}{(a-b)\sqrt{ab}}$	56
default	$\frac{-\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{a-b} + \frac{b \arctan\left(\frac{b \cot(dx+c)}{\sqrt{ab}}\right)}{(a-b)\sqrt{ab}}$	56
risch	$\frac{x}{a-b} + \frac{\sqrt{-ab} \ln\left(\frac{e^{2i(dx+c)} - 2i\sqrt{-ab} + a+b}{a-b}\right)}{2a(a-b)d} - \frac{\sqrt{-ab} \ln\left(\frac{e^{2i(dx+c)} + 2i\sqrt{-ab} - a-b}{a-b}\right)}{2a(a-b)d}$	120

[In] int(1/(a+b*cot(d*x+c)^2),x,method=_RETURNVERBOSE)

[Out] 1/d*(-1/(a-b)*(1/2*Pi-arccot(cot(d*x+c)))+b/(a-b)/(a*b)^(1/2)*arctan(b*cot(d*x+c)/(a*b)^(1/2)))

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 252, normalized size of antiderivative = 5.14

$$\int \frac{1}{a + b \cot^2(c + dx)} dx$$

$$= \frac{4 dx - \sqrt{-\frac{b}{a}} \log\left(\frac{(a^2+6ab+b^2) \cos(2dx+2c)^2 + 4(a^2-ab-(a^2+ab) \cos(2dx+2c)) \sqrt{-\frac{b}{a}} \sin(2dx+2c) + a^2-6ab+b^2-2(a^2-b^2) \cos(2dx+2c)}{(a^2-2ab+b^2) \cos(2dx+2c)^2 + a^2+2ab+b^2-2(a^2-b^2) \cos(2dx+2c)}\right)}{4(a-b)d}$$

[In] integrate(1/(a+b*cot(d*x+c)^2),x, algorithm="fricas")

[Out] [1/4*(4*d*x - sqrt(-b/a)*log(((a^2 + 6*a*b + b^2)*cos(2*d*x + 2*c))^2 + 4*(a^2 - a*b - (a^2 + a*b)*cos(2*d*x + 2*c))*sqrt(-b/a)*sin(2*d*x + 2*c) + a^2 - 6*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c)))/((a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c)))/((a - b)*d), 1/2*(2*d*x + sqrt(b/a)*arctan(1/2*((a + b)*cos(2*d*x + 2*c) - a + b)*sqrt(b/a)/(b*sin(2*d*x + 2*c))))/((a - b)*d)]

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 238 vs. 2(37) = 74.

Time = 0.74 (sec) , antiderivative size = 238, normalized size of antiderivative = 4.86

$$\int \frac{1}{a + b \cot^2(c + dx)} dx$$

$$= \begin{cases} \frac{\infty x}{\cot^2(c)} & \text{for } a = 0 \wedge b = 0 \wedge d = 0 \\ \frac{x}{a} & \text{for } b = 0 \\ \frac{-x + \frac{1}{d \cot(c+dx)}}{b} & \text{for } a = 0 \\ \frac{dx \cot^2(c+dx)}{2bd \cot^2(c+dx)+2bd} + \frac{dx}{2bd \cot^2(c+dx)+2bd} - \frac{\cot(c+dx)}{2bd \cot^2(c+dx)+2bd} & \text{for } a = b \\ \frac{x}{a+b \cot^2(c)} & \text{for } d = 0 \\ \frac{2dx\sqrt{-\frac{a}{b}}}{2ad\sqrt{-\frac{a}{b}}-2bd\sqrt{-\frac{a}{b}}} + \frac{\log\left(-\sqrt{-\frac{a}{b}}+\cot(c+dx)\right)}{2ad\sqrt{-\frac{a}{b}}-2bd\sqrt{-\frac{a}{b}}} - \frac{\log\left(\sqrt{-\frac{a}{b}}+\cot(c+dx)\right)}{2ad\sqrt{-\frac{a}{b}}-2bd\sqrt{-\frac{a}{b}}} & \text{otherwise} \end{cases}$$

[In] integrate(1/(a+b*cot(d*x+c)**2),x)

[Out] Piecewise((zoo*x/cot(c)**2, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a, Eq(b, 0)), ((-x + 1/(d*cot(c + d*x)))/b, Eq(a, 0)), (d*x*cot(c + d*x)**2/(2*b*d*cot(c + d*x)**2 + 2*b*d) + d*x/(2*b*d*cot(c + d*x)**2 + 2*b*d) - cot(c + d*x)/(2*b*d*cot(c + d*x)**2 + 2*b*d), Eq(a, b)), (x/(a + b*cot(c)**2), Eq(d, 0)), (2*d*x*sqrt(-a/b)/(2*a*d*sqrt(-a/b) - 2*b*d*sqrt(-a/b)) + log(-sqrt(-a/b) + cot(c + d*x))/(2*a*d*sqrt(-a/b) - 2*b*d*sqrt(-a/b)) - log(sqrt(-a/b) + cot(c + d*x))/(2*a*d*sqrt(-a/b) - 2*b*d*sqrt(-a/b)), True))

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.98

$$\int \frac{1}{a + b \cot^2(c + dx)} dx = -\frac{b \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right)}{\sqrt{ab}(a-b)} - \frac{dx+c}{a-b}$$

[In] integrate(1/(a+b*cot(d*x+c)^2),x, algorithm="maxima")

[Out] -(b*arctan(a*tan(d*x + c)/sqrt(a*b))/(sqrt(a*b)*(a - b)) - (d*x + c)/(a - b))/d

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.33

$$\int \frac{1}{a + b \cot^2(c + dx)} dx = -\frac{\left(\pi \left\lfloor \frac{dx+c}{\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right)\right) b}{\sqrt{ab}(a-b)} - \frac{dx+c}{a-b}$$

[In] integrate(1/(a+b*cot(d*x+c)^2),x, algorithm="giac")

[Out] -((pi*floor((d*x + c)/pi + 1/2)*sgn(a) + arctan(a*tan(d*x + c)/sqrt(a*b)))*
b/(sqrt(a*b)*(a - b)) - (d*x + c)/(a - b))/d**Mupad [B] (verification not implemented)**

Time = 0.12 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.84

$$\int \frac{1}{a + b \cot^2(c + dx)} dx = \frac{x}{a - b} + \frac{b \operatorname{atan}\left(\frac{b \cot(c+dx)}{\sqrt{ab}}\right)}{d \sqrt{ab} (a - b)}$$

[In] int(1/(a + b*cot(c + d*x)^2),x)

[Out] x/(a - b) + (b*atan((b*cot(c + d*x))/(a*b)^(1/2)))/(d*(a*b)^(1/2)*(a - b))

3.6 $\int \frac{1}{(a+b \cot^2(c+dx))^2} dx$

Optimal result	74
Rubi [A] (verified)	74
Mathematica [A] (verified)	76
Maple [A] (verified)	76
Fricas [B] (verification not implemented)	77
Sympy [B] (verification not implemented)	77
Maxima [A] (verification not implemented)	79
Giac [A] (verification not implemented)	79
Mupad [B] (verification not implemented)	79

Optimal result

Integrand size = 14, antiderivative size = 97

$$\int \frac{1}{(a+b \cot^2(c+dx))^2} dx = \frac{x}{(a-b)^2} + \frac{(3a-b)\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{2a^{3/2}(a-b)^2 d} + \frac{b \cot(c+dx)}{2a(a-b)d(a+b \cot^2(c+dx))}$$

[Out] $x/(a-b)^2 + 1/2*b*\cot(d*x+c)/a/(a-b)/d/(a+b*\cot(d*x+c)^2) + 1/2*(3*a-b)*\arctan(\cot(d*x+c)*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/a^{(3/2)}/(a-b)^2/d$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3742, 425, 536, 209, 211}

$$\int \frac{1}{(a+b \cot^2(c+dx))^2} dx = \frac{\sqrt{b}(3a-b) \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{2a^{3/2}d(a-b)^2} + \frac{b \cot(c+dx)}{2ad(a-b)(a+b \cot^2(c+dx))} + \frac{x}{(a-b)^2}$$

[In] Int[(a + b*Cot[c + d*x]^2)^(-2), x]

[Out] $x/(a-b)^2 + ((3*a-b)*\text{Sqrt}[b]*\text{ArcTan}[(\text{Sqrt}[b]*\text{Cot}[c+d*x])/\text{Sqrt}[a]])/(2*a^{(3/2)}*(a-b)^2*d) + (b*\text{Cot}[c+d*x])/(2*a*(a-b)*d*(a+b*\text{Cot}[c+d*x]^2))$

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 211

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 425

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (!(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x])
```

Rule 536

```
Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 3742

```
Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x^2)(a+bx^2)^2} dx, x, \cot(c+dx)\right)}{d} \\ &= \frac{b \cot(c+dx)}{2a(a-b)d(a+b \cot^2(c+dx))} - \frac{\text{Subst}\left(\int \frac{2a-b-bx^2}{(1+x^2)(a+bx^2)} dx, x, \cot(c+dx)\right)}{2a(a-b)d} \\ &= \frac{b \cot(c+dx)}{2a(a-b)d(a+b \cot^2(c+dx))} - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(c+dx)\right)}{(a-b)^2d} \\ &\quad + \frac{((3a-b)b)\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cot(c+dx)\right)}{2a(a-b)^2d} \end{aligned}$$

$$= \frac{x}{(a-b)^2} + \frac{(3a-b)\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{2a^{3/2}(a-b)^2 d} + \frac{b \cot(c+dx)}{2a(a-b)d(a+b \cot^2(c+dx))}$$

Mathematica [A] (verified)

Time = 1.02 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.93

$$\int \frac{1}{(a+b \cot^2(c+dx))^2} dx$$

$$= \frac{-2 \arctan(\cot(c+dx)) + \frac{(3a-b)\sqrt{b} \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{a^{3/2}} + \frac{(a-b)b \cot(c+dx)}{a(a+b \cot^2(c+dx))}}{2(a-b)^2 d}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(-2), x]

[Out] (-2*ArcTan[Cot[c + d*x]] + ((3*a - b)*Sqrt[b]*ArcTan[(Sqrt[b]*Cot[c + d*x])/Sqrt[a]])/a^(3/2) + ((a - b)*b*Cot[c + d*x])/(a*(a + b*Cot[c + d*x]^2)))/(2*(a - b)^2*d)

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.02

method	result
derivativedivides	$\frac{-\frac{\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{(a-b)^2} + \frac{b \left(\frac{(a-b) \cot(dx+c)}{2a(a+b \cot(dx+c)^2)} + \frac{(3a-b) \arctan\left(\frac{b \cot(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a-b)^2}}{d}$
default	$\frac{-\frac{\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{(a-b)^2} + \frac{b \left(\frac{(a-b) \cot(dx+c)}{2a(a+b \cot(dx+c)^2)} + \frac{(3a-b) \arctan\left(\frac{b \cot(dx+c)}{\sqrt{ab}}\right)}{2a\sqrt{ab}} \right)}{(a-b)^2}}{d}$
risch	$\frac{x}{a^2-2ab+b^2} - \frac{ib(e^{2i(dx+c)}a + e^{2i(dx+c)}b - a + b)}{da(-a+b)^2(-ae^{4i(dx+c)} + be^{4i(dx+c)} + 2e^{2i(dx+c)}a + 2e^{2i(dx+c)}b - a + b)} + \frac{3\sqrt{-ab} \ln\left(e^{2i(dx+c)} - \frac{2i\sqrt{-ab}}{a}\right)}{4a(a-b)^2 d}$

[In] int(1/(a+b*cot(d*x+c)^2)^2,x,method=_RETURNVERBOSE)

[Out] 1/d*(-1/(a-b)^2*(1/2*Pi-arccot(cot(d*x+c)))+1/(a-b)^2*b*(1/2*(a-b)/a*cot(d*x+c)/(a+b*cot(d*x+c)^2)+1/2*(3*a-b)/a/(a*b)^(1/2)*arctan(b*cot(d*x+c)/(a*b)^(1/2))))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(85) = 170.

Time = 0.31 (sec) , antiderivative size = 534, normalized size of antiderivative = 5.51

$$\int \frac{1}{(a + b \cot^2(c + dx))^2} dx$$

$$= \frac{8(a^2 - ab)dx \cos(2dx + 2c) - 8(a^2 + ab)dx + (3a^2 + 2ab - b^2 - (3a^2 - 4ab + b^2) \cos(2dx + 2c)) \sqrt{8((a^4 - 3a^3b + 3a^2b^2 -$$

```
[In] integrate(1/(a+b*cot(d*x+c)^2)^2,x, algorithm="fricas")
```

```
[Out] [1/8*(8*(a^2 - a*b)*d*x*cos(2*d*x + 2*c) - 8*(a^2 + a*b)*d*x + (3*a^2 + 2*a*b - b^2 - (3*a^2 - 4*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(-b/a)*log(((a^2 + 6*a*b + b^2)*cos(2*d*x + 2*c)^2 + 4*(a^2 - a*b - (a^2 + a*b)*cos(2*d*x + 2*c))*sqrt(-b/a)*sin(2*d*x + 2*c) + a^2 - 6*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c))/((a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c))) - 4*(a*b - b^2)*sin(2*d*x + 2*c))/((a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3)*d*cos(2*d*x + 2*c) - (a^4 - a^3*b - a^2*b^2 + a*b^3)*d), 1/4*(4*(a^2 - a*b)*d*x*cos(2*d*x + 2*c) - 4*(a^2 + a*b)*d*x - (3*a^2 + 2*a*b - b^2 - (3*a^2 - 4*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(b/a)*arctan(1/2*((a + b)*cos(2*d*x + 2*c) - a + b)*sqrt(b/a)/(b*sin(2*d*x + 2*c))) - 2*(a*b - b^2)*sin(2*d*x + 2*c))/((a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3)*d*cos(2*d*x + 2*c) - (a^4 - a^3*b - a^2*b^2 + a*b^3)*d)]
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2125 vs. 2(78) = 156.

Time = 9.07 (sec) , antiderivative size = 2125, normalized size of antiderivative = 21.91

$$\int \frac{1}{(a + b \cot^2(c + dx))^2} dx = \text{Too large to display}$$

```
[In] integrate(1/(a+b*cot(d*x+c)**2)**2,x)
```

```
[Out] Piecewise((zoo*x/cot(c)**4, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**2, Eq(b, 0)), ((x - 1/(d*cot(c + d*x)) + 1/(3*d*cot(c + d*x)**3))/b**2, Eq(a, 0)), (3*d*x*cot(c + d*x)**4/(8*b**2*d*cot(c + d*x)**4 + 16*b**2*d*cot(c + d*x)**2 + 8*b**2*d) + 6*d*x*cot(c + d*x)**2/(8*b**2*d*cot(c + d*x)**4 + 16*b**2*d*cot(c + d*x)**2 + 8*b**2*d) + 3*d*x/(8*b**2*d*cot(c + d*x)**4 + 16*b**2*d*cot(c + d*x)**2 + 8*b**2*d) - 3*cot(c + d*x)**3/(8*b**2*d*cot(c + d*x)**4 + 16*b**2*d*cot(c + d*x)**2 + 8*b**2*d) - 5*cot(c + d*x)/(8*b**2*d*cot(c + d
```

```

*x)**4 + 16*b**2*d*cot(c + d*x)**2 + 8*b**2*d), Eq(a, b)), (x/(a + b*cot(c)
**2)**2, Eq(d, 0)), (4*a**2*d*x*sqrt(-a/b)/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*
d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-
a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot
(c + d*x)**2) + 3*a**2*log(-sqrt(-a/b) + cot(c + d*x))/(4*a**4*d*sqrt(-a/b)
+ 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b
**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sq
rt(-a/b)*cot(c + d*x)**2) - 3*a**2*log(sqrt(-a/b) + cot(c + d*x))/(4*a**4*d
*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b)
- 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*
a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) + 4*a*b*d*x*sqrt(-a/b)*cot(c + d*x)**2
/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*
sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(
-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) + 2*a*b*sqrt(-a/b)*cot(c + d
*x)/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b
*d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sq
rt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) + 3*a*b*log(-sqrt(-a/b) +
cot(c + d*x))*cot(c + d*x)**2/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)
*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c +
d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2
) - a*b*log(-sqrt(-a/b) + cot(c + d*x))/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*s
qrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b
)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c
+ d*x)**2) - 3*a*b*log(sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**2/(4*a**4*d
*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b)
- 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*
a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) + a*b*log(sqrt(-a/b) + cot(c + d*x))/(
4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sq
rt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a
/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) - 2*b**2*sqrt(-a/b)*cot(c + d*
x)/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*
d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sq
rt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2) - b**2*log(-sqrt(-a/b) + c
ot(c + d*x))*cot(c + d*x)**2/(4*a**4*d*sqrt(-a/b) + 4*a**3*b*d*sqrt(-a/b)*c
ot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b**2*d*sqrt(-a/b)*cot(c + d
*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sqrt(-a/b)*cot(c + d*x)**2)
+ b**2*log(sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**2/(4*a**4*d*sqrt(-a/b)
+ 4*a**3*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 8*a**3*b*d*sqrt(-a/b) - 8*a**2*b*
**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 4*a**2*b**2*d*sqrt(-a/b) + 4*a*b**3*d*sq
rt(-a/b)*cot(c + d*x)**2), True))

```

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.19

$$\int \frac{1}{(a + b \cot^2(c + dx))^2} dx = \frac{\frac{b \tan(dx+c)}{a^2 b - ab^2 + (a^3 - a^2 b) \tan(dx+c)^2} - \frac{(3ab - b^2) \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right)}{(a^3 - 2a^2 b + ab^2) \sqrt{ab}} + \frac{2(dx+c)}{a^2 - 2ab + b^2}}{2d}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^2,x, algorithm="maxima")

```
[Out] 1/2*(b*tan(d*x + c)/(a^2*b - a*b^2 + (a^3 - a^2*b)*tan(d*x + c)^2) - (3*a*b - b^2)*arctan(a*tan(d*x + c)/sqrt(a*b)))/((a^3 - 2*a^2*b + a*b^2)*sqrt(a*b)) + 2*(d*x + c)/(a^2 - 2*a*b + b^2))/d
```

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 123, normalized size of antiderivative = 1.27

$$\int \frac{1}{(a + b \cot^2(c + dx))^2} dx = \frac{\left(\pi \left\lfloor \frac{dx+c}{\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right)\right) (3ab - b^2)}{(a^3 - 2a^2 b + ab^2) \sqrt{ab}} - \frac{2(dx+c)}{a^2 - 2ab + b^2} - \frac{b \tan(dx+c)}{(a \tan(dx+c)^2 + b)(a^2 - ab)}$$

$$= - \frac{\dots}{2d}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^2,x, algorithm="giac")

```
[Out] -1/2*((pi*floor((d*x + c)/pi + 1/2)*sgn(a) + arctan(a*tan(d*x + c)/sqrt(a*b)))*(3*a*b - b^2)/((a^3 - 2*a^2*b + a*b^2)*sqrt(a*b)) - 2*(d*x + c)/(a^2 - 2*a*b + b^2) - b*tan(d*x + c)/((a*tan(d*x + c)^2 + b)*(a^2 - a*b)))/d
```

Mupad [B] (verification not implemented)

Time = 12.40 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.23

$$\int \frac{1}{(a + b \cot^2(c + dx))^2} dx = \frac{\frac{ax}{(a-b)^2} + \frac{bx \cot(c+dx)^2}{(a-b)^2} + \frac{b \cot(c+dx)}{2ad(a-b)}}{b \cot(c + dx)^2 + a} + \frac{\operatorname{atan}\left(\frac{b \cot(c+dx)}{\sqrt{ab}}\right) (3ab - b^2)}{\sqrt{ab} (2a^3 d - ab (4ad - 2bd))}$$

[In] int(1/(a + b*cot(c + d*x)^2)^2,x)

```
[Out] ((a*x)/(a - b)^2 + (b*x*cot(c + d*x)^2)/(a - b)^2 + (b*cot(c + d*x))/(2*a*d*(a - b)))/(a + b*cot(c + d*x)^2) + (atan((b*cot(c + d*x))/(a*b)^(1/2)))*(3*a*b - b^2))/((a*b)^(1/2)*(2*a^3*d - a*b*(4*a*d - 2*b*d)))
```

3.7 $\int \frac{1}{(a+b \cot^2(c+dx))^3} dx$

Optimal result	80
Rubi [A] (verified)	80
Mathematica [A] (verified)	82
Maple [A] (verified)	83
Fricas [B] (verification not implemented)	83
Sympy [B] (verification not implemented)	84
Maxima [A] (verification not implemented)	89
Giac [A] (verification not implemented)	89
Mupad [B] (verification not implemented)	90

Optimal result

Integrand size = 14, antiderivative size = 150

$$\int \frac{1}{(a+b \cot^2(c+dx))^3} dx = \frac{x}{(a-b)^3} + \frac{\sqrt{b}(15a^2 - 10ab + 3b^2) \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{8a^{5/2}(a-b)^3 d}$$

$$+ \frac{b \cot(c+dx)}{4a(a-b)d (a+b \cot^2(c+dx))^2}$$

$$+ \frac{(7a-3b)b \cot(c+dx)}{8a^2(a-b)^2 d (a+b \cot^2(c+dx))}$$

[Out] x/(a-b)^3+1/4*b*cot(d*x+c)/a/(a-b)/d/(a+b*cot(d*x+c)^2)^2+1/8*(7*a-3*b)*b*cot(d*x+c)/a^2/(a-b)^2/d/(a+b*cot(d*x+c)^2)+1/8*(15*a^2-10*a*b+3*b^2)*arctan(cot(d*x+c)*b^(1/2)/a^(1/2))*b^(1/2)/a^(5/2)/(a-b)^3/d

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3742, 425, 541, 536, 209, 211}

$$\int \frac{1}{(a+b \cot^2(c+dx))^3} dx = \frac{b(7a-3b) \cot(c+dx)}{8a^2 d (a-b)^2 (a+b \cot^2(c+dx))}$$

$$+ \frac{\sqrt{b}(15a^2 - 10ab + 3b^2) \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{8a^{5/2} d (a-b)^3}$$

$$+ \frac{b \cot(c+dx)}{4ad(a-b) (a+b \cot^2(c+dx))^2} + \frac{x}{(a-b)^3}$$

[In] Int[(a + b*Cot[c + d*x]^2)^(-3), x]

[Out] x/(a - b)^3 + (Sqrt[b]*(15*a^2 - 10*a*b + 3*b^2)*ArcTan[(Sqrt[b]*Cot[c + d*x])/Sqrt[a]])/(8*a^(5/2)*(a - b)^3*d) + (b*Cot[c + d*x])/(4*a*(a - b)*d*(a + b*Cot[c + d*x]^2)^2) + ((7*a - 3*b)*b*Cot[c + d*x])/(8*a^2*(a - b)^2*d*(a + b*Cot[c + d*x]^2))

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 425

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && (!(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1])) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 536

Int[((e_) + (f_)*(x_)^(n_))/((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 541

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 3742

Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(

$\text{ff*x})^n)^p/(c^2 + \text{ff}^2*x^2), x], x, c*(\text{Tan}[e + f*x]/\text{ff}], x]] /; \text{FreeQ}\{a, b, c, e, f, n, p\}, x] \&\& (\text{IntegersQ}[n, p] \parallel \text{IGtQ}[p, 0] \parallel \text{EqQ}[n^2, 4] \parallel \text{EqQ}[n^2, 16])$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x^2)(a+bx^2)^3} dx, x, \cot(c+dx)\right)}{d} \\
 &= \frac{b \cot(c+dx)}{4a(a-b)d(a+b \cot^2(c+dx))^2} - \frac{\text{Subst}\left(\int \frac{4a-3b-3bx^2}{(1+x^2)(a+bx^2)^2} dx, x, \cot(c+dx)\right)}{4a(a-b)d} \\
 &= \frac{b \cot(c+dx)}{4a(a-b)d(a+b \cot^2(c+dx))^2} + \frac{(7a-3b)b \cot(c+dx)}{8a^2(a-b)^2d(a+b \cot^2(c+dx))} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{8a^2-7ab+3b^2-(7a-3b)bx^2}{(1+x^2)(a+bx^2)} dx, x, \cot(c+dx)\right)}{8a^2(a-b)^2d} \\
 &= \frac{b \cot(c+dx)}{4a(a-b)d(a+b \cot^2(c+dx))^2} + \frac{(7a-3b)b \cot(c+dx)}{8a^2(a-b)^2d(a+b \cot^2(c+dx))} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(c+dx)\right)}{(a-b)^3d} \\
 &\quad + \frac{(b(15a^2-10ab+3b^2)) \text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cot(c+dx)\right)}{8a^2(a-b)^3d} \\
 &= \frac{x}{(a-b)^3} + \frac{\sqrt{b}(15a^2-10ab+3b^2) \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{8a^{5/2}(a-b)^3d} \\
 &\quad + \frac{b \cot(c+dx)}{4a(a-b)d(a+b \cot^2(c+dx))^2} + \frac{(7a-3b)b \cot(c+dx)}{8a^2(a-b)^2d(a+b \cot^2(c+dx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.92

$$\begin{aligned}
 &\int \frac{1}{(a+b \cot^2(c+dx))^3} dx \\
 &= \frac{-8 \arctan(\cot(c+dx)) + \frac{\sqrt{b}(15a^2-10ab+3b^2) \arctan\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a}}\right)}{a^{5/2}} + \frac{2(a-b)^2b \cot(c+dx)}{a(a+b \cot^2(c+dx))^2} + \frac{(7a-3b)(a-b)b \cot(c+dx)}{a^2(a+b \cot^2(c+dx))}}{8(a-b)^3d}
 \end{aligned}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(-3), x]

[Out] (-8*ArcTan[Cot[c + d*x]] + (Sqrt[b]*(15*a^2 - 10*a*b + 3*b^2)*ArcTan[(Sqrt[b]*Cot[c + d*x])/Sqrt[a]])/a^(5/2) + (2*(a - b)^2*b*Cot[c + d*x])/(a*(a + b*Cot[c + d*x]^2)^2) + ((7*a - 3*b)*(a - b)*b*Cot[c + d*x])/(a^2*(a + b*Cot[c + d*x]^2)))/(8*(a - b)^3*d)

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 148, normalized size of antiderivative = 0.99

method	result
derivativedivides	$-\frac{\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{(a-b)^3} + \frac{b \left(\frac{b(7a^2-10ab+3b^2)\cot(dx+c)^3}{8a^2} + \frac{(9a^2-14ab+5b^2)\cot(dx+c)}{8a} \right) + \frac{(15a^2-10ab+3b^2)\arctan\left(\frac{b\cot(dx+c)}{\sqrt{ab}}\right)}{8a^2\sqrt{ab}}}{(a-b)^3}$
default	$-\frac{\frac{\pi}{2} - \operatorname{arccot}(\cot(dx+c))}{(a-b)^3} + \frac{b \left(\frac{b(7a^2-10ab+3b^2)\cot(dx+c)^3}{8a^2} + \frac{(9a^2-14ab+5b^2)\cot(dx+c)}{8a} \right) + \frac{(15a^2-10ab+3b^2)\arctan\left(\frac{b\cot(dx+c)}{\sqrt{ab}}\right)}{8a^2\sqrt{ab}}}{d(a-b)^3}$
risch	$\frac{x}{a^3-3a^2b+3ab^2-b^3} - \frac{ib(9a^3e^{6i(dx+c)}+a^2be^{6i(dx+c)}-13ab^2e^{6i(dx+c)}+3b^3e^{6i(dx+c)}-27a^3e^{4i(dx+c)}-9a^2be^{4i(dx+c)}-4(-ae^{4i(dx+c)}+be^{4i(dx+c)}))}{4(-ae^{4i(dx+c)}+be^{4i(dx+c)})}$

[In] int(1/(a+b*cot(d*x+c)^2)^3,x,method=_RETURNVERBOSE)

[Out] 1/d*(-1/(a-b)^3*(1/2*Pi-arccot(cot(d*x+c)))+b/(a-b)^3*((1/8*b*(7*a^2-10*a*b+3*b^2)/a^2*cot(d*x+c)^3+1/8*(9*a^2-14*a*b+5*b^2)/a*cot(d*x+c))/(a+b*cot(d*x+c)^2)^2+1/8*(15*a^2-10*a*b+3*b^2)/a^2/(a*b)^(1/2)*arctan(b*cot(d*x+c)/(a*b)^(1/2))))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 475 vs. 2(136) = 272.

Time = 0.33 (sec) , antiderivative size = 1068, normalized size of antiderivative = 7.12

$$\int \frac{1}{(a+b\cot^2(c+dx))^3} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^3,x, algorithm="fricas")

[Out] [1/32*(32*(a^4 - 2*a^3*b + a^2*b^2)*d*x*cos(2*d*x + 2*c)^2 - 64*(a^4 - a^2*b^2)*d*x*cos(2*d*x + 2*c) + 32*(a^4 + 2*a^3*b + a^2*b^2)*d*x - (15*a^4 + 20*a^3*b - 2*a^2*b^2 - 4*a*b^3 + 3*b^4 + (15*a^4 - 40*a^3*b + 38*a^2*b^2 - 16*a*b^3 + 3*b^4)*cos(2*d*x + 2*c)^2 - 2*(15*a^4 - 10*a^3*b - 12*a^2*b^2 + 10*a*b^3 - 3*b^4)*cos(2*d*x + 2*c))*sqrt(-b/a)*log(((a^2 + 6*a*b + b^2)*cos(2*d*x + 2*c)^2 + 4*(a^2 - a*b - (a^2 + a*b)*cos(2*d*x + 2*c))*sqrt(-b/a)*sin(2*d*x + 2*c) + a^2 - 6*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c)))/((a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*d*x + 2*c)) + 4*(9*a^3*b - 7*a^2*b^2 - 5*a*b^3 + 3*b^4 - 3*(3*a^3*b - 7*a^2*b^2 + 5*a*b^3 - b^4)*cos(2*d*x + 2*c))*sin(2*d*x + 2*c))/((a^7 - 5*a^6*b + 10*a^5*b^2 - 10*a^4*b^3 + 5*a^3*b^4 - a^2*b^5)*d*cos(2*d*x + 2*c)^2 - 2*(a^7 - 3*a^6*b + 2*a^5*b^2 + 2*a^4*b^3 - 3*a^3*b^4 + a^2*b^5)*d*cos(2*d*x +

$2*c) + (a^7 - a^6*b - 2*a^5*b^2 + 2*a^4*b^3 + a^3*b^4 - a^2*b^5)*d)$, $1/16*($
 $16*(a^4 - 2*a^3*b + a^2*b^2)*d*x*\cos(2*d*x + 2*c)^2 - 32*(a^4 - a^2*b^2)*d*$
 $x*\cos(2*d*x + 2*c) + 16*(a^4 + 2*a^3*b + a^2*b^2)*d*x + (15*a^4 + 20*a^3*b$
 $- 2*a^2*b^2 - 4*a*b^3 + 3*b^4 + (15*a^4 - 40*a^3*b + 38*a^2*b^2 - 16*a*b^3$
 $+ 3*b^4)*\cos(2*d*x + 2*c)^2 - 2*(15*a^4 - 10*a^3*b - 12*a^2*b^2 + 10*a*b^3$
 $- 3*b^4)*\cos(2*d*x + 2*c))*\sqrt{b/a}*\arctan(1/2*((a + b)*\cos(2*d*x + 2*c) -$
 $a + b)*\sqrt{b/a}/(b*\sin(2*d*x + 2*c))) + 2*(9*a^3*b - 7*a^2*b^2 - 5*a*b^3$
 $+ 3*b^4 - 3*(3*a^3*b - 7*a^2*b^2 + 5*a*b^3 - b^4)*\cos(2*d*x + 2*c))*\sin(2*d$
 $*x + 2*c))/((a^7 - 5*a^6*b + 10*a^5*b^2 - 10*a^4*b^3 + 5*a^3*b^4 - a^2*b^5)$
 $*d*\cos(2*d*x + 2*c)^2 - 2*(a^7 - 3*a^6*b + 2*a^5*b^2 + 2*a^4*b^3 - 3*a^3*b^4$
 $+ a^2*b^5)*d*\cos(2*d*x + 2*c) + (a^7 - a^6*b - 2*a^5*b^2 + 2*a^4*b^3 + a^3$
 $*b^4 - a^2*b^5)*d)]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 8964 vs. $2(133) = 266$.

Time = 48.20 (sec) , antiderivative size = 8964, normalized size of antiderivative = 59.76

$$\int \frac{1}{(a + b \cot^2(c + dx))^3} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cot(d*x+c)**2)**3,x)

[Out] Piecewise((zoo*x/cot(c)**6, Eq(a, 0) & Eq(b, 0) & Eq(d, 0)), (x/a**3, Eq(b, 0)), ((-x + 1/(d*cot(c + d*x)) - 1/(3*d*cot(c + d*x)**3) + 1/(5*d*cot(c + d*x)**5))/b**3, Eq(a, 0)), (15*d*x*cot(c + d*x)**6/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) + 45*d*x*cot(c + d*x)**4/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) + 45*d*x*cot(c + d*x)**2/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) + 15*d*x/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) - 15*cot(c + d*x)**5/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) - 40*cot(c + d*x)**3/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d) - 33*cot(c + d*x)/(48*b**3*d*cot(c + d*x)**6 + 144*b**3*d*cot(c + d*x)**4 + 144*b**3*d*cot(c + d*x)**2 + 48*b**3*d), Eq(a, b)), (x/(a + b*cot(c)**2)**3, Eq(d, 0)), (16*a**4*d*x*sqrt(-a/b)/(16*a**7*d*sqrt(-a/b) + 32*a**6*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48*a**6*b*d*sqrt(-a/b) + 16*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**4 + 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b**3*d*sqrt(-a/b) + 48*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4) + 15*a**4*log(-sqrt(-a/b) + cot(c + d*x))/(1

$$\begin{aligned}
& 6a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b*d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 48a^{**6}b*d* \\
& \text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5}b^{**2}d*\text{sqrt} \\
& \text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5}b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4}b^{**3}d*\text{sqrt}(- \\
& a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**4}b^{**3} \\
& b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 32a^{**3}b^{**4} \\
& d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4) \\
& - 15a^{**4}*\log(\text{sqrt}(-a/b) + \cot(c + d*x))/(16a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b \\
& *d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 48a^{**6}b*d*\text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt} \\
& \text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5} \\
& b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3} \\
& b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**4}b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4} \\
& d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 32a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 \\
& - 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4) + 32a^{**3}b*d*x*\text{sqrt}(-a/b)*\cot \\
& (c + d*x)**2/(16a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b*d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 \\
& - 48a^{**6}b*d*\text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5} \\
& b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5}b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4} \\
& b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x) \\
&)**2 - 16a^{**4}b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)** \\
& 4 - 32a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot \\
& (c + d*x)**4) + 18a^{**3}b*\text{sqrt}(-a/b)*\cot(c + d*x)/(16a^{**7}d*\text{sqrt}(-a/b) + \\
& 32a^{**6}b*d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 48a^{**6}b*d*\text{sqrt}(-a/b) + 16a^{**5}b^{**2} \\
& b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)** \\
& 2 + 48a^{**5}b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 \\
& + 96a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**4}b^{**3}d*\text{sqrt}(-a/b) + 4 \\
& 8a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 32a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c \\
& + d*x)**2 - 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4) + 30a^{**3}b*\log(-\text{sqrt} \\
& \text{sqrt}(-a/b) + \cot(c + d*x))*\cot(c + d*x)**2/(16a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b*d \\
& *\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 48a^{**6}b*d*\text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt} \\
& (-a/b)*\cot(c + d*x)**4 - 96a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5} \\
& b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3} \\
& b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**4}b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4} \\
& d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 32a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - \\
& 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4) - 10a^{**3}b*\log(-\text{sqrt}(-a/b) + \cot \\
& (c + d*x))/(16a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b*d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 \\
& - 48a^{**6}b*d*\text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5} \\
& b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5}b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4} \\
& b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x) \\
&)**2 - 16a^{**4}b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 \\
& - 32a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**2}b^{**5}d*\text{sqrt}(-a/b)*\cot \\
& (c + d*x)**4) - 30a^{**3}b*\log(\text{sqrt}(-a/b) + \cot(c + d*x))*\cot(c + d*x)**2/(\\
& 16a^{**7}d*\text{sqrt}(-a/b) + 32a^{**6}b*d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 48a^{**6}b*d \\
& *\text{sqrt}(-a/b) + 16a^{**5}b^{**2}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 96a^{**5}b^{**2}d*\text{sqrt} \\
& \text{sqrt}(-a/b)*\cot(c + d*x)**2 + 48a^{**5}b^{**2}d*\text{sqrt}(-a/b) - 48a^{**4}b^{**3}d*\text{sqrt} \\
& (-a/b)*\cot(c + d*x)**4 + 96a^{**4}b^{**3}d*\text{sqrt}(-a/b)*\cot(c + d*x)**2 - 16a^{**4} \\
& b^{**3}d*\text{sqrt}(-a/b) + 48a^{**3}b^{**4}d*\text{sqrt}(-a/b)*\cot(c + d*x)**4 - 32a^{**3}b^{**4}
\end{aligned}$$

$$\begin{aligned}
& + d*x)^{**4} + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} \\
& (-a/b) + 48*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b} \\
& /b)*\cot(c + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4}) - 15*a^{**2} \\
& b^{**2}*\log(\sqrt{-a/b} + \cot(c + d*x))*\cot(c + d*x)^{**4}/(16*a^{**7}*d*\sqrt{-a/b} + \\
& 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 48*a^{**6}*b*d*\sqrt{-a/b} + 16*a^{**5} \\
& b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 96*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x) \\
& **2 + 48*a^{**5}*b^{**2}*d*\sqrt{-a/b} - 48*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} \\
& + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} + 4 \\
& 8*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c \\
& + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4}) + 20*a^{**2}*b^{**2}*\log(s \\
& \sqrt{-a/b} + \cot(c + d*x))*\cot(c + d*x)^{**2}/(16*a^{**7}*d*\sqrt{-a/b} + 32*a^{**6}*b \\
& *d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 48*a^{**6}*b*d*\sqrt{-a/b} + 16*a^{**5}*b^{**2}*d*\sqrt{-a/b} \\
& *d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 96*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} + 48*a \\
& *5*b^{**2}*d*\sqrt{-a/b} - 48*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} + 96*a^{**4} \\
& b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} + 48*a^{**3}*b^{** \\
& 4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} \\
& - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4}) - 3*a^{**2}*b^{**2}*\log(\sqrt{-a/b} + \\
& \cot(c + d*x))/(16*a^{**7}*d*\sqrt{-a/b} + 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + d*x) \\
& **2 - 48*a^{**6}*b*d*\sqrt{-a/b} + 16*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 9 \\
& 6*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} + 48*a^{**5}*b^{**2}*d*\sqrt{-a/b} - 48*a \\
& **4*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d \\
& *x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} + 48*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x) \\
& **4 - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b} \\
& *\cot(c + d*x)^{**4}) - 20*a*b^{**3}*\sqrt{-a/b}*\cot(c + d*x)^{**3}/(16*a^{**7}*d*\sqrt{-a \\
& /b) + 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 48*a^{**6}*b*d*\sqrt{-a/b} + 16* \\
& a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 96*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + \\
& d*x)^{**2} + 48*a^{**5}*b^{**2}*d*\sqrt{-a/b} - 48*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x) \\
&)^{**4} + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} \\
&) + 48*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot \\
& (c + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4}) + 10*a*b^{**3}*\sqrt{-a/b} \\
& *\cot(c + d*x)/(16*a^{**7}*d*\sqrt{-a/b} + 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + \\
& d*x)^{**2} - 48*a^{**6}*b*d*\sqrt{-a/b} + 16*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x) \\
& **4 - 96*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} + 48*a^{**5}*b^{**2}*d*\sqrt{-a/b} \\
& - 48*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot \\
& (c + d*x)^{**2} - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} + 48*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c \\
& + d*x)^{**4} - 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b} \\
& *\cot(c + d*x)^{**4}) - 10*a*b^{**3}*\log(-\sqrt{-a/b} + \cot(c + d*x))*\cot(c + \\
& d*x)^{**4}/(16*a^{**7}*d*\sqrt{-a/b} + 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 4 \\
& 8*a^{**6}*b*d*\sqrt{-a/b} + 16*a^{**5}*b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - 96*a^{**5} \\
& *b^{**2}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} + 48*a^{**5}*b^{**2}*d*\sqrt{-a/b} - 48*a^{**4}*b \\
& **3*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} + 96*a^{**4}*b^{**3}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} \\
& - 16*a^{**4}*b^{**3}*d*\sqrt{-a/b} + 48*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**4} - \\
& 32*a^{**3}*b^{**4}*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 16*a^{**2}*b^{**5}*d*\sqrt{-a/b}*\cot(c \\
& + d*x)^{**4}) + 6*a*b^{**3}*\log(-\sqrt{-a/b} + \cot(c + d*x))*\cot(c + d*x)^{**2}/(16* \\
& a^{**7}*d*\sqrt{-a/b} + 32*a^{**6}*b*d*\sqrt{-a/b}*\cot(c + d*x)^{**2} - 48*a^{**6}*b*d*\sqrt{-a/b}
\end{aligned}$$

```

rt(-a/b) + 16*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(
-a/b)*cot(c + d*x)**2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/
b)*cot(c + d*x)**4 + 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b*
*3*d*sqrt(-a/b) + 48*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*
d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4) +
 10*a*b**3*log(sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**4/(16*a**7*d*sqrt(-
a/b) + 32*a**6*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48*a**6*b*d*sqrt(-a/b) + 16
*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(-a/b)*cot(c +
d*x)**2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/b)*cot(c + d*
x)**4 + 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b**3*d*sqrt(-a/
b) + 48*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*d*sqrt(-a/b)*
cot(c + d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4) - 6*a*b**3*log
(sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**2/(16*a**7*d*sqrt(-a/b) + 32*a**6
*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48*a**6*b*d*sqrt(-a/b) + 16*a**5*b**2*d*s
qrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 48*
a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**4 + 96*a**
4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b**3*d*sqrt(-a/b) + 48*a**3*b
**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**
2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4) + 6*b**4*sqrt(-a/b)*cot(c +
d*x)**3/(16*a**7*d*sqrt(-a/b) + 32*a**6*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48
*a**6*b*d*sqrt(-a/b) + 16*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*
b**2*d*sqrt(-a/b)*cot(c + d*x)**2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**
3*d*sqrt(-a/b)*cot(c + d*x)**4 + 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2
- 16*a**4*b**3*d*sqrt(-a/b) + 48*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 3
2*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c
+ d*x)**4) + 3*b**4*log(-sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**4/(16*a**
7*d*sqrt(-a/b) + 32*a**6*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48*a**6*b*d*sqrt(
-a/b) + 16*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(-a/
b)*cot(c + d*x)**2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/b)*
cot(c + d*x)**4 + 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b**3*
d*sqrt(-a/b) + 48*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*d*s
qrt(-a/b)*cot(c + d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4) - 3*
b**4*log(sqrt(-a/b) + cot(c + d*x))*cot(c + d*x)**4/(16*a**7*d*sqrt(-a/b) +
 32*a**6*b*d*sqrt(-a/b)*cot(c + d*x)**2 - 48*a**6*b*d*sqrt(-a/b) + 16*a**5*
b**2*d*sqrt(-a/b)*cot(c + d*x)**4 - 96*a**5*b**2*d*sqrt(-a/b)*cot(c + d*x)*
*2 + 48*a**5*b**2*d*sqrt(-a/b) - 48*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**4
+ 96*a**4*b**3*d*sqrt(-a/b)*cot(c + d*x)**2 - 16*a**4*b**3*d*sqrt(-a/b) + 4
8*a**3*b**4*d*sqrt(-a/b)*cot(c + d*x)**4 - 32*a**3*b**4*d*sqrt(-a/b)*cot(c
+ d*x)**2 - 16*a**2*b**5*d*sqrt(-a/b)*cot(c + d*x)**4), True))

```


Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.52

$$\int \frac{1}{(a + b \cot^2(c + dx))^3} dx = \frac{(15a^2b - 10ab^2 + 3b^3) \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right)}{(a^5 - 3a^4b + 3a^3b^2 - a^2b^3)\sqrt{ab}} - \frac{(9a^2b - 5ab^2) \tan(dx+c)^3 + (7ab^2 - 3b^3) \tan(dx+c)}{a^4b^2 - 2a^3b^3 + a^2b^4 + (a^6 - 2a^5b + a^4b^2) \tan(dx+c)^4 + 2(a^5b - 2a^4b^2 + a^3b^3) \tan(dx+c)^2 - a^3b^3} - \frac{8(dx+c)}{8d}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^3,x, algorithm="maxima")

[Out] -1/8*((15*a^2*b - 10*a*b^2 + 3*b^3)*arctan(a*tan(d*x + c)/sqrt(a*b)))/((a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*sqrt(a*b)) - ((9*a^2*b - 5*a*b^2)*tan(d*x + c)^3 + (7*a*b^2 - 3*b^3)*tan(d*x + c))/(a^4*b^2 - 2*a^3*b^3 + a^2*b^4 + (a^6 - 2*a^5*b + a^4*b^2)*tan(d*x + c)^4 + 2*(a^5*b - 2*a^4*b^2 + a^3*b^3)*tan(d*x + c)^2 - 8*(d*x + c)/(a^3 - 3*a^2*b + 3*a*b^2 - b^3))/d

Giac [A] (verification not implemented)

none

Time = 0.40 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.37

$$\int \frac{1}{(a + b \cot^2(c + dx))^3} dx = \frac{(15a^2b - 10ab^2 + 3b^3) \left(\pi \left\lfloor \frac{dx+c}{\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(a) + \arctan\left(\frac{a \tan(dx+c)}{\sqrt{ab}}\right) \right)}{(a^5 - 3a^4b + 3a^3b^2 - a^2b^3)\sqrt{ab}} - \frac{8(dx+c)}{a^3 - 3a^2b + 3ab^2 - b^3} - \frac{9a^2b \tan(dx+c)^3 - 5ab^2 \tan(dx+c)^3 + 7ab^2 \tan(dx+c)}{(a^4 - 2a^3b + a^2b^2)(a \tan(dx+c) + b)} - \frac{8(dx+c)}{8d}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^3,x, algorithm="giac")

[Out] -1/8*((15*a^2*b - 10*a*b^2 + 3*b^3)*(pi*floor((d*x + c)/pi + 1/2)*sgn(a) + arctan(a*tan(d*x + c)/sqrt(a*b)))/((a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*sqrt(a*b)) - 8*(d*x + c)/(a^3 - 3*a^2*b + 3*a*b^2 - b^3) - (9*a^2*b*tan(d*x + c)^3 - 5*a*b^2*tan(d*x + c)^3 + 7*a*b^2*tan(d*x + c))/(a^4 - 2*a^3*b + a^2*b^2)*(a*tan(d*x + c) + b)^2))/d

$$\begin{aligned}
& (5280*a^8*b^4*d^2 - 1760*a^9*b^3*d^2 + 256*a^{10}*b^2*d^2)/(64*(a^{10}*d^3 - 6* \\
& a^9*b*d^3 + a^4*b^6*d^3 - 6*a^5*b^5*d^3 + 15*a^6*b^4*d^3 - 20*a^7*b^3*d^3 + \\
& 15*a^8*b^2*d^3)) - (\cot(c + d*x)*(-a^5*b)^{(1/2)}*(15*a^2 - 10*a*b + 3*b^2)* \\
& (256*a^4*b^9*d^2 - 1280*a^5*b^8*d^2 + 2304*a^6*b^7*d^2 - 1280*a^7*b^6*d^2 - \\
& 1280*a^8*b^5*d^2 + 2304*a^9*b^4*d^2 - 1280*a^{10}*b^3*d^2 + 256*a^{11}*b^2*d^2 \\
&))/(512*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d)*(a^8*d^2 - 4*a^7*b*d^2 \\
& + a^4*b^4*d^2 - 4*a^5*b^3*d^2 + 6*a^6*b^2*d^2))*(-a^5*b)^{(1/2)}*(15*a^2 - \\
& 10*a*b + 3*b^2))/(16*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d))*(15*a^2 \\
& - 10*a*b + 3*b^2))/(16*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d)) + \\
& ((-a^5*b)^{(1/2)}*((\cot(c + d*x)*(9*b^7 - 60*a*b^6 + 190*a^2*b^5 - 300*a^3*b^4 \\
& + 289*a^4*b^3)))/(32*(a^8*d^2 - 4*a^7*b*d^2 + a^4*b^4*d^2 - 4*a^5*b^3*d^2 \\
& + 6*a^6*b^2*d^2)) + (((96*a^2*b^{10}*d^2 - 800*a^3*b^9*d^2 + 3040*a^4*b^8*d^2 \\
& - 6816*a^5*b^7*d^2 + 9760*a^6*b^6*d^2 - 9056*a^7*b^5*d^2 + 5280*a^8*b^4*d^2 \\
& - 1760*a^9*b^3*d^2 + 256*a^{10}*b^2*d^2)/(64*(a^{10}*d^3 - 6*a^9*b*d^3 + a^4* \\
& b^6*d^3 - 6*a^5*b^5*d^3 + 15*a^6*b^4*d^3 - 20*a^7*b^3*d^3 + 15*a^8*b^2*d^3) \\
&) + (\cot(c + d*x)*(-a^5*b)^{(1/2)}*(15*a^2 - 10*a*b + 3*b^2)*(256*a^4*b^9*d^2 \\
& - 1280*a^5*b^8*d^2 + 2304*a^6*b^7*d^2 - 1280*a^7*b^6*d^2 - 1280*a^8*b^5*d^2 \\
& + 2304*a^9*b^4*d^2 - 1280*a^{10}*b^3*d^2 + 256*a^{11}*b^2*d^2))/(512*(a^8*d - \\
& a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d)*(a^8*d^2 - 4*a^7*b*d^2 + a^4*b^4*d^2 \\
& - 4*a^5*b^3*d^2 + 6*a^6*b^2*d^2))*(-a^5*b)^{(1/2)}*(15*a^2 - 10*a*b + 3*b^2) \\
&))/(16*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d))*(15*a^2 - 10*a*b + 3* \\
& b^2))/(16*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b*d)))*(-a^5*b)^{(1/2)}*(\\
& 15*a^2 - 10*a*b + 3*b^2)*i)/(8*(a^8*d - a^5*b^3*d + 3*a^6*b^2*d - 3*a^7*b* \\
& d))
\end{aligned}$$

3.8 $\int (1 + \cot^2(x))^{3/2} dx$

Optimal result	93
Rubi [A] (verified)	93
Mathematica [B] (verified)	94
Maple [A] (verified)	95
Fricas [B] (verification not implemented)	95
Sympy [F]	95
Maxima [B] (verification not implemented)	96
Giac [A] (verification not implemented)	96
Mupad [B] (verification not implemented)	96

Optimal result

Integrand size = 10, antiderivative size = 22

$$\int (1 + \cot^2(x))^{3/2} dx = -\frac{1}{2} \operatorname{arcsinh}(\cot(x)) - \frac{1}{2} \cot(x) \sqrt{\csc^2(x)}$$

[Out] $-1/2*\operatorname{arcsinh}(\cot(x))-1/2*\cot(x)*(\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3738, 4207, 201, 221}

$$\int (1 + \cot^2(x))^{3/2} dx = -\frac{1}{2} \operatorname{arcsinh}(\cot(x)) - \frac{1}{2} \cot(x) \sqrt{\csc^2(x)}$$

[In] $\operatorname{Int}[(1 + \operatorname{Cot}[x]^2)^{(3/2)}, x]$

[Out] $-1/2*\operatorname{ArcSinh}[\operatorname{Cot}[x]] - (\operatorname{Cot}[x]*\operatorname{Sqrt}[\operatorname{Csc}[x]^2])/2$

Rule 201

$\operatorname{Int}[(a_+ + (b_+)(x_+)^{(n_+)})^{(p_+)}, x_Symbol] \rightarrow \operatorname{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \operatorname{Dist}[a*n*(p/(n*p + 1)), \operatorname{Int}[(a + b*x^n)^{(p - 1)}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a, b]
```

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \csc^2(x)^{3/2} dx \\
 &= -\text{Subst}\left(\int \sqrt{1+x^2} dx, x, \cot(x)\right) \\
 &= -\frac{1}{2} \cot(x) \sqrt{\csc^2(x)} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \cot(x)\right) \\
 &= -\frac{1}{2} \text{arcsinh}(\cot(x)) - \frac{1}{2} \cot(x) \sqrt{\csc^2(x)}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 51 vs. 2(22) = 44.

Time = 0.13 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.32

$$\begin{aligned}
 \int (1 + \cot^2(x))^{3/2} dx &= \frac{1}{8} \sqrt{\csc^2(x)} \left(-\csc^2\left(\frac{x}{2}\right) \right. \\
 &\quad \left. - 4 \log\left(\cos\left(\frac{x}{2}\right)\right) + 4 \log\left(\sin\left(\frac{x}{2}\right)\right) + \sec^2\left(\frac{x}{2}\right) \right) \sin(x)
 \end{aligned}$$

```
[In] Integrate[(1 + Cot[x]^2)^(3/2), x]
```

```
[Out] (Sqrt[Csc[x]^2]*(-Csc[x/2]^2 - 4*Log[Cos[x/2]] + 4*Log[Sin[x/2]] + Sec[x/2]^2)*Sin[x])/8
```

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.86

method	result
derivativedivides	$-\frac{\cot(x)\sqrt{\cot(x)^2+1}}{2} - \frac{\operatorname{arcsinh}(\cot(x))}{2}$
default	$-\frac{\cot(x)\sqrt{\cot(x)^2+1}}{2} - \frac{\operatorname{arcsinh}(\cot(x))}{2}$
risch	$-\frac{i\sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}+1)}{e^{2ix}-1} - \sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}+1)\sin(x) + \sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}-1)\sin(x)$

[In] `int((cot(x)^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] `-1/2*cot(x)*(cot(x)^2+1)^(1/2)-1/2*arcsinh(cot(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 91 vs. $2(16) = 32$.

Time = 0.27 (sec) , antiderivative size = 91, normalized size of antiderivative = 4.14

$$\int (1 + \cot^2(x))^{3/2} dx = \frac{2\sqrt{2}\sqrt{-\frac{1}{\cos(2x)-1}}(\cos(2x)+1) + \log\left(\frac{1}{2}\sqrt{2}\sqrt{-\frac{1}{\cos(2x)-1}}\sin(2x)+1\right)\sin(2x) - \log\left(-\frac{1}{2}\sqrt{2}\sqrt{-\frac{1}{\cos(2x)-1}}\sin(2x)+1\right)\sin(2x)}{4\sin(2x)}$$

[In] `integrate((1+cot(x)^2)^(3/2),x, algorithm="fricas")`

[Out] `-1/4*(2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*(cos(2*x) + 1) + log(1/2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*sin(2*x) + 1)*sin(2*x) - log(-1/2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*sin(2*x) + 1)*sin(2*x))/sin(2*x)`

Sympy [F]

$$\int (1 + \cot^2(x))^{3/2} dx = \int (\cot^2(x) + 1)^{\frac{3}{2}} dx$$

[In] `integrate((1+cot(x)**2)**(3/2),x)`

[Out] `Integral((cot(x)**2 + 1)**(3/2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 300 vs. $2(16) = 32$.

Time = 0.37 (sec) , antiderivative size = 300, normalized size of antiderivative = 13.64

$$\int (1 + \cot^2(x))^{3/2} dx = \frac{-4(\cos(3x) + \cos(x))\cos(4x) - 4(2\cos(2x) - 1)\cos(3x) - 8\cos(2x)\cos(x) + (2(2\cos(2x) - 1)\cos($$

[In] integrate((1+cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] $-1/4*(4*(\cos(3*x) + \cos(x))*\cos(4*x) - 4*(2*\cos(2*x) - 1)*\cos(3*x) - 8*\cos(2*x)*\cos(x) + (2*(2*\cos(2*x) - 1)*\cos(4*x) - \cos(4*x)^2 - 4*\cos(2*x)^2 - \sin(4*x)^2 + 4*\sin(4*x)*\sin(2*x) - 4*\sin(2*x)^2 + 4*\cos(2*x) - 1)*\log(\cos(x)^2 + \sin(x)^2 + 2*\cos(x) + 1) - (2*(2*\cos(2*x) - 1)*\cos(4*x) - \cos(4*x)^2 - 4*\cos(2*x)^2 - \sin(4*x)^2 + 4*\sin(4*x)*\sin(2*x) - 4*\sin(2*x)^2 + 4*\cos(2*x) - 1)*\log(\cos(x)^2 + \sin(x)^2 - 2*\cos(x) + 1) + 4*(\sin(3*x) + \sin(x))*\sin(4*x) - 8*\sin(3*x)*\sin(2*x) - 8*\sin(2*x)*\sin(x) + 4*\cos(x))/(2*(2*\cos(2*x) - 1)*\cos(4*x) - \cos(4*x)^2 - 4*\cos(2*x)^2 - \sin(4*x)^2 + 4*\sin(4*x)*\sin(2*x) - 4*\sin(2*x)^2 + 4*\cos(2*x) - 1)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.45

$$\int (1 + \cot^2(x))^{3/2} dx = \frac{1}{4} \left(\frac{2 \cos(x)}{\cos(x)^2 - 1} - \log(\cos(x) + 1) + \log(-\cos(x) + 1) \right) \operatorname{sgn}(\sin(x))$$

[In] integrate((1+cot(x)^2)^(3/2),x, algorithm="giac")

[Out] $1/4*(2*\cos(x)/(\cos(x)^2 - 1) - \log(\cos(x) + 1) + \log(-\cos(x) + 1))*\operatorname{sgn}(\sin(x))$

Mupad [B] (verification not implemented)

Time = 13.18 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.82

$$\int (1 + \cot^2(x))^{3/2} dx = -\frac{\operatorname{asinh}(\cot(x))}{2} - \frac{\cot(x) \sqrt{\cot(x)^2 + 1}}{2}$$

[In] int((cot(x)^2 + 1)^(3/2),x)

[Out] $-\operatorname{asinh}(\cot(x))/2 - (\cot(x)*(\cot(x)^2 + 1)^(1/2))/2$

3.9 $\int \sqrt{1 + \cot^2(x)} dx$

Optimal result	97
Rubi [A] (verified)	97
Mathematica [B] (verified)	98
Maple [A] (verified)	98
Fricas [B] (verification not implemented)	99
Sympy [F]	99
Maxima [B] (verification not implemented)	99
Giac [A] (verification not implemented)	100
Mupad [B] (verification not implemented)	100

Optimal result

Integrand size = 10, antiderivative size = 5

$$\int \sqrt{1 + \cot^2(x)} dx = -\operatorname{arcsinh}(\cot(x))$$

[Out] $-\operatorname{arcsinh}(\cot(x))$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 5, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3738, 4207, 221}

$$\int \sqrt{1 + \cot^2(x)} dx = -\operatorname{arcsinh}(\cot(x))$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[1 + \operatorname{Cot}[x]^2], x]$

[Out] $-\operatorname{ArcSinh}[\operatorname{Cot}[x]]$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /;$ $\operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ \operatorname{GtQ}[a, 0] \ \&\& \ \operatorname{PosQ}[b]$

Rule 3738

$\operatorname{Int}[(u_)*((a_) + (b_)*\tan[(e_) + (f_)*(x_)]^2)^{(p_)}, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ActivateTrig}[u*(a*\sec[e + f*x]^2)^p], x] /;$ $\operatorname{FreeQ}[\{a, b, e, f, p\}, x] \ \&\& \ \operatorname{EqQ}[a, b]$

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \sqrt{\csc^2(x)} dx \\ &= -\text{Subst}\left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \cot(x)\right) \\ &= -\text{arcsinh}(\cot(x)) \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 28 vs. 2(5) = 10.

Time = 0.03 (sec) , antiderivative size = 28, normalized size of antiderivative = 5.60

$$\int \sqrt{1 + \cot^2(x)} dx = \sqrt{\csc^2(x)} \left(-\log\left(\cos\left(\frac{x}{2}\right)\right) + \log\left(\sin\left(\frac{x}{2}\right)\right) \right) \sin(x)$$

[In] Integrate[Sqrt[1 + Cot[x]^2], x]

[Out] Sqrt[Csc[x]^2]*(-Log[Cos[x/2]] + Log[Sin[x/2]])*Sin[x]

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.20

method	result	size
derivativdivides	$-\text{arcsinh}(\cot(x))$	6
default	$-\text{arcsinh}(\cot(x))$	6
risch	$-2\sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix} + 1) \sin(x) + 2\sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix} - 1) \sin(x)$	62

[In] int((cot(x)^2+1)^(1/2), x, method=_RETURNVERBOSE)

[Out] -arcsinh(cot(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(5) = 10.

Time = 0.29 (sec) , antiderivative size = 53, normalized size of antiderivative = 10.60

$$\int \sqrt{1 + \cot^2(x)} dx = -\frac{1}{2} \log \left(\frac{1}{2} \sqrt{2} \sqrt{-\frac{1}{\cos(2x) - 1} \sin(2x) + 1} \right) + \frac{1}{2} \log \left(-\frac{1}{2} \sqrt{2} \sqrt{-\frac{1}{\cos(2x) - 1} \sin(2x) + 1} \right)$$

[In] integrate((1+cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/2*log(1/2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*sin(2*x) + 1) + 1/2*log(-1/2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*sin(2*x) + 1)

Sympy [F]

$$\int \sqrt{1 + \cot^2(x)} dx = \int \sqrt{\cot^2(x) + 1} dx$$

[In] integrate((1+cot(x)**2)**(1/2),x)

[Out] Integral(sqrt(cot(x)**2 + 1), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(5) = 10.

Time = 0.36 (sec) , antiderivative size = 35, normalized size of antiderivative = 7.00

$$\int \sqrt{1 + \cot^2(x)} dx = -\frac{1}{2} \log(\cos(x)^2 + \sin(x)^2 + 2 \cos(x) + 1) + \frac{1}{2} \log(\cos(x)^2 + \sin(x)^2 - 2 \cos(x) + 1)$$

[In] integrate((1+cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/2*log(cos(x)^2 + sin(x)^2 + 2*cos(x) + 1) + 1/2*log(cos(x)^2 + sin(x)^2 - 2*cos(x) + 1)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 2.00

$$\int \sqrt{1 + \cot^2(x)} dx = \log \left(\left| \tan \left(\frac{1}{2} x \right) \right| \right) \operatorname{sgn}(\sin(x))$$

[In] integrate((1+cot(x)^2)^(1/2),x, algorithm="giac")

[Out] log(abs(tan(1/2*x)))*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 12.97 (sec) , antiderivative size = 5, normalized size of antiderivative = 1.00

$$\int \sqrt{1 + \cot^2(x)} dx = -\operatorname{asinh}(\cot(x))$$

[In] int((cot(x)^2 + 1)^(1/2),x)

[Out] -asinh(cot(x))

3.10 $\int \frac{1}{\sqrt{1+\cot^2(x)}} dx$

Optimal result	101
Rubi [A] (verified)	101
Mathematica [A] (verified)	102
Maple [A] (verified)	102
Fricas [B] (verification not implemented)	103
Sympy [A] (verification not implemented)	103
Maxima [A] (verification not implemented)	103
Giac [B] (verification not implemented)	104
Mupad [B] (verification not implemented)	104

Optimal result

Integrand size = 10, antiderivative size = 12

$$\int \frac{1}{\sqrt{1+\cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{\csc^2(x)}}$$

[Out] $-\cot(x)/(\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3738, 4207, 197}

$$\int \frac{1}{\sqrt{1+\cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{\csc^2(x)}}$$

[In] `Int[1/Sqrt[1 + Cot[x]^2], x]`

[Out] `-(Cot[x]/Sqrt[Csc[x]^2])`

Rule 197

`Int[((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 3738

`Int[(u_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_.)]^2)^(p_), x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{\sqrt{\csc^2(x)}} dx \\ &= -\text{Subst}\left(\int \frac{1}{(1+x^2)^{3/2}} dx, x, \cot(x)\right) \\ &= -\frac{\cot(x)}{\sqrt{\csc^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{\csc^2(x)}}$$

[In] Integrate[1/Sqrt[1 + Cot[x]^2],x]

[Out] -(Cot[x]/Sqrt[Csc[x]^2])

Maple [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.08

method	result	size
derivativedivides	$-\frac{\cot(x)}{\sqrt{\cot(x)^2+1}}$	13
default	$-\frac{\cot(x)}{\sqrt{\cot(x)^2+1}}$	13
risch	$-\frac{i e^{2ix}}{2 \sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}} (e^{2ix}-1)}} - \frac{i}{2(e^{2ix}-1) \sqrt{-\frac{e^{2ix}}{(e^{2ix}-1)^2}}}}$	67

[In] int(1/(cot(x)^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] -cot(x)/(cot(x)^2+1)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21 vs. 2(10) = 20.

Time = 0.26 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.75

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = -\frac{1}{2} \sqrt{2} \sqrt{-\frac{1}{\cos(2x) - 1}} \sin(2x)$$

[In] integrate(1/(1+cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/2*sqrt(2)*sqrt(-1/(cos(2*x) - 1))*sin(2*x)

Sympy [A] (verification not implemented)

Time = 0.19 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{\cot^2(x) + 1}}$$

[In] integrate(1/(1+cot(x)**2)**(1/2),x)

[Out] -cot(x)/sqrt(cot(x)**2 + 1)

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = -\frac{1}{\sqrt{\tan(x)^2 + 1}}$$

[In] integrate(1/(1+cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/sqrt(tan(x)^2 + 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 28 vs. 2(10) = 20.

Time = 0.28 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.33

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = \frac{2}{\left(\frac{\cos(x)-1}{\cos(x)+1} - 1\right) \operatorname{sgn}(\sin(x))} + 2 \operatorname{sgn}(\sin(x))$$

[In] integrate(1/(1+cot(x)^2)^(1/2),x, algorithm="giac")

[Out] 2/(((cos(x) - 1)/(cos(x) + 1) - 1)*sgn(sin(x))) + 2*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{1 + \cot^2(x)}} dx = -\frac{\sin(2x)}{2\sqrt{\sin(x)^2}}$$

[In] int(1/(cot(x)^2 + 1)^(1/2),x)

[Out] -sin(2*x)/(2*(sin(x)^2)^(1/2))

3.11 $\int (-1 - \cot^2(x))^{3/2} dx$

Optimal result	105
Rubi [A] (verified)	105
Mathematica [A] (verified)	107
Maple [A] (verified)	107
Fricas [C] (verification not implemented)	107
Sympy [F]	108
Maxima [B] (verification not implemented)	108
Giac [C] (verification not implemented)	108
Mupad [B] (verification not implemented)	109

Optimal result

Integrand size = 12, antiderivative size = 35

$$\int (-1 - \cot^2(x))^{3/2} dx = -\frac{1}{2} \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right) + \frac{1}{2} \cot(x) \sqrt{-\csc^2(x)}$$

[Out] $-1/2*\arctan(\cot(x)/(-\csc(x)^2)^{(1/2)})+1/2*\cot(x)*(-\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3738, 4207, 201, 223, 209}

$$\int (-1 - \cot^2(x))^{3/2} dx = \frac{1}{2} \cot(x) \sqrt{-\csc^2(x)} - \frac{1}{2} \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right)$$

[In] $\text{Int}[(-1 - \text{Cot}[x]^2)^{(3/2)}, x]$

[Out] $-1/2*\text{ArcTan}[\text{Cot}[x]/\text{Sqrt}[-\text{Csc}[x]^2]] + (\text{Cot}[x]*\text{Sqrt}[-\text{Csc}[x]^2])/2$

Rule 201

$\text{Int}[(a + b*x^n)^p, x_Symbol] \rightarrow \text{Simp}[x*(a + b*x^n)^p/(n*p + 1), x] + \text{Dist}[a*n*(p/(n*p + 1)), \text{Int}[(a + b*x^n)^{p-1}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]
```

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int (-\csc^2(x))^{3/2} dx \\
 &= \text{Subst}\left(\int \sqrt{-1-x^2} dx, x, \cot(x)\right) \\
 &= \frac{1}{2} \cot(x) \sqrt{-\csc^2(x)} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{\sqrt{-1-x^2}} dx, x, \cot(x)\right) \\
 &= \frac{1}{2} \cot(x) \sqrt{-\csc^2(x)} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right) \\
 &= -\frac{1}{2} \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right) + \frac{1}{2} \cot(x) \sqrt{-\csc^2(x)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.37

$$\int (-1 - \cot^2(x))^{3/2} dx = -\frac{\csc\left(\frac{x}{2}\right) \left(\cot(x) \csc(x) + \log\left(\cos\left(\frac{x}{2}\right)\right) - \log\left(\sin\left(\frac{x}{2}\right)\right)\right) \sec\left(\frac{x}{2}\right)}{4\sqrt{-\csc^2(x)}}$$

[In] Integrate[(-1 - Cot[x]^2)^(3/2), x]

[Out] -1/4*(Csc[x/2]*(Cot[x]*Csc[x] + Log[Cos[x/2]] - Log[Sin[x/2]])*Sec[x/2])/Sqrt[-Csc[x]^2]

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.91

method	result	size
derivativedivides	$\frac{\cot(x)\sqrt{-1-\cot(x)^2}}{2} - \frac{\arctan\left(\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}\right)}{2}$	32
default	$\frac{\cot(x)\sqrt{-1-\cot(x)^2}}{2} - \frac{\arctan\left(\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}\right)}{2}$	32
risch	$\frac{i\sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}+1)}{e^{2ix}-1} - \sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}-1)\sin(x) + \sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}+1)\sin(x)$	95

[In] int((-1-cot(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/2*cot(x)*(-1-cot(x)^2)^(1/2)-1/2*arctan(cot(x)/(-1-cot(x)^2)^(1/2))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 73, normalized size of antiderivative = 2.09

$$\int (-1 - \cot^2(x))^{3/2} dx = \frac{(-i e^{(4ix)} + 2i e^{(2ix)} - i) \log(e^{(ix)} + 1) + (i e^{(4ix)} - 2i e^{(2ix)} + i) \log(e^{(ix)} - 1) + 2i e^{(3ix)}}{2(e^{(4ix)} - 2e^{(2ix)} + 1)}$$

[In] integrate((-1-cot(x)^2)^(3/2), x, algorithm="fricas")

[Out] 1/2*((-I*e^(4*I*x) + 2*I*e^(2*I*x) - I)*log(e^(I*x) + 1) + (I*e^(4*I*x) - 2*I*e^(2*I*x) + I)*log(e^(I*x) - 1) + 2*I*e^(3*I*x) + 2*I*e^(I*x))/(e^(4*I*x) - 2*e^(2*I*x) + 1)

Sympy [F]

$$\int (-1 - \cot^2(x))^{3/2} dx = \int (-\cot^2(x) - 1)^{\frac{3}{2}} dx$$

[In] integrate((-1-cot(x)**2)**(3/2),x)

[Out] Integral((-cot(x)**2 - 1)**(3/2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 284 vs. 2(27) = 54.

Time = 0.39 (sec) , antiderivative size = 284, normalized size of antiderivative = 8.11

$$\int (-1 - \cot^2(x))^{3/2} dx = \frac{(2(2 \cos(2x) - 1) \cos(4x) - \cos(4x)^2 - 4 \cos(2x)^2 - \sin(4x)^2 + 4 \sin(4x) \sin(2x) - \dots}{\dots}$$

[In] integrate((-1-cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] 1/2*((2*(2*cos(2*x) - 1)*cos(4*x) - cos(4*x)^2 - 4*cos(2*x)^2 - sin(4*x)^2 + 4*sin(4*x)*sin(2*x) - 4*sin(2*x)^2 + 4*cos(2*x) - 1)*arctan2(sin(x), cos(x) + 1) - (2*(2*cos(2*x) - 1)*cos(4*x) - cos(4*x)^2 - 4*cos(2*x)^2 - sin(4*x)^2 + 4*sin(4*x)*sin(2*x) - 4*sin(2*x)^2 + 4*cos(2*x) - 1)*arctan2(sin(x), cos(x) - 1) + 2*(sin(3*x) + sin(x))*cos(4*x) - 2*(cos(3*x) + cos(x))*sin(4*x) - 2*(2*cos(2*x) - 1)*sin(3*x) + 4*cos(3*x)*sin(2*x) + 4*cos(x)*sin(2*x) - 4*cos(2*x)*sin(x) + 2*sin(x))/(2*(2*cos(2*x) - 1)*cos(4*x) - cos(4*x)^2 - 4*cos(2*x)^2 - sin(4*x)^2 + 4*sin(4*x)*sin(2*x) - 4*sin(2*x)^2 + 4*cos(2*x) - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.97

$$\int (-1 - \cot^2(x))^{3/2} dx = -\frac{1}{4} \left(\frac{2i \cos(x)}{\cos(x)^2 - 1} - i \log(\cos(x) + 1) + i \log(-\cos(x) + 1) \right) \operatorname{sgn}(\sin(x))$$

[In] integrate((-1-cot(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/4*(2*I*cos(x)/(cos(x)^2 - 1) - I*log(cos(x) + 1) + I*log(-cos(x) + 1))*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.89

$$\int (-1 - \cot^2(x))^{3/2} dx = \frac{\cot(x) \sqrt{-\cot(x)^2 - 1}}{2} - \frac{\operatorname{atan}\left(\frac{\cot(x)}{\sqrt{-\cot(x)^2 - 1}}\right)}{2}$$

[In] `int((- cot(x)^2 - 1)^(3/2),x)`

[Out] `(cot(x)*(- cot(x)^2 - 1)^(1/2))/2 - atan(cot(x)/(- cot(x)^2 - 1)^(1/2))/2`

3.12 $\int \sqrt{-1 - \cot^2(x)} dx$

Optimal result	110
Rubi [A] (verified)	110
Mathematica [B] (verified)	111
Maple [A] (verified)	112
Fricas [C] (verification not implemented)	112
Sympy [F]	112
Maxima [A] (verification not implemented)	113
Giac [C] (verification not implemented)	113
Mupad [B] (verification not implemented)	113

Optimal result

Integrand size = 12, antiderivative size = 14

$$\int \sqrt{-1 - \cot^2(x)} dx = \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right)$$

[Out] $\arctan(\cot(x)/(-\csc(x)^2)^{(1/2)})$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3738, 4207, 223, 209}

$$\int \sqrt{-1 - \cot^2(x)} dx = \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right)$$

[In] $\text{Int}[\text{Sqrt}[-1 - \text{Cot}[x]^2], x]$

[Out] $\text{ArcTan}[\text{Cot}[x]/\text{Sqrt}[-\text{Csc}[x]^2]]$

Rule 209

$\text{Int}[(a_ + (b_ \cdot (x_)^2)^{-1}), x_Symbol] := \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 223

$\text{Int}[1/\text{Sqrt}[(a_ + (b_ \cdot (x_)^2)], x_Symbol] := \text{Subst}[\text{Int}[1/(1 - b \cdot x^2), x], x, x/\text{Sqrt}[a + b \cdot x^2]] /; \text{FreeQ}\{a, b\}, x \ \&\& \ !\text{GtQ}[a, 0]$

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]
```

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \sqrt{-\csc^2(x)} \, dx \\
 &= \text{Subst}\left(\int \frac{1}{\sqrt{-1-x^2}} \, dx, x, \cot(x)\right) \\
 &= \text{Subst}\left(\int \frac{1}{1+x^2} \, dx, x, \frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right) \\
 &= \arctan\left(\frac{\cot(x)}{\sqrt{-\csc^2(x)}}\right)
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 30 vs. 2(14) = 28.

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 2.14

$$\int \sqrt{-1 - \cot^2(x)} \, dx = \frac{\csc(x) (\log(\cos(\frac{x}{2})) - \log(\sin(\frac{x}{2})))}{\sqrt{-\csc^2(x)}}$$

```
[In] Integrate[Sqrt[-1 - Cot[x]^2], x]
```

```
[Out] (Csc[x]*(Log[Cos[x/2]] - Log[Sin[x/2]]))/Sqrt[-Csc[x]^2]
```

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$\arctan\left(\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}\right)$	15
default	$\arctan\left(\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}\right)$	15
risch	$-2\sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}+1)\sin(x) + 2\sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}} \ln(e^{ix}-1)\sin(x)$	60

[In] `int((-1-cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `arctan(cot(x)/(-1-cot(x)^2)^(1/2))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.36

$$\int \sqrt{-1 - \cot^2(x)} dx = i \log(e^{ix} + 1) - i \log(e^{ix} - 1)$$

[In] `integrate((-1-cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `I*log(e^(I*x) + 1) - I*log(e^(I*x) - 1)`

Sympy [F]

$$\int \sqrt{-1 - \cot^2(x)} dx = \int \sqrt{-\cot^2(x) - 1} dx$$

[In] `integrate((-1-cot(x)**2)**(1/2),x)`

[Out] `Integral(sqrt(-cot(x)**2 - 1), x)`

Maxima [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.21

$$\int \sqrt{-1 - \cot^2(x)} dx = -\arctan(\sin(x), \cos(x) + 1) + \arctan(\sin(x), \cos(x) - 1)$$

[In] integrate((-1-cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -arctan2(sin(x), cos(x) + 1) + arctan2(sin(x), cos(x) - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.79

$$\int \sqrt{-1 - \cot^2(x)} dx = i \log \left(\left| \tan \left(\frac{1}{2} x \right) \right| \right) \operatorname{sgn}(\sin(x))$$

[In] integrate((-1-cot(x)^2)^(1/2),x, algorithm="giac")

[Out] I*log(abs(tan(1/2*x)))*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 12.87 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 - \cot^2(x)} dx = \operatorname{atan} \left(\frac{\cot(x)}{\sqrt{-\cot(x)^2 - 1}} \right)$$

[In] int((-cot(x)^2 - 1)^(1/2),x)

[Out] atan(cot(x)/(-cot(x)^2 - 1)^(1/2))

3.13 $\int \frac{1}{\sqrt{-1-\cot^2(x)}} dx$

Optimal result	114
Rubi [A] (verified)	114
Mathematica [A] (verified)	115
Maple [A] (verified)	115
Fricas [C] (verification not implemented)	116
Sympy [F]	116
Maxima [A] (verification not implemented)	116
Giac [C] (verification not implemented)	116
Mupad [B] (verification not implemented)	117

Optimal result

Integrand size = 12, antiderivative size = 14

$$\int \frac{1}{\sqrt{-1-\cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{-\csc^2(x)}}$$

[Out] $-\cot(x)/(-\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3738, 4207, 197}

$$\int \frac{1}{\sqrt{-1-\cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{-\csc^2(x)}}$$

[In] `Int[1/Sqrt[-1 - Cot[x]^2], x]`

[Out] `-(Cot[x]/Sqrt[-Csc[x]^2])`

Rule 197

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^(p + 1)/a), x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 3738

`Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]`

Rule 4207

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{\sqrt{-\csc^2(x)}} dx \\ &= \text{Subst}\left(\int \frac{1}{(-1-x^2)^{3/2}} dx, x, \cot(x)\right) \\ &= -\frac{\cot(x)}{\sqrt{-\csc^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = -\frac{\cot(x)}{\sqrt{-\csc^2(x)}}$$

[In] Integrate[1/Sqrt[-1 - Cot[x]^2],x]

[Out] -(Cot[x]/Sqrt[-Csc[x]^2])

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.07

method	result	size
derivativeldivides	$-\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}$	15
default	$-\frac{\cot(x)}{\sqrt{-1-\cot(x)^2}}$	15
risch	$-\frac{ie^{2ix}}{2\sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} - \frac{i}{2(e^{2ix}-1)\sqrt{\frac{e^{2ix}}{(e^{2ix}-1)^2}}}}$	65

[In] int(1/(-1-cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -cot(x)/(-1-cot(x)^2)^(1/2)

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = \frac{1}{2} (-i e^{(2ix)} - i) e^{(-ix)}$$

[In] integrate(1/(-1-cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/2*(-I*e^(2*I*x) - I)*e^(-I*x)

Sympy [F]

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = \int \frac{1}{\sqrt{-\cot^2(x) - 1}} dx$$

[In] integrate(1/(-1-cot(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(-cot(x)**2 - 1), x)

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = -\frac{1}{\sqrt{-\tan(x)^2 - 1}}$$

[In] integrate(1/(-1-cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/sqrt(-tan(x)^2 - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 28, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = -\frac{2i}{\left(\frac{\cos(x)-1}{\cos(x)+1} - 1\right) \operatorname{sgn}(\sin(x))} - 2i \operatorname{sgn}(\sin(x))$$

[In] integrate(1/(-1-cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -2*I/(((cos(x) - 1)/(cos(x) + 1) - 1)*sgn(sin(x))) - 2*I*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.20 (sec) , antiderivative size = 13, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sqrt{-1 - \cot^2(x)}} dx = \frac{\sin(2x) \operatorname{li}}{2 \sqrt{\sin(x)^2}}$$

[In] int(1/(-cot(x)^2 - 1)^(1/2),x)

[Out] (sin(2*x)*li)/(2*(sin(x)^2)^(1/2))

3.14 $\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx$

Optimal result	118
Rubi [A] (verified)	118
Mathematica [A] (verified)	119
Maple [A] (verified)	119
Fricas [A] (verification not implemented)	120
Sympy [F]	120
Maxima [A] (verification not implemented)	121
Giac [A] (verification not implemented)	121
Mupad [B] (verification not implemented)	121

Optimal result

Integrand size = 17, antiderivative size = 28

$$\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx = -\frac{1}{\sqrt{a \csc^2(x)}} - \frac{\sqrt{a \csc^2(x)}}{a}$$

[Out] $-1/(a*\csc(x)^2)^{(1/2)}-(a*\csc(x)^2)^{(1/2)}/a$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {3738, 4209, 45}

$$\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx = -\frac{\sqrt{a \csc^2(x)}}{a} - \frac{1}{\sqrt{a \csc^2(x)}}$$

[In] `Int[Cot[x]^3/Sqrt[a + a*Cot[x]^2],x]`

[Out] $-(1/\text{Sqrt}[a*\text{Csc}[x]^2]) - \text{Sqrt}[a*\text{Csc}[x]^2]/a$

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
```

[a, b]

Rule 4209

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.),
x_Symbol] :> Dist[b/(2*f), Subst[Int[(-1 + x)^((m - 1)/2)*(b*x)^(p - 1), x]
, x, Sec[e + f*x]^2], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\cot^3(x)}{\sqrt{a \csc^2(x)}} dx \\
&= -\left(\frac{1}{2}a \text{Subst}\left(\int \frac{-1+x}{(ax)^{3/2}} dx, x, \csc^2(x)\right)\right) \\
&= -\left(\frac{1}{2}a \text{Subst}\left(\int \left(-\frac{1}{(ax)^{3/2}} + \frac{1}{a\sqrt{ax}}\right) dx, x, \csc^2(x)\right)\right) \\
&= -\frac{1}{\sqrt{a \csc^2(x)}} - \frac{\sqrt{a \csc^2(x)}}{a}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.68

$$\int \frac{\cot^3(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{-1 - \csc^2(x)}{\sqrt{a \csc^2(x)}}$$

[In] Integrate[Cot[x]^3/Sqrt[a + a*Cot[x]^2], x]

[Out] (-1 - Csc[x]^2)/Sqrt[a*Csc[x]^2]

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.04

method	result	size
derivativedivides	$-\frac{\sqrt{a+a \cot(x)^2}}{a} - \frac{1}{\sqrt{a+a \cot(x)^2}}$	29
default	$-\frac{\sqrt{a+a \cot(x)^2}}{a} - \frac{1}{\sqrt{a+a \cot(x)^2}}$	29
risch	$-\frac{e^{4ix}-6e^{2ix}+1}{2\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)^2}$	45

[In] `int(cot(x)^3/(a+a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/a*(a+a*cot(x)^2)^(1/2)-1/(a+a*cot(x)^2)^(1/2)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.96

$$\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\sqrt{2}\sqrt{-\frac{a}{\cos(2x)-1}}(\cos(2x)-3)}{2a}$$

[In] `integrate(cot(x)^3/(a+a*cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `1/2*sqrt(2)*sqrt(-a/(cos(2*x) - 1))*(cos(2*x) - 3)/a`

Sympy [F]

$$\int \frac{\cot^3(x)}{\sqrt{a+a \cot^2(x)}} dx = \int \frac{\cot^3(x)}{\sqrt{a(\cot^2(x)+1)}} dx$$

[In] `integrate(cot(x)**3/(a+a*cot(x)**2)**(1/2),x)`

[Out] `Integral(cot(x)**3/sqrt(a*(cot(x)**2 + 1)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.86

$$\int \frac{\cot^3(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{1}{\sqrt{\frac{a}{\sin(x)^2}}} - \frac{\sqrt{\frac{a}{\sin(x)^2}}}{a}$$

[In] integrate(cot(x)^3/(a+a*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/sqrt(a/sin(x)^2) - sqrt(a/sin(x)^2)/a

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.89

$$\int \frac{\cot^3(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{\sqrt{a} \sin(x) + \frac{\sqrt{a}}{\sin(x)}}{a \operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)^3/(a+a*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -(sqrt(a)*sin(x) + sqrt(a)/sin(x))/(a*sgn(sin(x)))

Mupad [B] (verification not implemented)

Time = 13.10 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.61

$$\int \frac{\cot^3(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{\sin(x)^2 + 1}{\sqrt{a} \sqrt{\sin(x)^2}}$$

[In] int(cot(x)^3/(a + a*cot(x)^2)^(1/2),x)

[Out] -(sin(x)^2 + 1)/(a^(1/2)*(sin(x)^2)^(1/2))

3.15 $\int \frac{\cot^2(x)}{\sqrt{a+a \cot^2(x)}} dx$

Optimal result	122
Rubi [A] (verified)	122
Mathematica [A] (verified)	124
Maple [A] (verified)	124
Fricas [B] (verification not implemented)	124
Sympy [F]	125
Maxima [A] (verification not implemented)	125
Giac [A] (verification not implemented)	125
Mupad [F(-1)]	125

Optimal result

Integrand size = 17, antiderivative size = 31

$$\int \frac{\cot^2(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\cot(x)}{\sqrt{a \csc^2(x)}} - \frac{\operatorname{arctanh}(\cos(x)) \csc(x)}{\sqrt{a \csc^2(x)}}$$

[Out] $\cot(x)/(a*\csc(x)^2)^{(1/2)}-\operatorname{arctanh}(\cos(x))*\csc(x)/(a*\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3738, 4210, 2672, 327, 212}

$$\int \frac{\cot^2(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\cot(x)}{\sqrt{a \csc^2(x)}} - \frac{\csc(x)\operatorname{arctanh}(\cos(x))}{\sqrt{a \csc^2(x)}}$$

[In] $\text{Int}[\text{Cot}[x]^2/\text{Sqrt}[a + a*\text{Cot}[x]^2], x]$

[Out] $\text{Cot}[x]/\text{Sqrt}[a*\text{Csc}[x]^2] - (\text{ArcTanh}[\text{Cos}[x]]*\text{Csc}[x])/ \text{Sqrt}[a*\text{Csc}[x]^2]$

Rule 212

$\text{Int}[\left((a_) + (b_)*(x_)^2\right)^{-1}, x_Symbol] \rightarrow \text{Simp}[\left(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2])\right)*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 327

$\text{Int}[\left((c_)*(x_)\right)^{m_*}\left((a_) + (b_)*(x_)^n\right)^{p_}, x_Symbol] \rightarrow \text{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1})/(b*(m+n*p+1))), x] - \text{Dist}[\dots]$

```
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2672

```
Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a, b]
```

Rule 4210

```
Int[(u_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] :> With[{ff
= FreeFactors[Sec[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*(b*Sec[e + f*x]^
n)^FracPart[p]/(Sec[e + f*x]/ff)^(n*FracPart[p])], Int[ActivateTrig[u]*(Sec
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\cot^2(x)}{\sqrt{a \csc^2(x)}} dx \\
&= \frac{\csc(x) \int \cos(x) \cot(x) dx}{\sqrt{a \csc^2(x)}} \\
&= -\frac{\csc(x) \text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \cos(x)\right)}{\sqrt{a \csc^2(x)}} \\
&= \frac{\cot(x)}{\sqrt{a \csc^2(x)}} - \frac{\csc(x) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(x)\right)}{\sqrt{a \csc^2(x)}} \\
&= \frac{\cot(x)}{\sqrt{a \csc^2(x)}} - \frac{\text{arctanh}(\cos(x)) \csc(x)}{\sqrt{a \csc^2(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.03

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\csc(x) (\cos(x) - \log(\cos(\frac{x}{2})) + \log(\sin(\frac{x}{2})))}{\sqrt{a \csc^2(x)}}$$

[In] Integrate[Cot[x]^2/Sqrt[a + a*Cot[x]^2],x]

[Out] (Csc[x]*(Cos[x] - Log[Cos[x/2]] + Log[Sin[x/2]]))/Sqrt[a*Csc[x]^2]

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.23

method	result	size
derivativedivides	$-\frac{\ln(\sqrt{a} \cot(x) + \sqrt{a + a \cot^2(x)})}{\sqrt{a}} + \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}}$	38
default	$-\frac{\ln(\sqrt{a} \cot(x) + \sqrt{a + a \cot^2(x)})}{\sqrt{a}} + \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}}$	38
risch	$\frac{ie^{2ix}}{2\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} + \frac{i}{2(e^{2ix}-1)\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}} - \frac{ie^{ix} \ln(e^{ix}+1)}{\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} + \frac{ie^{ix} \ln(e^{ix}-1)}{\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)}$	157

[In] int(cot(x)^2/(a+a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -ln(a^(1/2)*cot(x)+(a+a*cot(x)^2)^(1/2))/a^(1/2)+cot(x)/(a+a*cot(x)^2)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(27) = 54.

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 2.48

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\sqrt{2} \sqrt{-\frac{a}{\cos(2x)-1}} \sin(2x) + \sqrt{a} \log\left(\frac{2\sqrt{2}\sqrt{a}\sqrt{-\frac{a}{\cos(2x)-1}} \sin(2x) - a \cos(2x) - 3a}{\cos(2x)-1}\right)}{2a}$$

[In] integrate(cot(x)^2/(a+a*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/2*(sqrt(2)*sqrt(-a/(cos(2*x) - 1))*sin(2*x) + sqrt(a)*log((2*sqrt(2)*sqrt(a)*sqrt(-a/(cos(2*x) - 1))*sin(2*x) - a*cos(2*x) - 3*a)/(cos(2*x) - 1)))/a

Sympy [F]

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \int \frac{\cot^2(x)}{\sqrt{a(\cot^2(x) + 1)}} dx$$

[In] integrate(cot(x)**2/(a+a*cot(x)**2)**(1/2), x)

[Out] Integral(cot(x)**2/sqrt(a*(cot(x)**2 + 1)), x)

Maxima [A] (verification not implemented)

none

Time = 0.37 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.87

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{\sqrt{-a}(\arctan(\sin(x), \cos(x) + 1) - \arctan(\sin(x), \cos(x) - 1))}{a}$$

[In] integrate(cot(x)^2/(a+a*cot(x)^2)^(1/2), x, algorithm="maxima")

[Out] -sqrt(-a)*(arctan2(sin(x), cos(x) + 1) - arctan2(sin(x), cos(x) - 1))/a

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.58

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{1}{2} \sqrt{a} \left(\frac{2 \cos(x)}{\operatorname{asgn}(\sin(x))} - \frac{\log(\cos(x) + 1)}{\operatorname{asgn}(\sin(x))} + \frac{\log(-\cos(x) + 1)}{\operatorname{asgn}(\sin(x))} \right)$$

[In] integrate(cot(x)^2/(a+a*cot(x)^2)^(1/2), x, algorithm="giac")

[Out] 1/2*sqrt(a)*(2*cos(x)/(a*sgn(sin(x))) - log(cos(x) + 1)/(a*sgn(sin(x))) + log(-cos(x) + 1)/(a*sgn(sin(x))))

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \int \frac{\cot(x)^2}{\sqrt{a \cot(x)^2 + a}} dx$$

[In] int(cot(x)^2/(a + a*cot(x)^2)^(1/2), x)

[Out] int(cot(x)^2/(a + a*cot(x)^2)^(1/2), x)

3.16 $\int \frac{\cot(x)}{\sqrt{a+a \cot^2(x)}} dx$

Optimal result	126
Rubi [A] (verified)	126
Mathematica [A] (verified)	127
Maple [A] (verified)	127
Fricas [B] (verification not implemented)	128
Sympy [A] (verification not implemented)	128
Maxima [A] (verification not implemented)	128
Giac [A] (verification not implemented)	129
Mupad [B] (verification not implemented)	129

Optimal result

Integrand size = 15, antiderivative size = 10

$$\int \frac{\cot(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{1}{\sqrt{a \csc^2(x)}}$$

[Out] $1/(a*\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3738, 4209, 32}

$$\int \frac{\cot(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{1}{\sqrt{a \csc^2(x)}}$$

[In] Int[Cot[x]/Sqrt[a + a*Cot[x]^2],x]

[Out] 1/Sqrt[a*Csc[x]^2]

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 3738

Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]

Rule 4209

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.),
x_Symbol] :> Dist[b/(2*f), Subst[Int[(-1 + x)^((m - 1)/2)*(b*x)^(p - 1), x]
, x, Sec[e + f*x]^2], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{\cot(x)}{\sqrt{a \csc^2(x)}} dx \\ &= -\left(\frac{1}{2}a \text{Subst}\left(\int \frac{1}{(ax)^{3/2}} dx, x, \csc^2(x)\right)\right) \\ &= \frac{1}{\sqrt{a \csc^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{1}{\sqrt{a \csc^2(x)}}$$

[In] Integrate[Cot[x]/Sqrt[a + a*Cot[x]^2],x]

[Out] 1/Sqrt[a*Csc[x]^2]

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

method	result	size
derivativedivides	$\frac{1}{\sqrt{a+a \cot(x)^2}}$	11
default	$\frac{1}{\sqrt{a+a \cot(x)^2}}$	11
risch	$\frac{e^{2ix}}{2\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} - \frac{1}{2(e^{2ix}-1)\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}}$	67

[In] int(cot(x)/(a+a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/(a+a*cot(x)^2)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(8) = 16.

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 2.70

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{\sqrt{2} \sqrt{-\frac{a}{\cos(2x)-1}} (\cos(2x) - 1)}{2a}$$

[In] integrate(cot(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/2*sqrt(2)*sqrt(-a/(cos(2*x) - 1))*(cos(2*x) - 1)/a

Sympy [A] (verification not implemented)

Time = 0.54 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{1}{\sqrt{a \cot^2(x) + a}}$$

[In] integrate(cot(x)/(a+a*cot(x)**2)**(1/2),x)

[Out] 1/sqrt(a*cot(x)**2 + a)

Maxima [A] (verification not implemented)

none

Time = 0.23 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{1}{\sqrt{\frac{a}{\sin(x)^2}}}$$

[In] integrate(cot(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/sqrt(a/sin(x)^2)

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.10

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\sin(x)}{\sqrt{a} \operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] sin(x)/(sqrt(a)*sgn(sin(x)))

Mupad [B] (verification not implemented)

Time = 12.98 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\sqrt{\sin(x)^2}}{\sqrt{a}}$$

[In] int(cot(x)/(a + a*cot(x)^2)^(1/2),x)

[Out] (sin(x)^2)^(1/2)/a^(1/2)

$$3.17 \quad \int \frac{\tan(x)}{\sqrt{a+a \cot^2(x)}} dx$$

Optimal result	130
Rubi [A] (verified)	130
Mathematica [A] (verified)	132
Maple [A] (verified)	132
Fricas [B] (verification not implemented)	132
Sympy [F]	133
Maxima [A] (verification not implemented)	133
Giac [A] (verification not implemented)	133
Mupad [B] (verification not implemented)	134

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \frac{\tan(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a \csc^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{1}{\sqrt{a \csc^2(x)}}$$

[Out] $\operatorname{arctanh}((a*\csc(x)^2)^{(1/2)}/a^{(1/2)})/a^{(1/2)}-1/(a*\csc(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3738, 4209, 53, 65, 213}

$$\int \frac{\tan(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a \csc^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{1}{\sqrt{a \csc^2(x)}}$$

[In] $\text{Int}[\text{Tan}[x]/\text{Sqrt}[a + a*\text{Cot}[x]^2], x]$

[Out] $\text{ArcTanh}[\text{Sqrt}[a*\text{Csc}[x]^2]/\text{Sqrt}[a]]/\text{Sqrt}[a] - 1/\text{Sqrt}[a*\text{Csc}[x]^2]$

Rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ[
n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 213

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-
1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] &&
(LtQ[a, 0] || GtQ[b, 0])
```

Rule 3738

```
Int[(u_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Int[A
ctivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a, b]
```

Rule 4209

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.),
x_Symbol] := Dist[b/(2*f), Subst[Int[(-1 + x)^((m - 1)/2)*(b*x)^(p - 1), x]
, x, Sec[e + f*x]^2], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p] && Inte
gerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\tan(x)}{\sqrt{a \csc^2(x)}} dx \\
&= -\left(\frac{1}{2}a \text{Subst}\left(\int \frac{1}{(-1+x)(ax)^{3/2}} dx, x, \csc^2(x)\right)\right) \\
&= -\frac{1}{\sqrt{a \csc^2(x)}} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{(-1+x)\sqrt{ax}} dx, x, \csc^2(x)\right) \\
&= -\frac{1}{\sqrt{a \csc^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{-1+\frac{x^2}{a}} dx, x, \sqrt{a \csc^2(x)}\right)}{a} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a \csc^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{1}{\sqrt{a \csc^2(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.53

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{-1 + \operatorname{arctanh}(\sin(x)) \csc(x)}{\sqrt{a \csc^2(x)}}$$

[In] Integrate[Tan[x]/Sqrt[a + a*Cot[x]^2],x]

[Out] (-1 + ArcTanh[Sin[x]]*Csc[x])/Sqrt[a*Csc[x]^2]

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.08

method	result	size
default	$-\frac{\sqrt{4}(\sin(x)+\ln(-\cot(x)+\csc(x)-1)-\ln(-\cot(x)+\csc(x)+1))\csc(x)}{2\sqrt{a}\csc(x)^2}$	39
risch	$-\frac{e^{2ix}}{2\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} + \frac{1}{2(e^{2ix}-1)\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}} - \frac{ie^{ix}\ln(e^{ix}-i)}{\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)} + \frac{ie^{ix}\ln(e^{ix}+i)}{\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)}$	157

[In] int(tan(x)/(a+a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/2*4^(1/2)*(sin(x)+ln(-cot(x)+csc(x)-1)-ln(-cot(x)+csc(x)+1))/(a*csc(x)^2)^(1/2)*csc(x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(28) = 56.

Time = 0.28 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.17

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{(\tan(x)^2 + 1)\sqrt{a} \log\left(2a \tan(x)^2 + 2\sqrt{a}\sqrt{\frac{a \tan(x)^2 + a}{\tan(x)^2}} \tan(x)^2 + a\right) - 2\sqrt{\frac{a \tan(x)^2 + a}{\tan(x)^2}} \tan(x)^2}{2(a \tan(x)^2 + a)}$$

[In] integrate(tan(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/2*((tan(x)^2 + 1)*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + a)/tan(x)^2)*tan(x)^2 + a) - 2*sqrt((a*tan(x)^2 + a)/tan(x)^2)*tan(x)^2)/(a*tan(x)^2 + a)

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{a (\cot^2(x) + 1)}} dx$$

[In] integrate(tan(x)/(a+a*cot(x)**2)**(1/2),x)

[Out] Integral(tan(x)/sqrt(a*(cot(x)**2 + 1)), x)

Maxima [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.44

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{1}{2} a \left(\frac{\log \left(-\frac{\sqrt{a} - \sqrt{\frac{a}{\sin(x)^2}}}{\sqrt{a} + \sqrt{\frac{a}{\sin(x)^2}}} \right)}{a^{\frac{3}{2}}} + \frac{2}{a \sqrt{\frac{a}{\sin(x)^2}}} \right)$$

[In] integrate(tan(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/2*a*(log(-(sqrt(a) - sqrt(a/sin(x)^2))/(sqrt(a) + sqrt(a/sin(x)^2)))/a^(3/2) + 2/(a*sqrt(a/sin(x)^2)))

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.33

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{\sin(x)}{\sqrt{a} \operatorname{sgn}(\sin(x))}$$

[In] integrate(tan(x)/(a+a*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -sin(x)/(sqrt(a)*sgn(sin(x)))

Mupad [B] (verification not implemented)

Time = 13.19 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int \frac{\tan(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\operatorname{atanh}\left(\sqrt{\frac{1}{\sin(x)^2}}\right) - \sqrt{\sin(x)^2}}{\sqrt{a}}$$

[In] `int(tan(x)/(a + a*cot(x)^2)^(1/2),x)`

[Out] `(atanh((1/sin(x)^2)^(1/2)) - (sin(x)^2)^(1/2))/a^(1/2)`

3.18 $\int \frac{\tan^2(x)}{\sqrt{a+a \cot^2(x)}} dx$

Optimal result	135
Rubi [A] (verified)	135
Mathematica [A] (verified)	136
Maple [A] (verified)	137
Fricas [A] (verification not implemented)	137
Sympy [F]	137
Maxima [A] (verification not implemented)	138
Giac [A] (verification not implemented)	138
Mupad [B] (verification not implemented)	138

Optimal result

Integrand size = 17, antiderivative size = 29

$$\int \frac{\tan^2(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\cot(x)}{\sqrt{a \csc^2(x)}} + \frac{\csc(x) \sec(x)}{\sqrt{a \csc^2(x)}}$$

[Out] $\cot(x)/(\sqrt{a \csc(x)^2})^{1/2} + \csc(x) \sec(x)/(\sqrt{a \csc(x)^2})^{1/2}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3738, 4210, 2670, 14}

$$\int \frac{\tan^2(x)}{\sqrt{a+a \cot^2(x)}} dx = \frac{\cot(x)}{\sqrt{a \csc^2(x)}} + \frac{\csc(x) \sec(x)}{\sqrt{a \csc^2(x)}}$$

[In] Int[Tan[x]^2/Sqrt[a + a*Cot[x]^2],x]

[Out] Cot[x]/Sqrt[a*Csc[x]^2] + (Csc[x]*Sec[x])/Sqrt[a*Csc[x]^2]

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2670

Int[sin[(e_)+(f_)*(x_)]^(m_)*tan[(e_)+(f_)*(x_)]^(n_), x_Symbol] := Dist[-f^(-1), Subst[Int[(1-x^2)^((m+n-1)/2)/x^n, x], x, Cos[e+f*

x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rule 3738

Int[(u_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^2)^p], x_Symbol] :> Int[ActivateTrig[u*(a*sec[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a, b]

Rule 4210

Int[(u_.)*((b_.)*sec[(e_.) + (f_.)*(x_)]^n)^p], x_Symbol] :> With[{ff = FreeFactors[Sec[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sec[e + f*x]^n)^FracPart[p]/(Sec[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sec[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\tan^2(x)}{\sqrt{a \csc^2(x)}} dx \\
 &= \frac{\csc(x) \int \sin(x) \tan^2(x) dx}{\sqrt{a \csc^2(x)}} \\
 &= -\frac{\csc(x) \text{Subst}\left(\int \frac{1-x^2}{x^2} dx, x, \cos(x)\right)}{\sqrt{a \csc^2(x)}} \\
 &= -\frac{\csc(x) \text{Subst}\left(\int \left(-1 + \frac{1}{x^2}\right) dx, x, \cos(x)\right)}{\sqrt{a \csc^2(x)}} \\
 &= \frac{\cot(x)}{\sqrt{a \csc^2(x)}} + \frac{\csc(x) \sec(x)}{\sqrt{a \csc^2(x)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\cot(x) + \csc(x) \sec(x)}{\sqrt{a \csc^2(x)}}$$

[In] Integrate[Tan[x]^2/Sqrt[a + a*Cot[x]^2],x]

[Out] (Cot[x] + Csc[x]*Sec[x])/Sqrt[a*Csc[x]^2]

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
default	$\frac{\sqrt{4}(\cos(x)+1)^2 \sec(x) \csc(x)}{2\sqrt{a} \csc(x)^2}$	24
risch	$\frac{i(e^{4ix}+6e^{2ix}+1)}{2\sqrt{-\frac{ae^{2ix}}{(e^{2ix}-1)^2}}(e^{2ix}-1)(e^{2ix}+1)}$	55

[In] `int(tan(x)^2/(a+a*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `1/2*4^(1/2)*(cos(x)+1)^2/(a*csc(x)^2)^(1/2)*sec(x)*csc(x)`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.21

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{(\tan(x)^3 + 2 \tan(x)) \sqrt{\frac{a \tan(x)^2 + a}{\tan(x)^2}}}{a \tan(x)^2 + a}$$

[In] `integrate(tan(x)^2/(a+a*cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `(tan(x)^3 + 2*tan(x))*sqrt((a*tan(x)^2 + a)/tan(x)^2)/(a*tan(x)^2 + a)`

Sympy [F]

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \int \frac{\tan^2(x)}{\sqrt{a(\cot^2(x) + 1)}} dx$$

[In] `integrate(tan(x)**2/(a+a*cot(x)**2)**(1/2),x)`

[Out] `Integral(tan(x)**2/sqrt(a*(cot(x)**2 + 1)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.62

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\tan(x)^2 + 2}{\sqrt{\tan(x)^2 + 1} \sqrt{a}}$$

[In] integrate(tan(x)^2/(a+a*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] (tan(x)^2 + 2)/(sqrt(tan(x)^2 + 1)*sqrt(a))

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.86

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = -\frac{2 \operatorname{sgn}(\sin(x))}{\sqrt{a}} + \frac{\frac{1}{\cos(x)} + \cos(x)}{\sqrt{a} \operatorname{sgn}(\sin(x))}$$

[In] integrate(tan(x)^2/(a+a*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -2*sgn(sin(x))/sqrt(a) + (1/cos(x) + cos(x))/(sqrt(a)*sgn(sin(x)))

Mupad [B] (verification not implemented)

Time = 13.35 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.17

$$\int \frac{\tan^2(x)}{\sqrt{a + a \cot^2(x)}} dx = \frac{\tan(x)^3 \sqrt{\frac{1}{\tan(x)^2} + 2 \tan(x)} \sqrt{\frac{1}{\tan(x)^2}}}{\sqrt{a} \sqrt{\tan(x)^2 + 1}}$$

[In] int(tan(x)^2/(a + a*cot(x)^2)^(1/2),x)

[Out] (tan(x)^3*(1/tan(x)^2)^(1/2) + 2*tan(x)*(1/tan(x)^2)^(1/2))/(a^(1/2)*(tan(x)^2 + 1)^(1/2))

3.19 $\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx$

Optimal result	139
Rubi [A] (verified)	139
Mathematica [A] (verified)	141
Maple [A] (verified)	141
Fricas [B] (verification not implemented)	142
Sympy [F]	142
Maxima [F(-2)]	143
Giac [F(-2)]	143
Mupad [B] (verification not implemented)	143

Optimal result

Integrand size = 17, antiderivative size = 66

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = -\sqrt{a-b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a-b}}\right) + \sqrt{a + b \cot^2(x)} - \frac{(a + b \cot^2(x))^{3/2}}{3b}$$

[Out] $-1/3*(a+b*\cot(x)^2)^{(3/2)}/b-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)))*(a-b)^{(1/2)}+(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 81, 52, 65, 214}

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = -\sqrt{a-b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a-b}}\right) - \frac{(a + b \cot^2(x))^{3/2}}{3b} + \sqrt{a + b \cot^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]^3 \operatorname{Sqrt}[a + b \operatorname{Cot}[x]^2], x]$

[Out] $-(\operatorname{Sqrt}[a - b] \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b \operatorname{Cot}[x]^2] / \operatorname{Sqrt}[a - b]]) + \operatorname{Sqrt}[a + b \operatorname{Cot}[x]^2] - (a + b \operatorname{Cot}[x]^2)^{(3/2)} / (3*b)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_. + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}((c + d*x)^n / (b*(m + n + 1))), x] + \operatorname{Dist}[n*((b*c - a*d)/($

```
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
.), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \frac{x^3\sqrt{a+bx^2}}{1+x^2} dx, x, \cot(x)\right)$$

$$\begin{aligned}
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{x\sqrt{a+bx}}{1+x} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{(a+b\cot^2(x))^{3/2}}{3b} + \frac{1}{2}\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1+x} dx, x, \cot^2(x)\right) \\
&= \sqrt{a+b\cot^2(x)} - \frac{(a+b\cot^2(x))^{3/2}}{3b} + \frac{1}{2}(a-b)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= \sqrt{a+b\cot^2(x)} - \frac{(a+b\cot^2(x))^{3/2}}{3b} + \frac{(a-b)\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&= -\sqrt{a-b}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) + \sqrt{a+b\cot^2(x)} - \frac{(a+b\cot^2(x))^{3/2}}{3b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.20 (sec) , antiderivative size = 65, normalized size of antiderivative = 0.98

$$\begin{aligned}
\int \cot^3(x)\sqrt{a+b\cot^2(x)} dx &= -\sqrt{a-b}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) \\
&\quad - \frac{\sqrt{a+b\cot^2(x)}(a-3b+b\cot^2(x))}{3b}
\end{aligned}$$

[In] Integrate[Cot[x]^3*Sqrt[a + b*Cot[x]^2], x]

[Out] -(Sqrt[a - b]*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]]) - (Sqrt[a + b*Cot[x]^2]*(a - 3*b + b*Cot[x]^2))/(3*b)

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.27

method	result	size
derivativedivides	$-\frac{(a+b\cot(x)^2)^{\frac{3}{2}}}{3b} + \sqrt{a+b\cot(x)^2} - \frac{b\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} + \frac{a\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$	84
default	$-\frac{(a+b\cot(x)^2)^{\frac{3}{2}}}{3b} + \sqrt{a+b\cot(x)^2} - \frac{b\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} + \frac{a\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$	84

[In] int(cot(x)^3*(a+b*cot(x)^2)^(1/2), x, method=_RETURNVERBOSE)

```
[Out] -1/3*(a+b*cot(x)^2)^(3/2)/b+(a+b*cot(x)^2)^(1/2)-b/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))+a/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(54) = 108.

Time = 0.34 (sec) , antiderivative size = 330, normalized size of antiderivative = 5.00

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx$$

$$= \frac{3(b \cos(2x) - b) \sqrt{a - b} \log\left(-2(a^2 - 2ab + b^2) \cos(2x)^2 - 2a^2 + b^2 + 2((a - b) \cos(2x))^2 - (2a - b) \cos(2x) + a\right) \sqrt{a - b}}{12(b \cos(2x) - b)} + \frac{3(b \cos(2x) - b) \sqrt{-a + b} \arctan\left(-\frac{\sqrt{-a + b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) - 1)}{(a - b) \cos(2x) - a}\right) + 2((a - 4b) \cos(2x) - a + 2b) \sqrt{-a + b}}{6(b \cos(2x) - b)}$$

```
[In] integrate(cot(x)^3*(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] [1/12*(3*(b*cos(2*x) - b)*sqrt(a - b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 - 2*a^2 + b^2 + 2*((a - b)*cos(2*x)^2 - (2*a - b)*cos(2*x) + a)*sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)) + 4*(a^2 - a*b)*cos(2*x) - 4*((a - 4*b)*cos(2*x) - a + 2*b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(b*cos(2*x) - b), -1/6*(3*(b*cos(2*x) - b)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1)/((a - b)*cos(2*x) - a)) + 2*((a - 4*b)*cos(2*x) - a + 2*b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(b*cos(2*x) - b)]
```

Sympy [F]

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = \int \sqrt{a + b \cot^2(x)} \cot^3(x) dx$$

```
[In] integrate(cot(x)**3*(a+b*cot(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*cot(x)**2)*cot(x)**3, x)
```

Maxima [F(-2)]

Exception generated.

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)^3*(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)

Giac [F(-2)]

Exception generated.

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(x)^3*(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to convert to real sageVARb Error: Bad Argument ValueUnable to convert to real sageVARb Error: Bad Argument Val

Mupad [B] (verification not implemented)

Time = 15.11 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.00

$$\int \cot^3(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{b \cot^2(x) + a} - \frac{(b \cot^2(x) + a)^{3/2}}{3b} + 2 \operatorname{atan} \left(\frac{2 \sqrt{b \cot^2(x) + a} \sqrt{\frac{b}{4} - \frac{a}{4}}}{a - b} \right) \sqrt{\frac{b}{4} - \frac{a}{4}}$$

[In] int(cot(x)^3*(a + b*cot(x)^2)^(1/2),x)

[Out] (a + b*cot(x)^2)^(1/2) - (a + b*cot(x)^2)^(3/2)/(3*b) + 2*atan((2*(a + b*cot(x)^2)^(1/2)*(b/4 - a/4)^(1/2))/(a - b))*(b/4 - a/4)^(1/2)

3.20 $\int \cot(x) \sqrt{a + b \cot^2(x)} dx$

Optimal result	144
Rubi [A] (verified)	144
Mathematica [A] (verified)	146
Maple [A] (verified)	146
Fricas [B] (verification not implemented)	147
Sympy [F]	147
Maxima [F(-2)]	148
Giac [B] (verification not implemented)	148
Mupad [B] (verification not implemented)	148

Optimal result

Integrand size = 15, antiderivative size = 48

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{a-b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a-b}}\right) - \sqrt{a + b \cot^2(x)}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)/(a-b)^{(1/2)})*(a-b)^{(1/2)}-(a+b*\cot(x)^2)^{(1/2)})$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 52, 65, 214}

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{a-b} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a-b}}\right) - \sqrt{a + b \cot^2(x)}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]*\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2], x]$

[Out] $\operatorname{Sqrt}[a - b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a - b]] - \operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```


Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x\sqrt{a+bx^2}}{1+x^2} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1+x} dx, x, \cot^2(x)\right)\right) \\
&= -\sqrt{a+b\cot^2(x)} - \frac{1}{2}(a-b)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= -\sqrt{a+b\cot^2(x)} - \frac{(a-b)\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&= \sqrt{a-b}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) - \sqrt{a+b\cot^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.00

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{a - b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}} \right) - \sqrt{a + b \cot^2(x)}$$

[In] Integrate[Cot[x]*Sqrt[a + b*Cot[x]^2],x]

[Out] Sqrt[a - b]*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]] - Sqrt[a + b*Cot[x]^2]

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.48

method	result	size
derivativedivides	$-\sqrt{a + b \cot(x)^2} + \frac{b \arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}} - \frac{a \arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}}$	71
default	$-\sqrt{a + b \cot(x)^2} + \frac{b \arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}} - \frac{a \arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}}$	71

[In] int(cot(x)*(a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -(a+b*cot(x)^2)^(1/2)+b/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))-a/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(40) = 80$.

Time = 0.34 (sec) , antiderivative size = 248, normalized size of antiderivative = 5.17

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \left[\frac{1}{4} \sqrt{a-b} \log \left(-2(a^2 - 2ab + b^2) \cos(2x)^2 - 2a^2 + b^2 \right. \right. \\ \left. \left. - 2((a-b) \cos(2x)^2 - (2a-b) \cos(2x) + a) \sqrt{a-b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} \right. \right. \\ \left. \left. + 4(a^2 - ab) \cos(2x) \right) \right. \\ \left. - \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}}, \frac{1}{2} \sqrt{-a+b} \arctan \left(-\frac{\sqrt{-a+b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) - 1)}{(a-b) \cos(2x) - a} \right) \right. \\ \left. - \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} \right]$$

[In] integrate(cot(x)*(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*sqrt(a - b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 - 2*a^2 + b^2 - 2*((a - b)*cos(2*x)^2 - (2*a - b)*cos(2*x) + a)*sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)) + 4*(a^2 - a*b)*cos(2*x)) - sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)), 1/2*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1)/((a - b)*cos(2*x) - a)) - sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))]

Sympy [F]

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \int \sqrt{a + b \cot^2(x)} \cot(x) dx$$

[In] integrate(cot(x)*(a+b*cot(x)**2)**(1/2),x)

[Out] Integral(sqrt(a + b*cot(x)**2)*cot(x), x)

Maxima [F(-2)]

Exception generated.

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)*(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(40) = 80.

Time = 0.34 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.98

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx =$$

$$-\frac{1}{2} \left(\sqrt{a-b} \log \left(\left(\sqrt{a-b} \sin(x) - \sqrt{a \sin^2(x) - b \sin^2(x) + b} \right)^2 \right) - \frac{4 \sqrt{a-b} b}{\left(\sqrt{a-b} \sin(x) - \sqrt{a \sin^2(x) - b \sin^2(x) + b} \right)} \right)$$

[In] integrate(cot(x)*(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(sqrt(a - b)*log((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2) - 4*sqrt(a - b)*b/((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2 - b))*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.96 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.10

$$\int \cot(x) \sqrt{a + b \cot^2(x)} dx$$

$$= -\sqrt{b \cot^2(x) + a} - 2 \operatorname{atan} \left(\frac{2 \sqrt{b \cot^2(x) + a} \sqrt{\frac{b}{4} - \frac{a}{4}}}{a - b} \right) \sqrt{\frac{b}{4} - \frac{a}{4}}$$

[In] int(cot(x)*(a + b*cot(x)^2)^(1/2),x)

[Out] -(a + b*cot(x)^2)^(1/2) - 2*atan((2*(a + b*cot(x)^2)^(1/2)*(b/4 - a/4)^(1/2))/(a - b))*(b/4 - a/4)^(1/2)

3.21 $\int \sqrt{a + b \cot^2(x)} \tan(x) dx$

Optimal result	149
Rubi [A] (verified)	149
Mathematica [A] (verified)	151
Maple [B] (verified)	151
Fricas [A] (verification not implemented)	152
Sympy [F]	153
Maxima [F]	153
Giac [B] (verification not implemented)	153
Mupad [B] (verification not implemented)	154

Optimal result

Integrand size = 15, antiderivative size = 60

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx = \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a}} \right) - \sqrt{a - b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}} \right)$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/a^{(1/2)})*a^{(1/2)}-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})*(a-b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 457, 85, 65, 214}

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx = \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a}} \right) - \sqrt{a - b} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}} \right)$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]*\operatorname{Tan}[x], x]$

[Out] $\operatorname{Sqrt}[a]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a]] - \operatorname{Sqrt}[a - b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a - b]]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 85

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(a + b*x),
x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[(e + f*x)^(p - 1)/(c + d*x), x]
, x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[0, p, 1]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{x(1+x^2)} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{\sqrt{a+bx}}{x(1+x)} dx, x, \cot^2(x)\right)\right) \\
&= -\left(\frac{1}{2}a\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cot^2(x)\right)\right) \\
&\quad - \frac{1}{2}(-a+b)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)
\end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 351, normalized size of antiderivative = 5.85

$$\begin{aligned}
& \int \sqrt{a + b \cot^2(x)} \tan(x) dx \\
&= \left[\frac{1}{2} \sqrt{a} \log \left(2a \tan(x)^2 + 2\sqrt{a} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b \right) \right. \\
&\quad + \frac{1}{2} \sqrt{a-b} \log \left(\frac{(2a-b) \tan(x)^2 - 2\sqrt{a-b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b}{\tan(x)^2 + 1} \right), \\
&\quad \quad \quad -\sqrt{-a+b} \arctan \left(-\frac{\sqrt{-a+b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a-b} \right) \\
&\quad + \frac{1}{2} \sqrt{a} \log \left(2a \tan(x)^2 + 2\sqrt{a} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b \right), \\
&\quad \quad \quad -\sqrt{-a} \arctan \left(\frac{\sqrt{-a} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a} \right) \\
&\quad + \frac{1}{2} \sqrt{a-b} \log \left(\frac{(2a-b) \tan(x)^2 - 2\sqrt{a-b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b}{\tan(x)^2 + 1} \right), \\
&\quad \quad \quad \left. -\sqrt{-a} \arctan \left(\frac{\sqrt{-a} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a} \right) - \sqrt{-a+b} \arctan \left(-\frac{\sqrt{-a+b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a-b} \right) \right]
\end{aligned}$$

```
[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x),x, algorithm="fricas")
```

```
[Out] [1/2*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b) + 1/2*sqrt(a - b)*log(((2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(tan(x)^2 + 1)), -sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a - b)) + 1/2*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b), -sqrt(-a)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/a) + 1/2*sqrt(a - b)*log(((2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(tan(x)^2 + 1)), -sqrt(-a)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/a) - sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a - b)]]
```


Sympy [F]

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx = \int \sqrt{a + b \cot^2(x)} \tan(x) dx$$

[In] integrate((a+b*cot(x)**2)**(1/2)*tan(x), x)

[Out] Integral(sqrt(a + b*cot(x)**2)*tan(x), x)

Maxima [F]

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx = \int \sqrt{b \cot^2(x) + a} \tan(x) dx$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x), x, algorithm="maxima")

[Out] integrate(sqrt(b*cot(x)^2 + a)*tan(x), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(48) = 96.

Time = 0.33 (sec) , antiderivative size = 187, normalized size of antiderivative = 3.12

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx$$

$$= \frac{1}{2} \left(\frac{2 \sqrt{a - b} a \arctan \left(\frac{(\sqrt{a - b} \sin(x) - \sqrt{a \sin(x)^2 - b \sin(x)^2 + b})^2 - 2a + b}{2 \sqrt{-a^2 + ab}} \right)}{\sqrt{-a^2 + ab}} + \sqrt{a - b} \log \left(\left(\sqrt{a - b} \sin(x) - \sqrt{a \sin(x)^2 - b \sin(x)^2 + b} \right)^2 - 2a + b \right) \right)$$

$$- \frac{\left(2 \sqrt{a - b} a \arctan \left(-\frac{a - b}{\sqrt{-a^2 + ab}} \right) + \sqrt{-a^2 + ab} \sqrt{a - b} \log(b) \right) \operatorname{sgn}(\sin(x))}{2 \sqrt{-a^2 + ab}}$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x), x, algorithm="giac")

[Out] 1/2*(2*sqrt(a - b)*a*arctan(1/2*((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2 - 2*a + b)/sqrt(-a^2 + a*b))/sqrt(-a^2 + a*b) + sqrt(a - b)*log((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2)*sgn(sin(x)) - 1/2*(2*sqrt(a - b)*a*arctan(-(a - b)/sqrt(-a^2 + a*b)) + sqrt(-a^2 + a*b)*sqrt(a - b)*log(b))*sgn(sin(x))/sqrt(-a^2 + a*b)

Mupad [B] (verification not implemented)

Time = 13.78 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.15

$$\int \sqrt{a + b \cot^2(x)} \tan(x) dx = \operatorname{atanh}\left(\frac{2 a b^3 \sqrt{a-b} \sqrt{a + \frac{b}{\tan(x)^2}}}{2 a b^4 - 2 a^2 b^3}\right) \sqrt{a-b} \\ + \sqrt{a} \operatorname{atanh}\left(\frac{\sqrt{a + \frac{b}{\tan(x)^2}}}{\sqrt{a}}\right)$$

`[In] int(tan(x)*(a + b*cot(x)^2)^(1/2),x)`

```
[Out] atanh((2*a*b^3*(a - b)^(1/2)*(a + b/tan(x)^2)^(1/2))/(2*a*b^4 - 2*a^2*b^3))
*(a - b)^(1/2) + a^(1/2)*atanh((a + b/tan(x)^2)^(1/2)/a^(1/2))
```

3.22 $\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx$

Optimal result	155
Rubi [A] (verified)	155
Mathematica [A] (verified)	157
Maple [B] (verified)	158
Fricas [B] (verification not implemented)	158
Sympy [F]	159
Maxima [F]	159
Giac [F(-2)]	159
Mupad [F(-1)]	160

Optimal result

Integrand size = 17, antiderivative size = 89

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{a - b} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - \frac{(a - 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right)}{2\sqrt{b}} - \frac{1}{2} \cot(x) \sqrt{a + b \cot^2(x)}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})*(a-b)^{(1/2)-1/2*(a-2*b)*\operatorname{arctanh}(\cot(x)*b^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/b^{(1/2)-1/2*\cot(x)*(a+b*\cot(x)^2)^{(1/2)}}$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3751, 489, 537, 223, 212, 385, 209}

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \sqrt{a - b} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - \frac{(a - 2b) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right)}{2\sqrt{b}} - \frac{1}{2} \cot(x) \sqrt{a + b \cot^2(x)}$$

[In] $\text{Int}[\text{Cot}[x]^2*\text{Sqrt}[a + b*\text{Cot}[x]^2], x]$

[Out] $\text{Sqrt}[a - b]*\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[x])/(\text{Sqrt}[a + b*\text{Cot}[x]^2])] - ((a - 2*b)*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Cot}[x])/(\text{Sqrt}[a + b*\text{Cot}[x]^2])]/(2*\text{Sqrt}[b])) - (\text{Cot}[x]*\text{Sqrt}[a + b*\text{Cot}[x]^2])/2$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 489

```
Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(b*(m + n*(p + q) + 1))), x] - Dist[e^n/(b*(m + n*(p + q) + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^p*(c + d*x^n)^(q - 1)*Simp[a*c*(m - n + 1) + (a*d*(m - n + 1) - n*q*(b*c - a*d))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && GtQ[q, 0] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 537

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
```

a1Q[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{x^2\sqrt{a+bx^2}}{1+x^2} dx, x, \cot(x)\right) \\
 &= -\frac{1}{2}\cot(x)\sqrt{a+b\cot^2(x)} + \frac{1}{2}\text{Subst}\left(\int \frac{a+(-a+2b)x^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= -\frac{1}{2}\cot(x)\sqrt{a+b\cot^2(x)} + (a-b)\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &\quad + \frac{1}{2}(-a+2b)\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= -\frac{1}{2}\cot(x)\sqrt{a+b\cot^2(x)} \\
 &\quad + (a-b)\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
 &\quad + \frac{1}{2}(-a+2b)\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
 &= \sqrt{a-b}\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) - \frac{(a-2b)\text{arctanh}\left(\frac{\sqrt{b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{2\sqrt{b}} - \frac{1}{2}\cot(x)\sqrt{a+b\cot^2(x)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 4.41 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.57

$$\int \cot^2(x)\sqrt{a+b\cot^2(x)} dx = -\frac{1}{2}\sqrt{\frac{-a-b+a\cos(2x)-b\cos(2x)}{-1+\cos(2x)}}\cot(x) \\
 - \frac{\left(2\sqrt{a-b}\sqrt{b}\arctan\left(\frac{\sqrt{b+a\tan^2(x)}}{\sqrt{a-b}}\right) + (a-2b)\text{arctanh}\left(\frac{\sqrt{b+a\tan^2(x)}}{\sqrt{b}}\right)\right)\sqrt{a+b\cot^2(x)}\tan(x)}{2\sqrt{b}\sqrt{b+a\tan^2(x)}}$$

[In] Integrate[Cot[x]^2*Sqrt[a + b*Cot[x]^2], x]

[Out] -1/2*(Sqrt[(-a - b + a*Cos[2*x] - b*Cos[2*x])/(-1 + Cos[2*x])]*Cot[x]) - ((2*Sqrt[a - b]*Sqrt[b]*ArcTan[Sqrt[b + a*Tan[x]^2]/Sqrt[a - b]] + (a - 2*b)*ArcTanh[Sqrt[b + a*Tan[x]^2]/Sqrt[b]])*Sqrt[a + b*Cot[x]^2]*Tan[x])/(2*Sqrt[b]*Sqrt[b + a*Tan[x]^2])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(71) = 142.

Time = 0.06 (sec) , antiderivative size = 174, normalized size of antiderivative = 1.96

method	result
derivativedivides	$-\frac{\cot(x)\sqrt{a+b\cot(x)^2}}{2} - \frac{a \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b\cot(x)^2}\right)}{2\sqrt{b}} + \sqrt{b} \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b\cot(x)^2}\right) -$
default	$-\frac{\cot(x)\sqrt{a+b\cot(x)^2}}{2} - \frac{a \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b\cot(x)^2}\right)}{2\sqrt{b}} + \sqrt{b} \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b\cot(x)^2}\right) -$

[In] `int(cot(x)^2*(a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2*\cot(x)*(a+b*\cot(x)^2)^(1/2)-1/2*a/b^(1/2)*\ln(b^(1/2)*\cot(x)+(a+b*\cot(x)^2)^(1/2))+b^(1/2)*\ln(b^(1/2)*\cot(x)+(a+b*\cot(x)^2)^(1/2))- (b^4*(a-b))^(1/2)/b/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*\cot(x)^2)^(1/2)*\cot(x))+ a*(b^4*(a-b))^(1/2)/b^2/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*\cot(x)^2)^(1/2)*\cot(x))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(71) = 142.

Time = 0.31 (sec) , antiderivative size = 768, normalized size of antiderivative = 8.63

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx$$

$$= \frac{2\sqrt{-a+bb} \log\left(- (a-b) \cos(2x) - \sqrt{-a+b} \sqrt{\frac{(a-b)\cos(2x)-a-b}{\cos(2x)-1}} \sin(2x) + b\right) \sin(2x) - (a-2b)\sqrt{b} \log(4b \sin(2x))}{4b \sin(2x)}$$

[In] `integrate(cot(x)^2*(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$[1/4*(2*\sqrt{-a+b}*b*\log(-(a-b)*\cos(2*x) - \sqrt{-a+b}*\sqrt{((a-b)*\cos(2*x) - a - b)/(\cos(2*x) - 1))*\sin(2*x) + b)*\sin(2*x) - (a-2*b)*\sqrt{b}*\log(((a-2*b)*\cos(2*x) - 2*\sqrt{b}*\sqrt{((a-b)*\cos(2*x) - a - b)/(\cos(2*x) - 1))*\sin(2*x) - a - 2*b)/(\cos(2*x) - 1))*\sin(2*x) - 2*(b*\cos(2*x) + b)*\sqrt{((a-b)*\cos(2*x) - a - b)/(\cos(2*x) - 1)))/(b*\sin(2*x)), 1/4*(4*\sqrt{(a-b)*b*\arctan(-\sqrt{a-b}*\sqrt{((a-b)*\cos(2*x) - a - b)/(\cos(2*x) - 1))*\sin(2*x)/((a-b)*\cos(2*x) + a - b))*\sin(2*x) - (a-2*b)*\sqrt{b}*\log(((a-2*b)*\cos(2*x) - 2*\sqrt{b}*\sqrt{((a-b)*\cos(2*x) - a - b)/(\cos(2*x) - 1))*\sin(2*x) - a - 2*b)/(\cos(2*x) - 1))*\sin(2*x) - 2*(b*\cos(2*x) + b)*\sqrt{(($$

```
(a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(b*sin(2*x)), 1/2*((a - 2*b)*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b))*sin(2*x) + sqrt(-a + b)*b*log(-(a - b)*cos(2*x) - sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - (b*cos(2*x) + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(b*sin(2*x)), 1/2*(2*sqrt(a - b)*b*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b))*sin(2*x) + (a - 2*b)*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b))*sin(2*x) - (b*cos(2*x) + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(b*sin(2*x))]
```

Sympy [F]

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \int \sqrt{a + b \cot^2(x)} \cot^2(x) dx$$

```
[In] integrate(cot(x)**2*(a+b*cot(x)**2)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*cot(x)**2)*cot(x)**2, x)
```

Maxima [F]

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \int \sqrt{b \cot^2(x) + a} \cot^2(x) dx$$

```
[In] integrate(cot(x)^2*(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cot(x)^2 + a)*cot(x)^2, x)
```

Giac [F(-2)]

Exception generated.

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(cot(x)^2*(a+b*cot(x)^2)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \cot^2(x) \sqrt{a + b \cot^2(x)} dx = \int \cot(x)^2 \sqrt{b \cot(x)^2 + a} dx$$

```
[In] int(cot(x)^2*(a + b*cot(x)^2)^(1/2),x)
```

```
[Out] int(cot(x)^2*(a + b*cot(x)^2)^(1/2), x)
```


3.23 $\int \sqrt{a + b \cot^2(x)} dx$

Optimal result	161
Rubi [A] (verified)	161
Mathematica [A] (verified)	163
Maple [B] (verified)	163
Fricas [B] (verification not implemented)	164
Sympy [F]	165
Maxima [F(-2)]	165
Giac [B] (verification not implemented)	165
Mupad [F(-1)]	166

Optimal result

Integrand size = 12, antiderivative size = 65

$$\int \sqrt{a + b \cot^2(x)} dx = -\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right) - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)$$

[Out] $-\arctan(\cot(x) \cdot (a-b)^{(1/2)} / (a+b \cdot \cot(x)^2)^{(1/2)}) \cdot (a-b)^{(1/2)} - \operatorname{arctanh}(\cot(x) \cdot b^{(1/2)} / (a+b \cdot \cot(x)^2)^{(1/2)}) \cdot b^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec), antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 399, 223, 212, 385, 209}

$$\int \sqrt{a + b \cot^2(x)} dx = -\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right) - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)$$

[In] `Int[Sqrt[a + b*Cot[x]^2],x]`

[Out] $-(\operatorname{Sqrt}[a-b] \cdot \operatorname{ArcTan}[(\operatorname{Sqrt}[a-b] \cdot \operatorname{Cot}[x]) / \operatorname{Sqrt}[a+b \cdot \operatorname{Cot}[x]^2]]) - \operatorname{Sqrt}[b] \cdot \operatorname{ArcTanh}[(\operatorname{Sqrt}[b] \cdot \operatorname{Cot}[x]) / \operatorname{Sqrt}[a+b \cdot \operatorname{Cot}[x]^2]]$

Rule 209

`Int[((a_) + (b_) * (x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2] * Rt[b, 2])) * ArcTan[Rt[b, 2] * (x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Di
st[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^
n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*
d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{1+x^2} dx, x, \cot(x)\right) \\
&= -\left(b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(x)\right)\right) \\
&\quad + (-a+b)\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= -\left(b\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)\right) \\
&\quad + (-a+b)\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)
\end{aligned}$$

$$= -\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right) - \sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.20

$$\int \sqrt{a+b \cot^2(x)} dx = \sqrt{a-b} \arctan\left(\frac{-\cot(x)\sqrt{a+b \cot^2(x)} + \sqrt{b} \csc^2(x)}{\sqrt{a-b}}\right) + \sqrt{b} \log\left(-\sqrt{b} \cot(x) + \sqrt{a+b \cot^2(x)}\right)$$

[In] Integrate[Sqrt[a + b*Cot[x]^2],x]

[Out] Sqrt[a - b]*ArcTan[(-(Cot[x]*Sqrt[a + b*Cot[x]^2]) + Sqrt[b]*Csc[x]^2)/Sqrt[a - b]] + Sqrt[b]*Log[-(Sqrt[b]*Cot[x]) + Sqrt[a + b*Cot[x]^2]]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 136 vs. 2(53) = 106.

Time = 0.05 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.11

method	result
derivativedivides	$-\sqrt{b} \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b \cot(x)^2}\right) + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(x)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(x)^2}}\right)}{b(a-b)} - \frac{a\sqrt{b^4(a-b)}}{b(a-b)}$
default	$-\sqrt{b} \ln\left(\sqrt{b} \cot(x) + \sqrt{a+b \cot(x)^2}\right) + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(x)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(x)^2}}\right)}{b(a-b)} - \frac{a\sqrt{b^4(a-b)}}{b(a-b)}$

[In] int((a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-b^{(1/2)}*\ln(b^{(1/2)}*\cot(x)+(a+b*\cot(x)^2)^{(1/2)})+(b^4*(a-b))^{(1/2)}/b/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)}*\cot(x))-a*(b^4*(a-b))^{(1/2)}/b^2/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)}*\cot(x))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(53) = 106.

Time = 0.28 (sec) , antiderivative size = 515, normalized size of antiderivative = 7.92

$$\begin{aligned}
 & \int \sqrt{a + b \cot^2(x)} dx \\
 &= \left[\frac{1}{2} \sqrt{-a + b} \log \left(-(a - b) \cos(2x) + \sqrt{-a + b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x) + b \right) \right. \\
 & \quad + \frac{1}{2} \sqrt{b} \log \left(\frac{(a - 2b) \cos(2x) + 2\sqrt{b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x) - a - 2b}{\cos(2x) - 1} \right), \\
 & \quad - \sqrt{a - b} \arctan \left(-\frac{\sqrt{a - b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{(a - b) \cos(2x) + a - b} \right) \\
 & \quad + \frac{1}{2} \sqrt{b} \log \left(\frac{(a - 2b) \cos(2x) + 2\sqrt{b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x) - a - 2b}{\cos(2x) - 1} \right), \sqrt{-b} \arctan \left(\frac{\sqrt{-b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{b \cos(2x) + b} \right) \\
 & \quad + \frac{1}{2} \sqrt{-a + b} \log \left(-(a - b) \cos(2x) + \sqrt{-a + b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x) + b \right), \\
 & \quad - \sqrt{a - b} \arctan \left(-\frac{\sqrt{a - b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{(a - b) \cos(2x) + a - b} \right) \\
 & \quad \left. + \sqrt{-b} \arctan \left(\frac{\sqrt{-b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{b \cos(2x) + b} \right) \right]
 \end{aligned}$$

[In] integrate((a+b*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/2*sqrt(-a + b)*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b) + 1/2*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1), -sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b)) + 1/2*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1), sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b)) + 1/2*sqrt(-a + b)*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b), -sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b)) + sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b))]

Sympy [F]

$$\int \sqrt{a + b \cot^2(x)} dx = \int \sqrt{a + b \cot^2(x)} dx$$

[In] integrate((a+b*cot(x)**2)**(1/2),x)

[Out] Integral(sqrt(a + b*cot(x)**2), x)

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a + b \cot^2(x)} dx = \text{Exception raised: ValueError}$$

[In] integrate((a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(53) = 106.

Time = 0.50 (sec) , antiderivative size = 210, normalized size of antiderivative = 3.23

$$\int \sqrt{a + b \cot^2(x)} dx =$$

$$-\frac{1}{2} \left(\frac{2\sqrt{-a+bb} \arctan \left(\frac{(\sqrt{-a+b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a})^2 + a - 2b}{2\sqrt{ab-b^2}} \right)}{\sqrt{ab-b^2}} + \sqrt{-a+b} \log \left(\left(\sqrt{-a+b} \cos(x) \right. \right. \right.$$

$$\left. \left. \left. \left(2\sqrt{-a+bb} \arctan \left(\frac{\sqrt{-a+b}\sqrt{b}}{\sqrt{ab-b^2}} \right) - \sqrt{ab-b^2} \sqrt{-a+b} \log(-a - 2\sqrt{-a+b}\sqrt{b} + 2b) \right) \right) \operatorname{sgn}(\sin(x)) \right)}{2\sqrt{ab-b^2}} \right)$$

[In] integrate((a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(2*sqrt(-a + b)*b*arctan(1/2*((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 + a - 2*b)/sqrt(a*b - b^2))/sqrt(a*b - b^2) + sqrt(-a + b)*log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2))*sgn(sin(x)) - 1/2*(2*sqrt(-a + b)*b*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) - sqrt(a*b - b^2)*sqrt(-a + b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b))*sgn(sin(x))/sqrt(a*b - b^2)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(x)} dx = \int \sqrt{b \cot(x)^2 + a} dx$$

```
[In] int((a + b*cot(x)^2)^(1/2), x)
```

```
[Out] int((a + b*cot(x)^2)^(1/2), x)
```

3.24 $\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx$

Optimal result	167
Rubi [A] (verified)	167
Mathematica [C] (verified)	169
Maple [B] (warning: unable to verify)	169
Fricas [A] (verification not implemented)	170
Sympy [F]	170
Maxima [F]	171
Giac [B] (verification not implemented)	171
Mupad [F(-1)]	171

Optimal result

Integrand size = 17, antiderivative size = 51

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \sqrt{a - b} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) + \sqrt{a + b \cot^2(x)} \tan(x)$$

[Out] $\arctan(\cot(x) * (a-b)^{(1/2)} / (a+b*\cot(x)^2)^{(1/2)}) * (a-b)^{(1/2)} + (a+b*\cot(x)^2)^{(1/2)} * \tan(x)$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 486, 12, 385, 209}

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \sqrt{a - b} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) + \tan(x) \sqrt{a + b \cot^2(x)}$$

[In] $\text{Int}[\text{Sqrt}[a + b*\text{Cot}[x]^2]*\text{Tan}[x]^2, x]$

[Out] $\text{Sqrt}[a - b]*\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[x])/\text{Sqrt}[a + b*\text{Cot}[x]^2]] + \text{Sqrt}[a + b*\text{Cot}[x]^2]*\text{Tan}[x]$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[u, (b_)*(v_)] /; \text{FreeQ}[b, x]$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 486

Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e*x)^(m+1)*(a + b*x^n)^(p+1)*((c + d*x^n)^q/(a*e*(m+1))), x] - Dist[1/(a*e^n*(m+1)), Int[(e*x)^(m+n)*(a + b*x^n)^p*(c + d*x^n)^(q-1)*Simp[c*b*(m+1) + n*(b*c*(p+1) + a*d*q) + d*(b*(m+1) + b*n*(p+q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[0, q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{x^2(1+x^2)} dx, x, \cot(x)\right) \\
 &= \sqrt{a+b\cot^2(x)} \tan(x) - \text{Subst}\left(\int \frac{-a+b}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= \sqrt{a+b\cot^2(x)} \tan(x) - (-a+b)\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= \sqrt{a+b\cot^2(x)} \tan(x) - (-a+b)\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
 &= \sqrt{a-b} \arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) + \sqrt{a+b\cot^2(x)} \tan(x)
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.10 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.86

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \sqrt{a + b \cot^2(x)} \operatorname{Hypergeometric2F1} \left(-\frac{1}{2}, 1, \frac{1}{2}, -\frac{(a-b) \cot^2(x)}{a + b \cot^2(x)} \right) \tan(x)$$

[In] Integrate[Sqrt[a + b*Cot[x]^2]*Tan[x]^2,x]

[Out] Sqrt[a + b*Cot[x]^2]*Hypergeometric2F1[-1/2, 1, 1/2, -(((a - b)*Cot[x]^2)/(a + b*Cot[x]^2))]*Tan[x]

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 311 vs. 2(43) = 86.

Time = 0.87 (sec) , antiderivative size = 312, normalized size of antiderivative = 6.12

method	result
default	$\frac{\sqrt{4} \sqrt{a+b \cot(x)^2} \left(\ln \left(4 \cos(x) \sqrt{-a+b} \sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} - 4 \cos(x) a + 4 b \cos(x) + 4 \sqrt{-a+b} \sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} \right) a \sin(x) \right)}{1}$

[In] int((a+b*cot(x)^2)^(1/2)*tan(x)^2,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{2} 4^{(1/2)} / (-a+b)^{(1/2)} * (a+b*\cot(x)^2)^{(1/2)} / (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2)^{(1/2)} / (\cos(x)+1) * (\ln(4*\cos(x)*(-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) - 4*\cos(x)*a + 4*b*\cos(x) + 4*(-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) * a*\sin(x) - \ln(4*\cos(x)*(-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) - 4*\cos(x)*a + 4*b*\cos(x) + 4*(-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) * b*\sin(x) + (-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) * \sin(x) + (-a+b)^{(1/2)} * (- (a*\cos(x)^2 - \cos(x)^2*b - a) / (\cos(x)+1)^2) * \tan(x)))$

Fricas [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 193, normalized size of antiderivative = 3.78

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx$$

$$= \left[\frac{1}{4} \sqrt{-a + b} \log \left(-\frac{a^2 \tan(x)^4 - 2(3a^2 - 4ab) \tan(x)^2 + a^2 - 8ab + 8b^2 - 4(a \tan(x)^3 - (a - 2b) \tan(x))}{\tan(x)^4 + 2 \tan(x)^2 + 1} \right) \right. \\ \left. + \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x), \frac{1}{2} \sqrt{a - b} \arctan \left(\frac{2 \sqrt{a - b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)}{a \tan(x)^2 - a + 2b} \right) \right. \\ \left. + \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x) \right]$$

```
[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^2,x, algorithm="fricas")
```

```
[Out] [1/4*sqrt(-a + b)*log(-(a^2*tan(x)^4 - 2*(3*a^2 - 4*a*b)*tan(x)^2 + a^2 - 8
*a*b + 8*b^2 - 4*(a*tan(x)^3 - (a - 2*b)*tan(x))*sqrt(-a + b)*sqrt((a*tan(x)
)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) + sqrt((a*tan(x)^2 + b)/ta
n(x)^2)*tan(x), 1/2*sqrt(a - b)*arctan(2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/
tan(x)^2)*tan(x)/(a*tan(x)^2 - a + 2*b)) + sqrt((a*tan(x)^2 + b)/tan(x)^2)*
tan(x)]
```

Sympy [F]

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \int \sqrt{a + b \cot^2(x)} \tan^2(x) dx$$

```
[In] integrate((a+b*cot(x)**2)**(1/2)*tan(x)**2,x)
```

```
[Out] Integral(sqrt(a + b*cot(x)**2)*tan(x)**2, x)
```

Maxima [F]

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \int \sqrt{b \cot(x)^2 + a} \tan(x)^2 dx$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^2,x, algorithm="maxima")

[Out] integrate(sqrt(b*cot(x)^2 + a)*tan(x)^2, x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 239 vs. 2(43) = 86.

Time = 0.29 (sec) , antiderivative size = 239, normalized size of antiderivative = 4.69

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx$$

$$= \frac{1}{2} \left(\sqrt{-a + b} \log \left(\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a} \right)^2 \right) - \frac{4a\sqrt{-a + b}}{\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a} \right)} \right. \\ \left. - \frac{\left(a\sqrt{-a + b} \log(-a - 2\sqrt{-a + b}\sqrt{b} + 2b) - a\sqrt{b} \log(-a - 2\sqrt{-a + b}\sqrt{b} + 2b) - \sqrt{-a + b} \log(-a - 2\sqrt{-a + b}\sqrt{b} + 2b) \right)}{2 \left(a + \sqrt{-a + b}\sqrt{b} \right)} \right)$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^2,x, algorithm="giac")

[Out] 1/2*(sqrt(-a + b)*log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2) - 4*a*sqrt(-a + b)/((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a))*sgn(sin(x)) - 1/2*(a*sqrt(-a + b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - a*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - sqrt(-a + b)*b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + b^(3/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 2*a*sqrt(-a + b))*sgn(sin(x))/(a + sqrt(-a + b)*sqrt(b) - b)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(x)} \tan^2(x) dx = \int \tan(x)^2 \sqrt{b \cot(x)^2 + a} dx$$

[In] int(tan(x)^2*(a + b*cot(x)^2)^(1/2),x)

[Out] int(tan(x)^2*(a + b*cot(x)^2)^(1/2), x)

3.25 $\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx$

Optimal result	172
Rubi [A] (verified)	172
Mathematica [C] (warning: unable to verify)	174
Maple [B] (verified)	175
Fricas [A] (verification not implemented)	176
Sympy [F]	176
Maxima [F]	177
Giac [B] (verification not implemented)	177
Mupad [F(-1)]	178

Optimal result

Integrand size = 17, antiderivative size = 85

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx = -\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right) - \frac{(3a-b)\sqrt{a+b \cot^2(x)} \tan(x)}{3a} + \frac{1}{3} \sqrt{a+b \cot^2(x)} \tan^3(x)$$

[Out] $-\arctan(\cot(x) \cdot (a-b)^{1/2} / (a+b \cdot \cot(x)^2)^{1/2}) \cdot (a-b)^{1/2} - 1/3 \cdot (3a-b) \cdot (a+b \cdot \cot(x)^2)^{1/2} \cdot \tan(x) / a + 1/3 \cdot (a+b \cdot \cot(x)^2)^{1/2} \cdot \tan(x)^3$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 486, 597, 12, 385, 209}

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx = -\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right) + \frac{1}{3} \tan^3(x) \sqrt{a+b \cot^2(x)} - \frac{(3a-b) \tan(x) \sqrt{a+b \cot^2(x)}}{3a}$$

[In] $\text{Int}[\text{Sqrt}[a + b \cdot \text{Cot}[x]^2] \cdot \text{Tan}[x]^4, x]$

[Out] $-(\text{Sqrt}[a - b] \cdot \text{ArcTan}[(\text{Sqrt}[a - b] \cdot \text{Cot}[x]) / \text{Sqrt}[a + b \cdot \text{Cot}[x]^2]]) - ((3a - b) \cdot \text{Sqrt}[a + b \cdot \text{Cot}[x]^2] \cdot \text{Tan}[x]) / (3a) + (\text{Sqrt}[a + b \cdot \text{Cot}[x]^2] \cdot \text{Tan}[x]^3) / 3$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 486

Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e*x)^(m+1)*(a + b*x^n)^(p+1)*((c + d*x^n)^q/(a*e^(m+1))), x] - Dist[1/(a*e^n*(m+1)), Int[(e*x)^(m+n)*(a + b*x^n)^p*(c + d*x^n)^(q-1)*Simp[c*b*(m+1) + n*(b*c*(p+1) + a*d*q) + d*(b*(m+1) + b*n*(p+q+1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[0, q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 597

Int[((g_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[e*(g*x)^(m+1)*(a + b*x^n)^(p+1)*((c + d*x^n)^(q+1)/(a*c*g*(m+1))), x] + Dist[1/(a*c*g^n*(m+1)), Int[(g*x)^(m+n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m+1) - e*(b*c + a*d)*(m+n+1) - e*n*(b*c*p + a*d*q) - b*e*d*(m+n*(p+q+2)+1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0] && LtQ[m, -1]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{x^4(1+x^2)} dx, x, \cot(x)\right) \\
&= \frac{1}{3}\sqrt{a+b\cot^2(x)}\tan^3(x) - \frac{1}{3}\text{Subst}\left(\int \frac{-3a+b-2bx^2}{x^2(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= -\frac{(3a-b)\sqrt{a+b\cot^2(x)}\tan(x)}{3a} + \frac{1}{3}\sqrt{a+b\cot^2(x)}\tan^3(x) \\
&\quad + \frac{\text{Subst}\left(\int -\frac{3a(a-b)}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{3a} \\
&= -\frac{(3a-b)\sqrt{a+b\cot^2(x)}\tan(x)}{3a} + \frac{1}{3}\sqrt{a+b\cot^2(x)}\tan^3(x) \\
&\quad + (-a+b)\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= -\frac{(3a-b)\sqrt{a+b\cot^2(x)}\tan(x)}{3a} + \frac{1}{3}\sqrt{a+b\cot^2(x)}\tan^3(x) \\
&\quad + (-a+b)\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&= -\sqrt{a-b}\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&\quad - \frac{(3a-b)\sqrt{a+b\cot^2(x)}\tan(x)}{3a} + \frac{1}{3}\sqrt{a+b\cot^2(x)}\tan^3(x)
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 1.76 (sec) , antiderivative size = 174, normalized size of antiderivative = 2.05

$$\begin{aligned}
\int \sqrt{a+b\cot^2(x)}\tan^4(x) dx &= \frac{1}{3}\sqrt{a+b\cot^2(x)}\left(1\right. \\
&\quad \left. + \frac{b\cot^2(x)}{a}\right)\sin^2(x)\left(-\frac{4(a-b)\cos^2(x)(a+b\cot^2(x))\text{Hypergeometric2F1}\left(2, 2, \frac{3}{2}, \frac{(a-b)\cos^2(x)}{a}\right)}{a^2}\right. \\
&\quad \left. + \frac{(a-2b\cot^2(x))\csc^2(x)\left(\arcsin\left(\sqrt{\frac{(a-b)\cos^2(x)}{a}}\right)\sqrt{\frac{(a-b)\cos^2(x)}{a}} + \sqrt{\frac{b\cos^2(x)}{a} + \sin^2(x)}\right)}{(a+b\cot^2(x))\sqrt{\frac{b\cos^2(x)}{a} + \sin^2(x)}}\right)\tan^3(x)
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Cot[x]^2]*Tan[x]^4,x]

[Out] (Sqrt[a + b*Cot[x]^2]*(1 + (b*Cot[x]^2)/a)*Sin[x]^2*((-4*(a - b)*Cos[x]^2*(a + b*Cot[x]^2)*Hypergeometric2F1[2, 2, 3/2, ((a - b)*Cos[x]^2)/a])/a^2 + (a - 2*b*Cot[x]^2)*Csc[x]^2*(ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*Sqrt[((a - b)*Cos[x]^2)/a + Sqrt[(b*Cos[x]^2)/a + Sin[x]^2]])/((a + b*Cot[x]^2)*Sqrt[(b*Cos[x]^2)/a + Sin[x]^2]))*Tan[x]^3)/3

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 494 vs. 2(71) = 142.

Time = 1.51 (sec) , antiderivative size = 495, normalized size of antiderivative = 5.82

method	result
default	$-\frac{\sqrt{4}\left(4\sqrt{-a+b}\sqrt{-\frac{a\cos(x)^2-\cos(x)^2b-a}{(\cos(x)+1)^2}}a\cos(x)^3-\cos(x)^3\sqrt{-\frac{a\cos(x)^2-\cos(x)^2b-a}{(\cos(x)+1)^2}}\sqrt{-a+b}b+3\ln\left(4\cos(x)\sqrt{-a+b}\sqrt{-\frac{a\cos(x)^2-\cos(x)^2b-a}{(\cos(x)+1)^2}}\right)\right)}{\dots}$

[In] int((a+b*cot(x)^2)^(1/2)*tan(x)^4,x,method=_RETURNVERBOSE)

[Out]
$$-1/6*4^{(1/2)}/a/(-a+b)^{(1/2)}*(4*(-a+b)^{(1/2)}*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*a*\cos(x)^3-\cos(x)^3*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*(-a+b)^{(1/2)}*b+3*\ln(4*\cos(x)*(-a+b)^{(1/2)}*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}-4*\cos(x)*a+4*b*\cos(x)+4*(-a+b)^{(1/2)}*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)})*\cos(x)^3*a^2-3*\ln(4*\cos(x)*(-a+b)^{(1/2)}*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}-4*\cos(x)*a+4*b*\cos(x)+4*(-a+b)^{(1/2)}*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)})*\cos(x)^3*a*b+4*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*\cos(x)^2*(-a+b)^{(1/2)}*a-\cos(x)^2*(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*(-a+b)^{(1/2)}*b-(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*(-a+b)^{(1/2)}*a*\cos(x)-(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*(-a+b)^{(1/2)}*a)*(a+b*cot(x)^2)^(1/2)/(\cos(x)+1)/(-(a*\cos(x)^2-\cos(x)^2*b-a)/(\cos(x)+1)^2)^{(1/2)}*\tan(x)*\sec(x)^2$$

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 239, normalized size of antiderivative = 2.81

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx$$

$$= \left[\frac{3 a \sqrt{-a + b} \log \left(-\frac{a^2 \tan(x)^4 - 2(3a^2 - 4ab) \tan(x)^2 + a^2 - 8ab + 8b^2 + 4(a \tan(x)^3 - (a - 2b) \tan(x)) \sqrt{-a + b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{\tan(x)^4 + 2 \tan(x)^2 + 1} \right) + 4(a \tan(x)^3 - (a - 2b) \tan(x)) \sqrt{-a + b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{12 a} \right. \\ \left. - \frac{3 \sqrt{a - b} a \arctan \left(\frac{2 \sqrt{a - b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)}{a \tan(x)^2 - a + 2b} \right) - 2(a \tan(x)^3 - (3a - b) \tan(x)) \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{6 a} \right]$$

```
[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^4,x, algorithm="fricas")
```

```
[Out] [1/12*(3*a*sqrt(-a + b)*log(-(a^2*tan(x)^4 - 2*(3*a^2 - 4*a*b)*tan(x)^2 + a^2 - 8*a*b + 8*b^2 + 4*(a*tan(x)^3 - (a - 2*b)*tan(x))*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) + 4*(a*tan(x)^3 - (3*a - b)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/a, -1/6*(3*sqrt(a - b)*a*arctan(2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)/(a*tan(x)^2 - a + 2*b)) - 2*(a*tan(x)^3 - (3*a - b)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/a]
```

Sympy [F]

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx = \int \sqrt{a + b \cot^2(x)} \tan^4(x) dx$$

```
[In] integrate((a+b*cot(x)**2)**(1/2)*tan(x)**4,x)
```

```
[Out] Integral(sqrt(a + b*cot(x)**2)*tan(x)**4, x)
```


Maxima [F]

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx = \int \sqrt{b \cot(x)^2 + a} \tan(x)^4 dx$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^4,x, algorithm="maxima")

[Out] integrate(sqrt(b*cot(x)^2 + a)*tan(x)^4, x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 476 vs. 2(71) = 142.

Time = 0.33 (sec) , antiderivative size = 476, normalized size of antiderivative = 5.60

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx =$$

$$-\frac{1}{6} \left(3 \sqrt{-a + b} \log \left(\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a} \right)^2 \right) - \frac{4 \left(3 \left(\sqrt{-a + b} \cos(x) \right. \right. \right.}{\left. \left. \left. \left(3 a^2 \sqrt{-a + b} \log \left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2b \right) - 9 a^2 \sqrt{b} \log \left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2b \right) - 15 a \sqrt{-a + b} \right. \right. \right. \right.$$

[In] integrate((a+b*cot(x)^2)^(1/2)*tan(x)^4,x, algorithm="giac")

[Out] -1/6*(3*sqrt(-a + b)*log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2) - 4*(3*(sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^4*(2*a - b)*sqrt(-a + b) - 6*(sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2*a^2*sqrt(-a + b) + (4*a^3 - a^2*b)*sqrt(-a + b))/((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a)^3)*sgn(sin(x)) + 1/6*(3*a^2*sqrt(-a + b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - 9*a^2*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - 15*a*sqrt(-a + b)*b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 21*a*b^(3/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 12*sqrt(-a + b)*b^2*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - 12*b^(5/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 8*a^2*sqrt(-a + b) - 18*a^2*sqrt(b) - 24*a*sqrt(-a + b)*b + 30*a*b^(3/2) + 12*sqrt(-a + b)*b^2 - 12*b^(5/2))*sgn(sin(x))/(a^2 + 3*a*sqrt(-a + b)*sqrt(b) - 5*a*b - 4*sqrt(-a + b)*b^(3/2) + 4*b^2)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(x)} \tan^4(x) dx = \int \tan(x)^4 \sqrt{b \cot(x)^2 + a} dx$$

```
[In] int(tan(x)^4*(a + b*cot(x)^2)^(1/2),x)
```

```
[Out] int(tan(x)^4*(a + b*cot(x)^2)^(1/2), x)
```

3.26 $\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx$

Optimal result	179
Rubi [A] (verified)	179
Mathematica [A] (verified)	181
Maple [B] (verified)	182
Fricas [B] (verification not implemented)	182
Sympy [F]	183
Maxima [F(-2)]	183
Giac [F(-2)]	183
Mupad [B] (verification not implemented)	184

Optimal result

Integrand size = 17, antiderivative size = 88

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = -(a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) + (a - b) \sqrt{a + b \cot^2(x)} + \frac{1}{3} (a + b \cot^2(x))^{3/2} - \frac{(a + b \cot^2(x))^{5/2}}{5b}$$

[Out] $-(a-b)^{(3/2)}*\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})+1/3*(a+b*\cot(x)^2)^{(3/2)}-1/5*(a+b*\cot(x)^2)^{(5/2)}/b+(a-b)*(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 81, 52, 65, 214}

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = -(a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) - \frac{(a + b \cot^2(x))^{5/2}}{5b} + \frac{1}{3} (a + b \cot^2(x))^{3/2} + (a - b) \sqrt{a + b \cot^2(x)}$$

[In] $\operatorname{Int}[\cot[x]^3*(a + b*\cot[x]^2)^{(3/2)}, x]$

[Out] $-\left((a - b)^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\cot[x]^2]/\operatorname{Sqrt}[a - b]]\right) + (a - b)*\operatorname{Sqrt}[a + b*\cot[x]^2] + (a + b*\cot[x]^2)^{(3/2)}/3 - (a + b*\cot[x]^2)^{(5/2)}/(5*b)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_. + (d_.)*(x_))^{(n_)}), x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))], x] + \operatorname{Dist}[n*((b*c - a*d)/($

```
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
.), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \frac{x^3(a + bx^2)^{3/2}}{1 + x^2} dx, x, \cot(x)\right)$$

$$\begin{aligned}
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{x(a+bx)^{3/2}}{1+x} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{(a+b\cot^2(x))^{5/2}}{5b} + \frac{1}{2}\text{Subst}\left(\int \frac{(a+bx)^{3/2}}{1+x} dx, x, \cot^2(x)\right) \\
&= \frac{1}{3}(a+b\cot^2(x))^{3/2} - \frac{(a+b\cot^2(x))^{5/2}}{5b} + \frac{1}{2}(a-b)\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1+x} dx, x, \cot^2(x)\right) \\
&= (a-b)\sqrt{a+b\cot^2(x)} + \frac{1}{3}(a+b\cot^2(x))^{3/2} - \frac{(a+b\cot^2(x))^{5/2}}{5b} \\
&\quad + \frac{1}{2}(a-b)^2\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= (a-b)\sqrt{a+b\cot^2(x)} + \frac{1}{3}(a+b\cot^2(x))^{3/2} - \frac{(a+b\cot^2(x))^{5/2}}{5b} \\
&\quad + \frac{(a-b)^2\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&= -(a-b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) + (a-b)\sqrt{a+b\cot^2(x)} \\
&\quad + \frac{1}{3}(a+b\cot^2(x))^{3/2} - \frac{(a+b\cot^2(x))^{5/2}}{5b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.61 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.03

$$\begin{aligned}
\int \cot^3(x) (a+b\cot^2(x))^{3/2} dx &= -(a-b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) \\
&\quad - \frac{\sqrt{a+b\cot^2(x)}(3a^2-20ab+15b^2+(6a-5b)b\cot^2(x)+3b^2\cot^4(x))}{15b}
\end{aligned}$$

[In] Integrate[Cot[x]^3*(a+b*Cot[x]^2)^(3/2),x]

[Out] -((a-b)^(3/2)*ArcTanh[Sqrt[a+b*Cot[x]^2]/Sqrt[a-b]]) - (Sqrt[a+b*Cot[x]^2]*(3*a^2-20*a*b+15*b^2+(6*a-5*b)*b*Cot[x]^2+3*b^2*Cot[x]^4))/(15*b)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(72) = 144$.

Time = 0.04 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.70

method	result
derivativedivides	$-\frac{(a+b \cot(x)^2)^{\frac{5}{2}}}{5b} + \frac{b \cot(x)^2 \sqrt{a+b \cot(x)^2}}{3} + \frac{4a \sqrt{a+b \cot(x)^2}}{3} - b \sqrt{a+b \cot(x)^2} + \frac{b^2 \arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$
default	$-\frac{(a+b \cot(x)^2)^{\frac{5}{2}}}{5b} + \frac{b \cot(x)^2 \sqrt{a+b \cot(x)^2}}{3} + \frac{4a \sqrt{a+b \cot(x)^2}}{3} - b \sqrt{a+b \cot(x)^2} + \frac{b^2 \arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$

[In] `int(cot(x)^3*(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/5*(a+b*\cot(x)^2)^{(5/2)}/b+1/3*b*\cot(x)^2*(a+b*\cot(x)^2)^{(1/2)}+4/3*a*(a+b*\cot(x)^2)^{(1/2)}-b*(a+b*\cot(x)^2)^{(1/2)}+b^2/(-a+b)^{(1/2)}*\arctan((a+b*\cot(x)^2)^{(1/2)}/(-a+b)^{(1/2)})-2*a*b/(-a+b)^{(1/2)}*\arctan((a+b*\cot(x)^2)^{(1/2)}/(-a+b)^{(1/2)})+a^2/(-a+b)^{(1/2)}*\arctan((a+b*\cot(x)^2)^{(1/2)}/(-a+b)^{(1/2)})$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. $2(72) = 144$.

Time = 0.35 (sec) , antiderivative size = 486, normalized size of antiderivative = 5.52

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = \left[\frac{15 ((ab - b^2) \cos(2x)^2 + ab - b^2 - 2(ab - b^2) \cos(2x)) \sqrt{a - b} \log\left(-2(a^2 - 2ab + b^2) \cos(2x) - a - b\right)}{15 ((ab - b^2) \cos(2x)^2 + ab - b^2 - 2(ab - b^2) \cos(2x)) \sqrt{-a + b} \arctan\left(-\frac{\sqrt{-a+b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) - 1)}{(a-b) \cos(2x) - a}\right)} \right] + \frac{1}{30} (b \cos(2x))$$

[In] `integrate(cot(x)^3*(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")`

[Out]
$$[-1/60*(15*((a*b - b^2)*\cos(2*x)^2 + a*b - b^2 - 2*(a*b - b^2)*\cos(2*x))*\sqrt{a - b}*\log(-2*(a^2 - 2*a*b + b^2)*\cos(2*x)^2 - 2*a^2 + b^2 - 2*((a - b)*\cos(2*x)^2 - (2*a - b)*\cos(2*x) + a)*\sqrt{a - b}*\sqrt{((a - b)*\cos(2*x) - a - b)/(\cos(2*x) - 1)} + 4*(a^2 - a*b)*\cos(2*x)) + 4*((3*a^2 - 26*a*b + 23*b^2)*\cos(2*x)^2 + 3*a^2 - 14*a*b + 13*b^2 - 2*(3*a^2 - 20*a*b + 12*b^2)*\cos(2*x))*\sqrt{((a - b)*\cos(2*x) - a - b)/(\cos(2*x) - 1))]/(b*\cos(2*x)^2 - 2*b*$$

$\cos(2x) + b)$, $-1/30*(15*((a*b - b^2)*\cos(2x)^2 + a*b - b^2 - 2*(a*b - b^2)*\cos(2x))*\sqrt{-a + b}*\arctan(-\sqrt{-a + b}*\sqrt{((a - b)*\cos(2x) - a - b)/(\cos(2x) - 1)}*(\cos(2x) - 1)/((a - b)*\cos(2x) - a)) + 2*((3*a^2 - 26*a*b + 23*b^2)*\cos(2x)^2 + 3*a^2 - 14*a*b + 13*b^2 - 2*(3*a^2 - 20*a*b + 12*b^2)*\cos(2x))*\sqrt{((a - b)*\cos(2x) - a - b)/(\cos(2x) - 1))}/(b*\cos(2x)^2 - 2*b*\cos(2x) + b)]$

Sympy [F]

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = \int (a + b \cot^2(x))^{\frac{3}{2}} \cot^3(x) dx$$

[In] `integrate(cot(x)**3*(a+b*cot(x)**2)**(3/2),x)`

[Out] `Integral((a + b*cot(x)**2)**(3/2)*cot(x)**3, x)`

Maxima [F(-2)]

Exception generated.

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = \text{Exception raised: ValueError}$$

[In] `integrate(cot(x)^3*(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail

Giac [F(-2)]

Exception generated.

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] `integrate(cot(x)^3*(a+b*cot(x)^2)^(3/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to convert to real sageVARb Error: Bad Argument ValueUnable to convert to real sageVARb Error: Bad Argument Val

Mupad [B] (verification not implemented)

Time = 24.41 (sec) , antiderivative size = 120, normalized size of antiderivative = 1.36

$$\int \cot^3(x) (a + b \cot^2(x))^{3/2} dx = \left(\frac{a}{3b} - \frac{a-b}{3b} \right) (b \cot(x)^2 + a)^{3/2} - \frac{(b \cot(x)^2 + a)^{5/2}}{5b}$$

$$+ (a-b) \left(\frac{a}{b} - \frac{a-b}{b} \right) \sqrt{b \cot(x)^2 + a} + \operatorname{atan} \left(\frac{(a-b)^{3/2} \sqrt{b \cot(x)^2 + a} \operatorname{li}}{a^2 - 2ab + b^2} \right) (a-b)^{3/2} \operatorname{li}$$

[In] `int(cot(x)^3*(a + b*cot(x)^2)^(3/2),x)`

[Out] `atan(((a - b)^(3/2)*(a + b*cot(x)^2)^(1/2)*1i)/(a^2 - 2*a*b + b^2))*(a - b)^(3/2)*1i - (a + b*cot(x)^2)^(5/2)/(5*b) + (a/(3*b) - (a - b)/(3*b))*(a + b*cot(x)^2)^(3/2) + (a - b)*(a/b - (a - b)/b)*(a + b*cot(x)^2)^(1/2)`

3.27 $\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx$

Optimal result	185
Rubi [A] (verified)	185
Mathematica [A] (verified)	188
Maple [B] (verified)	188
Fricas [B] (verification not implemented)	189
Sympy [F]	190
Maxima [F]	190
Giac [F(-2)]	190
Mupad [F(-1)]	190

Optimal result

Integrand size = 17, antiderivative size = 127

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = (a - b)^{3/2} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - \frac{(3a^2 - 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right)}{8\sqrt{b}} - \frac{1}{8}(5a - 4b) \cot(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4} b \cot^3(x) \sqrt{a + b \cot^2(x)}$$

[Out] (a-b)^(3/2)*arctan(cot(x)*(a-b)^(1/2)/(a+b*cot(x)^2)^(1/2))-1/8*(3*a^2-12*a*b+8*b^2)*arctanh(cot(x)*b^(1/2)/(a+b*cot(x)^2)^(1/2))/b^(1/2)-1/8*(5*a-4*b)*cot(x)*(a+b*cot(x)^2)^(1/2)-1/4*b*cot(x)^3*(a+b*cot(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.471$, Rules used = {3751, 488, 596, 537, 223, 212, 385, 209}

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = -\frac{(3a^2 - 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right)}{8\sqrt{b}} + (a - b)^{3/2} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - \frac{1}{8}(5a - 4b) \cot(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4} b \cot^3(x) \sqrt{a + b \cot^2(x)}$$

[In] Int[Cot[x]^2*(a + b*Cot[x]^2)^(3/2),x]

[Out] $(a - b)^{3/2} \text{ArcTan}[\text{Sqrt}[a - b] \text{Cot}[x]] / \text{Sqrt}[a + b \text{Cot}[x]^2] - ((3a^2 - 12ab + 8b^2) \text{ArcTanh}[\text{Sqrt}[b] \text{Cot}[x]] / \text{Sqrt}[a + b \text{Cot}[x]^2]) / (8 \text{Sqrt}[b]) - ((5a - 4b) \text{Cot}[x] \text{Sqrt}[a + b \text{Cot}[x]^2]) / 8 - (b \text{Cot}[x]^3 \text{Sqrt}[a + b \text{Cot}[x]^2]) / 4$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 488

Int[((e_.)*(x_)^(m_.))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(b*e*(m + n*(p + q) + 1))), x] + Dist[1/(b*(m + n*(p + q) + 1)), Int[(e*x)^m*(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp[c*((c*b - a*d)*(m + 1) + c*b*n*(p + q)) + (d*(c*b - a*d)*(m + 1) + d*n*(q - 1)*(b*c - a*d) + c*b*d*n*(p + q))*x^n, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && GtQ[q, 1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 537

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 596

```

Int[((g_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[f*g^(n - 1)*(g*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(b*d*(m + n*(p + q + 1) + 1))), x] - Dist[g^n/(b*d*(m + n*(p + q + 1) + 1)), Int[(g*x)^(m - n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m - n + 1) + (a*f*d*(m + n*q + 1) + b*(f*c*(m + n*p + 1) - e*d*(m + n*(p + q + 1) + 1))]*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0] && GtQ[m, n - 1]

```

Rule 3751

```

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^2(a + bx^2)^{3/2}}{1 + x^2} dx, x, \cot(x)\right) \\
&= -\frac{1}{4}b \cot^3(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4} \text{Subst}\left(\int \frac{x^2(a(4a - 3b) + (5a - 4b)bx^2)}{(1 + x^2) \sqrt{a + bx^2}} dx, x, \cot(x)\right) \\
&= -\frac{1}{8}(5a - 4b) \cot(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4}b \cot^3(x) \sqrt{a + b \cot^2(x)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{a(5a - 4b)b - b(3a^2 - 12ab + 8b^2)x^2}{(1 + x^2) \sqrt{a + bx^2}} dx, x, \cot(x)\right)}{8b} \\
&= -\frac{1}{8}(5a - 4b) \cot(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4}b \cot^3(x) \sqrt{a + b \cot^2(x)} \\
&\quad + (a - b)^2 \text{Subst}\left(\int \frac{1}{(1 + x^2) \sqrt{a + bx^2}} dx, x, \cot(x)\right) \\
&\quad + \frac{1}{8}(-3a^2 + 12ab - 8b^2) \text{Subst}\left(\int \frac{1}{\sqrt{a + bx^2}} dx, x, \cot(x)\right) \\
&= -\frac{1}{8}(5a - 4b) \cot(x) \sqrt{a + b \cot^2(x)} - \frac{1}{4}b \cot^3(x) \sqrt{a + b \cot^2(x)} \\
&\quad + (a - b)^2 \text{Subst}\left(\int \frac{1}{1 - (-a + b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}}\right) \\
&\quad + \frac{1}{8}(-3a^2 + 12ab - 8b^2) \text{Subst}\left(\int \frac{1}{1 - bx^2} dx, x, \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}}\right)
\end{aligned}$$

$$= (a-b)^{3/2} \arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) - \frac{(3a^2 - 12ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{8\sqrt{b}}$$

$$- \frac{1}{8}(5a-4b)\cot(x)\sqrt{a+b\cot^2(x)} - \frac{1}{4}b\cot^3(x)\sqrt{a+b\cot^2(x)}$$

Mathematica [A] (verified)

Time = 1.32 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.99

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \frac{\sqrt{-a-b+(a-b)\cos(2x)} \csc(x) \left(8\sqrt{2}(a-b)^2 \sqrt{-b} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{a-b}\cos(x)}{\sqrt{-a-b+(a-b)\cos(2x)}}\right) + \sqrt{a-b}\right)}{8\sqrt{2}\sqrt{a-b}}$$

[In] Integrate[Cot[x]^2*(a + b*Cot[x]^2)^(3/2), x]

[Out] (Sqrt[-a - b + (a - b)*Cos[2*x]]*Csc[x]*(8*Sqrt[2]*(a - b)^2*Sqrt[-b]*ArcTanh[(Sqrt[2]*Sqrt[a - b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]]] + Sqrt[a - b]*(-(Sqrt[2]*(3*a^2 - 12*a*b + 8*b^2)*ArcTanh[(Sqrt[2]*Sqrt[-b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]]]) + Sqrt[-b]*Sqrt[-a - b + (a - b)*Cos[2*x]]*Cot[x]*Csc[x]*(5*a - 6*b + 2*b*Csc[x]^2)))/(8*Sqrt[2]*Sqrt[a - b]*Sqrt[-b]*Sqrt[-((a - b)*Cos[2*x])*Csc[x]^2])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(105) = 210.

Time = 0.03 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.25

method	result
derivativedivides	$-\frac{\cot(x)(a+b\cot(x)^2)^{\frac{3}{2}}}{4} - \frac{3a\cot(x)\sqrt{a+b\cot(x)^2}}{8} - \frac{3a^2 \ln\left(\sqrt{b}\cot(x) + \sqrt{a+b\cot(x)^2}\right)}{8\sqrt{b}} - b^{\frac{3}{2}} \ln\left(\sqrt{b}\cot(x) + \sqrt{a+b\cot(x)^2}\right)$
default	$-\frac{\cot(x)(a+b\cot(x)^2)^{\frac{3}{2}}}{4} - \frac{3a\cot(x)\sqrt{a+b\cot(x)^2}}{8} - \frac{3a^2 \ln\left(\sqrt{b}\cot(x) + \sqrt{a+b\cot(x)^2}\right)}{8\sqrt{b}} - b^{\frac{3}{2}} \ln\left(\sqrt{b}\cot(x) + \sqrt{a+b\cot(x)^2}\right)$

[In] int(cot(x)^2*(a+b*cot(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/4*cot(x)*(a+b*cot(x)^2)^(3/2)-3/8*a*cot(x)*(a+b*cot(x)^2)^(1/2)-3/8*a^2/b^(1/2)*ln(b^(1/2)*cot(x)+(a+b*cot(x)^2)^(1/2))-b^(3/2)*ln(b^(1/2)*cot(x)+(a+b*cot(x)^2)^(1/2))+1/2*b*cot(x)*(a+b*cot(x)^2)^(1/2)+3/2*b^(1/2)*a*ln(b^(1/2)*cot(x)+(a+b*cot(x)^2)^(1/2))+(b^4*(a-b))^(1/2)/(a-b)*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(x)^2)^(1/2)*cot(x))-2*a/b*(b^4*(a-b))^(1/2)/(a-b)

)*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(x)^2)^(1/2)*cot(x))+a^2*(b^4*(a-b))^(1/2)/b^2/(a-b)*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(x)^2)^(1/2)*cot(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(105) = 210.

Time = 0.33 (sec) , antiderivative size = 1134, normalized size of antiderivative = 8.93

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \text{Too large to display}$$

[In] integrate(cot(x)^2*(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [1/16*(8*(a*b - b^2 - (a*b - b^2)*cos(2*x))*sqrt(-a + b)*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b)*sin(2*x) - (3*a^2 - 12*a*b + 8*b^2 - (3*a^2 - 12*a*b + 8*b^2)*cos(2*x)))*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1))*sin(2*x) + 2*(4*b^2*cos(2*x) - (5*a*b - 6*b^2)*cos(2*x)^2 + 5*a*b - 2*b^2)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/((b*cos(2*x) - b)*sin(2*x)), -1/8*((3*a^2 - 12*a*b + 8*b^2 - (3*a^2 - 12*a*b + 8*b^2)*cos(2*x))*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b))*sin(2*x) - 4*(a*b - b^2 - (a*b - b^2)*cos(2*x))*sqrt(-a + b)*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b)*sin(2*x) - (4*b^2*cos(2*x) - (5*a*b - 6*b^2)*cos(2*x)^2 + 5*a*b - 2*b^2)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/((b*cos(2*x) - b)*sin(2*x)), -1/16*(16*(a*b - b^2 - (a*b - b^2)*cos(2*x))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b))*sin(2*x) + (3*a^2 - 12*a*b + 8*b^2 - (3*a^2 - 12*a*b + 8*b^2)*cos(2*x))*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1))*sin(2*x) - 2*(4*b^2*cos(2*x) - (5*a*b - 6*b^2)*cos(2*x)^2 + 5*a*b - 2*b^2)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/((b*cos(2*x) - b)*sin(2*x)), -1/8*(8*(a*b - b^2 - (a*b - b^2)*cos(2*x))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b))*sin(2*x) + (3*a^2 - 12*a*b + 8*b^2 - (3*a^2 - 12*a*b + 8*b^2)*cos(2*x))*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b))*sin(2*x) - (4*b^2*cos(2*x) - (5*a*b - 6*b^2)*cos(2*x)^2 + 5*a*b - 2*b^2)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/((b*cos(2*x) - b)*sin(2*x))]

Sympy [F]

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \int (a + b \cot^2(x))^{\frac{3}{2}} \cot^2(x) dx$$

```
[In] integrate(cot(x)**2*(a+b*cot(x)**2)**(3/2),x)
```

```
[Out] Integral((a + b*cot(x)**2)**(3/2)*cot(x)**2, x)
```

Maxima [F]

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \int (b \cot(x)^2 + a)^{\frac{3}{2}} \cot(x)^2 dx$$

```
[In] integrate(cot(x)^2*(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cot(x)^2 + a)^(3/2)*cot(x)^2, x)
```

Giac [F(-2)]

Exception generated.

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(cot(x)^2*(a+b*cot(x)^2)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \cot^2(x) (a + b \cot^2(x))^{3/2} dx = \int \cot(x)^2 (b \cot(x)^2 + a)^{3/2} dx$$

```
[In] int(cot(x)^2*(a + b*cot(x)^2)^(3/2),x)
```

```
[Out] int(cot(x)^2*(a + b*cot(x)^2)^(3/2), x)
```

3.28 $\int \cot(x) (a + b \cot^2(x))^{3/2} dx$

Optimal result	191
Rubi [A] (verified)	191
Mathematica [A] (verified)	193
Maple [B] (verified)	193
Fricas [B] (verification not implemented)	194
Sympy [F]	194
Maxima [F(-2)]	195
Giac [F(-2)]	195
Mupad [B] (verification not implemented)	195

Optimal result

Integrand size = 15, antiderivative size = 69

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = (a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) - (a - b) \sqrt{a + b \cot^2(x)} - \frac{1}{3} (a + b \cot^2(x))^{3/2}$$

[Out] (a-b)^(3/2)*arctanh((a+b*cot(x)^2)^(1/2)/(a-b)^(1/2))-1/3*(a+b*cot(x)^2)^(3/2)-(a-b)*(a+b*cot(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 52, 65, 214}

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = (a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) - (a - b) \sqrt{a + b \cot^2(x)} - \frac{1}{3} (a + b \cot^2(x))^{3/2}$$

[In] Int[Cot[x]*(a + b*Cot[x]^2)^(3/2),x]

[Out] (a - b)^(3/2)*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]] - (a - b)*Sqrt[a + b*Cot[x]^2] - (a + b*Cot[x]^2)^(3/2)/3

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(

```
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x(a+bx^2)^{3/2}}{1+x^2} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{(a+bx)^{3/2}}{1+x} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{1}{3}(a+b\cot^2(x))^{3/2} - \frac{1}{2}(a-b)\text{Subst}\left(\int \frac{\sqrt{a+bx}}{1+x} dx, x, \cot^2(x)\right)
\end{aligned}$$

$$\begin{aligned}
&= -\left((a-b)\sqrt{a+b\cot^2(x)}\right) - \frac{1}{3}(a+b\cot^2(x))^{3/2} \\
&\quad - \frac{1}{2}(a-b)^2 \text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= -\left((a-b)\sqrt{a+b\cot^2(x)}\right) - \frac{1}{3}(a+b\cot^2(x))^{3/2} \\
&\quad - \frac{(a-b)^2 \text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&= (a-b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) - (a-b)\sqrt{a+b\cot^2(x)} - \frac{1}{3}(a+b\cot^2(x))^{3/2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 63, normalized size of antiderivative = 0.91

$$\begin{aligned}
\int \cot(x) (a+b\cot^2(x))^{3/2} dx &= (a-b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) \\
&\quad - \frac{1}{3}\sqrt{a+b\cot^2(x)}(4a-3b+b\cot^2(x))
\end{aligned}$$

[In] Integrate[Cot[x]*(a+b*Cot[x]^2)^(3/2),x]

[Out] (a-b)^(3/2)*ArcTanh[Sqrt[a+b*Cot[x]^2]/Sqrt[a-b]] - (Sqrt[a+b*Cot[x]^2]*(4*a-3*b+b*Cot[x]^2))/3

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 135 vs. 2(57) = 114.

Time = 0.02 (sec) , antiderivative size = 136, normalized size of antiderivative = 1.97

method	result
derivativedivides	$ -\frac{b\cot(x)^2\sqrt{a+b\cot(x)^2}}{3} - \frac{4a\sqrt{a+b\cot(x)^2}}{3} + b\sqrt{a+b\cot(x)^2} - \frac{b^2\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} + \frac{2ab\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} $
default	$ -\frac{b\cot(x)^2\sqrt{a+b\cot(x)^2}}{3} - \frac{4a\sqrt{a+b\cot(x)^2}}{3} + b\sqrt{a+b\cot(x)^2} - \frac{b^2\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} + \frac{2ab\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}} $

[In] int(cot(x)*(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/3*b*cot(x)^2*(a+b*cot(x)^2)^(1/2)-4/3*a*(a+b*cot(x)^2)^(1/2)+b*(a+b*cot(x)^2)^(1/2)-b^2/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))+2*a*

$b/(-a+b)^{(1/2)}*\arctan((a+b*\cot(x)^2)^{(1/2)/(-a+b)^{(1/2)})-a^2/(-a+b)^{(1/2)}*a$
 $rctan((a+b*\cot(x)^2)^{(1/2)/(-a+b)^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(57) = 114.

Time = 0.33 (sec) , antiderivative size = 330, normalized size of antiderivative = 4.78

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = \left[-\frac{3((a-b)\cos(2x) - a + b)\sqrt{a-b}\log\left(-2(a^2 - 2ab + b^2)\cos(2x)^2 - 2a^2 + b^2 + 2\right)}{\dots} \right]$$

[In] integrate(cot(x)*(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [-1/12*(3*((a - b)*cos(2*x) - a + b)*sqrt(a - b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 - 2*a^2 + b^2 + 2*((a - b)*cos(2*x)^2 - (2*a - b)*cos(2*x) + a)*sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)) + 4*(a^2 - a*b)*cos(2*x)) + 8*(2*(a - b)*cos(2*x) - 2*a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(cos(2*x) - 1), 1/6*(3*((a - b)*cos(2*x) - a + b)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))*(cos(2*x) - 1)/((a - b)*cos(2*x) - a)) - 4*(2*(a - b)*cos(2*x) - 2*a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(cos(2*x) - 1)]

Sympy [F]

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = \int (a + b \cot^2(x))^{\frac{3}{2}} \cot(x) dx$$

[In] integrate(cot(x)*(a+b*cot(x)**2)**(3/2),x)

[Out] Integral((a + b*cot(x)**2)**(3/2)*cot(x), x)

Maxima [F(-2)]

Exception generated.

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = \text{Exception raised: ValueError}$$

[In] `integrate(cot(x)*(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)

Giac [F(-2)]

Exception generated.

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] `integrate(cot(x)*(a+b*cot(x)^2)^(3/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Unable to convert to real sageVARb Error: Bad Argument ValueUnable to convert to real sageVARb Error: Bad Argument Val

Mupad [B] (verification not implemented)

Time = 16.99 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.01

$$\int \cot(x) (a + b \cot^2(x))^{3/2} dx = \operatorname{atanh}\left(\frac{(a-b)^{3/2} \sqrt{b \cot(x)^2 + a}}{a^2 - 2ab + b^2}\right) (a-b)^{3/2} - \frac{(b \cot(x)^2 + a)^{3/2}}{3} - (a-b) \sqrt{b \cot(x)^2 + a}$$

[In] `int(cot(x)*(a + b*cot(x)^2)^(3/2),x)`

[Out] `atanh(((a - b)^(3/2)*(a + b*cot(x)^2)^(1/2))/(a^2 - 2*a*b + b^2))*(a - b)^(3/2) - (a + b*cot(x)^2)^(3/2)/3 - (a - b)*(a + b*cot(x)^2)^(1/2)`

3.29 $\int (a + b \cot^2(x))^{3/2} \tan(x) dx$

Optimal result	196
Rubi [A] (verified)	196
Mathematica [A] (verified)	198
Maple [B] (verified)	198
Fricas [A] (verification not implemented)	199
Sympy [F]	201
Maxima [F]	201
Giac [F(-2)]	201
Mupad [B] (verification not implemented)	202

Optimal result

Integrand size = 15, antiderivative size = 75

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a}}\right) - (a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) - b\sqrt{a + b \cot^2(x)}$$

[Out] $a^{(3/2)}*\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/a^{(1/2)})-(a-b)^{(3/2)}*\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})-b*(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 457, 86, 162, 65, 214}

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a}}\right) - (a - b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right) - b\sqrt{a + b \cot^2(x)}$$

[In] $\operatorname{Int}[(a + b*\cot[x]^2)^{(3/2)}*\tan[x], x]$

[Out] $a^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\cot[x]^2]/\operatorname{Sqrt}[a]] - (a - b)^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\cot[x]^2]/\operatorname{Sqrt}[a - b]] - b*\operatorname{Sqrt}[a + b*\cot[x]^2]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 86

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Simp[f*(e + f*x)^(p - 1)/(b*d*(p - 1)), x] + Dist[1/(b*d), I
nt[(b*d*e^2 - a*c*f^2 + f*(2*b*d*e - b*c*f - a*d*f)*x)*((e + f*x)^(p - 2)/(
(a + b*x)*(c + d*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 1]
```

Rule 162

```
Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*
((c_.) + (d_.)*(x_))), x_Symbol] := Dist[(b*g - a*h)/(b*c - a*d), Int[(e +
f*x)^p/(a + b*x), x], x] - Dist[(d*g - c*h)/(b*c - a*d), Int[(e + f*x)^p/(c
+ d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[(((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \frac{(a + bx^2)^{3/2}}{x(1 + x^2)} dx, x, \cot(x)\right)$$

$$\begin{aligned}
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{(a+bx)^{3/2}}{x(1+x)} dx, x, \cot^2(x)\right)\right) \\
&= -b\sqrt{a+b\cot^2(x)} - \frac{1}{2}\text{Subst}\left(\int \frac{a^2+(2a-b)bx}{x(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= -b\sqrt{a+b\cot^2(x)} - \frac{1}{2}a^2\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&\quad + \frac{1}{2}(a-b)^2\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
&= -b\sqrt{a+b\cot^2(x)} - \frac{a^2\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&\quad + \frac{(a-b)^2\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
&= a^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a}}\right) - (a-b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) - b\sqrt{a+b\cot^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int (a+b\cot^2(x))^{3/2} \tan(x) dx &= a^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a}}\right) \\
&\quad - (a-b)^{3/2}\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right) - b\sqrt{a+b\cot^2(x)}
\end{aligned}$$

[In] Integrate[(a + b*Cot[x]^2)^(3/2)*Tan[x], x]

[Out] a^(3/2)*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a]] - (a - b)^(3/2)*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]] - b*Sqrt[a + b*Cot[x]^2]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 340 vs. 2(61) = 122.

Time = 1.28 (sec) , antiderivative size = 341, normalized size of antiderivative = 4.55

method	result
default	$\sqrt{4}(\cos(x)-1)(a+b\cot(x)^2)^{\frac{3}{2}} \left(a^{\frac{3}{2}}\sqrt{-a+b} \operatorname{arctanh} \left(\frac{\sqrt{-\frac{a\cos(x)^2-\cos(x)^2b-a}{(\cos(x)+1)^2}}(\cot(x)+\csc(x))}{\sqrt{a}} \right) \right) \sin(x)-\cos(x)\sqrt{-\frac{a\cos(x)^2-\cos(x)^2b-a}{(\cos(x)+1)^2}}$

[In] `int((a+b*cot(x)^2)^(3/2)*tan(x),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}4^{(1/2)} / (-a+b)^{(1/2)} * (\cos(x)-1) * (a+b*\cot(x)^2)^{(3/2)} * (a^{(3/2)} * (-a+b)^{(1/2)} * \operatorname{arctanh}(1/a^{(1/2)} * (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (\cot(x)+\csc(x))) * \sin(x) - \cos(x) * (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (-a+b)^{(1/2)} * b + \operatorname{arctan}(1/(-a+b)^{(1/2)} * (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (\cot(x)+\csc(x))) * a^2 * \sin(x) - 2 * \operatorname{arctan}(1/(-a+b)^{(1/2)} * (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (\cot(x)+\csc(x))) * a * b * \sin(x) + \operatorname{arctan}(1/(-a+b)^{(1/2)} * (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (\cot(x)+\csc(x))) * b^2 * \sin(x) - (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)} * (-a+b)^{(1/2)} * b) / (a*\cos(x)^2-\cos(x)^2*b-a) / (-(a*\cos(x)^2-\cos(x)^2*b-a) / (\cos(x)+1)^2)^{(1/2)}$

Fricas [A] (verification not implemented)

none

Time = 0.79 (sec) , antiderivative size = 565, normalized size of antiderivative = 7.53

$$\begin{aligned}
\int (a+b \cot^2(x))^{3/2} \tan(x) dx = & \left[\frac{1}{2} a^{3/2} \log \left(2 a \tan (x)^2 + 2 \sqrt{a} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2 + b \right) \right. \\
& - \frac{1}{4} (a-b)^{3/2} \log \left(-\frac{(8 a^2 - 8 a b + b^2) \tan (x)^4 + 2 (4 a b - 3 b^2) \tan (x)^2 + b^2 + 4 ((2 a - b) \tan (x)^4 + b \tan (x)^2)}{\tan (x)^4 + 2 \tan (x)^2 + 1} \right. \\
& - b \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}}, -\sqrt{-a} a \arctan \left(\frac{\sqrt{-a} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2}{a \tan (x)^2 + b} \right) \\
& - \frac{1}{4} (a-b)^{3/2} \log \left(-\frac{(8 a^2 - 8 a b + b^2) \tan (x)^4 + 2 (4 a b - 3 b^2) \tan (x)^2 + b^2 + 4 ((2 a - b) \tan (x)^4 + b \tan (x)^2)}{\tan (x)^4 + 2 \tan (x)^2 + 1} \right. \\
& - b \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}}, \frac{1}{2} (-a+b)^{3/2} \arctan \left(-\frac{2 \sqrt{-a+b} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2}{(2 a - b) \tan (x)^2 + b} \right) \\
& + \frac{1}{2} a^{3/2} \log \left(2 a \tan (x)^2 + 2 \sqrt{a} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2 + b \right) \\
& - b \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}}, -\sqrt{-a} a \arctan \left(\frac{\sqrt{-a} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2}{a \tan (x)^2 + b} \right) \\
& \left. + \frac{1}{2} (-a+b)^{3/2} \arctan \left(-\frac{2 \sqrt{-a+b} \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \tan (x)^2}{(2 a - b) \tan (x)^2 + b} \right) - b \sqrt{\frac{a \tan (x)^2 + b}{\tan (x)^2}} \right]
\end{aligned}$$

[In] integrate((a+b*cot(x)^2)^(3/2)*tan(x),x, algorithm="fricas")

[Out] [1/2*a^(3/2)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2))*tan(x)^2 + b) - 1/4*(a - b)^(3/2)*log(-((8*a^2 - 8*a*b + b^2)*tan(x)^4 + 2*(4*a*b - 3*b^2)*tan(x)^2 + b^2 + 4*((2*a - b)*tan(x)^4 + b*tan(x)^2)*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) - b*sqrt((a*tan(x)^2 + b)/tan(x)^2), -sqrt(-a)*a*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2/(a*tan(x)^2 + b)) - 1/4*(a - b)^(3/2)*log(-((8*a^2 - 8*a*b + b^2)*tan(x)^4 + 2*(4*a*b - 3*b^2)*tan(x)^2 + b^2 + 4*((2*a - b)*tan(x)^4 + b*tan(x)^2)*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) - b*sqrt((a*tan(x)^2 + b)/tan(x)^2), 1/2*(-a + b)^(3/2)*arctan(-2*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2/((2*a - b)*tan(x)^2 + b)) + 1/2*a^(3/2)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b) - b*sqrt((a*tan(x)^2 + b)/tan(x)^2), -sqrt(-a)*a*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2/(a*tan(x)^2 + b)) + 1/2*(-a + b)^(3/2)*arctan(-2*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2/((2*a - b)*tan(x)^2 + b)) - b*sqrt((a*tan(x)^2 + b)/tan(x)^2)]


```
)^2)*tan(x)^2/((2*a - b)*tan(x)^2 + b)) - b*sqrt((a*tan(x)^2 + b)/tan(x)^2)
]
```

Sympy [F]

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = \int (a + b \cot^2(x))^{\frac{3}{2}} \tan(x) dx$$

```
[In] integrate((a+b*cot(x)**2)**(3/2)*tan(x),x)
```

```
[Out] Integral((a + b*cot(x)**2)**(3/2)*tan(x), x)
```

Maxima [F]

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = \int (b \cot(x)^2 + a)^{\frac{3}{2}} \tan(x) dx$$

```
[In] integrate((a+b*cot(x)^2)^(3/2)*tan(x),x, algorithm="maxima")
```

```
[Out] integrate((b*cot(x)^2 + a)^(3/2)*tan(x), x)
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = \text{Exception raised: TypeError}$$

```
[In] integrate((a+b*cot(x)^2)^(3/2)*tan(x),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [B] (verification not implemented)

Time = 0.26 (sec) , antiderivative size = 506, normalized size of antiderivative = 6.75

$$\int (a + b \cot^2(x))^{3/2} \tan(x) dx = \operatorname{atanh} \left(\frac{2 b^6 \sqrt{a^3} \sqrt{a + \frac{b}{\tan(x)^2}}}{-6 a^5 b^3 + 12 a^4 b^4 - 8 a^3 b^5 + 2 a^2 b^6} \right. \\ \left. - \frac{8 a b^5 \sqrt{a^3} \sqrt{a + \frac{b}{\tan(x)^2}}}{-6 a^5 b^3 + 12 a^4 b^4 - 8 a^3 b^5 + 2 a^2 b^6} + \frac{12 a^2 b^4 \sqrt{a^3} \sqrt{a + \frac{b}{\tan(x)^2}}}{-6 a^5 b^3 + 12 a^4 b^4 - 8 a^3 b^5 + 2 a^2 b^6} \right. \\ \left. - \frac{6 a^3 b^3 \sqrt{a^3} \sqrt{a + \frac{b}{\tan(x)^2}}}{-6 a^5 b^3 + 12 a^4 b^4 - 8 a^3 b^5 + 2 a^2 b^6} \right) \sqrt{a^3} \\ - \operatorname{atanh} \left(\frac{2 a b^5 \sqrt{a + \frac{b}{\tan(x)^2}} \sqrt{a^3 - 3 a^2 b + 3 a b^2 - b^3}}{6 a^5 b^3 - 18 a^4 b^4 + 20 a^3 b^5 - 10 a^2 b^6 + 2 a b^7} \right. \\ \left. - \frac{6 a^2 b^4 \sqrt{a + \frac{b}{\tan(x)^2}} \sqrt{a^3 - 3 a^2 b + 3 a b^2 - b^3}}{6 a^5 b^3 - 18 a^4 b^4 + 20 a^3 b^5 - 10 a^2 b^6 + 2 a b^7} \right. \\ \left. + \frac{6 a^3 b^3 \sqrt{a + \frac{b}{\tan(x)^2}} \sqrt{a^3 - 3 a^2 b + 3 a b^2 - b^3}}{6 a^5 b^3 - 18 a^4 b^4 + 20 a^3 b^5 - 10 a^2 b^6 + 2 a b^7} \right) \sqrt{(a - b)^3} - b \sqrt{a + \frac{b}{\tan(x)^2}}$$

[In] `int(tan(x)*(a + b*cot(x)^2)^(3/2),x)`

[Out] `atanh((2*b^6*(a^3)^(1/2)*(a + b/tan(x)^2)^(1/2))/(2*a^2*b^6 - 8*a^3*b^5 + 12*a^4*b^4 - 6*a^5*b^3) - (8*a*b^5*(a^3)^(1/2)*(a + b/tan(x)^2)^(1/2))/(2*a^2*b^6 - 8*a^3*b^5 + 12*a^4*b^4 - 6*a^5*b^3) + (12*a^2*b^4*(a^3)^(1/2)*(a + b/tan(x)^2)^(1/2))/(2*a^2*b^6 - 8*a^3*b^5 + 12*a^4*b^4 - 6*a^5*b^3) - (6*a^3*b^3*(a^3)^(1/2)*(a + b/tan(x)^2)^(1/2))/(2*a^2*b^6 - 8*a^3*b^5 + 12*a^4*b^4 - 6*a^5*b^3))*(a^3)^(1/2) - atanh((2*a*b^5*(a + b/tan(x)^2)^(1/2)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)^(1/2))/(2*a*b^7 - 10*a^2*b^6 + 20*a^3*b^5 - 18*a^4*b^4 + 6*a^5*b^3) - (6*a^2*b^4*(a + b/tan(x)^2)^(1/2)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)^(1/2))/(2*a*b^7 - 10*a^2*b^6 + 20*a^3*b^5 - 18*a^4*b^4 + 6*a^5*b^3) + (6*a^3*b^3*(a + b/tan(x)^2)^(1/2)*(3*a*b^2 - 3*a^2*b + a^3 - b^3)^(1/2))/(2*a*b^7 - 10*a^2*b^6 + 20*a^3*b^5 - 18*a^4*b^4 + 6*a^5*b^3))*((a - b)^3)^(1/2) - b*(a + b/tan(x)^2)^(1/2)`

3.30 $\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx$

Optimal result	203
Rubi [A] (verified)	203
Mathematica [B] (verified)	205
Maple [B] (warning: unable to verify)	206
Fricas [A] (verification not implemented)	206
Sympy [F]	208
Maxima [F]	208
Giac [B] (verification not implemented)	208
Mupad [F(-1)]	209

Optimal result

Integrand size = 17, antiderivative size = 80

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx = (a - b)^{3/2} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) + a \sqrt{a + b \cot^2(x)} \tan(x)$$

[Out] (a-b)^(3/2)*arctan(cot(x)*(a-b)^(1/2)/(a+b*cot(x)^2)^(1/2))-b^(3/2)*arctanh(cot(x)*b^(1/2)/(a+b*cot(x)^2)^(1/2))+a*(a+b*cot(x)^2)^(1/2)*tan(x)

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3751, 485, 537, 223, 212, 385, 209}

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx = (a - b)^{3/2} \arctan\left(\frac{\sqrt{a - b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) - b^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a + b \cot^2(x)}}\right) + a \tan(x) \sqrt{a + b \cot^2(x)}$$

[In] Int[(a + b*Cot[x]^2)^(3/2)*Tan[x]^2,x]

[Out] (a - b)^(3/2)*ArcTan[(Sqrt[a - b]*Cot[x])/Sqrt[a + b*Cot[x]^2]] - b^(3/2)*ArcTanh[(Sqrt[b]*Cot[x])/Sqrt[a + b*Cot[x]^2]] + a*Sqrt[a + b*Cot[x]^2]*Tan[x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 485

Int[((e_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[c*(e*x)^(m+1)*(a + b*x^n)^(p+1)*((c + d*x^n)^(q-1)/(a*e*(m+1))), x] - Dist[1/(a*e^n*(m+1)), Int[(e*x)^(m+n)*(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(c*b - a*d)*(m+1) + c*n*(b*c*(p+1) + a*d*(q-1)) + d*((c*b - a*d)*(m+1) + c*b*n*(p+q))*x^n, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && GtQ[q, 1] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 537

Int[((e_) + (f_.)*(x_)^(n_))/(((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)]), x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{(a+bx^2)^{3/2}}{x^2(1+x^2)} dx, x, \cot(x)\right) \\
&= a\sqrt{a+b\cot^2(x)}\tan(x) - \text{Subst}\left(\int \frac{-a(a-2b)+b^2x^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= a\sqrt{a+b\cot^2(x)}\tan(x) + (a-b)^2\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&\quad - b^2\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= a\sqrt{a+b\cot^2(x)}\tan(x) + (a-b)^2\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&\quad - b^2\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&= (a-b)^{3/2}\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&\quad - b^{3/2}\text{arctanh}\left(\frac{\sqrt{b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) + a\sqrt{a+b\cot^2(x)}\tan(x)
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 222 vs. 2(80) = 160.

Time = 0.82 (sec) , antiderivative size = 222, normalized size of antiderivative = 2.78

$$\int (a + b\cot^2(x))^{3/2} \tan^2(x) dx = \frac{\sqrt{-((-a-b+(a-b)\cos(2x))\csc^2(x))}(-\sqrt{2}(a-b)^2\sqrt{-b}\text{arctanh}\left(\frac{\sqrt{2}\sqrt{a-b}}{\sqrt{-a-b+(a-b)\cos(2x)}}\right) + \sqrt{a-b}(\sqrt{2}\sqrt{b}\text{arctanh}\left(\frac{\sqrt{2}\sqrt{-b}\cos(x)}{\sqrt{-a-b+(a-b)\cos(2x)}}\right) + a\sqrt{-b}\sqrt{-a-b+(a-b)\cos(2x)}\sec(x))\sin(x)}{\sqrt{2}\sqrt{a-b}}$$

[In] Integrate[(a + b*Cot[x]^2)^(3/2)*Tan[x]^2, x]

[Out] (Sqrt[-((-a - b + (a - b)*Cos[2*x])*Csc[x]^2)]*(-(Sqrt[2]*(a - b)^2*Sqrt[-b]*ArcTanh[(Sqrt[2]*Sqrt[a - b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]])] + Sqrt[a - b]*(Sqrt[2]*b^2*ArcTanh[(Sqrt[2]*Sqrt[-b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]])] + a*Sqrt[-b]*Sqrt[-a - b + (a - b)*Cos[2*x]]*Sec[x]))*Sin[x]/(Sqrt[2]*Sqrt[a - b]*Sqrt[-b]*Sqrt[-a - b + (a - b)*Cos[2*x]])

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 641 vs. 2(66) = 132.

Time = 0.82 (sec) , antiderivative size = 642, normalized size of antiderivative = 8.02

method	result
default	$\sqrt{4} \left(2 \cos(x) b^{\frac{7}{2}} \ln \left(4 \cos(x) \sqrt{-a+b} \sqrt{\frac{-a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} - 4 \cos(x) a + 4 b \cos(x) + 4 \sqrt{-a+b} \sqrt{\frac{-a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} \right) - 4 \cos(x) b^{\frac{5}{2}} \right)$

[In] `int((a+b*cot(x)^2)^(3/2)*tan(x)^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4} 4^{1/2} / b^{3/2} / (-a+b)^{1/2} * (2 \cos(x) * b^{7/2} * \ln(4 \cos(x) * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} - 4 \cos(x) * a + 4 * b \cos(x) + 4 * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} - 4 \cos(x) * b^{5/2} * \ln(4 \cos(x) * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} - 4 \cos(x) * a + 4 * b \cos(x) + 4 * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * a + 2 \cos(x) * b^{3/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * (-a+b)^{1/2} * a + 2 \cos(x) * b^{3/2} * \ln(4 \cos(x) * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} - 4 \cos(x) * a + 4 * b \cos(x) + 4 * (-a+b)^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * a^2 + 2 * b^{3/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * (-a+b)^{1/2} * a - \cos(x) * \ln(-4 * (b^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * \cos(x) - \cos(x) * a + b \cos(x) + (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * b^{1/2} + a) / (\cos(x)-1) * (-a+b)^{1/2} * b^3 + \cos(x) * \ln(2 / b^{1/2} * (b^{1/2} * (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * \cos(x) + (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * b^{1/2} + \cos(x) * a - b \cos(x) + a) / (\cos(x)+1) * (-a+b)^{1/2} * b^3 * (\cos(x)-1) * (a+b \cot(x)^2)^{3/2} / (a \cos(x)^2 - \cos(x)^2 * b - a) / (-a \cos(x)^2 - \cos(x)^2 * b - a) / (\cos(x)+1)^2)^{1/2} * \tan(x)$

Fricas [A] (verification not implemented)

none

$\frac{a \tan^2(x) + 2b}{\tan^2(x)} + a \sqrt{\frac{a \tan^2(x) + b}{\tan^2(x)}} \tan(x)$
 $, \frac{1}{2} (a - b)^{3/2} \arctan\left(\frac{2 \sqrt{a - b} \sqrt{a \tan^2(x) + b}}{\tan^2(x) + a + 2b}\right) + \sqrt{-b} b \arctan\left(\frac{\sqrt{-b} \sqrt{a \tan^2(x) + b}}{\tan^2(x) + a + 2b}\right) \tan(x) / b + a \sqrt{\frac{a \tan^2(x) + b}{\tan^2(x)}} \tan(x)$

Sympy [F]

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx = \int (a + b \cot^2(x))^{3/2} \tan^2(x) dx$$

[In] integrate((a+b*cot(x)**2)**(3/2)*tan(x)**2,x)

[Out] Integral((a + b*cot(x)**2)**(3/2)*tan(x)**2, x)

Maxima [F]

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx = \int (b \cot^2(x) + a)^{3/2} \tan^2(x) dx$$

[In] integrate((a+b*cot(x)^2)^(3/2)*tan(x)^2,x, algorithm="maxima")

[Out] integrate((b*cot(x)^2 + a)^(3/2)*tan(x)^2, x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 625 vs. 2(66) = 132.

Time = 18.70 (sec) , antiderivative size = 625, normalized size of antiderivative = 7.81

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx =$$

$$-\frac{1}{2} \left(\frac{2 \sqrt{-a + b} b^2 \arctan\left(\frac{(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos^2(x) + b \cos^2(x) + a})^2 + a - 2b}{2 \sqrt{ab - b^2}}\right)}{\sqrt{ab - b^2}} - (a - b) \sqrt{-a + b} \log\left(\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos^2(x) + b \cos^2(x) + a}\right)\right) \right)$$

$$\frac{(2 a \sqrt{-a + b} b^2 \arctan\left(\frac{\sqrt{-a + b} \sqrt{b}}{\sqrt{ab - b^2}}\right) - 2 a b^{5/2} \arctan\left(\frac{\sqrt{-a + b} \sqrt{b}}{\sqrt{ab - b^2}}\right) - 2 \sqrt{-a + b} b^3 \arctan\left(\frac{\sqrt{-a + b} \sqrt{b}}{\sqrt{ab - b^2}}\right) + 2 b^{7/2} \arctan\left(\frac{\sqrt{-a + b} \sqrt{b}}{\sqrt{ab - b^2}}\right))}{\sqrt{ab - b^2}}$$

[In] integrate((a+b*cot(x)^2)^(3/2)*tan(x)^2,x, algorithm="giac")


```
[Out] -1/2*(2*sqrt(-a + b)*b^2*arctan(1/2*((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 + a - 2*b)/sqrt(a*b - b^2))/sqrt(a*b - b^2) - (a - b)*sqrt(-a + b)*log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2) + 4*a^2*sqrt(-a + b)/((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a)*sgn(sin(x)) - 1/2*(2*a*sqrt(-a + b)*b^2*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) - 2*a*b^(5/2)*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) - 2*sqrt(-a + b)*b^3*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) + 2*b^(7/2)*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) + sqrt(a*b - b^2)*a^2*sqrt(-a + b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - sqrt(a*b - b^2)*a^2*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - 2*sqrt(a*b - b^2)*a*sqrt(-a + b)*b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 2*sqrt(a*b - b^2)*a*b^(3/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + sqrt(a*b - b^2)*sqrt(-a + b)*b^2*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - sqrt(a*b - b^2)*b^(5/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 2*sqrt(a*b - b^2)*a^2*sqrt(-a + b)*sgn(sin(x))/(sqrt(a*b - b^2)*a + sqrt(a*b - b^2)*sqrt(-a + b)*sqrt(b) - sqrt(a*b - b^2)*b)
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \cot^2(x))^{3/2} \tan^2(x) dx = \int \tan(x)^2 (b \cot(x)^2 + a)^{3/2} dx$$

```
[In] int(tan(x)^2*(a + b*cot(x)^2)^(3/2),x)
```

```
[Out] int(tan(x)^2*(a + b*cot(x)^2)^(3/2), x)
```

3.31 $\int (a + b \cot^2(c + dx))^{5/2} dx$

Optimal result	210
Rubi [A] (verified)	210
Mathematica [A] (verified)	213
Maple [B] (verified)	213
Fricas [B] (verification not implemented)	214
Sympy [F]	215
Maxima [F]	215
Giac [F(-2)]	215
Mupad [F(-1)]	216

Optimal result

Integrand size = 16, antiderivative size = 171

$$\int (a + b \cot^2(c + dx))^{5/2} dx = -\frac{(a - b)^{5/2} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b}(15a^2 - 20ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{8d} - \frac{(7a - 4b)b \cot(c + dx) \sqrt{a + b \cot^2(c + dx)}}{8d} - \frac{b \cot(c + dx) (a + b \cot^2(c + dx))^{3/2}}{4d}$$

[Out] $-(a-b)^{(5/2)} * \arctan(\cot(d*x+c) * (a-b)^{(1/2)} / (a+b*\cot(d*x+c)^2)^{(1/2)}) / d - 1/4 * b * \cot(d*x+c) * (a+b*\cot(d*x+c)^2)^{(3/2)} / d - 1/8 * (15*a^2 - 20*a*b + 8*b^2) * \operatorname{arctanh}(\cot(d*x+c) * b^{(1/2)} / (a+b*\cot(d*x+c)^2)^{(1/2)}) * b^{(1/2)} / d - 1/8 * (7*a - 4*b) * b * \cot(d*x+c) * (a+b*\cot(d*x+c)^2)^{(1/2)} / d$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 427, 542, 537, 223, 212, 385, 209}

$$\int (a + b \cot^2(c + dx))^{5/2} dx = -\frac{\sqrt{b}(15a^2 - 20ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{8d} - \frac{(a - b)^{5/2} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{b \cot(c + dx) (a + b \cot^2(c + dx))^{3/2}}{4d} - \frac{b(7a - 4b) \cot(c + dx) \sqrt{a + b \cot^2(c + dx)}}{8d}$$

[In] Int[(a + b*Cot[c + d*x]^2)^(5/2), x]

[Out] -(((a - b)^(5/2)*ArcTan[(Sqrt[a - b]*Cot[c + d*x])/Sqrt[a + b*Cot[c + d*x]^2]])/d - (Sqrt[b]*(15*a^2 - 20*a*b + 8*b^2)*ArcTanh[(Sqrt[b]*Cot[c + d*x])/Sqrt[a + b*Cot[c + d*x]^2]])/(8*d) - ((7*a - 4*b)*b*Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2])/(8*d) - (b*Cot[c + d*x]*(a + b*Cot[c + d*x]^2)^(3/2))/(4*d)

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 427

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*x*(a + b*x^n)^(p+1)*((c + d*x^n)^(q-1)/(b*(n*(p+q) + 1))), x] + Dist[1/(b*(n*(p+q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 537

Int[((e_) + (f_)*(x_)^(n_))/((a_) + (b_)*(x_)^(n_))*Sqrt[(c_) + (d_)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 542

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (
f_)*(x_)^(n_)), x_Symbol] := Simp[f*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^q/(
b*(n*(p + q + 1) + 1)), x] + Dist[1/(b*(n*(p + q + 1) + 1)), Int[(a + b*x^
n)^p*(c + d*x^n)^(q - 1)*Simp[c*(b*e - a*f + b*e*n*(p + q + 1)) + (d*(b*e -
a*f) + f*n*q*(b*c - a*d) + b*d*e*n*(p + q + 1))*x^n, x], x] /; FreeQ[{
a, b, c, d, e, f, n, p}, x] && GtQ[q, 0] && NeQ[n*(p + q + 1) + 1, 0]
```

Rule 3742

```
Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= - \frac{\text{Subst}\left(\int \frac{(a+bx^2)^{5/2}}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\
&= - \frac{b \cot(c+dx) (a+b \cot^2(c+dx))^{3/2}}{4d} - \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx^2}(a(4a-b)+(7a-4b)bx^2)}{1+x^2} dx, x, \cot(c+dx)\right)}{4d} \\
&= - \frac{(7a-4b)b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{8d} - \frac{b \cot(c+dx) (a+b \cot^2(c+dx))^{3/2}}{4d} \\
&\quad - \frac{\text{Subst}\left(\int \frac{a(8a^2-9ab+4b^2)+b(15a^2-20ab+8b^2)x^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{8d} \\
&= - \frac{(7a-4b)b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{8d} - \frac{b \cot(c+dx) (a+b \cot^2(c+dx))^{3/2}}{4d} \\
&\quad - \frac{(a-b)^3 \text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{d} \\
&\quad - \frac{(b(15a^2-20ab+8b^2)) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{8d} \\
&= - \frac{(7a-4b)b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{8d} - \frac{b \cot(c+dx) (a+b \cot^2(c+dx))^{3/2}}{4d} \\
&\quad - \frac{(a-b)^3 \text{Subst}\left(\int \frac{1}{1-(a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} \\
&\quad - \frac{(b(15a^2-20ab+8b^2)) \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{8d}
\end{aligned}$$

$$= -\frac{(a-b)^{5/2} \arctan\left(\frac{\sqrt{a-b}\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b}(15a^2 - 20ab + 8b^2) \operatorname{arctanh}\left(\frac{\sqrt{b}\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{8d} - \frac{(7a-4b)b\cot(c+dx)\sqrt{a+b\cot^2(c+dx)}}{8d} - \frac{b\cot(c+dx)(a+b\cot^2(c+dx))^{3/2}}{4d}$$

Mathematica [A] (verified)

Time = 0.88 (sec) , antiderivative size = 169, normalized size of antiderivative = 0.99

$$\int (a + b\cot^2(c + dx))^{5/2} dx = \frac{8(a-b)^{5/2} \arctan\left(\frac{\sqrt{b} + \sqrt{b}\cot^2(c+dx) - \cot(c+dx)\sqrt{a+b\cot^2(c+dx)}}{\sqrt{a-b}}\right) - b\cot(c+dx)\sqrt{a+b\cot^2(c+dx)}}{8d}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(5/2), x]

[Out] (8*(a - b)^(5/2)*ArcTan[(Sqrt[b] + Sqrt[b]*Cot[c + d*x]^2 - Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2])/Sqrt[a - b]] - b*Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2]*(9*a - 4*b + 2*b*Cot[c + d*x]^2) + Sqrt[b]*(15*a^2 - 20*a*b + 8*b^2)*Log[-(Sqrt[b]*Cot[c + d*x]) + Sqrt[a + b*Cot[c + d*x]^2]])/(8*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 461 vs. 2(149) = 298.

Time = 0.18 (sec) , antiderivative size = 462, normalized size of antiderivative = 2.70

method	result
derivativedivides	$-\frac{b^{\frac{5}{2}} \ln\left(\sqrt{b}\cot(dx+c) + \sqrt{a+b\cot(dx+c)^2}\right)}{d} - \frac{b^2 \cot(dx+c)^3 \sqrt{a+b\cot(dx+c)^2}}{4d} - \frac{9ba \cot(dx+c) \sqrt{a+b\cot(dx+c)^2}}{8d}$
default	$-\frac{b^{\frac{5}{2}} \ln\left(\sqrt{b}\cot(dx+c) + \sqrt{a+b\cot(dx+c)^2}\right)}{d} - \frac{b^2 \cot(dx+c)^3 \sqrt{a+b\cot(dx+c)^2}}{4d} - \frac{9ba \cot(dx+c) \sqrt{a+b\cot(dx+c)^2}}{8d}$

[In] int((a+b*cot(d*x+c)^2)^(5/2), x, method=_RETURNVERBOSE)

[Out]
$$-1/d*b^{5/2}*\ln(b^{1/2}*\cot(d*x+c)+(a+b*\cot(d*x+c)^2)^{1/2})-1/4/d*b^2*\cot(d*x+c)^3*(a+b*\cot(d*x+c)^2)^{1/2}-9/8/d*b*a*\cot(d*x+c)*(a+b*\cot(d*x+c)^2)^{1/2}-15/8/d*b^{1/2}*a^2*\ln(b^{1/2}*\cot(d*x+c)+(a+b*\cot(d*x+c)^2)^{1/2})+1/2/d*b^2*\cot(d*x+c)*(a+b*\cot(d*x+c)^2)^{1/2}+5/2/d*b^{3/2}*a*\ln(b^{1/2}*\cot(d*x+c)+(a+b*\cot(d*x+c)^2)^{1/2})+1/d*b*(b^4*(a-b))^{1/2}/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{1/2}/(a+b*\cot(d*x+c)^2)^{1/2}*\cot(d*x+c))-3/d*a*(b^4*(a-b))^{1/2}/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{1/2}/(a+b*\cot(d*x+c)^2)^{1/2}*\cot(d*x+c))+3/d*a^2/b*(b^4*(a-b))^{1/2}/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{1/2}/(a+b*\cot(d*x+c)^2)^{1/2}*\cot(d*x+c))$$

$1/2)/(a+b*\cot(d*x+c)^2)^{(1/2)*\cot(d*x+c)}-1/d*a^3*(b^4*(a-b))^{(1/2)}/b^2/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{(1/2)})/(a+b*\cot(d*x+c)^2)^{(1/2)*\cot(d*x+c)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 373 vs. 2(149) = 298.

Time = 0.32 (sec) , antiderivative size = 1520, normalized size of antiderivative = 8.89

$$\int (a + b \cot^2(c + dx))^{5/2} dx = \text{Too large to display}$$

[In] integrate((a+b*cot(d*x+c)^2)^(5/2),x, algorithm="fricas")

[Out] [-1/16*(8*(a^2 - 2*a*b + b^2 - (a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(-a + b)*log(-(a - b)*cos(2*d*x + 2*c) + sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + b)*sin(2*d*x + 2*c) + (15*a^2 - 20*a*b + 8*b^2 - (15*a^2 - 20*a*b + 8*b^2)*cos(2*d*x + 2*c))*sqrt(b)*log(((a - 2*b)*cos(2*d*x + 2*c) + 2*sqrt(b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - a - 2*b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - 2*(4*b^2*cos(2*d*x + 2*c) - 3*(3*a*b - 2*b^2)*cos(2*d*x + 2*c)^2 + 9*a*b - 2*b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/((d*cos(2*d*x + 2*c) - d)*sin(2*d*x + 2*c)), 1/16*(16*(a^2 - 2*a*b + b^2 - (a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) + a - b))*sin(2*d*x + 2*c) - (15*a^2 - 20*a*b + 8*b^2 - (15*a^2 - 20*a*b + 8*b^2)*cos(2*d*x + 2*c))*sqrt(b)*log(((a - 2*b)*cos(2*d*x + 2*c) + 2*sqrt(b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - a - 2*b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + 2*(4*b^2*cos(2*d*x + 2*c) - 3*(3*a*b - 2*b^2)*cos(2*d*x + 2*c)^2 + 9*a*b - 2*b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/((d*cos(2*d*x + 2*c) - d)*sin(2*d*x + 2*c)), -1/8*((15*a^2 - 20*a*b + 8*b^2 - (15*a^2 - 20*a*b + 8*b^2)*cos(2*d*x + 2*c))*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(b*cos(2*d*x + 2*c) + b))*sin(2*d*x + 2*c) + 4*(a^2 - 2*a*b + b^2 - (a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(-a + b)*log(-(a - b)*cos(2*d*x + 2*c) + sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + b)*sin(2*d*x + 2*c) - (4*b^2*cos(2*d*x + 2*c) - 3*(3*a*b - 2*b^2)*cos(2*d*x + 2*c)^2 + 9*a*b - 2*b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/((d*cos(2*d*x + 2*c) - d)*sin(2*d*x + 2*c)), 1/8*(8*(a^2 - 2*a*b + b^2 - (a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) + a - b))*sin(2*d*x + 2*c) - (15*a^2 - 20*a*b + 8*b^2 - (15*a^2 - 20*a*b + 8*b^2)*cos(2*d*x + 2*c))*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) -

```
1))*sin(2*d*x + 2*c)/(b*cos(2*d*x + 2*c) + b))*sin(2*d*x + 2*c) + (4*b^2*cos(2*d*x + 2*c) - 3*(3*a*b - 2*b^2)*cos(2*d*x + 2*c)^2 + 9*a*b - 2*b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/((d*cos(2*d*x + 2*c) - d)*sin(2*d*x + 2*c))]
```

Sympy [F]

$$\int (a + b \cot^2(c + dx))^{5/2} dx = \int (a + b \cot^2(c + dx))^{\frac{5}{2}} dx$$

```
[In] integrate((a+b*cot(d*x+c)**2)**(5/2),x)
```

```
[Out] Integral((a + b*cot(c + d*x)**2)**(5/2), x)
```

Maxima [F]

$$\int (a + b \cot^2(c + dx))^{5/2} dx = \int (b \cot(dx + c)^2 + a)^{\frac{5}{2}} dx$$

```
[In] integrate((a+b*cot(d*x+c)^2)^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cot(d*x + c)^2 + a)^(5/2), x)
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \cot^2(c + dx))^{5/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate((a+b*cot(d*x+c)^2)^(5/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \cot^2(c + dx))^{5/2} dx = \int (b \cot(c + dx)^2 + a)^{5/2} dx$$

```
[In] int((a + b*cot(c + d*x)^2)^(5/2),x)
```

```
[Out] int((a + b*cot(c + d*x)^2)^(5/2), x)
```


3.32 $\int (a + b \cot^2(c + dx))^{3/2} dx$

Optimal result	217
Rubi [A] (verified)	217
Mathematica [A] (verified)	219
Maple [B] (verified)	220
Fricas [B] (verification not implemented)	220
Sympy [F]	221
Maxima [F]	221
Giac [F(-2)]	221
Mupad [F(-1)]	222

Optimal result

Integrand size = 16, antiderivative size = 126

$$\int (a + b \cot^2(c + dx))^{3/2} dx = -\frac{(a - b)^{3/2} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{(3a - 2b)\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{2d} - \frac{b \cot(c + dx) \sqrt{a + b \cot^2(c + dx)}}{2d}$$

[Out] $-(a-b)^{(3/2)}*\arctan(\cot(d*x+c)*(a-b)^{(1/2)/(a+b*\cot(d*x+c)^2)^{(1/2))}/d-1/2*(3*a-2*b)*\operatorname{arctanh}(\cot(d*x+c)*b^{(1/2)/(a+b*\cot(d*x+c)^2)^{(1/2))}*b^{(1/2)}/d-1/2*b*\cot(d*x+c)*(a+b*\cot(d*x+c)^2)^{(1/2)}/d$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3742, 427, 537, 223, 212, 385, 209}

$$\int (a + b \cot^2(c + dx))^{3/2} dx = -\frac{(a - b)^{3/2} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b}(3a - 2b)\operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{2d} - \frac{b \cot(c + dx) \sqrt{a + b \cot^2(c + dx)}}{2d}$$

[In] $\text{Int}[(a + b*\text{Cot}[c + d*x]^2)^{(3/2)}, x]$

[Out] $-(((a - b)^{(3/2)}*\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[c + d*x])/(\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2])])/d - (((3*a - 2*b)*\text{Sqrt}[b]*\text{ArcTanh}[(\text{Sqrt}[b]*\text{Cot}[c + d*x])/(\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2])])/(2*d) - (b*\text{Cot}[c + d*x]*\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2])/(2*d)$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 427

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*x*(a + b*x^n)^(p+1)*((c + d*x^n)^(q-1)/(b*(n*(p+q) + 1))), x] + Dist[1/(b*(n*(p+q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q) + 1) - a*d) + d*(b*c*(n*(p+2*q-1) + 1) - a*d*(n*(q-1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 537

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
```

qq[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{(a+bx^2)^{3/2}}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\
 &= -\frac{b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{2d} - \frac{\text{Subst}\left(\int \frac{a(2a-b)+(3a-2b)bx^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{2d} \\
 &= -\frac{b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{2d} \\
 &\quad - \frac{(a-b)^2 \text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{d} \\
 &\quad - \frac{((3a-2b)b) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{2d} \\
 &= -\frac{b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{2d} \\
 &\quad - \frac{(a-b)^2 \text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} \\
 &\quad - \frac{((3a-2b)b) \text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{2d} \\
 &= -\frac{(a-b)^{3/2} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{(3a-2b)\sqrt{b} \text{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{2d} \\
 &\quad - \frac{b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.13

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \frac{2(a-b)^{3/2} \arctan\left(\frac{\sqrt{b} + \sqrt{b} \cot^2(c+dx) - \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{\sqrt{a-b}}\right) - b \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{2d}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(3/2), x]

[Out] (2*(a - b)^(3/2)*ArcTan[(Sqrt[b] + Sqrt[b]*Cot[c + d*x]^2 - Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2])/Sqrt[a - b]] - b*Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2] + (3*a - 2*b)*Sqrt[b]*Log[-(Sqrt[b]*Cot[c + d*x]) + Sqrt[a + b*Cot[c + d*x]^2]])/(2*d)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. 2(108) = 216.

Time = 0.04 (sec) , antiderivative size = 298, normalized size of antiderivative = 2.37

method	result
derivativedivides	$\frac{b^{\frac{3}{2}} \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{d} - \frac{b \cot(dx+c) \sqrt{a+b \cot(dx+c)^2}}{2d} - \frac{3\sqrt{b} a \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{2d}$
default	$\frac{b^{\frac{3}{2}} \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{d} - \frac{b \cot(dx+c) \sqrt{a+b \cot(dx+c)^2}}{2d} - \frac{3\sqrt{b} a \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{2d}$

[In] `int((a+b*cot(d*x+c)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{d} b^{3/2} \ln(b^{1/2} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}) - \frac{1}{2} b \cot(dx+c) \sqrt{a+b \cot(dx+c)^2} - \frac{3\sqrt{b} a \ln(b^{1/2} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2})}{2d}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 258 vs. 2(108) = 216.

Time = 0.32 (sec) , antiderivative size = 1071, normalized size of antiderivative = 8.50

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \text{Too large to display}$$

[In] `integrate((a+b*cot(d*x+c)^2)^(3/2),x, algorithm="fricas")`

[Out] $[-1/4 * (2 * (a - b) * \sqrt{-a + b} * \log(-(a - b) * \cos(2 * d * x + 2 * c) - \sqrt{-a + b}) * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1)}) * \sin(2 * d * x + 2 * c) + b) * \sin(2 * d * x + 2 * c) + (3 * a - 2 * b) * \sqrt{b} * \log(((a - 2 * b) * \cos(2 * d * x + 2 * c) - 2 * \sqrt{b}) * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1)}) * \sin(2 * d * x + 2 * c) - a - 2 * b) / (\cos(2 * d * x + 2 * c) - 1) * \sin(2 * d * x + 2 * c) + 2 * (b * \cos(2 * d * x + 2 * c) + b) * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1))} / (d * \sin(2 * d * x + 2 * c)), 1/2 * ((3 * a - 2 * b) * \sqrt{-b}) * \arctan(\sqrt{-b} * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1)}) * \sin(2 * d * x + 2 * c) / (b * \cos(2 * d * x + 2 * c) + b) * \sin(2 * d * x + 2 * c) - (a - b) * \sqrt{-a + b} * \log(-(a - b) * \cos(2 * d * x + 2 * c) - \sqrt{-a + b}) * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1)}) * \sin(2 * d * x + 2 * c) + b) * \sin(2 * d * x + 2 * c) - (b * \cos(2 * d * x + 2 * c) + b) * \sqrt{((a - b) * \cos(2 * d * x + 2 * c) - a - b) / (\cos(2 * d * x + 2 * c) - 1))} / (d * \sin(2 * d * x + 2 * c)), -1/4 * (4 * (a - b)^(3/2) * \arctan(-s$

```

qrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*
sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) + a - b))*sin(2*d*x + 2*c) + (3*
a - 2*b)*sqrt(b)*log(((a - 2*b)*cos(2*d*x + 2*c) - 2*sqrt(b)*sqrt(((a - b)*
cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - a - 2*
b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + 2*(b*cos(2*d*x + 2*c) + b)*sq
rt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/(d*sin(2*d*x
+ 2*c)), -1/2*(2*(a - b)^(3/2)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x
+ 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*
x + 2*c) + a - b))*sin(2*d*x + 2*c) - (3*a - 2*b)*sqrt(-b)*arctan(sqrt(-b)*
sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x +
2*c)/(b*cos(2*d*x + 2*c) + b))*sin(2*d*x + 2*c) + (b*cos(2*d*x + 2*c) + b)
*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1)))/(d*sin(2*
d*x + 2*c))]

```

Sympy [F]

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \int (a + b \cot^2(c + dx))^{\frac{3}{2}} dx$$

```
[In] integrate((a+b*cot(d*x+c)**2)**(3/2),x)
```

```
[Out] Integral((a + b*cot(c + d*x)**2)**(3/2), x)
```

Maxima [F]

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \int (b \cot(dx + c)^2 + a)^{\frac{3}{2}} dx$$

```
[In] integrate((a+b*cot(d*x+c)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*cot(d*x + c)^2 + a)^(3/2), x)
```

Giac [F(-2)]

Exception generated.

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate((a+b*cot(d*x+c)^2)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Error: Bad Argument Type
```

Mupad [F(-1)]

Timed out.

$$\int (a + b \cot^2(c + dx))^{3/2} dx = \int (b \cot(c + dx)^2 + a)^{3/2} dx$$

```
[In] int((a + b*cot(c + d*x)^2)^(3/2),x)
```

```
[Out] int((a + b*cot(c + d*x)^2)^(3/2), x)
```

3.33 $\int \sqrt{a + b \cot^2(c + dx)} dx$

Optimal result	223
Rubi [A] (verified)	223
Mathematica [A] (verified)	225
Maple [B] (verified)	225
Fricas [B] (verification not implemented)	226
Sympy [F]	227
Maxima [F(-2)]	227
Giac [F(-2)]	227
Mupad [F(-1)]	227

Optimal result

Integrand size = 16, antiderivative size = 87

$$\int \sqrt{a + b \cot^2(c + dx)} dx = -\frac{\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d}$$

[Out] $-\arctan(\cot(d*x+c)*(a-b)^{(1/2)/(a+b*\cot(d*x+c)^2)^{(1/2))}*(a-b)^{(1/2)/d}-\operatorname{arctanh}(\cot(d*x+c)*b^{(1/2)/(a+b*\cot(d*x+c)^2)^{(1/2))}*b^{(1/2)/d}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3742, 399, 223, 212, 385, 209}

$$\int \sqrt{a + b \cot^2(c + dx)} dx = -\frac{\sqrt{a-b} \arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b} \operatorname{arctanh}\left(\frac{\sqrt{b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d}$$

[In] `Int[Sqrt[a + b*Cot[c + d*x]^2],x]`

[Out] $-\left(\frac{\sqrt{a-b} \operatorname{ArcTan}\left[\frac{\sqrt{a-b} \cot(c+d*x)}{\sqrt{a+b \cot^2(c+d*x)^2}}\right]}{d} - \frac{\sqrt{b} \operatorname{ArcTanh}\left[\frac{\sqrt{b} \cot(c+d*x)}{\sqrt{a+b \cot^2(c+d*x)^2}}\right]}{d}\right)$

Rule 209

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 399

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[b/d, Int[(a + b*x^n)^(p-1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^n)^(p-1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p-1) + 1, 0] && IntegerQ[n]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{1+x^2} dx, x, \cot(c+dx)\right)}{d} \\
 &= -\frac{(a-b)\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{d} - \frac{b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{d} \\
 &= -\frac{(a-b)\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d} - \frac{b\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d} \\
 &= -\frac{\sqrt{a-b}\arctan\left(\frac{\sqrt{a-b}\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d} - \frac{\sqrt{b}\arctanh\left(\frac{\sqrt{b}\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.23

$$\int \sqrt{a + b \cot^2(c + dx)} dx$$

$$= \frac{\sqrt{a-b} \arctan\left(\frac{\sqrt{b} + \sqrt{b} \cot^2(c+dx) - \cot(c+dx) \sqrt{a+b \cot^2(c+dx)}}{\sqrt{a-b}}\right) + \sqrt{b} \log\left(-\sqrt{b} \cot(c+dx) + \sqrt{a+b \cot^2(c+dx)}\right)}{d}$$

[In] Integrate[Sqrt[a + b*Cot[c + d*x]^2], x]

[Out] (Sqrt[a - b]*ArcTan[(Sqrt[b] + Sqrt[b]*Cot[c + d*x]^2 - Cot[c + d*x]*Sqrt[a + b*Cot[c + d*x]^2])/Sqrt[a - b]] + Sqrt[b]*Log[-(Sqrt[b]*Cot[c + d*x]) + Sqrt[a + b*Cot[c + d*x]^2]])/d

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(75) = 150.

Time = 0.05 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.95

method	result
derivativedivides	$-\frac{\sqrt{b} \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{d} + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{db(a-b)} - \frac{a\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{db(a-b)}$
default	$-\frac{\sqrt{b} \ln\left(\sqrt{b} \cot(dx+c) + \sqrt{a+b \cot(dx+c)^2}\right)}{d} + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{db(a-b)} - \frac{a\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{db(a-b)}$

[In] int((a+b*cot(d*x+c)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/d*b^(1/2)*ln(b^(1/2)*cot(d*x+c)+(a+b*cot(d*x+c)^2)^(1/2))+1/d*(b^4*(a-b))^(1/2)/b/(a-b)*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(d*x+c)^2)^(1/2)*cot(d*x+c))-1/d*a*(b^4*(a-b))^(1/2)/b^2/(a-b)*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(d*x+c)^2)^(1/2)*cot(d*x+c))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(75) = 150.

Time = 0.30 (sec) , antiderivative size = 703, normalized size of antiderivative = 8.08

$$\int \sqrt{a + b \cot^2(c + dx)} dx$$

$$= \frac{\sqrt{-a + b} \log\left(- (a - b) \cos(2 dx + 2 c) + \sqrt{-a + b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c) + b\right) + \sqrt{b} \log\left(\frac{(a - 2 b) \cos(2 dx + 2 c) + 2 \sqrt{b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c) - a - 2 b}{\cos(2 dx + 2 c) - 1}\right)}{2 d}$$

$$- \frac{2 \sqrt{a - b} \arctan\left(- \frac{\sqrt{a - b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c)}{(a - b) \cos(2 dx + 2 c) + a - b}\right) - \sqrt{b} \log\left(\frac{(a - 2 b) \cos(2 dx + 2 c) + 2 \sqrt{b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c) - a - 2 b}{\cos(2 dx + 2 c) - 1}\right)}{2 d}$$

$$- \frac{\sqrt{a - b} \arctan\left(- \frac{\sqrt{a - b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c)}{(a - b) \cos(2 dx + 2 c) + a - b}\right) - \sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{(a-b) \cos(2 dx + 2 c) - a - b}{\cos(2 dx + 2 c) - 1}} \sin(2 dx + 2 c)}{b \cos(2 dx + 2 c) + b}\right)}{d}$$

[In] integrate((a+b*cot(d*x+c)^2)^(1/2),x, algorithm="fricas")

[Out] [1/2*(sqrt(-a + b)*log(-(a - b)*cos(2*d*x + 2*c) + sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + b) + sqrt(b)*log(((a - 2*b)*cos(2*d*x + 2*c) + 2*sqrt(b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - a - 2*b)/(cos(2*d*x + 2*c) - 1))/d, -1/2*(2*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(a - b)*cos(2*d*x + 2*c) + a - b) - sqrt(b)*log(((a - 2*b)*cos(2*d*x + 2*c) + 2*sqrt(b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) - a - 2*b)/(cos(2*d*x + 2*c) - 1))/d, 1/2*(2*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(b*cos(2*d*x + 2*c) + b)) + sqrt(-a + b)*log(-(a - b)*cos(2*d*x + 2*c) + sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + b))/d, -(sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(a - b)*cos(2*d*x + 2*c) + a - b) - sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/(b*cos(2*d*x + 2*c) + b)))/d]

Sympy [F]

$$\int \sqrt{a + b \cot^2(c + dx)} dx = \int \sqrt{a + b \cot^2(c + dx)} dx$$

[In] `integrate((a+b*cot(d*x+c)**2)**(1/2),x)`

[Out] `Integral(sqrt(a + b*cot(c + d*x)**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a + b \cot^2(c + dx)} dx = \text{Exception raised: ValueError}$$

[In] `integrate((a+b*cot(d*x+c)^2)^(1/2),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [F(-2)]

Exception generated.

$$\int \sqrt{a + b \cot^2(c + dx)} dx = \text{Exception raised: TypeError}$$

[In] `integrate((a+b*cot(d*x+c)^2)^(1/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx)::OUTPUT:Error: Bad Argument Type

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(c + dx)} dx = \int \sqrt{b \cot^2(c + dx) + a} dx$$

[In] `int((a + b*cot(c + d*x)^2)^(1/2),x)`

[Out] `int((a + b*cot(c + d*x)^2)^(1/2), x)`

3.34 $\int \frac{1}{\sqrt{a+b \cot^2(c+dx)}} dx$

Optimal result	228
Rubi [A] (verified)	228
Mathematica [B] (verified)	229
Maple [A] (verified)	230
Fricas [B] (verification not implemented)	230
Sympy [F]	231
Maxima [F(-2)]	231
Giac [B] (verification not implemented)	231
Mupad [B] (verification not implemented)	232

Optimal result

Integrand size = 16, antiderivative size = 47

$$\int \frac{1}{\sqrt{a+b \cot^2(c+dx)}} dx = -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{\sqrt{a-bd}}$$

[Out] $-\arctan(\cot(d*x+c)*(a-b)^{(1/2)}/(a+b*\cot(d*x+c)^2)^{(1/2)})/d/(a-b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3742, 385, 209}

$$\int \frac{1}{\sqrt{a+b \cot^2(c+dx)}} dx = -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d\sqrt{a-b}}$$

[In] `Int[1/Sqrt[a + b*Cot[c + d*x]^2],x]`

[Out] `-(ArcTan[(Sqrt[a - b]*Cot[c + d*x])/Sqrt[a + b*Cot[c + d*x]^2]]/(Sqrt[a - b]*d))`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{d} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{d} \\ &= -\frac{\arctan\left(\frac{\sqrt{a-b}\cot(c+dx)}{\sqrt{a+b\cot^2(c+dx)}}\right)}{\sqrt{a-bd}} \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 111 vs. $2(47) = 94$.

Time = 0.48 (sec) , antiderivative size = 111, normalized size of antiderivative = 2.36

$$\int \frac{1}{\sqrt{a+b\cot^2(c+dx)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{-\frac{(a-b)\cot^2(c+dx)}{a}}}{\sqrt{1+\frac{b\cot^2(c+dx)}{a}}}\right) \cot(c+dx) \sqrt{1+\frac{b\cot^2(c+dx)}{a}}}{d\sqrt{-\frac{(a-b)\cot^2(c+dx)}{a}} \sqrt{a+b\cot^2(c+dx)}}$$

```
[In] Integrate[1/Sqrt[a + b*Cot[c + d*x]^2], x]
```

```
[Out] -((ArcTanh[Sqrt[-((a - b)*Cot[c + d*x]^2)/a]]/Sqrt[1 + (b*Cot[c + d*x]^2)/a])*Cot[c + d*x]*Sqrt[1 + (b*Cot[c + d*x]^2)/a])/(d*Sqrt[-((a - b)*Cot[c + d*x]^2)/a])*Sqrt[a + b*Cot[c + d*x]^2])
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.45

method	result	size
derivativedivides	$-\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{d b^2(a-b)}$	68
default	$-\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{d b^2(a-b)}$	68

[In] `int(1/(a+b*cot(d*x+c)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/d*(b^4*(a-b))^(1/2)/b^2/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*\cot(d*x+c)^2)^(1/2)*\cot(d*x+c))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. $2(41) = 82$.

Time = 0.31 (sec) , antiderivative size = 239, normalized size of antiderivative = 5.09

$$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx$$

$$= \left[\begin{aligned} &-\frac{\sqrt{-a + b} \log\left(-2(a^2 - 2ab + b^2) \cos(2dx + 2c)^2 - 2((a - b) \cos(2dx + 2c) - b) \sqrt{-a + b} \sqrt{\frac{(a-b) \cos(2dx + 2c) - a - b}{\cos(2dx + 2c) - 1}}\right)}{4(a - b)d} \\ &-\frac{\arctan\left(-\frac{\sqrt{a-b} \sqrt{\frac{(a-b) \cos(2dx + 2c) - a - b}{\cos(2dx + 2c) - 1}} \sin(2dx + 2c)}{(a-b) \cos(2dx + 2c) - b}\right)}{2\sqrt{a - bd}} \end{aligned} \right]$$

[In] `integrate(1/(a+b*cot(d*x+c)^2)^(1/2),x, algorithm="fricas")`

[Out]
$$\left[-1/4*\sqrt{-a + b}*\log(-2*(a^2 - 2*a*b + b^2)*\cos(2*d*x + 2*c)^2 - 2*((a - b)*\cos(2*d*x + 2*c) - b)*\sqrt{-a + b}*\sqrt{\frac{(a - b)*\cos(2*d*x + 2*c) - a - b}{(\cos(2*d*x + 2*c) - 1))*\sin(2*d*x + 2*c)} + a^2 - 2*b^2 + 4*(a*b - b^2)*\cos(2*d*x + 2*c))/((a - b)*d), -1/2*\arctan(-\sqrt{a - b}*\sqrt{\frac{(a - b)*\cos(2*d*x + 2*c) - a - b}{(\cos(2*d*x + 2*c) - 1))*\sin(2*d*x + 2*c)}}/((a - b)*\cos(2*d*x + 2*c) - b))/(\sqrt{a - b}*d)\right]$$

Sympy [F]

$$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx = \int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx$$

[In] integrate(1/(a+b*cot(d*x+c)**2)**(1/2),x)

[Out] Integral(1/sqrt(a + b*cot(c + d*x)**2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(41) = 82.

Time = 1.32 (sec) , antiderivative size = 97, normalized size of antiderivative = 2.06

$$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx$$

$$= \frac{2 \arctan\left(-\frac{\sqrt{b} \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - \sqrt{b \tan(\frac{1}{2} dx + \frac{1}{2} c)^4 + 4a \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 - 2b \tan(\frac{1}{2} dx + \frac{1}{2} c)^2 + b + \sqrt{b}}{2\sqrt{a-b}}}\right)}{\sqrt{a-b} \operatorname{sgn}(\sin(dx+c))}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(1/2),x, algorithm="giac")

[Out] 2*arctan(-1/2*(sqrt(b)*tan(1/2*d*x + 1/2*c)^2 - sqrt(b*tan(1/2*d*x + 1/2*c)^4 + 4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d*x + 1/2*c)^2 + b) + sqrt(b))/sqrt(a - b))/(sqrt(a - b)*d*sgn(sin(d*x + c)))

Mupad [B] (verification not implemented)

Time = 13.73 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.87

$$\int \frac{1}{\sqrt{a + b \cot^2(c + dx)}} dx = -\frac{\operatorname{atan}\left(\frac{\cot(c+dx)\sqrt{a-b}}{\sqrt{b \cot^2(c+dx)+a}}\right)}{d\sqrt{a-b}}$$

[In] int(1/(a + b*cot(c + d*x)^2)^(1/2),x)

[Out] -atan((cot(c + d*x)*(a - b)^(1/2))/(a + b*cot(c + d*x)^2)^(1/2))/(d*(a - b)^(1/2))

$$3.35 \quad \int \frac{1}{(a+b \cot^2(c+dx))^{3/2}} dx$$

Optimal result	233
Rubi [A] (verified)	233
Mathematica [C] (warning: unable to verify)	235
Maple [A] (verified)	235
Fricas [B] (verification not implemented)	236
Sympy [F]	236
Maxima [F(-2)]	236
Giac [B] (verification not implemented)	237
Mupad [F(-1)]	237

Optimal result

Integrand size = 16, antiderivative size = 85

$$\int \frac{1}{(a+b \cot^2(c+dx))^{3/2}} dx = -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{3/2}d} + \frac{b \cot(c+dx)}{a(a-b)d\sqrt{a+b \cot^2(c+dx)}}$$

[Out] $-\arctan(\cot(d*x+c)*(a-b)^{(1/2)/(a+b*\cot(d*x+c)^2)^{(1/2)})/(a-b)^{(3/2)/d+b*\cot(d*x+c)/a/(a-b)/d/(a+b*\cot(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3742, 390, 385, 209}

$$\int \frac{1}{(a+b \cot^2(c+dx))^{3/2}} dx = \frac{b \cot(c+dx)}{ad(a-b)\sqrt{a+b \cot^2(c+dx)}} - \frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d(a-b)^{3/2}}$$

[In] $\text{Int}[(a + b*\text{Cot}[c + d*x]^2)^{-3/2}, x]$

[Out] $-(\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[c + d*x])/(\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2])]/((a - b)^{(3/2)*d})) + (b*\text{Cot}[c + d*x])/(a*(a - b)*d*\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2])$

Rule 209

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 390

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d)), x] + Dist[(b*c + n*(p + 1)*(b*c - a*d))/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p + q + 2) + 1, 0] && (LtQ[p, -1] || !LtQ[q, -1]) && NeQ[p, -1]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n)^(p)/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(c+dx)\right)}{d} \\
 &= \frac{b \cot(c+dx)}{a(a-b)d\sqrt{a+b \cot^2(c+dx)}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{(a-b)d} \\
 &= \frac{b \cot(c+dx)}{a(a-b)d\sqrt{a+b \cot^2(c+dx)}} - \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)d} \\
 &= -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{3/2}d} + \frac{b \cot(c+dx)}{a(a-b)d\sqrt{a+b \cot^2(c+dx)}}
 \end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 3.72 (sec) , antiderivative size = 231, normalized size of antiderivative = 2.72

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx =$$

$$\cos^2(c + dx) \cot(c + dx) \left(4(a - b)^2 \cos^2(c + dx) \operatorname{Hypergeometric2F1} \left(2, 2, \frac{7}{2}, \frac{(a-b) \cos^2(c+dx)}{a} \right) (b + a \tan^2(c + dx)) \right)$$

$$15a^3(a - b)d\sqrt{a}$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(-3/2), x]

[Out] -1/15*(Cos[c + d*x]^2*Cot[c + d*x]*(4*(a - b)^2*Cos[c + d*x]^2*Hypergeometric2F1[2, 2, 7/2, ((a - b)*Cos[c + d*x]^2)/a]*(b + a*Tan[c + d*x]^2) - (15*a*(2*b + 3*a*Tan[c + d*x]^2)*(ArcSin[Sqrt[((a - b)*Cos[c + d*x]^2)/a]]*(b + a*Tan[c + d*x]^2) - a*Sec[c + d*x]^2*Sqrt[((a - b)*Cos[c + d*x]^4*(b + a*Tan[c + d*x]^2)/a^2]))/Sqrt[((a - b)*Cos[c + d*x]^4*(b + a*Tan[c + d*x]^2)/a^2)))/(a^3*(a - b)*d*Sqrt[a + b*Cot[c + d*x]^2])

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.20

method	result	size
derivativedivides	$\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^2 b^2} + \frac{b \cot(dx+c)}{(a-b)a \sqrt{a+b \cot(dx+c)^2}}$ $\frac{\hspace{10em}}{d}$	102
default	$\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^2 b^2} + \frac{b \cot(dx+c)}{(a-b)a \sqrt{a+b \cot(dx+c)^2}}$ $\frac{\hspace{10em}}{d}$	102

[In] int(1/(a+b*cot(d*x+c)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/d*(-1/(a-b)^2*(b^4*(a-b))^(1/2)/b^2*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(d*x+c)^2)^(1/2)*cot(d*x+c))+b/(a-b)*cot(d*x+c)/a/(a+b*cot(d*x+c)^2)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(77) = 154.

Time = 0.34 (sec) , antiderivative size = 526, normalized size of antiderivative = 6.19

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx = \left[-\frac{(a^2 + ab - (a^2 - ab) \cos(2dx + 2c))\sqrt{-a + b} \log\left(-2(a^2 - 2ab + b^2) \cos(2dx + 2c) + a^2 - 2ab + b^2\right)}{(a^2 + ab - (a^2 - ab) \cos(2dx + 2c))\sqrt{-a + b} \log\left(-2(a^2 - 2ab + b^2) \cos(2dx + 2c) + a^2 - 2ab + b^2\right)} \right]$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(3/2),x, algorithm="fricas")

[Out] [-1/4*((a^2 + a*b - (a^2 - a*b)*cos(2*d*x + 2*c))*sqrt(-a + b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c)^2 + 2*((a - b)*cos(2*d*x + 2*c) - b)*sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + a^2 - 2*b^2 + 4*(a*b - b^2)*cos(2*d*x + 2*c)) + 4*(a*b - b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c))/((a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3)*d*cos(2*d*x + 2*c) - (a^4 - a^3*b - a^2*b^2 + a*b^3)*d), 1/2*((a^2 + a*b - (a^2 - a*b)*cos(2*d*x + 2*c))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) - b)) - 2*(a*b - b^2)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c))/((a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3)*d*cos(2*d*x + 2*c) - (a^4 - a^3*b - a^2*b^2 + a*b^3)*d)]

Sympy [F]

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx = \int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx$$

[In] integrate(1/(a+b*cot(d*x+c)**2)**(3/2),x)

[Out] Integral((a + b*cot(c + d*x)**2)**(-3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 300 vs. 2(77) = 154.

Time = 0.85 (sec) , antiderivative size = 300, normalized size of antiderivative = 3.53

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx =$$

$$\frac{\frac{(a^2 b \operatorname{sgn}(\sin(dx+c)) - 2 a b^2 \operatorname{sgn}(\sin(dx+c)) + b^3 \operatorname{sgn}(\sin(dx+c))) \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - a^2 b \operatorname{sgn}(\sin(dx+c)) - 2 a b^2 \operatorname{sgn}(\sin(dx+c)) + b^3 \operatorname{sgn}(\sin(dx+c))}{a^4 - 3 a^3 b + 3 a^2 b^2 - a b^3} - \frac{2 \arctan\left(\frac{\sqrt{b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 4 a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 2 b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + b}}{a - b}\right)}{d}}{\sqrt{b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^4 + 4 a \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 - 2 b \tan\left(\frac{1}{2} dx + \frac{1}{2} c\right)^2 + b}}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(3/2),x, algorithm="giac")

[Out] -(((a^2*b*sgn(sin(d*x + c)) - 2*a*b^2*sgn(sin(d*x + c)) + b^3*sgn(sin(d*x + c))) * tan(1/2*d*x + 1/2*c)^2 / (a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3) - (a^2*b*sgn(sin(d*x + c)) - 2*a*b^2*sgn(sin(d*x + c)) + b^3*sgn(sin(d*x + c))) / (a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3)) / sqrt(b*tan(1/2*d*x + 1/2*c)^4 + 4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d*x + 1/2*c)^2 + b) - 2*arctan(-1/2*(sqrt(b)*tan(1/2*d*x + 1/2*c)^2 - sqrt(b*tan(1/2*d*x + 1/2*c)^4 + 4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d*x + 1/2*c)^2 + b) + sqrt(b)) / sqrt(a - b)) / ((a*sgn(sin(d*x + c)) - b*sgn(sin(d*x + c))) * sqrt(a - b))) / d

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{3/2}} dx = \int \frac{1}{(b \cot^2(c + dx) + a)^{3/2}} dx$$

[In] int(1/(a + b*cot(c + d*x)^2)^(3/2),x)

[Out] int(1/(a + b*cot(c + d*x)^2)^(3/2), x)

3.36 $\int \frac{1}{(a+b \cot^2(c+dx))^{5/2}} dx$

Optimal result	238
Rubi [A] (verified)	238
Mathematica [C] (warning: unable to verify)	240
Maple [A] (verified)	241
Fricas [B] (verification not implemented)	242
Sympy [F]	243
Maxima [F(-2)]	243
Giac [B] (verification not implemented)	243
Mupad [F(-1)]	244

Optimal result

Integrand size = 16, antiderivative size = 135

$$\int \frac{1}{(a+b \cot^2(c+dx))^{5/2}} dx = -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{5/2}d} + \frac{b \cot(c+dx)}{3a(a-b)d(a+b \cot^2(c+dx))^{3/2}} + \frac{(5a-2b)b \cot(c+dx)}{3a^2(a-b)^2d\sqrt{a+b \cot^2(c+dx)}}$$

[Out] $-\arctan(\cot(d*x+c)*(a-b)^{(1/2)}/(a+b*\cot(d*x+c)^2)^{(1/2)})/(a-b)^{(5/2)}/d+1/3*b*\cot(d*x+c)/a/(a-b)/d/(a+b*\cot(d*x+c)^2)^{(3/2)}+1/3*(5*a-2*b)*b*\cot(d*x+c)/a^2/(a-b)^2/d/(a+b*\cot(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3742, 425, 541, 12, 385, 209}

$$\int \frac{1}{(a+b \cot^2(c+dx))^{5/2}} dx = \frac{b(5a-2b) \cot(c+dx)}{3a^2d(a-b)^2\sqrt{a+b \cot^2(c+dx)}} - \frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d(a-b)^{5/2}} + \frac{b \cot(c+dx)}{3ad(a-b)(a+b \cot^2(c+dx))^{3/2}}$$

[In] Int[(a + b*Cot[c + d*x]^2)^(-5/2), x]

[Out] $-(\text{ArcTan}[(\text{Sqrt}[a-b]*\text{Cot}[c+d*x])/\text{Sqrt}[a+b*\text{Cot}[c+d*x]^2]]/((a-b)^{(5/2)*d})) + (b*\text{Cot}[c+d*x])/(3*a*(a-b)*d*(a+b*\text{Cot}[c+d*x]^2)^{(3/2)}) + ((5*a-2*b)*b*\text{Cot}[c+d*x])/(3*a^2*(a-b)^2*d*\text{Sqrt}[a+b*\text{Cot}[c+d*x]^2])$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 425

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c -
a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c
+ d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n,
x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -
1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b,
c, d, n, p, q, x]
```

Rule 541

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f
_)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c
+ d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(
p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*
c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; Fre
eQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 3742

```
Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] :=>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= - \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(c+dx)\right)}{d} \\
&= \frac{b \cot(c+dx)}{3a(a-b)d (a+b \cot^2(c+dx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{3a-2b-2bx^2}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(c+dx)\right)}{3a(a-b)d} \\
&= \frac{b \cot(c+dx)}{3a(a-b)d (a+b \cot^2(c+dx))^{3/2}} + \frac{(5a-2b)b \cot(c+dx)}{3a^2(a-b)^2 d \sqrt{a+b \cot^2(c+dx)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{3a^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{3a^2(a-b)^2 d} \\
&= \frac{b \cot(c+dx)}{3a(a-b)d (a+b \cot^2(c+dx))^{3/2}} + \frac{(5a-2b)b \cot(c+dx)}{3a^2(a-b)^2 d \sqrt{a+b \cot^2(c+dx)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{(a-b)^2 d} \\
&= \frac{b \cot(c+dx)}{3a(a-b)d (a+b \cot^2(c+dx))^{3/2}} + \frac{(5a-2b)b \cot(c+dx)}{3a^2(a-b)^2 d \sqrt{a+b \cot^2(c+dx)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^2 d} \\
&= - \frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{5/2} d} + \frac{b \cot(c+dx)}{3a(a-b)d (a+b \cot^2(c+dx))^{3/2}} \\
&\quad + \frac{(5a-2b)b \cot(c+dx)}{3a^2(a-b)^2 d \sqrt{a+b \cot^2(c+dx)}}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.14 (sec) , antiderivative size = 367, normalized size of antiderivative = 2.72

$$\int \frac{1}{(a+b \cot^2(c+dx))^{5/2}} dx =$$

$$\cot^5(c+dx) \left(24(a-b)^3 \cos^2(c+dx) {}_3F_2\left(2, 2, 2; 1, \frac{9}{2}; \frac{(a-b) \cos^2(c+dx)}{a}\right) (b+a \tan^2(c+dx))^2 + 24(a-b)^3 \cos^2(c+dx) \right)$$

[In] Integrate[(a + b*Cot[c + d*x]^2)^(-5/2),x]

[Out]
$$-1/315*(\text{Cot}[c + d*x]^5*(24*(a - b)^3*\text{Cos}[c + d*x]^2*\text{HypergeometricPFQ}[\{2, 2\}, \{1, 9/2\}, ((a - b)*\text{Cos}[c + d*x]^2)/a]*(b + a*\text{Tan}[c + d*x]^2)^2 + 24*(a - b)^3*\text{Cos}[c + d*x]^2*\text{Hypergeometric2F1}[2, 2, 9/2, ((a - b)*\text{Cos}[c + d*x]^2)/a]*(3*b^2 + 7*a*b*\text{Tan}[c + d*x]^2 + 4*a^2*\text{Tan}[c + d*x]^4) - (35*a*(8*b^2 + 20*a*b*\text{Tan}[c + d*x]^2 + 15*a^2*\text{Tan}[c + d*x]^4)*(-3*\text{ArcSin}[\text{Sqrt}[(a - b)*\text{Cos}[c + d*x]^2)/a]]*(b + a*\text{Tan}[c + d*x]^2)^2 + a*\text{Sec}[c + d*x]^2*\text{Sqrt}[(a - b)*\text{Cos}[c + d*x]^4*(b + a*\text{Tan}[c + d*x]^2))/a^2*(4*b + a*(-1 + 3*\text{Tan}[c + d*x]^2))))/\text{Sqrt}[(a - b)*\text{Cos}[c + d*x]^4*(b + a*\text{Tan}[c + d*x]^2)/a^2]))/(a^5*(a - b)^2*d*(1 + \text{Cot}[c + d*x]^2)*\text{Sqrt}[a + b*\text{Cot}[c + d*x]^2]*(1 + (b*\text{Cot}[c + d*x]^2)/a))$$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.20

method	result
derivativedivides	$-\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^3 b^2} + \frac{b \left(\frac{\cot(dx+c)}{3a(a+b \cot(dx+c)^2)^{\frac{3}{2}}} + \frac{2 \cot(dx+c)}{3a^2 \sqrt{a+b \cot(dx+c)^2}} \right)}{a-b} + \frac{b \cot(dx+c)}{(a-b)^2 a \sqrt{a+b \cot(dx+c)^2}}$
default	$-\frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^3 b^2} + \frac{b \left(\frac{\cot(dx+c)}{3a(a+b \cot(dx+c)^2)^{\frac{3}{2}}} + \frac{2 \cot(dx+c)}{3a^2 \sqrt{a+b \cot(dx+c)^2}} \right)}{a-b} + \frac{b \cot(dx+c)}{(a-b)^2 a \sqrt{a+b \cot(dx+c)^2}}$

[In] int(1/(a+b*cot(d*x+c)^2)^(5/2),x,method=_RETURNVERBOSE)

[Out]
$$1/d*(-1/(a-b)^3*(b^4*(a-b))^{(1/2)}/b^2*\arctan(b^2*(a-b)/(b^4*(a-b))^{(1/2)})/(a + b*\text{cot}(d*x+c)^2)^{(1/2)}*\text{cot}(d*x+c))+1/(a-b)*b*(1/3*\text{cot}(d*x+c)/a/(a+b*\text{cot}(d*x+c)^2)^{(3/2)}+2/3/a^2*\text{cot}(d*x+c)/(a+b*\text{cot}(d*x+c)^2)^{(1/2)})+1/(a-b)^2*b*\text{cot}(d*x+c)/a/(a+b*\text{cot}(d*x+c)^2)^{(1/2)}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 423 vs. 2(121) = 242.

Time = 0.35 (sec) , antiderivative size = 898, normalized size of antiderivative = 6.65

$$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx = \frac{3(a^4 + 2a^3b + a^2b^2 + (a^4 - 2a^3b + a^2b^2) \cos(2dx + 2c)^2 - 2(a^4 - a^2b^2) \cos(2dx + 2c)) \sqrt{a-b} \arctan\left(\frac{3(a^4 + 2a^3b + a^2b^2 + (a^4 - 2a^3b + a^2b^2) \cos(2dx + 2c)^2 - 2(a^4 - a^2b^2) \cos(2dx + 2c)) \sqrt{a-b}}{6((a^7 - 5a^6b + 10a^5b^2 - 10a^4b^3 + 5a^3b^4 - a^2b^5)d \cos(2dx + 2c))^2}\right)}{6((a^7 - 5a^6b + 10a^5b^2 - 10a^4b^3 + 5a^3b^4 - a^2b^5)d \cos(2dx + 2c))^2}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(5/2),x, algorithm="fricas")

```
[Out] [-1/12*(3*(a^4 + 2*a^3*b + a^2*b^2 + (a^4 - 2*a^3*b + a^2*b^2)*cos(2*d*x + 2*c)^2 - 2*(a^4 - a^2*b^2)*cos(2*d*x + 2*c))*sqrt(-a + b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*d*x + 2*c)^2 - 2*((a - b)*cos(2*d*x + 2*c) - b)*sqrt(-a + b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c) + a^2 - 2*b^2 + 4*(a*b - b^2)*cos(2*d*x + 2*c)) - 8*(3*a^3*b - 2*a^2*b^2 - 2*a*b^3 + b^4 - (3*a^3*b - 7*a^2*b^2 + 5*a*b^3 - b^4)*cos(2*d*x + 2*c))*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c))/((a^7 - 5*a^6*b + 10*a^5*b^2 - 10*a^4*b^3 + 5*a^3*b^4 - a^2*b^5)*d*cos(2*d*x + 2*c)^2 - 2*(a^7 - 3*a^6*b + 2*a^5*b^2 + 2*a^4*b^3 - 3*a^3*b^4 + a^2*b^5)*d*cos(2*d*x + 2*c) + (a^7 - a^6*b - 2*a^5*b^2 + 2*a^4*b^3 + a^3*b^4 - a^2*b^5)*d), -1/6*(3*(a^4 + 2*a^3*b + a^2*b^2 + (a^4 - 2*a^3*b + a^2*b^2)*cos(2*d*x + 2*c)^2 - 2*(a^4 - a^2*b^2)*cos(2*d*x + 2*c))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) - b)) - 4*(3*a^3*b - 2*a^2*b^2 - 2*a*b^3 + b^4 - (3*a^3*b - 7*a^2*b^2 + 5*a*b^3 - b^4)*cos(2*d*x + 2*c))*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c))/((a^7 - 5*a^6*b + 10*a^5*b^2 - 10*a^4*b^3 + 5*a^3*b^4 - a^2*b^5)*d*cos(2*d*x + 2*c)^2 - 2*(a^7 - 3*a^6*b + 2*a^5*b^2 + 2*a^4*b^3 - 3*a^3*b^4 + a^2*b^5)*d*cos(2*d*x + 2*c) + (a^7 - a^6*b - 2*a^5*b^2 + 2*a^4*b^3 + a^3*b^4 - a^2*b^5)*d)]
```

Sympy [F]

$$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx = \int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx$$

```
[In] integrate(1/(a+b*cot(d*x+c)**2)**(5/2),x)
```

```
[Out] Integral((a + b*cot(c + d*x)**2)**(-5/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx = \text{Exception raised: ValueError}$$

```
[In] integrate(1/(a+b*cot(d*x+c)^2)^(5/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more det
ails)Is
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1160 vs. 2(121) = 242.

Time = 1.14 (sec) , antiderivative size = 1160, normalized size of antiderivative = 8.59

$$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx = \text{Too large to display}$$

```
[In] integrate(1/(a+b*cot(d*x+c)^2)^(5/2),x, algorithm="giac")
```

```
[Out] -1/3*(((5*a^9*b^2*sgn(sin(d*x + c)) - 42*a^8*b^3*sgn(sin(d*x + c)) + 156*
a^7*b^4*sgn(sin(d*x + c)) - 336*a^6*b^5*sgn(sin(d*x + c)) + 462*a^5*b^6*sgn
(sin(d*x + c)) - 420*a^4*b^7*sgn(sin(d*x + c)) + 252*a^3*b^8*sgn(sin(d*x +
c)) - 96*a^2*b^9*sgn(sin(d*x + c)) + 21*a*b^10*sgn(sin(d*x + c)) - 2*b^11*s
gn(sin(d*x + c)))*tan(1/2*d*x + 1/2*c)^2/(a^12 - 10*a^11*b + 45*a^10*b^2 -
120*a^9*b^3 + 210*a^8*b^4 - 252*a^7*b^5 + 210*a^6*b^6 - 120*a^5*b^7 + 45*a^
4*b^8 - 10*a^3*b^9 + a^2*b^10) + 3*(8*a^10*b*sgn(sin(d*x + c)) - 73*a^9*b^2
*sgn(sin(d*x + c)) + 298*a^8*b^3*sgn(sin(d*x + c)) - 716*a^7*b^4*sgn(sin(d*
x + c)) + 1120*a^6*b^5*sgn(sin(d*x + c)) - 1190*a^5*b^6*sgn(sin(d*x + c)) +
868*a^4*b^7*sgn(sin(d*x + c)) - 428*a^3*b^8*sgn(sin(d*x + c)) + 136*a^2*b^
```

```

9*sgn(sin(d*x + c)) - 25*a*b^10*sgn(sin(d*x + c)) + 2*b^11*sgn(sin(d*x + c))
)/(a^12 - 10*a^11*b + 45*a^10*b^2 - 120*a^9*b^3 + 210*a^8*b^4 - 252*a^7*b^5
+ 210*a^6*b^6 - 120*a^5*b^7 + 45*a^4*b^8 - 10*a^3*b^9 + a^2*b^10))*tan(1/
2*d*x + 1/2*c)^2 - 3*(8*a^10*b*sgn(sin(d*x + c)) - 73*a^9*b^2*sgn(sin(d*x +
c)) + 298*a^8*b^3*sgn(sin(d*x + c)) - 716*a^7*b^4*sgn(sin(d*x + c)) + 1120
*a^6*b^5*sgn(sin(d*x + c)) - 1190*a^5*b^6*sgn(sin(d*x + c)) + 868*a^4*b^7*sg
n(sin(d*x + c)) - 428*a^3*b^8*sgn(sin(d*x + c)) + 136*a^2*b^9*sgn(sin(d*x
+ c)) - 25*a*b^10*sgn(sin(d*x + c)) + 2*b^11*sgn(sin(d*x + c)))/(a^12 - 10*
a^11*b + 45*a^10*b^2 - 120*a^9*b^3 + 210*a^8*b^4 - 252*a^7*b^5 + 210*a^6*b^6
- 120*a^5*b^7 + 45*a^4*b^8 - 10*a^3*b^9 + a^2*b^10))*tan(1/2*d*x + 1/2*c)
^2 - (5*a^9*b^2*sgn(sin(d*x + c)) - 42*a^8*b^3*sgn(sin(d*x + c)) + 156*a^7*
b^4*sgn(sin(d*x + c)) - 336*a^6*b^5*sgn(sin(d*x + c)) + 462*a^5*b^6*sgn(sin
(d*x + c)) - 420*a^4*b^7*sgn(sin(d*x + c)) + 252*a^3*b^8*sgn(sin(d*x + c))
- 96*a^2*b^9*sgn(sin(d*x + c)) + 21*a*b^10*sgn(sin(d*x + c)) - 2*b^11*sgn(s
in(d*x + c)))/(a^12 - 10*a^11*b + 45*a^10*b^2 - 120*a^9*b^3 + 210*a^8*b^4 -
252*a^7*b^5 + 210*a^6*b^6 - 120*a^5*b^7 + 45*a^4*b^8 - 10*a^3*b^9 + a^2*b^
10))/(b*tan(1/2*d*x + 1/2*c)^4 + 4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d
*x + 1/2*c)^2 + b)^(3/2) - 6*arctan(-1/2*(sqrt(b)*tan(1/2*d*x + 1/2*c)^2 -
sqrt(b*tan(1/2*d*x + 1/2*c)^4 + 4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d
*x + 1/2*c)^2 + b) + sqrt(b))/sqrt(a - b))/((a^2*sgn(sin(d*x + c)) - 2*a*b*sg
n(sin(d*x + c)) + b^2*sgn(sin(d*x + c)))*sqrt(a - b))/d

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{5/2}} dx = \int \frac{1}{(b \cot(c + dx)^2 + a)^{5/2}} dx$$

[In] int(1/(a + b*cot(c + d*x)^2)^(5/2),x)

[Out] int(1/(a + b*cot(c + d*x)^2)^(5/2), x)

$$3.37 \quad \int \frac{1}{(a+b \cot^2(c+dx))^{7/2}} dx$$

Optimal result	245
Rubi [A] (verified)	245
Mathematica [C] (warning: unable to verify)	248
Maple [A] (verified)	249
Fricas [B] (verification not implemented)	250
Sympy [F]	251
Maxima [F(-2)]	251
Giac [B] (verification not implemented)	252
Mupad [F(-1)]	254

Optimal result

Integrand size = 16, antiderivative size = 190

$$\int \frac{1}{(a+b \cot^2(c+dx))^{7/2}} dx = -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{7/2}d} + \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3(a-b)^3d\sqrt{a+b \cot^2(c+dx)}}$$

[Out] $-\arctan(\cot(d*x+c)*(a-b)^{(1/2)}/(a+b*\cot(d*x+c)^2)^{(1/2)})/(a-b)^{(7/2)}/d+1/5*b*\cot(d*x+c)/a/(a-b)/d/(a+b*\cot(d*x+c)^2)^{(5/2)}+1/15*(9*a-4*b)*b*\cot(d*x+c)/a^2/(a-b)^2/d/(a+b*\cot(d*x+c)^2)^{(3/2)}+1/15*b*(33*a^2-26*a*b+8*b^2)*\cot(d*x+c)/a^3/(a-b)^3/d/(a+b*\cot(d*x+c)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3742, 425, 541, 12, 385, 209}

$$\int \frac{1}{(a+b \cot^2(c+dx))^{7/2}} dx = \frac{b(9a-4b) \cot(c+dx)}{15a^2d(a-b)^2(a+b \cot^2(c+dx))^{3/2}} + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3d(a-b)^3\sqrt{a+b \cot^2(c+dx)}} - \frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{d(a-b)^{7/2}} + \frac{b \cot(c+dx)}{5ad(a-b)(a+b \cot^2(c+dx))^{5/2}}$$

[In] Int[(a + b*Cot[c + d*x]^2)^(-7/2), x]

[Out] -(ArcTan[(Sqrt[a - b]*Cot[c + d*x])/Sqrt[a + b*Cot[c + d*x]^2]]/((a - b)^(7/2)*d)) + (b*Cot[c + d*x])/(5*a*(a - b)*d*(a + b*Cot[c + d*x]^2)^(5/2)) + ((9*a - 4*b)*b*Cot[c + d*x])/(15*a^2*(a - b)^2*d*(a + b*Cot[c + d*x]^2)^(3/2)) + (b*(33*a^2 - 26*a*b + 8*b^2)*Cot[c + d*x])/(15*a^3*(a - b)^3*d*Sqrt[a + b*Cot[c + d*x]^2])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 425

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 541

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(

$\text{ff*x})^n)^p/(c^2 + \text{ff}^2*x^2), x], x, c*(\text{Tan}[e + f*x]/\text{ff}), x]] /; \text{FreeQ}\{a, b, c, e, f, n, p\}, x\} \&\& (\text{IntegersQ}[n, p] \parallel \text{IGtQ}[p, 0] \parallel \text{EqQ}[n^2, 4] \parallel \text{E} \text{qQ}[n^2, 16])$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x^2)(a+bx^2)^{7/2}} dx, x, \cot(c+dx)\right)}{d} \\
 &= \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} - \frac{\text{Subst}\left(\int \frac{5a-4b-4bx^2}{(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(c+dx)\right)}{5a(a-b)d} \\
 &= \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{15a^2-18ab+8b^2-2(9a-4b)bx^2}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(c+dx)\right)}{15a^2(a-b)^2d} \\
 &= \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} \\
 &\quad + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3(a-b)^3d\sqrt{a+b \cot^2(c+dx)}} - \frac{\text{Subst}\left(\int \frac{15a^3}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{15a^3(a-b)^3d} \\
 &= \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} \\
 &\quad + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3(a-b)^3d\sqrt{a+b \cot^2(c+dx)}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(c+dx)\right)}{(a-b)^3d} \\
 &= \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} \\
 &\quad + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3(a-b)^3d\sqrt{a+b \cot^2(c+dx)}} - \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^3d} \\
 &= -\frac{\arctan\left(\frac{\sqrt{a-b} \cot(c+dx)}{\sqrt{a+b \cot^2(c+dx)}}\right)}{(a-b)^{7/2}d} + \frac{b \cot(c+dx)}{5a(a-b)d(a+b \cot^2(c+dx))^{5/2}} \\
 &\quad + \frac{(9a-4b)b \cot(c+dx)}{15a^2(a-b)^2d(a+b \cot^2(c+dx))^{3/2}} + \frac{b(33a^2-26ab+8b^2) \cot(c+dx)}{15a^3(a-b)^3d\sqrt{a+b \cot^2(c+dx)}}
 \end{aligned}$$

$$\begin{aligned}
& (\cos[c + dx]^2(b + a \tan[c + dx]^2)/a)/a^2 + (880b^3(((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^6 \text{Hypergeometric2F1}[2, 2, 11/2, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a^3 + 600 \\
& *(((a - b)\cos[c + dx]^2)/a)^{9/2} \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a \\
& + (1680b * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^2 \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a + (1560b^2 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^4 \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a^2 + (480b^3 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^6 \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a^3 + 80 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a + (240b^2 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^2 \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a + (240b^2 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^4 \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a^2 + (80b^3 * (((a - b)\cos[c + dx]^2)/a)^{9/2} \cot[c + dx]^6 \text{HypergeometricPFQ}[\{2, 2, 2, 2\}, \{1, 1, 11/2\}, ((a - b)\cos[c + dx]^2)/a] \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}/a^3 + 33075 * \sqrt{((a - b)\cos[c + dx]^4(b + a \tan[c + dx]^2))/a^2} + (66150b \cot[c + dx]^2 \sqrt{((a - b)\cos[c + dx]^4(b + a \tan[c + dx]^2))/a^2})/a + (52920b^2 \cot[c + dx]^4 \sqrt{((a - b)\cos[c + dx]^4(b + a \tan[c + dx]^2))/a^2})/a^2 + (15120b^3 \cot[c + dx]^6 \sqrt{((a - b)\cos[c + dx]^4(b + a \tan[c + dx]^2))/a^2})/a^3 - (198450(a - b)^2 b \text{ArcSin}[\sqrt{((a - b)\cos[c + dx]^2)/a}])/(a^3 * (\tan[c + dx] + \tan[c + dx]^3)^2)/(a^3 * d * (((a - b)\cos[c + dx]^2)/a)^{7/2} * (1 + \cot[c + dx]^2) \sqrt{a + b \cot[c + dx]^2} * (1 + (b \cot[c + dx]^2)/a)^2 \sqrt{((\cos[c + dx]^2(b + a \tan[c + dx]^2))/a)}
\end{aligned}$$

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 253, normalized size of antiderivative = 1.33

method	result
derivativedivides	$b \left(\frac{\cot(dx+c)}{5a(a+b \cot(dx+c))^{\frac{5}{2}}} + \frac{\frac{4 \cot(dx+c)}{15a(a+b \cot(dx+c))^{\frac{3}{2}}} + \frac{8 \cot(dx+c)}{15a^2 \sqrt{a+b \cot(dx+c)^2}}}{a} \right) - \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^4 b^2}$
default	$b \left(\frac{\cot(dx+c)}{5a(a+b \cot(dx+c))^{\frac{5}{2}}} + \frac{\frac{4 \cot(dx+c)}{15a(a+b \cot(dx+c))^{\frac{3}{2}}} + \frac{8 \cot(dx+c)}{15a^2 \sqrt{a+b \cot(dx+c)^2}}}{a} \right) - \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b) \cot(dx+c)}{\sqrt{b^4(a-b)} \sqrt{a+b \cot(dx+c)^2}}\right)}{(a-b)^4 b^2}$

[In] `int(1/(a+b*cot(d*x+c)^2)^(7/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{d} \left(\frac{1}{(a-b)} b \left(\frac{1}{5} \frac{\cot(dx+c)}{a(a+b \cot(dx+c)^2)^{5/2}} + \frac{4}{5} \frac{1}{a} \left(\frac{1}{3} \cot(dx+c) \right) \frac{1}{(a+b \cot(dx+c)^2)^{3/2}} + \frac{2}{3} \frac{1}{a^2} \frac{\cot(dx+c)}{(a+b \cot(dx+c)^2)^{1/2}} \right) - \frac{1}{(a-b)^4} \frac{b^4 (a-b)^{1/2}}{b^2} \arctan\left(\frac{b^2 (a-b)}{b^4 (a-b)} \frac{\cot(dx+c)}{\sqrt{a+b \cot(dx+c)^2}}\right) \right) + \frac{1}{(a-b)^2} b \left(\frac{1}{3} \frac{\cot(dx+c)}{a(a+b \cot(dx+c)^2)^{3/2}} + \frac{2}{3} \frac{1}{a^2} \frac{\cot(dx+c)}{(a+b \cot(dx+c)^2)^{1/2}} \right) + \frac{b}{(a-b)^3} \frac{\cot(dx+c)}{a(a+b \cot(dx+c)^2)^{1/2}}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 700 vs. $2(172) = 344$.

Time = 0.42 (sec) , antiderivative size = 1452, normalized size of antiderivative = 7.64

$$\int \frac{1}{(a+b \cot^2(c+dx))^{7/2}} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*cot(d*x+c)^2)^(7/2),x, algorithm="fricas")`

[Out] $[-1/60*(15*(a^6 + 3*a^5*b + 3*a^4*b^2 + a^3*b^3 - (a^6 - 3*a^5*b + 3*a^4*b^2 - a^3*b^3)*\cos(2*d*x + 2*c))^3 + 3*(a^6 - a^5*b - a^4*b^2 + a^3*b^3)*\cos(2*d*x + 2*c)^2 - 3*(a^6 + a^5*b - a^4*b^2 - a^3*b^3)*\cos(2*d*x + 2*c)]*\sqrt{(-a + b)*\log(-2*(a^2 - 2*a*b + b^2)*\cos(2*d*x + 2*c)^2 + 2*((a - b)*\cos(2*d*x + 2*c) - b)*\sqrt{-a + b}*\sqrt{((a - b)*\cos(2*d*x + 2*c) - a - b)/(\cos(2*d*x + 2*c) - 1)}*\sin(2*d*x + 2*c) + a^2 - 2*b^2 + 4*(a*b - b^2)*\cos(2*d*x + 2*c)} + 4*(45*a^5*b - 15*a^4*b^2 - 47*a^3*b^3 + 11*a^2*b^4 + 14*a*b^5 - 8*b^6 + (45*a^5*b - 165*a^4*b^2 + 233*a^3*b^3 - 159*a^2*b^4 + 54*a*b^5 - 8*b^6)*\cos(2*d*x + 2*c)^2 - 2*(45*a^5*b - 90*a^4*b^2 + 27*a^3*b^3 + 44*a^2*b^4 - 34*a*b^5 + 8*b^6)*\cos(2*d*x + 2*c)]*\sqrt{((a - b)*\cos(2*d*x + 2*c) - a - b)/(\cos(2*d*x + 2*c) - 1)}*\sin(2*d*x + 2*c)]/((a^10 - 7*a^9*b + 21*a^8*b^2 - 35*a^7*b^3 + 35*a^6*b^4 - 21*a^5*b^5 + 7*a^4*b^6 - a^3*b^7)*d*\cos(2*d*x + 2*c)^3 - 3*(a^10 - 5*a^9*b + 9*a^8*b^2 - 5*a^7*b^3 - 5*a^6*b^4 + 9*a^5*b^5 - 5*a^4*b^6 + a^3*b^7)*d*\cos(2*d*x + 2*c)^2 + 3*(a^10 - 3*a^9*b + a^8*b^2 +$

```

5*a^7*b^3 - 5*a^6*b^4 - a^5*b^5 + 3*a^4*b^6 - a^3*b^7)*d*cos(2*d*x + 2*c)
- (a^10 - a^9*b - 3*a^8*b^2 + 3*a^7*b^3 + 3*a^6*b^4 - 3*a^5*b^5 - a^4*b^6 +
a^3*b^7)*d), 1/30*(15*(a^6 + 3*a^5*b + 3*a^4*b^2 + a^3*b^3 - (a^6 - 3*a^5*
b + 3*a^4*b^2 - a^3*b^3)*cos(2*d*x + 2*c)^3 + 3*(a^6 - a^5*b - a^4*b^2 + a^
3*b^3)*cos(2*d*x + 2*c)^2 - 3*(a^6 + a^5*b - a^4*b^2 - a^3*b^3)*cos(2*d*x +
2*c))*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*d*x + 2*c) - a -
b)/(cos(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c)/((a - b)*cos(2*d*x + 2*c) - b)
) - 2*(45*a^5*b - 15*a^4*b^2 - 47*a^3*b^3 + 11*a^2*b^4 + 14*a*b^5 - 8*b^6 +
(45*a^5*b - 165*a^4*b^2 + 233*a^3*b^3 - 159*a^2*b^4 + 54*a*b^5 - 8*b^6)*co
s(2*d*x + 2*c)^2 - 2*(45*a^5*b - 90*a^4*b^2 + 27*a^3*b^3 + 44*a^2*b^4 - 34*
a*b^5 + 8*b^6)*cos(2*d*x + 2*c))*sqrt(((a - b)*cos(2*d*x + 2*c) - a - b)/(c
os(2*d*x + 2*c) - 1))*sin(2*d*x + 2*c))/((a^10 - 7*a^9*b + 21*a^8*b^2 - 35*
a^7*b^3 + 35*a^6*b^4 - 21*a^5*b^5 + 7*a^4*b^6 - a^3*b^7)*d*cos(2*d*x + 2*c)
^3 - 3*(a^10 - 5*a^9*b + 9*a^8*b^2 - 5*a^7*b^3 - 5*a^6*b^4 + 9*a^5*b^5 - 5*
a^4*b^6 + a^3*b^7)*d*cos(2*d*x + 2*c)^2 + 3*(a^10 - 3*a^9*b + a^8*b^2 + 5*a
^7*b^3 - 5*a^6*b^4 - a^5*b^5 + 3*a^4*b^6 - a^3*b^7)*d*cos(2*d*x + 2*c) - (a
^10 - a^9*b - 3*a^8*b^2 + 3*a^7*b^3 + 3*a^6*b^4 - 3*a^5*b^5 - a^4*b^6 + a^3
*b^7)*d)]

```

Sympy [F]

$$\int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx = \int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx$$

```
[In] integrate(1/(a+b*cot(d*x+c)**2)**(7/2),x)
```

```
[Out] Integral((a + b*cot(c + d*x)**2)**(-7/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx = \text{Exception raised: ValueError}$$

```
[In] integrate(1/(a+b*cot(d*x+c)^2)^(7/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested a
dditional constraints; using the 'assume' command before evaluation *may* h
elp (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more det
ails)Is
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3249 vs. $2(172) = 344$.

Time = 2.09 (sec) , antiderivative size = 3249, normalized size of antiderivative = 17.10

$$\int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cot(d*x+c)^2)^(7/2),x, algorithm="giac")

[Out] $\frac{1}{15} \cdot (30 \arctan(-\frac{1}{2} \cdot (\sqrt{b} \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c))^2 - \sqrt{b \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c))^4 + 4 \cdot a \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c))^2 - 2 \cdot b \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c)^2 + b) + \sqrt{b}) / \sqrt{a - b}) / ((a^3 \cdot \text{sgn}(\sin(d \cdot x + c))) - 3 \cdot a^2 \cdot b \cdot \text{sgn}(\sin(d \cdot x + c))) + 3 \cdot a \cdot b^2 \cdot \text{sgn}(\sin(d \cdot x + c)) - b^3 \cdot \text{sgn}(\sin(d \cdot x + c))) \cdot \sqrt{a - b}) - ((((((33 \cdot a^{20} \cdot b^3 \cdot \text{sgn}(\sin(d \cdot x + c))) - 620 \cdot a^{19} \cdot b^4 \cdot \text{sgn}(\sin(d \cdot x + c))) + 5525 \cdot a^{18} \cdot b^5 \cdot \text{sgn}(\sin(d \cdot x + c))) - 31050 \cdot a^{17} \cdot b^6 \cdot \text{sgn}(\sin(d \cdot x + c))) + 123420 \cdot a^{16} \cdot b^7 \cdot \text{sgn}(\sin(d \cdot x + c))) - 368832 \cdot a^{15} \cdot b^8 \cdot \text{sgn}(\sin(d \cdot x + c))) + 859860 \cdot a^{14} \cdot b^9 \cdot \text{sgn}(\sin(d \cdot x + c))) - 1601400 \cdot a^{13} \cdot b^{10} \cdot \text{sgn}(\sin(d \cdot x + c))) + 2419950 \cdot a^{12} \cdot b^{11} \cdot \text{sgn}(\sin(d \cdot x + c))) - 2996760 \cdot a^{11} \cdot b^{12} \cdot \text{sgn}(\sin(d \cdot x + c))) + 3058198 \cdot a^{10} \cdot b^{13} \cdot \text{sgn}(\sin(d \cdot x + c))) - 2576860 \cdot a^9 \cdot b^{14} \cdot \text{sgn}(\sin(d \cdot x + c))) + 1790100 \cdot a^8 \cdot b^{15} \cdot \text{sgn}(\sin(d \cdot x + c))) - 1020000 \cdot a^7 \cdot b^{16} \cdot \text{sgn}(\sin(d \cdot x + c))) + 472260 \cdot a^6 \cdot b^{17} \cdot \text{sgn}(\sin(d \cdot x + c))) - 175032 \cdot a^5 \cdot b^{18} \cdot \text{sgn}(\sin(d \cdot x + c))) + 50745 \cdot a^4 \cdot b^{19} \cdot \text{sgn}(\sin(d \cdot x + c))) - 11100 \cdot a^3 \cdot b^{20} \cdot \text{sgn}(\sin(d \cdot x + c))) + 1725 \cdot a^2 \cdot b^{21} \cdot \text{sgn}(\sin(d \cdot x + c))) - 170 \cdot a \cdot b^{22} \cdot \text{sgn}(\sin(d \cdot x + c))) + 8 \cdot b^{23} \cdot \text{sgn}(\sin(d \cdot x + c))) \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c))^2 / (a^{24} - 21 \cdot a^{23} \cdot b + 210 \cdot a^{22} \cdot b^2 - 1330 \cdot a^{21} \cdot b^3 + 5985 \cdot a^{20} \cdot b^4 - 20349 \cdot a^{19} \cdot b^5 + 54264 \cdot a^{18} \cdot b^6 - 116280 \cdot a^{17} \cdot b^7 + 203490 \cdot a^{16} \cdot b^8 - 293930 \cdot a^{15} \cdot b^9 + 352716 \cdot a^{14} \cdot b^{10} - 352716 \cdot a^{13} \cdot b^{11} + 293930 \cdot a^{12} \cdot b^{12} - 203490 \cdot a^{11} \cdot b^{13} + 116280 \cdot a^{10} \cdot b^{14} - 54264 \cdot a^9 \cdot b^{15} + 20349 \cdot a^8 \cdot b^{16} - 5985 \cdot a^7 \cdot b^{17} + 1330 \cdot a^6 \cdot b^{18} - 210 \cdot a^5 \cdot b^{19} + 21 \cdot a^4 \cdot b^{20} - a^3 \cdot b^{21}) + 5 \cdot (60 \cdot a^{21} \cdot b^2 \cdot \text{sgn}(\sin(d \cdot x + c))) - 1165 \cdot a^{20} \cdot b^3 \cdot \text{sgn}(\sin(d \cdot x + c))) + 10752 \cdot a^{19} \cdot b^4 \cdot \text{sgn}(\sin(d \cdot x + c))) - 62729 \cdot a^{18} \cdot b^5 \cdot \text{sgn}(\sin(d \cdot x + c))) + 259530 \cdot a^{17} \cdot b^6 \cdot \text{sgn}(\sin(d \cdot x + c))) - 809676 \cdot a^{16} \cdot b^7 \cdot \text{sgn}(\sin(d \cdot x + c))) + 1977168 \cdot a^{15} \cdot b^8 \cdot \text{sgn}(\sin(d \cdot x + c))) - 3871716 \cdot a^{14} \cdot b^9 \cdot \text{sgn}(\sin(d \cdot x + c))) + 6178752 \cdot a^{13} \cdot b^{10} \cdot \text{sgn}(\sin(d \cdot x + c))) - 8121750 \cdot a^{12} \cdot b^{11} \cdot \text{sgn}(\sin(d \cdot x + c))) + 8850608 \cdot a^{11} \cdot b^{12} \cdot \text{sgn}(\sin(d \cdot x + c))) - 8020974 \cdot a^{10} \cdot b^{13} \cdot \text{sgn}(\sin(d \cdot x + c))) + 6045676 \cdot a^9 \cdot b^{14} \cdot \text{sgn}(\sin(d \cdot x + c))) - 3778692 \cdot a^8 \cdot b^{15} \cdot \text{sgn}(\sin(d \cdot x + c))) + 1946160 \cdot a^7 \cdot b^{16} \cdot \text{sgn}(\sin(d \cdot x + c))) - 817428 \cdot a^6 \cdot b^{17} \cdot \text{sgn}(\sin(d \cdot x + c))) + 275604 \cdot a^5 \cdot b^{18} \cdot \text{sgn}(\sin(d \cdot x + c))) - 72837 \cdot a^4 \cdot b^{19} \cdot \text{sgn}(\sin(d \cdot x + c))) + 14544 \cdot a^3 \cdot b^{20} \cdot \text{sgn}(\sin(d \cdot x + c))) - 2065 \cdot a^2 \cdot b^{21} \cdot \text{sgn}(\sin(d \cdot x + c))) + 186 \cdot a \cdot b^{22} \cdot \text{sgn}(\sin(d \cdot x + c))) - 8 \cdot b^{23} \cdot \text{sgn}(\sin(d \cdot x + c))) / (a^{24} - 21 \cdot a^{23} \cdot b + 210 \cdot a^{22} \cdot b^2 - 1330 \cdot a^{21} \cdot b^3 + 5985 \cdot a^{20} \cdot b^4 - 20349 \cdot a^{19} \cdot b^5 + 54264 \cdot a^{18} \cdot b^6 - 116280 \cdot a^{17} \cdot b^7 + 203490 \cdot a^{16} \cdot b^8 - 293930 \cdot a^{15} \cdot b^9 + 352716 \cdot a^{14} \cdot b^{10} - 352716 \cdot a^{13} \cdot b^{11} + 293930 \cdot a^{12} \cdot b^{12} - 203490 \cdot a^{11} \cdot b^{13} + 116280 \cdot a^{10} \cdot b^{14} - 54264 \cdot a^9 \cdot b^{15} + 20349 \cdot a^8 \cdot b^{16} - 5985 \cdot a^7 \cdot b^{17} + 1330 \cdot a^6 \cdot b^{18} - 210 \cdot a^5 \cdot b^{19} + 21 \cdot a^4 \cdot b^{20} - a^3 \cdot b^{21})) \cdot \tan(\frac{1}{2} \cdot d \cdot x + \frac{1}{2} \cdot c))^2 + 10 \cdot (72 \cdot a^{22} \cdot b \cdot \text{sgn}(\sin(d \cdot x + c))) - 1458 \cdot a^{21} \cdot b^2$

$$\begin{aligned}
& * \operatorname{sgn}(\sin(dx + c)) + 14067a^{20}b^3 \operatorname{sgn}(\sin(dx + c)) - 86018a^{19}b^4 \operatorname{sgn}(\sin(dx + c)) + 374075a^{18}b^5 \operatorname{sgn}(\sin(dx + c)) - 1230570a^{17}b^6 \operatorname{sgn}(\sin(dx + c)) + 3179748a^{16}b^7 \operatorname{sgn}(\sin(dx + c)) - 6614904a^{15}b^8 \operatorname{sgn}(\sin(dx + c)) + 11265084a^{14}b^9 \operatorname{sgn}(\sin(dx + c)) - 15882420a^{13}b^{10} \operatorname{sgn}(\sin(dx + c)) + 18674058a^{12}b^{11} \operatorname{sgn}(\sin(dx + c)) - 18386316a^{11}b^{12} \operatorname{sgn}(\sin(dx + c)) + 15180490a^{10}b^{13} \operatorname{sgn}(\sin(dx + c)) - 10497364a^9b^{14} \operatorname{sgn}(\sin(dx + c)) + 6055740a^8b^{15} \operatorname{sgn}(\sin(dx + c)) - 2893944a^7b^{16} \operatorname{sgn}(\sin(dx + c)) + 1133220a^6b^{17} \operatorname{sgn}(\sin(dx + c)) - 357786a^5b^{18} \operatorname{sgn}(\sin(dx + c)) + 88923a^4b^{19} \operatorname{sgn}(\sin(dx + c)) - 16770a^3b^{20} \operatorname{sgn}(\sin(dx + c)) + 2259a^2b^{21} \operatorname{sgn}(\sin(dx + c)) - 194ab^{22} \operatorname{sgn}(\sin(dx + c)) + 8b^{23} \operatorname{sgn}(\sin(dx + c)) / (a^{24} - 21a^{23}b + 210a^{22}b^2 - 1330a^{21}b^3 + 5985a^{20}b^4 - 20349a^{19}b^5 + 54264a^{18}b^6 - 116280a^{17}b^7 + 203490a^{16}b^8 - 293930a^{15}b^9 + 352716a^{14}b^{10} - 352716a^{13}b^{11} + 293930a^{12}b^{12} - 203490a^{11}b^{13} + 116280a^{10}b^{14} - 54264a^9b^{15} + 20349a^8b^{16} - 5985a^7b^{17} + 1330a^6b^{18} - 210a^5b^{19} + 21a^4b^{20} - a^3b^{21}) * \tan(1/2dx + 1/2c)^2 - 10(72a^{22}b \operatorname{sgn}(\sin(dx + c)) - 1458a^{21}b^2 \operatorname{sgn}(\sin(dx + c)) + 14067a^{20}b^3 \operatorname{sgn}(\sin(dx + c)) - 86018a^{19}b^4 \operatorname{sgn}(\sin(dx + c)) + 374075a^{18}b^5 \operatorname{sgn}(\sin(dx + c)) - 1230570a^{17}b^6 \operatorname{sgn}(\sin(dx + c)) + 3179748a^{16}b^7 \operatorname{sgn}(\sin(dx + c)) - 6614904a^{15}b^8 \operatorname{sgn}(\sin(dx + c)) + 11265084a^{14}b^9 \operatorname{sgn}(\sin(dx + c)) - 15882420a^{13}b^{10} \operatorname{sgn}(\sin(dx + c)) + 18674058a^{12}b^{11} \operatorname{sgn}(\sin(dx + c)) - 18386316a^{11}b^{12} \operatorname{sgn}(\sin(dx + c)) + 15180490a^{10}b^{13} \operatorname{sgn}(\sin(dx + c)) - 10497364a^9b^{14} \operatorname{sgn}(\sin(dx + c)) + 6055740a^8b^{15} \operatorname{sgn}(\sin(dx + c)) - 2893944a^7b^{16} \operatorname{sgn}(\sin(dx + c)) + 1133220a^6b^{17} \operatorname{sgn}(\sin(dx + c)) - 357786a^5b^{18} \operatorname{sgn}(\sin(dx + c)) + 88923a^4b^{19} \operatorname{sgn}(\sin(dx + c)) - 16770a^3b^{20} \operatorname{sgn}(\sin(dx + c)) + 2259a^2b^{21} \operatorname{sgn}(\sin(dx + c)) - 194ab^{22} \operatorname{sgn}(\sin(dx + c)) + 8b^{23} \operatorname{sgn}(\sin(dx + c))) / (a^{24} - 21a^{23}b + 210a^{22}b^2 - 1330a^{21}b^3 + 5985a^{20}b^4 - 20349a^{19}b^5 + 54264a^{18}b^6 - 116280a^{17}b^7 + 203490a^{16}b^8 - 293930a^{15}b^9 + 352716a^{14}b^{10} - 352716a^{13}b^{11} + 293930a^{12}b^{12} - 203490a^{11}b^{13} + 116280a^{10}b^{14} - 54264a^9b^{15} + 20349a^8b^{16} - 5985a^7b^{17} + 1330a^6b^{18} - 210a^5b^{19} + 21a^4b^{20} - a^3b^{21}) * \tan(1/2dx + 1/2c)^2 - 5(60a^{21}b^2 \operatorname{sgn}(\sin(dx + c)) - 1165a^{20}b^3 \operatorname{sgn}(\sin(dx + c)) + 10752a^{19}b^4 \operatorname{sgn}(\sin(dx + c)) - 62729a^{18}b^5 \operatorname{sgn}(\sin(dx + c)) + 259530a^{17}b^6 \operatorname{sgn}(\sin(dx + c)) - 809676a^{16}b^7 \operatorname{sgn}(\sin(dx + c)) + 1977168a^{15}b^8 \operatorname{sgn}(\sin(dx + c)) - 3871716a^{14}b^9 \operatorname{sgn}(\sin(dx + c)) + 6178752a^{13}b^{10} \operatorname{sgn}(\sin(dx + c)) - 8121750a^{12}b^{11} \operatorname{sgn}(\sin(dx + c)) + 8850608a^{11}b^{12} \operatorname{sgn}(\sin(dx + c)) - 8020974a^{10}b^{13} \operatorname{sgn}(\sin(dx + c)) + 6045676a^9b^{14} \operatorname{sgn}(\sin(dx + c)) - 3778692a^8b^{15} \operatorname{sgn}(\sin(dx + c)) + 1946160a^7b^{16} \operatorname{sgn}(\sin(dx + c)) - 817428a^6b^{17} \operatorname{sgn}(\sin(dx + c)) + 275604a^5b^{18} \operatorname{sgn}(\sin(dx + c)) - 72837a^4b^{19} \operatorname{sgn}(\sin(dx + c)) + 14544a^3b^{20} \operatorname{sgn}(\sin(dx + c)) - 2065a^2b^{21} \operatorname{sgn}(\sin(dx + c)) + 186ab^{22} \operatorname{sgn}(\sin(dx + c)) - 8b^{23} \operatorname{sgn}(\sin(dx + c))) / (a^{24} - 21a^{23}b + 210a^{22}b^2 - 1330a^{21}b^3 + 5985a^{20}b^4 - 20349a^{19}b^5 + 54264a^{18}b^6 - 116280a^{17}b^7 + 203490a^{16}b^8 - 293930a^{15}b^9 + 352716a^{14}b^{10} - 352716a^{13}b^{11} + 293930a^{12}b^{12} - 203490a^{11}b^{13}
\end{aligned}$$

```

+ 116280*a^10*b^14 - 54264*a^9*b^15 + 20349*a^8*b^16 - 5985*a^7*b^17 + 1330
*a^6*b^18 - 210*a^5*b^19 + 21*a^4*b^20 - a^3*b^21))*tan(1/2*d*x + 1/2*c)^2
- (33*a^20*b^3*sgn(sin(d*x + c)) - 620*a^19*b^4*sgn(sin(d*x + c)) + 5525*a^
18*b^5*sgn(sin(d*x + c)) - 31050*a^17*b^6*sgn(sin(d*x + c)) + 123420*a^16*b
^7*sgn(sin(d*x + c)) - 368832*a^15*b^8*sgn(sin(d*x + c)) + 859860*a^14*b^9*
sgn(sin(d*x + c)) - 1601400*a^13*b^10*sgn(sin(d*x + c)) + 2419950*a^12*b^11
*sgn(sin(d*x + c)) - 2996760*a^11*b^12*sgn(sin(d*x + c)) + 3058198*a^10*b^1
3*sgn(sin(d*x + c)) - 2576860*a^9*b^14*sgn(sin(d*x + c)) + 1790100*a^8*b^15
*sgn(sin(d*x + c)) - 1020000*a^7*b^16*sgn(sin(d*x + c)) + 472260*a^6*b^17*s
gn(sin(d*x + c)) - 175032*a^5*b^18*sgn(sin(d*x + c)) + 50745*a^4*b^19*sgn(s
in(d*x + c)) - 11100*a^3*b^20*sgn(sin(d*x + c)) + 1725*a^2*b^21*sgn(sin(d*x
+ c)) - 170*a*b^22*sgn(sin(d*x + c)) + 8*b^23*sgn(sin(d*x + c)))/(a^24 - 2
1*a^23*b + 210*a^22*b^2 - 1330*a^21*b^3 + 5985*a^20*b^4 - 20349*a^19*b^5 +
54264*a^18*b^6 - 116280*a^17*b^7 + 203490*a^16*b^8 - 293930*a^15*b^9 + 3527
16*a^14*b^10 - 352716*a^13*b^11 + 293930*a^12*b^12 - 203490*a^11*b^13 + 116
280*a^10*b^14 - 54264*a^9*b^15 + 20349*a^8*b^16 - 5985*a^7*b^17 + 1330*a^6*
b^18 - 210*a^5*b^19 + 21*a^4*b^20 - a^3*b^21))/(b*tan(1/2*d*x + 1/2*c)^4 +
4*a*tan(1/2*d*x + 1/2*c)^2 - 2*b*tan(1/2*d*x + 1/2*c)^2 + b)^(5/2))/d

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cot^2(c + dx))^{7/2}} dx = \int \frac{1}{(b \cot(c + dx)^2 + a)^{7/2}} dx$$

[In] int(1/(a + b*cot(c + d*x)^2)^(7/2),x)

[Out] int(1/(a + b*cot(c + d*x)^2)^(7/2), x)

3.38 $\int (1 - \cot^2(x))^{3/2} dx$

Optimal result	255
Rubi [A] (verified)	255
Mathematica [B] (verified)	257
Maple [A] (verified)	257
Fricas [B] (verification not implemented)	258
Sympy [F]	258
Maxima [F]	258
Giac [B] (verification not implemented)	259
Mupad [B] (verification not implemented)	259

Optimal result

Integrand size = 12, antiderivative size = 54

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{5}{2} \arcsin(\cot(x)) - 2\sqrt{2} \arctan\left(\frac{\sqrt{2} \cot(x)}{\sqrt{1 - \cot^2(x)}}\right) + \frac{1}{2} \cot(x) \sqrt{1 - \cot^2(x)}$$

[Out] 5/2*arcsin(cot(x))-2*arctan(cot(x)*2^(1/2)/(1-cot(x)^2)^(1/2))*2^(1/2)+1/2*cot(x)*(1-cot(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 427, 537, 222, 385, 209}

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{5}{2} \arcsin(\cot(x)) - 2\sqrt{2} \arctan\left(\frac{\sqrt{2} \cot(x)}{\sqrt{1 - \cot^2(x)}}\right) + \frac{1}{2} \cot(x) \sqrt{1 - \cot^2(x)}$$

[In] Int[(1 - Cot[x]^2)^(3/2), x]

[Out] (5*ArcSin[Cot[x]])/2 - 2*Sqrt[2]*ArcTan[(Sqrt[2]*Cot[x])/Sqrt[1 - Cot[x]^2]] + (Cot[x]*Sqrt[1 - Cot[x]^2])/2

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 222

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 427

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[d*x*(a + b*x^n)^(p+1)*((c + d*x^n)^(q-1)/(b*(n*(p+q)+1))), x] + Dist[1/(b*(n*(p+q)+1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q-2)*Simp[c*(b*c*(n*(p+q)+1) - a*d) + d*(b*c*(n*(p+2*q-1)+1) - a*d*(n*(q-1)+1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d, 0] && GtQ[q, 1] && NeQ[n*(p+q)+1, 0] && !IGtQ[p, 1] && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 537

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e - a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{(1-x^2)^{3/2}}{1+x^2} dx, x, \cot(x)\right) \\ &= \frac{1}{2} \cot(x) \sqrt{1-\cot^2(x)} - \frac{1}{2} \text{Subst}\left(\int \frac{3-5x^2}{\sqrt{1-x^2}(1+x^2)} dx, x, \cot(x)\right) \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \cot(x) \sqrt{1 - \cot^2(x)} + \frac{5}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1-x^2}} dx, x, \cot(x) \right) \\
&\quad - 4 \text{Subst} \left(\int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx, x, \cot(x) \right) \\
&= \frac{5}{2} \arcsin(\cot(x)) + \frac{1}{2} \cot(x) \sqrt{1 - \cot^2(x)} - 4 \text{Subst} \left(\int \frac{1}{1+2x^2} dx, x, \frac{\cot(x)}{\sqrt{1 - \cot^2(x)}} \right) \\
&= \frac{5}{2} \arcsin(\cot(x)) - 2\sqrt{2} \arctan \left(\frac{\sqrt{2} \cot(x)}{\sqrt{1 - \cot^2(x)}} \right) + \frac{1}{2} \cot(x) \sqrt{1 - \cot^2(x)}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 123 vs. $2(54) = 108$.

Time = 0.51 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.28

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{1}{2} (1 - \cot^2(x))^{3/2} \sec^2(2x) \left(\arctan \left(\frac{\cos(x)}{\sqrt{-\cos(2x)}} \right) \sqrt{-\cos(2x)} \sin^3(x) + 4 \operatorname{arctanh} \left(\frac{\cos(x)}{\sqrt{\cos(2x)}} \right) \sqrt{\cos(2x)} \sin^3(x) \right)$$

[In] Integrate[(1 - Cot[x]^2)^(3/2), x]

[Out] ((1 - Cot[x]^2)^(3/2)*Sec[2*x]^2*(ArcTan[Cos[x]/Sqrt[-Cos[2*x]])*Sqrt[-Cos[2*x]]*Sin[x]^3 + 4*ArcTanh[Cos[x]/Sqrt[Cos[2*x]]*Sqrt[Cos[2*x]]*Sin[x]^3 - 4*Sqrt[2]*Sqrt[Cos[2*x]]*Log[Sqrt[2]*Cos[x] + Sqrt[Cos[2*x]]]*Sin[x]^3 - Sin[4*x]/4)/2

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{5 \arcsin(\cot(x))}{2} + \frac{\cot(x) \sqrt{1 - \cot^2(x)}}{2} + 2\sqrt{2} \arctan \left(\frac{\sqrt{2} \sqrt{1 - \cot^2(x)} \cot(x)}{-1 + \cot(x)^2} \right)$	51
default	$\frac{5 \arcsin(\cot(x))}{2} + \frac{\cot(x) \sqrt{1 - \cot^2(x)}}{2} + 2\sqrt{2} \arctan \left(\frac{\sqrt{2} \sqrt{1 - \cot^2(x)} \cot(x)}{-1 + \cot(x)^2} \right)$	51

[In] int((1-cot(x)^2)^(3/2), x, method=_RETURNVERBOSE)

[Out] 5/2*arcsin(cot(x))+1/2*cot(x)*(1-cot(x)^2)^(1/2)+2*2^(1/2)*arctan(2^(1/2)*(1-cot(x)^2)^(1/2)/(-1+cot(x)^2)*cot(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(42) = 84.

Time = 0.31 (sec) , antiderivative size = 110, normalized size of antiderivative = 2.04

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{4\sqrt{2} \arctan\left(\frac{\sqrt{\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x)}{\cos(2x)+1}\right) \sin(2x) + \sqrt{2} \sqrt{\frac{\cos(2x)}{\cos(2x)-1}} (\cos(2x) + 1) - 5 \arctan\left(\frac{\sqrt{2} \sqrt{\frac{\cos(2x)}{\cos(2x)-1}}}{\cos(2x)+1}\right) \sin(2x)}{2 \sin(2x)}$$

[In] integrate((1-cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] 1/2*(4*sqrt(2)*arctan(sqrt(cos(2*x)/(cos(2*x) - 1))*sin(2*x)/(cos(2*x) + 1))*sin(2*x) + sqrt(2)*sqrt(cos(2*x)/(cos(2*x) - 1))*(cos(2*x) + 1) - 5*arctan(sqrt(2)*sqrt(cos(2*x)/(cos(2*x) - 1))*sin(2*x)/(cos(2*x) + 1))*sin(2*x))/sin(2*x)

Sympy [F]

$$\int (1 - \cot^2(x))^{3/2} dx = \int (1 - \cot^2(x))^{\frac{3}{2}} dx$$

[In] integrate((1-cot(x)**2)**(3/2),x)

[Out] Integral((1 - cot(x)**2)**(3/2), x)

Maxima [F]

$$\int (1 - \cot^2(x))^{3/2} dx = \int (-\cot(x)^2 + 1)^{\frac{3}{2}} dx$$

[In] integrate((1-cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate((-cot(x)^2 + 1)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 257 vs. $2(42) = 84$.

Time = 0.33 (sec) , antiderivative size = 257, normalized size of antiderivative = 4.76

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{1}{4} \left(5 \pi \operatorname{sgn}(\cos(x)) - 4 \sqrt{2} \left(\pi \operatorname{sgn}(\cos(x)) + 2 \arctan \left(- \frac{\left(\frac{\sqrt{2} \sqrt{-2 \cos(x)^2 + 1} - \sqrt{2}}{\cos(x)^2} \right)^2 - 4}{4 \left(\sqrt{2} \sqrt{-2 \cos(x)^2 + 1} - \right)} \right) \right) \right)$$

[In] integrate((1-cot(x)^2)^(3/2),x, algorithm="giac")

[Out] 1/4*(5*pi*sgn(cos(x)) - 4*sqrt(2)*(pi*sgn(cos(x)) + 2*arctan(-1/4*((sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2))/cos(x)^2 - 4)*cos(x)/(sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2)))) + 4*sqrt(2)*((sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2))/cos(x) - 4*cos(x)/(sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2)))/(((sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2))/cos(x) - 4*cos(x)/(sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2)))^2 + 8) + 10*arctan(-1/4*sqrt(2)*((sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2))/cos(x)^2 - 4)*cos(x)/(sqrt(2)*sqrt(-2*cos(x)^2 + 1) - sqrt(2))))*sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 14.22 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.93

$$\int (1 - \cot^2(x))^{3/2} dx = \frac{5 \operatorname{asin}(\cot(x))}{2} + \frac{\cot(x) \sqrt{1 - \cot(x)^2}}{2} - \sqrt{2} \ln \left(\frac{\frac{\sqrt{2}(-1+\cot(x) \operatorname{li}) \operatorname{li}}{2} - \sqrt{1 - \cot(x)^2} \operatorname{li}}{\cot(x) - i} \right) \operatorname{li} + \sqrt{2} \ln \left(\frac{\frac{\sqrt{2}(1+\cot(x) \operatorname{li}) \operatorname{li}}{2} + \sqrt{1 - \cot(x)^2} \operatorname{li}}{\cot(x) + i} \right) \operatorname{li}$$

[In] int((1 - cot(x)^2)^(3/2),x)

[Out] (5*asin(cot(x)))/2 + (cot(x)*(1 - cot(x)^2)^(1/2))/2 - 2^(1/2)*log(((2^(1/2)*(cot(x)*1i - 1)*1i)/2 - (1 - cot(x)^2)^(1/2)*1i)/(cot(x) - 1i))*1i + 2^(1/2)*log(((2^(1/2)*(cot(x)*1i + 1)*1i)/2 + (1 - cot(x)^2)^(1/2)*1i)/(cot(x) + 1i))*1i

3.39 $\int \sqrt{1 - \cot^2(x)} dx$

Optimal result	260
Rubi [A] (verified)	260
Mathematica [A] (verified)	261
Maple [A] (verified)	262
Fricas [B] (verification not implemented)	262
Sympy [F]	262
Maxima [C] (verification not implemented)	263
Giac [C] (verification not implemented)	263
Mupad [B] (verification not implemented)	264

Optimal result

Integrand size = 12, antiderivative size = 32

$$\int \sqrt{1 - \cot^2(x)} dx = \arcsin(\cot(x)) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \cot(x)}{\sqrt{1 - \cot^2(x)}}\right)$$

[Out] $\arcsin(\cot(x)) - \arctan(\cot(x) * 2^{(1/2)} / (1 - \cot(x)^2)^{(1/2)}) * 2^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {3742, 399, 222, 385, 209}

$$\int \sqrt{1 - \cot^2(x)} dx = \arcsin(\cot(x)) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \cot(x)}{\sqrt{1 - \cot^2(x)}}\right)$$

[In] `Int[Sqrt[1 - Cot[x]^2], x]`

[Out] `ArcSin[Cot[x]] - Sqrt[2]*ArcTan[(Sqrt[2]*Cot[x])/Sqrt[1 - Cot[x]^2]]`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 222

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 399

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]
```

Rule 3742

```
Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{1-x^2}}{1+x^2} dx, x, \cot(x)\right) \\
&= -\left(2\text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx, x, \cot(x)\right)\right) + \text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}} dx, x, \cot(x)\right) \\
&= \arcsin(\cot(x)) - 2\text{Subst}\left(\int \frac{1}{1+2x^2} dx, x, \frac{\cot(x)}{\sqrt{1-\cot^2(x)}}\right) \\
&= \arcsin(\cot(x)) - \sqrt{2} \arctan\left(\frac{\sqrt{2} \cot(x)}{\sqrt{1-\cot^2(x)}}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.94

$$\begin{aligned}
&\int \sqrt{1-\cot^2(x)} dx \\
&= \frac{\sqrt{1-\cot^2(x)} \left(-\operatorname{arctanh}\left(\frac{\cos(x)}{\sqrt{\cos(2x)}}\right) + \sqrt{2} \log\left(\sqrt{2} \cos(x) + \sqrt{\cos(2x)}\right)\right) \sin(x)}{\sqrt{\cos(2x)}}
\end{aligned}$$

[In] Integrate[Sqrt[1 - Cot[x]^2], x]

[Out] (Sqrt[1 - Cot[x]^2]*(-ArcTanh[Cos[x]/Sqrt[Cos[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cos[x] + Sqrt[Cos[2*x]]])*Sin[x])/Sqrt[Cos[2*x]]

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.06

method	result	size
derivativedivides	$\arcsin(\cot(x)) + \sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{1-\cot(x)^2}\cot(x)}{-1+\cot(x)^2}\right)$	34
default	$\arcsin(\cot(x)) + \sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{1-\cot(x)^2}\cot(x)}{-1+\cot(x)^2}\right)$	34

[In] int((1-cot(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] arcsin(cot(x))+2^(1/2)*arctan(2^(1/2)*(1-cot(x)^2)^(1/2)/(-1+cot(x)^2)*cot(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. 2(26) = 52.

Time = 0.28 (sec) , antiderivative size = 68, normalized size of antiderivative = 2.12

$$\int \sqrt{1 - \cot^2(x)} dx = \sqrt{2} \arctan\left(\frac{\sqrt{\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x)}{\cos(2x) + 1}\right) - \arctan\left(\frac{\sqrt{2}\sqrt{\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x)}{\cos(2x) + 1}\right)$$

[In] integrate((1-cot(x)^2)^(1/2), x, algorithm="fricas")

[Out] sqrt(2)*arctan(sqrt(cos(2*x)/(cos(2*x) - 1))*sin(2*x)/(cos(2*x) + 1)) - arctan(sqrt(2)*sqrt(cos(2*x)/(cos(2*x) - 1))*sin(2*x)/(cos(2*x) + 1))

Sympy [F]

$$\int \sqrt{1 - \cot^2(x)} dx = \int \sqrt{1 - \cot^2(x)} dx$$

[In] integrate((1-cot(x)**2)**(1/2), x)

[Out] Integral(sqrt(1 - cot(x)**2), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.85 (sec) , antiderivative size = 507, normalized size of antiderivative = 15.84

$$\int \sqrt{1 - \cot^2(x)} dx = -\frac{1}{2} \sqrt{2} \left(\sqrt{2} \arctan \left(\frac{(|2e^{2ix} - 2|^4 + 16 \cos(2x)^4 + 16 \sin(2x)^4 + 8(\cos(2x)^2 - \sin(2x)^2 + 2 \cos(2x) + 1) \operatorname{abs}(2e^{2ix} - 2)^2 + 64 \cos(2x)^3 + 32(\cos(2x)^2 + 2 \cos(2x) + 1) \sin(2x)^2 + 96 \cos(2x)^2 + 64 \cos(2x) + 16)^{1/4} \sin(1/2 \arctan(8(\cos(2x) + 1) \sin(2x) / \operatorname{abs}(2e^{2ix} - 2)^2, (\operatorname{abs}(2e^{2ix} - 2)^2 + 4 \cos(2x)^2 - 4 \sin(2x)^2 + 8 \cos(2x) + 4) / \operatorname{abs}(2e^{2ix} - 2)^2)) + 2 \sin(2x)) / \operatorname{abs}(2e^{2ix} - 2), ((\operatorname{abs}(2e^{2ix} - 2)^4 + 16 \cos(2x)^4 + 16 \sin(2x)^4 + 8(\cos(2x)^2 - \sin(2x)^2 + 2 \cos(2x) + 1) \operatorname{abs}(2e^{2ix} - 2)^2 + 64 \cos(2x)^3 + 32(\cos(2x)^2 + 2 \cos(2x) + 1) \sin(2x)^2 + 96 \cos(2x)^2 + 64 \cos(2x) + 16)^{1/4} \cos(1/2 \arctan(8(\cos(2x) + 1) \sin(2x) / \operatorname{abs}(2e^{2ix} - 2)^2, (\operatorname{abs}(2e^{2ix} - 2)^2 + 4 \cos(2x)^2 - 4 \sin(2x)^2 + 8 \cos(2x) + 4) / \operatorname{abs}(2e^{2ix} - 2)^2)) + 2 \cos(2x) + 2) / \operatorname{abs}(2e^{2ix} - 2)) - \arctan(2 \arctan((\cos(4x)^2 + \sin(4x)^2 + 2 \cos(4x) + 1)^{1/4} \sin(1/2 \arctan(2 \sin(4x) / (\cos(4x) + 1)), \cos(4x) + 1)) + \sin(2x), (\cos(4x)^2 + \sin(4x)^2 + 2 \cos(4x) + 1)^{1/4} \cos(1/2 \arctan(2 \sin(4x) / (\cos(4x) + 1)), \cos(4x) + 1)) + \cos(2x)) \right)$$

[In] integrate((1-cot(x)^2)^(1/2),x, algorithm="maxima")

[Out]
$$-1/2 \sqrt{2} (\sqrt{2} \arctan(2 \arctan((\cos(4x)^2 + \sin(4x)^2 + 2 \cos(4x) + 1)^{1/4} \sin(1/2 \arctan(2 \sin(4x) / (\cos(4x) + 1)), \cos(4x) + 1)) + \sin(2x), (\cos(4x)^2 + \sin(4x)^2 + 2 \cos(4x) + 1)^{1/4} \cos(1/2 \arctan(2 \sin(4x) / (\cos(4x) + 1)), \cos(4x) + 1)) + \cos(2x)))$$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 170, normalized size of antiderivative = 5.31

$$\int \sqrt{1 - \cot^2(x)} dx = -\frac{1}{2} \left(\pi - \sqrt{2} \pi - 2 \sqrt{2} \arctan \left(-\frac{1}{2} i \sqrt{2} \right) + 2 \arctan(-i) \right) \operatorname{sgn}(\sin(x)) + \frac{1}{2} \left(\pi \operatorname{sgn}(\cos(x)) - \sqrt{2} \left(\pi \operatorname{sgn}(\cos(x)) + 2 \arctan \left(-\frac{\left(\frac{(\sqrt{2} \sqrt{-2 \cos(x)^2 + 1 - \sqrt{2}})^2}{\cos(x)^2} - 4 \right) \cos(x)}{4 \left(\sqrt{2} \sqrt{-2 \cos(x)^2 + 1 - \sqrt{2}} \right)} \right) \right) \right) + 2 a$$

[In] integrate((1-cot(x)^2)^(1/2),x, algorithm="giac")

[Out] $-1/2*(\pi - \sqrt{2}*\pi - 2*\sqrt{2}*\arctan(-1/2*I*\sqrt{2})) + 2*\arctan(-I))*\operatorname{sgn}(\sin(x)) + 1/2*(\pi*\operatorname{sgn}(\cos(x)) - \sqrt{2}*(\pi*\operatorname{sgn}(\cos(x)) + 2*\arctan(-1/4*(\sqrt{2}*\sqrt{-2*\cos(x)^2 + 1} - \sqrt{2}))^2/\cos(x)^2 - 4)*\cos(x)/(\sqrt{2}*\sqrt{-2*\cos(x)^2 + 1} - \sqrt{2}))) + 2*\arctan(-1/4*\sqrt{2}*((\sqrt{2}*\sqrt{-2*\cos(x)^2 + 1} - \sqrt{2}))^2/\cos(x)^2 - 4)*\cos(x)/(\sqrt{2}*\sqrt{-2*\cos(x)^2 + 1} - \sqrt{2})))*\operatorname{sgn}(\sin(x))$

Mupad [B] (verification not implemented)

Time = 14.19 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.75

$$\int \sqrt{1 - \cot^2(x)} dx = \operatorname{asin}(\cot(x)) - \frac{\sqrt{2} \ln \left(\frac{\frac{\sqrt{2}(-1+\cot(x)1i)1i - \sqrt{1-\cot(x)^2}1i}{2}}{\cot(x)-i} \right) 1i}{2} + \frac{\sqrt{2} \ln \left(\frac{\frac{\sqrt{2}(1+\cot(x)1i)1i + \sqrt{1-\cot(x)^2}1i}{2}}{\cot(x)+1i} \right) 1i}{2}$$

[In] int((1 - cot(x)^2)^(1/2),x)

[Out] $\operatorname{asin}(\cot(x)) - (2^{1/2}*\log(((2^{1/2}*(\cot(x)*1i - 1)*1i)/2 - (1 - \cot(x)^2)^{1/2}*1i)/(\cot(x) - 1i))*1i)/2 + (2^{1/2}*\log(((2^{1/2}*(\cot(x)*1i + 1)*1i)/2 + (1 - \cot(x)^2)^{1/2}*1i)/(\cot(x) + 1i))*1i)/2$

3.40 $\int \frac{1}{\sqrt{1-\cot^2(x)}} dx$

Optimal result	265
Rubi [A] (verified)	265
Mathematica [A] (warning: unable to verify)	266
Maple [A] (verified)	266
Fricas [B] (verification not implemented)	267
Sympy [F]	267
Maxima [B] (verification not implemented)	267
Giac [C] (verification not implemented)	268
Mupad [B] (verification not implemented)	268

Optimal result

Integrand size = 12, antiderivative size = 28

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx = -\frac{\arctan\left(\frac{\sqrt{2}\cot(x)}{\sqrt{1-\cot^2(x)}}\right)}{\sqrt{2}}$$

[Out] $-1/2*\arctan(\cot(x)*2^{(1/2)/(1-\cot(x)^2)^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3742, 385, 209}

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx = -\frac{\arctan\left(\frac{\sqrt{2}\cot(x)}{\sqrt{1-\cot^2(x)}}\right)}{\sqrt{2}}$$

[In] `Int[1/Sqrt[1 - Cot[x]^2], x]`

[Out] `-(ArcTan[(Sqrt[2]*Cot[x])/Sqrt[1 - Cot[x]^2]]/Sqrt[2])`

Rule 209

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 385

`Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b`

, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1}{\sqrt{1-x^2}(1+x^2)} dx, x, \cot(x)\right) \\ &= -\text{Subst}\left(\int \frac{1}{1+2x^2} dx, x, \frac{\cot(x)}{\sqrt{1-\cot^2(x)}}\right) \\ &= -\frac{\arctan\left(\frac{\sqrt{2}\cot(x)}{\sqrt{1-\cot^2(x)}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.93

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx = -\frac{\arcsin\left(\frac{\sqrt{2}\cot(x)}{\sqrt{1+\cot^2(x)}}\right)}{\sqrt{2}}$$

[In] Integrate[1/Sqrt[1 - Cot[x]^2], x]

[Out] -(ArcSin[(Sqrt[2]*Cot[x])/Sqrt[1 + Cot[x]^2]]/Sqrt[2])

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.11

method	result	size
derivativedivides	$\frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{1-\cot(x)^2} \cot(x)}{-1+\cot(x)^2}\right)}{2}$	31
default	$\frac{\sqrt{2} \arctan\left(\frac{\sqrt{2}\sqrt{1-\cot(x)^2} \cot(x)}{-1+\cot(x)^2}\right)}{2}$	31

[In] `int(1/(1-cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2} \sqrt{2} \arctan\left(\frac{\sqrt{2} \sqrt{1-\cot(x)^2}}{-1+\cot(x)^2} \cot(x)\right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(22) = 44$.

Time = 0.31 (sec) , antiderivative size = 56, normalized size of antiderivative = 2.00

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx = \frac{1}{4} \sqrt{2} \arctan\left(\frac{\sqrt{2}(2\sqrt{2}\cos(2x) + \sqrt{2})\sqrt{\frac{\cos(2x)}{\cos(2x)-1}}\sin(2x)}{4(\cos(2x)^2 + \cos(2x))}\right)$$

[In] `integrate(1/(1-cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{4}\sqrt{2}\arctan\left(\frac{1}{4}\sqrt{2}\arctan\left(\frac{2\sqrt{2}\cos(2x) + \sqrt{2}}{\cos(2x)-1}\right)\sqrt{\frac{\cos(2x)}{\cos(2x)-1}}\sin(2x)\right) + \sqrt{2}\sqrt{\cos(2x)}$
 $\left/\left(\cos(2x) - 1\right)\sin(2x)\left/\left(\cos(2x)^2 + \cos(2x)\right)\right)\right)$

Sympy [F]

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx = \int \frac{1}{\sqrt{1-\cot^2(x)}} dx$$

[In] `integrate(1/(1-cot(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(1 - cot(x)**2), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 90 vs. $2(22) = 44$.

Time = 0.48 (sec) , antiderivative size = 90, normalized size of antiderivative = 3.21

$$\int \frac{1}{\sqrt{1-\cot^2(x)}} dx$$

$$= \frac{1}{4} \sqrt{2} \arctan\left(\left(\cos(4x)^2 + \sin(4x)^2 + 2\cos(4x) + 1\right)^{\frac{1}{4}} \sin\left(\frac{1}{2} \arctan(\sin(4x), \cos(4x)+1)\right)\right)$$

$$+ \sin(2x), \left(\cos(4x)^2 + \sin(4x)^2 + 2\cos(4x) + 1\right)^{\frac{1}{4}} \cos\left(\frac{1}{2} \arctan(\sin(4x), \cos(4x)+1)\right)$$

$$+ \cos(2x)$$

[In] `integrate(1/(1-cot(x)^2)^(1/2),x, algorithm="maxima")`

[Out] $\frac{1}{4}\sqrt{2}\arctan2\left(\left(\cos(4x)^2 + \sin(4x)^2 + 2\cos(4x) + 1\right)^{\frac{1}{4}}\sin\left(\frac{1}{2}\arctan2(\sin(4x), \cos(4x) + 1)\right) + \sin(2x), \left(\cos(4x)^2 + \sin(4x)^2 + 2\cos(4x) + 1\right)^{\frac{1}{4}}\cos\left(\frac{1}{2}\arctan2(\sin(4x), \cos(4x) + 1)\right) + \cos(2x)\right)$

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 34, normalized size of antiderivative = 1.21

$$\int \frac{1}{\sqrt{1 - \cot^2(x)}} dx = -\frac{1}{2}i\sqrt{2} \log(i\sqrt{2} + i) \operatorname{sgn}(\sin(x)) - \frac{\sqrt{2} \arcsin(\sqrt{2} \cos(x))}{2 \operatorname{sgn}(\sin(x))}$$

[In] integrate(1/(1-cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*I*sqrt(2)*log(I*sqrt(2) + I)*sgn(sin(x)) - 1/2*sqrt(2)*arcsin(sqrt(2)*cos(x))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.18 (sec) , antiderivative size = 85, normalized size of antiderivative = 3.04

$$\int \frac{1}{\sqrt{1 - \cot^2(x)}} dx = -\frac{\sqrt{2} \ln\left(\frac{\frac{\sqrt{2}(-1+\cot(x)1i)1i}{2} - \sqrt{1-\cot(x)^2}1i}{\cot(x)-i}\right) 1i}{4} + \frac{\sqrt{2} \ln\left(\frac{\frac{\sqrt{2}(1+\cot(x)1i)1i}{2} + \sqrt{1-\cot(x)^2}1i}{\cot(x)+1i}\right) 1i}{4}$$

[In] int(1/(1 - cot(x)^2)^(1/2),x)

[Out] (2^(1/2)*log(((2^(1/2)*(cot(x)*1i + 1)*1i)/2 + (1 - cot(x)^2)^(1/2)*1i)/(cot(x) + 1i))*1i)/4 - (2^(1/2)*log(((2^(1/2)*(cot(x)*1i - 1)*1i)/2 - (1 - cot(x)^2)^(1/2)*1i)/(cot(x) - 1i))*1i)/4

3.41 $\int (-1 + \cot^2(x))^{3/2} dx$

Optimal result	269
Rubi [A] (verified)	269
Mathematica [A] (verified)	271
Maple [A] (verified)	271
Fricas [B] (verification not implemented)	272
Sympy [F]	272
Maxima [F]	272
Giac [B] (verification not implemented)	273
Mupad [F(-1)]	273

Optimal result

Integrand size = 10, antiderivative size = 61

$$\int (-1 + \cot^2(x))^{3/2} dx = \frac{5}{2} \operatorname{arctanh}\left(\frac{\cot(x)}{\sqrt{-1 + \cot^2(x)}}\right) - 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \cot(x)}{\sqrt{-1 + \cot^2(x)}}\right) - \frac{1}{2} \cot(x) \sqrt{-1 + \cot^2(x)}$$

[Out] 5/2*arctanh(cot(x)/(-1+cot(x)^2)^(1/2))-2*arctanh(cot(x)*2^(1/2)/(-1+cot(x)^2)^(1/2))*2^(1/2)-1/2*cot(x)*(-1+cot(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3742, 427, 537, 223, 212, 385}

$$\int (-1 + \cot^2(x))^{3/2} dx = \frac{5}{2} \operatorname{arctanh}\left(\frac{\cot(x)}{\sqrt{\cot^2(x) - 1}}\right) - 2\sqrt{2} \operatorname{arctanh}\left(\frac{\sqrt{2} \cot(x)}{\sqrt{\cot^2(x) - 1}}\right) - \frac{1}{2} \cot(x) \sqrt{\cot^2(x) - 1}$$

[In] Int[(-1 + Cot[x]^2)^(3/2), x]

[Out] (5*ArcTanh[Cot[x]/Sqrt[-1 + Cot[x]^2]])/2 - 2*Sqrt[2]*ArcTanh[(Sqrt[2]*Cot[x])/Sqrt[-1 + Cot[x]^2]] - (Cot[x]*Sqrt[-1 + Cot[x]^2])/2

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 427

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[d*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q - 1)/(b*(n*(p + q) + 1))),
x] + Dist[1/(b*(n*(p + q) + 1)), Int[(a + b*x^n)^p*(c + d*x^n)^(q - 2)*Simp
[c*(b*c*(n*(p + q) + 1) - a*d) + d*(b*c*(n*(p + 2*q - 1) + 1) - a*d*(n*(q -
1) + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, n, p}, x] && NeQ[b*c - a*d,
0] && GtQ[q, 1] && NeQ[n*(p + q) + 1, 0] && !IGtQ[p, 1] && IntBinomialQ[a
, b, c, d, n, p, q, x]
```

Rule 537

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*Sqrt[(c_) + (d_.)*(x
_)^(n_)], x_Symbol] := Dist[f/b, Int[1/Sqrt[c + d*x^n], x], x] + Dist[(b*e
- a*f)/b, Int[1/((a + b*x^n)*Sqrt[c + d*x^n]), x], x] /; FreeQ[{a, b, c, d
, e, f, n}, x]
```

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \frac{(-1+x^2)^{3/2}}{1+x^2} dx, x, \cot(x)\right)$$

$$\begin{aligned}
&= -\frac{1}{2} \cot(x) \sqrt{-1 + \cot^2(x)} - \frac{1}{2} \text{Subst} \left(\int \frac{3 - 5x^2}{\sqrt{-1 + x^2} (1 + x^2)} dx, x, \cot(x) \right) \\
&= -\frac{1}{2} \cot(x) \sqrt{-1 + \cot^2(x)} + \frac{5}{2} \text{Subst} \left(\int \frac{1}{\sqrt{-1 + x^2}} dx, x, \cot(x) \right) \\
&\quad - 4 \text{Subst} \left(\int \frac{1}{\sqrt{-1 + x^2} (1 + x^2)} dx, x, \cot(x) \right) \\
&= -\frac{1}{2} \cot(x) \sqrt{-1 + \cot^2(x)} + \frac{5}{2} \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, \frac{\cot(x)}{\sqrt{-1 + \cot^2(x)}} \right) \\
&\quad - 4 \text{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \frac{\cot(x)}{\sqrt{-1 + \cot^2(x)}} \right) \\
&= \frac{5}{2} \operatorname{arctanh} \left(\frac{\cot(x)}{\sqrt{-1 + \cot^2(x)}} \right) - 2\sqrt{2} \operatorname{arctanh} \left(\frac{\sqrt{2} \cot(x)}{\sqrt{-1 + \cot^2(x)}} \right) - \frac{1}{2} \cot(x) \sqrt{-1 + \cot^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 121, normalized size of antiderivative = 1.98

$$\int (-1 + \cot^2(x))^{3/2} dx = \frac{1}{2} (-1 + \cot^2(x))^{3/2} \sec^2(2x) \left(\arctan \left(\frac{\cos(x)}{\sqrt{-\cos(2x)}} \right) \sqrt{-\cos(2x)} \sin^3(x) + 4 \operatorname{arctanh} \left(\frac{\cos(x)}{\sqrt{\cos(2x)}} \right) \sqrt{\cos(2x)} \sin^3(x) - \sin(4x)/4 \right) / 2$$

[In] Integrate[(-1 + Cot[x]^2)^(3/2), x]

[Out] ((-1 + Cot[x]^2)^(3/2)*Sec[2*x]^2*(ArcTan[Cos[x]/Sqrt[-Cos[2*x]]]*Sqrt[-Cos[2*x]]*Sin[x]^3 + 4*ArcTanh[Cos[x]/Sqrt[Cos[2*x]]]*Sqrt[Cos[2*x]]*Sin[x]^3 - 4*Sqrt[2]*Sqrt[Cos[2*x]]*Log[Sqrt[2]*Cos[x] + Sqrt[Cos[2*x]]]*Sin[x]^3 - Sin[4*x]/4))/2

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.79

method	result	size
derivativedivides	$\frac{5 \ln \left(\cot(x) + \sqrt{-1 + \cot(x)^2} \right)}{2} - \frac{\cot(x) \sqrt{-1 + \cot(x)^2}}{2} - 2 \operatorname{arctanh} \left(\frac{\cot(x) \sqrt{2}}{\sqrt{-1 + \cot(x)^2}} \right) \sqrt{2}$	48
default	$\frac{5 \ln \left(\cot(x) + \sqrt{-1 + \cot(x)^2} \right)}{2} - \frac{\cot(x) \sqrt{-1 + \cot(x)^2}}{2} - 2 \operatorname{arctanh} \left(\frac{\cot(x) \sqrt{2}}{\sqrt{-1 + \cot(x)^2}} \right) \sqrt{2}$	48

[In] `int((-1+cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $5/2*\ln(\cot(x)+(-1+\cot(x)^2)^{(1/2)})-1/2*\cot(x)*(-1+\cot(x)^2)^{(1/2)}-2*\operatorname{arctanh}(\cot(x)*2^{(1/2)/(-1+\cot(x)^2)^{(1/2)}*2^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(47) = 94$.

Time = 0.28 (sec) , antiderivative size = 170, normalized size of antiderivative = 2.79

$$\int (-1 + \cot^2(x))^{3/2} dx = \frac{4\sqrt{2} \log\left(2\sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) - 2\cos(2x) - 1\right) \sin(2x) - 2\sqrt{2}\sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} (\cos(2x) + 1) \sin(2x) - 2\sqrt{2}\sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} (\cos(2x) - 1) \sin(2x) + 5 \log\left(\frac{\sqrt{2}\sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) + \cos(2x) + 1}{\cos(2x) + 1}\right) \sin(2x) - 5 \log\left(\frac{\sqrt{2}\sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) - \cos(2x) - 1}{\cos(2x) + 1}\right) \sin(2x)}{\sin(2x)}$$

[In] `integrate((-1+cot(x)^2)^(3/2),x, algorithm="fricas")`

[Out] $1/4*(4*\sqrt{2}*\log(2*\sqrt{-\cos(2*x)/(\cos(2*x)-1)}*\sin(2*x) - 2*\cos(2*x) - 1)*\sin(2*x) - 2*\sqrt{2}*\sqrt{-\cos(2*x)/(\cos(2*x)-1)}*(\cos(2*x) + 1) + 5*\log((\sqrt{2}*\sqrt{-\cos(2*x)/(\cos(2*x)-1)}*\sin(2*x) + \cos(2*x) + 1)/(\cos(2*x) + 1))*\sin(2*x) - 5*\log((\sqrt{2}*\sqrt{-\cos(2*x)/(\cos(2*x)-1)}*\sin(2*x) - \cos(2*x) - 1)/(\cos(2*x) + 1))*\sin(2*x))/\sin(2*x)$

Sympy [F]

$$\int (-1 + \cot^2(x))^{3/2} dx = \int (\cot^2(x) - 1)^{\frac{3}{2}} dx$$

[In] `integrate((-1+cot(x)**2)**(3/2),x)`

[Out] `Integral((cot(x)**2 - 1)**(3/2), x)`

Maxima [F]

$$\int (-1 + \cot^2(x))^{3/2} dx = \int (\cot(x)^2 - 1)^{\frac{3}{2}} dx$$

[In] `integrate((-1+cot(x)^2)^(3/2),x, algorithm="maxima")`

[Out] `integrate((cot(x)^2 - 1)^(3/2), x)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 179 vs. 2(47) = 94.

Time = 0.52 (sec) , antiderivative size = 179, normalized size of antiderivative = 2.93

$$\int (-1 + \cot^2(x))^{3/2} dx = \frac{1}{4} \left(4\sqrt{2} \log \left(\left(\sqrt{2} \cos(x) - \sqrt{2 \cos(x)^2 - 1} \right)^2 \right) - \frac{4\sqrt{2} \left(3 \left(\sqrt{2} \cos(x) - \sqrt{2 \cos(x)^2 - 1} \right)^4 - 6 \left(\sqrt{2} \cos(x) - \sqrt{2 \cos(x)^2 - 1} \right)^2 + 1 \right)}{\left(\sqrt{2} \cos(x) - \sqrt{2 \cos(x)^2 - 1} \right)^4} \right)$$

[In] integrate((-1+cot(x)^2)^(3/2),x, algorithm="giac")

[Out] 1/4*(4*sqrt(2)*log((sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^2) - 4*sqrt(2)*(3*(sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^2 - 1)/((sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^4 - 6*(sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^2 + 1) + 5*log(abs(2*(sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^2 - 4*sqrt(2) - 6)/abs(2*(sqrt(2)*cos(x) - sqrt(2*cos(x)^2 - 1))^2 + 4*sqrt(2) - 6)))*sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int (-1 + \cot^2(x))^{3/2} dx = \int (\cot(x)^2 - 1)^{3/2} dx$$

[In] int((cot(x)^2 - 1)^(3/2),x)

[Out] int((cot(x)^2 - 1)^(3/2), x)

3.42 $\int \sqrt{-1 + \cot^2(x)} dx$

Optimal result	274
Rubi [A] (verified)	274
Mathematica [A] (verified)	276
Maple [A] (verified)	276
Fricas [B] (verification not implemented)	276
Sympy [F]	277
Maxima [C] (verification not implemented)	277
Giac [F(-1)]	278
Mupad [B] (verification not implemented)	278

Optimal result

Integrand size = 10, antiderivative size = 42

$$\int \sqrt{-1 + \cot^2(x)} dx = -\operatorname{arctanh}\left(\frac{\cot(x)}{\sqrt{-1 + \cot^2(x)}}\right) + \sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{-1 + \cot^2(x)}}\right)$$

[Out] $-\operatorname{arctanh}(\cot(x)/(-1+\cot(x)^2)^{(1/2)})+\operatorname{arctanh}(\cot(x)*2^{(1/2)/(-1+\cot(x)^2)^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3742, 399, 223, 212, 385}

$$\int \sqrt{-1 + \cot^2(x)} dx = \sqrt{2}\operatorname{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{\cot^2(x) - 1}}\right) - \operatorname{arctanh}\left(\frac{\cot(x)}{\sqrt{\cot^2(x) - 1}}\right)$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[-1 + \operatorname{Cot}[x]^2], x]$

[Out] $-\operatorname{ArcTanh}[\operatorname{Cot}[x]/\operatorname{Sqrt}[-1 + \operatorname{Cot}[x]^2]] + \operatorname{Sqrt}[2]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[2]*\operatorname{Cot}[x])/\operatorname{Sqrt}[-1 + \operatorname{Cot}[x]^2]]$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 385

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 399

Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[b/d, Int[(a + b*x^n)^(p - 1), x], x] - Dist[(b*c - a*d)/d, Int[(a + b*x^n)^(p - 1)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b*c - a*d, 0] && EqQ[n*(p - 1) + 1, 0] && IntegerQ[n]

Rule 3742

Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || EqQ[n^2, 16])

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{\sqrt{-1+x^2}}{1+x^2} dx, x, \cot(x)\right) \\
 &= 2\text{Subst}\left(\int \frac{1}{\sqrt{-1+x^2}(1+x^2)} dx, x, \cot(x)\right) - \text{Subst}\left(\int \frac{1}{\sqrt{-1+x^2}} dx, x, \cot(x)\right) \\
 &= 2\text{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \frac{\cot(x)}{\sqrt{-1+\cot^2(x)}}\right) - \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\cot(x)}{\sqrt{-1+\cot^2(x)}}\right) \\
 &= -\text{arctanh}\left(\frac{\cot(x)}{\sqrt{-1+\cot^2(x)}}\right) + \sqrt{2}\text{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{-1+\cot^2(x)}}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.43

$$\int \sqrt{-1 + \cot^2(x)} dx = \frac{\sqrt{-1 + \cot^2(x)} \left(-\operatorname{arctanh} \left(\frac{\cos(x)}{\sqrt{\cos(2x)}} \right) + \sqrt{2} \log \left(\sqrt{2} \cos(x) + \sqrt{\cos(2x)} \right) \right) \sin(x)}{\sqrt{\cos(2x)}}$$

[In] Integrate[Sqrt[-1 + Cot[x]^2], x]

[Out] (Sqrt[-1 + Cot[x]^2]*(-ArcTanh[Cos[x]/Sqrt[Cos[2*x]]] + Sqrt[2]*Log[Sqrt[2]*Cos[x] + Sqrt[Cos[2*x]]])*Sin[x])/Sqrt[Cos[2*x]]

Maple [A] (verified)

Time = 0.03 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\ln \left(\cot(x) + \sqrt{-1 + \cot(x)^2} \right) + \operatorname{arctanh} \left(\frac{\cot(x)\sqrt{2}}{\sqrt{-1 + \cot(x)^2}} \right) \sqrt{2}$	35
default	$-\ln \left(\cot(x) + \sqrt{-1 + \cot(x)^2} \right) + \operatorname{arctanh} \left(\frac{\cot(x)\sqrt{2}}{\sqrt{-1 + \cot(x)^2}} \right) \sqrt{2}$	35

[In] int((-1+cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -ln(cot(x)+(-1+cot(x)^2)^(1/2))+arctanh(cot(x)*2^(1/2)/(-1+cot(x)^2)^(1/2))*2^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 123 vs. 2(34) = 68.

Time = 0.29 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.93

$$\int \sqrt{-1 + \cot^2(x)} dx = \frac{1}{2} \sqrt{2} \log \left(-2 \sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) - 2 \cos(2x) - 1 \right) - \frac{1}{2} \log \left(\frac{\sqrt{2} \sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) + \cos(2x) + 1}{\cos(2x) + 1} \right) + \frac{1}{2} \log \left(\frac{\sqrt{2} \sqrt{-\frac{\cos(2x)}{\cos(2x)-1}} \sin(2x) - \cos(2x) - 1}{\cos(2x) + 1} \right)$$

[In] integrate((-1+cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/2*sqrt(2)*log(-2*sqrt(-cos(2*x)/(cos(2*x) - 1))*sin(2*x) - 2*cos(2*x) - 1) - 1/2*log((sqrt(2)*sqrt(-cos(2*x)/(cos(2*x) - 1))*sin(2*x) + cos(2*x) + 1)/(cos(2*x) + 1)) + 1/2*log((sqrt(2)*sqrt(-cos(2*x)/(cos(2*x) - 1))*sin(2*x) - cos(2*x) - 1)/(cos(2*x) + 1))

Sympy [F]

$$\int \sqrt{-1 + \cot^2(x)} dx = \int \sqrt{\cot^2(x) - 1} dx$$

[In] integrate((-1+cot(x)**2)**(1/2),x)

[Out] Integral(sqrt(cot(x)**2 - 1), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.69 (sec) , antiderivative size = 941, normalized size of antiderivative = 22.40

$$\int \sqrt{-1 + \cot^2(x)} dx = \text{Too large to display}$$

[In] integrate((-1+cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(2)*arcsinh(1) + 1/4*sqrt(2)*log(cos(2*x)^2 + sin(2*x)^2 + sqrt(cos(4*x)^2 + sin(4*x)^2 + 2*cos(4*x) + 1)*(cos(1/2*arctan2(sin(4*x), cos(4*x) + 1))^2 + sin(1/2*arctan2(sin(4*x), cos(4*x) + 1))^2) + 2*(cos(4*x)^2 + sin(4*x)^2 + 2*cos(4*x) + 1)^(1/4)*(cos(2*x)*cos(1/2*arctan2(sin(4*x), cos(4*x) + 1)) + sin(2*x)*sin(1/2*arctan2(sin(4*x), cos(4*x) + 1)))) - 1/2*log((sqrt(abs(2*e^(2*I*x) - 2))^4 + 16*cos(2*x)^4 + 16*sin(2*x)^4 + 8*(cos(2*x)^2 - sin(2*x)^2 + 2*cos(2*x) + 1)*abs(2*e^(2*I*x) - 2)^2 + 64*cos(2*x)^3 + 32*(cos(2*x)^2 + 2*cos(2*x) + 1)*sin(2*x)^2 + 96*cos(2*x)^2 + 64*cos(2*x) + 16)*cos(1/2*arctan2(8*(cos(2*x) + 1)*sin(2*x)/abs(2*e^(2*I*x) - 2)^2, (abs(2*e^(2*I*x) - 2)^2 + 4*cos(2*x)^2 - 4*sin(2*x)^2 + 8*cos(2*x) + 4)/abs(2*e^(2*I*x) - 2)^2))^2 + sqrt(abs(2*e^(2*I*x) - 2))^4 + 16*cos(2*x)^4 + 16*sin(2*x)^4 + 8*(cos(2*x)^2 - sin(2*x)^2 + 2*cos(2*x) + 1)*abs(2*e^(2*I*x) - 2)^2 + 64*cos(2*x)^3 + 32*(cos(2*x)^2 + 2*cos(2*x) + 1)*sin(2*x)^2 + 96*cos(2*x)^2 + 64*cos(2*x) + 16)*sin(1/2*arctan2(8*(cos(2*x) + 1)*sin(2*x)/abs(2*e^(2*I*x) - 2)^2, (abs(2*e^(2*I*x) - 2)^2 + 4*cos(2*x)^2 - 4*sin(2*x)^2 + 8*cos(2*x) + 4)/abs(2*e^(2*I*x) - 2)^2))^2 + 4*(abs(2*e^(2*I*x) - 2))^4 + 16*cos(2*x)^4 + 16*sin(2*x)^4 + 8*(cos(2*x)^2 - sin(2*x)^2 + 2*cos(2*x) + 1)*abs(2*e^(2*I*x) - 2)^2 + 64*cos(2*x)^3 + 32*(cos(2*x)^2 + 2*cos(2*x) + 1)*sin(2*x)

$$\begin{aligned} &^2 + 96\cos(2x)^2 + 64\cos(2x) + 16)^{1/4}(\cos(2x) + 1)\cos(1/2\arctan2 \\ &(8(\cos(2x) + 1)\sin(2x)/\operatorname{abs}(2e^{2Ix} - 2)^2, (\operatorname{abs}(2e^{2Ix} - 2)^2 \\ &+ 4\cos(2x)^2 - 4\sin(2x)^2 + 8\cos(2x) + 4)/\operatorname{abs}(2e^{2Ix} - 2)^2)) + \\ &4(\operatorname{abs}(2e^{2Ix} - 2)^4 + 16\cos(2x)^4 + 16\sin(2x)^4 + 8(\cos(2x)^2 - \\ &\sin(2x)^2 + 2\cos(2x) + 1)\operatorname{abs}(2e^{2Ix} - 2)^2 + 64\cos(2x)^3 + 32(\cos(2x)^2 \\ &+ 2\cos(2x) + 1)\sin(2x)^2 + 96\cos(2x)^2 + 64\cos(2x) + 16) \\ &^{1/4}\sin(2x)\sin(1/2\arctan2(8(\cos(2x) + 1)\sin(2x)/\operatorname{abs}(2e^{2Ix} - 2)^2, \\ &(\operatorname{abs}(2e^{2Ix} - 2)^2 + 4\cos(2x)^2 - 4\sin(2x)^2 + 8\cos(2x) + 4)/\operatorname{abs}(2e^{2Ix} - 2)^2)) \\ &+ 4\cos(2x)^2 + 4\sin(2x)^2 + 8\cos(2x) + 4)/\operatorname{abs}(2e^{2Ix} - 2)^2 \end{aligned}$$

Giac [F(-1)]

Timed out.

$$\int \sqrt{-1 + \cot^2(x)} dx = \text{Timed out}$$

[In] integrate((-1+cot(x)^2)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [B] (verification not implemented)

Time = 13.92 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.81

$$\int \sqrt{-1 + \cot^2(x)} dx = \sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \cot(x)}{\sqrt{\cot(x)^2 - 1}}\right) - \ln\left(\cot(x) + \sqrt{\cot(x)^2 - 1}\right)$$

[In] int((cot(x)^2 - 1)^(1/2),x)

[Out] 2^(1/2)*atanh((2^(1/2)*cot(x))/(cot(x)^2 - 1)^(1/2)) - log(cot(x) + (cot(x)^2 - 1)^(1/2))

3.43 $\int \frac{1}{\sqrt{-1+\cot^2(x)}} dx$

Optimal result	279
Rubi [A] (verified)	279
Mathematica [A] (warning: unable to verify)	280
Maple [A] (verified)	280
Fricas [B] (verification not implemented)	281
Sympy [F]	281
Maxima [B] (verification not implemented)	281
Giac [B] (verification not implemented)	282
Mupad [B] (verification not implemented)	282

Optimal result

Integrand size = 10, antiderivative size = 26

$$\int \frac{1}{\sqrt{-1+\cot^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{-1+\cot^2(x)}}\right)}{\sqrt{2}}$$

[Out] $-1/2*\operatorname{arctanh}(\cot(x)*2^{(1/2)/(-1+\cot(x)^2)^{(1/2)})*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3742, 385, 212}

$$\int \frac{1}{\sqrt{-1+\cot^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{\cot^2(x)-1}}\right)}{\sqrt{2}}$$

[In] $\operatorname{Int}[1/\operatorname{Sqrt}[-1 + \operatorname{Cot}[x]^2], x]$

[Out] $-(\operatorname{ArcTanh}[(\operatorname{Sqrt}[2]*\operatorname{Cot}[x])/\operatorname{Sqrt}[-1 + \operatorname{Cot}[x]^2]])/\operatorname{Sqrt}[2]$

Rule 212

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 385

$\operatorname{Int}[(a + (b \cdot x)^n)^p / ((c + (d \cdot x)^n)), x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(c - (b \cdot c - a \cdot d) \cdot x^n), x], x, x/(a + b \cdot x^n)^{(1/n)}] /; \operatorname{FreeQ}\{a, b$

, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]

Rule 3742

```
Int[((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegersQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1}{\sqrt{-1+x^2}(1+x^2)} dx, x, \cot(x)\right) \\ &= -\text{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \frac{\cot(x)}{\sqrt{-1+\cot^2(x)}}\right) \\ &= -\frac{\text{arctanh}\left(\frac{\sqrt{2}\cot(x)}{\sqrt{-1+\cot^2(x)}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 0.07 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.85

$$\int \frac{1}{\sqrt{-1+\cot^2(x)}} dx = -\frac{\arcsin\left(\frac{\sqrt{2}\cot(x)}{\sqrt{1+\cot^2(x)}}\right)\sqrt{1-\cot^2(x)}}{\sqrt{2}\sqrt{-1+\cot^2(x)}}$$

[In] Integrate[1/Sqrt[-1 + Cot[x]^2],x]

[Out] -((ArcSin[(Sqrt[2]*Cot[x])/Sqrt[1 + Cot[x]^2]]*Sqrt[1 - Cot[x]^2])/(Sqrt[2]*Sqrt[-1 + Cot[x]^2]))

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$-\frac{\text{arctanh}\left(\frac{\cot(x)\sqrt{2}}{\sqrt{-1+\cot(x)^2}}\right)\sqrt{2}}{2}$	21
default	$-\frac{\text{arctanh}\left(\frac{\cot(x)\sqrt{2}}{\sqrt{-1+\cot(x)^2}}\right)\sqrt{2}}{2}$	21

[In] `int(1/(-1+cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-1/2*\operatorname{arctanh}(\cot(x)*2^{(1/2)/(-1+cot(x)^2)^{(1/2)})*2^{(1/2)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(20) = 40$.

Time = 0.30 (sec) , antiderivative size = 60, normalized size of antiderivative = 2.31

$$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx = \frac{1}{8} \sqrt{2} \log \left(2 \sqrt{2} \left(2 \sqrt{2} \cos(2x) + \sqrt{2} \right) \sqrt{-\frac{\cos(2x)}{\cos(2x) - 1}} \sin(2x) - 8 \cos(2x)^2 - 8 \cos(2x) - 1 \right)$$

[In] `integrate(1/(-1+cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] $1/8*\sqrt{2}*\log(2*\sqrt{2}*(2*\sqrt{2}*\cos(2*x) + \sqrt{2})*\sqrt{-\cos(2*x)/(\cos(2*x) - 1)}*\sin(2*x) - 8*\cos(2*x)^2 - 8*\cos(2*x) - 1)$

Sympy [F]

$$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx = \int \frac{1}{\sqrt{\cot^2(x) - 1}} dx$$

[In] `integrate(1/(-1+cot(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(cot(x)**2 - 1), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 143 vs. $2(20) = 40$.

Time = 0.53 (sec) , antiderivative size = 143, normalized size of antiderivative = 5.50

$$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx = -\frac{1}{8} \sqrt{2} \left(2 \operatorname{arsinh}(1) + \log \left(\cos(2x)^2 + \sin(2x)^2 + \sqrt{\cos(4x)^2 + \sin(4x)^2 + 2 \cos(4x) + 1} \left(\cos \left(\frac{1}{2} \arctan 2 \left(\frac{\sin(4x)}{\cos(4x) + 1} \right) \right) \right) \right) \right)$$

[In] `integrate(1/(-1+cot(x)^2)^(1/2),x, algorithm="maxima")`

[Out] $-1/8*\sqrt{2}*(2*\operatorname{arcsinh}(1) + \log(\cos(2*x)^2 + \sin(2*x)^2 + \sqrt{\cos(4*x)^2 + \sin(4*x)^2 + 2*\cos(4*x) + 1}*(\cos(1/2*\arctan 2(\sin(4*x)/(\cos(4*x) + 1)))^2 + \sin(1/2*\arctan 2(\sin(4*x)/(\cos(4*x) + 1)))^2) + 2*(\cos(4*x)^2 + \sin(4*x)^2 + 2*\cos(4*x) + 1)^{(1/4)}*(\cos(2*x)*\cos(1/2*\arctan 2(\sin(4*x)/(\cos(4*x) + 1))) + \sin(2*x)*\sin(1/2*\arctan 2(\sin(4*x)/(\cos(4*x) + 1))))$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(20) = 40.

Time = 0.30 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.73

$$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx = -\frac{1}{2} \sqrt{2} \log(\sqrt{2} - 1) \operatorname{sgn}(\sin(x)) + \frac{\sqrt{2} \log\left(\left| -\sqrt{2} \cos(x) + \sqrt{2 \cos(x)^2 - 1} \right| \right)}{2 \operatorname{sgn}(\sin(x))}$$

[In] integrate(1/(-1+cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*sqrt(2)*log(sqrt(2) - 1)*sgn(sin(x)) + 1/2*sqrt(2)*log(abs(-sqrt(2)*cos(x) + sqrt(2*cos(x)^2 - 1)))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.91 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.77

$$\int \frac{1}{\sqrt{-1 + \cot^2(x)}} dx = -\frac{\sqrt{2} \operatorname{atanh}\left(\frac{\sqrt{2} \cot(x)}{\sqrt{\cot(x)^2 - 1}}\right)}{2}$$

[In] int(1/(cot(x)^2 - 1)^(1/2),x)

[Out] -(2^(1/2)*atanh((2^(1/2)*cot(x))/(cot(x)^2 - 1)^(1/2)))/2

3.44 $\int \frac{\cot^3(x)}{\sqrt{a+b \cot^2(x)}} dx$

Optimal result	283
Rubi [A] (verified)	283
Mathematica [A] (verified)	285
Maple [A] (verified)	285
Fricas [B] (verification not implemented)	286
Sympy [F]	286
Maxima [F(-2)]	287
Giac [B] (verification not implemented)	287
Mupad [B] (verification not implemented)	287

Optimal result

Integrand size = 17, antiderivative size = 52

$$\int \frac{\cot^3(x)}{\sqrt{a+b \cot^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}} - \frac{\sqrt{a+b \cot^2(x)}}{b}$$

[Out] $-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)/(a-b)^{(1/2)})/(a-b)^{(1/2)}-(a+b*\cot(x)^2)^{(1/2)})/b$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 457, 81, 65, 214}

$$\int \frac{\cot^3(x)}{\sqrt{a+b \cot^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}} - \frac{\sqrt{a+b \cot^2(x)}}{b}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]^3/\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2], x]$

[Out] $-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a - b]]/\operatorname{Sqrt}[a - b]) - \operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/b$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^{n}, x], x, (a+b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 457

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{x^3}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{x}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)\right) \\
 &= -\frac{\sqrt{a+b\cot^2(x)}}{b} + \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
 &= -\frac{\sqrt{a+b\cot^2(x)}}{b} + \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
 &= -\frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}} - \frac{\sqrt{a+b\cot^2(x)}}{b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.00

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a + b \cot^2(x)}}{\sqrt{a - b}}\right)}{\sqrt{a - b}} + \sqrt{a + b \cot^2(x)}$$

[In] Integrate[Cot[x]^3/Sqrt[a + b*Cot[x]^2],x]

[Out] -(((b*ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]])/Sqrt[a - b] + Sqrt[a + b*Cot[x]^2])/b)

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.85

method	result	size
derivativedivides	$-\frac{\sqrt{a + b \cot(x)^2}}{b} + \frac{\arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}}$	44
default	$-\frac{\sqrt{a + b \cot(x)^2}}{b} + \frac{\arctan\left(\frac{\sqrt{a + b \cot(x)^2}}{\sqrt{-a + b}}\right)}{\sqrt{-a + b}}$	44

[In] int(cot(x)^3/(a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -(a+b*cot(x)^2)^(1/2)/b+1/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 118 vs. 2(44) = 88.

Time = 0.35 (sec) , antiderivative size = 284, normalized size of antiderivative = 5.46

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \left[\frac{\sqrt{a - b} \log \left(-2(a^2 - 2ab + b^2) \cos(2x)^2 - 2a^2 + b^2 + 2((a - b) \cos(2x))^2 - (2a - b) \cos(2x) + a \right) \sqrt{a - b}}{4(ab - b^2)} \right. \\ \left. - \frac{\sqrt{-a + b} \arctan \left(-\frac{\sqrt{-a + b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) - 1)}{(a - b) \cos(2x) - a} \right) + 2(a - b) \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}}}{2(ab - b^2)} \right]$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(a - b)*b*log(-2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 - 2*a^2 + b^2 + 2*((a - b)*cos(2*x)^2 - (2*a - b)*cos(2*x) + a)*sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)) + 4*(a^2 - a*b)*cos(2*x)) - 4*(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a*b - b^2), -1/2*(sqrt(-a + b)*b*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1)/((a - b)*cos(2*x) - a)) + 2*(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a*b - b^2)]

Sympy [F]

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx$$

[In] integrate(cot(x)**3/(a+b*cot(x)**2)**(1/2),x)

[Out] Integral(cot(x)**3/sqrt(a + b*cot(x)**2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs. 2(44) = 88.

Time = 0.36 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.85

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\log\left(\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2\right)}{\sqrt{a-b}} + \frac{4\sqrt{a-b}}{\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2 - b}$$

$$= \frac{\log\left(\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2\right)}{2 \operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*(log((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2)/sqrt(a - b) + 4*sqrt(a - b)/((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2 - b))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 14.61 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.85

$$\int \frac{\cot^3(x)}{\sqrt{a + b \cot^2(x)}} dx = -\frac{\sqrt{b \cot^2(x) + a}}{b} - \frac{\operatorname{atanh}\left(\frac{\sqrt{b \cot^2(x) + a}}{\sqrt{a - b}}\right)}{\sqrt{a - b}}$$

[In] int(cot(x)^3/(a + b*cot(x)^2)^(1/2),x)

[Out] -(a + b*cot(x)^2)^(1/2)/b - atanh((a + b*cot(x)^2)^(1/2)/(a - b)^(1/2))/(a - b)^(1/2)

3.45 $\int \frac{\cot^2(x)}{\sqrt{a+b \cot^2(x)}} dx$

Optimal result	288
Rubi [A] (verified)	288
Mathematica [B] (verified)	290
Maple [A] (verified)	290
Fricas [B] (verification not implemented)	291
Sympy [F]	291
Maxima [F]	292
Giac [B] (verification not implemented)	292
Mupad [F(-1)]	292

Optimal result

Integrand size = 17, antiderivative size = 64

$$\int \frac{\cot^2(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{a-b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{b}}$$

[Out] $\arctan(\cot(x) \cdot (a-b)^{(1/2)} / (a+b \cdot \cot(x)^2)^{(1/2)}) / (a-b)^{(1/2)} - \operatorname{arctanh}(\cot(x) \cdot b^{(1/2)} / (a+b \cdot \cot(x)^2)^{(1/2)}) / b^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 494, 223, 212, 385, 209}

$$\int \frac{\cot^2(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{a-b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{b}}$$

[In] $\text{Int}[\text{Cot}[x]^2/\text{Sqrt}[a + b \cdot \text{Cot}[x]^2], x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a - b] \cdot \text{Cot}[x]) / \text{Sqrt}[a + b \cdot \text{Cot}[x]^2]] / \text{Sqrt}[a - b] - \text{ArcTanh}[(\text{Sqrt}[b] \cdot \text{Cot}[x]) / \text{Sqrt}[a + b \cdot \text{Cot}[x]^2]] / \text{Sqrt}[b]$

Rule 209

$\text{Int}[(a_ + (b_ \cdot x)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \cdot \text{Rt}[b, 2])) \cdot \text{ArcTan}[\text{Rt}[b, 2] \cdot (x/\text{Rt}[a, 2])], x] / ; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 223

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x],
x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]
```

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 494

```
Int[(((e_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_))^(q_))/((a_) + (b_)*(x_)^(
n_)), x_Symbol] := Dist[e^n/b, Int[(e*x)^(m - n)*(c + d*x^n)^q, x], x] - Di
st[a*(e^n/b), Int[(e*x)^(m - n)*((c + d*x^n)^q/(a + b*x^n)), x], x] /; Free
Q[{a, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LeQ[n, m,
2*n - 1] && IntBinomialQ[a, b, c, d, e, m, n, -1, q, x]
```

Rule 3751

```
Int[(((d_)*tan[(e_) + (f_)*(x_)])^(m_))*((a_) + (b_))*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= -\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot(x)\right) + \text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
&= -\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
&\quad + \text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)
\end{aligned}$$

$$= \frac{\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b}\cot^2(x)}\right)}{\sqrt{a-b}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{b}\cot(x)}{\sqrt{a+b}\cot^2(x)}\right)}{\sqrt{b}}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 158 vs. $2(64) = 128$.

Time = 0.35 (sec) , antiderivative size = 158, normalized size of antiderivative = 2.47

$$\int \frac{\cot^2(x)}{\sqrt{a+b}\cot^2(x)} dx$$

$$= \frac{\left(-\sqrt{-b}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{a-b}\cos(x)}{\sqrt{-a-b+(a-b)}\cos(2x)}\right) + \sqrt{a-b}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{-b}\cos(x)}{\sqrt{-a-b+(a-b)}\cos(2x)}\right)\right) \sqrt{(a+b+(-a+b)\cos(2x))}}{\sqrt{a-b}\sqrt{-b}\sqrt{-a-b+(a-b)\cos(2x)}}$$

[In] Integrate[Cot[x]^2/Sqrt[a + b*Cot[x]^2], x]

[Out] ((- (Sqrt[-b]*ArcTanh[(Sqrt[2]*Sqrt[a - b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]])] + Sqrt[a - b]*ArcTanh[(Sqrt[2]*Sqrt[-b]*Cos[x])/Sqrt[-a - b + (a - b)*Cos[2*x]])]*Sqrt[(a + b + (-a + b)*Cos[2*x])*Csc[x]^2]*Sin[x])/(Sqrt[a - b]*Sqrt[-b]*Sqrt[-a - b + (a - b)*Cos[2*x]])

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.25

method	result	size
derivativedivides	$-\frac{\ln\left(\sqrt{b}\cot(x)+\sqrt{a+b}\cot^2(x)\right)}{\sqrt{b}} + \frac{\sqrt{b^4(a-b)}\arctan\left(\frac{b^2(a-b)\cot(x)}{\sqrt{b^4(a-b)}\sqrt{a+b}\cot^2(x)}\right)}{b^2(a-b)}$	80
default	$-\frac{\ln\left(\sqrt{b}\cot(x)+\sqrt{a+b}\cot^2(x)\right)}{\sqrt{b}} + \frac{\sqrt{b^4(a-b)}\arctan\left(\frac{b^2(a-b)\cot(x)}{\sqrt{b^4(a-b)}\sqrt{a+b}\cot^2(x)}\right)}{b^2(a-b)}$	80

[In] int(cot(x)^2/(a+b*cot(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] $-\ln(b^{(1/2)}*\cot(x)+(a+b*\cot(x)^2)^{(1/2)})/b^{(1/2)}+(b^4*(a-b))^{(1/2)}/b^2/(a-b)*\arctan(b^2*(a-b)/(b^4*(a-b))^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)}*\cot(x))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(52) = 104.

Time = 0.32 (sec) , antiderivative size = 588, normalized size of antiderivative = 9.19

$$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \left[\frac{\sqrt{-a + b} \log\left(- (a - b) \cos(2x) + \sqrt{-a + b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x) + b\right) - (a - b) \sqrt{b} \log\left(\frac{(a-2b) \cos(2x) - a - b}{(a-b) \cos(2x) + a - b}\right)}{2(ab - b^2)} \right]$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")

[Out] [-1/2*(sqrt(-a + b)*b*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b) - (a - b)*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1)))/(a*b - b^2), 1/2*(2*(a - b)*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b)) - sqrt(-a + b)*b*log(-(a - b)*cos(2*x) + sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + b))/(a*b - b^2), 1/2*(2*sqrt(a - b)*b*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b)) + (a - b)*sqrt(b)*log(((a - 2*b)*cos(2*x) + 2*sqrt(b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) - a - 2*b)/(cos(2*x) - 1)))/(a*b - b^2), (sqrt(a - b)*b*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) + a - b)) + (a - b)*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/(b*cos(2*x) + b)))/(a*b - b^2)]

Sympy [F]

$$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

[In] integrate(cot(x)**2/(a+b*cot(x)**2)**(1/2),x)

[Out] Integral(cot(x)**2/sqrt(a + b*cot(x)**2), x)

Maxima [F]

$$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\cot(x)^2}{\sqrt{b \cot(x)^2 + a}} dx$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(cot(x)^2/sqrt(b*cot(x)^2 + a), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 229 vs. 2(52) = 104.

Time = 0.54 (sec) , antiderivative size = 229, normalized size of antiderivative = 3.58

$$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\left(2a \arctan\left(\frac{\sqrt{-a+b}\sqrt{b}}{\sqrt{ab-b^2}}\right) - 2b \arctan\left(\frac{\sqrt{-a+b}\sqrt{b}}{\sqrt{ab-b^2}}\right) + \sqrt{ab-b^2} \log\left(-a - 2\sqrt{-a+b}\sqrt{b} + 2b\right)\right) \operatorname{sgn}(\sin(x))}{2\sqrt{ab-b^2}\sqrt{-a+b}} + \frac{2\sqrt{-a+b} \arctan\left(\frac{\left(\sqrt{-a+b}\cos(x) - \sqrt{-a\cos(x)^2 + b\cos(x)^2 + a}\right)^2 + a - 2b}{2\sqrt{ab-b^2}}\right)}{\sqrt{ab-b^2}} + \frac{\log\left(\left(\sqrt{-a+b}\cos(x) - \sqrt{-a\cos(x)^2 + b\cos(x)^2 + a}\right)^2\right)}{\sqrt{-a+b}}$$

$$= \frac{\left(2a \arctan\left(\frac{\sqrt{-a+b}\sqrt{b}}{\sqrt{ab-b^2}}\right) - 2b \arctan\left(\frac{\sqrt{-a+b}\sqrt{b}}{\sqrt{ab-b^2}}\right) + \sqrt{ab-b^2} \log\left(-a - 2\sqrt{-a+b}\sqrt{b} + 2b\right)\right) \operatorname{sgn}(\sin(x))}{2\sqrt{ab-b^2}\sqrt{-a+b}} + \frac{2\sqrt{-a+b} \arctan\left(\frac{\left(\sqrt{-a+b}\cos(x) - \sqrt{-a\cos(x)^2 + b\cos(x)^2 + a}\right)^2 + a - 2b}{2\sqrt{ab-b^2}}\right)}{\sqrt{ab-b^2}} + \frac{\log\left(\left(\sqrt{-a+b}\cos(x) - \sqrt{-a\cos(x)^2 + b\cos(x)^2 + a}\right)^2\right)}{\sqrt{-a+b}}$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*(2*a*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) - 2*b*arctan(sqrt(-a + b)*sqrt(b)/sqrt(a*b - b^2)) + sqrt(a*b - b^2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b))*sgn(sin(x))/(sqrt(a*b - b^2)*sqrt(-a + b)) - 1/2*(2*sqrt(-a + b)*arctan(1/2*((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 + a - 2*b)/sqrt(a*b - b^2))/sqrt(a*b - b^2) + log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2)/sqrt(-a + b))/sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\cot(x)^2}{\sqrt{b \cot(x)^2 + a}} dx$$

[In] int(cot(x)^2/(a + b*cot(x)^2)^(1/2),x)

[Out] int(cot(x)^2/(a + b*cot(x)^2)^(1/2), x)

$$3.46 \quad \int \frac{\cot(x)}{\sqrt{a+b \cot^2(x)}} dx$$

Optimal result	293
Rubi [A] (verified)	293
Mathematica [A] (verified)	294
Maple [A] (verified)	295
Fricas [B] (verification not implemented)	295
Sympy [F]	295
Maxima [F(-2)]	296
Giac [B] (verification not implemented)	296
Mupad [B] (verification not implemented)	296

Optimal result

Integrand size = 15, antiderivative size = 33

$$\int \frac{\cot(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)/(a-b)^{(1/2)})/(a-b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3751, 455, 65, 214}

$$\int \frac{\cot(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]/\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2], x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a - b]]/\operatorname{Sqrt}[a - b]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)\right) \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
 &= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}} dx = \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

```
[In] Integrate[Cot[x]/Sqrt[a + b*Cot[x]^2], x]
```

```
[Out] ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]]/Sqrt[a - b]
```

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.88

method	result	size
derivativedivides	$-\frac{\arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$	29
default	$-\frac{\arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{\sqrt{-a+b}}$	29

[In] `int(cot(x)/(a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. 2(27) = 54.

Time = 0.29 (sec) , antiderivative size = 127, normalized size of antiderivative = 3.85

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \left[\frac{\log\left(-\sqrt{a-b} \sqrt{\frac{(a-b)\cos(2x)-a-b}{\cos(2x)-1}} (\cos(2x)-1) - (a-b)\cos(2x) + a\right)}{2\sqrt{a-b}}, \frac{\sqrt{-a+b} \arctan\left(-\frac{\sqrt{-a+b} \sqrt{\frac{(a-b)\cos(2x)-a-b}{\cos(2x)-1}}}{a-b}\right)}{a-b} \right]$$

[In] `integrate(cot(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `[1/2*log(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1) - (a - b)*cos(2*x) + a)/sqrt(a - b), sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))/(a - b))/(a - b)]`

Sympy [F]

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx$$

[In] `integrate(cot(x)/(a+b*cot(x)**2)**(1/2),x)`

[Out] `Integral(cot(x)/sqrt(a + b*cot(x)**2), x)`

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(27) = 54.

Time = 0.33 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.85

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\log(|b| \operatorname{sgn}(\sin(x)))}{2\sqrt{a-b}} - \frac{\log\left(\left|-\sqrt{a-b}\sin(x) + \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right|\right)}{\sqrt{a-b}\operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*log(abs(b))*sgn(sin(x))/sqrt(a - b) - log(abs(-sqrt(a - b)*sin(x) + sqrt(a*sin(x)^2 - b*sin(x)^2 + b)))/(sqrt(a - b)*sgn(sin(x)))

Mupad [B] (verification not implemented)

Time = 13.51 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.82

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \cot^2(x) + a}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[In] int(cot(x)/(a + b*cot(x)^2)^(1/2),x)

[Out] atanh((a + b*cot(x)^2)^(1/2)/(a - b)^(1/2))/(a - b)^(1/2)

$$3.47 \quad \int \frac{\tan(x)}{\sqrt{a+b \cot^2(x)}} dx$$

Optimal result	297
Rubi [A] (verified)	297
Mathematica [A] (verified)	299
Maple [B] (verified)	299
Fricas [A] (verification not implemented)	300
Sympy [F]	301
Maxima [F]	301
Giac [B] (verification not implemented)	301
Mupad [B] (verification not implemented)	302

Optimal result

Integrand size = 15, antiderivative size = 60

$$\int \frac{\tan(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/a^{(1/2)})/a^{(1/2)}-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 457, 88, 65, 214}

$$\int \frac{\tan(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]/\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2], x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a]]/\operatorname{Sqrt}[a] - \operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a - b]]/\operatorname{Sqrt}[a - b]$

Rule 65

$\operatorname{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 88

Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Dist[b/(b*c - a*d), Int[(e + f*x)^p/(a + b*x), x], x] - Dist[d/
(b*c - a*d), Int[(e + f*x)^p/(c + d*x), x], x] /; FreeQ[{a, b, c, d, e, f,
p}, x] && !IntegerQ[p]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 457

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
) , x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n,
p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{1}{x(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{x(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cot^2(x)\right)\right) + \frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right) \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{b} \\
 &= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{\sqrt{a-b}}$$

[In] Integrate[Tan[x]/Sqrt[a + b*Cot[x]^2],x]

[Out] ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a]]/Sqrt[a] - ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]]/Sqrt[a - b]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(48) = 96.

Time = 0.80 (sec) , antiderivative size = 149, normalized size of antiderivative = 2.48

method	result
default	$\frac{\sqrt{4} \left(\arctan \left(\frac{\sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} (\cot(x) + \csc(x))}{\sqrt{-a+b}} \right) \sqrt{a} + \operatorname{arctanh} \left(\frac{\sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} (\cot(x) + \csc(x))}{\sqrt{a}} \right) \sqrt{-a+b} \right) \sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}}}{2\sqrt{-a+b} \sqrt{a} \sqrt{a+b \cot(x)^2}}$

[In] int(tan(x)/(a+b*cot(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2*4^(1/2)/(-a+b)^(1/2)/a^(1/2)*(arctan(1/(-a+b)^(1/2)*(-(a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)^(1/2)*(cot(x)+csc(x))))*a^(1/2)+arctanh(1/a^(1/2)*(-(a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)^(1/2)*(cot(x)+csc(x)))*(-a+b)^(1/2))*(-(a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)^(1/2)/(a+b*cot(x)^2)^(1/2)*(cot(x)+csc(x))

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 419, normalized size of antiderivative = 6.98

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \left[\frac{(a - b)\sqrt{a} \log\left(2a \tan(x)^2 + 2\sqrt{a}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b\right) + \sqrt{a - b} a \log\left(\frac{(2a - b) \tan(x)^2 - 2\sqrt{a - b}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b}{\tan(x)^2 + 1}\right)}{2(a^2 - ab)} \right.$$

$$- \frac{2a\sqrt{-a + b} \arctan\left(-\frac{\sqrt{-a + b}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a - b}\right) - (a - b)\sqrt{a} \log\left(2a \tan(x)^2 + 2\sqrt{a}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b\right)}{2(a^2 - ab)}$$

$$- \frac{2\sqrt{-a}(a - b) \arctan\left(\frac{\sqrt{-a}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a}\right) - \sqrt{a - b} a \log\left(\frac{(2a - b) \tan(x)^2 - 2\sqrt{a - b}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}} \tan(x)^2 + b}{\tan(x)^2 + 1}\right)}{2(a^2 - ab)},$$

$$\left. - \frac{\sqrt{-a}(a - b) \arctan\left(\frac{\sqrt{-a}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a}\right) + a\sqrt{-a + b} \arctan\left(-\frac{\sqrt{-a + b}\sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{a - b}\right)}{a^2 - ab} \right]$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")

```
[Out] [1/2*((a - b)*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b) + sqrt(a - b)*a*log(((2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(tan(x)^2 + 1)))/(a^2 - a*b), -1/2*(2*a*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a - b)) - (a - b)*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(a^2 - a*b), -1/2*(2*sqrt(-a)*(a - b)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/a) - sqrt(a - b)*a*log(((2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(tan(x)^2 + 1)))/(a^2 - a*b), -(sqrt(-a)*(a - b)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/a) + a*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a - b)))/(a^2 - a*b)]
```

Sympy [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx$$

[In] integrate(tan(x)/(a+b*cot(x)**2)**(1/2),x)

[Out] Integral(tan(x)/sqrt(a + b*cot(x)**2), x)

Maxima [F]

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\tan(x)}{\sqrt{b \cot(x)^2 + a}} dx$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tan(x)/sqrt(b*cot(x)^2 + a), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(48) = 96.

Time = 0.37 (sec) , antiderivative size = 203, normalized size of antiderivative = 3.38

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= -\frac{\left(2a \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right) - 2b \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right) + \sqrt{-a^2+ab} \log(b)\right) \operatorname{sgn}(\sin(x))}{2\sqrt{-a^2+ab}\sqrt{a-b}}$$

$$+ \frac{2\sqrt{a-b} \arctan\left(\frac{\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2 - 2a + b}{2\sqrt{-a^2+ab}}\right)}{\sqrt{-a^2+ab}} + \frac{\log\left(\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2\right)}{\sqrt{a-b}}$$

$$2 \operatorname{sgn}(\sin(x))$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*(2*a*arctan(-(a - b)/sqrt(-a^2 + a*b)) - 2*b*arctan(-(a - b)/sqrt(-a^2 + a*b)) + sqrt(-a^2 + a*b)*log(b))*sgn(sin(x))/(sqrt(-a^2 + a*b)*sqrt(a - b)) + 1/2*(2*sqrt(a - b)*arctan(1/2*((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2 - 2*a + b)/sqrt(-a^2 + a*b))/sqrt(-a^2 + a*b) + log((sqrt(a - b)*sin(x) - sqrt(a*sin(x)^2 - b*sin(x)^2 + b))^2)/sqrt(a - b))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 13.10 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.55

$$\int \frac{\tan(x)}{\sqrt{a + b \cot^2(x)}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{a + \frac{b}{\tan(x)^2}}}{\sqrt{a-b}} + \frac{2\sqrt{a-b}\sqrt{a + \frac{b}{\tan(x)^2}}}{b} - \frac{2a\sqrt{a + \frac{b}{\tan(x)^2}}}{b\sqrt{a-b}}\right)}{\sqrt{a-b}} + \frac{\operatorname{atanh}\left(\frac{\sqrt{a + \frac{b}{\tan(x)^2}}}{\sqrt{a}}\right)}{\sqrt{a}}$$

[In] int(tan(x)/(a + b*cot(x)^2)^(1/2),x)

[Out] atanh((a + b/tan(x)^2)^(1/2)/(a - b)^(1/2) + (2*(a - b)^(1/2)*(a + b/tan(x)^2)^(1/2))/b - (2*a*(a + b/tan(x)^2)^(1/2))/(b*(a - b)^(1/2)))/(a - b)^(1/2) + atanh((a + b/tan(x)^2)^(1/2)/a^(1/2))/a^(1/2)

3.48 $\int \frac{\tan^2(x)}{\sqrt{a+b \cot^2(x)}} dx$

Optimal result	303
Rubi [A] (verified)	303
Mathematica [C] (warning: unable to verify)	305
Maple [B] (verified)	305
Fricas [A] (verification not implemented)	306
Sympy [F]	306
Maxima [F]	306
Giac [B] (verification not implemented)	307
Mupad [F(-1)]	307

Optimal result

Integrand size = 17, antiderivative size = 54

$$\int \frac{\tan^2(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{a-b}} + \frac{\sqrt{a+b \cot^2(x)} \tan(x)}{a}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/(a-b)^{(1/2)}+(a+b*\cot(x)^2)^{(1/2)*\tan(x)/a}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 491, 12, 385, 209}

$$\int \frac{\tan^2(x)}{\sqrt{a+b \cot^2(x)}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{\sqrt{a-b}} + \frac{\tan(x) \sqrt{a+b \cot^2(x)}}{a}$$

[In] $\text{Int}[\text{Tan}[x]^2/\text{Sqrt}[a + b*\text{Cot}[x]^2], x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[x])/\text{Sqrt}[a + b*\text{Cot}[x]^2]]/\text{Sqrt}[a - b] + (\text{Sqrt}[a + b*\text{Cot}[x]^2]*\text{Tan}[x])/a$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match}[\text{Q}[u, (b_)*(\text{v}_)] /; \text{FreeQ}[b, x]]$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 385

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)/((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 491

```
Int[((e_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*c*e*(m + 1))), x] - Dist[1/(a*c*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[(b*c + a*d)*(m + n + 1) + n*(b*c*p + a*d*q) + b*d*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{1}{x^2(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= \frac{\sqrt{a+b\cot^2(x)}\tan(x)}{a} + \frac{\text{Subst}\left(\int \frac{a}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{a} \\
 &= \frac{\sqrt{a+b\cot^2(x)}\tan(x)}{a} + \text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right) \\
 &= \frac{\sqrt{a+b\cot^2(x)}\tan(x)}{a} + \text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right) \\
 &= \frac{\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{\sqrt{a-b}} + \frac{\sqrt{a+b\cot^2(x)}\tan(x)}{a}
 \end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.85 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.48

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \frac{\left(1 + \frac{b \cot^2(x)}{a}\right) \sin^2(x) \left(\frac{4(a-b) \cos^2(x) (a+b \cot^2(x)) \operatorname{Hypergeometric2F1}\left(2, 2, \frac{5}{2}, \frac{(a-b) \cos^2(x)}{a}\right)}{3a^2} + \frac{\arcsin\left(\sqrt{\frac{(a-b) \cos^2(x)}{a}}\right) (a+2b \cot^2(x))}{a \sqrt{\frac{(a-b) \cos^2(x) (a+b \cot^2(x)) \sin^2(x)}{a^2}}}\right)}{\sqrt{a + b \cot^2(x)}}$$

[In] Integrate[Tan[x]^2/Sqrt[a + b*Cot[x]^2], x]

[Out] ((1 + (b*Cot[x]^2)/a)*Sin[x]^2*((4*(a - b)*Cos[x]^2*(a + b*Cot[x]^2)*Hypergeometric2F1[2, 2, 5/2, ((a - b)*Cos[x]^2)/a])/(3*a^2) + (ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*(a + 2*b*Cot[x]^2))/(a*Sqrt[((a - b)*Cos[x]^2*(a + b*Cot[x]^2)*Sin[x]^2)/a^2]))*Tan[x])/Sqrt[a + b*Cot[x]^2]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 282 vs. 2(46) = 92.

Time = 0.91 (sec) , antiderivative size = 283, normalized size of antiderivative = 5.24

method	result
default	$\frac{\sqrt{4} \left(\sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} \ln\left(4 \cos(x) \sqrt{-a+b} \sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}} - 4 \cos(x) a + 4 b \cos(x) + 4 \sqrt{-a+b} \sqrt{-\frac{a \cos(x)^2 - \cos(x)^2 b - a}{(\cos(x)+1)^2}}\right) \right)}{\dots}$

[In] int(tan(x)^2/(a+b*cot(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/2*4^(1/2)/a/(-a+b)^(1/2)/(a+b*cot(x)^2)^(1/2)*((-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)^(1/2)*ln(4*cos(x)*(-a+b)^(1/2)*(-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2-4*cos(x)*a+4*b*cos(x)+4*(-a+b)^(1/2)*(-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)*a*cot(x)+(-a+b)^(1/2)*a*tan(x)+(-a+b)^(1/2)*b*cot(x)+(-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)^(1/2)*ln(4*cos(x)*(-a+b)^(1/2)*(-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2-4*cos(x)*a+4*b*cos(x)+4*(-a+b)^(1/2)*(-a*cos(x)^2-cos(x)^2*b-a)/(cos(x)+1)^2)*a*csc(x)

Fricas [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 229, normalized size of antiderivative = 4.24

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \left[\frac{a\sqrt{-a+b} \log\left(-\frac{a^2 \tan(x)^4 - 2(3a^2 - 4ab) \tan(x)^2 + a^2 - 8ab + 8b^2 + 4(a \tan(x)^3 - (a-2b) \tan(x))\sqrt{-a+b} \sqrt{\frac{a \tan(x)^2 + b}{\tan(x)^2}}}{\tan(x)^4 + 2 \tan(x)^2 + 1}\right) - 4(a - b) \sqrt{-a+b} \arctan\left(\frac{\tan(x) \sqrt{a+b \cot^2(x)}}{a - b}\right)}{4(a^2 - ab)} \right]$$

```
[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] [-1/4*(a*sqrt(-a + b)*log(-(a^2*tan(x)^4 - 2*(3*a^2 - 4*a*b)*tan(x)^2 + a^2 - 8*a*b + 8*b^2 + 4*(a*tan(x)^3 - (a - 2*b)*tan(x))*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) - 4*(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x))/(a^2 - a*b), 1/2*(sqrt(a - b)*a*arctan(2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)/(a*tan(x)^2 - a + 2*b)) + 2*(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x))/(a^2 - a*b)]
```

Sympy [F]

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

```
[In] integrate(tan(x)**2/(a+b*cot(x)**2)^(1/2),x)
```

```
[Out] Integral(tan(x)**2/sqrt(a + b*cot(x)**2), x)
```

Maxima [F]

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\tan(x)^2}{\sqrt{b \cot(x)^2 + a}} dx$$

```
[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(tan(x)^2/sqrt(b*cot(x)^2 + a), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(46) = 92.

Time = 0.34 (sec) , antiderivative size = 216, normalized size of antiderivative = 4.00

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx$$

$$= \frac{\left(a \log \left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2b \right) + \sqrt{-a + b} \sqrt{b} \log \left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2b \right) - b \log \left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2b \right) \right)}{2 \left(a \sqrt{-a + b} - a \sqrt{b} - \sqrt{-a + b} b + b^{\frac{3}{2}} \right)}$$

$$- \frac{\log \left(\left(\frac{\sqrt{-a + b} \cos(x) - \sqrt{-a \cos^2(x) + b \cos^2(x) + a}}{\sqrt{-a + b}} \right)^2 \right)}{2 \operatorname{sgn}(\sin(x))} + \frac{4 \sqrt{-a + b}}{\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos^2(x) + b \cos^2(x) + a} \right)^2 - a}$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*(a*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + sqrt(-a + b)*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 2*a - 2*b)*sgn(sin(x))/(a*sqrt(-a + b) - a*sqrt(b) - sqrt(-a + b)*b + b^(3/2)) - 1/2*(log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2)/sqrt(-a + b) + 4*sqrt(-a + b)/((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a))/sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^2(x)}{\sqrt{a + b \cot^2(x)}} dx = \int \frac{\tan(x)^2}{\sqrt{b \cot(x)^2 + a}} dx$$

[In] int(tan(x)^2/(a + b*cot(x)^2)^(1/2),x)

[Out] int(tan(x)^2/(a + b*cot(x)^2)^(1/2), x)

3.49 $\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{3/2}} dx$

Optimal result	308
Rubi [A] (verified)	308
Mathematica [A] (verified)	310
Maple [A] (verified)	310
Fricas [B] (verification not implemented)	311
Sympy [F]	311
Maxima [F(-2)]	312
Giac [B] (verification not implemented)	312
Mupad [B] (verification not implemented)	312

Optimal result

Integrand size = 17, antiderivative size = 59

$$\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{a}{(a-b)b\sqrt{a+b \cot^2(x)}}$$

[Out] $-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2))/(a-b)^{(1/2)})/(a-b)^{(3/2)}+a/(a-b)/b/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.294$, Rules used = {3751, 457, 79, 65, 214}

$$\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{a}{b(a-b)\sqrt{a+b \cot^2(x)}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}}$$

[In] $\text{Int}[\text{Cot}[x]^3/(a + b*\text{Cot}[x]^2)^{(3/2)}, x]$

[Out] $-(\text{ArcTanh}[\text{Sqrt}[a + b*\text{Cot}[x]^2]/\text{Sqrt}[a - b]]/(a - b)^{(3/2)}) + a/((a - b)*b*\text{Sqrt}[a + b*\text{Cot}[x]^2])$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 79

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_.)*((c_) + (d_.)*(x_)^(n_.))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{x^3}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right) \\ &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{x}{(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)\right) \\ &= \frac{a}{(a-b)b\sqrt{a+b\cot^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)} \end{aligned}$$

$$= \frac{a}{(a-b)b\sqrt{a+b\cot^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)b}$$

$$= -\frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{a}{(a-b)b\sqrt{a+b\cot^2(x)}}$$

Mathematica [A] (verified)

Time = 0.26 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\cot^3(x)}{(a+b\cot^2(x))^{3/2}} dx = -\frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{a}{(a-b)b\sqrt{a+b\cot^2(x)}}$$

[In] Integrate[Cot[x]^3/(a + b*Cot[x]^2)^(3/2),x]

[Out] -(ArcTanh[Sqrt[a + b*Cot[x]^2]/Sqrt[a - b]]/(a - b)^(3/2)) + a/((a - b)*b*Sqrt[a + b*Cot[x]^2])

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.15

method	result	size
derivativedivides	$\frac{1}{b\sqrt{a+b\cot(x)^2}} + \frac{1}{(a-b)\sqrt{a+b\cot(x)^2}} + \frac{\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)\sqrt{-a+b}}$	68
default	$\frac{1}{b\sqrt{a+b\cot(x)^2}} + \frac{1}{(a-b)\sqrt{a+b\cot(x)^2}} + \frac{\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)\sqrt{-a+b}}$	68

[In] int(cot(x)^3/(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] 1/b/(a+b*cot(x)^2)^(1/2)+1/(a-b)/(a+b*cot(x)^2)^(1/2)+1/(a-b)/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(51) = 102.

Time = 0.31 (sec) , antiderivative size = 385, normalized size of antiderivative = 6.53

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx = \left[\frac{(ab + b^2 - (ab - b^2) \cos(2x)) \sqrt{a-b} \log\left(-\sqrt{a-b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) + a) - 2(a^2 - ab - (a^2 - ab) \cos(2x)) \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}}\right)}{2(a^3b - a^2b^2 - ab^3 + b^4 - (a^3b - 3a^2b^2 + 3ab^3 - b^4) \cos(2x))} \right. \\ \left. - \frac{(ab + b^2 - (ab - b^2) \cos(2x)) \sqrt{-a+b} \arctan\left(-\frac{\sqrt{-a+b} \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}}}{a-b}\right) - (a^2 - ab - (a^2 - ab) \cos(2x)) \sqrt{\frac{(a-b) \cos(2x) - a - b}{\cos(2x) - 1}}}{a^3b - a^2b^2 - ab^3 + b^4 - (a^3b - 3a^2b^2 + 3ab^3 - b^4) \cos(2x)} \right]$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [-1/2*((a*b + b^2 - (a*b - b^2)*cos(2*x))*sqrt(a - b)*log(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1) - (a - b)*cos(2*x) + a) - 2*(a^2 - a*b - (a^2 - a*b)*cos(2*x))*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^3*b - a^2*b^2 - a*b^3 + b^4 - (a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4)*cos(2*x)), -((a*b + b^2 - (a*b - b^2)*cos(2*x))*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))/(a - b)) - (a^2 - a*b - (a^2 - a*b)*cos(2*x))*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^3*b - a^2*b^2 - a*b^3 + b^4 - (a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4)*cos(2*x)]]

Sympy [F]

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\cot^3(x)}{(a + b \cot^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(cot(x)**3/(a+b*cot(x)**2)**(3/2),x)

[Out] Integral(cot(x)**3/(a + b*cot(x)**2)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(51) = 102.

Time = 0.32 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.85

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx = -\frac{\log(|b|) \operatorname{sgn}(\sin(x))}{2(\sqrt{a-ba} - \sqrt{a-bb})} + \frac{\frac{a \sin(x)}{\sqrt{a \sin(x)^2 - b \sin(x)^2 + b(ab-b^2)}} + \frac{\log\left(-\sqrt{a-b} \sin(x) + \sqrt{a \sin(x)^2 - b \sin(x)^2 + b}\right)}{(a-b)^{3/2}}}{\operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/2*log(abs(b))*sgn(sin(x))/(sqrt(a-b)*a - sqrt(a-b)*b) + (a*sin(x)/(sqrt(a*sin(x)^2 - b*sin(x)^2 + b)*(a*b - b^2)) + log(abs(-sqrt(a-b)*sin(x) + sqrt(a*sin(x)^2 - b*sin(x)^2 + b)))/(a-b)^(3/2))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 14.40 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{a}{(ab - b^2) \sqrt{b \cot(x)^2 + a}} - \frac{\operatorname{atanh}\left(\frac{\sqrt{b \cot(x)^2 + a}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}}$$

[In] int(cot(x)^3/(a + b*cot(x)^2)^(3/2),x)

[Out] a/((a*b - b^2)*(a + b*cot(x)^2)^(1/2)) - atanh((a + b*cot(x)^2)^(1/2)/(a - b)^(1/2))/(a - b)^(3/2)

$$3.50 \quad \int \frac{\cot^2(x)}{(a+b \cot^2(x))^{3/2}} dx$$

Optimal result	313
Rubi [A] (verified)	313
Mathematica [B] (verified)	315
Maple [A] (verified)	315
Fricas [B] (verification not implemented)	316
Sympy [F]	316
Maxima [F(-2)]	317
Giac [B] (verification not implemented)	317
Mupad [F(-1)]	317

Optimal result

Integrand size = 17, antiderivative size = 59

$$\int \frac{\cot^2(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{3/2}} - \frac{\cot(x)}{(a-b)\sqrt{a+b \cot^2(x)}}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/(a-b)^{(3/2)}-\cot(x)/(a-b)/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {3751, 482, 385, 209}

$$\int \frac{\cot^2(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{3/2}} - \frac{\cot(x)}{(a-b)\sqrt{a+b \cot^2(x)}}$$

[In] $\text{Int}[\text{Cot}[x]^2/(a + b*\text{Cot}[x]^2)^{(3/2)}, x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a - b]*\text{Cot}[x])/(\text{Sqrt}[a + b*\text{Cot}[x]^2])]/(a - b)^{(3/2)} - \text{Cot}[x]/((a - b)*\text{Sqrt}[a + b*\text{Cot}[x]^2])$

Rule 209

$\text{Int}[(a_+ + (b_-)*(x_-)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Subst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 482

```
Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(n*(b*c - a*d)*(p + 1))), x] - Dist[e^n/(n*(b*c - a*d)*(p + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(m - n + 1) + d*(m + n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GeQ[n, m - n + 1] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^2}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right) \\
&= -\frac{\cot(x)}{(a-b)\sqrt{a+b\cot^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{a-b} \\
&= -\frac{\cot(x)}{(a-b)\sqrt{a+b\cot^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{a-b} \\
&= \frac{\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{(a-b)^{3/2}} - \frac{\cot(x)}{(a-b)\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 137 vs. $2(59) = 118$.

Time = 0.81 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.32

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{(-a + b) \cot(x) \sqrt{1 + \frac{b \cot^2(x)}{a}} + \frac{1}{2} \operatorname{arctanh}\left(\frac{\sqrt{-\frac{(a-b) \cot^2(x)}{a}}}{\sqrt{1 + \frac{b \cot^2(x)}{a}}}\right) (-a - b + (a - b) \cos(2x))}{(a - b)^2 \sqrt{a + b \cot^2(x)} \sqrt{1 + \frac{b \cot^2(x)}{a}}}$$

[In] Integrate[Cot[x]^2/(a + b*Cot[x]^2)^(3/2),x]

[Out] $((-a + b) \operatorname{Cot}[x] \operatorname{Sqrt}[1 + (b \operatorname{Cot}[x]^2)/a] + (\operatorname{ArcTanh}[\operatorname{Sqrt}[-((a - b) \operatorname{Cot}[x]^2)/a]]/\operatorname{Sqrt}[1 + (b \operatorname{Cot}[x]^2)/a]) * (-a - b + (a - b) \operatorname{Cos}[2*x]) \operatorname{Sqrt}[-((a - b) \operatorname{Cot}[x]^2)/a]) * \operatorname{Csc}[x] \operatorname{Sec}[x])/2)/((a - b)^2 \operatorname{Sqrt}[a + b \operatorname{Cot}[x]^2] \operatorname{Sqrt}[1 + (b \operatorname{Cot}[x]^2)/a])$

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.68

method	result	size
derivativedivides	$-\frac{\cot(x)}{a\sqrt{a+b\cot(x)^2}} - \frac{b\cot(x)}{(a-b)a\sqrt{a+b\cot(x)^2}} + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b)\cot(x)}{\sqrt{b^4(a-b)}\sqrt{a+b\cot(x)^2}}\right)}{(a-b)^2 b^2}$	99
default	$-\frac{\cot(x)}{a\sqrt{a+b\cot(x)^2}} - \frac{b\cot(x)}{(a-b)a\sqrt{a+b\cot(x)^2}} + \frac{\sqrt{b^4(a-b)} \arctan\left(\frac{b^2(a-b)\cot(x)}{\sqrt{b^4(a-b)}\sqrt{a+b\cot(x)^2}}\right)}{(a-b)^2 b^2}$	99

[In] int(cot(x)^2/(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] $-\cot(x)/a/(a+b*\cot(x)^2)^(1/2)-b/(a-b)*\cot(x)/a/(a+b*\cot(x)^2)^(1/2)+1/(a-b)^2*(b^4*(a-b))^(1/2)/b^2*\arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*\cot(x)^2)^(1/2)*\cot(x))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 173 vs. 2(51) = 102.

Time = 0.37 (sec) , antiderivative size = 388, normalized size of antiderivative = 6.58

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \left[\frac{((a - b) \cos(2x) - a - b) \sqrt{-a + b} \log\left(-2(a^2 - 2ab + b^2) \cos(2x)^2 - 2((a - b) \cos(2x) - a - b) \sqrt{-a + b}\right)}{4(a^3 - a^2b - ab^2 + b^3 - (a^3 - 3a^2b + 3ab^2 - b^3) \cos(2x))} \right. \\ \left. - \frac{((a - b) \cos(2x) - a - b) \sqrt{a - b} \arctan\left(-\frac{\sqrt{a - b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{(a - b) \cos(2x) - b}\right) + 2(a - b) \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \sin(2x)}{2(a^3 - a^2b - ab^2 + b^3 - (a^3 - 3a^2b + 3ab^2 - b^3) \cos(2x))} \right]$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [-1/4*(((a - b)*cos(2*x) - a - b)*sqrt(-a + b)*log(-2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 - 2*((a - b)*cos(2*x) - b)*sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x) + a^2 - 2*b^2 + 4*(a*b - b^2)*cos(2*x)) + 4*(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x))/(a^3 - a^2*b - a*b^2 + b^3 - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*cos(2*x)), -1/2*(((a - b)*cos(2*x) - a - b)*sqrt(a - b)*arctan(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x)/((a - b)*cos(2*x) - b)) + 2*(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*sin(2*x))/(a^3 - a^2*b - a*b^2 + b^3 - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*cos(2*x))]

Sympy [F]

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\cot^2(x)}{(a + b \cot^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(cot(x)**2/(a+b*cot(x)**2)**(3/2),x)

[Out] Integral(cot(x)**2/(a + b*cot(x)**2)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 159 vs. 2(51) = 102.

Time = 0.34 (sec) , antiderivative size = 159, normalized size of antiderivative = 2.69

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\left(\sqrt{b} \log\left(\left|-\sqrt{-a+b} + \sqrt{b}\right|\right) + \sqrt{-a+b}\right) \operatorname{sgn}(\sin(x))}{a\sqrt{-a+b}\sqrt{b} - \sqrt{-a+b}b^{3/2}} + \frac{\frac{\sqrt{-a \cos(x)^2 + b \cos(x)^2 + a \cos(x)}}{(a \cos(x)^2 - b \cos(x)^2 - a)(a-b)} - \frac{\log\left(\left|-\sqrt{-a+b} \cos(x) + \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}\right|\right)}{(a-b)\sqrt{-a+b}}}{\operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="giac")

[Out] (sqrt(b)*log(abs(-sqrt(-a + b) + sqrt(b))) + sqrt(-a + b))*sgn(sin(x))/(a*sqrt(-a + b)*sqrt(b) - sqrt(-a + b)*b^(3/2)) + (sqrt(-a*cos(x)^2 + b*cos(x)^2 + a)*cos(x)/((a*cos(x)^2 - b*cos(x)^2 - a)*(a - b)) - log(abs(-sqrt(-a + b)*cos(x) + sqrt(-a*cos(x)^2 + b*cos(x)^2 + a)))/((a - b)*sqrt(-a + b)))/sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\cot(x)^2}{(b \cot(x)^2 + a)^{3/2}} dx$$

[In] int(cot(x)^2/(a + b*cot(x)^2)^(3/2),x)

[Out] int(cot(x)^2/(a + b*cot(x)^2)^(3/2), x)

3.51 $\int \frac{\cot(x)}{(a+b \cot^2(x))^{3/2}} dx$

Optimal result	318
Rubi [A] (verified)	318
Mathematica [C] (verified)	320
Maple [A] (verified)	320
Fricas [B] (verification not implemented)	321
Sympy [A] (verification not implemented)	321
Maxima [F(-2)]	322
Giac [B] (verification not implemented)	322
Mupad [B] (verification not implemented)	322

Optimal result

Integrand size = 15, antiderivative size = 55

$$\int \frac{\cot(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} - \frac{1}{(a-b)\sqrt{a+b \cot^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(3/2)}-1/(a-b)/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 53, 65, 214}

$$\int \frac{\cot(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} - \frac{1}{(a-b)\sqrt{a+b \cot^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]/(a+b*\operatorname{Cot}[x]^2)^{(3/2)}, x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a-b]]/(a-b)^{(3/2)}-1/((a-b)*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2])$

Rule 53

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)}/((b*c - a*d)*(m + 1))), x] - \operatorname{Dist}[d*((m + n + 2)/((b*c - a*d)*(m + 1))), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, n\}, x\} \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{LtQ}[m, -1] \ \&\& \ !(\operatorname{LtQ}$

`[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n])) && IntLinearQ[a, b, c, d, m, n, x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 455

`Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n + 1, 0]`

Rule 3751

`Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))`

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)\right) \\
 &= -\frac{1}{(a-b)\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)} \\
 &= -\frac{1}{(a-b)\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)b}
 \end{aligned}$$

$$= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} - \frac{1}{(a-b)\sqrt{a+b \cot^2(x)}}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.05 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.80

$$\int \frac{\cot(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\operatorname{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b \cot^2(x)}{a-b}\right)}{(-a+b)\sqrt{a+b \cot^2(x)}}$$

[In] Integrate[Cot[x]/(a + b*Cot[x]^2)^(3/2),x]

[Out] Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Cot[x]^2)/(a - b)]/((-a + b)*Sqrt[a + b*Cot[x]^2])

Maple [A] (verified)

Time = 0.04 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.02

method	result	size
derivativedivides	$-\frac{1}{(a-b)\sqrt{a+b \cot(x)^2}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)\sqrt{-a+b}}$	56
default	$-\frac{1}{(a-b)\sqrt{a+b \cot(x)^2}} - \frac{\operatorname{arctan}\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)\sqrt{-a+b}}$	56

[In] int(cot(x)/(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/(a-b)/(a+b*cot(x)^2)^(1/2)-1/(a-b)/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 166 vs. 2(47) = 94.

Time = 0.31 (sec) , antiderivative size = 344, normalized size of antiderivative = 6.25

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\left(((a - b) \cos(2x) - a - b) \sqrt{a - b} \log \left(\sqrt{a - b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} (\cos(2x) - 1) - \right) - ((a - b) \cos(2x) - a + b) \sqrt{-a + b} \arctan \left(-\frac{\sqrt{-a + b} \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}}}{a - b} \right) - ((a - b) \cos(2x) - a + b) \sqrt{\frac{(a - b) \cos(2x) - a - b}{\cos(2x) - 1}} \right)}{a^3 - a^2 b - ab^2 + b^3 - (a^3 - 3a^2 b + 3ab^2 - b^3) \cos(2x)}$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [1/2*(((a - b)*cos(2*x) - a - b)*sqrt(a - b)*log(sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1) - (a - b)*cos(2*x) + a) + 2*(((a - b)*cos(2*x) - a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^3 - a^2*b - a*b^2 + b^3 - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*cos(2*x)), -(((a - b)*cos(2*x) - a - b)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a - b) - ((a - b)*cos(2*x) - a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^3 - a^2*b - a*b^2 + b^3 - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*cos(2*x))]

Sympy [A] (verification not implemented)

Time = 4.82 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.60

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx = - \begin{cases} \frac{2 \left(\frac{b}{2(a-b)\sqrt{a+b \cot^2(x)}} + \frac{b \operatorname{atan} \left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{-a+b}} \right)}{2\sqrt{-a+b}(a-b)} \right)}{b} & \text{for } b \neq 0 \\ \infty \cot^2(x) & \text{for } a^{\frac{3}{2}} = 0 \\ \frac{\log(2a^{\frac{3}{2}} \cot^2(x) + 2a^{\frac{3}{2}})}{2a^{\frac{3}{2}}} & \text{otherwise} \end{cases}$$

[In] integrate(cot(x)/(a+b*cot(x)**2)**(3/2),x)

[Out] -Piecewise((2*(b/(2*(a - b)*sqrt(a + b*cot(x)**2)) + b*atan(sqrt(a + b*cot(x)**2)/sqrt(-a + b))/(2*sqrt(-a + b)*(a - b)))/b, Ne(b, 0)), (Piecewise((zoo*cot(x)**2, Eq(a**(3/2), 0)), (log(2*a**(3/2)*cot(x)**2 + 2*a**(3/2))/(2*a**(3/2)), True)), True))

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(47) = 94.

Time = 0.30 (sec) , antiderivative size = 105, normalized size of antiderivative = 1.91

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\log(|b|) \operatorname{sgn}(\sin(x))}{2(\sqrt{a-ba} - \sqrt{a-bb})} - \frac{\log\left(\left|-\sqrt{a-b}\sin(x) + \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right|\right)}{(a-b)^{3/2}} + \frac{\sin(x)}{\sqrt{a\sin(x)^2 - b\sin(x)^2 + b(a-b)}} \operatorname{sgn}(\sin(x))$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(3/2),x, algorithm="giac")

[Out] 1/2*log(abs(b))*sgn(sin(x))/(sqrt(a-b)*a - sqrt(a-b)*b) - (log(abs(-sqrt(a-b)*sin(x) + sqrt(a*sin(x)^2 - b*sin(x)^2 + b)))/(a-b)^(3/2) + sin(x)/(sqrt(a*sin(x)^2 - b*sin(x)^2 + b)*(a-b)))/sgn(sin(x))

Mupad [B] (verification not implemented)

Time = 14.41 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.85

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \cot(x)^2 + a}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} - \frac{1}{(a-b) \sqrt{b \cot(x)^2 + a}}$$

[In] int(cot(x)/(a + b*cot(x)^2)^(3/2),x)

[Out] atanh((a + b*cot(x)^2)^(1/2)/(a - b)^(1/2))/(a - b)^(3/2) - 1/((a - b)*(a + b*cot(x)^2)^(1/2))

$$3.52 \quad \int \frac{\tan(x)}{(a+b \cot^2(x))^{3/2}} dx$$

Optimal result	323
Rubi [A] (verified)	323
Mathematica [C] (verified)	325
Maple [B] (warning: unable to verify)	326
Fricas [B] (verification not implemented)	326
Sympy [F]	327
Maxima [F]	327
Giac [B] (verification not implemented)	327
Mupad [B] (verification not implemented)	328

Optimal result

Integrand size = 15, antiderivative size = 84

$$\int \frac{\tan(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{b}{a(a-b)\sqrt{a+b \cot^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/a^{(1/2)})/a^{(3/2)}-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(3/2)}+b/a/(a-b)/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 457, 87, 162, 65, 214}

$$\int \frac{\tan(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{b}{a(a-b)\sqrt{a+b \cot^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]/(a+b*\operatorname{Cot}[x]^2)^{(3/2)},x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a]]/a^{(3/2)}-\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a-b]]/(a-b)^{(3/2)}+b/(a*(a-b)*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2])$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 87

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Simp[f*((e + f*x)^(p + 1)/((p + 1)*(b*e - a*f)*(d*e - c*f))),
x] + Dist[1/((b*e - a*f)*(d*e - c*f)), Int[(b*d*e - b*c*f - a*d*f - b*d*f*x
)*(e + f*x)^(p + 1)/((a + b*x)*(c + d*x))], x], x] /; FreeQ[{a, b, c, d, e
, f}, x] && LtQ[p, -1]
```

Rule 162

```
Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*
((c_.) + (d_.)*(x_))), x_Symbol] := Dist[(b*g - a*h)/(b*c - a*d), Int[(e +
f*x)^p/(a + b*x), x], x] - Dist[(d*g - c*h)/(b*c - a*d), Int[(e + f*x)^p/(c
+ d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[(((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{1}{x(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{x(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)\right) \\
&= \frac{b}{a(a-b)\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{a-b-bx}{x(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2a(a-b)} \\
&= \frac{b}{a(a-b)\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2a} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)} \\
&= \frac{b}{a(a-b)\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{ab} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)b} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a}}\right)}{a^{3/2}} - \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{3/2}} + \frac{b}{a(a-b)\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.89

$$\int \frac{\tan(x)}{(a+b\cot^2(x))^{3/2}} dx = \frac{a \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b\cot^2(x)}{a-b}\right) + (-a+b) \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, 1 + \frac{b\cot^2(x)}{a}\right)}{a(a-b)\sqrt{a+b\cot^2(x)}}$$

[In] Integrate[Tan[x]/(a + b*Cot[x]^2)^(3/2), x]

[Out] (a*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Cot[x]^2)/(a - b)] + (-a + b)*Hypergeometric2F1[-1/2, 1, 1/2, 1 + (b*Cot[x]^2)/a])/(a*(a - b)*Sqrt[a + b*Cot[x]^2])

$$2 + b^3 + (a^3 - 2a^2b + ab^2)\tan(x)^2\sqrt{a}\log(2a\tan(x)^2 + 2\sqrt{a}\sqrt{(a\tan(x)^2 + b)/\tan(x)^2}\tan(x)^2 + b)/(a^4b - 2a^3b^2 + a^2b^3 + (a^5 - 2a^4b + a^3b^2)\tan(x)^2), 1/2(2(a^2b - ab^2)\sqrt{(a\tan(x)^2 + b)/\tan(x)^2}\tan(x)^2 - 2(a^2b - 2ab^2 + b^3 + (a^3 - 2a^2b + ab^2)\tan(x)^2)\sqrt{-a}\arctan(\sqrt{-a}\sqrt{(a\tan(x)^2 + b)/\tan(x)^2})/a) - (a^3\tan(x)^2 + a^2b)\sqrt{a - b}\log(((2a - b)\tan(x)^2 + 2\sqrt{a - b}\sqrt{(a\tan(x)^2 + b)/\tan(x)^2}\tan(x)^2 + b)/(\tan(x)^2 + 1)))/(a^4b - 2a^3b^2 + a^2b^3 + (a^5 - 2a^4b + a^3b^2)\tan(x)^2), ((a^2b - ab^2)\sqrt{(a\tan(x)^2 + b)/\tan(x)^2}\tan(x)^2 - (a^2b - 2ab^2 + b^3 + (a^3 - 2a^2b + ab^2)\tan(x)^2)\sqrt{-a}\arctan(\sqrt{-a}\sqrt{(a\tan(x)^2 + b)/\tan(x)^2})/a) - (a^3\tan(x)^2 + a^2b)\sqrt{-a + b}\arctan(-\sqrt{-a + b}\sqrt{(a\tan(x)^2 + b)/\tan(x)^2})/(a - b))/(a^4b - 2a^3b^2 + a^2b^3 + (a^5 - 2a^4b + a^3b^2)\tan(x)^2)]$$

Sympy [F]

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\tan(x)}{(a + b \cot^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(tan(x)/(a+b*cot(x)**2)**(3/2),x)

[Out] Integral(tan(x)/(a + b*cot(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\tan(x)}{(b \cot(x)^2 + a)^{\frac{3}{2}}} dx$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate(tan(x)/(b*cot(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 295 vs. 2(70) = 140.

$$\begin{aligned}
& /2 + (((a - b)^3)^{(1/2)} * (12*a^5*b^7 - 2*a^4*b^8 - 28*a^6*b^6 + 32*a^7*b^5 - \\
& 18*a^8*b^4 + 4*a^9*b^3 + ((a + b/\tan(x)^2)^{(1/2)} * ((a - b)^3)^{(1/2)} * (8*a^5* \\
& b^8 - 56*a^6*b^7 + 160*a^7*b^6 - 240*a^8*b^5 + 200*a^9*b^4 - 88*a^{10}*b^3 + \\
& 16*a^{11}*b^2)) / (4*(a - b)^3)) / (2*(a - b)^3) * 1i) / (a - b)^3 + (((a - b)^3)^{(1/2)} * \\
& (((a + b/\tan(x)^2)^{(1/2)} * (2*a^3*b^7 - 10*a^4*b^6 + 22*a^5*b^5 - 26*a^6* \\
& b^4 + 16*a^7*b^3 - 4*a^8*b^2)) / 2 + (((a - b)^3)^{(1/2)} * (2*a^4*b^8 - 12*a^5* \\
& b^7 + 28*a^6*b^6 - 32*a^7*b^5 + 18*a^8*b^4 - 4*a^9*b^3 + ((a + b/\tan(x)^2)^{(1/2)} * \\
& ((a - b)^3)^{(1/2)} * (8*a^5*b^8 - 56*a^6*b^7 + 160*a^7*b^6 - 240*a^8*b^5 \\
& + 200*a^9*b^4 - 88*a^{10}*b^3 + 16*a^{11}*b^2)) / (4*(a - b)^3)) / (2*(a - b)^3)) \\
& * 1i) / (a - b)^3 / (2*a^3*b^6 - 6*a^4*b^5 + 6*a^5*b^4 - 2*a^6*b^3 - (((a - b)^3)^{(1/2)} * \\
& (((a + b/\tan(x)^2)^{(1/2)} * (2*a^3*b^7 - 10*a^4*b^6 + 22*a^5*b^5 - 26* \\
& a^6*b^4 + 16*a^7*b^3 - 4*a^8*b^2)) / 2 + (((a - b)^3)^{(1/2)} * (12*a^5*b^7 - 2* \\
& a^4*b^8 - 28*a^6*b^6 + 32*a^7*b^5 - 18*a^8*b^4 + 4*a^9*b^3 + ((a + b/\tan(x)^2)^{(1/2)} * \\
& ((a - b)^3)^{(1/2)} * (8*a^5*b^8 - 56*a^6*b^7 + 160*a^7*b^6 - 240*a^8* \\
& b^5 + 200*a^9*b^4 - 88*a^{10}*b^3 + 16*a^{11}*b^2)) / (4*(a - b)^3)) / (2*(a - b)^3)) / (a - b)^3 + \\
& (((a - b)^3)^{(1/2)} * (((a + b/\tan(x)^2)^{(1/2)} * (2*a^3*b^7 - 10*a^4*b^6 + 22*a^5*b^5 - 26* \\
& a^6*b^4 + 16*a^7*b^3 - 4*a^8*b^2)) / 2 + (((a - b)^3)^{(1/2)} * (12*a^5*b^7 - 2* \\
& a^4*b^8 - 28*a^6*b^6 + 32*a^7*b^5 - 18*a^8*b^4 + 4*a^9*b^3 + ((a + b/\tan(x)^2)^{(1/2)} * \\
& ((a - b)^3)^{(1/2)} * (8*a^5*b^8 - 56*a^6*b^7 + 160*a^7*b^6 - 240*a^8* \\
& b^5 + 200*a^9*b^4 - 88*a^{10}*b^3 + 16*a^{11}*b^2)) / (4*(a - b)^3)) / (2*(a - b)^3)) / (a - b)^3 + \\
& (((a - b)^3)^{(1/2)} * (2*a^4*b^8 - 12*a^5*b^7 + 28*a^6*b^6 - 32*a^7*b^5 + 18*a^8*b^4 \\
& - 4*a^9*b^3 + ((a + b/\tan(x)^2)^{(1/2)} * ((a - b)^3)^{(1/2)} * (8*a^5*b^8 - 56*a^6* \\
& b^7 + 160*a^7*b^6 - 240*a^8*b^5 + 200*a^9*b^4 - 88*a^{10}*b^3 + 16*a^{11}*b^2) \\
&) / (4*(a - b)^3)) / (2*(a - b)^3)) / (a - b)^3) * ((a - b)^3)^{(1/2)} * 1i) / (a - b)^3 - b / ((a*b - a^2) * (a + b/\tan(x)^2)^{(1/2)})
\end{aligned}$$

$$3.53 \quad \int \frac{\tan^2(x)}{(a+b \cot^2(x))^{3/2}} dx$$

Optimal result	330
Rubi [A] (verified)	330
Mathematica [C] (warning: unable to verify)	332
Maple [B] (warning: unable to verify)	333
Fricas [B] (verification not implemented)	334
Sympy [F]	334
Maxima [F]	334
Giac [B] (verification not implemented)	335
Mupad [F(-1)]	335

Optimal result

Integrand size = 17, antiderivative size = 92

$$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{3/2}} + \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} + \frac{(a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{a^2(a-b)}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/(a-b)^{(3/2)}+b*\tan(x)/a/(a-b)/(a+b*\cot(x)^2)^{(1/2)}+(a-2*b)*(a+b*\cot(x)^2)^{(1/2)}*\tan(x)/a^2/(a-b)$

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 483, 597, 12, 385, 209}

$$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{(a-2b) \tan(x) \sqrt{a+b \cot^2(x)}}{a^2(a-b)} + \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{3/2}} + \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}}$$

[In] $\text{Int}[\text{Tan}[x]^2/(a+b*\text{Cot}[x]^2)^{(3/2)},x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a-b]*\text{Cot}[x])/(\text{Sqrt}[a+b*\text{Cot}[x]^2])]/(a-b)^{(3/2)}+(b*\text{Tan}[x])/((a*(a-b)*\text{Sqrt}[a+b*\text{Cot}[x]^2))+((a-2*b)*\text{Sqrt}[a+b*\text{Cot}[x]^2]*\text{Tan}[x])/(a^2*(a-b))$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 483

```
Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_
))^(-q_), x_Symbol] := Simp[(-b)*(e*x)^(m + 1)*(a + b*x^n)^(p + 1)*((c + d*x
^n)^(q + 1)/(a*e*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p +
1)), Int[(e*x)^m*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*b*(m + 1) + n*(b
*c - a*d)*(p + 1) + d*b*(m + n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a
, b, c, d, e, m, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] &&
IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 597

```
Int[((g_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_
))^(-q_)*((e_) + (f_)*(x_)^(n_)), x_Symbol] := Simp[e*(g*x)^(m + 1)*(a + b
*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*c*g*(m + 1))), x] + Dist[1/(a*c*g^n*(
m + 1)), Int[(g*x)^(m + n)*(a + b*x^n)^p*(c + d*x^n)^q*Simp[a*f*c*(m + 1) -
e*(b*c + a*d)*(m + n + 1) - e*n*(b*c*p + a*d*q) - b*e*d*(m + n*(p + q + 2)
+ 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0
] && LtQ[m, -1]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_))*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{1}{x^2(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right) \\
&= \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{a-2b-2bx^2}{x^2(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{a(a-b)} \\
&= \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} + \frac{(a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{a^2(a-b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{a^2}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{a^2(a-b)} \\
&= \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} + \frac{(a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{a^2(a-b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{a-b} \\
&= \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} + \frac{(a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{a^2(a-b)} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{a-b} \\
&= \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{3/2}} + \frac{b \tan(x)}{a(a-b)\sqrt{a+b \cot^2(x)}} + \frac{(a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{a^2(a-b)}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.95 (sec) , antiderivative size = 674, normalized size of antiderivative = 7.33

$$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{3/2}} dx = \frac{\sin^2(x) \left(\frac{12b \csc^2(x)}{a-b} + \frac{8b^2 \cot^2(x) \csc^2(x)}{a(a-b)} + \frac{16(a-b) \cos^2(x) \text{Hypergeometric2F1}\left(2, 2, \frac{7}{2}, \frac{(a-b) \cos^2(x)}{a}\right)}{15a} \right)}{1}$$

[In] Integrate[Tan[x]^2/(a + b*Cot[x]^2)^(3/2), x]

[Out] (Sin[x]^2*((12*b*Csc[x]^2)/(a - b) + (8*b^2*Cot[x]^2*Csc[x]^2)/(a*(a - b)) + (16*(a - b)*Cos[x]^2*Hypergeometric2F1[2, 2, 7/2, ((a - b)*Cos[x]^2)/a]))/(15*a) + (8*(a - b)*b*Cos[x]^2*Cot[x]^2*Hypergeometric2F1[2, 2, 7/2, ((a -

$$\begin{aligned}
& b) \cdot \cos[x]^2/a) / (3 \cdot a^2) + (8 \cdot (a - b) \cdot b^2 \cdot \cos[x]^2 \cdot \cot[x]^4 \cdot \text{Hypergeometric2} \\
& \text{F1}[2, 2, 7/2, ((a - b) \cdot \cos[x]^2/a)] / (5 \cdot a^3) + (8 \cdot (a - b) \cdot \cos[x]^2 \cdot \text{Hypergeo} \\
& \text{metricPFQ}[\{2, 2, 2\}, \{1, 7/2\}, ((a - b) \cdot \cos[x]^2/a)] / (15 \cdot a) + (16 \cdot (a - b) \cdot \\
& b \cdot \cos[x]^2 \cdot \cot[x]^2 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 7/2\}, ((a - b) \cdot \cos[x]^2/a)] / (15 \cdot a^2) \\
& + (8 \cdot (a - b) \cdot b^2 \cdot \cos[x]^2 \cdot \cot[x]^4 \cdot \text{HypergeometricPFQ}[\{2, 2, 2\}, \{1, 7/2\}, ((a - b) \cdot \cos[x]^2/a)] / (15 \cdot a^3) \\
& + (3 \cdot a \cdot \sec[x]^2) / (a - b) - (3 \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] / (((a - b) \cdot \cos[x]^2/a)^{3/2} \cdot \text{Sqrt}[(a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a]) \\
& - (12 \cdot b \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] \cdot \cot[x]^2) / (a \cdot ((a - b) \cdot \cos[x]^2/a)^{3/2} \cdot \text{Sqrt}[(a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a]) \\
& - (8 \cdot b^2 \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] \cdot \cot[x]^4) / (a^2 \cdot ((a - b) \cdot \cos[x]^2/a)^{3/2} \cdot \text{Sqrt}[(a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a]) \\
& + (3 \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] / \text{Sqrt}[(a - b) \cdot \cos[x]^2 \cdot (a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a^2] \\
& + (12 \cdot b \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] \cdot \cot[x]^2) / (a \cdot \text{Sqrt}[(a - b) \cdot \cos[x]^2 \cdot (a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a^2]) \\
& + (8 \cdot b^2 \cdot \text{ArcSin}[\text{Sqrt}[(a - b) \cdot \cos[x]^2/a]] \cdot \cot[x]^4) / (a^2 \cdot \text{Sqrt}[(a - b) \cdot \cos[x]^2 \cdot (a + b \cdot \cot[x]^2) \cdot \sin[x]^2/a^2]) \\
& \cdot \tan[x] / (a \cdot \text{Sqrt}[a + b \cdot \cot[x]^2])
\end{aligned}$$

Maple [B] (warning: unable to verify)

Leaf count of result is larger than twice the leaf count of optimal. 630 vs. $2(82) = 164$.

Time = 2.93 (sec) , antiderivative size = 631, normalized size of antiderivative = 6.86

method	result
default	$\frac{\sqrt{4} \left(-\sqrt{-a+ab} ab(1-\cos(x))^4 \csc(x)^4 + 2\sqrt{-a+bb^2} b^2(1-\cos(x))^4 \csc(x)^4 + \sqrt{b(1-\cos(x))^4} \csc(x)^4 + 4a(1-\cos(x))^2 \csc(x)^2 - 2b(1-\cos(x))^2 \right)}{\dots}$

[In] `int(tan(x)^2/(a+b*cot(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned}
& 1/2 \cdot 4^{1/2} / (-a+b)^{1/2} / a^2 / (a-b) \cdot (-(-a+b)^{1/2} \cdot a \cdot b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 \\
& + 2 \cdot (-a+b)^{1/2} \cdot b^2 \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + (b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + 4 \cdot a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 2 \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + b)^{1/2} \cdot \ln(4 \cdot (a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + (-a+b)^{1/2} \cdot (b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + 4 \cdot a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 2 \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + b)^{1/2} - a+b) / ((1-\cos(x))^2 \cdot \csc(x)^2 + 1)) \cdot a^2 \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 4 \cdot (-a+b)^{1/2} \cdot a^2 \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + 6 \cdot (-a+b)^{1/2} \cdot a \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 4 \cdot (-a+b)^{1/2} \cdot b^2 \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - \ln(4 \cdot (a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + (-a+b)^{1/2} \cdot (b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + 4 \cdot a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 2 \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + b)^{1/2} - a+b) / ((1-\cos(x))^2 \cdot \csc(x)^2 + 1)) \cdot (b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + 4 \cdot a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 2 \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + b)^{1/2} \cdot a^2 - a \cdot (-a+b)^{1/2} \cdot b + 2 \cdot b^2 \cdot (-a+b)^{1/2} \cdot (b \cdot (1-\cos(x))^4 \cdot \csc(x)^4 + 4 \cdot a \cdot (1-\cos(x))^2 \cdot \csc(x)^2 - 2 \cdot b \cdot (1-\cos(x))^2 \cdot \csc(x)^2 + b) / ((1-\cos(x))^2 \cdot \csc(x)^2 - 1) / (1-\cos(x))^3 \cdot \sin(x)^3 / (1/(1-\cos(x))^2 \cdot (b \cdot (1-\cos(x))^4 \cdot \csc(x)^2 + 4 \cdot a \cdot (1-\cos(x))^2 - 2 \cdot b \cdot (1-\cos(x))^2 + b \cdot \sin(x)^2))^{3/2}
\end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(82) = 164$.

Time = 0.35 (sec) , antiderivative size = 393, normalized size of antiderivative = 4.27

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\left((a^3 \tan(x)^2 + a^2 b) \sqrt{-a + b} \log \left(-\frac{a^2 \tan(x)^4 - 2(3a^2 - 4ab) \tan(x)^2 + a^2 - 8ab + 8b^2 - 4(a \tan(x)^4 + 2 \tan(x)^2 + 1)}{\tan(x)^4 + 2 \tan(x)^2 + 1} \right) \right)}{4(a^4 b - 2a^3 b)}$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="fricas")

[Out] [1/4*((a^3*tan(x)^2 + a^2*b)*sqrt(-a + b)*log(-(a^2*tan(x)^4 - 2*(3*a^2 - 4*a*b)*tan(x)^2 + a^2 - 8*a*b + 8*b^2 - 4*(a*tan(x)^3 - (a - 2*b)*tan(x))*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) + 4*((a^3 - 2*a^2*b + a*b^2)*tan(x)^3 + (a^2*b - 3*a*b^2 + 2*b^3)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(a^4*b - 2*a^3*b^2 + a^2*b^3 + (a^5 - 2*a^4*b + a^3*b^2)*tan(x)^2), 1/2*((a^3*tan(x)^2 + a^2*b)*sqrt(a - b)*arctan(2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)/(a*tan(x)^2 - a + 2*b)) + 2*((a^3 - 2*a^2*b + a*b^2)*tan(x)^3 + (a^2*b - 3*a*b^2 + 2*b^3)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(a^4*b - 2*a^3*b^2 + a^2*b^3 + (a^5 - 2*a^4*b + a^3*b^2)*tan(x)^2)]

Sympy [F]

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\tan^2(x)}{(a + b \cot^2(x))^{\frac{3}{2}}} dx$$

[In] integrate(tan(x)**2/(a+b*cot(x)**2)**(3/2),x)

[Out] Integral(tan(x)**2/(a + b*cot(x)**2)**(3/2), x)

Maxima [F]

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\tan(x)^2}{(b \cot(x)^2 + a)^{\frac{3}{2}}} dx$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate(tan(x)^2/(b*cot(x)^2 + a)^(3/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. 2(82) = 164.

Time = 0.33 (sec) , antiderivative size = 359, normalized size of antiderivative = 3.90

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \frac{\left(a^3 \log\left(-a - 2\sqrt{-a + b}\sqrt{b} + 2b\right) + a^2\sqrt{-a + b}\sqrt{b} \log\left(-a - 2\sqrt{-a + b}\sqrt{b} + 2b\right)\right)}{2\left(a^4\sqrt{-a + b} - a^4\sqrt{b}\right)} + \frac{2\sqrt{-a \cos(x)^2 + b \cos(x)^2 + ab^2 \cos(x)}}{(a^3 - a^2b)(a \cos(x)^2 - b \cos(x)^2 - a)} - \frac{\log\left(\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}\right)^2\right)}{(a - b)\sqrt{-a + b}} - \frac{4\sqrt{-a + b}}{\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}\right)^2} + \frac{2 \operatorname{sgn}(\sin(x))}{2 \operatorname{sgn}(\sin(x))}$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(3/2),x, algorithm="giac")

[Out] 1/2*(a^3*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + a^2*sqrt(-a + b)*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - a^2*b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 2*a^3 - 4*a^2*b + 2*a*sqrt(-a + b)*b^(3/2) - 2*sqrt(-a + b)*b^(5/2) + 2*b^3)*sgn(sin(x))/(a^4*sqrt(-a + b) - a^4*sqrt(b) - 2*a^3*sqrt(-a + b)*b + 2*a^3*b^(3/2) + a^2*sqrt(-a + b)*b^2 - a^2*b^(5/2)) + 1/2*(2*sqrt(-a*cos(x)^2 + b*cos(x)^2 + a)*b^2*cos(x)/((a^3 - a^2*b)*(a*cos(x)^2 - b*cos(x)^2 - a)) - log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2)/((a - b)*sqrt(-a + b)) - 4*sqrt(-a + b)/(((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a)*a))/sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{3/2}} dx = \int \frac{\tan(x)^2}{(b \cot(x)^2 + a)^{3/2}} dx$$

[In] int(tan(x)^2/(a + b*cot(x)^2)^(3/2),x)

[Out] int(tan(x)^2/(a + b*cot(x)^2)^(3/2), x)

3.54 $\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{5/2}} dx$

Optimal result	336
Rubi [A] (verified)	336
Mathematica [C] (verified)	338
Maple [A] (verified)	339
Fricas [B] (verification not implemented)	339
Sympy [F]	340
Maxima [F(-2)]	340
Giac [B] (verification not implemented)	340
Mupad [B] (verification not implemented)	341

Optimal result

Integrand size = 17, antiderivative size = 82

$$\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{5/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} + \frac{a}{3(a-b)b(a+b \cot^2(x))^{3/2}} + \frac{1}{(a-b)^2 \sqrt{a+b \cot^2(x)}}$$

[Out] $-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(5/2)}+1/3*a/(a-b)/b/(a+b*\cot(x)^2)^{(3/2)}+1/(a-b)^2/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 457, 79, 53, 65, 214}

$$\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{5/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} + \frac{a}{3b(a-b)(a+b \cot^2(x))^{3/2}} + \frac{1}{(a-b)^2 \sqrt{a+b \cot^2(x)}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]^3/(a+b*\operatorname{Cot}[x]^2)^{(5/2)},x]$

[Out] $-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a-b]]/(a-b)^{(5/2)})+a/(3*(a-b)*b*(a+b*\operatorname{Cot}[x]^2)^{(3/2)})+1/((a-b)^2*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2])$

Rule 53


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 79

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/
(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c
*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x],
x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || I
ntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
.), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^3}{(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{x}{(1+x)(a+bx)^{5/2}} dx, x, \cot^2(x)\right)\right) \\
&= \frac{a}{3(a-b)b(a+b\cot^2(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1}{(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)}{2(a-b)} \\
&= \frac{a}{3(a-b)b(a+b\cot^2(x))^{3/2}} + \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}} + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)^2} \\
&= \frac{a}{3(a-b)b(a+b\cot^2(x))^{3/2}} + \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)^2b} \\
&= -\frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} + \frac{a}{3(a-b)b(a+b\cot^2(x))^{3/2}} + \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.11 (sec) , antiderivative size = 69, normalized size of antiderivative = 0.84

$$\int \frac{\cot^3(x)}{(a+b\cot^2(x))^{5/2}} dx = \frac{a(a-b) + 3b(a+b\cot^2(x)) \text{Hypergeometric2F1}\left(-\frac{1}{2}, 1, \frac{1}{2}, \frac{a+b\cot^2(x)}{a-b}\right)}{3(a-b)^2b(a+b\cot^2(x))^{3/2}}$$

[In] Integrate[Cot[x]^3/(a + b*Cot[x]^2)^(5/2), x]

[Out] (a*(a - b) + 3*b*(a + b*Cot[x]^2)*Hypergeometric2F1[-1/2, 1, 1/2, (a + b*Cot[x]^2)/(a - b)])/(3*(a - b)^2*b*(a + b*Cot[x]^2)^(3/2))

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.07

method	result	size
derivativedivides	$\frac{1}{3b(a+b \cot(x)^2)^{\frac{3}{2}}} + \frac{1}{3(a-b)(a+b \cot(x)^2)^{\frac{3}{2}}} + \frac{1}{(a-b)^2 \sqrt{a+b \cot(x)^2}} + \frac{\arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)^2 \sqrt{-a+b}}$	88
default	$\frac{1}{3b(a+b \cot(x)^2)^{\frac{3}{2}}} + \frac{1}{3(a-b)(a+b \cot(x)^2)^{\frac{3}{2}}} + \frac{1}{(a-b)^2 \sqrt{a+b \cot(x)^2}} + \frac{\arctan\left(\frac{\sqrt{a+b \cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)^2 \sqrt{-a+b}}$	88

[In] int(cot(x)^3/(a+b*cot(x)^2)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/3/b/(a+b*cot(x)^2)^(3/2)+1/3/(a-b)/(a+b*cot(x)^2)^(3/2)+1/(a-b)^2/(a+b*cot(x)^2)^(1/2)+1/(a-b)^2/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 343 vs. 2(70) = 140.

Time = 0.31 (sec) , antiderivative size = 698, normalized size of antiderivative = 8.51

$$\int \frac{\cot^3(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\left[\frac{3(a^2b + 2ab^2 + b^3 + (a^2b - 2ab^2 + b^3) \cos(2x))^2 - 2(a^2b - b^3) \cos(2x) \sqrt{a-b}}{6(a^5b - a^4b^2 - 2a^3b^3 + 2a^2b^4 + ab^5 - b^6 + (a^5b - 5a^4b^2 + 10a^3b^3 - 10a^2b^4 + 5ab^5 - b^6) \cos(2x))} \sqrt{a-b} \arctan\left(-\frac{\sqrt{-a+b} \sqrt{\frac{(a-b) \cos(2x)}{a-b}}}{a-b}\right) \right]}{3(a^5b - a^4b^2 - 2a^3b^3 + 2a^2b^4 + ab^5 - b^6 + (a^5b - 5a^4b^2 + 10a^3b^3 - 10a^2b^4 + 5ab^5 - b^6) \cos(2x))}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(5/2),x, algorithm="fricas")

[Out] [1/6*(3*(a^2*b + 2*a*b^2 + b^3 + (a^2*b - 2*a*b^2 + b^3)*cos(2*x))^2 - 2*(a^2*b - b^3)*cos(2*x))*sqrt(a-b)*log(sqrt(a-b)*sqrt(((a-b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1) - (a-b)*cos(2*x) + a) + 2*(a^3 + a^2*b + a*b^2 - 3*b^3 + (a^3 + a^2*b - 5*a*b^2 + 3*b^3)*cos(2*x))^2 - 2*(a^3 + a^2*b - 2*a*b^2)*cos(2*x))*sqrt(((a-b)*cos(2*x) - a - b)/(cos(2*x) - 1)) / (a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6 + (a^5*b - 5*a^4*b^2 + 10*a^3*b^3 - 10*a^2*b^4 + 5*a*b^5 - b^6)*cos(2*x))^2 - 2*(a^5*b - 3*a^4*b^2 + 2*a^3*b^3 + 2*a^2*b^4 - 3*a*b^5 + b^6)*cos(2*x)), -1/3*(3*(a^2*b + 2*a*b^2 + b^3 + (a^2*b - 2*a*b^2 + b^3)*cos(2*x))^2 - 2*(a^2*b - b^3)*cos(2*x))

```
)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a - b)) - (a^3 + a^2*b + a*b^2 - 3*b^3 + (a^3 + a^2*b - 5*a*b^2 + 3*b^3)*cos(2*x)^2 - 2*(a^3 + a^2*b - 2*a*b^2)*cos(2*x))*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^5*b - a^4*b^2 - 2*a^3*b^3 + 2*a^2*b^4 + a*b^5 - b^6 + (a^5*b - 5*a^4*b^2 + 10*a^3*b^3 - 10*a^2*b^4 + 5*a*b^5 - b^6)*cos(2*x)^2 - 2*(a^5*b - 3*a^4*b^2 + 2*a^3*b^3 + 2*a^2*b^4 - 3*a*b^5 + b^6)*cos(2*x))]
```

Sympy [F]

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx$$

```
[In] integrate(cot(x)**3/(a+b*cot(x)**2)**(5/2),x)
```

```
[Out] Integral(cot(x)**3/(a + b*cot(x)**2)**(5/2), x)
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx = \text{Exception raised: ValueError}$$

```
[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(5/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 219 vs. 2(70) = 140.

Time = 0.29 (sec) , antiderivative size = 219, normalized size of antiderivative = 2.67

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx = -\frac{\log(|b|) \operatorname{sgn}(\sin(x))}{2(\sqrt{a - ba^2} - 2\sqrt{a - bab} + \sqrt{a - bb^2})} + \frac{\left(\frac{(a^3 + a^2b - 5ab^2 + 3b^3) \sin(x)^2}{a^3b - 3a^2b^2 + 3ab^3 - b^4} + \frac{3(ab^2 - b^3)}{a^3b - 3a^2b^2 + 3ab^3 - b^4}\right) \sin(x)}{(a \sin(x)^2 - b \sin(x)^2 + b)^{3/2}} + \frac{3 \log\left(\left|-\sqrt{a-b} \sin(x) + \sqrt{a \sin(x)^2 - b \sin(x)^2 + b}\right|\right)}{(a^2 - 2ab + b^2)\sqrt{a-b}} + \frac{3 \operatorname{sgn}(\sin(x))}{3 \operatorname{sgn}(\sin(x))}$$

[In] integrate(cot(x)^3/(a+b*cot(x)^2)^(5/2),x, algorithm="giac")

[Out]
$$-1/2*\log(\text{abs}(b))*\text{sgn}(\sin(x))/(\sqrt{a-b})*a^2 - 2*\sqrt{a-b}*a*b + \sqrt{(a-b)*b^2}) + 1/3*((a^3 + a^2*b - 5*a*b^2 + 3*b^3)*\sin(x)^2/(a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4) + 3*(a*b^2 - b^3)/(a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4))*\sin(x)/(a*\sin(x)^2 - b*\sin(x)^2 + b)^{(3/2)} + 3*\log(\text{abs}(-\sqrt{a-b})*\sin(x) + \sqrt{a*\sin(x)^2 - b*\sin(x)^2 + b}))/((a^2 - 2*a*b + b^2)*\sqrt{a-b}))/\text{sgn}(\sin(x))$$

Mupad [B] (verification not implemented)

Time = 16.21 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.07

$$\int \frac{\cot^3(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{\frac{a}{3(a-b)} + \frac{b(b \cot(x)^2 + a)}{(a-b)^2}}{b(b \cot(x)^2 + a)^{3/2}} - \frac{\text{atanh}\left(\frac{\sqrt{b \cot(x)^2 + a}(2a^2 - 4ab + 2b^2)}{2(a-b)^{5/2}}\right)}{(a-b)^{5/2}}$$

[In] int(cot(x)^3/(a + b*cot(x)^2)^(5/2),x)

[Out]
$$(a/(3*(a-b)) + (b*(a + b*\cot(x)^2))/(a-b)^2)/(b*(a + b*\cot(x)^2)^(3/2)) - \text{atanh}(((a + b*\cot(x)^2)^(1/2)*(2*a^2 - 4*a*b + 2*b^2))/(2*(a-b)^(5/2)))/(a-b)^(5/2)$$

$$3.55 \quad \int \frac{\cot^2(x)}{(a+b \cot^2(x))^{5/2}} dx$$

Optimal result	342
Rubi [A] (verified)	342
Mathematica [C] (warning: unable to verify)	344
Maple [A] (verified)	345
Fricas [B] (verification not implemented)	345
Sympy [F]	346
Maxima [F(-2)]	346
Giac [B] (verification not implemented)	346
Mupad [F(-1)]	347

Optimal result

Integrand size = 17, antiderivative size = 94

$$\int \frac{\cot^2(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{5/2}} - \frac{\cot(x)}{3(a-b)(a+b \cot^2(x))^{3/2}} - \frac{(2a+b) \cot(x)}{3a(a-b)^2 \sqrt{a+b \cot^2(x)}}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/(a-b)^{(5/2)}-1/3*\cot(x)/(a-b)/(a+b*\cot(x)^2)^{(3/2)}-1/3*(2*a+b)*\cot(x)/a/(a-b)^2/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.353$, Rules used = {3751, 482, 541, 12, 385, 209}

$$\int \frac{\cot^2(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{5/2}} - \frac{(2a+b) \cot(x)}{3a(a-b)^2 \sqrt{a+b \cot^2(x)}} - \frac{\cot(x)}{3(a-b)(a+b \cot^2(x))^{3/2}}$$

[In] $\text{Int}[\text{Cot}[x]^2/(a+b*\text{Cot}[x]^2)^{(5/2)},x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a-b]*\text{Cot}[x])/(\text{Sqrt}[a+b*\text{Cot}[x]^2])]/(a-b)^{(5/2)}-\text{Cot}[x]/(3*(a-b)*(a+b*\text{Cot}[x]^2)^{(3/2)})-((2*a+b)*\text{Cot}[x])/(3*a*(a-b)^2*\text{Sqrt}[a+b*\text{Cot}[x]^2])]$

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 385

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)/((c_) + (d_)*(x_)^(n_)), x_Symbol] := Su
bst[Int[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^(1/n)] /; FreeQ[{a, b
, c, d}, x] && NeQ[b*c - a*d, 0] && EqQ[n*p + 1, 0] && IntegerQ[n]
```

Rule 482

```
Int[((e_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_
))^(-q_), x_Symbol] := Simp[e^(n - 1)*(e*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*
((c + d*x^n)^(q + 1)/(n*(b*c - a*d)*(p + 1))), x] - Dist[e^n/(n*(b*c - a*d)
*(p + 1)), Int[(e*x)^(m - n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(m -
n + 1) + d*(m + n*(p + q + 1) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e,
q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GeQ[n, m - n +
1] && GtQ[m - n + 1, 0] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]
```

Rule 541

```
Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_)*((e_) + (f
_)*(x_)^(n_)), x_Symbol] := Simp[(-(b*e - a*f))*x*(a + b*x^n)^(p + 1)*((c
+ d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(
p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*
c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x] /; Fre
eQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_))*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^2}{(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(x)\right) \\
&= -\frac{\cot(x)}{3(a-b)(a+b\cot^2(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1-2x^2}{(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right)}{3(a-b)} \\
&= -\frac{\cot(x)}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{(2a+b)\cot(x)}{3a(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{3a}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{3a(a-b)^2} \\
&= -\frac{\cot(x)}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{(2a+b)\cot(x)}{3a(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{(a-b)^2} \\
&= -\frac{\cot(x)}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{(2a+b)\cot(x)}{3a(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{(a-b)^2} \\
&= \frac{\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b\cot^2(x)}}\right)}{(a-b)^{5/2}} - \frac{\cot(x)}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{(2a+b)\cot(x)}{3a(a-b)^2\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 6.34 (sec) , antiderivative size = 197, normalized size of antiderivative = 2.10

$$\int \frac{\cot^2(x)}{(a+b\cot^2(x))^{5/2}} dx = \frac{\cos(x) \left(-12(a-b)^3 \cos^3(x) \cot(x) (a+b\cot^2(x)) \text{Hypergeometric2F1}\left(2, 2, \frac{9}{2}, \frac{(a-b)\cos^2(x)}{a}\right) - (35a(5a+2b\cot^2(x))\sin(x)(a((a-b)\cos^2(x))^{5/2}) - 12(a-b)^3 \cos^3(x) \cot(x) (a+b\cot^2(x)) \text{Hypergeometric2F1}\left(2, 2, \frac{9}{2}, \frac{(a-b)\cos^2(x)}{a}\right) \right)}{(a-b)^{5/2}}$$

[In] Integrate[Cot[x]^2/(a + b*Cot[x]^2)^(5/2), x]

[Out] (Cos[x]*(-12*(a - b)^3*Cos[x]^3*Cot[x]*(a + b*Cot[x]^2)*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a] - (35*a*(5*a + 2*b*Cot[x]^2)*Sin[x]*(a*((a - b)*Cos[x]^2)/a)^(5/2) - 12*(a - b)^3*Cos[x]^3*Cot[x]*(a + b*Cot[x]^2)*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a])/(a - b)^(5/2)

- 4*b)*Csc[x]^2 - 3*a*Sec[x]^2)*Sqrt[((a - b)*Cos[x]^2*(a + b*Cot[x]^2)*Sin[x]^2)/a^2] + 3*ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*(b*Cot[x] + a*Tan[x])^2)/Sqrt[((a - b)*Cos[x]^2*(a + b*Cot[x]^2)*Sin[x]^2)/a^2])/(315*a^3*(a - b)^2*(a + b*Cot[x]^2)^(3/2))

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.71

method	result
derivativedivides	$-\frac{\cot(x)}{3a(a+b\cot(x)^2)^{\frac{3}{2}}} - \frac{2\cot(x)}{3a^2\sqrt{a+b\cot(x)^2}} - \frac{b\left(\frac{\cot(x)}{3a(a+b\cot(x)^2)^{\frac{3}{2}}} + \frac{2\cot(x)}{3a^2\sqrt{a+b\cot(x)^2}}\right)}{a-b} - \frac{b\cot(x)}{(a-b)^2a\sqrt{a+b\cot(x)^2}}$
default	$-\frac{\cot(x)}{3a(a+b\cot(x)^2)^{\frac{3}{2}}} - \frac{2\cot(x)}{3a^2\sqrt{a+b\cot(x)^2}} - \frac{b\left(\frac{\cot(x)}{3a(a+b\cot(x)^2)^{\frac{3}{2}}} + \frac{2\cot(x)}{3a^2\sqrt{a+b\cot(x)^2}}\right)}{a-b} - \frac{b\cot(x)}{(a-b)^2a\sqrt{a+b\cot(x)^2}}$

[In] int(cot(x)^2/(a+b*cot(x)^2)^(5/2),x,method=_RETURNVERBOSE)

[Out] -1/3*cot(x)/a/(a+b*cot(x)^2)^(3/2)-2/3/a^2*cot(x)/(a+b*cot(x)^2)^(1/2)-1/(a-b)*b*(1/3*cot(x)/a/(a+b*cot(x)^2)^(3/2)+2/3/a^2*cot(x)/(a+b*cot(x)^2)^(1/2))-1/(a-b)^2*b*cot(x)/a/(a+b*cot(x)^2)^(1/2)+1/(a-b)^3*(b^4*(a-b))^(1/2)/b^2*arctan(b^2*(a-b)/(b^4*(a-b))^(1/2)/(a+b*cot(x)^2)^(1/2)*cot(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 339 vs. 2(80) = 160.

Time = 0.39 (sec) , antiderivative size = 720, normalized size of antiderivative = 7.66

$$\int \frac{\cot^2(x)}{(a+b\cot^2(x))^{5/2}} dx = \left[\frac{3(a^3+2a^2b+ab^2+(a^3-2a^2b+ab^2)\cos(2x)^2-2(a^3-ab^2)\cos(2x))\sqrt{-}}{12(a^6} \right.$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="fricas")

[Out] [-1/12*(3*(a^3+2*a^2*b+a*b^2+(a^3-2*a^2*b+a*b^2)*cos(2*x))^2-2*(a^3-a*b^2)*cos(2*x))*sqrt(-a+b)*log(-2*(a^2-2*a*b+b^2)*cos(2*x)^2+2*((a-b)*cos(2*x)-b)*sqrt(-a+b)*sqrt(((a-b)*cos(2*x)-a-b)/(cos(2*x)-1))*sin(2*x)+a^2-2*b^2+4*(a*b-b^2)*cos(2*x))+4*(3*a^3-a^2*b-a*b^2-b^3-(3*a^3-5*a^2*b+a*b^2+b^3)*cos(2*x))*sqrt(((a-b)*cos(2*x)-a-b)/(cos(2*x)-1))*sin(2*x))/(a^6-a^5*b-2*a^4*b^2+2*a^3*b^3+a^2*b^4-a*b^5+(a^6-5*a^5*b+10*a^4*b^2-10*a^3*b^3+5*a^

$2*b^4 - a*b^5)*\cos(2*x)^2 - 2*(a^6 - 3*a^5*b + 2*a^4*b^2 + 2*a^3*b^3 - 3*a^2*b^4 + a*b^5)*\cos(2*x)$, $1/6*(3*(a^3 + 2*a^2*b + a*b^2 + (a^3 - 2*a^2*b + a*b^2)*\cos(2*x)^2 - 2*(a^3 - a*b^2)*\cos(2*x))*\sqrt{a - b}*\arctan(-\sqrt{a - b}*\sqrt{((a - b)*\cos(2*x) - a - b)/(\cos(2*x) - 1)}*\sin(2*x)/((a - b)*\cos(2*x) - b)) - 2*(3*a^3 - a^2*b - a*b^2 - b^3 - (3*a^3 - 5*a^2*b + a*b^2 + b^3)*\cos(2*x))*\sqrt{((a - b)*\cos(2*x) - a - b)/(\cos(2*x) - 1)}*\sin(2*x))/(a^6 - a^5*b - 2*a^4*b^2 + 2*a^3*b^3 + a^2*b^4 - a*b^5 + (a^6 - 5*a^5*b + 10*a^4*b^2 - 10*a^3*b^3 + 5*a^2*b^4 - a*b^5)*\cos(2*x)^2 - 2*(a^6 - 3*a^5*b + 2*a^4*b^2 + 2*a^3*b^3 - 3*a^2*b^4 + a*b^5)*\cos(2*x))]$

Sympy [F]

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\cot^2(x)}{(a + b \cot^2(x))^{\frac{5}{2}}} dx$$

[In] integrate(cot(x)**2/(a+b*cot(x)**2)**(5/2),x)

[Out] Integral(cot(x)**2/(a + b*cot(x)**2)**(5/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(b-a>0)', see 'assume?' for more details)Is

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 281 vs. 2(80) = 160.

Time = 0.34 (sec) , antiderivative size = 281, normalized size of antiderivative = 2.99

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{\left(3a\sqrt{b} \log\left(\left|-\sqrt{-a+b} + \sqrt{b}\right|\right) + 2a\sqrt{-a+b} + \sqrt{-a+bb}\right) \operatorname{sgn}(\sin(x))}{3\left(a^3\sqrt{-a+b}\sqrt{b} - 2a^2\sqrt{-a+bb}^{\frac{3}{2}} + a\sqrt{-a+bb}^{\frac{5}{2}}\right)} + \frac{\left(\frac{(3a^3-5a^2b+ab^2+b^3)\cos(x)^2}{a^4-3a^3b+3a^2b^2-ab^3} - \frac{3(a^3-a^2b)}{a^4-3a^3b+3a^2b^2-ab^3}\right)\cos(x)}{(a\cos(x)^2-b\cos(x)^2-a)\sqrt{-a\cos(x)^2+b\cos(x)^2+a}} + \frac{3\log\left(\left|-\sqrt{-a+b}\cos(x)+\sqrt{-a\cos(x)^2+b\cos(x)^2+a}\right|\right)}{(a^2-2ab+b^2)\sqrt{-a+b}}$$

$3 \operatorname{sgn}(\sin(x))$

[In] integrate(cot(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="giac")

[Out] $\frac{1}{3}*(3*a*\sqrt{b}*\log(\text{abs}(-\sqrt{-a+b} + \sqrt{b})) + 2*a*\sqrt{-a+b} + \sqrt{-a+b}*b)*\text{sgn}(\sin(x))/(a^3*\sqrt{-a+b}*\sqrt{b} - 2*a^2*\sqrt{-a+b}*b^{3/2} + a*\sqrt{-a+b}*b^{5/2}) - \frac{1}{3}*(((3*a^3 - 5*a^2*b + a*b^2 + b^3)*\cos(x)^2/(a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3) - 3*(a^3 - a^2*b)/(a^4 - 3*a^3*b + 3*a^2*b^2 - a*b^3))*\cos(x)/((a*\cos(x)^2 - b*\cos(x)^2 - a)*\sqrt{-a*\cos(x)^2 + b*\cos(x)^2 + a}) + 3*\log(\text{abs}(-\sqrt{-a+b})*\cos(x) + \sqrt{-a*\cos(x)^2 + b*\cos(x)^2 + a}))/((a^2 - 2*a*b + b^2)*\sqrt{-a+b}))/\text{sgn}(\sin(x))$

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\cot(x)^2}{(b \cot(x)^2 + a)^{5/2}} dx$$

[In] int(cot(x)^2/(a + b*cot(x)^2)^(5/2),x)

[Out] int(cot(x)^2/(a + b*cot(x)^2)^(5/2), x)

$$3.56 \quad \int \frac{\cot(x)}{(a+b \cot^2(x))^{5/2}} dx$$

Optimal result	348
Rubi [A] (verified)	348
Mathematica [C] (verified)	350
Maple [A] (verified)	350
Fricas [B] (verification not implemented)	351
Sympy [A] (verification not implemented)	351
Maxima [F(-2)]	352
Giac [B] (verification not implemented)	352
Mupad [B] (verification not implemented)	353

Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\cot(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} - \frac{1}{3(a-b)(a+b \cot^2(x))^{3/2}} - \frac{1}{(a-b)^2 \sqrt{a+b \cot^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(5/2)}-1/3/(a-b)/(a+b*\cot(x)^2)^{(3/2)}-1/(a-b)^2/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3751, 455, 53, 65, 214}

$$\int \frac{\cot(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} - \frac{1}{(a-b)^2 \sqrt{a+b \cot^2(x)}} - \frac{1}{3(a-b)(a+b \cot^2(x))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]/(a+b*\operatorname{Cot}[x]^2)^{(5/2)},x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a-b]]/(a-b)^{(5/2)}-1/(3*(a-b)*(a+b*\operatorname{Cot}[x]^2)^{(3/2)})-1/((a-b)^2*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2])$

Rule 53

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x]
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && LtQ[m, -1] && !(LtQ
[n, -1] && (EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && I
ntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 455

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.)*((c_) + (d_.)*(x_)^(n_))^(q_.
), x_Symbol] := Dist[1/n, Subst[Int[(a + b*x)^p*(c + d*x)^q, x], x, x^n], x]
] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[b*c - a*d, 0] && EqQ[m - n +
1, 0]
```

Rule 3751

```
Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) +
(f_.)*(x_)])^(n_))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)(a+bx)^{5/2}} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{1}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)}{2(a-b)}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)^2} \\
&= -\frac{1}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{1-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)^2b} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} - \frac{1}{3(a-b)(a+b\cot^2(x))^{3/2}} - \frac{1}{(a-b)^2\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.60

$$\int \frac{\cot(x)}{(a+b\cot^2(x))^{5/2}} dx = -\frac{\text{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b\cot^2(x)}{a-b}\right)}{3(a-b)(a+b\cot^2(x))^{3/2}}$$

[In] Integrate[Cot[x]/(a + b*Cot[x]^2)^(5/2), x]

[Out] -1/3*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Cot[x]^2)/(a - b)]/((a - b)*(a + b*Cot[x]^2)^(3/2))

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.96

method	result	size
derivativedivides	$-\frac{1}{3(a-b)(a+b\cot(x)^2)^{3/2}} - \frac{1}{(a-b)^2\sqrt{a+b\cot(x)^2}} - \frac{\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)^2\sqrt{-a+b}}$	75
default	$-\frac{1}{3(a-b)(a+b\cot(x)^2)^{3/2}} - \frac{1}{(a-b)^2\sqrt{a+b\cot(x)^2}} - \frac{\arctan\left(\frac{\sqrt{a+b\cot(x)^2}}{\sqrt{-a+b}}\right)}{(a-b)^2\sqrt{-a+b}}$	75

[In] int(cot(x)/(a+b*cot(x)^2)^(5/2), x, method=_RETURNVERBOSE)

[Out] -1/3/(a-b)/(a+b*cot(x)^2)^(3/2)-1/(a-b)^2/(a+b*cot(x)^2)^(1/2)-1/(a-b)^2/(-a+b)^(1/2)*arctan((a+b*cot(x)^2)^(1/2)/(-a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 307 vs. 2(66) = 132.

Time = 0.34 (sec) , antiderivative size = 627, normalized size of antiderivative = 8.04

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx = \left[\frac{3 \left((a^2 - 2ab + b^2) \cos(2x)^2 + a^2 + 2ab + b^2 - 2(a^2 - b^2) \cos(2x) \right) \sqrt{a-b} \log}{6 (a^5 - a^4b - 2a^3b^2 + 2a^2b^3 + ab^4)} \right]$$

[In] integrate(cot(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="fricas")

[Out] [1/6*(3*((a^2 - 2*a*b + b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*x))*sqrt(a - b)*log(-sqrt(a - b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1))*(cos(2*x) - 1) - (a - b)*cos(2*x) + a) - 4*(2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 + 2*a^2 - a*b - b^2 - (4*a^2 - 5*a*b + b^2)*cos(2*x))*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*cos(2*x)^2 - 2*(a^5 - 3*a^4*b + 2*a^3*b^2 + 2*a^2*b^3 - 3*a*b^4 + b^5)*cos(2*x)), 1/3*(3*((a^2 - 2*a*b + b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*x))*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a - b) - 2*(2*(a^2 - 2*a*b + b^2)*cos(2*x)^2 + 2*a^2 - a*b - b^2 - (4*a^2 - 5*a*b + b^2)*cos(2*x))*sqrt(((a - b)*cos(2*x) - a - b)/(cos(2*x) - 1)))/(a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5 + (a^5 - 5*a^4*b + 10*a^3*b^2 - 10*a^2*b^3 + 5*a*b^4 - b^5)*cos(2*x)^2 - 2*(a^5 - 3*a^4*b + 2*a^3*b^2 + 2*a^2*b^3 - 3*a*b^4 + b^5)*cos(2*x))]

Sympy [A] (verification not implemented)

Time = 7.39 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.41

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx = \begin{cases} 2 \left(\frac{b}{6(a-b)(a+b \cot^2(x))^{\frac{3}{2}}} + \frac{b}{2(a-b)^2 \sqrt{a+b \cot^2(x)}} + \frac{b \operatorname{atan} \left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{-a+b}} \right)}{2\sqrt{-a+b}(a-b)^2} \right) & \text{for } b \neq 0 \\ \begin{cases} \infty \cot^2(x) & \text{for } a^{\frac{5}{2}} = 0 \\ \frac{\log(2a^{\frac{5}{2}} \cot^2(x) + 2a^{\frac{5}{2}})}{2a^{\frac{5}{2}}} & \text{otherwise} \end{cases} & \text{otherwise} \end{cases}$$

[In] integrate(cot(x)/(a+b*cot(x)**2)**(5/2),x)

```
[Out] -Piecewise((2*(b/(6*(a - b)*(a + b*cot(x)**2)**(3/2)) + b/(2*(a - b)**2*sqrt(a + b*cot(x)**2)) + b*atan(sqrt(a + b*cot(x)**2)/sqrt(-a + b))/(2*sqrt(-a + b)*(a - b)**2))/b, Ne(b, 0)), (Piecewise((zoo*cot(x)**2, Eq(a**(5/2), 0)), (log(2*a**(5/2)*cot(x)**2 + 2*a**(5/2))/(2*a**(5/2))), True)), True))
```

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx = \text{Exception raised: ValueError}$$

```
[In] integrate(cot(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a-4*b>0)', see 'assume?' for more detail)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(66) = 132.
Time = 0.32 (sec) , antiderivative size = 215, normalized size of antiderivative = 2.76

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{\log(|b|) \operatorname{sgn}(\sin(x))}{2(\sqrt{a-b}a^2 - 2\sqrt{a-b}ab + \sqrt{a-b}b^2)} + \frac{\left(\frac{4(a^2b-2ab^2+b^3)\sin(x)^2}{a^3b-3a^2b^2+3ab^3-b^4} + \frac{3(ab^2-b^3)}{a^3b-3a^2b^2+3ab^3-b^4}\right)\sin(x)}{(a\sin(x)^2 - b\sin(x)^2 + b)^{3/2}} + \frac{3 \log\left(\left|-\sqrt{a-b}\sin(x) + \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right|\right)}{(a^2 - 2ab + b^2)\sqrt{a-b}}$$

$$3 \operatorname{sgn}(\sin(x))$$

```
[In] integrate(cot(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="giac")
```

```
[Out] 1/2*log(abs(b))*sgn(sin(x))/(sqrt(a - b)*a^2 - 2*sqrt(a - b)*a*b + sqrt(a - b)*b^2) - 1/3*((4*(a^2*b - 2*a*b^2 + b^3)*sin(x)^2/(a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4) + 3*(a*b^2 - b^3)/(a^3*b - 3*a^2*b^2 + 3*a*b^3 - b^4))*sin(x)/(a*sin(x)^2 - b*sin(x)^2 + b)^(3/2) + 3*log(abs(-sqrt(a - b)*sin(x) + sqrt(a*sin(x)^2 - b*sin(x)^2 + b)))/((a^2 - 2*a*b + b^2)*sqrt(a - b))/sgn(sin(x))
```


Mupad [B] (verification not implemented)

Time = 17.27 (sec) , antiderivative size = 82, normalized size of antiderivative = 1.05

$$\int \frac{\cot(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{\operatorname{atanh}\left(\frac{\sqrt{b \cot(x)^2 + a} (2a^2 - 4ab + 2b^2)}{2(a-b)^{5/2}}\right)}{(a-b)^{5/2}} - \frac{\frac{1}{3(a-b)} + \frac{b \cot(x)^2 + a}{(a-b)^2}}{(b \cot(x)^2 + a)^{3/2}}$$

`[In] int(cot(x)/(a + b*cot(x)^2)^(5/2),x)`

```
[Out] atanh(((a + b*cot(x)^2)^(1/2)*(2*a^2 - 4*a*b + 2*b^2))/(2*(a - b)^(5/2)))/(a - b)^(5/2) - (1/(3*(a - b)) + (a + b*cot(x)^2)/(a - b)^2)/(a + b*cot(x)^2)^(3/2)
```

$$3.57 \quad \int \frac{\tan(x)}{(a+b \cot^2(x))^{5/2}} dx$$

Optimal result	354
Rubi [A] (verified)	354
Mathematica [C] (verified)	357
Maple [F]	357
Fricas [B] (verification not implemented)	357
Sympy [F]	358
Maxima [F]	358
Giac [B] (verification not implemented)	359
Mupad [B] (verification not implemented)	359

Optimal result

Integrand size = 15, antiderivative size = 118

$$\int \frac{\tan(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} + \frac{b}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(2a-b)b}{a^2(a-b)^2 \sqrt{a+b \cot^2(x)}}$$

[Out] $\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/a^{(1/2)})/a^{(5/2)}-\operatorname{arctanh}((a+b*\cot(x)^2)^{(1/2)}/(a-b)^{(1/2)})/(a-b)^{(5/2)}+1/3*b/a/(a-b)/(a+b*\cot(x)^2)^{(3/2)}+(2*a-b)*b/a^2/(a-b)^2/(a+b*\cot(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3751, 457, 87, 157, 162, 65, 214}

$$\int \frac{\tan(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} + \frac{b(2a-b)}{a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} + \frac{b}{3a(a-b)(a+b \cot^2(x))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Tan}[x]/(a+b*\operatorname{Cot}[x]^2)^{(5/2)}, x]$

[Out] $\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a]]/a^{(5/2)}-\operatorname{ArcTanh}[\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2]/\operatorname{Sqrt}[a-b]]/(a-b)^{(5/2)}+b/(3*a*(a-b)*(a+b*\operatorname{Cot}[x]^2)^{(3/2)})+((2*a-b)*b)/(a^2*(a-b)^2*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^2])$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 87

```
Int[((e_.) + (f_.)*(x_))^(p_)/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))),
x_Symbol] := Simp[f*(e + f*x)^(p + 1)/((p + 1)*(b*e - a*f)*(d*e - c*f)),
x] + Dist[1/((b*e - a*f)*(d*e - c*f)), Int[(b*d*e - b*c*f - a*d*f - b*d*f*x
)*(e + f*x)^(p + 1)/((a + b*x)*(c + d*x)), x], x] /; FreeQ[{a, b, c, d, e
, f}, x] && LtQ[p, -1]
```

Rule 157

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n + 1)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && Integ
ersQ[2*m, 2*n, 2*p]
```

Rule 162

```
Int[(((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)))/(((a_.) + (b_.)*(x_))*
((c_.) + (d_.)*(x_))), x_Symbol] := Dist[(b*g - a*h)/(b*c - a*d), Int[(e +
f*x)^p/(a + b*x), x], x] - Dist[(d*g - c*h)/(b*c - a*d), Int[(e + f*x)^p/(c
+ d*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, h}, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 457

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_
), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p
*(c + d*x)^q, x], x, x^n], x] /; FreeQ[{a, b, c, d, m, n, p, q}, x] && NeQ[
b*c - a*d, 0] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_))*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{1}{x(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{x(1+x)(a+bx)^{5/2}} dx, x, \cot^2(x)\right)\right) \\
&= \frac{b}{3a(a-b)(a+b\cot^2(x))^{3/2}} - \frac{\text{Subst}\left(\int \frac{a-b-bx}{x(1+x)(a+bx)^{3/2}} dx, x, \cot^2(x)\right)}{2a(a-b)} \\
&= \frac{b}{3a(a-b)(a+b\cot^2(x))^{3/2}} + \frac{(2a-b)b}{a^2(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-\frac{1}{2}(a-b)^2 + \frac{1}{2}(2a-b)bx}{x(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{a^2(a-b)^2} \\
&= \frac{b}{3a(a-b)(a+b\cot^2(x))^{3/2}} + \frac{(2a-b)b}{a^2(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2a^2} + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx}} dx, x, \cot^2(x)\right)}{2(a-b)^2} \\
&= \frac{b}{3a(a-b)(a+b\cot^2(x))^{3/2}} + \frac{(2a-b)b}{a^2(a-b)^2\sqrt{a+b\cot^2(x)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{a^2b} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{1 - \frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b\cot^2(x)}\right)}{(a-b)^2b} \\
&= \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a}}\right)}{a^{5/2}} - \frac{\text{arctanh}\left(\frac{\sqrt{a+b\cot^2(x)}}{\sqrt{a-b}}\right)}{(a-b)^{5/2}} \\
&\quad + \frac{b}{3a(a-b)(a+b\cot^2(x))^{3/2}} + \frac{(2a-b)b}{a^2(a-b)^2\sqrt{a+b\cot^2(x)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 0.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.66

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{a \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \cot^2(x)}{a-b}\right) + (-a + b) \operatorname{Hypergeometric2F1}\left(-\frac{3}{2}, 1, -\frac{1}{2}, \frac{a+b \cot^2(x)}{a-b}\right)}{3a(a-b)(a+b \cot^2(x))^{3/2}}$$

[In] Integrate[Tan[x]/(a + b*Cot[x]^2)^(5/2),x]

[Out] (a*Hypergeometric2F1[-3/2, 1, -1/2, (a + b*Cot[x]^2)/(a - b)] + (-a + b)*Hypergeometric2F1[-3/2, 1, -1/2, 1 + (b*Cot[x]^2)/a])/(3*a*(a - b)*(a + b*Cot[x]^2)^(3/2))

Maple [F]

$$\int \frac{\tan(x)}{(a + b \cot(x)^2)^{5/2}} dx$$

[In] int(tan(x)/(a+b*cot(x)^2)^(5/2),x)

[Out] int(tan(x)/(a+b*cot(x)^2)^(5/2),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 369 vs. 2(100) = 200.

Time = 0.49 (sec) , antiderivative size = 1531, normalized size of antiderivative = 12.97

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="fricas")

[Out] [1/6*(3*(a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 + (a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*tan(x)^4 + 2*(a^4*b - 3*a^3*b^2 + 3*a^2*b^3 - a*b^4)*tan(x)^2)*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b) + 3*(a^5*tan(x)^4 + 2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(a - b)*log(((2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b)/(tan(x)^2 + 1)) + 2*((7*a^4*b - 11*a^3*b^2 + 4*a^2*b^3)*tan(x)^4 + 3*(2*a^3*b^2 - 3*a^2*b^3 + a*b^4)*tan(x)^2)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4)*tan(x)^2), -1/6*(6*(a^5*tan(x)^4 + 2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/(a - b)) - 3*(a^3*b^2 - 3*a^2*b^3 + 3

```

*a*b^4 - b^5 + (a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*tan(x)^4 + 2*(a^4*b -
3*a^3*b^2 + 3*a^2*b^3 - a*b^4)*tan(x)^2)*sqrt(a)*log(2*a*tan(x)^2 + 2*sqrt(
a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2 + b) - 2*((7*a^4*b - 11*a^3*b^2
+ 4*a^2*b^3)*tan(x)^4 + 3*(2*a^3*b^2 - 3*a^2*b^3 + a*b^4)*tan(x)^2)*sqrt((
a*tan(x)^2 + b)/tan(x)^2))/(a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^
8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*
b^3 - a^4*b^4)*tan(x)^2), -1/6*(6*(a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 + (a
^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*tan(x)^4 + 2*(a^4*b - 3*a^3*b^2 + 3*a^2
*b^3 - a*b^4)*tan(x)^2)*sqrt(-a)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(
x)^2)/a) - 3*(a^5*tan(x)^4 + 2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(a - b)*log(((
2*a - b)*tan(x)^2 - 2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)^2
+ b)/(tan(x)^2 + 1)) - 2*((7*a^4*b - 11*a^3*b^2 + 4*a^2*b^3)*tan(x)^4 + 3*(
2*a^3*b^2 - 3*a^2*b^3 + a*b^4)*tan(x)^2)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(
a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^8 - 3*a^7*b + 3*a^6*b^2 - a^
5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4)*tan(x)^2), -1
/3*(3*(a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 + (a^5 - 3*a^4*b + 3*a^3*b^2 - a
^2*b^3)*tan(x)^4 + 2*(a^4*b - 3*a^3*b^2 + 3*a^2*b^3 - a*b^4)*tan(x)^2)*sqrt
(-a)*arctan(sqrt(-a)*sqrt((a*tan(x)^2 + b)/tan(x)^2)/a) + 3*(a^5*tan(x)^4 +
2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(-a + b)*arctan(-sqrt(-a + b)*sqrt((a*tan(
x)^2 + b)/tan(x)^2)/(a - b)) - ((7*a^4*b - 11*a^3*b^2 + 4*a^2*b^3)*tan(x)^4
+ 3*(2*a^3*b^2 - 3*a^2*b^3 + a*b^4)*tan(x)^2)*sqrt((a*tan(x)^2 + b)/tan(x)
^2))/(a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^8 - 3*a^7*b + 3*a^6*b^
2 - a^5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4)*tan(x)^
2)]

```

Sympy [F]

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx$$

```
[In] integrate(tan(x)/(a+b*cot(x)**2)**(5/2),x)
```

```
[Out] Integral(tan(x)/(a + b*cot(x)**2)**(5/2), x)
```

Maxima [F]

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\tan(x)}{(b \cot^2(x) + a)^{5/2}} dx$$

```
[In] integrate(tan(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(tan(x)/(b*cot(x)^2 + a)^(5/2), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 483 vs. $2(100) = 200$.

Time = 0.38 (sec) , antiderivative size = 483, normalized size of antiderivative = 4.09

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx =$$

$$\frac{\left(2a^3 \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right) - 6a^2b \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right) + 6ab^2 \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right) - 2b^3 \arctan\left(-\frac{a-b}{\sqrt{-a^2+ab}}\right)\right)}{2\left(\sqrt{-a^2+ab}\sqrt{a-b}a^4 - 2\sqrt{-a^2+ab}\sqrt{a-b}a^3b + \sqrt{-a^2+ab}\sqrt{a-b}ba^3 - 2\sqrt{-a^2+ab}\sqrt{a-b}b^3\right)}$$

$$+ \frac{2\left(\frac{(7a^5b^2 - 17a^4b^3 + 13a^3b^4 - 3a^2b^5)\sin(x)^2}{a^7b - 3a^6b^2 + 3a^5b^3 - a^4b^4} + \frac{3(2a^4b^3 - 3a^3b^4 + a^2b^5)}{a^7b - 3a^6b^2 + 3a^5b^3 - a^4b^4}\right)\sin(x)}{(a\sin(x)^2 - b\sin(x)^2 + b)^{3/2}} + \frac{3\log\left(\left(\sqrt{a-b}\sin(x) - \sqrt{a\sin(x)^2 - b\sin(x)^2 + b}\right)^2\right)}{(a^2 - 2ab + b^2)\sqrt{a-b}} + \frac{6\sqrt{a-b}}{6\operatorname{sgn}(\sin(x))}$$

[In] integrate(tan(x)/(a+b*cot(x)^2)^(5/2),x, algorithm="giac")

[Out] $-1/2*(2*a^3*\arctan(-(a - b)/\sqrt{-a^2 + a*b}) - 6*a^2*b*\arctan(-(a - b)/\sqrt{-a^2 + a*b}) + 6*a*b^2*\arctan(-(a - b)/\sqrt{-a^2 + a*b}) - 2*b^3*\arctan(-(a - b)/\sqrt{-a^2 + a*b}) + \sqrt{-a^2 + a*b}*a^2*\log(b))*\operatorname{sgn}(\sin(x))/(\sqrt{-a^2 + a*b}*\sqrt{a - b}*a^4 - 2*\sqrt{-a^2 + a*b}*\sqrt{a - b}*a^3*b + \sqrt{-a^2 + a*b}*\sqrt{a - b}*a^2*b^2) + 1/6*(2*((7*a^5*b^2 - 17*a^4*b^3 + 13*a^3*b^4 - 3*a^2*b^5)*\sin(x)^2/(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4) + 3*(2*a^4*b^3 - 3*a^3*b^4 + a^2*b^5)/(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4))*\sin(x)/(a*\sin(x)^2 - b*\sin(x)^2 + b)^{3/2} + 3*\log((\sqrt{a - b}*\sin(x) - \sqrt{a*\sin(x)^2 - b*\sin(x)^2 + b})^2)/((a^2 - 2*a*b + b^2)*\sqrt{a - b}) + 6*\sqrt{a - b}*\arctan(1/2*((\sqrt{a - b}*\sin(x) - \sqrt{a*\sin(x)^2 - b*\sin(x)^2 + b})^2 - 2*a + b)/\sqrt{-a^2 + a*b}))/(\sqrt{-a^2 + a*b}*a^2))/\operatorname{sgn}(\sin(x))$

Mupad [B] (verification not implemented)

Time = 13.68 (sec) , antiderivative size = 2817, normalized size of antiderivative = 23.87

$$\int \frac{\tan(x)}{(a + b \cot^2(x))^{5/2}} dx = \text{Too large to display}$$

[In] int(tan(x)/(a + b*cot(x)^2)^(5/2),x)

[Out] $\operatorname{atanh}\left(\frac{(2*a^5*b^{13}*(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}*(2*a^3*b^{13} - 22*a^4*b^{12} + 110*a^5*b^{11} - 330*a^6*b^{10} + 660*a^7*b^9 - 922*a^8*b^8 + 912*a^9*b^7 - 630*a^{10}*b^6 + 290*a^{11}*b^5 - 80*a^{12}*b^4 + 10*a^{13}*b^3)) - (22*a^6*b^{12}*(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}*(2*a^3*b^{13} - 22*a^4*b^{12} + 110*a^5*b^{11} - 330*a^6*b^{10} + 660*a^7*b^9 - 922*a^8*b^8 + 912*a^9*b^7 - 630*a^{10}*b^6))}{(a^5)^{(1/2)}*(2*a^3*b^{13} - 22*a^4*b^{12} + 110*a^5*b^{11} - 330*a^6*b^{10} + 660*a^7*b^9 - 922*a^8*b^8 + 912*a^9*b^7 - 630*a^{10}*b^6)}\right)$

$$\begin{aligned}
&^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) + (110a^7b^{11}(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) - (330a^8b^{10}(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) + (660a^9b^9(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) - (922a^{10}b^8(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) + (912a^{11}b^7(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) - (630a^{12}b^6(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) + (290a^{13}b^5(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) - (80a^{14}b^4(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)) + (10a^{15}b^3(a + b/\tan(x)^2)^{(1/2)})/((a^5)^{(1/2)}(2a^3b^{13} - 22a^4b^{12} + 110a^5b^{11} - 330a^6b^{10} + 660a^7b^9 - 922a^8b^8 + 912a^9b^7 - 630a^{10}b^6 + 290a^{11}b^5 - 80a^{12}b^4 + 10a^{13}b^3)))/(a^5)^{(1/2)} - (b/(3*(a*b - a^2))) - (b*(a + b/\tan(x)^2)*(2*a - b))/(a*b - a^2)^2)/(a + b/\tan(x)^2)^{(3/2)} + (atan((((a + b/\tan(x)^2)^{(1/2)}(2a^6b^{12} - 20a^7b^{11} + 90a^8b^{10} - 240a^9b^9 + 422a^{10}b^8 - 516a^{11}b^7 + 450a^{12}b^6 - 280a^{13}b^5 + 120a^{14}b^4 - 32a^{15}b^3 + 4a^{16}b^2)))/2 - (((a - b)^5)^{(1/2)}(2a^8b^{13} - 22a^9b^{12} + 110a^{10}b^{11} - 328a^{11}b^{10} + 644a^{12}b^9 - 868a^{13}b^8 + 812a^{14}b^7 - 520a^{15}b^6 + 218a^{16}b^5 - 54a^{17}b^4 + 6a^{18}b^3 - ((a + b/\tan(x)^2)^{(1/2)}((a - b)^5)^{(1/2)}(8a^{10}b^{13} - 96a^{11}b^{12} + 520a^{12}b^{11} - 1680a^{13}b^{10} + 3600a^{14}b^9 - 5376a^{15}b^8 + 5712a^{16}b^7 - 4320a^{17}b^6 + 2280a^{18}b^5 - 800a^{19}b^4 + 168a^{20}b^3 - 16a^{21}b^2)))/(4*(a - b)^5)))/(2*(a - b)^5))*((a - b)^5)^{(1/2)*1i)/(a - b)^5 + (((a + b/\tan(x)^2)^{(1/2)}(2a^6b^{12} - 20a^7b^{11} + 90a^8b^{10} - 240a^9b^9 + 422a^{10}b^8 - 516a^{11}b^7 + 450a^{12}b^6 - 280a^{13}b^5 + 120a^{14}b^4 - 32a^{15}b^3 + 4a^{16}b^2)))/2 + (((a - b)^5)^{(1/2)}(2a^8b^{13} - 22a^9b^{12} + 110a^{10}b^{11} - 328a^{11}b^{10} + 644a^{12}b^9 - 868a^{13}b^8 + 812a^{14}b^7 - 520a^{15}b^6 + 218a^{16}b^5 - 54a^{17}b^4 + 6a^{18}b^3 + ((a + b/\tan(x)^2)^{(1/2)}((a - b)^5)^{(1/2)}(8a^{10}b^{13} - 96a^{11}b^{12} + 520a^{12}b^{11} - 1680a^{13}b^{10} + 3600a^{14}b^9 - 5376a^{15}b^8 + 5712a^{16}b^7 - 4320a^{17}b^6 + 2280a^{18}b^5 - 800a^{19}b^4 + 168a^{20}b^3 - 16a^{21}b^2)))/(4*(a - b)^5)))/(2*(a - b)^5))*((a - b)^5)^{(1/2)*1i)/(a
\end{aligned}$$

$$\begin{aligned}
& - b)^5)/(2*a^6*b^10 - 16*a^7*b^9 + 54*a^8*b^8 - 100*a^9*b^7 + 110*a^10*b^6 \\
& - 72*a^11*b^5 + 26*a^12*b^4 - 4*a^13*b^3 + (((a + b/\tan(x)^2)^{(1/2)}*(2*a^6 \\
& *b^12 - 20*a^7*b^11 + 90*a^8*b^10 - 240*a^9*b^9 + 422*a^10*b^8 - 516*a^11*b \\
& ^7 + 450*a^12*b^6 - 280*a^13*b^5 + 120*a^14*b^4 - 32*a^15*b^3 + 4*a^16*b^2) \\
&)/2 - (((a - b)^5)^{(1/2)}*(2*a^8*b^13 - 22*a^9*b^12 + 110*a^10*b^11 - 328*a^ \\
& 11*b^10 + 644*a^12*b^9 - 868*a^13*b^8 + 812*a^14*b^7 - 520*a^15*b^6 + 218*a \\
& ^16*b^5 - 54*a^17*b^4 + 6*a^18*b^3 - ((a + b/\tan(x)^2)^{(1/2)}*((a - b)^5)^{(1 \\
& /2)}*(8*a^10*b^13 - 96*a^11*b^12 + 520*a^12*b^11 - 1680*a^13*b^10 + 3600*a^1 \\
& 4*b^9 - 5376*a^15*b^8 + 5712*a^16*b^7 - 4320*a^17*b^6 + 2280*a^18*b^5 - 800 \\
& *a^19*b^4 + 168*a^20*b^3 - 16*a^21*b^2))/(4*(a - b)^5)))/(2*(a - b)^5))*((a \\
& - b)^5)^{(1/2)}/(a - b)^5 - (((a + b/\tan(x)^2)^{(1/2)}*(2*a^6*b^12 - 20*a^7*b \\
& ^11 + 90*a^8*b^10 - 240*a^9*b^9 + 422*a^10*b^8 - 516*a^11*b^7 + 450*a^12*b \\
& ^6 - 280*a^13*b^5 + 120*a^14*b^4 - 32*a^15*b^3 + 4*a^16*b^2))/2 + (((a - b) \\
& ^5)^{(1/2)}*(2*a^8*b^13 - 22*a^9*b^12 + 110*a^10*b^11 - 328*a^11*b^10 + 644*a \\
& ^12*b^9 - 868*a^13*b^8 + 812*a^14*b^7 - 520*a^15*b^6 + 218*a^16*b^5 - 54*a^ \\
& 17*b^4 + 6*a^18*b^3 + ((a + b/\tan(x)^2)^{(1/2)}*((a - b)^5)^{(1/2)}*(8*a^10*b^1 \\
& 3 - 96*a^11*b^12 + 520*a^12*b^11 - 1680*a^13*b^10 + 3600*a^14*b^9 - 5376*a^ \\
& 15*b^8 + 5712*a^16*b^7 - 4320*a^17*b^6 + 2280*a^18*b^5 - 800*a^19*b^4 + 168 \\
& *a^20*b^3 - 16*a^21*b^2))/(4*(a - b)^5)))/(2*(a - b)^5))*((a - b)^5)^{(1/2)} \\
& /((a - b)^5))*((a - b)^5)^{(1/2)*1i)/(a - b)^5
\end{aligned}$$

$$3.58 \quad \int \frac{\tan^2(x)}{(a+b \cot^2(x))^{5/2}} dx$$

Optimal result	362
Rubi [A] (verified)	362
Mathematica [C] (warning: unable to verify)	365
Maple [B] (warning: unable to verify)	366
Fricas [B] (verification not implemented)	367
Sympy [F]	367
Maxima [F]	368
Giac [B] (verification not implemented)	368
Mupad [F(-1)]	369

Optimal result

Integrand size = 17, antiderivative size = 141

$$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{5/2}} + \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} + \frac{(a-4b)(3a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{3a^3(a-b)^2}$$

[Out] $\arctan(\cot(x)*(a-b)^{(1/2)}/(a+b*\cot(x)^2)^{(1/2)})/(a-b)^{(5/2)}+1/3*b*\tan(x)/a/(a-b)/(a+b*\cot(x)^2)^{(3/2)}+1/3*(7*a-4*b)*b*\tan(x)/a^2/(a-b)^2/(a+b*\cot(x)^2)^{(1/2)}+1/3*(a-4*b)*(3*a-2*b)*(a+b*\cot(x)^2)^{(1/2)}*\tan(x)/a^3/(a-b)^2$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.412$, Rules used = {3751, 483, 593, 597, 12, 385, 209}

$$\int \frac{\tan^2(x)}{(a+b \cot^2(x))^{5/2}} dx = \frac{(a-4b)(3a-2b) \tan(x) \sqrt{a+b \cot^2(x)}}{3a^3(a-b)^2} + \frac{b(7a-4b) \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} + \frac{\arctan\left(\frac{\sqrt{a-b} \cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^{5/2}} + \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}}$$

[In] $\text{Int}[\text{Tan}[x]^2/(a+b*\text{Cot}[x]^2)^{(5/2)},x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[a-b]*\text{Cot}[x])/(\text{Sqrt}[a+b*\text{Cot}[x]^2])]/(a-b)^{(5/2)}+(b*\text{Tan}[x])/(3*a*(a-b)*(a+b*\text{Cot}[x]^2)^{(3/2)})+((7*a-4*b)*b*\text{Tan}[x])/(3*a^2*(a-$

$b^2 \sqrt{a + b \cot[x]^2} + ((a - 4b)(3a - 2b) \sqrt{a + b \cot[x]^2} \tan[x]) / (3a^3(a - b)^2)$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match}[\text{Q}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]]$

Rule 209

$\text{Int}[(a_*) + (b_*)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2] \text{Rt}[b, 2])) * \text{ArcTan}[\text{Rt}[b, 2] * (x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 385

$\text{Int}[(a_*) + (b_*)(x_)^{(n_)}])^{(p_)} / ((c_*) + (d_*)(x_)^{(n_)}), x_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(c - (b*c - a*d)*x^n), x], x, x/(a + b*x^n)^{1/n}] /; \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{EqQ}[n*p + 1, 0] \ \&\& \ \text{IntegerQ}[n]$

Rule 483

$\text{Int}[(e_*)(x_)^{(m_)} * ((a_*) + (b_*)(x_)^{(n_)}])^{(p_)} * ((c_*) + (d_*)(x_)^{(n_)}])^{(q_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*(e*x)^{(m+1)} * (a + b*x^n)^{(p+1)} * ((c + d*x^n)^{(q+1)} / (a*e*n*(b*c - a*d)*(p+1))), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(e*x)^m * (a + b*x^n)^{(p+1)} * (c + d*x^n)^q * \text{Simp}[c*b*(m+1) + n*(b*c - a*d)*(p+1) + d*b*(m + n*(p+q+2) + 1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, m, q\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, d, e, m, n, p, q, x]$

Rule 593

$\text{Int}[(g_*)(x_)^{(m_)} * ((a_*) + (b_*)(x_)^{(n_)}])^{(p_)} * ((c_*) + (d_*)(x_)^{(n_)}])^{(q_)} * ((e_*) + (f_*)(x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(-b*e - a*f) * (g*x)^{(m+1)} * (a + b*x^n)^{(p+1)} * ((c + d*x^n)^{(q+1)} / (a*g*n*(b*c - a*d)*(p+1))), x] + \text{Dist}[1/(a*n*(b*c - a*d)*(p+1)), \text{Int}[(g*x)^m * (a + b*x^n)^{(p+1)} * (c + d*x^n)^q * \text{Simp}[c*(b*e - a*f)*(m+1) + e*n*(b*c - a*d)*(p+1) + d*(b*e - a*f)*(m + n*(p+q+2) + 1)*x^n, x], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g, m, q\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1]$

Rule 597

$\text{Int}[(g_*)(x_)^{(m_)} * ((a_*) + (b_*)(x_)^{(n_)}])^{(p_)} * ((c_*) + (d_*)(x_)^{(n_)}])^{(q_)} * ((e_*) + (f_*)(x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[e * (g*x)^{(m+1)} * (a + b*x^n)^{(p+1)} * ((c + d*x^n)^{(q+1)} / (a*c*g*(m+1))), x] + \text{Dist}[1/(a*c*g*(m+1)), \text{Int}[(g*x)^{(m+n)} * (a + b*x^n)^p * (c + d*x^n)^q * \text{Simp}[a*f*c*(m+1) - e*(b*c + a*d)*(m+n+1) - e*n*(b*c*p + a*d*q) - b*e*d*(m+n*(p+q+2))$

+ 1)*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, p, q}, x] && IGtQ[n, 0] && LtQ[m, -1]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_.)])^(m_.)*((a_.) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.))^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{1}{x^2(1+x^2)(a+bx^2)^{5/2}} dx, x, \cot(x)\right) \\
 &= \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} - \frac{\text{Subst}\left(\int \frac{3a-4b-4bx^2}{x^2(1+x^2)(a+bx^2)^{3/2}} dx, x, \cot(x)\right)}{3a(a-b)} \\
 &= \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{(a-4b)(3a-2b)-2(7a-4b)bx^2}{x^2(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{3a^2(a-b)^2} \\
 &= \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} \\
 &\quad + \frac{(a-4b)(3a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{3a^3(a-b)^2} + \frac{\text{Subst}\left(\int \frac{3a^3}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{3a^3(a-b)^2} \\
 &= \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} \\
 &\quad + \frac{(a-4b)(3a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{3a^3(a-b)^2} + \frac{\text{Subst}\left(\int \frac{1}{(1+x^2)\sqrt{a+bx^2}} dx, x, \cot(x)\right)}{(a-b)^2} \\
 &= \frac{b \tan(x)}{3a(a-b)(a+b \cot^2(x))^{3/2}} + \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2 \sqrt{a+b \cot^2(x)}} \\
 &\quad + \frac{(a-4b)(3a-2b)\sqrt{a+b \cot^2(x)} \tan(x)}{3a^3(a-b)^2} + \frac{\text{Subst}\left(\int \frac{1}{1-(-a+b)x^2} dx, x, \frac{\cot(x)}{\sqrt{a+b \cot^2(x)}}\right)}{(a-b)^2}
 \end{aligned}$$

$$= \frac{\arctan\left(\frac{\sqrt{a-b}\cot(x)}{\sqrt{a+b}\cot^2(x)}\right)}{(a-b)^{5/2}} + \frac{b \tan(x)}{3a(a-b)(a+b\cot^2(x))^{3/2}}$$

$$+ \frac{(7a-4b)b \tan(x)}{3a^2(a-b)^2\sqrt{a+b\cot^2(x)}} + \frac{(a-4b)(3a-2b)\sqrt{a+b\cot^2(x)} \tan(x)}{3a^3(a-b)^2}$$

Mathematica [C] (warning: unable to verify)

Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

Time = 8.16 (sec) , antiderivative size = 1450, normalized size of antiderivative = 10.28

$$\int \frac{\tan^2(x)}{(a+b\cot^2(x))^{5/2}} dx = \frac{\sin^2(x) \left(-\frac{16b^3(\cot(x)+\cot^3(x))^2}{a(a-b)^2} + \frac{40b \csc^2(x)}{a-b} + \frac{160b^2 \cot^2(x) \csc^2(x)}{3a(a-b)} + \frac{64b^3 \cot^4(x) \csc^2(x)}{3a^2(a-b)} - 4 \right)}{\dots}$$

[In] Integrate[Tan[x]^2/(a + b*Cot[x]^2)^(5/2), x]

[Out] (Sin[x]^2*((-16*b^3*(Cot[x] + Cot[x]^3)^2)/(a*(a - b)^2) + (40*b*Csc[x]^2)/(a - b) + (160*b^2*Cot[x]^2*Csc[x]^2)/(3*a*(a - b)) + (64*b^3*Cot[x]^4*Csc[x]^2)/(3*a^2*(a - b)) - (40*b^2*Csc[x]^4)/(a - b)^2 + (92*(a - b)*Cos[x]^2*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a])/(105*a) + (124*(a - b)*b*Cos[x]^2*Cot[x]^2*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a])/(35*a^2) + (152*(a - b)*b^2*Cos[x]^2*Cot[x]^4*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a])/(35*a^3) + (176*(a - b)*b^3*Cos[x]^2*Cot[x]^6*Hypergeometric2F1[2, 2, 9/2, ((a - b)*Cos[x]^2)/a])/(105*a^4) + (24*(a - b)*Cos[x]^2*HypergeometricPFQ[{2, 2, 2}, {1, 9/2}, ((a - b)*Cos[x]^2)/a])/(35*a) + (16*(a - b)*b*Cos[x]^2*Cot[x]^2*HypergeometricPFQ[{2, 2, 2}, {1, 9/2}, ((a - b)*Cos[x]^2)/a])/(7*a^2) + (88*(a - b)*b^2*Cos[x]^2*Cot[x]^4*HypergeometricPFQ[{2, 2, 2}, {1, 9/2}, ((a - b)*Cos[x]^2)/a])/(35*a^3) + (32*(a - b)*b^3*Cos[x]^2*Cot[x]^6*HypergeometricPFQ[{2, 2, 2}, {1, 9/2}, ((a - b)*Cos[x]^2)/a])/(35*a^4) + (16*(a - b)*Cos[x]^2*HypergeometricPFQ[{2, 2, 2, 2}, {1, 1, 9/2}, ((a - b)*Cos[x]^2)/a])/(105*a) + (16*(a - b)*b*Cos[x]^2*Cot[x]^2*HypergeometricPFQ[{2, 2, 2, 2}, {1, 1, 9/2}, ((a - b)*Cos[x]^2)/a])/(35*a^2) + (16*(a - b)*b^2*Cos[x]^2*Cot[x]^4*HypergeometricPFQ[{2, 2, 2, 2}, {1, 1, 9/2}, ((a - b)*Cos[x]^2)/a])/(35*a^3) + (16*(a - b)*b^3*Cos[x]^2*Cot[x]^6*HypergeometricPFQ[{2, 2, 2, 2}, {1, 1, 9/2}, ((a - b)*Cos[x]^2)/a])/(105*a^4) + (20*a*Sec[x]^2)/(3*(a - b)) - (30*a*b*Csc[x]^2*Sec[x]^2)/(a - b)^2 - (5*a^2*Sec[x]^4)/(a - b)^2 + (5*ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]])/(((a - b)*Cos[x]^2)/a)^(5/2)*Sqrt[((a + b*Cot[x]^2)*Sin[x]^2)/a]) + (30*b*ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*Cot[x]^2)/(a*((a - b)*Cos[x]^2)/a)^(5/2)*Sqrt[((a + b*Cot[x]^2)*Sin[x]^2)/a]) + (40*b^2*ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*Cot[x]^4)/(a^2*((a - b)*Cos[x]^2)/a)^(5/2)*Sqrt[((a + b*Cot[x]^2)*Sin[x]^2)/a]) + (16*b^3*ArcSin[Sqrt[((a - b)*Cos[x]^2)/a]]*Cot[x]^6)/(a^3*((a - b)*C

$$2*\csc(x)^2-2*b*(1-\cos(x))^2*\csc(x)^2+b)/((1-\cos(x))^2*\csc(x)^2-1)/(1-\cos(x))^5*\sin(x)^5/(1/(1-\cos(x))^2*(b*(1-\cos(x))^4*\csc(x)^2+4*a*(1-\cos(x))^2-2*b*(1-\cos(x))^2+b*\sin(x)^2))^(5/2)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 297 vs. 2(123) = 246.

Time = 0.34 (sec) , antiderivative size = 647, normalized size of antiderivative = 4.59

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \left[\frac{3(a^5 \tan(x)^4 + 2a^4 b \tan(x)^2 + a^3 b^2) \sqrt{-a + b} \log\left(-\frac{a^2 \tan(x)^4 - 2(3a^2 - 4ab) \tan(x)^2 + a^2}{(a \tan(x)^2 + b)/\tan(x)^2}\right)}{(a \tan(x)^2 + b)/\tan(x)^2} \right]$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="fricas")

[Out] [-1/12*(3*(a^5*tan(x)^4 + 2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(-a + b)*log(-(a^2*tan(x)^4 - 2*(3*a^2 - 4*a*b)*tan(x)^2 + a^2 - 8*a*b + 8*b^2 + 4*(a*tan(x)^3 - (a - 2*b)*tan(x))*sqrt(-a + b)*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(tan(x)^4 + 2*tan(x)^2 + 1)) - 4*(3*(a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*tan(x)^5 + 3*(2*a^4*b - 9*a^3*b^2 + 11*a^2*b^3 - 4*a*b^4)*tan(x)^3 + (3*a^3*b^2 - 17*a^2*b^3 + 22*a*b^4 - 8*b^5)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4)*tan(x)^2), 1/6*(3*(a^5*tan(x)^4 + 2*a^4*b*tan(x)^2 + a^3*b^2)*sqrt(a - b)*arctan(2*sqrt(a - b)*sqrt((a*tan(x)^2 + b)/tan(x)^2)*tan(x)/(a*tan(x)^2 - a + 2*b)) + 2*(3*(a^5 - 3*a^4*b + 3*a^3*b^2 - a^2*b^3)*tan(x)^5 + 3*(2*a^4*b - 9*a^3*b^2 + 11*a^2*b^3 - 4*a*b^4)*tan(x)^3 + (3*a^3*b^2 - 17*a^2*b^3 + 22*a*b^4 - 8*b^5)*tan(x))*sqrt((a*tan(x)^2 + b)/tan(x)^2))/(a^6*b^2 - 3*a^5*b^3 + 3*a^4*b^4 - a^3*b^5 + (a^8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3)*tan(x)^4 + 2*(a^7*b - 3*a^6*b^2 + 3*a^5*b^3 - a^4*b^4)*tan(x)^2)]

Sympy [F]

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx$$

[In] integrate(tan(x)**2/(a+b*cot(x)**2)**(5/2),x)

[Out] Integral(tan(x)**2/(a + b*cot(x)**2)**(5/2), x)

Maxima [F]

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\tan(x)^2}{(b \cot(x)^2 + a)^{5/2}} dx$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="maxima")

[Out] integrate(tan(x)^2/(b*cot(x)^2 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 537 vs. 2(123) = 246.

Time = 0.36 (sec) , antiderivative size = 537, normalized size of antiderivative = 3.81

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \frac{\left(3 a^4 \sqrt{b} \log\left(-a - 2 \sqrt{-a + b} \sqrt{b} + 2 b\right) + 3 a^3 \sqrt{-a + b} b \log\left(-a - 2 \sqrt{-a + b} \sqrt{b}\right)\right)}{6 \left(a^6 \sqrt{-a + b}\right)} + \frac{2 \left(\frac{\left(9 a^5 b^2 - 23 a^4 b^3 + 19 a^3 b^4 - 5 a^2 b^5\right) \cos(x)^2 - 3 \left(3 a^5 b^2 - 5 a^4 b^3 + 2 a^3 b^4\right)}{a^8 - 3 a^7 b + 3 a^6 b^2 - a^5 b^3}\right) \cos(x)}{\left(a \cos(x)^2 - b \cos(x)^2 - a\right) \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}} + \frac{3 \log\left(\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}\right)^2\right)}{\left(a^2 - 2 a b + b^2\right) \sqrt{-a + b}} + \frac{\left(\sqrt{-a + b} \cos(x) - \sqrt{-a \cos(x)^2 + b \cos(x)^2 + a}\right)}{6 \operatorname{sgn}(\sin(x))}$$

[In] integrate(tan(x)^2/(a+b*cot(x)^2)^(5/2),x, algorithm="giac")

[Out] 1/6*(3*a^4*sqrt(b)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 3*a^3*sqrt(-a + b)*b*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) - 3*a^3*b^(3/2)*log(-a - 2*sqrt(-a + b)*sqrt(b) + 2*b) + 6*a^4*sqrt(b) - 18*a^3*b^(3/2) + 16*a^2*sqrt(-a + b)*b^2 + 2*a^2*b^(5/2) - 26*a*sqrt(-a + b)*b^3 + 20*a*b^(7/2) + 10*sqrt(-a + b)*b^4 - 10*b^(9/2))*sgn(sin(x))/(a^6*sqrt(-a + b)*sqrt(b) - a^6*b - 3*a^5*sqrt(-a + b)*b^(3/2) + 3*a^5*b^2 + 3*a^4*sqrt(-a + b)*b^(5/2) - 3*a^4*b^3 - a^3*sqrt(-a + b)*b^(7/2) + a^3*b^4) - 1/6*(2*((9*a^5*b^2 - 23*a^4*b^3 + 19*a^3*b^4 - 5*a^2*b^5)*cos(x)^2/(a^8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3) - 3*(3*a^5*b^2 - 5*a^4*b^3 + 2*a^3*b^4)/(a^8 - 3*a^7*b + 3*a^6*b^2 - a^5*b^3))*cos(x)/((a*cos(x)^2 - b*cos(x)^2 - a)*sqrt(-a*cos(x)^2 + b*cos(x)^2 + a)) + 3*log((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2)/((a^2 - 2*a*b + b^2)*sqrt(-a + b)) + 12*sqrt(-a + b)/(((sqrt(-a + b)*cos(x) - sqrt(-a*cos(x)^2 + b*cos(x)^2 + a))^2 - a)*a^2))/sgn(sin(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^2(x)}{(a + b \cot^2(x))^{5/2}} dx = \int \frac{\tan(x)^2}{(b \cot(x)^2 + a)^{5/2}} dx$$

```
[In] int(tan(x)^2/(a + b*cot(x)^2)^(5/2), x)
```

```
[Out] int(tan(x)^2/(a + b*cot(x)^2)^(5/2), x)
```

3.59 $\int \frac{1}{1+\cot^3(x)} dx$

Optimal result	370
Rubi [A] (verified)	370
Mathematica [C] (verified)	372
Maple [A] (verified)	372
Fricas [A] (verification not implemented)	373
Sympy [A] (verification not implemented)	373
Maxima [A] (verification not implemented)	373
Giac [A] (verification not implemented)	374
Mupad [B] (verification not implemented)	374

Optimal result

Integrand size = 8, antiderivative size = 37

$$\int \frac{1}{1+\cot^3(x)} dx = \frac{x}{2} - \frac{1}{6} \log(1+\cot(x)) + \frac{1}{3} \log(1-\cot(x)+\cot^2(x)) + \frac{1}{2} \log(\sin(x))$$

[Out] 1/2*x-1/6*ln(1+cot(x))+1/3*ln(1-cot(x)+cot(x)^2)+1/2*ln(sin(x))

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3742, 2099, 649, 209, 266, 642}

$$\int \frac{1}{1+\cot^3(x)} dx = \frac{x}{2} + \frac{1}{2} \log(\sin(x)) + \frac{1}{3} \log(\cot^2(x) - \cot(x) + 1) - \frac{1}{6} \log(\cot(x) + 1)$$

[In] Int[(1 + Cot[x]^3)^(-1),x]

[Out] x/2 - Log[1 + Cot[x]]/6 + Log[1 - Cot[x] + Cot[x]^2]/3 + Log[Sin[x]]/2

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 266

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 649

```
Int[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] := Dist[d, Int[1/(
a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e
}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 2099

```
Int[(P_)^(p_)*(Q_)^(q_), x_Symbol] := With[{PP = Factor[P]}, Int[ExpandInt
egrand[PP^p*Q^q, x], x] /; !SumQ[NonfreeFactors[PP, x]] /; FreeQ[q, x] &&
PolyQ[P, x] && PolyQ[Q, x] && IntegerQ[p] && NeQ[P, x]
```

Rule 3742

```
Int[((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] :=
With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(a + b*(
ff*x)^n]^p/(c^2 + ff^2*x^2), x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a,
b, c, e, f, n, p}, x] && (IntegerQ[n, p] || IGtQ[p, 0] || EqQ[n^2, 4] || E
qQ[n^2, 16])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{1}{(1+x^2)(1+x^3)} dx, x, \cot(x)\right) \\
&= -\text{Subst}\left(\int \left(\frac{1}{6(1+x)} + \frac{1+x}{2(1+x^2)} + \frac{1-2x}{3(1-x+x^2)}\right) dx, x, \cot(x)\right) \\
&= -\frac{1}{6} \log(1 + \cot(x)) - \frac{1}{3} \text{Subst}\left(\int \frac{1-2x}{1-x+x^2} dx, x, \cot(x)\right) \\
&\quad - \frac{1}{2} \text{Subst}\left(\int \frac{1+x}{1+x^2} dx, x, \cot(x)\right) \\
&= -\frac{1}{6} \log(1 + \cot(x)) + \frac{1}{3} \log(1 - \cot(x) + \cot^2(x)) \\
&\quad - \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(x)\right) - \frac{1}{2} \text{Subst}\left(\int \frac{x}{1+x^2} dx, x, \cot(x)\right) \\
&= \frac{x}{2} - \frac{1}{6} \log(1 + \cot(x)) + \frac{1}{3} \log(1 - \cot(x) + \cot^2(x)) + \frac{1}{2} \log(\sin(x))
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.05 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.54

$$\int \frac{1}{1 + \cot^3(x)} dx = \left(-\frac{1}{4} - \frac{i}{4}\right) \log(i - \tan(x)) - \left(\frac{1}{4} - \frac{i}{4}\right) \log(i + \tan(x)) - \frac{1}{6} \log(1 + \tan(x)) + \frac{1}{3} \log(1 - \tan(x) + \tan^2(x))$$

[In] Integrate[(1 + Cot[x]^3)^(-1),x]

[Out] (-1/4 - I/4)*Log[I - Tan[x]] - (1/4 - I/4)*Log[I + Tan[x]] - Log[1 + Tan[x]]/6 + Log[1 - Tan[x] + Tan[x]^2]/3

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 0.84

method	result	size
parallelrisc	$\frac{x}{2} + \ln\left(\frac{1}{(\tan(x)+1)^{\frac{1}{6}}}\right) + \ln\left(\frac{1}{(\sec(x)^2)^{\frac{1}{4}}}\right) + \ln\left((- \tan(x) + \sec(x)^2)^{\frac{1}{3}}\right)$	31
norman	$\frac{x}{2} - \frac{\ln(\tan(x)+1)}{6} - \frac{\ln(\tan(x)^2+1)}{4} + \frac{\ln(\tan(x)^2-\tan(x)+1)}{3}$	34
risc	$\frac{x}{2} - \frac{ix}{2} - \frac{\ln(e^{2ix}+i)}{6} + \frac{\ln(e^{4ix}-4ie^{2ix}-1)}{3}$	38
derivativdivides	$\frac{\ln(1-\cot(x)+\cot(x)^2)}{3} - \frac{\ln(\cot(x)^2+1)}{4} - \frac{\pi}{4} + \frac{\operatorname{arccot}(\cot(x))}{2} - \frac{\ln(1+\cot(x))}{6}$	39
default	$\frac{\ln(1-\cot(x)+\cot(x)^2)}{3} - \frac{\ln(\cot(x)^2+1)}{4} - \frac{\pi}{4} + \frac{\operatorname{arccot}(\cot(x))}{2} - \frac{\ln(1+\cot(x))}{6}$	39

[In] int(1/(1+cot(x)^3),x,method=_RETURNVERBOSE)

[Out] 1/2*x+ln(1/(tan(x)+1)^(1/6))+ln(1/(sec(x)^2)^(1/4))+ln((-tan(x)+sec(x)^2)^(1/3))

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.65

$$\int \frac{1}{1 + \cot^3(x)} dx = \frac{1}{2}x - \frac{1}{12} \log(\sin(2x) + 1) + \frac{1}{3} \log\left(-\frac{1}{2} \sin(2x) + 1\right)$$

[In] integrate(1/(1+cot(x)^3),x, algorithm="fricas")

[Out] 1/2*x - 1/12*log(sin(2*x) + 1) + 1/3*log(-1/2*sin(2*x) + 1)

Sympy [A] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int \frac{1}{1 + \cot^3(x)} dx = \frac{x}{2} - \frac{\log(\tan(x) + 1)}{6} - \frac{\log(\tan^2(x) + 1)}{4} + \frac{\log(\tan^2(x) - \tan(x) + 1)}{3}$$

[In] integrate(1/(1+cot(x)**3),x)

[Out] x/2 - log(tan(x) + 1)/6 - log(tan(x)**2 + 1)/4 + log(tan(x)**2 - tan(x) + 1)/3

Maxima [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.89

$$\int \frac{1}{1 + \cot^3(x)} dx = \frac{1}{2}x + \frac{1}{3} \log(\tan(x)^2 - \tan(x) + 1) - \frac{1}{4} \log(\tan(x)^2 + 1) - \frac{1}{6} \log(\tan(x) + 1)$$

[In] integrate(1/(1+cot(x)^3),x, algorithm="maxima")

[Out] 1/2*x + 1/3*log(tan(x)^2 - tan(x) + 1) - 1/4*log(tan(x)^2 + 1) - 1/6*log(tan(x) + 1)

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.92

$$\int \frac{1}{1 + \cot^3(x)} dx = \frac{1}{2} x + \frac{1}{3} \log(\tan(x)^2 - \tan(x) + 1) - \frac{1}{4} \log(\tan(x)^2 + 1) - \frac{1}{6} \log(|\tan(x) + 1|)$$

[In] integrate(1/(1+cot(x)^3),x, algorithm="giac")

[Out] 1/2*x + 1/3*log(tan(x)^2 - tan(x) + 1) - 1/4*log(tan(x)^2 + 1) - 1/6*log(abs(tan(x) + 1))

Mupad [B] (verification not implemented)

Time = 13.79 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \cot^3(x)} dx = x \left(\frac{1}{2} - \frac{1}{2}i \right) - \frac{\ln(12e^{x2i} + 12i)}{6} + \frac{\ln(e^{x4i} - 1 - e^{x2i}4i)}{3}$$

[In] int(1/(cot(x)^3 + 1),x)

[Out] x*(1/2 - 1i/2) - log(12*exp(x*2i) + 12i)/6 + log(exp(x*4i) - exp(x*2i)*4i - 1)/3

3.60 $\int \cot(x) \sqrt{a + b \cot^4(x)} dx$

Optimal result	375
Rubi [A] (verified)	375
Mathematica [A] (verified)	377
Maple [A] (verified)	378
Fricas [B] (verification not implemented)	378
Sympy [F]	379
Maxima [F]	379
Giac [B] (verification not implemented)	380
Mupad [F(-1)]	380

Optimal result

Integrand size = 15, antiderivative size = 90

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \frac{1}{2} \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a + b \cot^4(x)}} \right) + \frac{1}{2} \sqrt{a + b} \operatorname{arctanh} \left(\frac{a - b \cot^2(x)}{\sqrt{a + b} \sqrt{a + b \cot^4(x)}} \right) - \frac{1}{2} \sqrt{a + b \cot^4(x)}$$

[Out] 1/2*arctanh(cot(x)^2*b^(1/2)/(a+b*cot(x)^4)^(1/2))*b^(1/2)+1/2*arctanh((a-b*cot(x)^2)/(a+b)^(1/2)/(a+b*cot(x)^4)^(1/2))*(a+b)^(1/2)-1/2*(a+b*cot(x)^4)^(1/2)

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3751, 1262, 749, 858, 223, 212, 739}

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \frac{1}{2} \sqrt{b} \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a + b \cot^4(x)}} \right) + \frac{1}{2} \sqrt{a + b} \operatorname{arctanh} \left(\frac{a - b \cot^2(x)}{\sqrt{a + b} \sqrt{a + b \cot^4(x)}} \right) - \frac{1}{2} \sqrt{a + b \cot^4(x)}$$

[In] Int[Cot[x]*Sqrt[a + b*Cot[x]^4],x]

[Out] $(\sqrt{b} \operatorname{ArcTanh}[(\sqrt{b} \cot[x]^2)/\sqrt{a + b \cot[x]^4}])/2 + (\sqrt{a + b} \operatorname{ArcTanh}[(a - b \cot[x]^2)/(\sqrt{a + b} \sqrt{a + b \cot[x]^4})])/2 - \sqrt{a + b \cot[x]^4}/2$

Rule 212

$\operatorname{Int}[(a_ + (b_)(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2](x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\sqrt{(a_ + (b_)(x_)^2)}, x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b x^2), x], x, x/\sqrt{a + b x^2}] /; \operatorname{FreeQ}[\{a, b\}, x] \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 739

$\operatorname{Int}[1/(((d_ + (e_)(x_)) \sqrt{(a_ + (c_)(x_)^2)}), x_Symbol] \rightarrow -\operatorname{Subst}[\operatorname{Int}[1/(c d^2 + a e^2 - x^2), x], x, (a e - c d x)/\sqrt{a + c x^2}] /; \operatorname{FreeQ}[\{a, c, d, e\}, x]$

Rule 749

$\operatorname{Int}(((d_ + (e_)(x_))^{(m_)}((a_ + (c_)(x_)^2)^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[(d + e x)^{(m + 1)}((a + c x^2)^p/(e(m + 2p + 1))), x] + \operatorname{Dist}[2(p/(e(m + 2p + 1))), \operatorname{Int}[(d + e x)^m \operatorname{Simp}[a e - c d x, x](a + c x^2)^{(p - 1)}, x], x] /; \operatorname{FreeQ}[\{a, c, d, e, m\}, x] \ \&\& \operatorname{NeQ}[c d^2 + a e^2, 0] \ \&\& \operatorname{GtQ}[p, 0] \ \&\& \operatorname{NeQ}[m + 2p + 1, 0] \ \&\& (\ !\operatorname{RationalQ}[m] \ || \operatorname{LtQ}[m, 1]) \ \&\& \ !\operatorname{ILtQ}[m + 2p, 0] \ \&\& \operatorname{IntQuadraticQ}[a, 0, c, d, e, m, p, x]$

Rule 858

$\operatorname{Int}(((d_ + (e_)(x_))^{(m_)}((f_ + (g_)(x_))((a_ + (c_)(x_)^2)^{(p_)}), x_Symbol] \rightarrow \operatorname{Dist}[g/e, \operatorname{Int}[(d + e x)^{(m + 1)}(a + c x^2)^p, x], x] + \operatorname{Dist}[(e f - d g)/e, \operatorname{Int}[(d + e x)^m (a + c x^2)^p, x], x] /; \operatorname{FreeQ}[\{a, c, d, e, f, g, m, p\}, x] \ \&\& \operatorname{NeQ}[c d^2 + a e^2, 0] \ \&\& \ !\operatorname{IGtQ}[m, 0]$

Rule 1262

$\operatorname{Int}(x_)((d_ + (e_)(x_)^2)^{(q_)}((a_ + (c_)(x_)^4)^{(p_)}), x_Symbol] \rightarrow \operatorname{Dist}[1/2, \operatorname{Subst}[\operatorname{Int}[(d + e x)^q (a + c x^2)^p, x], x, x^2], x] /; \operatorname{FreeQ}[\{a, c, d, e, p, q\}, x]$

Rule 3751

$\operatorname{Int}(((d_)\tan[(e_ + (f_)(x_))]^{(m_)}((a_ + (b_)((c_)\tan[(e_ + (f_)(x_)]^{(n_)}))^{(p_)}), x_Symbol] \rightarrow \operatorname{With}[\{ff = \operatorname{FreeFactors}[\operatorname{Tan}[e + f x],$

$x\}}, \text{Dist}[c*(ff/f), \text{Subst}[\text{Int}[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)), x], x, c*(\text{Tan}[e + f*x]/ff)], x]] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& (\text{IGtQ}[p, 0] \|\ \text{EqQ}[n, 2] \|\ \text{EqQ}[n, 4] \|\ (\text{IntegerQ}[p] \&\& \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x\sqrt{a+bx^4}}{1+x^2} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{\sqrt{a+bx^2}}{1+x} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{1}{2}\sqrt{a+b\cot^4(x)} - \frac{1}{2}\text{Subst}\left(\int \frac{a-bx}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right) \\
&= -\frac{1}{2}\sqrt{a+b\cot^4(x)} + \frac{1}{2}b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot^2(x)\right) \\
&\quad - \frac{1}{2}(a+b)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right) \\
&= -\frac{1}{2}\sqrt{a+b\cot^4(x)} - \frac{1}{2}(-a-b)\text{Subst}\left(\int \frac{1}{a+b-x^2} dx, x, \frac{a-b\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right) \\
&\quad + \frac{1}{2}b\text{Subst}\left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right) \\
&= \frac{1}{2}\sqrt{b}\text{arctanh}\left(\frac{\sqrt{b}\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right) \\
&\quad + \frac{1}{2}\sqrt{a+b}\text{arctanh}\left(\frac{a-b\cot^2(x)}{\sqrt{a+b}\sqrt{a+b\cot^4(x)}}\right) - \frac{1}{2}\sqrt{a+b\cot^4(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.19 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.96

$$\begin{aligned}
\int \cot(x)\sqrt{a+b\cot^4(x)} dx &= \frac{1}{2}\left(\sqrt{b}\text{arctanh}\left(\frac{\sqrt{b}\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right)\right. \\
&\quad \left. + \sqrt{a+b}\text{arctanh}\left(\frac{a-b\cot^2(x)}{\sqrt{a+b}\sqrt{a+b\cot^4(x)}}\right)\right. \\
&\quad \left. - \sqrt{a+b\cot^4(x)}\right)
\end{aligned}$$

[In] Integrate[Cot[x]*Sqrt[a + b*Cot[x]^4],x]

[Out] (Sqrt[b]*ArcTanh[(Sqrt[b]*Cot[x]^2)/Sqrt[a + b*Cot[x]^4]] + Sqrt[a + b]*ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])) - Sqrt[a + b*Cot[x]^4])/2

Maple [A] (verified)

Time = 0.24 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.54

method	result
derivativedivides	$-\frac{\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(\cot(x)^2+1)-b}{\sqrt{b}} + \sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}\right)}{2} +$
default	$-\frac{\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{2} + \frac{\sqrt{b} \ln\left(\frac{b(\cot(x)^2+1)-b}{\sqrt{b}} + \sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}\right)}{2} +$

[In] int(cot(x)*(a+b*cot(x)^4)^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/2*(b*(cot(x)^2+1)^2-2*b*(cot(x)^2+1)+a+b)^(1/2)+1/2*b^(1/2)*ln((b*(cot(x)^2+1)-b)/b^(1/2)+(b*(cot(x)^2+1)^2-2*b*(cot(x)^2+1)+a+b)^(1/2))+1/2*(a+b)^(1/2)*ln((2*a+2*b-2*b*(cot(x)^2+1)+2*(a+b)^(1/2)*(b*(cot(x)^2+1)^2-2*b*(cot(x)^2+1)+a+b)^(1/2))/(cot(x)^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(72) = 144.

Time = 0.45 (sec) , antiderivative size = 1063, normalized size of antiderivative = 11.81

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \text{Too large to display}$$

[In] integrate(cot(x)*(a+b*cot(x)^4)^(1/2),x, algorithm="fricas")

[Out] [1/4*sqrt(a + b)*log(1/2*(a^2 + 2*a*b + b^2)*cos(2*x)^2 + 1/2*a^2 + 1/2*b^2 + 1/2*((a + b)*cos(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt(a + b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - (a^2 - b^2)*cos(2*x)) + 1/4*sqrt(b)*log(-((a + 2*b)*cos(2*x)^2 - 2*(cos(2*x)^2 - 1)*sqrt(b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - 2*(a - 2*b)*cos(2*x) + a + 2*b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - 1/2*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)), 1/2*sqrt(-b)*arctan(sqrt(-b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)))*(cos(2*x) - 1)/(b*cos(2*x) + b)) + 1/4*sqrt(a + b)*log(1/2*(a^2 + 2*a*b + b^2)*cos(2*x)^2 + 1/2*a^2 + 1/2*b^2 + 1/2*((a + b)*cos(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt(a + b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - (a^2 - b^2)*cos(2*x))

```

x) + a - b)*sqrt(a + b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a +
b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - (a^2 - b^2)*cos(2*x)) - 1/2*sqrt(((a +
b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1))
, -1/2*sqrt(-a - b)*arctan(((a + b)*cos(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt
(-a - b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2
- 2*cos(2*x) + 1)))/((a^2 + 2*a*b + b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2
*(a^2 - b^2)*cos(2*x))) + 1/4*sqrt(b)*log(-((a + 2*b)*cos(2*x)^2 - 2*(cos(2
*x)^2 - 1)*sqrt(b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(
cos(2*x)^2 - 2*cos(2*x) + 1)) - 2*(a - 2*b)*cos(2*x) + a + 2*b)/(cos(2*x)^2
- 2*cos(2*x) + 1)) - 1/2*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a
+ b)/(cos(2*x)^2 - 2*cos(2*x) + 1)), -1/2*sqrt(-a - b)*arctan(((a + b)*cos
(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt(-a - b)*sqrt(((a + b)*cos(2*x)^2 - 2*(
a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)))/((a^2 + 2*a*b + b^2
)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*x))) + 1/2*sqrt(-b)*
arctan(sqrt(-b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos
(2*x)^2 - 2*cos(2*x) + 1))*(cos(2*x) - 1)/(b*cos(2*x) + b)) - 1/2*sqrt(((a
+ b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)
)]

```

Sympy [F]

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \int \sqrt{a + b \cot^4(x)} \cot(x) dx$$

```
[In] integrate(cot(x)*(a+b*cot(x)**4)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*cot(x)**4)*cot(x), x)
```

Maxima [F]

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \int \sqrt{b \cot^4(x) + a} \cot(x) dx$$

```
[In] integrate(cot(x)*(a+b*cot(x)^4)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*cot(x)^4 + a)*cot(x), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(72) = 144.

Time = 0.30 (sec) , antiderivative size = 204, normalized size of antiderivative = 2.27

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = -\frac{b \arctan\left(-\frac{\sqrt{a+b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b}}{\sqrt{-b}}\right)}{\sqrt{-b}} - \frac{\frac{1}{2} \sqrt{a+b} \log\left(\left|-\left(\sqrt{a+b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b}\right)(a+b) + \sqrt{a+bb}\right|\right)}{\left(\sqrt{a+b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b}\right)b - \sqrt{a+bb}} - \frac{\left(\sqrt{a+b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b}\right)^2 - b}{\left(\sqrt{a+b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b}\right)^2 - b}$$

[In] integrate(cot(x)*(a+b*cot(x)^4)^(1/2),x, algorithm="giac")

[Out] -b*arctan(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))/sqrt(-b))/sqrt(-b) - 1/2*sqrt(a + b)*log(abs(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))*(a + b) + sqrt(a + b)*b)) - ((sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))*b - sqrt(a + b)*b)/((sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^2 - b)

Mupad [F(-1)]

Timed out.

$$\int \cot(x) \sqrt{a + b \cot^4(x)} dx = \int \cot(x) \sqrt{b \cot^4(x) + a} dx$$

[In] int(cot(x)*(a + b*cot(x)^4)^(1/2),x)

[Out] int(cot(x)*(a + b*cot(x)^4)^(1/2), x)

3.61 $\int \cot(x) (a + b \cot^4(x))^{3/2} dx$

Optimal result	381
Rubi [A] (verified)	381
Mathematica [A] (verified)	384
Maple [B] (verified)	384
Fricas [B] (verification not implemented)	385
Sympy [F]	386
Maxima [F]	386
Giac [B] (verification not implemented)	387
Mupad [F(-1)]	387

Optimal result

Integrand size = 15, antiderivative size = 126

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \frac{1}{4} \sqrt{b} (3a + 2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a + b \cot^4(x)}} \right) + \frac{1}{2} (a + b)^{3/2} \operatorname{arctanh} \left(\frac{a - b \cot^2(x)}{\sqrt{a + b} \sqrt{a + b \cot^4(x)}} \right) - \frac{1}{4} (2(a + b) - b \cot^2(x)) \sqrt{a + b \cot^4(x)} - \frac{1}{6} (a + b \cot^4(x))^{3/2}$$

[Out] $1/2*(a+b)^{(3/2)}*\operatorname{arctanh}((a-b*\cot(x)^2)/(a+b)^{(1/2)}/(a+b*\cot(x)^4)^{(1/2)})-1/6*(a+b*\cot(x)^4)^{(3/2)}+1/4*(3*a+2*b)*\operatorname{arctanh}(\cot(x)^2*b^{(1/2)}/(a+b*\cot(x)^4)^{(1/2)})*b^{(1/2)}-1/4*(2*a+2*b-b*\cot(x)^2)*(a+b*\cot(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.533$, Rules used = {3751, 1262, 749, 829, 858, 223, 212, 739}

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \frac{1}{2} (a + b)^{3/2} \operatorname{arctanh} \left(\frac{a - b \cot^2(x)}{\sqrt{a + b} \sqrt{a + b \cot^4(x)}} \right) + \frac{1}{4} \sqrt{b} (3a + 2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a + b \cot^4(x)}} \right) - \frac{1}{6} (a + b \cot^4(x))^{3/2} - \frac{1}{4} (2(a + b) - b \cot^2(x)) \sqrt{a + b \cot^4(x)}$$

[In] $\text{Int}[\text{Cot}[x]*(a + b*\text{Cot}[x]^4)^{(3/2)}, x]$

[Out] $(\sqrt{b}*(3*a + 2*b)*\text{ArcTanh}[(\sqrt{b}*\text{Cot}[x]^2)/\sqrt{a + b*\text{Cot}[x]^4}])/4 + ((a + b)^{(3/2)}*\text{ArcTanh}[(a - b*\text{Cot}[x]^2)/(\sqrt{a + b}*\sqrt{a + b*\text{Cot}[x]^4})])/2 - ((2*(a + b) - b*\text{Cot}[x]^2)*\sqrt{a + b*\text{Cot}[x]^4})/4 - (a + b*\text{Cot}[x]^4)^{(3/2)}/6$

Rule 212

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 223

$\text{Int}[1/\sqrt{(a_ + (b_)*(x_)^2)}, x_Symbol] \rightarrow \text{Subst}[\text{Int}[1/(1 - b*x^2), x], x, x/\sqrt{a + b*x^2}] /; \text{FreeQ}\{a, b, x\} \ \&\& \ !\text{GtQ}[a, 0]$

Rule 739

$\text{Int}[1/(((d_ + (e_)*(x_))*\sqrt{(a_ + (c_)*(x_)^2)}), x_Symbol] \rightarrow -\text{Subst}[\text{Int}[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/\sqrt{a + c*x^2}] /; \text{FreeQ}\{a, c, d, e, x\}$

Rule 749

$\text{Int}(((d_ + (e_)*(x_))^{(m_)}*((a_ + (c_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[(d + e*x)^{(m + 1)}*((a + c*x^2)^p/(e*(m + 2*p + 1))), x] + \text{Dist}[2*(p/(e*(m + 2*p + 1))), \text{Int}[(d + e*x)^m*\text{Simp}[a*e - c*d*x, x]*(a + c*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}\{a, c, d, e, m, x\} \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{NeQ}[m + 2*p + 1, 0] \ \&\& \ (!\text{RationalQ}[m] \ || \ \text{LtQ}[m, 1]) \ \&\& \ !\text{ILtQ}[m + 2*p, 0] \ \&\& \ \text{IntQuadraticQ}[a, 0, c, d, e, m, p, x]$

Rule 829

$\text{Int}(((d_ + (e_)*(x_))^{(m_)}*((f_ + (g_)*(x_))*((a_ + (c_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[(d + e*x)^{(m + 1)}*(c*e*f*(m + 2*p + 2) - g*c*d*(2*p + 1) + g*c*e*(m + 2*p + 1)*x)*((a + c*x^2)^p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), x] + \text{Dist}[2*(p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), \text{Int}[(d + e*x)^m*(a + c*x^2)^{(p - 1)}*\text{Simp}[f*a*c*e^2*(m + 2*p + 2) + a*c*d*e*g*m - (c^2*f*d*e*(m + 2*p + 2) - g*(c^2*d^2*(2*p + 1) + a*c*e^2*(m + 2*p + 1)))*x, x], x] /; \text{FreeQ}\{a, c, d, e, f, g, m, x\} \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ (\text{IntegerQ}[p] \ || \ !\text{RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 0])) \ \&\& \ !\text{ILtQ}[m + 2*p, 0] \ \&\& \ (\text{IntegerQ}[m] \ || \ \text{IntegerQ}[p] \ || \ \text{IntegersQ}[2*m, 2*p])$

Rule 858

$\text{Int}(((d_ + (e_)*(x_))^{(m_)}*((f_ + (g_)*(x_))*((a_ + (c_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Dist}[g/e, \text{Int}[(d + e*x)^{(m + 1)}*(a + c*x^2)^p, x], x] + D$

ist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && !IGtQ[m, 0]

Rule 1262

Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]

Rule 3751

Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*((c_)*tan[(e_) + (f_)*(x_)]^(n_))^(p_)), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && RationalQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{x(a+bx^4)^{3/2}}{1+x^2} dx, x, \cot(x)\right) \\
 &= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{(a+bx^2)^{3/2}}{1+x} dx, x, \cot^2(x)\right)\right) \\
 &= -\frac{1}{6}(a+b\cot^4(x))^{3/2} - \frac{1}{2}\text{Subst}\left(\int \frac{(a-bx)\sqrt{a+bx^2}}{1+x} dx, x, \cot^2(x)\right) \\
 &= -\frac{1}{4}(2(a+b) - b\cot^2(x))\sqrt{a+b\cot^4(x)} - \frac{1}{6}(a+b\cot^4(x))^{3/2} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{ab(2a+b)-b^2(3a+2b)x}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)}{4b} \\
 &= -\frac{1}{4}(2(a+b) - b\cot^2(x))\sqrt{a+b\cot^4(x)} - \frac{1}{6}(a+b\cot^4(x))^{3/2} \\
 &\quad - \frac{1}{2}(a+b)^2\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right) \\
 &\quad + \frac{1}{4}(b(3a+2b))\text{Subst}\left(\int \frac{1}{\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4}(2(a+b) - b \cot^2(x)) \sqrt{a + b \cot^4(x)} - \frac{1}{6}(a + b \cot^4(x))^{3/2} \\
&\quad + \frac{1}{2}(a+b)^2 \text{Subst} \left(\int \frac{1}{a+b-x^2} dx, x, \frac{a-b \cot^2(x)}{\sqrt{a+b \cot^4(x)}} \right) \\
&\quad\quad + \frac{1}{4}(b(3a+2b)) \text{Subst} \left(\int \frac{1}{1-bx^2} dx, x, \frac{\cot^2(x)}{\sqrt{a+b \cot^4(x)}} \right) \\
&= \frac{1}{4} \sqrt{b}(3a+2b) \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a+b \cot^4(x)}} \right) \\
&\quad + \frac{1}{2}(a+b)^{3/2} \operatorname{arctanh} \left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}} \right) \\
&\quad - \frac{1}{4}(2(a+b) - b \cot^2(x)) \sqrt{a + b \cot^4(x)} - \frac{1}{6}(a + b \cot^4(x))^{3/2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 4.81 (sec) , antiderivative size = 167, normalized size of antiderivative = 1.33

$$\begin{aligned}
\int \cot(x) (a + b \cot^4(x))^{3/2} dx &= \frac{1}{12} \left(6\sqrt{b}(a+b) \operatorname{arctanh} \left(\frac{\sqrt{b} \cot^2(x)}{\sqrt{a+b \cot^4(x)}} \right) \right. \\
&\quad \left. + 6(a+b)^{3/2} \operatorname{arctanh} \left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}} \right) - \sqrt{a+b \cot^4(x)} (8a+6b-3b \cot^2(x)+2b \cot^4(x)) + \frac{3\sqrt{a}\sqrt{b} \operatorname{arcsinh} \left(\frac{\cot^2(x)}{\sqrt{a+b \cot^4(x)}} \right)}{12} \right)
\end{aligned}$$

[In] Integrate[Cot[x]*(a + b*Cot[x]^4)^(3/2),x]

[Out] (6*Sqrt[b]*(a + b)*ArcTanh[(Sqrt[b]*Cot[x]^2)/Sqrt[a + b*Cot[x]^4]] + 6*(a + b)^(3/2)*ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])] - Sqrt[a + b*Cot[x]^4]*(8*a + 6*b - 3*b*Cot[x]^2 + 2*b*Cot[x]^4) + (3*Sqrt[a]*Sqrt[b]*ArcSinh[(Sqrt[b]*Cot[x]^2)/Sqrt[a]]*Sqrt[a + b*Cot[x]^4])/Sqrt[1 + (b*Cot[x]^4)/a])/12

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 244 vs. 2(103) = 206.

Time = 0.07 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.94

method	result
derivativedivides	$\frac{b^{\frac{3}{2}} \ln\left(\sqrt{b} \cot(x)^2 + \sqrt{a+b \cot(x)^4}\right)}{2} - \frac{b^2 \left(\frac{\cot(x)^4 \sqrt{a+b \cot(x)^4}}{3b} - \frac{2a \sqrt{a+b \cot(x)^4}}{3b^2}\right)}{2} - \frac{b \sqrt{a+b \cot(x)^4}}{2} + \sqrt{b} a \ln$
default	$\frac{b^{\frac{3}{2}} \ln\left(\sqrt{b} \cot(x)^2 + \sqrt{a+b \cot(x)^4}\right)}{2} - \frac{b^2 \left(\frac{\cot(x)^4 \sqrt{a+b \cot(x)^4}}{3b} - \frac{2a \sqrt{a+b \cot(x)^4}}{3b^2}\right)}{2} - \frac{b \sqrt{a+b \cot(x)^4}}{2} + \sqrt{b} a \ln$

[In] `int(cot(x)*(a+b*cot(x)^4)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}b^{3/2} \ln(b^{1/2} \cot(x)^2 + (a+b \cot(x)^4)^{1/2}) - \frac{1}{2}b^2 \left(\frac{1}{3} \cot(x)^4 / b (a+b \cot(x)^4)^{1/2} - \frac{2}{3} a/b^2 (a+b \cot(x)^4)^{1/2}\right) - \frac{1}{2}b (a+b \cot(x)^4)^{1/2} + b^{1/2} a \ln(b^{1/2} \cot(x)^2 + (a+b \cot(x)^4)^{1/2}) + \frac{1}{2}b^2 \left(\frac{1}{2} \cot(x)^2 / b (a+b \cot(x)^4)^{1/2} - \frac{1}{2} a/b^{3/2} \ln(b^{1/2} \cot(x)^2 + (a+b \cot(x)^4)^{1/2})\right) - a (a+b \cot(x)^4)^{1/2} + \frac{1}{2} (a^2 + 2ab + b^2) / (a+b)^{1/2} \ln((2a + 2b - 2b \cot(x)^2 + 1) + 2(a+b)^{1/2} (b \cot(x)^2 + 1)^2 - 2b \cot(x)^2 + 1) / (a+b)^{1/2} / (b \cot(x)^2 + 1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 357 vs. $2(104) = 208$.

Time = 0.48 (sec) , antiderivative size = 1486, normalized size of antiderivative = 11.79

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \text{Too large to display}$$

[In] `integrate(cot(x)*(a+b*cot(x)^4)^(3/2),x, algorithm="fricas")`

[Out] $\frac{1}{24} (6((a+b)\cos(2x)^2 - 2(a+b)\cos(2x) + a+b)\sqrt{a+b}) \log\left(\frac{1}{2}(a^2 + 2ab + b^2)\cos(2x)^2 + \frac{1}{2}a^2 + \frac{1}{2}b^2 + \frac{1}{2}((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{a+b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}} - \frac{(a^2 - b^2)\cos(2x)}{\cos(2x)^2 - 2\cos(2x) + 1}\right) + 3\left(\frac{(3a+2b)\cos(2x)^2 - 2(3a+2b)\cos(2x) + 3a+2b}{\cos(2x)^2 - 2\cos(2x) + 1}\right)\sqrt{b} \log\left(-\frac{(a+2b)\cos(2x)^2 - 2(\cos(2x)^2 - 1)\sqrt{b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}} - 2(a-2b)\cos(2x) + a+2b}{\cos(2x)^2 - 2\cos(2x) + 1} - \frac{2((8a+11b)\cos(2x)^2 - 8(2a+b)\cos(2x) + 8a+5b)\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}}}{\cos(2x)^2 - 2\cos(2x) + 1}\right) + \frac{1}{12} \left(3\left(\frac{(3a+2b)\cos(2x)^2 - 2(3a+2b)\cos(2x) + 3a+2b}{\cos(2x)^2 - 2\cos(2x) + 1}\right)\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}}}{\cos(2x) - 1}\right) / (b\cos(2x) + b) + 3\left(\frac{(a+b)\cos(2x)^2 - 2(a+b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}\right)\sqrt{a+b} \log\left(\frac{1}{2}(a^2 + 2ab + b^2)\cos(2x)^2 + \frac{1}{2}a^2 + \frac{1}{2}b^2 + \frac{1}{2}((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{a+b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}}\right)$

$$\begin{aligned}
& b) \cos(2x) + a + b) / (\cos(2x)^2 - 2\cos(2x) + 1)) - (a^2 - b^2) \cos(2x) \\
&) - ((8a + 11b) \cos(2x)^2 - 8(2a + b) \cos(2x) + 8a + 5b) \sqrt{((a + \\
& b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + b) / (\cos(2x)^2 - 2\cos(2x) + 1))} \\
&) / (\cos(2x)^2 - 2\cos(2x) + 1), -1/24 * (12 * ((a + b) \cos(2x)^2 - 2(a + b) * \\
& \cos(2x) + a + b) \sqrt{-a - b} \arctan(((a + b) \cos(2x)^2 - 2a \cos(2x) + \\
& a - b) \sqrt{-a - b} \sqrt{((a + b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + b) / \\
& (\cos(2x)^2 - 2\cos(2x) + 1)) / ((a^2 + 2ab + b^2) \cos(2x)^2 + a^2 + 2ab \\
& + b^2 - 2(a^2 - b^2) \cos(2x)) - 3 * ((3a + 2b) \cos(2x)^2 - 2(3a + 2 \\
& * b) \cos(2x) + 3a + 2b) \sqrt{b} \log(-((a + 2b) \cos(2x)^2 - 2(\cos(2x) \\
& ^2 - 1) \sqrt{b} \sqrt{((a + b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + b) / (\cos(\\
& 2x)^2 - 2\cos(2x) + 1)) - 2(a - 2b) \cos(2x) + a + 2b) / (\cos(2x)^2 - 2 \\
& * \cos(2x) + 1)) + 2 * ((8a + 11b) \cos(2x)^2 - 8(2a + b) \cos(2x) + 8a + \\
& 5b) \sqrt{((a + b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + b) / (\cos(2x)^2 - \\
& 2\cos(2x) + 1))} / (\cos(2x)^2 - 2\cos(2x) + 1), -1/12 * (6 * ((a + b) \cos(2x) \\
& ^2 - 2(a + b) \cos(2x) + a + b) \sqrt{-a - b} \arctan(((a + b) \cos(2x)^2 - \\
& 2a \cos(2x) + a - b) \sqrt{-a - b} \sqrt{((a + b) \cos(2x)^2 - 2(a - b) \cos \\
& (2x) + a + b) / (\cos(2x)^2 - 2\cos(2x) + 1)) / ((a^2 + 2ab + b^2) \cos(2x) \\
& ^2 + a^2 + 2ab + b^2 - 2(a^2 - b^2) \cos(2x)) - 3 * ((3a + 2b) \cos(2x) \\
& ^2 - 2(3a + 2b) \cos(2x) + 3a + 2b) \sqrt{-b} \arctan(\sqrt{-b} \sqrt{((a \\
& + b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + b) / (\cos(2x)^2 - 2\cos(2x) + 1) \\
&) * (\cos(2x) - 1) / (b \cos(2x) + b)) + ((8a + 11b) \cos(2x)^2 - 8(2a + b) \\
& * \cos(2x) + 8a + 5b) \sqrt{((a + b) \cos(2x)^2 - 2(a - b) \cos(2x) + a + \\
& b) / (\cos(2x)^2 - 2\cos(2x) + 1))} / (\cos(2x)^2 - 2\cos(2x) + 1)]
\end{aligned}$$

Sympy [F]

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \int (a + b \cot^4(x))^{\frac{3}{2}} \cot(x) dx$$

[In] integrate(cot(x)*(a+b*cot(x)**4)**(3/2),x)

[Out] Integral((a + b*cot(x)**4)**(3/2)*cot(x), x)

Maxima [F]

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \int (b \cot(x)^4 + a)^{\frac{3}{2}} \cot(x) dx$$

[In] integrate(cot(x)*(a+b*cot(x)^4)^(3/2),x, algorithm="maxima")

[Out] integrate((b*cot(x)^4 + a)^(3/2)*cot(x), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 445 vs. 2(104) = 208.

Time = 0.54 (sec) , antiderivative size = 445, normalized size of antiderivative = 3.53

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx =$$

$$\frac{(3ab + 2b^2) \arctan\left(-\frac{\sqrt{a+b}\sin(x)^2 - \sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}}{\sqrt{-b}}\right)}{2\sqrt{-b}}$$

$$- \frac{(a^2 + 2ab + b^2) \log\left(\left|-\left(\sqrt{a+b}\sin(x)^2 - \sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}\right)(a+b) + \sqrt{a+bb}\right|\right)}{2\sqrt{a+b}}$$

$$+ 3\left(\sqrt{a+b}\sin(x)^2 - \sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}\right)^5 (5ab + 6b^2) + 8\left(\sqrt{a+b}\sin(x)^2 - \sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}\right)^3$$

[In] integrate(cot(x)*(a+b*cot(x)^4)^(3/2),x, algorithm="giac")

[Out] -1/2*(3*a*b + 2*b^2)*arctan(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))/sqrt(-b))/sqrt(-b) - 1/2*(a^2 + 2*a*b + b^2)*log(abs(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))*(a + b) + sqrt(a + b)*b))/sqrt(a + b) - 1/6*(3*(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^5*(5*a*b + 6*b^2) + 8*(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^3*b^3 - 12*(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^4*(a*b + 3*b^2)*sqrt(a + b) + 12*(a*b^2 + b^3)*(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^2*sqrt(a + b) + 3*(3*a*b^3 + 2*b^4)*(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b)) - 8*(a*b^3 + b^4)*sqrt(a + b))/((sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))^2 - b)^3

Mupad [F(-1)]

Timed out.

$$\int \cot(x) (a + b \cot^4(x))^{3/2} dx = \int \cot(x) (b \cot^4(x) + a)^{3/2} dx$$

[In] int(cot(x)*(a + b*cot(x)^4)^(3/2),x)

[Out] int(cot(x)*(a + b*cot(x)^4)^(3/2), x)

3.62 $\int \frac{\cot(x)}{\sqrt{a+b \cot^4(x)}} dx$

Optimal result	388
Rubi [A] (verified)	388
Mathematica [A] (verified)	389
Maple [A] (verified)	390
Fricas [B] (verification not implemented)	390
Sympy [F]	391
Maxima [F]	391
Giac [A] (verification not implemented)	391
Mupad [F(-1)]	392

Optimal result

Integrand size = 15, antiderivative size = 41

$$\int \frac{\cot(x)}{\sqrt{a+b \cot^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2\sqrt{a+b}}$$

[Out] $1/2*\operatorname{arctanh}((a-b*\cot(x)^2)/(a+b)^{(1/2)/(a+b*\cot(x)^4)^{(1/2))}/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3751, 1262, 739, 212}

$$\int \frac{\cot(x)}{\sqrt{a+b \cot^4(x)}} dx = \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2\sqrt{a+b}}$$

[In] `Int[Cot[x]/Sqrt[a + b*Cot[x]^4],x]`

[Out] `ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])]/(2*Sqrt[a + b])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 739

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (c_)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ
[{a, c, d, e}, x]
```

Rule 1262

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
[{a, c, d, e, p, q}, x]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)]^(n_))^(p_)), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x]] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx^4}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)\right) \\
&= \frac{1}{2}\text{Subst}\left(\int \frac{1}{a+b-x^2} dx, x, \frac{a-b\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right) \\
&= \frac{\text{arctanh}\left(\frac{a-b\cot^2(x)}{\sqrt{a+b}\sqrt{a+b\cot^4(x)}}\right)}{2\sqrt{a+b}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00

$$\int \frac{\cot(x)}{\sqrt{a+b\cot^4(x)}} dx = \frac{\text{arctanh}\left(\frac{a-b\cot^2(x)}{\sqrt{a+b}\sqrt{a+b\cot^4(x)}}\right)}{2\sqrt{a+b}}$$

```
[In] Integrate[Cot[x]/Sqrt[a + b*Cot[x]^4], x]
```

```
[Out] ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])]/(2*Sqrt[a + b])
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.59

method	result	size
derivativedivides	$\frac{\ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2\sqrt{a+b}}$	65
default	$\frac{\ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2\sqrt{a+b}}$	65

[In] `int(cot(x)/(a+b*cot(x)^4)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}(a+b)^{1/2} \ln\left(\frac{(2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b})}{\cot(x)^2+1}\right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. $2(35) = 70$.

Time = 0.39 (sec) , antiderivative size = 264, normalized size of antiderivative = 6.44

$$\int \frac{\cot(x)}{\sqrt{a+b\cot^4(x)}} dx$$

$$= \left[\frac{\log\left(\frac{1}{2}(a^2+2ab+b^2)\cos(2x)^2 + \frac{1}{2}a^2 + \frac{1}{2}b^2 + \frac{1}{2}((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{a+b}\sqrt{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}}{4\sqrt{a+b}}\right)}{2(a+b)} \right]$$

$$+ \frac{\sqrt{-a-b} \arctan\left(\frac{((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{-a-b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}}}{(a^2+2ab+b^2)\cos(2x)^2 + a^2+2ab+b^2 - 2(a^2-b^2)\cos(2x)}\right)}{2(a+b)}$$

[In] `integrate(cot(x)/(a+b*cot(x)^4)^(1/2),x, algorithm="fricas")`

[Out] $\left[\frac{1}{4}\log\left(\frac{1}{2}(a^2+2ab+b^2)\cos(2x)^2 + \frac{1}{2}a^2 + \frac{1}{2}b^2 + \frac{1}{2}((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{a+b}\sqrt{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}}{(a^2+2ab+b^2)\cos(2x)^2 + a^2+2ab+b^2 - 2(a^2-b^2)\cos(2x)}\right), -\frac{1}{2}\sqrt{-a-b}\arctan\left(\frac{((a+b)\cos(2x)^2 - 2a\cos(2x) + a-b)\sqrt{-a-b}\sqrt{\frac{(a+b)\cos(2x)^2 - 2(a-b)\cos(2x) + a+b}{\cos(2x)^2 - 2\cos(2x) + 1}}}{(a^2+2ab+b^2)\cos(2x)^2 + a^2+2ab+b^2 - 2(a^2-b^2)\cos(2x)}\right)\right]/(a+b)$

Sympy [F]

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^4(x)}} dx = \int \frac{\cot(x)}{\sqrt{a + b \cot^4(x)}} dx$$

```
[In] integrate(cot(x)/(a+b*cot(x)**4)**(1/2),x)
```

```
[Out] Integral(cot(x)/sqrt(a + b*cot(x)**4), x)
```

Maxima [F]

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^4(x)}} dx = \int \frac{\cot(x)}{\sqrt{b \cot^4(x) + a}} dx$$

```
[In] integrate(cot(x)/(a+b*cot(x)^4)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(cot(x)/sqrt(b*cot(x)^4 + a), x)
```

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.41

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^4(x)}} dx = \frac{\log \left(\left| - \left(\sqrt{a + b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b} \right) (a + b) + \sqrt{a + bb} \right| \right)}{2\sqrt{a + b}}$$

```
[In] integrate(cot(x)/(a+b*cot(x)^4)^(1/2),x, algorithm="giac")
```

```
[Out] -1/2*log(abs(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*si
n(x)^2 + b))*(a + b) + sqrt(a + b)*b))/sqrt(a + b)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(x)}{\sqrt{a + b \cot^4(x)}} dx = \int \frac{\cot(x)}{\sqrt{b \cot^4(x) + a}} dx$$

```
[In] int(cot(x)/(a + b*cot(x)^4)^(1/2),x)
```

```
[Out] int(cot(x)/(a + b*cot(x)^4)^(1/2), x)
```


$$3.63 \quad \int \frac{\cot(x)}{(a+b \cot^4(x))^{3/2}} dx$$

Optimal result	393
Rubi [A] (verified)	393
Mathematica [A] (verified)	395
Maple [B] (verified)	395
Fricas [B] (verification not implemented)	396
Sympy [F]	397
Maxima [F]	397
Giac [A] (verification not implemented)	397
Mupad [F(-1)]	398

Optimal result

Integrand size = 15, antiderivative size = 74

$$\int \frac{\cot(x)}{(a+b \cot^4(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a+b \cot^2(x)}{2a(a+b)\sqrt{a+b \cot^4(x)}}$$

[Out] 1/2*arctanh((a-b*cot(x)^2)/(a+b)^(1/2)/(a+b*cot(x)^4)^(1/2))/(a+b)^(3/2)+1/2*(-a-b*cot(x)^2)/a/(a+b)/(a+b*cot(x)^4)^(1/2)

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3751, 1262, 755, 12, 739, 212}

$$\int \frac{\cot(x)}{(a+b \cot^4(x))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a+b \cot^2(x)}{2a(a+b)\sqrt{a+b \cot^4(x)}}$$

[In] Int[Cot[x]/(a + b*Cot[x]^4)^(3/2),x]

[Out] ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])]/(2*(a + b)^(3/2)) - (a + b*Cot[x]^2)/(2*a*(a + b)*Sqrt[a + b*Cot[x]^4])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 739

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (c_)*(x_)^2]), x_Symbol] := -Subst[
Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ
[{a, c, d, e}, x]
```

Rule 755

```
Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[
(-(d + e*x)^(m + 1))*(a*e + c*d*x)*((a + c*x^2)^(p + 1)/(2*a*(p + 1)*(c*d^2
+ a*e^2))), x] + Dist[1/(2*a*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*Sim
p[c*d^2*(2*p + 3) + a*e^2*(m + 2*p + 3) + c*e*d*(m + 2*p + 4)*x, x]*(a + c*
x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0]
&& LtQ[p, -1] && IntQuadraticQ[a, 0, c, d, e, m, p, x]
```

Rule 1262

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
[{a, c, d, e, p, q}, x]
```

Rule 3751

```
Int[((d_)*tan[(e_) + (f_)*(x_)])^(m_)*((a_) + (b_)*((c_)*tan[(e_) +
(f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],
x]}, Dist[c*(ff/f), Subst[Int[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff
^2*x^2)], x], x, c*(Tan[e + f*x]/ff)], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && (IGtQ[p, 0] || EqQ[n, 2] || EqQ[n, 4] || (IntegerQ[p] && Ration
alQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx^4)^{3/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)(a+bx^2)^{3/2}} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{a+b\cot^2(x)}{2a(a+b)\sqrt{a+b\cot^4(x)}} - \frac{\text{Subst}\left(\int \frac{a}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)}{2a(a+b)}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{a + b \cot^2(x)}{2a(a+b)\sqrt{a+b \cot^4(x)}} - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)}{2(a+b)} \\
&= -\frac{a + b \cot^2(x)}{2a(a+b)\sqrt{a+b \cot^4(x)}} + \frac{\text{Subst}\left(\int \frac{1}{a+b-x^2} dx, x, \frac{a-b \cot^2(x)}{\sqrt{a+b \cot^4(x)}}\right)}{2(a+b)} \\
&= \frac{\text{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b}\sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{3/2}} - \frac{a + b \cot^2(x)}{2a(a+b)\sqrt{a+b \cot^4(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.33 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.99

$$\int \frac{\cot(x)}{(a+b \cot^4(x))^{3/2}} dx = \frac{1}{2} \left(\frac{\text{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b}\sqrt{a+b \cot^4(x)}}\right)}{(a+b)^{3/2}} - \frac{a + b \cot^2(x)}{a(a+b)\sqrt{a+b \cot^4(x)}} \right)$$

[In] Integrate[Cot[x]/(a + b*Cot[x]^4)^(3/2), x]

[Out] (ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4]])/(a + b)^(3/2) - (a + b*Cot[x]^2)/(a*(a + b)*Sqrt[a + b*Cot[x]^4]))/2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. 2(65) = 130.

Time = 1.02 (sec) , antiderivative size = 248, normalized size of antiderivative = 3.35

method	result
derivativedivides	$ -\frac{b \ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2(b+\sqrt{-ab})(-b+\sqrt{-ab})\sqrt{a+b}} + \frac{\sqrt{(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2 b-2\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-a}}{b})}}{4a(-b+\sqrt{-ab})(\cot(x)^2+\frac{\sqrt{-a}}{b})} $
default	$ -\frac{b \ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2(b+\sqrt{-ab})(-b+\sqrt{-ab})\sqrt{a+b}} + \frac{\sqrt{(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2 b-2\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-a}}{b})}}{4a(-b+\sqrt{-ab})(\cot(x)^2+\frac{\sqrt{-a}}{b})} $

[In] int(cot(x)/(a+b*cot(x)^4)^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/2*b/(b+(-a*b)^(1/2))/(-b+(-a*b)^(1/2))/(a+b)^(1/2)*ln((2*a+2*b-2*b*(cot(x)^2+1)+2*(a+b)^(1/2)*(b*(cot(x)^2+1)^2-2*b*(cot(x)^2+1)+a+b)^(1/2))/(cot(x)^2+1))+1/4/a/(-b+(-a*b)^(1/2))/(cot(x)^2+(-a*b)^(1/2)/b)*((cot(x)^2+(-a*b)^(1/2)/b)^2*b-2*(-a*b)^(1/2)*(cot(x)^2+(-a*b)^(1/2)/b))^(1/2)-1/4/a/(b+(-a*

$$b^{1/2})/(\cot(x)^2 - (-a*b)^{1/2}/b)*((\cot(x)^2 - (-a*b)^{1/2}/b)^2*b + 2*(-a*b)^{1/2}*(\cot(x)^2 - (-a*b)^{1/2}/b))^{1/2}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 338 vs. 2(64) = 128.

Time = 0.45 (sec) , antiderivative size = 670, normalized size of antiderivative = 9.05

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{3/2}} dx = \left[\frac{((a^2 + ab) \cos(2x))^2 + a^2 + ab - 2(a^2 - ab) \cos(2x)) \sqrt{a+b} \log\left(\frac{1}{2}(a^2 + 2ab + b^2) \cos(2x) + a - b\right)}{2(a^4 + 3a^3b + 3a^2b^2 + ab^3 + (a^4 + 3a^3b + 3a^2b^2 + ab^3) \cos(2x)^2 - 2(a^4 + a^3b - a^2b^2 - ab^3) \cos(2x))} \sqrt{-a-b} \arctan\left(\frac{((a+b) \cos(2x)^2 - 2a \cos(2x) + a-b) \sqrt{-a-b}}{(a^2 + 2ab + b^2) \cos(2x)^2 + a^2 + 2ab + b^2}\right) \right]$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(3/2),x, algorithm="fricas")

[Out] [1/4*(((a^2 + a*b)*cos(2*x)^2 + a^2 + a*b - 2*(a^2 - a*b)*cos(2*x))*sqrt(a + b)*log(1/2*(a^2 + 2*a*b + b^2)*cos(2*x)^2 + 1/2*a^2 + 1/2*b^2 + 1/2*((a + b)*cos(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt(a + b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)) - (a^2 - b^2)*cos(2*x)) - 2*((a^2 - b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 + a*b)*cos(2*x))*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)))/(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cos(2*x)^2 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3)*cos(2*x)), -1/2*(((a^2 + a*b)*cos(2*x)^2 + a^2 + a*b - 2*(a^2 - a*b)*cos(2*x))*sqrt(-a - b)*arctan(((a + b)*cos(2*x)^2 - 2*a*cos(2*x) + a - b)*sqrt(-a - b)*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)))/((a^2 + 2*a*b + b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*cos(2*x)) + ((a^2 - b^2)*cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 + a*b)*cos(2*x))*sqrt(((a + b)*cos(2*x)^2 - 2*(a - b)*cos(2*x) + a + b)/(cos(2*x)^2 - 2*cos(2*x) + 1)))/(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cos(2*x)^2 - 2*(a^4 + a^3*b - a^2*b^2 - a*b^3)*cos(2*x))]

Sympy [F]

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{3/2}} dx = \int \frac{\cot(x)}{(a + b \cot^4(x))^{\frac{3}{2}}} dx$$

[In] integrate(cot(x)/(a+b*cot(x)**4)**(3/2),x)

[Out] Integral(cot(x)/(a + b*cot(x)**4)**(3/2), x)

Maxima [F]

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{3/2}} dx = \int \frac{\cot(x)}{(b \cot^4(x) + a)^{\frac{3}{2}}} dx$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(3/2),x, algorithm="maxima")

[Out] integrate(cot(x)/(b*cot(x)^4 + a)^(3/2), x)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.50

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{3/2}} dx = -\frac{\frac{(a-b)\sin(x)^2}{a^2+ab} + \frac{b}{a^2+ab}}{2\sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}} - \frac{\log\left(\left|-\left(\sqrt{a+b}\sin(x)^2 - \sqrt{a\sin(x)^4 + b\sin(x)^4 - 2b\sin(x)^2 + b}\right)\sqrt{a+b} + b\right|\right)}{2(a+b)^{\frac{3}{2}}}$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(3/2),x, algorithm="giac")

[Out] -1/2*((a - b)*sin(x)^2/(a^2 + a*b) + b/(a^2 + a*b))/sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b) - 1/2*log(abs(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))*sqrt(a + b) + b))/(a + b)^(3/2)

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{3/2}} dx = \int \frac{\cot(x)}{(b \cot(x)^4 + a)^{3/2}} dx$$

```
[In] int(cot(x)/(a + b*cot(x)^4)^(3/2),x)
```

```
[Out] int(cot(x)/(a + b*cot(x)^4)^(3/2), x)
```

$$3.64 \quad \int \frac{\cot(x)}{(a+b \cot^4(x))^{5/2}} dx$$

Optimal result	399
Rubi [A] (verified)	399
Mathematica [A] (verified)	402
Maple [B] (verified)	402
Fricas [B] (verification not implemented)	403
Sympy [F]	404
Maxima [F]	404
Giac [B] (verification not implemented)	404
Mupad [F(-1)]	405

Optimal result

Integrand size = 15, antiderivative size = 117

$$\int \frac{\cot(x)}{(a+b \cot^4(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{5/2}} - \frac{a+b \cot^2(x)}{6a(a+b)(a+b \cot^4(x))^{3/2}} - \frac{3a^2+b(5a+2b) \cot^2(x)}{6a^2(a+b)^2 \sqrt{a+b \cot^4(x)}}$$

[Out] $1/2*\operatorname{arctanh}((a-b*\cot(x)^2)/(a+b)^{(1/2)/(a+b*\cot(x)^4)^{(1/2)})/(a+b)^{(5/2)+1/6*(-a-b*\cot(x)^2)/a/(a+b)/(a+b*\cot(x)^4)^{(3/2)+1/6*(-3*a^2-b*(5*a+2*b))*\cot(x)^2)/a^2/(a+b)^2/(a+b*\cot(x)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.22 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.467$, Rules used = {3751, 1262, 755, 837, 12, 739, 212}

$$\int \frac{\cot(x)}{(a+b \cot^4(x))^{5/2}} dx = -\frac{3a^2+b(5a+2b) \cot^2(x)}{6a^2(a+b)^2 \sqrt{a+b \cot^4(x)}} + \frac{\operatorname{arctanh}\left(\frac{a-b \cot^2(x)}{\sqrt{a+b} \sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{5/2}} - \frac{a+b \cot^2(x)}{6a(a+b)(a+b \cot^4(x))^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[x]/(a+b*\operatorname{Cot}[x]^4)^{(5/2)}, x]$

[Out] $\operatorname{ArcTanh}[(a-b*\operatorname{Cot}[x]^2)/(\operatorname{Sqrt}[a+b]*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^4])]/(2*(a+b)^{(5/2)}) - (a+b*\operatorname{Cot}[x]^2)/(6*a*(a+b)*(a+b*\operatorname{Cot}[x]^4)^{(3/2)}) - (3*a^2+b*(5*a+2*b)*\operatorname{Cot}[x]^2)/(6*a^2*(a+b)^2*\operatorname{Sqrt}[a+b*\operatorname{Cot}[x]^4])$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 739

Int[1/(((d_) + (e_.)*(x_))*Sqrt[(a_) + (c_.)*(x_)^2]), x_Symbol] := -Subst[Int[1/(c*d^2 + a*e^2 - x^2), x], x, (a*e - c*d*x)/Sqrt[a + c*x^2]] /; FreeQ[{a, c, d, e}, x]

Rule 755

Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-d + e*x)^(m + 1)*(a*e + c*d*x)*((a + c*x^2)^(p + 1)/(2*a*(p + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*a*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*Simp[c*d^2*(2*p + 3) + a*e^2*(m + 2*p + 3) + c*e*d*(m + 2*p + 4)*x, x]*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntQuadraticQ[a, 0, c, d, e, m, p, x]

Rule 837

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-d + e*x)^(m + 1)*(f*a*c*e - a*g*c*d + c*(c*d*f + a*e*g)*x)*((a + c*x^2)^(p + 1)/(2*a*c*(p + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*a*c*(p + 1)*(c*d^2 + a*e^2)), Int[(d + e*x)^m*(a + c*x^2)^(p + 1)*Simp[f*(c^2*d^2*(2*p + 3) + a*c*e^2*(m + 2*p + 3)) - a*c*d*e*g*m + c*e*(c*d*f + a*e*g)*(m + 2*p + 4)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 1262

Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]

Rule 3751

Int[((d_.)*tan[(e_.) + (f_.)*(x_)])^(m_.)*((a_) + (b_.)*((c_.)*tan[(e_.) + (f_.)*(x_)])^(n_.))^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x],

$x\}}, \text{Dist}[c*(ff/f), \text{Subst}[\text{Int}[(d*ff*(x/c))^m*((a + b*(ff*x)^n)^p/(c^2 + ff^2*x^2)], x], x, c*(\text{Tan}[e + f*x]/ff)], x]] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& (\text{IGtQ}[p, 0] \|\ \text{EqQ}[n, 2] \|\ \text{EqQ}[n, 4] \|\ (\text{IntegerQ}[p] \&\& \text{RationalQ}[n]))$

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx^4)^{5/2}} dx, x, \cot(x)\right) \\
&= -\left(\frac{1}{2}\text{Subst}\left(\int \frac{1}{(1+x)(a+bx^2)^{5/2}} dx, x, \cot^2(x)\right)\right) \\
&= -\frac{a+b\cot^2(x)}{6a(a+b)(a+b\cot^4(x))^{3/2}} + \frac{\text{Subst}\left(\int \frac{-3a-2b-2bx}{(1+x)(a+bx^2)^{3/2}} dx, x, \cot^2(x)\right)}{6a(a+b)} \\
&= -\frac{a+b\cot^2(x)}{6a(a+b)(a+b\cot^4(x))^{3/2}} - \frac{3a^2+b(5a+2b)\cot^2(x)}{6a^2(a+b)^2\sqrt{a+b\cot^4(x)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{3a^2b}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)}{6a^2b(a+b)^2} \\
&= -\frac{a+b\cot^2(x)}{6a(a+b)(a+b\cot^4(x))^{3/2}} - \frac{3a^2+b(5a+2b)\cot^2(x)}{6a^2(a+b)^2\sqrt{a+b\cot^4(x)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx^2}} dx, x, \cot^2(x)\right)}{2(a+b)^2} \\
&= -\frac{a+b\cot^2(x)}{6a(a+b)(a+b\cot^4(x))^{3/2}} - \frac{3a^2+b(5a+2b)\cot^2(x)}{6a^2(a+b)^2\sqrt{a+b\cot^4(x)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{a+b-x^2} dx, x, \frac{a-b\cot^2(x)}{\sqrt{a+b\cot^4(x)}}\right)}{2(a+b)^2} \\
&= \frac{\text{arctanh}\left(\frac{a-b\cot^2(x)}{\sqrt{a+b}\sqrt{a+b\cot^4(x)}}\right)}{2(a+b)^{5/2}} - \frac{a+b\cot^2(x)}{6a(a+b)(a+b\cot^4(x))^{3/2}} - \frac{3a^2+b(5a+2b)\cot^2(x)}{6a^2(a+b)^2\sqrt{a+b\cot^4(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.97

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \frac{\operatorname{arctanh}\left(\frac{a - b \cot^2(x)}{\sqrt{a+b}\sqrt{a+b \cot^4(x)}}\right)}{2(a+b)^{5/2}} - \frac{a^2(4a+b) + 3ab(2a+b)\cot^2(x) + 3a^2b \cot^4(x) + b^2(5a+2b)\cot^6(x)}{6a^2(a+b)^2(a+b \cot^4(x))^{3/2}}$$

[In] Integrate[Cot[x]/(a + b*Cot[x]^4)^(5/2),x]

[Out] ArcTanh[(a - b*Cot[x]^2)/(Sqrt[a + b]*Sqrt[a + b*Cot[x]^4])]/(2*(a + b)^(5/2)) - (a^2*(4*a + b) + 3*a*b*(2*a + b)*Cot[x]^2 + 3*a^2*b*Cot[x]^4 + b^2*(5*a + 2*b)*Cot[x]^6)/(6*a^2*(a + b)^2*(a + b*Cot[x]^4)^(3/2))

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 585 vs. 2(105) = 210.

Time = 0.14 (sec) , antiderivative size = 586, normalized size of antiderivative = 5.01

method	result
derivativedivides	$\frac{b^2 \ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2(b+\sqrt{-ab})^2(-b+\sqrt{-ab})^2\sqrt{a+b}} - \frac{\sqrt{(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2b-2\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2}}{3\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2}$
default	$\frac{b^2 \ln\left(\frac{2a+2b-2b(\cot(x)^2+1)+2\sqrt{a+b}\sqrt{b(\cot(x)^2+1)^2-2b(\cot(x)^2+1)+a+b}}{\cot(x)^2+1}\right)}{2(b+\sqrt{-ab})^2(-b+\sqrt{-ab})^2\sqrt{a+b}} - \frac{\sqrt{(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2b-2\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2}}{3\sqrt{-ab}(\cot(x)^2+\frac{\sqrt{-ab}}{b})^2}$

[In] int(cot(x)/(a+b*cot(x)^4)^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/2*b^2/(b+(-a*b)^(1/2))^2/(-b+(-a*b)^(1/2))^2/(a+b)^(1/2)*ln((2*a+2*b-2*b*(cot(x)^2+1)+2*(a+b)^(1/2)*(b*(cot(x)^2+1)^2-2*b*(cot(x)^2+1)+a+b)^(1/2))/(cot(x)^2+1))-1/8/(-b+(-a*b)^(1/2))/a*(1/3/(-a*b)^(1/2)/(cot(x)^2+(-a*b)^(1/2)/b)^2*((cot(x)^2+(-a*b)^(1/2)/b)^2*b-2*(-a*b)^(1/2)*(cot(x)^2+(-a*b)^(1/2)/b))^(1/2))-1/3/a/(cot(x)^2+(-a*b)^(1/2)/b)*((cot(x)^2+(-a*b)^(1/2)/b)^2*b-2*(-a*b)^(1/2)*(cot(x)^2+(-a*b)^(1/2)/b))^(1/2))+1/8/(b+(-a*b)^(1/2))/a*(-1/3/(-a*b)^(1/2)/(cot(x)^2-(-a*b)^(1/2)/b)^2*((cot(x)^2-(-a*b)^(1/2)/b)^2*b+2*(-a*b)^(1/2)*(cot(x)^2-(-a*b)^(1/2)/b))^(1/2))-1/3/a/(cot(x)^2-(-a*b)^(1/2)/b)*((cot(x)^2-(-a*b)^(1/2)/b)^2*b+2*(-a*b)^(1/2)*(cot(x)^2-(-a*b)^(1/2)/b))^(1/2))+1/8*(2*(-a*b)^(1/2)-b)/(-b+(-a*b)^(1/2))^2/a^2/(cot(x)^2+(-a*b)^(1/2)/b)*((cot(x)^2+(-a*b)^(1/2)/b)^2*b-2*(-a*b)^(1/2)*(cot(x)^2+(-a*b)^(1/2)/b))^(1/2))-1/8*(2*(-a*b)^(1/2)+b)/(b+(-a*b)^(1/2))^2/a^2/(cot(x)^2-(-a*b)^(1/2)/b)*((cot(x)^2-(-a*b)^(1/2)/b)^2*b+2*(-a*b)^(1/2)*(cot(x)^2-(-a*b)^(1/2)/b))^(1/2))

$(1/2)/b * ((\cot(x)^2 - (-a*b)^{(1/2)/b})^{2*b+2} * (-a*b)^{(1/2)} * (\cot(x)^2 - (-a*b)^{(1/2)/b})^{(1/2)})^{(1/2)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 686 vs. 2(103) = 206.

Time = 0.52 (sec) , antiderivative size = 1365, normalized size of antiderivative = 11.67

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \text{Too large to display}$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(5/2),x, algorithm="fricas")

[Out] $[1/12*(3*((a^4 + 2*a^3*b + a^2*b^2)*\cos(2*x)^4 + a^4 + 2*a^3*b + a^2*b^2 - 4*(a^4 - a^2*b^2)*\cos(2*x)^3 + 2*(3*a^4 - 2*a^3*b + 3*a^2*b^2)*\cos(2*x)^2 - 4*(a^4 - a^2*b^2)*\cos(2*x))*\sqrt{a + b}*\log(1/2*(a^2 + 2*a*b + b^2)*\cos(2*x)^2 + 1/2*a^2 + 1/2*b^2 + 1/2*((a + b)*\cos(2*x)^2 - 2*a*\cos(2*x) + a - b)*\sqrt{a + b}*\sqrt{((a + b)*\cos(2*x)^2 - 2*(a - b)*\cos(2*x) + a + b)/(\cos(2*x)^2 - 2*\cos(2*x) + 1)}) - (a^2 - b^2)*\cos(2*x)) - 4*((2*a^4 + a^3*b - 5*a^2*b^2 - 5*a*b^3 - b^4)*\cos(2*x)^4 + 2*a^4 + 7*a^3*b + 9*a^2*b^2 + 5*a*b^3 + b^4 - 2*(4*a^4 + 2*a^3*b - a^2*b^2 + 2*a*b^3 + b^4)*\cos(2*x)^3 + 12*(a^4 + a^3*b)*\cos(2*x)^2 - 2*(4*a^4 + 8*a^3*b + 3*a^2*b^2 - 2*a*b^3 - b^4)*\cos(2*x))*\sqrt{((a + b)*\cos(2*x)^2 - 2*(a - b)*\cos(2*x) + a + b)/(\cos(2*x)^2 - 2*\cos(2*x) + 1))}/(a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5 + (a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5)*\cos(2*x))^4 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*\cos(2*x)^3 + 2*(3*a^7 + 7*a^6*b + 6*a^5*b^2 + 6*a^4*b^3 + 7*a^3*b^4 + 3*a^2*b^5)*\cos(2*x)^2 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*\cos(2*x)), -1/6*(3*((a^4 + 2*a^3*b + a^2*b^2)*\cos(2*x)^4 + a^4 + 2*a^3*b + a^2*b^2 - 4*(a^4 - a^2*b^2)*\cos(2*x)^3 + 2*(3*a^4 - 2*a^3*b + 3*a^2*b^2)*\cos(2*x)^2 - 4*(a^4 - a^2*b^2)*\cos(2*x))*\sqrt{-a - b}*\arctan(((a + b)*\cos(2*x)^2 - 2*a*\cos(2*x) + a - b)*\sqrt{-a - b}*\sqrt{((a + b)*\cos(2*x)^2 - 2*(a - b)*\cos(2*x) + a + b)/(\cos(2*x)^2 - 2*\cos(2*x) + 1)})/((a^2 + 2*a*b + b^2)*\cos(2*x)^2 + a^2 + 2*a*b + b^2 - 2*(a^2 - b^2)*\cos(2*x))) + 2*((2*a^4 + a^3*b - 5*a^2*b^2 - 5*a*b^3 - b^4)*\cos(2*x)^4 + 2*a^4 + 7*a^3*b + 9*a^2*b^2 + 5*a*b^3 + b^4 - 2*(4*a^4 + 2*a^3*b - a^2*b^2 + 2*a*b^3 + b^4)*\cos(2*x)^3 + 12*(a^4 + a^3*b)*\cos(2*x)^2 - 2*(4*a^4 + 8*a^3*b + 3*a^2*b^2 - 2*a*b^3 - b^4)*\cos(2*x))*\sqrt{((a + b)*\cos(2*x)^2 - 2*(a - b)*\cos(2*x) + a + b)/(\cos(2*x)^2 - 2*\cos(2*x) + 1))}/(a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5 + (a^7 + 5*a^6*b + 10*a^5*b^2 + 10*a^4*b^3 + 5*a^3*b^4 + a^2*b^5)*\cos(2*x))^4 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*\cos(2*x)^3 + 2*(3*a^7 + 7*a^6*b + 6*a^5*b^2 + 6*a^4*b^3 + 7*a^3*b^4 + 3*a^2*b^5)*\cos(2*x)^2 - 4*(a^7 + 3*a^6*b + 2*a^5*b^2 - 2*a^4*b^3 - 3*a^3*b^4 - a^2*b^5)*\cos(2*x))]$

Sympy [F]

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \int \frac{\cot(x)}{(a + b \cot^4(x))^{\frac{5}{2}}} dx$$

[In] integrate(cot(x)/(a+b*cot(x)**4)**(5/2),x)

[Out] Integral(cot(x)/(a + b*cot(x)**4)**(5/2), x)

Maxima [F]

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \int \frac{\cot(x)}{(b \cot^4(x) + a)^{\frac{5}{2}}} dx$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(5/2),x, algorithm="maxima")

[Out] integrate(cot(x)/(b*cot(x)^4 + a)^(5/2), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 276 vs. 2(103) = 206.

Time = 0.31 (sec) , antiderivative size = 276, normalized size of antiderivative = 2.36

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \frac{\left(2 \left(\frac{(2a^3b - a^2b^2 - 4ab^3 - b^4) \sin(x)^2}{a^4b + 2a^3b^2 + a^2b^3} + \frac{3(3ab^3 + b^4)}{a^4b + 2a^3b^2 + a^2b^3} \right) \sin(x)^2 + \frac{3(a^2b^2 - 5ab^3 - 2b^4)}{a^4b + 2a^3b^2 + a^2b^3} \sin(x)^2 + \frac{5ab^3 + 2b^4}{a^4b + 2a^3b^2 + a^2b^3} \right)}{6(a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b)^{\frac{3}{2}}} \log \left(\left| - \left(\sqrt{a + b} \sin(x)^2 - \sqrt{a \sin(x)^4 + b \sin(x)^4 - 2b \sin(x)^2 + b} \right) \sqrt{a + b} + b \right| \right) \frac{1}{2(a^2 + 2ab + b^2)\sqrt{a + b}}$$

[In] integrate(cot(x)/(a+b*cot(x)^4)^(5/2),x, algorithm="giac")

[Out] -1/6*((2*((2*a^3*b - a^2*b^2 - 4*a*b^3 - b^4)*sin(x)^2/(a^4*b + 2*a^3*b^2 + a^2*b^3) + 3*(3*a*b^3 + b^4)/(a^4*b + 2*a^3*b^2 + a^2*b^3))*sin(x)^2 + 3*(a^2*b^2 - 5*a*b^3 - 2*b^4)/(a^4*b + 2*a^3*b^2 + a^2*b^3))*sin(x)^2 + (5*a*b^3 + 2*b^4)/(a^4*b + 2*a^3*b^2 + a^2*b^3))/(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b)^(3/2) - 1/2*log(abs(-(sqrt(a + b)*sin(x)^2 - sqrt(a*sin(x)^4 + b*sin(x)^4 - 2*b*sin(x)^2 + b))*sqrt(a + b) + b))/((a^2 + 2*a*b + b^2)*sqrt(a + b))

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(x)}{(a + b \cot^4(x))^{5/2}} dx = \int \frac{\cot(x)}{(b \cot(x)^4 + a)^{5/2}} dx$$

```
[In] int(cot(x)/(a + b*cot(x)^4)^(5/2),x)
```

```
[Out] int(cot(x)/(a + b*cot(x)^4)^(5/2), x)
```

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 407

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_coun
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;
end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```



```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+str(ExpnType_optimal)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```