

Computer Algebra Independent Integration Tests

Summer 2023 edition

4-Trig-functions/4.4-Cotangent/114-4.4.9-trig^{m-a+b-cot^{n+c-}}
cot^{-2-n-^p}

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 4:56pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	35
4	Appendix	303

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [32]. This is test number [114].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (32)	0.00 (0)
Mathematica	100.00 (32)	0.00 (0)
Fricas	100.00 (32)	0.00 (0)
Maple	62.50 (20)	37.50 (12)
Mupad	0.00 (0)	100.00 (32)
Giac	0.00 (0)	100.00 (32)
Maxima	0.00 (0)	100.00 (32)
Sympy	0.00 (0)	100.00 (32)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	43.750	6.250	50.000	0.000
Maple	18.750	43.750	0.000	37.500
Fricas	9.375	90.625	0.000	0.000
Giac	0.000	0.000	0.000	100.000
Mupad	0.000	0.000	0.000	100.000
Maxima	0.000	0.000	0.000	100.000
Sympy	0.000	0.000	0.000	100.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	12	50.00	50.00	0.00
Mupad	32	0.00	100.00	0.00
Giac	32	15.62	50.00	34.38
Maxima	32	59.38	21.88	18.75
Sympy	32	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maple	1.45
Mathematica	4.59
Fricas	4.90
Rubi	5.18
Sympy	-nan(ind)
Maxima	-nan(ind)
Giac	-nan(ind)
Mupad	-nan(ind)

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Rubi	448.28	1.00	366.50	1.00
Mathematica	569.22	1.36	347.50	1.26
Fricas	14244.88	23.35	5146.00	12.73
Maple	6703958.85	10574.14	4669917.00	5496.79
Sympy	-nan(ind)	-nan(ind)	nan	nan
Maxima	-nan(ind)	-nan(ind)	nan	nan
Giac	-nan(ind)	-nan(ind)	nan	nan
Mupad	-nan(ind)	-nan(ind)	nan	nan

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {32}

Maple {1, 2, 3, 6, 7, 8, 11, 12, 13, 14}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	32

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	23
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 17, 18, 19, 20, 21, 24, 25, 26, 27, 28, 29, 30, 31, 32 }

B grade { 22, 23 }

C grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16 }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Maple**A grade** { 17, 18, 19, 22, 23, 24 }**B grade** { 1, 2, 3, 6, 7, 8, 11, 12, 13, 14, 27, 28, 29, 30 }**C grade** { }**F normal fail** { 20, 21, 25, 26, 31, 32 }**F(-1) timeout fail** { 4, 5, 9, 10, 15, 16 }**F(-2) exception fail** { }**Fricas****A grade** { 21, 25, 26 }**B grade** { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 27, 28, 29, 30, 31, 32 }**C grade** { }**F normal fail** { }**F(-1) timeout fail** { }**F(-2) exception fail** { }**Maxima****A grade** { }**B grade** { }**C grade** { }**F normal fail** { 1, 2, 4, 5, 6, 7, 8, 10, 18, 19, 20, 22, 23, 24, 25, 26, 28, 29, 30 }**F(-1) timeout fail** { 11, 12, 15, 16, 17, 21, 32 }**F(-2) exception fail** { 3, 9, 13, 14, 27, 31 }**Giac****A grade** { }**B grade** { }**C grade** { }**F normal fail** { 6, 7, 8, 9, 10 }**F(-1) timeout fail** { 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32 }**F(-2) exception fail** { 1, 2, 3, 4, 5, 11, 12, 13, 14, 15, 16 }

Mupad

A grade { }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32 }

F(-2) exception fail { }

Sympy

A grade { }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32 }

F(-1) timeout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	547	547	735	9581948	0	24241	0	0	0
N.S.	1	1.00	1.34	17517.27	0.00	44.32	0.00	0.00	0.00
time (sec)	N/A	1.415	6.770	3.548	0.000	6.095	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	384	384	397	9581108	0	23719	0	0	0
N.S.	1	1.00	1.03	24950.80	0.00	61.77	0.00	0.00	0.00
time (sec)	N/A	1.039	1.839	1.742	0.000	5.734	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	294	294	256	9339148	0	5853	0	0	0
N.S.	1	1.00	0.87	31765.81	0.00	19.91	0.00	0.00	0.00
time (sec)	N/A	0.301	0.612	1.041	0.000	2.134	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	349	349	344	0	0	10333	0	0	0
N.S.	1	1.00	0.99	0.00	0.00	29.61	0.00	0.00	0.00
time (sec)	N/A	0.733	0.738	180.000	0.000	1.429	0.000	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	501	501	422	0	0	10495	0	0	0
N.S.	1	1.00	0.84	0.00	0.00	20.95	0.00	0.00	0.00
time (sec)	N/A	0.916	4.501	180.000	0.000	1.633	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	976	976	1214	17768513	0	12721	0	0	0
N.S.	1	1.00	1.24	18205.44	0.00	13.03	0.00	0.00	0.00
time (sec)	N/A	25.180	6.709	6.857	0.000	3.017	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	747	747	381	17768080	0	11897	0	0	0
N.S.	1	1.00	0.51	23785.92	0.00	15.93	0.00	0.00	0.00
time (sec)	N/A	24.327	6.297	3.965	0.000	2.413	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	602	602	324	17767874	0	11351	0	0	0
N.S.	1	1.00	0.54	29514.74	0.00	18.86	0.00	0.00	0.00
time (sec)	N/A	23.711	1.433	2.437	0.000	2.295	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F(-2)	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	570	570	283	0	0	4847	0	0	0
N.S.	1	1.00	0.50	0.00	0.00	8.50	0.00	0.00	0.00
time (sec)	N/A	24.184	0.733	0.000	0.000	0.676	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	691	691	363	0	0	5018	0	0	0
N.S.	1	1.00	0.53	0.00	0.00	7.26	0.00	0.00	0.00
time (sec)	N/A	24.415	2.479	180.000	0.000	0.979	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	1189	1189	2097	13067599	0	89177	0	0	0
N.S.	1	1.00	1.76	10990.41	0.00	75.00	0.00	0.00	0.00
time (sec)	N/A	7.143	7.236	1.567	0.000	35.377	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	865	865	1674	13067692	0	84903	0	0	0
N.S.	1	1.00	1.94	15107.16	0.00	98.15	0.00	0.00	0.00
time (sec)	N/A	5.227	6.909	2.906	0.000	29.832	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	686	686	1419	13067316	0	21092	0	0	0
N.S.	1	1.00	2.07	19048.57	0.00	30.75	0.00	0.00	0.00
time (sec)	N/A	5.095	6.681	1.534	0.000	12.127	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F(-2)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	No	TBD	TBD	TBD	TBD	TBD
size	635	635	460	13066366	0	21090	0	0	0
N.S.	1	1.00	0.72	20576.95	0.00	33.21	0.00	0.00	0.00
time (sec)	N/A	4.229	6.398	1.243	0.000	11.870	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	749	749	934	0	0	39885	0	0	0
N.S.	1	1.00	1.25	0.00	0.00	53.25	0.00	0.00	0.00
time (sec)	N/A	5.126	6.547	180.000	0.000	8.077	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F(-1)	F(-1)	B	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	1008	1008	1401	0	0	40633	0	0	0
N.S.	1	1.00	1.39	0.00	0.00	40.31	0.00	0.00	0.00
time (sec)	N/A	5.549	6.660	180.000	0.000	7.929	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	182	182	261	232	0	2100	0	0	0
N.S.	1	1.00	1.43	1.27	0.00	11.54	0.00	0.00	0.00
time (sec)	N/A	0.670	3.623	0.716	0.000	1.312	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	141	141	212	153	0	1695	0	0	0
N.S.	1	1.00	1.50	1.09	0.00	12.02	0.00	0.00	0.00
time (sec)	N/A	0.348	0.838	0.098	0.000	1.035	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	79	79	139	102	0	433	0	0	0
N.S.	1	1.00	1.76	1.29	0.00	5.48	0.00	0.00	0.00
time (sec)	N/A	0.191	0.576	0.203	0.000	0.462	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	142	142	197	0	0	1141	0	0	0
N.S.	1	1.00	1.39	0.00	0.00	8.04	0.00	0.00	0.00
time (sec)	N/A	0.459	1.185	0.000	0.000	1.009	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-1)	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	249	249	258	0	0	1444	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	5.80	0.00	0.00	0.00
time (sec)	N/A	0.461	1.379	0.000	0.000	1.377	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	270	270	1017	455	0	3019	0	0	0
N.S.	1	1.00	3.77	1.69	0.00	11.18	0.00	0.00	0.00
time (sec)	N/A	0.679	6.499	0.122	0.000	2.950	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	209	777	318	0	2344	0	0	0
N.S.	1	1.00	3.72	1.52	0.00	11.22	0.00	0.00	0.00
time (sec)	N/A	0.406	6.358	0.102	0.000	2.337	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	179	246	217	0	1932	0	0	0
N.S.	1	1.00	1.37	1.21	0.00	10.79	0.00	0.00	0.00
time (sec)	N/A	0.276	2.464	0.086	0.000	1.748	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	203	203	254	0	0	2522	0	0	0
N.S.	1	1.00	1.25	0.00	0.00	12.42	0.00	0.00	0.00
time (sec)	N/A	0.315	0.768	0.000	0.000	0.705	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	435	435	235	0	0	1282	0	0	0
N.S.	1	1.00	0.54	0.00	0.00	2.95	0.00	0.00	0.00
time (sec)	N/A	0.646	1.290	0.000	0.000	1.759	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F(-2)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	236	236	339	686	0	6011	0	0	0
N.S.	1	1.00	1.44	2.91	0.00	25.47	0.00	0.00	0.00
time (sec)	N/A	0.641	5.141	0.590	0.000	2.596	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	160	160	204	507	0	1743	0	0	0
N.S.	1	1.00	1.28	3.17	0.00	10.89	0.00	0.00	0.00
time (sec)	N/A	0.467	1.974	0.087	0.000	0.630	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	153	195	457	0	1717	0	0	0
N.S.	1	1.00	1.27	2.99	0.00	11.22	0.00	0.00	0.00
time (sec)	N/A	0.336	1.709	0.109	0.000	0.622	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	156	264	406	0	1771	0	0	0
N.S.	1	1.00	1.69	2.60	0.00	11.35	0.00	0.00	0.00
time (sec)	N/A	0.277	10.901	0.093	0.000	0.660	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-2)	B	F	F(-1)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	280	280	351	0	0	4153	0	0	0
N.S.	1	1.00	1.25	0.00	0.00	14.83	0.00	0.00	0.00
time (sec)	N/A	0.475	13.755	0.000	0.000	2.669	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F(-1)	B	F	F(-1)	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	478	478	562	0	0	5274	0	0	0
N.S.	1	1.00	1.18	0.00	0.00	11.03	0.00	0.00	0.00
time (sec)	N/A	0.667	15.974	0.000	0.000	3.339	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [6] had the largest ratio of [.424200000000000021]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	15	10	1.00	33	0.303
2	A	11	8	1.00	33	0.242
3	A	6	4	1.00	31	0.129
4	A	10	7	1.00	31	0.226
5	A	14	9	1.00	33	0.273
6	A	21	14	1.00	33	0.424
7	A	16	12	1.00	33	0.364
8	A	10	9	1.00	31	0.290
9	A	18	13	1.00	31	0.419
10	A	21	14	1.00	33	0.424
11	A	20	12	1.00	33	0.364
12	A	14	11	1.00	33	0.333
13	A	10	7	1.00	33	0.212
14	A	7	5	1.00	31	0.161
15	A	13	10	1.00	31	0.323
16	A	18	12	1.00	33	0.364
17	A	8	7	1.00	35	0.200
18	A	7	6	1.00	35	0.171
19	A	4	4	1.00	33	0.121
20	A	8	5	1.00	33	0.152
21	A	11	6	1.00	35	0.171
22	A	9	8	1.00	35	0.229
23	A	8	7	1.00	35	0.200
24	A	8	7	1.00	33	0.212

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	10	7	1.00	33	0.212
26	A	22	9	1.00	35	0.257
27	A	8	7	1.00	35	0.200
28	A	6	6	1.00	35	0.171
29	A	6	6	1.00	35	0.171
30	A	6	6	1.00	33	0.182
31	A	12	7	1.00	33	0.212
32	A	16	8	1.00	35	0.229

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$	37
3.2	$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$	46
3.3	$\int \frac{\cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$	52
3.4	$\int \frac{\tan(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$	57
3.5	$\int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$	63
3.6	$\int \cot^5(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$	71
3.7	$\int \cot^3(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$	86
3.8	$\int \cot(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$	95
3.9	$\int \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan(d+ex) dx$	102
3.10	$\int \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan^3(d+ex) dx$	112
3.11	$\int \frac{\cot^7(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	121
3.12	$\int \frac{\cot^5(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	137
3.13	$\int \frac{\cot^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	148
3.14	$\int \frac{\cot(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	157
3.15	$\int \frac{\tan(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	164
3.16	$\int \frac{\tan^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$	172
3.17	$\int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$	186
3.18	$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$	193
3.19	$\int \frac{\cot(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$	199
3.20	$\int \frac{\tan(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$	204

3.21	$\int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$	209
3.22	$\int \cot^5(d+ex) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} dx$	216
3.23	$\int \cot^3(d+ex) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} dx$	226
3.24	$\int \cot(d+ex) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} dx$	234
3.25	$\int \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} \tan(d+ex) dx$	241
3.26	$\int \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} \tan^3(d+ex) dx$	249
3.27	$\int \frac{\cot^7(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	258
3.28	$\int \frac{\cot^5(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	264
3.29	$\int \frac{\cot^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	271
3.30	$\int \frac{\cot(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	278
3.31	$\int \frac{\tan(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	284
3.32	$\int \frac{\tan^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$	293

$$3.1 \quad \int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

Optimal result	37
Rubi [A] (verified)	38
Mathematica [C] (verified)	43
Maple [B] (warning: unable to verify)	44
Fricas [B] (verification not implemented)	44
Sympy [F]	44
Maxima [F]	44
Giac [F(-2)]	45
Mupad [F(-1)]	45

Optimal result

Integrand size = 33, antiderivative size = 547

$$\begin{aligned} & \int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx \\ &= -\frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\ &+ \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\ &- \frac{\operatorname{barctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{3/2}e} \\ &+ \frac{b(5b^2-12ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{7/2}e} \\ &+ \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{ce} \\ &- \frac{\cot^2(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{3ce} \\ &- \frac{(15b^2-16ac-10bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{24c^3e} \end{aligned}$$

[Out] $-1/2*b*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})/c^{(3/2)}/e+1/16*b*(-12*a*c+5*b^2)*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})/c^{(7/2)}/e+(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/c/e-1/3*\cot(e*x+d)^2*(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/c/e-1/24*(15*b^2-16*a*c-10*b*c*\cot(e*x+d))*(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/c^3/e-1/2*\operatorname{arctanh}(1/2*(a-c+b*\cot(e*x+d)-(a^2-2*a*c+b^2+c^2)^{(1/2)})$

$$\begin{aligned} & *2^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)/(a-c-(a^2-2*a*c+b^2+c^2)^{(1/2)})^{(1/2)}} \\ & *(a-c-(a^2-2*a*c+b^2+c^2)^{(1/2)})^{(1/2)}/e*2^{(1/2)/(a^2-2*a*c+b^2+c^2)^{(1/2)}} \\ & *(1/2)*\arctanh(1/2*(a-c+b*\cot(e*x+d)+(a^2-2*a*c+b^2+c^2)^{(1/2)})^{(1/2)}) \\ & *(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)/(a-c+(a^2-2*a*c+b^2+c^2)^{(1/2)})^{(1/2)}} \\ & *(a-c+(a^2-2*a*c+b^2+c^2)^{(1/2)})^{(1/2)}/e*2^{(1/2)/(a^2-2*a*c+b^2+c^2)^{(1/2)}} \end{aligned}$$

Rubi [A] (verified)

Time = 1.41 (sec) , antiderivative size = 547, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {3782, 6857, 654, 635, 212, 756, 793, 1050, 1044, 214}

$$\begin{aligned} & \int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \\ & \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-\operatorname{arctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}} \\ & + \frac{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-\operatorname{arctanh}\left(\frac{\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}} \\ & + \frac{b(5b^2-12ac)\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{16c^{7/2}e} \\ & - \frac{\operatorname{barctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2c^{3/2}e} \\ & - \frac{(-16ac+15b^2-10bc\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{24c^3e} \\ & - \frac{\cot^2(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{3ce} + \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{ce} \end{aligned}$$

[In] Int[Cot[d + e*x]^5/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] -((Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e)) + (Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e) - (b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(2*c^(3/2)*e) + (b*(5*b^2 - 12*a*c)*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(16*c^(7/2)*e) + Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]/(c*e) - (Cot[d + e*x]^2*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])/(c*e)

$\text{ot}[d + e*x] + c*\text{Cot}[d + e*x]^2]/(3*c*e) - ((15*b^2 - 16*a*c - 10*b*c*\text{Cot}[d + e*x])* \text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2])/(24*c^3*e)$

Rule 212

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] := \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 214

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] := \text{Simp}[(\text{Rt}[-a/b, 2]/a)* \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

Rule 635

$\text{Int}[1/\text{Sqrt}[(a_ + (b_)*(x_) + (c_)*(x_)^2)], x_Symbol] := \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 654

$\text{Int}(((d_ + (e_)*(x_))*((a_ + (b_)*(x_) + (c_)*(x_)^2)^{p_}), x_Symbol] := \text{Simp}[e*((a + b*x + c*x^2)^{(p + 1)}/(2*c*(p + 1))), x] + \text{Dist}[(2*c*d - b*e)/(2*c), \text{Int}[(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0] \ \&\& \ \text{NeQ}[p, -1]$

Rule 756

$\text{Int}(((d_ + (e_)*(x_))^{m_})*((a_ + (b_)*(x_) + (c_)*(x_)^2)^{p_}), x_Symbol] := \text{Simp}[e*(d + e*x)^{(m - 1)}*((a + b*x + c*x^2)^{(p + 1)}/(c*(m + 2*p + 1))), x] + \text{Dist}[1/(c*(m + 2*p + 1)), \text{Int}[(d + e*x)^{(m - 2)}*\text{Simp}[c*d^2*(m + 2*p + 1) - e*(a*e*(m - 1) + b*d*(p + 1)) + e*(2*c*d - b*e)*(m + p)*x, x]*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, p\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0] \ \&\& \ \text{If}[\text{RationalQ}[m], \text{GtQ}[m, 1], \text{SumSimplerQ}[m, -2]] \ \&\& \ \text{NeQ}[m + 2*p + 1, 0] \ \&\& \ \text{IntQuadraticQ}[a, b, c, d, e, m, p, x]$

Rule 793

$\text{Int}(((d_ + (e_)*(x_))*((f_ + (g_)*(x_))*((a_ + (b_)*(x_) + (c_)*(x_)^2)^{p_}), x_Symbol] := \text{Simp}[(-b*e*g*(p + 2) - c*(e*f + d*g)*(2*p + 3) - 2*c*e*g*(p + 1)*x)*((a + b*x + c*x^2)^{(p + 1)}/(2*c^2*(p + 1)*(2*p + 3))), x] + \text{Dist}[(b^2*e*g*(p + 2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p + 3))/(2*c^2*(2*p + 3)), \text{Int}[(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ !\text{LeQ}[p, -1]$

Rule 1044

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]
```

Rule 1050

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]
```

Rule 3782

```
Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n_)) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_)]^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rule 6857

```
Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \left(-\frac{x}{\sqrt{a+bx+cx^2}} + \frac{x^3}{\sqrt{a+bx+cx^2}} + \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}}\right) dx, x, \cot(d+ex)\right)}{e} \\ &= \frac{\text{Subst}\left(\int \frac{x}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} - \frac{\text{Subst}\left(\int \frac{x^3}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{\cot^2(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{ce}{3ce} \\
&\quad \text{Subst} \left(\int \frac{x \left(-2a - \frac{5bx}{2} \right)}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&\quad - \frac{3ce}{3ce} \\
&\quad b \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&\quad - \frac{2ce}{2ce} \\
&\quad \text{Subst} \left(\int \frac{-b + \left(a - c - \sqrt{a^2 + b^2 - 2ac + c^2} \right) x}{(1+x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&+ \frac{2\sqrt{a^2 + b^2 - 2ac + c^2} e}{2\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&\quad \text{Subst} \left(\int \frac{-b + \left(a - c + \sqrt{a^2 + b^2 - 2ac + c^2} \right) x}{(1+x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&\quad - \frac{2\sqrt{a^2 + b^2 - 2ac + c^2} e}{2\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&= \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{\cot^2(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{ce}{3ce} \\
&\quad (15b^2 - 16ac - 10bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \\
&\quad - \frac{24c^3 e}{24c^3 e} \\
&\quad b \text{Subst} \left(\int \frac{1}{4c - x^2} dx, x, \frac{b + 2c \cot(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \right) \\
&\quad - \frac{ce}{16c^3 e} \\
&\quad (b(5b^2 - 12ac)) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&+ \frac{(b(a - c - \sqrt{a^2 + b^2 - 2ac + c^2})) \text{Subst} \left(\int \frac{1}{-2b(a - c - \sqrt{a^2 + b^2 - 2ac + c^2}) + bx^2} dx, x, \frac{a - c - \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \right)}{\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&+ \frac{(b(a - c + \sqrt{a^2 + b^2 - 2ac + c^2})) \text{Subst} \left(\int \frac{1}{-2b(a - c + \sqrt{a^2 + b^2 - 2ac + c^2}) + bx^2} dx, x, \frac{a - c + \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \right)}{\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&\quad - \frac{\sqrt{a^2 + b^2 - 2ac + c^2} e}{\sqrt{a^2 + b^2 - 2ac + c^2} e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&+ \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&- \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{3/2}e} + \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{ce} \\
&- \frac{\cot^2(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{3ce} \\
&- \frac{(15b^2-16ac-10bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{24c^3e} \\
&+ \frac{(b(5b^2-12ac)) \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8c^3e} \\
&= \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&+ \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&- \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{3/2}e} \\
&+ \frac{b(5b^2-12ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{7/2}e} \\
&+ \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{ce} \\
&- \frac{\cot^2(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{3ce} \\
&- \frac{(15b^2-16ac-10bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{24c^3e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.77 (sec) , antiderivative size = 735, normalized size of antiderivative = 1.34

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

$$= \frac{\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{ce}$$

$$\frac{i \arctan\left(\frac{ib-2c+(2ia-b)\tan(d+ex)}{2\sqrt{a+ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{2\sqrt{a+ib-ce}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

$$\frac{i \arctan\left(\frac{ib+2c+(2ia+b)\tan(d+ex)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{2\sqrt{a-ib-ce}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

$$\frac{\operatorname{barctanh}\left(\frac{2c+b\tan(d+ex)}{2\sqrt{c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{2c^{3/2}e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

$$\frac{\cot(d+ex) \sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{48e\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}} \left(\frac{16\cot^3(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{c} - \frac{20b\cot^2(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{c} \right)$$

[In] Integrate[Cot[d + e*x]^5/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]/(c*e) - ((I/2) *ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Tan[d + e*x]*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))]/(Sqrt[a + I*b - c]*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - ((I/2)*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Tan[d + e*x]*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))]/(Sqrt[a - I*b - c]*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - (b*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Tan[d + e*x]*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))]/(2*c^(3/2)*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - (Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((16*Cot[d + e*x]^3*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/c - ((20*b*Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/c + ((3*b*(5*b^2 - 12*a*c)*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))])/c^(3/2) - (2*(15*b^2 - 16*a*c)*Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])/c)/c)/(48*e*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)])

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 3.55 (sec) , antiderivative size = 9581948, normalized size of antiderivative = 17517.27

output too large to display

[In] `int(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)`

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 12095 vs. 2(484) = 968.

Time = 6.09 (sec) , antiderivative size = 24241, normalized size of antiderivative = 44.32

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Too large to display}$$

[In] `integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

[In] `integrate(cot(e*x+d)**5/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)`

[Out] `Integral(cot(d + e*x)**5/sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2), x)`

Maxima [F]

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot^5(ex+d)}{\sqrt{c\cot^2(ex+d)+b\cot(ex+d)+a}} dx$$

[In] `integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")`

[Out] `integrate(cot(e*x + d)^5/sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a), x)`

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot(d+ex)^5}{\sqrt{c\cot(d+ex)^2+b\cot(d+ex)+a}} dx$$

[In] int(cot(d + e*x)^5/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(cot(d + e*x)^5/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

$$3.2 \quad \int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

Optimal result	46
Rubi [A] (verified)	47
Mathematica [C] (verified)	49
Maple [B] (warning: unable to verify)	50
Fricas [B] (verification not implemented)	50
Sympy [F]	50
Maxima [F]	51
Giac [F(-2)]	51
Mupad [F(-1)]	51

Optimal result

Integrand size = 33, antiderivative size = 384

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

$$= \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} - \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} + \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{3/2}e} - \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{ce}$$

[Out] 1/2*b*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/c^(3/2)/e-(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/c/e+1/2*arctanh(1/2*(a-c+b*cot(e*x+d)-(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)-1/2*arctanh(1/2*(a-c+b*cot(e*x+d)+(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)

Rubi [A] (verified)

Time = 1.04 (sec) , antiderivative size = 384, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.242$, Rules used = {3782, 6857, 654, 635, 212, 1050, 1044, 214}

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

$$= \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-\operatorname{arctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}}$$

$$- \frac{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-\operatorname{arctanh}\left(\frac{\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}}$$

$$+ \frac{\operatorname{barctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2c^{3/2}e} - \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{ce}$$

[In] Int[Cot[d + e*x]^3/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] (Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]])*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e) - (Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]])*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e) + (b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(2*c^(3/2)*e) - Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]/(c*e)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 635

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 654

```
Int[((d._) + (e._)*(x_))*((a._) + (b._)*(x_) + (c._)*(x_)^2)^(p_), x_Symbol]
:= Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[(2*c*d - b*e)/(2*c),
Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x]
&& NeQ[2*c*d - b*e, 0] && NeQ[p, -1]
```

Rule 1044

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]),
x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x,
Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x]
&& EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]
```

Rule 1050

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]),
x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q),
Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q),
Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x]
&& NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]
```

Rule 3782

```
Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x]
&& EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rule 6857

```
Int[(u_)/((a_) + (b_)*(x_)^n), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x]
&& IGtQ[n, 0]
```

Rubi steps

$$\text{integral} = \frac{\text{Subst}\left(\int \frac{x^3}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e}$$

$$= \frac{\text{Subst}\left(\int \left(\frac{x}{\sqrt{a+bx+cx^2}} - \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}}\right) dx, x, \cot(d+ex)\right)}{e}$$

$$\begin{aligned}
&= -\frac{\text{Subst}\left(\int \frac{x}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{ce} + \frac{b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2ce} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-b+(a-c-\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-b+(a-c+\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&= -\frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{ce} + \frac{b\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{ce} \\
&\quad - \frac{(b(a-c-\sqrt{a^2+b^2-2ac+c^2}))\text{Subst}\left(\int \frac{1}{-2b(a-c-\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c-\sqrt{a^2+b^2-2ac+c^2}}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{(b(a-c+\sqrt{a^2+b^2-2ac+c^2}))\text{Subst}\left(\int \frac{1}{-2b(a-c+\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c+\sqrt{a^2+b^2-2ac+c^2}}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&= \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\text{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad - \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\text{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{b\text{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2c^{3/2}e} - \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{ce}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.84 (sec) , antiderivative size = 397, normalized size of antiderivative = 1.03

$$\begin{aligned}
&\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx \\
&= \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} \tan(d+ex) \left(b\sqrt{a-ib-c\sqrt{a+ib}} - c\text{arctanh}\left(\frac{2c+b\tan(d+ex)}{2\sqrt{c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \right)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}
\end{aligned}$$

[In] Integrate[Cot[d + e*x]^3/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

```
[Out] (Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]*(b*Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + I*Sqrt[c]*(Sqrt[a - I*b - c]*c*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + Sqrt[a + I*b - c]*(c*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + (2*I)*Sqrt[a - I*b - c]*Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))/(2*Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*c^(3/2)*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])
```

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.74 (sec) , antiderivative size = 9581108, normalized size of antiderivative = 24950.80

output too large to display

```
[In] int(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

```
[Out] result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11834 vs. 2(339) = 678.

Time = 5.73 (sec) , antiderivative size = 23719, normalized size of antiderivative = 61.77

$$\int \frac{\cot^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \text{Too large to display}$$

```
[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] Too large to include
```

Sympy [F]

$$\int \frac{\cot^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \int \frac{\cot^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx$$

```
[In] integrate(cot(e*x+d)**3/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)
```

```
[Out] Integral(cot(d + e*x)**3/sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2), x)
```

Maxima [F]

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot(ex+d)^3}{\sqrt{c\cot(ex+d)^2+b\cot(ex+d)+a}} dx$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)^3/sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot(d+ex)^3}{\sqrt{c\cot(d+ex)^2+b\cot(d+ex)+a}} dx$$

[In] int(cot(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(cot(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

3.3 $\int \frac{\cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$

Optimal result	52
Rubi [A] (verified)	52
Mathematica [C] (verified)	54
Maple [B] (warning: unable to verify)	55
Fricas [B] (verification not implemented)	55
Sympy [F]	55
Maxima [F(-2)]	56
Giac [F(-2)]	56
Mupad [F(-1)]	56

Optimal result

Integrand size = 31, antiderivative size = 294

$$\int \frac{\cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

$$= -\frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}$$

$$+ \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(a-c+b*\cot(e*x+d)-(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^(1/2)/(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)+1/2*\operatorname{arctanh}(1/2*(a-c+b*\cot(e*x+d)+(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^(1/2)/(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used

= {3782, 1050, 1044, 214}

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

$$= \frac{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a} - \operatorname{carctanh}\left(\frac{\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}}$$

$$- \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a} - \operatorname{carctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}e\sqrt{a^2-2ac+b^2+c^2}}$$

[In] Int[Cot[d + e*x]/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] -((Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e)) + (Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e)

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 1044

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n2_)]^(p_), x_Symbol]

```

:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \frac{-b+(a-c-\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-b+(a-c+\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&= \frac{(b(a-c-\sqrt{a^2+b^2-2ac+c^2})) \text{Subst}\left(\int \frac{1}{-2b(a-c-\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad - \frac{(b(a-c+\sqrt{a^2+b^2-2ac+c^2})) \text{Subst}\left(\int \frac{1}{-2b(a-c+\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&= -\frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \text{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \text{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b\cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.61 (sec) , antiderivative size = 256, normalized size of antiderivative = 0.87

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \frac{i\left(\sqrt{a-ib-c} \arctan\left(\frac{ib-2c+(2ia-b)\tan(d+ex)}{2\sqrt{a+ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) + \sqrt{a+ib-c} \arctan\left(\frac{ib+2c+(2ia+b)\tan(d+ex)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right)\right)}{2\sqrt{a-ib-c}\sqrt{a+ib-c}e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

[In] Integrate[Cot[d + e*x]/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

```
[Out] ((-1/2*I)*(Sqrt[a - I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])]/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + Sqrt[a + I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])]/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x])/(Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])
```

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.04 (sec) , antiderivative size = 9339148, normalized size of antiderivative = 31765.81

output too large to display

```
[In] int(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5853 vs. $2(261) = 522$.

Time = 2.13 (sec) , antiderivative size = 5853, normalized size of antiderivative = 19.91

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Too large to display}$$

```
[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")
```

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

```
[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)
```

[Out] Integral(cot(d + e*x)/sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see 'assume?' for more details)

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\cot(d+ex)}{\sqrt{c\cot(d+ex)^2+b\cot(d+ex)+a}} dx$$

[In] int(cot(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(cot(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

$$3.4 \quad \int \frac{\tan(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

Optimal result	57
Rubi [A] (verified)	58
Mathematica [C] (verified)	60
Maple [F(-1)]	61
Fricas [B] (verification not implemented)	61
Sympy [F]	61
Maxima [F]	62
Giac [F(-2)]	62
Mupad [F(-1)]	62

Optimal result

Integrand size = 31, antiderivative size = 349

$$\int \frac{\tan(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

$$= \frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}}$$

$$+ \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}$$

$$- \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}$$

```
[Out] arctanh(1/2*(2*a+b*cot(e*x+d))/a^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/e/a^(1/2)+1/2*arctanh(1/2*(a-c+b*cot(e*x+d)-(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)-1/2*arctanh(1/2*(a-c+b*cot(e*x+d)+(a^2-2*a*c+b^2+c^2)^(1/2))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/e*2^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/2)
```

Rubi [A] (verified)

Time = 0.73 (sec) , antiderivative size = 349, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.226$, Rules used = {3782, 6857, 738, 212, 1050, 1044, 214}

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

$$= \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a} - \operatorname{arctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2e\sqrt{a^2-2ac+b^2+c^2}}}$$

$$- \frac{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a} - \operatorname{arctanh}\left(\frac{\sqrt{a^2-2ac+b^2+c^2}+a+b\cot(d+ex)-c}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+a-c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{2e\sqrt{a^2-2ac+b^2+c^2}}}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{\sqrt{ae}}$$

[In] Int[Tan[d + e*x]/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[a]*e) + (Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e) - (Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2] + b*Cot[d + e*x])/(Sqrt[2]*Sqrt[a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(Sqrt[2]*Sqrt[a^2 + b^2 - 2*a*c + c^2]*e)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 738

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 1044

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]
```

Rule 1050

```
Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]
```

Rule 3782

```
Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n_))^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rule 6857

```
Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x\sqrt{a+bx+cx^2}} - \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}}\right) dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \end{aligned}$$

$$\begin{aligned}
&= \frac{2 \operatorname{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&\quad - \frac{\operatorname{Subst}\left(\int \frac{-b+(a-c-\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{\operatorname{Subst}\left(\int \frac{-b+(a-c+\sqrt{a^2+b^2-2ac+c^2})x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&= \frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}} \\
&\quad - \frac{(b(a-c-\sqrt{a^2+b^2-2ac+c^2})) \operatorname{Subst}\left(\int \frac{1}{-2b(a-c-\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{(b(a-c+\sqrt{a^2+b^2-2ac+c^2})) \operatorname{Subst}\left(\int \frac{1}{-2b(a-c+\sqrt{a^2+b^2-2ac+c^2})+bx^2} dx, x, \frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{a^2+b^2-2ac+c^2}e} \\
&= \frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}} \\
&\quad + \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad - \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.74 (sec) , antiderivative size = 344, normalized size of antiderivative = 0.99

$$\begin{aligned}
&\int \frac{\tan(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx \\
&= \frac{\left(i\sqrt{a}\left(\sqrt{a-ib-c} \operatorname{arctan}\left(\frac{ib-2c+(2ia-b)\tan(d+ex)}{2\sqrt{a+ib-c}\sqrt{c+b \tan(d+ex)+a \tan^2(d+ex)}}\right)\right) + \sqrt{a+ib-c} \operatorname{arctan}\left(\frac{ib+2c+(2ia+b)\tan(d+ex)}{2\sqrt{a-ib-c}\sqrt{c+b \tan(d+ex)+a \tan^2(d+ex)}}\right)\right)}{2\sqrt{a}\sqrt{a-ib-c}\sqrt{a+ib-c}}
\end{aligned}$$

[In] Integrate[Tan[d + e*x]/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] ((I*Sqrt[a]*(Sqrt[a - I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])]/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + Sqrt[a + I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])]/(2*Sqrt[a -

```
I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + 2*Sqrt[a - I*b -
c]*Sqrt[a + I*b - c]*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*
Tan[d + e*x] + a*Tan[d + e*x]^2]))]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x
]^2]*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*e*Sqrt[c
+ b*Tan[d + e*x] + a*Tan[d + e*x]^2])
```

Maple **[F(-1)]**

Timed out.

hanged

```
[In] int(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

```
[Out] int(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

Fricas **[B]** (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5166 vs. 2(308) = 616.

Time = 1.43 (sec) , antiderivative size = 10333, normalized size of antiderivative = 29.61

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Too large to display}$$

```
[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fr
icas")
```

```
[Out] Too large to include
```

Sympy **[F]**

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx$$

```
[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)
```

```
[Out] Integral(tan(d + e*x)/sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2), x)
```

Maxima [F]

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\tan(ex+d)}{\sqrt{c\cot(ex+d)^2+b\cot(ex+d)+a}} dx$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tan(e*x + d)/sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} dx = \int \frac{\tan(d+ex)}{\sqrt{c\cot(d+ex)^2+b\cot(d+ex)+a}} dx$$

[In] int(tan(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(tan(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

$$3.5 \quad \int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx$$

Optimal result	63
Rubi [A] (verified)	64
Mathematica [C] (verified)	68
Maple [F(-1)]	69
Fricas [B] (verification not implemented)	69
Sympy [F]	69
Maxima [F]	70
Giac [F(-2)]	70
Mupad [F(-1)]	70

Optimal result

Integrand size = 33, antiderivative size = 501

$$\begin{aligned} & \int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx \\ &= -\frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}} \\ &+ \frac{(3b^2-4ac) \operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8a^{5/2}e} \\ &- \frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\ &+ \frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\ &- \frac{3b\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan(d+ex)}{4a^2e} \\ &+ \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan^2(d+ex)}{2ae} \end{aligned}$$

[Out] $\frac{1}{8}(-4ac+3b^2) \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b \cot(ex+d)}{a^{1/2} \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)}}\right) / a^{5/2} e - \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b \cot(ex+d)}{a^{1/2} \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)}}\right) / a^{5/2} e - \operatorname{arctanh}\left(\frac{1}{2} \frac{2a+b \cot(ex+d)}{a^{1/2} \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)}}\right) / a^{5/2} e - \frac{1}{2} \operatorname{arctanh}\left(\frac{1}{2} \frac{a-c+b \cot(ex+d)}{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}} \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)}}\right) - \frac{1}{2} \operatorname{arctanh}\left(\frac{1}{2} \frac{a-c+b \cot(ex+d)}{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}} \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)}}\right) + \frac{3b \sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)} \tan(ex+d)}{4a^2e} + \frac{\sqrt{a+b \cot(ex+d)+c \cot^2(ex+d)} \tan^2(ex+d)}{2ae}$

$$\frac{2^{1/2}}{(a^2 - 2ac + b^2 + c^2)^{1/2}} - \frac{3}{4} \frac{b(a + b \cot(ex + d) + c \cot(ex + d)^2)^{1/2} \tan(ex + d)}{a^2/e + \frac{1}{2}(a + b \cot(ex + d) + c \cot(ex + d)^2)^{1/2} \tan(ex + d)^2/a/e}$$

Rubi [A] (verified)

Time = 0.92 (sec) , antiderivative size = 501, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3782, 6857, 758, 820, 738, 212, 1050, 1044, 214}

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx$$

$$= \frac{(3b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{8a^{5/2}e}$$

$$- \frac{\sqrt{-\sqrt{a^2 - 2ac + b^2 + c^2} + a} \operatorname{carctanh}\left(\frac{-\sqrt{a^2 - 2ac + b^2 + c^2} + a + b \cot(d + ex) - c}{\sqrt{2}\sqrt{-\sqrt{a^2 - 2ac + b^2 + c^2} + a} - c\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{2}e\sqrt{a^2 - 2ac + b^2 + c^2}}$$

$$+ \frac{\sqrt{\sqrt{a^2 - 2ac + b^2 + c^2} + a} \operatorname{carctanh}\left(\frac{\sqrt{a^2 - 2ac + b^2 + c^2} + a + b \cot(d + ex) - c}{\sqrt{2}\sqrt{\sqrt{a^2 - 2ac + b^2 + c^2} + a} - c\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{2}e\sqrt{a^2 - 2ac + b^2 + c^2}}$$

$$- \frac{3b \tan(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{4a^2e}$$

$$- \frac{\operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{ae}}$$

$$+ \frac{\tan^2(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{2ae}$$

[In] Int[Tan[d + e*x]^3/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] $-(\operatorname{ArcTanh}[(2a + b \cot(d + ex))/(2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)})]) / (\sqrt{a}e) + ((3b^2 - 4ac) \operatorname{ArcTanh}[(2a + b \cot(d + ex))/(2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)})]) / (8a^{5/2}e) - (\sqrt{a - c - \sqrt{a^2 + b^2 - 2ac + c^2}}) \operatorname{ArcTanh}[(a - c - \sqrt{a^2 + b^2 - 2ac + c^2} + b \cot(d + ex)) / (\sqrt{2}\sqrt{a - c - \sqrt{a^2 + b^2 - 2ac + c^2}}) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)})] / (\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}e) + (\sqrt{a - c + \sqrt{a^2 + b^2 - 2ac + c^2}}) \operatorname{ArcTanh}[(a - c + \sqrt{a^2 + b^2 - 2ac + c^2} + b \cot(d + ex)) / (\sqrt{2}\sqrt{a - c + \sqrt{a^2 + b^2 - 2ac + c^2}}) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)})] / (\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}e) - (3b \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex)) / (4a^2e) + (\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex)^2) / (2ae)$

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 758

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*Simp[c*d*(m + 1) - b*e*(m + p + 2) - c*e*(m + 2*p + 3)*x, x]*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && NeQ[m, -1] && ((LtQ[m, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]) || (SumSimplerQ[m, 1] && IntegerQ[p]) || ILtQ[Simplify[m + 2*p + 3], 0])

Rule 820

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(-(e*f - d*g))*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/(2*(p + 1)*(c*d^2 - b*d*e + a*e^2))), x] - Dist[(b*(e*f + d*g) - 2*(c*d*f + a*e*g))/(2*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 1044

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

```

Int[((g_.) + (h_.)*(x_))/((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (
f_.)*(x_)^2]], x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist
[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*
e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Si
mp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c
*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] &&
NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

```

Rule 3782

```

Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rule 6857

```

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \frac{1}{x^3(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \left(\frac{1}{x^3\sqrt{a+bx+cx^2}} - \frac{1}{x\sqrt{a+bx+cx^2}} + \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}}\right) dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \frac{1}{x^3\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{2ae} \\
&\quad - \frac{2 \operatorname{Subst}\left(\int \frac{1}{4a - x^2} dx, x, \frac{2a + b \cot(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{e} \\
&\quad + \frac{\operatorname{Subst}\left(\int \frac{\frac{3b}{2} + cx}{x^2 \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex)\right)}{2ae} \\
&\quad + \frac{\operatorname{Subst}\left(\int \frac{-b + (a - c - \sqrt{a^2 + b^2 - 2ac + c^2})x}{(1 + x^2)\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&\quad - \frac{\operatorname{Subst}\left(\int \frac{-b + (a - c + \sqrt{a^2 + b^2 - 2ac + c^2})x}{(1 + x^2)\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&= - \frac{\operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{ae}} \\
&\quad - \frac{3b\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex)}{4a^2e} \\
&\quad + \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{2ae} \\
&\quad - \frac{(3b^2 - 4ac) \operatorname{Subst}\left(\int \frac{1}{x\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex)\right)}{8a^2e} \\
&\quad + \frac{(b(a - c - \sqrt{a^2 + b^2 - 2ac + c^2})) \operatorname{Subst}\left(\int \frac{1}{-2b(a - c - \sqrt{a^2 + b^2 - 2ac + c^2}) + bx^2} dx, x, \frac{a - c - \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&\quad - \frac{(b(a - c + \sqrt{a^2 + b^2 - 2ac + c^2})) \operatorname{Subst}\left(\int \frac{1}{-2b(a - c + \sqrt{a^2 + b^2 - 2ac + c^2}) + bx^2} dx, x, \frac{a - c + \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{\sqrt{a^2 + b^2 - 2ac + c^2}e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}} \\
&\quad -\frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad +\frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad -\frac{3b\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}\tan(d+ex)}{4a^2e} \\
&\quad +\frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}\tan^2(d+ex)}{2ae} \\
&\quad +\frac{(3b^2-4ac)\operatorname{Subst}\left(\int\frac{1}{4a-x^2}dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{4a^2e} \\
&= -\frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{ae}} \\
&\quad +\frac{(3b^2-4ac)\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8a^{5/2}e} \\
&\quad -\frac{\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{a-c-\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad +\frac{\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{a-c+\sqrt{a^2+b^2-2ac+c^2}+b \cot(d+ex)}{\sqrt{2}\sqrt{a-c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad -\frac{3b\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}\tan(d+ex)}{4a^2e} \\
&\quad +\frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}\tan^2(d+ex)}{2ae}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.50 (sec) , antiderivative size = 422, normalized size of antiderivative = 0.84

$$\int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} dx = \frac{\cot(d+ex)\sqrt{c+b \tan(d+ex)+a \tan^2(d+ex)}\left(\sqrt{a-ib-c}\sqrt{a+ib-c}(8a^2-3b^2+4ac)\operatorname{arctanh}\left(\frac{\sqrt{a-ib-c}\sqrt{a+ib-c}}{2\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)\right)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

[In] Integrate[Tan[d + e*x]^3/Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

```
[Out] -1/8*(Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*(Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*(8*a^2 - 3*b^2 + 4*a*c)*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + 2*Sqrt[a]*(2*I)*a^2*Sqrt[a - I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + Sqrt[a + I*b - c]*((2*I)*a^2*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + Sqrt[a - I*b - c]*(3*b - 2*a*Tan[d + e*x])*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))/(a^(5/2)*Sqrt[a - I*b - c]*Sqrt[a + I*b - c]*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])
```

Maple **[F(-1)]**

Timed out.

hanged

```
[In] int(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

```
[Out] int(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)
```

Fricas **[B]** (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5247 vs. 2(442) = 884.

Time = 1.63 (sec) , antiderivative size = 10495, normalized size of antiderivative = 20.95

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \text{Too large to display}$$

```
[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")
```

```
[Out] Too large to include
```

Sympy **[F]**

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx$$

```
[In] integrate(tan(e*x+d)**3/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)
```

```
[Out] Integral(tan(d + e*x)**3/sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2), x)
```

Maxima [F]

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \int \frac{\tan^3(ex + d)}{\sqrt{c \cot^2(ex + d) + b \cot(ex + d) + a}} dx$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tan(e*x + d)^3/sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} dx = \int \frac{\tan^3(d + ex)}{\sqrt{c \cot^2(d + ex) + b \cot(d + ex) + a}} dx$$

[In] int(tan(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(tan(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

3.6 $\int \cot^5(d+ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} dx$

Optimal result	72
Rubi [A] (verified)	73
Mathematica [C] (verified)	82
Maple [B] (warning: unable to verify)	83
Fricas [B] (verification not implemented)	84
Sympy [F]	84
Maxima [F]	84
Giac [F]	85
Mupad [F(-1)]	85

Optimal result

Integrand size = 33, antiderivative size = 976

$$\begin{aligned}
 & \int \cot^5(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx = \\
 & \frac{\sqrt{a^2+b^2+c}(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2}) \arctan\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}e} \\
 & - \frac{b \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
 & + \frac{b(b^2-4ac) \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{16c^{5/2}e} \\
 & - \frac{b(7b^2-12ac)(b^2-4ac) \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{256c^{9/2}e} \\
 & + \frac{\sqrt{a^2+b^2+c}(c-\sqrt{a^2+b^2-2ac+c^2})-a(2c-\sqrt{a^2+b^2-2ac+c^2}) \operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}e} \\
 & - \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} \\
 & - \frac{b(b+2c\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{8c^2e} \\
 & + \frac{b(7b^2-12ac)(b+2c\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{128c^4e} \\
 & + \frac{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}}{3ce} \\
 & - \frac{\cot^2(d+ex)(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}}{5ce} \\
 & - \frac{(35b^2-32ac-42bc\cot(d+ex))(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}}{240c^3e}
 \end{aligned}$$

[Out] 1/16*b*(-4*a*c+b^2)*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/c^(5/2)/e-1/256*b*(-12*a*c+7*b^2)*(-4*a*c+b^2)*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/c^(9/2)/e+1/3*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2)/c/e-1/5*cot(e*x+d)^2*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2)/c/e-1/240*(35*b^2-32*a*c-42*b*c*cot(e*x+d))*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2)/c^3/e-1/2*b*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/e/c^(1/2)-(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/e-1/8*b*(b+2*c*cot(e*x+d))*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/c^2/e+1/128*b*(-12*a*c+7*b^2)*(b+2*c*cot(e*x+d))*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/c^2/e

$$\begin{aligned}
& t(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/c^4/e+1/2*\operatorname{arctanh}(1/2*(b^2+b*\cot(e*x+d)*(a^2 \\
& -2*a*c+b^2+c^2)^{(1/2)}+(a-c)*(a-c+(a^2-2*a*c+b^2+c^2)^{(1/2)})))/(a^2-2*a*c+b^2 \\
& +c^2)^{(1/4)}*2^{(1/2)}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/(a^2+b^2+c*(c-(a^2 \\
& -2*a*c+b^2+c^2)^{(1/2)})-a*(2*c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)}*(a^2+b^2+ \\
& c*(c-(a^2-2*a*c+b^2+c^2)^{(1/2)})-a*(2*c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)}/(a \\
& ^2-2*a*c+b^2+c^2)^{(1/4)}/e*2^{(1/2)}-1/2*\operatorname{arctan}(1/2*(b^2+(a-c)*(a-c-(a^2-2*a*c \\
& +b^2+c^2)^{(1/2)})-b*\cot(e*x+d)*(a^2-2*a*c+b^2+c^2)^{(1/2)}))/(a^2-2*a*c+b^2+c^2 \\
&)^{(1/4)}*2^{(1/2)}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/(a^2+b^2+c*(c+(a^2-2* \\
& a*c+b^2+c^2)^{(1/2)})-a*(2*c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)}*(a^2+b^2+c*(c \\
& +(a^2-2*a*c+b^2+c^2)^{(1/2)})-a*(2*c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)}/(a^2-2 \\
& *a*c+b^2+c^2)^{(1/4)}/e*2^{(1/2)}
\end{aligned}$$

Rubi [A] (verified)

Time = 25.18 (sec) , antiderivative size = 976, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules

used = {3782, 6857, 654, 626, 635, 212, 756, 793, 1035, 1092, 1050, 1044, 214, 211}

$$\begin{aligned}
 & \int \cot^5(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx \\
 = & -\frac{(c\cot^2(d+ex)+b\cot(d+ex)+a)^{3/2} \cot^2(d+ex)}{5ce} \\
 & -\frac{(35b^2-42c\cot(d+ex)b-32ac)(c\cot^2(d+ex)+b\cot(d+ex)+a)^{3/2}}{240c^3e} \\
 & +\frac{(c\cot^2(d+ex)+b\cot(d+ex)+a)^{3/2}}{3ce} \\
 & -\frac{\sqrt{a^2-(2c+\sqrt{a^2-2ca+b^2+c^2})a+b^2+c}(c+\sqrt{a^2-2ca+b^2+c^2}) \arctan\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2-2ca+b^2+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2-2ca+b^2+c^2}e} \\
 & -\frac{b(7b^2-12ac)(b^2-4ac) \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{256c^{9/2}e} \\
 & +\frac{b(b^2-4ac) \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{16c^{5/2}e} \\
 & -\frac{b \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{2\sqrt{ce}} \\
 & +\frac{\sqrt{a^2-(2c-\sqrt{a^2-2ca+b^2+c^2})a+b^2+c}(c-\sqrt{a^2-2ca+b^2+c^2}) \operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2-2ca+b^2+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2-2ca+b^2+c^2}e} \\
 & +\frac{b(7b^2-12ac)(b+2c\cot(d+ex))\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{128c^4e} \\
 & -\frac{b(b+2c\cot(d+ex))\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{8c^2e} \\
 & -\frac{\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{e}
 \end{aligned}$$

[In] Int[Cot[d + e*x]^5*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] -((Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*ArcTan[(b^2 + (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*Sqrt[a^2 + b^2 - 2*a*c + c^2]*Cot[d + e*x])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e)) - (b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(2*Sqrt[c]*e) + (b*(b^2 - 4*a*c)*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(16*c^(5/2)*e) -

$$\begin{aligned} & (b*(7*b^2 - 12*a*c)*(b^2 - 4*a*c)*\text{ArcTanh}[(b + 2*c*\text{Cot}[d + e*x])/(2*\text{Sqrt}[c \\ & * \text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2])])/(256*c^{(9/2)*e}) + (\text{Sqrt}[a^2 \\ & + b^2 + c*(c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - \text{Sqrt}[a^2 + b^2 - \\ & 2*a*c + c^2])] * \text{ArcTanh}[(b^2 + (a - c)*(a - c + \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2 \\ &]) + b*\text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]*\text{Cot}[d + e*x])/(\text{Sqrt}[2]*(a^2 + b^2 - 2* \\ & a*c + c^2)^{(1/4)}*\text{Sqrt}[a^2 + b^2 + c*(c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]) - a \\ & *(2*c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2])] * \text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + \\ & e*x]^2])])/(\text{Sqrt}[2]*(a^2 + b^2 - 2*a*c + c^2)^{(1/4)*e}) - \text{Sqrt}[a + b*\text{Cot}[d \\ & + e*x] + c*\text{Cot}[d + e*x]^2]/e - (b*(b + 2*c*\text{Cot}[d + e*x])* \text{Sqrt}[a + b*\text{Cot}[d + \\ & e*x] + c*\text{Cot}[d + e*x]^2])/(8*c^2*e) + (b*(7*b^2 - 12*a*c)*(b + 2*c*\text{Cot}[d + \\ & e*x])* \text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2])/(128*c^4*e) + (a + b*\text{Co \\ & t}[d + e*x] + c*\text{Cot}[d + e*x]^2)^{(3/2)}/(3*c*e) - (\text{Cot}[d + e*x]^2*(a + b*\text{Cot}[d \\ & + e*x] + c*\text{Cot}[d + e*x]^2)^{(3/2)})/(5*c*e) - ((35*b^2 - 32*a*c - 42*b*c*\text{Cot} \\ & [d + e*x])*(a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2)^{(3/2)})/(240*c^3*e) \end{aligned}$$
Rule 211

$$\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{PosQ}[a/b]$$
Rule 212

$$\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{LtQ}[b, 0])$$
Rule 214

$$\text{Int}[\{(a_)+(b_)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x\} \&\& \text{NegQ}[a/b]$$
Rule 626

$$\text{Int}[\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(b + 2*c*x) * \{(a + b*x + c*x^2)^p/(2*c*(2*p + 1))\}, x] - \text{Dist}[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), \text{Int}[(a + b*x + c*x^2)^{(p - 1)}, x], x] /; \text{FreeQ}\{a, b, c\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{GtQ}[p, 0] \&\& \text{IntegerQ}[4*p]$$
Rule 635

$$\text{Int}[1/\text{Sqrt}[(a_)+(b_)*(x_)+(c_)*(x_)^2], x_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0]$$
Rule 654

$$\text{Int}[\{(d_)+(e_)*(x_)*\{(a_)+(b_)*(x_)+(c_)*(x_)^2\}^{(p_)}\}, x_Symbol] \rightarrow \text{Simp}[e*\{(a + b*x + c*x^2)^{(p + 1)}/(2*c*(p + 1))\}, x] + \text{Dist}[(2*c*d - b$$

$e)/(2c), \text{Int}[(a + b*x + c*x^2)^p, x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x\}$
 $\&\& \text{NeQ}[2*c*d - b*e, 0] \&\& \text{NeQ}[p, -1]$

Rule 756

$\text{Int}[(d + e*x)^m * (a + b*x + c*x^2)^p, x_Symbol] :> \text{Simp}[e*(d + e*x)^{m-1} * (a + b*x + c*x^2)^{p+1} / (c*(m + 2*p + 1)), x] + \text{Dist}[1/(c*(m + 2*p + 1)), \text{Int}[(d + e*x)^{m-2} * \text{Simp}[c*d^2*(m + 2*p + 1) - e*(a*e*(m-1) + b*d*(p+1)) + e*(2*c*d - b*e)*(m+p)*x, x] * (a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, m, p\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NeQ}[2*c*d - b*e, 0] \&\& \text{If}[\text{RationalQ}[m], \text{GtQ}[m, 1], \text{SumSimplerQ}[m, -2]] \&\& \text{NeQ}[m + 2*p + 1, 0] \&\& \text{IntQuadraticQ}[a, b, c, d, e, m, p, x]$

Rule 793

$\text{Int}[(d + e*x)*(f + g*x)*(a + b*x + c*x^2)^p, x_Symbol] :> \text{Simp}[(-b*e*g*(p+2) - c*(e*f + d*g)*(2*p+3) - 2*c*e*g*(p+1)*x) * (a + b*x + c*x^2)^{p+1} / (2*c^2*(p+1)*(2*p+3)), x] + \text{Dist}[(b^2*e*g*(p+2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p+3)) / (2*c^2*(2*p+3)), \text{Int}[(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, p\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{!LeQ}[p, -1]$

Rule 1035

$\text{Int}[(g + h*x)*(a + b*x + c*x^2)^p * (d + f*x^2)^q, x_Symbol] :> \text{Simp}[h*(a + b*x + c*x^2)^p * (d + f*x^2)^{q+1} / (2*f*(p+q+1)), x] - \text{Dist}[1/(2*f*(p+q+1)), \text{Int}[(a + b*x + c*x^2)^{p-1} * (d + f*x^2)^q * \text{Simp}[h*p*(b*d) + a*(-2*g*f)*(p+q+1) + (2*h*p*(c*d - a*f) + b*(-2*g*f)*(p+q+1))*x + (h*p*(-b)*f) + c*(-2*g*f)*(p+q+1))*x^2, x], x] /; \text{FreeQ}\{a, b, c, d, f, g, h, q\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{GtQ}[p, 0] \&\& \text{NeQ}[p+q+1, 0]$

Rule 1044

$\text{Int}[(g + h*x) / ((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x_Symbol] :> \text{Dist}[-2*a*g*h, \text{Subst}[\text{Int}[1/\text{Simp}[2*a^2*g*h*c + a*e*x^2, x], x], x, \text{Simp}[a*h - g*c*x, x] / \text{Sqrt}[d + e*x + f*x^2]], x] /; \text{FreeQ}\{a, c, d, e, f, g, h\}, x\} \&\& \text{EqQ}[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]$

Rule 1050

$\text{Int}[(g + h*x) / ((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x_Symbol] :> \text{With}\{q = \text{Rt}[(c*d - a*f)^2 + a*c*e^2, 2]\}, \text{Dist}[1/(2*q), \text{Int}[\text{Simp}[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x] / ((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x], x] - \text{Dist}[1/(2*q), \text{Int}[\text{Si}$

```
mp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c
*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] &&
NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]
```

Rule 1092

```
Int[((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)/(((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) +
(e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := Dist[C/c, Int[1/Sqrt[d + e*x + f
*x^2], x], x] + Dist[1/c, Int[(A*c - a*C + B*c*x)/((a + c*x^2)*Sqrt[d + e*x
+ f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, A, B, C}, x] && NeQ[e^2 - 4*d*f
, 0]
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \left(-x\sqrt{a+bx+cx^2} + x^3\sqrt{a+bx+cx^2} + \frac{x\sqrt{a+bx+cx^2}}{1+x^2}\right) dx, x, \cot(d+ex)\right)}{e} \\
 &= \frac{\text{Subst}\left(\int x\sqrt{a+bx+cx^2} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int x^3\sqrt{a+bx+cx^2} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} + \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&\quad - \frac{\cot^2(d + ex) (a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{5ce} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{b}{2} - (a-c)x - \frac{bx^2}{2}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{e} \\
&\quad - \frac{\text{Subst}\left(\int x\left(-2a - \frac{7bx}{2}\right) \sqrt{a + bx + cx^2} dx, x, \cot(d + ex)\right)}{5ce} \\
&\quad - \frac{b \text{Subst}\left(\int \sqrt{a + bx + cx^2} dx, x, \cot(d + ex)\right)}{2ce} \\
&= -\frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&\quad - \frac{b(b + 2c \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2e} \\
&\quad + \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&\quad - \frac{\cot^2(d + ex) (a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{5ce} \\
&\quad - \frac{(35b^2 - 32ac - 42bc \cot(d + ex)) (a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{240c^3e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{b + (-a+c)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{e} \\
&\quad - \frac{b \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2e} \\
&\quad + \frac{(b(7b^2 - 12ac)) \text{Subst}\left(\int \sqrt{a + bx + cx^2} dx, x, \cot(d + ex)\right)}{32c^3e} \\
&\quad + \frac{(b(b^2 - 4ac)) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{16c^2e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&\quad - \frac{b(b + 2c \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2e} \\
&\quad + \frac{b(7b^2 - 12ac) (b + 2c \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{128c^4e} \\
&\quad + \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&\quad - \frac{\cot^2(d + ex) (a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{5ce} \\
&\quad - \frac{(35b^2 - 32ac - 42bc \cot(d + ex)) (a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{240c^3e} \\
&\quad - \frac{b \text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&\quad + \frac{(b(b^2 - 4ac)) \text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8c^2e} \\
&\quad - \frac{(b(7b^2 - 12ac) (b^2 - 4ac)) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{256c^4e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{\operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
&+ \frac{b(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{5/2}e} \\
&- \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&- \frac{b(b + 2c \cot(d + ex))\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2e} \\
&+ \frac{b(7b^2 - 12ac)(b + 2c \cot(d + ex))\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{128c^4e} \\
&+ \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&- \frac{\cot^2(d + ex)(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{5ce} \\
&- \frac{(35b^2 - 32ac - 42bc \cot(d + ex))(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{240c^3e} \\
&- \frac{(b(7b^2 - 12ac)(b^2 - 4ac)) \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{128c^4e} \\
&+ \frac{(b(b^2 + (a - c)(a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \operatorname{Subst}\left(\int \frac{1}{2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&+ \frac{(b(b^2 + (a - c)(a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \operatorname{Subst}\left(\int \frac{1}{-2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e}
\end{aligned}$$

$$\begin{aligned}
&= \\
&\frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})} \arctan\left(\frac{\sqrt{2}\sqrt[4]{a^2 + b^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac}} \\
&- \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
&+ \frac{b(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{5/2}e} \\
&- \frac{b(7b^2 - 12ac)(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{256c^{9/2}e} \\
&+ \frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a^2 + b^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac}} \\
&- \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&- \frac{b(b + 2c \cot(d + ex))\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2e} \\
&+ \frac{b(7b^2 - 12ac)(b + 2c \cot(d + ex))\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{128c^4e} \\
&+ \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&- \frac{\cot^2(d + ex)(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{5ce} \\
&- \frac{(35b^2 - 32ac - 42bc \cot(d + ex))(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{240c^3e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.71 (sec) , antiderivative size = 1214, normalized size of antiderivative = 1.24

$$\int \cot^5(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx =$$

$$\frac{i\sqrt{a+ib-c} \arctan\left(\frac{ib-2c+(2ia-b)\tan(d+ex)}{2\sqrt{a+ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{2e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} +$$

$$\frac{i\sqrt{a-ib-c} \arctan\left(\frac{ib+2c+(2ia+b)\tan(d+ex)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{2e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} +$$

$$\frac{\sqrt{a} \operatorname{arctanh}\left(\frac{b+2a\tan(d+ex)}{2\sqrt{a}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) \tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} +$$

$$\frac{\tan(d+ex) \sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))} \left(2\sqrt{a} \operatorname{arctanh}\left(\frac{b+2a\tan(d+ex)}{2\sqrt{a}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right) + \frac{16 \cot^3(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))^{3/2}}{c} + \frac{48e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{5c} - \frac{\cot^5(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))^{3/2}}{5c}\right)}{48e\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} +$$

[In] Integrate[Cot[d + e*x]^5*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] ((-1/2*I)*Sqrt[a + I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x]]/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))*Tan[d + e*x]*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]/(e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)) - ((I/2)*Sqrt[a - I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x]]/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))*Tan[d + e*x]*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]/(e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)) +

$$\begin{aligned} & n[d + e*x] + a*\text{Tan}[d + e*x]^2])*\text{Tan}[d + e*x]*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + e*x] \\ & + a*\text{Tan}[d + e*x]^2))]/(e*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)) - (\text{Sqrt}[a]*\text{ArcTanh}[(b + 2*a*\text{Tan}[d + e*x])/(2*\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d \\ & + e*x] + a*\text{Tan}[d + e*x]^2))]*\text{Tan}[d + e*x]*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d \\ & + e*x] + a*\text{Tan}[d + e*x]^2))]/(e*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2 \\ &]) + (\text{Tan}[d + e*x]*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)]*(2*\text{Sqrt}[a]*\text{ArcTanh}[(b + 2*a*\text{Tan}[d + e*x])/(2*\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d \\ & + e*x] + a*\text{Tan}[d + e*x]^2))]) - (b*\text{ArcTanh}[(2*c + b*\text{Tan}[d + e*x])/(2*\text{Sqrt}[c] \\ & *\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))])/(\text{Sqrt}[c] - 2*\text{Cot}[d + e*x]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)))/(2*e*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + \\ & a*\text{Tan}[d + e*x]^2)) + (\text{Tan}[d + e*x]*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)]*((16*\text{Cot}[d + e*x]^3*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)^(3/2))/c + (3*b*((b^2 - 4*a*c)*\text{ArcTanh}[(2*c + b*\text{Tan}[d + e*x])/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))])/c^(3/2) - (2*\text{Cot}[d + e*x]^2*(2*c + b*\text{Tan}[d + e*x])*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))/c))/c))/(48*e*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)) + (\text{Tan}[d + e*x]*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)]*(-1/5*(\text{Cot}[d + e*x]^5*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)^(3/2))/c - ((-7*b*\text{Cot}[d + e*x]^4*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)^(3/2))/(8*c) - (((-35*b^2 + 32*a*c)*\text{Cot}[d + e*x]^3*(c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2)^(3/2))/(12*c) - ((-7*a*b*c + (b*(35*b^2 - 32*a*c))/4)*((b^2 - 4*a*c)*\text{ArcTanh}[(2*c + b*\text{Tan}[d + e*x])/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))])/((8*c^(3/2)) + (\text{Cot}[d + e*x]^2*(-2*c - b*\text{Tan}[d + e*x])*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))/(4*c)))/(2*c))/(4*c))/(5*c)))/(e*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2]) \end{aligned}$$

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 6.86 (sec) , antiderivative size = 17768513, normalized size of antiderivative = 18205.44

output too large to display

[In] $\text{int}(\text{cot}(e*x+d)^5*(a+b*\text{cot}(e*x+d)+c*\text{cot}(e*x+d)^2)^(1/2), x)$

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6335 vs. $2(877) = 1754$.

Time = 3.02 (sec) , antiderivative size = 12721, normalized size of antiderivative = 13.03

$$\int \cot^5(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\begin{aligned} & \int \cot^5(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx \\ &= \int \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}\cot^5(d+ex)dx \end{aligned}$$

[In] integrate(cot(e*x+d)**5*(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)

[Out] Integral(sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2)*cot(d + e*x)**5, x)

Maxima [F]

$$\begin{aligned} & \int \cot^5(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx \\ &= \int \sqrt{c\cot(ex+d)^2+b\cot(ex+d)+a}\cot(ex+d)^5dx \end{aligned}$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d)^5, x)

Giac [F]

$$\int \cot^5(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx$$

$$= \int \sqrt{c\cot^2(ex+d)+b\cot(ex+d)+a} \cot^5(ex+d) dx$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d)^5, x)

Mupad [F(-1)]

Timed out.

$$\int \cot^5(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx = \text{Hanged}$$

[In] int(cot(d + e*x)^5*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] \text{Hanged}

3.7 $\int \cot^3(d+ex) \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)} dx$

Optimal result	86
Rubi [A] (verified)	87
Mathematica [C] (verified)	92
Maple [B] (warning: unable to verify)	93
Fricas [B] (verification not implemented)	93
Sympy [F]	93
Maxima [F]	94
Giac [F]	94
Mupad [F(-1)]	94

Optimal result

Integrand size = 33, antiderivative size = 747

$$\begin{aligned}
 & \int \cot^3(d+ex) \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)} dx \\
 = & \frac{\sqrt{a^2 + b^2 + c} (c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2}) \arctan\left(\frac{b^2 + \sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2}}{\sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2} e} \\
 & + \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c} \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{c} e} \\
 & - \frac{b(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c} \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{5/2} e} \\
 & - \frac{\sqrt{a^2 + b^2 + c} (c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c - \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{\sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2}}{\sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2} \sqrt[4]{a^2 + b^2 - 2ac + c^2} e} \\
 & + \frac{\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}}{e} \\
 & + \frac{b(b + 2c \cot(d+ex)) \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}}{8c^2 e} \\
 & - \frac{(a + b \cot(d+ex) + c \cot^2(d+ex))^{3/2}}{3ce}
 \end{aligned}$$

[Out] $-1/16*b*(-4*a*c+b^2)*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})/c^{(5/2)}/e-1/3*(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(3/2)}/c/e+1/2*b*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})/e/c^{(1/2)}+(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/e+1/8*b*(b+2*$

$$\begin{aligned}
& c \cot(ex+d) * (a+b \cot(ex+d)+c \cot(ex+d)^2)^{(1/2)} / c^2 / e - 1/2 * \operatorname{arctanh}(1/2 * (b^2+b \cot(ex+d) * (a^2-2*a*c+b^2+c^2)^{(1/2)}+(a-c) * (a-c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))) / (a^2-2*a*c+b^2+c^2)^{(1/4)} * 2^{(1/2)} / (a+b \cot(ex+d)+c \cot(ex+d)^2)^{(1/2)} / (a^2+b^2+c * (c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))-a * (2*c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)} * (a^2+b^2+c * (c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))-a * (2*c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)} / (a^2-2*a*c+b^2+c^2)^{(1/4)} / e * 2^{(1/2)} + 1/2 * \operatorname{arctan}(1/2 * (b^2+(a-c) * (a-c-(a^2-2*a*c+b^2+c^2)^{(1/2)}))-b \cot(ex+d) * (a^2-2*a*c+b^2+c^2)^{(1/2)})) / (a^2-2*a*c+b^2+c^2)^{(1/4)} * 2^{(1/2)} / (a+b \cot(ex+d)+c \cot(ex+d)^2)^{(1/2)} / (a^2+b^2+c * (c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))-a * (2*c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)} * (a^2+b^2+c * (c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))-a * (2*c+(a^2-2*a*c+b^2+c^2)^{(1/2)}))^{(1/2)} / (a^2-2*a*c+b^2+c^2)^{(1/4)} / e * 2^{(1/2)}
\end{aligned}$$

Rubi [A] (verified)

Time = 24.33 (sec) , antiderivative size = 747, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3782, 6857, 654, 626, 635, 212, 1035, 1092, 1050, 1044, 214, 211}

$$\begin{aligned}
& \int \cot^3(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx \\
& = \frac{\sqrt{-a(\sqrt{a^2-2ac+b^2+c^2}+2c)+c(\sqrt{a^2-2ac+b^2+c^2}+c)+a^2+b^2} \operatorname{arctan}\left(\frac{\sqrt{2}^4 \sqrt{a^2-2ac+b^2+c^2}}{\sqrt{2} e^4 \sqrt{a^2-2ac+b^2+c^2}}\right)}{\sqrt{2} e^4 \sqrt{a^2-2ac+b^2+c^2}} \\
& - \frac{\sqrt{-a(2c-\sqrt{a^2-2ac+b^2+c^2})+c(c-\sqrt{a^2-2ac+b^2+c^2})+a^2+b^2} \operatorname{arctanh}\left(\frac{\sqrt{2}^4 \sqrt{a^2-2ac+b^2+c^2}}{\sqrt{2} e^4 \sqrt{a^2-2ac+b^2+c^2}}\right)}{\sqrt{2} e^4 \sqrt{a^2-2ac+b^2+c^2}} \\
& - \frac{b(b^2-4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{5/2}e} \\
& + \frac{b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
& + \frac{b(b+2c \cot(d+ex)) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{8c^2e} \\
& - \frac{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}}{3ce} + \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{e}
\end{aligned}$$

[In] Int[Cot[d + e*x]^3*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] (Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*ArcTan[(b^2 + (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*Sqrt[a^2 + b^2 - 2*a*c + c^2]*Cot[d + e*x])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2])])]

$$\begin{aligned} &^2]) - a*(2*c + \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]))*\text{Sqrt}[a + b*\text{Cot}[d + e*x] + c \\ &*\text{Cot}[d + e*x]^2]]]/(\text{Sqrt}[2]*(a^2 + b^2 - 2*a*c + c^2)^{(1/4)}*e) + (b*\text{ArcTan} \\ &\text{h}[(b + 2*c*\text{Cot}[d + e*x])/(2*\text{Sqrt}[c]*\text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x \\ &]^2))]/(2*\text{Sqrt}[c]*e) - (b*(b^2 - 4*a*c)*\text{ArcTanh}[(b + 2*c*\text{Cot}[d + e*x])/(2* \\ &\text{Sqrt}[c]*\text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2))]/(16*c^{(5/2)}*e) - (\text{Sq} \\ &\text{rt}[a^2 + b^2 + c*(c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - \text{Sqrt}[a^2 + \\ &b^2 - 2*a*c + c^2]))*\text{ArcTanh}[(b^2 + (a - c)*(a - c + \text{Sqrt}[a^2 + b^2 - 2*a*c \\ &+ c^2]) + b*\text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]*\text{Cot}[d + e*x])]/(\text{Sqrt}[2]*(a^2 + b^ \\ &2 - 2*a*c + c^2)^{(1/4)}*\text{Sqrt}[a^2 + b^2 + c*(c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2 \\ &]) - a*(2*c - \text{Sqrt}[a^2 + b^2 - 2*a*c + c^2]))*\text{Sqrt}[a + b*\text{Cot}[d + e*x] + c*C \\ &\text{ot}[d + e*x]^2]]]/(\text{Sqrt}[2]*(a^2 + b^2 - 2*a*c + c^2)^{(1/4)}*e) + \text{Sqrt}[a + b* \\ &\text{Cot}[d + e*x] + c*\text{Cot}[d + e*x]^2]/e + (b*(b + 2*c*\text{Cot}[d + e*x])* \text{Sqrt}[a + b*C \\ &\text{ot}[d + e*x] + c*\text{Cot}[d + e*x]^2])/(8*c^2*e) - (a + b*\text{Cot}[d + e*x] + c*\text{Cot}[d \\ &+ e*x]^2)^{(3/2)}/(3*c*e) \end{aligned}$$
Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 626

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x) * ((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]
```

Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 654

```
Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[(2*c*d - b
```


$e)/(2*c), \text{Int}[(a + b*x + c*x^2)^p, x] /; \text{FreeQ}\{a, b, c, d, e, p\}, x\}$
 $\&\& \text{NeQ}[2*c*d - b*e, 0] \&\& \text{NeQ}[p, -1]$

Rule 1035

$\text{Int}[(g_. + (h_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^{(p_)}*((d_.) + (f_.)*(x_)^2)^{(q_)}, x_Symbol] \rightarrow \text{Simp}[h*(a + b*x + c*x^2)^p*((d + f*x^2)^{q+1}/(2*f*(p + q + 1))), x] - \text{Dist}[1/(2*f*(p + q + 1)), \text{Int}[(a + b*x + c*x^2)^{(p-1)}*(d + f*x^2)^q*\text{Simp}[h*p*(b*d) + a*(-2*g*f)*(p + q + 1) + (2*h*p*(c*d - a*f) + b*(-2*g*f)*(p + q + 1))*x + (h*p*((-b)*f) + c*(-2*g*f)*(p + q + 1))*x^2, x], x] /; \text{FreeQ}\{a, b, c, d, f, g, h, q\}, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{GtQ}[p, 0] \&\& \text{NeQ}[p + q + 1, 0]$

Rule 1044

$\text{Int}[(g_. + (h_.)*(x_))/((a_.) + (c_.)*(x_)^2)*\text{Sqrt}[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] \rightarrow \text{Dist}[-2*a*g*h, \text{Subst}[\text{Int}[1/\text{Simp}[2*a^2*g*h*c + a*e*x^2, x], x], x, \text{Simp}[a*h - g*c*x, x]/\text{Sqrt}[d + e*x + f*x^2]], x] /; \text{FreeQ}\{a, c, d, e, f, g, h\}, x\} \&\& \text{EqQ}[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]$

Rule 1050

$\text{Int}[(g_. + (h_.)*(x_))/((a_.) + (c_.)*(x_)^2)*\text{Sqrt}[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[(c*d - a*f)^2 + a*c*e^2, 2]\}, \text{Dist}[1/(2*q), \text{Int}[\text{Simp}[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x], x] - \text{Dist}[1/(2*q), \text{Int}[\text{Simp}[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x], x]] /; \text{FreeQ}\{a, c, d, e, f, g, h\}, x\} \&\& \text{NeQ}[e^2 - 4*d*f, 0] \&\& \text{NegQ}[(-a)*c]$

Rule 1092

$\text{Int}[(A_.) + (B_.)*(x_) + (C_.)*(x_)^2)/((a_.) + (c_.)*(x_)^2)*\text{Sqrt}[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] \rightarrow \text{Dist}[C/c, \text{Int}[1/\text{Sqrt}[d + e*x + f*x^2], x], x] + \text{Dist}[1/c, \text{Int}[(A*c - a*C + B*c*x)/((a + c*x^2)*\text{Sqrt}[d + e*x + f*x^2]), x], x] /; \text{FreeQ}\{a, c, d, e, f, A, B, C\}, x\} \&\& \text{NeQ}[e^2 - 4*d*f, 0]$

Rule 3782

$\text{Int}[\text{cot}[(d_.) + (e_.)*(x_)]^{(m_.)}*((a_.) + (b_.)*\text{cot}[(d_.) + (e_.)*(x_)]*(f_.))^{(n_.)} + (c_.)*\text{cot}[(d_.) + (e_.)*(x_)]*(f_.))^{(n2_.)}]^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[-f/e, \text{Subst}[\text{Int}[(x/f)^m*((a + b*x^n + c*x^{(2*n)})^p/(f^2 + x^2)), x], x, f*\text{Cot}[d + e*x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x\} \&\& \text{EqQ}[n^2, 2*n] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 6857

`Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]`

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^3\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \left(x\sqrt{a+bx+cx^2} - \frac{x\sqrt{a+bx+cx^2}}{1+x^2}\right) dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int x\sqrt{a+bx+cx^2} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} - \frac{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}}{3ce} \\
&\quad - \frac{\text{Subst}\left(\int \frac{\frac{b}{2} - (a-c)x - \frac{bx^2}{2}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&\quad + \frac{b\text{Subst}\left(\int \sqrt{a+bx+cx^2} dx, x, \cot(d+ex)\right)}{2ce} \\
&= \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} \\
&\quad + \frac{b(b+2c\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{8c^2e} \\
&\quad - \frac{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}}{3ce} \\
&\quad - \frac{\text{Subst}\left(\int \frac{b+(-a+c)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&\quad + \frac{b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2e} \\
&\quad - \frac{(b^2-4ac)\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{16c^2e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&+ \frac{b(b + 2c \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2 e} \\
&- \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&+ \frac{b \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&- \frac{(b(b^2 - 4ac)) \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8c^2 e} \\
&+ \frac{\operatorname{Subst}\left(\int \frac{-b\sqrt{a^2+b^2-2ac+c^2} + (-b^2 - (a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&- \frac{\operatorname{Subst}\left(\int \frac{b\sqrt{a^2+b^2-2ac+c^2} + (-b^2 - (a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2} e} \\
&= \frac{\operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
&- \frac{b(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{5/2} e} \\
&+ \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e} \\
&+ \frac{b(b + 2c \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{8c^2 e} \\
&- \frac{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}}{3ce} \\
&- \frac{(b(b^2 + (a - c)(a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \operatorname{Subst}\left(\int \frac{1}{2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d + ex)\right)}{e} \\
&- \frac{(b(b^2 + (a - c)(a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \operatorname{Subst}\left(\int \frac{1}{-2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d + ex)\right)}{e}
\end{aligned}$$

$$\begin{aligned}
& \frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})} \arctan\left(\frac{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}} \\
& + \frac{\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
& - \frac{b(b^2 - 4ac) \operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{16c^{5/2}e} \\
& - \frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}} \\
& + \frac{\sqrt{a + b\cot(d + ex) + c\cot^2(d + ex)}}{e} \\
& + \frac{b(b + 2c\cot(d + ex))\sqrt{a + b\cot(d + ex) + c\cot^2(d + ex)}}{8c^2e} \\
& - \frac{(a + b\cot(d + ex) + c\cot^2(d + ex))^{3/2}}{3ce}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.30 (sec) , antiderivative size = 381, normalized size of antiderivative = 0.51

$$\begin{aligned}
& \int \cot^3(d + ex) \sqrt{a + b\cot(d + ex) + c\cot^2(d + ex)} dx \\
& = \frac{\sqrt{a + b\cot(d + ex) + c\cot^2(d + ex)} \tan(d + ex) \left(-3b(b^2 - 4c(a + 2c)) \operatorname{arctanh}\left(\frac{2c + b\tan(d+ex)}{2\sqrt{c}\sqrt{c + b\tan(d+ex) + a\tan^2(d+ex)}}\right)\right)}{\sqrt{a + b\cot(d + ex) + c\cot^2(d + ex)}}
\end{aligned}$$

[In] Integrate[Cot[d + e*x]^3*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] (Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]*(-3*b*(b^2 - 4*c*(a + 2*c))*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])] + 2*Sqrt[c]*((1*I)*Sqrt[a + I*b - c])*c^2*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])] + (1*I)*Sqrt[a - I*b - c])*c^2*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])] + Cot[d + e*x]*(3*b^2 - 8*a*c + 24*c^2 - 2*b*c*Cot[d + e*x] - 8*c^2*Cot[d + e*x]^2)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))/(48*c^(5/2)*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 3.96 (sec) , antiderivative size = 17768080, normalized size of antiderivative = 23785.92

output too large to display

[In] `int(cot(e*x+d)^3*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)`

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5923 vs. 2(670) = 1340.

Time = 2.41 (sec) , antiderivative size = 11897, normalized size of antiderivative = 15.93

$$\int \cot^3(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx = \text{Too large to display}$$

[In] `integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\begin{aligned} & \int \cot^3(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx \\ &= \int \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} \cot^3(d+ex) dx \end{aligned}$$

[In] `integrate(cot(e*x+d)**3*(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)`

[Out] `Integral(sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2)*cot(d + e*x)**3, x)`

Maxima [F]

$$\int \cot^3(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx$$

$$= \int \sqrt{c\cot(ex+d)^2+b\cot(ex+d)+a\cot(ex+d)^3}dx$$

[In] integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d)^3, x)

Giac [F]

$$\int \cot^3(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx$$

$$= \int \sqrt{c\cot(ex+d)^2+b\cot(ex+d)+a\cot(ex+d)^3}dx$$

[In] integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \cot^3(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}dx$$

$$= \int \cot(d+ex)^3\sqrt{c\cot(d+ex)^2+b\cot(d+ex)+a}dx$$

[In] int(cot(d + e*x)^3*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(cot(d + e*x)^3*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

3.8 $\int \cot(d+ex) \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)} dx$

Optimal result	95
Rubi [A] (verified)	96
Mathematica [C] (verified)	99
Maple [B] (warning: unable to verify)	100
Fricas [B] (verification not implemented)	100
Sympy [F]	100
Maxima [F]	101
Giac [F]	101
Mupad [F(-1)]	101

Optimal result

Integrand size = 31, antiderivative size = 602

$$\int \cot(d+ex) \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)} dx =$$

$$\frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})} \arctan\left(\frac{\sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right) - \frac{\operatorname{barctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} + \frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})} \operatorname{arctanh}\left(\frac{\sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right) - \frac{\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}}{e}}$$

```
[Out] -1/2*b*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/e/c^(1/2)-(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/e+1/2*arctanh(1/2*(b^2+b*cot(e*x+d)*(a^2-2*a*c+b^2+c^2)^(1/2)+(a-c)*(a-c+(a^2-2*a*c+b^2+c^2)^(1/2))))/(a^2-2*a*c+b^2+c^2)^(1/4)*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a^2+b^2+c*(c-(a^2-2*a*c+b^2+c^2)^(1/2))-a*(2*c-(a^2-2*a*c+b^2+c^2)^(1/2)))^(1/2)*(a^2+b^2+c*(c-(a^2-2*a*c+b^2+c^2)^(1/2))-a*(2*c-(a^2-2*a*c+b^2+c^2)^(1/2)))^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/4)/e*2^(1/2)-1/2*arctan(1/2*(b^2+(a-c)*(a-c-(a^2-2*a*c+b^2+c^2)^(1/2))-b*cot(e*x+d)*(a^2-2*a*c+b^2+c^2)^(1/2))/(a^2-2*a*c+b^2+c^2)^(1/4)*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(a^2+b^2+c*(c+(a^2-2*a*c+b^2+c^2)^(1/2))-a*(2*c+(a^2-2*a*c+b^2+c^2)^(1/2)))^(1/2)*(a^2+b^2+c*(c+(a^2-2*a*c+b^2+c^2)^(1/2))-a*(2*c+(a^2-2*a*c+b^2+c^2)^(1/2)))^(1/2)/(a^2-2*a*c+b^2+c^2)^(1/4)/e*2^(1/2)
```

Rubi [A] (verified)

Time = 23.71 (sec) , antiderivative size = 602, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.290$, Rules used = {3782, 1035, 1092, 635, 212, 1050, 1044, 214, 211}

$$\int \cot(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx =$$

$$-\frac{\sqrt{-a(\sqrt{a^2-2ac+b^2+c^2}+2c)+c(\sqrt{a^2-2ac+b^2+c^2}+c)+a^2+b^2} \arctan\left(\frac{\sqrt{-a(\sqrt{a^2-2ac+b^2+c^2}+2c)+c(\sqrt{a^2-2ac+b^2+c^2}+c)+a^2+b^2}}{\sqrt{2} \sqrt[4]{a^2-2ac+b^2+c^2}}\right)}{\sqrt{2} \sqrt[4]{a^2-2ac+b^2+c^2}}$$

$$+\frac{\sqrt{-a(2c-\sqrt{a^2-2ac+b^2+c^2})+c(c-\sqrt{a^2-2ac+b^2+c^2})+a^2+b^2} \operatorname{arctanh}\left(\frac{\sqrt{-a(2c-\sqrt{a^2-2ac+b^2+c^2})+c(c-\sqrt{a^2-2ac+b^2+c^2})+a^2+b^2}}{\sqrt{2} \sqrt[4]{a^2-2ac+b^2+c^2}}\right)}{\sqrt{2} \sqrt[4]{a^2-2ac+b^2+c^2}}$$

$$-\frac{\operatorname{barctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} - \frac{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{e}$$

[In] Int[Cot[d + e*x]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2], x]

[Out] -((Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*ArcTan[(b^2 + (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e)) - (b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(2*Sqrt[c]*e) + (Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])]*ArcTanh[(b^2 + (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) + b*Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e) - Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]/e

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

$Q[a, 0] \parallel LtQ[b, 0]$

Rule 214

$\text{Int}[(a_ + (b_ \cdot)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a) \cdot \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b]$

Rule 635

$\text{Int}[1/\text{Sqrt}[(a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)], x_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 1035

$\text{Int}[(g_ + (h_ \cdot)(x_)) \cdot ((a_ + (b_ \cdot)(x_) + (c_ \cdot)(x_)^2)^{p_}) \cdot ((d_ + (f_ \cdot)(x_)^2)^{q_}), x_Symbol] \rightarrow \text{Simp}[h \cdot (a + b*x + c*x^2)^p \cdot (d + f*x^2)^{q+1} / (2*f*(p+q+1)), x] - \text{Dist}[1/(2*f*(p+q+1)), \text{Int}[(a + b*x + c*x^2)^{p-1} \cdot (d + f*x^2)^q \cdot \text{Simp}[h*p*(b*d) + a*(-2*g*f)*(p+q+1) + (2*h*p*(c*d - a*f) + b*(-2*g*f)*(p+q+1))*x + (h*p*(-b)*f) + c*(-2*g*f)*(p+q+1)]*x^2, x], x] /; \text{FreeQ}\{a, b, c, d, f, g, h, q\}, x \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{GtQ}[p, 0] \ \&\& \ \text{NeQ}[p+q+1, 0]$

Rule 1044

$\text{Int}[(g_ + (h_ \cdot)(x_)) / (((a_ + (c_ \cdot)(x_)^2) \cdot \text{Sqrt}[(d_ + (e_ \cdot)(x_) + (f_ \cdot)(x_)^2)]), x_Symbol] \rightarrow \text{Dist}[-2*a*g*h, \text{Subst}[\text{Int}[1/\text{Simp}[2*a^2*g*h*c + a*e*x^2, x], x], x, \text{Simp}[a*h - g*c*x, x]/\text{Sqrt}[d + e*x + f*x^2]], x] /; \text{FreeQ}\{a, c, d, e, f, g, h\}, x \ \&\& \ \text{EqQ}[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]$

Rule 1050

$\text{Int}[(g_ + (h_ \cdot)(x_)) / (((a_ + (c_ \cdot)(x_)^2) \cdot \text{Sqrt}[(d_ + (e_ \cdot)(x_) + (f_ \cdot)(x_)^2)]), x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[(c*d - a*f)^2 + a*c*e^2, 2]\}, \text{Dist}[1/(2*q), \text{Int}[\text{Simp}[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x] / ((a + c*x^2) \cdot \text{Sqrt}[d + e*x + f*x^2]), x], x] - \text{Dist}[1/(2*q), \text{Int}[\text{Simp}[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x] / ((a + c*x^2) \cdot \text{Sqrt}[d + e*x + f*x^2]), x], x]] /; \text{FreeQ}\{a, c, d, e, f, g, h\}, x \ \&\& \ \text{NeQ}[e^2 - 4*d*f, 0] \ \&\& \ \text{NegQ}[(-a)*c]$

Rule 1092

$\text{Int}[(A_ + (B_ \cdot)(x_) + (C_ \cdot)(x_)^2) / (((a_ + (c_ \cdot)(x_)^2) \cdot \text{Sqrt}[(d_ + (e_ \cdot)(x_) + (f_ \cdot)(x_)^2)]), x_Symbol] \rightarrow \text{Dist}[C/c, \text{Int}[1/\text{Sqrt}[d + e*x + f*x^2], x], x] + \text{Dist}[1/c, \text{Int}[(A*c - a*C + B*c*x) / ((a + c*x^2) \cdot \text{Sqrt}[d + e*x + f*x^2]), x], x] /; \text{FreeQ}\{a, c, d, e, f, A, B, C\}, x \ \&\& \ \text{NeQ}[e^2 - 4*d*f]$

, 0]

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_.)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n2_.))^(p_), x_Symbol]
  :> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} + \frac{\text{Subst}\left(\int \frac{\frac{b}{2}-(a-c)x-\frac{bx^2}{2}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{b+(-a+c)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&\quad - \frac{b\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2e} \\
&= -\frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} - \frac{b\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-b\sqrt{a^2+b^2-2ac+c^2}+(-b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{b\sqrt{a^2+b^2-2ac+c^2}+(-b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&= -\frac{\text{barctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2\sqrt{c}e} - \frac{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{e} \\
&\quad + \frac{(b(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2})))\text{Subst}\left(\int \frac{1}{2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d+ex)\right)}{e} \\
&\quad + \frac{(b(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2})))\text{Subst}\left(\int \frac{1}{-2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d+ex)\right)}{e}
\end{aligned}$$

$$\begin{aligned}
&= \\
&\frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})} \arctan\left(\frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})}}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac}}\right)}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac}} \\
&- \frac{\operatorname{barctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2\sqrt{ce}} \\
&+ \frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})} \operatorname{arctanh}\left(\frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})}}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac}}\right)}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac}} \\
&- \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.43 (sec) , antiderivative size = 324, normalized size of antiderivative = 0.54

$$\int \cot(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} dx = \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) \left(\operatorname{barctanh}\left(\frac{2c + b \tan(d + ex)}{2\sqrt{c}\sqrt{c + b \tan(d + ex) + a \tan^2(d + ex)}}\right) + \sqrt{c} \left(i\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \right) \right)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}$$

[In] Integrate[Cot[d + e*x]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2],x]

[Out] -1/2*(Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]*(b*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + Sqrt[c]*(I*Sqrt[a + I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + I*Sqrt[a - I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]) + 2*Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))/(Sqrt[c]*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.44 (sec) , antiderivative size = 17767874, normalized size of antiderivative = 29514.74

output too large to display

[In] `int(cot(e*x+d)*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x)`

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 5650 vs. 2(543) = 1086.

Time = 2.30 (sec) , antiderivative size = 11351, normalized size of antiderivative = 18.86

$$\int \cot(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx = \text{Too large to display}$$

[In] `integrate(cot(e*x+d)*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\begin{aligned} & \int \cot(d+ex) \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} dx \\ &= \int \sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)} \cot(d+ex) dx \end{aligned}$$

[In] `integrate(cot(e*x+d)*(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2),x)`

[Out] `Integral(sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2)*cot(d + e*x), x)`

Maxima [F]

$$\int \cot(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$$

$$= \int \sqrt{c \cot (ex+d)^2 + b \cot (ex+d) + a} \cot (ex+d) dx$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d), x)

Giac [F]

$$\int \cot(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$$

$$= \int \sqrt{c \cot (ex+d)^2 + b \cot (ex+d) + a} \cot (ex+d) dx$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*cot(e*x + d), x)

Mupad [F(-1)]

Timed out.

$$\int \cot(d+ex) \sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} dx$$

$$= \int \cot(d+ex) \sqrt{c \cot (d+ex)^2 + b \cot (d+ex) + a} dx$$

[In] int(cot(d + e*x)*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(cot(d + e*x)*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

3.9 $\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$

Optimal result	102
Rubi [A] (verified)	103
Mathematica [C] (verified)	107
Maple [F(-1)]	107
Fricas [B] (verification not implemented)	107
Sympy [F]	110
Maxima [F(-2)]	110
Giac [F]	111
Mupad [F(-1)]	111

Optimal result

Integrand size = 31, antiderivative size = 570

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$$

$$= \frac{\sqrt{a^2 + b^2 + c} (c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2}) \arctan\left(\frac{b^2 + \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2} e} + \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a} \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{e} - \frac{\sqrt{a^2 + b^2 + c} (c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c - \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{b^2 + \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2} e}$$

[Out] $\operatorname{arctanh}\left(\frac{1}{2} \frac{2a + b \cot(ex + d)}{\sqrt{a} \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)}}\right) / \sqrt{a} / \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)}^{1/2} - \frac{1}{2} \operatorname{arctanh}\left(\frac{1}{2} \frac{b^2 + b \cot(ex + d) \sqrt{a^2 - 2ac + b^2 + c^2}}{(a - c) \sqrt{a^2 - 2ac + b^2 + c^2}}\right) / \sqrt{a^2 - 2ac + b^2 + c^2}^{1/4} \sqrt{2}^{1/2} / \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)}^{1/2} / \sqrt{a^2 + b^2 + c} (c - \sqrt{a^2 - 2ac + b^2 + c^2})^{1/2} - a (2c - \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{b^2 + \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}\right) / \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2} e + \frac{1}{2} \operatorname{arctan}\left(\frac{1}{2} \frac{b^2 + (a - c) \sqrt{a^2 - 2ac + b^2 + c^2}}{(a - c) \sqrt{a^2 - 2ac + b^2 + c^2}}\right) - b \cot(ex + d) \sqrt{a^2 - 2ac + b^2 + c^2}^{1/2} / \sqrt{a^2 - 2ac + b^2 + c^2}^{1/4} \sqrt{2}^{1/2} / \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)}^{1/2} / \sqrt{a^2 + b^2 + c} (c + \sqrt{a^2 - 2ac + b^2 + c^2})^{1/2} - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{b^2 + \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}\right) / \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2} e + \frac{1}{2} \operatorname{arctan}\left(\frac{1}{2} \frac{b^2 + (a - c) \sqrt{a^2 - 2ac + b^2 + c^2}}{(a - c) \sqrt{a^2 - 2ac + b^2 + c^2}}\right) - b \cot(ex + d) \sqrt{a^2 - 2ac + b^2 + c^2}^{1/2} / \sqrt{a^2 - 2ac + b^2 + c^2}^{1/4} \sqrt{2}^{1/2} / \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)}^{1/2} / \sqrt{a^2 + b^2 + c} (c + \sqrt{a^2 - 2ac + b^2 + c^2})^{1/2} - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{b^2 + \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2}}\right) / \sqrt{2}^4 \sqrt{a^2 + b^2 - 2ac + c^2} e$

Rubi [A] (verified)

Time = 24.18 (sec) , antiderivative size = 570, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.419$, Rules used = {3782, 6857, 748, 857, 635, 212, 738, 1035, 1092, 1050, 1044, 214, 211}

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$$

$$= \frac{\sqrt{-a(\sqrt{a^2 - 2ac + b^2 + c^2} + 2c) + c(\sqrt{a^2 - 2ac + b^2 + c^2} + c) + a^2 + b^2} \arctan\left(\frac{\sqrt{a^2 - 2ac + b^2 + c^2}}{\sqrt{2}\sqrt[4]{a^2 - 2ac + b^2 + c^2}}\right) - \sqrt{-a(2c - \sqrt{a^2 - 2ac + b^2 + c^2}) + c(c - \sqrt{a^2 - 2ac + b^2 + c^2}) + a^2 + b^2} \operatorname{arctanh}\left(\frac{\sqrt{a^2 - 2ac + b^2 + c^2}}{\sqrt{2}\sqrt[4]{a^2 - 2ac + b^2 + c^2}}\right)}{\sqrt{2}e\sqrt[4]{a^2 - 2ac + b^2 + c^2}} + \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{e}$$

[In] Int[Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x], x]

[Out] (Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2])] - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*ArcTan[(b^2 + (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*Sqrt[a^2 + b^2 - 2*a*c + c^2]*Cot[d + e*x])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e) + (Sqrt[a]*ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/e - (Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2])] - a*(2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]))*ArcTanh[(b^2 + (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) + b*Sqrt[a^2 + b^2 - 2*a*c + c^2]*Cot[d + e*x])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e)

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

Q[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 748

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[p/(e*(m + 2*p + 1)), Int[(d + e*x)^m*Simp[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 857

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1035

Int[((g_) + (h_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (f_)*(x_)^2)^(q_), x_Symbol] := Simp[h*(a + b*x + c*x^2)^p*((d + f*x^2)^(q + 1)/(2*f*(p + q + 1))), x] - Dist[1/(2*f*(p + q + 1)), Int[(a + b*x + c*x^2)^(p - 1)*(d + f*x^2)^q*Simp[h*p*(b*d) + a*(-2*g*f)*(p + q + 1) + (2*h*p*(c*d - a*f) + b*(-2*g*f)*(p + q + 1))*x + (h*p*((-b)*f) + c*(-2*g*f)*(p + q + 1))*x^2, x], x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*a

*c, 0] && GtQ[p, 0] && NeQ[p + q + 1, 0]

Rule 1044

Int[((g_) + (h_)*(x_))/((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2], x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_) + (h_)*(x_))/((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2], x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 1092

Int[((A_) + (B_)*(x_) + (C_)*(x_)^2)/((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2], x_Symbol] := Dist[C/c, Int[1/Sqrt[d + e*x + f*x^2], x], x] + Dist[1/c, Int[(A*c - a*C + B*c*x)/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, A, B, C}, x] && NeQ[e^2 - 4*d*f, 0]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n2_)]^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

Int[(u_)/((a_) + (b_)*(x_)^n), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rubi steps

$$\text{integral} = -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x(1+x^2)} dx, x, \cot(d+ex)\right)}{e}$$

$$\begin{aligned}
&= \frac{\text{Subst}\left(\int \left(\frac{\sqrt{a+bx+cx^2}}{x} - \frac{x\sqrt{a+bx+cx^2}}{1+x^2}\right) dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \frac{-2a-bx}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2e} - \frac{\text{Subst}\left(\int \frac{\frac{b}{2} - (a-c)x - \frac{bx^2}{2}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\text{Subst}\left(\int \frac{b+(-a+c)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} - \frac{a\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{(2a)\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b\cot(d+ex)}{\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2\sqrt{a^2+b^2-2ac+c^2}e} \\
&= \frac{\sqrt{a}\text{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{e} \\
&\quad - \frac{(b(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2})))\text{Subst}\left(\int \frac{1}{2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d+ex)\right)}{e} \\
&\quad - \frac{(b(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2})))\text{Subst}\left(\int \frac{1}{-2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d+ex)\right)}{e} \\
&= \frac{\sqrt{a^2+b^2+c}(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2})\arctan\left(\frac{1}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}} \\
&\quad + \frac{\sqrt{a}\text{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{e} \\
&\quad - \frac{\sqrt{a^2+b^2+c}(c-\sqrt{a^2+b^2-2ac+c^2})-a(2c-\sqrt{a^2+b^2-2ac+c^2})\text{arctanh}\left(\frac{1}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2+b^2-2ac+c^2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.73 (sec) , antiderivative size = 283, normalized size of antiderivative = 0.50

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$$

$$= \frac{\left(i \left(\sqrt{a + ib - c} \arctan \left(\frac{ib - 2c + (2ia - b) \tan(d + ex)}{2\sqrt{a + ib - c} \sqrt{c + b \tan(d + ex)} + a \tan^2(d + ex)} \right) + \sqrt{a - ib - c} \arctan \left(\frac{ib + 2c + (2ia + b) \tan(d + ex)}{2\sqrt{a - ib - c} \sqrt{c + b \tan(d + ex)} + a \tan^2(d + ex)} \right) \right)}{2e \sqrt{c + b \tan(d + ex)}}$$

[In] Integrate[Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x],x]

[Out] ((I*(Sqrt[a + I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x]])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + Sqrt[a - I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x]])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])) + 2*Sqrt[a]*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x])/(2*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])

Maple [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot(ex + d) + c \cot^2(ex + d)} \tan(ex + d) dx$$

[In] int((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d),x)

[Out] int((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2423 vs. 2(515) = 1030.

Time = 0.68 (sec) , antiderivative size = 4847, normalized size of antiderivative = 8.50

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx = \text{Too large to display}$$

[In] integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d),x, algorithm="fricas")

[Out] [-1/4*(e*sqrt((e^2*sqrt(-b^2/e^4) + a - c)/e^2)*log((((2*a^2*b^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*tan(e*x + d)^2 + 4*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*tan(e*x + d) + (b^5 - 2*a*b^3*c + 6*b^3*c^2 - 8*a

$$\begin{aligned}
& *b*c^3 + 8*b*c^4)*e + ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^3 - (8*a^2*b - 3*b^3)*c \\
&)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4* \\
& a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c + 4*a*b*c^2 + 4*b*c^3)* \\
& e^3)*\sqrt{-b^2/e^4})*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2} + 2*((a^2*b^3 - \\
& b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + 3*b^3)*c^2)*\tan(e*x + d \\
&)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\tan(e*x + d) + (2*(a*b^3 \\
& + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - (a^2*b^2 - b^4 + 6*a*b^ \\
& 2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x + d))*\sqrt{-b^2/e^ \\
& 4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x + d)^2}))/(\tan(e*x \\
& + d)^2 + 1)) - e*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2})*\log(-(((2*a^2*b^3 - \\
& b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x + d)^2 + 4*(a*b^4 + \\
& 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 - 2*a*b^3*c + 6*b^3* \\
& c^2 - 8*a*b*c^3 + 8*b*c^4)*e + ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^3 - (8*a^2*b - \\
& 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4* \\
& c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c + 4*a*b*c^2 + \\
& 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2} - 2*((\\
& a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + 3*b^3)*c^2)*\tan \\
& (e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\tan(e*x + d) + (\\
& 2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - (a^2*b^2 - b^4 \\
& + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x + d))*\sqrt{ \\
& -b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x + d)^2}))/ \\
& (\tan(e*x + d)^2 + 1)) + e*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + c)/e^2})*\log((((2* \\
& a^2*b^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x + d)^2 + 4* \\
& (a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 - 2*a*b^3*c \\
& + 6*b^3*c^2 - 8*a*b*c^3 + 8*b*c^4)*e - ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^3 - (\\
& 8*a^2*b - 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c + 8*a \\
& *c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c + 4*a \\
& *b*c^2 + 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + c)/e \\
& ^2} + 2*((a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + 3*b^ \\
& 3)*c^2)*\tan(e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\tan(e* \\
& x + d) - (2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - (a^2 \\
& *b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x \\
& + d))*\sqrt{-b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x \\
& + d)^2}))/(\tan(e*x + d)^2 + 1)) - e*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + c)/e^2} \\
&)*\log(-((((2*a^2*b^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x \\
& + d)^2 + 4*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 \\
& - 2*a*b^3*c + 6*b^3*c^2 - 8*a*b*c^3 + 8*b*c^4)*e - ((3*a*b^3 - 4*a*b*c^2 + \\
& 4*b*c^3 - (8*a^2*b - 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a* \\
& b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + \\
& b^3*c + 4*a*b*c^2 + 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{-(e^2*\sqrt{-b^2/e^4} \\
& - a + c)/e^2} - 2*((a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a \\
& ^2*b + 3*b^3)*c^2)*\tan(e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4 \\
&)*c)*\tan(e*x + d) - (2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + \\
& d)^2 - (a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)* \\
& e^2*\tan(e*x + d))*\sqrt{-b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) +
\end{aligned}$$

$$\begin{aligned}
& c)/\tan(e*x + d)^2)/(\tan(e*x + d)^2 + 1)) - 2*\sqrt{a}*\log(8*a^2*\tan(e*x + \\
& d)^2 + 8*a*b*\tan(e*x + d) + b^2 + 4*a*c + 4*(2*a*\tan(e*x + d)^2 + b*\tan(e*x \\
& + d))*\sqrt{a}*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x + d)^2} \\
&))/e, -1/4*(e*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2})*\log((((2*a^2*b^3 - b^5 \\
& + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x + d)^2 + 4*(a*b^4 + 2*b^ \\
& 2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 - 2*a*b^3*c + 6*b^3*c^2 \\
& - 8*a*b*c^3 + 8*b*c^4)*e + ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^3 - (8*a^2*b - 3*b \\
& ^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 \\
& - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c + 4*a*b*c^2 + 4*b* \\
& c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2} + 2*((a^2*b \\
& ^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + 3*b^3)*c^2)*\tan(e* \\
& x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\tan(e*x + d) + (2*(a \\
& *b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - (a^2*b^2 - b^4 + 6 \\
& *a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x + d))*\sqrt{-b \\
& ^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x + d)^2)/(\tan \\
& (e*x + d)^2 + 1)) - e*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2})*\log(-(((2*a^2*b \\
& ^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x + d)^2 + 4*(a*b \\
& ^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 - 2*a*b^3*c + 6 \\
& *b^3*c^2 - 8*a*b*c^3 + 8*b*c^4)*e + ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^3 - (8*a^ \\
& 2*b - 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 \\
& - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c + 4*a*b*c \\
& ^2 + 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{(e^2*\sqrt{-b^2/e^4} + a - c)/e^2} - \\
& 2*((a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + 3*b^3)*c^ \\
& 2)*\tan(e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\tan(e*x + d \\
&) + (2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - (a^2*b^2 \\
& - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x + d) \\
&)*\sqrt{-b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan(e*x + d) \\
& ^2)/(\tan(e*x + d)^2 + 1)) + e*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + c)/e^2})*\log(\\
& (((2*a^2*b^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan(e*x + d)^2 \\
& + 4*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + (b^5 - 2*a* \\
& b^3*c + 6*b^3*c^2 - 8*a*b*c^3 + 8*b*c^4)*e - ((3*a*b^3 - 4*a*b*c^2 + 4*b*c^ \\
& 3 - (8*a^2*b - 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + 6*a*b^2*c \\
& + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b^3 + b^3*c \\
& + 4*a*b*c^2 + 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + \\
& c)/e^2} + 2*((a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - (4*a^2*b + \\
& 3*b^3)*c^2)*\tan(e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*\t \\
& an(e*x + d) - (2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e*x + d)^2 - \\
& (a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^2*\tan(e*x + d) \\
&)*\sqrt{-b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + d) + c)/\tan \\
& (e*x + d)^2)/(\tan(e*x + d)^2 + 1)) - e*\sqrt{-(e^2*\sqrt{-b^2/e^4} - a + c) \\
& /e^2})*\log(-(((2*a^2*b^3 - b^5 + 10*a*b^3*c - 8*a^2*b*c^2 + 8*a*b*c^3)*e*\tan \\
& (e*x + d)^2 + 4*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 - b^4)*c)*e*\tan(e*x + d) + \\
& (b^5 - 2*a*b^3*c + 6*b^3*c^2 - 8*a*b*c^3 + 8*b*c^4)*e - ((3*a*b^3 - 4*a*b*c \\
& ^2 + 4*b*c^3 - (8*a^2*b - 3*b^3)*c)*e^3*\tan(e*x + d)^2 - 2*(a^2*b^2 - b^4 + \\
& 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)*c^2)*e^3*\tan(e*x + d) - (a*b
\end{aligned}$$

$$\begin{aligned} &^3 + b^3*c + 4*a*b*c^2 + 4*b*c^3)*e^3)*\sqrt{-b^2/e^4})*\sqrt{-(e^2*\sqrt{-b^2} \\ &/e^4) - a + c)/e^2) - 2*((a^2*b^3 - b^5 + 6*a*b^3*c + 8*a*b*c^3 - 4*b*c^4 - \\ &(4*a^2*b + 3*b^3)*c^2)*\tan(e*x + d)^2 + 2*(a*b^4 + 2*b^2*c^3 - (2*a^2*b^2 \\ &- b^4)*c)*\tan(e*x + d) - (2*(a*b^3 + 2*b*c^3 - (2*a^2*b - b^3)*c)*e^2*\tan(e \\ &*x + d)^2 - (a^2*b^2 - b^4 + 6*a*b^2*c + 8*a*c^3 - 4*c^4 - (4*a^2 + 3*b^2)* \\ &c^2)*e^2*\tan(e*x + d))*\sqrt{-b^2/e^4})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(e*x + \\ &d) + c)/\tan(e*x + d)^2))/(\tan(e*x + d)^2 + 1)) + 4*\sqrt{-a}*\arctan(1/2*(2* \\ &a*\tan(e*x + d)^2 + b*\tan(e*x + d))*\sqrt{-a})*\sqrt{(a*\tan(e*x + d)^2 + b*\tan(\\ &e*x + d) + c)/\tan(e*x + d)^2)/(a^2*\tan(e*x + d)^2 + a*b*\tan(e*x + d) + a*c \\ &))/e] \end{aligned}$$

Sympy [F]

$$\begin{aligned} &\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex) \tan(d + ex)} dx \\ &= \int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex) \tan(d + ex)} dx \end{aligned}$$

[In] integrate((a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2)*tan(e*x+d),x)

[Out] Integral(sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2)*tan(d + e*x), x)

Maxima [F(-2)]

Exception generated.

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex) \tan(d + ex)} dx = \text{Exception raised: ValueError}$$

[In] integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume((c-b-a)*(c+b-a)>0)', see 'assume?' for mor

Giac [F]

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$$

$$= \int \sqrt{c \cot^2(ex + d) + b \cot(ex + d) + a} \tan(ex + d) dx$$

[In] integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d),x, algorithm="giac")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*tan(e*x + d), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex) dx$$

$$= \int \tan(d + ex) \sqrt{c \cot^2(d + ex) + b \cot(d + ex) + a} dx$$

[In] int(tan(d + e*x)*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(tan(d + e*x)*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

3.10 $\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx$

Optimal result	112
Rubi [A] (verified)	113
Mathematica [C] (verified)	118
Maple [F(-1)]	119
Fricas [B] (verification not implemented)	119
Sympy [F]	119
Maxima [F]	119
Giac [F]	120
Mupad [F(-1)]	120

Optimal result

Integrand size = 33, antiderivative size = 691

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx =$$

$$\frac{\sqrt{a^2 + b^2 + c} (c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2}) \arctan\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}e}$$

$$- \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{e}$$

$$- \frac{(b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{8a^{3/2}e}$$

$$+ \frac{\sqrt{a^2 + b^2 + c} (c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c - \sqrt{a^2 + b^2 - 2ac + c^2}) \operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt[4]{a^2 + b^2 - 2ac + c^2}e}$$

$$+ \frac{(2a + b \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{4ae}$$

[Out] $-1/8*(-4*a*c+b^2)*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d))/a^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})/a^{(3/2)}/e-\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d))/a^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)})*a^{(1/2)}/e+1/2*\operatorname{arctanh}(1/2*(b^2+b*\cot(e*x+d)*(a^2-2*a*c+b^2+c^2)^{(1/2)+(a-c)*(a-c+(a^2-2*a*c+b^2+c^2)^{(1/2))})/(a^2-2*a*c+b^2+c^2)^{(1/4)}*2^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)/(a^2+b^2+c*(c-(a^2-2*a*c+b^2+c^2)^{(1/2))}-a*(2*c-(a^2-2*a*c+b^2+c^2)^{(1/2))})^{(1/2)})*(a^2+b^2+c*(c-(a^2-2*a*c+b^2+c^2)^{(1/2))}-a*(2*c-(a^2-2*a*c+b^2+c^2)^{(1/2))})^{(1/2)})^{(1/2)}$

$$\frac{1/2}{(a^2-2ac+b^2+c^2)^{1/4}} \frac{1}{e^{1/2}} - \frac{1}{2} \arctan\left(\frac{1}{2}(b^2+(a-c)(a-c-(a^2-2ac+b^2+c^2)^{1/2})) - b \cot(ex+d) (a^2-2ac+b^2+c^2)^{1/2}\right) / (a^2-2ac+b^2+c^2)^{1/4} \cdot 2^{1/2} / (a+b \cot(ex+d) + c \cot(ex+d)^2)^{1/2} / (a^2+b^2+c(c+(a^2-2ac+b^2+c^2)^{1/2}) - a(2c+(a^2-2ac+b^2+c^2)^{1/2}))^{1/2} \cdot (a^2+b^2+c(c+(a^2-2ac+b^2+c^2)^{1/2}) - a(2c+(a^2-2ac+b^2+c^2)^{1/2}))^{1/2} / (a^2-2ac+b^2+c^2)^{1/4} \frac{1}{e^{1/2}} + \frac{1}{4} (2a+b \cot(ex+d)) (a+b \cot(ex+d) + c \cot(ex+d)^2)^{1/2} \tan(ex+d)^2 / a/e$$

Rubi [A] (verified)

Time = 24.41 (sec) , antiderivative size = 691, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.424$, Rules used = {3782, 6857, 734, 738, 212, 748, 857, 635, 1035, 1092, 1050, 1044, 214, 211}

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx$$

$$= -\frac{(b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{8a^{3/2}e}$$

$$- \frac{\sqrt{-a(\sqrt{a^2 - 2ac + b^2 + c^2} + 2c) + c(\sqrt{a^2 - 2ac + b^2 + c^2} + c) + a^2 + b^2} \operatorname{arctan}\left(\frac{\sqrt{2}^4 \sqrt{a^2 - 2ac + b^2}}{\sqrt{2}e^4 \sqrt{a^2 - 2ac + b^2 + c^2}}\right)}{\sqrt{2}e^4 \sqrt{a^2 - 2ac + b^2 + c^2}}$$

$$+ \frac{\sqrt{-a(2c - \sqrt{a^2 - 2ac + b^2 + c^2}) + c(c - \sqrt{a^2 - 2ac + b^2 + c^2}) + a^2 + b^2} \operatorname{arctanh}\left(\frac{\sqrt{2}^4 \sqrt{a^2 - 2ac + b^2}}{\sqrt{2}e^4 \sqrt{a^2 - 2ac + b^2 + c^2}}\right)}{\sqrt{2}e^4 \sqrt{a^2 - 2ac + b^2 + c^2}}$$

$$- \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot(d + ex)}{2\sqrt{a}\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{e}$$

$$+ \frac{\tan^2(d + ex)(2a + b \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{4ae}$$

[In] Int[Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]^3,x]

[Out] -((Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2])] - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*ArcTan[(b^2 + (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x]]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^{1/4}*Sqrt[a^2 + b^2 + c*(c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^{1/4}*e) - (Sqrt[a]*ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/e - ((b^2 - 4*a*c)*ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(8*a^{3/2}*e) + (Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - Sqrt[a^2 + b^2 -

$$2*a*c + c^2)]*ArcTanh[(b^2 + (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) + b*Sqrt[a^2 + b^2 - 2*a*c + c^2]*Cot[d + e*x])/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*Sqrt[a^2 + b^2 + c*(c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - a*(2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(1/4)*e) + ((2*a + b*Cot[d + e*x])*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]^2)/(4*a*e)$$
Rule 211

$$\text{Int}[\frac{(a_.) + (b_.)(x_)^2}{(x_)^2}, x_Symbol] \rightarrow \text{Simp}[\frac{\text{Rt}[a/b, 2]}{a} * \text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$$
Rule 212

$$\text{Int}[\frac{(a_.) + (b_.)(x_)^2}{(x_)^2}, x_Symbol] \rightarrow \text{Simp}[\frac{1}{\text{Rt}[a, 2]*\text{Rt}[-b, 2]} * \text{ArcTanh}[\frac{\text{Rt}[-b, 2]*x}{\text{Rt}[a, 2]}], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$$
Rule 214

$$\text{Int}[\frac{(a_.) + (b_.)(x_)^2}{(x_)^2}, x_Symbol] \rightarrow \text{Simp}[\frac{\text{Rt}[-a/b, 2]}{a} * \text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$$
Rule 635

$$\text{Int}[1/\text{Sqrt}[(a_.) + (b_.)(x_) + (c_.)(x_)^2], x_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$$
Rule 734

$$\text{Int}[\frac{(d_.) + (e_.)(x_)}{(x_)} * \frac{(a_.) + (b_.)(x_) + (c_.)(x_)^2}{(x_)^2}, x_Symbol] \rightarrow \text{Simp}[\frac{-(d + e*x)^{m+1} * (d*b - 2*a*e + (2*c*d - b*e)*x) * ((a + b*x + c*x^2)^p / (2*(m+1)*(c*d^2 - b*d*e + a*e^2)))}{(x_)} + \text{Dist}[p * ((b^2 - 4*a*c) / (2*(m+1)*(c*d^2 - b*d*e + a*e^2))), \text{Int}[(d + e*x)^{m+2} * (a + b*x + c*x^2)^{p-1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0] \ \&\& \ \text{EqQ}[m + 2*p + 2, 0] \ \&\& \ \text{GtQ}[p, 0]$$
Rule 738

$$\text{Int}[1/((d_.) + (e_.)(x_)) * \text{Sqrt}[(a_.) + (b_.)(x_) + (c_.)(x_)^2], x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c, d, e, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0]$$
Rule 748

```

Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[p/(e*(m + 2*p + 1)), Int[(d + e*x)^m*Simp[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0]
&& GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0]
&& IntQuadraticQ[a, b, c, d, e, m, p, x]

```

Rule 857

```

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

```

Rule 1035

```

Int[((g_.) + (h_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_.) + (f_.)*(x_)^2)^(q_), x_Symbol]
:= Simp[h*(a + b*x + c*x^2)^p*((d + f*x^2)^(q + 1)/(2*f*(p + q + 1))), x] - Dist[1/(2*f*(p + q + 1)), Int[(a + b*x + c*x^2)^(p - 1)*(d + f*x^2)^q*Simp[h*p*(b*d) + a*(-2*g*f)*(p + q + 1) + (2*h*p*(c*d - a*f) + b*(-2*g*f)*(p + q + 1))*x + (h*p*((-b)*f) + c*(-2*g*f)*(p + q + 1))*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x]
&& NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && NeQ[p + q + 1, 0]

```

Rule 1044

```

Int[((g_.) + (h_.)*(x_))/(((a_.) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol]
:= Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x]
&& EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

```

Rule 1050

```

Int[((g_.) + (h_.)*(x_))/(((a_.) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol]
:= With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x]
&& NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

```

Rule 1092

```

Int[((A_.) + (B_.)*(x_) + (C_.)*(x_)^2)/(((a_.) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol]
:= Dist[C/c, Int[1/Sqrt[d + e*x + f

```

*x^2], x], x] + Dist[1/c, Int[(A*c - a*C + B*c*x)/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, A, B, C}, x] && NeQ[e^2 - 4*d*f, 0]

Rule 3782

Int[cot[(d_.) + (e_.)*(x_.)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n2_.))^(p_), x_Symbol] :> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x^3(1+x^2)} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{\sqrt{a+bx+cx^2}}{x^3} - \frac{\sqrt{a+bx+cx^2}}{x} + \frac{x\sqrt{a+bx+cx^2}}{1+x^2}\right) dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x^3} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
 &= \frac{(2a + b \cot(d+ex))\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)} \tan^2(d+ex)}{4ae} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{-2a-bx}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{2e} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{\frac{b}{2} - (a-c)x - \frac{bx^2}{2}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad + \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{8ae}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(2a + b \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{4ae} \\
&+ \frac{\text{Subst}\left(\int \frac{b+(-a+c)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{e} \\
&+ \frac{a \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{e} \\
&- \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{4ae} \\
&= -\frac{(b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8a^{3/2}e} \\
&+ \frac{(2a + b \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{4ae} \\
&- \frac{(2a) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&- \frac{\text{Subst}\left(\int \frac{-b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&+ \frac{\text{Subst}\left(\int \frac{b\sqrt{a^2+b^2-2ac+c^2} + (-b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2\sqrt{a^2 + b^2 - 2ac + c^2}e} \\
&= -\frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{e} \\
&- \frac{(b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8a^{3/2}e} \\
&+ \frac{(2a + b \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{4ae} \\
&+ \frac{(b(b^2 + (a - c)(a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d + ex)\right)}{e} \\
&+ \frac{(b(b^2 + (a - c)(a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{-2b\sqrt{a^2+b^2-2ac+c^2}(b^2+(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))} dx, x, \cot(d + ex)\right)}{e}
\end{aligned}$$

=

$$\begin{aligned}
& \frac{\sqrt{a^2 + b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})} \arctan\left(\frac{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}} \\
& - \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot(d+ex)}{2\sqrt{a}\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}}\right)}{2\sqrt{a}\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
& - \frac{(b^2 - 4ac) \operatorname{arctanh}\left(\frac{e}{2\sqrt{a}\sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}}\right)}{8a^{3/2}e} \\
& + \frac{\sqrt{a^2 + b^2 + c(c - \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c - \sqrt{a^2 + b^2 - 2ac + c^2})} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}}\right)}{\sqrt{2}\sqrt{a^2 + b^2 - 2ac + c^2}} \\
& + \frac{(2a + b \cot(d + ex))\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{4ae}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.48 (sec) , antiderivative size = 363, normalized size of antiderivative = 0.53

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx = \frac{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex) \left((8a^2 + b^2 - 4ac) \operatorname{arctanh}\left(\frac{b + 2a \tan(d+ex)}{2\sqrt{a}\sqrt{c + b \tan(d+ex) + a \tan^2(d+ex)}}\right) \right)}{2\sqrt{a}\sqrt{c + b \tan(d+ex) + a \tan^2(d+ex)}}$$

[In] Integrate[Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]^3,x]

[Out] -1/8*(Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]*Tan[d + e*x]^2*((8*a^2 + b^2 - 4*a*c)*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Cot[d + e*x] - 2*Sqrt[a]*((-2*I)*a*Sqrt[a + I*b - c]*ArcTan[(I*b - 2*c + ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Cot[d + e*x] - (2*I)*a*Sqrt[a - I*b - c]*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]*Cot[d + e*x] + (2*a + b*Cot[d + e*x])*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])))/(a^(3/2)*e*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])

Maple [F(-1)]

Timed out.

hanged

[In] `int((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d)^3,x)`

[Out] `int((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d)^3,x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2508 vs. 2(622) = 1244.

Time = 0.98 (sec) , antiderivative size = 5018, normalized size of antiderivative = 7.26

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx = \text{Too large to display}$$

[In] `integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d)^3,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\begin{aligned} & \int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx \\ &= \int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx \end{aligned}$$

[In] `integrate((a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(1/2)*tan(e*x+d)**3,x)`

[Out] `Integral(sqrt(a + b*cot(d + e*x) + c*cot(d + e*x)**2)*tan(d + e*x)**3, x)`

Maxima [F]

$$\begin{aligned} & \int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx \\ &= \int \sqrt{c \cot(ex + d)^2 + b \cot(ex + d) + a} \tan(ex + d)^3 dx \end{aligned}$$

[In] `integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d)^3,x, algorithm="maxima")`

[Out] `integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*tan(e*x + d)^3, x)`

Giac [F]

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx$$

$$= \int \sqrt{c \cot^2(ex + d) + b \cot(ex + d) + a} \tan^3(ex + d) dx$$

[In] integrate((a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)*tan(e*x+d)^3,x, algorithm="giac")

[Out] integrate(sqrt(c*cot(e*x + d)^2 + b*cot(e*x + d) + a)*tan(e*x + d)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^3(d + ex) dx$$

$$= \int \tan^3(d + ex) \sqrt{c \cot^2(d + ex) + b \cot(d + ex) + a} dx$$

[In] int(tan(d + e*x)^3*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2),x)

[Out] int(tan(d + e*x)^3*(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(1/2), x)

$$3.11 \quad \int \frac{\cot^7(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	122
Rubi [A] (verified)	123
Mathematica [C] (verified)	133
Maple [B] (warning: unable to verify)	135
Fricas [B] (verification not implemented)	135
Sympy [F]	135
Maxima [F(-1)]	136
Giac [F(-2)]	136
Mupad [F(-1)]	136

Optimal result

Integrand size = 33, antiderivative size = 1189

$$\begin{aligned}
& \int \frac{\cot^7(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = -\frac{3b\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{2c^{5/2}e} \\
& + \frac{5b(7b^2-12ac)\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{16c^{9/2}e} \\
& + \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}+(a-c)\sqrt{a^2+b^2-2ac+c^2}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e} \\
& - \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}-(a-c)\sqrt{a^2+b^2-2ac+c^2}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e} \\
& - \frac{2(2a+b\cot(d+ex))}{(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
& + \frac{2\cot^2(d+ex)(2a+b\cot(d+ex))}{(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
& - \frac{2\cot^4(d+ex)(2a+b\cot(d+ex))}{(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
& + \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
& - \frac{(7b^2-16ac)\cot^2(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{3c^2(b^2-4ac)e} \\
& + \frac{2b\cot^3(d+ex)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{c(b^2-4ac)e} \\
& + \frac{(3b^2-8ac-2bc\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{c^2(b^2-4ac)e} \\
& - \frac{(105b^4-460ab^2c+256a^2c^2-2bc(35b^2-116ac)\cot(d+ex))\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}{24c^4(b^2-4ac)e}
\end{aligned}$$

[Out] $-3/2*b*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2})^{(1/2)}/c^{(5/2)}/e+5/16*b*(-12*a*c+7*b^2)*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d))/c^{(1/2)/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2})^{(1/2)}/c^{(9/2)}/e-2*(2*a+b*\cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2)^{(1/2)+2*\cot(e*x+d)^2*(2*a+b*\cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2)^{(1/2)-2*\cot(e*x+d)^4*(2*a+b*\cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2)^{(1/2)+2*(a*(b^2-2*(a-c)*c)+b*c*(a+c)*\cot(e*x+d))/(b^2+(a-c)^2)/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2)^{(1/2)-1/3*(-16*a*c+7*b^2)*\cot(e*x+d)^2*(a+b*\cot(e*x+d)+c*\cot(e*x+d))^2)^{(1/2)}/c^2/(-4*a*c+b^2)/e+2*b*\cot(e*x+d)^3*(a+b$

$$\begin{aligned}
& * \cot(e*x+d) + c * \cot(e*x+d)^2)^{1/2} / c / (-4*a*c + b^2) / e + (3*b^2 - 8*a*c - 2*b*c * \cot(e \\
& *x+d)) * (a + b * \cot(e*x+d) + c * \cot(e*x+d)^2)^{1/2} / c^2 / (-4*a*c + b^2) / e - 1/24 * (105*b \\
& ^4 - 460*a*b^2*c + 256*a^2*c^2 - 2*b*c * (-116*a*c + 35*b^2) * \cot(e*x+d)) * (a + b * \cot(e*x \\
& +d) + c * \cot(e*x+d)^2)^{1/2} / c^4 / (-4*a*c + b^2) / e - 1/2 * \operatorname{arctanh}(1/2 * (b^2 - (a-c) * (a- \\
& c - (a^2 - 2*a*c + b^2 + c^2)^{1/2})) - b * \cot(e*x+d) * (2*a - 2*c + (a^2 - 2*a*c + b^2 + c^2)^{1/2} \\
&)) * 2^{1/2} / (a + b * \cot(e*x+d) + c * \cot(e*x+d)^2)^{1/2} / (2*a - 2*c + (a^2 - 2*a*c + b^2 + c \\
& ^2)^{1/2})^{1/2} / (a^2 - b^2 - 2*a*c + c^2 - (a-c) * (a^2 - 2*a*c + b^2 + c^2)^{1/2})^{1/2}) \\
& * (2*a - 2*c + (a^2 - 2*a*c + b^2 + c^2)^{1/2})^{1/2} * (a^2 - b^2 - 2*a*c + c^2 - (a-c) * (a^2 - 2* \\
& a*c + b^2 + c^2)^{1/2})^{1/2} / (a^2 - 2*a*c + b^2 + c^2)^{3/2} / e * 2^{1/2} + 1/2 * \operatorname{arctanh}(1 \\
& /2 * (b^2 - b * \cot(e*x+d) * (2*a - 2*c - (a^2 - 2*a*c + b^2 + c^2)^{1/2}) - (a-c) * (a-c + (a^2 - 2* \\
& a*c + b^2 + c^2)^{1/2}))) * 2^{1/2} / (a + b * \cot(e*x+d) + c * \cot(e*x+d)^2)^{1/2} / (2*a - 2*c \\
& - (a^2 - 2*a*c + b^2 + c^2)^{1/2})^{1/2} / (a^2 - b^2 - 2*a*c + c^2 + (a-c) * (a^2 - 2*a*c + b^2 + c \\
& ^2)^{1/2})^{1/2}) * (2*a - 2*c - (a^2 - 2*a*c + b^2 + c^2)^{1/2})^{1/2} * (a^2 - b^2 - 2*a*c + \\
& c^2 + (a-c) * (a^2 - 2*a*c + b^2 + c^2)^{1/2})^{1/2} / (a^2 - 2*a*c + b^2 + c^2)^{3/2} / e * 2^{1 \\
& /2}
\end{aligned}$$

Rubi [A] (verified)

Time = 7.14 (sec) , antiderivative size = 1189, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules

used = {3782, 6857, 650, 752, 793, 635, 212, 846, 1032, 1050, 1044, 214}

$$\begin{aligned}
 & \int \frac{\cot^7(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \\
 & - \frac{2(2a+b\cot(d+ex))\cot^4(d+ex)}{(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} \\
 & + \frac{2b\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}\cot^3(d+ex)}{c(b^2-4ac)e} \\
 & - \frac{(7b^2-16ac)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}\cot^2(d+ex)}{3c^2(b^2-4ac)e} \\
 & + \frac{2(2a+b\cot(d+ex))\cot^2(d+ex)}{(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} \\
 & + \frac{5b(7b^2-12ac)\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{16c^{9/2}e} \\
 & - \frac{3b\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{2c^{5/2}e} \\
 & + \frac{\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2+(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e} \\
 & - \frac{\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2-(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e} \\
 & + \frac{(3b^2-2c\cot(d+ex)b-8ac)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{c^2(b^2-4ac)e} \\
 & - \frac{(105b^4-460acb^2-2c(35b^2-116ac)\cot(d+ex)b+256a^2c^2)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{24c^4(b^2-4ac)e} \\
 & - \frac{2(2a+b\cot(d+ex))}{(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} \\
 & + \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}
 \end{aligned}$$

[In] Int[Cot[d + e*x]^7/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2), x]

[Out] (-3*b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(2*c^(5/2)*e) + (5*b*(7*b^2 - 12*a*c)*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])])/(16*c^(9/2)*e) + (Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])]*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]])

$$\begin{aligned}
& 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (2*(2*a + b*Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) + (2*Cot[d + e*x]^2*(2*a + b*Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) - (2*Cot[d + e*x]^4*(2*a + b*Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) + (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) - ((7*b^2 - 16*a*c)*Cot[d + e*x]^2*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])/(3*c^2*(b^2 - 4*a*c)*e) + (2*b*Cot[d + e*x]^3*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])/(c*(b^2 - 4*a*c)*e) + ((3*b^2 - 8*a*c - 2*b*c*Cot[d + e*x])*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])/(c^2*(b^2 - 4*a*c)*e) - ((105*b^4 - 460*a*b^2*c + 256*a^2*c^2 - 2*b*c*(35*b^2 - 116*a*c)*Cot[d + e*x])*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])/(24*c^4*(b^2 - 4*a*c)*e)
\end{aligned}$$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 650

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(3/2), x_Symbol] := Simp[-2*((b*d - 2*a*e + (2*c*d - b*e)*x)/((b^2 - 4*a*c)*Sqrt[a + b*x + c*x^2])), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0]
```

Rule 752

```

Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x)^(m - 1)*(d*b - 2*a*e + (2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)), Int[(d + e*x)^(m - 2)*Simp[e*(2*a*e*(m - 1) + b*d*(2*p - m + 4)) - 2*c*d^2*(2*p + 3) + e*(b*e - 2*d*c)*(m + 2*p + 2)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && GtQ[m, 1] && IntQuadraticQ[a, b, c, d, e, m, p, x]

```

Rule 793

```

Int[((d_.) + (e_.)*(x_))*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(-b*e*g*(p + 2) - c*(e*f + d*g)*(2*p + 3) - 2*c*e*g*(p + 1)*x)*((a + b*x + c*x^2)^(p + 1)/(2*c^2*(p + 1)*(2*p + 3))), x] + Dist[(b^2*e*g*(p + 2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p + 3))/(2*c^2*(2*p + 3)), Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]

```

Rule 846

```

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[g*(d + e*x)^m*((a + b*x + c*x^2)^(p + 1)/(c*(m + 2*p + 2))), x] + Dist[1/(c*(m + 2*p + 2)), Int[(d + e*x)^(m - 1)*(a + b*x + c*x^2)^p*Simp[m*(c*d*f - a*e*g) + d*(2*c*f - b*g)*(p + 1) + (m*(c*e*f + c*d*g - b*e*g) + e*(p + 1)*(2*c*f - b*g))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[m, 0] && NeQ[m + 2*p + 2, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p]) && !(IGtQ[m, 0] && EqQ[f, 0])

```

Rule 1032

```

Int[((g_.) + (h_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_.) + (f_.)*(x_)^2)^(q_), x_Symbol]
:> Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2*f - c*(2*a*f)) - h*(b*c*d + a*b*f))*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Simp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))*(p + q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1))*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !(IntegerQ[p] && ILtQ[q, -1])

```

Rule 1044

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^7}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\ &= \\ &= -\frac{\text{Subst}\left(\int \left(\frac{x}{(a+bx+cx^2)^{3/2}} - \frac{x^3}{(a+bx+cx^2)^{3/2}} + \frac{x^5}{(a+bx+cx^2)^{3/2}} - \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot(d+ex)\right)}{e} \end{aligned}$$

$$\begin{aligned}
&= - \frac{\text{Subst}\left(\int \frac{x}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&+ \frac{\text{Subst}\left(\int \frac{x^3}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&- \frac{\text{Subst}\left(\int \frac{x^5}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&+ \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= - \frac{2(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&+ \frac{2 \cot^2(d+ex)(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&- \frac{2 \cot^4(d+ex)(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&+ \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&- \frac{2 \text{Subst}\left(\int \frac{x(4a+2bx)}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) e} \\
&+ \frac{2 \text{Subst}\left(\int \frac{x^3(8a+4bx)}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) e} \\
&- \frac{2 \text{Subst}\left(\int \frac{-\frac{1}{2}b(b^2-4ac) - \frac{1}{2}(a-c)(b^2-4ac)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 + (a-c)^2) (b^2 - 4ac) e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2 \cot^2(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&- \frac{2 \cot^4(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2b \cot^3(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c (b^2 - 4ac) e} \\
&+ \frac{(3b^2 - 8ac - 2bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c^2 (b^2 - 4ac) e} \\
&- \frac{(3b) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2c^2 e} \\
&+ \frac{\text{Subst}\left(\int \frac{x^2(-12ab-2(7b^2-16ac)x)}{\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{2c (b^2 - 4ac) e} \\
&+ \frac{\text{Subst}\left(\int \frac{\frac{1}{2}b(b^2-4ac)(2a-2c+\sqrt{a^2+b^2-2ac+c^2})+\frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&- \frac{\text{Subst}\left(\int \frac{\frac{1}{2}b(b^2-4ac)(2a-2c-\sqrt{a^2+b^2-2ac+c^2})+\frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d + ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2 \cot^2(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&- \frac{2 \cot^4(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&- \frac{(7b^2 - 16ac) \cot^2(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{3c^2 (b^2 - 4ac) e} \\
&+ \frac{2b \cot^3(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c (b^2 - 4ac) e} \\
&+ \frac{(3b^2 - 8ac - 2bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c^2 (b^2 - 4ac) e} \\
&- \frac{(3b) \text{Subst} \left(\int \frac{1}{4c - x^2} dx, x, \frac{b + 2c \cot(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \right)}{c^2 e} \\
&+ \frac{\text{Subst} \left(\int \frac{x(4a(7b^2 - 16ac) + b(35b^2 - 116ac)x)}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{6c^2 (b^2 - 4ac) e} \\
&\frac{(b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Sub}}{+} \\
&\frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Sub}}{+}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{5/2}e} \\
&+ \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{\sqrt{2}(a^2+b^2-2ac+c^2)}{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2})}}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{2(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&+ \frac{2 \cot^2(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&- \frac{2 \cot^4(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&+ \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&- \frac{(7b^2-16ac) \cot^2(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{3c^2(b^2-4ac)e} \\
&+ \frac{2b \cot^3(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c(b^2-4ac)e} \\
&+ \frac{(3b^2-8ac-2bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c^2(b^2-4ac)e} \\
&- \frac{(105b^4-460ab^2c+256a^2c^2-2bc(35b^2-116ac) \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{24c^4(b^2-4ac)e} \\
&+ \frac{(5b(7b^2-12ac)) \operatorname{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{16c^4e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{3b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{5/2}e} \\
&+ \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{-\sqrt{2}}{\sqrt{2}(a^2+b^2-2ac+c^2)}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2})}}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{2(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&+ \frac{2 \cot^2(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&- \frac{2 \cot^4(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&+ \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&- \frac{(7b^2-16ac) \cot^2(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{3c^2(b^2-4ac)e} \\
&+ \frac{2b \cot^3(d+ex)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c(b^2-4ac)e} \\
&+ \frac{(3b^2-8ac-2bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c^2(b^2-4ac)e} \\
&- \frac{(105b^4-460ab^2c+256a^2c^2-2bc(35b^2-116ac) \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{24c^4(b^2-4ac)e} \\
&+ \frac{(5b(7b^2-12ac)) \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8c^4e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{5/2}e} \\
&+ \frac{5b(7b^2 - 12ac) \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{16c^{9/2}e} \\
&+ \frac{\sqrt{2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 - 2ac + c^2 + (a - c)\sqrt{a^2 + b^2 - 2ac + c^2}} \operatorname{arctanh}\left(\frac{\sqrt{2}(a^2 + b^2 - 2ac + c^2)}{\sqrt{2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})}}\right)}{\sqrt{2}(a^2 + b^2 - 2ac + c^2)} \\
&- \frac{\sqrt{2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 + c(c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a(2c + \sqrt{a^2 + b^2 - 2ac + c^2})}}{\sqrt{2}(a^2 + b^2 - 2ac + c^2)} \\
&- \frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2 \cot^2(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&- \frac{2 \cot^4(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&- \frac{(7b^2 - 16ac) \cot^2(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{3c^2 (b^2 - 4ac) e} \\
&+ \frac{2b \cot^3(d + ex) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c (b^2 - 4ac) e} \\
&+ \frac{(3b^2 - 8ac - 2bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c^2 (b^2 - 4ac) e} \\
&- \frac{(105b^4 - 460ab^2c + 256a^2c^2 - 2bc(35b^2 - 116ac) \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{24c^4 (b^2 - 4ac) e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 7.24 (sec) , antiderivative size = 2097, normalized size of antiderivative = 1.76

$$\int \frac{\cot^7(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Result too large to show}$$

[In] Integrate[Cot[d + e*x]^7/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] (4*Cot[d + e*x]*(b + 2*a*Tan[d + e*x])*(-(a*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2 - 4*a*c)))^(3/2)/(a*e*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))

$$\begin{aligned}
& ^2) * \text{Sqrt}[\text{Cot}[d + e*x]^2 * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2)] * \text{Sqrt}[1 - (\\
& b^2 - 4*a*c) * (b / (b^2 - 4*a*c) + (2*a * \text{Tan}[d + e*x]) / (b^2 - 4*a*c))^2] - (\text{Co} \\
& t[d + e*x] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2] * ((4 * \text{Cot}[d + e*x] * (b^ \\
& 2 - 2*a*c + a * b * \text{Tan}[d + e*x])) / (c * (b^2 - 4*a*c) * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a \\
& * \text{Tan}[d + e*x]^2]) + ((3 * b * (b^2 - 4*a*c) * \text{ArcTanh}[(2*c + b * \text{Tan}[d + e*x]) / (2 * \text{S} \\
& \text{qrt}[c] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2])]) / c^{(3/2)} - (2 * (3 * b^2 - \\
& 8 * a * c) * \text{Cot}[d + e*x] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) / c) / (c * (b^ \\
& 2 - 4 * a * c))) / (2 * e * \text{Sqrt}[\text{Cot}[d + e*x]^2 * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x] \\
& ^2)]) - (\text{Cot}[d + e*x] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2] * ((-2 * \text{Tan}[\\
& d + e*x]^3 * (-b^2 + 2 * a * c - a * b * \text{Tan}[d + e*x])) / (c * (b^2 - 4 * a * c) * \text{Sqrt}[c + b * \text{T} \\
& \text{an}[d + e*x] + a * \text{Tan}[d + e*x]^2]) - (2 * (b * \text{Tan}[d + e*x]^2 * \text{Sqrt}[c + b * \text{Tan}[d + \\
& e*x] + a * \text{Tan}[d + e*x]^2] + (((-6 * a^2 * b^2 * c + 24 * a^3 * c^2) * \text{ArcTanh}[(b + 2 * a * \text{T} \\
& \text{an}[d + e*x]) / (2 * \text{Sqrt}[a] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2])]) / (4 * a \\
& ^{(5/2)}) + ((6 * a^2 * b * c - 12 * a^3 * c * \text{Tan}[d + e*x]) * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \\
& \text{Tan}[d + e*x]^2]) / (2 * a^2)) / (3 * a))) / (c * (b^2 - 4 * a * c))) / (e * \text{Sqrt}[\text{Cot}[d + e*x]^ \\
& 2 * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2)]) + (\text{Cot}[d + e*x] * \text{Sqrt}[c + b * \text{Tan}[\\
& d + e*x] + a * \text{Tan}[d + e*x]^2] * ((2 * ((-4 * \text{Sqrt}[a - I * b - c] * (-1/4 * (b * (b^2 - 4 * a \\
& * c)) + (I/4) * (a - c) * (b^2 - 4 * a * c)) * \text{ArcTan}[(I * b + 2 * c - ((-2 * I) * a - b) * \text{Tan}[\\
& d + e*x]) / (2 * \text{Sqrt}[a - I * b - c] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) \\
&]) / (-4 * a + (4 * I) * b + 4 * c) - (4 * \text{Sqrt}[a + I * b - c] * (-1/4 * (b * (b^2 - 4 * a * c)) - \\
& (I/4) * (a - c) * (b^2 - 4 * a * c)) * \text{ArcTan}[((-I) * b + 2 * c - ((2 * I) * a - b) * \text{Tan}[d + e \\
& * x]) / (2 * \text{Sqrt}[a + I * b - c] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2])]) / (- \\
& 4 * a - (4 * I) * b + 4 * c))) / ((b^2 + (a - c)^2) * (b^2 - 4 * a * c)) - (2 * \text{Tan}[d + e*x]^ \\
& 3 * (-b^2 + 2 * a * c - a * b * \text{Tan}[d + e*x])) / (c * (b^2 - 4 * a * c) * \text{Sqrt}[c + b * \text{Tan}[d + e \\
& x] + a * \text{Tan}[d + e*x]^2]) - (2 * (b^3 + a * b * (a - 3 * c) + a * (2 * a^2 + b^2 - 2 * a * c) \\
& * \text{Tan}[d + e*x])) / ((b^2 + (a - c)^2) * (b^2 - 4 * a * c) * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + \\
& a * \text{Tan}[d + e*x]^2]) + (4 * (b^2 - 4 * a * c) * (a^2 / ((b^2 - 4 * a * c) * ((a^2 * b^2) / (b^2 - \\
& 4 * a * c)^2 - (4 * a^3 * c) / (b^2 - 4 * a * c)^2)))^{(3/2)} * (-((a * b) / (b^2 - 4 * a * c)) - (2 \\
& * a^2 * \text{Tan}[d + e*x]) / (b^2 - 4 * a * c)) * (-((a * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x] \\
& ^2)) / (b^2 - 4 * a * c)))^{(3/2)} / (a^2 * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2)^{ \\
& (3/2)} * \text{Sqrt}[1 - (-((a * b) / (b^2 - 4 * a * c)) - (2 * a^2 * \text{Tan}[d + e*x]) / (b^2 - 4 * a * c)) \\
& ^2 / ((a^2 * b^2) / (b^2 - 4 * a * c)^2 - (4 * a^3 * c) / (b^2 - 4 * a * c)^2)]) - (2 * (b * \text{Tan}[d \\
& + e*x]^2 * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2] + (((-6 * a^2 * b^2 * c + 24 \\
& * a^3 * c^2) * \text{ArcTanh}[(b + 2 * a * \text{Tan}[d + e*x]) / (2 * \text{Sqrt}[a] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] \\
& + a * \text{Tan}[d + e*x]^2])]) / (4 * a^{(5/2)}) + ((6 * a^2 * b * c - 12 * a^3 * c * \text{Tan}[d + e*x]) * \\
& \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) / (2 * a^2)) / (3 * a))) / (c * (b^2 - 4 * a \\
& * c))) / (e * \text{Sqrt}[\text{Cot}[d + e*x]^2 * (c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2)]) + (\\
& \text{Cot}[d + e*x] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2] * ((-2 * \text{Cot}[d + e*x]^ \\
& 3 * (-b^2 + 2 * a * c - a * b * \text{Tan}[d + e*x])) / (c * (b^2 - 4 * a * c) * \text{Sqrt}[c + b * \text{Tan}[d + e \\
& x] + a * \text{Tan}[d + e*x]^2]) - (2 * (-1/6 * ((-7 * b^2 + 16 * a * c) * \text{Cot}[d + e*x]^3 * \text{Sqrt}[c \\
& + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) / c - ((b * (35 * b^2 - 116 * a * c) * \text{Cot}[d + e \\
& * x]^2 * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) / (8 * c) - (((a * b * c * (35 * b^ \\
& 2 - 116 * a * c)) / 2 + (b * (-105 * b^4 + 460 * a * b^2 * c - 256 * a^2 * c^2)) / 8) * \text{ArcTanh}[(2 * \\
& c + b * \text{Tan}[d + e*x]) / (2 * \text{Sqrt}[c] * \text{Sqrt}[c + b * \text{Tan}[d + e*x] + a * \text{Tan}[d + e*x]^2]) \\
&]) / (2 * c^{(3/2)}) + ((105 * b^4 - 460 * a * b^2 * c + 256 * a^2 * c^2) * \text{Cot}[d + e*x] * \text{Sqrt}[c
\end{aligned}$$

+ b*Tan[d + e*x] + a*Tan[d + e*x]^2])/(8*c))/(2*c))/(3*c)))/(c*(b^2 - 4*a*c)))/(e*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.57 (sec) , antiderivative size = 13067599, normalized size of antiderivative = 10990.41

output too large to display

[In] int(cot(e*x+d)^7/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 44563 vs. 2(1097) = 2194.

Time = 35.38 (sec) , antiderivative size = 89177, normalized size of antiderivative = 75.00

$$\int \frac{\cot^7(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot^7(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\cot^7(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(cot(e*x+d)**7/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2),x)

[Out] Integral(cot(d + e*x)**7/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)

Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^7(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="maxima")

[Out] Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^7(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^7(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Hanged}$$

[In] int(cot(d + e*x)^7/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2),x)

[Out] \text{Hanged}

$$3.12 \quad \int \frac{\cot^5(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	137
Rubi [A] (verified)	138
Mathematica [C] (verified)	143
Maple [B] (warning: unable to verify)	145
Fricas [B] (verification not implemented)	146
Sympy [F]	146
Maxima [F(-1)]	146
Giac [F(-2)]	146
Mupad [F(-1)]	147

Optimal result

Integrand size = 33, antiderivative size = 865

$$\int \frac{\cot^5(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx = \frac{3b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{5/2}e}$$

$$- \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}+(a-c)\sqrt{a^2+b^2-2ac+c^2} \operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$+ \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}-(a-c)\sqrt{a^2+b^2-2ac+c^2} \operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$+ \frac{2(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \frac{2 \cot^2(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$- \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$- \frac{(3b^2-8ac-2bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c^2(b^2-4ac)e}$$

```
[Out] 3/2*b*arctanh(1/2*(b+2*c*cot(e*x+d))/c^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2))/c^(5/2)/e+2*(2*a+b*cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)-2*cot(e*x+d)^2*(2*a+b*cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)-2*(a*(b^2-2*(a-c)*c)+b*c*(a+c)*cot(e*x+d))/(b^2+(a-c)^2)/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)-(3*b^2-8*a*c-2*b*c*cot(e*x+d))*(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/c^2/(-4*a*c+b^2)
```

$$\begin{aligned} & /e^{1/2} \operatorname{arctanh}\left(\frac{1}{2}(b^2 - (a-c)(a-c - (a^2 - 2ac + b^2 + c^2)^{1/2}) - b \cot(ex+d) * \right. \\ & (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2}))^{1/2} / (a + b \cot(ex+d) + c \cot(ex+d)^2 \\ &)^{1/2} / (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - b^2 - 2ac + c^2 - (a-c) * \\ & (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} * (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} \\ & * (a^2 - b^2 - 2ac + c^2 - (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - 2ac + b^2 + c^2 \\ &)^{3/2} / e^{1/2} - 1/2 \operatorname{arctanh}\left(\frac{1}{2}(b^2 - b \cot(ex+d) * (2a - 2c - (a^2 - 2ac + b^2 + c^2)^{1/2}) - \right. \\ & (a-c)(a-c + (a^2 - 2ac + b^2 + c^2)^{1/2}))^{1/2} / (a + b \cot(ex+d) + c \cot(ex+d)^2 \\ &)^{1/2} / (2a - 2c - (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - b^2 - 2ac + c^2 + (a-c) * \\ & (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} * (2a - 2c - (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} \\ & * (a^2 - b^2 - 2ac + c^2 + (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - 2ac + b^2 + c^2)^{3/2} / e^{1/2} \end{aligned}$$

Rubi [A] (verified)

Time = 5.23 (sec) , antiderivative size = 865, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3782, 6857, 650, 752, 793, 635, 212, 1032, 1050, 1044, 214}

$$\begin{aligned} & \int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \\ & \frac{2(2a+b\cot(d+ex))\cot^2(d+ex)}{(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} + \frac{3b\operatorname{arctanh}\left(\frac{b+2c\cot(d+ex)}{2\sqrt{c}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{2c^{5/2}e} \\ & - \frac{\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2+(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}\sqrt{a^2-2ca+b^2+c^2}}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e} \\ & + \frac{\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2-(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}\sqrt{a^2-2ca+b^2+c^2}}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e} \\ & - \frac{(3b^2-2c\cot(d+ex)b-8ac)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}{c^2(b^2-4ac)e} \\ & + \frac{2(2a+b\cot(d+ex))}{(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} \\ & - \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}} \end{aligned}$$

[In] Int[Cot[d + e*x]^5/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] (3*b*ArcTanh[(b + 2*c*Cot[d + e*x])/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2))]/(2*c^(5/2)*e) - (Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2

```
*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*
Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])
]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) + (Sqrt[2*a - 2*c + Sqrt[a^2
+ b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^
2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c +
c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x])/(Sqrt[2
]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c +
c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Co
t[d + e*x]^2])))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) + (2*(2*a + b*
Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]
) - (2*Cot[d + e*x]^2*(2*a + b*Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*C
ot[d + e*x] + c*Cot[d + e*x]^2]) - (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*
Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] +
c*Cot[d + e*x]^2]) - ((3*b^2 - 8*a*c - 2*b*c*Cot[d + e*x])*Sqrt[a + b*Cot[
d + e*x] + c*Cot[d + e*x]^2])/(c^2*(b^2 - 4*a*c)*e)
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 635

```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[In
t[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a,
b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 650

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(3/2), x_Symbo
l] := Simp[-2*((b*d - 2*a*e + (2*c*d - b*e)*x)/((b^2 - 4*a*c)*Sqrt[a + b*x
+ c*x^2])), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b
^2 - 4*a*c, 0]
```

Rule 752

```
Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_S
ymbol] := Simp[(d + e*x)^(m - 1)*(d*b - 2*a*e + (2*c*d - b*e)*x)*((a + b*x
+ c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Dist[1/((p + 1)*(b^2 - 4*a*
c)), Int[(d + e*x)^(m - 2)*Simp[e*(2*a*e*(m - 1) + b*d*(2*p - m + 4)) - 2*c
*d^2*(2*p + 3) + e*(b*e - 2*d*c)*(m + 2*p + 2)*x, x]*(a + b*x + c*x^2)^(p +
```

1), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && GtQ[m, 1] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 793

Int[((d_.) + (e_.)*(x_))*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(-b*e*g*(p + 2) - c*(e*f + d*g)*(2*p + 3) - 2*c*e*g*(p + 1)*x)*((a + b*x + c*x^2)^(p + 1)/(2*c^2*(p + 1)*(2*p + 3))), x] + Dist[(b^2*e*g*(p + 2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p + 3))/(2*c^2*(2*p + 3)), Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]

Rule 1032

Int[((g_.) + (h_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_.) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2*f - c*(2*a*f)) - h*(b*c*d + a*b*f))*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Simp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))*(p + q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1)))*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !(!IntegerQ[p] && ILtQ[q, -1])

Rule 1044

Int[((g_.) + (h_.)*(x_))/(((a_.) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_.) + (h_.)*(x_))/(((a_.) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] &&

NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 3782

Int[cot[(d_.) + (e_.)*(x_.)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n2_.))^(p_), x_Symbol] :> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \left(-\frac{x}{(a+bx+cx^2)^{3/2}} + \frac{x^3}{(a+bx+cx^2)^{3/2}} + \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot(d+ex)\right)}{e} \\
 &= \frac{\text{Subst}\left(\int \frac{x}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{x^3}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &= \frac{2(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
 &\quad - \frac{2 \cot^2(d+ex)(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
 &\quad - \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
 &\quad + \frac{2 \text{Subst}\left(\int \frac{x(4a+2bx)}{\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) e} \\
 &\quad + \frac{2 \text{Subst}\left(\int \frac{-\frac{1}{2}b(b^2-4ac) - \frac{1}{2}(a-c)(b^2-4ac)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 + (a-c)^2) (b^2 - 4ac) e}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2 \cot^2(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{(3b^2 - 8ac - 2bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c^2 (b^2 - 4ac) e} \\
&\quad + \frac{(3b) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{2c^2 e} \\
&\quad - \frac{\text{Subst} \left(\int \frac{\frac{1}{2} b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2} (b^2 - 4ac) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2})) x}{(1 + x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&\quad - \frac{\text{Subst} \left(\int \frac{\frac{1}{2} b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2} (b^2 - 4ac) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2})) x}{(1 + x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&= \frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2 \cot^2(d + ex)(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{(3b^2 - 8ac - 2bc \cot(d + ex)) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}{c^2 (b^2 - 4ac) e} \\
&\quad + \frac{(3b) \text{Subst} \left(\int \frac{1}{4c - x^2} dx, x, \frac{b + 2c \cot(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \right)}{c^2 e} \\
&\quad + \frac{(b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&\quad + \frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3b \operatorname{arctanh}\left(\frac{b+2c \cot(d+ex)}{2\sqrt{c}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{2c^{5/2}e} \\
&\quad - \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}} \operatorname{arctanh}\left(\frac{\sqrt{2}(a^2+b^2-2ac+c^2)}{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&\quad + \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})}-a(2c+\sqrt{a^2+b^2-2ac+c^2})}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&\quad + \frac{2(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad - \frac{2 \cot^2(d+ex)(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad - \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad - \frac{(3b^2-8ac-2bc \cot(d+ex))\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}{c^2(b^2-4ac)e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.91 (sec) , antiderivative size = 1674, normalized size of antiderivative = 1.94

$$\int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx =$$

$$\frac{4\cot(d+ex)(b+2a\tan(d+ex))\left(-\frac{a(c+b\tan(d+ex)+a\tan^2(d+ex))}{b^2-4ac}\right)^{3/2}}{ae(c+b\tan(d+ex)+a\tan^2(d+ex))\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}\sqrt{1-(b^2-4ac)}}$$

$$+ \frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}\left(\frac{4\cot(d+ex)(b^2-2ac+ab\tan(d+ex))}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} + \frac{3b(b^2-4ac)\operatorname{arctanh}\left(\frac{1}{2\sqrt{c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right)}{c^{3/2}}\right)}{2e\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}$$

$$+ \frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}\left(\frac{2\tan^3(d+ex)(-b^2+2ac-ab\tan(d+ex))}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} - \frac{2\left(b\tan^2(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}\right)}{c^{3/2}}\right)}{e\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}$$

$$+ \frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}\left(\frac{4\sqrt{a-ib-c}\left(-\frac{1}{4}b(b^2-4ac)+\frac{1}{4}i(a-c)(b^2-4ac)\right)\operatorname{arctan}\left(\frac{ib+2c-(-2ia-c)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right)}{-4a+4ib+4c}\right)}{c^{3/2}}$$

```
[In] Integrate[Cot[d + e*x]^5/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]
[Out] (-4*Cot[d + e*x]*(b + 2*a*Tan[d + e*x])*(-(a*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2 - 4*a*c)))^(3/2)/(a*e*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]*Sqrt[1 - (b^2 - 4*a*c)*(b/(b^2 - 4*a*c) + (2*a*Tan[d + e*x])/(b^2 - 4*a*c))^2]) + (Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((4*Cot[d + e*x]*(b^2 - 2*a*c + a*b*Tan[d + e*x]))/(c*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) + ((3*b*(b^2 - 4*a*c)*ArcTanh[(2*c + b*Tan[d + e*x])/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]))/c^(3/2) - (2*(3*b^2 - 8*a*c)*Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])/c)/(c*(b
```


$$\begin{aligned}
&^2 - 4ac)))/(2e\sqrt{\cot(d+ex)^2(c+b\tan(d+ex)+a\tan(d+ex)^2)} + (\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}*((-2\tan(d+ex)^3(-b^2+2ac-a*b\tan(d+ex)))/(c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}) - (2(b\tan(d+ex)^2\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2} + ((-6a^2b^2c+24a^3c^2)\operatorname{ArcTanh}[(b+2a\tan(d+ex))/(2\sqrt{a}\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2})]))/(4a^{5/2}) + ((6a^2b^2c-12a^3c^2)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2})/(2a^2))/(3a)))/(c(b^2-4ac)))/(e\sqrt{\cot(d+ex)^2(c+b\tan(d+ex)+a\tan(d+ex)^2)} - (\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}*((2((-4\sqrt{a-Ib-c})(-1/4(b(b^2-4ac)) + (I/4)(a-c)(b^2-4ac))\operatorname{ArcTan}[(Ib+2c-((-2I)a-b)\tan(d+ex)])/(2\sqrt{a-Ib-c}\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}])))/(-4a+(4I)b+4c) - (4\sqrt{a+Ib-c})(-1/4(b(b^2-4ac)) - (I/4)(a-c)(b^2-4ac))\operatorname{ArcTan}[(Ib+2c-((2I)a-b)\tan(d+ex)]/(2\sqrt{a+Ib-c}\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}))))/(-4a-(4I)b+4c))/((b^2+(a-c)^2)(b^2-4ac)) - (2\tan(d+ex)^3(-b^2+2ac-a*b\tan(d+ex)))/(c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}) - (2(b^3+a*b(a-3c)+a(2a^2+b^2-2ac))\tan(d+ex))/((b^2+(a-c)^2)(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2}) + (4(b^2-4ac)(a^2/((b^2-4ac)((a^2b^2)/(b^2-4ac)^2 - (4a^3c)/(b^2-4ac)^2)))^{3/2}*(-((a*b)/(b^2-4ac)) - (2a^2\tan(d+ex))/(b^2-4ac)))*(-(a(c+b\tan(d+ex)+a\tan(d+ex)^2))/(b^2-4ac)))^{3/2})/(a^2(c+b\tan(d+ex)+a\tan(d+ex)^2)^{3/2}\sqrt{1-((a*b)/(b^2-4ac)) - (2a^2\tan(d+ex))/(b^2-4ac)})^2/((a^2b^2)/(b^2-4ac)^2 - (4a^3c)/(b^2-4ac)^2)) - (2(b\tan(d+ex)^2\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2} + ((-6a^2b^2c+24a^3c^2)\operatorname{ArcTanh}[(b+2a\tan(d+ex))/(2\sqrt{a}\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2})]))/(4a^{5/2}) + ((6a^2b^2c-12a^3c^2)\sqrt{c+b\tan(d+ex)+a\tan(d+ex)^2})/(2a^2))/(3a)))/(c(b^2-4ac)))/(e\sqrt{\cot(d+ex)^2(c+b\tan(d+ex)+a\tan(d+ex)^2)}
\end{aligned}$$

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 2.91 (sec) , antiderivative size = 13067692, normalized size of antiderivative = 15107.16

output too large to display

[In] $\int \cot(ex+d)^5/(a+b\cot(ex+d)+c\cot(ex+d)^2)^{3/2}, x$

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42426 vs. 2(793) = 1586.
Time = 29.83 (sec) , antiderivative size = 84903, normalized size of antiderivative = 98.15

$$\int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx$$

[In] integrate(cot(e*x+d)**5/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2),x)

[Out] Integral(cot(d + e*x)**5/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)

Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="maxima")

[Out] Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^5(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^5(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \int \frac{\cot(d + ex)^5}{(c \cot(d + ex)^2 + b \cot(d + ex) + a)^{3/2}} dx$$

```
[In] int(cot(d + e*x)^5/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)
```

```
[Out] int(cot(d + e*x)^5/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)
```

$$3.13 \quad \int \frac{\cot^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	148
Rubi [A] (verified)	149
Mathematica [C] (verified)	152
Maple [B] (warning: unable to verify)	154
Fricas [B] (verification not implemented)	154
Sympy [F]	155
Maxima [F(-2)]	155
Giac [F(-2)]	155
Mupad [F(-1)]	156

Optimal result

Integrand size = 33, antiderivative size = 686

$$\int \frac{\cot^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx = \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}})^2} \arctanh\left(\frac{b^2-\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}\sqrt{a^2+b^2-2ac+c^2}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e} - \frac{2(2a+b \cot(d+ex))}{(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} + \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

```
[Out] -2*(2*a+b*cot(e*x+d))/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)+
2*(a*(b^2-2*(a-c)*c)+b*c*(a+c)*cot(e*x+d))/(b^2+(a-c)^2)/(-4*a*c+b^2)/e/(a+
b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)-1/2*arctanh(1/2*(b^2-(a-c)*(a-c-(a^2-2*a
*c+b^2+c^2)^(1/2))-b*cot(e*x+d)*(2*a-2*c+(a^2-2*a*c+b^2+c^2)^(1/2))))*2^(1/2
)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(2*a-2*c+(a^2-2*a*c+b^2+c^2)^(1/2))
^(1/2)/(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)*(2*a-2*c+
(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)*(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c^
2)^(1/2))^(1/2)/(a^2-2*a*c+b^2+c^2)^(3/2)/e*2^(1/2)+1/2*arctanh(1/2*(b^2-b*
cot(e*x+d)*(2*a-2*c-(a^2-2*a*c+b^2+c^2)^(1/2))-(a-c)*(a-c+(a^2-2*a*c+b^2+c^
2)^(1/2))))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(2*a-2*c-(a^2-2*a*
c+b^2+c^2)^(1/2))^(1/2)/(a^2-b^2-2*a*c+c^2+(a-c)*(a^2-2*a*c+b^2+c^2)^(1/2))
^(1/2)*(2*a-2*c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)*(a^2-b^2-2*a*c+c^2+(a-c)*
(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/(a^2-2*a*c+b^2+c^2)^(3/2)/e*2^(1/2)
```

Rubi [A] (verified)

Time = 5.09 (sec) , antiderivative size = 686, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3782, 6857, 650, 1032, 1050, 1044, 214}

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+2a-2c}\sqrt{(a-c)\sqrt{a^2-2ac+b^2}}}{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+2a-2c}\sqrt{-(a-c)\sqrt{a^2-2ac+b^2+c^2}+a^2-2ac-b^2+c^2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+2a-2c}}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+2a-2c}}\right)} + \frac{2(2a+b\cot(d+ex))}{e(b^2-4ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} + \frac{2(a(b^2-2c(a-c))+bc(a+c)\cot(d+ex))}{e((a-c)^2+b^2)(b^2-4ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}$$

[In] Int[Cot[d + e*x]^3/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] (Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]))/(Sqrt[2]*Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]])*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Cot[d + e*x]]/(Sqrt[2]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]])*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (2*(2*a + b*Cot[d + e*x]))/((b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) + (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 650

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2)^(3/2), x_Symbol] := Simp[-2*((b*d - 2*a*e + (2*c*d - b*e)*x)/((b^2 - 4*a*c)*Sqrt[a + b*x

+ c*x^2)), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0]

Rule 1032

Int[((g_.) + (h_.)*(x_))*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2*f - c*(2*a*f)) - h*(b*c*d + a*b*f))*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Simp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))*(p + q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1)))*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !(!IntegerQ[p] && !LtQ[q, -1])

Rule 1044

Int[((g_) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_.) + (h_.)*(x_))/(((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] && NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 3782

Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n^2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xexpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^3}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \left(\frac{x}{(a+bx+cx^2)^{3/2}} - \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x}{(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{2(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad - \frac{2 \text{Subst}\left(\int \frac{-\frac{1}{2}b(b^2-4ac) - \frac{1}{2}(a-c)(b^2-4ac)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 + (a-c)^2) (b^2 - 4ac) e} \\
&= -\frac{2(2a + b \cot(d+ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{1}{2}b(b^2-4ac)(2a-2c+\sqrt{a^2+b^2-2ac+c^2}) + \frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{\frac{1}{2}b(b^2-4ac)(2a-2c-\sqrt{a^2+b^2-2ac+c^2}) + \frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad (b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{ Sub} \\
&- \frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{ Sub}}{+} \\
&= \frac{\sqrt{2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 - 2ac + c^2 + (a - c) \sqrt{a^2 + b^2 - 2ac + c^2}} \operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{2}}{\sqrt{2}\sqrt{2}}\right)}{\sqrt{2} (a^2 + b^2 - 2ac + c^2)^{3/2}} \\
&- \frac{\sqrt{2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 + c (c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2})}}{\sqrt{2} (a^2 + b^2 - 2ac + c^2)^{3/2}} \\
&- \frac{2(2a + b \cot(d + ex))}{(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.68 (sec) , antiderivative size = 1419, normalized size of antiderivative = 2.07

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{4\cot(d+ex)(b+2a\tan(d+ex))}{ae(c+b\tan(d+ex)+a\tan^2(d+ex))\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}} + \frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} - \frac{2\tan^3(d+ex)(-b^2+2ac-ab\tan(d+ex))}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} - \frac{2\left(b\tan^2(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}\right)}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} + \frac{e\sqrt{\cot^2(d+ex)(c+b\tan(d+ex)+a\tan^2(d+ex))}}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} + \frac{2\left(\frac{4\sqrt{a-ib-c}\left(-\frac{1}{4}b(b^2-4ac)+\frac{1}{4}i(a-c)(b^2-4ac)\right)\arctan\left(\frac{ib+2c-(-2ia)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}\right)}{-4a+4ib+4c}\right)}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

```
[In] Integrate[Cot[d + e*x]^3/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]
[Out] (4*Cot[d + e*x]*(b + 2*a*Tan[d + e*x])*(-(a*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2 - 4*a*c)))^(3/2))/(a*e*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]*Sqrt[1 - (b^2 - 4*a*c)*(b/(b^2 - 4*a*c) + (2*a*Tan[d + e*x])/(b^2 - 4*a*c))^2]) - (Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((-2*Tan[d + e*x]^3*(-b^2 + 2*a*c - a*b*Tan[d + e*x]))/(c*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - (2*(b*Tan[d + e*x]^2*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2) + (((-6*a^2*b^2*c + 24*a^3*c^2)*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])))/(4*a^(5/2)) + ((6*a^2*b*c - 12*a^3*c*Tan[d + e*x])*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])/(2*a^2))/(3*a)))/(c*(b^2 - 4*a*c)))/(e*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)]) + (Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((2*((-4*Sqrt[a - I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) + (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[(I*b + 2*c - ((-2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]))/(-4*a + (4*I)*b + 4*c) - (4*Sqrt[a + I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) - (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[(-I)*b + 2*c - ((2*I)*a - b)*Tan[d + e*x])/(2*Sqr
```

```
t[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]])/(-4*a - (4*I
*b + 4*c))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)) - (2*Tan[d + e*x]^3*(-b^2 + 2
*a*c - a*b*Tan[d + e*x]))/(c*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[
d + e*x]^2]) - (2*(b^3 + a*b*(a - 3*c) + a*(2*a^2 + b^2 - 2*a*c)*Tan[d + e*
x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e
*x]^2]) + (4*(b^2 - 4*a*c)*(a^2/((b^2 - 4*a*c)*((a^2*b^2)/(b^2 - 4*a*c)^2 -
(4*a^3*c)/(b^2 - 4*a*c)^2)))^(3/2)*(-(a*b)/(b^2 - 4*a*c)) - (2*a^2*Tan[d
+ e*x])/(b^2 - 4*a*c))*(-(a*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2
- 4*a*c)))^(3/2))/(a^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)^(3/2)*Sqrt[1
- (-(a*b)/(b^2 - 4*a*c)) - (2*a^2*Tan[d + e*x])/(b^2 - 4*a*c))^2/((a^2*b^
2)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2 - 4*a*c)^2)]) - (2*(b*Tan[d + e*x]^2*Sq
rt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2] + (((-6*a^2*b^2*c + 24*a^3*c^2)*A
rcTanh[(b + 2*a*Tan[d + e*x])/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d
+ e*x]^2)])))/(4*a^(5/2)) + ((6*a^2*b*c - 12*a^3*c*Tan[d + e*x])*Sqrt[c + b*
Tan[d + e*x] + a*Tan[d + e*x]^2])/(2*a^2))/(3*a)))/(c*(b^2 - 4*a*c)))/(e*S
qrt[Cot[d + e*x]^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)])
```

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.53 (sec) , antiderivative size = 13067316, normalized size of antiderivative = 19048.57

output too large to display

```
[In] int(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)
```

```
[Out] result too large to display
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21092 vs. 2(629) = 1258.

Time = 12.13 (sec) , antiderivative size = 21092, normalized size of antiderivative = 30.75

$$\int \frac{\cot^3(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Too large to display}$$

```
[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="
fricas")
```

```
[Out] Too large to include
```

Sympy [F]

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(cot(e*x+d)**3/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2),x)

[Out] Integral(cot(d + e*x)**3/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see 'assume?' for more details)

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)^3}{(c\cot(d+ex)^2+b\cot(d+ex)+a)^{3/2}} dx$$

```
[In] int(cot(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)
```

```
[Out] int(cot(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)
```

$$3.14 \quad \int \frac{\cot(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	157
Rubi [A] (verified)	158
Mathematica [C] (verified)	161
Maple [B] (warning: unable to verify)	161
Fricas [B] (verification not implemented)	162
Sympy [F]	162
Maxima [F(-2)]	162
Giac [F(-2)]	163
Mupad [F(-1)]	163

Optimal result

Integrand size = 31, antiderivative size = 635

$$\int \frac{\cot(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx =$$

$$\frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$+ \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2-(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{b}{\sqrt{2}\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$- \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

```
[Out] -2*(a*(b^2-2*(a-c)*c)+b*c*(a+c)*cot(e*x+d))/(b^2+(a-c)^2)/(-4*a*c+b^2)/e/(a
+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)+1/2*arctanh(1/2*(b^2-(a-c)*(a-c-(a^2-2*
a*c+b^2+c^2)^(1/2))-b*cot(e*x+d)*(2*a-2*c+(a^2-2*a*c+b^2+c^2)^(1/2))))*2^(1/
2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(2*a-2*c+(a^2-2*a*c+b^2+c^2)^(1/2)
)^(1/2)/(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2))*(2*a-2*c
+(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)*(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c
^2)^(1/2))^(1/2)/(a^2-2*a*c+b^2+c^2)^(3/2)/e*2^(1/2)-1/2*arctanh(1/2*(b^2-b
*cot(e*x+d)*(2*a-2*c-(a^2-2*a*c+b^2+c^2)^(1/2))-(a-c)*(a-c+(a^2-2*a*c+b^2+c
^2)^(1/2))))*2^(1/2)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(1/2)/(2*a-2*c-(a^2-2*a
*c+b^2+c^2)^(1/2))^(1/2)/(a^2-b^2-2*a*c+c^2+(a-c)*(a^2-2*a*c+b^2+c^2)^(1/2)
)^(1/2))*(2*a-2*c-(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)*(a^2-b^2-2*a*c+c^2+(a-c)
*(a^2-2*a*c+b^2+c^2)^(1/2))^(1/2)/(a^2-2*a*c+b^2+c^2)^(3/2)/e*2^(1/2)
```

Rubi [A] (verified)

Time = 4.23 (sec) , antiderivative size = 635, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.161$, Rules used = {3782, 1032, 1050, 1044, 214}

$$\int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx =$$

$$\frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2+2a-2c}\sqrt{(a-c)\sqrt{a^2-2ac+b^2+c^2+a^2-2ac-b^2+c^2}\operatorname{arctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2+2a-2c}}{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2+2a-2c}}}\right)} + \sqrt{\sqrt{a^2-2ac+b^2+c^2+2a-2c}\sqrt{-(a-c)\sqrt{a^2-2ac+b^2+c^2+a^2-2ac-b^2+c^2}\operatorname{arctanh}\left(\frac{-\sqrt{a^2-2ac+b^2+c^2+2a-2c}}{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2+2a-2c}}}\right)}}}{\sqrt{2}e(a^2-2ac+b^2+c^2)^{3/2}}$$

$$+ \frac{2(a(b^2-2c(a-c))+bc(a+c)\cot(d+ex))}{e((a-c)^2+b^2)(b^2-4ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}$$

[In] Int[Cot[d + e*x]/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2), x]

[Out] -((Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) + (Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 1032

Int[((g_) + (h_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_)*((d_) + (f_)*(x_)^2)^(q_), x_Symbol] := Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c

```

d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2
*f - c*(2*a*f)) - h*(b*c*d + a*b*f))*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*
f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Si
mp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*
(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(
(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))*(p +
q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1
)))*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q
+ 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*
a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !( !IntegerQ[p
] && ILtQ[q, -1])

```

Rule 1044

```

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f
_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*
e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[
{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

```

Rule 1050

```

Int[((g_) + (h_)*(x_))/(((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (
f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist
[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*
e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Si
mp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c
*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] &&
NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

```

Rule 3782

```

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)])*(
f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)])*(f_)^(n2_)]^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\text{integral} = -\frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e}$$

$$\begin{aligned}
&= -\frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&+ \frac{2 \text{Subst} \left(\int \frac{-\frac{1}{2}b(b^2 - 4ac) - \frac{1}{2}(a - c)(b^2 - 4ac)x}{(1 + x^2)\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 + (a - c)^2) (b^2 - 4ac) e} \\
&= -\frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&\text{Subst} \left(\int \frac{\frac{1}{2}b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2}(b^2 - 4ac) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))x}{(1 + x^2)\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&- \frac{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&\text{Subst} \left(\int \frac{\frac{1}{2}b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2}(b^2 - 4ac) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))x}{(1 + x^2)\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right) \\
&+ \frac{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&= -\frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&\frac{(b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&+ \frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst} \left(\int \frac{1}{\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)} \\
&= \\
&\frac{\sqrt{2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 - 2ac + c^2} + (a - c) \sqrt{a^2 + b^2 - 2ac + c^2} \operatorname{arctanh} \left(\frac{1}{\sqrt{2}} \right)}{\sqrt{2} (a^2 + b^2 - 2ac + c^2)} \\
&- \frac{\sqrt{2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}} \sqrt{a^2 - b^2 + c} (c + \sqrt{a^2 + b^2 - 2ac + c^2}) - a (2c + \sqrt{a^2 + b^2 - 2ac + c^2})}{\sqrt{2} (a^2 + b^2 - 2ac + c^2)} \\
&+ \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex)} + c \cot^2(d + ex)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.40 (sec) , antiderivative size = 460, normalized size of antiderivative = 0.72

$$\int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{2\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{\dots} \left(-\frac{i(4a^2c+\dots)}{\dots} \right)$$

[In] Integrate[Cot[d + e*x]/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] (2*Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*(-1/4*((I*(4*a^2*c + b^2*(I*b + c) - a*(b^2 + (4*I)*b*c + 4*c^2))*ArcTan[((-I)*b + 2*c + (-2*I)*a + b)*Tan[d + e*x]]/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])))/Sqrt[a + I*b - c] + ((-(b^2*(b + I*c)) - (4*I)*a^2*c + a*(I*b^2 + 4*b*c + (4*I)*c^2))*ArcTan[(I*b + 2*c + ((2*I)*a + b)*Tan[d + e*x]]/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])))/Sqrt[a - I*b - c])/((b^2 + (a - c)^2)*(b^2 - 4*a*c)) + (b + 2*a*Tan[d + e*x])/((-b^2 + 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) + (b^3 + a*b*(a - 3*c) + a*(2*a^2 + b^2 - 2*a*c)*Tan[d + e*x])/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]))/(e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])

Maple [B] (warning: unable to verify)

result has leaf size over 500,000. Avoiding possible recursion issues.

Time = 1.24 (sec) , antiderivative size = 13066366, normalized size of antiderivative = 20576.95

output too large to display

[In] int(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)

[Out] result too large to display

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 21090 vs. 2(580) = 1160.

Time = 11.87 (sec) , antiderivative size = 21090, normalized size of antiderivative = 33.21

$$\int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2),x)

[Out] Integral(cot(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Exception raised: ValueError}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a*c-b^2>0)', see 'assume?' for more details)

Giac [F(-2)]

Exception generated.

$$\int \frac{\cot(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Not invertible Error: Bad Argument Value
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \int \frac{\cot(d + ex)}{(c \cot(d + ex)^2 + b \cot(d + ex) + a)^{3/2}} dx$$

```
[In] int(cot(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2),x)
```

```
[Out] int(cot(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)
```

$$3.15 \quad \int \frac{\tan(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	164
Rubi [A] (verified)	165
Mathematica [C] (verified)	169
Maple [F(-1)]	170
Fricas [B] (verification not implemented)	170
Sympy [F]	170
Maxima [F(-1)]	171
Giac [F(-2)]	171
Mupad [F(-1)]	171

Optimal result

Integrand size = 31, antiderivative size = 749

$$\int \frac{\tan(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{a^{3/2}e}$$

$$+ \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$- \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2-(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{b^2+\sqrt{2}\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$- \frac{2(b^2-2ac+bc \cot(d+ex))}{a(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$+ \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

[Out] $\operatorname{arctanh}\left(\frac{1/2*(2*a+b*\cot(e*x+d))/a^{1/2}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}}{a^{3/2}/e-2*(b^2-2*a*c+b*c*\cot(e*x+d))/a/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}+2*(a*(b^2-2*(a-c)*c)+b*c*(a+c)*\cot(e*x+d))/(b^2+(a-c)^2)/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}-1/2*\operatorname{arctanh}\left(\frac{1/2*(b^2-(a-c)*(a-c-(a^2-2*a*c+b^2+c^2)^{1/2})-b*\cot(e*x+d)*(2*a-2*c+(a^2-2*a*c+b^2+c^2)^{1/2}))}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)*2^{1/2}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}/(2*a-2*c+(a^2-2*a*c+b^2+c^2)^{1/2})^{1/2}/(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c^2)^{1/2})^{1/2}\right)*2^{1/2}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}/(2*a-2*c+(a^2-2*a*c+b^2+c^2)^{1/2})^{1/2}/(a^2-b^2-2*a*c+c^2-(a-c)*(a^2-2*a*c+b^2+c^2)^{1/2})^{1/2}/(a^2-2*a*c+b^2+c^2)^{3/2}/e*2^{1/2}+1/2*\operatorname{arctanh}\left(\frac{1/2*(b^2-b*\cot(e*x+d)*(2*a-2*c-(a^2-2*a*c+b^2+c^2)^{1/2})-(a-c)*(a-c+(a^2-2*a*c+b^2+c^2)^{1/2}))}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}\right)*2^{1/2}/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{1/2}$

$$\frac{1/2}{(2a-2c-(a^2-2ac+b^2+c^2)^{1/2})^{1/2}} \frac{1/2}{(a^2-b^2-2ac+c^2+(a-c)(a^2-2ac+b^2+c^2)^{1/2})^{1/2}} * (2a-2c-(a^2-2ac+b^2+c^2)^{1/2})^{1/2} * (a^2-b^2-2ac+c^2+(a-c)(a^2-2ac+b^2+c^2)^{1/2})^{1/2} / (a^2-2ac+b^2+c^2)^{3/2} / e^{1/2}$$

Rubi [A] (verified)

Time = 5.13 (sec) , antiderivative size = 749, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.323$, Rules used = {3782, 6857, 754, 12, 738, 212, 1032, 1050, 1044, 214}

$$\int \frac{\tan(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{a^{3/2}e}$$

$$+ \frac{\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+2a-2c}\sqrt{(a-c)\sqrt{a^2-2ac+b^2+c^2}+a^2-2ac-b^2+c^2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{-\sqrt{a^2-2ac+b^2+c^2}+2a-2c}}{\sqrt{2}e(a^2-2ac+b^2+c^2)^{3/2}}\right)}{\sqrt{2}e(a^2-2ac+b^2+c^2)^{3/2}}$$

$$- \frac{\sqrt{\sqrt{a^2-2ac+b^2+c^2}+2a-2c}\sqrt{-(a-c)\sqrt{a^2-2ac+b^2+c^2}+a^2-2ac-b^2+c^2}\operatorname{arctanh}\left(\frac{\sqrt{2}\sqrt{\sqrt{a^2-2ac+b^2+c^2}+2a-2c}}{\sqrt{2}e(a^2-2ac+b^2+c^2)^{3/2}}\right)}{\sqrt{2}e(a^2-2ac+b^2+c^2)^{3/2}}$$

$$- \frac{2(-2ac+b^2+bc\cot(d+ex))}{ae(b^2-4ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}$$

$$+ \frac{2(a(b^2-2c(a-c))+bc(a+c)\cot(d+ex))}{e((a-c)^2+b^2)(b^2-4ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}$$

[In] Int[Tan[d + e*x]/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2), x]

[Out] ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(a^(3/2)*e) + (Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2])*Cot[d + e*x])/(Sqrt[2]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])]/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) - (2*(b^2 - 2*a*c + b*c*Cot[d + e*x]))/(a*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) + (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 738

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 754

Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 1032

Int[((g_.) + (h_.)*(x_))*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2*f - c*(2*a*f)) - h*(b*c*d + a*b*f))*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Simp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))]*(p +

```

q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1)
)*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q
+ 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*
a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !( !IntegerQ[p
] && ILtQ[q, -1])

```

Rule 1044

```

Int[((g_) + (h_)*(x_))/((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (f
_)*(x_)^2]), x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*
e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[
{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

```

Rule 1050

```

Int[((g_) + (h_)*(x_))/((a_) + (c_)*(x_)^2)*Sqrt[(d_) + (e_)*(x_) + (
f_)*(x_)^2]), x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist
[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*
e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Si
mp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c
*x^2)*Sqrt[d + e*x + f*x^2]), x], x]] /; FreeQ[{a, c, d, e, f, g, h}, x] &&
NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

```

Rule 3782

```

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(
f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rule 6857

```

Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \left(\frac{1}{x(a+bx+cx^2)^{3/2}} - \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot(d+ex)\right)}{e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{\text{Subst}\left(\int \frac{1}{x(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} + \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= - \frac{2(b^2 - 2ac + bc \cot(d+ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2 \text{Subst}\left(\int \frac{-\frac{b^2}{2} + 2ac}{x \sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{a(b^2 - 4ac) e} \\
&\quad - \frac{2 \text{Subst}\left(\int \frac{-\frac{1}{2}b(b^2-4ac) - \frac{1}{2}(a-c)(b^2-4ac)x}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 + (a-c)^2) (b^2 - 4ac) e} \\
&= - \frac{2(b^2 - 2ac + bc \cot(d+ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{x \sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{ae} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{ae}{\frac{1}{2}b(b^2-4ac)(2a-2c+\sqrt{a^2+b^2-2ac+c^2}) + \frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c-\sqrt{a^2+b^2-2ac+c^2}))x}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{1}{2}b(b^2-4ac)(2a-2c-\sqrt{a^2+b^2-2ac+c^2}) + \frac{1}{2}(b^2-4ac)(b^2-(a-c)(a-c+\sqrt{a^2+b^2-2ac+c^2}))x}}{(1+x^2)\sqrt{a+bx+cx^2}} dx, x, \cot(d+ex)\right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&= - \frac{2(b^2 - 2ac + bc \cot(d+ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2(a(b^2 - 2(a-c)c) + bc(a+c) \cot(d+ex))}{(b^2 + (a-c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d+ex) + c \cot^2(d+ex)}} \\
&\quad + \frac{2 \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{ae} \\
&\quad - \frac{(b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a-c) (a-c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a-c) (a-c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)} \\
&\quad + \frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a-c) (a-c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a-c) (a-c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{a^{3/2}e} \\
&+ \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{\sqrt{2}(a^2+b^2-2ac+c^2)}{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2})}}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&- \frac{2(b^2-2ac+bc\cot(d+ex))}{a(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
&+ \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.55 (sec) , antiderivative size = 934, normalized size of antiderivative = 1.25

$$\int \frac{\tan(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{2\left(\frac{4\sqrt{a-ib}}{\dots}\right)}$$

[In] Integrate[Tan[d + e*x]/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] (Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((2*((-4*Sqrt[a - I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) + (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[(I*b + 2*c - ((-2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]))/(-4*a + (4*I)*b + 4*c) - (4*Sqrt[a + I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) - (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[((-I)*b + 2*c - ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])]))/(-4*a - (4*I)*b + 4*c))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)) - (2*Tan[d + e*x]^3*(-b^2 + 2*a*c - a*b*Tan[d + e*x]))/(c*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - (2*(b^3 + a*b*(a - 3*c) + a*(2*a^2 + b^2 - 2*a*c)*Tan[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) + (4*(b^2 - 4*a*c)*(a^2/((b^2 - 4*a*c)*((a^2*b^2)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2 - 4*a*c)^2))))^(3/2)

```
*(-((a*b)/(b^2 - 4*a*c)) - (2*a^2*Tan[d + e*x])/(b^2 - 4*a*c))*(-((a*(c + b
*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2 - 4*a*c)))^(3/2))/(a^2*(c + b*Tan[d
+ e*x] + a*Tan[d + e*x]^2)^(3/2)*Sqrt[1 - (-((a*b)/(b^2 - 4*a*c)) - (2*a^2
*Tan[d + e*x])/(b^2 - 4*a*c))^2/((a^2*b^2)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2
- 4*a*c)^2)]) - (2*(b*Tan[d + e*x]^2*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e
*x]^2] + (((-6*a^2*b^2*c + 24*a^3*c^2)*ArcTanh[(b + 2*a*Tan[d + e*x])/(2*Sq
rt[a]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)])))/(4*a^(5/2)) + ((6*a^2*
b*c - 12*a^3*c*Tan[d + e*x])*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])/(
2*a^2))/(3*a)))/(c*(b^2 - 4*a*c)))/(e*Sqrt[Cot[d + e*x]^2*(c + b*Tan[d + e
*x] + a*Tan[d + e*x]^2]))
```

Maple [F(-1)]

Timed out.

hanged

```
[In] int(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)
```

```
[Out] int(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 19942 vs. 2(684) = 1368.

Time = 8.08 (sec) , antiderivative size = 39885, normalized size of antiderivative = 53.25

$$\int \frac{\tan(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Too large to display}$$

```
[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="fr
icas")
```

```
[Out] Too large to include
```

Sympy [F]

$$\int \frac{\tan(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\tan(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx$$

```
[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2),x)
```

```
[Out] Integral(tan(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)
```

Maxima [F(-1)]

Timed out.

$$\int \frac{\tan(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="maxima")

[Out] Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{\tan(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \int \frac{\tan(d + ex)}{(c \cot(d + ex)^2 + b \cot(d + ex) + a)^{3/2}} dx$$

[In] int(tan(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2),x)

[Out] int(tan(d + e*x)/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2), x)

$$3.16 \quad \int \frac{\tan^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx$$

Optimal result	172
Rubi [A] (verified)	173
Mathematica [C] (verified)	181
Maple [F(-1)]	183
Fricas [B] (verification not implemented)	183
Sympy [F]	184
Maxima [F(-1)]	184
Giac [F(-2)]	184
Mupad [F(-1)]	185

Optimal result

Integrand size = 33, antiderivative size = 1008

$$\int \frac{\tan^3(d+ex)}{(a+b \cot(d+ex)+c \cot^2(d+ex))^{3/2}} dx = -\frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{a^{3/2}e}$$

$$+ \frac{3(5b^2 - 4ac) \operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{8a^{7/2}e}$$

$$- \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}+(a-c)\sqrt{a^2+b^2-2ac+c^2} \operatorname{arctanh}\left(\frac{b^2-\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$+ \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2}-(a-c)\sqrt{a^2+b^2-2ac+c^2} \operatorname{arctanh}\left(\frac{b^2-\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}{\sqrt{2}\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)^{3/2}e}$$

$$+ \frac{2(b^2-2ac+bc \cot(d+ex))}{a(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$- \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$- \frac{b(15b^2-52ac)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan(d+ex)}{4a^3(b^2-4ac)e}$$

$$- \frac{2(b^2-2ac+bc \cot(d+ex)) \tan^2(d+ex)}{a(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}$$

$$+ \frac{(5b^2-12ac)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan^2(d+ex)}{2a^2(b^2-4ac)e}$$

[Out] $-\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d))/a^{(1/2)})/(a+b*\cot(e*x+d)+c*\cot(e*x+d)^2)^{(1/2)}/a^{(3/2)}/e+3/8*(-4*a*c+5*b^2)*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d))/a^{(1/2)})/(a+$

$$\begin{aligned}
& b \cot(e^x+d) + c \cot(e^x+d)^2)^{1/2} / a^{7/2} / e + 2(b^2 - 2ac + bc \cot(e^x+d)) / \\
& a / (-4ac + b^2) / e / (a + b \cot(e^x+d) + c \cot(e^x+d)^2)^{1/2} - 2(a(b^2 - 2(a-c)c \\
& + bc(a+c) \cot(e^x+d)) / (b^2 + (a-c)^2) / (-4ac + b^2) / e / (a + b \cot(e^x+d) + c \cot(e \\
& *x+d)^2)^{1/2} + 1/2 \operatorname{arctanh}(1/2(b^2 - (a-c)(a-c - (a^2 - 2ac + b^2 + c^2)^{1/2})) - b \\
& * \cot(e^x+d) * (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2})) * 2^{1/2} / (a + b \cot(e^x+d) + c \\
& \cot(e^x+d)^2)^{1/2} / (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - b^2 - 2ac \\
& + c^2 - (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} * (2a - 2c + (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} * (a^2 - b^2 - 2ac + c^2 - (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 \\
& - 2ac + b^2 + c^2)^{3/2} / e * 2^{1/2} - 1/2 \operatorname{arctanh}(1/2(b^2 - b \cot(e^x+d) * (2a - 2c - \\
& (a^2 - 2ac + b^2 + c^2)^{1/2})) - (a-c)(a-c + (a^2 - 2ac + b^2 + c^2)^{1/2})) * 2^{1/2} / (\\
& a + b \cot(e^x+d) + c \cot(e^x+d)^2)^{1/2} / (2a - 2c - (a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - b^2 - 2ac + c^2 + (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} * (2a - 2c - (a^2 \\
& - 2ac + b^2 + c^2)^{1/2})^{1/2} * (a^2 - b^2 - 2ac + c^2 + (a-c)(a^2 - 2ac + b^2 + c^2)^{1/2})^{1/2} / (a^2 - 2ac + b^2 + c^2)^{3/2} / e * 2^{1/2} - 1/4 b * (-52ac + 15b^2) * (a + \\
& b \cot(e^x+d) + c \cot(e^x+d)^2)^{1/2} * \tan(e^x+d) / a^3 / (-4ac + b^2) / e - 2(b^2 - 2ac \\
& + bc \cot(e^x+d)) * \tan(e^x+d)^2 / a / (-4ac + b^2) / e / (a + b \cot(e^x+d) + c \cot(e^x + \\
& d)^2)^{1/2} + 1/2(-12ac + 5b^2) * (a + b \cot(e^x+d) + c \cot(e^x+d)^2)^{1/2} * \tan(e \\
& *x+d)^2 / a^2 / (-4ac + b^2) / e
\end{aligned}$$

Rubi [A] (verified)

Time = 5.55 (sec) , antiderivative size = 1008, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules

used = {3782, 6857, 754, 848, 820, 738, 212, 12, 1032, 1050, 1044, 214}

$$\int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \frac{(5b^2-12ac)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}\tan^2(d+ex)}{2a^2(b^2-4ac)e}$$

$$- \frac{2(b^2+c\cot(d+ex)b-2ac)\tan^2(d+ex)}{a(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}$$

$$- \frac{b(15b^2-52ac)\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}\tan(d+ex)}{4a^3(b^2-4ac)e}$$

$$+ \frac{3(5b^2-4ac)\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{8a^{7/2}e} - \frac{\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}\right)}{a^{3/2}e}$$

$$- \frac{\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2+(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}\sqrt{2a-2c-\sqrt{a^2-2ca+b^2+c^2}}}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e}$$

$$+ \frac{\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}\sqrt{a^2-2ca-b^2+c^2-(a-c)\sqrt{a^2-2ca+b^2+c^2}}\operatorname{arctanh}\left(\frac{b^2-\sqrt{2}\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}}{\sqrt{2}\sqrt{2a-2c+\sqrt{a^2-2ca+b^2+c^2}}}\right)}{\sqrt{2}(a^2-2ca+b^2+c^2)^{3/2}e}$$

$$+ \frac{2(b^2+c\cot(d+ex)b-2ac)}{a(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}$$

$$- \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{c\cot^2(d+ex)+b\cot(d+ex)+a}}$$

[In] Int[Tan[d + e*x]^3/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2), x]

[Out] -(ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2))]/(a^(3/2)*e)) + (3*(5*b^2 - 4*a*c)*ArcTanh[(2*a + b*Cot[d + e*x])/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2))]/(8*a^(7/2)*e) - (Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c + Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Cot[d + e*x])]/(Sqrt[2]*Sqrt[2*a - 2*c - Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 + (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) + (Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*ArcTanh[(b^2 - (a - c)*(a - c - Sqrt[a^2 + b^2 - 2*a*c + c^2]) - b*(2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]))*Cot[d + e*x])]/(Sqrt[2]*Sqrt[2*a - 2*c + Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a^2 - b^2 - 2*a*c + c^2 - (a - c)*Sqrt[a^2 + b^2 - 2*a*c + c^2]]*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]))/(Sqrt[2]*(a^2 + b^2 - 2*a*c + c^2)^(3/2)*e) + (2*(b^2 - 2*a*c + b*c*Cot[d + e*x]))/(a*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) - (2*(a*(b^2 - 2*(a - c)*c) + b*c*(a + c)*Cot[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x] + c*Cot[d + e*x]^2]) - (b*(15*b^2 - 52*a*c)*Sqrt[a

$$+ b \cot[d + e x] + c \cot[d + e x]^2 \tan[d + e x] / (4 a^3 (b^2 - 4 a c) e) - (2 (b^2 - 2 a c + b c \cot[d + e x]) \tan[d + e x]^2) / (a (b^2 - 4 a c) e \sqrt{a + b \cot[d + e x] + c \cot[d + e x]^2}) + ((5 b^2 - 12 a c) \sqrt{a + b \cot[d + e x] + c \cot[d + e x]^2} \tan[d + e x]^2) / (2 a^2 (b^2 - 4 a c) e)$$
Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 738

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 754

```
Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1))/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]
```

Rule 820

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(-e*f - d*g)*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1))/(2*(p + 1)*(c*d^2 - b*d*e + a*e^2)), x] - Dist[(b*(e*f + d*g) - 2*(c*d*f + a*e*g))/(2*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x]
```

&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 848

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*Simp[(c*d*f - f*b*e + a*e*g)*(m + 1) + b*(d*g - e*f)*(p + 1) - c*(e*f - d*g)*(m + 2*p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

Rule 1032

Int[((g_.) + (h_.)*(x_))*((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)*((d_) + (f_.)*(x_)^2)^(q_), x_Symbol] := Simp[(a + b*x + c*x^2)^(p + 1)*((d + f*x^2)^(q + 1)/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)))*((g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)) + c*(g*(2*c^2*d + b^2*f - c*(2*a*f)) - h*(b*c*d + a*b*f)*x), x] + Dist[1/((b^2 - 4*a*c)*(b^2*d*f + (c*d - a*f)^2)*(p + 1)), Int[(a + b*x + c*x^2)^(p + 1)*(d + f*x^2)^q*Simp[(b*h - 2*g*c)*((c*d - a*f)^2 - (b*d)*((-b)*f))*(p + 1) + (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(a*f*(p + 1) - c*d*(p + 2)) - (2*f*(g*c)*((-b)*(c*d + a*f)) + (g*b - a*h)*(2*c^2*d + b^2*f - c*(2*a*f)))*(p + q + 2) - (b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(b*f*(p + 1)))*x - c*f*(b^2*(g*f) - b*(h*c*d + a*h*f) + 2*(g*c*(c*d - a*f)))*(2*p + 2*q + 5)*x^2, x], x] /; FreeQ[{a, b, c, d, f, g, h, q}, x] && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1] && NeQ[b^2*d*f + (c*d - a*f)^2, 0] && !(IntegerQ[p] || ILtQ[q, -1])

Rule 1044

Int[((g_) + (h_.)*(x_))/((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2], x_Symbol] := Dist[-2*a*g*h, Subst[Int[1/Simp[2*a^2*g*h*c + a*e*x^2, x], x], x, Simp[a*h - g*c*x, x]/Sqrt[d + e*x + f*x^2]], x] /; FreeQ[{a, c, d, e, f, g, h}, x] && EqQ[a*h^2*e + 2*g*h*(c*d - a*f) - g^2*c*e, 0]

Rule 1050

Int[((g_.) + (h_.)*(x_))/((a_) + (c_.)*(x_)^2)*Sqrt[(d_.) + (e_.)*(x_) + (f_.)*(x_)^2], x_Symbol] := With[{q = Rt[(c*d - a*f)^2 + a*c*e^2, 2]}, Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f - q) + (h*(c*d - a*f + q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] - Dist[1/(2*q), Int[Simp[(-a)*h*e - g*(c*d - a*f + q) + (h*(c*d - a*f - q) - g*c*e)*x, x]/((a + c*x^2)*Sqrt[d + e*x + f*x^2]), x], x] /; FreeQ[{a, c, d, e, f, g, h}, x] &&

NeQ[e^2 - 4*d*f, 0] && NegQ[(-a)*c]

Rule 3782

Int[cot[(d_.) + (e_.)*(x_.)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_.)]*(f_.))^(n2_.))^(p_), x_Symbol]
 :> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
 xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
 [n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x^3(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^3(a+bx+cx^2)^{3/2}} - \frac{1}{x(a+bx+cx^2)^{3/2}} + \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^3(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{x(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx+cx^2)^{3/2}} dx, x, \cot(d+ex)\right)}{e}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{2(b^2 - 2ac + bc \cot(d + ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(b^2 - 2ac + bc \cot(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2 \text{Subst} \left(\int \frac{-\frac{b^2}{2} + 2ac}{x \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{a(b^2 - 4ac) e} \\
&\quad + \frac{2 \text{Subst} \left(\int \frac{\frac{1}{2}(-5b^2 + 12ac) - 2bcx}{x^3 \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{a(b^2 - 4ac) e} \\
&\quad + \frac{2 \text{Subst} \left(\int \frac{-\frac{1}{2}b(b^2 - 4ac) - \frac{1}{2}(a - c)(b^2 - 4ac)x}{(1 + x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 + (a - c)^2) (b^2 - 4ac) e} \\
&= \frac{2(b^2 - 2ac + bc \cot(d + ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(b^2 - 2ac + bc \cot(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad + \frac{(5b^2 - 12ac) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{2a^2 (b^2 - 4ac) e} \\
&\quad + \frac{\text{Subst} \left(\int \frac{1}{x \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{ae} \\
&\quad - \frac{\text{Subst} \left(\int \frac{-\frac{1}{4}b(15b^2 - 52ac) - \frac{1}{2}c(5b^2 - 12ac)x}{x^2 \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{a^2 (b^2 - 4ac) e} \\
&\quad - \frac{\text{Subst} \left(\int \frac{\frac{1}{2}b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2}(b^2 - 4ac) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2})) x}{(1 + x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e} \\
&\quad + \frac{\text{Subst} \left(\int \frac{\frac{1}{2}b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) + \frac{1}{2}(b^2 - 4ac) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2})) x}{(1 + x^2) \sqrt{a + bx + cx^2}} dx, x, \cot(d + ex) \right)}{(b^2 - 4ac) (a^2 + b^2 - 2ac + c^2)^{3/2} e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2(b^2 - 2ac + bc \cot(d + ex))}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{2(a(b^2 - 2(a - c)c) + bc(a + c) \cot(d + ex))}{(b^2 + (a - c)^2) (b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad - \frac{b(15b^2 - 52ac) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan(d + ex)}{4a^3 (b^2 - 4ac) e} \\
&\quad - \frac{2(b^2 - 2ac + bc \cot(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}} \\
&\quad + \frac{(5b^2 - 12ac) \sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)} \tan^2(d + ex)}{2a^2 (b^2 - 4ac) e} \\
&\quad - \frac{2 \text{Subst}\left(\int \frac{1}{4a - x^2} dx, x, \frac{2a + b \cot(d + ex)}{\sqrt{a + b \cot(d + ex) + c \cot^2(d + ex)}}\right)}{ae} \\
&\quad - \frac{(3(5b^2 - 4ac)) \text{Subst}\left(\int \frac{1}{x\sqrt{a + bx + cx^2}} dx, x, \cot(d + ex)\right)}{8a^3 e} \\
&\quad + \frac{(b(b^2 - 4ac) (2a - 2c + \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c - \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Su}}{---} \\
&\quad - \frac{(b(b^2 - 4ac) (2a - 2c - \sqrt{a^2 + b^2 - 2ac + c^2}) (b^2 - (a - c) (a - c + \sqrt{a^2 + b^2 - 2ac + c^2}))) \text{Su}}{---}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{\operatorname{arctanh}\left(\frac{2a+b \cot(d+ex)}{2\sqrt{a}\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{a^{3/2}e} \\
&\quad - \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{-}{\sqrt{2}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&\quad + \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})-a(2c+\sqrt{a^2+b^2-2ac+c^2})}}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&\quad + \frac{2(b^2-2ac+bc \cot(d+ex))}{a(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad - \frac{2(a(b^2-2(a-c)c)+bc(a+c) \cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad - \frac{b(15b^2-52ac)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan(d+ex)}{4a^3(b^2-4ac)e} \\
&\quad - \frac{2(b^2-2ac+bc \cot(d+ex)) \tan^2(d+ex)}{a(b^2-4ac)e\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}} \\
&\quad + \frac{(5b^2-12ac)\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)} \tan^2(d+ex)}{2a^2(b^2-4ac)e} \\
&\quad + \frac{(3(5b^2-4ac)) \operatorname{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot(d+ex)}{\sqrt{a+b \cot(d+ex)+c \cot^2(d+ex)}}\right)}{4a^3e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{a^{3/2}e} \\
&+ \frac{3(5b^2-4ac)\operatorname{arctanh}\left(\frac{2a+b\cot(d+ex)}{2\sqrt{a}\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}}\right)}{8a^{7/2}e} \\
&- \frac{\sqrt{2a-2c-\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2-2ac+c^2+(a-c)\sqrt{a^2+b^2-2ac+c^2}}\operatorname{arctanh}\left(\frac{\sqrt{2}(a^2+b^2-2ac+c^2)}{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}}\right)}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&+ \frac{\sqrt{2a-2c+\sqrt{a^2+b^2-2ac+c^2}}\sqrt{a^2-b^2+c(c+\sqrt{a^2+b^2-2ac+c^2})}-a(2c+\sqrt{a^2+b^2-2ac+c^2})}{\sqrt{2}(a^2+b^2-2ac+c^2)} \\
&+ \frac{2(b^2-2ac+bc\cot(d+ex))}{a(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
&- \frac{2(a(b^2-2(a-c)c)+bc(a+c)\cot(d+ex))}{(b^2+(a-c)^2)(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
&- \frac{b(15b^2-52ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}\tan(d+ex)}{4a^3(b^2-4ac)e} \\
&- \frac{2(b^2-2ac+bc\cot(d+ex))\tan^2(d+ex)}{a(b^2-4ac)e\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}} \\
&+ \frac{(5b^2-12ac)\sqrt{a+b\cot(d+ex)+c\cot^2(d+ex)}\tan^2(d+ex)}{2a^2(b^2-4ac)e}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 6.66 (sec) , antiderivative size = 1401, normalized size of antiderivative = 1.39

$$\int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx =$$

$$\frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{2} \left(\frac{4\sqrt{a-ib-c}\left(-\frac{1}{4}b(b^2-4ac)+\frac{1}{4}i(a-c)(b^2-4ac)\right)\arctan\left(\frac{ib+2c-(-2ia-)}{2\sqrt{a-ib-c}\sqrt{c+b\tan(d+ex)}}\right)}{-4a+4ib+4c} \right)$$

$$\frac{\cot(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} - \frac{2\tan^5(d+ex)(-b^2+2ac-ab\tan(d+ex))}{c(b^2-4ac)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}} - \frac{2}{b\tan^4(d+ex)\sqrt{c+b\tan(d+ex)+a\tan^2(d+ex)}}$$

+

[In] Integrate[Tan[d + e*x]^3/(a + b*Cot[d + e*x] + c*Cot[d + e*x]^2)^(3/2),x]

[Out] -((Cot[d + e*x]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]*((2*((-4*Sqrt[a - I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) + (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[(I*b + 2*c - ((-2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a - I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])])/(-4*a + (4*I)*b + 4*c) - (4*Sqrt[a + I*b - c]*(-1/4*(b*(b^2 - 4*a*c)) - (I/4)*(a - c)*(b^2 - 4*a*c))*ArcTan[((-I)*b + 2*c - ((2*I)*a - b)*Tan[d + e*x])/(2*Sqrt[a + I*b - c]*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2])])/(-4*a - (4*I)*b + 4*c)))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)) - (2*Tan[d + e*x]^3*(-b^2 + 2*a*c - a*b*Tan[d + e*x]))/(c*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) - (2*(b^3 + a*b*(a - 3*c) + a*(2*a^2 + b^2 - 2*a*c)*Tan[d + e*x]))/((b^2 + (a - c)^2)*(b^2 - 4*a*c)*Sqrt[c + b*Tan[d + e*x] + a*Tan[d + e*x]^2]) + (4*(b^2 - 4*a*c)*(a^2/(b^2 - 4*a*c)*((a^2*b^2)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2 - 4*a*c)^2)))^(3/2)*(-(a*b)/(b^2 - 4*a*c)) - (2*a^2*Tan[d + e*x])/(b^2 - 4*a*c))*(-(a*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2))/(b^2 - 4*a*c)))^(3/2))/(a^2*(c + b*Tan[d + e*x] + a*Tan[d + e*x]^2)^(3/2)*Sqrt[1 - ((a*b)/(b^2 - 4*a*c)) - (2*a^2*Tan[d + e*x])/(b^2 - 4*a*c)]^2/((a^2*b^2)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2 - 4*a*c)^2 - (4*a^3*c)/(b^2 - 4*a*c)^2))^(3/2))

$$\begin{aligned} &^2 - 4*a*c)^2]) - (2*(b*\text{Tan}[d + e*x]^2*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + \\ &e*x]^2] + (((-6*a^2*b^2*c + 24*a^3*c^2)*\text{ArcTanh}[(b + 2*a*\text{Tan}[d + e*x])/(2* \\ &\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2])))/(4*a^{(5/2)}) + ((6*a^ \\ &2*b*c - 12*a^3*c*\text{Tan}[d + e*x])* \text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2]) \\ &/(2*a^2))/(3*a)))/(c*(b^2 - 4*a*c)))/(e*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + \\ &e*x] + a*\text{Tan}[d + e*x]^2])) + (\text{Cot}[d + e*x]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan} \\ &[d + e*x]^2]*((-2*\text{Tan}[d + e*x]^5*(-b^2 + 2*a*c - a*b*\text{Tan}[d + e*x]))/(c*(b^ \\ &2 - 4*a*c)*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2]) - (2*(b*\text{Tan}[d + e*x] \\ &]^4*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2] + (-10*a*c*\text{Tan}[d + e*x]^3*\text{S} \\ &\text{qrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2] + (20*a*b*c*\text{Tan}[d + e*x]^2*\text{Sqrt}[\\ &c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2] + (((240*a^3*b^2*c^2 - 45*a^2*b^2*c* \\ &(5*b^2 - 12*a*c) + 60*a^3*c^2*(5*b^2 - 12*a*c))*\text{ArcTanh}[(b + 2*a*\text{Tan}[d + e* \\ &x])/(2*\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2])))/(4*a^{(5/2)}) + \\ &((-240*a^3*b*c^2 + 45*a^2*b*c*(5*b^2 - 12*a*c) - 30*a^3*c*(5*b^2 - 12*a*c) \\ &*\text{Tan}[d + e*x])* \text{Sqrt}[c + b*\text{Tan}[d + e*x] + a*\text{Tan}[d + e*x]^2))/(2*a^2))/(3*a) \\ &/(4*a))/(5*a)))/(c*(b^2 - 4*a*c)))/(e*\text{Sqrt}[\text{Cot}[d + e*x]^2*(c + b*\text{Tan}[d + e \\ &*x] + a*\text{Tan}[d + e*x]^2)]) \end{aligned}$$

Maple [F(-1)]

Timed out.

hanged

[In] `int(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)`

[Out] `int(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20316 vs. $2(923) = 1846$.

Time = 7.93 (sec) , antiderivative size = 40633, normalized size of antiderivative = 40.31

$$\int \frac{\tan^3(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Too large to display}$$

[In] `integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(tan(e*x+d)**3/(a+b*cot(e*x+d)+c*cot(e*x+d)**2)**(3/2), x)

[Out] Integral(tan(d + e*x)**3/(a + b*cot(d + e*x) + c*cot(d + e*x)**2)**(3/2), x)

Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2), x, algorithm="maxima")

[Out] Timed out

Giac [F(-2)]

Exception generated.

$$\int \frac{\tan^3(d+ex)}{(a+b\cot(d+ex)+c\cot^2(d+ex))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)+c*cot(e*x+d)^2)^(3/2), x, algorithm="giac")

[Out] Exception raised: TypeError >> an error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Not invertible Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^3(d + ex)}{(a + b \cot(d + ex) + c \cot^2(d + ex))^{3/2}} dx = \text{Hanged}$$

```
[In] int(tan(d + e*x)^3/(a + b*cot(d + e*x) + c*cot(d + e*x)^2)^(3/2),x)
```

```
[Out] \text{Hanged}
```

$$3.17 \quad \int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

Optimal result	186
Rubi [A] (verified)	186
Mathematica [A] (verified)	189
Maple [A] (verified)	189
Fricas [B] (verification not implemented)	190
Sympy [F]	191
Maxima [F(-1)]	192
Giac [F(-1)]	192
Mupad [F(-1)]	192

Optimal result

Integrand size = 35, antiderivative size = 182

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

$$= \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}}$$

$$+ \frac{(b+2c)\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4c^{3/2}e} - \frac{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}{2ce}$$

[Out] 1/4*(b+2*c)*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/c^(3/2)/e+1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/(a-b+c)^(1/2)-1/2*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)/c/e

Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 182, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3782, 1265, 1667, 857, 635, 212, 738}

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \frac{(b+2c)\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4c^{3/2}e}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e\sqrt{a-b+c}}$$

$$- \frac{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}{2ce}$$

[In] Int[Cot[d + e*x]^5/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*Sqrt[a - b + c]*e) + ((b + 2*c)*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]))/(4*c^(3/2)*e) - Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]/(2*c*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 857

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 1667

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*(a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1)), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b

```
*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)
)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*
d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q
, 1] && NeQ[m + q + 2*p + 1, 0]] /; FreeQ[{a, b, c, d, e, m, p}, x] && Poly
Q[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ
[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5}{(1+x^2)\sqrt{a+bx^2+cx^4}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x^2}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2ce} - \frac{\text{Subst}\left(\int \frac{-\frac{b}{2}-\frac{1}{2}(b+2c)x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2ce} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2ce} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&\quad + \frac{(b+2c)\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4ce} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2ce} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
&\quad + \frac{(b+2c)\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2ce}
\end{aligned}$$

$$= \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} + \frac{(b+2c)\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4c^{3/2}e} - \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2ce}$$

Mathematica [A] (verified)

Time = 3.62 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.43

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} \tan^2(d+ex) \left(2c^{3/2} \operatorname{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right) + \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} \right)}{4c^{3/2}\sqrt{a-b+ce}\sqrt{c}}$$

[In] Integrate[Cot[d + e*x]^5/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] (Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2*(2*c^(3/2)*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]]) + Sqrt[a - b + c]*((b + 2*c)*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]]) - 2*Sqrt[c]*Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))/(4*c^(3/2)*Sqrt[a - b + c]*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [A] (verified)

Time = 0.72 (sec) , antiderivative size = 232, normalized size of antiderivative = 1.27

method	result
derivativedivides	$-\frac{\sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2}+c\cot(ex+d)}{\sqrt{c}} + \sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}\right)}{4c^{3/2}} + \frac{\ln\left(\frac{\frac{b}{2}+c\cot(ex+d)}{\sqrt{c}} + \sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}\right)}{2\sqrt{c}}$
default	$-\frac{\sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}}{2c} + \frac{b \ln\left(\frac{\frac{b}{2}+c\cot(ex+d)}{\sqrt{c}} + \sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}\right)}{4c^{3/2}} + \frac{\ln\left(\frac{\frac{b}{2}+c\cot(ex+d)}{\sqrt{c}} + \sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}\right)}{2\sqrt{c}}$

[In] int(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2), x, method=_RETURNVE RBOSE)

```
[Out] 1/e*(-1/2/c*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)+1/4*b/c^(3/2)*ln((1/2*b+c*cot(e*x+d)^2)/c^(1/2)+(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))+1/2*ln((1/2*b+c*cot(e*x+d)^2)/c^(1/2)+(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/c^(1/2)+1/2/(a-b+c)^(1/2)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2))*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1)))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 508 vs. 2(158) = 316.

Time = 1.31 (sec) , antiderivative size = 2100, normalized size of antiderivative = 11.54

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Too large to display}$$

```
[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")
```

```
[Out] [1/8*(2*sqrt(a - b + c)*c^2*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d) + (a*b - b^2 + (2*a - b)*c + 2*c^2)*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 + 4*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*((a - b)*c + c^2)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c^2 + c^3)*e), 1/4*(sqrt(a - b + c)*c^2*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d) + (a*b - b^2 + (2*a - b)*c + 2*c^2)*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d)) - 2*((a - b)*c + c^2)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c^2 + c^3)*e), -1/8*(4*sqrt(-a + b - c)*c^2*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos
```

```
(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^
2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e
*x + 2*d) + 1))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2
+ a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d))) -
(a*b - b^2 + (2*a - b)*c + 2*c^2)*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^
2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 + 4*((b - 2*c)*cos(2*e*
x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(
2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^
2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(c
os(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) + 4*((a - b)*c + c^2)*sqrt(((a
- b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos
(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c^2 + c^3)*e), -1/4*(
2*sqrt(-a + b - c)*c^2*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*c
os(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d
)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2
*e*x + 2*d) + 1))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)
^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)))
- (a*b - b^2 + (2*a - b)*c + 2*c^2)*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*
e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*c
os(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*
d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a
+ b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d))) + 2*((a - b)*c + c^2)*sqrt(
((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(
cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c^2 + c^3)*e)]
```

Sympy [F]

$$\int \frac{\cot^5(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx = \int \frac{\cot^5(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx$$

[In] integrate(cot(e*x+d)**5/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(cot(d + e*x)**5/sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4), x)

Maxima [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] Timed out

Giac [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(d+ex)^5}{\sqrt{c\cot(d+ex)^4+b\cot(d+ex)^2+a}} dx$$

[In] int(cot(d + e*x)^5/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(cot(d + e*x)^5/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

$$3.18 \quad \int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

Optimal result	193
Rubi [A] (verified)	193
Mathematica [A] (verified)	195
Maple [A] (verified)	196
Fricas [B] (verification not implemented)	196
Sympy [F]	197
Maxima [F]	198
Giac [F(-1)]	198
Mupad [F(-1)]	198

Optimal result

Integrand size = 35, antiderivative size = 141

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = -\frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+c}} - \frac{\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ce}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d)^2)/c^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/e/c^{(1/2)}-1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/e/(a-b+c)^{(1/2)}$

Rubi [A] (verified)

Time = 0.35 (sec) , antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3782, 1265, 857, 635, 212, 738}

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = -\frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e\sqrt{a-b+c}} - \frac{\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ce}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[d+e*x]^3/\operatorname{Sqrt}[a+b*\operatorname{Cot}[d+e*x]^2+c*\operatorname{Cot}[d+e*x]^4],x]$

[Out] $-1/2*\operatorname{ArcTanh}[(2*a-b+(b-2*c)*\operatorname{Cot}[d+e*x]^2)/(2*\operatorname{Sqrt}[a-b+c]*\operatorname{Sqrt}[a+b*\operatorname{Cot}[d+e*x]^2+c*\operatorname{Cot}[d+e*x]^4)]/(\operatorname{Sqrt}[a-b+c]*e)-\operatorname{ArcTanh}[(b$

$+ 2*c*\cot[d + e*x]^2)/(2*\sqrt{c}*\sqrt{a + b*\cot[d + e*x]^2 + c*\cot[d + e*x]^4}))/ (2*\sqrt{c}*e)$

Rule 212

$\text{Int}[(a_.) + (b_.)*(x_.)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 635

$\text{Int}[1/\sqrt{(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2}], x_Symbol] \rightarrow \text{Dist}[2, \text{Subst}[\text{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\sqrt{a + b*x + c*x^2}], x] /; \text{FreeQ}\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 738

$\text{Int}[1/(((d_.) + (e_.)*(x_))*\sqrt{(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2})], x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\sqrt{a + b*x + c*x^2}], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0]$

Rule 857

$\text{Int}(((d_.) + (e_.)*(x_.)^m)*((f_.) + (g_.)*(x_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2)^{p_})), x_Symbol] \rightarrow \text{Dist}[g/e, \text{Int}[(d + e*x)^{m+1}*(a + b*x + c*x^2)^p, x], x] + \text{Dist}[(e*f - d*g)/e, \text{Int}[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, g, m, p\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ !\text{IGtQ}[m, 0]$

Rule 1265

$\text{Int}[(x_.)^{m_}]*((d_.) + (e_.)*(x_.)^2)^{q_}*((a_.) + (b_.)*(x_.)^2 + (c_.)*(x_.)^4)^{p_}, x_Symbol] \rightarrow \text{Dist}[1/2, \text{Subst}[\text{Int}[x^{(m-1)/2}*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, c, d, e, p, q\}, x] \ \&\& \ \text{IntegerQ}[(m-1)/2]$

Rule 3782

$\text{Int}[\cot[(d_.) + (e_.)*(x_.)]^{m_}]*((a_.) + (b_.)*(\cot[(d_.) + (e_.)*(x_.)]*(f_.)^{n_} + (c_.)*(\cot[(d_.) + (e_.)*(x_.)]*(f_.)^{n2_}))^{p_}), x_Symbol] \rightarrow \text{Dist}[-f/e, \text{Subst}[\text{Int}[(x/f)^m*((a + b*x^n + c*x^{2*n})^p/(f^2 + x^2)), x], x, f*\cot[d + e*x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \ \&\& \ \text{EqQ}[n, 2, 2*n] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= - \frac{\text{Subst}\left(\int \frac{x^3}{(1+x^2)\sqrt{a+bx^2+cx^4}} dx, x, \cot(d+ex)\right)}{e} \\
 &= - \frac{\text{Subst}\left(\int \frac{x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= - \frac{\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
 &= - \frac{\text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} - \frac{\text{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{ce}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.84 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.50

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \frac{\left(\sqrt{c}\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right) + \sqrt{a-b+c}\text{arctanh}\left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)\right)}{2\sqrt{c}\sqrt{a-b+ce}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}$$

[In] Integrate[Cot[d + e*x]^3/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] -1/2*((Sqrt[c]*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]) + Sqrt[a - b + c]*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)])]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/(Sqrt[c]*Sqrt[a - b + c]*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.09

method	result
derivativedivides	$\frac{\ln\left(\frac{\frac{b}{2}+c \cot(ex+d)^2}{\sqrt{c}}+\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{2\sqrt{c}} - \frac{\ln\left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c}\sqrt{(\cot(ex+d)^2+1)^2 c}}{\cot(ex+d)^2+1}\right)}{2\sqrt{a-b+c}}$
default	$\frac{\ln\left(\frac{\frac{b}{2}+c \cot(ex+d)^2}{\sqrt{c}}+\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{2\sqrt{c}} - \frac{\ln\left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c}\sqrt{(\cot(ex+d)^2+1)^2 c}}{\cot(ex+d)^2+1}\right)}{2\sqrt{a-b+c}}$

[In] `int(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x,method=_RETURNVE
RBOSE)`

[Out] $\frac{1}{e} \left(-\frac{1}{2} \ln\left(\frac{(1/2)b+c \cot(ex+d)^2}{c} + \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right) / c^{1/2} + (a+b \cot(ex+d)^2+c \cot(ex+d)^4)^{1/2} / c^{1/2} - \frac{1}{2} (a-b+c)^{1/2} \ln\left(\frac{(2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c}\sqrt{(\cot(ex+d)^2+1)^2 c}}{\cot(ex+d)^2+1}\right) + 2(a-b+c)^{1/2} \left(\frac{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}{(\cot(ex+d)^2+1)}\right) \right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 406 vs. 2(121) = 242.

Time = 1.03 (sec) , antiderivative size = 1695, normalized size of antiderivative = 12.02

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \text{Too large to display}$$

[In] `integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm
="fricas")`

[Out] $\frac{1}{4} \left(\sqrt{a-b+c} \log(2(a^2-2ab+b^2+2(a-b)c+c^2)) \cos(2ex+2d)^2 + 2a^2 - b^2 + 2c^2 - 2((a-b+c)\cos(2ex+2d)^2 - (2a-b)\cos(2ex+2d) + a-c)\sqrt{a-b+c} \sqrt{((a-b+c)\cos(2ex+2d)^2 - 2(a-c)\cos(2ex+2d) + a+b+c)/(\cos(2ex+2d)^2 - 2\cos(2ex+2d) + 1)} - 4(a^2 - ab + bc - c^2)\cos(2ex+2d) + (a-b+c)\sqrt{c} \log(((b^2+4(a-2b)c+8c^2)\cos(2ex+2d)^2 + b^2+4(a+2b)c+8c^2-4((b-2c)\cos(2ex+2d)^2-2b\cos(2ex+2d)+b+2c)\sqrt{c})\sqrt{((a-b+c)\cos(2ex+2d)^2-2(a-c)\cos(2ex+2d)+a+b+c)/(\cos(2ex+2d)^2-2\cos(2ex+2d)+1)} - 2(b^2+4ac-8c^2)\cos(2ex+2d) \right) / (((a-b)c+c^2)e), -\frac{1}{4} \left(2(a-b+c)\sqrt{-c} \arctan\left(-\frac{1}{2} \frac{(b-2c)\cos(2ex+2d)^2-2b\cos(2ex+2d)+b+2c}{(\cot(ex+d)^2+1)}\right) \sqrt{-c} \sqrt{((a-b+c)\cos(2ex+2d)^2-2(a-c)\cos(2ex+2d)+a+b+c)} \right)$

```

+ a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1))/(((a - b)*c +
c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d))
) - sqrt(a - b + c)*c*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e
*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 - 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*
a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*
x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 -
2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)))/(((
a - b)*c + c^2)*e), 1/4*(2*sqrt(-a + b - c)*c*arctan(((a - b + c)*cos(2*e*x
+ 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a -
b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2
*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c
+ c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c -
c^2)*cos(2*e*x + 2*d))) + (a - b + c)*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8
*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 - 4*((b - 2*c)*cos(2
*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*c
os(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*
d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))
/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c + c^2)*e), 1/2
*(sqrt(-a + b - c)*c*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos
(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^
2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e
*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2
+ a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d))) -
(a - b + c)*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2
*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a
- c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d
) + 1)))/(((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c -
c^2)*cos(2*e*x + 2*d)))/(((a - b)*c + c^2)*e)]

```

Sympy [F]

$$\int \frac{\cot^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx = \int \frac{\cot^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx$$

[In] integrate(cot(e*x+d)**3/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(cot(d + e*x)**3/sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4), x)

Maxima [F]

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(ex+d)^3}{\sqrt{c\cot(ex+d)^4+b\cot(ex+d)^2+a}} dx$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)^3/sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(d+ex)^3}{\sqrt{c\cot(d+ex)^4+b\cot(d+ex)^2+a}} dx$$

[In] int(cot(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(cot(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

$$3.19 \quad \int \frac{\cot(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

Optimal result	199
Rubi [A] (verified)	199
Mathematica [A] (verified)	200
Maple [A] (verified)	201
Fricas [B] (verification not implemented)	201
Sympy [F]	202
Maxima [F]	202
Giac [F(-1)]	203
Mupad [F(-1)]	203

Optimal result

Integrand size = 33, antiderivative size = 79

$$\int \frac{\cot(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+c}}$$

[Out] 1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/(a-b+c)^(1/2)

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$, Rules used = {3782, 1261, 738, 212}

$$\int \frac{\cot(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e\sqrt{a-b+c}}$$

[In] Int[Cot[d + e*x]/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*Sqrt[a - b + c]*e)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt Q[a, 0] || LtQ[b, 0])

Rule 738

```
Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol]
:> Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 1261

```
Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol]
:> Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)])*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)])*(f_.))^(n2_.))^(p_), x_Symbol]
:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n 2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x}{(1+x^2)\sqrt{a+bx^2+cx^4}} dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\ &= \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\ &= \frac{\text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.58 (sec) , antiderivative size = 139, normalized size of antiderivative = 1.76

$$\begin{aligned} &\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx \\ &= \frac{\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right) \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} \tan^2(d+ex)}{2\sqrt{a-b+ce}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}} \end{aligned}$$

[In] Integrate[Cot[d + e*x]/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4],x]

[Out] (ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 102, normalized size of antiderivative = 1.29

method	result	size
derivativedivides	$\frac{\ln\left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c}\sqrt{(\cot(ex+d)^2+1)^2c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}}{\cot(ex+d)^2+1}\right)}{2e\sqrt{a-b+c}}$	102
default	$\frac{\ln\left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c}\sqrt{(\cot(ex+d)^2+1)^2c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}}{\cot(ex+d)^2+1}\right)}{2e\sqrt{a-b+c}}$	102

[In] int(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x,method=_RETURNVERB
OSE)

[Out] 1/2/e/(a-b+c)^(1/2)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2)*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 222 vs. 2(69) = 138.

Time = 0.46 (sec) , antiderivative size = 433, normalized size of antiderivative = 5.48

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx$$

$$= \frac{\log\left(2(a^2-2ab+b^2+2(a-b)c+c^2)\cos(2ex+2d)^2+2a^2-b^2+2c^2+2((a-b+c)\cos(2ex+2d)+\sqrt{a+b+c})\cos(2ex+2d)\right)}{2(a-b+c)e} + \frac{\sqrt{-a+b-c}\arctan\left(\frac{((a-b+c)\cos(2ex+2d)^2-(2a-b)\cos(2ex+2d)+a-c)\sqrt{-a+b-c}\sqrt{\frac{(a-b+c)\cos(2ex+2d)^2-2(a-c)\cos(2ex+2d)+a-b+c}{\cos(2ex+2d)^2-2\cos(2ex+2d)+1}}}{(a^2-2ab+b^2+2(a-b)c+c^2)\cos(2ex+2d)^2+a^2-b^2+2ac+c^2-2(a^2-ab+bc-c^2)\cos(2ex+2d)}\right)}{2(a-b+c)e}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")

[Out] [1/4*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d))/(sqrt(a - b + c)*e), -1/2*sqrt(-a + b - c)*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)))/((a - b + c)*e)]

Sympy [F]

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(cot(d + e*x)/sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4), x)

Maxima [F]

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(ex+d)}{\sqrt{c\cot(ex+d)^4+b\cot(ex+d)^2+a}} dx$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)/sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

```
[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="
giac")
```

```
[Out] Timed out
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\cot(d+ex)}{\sqrt{c\cot^4(d+ex)+b\cot^2(d+ex)+a}} dx$$

```
[In] int(cot(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)
```

```
[Out] int(cot(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)
```

$$3.20 \quad \int \frac{\tan(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

Optimal result	204
Rubi [A] (verified)	204
Mathematica [A] (verified)	206
Maple [F]	206
Fricas [B] (verification not implemented)	207
Sympy [F]	208
Maxima [F]	208
Giac [F(-1)]	208
Mupad [F(-1)]	208

Optimal result

Integrand size = 33, antiderivative size = 142

$$\int \frac{\tan(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ae}} - \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}}$$

[Out] 1/2*arctanh(1/2*(2*a+b*cot(e*x+d)^2)/a^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/a^(1/2)-1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/(a-b+c)^(1/2)

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {3782, 1265, 974, 738, 212}

$$\int \frac{\tan(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ae}} - \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c) \cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e\sqrt{a-b+c}}$$

[In] Int[Tan[d + e*x]/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*Sqrt[a]*e) - ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a-b+c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*e*Sqrt[a-b+c])

2)/(2*sqrt[a - b + c]*sqrt[a + b*cot[d + e*x]^2 + c*cot[d + e*x]^4])/(2*sqrt[a - b + c]*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_) + (e_)*(x_))*sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 974

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n2_))^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x^2)\sqrt{a+bx^2+cx^4}} dx, x, \cot(d+ex)\right)}{e} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \end{aligned}$$

$$\begin{aligned}
&= - \frac{\text{Subst}\left(\int \left(\frac{1}{(-1-x)\sqrt{a+bx+cx^2}} + \frac{1}{x\sqrt{a+bx+cx^2}}\right) dx, x, \cot^2(d+ex)\right)}{2e} \\
&= - \frac{\text{Subst}\left(\int \frac{1}{(-1-x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= \frac{\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{-2a+b-(b-2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
&= \frac{\text{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{a}e} - \frac{\text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{a-b+c}e}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.19 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.39

$$\begin{aligned}
&\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx \\
&\left(\frac{\text{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{\sqrt{a}} - \frac{\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{\sqrt{a-b+c}} \right) \cot^2(d+ex)\sqrt{c+b\tan^2(d+ex)} \\
&= \frac{\left(\frac{\text{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{\sqrt{a}} - \frac{\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{\sqrt{a-b+c}} \right) \cot^2(d+ex)\sqrt{c+b\tan^2(d+ex)}}{2e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
\end{aligned}$$

[In] Integrate[Tan[d + e*x]/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] ((ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]/Sqrt[a] - ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]/Sqrt[a - b + c])*Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])/(2*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])

Maple [F]

$$\int \frac{\tan(ex+d)}{\sqrt{a+b\cot(ex+d)^2+c\cot(ex+d)^4}} dx$$

[In] int(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2), x)

[Out] int(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2), x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(122) = 244.

Time = 1.01 (sec) , antiderivative size = 1141, normalized size of antiderivative = 8.04

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Too large to display}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")

[Out] [1/4*((a - b + c)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)) + sqrt(a - b + c)*a*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)))/(a^2 - a*b + a*c)*e), -1/4*(2*sqrt(-a)*(a - b + c)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(a^2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)) - sqrt(a - b + c)*a*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)))/(a^2 - a*b + a*c)*e), -1/4*(2*a*sqrt(-a + b - c)*arctan(-1/2*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(-a + b - c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(a^2 - a*b + a*c)*tan(e*x + d)^4 + (a*b - b^2 + b*c)*tan(e*x + d)^2 + (a - b)*c + c^2)) - (a - b + c)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)))/(a^2 - a*b + a*c)*e), -1/2*(sqrt(-a)*(a - b + c)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(a^2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)) + a*sqrt(-a + b - c)*arctan(-1/2*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(-a + b - c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(a^2 - a*b + a*c)*tan(e*x + d)^4 + (a*b - b^2 + b*c)*tan(e*x + d)^2 + (a - b)*c + c^2)))/(a^2 - a*b + a*c)*e]

Sympy [F]

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(tan(d + e*x)/sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4), x)

Maxima [F]

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\tan(ex+d)}{\sqrt{c\cot(ex+d)^4+b\cot(ex+d)^2+a}} dx$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(tan(e*x + d)/sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\tan(d+ex)}{\sqrt{c\cot(d+ex)^4+b\cot(d+ex)^2+a}} dx$$

[In] int(tan(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(tan(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

$$3.21 \quad \int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx$$

Optimal result	209
Rubi [A] (verified)	209
Mathematica [A] (verified)	212
Maple [F]	213
Fricas [A] (verification not implemented)	213
Sympy [F]	214
Maxima [F(-1)]	214
Giac [F(-1)]	215
Mupad [F(-1)]	215

Optimal result

Integrand size = 35, antiderivative size = 249

$$\begin{aligned} & \int \frac{\tan^3(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}} dx \\ &= -\frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ae}} - \frac{b \operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4a^{3/2}e} \\ & \quad + \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} \\ & \quad + \frac{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} \tan^2(d+ex)}{2ae} \end{aligned}$$

[Out] $-1/4*b*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d)^2)/a^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/a^{(3/2)}/e-1/2*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d)^2)/a^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/e/a^{(1/2)}+1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/e/(a-b+c)^{(1/2)}+1/2*(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}*\tan(e*x+d)^2/a/e$

Rubi [A] (verified)

Time = 0.46 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used

= {3782, 1265, 974, 744, 738, 212}

$$\int \frac{\tan^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx$$

$$= -\frac{\operatorname{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4a^{3/2}e} - \frac{\operatorname{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2\sqrt{ae}}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e\sqrt{a-b+c}}$$

$$+ \frac{\tan^2(d+ex)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2ae}$$

[In] Int[Tan[d + e*x]^3/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] -1/2*ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(Sqrt[a]*e) - (b*ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])])/(4*a^(3/2)*e) + ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*Sqrt[a - b + c]*e) + (Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/(2*a*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 744

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/((m + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[(2*c*d - b*e)/(2*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && EqQ[m + 2*p + 3, 0]

Rule 974

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g

$*x)^n*(a + b*x + c*x^2)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, g\}, x] \&\& \text{NeQ}[e*f - d*g, 0] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& (\text{IntegerQ}[p] \mid\mid (\text{ILtQ}[m, 0] \&\& \text{ILtQ}[n, 0])) \&\& !(\text{IGtQ}[m, 0] \mid\mid \text{IGtQ}[n, 0])$

Rule 1265

$\text{Int}[(x_)^{(m_.)*((d_) + (e_.)*(x_)^2)^{(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^{(p_.)}, x_Symbol] := \text{Dist}[1/2, \text{Subst}[\text{Int}[x^{((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p}, x], x, x^2], x] /; \text{FreeQ}[\{a, b, c, d, e, p, q\}, x] \&\& \text{IntegerQ}[(m - 1)/2]$

Rule 3782

$\text{Int}[\cot[(d_.) + (e_.)*(x_)]^{(m_.)*((a_.) + (b_.)*\cot[(d_.) + (e_.)*(x_)]*(f_.))^{(n_.) + (c_.)*\cot[(d_.) + (e_.)*(x_)]*(f_.))^{(n2_.))^{(p_.)}, x_Symbol] := \text{Dist}[-f/e, \text{Subst}[\text{Int}[(x/f)^m*((a + b*x^n + c*x^{(2*n)})^p/(f^2 + x^2)), x], x, f*\text{Cot}[d + e*x]], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{EqQ}[n^2, 2*n] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x^3(1+x^2)\sqrt{a+bx^2+cx^4}} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^2(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^2\sqrt{a+bx+cx^2}} - \frac{1}{x\sqrt{a+bx+cx^2}} + \frac{1}{(1+x)\sqrt{a+bx+cx^2}}\right) dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^2\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2ae} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&\quad + \frac{b \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{4ae} \\
&= -\frac{\text{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ae}} \\
&\quad + \frac{\text{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} \\
&\quad + \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2ae} \\
&\quad - \frac{b \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2ae} \\
&= -\frac{\text{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{ae}} - \frac{b \text{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4a^{3/2}e} \\
&\quad + \frac{\text{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2\sqrt{a-b+ce}} \\
&\quad + \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2ae}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.38 (sec) , antiderivative size = 258, normalized size of antiderivative = 1.04

$$\begin{aligned}
&\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx \\
&= \frac{\cot^2(d + ex) \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)} \left(-\left((2a + b) \sqrt{a - b} + \text{arctanh}\left(\frac{b + 2a \tan^2(d + ex)}{2\sqrt{a}\sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}} \right) \right) \right)}{4a^{3/2} \sqrt{a - b + ce} \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}
\end{aligned}$$

[In] Integrate[Tan[d + e*x]^3/Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] (Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]*(-(2*a + b)*Sqrt[a - b + c]*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)])) + 2*Sqrt[a]*(a*ArcTanh[(b - 2*c + (2*a -

b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])) + Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))/(4*a^(3/2)*Sqrt[a - b + c]*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]))

Maple [F]

$$\int \frac{\tan^3(ex + d)}{\sqrt{a + b \cot^2(ex + d) + c \cot^4(ex + d)}} dx$$

[In] int(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x)

[Out] int(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x)

Fricas [A] (verification not implemented)

none

Time = 1.38 (sec) , antiderivative size = 1444, normalized size of antiderivative = 5.80

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx = \text{Too large to display}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")

[Out] [1/8*(2*sqrt(a - b + c)*a^2*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 + 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)) + 4*(a^2 - a*b + a*c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 + (2*a^2 - a*b - b^2 + (2*a + b)*c)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c - 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)))/((a^3 - a^2*b + a^2*c)*e), 1/4*(sqrt(a - b + c)*a^2*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 + 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)) + 2*(a^2 - a*b + a*c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 + (2*a^2 - a*b - b^2 + (2*a + b)*c)*sqrt(-a)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)/(a^2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)))/((a^3 - a^2*b + a^2*c)*e), 1/8*(4*a^2*sqrt(-a + b - c)*arctan(-1/2*((2*a - b)*tan(e*x +

$$d)^4 + (b - 2c) \tan(e^x + d)^2 \sqrt{-a + b - c} \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}} / \left((a^2 - a^2 b + a^2 c) \tan(e^x + d)^4 + (a^2 b - b^2 + b^2 c) \tan(e^x + d)^2 + (a - b) c + c^2 \right) + 4(a^2 - a^2 b + a^2 c) \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}} \tan(e^x + d)^2 + (2a^2 - a^2 b - b^2 + (2a + b)c) \sqrt{a} \log(8a^2 \tan(e^x + d)^4 + 8a^2 b \tan(e^x + d)^2 + b^2 + 4a^2 c - 4(2a \tan(e^x + d))^4 + b \tan(e^x + d)^2) \sqrt{a} \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}}) / \left((a^3 - a^2 b + a^2 c) e \right), \frac{1}{4} (2a^2 \sqrt{-a + b - c} \arctan(-1/2 * ((2a - b) \tan(e^x + d))^4 + (b - 2c) \tan(e^x + d)^2) \sqrt{-a + b - c} \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}} / \left((a^2 - a^2 b + a^2 c) \tan(e^x + d)^4 + (a^2 b - b^2 + b^2 c) \tan(e^x + d)^2 + (a - b) c + c^2 \right) + 2(a^2 - a^2 b + a^2 c) \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}} \tan(e^x + d)^2 + (2a^2 - a^2 b - b^2 + (2a + b)c) \sqrt{-a} \arctan(1/2 * (2a \tan(e^x + d))^4 + b \tan(e^x + d)^2) \sqrt{-a} \sqrt{\frac{(a \tan(e^x + d))^4 + b \tan(e^x + d)^2 + c}{\tan(e^x + d)^4}} / (a^2 \tan(e^x + d)^4 + a^2 b \tan(e^x + d)^2 + a^2 c)) / \left((a^3 - a^2 b + a^2 c) e \right)]$$

Sympy [F]

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx = \int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx$$

[In] integrate(tan(e*x+d)**3/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(tan(d + e*x)**3/sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4), x)

Maxima [F(-1)]

Timed out.

$$\int \frac{\tan^3(d + ex)}{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] Timed out

Giac [F(-1)]

Timed out.

$$\int \frac{\tan^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^3(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} dx = \int \frac{\tan(d+ex)^3}{\sqrt{c\cot(d+ex)^4+b\cot(d+ex)^2+a}} dx$$

[In] int(tan(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(tan(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

3.22 $\int \cot^5(d+ex) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)} dx$

Optimal result	216
Rubi [A] (verified)	217
Mathematica [B] (verified)	220
Maple [A] (verified)	221
Fricas [B] (verification not implemented)	222
Sympy [F]	224
Maxima [F]	224
Giac [F(-1)]	224
Mupad [F(-1)]	225

Optimal result

Integrand size = 35, antiderivative size = 270

$$\begin{aligned}
 & \int \cot^5(d+ex) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)} dx \\
 &= \frac{\sqrt{a-b+c} \operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
 & \quad - \frac{(b^3 + 2b^2c - 4b(a-2c)c - 8c^2(a+2c)) \operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{32c^{5/2}e} \\
 & \quad + \frac{((b-2c)(b+4c) + 2c(b+2c)\cot^2(d+ex)) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}}{16c^2e} \\
 & \quad - \frac{(a + b \cot^2(d+ex) + c \cot^4(d+ex))^{3/2}}{6ce}
 \end{aligned}$$

```
[Out] -1/32*(b^3+2*b^2*c-4*b*(a-2*c)*c-8*c^2*(a+2*c))*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/c^(5/2)/e-1/6*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2)/c/e+1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*(a-b+c)^(1/2)/e+1/16*((b-2*c)*(b+4*c)+2*c*(b+2*c)*cot(e*x+d)^2)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)/c^2/e
```


Rubi [A] (verified)

Time = 0.68 (sec) , antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3782, 1265, 1667, 828, 857, 635, 212, 738}

$$\int \cot^5(d+ex) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} dx$$

$$= -\frac{(-4bc(a-2c)-8c^2(a+2c)+b^3+2b^2c) \operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{32c^{5/2}e}$$

$$+ \frac{\sqrt{a-b} + c \operatorname{arctanh}\left(\frac{2a+(b-2c) \cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e}$$

$$+ \frac{(2c(b+2c) \cot^2(d+ex)+(b-2c)(b+4c)) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}{16c^2e}$$

$$- \frac{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}}{6ce}$$

[In] Int[Cot[d + e*x]^5*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] (Sqrt[a - b + c]*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]])/(2*e) - ((b^3 + 2*b^2*c - 4*b*(a - 2*c)*c - 8*c^2*(a + 2*c))*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]])/(32*c^(5/2)*e) + (((b - 2*c)*(b + 4*c) + 2*c*(b + 2*c)*Cot[d + e*x]^2)*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])/(16*c^2*e) - (a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2)/(6*c*e)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 828

```
Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*((a + b*x + c*x^2)^p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), x] - Dist[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Rule 857

```
Int[((d_.) + (e_.)*(x_))^(m_.)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1265

```
Int[(x_)^(m_.)*((d_.) + (e_.)*(x_)^2)^(q_.)*((a_.) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]
```

Rule 1667

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)])*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)])*(f_.))^(n2_.))^(p_), x_Symbol]
```

```

:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5 \sqrt{a+bx^2+cx^4}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x^2 \sqrt{a+bx+cx^2}}{1+x} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}{6ce} \\
&\quad -\frac{\text{Subst}\left(\int \frac{\left(-\frac{3b}{2}-\frac{3}{2}(b+2c)x\right)\sqrt{a+bx+cx^2}}{1+x} dx, x, \cot^2(d+ex)\right)}{6ce} \\
&= \frac{((b-2c)(b+4c)+2c(b+2c)\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{16c^2e} \\
&\quad -\frac{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}{6ce} \\
&\quad +\frac{\text{Subst}\left(\int \frac{-\frac{3}{4}(b-2c)(b^2-4ac+4bc)-\frac{3}{4}(b^3+2b^2c-4b(a-2c)c-8c^2(a+2c))x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{24c^2e} \\
&= \frac{((b-2c)(b+4c)+2c(b+2c)\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{16c^2e} \\
&\quad -\frac{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}{6ce} \\
&\quad -\frac{(a-b+c)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&\quad -\frac{(b^3+2b^2c-4b(a-2c)c-8c^2(a+2c))\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{32c^2e} \\
&= \frac{((b-2c)(b+4c)+2c(b+2c)\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{16c^2e} \\
&\quad -\frac{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}{6ce} \\
&\quad +\frac{(a-b+c)\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
&\quad -\frac{(b^3+2b^2c-4b(a-2c)c-8c^2(a+2c))\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{16c^2e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a-b} + \operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&\quad - \frac{(b^3+2b^2c-4b(a-2c)c-8c^2(a+2c))\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{32c^{5/2}e} \\
&\quad + \frac{((b-2c)(b+4c)+2c(b+2c)\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{16c^2e} \\
&\quad - \frac{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}{6ce}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1017 vs. $2(270) = 540$.

Time = 6.50 (sec) , antiderivative size = 1017, normalized size of antiderivative = 3.77

$$\begin{aligned}
&\int \cot^5(d+ex)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}dx = \\
&\quad \left(\frac{b\operatorname{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{2\sqrt{a}} - \sqrt{c}\operatorname{arctanh}\left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right) + \frac{1}{2}\left(\frac{(2a-b)\operatorname{arctanh}\left(\frac{\tan^2(d+ex)\sqrt{\cot^4(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))}}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{4e\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right.\right. \\
&\quad + \frac{\tan^2(d+ex)\sqrt{\cot^4(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))}}{16e\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\left.\left(\frac{(b^2-4ac)\operatorname{arctanh}\left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{c^{3/2}}\right)\right. \\
&\quad \left. + \frac{\tan^2(d+ex)\sqrt{\cot^4(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))}}{16e\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\left(\frac{16\cot^6(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))^{3/2}}{c}\right)\right) \\
&\quad \left. - \frac{96e\sqrt{c+b\tan^2(d+ex)}}{96e\sqrt{c+b\tan^2(d+ex)}}\right)
\end{aligned}$$

[In] Integrate[Cot[d + e*x]^5*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] -1/2*(((b*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/(2*Sqrt[a]) - Sqrt[c]*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])])

$$\begin{aligned}
& e^x]^2)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4]) + (((2*a \\
& - b)*\text{ArcTanh}[(b + 2*a*\text{Tan}[d + e^x]^2)/(2*\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 \\
& + a*\text{Tan}[d + e^x]^4))]/\text{Sqrt}[a] - (4*\text{Sqrt}[a - b + c]*(2*a - 2*b + 2*c)*\text{ArcT} \\
& \text{anh}[(b - 2*c - (-2*a + b)*\text{Tan}[d + e^x]^2)/(2*\text{Sqrt}[a - b + c]*\text{Sqrt}[c + b*\text{Tan} \\
& [d + e^x]^2 + a*\text{Tan}[d + e^x]^4))]/(4*a - 4*b + 4*c))/2)*\text{Tan}[d + e^x]^2*\text{Sqr} \\
& \text{t}[\text{Cot}[d + e^x]^4*(c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)]/(e*\text{Sqrt}[c + b* \\
& \text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)) + (\text{Tan}[d + e^x]^2*\text{Sqrt}[\text{Cot}[d + e^x]^4*(\\
& c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)]*(2*\text{Sqrt}[a]*\text{ArcTanh}[(b + 2*a*\text{Tan}[d \\
& + e^x]^2)/(2*\text{Sqrt}[a]*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)]) - (b* \\
& \text{ArcTanh}[(2*c + b*\text{Tan}[d + e^x]^2)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*T \\
& \text{an}[d + e^x]^4))]/\text{Sqrt}[c] - 2*\text{Cot}[d + e^x]^2*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a* \\
& \text{Tan}[d + e^x]^4)]/(4*e*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)) - (\text{Ta} \\
& \text{n}[d + e^x]^2*\text{Sqrt}[\text{Cot}[d + e^x]^4*(c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)] \\
& *(((b^2 - 4*a*c)*\text{ArcTanh}[(2*c + b*\text{Tan}[d + e^x]^2)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan} \\
& [d + e^x]^2 + a*\text{Tan}[d + e^x]^4)))/c^(3/2) - (2*\text{Cot}[d + e^x]^4*(2*c + b*\text{Tan} \\
& [d + e^x]^2)*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4))/c))/(16*e*\text{Sqrt}[\\
& c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)) - (\text{Tan}[d + e^x]^2*\text{Sqrt}[\text{Cot}[d + e \\
& x]^4*(c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)]*((16*\text{Cot}[d + e^x]^6*(c + b* \\
& \text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4)^(3/2))/c + (3*b*(((b^2 - 4*a*c)*\text{ArcTanh} \\
& (2*c + b*\text{Tan}[d + e^x]^2)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 + a*\text{Tan}[d + e \\
& *x]^4)))/c^(3/2) - (2*\text{Cot}[d + e^x]^4*(2*c + b*\text{Tan}[d + e^x]^2)*\text{Sqrt}[c + b*T \\
& \text{an}[d + e^x]^2 + a*\text{Tan}[d + e^x]^4))/c))/c))/(96*e*\text{Sqrt}[c + b*\text{Tan}[d + e^x]^2 \\
& + a*\text{Tan}[d + e^x]^4])
\end{aligned}$$

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 455, normalized size of antiderivative = 1.69

method	result
derivativedivides	$ -\frac{(a+b \cot (e x+d)^2+c \cot (e x+d)^4)^{\frac{3}{2}}}{6 c}+\frac{b\left(\frac{(b+2 c \cot (e x+d)^2) \sqrt{a+b \cot (e x+d)^2+c \cot (e x+d)^4}}{4 c}+\frac{(4 a c-b^2) \ln \left(\frac{\frac{b}{2}+c \cot (e x+d)^2}{\sqrt{c}}+\sqrt{\frac{a+b \cot (e x+d)^2+c \cot (e x+d)^4}{c}}\right)}{4 c}\right)}{4 c} $
default	$ -\frac{(a+b \cot (e x+d)^2+c \cot (e x+d)^4)^{\frac{3}{2}}}{6 c}+\frac{b\left(\frac{(b+2 c \cot (e x+d)^2) \sqrt{a+b \cot (e x+d)^2+c \cot (e x+d)^4}}{4 c}+\frac{(4 a c-b^2) \ln \left(\frac{\frac{b}{2}+c \cot (e x+d)^2}{\sqrt{c}}+\sqrt{\frac{a+b \cot (e x+d)^2+c \cot (e x+d)^4}{c}}\right)}{4 c}\right)}{4 c} $

[In] int(cot(e*x+d)^5*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x,method=_RETURNVE
RBOSE)

[Out] 1/e*(-1/6*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2)/c+1/4*b/c*(1/4*(b+2*c*cot
(e*x+d)^2)/c*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)+1/8*(4*a*c-b^2)/c^(3/2
)*ln(((1/2*b+c*cot(e*x+d)^2)/c^(1/2)+(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2
))+1/8*(b+2*c*cot(e*x+d)^2)/c*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)+1/16*

$$(4*a*c-b^2)/c^{(3/2)}*\ln((1/2*b+c*\cot(e*x+d)^2)/c^{(1/2)}+(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})-1/2*((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^{(1/2)}-1/4*(b-2*c)*\ln((1/2*b-c+(\cot(e*x+d)^2+1)*c)/c^{(1/2)}+((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^{(1/2)})/c^{(1/2)}+1/2*(a-b+c)^{(1/2)}*\ln((2*a-2*b+2*c+(b-2*c)*(\cot(e*x+d)^2+1)+2*(a-b+c)^{(1/2)}*((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^{(1/2)})/(\cot(e*x+d)^2+1))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 737 vs. 2(242) = 484.

Time = 2.95 (sec) , antiderivative size = 3019, normalized size of antiderivative = 11.18

$$\int \cot^5(d+ex)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")

[Out] [1/192*(48*(c^3*cos(2*e*x + 2*d)^2 - 2*c^3*cos(2*e*x + 2*d) + c^3)*sqrt(a - b + c)*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) - 3*(b^3 - 8*(a - b)*c^2 - 16*c^3 + (b^3 - 8*(a - b)*c^2 - 16*c^3 - 2*(2*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - 2*(2*a*b - b^2)*c - 2*(b^3 - 8*(a - b)*c^2 - 16*c^3 - 2*(2*a*b - b^2)*c)*cos(2*e*x + 2*d))*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 + 4*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) + 4*(3*b^2*c - 4*(2*a - b)*c^2 - 2*0*c^3 + (3*b^2*c - 8*(a - b)*c^2 - 44*c^3)*cos(2*e*x + 2*d)^2 - 2*(3*b^2*c - 2*(4*a - 3*b)*c^2 - 16*c^3)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c^3*e*cos(2*e*x + 2*d)^2 - 2*c^3*e*cos(2*e*x + 2*d) + c^3*e), -1/96*(3*(b^3 - 8*(a - b)*c^2 - 16*c^3 + (b^3 - 8*(a - b)*c^2 - 16*c^3 - 2*(2*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - 2*(2*a*b - b^2)*c - 2*(b^3 - 8*(a - b)*c^2 - 16*c^3 - 2*(2*a*b - b^2)*c)*cos(2*e*x + 2*d))*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d)) - 24*(c^3*cos(2*e*x + 2*d)^2 - 2*c^3*cos(2*e*x + 2*d) + c^3)*sqrt(a - b + c)*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a

$$\begin{aligned}
&^2 - b^2 + 2c^2 + 2*((a - b + c)*\cos(2ex + 2d))^2 - (2a - b)*\cos(2ex + 2d) + a - c)*\sqrt{a - b + c}*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)} - 4*(a^2 - ab + bc - c^2)*\cos(2ex + 2d) - 2*(3b^2c - 4*(2a - b)*c^2 - 20c^3 + (3b^2c - 8*(a - b)*c^2 - 44c^3)*\cos(2ex + 2d))^2 - 2*(3b^2c - 2*(4a - 3b)*c^2 - 16c^3)*\cos(2ex + 2d))*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/(c^3e*\cos(2ex + 2d))^2 - 2*c^3e*\cos(2ex + 2d) + c^3e), -1/192*(96*(c^3*\cos(2ex + 2d))^2 - 2*c^3*\cos(2ex + 2d) + c^3)*\sqrt{-a + b - c}*\arctan(((a - b + c)*\cos(2ex + 2d))^2 - (2a - b)*\cos(2ex + 2d) + a - c)*\sqrt{-a + b - c}*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/((a^2 - 2ab + b^2 + 2*(a - b)*c + c^2)*\cos(2ex + 2d))^2 + a^2 - b^2 + 2ac + c^2 - 2*(a^2 - ab + bc - c^2)*\cos(2ex + 2d))) + 3*(b^3 - 8*(a - b)*c^2 - 16c^3 + (b^3 - 8*(a - b)*c^2 - 16c^3 - 2*(2ab - b^2)*c)*\cos(2ex + 2d))^2 - 2*(2ab - b^2)*c - 2*(b^3 - 8*(a - b)*c^2 - 16c^3 - 2*(2ab - b^2)*c)*\cos(2ex + 2d))*\sqrt{c}*\log(((b^2 + 4*(a - 2b)*c + 8c^2)*\cos(2ex + 2d))^2 + b^2 + 4*(a + 2b)*c + 8c^2 + 4*((b - 2c)*\cos(2ex + 2d))^2 - 2b*\cos(2ex + 2d) + b + 2c)*\sqrt{c}*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)} - 2*(b^2 + 4ac - 8c^2)*\cos(2ex + 2d))/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)} - 4*(3b^2c - 4*(2a - b)*c^2 - 20c^3 + (3b^2c - 8*(a - b)*c^2 - 44c^3)*\cos(2ex + 2d))^2 - 2*(3b^2c - 2*(4a - 3b)*c^2 - 16c^3)*\cos(2ex + 2d))*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/(c^3e*\cos(2ex + 2d))^2 - 2*c^3e*\cos(2ex + 2d) + c^3e), -1/96*(48*(c^3*\cos(2ex + 2d))^2 - 2*c^3*\cos(2ex + 2d) + c^3)*\sqrt{-a + b - c}*\arctan(((a - b + c)*\cos(2ex + 2d))^2 - (2a - b)*\cos(2ex + 2d) + a - c)*\sqrt{-a + b - c}*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/((a^2 - 2ab + b^2 + 2*(a - b)*c + c^2)*\cos(2ex + 2d))^2 + a^2 - b^2 + 2ac + c^2 - 2*(a^2 - ab + bc - c^2)*\cos(2ex + 2d))) + 3*(b^3 - 8*(a - b)*c^2 - 16c^3 + (b^3 - 8*(a - b)*c^2 - 16c^3 - 2*(2ab - b^2)*c)*\cos(2ex + 2d))^2 - 2*(2ab - b^2)*c - 2*(b^3 - 8*(a - b)*c^2 - 16c^3 - 2*(2ab - b^2)*c)*\cos(2ex + 2d))*\sqrt{-c}*\arctan(-1/2*((b - 2c)*\cos(2ex + 2d))^2 - 2b*\cos(2ex + 2d) + b + 2c)*\sqrt{-c}*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/(((a - b)*c + c^2)*\cos(2ex + 2d))^2 + (a + b)*c + c^2 - 2*(ac - c^2)*\cos(2ex + 2d))) - 2*(3b^2c - 4*(2a - b)*c^2 - 20c^3 + (3b^2c - 8*(a - b)*c^2 - 44c^3)*\cos(2ex + 2d))^2 - 2*(3b^2c - 2*(4a - 3b)*c^2 - 16c^3)*\cos(2ex + 2d))*\sqrt{((a - b + c)*\cos(2ex + 2d))^2 - 2*(a - c)*\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d))^2 - 2*\cos(2ex + 2d) + 1)}}/(c^3e*\cos(2ex + 2d))^2 - 2*c^3e*\cos(2ex + 2d) + c^3e)]
\end{aligned}$$

Sympy [F]

$$\int \cot^5(d+ex) \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} dx$$

$$= \int \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} \cot^5(d+ex) dx$$

[In] integrate(cot(e*x+d)**5*(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)*cot(d + e*x)**5, x)

Maxima [F]

$$\int \cot^5(d+ex) \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} dx$$

$$= \int \sqrt{c\cot^4(ex+d)^4 + b\cot^2(ex+d)^2 + a} \cot^5(ex+d) dx$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)*cot(e*x + d)^5, x)

Giac [F(-1)]

Timed out.

$$\int \cot^5(d+ex) \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^5*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \cot^5(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx = \text{Hanged}$$

```
[In] int(cot(d + e*x)^5*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)
```

```
[Out] \text{Hanged}
```

3.23 $\int \cot^3(d+ex) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)} dx$

Optimal result	226
Rubi [A] (verified)	226
Mathematica [B] (verified)	229
Maple [A] (verified)	230
Fricas [B] (verification not implemented)	231
Sympy [F]	232
Maxima [F]	232
Giac [F(-1)]	233
Mupad [F(-1)]	233

Optimal result

Integrand size = 35, antiderivative size = 209

$$\int \cot^3(d+ex) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)} dx$$

$$= -\frac{\sqrt{a-b} \operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e}$$

$$+ \frac{(b^2 + 4bc - 4c(a + 2c)) \operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{16c^{3/2}e}$$

$$- \frac{(b - 4c + 2c\cot^2(d+ex)) \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}}{8ce}$$

[Out] 1/16*(b^2+4*b*c-4*c*(a+2*c))*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/c^(3/2)/e-1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*(a-b+c)^(1/2)/e-1/8*(b-4*c+2*c*cot(e*x+d)^2)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)/c/e

Rubi [A] (verified)

Time = 0.41 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used

= {3782, 1265, 828, 857, 635, 212, 738}

$$\int \cot^3(d+ex) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} dx$$

$$= \frac{(-4c(a+2c)+b^2+4bc) \operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{16c^{3/2}e}$$

$$- \frac{\sqrt{a-b} + \operatorname{arctanh}\left(\frac{2a+(b-2c) \cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e}$$

$$- \frac{(b+2c \cot^2(d+ex)-4c) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}{8ce}$$

[In] Int[Cot[d + e*x]^3*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4],x]

[Out] -1/2*(Sqrt[a - b + c]*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]])/e + ((b^2 + 4*b*c - 4*c*(a + 2*c))*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]])/(16*c^(3/2)*e) - ((b - 4*c + 2*c*Cot[d + e*x]^2)*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])/(8*c*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 828

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m+1)*(c*e*f*(m+2*p+2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m+2*p+1)*x)*((a + b*x + c*x^2)^p/(c*e^2*(m+2*p+1)*(m+2*p+2))), x] - Dist[p/(c*e^2*(m+2*p+1)*(m+2*p+2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p-1)*Simp[c*e*f*(b*d - 2*a*e)*(m+2*p+2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c

```
*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^
2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1))))*x, x], x], x]
/; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2
- b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[
m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p]
|| IntegersQ[2*m, 2*p])
```

Rule 857

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c
_.)*(x_)^2)^(p_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x +
c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p,
x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] &&
NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1265

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_
)^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a +
b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && Inte
gerQ[(m - 1)/2]
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^3\sqrt{a+bx^2+cx^4}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx+cx^2}}{1+x} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{(b-4c+2c\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{8ce} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{1}{2}(b^2+4ac-4bc)+\frac{1}{2}(b^2+4bc-4c(a+2c))x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{8ce}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(b-4c+2c\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{8ce} \\
&\quad + \frac{(a-b+c)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&\quad + \frac{(b^2+4bc-4c(a+2c))\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{16ce} \\
&= -\frac{(b-4c+2c\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{8ce} \\
&\quad - \frac{(a-b+c)\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&\quad + \frac{(b^2+4bc-4c(a+2c))\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{8ce} \\
&= -\frac{\sqrt{a-b} + \text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&\quad + \frac{(b^2+4bc-4c(a+2c))\text{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{16c^{3/2}e} \\
&\quad - \frac{(b-4c+2c\cot^2(d+ex))\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{8ce}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 777 vs. $2(209) = 418$.

Time = 6.36 (sec) , antiderivative size = 777, normalized size of antiderivative = 3.72

$$\begin{aligned}
&\int \cot^3(d+ex)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} dx \\
&= \left(\frac{\text{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{2\sqrt{a}} - \sqrt{c}\text{arctanh}\left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right) + \frac{1}{2} \left(\frac{(2a-b)\text{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{2\sqrt{a}} \right. \right. \\
&\quad \left. \left. - \frac{\tan^2(d+ex)\sqrt{\cot^4(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))}}{4e\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}} \right) \right) \\
&\quad + \frac{\tan^2(d+ex)\sqrt{\cot^4(d+ex)(c+b\tan^2(d+ex)+a\tan^4(d+ex))}}{16e\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}} \left(\frac{(b^2-4ac)\text{arctanh}\left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{c^{3/2}} \right)
\end{aligned}$$

[In] Integrate[Cot[d + e*x]^3*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] (((b*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/(2*Sqrt[a]) - Sqrt[c]*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]]) + (((2*a - b)*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/Sqrt[a] - (4*Sqrt[a - b + c]*(2*a - 2*b + 2*c)*ArcTanh[(b - 2*c - (-2*a + b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/(4*a - 4*b + 4*c))/2)*Tan[d + e*x]^2*Sqrt[Cot[d + e*x]^4*(c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]/(2*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)) - (Tan[d + e*x]^2*Sqrt[Cot[d + e*x]^4*(c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4))]*(2*Sqrt[a]*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)])) - (b*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/Sqrt[c] - 2*Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]/(4*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)) + (Tan[d + e*x]^2*Sqrt[Cot[d + e*x]^4*(c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4))*((b^2 - 4*a*c)*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]))/c^(3/2) - (2*Cot[d + e*x]^4*(2*c + b*Tan[d + e*x]^2)*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])/c)/(16*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 318, normalized size of antiderivative = 1.52

method	result
derivativedivides	$\frac{\frac{(b+2c \cot(ex+d)^2) \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{8c} - \frac{(4ac-b^2) \ln\left(\frac{\frac{b}{2}+c \cot(ex+d)}{\sqrt{c}} + \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{16c^{\frac{3}{2}}}}{\frac{(b+2c \cot(ex+d)^2) \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{8c} - \frac{(4ac-b^2) \ln\left(\frac{\frac{b}{2}+c \cot(ex+d)}{\sqrt{c}} + \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{16c^{\frac{3}{2}}}} + \sqrt{\cot(ex+d)^2 + a}}$
default	$\frac{\frac{(b+2c \cot(ex+d)^2) \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{8c} - \frac{(4ac-b^2) \ln\left(\frac{\frac{b}{2}+c \cot(ex+d)}{\sqrt{c}} + \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{16c^{\frac{3}{2}}}}{\frac{(b+2c \cot(ex+d)^2) \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{8c} - \frac{(4ac-b^2) \ln\left(\frac{\frac{b}{2}+c \cot(ex+d)}{\sqrt{c}} + \sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}\right)}{16c^{\frac{3}{2}}}} + \sqrt{\cot(ex+d)^2 + a}}$

[In] int(cot(e*x+d)^3*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2), x, method=_RETURNVE
RBOSE)

[Out] 1/e*(-1/8*(b+2*c*cot(e*x+d)^2)/c*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)-1/16*(4*a*c-b^2)/c^(3/2)*ln((1/2*b+c*cot(e*x+d)^2)/c^(1/2)+(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))+1/2*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2)+1/4*(b-2*c)*ln((1/2*b-c+(cot(e*x+d)^2+1)*c)/c^(1/2)+((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/c^(1/2)-1/2*(a-b+c)^(1/2)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2))*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 569 vs. 2(185) = 370.

Time = 2.34 (sec) , antiderivative size = 2344, normalized size of antiderivative = 11.22

$$\int \cot^3(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx = \text{Too large to display}$$

```
[In] integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm
="fricas")
```

```
[Out] [1/32*(8*(c^2*cos(2*e*x + 2*d) - c^2)*sqrt(a - b + c)*log(2*(a^2 - 2*a*b +
b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 - 2*((a -
b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b
+ c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a
+ b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b
*c - c^2)*cos(2*e*x + 2*d)) + (b^2 - 4*(a - b)*c - 8*c^2 - (b^2 - 4*(a - b)
*c - 8*c^2)*cos(2*e*x + 2*d))*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*co
s(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 - 4*((b - 2*c)*cos(2*e*x + 2
*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x
+ 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2
*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*
e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) + 4*(b*c - 2*c^2 - (b*c - 6*c^2)*co
s(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x
+ 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c^2*e*
cos(2*e*x + 2*d) - c^2*e), -1/16*((b^2 - 4*(a - b)*c - 8*c^2 - (b^2 - 4*(a
- b)*c - 8*c^2)*cos(2*e*x + 2*d))*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x
+ 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(
2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^
2 - 2*cos(2*e*x + 2*d) + 1)))/(((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b
)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d))) - 4*(c^2*cos(2*e*x + 2*d) - c^
2)*sqrt(a - b + c)*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x
+ 2*d)^2 + 2*a^2 - b^2 + 2*c^2 - 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a -
b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x +
2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*c
os(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) - 2*(b*
c - 2*c^2 - (b*c - 6*c^2)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2
*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos
(2*e*x + 2*d) + 1)))/(c^2*e*cos(2*e*x + 2*d) - c^2*e), 1/32*(16*(c^2*cos(2*
e*x + 2*d) - c^2)*sqrt(-a + b - c)*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 -
(2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos
(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)
^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(
2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*
e*x + 2*d))) + (b^2 - 4*(a - b)*c - 8*c^2 - (b^2 - 4*(a - b)*c - 8*c^2)*cos
```

```
(2*e*x + 2*d))*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 - 4*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) + 4*(b*c - 2*c^2 - (b*c - 6*c^2)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c^2*e*cos(2*e*x + 2*d) - c^2*e), 1/16*(8*(c^2*cos(2*e*x + 2*d) - c^2)*sqrt(-a + b - c)*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d))) - (b^2 - 4*(a - b)*c - 8*c^2 - (b^2 - 4*(a - b)*c - 8*c^2)*cos(2*e*x + 2*d))*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d))) + 2*(b*c - 2*c^2 - (b*c - 6*c^2)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c^2*e*cos(2*e*x + 2*d) - c^2*e)]
```

Sympy [F]

$$\int \cot^3(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \cot^3(d + ex) dx$$

```
[In] integrate(cot(e*x+d)**3*(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)
```

```
[Out] Integral(sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)*cot(d + e*x)**3, x)
```

Maxima [F]

$$\int \cot^3(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \int \sqrt{c \cot^4(ex + d) + b \cot^2(ex + d) + a} \cot^3(ex + d) dx$$

```
[In] integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")
```


[Out] integrate(sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)*cot(e*x + d)^3, x)

Giac **[F(-1)]**

Timed out.

$$\int \cot^3(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^3*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad **[F(-1)]**

Timed out.

$$\begin{aligned} & \int \cot^3(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx \\ &= \int \cot(d + ex)^3 \sqrt{c \cot^4(d + ex) + b \cot^2(d + ex) + a} dx \end{aligned}$$

[In] int(cot(d + e*x)^3*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(cot(d + e*x)^3*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

3.24 $\int \cot(d+ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$

Optimal result	234
Rubi [A] (verified)	234
Mathematica [A] (verified)	237
Maple [A] (verified)	237
Fricas [B] (verification not implemented)	238
Sympy [F]	239
Maxima [F]	239
Giac [F(-1)]	240
Mupad [F(-1)]	240

Optimal result

Integrand size = 33, antiderivative size = 179

$$\int \cot(d+ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \frac{\sqrt{a-b+c} \operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} - \frac{(b-2c)\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ce}} - \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e}$$

[Out] $-1/4*(b-2*c)*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d)^2)/c^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/e/c^{(1/2)}+1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})*(a-b+c)^{(1/2)}/e-1/2*(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}/e$

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3782, 1261, 748, 857, 635, 212, 738}

$$\int \cot(d+ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \frac{\sqrt{a-b+c} \operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} - \frac{(b-2c)\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ce}} - \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e}$$

[In] Int[Cot[d + e*x]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] (Sqrt[a - b + c]*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*e) - ((b - 2*c)*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])])/(4*Sqrt[c]*e) - Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]/(2*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 748

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[p/(e*(m + 2*p + 1)), Int[(d + e*x)^m*Simp[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 857

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1261

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(
p_), x_Symbol] := Dist[1/2, Subst[Int[(d + e*x)^q*(a + b*x + c*x^2)^p, x],
x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x]
```

Rule 3782

```
Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(
f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x\sqrt{a+bx^2+cx^4}}{1+x^2} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{1+x} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e} + \frac{\text{Subst}\left(\int \frac{-2a+b-(b-2c)x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4e} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e} \\
&\quad - \frac{(b-2c)\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4e} \\
&\quad - \frac{(a-b+c)\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e} \\
&\quad - \frac{(b-2c)\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&\quad + \frac{(a-b+c)\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{e} \\
&= \frac{\sqrt{a-b} + \text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&\quad - \frac{(b-2c)\text{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ce}} \\
&\quad - \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}{2e}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.46 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.37

$$\int \cot(d+ex) \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} dx = \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)} \tan^2(d+ex) \left((b-2c) \operatorname{arctanh} \left(\frac{2c+b\tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}} \right) + 2 \right)}{4\sqrt{ce}\sqrt{c+bt}}$$

[In] Integrate[Cot[d + e*x]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4], x]

[Out] $-1/4*(\text{Sqrt}[a + b*\text{Cot}[d + e*x]^2 + c*\text{Cot}[d + e*x]^4]*\text{Tan}[d + e*x]^2*((b - 2*c)*\text{ArcTanh}[(2*c + b*\text{Tan}[d + e*x]^2)/(2*\text{Sqrt}[c]*\text{Sqrt}[c + b*\text{Tan}[d + e*x]^2 + a*\text{Tan}[d + e*x]^4)]) + 2*\text{Sqrt}[c]*(-(\text{Sqrt}[a - b + c]*\text{ArcTanh}[(b - 2*c + (2*a - b)*\text{Tan}[d + e*x]^2)/(2*\text{Sqrt}[a - b + c]*\text{Sqrt}[c + b*\text{Tan}[d + e*x]^2 + a*\text{Tan}[d + e*x]^4)]) + \text{Cot}[d + e*x]^2*\text{Sqrt}[c + b*\text{Tan}[d + e*x]^2 + a*\text{Tan}[d + e*x]^4])))/(\text{Sqrt}[c]*e*\text{Sqrt}[c + b*\text{Tan}[d + e*x]^2 + a*\text{Tan}[d + e*x]^4])$

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.21

method	result
derivativedivides	$\frac{-\sqrt{\frac{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}{2}}}{4\sqrt{c}} - \frac{(b-2c) \ln\left(\frac{\frac{b}{2}-c+(\cot(ex+d)^2+1)c}{\sqrt{c}} + \sqrt{\frac{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}{2}}\right)}{4\sqrt{c}}$
default	$\frac{-\sqrt{\frac{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}{2}}}{4\sqrt{c}} - \frac{(b-2c) \ln\left(\frac{\frac{b}{2}-c+(\cot(ex+d)^2+1)c}{\sqrt{c}} + \sqrt{\frac{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}{2}}\right)}{4\sqrt{c}}$

[In] int(cot(e*x+d)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2), x, method=_RETURNVERB OSE)

[Out] $1/e*(-1/2*((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^(1/2)-1/4*(b-2*c)*\ln((1/2*b-c+(\cot(e*x+d)^2+1)*c)/c^(1/2)+((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^(1/2))/c^(1/2)+1/2*(a-b+c)^(1/2)*\ln((2*a-2*b+2*c+(b-2*c)*(\cot(e*x+d)^2+1)+2*(a-b+c)^(1/2))*((\cot(e*x+d)^2+1)^2*c+(b-2*c)*(\cot(e*x+d)^2+1)+a-b+c)^(1/2))/(\cot(e*x+d)^2+1))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 465 vs. 2(155) = 310.

Time = 1.75 (sec) , antiderivative size = 1932, normalized size of antiderivative = 10.79

$$\int \cot(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="fricas")

[Out] [1/8*(2*sqrt(a - b + c)*c*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) - (b - 2*c)*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 + 4*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*c*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c*e), -1/4*((b - 2*c)*sqrt(-c)*arctan(-1/2*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(-c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + (a + b)*c + c^2 - 2*(a*c - c^2)*cos(2*e*x + 2*d)) - sqrt(a - b + c)*c*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) + 2*c*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/(c*e), -1/8*(4*sqrt(-a + b - c)*c*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) + (b - 2*c)*sqrt(c)*log(((b^2 + 4*(a - 2*b)*c + 8*c^2)*cos(2*e*x + 2*d)^2 + b^2 + 4*(a + 2*b)*c + 8*c^2 + 4*((b - 2*c)*cos(2*e*x + 2*d)^2 - 2*b*cos(2*e*x + 2*d) + b + 2*c)*sqrt(c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 2*(b^2 + 4*a*c - 8*c^2)*cos(2*e*x + 2*d))/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) + 4*c*sqrt(((a - b + c)*c

$$\frac{\cos(2ex + 2d)^2 - 2(a - c)\cos(2ex + 2d) + a + b + c}{(\cos(2ex + 2d)^2 - 2\cos(2ex + 2d) + 1)} \cdot \frac{-1/4(2\sqrt{-a + b - c})c \arctan\left(\frac{(a - b + c)\cos(2ex + 2d)^2 - (2a - b)\cos(2ex + 2d) + a - c}{\sqrt{-a + b - c}}\right) + \frac{a + b + c}{(\cos(2ex + 2d)^2 - 2\cos(2ex + 2d) + 1)} \cdot \frac{1}{\sqrt{(a^2 - 2ab + b^2 + 2(a - b)c + c^2)\cos(2ex + 2d)^2 + a^2 - b^2 + 2ac + c^2 - 2(a^2 - ab + bc - c^2)\cos(2ex + 2d)}} + (b - 2c)\sqrt{-c} \arctan\left(\frac{-1/2((b - 2c)\cos(2ex + 2d)^2 - 2b\cos(2ex + 2d) + b + 2c)}{\sqrt{-c}}\right) + \frac{2c\sqrt{((a - b + c)\cos(2ex + 2d)^2 - 2(a - c)\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d)^2 - 2\cos(2ex + 2d) + 1)}}{\sqrt{((a - b)c + c^2)\cos(2ex + 2d)^2 + (a + b)c + c^2 - 2(ac - c^2)\cos(2ex + 2d)}} + \frac{2c\sqrt{((a - b + c)\cos(2ex + 2d)^2 - 2(a - c)\cos(2ex + 2d) + a + b + c)/(\cos(2ex + 2d)^2 - 2\cos(2ex + 2d) + 1)}}{c}$$

Sympy [F]

$$\int \cot(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \cot(d + ex) dx$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2),x)

[Out] Integral(sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)*cot(d + e*x), x)

Maxima [F]

$$\int \cot(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx$$

$$= \int \sqrt{c \cot^4(ex + d) + b \cot^2(ex + d) + a} \cot(ex + d) dx$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)*cot(e*x + d), x)

Giac [F(-1)]

Timed out.

$$\int \cot(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)*(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\begin{aligned} & \int \cot(d + ex) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} dx \\ &= \int \cot(d + ex) \sqrt{c \cot^4(d + ex) + b \cot^2(d + ex) + a} dx \end{aligned}$$

[In] int(cot(d + e*x)*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)

[Out] int(cot(d + e*x)*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)

3.25 $\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan(d + ex) dx$

Optimal result	241
Rubi [A] (verified)	242
Mathematica [A] (verified)	244
Maple [F]	245
Fricas [A] (verification not implemented)	245
Sympy [F]	247
Maxima [F]	247
Giac [F(-1)]	247
Mupad [F(-1)]	248

Optimal result

Integrand size = 33, antiderivative size = 203

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan(d + ex) dx$$

$$= \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot^2(d + ex)}{2\sqrt{a}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e}$$

$$- \frac{\sqrt{a - b} \operatorname{arctanh}\left(\frac{2a - b + (b - 2c) \cot^2(d + ex)}{2\sqrt{a - b} \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e}$$

$$- \frac{\sqrt{c} \operatorname{arctanh}\left(\frac{b + 2c \cot^2(d + ex)}{2\sqrt{c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e}$$

```
[Out] 1/2*arctanh(1/2*(2*a+b*cot(e*x+d)^2)/a^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*a^(1/2)/e-1/2*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*c^(1/2)/e-1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*(a-b+c)^(1/2)/e
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3782, 1265, 909, 738, 212, 857, 635}

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan(d + ex) dx$$

$$= \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot^2(d + ex)}{2\sqrt{a}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e} - \frac{\sqrt{a - b} + c \operatorname{arctanh}\left(\frac{2a + (b - 2c) \cot^2(d + ex) - b}{2\sqrt{a - b} + c\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e} - \frac{\sqrt{c} \operatorname{arctanh}\left(\frac{b + 2c \cot^2(d + ex)}{2\sqrt{c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e}$$

[In] Int[Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x],x]

[Out] (Sqrt[a]*ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*e) - (Sqrt[a - b + c]*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])])/(2*e) - (Sqrt[c]*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])])/(2*e)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 857

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x +

```
c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p,
x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] &&
NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 909

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_)/(((d_.) + (e_.)*(x_))*((f_.) +
(g_.)*(x_))), x_Symbol] := Dist[(c*d^2 - b*d*e + a*e^2)/(e*(e*f - d*g)), I
nt[(a + b*x + c*x^2)^(p - 1)/(d + e*x), x], x] - Dist[1/(e*(e*f - d*g)), In
t[Simp[c*d*f - b*e*f + a*e*g - c*(e*f - d*g)*x, x]*((a + b*x + c*x^2)^(p -
1)/(f + g*x)), x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g,
0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && FractionQ[p]
&& GtQ[p, 0]
```

Rule 1265

```
Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)
^4)^(p_.), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a +
b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && Inte
gerQ[(m - 1)/2]
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:= Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx^2+cx^4}}{x(1+x^2)} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x(1+x)} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= \frac{\text{Subst}\left(\int \frac{a-b-cx}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} - \frac{a\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{a \operatorname{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&\quad - \frac{c \operatorname{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&\quad + \frac{(a-b+c) \operatorname{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&\quad - \frac{c \operatorname{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&\quad - \frac{(a-b+c) \operatorname{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&= \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&\quad - \frac{\sqrt{a-b+c} \operatorname{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&\quad - \frac{\sqrt{c} \operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.77 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.25

$$\begin{aligned}
&\int \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} \tan(d+ex) dx \\
&= \frac{\left(\sqrt{a} \operatorname{arctanh}\left(\frac{b+2a \tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b \tan^2(d+ex)+a \tan^4(d+ex)}}\right) - \sqrt{a-b+c} \operatorname{arctanh}\left(\frac{b-2c+(2a-b) \tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b \tan^2(d+ex)+a \tan^4(d+ex)}}\right) - \sqrt{c} \operatorname{arctanh}\left(\frac{b+2c \tan^2(d+ex)}{2\sqrt{c}\sqrt{c+b \tan^2(d+ex)+a \tan^4(d+ex)}}\right)\right)}{2e\sqrt{c+b \tan^2(d+ex)+a \tan^4(d+ex)}}
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x],x]

[Out] ((Sqrt[a]*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]) - Sqrt[a - b + c]*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)]) - Sqrt[c]*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)])]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/(2*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [F]

$$\int \sqrt{a + b \cot(ex + d)^2 + c \cot(ex + d)^4} \tan(ex + d) dx$$

[In] int((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d),x)

[Out] int((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d),x)

Fricas [A] (verification not implemented)

none

Time = 0.70 (sec) , antiderivative size = 2522, normalized size of antiderivative = 12.42

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan(d + ex) dx = \text{Too large to display}$$

[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d),x, algorithm="fricas")

[Out] [1/4*(sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)) + sqrt(a - b + c)*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)) + sqrt(c)*log(((b^2 + 4*a*c)*tan(e*x + d)^4 + 8*b*c*tan(e*x + d)^2 + 8*c^2 - 4*(b*tan(e*x + d)^4 + 2*c*tan(e*x + d)^2)*sqrt(c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/tan(e*x + d)^4)/e, 1/4*(2*sqrt(-c)*arctan(2*sqrt(-c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2/(b*tan(e*x + d)^2 + 2*c)) + sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)) + sqrt(a - b + c)*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)))/e, -1/4*(2*sqrt(-a)*arctan(2*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2/(2*a*tan(e*x + d)^2 + b)) - sqrt(a - b + c)*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)) - sqrt(c)*log(((b^2 + 4*a*c)*tan(e*x + d)^4 + 8*b*c*tan(e*x + d)^2 + 8*c^2 - 4*(b*tan(e*x + d)^4 + 2

$$\begin{aligned}
& *c*\tan(e*x + d)^2*\sqrt{c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4))/\tan(e*x + d)^4)/e, -1/4*(2*\sqrt{-a}*\arctan(2*\sqrt{-a}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(2*a*\tan(e*x + d)^2 + b))} - 2*\sqrt{-c}*\arctan(2*\sqrt{-c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(b*\tan(e*x + d)^2 + 2*c))} - \sqrt{a - b + c}*\log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*\tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*\tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*\tan(e*x + d)^4 + (b - 2*c)*\tan(e*x + d)^2)*\sqrt{a - b + c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)))/(\tan(e*x + d)^4 + 2*\tan(e*x + d)^2 + 1)))/e, -1/4*(2*\sqrt{-a + b - c}*\arctan(-2*\sqrt{-a + b - c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/((2*a - b)*\tan(e*x + d)^2 + b - 2*c))} - \sqrt{a}*\log(8*a^2*\tan(e*x + d)^4 + 8*a*b*\tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2)*\sqrt{a}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4))} - \sqrt{c}*\log(((b^2 + 4*a*c)*\tan(e*x + d)^4 + 8*b*c*\tan(e*x + d)^2 + 8*c^2 - 4*(b*\tan(e*x + d)^4 + 2*c*\tan(e*x + d)^2)*\sqrt{c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)))/\tan(e*x + d)^4))/e, -1/4*(2*\sqrt{-a + b - c}*\arctan(-2*\sqrt{-a + b - c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/((2*a - b)*\tan(e*x + d)^2 + b - 2*c))} - 2*\sqrt{-c}*\arctan(2*\sqrt{-c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(b*\tan(e*x + d)^2 + 2*c))} - \sqrt{a}*\log(8*a^2*\tan(e*x + d)^4 + 8*a*b*\tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2)*\sqrt{a}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)))/e, -1/4*(2*\sqrt{-a}*\arctan(2*\sqrt{-a}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(2*a*\tan(e*x + d)^2 + b))} + 2*\sqrt{-a + b - c}*\arctan(-2*\sqrt{-a + b - c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/((2*a - b)*\tan(e*x + d)^2 + b - 2*c))} - \sqrt{c}*\log(((b^2 + 4*a*c)*\tan(e*x + d)^4 + 8*b*c*\tan(e*x + d)^2 + 8*c^2 - 4*(b*\tan(e*x + d)^4 + 2*c*\tan(e*x + d)^2)*\sqrt{c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)))/\tan(e*x + d)^4))/e, -1/2*(\sqrt{-a}*\arctan(2*\sqrt{-a}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(2*a*\tan(e*x + d)^2 + b))} + \sqrt{-a + b - c}*\arctan(-2*\sqrt{-a + b - c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/((2*a - b)*\tan(e*x + d)^2 + b - 2*c))} - \sqrt{-c}*\arctan(2*\sqrt{-c}*\sqrt{((a*\tan(e*x + d)^4 + b*\tan(e*x + d)^2 + c)/\tan(e*x + d)^4)*\tan(e*x + d)^2/(b*\tan(e*x + d)^2 + 2*c)))/e]
\end{aligned}$$

Sympy [F]

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex) \tan(d + ex)} dx$$

$$= \int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex) \tan(d + ex)} dx$$

[In] integrate((a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2)*tan(e*x+d),x)

[Out] Integral(sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)*tan(d + e*x), x)

Maxima [F]

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex) \tan(d + ex)} dx$$

$$= \int \sqrt{c \cot^4(ex + d) + b \cot^2(ex + d) + a} \tan(ex + d) dx$$

[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d),x, algorithm="maxima")

[Out] integrate(sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)*tan(e*x + d), x)

Giac [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex) \tan(d + ex)} dx = \text{Timed out}$$

[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan(d + ex) dx$$

$$= \int \tan(d + ex) \sqrt{c \cot^4(d + ex) + b \cot^2(d + ex) + a} dx$$

```
[In] int(tan(d + e*x)*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)
```

```
[Out] int(tan(d + e*x)*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)
```


3.26 $\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$

Optimal result	249
Rubi [A] (verified)	250
Mathematica [A] (verified)	254
Maple [F]	255
Fricas [A] (verification not implemented)	255
Sympy [F]	256
Maxima [F]	256
Giac [F(-1)]	256
Mupad [F(-1)]	257

Optimal result

Integrand size = 35, antiderivative size = 435

$$\begin{aligned}
 & \int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx \\
 &= -\frac{\sqrt{a} \operatorname{arctanh}\left(\frac{2a + b \cot^2(d + ex)}{2\sqrt{a}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e} + \frac{\operatorname{barctanh}\left(\frac{2a + b \cot^2(d + ex)}{2\sqrt{a}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{4\sqrt{ae}} \\
 &+ \frac{\sqrt{a - b} \operatorname{carctanh}\left(\frac{2a - b + (b - 2c) \cot^2(d + ex)}{2\sqrt{a - b + c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e} \\
 &+ \frac{\operatorname{barctanh}\left(\frac{b + 2c \cot^2(d + ex)}{2\sqrt{c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{4\sqrt{ce}} \\
 &- \frac{(b - 2c) \operatorname{arctanh}\left(\frac{b + 2c \cot^2(d + ex)}{2\sqrt{c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{4\sqrt{ce}} \\
 &- \frac{\sqrt{c} \operatorname{carctanh}\left(\frac{b + 2c \cot^2(d + ex)}{2\sqrt{c}\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}\right)}{2e} \\
 &+ \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2e}
 \end{aligned}$$

[Out] 1/4*b*arctanh(1/2*(2*a+b*cot(e*x+d)^2)/a^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/a^(1/2)-1/2*arctanh(1/2*(2*a+b*cot(e*x+d)^2)/a^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*a^(1/2)/e+1/4*b*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/c^(1/2)-1/4*(b-2*c)*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/e/c^(1/2)-1/2*arctanh(1/2*(b+2*c*cot(e*x+d)^2)/c^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))*c^(1/2)/e+1/2*arctanh(1/2*(2*a-b+(b-2*c)*co

$$t(e*x+d)^2/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}*(a-b+c)^{(1/2)}/e+1/2*(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}*\tan(e*x+d)^2/e$$

Rubi [A] (verified)

Time = 0.65 (sec) , antiderivative size = 435, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.257$, Rules used = {3782, 1265, 974, 746, 857, 635, 212, 738, 748}

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$$

$$= \frac{\operatorname{barctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4\sqrt{ae}} - \frac{\sqrt{a}\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e}$$

$$+ \frac{\sqrt{a-b} + \operatorname{carctanh}\left(\frac{2a+(b-2c) \cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e}$$

$$- \frac{\sqrt{c}\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} + \frac{\operatorname{barctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4\sqrt{ce}}$$

$$- \frac{(b-2c)\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4\sqrt{ce}}$$

$$+ \frac{\tan^2(d+ex)\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}{2e}$$

[In] Int[Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^3,x]

[Out] -1/2*(Sqrt[a]*ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/e + (b*ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/(4*Sqrt[a]*e) + (Sqrt[a - b + c]*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/(2*e) + (b*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/(4*Sqrt[c]*e) - ((b - 2*c)*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/(4*Sqrt[c]*e) - (Sqrt[c]*ArcTanh[(b + 2*c*Cot[d + e*x]^2)/(2*Sqrt[c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4))]/(2*e) + (Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/(2*e)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

```
Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 738

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 746

```
Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 1))), x] - Dist[p/(e*(m + 1)), Int[(d + e*x)^(m + 1)*(b + 2*c*x)*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && GtQ[p, 0] && (IntegerQ[p] || LtQ[m, -1]) && NeQ[m, -1] && !ILtQ[m + 2*p + 1, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]
```

Rule 748

```
Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + b*x + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[p/(e*(m + 2*p + 1)), Int[(d + e*x)^m*Simp[b*d - 2*a*e + (2*c*d - b*e)*x, x]*(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && GtQ[p, 0] && NeQ[m + 2*p + 1, 0] && (!RationalQ[m] || LtQ[m, 1]) && !ILtQ[m + 2*p, 0] && IntQuadraticQ[a, b, c, d, e, m, p, x]
```

Rule 857

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 974

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (
```

IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])

Rule 1265

Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol] :> Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 3782

Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*cot[(d_.) + (e_.)*(x_)])*(f_.)^(n_.) + (c_.)*cot[(d_.) + (e_.)*(x_)])*(f_.)^(n2_.))^(p_), x_Symbol] :> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x^3(1+x^2)} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x^2(1+x)} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{\sqrt{a+bx+cx^2}}{x^2} - \frac{\sqrt{a+bx+cx^2}}{x} + \frac{\sqrt{a+bx+cx^2}}{1+x}\right) dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x^2} dx, x, \cot^2(d+ex)\right)}{2e} + \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{x} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx+cx^2}}{1+x} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}\tan^2(d+ex)}{2e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{-2a-bx}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4e} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{-2a+b-(b-2c)x}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{b+2cx}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{4e}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2e} \\
&+ \frac{a \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{2e} \\
&+ \frac{b \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{4e} \\
&- \frac{b \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{4e} \\
&- \frac{(b - 2c) \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{4e} \\
&- \frac{c \text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{2e} \\
&- \frac{(a - b + c) \text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{2e} \\
&= \frac{\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2e} \\
&- \frac{a \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&+ \frac{b \text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&+ \frac{b \text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&- \frac{(b - 2c) \text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e} \\
&- \frac{c \text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e} \\
&+ \frac{(a - b + c) \text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{a}\operatorname{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&+ \frac{\operatorname{barctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ae}} \\
&+ \frac{\sqrt{a-b} + \operatorname{carctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&+ \frac{\operatorname{barctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ce}} \\
&- \frac{(b-2c)\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{4\sqrt{ce}} \\
&- \frac{\sqrt{c}\operatorname{arctanh}\left(\frac{b+2c\cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e} \\
&+ \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}\tan^2(d+ex)}{2e}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.29 (sec) , antiderivative size = 235, normalized size of antiderivative = 0.54

$$\begin{aligned}
&\int \sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}\tan^3(d+ex)dx \\
&= \frac{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}\tan^2(d+ex)\left((-2a+b)\operatorname{arctanh}\left(\frac{b+2a\tan^2(d+ex)}{2\sqrt{a}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)+2\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}\right)}{4\sqrt{ae}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}
\end{aligned}$$

[In] Integrate[Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^3,x]

[Out] (Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2*((-2*a + b)*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])] + 2*Sqrt[a]*(Sqrt[a - b + c]*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])]) + Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))/(4*Sqrt[a]*e*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])

Maple [F]

$$\int \sqrt{a + b \cot(ex + d)^2 + c \cot(ex + d)^4} \tan(ex + d)^3 dx$$

[In] int((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d)^3,x)

[Out] int((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d)^3,x)

Fricas [A] (verification not implemented)

none

Time = 1.76 (sec) , antiderivative size = 1282, normalized size of antiderivative = 2.95

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx = \text{Too large to display}$$

[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d)^3,x, algorithm="fricas")

[Out] [1/8*(4*a*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 - (2*a - b)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)) + 2*sqrt(a - b + c)*a*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 + 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)))/(a*e), 1/8*(4*a*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 + 4*a*sqrt(-a + b - c)*arctan(-1/2*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(-a + b - c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/((a^2 - a*b + a*c)*tan(e*x + d)^4 + (a*b - b^2 + b*c)*tan(e*x + d)^2 + (a - b)*c + c^2)) - (2*a - b)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)))/(a*e), 1/4*(2*a*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 + sqrt(-a)*(2*a - b)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)/(a^2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)) + sqrt(a - b + c)*a*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 + 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)))/(a*e), 1/4*(2*a*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)*tan(e*x + d)^2 + sqrt(-a)*(2*a - b)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*

```
sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)/(a^
2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)) + 2*a*sqrt(-a + b - c)*arctan
(-1/2*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(-a + b - c
)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)/((a^2 - a*
b + a*c)*tan(e*x + d)^4 + (a*b - b^2 + b*c)*tan(e*x + d)^2 + (a - b)*c + c^
2)))/(a*e)]
```

Sympy [F]

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$$

$$= \int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$$

```
[In] integrate((a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(1/2)*tan(e*x+d)**3,x)
```

```
[Out] Integral(sqrt(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)*tan(d + e*x)**3, x
)
```

Maxima [F]

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$$

$$= \int \sqrt{c \cot^4(ex + d) + b \cot^2(ex + d)^2 + a} \tan^3(ex + d)^3 dx$$

```
[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d)^3,x, algorithm
="maxima")
```

```
[Out] integrate(sqrt(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)*tan(e*x + d)^3, x)
```

Giac [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx = \text{Timed out}$$

```
[In] integrate((a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*tan(e*x+d)^3,x, algorithm
="giac")
```

```
[Out] Timed out
```


Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^3(d + ex) dx$$

$$= \int \tan(d + ex)^3 \sqrt{c \cot(d + ex)^4 + b \cot(d + ex)^2 + a} dx$$

```
[In] int(tan(d + e*x)^3*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2),x)
```

```
[Out] int(tan(d + e*x)^3*(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(1/2), x)
```

$$3.27 \quad \int \frac{\cot^7(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	258
Rubi [A] (verified)	258
Mathematica [A] (verified)	261
Maple [B] (verified)	262
Fricas [B] (verification not implemented)	262
Sympy [F]	263
Maxima [F(-2)]	263
Giac [F(-1)]	263
Mupad [F(-1)]	263

Optimal result

Integrand size = 35, antiderivative size = 236

$$\int \frac{\cot^7(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx =$$

$$\frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} - \frac{\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2c^{3/2}e}$$

$$- \frac{a(b^2 - a(b+2c)) + (b^3 + 2a^2c - ab(b+3c)) \cot^2(d+ex)}{c(a-b+c)(b^2 - 4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(b+2*c*\cot(e*x+d)^2)/c^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/c^{(3/2)}/e-1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/(a-b+c)^{(3/2)}/e+(-a*(b^2-a*(b+2*c))-(b^3+2*a^2*c-a*b*(b+3*c))*\cot(e*x+d)^2)/c/(a-b+c)/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.64 (sec) , antiderivative size = 236, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3782, 1265, 1660, 857, 635, 212, 738}

$$\int \frac{\cot^7(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx =$$

$$\frac{(2a^2c - ab(b+3c) + b^3) \cot^2(d+ex) + a(b^2 - a(b+2c))}{ce(a-b+c)(b^2 - 4ac)\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$- \frac{\operatorname{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2c^{3/2}e} - \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}}$$

[In] Int[Cot[d + e*x]^7/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2),x]

[Out]
$$-1/2 \operatorname{ArcTanh}\left[\frac{2a - b + (b - 2c)\operatorname{Cot}[d + e*x]^2}{2\sqrt{a - b + c}\sqrt{a + b\operatorname{Cot}[d + e*x]^2 + c\operatorname{Cot}[d + e*x]^4}}\right] / ((a - b + c)^{3/2}e) - \operatorname{ArcTanh}\left[\frac{b + 2c\operatorname{Cot}[d + e*x]^2}{2\sqrt{c}\sqrt{a + b\operatorname{Cot}[d + e*x]^2 + c\operatorname{Cot}[d + e*x]^4}}\right] / (2c^{3/2}e) - (a(b^2 - a(b + 2c)) + (b^3 + 2a^2c - ab(b + 3c))\operatorname{Cot}[d + e*x]^2) / (c(a - b + c)(b^2 - 4ac)e\sqrt{a + b\operatorname{Cot}[d + e*x]^2 + c\operatorname{Cot}[d + e*x]^4})$$

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 857

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 1660

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + b*x + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x + c*x^2, x],

```
x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x + c*x^2, x], x
, 1]}, Simp[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^(p + 1)/((p
+ 1)*(b^2 - 4*a*c))), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)), Int[(d + e*x)^m*
(a + b*x + c*x^2)^(p + 1)*ExpandToSum[((p + 1)*(b^2 - 4*a*c)*Q)/(d + e*x)^m
- ((2*p + 3)*(2*c*f - b*g))/(d + e*x)^m, x], x] /; FreeQ[{a, b, c, d,
e}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2
, 0] && LtQ[p, -1] && ILtQ[m, 0]
```

Rule 3782

```
Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(
f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^7}{(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x^3}{(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{a(b^2 - a(b+2c)) + (b^3 + 2a^2c - ab(b+3c)) \cot^2(d+ex)}{c(a-b+c)(b^2 - 4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{(a-b)(b^2-4ac)}{2c(a-b+c)} - \frac{(b^2-4ac)x}{2c}}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{(b^2 - 4ac)e} \\
&= -\frac{a(b^2 - a(b+2c)) + (b^3 + 2a^2c - ab(b+3c)) \cot^2(d+ex)}{c(a-b+c)(b^2 - 4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2ce} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2(a-b+c)e}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{a(b^2 - a(b + 2c)) + (b^3 + 2a^2c - ab(b + 3c)) \cot^2(d + ex)}{c(a - b + c)(b^2 - 4ac)e\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{4c-x^2} dx, x, \frac{b+2c \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{ce} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{(a-b+c)e} \\
&= -\frac{\text{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} \\
&\quad - \frac{\text{arctanh}\left(\frac{b+2c \cot^2(d+ex)}{2\sqrt{c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2c^{3/2}e} \\
&\quad - \frac{a(b^2 - a(b + 2c)) + (b^3 + 2a^2c - ab(b + 3c)) \cot^2(d + ex)}{c(a - b + c)(b^2 - 4ac)e\sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 5.14 (sec) , antiderivative size = 339, normalized size of antiderivative = 1.44

$$\int \frac{\cot^7(d + ex)}{(a + b \cot^2(d + ex) + c \cot^4(d + ex))^{3/2}} dx = \frac{\cot^2(d + ex) \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}}{(b^2 - 4ac)^{3/2}} \left(\frac{(b^2 - 4ac) \text{arctanh}\left(\frac{b - 2c + (2a - b) \tan^2(d + ex)}{2\sqrt{a - b + c} \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}}\right)}{(a - b + c)^{3/2}} + \frac{(-1/2 b^2 + 2ac) \text{arctanh}\left(\frac{2c + b \tan^2(d + ex)}{2\sqrt{c} \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}}\right)}{c^{3/2}} + \frac{(b^2 - 2ac + ab \tan^2(d + ex)) \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}}{(a - b + c) \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}} - \frac{(b^2 - a(b + 2c) + a(-2a + b) \tan^2(d + ex)) \sqrt{c + b \tan^2(d + ex) + a \tan^4(d + ex)}}{(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \right)$$

[In] Integrate[Cot[d + e*x]^7/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2),x]

[Out] (Cot[d + e*x]^2*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]*(-1/2*((b^2 - 4*a*c)*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]])))/(a - b + c)^(3/2) + ((-1/2*b^2 + 2*a*c)*ArcTanh[(2*c + b*Tan[d + e*x]^2)/(2*Sqrt[c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]]))/c^(3/2) + (b^2 - 2*a*c + a*b*Tan[d + e*x]^2)/(c*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]) - (b^2 - a*(b + 2*c) + a*(-2*a + b)*Tan[d + e*x]^2)/((a - b + c)*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))/(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 685 vs. $2(215) = 430$.

Time = 0.59 (sec) , antiderivative size = 686, normalized size of antiderivative = 2.91

method	result
derivativedivides	$-\frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{\cot(ex+d)^2}{2c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}} + b \left(-\frac{1}{c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}} - \frac{c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{4c} \right)$
default	$-\frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{\cot(ex+d)^2}{2c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}} + b \left(-\frac{1}{c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}} - \frac{c\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4}}{4c} \right)$

[In] `int(cot(e*x+d)^7/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x,method=_RETURNVE
RBOSE)`

[Out] $1/e*(-1/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2)*(b+2*c*\cot(e*x+d)^2)/(4*a*c-b^2)+1/2*\cot(e*x+d)^2/c/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2)+1/4*b/c*(-1/c/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2)-b/c/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2)*(b+2*c*\cot(e*x+d)^2)/(4*a*c-b^2))-1/2/c^(3/2)*\ln((1/2*b+c*\cot(e*x+d)^2)/c^(1/2)+(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2))-1/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^(1/2)*(2*a+b*\cot(e*x+d)^2)/(4*a*c-b^2)+2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/((-4*a*c+b^2)^(1/2)+b-2*c)/(a-b+c)^(1/2)*\ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2))*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1)+2*c/((-4*a*c+b^2)^(1/2)+b-2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)^2*c-(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c))^(1/2)-2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)^2*c+(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c))^(1/2))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1484 vs. $2(215) = 430$.

Time = 2.60 (sec) , antiderivative size = 6011, normalized size of antiderivative = 25.47

$$\int \frac{\cot^7(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] `integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{\cot^7(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot^7(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx$$

[In] integrate(cot(e*x+d)**7/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(cot(d + e*x)**7/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\cot^7(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

[In] integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Giac [F(-1)]

Timed out.

$$\int \frac{\cot^7(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^7/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^7(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)^7}{(c\cot(d+ex)^4+b\cot(d+ex)^2+a)^{3/2}} dx$$

[In] int(cot(d + e*x)^7/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2),x)

[Out] int(cot(d + e*x)^7/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)

$$3.28 \quad \int \frac{\cot^5(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	264
Rubi [A] (verified)	264
Mathematica [A] (verified)	266
Maple [B] (verified)	267
Fricas [B] (verification not implemented)	268
Sympy [F]	269
Maxima [F]	269
Giac [F(-1)]	269
Mupad [F(-1)]	270

Optimal result

Integrand size = 35, antiderivative size = 160

$$\int \frac{\cot^5(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} - \frac{a(2a-b) + ((a-b)b+2ac)\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[Out] 1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/(a-b+c)^(3/2)/e+(-a*(2*a-b)-((a-b)*b+2*a*c)*cot(e*x+d)^2)/(a-b+c)/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3782, 1265, 1660, 12, 738, 212}

$$\int \frac{\cot^5(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}} - \frac{(b(a-b)+2ac)\cot^2(d+ex)+a(2a-b)}{e(a-b+c)(b^2-4ac)\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[In] Int[Cot[d + e*x]^5/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

[Out] ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*sqrt[a - b + c]*sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*(a - b + c)^(3/2)*e) - (a*(2*a - b)

+ ((a - b)*b + 2*a*c)*Cot[d + e*x]^2)/((a - b + c)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 1660

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[(d + e*x)^m*Pq, a + b*x + c*x^2, x], f = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x + c*x^2, x], x, 0], g = Coeff[PolynomialRemainder[(d + e*x)^m*Pq, a + b*x + c*x^2, x], x, 1]}, Simp[(b*f - 2*a*g + (2*c*f - b*g)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c))), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p + 1)*ExpandToSum[((p + 1)*(b^2 - 4*a*c)*Q)/(d + e*x)^m - ((2*p + 3)*(2*c*f - b*g))/(d + e*x)^m, x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[p, -1] && ILtQ[m, 0]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)])*(f_))^(n2_)]^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n

2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^5}{(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{x^2}{(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{a(2a-b) + ((a-b)b + 2ac) \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
 &\quad + \frac{\text{Subst}\left(\int -\frac{b^2-4ac}{2(a-b+c)(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{(b^2-4ac)e} \\
 &= -\frac{a(2a-b) + ((a-b)b + 2ac) \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2(a-b+c)e} \\
 &= -\frac{a(2a-b) + ((a-b)b + 2ac) \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{(a-b+c)e} \\
 &= \frac{\text{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} \\
 &\quad - \frac{a(2a-b) + ((a-b)b + 2ac) \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.28

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \frac{2\sqrt{a-b+c}(a(2a-b) + (ab-b^2+2ac)\cot^2(d+ex)) - (b^2-4ac)\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{2(a-b+c)^{3/2}(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}$$

[In] Integrate[Cot[d + e*x]^5/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

```
[Out] -1/2*(2*Sqrt[a - b + c]*(a*(2*a - b) + (a*b - b^2 + 2*a*c)*Cot[d + e*x]^2)
- (b^2 - 4*a*c)*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b
+ c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))*Cot[d + e*x]^2*Sqrt[c
+ b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])/((a - b + c)^(3/2)*(b^2 - 4*a*c)*e*
Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 506 vs. $2(149) = 298$.

Time = 0.09 (sec) , antiderivative size = 507, normalized size of antiderivative = 3.17

method	result
derivativedivides	$\frac{\frac{2a+b \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} - \frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)}{\sqrt{-4ac}} \right)}{\sqrt{-4ac}}$
default	$\frac{\frac{2a+b \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} - \frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)}{\sqrt{-4ac}} \right)}{\sqrt{-4ac}}$

```
[In] int(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x,method=_RETURNVE
RBOSE)
```

```
[Out] 1/e*(1/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*(2*a+b*cot(e*x+d)^2)/(4*a*c-
b^2)+1/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*(b+2*c*cot(e*x+d)^2)/(4*a*c-
b^2)-2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/((-4*a*c+b^2)^(1/2)+b-2*c)/(a-b+c)^(1/2
)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2)*((cot(e*x+d)^2+1
)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1)-2*c/((-4*a*c
+b^2)^(1/2)+b-2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)
*((cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)^2*c-(-4*a*c+b^2)^(1/2)*(cot(e
*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c))^(1/2)+2*c/((-4*a*c+b^2)^(1/2)-b+2*c)
/(-4*a*c+b^2)/(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2-1
/2*(-b+(-4*a*c+b^2)^(1/2))/c)^2*c+(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2-1/2*(-b+
(-4*a*c+b^2)^(1/2))/c))^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 873 vs. 2(148) = 296.

Time = 0.63 (sec) , antiderivative size = 1743, normalized size of antiderivative = 10.89

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")

[Out] [1/4*((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*cos(2*e*x + 2*d))*sqrt(a - b + c)*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d) - 4*(2*a^3 - 2*a^2*b - a*b^2 + b^3 + 2*a*c^2 + (2*a^3 - 4*a^2*b + 3*a*b^2 - b^3 + b^2*c - 2*a*c^2)*cos(2*e*x + 2*d)^2 + (4*a^2 - 2*a*b - b^2)*c - 2*(2*a^3 - 3*a^2*b + a*b^2 + (2*a^2 - a*b)*c)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e), -1/2*((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*cos(2*e*x + 2*d))*sqrt(-a + b - c)*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d)) + 2*(2*a^3 - 2*a^2*b - a*b^2 + b^3 + 2*a*c^2 + (2*a^3 - 4*a^2*b + 3*a*b^2 - b^3 + b^2*c - 2*a*c^2)*cos(2*e*x + 2*d)^2 + (4*a^2 - 2*a*b - b^2)*c - 2*(2*a^3 - 3*a^2*b + a*b^2 + (2*a^2 - a*b)*c)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e)

$$2 - 8ab - b^2)c^3 - (4a^3 - 3ab^2 - 2b^3)c^2 - (4a^4 - 8a^3b + 3a^2b^2 + b^4)c)e\cos(2ex + 2d) + (a^3b^2 - a^2b^3 - ab^4 + b^5 - 4ac^4 - (12a^2 - 4ab - b^2)c^3 - (12a^3 - 8a^2b - 7ab^2 + b^3)c^2 - (4a^4 - 4a^3b - 7a^2b^2 + 6ab^3 + b^4)c)e]$$

Sympy [F]

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx$$

[In] integrate(cot(e*x+d)**5/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(cot(d + e*x)**5/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima [F]

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(ex+d)^5}{(c\cot(ex+d)^4+b\cot(ex+d)^2+a)^{3/2}} dx$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)^5/(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)^(3/2), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^5/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^5(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)^5}{(c\cot(d+ex)^4+b\cot(d+ex)^2+a)^{3/2}} dx$$

```
[In] int(cot(d + e*x)^5/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

```
[Out] int(cot(d + e*x)^5/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

$$3.29 \quad \int \frac{\cot^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	271
Rubi [A] (verified)	271
Mathematica [A] (verified)	273
Maple [B] (verified)	274
Fricas [B] (verification not implemented)	275
Sympy [F]	276
Maxima [F]	276
Giac [F(-1)]	276
Mupad [F(-1)]	277

Optimal result

Integrand size = 35, antiderivative size = 153

$$\int \frac{\cot^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx =$$

$$\frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e}$$

$$+ \frac{a(b-2c) + (2a-b)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/(a-b+c)^{(3/2)}/e+(a*(b-2*c)+(2*a-b)*c*\cot(e*x+d)^2)/(a-b+c)/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}$

Rubi [A] (verified)

Time = 0.34 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.171$, Rules used = {3782, 1265, 836, 12, 738, 212}

$$\int \frac{\cot^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{c(2a-b)\cot^2(d+ex)+a(b-2c)}{e(a-b+c)(b^2-4ac)\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$- \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}}$$

[In] $\operatorname{Int}[\operatorname{Cot}[d+e*x]^3/(a+b*\operatorname{Cot}[d+e*x]^2+c*\operatorname{Cot}[d+e*x]^4)^{(3/2)},x]$

```
[Out] -1/2*ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a
+ b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])/((a - b + c)^(3/2)*e) + (a*(b - 2
*c) + (2*a - b)*c*Cot[d + e*x]^2)/((a - b + c)*(b^2 - 4*a*c)*e*Sqrt[a + b*C
ot[d + e*x]^2 + c*Cot[d + e*x]^4)]
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 738

```
Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Sym
bol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2
*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c,
d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 836

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c
_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(f*(b*c*d - b^2*e + 2
*a*c*e) - a*g*(2*c*d - b*e) + c*(f*(2*c*d - b*e) - g*(b*d - 2*a*e))*x)*((a
+ b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x]
+ Dist[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^m*
(a + b*x + c*x^2)^(p + 1)*Simp[f*(b*c*d*e*(2*p - m + 2) + b^2*e^2*(p + m +
2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3)) - g*(a*e*(b*e - 2*c*d*m
+ b*e*m) - b*d*(3*c*d - b*e + 2*c*d*p - b*e*p)) + c*e*(g*(b*d - 2*a*e) - f
*(2*c*d - b*e))*(m + 2*p + 4)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g,
m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[p, -1
] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Rule 1265

```
Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)
^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a +
b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && Inte
gerQ[(m - 1)/2]
```

Rule 3782

```
Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(
f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol]
```



```

:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x
], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n
2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x^3}{(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{x}{(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= \frac{a(b-2c) + (2a-b)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{b^2-4ac}{2(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{(a-b+c)(b^2-4ac)e} \\
&= \frac{a(b-2c) + (2a-b)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2(a-b+c)e} \\
&= \frac{a(b-2c) + (2a-b)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{(a-b+c)e} \\
&= -\frac{\text{arctanh}\left(\frac{2a-b+(-b+2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} \\
&\quad + \frac{a(b-2c) + (2a-b)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.71 (sec) , antiderivative size = 195, normalized size of antiderivative = 1.27

$$\begin{aligned}
&\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \\
&\frac{2\sqrt{a-b+c}(-a(b-2c) + (-2a+b)c\cot^2(d+ex)) + (b^2-4ac)\text{arctanh}\left(\frac{b-2c+(2a-b)\tan^2(d+ex)}{2\sqrt{a-b+c}\sqrt{c+b\tan^2(d+ex)+a\tan^4(d+ex)}}\right)}{2(a-b+c)^{3/2}(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
\end{aligned}$$

```
[In] Integrate[Cot[d + e*x]^3/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2),x]
[Out] -1/2*(2*Sqrt[a - b + c]*(-(a*(b - 2*c)) + (-2*a + b)*c*Cot[d + e*x]^2) + (b
^2 - 4*a*c)*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]
*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]])*Cot[d + e*x]^2*Sqrt[c + b*
Tan[d + e*x]^2 + a*Tan[d + e*x]^4])/((a - b + c)^(3/2)*(b^2 - 4*a*c)*e*Sqrt
[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 456 vs. 2(141) = 282.

Time = 0.11 (sec) , antiderivative size = 457, normalized size of antiderivative = 2.99

method	result
derivativedivides	$-\frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c} \sqrt{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)}}{\cot(ex+d)^2+1} \right)}{(\sqrt{-4ac+b^2-b+2c})(\sqrt{-4ac+b^2+b-2c})\sqrt{a-b+c}}$
default	$-\frac{b+2c \cot(ex+d)^2}{\sqrt{a+b \cot(ex+d)^2+c \cot(ex+d)^4} (4ac-b^2)} + \frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c} \sqrt{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)}}{\cot(ex+d)^2+1} \right)}{(\sqrt{-4ac+b^2-b+2c})(\sqrt{-4ac+b^2+b-2c})\sqrt{a-b+c}}$

```
[In] int(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x,method=_RETURNVE
RBOSE)
```

```
[Out] 1/e*(-1/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)*(b+2*c*cot(e*x+d)^2)/(4*a*c
-b^2)+2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/((-4*a*c+b^2)^(1/2)+b-2*c)/(a-b+c)^(1/
2)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2)*((cot(e*x+d)^2+
1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1))+2*c/((-4*a*
c+b^2)^(1/2)+b-2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c
)*((cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)^2*c-(-4*a*c+b^2)^(1/2)*(cot(
e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c))^(1/2)-2*c/((-4*a*c+b^2)^(1/2)-b+2*c
)/(-4*a*c+b^2)/(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2-
1/2*(-b+(-4*a*c+b^2)^(1/2))/c)^2*c+(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2-1/2*(-b
+(-4*a*c+b^2)^(1/2))/c))^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 860 vs. $2(141) = 282$.

Time = 0.62 (sec) , antiderivative size = 1717, normalized size of antiderivative = 11.22

$$\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{4} * ((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*\cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*\cos(2*e*x + 2*d))*\sqrt{a - b + c} * \log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*\cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 - 2*((a - b + c)*\cos(2*e*x + 2*d)^2 - (2*a - b)*\cos(2*e*x + 2*d) + a - c)*\sqrt{a - b + c})*\sqrt{((a - b + c)*\cos(2*e*x + 2*d)^2 - 2*(a - c)*\cos(2*e*x + 2*d) + a + b + c)/(\cos(2*e*x + 2*d)^2 - 2*\cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*\cos(2*e*x + 2*d)) + 4*(a^2*b - a*b^2 + b^2*c - b*c^2 + (a^2*b - a*b^2 - (4*a - b)*c^2 - (4*a^2 - 6*a*b + b^2)*c)*\cos(2*e*x + 2*d)^2 - 2*(a^2*b - a*b^2 - 2*a*c^2 - (2*a^2 - 3*a*b)*c)*\cos(2*e*x + 2*d))*\sqrt{((a - b + c)*\cos(2*e*x + 2*d)^2 - 2*(a - c)*\cos(2*e*x + 2*d) + a + b + c)/(\cos(2*e*x + 2*d)^2 - 2*\cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*\cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*\cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e), 1/2*((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*\cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*\cos(2*e*x + 2*d))*\sqrt{-a + b - c} * \arctan(((a - b + c)*\cos(2*e*x + 2*d)^2 - (2*a - b)*\cos(2*e*x + 2*d) + a - c)*\sqrt{-a + b - c} * \sqrt{((a - b + c)*\cos(2*e*x + 2*d)^2 - 2*(a - c)*\cos(2*e*x + 2*d) + a + b + c)/(\cos(2*e*x + 2*d)^2 - 2*\cos(2*e*x + 2*d) + 1)))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*\cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*\cos(2*e*x + 2*d))) + 2*(a^2*b - a*b^2 + b^2*c - b*c^2 + (a^2*b - a*b^2 - (4*a - b)*c^2 - (4*a^2 - 6*a*b + b^2)*c)*\cos(2*e*x + 2*d)^2 - 2*(a^2*b - a*b^2 - 2*a*c^2 - (2*a^2 - 3*a*b)*c)*\cos(2*e*x + 2*d))*\sqrt{((a - b + c)*\cos(2*e*x + 2*d)^2 - 2*(a - c)*\cos(2*e*x + 2*d) + a + b + c)/(\cos(2*e*x + 2*d)^2 - 2*\cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*\cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4$

$*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*\cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e]$

Sympy [F]

$$\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(cot(e*x+d)**3/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(cot(d + e*x)**3/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima [F]

$$\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(ex+d)^3}{(c\cot(ex+d)^4+b\cot(ex+d)^2+a)^{\frac{3}{2}}} dx$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)^3/(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)^(3/2), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)^3}{(c\cot(d+ex)^4+b\cot(d+ex)^2+a)^{3/2}} dx$$

```
[In] int(cot(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

```
[Out] int(cot(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

$$3.30 \quad \int \frac{\cot(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	278
Rubi [A] (verified)	278
Mathematica [A] (verified)	280
Maple [B] (verified)	281
Fricas [B] (verification not implemented)	281
Sympy [F]	283
Maxima [F]	283
Giac [F(-1)]	283
Mupad [F(-1)]	283

Optimal result

Integrand size = 33, antiderivative size = 156

$$\int \frac{\cot(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} - \frac{b^2-2ac-bc+(b-2c)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[Out] 1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/(a-b+c)^(3/2)/e+(-b^2+2*a*c+b*c-(b-2*c)*c*cot(e*x+d)^2)/(a-b+c)/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 156, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3782, 1261, 754, 12, 738, 212}

$$\int \frac{\cot(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}} - \frac{-2ac+b^2+c(b-2c)\cot^2(d+ex)-bc}{e(a-b+c)(b^2-4ac)\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

[In] Int[Cot[d + e*x]/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

[Out] ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*(a - b + c)^(3/2)*e) - (b^2 - 2*a*c

$$- b*c + (b - 2*c)*c*\text{Cot}[d + e*x]^2)/((a - b + c)*(b^2 - 4*a*c)*e*\text{Sqrt}[a + b*\text{Cot}[d + e*x]^2 + c*\text{Cot}[d + e*x]^4])$$

Rule 12

$$\text{Int}[(a_)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match} Q[u, (b_)*(v_)] /; \text{FreeQ}[b, x]$$

Rule 212

$$\text{Int}[((a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{Gt} Q[a, 0] \ || \ \text{Lt} Q[b, 0])$$

Rule 738

$$\text{Int}[1/(((d_)+(e_)*(x_))*\text{Sqrt}[(a_)+(b_)*(x_)+(c_)*(x_)^2]), x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/\text{Sqrt}[a + b*x + c*x^2]], x] /; \text{FreeQ}\{a, b, c, d, e\}, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0]$$

Rule 754

$$\text{Int}[((d_)+(e_)*(x_))^m*((a_)+(b_)*(x_)+(c_)*(x_)^2)^p, x_Symbol] \rightarrow \text{Simp}[(d + e*x)^{m+1}*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^{p+1}/((p+1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + \text{Dist}[1/((p+1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), \text{Int}[(d + e*x)^m*\text{Simp}[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^{p+1}, x], x] /; \text{FreeQ}\{a, b, c, d, e, m\}, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{NeQ}[2*c*d - b*e, 0] \ \&\& \ \text{Lt} Q[p, -1] \ \&\& \ \text{IntQuadratic} Q[a, b, c, d, e, m, p, x]$$

Rule 1261

$$\text{Int}[(x_)*((d_)+(e_)*(x_)^2)^q*((a_)+(b_)*(x_)^2+(c_)*(x_)^4)^p, x_Symbol] \rightarrow \text{Dist}[1/2, \text{Subst}[\text{Int}[(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; \text{FreeQ}\{a, b, c, d, e, p, q\}, x]$$

Rule 3782

$$\text{Int}[\text{cot}[(d_)+(e_)*(x_)]^m*((a_)+(b_)*(\text{cot}[(d_)+(e_)*(x_)]*(f_))^{n_})+(c_)*(\text{cot}[(d_)+(e_)*(x_)]*(f_))^{n_2})^p, x_Symbol] \rightarrow \text{Dist}[-f/e, \text{Subst}[\text{Int}[(x/f)^m*((a + b*x^n + c*x^{2*n})^p/(f^2 + x^2)), x], x, f*\text{Cot}[d + e*x]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x\} \ \&\& \ \text{Eq} Q[n, 2, 2*n] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$$

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{x}{(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{1}{(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{b^2-2ac-bc+(b-2c)c\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-\frac{b^2}{2}+2ac}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{(a-b+c)(b^2-4ac)e} \\
&= -\frac{b^2-2ac-bc+(b-2c)c\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2(a-b+c)e} \\
&= -\frac{b^2-2ac-bc+(b-2c)c\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c)\cot^2(d+ex)}{\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{(a-b+c)e} \\
&= \frac{\text{arctanh}\left(\frac{2a-b+(-b+2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} \\
&\quad - \frac{b^2-2ac-bc+(b-2c)c\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 10.90 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.69

$$\int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \frac{2\sqrt{2}(b^2-2c(a+c)-(b^2-2bc+2c(-a+c))\cos(2(d+ex)))\csc^2(d+ex)}{(a-b+c)(-b^2+4ac)\sqrt{(3a+b+3c-4(a-c)\cos(2(d+ex)))+(a-b+c)\cos(4(d+ex))})\csc^2(d+ex)}$$

[In] Integrate[Cot[d + e*x]/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

[Out] ((2*Sqrt[2]*(b^2 - 2*c*(a + c) - (b^2 - 2*b*c + 2*c*(-a + c))*Cos[2*(d + e*x)])*Csc[d + e*x]^2)/((a - b + c)*(-b^2 + 4*a*c)*Sqrt[(3*a + b + 3*c - 4*(a - c)*Cos[2*(d + e*x)] + (a - b + c)*Cos[4*(d + e*x)])]*Csc[d + e*x]^4) - (

ArcTanh[(-b + 2*c + (-2*a + b)*Tan[d + e*x]^2)/(2*sqrt[a - b + c]*sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])]*sqrt[a + b*cot[d + e*x]^2 + c*cot[d + e*x]^4]*Tan[d + e*x]^2)/((a - b + c)^(3/2)*sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])/(2*e)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 405 vs. 2(145) = 290.

Time = 0.09 (sec) , antiderivative size = 406, normalized size of antiderivative = 2.60

method	result
derivativedivides	$\frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c} \sqrt{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}}{\cot(ex+d)^2+1} \right)}{(\sqrt{-4ac+b^2}-b+2c)(\sqrt{-4ac+b^2}+b-2c)\sqrt{a-b+c}} - \frac{2c \sqrt{(\cot(ex+d)^2+1)}}{(\sqrt{-4ac+b^2}-b+2c)}$
default	$\frac{2c \ln \left(\frac{2a-2b+2c+(b-2c)(\cot(ex+d)^2+1)+2\sqrt{a-b+c} \sqrt{(\cot(ex+d)^2+1)^2 c+(b-2c)(\cot(ex+d)^2+1)+a-b+c}}{\cot(ex+d)^2+1} \right)}{(\sqrt{-4ac+b^2}-b+2c)(\sqrt{-4ac+b^2}+b-2c)\sqrt{a-b+c}} - \frac{2c \sqrt{(\cot(ex+d)^2+1)}}{(\sqrt{-4ac+b^2}-b+2c)}$

[In] int(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x,method=_RETURNVERB OSE)

[Out] 1/e*(-2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/((-4*a*c+b^2)^(1/2)+b-2*c)/(a-b+c)^(1/2)*ln((2*a-2*b+2*c+(b-2*c)*(cot(e*x+d)^2+1)+2*(a-b+c)^(1/2)*((cot(e*x+d)^2+1)^2*c+(b-2*c)*(cot(e*x+d)^2+1)+a-b+c)^(1/2))/(cot(e*x+d)^2+1))-2*c/((-4*a*c+b^2)^(1/2)+b-2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c)^2*c-(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2+1/2*(b+(-4*a*c+b^2)^(1/2))/c))^(1/2)+2*c/((-4*a*c+b^2)^(1/2)-b+2*c)/(-4*a*c+b^2)/(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)*((cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c)^2*c+(-4*a*c+b^2)^(1/2)*(cot(e*x+d)^2-1/2*(-b+(-4*a*c+b^2)^(1/2))/c))^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 887 vs. 2(144) = 288.

Time = 0.66 (sec) , antiderivative size = 1771, normalized size of antiderivative = 11.35

$$\int \frac{\cot(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")

```
[Out] [1/4*((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*cos(2*e*x + 2*d))*sqrt(a - b + c)*log(2*(a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + 2*a^2 - b^2 + 2*c^2 + 2*((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(a - b + c)*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)) - 4*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d) - 4*(a*b^2 - b^3 - 2*(2*a - b)*c^2 - 2*c^3 + (a*b^2 - b^3 - 4*b*c^2 + 2*c^3 - (2*a^2 - 3*b^2)*c)*cos(2*e*x + 2*d)^2 - (2*a^2 - 2*a*b - b^2)*c - 2*(a*b^2 - b^3 - (2*a + b)*c^2 - (2*a^2 - a*b - 2*b^2)*c)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e), - 1/2*((a*b^2 + b^3 - 4*a*c^2 + (a*b^2 - b^3 - 4*a*c^2 - (4*a^2 - 4*a*b - b^2)*c)*cos(2*e*x + 2*d)^2 - (4*a^2 + 4*a*b - b^2)*c - 2*(a*b^2 + 4*a*c^2 - (4*a^2 + b^2)*c)*cos(2*e*x + 2*d))*sqrt(-a + b - c)*arctan(((a - b + c)*cos(2*e*x + 2*d)^2 - (2*a - b)*cos(2*e*x + 2*d) + a - c)*sqrt(-a + b - c)*sqrt((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1))/((a^2 - 2*a*b + b^2 + 2*(a - b)*c + c^2)*cos(2*e*x + 2*d)^2 + a^2 - b^2 + 2*a*c + c^2 - 2*(a^2 - a*b + b*c - c^2)*cos(2*e*x + 2*d))) + 2*(a*b^2 - b^3 - 2*(2*a - b)*c^2 - 2*c^3 + (a*b^2 - b^3 - 4*b*c^2 + 2*c^3 - (2*a^2 - 3*b^2)*c)*cos(2*e*x + 2*d)^2 - (2*a^2 - 2*a*b - b^2)*c - 2*(a*b^2 - b^3 - (2*a + b)*c^2 - (2*a^2 - a*b - 2*b^2)*c)*cos(2*e*x + 2*d))*sqrt(((a - b + c)*cos(2*e*x + 2*d)^2 - 2*(a - c)*cos(2*e*x + 2*d) + a + b + c)/(cos(2*e*x + 2*d)^2 - 2*cos(2*e*x + 2*d) + 1)))/((a^3*b^2 - 3*a^2*b^3 + 3*a*b^4 - b^5 - 4*a*c^4 - (12*a^2 - 12*a*b - b^2)*c^3 - 3*(4*a^3 - 8*a^2*b + 3*a*b^2 + b^3)*c^2 - (4*a^4 - 12*a^3*b + 9*a^2*b^2 + 2*a*b^3 - 3*b^4)*c)*e*cos(2*e*x + 2*d)^2 - 2*(a^3*b^2 - 2*a^2*b^3 + a*b^4 + 4*a*c^4 + (4*a^2 - 8*a*b - b^2)*c^3 - (4*a^3 - 3*a*b^2 - 2*b^3)*c^2 - (4*a^4 - 8*a^3*b + 3*a^2*b^2 + b^4)*c)*e*cos(2*e*x + 2*d) + (a^3*b^2 - a^2*b^3 - a*b^4 + b^5 - 4*a*c^4 - (12*a^2 - 4*a*b - b^2)*c^3 - (12*a^3 - 8*a^2*b - 7*a*b^2 + b^3)*c^2 - (4*a^4 - 4*a^3*b - 7*a^2*b^2 + 6*a*b^3 + b^4)*c)*e)]
```

Sympy [F]

$$\int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(cot(d + e*x)/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima [F]

$$\int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(ex+d)}{(c\cot(ex+d)^4+b\cot(ex+d)^2+a)^{3/2}} dx$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] integrate(cot(e*x + d)/(c*cot(e*x + d)^4 + b*cot(e*x + d)^2 + a)^(3/2), x)

Giac [F(-1)]

Timed out.

$$\int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(cot(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\cot(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\cot(d+ex)}{(c\cot(d+ex)^4+b\cot(d+ex)^2+a)^{3/2}} dx$$

[In] int(cot(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2),x)

[Out] int(cot(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)

$$3.31 \quad \int \frac{\tan(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	284
Rubi [A] (verified)	285
Mathematica [A] (verified)	288
Maple [F]	288
Fricas [B] (verification not implemented)	289
Sympy [F]	291
Maxima [F(-2)]	291
Giac [F(-1)]	291
Mupad [F(-1)]	292

Optimal result

Integrand size = 33, antiderivative size = 280

$$\int \frac{\tan(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2a^{3/2}e}$$

$$- \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e}$$

$$- \frac{b^2-2ac+bc \cot^2(d+ex)}{a(b^2-4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$+ \frac{b^2-2ac-bc+(b-2c)c \cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

```
[Out] 1/2*arctanh(1/2*(2*a+b*cot(e*x+d)^2)/a^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/a^(3/2)/e-1/2*arctanh(1/2*(2*a-b+(b-2*c)*cot(e*x+d)^2)/(a-b+c)^(1/2)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2))/(a-b+c)^(3/2)/e+(-b^2+2*a*c-b*c*cot(e*x+d)^2)/a/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)+(b^2-2*a*c-b*c+(b-2*c)*c*cot(e*x+d)^2)/(a-b+c)/(-4*a*c+b^2)/e/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(1/2)
```

Rubi [A] (verified)

Time = 0.47 (sec) , antiderivative size = 280, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {3782, 1265, 974, 754, 12, 738, 212}

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \frac{\operatorname{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2a^{3/2}e}$$

$$- \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c)\cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}}$$

$$- \frac{-2ac+b^2+bc\cot^2(d+ex)}{ae(b^2-4ac)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}$$

$$+ \frac{-2ac+b^2+c(b-2c)\cot^2(d+ex)-bc}{e(a-b+c)(b^2-4ac)\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}$$

[In] Int[Tan[d + e*x]/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

[Out] ArcTanh[(2*a + b*Cot[d + e*x]^2)/(2*Sqrt[a]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*a^(3/2)*e) - ArcTanh[(2*a - b + (b - 2*c)*Cot[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])]/(2*(a - b + c)^(3/2)*e) - (b^2 - 2*a*c + b*c*Cot[d + e*x]^2)/(a*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]) + (b^2 - 2*a*c - b*c + (b - 2*c)*c*Cot[d + e*x]^2)/((a - b + c)*(b^2 - 4*a*c)*e*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4])

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 754

```

Int[((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x]
+ Dist[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]

```

Rule 974

```

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])

```

Rule 1265

```

Int[(x_)^(m_.)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_.), x_Symbol]
:> Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

```

Rule 3782

```

Int[cot[(d_.) + (e_.)*(x_)]^(m_.)*((a_.) + (b_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n_.) + (c_.)*(cot[(d_.) + (e_.)*(x_)]*(f_.))^(n2_.))^(p_), x_Symbol]
:> Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
&= -\frac{\text{Subst}\left(\int \frac{1}{x(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= -\frac{\text{Subst}\left(\int \left(\frac{1}{(-1-x)(a+bx+cx^2)^{3/2}} + \frac{1}{x(a+bx+cx^2)^{3/2}}\right) dx, x, \cot^2(d+ex)\right)}{2e}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{\text{Subst}\left(\int \frac{1}{(-1-x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{x(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
&= - \frac{b^2 - 2ac + bc \cot^2(d+ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad + \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d+ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-\frac{b^2}{2} + 2ac}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{a(b^2 - 4ac) e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-\frac{b^2}{2} + 2ac}{(-1-x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{(a - b + c)(b^2 - 4ac) e} \\
&= - \frac{b^2 - 2ac + bc \cot^2(d+ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad + \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d+ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2ae} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(-1-x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d+ex)\right)}{2(a - b + c)e} \\
&= - \frac{b^2 - 2ac + bc \cot^2(d+ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad + \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d+ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d+ex) + c \cot^4(d+ex)}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{ae} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{-2a+b-(b-2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{(a - b + c)e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\operatorname{arctanh}\left(\frac{2a+b\cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2a^{3/2}e} \\
&- \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c)\cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e} \\
&- \frac{b^2-2ac+bc\cot^2(d+ex)}{a(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}} \\
&+ \frac{b^2-2ac-bc+(b-2c)c\cot^2(d+ex)}{(a-b+c)(b^2-4ac)e\sqrt{a+b\cot^2(d+ex)+c\cot^4(d+ex)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 13.75 (sec) , antiderivative size = 351, normalized size of antiderivative = 1.25

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \frac{2\sqrt{2}\sqrt{a}(-b^3+bc(3a+c)+(b^3-2b^2c+4ac^2+bc(-3a+c))\cos(2(d+ex)))\operatorname{csc}^2(d+ex)}{(a-b+c)(-b^2+4ac)\sqrt{(3a+b+3c-4(a-c)\cos(2(d+ex))+(a-b+c)\cos(4(d+ex)))}\operatorname{cs}}$$

[In] Integrate[Tan[d + e*x]/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2), x]

[Out] ((2*sqrt[2]*sqrt[a]*(-b^3 + b*c*(3*a + c) + (b^3 - 2*b^2*c + 4*a*c^2 + b*c*(-3*a + c))*Cos[2*(d + e*x)])*Csc[d + e*x]^2)/((a - b + c)*(-b^2 + 4*a*c)*Sqrt[(3*a + b + 3*c - 4*(a - c)*Cos[2*(d + e*x)] + (a - b + c)*Cos[4*(d + e*x)])]*Csc[d + e*x]^4) + (((a - b + c)^(3/2)*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*sqrt[a]*sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]]) + a^(3/2)*ArcTanh[(-b + 2*c + (-2*a + b)*Tan[d + e*x]^2)/(2*sqrt[a - b + c]*sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]])]*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/((a - b + c)^(3/2)*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4]))/(2*a^(3/2)*e)

Maple [F]

$$\int \frac{\tan(ex+d)}{(a+b\cot(ex+d)^2+c\cot(ex+d)^4)^{3/2}} dx$$

[In] int(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2), x)

[Out] int(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2), x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1020 vs. 2(256) = 512.

Time = 2.67 (sec) , antiderivative size = 4153, normalized size of antiderivative = 14.83

$$\int \frac{\tan(d + ex)}{(a + b \cot^2(d + ex) + c \cot^4(d + ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")

[Out] [-1/4*((4*a*c^4 - (a^3*b^2 - 2*a^2*b^3 + a*b^4 - 4*a^2*c^3 - (8*a^3 - 8*a^2*b - a*b^2)*c^2 - 2*(2*a^4 - 4*a^3*b + a^2*b^2 + a*b^3)*c)*tan(e*x + d)^4 + (8*a^2 - 8*a*b - b^2)*c^3 + 2*(2*a^3 - 4*a^2*b + a*b^2 + b^3)*c^2 - (a^2*b^3 - 2*a*b^4 + b^5 - 4*a*b*c^3 - (8*a^2*b - 8*a*b^2 - b^3)*c^2 - 2*(2*a^3*b - 4*a^2*b^2 + a*b^3 + b^4)*c)*tan(e*x + d)^2 - (a^2*b^2 - 2*a*b^3 + b^4)*c)*sqrt(a)*log(8*a^2*tan(e*x + d)^4 + 8*a*b*tan(e*x + d)^2 + b^2 + 4*a*c + 4*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)) - (a^2*b^2*c - 4*a^3*c^2 + (a^3*b^2 - 4*a^4*c)*tan(e*x + d)^4 + (a^2*b^3 - 4*a^3*b*c)*tan(e*x + d)^2)*sqrt(a - b + c)*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2)*sqrt(a - b + c)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/(tan(e*x + d)^4 + 2*tan(e*x + d)^2 + 1)) - 4*((a^2*b^3 - a*b^4 + 2*a^2*c^3 + (2*a^3 - 5*a^2*b - a*b^2)*c^2 - (3*a^3*b - 2*a^2*b^2 - 2*a*b^3)*c)*tan(e*x + d)^4 - ((2*a^2 + a*b)*c^3 + (2*a^3 - a^2*b - 2*a*b^2)*c^2 - (a^2*b^2 - a*b^3)*c)*tan(e*x + d)^2)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4))/((a^5*b^2 - 2*a^4*b^3 + a^3*b^4 - 4*a^4*c^3 - (8*a^5 - 8*a^4*b - a^3*b^2)*c^2 - 2*(2*a^6 - 4*a^5*b + a^4*b^2 + a^3*b^3)*c)*e*tan(e*x + d)^4 + (a^4*b^3 - 2*a^3*b^4 + a^2*b^5 - 4*a^3*b*c^3 - (8*a^4*b - 8*a^3*b^2 - a^2*b^3)*c^2 - 2*(2*a^5*b - 4*a^4*b^2 + a^3*b^3 + a^2*b^4)*c)*e*tan(e*x + d)^2 - (4*a^3*c^4 + (8*a^4 - 8*a^3*b - a^2*b^2)*c^3 + 2*(2*a^5 - 4*a^4*b + a^3*b^2 + a^2*b^3)*c^2 - (a^4*b^2 - 2*a^3*b^3 + a^2*b^4)*c)*e), 1/4*(2*(4*a*c^4 - (a^3*b^2 - 2*a^2*b^3 + a*b^4 - 4*a^2*c^3 - (8*a^3 - 8*a^2*b - a*b^2)*c^2 - 2*(2*a^4 - 4*a^3*b + a^2*b^2 + a*b^3)*c)*tan(e*x + d)^4 + (8*a^2 - 8*a*b - b^2)*c^3 + 2*(2*a^3 - 4*a^2*b + a*b^2 + b^3)*c^2 - (a^2*b^3 - 2*a*b^4 + b^5 - 4*a*b*c^3 - (8*a^2*b - 8*a*b^2 - b^3)*c^2 - 2*(2*a^3*b - 4*a^2*b^2 + a*b^3 + b^4)*c)*tan(e*x + d)^2 - (a^2*b^2 - 2*a*b^3 + b^4)*c)*sqrt(-a)*arctan(1/2*(2*a*tan(e*x + d)^4 + b*tan(e*x + d)^2)*sqrt(-a)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 + c)/tan(e*x + d)^4)/(a^2*tan(e*x + d)^4 + a*b*tan(e*x + d)^2 + a*c)) + (a^2*b^2*c - 4*a^3*c^2 + (a^3*b^2 - 4*a^4*c)*tan(e*x + d)^4 + (a^2*b^3 - 4*a^3*b*c)*tan(e*x + d)^2)*sqrt(a - b + c)*log(((8*a^2 - 8*a*b + b^2 + 4*a*c)*tan(e*x + d)^4 + 2*(4*a*b - 3*b^2 - 4*(a - b)*c)*tan(e*x + d)^2 + b^2 + 4*(a - 2*b)*c + 8*c^2 - 4*((2*a - b)*tan(e*x + d)^4 + (b - 2*c)*tan(e*x + d)^2))

$$\begin{aligned}
& \sqrt{a-b+c} \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / (\tan(e^x+d)^4 + 2 \tan(e^x+d)^2 + 1) + 4((a^2 b^3 - a b^4 + 2 a^2 c^3 + (2 a^3 - 5 a^2 b - a b^2) c^2 - (3 a^3 b - 2 a^2 b^2 - 2 a b^3) c) * \tan(e^x+d)^4 - ((2 a^2 + a b) c^3 + (2 a^3 - a^2 b - 2 a b^2) c^2 - (a^2 b^2 - a b^3) c) * \tan(e^x+d)^2) * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / ((a^5 b^2 - 2 a^4 b^3 + a^3 b^4 - 4 a^4 c^3 - (8 a^5 - 8 a^4 b - a^3 b^2) c^2 - 2(2 a^6 - 4 a^5 b + a^4 b^2 + a^3 b^3) c) * e \tan(e^x+d)^4 + (a^4 b^3 - 2 a^3 b^4 + a^2 b^5 - 4 a^3 b c^3 - (8 a^4 b - 8 a^3 b^2 - a^2 b^3) c^2 - 2(2 a^5 b - 4 a^4 b^2 + a^3 b^3 + a^2 b^4) c) * e \tan(e^x+d)^2 - (4 a^3 c^4 + (8 a^4 - 8 a^3 b - a^2 b^2) c^3 + 2(2 a^5 - 4 a^4 b + a^3 b^2 + a^2 b^3) c^2 - (a^4 b^2 - 2 a^3 b^3 + a^2 b^4) c) * e), -1/4 * (2(a^2 b^2 c - 4 a^3 c^2 + (a^3 b^2 - 4 a^4 c) \tan(e^x+d)^4 + (a^2 b^3 - 4 a^3 b c) \tan(e^x+d)^2) * \sqrt{-a+b-c} * \arctan(-1/2 * ((2 a - b) \tan(e^x+d)^4 + (b - 2 c) \tan(e^x+d)^2) * \sqrt{-a+b-c} * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / ((a^2 - a b + a c) \tan(e^x+d)^4 + (a b - b^2 + b c) \tan(e^x+d)^2 + (a - b) c + c^2)) + (4 a c^4 - (a^3 b^2 - 2 a^2 b^3 + a b^4 - 4 a^2 c^3 - (8 a^3 - 8 a^2 b - a b^2) c^2 - 2(2 a^4 - 4 a^3 b + a^2 b^2 + a b^3) c) * \tan(e^x+d)^4 + (8 a^2 - 8 a b - b^2) c^3 + 2(2 a^3 - 4 a^2 b + a b^2 + b^3) c^2 - (a^2 b^3 - 2 a b^4 + b^5 - 4 a b c^3 - (8 a^2 b - 8 a b^2 - b^3) c^2 - 2(2 a^3 b - 4 a^2 b^2 + a b^3 + b^4) c) * \tan(e^x+d)^2 - (a^2 b^2 - 2 a b^3 + b^4) c) * \sqrt{a} * \log(8 a^2 \tan(e^x+d)^4 + 8 a b \tan(e^x+d)^2 + b^2 + 4 a c + 4(2 a \tan(e^x+d)^4 + b \tan(e^x+d)^2) * \sqrt{a} * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4}) - 4((a^2 b^3 - a b^4 + 2 a^2 c^3 + (2 a^3 - 5 a^2 b - a b^2) c^2 - (3 a^3 b - 2 a^2 b^2 - 2 a b^3) c) * \tan(e^x+d)^4 - ((2 a^2 + a b) c^3 + (2 a^3 - a^2 b - 2 a b^2) c^2 - (a^2 b^2 - a b^3) c) * \tan(e^x+d)^2) * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / ((a^5 b^2 - 2 a^4 b^3 + a^3 b^4 - 4 a^4 c^3 - (8 a^5 - 8 a^4 b - a^3 b^2) c^2 - 2(2 a^6 - 4 a^5 b + a^4 b^2 + a^3 b^3) c) * e \tan(e^x+d)^4 + (a^4 b^3 - 2 a^3 b^4 + a^2 b^5 - 4 a^3 b c^3 - (8 a^4 b - 8 a^3 b^2 - a^2 b^3) c^2 - 2(2 a^5 b - 4 a^4 b^2 + a^3 b^3 + a^2 b^4) c) * e \tan(e^x+d)^2 - (4 a^3 c^4 + (8 a^4 - 8 a^3 b - a^2 b^2) c^3 + 2(2 a^5 - 4 a^4 b + a^3 b^2 + a^2 b^3) c^2 - (a^4 b^2 - 2 a^3 b^3 + a^2 b^4) c) * e), 1/2 * ((4 a c^4 - (a^3 b^2 - 2 a^2 b^3 + a b^4 - 4 a^2 c^3 - (8 a^3 - 8 a^2 b - a b^2) c^2 - 2(2 a^4 - 4 a^3 b + a^2 b^2 + a b^3) c) * \tan(e^x+d)^4 + (8 a^2 - 8 a b - b^2) c^3 + 2(2 a^3 - 4 a^2 b + a b^2 + b^3) c^2 - (a^2 b^3 - 2 a b^4 + b^5 - 4 a b c^3 - (8 a^2 b - 8 a b^2 - b^3) c^2 - 2(2 a^3 b - 4 a^2 b^2 + a b^3 + b^4) c) * \tan(e^x+d)^2 - (a^2 b^2 - 2 a b^3 + b^4) c) * \sqrt{-a} * \arctan(1/2 * (2 a \tan(e^x+d)^4 + b \tan(e^x+d)^2) * \sqrt{-a} * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / (a^2 \tan(e^x+d)^4 + a b \tan(e^x+d)^2 + a c)) - (a^2 b^2 c - 4 a^3 c^2 + (a^3 b^2 - 4 a^4 c) \tan(e^x+d)^4 + (a^2 b^3 - 4 a^3 b c) \tan(e^x+d)^2) * \sqrt{-a+b-c} * \arctan(-1/2 * ((2 a - b) \tan(e^x+d)^4 + (b - 2 c) \tan(e^x+d)^2) * \sqrt{-a+b-c} * \sqrt{(a \tan(e^x+d)^4 + b \tan(e^x+d)^2 + c) / \tan(e^x+d)^4} / ((a^2 - a b + a c) \tan(e^x+d)^4 + (a b - b^2 + b c) \tan(e^x+d)^2 + (a - b) c + c^2)) + 2((a^2 b^3 - a b^4 + 2 a
\end{aligned}$$

$$\begin{aligned} &^2*c^3 + (2*a^3 - 5*a^2*b - a*b^2)*c^2 - (3*a^3*b - 2*a^2*b^2 - 2*a*b^3)*c) \\ &*tan(e*x + d)^4 - ((2*a^2 + a*b)*c^3 + (2*a^3 - a^2*b - 2*a*b^2)*c^2 - (a^2 \\ &*b^2 - a*b^3)*c)*tan(e*x + d)^2)*sqrt((a*tan(e*x + d)^4 + b*tan(e*x + d)^2 \\ &+ c)/tan(e*x + d)^4))/((a^5*b^2 - 2*a^4*b^3 + a^3*b^4 - 4*a^4*c^3 - (8*a^5 \\ &- 8*a^4*b - a^3*b^2)*c^2 - 2*(2*a^6 - 4*a^5*b + a^4*b^2 + a^3*b^3)*c)*e*tan \\ &(e*x + d)^4 + (a^4*b^3 - 2*a^3*b^4 + a^2*b^5 - 4*a^3*b*c^3 - (8*a^4*b - 8*a \\ &^3*b^2 - a^2*b^3)*c^2 - 2*(2*a^5*b - 4*a^4*b^2 + a^3*b^3 + a^2*b^4)*c)*e*ta \\ &n(e*x + d)^2 - (4*a^3*c^4 + (8*a^4 - 8*a^3*b - a^2*b^2)*c^3 + 2*(2*a^5 - 4* \\ &a^4*b + a^3*b^2 + a^2*b^3)*c^2 - (a^4*b^2 - 2*a^3*b^3 + a^2*b^4)*c)*e)] \end{aligned}$$

Sympy [F]

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(tan(d + e*x)/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima [F(-2)]

Exception generated.

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Exception raised: RuntimeError}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Giac [F(-1)]

Timed out.

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\tan(d+ex)}{(c\cot(d+ex)^4+b\cot(d+ex)^2+a)^{3/2}} dx$$

```
[In] int(tan(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

```
[Out] int(tan(d + e*x)/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2), x)
```

$$3.32 \quad \int \frac{\tan^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx$$

Optimal result	293
Rubi [A] (verified)	294
Mathematica [A] (warning: unable to verify)	299
Maple [F]	299
Fricas [B] (verification not implemented)	299
Sympy [F]	300
Maxima [F(-1)]	300
Giac [F(-1)]	300
Mupad [F(-1)]	301

Optimal result

Integrand size = 35, antiderivative size = 478

$$\int \frac{\tan^3(d+ex)}{(a+b \cot^2(d+ex)+c \cot^4(d+ex))^{3/2}} dx =$$

$$\frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2a^{3/2}e}$$

$$- \frac{3b \operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4a^{5/2}e}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2}e}$$

$$+ \frac{b^2 - 2ac + bc \cot^2(d+ex)}{a(b^2 - 4ac) e \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$- \frac{b^2 - 2ac - bc + (b-2c)c \cot^2(d+ex)}{(a-b+c)(b^2 - 4ac) e \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$- \frac{(b^2 - 2ac + bc \cot^2(d+ex)) \tan^2(d+ex)}{a(b^2 - 4ac) e \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}$$

$$+ \frac{(3b^2 - 8ac) \sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)} \tan^2(d+ex)}{2a^2 (b^2 - 4ac) e}$$

[Out] $-1/2*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d)^2)/a^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/a^{(3/2)}/e-3/4*b*\operatorname{arctanh}(1/2*(2*a+b*\cot(e*x+d)^2)/a^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/a^{(5/2)}/e+1/2*\operatorname{arctanh}(1/2*(2*a-b+(b-2*c)*\cot(e*x+d)^2)/(a-b+c)^{(1/2)}/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)})/(a-b+c)^{(3/2)}/e+(b^2-2*a*c+b*c*\cot(e*x+d)^2)/a/(-4*a*c+b^2)/e/(a+b*\cot(e*x+d)^2+c*\cot(e*x+d)^4)^{(1/2)}+(-b^2+2*a*c+b*c-(b-2*c)*c*\cot(e*x+d)^2)/(a-b+c)/(-4*a$

$$\frac{c+b^2}{e} / (a+b \cot(e*x+d)^2 + c \cot(e*x+d)^4)^{1/2} - (b^2 - 2*a*c + b*c \cot(e*x+d)^2) * \tan(e*x+d)^2 / a / (-4*a*c + b^2) / e / (a+b \cot(e*x+d)^2 + c \cot(e*x+d)^4)^{1/2} + 1/2 * (-8*a*c + 3*b^2) * (a+b \cot(e*x+d)^2 + c \cot(e*x+d)^4)^{1/2} * \tan(e*x+d)^2 / a^2 / (-4*a*c + b^2) / e$$

Rubi [A] (verified)

Time = 0.67 (sec) , antiderivative size = 478, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.229$, Rules used = {3782, 1265, 974, 754, 820, 738, 212, 12}

$$\int \frac{\tan^3(d+ex)}{(a+b \cot^2(d+ex) + c \cot^4(d+ex))^{3/2}} dx =$$

$$\frac{3b \operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4a^{5/2}e}$$

$$- \frac{\operatorname{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2a^{3/2}e}$$

$$+ \frac{(3b^2 - 8ac) \tan^2(d+ex) \sqrt{a+b \cot^2(d+ex) + c \cot^4(d+ex)}}{2a^2e(b^2 - 4ac)}$$

$$+ \frac{\operatorname{arctanh}\left(\frac{2a+(b-2c) \cot^2(d+ex)-b}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2e(a-b+c)^{3/2}}$$

$$+ \frac{-2ac + b^2 + bc \cot^2(d+ex)}{ae(b^2 - 4ac) \sqrt{a+b \cot^2(d+ex) + c \cot^4(d+ex)}}$$

$$- \frac{-2ac + b^2 + c(b-2c) \cot^2(d+ex) - bc}{e(a-b+c)(b^2 - 4ac) \sqrt{a+b \cot^2(d+ex) + c \cot^4(d+ex)}}$$

$$- \frac{\tan^2(d+ex) (-2ac + b^2 + bc \cot^2(d+ex))}{ae(b^2 - 4ac) \sqrt{a+b \cot^2(d+ex) + c \cot^4(d+ex)}}$$

[In] Int[Tan[d + e*x]^3/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2),x]

[Out] $-1/2 * \operatorname{ArcTanh}[(2*a + b * \operatorname{Cot}[d + e*x]^2) / (2 * \operatorname{Sqrt}[a] * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4])] / (a^{3/2} * e) - (3 * b * \operatorname{ArcTanh}[(2*a + b * \operatorname{Cot}[d + e*x]^2) / (2 * \operatorname{Sqrt}[a] * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4)])] / (4 * a^{5/2} * e) + \operatorname{ArcTanh}[(2*a - b + (b - 2*c) * \operatorname{Cot}[d + e*x]^2) / (2 * \operatorname{Sqrt}[a - b + c] * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4))] / (2 * (a - b + c)^{3/2} * e) + (b^2 - 2*a*c + b*c * \operatorname{Cot}[d + e*x]^2) / (a * (b^2 - 4*a*c) * e * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4]) - (b^2 - 2*a*c - b*c + (b - 2*c) * c * \operatorname{Cot}[d + e*x]^2) / ((a - b + c) * (b^2 - 4*a*c) * e * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4]) - ((b^2 - 2*a*c + b*c * \operatorname{Cot}[d + e*x]^2) * \operatorname{Tan}[d + e*x]^2) / (a * (b^2 - 4*a*c) * e * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4]) + ((3 * b^2 - 8 * a * c) * \operatorname{Sqrt}[a + b * \operatorname{Cot}[d + e*x]^2 + c * \operatorname{Cot}[d + e*x]^4] * \operatorname{Tan}[d + e*x]^2) / (2 * a^2 * (b^2 - 4 * a * c) * e)$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 738

Int[1/(((d_) + (e_)*(x_))*Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2]), x_Symbol] := Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]

Rule 754

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*(b*c*d - b^2*e + 2*a*c*e + c*(2*c*d - b*e)*x)*((a + b*x + c*x^2)^(p + 1)/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/((p + 1)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^m*Simp[b*c*d*e*(2*p - m + 2) + b^2*e^2*(m + p + 2) - 2*c^2*d^2*(2*p + 3) - 2*a*c*e^2*(m + 2*p + 3) - c*e*(2*c*d - b*e)*(m + 2*p + 4)*x, x]*(a + b*x + c*x^2)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[2*c*d - b*e, 0] && LtQ[p, -1] && IntQuadraticQ[a, b, c, d, e, m, p, x]

Rule 820

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(-(e*f - d*g))*(d + e*x)^(m + 1)*((a + b*x + c*x^2)^(p + 1)/(2*(p + 1)*(c*d^2 - b*d*e + a*e^2))), x] - Dist[(b*(e*f + d*g) - 2*(c*d*f + a*e*g))/(2*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 974

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && (

IntegerQ[p] || (ILtQ[m, 0] && ILtQ[n, 0])) && !(IGtQ[m, 0] || IGtQ[n, 0])

Rule 1265

Int[(x_)^(m_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[x^((m - 1)/2)*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && IntegerQ[(m - 1)/2]

Rule 3782

Int[cot[(d_) + (e_)*(x_)]^(m_)*((a_) + (b_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n_) + (c_)*(cot[(d_) + (e_)*(x_)]*(f_))^(n2_))^(p_), x_Symbol] := Dist[-f/e, Subst[Int[(x/f)^m*((a + b*x^n + c*x^(2*n))^p/(f^2 + x^2)), x], x, f*Cot[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[n, 2, 2*n] && NeQ[b^2 - 4*a*c, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\text{Subst}\left(\int \frac{1}{x^3(1+x^2)(a+bx^2+cx^4)^{3/2}} dx, x, \cot(d+ex)\right)}{e} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^2(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^2(a+bx+cx^2)^{3/2}} - \frac{1}{x(a+bx+cx^2)^{3/2}} + \frac{1}{(1+x)(a+bx+cx^2)^{3/2}}\right) dx, x, \cot^2(d+ex)\right)}{2e} \\
 &= -\frac{\text{Subst}\left(\int \frac{1}{x^2(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &\quad + \frac{\text{Subst}\left(\int \frac{1}{x(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e} \\
 &\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)(a+bx+cx^2)^{3/2}} dx, x, \cot^2(d+ex)\right)}{2e}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 - 2ac + bc \cot^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d + ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{(b^2 - 2ac + bc \cot^2(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{-\frac{b^2}{2} + 2ac}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{a(b^2 - 4ac) e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{\frac{1}{2}(-3b^2 + 8ac) - bcx}{x^2\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{a(b^2 - 4ac) e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-\frac{b^2}{2} + 2ac}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{(a - b + c)(b^2 - 4ac) e} \\
&= \frac{b^2 - 2ac + bc \cot^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d + ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{(b^2 - 2ac + bc \cot^2(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad + \frac{(3b^2 - 8ac) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2a^2(b^2 - 4ac) e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{2ae} \\
&\quad + \frac{(3b)\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{4a^2 e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{(1+x)\sqrt{a+bx+cx^2}} dx, x, \cot^2(d + ex)\right)}{2(a - b + c)e}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^2 - 2ac + bc \cot^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d + ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{(b^2 - 2ac + bc \cot^2(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad + \frac{(3b^2 - 8ac) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2a^2 (b^2 - 4ac) e} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{ae} \\
&\quad - \frac{(3b)\text{Subst}\left(\int \frac{1}{4a-x^2} dx, x, \frac{2a+b \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2a^2 e} \\
&\quad + \frac{\text{Subst}\left(\int \frac{1}{4a-4b+4c-x^2} dx, x, \frac{2a-b-(-b+2c) \cot^2(d+ex)}{\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{(a-b+c)e} \\
&= - \frac{\text{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2a^{3/2} e} \\
&\quad - \frac{3b \text{arctanh}\left(\frac{2a+b \cot^2(d+ex)}{2\sqrt{a}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{4a^{5/2} e} \\
&\quad + \frac{\text{arctanh}\left(\frac{2a-b+(b-2c) \cot^2(d+ex)}{2\sqrt{a-b+c}\sqrt{a+b \cot^2(d+ex)+c \cot^4(d+ex)}}\right)}{2(a-b+c)^{3/2} e} \\
&\quad + \frac{b^2 - 2ac + bc \cot^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{b^2 - 2ac - bc + (b - 2c)c \cot^2(d + ex)}{(a - b + c)(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad - \frac{(b^2 - 2ac + bc \cot^2(d + ex)) \tan^2(d + ex)}{a(b^2 - 4ac) e \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)}} \\
&\quad + \frac{(3b^2 - 8ac) \sqrt{a + b \cot^2(d + ex) + c \cot^4(d + ex)} \tan^2(d + ex)}{2a^2 (b^2 - 4ac) e}
\end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 15.97 (sec) , antiderivative size = 562, normalized size of antiderivative = 1.18

$$\int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \frac{\sqrt{2}\sqrt{(3a+b+3c-4(a-c)\cos(2(d+ex))+(a-b+c)\cos(4(d+ex)))}}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}}$$

```
[In] Integrate[Tan[d + e*x]^3/(a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4)^(3/2),x]
[Out] (Sqrt[2]*Sqrt[(3*a + b + 3*c - 4*(a - c)*Cos[2*(d + e*x)] + (a - b + c)*Cos[4*(d + e*x)])*Csc[d + e*x]^4]*((3*b^2*(b - c)^2 - 4*a^3*c + a^2*(b^2 + 8*b*c - 4*c^2) - 2*a*(b^3 + 5*b^2*c - 10*b*c^2 + 4*c^3))/((a - b + c)^2*(-b^2 + 4*a*c)) + (8*(-b^5 + b^4*c + 2*a*c^3*(a + c) - b^2*c^2*(4*a + c) + b^3*c*(5*a + c) - a*b*c^2*(5*a + 3*c) + (b^5 - 3*b^4*c + a*b*(5*a - 9*c)*c^2 + b^2*(12*a - c)*c^2 + 2*a*c^3*(-3*a + c) + b^3*c*(-5*a + 3*c))*Cos[2*(d + e*x)]))/((a - b + c)^2*(-b^2 + 4*a*c)*(3*a + b + 3*c - 4*(a - c)*Cos[2*(d + e*x)] + (a - b + c)*Cos[4*(d + e*x)])) + Sec[d + e*x]^2 - (2*(((2*a + 3*b)*(a - b + c)*ArcTanh[(b + 2*a*Tan[d + e*x]^2)/(2*Sqrt[a]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])])/Sqrt[a] - (2*a^2*ArcTanh[(b - 2*c + (2*a - b)*Tan[d + e*x]^2)/(2*Sqrt[a - b + c]*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4])])/Sqrt[a - b + c])*Sqrt[a + b*Cot[d + e*x]^2 + c*Cot[d + e*x]^4]*Tan[d + e*x]^2)/((a - b + c)*Sqrt[c + b*Tan[d + e*x]^2 + a*Tan[d + e*x]^4)))/(8*a^2*e)
```

Maple [F]

$$\int \frac{\tan^3(ex+d)}{(a+b\cot^2(ex+d)+c\cot^4(ex+d))^{3/2}} dx$$

```
[In] int(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x)
```

```
[Out] int(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1300 vs. 2(438) = 876.

Time = 3.34 (sec) , antiderivative size = 5274, normalized size of antiderivative = 11.03

$$\int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Too large to display}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="fricas")

[Out] Too large to include

Sympy **[F]**

$$\int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{\frac{3}{2}}} dx$$

[In] integrate(tan(e*x+d)**3/(a+b*cot(e*x+d)**2+c*cot(e*x+d)**4)**(3/2),x)

[Out] Integral(tan(d + e*x)**3/(a + b*cot(d + e*x)**2 + c*cot(d + e*x)**4)**(3/2), x)

Maxima **[F(-1)]**

Timed out.

$$\int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="maxima")

[Out] Timed out

Giac **[F(-1)]**

Timed out.

$$\int \frac{\tan^3(d+ex)}{(a+b\cot^2(d+ex)+c\cot^4(d+ex))^{3/2}} dx = \text{Timed out}$$

[In] integrate(tan(e*x+d)^3/(a+b*cot(e*x+d)^2+c*cot(e*x+d)^4)^(3/2),x, algorithm="giac")

[Out] Timed out

Mupad [F(-1)]

Timed out.

$$\int \frac{\tan^3(d + ex)}{(a + b \cot^2(d + ex) + c \cot^4(d + ex))^{3/2}} dx = \text{Hanged}$$

```
[In] int(tan(d + e*x)^3/(a + b*cot(d + e*x)^2 + c*cot(d + e*x)^4)^(3/2),x)
```

```
[Out] \text{Hanged}
```

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 303

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well");
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```



```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(max(expnType_result,expnType_optimal))

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
# Albert Rich to use with Sagemath. This is used to
# grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
# 'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
# issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```