

Computer Algebra Independent Integration Tests

Summer 2023 edition

5-Inverse-trig-functions/5.5-Inverse-secant/157-5.5.2-Inverse-secant-
functions

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 1:58pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	39
4	Appendix	327

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [50]. This is test number [157].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (50)	0.00 (0)
Mathematica	98.00 (49)	2.00 (1)
Maple	74.00 (37)	26.00 (13)
Fricas	56.00 (28)	44.00 (22)
Giac	54.00 (27)	46.00 (23)
Maxima	36.00 (18)	64.00 (32)
Sympy	26.00 (13)	74.00 (37)
Mupad	20.00 (10)	80.00 (40)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	70.000	14.000	14.000	2.000
Maple	68.000	6.000	0.000	26.000
Fricas	48.000	8.000	0.000	44.000
Giac	32.000	22.000	0.000	46.000
Maxima	32.000	4.000	0.000	64.000
Sympy	6.000	0.000	20.000	74.000
Mupad	0.000	20.000	0.000	80.000

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	1	0.00	100.00	0.00
Maple	13	100.00	0.00	0.00
Fricas	22	90.91	0.00	9.09
Giac	23	91.30	0.00	8.70
Maxima	32	100.00	0.00	0.00
Sympy	37	91.89	8.11	0.00
Mupad	40	0.00	100.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.10
Maxima	0.21
Fricas	0.29
Giac	0.31
Mathematica	0.60
Maple	0.79
Mupad	0.92
Sympy	21.08

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	37.50	0.91	36.50	0.90
Maxima	56.67	1.21	54.50	1.20
Sympy	76.46	1.63	61.00	1.67
Fricas	102.61	1.37	56.50	0.92
Giac	120.67	1.65	82.00	1.72
Rubi	120.76	1.00	69.00	1.00
Maple	174.38	1.36	76.00	1.31
Mathematica	195.20	1.84	107.00	1.00

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {27, 28, 31, 36}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design v1.0

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	36

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 15, 16, 18, 19, 20, 21, 23, 27, 28, 29, 30, 33, 34, 35, 43, 44, 45, 46, 47, 48, 49, 50 }

B grade { 14, 31, 32, 36, 40, 41, 42 }

C grade { 17, 22, 24, 25, 26, 38, 39 }

F normal fail { }

F(-1) timedout fail { 37 }

F(-2) exception fail { }

Maple

A grade { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 27, 28, 29, 30, 32, 33, 34, 38, 39, 40, 42, 50 }

B grade { 24, 25, 26 }

C grade { }

F normal fail { 1, 31, 35, 36, 37, 41, 43, 44, 45, 46, 47, 48, 49 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 2, 3, 4, 5, 7, 8, 9, 10, 11, 12, 15, 18, 19, 20, 21, 25, 26, 38, 39, 40, 41, 47, 48, 49 }

B grade { 14, 16, 22, 24 }

C grade { }

F normal fail { 1, 6, 13, 23, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 43, 44, 45, 46, 50 }

F(-1) timeout fail { }

F(-2) exception fail { 17, 42 }

Maxima

A grade { 2, 3, 4, 5, 7, 10, 11, 12, 14, 15, 16, 22, 38, 39, 40, 41 }

B grade { 8, 9 }

C grade { }

F normal fail { 1, 6, 13, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 47, 48, 49, 50 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac**A grade** { 7, 8, 9, 10, 11, 12, 15, 16, 21, 24, 38, 39, 40, 41, 47, 50 }**B grade** { 2, 3, 4, 5, 14, 18, 19, 20, 22, 25, 26 }**C grade** { }**F normal fail** { 1, 13, 17, 23, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 48, 49 }**F(-1) timedout fail** { }**F(-2) exception fail** { 6, 27 }**Mupad****A grade** { }**B grade** { 5, 7, 11, 12, 14, 22, 38, 39, 40, 41 }**C grade** { }**F normal fail** { }**F(-1) timedout fail** { 1, 2, 3, 4, 6, 8, 9, 10, 13, 15, 16, 17, 18, 19, 20, 21, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44, 45, 46, 47, 48, 49, 50 }**F(-2) exception fail** { }**Sympy****A grade** { 10, 11, 12 }**B grade** { }**C grade** { 2, 3, 4, 5, 7, 8, 9, 14, 15, 16 }**F normal fail** { 1, 6, 13, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 42, 43, 44, 45, 46, 47, 48, 49, 50 }**F(-1) timedout fail** { 39, 40, 41 }**F(-2) exception fail** { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	62	62	56	0	0	0	0	0	0
N.S.	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.070	0.029	0.000	0.000	0.000	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	40	40	66	32	119	152	0
N.S.	1	1.00	0.69	0.69	1.14	0.55	2.05	2.62	0.00
time (sec)	N/A	0.016	0.021	0.230	0.182	0.263	61.845	0.295	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	35	35	52	27	90	116	0
N.S.	1	1.00	0.74	0.74	1.11	0.57	1.91	2.47	0.00
time (sec)	N/A	0.013	0.017	0.212	0.187	0.253	18.696	0.292	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	28	28	38	20	61	80	0
N.S.	1	1.00	0.78	0.78	1.06	0.56	1.69	2.22	0.00
time (sec)	N/A	0.009	0.014	0.161	0.182	0.267	5.586	0.281	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	18	18	18	22	21	14	29	41	21
N.S.	1	1.00	1.00	1.22	1.17	0.78	1.61	2.28	1.17
time (sec)	N/A	0.003	0.003	0.214	0.181	0.261	1.954	0.288	1.357

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	56	54	63	0	0	0	0	0
N.S.	1	1.00	0.96	1.12	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.062	0.018	0.651	0.000	0.000	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	32	44	51	19	75	30	28
N.S.	1	1.00	0.84	1.16	1.34	0.50	1.97	0.79	0.74
time (sec)	N/A	0.013	0.017	0.252	0.263	0.264	12.657	0.280	0.872

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	C	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	55	57	80	29	144	44	0
N.S.	1	1.00	1.02	1.06	1.48	0.54	2.67	0.81	0.00
time (sec)	N/A	0.016	0.029	0.223	0.265	0.263	41.805	0.273	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	C	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	68	68	45	67	106	35	180	58	0
N.S.	1	1.00	0.66	0.99	1.56	0.51	2.65	0.85	0.00
time (sec)	N/A	0.017	0.041	0.201	0.270	0.275	127.916	0.283	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	56	42	56	54	39	51	47	0
N.S.	1	1.00	0.75	1.00	0.96	0.70	0.91	0.84	0.00
time (sec)	N/A	0.032	0.025	1.369	0.268	0.259	0.169	0.265	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	44	67	46	38	41	39	38
N.S.	1	1.00	0.94	1.43	0.98	0.81	0.87	0.83	0.81
time (sec)	N/A	0.017	0.017	1.369	0.263	0.258	0.142	0.264	0.734

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	26	28	24	27	22	28	24
N.S.	1	1.00	1.00	1.08	0.92	1.04	0.85	1.08	0.92
time (sec)	N/A	0.007	0.008	1.352	0.180	0.270	0.128	0.263	0.115

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	59	76	0	0	0	0	0
N.S.	1	1.00	1.00	1.29	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.051	0.016	1.773	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	A	B	C	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	93	30	52	107	29	61	29
N.S.	1	1.00	3.00	0.97	1.68	3.45	0.94	1.97	0.94
time (sec)	N/A	0.023	0.093	0.063	0.181	0.266	1.072	0.267	0.706

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	C	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	36	33	32	39	53	61	0
N.S.	1	1.00	0.95	0.87	0.84	1.03	1.39	1.61	0.00
time (sec)	N/A	0.018	0.016	1.352	0.261	0.254	0.619	0.292	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	C	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	60	60	69	54	64	142	100	80	0
N.S.	1	1.00	1.15	0.90	1.07	2.37	1.67	1.33	0.00
time (sec)	N/A	0.031	0.031	1.329	0.264	0.275	1.399	0.270	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	60	93	0	0	0	0	0
N.S.	1	1.00	0.87	1.35	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.076	0.065	1.443	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	197	197	173	329	0	152	0	409	0
N.S.	1	1.00	0.88	1.67	0.00	0.77	0.00	2.08	0.00
time (sec)	N/A	0.176	0.115	0.389	0.000	0.291	0.000	0.311	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	155	155	150	249	0	130	0	299	0
N.S.	1	1.00	0.97	1.61	0.00	0.84	0.00	1.93	0.00
time (sec)	N/A	0.107	0.181	0.310	0.000	0.298	0.000	0.306	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	116	116	131	190	0	117	0	204	0
N.S.	1	1.00	1.13	1.64	0.00	1.01	0.00	1.76	0.00
time (sec)	N/A	0.070	0.132	0.301	0.000	0.271	0.000	0.312	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	110	108	0	104	0	133	0
N.S.	1	1.00	1.41	1.38	0.00	1.33	0.00	1.71	0.00
time (sec)	N/A	0.040	0.084	0.307	0.000	0.266	0.000	0.309	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	B	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	37	37	468	45	55	73	0	82	35
N.S.	1	1.00	12.65	1.22	1.49	1.97	0.00	2.22	0.95
time (sec)	N/A	0.017	1.992	0.074	0.179	0.303	0.000	0.296	1.044

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	200	200	284	374	0	0	0	0	0
N.S.	1	1.00	1.42	1.87	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.231	0.238	1.642	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	B	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	70	70	112	126	0	281	0	94	0
N.S.	1	1.00	1.60	1.80	0.00	4.01	0.00	1.34	0.00
time (sec)	N/A	0.076	0.209	0.721	0.000	0.317	0.000	0.307	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	125	125	198	319	0	427	0	216	0
N.S.	1	1.00	1.58	2.55	0.00	3.42	0.00	1.73	0.00
time (sec)	N/A	0.156	0.711	0.704	0.000	0.302	0.000	0.322	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	B	F	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	181	181	241	528	0	548	0	451	0
N.S.	1	1.00	1.33	2.92	0.00	3.03	0.00	2.49	0.00
time (sec)	N/A	0.232	0.311	0.699	0.000	0.318	0.000	0.333	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	381	381	667	673	0	0	0	0	0
N.S.	1	1.00	1.75	1.77	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.228	8.394	1.435	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	No	Yes	TBD	TBD	TBD	TBD	TBD
size	288	288	473	498	0	0	0	0	0
N.S.	1	1.00	1.64	1.73	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.180	4.591	1.514	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	154	154	144	225	0	0	0	0	0
N.S.	1	1.00	0.94	1.46	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.097	0.102	0.949	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	111	162	0	0	0	0	0
N.S.	1	1.00	1.18	1.72	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.050	0.086	0.539	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	310	310	813	0	0	0	0	0	0
N.S.	1	1.00	2.62	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.395	1.815	0.000	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	244	244	686	336	0	0	0	0	0
N.S.	1	1.00	2.81	1.38	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.325	1.728	1.645	0.000	0.000	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	494	494	446	716	0	0	0	0	0
N.S.	1	1.00	0.90	1.45	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.306	0.304	1.256	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	278	278	257	379	0	0	0	0	0
N.S.	1	1.00	0.92	1.36	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.193	0.113	1.451	0.000	0.000	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	154	154	160	0	0	0	0	0	0
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.077	0.077	0.000	0.000	0.000	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	F	F	F	F	F	F(-1)
verified	N/A	Yes	No	N/A	TBD	TBD	TBD	TBD	TBD
size	430	430	1058	0	0	0	0	0	0
N.S.	1	1.00	2.46	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.482	2.916	0.000	0.000	0.000	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F(-1)	F	F	F	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	362	362	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.481	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	58	58	516	64	71	96	0	100	52
N.S.	1	1.00	8.90	1.10	1.22	1.66	0.00	1.72	0.90
time (sec)	N/A	0.057	2.310	0.327	0.178	0.301	0.000	0.390	1.209

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	91	91	107	0	0	0	0	0	0
N.S.	1	1.00	1.18	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.086	0.161	0.000	0.000	0.000	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	91	91	54	0	0	0	0	0	0
N.S.	1	1.00	0.59	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.069	0.046	0.000	0.000	0.000	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	45	45	79	0	0	0	0	0	0
N.S.	1	1.00	1.76	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.053	0.040	0.000	0.000	0.000	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	A	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	39	39	34	0	0	23	0	43	0
N.S.	1	1.00	0.87	0.00	0.00	0.59	0.00	1.10	0.00
time (sec)	N/A	0.023	0.030	0.000	0.000	0.415	0.000	0.288	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	41	41	30	0	0	30	0	0	0
N.S.	1	1.00	0.73	0.00	0.00	0.73	0.00	0.00	0.00
time (sec)	N/A	0.031	0.035	0.000	0.000	0.282	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	84	84	54	0	0	40	0	0	0
N.S.	1	1.00	0.64	0.00	0.00	0.48	0.00	0.00	0.00
time (sec)	N/A	0.051	0.113	0.000	0.000	0.282	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	59	99	0	0	0	115	0
N.S.	1	1.00	0.86	1.43	0.00	0.00	0.00	1.67	0.00
time (sec)	N/A	0.071	0.053	1.037	0.000	0.000	0.000	0.502	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. The column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [34] had the largest ratio of [1.1999999999999996]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	7	6	1.00	10	0.600
2	A	4	3	1.00	10	0.300
3	A	4	3	1.00	10	0.300
4	A	4	3	1.00	8	0.375
5	A	3	3	1.00	6	0.500
6	A	7	6	1.00	10	0.600
7	A	5	5	1.00	10	0.500
8	A	6	5	1.00	10	0.500
9	A	7	5	1.00	10	0.500
10	A	5	4	1.00	10	0.400
11	A	4	4	1.00	8	0.500
12	A	3	3	1.00	6	0.500
13	A	6	6	1.00	10	0.600
14	A	5	5	1.00	10	0.500
15	A	3	3	1.00	10	0.300
16	A	6	6	1.00	10	0.600
17	A	7	6	1.00	10	0.600
18	A	9	8	1.00	10	0.800
19	A	8	7	1.00	10	0.700
20	A	7	6	1.00	10	0.600
21	A	6	6	1.00	8	0.750
22	A	5	5	1.00	6	0.833
23	A	14	8	1.00	10	0.800
24	A	5	5	1.00	10	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	7	7	1.00	10	0.700
26	A	8	8	1.00	10	0.800
27	A	20	9	1.00	12	0.750
28	A	17	9	1.00	12	0.750
29	A	11	8	1.00	10	0.800
30	A	8	6	1.00	8	0.750
31	A	17	9	1.00	12	0.750
32	A	12	8	1.00	12	0.667
33	A	25	14	1.00	12	1.167
34	A	16	12	1.00	10	1.200
35	A	10	7	1.00	8	0.875
36	A	20	10	1.00	12	0.833
37	A	14	9	1.00	12	0.750
38	A	7	6	1.00	14	0.429
39	A	7	6	1.00	16	0.375
40	A	7	6	1.00	16	0.375
41	A	6	6	1.00	14	0.429
42	A	7	7	1.00	10	0.700
43	A	6	4	1.00	10	0.400
44	A	6	4	1.00	8	0.500
45	A	5	3	1.00	6	0.500
46	A	6	5	1.00	10	0.500
47	A	3	3	1.00	10	0.300
48	A	5	4	1.00	10	0.400
49	A	6	4	1.00	10	0.400
50	A	8	8	1.00	19	0.421

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\sec^{-1}(ax^5)}{x} dx$	41
3.2	$\int x^3 \sec^{-1}(\sqrt{x}) dx$	45
3.3	$\int x^2 \sec^{-1}(\sqrt{x}) dx$	50
3.4	$\int x \sec^{-1}(\sqrt{x}) dx$	55
3.5	$\int \sec^{-1}(\sqrt{x}) dx$	59
3.6	$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx$	63
3.7	$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx$	68
3.8	$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx$	73
3.9	$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$	78
3.10	$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx$	83
3.11	$\int x \sec^{-1}\left(\frac{a}{x}\right) dx$	87
3.12	$\int \sec^{-1}\left(\frac{a}{x}\right) dx$	92
3.13	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx$	96
3.14	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx$	101
3.15	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx$	106
3.16	$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx$	110
3.17	$\int \frac{\sec^{-1}(ax^n)}{x} dx$	115
3.18	$\int x^4 \sec^{-1}(a + bx) dx$	120
3.19	$\int x^3 \sec^{-1}(a + bx) dx$	127
3.20	$\int x^2 \sec^{-1}(a + bx) dx$	133
3.21	$\int x \sec^{-1}(a + bx) dx$	139
3.22	$\int \sec^{-1}(a + bx) dx$	144
3.23	$\int \frac{\sec^{-1}(a+bx)}{x} dx$	149
3.24	$\int \frac{\sec^{-1}(a+bx)}{x^2} dx$	156
3.25	$\int \frac{\sec^{-1}(a+bx)}{x^3} dx$	161

3.26	$\int \frac{\sec^{-1}(a+bx)}{x^4} dx$	167
3.27	$\int x^3 \sec^{-1}(a+bx)^2 dx$	174
3.28	$\int x^2 \sec^{-1}(a+bx)^2 dx$	183
3.29	$\int x \sec^{-1}(a+bx)^2 dx$	191
3.30	$\int \sec^{-1}(a+bx)^2 dx$	197
3.31	$\int \frac{\sec^{-1}(a+bx)^2}{x} dx$	202
3.32	$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx$	211
3.33	$\int x^2 \sec^{-1}(a+bx)^3 dx$	218
3.34	$\int x \sec^{-1}(a+bx)^3 dx$	232
3.35	$\int \sec^{-1}(a+bx)^3 dx$	241
3.36	$\int \frac{\sec^{-1}(a+bx)^3}{x} dx$	247
3.37	$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx$	260
3.38	$\int x(a+b \sec^{-1}(c+dx^2)) dx$	268
3.39	$\int x^2(a+b \sec^{-1}(c+dx^3)) dx$	273
3.40	$\int x^3(a+b \sec^{-1}(c+dx^4)) dx$	278
3.41	$\int x^{-1+n} \sec^{-1}(a+bx^n) dx$	283
3.42	$\int \sec^{-1}(ce^{a+bx}) dx$	288
3.43	$\int e^{\sec^{-1}(ax)} x^2 dx$	294
3.44	$\int e^{\sec^{-1}(ax)} x dx$	298
3.45	$\int e^{\sec^{-1}(ax)} dx$	302
3.46	$\int \frac{e^{\sec^{-1}(ax)}}{x} dx$	306
3.47	$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$	310
3.48	$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx$	314
3.49	$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$	318
3.50	$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$	322

3.1 $\int \frac{\sec^{-1}(ax^5)}{x} dx$

Optimal result	41
Rubi [A] (verified)	41
Mathematica [A] (verified)	43
Maple [F]	43
Fricas [F]	43
Sympy [F]	44
Maxima [F]	44
Giac [F]	44
Mupad [F(-1)]	44

Optimal result

Integrand size = 10, antiderivative size = 62

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right) + \frac{1}{10}i \text{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right)$$

[Out] 1/10*I*arcsec(a*x^5)^2-1/5*arcsec(a*x^5)*ln(1+(1/a/x^5+I*(1-1/a^2/x^10)^(1/2))^2)+1/10*I*polylog(2,-(1/a/x^5+I*(1-1/a^2/x^10)^(1/2))^2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5326, 4722, 3800, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \frac{1}{10}i \text{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right) + \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right)$$

[In] Int[ArcSec[a*x^5]/x,x]

[Out] (I/10)*ArcSec[a*x^5]^2 - (ArcSec[a*x^5]*Log[1 + E^((2*I)*ArcSec[a*x^5])])/5 + (I/10)*PolyLog[2, -E^((2*I)*ArcSec[a*x^5])]

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di

```
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4722

```
Int[((a_) + ArcCos[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5326

```
Int[((a_) + ArcSec[(c_)*(x_)]*(b_))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{5} \text{Subst} \left(\int \frac{\sec^{-1}(ax)}{x} dx, x, x^5 \right) \\
 &= - \left(\frac{1}{5} \text{Subst} \left(\int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx, x, \frac{1}{x^5} \right) \right) \\
 &= \frac{1}{5} \text{Subst} \left(\int x \tan(x) dx, x, \sec^{-1}(ax^5) \right) \\
 &= \frac{1}{10} i \sec^{-1}(ax^5)^2 - \frac{2}{5} i \text{Subst} \left(\int \frac{e^{2ix} x}{1 + e^{2ix}} dx, x, \sec^{-1}(ax^5) \right)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right) \\
&\quad + \frac{1}{5} \text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \sec^{-1}(ax^5)\right) \\
&= \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right) \\
&\quad - \frac{1}{10}i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \sec^{-1}(ax^5)}\right) \\
&= \frac{1}{10}i \sec^{-1}(ax^5)^2 - \frac{1}{5} \sec^{-1}(ax^5) \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right) + \frac{1}{10}i \text{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.90

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \frac{1}{10}i \left(\sec^{-1}(ax^5) \left(\sec^{-1}(ax^5) + 2i \log\left(1 + e^{2i \sec^{-1}(ax^5)}\right)\right) \right. \\
\left. + \text{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^5)}\right) \right)$$

[In] Integrate[ArcSec[a*x^5]/x,x]

[Out] (I/10)*(ArcSec[a*x^5]*(ArcSec[a*x^5] + (2*I)*Log[1 + E^((2*I)*ArcSec[a*x^5])]) + PolyLog[2, -E^((2*I)*ArcSec[a*x^5])])

Maple [F]

$$\int \frac{\text{arcsec}(ax^5)}{x} dx$$

[In] int(arcsec(a*x^5)/x,x)

[Out] int(arcsec(a*x^5)/x,x)

Fricas [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\text{arcsec}(ax^5)}{x} dx$$

[In] integrate(arcsec(a*x^5)/x,x, algorithm="fricas")

[Out] integral(arcsec(a*x^5)/x, x)

Sympy [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{asec}(ax^5)}{x} dx$$

[In] integrate(asec(a*x**5)/x,x)

[Out] Integral(asec(a*x**5)/x, x)

Maxima [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

[In] integrate(arcsec(a*x^5)/x,x, algorithm="maxima")

[Out] -5*a^2*integrate(sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1)*log(x)/(a^4*x^11 - a^2*x), x) - 5*I*a^2*integrate(log(x)/(a^4*x^11 - a^2*x), x) + arctan(sqrt(a*x^5 + 1)*sqrt(a*x^5 - 1))*log(x) - 1/2*I*log(a^2*x^10)*log(x) + 1/2*I*log(a*x^5 + 1)*log(x) + 1/2*I*log(-a*x^5 + 1)*log(x) + I*log(a)*log(x) + 5/2*I*log(x)^2 + 1/10*I*dilog(a*x^5) + 1/10*I*dilog(-a*x^5)

Giac [F]

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{arcsec}(ax^5)}{x} dx$$

[In] integrate(arcsec(a*x^5)/x,x, algorithm="giac")

[Out] integrate(arcsec(a*x^5)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(ax^5)}{x} dx = \int \frac{\operatorname{acos}\left(\frac{1}{ax^5}\right)}{x} dx$$

[In] int(acos(1/(a*x^5))/x,x)

[Out] int(acos(1/(a*x^5))/x, x)

3.2 $\int x^3 \sec^{-1}(\sqrt{x}) dx$

Optimal result	45
Rubi [A] (verified)	45
Mathematica [A] (verified)	46
Maple [A] (verified)	47
Fricas [A] (verification not implemented)	47
Sympy [C] (verification not implemented)	47
Maxima [A] (verification not implemented)	48
Giac [B] (verification not implemented)	48
Mupad [F(-1)]	49

Optimal result

Integrand size = 10, antiderivative size = 58

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{4}\sqrt{-1+x} - \frac{1}{4}(-1+x)^{3/2} - \frac{3}{20}(-1+x)^{5/2} - \frac{1}{28}(-1+x)^{7/2} + \frac{1}{4}x^4 \sec^{-1}(\sqrt{x})$$

[Out] $-1/4*(-1+x)^{(3/2)}-3/20*(-1+x)^{(5/2)}-1/28*(-1+x)^{(7/2)}+1/4*x^4*\text{arcsec}(x^{(1/2)})-1/4*(-1+x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5378, 12, 45}

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \frac{1}{4}x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{28}(x-1)^{7/2} - \frac{3}{20}(x-1)^{5/2} - \frac{1}{4}(x-1)^{3/2} - \frac{\sqrt{x-1}}{4}$$

[In] `Int[x^3*ArcSec[Sqrt[x]],x]`

[Out] $-1/4*\text{Sqrt}[-1+x] - (-1+x)^{(3/2)}/4 - (3*(-1+x)^{(5/2)})/20 - (-1+x)^{(7/2)}/28 + (x^4*\text{ArcSec}[\text{Sqrt}[x]])/4$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 5378

```
Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Sim
p[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m +
1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u
^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inverse
FunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Function
OfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{4}x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \frac{x^3}{2\sqrt{-1+x}} dx \\
&= \frac{1}{4}x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{8} \int \frac{x^3}{\sqrt{-1+x}} dx \\
&= \frac{1}{4}x^4 \sec^{-1}(\sqrt{x}) - \frac{1}{8} \int \left(\frac{1}{\sqrt{-1+x}} + 3\sqrt{-1+x} + 3(-1+x)^{3/2} + (-1+x)^{5/2} \right) dx \\
&= -\frac{1}{4}\sqrt{-1+x} - \frac{1}{4}(-1+x)^{3/2} - \frac{3}{20}(-1+x)^{5/2} - \frac{1}{28}(-1+x)^{7/2} + \frac{1}{4}x^4 \sec^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{140}\sqrt{-1+x}(16 + 8x + 6x^2 + 5x^3) + \frac{1}{4}x^4 \sec^{-1}(\sqrt{x})$$

```
[In] Integrate[x^3*ArcSec[Sqrt[x]], x]
```

```
[Out] -1/140*(Sqrt[-1 + x]*(16 + 8*x + 6*x^2 + 5*x^3)) + (x^4*ArcSec[Sqrt[x]])/4
```

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.69

method	result	size
parts	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{\sqrt{\frac{x-1}{x}} \sqrt{x} (5x^3+6x^2+8x+16)}{140}$	40
derivativedivides	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{(x-1)(5x^3+6x^2+8x+16)}{140\sqrt{\frac{x-1}{x}} \sqrt{x}}$	43
default	$\frac{x^4 \operatorname{arcsec}(\sqrt{x})}{4} - \frac{(x-1)(5x^3+6x^2+8x+16)}{140\sqrt{\frac{x-1}{x}} \sqrt{x}}$	43

[In] `int(x^3*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `1/4*x^4*arcsec(x^(1/2))-1/140*((x-1)/x)^(1/2)*x^(1/2)*(5*x^3+6*x^2+8*x+16)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.55

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \frac{1}{4} x^4 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{140} (5x^3 + 6x^2 + 8x + 16) \sqrt{x-1}$$

[In] `integrate(x^3*arcsec(x^(1/2)),x, algorithm="fricas")`

[Out] `1/4*x^4*arcsec(sqrt(x)) - 1/140*(5*x^3 + 6*x^2 + 8*x + 16)*sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 61.84 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.05

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \frac{x^4 \operatorname{asec}(\sqrt{x})}{4} - \frac{\begin{cases} \frac{2x^3\sqrt{x-1}}{7} + \frac{12x^2\sqrt{x-1}}{35} + \frac{16x\sqrt{x-1}}{35} + \frac{32\sqrt{x-1}}{35} & \text{for } |x| > 1 \\ \frac{2ix^3\sqrt{1-x}}{7} + \frac{12ix^2\sqrt{1-x}}{35} + \frac{16ix\sqrt{1-x}}{35} + \frac{32i\sqrt{1-x}}{35} & \text{otherwise} \end{cases}}{8}$$

[In] `integrate(x**3*asec(x**(1/2)),x)`

[Out] `x**4*asec(sqrt(x))/4 - Piecewise((2*x**3*sqrt(x - 1)/7 + 12*x**2*sqrt(x - 1)/35 + 16*x*sqrt(x - 1)/35 + 32*sqrt(x - 1)/35, Abs(x) > 1), (2*I*x**3*sqrt(1 - x)/7 + 12*I*x**2*sqrt(1 - x)/35 + 16*I*x*sqrt(1 - x)/35 + 32*I*sqrt(1 - x)/35, True))/8`

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.14

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{28} x^{\frac{7}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{7}{2}} - \frac{3}{20} x^{\frac{5}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{5}{2}} \\ + \frac{1}{4} x^4 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{4} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{4} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

[In] integrate(x^3*arcsec(x^(1/2)),x, algorithm="maxima")

[Out] -1/28*x^(7/2)*(-1/x + 1)^(7/2) - 3/20*x^(5/2)*(-1/x + 1)^(5/2) + 1/4*x^4*arcsec(sqrt(x)) - 1/4*x^(3/2)*(-1/x + 1)^(3/2) - 1/4*sqrt(x)*sqrt(-1/x + 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(38) = 76.

Time = 0.30 (sec) , antiderivative size = 152, normalized size of antiderivative = 2.62

$$\int x^3 \sec^{-1}(\sqrt{x}) dx \\ = -\frac{1}{3584} x^{\frac{7}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^7 - \frac{7}{2560} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^5 \\ + \frac{1}{4} x^4 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{7}{512} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^3 - \frac{35}{512} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1\right) \\ + \frac{1225 x^3 \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^6 + 245 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^4 + 49 x \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^2 + 5}{17920 x^{\frac{7}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^7}$$

[In] integrate(x^3*arcsec(x^(1/2)),x, algorithm="giac")

[Out] -1/3584*x^(7/2)*(sqrt(-1/x + 1) - 1)^7 - 7/2560*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 + 1/4*x^4*arccos(1/sqrt(x)) - 7/512*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 - 35/512*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/17920*(1225*x^3*(sqrt(-1/x + 1) - 1)^6 + 245*x^2*(sqrt(-1/x + 1) - 1)^4 + 49*x*(sqrt(-1/x + 1) - 1)^2 + 5)/(x^(7/2)*(sqrt(-1/x + 1) - 1)^7)

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(\sqrt{x}) dx = \int x^3 \arccos\left(\frac{1}{\sqrt{x}}\right) dx$$

```
[In] int(x^3*acos(1/x^(1/2)),x)
```

```
[Out] int(x^3*acos(1/x^(1/2)), x)
```

3.3 $\int x^2 \sec^{-1}(\sqrt{x}) dx$

Optimal result	50
Rubi [A] (verified)	50
Mathematica [A] (verified)	51
Maple [A] (verified)	51
Fricas [A] (verification not implemented)	52
Sympy [C] (verification not implemented)	52
Maxima [A] (verification not implemented)	53
Giac [B] (verification not implemented)	53
Mupad [F(-1)]	54

Optimal result

Integrand size = 10, antiderivative size = 47

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{3}\sqrt{-1+x} - \frac{2}{9}(-1+x)^{3/2} - \frac{1}{15}(-1+x)^{5/2} + \frac{1}{3}x^3 \sec^{-1}(\sqrt{x})$$

[Out] -2/9*(-1+x)^(3/2)-1/15*(-1+x)^(5/2)+1/3*x^3*arcsec(x^(1/2))-1/3*(-1+x)^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5378, 12, 45}

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \frac{1}{3}x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{15}(x-1)^{5/2} - \frac{2}{9}(x-1)^{3/2} - \frac{\sqrt{x-1}}{3}$$

[In] Int[x^2*ArcSec[Sqrt[x]],x]

[Out] -1/3*Sqrt[-1 + x] - (2*(-1 + x)^(3/2))/9 - (-1 + x)^(5/2)/15 + (x^3*ArcSec[Sqrt[x]])/3

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},

`x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 5378

```
Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_.))^(m_.), x_Symbol] := Sim
p[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m +
1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u
^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inverse
FunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Function
OfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3}x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{3} \int \frac{x^2}{2\sqrt{-1+x}} dx \\
 &= \frac{1}{3}x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{6} \int \frac{x^2}{\sqrt{-1+x}} dx \\
 &= \frac{1}{3}x^3 \sec^{-1}(\sqrt{x}) - \frac{1}{6} \int \left(\frac{1}{\sqrt{-1+x}} + 2\sqrt{-1+x} + (-1+x)^{3/2} \right) dx \\
 &= -\frac{1}{3}\sqrt{-1+x} - \frac{2}{9}(-1+x)^{3/2} - \frac{1}{15}(-1+x)^{5/2} + \frac{1}{3}x^3 \sec^{-1}(\sqrt{x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.74

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{45}\sqrt{-1+x}(8+4x+3x^2) + \frac{1}{3}x^3 \sec^{-1}(\sqrt{x})$$

[In] Integrate[x^2*ArcSec[Sqrt[x]],x]

[Out] -1/45*(Sqrt[-1 + x]*(8 + 4*x + 3*x^2)) + (x^3*ArcSec[Sqrt[x]])/3

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.74

method	result	size
parts	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{\sqrt{\frac{x-1}{x}} \sqrt{x} (3x^2+4x+8)}{45}$	35
derivativedivides	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{(x-1)(3x^2+4x+8)}{45\sqrt{\frac{x-1}{x}} \sqrt{x}}$	38
default	$\frac{x^3 \operatorname{arcsec}(\sqrt{x})}{3} - \frac{(x-1)(3x^2+4x+8)}{45\sqrt{\frac{x-1}{x}} \sqrt{x}}$	38

[In] `int(x^2*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] `1/3*x^3*arcsec(x^(1/2))-1/45*((x-1)/x)^(1/2)*x^(1/2)*(3*x^2+4*x+8)`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.57

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \frac{1}{3} x^3 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{45} (3x^2 + 4x + 8) \sqrt{x-1}$$

[In] `integrate(x^2*arcsec(x^(1/2)),x, algorithm="fricas")`

[Out] `1/3*x^3*arcsec(sqrt(x)) - 1/45*(3*x^2 + 4*x + 8)*sqrt(x - 1)`

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 18.70 (sec) , antiderivative size = 90, normalized size of antiderivative = 1.91

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \frac{x^3 \operatorname{asec}(\sqrt{x})}{3} - \frac{\begin{cases} \frac{2x^2\sqrt{x-1}}{5} + \frac{8x\sqrt{x-1}}{15} + \frac{16\sqrt{x-1}}{15} & \text{for } |x| > 1 \\ \frac{2ix^2\sqrt{1-x}}{5} + \frac{8ix\sqrt{1-x}}{15} + \frac{16i\sqrt{1-x}}{15} & \text{otherwise} \end{cases}}{6}$$

[In] `integrate(x**2*asec(x**(1/2)),x)`

[Out] `x**3*asec(sqrt(x))/3 - Piecewise((2*x**2*sqrt(x - 1)/5 + 8*x*sqrt(x - 1)/15 + 16*sqrt(x - 1)/15, Abs(x) > 1), (2*I*x**2*sqrt(1 - x)/5 + 8*I*x*sqrt(1 - x)/15 + 16*I*sqrt(1 - x)/15, True))/6`

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.11

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{15} x^{\frac{5}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{5}{2}} + \frac{1}{3} x^3 \operatorname{arcsec}(\sqrt{x}) - \frac{2}{9} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} - \frac{1}{3} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

[In] integrate(x^2*arcsec(x^(1/2)),x, algorithm="maxima")

[Out] -1/15*x^(5/2)*(-1/x + 1)^(5/2) + 1/3*x^3*arcsec(sqrt(x)) - 2/9*x^(3/2)*(-1/x + 1)^(3/2) - 1/3*sqrt(x)*sqrt(-1/x + 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(31) = 62.

Time = 0.29 (sec) , antiderivative size = 116, normalized size of antiderivative = 2.47

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = -\frac{1}{480} x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^5 - \frac{5}{288} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^3 + \frac{1}{3} x^3 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{5}{48} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1\right) + \frac{150 x^2 \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^4 + 25 x \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^2 + 3}{1440 x^{\frac{5}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1\right)^5}$$

[In] integrate(x^2*arcsec(x^(1/2)),x, algorithm="giac")

[Out] -1/480*x^(5/2)*(sqrt(-1/x + 1) - 1)^5 - 5/288*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 1/3*x^3*arccos(1/sqrt(x)) - 5/48*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/1440*(150*x^2*(sqrt(-1/x + 1) - 1)^4 + 25*x*(sqrt(-1/x + 1) - 1)^2 + 3)/(x^(5/2)*(sqrt(-1/x + 1) - 1)^5)

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(\sqrt{x}) dx = \int x^2 \arccos\left(\frac{1}{\sqrt{x}}\right) dx$$

```
[In] int(x^2*acos(1/x^(1/2)),x)
```

```
[Out] int(x^2*acos(1/x^(1/2)), x)
```

3.4 $\int x \sec^{-1}(\sqrt{x}) dx$

Optimal result	55
Rubi [A] (verified)	55
Mathematica [A] (verified)	56
Maple [A] (verified)	56
Fricas [A] (verification not implemented)	57
Sympy [C] (verification not implemented)	57
Maxima [A] (verification not implemented)	57
Giac [B] (verification not implemented)	58
Mupad [F(-1)]	58

Optimal result

Integrand size = 8, antiderivative size = 36

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{2}\sqrt{-1+x} - \frac{1}{6}(-1+x)^{3/2} + \frac{1}{2}x^2 \sec^{-1}(\sqrt{x})$$

[Out] $-1/6*(-1+x)^{(3/2)}+1/2*x^2*\text{arcsec}(x^{(1/2)})-1/2*(-1+x)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5378, 12, 45}

$$\int x \sec^{-1}(\sqrt{x}) dx = \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{6}(x-1)^{3/2} - \frac{\sqrt{x-1}}{2}$$

[In] `Int[x*ArcSec[Sqrt[x]],x]`

[Out] $-1/2*\text{Sqrt}[-1+x] - (-1+x)^{(3/2)}/6 + (x^2*\text{ArcSec}[\text{Sqrt}[x]])/2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]`

Rule 45

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 5378

```
Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Sim
p[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m +
1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u
^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && Inverse
FunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !Function
OfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{2} \int \frac{x}{2\sqrt{-1+x}} dx \\
&= \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \frac{x}{\sqrt{-1+x}} dx \\
&= \frac{1}{2}x^2 \sec^{-1}(\sqrt{x}) - \frac{1}{4} \int \left(\frac{1}{\sqrt{-1+x}} + \sqrt{-1+x} \right) dx \\
&= -\frac{1}{2}\sqrt{-1+x} - \frac{1}{6}(-1+x)^{3/2} + \frac{1}{2}x^2 \sec^{-1}(\sqrt{x})
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{6}\sqrt{-1+x}(2+x) + \frac{1}{2}x^2 \sec^{-1}(\sqrt{x})$$

[In] Integrate[x*ArcSec[Sqrt[x]],x]

[Out] -1/6*(Sqrt[-1 + x]*(2 + x)) + (x^2*ArcSec[Sqrt[x]])/2

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.78

method	result	size
parts	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{\sqrt{\frac{x-1}{x}} \sqrt{x} (2+x)}{6}$	28
derivativedivides	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{(x-1)(2+x)}{6\sqrt{\frac{x-1}{x}} \sqrt{x}}$	31
default	$\frac{x^2 \operatorname{arcsec}(\sqrt{x})}{2} - \frac{(x-1)(2+x)}{6\sqrt{\frac{x-1}{x}} \sqrt{x}}$	31

[In] `int(x*arcsec(x^(1/2)),x,method=_RETURNVERBOSE)`

[Out] $1/2*x^2*arcsec(x^(1/2))-1/6*((x-1)/x)^(1/2)*x^(1/2)*(2+x)$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 0.56

$$\int x \sec^{-1}(\sqrt{x}) dx = \frac{1}{2} x^2 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{6} (x+2) \sqrt{x-1}$$

[In] `integrate(x*arcsec(x^(1/2)),x, algorithm="fricas")`

[Out] $1/2*x^2*arcsec(sqrt(x)) - 1/6*(x + 2)*sqrt(x - 1)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 5.59 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.69

$$\int x \sec^{-1}(\sqrt{x}) dx = \frac{x^2 \operatorname{asec}(\sqrt{x})}{2} - \frac{\begin{cases} \frac{2x\sqrt{x-1}}{3} + \frac{4\sqrt{x-1}}{3} & \text{for } |x| > 1 \\ \frac{2ix\sqrt{1-x}}{3} + \frac{4i\sqrt{1-x}}{3} & \text{otherwise} \end{cases}}{4}$$

[In] `integrate(x*asec(x**(1/2)),x)`

[Out] $x**2*asec(sqrt(x))/2 - \operatorname{Piecewise}((2*x*sqrt(x - 1)/3 + 4*sqrt(x - 1)/3, \operatorname{Abs}(x) > 1), (2*I*x*sqrt(1 - x)/3 + 4*I*sqrt(1 - x)/3, \operatorname{True}))/4$

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.06

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{6} x^{\frac{3}{2}} \left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + \frac{1}{2} x^2 \operatorname{arcsec}(\sqrt{x}) - \frac{1}{2} \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

[In] `integrate(x*arcsec(x^(1/2)),x, algorithm="maxima")`

[Out] $-1/6*x^(3/2)*(-1/x + 1)^(3/2) + 1/2*x^2*arcsec(sqrt(x)) - 1/2*sqrt(x)*sqrt(-1/x + 1)$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(24) = 48.

Time = 0.28 (sec) , antiderivative size = 80, normalized size of antiderivative = 2.22

$$\int x \sec^{-1}(\sqrt{x}) dx = -\frac{1}{48} x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3 + \frac{1}{2} x^2 \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{3}{16} \sqrt{x} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right) + \frac{9x \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^2 + 1}{48 x^{\frac{3}{2}} \left(\sqrt{-\frac{1}{x} + 1} - 1 \right)^3}$$

[In] integrate(x*arcsec(x^(1/2)),x, algorithm="giac")

[Out] -1/48*x^(3/2)*(sqrt(-1/x + 1) - 1)^3 + 1/2*x^2*arccos(1/sqrt(x)) - 3/16*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/48*(9*x*(sqrt(-1/x + 1) - 1)^2 + 1)/(x^(3/2)*(sqrt(-1/x + 1) - 1)^3)

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(\sqrt{x}) dx = \int x \arccos\left(\frac{1}{\sqrt{x}}\right) dx$$

[In] int(x*acos(1/x^(1/2)),x)

[Out] int(x*acos(1/x^(1/2)), x)

3.5 $\int \sec^{-1}(\sqrt{x}) dx$

Optimal result	59
Rubi [A] (verified)	59
Mathematica [A] (verified)	60
Maple [A] (verified)	60
Fricas [A] (verification not implemented)	61
Sympy [C] (verification not implemented)	61
Maxima [A] (verification not implemented)	61
Giac [B] (verification not implemented)	62
Mupad [B] (verification not implemented)	62

Optimal result

Integrand size = 6, antiderivative size = 18

$$\int \sec^{-1}(\sqrt{x}) dx = -\sqrt{-1+x} + x \sec^{-1}(\sqrt{x})$$

[Out] $x \cdot \text{arcsec}(x^{1/2}) - (-1+x)^{1/2}$

Rubi [A] (verified)

Time = 0.00 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5376, 12, 32}

$$\int \sec^{-1}(\sqrt{x}) dx = x \sec^{-1}(\sqrt{x}) - \sqrt{x-1}$$

[In] `Int[ArcSec[Sqrt[x]],x]`

[Out] `-Sqrt[-1 + x] + x*ArcSec[Sqrt[x]]`

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 32

`Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]`

Rule 5376

```
Int[ArcSec[u_], x_Symbol] := Simp[x*ArcSec[u], x] - Dist[u/Sqrt[u^2], Int[S
implifyIntegrand[x*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; InverseFunci
onFreeQ[u, x] && !FunctionOfExponentialQ[u, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= x \sec^{-1}(\sqrt{x}) - \int \frac{1}{2\sqrt{-1+x}} dx \\ &= x \sec^{-1}(\sqrt{x}) - \frac{1}{2} \int \frac{1}{\sqrt{-1+x}} dx \\ &= -\sqrt{-1+x} + x \sec^{-1}(\sqrt{x}) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \sec^{-1}(\sqrt{x}) dx = -\sqrt{-1+x} + x \sec^{-1}(\sqrt{x})$$

```
[In] Integrate[ArcSec[Sqrt[x]], x]
```

```
[Out] -Sqrt[-1 + x] + x*ArcSec[Sqrt[x]]
```

Maple [A] (verified)

Time = 0.21 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.22

method	result	size
parts	$x \operatorname{arcsec}(\sqrt{x}) - \sqrt{\frac{x-1}{x}} \sqrt{x}$	22
derivativedivides	$x \operatorname{arcsec}(\sqrt{x}) - \frac{x-1}{\sqrt{\frac{x-1}{x}} \sqrt{x}}$	25
default	$x \operatorname{arcsec}(\sqrt{x}) - \frac{x-1}{\sqrt{\frac{x-1}{x}} \sqrt{x}}$	25

```
[In] int(arcsec(x^(1/2)), x, method=_RETURNVERBOSE)
```

```
[Out] x*arcsec(x^(1/2))-((x-1)/x)^(1/2)*x^(1/2)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.78

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{arcsec}(\sqrt{x}) - \sqrt{x-1}$$

[In] integrate(arcsec(x^(1/2)),x, algorithm="fricas")

[Out] x*arcsec(sqrt(x)) - sqrt(x - 1)

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.95 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.61

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{asec}(\sqrt{x}) - \frac{\begin{cases} 2\sqrt{x-1} & \text{for } |x| > 1 \\ 2i\sqrt{1-x} & \text{otherwise} \end{cases}}{2}$$

[In] integrate(asec(x**(1/2)),x)

[Out] x*asec(sqrt(x)) - Piecewise((2*sqrt(x - 1), Abs(x) > 1), (2*I*sqrt(1 - x), True))/2

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.17

$$\int \sec^{-1}(\sqrt{x}) dx = x \operatorname{arcsec}(\sqrt{x}) - \sqrt{x} \sqrt{-\frac{1}{x} + 1}$$

[In] integrate(arcsec(x^(1/2)),x, algorithm="maxima")

[Out] x*arcsec(sqrt(x)) - sqrt(x)*sqrt(-1/x + 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 41 vs. 2(14) = 28.

Time = 0.29 (sec) , antiderivative size = 41, normalized size of antiderivative = 2.28

$$\int \sec^{-1}(\sqrt{x}) dx = x \arccos\left(\frac{1}{\sqrt{x}}\right) - \frac{1}{2}\sqrt{x}\left(\sqrt{-\frac{1}{x} + 1} - 1\right) + \frac{1}{2\sqrt{x}\left(\sqrt{-\frac{1}{x} + 1} - 1\right)}$$

[In] integrate(arcsec(x^(1/2)),x, algorithm="giac")

[Out] x*arccos(1/sqrt(x)) - 1/2*sqrt(x)*(sqrt(-1/x + 1) - 1) + 1/2/(sqrt(x)*(sqrt(-1/x + 1) - 1))

Mupad [B] (verification not implemented)

Time = 1.36 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.17

$$\int \sec^{-1}(\sqrt{x}) dx = x \arccos\left(\frac{1}{\sqrt{x}}\right) - \sqrt{x}\sqrt{1 - \frac{1}{x}}$$

[In] int(acos(1/x^(1/2)),x)

[Out] x*acos(1/x^(1/2)) - x^(1/2)*(1 - 1/x)^(1/2)

3.6 $\int \frac{\sec^{-1}(\sqrt{x})}{x} dx$

Optimal result	63
Rubi [A] (verified)	63
Mathematica [A] (verified)	65
Maple [A] (verified)	65
Fricas [F]	66
Sympy [F]	66
Maxima [F]	66
Giac [F(-2)]	66
Mupad [F(-1)]	67

Optimal result

Integrand size = 10, antiderivative size = 56

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = i \sec^{-1}(\sqrt{x})^2 - 2 \sec^{-1}(\sqrt{x}) \log\left(1 + e^{2i \sec^{-1}(\sqrt{x})}\right) + i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(\sqrt{x})}\right)$$

[Out] $I*\operatorname{arcsec}(x^{(1/2)})^2 - 2*\operatorname{arcsec}(x^{(1/2)})*\ln(1 + (1/x^{(1/2)} + I*(1-1/x)^{(1/2)})^2) + I*\operatorname{polylog}(2, -(1/x^{(1/2)} + I*(1-1/x)^{(1/2)})^2)$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5326, 4722, 3800, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(\sqrt{x})}\right) + i \sec^{-1}(\sqrt{x})^2 - 2 \sec^{-1}(\sqrt{x}) \log\left(1 + e^{2i \sec^{-1}(\sqrt{x})}\right)$$

[In] $\operatorname{Int}[\operatorname{ArcSec}[\operatorname{Sqrt}[x]]/x, x]$

[Out] $I*\operatorname{ArcSec}[\operatorname{Sqrt}[x]]^2 - 2*\operatorname{ArcSec}[\operatorname{Sqrt}[x]]*\operatorname{Log}[1 + E^{((2*I)*\operatorname{ArcSec}[\operatorname{Sqrt}[x]])}] + I*\operatorname{PolyLog}[2, -E^{((2*I)*\operatorname{ArcSec}[\operatorname{Sqrt}[x]])}]$

Rule 2221

$\operatorname{Int}[(((F_.)^{((g_.)*((e_.) + (f_.)*(x_)))})^{(n_.)}*((c_.) + (d_.)*(x_))^{(m_.)})/((a_.) + (b_.)*((F_.)^{((g_.)*((e_.) + (f_.)*(x_)))})^{(n_.)}), x_Symbol] \rightarrow \operatorname{Simp}[\frac{(c + d*x)^m}{(b*f*g*n*\operatorname{Log}[F])}*\operatorname{Log}[1 + b*((F^{(g*(e + f*x)))^n/a}], x] - \operatorname{Di}$

```
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4722

```
Int[((a_) + ArcCos[(c_)*(x_)]*(b_))^(n_)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]
```

Rule 5326

```
Int[((a_) + ArcSec[(c_)*(x_)]*(b_))/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= 2\text{Subst}\left(\int \frac{\sec^{-1}(x)}{x} dx, x, \sqrt{x}\right) \\
 &= -\left(2\text{Subst}\left(\int \frac{\arccos(x)}{x} dx, x, \frac{1}{\sqrt{x}}\right)\right) \\
 &= 2\text{Subst}\left(\int x \tan(x) dx, x, \arccos\left(\frac{1}{\sqrt{x}}\right)\right) \\
 &= i \arccos\left(\frac{1}{\sqrt{x}}\right)^2 - 4i\text{Subst}\left(\int \frac{e^{2ix}x}{1 + e^{2ix}} dx, x, \arccos\left(\frac{1}{\sqrt{x}}\right)\right)
 \end{aligned}$$

$$\begin{aligned}
&= i \arccos\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad + 2 \text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \arccos\left(\frac{1}{\sqrt{x}}\right)\right) \\
&= i \arccos\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&\quad - i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) \\
&= i \arccos\left(\frac{1}{\sqrt{x}}\right)^2 - 2 \arccos\left(\frac{1}{\sqrt{x}}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right) + i \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{1}{\sqrt{x}}\right)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = i \left(\sec^{-1}(\sqrt{x}) \left(\sec^{-1}(\sqrt{x}) + 2i \log\left(1 + e^{2i \sec^{-1}(\sqrt{x})}\right) \right) \right. \\
\left. + \text{PolyLog}\left(2, -e^{2i \sec^{-1}(\sqrt{x})}\right) \right)$$

[In] Integrate[ArcSec[Sqrt[x]]/x,x]

[Out] I*(ArcSec[Sqrt[x]]*(ArcSec[Sqrt[x]] + (2*I)*Log[1 + E^((2*I)*ArcSec[Sqrt[x]])]) + PolyLog[2, -E^((2*I)*ArcSec[Sqrt[x]])])

Maple [A] (verified)

Time = 0.65 (sec) , antiderivative size = 63, normalized size of antiderivative = 1.12

method	result
derivativedivides	$i \operatorname{arcsec}(\sqrt{x})^2 - 2 \operatorname{arcsec}(\sqrt{x}) \ln\left(1 + \left(\frac{1}{\sqrt{x}} + i\sqrt{1 - \frac{1}{x}}\right)^2\right) + i \operatorname{polylog}\left(2, -\left(\frac{1}{\sqrt{x}} + i\sqrt{1 - \frac{1}{x}}\right)^2\right)$
default	$i \operatorname{arcsec}(\sqrt{x})^2 - 2 \operatorname{arcsec}(\sqrt{x}) \ln\left(1 + \left(\frac{1}{\sqrt{x}} + i\sqrt{1 - \frac{1}{x}}\right)^2\right) + i \operatorname{polylog}\left(2, -\left(\frac{1}{\sqrt{x}} + i\sqrt{1 - \frac{1}{x}}\right)^2\right)$

[In] int(arcsec(x^(1/2))/x,x,method=_RETURNVERBOSE)

[Out] I*arcsec(x^(1/2))^2-2*arcsec(x^(1/2))*ln(1+(1/x^(1/2)+I*(1-1/x)^(1/2))^2)+I*polylog(2,-(1/x^(1/2)+I*(1-1/x)^(1/2))^2)

Fricas [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsec}(\sqrt{x})}{x} dx$$

```
[In] integrate(arcsec(x^(1/2))/x,x, algorithm="fricas")
```

```
[Out] integral(arcsec(sqrt(x))/x, x)
```

Sympy [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{asec}(\sqrt{x})}{x} dx$$

```
[In] integrate(asec(x**(1/2))/x,x)
```

```
[Out] Integral(asec(sqrt(x))/x, x)
```

Maxima [F]

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\operatorname{arcsec}(\sqrt{x})}{x} dx$$

```
[In] integrate(arcsec(x^(1/2))/x,x, algorithm="maxima")
```

```
[Out] integrate(arcsec(sqrt(x))/x, x)
```

Giac [F(-2)]

Exception generated.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \text{Exception raised: NotImplementedError}$$

```
[In] integrate(arcsec(x^(1/2))/x,x, algorithm="giac")
```

```
[Out] Exception raised: NotImplementedError >> unable to parse Giac output: Inval  
id series expansion: non tractable function acos at +infinity
```

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x} dx$$

```
[In] int(acos(1/x^(1/2))/x,x)
```

```
[Out] int(acos(1/x^(1/2))/x, x)
```

3.7 $\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx$

Optimal result	68
Rubi [A] (verified)	68
Mathematica [A] (verified)	70
Maple [A] (verified)	70
Fricas [A] (verification not implemented)	70
Sympy [C] (verification not implemented)	71
Maxima [A] (verification not implemented)	71
Giac [A] (verification not implemented)	71
Mupad [B] (verification not implemented)	72

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1+x}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{2} \arctan(\sqrt{-1+x})$$

[Out] $-\text{arcsec}(x^{(1/2)})/x+1/2*\arctan((-1+x)^{(1/2)})+1/2*(-1+x)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5378, 12, 44, 65, 209}

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{1}{2} \arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x}$$

[In] $\text{Int}[\text{ArcSec}[\text{Sqrt}[x]]/x^2, x]$

[Out] $\text{Sqrt}[-1 + x]/(2*x) - \text{ArcSec}[\text{Sqrt}[x]]/x + \text{ArcTan}[\text{Sqrt}[-1 + x]]/2$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{Match}[\text{Q}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]]$

Rule 44

$\text{Int}[((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \text{Dist}[d*((m + n + 2) / ((b*c - a*d)*(m + 1))), \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x], x$

] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5378

Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sec^{-1}(\sqrt{x})}{x} + \int \frac{1}{2\sqrt{-1+xx^2}} dx \\
 &= -\frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{2} \int \frac{1}{\sqrt{-1+xx^2}} dx \\
 &= \frac{\sqrt{-1+x}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{4} \int \frac{1}{\sqrt{-1+xx}} dx \\
 &= \frac{\sqrt{-1+x}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{2} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x}\right) \\
 &= \frac{\sqrt{-1+x}}{2x} - \frac{\sec^{-1}(\sqrt{x})}{x} + \frac{1}{2} \arctan(\sqrt{-1+x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-1+x} - 2\sec^{-1}(\sqrt{x}) - x \arcsin\left(\frac{1}{\sqrt{x}}\right)}{2x}$$

[In] Integrate[ArcSec[Sqrt[x]]/x^2,x]

[Out] (Sqrt[-1 + x] - 2*ArcSec[Sqrt[x]] - x*ArcSin[1/Sqrt[x]])/(2*x)

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.16

method	result	size
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} + \frac{\sqrt{\frac{x-1}{x}} (\arctan(\sqrt{x-1})x + \sqrt{x-1})}{2\sqrt{x}\sqrt{x-1}}$	44
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} - \frac{\sqrt{x-1} \left(\arctan\left(\frac{1}{\sqrt{x-1}}\right)x - \sqrt{x-1} \right)}{2\sqrt{\frac{x-1}{x}} x^{\frac{3}{2}}}$	46
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{x} - \frac{\sqrt{x-1} \left(\arctan\left(\frac{1}{\sqrt{x-1}}\right)x - \sqrt{x-1} \right)}{2\sqrt{\frac{x-1}{x}} x^{\frac{3}{2}}}$	46

[In] int(arcsec(x^(1/2))/x^2,x,method=_RETURNVERBOSE)

[Out] -arcsec(x^(1/2))/x+1/2*((x-1)/x)^(1/2)/x^(1/2)*(arctan((x-1)^(1/2))*x+(x-1)^(1/2))/(x-1)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.50

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{(x-2) \operatorname{arcsec}(\sqrt{x}) + \sqrt{x-1}}{2x}$$

[In] integrate(arcsec(x^(1/2))/x^2,x, algorithm="fricas")

[Out] 1/2*((x - 2)*arcsec(sqrt(x)) + sqrt(x - 1))/x

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 12.66 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.97

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\begin{cases} i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right) - \frac{i}{\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{i}{x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\operatorname{asin}\left(\frac{1}{\sqrt{x}}\right) + \frac{\sqrt{1-\frac{1}{x}}}{\sqrt{x}} & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{asec}(\sqrt{x})}{x}$$

[In] integrate(asec(x**(1/2))/x**2,x)

[Out] Piecewise((I*acosh(1/sqrt(x)) - I/(sqrt(x)*sqrt(-1 + 1/x)) + I/(x**(3/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-asin(1/sqrt(x)) + sqrt(1 - 1/x)/sqrt(x), True))/2 - asecc(sqrt(x))/x

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.34

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = -\frac{\sqrt{x}\sqrt{-\frac{1}{x}+1}}{2\left(x\left(\frac{1}{x}-1\right)-1\right)} - \frac{\operatorname{arcsec}(\sqrt{x})}{x} + \frac{1}{2} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x}+1}\right)$$

[In] integrate(arcsec(x^(1/2))/x^2,x, algorithm="maxima")

[Out] -1/2*sqrt(x)*sqrt(-1/x + 1)/(x*(1/x - 1) - 1) - arcsec(sqrt(x))/x + 1/2*arctan(sqrt(x)*sqrt(-1/x + 1))

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.79

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{-\frac{1}{x}+1}}{2\sqrt{x}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x} + \frac{1}{2} \arccos\left(\frac{1}{\sqrt{x}}\right)$$

[In] integrate(arcsec(x^(1/2))/x^2,x, algorithm="giac")

[Out] 1/2*sqrt(-1/x + 1)/sqrt(x) - arccos(1/sqrt(x))/x + 1/2*arccos(1/sqrt(x))

Mupad [B] (verification not implemented)

Time = 0.87 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.74

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^2} dx = \frac{\sqrt{1 - \frac{1}{x}}}{2\sqrt{x}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right) \left(\frac{2}{x} - 1\right)}{2}$$

[In] int(acos(1/x^(1/2))/x^2,x)

[Out] (1 - 1/x)^(1/2)/(2*x^(1/2)) - (acos(1/x^(1/2))*(2/x - 1))/2

3.8 $\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx$

Optimal result	73
Rubi [A] (verified)	73
Mathematica [A] (verified)	75
Maple [A] (verified)	75
Fricas [A] (verification not implemented)	75
Sympy [C] (verification not implemented)	76
Maxima [B] (verification not implemented)	76
Giac [A] (verification not implemented)	77
Mupad [F(-1)]	77

Optimal result

Integrand size = 10, antiderivative size = 54

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{\sqrt{-1+x}}{8x^2} + \frac{3\sqrt{-1+x}}{16x} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \arctan(\sqrt{-1+x})$$

[Out] $-1/2*\text{arcsec}(x^{(1/2)})/x^2+3/16*\arctan((-1+x)^{(1/2)})+1/8*(-1+x)^{(1/2)}/x^2+3/16*(-1+x)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5378, 12, 44, 65, 209}

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{3}{16} \arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{8x^2} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3\sqrt{x-1}}{16x}$$

[In] Int[ArcSec[Sqrt[x]]/x^3,x]

[Out] $\text{Sqrt}[-1+x]/(8*x^2) + (3*\text{Sqrt}[-1+x])/(16*x) - \text{ArcSec}[\text{Sqrt}[x]]/(2*x^2) + (3*\text{ArcTan}[\text{Sqrt}[-1+x]])/16$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 44

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((

$m + n + 2)/((b*c - a*d)*(m + 1))$, Int[(a + b*x)^(m + 1)*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]

Rule 65

Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5378

Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_)^(m_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{1}{2} \int \frac{1}{2\sqrt{-1+xx^3}} dx \\
 &= -\frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{1}{4} \int \frac{1}{\sqrt{-1+xx^3}} dx \\
 &= \frac{\sqrt{-1+x}}{8x^2} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \int \frac{1}{\sqrt{-1+xx^2}} dx \\
 &= \frac{\sqrt{-1+x}}{8x^2} + \frac{3\sqrt{-1+x}}{16x} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{32} \int \frac{1}{\sqrt{-1+xx}} dx \\
 &= \frac{\sqrt{-1+x}}{8x^2} + \frac{3\sqrt{-1+x}}{16x} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x}\right) \\
 &= \frac{\sqrt{-1+x}}{8x^2} + \frac{3\sqrt{-1+x}}{16x} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} + \frac{3}{16} \arctan(\sqrt{-1+x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.02

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \left(\frac{1}{8x^{3/2}} + \frac{3}{16\sqrt{x}} \right) \sqrt{\frac{-1+x}{x}} - \frac{\sec^{-1}(\sqrt{x})}{2x^2} - \frac{3}{16} \arcsin\left(\frac{1}{\sqrt{x}}\right)$$

[In] Integrate[ArcSec[Sqrt[x]]/x^3,x]

[Out] (1/(8*x^(3/2)) + 3/(16*Sqrt[x]))*Sqrt[(-1 + x)/x] - ArcSec[Sqrt[x]]/(2*x^2) - (3*ArcSin[1/Sqrt[x]])/16

Maple [A] (verified)

Time = 0.22 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.06

method	result	size
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} - \frac{\sqrt{x-1} \left(3 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^2 - 3\sqrt{x-1} x - 2\sqrt{x-1} \right)}{16\sqrt{\frac{x-1}{x}} x^{\frac{5}{2}}}$	57
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} - \frac{\sqrt{x-1} \left(3 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^2 - 3\sqrt{x-1} x - 2\sqrt{x-1} \right)}{16\sqrt{\frac{x-1}{x}} x^{\frac{5}{2}}}$	57
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} + \frac{\sqrt{\frac{x-1}{x}} \left(3 \arctan(\sqrt{x-1}) x^2 + 3\sqrt{x-1} x + 2\sqrt{x-1} \right)}{16x^{\frac{3}{2}} \sqrt{x-1}}$	57

[In] int(arcsec(x^(1/2))/x^3,x,method=_RETURNVERBOSE)

[Out] -1/2*arcsec(x^(1/2))/x^2-1/16*(x-1)^(1/2)*(3*arctan(1/(x-1)^(1/2))*x^2-3*(x-1)^(1/2)*x-2*(x-1)^(1/2))/((x-1)/x)^(1/2)/x^(5/2)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.54

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{(3x^2 - 8) \operatorname{arcsec}(\sqrt{x}) + (3x + 2)\sqrt{x-1}}{16x^2}$$

[In] integrate(arcsec(x^(1/2))/x^3,x, algorithm="fricas")

[Out] 1/16*((3*x^2 - 8)*arcsec(sqrt(x)) + (3*x + 2)*sqrt(x - 1))/x^2

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 41.80 (sec) , antiderivative size = 144, normalized size of antiderivative = 2.67

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{\begin{cases} \frac{3i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{4} - \frac{3i}{4\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{i}{4x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{2x^{\frac{5}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\frac{3 \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{4} + \frac{3}{4\sqrt{x}\sqrt{1-\frac{1}{x}}} - \frac{1}{4x^{\frac{3}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{2x^{\frac{5}{2}}\sqrt{1-\frac{1}{x}}} & \text{otherwise} \end{cases}}{4} - \frac{\operatorname{asec}(\sqrt{x})}{2x^2}$$

[In] integrate(asec(x**(1/2))/x**3,x)

[Out] Piecewise((3*I*acosh(1/sqrt(x))/4 - 3*I/(4*sqrt(x)*sqrt(-1 + 1/x)) + I/(4*x**(3/2)*sqrt(-1 + 1/x)) + I/(2*x**(5/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-3*asin(1/sqrt(x))/4 + 3/(4*sqrt(x)*sqrt(1 - 1/x)) - 1/(4*x**(3/2)*sqrt(1 - 1/x)) - 1/(2*x**(5/2)*sqrt(1 - 1/x)), True))/4 - asecc(sqrt(x))/(2*x**2)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(38) = 76.

Time = 0.26 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.48

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{3x^{\frac{3}{2}}\left(-\frac{1}{x} + 1\right)^{\frac{3}{2}} + 5\sqrt{x}\sqrt{-\frac{1}{x} + 1}}{16\left(x^2\left(\frac{1}{x} - 1\right)^2 - 2x\left(\frac{1}{x} - 1\right) + 1\right)} - \frac{\operatorname{arcsec}(\sqrt{x})}{2x^2} + \frac{3}{16} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x} + 1}\right)$$

[In] integrate(arcsec(x^(1/2))/x^3,x, algorithm="maxima")

[Out] 1/16*(3*x^(3/2)*(-1/x + 1)^(3/2) + 5*sqrt(x)*sqrt(-1/x + 1))/(x^2*(1/x - 1)^2 - 2*x*(1/x - 1) + 1) - 1/2*arcsec(sqrt(x))/x^2 + 3/16*arctan(sqrt(x)*sqrt(-1/x + 1))

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.81

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \frac{3\sqrt{-\frac{1}{x}+1}}{16\sqrt{x}} + \frac{\sqrt{-\frac{1}{x}+1}}{8x^{\frac{3}{2}}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{2x^2} + \frac{3}{16}\arccos\left(\frac{1}{\sqrt{x}}\right)$$

[In] integrate(arcsec(x^(1/2))/x^3,x, algorithm="giac")

[Out] 3/16*sqrt(-1/x + 1)/sqrt(x) + 1/8*sqrt(-1/x + 1)/x^(3/2) - 1/2*arccos(1/sqrt(x))/x^2 + 3/16*arccos(1/sqrt(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^3} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x^3} dx$$

[In] int(acos(1/x^(1/2))/x^3,x)

[Out] int(acos(1/x^(1/2))/x^3, x)

3.9 $\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$

Optimal result	78
Rubi [A] (verified)	78
Mathematica [A] (verified)	80
Maple [A] (verified)	80
Fricas [A] (verification not implemented)	80
Sympy [C] (verification not implemented)	81
Maxima [B] (verification not implemented)	81
Giac [A] (verification not implemented)	82
Mupad [F(-1)]	82

Optimal result

Integrand size = 10, antiderivative size = 68

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} + \frac{5\sqrt{-1+x}}{48x} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{48} \arctan(\sqrt{-1+x})$$

[Out] $-1/3*\text{arcsec}(x^{(1/2)})/x^3+5/48*\text{arctan}((-1+x)^{(1/2)})+1/18*(-1+x)^{(1/2)}/x^3+5/72*(-1+x)^{(1/2)}/x^2+5/48*(-1+x)^{(1/2)}/x$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5378, 12, 44, 65, 209}

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{5}{48} \arctan(\sqrt{x-1}) + \frac{\sqrt{x-1}}{18x^3} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5\sqrt{x-1}}{72x^2} + \frac{5\sqrt{x-1}}{48x}$$

[In] Int[ArcSec[Sqrt[x]]/x^4,x]

[Out] Sqrt[-1 + x]/(18*x^3) + (5*Sqrt[-1 + x])/(72*x^2) + (5*Sqrt[-1 + x])/(48*x) - ArcSec[Sqrt[x]]/(3*x^3) + (5*ArcTan[Sqrt[-1 + x]])/48

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 44

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((

$m + n + 2)/((b*c - a*d)*(m + 1))$, Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5378

Int[((a_.) + ArcSec[u_]*(b_.))*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^(m + 1)*((a + b*ArcSec[u])/(d*(m + 1))), x] - Dist[b*(u/(d*(m + 1)*Sqrt[u^2])), Int[SimplifyIntegrand[(c + d*x)^(m + 1)*(D[u, x]/(u*Sqrt[u^2 - 1])), x], x], x] /; FreeQ[{a, b, c, d, m}, x] && NeQ[m, -1] && InverseFunctionFreeQ[u, x] && !FunctionOfQ[(c + d*x)^(m + 1), u, x] && !FunctionOfExponentialQ[u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{1}{3} \int \frac{1}{2\sqrt{-1+xx^4}} dx \\
 &= -\frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{1}{6} \int \frac{1}{\sqrt{-1+xx^4}} dx \\
 &= \frac{\sqrt{-1+x}}{18x^3} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{36} \int \frac{1}{\sqrt{-1+xx^3}} dx \\
 &= \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{48} \int \frac{1}{\sqrt{-1+xx^2}} dx \\
 &= \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} + \frac{5\sqrt{-1+x}}{48x} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{96} \int \frac{1}{\sqrt{-1+xx}} dx \\
 &= \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} + \frac{5\sqrt{-1+x}}{48x} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{48} \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sqrt{-1+x}\right) \\
 &= \frac{\sqrt{-1+x}}{18x^3} + \frac{5\sqrt{-1+x}}{72x^2} + \frac{5\sqrt{-1+x}}{48x} - \frac{\sec^{-1}(\sqrt{x})}{3x^3} + \frac{5}{48} \arctan(\sqrt{-1+x})
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.66

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{\sqrt{-1+x}(8+10x+15x^2) - 48\sec^{-1}(\sqrt{x}) - 15x^3 \arcsin\left(\frac{1}{\sqrt{x}}\right)}{144x^3}$$

[In] Integrate[ArcSec[Sqrt[x]]/x^4,x]

[Out] (Sqrt[-1 + x]*(8 + 10*x + 15*x^2) - 48*ArcSec[Sqrt[x]] - 15*x^3*ArcSin[1/Sqrt[x]])/(144*x^3)

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 67, normalized size of antiderivative = 0.99

method	result	size
derivativedivides	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} - \frac{\sqrt{x-1} \left(15 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^3 - 15\sqrt{x-1} x^2 - 10\sqrt{x-1} x - 8\sqrt{x-1} \right)}{144\sqrt{\frac{x-1}{x}} x^{\frac{7}{2}}}$	67
default	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} - \frac{\sqrt{x-1} \left(15 \arctan\left(\frac{1}{\sqrt{x-1}}\right) x^3 - 15\sqrt{x-1} x^2 - 10\sqrt{x-1} x - 8\sqrt{x-1} \right)}{144\sqrt{\frac{x-1}{x}} x^{\frac{7}{2}}}$	67
parts	$-\frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} + \frac{\sqrt{\frac{x-1}{x}} \left(15 \arctan(\sqrt{x-1}) x^3 + 15\sqrt{x-1} x^2 + 10\sqrt{x-1} x + 8\sqrt{x-1} \right)}{144x^{\frac{5}{2}} \sqrt{x-1}}$	67

[In] int(arcsec(x^(1/2))/x^4,x,method=_RETURNVERBOSE)

[Out] -1/3*arcsec(x^(1/2))/x^3-1/144*(x-1)^(1/2)*(15*arctan(1/(x-1)^(1/2))*x^3-15*(x-1)^(1/2)*x^2-10*(x-1)^(1/2)*x-8*(x-1)^(1/2))/((x-1)/x)^(1/2)/x^(7/2)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.51

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{3(5x^3 - 16) \operatorname{arcsec}(\sqrt{x}) + (15x^2 + 10x + 8)\sqrt{x-1}}{144x^3}$$

[In] integrate(arcsec(x^(1/2))/x^4,x, algorithm="fricas")

[Out] 1/144*(3*(5*x^3 - 16)*arcsec(sqrt(x)) + (15*x^2 + 10*x + 8)*sqrt(x - 1))/x^3

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 127.92 (sec) , antiderivative size = 180, normalized size of antiderivative = 2.65

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx$$

$$= \begin{cases} \frac{5i \operatorname{acosh}\left(\frac{1}{\sqrt{x}}\right)}{8} - \frac{5i}{8\sqrt{x}\sqrt{-1+\frac{1}{x}}} + \frac{5i}{24x^{\frac{3}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{12x^{\frac{5}{2}}\sqrt{-1+\frac{1}{x}}} + \frac{i}{3x^{\frac{7}{2}}\sqrt{-1+\frac{1}{x}}} & \text{for } \frac{1}{|x|} > 1 \\ -\frac{5 \operatorname{asin}\left(\frac{1}{\sqrt{x}}\right)}{8} + \frac{5}{8\sqrt{x}\sqrt{1-\frac{1}{x}}} - \frac{5}{24x^{\frac{3}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{12x^{\frac{5}{2}}\sqrt{1-\frac{1}{x}}} - \frac{1}{3x^{\frac{7}{2}}\sqrt{1-\frac{1}{x}}} & \text{otherwise} \end{cases}$$

$$- \frac{\operatorname{asec}(\sqrt{x})}{3x^3}$$

[In] integrate(asec(x**(1/2))/x**4,x)

[Out] Piecewise((5*I*acosh(1/sqrt(x))/8 - 5*I/(8*sqrt(x)*sqrt(-1 + 1/x)) + 5*I/(24*x**(3/2)*sqrt(-1 + 1/x)) + I/(12*x**(5/2)*sqrt(-1 + 1/x)) + I/(3*x**(7/2)*sqrt(-1 + 1/x)), 1/Abs(x) > 1), (-5*asin(1/sqrt(x))/8 + 5/(8*sqrt(x)*sqrt(1 - 1/x)) - 5/(24*x**(3/2)*sqrt(1 - 1/x)) - 1/(12*x**(5/2)*sqrt(1 - 1/x)) - 1/(3*x**(7/2)*sqrt(1 - 1/x)), True))/6 - asecc(sqrt(x))/(3*x**3)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. 2(48) = 96.

Time = 0.27 (sec) , antiderivative size = 106, normalized size of antiderivative = 1.56

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = -\frac{15x^{\frac{5}{2}}\left(-\frac{1}{x}+1\right)^{\frac{5}{2}} + 40x^{\frac{3}{2}}\left(-\frac{1}{x}+1\right)^{\frac{3}{2}} + 33\sqrt{x}\sqrt{-\frac{1}{x}+1}}{144\left(x^3\left(\frac{1}{x}-1\right)^3 - 3x^2\left(\frac{1}{x}-1\right)^2 + 3x\left(\frac{1}{x}-1\right) - 1\right)}$$

$$- \frac{\operatorname{arcsec}(\sqrt{x})}{3x^3} + \frac{5}{48} \arctan\left(\sqrt{x}\sqrt{-\frac{1}{x}+1}\right)$$

[In] integrate(arcsec(x^(1/2))/x^4,x, algorithm="maxima")

[Out] -1/144*(15*x^(5/2)*(-1/x + 1)^(5/2) + 40*x^(3/2)*(-1/x + 1)^(3/2) + 33*sqrt(x)*sqrt(-1/x + 1))/(x^3*(1/x - 1)^3 - 3*x^2*(1/x - 1)^2 + 3*x*(1/x - 1) - 1) - 1/3*arcsec(sqrt(x))/x^3 + 5/48*arctan(sqrt(x)*sqrt(-1/x + 1))

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.85

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \frac{5\sqrt{-\frac{1}{x}+1}}{48\sqrt{x}} + \frac{5\sqrt{-\frac{1}{x}+1}}{72x^{\frac{3}{2}}} + \frac{\sqrt{-\frac{1}{x}+1}}{18x^{\frac{5}{2}}} - \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{3x^3} + \frac{5}{48}\arccos\left(\frac{1}{\sqrt{x}}\right)$$

[In] integrate(arcsec(x^(1/2))/x^4,x, algorithm="giac")

[Out] 5/48*sqrt(-1/x + 1)/sqrt(x) + 5/72*sqrt(-1/x + 1)/x^(3/2) + 1/18*sqrt(-1/x + 1)/x^(5/2) - 1/3*arccos(1/sqrt(x))/x^3 + 5/48*arccos(1/sqrt(x))

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(\sqrt{x})}{x^4} dx = \int \frac{\arccos\left(\frac{1}{\sqrt{x}}\right)}{x^4} dx$$

[In] int(acos(1/x^(1/2))/x^4,x)

[Out] int(acos(1/x^(1/2))/x^4, x)

3.10 $\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	83
Rubi [A] (verified)	83
Mathematica [A] (verified)	84
Maple [A] (verified)	85
Fricas [A] (verification not implemented)	85
Sympy [A] (verification not implemented)	85
Maxima [A] (verification not implemented)	86
Giac [A] (verification not implemented)	86
Mupad [F(-1)]	86

Optimal result

Integrand size = 10, antiderivative size = 56

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{3}a^3 \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{9}a^3 \left(1 - \frac{x^2}{a^2}\right)^{3/2} + \frac{1}{3}x^3 \arccos \left(\frac{x}{a} \right)$$

[Out] $1/9*a^3*(1-x^2/a^2)^{(3/2)}+1/3*x^3*\arccos(x/a)-1/3*a^3*(1-x^2/a^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5372, 4724, 272, 45}

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{9}a^3 \left(1 - \frac{x^2}{a^2}\right)^{3/2} - \frac{1}{3}a^3 \sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{3}x^3 \arccos \left(\frac{x}{a} \right)$$

[In] Int[x^2*ArcSec[a/x],x]

[Out] $-1/3*(a^3*\text{Sqrt}[1 - x^2/a^2]) + (a^3*(1 - x^2/a^2)^{(3/2)})/9 + (x^3*\text{ArcCos}[x/a])/3$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4724

```
Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Dist[b*c*(n
/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*
x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5372

```
Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int x^2 \arccos\left(\frac{x}{a}\right) dx \\
&= \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) + \frac{\int \frac{x^3}{\sqrt{1-\frac{x^2}{a^2}}} dx}{3a} \\
&= \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) + \frac{\text{Subst}\left(\int \frac{x}{\sqrt{1-\frac{x}{a^2}}} dx, x, x^2\right)}{6a} \\
&= \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right) + \frac{\text{Subst}\left(\int \left(\frac{a^2}{\sqrt{1-\frac{x}{a^2}}} - a^2\sqrt{1-\frac{x}{a^2}}\right) dx, x, x^2\right)}{6a} \\
&= -\frac{1}{3}a^3\sqrt{1-\frac{x^2}{a^2}} + \frac{1}{9}a^3\left(1-\frac{x^2}{a^2}\right)^{3/2} + \frac{1}{3}x^3 \arccos\left(\frac{x}{a}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 42, normalized size of antiderivative = 0.75

$$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx = -\frac{1}{9}a(2a^2 + x^2)\sqrt{1-\frac{x^2}{a^2}} + \frac{1}{3}x^3 \sec^{-1}\left(\frac{a}{x}\right)$$

```
[In] Integrate[x^2*ArcSec[a/x],x]
```

```
[Out] -1/9*(a*(2*a^2 + x^2)*Sqrt[1 - x^2/a^2]) + (x^3*ArcSec[a/x])/3
```

Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00

method	result	size
parts	$\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3} + \frac{-\frac{x^2 a^2 \sqrt{1-\frac{x^2}{a^2}}}{3} - \frac{2a^4 \sqrt{1-\frac{x^2}{a^2}}}{3}}{3a}$	56
derivativedivides	$-a^3 \left(-\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3a^3} + \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66
default	$-a^3 \left(-\frac{x^3 \operatorname{arcsec}\left(\frac{a}{x}\right)}{3a^3} + \frac{\left(\frac{a^2}{x^2}-1\right)\left(\frac{2a^2}{x^2}+1\right)x^4}{9\sqrt{\frac{\left(\frac{a^2}{x^2}-1\right)x^2}{a^2}} a^4} \right)$	66

```
[In] int(x^2*arcsec(a/x),x,method=_RETURNVERBOSE)
```

```
[Out] 1/3*x^3*arcsec(a/x)+1/3/a*(-1/3*x^2*a^2*(1-x^2/a^2)^(1/2)-2/3*a^4*(1-x^2/a^2)^(1/2))
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.70

$$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx = \frac{1}{3} x^3 \operatorname{arcsec}\left(\frac{a}{x}\right) - \frac{1}{9} (2a^2 x + x^3) \sqrt{\frac{a^2 - x^2}{x^2}}$$

```
[In] integrate(x^2*arcsec(a/x),x, algorithm="fricas")
```

```
[Out] 1/3*x^3*arcsec(a/x) - 1/9*(2*a^2*x + x^3)*sqrt((a^2 - x^2)/x^2)
```

Sympy [A] (verification not implemented)

Time = 0.17 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.91

$$\int x^2 \sec^{-1}\left(\frac{a}{x}\right) dx = \begin{cases} -\frac{2a^3 \sqrt{1-\frac{x^2}{a^2}}}{9} - \frac{ax^2 \sqrt{1-\frac{x^2}{a^2}}}{9} + \frac{x^3 \operatorname{asec}\left(\frac{a}{x}\right)}{3} & \text{for } a \neq 0 \\ \infty x^3 & \text{otherwise} \end{cases}$$

```
[In] integrate(x**2*asec(a/x),x)
```

```
[Out] Piecewise((-2*a**3*sqrt(1 - x**2/a**2)/9 - a*x**2*sqrt(1 - x**2/a**2)/9 + x**3*asec(a/x)/3, Ne(a, 0)), (zoo*x**3, True))
```

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.96

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{3} x^3 \operatorname{arcsec} \left(\frac{a}{x} \right) - \frac{2 a^4 \sqrt{-\frac{x^2}{a^2} + 1} + a^2 x^2 \sqrt{-\frac{x^2}{a^2} + 1}}{9 a}$$

[In] integrate(x^2*arcsec(a/x),x, algorithm="maxima")

[Out] 1/3*x^3*arcsec(a/x) - 1/9*(2*a^4*sqrt(-x^2/a^2 + 1) + a^2*x^2*sqrt(-x^2/a^2 + 1))/a

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.84

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{3} x^3 \arccos \left(\frac{x}{a} \right) - \frac{2}{9} a^3 \sqrt{-\frac{x^2}{a^2} + 1} - \frac{1}{9} a x^2 \sqrt{-\frac{x^2}{a^2} + 1}$$

[In] integrate(x^2*arcsec(a/x),x, algorithm="giac")

[Out] 1/3*x^3*arccos(x/a) - 2/9*a^3*sqrt(-x^2/a^2 + 1) - 1/9*a*x^2*sqrt(-x^2/a^2 + 1)

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1} \left(\frac{a}{x} \right) dx = \begin{cases} \frac{x^3 \operatorname{acos} \left(\frac{x}{a} \right) - \sqrt{a^2 - x^2} (2a^2 + x^2)}{9} & \text{if } 0 < a \\ \int x^2 \operatorname{acos} \left(\frac{x}{a} \right) dx & \text{if } -0 < a \end{cases}$$

[In] int(x^2*acos(x/a),x)

[Out] piecewise(0 < a, (x^3*acos(x/a))/3 - ((a^2 - x^2)^(1/2)*(2*a^2 + x^2))/9, ~ 0 < a, int(x^2*acos(x/a), x))

3.11 $\int x \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	87
Rubi [A] (verified)	87
Mathematica [A] (verified)	88
Maple [A] (verified)	89
Fricas [A] (verification not implemented)	89
Sympy [A] (verification not implemented)	90
Maxima [A] (verification not implemented)	90
Giac [A] (verification not implemented)	90
Mupad [B] (verification not implemented)	91

Optimal result

Integrand size = 8, antiderivative size = 47

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{4}ax\sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{2}x^2 \arccos \left(\frac{x}{a} \right) + \frac{1}{4}a^2 \arcsin \left(\frac{x}{a} \right)$$

[Out] $1/2*x^2*\arccos(x/a)+1/4*a^2*\arcsin(x/a)-1/4*a*x*(1-x^2/a^2)^(1/2)$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5372, 4724, 327, 222}

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{4}a^2 \arcsin \left(\frac{x}{a} \right) - \frac{1}{4}ax\sqrt{1 - \frac{x^2}{a^2}} + \frac{1}{2}x^2 \arccos \left(\frac{x}{a} \right)$$

[In] `Int[x*ArcSec[a/x],x]`

[Out] $-1/4*(a*x*\text{Sqrt}[1 - x^2/a^2]) + (x^2*\text{ArcCos}[x/a])/2 + (a^2*\text{ArcSin}[x/a])/4$

Rule 222

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSin[Rt[-b, 2]*(x/Sqrt[a])]/Rt[-b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && NegQ[b]`

Rule 327

`Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p]`

+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 4724

```
Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:> Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Dist[b*c*(n
/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*
x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5372

```
Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int x \arccos\left(\frac{x}{a}\right) dx \\
 &= \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) + \frac{\int \frac{x^2}{\sqrt{1-\frac{x^2}{a^2}}} dx}{2a} \\
 &= -\frac{1}{4}ax\sqrt{1-\frac{x^2}{a^2}} + \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) + \frac{1}{4}a \int \frac{1}{\sqrt{1-\frac{x^2}{a^2}}} dx \\
 &= -\frac{1}{4}ax\sqrt{1-\frac{x^2}{a^2}} + \frac{1}{2}x^2 \arccos\left(\frac{x}{a}\right) + \frac{1}{4}a^2 \arcsin\left(\frac{x}{a}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.94

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = \frac{1}{4} \left(-ax\sqrt{1-\frac{x^2}{a^2}} + 2x^2 \sec^{-1}\left(\frac{a}{x}\right) + a^2 \arcsin\left(\frac{x}{a}\right) \right)$$

[In] Integrate[x*ArcSec[a/x],x]

[Out] (-a*x*Sqrt[1 - x^2/a^2]) + 2*x^2*ArcSec[a/x] + a^2*ArcSin[x/a])/4

Maple [A] (verified)

Time = 1.37 (sec) , antiderivative size = 67, normalized size of antiderivative = 1.43

method	result	size
parts	$\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2} + \frac{-\frac{x a^2 \sqrt{1-\frac{x^2}{a^2}}}{2} + \frac{a^2 \arctan\left(\frac{\sqrt{\frac{1}{a^2}} x}{\sqrt{1-\frac{x^2}{a^2}}}\right)}{2\sqrt{\frac{1}{a^2}}}}{2a}$	67
derivativeldivides	$-a^2 \left(-\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2a^2} - \frac{\left(\sqrt{\frac{a^2}{x^2}-1} \left(\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2}-1}}\right) a^2}{x^2} - \sqrt{\frac{a^2}{x^2}-1} \right) x^3 \right)}{4\sqrt{\frac{(a^2-1)x^2}{a^2}} a^3} \right)$	91
default	$-a^2 \left(-\frac{x^2 \operatorname{arcsec}\left(\frac{a}{x}\right)}{2a^2} - \frac{\left(\sqrt{\frac{a^2}{x^2}-1} \left(\frac{\arctan\left(\frac{1}{\sqrt{\frac{a^2}{x^2}-1}}\right) a^2}{x^2} - \sqrt{\frac{a^2}{x^2}-1} \right) x^3 \right)}{4\sqrt{\frac{(a^2-1)x^2}{a^2}} a^3} \right)$	91

[In] int(x*arcsec(a/x),x,method=_RETURNVERBOSE)

[Out] $\frac{1}{2}x^2 \operatorname{arcsec}\left(\frac{a}{x}\right) + \frac{1}{2}a \left(-\frac{1}{2}x a^2 \left(1 - \frac{x^2}{a^2}\right)^{\frac{1}{2}} + \frac{1}{2}a^2 \left(\frac{1}{a^2}\right)^{\frac{1}{2}} \right) \arctan\left(\frac{\left(\frac{1}{a^2}\right)^{\frac{1}{2}} x}{\left(1 - \frac{x^2}{a^2}\right)^{\frac{1}{2}}}\right)$
Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = -\frac{1}{4}x^2 \sqrt{\frac{a^2 - x^2}{x^2}} - \frac{1}{4}(a^2 - 2x^2) \operatorname{arcsec}\left(\frac{a}{x}\right)$$

[In] integrate(x*arcsec(a/x),x, algorithm="fricas")

[Out] $-\frac{1}{4}x^2 \sqrt{\frac{a^2 - x^2}{x^2}} - \frac{1}{4}(a^2 - 2x^2) \operatorname{arcsec}\left(\frac{a}{x}\right)$

Sympy [A] (verification not implemented)

Time = 0.14 (sec) , antiderivative size = 41, normalized size of antiderivative = 0.87

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \begin{cases} -\frac{a^2 \operatorname{asec} \left(\frac{a}{x} \right)}{4} - \frac{ax \sqrt{1 - \frac{x^2}{a^2}}}{4} + \frac{x^2 \operatorname{asec} \left(\frac{a}{x} \right)}{2} & \text{for } a \neq 0 \\ \infty x^2 & \text{otherwise} \end{cases}$$

[In] integrate(x*asec(a/x),x)

[Out] Piecewise((-a**2*asec(a/x)/4 - a*x*sqrt(1 - x**2/a**2)/4 + x**2*asec(a/x)/2, Ne(a, 0)), (zoo*x**2, True))

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.98

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = \frac{1}{2} x^2 \operatorname{arcsec} \left(\frac{a}{x} \right) + \frac{a^3 \arcsin \left(\frac{x}{a} \right) - a^2 x \sqrt{-\frac{x^2}{a^2} + 1}}{4a}$$

[In] integrate(x*arcsec(a/x),x, algorithm="maxima")

[Out] 1/2*x^2*arcsec(a/x) + 1/4*(a^3*arcsin(x/a) - a^2*x*sqrt(-x^2/a^2 + 1))/a

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 0.83

$$\int x \sec^{-1} \left(\frac{a}{x} \right) dx = -\frac{1}{4} a^2 \arccos \left(\frac{x}{a} \right) + \frac{1}{2} x^2 \arccos \left(\frac{x}{a} \right) - \frac{1}{4} ax \sqrt{-\frac{x^2}{a^2} + 1}$$

[In] integrate(x*arcsec(a/x),x, algorithm="giac")

[Out] -1/4*a^2*arccos(x/a) + 1/2*x^2*arccos(x/a) - 1/4*a*x*sqrt(-x^2/a^2 + 1)

Mupad [B] (verification not implemented)

Time = 0.73 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

$$\int x \sec^{-1}\left(\frac{a}{x}\right) dx = \frac{a^2 \arccos\left(\frac{x}{a}\right) \left(\frac{2x^2}{a^2} - 1\right)}{4} - \frac{ax \sqrt{1 - \frac{x^2}{a^2}}}{4}$$

[In] int(x*acos(x/a),x)

[Out] (a^2*acos(x/a)*((2*x^2)/a^2 - 1))/4 - (a*x*(1 - x^2/a^2)^(1/2))/4

3.12 $\int \sec^{-1} \left(\frac{a}{x} \right) dx$

Optimal result	92
Rubi [A] (verified)	92
Mathematica [A] (verified)	93
Maple [A] (verified)	93
Fricas [A] (verification not implemented)	94
Sympy [A] (verification not implemented)	94
Maxima [A] (verification not implemented)	94
Giac [A] (verification not implemented)	95
Mupad [B] (verification not implemented)	95

Optimal result

Integrand size = 6, antiderivative size = 26

$$\int \sec^{-1} \left(\frac{a}{x} \right) dx = -a\sqrt{1 - \frac{x^2}{a^2}} + x \arccos \left(\frac{x}{a} \right)$$

[Out] $x*\arccos(x/a)-a*(1-x^2/a^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5372, 4716, 267}

$$\int \sec^{-1} \left(\frac{a}{x} \right) dx = x \arccos \left(\frac{x}{a} \right) - a\sqrt{1 - \frac{x^2}{a^2}}$$

[In] $\text{Int}[\text{ArcSec}[a/x], x]$

[Out] $-(a*\text{Sqrt}[1 - x^2/a^2]) + x*\text{ArcCos}[x/a]$

Rule 267

$\text{Int}[(x_)^{(m_.)}*((a_) + (b_.)*(x_)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x^n)^{(p + 1)}/(b*n*(p + 1)), x] /;$ FreeQ[{a, b, m, n, p}, x] && EqQ[m, n - 1] && NeQ[p, -1]

Rule 4716

$\text{Int}[((a_.) + \text{ArcCos}[(c_.)*(x_)])*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[x*(a + b*\text{ArcCos}[c*x])^n, x] + \text{Dist}[b*c*n, \text{Int}[x*((a + b*\text{ArcCos}[c*x])^{(n - 1)})/\text{Sqrt}[1 - c^2*x^2]), x, x] /;$ FreeQ[{a, b, c}, x] && GtQ[n, 0]

Rule 5372

`Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] :> Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]`

Rubi steps

$$\begin{aligned} \text{integral} &= \int \arccos\left(\frac{x}{a}\right) dx \\ &= x \arccos\left(\frac{x}{a}\right) + \frac{\int \frac{x}{\sqrt{1-\frac{x^2}{a^2}}} dx}{a} \\ &= -a\sqrt{1-\frac{x^2}{a^2}} + x \arccos\left(\frac{x}{a}\right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = -a\sqrt{1-\frac{x^2}{a^2}} + x \sec^{-1}\left(\frac{a}{x}\right)$$

[In] `Integrate[ArcSec[a/x], x]`

[Out] `-(a*Sqrt[1 - x^2/a^2]) + x*ArcSec[a/x]`

Maple [A] (verified)

Time = 1.35 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

method	result	size
parts	$x \operatorname{arcsec}\left(\frac{a}{x}\right) - a\sqrt{\frac{a^2-x^2}{a^2}}$	28
derivativedivides	$-a\left(-\frac{x \operatorname{arcsec}\left(\frac{a}{x}\right)}{a} + \frac{x^2\left(\frac{a^2}{x^2}-1\right)}{\sqrt{\left(\frac{a^2}{x^2}-1\right)x^2} a^2}\right)$	51
default	$-a\left(-\frac{x \operatorname{arcsec}\left(\frac{a}{x}\right)}{a} + \frac{x^2\left(\frac{a^2}{x^2}-1\right)}{\sqrt{\left(\frac{a^2}{x^2}-1\right)x^2} a^2}\right)$	51

[In] `int(arcsec(a/x), x, method=_RETURNVERBOSE)`

[Out] `x*arcsec(a/x)-a*((a^2-x^2)/a^2)^(1/2)`

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.04

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \operatorname{arcsec}\left(\frac{a}{x}\right) - x \sqrt{\frac{a^2 - x^2}{x^2}}$$

[In] integrate(arcsec(a/x),x, algorithm="fricas")

[Out] x*arcsec(a/x) - x*sqrt((a^2 - x^2)/x^2)

Sympy [A] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 22, normalized size of antiderivative = 0.85

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = \begin{cases} -a\sqrt{1 - \frac{x^2}{a^2}} + x \operatorname{asec}\left(\frac{a}{x}\right) & \text{for } a \neq 0 \\ \infty & \text{otherwise} \end{cases}$$

[In] integrate(asec(a/x),x)

[Out] Piecewise((-a*sqrt(1 - x**2/a**2) + x*asec(a/x), Ne(a, 0)), (zoo*x, True))

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \operatorname{arcsec}\left(\frac{a}{x}\right) - a \sqrt{-\frac{x^2}{a^2} + 1}$$

[In] integrate(arcsec(a/x),x, algorithm="maxima")

[Out] x*arcsec(a/x) - a*sqrt(-x^2/a^2 + 1)

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.08

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = a\left(\frac{x \arccos\left(\frac{x}{a}\right)}{a} - \sqrt{-\frac{x^2}{a^2} + 1}\right)$$

[In] integrate(arcsec(a/x),x, algorithm="giac")

[Out] a*(x*arccos(x/a)/a - sqrt(-x^2/a^2 + 1))

Mupad [B] (verification not implemented)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.92

$$\int \sec^{-1}\left(\frac{a}{x}\right) dx = x \arccos\left(\frac{x}{a}\right) - a \sqrt{1 - \frac{x^2}{a^2}}$$

[In] int(acos(x/a),x)

[Out] x*acos(x/a) - a*(1 - x^2/a^2)^(1/2)

3.13 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx$

Optimal result	96
Rubi [A] (verified)	96
Mathematica [A] (verified)	98
Maple [A] (verified)	98
Fricas [F]	99
Sympy [F]	99
Maxima [F]	99
Giac [F]	99
Mupad [F(-1)]	100

Optimal result

Integrand size = 10, antiderivative size = 59

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x}{a}\right)}\right)$$

[Out] $-1/2*I*\arccos(x/a)^2 + \arccos(x/a)*\ln(1+(x/a+I*(1-x^2/a^2)^{(1/2)})^2) - 1/2*I*\operatorname{polylog}(2, -(x/a+I*(1-x^2/a^2)^{(1/2)})^2)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5372, 4722, 3800, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = -\frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right)$$

[In] $\operatorname{Int}[\operatorname{ArcSec}[a/x]/x, x]$

[Out] $(-1/2*I)*\operatorname{ArcCos}[x/a]^2 + \operatorname{ArcCos}[x/a]*\operatorname{Log}[1 + E^{((2*I)*\operatorname{ArcCos}[x/a])}] - (I/2)*\operatorname{PolyLog}[2, -E^{((2*I)*\operatorname{ArcCos}[x/a])}]$

Rule 2221

$\operatorname{Int}[(((F_)^{((g_)*(e_)+(f_)*(x_)))})^{(n_)*((c_)+(d_)*(x_))^{(m_)}))/((a_)+(b_)*((F_)^{((g_)*(e_)+(f_)*(x_)))})^{(n_)}), x_Symbol] :> \operatorname{Simp}$

$[(c + dx)^m / (bfgn \log[F]) * \log[1 + b((F^{g(e+fx)})^n/a)], x] - \text{Dist}[d(m / (bfgn \log[F])), \text{Int}[(c + dx)^{m-1} * \log[1 + b((F^{g(e+fx)})^n/a)], x], x] /;$ FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

$\text{Int}[\log[a + (b \cdot (F^{(e \cdot (c + dx))})^n)], x_Symbol] \rightarrow \text{Dist}[1 / (d \cdot e \cdot n \cdot \log[F]), \text{Subst}[\text{Int}[\log[a + b \cdot x] / x, x], x, (F^{e \cdot (c + dx)})^n], x] /;$ FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

$\text{Int}[\log[(c \cdot (d + (e \cdot x)^n))] / (x), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c) \cdot e \cdot x^n / n, x] /;$ FreeQ[{c, d, e, n}, x] && EqQ[c \cdot d, 1]

Rule 3800

$\text{Int}[(c + dx)^m \cdot \tan[e + f \cdot x], x_Symbol] \rightarrow \text{Simp}[I * ((c + dx)^{m+1} / (d \cdot (m+1))), x] - \text{Dist}[2 \cdot I, \text{Int}[(c + dx)^m \cdot (E^{2 \cdot I \cdot (e + fx)}) / (1 + E^{2 \cdot I \cdot (e + fx)})], x, x] /;$ FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

Rule 4722

$\text{Int}[(a + \arccos[(c \cdot x) \cdot b])^n / (x), x_Symbol] \rightarrow -\text{Subst}[\text{Int}[(a + b \cdot x)^n \cdot \tan[x], x], x, \arccos[c \cdot x]] /;$ FreeQ[{a, b, c}, x] && IGtQ[n, 0]

Rule 5372

$\text{Int}[\text{ArcSec}[(c \cdot (a + (b \cdot x)^n))]^m \cdot u, x_Symbol] \rightarrow \text{Int}[u \cdot \text{ArcCos}[a/c + b \cdot (x^n/c)]^m, x] /;$ FreeQ[{a, b, c, n, m}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx \\ &= -\text{Subst}\left(\int x \tan(x) dx, x, \arccos\left(\frac{x}{a}\right)\right) \\ &= -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + 2i \text{Subst}\left(\int \frac{e^{2ix} x}{1 + e^{2ix}} dx, x, \arccos\left(\frac{x}{a}\right)\right) \\ &= -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) \\ &\quad - \text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \arccos\left(\frac{x}{a}\right)\right) \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) \\
&\quad + \frac{1}{2}i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \arccos\left(\frac{x}{a}\right)}\right) \\
&= -\frac{1}{2}i \arccos\left(\frac{x}{a}\right)^2 + \arccos\left(\frac{x}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x}{a}\right)}\right) - \frac{1}{2}i \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x}{a}\right)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\begin{aligned}
\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx &= -\frac{1}{2}i \sec^{-1}\left(\frac{a}{x}\right)^2 + \sec^{-1}\left(\frac{a}{x}\right) \log\left(1 + e^{2i \sec^{-1}\left(\frac{a}{x}\right)}\right) \\
&\quad - \frac{1}{2}i \text{PolyLog}\left(2, -e^{2i \sec^{-1}\left(\frac{a}{x}\right)}\right)
\end{aligned}$$

[In] Integrate[ArcSec[a/x]/x,x]

[Out] $(-1/2*I)*\text{ArcSec}[a/x]^2 + \text{ArcSec}[a/x]*\text{Log}[1 + E^{\left((2*I)*\text{ArcSec}[a/x]\right)}] - (I/2)*\text{PolyLog}[2, -E^{\left((2*I)*\text{ArcSec}[a/x]\right)}]$

Maple [A] (verified)

Time = 1.77 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.29

method	result	size
derivativedivides	$-\frac{i \operatorname{arcsec}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arcsec}\left(\frac{a}{x}\right) \ln\left(1 + \left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right) - \frac{i \operatorname{polylog}\left(2, -\left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right)}{2}$	76
default	$-\frac{i \operatorname{arcsec}\left(\frac{a}{x}\right)^2}{2} + \operatorname{arcsec}\left(\frac{a}{x}\right) \ln\left(1 + \left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right) - \frac{i \operatorname{polylog}\left(2, -\left(\frac{x}{a} + i\sqrt{1 - \frac{x^2}{a^2}}\right)^2\right)}{2}$	76

[In] int(arcsec(a/x)/x,x,method=_RETURNVERBOSE)

[Out] $-1/2*I*\operatorname{arcsec}(a/x)^2 + \operatorname{arcsec}(a/x)*\ln\left(1 + \left(\frac{x}{a} + I*\left(1 - \frac{x^2}{a^2}\right)^{(1/2)}\right)^2\right) - 1/2*I*\operatorname{polylog}\left(2, -\left(\frac{x}{a} + I*\left(1 - \frac{x^2}{a^2}\right)^{(1/2)}\right)^2\right)$

Fricas [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

[In] integrate(arcsec(a/x)/x,x, algorithm="fricas")

[Out] integral(arcsec(a/x)/x, x)

Sympy [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x} dx$$

[In] integrate(asec(a/x)/x,x)

[Out] Integral(asec(a/x)/x, x)

Maxima [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

[In] integrate(arcsec(a/x)/x,x, algorithm="maxima")

[Out] integrate(arcsec(a/x)/x, x)

Giac [F]

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{x} dx$$

[In] integrate(arcsec(a/x)/x,x, algorithm="giac")

[Out] integrate(arcsec(a/x)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} dx = \int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx$$

```
[In] int(acos(x/a)/x,x)
```

```
[Out] int(acos(x/a)/x, x)
```

3.14 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx$

Optimal result	101
Rubi [A] (verified)	101
Mathematica [B] (verified)	103
Maple [A] (verified)	103
Fricas [B] (verification not implemented)	103
Sympy [C] (verification not implemented)	104
Maxima [A] (verification not implemented)	104
Giac [B] (verification not implemented)	104
Mupad [B] (verification not implemented)	105

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\arccos\left(\frac{x}{a}\right)}{x} + \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a}$$

[Out] $-\arccos(x/a)/x + \operatorname{arctanh}((1-x^2/a^2)^{(1/2)})/a$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5372, 4724, 272, 65, 214}

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a} - \frac{\arccos\left(\frac{x}{a}\right)}{x}$$

[In] $\text{Int}[\text{ArcSec}[a/x]/x^2, x]$

[Out] $-(\text{ArcCos}[x/a]/x) + \text{ArcTanh}[\text{Sqrt}[1 - x^2/a^2]]/a$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 4724

Int[((a_) + ArcCos[(c_)*(x_)])*(b_)^(n_)*((d_)*(x_)^(m_)), x_Symbol] := Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Dist[b*c*(n/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]

Rule 5372

Int[ArcSec[(c_)/((a_) + (b_)*(x_)^(n_))]^(m_)*(u_), x_Symbol] := Int[u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\arccos\left(\frac{x}{a}\right)}{x^2} dx \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{x} - \frac{\int \frac{1}{x\sqrt{1-\frac{x^2}{a^2}}} dx}{a} \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{x} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1-\frac{x^2}{a^2}}} dx, x, x^2\right)}{2a} \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{x} + a\text{Subst}\left(\int \frac{1}{a^2 - a^2x^2} dx, x, \sqrt{1 - \frac{x^2}{a^2}}\right) \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{x} + \frac{\text{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{a}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 93 vs. $2(31) = 62$.

Time = 0.09 (sec) , antiderivative size = 93, normalized size of antiderivative = 3.00

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\sec^{-1}\left(\frac{a}{x}\right)}{x} + \frac{\sqrt{-1 + \frac{a^2}{x^2}} x \left(-\log\left(1 - \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right) + \log\left(1 + \frac{a}{\sqrt{-1 + \frac{a^2}{x^2}} x}\right) \right)}{2a^2 \sqrt{1 - \frac{x^2}{a^2}}}$$

[In] Integrate[ArcSec[a/x]/x^2,x]

[Out] $-(\text{ArcSec}[a/x]/x) + (\text{Sqrt}[-1 + a^2/x^2]*x*(-\text{Log}[1 - a/(\text{Sqrt}[-1 + a^2/x^2]*x)] + \text{Log}[1 + a/(\text{Sqrt}[-1 + a^2/x^2]*x)]))/(2*a^2*\text{Sqrt}[1 - x^2/a^2])$

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.97

method	result	size
parts	$-\frac{\text{arcsec}\left(\frac{a}{x}\right)}{x} + \frac{\text{arctanh}\left(\frac{1}{\sqrt{1 - \frac{x^2}{a^2}}}\right)}{a}$	30
derivativedivides	$-\frac{\frac{a \text{ arcsec}\left(\frac{a}{x}\right)}{x} - \ln\left(\frac{a}{x} + \frac{a\sqrt{1 - \frac{x^2}{a^2}}}{x}\right)}{a}$	44
default	$-\frac{\frac{a \text{ arcsec}\left(\frac{a}{x}\right)}{x} - \ln\left(\frac{a}{x} + \frac{a\sqrt{1 - \frac{x^2}{a^2}}}{x}\right)}{a}$	44

[In] int(arcsec(a/x)/x^2,x,method=_RETURNVERBOSE)

[Out] $-\text{arcsec}(a/x)/x + 1/a * \text{arctanh}(1/(1 - x^2/a^2)^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(29) = 58$.

Time = 0.27 (sec) , antiderivative size = 107, normalized size of antiderivative = 3.45

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{2ax \arctan\left(-\frac{x^2 \sqrt{\frac{a^2 - x^2}{x^2}}}{a^2 - x^2}\right) - 2(ax - a) \text{arcsec}\left(\frac{a}{x}\right) - x \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} + a\right) + x \log\left(x \sqrt{\frac{a^2 - x^2}{x^2}} - a\right)}{2ax}$$

[In] integrate(arcsec(a/x)/x^2,x, algorithm="fricas")

[Out] -1/2*(2*a*x*arctan(-x^2*sqrt((a^2 - x^2)/x^2)/(a^2 - x^2)) - 2*(a*x - a)*arcsec(a/x) - x*log(x*sqrt((a^2 - x^2)/x^2) + a) + x*log(x*sqrt((a^2 - x^2)/x^2) - a))/(a*x)

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\operatorname{asec}\left(\frac{a}{x}\right)}{x} - \frac{\begin{cases} -\operatorname{acosh}\left(\frac{a}{x}\right) & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ i \operatorname{asin}\left(\frac{a}{x}\right) & \text{otherwise} \end{cases}}{a}$$

[In] integrate(asec(a/x)/x**2,x)

[Out] -asec(a/x)/x - Piecewise((-acosh(a/x), Abs(a**2/x**2) > 1), (I*asin(a/x), True))/a

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.68

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = -\frac{\frac{2a \operatorname{arcsec}\left(\frac{a}{x}\right)}{x} - \log\left(\sqrt{-\frac{x^2}{a^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{x^2}{a^2} + 1} + 1\right)}{2a}$$

[In] integrate(arcsec(a/x)/x^2,x, algorithm="maxima")

[Out] -1/2*(2*a*arcsec(a/x)/x - log(sqrt(-x^2/a^2 + 1) + 1) + log(-sqrt(-x^2/a^2 + 1) + 1))/a

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(29) = 58.

Time = 0.27 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.97

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{a\left(\frac{\log\left(|a+\sqrt{a^2-x^2}\right|}{a} - \frac{\log\left(|-a+\sqrt{a^2-x^2}\right|}{a}\right)}{2|a|} - \frac{\arccos\left(\frac{x}{a}\right)}{x}$$

[In] integrate(arcsec(a/x)/x^2,x, algorithm="giac")

[Out] 1/2*a*(log(abs(a + sqrt(a^2 - x^2)))/a - log(abs(-a + sqrt(a^2 - x^2)))/a)/abs(a) - arccos(x/a)/x

Mupad [B] (verification not implemented)

Time = 0.71 (sec) , antiderivative size = 29, normalized size of antiderivative = 0.94

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^2} dx = \frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1-\frac{x^2}{a^2}}}\right)}{a} - \frac{\operatorname{acos}\left(\frac{x}{a}\right)}{x}$$

[In] int(acos(x/a)/x^2,x)

[Out] atanh(1/(1 - x^2/a^2)^(1/2))/a - acos(x/a)/x

3.15 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx$

Optimal result	106
Rubi [A] (verified)	106
Mathematica [A] (verified)	107
Maple [A] (verified)	107
Fricas [A] (verification not implemented)	108
Sympy [C] (verification not implemented)	108
Maxima [A] (verification not implemented)	108
Giac [A] (verification not implemented)	109
Mupad [F(-1)]	109

Optimal result

Integrand size = 10, antiderivative size = 38

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2}$$

[Out] $-1/2*\arccos(x/a)/x^2+1/2*(1-x^2/a^2)^{(1/2)}/a/x$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5372, 4724, 270}

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2}$$

[In] Int[ArcSec[a/x]/x^3,x]

[Out] Sqrt[1 - x^2/a^2]/(2*a*x) - ArcCos[x/a]/(2*x^2)

Rule 270

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a + b*x^n)^(p+1)/(a*c*(m+1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m+1)/n + p + 1, 0] && NeQ[m, -1]

Rule 4724

Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol] := Simp[(d*x)^(m+1)*((a + b*ArcCos[c*x])^n/(d*(m+1))), x] + Dist[b*c*(n

$\int (d*(m + 1)) \int [(d*x)^{(m + 1)} * ((a + b*\text{ArcCos}[c*x])^{(n - 1)} / \text{Sqrt}[1 - c^2*x^2])], x], x] /; \text{FreeQ}\{a, b, c, d, m\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{NeQ}[m, -1]$

Rule 5372

$\text{Int}[\text{ArcSec}[(c_.) / ((a_.) + (b_.) * (x_)^{(n_.)})]^{(m_.)} * (u_.), x_Symbol] \rightarrow \text{Int}[u * \text{ArcCos}[a/c + b*(x^n/c)]^m, x] /; \text{FreeQ}\{a, b, c, n, m\}, x]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{\arccos\left(\frac{x}{a}\right)}{x^3} dx \\ &= -\frac{\arccos\left(\frac{x}{a}\right)}{2x^2} - \frac{\int \frac{1}{x^2 \sqrt{1 - \frac{x^2}{a^2}}} dx}{2a} \\ &= \frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.95

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{x \sqrt{1 - \frac{x^2}{a^2}} - a \sec^{-1}\left(\frac{a}{x}\right)}{2ax^2}$$

[In] Integrate[ArcSec[a/x]/x^3,x]

[Out] (x*Sqrt[1 - x^2/a^2] - a*ArcSec[a/x])/(2*a*x^2)

Maple [A] (verified)

Time = 1.35 (sec) , antiderivative size = 33, normalized size of antiderivative = 0.87

method	result	size
parts	$-\frac{\text{arcsec}\left(\frac{a}{x}\right)}{2x^2} + \frac{\sqrt{1 - \frac{x^2}{a^2}}}{2ax}$	33
derivativedivides	$-\frac{\frac{a^2 \text{arcsec}\left(\frac{a}{x}\right)}{2x^2} - \frac{x\left(\frac{a^2}{x^2} - 1\right)}{2\sqrt{\left(\frac{a^2}{x^2} - 1\right)x^2} a}}{a^2}$	54
default	$-\frac{\frac{a^2 \text{arcsec}\left(\frac{a}{x}\right)}{2x^2} - \frac{x\left(\frac{a^2}{x^2} - 1\right)}{2\sqrt{\left(\frac{a^2}{x^2} - 1\right)x^2} a}}{a^2}$	54

[In] `int(arcsec(a/x)/x^3,x,method=_RETURNVERBOSE)`

[Out] $-1/2*\text{arcsec}(a/x)/x^2+1/2*(1-x^2/a^2)^{(1/2)}/a/x$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.03

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{a^2 \text{arcsec}\left(\frac{a}{x}\right) - x^2 \sqrt{\frac{a^2-x^2}{x^2}}}{2 a^2 x^2}$$

[In] `integrate(arcsec(a/x)/x^3,x, algorithm="fricas")`

[Out] $-1/2*(a^2*\text{arcsec}(a/x) - x^2*\text{sqrt}((a^2 - x^2)/x^2))/(a^2*x^2)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.62 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.39

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{\text{asec}\left(\frac{a}{x}\right)}{2x^2} - \frac{\begin{cases} -\frac{\sqrt{\frac{a^2}{x^2}-1}}{a} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ -\frac{i\sqrt{-\frac{a^2}{x^2}+1}}{a} & \text{otherwise} \end{cases}}{2a}$$

[In] `integrate(asec(a/x)/x**3,x)`

[Out] $-\text{asec}(a/x)/(2*x**2) - \text{Piecewise}((- \text{sqrt}(a**2/x**2 - 1)/a, \text{Abs}(a**2/x**2) > 1), (-I*\text{sqrt}(-a**2/x**2 + 1)/a, \text{True}))/ (2*a)$

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 32, normalized size of antiderivative = 0.84

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = -\frac{\text{arcsec}\left(\frac{a}{x}\right)}{2 x^2} + \frac{\sqrt{-\frac{x^2}{a^2} + 1}}{2 a x}$$

[In] `integrate(arcsec(a/x)/x^3,x, algorithm="maxima")`

[Out] $-1/2*\text{arcsec}(a/x)/x^2 + 1/2*\text{sqrt}(-x^2/a^2 + 1)/(a*x)$

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.61

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \frac{a\left(\frac{a+\sqrt{a^2-x^2}}{a^2x} - \frac{x}{(a+\sqrt{a^2-x^2})a^2}\right)}{4|a|} - \frac{\arccos\left(\frac{x}{a}\right)}{2x^2}$$

[In] integrate(arcsec(a/x)/x^3,x, algorithm="giac")

[Out] 1/4*a*((a + sqrt(a^2 - x^2))/(a^2*x) - x/((a + sqrt(a^2 - x^2))*a^2))/abs(a) - 1/2*arccos(x/a)/x^2

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^3} dx = \int \frac{\arccos\left(\frac{x}{a}\right)}{x^3} dx$$

[In] int(acos(x/a)/x^3,x)

[Out] int(acos(x/a)/x^3, x)

3.16 $\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx$

Optimal result	110
Rubi [A] (verified)	110
Mathematica [A] (verified)	112
Maple [A] (verified)	112
Fricas [B] (verification not implemented)	113
Sympy [C] (verification not implemented)	113
Maxima [A] (verification not implemented)	114
Giac [A] (verification not implemented)	114
Mupad [F(-1)]	114

Optimal result

Integrand size = 10, antiderivative size = 60

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} + \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3}$$

[Out] $-1/3*\arccos(x/a)/x^3+1/6*\operatorname{arctanh}((1-x^2/a^2)^{(1/2)})/a^3+1/6*(1-x^2/a^2)^{(1/2)}/a/x^2$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5372, 4724, 272, 44, 65, 214}

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} + \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3}$$

[In] Int[ArcSec[a/x]/x^4,x]

[Out] Sqrt[1 - x^2/a^2]/(6*a*x^2) - ArcCos[x/a]/(3*x^3) + ArcTanh[Sqrt[1 - x^2/a^2]]/(6*a^3)

Rule 44

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((
m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x
] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !Int
```

egerQ[n] && LtQ[n, 0]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 4724

```
Int[((a_.) + ArcCos[(c_.)*(x_)]*(b_.))^(n_.)*((d_.)*(x_))^(m_.), x_Symbol]
:= Simp[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^n/(d*(m + 1))), x] + Dist[b*c*(n
/(d*(m + 1))), Int[(d*x)^(m + 1)*((a + b*ArcCos[c*x])^(n - 1)/Sqrt[1 - c^2*
x^2]), x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && NeQ[m, -1]
```

Rule 5372

```
Int[ArcSec[(c_.)/((a_.) + (b_.)*(x_)^(n_.))]^(m_.)*(u_.), x_Symbol] := Int[
u*ArcCos[a/c + b*(x^n/c)]^m, x] /; FreeQ[{a, b, c, n, m}, x]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \frac{\arccos\left(\frac{x}{a}\right)}{x^4} dx \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{3x^3} - \frac{\int \frac{1}{x^3\sqrt{1-\frac{x^2}{a^2}}} dx}{3a} \\
 &= -\frac{\arccos\left(\frac{x}{a}\right)}{3x^3} - \frac{\text{Subst}\left(\int \frac{1}{x^2\sqrt{1-\frac{x}{a^2}}} dx, x, x^2\right)}{6a}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} - \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{1 - \frac{x}{a^2}}} dx, x, x^2\right)}{12a^3} \\
&= \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} + \frac{\text{Subst}\left(\int \frac{1}{a^2 - a^2x^2} dx, x, \sqrt{1 - \frac{x^2}{a^2}}\right)}{6a} \\
&= \frac{\sqrt{1 - \frac{x^2}{a^2}}}{6ax^2} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3} + \frac{\text{arctanh}\left(\sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.15

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{a^2x\sqrt{1 - \frac{x^2}{a^2}} - 2a^3\sec^{-1}\left(\frac{a}{x}\right) - x^3\log(x) + x^3\log\left(1 + \sqrt{1 - \frac{x^2}{a^2}}\right)}{6a^3x^3}$$

[In] Integrate[ArcSec[a/x]/x^4,x]

[Out] (a^2*x*Sqrt[1 - x^2/a^2] - 2*a^3*ArcSec[a/x] - x^3*Log[x] + x^3*Log[1 + Sqrt[1 - x^2/a^2]])/(6*a^3*x^3)

Maple [A] (verified)

Time = 1.33 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.90

method	result	size
parts	$ -\frac{\text{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\frac{\sqrt{1 - \frac{x^2}{a^2}}}{2x^2} - \frac{\text{arctanh}\left(\frac{1}{\sqrt{1 - \frac{x^2}{a^2}}}\right)}{2a^2}}{3a} $	54
derivativedivides	$ -\frac{a^3 \text{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(a \frac{\sqrt{\frac{a^2}{x^2} - 1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2} - 1}\right) \right) x}{6\sqrt{\frac{\left(\frac{a^2}{x^2} - 1\right)x^2}{a^2}} a}{a^3} $	91
default	$ -\frac{a^3 \text{arcsec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\sqrt{\frac{a^2}{x^2} - 1} \left(a \frac{\sqrt{\frac{a^2}{x^2} - 1}}{x} + \ln\left(\frac{a}{x} + \sqrt{\frac{a^2}{x^2} - 1}\right) \right) x}{6\sqrt{\frac{\left(\frac{a^2}{x^2} - 1\right)x^2}{a^2}} a}{a^3} $	91

[In] int(arcsec(a/x)/x^4,x,method=_RETURNVERBOSE)

[Out] $-1/3 \operatorname{arcsec}(a/x)/x^3 - 1/3/a * (-1/2/x^2 * (1-x^2/a^2)^{(1/2)} - 1/2/a^2 * \operatorname{arctanh}(1/(1-x^2/a^2)^{(1/2)}))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 142 vs. 2(50) = 100.

Time = 0.28 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.37

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{4a^3x^3 \arctan\left(-\frac{x^2\sqrt{\frac{a^2-x^2}{x^2}}}{a^2-x^2}\right) - x^3 \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} + a\right) + x^3 \log\left(x\sqrt{\frac{a^2-x^2}{x^2}} - a\right) - 2ax^2\sqrt{\frac{a^2-x^2}{x^2}} - 4(a^2-x^2)\sqrt{\frac{a^2-x^2}{x^2}}}{12a^3x^3}$$

[In] `integrate(arcsec(a/x)/x^4,x, algorithm="fricas")`

[Out] $-1/12*(4*a^3*x^3*\arctan(-x^2*\sqrt{(a^2-x^2)/x^2})/(a^2-x^2)) - x^3*\log(x*\sqrt{(a^2-x^2)/x^2}+a) + x^3*\log(x*\sqrt{(a^2-x^2)/x^2}-a) - 2*a*x^2*\sqrt{(a^2-x^2)/x^2} - 4*(a^3*x^3-a^3)*\operatorname{arcsec}(a/x)/(a^3*x^3)$

Sympy [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.40 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.67

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = -\frac{\operatorname{asec}\left(\frac{a}{x}\right)}{3x^3} - \frac{\begin{cases} -\frac{\sqrt{\frac{a^2}{x^2}-1}}{2ax} - \frac{\operatorname{acosh}\left(\frac{a}{x}\right)}{2a^2} & \text{for } \left|\frac{a^2}{x^2}\right| > 1 \\ \frac{ia}{2x^3\sqrt{-\frac{a^2}{x^2}+1}} - \frac{i}{2ax\sqrt{-\frac{a^2}{x^2}+1}} + \frac{i\operatorname{asin}\left(\frac{a}{x}\right)}{2a^2} & \text{otherwise} \end{cases}}{3a}$$

[In] `integrate(asec(a/x)/x**4,x)`

[Out] $-\operatorname{asec}(a/x)/(3*x**3) - \operatorname{Piecewise}((- \sqrt{a**2/x**2 - 1})/(2*a*x) - \operatorname{acosh}(a/x)/(2*a**2), \operatorname{Abs}(a**2/x**2) > 1), (I*a/(2*x**3*\sqrt{-a**2/x**2 + 1}) - I/(2*a*x*\sqrt{-a**2/x**2 + 1}) + I*\operatorname{asin}(a/x)/(2*a**2), \operatorname{True}))/ (3*a)$

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.07

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{\log\left(\frac{2\sqrt{-\frac{x^2}{a^2}+1}}{|x|} + \frac{2}{|x|}\right)}{6a} + \frac{\sqrt{-\frac{x^2}{a^2}+1}}{x^2} - \frac{\operatorname{arcsec}\left(\frac{a}{x}\right)}{3x^3}$$

[In] integrate(arcsec(a/x)/x^4,x, algorithm="maxima")

[Out] 1/6*(log(2*sqrt(-x^2/a^2 + 1)/abs(x) + 2/abs(x))/a^2 + sqrt(-x^2/a^2 + 1)/x^2)/a - 1/3*arcsec(a/x)/x^3

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 80, normalized size of antiderivative = 1.33

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \frac{a\left(\frac{\log\left(|a+\sqrt{a^2-x^2}\right|}{a^3} - \frac{\log\left(|-a+\sqrt{a^2-x^2}\right|}{a^3} + \frac{2\sqrt{a^2-x^2}}{a^2x^2}\right)}{12|a|} - \frac{\arccos\left(\frac{x}{a}\right)}{3x^3}$$

[In] integrate(arcsec(a/x)/x^4,x, algorithm="giac")

[Out] 1/12*a*(log(abs(a + sqrt(a^2 - x^2)))/a^3 - log(abs(-a + sqrt(a^2 - x^2)))/a^3 + 2*sqrt(a^2 - x^2)/(a^2*x^2))/abs(a) - 1/3*arccos(x/a)/x^3

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}\left(\frac{a}{x}\right)}{x^4} dx = \int \frac{\operatorname{acos}\left(\frac{x}{a}\right)}{x^4} dx$$

[In] int(acos(x/a)/x^4,x)

[Out] int(acos(x/a)/x^4, x)

3.17 $\int \frac{\sec^{-1}(ax^n)}{x} dx$

Optimal result	115
Rubi [A] (verified)	115
Mathematica [C] (verified)	117
Maple [A] (verified)	117
Fricas [F(-2)]	118
Sympy [F]	118
Maxima [F]	118
Giac [F]	119
Mupad [F(-1)]	119

Optimal result

Integrand size = 10, antiderivative size = 69

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \frac{i \sec^{-1}(ax^n)^2}{2n} - \frac{\sec^{-1}(ax^n) \log\left(1 + e^{2i \sec^{-1}(ax^n)}\right)}{n} + \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^n)}\right)}{2n}$$

[Out] $1/2*I*\operatorname{arcsec}(a*x^n)^2/n - \operatorname{arcsec}(a*x^n)*\ln(1+(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)/n + 1/2*I*\operatorname{polylog}(2, -(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)/n$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5326, 4722, 3800, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(ax^n)}\right)}{2n} + \frac{i \sec^{-1}(ax^n)^2}{2n} - \frac{\sec^{-1}(ax^n) \log\left(1 + e^{2i \sec^{-1}(ax^n)}\right)}{n}$$

[In] $\operatorname{Int}[\operatorname{ArcSec}[a*x^n]/x, x]$

[Out] $((I/2)*\operatorname{ArcSec}[a*x^n]^2)/n - (\operatorname{ArcSec}[a*x^n]*\operatorname{Log}[1 + E^{((2*I)*\operatorname{ArcSec}[a*x^n])}])/n + ((I/2)*\operatorname{PolyLog}[2, -E^{((2*I)*\operatorname{ArcSec}[a*x^n])}])/n$

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[(((c_) + (d_)*(x_))^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
 + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 4722

```
Int[(((a_) + ArcCos[(c_)*(x_)])*(b_))^(n_)/(x_), x_Symbol] := -Subst[Int[
(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0
]
```

Rule 5326

```
Int[(((a_) + ArcSec[(c_)*(x_)])*(b_))/(x_), x_Symbol] := -Subst[Int[(a + b
*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int \frac{\sec^{-1}(ax)}{x} dx, x, x^n\right)}{n} \\ &= -\frac{\text{Subst}\left(\int \frac{\arccos\left(\frac{x}{a}\right)}{x} dx, x, x^{-n}\right)}{n} \\ &= \frac{\text{Subst}\left(\int x \tan(x) dx, x, \arccos\left(\frac{x^{-n}}{a}\right)\right)}{n} \end{aligned}$$

$$\begin{aligned}
&= \frac{i \arccos\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{(2i) \text{Subst}\left(\int \frac{e^{2ix} x}{1+e^{2ix}} dx, x, \arccos\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{i \arccos\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \arccos\left(\frac{x^{-n}}{a}\right)\right)}{n} \\
&= \frac{i \arccos\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)}{n} \\
&\quad - \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)}{2n} \\
&= \frac{i \arccos\left(\frac{x^{-n}}{a}\right)^2}{2n} - \frac{\arccos\left(\frac{x^{-n}}{a}\right) \log\left(1 + e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)}{n} + \frac{i \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{x^{-n}}{a}\right)}\right)}{2n}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

Time = 0.06 (sec) , antiderivative size = 60, normalized size of antiderivative = 0.87

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \frac{x^{-n} {}_3F_2\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}; \frac{3}{2}, \frac{3}{2}; \frac{x^{-2n}}{a^2}\right)}{an} + \left(\sec^{-1}(ax^n) + \arcsin\left(\frac{x^{-n}}{a}\right)\right) \log(x)$$

[In] Integrate[ArcSec[a*x^n]/x,x]

[Out] HypergeometricPFQ[{1/2, 1/2, 1/2}, {3/2, 3/2}, 1/(a^2*x^(2*n))]/(a*n*x^n) + (ArcSec[a*x^n] + ArcSin[1/(a*x^n)])*Log[x]

Maple [A] (verified)

Time = 1.44 (sec) , antiderivative size = 93, normalized size of antiderivative = 1.35

method	result	size
derivativedivides	$\frac{\frac{i \operatorname{arcsec}(a x^n)^2}{2} - \operatorname{arcsec}(a x^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + i\sqrt{1 - \frac{x^{-2n}}{a^2}}\right)^2\right)}{n} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + i\sqrt{1 - \frac{x^{-2n}}{a^2}}\right)^2\right)}{2}}$	93
default	$\frac{\frac{i \operatorname{arcsec}(a x^n)^2}{2} - \operatorname{arcsec}(a x^n) \ln\left(1 + \left(\frac{x^{-n}}{a} + i\sqrt{1 - \frac{x^{-2n}}{a^2}}\right)^2\right)}{n} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{x^{-n}}{a} + i\sqrt{1 - \frac{x^{-2n}}{a^2}}\right)^2\right)}{2}}$	93

[In] `int(arcsec(a*x^n)/x,x,method=_RETURNVERBOSE)`

[Out] $1/n*(1/2*I*\operatorname{arcsec}(a*x^n)^2 - \operatorname{arcsec}(a*x^n)*\ln(1+(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2)+1/2*I*\operatorname{polylog}(2,-(1/a/(x^n)+I*(1-1/a^2/(x^n)^2)^{(1/2)})^2))$

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \text{Exception raised: TypeError}$$

[In] `integrate(arcsec(a*x^n)/x,x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{asec}(ax^n)}{x} dx$$

[In] `integrate(asec(a*x**n)/x,x)`

[Out] `Integral(asec(a*x**n)/x, x)`

Maxima [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsec}(ax^n)}{x} dx$$

[In] `integrate(arcsec(a*x^n)/x,x, algorithm="maxima")`

[Out] $-a^{2n}*\operatorname{integrate}(\sqrt{a*x^n + 1}*\sqrt{a*x^n - 1}*\log(x)/(a^4*x*x^{(2n)} - a^{2*x}), x) + \operatorname{arctan}(\sqrt{a*x^n + 1}*\sqrt{a*x^n - 1})*\log(x)$

Giac [F]

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{arcsec}(ax^n)}{x} dx$$

[In] integrate(arcsec(a*x^n)/x,x, algorithm="giac")

[Out] integrate(arcsec(a*x^n)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(ax^n)}{x} dx = \int \frac{\operatorname{acos}\left(\frac{1}{ax^n}\right)}{x} dx$$

[In] int(acos(1/(a*x^n))/x,x)

[Out] int(acos(1/(a*x^n))/x, x)

3.18 $\int x^4 \sec^{-1}(a + bx) dx$

Optimal result	120
Rubi [A] (verified)	120
Mathematica [A] (verified)	124
Maple [A] (verified)	124
Fricas [A] (verification not implemented)	125
Sympy [F]	125
Maxima [F]	125
Giac [B] (verification not implemented)	126
Mupad [F(-1)]	126

Optimal result

Integrand size = 10, antiderivative size = 197

$$\int x^4 \sec^{-1}(a + bx) dx = \frac{a(20 + 53a^2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{30b^5} + \frac{11ax^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{60b^3}$$

$$- \frac{x^3(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{20b^2} - \frac{(9 + 58a^2)(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{120b^5}$$

$$+ \frac{a^5 \sec^{-1}(a + bx)}{5b^5} + \frac{1}{5}x^5 \sec^{-1}(a + bx)$$

$$- \frac{(3 + 40a^2 + 40a^4) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{40b^5}$$

[Out] $\frac{1}{5}a^5 \operatorname{arcsec}(b*x+a)/b^5 + \frac{1}{5}x^5 \operatorname{arcsec}(b*x+a) - \frac{1}{40} * (40*a^4 + 40*a^2 + 3) * \operatorname{arctanh}\left(\frac{1 - 1/(b*x+a)^2}{(1/2)}\right) / b^5 + \frac{1}{30} * a * (53*a^2 + 20) * (b*x+a) * \frac{1 - 1/(b*x+a)^2}{(1/2)} / b^5 + \frac{11}{60} * a * x^2 * (b*x+a) * \frac{1 - 1/(b*x+a)^2}{(1/2)} / b^3 - \frac{1}{20} * x^3 * (b*x+a) * \frac{1 - 1/(b*x+a)^2}{(1/2)} / b^2 - \frac{1}{120} * (58*a^2 + 9) * (b*x+a)^2 * \frac{1 - 1/(b*x+a)^2}{(1/2)} / b^5$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 197, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used

= {5366, 4511, 3867, 4141, 4133, 3855, 3852, 8}

$$\int x^4 \sec^{-1}(a + bx) dx = \frac{a^5 \sec^{-1}(a + bx)}{5b^5} + \frac{(53a^2 + 20) a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{30b^5} - \frac{(58a^2 + 9) (a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}}}{120b^5} - \frac{(40a^4 + 40a^2 + 3) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{40b^5} + \frac{11ax^2(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{20b^2} + \frac{1}{5}x^5 \sec^{-1}(a + bx)$$

[In] Int[x^4*ArcSec[a + b*x],x]

[Out] (a*(20 + 53*a^2)*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]/(30*b^5) + (11*a*x^2*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]/(60*b^3) - (x^3*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]/(20*b^2) - ((9 + 58*a^2)*(a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]/(120*b^5) + (a^5*ArcSec[a + b*x])/(5*b^5) + (x^5*ArcSec[a + b*x])/5 - ((3 + 40*a^2 + 40*a^4)*ArcTanh[Sqrt[1 - (a + b*x)^(-2)]])/(40*b^5)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4133

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*((csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Simp[(-b)*C*Csc[e +
f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A
+ C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a,
b, e, f, A, B, C}, x]
```

Rule 4141

```
Int[((A_.) + csc[(e_.) + (f_.)*(x_.)]*(B_.) + csc[(e_.) + (f_.)*(x_.)]^2*(C_.
))*((csc[(e_.) + (f_.)*(x_.)]*(b_.) + (a_.))^(m_.), x_Symbol] := Simp[(-C)*Cot
[e + f*x]*((a + b*Csc[e + f*x])^m/(f*(m + 1))), x] + Dist[1/(m + 1), Int[(a
+ b*Csc[e + f*x])^(m - 1)*Simp[a*A*(m + 1) + ((A*b + a*B)*(m + 1) + b*C*m)
*Csc[e + f*x] + (b*B*(m + 1) + a*C*m)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{
a, b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && IGtQ[2*m, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*Sec[(c_.) + (d_.)*(x_.)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_.)])^(n_.)*Tan[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x \sec(x) (-a + \sec(x))^4 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^5} \\
&= \frac{1}{5} x^5 \sec^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \sec(x))^5 dx, x, \sec^{-1}(a + bx)\right)}{5b^5} \\
&= -\frac{x^3(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}}}{20b^2} + \frac{1}{5} x^5 \sec^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-a + \sec(x))^2 (-4a^3 + 3(1 + 4a^2) \sec(x) - 11a \sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{20b^5}
\end{aligned}$$

$$\begin{aligned}
&= \frac{11ax^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{20b^2} + \frac{1}{5}x^5\sec^{-1}(a+bx) \\
&\quad \text{Subst}\left(\int(-a+\sec(x))(12a^4-a(31+48a^2)\sec(x)+(9+58a^2)\sec^2(x))dx, x, \sec^{-1}(a+bx)\right) \\
&\quad \frac{60b^5}{60b^5} \\
&= \frac{11ax^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{20b^2} \\
&\quad - \frac{(9+58a^2)(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}}{120b^5} + \frac{1}{5}x^5\sec^{-1}(a+bx) \\
&\quad \text{Subst}\left(\int(-24a^5+3(3+40a^2+40a^4)\sec(x)-4a(20+53a^2)\sec^2(x))dx, x, \sec^{-1}(a+bx)\right) \\
&\quad \frac{120b^5}{120b^5} \\
&= \frac{11ax^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{20b^2} \\
&\quad - \frac{(9+58a^2)(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}}{120b^5} + \frac{a^5\sec^{-1}(a+bx)}{5b^5} \\
&\quad + \frac{1}{5}x^5\sec^{-1}(a+bx) + \frac{(a(20+53a^2))\text{Subst}\left(\int\sec^2(x)dx, x, \sec^{-1}(a+bx)\right)}{30b^5} \\
&\quad - \frac{(3+40a^2+40a^4)\text{Subst}\left(\int\sec(x)dx, x, \sec^{-1}(a+bx)\right)}{40b^5} \\
&= \frac{11ax^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{60b^3} - \frac{x^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{20b^2} \\
&\quad - \frac{(9+58a^2)(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}}{120b^5} + \frac{a^5\sec^{-1}(a+bx)}{5b^5} \\
&\quad + \frac{1}{5}x^5\sec^{-1}(a+bx) - \frac{(3+40a^2+40a^4)\operatorname{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{40b^5} \\
&\quad - \frac{(a(20+53a^2))\text{Subst}\left(\int 1 dx, x, -\left((a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\right)\right)}{30b^5} \\
&= \frac{a(20+53a^2)(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{30b^5} + \frac{11ax^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{60b^3} \\
&\quad - \frac{x^3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{20b^2} - \frac{(9+58a^2)(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}}{120b^5} + \frac{a^5\sec^{-1}(a+bx)}{5b^5} \\
&\quad + \frac{1}{5}x^5\sec^{-1}(a+bx) - \frac{(3+40a^2+40a^4)\operatorname{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{40b^5}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 173, normalized size of antiderivative = 0.88

$$\int x^4 \sec^{-1}(a + bx) dx$$

$$= \frac{\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} (a^2(71 + 154a^2) + 2a(31 + 48a^2)bx - 9(1 + 4a^2)b^2x^2 + 16ab^3x^3 - 6b^4x^4) + 24b^5x^5 \sec^{-1}(a+bx)}{120b^5}$$

`[In] Integrate[x^4*ArcSec[a + b*x], x]`

```
[Out] (Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^2*(71 + 154*a^2) + 2*a
*(31 + 48*a^2)*b*x - 9*(1 + 4*a^2)*b^2*x^2 + 16*a*b^3*x^3 - 6*b^4*x^4) + 24
*b^5*x^5*ArcSec[a + b*x] - 24*a^5*ArcSin[(a + b*x)^(-1)] - 3*(3 + 40*a^2 +
40*a^4)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]
)])/ (120*b^5)
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 329, normalized size of antiderivative = 1.67

method	result
derivativedivides	$-\frac{\operatorname{arcsec}(bx+a)a^5}{5} + \operatorname{arcsec}(bx+a)a^4(bx+a) - 2 \operatorname{arcsec}(bx+a)a^3(bx+a)^2 + 2 \operatorname{arcsec}(bx+a)a^2(bx+a)^3 - \operatorname{arcsec}(bx+a)a(bx+a)$
default	$-\frac{\operatorname{arcsec}(bx+a)a^5}{5} + \operatorname{arcsec}(bx+a)a^4(bx+a) - 2 \operatorname{arcsec}(bx+a)a^3(bx+a)^2 + 2 \operatorname{arcsec}(bx+a)a^2(bx+a)^3 - \operatorname{arcsec}(bx+a)a(bx+a)$
parts	$\frac{x^5 \operatorname{arcsec}(bx+a)}{5} + \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(-6x^3\sqrt{b^2x^2+2abx+a^2-1}b^3\sqrt{b^2+22\sqrt{b^2x^2+2abx+a^2-1}}\sqrt{b^2}ab^2x^2-24a^5 \operatorname{arctan}\left(\frac{1}{(bx+a)^2-1}\right) \right)}{\dots}$

`[In] int(x^4*arcsec(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b^5*(-1/5*arcsec(b*x+a)*a^5+arcsec(b*x+a)*a^4*(b*x+a)-2*arcsec(b*x+a)*a^3
*(b*x+a)^2+2*arcsec(b*x+a)*a^2*(b*x+a)^3-arcsec(b*x+a)*a*(b*x+a)^4+1/5*arcs
ec(b*x+a)*(b*x+a)^5+1/120*((b*x+a)^2-1)^(1/2)*(-24*a^5*arctan(1/((b*x+a)^2-
1)^(1/2))-120*a^4*ln(b*x+a+((b*x+a)^2-1)^(1/2))+240*a^3*((b*x+a)^2-1)^(1/2)
-120*a^2*(b*x+a)*((b*x+a)^2-1)^(1/2)+40*a*(b*x+a)^2*((b*x+a)^2-1)^(1/2)-6*(
b*x+a)^3*((b*x+a)^2-1)^(1/2)-120*a^2*ln(b*x+a+((b*x+a)^2-1)^(1/2))+80*a*((b
*x+a)^2-1)^(1/2)-9*(b*x+a)*((b*x+a)^2-1)^(1/2)-9*ln(b*x+a+((b*x+a)^2-1)^(1/
2))))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 152, normalized size of antiderivative = 0.77

$$\int x^4 \sec^{-1}(a + bx) dx$$

$$= \frac{24 b^5 x^5 \operatorname{arcsec}(bx + a) + 48 a^5 \arctan(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1}) + 3(40 a^4 + 40 a^2 + 3) \log(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 - 1})}{b^5}$$

[In] integrate(x^4*arcsec(b*x+a),x, algorithm="fricas")

```
[Out] 1/120*(24*b^5*x^5*arcsec(b*x + a) + 48*a^5*arctan(-b*x - a + sqrt(b^2*x^2 +
2*a*b*x + a^2 - 1)) + 3*(40*a^4 + 40*a^2 + 3)*log(-b*x - a + sqrt(b^2*x^2
+ 2*a*b*x + a^2 - 1)) - (6*b^3*x^3 - 22*a*b^2*x^2 - 154*a^3 + (58*a^2 + 9)*
b*x - 71*a)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/b^5
```

Sympy [F]

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \operatorname{asec}(a + bx) dx$$

[In] integrate(x**4*asec(b*x+a),x)

[Out] Integral(x**4*asec(a + b*x), x)

Maxima [F]

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \operatorname{arcsec}(bx + a) dx$$

[In] integrate(x^4*arcsec(b*x+a),x, algorithm="maxima")

```
[Out] 1/5*x^5*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/5*(b^2*x^
6 + a*b*x^5)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a
*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x +
a - 1)) - 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 409 vs. 2(173) = 346.

Time = 0.31 (sec) , antiderivative size = 409, normalized size of antiderivative = 2.08

$$\int x^4 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{960} b \left(\frac{192 (bx + a)^5 \left(\frac{5a}{bx+a} - \frac{10a^2}{(bx+a)^2} + \frac{10a^3}{(bx+a)^3} - \frac{5a^4}{(bx+a)^4} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right) - \frac{3 (bx + a)^4}{b^6} \right)$$

[In] integrate(x^4*arcsec(b*x+a),x, algorithm="giac")

[Out] -1/960*b*(192*(b*x + a)^5*(5*a/(b*x + a) - 10*a^2/(b*x + a)^2 + 10*a^3/(b*x + a)^3 - 5*a^4/(b*x + a)^4 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^6 - (3*(b*x + a)^4*(sqrt(-1/(b*x + a)^2 + 1) - 1)^4 + 40*(b*x + a)^3*a*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 240*(b*x + a)^2*a^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 960*(b*x + a)*a^3*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 24*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 360*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 24*(40*a^4 + 40*a^2 + 3)*log(-(sqrt(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - (120*(8*a^3 + 3*a)*(b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 24*(10*a^2 + 1)*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 40*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 3)/((b*x + a)^4*(sqrt(-1/(b*x + a)^2 + 1) - 1)^4))/b^6)

Mupad [F(-1)]

Timed out.

$$\int x^4 \sec^{-1}(a + bx) dx = \int x^4 \arccos\left(\frac{1}{a + bx}\right) dx$$

[In] int(x^4*acos(1/(a + b*x)),x)

[Out] int(x^4*acos(1/(a + b*x)), x)

3.19 $\int x^3 \sec^{-1}(a + bx) dx$

Optimal result	127
Rubi [A] (verified)	127
Mathematica [A] (verified)	130
Maple [A] (verified)	130
Fricas [A] (verification not implemented)	131
Sympy [F]	131
Maxima [F]	131
Giac [B] (verification not implemented)	132
Mupad [F(-1)]	132

Optimal result

Integrand size = 10, antiderivative size = 155

$$\int x^3 \sec^{-1}(a + bx) dx = -\frac{(2 + 17a^2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^4} - \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2}$$

$$+ \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \sec^{-1}(a + bx)}{4b^4}$$

$$+ \frac{1}{4}x^4 \sec^{-1}(a + bx) + \frac{a(1 + 2a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{2b^4}$$

[Out] $-1/4*a^4*\operatorname{arcsec}(b*x+a)/b^4+1/4*x^4*\operatorname{arcsec}(b*x+a)+1/2*a*(2*a^2+1)*\operatorname{arctanh}((1-1/(b*x+a)^2)^{(1/2)})/b^4-1/12*(17*a^2+2)*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^4-1/12*x^2*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^2+1/3*a*(b*x+a)^2*(1-1/(b*x+a)^2)^{(1/2)}/b^4$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5366, 4511, 3867, 4133, 3855, 3852, 8}

$$\int x^3 \sec^{-1}(a + bx) dx = -\frac{a^4 \sec^{-1}(a + bx)}{4b^4} + \frac{(2a^2 + 1) a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{2b^4}$$

$$- \frac{(17a^2 + 2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^4} + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4}$$

$$- \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{1}{4}x^4 \sec^{-1}(a + bx)$$

[In] Int[x^3*ArcSec[a + b*x],x]

[Out] $-1/12*((2 + 17*a^2)*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}])/b^4 - (x^2*(a + b*x)*\text{Sqrt}[1 - (a + b*x)^{-2}])/(12*b^2) + (a*(a + b*x)^2*\text{Sqrt}[1 - (a + b*x)^{-2}])/(3*b^4) - (a^4*\text{ArcSec}[a + b*x])/(4*b^4) + (x^4*\text{ArcSec}[a + b*x])/4 + (a*(1 + 2*a^2)*\text{ArcTanh}[\text{Sqrt}[1 - (a + b*x)^{-2}]])/(2*b^4)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4133

Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.)), x_Symbol] := Simp[(-b)*C*Csc[e + f*x]*(Cot[e + f*x]/(2*f)), x] + Dist[1/2, Int[Simp[2*A*a + (2*B*a + b*(2*A + C))*Csc[e + f*x] + 2*(a*C + B*b)*Csc[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f, A, B, C}, x]

Rule 4511

Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)]^(n_.)*Tan[(c_.) + (d_.)*(x_)]), x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366


```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x \sec(x)(-a + \sec(x))^3 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^4} \\
&= \frac{1}{4}x^4 \sec^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \sec(x))^4 dx, x, \sec^{-1}(a + bx)\right)}{4b^4} \\
&= -\frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{1}{4}x^4 \sec^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-a + \sec(x))(-3a^3 + (2 + 9a^2)\sec(x) - 8a\sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{12b^4} \\
&= -\frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} + \frac{1}{4}x^4 \sec^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (6a^4 - 12a(1 + 2a^2)\sec(x) + 2(2 + 17a^2)\sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{24b^4} \\
&= -\frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \sec^{-1}(a + bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4 \sec^{-1}(a + bx) + \frac{(a(1 + 2a^2))\text{Subst}\left(\int \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{2b^4} \\
&\quad - \frac{(2 + 17a^2)\text{Subst}\left(\int \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{12b^4} \\
&= -\frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} - \frac{a^4 \sec^{-1}(a + bx)}{4b^4} \\
&\quad + \frac{1}{4}x^4 \sec^{-1}(a + bx) + \frac{a(1 + 2a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{2b^4} \\
&\quad + \frac{(2 + 17a^2)\text{Subst}\left(\int 1 dx, x, -\left((a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}\right)\right)}{12b^4} \\
&= -\frac{(2 + 17a^2)(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^4} - \frac{x^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{12b^2} + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}}}{3b^4} \\
&\quad - \frac{a^4 \sec^{-1}(a + bx)}{4b^4} + \frac{1}{4}x^4 \sec^{-1}(a + bx) + \frac{a(1 + 2a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{2b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.18 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.97

$$\int x^3 \sec^{-1}(a + bx) dx = \frac{-\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}(2a + 13a^3 + 2bx + 9a^2bx - 3ab^2x^2 + b^3x^3) + 3b^4x^4 \sec^{-1}(a + bx) + 3a^4 \arcsin\left(\frac{1}{a+bx}\right)}{12b^4}$$

`[In] Integrate[x^3*ArcSec[a + b*x], x]`

```
[Out] (-Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(2*a + 13*a^3 + 2*b*x + 9*a^2*b*x - 3*a*b^2*x^2 + b^3*x^3)) + 3*b^4*x^4*ArcSec[a + b*x] + 3*a^4*ArcSin[(a + b*x)^(-1)] + 6*a*(1 + 2*a^2)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])]/(12*b^4)
```

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.61

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsec}(bx+a)a^4}{4} - \operatorname{arcsec}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsec}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^4}{4} + \frac{\sqrt{(bx+a)^2 - 1}}{bx+a}}$
default	$\frac{\operatorname{arcsec}(bx+a)a^4}{4} - \operatorname{arcsec}(bx+a)a^3(bx+a) + \frac{3 \operatorname{arcsec}(bx+a)a^2(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^4}{4} + \frac{\sqrt{(bx+a)^2 - 1}}{bx+a}$
parts	$\frac{x^4 \operatorname{arcsec}(bx+a)}{4} + \frac{\sqrt{b^2x^2+2abx+a^2-1}}{12b^4} \left(3a^4 \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2-x^2} \sqrt{b^2x^2+2abx+a^2-1} b^2 \sqrt{b^2} + 12 \ln\left(\frac{\sqrt{b^2x^2+2abx+a^2-1}}{bx+a}\right) \right)$

`[In] int(x^3*arcsec(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b^4*(1/4*arcsec(b*x+a)*a^4-arcsec(b*x+a)*a^3*(b*x+a)+3/2*arcsec(b*x+a)*a^2*(b*x+a)^2-arcsec(b*x+a)*a*(b*x+a)^3+1/4*arcsec(b*x+a)*(b*x+a)^4+1/12*((b*x+a)^2-1)^(1/2)*(3*a^4*arctan(1/((b*x+a)^2-1)^(1/2))+12*a^3*ln(b*x+a+((b*x+a)^2-1)^(1/2))-18*a^2*((b*x+a)^2-1)^(1/2)+6*a*(b*x+a)*((b*x+a)^2-1)^(1/2)-(b*x+a)^2*((b*x+a)^2-1)^(1/2)+6*a*ln(b*x+a+((b*x+a)^2-1)^(1/2))-2*((b*x+a)^2-1)^(1/2))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 130, normalized size of antiderivative = 0.84

$$\int x^3 \sec^{-1}(a + bx) dx$$

$$= \frac{3b^4x^4 \operatorname{arcsec}(bx + a) - 6a^4 \arctan(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) - 6(2a^3 + a) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{12b^4}$$

[In] integrate(x^3*arcsec(b*x+a),x, algorithm="fricas")

```
[Out] 1/12*(3*b^4*x^4*arcsec(b*x + a) - 6*a^4*arctan(-b*x - a + sqrt(b^2*x^2 + 2*
a*b*x + a^2 - 1)) - 6*(2*a^3 + a)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a
^2 - 1)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(b^2*x^2 - 4*a*b*x + 13*a^2 +
2))/b^4
```

Sympy [F]

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \operatorname{asec}(a + bx) dx$$

[In] integrate(x**3*asec(b*x+a),x)

[Out] Integral(x**3*asec(a + b*x), x)

Maxima [F]

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \operatorname{arcsec}(bx + a) dx$$

[In] integrate(x^3*arcsec(b*x+a),x, algorithm="maxima")

```
[Out] 1/4*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/4*(b^2*x^
5 + a*b*x^4)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a
*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x +
a - 1)) - 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(135) = 270.

Time = 0.31 (sec) , antiderivative size = 299, normalized size of antiderivative = 1.93

$$\int x^3 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{96} b \left(\frac{24 (bx + a)^4 \left(\frac{4a}{bx+a} - \frac{6a^2}{(bx+a)^2} + \frac{4a^3}{(bx+a)^3} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right) + (bx + a)^3 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)^3 + 12 (bx + a)^2 a \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)^2 + 72 (bx + a) a^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + 9 (bx + a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + 48 (2a^3 + a) \log \left(-\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) \operatorname{abs}(bx + a) \right) - (9(8a^2 + 1)(bx + a)^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)^2 + 12 (bx + a) a \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + 1) / ((bx + a)^3 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)^3)}{b^5} \right)$$

[In] integrate(x^3*arcsec(b*x+a),x, algorithm="giac")

[Out] -1/96*b*(24*(b*x + a)^4*(4*a/(b*x + a) - 6*a^2/(b*x + a)^2 + 4*a^3/(b*x + a)^3 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^5 + ((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3 + 12*(b*x + a)^2*a*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 72*(b*x + a)*a^2*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 9*(b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 48*(2*a^3 + a)*log(-(sqrt(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - (9*(8*a^2 + 1)*(b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)/((b*x + a)^3*(sqrt(-1/(b*x + a)^2 + 1) - 1)^3))/b^5)

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(a + bx) dx = \int x^3 \operatorname{acos} \left(\frac{1}{a + bx} \right) dx$$

[In] int(x^3*acos(1/(a + b*x)),x)

[Out] int(x^3*acos(1/(a + b*x)), x)

3.20 $\int x^2 \sec^{-1}(a + bx) dx$

Optimal result	133
Rubi [A] (verified)	133
Mathematica [A] (verified)	135
Maple [A] (verified)	136
Fricas [A] (verification not implemented)	136
Sympy [F]	137
Maxima [F]	137
Giac [B] (verification not implemented)	137
Mupad [F(-1)]	138

Optimal result

Integrand size = 10, antiderivative size = 116

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{5a(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^3} - \frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2}$$

$$+ \frac{a^3 \sec^{-1}(a + bx)}{3b^3} + \frac{1}{3}x^3 \sec^{-1}(a + bx)$$

$$- \frac{(1 + 6a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{6b^3}$$

[Out] $1/3*a^3*\operatorname{arcsec}(b*x+a)/b^3+1/3*x^3*\operatorname{arcsec}(b*x+a)-1/6*(6*a^2+1)*\operatorname{arctanh}((1-1/(b*x+a)^2)^{(1/2)})/b^3+5/6*a*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^3-1/6*x*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {5366, 4511, 3867, 3855, 3852, 8}

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{a^3 \sec^{-1}(a + bx)}{3b^3} - \frac{(6a^2 + 1) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{6b^3}$$

$$+ \frac{5a(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^3}$$

$$- \frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{1}{3}x^3 \sec^{-1}(a + bx)$$

[In] Int[x^2*ArcSec[a + b*x],x]

[Out] (5*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]/(6*b^3) - (x*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]/(6*b^2) + (a^3*ArcSec[a + b*x])/(3*b^3) + (x^3*ArcSec[a + b*x])/3 - ((1 + 6*a^2)*ArcTanh[Sqrt[1 - (a + b*x)^(-2)]])/(6*b^3)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3867

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] := Simp[(-b^2)*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n - 2)/(d*(n - 1))), x] + Dist[1/(n - 1), Int[(a + b*Csc[c + d*x])^(n - 3)*Simp[a^3*(n - 1) + (b*(b^2*(n - 2) + 3*a^2*(n - 1)))*Csc[c + d*x] + (a*b^2*(3*n - 4))*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && GtQ[n, 2] && IntegerQ[2*n]

Rule 4511

Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366

Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x \sec(x)(-a + \sec(x))^2 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
&= \frac{1}{3}x^3 \sec^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \sec(x))^3 dx, x, \sec^{-1}(a + bx)\right)}{3b^3} \\
&= -\frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{1}{3}x^3 \sec^{-1}(a + bx) \\
&\quad - \frac{\text{Subst}\left(\int (-2a^3 + (1 + 6a^2)\sec(x) - 5a \sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{6b^3} \\
&= -\frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \sec^{-1}(a + bx)}{3b^3} + \frac{1}{3}x^3 \sec^{-1}(a + bx) \\
&\quad + \frac{(5a)\text{Subst}\left(\int \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{6b^3} \\
&\quad - \frac{(1 + 6a^2)\text{Subst}\left(\int \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{6b^3} \\
&= -\frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \sec^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3}x^3 \sec^{-1}(a + bx) - \frac{(1 + 6a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{6b^3} \\
&\quad - \frac{(5a)\text{Subst}\left(\int 1 dx, x, -\left((a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}\right)\right)}{6b^3} \\
&= \frac{5a(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^3} - \frac{x(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{6b^2} + \frac{a^3 \sec^{-1}(a + bx)}{3b^3} \\
&\quad + \frac{1}{3}x^3 \sec^{-1}(a + bx) - \frac{(1 + 6a^2) \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{6b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.13 (sec) , antiderivative size = 131, normalized size of antiderivative = 1.13

$$\begin{aligned}
&\int x^2 \sec^{-1}(a + bx) dx \\
&= \frac{(5a^2 + 4abx - b^2x^2) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} + 2b^3x^3 \sec^{-1}(a + bx) - 2a^3 \arcsin\left(\frac{1}{a+bx}\right) - (1 + 6a^2) \log\left((a + bx)\right)}{6b^3}
\end{aligned}$$

[In] Integrate[x^2*ArcSec[a + b*x], x]

```
[Out] ((5*a^2 + 4*a*b*x - b^2*x^2)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2] + 2*b^3*x^3*ArcSec[a + b*x] - 2*a^3*ArcSin[(a + b*x)^(-1)] - (1 + 6*a^2)*Log[(a + b*x)*(1 + Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2])]/(6*b^3)
```

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.64

method	result
derivativedivides	$-\frac{\operatorname{arcsec}\left(\frac{bx+a}{3}\right)a^3}{3} + \operatorname{arcsec}(bx+a)a^2(bx+a) - \operatorname{arcsec}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^3}{3} - \frac{\sqrt{(bx+a)^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \right)}{b^3}$
default	$-\frac{\operatorname{arcsec}\left(\frac{bx+a}{3}\right)a^3}{3} + \operatorname{arcsec}(bx+a)a^2(bx+a) - \operatorname{arcsec}(bx+a)a(bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)(bx+a)^3}{3} - \frac{\sqrt{(bx+a)^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \right)}{b^3}$
parts	$\frac{x^3 \operatorname{arcsec}(bx+a)}{3} - \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(2a^3 \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \right) \sqrt{b^2+6} \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2-1}\sqrt{b^2+ab}}{\sqrt{b^2}}\right)}{6b^3 \sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}}}$

```
[In] int(x^2*arcsec(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b^3*(-1/3*arcsec(b*x+a)*a^3+arcsec(b*x+a)*a^2*(b*x+a)-arcsec(b*x+a)*a*(b*x+a)^2+1/3*arcsec(b*x+a)*(b*x+a)^3-1/6*((b*x+a)^2-1)^(1/2)*(2*a^3*arctan(1/((b*x+a)^2-1)^(1/2))+6*a^2*ln(b*x+a+((b*x+a)^2-1)^(1/2))-6*a*((b*x+a)^2-1)^(1/2)+(b*x+a)*((b*x+a)^2-1)^(1/2)+ln(b*x+a+((b*x+a)^2-1)^(1/2)))/(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a))
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.01

$$\int x^2 \sec^{-1}(a + bx) dx = \frac{2b^3x^3 \operatorname{arcsec}(bx+a) + 4a^3 \arctan(-bx-a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) + (6a^2 + 1) \log(-bx-a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{6b^3}$$

```
[In] integrate(x^2*arcsec(b*x+a),x, algorithm="fricas")
```

```
[Out] 1/6*(2*b^3*x^3*arcsec(b*x + a) + 4*a^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (6*a^2 + 1)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(b*x - 5*a))/b^3
```


Sympy [F]

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \operatorname{asec}(a + bx) dx$$

```
[In] integrate(x**2*asec(b*x+a),x)
```

```
[Out] Integral(x**2*asec(a + b*x), x)
```

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \operatorname{arcsec}(bx + a) dx$$

```
[In] integrate(x^2*arcsec(b*x+a),x, algorithm="maxima")
```

```
[Out] 1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/3*(b^2*x^4 + a*b*x^3)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x + a - 1)) - 1), x)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(100) = 200.

Time = 0.31 (sec) , antiderivative size = 204, normalized size of antiderivative = 1.76

$$\int x^2 \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{24} b \left(\frac{8 (bx + a)^3 \left(\frac{3a}{bx+a} - \frac{3a^2}{(bx+a)^2} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right) - (bx + a)^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right)^2}{b^4} \right)$$

```
[In] integrate(x^2*arcsec(b*x+a),x, algorithm="giac")
```

```
[Out] -1/24*b*(8*(b*x + a)^3*(3*a/(b*x + a) - 3*a^2/(b*x + a)^2 - 1)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^4 - ((b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2 + 12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 4*(6*a^2 + 1)*log(-(sqrt(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - (12*(b*x + a)*a*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 1)/((b*x + a)^2*(sqrt(-1/(b*x + a)^2 + 1) - 1)^2))/b^4)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx) dx = \int x^2 \arccos\left(\frac{1}{a + bx}\right) dx$$

```
[In] int(x^2*acos(1/(a + b*x)),x)
```

```
[Out] int(x^2*acos(1/(a + b*x)), x)
```

3.21 $\int x \sec^{-1}(a + bx) dx$

Optimal result	139
Rubi [A] (verified)	139
Mathematica [A] (verified)	141
Maple [A] (verified)	141
Fricas [A] (verification not implemented)	142
Sympy [F]	142
Maxima [F]	142
Giac [A] (verification not implemented)	143
Mupad [F(-1)]	143

Optimal result

Integrand size = 8, antiderivative size = 78

$$\int x \sec^{-1}(a + bx) dx = -\frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \sec^{-1}(a + bx)}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx) + \frac{a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^2}$$

[Out] $-1/2*a^2*\operatorname{arcsec}(b*x+a)/b^2+1/2*x^2*\operatorname{arcsec}(b*x+a)+a*\operatorname{arctanh}((1-1/(b*x+a)^2)^{(1/2)})/b^2-1/2*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5366, 4511, 3858, 3855, 3852, 8}

$$\int x \sec^{-1}(a + bx) dx = -\frac{a^2 \sec^{-1}(a + bx)}{2b^2} + \frac{a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^2} - \frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx)$$

[In] $\operatorname{Int}[x*\operatorname{ArcSec}[a + b*x], x]$

[Out] $-1/2*((a + b*x)*\operatorname{Sqrt}[1 - (a + b*x)^{-2}])/b^2 - (a^2*\operatorname{ArcSec}[a + b*x])/(2*b^2) + (x^2*\operatorname{ArcSec}[a + b*x])/2 + (a*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - (a + b*x)^{-2}]])/b^2$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3852

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rule 3855

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rule 3858

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(2), x_Symbol] := Simp[a^2*x, x] + (Dist[2*a*b, Int[Csc[c + d*x], x], x] + Dist[b^2, Int[Csc[c + d*x]^2, x], x]) /; FreeQ[{a, b, c, d}, x]`

Rule 4511

`Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^(m+1)*((a + b*Sec[c + d*x])^(n+1)/(b*d*(n+1))), x] - Dist[f*(m/(b*d*(n+1))), Int[(e + f*x)^(m-1)*(a + b*Sec[c + d*x])^(n+1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]`

Rule 5366

`Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[1/d^(m+1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int x \sec(x)(-a + \sec(x)) \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
 &= \frac{1}{2}x^2 \sec^{-1}(a + bx) - \frac{\text{Subst}\left(\int (-a + \sec(x))^2 dx, x, \sec^{-1}(a + bx)\right)}{2b^2} \\
 &= -\frac{a^2 \sec^{-1}(a + bx)}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx) - \frac{\text{Subst}\left(\int \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{2b^2} \\
 &\quad + \frac{a \text{Subst}\left(\int \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{a^2 \sec^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a+bx) + \frac{a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^2} \\
&\quad + \frac{\operatorname{Subst}\left(\int 1 dx, x, -\left((a+bx)\sqrt{1 - \frac{1}{(a+bx)^2}}\right)\right)}{2b^2} \\
&= -\frac{(a+bx)\sqrt{1 - \frac{1}{(a+bx)^2}}}{2b^2} - \frac{a^2 \sec^{-1}(a+bx)}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a+bx) + \frac{a \operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 110, normalized size of antiderivative = 1.41

$$\begin{aligned}
&\int x \sec^{-1}(a+bx) dx \\
&= \frac{-\left((a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right) + b^2x^2 \sec^{-1}(a+bx) + a^2 \arcsin\left(\frac{1}{a+bx}\right) + 2a \log\left((a+bx)\left(1 + \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)\right)}{2b^2}
\end{aligned}$$

[In] Integrate[x*ArcSec[a + b*x],x]

[Out] $\frac{-((a + b*x)*\operatorname{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]) + b^2*x^2*\operatorname{ArcSec}[a + b*x] + a^2*\operatorname{ArcSin}[(a + b*x)^{-1}] + 2*a*\operatorname{Log}[(a + b*x)*(1 + \operatorname{Sqrt}[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]])]}{(2*b^2)}$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.38

method	result
derivativedivides	$\frac{\frac{\operatorname{arcsec}(bx+a)(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a) + \frac{\sqrt{(bx+a)^2-1}\left(2a \ln\left(bx+a+\sqrt{(bx+a)^2-1}\right) - \sqrt{(bx+a)^2-1}\right)}{2(bx+a)\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}}}{b^2}$
default	$\frac{\frac{\operatorname{arcsec}(bx+a)(bx+a)^2}{2} - \operatorname{arcsec}(bx+a)a(bx+a) + \frac{\sqrt{(bx+a)^2-1}\left(2a \ln\left(bx+a+\sqrt{(bx+a)^2-1}\right) - \sqrt{(bx+a)^2-1}\right)}{2(bx+a)\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}}}{b^2}$
parts	$\frac{x^2 \operatorname{arcsec}(bx+a)}{2} + \frac{\sqrt{b^2x^2+2abx+a^2-1}\left(a^2 \operatorname{arctan}\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right)\sqrt{b^2} + 2a \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2-1}\sqrt{b^2}+a}{\sqrt{b^2}}\right)\right)}{2b^2\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}}(bx+a)\sqrt{b^2}}$

[In] int(x*arcsec(b*x+a),x,method=_RETURNVERBOSE)

[Out] $\frac{1}{b^2}*\left(\frac{1}{2}*\operatorname{arcsec}(b*x+a)*(b*x+a)^2-\operatorname{arcsec}(b*x+a)*a*(b*x+a)+\frac{1}{2}*((b*x+a)^2-1)^{(1/2)}*(2*a*\ln(b*x+a+((b*x+a)^2-1)^{(1/2)})-((b*x+a)^2-1)^{(1/2)})/(b*x+a)/(((b*x+a)^2-1)/(b*x+a)^2)^{(1/2)}\right)$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.33

$$\int x \sec^{-1}(a + bx) dx$$

$$= \frac{b^2 x^2 \operatorname{arcsec}(bx + a) - 2a^2 \arctan(-bx - a + \sqrt{b^2 x^2 + 2abx + a^2 - 1}) - 2a \log(-bx - a + \sqrt{b^2 x^2 + 2abx + a^2 - 1})}{2b^2}$$

[In] integrate(x*arcsec(b*x+a),x, algorithm="fricas")

```
[Out] 1/2*(b^2*x^2*arcsec(b*x + a) - 2*a^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - 2*a*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/b^2
```

Sympy [F]

$$\int x \sec^{-1}(a + bx) dx = \int x \operatorname{asec}(a + bx) dx$$

[In] integrate(x*asec(b*x+a),x)

[Out] Integral(x*asec(a + b*x), x)

Maxima [F]

$$\int x \sec^{-1}(a + bx) dx = \int x \operatorname{arcsec}(bx + a) dx$$

[In] integrate(x*arcsec(b*x+a),x, algorithm="maxima")

```
[Out] 1/2*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) - integrate(1/2*(b^2*x^3 + a*b*x^2)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^2 + 2*a*b*x + a^2 + (b^2*x^2 + 2*a*b*x + a^2 - 1)*e^(log(b*x + a + 1) + log(b*x + a - 1)) - 1), x)
```

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.71

$$\int x \sec^{-1}(a + bx) dx =$$

$$-\frac{1}{4} b \left(\frac{2 (bx + a)^2 \left(\frac{2a}{bx+a} - 1 \right) \arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{b^3} + \frac{(bx + a) \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1 \right) + 4a \log \left(-\right)}{b^3} \right)$$

[In] integrate(x*arcsec(b*x+a),x, algorithm="giac")

```
[Out] -1/4*b*(2*(b*x + a)^2*(2*a/(b*x + a) - 1)*arccos(-1/((b*x + a)*(a/(b*x + a)
- 1) - a))/b^3 + ((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + 4*a*log(-(sqrt
t(-1/(b*x + a)^2 + 1) - 1)*abs(b*x + a)) - 1/((b*x + a)*(sqrt(-1/(b*x + a)^
2 + 1) - 1))))/b^3)
```

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx) dx = \int x \arccos \left(\frac{1}{a + bx} \right) dx$$

[In] int(x*acos(1/(a + b*x)),x)

[Out] int(x*acos(1/(a + b*x)), x)

3.22 $\int \sec^{-1}(a + bx) dx$

Optimal result	144
Rubi [A] (verified)	144
Mathematica [C] (verified)	146
Maple [A] (verified)	146
Fricas [B] (verification not implemented)	147
Sympy [F]	147
Maxima [A] (verification not implemented)	147
Giac [B] (verification not implemented)	148
Mupad [B] (verification not implemented)	148

Optimal result

Integrand size = 6, antiderivative size = 37

$$\int \sec^{-1}(a + bx) dx = \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b}$$

[Out] (b*x+a)*arcsec(b*x+a)/b-arctanh((1-1/(b*x+a)^2)^(1/2))/b

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5358, 379, 272, 65, 212}

$$\int \sec^{-1}(a + bx) dx = \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b}$$

[In] Int[ArcSec[a + b*x], x]

[Out] ((a + b*x)*ArcSec[a + b*x])/b - ArcTanh[Sqrt[1 - (a + b*x)^(-2)]]/b

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_)*((a_) + (b_)*(v_)^(n_))^(p_), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 5358

Int[ArcSec[(c_) + (d_)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \int \frac{1}{(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}} dx \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2}}} dx, x, a + bx\right)}{b} \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)}{b} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1 - xx}} dx, x, \frac{1}{(a+bx)^2}\right)}{2b} \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\text{Subst}\left(\int \frac{1}{1 - x^2} dx, x, \sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b} \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)}{b} - \frac{\text{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.99 (sec) , antiderivative size = 468, normalized size of antiderivative = 12.65

$$\int \sec^{-1}(a + bx) dx = x \sec^{-1}(a + bx) + \frac{(a + bx) \sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}} \left(\sqrt[4]{-1} (-i + \sqrt{-1+a^2}) \sqrt{2i - ia^2 + 2\sqrt{-1+a^2}} \arctan \left(\frac{(-1)^{3/4} \sqrt{2i - ia^2 + 2\sqrt{-1+a^2}}}{a\sqrt{-1+a^2} - a\sqrt{-1+a^2+2abx+b^2x^2}} \right) \right)}{b}$$

[In] Integrate[ArcSec[a + b*x],x]

[Out] x*ArcSec[a + b*x] + ((a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + a^2])*Sqrt[2*I - I*a^2 + 2*Sqrt[-1 + a^2]]*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*a^2 + 2*Sqrt[-1 + a^2]]*b*x)/(a*Sqrt[-1 + a^2] - a*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])]) + (-1)^(3/4)*(I + Sqrt[-1 + a^2])*Sqrt[-2*I + I*a^2 + 2*Sqrt[-1 + a^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*a^2 + 2*Sqrt[-1 + a^2]]*b*x)/(a*Sqrt[-1 + a^2] - a*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])]) + a*(a*ArcTan[(Sqrt[-1 + a^2]*b^2*x^2)/(a^4 + a^3*b*x + b^2*x^2 - a^2*(1 + Sqrt[-1 + a^2])*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])]) - Log[Sqrt[-1 + a^2] - b*x - Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2]] + Log[b^2*(Sqrt[-1 + a^2] + b*x - Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])])]/(a*b*Sqrt[-1 + a^2 + 2*a*b*x + b^2*x^2])

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.22

method	result
derivativedivides	$\frac{(bx+a) \operatorname{arcsec}(bx+a) - \ln\left(\frac{bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}}{b}\right)}{b}$
default	$\frac{(bx+a) \operatorname{arcsec}(bx+a) - \ln\left(\frac{bx+a+(bx+a)\sqrt{1-\frac{1}{(bx+a)^2}}}{b}\right)}{b}$
parts	$x \operatorname{arcsec}(bx+a) - \frac{\sqrt{b^2x^2+2abx+a^2-1} \left(a \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{b^2} + \ln\left(\frac{b^2x+\sqrt{b^2x^2+2abx+a^2-1}\sqrt{b^2}}{\sqrt{b^2}}\right) \right)}{b\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}}(bx+a)\sqrt{b^2}}$

[In] int(arcsec(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/b*((b*x+a)*arcsec(b*x+a)-ln(b*x+a+(b*x+a)*(1-1/(b*x+a)^2)^(1/2)))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(35) = 70.

Time = 0.30 (sec) , antiderivative size = 73, normalized size of antiderivative = 1.97

$$\int \sec^{-1}(a + bx) dx$$

$$= \frac{bx \operatorname{arcsec}(bx + a) + 2a \arctan(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1}) + \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 - 1})}{b}$$

[In] integrate(arcsec(b*x+a),x, algorithm="fricas")

[Out] (b*x*arcsec(b*x + a) + 2*a*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)))/b

Sympy [F]

$$\int \sec^{-1}(a + bx) dx = \int \operatorname{asec}(a + bx) dx$$

[In] integrate(asec(b*x+a),x)

[Out] Integral(asec(a + b*x), x)

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.49

$$\int \sec^{-1}(a + bx) dx$$

$$= \frac{2(bx + a) \operatorname{arcsec}(bx + a) - \log\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right)}{2b}$$

[In] integrate(arcsec(b*x+a),x, algorithm="maxima")

[Out] 1/2*(2*(b*x + a)*arcsec(b*x + a) - log(sqrt(-1/(b*x + a)^2 + 1) + 1) + log(-sqrt(-1/(b*x + a)^2 + 1) + 1))/b

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(35) = 70.

Time = 0.30 (sec) , antiderivative size = 82, normalized size of antiderivative = 2.22

$$\int \sec^{-1}(a + bx) dx$$

$$= \frac{1}{2} b \left(\frac{2(bx + a) \arccos\left(-\frac{1}{(bx+a)\left(\frac{a}{bx+a}-1\right)-a}\right)}{b^2} - \frac{\log\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx+a)^2} + 1} + 1\right)}{b^2} \right)$$

[In] integrate(arcsec(b*x+a),x, algorithm="giac")

[Out] 1/2*b*(2*(b*x + a)*arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/b^2 - (log(sqrt(-1/(b*x + a)^2 + 1) + 1) - log(-sqrt(-1/(b*x + a)^2 + 1) + 1))/b^2)

Mupad [B] (verification not implemented)

Time = 1.04 (sec) , antiderivative size = 35, normalized size of antiderivative = 0.95

$$\int \sec^{-1}(a + bx) dx = -\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1-\frac{1}{(a+bx)^2}}}\right) - \operatorname{acos}\left(\frac{1}{a+bx}\right) (a + bx)}{b}$$

[In] int(acos(1/(a + b*x)),x)

[Out] -(atanh(1/(1 - 1/(a + b*x)^2)^(1/2)) - acos(1/(a + b*x))*(a + b*x))/b

3.23 $\int \frac{\sec^{-1}(a+bx)}{x} dx$

Optimal result	149
Rubi [A] (verified)	150
Mathematica [A] (verified)	153
Maple [A] (verified)	154
Fricas [F]	154
Sympy [F]	154
Maxima [F]	155
Giac [F]	155
Mupad [F(-1)]	155

Optimal result

Integrand size = 10, antiderivative size = 200

$$\int \frac{\sec^{-1}(a+bx)}{x} dx = \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + \sec^{-1}(a+bx) \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) - \sec^{-1}(a+bx) \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) - i \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) - i \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) + \frac{1}{2} i \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right)$$

```
[Out] -arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+1/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-I*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))-I*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5366, 4647, 4626, 3800, 2221, 2317, 2438, 4616}

$$\int \frac{\sec^{-1}(a+bx)}{x} dx = -i \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) - i \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2} + 1}\right) \\ + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\ + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2} + 1}\right) \\ + \frac{1}{2}i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) \\ - \sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)$$

[In] Int[ArcSec[a + b*x]/x,x]

[Out] ArcSec[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + ArcSec[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] - I*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] - I*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + (I/2)*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])]

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3800

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 4616

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Sin[(c_.) + (d_.)*(x_)])/(Cos[(c_.) + (d_.)
*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[I*((e + f*x)^(m + 1)/(b*f*(m + 1)))
, x] + (-Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a - Rt[a^2 - b^2, 2] + b
*E^(I*(c + d*x))), x], x] - Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a +
Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x))), x], x]) /; FreeQ[{a, b, c, d, e, f}
, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

Rule 4626

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*Tan[(c_.) + (d_.)*(x_)])^(n_.)/(Cos[(c_.) +
(d_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[1/a, Int[(e + f*x)^m*Tan[c +
d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sin[c + d*x]*(Tan[c + d*x])^(n -
1)/(a + b*Cos[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m,
0] && IGtQ[n, 0]
```

Rule 4647

```
Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_)]^(p_.))/(a_ + (b_.)*Sec[(c_.) + (d_.)*(x_)]), x_Symbol] := In
t[(e + f*x)^m*Cos[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cos[c + d*x]))
, x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m
, n, p]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x \sec(x) \tan(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= \text{Subst}\left(\int \frac{x \tan(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= a \text{Subst}\left(\int \frac{x \sin(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) + \text{Subst}\left(\int x \tan(x) dx, x, \sec^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= -\left(2i\text{Subst}\left(\int \frac{e^{2ix}}{1+e^{2ix}} dx, x, \sec^{-1}(a+bx)\right)\right) \\
&\quad - (ia)\text{Subst}\left(\int \frac{e^{ix}}{1-\sqrt{1-a^2}-ae^{ix}} dx, x, \sec^{-1}(a+bx)\right) \\
&\quad - (ia)\text{Subst}\left(\int \frac{e^{ix}}{1+\sqrt{1-a^2}-ae^{ix}} dx, x, \sec^{-1}(a+bx)\right) \\
&= \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx) \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - \text{Subst}\left(\int \log\left(1 - \frac{ae^{ix}}{1-\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right) \\
&\quad - \text{Subst}\left(\int \log\left(1 - \frac{ae^{ix}}{1+\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right) \\
&\quad + \text{Subst}\left(\int \log(1+e^{2ix}) dx, x, \sec^{-1}(a+bx)\right) \\
&= \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx) \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - \frac{1}{2}i\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad + i\text{Subst}\left(\int \frac{\log\left(1 - \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right) \\
&\quad + i\text{Subst}\left(\int \frac{\log\left(1 - \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right) \\
&= \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx) \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) - i\text{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
&\quad - i\text{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right) + \frac{1}{2}i\text{PolyLog}\left(2, -e^{2i\sec^{-1}(a+bx)}\right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 284, normalized size of antiderivative = 1.42

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)}{x} dx = & -4i \arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right) \arctan\left(\frac{(1+a)\tan\left(\frac{1}{2}\sec^{-1}(a+bx)\right)}{\sqrt{1-a^2}}\right) \\
& + \left(\sec^{-1}(a+bx) - 2\arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right)\right) \log\left(1 + \frac{(-1+\sqrt{1-a^2})e^{i\sec^{-1}(a+bx)}}{a}\right) + \left(\sec^{-1}(a+bx) \right. \\
& \left. + 2\arcsin\left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}}\right)\right) \log\left(1 - \frac{(1+\sqrt{1-a^2})e^{i\sec^{-1}(a+bx)}}{a}\right) \\
& - \sec^{-1}(a+bx) \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
& - i\left(\text{PolyLog}\left(2, -\frac{(-1+\sqrt{1-a^2})e^{i\sec^{-1}(a+bx)}}{a}\right) \right. \\
& \left. + \text{PolyLog}\left(2, \frac{(1+\sqrt{1-a^2})e^{i\sec^{-1}(a+bx)}}{a}\right)\right) \\
& + \frac{1}{2}i \text{PolyLog}\left(2, -e^{2i\sec^{-1}(a+bx)}\right)
\end{aligned}$$

[In] Integrate[ArcSec[a + b*x]/x,x]

```

[Out] (-4*I)*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*ArcTan[((1 + a)*Tan[ArcSec[a + b*x]
/2])/Sqrt[1 - a^2]] + (ArcSec[a + b*x] - 2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]
)*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + (ArcSec[a + b*x]
+ 2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]])*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*Arc
Sec[a + b*x]))/a] - ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] -
I*(PolyLog[2, -(((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a)] + PolyLog[
2, ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a]) + (I/2)*PolyLog[2, -E^((
2*I)*ArcSec[a + b*x])]

```

Maple [A] (verified)

Time = 1.64 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.87

method	result
derivativedivides	$\operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1}+1}{1+\sqrt{-a^2+1}} \right) + \operatorname{arcsec}(bx+a) \ln \left(\frac{a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) - \sqrt{-a^2+1}-1}{-1+\sqrt{-a^2+1}} \right)$
default	$\operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1}+1}{1+\sqrt{-a^2+1}} \right) + \operatorname{arcsec}(bx+a) \ln \left(\frac{a \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) - \sqrt{-a^2+1}-1}{-1+\sqrt{-a^2+1}} \right)$

[In] int(arcsec(b*x+a)/x,x,method=_RETURNVERBOSE)

```
[Out] arcsec(b*x+a)*ln((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))+arcsec(b*x+a)*ln((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))-arcsec(b*x+a)*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-arcsec(b*x+a)*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-I*dilog((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))-I*dilog((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))
```

Fricas [F]

$$\int \frac{\sec^{-1}(a+bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx+a)}{x} dx$$

[In] integrate(arcsec(b*x+a)/x,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)/x, x)

Sympy [F]

$$\int \frac{\sec^{-1}(a+bx)}{x} dx = \int \frac{\operatorname{asec}(a+bx)}{x} dx$$

[In] integrate(asec(b*x+a)/x,x)

[Out] Integral(asec(a + b*x)/x, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x} dx$$

[In] integrate(arcsec(b*x+a)/x,x, algorithm="maxima")

[Out] integrate(arcsec(b*x + a)/x, x)

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x} dx$$

[In] integrate(arcsec(b*x+a)/x,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x} dx = \int \frac{\operatorname{acos}\left(\frac{1}{a+bx}\right)}{x} dx$$

[In] int(acos(1/(a + b*x))/x,x)

[Out] int(acos(1/(a + b*x))/x, x)

3.24 $\int \frac{\sec^{-1}(a+bx)}{x^2} dx$

Optimal result	156
Rubi [A] (verified)	156
Mathematica [C] (verified)	158
Maple [B] (verified)	158
Fricas [B] (verification not implemented)	159
Sympy [F]	159
Maxima [F]	160
Giac [A] (verification not implemented)	160
Mupad [F(-1)]	160

Optimal result

Integrand size = 10, antiderivative size = 70

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx = -\frac{b \sec^{-1}(a+bx)}{a} - \frac{\sec^{-1}(a+bx)}{x} + \frac{2b \arctan\left(\frac{\sqrt{1+a} \tan\left(\frac{1}{2} \sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}}$$

[Out] $-b*\text{arcsec}(b*x+a)/a-\text{arcsec}(b*x+a)/x+2*b*\text{arctan}((1+a)^{(1/2)}*\text{tan}(1/2*\text{arcsec}(b*x+a)))/(1-a)^{(1/2)}/a/(-a^2+1)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5366, 4511, 3868, 2738, 211}

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx = \frac{2b \arctan\left(\frac{\sqrt{a+1} \tan\left(\frac{1}{2} \sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{a\sqrt{1-a^2}} - \frac{b \sec^{-1}(a+bx)}{a} - \frac{\sec^{-1}(a+bx)}{x}$$

[In] Int[ArcSec[a + b*x]/x^2,x]

[Out] $-((b*\text{ArcSec}[a + b*x])/a) - \text{ArcSec}[a + b*x]/x + (2*b*\text{ArcTan}[(\text{Sqrt}[1 + a]*\text{Tan}[\text{ArcSec}[a + b*x]/2])/(\text{Sqrt}[1 - a])]/(a*\text{Sqrt}[1 - a^2]))$

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 2738

```
Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (
a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

Rule 3868

```
Int[(csc[(c_) + (d_)*(x_)]*(b_) + (a_))^-1, x_Symbol] := Simp[x/a, x]
- Dist[1/a, Int[1/(1 + (a/b)*Sin[c + d*x]), x], x] /; FreeQ[{a, b, c, d}, x]
] && NeQ[a^2 - b^2, 0]
```

Rule 4511

```
Int[((e_) + (f_)*(x_))^(m_)*Sec[(c_) + (d_)*(x_)]*((a_) + (b_)*Sec[(c
_) + (d_)*(x_)])^(n_)*Tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_) + ArcSec[(c_) + (d_)*(x_)]*(b_))^(p_)*((e_) + (f_)*(x_))^(m
_), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= b \text{Subst} \left(\int \frac{x \sec(x) \tan(x)}{(-a + \sec(x))^2} dx, x, \sec^{-1}(a + bx) \right) \\
&= -\frac{\sec^{-1}(a + bx)}{x} + b \text{Subst} \left(\int \frac{1}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right) \\
&= -\frac{b \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{x} + \frac{b \text{Subst} \left(\int \frac{1}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx) \right)}{a} \\
&= -\frac{b \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{x} + \frac{(2b) \text{Subst} \left(\int \frac{1}{1 - a + (1+a)x^2} dx, x, \tan \left(\frac{1}{2} \sec^{-1}(a + bx) \right) \right)}{a} \\
&= -\frac{b \sec^{-1}(a + bx)}{a} - \frac{\sec^{-1}(a + bx)}{x} + \frac{2b \arctan \left(\frac{\sqrt{1+a} \tan \left(\frac{1}{2} \sec^{-1}(a + bx) \right)}{\sqrt{1-a}} \right)}{a \sqrt{1 - a^2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.21 (sec) , antiderivative size = 112, normalized size of antiderivative = 1.60

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx = -\frac{\sec^{-1}(a+bx)}{x} + \frac{b \left(\arcsin\left(\frac{1}{a+bx}\right) - \frac{i \log\left(\frac{2\left(\frac{ia(-1+a^2+abx)}{\sqrt{1-a^2}} + a(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{bx}\right)}{\sqrt{1-a^2}} \right)}{a}$$

[In] Integrate[ArcSec[a + b*x]/x^2,x]

[Out] -(ArcSec[a + b*x]/x) + (b*(ArcSin[(a + b*x)^(-1)] - (I*Log[(2*((I*a*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] + a*(a + b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/(b*x)])/Sqrt[1 - a^2]))/a

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 125 vs. 2(62) = 124.

Time = 0.72 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.80

method	result
derivativedivides	$b \left(-\frac{\operatorname{arcsec}(bx+a)}{bx} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2\sqrt{a^2-1}\sqrt{(bx+a)^2-1}+2(bx+a)a-2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}} \right)$
default	$b \left(-\frac{\operatorname{arcsec}(bx+a)}{bx} + \frac{\sqrt{(bx+a)^2-1} \left(\arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2\sqrt{a^2-1}\sqrt{(bx+a)^2-1}+2(bx+a)a-2}{bx}\right) \right)}{\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}} \right)$
parts	$-\frac{\operatorname{arcsec}(bx+a)}{x} + \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left(\arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) \sqrt{a^2-1} - \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}\sqrt{b^2x^2+2abx+a^2-1}}{x}\right) \right)}{\sqrt{\frac{b^2x^2+2abx+a^2-1}{(bx+a)^2}} (bx+a)a\sqrt{a^2-1}}$

[In] int(arcsec(b*x+a)/x^2,x,method=_RETURNVERBOSE)

[Out] b*(-1/b/x*arcsec(b*x+a)+((b*x+a)^2-1)^(1/2)*(arctan(1/((b*x+a)^2-1)^(1/2))*(a^2-1)^(1/2)-ln(2*((a^2-1)^(1/2)*((b*x+a)^2-1)^(1/2)+(b*x+a)*a-1)/b/x))/((b*x+a)^2-1)/(b*x+a)^2)^(1/2)/(b*x+a)/a/(a^2-1)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(62) = 124$.

Time = 0.32 (sec) , antiderivative size = 281, normalized size of antiderivative = 4.01

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx$$

$$= \frac{\left[\begin{aligned} &2(a^2-1)bx \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) - \sqrt{a^2-1}bx \log\left(\frac{a^2bx+a^3+\sqrt{b^2x^2+2abx+a^2-1}(a^2-1)}{(a^3-a)x}\right) \right. \\ &\left. - 2(a^2-1)bx \arctan(-bx-a+\sqrt{b^2x^2+2abx+a^2-1}) - 2\sqrt{-a^2+1}bx \arctan\left(-\frac{\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}}{a^2-1}\right) \right]}{(a^3-a)x}$$

[In] integrate(arcsec(b*x+a)/x^2,x, algorithm="fricas")

[Out] $[-(2*(a^2-1)*b*x*\arctan(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1})-\sqrt{a^2-1}*b*x*\log((a^2*b*x+a^3+\sqrt{b^2*x^2+2*a*b*x+a^2-1})*(a^2-\sqrt{a^2-1}*a-1)-(a*b*x+a^2-1)*\sqrt{a^2-1}-a)/x)+(a^3-a)*\text{arcsec}(b*x+a))/((a^3-a)*x), -(2*(a^2-1)*b*x*\arctan(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2-1})-2*\sqrt{-a^2+1}*b*x*\arctan(-(\sqrt{-a^2+1}*b*x-\sqrt{b^2*x^2+2*a*b*x+a^2-1})*\sqrt{-a^2+1}))/((a^2-1))+(a^3-a)*\text{arcsec}(b*x+a))/((a^3-a)*x)]$

Sympy [F]

$$\int \frac{\sec^{-1}(a+bx)}{x^2} dx = \int \frac{\text{asec}(a+bx)}{x^2} dx$$

[In] integrate(asec(b*x+a)/x**2,x)

[Out] Integral(asec(a+b*x)/x**2, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x^2} dx$$

[In] integrate(arcsec(b*x+a)/x^2,x, algorithm="maxima")

[Out] (x*integrate((b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/
(b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x + (b^2*x^3 + 2*a*b*x^2 + (a^2 - 1)*x)*e^
(log(b*x + a + 1) + log(b*x + a - 1))), x) - arctan(sqrt(b*x + a + 1)*sqrt(
b*x + a - 1)))/x

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.34

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx$$

$$= b \left(\frac{2 \arctan \left(\frac{(bx+a) \left(\sqrt{\frac{1}{(bx+a)^2} + 1} - 1 \right) + a}{\sqrt{-a^2 + 1}} \right)}{\sqrt{-a^2 + 1} a} + \frac{\arccos \left(-\frac{1}{(bx+a) \left(\frac{a}{bx+a} - 1 \right) - a} \right)}{a \left(\frac{a}{bx+a} - 1 \right)} \right)$$

[In] integrate(arcsec(b*x+a)/x^2,x, algorithm="giac")

[Out] b*(2*arctan(((b*x + a)*(sqrt(-1/(b*x + a)^2 + 1) - 1) + a)/sqrt(-a^2 + 1))/
(sqrt(-a^2 + 1)*a) + arccos(-1/((b*x + a)*(a/(b*x + a) - 1) - a))/(a*(a/(b*
x + a) - 1)))

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)}{x^2} dx = \int \frac{\operatorname{acos}\left(\frac{1}{a+bx}\right)}{x^2} dx$$

[In] int(acos(1/(a + b*x))/x^2,x)

[Out] int(acos(1/(a + b*x))/x^2, x)

3.25 $\int \frac{\sec^{-1}(a+bx)}{x^3} dx$

Optimal result	161
Rubi [A] (verified)	161
Mathematica [C] (verified)	164
Maple [B] (verified)	164
Fricas [A] (verification not implemented)	165
Sympy [F]	165
Maxima [F]	166
Giac [B] (verification not implemented)	166
Mupad [F(-1)]	166

Optimal result

Integrand size = 10, antiderivative size = 125

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)x} + \frac{b^2 \sec^{-1}(a+bx)}{2a^2} - \frac{\sec^{-1}(a+bx)}{2x^2} - \frac{(1-2a^2)b^2 \arctan\left(\frac{\sqrt{1+a}\tan(\frac{1}{2}\sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}}$$

[Out] $1/2*b^2*arcsec(b*x+a)/a^2-1/2*arcsec(b*x+a)/x^2-(-2*a^2+1)*b^2*arctan((1+a)^{(1/2)*tan(1/2*arcsec(b*x+a))/(1-a)^{(1/2)})/a^2/(-a^2+1)^{(3/2)}+1/2*b*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/a/(-a^2+1)/x$

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {5366, 4511, 3870, 4004, 3916, 2738, 211}

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = -\frac{(1-2a^2)b^2 \arctan\left(\frac{\sqrt{a+1}\tan(\frac{1}{2}\sec^{-1}(a+bx))}{\sqrt{1-a}}\right)}{a^2(1-a^2)^{3/2}} + \frac{b^2 \sec^{-1}(a+bx)}{2a^2} + \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{2a(1-a^2)x} - \frac{\sec^{-1}(a+bx)}{2x^2}$$

[In] Int[ArcSec[a + b*x]/x^3,x]

[Out] $(b*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}])/(2*a*(1-a^2)*x) + (b^2*\text{ArcSec}[a+b*x])/(2*a^2) - \text{ArcSec}[a+b*x]/(2*x^2) - ((1-2*a^2)*b^2*\text{ArcTan}[(\text{Sqrt}[1+a]*\text{Tan}[\text{ArcSec}[a+b*x]/2])/\text{Sqrt}[1-a]])/(a^2*(1-a^2)^{(3/2)})$

Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 2738

```
Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3870

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] := Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]
```

Rule 3916

```
Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]
```

Rule 4004

```
Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
```

f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
\text{integral} &= b^2 \text{Subst} \left(\int \frac{x \sec(x) \tan(x)}{(-a + \sec(x))^3} dx, x, \sec^{-1}(a + bx) \right) \\
&= -\frac{\sec^{-1}(a + bx)}{2x^2} + \frac{1}{2} b^2 \text{Subst} \left(\int \frac{1}{(-a + \sec(x))^2} dx, x, \sec^{-1}(a + bx) \right) \\
&= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1 - a^2)x} - \frac{\sec^{-1}(a + bx)}{2x^2} - \frac{b^2 \text{Subst} \left(\int \frac{1 - a^2 - a \sec(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right)}{2a(1 - a^2)} \\
&= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1 - a^2)x} + \frac{b^2 \sec^{-1}(a + bx)}{2a^2} - \frac{\sec^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 - 2a^2)b^2) \text{Subst} \left(\int \frac{\sec(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right)}{2a^2(1 - a^2)} \\
&= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1 - a^2)x} + \frac{b^2 \sec^{-1}(a + bx)}{2a^2} - \frac{\sec^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 - 2a^2)b^2) \text{Subst} \left(\int \frac{1}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx) \right)}{2a^2(1 - a^2)} \\
&= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1 - a^2)x} + \frac{b^2 \sec^{-1}(a + bx)}{2a^2} - \frac{\sec^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{((1 - 2a^2)b^2) \text{Subst} \left(\int \frac{1}{1 - a + (1+a)x^2} dx, x, \tan \left(\frac{1}{2} \sec^{-1}(a + bx) \right) \right)}{a^2(1 - a^2)} \\
&= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{2a(1 - a^2)x} + \frac{b^2 \sec^{-1}(a + bx)}{2a^2} - \frac{\sec^{-1}(a + bx)}{2x^2} \\
&\quad - \frac{(1 - 2a^2)b^2 \arctan \left(\frac{\sqrt{1+a} \tan \left(\frac{1}{2} \sec^{-1}(a+bx) \right)}{\sqrt{1-a}} \right)}{a^2(1 - a^2)^{3/2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.71 (sec) , antiderivative size = 198, normalized size of antiderivative = 1.58

$$\int \frac{\sec^{-1}(a + bx)}{x^3} dx = \frac{\frac{bx(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}}{a(-1+a^2)} + \sec^{-1}(a + bx) + \frac{b^2x^2 \arcsin\left(\frac{1}{a+bx}\right)}{a^2} + \frac{i(-1+2a^2)b^2x^2 \log\left(\frac{4(-1+a)a^2(1+a)\left(-\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}\right) - (a+bx)}{(-1+2a^2)b^2x}\right)}{a^2(1-a^2)^{3/2}}}{2x^2}$$

[In] Integrate[ArcSec[a + b*x]/x^3,x]

[Out] $-\frac{1}{2} \left(\frac{b^2 x^2 \sqrt{-1+a^2+2abx+b^2x^2}}{(a+bx)^2} \right) / (a^2(-1+a^2)) + \text{ArcSec}[a + b*x] + \frac{b^2 x^2 \text{ArcSin}[(a + b*x)^{-1}]}{a^2} + \frac{I \left((-1 + 2a^2) b^2 x^2 \text{Log}\left[\frac{4(-1+a)a^2(1+a)\left(-\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}\right) - (a+bx)}{(-1+2a^2)b^2x}\right] \right)}{\text{Sqrt}[1 - a^2] - (a + b*x) \sqrt{-1+a^2+2abx+b^2x^2}} \right) / ((-1 + 2a^2) b^2 x) / (a^2 (1 - a^2)^{3/2}) / x^2$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 318 vs. 2(109) = 218.

Time = 0.70 (sec) , antiderivative size = 319, normalized size of antiderivative = 2.55

method	result
parts	$-\frac{\text{arcsec}(bx+a)}{2x^2} - \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left((a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) a^2bx - 2 \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}\sqrt{b^2x^2+2abx+a^2-1}}{x}\right) \right)}{2x^2}$
derivativedivides	$b^2 \left(-\frac{\text{arcsec}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2-1} \left((a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) a^3 - (a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) a^2(bx+a) \right)}{2b^2x^2} \right)$
default	$b^2 \left(-\frac{\text{arcsec}(bx+a)}{2b^2x^2} + \frac{\sqrt{(bx+a)^2-1} \left((a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) a^3 - (a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{(bx+a)^2-1}}\right) a^2(bx+a) \right)}{2b^2x^2} \right)$

[In] int(arcsec(b*x+a)/x^3,x,method=_RETURNVERBOSE)

[Out] $-\frac{1}{2} \text{arcsec}(b*x+a) / x^2 - \frac{1}{2} b \sqrt{b^2x^2+2abx+a^2-1} \left((a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) a^2bx - 2 \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}\sqrt{b^2x^2+2abx+a^2-1}}{x}\right) \right) / (a^2(1-a^2)^{3/2}) + \frac{b^2x^2 \text{ArcSin}[(a + b*x)^{-1}]}{a^2} + \frac{i(-1+2a^2)b^2x^2 \log\left(\frac{4(-1+a)a^2(1+a)\left(-\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}\right) - (a+bx)}{(-1+2a^2)b^2x}\right)}{a^2(1-a^2)^{3/2}}$

$a+3\ln(2*(a*b*x+(a^2-1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}+a^2-1)/x)*a^2*b$
 $*x-b*\ln(2*(a*b*x+(a^2-1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2-1)^{(1/2)}+a^2-1)/x)*x)/(($
 $b^2*x^2+2*a*b*x+a^2-1)/(b*x+a)^2)^{(1/2)}/(b*x+a)/a^2/(a^2-1)^{(5/2)}/x$

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 427, normalized size of antiderivative = 3.42

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx$$

$$= \left[\frac{(2a^2-1)\sqrt{a^2-1}b^2x^2 \log\left(\frac{a^2bx+a^3+\sqrt{b^2x^2+2abx+a^2-1}(a^2+\sqrt{a^2-1}a-1)+(abx+a^2-1)\sqrt{a^2-1}-a}{x}\right) + 2(a^4-2a^2+1)}{2(2a^2-1)\sqrt{-a^2+1}b^2x^2 \arctan\left(-\frac{\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}\sqrt{-a^2+1}}{a^2-1}\right) - 2(a^4-2a^2+1)b^2x^2 \arctan\left(-\frac{\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}\sqrt{-a^2+1}}{a^2-1}\right) - 2(a^4-2a^2+1)b^2x^2 \arctan\left(-\frac{\sqrt{-a^2+1}bx-\sqrt{b^2x^2+2abx+a^2-1}\sqrt{-a^2+1}}{a^2-1}\right)}$$

[In] integrate(arcsec(b*x+a)/x^3,x, algorithm="fricas")

[Out] [1/2*((2*a^2 - 1)*sqrt(a^2 - 1)*b^2*x^2*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 + sqrt(a^2 - 1)*a - 1) + (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) + 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) - (a^3 - a)*b^2*x^2 - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x - (a^6 - 2*a^4 + a^2)*arcsec(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2), -1/2*(2*(2*a^2 - 1)*sqrt(-a^2 + 1)*b^2*x^2*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) - 2*(a^4 - 2*a^2 + 1)*b^2*x^2*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (a^3 - a)*b^2*x^2 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^3 - a)*b*x + (a^6 - 2*a^4 + a^2)*arcsec(b*x + a))/((a^6 - 2*a^4 + a^2)*x^2)]

Sympy [F]

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = \int \frac{\operatorname{asec}(a+bx)}{x^3} dx$$

[In] integrate(asec(b*x+a)/x**3,x)

[Out] Integral(asec(a + b*x)/x**3, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = \int \frac{\operatorname{arcsec}(bx+a)}{x^3} dx$$

[In] integrate(arcsec(b*x+a)/x^3,x, algorithm="maxima")

[Out] $\frac{1}{2} \cdot (2 \cdot x^2 \cdot \int \frac{1}{2} \cdot (b^2 \cdot x + a \cdot b) \cdot e^{(1/2 \cdot \log(b \cdot x + a + 1) + 1/2 \cdot \log(b \cdot x + a - 1))} / (b^2 \cdot x^4 + 2 \cdot a \cdot b \cdot x^3 + (a^2 - 1) \cdot x^2 + (b^2 \cdot x^4 + 2 \cdot a \cdot b \cdot x^3 + (a^2 - 1) \cdot x^2) \cdot e^{(\log(b \cdot x + a + 1) + \log(b \cdot x + a - 1))}) dx - \arctan(\sqrt{(b \cdot x + a + 1) \cdot \sqrt{(b \cdot x + a - 1))})} / x^2$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(106) = 212.

Time = 0.32 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.73

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = -\frac{1}{2} b \left(\frac{2(2a^2b - b) \arctan\left(\frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + a}{\sqrt{-a^2 + 1}}\right)}{(a^4 - a^2)\sqrt{-a^2 + 1}} + \frac{2\left((bx+a)ab\left(\sqrt{-\frac{1}{(bx+a)^2}}\right)\right)}{\left((bx+a)^2\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)^2 + 2(bx+a)a\right)} \right)$$

[In] integrate(arcsec(b*x+a)/x^3,x, algorithm="giac")

[Out] $-\frac{1}{2} \cdot b \cdot (2 \cdot (2 \cdot a^2 \cdot b - b) \cdot \arctan\left(\frac{(b \cdot x + a) \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1) + a}{\sqrt{-a^2 + 1}}\right) / ((a^4 - a^2) \cdot \sqrt{-a^2 + 1}) + 2 \cdot ((b \cdot x + a) \cdot a \cdot b \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1) + b) / (((b \cdot x + a)^2 \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1)^2 + 2 \cdot (b \cdot x + a) \cdot a \cdot (\sqrt{-1/(b \cdot x + a)^2 + 1} - 1) + 1) \cdot (a^3 - a)) + (2 \cdot a \cdot b / (b \cdot x + a) - b) \cdot \arccos(-1 / ((b \cdot x + a) \cdot (a / (b \cdot x + a) - 1) - a)) / (a^2 \cdot (a / (b \cdot x + a) - 1)^2))$

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a+bx)}{x^3} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x^3} dx$$

[In] int(arccos(1/(a + b*x))/x^3,x)

[Out] int(arccos(1/(a + b*x))/x^3, x)

3.26 $\int \frac{\sec^{-1}(a+bx)}{x^4} dx$

Optimal result	167
Rubi [A] (verified)	167
Mathematica [C] (verified)	170
Maple [B] (verified)	171
Fricas [A] (verification not implemented)	171
Sympy [F]	172
Maxima [F]	172
Giac [B] (verification not implemented)	173
Mupad [F(-1)]	173

Optimal result

Integrand size = 10, antiderivative size = 181

$$\int \frac{\sec^{-1}(a+bx)}{x^4} dx = \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\sec^{-1}(a+bx)}{3a^3} - \frac{\sec^{-1}(a+bx)}{3x^3} + \frac{(2-5a^2+6a^4)b^3\arctan\left(\frac{\sqrt{1+a}\tan\left(\frac{1}{2}\sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}$$

[Out] $-1/3*b^3*arcsec(b*x+a)/a^3-1/3*arcsec(b*x+a)/x^3+1/3*(6*a^4-5*a^2+2)*b^3*arctan((1+a)^{(1/2)}*\tan(1/2*arcsec(b*x+a))/(1-a)^{(1/2)})/a^3/(-a^2+1)^{(5/2)}+1/6*b*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/a/(-a^2+1)/x^2-1/6*(-5*a^2+2)*b^2*(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/a^2/(-a^2+1)^2/x$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 181, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used = {5366, 4511, 3870, 4145, 4004, 3916, 2738, 211}

$$\int \frac{\sec^{-1}(a+bx)}{x^4} dx = -\frac{b^3\sec^{-1}(a+bx)}{3a^3} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} + \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} + \frac{(6a^4-5a^2+2)b^3\arctan\left(\frac{\sqrt{a+1}\tan\left(\frac{1}{2}\sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}} - \frac{\sec^{-1}(a+bx)}{3x^3}$$

[In] Int[ArcSec[a + b*x]/x^4,x]

[Out] (b*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)])/(6*a*(1 - a^2)*x^2) - ((2 - 5*a^2)*b^2*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)])/(6*a^2*(1 - a^2)^2*x) - (b^3*ArcSec[a + b*x])/(3*a^3) - ArcSec[a + b*x]/(3*x^3) + ((2 - 5*a^2 + 6*a^4)*b^3*ArcTan[(Sqrt[1 + a]*Tan[ArcSec[a + b*x]/2])/Sqrt[1 - a]])/(3*a^3*(1 - a^2)^(5/2))

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 2738

Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3870

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] := Simp[b^2*Cot[c + d*x]*((a + b*Csc[c + d*x])^(n + 1)/(a*d*(n + 1)*(a^2 - b^2))), x] + Dist[1/(a*(n + 1)*(a^2 - b^2)), Int[(a + b*Csc[c + d*x])^(n + 1)*Simp[(a^2 - b^2)*(n + 1) - a*b*(n + 1)*Csc[c + d*x] + b^2*(n + 2)*Csc[c + d*x]^2, x], x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3916

Int[csc[(e_.) + (f_.)*(x_)]/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Dist[1/b, Int[1/(1 + (a/b)*Sin[e + f*x]), x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a^2 - b^2, 0]

Rule 4004

Int[(csc[(e_.) + (f_.)*(x_)]*(d_.) + (c_))/(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_)), x_Symbol] := Simp[c*(x/a), x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]

Rule 4145

Int[((A_.) + csc[(e_.) + (f_.)*(x_)]*(B_.) + csc[(e_.) + (f_.)*(x_)]^2*(C_.))*(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_))^(m_), x_Symbol] := Simp[(A*b^2 - a*b*B + a^2*C)*Cot[e + f*x]*((a + b*Csc[e + f*x])^(m + 1)/(a*f*(m + 1)*(a^2 - b^2))), x] + Dist[1/(a*(m + 1)*(a^2 - b^2)), Int[(a + b*Csc[e + f*x])^(m

+ 1)*Simp[A*(a^2 - b^2)*(m + 1) - a*(A*b - a*B + b*C)*(m + 1)*Csc[e + f*x]
 + (A*b^2 - a*b*B + a^2*C)*(m + 2)*Csc[e + f*x]^2, x], x] /; FreeQ[{a,
 b, e, f, A, B, C}, x] && NeQ[a^2 - b^2, 0] && LtQ[m, -1]

Rule 4511

Int[((e_.) + (f_.)*(x_.))^(m_.)*Sec[(c_.) + (d_.)*(x_.)]*((a_.) + (b_.)*Sec[(c
 .) + (d.)*(x_.)]^(n_.)*Tan[(c_.) + (d_.)*(x_.)], x_Symbol] :> Simp[(e + f*
 x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
 {a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366

Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m
 _.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
 e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
 f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b^3 \text{Subst} \left(\int \frac{x \sec(x) \tan(x)}{(-a + \sec(x))^4} dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{\sec^{-1}(a + bx)}{3x^3} + \frac{1}{3} b^3 \text{Subst} \left(\int \frac{1}{(-a + \sec(x))^3} dx, x, \sec^{-1}(a + bx) \right) \\
 &= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1 - a^2)x^2} - \frac{\sec^{-1}(a + bx)}{3x^3} \\
 &\quad - \frac{b^3 \text{Subst} \left(\int \frac{2(1-a^2) - 2a \sec(x) - \sec^2(x)}{(-a + \sec(x))^2} dx, x, \sec^{-1}(a + bx) \right)}{6a(1 - a^2)} \\
 &= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1 - a^2)x^2} - \frac{(2 - 5a^2)b^2(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1 - a^2)^2 x} \\
 &\quad - \frac{\sec^{-1}(a + bx)}{3x^3} + \frac{b^3 \text{Subst} \left(\int \frac{2(1-a^2)^2 - a(1-4a^2)\sec(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right)}{6a^2(1 - a^2)^2} \\
 &= \frac{b(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a(1 - a^2)x^2} - \frac{(2 - 5a^2)b^2(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}}}{6a^2(1 - a^2)^2 x} - \frac{b^3 \sec^{-1}(a + bx)}{3a^3} \\
 &\quad - \frac{\sec^{-1}(a + bx)}{3x^3} + \frac{((2 - 5a^2 + 6a^4)b^3) \text{Subst} \left(\int \frac{\sec(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right)}{6a^3(1 - a^2)^2}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\sec^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\sec^{-1}(a+bx)}{3x^3} + \frac{((2-5a^2+6a^4)b^3)\text{Subst}\left(\int\frac{1}{1-a\cos(x)}dx, x, \sec^{-1}(a+bx)\right)}{6a^3(1-a^2)^2} \\
&= \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\sec^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\sec^{-1}(a+bx)}{3x^3} + \frac{((2-5a^2+6a^4)b^3)\text{Subst}\left(\int\frac{1}{1-a+(1+a)x^2}dx, x, \tan\left(\frac{1}{2}\sec^{-1}(a+bx)\right)\right)}{3a^3(1-a^2)^2} \\
&= \frac{b(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a(1-a^2)x^2} - \frac{(2-5a^2)b^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}}{6a^2(1-a^2)^2x} - \frac{b^3\sec^{-1}(a+bx)}{3a^3} \\
&\quad - \frac{\sec^{-1}(a+bx)}{3x^3} + \frac{(2-5a^2+6a^4)b^3\arctan\left(\frac{\sqrt{1+a}\tan\left(\frac{1}{2}\sec^{-1}(a+bx)\right)}{\sqrt{1-a}}\right)}{3a^3(1-a^2)^{5/2}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 241, normalized size of antiderivative = 1.33

$$\begin{aligned}
&\int \frac{\sec^{-1}(a+bx)}{x^4} dx \\
&= \frac{1}{6} \left(-\frac{b\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}(a^4+abx-4a^3bx+2b^2x^2-a^2(1+5b^2x^2))}{a^2(-1+a^2)^2x^2} - \frac{2\sec^{-1}(a+bx)}{x^3} \right. \\
&\quad \left. + \frac{2b^3\arcsin\left(\frac{1}{a+bx}\right)}{a^3} \right. \\
&\quad \left. - \frac{i(2-5a^2+6a^4)b^3\log\left(\frac{12a^3(-1+a^2)^2\left(\frac{i(-1+a^2+abx)}{\sqrt{1-a^2}}+(a+bx)\sqrt{\frac{-1+a^2+2abx+b^2x^2}{(a+bx)^2}}\right)}{(2-5a^2+6a^4)b^3x}\right)}{a^3(1-a^2)^{5/2}} \right)
\end{aligned}$$

[In] Integrate[ArcSec[a + b*x]/x^4,x]

```
[Out] (-((b*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]*(a^4 + a*b*x - 4*a^3
*b*x + 2*b^2*x^2 - a^2*(1 + 5*b^2*x^2)))/(a^2*(-1 + a^2)^2*x^2)) - (2*ArcSe
c[a + b*x])/x^3 + (2*b^3*ArcSin[(a + b*x)^(-1)]/a^3 - (I*(2 - 5*a^2 + 6*a^
4)*b^3*Log[(12*a^3*(-1 + a^2)^2*((I*(-1 + a^2 + a*b*x))/Sqrt[1 - a^2] + (a
+ b*x)*Sqrt[(-1 + a^2 + 2*a*b*x + b^2*x^2)/(a + b*x)^2]))/((2 - 5*a^2 + 6*a
^4)*b^3*x)))/(a^3*(1 - a^2)^(5/2)))/6
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 527 vs. 2(159) = 318.

Time = 0.70 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.92

method	result
parts	$-\frac{\operatorname{arcsec}(bx+a)}{3x^3} - \frac{b\sqrt{b^2x^2+2abx+a^2-1} \left(-2(a^2-1)^{\frac{3}{2}} \arctan\left(\frac{1}{\sqrt{b^2x^2+2abx+a^2-1}}\right) a^4 b^2 x^2 + 6 \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}}{b^2x^2+2abx+a^2-1}\right) \right)}{a^4 b^2 x^2 + 6 \ln\left(\frac{2a^2-2+2abx+2\sqrt{a^2-1}}{b^2x^2+2abx+a^2-1}\right)}$
derivativedivides	Expression too large to display
default	Expression too large to display

```
[In] int(arcsec(b*x+a)/x^4,x,method=_RETURNVERBOSE)
```

```
[Out] -1/3*arcsec(b*x+a)/x^3-1/6*b*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(-2*(a^2-1)^(3/2)
)*arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*a^4*b^2*x^2+6*ln(2*(a*b*x+(a^2-1)
)^(1/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*a^6*b^2*x^2+4*(a^2-1)^(3/2)*
arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*a^2*b^2*x^2-5*(b^2*x^2+2*a*b*x+a^2-
1)^(1/2)*(a^2-1)^(3/2)*a^3*b*x-11*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b*
x+a^2-1)^(1/2)+a^2-1)/x)*a^4*b^2*x^2+(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(
3/2)*a^4-2*b^2*arctan(1/(b^2*x^2+2*a*b*x+a^2-1)^(1/2))*x^2*(a^2-1)^(3/2)+2
*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)*(a^2-1)^(3/2)*a*b*x+7*ln(2*(a*b*x+(a^2-1)^(1
/2)*(b^2*x^2+2*a*b*x+a^2-1)^(1/2)+a^2-1)/x)*a^2*b^2*x^2-(b^2*x^2+2*a*b*x+a^
2-1)^(1/2)*(a^2-1)^(3/2)*a^2-2*b^2*ln(2*(a*b*x+(a^2-1)^(1/2)*(b^2*x^2+2*a*b
*x+a^2-1)^(1/2)+a^2-1)/x)*x^2)/((b^2*x^2+2*a*b*x+a^2-1)/(b*x+a)^2)^(1/2)/(b
*x+a)/a^3/(a^2-1)^(7/2)/x^2
```

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 548, normalized size of antiderivative = 3.03

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx$$

$$= \left[\frac{(6a^4 - 5a^2 + 2)\sqrt{a^2 - 1}b^3x^3 \log\left(\frac{a^2bx + a^3 + \sqrt{b^2x^2 + 2abx + a^2 - 1}(a^2 - \sqrt{a^2 - 1}a - 1) - (abx + a^2 - 1)\sqrt{a^2 - 1} - a}{x}\right) - 4(a^6 - 3}{\dots} \right]$$

[In] integrate(arcsec(b*x+a)/x^4,x, algorithm="fricas")

[Out] [1/6*((6*a^4 - 5*a^2 + 2)*sqrt(a^2 - 1)*b^3*x^3*log((a^2*b*x + a^3 + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*(a^2 - sqrt(a^2 - 1)*a - 1) - (a*b*x + a^2 - 1)*sqrt(a^2 - 1) - a)/x) - 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arcsec(b*x + a) + ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3), 1/6*(2*(6*a^4 - 5*a^2 + 2)*sqrt(-a^2 + 1)*b^3*x^3*arctan(-(sqrt(-a^2 + 1)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)*sqrt(-a^2 + 1))/(a^2 - 1)) - 4*(a^6 - 3*a^4 + 3*a^2 - 1)*b^3*x^3*arctan(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1)) + (5*a^5 - 7*a^3 + 2*a)*b^3*x^3 - 2*(a^9 - 3*a^7 + 3*a^5 - a^3)*arcsec(b*x + a) + ((5*a^5 - 7*a^3 + 2*a)*b^2*x^2 - (a^6 - 2*a^4 + a^2)*b*x)*sqrt(b^2*x^2 + 2*a*b*x + a^2 - 1))/((a^9 - 3*a^7 + 3*a^5 - a^3)*x^3)]

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{asec}(a + bx)}{x^4} dx$$

[In] integrate(asec(b*x+a)/x**4,x)

[Out] Integral(asec(a + b*x)/x**4, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)}{x^4} dx = \int \frac{\operatorname{arcsec}(bx + a)}{x^4} dx$$

[In] integrate(arcsec(b*x+a)/x^4,x, algorithm="maxima")

[Out] 1/3*(3*x^3*integrate(1/3*(b^2*x + a*b)*e^(1/2*log(b*x + a + 1) + 1/2*log(b*x + a - 1))/(b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3 + (b^2*x^5 + 2*a*b*x^4 + (a^2 - 1)*x^3)*e^(log(b*x + a + 1) + log(b*x + a - 1))), x) - arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/x^3

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 451 vs. $2(155) = 310$.

Time = 0.33 (sec) , antiderivative size = 451, normalized size of antiderivative = 2.49

$$\int \frac{\sec^{-1}(a+bx)}{x^4} dx = \frac{1}{3} b \left(\frac{(6a^4b^2 - 5a^2b^2 + 2b^2) \arctan\left(\frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + a}{\sqrt{-a^2+1}}\right)}{(a^7 - 2a^5 + a^3)\sqrt{-a^2+1}} + \frac{4(bx+a)^3 a^3 b^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)^3}{(a^7 - 2a^5 + a^3)\sqrt{-a^2+1}} \right)$$

[In] integrate(arcsec(b*x+a)/x^4,x, algorithm="giac")

[Out] $\frac{1}{3} b \left(\frac{(6a^4b^2 - 5a^2b^2 + 2b^2) \arctan\left(\frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right) + a}{\sqrt{-a^2+1}}\right)}{(a^7 - 2a^5 + a^3)\sqrt{-a^2+1}} + \frac{4(bx+a)^3 a^3 b^2 \left(\sqrt{-\frac{1}{(bx+a)^2} + 1} - 1\right)^3}{(a^7 - 2a^5 + a^3)\sqrt{-a^2+1}} \right)$

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a+bx)}{x^4} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{x^4} dx$$

[In] int(acos(1/(a + b*x))/x^4,x)

[Out] int(acos(1/(a + b*x))/x^4, x)

3.27 $\int x^3 \sec^{-1}(a + bx)^2 dx$

Optimal result	174
Rubi [A] (verified)	175
Mathematica [A] (warning: unable to verify)	180
Maple [A] (verified)	180
Fricas [F]	181
Sympy [F]	181
Maxima [F]	182
Giac [F(-2)]	182
Mupad [F(-1)]	182

Optimal result

Integrand size = 12, antiderivative size = 381

$$\begin{aligned}
 \int x^3 \sec^{-1}(a + bx)^2 dx = & -\frac{ax}{b^3} + \frac{(a + bx)^2}{12b^4} - \frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{3b^4} \\
 & - \frac{3a^2(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^4} \\
 & + \frac{a(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^4} \\
 & - \frac{(a + bx)^3\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{6b^4} - \frac{a^4 \sec^{-1}(a + bx)^2}{4b^4} \\
 & + \frac{1}{4}x^4 \sec^{-1}(a + bx)^2 - \frac{2ia \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{4ia^3 \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} + \frac{\log(a + bx)}{3b^4} \\
 & + \frac{3a^2 \log(a + bx)}{b^4} + \frac{ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & + \frac{2ia^3 \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{ia \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4} - \frac{2ia^3 \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4}
 \end{aligned}$$

[Out] $-a*x/b^3+1/12*(b*x+a)^2/b^4-1/4*a^4*\operatorname{arcsec}(b*x+a)^2/b^4+1/4*x^4*\operatorname{arcsec}(b*x+a)^2-2*I*a^3*\operatorname{polylog}(2,I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2))/b^4+I*a*\operatorname{polylog}(2,-I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^(1/2))/b^4+1/3*\ln(b*x+a)/b^4+3*a^2*\ln$

$(b*x+a)/b^4+2*I*a^3*\text{polylog}(2,-I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^{(1/2)})/b^4-I$
 $*a*\text{polylog}(2,I*(1/(b*x+a))+I*(1-1/(b*x+a)^2)^{(1/2)})/b^4-2*I*a*\text{arcsec}(b*x+a)$
 $*\arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/b^4-4*I*a^3*\text{arcsec}(b*x+a)*\arctan$
 $(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/b^4-1/3*(b*x+a)*\text{arcsec}(b*x+a)*(1-1/(b*x$
 $+a)^2)^{(1/2)}/b^4-3*a^2*(b*x+a)*\text{arcsec}(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^4+a*(b$
 $*x+a)^2*\text{arcsec}(b*x+a)*(1-1/(b*x+a)^2)^{(1/2)}/b^4-1/6*(b*x+a)^3*\text{arcsec}(b*x+a)$
 $*(1-1/(b*x+a)^2)^{(1/2)}/b^4$

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 381, normalized size of antiderivative = 1.00,
 number of steps used = 20, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used
 = {5366, 4511, 4275, 4266, 2317, 2438, 4269, 3556, 4270}

$$\begin{aligned}
 \int x^3 \sec^{-1}(a+bx)^2 dx = & -\frac{a^4 \sec^{-1}(a+bx)^2}{4b^4} - \frac{4ia^3 \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & + \frac{2ia^3 \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & - \frac{2ia^3 \text{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4} + \frac{3a^2 \log(a+bx)}{b^4} \\
 & - \frac{3a^2(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{b^4} \\
 & - \frac{2ia \sec^{-1}(a+bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & + \frac{ia \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^4} - \frac{ia \text{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^4} \\
 & + \frac{(a+bx)^2}{12b^4} + \frac{\log(a+bx)}{3b^4} + \frac{a(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{b^4} \\
 & - \frac{(a+bx)^3 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{6b^4} \\
 & - \frac{(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)}{3b^4} - \frac{ax}{b^3} + \frac{1}{4}x^4 \sec^{-1}(a+bx)^2
 \end{aligned}$$

[In] Int[x^3*ArcSec[a + b*x]^2,x]

[Out] $-\left(\frac{a*x}{b^3}\right) + \frac{(a+b*x)^2}{(12*b^4)} - \left(\frac{(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}]}{(3*b^4)} - \frac{(3*a^2*(a+b*x)*\text{Sqrt}[1-(a+b*x)^{-2}]*\text{ArcSec}[a+b*x]}{b^4} + \frac{a*(a+b*x)^2*\text{Sqrt}[1-(a+b*x)^{-2}]*\text{ArcSec}[a+b*x]}{b^4} - \frac{(a+b*x)^3*\text{Sqrt}[1-(a+b*x)^{-2}]*\text{ArcSec}[a+b*x]}{(6*b^4)} - \frac{a^4*A}{b^4}\right)$

$$\text{rcSec}[a + b*x]^2)/(4*b^4) + (x^4*\text{ArcSec}[a + b*x]^2)/4 - ((2*I)*a*\text{ArcSec}[a + b*x]*\text{ArcTan}[E^{(I*\text{ArcSec}[a + b*x])}])/b^4 - ((4*I)*a^3*\text{ArcSec}[a + b*x]*\text{ArcTan}[E^{(I*\text{ArcSec}[a + b*x])}])/b^4 + \text{Log}[a + b*x]/(3*b^4) + (3*a^2*\text{Log}[a + b*x])/b^4 + (I*a*\text{PolyLog}[2, (-I)*E^{(I*\text{ArcSec}[a + b*x])}])/b^4 + ((2*I)*a^3*\text{PolyLog}[2, (-I)*E^{(I*\text{ArcSec}[a + b*x])}])/b^4 - (I*a*\text{PolyLog}[2, I*E^{(I*\text{ArcSec}[a + b*x])}])/b^4 - ((2*I)*a^3*\text{PolyLog}[2, I*E^{(I*\text{ArcSec}[a + b*x])}])/b^4$$
Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 4266

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_)^(m_.), x_Symbol]
:= Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_)^(m_.), x_Symbol] := Simp[(-c + d*x)^m*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4270

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n*((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]
```

Rule 4275


```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x^2 \sec(x)(-a + \sec(x))^3 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^4} \\
&= \frac{1}{4}x^4 \sec^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x(-a + \sec(x))^4 dx, x, \sec^{-1}(a + bx)\right)}{2b^4} \\
&= \frac{1}{4}x^4 \sec^{-1}(a + bx)^2 \\
&\quad - \frac{\text{Subst}\left(\int (a^4x - 4a^3x \sec(x) + 6a^2x \sec^2(x) - 4ax \sec^3(x) + x \sec^4(x)) dx, x, \sec^{-1}(a + bx)\right)}{2b^4} \\
&= -\frac{a^4 \sec^{-1}(a + bx)^2}{4b^4} + \frac{1}{4}x^4 \sec^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x \sec^4(x) dx, x, \sec^{-1}(a + bx)\right)}{2b^4} \\
&\quad + \frac{(2a) \text{Subst}\left(\int x \sec^3(x) dx, x, \sec^{-1}(a + bx)\right)}{b^4} \\
&\quad - \frac{(3a^2) \text{Subst}\left(\int x \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{b^4} \\
&\quad + \frac{(2a^3) \text{Subst}\left(\int x \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^4}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} - \frac{3a^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} \\
&+ \frac{a(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} - \frac{(a+bx)^3\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{6b^4} \\
&- \frac{a^4\sec^{-1}(a+bx)^2}{4b^4} + \frac{1}{4}x^4\sec^{-1}(a+bx)^2 - \frac{4ia^3\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&- \frac{\text{Subst}\left(\int x\sec^2(x)dx, x, \sec^{-1}(a+bx)\right)}{3b^4} \\
&+ \frac{a\text{Subst}\left(\int x\sec(x)dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&+ \frac{(3a^2)\text{Subst}\left(\int \tan(x)dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&- \frac{(2a^3)\text{Subst}\left(\int \log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&+ \frac{(2a^3)\text{Subst}\left(\int \log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&= -\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} - \frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^4} \\
&- \frac{3a^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} + \frac{a(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} \\
&- \frac{(a+bx)^3\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{6b^4} - \frac{a^4\sec^{-1}(a+bx)^2}{4b^4} \\
&+ \frac{1}{4}x^4\sec^{-1}(a+bx)^2 - \frac{2ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&- \frac{4ia^3\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&+ \frac{3a^2\log(a+bx)}{b^4} + \frac{\text{Subst}\left(\int \tan(x)dx, x, \sec^{-1}(a+bx)\right)}{3b^4} \\
&- \frac{a\text{Subst}\left(\int \log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&+ \frac{a\text{Subst}\left(\int \log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^4} \\
&+ \frac{(2ia^3)\text{Subst}\left(\int \frac{\log(1-ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&- \frac{(2ia^3)\text{Subst}\left(\int \frac{\log(1+ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^4}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} - \frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^4} \\
&\quad - \frac{3a^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} + \frac{a(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} \\
&\quad - \frac{(a+bx)^3\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{6b^4} - \frac{a^4\sec^{-1}(a+bx)^2}{4b^4} \\
&\quad + \frac{1}{4}x^4\sec^{-1}(a+bx)^2 - \frac{2ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&\quad - \frac{4ia^3\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} + \frac{\log(a+bx)}{3b^4} \\
&\quad + \frac{3a^2\log(a+bx)}{b^4} + \frac{2ia^3\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&\quad - \frac{2ia^3\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^4} + \frac{(ia)\text{Subst}\left(\int\frac{\log(1-ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&\quad - \frac{(ia)\text{Subst}\left(\int\frac{\log(1+ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&= -\frac{ax}{b^3} + \frac{(a+bx)^2}{12b^4} - \frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^4} \\
&\quad - \frac{3a^2(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} + \frac{a(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^4} \\
&\quad - \frac{(a+bx)^3\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{6b^4} - \frac{a^4\sec^{-1}(a+bx)^2}{4b^4} \\
&\quad + \frac{1}{4}x^4\sec^{-1}(a+bx)^2 - \frac{2ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&\quad - \frac{4ia^3\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^4} + \frac{\log(a+bx)}{3b^4} + \frac{3a^2\log(a+bx)}{b^4} \\
&\quad + \frac{ia\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^4} + \frac{2ia^3\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^4} \\
&\quad - \frac{ia\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^4} - \frac{2ia^3\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^4}
\end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 8.39 (sec) , antiderivative size = 667, normalized size of antiderivative = 1.75

$$\int x^3 \sec^{-1}(a + bx)^2 dx$$

$$= \frac{\left(1 - \frac{a}{a+bx}\right)^3 \left(24a(2 + (1 + 2a^2) \sec^{-1}(a + bx)^2) + \frac{2+(-2+24a) \sec^{-1}(a+bx)+3(1-4a+12a^2) \sec^{-1}(a+bx)^2}{-1+\sqrt{1-\frac{1}{(a+bx)^2}}}\right) + 16(1 + 9a^2)}{}$$

[In] Integrate[x^3*ArcSec[a + b*x]^2,x]

[Out] $((1 - a/(a + b*x))^3*(24*a*(2 + (1 + 2*a^2)*ArcSec[a + b*x]^2) + (2 + (-2 + 24*a)*ArcSec[a + b*x] + 3*(1 - 4*a + 12*a^2)*ArcSec[a + b*x]^2)/(-1 + Sqrt[1 - (a + b*x)^(-2)])) + 16*(1 + 9*a^2)*Log[(a + b*x)^(-1)] - 24*a*(1 + 2*a^2)*((Pi - 2*ArcSec[a + b*x])*(Log[1 - I/E^(I*ArcSec[a + b*x])] - Log[1 + I/E^(I*ArcSec[a + b*x])]) - Pi*Log[Cot[(Pi + 2*ArcSec[a + b*x])/4]] + (2*I)*(PolyLog[2, (-I)/E^(I*ArcSec[a + b*x])] - PolyLog[2, I/E^(I*ArcSec[a + b*x])])) - (3*ArcSec[a + b*x]^2)/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2])^4 + (4*ArcSec[a + b*x]*(1 + 6*a*ArcSec[a + b*x])*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2])^3 + (8*(2*ArcSec[a + b*x] + 18*a^2*ArcSec[a + b*x] + 6*a^3*ArcSec[a + b*x]^2 + 3*a*(2 + ArcSec[a + b*x]^2))*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2]) - (3*ArcSec[a + b*x]^2)/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^4 + (4*ArcSec[a + b*x]*(1 - 6*a*ArcSec[a + b*x])*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^3 - (2 + (2 - 24*a)*ArcSec[a + b*x] + 3*(1 - 4*a + 12*a^2)*ArcSec[a + b*x]^2)/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^2 - (8*(-2*ArcSec[a + b*x] - 18*a^2*ArcSec[a + b*x] + 6*a^3*ArcSec[a + b*x]^2 + 3*a*(2 + ArcSec[a + b*x]^2))*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^3)/(48*b^4*(-1 + a/(a + b*x))^3)$

Maple [A] (verified)

Time = 1.44 (sec) , antiderivative size = 673, normalized size of antiderivative = 1.77

method	result
derivativedivides	$-\operatorname{arcsec}(bx+a)^2 a^3 (bx+a) + \frac{3 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a)^2}{2} - \operatorname{arcsec}(bx+a)^2 a (bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^4}{4} - 3 \operatorname{arcsec}(bx+a) \sqrt{1 - \frac{1}{(bx+a)^2}}$
default	$-\operatorname{arcsec}(bx+a)^2 a^3 (bx+a) + \frac{3 \operatorname{arcsec}(bx+a)^2 a^2 (bx+a)^2}{2} - \operatorname{arcsec}(bx+a)^2 a (bx+a)^3 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^4}{4} - 3 \operatorname{arcsec}(bx+a) \sqrt{1 - \frac{1}{(bx+a)^2}}$

[In] `int(x^3*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b^4}(-\operatorname{arcsec}(bx+a)^2 a^3(bx+a) + \frac{3}{2}\operatorname{arcsec}(bx+a)^2 a^2(bx+a)^2 - \operatorname{arcsec}(bx+a)^2 a(bx+a)^3 + \frac{1}{4}\operatorname{arcsec}(bx+a)^2(bx+a)^4 - 3\operatorname{arcsec}(bx+a) \left(\frac{(bx+a)^2-1}{(bx+a)^2} \right)^{1/2} a^2(bx+a) + \operatorname{arcsec}(bx+a) \left(\frac{(bx+a)^2-1}{(bx+a)^2} \right)^{1/2} a(bx+a)^2 - \frac{1}{6}\operatorname{arcsec}(bx+a) \left(\frac{(bx+a)^2-1}{(bx+a)^2} \right)^{1/2} (bx+a)^3 - 3I a^2 \operatorname{arcsec}(bx+a) - \frac{1}{3}\operatorname{arcsec}(bx+a) \left(\frac{(bx+a)^2-1}{(bx+a)^2} \right)^{1/2} (bx+a) + I \operatorname{dilog}(1 + I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a - (bx+a) a + \frac{1}{12}(bx+a)^2 - \frac{1}{3}\ln(1 + (1/(bx+a) + I(1-1/(bx+a)^2)^{1/2}))^2 + \frac{2}{3}\ln(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2}) - 3\ln(1 + (1/(bx+a) + I(1-1/(bx+a)^2)^{1/2}))^2 a^2 + 6\ln(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2}) a^2 - 2\ln(1 + I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a^3 \operatorname{arcsec}(bx+a) + 2\ln(1 - I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a^3 \operatorname{arcsec}(bx+a) - \frac{1}{3}I \operatorname{arcsec}(bx+a) - 2I \operatorname{dilog}(1 - I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a^3 - \ln(1 + I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a \operatorname{arcsec}(bx+a) + \ln(1 - I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a \operatorname{arcsec}(bx+a) - I \operatorname{dilog}(1 - I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a + 2I \operatorname{dilog}(1 + I(1/(bx+a) + I(1-1/(bx+a)^2)^{1/2})) a^3$

Fricas [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{arcsec}(bx + a)^2 dx$$

[In] `integrate(x^3*arcsec(b*x+a)^2,x, algorithm="fricas")`

[Out] `integral(x^3*arcsec(b*x + a)^2, x)`

Sympy [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{asec}^2(a + bx) dx$$

[In] `integrate(x**3*asec(b*x+a)**2,x)`

[Out] `Integral(x**3*asec(a + b*x)**2, x)`

Maxima [F]

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(x^3*arcsec(b*x+a)^2,x, algorithm="maxima")

[Out] 1/4*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/16*x^4*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/4*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^4*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 4*(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2 - 1)*b*x^4 + (a^3 - a)*x^3)*log(b*x + a)^2 - (b^3*x^6 + 2*a*b^2*x^5 + (a^2 - 1)*b*x^4 + 4*(b^3*x^6 + 3*a*b^2*x^5 + (3*a^2 - 1)*b*x^4 + (a^3 - a)*x^3)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)

Giac [F(-2)]

Exception generated.

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \text{Exception raised: RuntimeError}$$

[In] integrate(x^3*arcsec(b*x+a)^2,x, algorithm="giac")

[Out] Exception raised: RuntimeError >> an error occurred running a Giac command: INPUT:sage2OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vector & l) Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int x^3 \sec^{-1}(a + bx)^2 dx = \int x^3 \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

[In] int(x^3*acos(1/(a + b*x))^2,x)

[Out] int(x^3*acos(1/(a + b*x))^2, x)

3.28 $\int x^2 \sec^{-1}(a + bx)^2 dx$

Optimal result	183
Rubi [A] (verified)	184
Mathematica [A] (warning: unable to verify)	188
Maple [A] (verified)	188
Fricas [F]	189
Sympy [F]	189
Maxima [F]	189
Giac [F]	190
Mupad [F(-1)]	190

Optimal result

Integrand size = 12, antiderivative size = 288

$$\begin{aligned}
 \int x^2 \sec^{-1}(a + bx)^2 dx = & \frac{x}{3b^2} + \frac{2a(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3} \\
 & - \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{3b^3} + \frac{a^3 \sec^{-1}(a + bx)^2}{3b^3} \\
 & + \frac{1}{3}x^3 \sec^{-1}(a + bx)^2 + \frac{2i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{3b^3} \\
 & + \frac{4ia^2 \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{2a \log(a + bx)}{b^3} - \frac{i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{3b^3} \\
 & - \frac{2ia^2 \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{3b^3} + \frac{2ia^2 \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3}
 \end{aligned}$$

```

[Out] 1/3*x/b^2+1/3*a^3*arcsec(b*x+a)^2/b^3+1/3*x^3*arcsec(b*x+a)^2+2/3*I*arcsec(
b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^3+4*I*a^2*arcsec(b*x+a)*
arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^3-2*a*ln(b*x+a)/b^3-1/3*I*polyl
og(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3-2*I*a^2*polylog(2,-I*(1/(b
*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^3+1/3*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b
*x+a)^2)^(1/2)))/b^3+2*I*a^2*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))
)/b^3+2*a*(b*x+a)*arcsec(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^3-1/3*(b*x+a)^2*arc
sec(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^3

```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5366, 4511, 4275, 4266, 2317, 2438, 4269, 3556, 4270}

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \frac{a^3 \sec^{-1}(a + bx)^2}{3b^3} + \frac{4ia^2 \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b^3} - \frac{2ia^2 \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b^3} + \frac{2ia^2 \text{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b^3} + \frac{2i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{3b^3} - \frac{i \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{3b^3} + \frac{i \text{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{3b^3} - \frac{2a \log(a + bx)}{b^3} + \frac{2a(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \sec^{-1}(a + bx)}{b^3} - \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a + bx)^2}} \sec^{-1}(a + bx)}{3b^3} + \frac{1}{3} x^3 \sec^{-1}(a + bx)^2 + \frac{x}{3b^2}$$

[In] Int[x^2*ArcSec[a + b*x]^2,x]

[Out] x/(3*b^2) + (2*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x])/b^3 - ((a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x])/(3*b^3) + (a^3*ArcSec[a + b*x]^2)/(3*b^3) + (x^3*ArcSec[a + b*x]^2)/3 + (((2*I)/3)*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])])/b^3 + ((4*I)*a^2*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])])/b^3 - (2*a*Log[a + b*x])/b^3 - ((I/3)*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])])/b^3 - ((2*I)*a^2*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])])/b^3 + ((I/3)*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^3 + ((2*I)*a^2*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^3

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4266

Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269

Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 4270

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^n*((c_.) + (d_.)*(x_)), x_Symbol] := Simp[(-b^2)*(c + d*x)*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2]

Rule 4275

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^n*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 4511

Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366

Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^p*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,

f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int x^2 \sec(x)(-a + \sec(x))^2 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{1}{3}x^3 \sec^{-1}(a + bx)^2 - \frac{2\text{Subst}\left(\int x(-a + \sec(x))^3 dx, x, \sec^{-1}(a + bx)\right)}{3b^3} \\
 &= \frac{1}{3}x^3 \sec^{-1}(a + bx)^2 \\
 &\quad - \frac{2\text{Subst}\left(\int (-a^3x + 3a^2x \sec(x) - 3ax \sec^2(x) + x \sec^3(x)) dx, x, \sec^{-1}(a + bx)\right)}{3b^3} \\
 &= \frac{a^3 \sec^{-1}(a + bx)^2}{3b^3} + \frac{1}{3}x^3 \sec^{-1}(a + bx)^2 - \frac{2\text{Subst}\left(\int x \sec^3(x) dx, x, \sec^{-1}(a + bx)\right)}{3b^3} \\
 &\quad + \frac{(2a)\text{Subst}\left(\int x \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &\quad - \frac{(2a^2)\text{Subst}\left(\int x \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{x}{3b^2} + \frac{2a(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^3} - \frac{(a + bx)^2\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{3b^3} \\
 &\quad + \frac{a^3 \sec^{-1}(a + bx)^2}{3b^3} + \frac{1}{3}x^3 \sec^{-1}(a + bx)^2 + \frac{4ia^2 \sec^{-1}(a + bx) \arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
 &\quad - \frac{\text{Subst}\left(\int x \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{3b^3} \\
 &\quad - \frac{(2a)\text{Subst}\left(\int \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &\quad + \frac{(2a^2)\text{Subst}\left(\int \log(1 - ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &\quad - \frac{(2a^2)\text{Subst}\left(\int \log(1 + ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{x}{3b^2} + \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^3} \\
&\quad + \frac{a^3\sec^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^2 + \frac{2i\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{3b^3} \\
&\quad + \frac{4ia^2\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{2a\log(a+bx)}{b^3} \\
&\quad + \frac{\text{Subst}\left(\int \log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{3b^3} \\
&\quad - \frac{\text{Subst}\left(\int \log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{3b^3} \\
&\quad - \frac{(2ia^2)\text{Subst}\left(\int \frac{\log(1-ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{(2ia^2)\text{Subst}\left(\int \frac{\log(1+ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&= \frac{x}{3b^2} + \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^3} \\
&\quad + \frac{a^3\sec^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^2 + \frac{2i\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{3b^3} \\
&\quad + \frac{4ia^2\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{2a\log(a+bx)}{b^3} \\
&\quad - \frac{2ia^2\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{2ia^2\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad - \frac{i\text{Subst}\left(\int \frac{\log(1-ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{3b^3} + \frac{i\text{Subst}\left(\int \frac{\log(1+ix)}{x}dx, x, e^{i\sec^{-1}(a+bx)}\right)}{3b^3} \\
&= \frac{x}{3b^2} + \frac{2a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{3b^3} \\
&\quad + \frac{a^3\sec^{-1}(a+bx)^2}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^2 + \frac{2i\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{3b^3} \\
&\quad + \frac{4ia^2\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{2a\log(a+bx)}{b^3} \\
&\quad - \frac{i\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{3b^3} - \frac{2ia^2\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{i\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{3b^3} + \frac{2ia^2\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

Mathematica [A] (warning: unable to verify)

Time = 4.59 (sec) , antiderivative size = 473, normalized size of antiderivative = 1.64

$$\int x^2 \sec^{-1}(a + bx)^2 dx$$

$$= \frac{4 + 2(1 + 6a^2) \sec^{-1}(a + bx)^2 + \frac{\sec^{-1}(a+bx)(2+(-1+6a)\sec^{-1}(a+bx))}{-1+\sqrt{1-\frac{1}{(a+bx)^2}}} + 24a \log\left(\frac{1}{a+bx}\right) + 2(-1 - 6a^2) \left((\pi - 2 \sec^{-1}(a+bx)) \right)}{1}$$

`[In] Integrate[x^2*ArcSec[a + b*x]^2,x]`

```
[Out] (4 + 2*(1 + 6*a^2)*ArcSec[a + b*x]^2 + (ArcSec[a + b*x]*(2 + (-1 + 6*a)*ArcSec[a + b*x]))/(-1 + Sqrt[1 - (a + b*x)^(-2)]) + 24*a*Log[(a + b*x)^(-1)] + 2*(-1 - 6*a^2)*((Pi - 2*ArcSec[a + b*x])*(Log[1 - I/E^(I*ArcSec[a + b*x])] - Log[1 + I/E^(I*ArcSec[a + b*x])]) - Pi*Log[Cot[(Pi + 2*ArcSec[a + b*x])/4]] + (2*I)*(PolyLog[2, (-I)/E^(I*ArcSec[a + b*x])] - PolyLog[2, I/E^(I*ArcSec[a + b*x])])) + (2*ArcSec[a + b*x]^2*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2])^3 + (2*(2 + 12*a*ArcSec[a + b*x] + (1 + 6*a^2)*ArcSec[a + b*x]^2)*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] - Sin[ArcSec[a + b*x]/2]) - (2*ArcSec[a + b*x]^2*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^3 + (ArcSec[a + b*x]*(2 + (1 - 6*a)*ArcSec[a + b*x]))/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2])^2 - (2*(2 - 12*a*ArcSec[a + b*x] + (1 + 6*a^2)*ArcSec[a + b*x]^2)*Sin[ArcSec[a + b*x]/2])/(Cos[ArcSec[a + b*x]/2] + Sin[ArcSec[a + b*x]/2]))/(12*b^3)
```

Maple [A] (verified)

Time = 1.51 (sec) , antiderivative size = 498, normalized size of antiderivative = 1.73

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) - \frac{\operatorname{arcsec}(bx+a)}{3}}{1}$
default	$\frac{\operatorname{arcsec}(bx+a)^2 a^2 (bx+a) - \operatorname{arcsec}(bx+a)^2 a (bx+a)^2 + \frac{\operatorname{arcsec}(bx+a)^2 (bx+a)^3}{3} + 2 \operatorname{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) - \frac{\operatorname{arcsec}(bx+a)}{3}}{1}$

`[In] int(x^2*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/b^3*(arcsec(b*x+a)^2*a^2*(b*x+a)-arcsec(b*x+a)^2*a*(b*x+a)^2+1/3*arcsec(b*x+a)^2*(b*x+a)^3+2*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*a*(b*x+a)-1/3*arcsec(b*x+a)*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)^2+2*I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))*a^2+1/3*b*x+1/3*a+1/3*arcsec(b*x+a)*ln
```

$$(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))-1/3*\operatorname{arcsec}(b*x+a)*\ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))+1/3*I*\operatorname{dilog}(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))+2*I*\operatorname{arcsec}(b*x+a)*a+2*\ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})^2)*a-4*\ln(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})*a-2*I*\operatorname{dilog}(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2-1/3*I*\operatorname{dilog}(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))+2*\ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2*\operatorname{arcsec}(b*x+a)-2*\ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))*a^2*\operatorname{arcsec}(b*x+a))$$

Fricas [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

```
[In] integrate(x^2*arcsec(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] integral(x^2*arcsec(b*x + a)^2, x)
```

Sympy [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{asec}^2(a + bx) dx$$

```
[In] integrate(x**2*asec(b*x+a)**2,x)
```

```
[Out] Integral(x**2*asec(a + b*x)**2, x)
```

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

```
[In] integrate(x^2*arcsec(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] 1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/12*x^3*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/3*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2 - 1)*b*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a)*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)
```

Giac [F]

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(x^2*arcsec(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x^2*arcsec(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx)^2 dx = \int x^2 \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

[In] int(x^2*acos(1/(a + b*x))^2,x)

[Out] int(x^2*acos(1/(a + b*x))^2, x)

3.29 $\int x \sec^{-1}(a + bx)^2 dx$

Optimal result	191
Rubi [A] (verified)	191
Mathematica [A] (verified)	194
Maple [A] (verified)	195
Fricas [F]	195
Sympy [F]	195
Maxima [F]	196
Giac [F]	196
Mupad [F(-1)]	196

Optimal result

Integrand size = 10, antiderivative size = 154

$$\int x \sec^{-1}(a + bx)^2 dx = -\frac{(a + bx)\sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)}{b^2} - \frac{a^2 \sec^{-1}(a + bx)^2}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx)^2 - \frac{4ia \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{\log(a + bx)}{b^2} + \frac{2ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{2ia \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2}$$

```
[Out] -1/2*a^2*arcsec(b*x+a)^2/b^2+1/2*x^2*arcsec(b*x+a)^2-4*I*a*arcsec(b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^2+ln(b*x+a)/b^2+2*I*a*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2-2*I*a*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2-(b*x+a)*arcsec(b*x+a)*(1-1/(b*x+a)^2)^(1/2)/b^2
```

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.800$, Rules used

= {5366, 4511, 4275, 4266, 2317, 2438, 4269, 3556}

$$\int x \sec^{-1}(a + bx)^2 dx = -\frac{a^2 \sec^{-1}(a + bx)^2}{2b^2} - \frac{4ia \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b^2}$$

$$+ \frac{2ia \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b^2}$$

$$- \frac{2ia \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b^2} + \frac{\log(a + bx)}{b^2}$$

$$- \frac{(a + bx) \sqrt{1 - \frac{1}{(a + bx)^2}} \sec^{-1}(a + bx)}{b^2} + \frac{1}{2} x^2 \sec^{-1}(a + bx)^2$$

[In] Int[x*ArcSec[a + b*x]^2,x]

[Out] -(((a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x])/b^2) - (a^2*ArcSec[a + b*x]^2)/(2*b^2) + (x^2*ArcSec[a + b*x]^2)/2 - ((4*I)*a*ArcSec[a + b*x]*ArcTan[E^(I*ArcSec[a + b*x])])/b^2 + Log[a + b*x]/b^2 + ((2*I)*a*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])])/b^2 - ((2*I)*a*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^2

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4266

Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269


```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_)])^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x^2 \sec(x)(-a + \sec(x)) \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
&= \frac{1}{2} x^2 \sec^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x(-a + \sec(x))^2 dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
&= \frac{1}{2} x^2 \sec^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int (a^2 x - 2ax \sec(x) + x \sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
&= -\frac{a^2 \sec^{-1}(a + bx)^2}{2b^2} + \frac{1}{2} x^2 \sec^{-1}(a + bx)^2 - \frac{\text{Subst}\left(\int x \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
&\quad + \frac{(2a)\text{Subst}\left(\int x \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^2} - \frac{a^2\sec^{-1}(a+bx)^2}{2b^2} + \frac{1}{2}x^2\sec^{-1}(a+bx)^2 \\
&\quad - \frac{4ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^2} + \frac{\text{Subst}\left(\int \tan(x) dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&\quad - \frac{(2a)\text{Subst}\left(\int \log(1-ie^{ix}) dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&\quad + \frac{(2a)\text{Subst}\left(\int \log(1+ie^{ix}) dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&= -\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^2} - \frac{a^2\sec^{-1}(a+bx)^2}{2b^2} \\
&\quad + \frac{1}{2}x^2\sec^{-1}(a+bx)^2 - \frac{4ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&\quad + \frac{\log(a+bx)}{b^2} + \frac{(2ia)\text{Subst}\left(\int \frac{\log(1-ix)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&\quad - \frac{(2ia)\text{Subst}\left(\int \frac{\log(1+ix)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&= -\frac{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)}{b^2} - \frac{a^2\sec^{-1}(a+bx)^2}{2b^2} \\
&\quad + \frac{1}{2}x^2\sec^{-1}(a+bx)^2 - \frac{4ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^2} + \frac{\log(a+bx)}{b^2} \\
&\quad + \frac{2ia\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^2} - \frac{2ia\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 144, normalized size of antiderivative = 0.94

$$\int x \sec^{-1}(a+bx)^2 dx$$

$$= \frac{-\left((a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)\right) - a(a+bx)\sec^{-1}(a+bx)^2 + \frac{1}{2}(a+bx)^2\sec^{-1}(a+bx)^2 - 4ia\sec^{-1}(a+bx)\arctan\left(e^{i\sec^{-1}(a+bx)}\right) + \log(a+bx) + (2ia)\text{PolyLog}\left[2, (-I)E^{(I\sec^{-1}(a+bx))}\right] - (2ia)\text{PolyLog}\left[2, I E^{(I\sec^{-1}(a+bx))}\right]}{b^2}$$

[In] Integrate[x*ArcSec[a + b*x]^2, x]

[Out] $\frac{-((a+bx)\sqrt{1-(a+bx)^{-2}})\text{ArcSec}[a+bx] - a(a+bx)\text{ArcSec}[a+bx]^2 + ((a+bx)^2\text{ArcSec}[a+bx]^2)/2 - (4I)a\text{ArcSec}[a+bx]\text{ArcTan}[E^{(I\text{ArcSec}[a+bx])}] + \text{Log}[a+bx] + (2I)a\text{PolyLog}[2, (-I)E^{(I\text{ArcSec}[a+bx])}] - (2I)a\text{PolyLog}[2, I E^{(I\text{ArcSec}[a+bx])}]}{b^2}$

Maple [A] (verified)

Time = 0.95 (sec) , antiderivative size = 225, normalized size of antiderivative = 1.46

method	result
derivativedivides	$\frac{-a \left(\operatorname{arcsec}(bx+a)^2 (bx+a) + 2 \operatorname{arcsec}(bx+a) \ln \left(1 + i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) \right) - 2 \operatorname{arcsec}(bx+a) \ln \left(1 - i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right)}{1}$
default	$\frac{-a \left(\operatorname{arcsec}(bx+a)^2 (bx+a) + 2 \operatorname{arcsec}(bx+a) \ln \left(1 + i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right) \right) - 2 \operatorname{arcsec}(bx+a) \ln \left(1 - i \left(\frac{1}{bx+a} + i \sqrt{1 - \frac{1}{(bx+a)^2}} \right) \right)}{1}$

[In] `int(x*arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/b^2*(-a*(arcsec(b*x+a)^2*(b*x+a)+2*arcsec(b*x+a)*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-2*arcsec(b*x+a)*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-2*I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+2*I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))+1/2*arcsec(b*x+a)^2*(b*x+a)^2-arcsec(b*x+a)*((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)-ln(1/(b*x+a)))
```

Fricas [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

[In] `integrate(x*arcsec(b*x+a)^2,x, algorithm="fricas")`[Out] `integral(x*arcsec(b*x + a)^2, x)`**Sympy [F]**

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{asec}^2(a + bx) dx$$

[In] `integrate(x*asec(b*x+a)**2,x)`[Out] `Integral(x*asec(a + b*x)**2, x)`

Maxima [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(x*arcsec(b*x+a)^2,x, algorithm="maxima")

[Out] 1/2*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/8*x^2*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/2*(2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x^2*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a)^2 - (b^3*x^4 + 2*a*b^2*x^3 + (a^2 - 1)*b*x^2 + 2*(b^3*x^4 + 3*a*b^2*x^3 + (3*a^2 - 1)*b*x^2 + (a^3 - a)*x)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)

Giac [F]

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(x*arcsec(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x*arcsec(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx)^2 dx = \int x \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

[In] int(x*acos(1/(a + b*x))^2,x)

[Out] int(x*acos(1/(a + b*x))^2, x)

3.30 $\int \sec^{-1}(a + bx)^2 dx$

Optimal result	197
Rubi [A] (verified)	197
Mathematica [A] (verified)	199
Maple [A] (verified)	200
Fricas [F]	200
Sympy [F]	200
Maxima [F]	201
Giac [F]	201
Mupad [F(-1)]	201

Optimal result

Integrand size = 8, antiderivative size = 94

$$\int \sec^{-1}(a + bx)^2 dx = \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} + \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

[Out] (b*x+a)*arcsec(b*x+a)^2/b+4*I*arcsec(b*x+a)*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b-2*I*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+2*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5360, 5324, 3842, 4266, 2317, 2438}

$$\int \sec^{-1}(a + bx)^2 dx = \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{2i \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{2i \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{(a + bx) \sec^{-1}(a + bx)^2}{b}$$

[In] Int[ArcSec[a + b*x]^2,x]

[Out] $((a + b*x)*\text{ArcSec}[a + b*x]^2)/b + ((4*I)*\text{ArcSec}[a + b*x]*\text{ArcTan}[E^{(I*\text{ArcSec}[a + b*x])}])/b - ((2*I)*\text{PolyLog}[2, (-I)*E^{(I*\text{ArcSec}[a + b*x])}])/b + ((2*I)*\text{PolyLog}[2, I*E^{(I*\text{ArcSec}[a + b*x])}])/b$

Rule 2317

$\text{Int}[\text{Log}[(a_.) + (b_.)*(F_.)^{((e_.)*((c_.) + (d_.)*(x_)))})^{(n_.)}], x_Symbol] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x)})^n), x] /; \text{FreeQ}\{F, a, b, c, d, e, n\}, x] \&\& \text{GtQ}[a, 0]$

Rule 2438

$\text{Int}[\text{Log}[(c_.)*((d_.) + (e_.)*(x_.)^{(n_.)})]/(x_.), x_Symbol] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}\{c, d, e, n\}, x] \&\& \text{EqQ}[c*d, 1]$

Rule 3842

$\text{Int}[(x_.)^{(m_.)}*\text{Sec}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(p_.)}*\text{Tan}[(a_.) + (b_.)*(x_.)^{(n_.)}]^{(q_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m - n + 1)}*(\text{Sec}[a + b*x^n]^p/(b*n*p)), x] - \text{Dist}[(m - n + 1)/(b*n*p), \text{Int}[x^{(m - n)}*\text{Sec}[a + b*x^n]^p, x], x] /; \text{FreeQ}\{a, b, p\}, x] \&\& \text{IntegerQ}[n] \&\& \text{GeQ}[m, n] \&\& \text{EqQ}[q, 1]$

Rule 4266

$\text{Int}[\text{csc}[(e_.) + \text{Pi}*(k_.) + (f_.)*(x_.)]*((c_.) + (d_.)*(x_.)^{(m_.)}), x_Symbol] \rightarrow \text{Simp}[-2*(c + d*x)^m*(\text{ArcTanh}[E^{(I*k*Pi)}*E^{(I*(e + f*x))}]/f), x] + (-\text{Dist}[d*(m/f), \text{Int}[(c + d*x)^{(m - 1)}*\text{Log}[1 - E^{(I*k*Pi)}*E^{(I*(e + f*x))}], x], x] + \text{Dist}[d*(m/f), \text{Int}[(c + d*x)^{(m - 1)}*\text{Log}[1 + E^{(I*k*Pi)}*E^{(I*(e + f*x))}], x], x]) /; \text{FreeQ}\{c, d, e, f\}, x] \&\& \text{IntegerQ}[2*k] \&\& \text{IGtQ}[m, 0]$

Rule 5324

$\text{Int}[(a_.) + \text{ArcSec}[(c_.)*(x_.)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/c, \text{Subst}[\text{Int}[(a + b*x)^n*\text{Sec}[x]*\text{Tan}[x], x], x, \text{ArcSec}[c*x]], x] /; \text{FreeQ}\{a, b, c, n\}, x] \&\& \text{IGtQ}[n, 0]$

Rule 5360

$\text{Int}[(a_.) + \text{ArcSec}[(c_.) + (d_.)*(x_.)]*(b_.)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[1/d, \text{Subst}[\text{Int}[(a + b*\text{ArcSec}[x])^p, x], x, c + d*x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{IGtQ}[p, 0]$

Rubi steps

$$\text{integral} = \frac{\text{Subst}\left(\int \sec^{-1}(x)^2 dx, x, a + bx\right)}{b}$$

$$\begin{aligned}
&= \frac{\text{Subst}\left(\int x^2 \sec(x) \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} - \frac{2\text{Subst}\left(\int x \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} + \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} \\
&\quad + \frac{2\text{Subst}\left(\int \log(1 - ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
&\quad - \frac{2\text{Subst}\left(\int \log(1 + ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
&= \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} + \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} \\
&\quad - \frac{(2i)\text{Subst}\left(\int \frac{\log(1 - ix)}{x} dx, x, e^{i \sec^{-1}(a + bx)}\right)}{b} \\
&\quad + \frac{(2i)\text{Subst}\left(\int \frac{\log(1 + ix)}{x} dx, x, e^{i \sec^{-1}(a + bx)}\right)}{b} \\
&= \frac{(a + bx) \sec^{-1}(a + bx)^2}{b} + \frac{4i \sec^{-1}(a + bx) \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} \\
&\quad - \frac{2i \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{2i \text{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.18

$$\begin{aligned}
&\int \sec^{-1}(a + bx)^2 dx \\
&= \frac{\sec^{-1}(a + bx) \left((a + bx) \sec^{-1}(a + bx) - 2 \log\left(1 - ie^{i \sec^{-1}(a + bx)}\right) + 2 \log\left(1 + ie^{i \sec^{-1}(a + bx)}\right) \right) - 2i \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right) + 2i \text{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}
\end{aligned}$$

[In] Integrate[ArcSec[a + b*x]^2,x]

[Out] (ArcSec[a + b*x]*((a + b*x)*ArcSec[a + b*x] - 2*Log[1 - I*E^(I*ArcSec[a + b*x])]) + 2*Log[1 + I*E^(I*ArcSec[a + b*x])]) - (2*I)*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] + (2*I)*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b

Maple [A] (verified)

Time = 0.54 (sec) , antiderivative size = 162, normalized size of antiderivative = 1.72

method	result
derivativedivides	$\frac{\operatorname{arcsec}(bx+a)^2(bx+a)+2 \operatorname{arcsec}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)-2 \operatorname{arcsec}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)}{b}$
default	$\frac{\operatorname{arcsec}(bx+a)^2(bx+a)+2 \operatorname{arcsec}(bx+a) \ln\left(1+i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)-2 \operatorname{arcsec}(bx+a) \ln\left(1-i\left(\frac{1}{bx+a}+i\sqrt{1-\frac{1}{(bx+a)^2}}\right)\right)}{b}$

[In] int(arcsec(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(arcsec(b*x+a)^2*(b*x+a)+2*arcsec(b*x+a)*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-2*arcsec(b*x+a)*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))-2*I*dilog(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))+2*I*dilog(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))

Fricas [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(arcsec(b*x+a)^2,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)^2, x)

Sympy [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{asec}^2(a + bx) dx$$

[In] integrate(asec(b*x+a)**2,x)

[Out] Integral(asec(a + b*x)**2, x)

Maxima [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(arcsec(b*x+a)^2,x, algorithm="maxima")

[Out] x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - 1/4*x*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate((2*sqrt(b*x + a + 1)*sqrt(b*x + a - 1)*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)) + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)

Giac [F]

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{arcsec}(bx + a)^2 dx$$

[In] integrate(arcsec(b*x+a)^2,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(a + bx)^2 dx = \int \operatorname{acos}\left(\frac{1}{a + bx}\right)^2 dx$$

[In] int(acos(1/(a + b*x))^2,x)

[Out] int(acos(1/(a + b*x))^2, x)

3.31 $\int \frac{\sec^{-1}(a+bx)^2}{x} dx$

Optimal result	202
Rubi [A] (verified)	203
Mathematica [B] (warning: unable to verify)	207
Maple [F]	209
Fricas [F]	209
Sympy [F]	210
Maxima [F]	210
Giac [F]	210
Mupad [F(-1)]	210

Optimal result

Integrand size = 12, antiderivative size = 310

$$\begin{aligned}
 \int \frac{\sec^{-1}(a+bx)^2}{x} dx &= \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \sec^{-1}(a+bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
 &\quad - 2i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) \\
 &\quad - 2i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad + i \sec^{-1}(a+bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
 &\quad + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}} \right) + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
 &\quad - \frac{1}{2} \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right)
 \end{aligned}$$

```

[Out] -arcsec(b*x+a)^2*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+arcsec(b*x+a)^
2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x
+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+I*arcs
ec(b*x+a)*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-2*I*arcsec(b*x+
a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))-2*I*
arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(
1/2)))-1/2*polylog(3,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+2*polylog(3,a*

```

$(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/(1-(-a^2+1)^{(1/2)}))+2*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/(1+(-a^2+1)^{(1/2)}))$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 310, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5366, 4647, 4626, 3800, 2221, 2611, 2320, 6724, 4616}

$$\begin{aligned} \int \frac{\sec^{-1}(a+bx)^2}{x} dx = & -2i \sec^{-1}(a+bx) \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) \\ & - 2i \sec^{-1}(a+bx) \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right) \\ & + 2 \text{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) + 2 \text{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right) \\ & + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}} \right) \\ & + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1} \right) \\ & + i \sec^{-1}(a+bx) \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\ & - \frac{1}{2} \text{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) \\ & - \sec^{-1}(a+bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \end{aligned}$$

[In] Int[ArcSec[a + b*x]^2/x,x]

[Out] ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - ArcSec[a + b*x]^2*Log[1 + E^((2*I)*ArcSec[a + b*x])] - (2*I)*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] - (2*I)*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + I*ArcSec[a + b*x]*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] + 2*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + 2*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - PolyLog[3, -E^((2*I)*ArcSec[a + b*x])]/2

Rule 2221

Int[(((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a]], x]

)^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.))]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3800

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] :=> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

Rule 4616

Int[(((e_.) + (f_.)*(x_))^(m_.)*Sin[(c_.) + (d_.)*(x_)])/(Cos[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :=> Simp[I*((e + f*x)^(m + 1)/(b*f*(m + 1))), x] + (-Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a - Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x))), x], x] - Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a + Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x))), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]

Rule 4626

Int[(((e_.) + (f_.)*(x_))^(m_.)*Tan[(c_.) + (d_.)*(x_)])^(n_.)/(Cos[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :=> Dist[1/a, Int[(e + f*x)^m*Tan[c + d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sin[c + d*x]*(Tan[c + d*x])^(n - 1)/(a + b*Cos[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 4647

Int[(((e_.) + (f_.)*(x_))^(m_.)*(F_)[(c_.) + (d_.)*(x_)])^(n_.)*(G_)[(c_.) + (d_.)*(x_)]^(p_.)/((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)]), x_Symbol] :=> In

```
t[(e + f*x)^m*Cos[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cos[c + d*x]))
, x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m
, n, p]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^ (p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x^2 \sec(x) \tan(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= \text{Subst}\left(\int \frac{x^2 \tan(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= a \text{Subst}\left(\int \frac{x^2 \sin(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) + \text{Subst}\left(\int x^2 \tan(x) dx, x, \sec^{-1}(a + bx)\right) \\
&= -\left(2i \text{Subst}\left(\int \frac{e^{2ix} x^2}{1 + e^{2ix}} dx, x, \sec^{-1}(a + bx)\right)\right) \\
&\quad - (ia) \text{Subst}\left(\int \frac{e^{ix} x^2}{1 - \sqrt{1 - a^2} - ae^{ix}} dx, x, \sec^{-1}(a + bx)\right) \\
&\quad - (ia) \text{Subst}\left(\int \frac{e^{ix} x^2}{1 + \sqrt{1 - a^2} - ae^{ix}} dx, x, \sec^{-1}(a + bx)\right) \\
&= \sec^{-1}(a + bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a + bx)}}{1 - \sqrt{1 - a^2}}\right) + \sec^{-1}(a + bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a + bx)}}{1 + \sqrt{1 - a^2}}\right) \\
&\quad - \sec^{-1}(a + bx)^2 \log\left(1 + e^{2i \sec^{-1}(a + bx)}\right) \\
&\quad - 2 \text{Subst}\left(\int x \log\left(1 - \frac{ae^{ix}}{1 - \sqrt{1 - a^2}}\right) dx, x, \sec^{-1}(a + bx)\right) \\
&\quad - 2 \text{Subst}\left(\int x \log\left(1 - \frac{ae^{ix}}{1 + \sqrt{1 - a^2}}\right) dx, x, \sec^{-1}(a + bx)\right) \\
&\quad + 2 \text{Subst}\left(\int x \log(1 + e^{2ix}) dx, x, \sec^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx)^2 \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - 2i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 2i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - i\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -e^{2ix}\right) dx, x, \sec^{-1}(a+bx)\right) \\
&\quad + 2i\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{ae^{ix}}{1 - \sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right) \\
&\quad + 2i\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{ae^{ix}}{1 + \sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right) \\
&= \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx)^2 \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - 2i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 2i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + i\sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - \frac{1}{2}\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad + 2\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1 - \sqrt{1-a^2}}\right)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right) \\
&\quad + 2\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1 + \sqrt{1-a^2}}\right)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a + bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 2i \sec^{-1}(a + bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad - 2i \sec^{-1}(a + bx) \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad + i \sec^{-1}(a + bx) \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad + 2 \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) - \frac{1}{2} \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right)
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 813 vs. $2(310) = 620$.

Time = 1.82 (sec) , antiderivative size = 813, normalized size of antiderivative = 2.62

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^2}{x} dx = & \sec^{-1}(a+bx)^2 \log \left(1 + \frac{ae^{i \sec^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}} \right) \\
& + \sec^{-1}(a+bx)^2 \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) - 4 \sec^{-1}(a \\
& + bx) \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) \\
& + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
& + \sec^{-1}(a+bx)^2 \log \left(1 - \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) + 4 \sec^{-1}(a \\
& + bx) \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 - \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) \\
& - 2 \sec^{-1}(a+bx)^2 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
& + \sec^{-1}(a+bx)^2 \log \left(\frac{2 \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a+bx} \right) \\
& - \sec^{-1}(a+bx)^2 \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& + 4 \sec^{-1}(a+bx) \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 \right. \\
& \quad \left. + \frac{(-1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& - \sec^{-1}(a+bx)^2 \log \left(1 - \frac{(1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& - 4 \sec^{-1}(a+bx) \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 \right. \\
& \quad \left. - \frac{(1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& - 2i \sec^{-1}(a+bx) \text{PolyLog} \left(2, -\frac{ae^{i \sec^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}} \right) \\
& - 2i \sec^{-1}(a+bx) \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right)
\end{aligned}$$

[In] Integrate[ArcSec[a + b*x]^2/x,x]

[Out] ArcSec[a + b*x]^2*Log[1 + (a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])] + ArcSec[a + b*x]^2*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] - 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + ArcSec[a + b*x]^2*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] - 2*ArcSec[a + b*x]^2*Log[1 + E^((2*I)*ArcSec[a + b*x])] + ArcSec[a + b*x]^2*Log[(2*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/(a + b*x)] - ArcSec[a + b*x]^2*Log[1 + ((-1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] + 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] - ArcSec[a + b*x]^2*Log[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] - 4*ArcSec[a + b*x]*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] - (2*I)*ArcSec[a + b*x]*PolyLog[2, -(a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])] - (2*I)*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + I*ArcSec[a + b*x]*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] + 2*PolyLog[3, -(a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])] + 2*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - PolyLog[3, -E^((2*I)*ArcSec[a + b*x])]/2

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

[In] int(arcsec(b*x+a)^2/x,x)

[Out] int(arcsec(b*x+a)^2/x,x)

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

[In] integrate(arcsec(b*x+a)^2/x,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)^2/x, x)

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{asec}^2(a + bx)}{x} dx$$

[In] integrate(asec(b*x+a)**2/x,x)

[Out] Integral(asec(a + b*x)**2/x, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

[In] integrate(arcsec(b*x+a)^2/x,x, algorithm="maxima")

[Out] integrate(arcsec(b*x + a)^2/x, x)

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x} dx$$

[In] integrate(arcsec(b*x+a)^2/x,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^2/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^2}{x} dx = \int \frac{\operatorname{acos}\left(\frac{1}{a+bx}\right)^2}{x} dx$$

[In] int(acos(1/(a + b*x))^2/x,x)

[Out] int(acos(1/(a + b*x))^2/x, x)

3.32 $\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx$

Optimal result	211
Rubi [A] (verified)	212
Mathematica [B] (verified)	215
Maple [A] (verified)	216
Fricas [F]	216
Sympy [F]	217
Maxima [F]	217
Giac [F]	217
Mupad [F(-1)]	217

Optimal result

Integrand size = 12, antiderivative size = 244

$$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx = -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

```
[Out] -b*arcsec(b*x+a)^2/a-arcsec(b*x+a)^2/x-2*I*b*arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*I*b*arcsec(b*x+a)*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-2*b*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+2*b*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Rubi [A] (verified)

Time = 0.32 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5366, 4511, 4276, 3402, 2296, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx = -\frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x}$$

[In] Int[ArcSec[a + b*x]^2/x^2,x]

[Out] -((b*ArcSec[a + b*x]^2)/a) - ArcSec[a + b*x]^2/x - ((2*I)*b*ArcSec[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + ((2*I)*b*ArcSec[a + b*x]*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (2*b*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (2*b*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2])

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2296

Int[((F_)^(u_)*((f_) + (g_)*(x_))^(m_))/((a_) + (b_)*(F_)^(u_) + (c_)*(F_)^(v_)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v, 2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]

)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3402

Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(x_)]), x_Symbol] :> Dist[2, Int[(c + d*x)^m*E^(I*Pi*(k - 1/2))*(E^(I*(e + f*x)))/(b + 2*a*E^(I*Pi*(k - 1/2))*E^(I*(e + f*x)) - b*E^(2*I*k*Pi)*E^(2*I*(e + f*x))), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[2*k] && NeQ[a^2 - b^2, 0] && IGtQ[m, 0]

Rule 4276

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Sin[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGtQ[m, 0]

Rule 4511

Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_) + (b_.)*Sec[(c_.) + (d_.)*(x_)]^(n_.)*Tan[(c_.) + (d_.)*(x_)]), x_Symbol] :> Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366

Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rubi steps

$$\begin{aligned}
 \text{integral} &= b \text{Subst} \left(\int \frac{x^2 \sec(x) \tan(x)}{(-a + \sec(x))^2} dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{\sec^{-1}(a + bx)^2}{x} + (2b) \text{Subst} \left(\int \frac{x}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{\sec^{-1}(a + bx)^2}{x} + (2b) \text{Subst} \left(\int \left(-\frac{x}{a} + \frac{x}{a(1 - a \cos(x))} \right) dx, x, \sec^{-1}(a + bx) \right)
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} + \frac{(2b)\text{Subst}\left(\int \frac{x}{1-a \cos(x)} dx, x, \sec^{-1}(a+bx)\right)}{a} \\
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} + \frac{(4b)\text{Subst}\left(\int \frac{e^{ix}x}{-a+2e^{ix}-ae^{2ix}} dx, x, \sec^{-1}(a+bx)\right)}{a} \\
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} \\
&\quad - \frac{(4b)\text{Subst}\left(\int \frac{e^{ix}x}{2-2\sqrt{1-a^2}-2ae^{ix}} dx, x, \sec^{-1}(a+bx)\right)}{\sqrt{1-a^2}} \\
&\quad + \frac{(4b)\text{Subst}\left(\int \frac{e^{ix}x}{2+2\sqrt{1-a^2}-2ae^{ix}} dx, x, \sec^{-1}(a+bx)\right)}{\sqrt{1-a^2}} \\
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(2ib)\text{Subst}\left(\int \log\left(1 - \frac{2ae^{ix}}{2-2\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(2ib)\text{Subst}\left(\int \log\left(1 - \frac{2ae^{ix}}{2+2\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{(2b)\text{Subst}\left(\int \frac{\log\left(1 - \frac{2ax}{2-2\sqrt{1-a^2}}\right)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{(2b)\text{Subst}\left(\int \frac{\log\left(1 - \frac{2ax}{2+2\sqrt{1-a^2}}\right)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b \sec^{-1}(a+bx)^2}{a} - \frac{\sec^{-1}(a+bx)^2}{x} - \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{2ib \sec^{-1}(a+bx) \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{2b \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

Mathematica [B] (verified)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 686 vs. $2(244) = 488$.

Time = 1.73 (sec) , antiderivative size = 686, normalized size of antiderivative = 2.81

$$\int \frac{\sec^{-1}(a+bx)^2}{x^2} dx = \frac{(a+bx) \sec^{-1}(a+bx)^2}{x} + \frac{2b \left(2 \sec^{-1}(a+bx) \operatorname{arctanh}\left(\frac{(-1+a) \cot\left(\frac{1}{2} \sec^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) - 2 \arccos\left(\frac{1}{a}\right) \operatorname{arctanh}\left(\frac{(1+a) \tan\left(\frac{1}{2} \sec^{-1}(a+bx)\right)}{\sqrt{-1+a^2}}\right) \right)}{\sqrt{-1+a^2}}$$

[In] Integrate[ArcSec[a + b*x]^2/x^2,x]

[Out] -((((a + b*x)*ArcSec[a + b*x]^2)/x + (2*b*(2*ArcSec[a + b*x]*ArcTanh[((-1 + a)*Cot[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]] - 2*ArcCos[a^(-1)]*ArcTanh[((1 + a)*Tan[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]] + (ArcCos[a^(-1)] - (2*I)*ArcTanh[((-1 + a)*Cot[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]] + (2*I)*ArcTanh[((1 + a)*Tan[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]))*Log[Sqrt[-1 + a^2]/(Sqrt[2]*Sqrt[a]*E^((I/2)*ArcSec[a + b*x])*Sqrt[-((b*x)/(a + b*x))]) + (ArcCos[a^(-1)] + (2*I)*(ArcTanh[((-1 + a)*Cot[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]] - ArcTanh[((1 + a)*Tan[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]))*Log[(Sqrt[-1 + a^2]*E^((I/2)*ArcSec[a + b*x]))/(Sqrt[2]*Sqrt[a]*Sqrt[-((b*x)/(a + b*x))]) - (ArcCos[a^(-1)] - (2*I)*ArcTanh[((1 + a)*Tan[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]))*Log[((-1 + a)*(I + I*a + Sqrt[-1 + a^2])*(-I + Tan[ArcSec[a + b*x]/2]))/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] - (ArcCos[a^(-1)] + (2*I)*ArcTanh[((1 + a)*Tan[ArcSec[a + b*x]/2)]/Sqrt[-1 + a^2]))*Log[((-1 + a)*(-I - I*a + Sqrt[-1 + a^2])*(I + Tan[ArcSec[a + b*x]/2]))/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] + I*(-PolyLog[2, ((1 - I*Sqrt[-1 + a^2])*(1 - a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))] + PolyLog[2, ((1 + I*Sqrt[-1 + a^2])*(1 - a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))/(a*(-1 + a + Sqrt[-1 + a^2]*Tan[ArcSec[a + b*x]/2]))]))/Sqrt[-1 + a^2])/a

Maple [A] (verified)

Time = 1.64 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.38

method	result
derivativedivides	$b \left(-\frac{(bx+a) \operatorname{arcsec}(bx+a)^2}{abx} - \frac{2i\sqrt{-a^2+1} \operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i\sqrt{1-\frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} + \frac{2i\sqrt{-a^2+1}}{a(a^2-1)} \right)$
default	$b \left(-\frac{(bx+a) \operatorname{arcsec}(bx+a)^2}{abx} - \frac{2i\sqrt{-a^2+1} \operatorname{arcsec}(bx+a) \ln \left(\frac{-a \left(\frac{1}{bx+a} + i\sqrt{1-\frac{1}{(bx+a)^2}} \right) + \sqrt{-a^2+1} + 1}{1+\sqrt{-a^2+1}} \right)}{a(a^2-1)} + \frac{2i\sqrt{-a^2+1}}{a(a^2-1)} \right)$

```
[In] int(arcsec(b*x+a)^2/x^2,x,method=_RETURNVERBOSE)
```

```
[Out] b*(-(b*x+a)*arcsec(b*x+a)^2/a/b/x-2*I*(-a^2+1)^(1/2)/a/(a^2-1)*arcsec(b*x+a)
)*ln((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1+(-a^2+1)^(1/2)))
)+2*I*(-a^2+1)^(1/2)/a/(a^2-1)*arcsec(b*x+a)*ln((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))
)+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2)))
)-2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)+1)/(1
+(-a^2+1)^(1/2)))
)+2*(-a^2+1)^(1/2)/a/(a^2-1)*dilog((a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))+(-a^2+1)^(1/2)-1)/(-1+(-a^2+1)^(1/2))))
```

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

```
[In] integrate(arcsec(b*x+a)^2/x^2,x, algorithm="fricas")
```

```
[Out] integral(arcsec(b*x + a)^2/x^2, x)
```


Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{asec}^2(a + bx)}{x^2} dx$$

[In] integrate(asec(b*x+a)**2/x**2,x)

[Out] Integral(asec(a + b*x)**2/x**2, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

[In] integrate(arcsec(b*x+a)^2/x^2,x, algorithm="maxima")

[Out] $-1/4*(4*\arctan(\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1})^2 - 4*x*\operatorname{integrate}((2*\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1}*b*x*\arctan(\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1}) - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a))*\log(b^2*x^2 + 2*a*b*x + a^2))/(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2), x) - \log(b^2*x^2 + 2*a*b*x + a^2)^2)/x$

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^2}{x^2} dx$$

[In] integrate(arcsec(b*x+a)^2/x^2,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^2/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^2}{x^2} dx = \int \frac{\operatorname{acos}\left(\frac{1}{a+bx}\right)^2}{x^2} dx$$

[In] int(acos(1/(a + b*x))^2/x^2,x)

[Out] int(acos(1/(a + b*x))^2/x^2, x)

3.33 $\int x^2 \sec^{-1}(a + bx)^3 dx$

Optimal result	219
Rubi [A] (verified)	220
Mathematica [A] (verified)	228
Maple [A] (verified)	229
Fricas [F]	230
Sympy [F]	230
Maxima [F]	230
Giac [F]	231
Mupad [F(-1)]	231

Optimal result

Integrand size = 12, antiderivative size = 494

$$\begin{aligned}
 \int x^2 \sec^{-1}(a+bx)^3 dx = & \frac{(a+bx) \sec^{-1}(a+bx)}{b^3} - \frac{3ia \sec^{-1}(a+bx)^2}{b^3} \\
 & + \frac{3a(a+bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{b^3} \\
 & - \frac{(a+bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{2b^3} + \frac{a^3 \sec^{-1}(a+bx)^3}{3b^3} \\
 & + \frac{1}{3} x^3 \sec^{-1}(a+bx)^3 + \frac{i \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{6ia^2 \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^3} \\
 & + \frac{6a \sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{6ia^2 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{i \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{6ia^2 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{3ia \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{b^3} + \frac{\operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{6a^2 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{\operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} - \frac{6a^2 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3}
 \end{aligned}$$

```

[Out] (b*x+a)*arcsec(b*x+a)/b^3-I*arcsec(b*x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b
*x+a)^2)^(1/2)))/b^3+1/3*a^3*arcsec(b*x+a)^3/b^3+1/3*x^3*arcsec(b*x+a)^3-3*
I*a*arcsec(b*x+a)^2/b^3+6*I*a^2*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*
x+a)^2)^(1/2))/b^3-arctanh((1-1/(b*x+a)^2)^(1/2))/b^3+6*a*arcsec(b*x+a)*ln(

```

$$\begin{aligned}
& 1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})^2/b^3+I*\operatorname{arcsec}(b*x+a)^2*\arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/b^3-3*I*a*\operatorname{polylog}(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})^2)/b^3+I*\operatorname{arcsec}(b*x+a)*\operatorname{polylog}(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3+6*I*a^2*\operatorname{arcsec}(b*x+a)*\operatorname{polylog}(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3-6*I*a^2*\operatorname{arcsec}(b*x+a)*\operatorname{polylog}(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3+\operatorname{polylog}(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3+6*a^2*\operatorname{polylog}(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3-\operatorname{polylog}(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3-6*a^2*\operatorname{polylog}(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)}))/b^3+3*a*(b*x+a)*\operatorname{arcsec}(b*x+a)^2*(1-1/(b*x+a)^2)^{(1/2)}/b^3-1/2*(b*x+a)^2*\operatorname{arcsec}(b*x+a)^2*(1-1/(b*x+a)^2)^{(1/2)}/b^3
\end{aligned}$$

Rubi [A] (verified)

Time = 0.31 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 1.167$, Rules

used = {5366, 4511, 4275, 4266, 2611, 2320, 6724, 4269, 3800, 2221, 2317, 2438, 4271, 3855}

$$\begin{aligned}
 \int x^2 \sec^{-1}(a + bx)^3 dx = & \frac{a^3 \sec^{-1}(a + bx)^3}{3b^3} + \frac{6ia^2 \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{6ia^2 \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{6ia^2 \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{6a^2 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{6a^2 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{i \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx)^2}}\right)}{b^3} \\
 & - \frac{i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{3ia \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & + \frac{\operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^3} \\
 & - \frac{(a + bx)^2 \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{2b^3} \\
 & + \frac{3a(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{b^3} \\
 & - \frac{3ia \sec^{-1}(a + bx)^2}{b^3} + \frac{(a + bx) \sec^{-1}(a + bx)}{b^3} \\
 & + \frac{6a \sec^{-1}(a + bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^3} + \frac{1}{3} x^3 \sec^{-1}(a + bx)^3
 \end{aligned}$$

[In] Int[x^2*ArcSec[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSec[a + b*x])/b^3 - ((3*I)*a*ArcSec[a + b*x]^2)/b^3 + (3*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/b^3 - ((a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/(2*b^3) + (a^3*ArcSec[a + b*x]^3)/(3*b^3) + (x^3*ArcSec[a + b*x]^3)/3 + (I*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSe

```

c[a + b*x]))/b^3 + ((6*I)*a^2*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x
]]))/b^3 - ArcTanh[Sqrt[1 - (a + b*x)^(-2)]]/b^3 + (6*a*ArcSec[a + b*x]*Log
[1 + E^((2*I)*ArcSec[a + b*x])])/b^3 - (I*ArcSec[a + b*x]*PolyLog[2, (-I)*E
^(I*ArcSec[a + b*x])])/b^3 - ((6*I)*a^2*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(
I*ArcSec[a + b*x])])/b^3 + (I*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a +
b*x])])/b^3 + ((6*I)*a^2*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x]
])/b^3 - ((3*I)*a*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])])/b^3 + PolyLog[3,
(-I)*E^(I*ArcSec[a + b*x])]/b^3 + (6*a^2*PolyLog[3, (-I)*E^(I*ArcSec[a + b*
x])])/b^3 - PolyLog[3, I*E^(I*ArcSec[a + b*x])]/b^3 - (6*a^2*PolyLog[3, I*E
^(I*ArcSec[a + b*x])])/b^3

```

Rule 2221

```

Int[(((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_)))^(n_))), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

```

Rule 2317

```

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol]
:= Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

```

Rule 2320

```

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

```

Rule 2438

```

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

```

Rule 2611

```

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]

```

Rule 3800

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x)))], x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 4266

Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]

Rule 4269

Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]

Rule 4271

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(-b^2)*(c + d*x)^m*Cot[e + f*x]*((b*Csc[e + f*x])^(n - 2)/(f*(n - 1))), x] + (Dist[b^2*d^2*m*((m - 1)/(f^2*(n - 1)*(n - 2))), Int[(c + d*x)^(m - 2)*(b*Csc[e + f*x])^(n - 2), x], x] + Dist[b^2*((n - 2)/(n - 1)), Int[(c + d*x)^m*(b*Csc[e + f*x])^(n - 2), x], x] - Simp[b^2*d*m*(c + d*x)^(m - 1)*((b*Csc[e + f*x])^(n - 2)/(f^2*(n - 1)*(n - 2))), x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && NeQ[n, 2] && GtQ[m, 1]

Rule 4275

Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 4511

Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_)]^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n + 1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[

{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]

Rule 5366

Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_.)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_.))^(m_.), x_Symbol] :> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IntegerQ[m]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int x^3 \sec(x)(-a + \sec(x))^2 \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{1}{3} x^3 \sec^{-1}(a + bx)^3 - \frac{\text{Subst}\left(\int x^2(-a + \sec(x))^3 dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{1}{3} x^3 \sec^{-1}(a + bx)^3 \\
 &\quad - \frac{\text{Subst}\left(\int (-a^3 x^2 + 3a^2 x^2 \sec(x) - 3ax^2 \sec^2(x) + x^2 \sec^3(x)) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &= \frac{a^3 \sec^{-1}(a + bx)^3}{3b^3} + \frac{1}{3} x^3 \sec^{-1}(a + bx)^3 - \frac{\text{Subst}\left(\int x^2 \sec^3(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &\quad + \frac{(3a) \text{Subst}\left(\int x^2 \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3} \\
 &\quad - \frac{(3a^2) \text{Subst}\left(\int x^2 \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\sec^{-1}(a+bx)}{b^3} + \frac{3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} \\
&\quad - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^3} + \frac{a^3\sec^{-1}(a+bx)^3}{3b^3} \\
&\quad + \frac{1}{3}x^3\sec^{-1}(a+bx)^3 + \frac{6ia^2\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad - \frac{\text{Subst}\left(\int x^2\sec(x)dx, x, \sec^{-1}(a+bx)\right)}{2b^3} - \frac{\text{Subst}\left(\int \sec(x)dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad - \frac{(6a)\text{Subst}\left(\int x\tan(x)dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad + \frac{(6a^2)\text{Subst}\left(\int x\log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad - \frac{(6a^2)\text{Subst}\left(\int x\log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&= \frac{(a+bx)\sec^{-1}(a+bx)}{b^3} - \frac{3ia\sec^{-1}(a+bx)^2}{b^3} \\
&\quad + \frac{3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^3} \\
&\quad + \frac{a^3\sec^{-1}(a+bx)^3}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^3 + \frac{i\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{6ia^2\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{\text{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{b^3} \\
&\quad - \frac{6ia^2\sec^{-1}(a+bx)\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{6ia^2\sec^{-1}(a+bx)\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&\quad + \frac{\text{Subst}\left(\int x\log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad - \frac{\text{Subst}\left(\int x\log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad + \frac{(12ia)\text{Subst}\left(\int \frac{e^{2ix}x}{1+e^{2ix}}dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad + \frac{(6ia^2)\text{Subst}\left(\int \text{PolyLog}\left(2, -ie^{ix}\right)dx, x, \sec^{-1}(a+bx)\right)}{b^3} \\
&\quad - \frac{(6ia^2)\text{Subst}\left(\int \text{PolyLog}\left(2, ie^{ix}\right)dx, x, \sec^{-1}(a+bx)\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\sec^{-1}(a+bx)}{b^3} - \frac{3ia\sec^{-1}(a+bx)^2}{b^3} \\
&+ \frac{3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^3} \\
&+ \frac{a^3\sec^{-1}(a+bx)^3}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^3 + \frac{i\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{b^3} \\
&+ \frac{6a\sec^{-1}(a+bx)\log\left(1+e^{2i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{i\operatorname{Subst}\left(\int\operatorname{PolyLog}\left(2,-ie^{ix}\right)dx,x,\sec^{-1}(a+bx)\right)}{b^3} \\
&- \frac{i\operatorname{Subst}\left(\int\operatorname{PolyLog}\left(2,ie^{ix}\right)dx,x,\sec^{-1}(a+bx)\right)}{b^3} \\
&- \frac{(6a)\operatorname{Subst}\left(\int\log\left(1+e^{2ix}\right)dx,x,\sec^{-1}(a+bx)\right)}{b^3} \\
&+ \frac{(6a^2)\operatorname{Subst}\left(\int\frac{\operatorname{PolyLog}(2,-ix)}{x}dx,x,e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{(6a^2)\operatorname{Subst}\left(\int\frac{\operatorname{PolyLog}(2,ix)}{x}dx,x,e^{i\sec^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\sec^{-1}(a+bx)}{b^3} - \frac{3ia\sec^{-1}(a+bx)^2}{b^3} \\
&+ \frac{3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^3} \\
&+ \frac{a^3\sec^{-1}(a+bx)^3}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^3 + \frac{i\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{b^3} \\
&+ \frac{6a\sec^{-1}(a+bx)\log\left(1+e^{2i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{6a^2\operatorname{PolyLog}\left(3,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{6a^2\operatorname{PolyLog}\left(3,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{\operatorname{Subst}\left(\int\frac{\operatorname{PolyLog}(2,-ix)}{x}dx,x,e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{\operatorname{Subst}\left(\int\frac{\operatorname{PolyLog}(2,ix)}{x}dx,x,e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{(3ia)\operatorname{Subst}\left(\int\frac{\log(1+x)}{x}dx,x,e^{2i\sec^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\sec^{-1}(a+bx)}{b^3} - \frac{3ia\sec^{-1}(a+bx)^2}{b^3} \\
&+ \frac{3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} - \frac{(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^3} \\
&+ \frac{a^3\sec^{-1}(a+bx)^3}{3b^3} + \frac{1}{3}x^3\sec^{-1}(a+bx)^3 + \frac{i\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{arctanh}\left(\sqrt{1-\frac{1}{(a+bx)^2}}\right)}{b^3} \\
&+ \frac{6a\sec^{-1}(a+bx)\log\left(1+e^{2i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{6ia^2\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{3ia\operatorname{PolyLog}\left(2,-e^{2i\sec^{-1}(a+bx)}\right)}{b^3} \\
&+ \frac{\operatorname{PolyLog}\left(3,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{6a^2\operatorname{PolyLog}\left(3,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} \\
&- \frac{\operatorname{PolyLog}\left(3,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{6a^2\operatorname{PolyLog}\left(3,ie^{i\sec^{-1}(a+bx)}\right)}{b^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 446, normalized size of antiderivative = 0.90

$$\int x^2 \sec^{-1}(a+bx)^3 dx$$

$$= \frac{(a+bx)\sec^{-1}(a+bx) - 3ia\sec^{-1}(a+bx)^2 + 3a(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2 - \frac{1}{2}(a+bx)^2\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{b^3} - \frac{3ia\operatorname{PolyLog}\left(2,-e^{2i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{\operatorname{PolyLog}\left(3,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} + \frac{6a^2\operatorname{PolyLog}\left(3,-ie^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{\operatorname{PolyLog}\left(3,ie^{i\sec^{-1}(a+bx)}\right)}{b^3} - \frac{6a^2\operatorname{PolyLog}\left(3,ie^{i\sec^{-1}(a+bx)}\right)}{b^3}$$

[In] Integrate[x^2*ArcSec[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSec[a + b*x] - (3*I)*a*ArcSec[a + b*x]^2 + 3*a*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2 - ((a + b*x)^2*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/2 + (a^3*ArcSec[a + b*x]^3)/3 + (b^3*x^3*ArcSec[a + b*x]^3)/3 + I*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] + (6*I)*a^2*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] - ArcTanh[Sqrt[1 - (a + b*x)^(-2)]]

$$\begin{aligned} & \text{^(-2)]}] + 6*a*\text{ArcSec}[a + b*x]*\text{Log}[1 + \text{E}^{\text{((2*I)*\text{ArcSec}[a + b*x])}] - I*\text{ArcSec} \\ & [a + b*x]*\text{PolyLog}[2, (-I)*\text{E}^{\text{(I*\text{ArcSec}[a + b*x])}] - (6*I)*a^2*\text{ArcSec}[a + b*x] \\ &]*\text{PolyLog}[2, (-I)*\text{E}^{\text{(I*\text{ArcSec}[a + b*x])}] + I*\text{ArcSec}[a + b*x]*\text{PolyLog}[2, I*\text{E} \\ & \text{^{\text{(I*\text{ArcSec}[a + b*x])}] + (6*I)*a^2*\text{ArcSec}[a + b*x]*\text{PolyLog}[2, I*\text{E}^{\text{(I*\text{ArcSec}[} \\ & a + b*x])}] - (3*I)*a*\text{PolyLog}[2, -\text{E}^{\text{((2*I)*\text{ArcSec}[a + b*x])}] + \text{PolyLog}[3, (- \\ & I)*\text{E}^{\text{(I*\text{ArcSec}[a + b*x])}] + 6*a^2*\text{PolyLog}[3, (-I)*\text{E}^{\text{(I*\text{ArcSec}[a + b*x])}] - \\ & \text{PolyLog}[3, I*\text{E}^{\text{(I*\text{ArcSec}[a + b*x])}] - 6*a^2*\text{PolyLog}[3, I*\text{E}^{\text{(I*\text{ArcSec}[a + b*} \\ & x])}]])]/b^3 \end{aligned}$$

Maple [A] (verified)

Time = 1.26 (sec) , antiderivative size = 716, normalized size of antiderivative = 1.45

method	result
derivativedivides	$\frac{\text{arcsec}(bx+a) \left(6 \text{arcsec}(bx+a)^2 a^2 (bx+a) - 6 \text{arcsec}(bx+a)^2 a (bx+a)^2 + 2 \text{arcsec}(bx+a)^2 (bx+a)^3 + 18 \text{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) \right)}{6}$
default	$\frac{\text{arcsec}(bx+a) \left(6 \text{arcsec}(bx+a)^2 a^2 (bx+a) - 6 \text{arcsec}(bx+a)^2 a (bx+a)^2 + 2 \text{arcsec}(bx+a)^2 (bx+a)^3 + 18 \text{arcsec}(bx+a) \sqrt{\frac{(bx+a)^2 - 1}{(bx+a)^2}} a (bx+a) \right)}{6}$

[In] int(x^2*arcsec(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{b^3} \left(\frac{1}{6} \text{arcsec}(b*x+a) * (6*\text{arcsec}(b*x+a)^2*a^2*(b*x+a) - 6*\text{arcsec}(b*x+a)^2*a*(b*x+a)^2 + 2*\text{arcsec}(b*x+a)^2*(b*x+a)^3 + 18*\text{arcsec}(b*x+a)*\left(\frac{(b*x+a)^2 - 1}{(b*x+a)^2}\right)^{1/2} * a*(b*x+a) - 3*\text{arcsec}(b*x+a)*\left(\frac{(b*x+a)^2 - 1}{(b*x+a)^2}\right)^{1/2} * (b*x+a)^2 + 18*I*a*\text{arcsec}(b*x+a) + 6*b*x + 6*a) + 2*I*\text{arctan}\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2} + I*\text{arcsec}(b*x+a)*\text{polylog}\left(2, I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) + 3*\ln\left(1 + I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) \right) * a^2*\text{arcsec}(b*x+a)^2 + 6*\text{polylog}\left(3, -I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a^2 - I*\text{arcsec}(b*x+a)*\text{polylog}\left(2, -I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) - 3*\ln\left(1 - I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) \right) * a^2*\text{arcsec}(b*x+a)^2 + 6*I*\text{polylog}\left(2, I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a^2*\text{arcsec}(b*x+a) - 6*\text{polylog}\left(3, I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a^2 + 6*\ln\left(1 + \left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a*\text{arcsec}(b*x+a) - 6*I*\text{polylog}\left(2, -I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a^2*\text{arcsec}(b*x+a) - 3*I*\text{polylog}\left(2, -\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) * a + \frac{1}{2}*\text{arcsec}(b*x+a)^2*\ln\left(1 + I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) - 6*I*\text{arcsec}(b*x+a)^2*a + \text{polylog}\left(3, -I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) - \frac{1}{2}*\text{arcsec}(b*x+a)^2*\ln\left(1 - I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) - \text{polylog}\left(3, I*\left(\frac{1}{(b*x+a)} + I*(1 - 1/(b*x+a)^2)\right)^{1/2}\right) \right)$

Fricas [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

```
[In] integrate(x^2*arcsec(b*x+a)^3,x, algorithm="fricas")
```

```
[Out] integral(x^2*arcsec(b*x + a)^3, x)
```

Sympy [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{asec}^3(a + bx) dx$$

```
[In] integrate(x**2*asec(b*x+a)**3,x)
```

```
[Out] Integral(x**2*asec(a + b*x)**3, x)
```

Maxima [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

```
[In] integrate(x^2*arcsec(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] 1/3*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 1/4*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(1/4*((4*b*x^3*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x^3*log(b^2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) + 4*(3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a)^2 - (b^3*x^5 + 2*a*b^2*x^4 + (a^2 - 1)*b*x^3 + 3*(b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)
```

Giac [F]

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(x^2*arcsec(b*x+a)^3,x, algorithm="giac")

[Out] integrate(x^2*arcsec(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \sec^{-1}(a + bx)^3 dx = \int x^2 \operatorname{acos}\left(\frac{1}{a + bx}\right)^3 dx$$

[In] int(x^2*acos(1/(a + b*x))^3,x)

[Out] int(x^2*acos(1/(a + b*x))^3, x)

3.34 $\int x \sec^{-1}(a + bx)^3 dx$

Optimal result	232
Rubi [A] (verified)	233
Mathematica [A] (verified)	238
Maple [A] (verified)	239
Fricas [F]	239
Sympy [F]	239
Maxima [F]	240
Giac [F]	240
Mupad [F(-1)]	240

Optimal result

Integrand size = 10, antiderivative size = 278

$$\begin{aligned}
 \int x \sec^{-1}(a + bx)^3 dx = & \frac{3i \sec^{-1}(a + bx)^2}{2b^2} - \frac{3(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{2b^2} \\
 & - \frac{a^2 \sec^{-1}(a + bx)^3}{2b^2} + \frac{1}{2} x^2 \sec^{-1}(a + bx)^3 \\
 & - \frac{6ia \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{3 \sec^{-1}(a + bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{6ia \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & - \frac{6ia \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
 & + \frac{3i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2b^2} \\
 & - \frac{6a \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{6a \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^2}
 \end{aligned}$$

```

[Out] 3/2*I*arcsec(b*x+a)^2/b^2-1/2*a^2*arcsec(b*x+a)^3/b^2+1/2*x^2*arcsec(b*x+a)
^3-6*I*a*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b^2-3*ar
csec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/b^2+6*I*a*arcsec(b*
x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2-6*I*a*arcsec(b*x
+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b^2+3/2*I*polylog(2,-(
1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/b^2-6*a*polylog(3,-I*(1/(b*x+a)+I*(1-

```


$1/(b*x+a)^2)^{(1/2)})/b^2+6*a*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^{(1/2)})/b^2-3/2*(b*x+a)*arcsec(b*x+a)^2*(1-1/(b*x+a)^2)^{(1/2)}/b^2$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 278, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 1.200$, Rules used = {5366, 4511, 4275, 4266, 2611, 2320, 6724, 4269, 3800, 2221, 2317, 2438}

$$\int x \sec^{-1}(a + bx)^3 dx = -\frac{a^2 \sec^{-1}(a + bx)^3}{2b^2} - \frac{6ia \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{6ia \sec^{-1}(a + bx) \text{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{6ia \sec^{-1}(a + bx) \text{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{3i \text{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2b^2} - \frac{6a \text{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{6a \text{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{3(a + bx) \sqrt{1 - \frac{1}{(a+bx)^2}} \sec^{-1}(a + bx)^2}{2b^2} + \frac{3i \sec^{-1}(a + bx)^2}{2b^2} - \frac{3 \sec^{-1}(a + bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{1}{2} x^2 \sec^{-1}(a + bx)^3$$

[In] Int[x*ArcSec[a + b*x]^3,x]

[Out] (((3*I)/2)*ArcSec[a + b*x]^2)/b^2 - (3*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/(2*b^2) - (a^2*ArcSec[a + b*x]^3)/(2*b^2) + (x^2*ArcSec[a + b*x]^3)/2 - ((6*I)*a*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])])/b^2 - (3*ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])])/b^2 + ((6*I)*a*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])])/b^2 - ((6*I)*a*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b^2 + (((3*I)/2)*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])])/b^2 - (6*a*PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])])/b^2 + (6*a*PolyLog[3, I*E^(I*ArcSec[a + b*x])])/b^2

Rule 2221

Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a], x]

```
)^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] :> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] :> Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]
```

Rule 4266

```
Int[csc[(e_) + Pi*(k_) + (f_)*(x_)]*((c_) + (d_)*(x_)^(m_)), x_Symbol]
:> Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

Rule 4269

```
Int[csc[(e_.) + (f_.)*(x_)]^2*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp
[(-(c + d*x)^m)*(Cot[e + f*x]/f), x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*
Cot[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && GtQ[m, 0]
```

Rule 4275

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Csc[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_)]^(n_.)*Tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int x^3 \sec(x)(-a + \sec(x)) \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2} \\
&= \frac{1}{2}x^2 \sec^{-1}(a + bx)^3 - \frac{3\text{Subst}\left(\int x^2(-a + \sec(x))^2 dx, x, \sec^{-1}(a + bx)\right)}{2b^2} \\
&= \frac{1}{2}x^2 \sec^{-1}(a + bx)^3 - \frac{3\text{Subst}\left(\int (a^2x^2 - 2ax^2 \sec(x) + x^2 \sec^2(x)) dx, x, \sec^{-1}(a + bx)\right)}{2b^2} \\
&= -\frac{a^2 \sec^{-1}(a + bx)^3}{2b^2} + \frac{1}{2}x^2 \sec^{-1}(a + bx)^3 - \frac{3\text{Subst}\left(\int x^2 \sec^2(x) dx, x, \sec^{-1}(a + bx)\right)}{2b^2} \\
&\quad + \frac{(3a)\text{Subst}\left(\int x^2 \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^2} - \frac{a^2\sec^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{1}{2}x^2\sec^{-1}(a+bx)^3 - \frac{6ia\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{3\text{Subst}\left(\int x\tan(x)dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&- \frac{(6a)\text{Subst}\left(\int x\log(1-ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&+ \frac{(6a)\text{Subst}\left(\int x\log(1+ie^{ix})dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&= \frac{3i\sec^{-1}(a+bx)^2}{2b^2} - \frac{3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}\sec^{-1}(a+bx)^2}{2b^2} - \frac{a^2\sec^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{1}{2}x^2\sec^{-1}(a+bx)^3 - \frac{6ia\sec^{-1}(a+bx)^2\arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6ia\sec^{-1}(a+bx)\text{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{6ia\sec^{-1}(a+bx)\text{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{(6i)\text{Subst}\left(\int \frac{e^{2ix}x}{1+e^{2ix}}dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&- \frac{(6ia)\text{Subst}\left(\int \text{PolyLog}\left(2, -ie^{ix}\right)dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&+ \frac{(6ia)\text{Subst}\left(\int \text{PolyLog}\left(2, ie^{ix}\right)dx, x, \sec^{-1}(a+bx)\right)}{b^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3i \sec^{-1}(a+bx)^2}{2b^2} - \frac{3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{2b^2} - \frac{a^2 \sec^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{\frac{1}{2}x^2 \sec^{-1}(a+bx)^3}{b^2} - \frac{6ia \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{3 \sec^{-1}(a+bx) \log\left(1+e^{2i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{3 \operatorname{Subst}\left(\int \log(1+e^{2ix}) dx, x, \sec^{-1}(a+bx)\right)}{b^2} \\
&- \frac{(6a) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -ix)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{(6a) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, ix)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&= \frac{3i \sec^{-1}(a+bx)^2}{2b^2} - \frac{3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{2b^2} - \frac{a^2 \sec^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{\frac{1}{2}x^2 \sec^{-1}(a+bx)^3}{b^2} - \frac{6ia \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{3 \sec^{-1}(a+bx) \log\left(1+e^{2i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{6a \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6a \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} - \frac{(3i) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \sec^{-1}(a+bx)}\right)}{2b^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{3i \sec^{-1}(a+bx)^2}{2b^2} - \frac{3(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2}{2b^2} - \frac{a^2 \sec^{-1}(a+bx)^3}{2b^2} \\
&+ \frac{1}{2}x^2 \sec^{-1}(a+bx)^3 - \frac{6ia \sec^{-1}(a+bx)^2 \arctan\left(e^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{3 \sec^{-1}(a+bx) \log\left(1+e^{2i \sec^{-1}(a+bx)}\right)}{b^2} \\
&+ \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} \\
&- \frac{6ia \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{3i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2b^2} \\
&- \frac{6a \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a+bx)}\right)}{b^2} + \frac{6a \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a+bx)}\right)}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 257, normalized size of antiderivative = 0.92

$$\int x \sec^{-1}(a+bx)^3 dx$$

$$= \frac{\frac{3}{2}i \sec^{-1}(a+bx)^2 - \frac{3}{2}(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}} \sec^{-1}(a+bx)^2 - a(a+bx) \sec^{-1}(a+bx)^3 + \frac{1}{2}(a+bx)^2 \sec^{-1}(a+bx)}{b^2}$$

[In] Integrate[x*ArcSec[a + b*x]^3,x]

[Out] (((3*I)/2)*ArcSec[a + b*x]^2 - (3*(a + b*x)*Sqrt[1 - (a + b*x)^(-2)]*ArcSec[a + b*x]^2)/2 - a*(a + b*x)*ArcSec[a + b*x]^3 + ((a + b*x)^2*ArcSec[a + b*x]^3)/2 - (6*I)*a*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])] - 3*ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])] + (6*I)*a*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - (6*I)*a*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])] + ((3*I)/2)*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] - 6*a*PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])] + 6*a*PolyLog[3, I*E^(I*ArcSec[a + b*x])])]/b^2

Maple [A] (verified)

Time = 1.45 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.36

method	result
derivativedivides	$-\frac{\operatorname{arcsec}(bx+a)^2 \left(2 \operatorname{arcsec}(bx+a)a(bx+a) - \operatorname{arcsec}(bx+a)(bx+a)^2 + 3\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}(bx+a)+3i \right)}{2} - 3 \ln \left(1+i \left(\frac{1}{bx+a} + i \sqrt{1-\frac{1}{(bx+a)^2}} \right) \right)$
default	$-\frac{\operatorname{arcsec}(bx+a)^2 \left(2 \operatorname{arcsec}(bx+a)a(bx+a) - \operatorname{arcsec}(bx+a)(bx+a)^2 + 3\sqrt{\frac{(bx+a)^2-1}{(bx+a)^2}}(bx+a)+3i \right)}{2} - 3 \ln \left(1+i \left(\frac{1}{bx+a} + i \sqrt{1-\frac{1}{(bx+a)^2}} \right) \right)$

[In] int(x*arcsec(b*x+a)^3,x,method=_RETURNVERBOSE)

```
[Out] 1/b^2*(-1/2*arcsec(b*x+a)^2*(2*arcsec(b*x+a)*a*(b*x+a)-arcsec(b*x+a)*(b*x+a)^2+3*(((b*x+a)^2-1)/(b*x+a)^2)^(1/2)*(b*x+a)+3*I)-3*ln(1+I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a*arcsec(b*x+a)^2+6*I*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a*arcsec(b*x+a)-6*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a+3*ln(1-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a*arcsec(b*x+a)^2-6*I*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a*arcsec(b*x+a)+6*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))))*a+3*I*arcsec(b*x+a)^2-3*arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+3/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2))
```

Fricas [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(x*arcsec(b*x+a)^3,x, algorithm="fricas")

[Out] integral(x*arcsec(b*x + a)^3, x)

Sympy [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{asec}^3(a + bx) dx$$

[In] integrate(x*asec(b*x+a)**3,x)

[Out] Integral(x*asec(a + b*x)**3, x)

Maxima [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(x*arcsec(b*x+a)^3,x, algorithm="maxima")

[Out] $\frac{1}{2}x^2 \arctan(\sqrt{bx + a + 1} \sqrt{bx + a - 1})^3 - \frac{3}{8}x^2 \arctan(\sqrt{bx + a + 1} \sqrt{bx + a - 1}) \log(b^2x^2 + 2abx + a^2)^2 - \int \frac{3}{8}((4b^2x^2 \arctan(\sqrt{bx + a + 1} \sqrt{bx + a - 1})^2 - b^2x^2 \log(b^2x^2 + 2abx + a^2)^2) \sqrt{bx + a + 1} \sqrt{bx + a - 1} + 4(2(b^3x^4 + 3ab^2x^3 + (3a^2 - 1)bx^2 + (a^3 - a)x) \log(bx + a)^2 - (b^3x^4 + 2ab^2x^3 + (a^2 - 1)bx^2 + 2(b^3x^4 + 3ab^2x^3 + (3a^2 - 1)bx^2 + (a^3 - a)x) \log(bx + a)) \log(b^2x^2 + 2abx + a^2)) \arctan(\sqrt{bx + a + 1} \sqrt{bx + a - 1}))}{(b^3x^3 + 3ab^2x^2 + a^3 + (3a^2 - 1)bx - a), x)$

Giac [F]

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(x*arcsec(b*x+a)^3,x, algorithm="giac")

[Out] integrate(x*arcsec(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int x \sec^{-1}(a + bx)^3 dx = \int x \arccos\left(\frac{1}{a + bx}\right)^3 dx$$

[In] int(x*acos(1/(a + b*x))^3,x)

[Out] int(x*acos(1/(a + b*x))^3, x)

3.35 $\int \sec^{-1}(a + bx)^3 dx$

Optimal result	241
Rubi [A] (verified)	241
Mathematica [A] (verified)	245
Maple [F]	245
Fricas [F]	245
Sympy [F]	245
Maxima [F]	246
Giac [F]	246
Mupad [F(-1)]	246

Optimal result

Integrand size = 8, antiderivative size = 154

$$\int \sec^{-1}(a + bx)^3 dx = \frac{(a + bx) \sec^{-1}(a + bx)^3}{b} + \frac{6i \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$- \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$+ \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

$$+ \frac{6 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{6 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a + bx)}\right)}{b}$$

```
[Out] (b*x+a)*arcsec(b*x+a)^3/b+6*I*arcsec(b*x+a)^2*arctan(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/b-6*I*arcsec(b*x+a)*polylog(2,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+6*I*arcsec(b*x+a)*polylog(2,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b+6*polylog(3,-I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b-6*polylog(3,I*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2)))/b
```

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used

= {5360, 5324, 3842, 4266, 2611, 2320, 6724}

$$\int \sec^{-1}(a + bx)^3 dx = \frac{6i \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, -ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{6i \sec^{-1}(a + bx) \operatorname{PolyLog}\left(2, ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{6 \operatorname{PolyLog}\left(3, -ie^{i \sec^{-1}(a + bx)}\right)}{b} - \frac{6 \operatorname{PolyLog}\left(3, ie^{i \sec^{-1}(a + bx)}\right)}{b} + \frac{(a + bx) \sec^{-1}(a + bx)^3}{b}$$

[In] Int[ArcSec[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSec[a + b*x]^3)/b + ((6*I)*ArcSec[a + b*x]^2*ArcTan[E^(I*ArcSec[a + b*x])])/b - ((6*I)*ArcSec[a + b*x]*PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])])/b + ((6*I)*ArcSec[a + b*x]*PolyLog[2, I*E^(I*ArcSec[a + b*x])])/b + (6*PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])])/b - (6*PolyLog[3, I*E^(I*ArcSec[a + b*x])])/b

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_) /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3842

Int[(x_)^(m_.)*Sec[(a_.) + (b_.)*(x_)^(n_.)]^(p_.)*Tan[(a_.) + (b_.)*(x_)^(n_.)]^(q_.), x_Symbol] := Simp[x^(m - n + 1)*(Sec[a + b*x^n]^p/(b*n*p)), x] - Dist[(m - n + 1)/(b*n*p), Int[x^(m - n)*Sec[a + b*x^n]^p, x], x] /; FreeQ[{a, b, p}, x] && IntegerQ[n] && GeQ[m, n] && EqQ[q, 1]

Rule 4266

```
Int[csc[(e_.) + Pi*(k_.) + (f_.)*(x_)]*((c_.) + (d_.)*(x_))^(m_.), x_Symbol]
  := Simp[-2*(c + d*x)^m*(ArcTanh[E^(I*k*Pi)*E^(I*(e + f*x))]/f), x] + (-Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 - E^(I*k*Pi)*E^(I*(e + f*x))], x], x] + Dist[d*(m/f), Int[(c + d*x)^(m - 1)*Log[1 + E^(I*k*Pi)*E^(I*(e + f*x))], x], x]) /; FreeQ[{c, d, e, f}, x] && IntegerQ[2*k] && IGtQ[m, 0]
```

Rule 5324

```
Int[((a_.) + ArcSec[(c_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[1/c, Subst[Int[(a + b*x)^n*Sec[x]*Tan[x], x], x, ArcSec[c*x]], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[n, 0]
```

Rule 5360

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)]*(b_.))^(p_.), x_Symbol] := Dist[1/d, Subst[Int[(a + b*ArcSec[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d}, x] && IGtQ[p, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \sec^{-1}(x)^3 dx, x, a + bx\right)}{b} \\
 &= \frac{\text{Subst}\left(\int x^3 \sec(x) \tan(x) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)^3}{b} - \frac{3 \text{Subst}\left(\int x^2 \sec(x) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
 &= \frac{(a + bx) \sec^{-1}(a + bx)^3}{b} + \frac{6i \sec^{-1}(a + bx)^2 \arctan\left(e^{i \sec^{-1}(a + bx)}\right)}{b} \\
 &\quad + \frac{6 \text{Subst}\left(\int x \log(1 - ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b} \\
 &\quad - \frac{6 \text{Subst}\left(\int x \log(1 + ie^{ix}) dx, x, \sec^{-1}(a + bx)\right)}{b}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{(a+bx)\sec^{-1}(a+bx)^3}{b} + \frac{6i\sec^{-1}(a+bx)^2 \arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad - \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{(6i)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, -ie^{ix}\right) dx, x, \sec^{-1}(a+bx)\right)}{b} \\
&\quad - \frac{(6i)\operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, ie^{ix}\right) dx, x, \sec^{-1}(a+bx)\right)}{b} \\
&= \frac{(a+bx)\sec^{-1}(a+bx)^3}{b} + \frac{6i\sec^{-1}(a+bx)^2 \arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad - \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{6\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -ix)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad - \frac{6\operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, ix)}{x} dx, x, e^{i\sec^{-1}(a+bx)}\right)}{b} \\
&= \frac{(a+bx)\sec^{-1}(a+bx)^3}{b} + \frac{6i\sec^{-1}(a+bx)^2 \arctan\left(e^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad - \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, -ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{6i\sec^{-1}(a+bx)\operatorname{PolyLog}\left(2, ie^{i\sec^{-1}(a+bx)}\right)}{b} \\
&\quad + \frac{6\operatorname{PolyLog}\left(3, -ie^{i\sec^{-1}(a+bx)}\right)}{b} - \frac{6\operatorname{PolyLog}\left(3, ie^{i\sec^{-1}(a+bx)}\right)}{b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 160, normalized size of antiderivative = 1.04

$$\int \sec^{-1}(a + bx)^3 dx$$

$$= \frac{(a + bx) \sec^{-1}(a + bx)^3 - 3 \sec^{-1}(a + bx)^2 \left(\log \left(1 - ie^{i \sec^{-1}(a + bx)} \right) - \log \left(1 + ie^{i \sec^{-1}(a + bx)} \right) \right) - 6i \sec^{-1}}{b}$$

[In] Integrate[ArcSec[a + b*x]^3,x]

[Out] ((a + b*x)*ArcSec[a + b*x]^3 - 3*ArcSec[a + b*x]^2*(Log[1 - I*E^(I*ArcSec[a + b*x])] - Log[1 + I*E^(I*ArcSec[a + b*x])]) - (6*I)*ArcSec[a + b*x]*(PolyLog[2, (-I)*E^(I*ArcSec[a + b*x])] - PolyLog[2, I*E^(I*ArcSec[a + b*x])]) + 6*(PolyLog[3, (-I)*E^(I*ArcSec[a + b*x])] - PolyLog[3, I*E^(I*ArcSec[a + b*x])]))/b

Maple [F]

$$\int \operatorname{arcsec}(bx + a)^3 dx$$

[In] int(arcsec(b*x+a)^3,x)

[Out] int(arcsec(b*x+a)^3,x)

Fricas [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(arcsec(b*x+a)^3,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)^3, x)

Sympy [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{asec}^3(a + bx) dx$$

[In] integrate(asec(b*x+a)**3,x)

[Out] Integral(asec(a + b*x)**3, x)

Maxima [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(arcsec(b*x+a)^3,x, algorithm="maxima")

[Out] x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^3 - 3/4*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))*log(b^2*x^2 + 2*a*b*x + a^2)^2 - integrate(3/4*((4*b*x*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1))^2 - b*x*log(b^2*x^2 + 2*a*b*x + a^2)^2)*sqrt(b*x + a + 1)*sqrt(b*x + a - 1) + 4*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a)^2 - (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x + (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*log(b*x + a))*log(b^2*x^2 + 2*a*b*x + a^2))*arctan(sqrt(b*x + a + 1)*sqrt(b*x + a - 1)))/(b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a), x)

Giac [F]

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{arcsec}(bx + a)^3 dx$$

[In] integrate(arcsec(b*x+a)^3,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(a + bx)^3 dx = \int \operatorname{acos}\left(\frac{1}{a + bx}\right)^3 dx$$

[In] int(acos(1/(a + b*x))^3,x)

[Out] int(acos(1/(a + b*x))^3, x)

3.36 $\int \frac{\sec^{-1}(a+bx)^3}{x} dx$

Optimal result	247
Rubi [A] (verified)	248
Mathematica [B] (warning: unable to verify)	255
Maple [F]	257
Fricas [F]	258
Sympy [F]	258
Maxima [F]	258
Giac [F]	258
Mupad [F(-1)]	259

Optimal result

Integrand size = 12, antiderivative size = 430

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^3}{x} dx &= \sec^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + \sec^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \sec^{-1}(a+bx)^3 \log\left(1 + e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad - 3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad - 3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad + \frac{3}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad + 6 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) \\
&\quad + 6 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{2} \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2i\sec^{-1}(a+bx)}\right) \\
&\quad + 6i \operatorname{PolyLog}\left(4, \frac{ae^{i\sec^{-1}(a+bx)}}{1 - \sqrt{1-a^2}}\right) + 6i \operatorname{PolyLog}\left(4, \frac{ae^{i\sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}}\right) \\
&\quad - \frac{3}{4}i \operatorname{PolyLog}\left(4, -e^{2i\sec^{-1}(a+bx)}\right)
\end{aligned}$$

```
[Out] -arcsec(b*x+a)^3*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+arcsec(b*x+a)^
3*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))+arcsec(b*x
+a)^3*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))+3/2*I*
arcsec(b*x+a)^2*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)-3*I*arcse
c(b*x+a)^2*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2
)))-3*I*arcsec(b*x+a)^2*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+
(-a^2+1)^(1/2)))-3/2*arcsec(b*x+a)*polylog(3,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(
1/2))^2)+6*arcsec(b*x+a)*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(
1-(-a^2+1)^(1/2)))+6*arcsec(b*x+a)*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)
^(1/2))/(1+(-a^2+1)^(1/2)))-3/4*I*polylog(4,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(
1/2))^2)+6*I*polylog(4,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1
/2)))+6*I*polylog(4,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2
))
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 430, normalized size of antiderivative = 1.00,
number of steps used = 20, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules

used = {5366, 4647, 4626, 3800, 2221, 2611, 6744, 2320, 6724, 4616}

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^3}{x} dx = & -3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
& - 3i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
& + 6 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
& + 6 \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
& + 6i \operatorname{PolyLog}\left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) + 6i \operatorname{PolyLog}\left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
& + \sec^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right) \\
& + \sec^{-1}(a+bx)^3 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right) \\
& + \frac{3}{2}i \sec^{-1}(a+bx)^2 \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) \\
& - \frac{3}{2} \sec^{-1}(a+bx) \operatorname{PolyLog}\left(3, -e^{2i \sec^{-1}(a+bx)}\right) \\
& - \frac{3}{4}i \operatorname{PolyLog}\left(4, -e^{2i \sec^{-1}(a+bx)}\right) \\
& - \sec^{-1}(a+bx)^3 \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)
\end{aligned}$$

[In] Int[ArcSec[a + b*x]^3/x, x]

[Out] ArcSec[a + b*x]^3*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + ArcSec[a + b*x]^3*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - ArcSec[a + b*x]^3*Log[1 + E^((2*I)*ArcSec[a + b*x])] - (3*I)*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] - (3*I)*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + ((3*I)/2)*ArcSec[a + b*x]^2*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])] + 6*ArcSec[a + b*x]*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + 6*ArcSec[a + b*x]*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - (3*ArcSec[a + b*x]*PolyLog[3, -E^((2*I)*ArcSec[a + b*x])])/2 + (6*I)*PolyLog[4, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])] + (6*I)*PolyLog[4, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - ((3*I)/4)*PolyLog[4, -E^((2*I)*ArcSec[a + b*x])]

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*(f_) + (g_)
*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3800

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 4616

```
Int[(((e_) + (f_)*(x_))^(m_)*Sin[(c_) + (d_)*(x_)])/(Cos[(c_) + (d_)
*(x_)]*(b_) + (a_)), x_Symbol] := Simp[I*((e + f*x)^(m + 1)/(b*f*(m + 1))
, x] + (-Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a - Rt[a^2 - b^2, 2] + b
*E^(I*(c + d*x))), x], x] - Dist[I, Int[(e + f*x)^m*(E^(I*(c + d*x)))/(a +
Rt[a^2 - b^2, 2] + b*E^(I*(c + d*x))), x], x]) /; FreeQ[{a, b, c, d, e, f}
, x] && IGtQ[m, 0] && PosQ[a^2 - b^2]
```

Rule 4626

```
Int[(((e_) + (f_)*(x_))^(m_)*Tan[(c_) + (d_)*(x_)]^(n_))/(Cos[(c_) +
(d_)*(x_)]*(b_) + (a_)), x_Symbol] := Dist[1/a, Int[(e + f*x)^m*Tan[c +
d*x]^n, x], x] - Dist[b/a, Int[(e + f*x)^m*Sin[c + d*x]*(Tan[c + d*x]^(n -
1)/(a + b*Cos[c + d*x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[m,
0] && IGtQ[n, 0]
```

Rule 4647

```
Int[(((e_.) + (f_.)*(x_.))^(m_.)*(F_)[(c_.) + (d_.)*(x_.)]^(n_.)*(G_)[(c_.) +
(d_.)*(x_.)]^(p_.))/((a_.) + (b_.)*Sec[(c_.) + (d_.)*(x_.)]), x_Symbol] := In
t[(e + f*x)^m*Cos[c + d*x]*F[c + d*x]^n*(G[c + d*x]^p/(b + a*Cos[c + d*x]))
, x] /; FreeQ[{a, b, c, d, e, f}, x] && TrigQ[F] && TrigQ[G] && IntegersQ[m
, n, p]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_) + (d_.)*(x_)])*(b_.)^(p_.)*((e_.) + (f_.)*(x_.))^(m
_.), x_Symbol] := Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
e - c*f + f*Sec[x])^m, x], x, ArcSec[c + d*x]], x] /; FreeQ[{a, b, c, d, e,
f}, x] && IGtQ[p, 0] && IntegerQ[m]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{x^3 \sec(x) \tan(x)}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= \text{Subst}\left(\int \frac{x^3 \tan(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) \\
&= a \text{Subst}\left(\int \frac{x^3 \sin(x)}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx)\right) + \text{Subst}\left(\int x^3 \tan(x) dx, x, \sec^{-1}(a + bx)\right) \\
&= -\left(2i \text{Subst}\left(\int \frac{e^{2ix} x^3}{1 + e^{2ix}} dx, x, \sec^{-1}(a + bx)\right)\right) \\
&\quad - (ia) \text{Subst}\left(\int \frac{e^{ix} x^3}{1 - \sqrt{1 - a^2} - ae^{ix}} dx, x, \sec^{-1}(a + bx)\right) \\
&\quad - (ia) \text{Subst}\left(\int \frac{e^{ix} x^3}{1 + \sqrt{1 - a^2} - ae^{ix}} dx, x, \sec^{-1}(a + bx)\right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3 \text{Subst} \left(\int x^2 \log \left(1 - \frac{ae^{ix}}{1 - \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad - 3 \text{Subst} \left(\int x^2 \log \left(1 - \frac{ae^{ix}}{1 + \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad + 3 \text{Subst} \left(\int x^2 \log (1 + e^{2ix}) dx, x, \sec^{-1}(a + bx) \right) \\
&= \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad + \frac{3}{2} i \sec^{-1}(a + bx)^2 \text{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3i \text{Subst} \left(\int x \text{PolyLog} (2, -e^{2ix}) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad + 6i \text{Subst} \left(\int x \text{PolyLog} \left(2, \frac{ae^{ix}}{1 - \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad + 6i \text{Subst} \left(\int x \text{PolyLog} \left(2, \frac{ae^{ix}}{1 + \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad + \frac{3}{2} i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \frac{3}{2} \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + \frac{3}{2} \operatorname{Subst} \left(\int \operatorname{PolyLog} (3, -e^{2ix}) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad - 6 \operatorname{Subst} \left(\int \operatorname{PolyLog} \left(3, \frac{ae^{ix}}{1 - \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right) \\
&\quad - 6 \operatorname{Subst} \left(\int \operatorname{PolyLog} \left(3, \frac{ae^{ix}}{1 + \sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad + \frac{3}{2} i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \frac{3}{2} \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - \frac{3}{4} i \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog}(3, -x)}{x} dx, x, e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + 6i \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(3, \frac{ax}{1 - \sqrt{1 - a^2}} \right)}{x} dx, x, e^{i \sec^{-1}(a+bx)} \right) \\
&\quad + 6i \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(3, \frac{ax}{1 + \sqrt{1 - a^2}} \right)}{x} dx, x, e^{i \sec^{-1}(a+bx)} \right)
\end{aligned}$$

$$\begin{aligned}
&= \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) + \sec^{-1}(a + bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \sec^{-1}(a + bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad - 3i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad + \frac{3}{2}i \sec^{-1}(a + bx)^2 \operatorname{PolyLog} \left(2, -e^{2i \sec^{-1}(a+bx)} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad + 6 \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) \\
&\quad - \frac{3}{2} \sec^{-1}(a + bx) \operatorname{PolyLog} \left(3, -e^{2i \sec^{-1}(a+bx)} \right) + 6i \operatorname{PolyLog} \left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{1 - \sqrt{1 - a^2}} \right) \\
&\quad + 6i \operatorname{PolyLog} \left(4, \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1 - a^2}} \right) - \frac{3}{4}i \operatorname{PolyLog} \left(4, -e^{2i \sec^{-1}(a+bx)} \right)
\end{aligned}$$

Mathematica [B] (warning: unable to verify)

Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 1058 vs. $2(430) = 860$.

Time = 2.92 (sec) , antiderivative size = 1058, normalized size of antiderivative = 2.46

$$\begin{aligned}
\int \frac{\sec^{-1}(a+bx)^3}{x} dx = & 2 \sec^{-1}(a+bx)^3 \log \left(1 + \frac{ae^{i \sec^{-1}(a+bx)}}{-1 + \sqrt{1-a^2}} \right) \\
& + \sec^{-1}(a+bx)^3 \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) - 6 \sec^{-1}(a \\
& + bx)^2 \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) \\
& + 2 \sec^{-1}(a+bx)^3 \log \left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1 + \sqrt{1-a^2}} \right) \\
& + \sec^{-1}(a+bx)^3 \log \left(1 - \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) + 6 \sec^{-1}(a \\
& + bx)^2 \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 - \frac{(1 + \sqrt{1-a^2}) e^{i \sec^{-1}(a+bx)}}{a} \right) \\
& - 3 \sec^{-1}(a+bx)^3 \log \left(1 + e^{2i \sec^{-1}(a+bx)} \right) \\
& + 2 \sec^{-1}(a+bx)^3 \log \left(\frac{2 \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a+bx} \right) \\
& - \sec^{-1}(a+bx)^3 \log \left(1 + \frac{a \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{-1 + \sqrt{1-a^2}} \right) \\
& - \sec^{-1}(a+bx)^3 \log \left(1 + \frac{(-1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& + 6 \sec^{-1}(a+bx)^2 \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 \right. \\
& \quad \left. + \frac{(-1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& - \sec^{-1}(a+bx)^3 \log \left(1 - \frac{a \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{1 + \sqrt{1-a^2}} \right) \\
& - \sec^{-1}(a+bx)^3 \log \left(1 - \frac{(1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right) \\
& - 6 \sec^{-1}(a+bx)^2 \arcsin \left(\frac{\sqrt{\frac{-1+a}{a}}}{\sqrt{2}} \right) \log \left(1 \right. \\
& \quad \left. - \frac{(1 + \sqrt{1-a^2}) \left(\frac{1}{a+bx} + i \sqrt{1 - \frac{1}{(a+bx)^2}} \right)}{a} \right)
\end{aligned}$$


```
[In] Integrate[ArcSec[a + b*x]^3/x,x]
[Out] 2*ArcSec[a + b*x]^3*Log[1 + (a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2])]
+ ArcSec[a + b*x]^3*Log[1 + ((-1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a]
- 6*ArcSec[a + b*x]^2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqr
t[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 2*ArcSec[a + b*x]^3*Log[1 - (a*E^(I
*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] + ArcSec[a + b*x]^3*Log[1 - ((1 + S
qrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a] + 6*ArcSec[a + b*x]^2*ArcSin[Sqrt[(
-1 + a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt[1 - a^2])*E^(I*ArcSec[a + b*x]))/a]
- 3*ArcSec[a + b*x]^3*Log[1 + E^((2*I)*ArcSec[a + b*x])] + 2*ArcSec[a + b*x
]^3*Log[(2*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))/(a + b*x)] - ArcS
ec[a + b*x]^3*Log[1 + (a*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/(-1
+ Sqrt[1 - a^2])] - ArcSec[a + b*x]^3*Log[1 + ((-1 + Sqrt[1 - a^2])*((a +
b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] + 6*ArcSec[a + b*x]^2*ArcSin[Sq
rt[(-1 + a)/a]/Sqrt[2]]*Log[1 + ((-1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*S
qrt[1 - (a + b*x)^(-2)]))]/a] - ArcSec[a + b*x]^3*Log[1 - (a*((a + b*x)^(-1)
+ I*Sqrt[1 - (a + b*x)^(-2)]))]/(1 + Sqrt[1 - a^2])] - ArcSec[a + b*x]^3*Lo
g[1 - ((1 + Sqrt[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a
] - 6*ArcSec[a + b*x]^2*ArcSin[Sqrt[(-1 + a)/a]/Sqrt[2]]*Log[1 - ((1 + Sqrt
[1 - a^2])*((a + b*x)^(-1) + I*Sqrt[1 - (a + b*x)^(-2)]))]/a] - (3*I)*ArcSec
[a + b*x]^2*PolyLog[2, -((a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2]))] -
(3*I)*ArcSec[a + b*x]^2*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 -
a^2])] + ((3*I)/2)*ArcSec[a + b*x]^2*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])]
+ 6*ArcSec[a + b*x]*PolyLog[3, -((a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 -
a^2]))] + 6*ArcSec[a + b*x]*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt
[1 - a^2])] - (3*ArcSec[a + b*x]*PolyLog[3, -E^((2*I)*ArcSec[a + b*x]))]/2
+ (6*I)*PolyLog[4, -((a*E^(I*ArcSec[a + b*x]))/(-1 + Sqrt[1 - a^2]))] + (6*
I)*PolyLog[4, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])] - ((3*I)/4)*Po
lyLog[4, -E^((2*I)*ArcSec[a + b*x])]
```

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx+a)^3}{x} dx$$

```
[In] int(arcsec(b*x+a)^3/x,x)
```

```
[Out] int(arcsec(b*x+a)^3/x,x)
```

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

[In] integrate(arcsec(b*x+a)^3/x,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)^3/x, x)

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{asec}^3(a + bx)}{x} dx$$

[In] integrate(asec(b*x+a)**3/x,x)

[Out] Integral(asec(a + b*x)**3/x, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

[In] integrate(arcsec(b*x+a)^3/x,x, algorithm="maxima")

[Out] integrate(arcsec(b*x + a)^3/x, x)

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x} dx$$

[In] integrate(arcsec(b*x+a)^3/x,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^3/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^3}{x} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)^3}{x} dx$$

```
[In] int(acos(1/(a + b*x))^3/x,x)
```

```
[Out] int(acos(1/(a + b*x))^3/x, x)
```

3.37 $\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx$

Optimal result	260
Rubi [A] (verified)	261
Mathematica [F(-1)]	265
Maple [F]	265
Fricas [F]	265
Sympy [F]	266
Maxima [F]	266
Giac [F]	266
Mupad [F(-1)]	267

Optimal result

Integrand size = 12, antiderivative size = 362

$$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx = -\frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6b \sec^{-1}(a+bx) \text{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \sec^{-1}(a+bx) \text{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} - \frac{6ib \text{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6ib \text{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}$$

```
[Out] -b*arcsec(b*x+a)^3/a-arcsec(b*x+a)^3/x-3*I*b*arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+3*I*b*arcsec(b*x+a)^2*ln(1-a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*b*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*b*arcsec(b*x+a)*polylog(2,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)-6*I*b*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1-(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)+6*I*b*polylog(3,a*(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))/(1+(-a^2+1)^(1/2)))/a/(-a^2+1)^(1/2)
```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 362, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5366, 4511, 4276, 3402, 2296, 2221, 2611, 2320, 6724}

$$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx = -\frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{6ib \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6ib \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{\sqrt{1-a^2}+1}\right)}{a\sqrt{1-a^2}} - \frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x}$$

[In] Int[ArcSec[a + b*x]^3/x^2,x]

[Out] -((b*ArcSec[a + b*x]^3)/a) - ArcSec[a + b*x]^3/x - ((3*I)*b*ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + ((3*I)*b*ArcSec[a + b*x]^2*Log[1 - (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - (6*b*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + (6*b*ArcSec[a + b*x]*PolyLog[2, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) - ((6*I)*b*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 - Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2]) + ((6*I)*b*PolyLog[3, (a*E^(I*ArcSec[a + b*x]))/(1 + Sqrt[1 - a^2])])/(a*Sqrt[1 - a^2])

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*(F_)^((g_)*(e_) + (f_)*(x_)))^(n_), x_Symbol] :> Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2296

Int[(F_)^(u)*(f_) + (g_)*(x_))^(m_)]/((a_) + (b_)*(F_)^(u) + (c_)*(F_)^(v_)), x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[2*(c/q), Int[

```
(f + g*x)^m*(F^u/(b - q + 2*c*F^u)), x], x] - Dist[2*(c/q), Int[(f + g*x)^m
*(F^u/(b + q + 2*c*F^u)), x], x]] /; FreeQ[{F, a, b, c, f, g}, x] && EqQ[v,
2*u] && LinearQ[u, x] && NeQ[b^2 - 4*a*c, 0] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] :=> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3402

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*sin[(e_.) + Pi*(k_.) + (f_.)*(
x_)]), x_Symbol] :=> Dist[2, Int[(c + d*x)^m*E^(I*Pi*(k - 1/2))*E^(I*(e + f
*x))/(b + 2*a*E^(I*Pi*(k - 1/2))*E^(I*(e + f*x)) - b*E^(2*I*k*Pi)*E^(2*I*(e
+ f*x)))]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[2*k] && NeQ[
a^2 - b^2, 0] && IGtQ[m, 0]
```

Rule 4276

```
Int[(csc[(e_.) + (f_.)*(x_)]*(b_.) + (a_.))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)
, x_Symbol] :=> Int[ExpandIntegrand[(c + d*x)^m, 1/(Sin[e + f*x]^n/(b + a*Si
n[e + f*x]^n)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && ILtQ[n, 0] && IGt
Q[m, 0]
```

Rule 4511

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sec[(c_.) + (d_.)*(x_)]*((a_.) + (b_.)*Sec[(c
_.) + (d_.)*(x_)]^(n_.)*Tan[(c_.) + (d_.)*(x_)]), x_Symbol] :=> Simp[(e + f*
x)^m*((a + b*Sec[c + d*x])^(n + 1)/(b*d*(n + 1))), x] - Dist[f*(m/(b*d*(n +
1))), Int[(e + f*x)^(m - 1)*(a + b*Sec[c + d*x])^(n + 1), x], x] /; FreeQ[
{a, b, c, d, e, f, n}, x] && IGtQ[m, 0] && NeQ[n, -1]
```

Rule 5366

```
Int[((a_.) + ArcSec[(c_.) + (d_.)*(x_)]*(b_.))^(p_.)*((e_.) + (f_.)*(x_))^(m
_.), x_Symbol] :=> Dist[1/d^(m + 1), Subst[Int[(a + b*x)^p*Sec[x]*Tan[x]*(d*
```

$e - c*f + f*\text{Sec}[x]^m, x], x, \text{ArcSec}[c + d*x]], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[p, 0] \&\& \text{IntegerQ}[m]$

Rule 6724

$\text{Int}[\text{PolyLog}[n, (c_*)*((a_*) + (b_*)*(x_))^{(p_)}]/((d_*) + (e_*)*(x_)), x_Symbol] \rightarrow \text{Simp}[\text{PolyLog}[n + 1, c*(a + b*x)^p]/(e*p), x] /; \text{FreeQ}\{a, b, c, d, e, n, p\}, x] \&\& \text{EqQ}[b*d, a*e]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= b \text{Subst} \left(\int \frac{x^3 \sec(x) \tan(x)}{(-a + \sec(x))^2} dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{\sec^{-1}(a + bx)^3}{x} + (3b) \text{Subst} \left(\int \frac{x^2}{-a + \sec(x)} dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{\sec^{-1}(a + bx)^3}{x} + (3b) \text{Subst} \left(\int \left(-\frac{x^2}{a} + \frac{x^2}{a(1 - a \cos(x))} \right) dx, x, \sec^{-1}(a + bx) \right) \\
 &= -\frac{b \sec^{-1}(a + bx)^3}{a} - \frac{\sec^{-1}(a + bx)^3}{x} + \frac{(3b) \text{Subst} \left(\int \frac{x^2}{1 - a \cos(x)} dx, x, \sec^{-1}(a + bx) \right)}{a} \\
 &= -\frac{b \sec^{-1}(a + bx)^3}{a} - \frac{\sec^{-1}(a + bx)^3}{x} + \frac{(6b) \text{Subst} \left(\int \frac{e^{ix} x^2}{-a + 2e^{ix} - ae^{2ix}} dx, x, \sec^{-1}(a + bx) \right)}{a} \\
 &= -\frac{b \sec^{-1}(a + bx)^3}{a} - \frac{\sec^{-1}(a + bx)^3}{x} \\
 &\quad - \frac{(6b) \text{Subst} \left(\int \frac{e^{ix} x^2}{2 - 2\sqrt{1 - a^2} - 2ae^{ix}} dx, x, \sec^{-1}(a + bx) \right)}{\sqrt{1 - a^2}} \\
 &\quad + \frac{(6b) \text{Subst} \left(\int \frac{e^{ix} x^2}{2 + 2\sqrt{1 - a^2} - 2ae^{ix}} dx, x, \sec^{-1}(a + bx) \right)}{\sqrt{1 - a^2}} \\
 &= -\frac{b \sec^{-1}(a + bx)^3}{a} - \frac{\sec^{-1}(a + bx)^3}{x} - \frac{3ib \sec^{-1}(a + bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a + bx)}}{1 - \sqrt{1 - a^2}} \right)}{a\sqrt{1 - a^2}} \\
 &\quad + \frac{3ib \sec^{-1}(a + bx)^2 \log \left(1 - \frac{ae^{i \sec^{-1}(a + bx)}}{1 + \sqrt{1 - a^2}} \right)}{a\sqrt{1 - a^2}} \\
 &\quad + \frac{(6ib) \text{Subst} \left(\int x \log \left(1 - \frac{2ae^{ix}}{2 - 2\sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right)}{a\sqrt{1 - a^2}} \\
 &\quad - \frac{(6ib) \text{Subst} \left(\int x \log \left(1 - \frac{2ae^{ix}}{2 + 2\sqrt{1 - a^2}} \right) dx, x, \sec^{-1}(a + bx) \right)}{a\sqrt{1 - a^2}}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&- \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{(6b) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{2ae^{ix}}{2-2\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&- \frac{(6b) \operatorname{Subst}\left(\int \operatorname{PolyLog}\left(2, \frac{2ae^{ix}}{2+2\sqrt{1-a^2}}\right) dx, x, \sec^{-1}(a+bx)\right)}{a\sqrt{1-a^2}} \\
&= -\frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&- \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&- \frac{(6ib) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1-\sqrt{1-a^2}}\right)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}} \\
&+ \frac{(6ib) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{ax}{1+\sqrt{1-a^2}}\right)}{x} dx, x, e^{i \sec^{-1}(a+bx)}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b \sec^{-1}(a+bx)^3}{a} - \frac{\sec^{-1}(a+bx)^3}{x} - \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{3ib \sec^{-1}(a+bx)^2 \log\left(1 - \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad + \frac{6b \sec^{-1}(a+bx) \operatorname{PolyLog}\left(2, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} \\
&\quad - \frac{6ib \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1-\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}} + \frac{6ib \operatorname{PolyLog}\left(3, \frac{ae^{i \sec^{-1}(a+bx)}}{1+\sqrt{1-a^2}}\right)}{a\sqrt{1-a^2}}
\end{aligned}$$

Mathematica [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx = \$Aborted$$

[In] Integrate[ArcSec[a + b*x]^3/x^2,x]

[Out] \$Aborted

Maple [F]

$$\int \frac{\operatorname{arcsec}(bx+a)^3}{x^2} dx$$

[In] int(arcsec(b*x+a)^3/x^2,x)

[Out] int(arcsec(b*x+a)^3/x^2,x)

Fricas [F]

$$\int \frac{\sec^{-1}(a+bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx+a)^3}{x^2} dx$$

[In] integrate(arcsec(b*x+a)^3/x^2,x, algorithm="fricas")

[Out] integral(arcsec(b*x + a)^3/x^2, x)

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{asec}^3(a + bx)}{x^2} dx$$

[In] integrate(asec(b*x+a)**3/x**2,x)

[Out] Integral(asec(a + b*x)**3/x**2, x)

Maxima [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

[In] integrate(arcsec(b*x+a)^3/x^2,x, algorithm="maxima")

[Out]
$$\begin{aligned} & -1/4*(4*\arctan(\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1})^3 - 3*\arctan(\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1} \\ & * \log(b^2*x^2 + 2*a*b*x + a^2)^2 - 4*x*\integrate(3/4*((4*b*x*\arctan(\sqrt{b*x + a + 1})*\sqrt{b*x + a - 1})^2 - b*x*\log(b^2*x^2 + 2*a*b*x + a^2)^2) \\ & * \sqrt{b*x + a + 1}*\sqrt{b*x + a - 1} - 4*((b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a)^2 + (b^3*x^3 + 2*a*b^2*x^2 + (a^2 - 1)*b*x - (b^3*x^3 + 3*a*b^2*x^2 + a^3 + (3*a^2 - 1)*b*x - a)*\log(b*x + a))*\log(b^2*x^2 + 2*a*b*x + a^2))*\arctan(\sqrt{b*x + a + 1}*\sqrt{b*x + a - 1}))/ \\ & (b^3*x^5 + 3*a*b^2*x^4 + (3*a^2 - 1)*b*x^3 + (a^3 - a)*x^2), x) /x \end{aligned}$$

Giac [F]

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\operatorname{arcsec}(bx + a)^3}{x^2} dx$$

[In] integrate(arcsec(b*x+a)^3/x^2,x, algorithm="giac")

[Out] integrate(arcsec(b*x + a)^3/x^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a + bx)^3}{x^2} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)^3}{x^2} dx$$

```
[In] int(acos(1/(a + b*x))^3/x^2,x)
```

```
[Out] int(acos(1/(a + b*x))^3/x^2, x)
```

3.38 $\int x(a + b \sec^{-1}(c + dx^2)) dx$

Optimal result	268
Rubi [A] (verified)	268
Mathematica [C] (verified)	270
Maple [A] (verified)	271
Fricas [A] (verification not implemented)	271
Sympy [F]	271
Maxima [A] (verification not implemented)	272
Giac [A] (verification not implemented)	272
Mupad [B] (verification not implemented)	272

Optimal result

Integrand size = 14, antiderivative size = 58

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^2)^2}}\right)}{2d}$$

[Out] $1/2*a*x^2 + 1/2*b*(d*x^2 + c)*\operatorname{arcsec}(d*x^2 + c)/d - 1/2*b*\operatorname{arctanh}((1 - 1/(d*x^2 + c)^2)^{(1/2)})/d$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6847, 5358, 379, 272, 65, 212}

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^2)^2}}\right)}{2d} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d}$$

[In] $\operatorname{Int}[x*(a + b*\operatorname{ArcSec}[c + d*x^2]), x]$

[Out] $(a*x^2)/2 + (b*(c + d*x^2)*\operatorname{ArcSec}[c + d*x^2])/(2*d) - (b*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - (c + d*x^2)^{-2}]])/(2*d)$

Rule 65

$\operatorname{Int}[(a_.) + (b_.)*(x_)^{(m_)}*((c_.) + (d_.)*(x_)^{(n_)}, x_Symbol] :> \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /;$ $\operatorname{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{LeQ}[-1, n, 0] \ \&\& \ \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \ \&\& \ \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 5358

Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{2} \text{Subst} \left(\int (a + b \sec^{-1}(c + dx)) dx, x, x^2 \right) \\
 &= \frac{ax^2}{2} + \frac{1}{2} b \text{Subst} \left(\int \sec^{-1}(c + dx) dx, x, x^2 \right) \\
 &= \frac{ax^2}{2} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d} - \frac{1}{2} b \text{Subst} \left(\int \frac{1}{(c + dx) \sqrt{1 - \frac{1}{(c + dx)^2}}} dx, x, x^2 \right) \\
 &= \frac{ax^2}{2} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d} - \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2} x}} dx, x, c + dx^2 \right)}{2d} \\
 &= \frac{ax^2}{2} + \frac{b(c + dx^2) \sec^{-1}(c + dx^2)}{2d} + \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - xx}} dx, x, \frac{1}{(c + dx^2)^2} \right)}{4d}
 \end{aligned}$$

$$= \frac{ax^2}{2} + \frac{b(c+dx^2)\sec^{-1}(c+dx^2)}{2d} - \frac{b\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1 - \frac{1}{(c+dx^2)^2}}\right)}{2d}$$

$$= \frac{ax^2}{2} + \frac{b(c+dx^2)\sec^{-1}(c+dx^2)}{2d} - \frac{b\text{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^2)^2}}\right)}{2d}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 2.31 (sec) , antiderivative size = 516, normalized size of antiderivative = 8.90

$$\int x(a + b\sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} + \frac{1}{2}bx^2\sec^{-1}(c + dx^2)$$

$$+ \frac{b(c + dx^2)\sqrt{\frac{-1+c^2+2cdx^2+d^2x^4}{(c+dx^2)^2}}\left(\sqrt[4]{-1}(-i + \sqrt{-1+c^2})\sqrt{2i-ic^2+2\sqrt{-1+c^2}}\arctan\left(\frac{(-1)^{3/4}\sqrt{2i-ic^2+2\sqrt{-1+c^2}}}{c\sqrt{-1+c^2}-c\sqrt{-1+c^2}}\right)\right)}{2d}$$

[In] Integrate[x*(a + b*ArcSec[c + d*x^2]),x]

[Out] (a*x^2)/2 + (b*x^2*ArcSec[c + d*x^2])/2 + (b*(c + d*x^2)*Sqrt[(-1 + c^2 + 2*c*d*x^2 + d^2*x^4)/(c + d*x^2)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + c^2])*Sqrt[2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^2)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4]]) + (-1)^(3/4)*(I + Sqrt[-1 + c^2])*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^2)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4]]) + c*(c*ArcTan[(Sqrt[-1 + c^2]*d^2*x^4)/(c^4 + c^3*d*x^2 + d^2*x^4 - c^2*(1 + Sqrt[-1 + c^2])*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4])]) - Log[Sqrt[-1 + c^2] - d*x^2 - Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4]] + Log[d^2*(Sqrt[-1 + c^2] + d*x^2 - Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4])])]/(2*c*d*Sqrt[-1 + c^2 + 2*c*d*x^2 + d^2*x^4])

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{ax^2}{2} + \frac{b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	64
derivativedivides	$\frac{(dx^2+c)a+b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	68
default	$\frac{(dx^2+c)a+b \left((dx^2+c) \operatorname{arcsec}(dx^2+c) - \ln \left(dx^2+c + (dx^2+c) \sqrt{1 - \frac{1}{(dx^2+c)^2}} \right) \right)}{2d}$	68

[In] `int(x*(a+b*arcsec(d*x^2+c)),x,method=_RETURNVERBOSE)`[Out] $\frac{1}{2}ax^2 + \frac{1}{2}b/d * ((dx^2+c) * \operatorname{arcsec}(dx^2+c) - \ln(dx^2+c + (dx^2+c) * (1 - 1/(dx^2+c)^2)^{(1/2)}))$ **Fricas [A] (verification not implemented)**

none

Time = 0.30 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{bdx^2 \operatorname{arcsec}(dx^2 + c) + adx^2 + 2bc \arctan(-dx^2 - c + \sqrt{d^2x^4 + 2cdx^2 + c^2 - 1}) + b \log(-dx^2 - c + \sqrt{d^2x^4 + 2cdx^2 + c^2 - 1})}{2d}$$

[In] `integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="fricas")`[Out] $\frac{1}{2} * (b * dx^2 * \operatorname{arcsec}(dx^2 + c) + a * dx^2 + 2 * b * c * \arctan(-dx^2 - c + \sqrt{d^2 * x^4 + 2 * c * dx^2 + c^2 - 1}) + b * \log(-dx^2 - c + \sqrt{d^2 * x^4 + 2 * c * dx^2 + c^2 - 1})) / d$ **Sympy [F]**

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \int x(a + b \operatorname{asec}(c + dx^2)) dx$$

[In] `integrate(x*(a+b*asec(d*x**2+c)),x)`[Out] `Integral(x*(a + b*asec(c + d*x**2)), x)`

Maxima [A] (verification not implemented)

none

Time = 0.18 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{1}{2} ax^2 + \frac{\left(2(dx^2 + c) \operatorname{arcsec}(dx^2 + c) - \log\left(\sqrt{-\frac{1}{(dx^2+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^2+c)^2} + 1}\right)\right)b}{4d}$$

[In] integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="maxima")

[Out] 1/2*a*x^2 + 1/4*(2*(d*x^2 + c)*arcsec(d*x^2 + c) - log(sqrt(-1/(d*x^2 + c)^2 + 1) + 1) + log(-sqrt(-1/(d*x^2 + c)^2 + 1)))*b/d

Giac [A] (verification not implemented)

none

Time = 0.39 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{1}{2} ax^2 + \frac{1}{4} bd \left(\frac{2(dx^2 + c) \arccos\left(-\frac{1}{(dx^2+c)\left(\frac{c}{dx^2+c}-1\right)-c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^2+c)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(dx^2+c)^2} + 1}\right)}{d^2} \right)$$

[In] integrate(x*(a+b*arcsec(d*x^2+c)),x, algorithm="giac")

[Out] 1/2*a*x^2 + 1/4*b*d*(2*(d*x^2 + c)*arccos(-1/((d*x^2 + c)*(c/(d*x^2 + c) - 1) - c))/d^2 - (log(sqrt(-1/(d*x^2 + c)^2 + 1) + 1) - log(-sqrt(-1/(d*x^2 + c)^2 + 1) + 1))/d^2)

Mupad [B] (verification not implemented)

Time = 1.21 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x(a + b \sec^{-1}(c + dx^2)) dx = \frac{ax^2}{2} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(dx^2+c)^2}}}\right)}{2d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^2+c}\right) (dx^2 + c)}{2d}$$

[In] int(x*(a + b*acos(1/(c + d*x^2))),x)

[Out] (a*x^2)/2 - (b*atanh(1/(1 - 1/(c + d*x^2)^2)^(1/2)))/(2*d) + (b*acos(1/(c + d*x^2))*(c + d*x^2))/(2*d)

3.39 $\int x^2(a + b \sec^{-1}(c + dx^3)) dx$

Optimal result	273
Rubi [A] (verified)	273
Mathematica [C] (verified)	275
Maple [A] (verified)	276
Fricas [A] (verification not implemented)	276
Sympy [F(-1)]	276
Maxima [A] (verification not implemented)	277
Giac [A] (verification not implemented)	277
Mupad [B] (verification not implemented)	277

Optimal result

Integrand size = 16, antiderivative size = 58

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^3)^2}}\right)}{3d}$$

[Out] $1/3*a*x^3 + 1/3*b*(d*x^3 + c)*\operatorname{arcsec}(d*x^3 + c)/d - 1/3*b*\operatorname{arctanh}((1 - 1/(d*x^3 + c)^2)^{(1/2)})/d$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6847, 5358, 379, 272, 65, 212}

$$\int x^2(a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^3)^2}}\right)}{3d} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d}$$

[In] $\operatorname{Int}[x^2*(a + b*\operatorname{ArcSec}[c + d*x^3]), x]$

[Out] $(a*x^3)/3 + (b*(c + d*x^3)*\operatorname{ArcSec}[c + d*x^3])/(3*d) - (b*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - (c + d*x^3)^{-2}]])/(3*d)$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b)^n, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_.), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 5358

Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3} \text{Subst} \left(\int (a + b \sec^{-1}(c + dx)) dx, x, x^3 \right) \\
 &= \frac{ax^3}{3} + \frac{1}{3} b \text{Subst} \left(\int \sec^{-1}(c + dx) dx, x, x^3 \right) \\
 &= \frac{ax^3}{3} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d} - \frac{1}{3} b \text{Subst} \left(\int \frac{1}{(c + dx) \sqrt{1 - \frac{1}{(c + dx)^2}}} dx, x, x^3 \right) \\
 &= \frac{ax^3}{3} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d} - \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2} x}} dx, x, c + dx^3 \right)}{3d} \\
 &= \frac{ax^3}{3} + \frac{b(c + dx^3) \sec^{-1}(c + dx^3)}{3d} + \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - xx}} dx, x, \frac{1}{(c + dx^3)^2} \right)}{6d}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{ax^3}{3} + \frac{b(c+dx^3)\sec^{-1}(c+dx^3)}{3d} - \frac{b\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1 - \frac{1}{(c+dx^3)^2}}\right)}{3d} \\
&= \frac{ax^3}{3} + \frac{b(c+dx^3)\sec^{-1}(c+dx^3)}{3d} - \frac{b\text{arctanh}\left(\sqrt{1 - \frac{1}{(c+dx^3)^2}}\right)}{3d}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 516, normalized size of antiderivative = 8.90

$$\int x^2(a + b\sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} + \frac{1}{3}bx^3\sec^{-1}(c + dx^3) + \frac{b(c + dx^3)\sqrt{\frac{-1+c^2+2cdx^3+d^2x^6}{(c+dx^3)^2}}\left(\sqrt[4]{-1}(-i + \sqrt{-1+c^2})\sqrt{2i-ic^2+2\sqrt{-1+c^2}}\arctan\left(\frac{(-1)^{3/4}\sqrt{2i-ic^2+2\sqrt{-1+c^2}}}{c\sqrt{-1+c^2}-c\sqrt{-1+c^2}}\right)\right)}{3}$$

[In] Integrate[x^2*(a + b*ArcSec[c + d*x^3]),x]

[Out] (a*x^3)/3 + (b*x^3*ArcSec[c + d*x^3])/3 + (b*(c + d*x^3)*Sqrt[(-1 + c^2 + 2*c*d*x^3 + d^2*x^6)/(c + d*x^3)^2]*((-1)^(1/4)*(-I + Sqrt[-1 + c^2])*Sqrt[2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(3/4)*Sqrt[2*I - I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^3)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]]) + (-1)^(3/4)*(I + Sqrt[-1 + c^2])*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*ArcTan[((-1)^(1/4)*Sqrt[-2*I + I*c^2 + 2*Sqrt[-1 + c^2]]*d*x^3)/(c*Sqrt[-1 + c^2] - c*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]]) + c*(c*ArcTan[(Sqrt[-1 + c^2]*d^2*x^6)/(c^4 + c^3*d*x^3 + d^2*x^6 - c^2*(1 + Sqrt[-1 + c^2])*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]])] - Log[Sqrt[-1 + c^2] - d*x^3 - Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6]] + Log[d^2*(Sqrt[-1 + c^2] + d*x^3 - Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6])])]/(3*c*d*Sqrt[-1 + c^2 + 2*c*d*x^3 + d^2*x^6])

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{x^3 a}{3} + \frac{b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	64
derivativedivides	$\frac{(dx^3+c)a + b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	68
default	$\frac{(dx^3+c)a + b \left((dx^3+c) \operatorname{arcsec}(dx^3+c) - \ln \left(dx^3+c + (dx^3+c) \sqrt{1 - \frac{1}{(dx^3+c)^2}} \right) \right)}{3d}$	68

```
[In] int(x^2*(a+b*arcsec(d*x^3+c)),x,method=_RETURNVERBOSE)
```

```
[Out] 1/3*x^3*a+1/3*b/d*((d*x^3+c)*arcsec(d*x^3+c)-ln(d*x^3+c+(d*x^3+c)*(1-1/(d*x^3+c)^2)^(1/2)))
```

Fricas [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx$$

$$= \frac{bdx^3 \operatorname{arcsec}(dx^3 + c) + adx^3 + 2bc \arctan(-dx^3 - c + \sqrt{d^2x^6 + 2cdx^3 + c^2 - 1}) + b \log(-dx^3 - c + \sqrt{d^2x^6 + 2cdx^3 + c^2 - 1})}{3d}$$

```
[In] integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="fricas")
```

```
[Out] 1/3*(b*d*x^3*arcsec(d*x^3 + c) + a*d*x^3 + 2*b*c*arctan(-d*x^3 - c + sqrt(d^2*x^6 + 2*c*d*x^3 + c^2 - 1)) + b*log(-d*x^3 - c + sqrt(d^2*x^6 + 2*c*d*x^3 + c^2 - 1)))/d
```

Sympy [F(-1)]

Timed out.

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \text{Timed out}$$

```
[In] integrate(x**2*(a+b*asec(d*x**3+c)),x)
```

```
[Out] Timed out
```

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \frac{1}{3} ax^3 + \frac{\left(2(dx^3 + c) \operatorname{arcsec}(dx^3 + c) - \log\left(\sqrt{-\frac{1}{(dx^3+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^3+c)^2} + 1}\right)\right) b}{6d}$$

[In] integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="maxima")

[Out] 1/3*a*x^3 + 1/6*(2*(d*x^3 + c)*arcsec(d*x^3 + c) - log(sqrt(-1/(d*x^3 + c)^2 + 1) + 1) + log(-sqrt(-1/(d*x^3 + c)^2 + 1)))*b/d

Giac [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \frac{1}{3} ax^3 + \frac{1}{6} bd \left(\frac{2(dx^3 + c) \arccos\left(\frac{1}{(dx^3+c)\left(\frac{c}{dx^3+c}-1\right)-c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^3+c)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(dx^3+c)^2} + 1}\right)}{d^2} \right)$$

[In] integrate(x^2*(a+b*arcsec(d*x^3+c)),x, algorithm="giac")

[Out] 1/3*a*x^3 + 1/6*b*d*(2*(d*x^3 + c)*arccos(-1/((d*x^3 + c)*(c/(d*x^3 + c) - 1) - c))/d^2 - (log(sqrt(-1/(d*x^3 + c)^2 + 1) + 1) - log(-sqrt(-1/(d*x^3 + c)^2 + 1)))/d^2)

Mupad [B] (verification not implemented)

Time = 0.93 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x^2 (a + b \sec^{-1}(c + dx^3)) dx = \frac{ax^3}{3} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(dx^3+c)^2}}}\right)}{3d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^3+c}\right) (dx^3 + c)}{3d}$$

[In] int(x^2*(a + b*acos(1/(c + d*x^3))),x)

[Out] (a*x^3)/3 - (b*atanh(1/(1 - 1/(c + d*x^3)^2)^(1/2)))/(3*d) + (b*acos(1/(c + d*x^3))*(c + d*x^3))/(3*d)

3.40 $\int x^3(a + b \sec^{-1}(c + dx^4)) dx$

Optimal result	278
Rubi [A] (verified)	278
Mathematica [B] (verified)	280
Maple [A] (verified)	280
Fricas [A] (verification not implemented)	281
Sympy [F(-1)]	281
Maxima [A] (verification not implemented)	281
Giac [A] (verification not implemented)	282
Mupad [B] (verification not implemented)	282

Optimal result

Integrand size = 16, antiderivative size = 58

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^4)^2}}\right)}{4d}$$

[Out] $1/4*a*x^4 + 1/4*b*(d*x^4 + c)*\operatorname{arcsec}(d*x^4 + c)/d - 1/4*b*\operatorname{arctanh}((1 - 1/(d*x^4 + c)^2)^{(1/2)})/d$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {6847, 5358, 379, 272, 65, 212}

$$\int x^3(a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} - \frac{\operatorname{barctanh}\left(\sqrt{1 - \frac{1}{(c + dx^4)^2}}\right)}{4d} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d}$$

[In] $\operatorname{Int}[x^3*(a + b*\operatorname{ArcSec}[c + d*x^4]), x]$

[Out] $(a*x^4)/4 + (b*(c + d*x^4)*\operatorname{ArcSec}[c + d*x^4])/(4*d) - (b*\operatorname{ArcTanh}[\operatorname{Sqrt}[1 - (c + d*x^4)^{-2}]])/ (4*d)$

Rule 65

$\operatorname{Int}[(a_.) + (b_.)*(x_)^m*((c_.) + (d_.)*(x_)^n), x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{p*(m+1)-1}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x]] /;$ $\operatorname{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{LeQ}[-1, n, 0] \ \&\& \ \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \ \&\& \ \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 5358

Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} \text{Subst} \left(\int (a + b \sec^{-1}(c + dx)) dx, x, x^4 \right) \\
 &= \frac{ax^4}{4} + \frac{1}{4} b \text{Subst} \left(\int \sec^{-1}(c + dx) dx, x, x^4 \right) \\
 &= \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} - \frac{1}{4} b \text{Subst} \left(\int \frac{1}{(c + dx) \sqrt{1 - \frac{1}{(c + dx)^2}}} dx, x, x^4 \right) \\
 &= \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} - \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - \frac{1}{x^2}} dx, x, c + dx^4 \right)}{4d} \\
 &= \frac{ax^4}{4} + \frac{b(c + dx^4) \sec^{-1}(c + dx^4)}{4d} + \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{1 - xx}} dx, x, \frac{1}{(c + dx^4)^2} \right)}{8d}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{ax^4}{4} + \frac{b(c+dx^4)\sec^{-1}(c+dx^4)}{4d} - \frac{b\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-\frac{1}{(c+dx^4)^2}}\right)}{4d} \\
&= \frac{ax^4}{4} + \frac{b(c+dx^4)\sec^{-1}(c+dx^4)}{4d} - \frac{b\text{arctanh}\left(\sqrt{1-\frac{1}{(c+dx^4)^2}}\right)}{4d}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 137 vs. 2(58) = 116.

Time = 0.24 (sec) , antiderivative size = 137, normalized size of antiderivative = 2.36

$$\begin{aligned}
&\int x^3(a+b\sec^{-1}(c+dx^4)) dx \\
&= \frac{ax^4}{4} + \frac{b(c+dx^4)\sec^{-1}(c+dx^4)}{4d} \\
&\quad - \frac{b\sqrt{-1+(c+dx^4)^2}\left(-\log\left(1-\frac{c+dx^4}{\sqrt{-1+(c+dx^4)^2}}\right) + \log\left(1+\frac{c+dx^4}{\sqrt{-1+(c+dx^4)^2}}\right)\right)}{8d(c+dx^4)\sqrt{1-\frac{1}{(c+dx^4)^2}}}
\end{aligned}$$

[In] Integrate[x^3*(a + b*ArcSec[c + d*x^4]),x]

[Out] (a*x^4)/4 + (b*(c + d*x^4)*ArcSec[c + d*x^4])/(4*d) - (b*Sqrt[-1 + (c + d*x^4)^2]*(-Log[1 - (c + d*x^4)/Sqrt[-1 + (c + d*x^4)^2]] + Log[1 + (c + d*x^4)/Sqrt[-1 + (c + d*x^4)^2]]))/(8*d*(c + d*x^4)*Sqrt[1 - (c + d*x^4)^(-2)])

Maple [A] (verified)

Time = 0.32 (sec) , antiderivative size = 64, normalized size of antiderivative = 1.10

method	result	size
parts	$\frac{x^4 a}{4} + \frac{b\left((dx^4+c)\operatorname{arcsec}(dx^4+c) - \ln\left(dx^4+c+(dx^4+c)\sqrt{1-\frac{1}{(dx^4+c)^2}}\right)\right)}{4d}$	64
derivativedivides	$\frac{(dx^4+c)a + b\left((dx^4+c)\operatorname{arcsec}(dx^4+c) - \ln\left(dx^4+c+(dx^4+c)\sqrt{1-\frac{1}{(dx^4+c)^2}}\right)\right)}{4d}$	68
default	$\frac{(dx^4+c)a + b\left((dx^4+c)\operatorname{arcsec}(dx^4+c) - \ln\left(dx^4+c+(dx^4+c)\sqrt{1-\frac{1}{(dx^4+c)^2}}\right)\right)}{4d}$	68

[In] int(x^3*(a+b*arcsec(d*x^4+c)),x,method=_RETURNVERBOSE)

[Out] 1/4*x^4*a+1/4*b/d*((d*x^4+c)*arcsec(d*x^4+c)-ln(d*x^4+c+(d*x^4+c)*(1-1/(d*x^4+c)^2)^(1/2)))

Fricas [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 96, normalized size of antiderivative = 1.66

$$\int x^3 (a + b \sec^{-1}(c + dx^4)) dx$$

$$= \frac{bdx^4 \operatorname{arcsec}(dx^4 + c) + adx^4 + 2bc \arctan(-dx^4 - c + \sqrt{d^2x^8 + 2cdx^4 + c^2 - 1}) + b \log(-dx^4 - c + \sqrt{d^2x^8 + 2cdx^4 + c^2 - 1})}{4d}$$

[In] integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="fricas")

```
[Out] 1/4*(b*d*x^4*arcsec(d*x^4 + c) + a*d*x^4 + 2*b*c*arctan(-d*x^4 - c + sqrt(d^2*x^8 + 2*c*d*x^4 + c^2 - 1)) + b*log(-d*x^4 - c + sqrt(d^2*x^8 + 2*c*d*x^4 + c^2 - 1)))/d
```

Sympy [F(-1)]

Timed out.

$$\int x^3 (a + b \sec^{-1}(c + dx^4)) dx = \text{Timed out}$$

[In] integrate(x**3*(a+b*asec(d*x**4+c)),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.22

$$\int x^3 (a + b \sec^{-1}(c + dx^4)) dx = \frac{1}{4} ax^4$$

$$+ \frac{\left(2(dx^4 + c) \operatorname{arcsec}(dx^4 + c) - \log\left(\sqrt{-\frac{1}{(dx^4+c)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(dx^4+c)^2} + 1} + 1\right)\right) b}{8d}$$

[In] integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="maxima")

```
[Out] 1/4*a*x^4 + 1/8*(2*(d*x^4 + c)*arcsec(d*x^4 + c) - log(sqrt(-1/(d*x^4 + c)^2 + 1) + 1) + log(-sqrt(-1/(d*x^4 + c)^2 + 1) + 1))*b/d
```

Giac [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.72

$$\int x^3 (a + b \sec^{-1}(c + dx^4)) dx = \frac{1}{4} ax^4 + \frac{1}{8} bd \left(\frac{2(dx^4 + c) \arccos\left(-\frac{1}{(dx^4 + c)\left(\frac{c}{dx^4 + c} - 1\right) - c}\right)}{d^2} - \frac{\log\left(\sqrt{-\frac{1}{(dx^4 + c)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(dx^4 + c)^2} + 1} + 1\right)}{d^2} \right)$$

[In] integrate(x^3*(a+b*arcsec(d*x^4+c)),x, algorithm="giac")

[Out] 1/4*a*x^4 + 1/8*b*d*(2*(d*x^4 + c)*arccos(-1/((d*x^4 + c)*(c/(d*x^4 + c) - 1) - c))/d^2 - (log(sqrt(-1/(d*x^4 + c)^2 + 1) + 1) - log(-sqrt(-1/(d*x^4 + c)^2 + 1) + 1))/d^2)

Mupad [B] (verification not implemented)

Time = 0.94 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.90

$$\int x^3 (a + b \sec^{-1}(c + dx^4)) dx = \frac{ax^4}{4} - \frac{b \operatorname{atanh}\left(\frac{1}{\sqrt{1 - \frac{1}{(dx^4 + c)^2}}}\right)}{4d} + \frac{b \operatorname{acos}\left(\frac{1}{dx^4 + c}\right) (dx^4 + c)}{4d}$$

[In] int(x^3*(a + b*acos(1/(c + d*x^4))),x)

[Out] (a*x^4)/4 - (b*atanh(1/(1 - 1/(c + d*x^4)^2)^(1/2)))/(4*d) + (b*acos(1/(c + d*x^4))*(c + d*x^4))/(4*d)

3.41 $\int x^{-1+n} \sec^{-1}(a + bx^n) dx$

Optimal result	283
Rubi [A] (verified)	283
Mathematica [B] (verified)	285
Maple [F]	285
Fricas [A] (verification not implemented)	285
Sympy [F(-1)]	286
Maxima [A] (verification not implemented)	286
Giac [A] (verification not implemented)	286
Mupad [B] (verification not implemented)	287

Optimal result

Integrand size = 14, antiderivative size = 49

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

[Out] (a+b*x^n)*arcsec(a+b*x^n)/b/n-arcTanh((1-1/(a+b*x^n)^2)^(1/2))/b/n

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {6847, 5358, 379, 272, 65, 212}

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

[In] Int[x^(-1 + n)*ArcSec[a + b*x^n],x]

[Out] ((a + b*x^n)*ArcSec[a + b*x^n])/(b*n) - ArcTanh[Sqrt[1 - (a + b*x^n)^(-2)]]/(b*n)

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 379

Int[(u_)^(m_.)*((a_) + (b_.)*(v_)^(n_))^(p_.), x_Symbol] := Dist[u^m/(Coefficient[v, x, 1]*v^m), Subst[Int[x^m*(a + b*x^n)^p, x], x, v], x] /; FreeQ[{a, b, m, n, p}, x] && LinearPairQ[u, v, x]

Rule 5358

Int[ArcSec[(c_) + (d_.)*(x_)], x_Symbol] := Simp[(c + d*x)*(ArcSec[c + d*x]/d), x] - Int[1/((c + d*x)*Sqrt[1 - 1/(c + d*x)^2]), x] /; FreeQ[{c, d}, x]

Rule 6847

Int[(u_)*(x_)^(m_.), x_Symbol] := Dist[1/(m + 1), Subst[Int[SubstFor[x^(m + 1), u, x], x], x, x^(m + 1)], x] /; FreeQ[m, x] && NeQ[m, -1] && FunctionOfQ[x^(m + 1), u, x]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \sec^{-1}(a + bx) dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{(a+bx)\sqrt{1-\frac{1}{(a+bx)^2}}} dx, x, x^n\right)}{n} \\
 &= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-\frac{1}{x^2}}} dx, x, a + bx^n\right)}{bn} \\
 &= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1-xx}} dx, x, \frac{1}{(a+bx^n)^2}\right)}{2bn} \\
 &= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{1-\frac{1}{(a+bx^n)^2}}\right)}{bn}
 \end{aligned}$$

$$= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\operatorname{arctanh}\left(\sqrt{1 - \frac{1}{(a+bx^n)^2}}\right)}{bn}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 130 vs. $2(49) = 98$.

Time = 0.23 (sec) , antiderivative size = 130, normalized size of antiderivative = 2.65

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{(a + bx^n) \sec^{-1}(a + bx^n)}{bn} - \frac{\sqrt{-1 + (a + bx^n)^2} \left(-\log\left(1 - \frac{a+bx^n}{\sqrt{-1+(a+bx^n)^2}}\right) + \log\left(1 + \frac{a+bx^n}{\sqrt{-1+(a+bx^n)^2}}\right) \right)}{2bn(a + bx^n) \sqrt{1 - \frac{1}{(a+bx^n)^2}}}$$

[In] Integrate[x^(-1 + n)*ArcSec[a + b*x^n], x]

[Out] ((a + b*x^n)*ArcSec[a + b*x^n])/(b*n) - (Sqrt[-1 + (a + b*x^n)^2]*(-Log[1 - (a + b*x^n)/Sqrt[-1 + (a + b*x^n)^2]] + Log[1 + (a + b*x^n)/Sqrt[-1 + (a + b*x^n)^2]]))/(2*b*n*(a + b*x^n)*Sqrt[1 - (a + b*x^n)^(-2)])

Maple [F]

$$\int x^{-1+n} \operatorname{arcsec}(a + bx^n) dx$$

[In] int(x^(-1+n)*arcsec(a+b*x^n), x)

[Out] int(x^(-1+n)*arcsec(a+b*x^n), x)

Fricas [A] (verification not implemented)

none

Time = 0.36 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.88

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{bx^n \operatorname{arcsec}(bx^n + a) + 2a \arctan(-bx^n - a + \sqrt{b^2x^{2n} + 2abx^n + a^2 - 1}) + \log(-bx^n - a + \sqrt{b^2x^{2n} + 2abx^n + a^2 - 1})}{bn}$$

[In] integrate(x^(-1+n)*arcsec(a+b*x^n), x, algorithm="fricas")

[Out] (b*x^n*arcsec(b*x^n + a) + 2*a*arctan(-b*x^n - a + sqrt(b^2*x^(2*n) + 2*a*b*x^n + a^2 - 1)) + log(-b*x^n - a + sqrt(b^2*x^(2*n) + 2*a*b*x^n + a^2 - 1)))/(b*n)

Sympy [F(-1)]

Timed out.

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = \text{Timed out}$$

[In] integrate(x**(-1+n)*asec(a+b*x**n),x)

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.35

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{2(bx^n + a) \operatorname{arcsec}(bx^n + a) - \log\left(\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right) + \log\left(-\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right)}{2bn}$$

[In] integrate(x^(-1+n)*arcsec(a+b*x^n),x, algorithm="maxima")

[Out] 1/2*(2*(b*x^n + a)*arcsec(b*x^n + a) - log(sqrt(-1/(b*x^n + a)^2 + 1) + 1) + log(-sqrt(-1/(b*x^n + a)^2 + 1) + 1))/(b*n)

Giac [A] (verification not implemented)

none

Time = 0.34 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.53

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx$$

$$= \frac{b \left(\frac{2(bx^n+a) \arccos\left(\frac{1}{bx^n+a}\right)}{b^2} - \frac{\log\left(\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right) - \log\left(-\sqrt{-\frac{1}{(bx^n+a)^2} + 1} + 1\right)}{b^2} \right)}{2n}$$

[In] integrate(x^(-1+n)*arcsec(a+b*x^n),x, algorithm="giac")

[Out] 1/2*b*(2*(b*x^n + a)*arccos(1/(b*x^n + a))/b^2 - (log(sqrt(-1/(b*x^n + a)^2 + 1) + 1) - log(-sqrt(-1/(b*x^n + a)^2 + 1) + 1))/b^2)/n

Mupad [B] (verification not implemented)

Time = 1.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 0.90

$$\int x^{-1+n} \sec^{-1}(a + bx^n) dx = -\frac{\operatorname{atanh}\left(\frac{1}{\sqrt{1-\frac{1}{(a+bx^n)^2}}}\right) - \operatorname{acos}\left(\frac{1}{a+bx^n}\right) (a + bx^n)}{bn}$$

[In] `int(x^(n - 1)*acos(1/(a + b*x^n)),x)`

[Out] `-(atanh(1/(1 - 1/(a + b*x^n)^2)^(1/2)) - acos(1/(a + b*x^n))*(a + b*x^n))/(b*n)`

3.42 $\int \sec^{-1}(ce^{a+bx}) dx$

Optimal result	288
Rubi [A] (verified)	288
Mathematica [B] (verified)	290
Maple [A] (verified)	291
Fricas [F(-2)]	291
Sympy [F]	292
Maxima [F]	292
Giac [F]	292
Mupad [F(-1)]	293

Optimal result

Integrand size = 10, antiderivative size = 85

$$\int \sec^{-1}(ce^{a+bx}) dx = \frac{i \sec^{-1}(ce^{a+bx})^2}{2b} - \frac{\sec^{-1}(ce^{a+bx}) \log(1 + e^{2i \sec^{-1}(ce^{a+bx})})}{b} + \frac{i \operatorname{PolyLog}(2, -e^{2i \sec^{-1}(ce^{a+bx})})}{2b}$$

[Out] 1/2*I*arcsec(c*exp(b*x+a))^2/b-arcsec(c*exp(b*x+a))*ln(1+(1/c/exp(b*x+a)+I*(1-1/c^2/exp(b*x+a)^2)^(1/2))^2)/b+1/2*I*polylog(2,-(1/c/exp(b*x+a)+I*(1-1/c^2/exp(b*x+a)^2)^(1/2))^2)/b

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {2320, 5326, 4722, 3800, 2221, 2317, 2438}

$$\int \sec^{-1}(ce^{a+bx}) dx = \frac{i \operatorname{PolyLog}(2, -e^{2i \sec^{-1}(ce^{a+bx})})}{2b} + \frac{i \sec^{-1}(ce^{a+bx})^2}{2b} - \frac{\sec^{-1}(ce^{a+bx}) \log(1 + e^{2i \sec^{-1}(ce^{a+bx})})}{b}$$

[In] Int[ArcSec[c*E^(a + b*x)],x]

[Out] ((I/2)*ArcSec[c*E^(a + b*x)]^2)/b - (ArcSec[c*E^(a + b*x)]*Log[1 + E^((2*I)*ArcSec[c*E^(a + b*x)])])/b + ((I/2)*PolyLog[2, -E^((2*I)*ArcSec[c*E^(a + b*x)])])/b

Rule 2221

```
Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*(c_) + (d_)*(x_)))^(n_)], x_Symbol]
:=> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)
))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :=> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*
(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :=> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3800

```
Int[(((c_) + (d_)*(x_))^(m_))*tan[(e_) + (f_)*(x_)], x_Symbol] :=> Simp[I
*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e
+ f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ
[m, 0]
```

Rule 4722

```
Int[((a_) + ArcCos[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] :=> -Subst[Int[
(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0
]
```

Rule 5326

```
Int[((a_) + ArcSec[(c_)*(x_)])*(b_)/(x_), x_Symbol] :=> -Subst[Int[(a + b
*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int \frac{\sec^{-1}(cx)}{x} dx, x, e^{a+bx}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{\arccos\left(\frac{x}{c}\right)}{x} dx, x, e^{-a-bx}\right)}{b} \\
&= \frac{\text{Subst}\left(\int x \tan(x) dx, x, \arccos\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \arccos\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{(2i)\text{Subst}\left(\int \frac{e^{2ix}x}{1+e^{2ix}} dx, x, \arccos\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \arccos\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \arccos\left(\frac{e^{-a-bx}}{c}\right)\right)}{b} \\
&= \frac{i \arccos\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad - \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b} \\
&= \frac{i \arccos\left(\frac{e^{-a-bx}}{c}\right)^2}{2b} - \frac{\arccos\left(\frac{e^{-a-bx}}{c}\right) \log\left(1 + e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right)}{b} \\
&\quad + \frac{i \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{e^{-a-bx}}{c}\right)}\right)}{2b}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 280 vs. 2(85) = 170.

Time = 0.92 (sec) , antiderivative size = 280, normalized size of antiderivative = 3.29

$$\int \sec^{-1}(ce^{a+bx}) dx = x \sec^{-1}(ce^{a+bx}) - \frac{e^{-a-bx} \left(4\sqrt{-1 + c^2 e^{2(a+bx)}} \arctan\left(\sqrt{-1 + c^2 e^{2(a+bx)}}\right) (2bx - \log(c^2 e^{2(a+bx)})) + \sqrt{1 - c^2 e^{2(a+bx)}} \left(\log^2(c\right)\right)}{2b}$$

```
[In] Integrate[ArcSec[c*E^(a + b*x)],x]
```

```
[Out] x*ArcSec[c*E^(a + b*x)] - (E^(-a - b*x)*(4*Sqrt[-1 + c^2*E^(2*(a + b*x))]*ArcTan[Sqrt[-1 + c^2*E^(2*(a + b*x))]]*(2*b*x - Log[c^2*E^(2*(a + b*x))]) + Sqrt[1 - c^2*E^(2*(a + b*x))]*(Log[c^2*E^(2*(a + b*x))]^2 - 4*Log[c^2*E^(2*(a + b*x))]*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2] + 2*Log[(1 + Sqrt[1 - c^2*E^(2*(a + b*x))])/2]^2) - 4*Sqrt[1 - c^2*E^(2*(a + b*x))]*PolyLog[2, (1 - Sqrt[1 - c^2*E^(2*(a + b*x))])/2]))/(8*b*c*Sqrt[1 - 1/(c^2*E^(2*(a + b*x)))]))
```

Maple [A] (verified)

Time = 1.23 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.31

method	result
derivativedivides	$\frac{i \operatorname{arcsec}\left(\frac{e^{bx+a}c}{2}\right)^2 - \operatorname{arcsec}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{b} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{2}$
default	$\frac{i \operatorname{arcsec}\left(\frac{e^{bx+a}c}{2}\right)^2 - \operatorname{arcsec}(e^{bx+a}c) \ln\left(1 + \left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{b} + \frac{i \operatorname{polylog}\left(2, -\left(\frac{e^{-bx-a}}{c} + i\sqrt{1 - \frac{e^{-2bx-2a}}{c^2}}\right)^2\right)}{2}$

```
[In] int(arcsec(exp(b*x+a)*c),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(1/2*I*arcsec(exp(b*x+a)*c)^2-arcsec(exp(b*x+a)*c)*ln(1+(1/c/exp(b*x+a)+I*(1-1/c^2/exp(b*x+a)^2)^(1/2))^2)+1/2*I*polylog(2,-(1/c/exp(b*x+a)+I*(1-1/c^2/exp(b*x+a)^2)^(1/2))^2))
```

Fricas [F(-2)]

Exception generated.

$$\int \sec^{-1}(ce^{a+bx}) dx = \text{Exception raised: TypeError}$$

```
[In] integrate(arcsec(c*exp(b*x+a)),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)
```

Sympy [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{asec}(ce^{a+bx}) dx$$

[In] integrate(asec(c*exp(b*x+a)),x)

[Out] Integral(asec(c*exp(a + b*x)), x)

Maxima [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsec}(ce^{(bx+a)}) dx$$

[In] integrate(arcsec(c*exp(b*x+a)),x, algorithm="maxima")

[Out]
$$\begin{aligned} & -1/2*(2*b^2*c^2*\integrate(x*e^{(2*b*x + 2*a + 1/2*\log(c*e^{(b*x + a) + 1}) + 1} \\ & /2*\log(c*e^{(b*x + a) - 1}))/c^2*e^{(2*b*x + 2*a) + (c^2*e^{(2*b*x + 2*a) - 1} \\ & *e^{(\log(c*e^{(b*x + a) + 1}) + \log(c*e^{(b*x + a) - 1})) - 1}, x) + 2*I*b^2*c^2 \\ & *\integrate(x*e^{(2*b*x + 2*a)}/(c^2*e^{(2*b*x + 2*a) + (c^2*e^{(2*b*x + 2*a) - 1} \\ &)*e^{(\log(c*e^{(b*x + a) + 1}) + \log(c*e^{(b*x + a) - 1})) - 1}, x) - I*b^2*x^2 \\ & - 2*b*x*\arctan(\sqrt{c*e^{(b*x + a) + 1}*\sqrt{c*e^{(b*x + a) - 1}}) + I*b*x*\log \\ & (c^2*e^{(2*b*x + 2*a)}) - I*b*x*\log(c*e^{(b*x + a) + 1}) - I*b*x*\log(-c*e^{(b*x \\ & + a) + 1}) - 2*(I*a*b + I*b*\log(c))*x - I*\operatorname{dilog}(c*e^{(b*x + a)}) - I*\operatorname{dilog}(-c \\ & *e^{(b*x + a)}))/b \end{aligned}$$

Giac [F]

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{arcsec}(ce^{(bx+a)}) dx$$

[In] integrate(arcsec(c*exp(b*x+a)),x, algorithm="giac")

[Out] integrate(arcsec(c*e^{(b*x + a)}), x)

Mupad [F(-1)]

Timed out.

$$\int \sec^{-1}(ce^{a+bx}) dx = \int \operatorname{acos}\left(\frac{e^{-a-bx}}{c}\right) dx$$

```
[In] int(acos(exp(- a - b*x)/c),x)
```

```
[Out] int(acos(exp(- a - b*x)/c), x)
```

3.43 $\int e^{\sec^{-1}(ax)} x^2 dx$

Optimal result	294
Rubi [A] (verified)	294
Mathematica [A] (verified)	296
Maple [F]	296
Fricas [F]	296
Sympy [F]	297
Maxima [F]	297
Giac [F]	297
Mupad [F(-1)]	297

Optimal result

Integrand size = 10, antiderivative size = 99

$$\int e^{\sec^{-1}(ax)} x^2 dx$$

$$= -\frac{\left(\frac{12}{5} + \frac{4i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 3, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

$$+ \frac{\left(\frac{24}{5} + \frac{8i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

[Out] $(-12/5-4/5*I)*\exp((1+3*I)*\operatorname{arcsec}(a*x))*\operatorname{hypergeom}([3, 3/2-1/2*I], [5/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^{(1/2)})^2)/a^3+(24/5+8/5*I)*\exp((1+3*I)*\operatorname{arcsec}(a*x))*\operatorname{hypergeom}([4, 3/2-1/2*I], [5/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^{(1/2)})^2)/a^3$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5374, 12, 4559, 2283}

$$\int e^{\sec^{-1}(ax)} x^2 dx$$

$$= \frac{\left(\frac{24}{5} + \frac{8i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

$$- \frac{\left(\frac{12}{5} + \frac{4i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 3, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

[In] Int[E^ArcSec[a*x]*x^2,x]

[Out] ((-12/5 - (4*I)/5)*E^((1 + 3*I)*ArcSec[a*x])*Hypergeometric2F1[3/2 - I/2, 3, 5/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a^3 + ((24/5 + (8*I)/5)*E^((1 + 3*I)*ArcSec[a*x])*Hypergeometric2F1[3/2 - I/2, 4, 5/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a^3

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2283

Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x))/(g*h*Log[G]))*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 4559

Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^(c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]

Rule 5374

Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Dist[1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{e^x \sec^3(x) \tan(x)}{a^2} dx, x, \sec^{-1}(ax)\right)}{a} \\
 &= \frac{\text{Subst}\left(\int e^x \sec^3(x) \tan(x) dx, x, \sec^{-1}(ax)\right)}{a^3} \\
 &= \frac{\text{Subst}\left(\int \left(\frac{16ie^{(1+3i)x}}{(1+e^{2ix})^4} - \frac{8ie^{(1+3i)x}}{(1+e^{2ix})^3}\right) dx, x, \sec^{-1}(ax)\right)}{a^3} \\
 &= -\frac{(8i)\text{Subst}\left(\int \frac{e^{(1+3i)x}}{(1+e^{2ix})^3} dx, x, \sec^{-1}(ax)\right)}{a^3} + \frac{(16i)\text{Subst}\left(\int \frac{e^{(1+3i)x}}{(1+e^{2ix})^4} dx, x, \sec^{-1}(ax)\right)}{a^3}
 \end{aligned}$$

$$= -\frac{\left(\frac{12}{5} + \frac{4i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 3, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3} \\ + \frac{\left(\frac{24}{5} + \frac{8i}{5}\right) e^{(1+3i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{3}{2} - \frac{i}{2}, 4, \frac{5}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^3}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 95, normalized size of antiderivative = 0.96

$$\int e^{\sec^{-1}(ax)} x^2 dx \\ = \frac{e^{\sec^{-1}(ax)} \left((-4 - 4i) \left(-i + a\sqrt{1 - \frac{1}{a^2 x^2}} \right) \operatorname{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right) + a^4 x^4 (5 + \cos[2\operatorname{ArcSec}[a*x]] - \sin[2\operatorname{ArcSec}[a*x]]) \right)}{12a^4 x}$$

[In] Integrate[E^ArcSec[a*x]*x^2,x]

[Out] (E^ArcSec[a*x]*((-4 - 4*I)*(-I + a*Sqrt[1 - 1/(a^2*x^2)]*x)*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])] + a^4*x^4*(5 + Cos[2*ArcSec[a*x]] - Sin[2*ArcSec[a*x]])))/(12*a^4*x)

Maple [F]

$$\int e^{\operatorname{arcsec}(ax)} x^2 dx$$

[In] int(exp(arcsec(a*x))*x^2,x)

[Out] int(exp(arcsec(a*x))*x^2,x)

Fricas [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{(\operatorname{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x))*x^2,x, algorithm="fricas")

[Out] integral(x^2*e^(arcsec(a*x)), x)

Sympy [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\operatorname{asec}(ax)} dx$$

[In] integrate(exp(asec(a*x))*x**2,x)

[Out] Integral(x**2*exp(asec(a*x)), x)

Maxima [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\operatorname{arcsec}(ax)} dx$$

[In] integrate(exp(arcsec(a*x))*x^2,x, algorithm="maxima")

[Out] integrate(x^2*e^(arcsec(a*x)), x)

Giac [F]

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\operatorname{arcsec}(ax)} dx$$

[In] integrate(exp(arcsec(a*x))*x^2,x, algorithm="giac")

[Out] integrate(x^2*e^(arcsec(a*x)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} x^2 dx = \int x^2 e^{\operatorname{acos}(\frac{1}{ax})} dx$$

[In] int(x^2*exp(acos(1/(a*x))),x)

[Out] int(x^2*exp(acos(1/(a*x))), x)

3.44 $\int e^{\sec^{-1}(ax)} x dx$

Optimal result	298
Rubi [A] (verified)	298
Mathematica [A] (verified)	300
Maple [F]	300
Fricas [F]	300
Sympy [F]	300
Maxima [F]	301
Giac [F]	301
Mupad [F(-1)]	301

Optimal result

Integrand size = 8, antiderivative size = 91

$$\int e^{\sec^{-1}(ax)} x dx$$

$$= -\frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 2, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}$$

$$+ \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}$$

[Out] $(-8/5-4/5*I)*\exp((1+2*I)*\operatorname{arcsec}(a*x))*\operatorname{hypergeom}([2, 1-1/2*I], [2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^{(1/2)})^2/a^2+(16/5+8/5*I)*\exp((1+2*I)*\operatorname{arcsec}(a*x))*\operatorname{hypergeom}([3, 1-1/2*I], [2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^{(1/2)})^2/a^2)$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5374, 12, 4559, 2283}

$$\int e^{\sec^{-1}(ax)} x dx$$

$$= \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}$$

$$- \frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(1 - \frac{i}{2}, 2, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}$$

[In] Int[E^ArcSec[a*x]*x,x]

[Out] $((-8/5 - (4*I)/5)*E^{((1 + 2*I)*ArcSec[a*x])}*Hypergeometric2F1[1 - I/2, 2, 2 - I/2, -E^{((2*I)*ArcSec[a*x])}])/a^2 + ((16/5 + (8*I)/5)*E^{((1 + 2*I)*ArcSec[a*x])}*Hypergeometric2F1[1 - I/2, 3, 2 - I/2, -E^{((2*I)*ArcSec[a*x])}])/a^2$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2283

Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 4559

Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^(c*(a + b*x)), G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]

Rule 5374

Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Dist[1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{e^x \sec^2(x) \tan(x)}{a} dx, x, \sec^{-1}(ax)\right)}{a} \\
 &= \frac{\text{Subst}\left(\int e^x \sec^2(x) \tan(x) dx, x, \sec^{-1}(ax)\right)}{a^2} \\
 &= \frac{\text{Subst}\left(\int \left(\frac{8ie^{(1+2i)x}}{(1+e^{2ix})^3} - \frac{4ie^{(1+2i)x}}{(1+e^{2ix})^2}\right) dx, x, \sec^{-1}(ax)\right)}{a^2} \\
 &= -\frac{(4i)\text{Subst}\left(\int \frac{e^{(1+2i)x}}{(1+e^{2ix})^2} dx, x, \sec^{-1}(ax)\right)}{a^2} + \frac{(8i)\text{Subst}\left(\int \frac{e^{(1+2i)x}}{(1+e^{2ix})^3} dx, x, \sec^{-1}(ax)\right)}{a^2} \\
 &= -\frac{\left(\frac{8}{5} + \frac{4i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(1 - \frac{i}{2}, 2, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2} \\
 &\quad + \frac{\left(\frac{16}{5} + \frac{8i}{5}\right) e^{(1+2i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(1 - \frac{i}{2}, 3, 2 - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.18

$$\int e^{\sec^{-1}(ax)} x dx = \frac{\left(\frac{1}{5} + \frac{i}{10}\right) e^{\sec^{-1}(ax)} \left((-2 + i)ax \left(\sqrt{1 - \frac{1}{a^2 x^2}} - ax\right) + (1 + 2i) \text{Hypergeometric2F1}\left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)}\right)\right)}{a^2}$$

[In] Integrate[E^ArcSec[a*x]*x,x]

[Out] ((1/5 + I/10)*E^ArcSec[a*x]*((-2 + I)*a*x*(Sqrt[1 - 1/(a^2*x^2)] - a*x) + (1 + 2*I)*Hypergeometric2F1[-1/2*I, 1, 1 - I/2, -E^((2*I)*ArcSec[a*x])]) - E^((2*I)*ArcSec[a*x])*Hypergeometric2F1[1, 1 - I/2, 2 - I/2, -E^((2*I)*ArcSec[a*x])]))/a^2

Maple [F]

$$\int e^{\text{arcsec}(ax)} x dx$$

[In] int(exp(arcsec(a*x))*x,x)

[Out] int(exp(arcsec(a*x))*x,x)

Fricas [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\text{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x))*x,x, algorithm="fricas")

[Out] integral(x*e^(arcsec(a*x)), x)

Sympy [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{\text{asec}(ax)} dx$$

[In] integrate(exp(asec(a*x))*x,x)

[Out] Integral(x*exp(asec(a*x)), x)

Maxima [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\operatorname{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x))*x,x, algorithm="maxima")

[Out] integrate(x*e^(arcsec(a*x)), x)

Giac [F]

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{(\operatorname{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x))*x,x, algorithm="giac")

[Out] integrate(x*e^(arcsec(a*x)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} x dx = \int x e^{\operatorname{acos}(\frac{1}{ax})} dx$$

[In] int(x*exp(acos(1/(a*x))),x)

[Out] int(x*exp(acos(1/(a*x))), x)

3.45 $\int e^{\sec^{-1}(ax)} dx$

Optimal result	302
Rubi [A] (verified)	302
Mathematica [A] (verified)	304
Maple [F]	304
Fricas [F]	304
Sympy [F]	304
Maxima [F]	305
Giac [F]	305
Mupad [F(-1)]	305

Optimal result

Integrand size = 6, antiderivative size = 91

$$\int e^{\sec^{-1}(ax)} dx = -\frac{(1+i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 1, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a} + \frac{(2+2i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 2, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

[Out] (-1-I)*exp((1+I)*arcsec(a*x))*hypergeom([1, 1/2-1/2*I], [3/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a+(2+2*I)*exp((1+I)*arcsec(a*x))*hypergeom([2, 1/2-1/2*I], [3/2-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^(1/2))^2)/a

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5374, 4559, 2283}

$$\int e^{\sec^{-1}(ax)} dx = \frac{(2+2i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 2, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a} - \frac{(1+i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2}-\frac{i}{2}, 1, \frac{3}{2}-\frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

[In] Int[E^ArcSec[a*x], x]

[Out] ((-1 - I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a + ((2 + 2*I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 2, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a

Rule 2283

```
Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_)))^(p_)*(G_)^((h_)*((f_
) + (g_)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hype
rgeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1,
Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g,
h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])
```

Rule 4559

```
Int[(F_)^((c_)*((a_) + (b_)*(x_)))*(G_)[(d_) + (e_)*(x_)]^(m_)*(H_)[(
d_) + (e_)*(x_)]^(n_), x_Symbol] := Int[ExpandTrigToExp[F^(c*(a + b*x)),
G[d + e*x]^m*H[d + e*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IGtQ[
m, 0] && IGtQ[n, 0] && TrigQ[G] && TrigQ[H]
```

Rule 5374

```
Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Dist[
1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x,
ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{\text{Subst}\left(\int e^x \sec(x) \tan(x) dx, x, \sec^{-1}(ax)\right)}{a} \\
&= \frac{\text{Subst}\left(\int \left(\frac{4ie^{(1+i)x}}{(1+e^{2ix})^2} - \frac{2ie^{(1+i)x}}{1+e^{2ix}}\right) dx, x, \sec^{-1}(ax)\right)}{a} \\
&= -\frac{(2i)\text{Subst}\left(\int \frac{e^{(1+i)x}}{1+e^{2ix}} dx, x, \sec^{-1}(ax)\right)}{a} + \frac{(4i)\text{Subst}\left(\int \frac{e^{(1+i)x}}{(1+e^{2ix})^2} dx, x, \sec^{-1}(ax)\right)}{a} \\
&= -\frac{(1+i)e^{(1+i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a} \\
&\quad + \frac{(2+2i)e^{(1+i)\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 2, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.59

$$\int e^{\sec^{-1}(ax)} dx = e^{\sec^{-1}(ax)} x - \frac{(1-i)e^{(1+i)\sec^{-1}(ax)} \operatorname{Hypergeometric2F1}\left(\frac{1}{2} - \frac{i}{2}, 1, \frac{3}{2} - \frac{i}{2}, -e^{2i\sec^{-1}(ax)}\right)}{a}$$

[In] Integrate[E^ArcSec[a*x],x]

[Out] E^ArcSec[a*x]*x - ((1 - I)*E^((1 + I)*ArcSec[a*x])*Hypergeometric2F1[1/2 - I/2, 1, 3/2 - I/2, -E^((2*I)*ArcSec[a*x])])/a

Maple [F]

$$\int e^{\operatorname{arcsec}(ax)} dx$$

[In] int(exp(arcsec(a*x)),x)

[Out] int(exp(arcsec(a*x)),x)

Fricas [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\operatorname{arcsec}(ax)} dx$$

[In] integrate(exp(arcsec(a*x)),x, algorithm="fricas")

[Out] integral(e^(arcsec(a*x)), x)

Sympy [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\operatorname{asec}(ax)} dx$$

[In] integrate(exp(asec(a*x)),x)

[Out] Integral(exp(asec(a*x)), x)

Maxima [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{(\operatorname{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x)),x, algorithm="maxima")

[Out] integrate(e^(arcsec(a*x)), x)

Giac [F]

$$\int e^{\sec^{-1}(ax)} dx = \int e^{(\operatorname{arcsec}(ax))} dx$$

[In] integrate(exp(arcsec(a*x)),x, algorithm="giac")

[Out] integrate(e^(arcsec(a*x)), x)

Mupad [F(-1)]

Timed out.

$$\int e^{\sec^{-1}(ax)} dx = \int e^{\operatorname{acos}(\frac{1}{ax})} dx$$

[In] int(exp(acos(1/(a*x))),x)

[Out] int(exp(acos(1/(a*x))), x)

3.46 $\int \frac{e^{\sec^{-1}(ax)}}{x} dx$

Optimal result	306
Rubi [A] (verified)	306
Mathematica [A] (verified)	308
Maple [F]	308
Fricas [F]	308
Sympy [F]	308
Maxima [F]	309
Giac [F]	309
Mupad [F(-1)]	309

Optimal result

Integrand size = 10, antiderivative size = 45

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = -ie^{\sec^{-1}(ax)} + 2ie^{\sec^{-1}(ax)} \text{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right)$$

[Out] $-I*\exp(\text{arcsec}(a*x))+2*I*\exp(\text{arcsec}(a*x))*\text{hypergeom}([1, -1/2*I], [1-1/2*I], -(1/a/x+I*(1-1/a^2/x^2)^{(1/2)})^2)$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5374, 12, 4527, 2225, 2283}

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = 2ie^{\sec^{-1}(ax)} \text{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) - ie^{\sec^{-1}(ax)}$$

[In] $\text{Int}[E^{\text{ArcSec}[a*x]}/x, x]$

[Out] $(-I)*E^{\text{ArcSec}[a*x]} + (2*I)*E^{\text{ArcSec}[a*x]}*\text{Hypergeometric2F1}[-1/2*I, 1, 1 - I/2, -E^{((2*I)*\text{ArcSec}[a*x])}]$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_) /; \text{FreeQ}[b, x]]$

Rule 2225

Int[((F_)^((c_)*(a_) + (b_)*(x_)))^(n_), x_Symbol] := Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 2283

Int[((a_) + (b_)*(F_)^((e_)*((c_) + (d_)*(x_))))^(p_)*(G_)^((h_)*((f_) + (g_)*(x_))), x_Symbol] := Simp[a^p*(G^(h*(f + g*x)))/(g*h*Log[G])*Hypergeometric2F1[-p, g*h*(Log[G]/(d*e*Log[F])), g*h*(Log[G]/(d*e*Log[F])) + 1, Simplify[(-b/a)*F^(e*(c + d*x))], x] /; FreeQ[{F, G, a, b, c, d, e, f, g, h, p}, x] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 4527

Int[(F_)^((c_)*((a_) + (b_)*(x_)))*Tan[(d_) + (e_)*(x_)]^(n_), x_Symbol] := Dist[I^n, Int[ExpandIntegrand[F^(c*(a + b*x))*((1 - E^(2*I*(d + e*x)))^n/(1 + E^(2*I*(d + e*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e}, x] && IntegerQ[n]

Rule 5374

Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Dist[1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int a e^x \tan(x) dx, x, \sec^{-1}(ax)\right)}{a} \\
 &= \text{Subst}\left(\int e^x \tan(x) dx, x, \sec^{-1}(ax)\right) \\
 &= i \text{Subst}\left(\int \left(-e^x + \frac{2e^x}{1 + e^{2ix}}\right) dx, x, \sec^{-1}(ax)\right) \\
 &= -\left(i \text{Subst}\left(\int e^x dx, x, \sec^{-1}(ax)\right)\right) + 2i \text{Subst}\left(\int \frac{e^x}{1 + e^{2ix}} dx, x, \sec^{-1}(ax)\right) \\
 &= -ie^{\sec^{-1}(ax)} + 2ie^{\sec^{-1}(ax)} \text{Hypergeometric2F1}\left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)}\right)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 79, normalized size of antiderivative = 1.76

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = -i \left(-e^{\sec^{-1}(ax)} \operatorname{Hypergeometric2F1} \left(-\frac{i}{2}, 1, 1 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) + \left(\frac{1}{5} - \frac{2i}{5} \right) e^{(1+2i) \sec^{-1}(ax)} \operatorname{Hypergeometric2F1} \left(1, 1 - \frac{i}{2}, 2 - \frac{i}{2}, -e^{2i \sec^{-1}(ax)} \right) \right)$$

[In] Integrate[E^ArcSec[a*x]/x,x]

[Out] (-I)*(-(E^ArcSec[a*x]*Hypergeometric2F1[-1/2*I, 1, 1 - I/2, -E^((2*I)*ArcSec[a*x])])) + (1/5 - (2*I)/5)*E^((1 + 2*I)*ArcSec[a*x])*Hypergeometric2F1[1, 1 - I/2, 2 - I/2, -E^((2*I)*ArcSec[a*x])]]

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x} dx$$

[In] int(exp(arcsec(a*x))/x,x)

[Out] int(exp(arcsec(a*x))/x,x)

Fricas [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x} dx$$

[In] integrate(exp(arcsec(a*x))/x,x, algorithm="fricas")

[Out] integral(e^(arcsec(a*x))/x, x)

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x} dx$$

[In] integrate(exp(asec(a*x))/x,x)

[Out] Integral(exp(asec(a*x))/x, x)

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x} dx$$

[In] integrate(exp(arcsec(a*x))/x,x, algorithm="maxima")

[Out] integrate(e^(arcsec(a*x))/x, x)

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x} dx$$

[In] integrate(exp(arcsec(a*x))/x,x, algorithm="giac")

[Out] integrate(e^(arcsec(a*x))/x, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x} dx = \int \frac{e^{\operatorname{acos}(\frac{1}{ax})}}{x} dx$$

[In] int(exp(acos(1/(a*x)))/x,x)

[Out] int(exp(acos(1/(a*x)))/x, x)

3.47 $\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx$

Optimal result	310
Rubi [A] (verified)	310
Mathematica [A] (verified)	311
Maple [F]	311
Fricas [A] (verification not implemented)	312
Sympy [F]	312
Maxima [F]	312
Giac [A] (verification not implemented)	312
Mupad [F(-1)]	313

Optimal result

Integrand size = 10, antiderivative size = 39

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} a e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{e^{\sec^{-1}(ax)}}{2x}$$

[Out] $-1/2*\exp(\operatorname{arcsec}(a*x))/x+1/2*a*\exp(\operatorname{arcsec}(a*x))*(1-1/a^2/x^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5374, 12, 4517}

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} a \sqrt{1 - \frac{1}{a^2 x^2}} e^{\sec^{-1}(ax)} - \frac{e^{\sec^{-1}(ax)}}{2x}$$

[In] $\text{Int}[E^{\text{ArcSec}[a*x]}/x^2, x]$

[Out] $(a * E^{\text{ArcSec}[a*x]} * \text{Sqrt}[1 - 1/(a^2 * x^2)]) / 2 - E^{\text{ArcSec}[a*x]} / (2 * x)$

Rule 12

$\text{Int}[(a_*) * (u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*) * (v_)] /; \text{FreeQ}[b, x]$

Rule 4517

$\text{Int}[(F_)^c * ((c_*) * ((a_*) + (b_*) * (x_))) * \text{Sin}[(d_*) + (e_*) * (x_)], x_Symbol] \rightarrow \text{Simp}[b * c * \text{Log}[F] * F^{c * (a + b * x)} * (\text{Sin}[d + e * x] / (e^2 + b^2 * c^2 * \text{Log}[F]^2)), x] - \text{Simp}[e * F^{c * (a + b * x)} * (\text{Cos}[d + e * x] / (e^2 + b^2 * c^2 * \text{Log}[F]^2)), x] /; F$

```
reeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Rule 5374

```
Int[(u_.)*(f_)^(ArcSec[(a_.) + (b_.)*(x_)]^(n_.)*(c_.)), x_Symbol] := Dist[
1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x,
ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int a^2 e^x \sin(x) dx, x, \sec^{-1}(ax)\right)}{a} \\ &= a \text{Subst}\left(\int e^x \sin(x) dx, x, \sec^{-1}(ax)\right) \\ &= \frac{1}{2} a e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{e^{\sec^{-1}(ax)}}{2x} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.87

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} a e^{\sec^{-1}(ax)} \left(\sqrt{1 - \frac{1}{a^2 x^2}} - \frac{1}{ax} \right)$$

```
[In] Integrate[E^ArcSec[a*x]/x^2,x]
```

```
[Out] (a*E^ArcSec[a*x]*(Sqrt[1 - 1/(a^2*x^2)] - 1/(a*x)))/2
```

Maple [F]

$$\int \frac{e^{\text{arcsec}(ax)}}{x^2} dx$$

```
[In] int(exp(arcsec(a*x))/x^2,x)
```

```
[Out] int(exp(arcsec(a*x))/x^2,x)
```

Fricas [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.59

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{(\sqrt{a^2x^2 - 1} - 1)e^{\operatorname{arcsec}(ax)}}{2x}$$

[In] integrate(exp(arcsec(a*x))/x^2,x, algorithm="fricas")

[Out] 1/2*(sqrt(a^2*x^2 - 1) - 1)*e^(arcsec(a*x))/x

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^2} dx$$

[In] integrate(exp(asec(a*x))/x**2,x)

[Out] Integral(exp(asec(a*x))/x**2, x)

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^2} dx$$

[In] integrate(exp(arcsec(a*x))/x^2,x, algorithm="maxima")

[Out] integrate(e^(arcsec(a*x))/x^2, x)

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.10

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \frac{1}{2} \left(\sqrt{-\frac{1}{a^2x^2} + 1} e^{\operatorname{arccos}(\frac{1}{ax})} - \frac{e^{\operatorname{arccos}(\frac{1}{ax})}}{ax} \right) a$$

[In] integrate(exp(arcsec(a*x))/x^2,x, algorithm="giac")

[Out] 1/2*(sqrt(-1/(a^2*x^2) + 1)*e^(arccos(1/(a*x))) - e^(arccos(1/(a*x)))/(a*x)) * a

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^2} dx = \int \frac{e^{\arccos(\frac{1}{ax})}}{x^2} dx$$

```
[In] int(exp(acos(1/(a*x)))/x^2,x)
```

```
[Out] int(exp(acos(1/(a*x)))/x^2, x)
```

3.48 $\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx$

Optimal result	314
Rubi [A] (verified)	314
Mathematica [A] (verified)	315
Maple [F]	316
Fricas [A] (verification not implemented)	316
Sympy [F]	316
Maxima [F]	316
Giac [F]	317
Mupad [F(-1)]	317

Optimal result

Integrand size = 10, antiderivative size = 41

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = -\frac{1}{5}a^2 e^{\sec^{-1}(ax)} \cos(2 \sec^{-1}(ax)) + \frac{1}{10}a^2 e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax))$$

[Out] $-1/5*a^2*\exp(\operatorname{arcsec}(a*x))*\cos(2*\operatorname{arcsec}(a*x))+1/10*a^2*\exp(\operatorname{arcsec}(a*x))*\sin(2*\operatorname{arcsec}(a*x))$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5374, 12, 4557, 4517}

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \frac{1}{10}a^2 e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax)) - \frac{1}{5}a^2 e^{\sec^{-1}(ax)} \cos(2 \sec^{-1}(ax))$$

[In] `Int[E^ArcSec[a*x]/x^3,x]`

[Out] $-1/5*(a^2*E^{\operatorname{ArcSec}[a*x]}*\operatorname{Cos}[2*\operatorname{ArcSec}[a*x]]) + (a^2*E^{\operatorname{ArcSec}[a*x]}*\operatorname{Sin}[2*\operatorname{ArcSec}[a*x]])/10$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 4517

`Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))*Sin[(d_.) + (e_.)*(x_)], x_Symbol] := Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x`

```
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]
```

Rule 4557

```
Int[Cos[(f_.) + (g_.)*(x_.)]^(n_.)*(F_)^((c_.)*((a_.) + (b_.)*(x_.)))*Sin[(d_.) + (e_.)*(x_.)]^(m_.), x_Symbol] := Int[ExpandTrigReduce[F^(c*(a + b*x)), Sin[d + e*x]^m*Cos[f + g*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e, f, g}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 5374

```
Int[(u_.)*(f_)^(ArcSec[(a_.) + (b_.)*(x_.)]^(n_.)*(c_.)), x_Symbol] := Dist[1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int a^3 e^x \cos(x) \sin(x) dx, x, \sec^{-1}(ax)\right)}{a} \\
 &= a^2 \text{Subst}\left(\int e^x \cos(x) \sin(x) dx, x, \sec^{-1}(ax)\right) \\
 &= a^2 \text{Subst}\left(\int \frac{1}{2} e^x \sin(2x) dx, x, \sec^{-1}(ax)\right) \\
 &= \frac{1}{2} a^2 \text{Subst}\left(\int e^x \sin(2x) dx, x, \sec^{-1}(ax)\right) \\
 &= -\frac{1}{5} a^2 e^{\sec^{-1}(ax)} \cos(2 \sec^{-1}(ax)) + \frac{1}{10} a^2 e^{\sec^{-1}(ax)} \sin(2 \sec^{-1}(ax))
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \frac{1}{10} a^2 e^{\sec^{-1}(ax)} (-2 \cos(2 \sec^{-1}(ax)) + \sin(2 \sec^{-1}(ax)))$$

```
[In] Integrate[E^ArcSec[a*x]/x^3, x]
```

```
[Out] (a^2*E^ArcSec[a*x]*(-2*Cos[2*ArcSec[a*x]] + Sin[2*ArcSec[a*x]]))/10
```

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x^3} dx$$

[In] int(exp(arcsec(a*x))/x^3,x)

[Out] int(exp(arcsec(a*x))/x^3,x)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 30, normalized size of antiderivative = 0.73

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \frac{(a^2x^2 + \sqrt{a^2x^2 - 1} - 2)e^{\operatorname{arcsec}(ax)}}{5x^2}$$

[In] integrate(exp(arcsec(a*x))/x^3,x, algorithm="fricas")

[Out] 1/5*(a^2*x^2 + sqrt(a^2*x^2 - 1) - 2)*e^(arcsec(a*x))/x^2

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^3} dx$$

[In] integrate(exp(asec(a*x))/x**3,x)

[Out] Integral(exp(asec(a*x))/x**3, x)

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{arcsec}(ax)}}{x^3} dx$$

[In] integrate(exp(arcsec(a*x))/x^3,x, algorithm="maxima")

[Out] integrate(e^(arcsec(a*x))/x^3, x)

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x^3} dx$$

[In] integrate(exp(arcsec(a*x))/x^3,x, algorithm="giac")

[Out] integrate(e^(arcsec(a*x))/x^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^3} dx = \int \frac{e^{\operatorname{acos}(\frac{1}{ax})}}{x^3} dx$$

[In] int(exp(acos(1/(a*x)))/x^3,x)

[Out] int(exp(acos(1/(a*x)))/x^3, x)

3.49 $\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx$

Optimal result	318
Rubi [A] (verified)	318
Mathematica [A] (verified)	320
Maple [F]	320
Fricas [A] (verification not implemented)	320
Sympy [F]	320
Maxima [F]	321
Giac [F]	321
Mupad [F(-1)]	321

Optimal result

Integrand size = 10, antiderivative size = 84

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{1}{8} a^3 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{a^2 e^{\sec^{-1}(ax)}}{8x} - \frac{3}{40} a^3 e^{\sec^{-1}(ax)} \cos(3 \sec^{-1}(ax)) + \frac{1}{40} a^3 e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax))$$

[Out] $-1/8*a^2*\exp(\operatorname{arcsec}(a*x))/x-3/40*a^3*\exp(\operatorname{arcsec}(a*x))*\cos(3*\operatorname{arcsec}(a*x))+1/40*a^3*\exp(\operatorname{arcsec}(a*x))*\sin(3*\operatorname{arcsec}(a*x))+1/8*a^3*\exp(\operatorname{arcsec}(a*x))*(1-1/a^2/x^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {5374, 12, 4557, 4517}

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = -\frac{3}{40} a^3 e^{\sec^{-1}(ax)} \cos(3 \sec^{-1}(ax)) + \frac{1}{40} a^3 e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax)) - \frac{a^2 e^{\sec^{-1}(ax)}}{8x} + \frac{1}{8} a^3 \sqrt{1 - \frac{1}{a^2 x^2}} e^{\sec^{-1}(ax)}$$

[In] Int[E^ArcSec[a*x]/x^4,x]

[Out] $(a^3 * E^{\operatorname{ArcSec}[a*x]} * \operatorname{Sqrt}[1 - 1/(a^2 * x^2)]) / 8 - (a^2 * E^{\operatorname{ArcSec}[a*x]}) / (8 * x) - (3 * a^3 * E^{\operatorname{ArcSec}[a*x]} * \operatorname{Cos}[3 * \operatorname{ArcSec}[a*x]]) / 40 + (a^3 * E^{\operatorname{ArcSec}[a*x]} * \operatorname{Sin}[3 * \operatorname{ArcSec}[a*x]]) / 40$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 4517

Int[(F_)^((c_)*((a_) + (b_)*(x_)))*Sin[(d_) + (e_)*(x_)], x_Symbol] :=
Simp[b*c*Log[F]*F^(c*(a + b*x))*(Sin[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x]
] - Simp[e*F^(c*(a + b*x))*(Cos[d + e*x]/(e^2 + b^2*c^2*Log[F]^2)), x] /; FreeQ[{F, a, b, c, d, e}, x] && NeQ[e^2 + b^2*c^2*Log[F]^2, 0]

Rule 4557

Int[Cos[(f_) + (g_)*(x_)]^(n_)*(F_)^((c_)*((a_) + (b_)*(x_)))*Sin[(d_) + (e_)*(x_)]^(m_), x_Symbol] := Int[ExpandTrigReduce[F^(c*(a + b*x)), Sin[d + e*x]^m*Cos[f + g*x]^n, x], x] /; FreeQ[{F, a, b, c, d, e, f, g}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 5374

Int[(u_)*(f_)^(ArcSec[(a_) + (b_)*(x_)]^(n_)*(c_)), x_Symbol] := Dist[1/b, Subst[Int[(u /. x -> -a/b + Sec[x]/b)*f^(c*x^n)*Sec[x]*Tan[x], x], x, ArcSec[a + b*x]], x] /; FreeQ[{a, b, c, f}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\text{Subst}\left(\int a^4 e^x \cos^2(x) \sin(x) dx, x, \sec^{-1}(ax)\right)}{a} \\ &= a^3 \text{Subst}\left(\int e^x \cos^2(x) \sin(x) dx, x, \sec^{-1}(ax)\right) \\ &= a^3 \text{Subst}\left(\int \left(\frac{1}{4} e^x \sin(x) + \frac{1}{4} e^x \sin(3x)\right) dx, x, \sec^{-1}(ax)\right) \\ &= \frac{1}{4} a^3 \text{Subst}\left(\int e^x \sin(x) dx, x, \sec^{-1}(ax)\right) + \frac{1}{4} a^3 \text{Subst}\left(\int e^x \sin(3x) dx, x, \sec^{-1}(ax)\right) \\ &= \frac{1}{8} a^3 e^{\sec^{-1}(ax)} \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{a^2 e^{\sec^{-1}(ax)}}{8x} \\ &\quad - \frac{3}{40} a^3 e^{\sec^{-1}(ax)} \cos(3 \sec^{-1}(ax)) + \frac{1}{40} a^3 e^{\sec^{-1}(ax)} \sin(3 \sec^{-1}(ax)) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 0.64

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{1}{40} a^3 e^{\sec^{-1}(ax)} \left(5 \sqrt{1 - \frac{1}{a^2 x^2}} - \frac{5}{ax} - 3 \cos(3 \sec^{-1}(ax)) + \sin(3 \sec^{-1}(ax)) \right)$$

[In] Integrate[E^ArcSec[a*x]/x^4,x]

[Out] (a^3*E^ArcSec[a*x]*(5*Sqrt[1 - 1/(a^2*x^2)] - 5/(a*x) - 3*Cos[3*ArcSec[a*x]] + Sin[3*ArcSec[a*x]]))/40

Maple [F]

$$\int \frac{e^{\operatorname{arcsec}(ax)}}{x^4} dx$$

[In] int(exp(arcsec(a*x))/x^4,x)

[Out] int(exp(arcsec(a*x))/x^4,x)

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 40, normalized size of antiderivative = 0.48

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \frac{(a^2 x^2 + (a^2 x^2 + 1) \sqrt{a^2 x^2 - 1} - 3) e^{\operatorname{arcsec}(ax)}}{10 x^3}$$

[In] integrate(exp(arcsec(a*x))/x^4,x, algorithm="fricas")

[Out] 1/10*(a^2*x^2 + (a^2*x^2 + 1)*sqrt(a^2*x^2 - 1) - 3)*e^(arcsec(a*x))/x^3

Sympy [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{asec}(ax)}}{x^4} dx$$

[In] integrate(exp(asec(a*x))/x**4,x)

[Out] Integral(exp(asec(a*x))/x**4, x)

Maxima [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x^4} dx$$

[In] integrate(exp(arcsec(a*x))/x^4,x, algorithm="maxima")

[Out] integrate(e^(arcsec(a*x))/x^4, x)

Giac [F]

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{(\operatorname{arcsec}(ax))}}{x^4} dx$$

[In] integrate(exp(arcsec(a*x))/x^4,x, algorithm="giac")

[Out] integrate(e^(arcsec(a*x))/x^4, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{e^{\sec^{-1}(ax)}}{x^4} dx = \int \frac{e^{\operatorname{acos}(\frac{1}{ax})}}{x^4} dx$$

[In] int(exp(acos(1/(a*x)))/x^4,x)

[Out] int(exp(acos(1/(a*x)))/x^4, x)

3.50 $\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx$

Optimal result	322
Rubi [A] (verified)	322
Mathematica [A] (verified)	324
Maple [A] (verified)	325
Fricas [F]	325
Sympy [F]	325
Maxima [F]	326
Giac [A] (verification not implemented)	326
Mupad [F(-1)]	326

Optimal result

Integrand size = 19, antiderivative size = 69

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{i \sec^{-1}(a+bx)^2}{2d} - \frac{\sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{d} + \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2d}$$

[Out] 1/2*I*arcsec(b*x+a)^2/d-arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/d+1/2*I*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)/d

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.421$, Rules used = {5364, 12, 5326, 4722, 3800, 2221, 2317, 2438}

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \frac{i \operatorname{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right)}{2d} + \frac{i \sec^{-1}(a+bx)^2}{2d} - \frac{\sec^{-1}(a+bx) \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)}{d}$$

[In] Int[ArcSec[a + b*x]/((a*d)/b + d*x), x]

[Out] ((I/2)*ArcSec[a + b*x]^2)/d - (ArcSec[a + b*x]*Log[1 + E^((2*I)*ArcSec[a + b*x])])/d + ((I/2)*PolyLog[2, -E^((2*I)*ArcSec[a + b*x])])/d

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3800

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (f_)*(x_)], x_Symbol] := Simp[I*((c + d*x)^(m + 1)/(d*(m + 1))), x] - Dist[2*I, Int[(c + d*x)^m*(E^(2*I*(e + f*x)))/(1 + E^(2*I*(e + f*x))), x], x] /; FreeQ[{c, d, e, f}, x] && IGtQ[m, 0]

Rule 4722

Int[((a_) + ArcCos[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := -Subst[Int[(a + b*x)^n*Tan[x], x], x, ArcCos[c*x]] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0]

Rule 5326

Int[((a_) + ArcSec[(c_)*(x_)])*(b_)^(n_)/(x_), x_Symbol] := -Subst[Int[(a + b*ArcCos[x/c])/x, x], x, 1/x] /; FreeQ[{a, b, c}, x]

Rule 5364

Int[((a_) + ArcSec[(c_) + (d_)*(x_)])*(b_)^(p_)*((e_) + (f_)*(x_))^(m_), x_Symbol] := Dist[1/d, Subst[Int[(f*(x/d))^m*(a + b*ArcSec[x])^p, x], x, c + d*x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[d*e - c*f, 0] && IGtQ[p, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\text{Subst}\left(\int \frac{b \sec^{-1}(x)}{dx} dx, x, a + bx\right)}{b} \\
 &= \frac{\text{Subst}\left(\int \frac{\sec^{-1}(x)}{x} dx, x, a + bx\right)}{d} \\
 &= -\frac{\text{Subst}\left(\int \frac{\arccos(x)}{x} dx, x, \frac{1}{a+bx}\right)}{d} \\
 &= \frac{\text{Subst}\left(\int x \tan(x) dx, x, \arccos\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{i \arccos\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{(2i) \text{Subst}\left(\int \frac{e^{2ix} x}{1+e^{2ix}} dx, x, \arccos\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{i \arccos\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\arccos\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad + \frac{\text{Subst}\left(\int \log(1 + e^{2ix}) dx, x, \arccos\left(\frac{1}{a+bx}\right)\right)}{d} \\
 &= \frac{i \arccos\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\arccos\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)}\right)}{d} \\
 &\quad - \frac{i \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2i \arccos\left(\frac{1}{a+bx}\right)}\right)}{2d} \\
 &= \frac{i \arccos\left(\frac{1}{a+bx}\right)^2}{2d} - \frac{\arccos\left(\frac{1}{a+bx}\right) \log\left(1 + e^{2i \arccos\left(\frac{1}{a+bx}\right)}\right)}{d} + \frac{i \text{PolyLog}\left(2, -e^{2i \arccos\left(\frac{1}{a+bx}\right)}\right)}{2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 59, normalized size of antiderivative = 0.86

$$\begin{aligned}
 &\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx \\
 &= \frac{i \left(\sec^{-1}(a + bx) \left(\sec^{-1}(a + bx) + 2i \log\left(1 + e^{2i \sec^{-1}(a+bx)}\right)\right) + \text{PolyLog}\left(2, -e^{2i \sec^{-1}(a+bx)}\right) \right)}{2d}
 \end{aligned}$$

[In] Integrate[ArcSec[a + b*x]/((a*d)/b + d*x), x]

[Out] ((I/2)*(ArcSec[a + b*x]*(ArcSec[a + b*x] + (2*I)*Log[1 + E^((2*I)*ArcSec[a + b*x])]) + PolyLog[2, -E^((2*I)*ArcSec[a + b*x])]))/d

Maple [A] (verified)

Time = 1.04 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.43

method	result	size
derivativedivides	$\frac{\frac{ib \operatorname{arcsec}(bx+a)^2}{2d} - \frac{b \operatorname{arcsec}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{d}}{b} + \frac{ib \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{2d}$	99
default	$\frac{\frac{ib \operatorname{arcsec}(bx+a)^2}{2d} - \frac{b \operatorname{arcsec}(bx+a) \ln\left(1 + \left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{d}}{b} + \frac{ib \operatorname{polylog}\left(2, -\left(\frac{1}{bx+a} + i\sqrt{1 - \frac{1}{(bx+a)^2}}\right)^2\right)}{2d}$	99

```
[In] int(arcsec(b*x+a)/(a*d/b+d*x),x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(1/2*I*b/d*arcsec(b*x+a)^2-b/d*arcsec(b*x+a)*ln(1+(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2)+1/2*I*b/d*polylog(2,-(1/(b*x+a)+I*(1-1/(b*x+a)^2)^(1/2))^2))
```

Fricas [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \int \frac{\operatorname{arcsec}(bx + a)}{dx + \frac{ad}{b}} dx$$

```
[In] integrate(arcsec(b*x+a)/(a*d/b+d*x),x, algorithm="fricas")
```

```
[Out] integral(b*arcsec(b*x + a)/(b*d*x + a*d), x)
```

Sympy [F]

$$\int \frac{\sec^{-1}(a + bx)}{\frac{ad}{b} + dx} dx = \frac{b \int \frac{\operatorname{asec}(a+bx)}{a+bx} dx}{d}$$

```
[In] integrate(asec(b*x+a)/(a*d/b+d*x),x)
```

```
[Out] b*Integral(asec(a + b*x)/(a + b*x), x)/d
```

Maxima [F]

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\operatorname{arcsec}(bx+a)}{dx+\frac{ad}{b}} dx$$

[In] integrate(arcsec(b*x+a)/(a*d/b+d*x),x, algorithm="maxima")

[Out] $-\frac{1}{2}*(2*b*d*\int \sqrt{b*x+a+1}*\sqrt{b*x+a-1}*\log(b*x+a)/(b^3*d*x^3+3*a*b^2*d*x^2+(3*a^2-1)*b*d*x+(a^3-a)*d),x)+2*I*b*d*\int \log(b*x+a)/(b^3*d*x^3+3*a*b^2*d*x^2+(3*a^2-1)*b*d*x+(a^3-a)*d),x)-2*\arctan(\sqrt{b*x+a+1}*\sqrt{b*x+a-1})*\log(b*x+a)+I*\log(b^2*x^2+2*a*b*x+a^2)*\log(b*x+a)-I*\log(b*x+a+1)*\log(b*x+a)-I*\log(b*x+a)^2-I*\log(b*x+a)*\log(-b*x-a+1)-I*\operatorname{dilog}(b*x+a)-I*\operatorname{dilog}(-b*x-a))/d$

Giac [A] (verification not implemented)

none

Time = 0.50 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.67

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = -\frac{1}{4}b^2 \left(\frac{2(bx+a)^2 \arccos\left(\frac{1}{((bx+a)(\frac{a}{bx+a}-1)-a)(\frac{a}{bx+a}-1)+a}\right)}{b^3d} - \frac{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2}+1}-1\right)}{b^3d} - \frac{1}{(bx+a)\left(\sqrt{-\frac{1}{(bx+a)^2}+1}\right)} \right)$$

[In] integrate(arcsec(b*x+a)/(a*d/b+d*x),x, algorithm="giac")

[Out] $-\frac{1}{4}*b^2*(2*(b*x+a)^2*\arccos(1/(((b*x+a)*(a/(b*x+a)-1)-a)*(a/(b*x+a)-1)+a)))/(b^3*d)-((b*x+a)*(sqrt(-1/(b*x+a)^2+1)-1)-1/(b*x+a)*(sqrt(-1/(b*x+a)^2+1)-1)))/(b^3*d)$

Mupad [F(-1)]

Timed out.

$$\int \frac{\sec^{-1}(a+bx)}{\frac{ad}{b}+dx} dx = \int \frac{\arccos\left(\frac{1}{a+bx}\right)}{dx+\frac{ad}{b}} dx$$

[In] int(acos(1/(a+b*x))/(d*x+(a*d)/b),x)

[Out] int(acos(1/(a+b*x))/(d*x+(a*d)/b),x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 327

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
            If[HypergeometricFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
            If[AppellFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
            If[Head[expn]===RootSum,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
            If[Head[expn]===Integrate || Head[expn]===Int,
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
            9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```



```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+"/"+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+"/"+str(ExpnType_optimal)

```

```

#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#    Albert Rich to use with Sagemath. This is used to
#    grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#    'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#    issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr, Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```