

Computer Algebra Independent Integration Tests

Summer 2023 edition

6-Hyperbolic-functions/6.2-Hyperbolic-cosine/170-6.2.7-hyper^m-
a+b-coshⁿ-^p

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 5:15pm

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	47
4	Appendix	543

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [85]. This is test number [170].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (85)	0.00 (0)
Mathematica	97.65 (83)	2.35 (2)
Maple	97.65 (83)	2.35 (2)
Fricas	89.41 (76)	10.59 (9)
Mupad	64.71 (55)	35.29 (30)
Giac	48.24 (41)	51.76 (44)
Maxima	40.00 (34)	60.00 (51)
Sympy	21.18 (18)	78.82 (67)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.000	0.000	0.000	0.000
Mathematica	71.765	0.000	25.882	2.353
Maple	37.647	34.118	25.882	2.353
Giac	23.529	18.824	5.882	51.765
Fricas	12.941	64.706	11.765	10.588
Maxima	8.235	27.059	4.706	60.000
Sympy	3.529	17.647	0.000	78.824
Mupad	0.000	64.706	0.000	35.294

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	2	0.00	100.00	0.00
Maple	2	100.00	0.00	0.00
Fricas	9	66.67	0.00	33.33
Mupad	30	0.00	100.00	0.00
Giac	44	97.73	0.00	2.27
Maxima	51	100.00	0.00	0.00
Sympy	67	55.22	43.28	1.49

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.10
Maxima	0.27
Giac	0.39
Fricas	0.40
Mathematica	1.04
Maple	3.74
Mupad	5.67
Sympy	9.85

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mathematica	72.94	1.10	51.00	1.00
Rubi	83.32	1.00	49.00	1.00
Maple	85.66	1.47	66.00	1.27
Maxima	124.91	2.79	56.00	2.11
Giac	154.07	2.17	45.00	1.64
Mupad	443.51	5.79	243.00	3.33
Sympy	4434.50	151.69	119.00	3.81
Fricas	18607.54	89.10	353.00	8.70

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	43

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	24
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 4, 5, 9, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 61, 63, 75, 76, 77, 78, 79, 80, 82, 83, 84, 85 }

B grade { }

C grade { 6, 7, 8, 10, 11, 12, 57, 58, 60, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 81 }

F normal fail { }

F(-1) timedout fail { 56, 59 }

F(-2) exception fail { }

Maple

A grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 37, 38, 39, 40, 43, 44, 45, 47, 48, 49, 50, 51, 54, 55, 76, 77, 78, 79, 84, 85 }

B grade { 13, 14, 15, 16, 17, 18, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 41, 42, 46, 52, 53, 63 }

C grade { 19, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 80, 81 }

F normal fail { 82, 83 }

F(-1) timedout fail { }

F(-2) exception fail { }

Fricas

A grade { 1, 2, 3, 15, 25, 44, 49, 54, 55, 84, 85 }

B grade { 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 43, 48, 51, 52, 53, 60, 61, 63, 66, 69, 70, 72, 73, 75, 76, 77, 78, 79, 80, 83 }

C grade { 56, 57, 58, 59, 62, 65, 68, 71, 74, 81 }

F normal fail { 41, 42, 45, 46, 47, 50 }

F(-1) timedout fail { }

F(-2) exception fail { 64, 67, 82 }

Maxima

A grade { 1, 3, 37, 44, 49, 54, 76 }

B grade { 2, 4, 5, 13, 14, 15, 16, 17, 18, 21, 23, 25, 27, 29, 31, 33, 34, 35, 36, 38, 39, 40, 63 }

C grade { 43, 48, 53, 80 }

F normal fail { 6, 7, 8, 9, 10, 11, 12, 19, 20, 22, 24, 26, 28, 30, 32, 41, 42, 45, 46, 47, 50, 51, 52, 55, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 77, 78, 79, 81, 82, 83, 84, 85 }

F(-1) timedout fail { }

F(-2) exception fail { }

Giac

A grade { 1, 2, 3, 4, 5, 15, 16, 19, 25, 27, 29, 33, 37, 39, 40, 72, 74, 75, 76, 81 }

B grade { 13, 14, 21, 23, 35, 36, 38, 44, 49, 54, 58, 59, 60, 61, 63, 71 }

C grade { 43, 48, 53, 62, 80 }

F normal fail { 6, 7, 8, 9, 10, 11, 12, 17, 18, 20, 22, 24, 26, 28, 30, 31, 32, 34, 41, 42, 45, 47, 50, 51, 52, 55, 56, 57, 64, 65, 66, 67, 68, 69, 70, 73, 77, 78, 79, 82, 83, 84, 85 }

F(-1) timedout fail { }

F(-2) exception fail { 46 }

Mupad

A grade { }

B grade { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 35, 36, 37, 38, 39, 40, 43, 44, 56, 57, 58, 59, 60, 61, 62, 63, 65, 68, 71, 74, 75, 76, 81 }

C grade { }

F normal fail { }

F(-1) timedout fail { 33, 34, 41, 42, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 64, 66, 67, 69, 70, 72, 73, 77, 78, 79, 80, 82, 83, 84, 85 }

F(-2) exception fail { }

Sympy

A grade { 2, 3, 19 }

B grade { 1, 9, 16, 26, 27, 35, 36, 37, 38, 39, 40, 58, 59, 63, 74 }

C grade { }

F normal fail { 4, 5, 10, 11, 17, 28, 29, 30, 31, 32, 41, 42, 43, 44, 45, 46, 51, 52, 53, 54, 55, 64, 65, 66, 67, 68, 69, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85 }

F(-1) timedout fail { 6, 7, 8, 12, 13, 14, 15, 18, 20, 21, 22, 23, 24, 25, 33, 34, 47, 48, 49, 50, 56, 57, 60, 61, 62, 71, 72, 73, 75 }

F(-2) exception fail { 70 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	20	20	19	26	25	14	153	26	25
N.S.	1	1.00	0.95	1.30	1.25	0.70	7.65	1.30	1.25
time (sec)	N/A	0.032	0.003	1.776	0.207	0.244	0.744	0.261	1.780

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	7	7	7	8	17	7	10	12	7
N.S.	1	1.00	1.00	1.14	2.43	1.00	1.43	1.71	1.00
time (sec)	N/A	0.033	0.002	0.523	0.203	0.252	0.444	0.263	1.706

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	6	6	6	7	6	6	3	6	6
N.S.	1	1.00	1.00	1.17	1.00	1.00	0.50	1.00	1.00
time (sec)	N/A	0.030	0.000	0.167	0.190	0.255	0.273	0.260	0.030

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	22	22	61	100	0	21	21
N.S.	1	1.00	1.16	1.16	3.21	5.26	0.00	1.11	1.11
time (sec)	N/A	0.038	0.005	2.949	0.199	0.245	0.000	0.266	1.703

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	32	28	135	216	0	27	27
N.S.	1	1.00	1.10	0.97	4.66	7.45	0.00	0.93	0.93
time (sec)	N/A	0.040	0.004	14.812	0.197	0.253	0.000	0.261	1.709

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	148	94	0	2346	0	0	805
N.S.	1	1.00	1.90	1.21	0.00	30.08	0.00	0.00	10.32
time (sec)	N/A	0.104	0.251	0.057	0.000	0.313	0.000	0.000	2.382

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	54	54	120	54	0	1064	0	0	548
N.S.	1	1.00	2.22	1.00	0.00	19.70	0.00	0.00	10.15
time (sec)	N/A	0.067	0.181	112.590	0.000	0.287	0.000	0.000	2.115

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	36	36	83	34	0	416	0	0	257
N.S.	1	1.00	2.31	0.94	0.00	11.56	0.00	0.00	7.14
time (sec)	N/A	0.049	0.193	4.456	0.000	0.278	0.000	0.000	2.186

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	25	17	0	300	66	0	16
N.S.	1	1.00	1.00	0.68	0.00	12.00	2.64	0.00	0.64
time (sec)	N/A	0.025	0.037	0.434	0.000	0.274	0.347	0.000	1.782

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	42	42	106	52	0	349	0	0	462
N.S.	1	1.00	2.52	1.24	0.00	8.31	0.00	0.00	11.00
time (sec)	N/A	0.052	0.157	1.907	0.000	0.280	0.000	0.000	2.210

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	61	61	154	87	0	1332	0	0	2225
N.S.	1	1.00	2.52	1.43	0.00	21.84	0.00	0.00	36.48
time (sec)	N/A	0.084	0.308	6.491	0.000	0.296	0.000	0.000	8.262

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	94	94	229	135	0	5326	0	0	5056
N.S.	1	1.00	2.44	1.44	0.00	56.66	0.00	0.00	53.79
time (sec)	N/A	0.152	0.536	29.974	0.000	0.352	0.000	0.000	17.032

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	76	289	651	1308	0	166	248
N.S.	1	1.00	0.86	3.28	7.40	14.86	0.00	1.89	2.82
time (sec)	N/A	0.125	0.153	0.078	0.307	0.291	0.000	0.279	2.571

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	52	185	348	568	0	103	146
N.S.	1	1.00	0.88	3.14	5.90	9.63	0.00	1.75	2.47
time (sec)	N/A	0.094	0.098	26.566	0.292	0.281	0.000	0.266	2.115

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	36	88	120	300	0	52	79
N.S.	1	1.00	0.92	2.26	3.08	7.69	0.00	1.33	2.03
time (sec)	N/A	0.054	0.083	0.729	0.295	0.276	0.000	0.263	0.250

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	78	53	293	10924	39	267
N.S.	1	1.00	1.00	2.69	1.83	10.10	376.69	1.34	9.21
time (sec)	N/A	0.017	0.465	0.109	0.306	0.273	24.042	0.274	0.395

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	59	162	161	1875	0	0	245
N.S.	1	1.00	1.00	2.75	2.73	31.78	0.00	0.00	4.15
time (sec)	N/A	0.078	0.170	14.045	0.331	0.281	0.000	0.000	2.380

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	89	89	92	248	307	4977	0	0	333
N.S.	1	1.00	1.03	2.79	3.45	55.92	0.00	0.00	3.74
time (sec)	N/A	0.083	0.300	54.962	0.313	0.310	0.000	0.000	2.497

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	A	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	98	98	77	26	0	305	85	80	205
N.S.	1	1.00	0.79	0.27	0.00	3.11	0.87	0.82	2.09
time (sec)	N/A	0.105	0.164	0.532	0.000	0.256	0.482	0.277	4.709

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	78	78	86	206	0	2508	0	0	293
N.S.	1	1.00	1.10	2.64	0.00	32.15	0.00	0.00	3.76
time (sec)	N/A	0.072	0.233	1.423	0.000	0.283	0.000	0.000	2.345

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	76	170	651	1245	0	150	178
N.S.	1	1.00	0.86	1.93	7.40	14.15	0.00	1.70	2.02
time (sec)	N/A	0.158	0.163	0.843	0.324	0.283	0.000	0.269	2.260

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	56	56	61	146	0	1184	0	0	243
N.S.	1	1.00	1.09	2.61	0.00	21.14	0.00	0.00	4.34
time (sec)	N/A	0.053	0.137	0.518	0.000	0.280	0.000	0.000	2.180

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	52	120	347	573	0	95	142
N.S.	1	1.00	0.88	2.03	5.88	9.71	0.00	1.61	2.41
time (sec)	N/A	0.074	0.095	0.322	0.294	0.282	0.000	0.309	2.068

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	101	0	498	0	0	204
N.S.	1	1.00	1.00	2.66	0.00	13.11	0.00	0.00	5.37
time (sec)	N/A	0.043	0.028	0.214	0.000	0.278	0.000	0.000	2.057

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	36	92	120	317	0	50	376
N.S.	1	1.00	0.92	2.36	3.08	8.13	0.00	1.28	9.64
time (sec)	N/A	0.052	0.451	0.109	0.285	0.265	0.000	0.278	2.328

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	B	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	66	0	337	55498	0	87
N.S.	1	1.00	1.00	2.28	0.00	11.62	1913.72	0.00	3.00
time (sec)	N/A	0.023	0.012	0.104	0.000	0.263	106.055	0.000	2.072

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	78	53	293	10924	39	267
N.S.	1	1.00	1.00	2.69	1.83	10.10	376.69	1.34	9.21
time (sec)	N/A	0.018	0.019	0.000	0.280	0.265	24.281	0.261	0.003

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	41	41	45	85	0	360	0	0	208
N.S.	1	1.00	1.10	2.07	0.00	8.78	0.00	0.00	5.07
time (sec)	N/A	0.039	0.097	0.313	0.000	0.272	0.000	0.000	2.234

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	38	38	38	107	70	457	0	58	108
N.S.	1	1.00	1.00	2.82	1.84	12.03	0.00	1.53	2.84
time (sec)	N/A	0.051	0.366	0.534	0.284	0.261	0.000	0.276	0.311

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	59	59	58	123	0	963	0	0	447
N.S.	1	1.00	0.98	2.08	0.00	16.32	0.00	0.00	7.58
time (sec)	N/A	0.075	0.146	0.836	0.000	0.276	0.000	0.000	2.536

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	55	142	119	1377	0	0	239
N.S.	1	1.00	1.00	2.58	2.16	25.04	0.00	0.00	4.35
time (sec)	N/A	0.073	0.379	1.194	0.307	0.289	0.000	0.000	2.260

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	90	90	86	183	0	3239	0	0	1305
N.S.	1	1.00	0.96	2.03	0.00	35.99	0.00	0.00	14.50
time (sec)	N/A	0.112	0.245	1.742	0.000	0.303	0.000	0.000	43.632

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	68	171	134	1239	0	104	0
N.S.	1	1.00	1.05	2.63	2.06	19.06	0.00	1.60	0.00
time (sec)	N/A	0.047	5.084	0.356	0.295	0.271	0.000	0.269	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	107	107	106	264	344	5117	0	0	0
N.S.	1	1.00	0.99	2.47	3.21	47.82	0.00	0.00	0.00
time (sec)	N/A	0.104	5.343	0.852	0.304	0.335	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	36	34	66	60	34	50
N.S.	1	1.00	1.00	2.40	2.27	4.40	4.00	2.27	3.33
time (sec)	N/A	0.009	0.062	0.084	0.280	0.259	0.279	0.264	0.149

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	35	35	35	60	59	214	211	59	76
N.S.	1	1.00	1.00	1.71	1.69	6.11	6.03	1.69	2.17
time (sec)	N/A	0.018	0.102	0.210	0.283	0.256	1.162	0.272	1.886

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	51	51	51	72	83	575	428	71	112
N.S.	1	1.00	1.00	1.41	1.63	11.27	8.39	1.39	2.20
time (sec)	N/A	0.039	0.169	0.456	0.284	0.256	4.101	0.261	1.841

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	2	2	2	3	10	20	14	10	10
N.S.	1	1.00	1.00	1.50	5.00	10.00	7.00	5.00	5.00
time (sec)	N/A	0.014	0.003	0.046	0.195	0.248	0.212	0.267	0.067

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	17	16	49	84	34	18	18
N.S.	1	1.00	1.55	1.45	4.45	7.64	3.09	1.64	1.64
time (sec)	N/A	0.016	0.003	0.079	0.191	0.243	0.524	0.297	1.831

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	19	19	27	21	111	185	54	24	24
N.S.	1	1.00	1.42	1.11	5.84	9.74	2.84	1.26	1.26
time (sec)	N/A	0.032	0.003	0.092	0.198	0.240	1.572	0.275	0.072

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	53	114	0	0	0	0	0
N.S.	1	1.00	1.08	2.33	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.031	0.067	1.100	0.000	0.000	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	18	58	0	0	0	0	0
N.S.	1	1.00	1.06	3.41	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.008	0.021	0.623	0.000	0.000	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	15	11	33	0	31	13
N.S.	1	1.00	1.00	1.15	0.85	2.54	0.00	2.38	1.00
time (sec)	N/A	0.017	0.024	0.133	0.291	0.266	0.000	0.286	1.860

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	11	11	11	14	11	2	0	31	11
N.S.	1	1.00	1.00	1.27	1.00	0.18	0.00	2.82	1.00
time (sec)	N/A	0.014	0.023	0.109	0.280	0.256	0.000	0.280	1.779

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	40	62	0	0	0	0	0
N.S.	1	1.00	1.03	1.59	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.016	0.035	0.633	0.000	0.000	0.000	0.000	0.000

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	F	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	133	135	321	0	0	0	0	0
N.S.	1	1.00	1.02	2.41	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.122	0.377	1.327	0.000	0.000	0.000	0.000	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	55	55	51	99	0	0	0	0	0
N.S.	1	1.00	0.93	1.80	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.050	0.041	0.502	0.000	0.000	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	C	B	F(-1)	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	33	33	25	21	23	53	0	66	0
N.S.	1	1.00	0.76	0.64	0.70	1.61	0.00	2.00	0.00
time (sec)	N/A	0.027	0.050	0.124	0.277	0.257	0.000	0.300	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	23	21	23	19	0	66	0
N.S.	1	1.00	0.79	0.72	0.79	0.66	0.00	2.28	0.00
time (sec)	N/A	0.022	0.040	0.122	0.301	0.247	0.000	0.299	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	F	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	78	96	0	0	0	0	0
N.S.	1	1.00	0.77	0.95	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.076	0.066	0.425	0.000	0.000	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	49	49	53	66	0	132	0	0	0
N.S.	1	1.00	1.08	1.35	0.00	2.69	0.00	0.00	0.00
time (sec)	N/A	0.031	0.055	0.179	0.000	0.079	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	18	45	0	42	0	0	0
N.S.	1	1.00	1.06	2.65	0.00	2.47	0.00	0.00	0.00
time (sec)	N/A	0.008	0.030	0.143	0.000	0.078	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	C	B	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	17	17	30	34	19	49	0	40	0
N.S.	1	1.00	1.76	2.00	1.12	2.88	0.00	2.35	0.00
time (sec)	N/A	0.021	0.035	0.119	0.283	0.253	0.000	0.290	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	28	16	17	17	0	39	0
N.S.	1	1.00	1.87	1.07	1.13	1.13	0.00	2.60	0.00
time (sec)	N/A	0.017	0.037	0.121	0.305	0.257	0.000	0.300	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	40	61	0	39	0	0	0
N.S.	1	1.00	1.03	1.56	0.00	1.00	0.00	0.00	0.00
time (sec)	N/A	0.017	0.036	0.173	0.000	0.074	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F(-1)	C	F	C	F(-1)	F	B
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	288	288	0	100	0	18612	0	0	633
N.S.	1	1.00	0.00	0.35	0.00	64.62	0.00	0.00	2.20
time (sec)	N/A	0.391	0.000	0.550	0.000	0.976	0.000	0.000	7.029

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F(-1)	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	288	288	105	94	0	18612	0	0	633
N.S.	1	1.00	0.36	0.33	0.00	64.62	0.00	0.00	2.20
time (sec)	N/A	0.198	5.055	0.532	0.000	0.983	0.000	0.000	6.814

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	91	91	133	46	0	226	330	275	291
N.S.	1	1.00	1.46	0.51	0.00	2.48	3.63	3.02	3.20
time (sec)	N/A	0.112	5.498	0.070	0.000	0.275	1.513	0.346	4.324

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F(-1)	C	F	C	B	B	B
verified	N/A	Yes	N/A	Yes	TBD	TBD	TBD	TBD	TBD
size	95	95	0	46	0	234	320	275	295
N.S.	1	1.00	0.00	0.48	0.00	2.46	3.37	2.89	3.11
time (sec)	N/A	0.107	0.000	0.072	0.000	0.265	1.560	0.322	4.323

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	361	485	121	96	0	771	0	1781	1563
N.S.	1	1.34	0.34	0.27	0.00	2.14	0.00	4.93	4.33
time (sec)	N/A	0.852	1.984	0.334	0.000	0.291	0.000	2.365	8.766

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	109	96	0	779	0	1697	1487
N.S.	1	1.00	1.08	0.95	0.00	7.71	0.00	16.80	14.72
time (sec)	N/A	0.096	0.353	0.338	0.000	0.296	0.000	2.464	10.275

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F(-1)	C	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	176	176	45	36	0	133	0	281	205
N.S.	1	1.00	0.26	0.20	0.00	0.76	0.00	1.60	1.16
time (sec)	N/A	0.118	4.075	0.223	0.000	0.263	0.000	0.312	1.106

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	25	25	24	44	45	115	75	43	61
N.S.	1	1.00	0.96	1.76	1.80	4.60	3.00	1.72	2.44
time (sec)	N/A	0.014	1.068	0.187	0.302	0.255	0.634	0.256	0.133

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	494	494	139	156	0	0	0	0	0
N.S.	1	1.00	0.28	0.32	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.632	5.052	0.914	0.000	0.000	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	171	171	132	140	0	15201	0	0	844
N.S.	1	1.00	0.77	0.82	0.00	88.89	0.00	0.00	4.94
time (sec)	N/A	0.187	5.040	0.940	0.000	1.206	0.000	0.000	61.873

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	245	245	158	184	0	661324	0	0	0
N.S.	1	1.00	0.64	0.75	0.00	2699.28	0.00	0.00	0.00
time (sec)	N/A	0.370	5.068	2.375	0.000	3.140	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	494	494	139	150	0	0	0	0	0
N.S.	1	1.00	0.28	0.30	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.373	5.045	0.889	0.000	0.000	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	F	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	175	175	132	140	0	16379	0	0	855
N.S.	1	1.00	0.75	0.80	0.00	93.59	0.00	0.00	4.89
time (sec)	N/A	0.191	5.041	0.937	0.000	1.159	0.000	0.000	62.083

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	213	213	158	184	0	631813	0	0	0
N.S.	1	1.00	0.74	0.86	0.00	2966.26	0.00	0.00	0.00
time (sec)	N/A	0.178	2.635	2.384	0.000	3.107	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	223	223	445	62	0	836	0	0	0
N.S.	1	1.00	2.00	0.28	0.00	3.75	0.00	0.00	0.00
time (sec)	N/A	0.441	5.028	0.134	0.000	0.298	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F(-1)	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	83	83	68	71	0	315	0	140	337
N.S.	1	1.00	0.82	0.86	0.00	3.80	0.00	1.69	4.06
time (sec)	N/A	0.071	5.049	0.861	0.000	0.266	0.000	0.272	2.667

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F(-1)	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	129	129	127	47	0	737	0	1	0
N.S.	1	1.00	0.98	0.36	0.00	5.71	0.00	0.01	0.00
time (sec)	N/A	0.146	5.020	1.829	0.000	0.282	0.000	0.280	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	B	F(-1)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	205	205	445	64	0	852	0	0	0
N.S.	1	1.00	2.17	0.31	0.00	4.16	0.00	0.00	0.00
time (sec)	N/A	0.365	5.028	0.130	0.000	0.297	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	71	71	111	47	0	358	632	10	329
N.S.	1	1.00	1.56	0.66	0.00	5.04	8.90	0.14	4.63
time (sec)	N/A	0.098	1.235	0.179	0.000	0.277	9.151	0.266	4.603

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	F	B	F(-1)	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	69	69	64	81	0	358	0	45	271
N.S.	1	1.00	0.93	1.17	0.00	5.19	0.00	0.65	3.93
time (sec)	N/A	0.067	1.612	1.665	0.000	0.262	0.000	0.397	3.162

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	23	47	0	23	27
N.S.	1	1.00	1.00	1.47	1.53	3.13	0.00	1.53	1.80
time (sec)	N/A	0.022	0.007	0.425	0.194	0.253	0.000	0.255	0.096

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	39	39	39	42	0	357	0	0	0
N.S.	1	1.00	1.00	1.08	0.00	9.15	0.00	0.00	0.00
time (sec)	N/A	0.051	0.019	0.163	0.000	0.413	0.000	0.000	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	26	26	26	31	0	248	0	0	0
N.S.	1	1.00	1.00	1.19	0.00	9.54	0.00	0.00	0.00
time (sec)	N/A	0.057	0.012	0.140	0.000	0.304	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	13	12	0	63	0	0	0
N.S.	1	1.00	1.00	0.92	0.00	4.85	0.00	0.00	0.00
time (sec)	N/A	0.028	0.008	0.113	0.000	0.274	0.000	0.000	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	C	C	B	F	C	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	13	13	16	19	7	112	0	38	0
N.S.	1	1.00	1.23	1.46	0.54	8.62	0.00	2.92	0.00
time (sec)	N/A	0.041	0.010	0.164	0.292	0.268	0.000	0.276	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	C	F	C	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	153	153	145	93	0	1435	0	191	1173
N.S.	1	1.00	0.95	0.61	0.00	9.38	0.00	1.25	7.67
time (sec)	N/A	0.168	0.352	4.073	0.000	0.955	0.000	0.284	0.879

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	28	28	28	0	0	0	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.057	0.013	0.000	0.000	0.000	0.000	0.000	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	45	45	45	0	0	1648	0	0	0
N.S.	1	1.00	1.00	0.00	0.00	36.62	0.00	0.00	0.00
time (sec)	N/A	0.056	0.016	0.000	0.000	0.786	0.000	0.000	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	29	29	29	24	0	113	0	0	0
N.S.	1	1.00	1.00	0.83	0.00	3.90	0.00	0.00	0.00
time (sec)	N/A	0.066	0.016	0.118	0.000	0.257	0.000	0.000	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	47	47	45	38	0	156	0	0	0
N.S.	1	1.00	0.96	0.81	0.00	3.32	0.00	0.00	0.00
time (sec)	N/A	0.062	0.016	0.067	0.000	0.263	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [62] had the largest ratio of [.7500000000000000000]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	3	3	1.00	16	0.188
2	A	2	2	1.00	16	0.125
3	A	2	2	1.00	16	0.125
4	A	3	2	1.00	16	0.125
5	A	3	2	1.00	16	0.125
6	A	4	3	1.00	15	0.200
7	A	4	3	1.00	15	0.200
8	A	3	3	1.00	15	0.200
9	A	2	2	1.00	13	0.154
10	A	4	4	1.00	13	0.308
11	A	5	5	1.00	15	0.333
12	A	6	6	1.00	15	0.400
13	A	6	6	1.00	15	0.400
14	A	5	5	1.00	15	0.333
15	A	4	4	1.00	15	0.267
16	A	2	2	1.00	10	0.200
17	A	4	3	1.00	15	0.200
18	A	4	3	1.00	15	0.200
19	A	7	7	1.00	13	0.538
20	A	4	3	1.00	15	0.200
21	A	6	6	1.00	15	0.400
22	A	4	3	1.00	15	0.200
23	A	5	5	1.00	15	0.333
24	A	3	3	1.00	15	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
25	A	3	3	1.00	15	0.200
26	A	2	2	1.00	13	0.154
27	A	2	2	1.00	10	0.200
28	A	4	4	1.00	13	0.308
29	A	3	3	1.00	15	0.200
30	A	5	5	1.00	15	0.333
31	A	4	3	1.00	15	0.200
32	A	6	6	1.00	15	0.400
33	A	4	4	1.00	10	0.400
34	A	5	5	1.00	10	0.500
35	A	2	2	1.00	8	0.250
36	A	4	4	1.00	8	0.500
37	A	5	5	1.00	8	0.625
38	A	3	3	1.00	10	0.300
39	A	3	2	1.00	10	0.200
40	A	3	2	1.00	10	0.200
41	A	2	2	1.00	12	0.167
42	A	1	1	1.00	10	0.100
43	A	3	3	1.00	12	0.250
44	A	3	3	1.00	10	0.300
45	A	2	2	1.00	12	0.167
46	A	6	6	1.00	12	0.500
47	A	4	4	1.00	10	0.400
48	A	4	4	1.00	12	0.333
49	A	4	4	1.00	10	0.400
50	A	6	6	1.00	12	0.500
51	A	2	2	1.00	12	0.167
52	A	1	1	1.00	10	0.100
53	A	3	3	1.00	12	0.250
54	A	3	3	1.00	10	0.300
55	A	2	2	1.00	12	0.167
56	A	8	3	1.00	10	0.300
57	A	8	3	1.00	11	0.273
58	A	7	5	1.00	8	0.625
59	A	7	5	1.00	10	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
60	A	10	6	1.34	10	0.600
61	A	4	3	1.00	11	0.273
62	A	10	6	1.00	8	0.750
63	A	3	3	1.00	10	0.300
64	A	12	3	1.00	10	0.300
65	A	7	3	1.00	10	0.300
66	A	9	3	1.00	10	0.300
67	A	12	3	1.00	11	0.273
68	A	7	3	1.00	11	0.273
69	A	9	3	1.00	11	0.273
70	A	11	5	1.00	8	0.625
71	A	7	3	1.00	8	0.375
72	A	9	3	1.00	8	0.375
73	A	11	5	1.00	10	0.500
74	A	8	6	1.00	10	0.600
75	A	10	6	1.00	10	0.600
76	A	4	4	1.00	11	0.364
77	A	4	4	1.00	15	0.267
78	A	3	3	1.00	15	0.200
79	A	3	3	1.00	13	0.231
80	A	4	4	1.00	15	0.267
81	A	11	10	1.00	15	0.667
82	A	4	4	1.00	15	0.267
83	A	5	5	1.00	15	0.333
84	A	4	4	1.00	15	0.267
85	A	5	5	1.00	15	0.333

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int \frac{\sinh^4(x)}{a-a \cosh^2(x)} dx$	51
3.2	$\int \frac{\sinh^3(x)}{a-a \cosh^2(x)} dx$	55
3.3	$\int \frac{\sinh^2(x)}{a-a \cosh^2(x)} dx$	58
3.4	$\int \frac{\operatorname{csch}^2(x)}{a-a \cosh^2(x)} dx$	61
3.5	$\int \frac{\operatorname{csch}^4(x)}{a-a \cosh^2(x)} dx$	65
3.6	$\int \frac{\sinh^7(x)}{a+b \cosh^2(x)} dx$	69
3.7	$\int \frac{\sinh^5(x)}{a+b \cosh^2(x)} dx$	75
3.8	$\int \frac{\sinh^3(x)}{a+b \cosh^2(x)} dx$	80
3.9	$\int \frac{\sinh(x)}{a+b \cosh^2(x)} dx$	85
3.10	$\int \frac{\operatorname{csch}(x)}{a+b \cosh^2(x)} dx$	89
3.11	$\int \frac{\operatorname{csch}^3(x)}{a+b \cosh^2(x)} dx$	94
3.12	$\int \frac{\operatorname{csch}^5(x)}{a+b \cosh^2(x)} dx$	101
3.13	$\int \frac{\sinh^6(x)}{a+b \cosh^2(x)} dx$	109
3.14	$\int \frac{\sinh^4(x)}{a+b \cosh^2(x)} dx$	117
3.15	$\int \frac{\sinh^2(x)}{a+b \cosh^2(x)} dx$	123
3.16	$\int \frac{1}{a+b \cosh^2(x)} dx$	128
3.17	$\int \frac{\operatorname{csch}^4(x)}{a+b \cosh^2(x)} dx$	138
3.18	$\int \frac{\operatorname{csch}^6(x)}{a+b \cosh^2(x)} dx$	144
3.19	$\int \frac{\sinh(x)}{4-3 \cosh^3(x)} dx$	152

3.20	$\int \frac{\cosh^7(x)}{a+b \cosh^2(x)} dx$	159
3.21	$\int \frac{\cosh^6(x)}{a+b \cosh^2(x)} dx$	165
3.22	$\int \frac{\cosh^5(x)}{a+b \cosh^2(x)} dx$	173
3.23	$\int \frac{\cosh^4(x)}{a+b \cosh^2(x)} dx$	178
3.24	$\int \frac{\cosh^3(x)}{a+b \cosh^2(x)} dx$	184
3.25	$\int \frac{\cosh^2(x)}{a+b \cosh^2(x)} dx$	189
3.26	$\int \frac{\cosh(x)}{a+b \cosh^2(x)} dx$	194
3.27	$\int \frac{1}{a+b \cosh^2(x)} dx$	228
3.28	$\int \frac{\operatorname{sech}(x)}{a+b \cosh^2(x)} dx$	238
3.29	$\int \frac{\operatorname{sech}^2(x)}{a+b \cosh^2(x)} dx$	243
3.30	$\int \frac{\operatorname{sech}^3(x)}{a+b \cosh^2(x)} dx$	248
3.31	$\int \frac{\operatorname{sech}^4(x)}{a+b \cosh^2(x)} dx$	254
3.32	$\int \frac{\operatorname{sech}^5(x)}{a+b \cosh^2(x)} dx$	259
3.33	$\int \frac{1}{(a+b \cosh^2(x))^2} dx$	267
3.34	$\int \frac{1}{(a+b \cosh^2(x))^3} dx$	272
3.35	$\int \frac{1}{1+\cosh^2(x)} dx$	277
3.36	$\int \frac{1}{(1+\cosh^2(x))^2} dx$	281
3.37	$\int \frac{1}{(1+\cosh^2(x))^3} dx$	286
3.38	$\int \frac{1}{1-\cosh^2(x)} dx$	293
3.39	$\int \frac{1}{(1-\cosh^2(x))^2} dx$	297
3.40	$\int \frac{1}{(1-\cosh^2(x))^3} dx$	301
3.41	$\int \sqrt{a+b \cosh^2(x)} dx$	305
3.42	$\int \sqrt{1+\cosh^2(x)} dx$	309
3.43	$\int \sqrt{1-\cosh^2(x)} dx$	312
3.44	$\int \sqrt{-1+\cosh^2(x)} dx$	316
3.45	$\int \sqrt{-1-\cosh^2(x)} dx$	320
3.46	$\int (a+b \cosh^2(x))^{3/2} dx$	324
3.47	$\int (1+\cosh^2(x))^{3/2} dx$	329
3.48	$\int (1-\cosh^2(x))^{3/2} dx$	333
3.49	$\int (-1+\cosh^2(x))^{3/2} dx$	337
3.50	$\int (-1-\cosh^2(x))^{3/2} dx$	341
3.51	$\int \frac{1}{\sqrt{a+b \cosh^2(x)}} dx$	346

3.52	$\int \frac{1}{\sqrt{1+\cosh^2(x)}} dx$	350
3.53	$\int \frac{1}{\sqrt{1-\cosh^2(x)}} dx$	353
3.54	$\int \frac{1}{\sqrt{-1+\cosh^2(x)}} dx$	357
3.55	$\int \frac{1}{\sqrt{-1-\cosh^2(x)}} dx$	361
3.56	$\int \frac{1}{a+b \cosh^3(x)} dx$	365
3.57	$\int \frac{1}{a-b \cosh^3(x)} dx$	371
3.58	$\int \frac{1}{1+\cosh^3(x)} dx$	377
3.59	$\int \frac{1}{1-\cosh^3(x)} dx$	384
3.60	$\int \frac{1}{a+b \cosh^4(x)} dx$	391
3.61	$\int \frac{1}{a-b \cosh^4(x)} dx$	402
3.62	$\int \frac{1}{1+\cosh^4(x)} dx$	410
3.63	$\int \frac{1}{1-\cosh^4(x)} dx$	417
3.64	$\int \frac{1}{a+b \cosh^5(x)} dx$	421
3.65	$\int \frac{1}{a+b \cosh^6(x)} dx$	428
3.66	$\int \frac{1}{a+b \cosh^8(x)} dx$	434
3.67	$\int \frac{1}{a-b \cosh^5(x)} dx$	439
3.68	$\int \frac{1}{a-b \cosh^6(x)} dx$	446
3.69	$\int \frac{1}{a-b \cosh^8(x)} dx$	452
3.70	$\int \frac{1}{1+\cosh^5(x)} dx$	457
3.71	$\int \frac{1}{1+\cosh^6(x)} dx$	463
3.72	$\int \frac{1}{1+\cosh^8(x)} dx$	469
3.73	$\int \frac{1}{1-\cosh^5(x)} dx$	476
3.74	$\int \frac{1}{1-\cosh^6(x)} dx$	482
3.75	$\int \frac{1}{1-\cosh^8(x)} dx$	490
3.76	$\int \frac{\tanh(x)}{1+\cosh^2(x)} dx$	496
3.77	$\int \sqrt{a+b \cosh^2(x)} \tanh(x) dx$	500
3.78	$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^2(x)}} dx$	504
3.79	$\int \frac{\tanh(x)}{\sqrt{1+\cosh^2(x)}} dx$	508
3.80	$\int \frac{\tanh(x)}{\sqrt{1-\cosh^2(x)}} dx$	512
3.81	$\int \frac{\tanh^3(x)}{a+b \cosh^3(x)} dx$	517
3.82	$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^3(x)}} dx$	526
3.83	$\int \sqrt{a+b \cosh^3(x)} \tanh(x) dx$	530
3.84	$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^n(x)}} dx$	535

3.85	$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx$	539
------	--	-----

3.1 $\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx$

Optimal result	51
Rubi [A] (verified)	51
Mathematica [A] (verified)	52
Maple [A] (verified)	52
Fricas [A] (verification not implemented)	53
Sympy [B] (verification not implemented)	53
Maxima [A] (verification not implemented)	54
Giac [A] (verification not implemented)	54
Mupad [B] (verification not implemented)	54

Optimal result

Integrand size = 16, antiderivative size = 20

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = \frac{x}{2a} - \frac{\cosh(x) \sinh(x)}{2a}$$

[Out] 1/2*x/a-1/2*cosh(x)*sinh(x)/a

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.188$, Rules used = {3254, 2715, 8}

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = \frac{x}{2a} - \frac{\sinh(x) \cosh(x)}{2a}$$

[In] Int[Sinh[x]^4/(a - a*Cosh[x]^2),x]

[Out] x/(2*a) - (Cosh[x]*Sinh[x])/(2*a)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n-1)/(d*n), x] + Dist[b^2*((n-1)/n), Int[(b*Sin[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3254

```
Int[(u_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^p, x_Symbol] :> Dist[
a^p, Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p
}, x] && EqQ[a + b, 0] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\int \sinh^2(x) dx}{a} \\ &= -\frac{\cosh(x) \sinh(x)}{2a} + \frac{\int 1 dx}{2a} \\ &= \frac{x}{2a} - \frac{\cosh(x) \sinh(x)}{2a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = -\frac{-\frac{x}{2} + \frac{1}{4} \sinh(2x)}{a}$$

```
[In] Integrate[Sinh[x]^4/(a - a*Cosh[x]^2),x]
```

```
[Out] -((-1/2*x + Sinh[2*x]/4)/a)
```

Maple [A] (verified)

Time = 1.78 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30

method	result	size
risch	$\frac{x}{2a} - \frac{e^{2x}}{8a} + \frac{e^{-2x}}{8a}$	26
default	$\frac{\frac{1}{2(\tanh(\frac{x}{2})+1)^2} - \frac{1}{2(\tanh(\frac{x}{2})+1)} + \frac{\ln(\tanh(\frac{x}{2})+1)}{2} - \frac{1}{2(\tanh(\frac{x}{2})-1)^2} - \frac{1}{2(\tanh(\frac{x}{2})-1)} - \frac{\ln(\tanh(\frac{x}{2})-1)}{2}}{a}$	65

```
[In] int(sinh(x)^4/(a-a*cosh(x)^2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*x/a-1/8/a*exp(2*x)+1/8/a*exp(-2*x)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.70

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x) \sinh(x) - x}{2a}$$

[In] integrate(sinh(x)^4/(a-a*cosh(x)^2),x, algorithm="fricas")

[Out] -1/2*(cosh(x)*sinh(x) - x)/a

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(14) = 28.

Time = 0.74 (sec) , antiderivative size = 153, normalized size of antiderivative = 7.65

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = \frac{x \tanh^4\left(\frac{x}{2}\right)}{2a \tanh^4\left(\frac{x}{2}\right) - 4a \tanh^2\left(\frac{x}{2}\right) + 2a} - \frac{2x \tanh^2\left(\frac{x}{2}\right)}{2a \tanh^4\left(\frac{x}{2}\right) - 4a \tanh^2\left(\frac{x}{2}\right) + 2a} + \frac{x}{2a \tanh^4\left(\frac{x}{2}\right) - 4a \tanh^2\left(\frac{x}{2}\right) + 2a} - \frac{2 \tanh^3\left(\frac{x}{2}\right)}{2a \tanh^4\left(\frac{x}{2}\right) - 4a \tanh^2\left(\frac{x}{2}\right) + 2a} - \frac{2 \tanh\left(\frac{x}{2}\right)}{2a \tanh^4\left(\frac{x}{2}\right) - 4a \tanh^2\left(\frac{x}{2}\right) + 2a}$$

[In] integrate(sinh(x)**4/(a-a*cosh(x)**2),x)

```
[Out] x*tanh(x/2)**4/(2*a*tanh(x/2)**4 - 4*a*tanh(x/2)**2 + 2*a) - 2*x*tanh(x/2)*
*2/(2*a*tanh(x/2)**4 - 4*a*tanh(x/2)**2 + 2*a) + x/(2*a*tanh(x/2)**4 - 4*a*
tanh(x/2)**2 + 2*a) - 2*tanh(x/2)**3/(2*a*tanh(x/2)**4 - 4*a*tanh(x/2)**2 +
2*a) - 2*tanh(x/2)/(2*a*tanh(x/2)**4 - 4*a*tanh(x/2)**2 + 2*a)
```

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.25

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = \frac{x}{2a} - \frac{e^{2x}}{8a} + \frac{e^{-2x}}{8a}$$

[In] integrate(sinh(x)^4/(a-a*cosh(x)^2),x, algorithm="maxima")

[Out] 1/2*x/a - 1/8*e^(2*x)/a + 1/8*e^(-2*x)/a

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.30

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = -\frac{(2e^{2x} - 1)e^{-2x} - 4x + e^{2x}}{8a}$$

[In] integrate(sinh(x)^4/(a-a*cosh(x)^2),x, algorithm="giac")

[Out] -1/8*((2*e^(2*x) - 1)*e^(-2*x) - 4*x + e^(2*x))/a

Mupad [B] (verification not implemented)

Time = 1.78 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.25

$$\int \frac{\sinh^4(x)}{a - a \cosh^2(x)} dx = \frac{e^{-2x}}{8a} - \frac{e^{2x}}{8a} + \frac{x}{2a}$$

[In] int(sinh(x)^4/(a - a*cosh(x)^2),x)

[Out] exp(-2*x)/(8*a) - exp(2*x)/(8*a) + x/(2*a)

3.2 $\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx$

Optimal result	55
Rubi [A] (verified)	55
Mathematica [A] (verified)	56
Maple [A] (verified)	56
Fricas [A] (verification not implemented)	56
Sympy [A] (verification not implemented)	57
Maxima [B] (verification not implemented)	57
Giac [A] (verification not implemented)	57
Mupad [B] (verification not implemented)	57

Optimal result

Integrand size = 16, antiderivative size = 7

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x)}{a}$$

[Out] $-\cosh(x)/a$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3254, 2718}

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x)}{a}$$

[In] $\text{Int}[\text{Sinh}[x]^3/(a - a*\text{Cosh}[x]^2), x]$

[Out] $-(\text{Cosh}[x]/a)$

Rule 2718

$\text{Int}[\sin[(c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rule 3254

$\text{Int}[(u_.)*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[a^p, \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /;$ FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\int \sinh(x) dx}{a} \\ &= -\frac{\cosh(x)}{a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x)}{a}$$

[In] Integrate[Sinh[x]^3/(a - a*Cosh[x]^2),x]

[Out] -(Cosh[x]/a)

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 8, normalized size of antiderivative = 1.14

method	result	size
derivativedivides	$-\frac{\cosh(x)}{a}$	8
default	$-\frac{\cosh(x)}{a}$	8
risch	$-\frac{e^x}{2a} - \frac{e^{-x}}{2a}$	18

[In] int(sinh(x)^3/(a-a*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] -cosh(x)/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x)}{a}$$

[In] integrate(sinh(x)^3/(a-a*cosh(x)^2),x, algorithm="fricas")

[Out] -cosh(x)/a

Sympy [A] (verification not implemented)

Time = 0.44 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.43

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = \frac{2}{a \tanh^2\left(\frac{x}{2}\right) - a}$$

[In] integrate(sinh(x)**3/(a-a*cosh(x)**2),x)

[Out] 2/(a*tanh(x/2)**2 - a)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 17 vs. 2(7) = 14.

Time = 0.20 (sec) , antiderivative size = 17, normalized size of antiderivative = 2.43

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{e^{(-x)}}{2a} - \frac{e^x}{2a}$$

[In] integrate(sinh(x)^3/(a-a*cosh(x)^2),x, algorithm="maxima")

[Out] -1/2*e^(-x)/a - 1/2*e^x/a

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.71

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{e^{(-x)} + e^x}{2a}$$

[In] integrate(sinh(x)^3/(a-a*cosh(x)^2),x, algorithm="giac")

[Out] -1/2*(e^(-x) + e^x)/a

Mupad [B] (verification not implemented)

Time = 1.71 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^3(x)}{a - a \cosh^2(x)} dx = -\frac{\cosh(x)}{a}$$

[In] int(sinh(x)^3/(a - a*cosh(x)^2),x)

[Out] -cosh(x)/a

3.3 $\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx$

Optimal result	58
Rubi [A] (verified)	58
Mathematica [A] (verified)	59
Maple [A] (verified)	59
Fricas [A] (verification not implemented)	59
Sympy [A] (verification not implemented)	60
Maxima [A] (verification not implemented)	60
Giac [A] (verification not implemented)	60
Mupad [B] (verification not implemented)	60

Optimal result

Integrand size = 16, antiderivative size = 6

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[Out] $-x/a$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3254, 8}

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] `Int[Sinh[x]^2/(a - a*Cosh[x]^2),x]`

[Out] $-(x/a)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3254

`Int[(u_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Dist[a^p, Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]`

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\int 1 dx}{a} \\ &= -\frac{x}{a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] Integrate[Sinh[x]^2/(a - a*Cosh[x]^2),x]

[Out] -(x/a)

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 7, normalized size of antiderivative = 1.17

method	result	size
risch	$-\frac{x}{a}$	7
default	$-\frac{2 \operatorname{arctanh}(\tanh(\frac{x}{2}))}{a}$	11

[In] int(sinh(x)^2/(a-a*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] -x/a

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] integrate(sinh(x)^2/(a-a*cosh(x)^2),x, algorithm="fricas")

[Out] -x/a

Sympy [A] (verification not implemented)

Time = 0.27 (sec) , antiderivative size = 3, normalized size of antiderivative = 0.50

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] integrate(sinh(x)**2/(a-a*cosh(x)**2),x)

[Out] -x/a

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] integrate(sinh(x)^2/(a-a*cosh(x)^2),x, algorithm="maxima")

[Out] -x/a

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] integrate(sinh(x)^2/(a-a*cosh(x)^2),x, algorithm="giac")

[Out] -x/a

Mupad [B] (verification not implemented)

Time = 0.03 (sec) , antiderivative size = 6, normalized size of antiderivative = 1.00

$$\int \frac{\sinh^2(x)}{a - a \cosh^2(x)} dx = -\frac{x}{a}$$

[In] int(sinh(x)^2/(a - a*cosh(x)^2),x)

[Out] -x/a

3.4 $\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx$

Optimal result	61
Rubi [A] (verified)	61
Mathematica [A] (verified)	62
Maple [A] (verified)	62
Fricas [B] (verification not implemented)	63
Sympy [F]	63
Maxima [B] (verification not implemented)	63
Giac [A] (verification not implemented)	64
Mupad [B] (verification not implemented)	64

Optimal result

Integrand size = 16, antiderivative size = 19

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = -\frac{\operatorname{coth}(x)}{a} + \frac{\operatorname{coth}^3(x)}{3a}$$

[Out] $-\operatorname{coth}(x)/a + 1/3 * \operatorname{coth}(x)^3/a$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3254, 3852}

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = \frac{\operatorname{coth}^3(x)}{3a} - \frac{\operatorname{coth}(x)}{a}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]^2/(a - a * \operatorname{Cosh}[x]^2), x]$

[Out] $-(\operatorname{Coth}[x]/a) + \operatorname{Coth}[x]^3/(3*a)$

Rule 3254

$\operatorname{Int}[(u_.)*((a_) + (b_.)*\sin[(e_.) + (f_.)*(x_)]^2)^{(p_)}, x_Symbol] \rightarrow \operatorname{Dist}[a^p, \operatorname{Int}[\operatorname{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, p\}, x \ \&\& \operatorname{EqQ}[a + b, 0] \ \&\& \operatorname{IntegerQ}[p]$

Rule 3852

$\operatorname{Int}[\operatorname{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \operatorname{Dist}[-d^{(-1)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \operatorname{Cot}[c + d*x]], x] /;$ $\operatorname{FreeQ}\{c,$

d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\int \operatorname{csch}^4(x) dx}{a} \\ &= -\frac{i\operatorname{Subst}\left(\int (1+x^2) dx, x, -i \operatorname{coth}(x)\right)}{a} \\ &= -\frac{\operatorname{coth}(x)}{a} + \frac{\operatorname{coth}^3(x)}{3a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = -\frac{\frac{2 \operatorname{coth}(x)}{3} - \frac{1}{3} \operatorname{coth}(x) \operatorname{csch}^2(x)}{a}$$

[In] Integrate[Csch[x]^2/(a - a*Cosh[x]^2), x]

[Out] -(((2*Coth[x])/3 - (Coth[x]*Csch[x]^2)/3)/a)

Maple [A] (verified)

Time = 2.95 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.16

method	result	size
risch	$\frac{4 e^{2x} - \frac{4}{3}}{(e^{2x} - 1)^{\frac{3}{2}} a}$	22
default	$\frac{\frac{\tanh\left(\frac{x}{2}\right)^3}{3} - 3 \tanh\left(\frac{x}{2}\right) + \frac{1}{3 \tanh\left(\frac{x}{2}\right)^3} - \frac{3}{\tanh\left(\frac{x}{2}\right)}}{8a}$	37

[In] int(csch(x)^2/(a-a*cosh(x)^2), x, method=_RETURNVERBOSE)

[Out] 4/3*(3*exp(2*x)-1)/(exp(2*x)-1)^3/a

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(17) = 34.

Time = 0.24 (sec) , antiderivative size = 100, normalized size of antiderivative = 5.26

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx$$

$$= \frac{8(\cosh(x) + 2 \sinh(x))}{3(a \cosh(x)^5 + 5a \cosh(x) \sinh(x)^4 + a \sinh(x)^5 - 3a \cosh(x)^3 + (10a \cosh(x)^2 - 3a) \sinh(x)^3 + (10a \cosh(x) - 3a) \sinh(x)^2 + 2a \cosh(x) + 5a \sinh(x) - a)}$$

[In] integrate(csch(x)^2/(a-a*cosh(x)^2),x, algorithm="fricas")

[Out] 8/3*(cosh(x) + 2*sinh(x))/(a*cosh(x)^5 + 5*a*cosh(x)*sinh(x)^4 + a*sinh(x)^5 - 3*a*cosh(x)^3 + (10*a*cosh(x)^2 - 3*a)*sinh(x)^3 + (10*a*cosh(x)^2 - 9*a*cosh(x))*sinh(x)^2 + 2*a*cosh(x) + (5*a*cosh(x)^4 - 9*a*cosh(x)^2 + 4*a)*sinh(x))

Sympy [F]

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = -\frac{\int \frac{\operatorname{csch}^2(x)}{\cosh^2(x)-1} dx}{a}$$

[In] integrate(csch(x)**2/(a-a*cosh(x)**2),x)

[Out] -Integral(csch(x)**2/(cosh(x)**2 - 1), x)/a

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(17) = 34.

Time = 0.20 (sec) , antiderivative size = 61, normalized size of antiderivative = 3.21

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = -\frac{4e^{(-2x)}}{3ae^{(-2x)} - 3ae^{(-4x)} + ae^{(-6x)} - a} + \frac{4}{3(3ae^{(-2x)} - 3ae^{(-4x)} + ae^{(-6x)} - a)}$$

[In] integrate(csch(x)^2/(a-a*cosh(x)^2),x, algorithm="maxima")

[Out] -4*e^(-2*x)/(3*a*e^(-2*x) - 3*a*e^(-4*x) + a*e^(-6*x) - a) + 4/3/(3*a*e^(-2*x) - 3*a*e^(-4*x) + a*e^(-6*x) - a)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = \frac{4(3e^{2x} - 1)}{3a(e^{2x} - 1)^3}$$

[In] integrate(csch(x)^2/(a-a*cosh(x)^2),x, algorithm="giac")

[Out] 4/3*(3*e^(2*x) - 1)/(a*(e^(2*x) - 1)^3)

Mupad [B] (verification not implemented)

Time = 1.70 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

$$\int \frac{\operatorname{csch}^2(x)}{a - a \cosh^2(x)} dx = \frac{4(3e^{2x} - 1)}{3a(e^{2x} - 1)^3}$$

[In] int(1/(sinh(x)^2*(a - a*cosh(x)^2)),x)

[Out] (4*(3*exp(2*x) - 1))/(3*a*(exp(2*x) - 1)^3)

3.5 $\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx$

Optimal result	65
Rubi [A] (verified)	65
Mathematica [A] (verified)	66
Maple [A] (verified)	66
Fricas [B] (verification not implemented)	67
Sympy [F]	67
Maxima [B] (verification not implemented)	67
Giac [A] (verification not implemented)	68
Mupad [B] (verification not implemented)	68

Optimal result

Integrand size = 16, antiderivative size = 29

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = \frac{\operatorname{coth}(x)}{a} - \frac{2 \operatorname{coth}^3(x)}{3a} + \frac{\operatorname{coth}^5(x)}{5a}$$

[Out] $\operatorname{coth}(x)/a - 2/3 * \operatorname{coth}(x)^3/a + 1/5 * \operatorname{coth}(x)^5/a$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3254, 3852}

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = \frac{\operatorname{coth}^5(x)}{5a} - \frac{2 \operatorname{coth}^3(x)}{3a} + \frac{\operatorname{coth}(x)}{a}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]^4/(a - a * \operatorname{Cosh}[x]^2), x]$

[Out] $\operatorname{Coth}[x]/a - (2 * \operatorname{Coth}[x]^3)/(3 * a) + \operatorname{Coth}[x]^5/(5 * a)$

Rule 3254

$\operatorname{Int}[(u_.) * ((a_.) + (b_.) * \sin[(e_.) + (f_.) * (x_.)]^2)^{(p_.)}, x_Symbol] := \operatorname{Dist}[a^p, \operatorname{Int}[\operatorname{ActivateTrig}[u * \cos[e + f * x]^{(2 * p)}], x], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, p, x\}$ && $\operatorname{EqQ}[a + b, 0]$ && $\operatorname{IntegerQ}[p]$

Rule 3852

$\operatorname{Int}[\operatorname{csc}[(c_.) + (d_.) * (x_.)]^{(n_.)}, x_Symbol] := \operatorname{Dist}[-d^{(-1)}, \operatorname{Subst}[\operatorname{Int}[\operatorname{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \operatorname{Cot}[c + d * x]], x] /;$ $\operatorname{FreeQ}\{c,$

d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\int \operatorname{csch}^6(x) dx}{a} \\ &= \frac{i\text{Subst}\left(\int (1 + 2x^2 + x^4) dx, x, -i \operatorname{coth}(x)\right)}{a} \\ &= \frac{\operatorname{coth}(x)}{a} - \frac{2 \operatorname{coth}^3(x)}{3a} + \frac{\operatorname{coth}^5(x)}{5a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 32, normalized size of antiderivative = 1.10

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = -\frac{-\frac{8 \operatorname{coth}(x)}{15} + \frac{4}{15} \operatorname{coth}(x) \operatorname{csch}^2(x) - \frac{1}{5} \operatorname{coth}(x) \operatorname{csch}^4(x)}{a}$$

[In] Integrate[Csch[x]^4/(a - a*Cosh[x]^2),x]

[Out] -(((8*Coth[x])/15 + (4*Coth[x]*Csch[x]^2)/15 - (Coth[x]*Csch[x]^4)/5)/a)

Maple [A] (verified)

Time = 14.81 (sec) , antiderivative size = 28, normalized size of antiderivative = 0.97

method	result	size
risch	$\frac{32 e^{4x}}{3} - \frac{16 e^{2x}}{3} + \frac{16}{15}$ $(e^{2x}-1)^5 a$	28
default	$\frac{\tanh\left(\frac{x}{2}\right)^5}{5} - \frac{5 \tanh\left(\frac{x}{2}\right)^3}{3} + 10 \tanh\left(\frac{x}{2}\right) + \frac{1}{5 \tanh\left(\frac{x}{2}\right)^5} + \frac{10}{\tanh\left(\frac{x}{2}\right)} - \frac{5}{3 \tanh\left(\frac{x}{2}\right)^3}$ $32a$	53

[In] int(csch(x)^4/(a-a*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 16/15*(10*exp(4*x)-5*exp(2*x)+1)/(exp(2*x)-1)^5/a

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(25) = 50.

Time = 0.25 (sec) , antiderivative size = 216, normalized size of antiderivative = 7.45

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx$$

$$= \frac{15 (a \cosh(x))^8 + 8 a \cosh(x) \sinh(x)^7 + a \sinh(x)^8 - 5 a \cosh(x)^6 + (28 a \cosh(x)^2 - 5 a) \sinh(x)^6 + 2 ($$

[In] integrate(csch(x)^4/(a-a*cosh(x)^2),x, algorithm="fricas")

[Out] 16/15*(11*cosh(x)^2 + 18*cosh(x)*sinh(x) + 11*sinh(x)^2 - 5)/(a*cosh(x)^8 + 8*a*cosh(x)*sinh(x)^7 + a*sinh(x)^8 - 5*a*cosh(x)^6 + (28*a*cosh(x)^2 - 5*a)*sinh(x)^6 + 2*(28*a*cosh(x)^3 - 15*a*cosh(x))*sinh(x)^5 + 10*a*cosh(x)^4 + 5*(14*a*cosh(x)^4 - 15*a*cosh(x)^2 + 2*a)*sinh(x)^4 + 4*(14*a*cosh(x)^5 - 25*a*cosh(x)^3 + 10*a*cosh(x))*sinh(x)^3 - 11*a*cosh(x)^2 + (28*a*cosh(x)^6 - 75*a*cosh(x)^4 + 60*a*cosh(x)^2 - 11*a)*sinh(x)^2 + 2*(4*a*cosh(x)^7 - 15*a*cosh(x)^5 + 20*a*cosh(x)^3 - 9*a*cosh(x))*sinh(x) + 5*a)

Sympy [F]

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = -\frac{\int \frac{\operatorname{csch}^4(x)}{\cosh^2(x)-1} dx}{a}$$

[In] integrate(csch(x)**4/(a-a*cosh(x)**2),x)

[Out] -Integral(csch(x)**4/(cosh(x)**2 - 1), x)/a

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 135 vs. 2(25) = 50.

Time = 0.20 (sec) , antiderivative size = 135, normalized size of antiderivative = 4.66

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = \frac{16 e^{(-2x)}}{3 (5 a e^{(-2x)} - 10 a e^{(-4x)} + 10 a e^{(-6x)} - 5 a e^{(-8x)} + a e^{(-10x)} - a)}$$

$$- \frac{32 e^{(-4x)}}{3 (5 a e^{(-2x)} - 10 a e^{(-4x)} + 10 a e^{(-6x)} - 5 a e^{(-8x)} + a e^{(-10x)} - a)}$$

$$- \frac{16}{15 (5 a e^{(-2x)} - 10 a e^{(-4x)} + 10 a e^{(-6x)} - 5 a e^{(-8x)} + a e^{(-10x)} - a)}$$

[In] integrate(csch(x)^4/(a-a*cosh(x)^2),x, algorithm="maxima")

[Out] $16/3*e^{(-2*x)}/(5*a*e^{(-2*x)} - 10*a*e^{(-4*x)} + 10*a*e^{(-6*x)} - 5*a*e^{(-8*x)} + a*e^{(-10*x)} - a) - 32/3*e^{(-4*x)}/(5*a*e^{(-2*x)} - 10*a*e^{(-4*x)} + 10*a*e^{(-6*x)} - 5*a*e^{(-8*x)} + a*e^{(-10*x)} - a) - 16/15/(5*a*e^{(-2*x)} - 10*a*e^{(-4*x)} + 10*a*e^{(-6*x)} - 5*a*e^{(-8*x)} + a*e^{(-10*x)} - a)$

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = \frac{16 (10 e^{4x} - 5 e^{2x} + 1)}{15 a (e^{2x} - 1)^5}$$

[In] `integrate(csch(x)^4/(a-a*cosh(x)^2),x, algorithm="giac")`

[Out] $16/15*(10*e^{(4*x)} - 5*e^{(2*x)} + 1)/(a*(e^{(2*x)} - 1)^5)$

Mupad [B] (verification not implemented)

Time = 1.71 (sec) , antiderivative size = 27, normalized size of antiderivative = 0.93

$$\int \frac{\operatorname{csch}^4(x)}{a - a \cosh^2(x)} dx = \frac{16 (10 e^{4x} - 5 e^{2x} + 1)}{15 a (e^{2x} - 1)^5}$$

[In] `int(1/(sinh(x)^4*(a - a*cosh(x)^2)),x)`

[Out] $(16*(10*\exp(4*x) - 5*\exp(2*x) + 1))/(15*a*(\exp(2*x) - 1)^5)$

3.6 $\int \frac{\sinh^7(x)}{a+b \cosh^2(x)} dx$

Optimal result	69
Rubi [A] (verified)	69
Mathematica [C] (verified)	70
Maple [A] (verified)	71
Fricas [B] (verification not implemented)	71
Sympy [F(-1)]	73
Maxima [F]	73
Giac [F]	73
Mupad [B] (verification not implemented)	74

Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\sinh^7(x)}{a+b \cosh^2(x)} dx = -\frac{(a+b)^3 \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} + \frac{(a^2+3ab+3b^2) \cosh(x)}{b^3} - \frac{(a+3b) \cosh^3(x)}{3b^2} + \frac{\cosh^5(x)}{5b}$$

[Out] (a^2+3*a*b+3*b^2)*cosh(x)/b^3-1/3*(a+3*b)*cosh(x)^3/b^2+1/5*cosh(x)^5/b-(a+b)^3*arctan(cosh(x)*b^(1/2)/a^(1/2))/b^(7/2)/a^(1/2)

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3269, 398, 211}

$$\int \frac{\sinh^7(x)}{a+b \cosh^2(x)} dx = \frac{(a^2+3ab+3b^2) \cosh(x)}{b^3} - \frac{(a+b)^3 \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} - \frac{(a+3b) \cosh^3(x)}{3b^2} + \frac{\cosh^5(x)}{5b}$$

[In] Int[Sinh[x]^7/(a + b*Cosh[x]^2),x]

[Out] -(((a + b)^3*ArcTan[(Sqrt[b]*Cosh[x])/Sqrt[a]])/(Sqrt[a]*b^(7/2))) + ((a^2 + 3*a*b + 3*b^2)*Cosh[x])/b^3 - ((a + 3*b)*Cosh[x]^3)/(3*b^2) + Cosh[x]^5/(5*b)

Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]
```

Rule 3269

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol]
:= With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{(1-x^2)^3}{a+bx^2} dx, x, \cosh(x)\right) \\
&= -\text{Subst}\left(\int \left(-\frac{a^2+3ab+3b^2}{b^3} + \frac{(a+3b)x^2}{b^2} - \frac{x^4}{b} + \frac{a^3+3a^2b+3ab^2+b^3}{b^3(a+bx^2)}\right) dx, x, \cosh(x)\right) \\
&= \frac{(a^2+3ab+3b^2)\cosh(x)}{b^3} - \frac{(a+3b)\cosh^3(x)}{3b^2} \\
&\quad + \frac{\cosh^5(x)}{5b} - \frac{(a+b)^3\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cosh(x)\right)}{b^3} \\
&= -\frac{(a+b)^3 \arctan\left(\frac{\sqrt{b}\cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} + \frac{(a^2+3ab+3b^2)\cosh(x)}{b^3} - \frac{(a+3b)\cosh^3(x)}{3b^2} + \frac{\cosh^5(x)}{5b}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.25 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.90

$$\begin{aligned}
\int \frac{\sinh^7(x)}{a+b\cosh^2(x)} dx &= -\frac{(a+b)^3 \arctan\left(\frac{\sqrt{b-i\sqrt{a+b}}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} \\
&\quad - \frac{(a+b)^3 \arctan\left(\frac{\sqrt{b+i\sqrt{a+b}}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{ab}^{7/2}} \\
&\quad + \frac{(8a^2+22ab+19b^2)\cosh(x)}{8b^3} - \frac{(4a+9b)\cosh(3x)}{48b^2} + \frac{\cosh(5x)}{80b}
\end{aligned}$$

[In] Integrate[Sinh[x]^7/(a + b*Cosh[x]^2),x]

[Out] -(((a + b)^3*ArcTan[(Sqrt[b] - I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/(Sqrt[a]*b^(7/2))) - ((a + b)^3*ArcTan[(Sqrt[b] + I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/(Sqrt[a]*b^(7/2)) + ((8*a^2 + 22*a*b + 19*b^2)*Cosh[x])/(8*b^3) - ((4*a + 9*b)*Cosh[3*x])/(48*b^2) + Cosh[5*x]/(80*b)

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 94, normalized size of antiderivative = 1.21

$$\frac{\frac{\cosh(x)^5 b^2}{5} - \frac{ab \cosh(x)^3}{3} - b^2 \cosh(x)^3 + a^2 \cosh(x) + 3ab \cosh(x) + 3b^2 \cosh(x)}{b^3} + \frac{(-a^3 - 3a^2 b - 3ab^2 - b^3) \arctan\left(\frac{\cosh(x)}{a*b}\right)}{b^3 \sqrt{ab}}$$

[In] int(sinh(x)^7/(a+b*cosh(x)^2),x)

[Out] 1/b^3*(1/5*cosh(x)^5*b^2-1/3*a*b*cosh(x)^3-b^2*cosh(x)^3+a^2*cosh(x)+3*a*b*cosh(x)+3*b^2*cosh(x))+(-a^3-3*a^2*b-3*a*b^2-b^3)/b^3/(a*b)^(1/2)*arctan(b*cosh(x)/(a*b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1199 vs. 2(66) = 132.

Time = 0.31 (sec) , antiderivative size = 2346, normalized size of antiderivative = 30.08

$$\int \frac{\sinh^7(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(sinh(x)^7/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/480*(3*a*b^3*cosh(x)^10 + 30*a*b^3*cosh(x)*sinh(x)^9 + 3*a*b^3*sinh(x)^10 - 5*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^8 + 5*(27*a*b^3*cosh(x)^2 - 4*a^2*b^2 - 9*a*b^3)*sinh(x)^8 + 40*(9*a*b^3*cosh(x)^3 - (4*a^2*b^2 + 9*a*b^3)*cosh(x))*sinh(x)^7 + 30*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^6 + 10*(63*a*b^3*cosh(x)^4 + 24*a^3*b + 66*a^2*b^2 + 57*a*b^3 - 14*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^2)*sinh(x)^6 + 4*(189*a*b^3*cosh(x)^5 - 70*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^3 + 45*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x))*sinh(x)^5 + 30*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^4 + 10*(63*a*b^3*cosh(x)^6 - 35*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^4 + 24*a^3*b + 66*a^2*b^2 + 57*a*b^3 + 45*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^2)*sinh(x)^4 + 3*a*b^3 + 40*(9*a*b^3*cosh(x))^7 - 7*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^5 + 15*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^3 + 3*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x))*sinh(x)^3 - 5*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^2 + 5*(27*a*b^3*cosh(x)^8 - 28*(4*a^2*b^2 + 9*a*b^3)*cosh(x)^6 + 90*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^4 - 4*a^2*b^2 - 9*a*b^3 + 36*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*cosh(x)^2)*sinh(x)^2 -

$$\begin{aligned}
& 240*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^5 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^4*\sinh(x) + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^3*\sinh(x)^2 + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^2*\sinh(x)^3 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)*\sinh(x)^4 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\sinh(x)^5)*\sqrt{-a*b}*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a - b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a - b)*\cosh(x))*\sinh(x) + 4*(\cosh(x)^3 + 3*\cosh(x)*\sinh(x)^2 + \sinh(x)^3 + (3*\cosh(x)^2 + 1)*\sinh(x) + \cosh(x))*\sqrt{-a*b} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 10*(3*a*b^3*\cosh(x)^9 - 4*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^7 + 18*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^5 + 12*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^3 - (4*a^2*b^2 + 9*a*b^3)*\cosh(x))*\sinh(x))/(a*b^4*\cosh(x)^5 + 5*a*b^4*\cosh(x)^4*\sinh(x) + 10*a*b^4*\cosh(x)^3*\sinh(x)^2 + 10*a*b^4*\cosh(x)^2*\sinh(x)^3 + 5*a*b^4*\cosh(x)*\sinh(x)^4 + a*b^4*\sinh(x)^5), 1/480*(3*a*b^3*\cosh(x)^10 + 30*a*b^3*\cosh(x)*\sinh(x)^9 + 3*a*b^3*\sinh(x)^10 - 5*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^8 + 5*(27*a*b^3*\cosh(x)^2 - 4*a^2*b^2 - 9*a*b^3)*\sinh(x)^8 + 40*(9*a*b^3*\cosh(x)^3 - (4*a^2*b^2 + 9*a*b^3)*\cosh(x))*\sinh(x)^7 + 30*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^6 + 10*(63*a*b^3*\cosh(x)^4 + 24*a^3*b + 66*a^2*b^2 + 57*a*b^3 - 14*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^2)*\sinh(x)^6 + 4*(189*a*b^3*\cosh(x)^5 - 70*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^3 + 45*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x))*\sinh(x)^5 + 30*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^4 + 10*(63*a*b^3*\cosh(x)^6 - 35*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^4 + 24*a^3*b + 66*a^2*b^2 + 57*a*b^3 + 45*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^2)*\sinh(x)^4 + 3*a*b^3 + 40*(9*a*b^3*\cosh(x)^7 - 7*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^5 + 15*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^3 + 3*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x))*\sinh(x)^3 - 5*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^2 + 5*(27*a*b^3*\cosh(x)^8 - 28*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^6 + 90*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^4 - 4*a^2*b^2 - 9*a*b^3 + 36*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^2)*\sinh(x)^2 - 480*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^5 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^4*\sinh(x) + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^3*\sinh(x)^2 + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^2*\sinh(x)^3 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)*\sinh(x)^4 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\sinh(x)^5)*\sqrt{a*b}*\arctan(1/2*\sqrt{a*b}*(\cosh(x) + \sinh(x))/a) + 480*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^5 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^4*\sinh(x) + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^3*\sinh(x)^2 + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)^2*\sinh(x)^3 + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\cosh(x)*\sinh(x)^4 + (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\sinh(x)^5)*\sqrt{a*b}*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x)^3 + (4*a + b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + b)*\sinh(x))*\sqrt{a*b}/(a*b)) + 10*(3*a*b^3*\cosh(x)^9 - 4*(4*a^2*b^2 + 9*a*b^3)*\cosh(x)^7 + 18*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^5 + 12*(8*a^3*b + 22*a^2*b^2 + 19*a*b^3)*\cosh(x)^3 - (4*a^2*b^2 + 9*a*b^3)*\cosh(x))*\sinh(x))/(a*b^4*\cosh(x)^5 + 5*a*b^4*\cosh(x)^4*\sinh(x) + 10*a*b^4*\cosh(x)^3*\sinh(x)^2 + 10*a*b^4*\cosh(x)^2*\sinh(x)
\end{aligned}$$

)^3 + 5*a*b^4*cosh(x)*sinh(x)^4 + a*b^4*sinh(x)^5]

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^7(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(sinh(x)**7/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sinh^7(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^7}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^7/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/480*(3*b^2*e^(10*x) + 3*b^2 - 5*(4*a*b + 9*b^2)*e^(8*x) + 30*(8*a^2 + 22*a*b + 19*b^2)*e^(6*x) + 30*(8*a^2 + 22*a*b + 19*b^2)*e^(4*x) - 5*(4*a*b + 9*b^2)*e^(2*x))*e^(-5*x)/b^3 - 1/128*integrate(256*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(3*x) - (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^x)/(b^4*e^(4*x) + b^4 + 2*(2*a*b^3 + b^4)*e^(2*x)), x)

Giac [F]

$$\int \frac{\sinh^7(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^7}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^7/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.38 (sec) , antiderivative size = 805, normalized size of antiderivative = 10.32

$$\int \frac{\sinh^7(x)}{a + b \cosh^2(x)} dx = \frac{e^{-5x}}{160b} + \frac{e^{5x}}{160b} + \frac{e^{-x}(8a^2 + 22ab + 19b^2)}{16b^3}$$

$$\left(2 \operatorname{atan} \left(\frac{e^x (a+b)^3 \sqrt{ab^7}}{2ab^3 \sqrt{(a+b)^6}} \right) - 2 \operatorname{atan} \left(\frac{2e^{3x} (a^7 \sqrt{ab^7} + b^7 \sqrt{ab^7} + 7ab^6 \sqrt{ab^7} + 7a^6 b \sqrt{ab^7} + 21a^2 b^5 \sqrt{ab^7} + 35a^3 b^4 \sqrt{ab^7} + 35a^4 b^3 \sqrt{ab^7} + 35a^5 b^2 \sqrt{ab^7} + 35a^6 b \sqrt{ab^7} + 35a^7 \sqrt{ab^7})}{ab^3 \sqrt{(a+b)^6} (4a^4 + 16a^3 b + 24a^2 b^2 + 16ab^3 + 4b^4)} \right) \right)$$

$$- \frac{e^{-3x}(4a + 9b)}{96b^2} - \frac{e^{3x}(4a + 9b)}{96b^2} + \frac{e^x(8a^2 + 22ab + 19b^2)}{16b^3}$$

[In] int(sinh(x)^7/(a + b*cosh(x)^2),x)

[Out] exp(-5*x)/(160*b) + exp(5*x)/(160*b) + (exp(-x)*(22*a*b + 8*a^2 + 19*b^2))/(16*b^3) - ((2*atan((exp(x)*(a + b)^3*(a*b^7)^(1/2))/(2*a*b^3*((a + b)^6)^(1/2))) - 2*atan((2*exp(3*x)*(a^7*(a*b^7)^(1/2) + b^7*(a*b^7)^(1/2) + 7*a*b^6*(a*b^7)^(1/2) + 7*a^6*b*(a*b^7)^(1/2) + 21*a^2*b^5*(a*b^7)^(1/2) + 35*a^3*b^4*(a*b^7)^(1/2) + 35*a^4*b^3*(a*b^7)^(1/2) + 21*a^5*b^2*(a*b^7)^(1/2)))/(a*b^3*((a + b)^6)^(1/2)*(16*a*b^3 + 16*a^3*b + 4*a^4 + 4*b^4 + 24*a^2*b^2)) + (a*b^8*exp(x)*(a*b^7)^(1/2)*((4*(2*a*b^7*(6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2))^(1/2) + 8*a^2*b^6*(6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2))^(1/2) + 12*a^3*b^5*(6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2))^(1/2) + 8*a^4*b^4*(6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2))^(1/2) + 2*a^5*b^3*(6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2))^(1/2)))/(a^2*b^15*(a + b)^3) + (2*(a^7*(a*b^7)^(1/2) + b^7*(a*b^7)^(1/2) + 7*a*b^6*(a*b^7)^(1/2) + 7*a^6*b*(a*b^7)^(1/2) + 21*a^2*b^5*(a*b^7)^(1/2) + 35*a^3*b^4*(a*b^7)^(1/2) + 35*a^4*b^3*(a*b^7)^(1/2) + 21*a^5*b^2*(a*b^7)^(1/2)))/(a^2*b^11*(a*b^7)^(1/2)*((a + b)^6)^(1/2)))/(16*a*b^3 + 16*a^3*b + 4*a^4 + 4*b^4 + 24*a^2*b^2))*((6*a*b^5 + 6*a^5*b + a^6 + b^6 + 15*a^2*b^4 + 20*a^3*b^3 + 15*a^4*b^2)^(1/2))/(2*(a*b^7)^(1/2)) - (exp(-3*x)*(4*a + 9*b))/(96*b^2) - (exp(3*x)*(4*a + 9*b))/(96*b^2) + (exp(x)*(22*a*b + 8*a^2 + 19*b^2))/(16*b^3)

3.7 $\int \frac{\sinh^5(x)}{a+b \cosh^2(x)} dx$

Optimal result	75
Rubi [A] (verified)	75
Mathematica [C] (verified)	76
Maple [A] (verified)	77
Fricas [B] (verification not implemented)	77
Sympy [F(-1)]	78
Maxima [F]	78
Giac [F]	78
Mupad [B] (verification not implemented)	79

Optimal result

Integrand size = 15, antiderivative size = 54

$$\int \frac{\sinh^5(x)}{a+b \cosh^2(x)} dx = \frac{(a+b)^2 \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{5/2}} - \frac{(a+2b) \cosh(x)}{b^2} + \frac{\cosh^3(x)}{3b}$$

[Out] $-(a+2*b)*\cosh(x)/b^2+1/3*\cosh(x)^3/b+(a+b)^2*\arctan(\cosh(x)*b^{(1/2)}/a^{(1/2)})/b^{(5/2)}/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3269, 398, 211}

$$\int \frac{\sinh^5(x)}{a+b \cosh^2(x)} dx = \frac{(a+b)^2 \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{5/2}} - \frac{(a+2b) \cosh(x)}{b^2} + \frac{\cosh^3(x)}{3b}$$

[In] $\text{Int}[\text{Sinh}[x]^5/(a + b*\text{Cosh}[x]^2), x]$

[Out] $((a + b)^2*\text{ArcTan}[(\text{Sqrt}[b]*\text{Cosh}[x])/(\text{Sqrt}[a])]/(\text{Sqrt}[a]*b^{(5/2)}) - ((a + 2*b)*\text{Cosh}[x])/b^2 + \text{Cosh}[x]^3/(3*b))$

Rule 211

$\text{Int}[(a_0 + (b_0)*(x_0)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b]$

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3269

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_.), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Su
bst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/
ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{(1-x^2)^2}{a+bx^2} dx, x, \cosh(x) \right) \\
&= \text{Subst} \left(\int \left(-\frac{a+2b}{b^2} + \frac{x^2}{b} + \frac{a^2+2ab+b^2}{b^2(a+bx^2)} \right) dx, x, \cosh(x) \right) \\
&= -\frac{(a+2b)\cosh(x)}{b^2} + \frac{\cosh^3(x)}{3b} + \frac{(a+b)^2 \text{Subst} \left(\int \frac{1}{a+bx^2} dx, x, \cosh(x) \right)}{b^2} \\
&= \frac{(a+b)^2 \arctan \left(\frac{\sqrt{b}\cosh(x)}{\sqrt{a}} \right)}{\sqrt{ab^{5/2}}} - \frac{(a+2b)\cosh(x)}{b^2} + \frac{\cosh^3(x)}{3b}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.18 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.22

$$\begin{aligned}
&\int \frac{\sinh^5(x)}{a+b\cosh^2(x)} dx \\
&= \frac{12(a+b)^2 \arctan \left(\frac{\sqrt{b-i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}} \right)}{\sqrt{a}} + \frac{12(a+b)^2 \arctan \left(\frac{\sqrt{b+i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}} \right)}{\sqrt{a}} - 3\sqrt{b}(4a+7b)\cosh(x) + b^{3/2}\cosh(3x) \\
&= \frac{\hspace{15em}}{12b^{5/2}}
\end{aligned}$$

```
[In] Integrate[Sinh[x]^5/(a + b*Cosh[x]^2), x]
```

```
[Out] ((12*(a + b)^2*ArcTan[(Sqrt[b] - I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/Sqrt[a]
+ (12*(a + b)^2*ArcTan[(Sqrt[b] + I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/Sqrt[
a] - 3*Sqrt[b]*(4*a + 7*b)*Cosh[x] + b^(3/2)*Cosh[3*x])/(12*b^(5/2))
```

Maple [A] (verified)

Time = 112.59 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.00

method	result
derivativedivides	$-\frac{-\frac{b \cosh(x)^3}{3} + a \cosh(x) + 2b \cosh(x)}{b^2} + \frac{(a^2 + 2ab + b^2) \arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{b^2 \sqrt{ab}}$
default	$-\frac{-\frac{b \cosh(x)^3}{3} + a \cosh(x) + 2b \cosh(x)}{b^2} + \frac{(a^2 + 2ab + b^2) \arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{b^2 \sqrt{ab}}$
risch	$\frac{e^{3x}}{24b} - \frac{a e^x}{2b^2} - \frac{7e^x}{8b} - \frac{a e^{-x}}{2b^2} - \frac{7e^{-x}}{8b} + \frac{e^{-3x}}{24b} - \frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right) a^2}{2\sqrt{-ab} b^2} - \frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right) a}{\sqrt{-ab} b} - \frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right)}{2\sqrt{-ab}}$

[In] `int(sinh(x)^5/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/b^2*(-1/3*b*cosh(x)^3+a*cosh(x)+2*b*cosh(x))+(a^2+2*a*b+b^2)/b^2/(a*b)^(1/2)*arctan(b*cosh(x)/(a*b)^(1/2))$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 513 vs. 2(44) = 88.

Time = 0.29 (sec) , antiderivative size = 1064, normalized size of antiderivative = 19.70

$$\int \frac{\sinh^5(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] `integrate(sinh(x)^5/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/24*(a*b^2*cosh(x)^6 + 6*a*b^2*cosh(x)*sinh(x)^5 + a*b^2*sinh(x)^6 - 3*(4 \\ & *a^2*b + 7*a*b^2)*cosh(x)^4 + 3*(5*a*b^2*cosh(x)^2 - 4*a^2*b - 7*a*b^2)*sin \\ & h(x)^4 + 4*(5*a*b^2*cosh(x)^3 - 3*(4*a^2*b + 7*a*b^2)*cosh(x))*sinh(x)^3 + \\ & a*b^2 - 3*(4*a^2*b + 7*a*b^2)*cosh(x)^2 + 3*(5*a*b^2*cosh(x)^4 - 4*a^2*b - \\ & 7*a*b^2 - 6*(4*a^2*b + 7*a*b^2)*cosh(x)^2)*sinh(x)^2 - 12*((a^2 + 2*a*b + b \\ & ^2)*cosh(x)^3 + 3*(a^2 + 2*a*b + b^2)*cosh(x)^2*sinh(x) + 3*(a^2 + 2*a*b + \\ & b^2)*cosh(x)*sinh(x)^2 + (a^2 + 2*a*b + b^2)*sinh(x)^3)*sqrt(-a*b)*log((b*c \\ & osh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2* \\ & (3*b*cosh(x)^2 - 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a - b)*cosh(x))*s \\ & inh(x) - 4*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + sinh(x)^3 + (3*cosh(x)^2 + 1) \\ & *sinh(x) + cosh(x))*sqrt(-a*b) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + \\ & b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 \\ & + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b) + 6*(a*b^2*cosh(x)^5 - \\ & 2*(4*a^2*b + 7*a*b^2)*cosh(x)^3 - (4*a^2*b + 7*a*b^2)*cosh(x))*sinh(x))/(a \\ & *b^3*cosh(x)^3 + 3*a*b^3*cosh(x)^2*sinh(x) + 3*a*b^3*cosh(x)*sinh(x)^2 + a* \\ & b^3*sinh(x)^3), 1/24*(a*b^2*cosh(x)^6 + 6*a*b^2*cosh(x)*sinh(x)^5 + a*b^2*s \\ & inh(x)^6 - 3*(4*a^2*b + 7*a*b^2)*cosh(x)^4 + 3*(5*a*b^2*cosh(x)^2 - 4*a^2*b \\ & - 7*a*b^2)*sinh(x)^4 + 4*(5*a*b^2*cosh(x)^3 - 3*(4*a^2*b + 7*a*b^2)*cosh(x) \end{aligned}$$

) * sinh(x)^3 + a*b^2 - 3*(4*a^2*b + 7*a*b^2)*cosh(x)^2 + 3*(5*a*b^2*cosh(x)^4 - 4*a^2*b - 7*a*b^2 - 6*(4*a^2*b + 7*a*b^2)*cosh(x)^2)*sinh(x)^2 + 24*((a^2 + 2*a*b + b^2)*cosh(x)^3 + 3*(a^2 + 2*a*b + b^2)*cosh(x)^2*sinh(x) + 3*(a^2 + 2*a*b + b^2)*cosh(x)*sinh(x)^2 + (a^2 + 2*a*b + b^2)*sinh(x)^3)*sqrt(a*b)*arctan(1/2*sqrt(a*b)*(cosh(x) + sinh(x))/a) - 24*((a^2 + 2*a*b + b^2)*cosh(x)^3 + 3*(a^2 + 2*a*b + b^2)*cosh(x)^2*sinh(x) + 3*(a^2 + 2*a*b + b^2)*cosh(x)*sinh(x)^2 + (a^2 + 2*a*b + b^2)*sinh(x)^3)*sqrt(a*b)*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + b)*cosh(x) + (3*b*cosh(x)^2 + 4*a + b)*sinh(x))*sqrt(a*b)/(a*b)) + 6*(a*b^2*cosh(x)^5 - 2*(4*a^2*b + 7*a*b^2)*cosh(x)^3 - (4*a^2*b + 7*a*b^2)*cosh(x))*sinh(x))/(a*b^3*cosh(x)^3 + 3*a*b^3*cosh(x)^2*sinh(x) + 3*a*b^3*cosh(x)*sinh(x)^2 + a*b^3*sinh(x)^3)]

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^5(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(sinh(x)**5/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sinh^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^5/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/24*(b*e^(6*x) - 3*(4*a + 7*b)*e^(4*x) - 3*(4*a + 7*b)*e^(2*x) + b)*e^(-3*x)/b^2 + 1/32*integrate(64*((a^2 + 2*a*b + b^2)*e^(3*x) - (a^2 + 2*a*b + b^2)*e^x)/(b^3*e^(4*x) + b^3 + 2*(2*a*b^2 + b^3)*e^(2*x)), x)

Giac [F]

$$\int \frac{\sinh^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^5/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.12 (sec) , antiderivative size = 548, normalized size of antiderivative = 10.15

$$\int \frac{\sinh^5(x)}{a + b \cosh^2(x)} dx = \frac{e^{-3x}}{24b} + \frac{e^{3x}}{24b} - \frac{e^{-x}(4a + 7b)}{8b^2} + \frac{\left(2 \operatorname{atan}\left(\frac{e^x(a+b)^2 \sqrt{ab^5}}{2ab^2 \sqrt{(a+b)^4}}\right) - 2 \operatorname{atan}\left(\frac{ab^6 e^x \left(4(6a^2 b^4 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}+6a^3 b^3 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}+2a^4 b^2 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}\right)}{a^2 b^{11} (a+b)^2}\right)}{e^x(4a+7b)}\right)}{8b^2}$$

[In] int(sinh(x)^5/(a + b*cosh(x)^2), x)

[Out] $\frac{\exp(-3x)}{24b} + \frac{\exp(3x)}{24b} - \frac{(\exp(-x)(4a + 7b))}{(8b^2)} + \left(\frac{2 \operatorname{atan}\left(\frac{\exp(x)(a+b)^2(a^2 b^5)^{1/2}}{2ab^2(a+b)^4(a+b)^{1/2}}\right) - 2 \operatorname{atan}\left(\frac{ab^6 \exp(x) \left(4(6a^2 b^4 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}+6a^3 b^3 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}+2a^4 b^2 \sqrt{a^4+4a^3 b+6a^2 b^2+4ab^3+b^4}\right)}{a^2 b^{11} (a+b)^2}\right)}{e^x(4a+7b)}\right)}{8b^2}$

3.8 $\int \frac{\sinh^3(x)}{a+b \cosh^2(x)} dx$

Optimal result	80
Rubi [A] (verified)	80
Mathematica [C] (verified)	81
Maple [A] (verified)	82
Fricas [B] (verification not implemented)	82
Sympy [F(-1)]	83
Maxima [F]	83
Giac [F]	83
Mupad [B] (verification not implemented)	83

Optimal result

Integrand size = 15, antiderivative size = 36

$$\int \frac{\sinh^3(x)}{a+b \cosh^2(x)} dx = -\frac{(a+b) \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{3/2}} + \frac{\cosh(x)}{b}$$

[Out] $\cosh(x)/b - (a+b) \cdot \arctan(\cosh(x) \cdot b^{1/2}/a^{1/2})/b^{3/2}/a^{1/2}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3269, 396, 211}

$$\int \frac{\sinh^3(x)}{a+b \cosh^2(x)} dx = \frac{\cosh(x)}{b} - \frac{(a+b) \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab}^{3/2}}$$

[In] $\text{Int}[\text{Sinh}[x]^3/(a + b \cdot \text{Cosh}[x]^2), x]$

[Out] $-\left(\frac{(a+b) \cdot \text{ArcTan}\left[\frac{\text{Sqrt}[b] \cdot \text{Cosh}[x]}{\text{Sqrt}[a]}\right]}{\text{Sqrt}[a] \cdot b^{3/2}}\right) + \text{Cosh}[x]/b$

Rule 211

$\text{Int}[\left((a_) + (b_) \cdot (x_)^2\right)^{-1}, x_Symbol] \rightarrow \text{Simp}\left[\frac{\text{Rt}[a/b, 2]}{a} \cdot \text{ArcTan}[x/\text{Rt}[a/b, 2]], x\right] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$

Rule 396

$\text{Int}[\left((a_) + (b_) \cdot (x_)^{(n_)}\right)^{(p_)} \cdot \left((c_) + (d_) \cdot (x_)^{(n_)}\right), x_Symbol] \rightarrow \text{Simp}[d \cdot x \cdot \left((a + b \cdot x^n)^{(p+1)} / (b \cdot (n \cdot (p+1) + 1)\right)], x] - \text{Dist}[(a \cdot d - b \cdot c \cdot (n \cdot (p+1) + 1)) / (b \cdot (n \cdot (p+1) + 1)), \text{Int}[(a + b \cdot x^n)^{(p+1)}], x]$

$p + 1) + 1)) / (b * (n * (p + 1) + 1))$, Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 3269

Int[cos[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_.), x_Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1-x^2}{a+bx^2} dx, x, \cosh(x)\right) \\ &= \frac{\cosh(x)}{b} - \frac{(a+b)\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cosh(x)\right)}{b} \\ &= -\frac{(a+b)\arctan\left(\frac{\sqrt{b}\cosh(x)}{\sqrt{a}}\right)}{\sqrt{ab^{3/2}}} + \frac{\cosh(x)}{b} \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.19 (sec) , antiderivative size = 83, normalized size of antiderivative = 2.31

$$\int \frac{\sinh^3(x)}{a+b\cosh^2(x)} dx = -\frac{(a+b)\left(\arctan\left(\frac{\sqrt{b-i\sqrt{a+b}}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right) + \arctan\left(\frac{\sqrt{b+i\sqrt{a+b}}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)\right)}{\sqrt{ab^{3/2}}} + \frac{\cosh(x)}{b}$$

[In] Integrate[Sinh[x]^3/(a + b*Cosh[x]^2),x]

[Out] -(((a + b)*(ArcTan[(Sqrt[b] - I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]] + ArcTan[(Sqrt[b] + I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]]))/(Sqrt[a]*b^(3/2))) + Cosh[x]/b

Maple [A] (verified)

Time = 4.46 (sec) , antiderivative size = 34, normalized size of antiderivative = 0.94

method	result	size
derivativedivides	$\frac{\cosh(x)}{b} + \frac{(-a-b) \arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	34
default	$\frac{\cosh(x)}{b} + \frac{(-a-b) \arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{b\sqrt{ab}}$	34
risch	$\frac{e^x}{2b} + \frac{e^{-x}}{2b} - \frac{\ln\left(e^{2x} + \frac{2a e^x}{\sqrt{-ab}} + 1\right)a}{2\sqrt{-ab}b} - \frac{\ln\left(e^{2x} + \frac{2a e^x}{\sqrt{-ab}} + 1\right)}{2\sqrt{-ab}} + \frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right)a}{2\sqrt{-ab}b} + \frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right)}{2\sqrt{-ab}}$	130

[In] int(sinh(x)^3/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] cosh(x)/b+(-a-b)/b/(a*b)^(1/2)*arctan(b*cosh(x)/(a*b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(28) = 56.

Time = 0.28 (sec) , antiderivative size = 416, normalized size of antiderivative = 11.56

$$\int \frac{\sinh^3(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{ab \cosh(x)^2 + 2ab \cosh(x) \sinh(x) + ab \sinh(x)^2 - \sqrt{-ab}((a+b) \cosh(x) + (a+b) \sinh(x)) \log\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{\dots}$$

[In] integrate(sinh(x)^3/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/2*(a*b*cosh(x)^2 + 2*a*b*cosh(x)*sinh(x) + a*b*sinh(x)^2 - sqrt(-a*b)*((a + b)*cosh(x) + (a + b)*sinh(x))*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a - b)*cosh(x))*sinh(x) + 4*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + sinh(x)^3 + (3*cosh(x)^2 + 1)*sinh(x) + cosh(x))*sqrt(-a*b) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x)*sinh(x) + b)) + a*b)/(a*b^2*cosh(x) + a*b^2*sinh(x)), 1/2*(a*b*cosh(x)^2 + 2*a*b*cosh(x)*sinh(x) + a*b*sinh(x)^2 - 2*sqrt(a*b)*((a + b)*cosh(x) + (a + b)*sinh(x))*arctan(1/2*sqrt(a*b)*(cosh(x) + sinh(x))/a) + 2*sqrt(a*b)*((a + b)*cosh(x) + (a + b)*sinh(x))*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + b)*cosh(x) + (3*b*cosh(x)^2 + 4*a + b)*sinh(x))*sqrt(a*b)/(a*b)) + a*b)/(a*b^2*cosh(x) + a*b^2*sinh(x))]

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^3(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(sinh(x)**3/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\sinh^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^3}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^3/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/2*(e^(2*x) + 1)*e^(-x)/b - 1/8*integrate(16*((a + b)*e^(3*x) - (a + b)*e^x)/(b^2*e^(4*x) + b^2 + 2*(2*a*b + b^2)*e^(2*x)), x)

Giac [F]

$$\int \frac{\sinh^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)^3}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)^3/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.19 (sec) , antiderivative size = 257, normalized size of antiderivative = 7.14

$$\int \frac{\sinh^3(x)}{a + b \cosh^2(x)} dx = \frac{e^{-x}}{2b} + \frac{e^x}{2b} + \frac{2 \operatorname{atan} \left(\frac{e^{3x} (a^3 \sqrt{ab^3} + b^3 \sqrt{ab^3} + 3ab^2 \sqrt{ab^3} + 3a^2b \sqrt{ab^3})}{2ab((a+b)^2)} \right) + \frac{ab^4 e^x \sqrt{ab^3} \left(\frac{8(a^2 + 2ab + b^2)^{3/2}}{ab^6(a+b)^3} + \frac{2(a^3 \sqrt{ab^3} + b^3 \sqrt{ab^3} + 3ab^2 \sqrt{ab^3} + 3a^2b \sqrt{ab^3})}{a^2 b^5 \sqrt{ab^3} ((a+b)^2)^{3/2}} \right)}{4}}{2\sqrt{ab^3}}$$

[In] int(sinh(x)^3/(a + b*cosh(x)^2),x)

```
[Out] exp(-x)/(2*b) + exp(x)/(2*b) + ((2*atan((exp(3*x)*(a^3*(a*b^3)^(1/2) + b^3*(a*b^3)^(1/2) + 3*a*b^2*(a*b^3)^(1/2) + 3*a^2*b*(a*b^3)^(1/2)))/(2*a*b*((a + b)^2)^(3/2)) + (a*b^4*exp(x)*(a*b^3)^(1/2)*((8*(2*a*b + a^2 + b^2)^(3/2))/(a*b^6*(a + b)^3) + (2*(a^3*(a*b^3)^(1/2) + b^3*(a*b^3)^(1/2) + 3*a*b^2*(a*b^3)^(1/2) + 3*a^2*b*(a*b^3)^(1/2)))/(a^2*b^5*(a*b^3)^(1/2)*((a + b)^2)^(3/2))))/4 - 2*atan((exp(x)*(a + b)^3*(a*b^3)^(1/2))/(2*a*b*((a + b)^2)^(3/2))))*(2*a*b + a^2 + b^2)^(1/2))/(2*(a*b^3)^(1/2))
```

3.9 $\int \frac{\sinh(x)}{a+b \cosh^2(x)} dx$

Optimal result	85
Rubi [A] (verified)	85
Mathematica [A] (verified)	86
Maple [A] (verified)	86
Fricas [B] (verification not implemented)	87
Sympy [B] (verification not implemented)	87
Maxima [F]	88
Giac [F]	88
Mupad [B] (verification not implemented)	88

Optimal result

Integrand size = 13, antiderivative size = 25

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

[Out] $\arctan(\cosh(x)*b^{(1/2)}/a^{(1/2)})/a^{(1/2)}/b^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3269, 211}

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}\sqrt{b}}$$

[In] $\text{Int}[\text{Sinh}[x]/(a + b*\text{Cosh}[x]^2), x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[b]*\text{Cosh}[x])/(\text{Sqrt}[a])]/(\text{Sqrt}[a]*\text{Sqrt}[b])$

Rule 211

$\text{Int}[(a_1 + (b_1)*(x_1)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b]$

Rule 3269

$\text{Int}[\cos[(e_1) + (f_1)*(x_1)]^{(m_1)}*((a_1) + (b_1)*\sin[(e_1) + (f_1)*(x_1)]^2)^{(p_1)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Dist}[ff/f, \text{Subst}[\text{Int}[(1 - ff^2*x^2)^{(m-1)/2}*(a + b*ff^2*x^2)^p, x], x, \text{Sin}[e + f*x]/$

```
ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{1}{a + bx^2} dx, x, \cosh(x) \right) \\ &= \frac{\arctan \left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}} \right)}{\sqrt{a}\sqrt{b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 25, normalized size of antiderivative = 1.00

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \frac{\arctan \left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}} \right)}{\sqrt{a}\sqrt{b}}$$

```
[In] Integrate[Sinh[x]/(a + b*Cosh[x]^2),x]
```

```
[Out] ArcTan[(Sqrt[b]*Cosh[x])/Sqrt[a]]/(Sqrt[a]*Sqrt[b])
```

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.68

method	result	size
derivativedivides	$\frac{\arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$	17
default	$\frac{\arctan\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$	17
risch	$-\frac{\ln\left(e^{2x} - \frac{2a e^x}{\sqrt{-ab}} + 1\right)}{2\sqrt{-ab}} + \frac{\ln\left(e^{2x} + \frac{2a e^x}{\sqrt{-ab}} + 1\right)}{2\sqrt{-ab}}$	54

```
[In] int(sinh(x)/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/(a*b)^(1/2)*arctan(b*cosh(x)/(a*b)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 96 vs. $2(17) = 34$.

Time = 0.27 (sec) , antiderivative size = 300, normalized size of antiderivative = 12.00

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \left[\frac{\sqrt{-ab} \log \left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 - 2(2a-b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a+b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a-b) \cosh(x) \sinh(x) + b \sinh(x)^2)}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a+b) \sinh(x)^2} \right)}{2ab} \right]$$

[In] integrate(sinh(x)/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] $[-1/2*\sqrt{-a*b}*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a - b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a - b)*\cosh(x))*\sinh(x) - 4*(\cosh(x)^3 + 3*\cosh(x)*\sinh(x)^2 + \sinh(x)^3 + (3*\cosh(x)^2 + 1)*\sinh(x) + \cosh(x))*\sqrt{-a*b} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b))/a*b, (\sqrt{a*b}*\arctan(1/2*\sqrt{a*b}*(\cosh(x) + \sinh(x))/a) - \sqrt{a*b})*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x)^3 + (4*a + b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + b)*\sinh(x))*\sqrt{a*b}/(a*b)))/a*b]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(24) = 48$.

Time = 0.35 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.64

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \begin{cases} \frac{\infty}{\cosh(x)} & \text{for } a = 0 \wedge b = 0 \\ \frac{\cosh(x)}{a} & \text{for } b = 0 \\ -\frac{1}{b \cosh(x)} & \text{for } a = 0 \\ \frac{\log\left(-\sqrt{-\frac{a}{b}} + \cosh(x)\right)}{2b\sqrt{-\frac{a}{b}}} - \frac{\log\left(\sqrt{-\frac{a}{b}} + \cosh(x)\right)}{2b\sqrt{-\frac{a}{b}}} & \text{otherwise} \end{cases}$$

[In] integrate(sinh(x)/(a+b*cosh(x)**2),x)

[Out] Piecewise((zoo/cosh(x), Eq(a, 0) & Eq(b, 0)), (cosh(x)/a, Eq(b, 0)), (-1/(b*cosh(x)), Eq(a, 0)), (log(-sqrt(-a/b) + cosh(x))/(2*b*sqrt(-a/b)) - log(sqrt(-a/b) + cosh(x))/(2*b*sqrt(-a/b)), True))

Maxima [F]

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] integrate(sinh(x)/(b*cosh(x)^2 + a), x)

Giac [F]

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \int \frac{\sinh(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(sinh(x)/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 1.78 (sec) , antiderivative size = 16, normalized size of antiderivative = 0.64

$$\int \frac{\sinh(x)}{a + b \cosh^2(x)} dx = \frac{\operatorname{atan}\left(\frac{b \cosh(x)}{\sqrt{ab}}\right)}{\sqrt{ab}}$$

[In] int(sinh(x)/(a + b*cosh(x)^2),x)

[Out] atan((b*cosh(x))/(a*b)^(1/2))/(a*b)^(1/2)

3.10 $\int \frac{\operatorname{csch}(x)}{a+b \cosh^2(x)} dx$

Optimal result	89
Rubi [A] (verified)	89
Mathematica [C] (verified)	90
Maple [A] (verified)	91
Fricas [B] (verification not implemented)	91
Sympy [F]	92
Maxima [F]	92
Giac [F]	92
Mupad [B] (verification not implemented)	92

Optimal result

Integrand size = 13, antiderivative size = 42

$$\int \frac{\operatorname{csch}(x)}{a+b \cosh^2(x)} dx = -\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} - \frac{\operatorname{arctanh}(\cosh(x))}{a+b}$$

[Out] $-\operatorname{arctanh}(\cosh(x))/(a+b)-\arctan(\cosh(x)*b^{(1/2)}/a^{(1/2)})*b^{(1/2)}/(a+b)/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3269, 400, 212, 211}

$$\int \frac{\operatorname{csch}(x)}{a+b \cosh^2(x)} dx = -\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} - \frac{\operatorname{arctanh}(\cosh(x))}{a+b}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]/(a+b*\operatorname{Cosh}[x]^2),x]$

[Out] $-\left(\frac{\sqrt{b}*\operatorname{ArcTan}[\sqrt{b}*\operatorname{Cosh}[x]]/\sqrt{a}}{\sqrt{a}*(a+b)}\right)-\operatorname{ArcTanh}[\operatorname{Cosh}[x]]/(a+b)$

Rule 211

$\operatorname{Int}[\left((a_+)+(b_-)*(x_-)^2\right)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[\left(\operatorname{Rt}[a/b, 2]/a\right)*\operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ \operatorname{PosQ}[a/b]$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 400

```
Int[1/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dis
t[b/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[d/(b*c - a*d), Int[1/(c +
d*x^n), x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0]
```

Rule 3269

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^
(p_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Su
bst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/
ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1}{(1-x^2)(a+bx^2)} dx, x, \cosh(x)\right) \\ &= -\frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cosh(x)\right)}{a+b} - \frac{b\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cosh(x)\right)}{a+b} \\ &= -\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)} - \frac{\text{arctanh}(\cosh(x))}{a+b} \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.16 (sec) , antiderivative size = 106, normalized size of antiderivative = 2.52

$$\begin{aligned} &\int \frac{\text{csch}(x)}{a+b \cosh^2(x)} dx \\ &= -\frac{\frac{\sqrt{b} \arctan\left(\frac{\sqrt{b-i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{a}} + \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b+i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)}{\sqrt{a}} + \log\left(\cosh\left(\frac{x}{2}\right)\right) - \log\left(\sinh\left(\frac{x}{2}\right)\right)}{a+b} \end{aligned}$$

```
[In] Integrate[Csch[x]/(a + b*Cosh[x]^2), x]
```

```
[Out] -(((Sqrt[b]*ArcTan[(Sqrt[b] - I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/Sqrt[a] +
(Sqrt[b]*ArcTan[(Sqrt[b] + I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]])/Sqrt[a] + Log
[Cosh[x/2]] - Log[Sinh[x/2]])/(a + b))
```

Maple [A] (verified)

Time = 1.91 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.24

method	result	size
default	$\frac{\ln(\tanh(\frac{x}{2}))}{a+b} - \frac{b \arctan\left(\frac{2(a+b) \tanh(\frac{x}{2})^2 - 2a + 2b}{4\sqrt{ab}}\right)}{(a+b)\sqrt{ab}}$	52
risch	$\frac{\ln(e^x - 1)}{a+b} - \frac{\ln(e^x + 1)}{a+b} + \frac{\sqrt{-ab} \ln\left(e^{2x} - \frac{2\sqrt{-ab} e^x}{b} + 1\right)}{2a(a+b)} - \frac{\sqrt{-ab} \ln\left(e^{2x} + \frac{2\sqrt{-ab} e^x}{b} + 1\right)}{2a(a+b)}$	97

[In] `int(csch(x)/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] `1/(a+b)*ln(tanh(1/2*x))-b/(a+b)/(a*b)^(1/2)*arctan(1/4*(2*(a+b)*tanh(1/2*x)^2-2*a+2*b)/(a*b)^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(34) = 68$.

Time = 0.28 (sec) , antiderivative size = 349, normalized size of antiderivative = 8.31

$$\int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{\sqrt{-\frac{b}{a}} \log\left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 - 2(2a-b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a + b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a-b) \cosh(x) \sinh(x) + b) \sinh(x)}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a + b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a-b) \cosh(x) \sinh(x) + b) \sinh(x)}\right)}{a + b}$$

$$- \frac{\sqrt{\frac{b}{a}} \arctan\left(\frac{1}{2} \sqrt{\frac{b}{a}} (\cosh(x) + \sinh(x))\right) - \sqrt{\frac{b}{a}} \arctan\left(\frac{(b \cosh(x)^3 + 3b \cosh(x) \sinh(x)^2 + b \sinh(x)^3 + (4a+b) \cosh(x) \sinh(x) + b)}{2b}\right)}{a + b}$$

[In] `integrate(csch(x)/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] `[1/2*(sqrt(-b/a)*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a - b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a - b)*cosh(x))*sinh(x) - 4*(a*cosh(x)^3 + 3*a*cosh(x)*sinh(x)^2 + a*sinh(x)^3 + a*cosh(x) + (3*a*cosh(x)^2 + a)*sinh(x))*sqrt(-b/a) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) - 2*log(cosh(x) + sinh(x) + 1) + 2*log(cosh(x) + sinh(x) - 1))/(a + b), -(sqrt(b/a)*arctan(1/2*sqrt(b/a)*(cosh(x) + sinh(x)))) - sqrt(b/a)*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + b)*`

$\cosh(x) + (3*b*\cosh(x)^2 + 4*a + b)*\sinh(x))*\sqrt{b/a)/b) + \log(\cosh(x) + \sinh(x) + 1) - \log(\cosh(x) + \sinh(x) - 1))/(a + b)]$

Sympy [F]

$$\int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx$$

[In] integrate(csch(x)/(a+b*cosh(x)**2),x)

[Out] Integral(csch(x)/(a + b*cosh(x)**2), x)

Maxima [F]

$$\int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-\log(e^x + 1)/(a + b) + \log(e^x - 1)/(a + b) - 2*\integrate((b*e^{(3*x)} - b*e^{-x})/(a*b + b^2 + (a*b + b^2)*e^{(4*x)} + 2*(2*a^2 + 3*a*b + b^2)*e^{(2*x)}), x)$

Giac [F]

$$\int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.21 (sec) , antiderivative size = 462, normalized size of antiderivative = 11.00

$$\int \frac{\operatorname{csch}(x)}{a + b \cosh^2(x)} dx = -\frac{2 \operatorname{atan}\left(\frac{e^x (16 a^2 \sqrt{-a^2 - 2 a b - b^2} + b^2 \sqrt{-a^2 - 2 a b - b^2} + 8 a b \sqrt{-a^2 - 2 a b - b^2})}{16 a^3 + 24 a^2 b + 9 a b^2 + b^3}\right)}{\sqrt{-a^2 - 2 a b - b^2}} - \frac{\sqrt{b} \left(2 \operatorname{atan}\left(\frac{\sqrt{b} e^x \sqrt{a(a+b)^2}}{2 a(a+b)}\right) - 2 \operatorname{atan}\left(\frac{(a^3 b^{5/2} \sqrt{a^3 + 2 a^2 b + a b^2} + a^2 b^{7/2} \sqrt{a^3 + 2 a^2 b + a b^2})}{a b^3 \sqrt{a(a+b)^2} (a^2 + b a) \sqrt{a^3 + 2 a^2 b + a b^2}}\right) \right)}{2 \sqrt{a^3 + 2 a^2 b + a b^2}}$$

[In] `int(1/(sinh(x)*(a + b*cosh(x)^2)),x)`

[Out]
$$-\frac{(2*\operatorname{atan}(\exp(x)*(16*a^2*(-2*a*b - a^2 - b^2)^{1/2} + b^2*(-2*a*b - a^2 - b^2)^{1/2} + 8*a*b*(-2*a*b - a^2 - b^2)^{1/2}))/((9*a*b^2 + 24*a^2*b + 16*a^3 + b^3)))/(-2*a*b - a^2 - b^2)^{1/2} - (b^{1/2}*(2*\operatorname{atan}(b^{1/2}*\exp(x)*(a*(a + b)^2)^{1/2}))/((2*a*(a + b))) - 2*\operatorname{atan}(((a^3*b^{5/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2} + a^2*b^{7/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2})*(\exp(x)*((64*(2*a*b^2 + 10*a^2*b + 8*a^3))/(a*b^3*(a*(a + b)^2)^{1/2}*(a*b + a^2)*(a*b^2 + 2*a^2*b + a^3)^{1/2}) + (32*(b^{3/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2} + 4*a*b^{1/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2}))/((a^2*b^{5/2}*(a + b)*(a*b + a^2)*(a*b^2 + 2*a^2*b + a^3)^{1/2})) + (32*\exp(3*x)*(b^{3/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2} + 4*a*b^{1/2}*(a*b^2 + 2*a^2*b + a^3)^{1/2}))/((a^2*b^{5/2}*(a + b)*(a*b + a^2)*(a*b^2 + 2*a^2*b + a^3)^{1/2}))))/(256*a + 64*b)))/((2*(a*b^2 + 2*a^2*b + a^3)^{1/2}))$$

3.11 $\int \frac{\operatorname{csch}^3(x)}{a+b \cosh^2(x)} dx$

Optimal result	94
Rubi [A] (verified)	94
Mathematica [C] (verified)	96
Maple [A] (verified)	96
Fricas [B] (verification not implemented)	97
Sympy [F]	98
Maxima [F]	98
Giac [F]	98
Mupad [B] (verification not implemented)	99

Optimal result

Integrand size = 15, antiderivative size = 61

$$\int \frac{\operatorname{csch}^3(x)}{a+b \cosh^2(x)} dx = \frac{b^{3/2} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^2} + \frac{(a+3b) \operatorname{arctanh}(\cosh(x))}{2(a+b)^2} - \frac{\operatorname{coth}(x) \operatorname{csch}(x)}{2(a+b)}$$

[Out] $1/2*(a+3*b)*\operatorname{arctanh}(\cosh(x))/(a+b)^2 - 1/2*\operatorname{coth}(x)*\operatorname{csch}(x)/(a+b) + b^{3/2}*\operatorname{arctan}(\cosh(x)*b^{1/2}/a^{1/2})/(a+b)^2/a^{1/2}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3269, 425, 536, 212, 211}

$$\int \frac{\operatorname{csch}^3(x)}{a+b \cosh^2(x)} dx = \frac{b^{3/2} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^2} + \frac{(a+3b) \operatorname{arctanh}(\cosh(x))}{2(a+b)^2} - \frac{\operatorname{coth}(x) \operatorname{csch}(x)}{2(a+b)}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]^3/(a+b*\operatorname{Cosh}[x]^2), x]$

[Out] $(b^{3/2}*\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*\operatorname{Cosh}[x])/(\operatorname{Sqrt}[a])])/((\operatorname{Sqrt}[a]*(a+b)^2) + ((a+3*b)*\operatorname{ArcTanh}[\operatorname{Cosh}[x]])/(2*(a+b)^2) - (\operatorname{Coth}[x]*\operatorname{Csch}[x])/(2*(a+b))$

Rule 211

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2]/a)*\operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b, x\} \&\& \operatorname{PosQ}[a/b]$

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 425

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c -
a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c
+ d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n,
x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -
1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b,
c, d, n, p, q, x]
```

Rule 536

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(
n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x]
- Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b,
c, d, e, f, n}, x]
```

Rule 3269

```
Int[cos[(e_.) + (f_.)*(x_)^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)^2]^(
p_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Su
bst[Int[(1 - ff^2*x^2)^(m - 1)/2]*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/
ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1}{(1-x^2)^2(a+bx^2)} dx, x, \cosh(x)\right) \\
&= -\frac{\coth(x)\text{csch}(x)}{2(a+b)} + \frac{\text{Subst}\left(\int \frac{a+2b+bx^2}{(1-x^2)(a+bx^2)} dx, x, \cosh(x)\right)}{2(a+b)} \\
&= -\frac{\coth(x)\text{csch}(x)}{2(a+b)} + \frac{b^2\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cosh(x)\right)}{(a+b)^2} + \frac{(a+3b)\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cosh(x)\right)}{2(a+b)^2} \\
&= \frac{b^{3/2}\arctan\left(\frac{\sqrt{b}\cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^2} + \frac{(a+3b)\text{arctanh}(\cosh(x))}{2(a+b)^2} - \frac{\coth(x)\text{csch}(x)}{2(a+b)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 154, normalized size of antiderivative = 2.52

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \frac{8b^{3/2} \arctan\left(\frac{\sqrt{b-i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right) + 8b^{3/2} \arctan\left(\frac{\sqrt{b+i\sqrt{a+b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right) - \sqrt{a}(a+b) \operatorname{csch}^2\left(\frac{x}{2}\right) + 4\sqrt{a}(a+3b) \left(\log\left[\cosh\left(\frac{x}{2}\right)\right] - \log\left[\sinh\left(\frac{x}{2}\right)\right]\right) - \sqrt{a}(a+b) \operatorname{sech}^2\left(\frac{x}{2}\right)}{8\sqrt{a}(a+b)^2}$$

[In] Integrate[Csch[x]^3/(a + b*Cosh[x]^2),x]

[Out] (8*b^(3/2)*ArcTan[(Sqrt[b] - I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]] + 8*b^(3/2)*ArcTan[(Sqrt[b] + I*Sqrt[a + b]*Tanh[x/2])/Sqrt[a]] - Sqrt[a]*(a + b)*Csch[x/2]^2 + 4*Sqrt[a]*(a + 3*b)*(Log[Cosh[x/2]] - Log[Sinh[x/2]]) - Sqrt[a]*(a + b)*Sech[x/2]^2)/(8*Sqrt[a]*(a + b)^2)

Maple [A] (verified)

Time = 6.49 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.43

method	result
default	$\frac{\tanh\left(\frac{x}{2}\right)^2}{8a+8b} - \frac{1}{8(a+b)\tanh\left(\frac{x}{2}\right)^2} + \frac{(-2a-6b)\ln\left(\tanh\left(\frac{x}{2}\right)\right)}{4(a+b)^2} + \frac{b^2 \arctan\left(\frac{2(a+b)\tanh\left(\frac{x}{2}\right)^2 - 2a + 2b}{4\sqrt{ab}}\right)}{(a+b)^2\sqrt{ab}}$
risch	$-\frac{e^x(1+e^{2x})}{(e^{2x}-1)^2(a+b)} + \frac{a \ln(e^x+1)}{2a^2+4ab+2b^2} + \frac{3 \ln(e^x+1)b}{2(a^2+2ab+b^2)} - \frac{\ln(e^x-1)a}{2(a^2+2ab+b^2)} - \frac{3 \ln(e^x-1)b}{2(a^2+2ab+b^2)} + \frac{\sqrt{-ab}b \ln\left(e^{2x} + \frac{2\sqrt{-ab}e^x}{b} + 1\right)}{2a(a+b)^2} -$

[In] int(csch(x)^3/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/8*tanh(1/2*x)^2/(a+b)-1/8/(a+b)/tanh(1/2*x)^2+1/4/(a+b)^2*(-2*a-6*b)*ln(tanh(1/2*x))+b^2/(a+b)^2/(a*b)^(1/2)*arctan(1/4*(2*(a+b)*tanh(1/2*x)^2-2*a+2*b)/(a*b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 638 vs. 2(49) = 98.

Time = 0.30 (sec) , antiderivative size = 1332, normalized size of antiderivative = 21.84

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(csch(x)^3/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out]
$$\begin{aligned} & [-1/2*(2*(a + b)*\cosh(x)^3 + 6*(a + b)*\cosh(x)*\sinh(x)^2 + 2*(a + b)*\sinh(x)^3 - (b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + b)*\sqrt{-b/a}*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a - b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a - b)*\cosh(x))*\sinh(x) + 4*(a*\cosh(x)^3 + 3*a*\cosh(x)*\sinh(x)^2 + a*\sinh(x)^3 + a*\cosh(x) + (3*a*\cosh(x)^2 + a)*\sinh(x))*\sqrt{-b/a} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 2*(a + b)*\cosh(x) - ((a + 3*b)*\cosh(x)^4 + 4*(a + 3*b)*\cosh(x)*\sinh(x)^3 + (a + 3*b)*\sinh(x)^4 - 2*(a + 3*b)*\cosh(x)^2 + 2*(3*(a + 3*b)*\cosh(x)^2 - a - 3*b)*\sinh(x)^2 + 4*((a + 3*b)*\cosh(x)^3 - (a + 3*b)*\cosh(x))*\sinh(x) + a + 3*b)*\log(\cosh(x) + \sinh(x) + 1) + ((a + 3*b)*\cosh(x)^4 + 4*(a + 3*b)*\cosh(x)*\sinh(x)^3 + (a + 3*b)*\sinh(x)^4 - 2*(a + 3*b)*\cosh(x)^2 + 2*(3*(a + 3*b)*\cosh(x)^2 - a - 3*b)*\sinh(x)^2 + 4*((a + 3*b)*\cosh(x)^3 - (a + 3*b)*\cosh(x))*\sinh(x) + a + 3*b)*\log(\cosh(x) + \sinh(x) - 1) + 2*(3*(a + b)*\cosh(x)^2 + a + b)*\sinh(x))/((a^2 + 2*a*b + b^2)*\cosh(x)^4 + 4*(a^2 + 2*a*b + b^2)*\cosh(x)*\sinh(x)^3 + (a^2 + 2*a*b + b^2)*\sinh(x)^4 - 2*(a^2 + 2*a*b + b^2)*\cosh(x)^2 + 2*(3*(a^2 + 2*a*b + b^2)*\cosh(x)^2 - a^2 - 2*a*b - b^2)*\sinh(x)^2 + a^2 + 2*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*\cosh(x)^3 - (a^2 + 2*a*b + b^2)*\cosh(x))*\sinh(x)), -1/2*(2*(a + b)*\cosh(x)^3 + 6*(a + b)*\cosh(x)*\sinh(x)^2 + 2*(a + b)*\sinh(x)^3 - 2*(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + b)*\sqrt{b/a}*\arctan(1/2*\sqrt{b/a}*(\cosh(x) + \sinh(x))) + 2*(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*b*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - b*\cosh(x))*\sinh(x) + b)*\sqrt{b/a}*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x)^3 + (4*a + b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + b)*\sinh(x))*\sqrt{b/a}/b) + 2*(a + b)*\cosh(x) - ((a + 3*b)*\cosh(x)^4 + 4*(a + 3*b)*\cosh(x)*\sinh(x)^3 + (a + 3*b)*\sinh(x)^4 - 2*(a + 3*b)*\cosh(x)^2 + 2*(3*(a + 3*b)*\cosh(x)^2 - a - 3*b)*\sinh(x)^2 + 4*((a + 3*b)*\cosh(x)^3 - (a + 3*b)*\cosh(x))*\sinh(x) + a + 3*b)*\log(\cosh(x) + \sinh(x) + 1) + ((a + 3*b)*\cosh(x)^4 + 4*(a + 3*b)*\cosh(x)*\sinh(x)^3 + (a + 3*b)*\sinh(x)^4 - 2*(a + 3*b)*\cosh(x)^2 + 2*(3*(a + 3*b)*\cosh(x)^2 - a - 3*b)*\sinh(x)^2 + 4*((a + 3*b)*\cosh(x)^3 - (a + 3*b)*\cosh(x))*\sinh(x) + a + 3*b)*\log(\cosh(x) + \sinh(x) - 1) + 2*(3*(a + b)*\cosh(x)^2$$

```
+ a + b)*sinh(x))/((a^2 + 2*a*b + b^2)*cosh(x)^4 + 4*(a^2 + 2*a*b + b^2)*co
sh(x)*sinh(x)^3 + (a^2 + 2*a*b + b^2)*sinh(x)^4 - 2*(a^2 + 2*a*b + b^2)*cos
h(x)^2 + 2*(3*(a^2 + 2*a*b + b^2)*cosh(x)^2 - a^2 - 2*a*b - b^2)*sinh(x)^2
+ a^2 + 2*a*b + b^2 + 4*((a^2 + 2*a*b + b^2)*cosh(x)^3 - (a^2 + 2*a*b + b^2
)*cosh(x))*sinh(x))]
```

Sympy [F]

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx$$

```
[In] integrate(csch(x)**3/(a+b*cosh(x)**2),x)
```

```
[Out] Integral(csch(x)**3/(a + b*cosh(x)**2), x)
```

Maxima [F]

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^3}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(csch(x)^3/(a+b*cosh(x)^2),x, algorithm="maxima")
```

```
[Out] 1/2*(a + 3*b)*log(e^x + 1)/(a^2 + 2*a*b + b^2) - 1/2*(a + 3*b)*log(e^x - 1)
/(a^2 + 2*a*b + b^2) - (e^(3*x) + e^x)/((a + b)*e^(4*x) - 2*(a + b)*e^(2*x)
+ a + b) + 8*integrate(1/4*(b^2*e^(3*x) - b^2*e^x)/(a^2*b + 2*a*b^2 + b^3
+ (a^2*b + 2*a*b^2 + b^3)*e^(4*x) + 2*(2*a^3 + 5*a^2*b + 4*a*b^2 + b^3)*e^(
2*x)), x)
```

Giac [F]

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^3}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(csch(x)^3/(a+b*cosh(x)^2),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [B] (verification not implemented)

Time = 8.26 (sec) , antiderivative size = 2225, normalized size of antiderivative = 36.48

$$\int \frac{\operatorname{csch}^3(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] int(1/(sinh(x)^3*(a + b*cosh(x)^2)),x)

[Out] ((2*atan((b^2*exp(x)*(a*(a + b)^4)^(1/2))/(2*a*(a + b)^2*(b^3)^(1/2))) - 2*atan((exp(x)*((32*(b^8*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 36*a^2*b^6*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 47*a^3*b^5*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 30*a^4*b^4*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 9*a^5*b^3*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + a^6*b^2*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 12*a*b^7*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2)))/(a^2*b^2*(a + b)^7*(a*b + a^2)*(b^3)^(1/2)*(2*a*b + a^2 + b^2)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)*(9*a*b^2 + 6*a^2*b + a^3 + b^3)*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2)) + (64*(20*a^3*(b^3)^(5/2) + 232*a^6*(b^3)^(3/2) + 2*a^9*(b^3)^(1/2) + 10*a^2*b^4*(b^3)^(3/2) + 20*a^4*b^2*(b^3)^(3/2) + 18*a^2*b^7*(b^3)^(1/2) + 102*a^3*b^6*(b^3)^(1/2) + 242*a^4*b^5*(b^3)^(1/2) + 310*a^5*b^4*(b^3)^(1/2) + 98*a^7*b^2*(b^3)^(1/2) + 2*a*b^5*(b^3)^(3/2) + 10*a^5*b*(b^3)^(3/2) + 22*a^8*b*(b^3)^(1/2)))/(a*b^4*(a + b)^5*(a*b + a^2)*(a*(a + b)^4)^(1/2)*(2*a*b + a^2 + b^2)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)*(9*a*b^2 + 6*a^2*b + a^3 + b^3)*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))) + (32*exp(3*x)*(b^8*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 36*a^2*b^6*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 47*a^3*b^5*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 30*a^4*b^4*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 9*a^5*b^3*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + a^6*b^2*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2) + 12*a*b^7*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2)))/(a^2*b^2*(a + b)^7*(a*b + a^2)*(b^3)^(1/2)*(2*a*b + a^2 + b^2)*(3*a*b^2 + 3*a^2*b + a^3 + b^3)*(9*a*b^2 + 6*a^2*b + a^3 + b^3)*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2)))*((a^2*b^10*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/64 + (a^3*b^9*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/8 + (7*a^4*b^8*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/16 + (7*a^5*b^7*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/8 + (35*a^6*b^6*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/32 + (7*a^7*b^5*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/8 + (7*a^8*b^4*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/16 + (a^9*b^3*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/8 + (a^10*b^2*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2))/64)))*(b^3)^(1/2))/(2*(a*b^4 + 4*a^4*b + a^5 + 4*a^2*b^3 + 6*a^3*b^2)^(1/2)) - (2*exp(x))/((a + b)*(exp(4*x) - 2*exp(2*x) + 1)) - exp(x)/((a + b)*(exp(2*x) - 1)) - (atan((exp(x)*(a^7

$$\begin{aligned}
& *(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} + 3b^7(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} + 55a^6b^6(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} \\
& + 15a^6b^6(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} + 297a^2b^5(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} + 423a^3b^4(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} \\
& + 272a^4b^3(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)} + 90a^5b^2(-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(3/2)}) / (a^{12}(6ab + a^2 + 9b^2)^{(1/2)} \\
& + b^{12}(6ab + a^2 + 9b^2)^{(1/2)} + 24a^{11}b(6ab + a^2 + 9b^2)^{(1/2)} + 18a^{11}b(6ab + a^2 + 9b^2)^{(1/2)} + 216a^2b^{10}(6ab + a^2 + 9b^2)^{(1/2)} \\
& + 958a^3b^9(6ab + a^2 + 9b^2)^{(1/2)} + 2484a^4b^8(6ab + a^2 + 9b^2)^{(1/2)} + 4122a^5b^7(6ab + a^2 + 9b^2)^{(1/2)} + 4587a^6b^6(6ab + a^2 + 9b^2)^{(1/2)} \\
& + 3492a^7b^5(6ab + a^2 + 9b^2)^{(1/2)} + 1818a^8b^4(6ab + a^2 + 9b^2)^{(1/2)} + 634a^9b^3(6ab + a^2 + 9b^2)^{(1/2)} + 141a^{10}b^2(6ab + a^2 + 9b^2)^{(1/2)}) \\
& * (6ab + a^2 + 9b^2)^{(1/2)} / (-4ab^3 - 4a^3b - a^4 - b^4 - 6a^2b^2)^{(1/2)}
\end{aligned}$$

3.12 $\int \frac{\operatorname{csch}^5(x)}{a+b \cosh^2(x)} dx$

Optimal result	101
Rubi [A] (verified)	101
Mathematica [C] (verified)	103
Maple [A] (verified)	104
Fricas [B] (verification not implemented)	104
Sympy [F(-1)]	104
Maxima [F]	105
Giac [F]	105
Mupad [B] (verification not implemented)	105

Optimal result

Integrand size = 15, antiderivative size = 94

$$\int \frac{\operatorname{csch}^5(x)}{a+b \cosh^2(x)} dx = -\frac{b^{5/2} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^3} - \frac{(3a^2 + 10ab + 15b^2) \operatorname{arctanh}(\cosh(x))}{8(a+b)^3} + \frac{(3a+7b) \operatorname{coth}(x) \operatorname{csch}(x)}{8(a+b)^2} - \frac{\operatorname{coth}(x) \operatorname{csch}^3(x)}{4(a+b)}$$

[Out] $-1/8*(3*a^2+10*a*b+15*b^2)*\operatorname{arctanh}(\cosh(x))/(a+b)^3+1/8*(3*a+7*b)*\operatorname{coth}(x)*\operatorname{csch}(x)/(a+b)^2-1/4*\operatorname{coth}(x)*\operatorname{csch}(x)^3/(a+b)-b^{(5/2)}*\operatorname{arctan}(\cosh(x)*b^{(1/2)}/a^{(1/2)})/(a+b)^3/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.15 (sec), antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3269, 425, 541, 536, 212, 211}

$$\int \frac{\operatorname{csch}^5(x)}{a+b \cosh^2(x)} dx = -\frac{(3a^2 + 10ab + 15b^2) \operatorname{arctanh}(\cosh(x))}{8(a+b)^3} - \frac{b^{5/2} \arctan\left(\frac{\sqrt{b} \cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^3} - \frac{\operatorname{coth}(x) \operatorname{csch}^3(x)}{4(a+b)} + \frac{(3a+7b) \operatorname{coth}(x) \operatorname{csch}(x)}{8(a+b)^2}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]^5/(a+b*\operatorname{Cosh}[x]^2),x]$

[Out] $-((b^{(5/2)}*\operatorname{ArcTan}[(\operatorname{Sqrt}[b]*\operatorname{Cosh}[x])/(\operatorname{Sqrt}[a])])/(a+b)^3) - ((3*a^2 + 10*a*b + 15*b^2)*\operatorname{ArcTanh}[\operatorname{Cosh}[x]])/(8*(a+b)^3) + ((3*a + 7*b)*\operatorname{Coth}[x]*\operatorname{Csch}[x])/(8*(a+b)^2) - (\operatorname{Coth}[x]*\operatorname{Csch}[x]^3)/(4*(a+b))$

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 425

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(!IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]

Rule 536

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 541

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]

Rule 3269

Int[cos[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*ff^2*x^2)^p, x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{1}{(1-x^2)^3(a+bx^2)} dx, x, \cosh(x)\right) \\
&= -\frac{\coth(x)\text{csch}^3(x)}{4(a+b)} - \frac{\text{Subst}\left(\int \frac{3a+4b+3bx^2}{(1-x^2)^2(a+bx^2)} dx, x, \cosh(x)\right)}{4(a+b)} \\
&= \frac{(3a+7b)\coth(x)\text{csch}(x)}{8(a+b)^2} - \frac{\coth(x)\text{csch}^3(x)}{4(a+b)} - \frac{\text{Subst}\left(\int \frac{3a^2+7ab+8b^2+b(3a+7b)x^2}{(1-x^2)(a+bx^2)} dx, x, \cosh(x)\right)}{8(a+b)^2} \\
&= \frac{(3a+7b)\coth(x)\text{csch}(x)}{8(a+b)^2} - \frac{\coth(x)\text{csch}^3(x)}{4(a+b)} - \frac{b^3\text{Subst}\left(\int \frac{1}{a+bx^2} dx, x, \cosh(x)\right)}{(a+b)^3} \\
&\quad - \frac{(3a^2+10ab+15b^2)\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cosh(x)\right)}{8(a+b)^3} \\
&= -\frac{b^{5/2}\arctan\left(\frac{\sqrt{b}\cosh(x)}{\sqrt{a}}\right)}{\sqrt{a}(a+b)^3} - \frac{(3a^2+10ab+15b^2)\text{arctanh}(\cosh(x))}{8(a+b)^3} \\
&\quad + \frac{(3a+7b)\coth(x)\text{csch}(x)}{8(a+b)^2} - \frac{\coth(x)\text{csch}^3(x)}{4(a+b)}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 0.54 (sec) , antiderivative size = 229, normalized size of antiderivative = 2.44

$$\int \frac{\text{csch}^5(x)}{a+b\cosh^2(x)} dx$$

$$2\sqrt{a}(3a^2+10ab+7b^2)\text{csch}^2\left(\frac{x}{2}\right) - \sqrt{a}(a+b)^2\text{csch}^4\left(\frac{x}{2}\right) - 8\left(8b^{5/2}\arctan\left(\frac{\sqrt{b}-i\sqrt{a+b}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right) + 8b^{5/2}\arctan\left(\frac{\sqrt{b}+i\sqrt{a+b}\tanh\left(\frac{x}{2}\right)}{\sqrt{a}}\right)\right)$$

[In] Integrate[Csch[x]^5/(a + b*Cosh[x]^2), x]

[Out] (2*sqrt[a]*(3*a^2 + 10*a*b + 7*b^2)*Csch[x/2]^2 - sqrt[a]*(a + b)^2*Csch[x/2]^4 - 8*(8*b^(5/2)*ArcTan[(sqrt[b] - I*sqrt[a + b]*Tanh[x/2])/sqrt[a]] + 8*b^(5/2)*ArcTan[(sqrt[b] + I*sqrt[a + b]*Tanh[x/2])/sqrt[a]] + sqrt[a]*(3*a^2 + 10*a*b + 15*b^2)*(Log[Cosh[x/2]] - Log[Sinh[x/2]])) + 2*sqrt[a]*(3*a^2 + 10*a*b + 7*b^2)*Sech[x/2]^2 + sqrt[a]*(a + b)^2*Sech[x/2]^4)/(64*sqrt[a]*(a + b)^3)

Maple [A] (verified)

Time = 29.97 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.44

method	result
default	$\frac{\left(\tanh\left(\frac{x}{2}\right)^2 a + \tanh\left(\frac{x}{2}\right)^2 b - 4a - 8b\right)^2}{64(a+b)^3} - \frac{1}{64(a+b)\tanh\left(\frac{x}{2}\right)^4} - \frac{-4a-8b}{32(a+b)^2\tanh\left(\frac{x}{2}\right)^2} + \frac{(6a^2+20ab+30b^2)\ln\left(\tanh\left(\frac{x}{2}\right)\right)}{16(a+b)^3} - \frac{b^3\arctan\left(\frac{1}{4}\left(2(a+b)\tanh\left(\frac{x}{2}\right)\right)^2-2a+2b\right)}{(a+b)^3(a+b)^2}$
risch	$\frac{e^x(3ae^{6x}+7be^{6x}-11ae^{4x}-15be^{4x}-11ae^{2x}-15be^{2x}+3a+7b)}{4(e^{2x}-1)^4(a+b)^2} + \frac{3\ln(e^x-1)a^2}{8(a^3+3a^2b+3ab^2+b^3)} + \frac{5\ln(e^x-1)ab}{4(a^3+3a^2b+3ab^2+b^3)} + \frac{15\ln(e^x-1)b^3}{8(a^3+3a^2b+3ab^2+b^3)}$

[In] int(csch(x)^5/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/64*(tanh(1/2*x)^2*a+tanh(1/2*x)^2*b-4*a-8*b)^2/(a+b)^3-1/64/(a+b)/tanh(1/2*x)^4-1/32*(-4*a-8*b)/(a+b)^2/tanh(1/2*x)^2+1/16/(a+b)^3*(6*a^2+20*a*b+30*b^2)*ln(tanh(1/2*x))-b^3/(a+b)^3/(a*b)^(1/2)*arctan(1/4*(2*(a+b)*tanh(1/2*x))^2-2*a+2*b)/(a*b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2724 vs. 2(80) = 160.

Time = 0.35 (sec) , antiderivative size = 5326, normalized size of antiderivative = 56.66

$$\int \frac{\operatorname{csch}^5(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(csch(x)^5/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^5(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(csch(x)**5/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\operatorname{csch}^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)^5/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-1/8*(3*a^2 + 10*a*b + 15*b^2)*\log(e^x + 1)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3)$
 $+ 1/8*(3*a^2 + 10*a*b + 15*b^2)*\log(e^x - 1)/(a^3 + 3*a^2*b + 3*a*b^2 + b^3)$
 $+ 1/4*((3*a + 7*b)*e^{7*x} - (11*a + 15*b)*e^{5*x} - (11*a + 15*b)*e^{3*x})$
 $+ (3*a + 7*b)*e^x/(a^2 + 2*a*b + b^2 + (a^2 + 2*a*b + b^2)*e^{8*x} - 4*(a^2 + 2*a*b + b^2)*e^{6*x} + 6*(a^2 + 2*a*b + b^2)*e^{4*x} - 4*(a^2 + 2*a*b + b^2)*e^{2*x})$
 $- 32*\integrate(1/16*(b^3*e^{3*x} - b^3*e^x)/(a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4 + (a^3*b + 3*a^2*b^2 + 3*a*b^3 + b^4)*e^{4*x} + 2*(2*a^4 + 7*a^3*b + 9*a^2*b^2 + 5*a*b^3 + b^4)*e^{2*x}), x)$

Giac [F]

$$\int \frac{\operatorname{csch}^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)^5/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 17.03 (sec) , antiderivative size = 5056, normalized size of antiderivative = 53.79

$$\int \frac{\operatorname{csch}^5(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] int(1/(sinh(x)^5*(a + b*cosh(x)^2)),x)

[Out] $(\operatorname{atan}((\exp(x)*(243*a^{12}*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 3840*b^{12}*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 110560*a*b^{11}*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 4050*a^{11}*b*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 976143*a^2*b^{10}*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 2740050*a^3*b^9*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2))^{3/2} + 4252775*a^4*b^8*(-6*a$

$$\begin{aligned}
& *b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2)^{(3/2)} + \\
& 4316760*a^5*b^7*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 \\
& - 15*a^4*b^2)^{(3/2)} + 3087390*a^6*b^6*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15 \\
& *a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2)^{(3/2)} + 1608364*a^7*b^5*(-6*a*b^5 - 6* \\
& a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2)^{(3/2)} + 615750*a^ \\
& 8*b^4*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b \\
& ^2)^{(3/2)} + 171000*a^9*b^3*(-6*a*b^5 - 6*a^5*b - a^6 - b^6 - 15*a^2*b^4 - \\
& 20*a^3*b^3 - 15*a^4*b^2)^{(3/2)} + 33075*a^10*b^2*(-6*a*b^5 - 6*a^5*b - a^6 \\
& - b^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2)^{(3/2)))/(81*a^19*(300*a*b^3 + \\
& 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 256*b^19*(300*a*b^3 + 60 \\
& *a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 9504*a*b^18*(300*a*b^3 + 60 \\
& *a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 1809*a^18*b*(300*a*b^3 + 60 \\
& *a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 134241*a^2*b^17*(300*a*b^3 \\
& + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 963809*a^3*b^16*(300*a* \\
& b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 4252296*a^4*b^15*(3 \\
& 00*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 12815304*a^5*b \\
& ^14*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 28102636 \\
& *a^6*b^13*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 46 \\
& 681644*a^7*b^12*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} \\
&) + 60321816*a^8*b^11*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2 \\
&)^((1/2) + 61717144*a^9*b^10*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a \\
& ^2*b^2)^{(1/2)} + 50559894*a^10*b^9*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225*b^4 + \\
& 190*a^2*b^2)^{(1/2)} + 33362646*a^11*b^8*(300*a*b^3 + 60*a^3*b + 9*a^4 + 225 \\
& *b^4 + 190*a^2*b^2)^{(1/2)} + 17752184*a^12*b^7*(300*a*b^3 + 60*a^3*b + 9*a^4 \\
& + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 7586616*a^13*b^6*(300*a*b^3 + 60*a^3*b + \\
& 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 2577804*a^14*b^5*(300*a*b^3 + 60*a^3 \\
& *b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 683596*a^15*b^4*(300*a*b^3 + 60 \\
& *a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 137064*a^16*b^3*(300*a*b^3 \\
& + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)} + 19656*a^17*b^2*(300*a*b \\
& ^3 + 60*a^3*b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)))*(300*a*b^3 + 60*a^3* \\
& b + 9*a^4 + 225*b^4 + 190*a^2*b^2)^{(1/2)))/(4*(-6*a*b^5 - 6*a^5*b - a^6 - b \\
& ^6 - 15*a^2*b^4 - 20*a^3*b^3 - 15*a^4*b^2)^{(1/2)) - (4*exp(x))/((a + b)*(6* \\
& exp(4*x) - 4*exp(2*x) - 4*exp(6*x) + exp(8*x) + 1)) - ((b^5)^{(1/2)}*(2*atan(\\
& (b^3*exp(x)*(a*(a + b)^6)^{(1/2)))/(2*a*(a + b)^3*(b^5)^{(1/2)))) - 2*atan((exp \\
& (x))*((2*(16*b^14*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4*b \\
& ^3 + 15*a^5*b^2)^{(1/2)} + 321*a*b^13*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15 \\
& *a^3*b^4 + 20*a^4*b^3 + 15*a^5*b^2)^{(1/2)} + 1890*a^2*b^12*(a*b^6 + 6*a^6*b \\
& + a^7 + 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4*b^3 + 15*a^5*b^2)^{(1/2)} + 5685*a^3* \\
& b^11*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4*b^3 + 15*a^5* \\
& b^2)^{(1/2)} + 10440*a^4*b^10*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15*a^3*b^4 \\
& + 20*a^4*b^3 + 15*a^5*b^2)^{(1/2)} + 12690*a^5*b^9*(a*b^6 + 6*a^6*b + a^7 + \\
& 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4*b^3 + 15*a^5*b^2)^{(1/2)} + 10620*a^6*b^8*(a \\
& b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4*b^3 + 15*a^5*b^2)^{(1/ \\
& 2)} + 6210*a^7*b^7*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 + 15*a^3*b^4 + 20*a^4* \\
& b^3 + 15*a^5*b^2)^{(1/2)} + 2520*a^8*b^6*(a*b^6 + 6*a^6*b + a^7 + 6*a^2*b^5 +
\end{aligned}$$

$$\begin{aligned}
& 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 685a^9b^5(a^6b + 6a^6b \\
& + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 114a^{10} \\
& *b^4(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 9a^{11}b^3(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 2 \\
& 0a^4b^3 + 15a^5b^2)^{(1/2)))/(a^2b(a+b)^{10}(a+b+a^2)(b^5)^{(1/2)}*(\\
& 3a^2b + 3a^2b + a^3 + b^3)*(4a^3b + 4a^3b + a^4 + b^4 + 6a^2b^2)* \\
& (225a^4b + 60a^4b + 9a^5 + 16b^5 + 300a^2b^3 + 190a^3b^2)*(6a^5b + 6a^5b + a^6 + b^6 + 15a^2b^4 + 20a^3b^3 + 15a^4b^2)*(a^6b + 6a^6b \\
& + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + (4 \\
& *(4032a^5(b^5)^{(5/2)} + 74990a^{10}(b^5)^{(3/2)} + 18a^{15}(b^5)^{(1/2)} + 288 \\
& *a^2b^8(b^5)^{(3/2)} + 1152a^3b^7(b^5)^{(3/2)} + 2688a^4b^6(b^5)^{(3/2)} \\
& + 4032a^6b^4(b^5)^{(3/2)} + 2688a^7b^3(b^5)^{(3/2)} + 1152a^8b^2(b^5)^{(3/2)} \\
& + 450a^2b^{13}(b^5)^{(1/2)} + 4650a^3b^{12}(b^5)^{(1/2)} + 21980a^4b^{11}(b^5)^{(1/2)} + 62940a^5b^{10}(b^5)^{(1/2)} + 121878a^6b^9(b^5)^{(1/2)} + \\
& 168702a^7b^8(b^5)^{(1/2)} + 172008a^8b^7(b^5)^{(1/2)} + 131112a^9b^6(b^5)^{(1/2)} + 31878a^{11}b^4(b^5)^{(1/2)} + 9852a^{12}b^3(b^5)^{(1/2)} + 2108a^{13}b^2(b^5)^{(1/2)} + 32a^9b^9(b^5)^{(3/2)} + 288a^9b^9(b^5)^{(3/2)} + 282a^{14}b^9(b^5)^{(1/2)))/(a^4b^4(a+b)^7(a+b+a^2)(a(a+b)^6)^{(1/2)}*(3a^2b + 3a^2b + a^3 + b^3)*(4a^3b + 4a^3b + a^4 + b^4 + 6a^2b^2)*(225a^4b + 60a^4b + 9a^5 + 16b^5 + 300a^2b^3 + 190a^3b^2)*(6a^5b + 6a^5b + a^6 + b^6 + 15a^2b^4 + 20a^3b^3 + 15a^4b^2)*(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)) + (2*exp(3*x)*(16b^{14}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 321a^9b^{13}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 1890a^2b^{12}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 5685a^3b^{11}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 10440a^4b^{10}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 12690a^5b^9(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 10620a^6b^8(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 6210a^7b^7(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 2520a^8b^6(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 685a^9b^5(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 114a^{10}b^4(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)} + 9a^{11}b^3(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))/(a^2b(a+b)^{10}(a+b+a^2)(b^5)^{(1/2)}*(3a^2b + 3a^2b + a^3 + b^3)*(4a^3b + 4a^3b + a^4 + b^4 + 6a^2b^2)*(225a^4b + 60a^4b + 9a^5 + 16b^5 + 300a^2b^3 + 190a^3b^2)*(6a^5b + 6a^5b + a^6 + b^6 + 15a^2b^4 + 20a^3b^3 + 15a^4b^2)*(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))*((a^{17}b^9(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))/4 + (a^2b^{16}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))/4 + (15a^3b^{15}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))/4 + (15a^3b^{15}(a^6b + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2)))/4
\end{aligned}$$

$$\begin{aligned}
& a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1/2))/4 + (105a^4b^{14}(a \\
& *b^6 + 6a^6b + a^7 + 6a^2b^5 + 15a^3b^4 + 20a^4b^3 + 15a^5b^2)^{(1 \\
& /2))/4 + (455a^5b^{13}(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20 \\
& *a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (1365a^6*b^{12}(a*b^6 + 6a^6*b + a^7 + 6 \\
& *a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (3003a^7*b^{11}* \\
& (a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2)^{(\\
& 1/2))/4 + (5005a^8*b^{10}(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + \\
& 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (6435a^9*b^9*(a*b^6 + 6a^6*b + a^7 + \\
& 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (6435a^{10}*b^8 \\
& 8*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2 \\
&)^{(1/2))/4 + (5005a^{11}*b^7*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 \\
& + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (3003a^{12}*b^6*(a*b^6 + 6a^6*b + a^ \\
& 7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (1365a^{13} \\
& *b^5*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5* \\
& b^2)^{(1/2))/4 + (455a^{14}*b^4*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b \\
& ^4 + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (105a^{15}*b^3*(a*b^6 + 6a^6*b + a \\
& ^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b^2)^{(1/2))/4 + (15a^{16} \\
& b^2*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^4*b^3 + 15a^5*b \\
& ^2)^{(1/2))/4))))/(2*(a*b^6 + 6a^6*b + a^7 + 6a^2*b^5 + 15a^3*b^4 + 20a^ \\
& 4*b^3 + 15a^5*b^2)^{(1/2)) - (6*exp(x))/((a + b)*(3*exp(2*x) - 3*exp(4*x) + \\
& exp(6*x) - 1)) + (exp(x)*(10*a*b + 3*a^2 + 7*b^2))/(4*(a + b)^3*(exp(2*x) \\
& - 1)) - (exp(x)*(a - 3*b))/(2*(a + b)^2*(exp(4*x) - 2*exp(2*x) + 1))
\end{aligned}$$

3.13 $\int \frac{\sinh^6(x)}{a+b \cosh^2(x)} dx$

Optimal result	109
Rubi [A] (verified)	109
Mathematica [A] (verified)	111
Maple [B] (verified)	112
Fricas [B] (verification not implemented)	112
Sympy [F(-1)]	113
Maxima [B] (verification not implemented)	114
Giac [B] (verification not implemented)	115
Mupad [B] (verification not implemented)	116

Optimal result

Integrand size = 15, antiderivative size = 88

$$\int \frac{\sinh^6(x)}{a+b \cosh^2(x)} dx = \frac{(8a^2 + 20ab + 15b^2)x}{8b^3} - \frac{(a+b)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^3}} - \frac{(4a+7b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh(x) \sinh^3(x)}{4b}$$

[Out] 1/8*(8*a^2+20*a*b+15*b^2)*x/b^3-1/8*(4*a+7*b)*cosh(x)*sinh(x)/b^2+1/4*cosh(x)*sinh(x)^3/b-(a+b)^(5/2)*arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))/b^3/a^(1/2)

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3270, 425, 541, 536, 212, 214}

$$\int \frac{\sinh^6(x)}{a+b \cosh^2(x)} dx = \frac{x(8a^2 + 20ab + 15b^2)}{8b^3} - \frac{(a+b)^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^3}} - \frac{(4a+7b) \sinh(x) \cosh(x)}{8b^2} + \frac{\sinh^3(x) \cosh(x)}{4b}$$

[In] Int[Sinh[x]^6/(a + b*Cosh[x]^2),x]

[Out] ((8*a^2 + 20*a*b + 15*b^2)*x)/(8*b^3) - ((a + b)^(5/2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(Sqrt[a]*b^3) - ((4*a + 7*b)*Cosh[x]*Sinh[x])/(8*b^2) + (Cosh[x]*Sinh[x]^3)/(4*b)

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 425

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c -
a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c
+ d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n,
x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -
1] && !( !IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b,
c, d, n, p, q, x]
```

Rule 536

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(
n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x]
- Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b,
c, d, e, f, n}, x]
```

Rule 541

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_.)*((e_) + (f
_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c
+ d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(
p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*
c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; Fre
eQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 3270

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Sub
st[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e
+ f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\text{integral} = -\text{Subst}\left(\int \frac{1}{(1-x^2)^3(a-(a+b)x^2)} dx, x, \coth(x)\right)$$

$$\begin{aligned}
&= \frac{\cosh(x) \sinh^3(x)}{4b} + \frac{\text{Subst}\left(\int \frac{-a-4b-3(a+b)x^2}{(1-x^2)^2(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{4b} \\
&= -\frac{(4a+7b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh(x) \sinh^3(x)}{4b} \\
&\quad - \frac{\text{Subst}\left(\int \frac{4a^2+9ab+8b^2+(a+b)(4a+7b)x^2}{(1-x^2)(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{8b^2} \\
&= -\frac{(4a+7b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh(x) \sinh^3(x)}{4b} \\
&\quad - \frac{(a+b)^3 \text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{b^3} \\
&\quad + \frac{(8a^2+20ab+15b^2) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(x)\right)}{8b^3} \\
&= \frac{(8a^2+20ab+15b^2)x}{8b^3} - \frac{(a+b)^{5/2} \text{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^3}} \\
&\quad - \frac{(4a+7b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh(x) \sinh^3(x)}{4b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.86

$$\begin{aligned}
&\int \frac{\sinh^6(x)}{a+b \cosh^2(x)} dx \\
&= \frac{4(8a^2+20ab+15b^2)x - \frac{32(a+b)^{5/2} \text{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}} - 8b(a+2b) \sinh(2x) + b^2 \sinh(4x)}{32b^3}
\end{aligned}$$

[In] Integrate[Sinh[x]^6/(a + b*Cosh[x]^2), x]

[Out] (4*(8*a^2 + 20*a*b + 15*b^2)*x - (32*(a + b)^(5/2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a] - 8*b*(a + 2*b)*Sinh[2*x] + b^2*Sinh[4*x])/(32*b^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 288 vs. $2(74) = 148$.

Time = 0.08 (sec) , antiderivative size = 289, normalized size of antiderivative = 3.28

$$\frac{2(a^3 + 3a^2b + 3ab^2 + b^3) \left(-\frac{\ln(\sqrt{a+b} \tanh(\frac{x}{2})^2 + 2 \tanh(\frac{x}{2}) \sqrt{a+b})}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln(\sqrt{a+b} \tanh(\frac{x}{2})^2 - 2 \tanh(\frac{x}{2}) \sqrt{a+b})}{4\sqrt{a}\sqrt{a+b}} \right)}{b^3} + \frac{4b \left(\tanh(\frac{x}{2}) - 1 \right)^4}{b^3}$$

[In] `int(sinh(x)^6/(a+b*cosh(x)^2),x)`

[Out] $\frac{2}{b^3} (a^3 + 3a^2b + 3ab^2 + b^3) \left(-\frac{1}{4} \frac{1}{a^{1/2}} \frac{1}{(a+b)^{1/2}} \ln((a+b)^{1/2} \tanh(1/2*x)^2 + 2 \tanh(1/2*x) \sqrt{a+b}) + \frac{1}{4} \frac{1}{a^{1/2}} \frac{1}{(a+b)^{1/2}} \ln((a+b)^{1/2} \tanh(1/2*x)^2 - 2 \tanh(1/2*x) \sqrt{a+b}) \right) + \frac{1}{4} \frac{b}{(\tanh(1/2*x) - 1)^4} + \frac{1}{8} \frac{b}{(\tanh(1/2*x) - 1)^3} - \frac{1}{8} \frac{(4a+7b)}{b^2} \frac{1}{(\tanh(1/2*x) - 1)} - \frac{1}{8} \frac{(5b+4a)}{b^2} \frac{1}{(\tanh(1/2*x) - 1)^2} + \frac{1}{8} \frac{b^3}{b^3} (-8a^2 - 20ab - 15b^2) \ln(\tanh(1/2*x) - 1) - \frac{1}{4} \frac{b}{(\tanh(1/2*x) + 1)^4} + \frac{1}{8} \frac{b}{(\tanh(1/2*x) + 1)^3} - \frac{1}{8} \frac{(4a+7b)}{b^2} \frac{1}{(\tanh(1/2*x) + 1)} - \frac{1}{8} \frac{(-5b-4a)}{b^2} \frac{1}{(\tanh(1/2*x) + 1)^2} + \frac{1}{8} \frac{(8a^2 + 20ab + 15b^2)}{b^3} \ln(\tanh(1/2*x) + 1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 566 vs. $2(74) = 148$.

Time = 0.29 (sec) , antiderivative size = 1308, normalized size of antiderivative = 14.86

$$\int \frac{\sinh^6(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] `integrate(sinh(x)^6/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $\frac{1}{64} (b^2 \cosh(x)^8 + 8b^2 \cosh(x) \sinh(x)^7 + b^2 \sinh(x)^8 - 8(a*b + 2*b^2) \cosh(x)^6 + 4(7b^2 \cosh(x)^2 - 2a*b - 4b^2) \sinh(x)^6 + 8(8a^2 + 20a*b + 15b^2) x \cosh(x)^4 + 8(7b^2 \cosh(x)^3 - 6(a*b + 2b^2) \cosh(x)) \sinh(x)^5 + 2(35b^2 \cosh(x)^4 - 60(a*b + 2b^2) \cosh(x)^2 + 4(8a^2 + 20a*b + 15b^2) x) \sinh(x)^4 + 8(7b^2 \cosh(x)^5 - 20(a*b + 2b^2) \cosh(x)^3 + 4(8a^2 + 20a*b + 15b^2) x \cosh(x)) \sinh(x)^3 + 8(a*b + 2b^2) \cosh(x)^2 + 4(7b^2 \cosh(x)^6 - 30(a*b + 2b^2) \cosh(x)^4 + 12(8a^2 + 20a*b + 15b^2) x \cosh(x)^2 + 2a*b + 4b^2) \sinh(x)^2 + 32((a^2 + 2a*b + b^2) \cosh(x)^4 + 4(a^2 + 2a*b + b^2) \cosh(x)^3 \sinh(x) + 6(a^2 + 2a*b + b^2) \cosh(x)^2 \sinh(x)^2 + 4(a^2 + 2a*b + b^2) \cosh(x) \sinh(x)^3 + (a^2 + 2a*b + b^2) \sinh(x)^4) \sqrt{(a+b)/a} \log((b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2a*b + b^2) \cosh(x)^2 + 2(3b^2 \cosh(x)^2 + 2a*b + b^2) \sinh(x)^2 + 8a^2 + 8a*b + b^2 + 4(b^2 \cosh(x)^3 + (2a*b + b^2) \cosh(x)) \sinh(x) + 4(a*b \cosh(x)^2 + 2a*b \cosh(x) \sinh(x) + a$


```

*b*sinh(x)^2 + 2*a^2 + a*b)*sqrt((a + b)/a))/(b*cosh(x)^4 + 4*b*cosh(x)*sin
h(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*
sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) - b^2 + 8*(b^
2*cosh(x)^7 - 6*(a*b + 2*b^2)*cosh(x)^5 + 4*(8*a^2 + 20*a*b + 15*b^2)*x*cos
h(x)^3 + 2*(a*b + 2*b^2)*cosh(x))*sinh(x))/(b^3*cosh(x)^4 + 4*b^3*cosh(x)^3
*sinh(x) + 6*b^3*cosh(x)^2*sinh(x)^2 + 4*b^3*cosh(x)*sinh(x)^3 + b^3*sinh(x
)^4), 1/64*(b^2*cosh(x)^8 + 8*b^2*cosh(x)*sinh(x)^7 + b^2*sinh(x)^8 - 8*(a*
b + 2*b^2)*cosh(x)^6 + 4*(7*b^2*cosh(x)^2 - 2*a*b - 4*b^2)*sinh(x)^6 + 8*(8
*a^2 + 20*a*b + 15*b^2)*x*cosh(x)^4 + 8*(7*b^2*cosh(x)^3 - 6*(a*b + 2*b^2)*
cosh(x))*sinh(x)^5 + 2*(35*b^2*cosh(x)^4 - 60*(a*b + 2*b^2)*cosh(x)^2 + 4*(
8*a^2 + 20*a*b + 15*b^2)*x)*sinh(x)^4 + 8*(7*b^2*cosh(x)^5 - 20*(a*b + 2*b^
2)*cosh(x)^3 + 4*(8*a^2 + 20*a*b + 15*b^2)*x*cosh(x))*sinh(x)^3 + 8*(a*b +
2*b^2)*cosh(x)^2 + 4*(7*b^2*cosh(x)^6 - 30*(a*b + 2*b^2)*cosh(x)^4 + 12*(8*
a^2 + 20*a*b + 15*b^2)*x*cosh(x)^2 + 2*a*b + 4*b^2)*sinh(x)^2 - 64*((a^2 +
2*a*b + b^2)*cosh(x)^4 + 4*(a^2 + 2*a*b + b^2)*cosh(x)^3*sinh(x) + 6*(a^2 +
2*a*b + b^2)*cosh(x)^2*sinh(x)^2 + 4*(a^2 + 2*a*b + b^2)*cosh(x)*sinh(x)^3
+ (a^2 + 2*a*b + b^2)*sinh(x)^4)*sqrt(-(a + b)/a)*arctan(1/2*(b*cosh(x)^2
+ 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-(a + b)/a)/(a + b)) -
b^2 + 8*(b^2*cosh(x)^7 - 6*(a*b + 2*b^2)*cosh(x)^5 + 4*(8*a^2 + 20*a*b + 15
*b^2)*x*cosh(x)^3 + 2*(a*b + 2*b^2)*cosh(x))*sinh(x))/(b^3*cosh(x)^4 + 4*b^
3*cosh(x)^3*sinh(x) + 6*b^3*cosh(x)^2*sinh(x)^2 + 4*b^3*cosh(x)*sinh(x)^3 +
b^3*sinh(x)^4)]

```

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^6(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

```
[In] integrate(sinh(x)**6/(a+b*cosh(x)**2),x)
```

```
[Out] Timed out
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 651 vs. 2(74) = 148.

Time = 0.31 (sec) , antiderivative size = 651, normalized size of antiderivative = 7.40

$$\begin{aligned}
 \int \frac{\sinh^6(x)}{a + b \cosh^2(x)} dx = & -\frac{15(2a + b) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab}} \\
 & + \frac{5 \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)a}} + \frac{3(2a + b)x}{2b^2} \\
 & + \frac{15x}{16b} - \frac{(4(2a + b)e^{(-2x)} - b)e^{(4x)}}{64b^2} - \frac{3e^{(2x)}}{16b} \\
 & + \frac{3e^{(-2x)}}{16b} + \frac{(4(2a + b)e^{(2x)} - b)e^{(-4x)}}{64b^2} \\
 & - \frac{3(2a + b) \log\left(\frac{be^{(4x)} + 2(2a + b)e^{(2x)} + b}{16b^2}\right)}{16b^2} \\
 & + \frac{3(2a + b) \log\left(\frac{2(2a + b)e^{(-2x)} + be^{(-4x)} + b}{16b^2}\right)}{16b^2} \\
 & - \frac{3(8a^2 + 8ab + b^2) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab^2}} \\
 & + \frac{3(8a^2 + 8ab + b^2) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab^2}} \\
 & + \frac{(16a^2 + 16ab + 3b^2)x}{8b^3} \\
 & - \frac{(16a^2 + 16ab + 3b^2) \log\left(\frac{be^{(4x)} + 2(2a + b)e^{(2x)} + b}{64b^3}\right)}{64b^3} \\
 & + \frac{(16a^2 + 16ab + 3b^2) \log\left(\frac{2(2a + b)e^{(-2x)} + be^{(-4x)} + b}{64b^3}\right)}{64b^3} \\
 & - \frac{(32a^3 + 48a^2b + 18ab^2 + b^3) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{128\sqrt{(a+b)ab^3}} \\
 & + \frac{(32a^3 + 48a^2b + 18ab^2 + b^3) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{128\sqrt{(a+b)ab^3}}
 \end{aligned}$$

[In] integrate(sinh(x)^6/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] -15/64*(2*a + b)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b) + 5/32*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a) + 3/2*(2*a + b)*x/b^2 + 15/16*x/b - 1/64*(4*(2*a + b)*e^(-2*x) - b)*e^(4*x)/b^2 - 3/16*e^(2*x)/b + 3/16*e^(-2*x)/b + 1/64*(4*(2*a + b)*

$$\begin{aligned}
& e^{(2*x) - b}*e^{(-4*x)}/b^2 - 3/16*(2*a + b)*\log(b*e^{(4*x) + 2*(2*a + b)*e^{(2*x) + b)}/b^2 + 3/16*(2*a + b)*\log(2*(2*a + b)*e^{(-2*x) + b*e^{(-4*x) + b)}/b^2 \\
& - 3/64*(8*a^2 + 8*a*b + b^2)*\log((b*e^{(2*x) + 2*a + b - 2*\sqrt{(a + b)*a})})/(b*e^{(2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})/(\sqrt{(a + b)*a}*b^2) + 3/64*(8*a^2 + 8*a*b + b^2)*\log((b*e^{(-2*x) + 2*a + b - 2*\sqrt{(a + b)*a})})/(b*e^{(-2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})/(\sqrt{(a + b)*a}*b^2) + 1/8*(16*a^2 + 16*a*b + 3*b^2)*x/b^3 \\
& - 1/64*(16*a^2 + 16*a*b + 3*b^2)*\log(b*e^{(4*x) + 2*(2*a + b)*e^{(2*x) + b)}/b^3 + 1/64*(16*a^2 + 16*a*b + 3*b^2)*\log(2*(2*a + b)*e^{(-2*x) + b*e^{(-4*x) + b)}/b^3 \\
& - 1/128*(32*a^3 + 48*a^2*b + 18*a*b^2 + b^3)*\log((b*e^{(2*x) + 2*a + b - 2*\sqrt{(a + b)*a})})/(b*e^{(2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})/(\sqrt{(a + b)*a}*b^3) + 1/128*(32*a^3 + 48*a^2*b + 18*a*b^2 + b^3)*\log((b*e^{(-2*x) + 2*a + b - 2*\sqrt{(a + b)*a})})/(b*e^{(-2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})/(\sqrt{(a + b)*a}*b^3)
\end{aligned}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(74) = 148$.

Time = 0.28 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.89

$$\begin{aligned}
& \int \frac{\sinh^6(x)}{a + b \cosh^2(x)} dx \\
& = \frac{be^{(4x)} - 8ae^{(2x)} - 16be^{(2x)}}{64b^2} + \frac{(8a^2 + 20ab + 15b^2)x}{8b^3} \\
& - \frac{(48a^2e^{(4x)} + 120abe^{(4x)} + 90b^2e^{(4x)} - 8abe^{(2x)} - 16b^2e^{(2x)} + b^2)e^{(-4x)}}{64b^3} \\
& - \frac{(a^3 + 3a^2b + 3ab^2 + b^3) \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - abb^3}}
\end{aligned}$$

[In] integrate(sinh(x)^6/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $1/64*(b*e^{(4*x)} - 8*a*e^{(2*x)} - 16*b*e^{(2*x)})/b^2 + 1/8*(8*a^2 + 20*a*b + 15*b^2)*x/b^3 - 1/64*(48*a^2*e^{(4*x)} + 120*a*b*e^{(4*x)} + 90*b^2*e^{(4*x)} - 8*a*b*e^{(2*x)} - 16*b^2*e^{(2*x)} + b^2)*e^{(-4*x)}/b^3 - (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*\arctan(1/2*(b*e^{(2*x)} + 2*a + b)/\sqrt{-a^2 - a*b})/(\sqrt{-a^2 - a*b})*b^3)$

Mupad [B] (verification not implemented)

Time = 2.57 (sec) , antiderivative size = 248, normalized size of antiderivative = 2.82

$$\int \frac{\sinh^6(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{e^{4x}}{64b} - \frac{e^{-4x}}{64b} + \frac{x(8a^2 + 20ab + 15b^2)}{8b^3} + \frac{e^{-2x}(a+2b)}{8b^2} - \frac{e^{2x}(a+2b)}{8b^2}$$

$$+ \frac{\ln\left(\frac{4(a+b)^5(2ab+8a^2e^{2x}+b^2e^{2x}+b^2+8abe^{2x})}{ab^8} - \frac{8(a+b)^{11/2}(b+4ae^{2x}+2be^{2x})}{\sqrt{ab^8}}\right)(a+b)^{5/2}}{2\sqrt{ab^3}}$$

$$- \frac{\ln\left(\frac{8(a+b)^{11/2}(b+4ae^{2x}+2be^{2x})}{\sqrt{ab^8}} + \frac{4(a+b)^5(2ab+8a^2e^{2x}+b^2e^{2x}+b^2+8abe^{2x})}{ab^8}\right)(a+b)^{5/2}}{2\sqrt{ab^3}}$$

[In] int(sinh(x)^6/(a + b*cosh(x)^2),x)

[Out] exp(4*x)/(64*b) - exp(-4*x)/(64*b) + (x*(20*a*b + 8*a^2 + 15*b^2))/(8*b^3) + (exp(-2*x)*(a + 2*b))/(8*b^2) - (exp(2*x)*(a + 2*b))/(8*b^2) + (log((4*(a + b)^5*(2*a*b + 8*a^2*exp(2*x) + b^2*exp(2*x) + b^2 + 8*a*b*exp(2*x)))/(a*b^8) - (8*(a + b)^(11/2)*(b + 4*a*exp(2*x) + 2*b*exp(2*x)))/(a^(1/2)*b^8))*(a + b)^(5/2))/(2*a^(1/2)*b^3) - (log((8*(a + b)^(11/2)*(b + 4*a*exp(2*x) + 2*b*exp(2*x)))/(a^(1/2)*b^8) + (4*(a + b)^5*(2*a*b + 8*a^2*exp(2*x) + b^2*exp(2*x) + b^2 + 8*a*b*exp(2*x)))/(a*b^8)))/(a^(1/2)*b^3) + (4*(a + b)^5*(2*a*b + 8*a^2*exp(2*x) + b^2*exp(2*x) + b^2 + 8*a*b*exp(2*x)))/(a*b^8)*(a + b)^(5/2))/(2*a^(1/2)*b^3)

3.14 $\int \frac{\sinh^4(x)}{a+b \cosh^2(x)} dx$

Optimal result	117
Rubi [A] (verified)	117
Mathematica [A] (verified)	119
Maple [B] (verified)	119
Fricas [B] (verification not implemented)	119
Sympy [F(-1)]	120
Maxima [B] (verification not implemented)	121
Giac [B] (verification not implemented)	121
Mupad [B] (verification not implemented)	122

Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{\sinh^4(x)}{a+b \cosh^2(x)} dx = -\frac{(2a+3b)x}{2b^2} + \frac{(a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^2}} + \frac{\cosh(x) \sinh(x)}{2b}$$

[Out] $-1/2*(2*a+3*b)*x/b^2+1/2*\cosh(x)*\sinh(x)/b+(a+b)^{(3/2)}*\operatorname{arctanh}(a^{(1/2)}*\tanh(x)/(a+b)^{(1/2)})/b^2/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.09 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3270, 425, 536, 212, 214}

$$\int \frac{\sinh^4(x)}{a+b \cosh^2(x)} dx = \frac{(a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^2}} - \frac{x(2a+3b)}{2b^2} + \frac{\sinh(x) \cosh(x)}{2b}$$

[In] $\operatorname{Int}[\operatorname{Sinh}[x]^4/(a+b*\operatorname{Cosh}[x]^2),x]$

[Out] $-1/2*((2*a+3*b)*x)/b^2 + ((a+b)^{(3/2)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tanh}[x])/(\operatorname{Sqrt}[a+b])])/(b^2) + (\operatorname{Cosh}[x]*\operatorname{Sinh}[x])/(2*b)$

Rule 212

$\operatorname{Int}[(a_0 + (b_0*x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{Gt}Q[a, 0] \ || \ \operatorname{Lt}Q[b, 0])$

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 425

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c -
a*d)), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c
+ d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n,
x], x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -
1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b,
c, d, n, p, q, x]
```

Rule 536

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(
n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x]
- Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b,
c, d, e, f, n}, x]
```

Rule 3270

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Sub
st[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e
+ f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1}{(1-x^2)^2(a-(a+b)x^2)} dx, x, \coth(x)\right) \\
&= \frac{\cosh(x) \sinh(x)}{2b} - \frac{\text{Subst}\left(\int \frac{-a-2b+(-a-b)x^2}{(1-x^2)(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{2b} \\
&= \frac{\cosh(x) \sinh(x)}{2b} + \frac{(a+b)^2 \text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{b^2} \\
&\quad - \frac{(2a+3b) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(x)\right)}{2b^2} \\
&= -\frac{(2a+3b)x}{2b^2} + \frac{(a+b)^{3/2} \arctanh\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab^2}} + \frac{\cosh(x) \sinh(x)}{2b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = \frac{-4ax - 6bx + \frac{4(a+b)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}} + b \sinh(2x)}{4b^2}$$

[In] Integrate[Sinh[x]^4/(a + b*Cosh[x]^2),x]

[Out] $(-4*a*x - 6*b*x + (4*(a + b)^{(3/2)}*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a + b]*Sinh[2*x])/(4*b^2)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(47) = 94.

Time = 26.57 (sec) , antiderivative size = 185, normalized size of antiderivative = 3.14

method	result
default	$\frac{2(a^2+2ab+b^2) \left(\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} - \frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 - 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} \right)}{b^2} - \frac{1}{2b(\tanh\left(\frac{x}{2}\right)+1)^2} + \frac{1}{2b(\tanh\left(\frac{x}{2}\right)-1)^2}$
risch	$-\frac{ax}{b^2} - \frac{3x}{2b} + \frac{e^{2x}}{8b} - \frac{e^{-2x}}{8b} + \frac{\sqrt{a(a+b)} \ln\left(e^{2x} - \frac{2\sqrt{a(a+b)}-2a-b}{b}\right)}{2b^2} + \frac{\sqrt{a(a+b)} \ln\left(e^{2x} - \frac{2\sqrt{a(a+b)}-2a-b}{b}\right)}{2ab} - \frac{\sqrt{a(a+b)} \ln\left(e^{2x} - \frac{2\sqrt{a(a+b)}-2a-b}{b}\right)}{2ab}$

[In] int(sinh(x)^4/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] $2/b^2*(a^2+2*a*b+b^2)*(1/4/a^{(1/2)/(a+b)^{(1/2)}*ln((a+b)^{(1/2)}*tanh(1/2*x)^2+2*tanh(1/2*x)*a^{(1/2)+(a+b)^{(1/2)})-1/4/a^{(1/2)/(a+b)^{(1/2)}*ln((a+b)^{(1/2)}*tanh(1/2*x)^2-2*tanh(1/2*x)*a^{(1/2)+(a+b)^{(1/2)})})-1/2/b/(tanh(1/2*x)+1)^2+1/2/b/(tanh(1/2*x)+1)+1/2/b^2*(-2*a-3*b)*ln(tanh(1/2*x)+1)+1/2/b/(tanh(1/2*x)-1)^2+1/2/b/(tanh(1/2*x)-1)+1/2*(2*a+3*b)/b^2*ln(tanh(1/2*x)-1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 196 vs. 2(47) = 94.

Time = 0.28 (sec) , antiderivative size = 568, normalized size of antiderivative = 9.63

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = \frac{b \cosh^4(x) + 4b \cosh(x) \sinh^3(x) + b \sinh^4(x) - 4(2a + 3b)x \cosh^2(x) + 2(3b \cosh(x)^2 - 2(2a + 3b) \sinh(x)) \operatorname{arctanh}\left(\frac{\sinh(x)}{\cosh(x)}\right)}{4b^2}$$

[In] integrate(sinh(x)^4/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/8*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 4*(2*a + 3*b)*x*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*(2*a + 3*b)*x)*sinh(x)^2 + 4*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2)*sqrt((a + b)/a)*log((b^2*cosh(x)^4 + 4*b^2*cosh(x)*sinh(x)^3 + b^2*sinh(x)^4 + 2*(2*a*b + b^2)*cosh(x)^2 + 2*(3*b^2*cosh(x)^2 + 2*a*b + b^2)*sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*cosh(x)^3 + (2*a*b + b^2)*cosh(x))*sinh(x) - 4*(a*b*cosh(x)^2 + 2*a*b*cosh(x)*sinh(x) + a*b*sinh(x)^2 + 2*a^2 + a*b)*sqrt((a + b)/a))/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 4*(b*cosh(x)^3 - 2*(2*a + 3*b)*x*cosh(x))*sinh(x) - b)/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2), 1/8*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 4*(2*a + 3*b)*x*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*(2*a + 3*b)*x)*sinh(x)^2 + 8*((a + b)*cosh(x)^2 + 2*(a + b)*cosh(x)*sinh(x) + (a + b)*sinh(x)^2)*sqrt(-(a + b)/a)*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-(a + b)/a)/(a + b)) + 4*(b*cosh(x)^3 - 2*(2*a + 3*b)*x*cosh(x))*sinh(x) - b)/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2)]

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(sinh(x)**4/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 348 vs. 2(47) = 94.

Time = 0.29 (sec) , antiderivative size = 348, normalized size of antiderivative = 5.90

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = \frac{(2a + b) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)ab}} - \frac{3 \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{16\sqrt{(a+b)a}} - \frac{(2a + b)x}{b^2} - \frac{x}{b} + \frac{e^{(2x)}}{8b} - \frac{e^{(-2x)}}{8b} + \frac{(2a + b) \log\left(\frac{be^{(4x)} + 2(2a + b)e^{(2x)} + b}{2(2a + b)e^{(-2x)} + be^{(-4x)} + b}\right)}{8b^2} - \frac{(8a^2 + 8ab + b^2) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)ab^2}} + \frac{(8a^2 + 8ab + b^2) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)ab^2}}$$

[In] integrate(sinh(x)^4/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/4*(2*a + b)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b) - 3/16*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a) - (2*a + b)*x/b^2 - x/b + 1/8*e^(2*x)/b - 1/8*e^(-2*x)/b + 1/8*(2*a + b)*log(b*e^(4*x) + 2*(2*a + b)*e^(2*x) + b)/b^2 - 1/8*(2*a + b)*log(2*(2*a + b)*e^(-2*x) + b*e^(-4*x) + b)/b^2 + 1/32*(8*a^2 + 8*a*b + b^2)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b^2) - 1/32*(8*a^2 + 8*a*b + b^2)*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b^2)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 103 vs. 2(47) = 94.

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.75

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = -\frac{(2a + 3b)x}{2b^2} + \frac{e^{(2x)}}{8b} + \frac{(4ae^{(2x)} + 6be^{(2x)} - b)e^{(-2x)}}{8b^2} + \frac{(a^2 + 2ab + b^2) \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - abb^2}}$$

[In] integrate(sinh(x)^4/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $-1/2*(2*a + 3*b)*x/b^2 + 1/8*e^{(2*x)}/b + 1/8*(4*a*e^{(2*x)} + 6*b*e^{(2*x)} - b)*e^{(-2*x)}/b^2 + (a^2 + 2*a*b + b^2)*\arctan(1/2*(b*e^{(2*x)} + 2*a + b)/\sqrt{-a^2 - a*b})/(\sqrt{-a^2 - a*b})*b^2$

Mupad [B] (verification not implemented)

Time = 2.12 (sec) , antiderivative size = 146, normalized size of antiderivative = 2.47

$$\int \frac{\sinh^4(x)}{a + b \cosh^2(x)} dx = \frac{e^{2x}}{8b} - \frac{e^{-2x}}{8b} - \frac{x(2a + 3b)}{2b^2} + \frac{\ln\left(-\frac{4e^{2x}(a+b)^2}{b^3} - \frac{2(a+b)^{3/2}(b+2ae^{2x}+be^{2x})}{\sqrt{ab^3}}\right)(a+b)^{3/2}}{2\sqrt{ab^2}} - \frac{\ln\left(\frac{2(a+b)^{3/2}(b+2ae^{2x}+be^{2x})}{\sqrt{ab^3}} - \frac{4e^{2x}(a+b)^2}{b^3}\right)(a+b)^{3/2}}{2\sqrt{ab^2}}$$

[In] int(sinh(x)^4/(a + b*cosh(x)^2),x)

[Out] $\exp(2*x)/(8*b) - \exp(-2*x)/(8*b) - (x*(2*a + 3*b))/(2*b^2) + (\log(- (4*\exp(2*x)*(a + b)^2)/b^3 - (2*(a + b)^{(3/2)}*(b + 2*a*\exp(2*x) + b*\exp(2*x))))/(a^{(1/2)*b^3})*(a + b)^{(3/2)}/(2*a^{(1/2)*b^2}) - (\log((2*(a + b)^{(3/2)}*(b + 2*a*\exp(2*x) + b*\exp(2*x)))/(a^{(1/2)*b^3}) - (4*\exp(2*x)*(a + b)^2)/b^3)*(a + b)^{(3/2)}/(2*a^{(1/2)*b^2})$

3.15 $\int \frac{\sinh^2(x)}{a+b \cosh^2(x)} dx$

Optimal result	123
Rubi [A] (verified)	123
Mathematica [A] (verified)	124
Maple [B] (verified)	125
Fricas [A] (verification not implemented)	125
Sympy [F(-1)]	126
Maxima [B] (verification not implemented)	126
Giac [A] (verification not implemented)	126
Mupad [B] (verification not implemented)	127

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \frac{\sinh^2(x)}{a+b \cosh^2(x)} dx = \frac{x}{b} - \frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab}}$$

[Out] $x/b - \operatorname{arctanh}(a^{1/2} \tanh(x) / (a+b)^{1/2}) * (a+b)^{1/2} / b a^{1/2}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3270, 400, 212, 214}

$$\int \frac{\sinh^2(x)}{a+b \cosh^2(x)} dx = \frac{x}{b} - \frac{\sqrt{a+b} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab}}$$

[In] `Int[Sinh[x]^2/(a + b*Cosh[x]^2),x]`

[Out] $x/b - (\operatorname{Sqrt}[a + b] * \operatorname{ArcTanh}[(\operatorname{Sqrt}[a] * \operatorname{Tanh}[x]) / \operatorname{Sqrt}[a + b]]) / (\operatorname{Sqrt}[a] * b)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 400

```
Int[1/(((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_))), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0]
```

Rule 3270

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1}{(1-x^2)(a-(a+b)x^2)} dx, x, \coth(x)\right) \\ &= \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(x)\right)}{b} - \frac{(a+b)\text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{b} \\ &= \frac{x}{b} - \frac{\sqrt{a+b}\text{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{ab}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int \frac{\sinh^2(x)}{a+b\cosh^2(x)} dx = \frac{x - \frac{\sqrt{a+b}\text{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}}}{b}$$

```
[In] Integrate[Sinh[x]^2/(a + b*Cosh[x]^2),x]
```

```
[Out] (x - (Sqrt[a + b]*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a])/b
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(31) = 62$.

Time = 0.73 (sec) , antiderivative size = 88, normalized size of antiderivative = 2.26

method	result
risch	$\frac{x}{b} + \frac{\sqrt{a(a+b)} \ln\left(e^{2x + \frac{2\sqrt{a(a+b)+2a+b}}{b}}\right)}{2ab} - \frac{\sqrt{a(a+b)} \ln\left(e^{2x - \frac{2\sqrt{a(a+b)-2a-b}}{b}}\right)}{2ab}$
default	$\frac{2(a+b) \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 - 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} \right)}{b} + \frac{\ln\left(\tanh\left(\frac{x}{2}\right)+1\right)}{b} - \frac{\ln\left(\tanh\left(\frac{x}{2}\right)-1\right)}{b}$

[In] `int(sinh(x)^2/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $x/b + 1/2/a * (a*(a+b))^{1/2} / b * \ln(\exp(2*x) + (2*(a*(a+b))^{1/2} + 2*a+b)/b) - 1/2/a * (a*(a+b))^{1/2} / b * \ln(\exp(2*x) - (2*(a*(a+b))^{1/2} - 2*a-b)/b)$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 300, normalized size of antiderivative = 7.69

$$\int \frac{\sinh^2(x)}{a + b \cosh^2(x)} dx$$

$$= \left[\frac{\sqrt{\frac{a+b}{a}} \log\left(\frac{b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2ab+b^2) \cosh(x)^2 + 2(3b^2 \cosh(x)^2 + 2ab+b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4a^2}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 2ab+b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4a^2}\right)}{2b} \right.$$

$$\left. - \frac{\sqrt{-\frac{a+b}{a}} \arctan\left(\frac{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + 2a+b) \sqrt{-\frac{a+b}{a}}}{2(a+b)}\right) - x}{b} \right]$$

[In] `integrate(sinh(x)^2/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/2*(\sqrt{(a+b)/a})*\log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x)^2 + 2*(3*b^2*\cosh(x)^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) + 4*(a*b*\cosh(x)^2 + 2*a*b*\cosh(x)*\sinh(x) + a*b*\sinh(x)^2 + 2*a^2 + a*b)*\sqrt{(a+b)/a})/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x))^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 2*x)/b, -(\sqrt{-(a+b)/a})*\arctan($

$1/2*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{-(a + b)/a}/(a + b) - x)/b]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\sinh^2(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(sinh(x)**2/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(31) = 62.

Time = 0.29 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.08

$$\int \frac{\sinh^2(x)}{a + b \cosh^2(x)} dx = -\frac{(2a + b) \log\left(\frac{be^{2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)ab}} + \frac{\log\left(\frac{be^{-2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{-2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)a}} + \frac{x}{b}$$

[In] integrate(sinh(x)^2/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-1/4*(2*a + b)*\log((b*e^{2*x} + 2*a + b - 2*\sqrt{(a + b)*a})/(b*e^{2*x} + 2*a + b + 2*\sqrt{(a + b)*a}))/(\sqrt{(a + b)*a}*b) + 1/4*\log((b*e^{-2*x} + 2*a + b - 2*\sqrt{(a + b)*a})/(b*e^{-2*x} + 2*a + b + 2*\sqrt{(a + b)*a}))/\sqrt{(a + b)*a} + x/b$

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 52, normalized size of antiderivative = 1.33

$$\int \frac{\sinh^2(x)}{a + b \cosh^2(x)} dx = -\frac{(a + b) \arctan\left(\frac{be^{2x} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - ab}b} + \frac{x}{b}$$

[In] integrate(sinh(x)^2/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $-(a + b)*\arctan(1/2*(b*e^{2*x} + 2*a + b)/\sqrt{-a^2 - a*b})/(\sqrt{-a^2 - a*b}*b) + x/b$

Mupad [B] (verification not implemented)

Time = 0.25 (sec) , antiderivative size = 79, normalized size of antiderivative = 2.03

$$\int \frac{\sinh^2(x)}{a + b \cosh^2(x)} dx = \frac{x}{b} + \frac{\operatorname{atan}\left(\frac{\sqrt{-ab^2}}{2a\sqrt{a+b}} + \frac{\sqrt{-ab^2}}{b\sqrt{a+b}} + \frac{e^{2x}\sqrt{-ab^2}}{2a\sqrt{a+b}}\right) \sqrt{a+b}}{\sqrt{-ab^2}}$$

```
[In] int(sinh(x)^2/(a + b*cosh(x)^2),x)
```

```
[Out] x/b + (atan((-a*b^2)^(1/2)/(2*a*(a + b)^(1/2)) + (-a*b^2)^(1/2)/(b*(a + b)^(1/2)) + (exp(2*x)*(-a*b^2)^(1/2))/(2*a*(a + b)^(1/2)))*(a + b)^(1/2)/(-a*b^2)^(1/2)
```

3.16 $\int \frac{1}{a+b \cosh^2(x)} dx$

Optimal result	128
Rubi [A] (verified)	128
Mathematica [A] (verified)	129
Maple [B] (verified)	129
Fricas [B] (verification not implemented)	130
Sympy [B] (verification not implemented)	130
Maxima [B] (verification not implemented)	136
Giac [A] (verification not implemented)	137
Mupad [B] (verification not implemented)	137

Optimal result

Integrand size = 10, antiderivative size = 29

$$\int \frac{1}{a+b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[Out] $\operatorname{arctanh}(a^{(1/2)}*\tanh(x)/(a+b)^{(1/2)})/a^{(1/2)}/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3260, 214}

$$\int \frac{1}{a+b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[In] $\operatorname{Int}[(a + b*\operatorname{Cosh}[x]^2)^{-1}, x]$

[Out] $\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tanh}[x])/\operatorname{Sqrt}[a + b]]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[a + b])$

Rule 214

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b]$

Rule 3260

$\operatorname{Int}[(a_ + (b_)*\sin[(e_ + (f_)*(x_)]^2)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{\{ff = \operatorname{FreeFactors}[\operatorname{Tan}[e + f*x], x]\}, \operatorname{Dist}[ff/f, \operatorname{Subst}[\operatorname{Int}[1/(a + (a + b)*ff^2*x^2$

), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{a - (a + b)x^2} dx, x, \coth(x)\right) \\ &= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.47 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[In] Integrate[(a + b*Cosh[x]^2)^(-1),x]

[Out] ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]]/(Sqrt[a]*Sqrt[a + b])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(21) = 42.

Time = 0.11 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.69

method	result	size
default	$\frac{\ln\left(\sqrt{a+b}\tanh\left(\frac{x}{2}\right)^2 + 2\tanh\left(\frac{x}{2}\right)\sqrt{a}\sqrt{a+b}\right)}{2\sqrt{a}\sqrt{a+b}} - \frac{\ln\left(\sqrt{a+b}\tanh\left(\frac{x}{2}\right)^2 - 2\tanh\left(\frac{x}{2}\right)\sqrt{a}\sqrt{a+b}\right)}{2\sqrt{a}\sqrt{a+b}}$	78
risch	$\frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} - 2a^2 - 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}} - \frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} + 2a^2 + 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}}$	128

[In] int(1/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/2/a^(1/2)/(a+b)^(1/2)*ln((a+b)^(1/2)*tanh(1/2*x)^2+2*tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))-1/2/a^(1/2)/(a+b)^(1/2)*ln((a+b)^(1/2)*tanh(1/2*x)^2-2*tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(21) = 42.

Time = 0.27 (sec) , antiderivative size = 293, normalized size of antiderivative = 10.10

$$\int \frac{1}{a + b \cosh^2(x)} dx$$

$$= \left[\frac{\log \left(\frac{b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2ab + b^2) \cosh(x)^2 + 2(3b^2 \cosh(x)^2 + 2ab + b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4(b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2a + b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 2a + b) \sinh(x)^2)}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a + b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 2a + b) \sinh(x)^2} \right)}{2\sqrt{a^2 + ab}} \right]$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/2*log((b^2*cosh(x)^4 + 4*b^2*cosh(x)*sinh(x)^3 + b^2*sinh(x)^4 + 2*(2*a*b + b^2)*cosh(x)^2 + 2*(3*b^2*cosh(x)^2 + 2*a*b + b^2)*sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*cosh(x)^3 + (2*a*b + b^2)*cosh(x))*sinh(x) - 4*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(a^2 + a*b))/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b))/sqrt(a^2 + a*b), sqrt(-a^2 - a*b)*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-a^2 - a*b)/(a^2 + a*b))/(a^2 + a*b)]

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10924 vs. 2(27) = 54.

Time = 24.04 (sec) , antiderivative size = 10924, normalized size of antiderivative = 376.69

$$\int \frac{1}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cosh(x)**2),x)

[Out] Piecewise((zoo*tanh(x/2)/(tanh(x/2)**2 + 1), Eq(a, 0) & Eq(b, 0)), (-tanh(x/2)/(2*b) - 1/(2*b*tanh(x/2)), Eq(a, -b)), (2*tanh(x/2)/(b*(tanh(x/2)**2 + 1)), Eq(a, 0)), (-a**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))

$$\begin{aligned}
& + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a \\
& + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq \\
& rt(-a*b)/(a + b))) + a**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b) \\
&)*log(sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a* \\
& *4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a \\
& + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sq \\
& rt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a* \\
& *3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a \\
& + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/ \\
& (a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(\\
& a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(\\
& a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/ \\
& (a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2* \\
& sqrt(-a*b)/(a + b))) - a**3*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + \\
& b))*log(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2 \\
& *a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b \\
& /(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2 \\
& *sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8 \\
& *a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - \\
& b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b \\
&)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sq \\
& rt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt \\
& (a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + \\
& 2*sqrt(-a*b)/(a + b))) + a**3*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b))*log(sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(\\
& (2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - \\
& b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - \\
& 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + \\
& 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(\\
& a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) \\
& - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a \\
& *b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))* \\
& sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sq \\
& rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) \\
& + 2*sqrt(-a*b)/(a + b))) + 10*a**2*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(- \\
& a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan \\
& h(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(\\
& a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(\\
& a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b \\
&))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a \\
& /(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2 \\
& *sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(
\end{aligned}$$

$$\begin{aligned}
& 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt \\
& (a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + 5*a*b**2*sqrt(a/(a + b) - \\
& b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a* \\
& b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a* \\
& b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b* \\
& sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - \\
& 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - \\
& 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a \\
& + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + \\
& b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + \\
& b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b \\
&))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + 3*a*b**2*sqrt(a/(a \\
& + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) - \\
& 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*s \\
& qrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10* \\
& a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - \\
& b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a \\
& + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sq \\
& rt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - \\
& b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b \\
&))/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/ \\
& (a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) - 3*a*b**2*sq \\
& rt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))*log(sqrt(a/(a + b) - b/(a + \\
& b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) \\
& - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) \\
& - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + \\
& b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) \\
& - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a* \\
& b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b \\
&))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a \\
& + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq \\
& rt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(\\
& -a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + 10*a*b \\
& *sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a \\
& + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq \\
& rt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + \\
& b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b) \\
& *sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*s \\
& qrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a \\
& *b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b \\
& /(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(
\end{aligned}$$

$$\begin{aligned} & a + b) - 2\sqrt{-ab}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} \\ & + b)) - 10ab\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)} \\ &))\log(\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} + \tanh(x/2))/(2a \\ & a^{*4}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/ \\ & (a + b) + 2\sqrt{-ab}/(a + b)} - 10a^{*3}b\sqrt{a/(a + b) - b/(a + b) - 2 \\ & \sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} + 8 \\ & a^{*3}\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(\\ & a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 10a^{*2}b^{*2}\sqrt{a/(a + b) - \\ & b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab} \\ & / (a + b)} + 2ab^{*3}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{ \\ & t(a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)) - 8ab^{*2}\sqrt{-ab}\sqrt{ \\ & a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + \\ & 2\sqrt{-ab}/(a + b)) - 2ab\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{ \\ & rt(-ab)/(a + b)}\log(-\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)} + \\ & \tanh(x/2))/(2a^{*4}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{ \\ & a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 10a^{*3}b\sqrt{a/(a + b) - \\ & b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab} \\ &)/(a + b)} + 8a^{*3}\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a \\ & + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 10a^{*2}b^{*2}\sqrt{ \\ & rt(a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b))\sqrt{a/(a + b) - b/(a + b) \\ & + 2\sqrt{-ab}/(a + b)} + 2ab^{*3}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab} \\ & b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 8ab^{*2}\sqrt{ \\ & rt(-ab)\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) \\ & - b/(a + b) + 2\sqrt{-ab}/(a + b))} + 2ab\sqrt{-ab}\sqrt{a/(a + b) - b \\ & / (a + b) + 2\sqrt{-ab}/(a + b)}\log(\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab} \\ & *b)/(a + b)} + \tanh(x/2))/(2a^{*4}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab} \\ & / (a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 10a^{*3}b\sqrt{ \\ & rt(a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b))\sqrt{a/(a + b) - b/(a + b) \\ & + 2\sqrt{-ab}/(a + b)} + 8a^{*3}\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2 \\ & *sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} - 1 \\ & 0a^{*2}b^{*2}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + \\ & b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} + 2ab^{*3}\sqrt{a/(a + b) - b/(a + b) \\ &) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b) \\ &) - 8ab^{*2}\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{ \\ & sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b))} - b^{*2}\sqrt{-ab}\sqrt{ \\ & a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\log(-\sqrt{a/(a + b) - b/(a + \\ & b) + 2\sqrt{-ab}/(a + b)} + \tanh(x/2))/(2a^{*4}\sqrt{a/(a + b) - b/(a + b) \\ & - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} \\ & - 10a^{*3}b\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + \\ & b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} + 8a^{*3}\sqrt{-ab}\sqrt{a/(a + b) - \\ & b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab} \\ &)/(a + b)} - 10a^{*2}b^{*2}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-ab}/(a + b) \\ &)\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-ab}/(a + b)} + 2ab^{*3}\sqrt{a/(a + \\ & b) - b/(a + b) - 2\sqrt{-ab}/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{ \\ & (-ab)/(a + b)} - 8ab^{*2}\sqrt{-ab}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-} \end{aligned}$$

```

a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + b**2*sq
rt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(a/(a +
b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b)
- b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a
*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))
*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sq
rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b)
+ 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(
-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**
3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a
+ b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a +
b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b
))) + b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))*lo
g(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*
sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a +
b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(
-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*
sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b
) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a
+ b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a +
b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(
a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a
+ b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq
rt(-a*b)/(a + b))) - b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*
b)/(a + b))*log(sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x
/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a +
b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a +
b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a +
b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*
sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a
+ b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq
rt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a +
b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*
b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a
+ b) + 2*sqrt(-a*b)/(a + b))), True))

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(21) = 42.

Time = 0.31 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.83

$$\int \frac{1}{a + b \cosh^2(x)} dx = -\frac{\log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{2\sqrt{(a+b)a}}$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-1/2*\log((b*e^{(-2*x)} + 2*a + b - 2*sqrt((a + b)*a))/(b*e^{(-2*x)} + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.34

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\arctan\left(\frac{be^{(2x)}+2a+b}{2\sqrt{-a^2-ab}}\right)}{\sqrt{-a^2-ab}}$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $\arctan(1/2*(b*e^{(2*x)} + 2*a + b)/sqrt(-a^2 - a*b))/sqrt(-a^2 - a*b)$

Mupad [B] (verification not implemented)

Time = 0.40 (sec) , antiderivative size = 267, normalized size of antiderivative = 9.21

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\operatorname{atan}\left(\frac{b^2 e^{2x} (-a^2 - ba)^{3/2} \left(\frac{4(4a+2b)(8a^3+12a^2b+4ab^2)}{b^5(-a^2-ba)^{3/2}\sqrt{-a(a+b)}} + \frac{2(8a^2+8ab+b^2)(8a^2\sqrt{-a^2-ba}+b^2\sqrt{-a^2-ba}+8ab\sqrt{-a^2-ba})}{ab^5(a+b)(-a^2-ba)^{3/2}} \right)}{4} + \frac{(2a^2b)}{b^3}\right)}{\sqrt{-a^2 - ba}}$$

[In] int(1/(a + b*cosh(x)^2),x)

[Out] $-\operatorname{atan}((b^2*\exp(2*x)*(-a*b - a^2)^{(3/2)}*((4*(4*a + 2*b)*(4*a*b^2 + 12*a^2*b + 8*a^3))/(b^5*(-a*b - a^2)^{(3/2)}*(-a*(a + b))^{(1/2)}) + (2*(8*a*b + 8*a^2 + b^2)*(8*a^2*(-a*b - a^2)^{(1/2)} + b^2*(-a*b - a^2)^{(1/2)} + 8*a*b*(-a*b - a^2)^{(1/2)}))/(a*b^5*(a + b)*(-a*b - a^2)^{(3/2)}))/4 + ((2*a*b^2 + 2*a^2*b)*(4*a + 2*b))/(b^3*(-a*(a + b))^{(1/2)}) + ((b^2*(-a*b - a^2)^{(1/2)} + 2*a*b*(-a*b - a^2)^{(1/2)})*(8*a*b + 8*a^2 + b^2))/(2*a*b^3*(a + b)))/(-a*b - a^2)^{(1/2)}$

3.17 $\int \frac{\operatorname{csch}^4(x)}{a+b \cosh^2(x)} dx$

Optimal result	138
Rubi [A] (verified)	138
Mathematica [A] (verified)	139
Maple [B] (verified)	140
Fricas [B] (verification not implemented)	140
Sympy [F]	141
Maxima [B] (verification not implemented)	142
Giac [F]	142
Mupad [B] (verification not implemented)	142

Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{\operatorname{csch}^4(x)}{a+b \cosh^2(x)} dx = \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{5/2}} + \frac{(a+2b) \operatorname{coth}(x)}{(a+b)^2} - \frac{\operatorname{coth}^3(x)}{3(a+b)}$$

[Out] (a+2*b)*coth(x)/(a+b)^2-1/3*coth(x)^3/(a+b)+b^2*arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))/(a+b)^(5/2)/a^(1/2)

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3270, 398, 214}

$$\int \frac{\operatorname{csch}^4(x)}{a+b \cosh^2(x)} dx = \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{5/2}} - \frac{\operatorname{coth}^3(x)}{3(a+b)} + \frac{(a+2b) \operatorname{coth}(x)}{(a+b)^2}$$

[In] Int[Csch[x]^4/(a + b*Cosh[x]^2), x]

[Out] (b^2*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]]/(Sqrt[a]*(a + b)^(5/2)) + ((a + 2*b)*Coth[x])/(a + b)^2 - Coth[x]^3/(3*(a + b)))

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3270

```
Int[cos[(e_.) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Sub
st[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e
+ f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst} \left(\int \frac{(1-x^2)^2}{a-(a+b)x^2} dx, x, \coth(x) \right) \\
&= \text{Subst} \left(\int \left(\frac{a+2b}{(a+b)^2} - \frac{x^2}{a+b} + \frac{b^2}{(a+b)^2(a-(a+b)x^2)} \right) dx, x, \coth(x) \right) \\
&= \frac{(a+2b)\coth(x)}{(a+b)^2} - \frac{\coth^3(x)}{3(a+b)} + \frac{b^2 \text{Subst} \left(\int \frac{1}{a-(a+b)x^2} dx, x, \coth(x) \right)}{(a+b)^2} \\
&= \frac{b^2 \operatorname{arctanh} \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}} \right)}{\sqrt{a}(a+b)^{5/2}} + \frac{(a+2b)\coth(x)}{(a+b)^2} - \frac{\coth^3(x)}{3(a+b)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{csch}^4(x)}{a+b \operatorname{cosh}^2(x)} dx = \frac{b^2 \operatorname{arctanh} \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}} \right)}{\sqrt{a}(a+b)^{5/2}} - \frac{\coth(x) (-2a - 5b + (a+b) \operatorname{csch}^2(x))}{3(a+b)^2}$$

```
[In] Integrate[Csch[x]^4/(a + b*Cosh[x]^2), x]
```

```
[Out] (b^2*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(Sqrt[a]*(a + b)^(5/2)) - (Cot
h[x]*(-2*a - 5*b + (a + b)*Csch[x]^2))/(3*(a + b)^2)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 161 vs. $2(49) = 98$.

Time = 14.04 (sec) , antiderivative size = 162, normalized size of antiderivative = 2.75

method	result
default	$-\frac{\frac{a \tanh\left(\frac{x}{2}\right)^3}{3} + \frac{b \tanh\left(\frac{x}{2}\right)^3}{3} - 3a \tanh\left(\frac{x}{2}\right) - 7b \tanh\left(\frac{x}{2}\right)}{8(a+b)^2} - \frac{1}{24(a+b) \tanh\left(\frac{x}{2}\right)^3} - \frac{-3a-7b}{8(a+b)^2 \tanh\left(\frac{x}{2}\right)} - \frac{2b^2 \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)\right)^2 + 2 \tanh\left(\frac{x}{2}\right)}{4\sqrt{a+b}} \right)}{8(a+b)^2 \tanh\left(\frac{x}{2}\right)}$
risch	$-\frac{2(-3be^{4x}+6ae^{2x}+12be^{2x}-2a-5b)}{3(e^{2x}-1)^3(a+b)^2} + \frac{b^2 \ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab}+b\sqrt{a^2+ab}-2a^2-2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}(a+b)^2} - \frac{b^2 \ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab}+b\sqrt{a^2+ab}+2a^2+2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}(a+b)^2}$

[In] `int(csch(x)^4/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $-1/8/(a+b)^2*(1/3*a*\tanh(1/2*x)^3+1/3*b*\tanh(1/2*x)^3-3*a*\tanh(1/2*x)-7*b*\tanh(1/2*x))-1/24/(a+b)/\tanh(1/2*x)^3-1/8/(a+b)^2*(-3*a-7*b)/\tanh(1/2*x)-2*b^2/(a+b)^2*(-1/4/a^{1/2}/(a+b)^{1/2}*\ln((a+b)^{1/2}*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^{1/2}+(a+b)^{1/2}))+1/4/a^{1/2}/(a+b)^{1/2}*\ln((a+b)^{1/2}*\tanh(1/2*x)^2-2*\tanh(1/2*x)*a^{1/2}+(a+b)^{1/2}))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 857 vs. $2(49) = 98$.

Time = 0.28 (sec) , antiderivative size = 1875, normalized size of antiderivative = 31.78

$$\int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] `integrate(csch(x)^4/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/6*(12*(a^2*b + a*b^2)*\cosh(x)^4 + 48*(a^2*b + a*b^2)*\cosh(x)*\sinh(x)^3 + 12*(a^2*b + a*b^2)*\sinh(x)^4 + 8*a^3 + 28*a^2*b + 20*a*b^2 - 24*(a^3 + 3*a^2*b + 2*a*b^2)*\cosh(x)^2 - 24*(a^3 + 3*a^2*b + 2*a*b^2 - 3*(a^2*b + a*b^2)*\cosh(x)^2)*\sinh(x)^2 + 3*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 - 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 - b^2)*\sinh(x)^4 + 3*b^2*\cosh(x)^2 + 4*(5*b^2*\cosh(x)^3 - 3*b^2*\cosh(x))*\sinh(x)^3 + 3*(5*b^2*\cosh(x)^4 - 6*b^2*\cosh(x)^2 + b^2)*\sinh(x)^2 - b^2 + 6*(b^2*\cosh(x)^5 - 2*b^2*\cosh(x))^3 + b^2*\cosh(x))*\sinh(x))*\sqrt{a^2 + a*b}*\log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x)^2 + 2*(3*b^2*\cosh(x))^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) - 4*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{a^2 + a*b})/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 48*((a^2*b + a*b^2)*co$

```

sh(x)^3 - (a^3 + 3*a^2*b + 2*a*b^2)*cosh(x))*sinh(x))/((a^4 + 3*a^3*b + 3*a
^2*b^2 + a*b^3)*cosh(x)^6 + 6*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)*s
inh(x)^5 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*sinh(x)^6 - 3*(a^4 + 3*a^3*b
+ 3*a^2*b^2 + a*b^3)*cosh(x)^4 - 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - 5*
(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^2)*sinh(x)^4 - a^4 - 3*a^3*b -
3*a^2*b^2 - a*b^3 + 4*(5*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^3 - 3*
(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x))*sinh(x)^3 + 3*(a^4 + 3*a^3*b +
3*a^2*b^2 + a*b^3)*cosh(x)^2 + 3*(5*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*co
sh(x)^4 + a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - 6*(a^4 + 3*a^3*b + 3*a^2*b^2
+ a*b^3)*cosh(x)^2)*sinh(x)^2 + 6*((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh
(x)^5 - 2*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^3 + (a^4 + 3*a^3*b +
3*a^2*b^2 + a*b^3)*cosh(x))*sinh(x)), 1/3*(6*(a^2*b + a*b^2)*cosh(x)^4 + 24
*(a^2*b + a*b^2)*cosh(x)*sinh(x)^3 + 6*(a^2*b + a*b^2)*sinh(x)^4 + 4*a^3 +
14*a^2*b + 10*a*b^2 - 12*(a^3 + 3*a^2*b + 2*a*b^2)*cosh(x)^2 - 12*(a^3 + 3*
a^2*b + 2*a*b^2 - 3*(a^2*b + a*b^2)*cosh(x)^2)*sinh(x)^2 + 3*(b^2*cosh(x)^6
+ 6*b^2*cosh(x)*sinh(x)^5 + b^2*sinh(x)^6 - 3*b^2*cosh(x)^4 + 3*(5*b^2*cos
h(x)^2 - b^2)*sinh(x)^4 + 3*b^2*cosh(x)^2 + 4*(5*b^2*cosh(x)^3 - 3*b^2*cosh
(x))*sinh(x)^3 + 3*(5*b^2*cosh(x)^4 - 6*b^2*cosh(x)^2 + b^2)*sinh(x)^2 - b^
2 + 6*(b^2*cosh(x)^5 - 2*b^2*cosh(x)^3 + b^2*cosh(x))*sinh(x))*sqrt(-a^2 -
a*b)*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)
*sqrt(-a^2 - a*b)/(a^2 + a*b)) + 24*((a^2*b + a*b^2)*cosh(x)^3 - (a^3 + 3*a
^2*b + 2*a*b^2)*cosh(x))*sinh(x))/((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh
(x)^6 + 6*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)*sinh(x)^5 + (a^4 + 3*
a^3*b + 3*a^2*b^2 + a*b^3)*sinh(x)^6 - 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3
)*cosh(x)^4 - 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3 - 5*(a^4 + 3*a^3*b + 3*a
^2*b^2 + a*b^3)*cosh(x)^2)*sinh(x)^4 - a^4 - 3*a^3*b - 3*a^2*b^2 - a*b^3 +
4*(5*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^3 - 3*(a^4 + 3*a^3*b + 3*a
^2*b^2 + a*b^3)*cosh(x))*sinh(x)^3 + 3*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*
cosh(x)^2 + 3*(5*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^4 + a^4 + 3*a^
3*b + 3*a^2*b^2 + a*b^3 - 6*(a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^2)*
sinh(x)^2 + 6*((a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^5 - 2*(a^4 + 3*a
^3*b + 3*a^2*b^2 + a*b^3)*cosh(x)^3 + (a^4 + 3*a^3*b + 3*a^2*b^2 + a*b^3)*c
osh(x))*sinh(x))]

```

Sympy [F]

$$\int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx$$

[In] integrate(csch(x)**4/(a+b*cosh(x)**2),x)

[Out] Integral(csch(x)**4/(a + b*cosh(x)**2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 161 vs. $2(49) = 98$.

Time = 0.33 (sec) , antiderivative size = 161, normalized size of antiderivative = 2.73

$$\int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx = -\frac{b^2 \log\left(\frac{be^{(-2x)}+2a+b-2\sqrt{(a+b)a}}{be^{(-2x)}+2a+b+2\sqrt{(a+b)a}}\right)}{2\sqrt{(a+b)a}(a^2+2ab+b^2)} - \frac{2(6(a+2b)e^{(-2x)}-3be^{(-4x)}-2a-5b)}{3(a^2+2ab+b^2-3(a^2+2ab+b^2)e^{(-2x)}+3(a^2+2ab+b^2)e^{(-4x)}-(a^2+2ab+b^2)e^{(-6x)})}$$

[In] integrate(csch(x)^4/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-1/2*b^2*\log((b*e^{(-2*x)} + 2*a + b - 2*sqrt((a + b)*a))/(b*e^{(-2*x)} + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*(a^2 + 2*a*b + b^2)) - 2/3*(6*(a + 2*b)*e^{(-2*x)} - 3*b*e^{(-4*x)} - 2*a - 5*b)/(a^2 + 2*a*b + b^2 - 3*(a^2 + 2*a*b + b^2)*e^{(-2*x)} + 3*(a^2 + 2*a*b + b^2)*e^{(-4*x)} - (a^2 + 2*a*b + b^2)*e^{(-6*x)})$

Giac [F]

$$\int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^4}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)^4/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.38 (sec) , antiderivative size = 245, normalized size of antiderivative = 4.15

$$\int \frac{\operatorname{csch}^4(x)}{a + b \cosh^2(x)} dx = \frac{2b}{(a+b)^2(e^{2x}-1)} - \frac{4}{(a+b)(e^{4x}-2e^{2x}+1)} - \frac{3(a+b)(3e^{2x}-3e^{4x}+e^{6x}-1)}{8} - \frac{b^2 \ln\left(\frac{4b^2(2ab+8a^2e^{2x}+b^2e^{2x}+b^2+8abe^{2x})}{a(a+b)^5} - \frac{8b^2(b+4ae^{2x}+2be^{2x})}{\sqrt{a}(a+b)^{9/2}}\right)}{2\sqrt{a}(a+b)^{5/2}} + \frac{b^2 \ln\left(\frac{8b^2(b+4ae^{2x}+2be^{2x})}{\sqrt{a}(a+b)^{9/2}} + \frac{4b^2(2ab+8a^2e^{2x}+b^2e^{2x}+b^2+8abe^{2x})}{a(a+b)^5}\right)}{2\sqrt{a}(a+b)^{5/2}}$$

[In] `int(1/(sinh(x)^4*(a + b*cosh(x)^2)),x)`

[Out]
$$\begin{aligned} & (2*b)/((a + b)^2*(\exp(2*x) - 1)) - 4/((a + b)*(\exp(4*x) - 2*\exp(2*x) + 1)) \\ & - 8/(3*(a + b)*(3*\exp(2*x) - 3*\exp(4*x) + \exp(6*x) - 1)) - (b^2*\log((4*b^2* \\ & (2*a*b + 8*a^2*\exp(2*x) + b^2*\exp(2*x) + b^2 + 8*a*b*\exp(2*x)))/(a*(a + b)^5 \\ & - (8*b^2*(b + 4*a*\exp(2*x) + 2*b*\exp(2*x)))/(a^{1/2}*(a + b)^{9/2}))))/(2 \\ & *a^{1/2}*(a + b)^{5/2}) + (b^2*\log((8*b^2*(b + 4*a*\exp(2*x) + 2*b*\exp(2*x)) \\ &)/(a^{1/2}*(a + b)^{9/2}) + (4*b^2*(2*a*b + 8*a^2*\exp(2*x) + b^2*\exp(2*x) + \\ & b^2 + 8*a*b*\exp(2*x)))/(a*(a + b)^5)))/(2*a^{1/2}*(a + b)^{5/2}) \end{aligned}$$

3.18 $\int \frac{\operatorname{csch}^6(x)}{a+b \cosh^2(x)} dx$

Optimal result	144
Rubi [A] (verified)	144
Mathematica [A] (verified)	146
Maple [B] (verified)	146
Fricas [B] (verification not implemented)	147
Sympy [F(-1)]	150
Maxima [B] (verification not implemented)	150
Giac [F]	150
Mupad [B] (verification not implemented)	151

Optimal result

Integrand size = 15, antiderivative size = 89

$$\int \frac{\operatorname{csch}^6(x)}{a+b \cosh^2(x)} dx = -\frac{b^3 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{(a^2 + 3ab + 3b^2) \operatorname{coth}(x)}{(a+b)^3} + \frac{(2a+3b) \operatorname{coth}^3(x)}{3(a+b)^2} - \frac{\operatorname{coth}^5(x)}{5(a+b)}$$

[Out] $-(a^2+3*a*b+3*b^2)*\operatorname{coth}(x)/(a+b)^3+1/3*(2*a+3*b)*\operatorname{coth}(x)^3/(a+b)^2-1/5*\operatorname{coth}(x)^5/(a+b)-b^3*\operatorname{arctanh}(a^{1/2}*\tanh(x)/(a+b)^{1/2})/(a+b)^{7/2}/a^{1/2}$

Rubi [A] (verified)

Time = 0.08 (sec), antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3270, 398, 214}

$$\int \frac{\operatorname{csch}^6(x)}{a+b \cosh^2(x)} dx = -\frac{(a^2 + 3ab + 3b^2) \operatorname{coth}(x)}{(a+b)^3} - \frac{b^3 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{\operatorname{coth}^5(x)}{5(a+b)} + \frac{(2a+3b) \operatorname{coth}^3(x)}{3(a+b)^2}$$

[In] $\operatorname{Int}[\operatorname{Csch}[x]^6/(a+b*\operatorname{Cosh}[x]^2), x]$

[Out] $-(b^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tanh}[x])/(\operatorname{Sqrt}[a+b])]/(\operatorname{Sqrt}[a]*(a+b)^{7/2})) - ((a^2+3*a*b+3*b^2)*\operatorname{Coth}[x])/(a+b)^3 + ((2*a+3*b)*\operatorname{Coth}[x]^3)/(3*(a+b)^2) - \operatorname{Coth}[x]^5/(5*(a+b))$

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 398

Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]

Rule 3270

Int[cos[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{(1-x^2)^3}{a-(a+b)x^2} dx, x, \coth(x)\right) \\
 &= -\text{Subst}\left(\int \left(\frac{a^2+3ab+3b^2}{(a+b)^3} - \frac{(2a+3b)x^2}{(a+b)^2} + \frac{x^4}{a+b} + \frac{b^3}{(a+b)^3(a-(a+b)x^2)}\right) dx, x, \coth(x)\right) \\
 &= -\frac{(a^2+3ab+3b^2)\coth(x)}{(a+b)^3} + \frac{(2a+3b)\coth^3(x)}{3(a+b)^2} \\
 &\quad - \frac{\coth^5(x)}{5(a+b)} - \frac{b^3\text{Subst}\left(\int \frac{1}{a-(a+b)x^2} dx, x, \coth(x)\right)}{(a+b)^3} \\
 &= -\frac{b^3\text{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{(a^2+3ab+3b^2)\coth(x)}{(a+b)^3} + \frac{(2a+3b)\coth^3(x)}{3(a+b)^2} - \frac{\coth^5(x)}{5(a+b)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.30 (sec) , antiderivative size = 92, normalized size of antiderivative = 1.03

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx$$

$$= -\frac{b^3 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}(a+b)^{7/2}} - \frac{\operatorname{coth}(x) (8a^2 + 26ab + 33b^2 - (4a^2 + 13ab + 9b^2) \operatorname{csch}^2(x) + 3(a+b)^2 \operatorname{csch}^4(x))}{15(a+b)^3}$$

[In] Integrate[Csch[x]^6/(a + b*Cosh[x]^2), x]

[Out] $-\left(\frac{b^3 \operatorname{ArcTanh}\left[\frac{\sqrt{a} \operatorname{Tanh}[x]}{\sqrt{a+b}}\right]}{\sqrt{a}(a+b)^{7/2}}\right) - \left(\operatorname{Coth}[x] \cdot (8a^2 + 26ab + 33b^2 - (4a^2 + 13ab + 9b^2) \operatorname{Csch}[x]^2 + 3(a+b)^2 \operatorname{Csch}[x]^4)\right) / (15(a+b)^3)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 247 vs. 2(77) = 154.

Time = 54.96 (sec) , antiderivative size = 248, normalized size of antiderivative = 2.79

method	result
default	$-\frac{a^2 \tanh\left(\frac{x}{2}\right)^5}{5} + \frac{2ab \tanh\left(\frac{x}{2}\right)^5}{5} + \frac{b^2 \tanh\left(\frac{x}{2}\right)^5}{5} - \frac{5a^2 \tanh\left(\frac{x}{2}\right)^3}{3} - \frac{14ab \tanh\left(\frac{x}{2}\right)^3}{3} - \frac{3b^2 \tanh\left(\frac{x}{2}\right)^3}{3} + 10a^2 \tanh\left(\frac{x}{2}\right) + 32ab \tanh\left(\frac{x}{2}\right) + 38b^2 \tanh\left(\frac{x}{2}\right) - \frac{b^3 \ln\left(e^{2x} + 2\right)}{32(a+b)^3}$
risch	$-\frac{2(15b^2 e^{8x} - 30ab e^{6x} - 90b^2 e^{6x} + 80a^2 e^{4x} + 230ab e^{4x} + 240b^2 e^{4x} - 40a^2 e^{2x} - 130b e^{2x} a - 150b^2 e^{2x} + 8a^2 + 26ab + 33b^2)}{15(e^{2x} - 1)^5 (a+b)^3} + \frac{b^3 \ln\left(e^{2x} + 2\right)}{32(a+b)^3}$

[In] int(csch(x)^6/(a+b*cosh(x)^2), x, method=_RETURNVERBOSE)

[Out] $-1/32/(a+b)^3 \cdot (1/5 a^2 \tanh(1/2 x)^5 + 2/5 a b \tanh(1/2 x)^5 + 1/5 b^2 \tanh(1/2 x)^5 - 5/3 a^2 \tanh(1/2 x)^3 - 14/3 a b \tanh(1/2 x)^3 - 3 b^2 \tanh(1/2 x)^3 + 10 a^2 \tanh(1/2 x) + 32 a b \tanh(1/2 x) + 38 b^2 \tanh(1/2 x)) - 1/160/(a+b) \cdot \tanh(1/2 x)^5 - 1/96 \cdot (-5 a - 9 b) / (a+b)^2 \cdot \tanh(1/2 x)^3 - 1/32/(a+b)^3 \cdot (10 a^2 + 32 a b + 38 b^2) / \tanh(1/2 x) + 2 b^3 / (a+b)^3 \cdot (-1/4 a^{1/2} / (a+b)^{1/2} \cdot \ln((a+b)^{1/2} \tanh(1/2 x)^2 + 2 \tanh(1/2 x) a^{1/2} + (a+b)^{1/2}) + 1/4 a^{1/2} / (a+b)^{1/2} \cdot \ln((a+b)^{1/2} \tanh(1/2 x)^2 - 2 \tanh(1/2 x) a^{1/2} + (a+b)^{1/2}))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2408 vs. 2(77) = 154.

Time = 0.31 (sec) , antiderivative size = 4977, normalized size of antiderivative = 55.92

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(csch(x)^6/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [-1/30*(60*(a^2*b^2 + a*b^3)*cosh(x)^8 + 480*(a^2*b^2 + a*b^3)*cosh(x)*sinh(x)^7 + 60*(a^2*b^2 + a*b^3)*sinh(x)^8 - 120*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x)^6 - 120*(a^3*b + 4*a^2*b^2 + 3*a*b^3 - 14*(a^2*b^2 + a*b^3)*cosh(x)^2)*sinh(x)^6 + 240*(14*(a^2*b^2 + a*b^3)*cosh(x)^3 - 3*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x))*sinh(x)^5 + 40*(8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*cosh(x)^4 + 40*(105*(a^2*b^2 + a*b^3)*cosh(x)^4 + 8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3 - 45*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x)^2)*sinh(x)^4 + 32*a^4 + 136*a^3*b + 236*a^2*b^2 + 132*a*b^3 + 160*(21*(a^2*b^2 + a*b^3)*cosh(x)^5 - 15*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x)^3 + (8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*cosh(x))*sinh(x)^3 - 40*(4*a^4 + 17*a^3*b + 28*a^2*b^2 + 15*a*b^3)*cosh(x)^2 + 40*(42*(a^2*b^2 + a*b^3)*cosh(x)^6 - 45*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x)^4 - 4*a^4 - 17*a^3*b - 28*a^2*b^2 - 15*a*b^3 + 6*(8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*cosh(x)^2)*sinh(x)^2 - 15*(b^3*cosh(x)^10 + 10*b^3*cosh(x)*sinh(x)^9 + b^3*sinh(x)^10 - 5*b^3*cosh(x)^8 + 10*b^3*cosh(x)^6 + 5*(9*b^3*cosh(x)^2 - b^3)*sinh(x)^8 + 40*(3*b^3*cosh(x)^3 - b^3*cosh(x))*sinh(x)^7 - 10*b^3*cosh(x)^4 + 10*(21*b^3*cosh(x)^4 - 14*b^3*cosh(x)^2 + b^3)*sinh(x)^6 + 4*(63*b^3*cosh(x)^5 - 70*b^3*cosh(x)^3 + 15*b^3*cosh(x))*sinh(x)^5 + 5*b^3*cosh(x)^2 + 10*(21*b^3*cosh(x)^6 - 35*b^3*cosh(x)^4 + 15*b^3*cosh(x)^2 - b^3)*sinh(x)^4 + 40*(3*b^3*cosh(x)^7 - 7*b^3*cosh(x)^5 + 5*b^3*cosh(x)^3 - b^3*cosh(x))*sinh(x)^3 - b^3 + 5*(9*b^3*cosh(x)^8 - 28*b^3*cosh(x)^6 + 30*b^3*cosh(x)^4 - 12*b^3*cosh(x)^2 + b^3)*sinh(x)^2 + 10*(b^3*cosh(x)^9 - 4*b^3*cosh(x)^7 + 6*b^3*cosh(x)^5 - 4*b^3*cosh(x)^3 + b^3*cosh(x))*sinh(x))*sqrt(a^2 + a*b)*log((b^2*cosh(x)^4 + 4*b^2*cosh(x)*sinh(x)^3 + b^2*sinh(x)^4 + 2*(2*a*b + b^2)*cosh(x)^2 + 2*(3*b^2*cosh(x)^2 + 2*a*b + b^2)*sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*cosh(x)^3 + (2*a*b + b^2)*cosh(x))*sinh(x) + 4*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(a^2 + a*b))/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 80*(6*(a^2*b^2 + a*b^3)*cosh(x)^7 - 9*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*cosh(x)^5 + 2*(8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*cosh(x)^3 - (4*a^4 + 17*a^3*b + 28*a^2*b^2 + 15*a*b^3)*cosh(x))*sinh(x))/((a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*cosh(x)^10 + 10*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*cosh(x)*sinh(x)^9 + (a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*sinh(x)^10 - 5*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*cosh(x)^8 - 5*(a^5 + 4*a^4

$$\begin{aligned}
& *b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4 - 9*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^2*\sinh(x)^8 + 40*(3*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^3 - (a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x))*\sinh(x)^7 + 10*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^6 + 10*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4 + 21*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^4 - 14*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^2)*\sinh(x)^6 + 4*(63*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^5 - 70*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^3 + 15*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x))*\sinh(x)^5 - a^5 - 4*a^4*b - 6*a^3*b^2 - 4*a^2*b^3 - a*b^4 - 10*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^4 + 10*(21*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^6 - a^5 - 4*a^4*b - 6*a^3*b^2 - 4*a^2*b^3 - a*b^4 - 35*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^4 + 15*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^2)*\sinh(x)^4 + 40*(3*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^7 - 7*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^5 + 5*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^3 - (a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x))*\sinh(x)^3 + 5*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^2 + 5*(9*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^8 - 28*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^6 + a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4 + 30*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^4 - 12*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^2)*\sinh(x)^2 + 10*((a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^9 - 4*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^7 + 6*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^5 - 4*(a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x)^3 + (a^5 + 4*a^4*b + 6*a^3*b^2 + 4*a^2*b^3 + a*b^4)*\cosh(x))*\sinh(x)), -1/15*(30*(a^2*b^2 + a*b^3)*\cosh(x)^8 + 240*(a^2*b^2 + a*b^3)*\cosh(x)*\sinh(x)^7 + 30*(a^2*b^2 + a*b^3)*\sinh(x)^8 - 60*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*\cosh(x)^6 - 60*(a^3*b + 4*a^2*b^2 + 3*a*b^3 - 14*(a^2*b^2 + a*b^3)*\cosh(x)^2)*\sinh(x)^6 + 120*(14*(a^2*b^2 + a*b^3)*\cosh(x)^3 - 3*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*\cosh(x))*\sinh(x)^5 + 20*(8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*\cosh(x)^4 + 20*(105*(a^2*b^2 + a*b^3)*\cosh(x)^4 + 8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3 - 45*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*\cosh(x)^2)*\sinh(x)^4 + 16*a^4 + 68*a^3*b + 118*a^2*b^2 + 66*a*b^3 + 80*(21*(a^2*b^2 + a*b^3)*\cosh(x)^5 - 15*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*\cosh(x)^3 + (8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*\cosh(x))*\sinh(x)^3 - 20*(4*a^4 + 17*a^3*b + 28*a^2*b^2 + 15*a*b^3)*\cosh(x)^2 + 20*(42*(a^2*b^2 + a*b^3)*\cosh(x)^6 - 45*(a^3*b + 4*a^2*b^2 + 3*a*b^3)*\cosh(x)^4 - 4*a^4 - 17*a^3*b - 28*a^2*b^2 - 15*a*b^3 + 6*(8*a^4 + 31*a^3*b + 47*a^2*b^2 + 24*a*b^3)*\cosh(x)^2)*\sinh(x)^2 + 15*(b^3*\cosh(x)^10 + 10*b^3*\cosh(x)*\sinh(x)^9 + b^3*\sinh(x)^10 - 5*b^3*\cosh(x)^8 + 10*b^3*\cosh(x)^6 + 5*(9*b^3*\cosh(x)^2 - b^3)*\sinh(x)^8 + 40*(3*b^3*\cosh(x)^3 - b^3*\cosh(x))*\sinh(x)^7 - 10*b^3*\cosh(x)^4 + 10*(21*b^3*\cosh(x)^4 - 14*b^3*\cosh(x)^2 + b^3)*\sinh(x)^6 + 4*(63*b^3*\cosh(x)^5 - 70*b^3*\cosh(x)^3 + 15*b^3*\cosh(x))*\sinh(x)^5 + 5*b^3*\cosh(x)^2 + 10*(21*b^3*\cosh(x)^6 - 35*b^3*\cosh(x)^4 + 15*
\end{aligned}$$

$$\begin{aligned}
& b^3 \cosh(x)^2 - b^3 \sinh(x)^4 + 40(3b^3 \cosh(x)^7 - 7b^3 \cosh(x)^5 + 5b^3 \cosh(x)^3 - b^3 \cosh(x)) \sinh(x)^3 - b^3 + 5(9b^3 \cosh(x)^8 - 28b^3 \cosh(x)^6 + 30b^3 \cosh(x)^4 - 12b^3 \cosh(x)^2 + b^3) \sinh(x)^2 + 10(b^3 \cosh(x)^9 - 4b^3 \cosh(x)^7 + 6b^3 \cosh(x)^5 - 4b^3 \cosh(x)^3 + b^3 \cosh(x)) \sinh(x) \\
& \cdot \sqrt{-a^2 - ab} \arctan\left(\frac{1}{2}(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + 2a + b) \sqrt{-a^2 - ab} / (a^2 + ab)\right) + 40(6(a^2 b^2 + a^3 b) \cosh(x)^7 - 9(a^3 b + 4a^2 b^2 + 3a^3 b^3) \cosh(x)^5 + 2(8a^4 + 31a^3 b + 47a^2 b^2 + 24a^3 b^3) \cosh(x)^3 - (4a^4 + 17a^3 b + 28a^2 b^2 + 15a^3 b^3) \cosh(x)) \sinh(x) \\
& / ((a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^{10} + 10(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x) \sinh(x)^9 + (a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \sinh(x)^{10} - 5(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^8 - 5(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4 - 9(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^2) \sinh(x)^8 + 40(3(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^3 - (a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)) \sinh(x)^7 + 10(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^6 + 10(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4 + 21(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^4 - 14(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^2) \sinh(x)^6 + 4(63(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^5 - 70(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^3 + 15(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)) \sinh(x)^5 - a^5 - 4a^4 b - 6a^3 b^2 - 4a^2 b^3 - a^3 b^4 - 10(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^4 + 10(21(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^6 - a^5 - 4a^4 b - 6a^3 b^2 - 4a^2 b^3 - a^3 b^4 - 35(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^4 + 15(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^2) \sinh(x)^4 + 40(3(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^7 - 7(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^5 + 5(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^3 - (a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)) \sinh(x)^3 + 5(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^2 + 5(9(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^8 - 28(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^6 + a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4 + 30(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^4 - 12(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^2) \sinh(x)^2 + 10((a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^9 - 4(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^7 + 6(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^5 - 4(a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)^3 + (a^5 + 4a^4 b + 6a^3 b^2 + 4a^2 b^3 + a^3 b^4) \cosh(x)) \sinh(x)]
\end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(csch(x)**6/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 307 vs. 2(77) = 154.

Time = 0.31 (sec) , antiderivative size = 307, normalized size of antiderivative = 3.45

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx = \frac{b^3 \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{2(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{(a+b)a}} - \frac{2(15b^2e^{(-8x)} + 8a^2 + 26ab + 33b^2 - 10(4a^2 + 13ab + 15b^2)e^{(-2x)} - 15(a^3 + 3a^2b + 3ab^2 + b^3 - 5(a^3 + 3a^2b + 3ab^2 + b^3)e^{(-2x)} + 10(a^3 + 3a^2b + 3ab^2 + b^3)e^{(-4x)} - 10(a^3 + 3a^2b + 3ab^2 + b^3)e^{(-6x)} + 5(a^3 + 3a^2b + 3ab^2 + b^3)e^{(-8x)} - (a^3 + 3a^2b + 3ab^2 + b^3)e^{(-10x)}))}{2(a^3 + 3a^2b + 3ab^2 + b^3)\sqrt{(a+b)a}}$$

[In] integrate(csch(x)^6/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/2*b^3*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*sqrt((a + b)*a)) - 2/15*(15*b^2*e^(-8*x) + 8*a^2 + 26*a*b + 33*b^2 - 10*(4*a^2 + 13*a*b + 15*b^2)*e^(-2*x) + 10*(8*a^2 + 23*a*b + 24*b^2)*e^(-4*x) - 30*(a*b + 3*b^2)*e^(-6*x))/(a^3 + 3*a^2*b + 3*a*b^2 + b^3 - 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(-2*x) + 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(-4*x) - 10*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(-6*x) + 5*(a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(-8*x) - (a^3 + 3*a^2*b + 3*a*b^2 + b^3)*e^(-10*x))

Giac [F]

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{csch}(x)^6}{b \cosh(x)^2 + a} dx$$

[In] integrate(csch(x)^6/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.50 (sec) , antiderivative size = 333, normalized size of antiderivative = 3.74

$$\int \frac{\operatorname{csch}^6(x)}{a + b \cosh^2(x)} dx = \frac{4(b^2 + ab)}{(a+b)^3 (e^{4x} - 2e^{2x} + 1)} - \frac{16}{(a+b)(6e^{4x} - 4e^{2x} - 4e^{6x} + e^{8x} + 1)} - \frac{2b^2}{(a+b)^3 (e^{2x} - 1)} - \frac{32}{5(a+b)(5e^{2x} - 10e^{4x} + 10e^{6x} - 5e^{8x} + e^{10x} - 1)} - \frac{8(4a + 3b)}{3(a+b)^2 (3e^{2x} - 3e^{4x} + e^{6x} - 1)} + \frac{b^3 \ln\left(\frac{4b^4(2ab + 8a^2e^{2x} + b^2e^{2x} + b^2 + 8abe^{2x})}{a(a+b)^7} - \frac{8b^4(b + 4ae^{2x} + 2be^{2x})}{\sqrt{a}(a+b)^{13/2}}\right)}{2\sqrt{a}(a+b)^{7/2}} - \frac{b^3 \ln\left(\frac{8b^4(b + 4ae^{2x} + 2be^{2x})}{\sqrt{a}(a+b)^{13/2}} + \frac{4b^4(2ab + 8a^2e^{2x} + b^2e^{2x} + b^2 + 8abe^{2x})}{a(a+b)^7}\right)}{2\sqrt{a}(a+b)^{7/2}}$$

[In] int(1/(sinh(x)^6*(a + b*cosh(x)^2)),x)

```
[Out] (4*(a*b + b^2))/((a + b)^3*(exp(4*x) - 2*exp(2*x) + 1)) - 16/((a + b)*(6*exp(4*x) - 4*exp(2*x) - 4*exp(6*x) + exp(8*x) + 1)) - (2*b^2)/((a + b)^3*(exp(2*x) - 1)) - 32/(5*(a + b)*(5*exp(2*x) - 10*exp(4*x) + 10*exp(6*x) - 5*exp(8*x) + exp(10*x) - 1)) - (8*(4*a + 3*b))/(3*(a + b)^2*(3*exp(2*x) - 3*exp(4*x) + exp(6*x) - 1)) + (b^3*log((4*b^4*(2*a*b + 8*a^2*exp(2*x) + b^2*exp(2*x) + b^2 + 8*a*b*exp(2*x)))/(a*(a + b)^7) - (8*b^4*(b + 4*a*exp(2*x) + 2*b*exp(2*x)))/(a^(1/2)*(a + b)^(13/2))))/(2*a^(1/2)*(a + b)^(7/2)) - (b^3*log((8*b^4*(b + 4*a*exp(2*x) + 2*b*exp(2*x)))/(a^(1/2)*(a + b)^(13/2)) + (4*b^4*(2*a*b + 8*a^2*exp(2*x) + b^2*exp(2*x) + b^2 + 8*a*b*exp(2*x)))/(a*(a + b)^7)))/(2*a^(1/2)*(a + b)^(7/2))
```

3.19 $\int \frac{\sinh(x)}{4-3 \cosh^3(x)} dx$

Optimal result	152
Rubi [A] (verified)	152
Mathematica [A] (verified)	154
Maple [C] (verified)	155
Fricas [B] (verification not implemented)	155
Sympy [A] (verification not implemented)	157
Maxima [F]	157
Giac [A] (verification not implemented)	157
Mupad [B] (verification not implemented)	158

Optimal result

Integrand size = 13, antiderivative size = 98

$$\int \frac{\sinh(x)}{4-3 \cosh^3(x)} dx = \frac{\arctan\left(\frac{1+\sqrt[3]{6}\cosh(x)}{\sqrt{3}}\right)}{2\sqrt[3]{2}3^{5/6}} - \frac{\log\left(2^{2/3} - \sqrt[3]{3}\cosh(x)\right)}{6\sqrt[3]{6}} + \frac{\log\left(2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}\cosh(x) + 3^{2/3}\cosh^2(x)\right)}{12\sqrt[3]{6}}$$

[Out] 1/12*arctan(1/3*(1+6^(1/3)*cosh(x))*3^(1/2))*2^(2/3)*3^(1/6)-1/36*ln(2^(2/3)-3^(1/3)*cosh(x))*6^(2/3)+1/72*ln(2*2^(1/3)+2^(2/3)*3^(1/3)*cosh(x)+3^(2/3)*cosh(x)^2)*6^(2/3)

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {3302, 206, 31, 648, 631, 210, 642}

$$\int \frac{\sinh(x)}{4-3 \cosh^3(x)} dx = \frac{\arctan\left(\frac{\sqrt[3]{6}\cosh(x)+1}{\sqrt{3}}\right)}{2\sqrt[3]{2}3^{5/6}} + \frac{\log\left(3^{2/3}\cosh^2(x) + 2^{2/3}\sqrt[3]{3}\cosh(x) + 2\sqrt[3]{2}\right)}{12\sqrt[3]{6}} - \frac{\log\left(2^{2/3} - \sqrt[3]{3}\cosh(x)\right)}{6\sqrt[3]{6}}$$

[In] Int[Sinh[x]/(4 - 3*Cosh[x]^3), x]

[Out] ArcTan[(1 + 6^(1/3)*Cosh[x])/Sqrt[3]]/(2*2^(1/3)*3^(5/6)) - Log[2^(2/3) - 3^(1/3)*Cosh[x]]/(6*6^(1/3)) + Log[2*2^(1/3) + 2^(2/3)*3^(1/3)*Cosh[x] + 3^(2/3)*Cosh[x]^2]/(12*6^(1/3))

Rule 31

Int[((a_) + (b_)*(x_))^-1, x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 206

Int[((a_) + (b_)*(x_)^3)^-1, x_Symbol] := Dist[1/(3*Rt[a, 3]^2), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]^2), Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_)*(x_)^2)^-1, x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^-1, x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 3302

Int[cos[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*((c_)*sin[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Di

```

st[ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b*(c*ff*x)^n)^p, x], x,
Sin[e + f*x]/ff], x]] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)
)/2] && (EqQ[n, 4] || GtQ[m, 0] || IGtQ[p, 0] || IntegersQ[m, p])

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1}{4 - 3x^3} dx, x, \cosh(x)\right) \\
&= \frac{\text{Subst}\left(\int \frac{1}{2^{2/3} - \sqrt[3]{3}x} dx, x, \cosh(x)\right)}{6\sqrt[3]{2}} + \frac{\text{Subst}\left(\int \frac{2^{2/3} + \sqrt[3]{3}x}{2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}x + 3^{2/3}x^2} dx, x, \cosh(x)\right)}{6\sqrt[3]{2}} \\
&= -\frac{\log\left(2^{2/3} - \sqrt[3]{3}\cosh(x)\right)}{6\sqrt[3]{6}} + \frac{\text{Subst}\left(\int \frac{1}{2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}x + 3^{2/3}x^2} dx, x, \cosh(x)\right)}{2^{2/3}} \\
&\quad + \frac{\text{Subst}\left(\int \frac{2^{2/3}\sqrt[3]{3} + 2^{2/3}x}{2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}x + 3^{2/3}x^2} dx, x, \cosh(x)\right)}{12\sqrt[3]{6}} \\
&= -\frac{\log\left(2^{2/3} - \sqrt[3]{3}\cosh(x)\right)}{6\sqrt[3]{6}} + \frac{\log\left(2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}\cosh(x) + 3^{2/3}\cosh^2(x)\right)}{12\sqrt[3]{6}} \\
&\quad - \frac{\text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, 1 + \sqrt[3]{6}\cosh(x)\right)}{2\sqrt[3]{6}} \\
&= \frac{\arctan\left(\frac{1 + \sqrt[3]{6}\cosh(x)}{\sqrt{3}}\right)}{2\sqrt[3]{23^{5/6}}} - \frac{\log\left(2^{2/3} - \sqrt[3]{3}\cosh(x)\right)}{6\sqrt[3]{6}} \\
&\quad + \frac{\log\left(2\sqrt[3]{2} + 2^{2/3}\sqrt[3]{3}\cosh(x) + 3^{2/3}\cosh^2(x)\right)}{12\sqrt[3]{6}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.79

$$\begin{aligned}
&\int \frac{\sinh(x)}{4 - 3\cosh^3(x)} dx \\
&= \frac{1}{72} \left(6^{2/3} \sqrt[3]{3} \arctan\left(\frac{1 + \sqrt[3]{6}\cosh(x)}{\sqrt{3}}\right) \right. \\
&\quad \left. + 6^{2/3} \left(-2\log\left(2 - \sqrt[3]{6}\cosh(x)\right) + \log\left(4 + 2\sqrt[3]{6}\cosh(x) + 6^{2/3}\cosh^2(x)\right) \right) \right)
\end{aligned}$$

[In] Integrate[Sinh[x]/(4 - 3*Cosh[x]^3),x]

[Out] $(6 \cdot 2^{2/3} \cdot 3^{1/6} \cdot \text{ArcTan}[(1 + 6^{1/3} \cdot \text{Cosh}[x])/\text{Sqrt}[3]] + 6^{2/3} \cdot (-2 \cdot \text{Log}[2 - 6^{1/3} \cdot \text{Cosh}[x]] + \text{Log}[4 + 2 \cdot 6^{1/3} \cdot \text{Cosh}[x] + 6^{2/3} \cdot \text{Cosh}[x]^2]))/72$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.53 (sec) , antiderivative size = 26, normalized size of antiderivative = 0.27

method	result
risch	$\sum_{R=\text{RootOf}(1296Z^3+1)} -R \ln(24_R e^x + e^{2x} + 1)$
derivativdivides	$-\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cosh(x) - \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}}}{3}\right)}{36} + \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cosh(x)^2 + \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \cosh(x)}{3} + \frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{72} + \frac{4^{\frac{1}{3}} 3^{\frac{1}{6}} \arctan\left(\frac{\sqrt{3} \left(\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}} \cosh(x)}{2} + \frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{3}\right)}{12}$
default	$-\frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cosh(x) - \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}}}{3}\right)}{36} + \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \ln\left(\cosh(x)^2 + \frac{4^{\frac{1}{3}} 3^{\frac{2}{3}} \cosh(x)}{3} + \frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{72} + \frac{4^{\frac{1}{3}} 3^{\frac{1}{6}} \arctan\left(\frac{\sqrt{3} \left(\frac{4^{\frac{2}{3}} 3^{\frac{1}{3}} \cosh(x)}{2} + \frac{4^{\frac{2}{3}} 3^{\frac{1}{3}}}{3}\right)}{3}\right)}{12}$

[In] int(sinh(x)/(4-3*cosh(x)^3),x,method=_RETURNVERBOSE)

[Out] sum(_R*ln(24*_R*exp(x)+exp(2*x)+1),_R=RootOf(1296*_Z^3+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 305 vs. 2(69) = 138.

Time = 0.26 (sec) , antiderivative size = 305, normalized size of antiderivative = 3.11

$$\int \frac{\sinh(x)}{4 - 3 \cosh^3(x)} dx = \frac{1}{12} \cdot 6^{\frac{1}{6}} \sqrt{2} (-1)^{\frac{1}{3}} \arctan \left(\frac{1}{12} \cdot 6^{\frac{1}{6}} \left(6^{\frac{2}{3}} \sqrt{2} (-1)^{\frac{2}{3}} \cosh(x)^3 + 6^{\frac{2}{3}} \sqrt{2} (-1)^{\frac{2}{3}} \sinh(x)^3 + \left(3 \cdot 6^{\frac{2}{3}} \sqrt{2} (-1)^{\frac{2}{3}} \cosh(x) + 4 \cdot 6^{\frac{1}{3}} \sqrt{2} \right) \sinh(x)^2 + 4 \cdot 6^{\frac{1}{3}} \sqrt{2} \right) \right) - \frac{1}{12} \cdot 6^{\frac{1}{6}} \sqrt{2} (-1)^{\frac{1}{3}} \arctan \left(\frac{1}{12} \cdot 6^{\frac{1}{6}} \left(6^{\frac{2}{3}} \sqrt{2} (-1)^{\frac{2}{3}} \cosh(x) + 6^{\frac{2}{3}} \sqrt{2} (-1)^{\frac{2}{3}} \sinh(x) + 2 \cdot 6^{\frac{1}{3}} \sqrt{2} \right) \right) - \frac{1}{72} \cdot 6^{\frac{2}{3}} (-1)^{\frac{1}{3}} \log \left(-\frac{2 \left(2 \cdot 6^{\frac{2}{3}} (-1)^{\frac{1}{3}} \cosh(x) - 3 \cosh(x)^2 - 3 \sinh(x)^2 - 4 \cdot 6^{\frac{1}{3}} (-1)^{\frac{2}{3}} - 3 \right)}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2} \right) + \frac{1}{36} \cdot 6^{\frac{2}{3}} (-1)^{\frac{1}{3}} \log \left(\frac{2 \left(6^{\frac{2}{3}} (-1)^{\frac{1}{3}} + 3 \cosh(x) \right)}{\cosh(x) - \sinh(x)} \right)$$

[In] integrate(sinh(x)/(4-3*cosh(x)^3),x, algorithm="fricas")

[Out] 1/12*6^(1/6)*sqrt(2)*(-1)^(1/3)*arctan(1/12*6^(1/6)*(6^(2/3)*sqrt(2)*(-1)^(2/3)*cosh(x)^3 + 6^(2/3)*sqrt(2)*(-1)^(2/3)*sinh(x)^3 + (3*6^(2/3)*sqrt(2)*(-1)^(2/3)*cosh(x) + 4*6^(1/3)*sqrt(2))*sinh(x)^2 + 4*6^(1/3)*sqrt(2)*cosh(x)^2 + (6^(2/3)*sqrt(2)*(-1)^(2/3) - 16*sqrt(2)*(-1)^(1/3))*cosh(x) + (3*6^(2/3)*sqrt(2)*(-1)^(2/3)*cosh(x)^2 + 6^(2/3)*sqrt(2)*(-1)^(2/3) + 8*6^(1/3)*sqrt(2)*cosh(x) - 16*sqrt(2)*(-1)^(1/3))*sinh(x) + 2*6^(1/3)*sqrt(2)) - 1/12*6^(1/6)*sqrt(2)*(-1)^(1/3)*arctan(1/12*6^(1/6)*(6^(2/3)*sqrt(2)*(-1)^(2/3)*cosh(x) + 6^(2/3)*sqrt(2)*(-1)^(2/3)*sinh(x) + 2*6^(1/3)*sqrt(2)) - 1/72*6^(2/3)*(-1)^(1/3)*log(-2*(2*6^(2/3)*(-1)^(1/3)*cosh(x) - 3*cosh(x)^2 - 3*sinh(x)^2 - 4*6^(1/3)*(-1)^(2/3) - 3)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 1/36*6^(2/3)*(-1)^(1/3)*log(2*(6^(2/3)*(-1)^(1/3) + 3*cosh(x))/(cosh(x) - sinh(x)))

Sympy [A] (verification not implemented)

Time = 0.48 (sec) , antiderivative size = 85, normalized size of antiderivative = 0.87

$$\int \frac{\sinh(x)}{4 - 3 \cosh^3(x)} dx = -\frac{6^{\frac{2}{3}} \log\left(\cosh(x) - \frac{6^{\frac{2}{3}}}{3}\right)}{36} + \frac{6^{\frac{2}{3}} \log\left(36 \cosh^2(x) + 12 \cdot 6^{\frac{2}{3}} \cosh(x) + 24 \cdot \sqrt[3]{6}\right)}{72} + \frac{2^{\frac{2}{3}} \cdot \sqrt[6]{3} \operatorname{atan}\left(\frac{\sqrt[3]{2} \cdot 3^{\frac{5}{6}} \cosh(x)}{3} + \frac{\sqrt{3}}{3}\right)}{12}$$

```
[In] integrate(sinh(x)/(4-3*cosh(x)**3),x)
```

```
[Out] -6**(2/3)*log(cosh(x) - 6**(2/3)/3)/36 + 6**(2/3)*log(36*cosh(x)**2 + 12*6*
*(2/3)*cosh(x) + 24*6**(1/3))/72 + 2**(2/3)*3**(1/6)*atan(2**(1/3)*3**(5/6)
*cosh(x)/3 + sqrt(3)/3)/12
```

Maxima [F]

$$\int \frac{\sinh(x)}{4 - 3 \cosh^3(x)} dx = \int -\frac{\sinh(x)}{3 \cosh^3(x) - 4} dx$$

```
[In] integrate(sinh(x)/(4-3*cosh(x)^3),x, algorithm="maxima")
```

```
[Out] -integrate(sinh(x)/(3*cosh(x)^3 - 4), x)
```

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 80, normalized size of antiderivative = 0.82

$$\int \frac{\sinh(x)}{4 - 3 \cosh^3(x)} dx = \frac{1}{12} \sqrt{3} \left(\frac{4}{3}\right)^{\frac{1}{3}} \arctan\left(\frac{1}{4} \sqrt{3} \left(\frac{4}{3}\right)^{\frac{2}{3}} \left(\left(\frac{4}{3}\right)^{\frac{1}{3}} + e^{-x} + e^x\right)\right) + \frac{1}{72} \cdot 36^{\frac{1}{3}} \log\left(\left(e^{-x} + e^x\right)^2 + 2 \left(\frac{4}{3}\right)^{\frac{1}{3}} \left(e^{-x} + e^x\right) + 4 \left(\frac{4}{3}\right)^{\frac{2}{3}}\right) - \frac{1}{12} \left(\frac{4}{3}\right)^{\frac{1}{3}} \log\left(\left|-2 \left(\frac{4}{3}\right)^{\frac{1}{3}} + e^{-x} + e^x\right|\right)$$

```
[In] integrate(sinh(x)/(4-3*cosh(x)^3),x, algorithm="giac")
```

[Out] $1/12*\sqrt{3}*(4/3)^{(1/3)}*\arctan(1/4*\sqrt{3}*(4/3)^{(2/3)}*((4/3)^{(1/3)} + e^{-x}) + e^x) + 1/72*36^{(1/3)}*\log((e^{-x} + e^x)^2 + 2*(4/3)^{(1/3)}*(e^{-x} + e^x) + 4*(4/3)^{(2/3)}) - 1/12*(4/3)^{(1/3)}*\log(\text{abs}(-2*(4/3)^{(1/3)} + e^{-x} + e^x))$

Mupad [B] (verification not implemented)

Time = 4.71 (sec) , antiderivative size = 205, normalized size of antiderivative = 2.09

$$\int \frac{\sinh(x)}{4 - 3 \cosh^3(x)} dx$$

$$= \frac{6^{2/3} \ln \left(\frac{256 e^{2x}}{81} - \frac{128 e^x}{27} + \frac{6^{2/3} \left(\frac{256 e^{2x}}{9} - \frac{2048 e^x}{27} + \frac{6^{2/3} (256 e^{2x} - \frac{2048 e^x}{3} + 256) + 256}{36} \right) + \frac{256}{9}}{36} + \frac{256}{81} \right)}{36}$$

$$- \frac{6^{2/3} \ln \left(\frac{256 e^{2x}}{81} - \frac{128 e^x}{27} + \frac{6^{2/3} \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) \left(\frac{256 e^{2x}}{9} - \frac{2048 e^x}{27} + \frac{6^{2/3} \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) (256 e^{2x} - \frac{2048 e^x}{3} + 256) + \frac{256}{9}}{36} + \frac{256}{9} \right) + \frac{256}{81}}{36} \right) \left(-\frac{1}{2} + \frac{\sqrt{3} i}{2} \right)}{36}$$

$$+ \frac{6^{2/3} \ln \left(\frac{256 e^{2x}}{81} - \frac{128 e^x}{27} - \frac{6^{2/3} \left(\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) \left(\frac{256 e^{2x}}{9} - \frac{2048 e^x}{27} - \frac{6^{2/3} \left(\frac{1}{2} + \frac{\sqrt{3} i}{2} \right) (256 e^{2x} - \frac{2048 e^x}{3} + 256) + \frac{256}{9}}{36} + \frac{256}{9} \right) + \frac{256}{81}}{36} \right) \left(\frac{1}{2} + \frac{\sqrt{3} i}{2} \right)}{36}$$

[In] $\text{int}(-\sinh(x)/(3*\cosh(x)^3 - 4), x)$

[Out] $(6^{(2/3)}*\log((256*\exp(2*x))/81 - (128*\exp(x))/27 - (6^{(2/3)}*((3^{(1/2)}*i)/2 + 1/2)*((256*\exp(2*x))/9 - (2048*\exp(x))/27 - (6^{(2/3)}*((3^{(1/2)}*i)/2 + 1/2)*(256*\exp(2*x) - (2048*\exp(x))/3 + 256))/36 + 256/9))/36 + 256/81)*((3^{(1/2)}*i)/2 + 1/2))/36 - (6^{(2/3)}*\log((256*\exp(2*x))/81 - (128*\exp(x))/27 + (6^{(2/3)}*((3^{(1/2)}*i)/2 - 1/2)*((256*\exp(2*x))/9 - (2048*\exp(x))/27 + (6^{(2/3)}*((3^{(1/2)}*i)/2 - 1/2)*(256*\exp(2*x) - (2048*\exp(x))/3 + 256))/36 + 256/9))/36 + 256/81)*((3^{(1/2)}*i)/2 - 1/2))/36 - (6^{(2/3)}*\log((256*\exp(2*x))/81 - (128*\exp(x))/27 + (6^{(2/3)}*((256*\exp(2*x))/9 - (2048*\exp(x))/27 + (6^{(2/3)}*(256*\exp(2*x) - (2048*\exp(x))/3 + 256))/36 + 256/9))/36 + 256/81))/36$

3.20 $\int \frac{\cosh^7(x)}{a+b \cosh^2(x)} dx$

Optimal result	159
Rubi [A] (verified)	159
Mathematica [A] (verified)	160
Maple [B] (verified)	161
Fricas [B] (verification not implemented)	161
Sympy [F(-1)]	163
Maxima [F]	163
Giac [F]	163
Mupad [B] (verification not implemented)	164

Optimal result

Integrand size = 15, antiderivative size = 78

$$\int \frac{\cosh^7(x)}{a+b \cosh^2(x)} dx = -\frac{a^3 \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{7/2} \sqrt{a+b}} + \frac{(a^2 - ab + b^2) \sinh(x)}{b^3} - \frac{(a-2b) \sinh^3(x)}{3b^2} + \frac{\sinh^5(x)}{5b}$$

[Out] (a^2-a*b+b^2)*sinh(x)/b^3-1/3*(a-2*b)*sinh(x)^3/b^2+1/5*sinh(x)^5/b-a^3*arctan(sinh(x)*b^(1/2)/(a+b)^(1/2))/b^(7/2)/(a+b)^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3265, 398, 211}

$$\int \frac{\cosh^7(x)}{a+b \cosh^2(x)} dx = -\frac{a^3 \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{7/2} \sqrt{a+b}} + \frac{(a^2 - ab + b^2) \sinh(x)}{b^3} - \frac{(a-2b) \sinh^3(x)}{3b^2} + \frac{\sinh^5(x)}{5b}$$

[In] Int[Cosh[x]^7/(a + b*Cosh[x]^2),x]

[Out] -((a^3*ArcTan[(Sqrt[b]*Sinh[x])/Sqrt[a + b]])/(b^(7/2)*Sqrt[a + b])) + ((a^2 - a*b + b^2)*Sinh[x])/b^3 - ((a - 2*b)*Sinh[x]^3)/(3*b^2) + Sinh[x]^5/(5*b)

Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:= Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q, 0] && GeQ[p, -q]
```

Rule 3265

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol]
:= With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{(1+x^2)^3}{a+b+bx^2} dx, x, \sinh(x)\right) \\
 &= \text{Subst}\left(\int \left(\frac{a^2-ab+b^2}{b^3} - \frac{(a-2b)x^2}{b^2} + \frac{x^4}{b} - \frac{a^3}{b^3(a+b+bx^2)}\right) dx, x, \sinh(x)\right) \\
 &= \frac{(a^2-ab+b^2)\sinh(x)}{b^3} - \frac{(a-2b)\sinh^3(x)}{3b^2} + \frac{\sinh^5(x)}{5b} - \frac{a^3 \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{b^3} \\
 &= -\frac{a^3 \arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{b^{7/2}\sqrt{a+b}} + \frac{(a^2-ab+b^2)\sinh(x)}{b^3} - \frac{(a-2b)\sinh^3(x)}{3b^2} + \frac{\sinh^5(x)}{5b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.23 (sec) , antiderivative size = 86, normalized size of antiderivative = 1.10

$$\int \frac{\cosh^7(x)}{a+b\cosh^2(x)} dx = \frac{a^3 \arctan\left(\frac{\sqrt{a+b}\text{csch}(x)}{\sqrt{b}}\right)}{b^{7/2}\sqrt{a+b}} + \frac{(8a^2-6ab+5b^2)\sinh(x)}{8b^3} - \frac{(4a-5b)\sinh(3x)}{48b^2} + \frac{\sinh(5x)}{80b}$$

```
[In] Integrate[Cosh[x]^7/(a + b*Cosh[x]^2), x]
```

```
[Out] (a^3*ArcTan[(Sqrt[a + b]*Csch[x])/Sqrt[b]])/(b^(7/2)*Sqrt[a + b]) + ((8*a^2 - 6*a*b + 5*b^2)*Sinh[x])/(8*b^3) - ((4*a - 5*b)*Sinh[3*x])/(48*b^2) + Sinh[5*x]/(80*b)
```


Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 205 vs. 2(66) = 132.

Time = 1.42 (sec) , antiderivative size = 206, normalized size of antiderivative = 2.64

method	result
risch	$\frac{e^{5x}}{160b} + \frac{5e^{3x}}{96b} - \frac{e^{3x}a}{24b^2} + \frac{e^x a^2}{2b^3} - \frac{3ae^x}{8b^2} + \frac{5e^x}{16b} - \frac{e^{-x}a^2}{2b^3} + \frac{3ae^{-x}}{8b^2} - \frac{5e^{-x}}{16b} - \frac{5e^{-3x}}{96b} + \frac{e^{-3x}a}{24b^2} - \frac{e^{-5x}}{160b} - \frac{a^3 \ln\left(\frac{e^{2x} + \sqrt{a+b} \cosh(x)}{e^{2x} - \sqrt{a+b} \cosh(x)}\right)}{2\sqrt{a+b}}$
default	$-\frac{2a^3 \left(\frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b} + 2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b} \sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b} - 2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b} \sqrt{b}} \right)}{b^3} - \frac{1}{5b(\tanh\left(\frac{x}{2}\right) + 1)^5} + \frac{1}{2b(\tanh\left(\frac{x}{2}\right) + 1)^4} - \frac{-7b + \sqrt{a+b}}{8b^2(\tanh\left(\frac{x}{2}\right) + 1)}$

[In] int(cosh(x)^7/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/160/b*exp(5*x)+5/96/b*exp(3*x)-1/24/b^2*exp(3*x)*a+1/2/b^3*exp(x)*a^2-3/8*a/b^2*exp(x)+5/16/b*exp(x)-1/2/b^3*exp(-x)*a^2+3/8*a/b^2*exp(-x)-5/16/b*exp(-x)-5/96/b*exp(-3*x)+1/24/b^2*exp(-3*x)*a-1/160/b*exp(-5*x)-1/2/(-a*b-b^2)^(1/2)*a^3/b^3*ln(exp(2*x)+2*(a+b)/(-a*b-b^2)^(1/2)*exp(x)-1)+1/2/(-a*b-b^2)^(1/2)*a^3/b^3*ln(exp(2*x)-2*(a+b)/(-a*b-b^2)^(1/2)*exp(x)-1)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1232 vs. 2(66) = 132.

Time = 0.28 (sec) , antiderivative size = 2508, normalized size of antiderivative = 32.15

$$\int \frac{\cosh^7(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(cosh(x)^7/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/480*(3*(a*b^3 + b^4)*cosh(x)^10 + 30*(a*b^3 + b^4)*cosh(x)*sinh(x)^9 + 3*(a*b^3 + b^4)*sinh(x)^10 - 5*(4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x)^8 - 5*(4*a^2*b^2 - a*b^3 - 5*b^4 - 27*(a*b^3 + b^4)*cosh(x)^2)*sinh(x)^8 + 40*(9*(a*b^3 + b^4)*cosh(x)^3 - (4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x))*sinh(x)^7 + 30*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*cosh(x)^6 + 10*(63*(a*b^3 + b^4)*cosh(x)^4 + 24*a^3*b + 6*a^2*b^2 - 3*a*b^3 + 15*b^4 - 14*(4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x)^2)*sinh(x)^6 + 4*(189*(a*b^3 + b^4)*cosh(x)^5 - 70*(4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x)^3 + 45*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*cosh(x))*sinh(x)^5 - 30*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*cosh(x)^4 + 10*(63*(a*b^3 + b^4)*cosh(x)^6 - 35*(4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x)^4 - 24*a^3*b - 6*a^2*b^2 + 3*a*b^3 - 15*b^4 + 45*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*cosh(x)^2)*sinh(x)^4 - 3*a*b^3 - 3*b^4 + 40*(9*(a*b^3 + b^4)*cosh(x)^7 - 7*(4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x)^5 + 15*(8*a^3*b + 2*a^2*b^2 - a*b^3

$$\begin{aligned}
& + 5*b^4)*\cosh(x)^3 - 3*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x))*\sinh \\
& (x)^3 + 5*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^2 + 5*(27*(a*b^3 + b^4)*\cosh(x) \\
& ^8 - 28*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^6 + 90*(8*a^3*b + 2*a^2*b^2 - \\
& a*b^3 + 5*b^4)*\cosh(x)^4 + 4*a^2*b^2 - a*b^3 - 5*b^4 - 36*(8*a^3*b + 2*a^2 \\
& *b^2 - a*b^3 + 5*b^4)*\cosh(x)^2)*\sinh(x)^2 - 240*(a^3*\cosh(x)^5 + 5*a^3*\cos \\
& h(x)^4*\sinh(x) + 10*a^3*\cosh(x)^3*\sinh(x)^2 + 10*a^3*\cosh(x)^2*\sinh(x)^3 + \\
& 5*a^3*\cosh(x)*\sinh(x)^4 + a^3*\sinh(x)^5)*\sqrt{-a*b - b^2}*\log((b*\cosh(x)^4 \\
& + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a + 3*b)*\cosh(x)^2 + 2*(3*b*\co \\
& sh(x)^2 - 2*a - 3*b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a + 3*b)*\cosh(x))*\sinh \\
& (x) + 4*(\cosh(x)^3 + 3*\cosh(x)*\sinh(x)^2 + \sinh(x)^3 + (3*\cosh(x)^2 - 1)*\si \\
& nh(x) - \cosh(x))*\sqrt{-a*b - b^2} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 \\
& + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x) \\
&)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 10*(3*(a*b^3 + b^ \\
& 4)*\cosh(x)^9 - 4*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^7 + 18*(8*a^3*b + 2*a^ \\
& 2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^5 - 12*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4) \\
& *\cosh(x)^3 + (4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x))*\sinh(x))/((a*b^4 + b^5)*c \\
& osh(x)^5 + 5*(a*b^4 + b^5)*\cosh(x)^4*\sinh(x) + 10*(a*b^4 + b^5)*\cosh(x)^3*s \\
& inh(x)^2 + 10*(a*b^4 + b^5)*\cosh(x)^2*\sinh(x)^3 + 5*(a*b^4 + b^5)*\cosh(x)*s \\
& inh(x)^4 + (a*b^4 + b^5)*\sinh(x)^5), 1/480*(3*(a*b^3 + b^4)*\cosh(x)^10 + 30 \\
& *(a*b^3 + b^4)*\cosh(x)*\sinh(x)^9 + 3*(a*b^3 + b^4)*\sinh(x)^10 - 5*(4*a^2*b^ \\
& 2 - a*b^3 - 5*b^4)*\cosh(x)^8 - 5*(4*a^2*b^2 - a*b^3 - 5*b^4 - 27*(a*b^3 + b \\
& ^4)*\cosh(x)^2)*\sinh(x)^8 + 40*(9*(a*b^3 + b^4)*\cosh(x)^3 - (4*a^2*b^2 - a*b \\
& ^3 - 5*b^4)*\cosh(x))*\sinh(x)^7 + 30*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*c \\
& osh(x)^6 + 10*(63*(a*b^3 + b^4)*\cosh(x)^4 + 24*a^3*b + 6*a^2*b^2 - 3*a*b^3 \\
& + 15*b^4 - 14*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^2)*\sinh(x)^6 + 4*(189*(a* \\
& b^3 + b^4)*\cosh(x)^5 - 70*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^3 + 45*(8*a^3 \\
& *b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x))*\sinh(x)^5 - 30*(8*a^3*b + 2*a^2*b^ \\
& 2 - a*b^3 + 5*b^4)*\cosh(x)^4 + 10*(63*(a*b^3 + b^4)*\cosh(x)^6 - 35*(4*a^2*b \\
& ^2 - a*b^3 - 5*b^4)*\cosh(x)^4 - 24*a^3*b - 6*a^2*b^2 + 3*a*b^3 - 15*b^4 + 4 \\
& 5*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^2)*\sinh(x)^4 - 3*a*b^3 - 3* \\
& b^4 + 40*(9*(a*b^3 + b^4)*\cosh(x)^7 - 7*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x) \\
& ^5 + 15*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^3 - 3*(8*a^3*b + 2*a^ \\
& 2*b^2 - a*b^3 + 5*b^4)*\cosh(x))*\sinh(x)^3 + 5*(4*a^2*b^2 - a*b^3 - 5*b^4)*c \\
& osh(x)^2 + 5*(27*(a*b^3 + b^4)*\cosh(x)^8 - 28*(4*a^2*b^2 - a*b^3 - 5*b^4)*c \\
& osh(x)^6 + 90*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^4 + 4*a^2*b^2 - \\
& a*b^3 - 5*b^4 - 36*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^2)*\sinh(x) \\
&)^2 - 480*(a^3*\cosh(x)^5 + 5*a^3*\cosh(x)^4*\sinh(x) + 10*a^3*\cosh(x)^3*\sinh(x) \\
& ^2 + 10*a^3*\cosh(x)^2*\sinh(x)^3 + 5*a^3*\cosh(x)*\sinh(x)^4 + a^3*\sinh(x)^5 \\
&)*\sqrt{a*b + b^2}*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x) \\
& ^3 + (4*a + 3*b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + 3*b)*\sinh(x))/\sqrt{a*b \\
& + b^2})) - 480*(a^3*\cosh(x)^5 + 5*a^3*\cosh(x)^4*\sinh(x) + 10*a^3*\cosh(x)^3*s \\
& inh(x)^2 + 10*a^3*\cosh(x)^2*\sinh(x)^3 + 5*a^3*\cosh(x)*\sinh(x)^4 + a^3*\sinh(x) \\
& ^5)*\sqrt{a*b + b^2}*\arctan(1/2*\sqrt{a*b + b^2}*(\cosh(x) + \sinh(x))/(a + b \\
&)) + 10*(3*(a*b^3 + b^4)*\cosh(x)^9 - 4*(4*a^2*b^2 - a*b^3 - 5*b^4)*\cosh(x)^ \\
& 7 + 18*(8*a^3*b + 2*a^2*b^2 - a*b^3 + 5*b^4)*\cosh(x)^5 - 12*(8*a^3*b + 2*a^
\end{aligned}$$

```
2*b^2 - a*b^3 + 5*b^4)*cosh(x)^3 + (4*a^2*b^2 - a*b^3 - 5*b^4)*cosh(x))*sinh(x))/((a*b^4 + b^5)*cosh(x)^5 + 5*(a*b^4 + b^5)*cosh(x)^4*sinh(x) + 10*(a*b^4 + b^5)*cosh(x)^3*sinh(x)^2 + 10*(a*b^4 + b^5)*cosh(x)^2*sinh(x)^3 + 5*(a*b^4 + b^5)*cosh(x)*sinh(x)^4 + (a*b^4 + b^5)*sinh(x)^5]
```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^7(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

```
[In] integrate(cosh(x)**7/(a+b*cosh(x)**2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \frac{\cosh^7(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^7}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(cosh(x)^7/(a+b*cosh(x)^2),x, algorithm="maxima")
```

```
[Out] 1/480*(3*b^2*e^(10*x) - 3*b^2 - 5*(4*a*b - 5*b^2)*e^(8*x) + 30*(8*a^2 - 6*a*b + 5*b^2)*e^(6*x) - 30*(8*a^2 - 6*a*b + 5*b^2)*e^(4*x) + 5*(4*a*b - 5*b^2)*e^(2*x))*e^(-5*x)/b^3 - 1/128*integrate(256*(a^3*e^(3*x) + a^3*e^x)/(b^4*e^(4*x) + b^4 + 2*(2*a*b^3 + b^4)*e^(2*x)), x)
```

Giac [F]

$$\int \frac{\cosh^7(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^7}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(cosh(x)^7/(a+b*cosh(x)^2),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [B] (verification not implemented)

Time = 2.34 (sec) , antiderivative size = 293, normalized size of antiderivative = 3.76

$$\int \frac{\cosh^7(x)}{a + b \cosh^2(x)} dx = \frac{e^{5x}}{160b} - \frac{e^{-5x}}{160b} - \frac{e^{-x}(8a^2 - 6ab + 5b^2)}{16b^3} + \frac{\left(2 \operatorname{atan} \left(\frac{(b^9 \sqrt{b^8 + ab^7} + ab^8 \sqrt{b^8 + ab^7}) \left(e^x \left(\frac{2a^7}{b^{11}(a+b)^2 \sqrt{a^6}} - \frac{4(2a^4 b^4 \sqrt{a^6} + 2a^5 b^3 \sqrt{a^6})}{a^3 b^8 (a+b) \sqrt{b^7 (a+b) \sqrt{b^8 + ab^7}}} \right) - \frac{2a^7 e^{3x}}{b^{11}(a+b)^2 \sqrt{a^6}} \right)}{4a^4} \right) - 2 \operatorname{atan} \left(\frac{a^3 e^x}{2b^3} \right)}{2\sqrt{b^8 + ab^7}} + \frac{e^{-3x}(4a - 5b)}{96b^2} - \frac{e^{3x}(4a - 5b)}{96b^2} + \frac{e^x(8a^2 - 6ab + 5b^2)}{16b^3}$$

[In] int(cosh(x)^7/(a + b*cosh(x)^2),x)

[Out] $\exp(5x)/(160*b) - \exp(-5x)/(160*b) - (\exp(-x)*(8*a^2 - 6*a*b + 5*b^2))/(16*b^3) + ((2*\operatorname{atan}(((b^9*(a*b^7 + b^8))^{1/2} + a*b^8*(a*b^7 + b^8))^{1/2}))*(\exp(x)*((2*a^7)/(b^{11}*(a + b)^2*(a^6)^{1/2}) - (4*(2*a^4*b^4*(a^6)^{1/2} + 2*a^5*b^3*(a^6)^{1/2}))/((a^3*b^8*(a + b)*(b^7*(a + b))^{1/2}*(a*b^7 + b^8)^{1/2}))) - (2*a^7*\exp(3*x))/(b^{11}*(a + b)^2*(a^6)^{1/2}))/((4*a^4)) - 2*\operatorname{atan}((a^3*\exp(x)*(b^7*(a + b))^{1/2})/(2*b^3*(a + b)*(a^6)^{1/2}))*((a^6)^{1/2})/(2*(a*b^7 + b^8)^{1/2})) + (\exp(-3*x)*(4*a - 5*b))/(96*b^2) - (\exp(3*x)*(4*a - 5*b))/(96*b^2) + (\exp(x)*(8*a^2 - 6*a*b + 5*b^2))/(16*b^3)$

3.21 $\int \frac{\cosh^6(x)}{a+b \cosh^2(x)} dx$

Optimal result	165
Rubi [A] (verified)	165
Mathematica [A] (verified)	167
Maple [B] (verified)	168
Fricas [B] (verification not implemented)	168
Sympy [F(-1)]	169
Maxima [B] (verification not implemented)	170
Giac [B] (verification not implemented)	171
Mupad [B] (verification not implemented)	171

Optimal result

Integrand size = 15, antiderivative size = 88

$$\int \frac{\cosh^6(x)}{a+b \cosh^2(x)} dx = \frac{(8a^2 - 4ab + 3b^2)x}{8b^3} - \frac{a^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^3 \sqrt{a+b}} - \frac{(4a - 3b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh^3(x) \sinh(x)}{4b}$$

[Out] 1/8*(8*a^2-4*a*b+3*b^2)*x/b^3-1/8*(4*a-3*b)*cosh(x)*sinh(x)/b^2+1/4*cosh(x)^3*sinh(x)/b-a^(5/2)*arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))/b^3/(a+b)^(1/2)

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3266, 481, 592, 536, 212, 214}

$$\int \frac{\cosh^6(x)}{a+b \cosh^2(x)} dx = -\frac{a^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^3 \sqrt{a+b}} + \frac{x(8a^2 - 4ab + 3b^2)}{8b^3} - \frac{(4a - 3b) \sinh(x) \cosh(x)}{8b^2} + \frac{\sinh(x) \cosh^3(x)}{4b}$$

[In] Int[Cosh[x]^6/(a + b*Cosh[x]^2),x]

[Out] ((8*a^2 - 4*a*b + 3*b^2)*x)/(8*b^3) - (a^(5/2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(b^3*Sqrt[a + b]) - ((4*a - 3*b)*Cosh[x]*Sinh[x])/(8*b^2) + (Cosh[x]^3*Sinh[x])/(4*b)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 481

Int[((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-a)*e^(2*n - 1)*(e*x)^(m - 2*n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(b*n*(b*c - a*d)*(p + 1))), x] + Dist[e^(2*n)/(b*n*(b*c - a*d)*(p + 1)), Int[(e*x)^(m - 2*n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[a*c*(m - 2*n + 1) + (a*d*(m - n + n*q + 1) + b*c*n*(p + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m - n + 1, n] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 536

Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 592

Int[((g_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[g^(n - 1)*(b*e - a*f)*(g*x)^(m - n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(b*n*(b*c - a*d)*(p + 1))), x] - Dist[g^n/(b*n*(b*c - a*d)*(p + 1)), Int[(g*x)^(m - n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f)*(m - n + 1) + (d*(b*e - a*f)*(m + n*q + 1) - b*n*(c*f - d*e)*(p + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, q}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m - n + 1, 0]

Rule 3266

Int[sin[(e_) + (f_.)*(x_)]^(m_)*((a_) + (b_.)*sin[(e_) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\text{Subst}\left(\int \frac{x^6}{(1-x^2)^3(a-(a+b)x^2)} dx, x, \coth(x)\right) \\
&= \frac{\cosh^3(x) \sinh(x)}{4b} - \frac{\text{Subst}\left(\int \frac{x^2(3a+(a-3b)x^2)}{(1-x^2)^2(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{4b} \\
&= -\frac{(4a-3b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh^3(x) \sinh(x)}{4b} \\
&\quad + \frac{\text{Subst}\left(\int \frac{-a(4a-3b)+(-4a^2+ab-3b^2)x^2}{(1-x^2)(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{8b^2} \\
&= -\frac{(4a-3b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh^3(x) \sinh(x)}{4b} \\
&\quad - \frac{a^3 \text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{b^3} \\
&\quad + \frac{(8a^2-4ab+3b^2) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(x)\right)}{8b^3} \\
&= \frac{(8a^2-4ab+3b^2)x}{8b^3} - \frac{a^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^3 \sqrt{a+b}} \\
&\quad - \frac{(4a-3b) \cosh(x) \sinh(x)}{8b^2} + \frac{\cosh^3(x) \sinh(x)}{4b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.16 (sec) , antiderivative size = 76, normalized size of antiderivative = 0.86

$$\begin{aligned}
&\int \frac{\cosh^6(x)}{a+b \cosh^2(x)} dx \\
&= \frac{4(8a^2-4ab+3b^2)x - \frac{32a^{5/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}} - 8(a-b)b \sinh(2x) + b^2 \sinh(4x)}{32b^3}
\end{aligned}$$

[In] Integrate[Cosh[x]^6/(a + b*Cosh[x]^2), x]

[Out] (4*(8*a^2 - 4*a*b + 3*b^2)*x - (32*a^(5/2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a + b] - 8*(a - b)*b*Sinh[2*x] + b^2*Sinh[4*x])/(32*b^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(74) = 148$.

Time = 0.84 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.93

method	result
risch	$\frac{x a^2}{b^3} - \frac{ax}{2b^2} + \frac{3x}{8b} + \frac{e^{4x}}{64b} - \frac{ae^{2x}}{8b^2} + \frac{e^{2x}}{8b} + \frac{ae^{-2x}}{8b^2} - \frac{e^{-2x}}{8b} - \frac{e^{-4x}}{64b} + \frac{\sqrt{a(a+b)} a^2 \ln\left(\frac{e^{2x} + 2\sqrt{a(a+b)+2a+b}}{b}\right) - \sqrt{a(a+b)}}{2(a+b)b^3}$
default	$\frac{2a^3 \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln\left(-\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a-\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} \right)}{b^3} - \frac{1}{4b(\tanh\left(\frac{x}{2}\right)+1)^4} + \frac{1}{2b(\tanh\left(\frac{x}{2}\right))}$

[In] `int(cosh(x)^6/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $x/b^3 a^2 - 1/2 a x/b^2 + 3/8 x/b + 1/64/b \exp(4x) - 1/8/b^2 a \exp(2x) + 1/8/b \exp(2x) + 1/8/b^2 a \exp(-2x) - 1/8/b \exp(-2x) - 1/64/b \exp(-4x) + 1/2 (a(a+b))^{1/2} / (a+b) a^2/b^3 \ln(\exp(2x) + (2(a(a+b))^{1/2} + 2a+b)/b) - 1/2 (a(a+b))^{1/2} / (a+b) a^2/b^3 \ln(\exp(2x) - (2(a(a+b))^{1/2} - 2a-b)/b)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 524 vs. $2(74) = 148$.

Time = 0.28 (sec) , antiderivative size = 1245, normalized size of antiderivative = 14.15

$$\int \frac{\cosh^6(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] `integrate(cosh(x)^6/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/64*(b^2*\cosh(x)^8 + 8*b^2*\cosh(x)*\sinh(x)^7 + b^2*\sinh(x)^8 - 8*(a*b - b^2)*\cosh(x)^6 + 4*(7*b^2*\cosh(x)^2 - 2*a*b + 2*b^2)*\sinh(x)^6 + 8*(8*a^2 - 4*a*b + 3*b^2)*x*\cosh(x)^4 + 8*(7*b^2*\cosh(x)^3 - 6*(a*b - b^2)*\cosh(x))*\sinh(x)^5 + 2*(35*b^2*\cosh(x)^4 - 60*(a*b - b^2)*\cosh(x)^2 + 4*(8*a^2 - 4*a*b + 3*b^2)*x)*\sinh(x)^4 + 8*(7*b^2*\cosh(x)^5 - 20*(a*b - b^2)*\cosh(x)^3 + 4*(8*a^2 - 4*a*b + 3*b^2)*x*\cosh(x))*\sinh(x)^3 + 8*(a*b - b^2)*\cosh(x)^2 + 4*(7*b^2*\cosh(x)^6 - 30*(a*b - b^2)*\cosh(x)^4 + 12*(8*a^2 - 4*a*b + 3*b^2)*x*\cosh(x)^2 + 2*a*b - 2*b^2)*\sinh(x)^2 + 32*(a^2*\cosh(x)^4 + 4*a^2*\cosh(x)^3*\sinh(x) + 6*a^2*\cosh(x)^2*\sinh(x)^2 + 4*a^2*\cosh(x)*\sinh(x)^3 + a^2*\sinh(x)^4)*\sqrt{a/(a + b)}*\log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x)^2 + 2*(3*b^2*\cosh(x)^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) + 4*((a*b + b^2)*\cosh(x)^2 + 2*(a*b + b^2)*\cosh(x)*\sinh(x) + (a*b + b^2)*\sinh(x)^2 + 2*a^2 + 3*a*b + b^2)*\sqrt{a/(a + b)})/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*$


```

a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) - b^2
+ 8*(b^2*cosh(x)^7 - 6*(a*b - b^2)*cosh(x)^5 + 4*(8*a^2 - 4*a*b + 3*b^2)*x*
cosh(x)^3 + 2*(a*b - b^2)*cosh(x))*sinh(x))/(b^3*cosh(x)^4 + 4*b^3*cosh(x)^
3*sinh(x) + 6*b^3*cosh(x)^2*sinh(x)^2 + 4*b^3*cosh(x)*sinh(x)^3 + b^3*sinh(
x)^4), 1/64*(b^2*cosh(x)^8 + 8*b^2*cosh(x)*sinh(x)^7 + b^2*sinh(x)^8 - 8*(a
*b - b^2)*cosh(x)^6 + 4*(7*b^2*cosh(x)^2 - 2*a*b + 2*b^2)*sinh(x)^6 + 8*(8*
a^2 - 4*a*b + 3*b^2)*x*cosh(x)^4 + 8*(7*b^2*cosh(x)^3 - 6*(a*b - b^2)*cosh(
x))*sinh(x)^5 + 2*(35*b^2*cosh(x)^4 - 60*(a*b - b^2)*cosh(x)^2 + 4*(8*a^2 -
4*a*b + 3*b^2)*x)*sinh(x)^4 + 8*(7*b^2*cosh(x)^5 - 20*(a*b - b^2)*cosh(x)^
3 + 4*(8*a^2 - 4*a*b + 3*b^2)*x*cosh(x))*sinh(x)^3 + 8*(a*b - b^2)*cosh(x)^
2 + 4*(7*b^2*cosh(x)^6 - 30*(a*b - b^2)*cosh(x)^4 + 12*(8*a^2 - 4*a*b + 3*b
^2)*x*cosh(x)^2 + 2*a*b - 2*b^2)*sinh(x)^2 - 64*(a^2*cosh(x)^4 + 4*a^2*cosh
(x)^3*sinh(x) + 6*a^2*cosh(x)^2*sinh(x)^2 + 4*a^2*cosh(x)*sinh(x)^3 + a^2*s
inh(x)^4)*sqrt(-a/(a + b))*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) +
b*sinh(x)^2 + 2*a + b)*sqrt(-a/(a + b)))/a - b^2 + 8*(b^2*cosh(x)^7 - 6*(a*
b - b^2)*cosh(x)^5 + 4*(8*a^2 - 4*a*b + 3*b^2)*x*cosh(x)^3 + 2*(a*b - b^2)*
cosh(x))*sinh(x))/(b^3*cosh(x)^4 + 4*b^3*cosh(x)^3*sinh(x) + 6*b^3*cosh(x)^
2*sinh(x)^2 + 4*b^3*cosh(x)*sinh(x)^3 + b^3*sinh(x)^4)]

```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^6(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

```
[In] integrate(cosh(x)**6/(a+b*cosh(x)**2),x)
```

```
[Out] Timed out
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 651 vs. 2(74) = 148.

Time = 0.32 (sec) , antiderivative size = 651, normalized size of antiderivative = 7.40

$$\int \frac{\cosh^6(x)}{a + b \cosh^2(x)} dx = -\frac{15(2a + b) \log\left(\frac{be^{2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab}}$$

$$- \frac{5 \log\left(\frac{be^{-2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{-2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)a}} - \frac{3(2a + b)x}{2b^2}$$

$$+ \frac{15x}{16b} - \frac{(4(2a + b)e^{-2x} - b)e^{4x}}{64b^2} + \frac{3e^{2x}}{16b}$$

$$- \frac{3e^{-2x}}{16b} + \frac{(4(2a + b)e^{2x} - b)e^{-4x}}{64b^2}$$

$$+ \frac{3(2a + b) \log\left(\frac{be^{4x} + 2(2a + b)e^{2x} + b}{16b^2}\right)}{16b^2}$$

$$- \frac{3(2a + b) \log\left(\frac{2(2a + b)e^{-2x} + be^{-4x} + b}{16b^2}\right)}{16b^2}$$

$$+ \frac{3(8a^2 + 8ab + b^2) \log\left(\frac{be^{2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab^2}}$$

$$- \frac{3(8a^2 + 8ab + b^2) \log\left(\frac{be^{-2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{-2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{64\sqrt{(a+b)ab^2}}$$

$$+ \frac{(16a^2 + 16ab + 3b^2)x}{8b^3}$$

$$- \frac{(16a^2 + 16ab + 3b^2) \log\left(\frac{be^{4x} + 2(2a + b)e^{2x} + b}{64b^3}\right)}{64b^3}$$

$$+ \frac{(16a^2 + 16ab + 3b^2) \log\left(\frac{2(2a + b)e^{-2x} + be^{-4x} + b}{64b^3}\right)}{64b^3}$$

$$- \frac{(32a^3 + 48a^2b + 18ab^2 + b^3) \log\left(\frac{be^{2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{128\sqrt{(a+b)ab^3}}$$

$$+ \frac{(32a^3 + 48a^2b + 18ab^2 + b^3) \log\left(\frac{be^{-2x} + 2a + b - 2\sqrt{(a+b)a}}{be^{-2x} + 2a + b + 2\sqrt{(a+b)a}}\right)}{128\sqrt{(a+b)ab^3}}$$

[In] integrate(cosh(x)^6/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] -15/64*(2*a + b)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a)*b - 5/32*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a) - 3/2*(2*a + b)*x/b^2 + 15/16*x/b - 1/64*(4*(2*a + b)*e^(-2*x) - b)*e^(4*x)/b^2 + 3/16*e^(2*x)/b - 3/16*e^(-2*x)/b + 1/64*(4*(2*a + b)*

$$e^{(2*x) - b}*e^{(-4*x)}/b^2 + 3/16*(2*a + b)*\log(b*e^{(4*x) + 2*(2*a + b)*e^{(2*x) + b)}/b^2 - 3/16*(2*a + b)*\log(2*(2*a + b)*e^{(-2*x) + b*e^{(-4*x) + b)}/b^2 + 3/64*(8*a^2 + 8*a*b + b^2)*\log((b*e^{(2*x) + 2*a + b - 2*\sqrt{(a + b)*a)}}/(b*e^{(2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})))/(\sqrt{(a + b)*a}*b^2) - 3/64*(8*a^2 + 8*a*b + b^2)*\log((b*e^{(-2*x) + 2*a + b - 2*\sqrt{(a + b)*a)}}/(b*e^{(-2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})))/(\sqrt{(a + b)*a}*b^2) + 1/8*(16*a^2 + 16*a*b + 3*b^2)*x/b^3 - 1/64*(16*a^2 + 16*a*b + 3*b^2)*\log(b*e^{(4*x) + 2*(2*a + b)*e^{(2*x) + b)}/b^3 + 1/64*(16*a^2 + 16*a*b + 3*b^2)*\log(2*(2*a + b)*e^{(-2*x) + b*e^{(-4*x) + b)}/b^3 - 1/128*(32*a^3 + 48*a^2*b + 18*a*b^2 + b^3)*\log((b*e^{(2*x) + 2*a + b - 2*\sqrt{(a + b)*a)}}/(b*e^{(2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})))/(\sqrt{(a + b)*a}*b^3) + 1/128*(32*a^3 + 48*a^2*b + 18*a*b^2 + b^3)*\log((b*e^{(-2*x) + 2*a + b - 2*\sqrt{(a + b)*a)}}/(b*e^{(-2*x) + 2*a + b + 2*\sqrt{(a + b)*a}})))/(\sqrt{(a + b)*a}*b^3)$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 150 vs. $2(74) = 148$.

Time = 0.27 (sec) , antiderivative size = 150, normalized size of antiderivative = 1.70

$$\int \frac{\cosh^6(x)}{a + b \cosh^2(x)} dx$$

$$= -\frac{a^3 \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - ab}b^3} + \frac{be^{(4x)} - 8ae^{(2x)} + 8be^{(2x)}}{64b^2} + \frac{(8a^2 - 4ab + 3b^2)x}{8b^3}$$

$$- \frac{(48a^2e^{(4x)} - 24abe^{(4x)} + 18b^2e^{(4x)} - 8abe^{(2x)} + 8b^2e^{(2x)} + b^2)e^{(-4x)}}{64b^3}$$

[In] integrate(cosh(x)^6/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $-a^3*\arctan(1/2*(b*e^{(2*x) + 2*a + b)}/\sqrt{-a^2 - a*b}))/(\sqrt{-a^2 - a*b})*b^3 + 1/64*(b*e^{(4*x) - 8*a*e^{(2*x) + 8*b*e^{(2*x)}})/b^2 + 1/8*(8*a^2 - 4*a*b + 3*b^2)*x/b^3 - 1/64*(48*a^2*e^{(4*x) - 24*a*b*e^{(4*x) + 18*b^2*e^{(4*x) - 8*a*b*e^{(2*x) + 8*b^2*e^{(2*x) + b^2}})*e^{(-4*x)}/b^3$

Mupad [B] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 178, normalized size of antiderivative = 2.02

$$\int \frac{\cosh^6(x)}{a + b \cosh^2(x)} dx = \frac{e^{4x}}{64b} - \frac{e^{-4x}}{64b} + \frac{x(8a^2 - 4ab + 3b^2)}{8b^3} + \frac{e^{-2x}(a - b)}{8b^2}$$

$$- \frac{e^{2x}(a - b)}{8b^2} + \frac{a^{5/2} \ln\left(\frac{4a^3 e^{2x}}{b^4} - \frac{2a^{5/2}(b + 2ae^{2x} + be^{2x})}{b^4 \sqrt{a+b}}\right)}{2b^3 \sqrt{a+b}}$$

$$- \frac{a^{5/2} \ln\left(\frac{4a^3 e^{2x}}{b^4} + \frac{2a^{5/2}(b + 2ae^{2x} + be^{2x})}{b^4 \sqrt{a+b}}\right)}{2b^3 \sqrt{a+b}}$$

[In] `int(cosh(x)^6/(a + b*cosh(x)^2),x)`

[Out] $\frac{\exp(4x)}{64b} - \frac{\exp(-4x)}{64b} + \frac{x(8a^2 - 4ab + 3b^2)}{8b^3} + \frac{\exp(-2x)(a-b)}{8b^2} - \frac{\exp(2x)(a-b)}{8b^2} + \frac{a^{5/2} \log\left(\frac{4a^3 \exp(2x)}{b^4} - \frac{2a^{5/2}(b + 2a \exp(2x) + b \exp(2x))}{b^4(a+b)^{1/2}}\right)}{2b^3(a+b)^{1/2}} - \frac{a^{5/2} \log\left(\frac{4a^3 \exp(2x)}{b^4} + \frac{2a^{5/2}(b + 2a \exp(2x) + b \exp(2x))}{b^4(a+b)^{1/2}}\right)}{2b^3(a+b)^{1/2}}$

3.22 $\int \frac{\cosh^5(x)}{a+b \cosh^2(x)} dx$

Optimal result	173
Rubi [A] (verified)	173
Mathematica [A] (verified)	174
Maple [B] (verified)	175
Fricas [B] (verification not implemented)	175
Sympy [F(-1)]	176
Maxima [F]	176
Giac [F]	177
Mupad [B] (verification not implemented)	177

Optimal result

Integrand size = 15, antiderivative size = 56

$$\int \frac{\cosh^5(x)}{a+b \cosh^2(x)} dx = \frac{a^2 \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{5/2} \sqrt{a+b}} - \frac{(a-b) \sinh(x)}{b^2} + \frac{\sinh^3(x)}{3b}$$

[Out] $-(a-b)*\sinh(x)/b^2+1/3*\sinh(x)^3/b+a^2*\arctan(\sinh(x)*b^{(1/2)}/(a+b)^{(1/2)})/b^{(5/2)}/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3265, 398, 211}

$$\int \frac{\cosh^5(x)}{a+b \cosh^2(x)} dx = \frac{a^2 \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{5/2} \sqrt{a+b}} - \frac{(a-b) \sinh(x)}{b^2} + \frac{\sinh^3(x)}{3b}$$

[In] Int[Cosh[x]^5/(a + b*Cosh[x]^2),x]

[Out] $(a^2*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sinh}[x])/(\text{Sqrt}[a + b])])/(b^{(5/2)}*\text{Sqrt}[a + b]) - ((a - b)*\text{Sinh}[x])/b^2 + \text{Sinh}[x]^3/(3*b)$

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 398

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Int[PolynomialDivide[(a + b*x^n)^p, (c + d*x^n)^(-q), x], x] /; FreeQ[{a
, b, c, d}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && ILtQ[q,
0] && GeQ[p, -q]
```

Rule 3265

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(
p_.), x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, S
ubst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e +
f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{(1+x^2)^2}{a+b+bx^2} dx, x, \sinh(x)\right) \\
&= \text{Subst}\left(\int \left(-\frac{a-b}{b^2} + \frac{x^2}{b} + \frac{a^2}{b^2(a+b+bx^2)}\right) dx, x, \sinh(x)\right) \\
&= -\frac{(a-b)\sinh(x)}{b^2} + \frac{\sinh^3(x)}{3b} + \frac{a^2 \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{b^2} \\
&= \frac{a^2 \arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{b^{5/2}\sqrt{a+b}} - \frac{(a-b)\sinh(x)}{b^2} + \frac{\sinh^3(x)}{3b}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.14 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.09

$$\int \frac{\cosh^5(x)}{a+b\cosh^2(x)} dx = -\frac{a^2 \arctan\left(\frac{\sqrt{a+b}\text{csch}(x)}{\sqrt{b}}\right)}{b^{5/2}\sqrt{a+b}} - \frac{(4a-3b)\sinh(x)}{4b^2} + \frac{\sinh(3x)}{12b}$$

```
[In] Integrate[Cosh[x]^5/(a + b*Cosh[x]^2), x]
```

```
[Out] -((a^2*ArcTan[(Sqrt[a + b]*Csch[x])/Sqrt[b]])/(b^(5/2)*Sqrt[a + b])) - ((4*
a - 3*b)*Sinh[x])/(4*b^2) + Sinh[3*x]/(12*b)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 145 vs. 2(46) = 92.

Time = 0.52 (sec) , antiderivative size = 146, normalized size of antiderivative = 2.61

method	result
risch	$\frac{e^{3x}}{24b} - \frac{ae^x}{2b^2} + \frac{3e^x}{8b} + \frac{ae^{-x}}{2b^2} - \frac{3e^{-x}}{8b} - \frac{e^{-3x}}{24b} - \frac{a^2 \ln\left(e^{2x} - \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1\right)}{2\sqrt{-ab-b^2}b^2} + \frac{a^2 \ln\left(e^{2x} + \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1\right)}{2\sqrt{-ab-b^2}b^2}$
default	$\frac{2a^2 \left(\frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right)\sqrt{a+b+2\sqrt{a}}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right)\sqrt{a+b-2\sqrt{a}}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} \right)}{b^2} - \frac{1}{3b(\tanh(\frac{x}{2})+1)^3} + \frac{1}{2b(\tanh(\frac{x}{2})+1)^2} - \frac{-a+b}{b^2(\tanh(\frac{x}{2})+1)}$

[In] int(cosh(x)^5/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/24/b*exp(3*x)-1/2*a/b^2*exp(x)+3/8/b*exp(x)+1/2*a/b^2*exp(-x)-3/8/b*exp(-x)-1/24/b*exp(-3*x)-1/2/(-a*b-b^2)^(1/2)*a^2/b^2*ln(exp(2*x)-2*(a+b)/(-a*b-b^2)^(1/2)*exp(x)-1)+1/2/(-a*b-b^2)^(1/2)*a^2/b^2*ln(exp(2*x)+2*(a+b)/(-a*b-b^2)^(1/2)*exp(x)-1)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 557 vs. 2(46) = 92.

Time = 0.28 (sec) , antiderivative size = 1184, normalized size of antiderivative = 21.14

$$\int \frac{\cosh^5(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(cosh(x)^5/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/24*((a*b^2 + b^3)*cosh(x)^6 + 6*(a*b^2 + b^3)*cosh(x)*sinh(x)^5 + (a*b^2 + b^3)*sinh(x)^6 - 3*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x)^4 - 3*(4*a^2*b + a*b^2 - 3*b^3 - 5*(a*b^2 + b^3)*cosh(x)^2)*sinh(x)^4 + 4*(5*(a*b^2 + b^3)*cosh(x)^3 - 3*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x))*sinh(x)^3 - a*b^2 - b^3 + 3*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x)^2 + 3*(5*(a*b^2 + b^3)*cosh(x)^4 + 4*a^2*b + a*b^2 - 3*b^3 - 6*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x)^2)*sinh(x)^2 - 12*(a^2*cosh(x)^3 + 3*a^2*cosh(x)^2*sinh(x) + 3*a^2*cosh(x)*sinh(x)^2 + a^2*sinh(x)^3)*sqrt(-a*b - b^2)*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a + 3*b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*a - 3*b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a + 3*b)*cosh(x))*sinh(x) - 4*(cosh(x)^3 + 3*cosh(x)*sinh(x)^2 + sinh(x)^3 + (3*cosh(x)^2 - 1)*sinh(x) - cosh(x))*sqrt(-a*b - b^2) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 6*((a*b^2 + b^3)*cosh(x)^5 - 2*(4*a^2*b + a*b^2 - 3

```

*b^3)*cosh(x)^3 + (4*a^2*b + a*b^2 - 3*b^3)*cosh(x))*sinh(x))/((a*b^3 + b^4
)*cosh(x)^3 + 3*(a*b^3 + b^4)*cosh(x)^2*sinh(x) + 3*(a*b^3 + b^4)*cosh(x)*s
inh(x)^2 + (a*b^3 + b^4)*sinh(x)^3), 1/24*((a*b^2 + b^3)*cosh(x)^6 + 6*(a*b
^2 + b^3)*cosh(x)*sinh(x)^5 + (a*b^2 + b^3)*sinh(x)^6 - 3*(4*a^2*b + a*b^2
- 3*b^3)*cosh(x)^4 - 3*(4*a^2*b + a*b^2 - 3*b^3 - 5*(a*b^2 + b^3)*cosh(x)^2
)*sinh(x)^4 + 4*(5*(a*b^2 + b^3)*cosh(x)^3 - 3*(4*a^2*b + a*b^2 - 3*b^3)*co
sh(x))*sinh(x)^3 - a*b^2 - b^3 + 3*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x)^2 + 3*
(5*(a*b^2 + b^3)*cosh(x)^4 + 4*a^2*b + a*b^2 - 3*b^3 - 6*(4*a^2*b + a*b^2 -
3*b^3)*cosh(x)^2)*sinh(x)^2 + 24*(a^2*cosh(x)^3 + 3*a^2*cosh(x)^2*sinh(x)
+ 3*a^2*cosh(x)*sinh(x)^2 + a^2*sinh(x)^3)*sqrt(a*b + b^2)*arctan(1/2*(b*co
sh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + 3*b)*cosh(x) + (3*b*
cosh(x)^2 + 4*a + 3*b)*sinh(x))/sqrt(a*b + b^2)) + 24*(a^2*cosh(x)^3 + 3*a^
2*cosh(x)^2*sinh(x) + 3*a^2*cosh(x)*sinh(x)^2 + a^2*sinh(x)^3)*sqrt(a*b + b
^2)*arctan(1/2*sqrt(a*b + b^2)*(cosh(x) + sinh(x))/(a + b)) + 6*((a*b^2 + b
^3)*cosh(x)^5 - 2*(4*a^2*b + a*b^2 - 3*b^3)*cosh(x)^3 + (4*a^2*b + a*b^2 -
3*b^3)*cosh(x))*sinh(x))/((a*b^3 + b^4)*cosh(x)^3 + 3*(a*b^3 + b^4)*cosh(x)
^2*sinh(x) + 3*(a*b^3 + b^4)*cosh(x)*sinh(x)^2 + (a*b^3 + b^4)*sinh(x)^3)]

```

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^5(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

```
[In] integrate(cosh(x)**5/(a+b*cosh(x)**2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \frac{\cosh^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^5}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(cosh(x)^5/(a+b*cosh(x)^2),x, algorithm="maxima")
```

```
[Out] 1/24*(b*e^(6*x) - 3*(4*a - 3*b)*e^(4*x) + 3*(4*a - 3*b)*e^(2*x) - b)*e^(-3*
x)/b^2 + 1/32*integrate(64*(a^2*e^(3*x) + a^2*e^x)/(b^3*e^(4*x) + b^3 + 2*(
2*a*b^2 + b^3)*e^(2*x)), x)
```


Giac [F]

$$\int \frac{\cosh^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(cosh(x)^5/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.18 (sec) , antiderivative size = 243, normalized size of antiderivative = 4.34

$$\int \frac{\cosh^5(x)}{a + b \cosh^2(x)} dx = \frac{e^{3x}}{24b} - \frac{e^{-3x}}{24b} + \frac{e^{-x}(4a - 3b)}{8b^2}$$

$$+ \frac{\sqrt{a^4} \left(2 \operatorname{atan} \left(\frac{a^2 e^x \sqrt{b^5(a+b)}}{2b^2(a+b)\sqrt{a^4}} \right) - 2 \operatorname{atan} \left(\left(\frac{b^7 \sqrt{b^6+ab^5}}{4} + \frac{ab^6 \sqrt{b^6+ab^5}}{4} \right) \left(e^x \left(\frac{2a^2}{b^8(a+b)^2 \sqrt{a^4}} - \frac{4(2a^3 b^3 \sqrt{a^4+2a^4 b^5})}{a^5 b^6 (a+b) \sqrt{b^5(a+b)}} \right) \right) \right)}{2\sqrt{b^6+ab^5}}$$

$$- \frac{e^x(4a - 3b)}{8b^2}$$

[In] int(cosh(x)^5/(a + b*cosh(x)^2),x)

[Out] exp(3*x)/(24*b) - exp(-3*x)/(24*b) + (exp(-x)*(4*a - 3*b))/(8*b^2) + ((a^4)^(1/2)*(2*atan((a^2*exp(x)*(b^5*(a + b))^(1/2))/(2*b^2*(a + b)*(a^4)^(1/2))) - 2*atan(((b^7*(a*b^5 + b^6)^(1/2))/4 + (a*b^6*(a*b^5 + b^6)^(1/2))/4)*(exp(x)*((2*a^2)/(b^8*(a + b)^2*(a^4)^(1/2)) - (4*(2*a^3*b^3*(a^4)^(1/2) + 2*a^4*b^2*(a^4)^(1/2)))/(a^5*b^6*(a + b)*(b^5*(a + b))^(1/2)*(a*b^5 + b^6)^(1/2))) - (2*a^2*exp(3*x))/(b^8*(a + b)^2*(a^4)^(1/2)))))/(2*(a*b^5 + b^6)^(1/2)) - (exp(x)*(4*a - 3*b))/(8*b^2)

3.23 $\int \frac{\cosh^4(x)}{a+b \cosh^2(x)} dx$

Optimal result	178
Rubi [A] (verified)	178
Mathematica [A] (verified)	180
Maple [B] (verified)	180
Fricas [B] (verification not implemented)	180
Sympy [F(-1)]	181
Maxima [B] (verification not implemented)	182
Giac [B] (verification not implemented)	182
Mupad [B] (verification not implemented)	183

Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{\cosh^4(x)}{a+b \cosh^2(x)} dx = -\frac{(2a-b)x}{2b^2} + \frac{a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^2 \sqrt{a+b}} + \frac{\cosh(x) \sinh(x)}{2b}$$

[Out] $-1/2*(2*a-b)*x/b^2+1/2*\cosh(x)*\sinh(x)/b+a^{(3/2)*\operatorname{arctanh}(a^{(1/2)*\tanh(x)/(a+b)^{(1/2)})/b^2/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3266, 481, 536, 212, 214}

$$\int \frac{\cosh^4(x)}{a+b \cosh^2(x)} dx = \frac{a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^2 \sqrt{a+b}} - \frac{x(2a-b)}{2b^2} + \frac{\sinh(x) \cosh(x)}{2b}$$

[In] `Int[Cosh[x]^4/(a + b*Cosh[x]^2),x]`

[Out] $-1/2*((2*a - b)*x)/b^2 + (a^{(3/2)*\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tanh}[x])/(\operatorname{Sqrt}[a + b])])/(b^2*\operatorname{Sqrt}[a + b]) + (\operatorname{Cosh}[x]*\operatorname{Sinh}[x])/(2*b)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 481

Int[((e_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-a)*e^(2*n - 1)*(e*x)^(m - 2*n + 1)*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(b*n*(b*c - a*d)*(p + 1))), x] + Dist[e^(2*n)/(b*n*(b*c - a*d)*(p + 1)), Int[(e*x)^(m - 2*n)*(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[a*c*(m - 2*n + 1) + (a*d*(m - n + n*q + 1) + b*c*n*(p + 1))*x^n, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m - n + 1, n] && IntBinomialQ[a, b, c, d, e, m, n, p, q, x]

Rule 536

Int[((e_) + (f_)*(x_)^(n_))/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]

Rule 3266

Int[sin[(e_) + (f_)*(x_)^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)^2]^(p_)), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{x^4}{(1-x^2)^2(a-(a+b)x^2)} dx, x, \coth(x)\right) \\
 &= \frac{\cosh(x) \sinh(x)}{2b} + \frac{\text{Subst}\left(\int \frac{a+(a-b)x^2}{(1-x^2)(a+(-a-b)x^2)} dx, x, \coth(x)\right)}{2b} \\
 &= \frac{\cosh(x) \sinh(x)}{2b} + \frac{a^2 \text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{b^2} \\
 &\quad - \frac{(2a-b) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \coth(x)\right)}{2b^2} \\
 &= -\frac{(2a-b)x}{2b^2} + \frac{a^{3/2} \arctanh\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b^2 \sqrt{a+b}} + \frac{\cosh(x) \sinh(x)}{2b}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.09 (sec) , antiderivative size = 52, normalized size of antiderivative = 0.88

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = \frac{2(-2a + b)x + \frac{4a^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}} + b \sinh(2x)}{4b^2}$$

[In] Integrate[Cosh[x]^4/(a + b*Cosh[x]^2),x]

[Out] (2*(-2*a + b)*x + (4*a^(3/2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a + b] + b*Sinh[2*x])/(4*b^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(47) = 94.

Time = 0.32 (sec) , antiderivative size = 120, normalized size of antiderivative = 2.03

method	result
risch	$-\frac{ax}{b^2} + \frac{x}{2b} + \frac{e^{2x}}{8b} - \frac{e^{-2x}}{8b} + \frac{\sqrt{a(a+b)} a \ln\left(\frac{e^{2x} - 2\sqrt{a(a+b)} - 2a - b}{b}\right)}{2(a+b)b^2} - \frac{\sqrt{a(a+b)} a \ln\left(\frac{e^{2x} + 2\sqrt{a(a+b)} + 2a + b}{b}\right)}{2(a+b)b^2}$
default	$-\frac{2a^2 \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b}\right)}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln\left(-\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a-b}\right)}{4\sqrt{a}\sqrt{a+b}} \right)}{b^2} - \frac{1}{2b(\tanh\left(\frac{x}{2}\right)+1)^2} + \frac{1}{2b(\tanh\left(\frac{x}{2}\right)-1)^2}$

[In] int(cosh(x)^4/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] -a*x/b^2+1/2*x/b+1/8/b*exp(2*x)-1/8/b*exp(-2*x)+1/2*(a*(a+b))^(1/2)/(a+b)*a/b^2*ln(exp(2*x)-(2*(a*(a+b))^(1/2)-2*a-b)/b)-1/2*(a*(a+b))^(1/2)/(a+b)*a/b^2*ln(exp(2*x)+(2*(a*(a+b))^(1/2)+2*a+b)/b)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(47) = 94.

Time = 0.28 (sec) , antiderivative size = 573, normalized size of antiderivative = 9.71

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = \frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 - 4(2a - b)x \cosh(x)^2 + 2(3b \cosh(x)^2 - 2(2a - b)x) \sinh(x)}{4b^2}$$

[In] integrate(cosh(x)^4/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/8*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 4*(2*a - b)*x*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*(2*a - b)*x)*sinh(x)^2 + 4*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2)*sqrt(a/(a + b))*log((b^2*cosh(x)^4 + 4*b^2*cosh(x)*sinh(x)^3 + b^2*sinh(x)^4 + 2*(2*a*b + b^2)*cosh(x)^2 + 2*(3*b^2*cosh(x)^2 + 2*a*b + b^2)*sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*cosh(x)^3 + (2*a*b + b^2)*cosh(x))*sinh(x) - 4*((a*b + b^2)*cosh(x)^2 + 2*(a*b + b^2)*cosh(x)*sinh(x) + (a*b + b^2)*sinh(x)^2 + 2*a^2 + 3*a*b + b^2)*sqrt(a/(a + b)))/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 4*(b*cosh(x)^3 - 2*(2*a - b)*x*cosh(x))*sinh(x) - b)/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2), 1/8*(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 4*(2*a - b)*x*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*(2*a - b)*x)*sinh(x)^2 + 8*(a*cosh(x)^2 + 2*a*cosh(x)*sinh(x) + a*sinh(x)^2)*sqrt(-a/(a + b))*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-a/(a + b))/a) + 4*(b*cosh(x)^3 - 2*(2*a - b)*x*cosh(x))*sinh(x) - b)/(b^2*cosh(x)^2 + 2*b^2*cosh(x)*sinh(x) + b^2*sinh(x)^2)]

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(cosh(x)**4/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 347 vs. 2(47) = 94.

Time = 0.29 (sec) , antiderivative size = 347, normalized size of antiderivative = 5.88

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = -\frac{(2a + b) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)ab}} - \frac{3 \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{16\sqrt{(a+b)a}} - \frac{(2a + b)x}{b^2} + \frac{x}{b} + \frac{e^{(2x)}}{8b} - \frac{e^{(-2x)}}{8b} + \frac{(2a + b) \log\left(\frac{be^{(4x)} + 2(2a + b)e^{(2x)} + b}{2(2a + b)e^{(-2x)} + be^{(-4x)} + b}\right)}{8b^2} - \frac{(2a + b) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)ab^2}} + \frac{(8a^2 + 8ab + b^2) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{32\sqrt{(a+b)ab^2}}$$

[In] integrate(cosh(x)^4/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] -1/4*(2*a + b)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b) - 3/16*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a) - (2*a + b)*x/b^2 + x/b + 1/8*e^(2*x)/b - 1/8*e^(-2*x)/b + 1/8*(2*a + b)*log(b*e^(4*x) + 2*(2*a + b)*e^(2*x) + b)/b^2 - 1/8*(2*a + b)*log(2*(2*a + b)*e^(-2*x) + b*e^(-4*x) + b)/b^2 + 1/32*(8*a^2 + 8*a*b + b^2)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b^2) - 1/32*(8*a^2 + 8*a*b + b^2)*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b^2)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 95 vs. 2(47) = 94.

Time = 0.31 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.61

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = \frac{a^2 \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - abb^2}}\right)}{\sqrt{-a^2 - abb^2}} - \frac{(2a - b)x}{2b^2} + \frac{e^{(2x)}}{8b} + \frac{(4ae^{(2x)} - 2be^{(2x)} - b)e^{(-2x)}}{8b^2}$$

[In] integrate(cosh(x)^4/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $a^2 \arctan(1/2*(b*e^{2*x} + 2*a + b)/\sqrt{-a^2 - a*b})/(\sqrt{-a^2 - a*b})*b^2 - 1/2*(2*a - b)*x/b^2 + 1/8*e^{2*x}/b + 1/8*(4*a*e^{2*x} - 2*b*e^{2*x} - b)*e^{-2*x}/b^2$

Mupad [B] (verification not implemented)

Time = 2.07 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.41

$$\int \frac{\cosh^4(x)}{a + b \cosh^2(x)} dx = \frac{e^{2x}}{8b} - \frac{e^{-2x}}{8b} - \frac{x(2a - b)}{2b^2} + \frac{a^{3/2} \ln\left(-\frac{4a^2 e^{2x}}{b^3} - \frac{2a^{3/2}(b + 2ae^{2x} + be^{2x})}{b^3 \sqrt{a+b}}\right)}{2b^2 \sqrt{a+b}} - \frac{a^{3/2} \ln\left(\frac{2a^{3/2}(b + 2ae^{2x} + be^{2x})}{b^3 \sqrt{a+b}} - \frac{4a^2 e^{2x}}{b^3}\right)}{2b^2 \sqrt{a+b}}$$

[In] int(cosh(x)^4/(a + b*cosh(x)^2),x)

[Out] $\exp(2*x)/(8*b) - \exp(-2*x)/(8*b) - (x*(2*a - b))/(2*b^2) + (a^{3/2})*\log(- (4*a^2*\exp(2*x))/b^3 - (2*a^{3/2})*(b + 2*a*\exp(2*x) + b*\exp(2*x)))/(b^3*(a + b)^{1/2}))/ (2*b^2*(a + b)^{1/2}) - (a^{3/2})*\log((2*a^{3/2})*(b + 2*a*\exp(2*x) + b*\exp(2*x)))/(b^3*(a + b)^{1/2}) - (4*a^2*\exp(2*x))/b^3))/ (2*b^2*(a + b)^{1/2})$

3.24 $\int \frac{\cosh^3(x)}{a+b \cosh^2(x)} dx$

Optimal result	184
Rubi [A] (verified)	184
Mathematica [A] (verified)	185
Maple [B] (verified)	185
Fricas [B] (verification not implemented)	186
Sympy [F(-1)]	187
Maxima [F]	187
Giac [F]	187
Mupad [B] (verification not implemented)	187

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \frac{\cosh^3(x)}{a+b \cosh^2(x)} dx = -\frac{a \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}} + \frac{\sinh(x)}{b}$$

[Out] $\sinh(x)/b - a \arctan(\sinh(x) * b^{(1/2)} / (a+b)^{(1/2)}) / b^{(3/2)} / (a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3265, 396, 211}

$$\int \frac{\cosh^3(x)}{a+b \cosh^2(x)} dx = \frac{\sinh(x)}{b} - \frac{a \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{b^{3/2} \sqrt{a+b}}$$

[In] $\text{Int}[\text{Cosh}[x]^3/(a + b*\text{Cosh}[x]^2), x]$

[Out] $-((a*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sinh}[x])/(\text{Sqrt}[a + b])])/(b^{(3/2)}*\text{Sqrt}[a + b])) + \text{Sinh}[x]/b$

Rule 211

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$

Rule 396

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}*((c_ + (d_)*(x_)^{(n_)})), x_Symbol] \rightarrow \text{Simp}[d*x*((a + b*x^n)^{(p+1)} / (b*(n*(p+1) + 1))), x] - \text{Dist}[(a*d - b*c*(n*($

$p + 1) + 1)) / (b * (n * (p + 1) + 1))$, Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 3265

Int[sin[(e_.) + (f_.)*(x_.)]^(m_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_.), x_Symbol] :> With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1+x^2}{a+b+bx^2} dx, x, \sinh(x)\right) \\ &= \frac{\sinh(x)}{b} - \frac{a \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{b} \\ &= -\frac{a \arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{b^{3/2}\sqrt{a+b}} + \frac{\sinh(x)}{b} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{\cosh^3(x)}{a+b\cosh^2(x)} dx = -\frac{a \arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{b^{3/2}\sqrt{a+b}} + \frac{\sinh(x)}{b}$$

[In] Integrate[Cosh[x]^3/(a + b*Cosh[x]^2), x]

[Out] -((a*ArcTan[(Sqrt[b]*Sinh[x])/Sqrt[a + b]])/(b^(3/2)*Sqrt[a + b])) + Sinh[x]/b

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(30) = 60.

Time = 0.21 (sec) , antiderivative size = 101, normalized size of antiderivative = 2.66

method	result	size
default	$-\frac{1}{b(\tanh(\frac{x}{2})+1)} - \frac{2a \left(\frac{\arctan\left(\frac{2 \tanh(\frac{x}{2})\sqrt{a+b}+2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh(\frac{x}{2})\sqrt{a+b}-2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} \right)}{b} - \frac{1}{b(\tanh(\frac{x}{2})-1)}$	101
risch	$\frac{e^x}{2b} - \frac{e^{-x}}{2b} - \frac{a \ln\left(\frac{e^{2x} + \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1}{2\sqrt{-ab-b^2}b}\right)}{2\sqrt{-ab-b^2}b} + \frac{a \ln\left(\frac{e^{2x} - \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1}{2\sqrt{-ab-b^2}b}\right)}{2\sqrt{-ab-b^2}b}$	106

[In] `int(cosh(x)^3/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $-1/b/(\tanh(1/2*x)+1)-2*a/b*(1/2/(a+b)^(1/2)/b^(1/2)*\arctan(1/2*(2*\tanh(1/2*x)*(a+b)^(1/2)+2*a^(1/2))/b^(1/2))+1/2/(a+b)^(1/2)/b^(1/2)*\arctan(1/2*(2*\tanh(1/2*x)*(a+b)^(1/2)-2*a^(1/2))/b^(1/2)))-1/b/(\tanh(1/2*x)-1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 199 vs. $2(30) = 60$.

Time = 0.28 (sec) , antiderivative size = 498, normalized size of antiderivative = 13.11

$$\int \frac{\cosh^3(x)}{a + b \cosh^2(x)} dx$$

$$= \left[\frac{(ab + b^2) \cosh(x)^2 + 2(ab + b^2) \cosh(x) \sinh(x) + (ab + b^2) \sinh(x)^2 - \sqrt{-ab - b^2}(a \cosh(x) + a \sinh(x))}{(ab + b^2) \cosh(x)^2 + 2(ab + b^2) \cosh(x) \sinh(x) + (ab + b^2) \sinh(x)^2 - \sqrt{-ab - b^2}(a \cosh(x) + a \sinh(x))} \right]$$

[In] `integrate(cosh(x)^3/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/2*((a*b + b^2)*\cosh(x)^2 + 2*(a*b + b^2)*\cosh(x)*\sinh(x) + (a*b + b^2)*\sinh(x)^2 - \sqrt{-a*b - b^2}*(a*\cosh(x) + a*\sinh(x)))*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a + 3*b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a - 3*b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a + 3*b)*\cosh(x))*\sinh(x) + 4*(\cosh(x)^3 + 3*\cosh(x)*\sinh(x)^2 + \sinh(x)^3 + (3*\cosh(x)^2 - 1)*\sinh(x) - \cosh(x))*\sqrt{-a*b - b^2} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) - a*b - b^2)/((a*b^2 + b^3)*\cosh(x) + (a*b^2 + b^3)*\sinh(x)), 1/2*((a*b + b^2)*\cosh(x)^2 + 2*(a*b + b^2)*\cosh(x)*\sinh(x) + (a*b + b^2)*\sinh(x)^2 - 2*\sqrt{a*b + b^2}*(a*\cosh(x) + a*\sinh(x)))*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x)^3 + (4*a + 3*b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + 3*b)*\sinh(x))/\sqrt{a*b + b^2}) - 2*\sqrt{a*b + b^2}*(a*\cosh(x) + a*\sinh(x))*\arctan(1/2*\sqrt{a*b + b^2}*(\cosh(x) + \sinh(x))/(a + b)) - a*b - b^2)/((a*b^2 + b^3)*\cosh(x) + (a*b^2 + b^3)*\sinh(x))]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^3(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(cosh(x)**3/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [F]

$$\int \frac{\cosh^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^3}{b \cosh(x)^2 + a} dx$$

[In] integrate(cosh(x)^3/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/2*(e^(2*x) - 1)*e^(-x)/b - 1/8*integrate(16*(a*e^(3*x) + a*e^x)/(b^2*e^(4*x) + b^2 + 2*(2*a*b + b^2)*e^(2*x)), x)

Giac [F]

$$\int \frac{\cosh^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)^3}{b \cosh(x)^2 + a} dx$$

[In] integrate(cosh(x)^3/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.06 (sec) , antiderivative size = 204, normalized size of antiderivative = 5.37

$$\int \frac{\cosh^3(x)}{a + b \cosh^2(x)} dx = \frac{e^x}{2b} - \frac{e^{-x}}{2b} - \frac{\left(2 \operatorname{atan}\left(\frac{a^3 e^x \sqrt{b^3(a+b)}}{2b(a+b)(a^2)^{3/2}}\right) - 2 \operatorname{atan}\left(\left(\frac{b^5 \sqrt{b^4+ab^3}}{4} + \frac{ab^4 \sqrt{b^4+ab^3}}{4}\right)\right) \left(e^x \left(\frac{2a^3}{b^5(a+b)^2(a^2)^{3/2}} - \frac{4(2b^2(a^2)^{3/2} + 2ab(a^2))}{a^3 b^4(a+b)\sqrt{b^3(a+b)}\sqrt{b^4+a}} \right) \right)}{2\sqrt{b^4+ab^3}}$$

[In] int(cosh(x)^3/(a + b*cosh(x)^2),x)

```
[Out] exp(x)/(2*b) - exp(-x)/(2*b) - ((2*atan((a^3*exp(x)*(b^3*(a + b))^(1/2))/(2
*b*(a + b)*(a^2)^(3/2))) - 2*atan(((b^5*(a*b^3 + b^4)^(1/2))/4 + (a*b^4*(a*
b^3 + b^4)^(1/2))/4)*(exp(x)*((2*a^3)/(b^5*(a + b)^2*(a^2)^(3/2)) - (4*(2*b
^2*(a^2)^(3/2) + 2*a*b*(a^2)^(3/2)))/(a^3*b^4*(a + b)*(b^3*(a + b))^(1/2)*(
a*b^3 + b^4)^(1/2))) - (2*a^3*exp(3*x))/(b^5*(a + b)^2*(a^2)^(3/2))))*(a^2
)^(1/2))/(2*(a*b^3 + b^4)^(1/2))
```

3.25 $\int \frac{\cosh^2(x)}{a+b \cosh^2(x)} dx$

Optimal result	189
Rubi [A] (verified)	189
Mathematica [A] (verified)	190
Maple [B] (verified)	190
Fricas [A] (verification not implemented)	191
Sympy [F(-1)]	191
Maxima [B] (verification not implemented)	192
Giac [A] (verification not implemented)	192
Mupad [B] (verification not implemented)	192

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \frac{\cosh^2(x)}{a+b \cosh^2(x)} dx = \frac{x}{b} - \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b\sqrt{a+b}}$$

[Out] x/b-arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))*a^(1/2)/b/(a+b)^(1/2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3250, 3260, 214}

$$\int \frac{\cosh^2(x)}{a+b \cosh^2(x)} dx = \frac{x}{b} - \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b\sqrt{a+b}}$$

[In] Int[Cosh[x]^2/(a + b*Cosh[x]^2),x]

[Out] x/b - (Sqrt[a]*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(b*Sqrt[a + b])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 3250

Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^2/((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2), x_Symbol] := Simp[B*(x/b), x] + Dist[(A*b - a*B)/b, Int[1/(a +

`b*Sin[e + f*x]^2), x], x] /; FreeQ[{a, b, e, f, A, B}, x]`

Rule 3260

`Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]`

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{x}{b} - \frac{a \int \frac{1}{a+b \cosh^2(x)} dx}{b} \\ &= \frac{x}{b} - \frac{a \text{Subst}\left(\int \frac{1}{a-(a+b)x^2} dx, x, \coth(x)\right)}{b} \\ &= \frac{x}{b} - \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{b\sqrt{a+b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.92

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx = \frac{x - \frac{\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a+b}}}{b}$$

[In] `Integrate[Cosh[x]^2/(a + b*Cosh[x]^2),x]`

[Out] `(x - (Sqrt[a]*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/Sqrt[a + b])/b`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 91 vs. 2(31) = 62.

Time = 0.11 (sec) , antiderivative size = 92, normalized size of antiderivative = 2.36

method	result
risch	$\frac{x}{b} + \frac{\sqrt{a(a+b)} \ln\left(e^{2x + \frac{2\sqrt{a(a+b)}+2a+b}{b}}\right)}{2(a+b)b} - \frac{\sqrt{a(a+b)} \ln\left(e^{2x - \frac{2\sqrt{a(a+b)}-2a-b}{b}}\right)}{2(a+b)b}$
default	$\frac{2a \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 - 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+\sqrt{a+b}}\right)}{4\sqrt{a}\sqrt{a+b}} \right)}{b} + \frac{\ln\left(\tanh\left(\frac{x}{2}\right)+1\right)}{b} - \frac{\ln\left(\tanh\left(\frac{x}{2}\right)-1\right)}{b}$

[In] `int(cosh(x)^2/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $x/b + 1/2 * (a * (a+b))^{(1/2)} / (a+b) / b * \ln(\exp(2*x) + (2 * (a * (a+b))^{(1/2)} + 2 * a + b) / b) - 1 / 2 * (a * (a+b))^{(1/2)} / (a+b) / b * \ln(\exp(2*x) - (2 * (a * (a+b))^{(1/2)} - 2 * a - b) / b)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 317, normalized size of antiderivative = 8.13

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx$$

$$= \left[\frac{\sqrt{\frac{a}{a+b}} \log \left(\frac{b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2ab + b^2) \cosh(x)^2 + 2(3b^2 \cosh(x)^2 + 2ab + b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4a^2}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 2ab + b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4a^2} \right)}{b} - \frac{\sqrt{-\frac{a}{a+b}} \arctan \left(\frac{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + 2a + b) \sqrt{-\frac{a}{a+b}}}{2a} \right) - x}{b} \right]$$

[In] integrate(cosh(x)^2/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] $[1/2 * (\sqrt{a/(a+b)}) * \log((b^2 * \cosh(x)^4 + 4 * b^2 * \cosh(x) * \sinh(x)^3 + b^2 * \sinh(x)^4 + 2 * (2 * a * b + b^2) * \cosh(x)^2 + 2 * (3 * b^2 * \cosh(x)^2 + 2 * a * b + b^2) * \sinh(x)^2 + 8 * a^2 + 8 * a * b + b^2 + 4 * (b^2 * \cosh(x)^3 + (2 * a * b + b^2) * \cosh(x)) * \sinh(x) + 4 * ((a * b + b^2) * \cosh(x)^2 + 2 * (a * b + b^2) * \cosh(x) * \sinh(x) + (a * b + b^2) * \sinh(x)^2 + 2 * a^2 + 3 * a * b + b^2) * \sqrt{a/(a+b)}) / (b * \cosh(x)^4 + 4 * b * \cosh(x) * \sinh(x)^3 + b * \sinh(x)^4 + 2 * (2 * a + b) * \cosh(x)^2 + 2 * (3 * b * \cosh(x)^2 + 2 * a + b) * \sinh(x)^2 + 4 * (b * \cosh(x)^3 + (2 * a + b) * \cosh(x)) * \sinh(x) + b) + 2 * x) / b, -(\sqrt{-a/(a+b)}) * \arctan(1/2 * (b * \cosh(x)^2 + 2 * b * \cosh(x) * \sinh(x) + b * \sinh(x)^2 + 2 * a + b) * \sqrt{-a/(a+b)}) / a - x) / b]$

Sympy [F(-1)]

Timed out.

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx = \text{Timed out}$$

[In] integrate(cosh(x)**2/(a+b*cosh(x)**2),x)

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 120 vs. 2(31) = 62.

Time = 0.28 (sec) , antiderivative size = 120, normalized size of antiderivative = 3.08

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx = -\frac{(2a + b) \log\left(\frac{be^{(2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)ab}} - \frac{\log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)a}} + \frac{x}{b}$$

[In] integrate(cosh(x)^2/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] -1/4*(2*a + b)*log((b*e^(2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*b) - 1/4*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/sqrt((a + b)*a) + x/b

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 50, normalized size of antiderivative = 1.28

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx = -\frac{a \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - ab}} + \frac{x}{b}$$

[In] integrate(cosh(x)^2/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] -a*arctan(1/2*(b*e^(2*x) + 2*a + b)/sqrt(-a^2 - a*b))/(sqrt(-a^2 - a*b)*b) + x/b

Mupad [B] (verification not implemented)

Time = 2.33 (sec) , antiderivative size = 376, normalized size of antiderivative = 9.64

$$\int \frac{\cosh^2(x)}{a + b \cosh^2(x)} dx = \frac{x}{b} + \frac{\sqrt{a} \operatorname{atan}\left(\frac{(b^5 \sqrt{-b^3 - ab^2} + ab^4 \sqrt{-b^3 - ab^2}) \left(e^{2x} \left(\frac{2(8a^{5/2} \sqrt{-b^3 - ab^2} + \sqrt{a} b^2 \sqrt{-b^3 - ab^2} + 8a^{3/2} b \sqrt{-b^3 - ab^2}) (8a^2 + 8ab + b^2)\right)}{b^8 (a+b)^2 \sqrt{-b^3 - ab^2}}\right) + \frac{4\sqrt{a}(4a+2)}{b^7 (a+b)}}{4a}\right)}{\sqrt{-b^3 - ab^2}}$$

[In] $\text{int}(\cosh(x)^2/(a + b*\cosh(x)^2),x)$

[Out] $x/b + (a^{1/2})*\text{atan}(((b^5*(-a*b^2 - b^3)^{1/2} + a*b^4*(-a*b^2 - b^3)^{1/2})*(\exp(2*x))*((2*(8*a^{5/2})*(-a*b^2 - b^3)^{1/2} + a^{1/2}*b^2*(-a*b^2 - b^3)^{1/2} + 8*a^{3/2}*b*(-a*b^2 - b^3)^{1/2})*(8*a*b + 8*a^2 + b^2))/(b^8*(a + b)^2*(-a*b^2 - b^3)^{1/2}) + (4*a^{1/2}*(4*a + 2*b)*(4*a*b^3 + 8*a^3*b + 12*a^2*b^2))/(b^7*(a + b)*(-b^2*(a + b))^{1/2}*(-a*b^2 - b^3)^{1/2})) + (2*(a^{1/2}*b^2*(-a*b^2 - b^3)^{1/2} + 2*a^{3/2}*b*(-a*b^2 - b^3)^{1/2})*(8*a*b + 8*a^2 + b^2))/(b^8*(a + b)^2*(-a*b^2 - b^3)^{1/2}) + (4*a^{1/2}*(2*a*b^3 + 2*a^2*b^2)*(4*a + 2*b))/(b^7*(a + b)*(-b^2*(a + b))^{1/2}*(-a*b^2 - b^3)^{1/2}))/((4*a)))/(-a*b^2 - b^3)^{1/2}$

3.26 $\int \frac{\cosh(x)}{a+b \cosh^2(x)} dx$

Optimal result	194
Rubi [A] (verified)	194
Mathematica [A] (verified)	195
Maple [B] (verified)	195
Fricas [B] (verification not implemented)	196
Sympy [B] (verification not implemented)	196
Maxima [F]	227
Giac [F]	227
Mupad [B] (verification not implemented)	227

Optimal result

Integrand size = 13, antiderivative size = 29

$$\int \frac{\cosh(x)}{a+b \cosh^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{\sqrt{b}\sqrt{a+b}}$$

[Out] $\arctan(\sinh(x)*b^{(1/2)/(a+b)^{(1/2)})/b^{(1/2)/(a+b)^{(1/2)}}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {3265, 211}

$$\int \frac{\cosh(x)}{a+b \cosh^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{\sqrt{b}\sqrt{a+b}}$$

[In] $\text{Int}[\text{Cosh}[x]/(a + b*\text{Cosh}[x]^2), x]$

[Out] $\text{ArcTan}[(\text{Sqrt}[b]*\text{Sinh}[x])/(\text{Sqrt}[a + b])]/(\text{Sqrt}[b]*\text{Sqrt}[a + b])$

Rule 211

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

Rule 3265

$\text{Int}[\sin[(e_ + (f_)*(x_)]^{(m_)}*((a_ + (b_)*\sin[(e_ + (f_)*(x_)]^2)^{(p_)}), x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\text{Cos}[e + f*x], x]\}, \text{Dist}[-\text{ff}/f, \text{Subst}[\text{Int}[(1 - \text{ff}^2*x^2)^{(m-1)/2}*(a + b - b*\text{ff}^2*x^2)^p, x], x, \text{Cos}[e +$

`f*x]/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]`

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right) \\ &= \frac{\arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{\sqrt{b}\sqrt{a+b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\cosh(x)}{a+b\cosh^2(x)} dx = \frac{\arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{\sqrt{b}\sqrt{a+b}}$$

[In] `Integrate[Cosh[x]/(a + b*Cosh[x]^2), x]`

[Out] `ArcTan[(Sqrt[b]*Sinh[x])/Sqrt[a + b]]/(Sqrt[b]*Sqrt[a + b])`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 65 vs. 2(21) = 42.

Time = 0.10 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.28

method	result	size
default	$\frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right)\sqrt{a+b+2\sqrt{a}}}{2\sqrt{b}}\right)}{\sqrt{a+b}\sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right)\sqrt{a+b-2\sqrt{a}}}{2\sqrt{b}}\right)}{\sqrt{a+b}\sqrt{b}}$	66
risch	$-\frac{\ln\left(e^{2x} - \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1\right)}{2\sqrt{-ab-b^2}} + \frac{\ln\left(e^{2x} + \frac{2(a+b)e^x}{\sqrt{-ab-b^2}} - 1\right)}{2\sqrt{-ab-b^2}}$	82

[In] `int(cosh(x)/(a+b*cosh(x)^2), x, method=_RETURNVERBOSE)`

[Out] `1/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)+2*a^(1/2))/b^(1/2))+1/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)-2*a^(1/2))/b^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 114 vs. $2(21) = 42$.

Time = 0.26 (sec) , antiderivative size = 337, normalized size of antiderivative = 11.62

$$\int \frac{\cosh(x)}{a + b \cosh^2(x)} dx$$

$$= \left[\frac{\sqrt{-ab - b^2} \log \left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 - 2(2a+3b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a - 3b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a+3b) \cosh(x) \sinh(x) + b \sinh(x)^2) \sinh(x) + b}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a - 3b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a+3b) \cosh(x) \sinh(x) + b \sinh(x)^2) \sinh(x) + b} \right)}{2(ab + b^2)} \right]$$

[In] integrate(cosh(x)/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] $[-1/2\sqrt{-a*b - b^2}*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2*a + 3*b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a - 3*b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 - (2*a + 3*b)*\cosh(x))*\sinh(x) - 4*(\cosh(x)^3 + 3*\cosh(x)*\sinh(x)^2 + \sinh(x)^3 + (3*\cosh(x)^2 - 1)*\sinh(x) - \cosh(x))*\sqrt{-a*b - b^2} + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b))/(a*b + b^2), (\sqrt{a*b + b^2}*\arctan(1/2*(b*\cosh(x)^3 + 3*b*\cosh(x)*\sinh(x)^2 + b*\sinh(x)^3 + (4*a + 3*b)*\cosh(x) + (3*b*\cosh(x)^2 + 4*a + 3*b)*\sinh(x))/\sqrt{a*b + b^2})) + \sqrt{a*b + b^2}*\arctan(1/2*\sqrt{a*b + b^2}*(\cosh(x) + \sinh(x))/(a + b)))/(a*b + b^2)]$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 55498 vs. $2(27) = 54$.

Time = 106.06 (sec) , antiderivative size = 55498, normalized size of antiderivative = 1913.72

$$\int \frac{\cosh(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(cosh(x)/(a+b*cosh(x)**2),x)

[Out] Piecewise((zoo*atan(tanh(x/2)), Eq(a, 0) & Eq(b, 0)), (tanh(x/2)/(2*b) - 1/(2*b*tanh(x/2)), Eq(a, -b)), (sinh(x)/a, Eq(b, 0)), (13*a**6*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**7*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a**6*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 24*a**6*b*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**5*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))

$$\begin{aligned}
& b/(a + b) - 2\sqrt{-a*b}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}} \\
& / (a + b)) + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/ \\
& (a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + 858*a**5*b**3 \\
& *\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + \\
& b) + 2\sqrt{-a*b}}/(a + b)) - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(\\
& a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a \\
& + b)) - 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))*s \\
& qrt(a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + 1144*a**4*b**3*\sqrt{-a* \\
& b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a \\
& + b) + 2\sqrt{-a*b}}/(a + b)) - 858*a**3*b**5*\sqrt{a/(a + b) - b/(a + b) - \\
& 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + \\
& 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a \\
& + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a} \\
& / (a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2 \\
& *\sqrt{-a*b}}/(a + b)) - 130*a*b**7*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}} \\
& / (a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + 416*a*b**6*s \\
& qrt(-a*b)*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) \\
& - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + 2*b**8*\sqrt{a/(a + b) - b/(a + b) - \\
& 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - \\
& 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{ \\
& a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b))) + 11*a**6*b*\sqrt{a/(a + b) - \\
& b/(a + b) + 2\sqrt{-a*b}}/(a + b))*\log(\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{- \\
& a*b}}/(a + b)) + \tanh(x/2))/(2*a**7*b*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{- \\
& a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 130*a**6 \\
& *b**2*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b \\
& / (a + b) + 2\sqrt{-a*b}}/(a + b)) + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/ \\
& (a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(\\
& a + b)) + 858*a**5*b**3*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))* \\
& \sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 416*a**5*b**2*\sqrt{-a* \\
& b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a \\
& + b) + 2\sqrt{-a*b}}/(a + b)) - 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - \\
& 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + \\
& 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b) \\
&)\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 858*a**3*b**5*\sqrt{a} \\
& / (a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2 \\
& *\sqrt{-a*b}}/(a + b)) + 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a \\
& *b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 1144*a**2 \\
& *b**5*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/ \\
& (a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) - 130*a*b**7*\sqrt{a/(a + b) - b \\
& / (a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/ \\
& (a + b)) + 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b}}/ \\
& (a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b)) + 2*b**8*\sqrt{a} \\
& / (a + b) - b/(a + b) - 2\sqrt{-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2 \\
& *\sqrt{-a*b}}/(a + b)) - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*sqr \\
& rt(-a*b}}/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b}}/(a + b))) - a**
\end{aligned}$$

$$\begin{aligned}
& (a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)) + 2*b**8*\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b))) + a**6*\sqrt{-a*b}*\sqrt{ \\
& t(a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b))*\log(-\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b)) + \tanh(x/2))/(2*a**7*b*\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
&) - 130*a**6*b**2*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{ \\
& (a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)) + 24*a**6*b*\sqrt{-a*b}*\sqrt{ \\
& a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + \\
& 2*\sqrt{-a*b}/(a + b)} + 858*a**5*b**3*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 416*a**5 \\
& *b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/ \\
& (a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 858*a**4*b**4*\sqrt{a/(a + b) \\
& - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a* \\
& b)/(a + b)} + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{ \\
& (-a*b)/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 858*a* \\
& **3*b**5*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - \\
& b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + \\
& b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
&) - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a*b**7*\sqrt{ \\
& (a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + \\
& 2*\sqrt{-a*b}/(a + b)} + 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - \\
& 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + \\
& 2*b**8*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - \\
& b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/ \\
& (a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(\\
& a + b))) - a**6*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
&)*\log(\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} + \tanh(x/2))/(2*a \\
& **7*b*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b \\
& /(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a**6*b**2*\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))* \\
& \sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**5*b**3*\sqrt{a/(\\
& a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2* \\
& \sqrt{-a*b}/(a + b)} - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - \\
& 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - \\
& 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + \\
& b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a \\
& /(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2 \\
& *\sqrt{-a*b}/(a + b)} - 858*a**3*b**5*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a \\
& *b)/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**2* \\
& b**6*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/ \\
& (a + b) + 2*\sqrt{-a*b}/(a + b)} - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a + b)
\end{aligned}$$

$$\begin{aligned}
& -b/(a+b) - 2\sqrt{-ab}/(a+b))\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} \\
& b)/(a+b)) - 130ab^{*7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 416ab^{*6}\sqrt{-ab} \\
&)\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}}\sqrt{a/(a+b) - b/(a+b)} \\
& + 2\sqrt{-ab}/(a+b) + 2b^{*8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 24b^{*7}\sqrt{-ab} \\
&)\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}}\sqrt{a/(a+b) - b/(a+b)} \\
& + 2\sqrt{-ab}/(a+b)) - 286a^{*5}b^{*2}\sqrt{a/(a+b) - b/(a+b)} \\
& - 2\sqrt{-ab}/(a+b))\log(-\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} \\
&)/(a+b) + \tanh(x/2))/(2a^{*7}b\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 130a^{*6}b^{*2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 24a^{*6}b\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b)} \\
& - 2\sqrt{-ab}/(a+b))\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} \\
&) + 858a^{*5}b^{*3}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} \\
& - 416a^{*5}b^{*2}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b)} + 2\sqrt{-ab}/(a+b) - 858a^{*4}b^{*4}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 1144a^{*4}b^{*3}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 858a^{*3}b^{*5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 858a^{*2}b^{*6}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 1144a^{*2}b^{*5}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 130ab^{*7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 416ab^{*6}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 2b^{*8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 24b^{*7}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 286a^{*5}b^{*2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\log(\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}}/(a+b) + \tanh(x/2))/(2a^{*7}b\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 130a^{*6}b^{*2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 24a^{*6}b\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 858a^{*5}b^{*3}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 416a^{*5}b^{*2}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} - 858a^{*4}b^{*4}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}} + 1144a^{*4}b^{*3}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}} \\
&)\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}}
\end{aligned}$$

$$\begin{aligned}
& - 858*a**3*b**5*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**2*b**6*sqrt(a/(a + b) \\
& - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a* \\
& b)/(a + b)) - 1144*a**2*b**5*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt \\
& (-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a* \\
& b**7*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/ \\
& (a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a*b**6*sqrt(-a*b)*sqrt(a/(a + b) - b/ \\
& (a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(\\
& a + b)) + 2*b**8*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 24*b**7*sqrt(-a*b)*sqrt(a/(a \\
& + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt \\
& (-a*b)/(a + b)) + 154*a**5*b**2*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b) \\
& /(a + b))*log(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/ \\
& 2))/(2*a**7*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a \\
& + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a**6*b**2*sqrt(a/(a + b) - b \\
& /(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(\\
& (a + b)) + 24*a**6*b*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(\\
& a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**5*b**3* \\
& sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b)) - 416*a**5*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a \\
& + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b)) - 858*a**4*b**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt \\
& (a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 1144*a**4*b**3*sqrt(-a*b) \\
&)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a \\
& + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**3*b**5*sqrt(a/(a + b) - b/(a + b) - 2 \\
& *sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8 \\
& 58*a**2*b**6*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + \\
& b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 1144*a**2*b**5*sqrt(-a*b)*sqrt(a/ \\
& (a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2* \\
& sqrt(-a*b)/(a + b)) - 130*a*b**7*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/ \\
& (a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a*b**6*sqrt \\
& (-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) \\
& - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*b**8*sqrt(a/(a + b) - b/(a + b) - 2 \\
& *sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 2 \\
& 4*b**7*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a \\
& /(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 154*a**5*b**2*sqrt(a/(a + b) \\
&) - b/(a + b) + 2*sqrt(-a*b)/(a + b))*log(sqrt(a/(a + b) - b/(a + b) - 2*sqrt \\
& (-a*b)/(a + b)) + tanh(x/2))/(2*a**7*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt \\
& (-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a \\
& **6*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) \\
& - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 24*a**6*b*sqrt(-a*b)*sqrt(a/(a + b) - \\
& b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b) \\
&)/(a + b)) + 858*a**5*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b \\
&))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 416*a**5*b**2*sqrt(\\
& -a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b
\end{aligned}$$

$$\begin{aligned}
& + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)) - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b)} \\
&) - b/(a + b) - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& - 1287*a**4*b**3*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\log(\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + \tanh(x/2)) \\
& /((2*a**7*b*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b)} \\
&) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)) - 130*a**6*b**2*\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**5*b**3*\sqrt{a/(a + b) - b/(a + b)} \\
& - 2*\sqrt{-a*b}/(a + b))*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
&) - 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b)} \\
& + 2*\sqrt{-a*b}/(a + b) + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b)} \\
& + 2*\sqrt{-a*b}/(a + b)) - 858*a**3*b**5*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 130*a*b**7*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 2*b**8*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} \\
&)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
&) - 297*a**4*b**3*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)}*\log(-\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)} + \tanh(x/2)) \\
& /((2*a**7*b*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a**6*b**2*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**5*b**3*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 858*a**3*b**5*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a*b**7*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} -
\end{aligned}$$

$$\begin{aligned}
& *b)/(a + b)) + 858*a**2*b**6*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + \\
& b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 1144*a**2*b**5*sq \\
& rt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) \\
& - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a*b**7*sqrt(a/(a + b) - b/(a + b) \\
& - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) \\
& + 416*a*b**6*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) \\
& *sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*b**8*sqrt(a/(a + b) \\
& - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a \\
& *b)/(a + b)) - 24*b**7*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b) \\
& /(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + 715*a**4*b* \\
& *2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(a \\
& /(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**7*b*sqrt(a/ \\
& (a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2* \\
& sqrt(-a*b)/(a + b)) - 130*a**6*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a* \\
& b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 24*a**6*b* \\
& sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) \\
&) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**5*b**3*sqrt(a/(a + b) - b/(a \\
& + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b)) - 416*a**5*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/ \\
& (a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 858*a**4*b**4 \\
& *sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b)) + 1144*a**4*b**3*sqrt(-a*b)*sqrt(a/(a + b) - b/ \\
& (a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(\\
& a + b)) - 858*a**3*b**5*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))* \\
& sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 858*a**2*b**6*sqrt(a/(\\
& a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*s \\
& qrt(-a*b)/(a + b)) - 1144*a**2*b**5*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - \\
& 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - \\
& 130*a*b**7*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + \\
& b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a*b**6*sqrt(-a*b)*sqrt(a/(a + \\
& b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(\\
& -a*b)/(a + b)) + 2*b**8*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))* \\
& sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 24*b**7*sqrt(-a*b)*sq \\
& rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) \\
& + 2*sqrt(-a*b)/(a + b))) + 275*a**4*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/ \\
& (a + b)) + tanh(x/2))/(2*a**7*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(\\
& a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 130*a**6*b**2* \\
& sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + \\
& b) + 2*sqrt(-a*b)/(a + b)) + 24*a**6*b*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b \\
&) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b) \\
&) + 858*a**5*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a \\
& /(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 416*a**5*b**2*sqrt(-a*b)*sq \\
& rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) \\
& + 2*sqrt(-a*b)/(a + b)) - 858*a**4*b**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt
\end{aligned}$$

$$\begin{aligned}
& (-a*b)/(a + b)*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 1144*a \\
& **4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{ \\
& (a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)) - 858*a**3*b**5*\sqrt{a/(a + \\
& b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{ \\
& -a*b}/(a + b)} + 858*a**2*b**6*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a \\
& + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 1144*a**2*b**5* \\
& \sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b \\
&) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a*b**7*\sqrt{a/(a + b) - b/(a + \\
& b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b \\
&)} + 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b \\
&)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 2*b**8*\sqrt{a/(a + \\
& b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{ \\
& -a*b}/(a + b)} - 24*b**7*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a* \\
& b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b))} - 275*a**4* \\
& b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)}*\log(\sqrt{ \\
& (a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)) + \tanh(x/2)})/(2*a**7*b*\sqrt{ \\
& a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + \\
& 2*\sqrt{-a*b}/(a + b)} - 130*a**6*b**2*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{- \\
& a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 24*a**6* \\
& b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + \\
& b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**5*b**3*\sqrt{a/(a + b) - b/ \\
& (a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(\\
& a + b)} - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b} \\
&)/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 858*a**4*b* \\
& **4*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a \\
& + b) + 2*\sqrt{-a*b}/(a + b)} + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(a + b) - \\
& b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b} \\
& / (a + b)} - 858*a**3*b**5*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b \\
&)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*a**2*b**6*\sqrt{a \\
& / (a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2 \\
& *\sqrt{-a*b}/(a + b)} - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) \\
& - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& - 130*a*b**7*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a \\
& + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a \\
& + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*sqr \\
& t(-a*b)/(a + b)} + 2*b**8*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b \\
&)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 24*b**7*\sqrt{-a*b}*s \\
& \sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b \\
&) + 2*\sqrt{-a*b}/(a + b))} - 1716*a**3*b**4*\sqrt{a/(a + b) - b/(a + b) - 2* \\
& \sqrt{-a*b}/(a + b)}*\log(-\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} \\
& + \tanh(x/2)})/(2*a**7*b*\sqrt{a/(a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}* \\
& \sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} - 130*a**6*b**2*\sqrt{a/(\\
& a + b) - b/(a + b) - 2*\sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*s \\
& \sqrt{-a*b}/(a + b)} + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a + b) - b/(a + b) - 2*s \\
& \sqrt{-a*b}/(a + b)}*\sqrt{a/(a + b) - b/(a + b) + 2*\sqrt{-a*b}/(a + b)} + 858*
\end{aligned}$$

$$\begin{aligned}
& *b)/(a + b))\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 1144*a**2 \\
& *b**5\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/ \\
& (a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 130*a*b**7\sqrt{a/(a + b) - b \\
& / (a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/ \\
& (a + b)} + 416*a*b**6\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/ \\
& (a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 2*b**8\sqrt{a \\
& / (a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2 \\
& * \sqrt{-a*b)/(a + b)} - 24*b**7\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2* \sqrt{ \\
& rt(-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)}} + 715 \\
& *a**2*b**5\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}*\log(-\sqrt{a/(\\
& a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + \tanh(x/2))/(2*a**7*b*\sqrt{a/(a \\
& + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2* \sqrt{ \\
& rt(-a*b)/(a + b)} - 130*a**6*b**2\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b) \\
& / (a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 24*a**6*b* \sqrt{ \\
& rt(-a*b)}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) \\
& - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 858*a**5*b**3\sqrt{a/(a + b) - b/(a + \\
& b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + \\
& b)} - 416*a**5*b**2\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a \\
& + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 858*a**4*b**4* \sqrt{ \\
& rt(a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) \\
&) + 2\sqrt{-a*b)/(a + b)} + 1144*a**4*b**3\sqrt{-a*b}\sqrt{a/(a + b) - b/(a \\
& + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a \\
& + b)} - 858*a**3*b**5\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}* \sqrt{ \\
& rt(a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 858*a**2*b**6\sqrt{a/(a \\
& + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2* \sqrt{ \\
& rt(-a*b)/(a + b)} - 1144*a**2*b**5\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2 \\
& * \sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 1 \\
& 30*a*b**7\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) \\
& - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 416*a*b**6\sqrt{-a*b}\sqrt{a/(a + b) \\
& - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a \\
& *b)/(a + b)} + 2*b**8\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}* \sqrt{ \\
& rt(a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 24*b**7\sqrt{-a*b}\sqrt{ \\
& a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + \\
& 2\sqrt{-a*b)/(a + b)}} - 715*a**2*b**5\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{ \\
& -a*b)/(a + b)}*\log(\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + \tan \\
& h(x/2))/(2*a**7*b*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a \\
& / (a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} - 130*a**6*b**2\sqrt{a/(a + b) \\
& - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a \\
& *b)/(a + b)} + 24*a**6*b*\sqrt{-a*b}\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a* \\
& b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 858*a**5*b \\
& **3\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(\\
& a + b) + 2\sqrt{-a*b)/(a + b)} - 416*a**5*b**2\sqrt{-a*b}\sqrt{a/(a + b) - \\
& b/(a + b) - 2\sqrt{-a*b)/(a + b)}\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b) \\
& / (a + b)} - 858*a**4*b**4*\sqrt{a/(a + b) - b/(a + b) - 2\sqrt{-a*b)/(a + b) \\
&)\sqrt{a/(a + b) - b/(a + b) + 2\sqrt{-a*b)/(a + b)} + 1144*a**4*b**3\sqrt{
\end{aligned}$$

$$\begin{aligned}
& -a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/ \\
& / (a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 858*a**3*b**5*\text{sqrt}(a/(a + b) - b/(a + b) \\
& - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) \\
& + 858*a**2*b**6*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/ \\
& (a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 1144*a**2*b**5*\text{sqrt}(-a*b)*\text{sqr} \\
& t(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) \\
& + 2*\text{sqrt}(-a*b)/(a + b)) - 130*a*b**7*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a \\
& *b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 416*a*b** \\
& 6*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + \\
& b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 2*b**8*\text{sqrt}(a/(a + b) - b/(a + b) \\
& - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) \\
& - 24*b**7*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sq} \\
& rt(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b))) + 275*a**2*b**5*\text{sqrt}(a/(a \\
& + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b))*\text{log}(-\text{sqrt}(a/(a + b) - b/(a + b) - \\
& 2*\text{sqrt}(-a*b)/(a + b)) + \text{tanh}(x/2))/(2*a**7*b*\text{sqrt}(a/(a + b) - b/(a + b) - \\
& 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - \\
& 130*a**6*b**2*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a \\
& + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 24*a**6*b*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + \\
& b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt} \\
& (-a*b)/(a + b)) + 858*a**5*b**3*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(\\
& a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 416*a**5*b**2* \\
& \text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b \\
&) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 858*a**4*b**4*\text{sqrt}(a/(a + b) - b/(a \\
& + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a \\
& + b)) + 1144*a**4*b**3*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b) \\
& / (a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 858*a**3*b** \\
& 5*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a \\
& + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 858*a**2*b**6*\text{sqrt}(a/(a + b) - b/(a + b) - 2 \\
& *\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 1 \\
& 144*a**2*b**5*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b)) \\
& *\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 130*a*b**7*\text{sqrt}(a/(a \\
& + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqr} \\
& t(-a*b)/(a + b)) + 416*a*b**6*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqr} \\
& t(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 2*b** \\
& 8*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a \\
& + b) + 2*\text{sqrt}(-a*b)/(a + b)) - 24*b**7*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b \\
&) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b \\
&)) - 275*a**2*b**5*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b))*\text{log}(s \\
& \text{qrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b)) + \text{tanh}(x/2))/(2*a**7*b*\text{sq} \\
& rt(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) \\
& + 2*\text{sqrt}(-a*b)/(a + b)) - 130*a**6*b**2*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqr} \\
& t(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 24*a* \\
& *6*b*\text{sqrt}(-a*b)*\text{sqrt}(a/(a + b) - b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(\\
& a + b) - b/(a + b) + 2*\text{sqrt}(-a*b)/(a + b)) + 858*a**5*b**3*\text{sqrt}(a/(a + b) - \\
& b/(a + b) - 2*\text{sqrt}(-a*b)/(a + b))*\text{sqrt}(a/(a + b) - b/(a + b) + 2*\text{sqrt}(-a*b
\end{aligned}$$

$$\begin{aligned}
& \sqrt{-ab}/(a+b) \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2 \\
& 4a^6b \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^5b^3 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 416a^5b^2 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^4b^4 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 1144a^4b^3 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^3b^5 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^2b^6 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 1144a^2b^5 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 130ab^7 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 416ab^6 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2b^8 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 24b^7 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 297a^2b^4 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \log(-\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} + \tanh(x/2)) / (2a^7b \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}) - 130a^6b^2 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 24a^6b \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^5b^3 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 416a^5b^2 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^4b^4 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 1144a^4b^3 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^3b^5 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^2b^6 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 1144a^2b^5 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 130ab^7 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 416ab^6 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2b^8 \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 24b^7 \sqrt{-ab} \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2\sqrt{-ab}/(a+b) \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}
\end{aligned}$$

$$\begin{aligned}
& 297*a**2*b**4*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} \\
& * \log(\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} + \tanh(x/2))/(2*a** \\
& 7*b*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(\\
& a+b) + 2*\sqrt{-a*b}/(a+b)} - 130*a**6*b**2*\sqrt{a/(a+b) - b/(a+b) - \\
& 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + \\
& 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{ \\
& a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 858*a**5*b**3*\sqrt{a/(a \\
& + b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) - 2* \\
& \sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 85 \\
& 8*a**4*b**4*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) \\
& - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 1144*a**4*b**3*\sqrt{-a*b}*\sqrt{a/(\\
& a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} - 858*a**3*b**5*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
&)/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 858*a**2*b* \\
& *6*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a \\
& + b) + 2*\sqrt{-a*b}/(a+b)} - 1144*a**2*b**5*\sqrt{-a*b}*\sqrt{a/(a+b) - \\
& b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b} \\
& / (a+b)} - 130*a*b**7*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{ \\
& a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 416*a*b**6*\sqrt{-a*b}*\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b} \\
&)/(a+b)} + 2*\sqrt{-a*b}/(a+b)} + 2*b**8*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
&)/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 24*b**7*\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b} \\
& / (a+b) + 2*\sqrt{-a*b}/(a+b)} - 78*a*b**6*\sqrt{a/(a+b) - b/(a+b) \\
& - 2*\sqrt{-a*b}/(a+b)}*\log(-\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} \\
& + \tanh(x/2))/(2*a**7*b*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} \\
&)*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 130*a**6*b**2*\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2* \\
& \sqrt{-a*b}/(a+b)} + 24*a**6*b*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
& / (a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 858*a**5*b**3*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
&)/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 416*a**5*b**2*\sqrt{-a*b}*\sqrt{a/(\\
& a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2* \\
& \sqrt{-a*b}/(a+b)} - 858*a**4*b**4*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
& / (a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 1144*a**4* \\
& b**3*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(\\
& a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 858*a**3*b**5*\sqrt{a/(a+b) - \\
& b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b} \\
&)/(a+b)} + 858*a**2*b**6*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b) \\
&)*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 1144*a**2*b**5*\sqrt{ \\
& -a*b}*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - \\
& b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 130*a*b**7*\sqrt{a/(a+b) - b/(a+b) - 2* \\
& \sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + \\
& 416*a*b**6*\sqrt{-a*b}*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*s
\end{aligned}$$

$$\begin{aligned}
& \text{qrt}(a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)) + 2b^{**8}\sqrt{a/(a+b) - } \\
& \quad b/(a+b) - 2\sqrt{-ab}/(a+b))\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab} } \\
& \quad)/(a+b)) - 24b^{**7}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(} \\
& \quad a+b))\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b))} + 78ab^{**6}\sqrt{ } \\
& \quad t(a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b))*\log(\sqrt{a/(a+b) - b/(a+ } \\
& \quad b) + 2\sqrt{-ab}/(a+b) + \tanh(x/2))/(2a^{**7}b\sqrt{a/(a+b) - b/(a+ } \\
& \quad b) - 2\sqrt{-ab}/(a+b))\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b } \\
& \quad)) - 130a^{**6}b^{**2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{ } \\
& \quad a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)) + 24a^{**6}b\sqrt{-ab}\sqrt{a } \\
& \quad /(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b))\sqrt{a/(a+b) - b/(a+b) + 2 } \\
& \quad *\sqrt{-ab}/(a+b)} + 858a^{**5}b^{**3}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a } \\
& \quad *b)/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 416a^{**5} } \\
& \quad b^{**2}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(} \\
& \quad a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)) - 858a^{**4}b^{**4}\sqrt{a/(a+b) - } \\
& \quad b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab} } \\
& \quad)/(a+b)} + 1144a^{**4}b^{**3}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{- } \\
& \quad -ab)/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^{** } \\
& \quad 3b^{**5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - } \\
& \quad b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^{**2}b^{**6}\sqrt{a/(a+b) - b/(a+b } \\
& \quad) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b } \\
& \quad) - 1144a^{**2}b^{**5}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a } \\
& \quad +b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 130ab^{**7}\sqrt{ } \\
& \quad a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + } \\
& \quad 2\sqrt{-ab}/(a+b)} + 416ab^{**6}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - } \\
& \quad 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + } \\
& \quad 2b^{**8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - } \\
& \quad b/(a+b) + 2\sqrt{-ab}/(a+b)} - 24b^{**7}\sqrt{-ab}\sqrt{a/(a+b) - b/(} \\
& \quad a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a } \\
& \quad +b)) - 54ab^{**6}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}\log(} \\
& \quad -\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} + \tanh(x/2))/(2a^{**7}b } \\
& \quad \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) } \\
& \quad + 2\sqrt{-ab}/(a+b)} - 130a^{**6}b^{**2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{ } \\
& \quad \text{qrt}(-ab)/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 24 } \\
& \quad a^{**6}b\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a } \\
& \quad /(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^{**5}b^{**3}\sqrt{a/(a+b) } \\
& \quad - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a } \\
& \quad *b)/(a+b)} - 416a^{**5}b^{**2}\sqrt{-ab}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{ } \\
& \quad (-ab)/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^{** } \\
& \quad *4b^{**4}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - } \\
& \quad b/(a+b) + 2\sqrt{-ab}/(a+b)} + 1144a^{**4}b^{**3}\sqrt{-ab}\sqrt{a/(a+ } \\
& \quad b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{- } \\
& \quad -ab)/(a+b)} - 858a^{**3}b^{**5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a } \\
& \quad +b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^{**2}b^{**6} } \\
& \quad \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}\sqrt{a/(a+b) - b/(a+b } \\
& \quad) + 2\sqrt{-ab}/(a+b)} - 1144a^{**2}b^{**5}\sqrt{-ab}\sqrt{a/(a+b) - b/(a }
\end{aligned}$$

$$\begin{aligned}
& 858a^{*3}b^{*5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 858a^{*2}b^{*6}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} \\
& / (a+b) - 1144a^{*2}b^{*5}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 130a^{*b} \\
& *7\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 416a^{*b}b^{*6}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} \\
& + 2b^{*8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 24b^{*7}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} \\
& - 286a^{*b}b^{*5}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\log(\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)}) + \tanh(x/2)/(2a^{*7}b^{*}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 130a^{*6}b^{*2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 24a^{*6}b^{*}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 858a^{*5}b^{*3}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 416a^{*5}b^{*2}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 858a^{*4}b^{*4}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 1144a^{*4}b^{*3}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 858a^{*3}b^{*5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 858a^{*2}b^{*6}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 1144a^{*2}b^{*5}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 130a^{*b}b^{*7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 416a^{*b}b^{*6}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 2b^{*8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 24b^{*7}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 154a^{*b}b^{*5}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)}\log(-\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)} + \tanh(x/2))/(2a^{*7}b^{*}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 130a^{*6}b^{*2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 24a^{*6}b^{*}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 858a^{*5}b^{*3}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} - 416a^{*5}b^{*2}\sqrt{-a*b}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)}
\end{aligned}$$

$$\begin{aligned} & (a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)) - 858a^{**4}b^{**4}\sqrt{a/(a+b)} - b/(a+b) - 2\sqrt{-ab}/(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 1144a^{**4}b^{**3}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858 \\ & *a^{**3}b^{**5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 858a^{**2}b^{**6}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & - 1144a^{**2}b^{**5}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & /(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 130a^{**6}b^{**7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 416a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2b^{**8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & - 24b^{**7}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & - 154a^{**5}b^{**5}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}*\log(\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}) + \tanh(x/2)) / (2a^{**7}b^{**7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}) \\ & - 130a^{**6}b^{**2}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 24a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & /(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^{**5}b^{**3} \\ & *\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 416a^{**5}b^{**2}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^{**4}b^{**4}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 1144a^{**4}b^{**3}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 858a^{**3}b^{**5}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 858a^{**2}b^{**6}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 1144a^{**2}b^{**5}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 130a^{**6}b^{**7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 416a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 2b^{**8}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} - 24b^{**7}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + b^{**7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\log(-\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}) / (2a^{**7}b^{**7}\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)}) - 130a^{**6}b^{**2} \\ & \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} + 24a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 24a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 24a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \\ & + 24a^{**6}b^{**6}\sqrt{-ab}*\sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-ab}/(a+b)} \\ & *\sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-ab}/(a+b)} \end{aligned}$$

$$\begin{aligned}
& \sqrt{a/(a+b) - b/(a+b) + 2\sqrt{-a*b}/(a+b)} + 416*a*b**6*\sqrt{-a*b}* \\
& \sqrt{a/(a+b) - b/(a+b) - 2\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b)} \\
& + 2*\sqrt{-a*b}/(a+b)) + 2*b**8*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}} \\
& /(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 24*b**7*\sqrt{ \\
& t(-a*b)*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - \\
& b/(a+b) + 2*\sqrt{-a*b}/(a+b))} - 11*b**6*\sqrt{-a*b}* \sqrt{a/(a+b) - b \\
& /(a+b) + 2*\sqrt{-a*b}/(a+b)}*\log(-\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a}} \\
& *b)/(a+b)) + \tanh(x/2)/(2*a**7*b*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a}} \\
& *b)/(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 130*a**6* \\
& b**2*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/ \\
& (a+b) + 2*\sqrt{-a*b}/(a+b)} + 24*a**6*b*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(\\
& a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a \\
& +b))} + 858*a**5*b**3*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*s \\
& \sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 416*a**5*b**2*\sqrt{-a*b} \\
&)*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a \\
& +b) + 2*\sqrt{-a*b}/(a+b)} - 858*a**4*b**4*\sqrt{a/(a+b) - b/(a+b) - 2} \\
& *\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 1 \\
& 144*a**4*b**3*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} \\
& *\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 858*a**3*b**5*\sqrt{a/ \\
& (a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2* \\
& \sqrt{-a*b}/(a+b)} + 858*a**2*b**6*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a* \\
& b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 1144*a**2* \\
& b**5*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(\\
& a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 130*a*b**7*\sqrt{a/(a+b) - b/ \\
& (a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(\\
& a+b)} + 416*a*b**6*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(\\
& a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 2*b**8*\sqrt{a/ \\
& (a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2* \\
& \sqrt{-a*b}/(a+b)} - 24*b**7*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{ \\
& t(-a*b)/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b))} + 11*b \\
& **6*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)}*\log(\sqrt{ \\
& a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} + \tanh(x/2)/(2*a**7*b*\sqrt{a \\
& /(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2} \\
& *\sqrt{-a*b}/(a+b)} - 130*a**6*b**2*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a}} \\
& *b)/(a+b))*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 24*a**6*b \\
& *\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+ \\
& b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 858*a**5*b**3*\sqrt{a/(a+b) - b/(\\
& a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a \\
& +b)} - 416*a**5*b**2*\sqrt{-a*b}* \sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b} \\
& /(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} - 858*a**4*b** \\
& 4*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a \\
& +b) + 2*\sqrt{-a*b}/(a+b)} + 1144*a**4*b**3*\sqrt{-a*b}* \sqrt{a/(a+b) - b \\
& /(a+b) - 2*\sqrt{-a*b}/(a+b)}*\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/ \\
& (a+b)} - 858*a**3*b**5*\sqrt{a/(a+b) - b/(a+b) - 2*\sqrt{-a*b}/(a+b)} \\
& *\sqrt{a/(a+b) - b/(a+b) + 2*\sqrt{-a*b}/(a+b)} + 858*a**2*b**6*\sqrt{a/
\end{aligned}$$

```
(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*
sqrt(-a*b)/(a + b)) - 1144*a**2*b**5*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b)
- 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))
- 130*a*b**7*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a +
b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 416*a*b**6*sqrt(-a*b)*sqrt(a/(a +
b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt
(-a*b)/(a + b)) + 2*b**8*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))
*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 24*b**7*sqrt(-a*b)*sq
rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b)
+ 2*sqrt(-a*b)/(a + b))), True))
```

Maxima [F]

$$\int \frac{\cosh(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(cosh(x)/(a+b*cosh(x)^2),x, algorithm="maxima")
```

```
[Out] integrate(cosh(x)/(b*cosh(x)^2 + a), x)
```

Giac [F]

$$\int \frac{\cosh(x)}{a + b \cosh^2(x)} dx = \int \frac{\cosh(x)}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(cosh(x)/(a+b*cosh(x)^2),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [B] (verification not implemented)

Time = 2.07 (sec) , antiderivative size = 87, normalized size of antiderivative = 3.00

$$\int \frac{\cosh(x)}{a + b \cosh^2(x)} dx = -\frac{\ln\left(-\frac{4(a - ae^{2x})}{b^2(a+b)} - \frac{8ae^x}{(-b)^{5/2}\sqrt{a+b}}\right) - \ln\left(\frac{8ae^x}{(-b)^{5/2}\sqrt{a+b}} - \frac{4(a - ae^{2x})}{b^2(a+b)}\right)}{2\sqrt{-b}\sqrt{a+b}}$$

```
[In] int(cosh(x)/(a + b*cosh(x)^2),x)
```

```
[Out] -(log(- (4*(a - a*exp(2*x)))/(b^2*(a + b)) - (8*a*exp(x))/((-b)^(5/2)*(a +
b)^(1/2)))) - log((8*a*exp(x))/((-b)^(5/2)*(a + b)^(1/2)) - (4*(a - a*exp(2*
x)))/(b^2*(a + b))))/(2*(-b)^(1/2)*(a + b)^(1/2))
```

3.27 $\int \frac{1}{a+b \cosh^2(x)} dx$

Optimal result	228
Rubi [A] (verified)	228
Mathematica [A] (verified)	229
Maple [B] (verified)	229
Fricas [B] (verification not implemented)	230
Sympy [B] (verification not implemented)	230
Maxima [B] (verification not implemented)	236
Giac [A] (verification not implemented)	237
Mupad [B] (verification not implemented)	237

Optimal result

Integrand size = 10, antiderivative size = 29

$$\int \frac{1}{a+b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[Out] $\operatorname{arctanh}(a^{(1/2)}*\tanh(x)/(a+b)^{(1/2)})/a^{(1/2)}/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3260, 214}

$$\int \frac{1}{a+b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[In] $\operatorname{Int}[(a + b*\operatorname{Cosh}[x]^2)^{-1}, x]$

[Out] $\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tanh}[x])/\operatorname{Sqrt}[a + b]]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[a + b])$

Rule 214

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b]$

Rule 3260

$\operatorname{Int}[(a_ + (b_)*\sin[(e_ + (f_)*(x_)]^2)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{\{ff = \operatorname{FreeFactors}[\operatorname{Tan}[e + f*x], x]\}, \operatorname{Dist}[ff/f, \operatorname{Subst}[\operatorname{Int}[1/(a + (a + b)*ff^2*x^2$

), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{a - (a + b)x^2} dx, x, \coth(x)\right) \\ &= \frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{\sqrt{a}\sqrt{a+b}}$$

[In] Integrate[(a + b*Cosh[x]^2)^(-1),x]

[Out] ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]]/(Sqrt[a]*Sqrt[a + b])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(21) = 42.

Time = 0.00 (sec) , antiderivative size = 78, normalized size of antiderivative = 2.69

method	result	size
default	$\frac{\ln\left(\sqrt{a+b}\tanh\left(\frac{x}{2}\right)^2 + 2\tanh\left(\frac{x}{2}\right)\sqrt{a+b}\right)}{2\sqrt{a}\sqrt{a+b}} - \frac{\ln\left(\sqrt{a+b}\tanh\left(\frac{x}{2}\right)^2 - 2\tanh\left(\frac{x}{2}\right)\sqrt{a+b}\right)}{2\sqrt{a}\sqrt{a+b}}$	78
risch	$\frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} - 2a^2 - 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}} - \frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} + 2a^2 + 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}}$	128

[In] int(1/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 1/2/a^(1/2)/(a+b)^(1/2)*ln((a+b)^(1/2)*tanh(1/2*x)^2+2*tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))-1/2/a^(1/2)/(a+b)^(1/2)*ln((a+b)^(1/2)*tanh(1/2*x)^2-2*tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(21) = 42.

Time = 0.26 (sec) , antiderivative size = 293, normalized size of antiderivative = 10.10

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\log \left(\frac{b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2(2ab + b^2) \cosh(x)^2 + 2(3b^2 \cosh(x)^2 + 2ab + b^2) \sinh(x)^2 + 8a^2 + 8ab + b^2 + 4(b^2 \cosh(x)^2 + 2ab + b^2) \sinh(x)}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a + b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 2a + b) \sinh(x)} \right)}{2\sqrt{a^2 + ab}}$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/2*log((b^2*cosh(x)^4 + 4*b^2*cosh(x)*sinh(x)^3 + b^2*sinh(x)^4 + 2*(2*a*b + b^2)*cosh(x)^2 + 2*(3*b^2*cosh(x)^2 + 2*a*b + b^2)*sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*cosh(x)^3 + (2*a*b + b^2)*cosh(x))*sinh(x) - 4*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(a^2 + a*b))/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b))/sqrt(a^2 + a*b), sqrt(-a^2 - a*b)*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*sinh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-a^2 - a*b)/(a^2 + a*b))/(a^2 + a*b)]

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10924 vs. 2(27) = 54.

Time = 24.28 (sec) , antiderivative size = 10924, normalized size of antiderivative = 376.69

$$\int \frac{1}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cosh(x)**2),x)

[Out] Piecewise((zoo*tanh(x/2)/(tanh(x/2)**2 + 1), Eq(a, 0) & Eq(b, 0)), (-tanh(x/2)/(2*b) - 1/(2*b*tanh(x/2)), Eq(a, -b)), (2*tanh(x/2)/(b*(tanh(x/2)**2 + 1)), Eq(a, 0)), (-a**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))

$$\begin{aligned}
& + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a \\
& + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq \\
& rt(-a*b)/(a + b))) + a**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b) \\
&)*log(sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a* \\
& *4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a \\
& + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sq \\
& rt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a* \\
& *3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a \\
& + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/ \\
& (a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(\\
& a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(\\
& a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/ \\
& (a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2* \\
& sqrt(-a*b)/(a + b))) - a**3*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + \\
& b))*log(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2 \\
& *a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b \\
& /(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2 \\
& *sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8 \\
& *a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/ \\
& (a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - \\
& b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b \\
&)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sq \\
& rt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt \\
& (a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + \\
& 2*sqrt(-a*b)/(a + b))) + a**3*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b))*log(sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/ \\
& (2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - \\
& b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - \\
& 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + \\
& 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(\\
& a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) \\
& - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a \\
& *b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))* \\
& sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sq \\
& rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) \\
& + 2*sqrt(-a*b)/(a + b))) + 10*a**2*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(- \\
& a*b)/(a + b))*log(-sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tan \\
& h(x/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(\\
& a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(\\
& a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a \\
& + b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b \\
&))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a \\
& /(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2 \\
& *sqrt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(
\end{aligned}$$


```

a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))) + b**2*sq
rt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*log(sqrt(a/(a +
b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*sqrt(a/(a + b)
- b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a
*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))
*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*sqrt(-a*b)*sq
rt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b)
+ 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(
-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 2*a*b**
3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a
+ b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a +
b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b
))) + b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b))*lo
g(-sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x/2))/(2*a**4*
sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a +
b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(
-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) + 8*a**3*
sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b
) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a + b) - b/(a
+ b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a +
b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(
a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*b)*sqrt(a/(a
+ b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq
rt(-a*b)/(a + b))) - b**2*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*
b)/(a + b))*log(sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b)) + tanh(x
/2))/(2*a**4*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a +
b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**3*b*sqrt(a/(a + b) - b/(a +
b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a +
b)) + 8*a**3*sqrt(-a*b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*
sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 10*a**2*b**2*sqrt(a/(a
+ b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a + b) + 2*sq
rt(-a*b)/(a + b)) + 2*a*b**3*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a +
b))*sqrt(a/(a + b) - b/(a + b) + 2*sqrt(-a*b)/(a + b)) - 8*a*b**2*sqrt(-a*
b)*sqrt(a/(a + b) - b/(a + b) - 2*sqrt(-a*b)/(a + b))*sqrt(a/(a + b) - b/(a
+ b) + 2*sqrt(-a*b)/(a + b))), True))

```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(21) = 42.

Time = 0.28 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.83

$$\int \frac{1}{a + b \cosh^2(x)} dx = -\frac{\log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{2\sqrt{(a+b)a}}$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $-1/2*\log((b*e^{(-2*x)} + 2*a + b - 2*\sqrt{(a + b)*a}))/ (b*e^{(-2*x)} + 2*a + b + 2*\sqrt{(a + b)*a}))/\sqrt{(a + b)*a}$

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.34

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\arctan\left(\frac{be^{(2x)}+2a+b}{2\sqrt{-a^2-ab}}\right)}{\sqrt{-a^2-ab}}$$

[In] integrate(1/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] $\arctan(1/2*(b*e^{(2*x)} + 2*a + b)/\sqrt{-a^2 - a*b}))/\sqrt{-a^2 - a*b}$

Mupad [B] (verification not implemented)

Time = 0.00 (sec) , antiderivative size = 267, normalized size of antiderivative = 9.21

$$\int \frac{1}{a + b \cosh^2(x)} dx = \frac{\operatorname{atan}\left(\frac{b^2 e^{2x} (-a^2 - ba)^{3/2} \left(\frac{4(4a+2b)(8a^3+12a^2b+4ab^2)}{b^5(-a^2-ba)^{3/2}\sqrt{-a(a+b)}} + \frac{2(8a^2+8ab+b^2)(8a^2\sqrt{-a^2-ba}+b^2\sqrt{-a^2-ba}+8ab\sqrt{-a^2-ba})}{ab^5(a+b)(-a^2-ba)^{3/2}} \right)}{4} + \frac{(2a^2b)}{b^3}\right)}{\sqrt{-a^2 - ba}}$$

[In] int(1/(a + b*cosh(x)^2),x)

[Out] $-\operatorname{atan}((b^2*\exp(2*x))*(-a*b - a^2)^{(3/2)}*((4*(4*a + 2*b))*(4*a*b^2 + 12*a^2*b + 8*a^3)))/(b^5*(-a*b - a^2)^{(3/2)}*(-a*(a + b))^{(1/2)}) + (2*(8*a*b + 8*a^2 + b^2)*(8*a^2*(-a*b - a^2)^{(1/2)} + b^2*(-a*b - a^2)^{(1/2)} + 8*a*b*(-a*b - a^2)^{(1/2)}))/(a*b^5*(a + b)*(-a*b - a^2)^{(3/2)})/4 + ((2*a*b^2 + 2*a^2*b)*(4*a + 2*b))/(b^3*(-a*(a + b))^{(1/2)}) + ((b^2*(-a*b - a^2)^{(1/2)} + 2*a*b*(-a*b - a^2)^{(1/2)})*(8*a*b + 8*a^2 + b^2))/(2*a*b^3*(a + b))/(-a*b - a^2)^{(1/2)}$

3.28 $\int \frac{\operatorname{sech}(x)}{a+b \cosh^2(x)} dx$

Optimal result	238
Rubi [A] (verified)	238
Mathematica [A] (verified)	239
Maple [B] (verified)	240
Fricas [B] (verification not implemented)	240
Sympy [F]	241
Maxima [F]	241
Giac [F]	241
Mupad [B] (verification not implemented)	242

Optimal result

Integrand size = 13, antiderivative size = 41

$$\int \frac{\operatorname{sech}(x)}{a+b \cosh^2(x)} dx = \frac{\arctan(\sinh(x))}{a} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}}$$

[Out] $\arctan(\sinh(x))/a - \arctan(\sinh(x)*b^{(1/2)}/(a+b)^{(1/2)})*b^{(1/2)}/a/(a+b)^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.308$, Rules used = {3265, 400, 209, 211}

$$\int \frac{\operatorname{sech}(x)}{a+b \cosh^2(x)} dx = \frac{\arctan(\sinh(x))}{a} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}}$$

[In] `Int[Sech[x]/(a + b*Cosh[x]^2), x]`

[Out] `ArcTan[Sinh[x]]/a - (Sqrt[b]*ArcTan[(Sqrt[b]*Sinh[x])/Sqrt[a + b]])/(a*Sqrt[a + b])`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 400

Int[1/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[b/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0]

Rule 3265

Int[sin[(e_) + (f_)*(x_)]^(m_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{(1+x^2)(a+b+bx^2)} dx, x, \sinh(x)\right) \\ &= \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sinh(x)\right)}{a} - \frac{b \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{a} \\ &= \frac{\arctan(\sinh(x))}{a} - \frac{\sqrt{b} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a\sqrt{a+b}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.10

$$\int \frac{\text{sech}(x)}{a + b \cosh^2(x)} dx = \frac{\sqrt{b} \arctan\left(\frac{\sqrt{a+b} \text{csch}(x)}{\sqrt{b}}\right)}{a\sqrt{a+b}} + \frac{2 \arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{a}$$

[In] Integrate[Sech[x]/(a + b*Cosh[x]^2), x]

[Out] (Sqrt[b]*ArcTan[(Sqrt[a + b]*Csch[x])/Sqrt[b]])/(a*Sqrt[a + b]) + (2*ArcTan[Tanh[x/2]])/a

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(33) = 66.

Time = 0.31 (sec) , antiderivative size = 85, normalized size of antiderivative = 2.07

method	result	size
default	$\frac{2 \arctan(\tanh(\frac{x}{2}))}{a} - \frac{2b \left(\frac{\arctan\left(\frac{2 \tanh(\frac{x}{2}) \sqrt{a+b+2\sqrt{a}}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh(\frac{x}{2}) \sqrt{a+b-2\sqrt{a}}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} \right)}{a}$	85
risch	$\frac{i \ln(e^x+i)}{a} - \frac{i \ln(e^x-i)}{a} + \frac{\sqrt{-(a+b)b} \ln\left(e^{2x} - \frac{2\sqrt{-(a+b)b} e^x}{b} - 1\right)}{2(a+b)a} - \frac{\sqrt{-(a+b)b} \ln\left(e^{2x} + \frac{2\sqrt{-(a+b)b} e^x}{b} - 1\right)}{2(a+b)a}$	106

[In] int(sech(x)/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 2/a*arctan(tanh(1/2*x))-2*b/a*(1/2/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)+2*a^(1/2))/b^(1/2))+1/2/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)-2*a^(1/2))/b^(1/2)))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. 2(33) = 66.

Time = 0.27 (sec) , antiderivative size = 360, normalized size of antiderivative = 8.78

$$\int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{\sqrt{-\frac{b}{a+b}} \log\left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 - 2(2a+3b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a - 3b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a+3b) \cosh(x) \sinh(x)^2 - a - b) \sinh(x)}{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(2a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 - 2a - 3b) \sinh(x)^2 + 4(b \cosh(x)^3 - (2a+3b) \cosh(x) \sinh(x)^2 - a - b) \sinh(x)}\right)}{a}$$

$$+ \frac{\sqrt{\frac{b}{a+b}} \arctan\left(\frac{1}{2} \sqrt{\frac{b}{a+b}} (\cosh(x) + \sinh(x))\right) + \sqrt{\frac{b}{a+b}} \arctan\left(\frac{(b \cosh(x)^3 + 3b \cosh(x) \sinh(x)^2 + b \sinh(x)^3 + (4a+3b) \cosh(x) \sinh(x)^2 - a - b) \sinh(x)}{2b}\right)}{a}$$

[In] integrate(sech(x)/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/2*(sqrt(-b/(a + b))*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a + 3*b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*a - 3*b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a + 3*b)*cosh(x))*sinh(x) - 4*((a + b)*cosh(x)^3 + 3*(a + b)*cosh(x)*sinh(x)^2 + (a + b)*sinh(x)^3 - (a + b)*cosh(x) + (3*(a + b)*cosh(x)*sinh(x)^2 - a - b)*sinh(x))*sqrt(-b/(a + b)) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a +

b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 4*arctan(cosh(x) + sinh(x))/a, -(sqrt(b/(a + b))*arctan(1/2*sqrt(b/(a + b))*(cosh(x) + sinh(x))) + sqrt(b/(a + b))*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + 3*b)*cosh(x) + (3*b*cosh(x)^2 + 4*a + 3*b)*sinh(x))*sqrt(b/(a + b)))/b) - 2*arctan(cosh(x) + sinh(x))/a]

Sympy [F]

$$\int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx$$

[In] integrate(sech(x)/(a+b*cosh(x)**2),x)

[Out] Integral(sech(x)/(a + b*cosh(x)**2), x)

Maxima [F]

$$\int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(sech(x)/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 2*arctan(e^x)/a - 2*integrate((b*e^(3*x) + b*e^x)/(a*b*e^(4*x) + a*b + 2*(2*a^2 + a*b)*e^(2*x)), x)

Giac [F]

$$\int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)}{b \cosh(x)^2 + a} dx$$

[In] integrate(sech(x)/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 2.23 (sec) , antiderivative size = 208, normalized size of antiderivative = 5.07

$$\int \frac{\operatorname{sech}(x)}{a + b \cosh^2(x)} dx = \frac{2 \operatorname{atan}\left(\frac{e^x (16 (a^2)^{3/2} + 9 b^2 \sqrt{a^2} + 24 a b \sqrt{a^2})}{16 a^3 + 24 a^2 b + 9 a b^2}\right)}{\sqrt{a^2}} - \frac{\sqrt{b} \left(2 \operatorname{atan}\left(\frac{\sqrt{b} e^x \sqrt{a^2 (a+b)}}{2 a (a+b)}\right) + 2 \operatorname{atan}\left(\frac{4 a^4 e^x + 8 a^3 b e^x + 4 a^2 b^2 e^x - b e^x \sqrt{a^2 (a+b)} \sqrt{a^3 + b a^2} + b e^{3x} \sqrt{a^2 (a+b)} \sqrt{a^3 + b a^2}}{\sqrt{b} \sqrt{a^2 (a+b)} (2 a^2 + 2 b a)}\right) \right)}{2 \sqrt{a^3 + b a^2}}$$

[In] int(1/(cosh(x)*(a + b*cosh(x)^2)),x)

```
[Out] (2*atan((exp(x)*(16*(a^2)^(3/2) + 9*b^2*(a^2)^(1/2) + 24*a*b*(a^2)^(1/2)))/
(9*a*b^2 + 24*a^2*b + 16*a^3)))/(a^2)^(1/2) - (b^(1/2)*(2*atan((b^(1/2)*exp
(x)*(a^2*(a + b))^(1/2))/(2*a*(a + b))) + 2*atan((4*a^4*exp(x) + 8*a^3*b*ex
p(x) + 4*a^2*b^2*exp(x) - b*exp(x)*(a^2*(a + b))^(1/2)*(a^2*b + a^3)^(1/2)
+ b*exp(3*x)*(a^2*(a + b))^(1/2)*(a^2*b + a^3)^(1/2)))/(b^(1/2)*(a^2*(a + b)
)^(1/2)*(2*a*b + 2*a^2)))))/(2*(a^2*b + a^3)^(1/2))
```

3.29 $\int \frac{\operatorname{sech}^2(x)}{a+b \cosh^2(x)} dx$

Optimal result	243
Rubi [A] (verified)	243
Mathematica [A] (verified)	244
Maple [B] (verified)	245
Fricas [B] (verification not implemented)	245
Sympy [F]	246
Maxima [B] (verification not implemented)	246
Giac [A] (verification not implemented)	246
Mupad [B] (verification not implemented)	247

Optimal result

Integrand size = 15, antiderivative size = 38

$$\int \frac{\operatorname{sech}^2(x)}{a+b \cosh^2(x)} dx = -\frac{\operatorname{barctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{a^{3/2} \sqrt{a+b}} + \frac{\tanh(x)}{a}$$

[Out] $-b \cdot \operatorname{arctanh}(a^{1/2} \cdot \tanh(x) / (a+b)^{1/2}) / a^{3/2} / (a+b)^{1/2} + \tanh(x) / a$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3266, 464, 214}

$$\int \frac{\operatorname{sech}^2(x)}{a+b \cosh^2(x)} dx = \frac{\tanh(x)}{a} - \frac{\operatorname{barctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{a^{3/2} \sqrt{a+b}}$$

[In] $\operatorname{Int}[\operatorname{Sech}[x]^2 / (a + b \cdot \operatorname{Cosh}[x]^2), x]$

[Out] $-((b \cdot \operatorname{ArcTanh}[(\operatorname{Sqrt}[a] \cdot \operatorname{Tanh}[x]) / \operatorname{Sqrt}[a + b]]) / (a^{3/2} \cdot \operatorname{Sqrt}[a + b])) + \operatorname{Tanh}[x] / a$

Rule 214

$\operatorname{Int}[(a_0 + (b_0) \cdot (x_0)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2] / a) \cdot \operatorname{ArcTanh}[x / \operatorname{Rt}[-a/b, 2]], x] / ; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b]$

Rule 464

$\operatorname{Int}[(e_0 \cdot (x_0))^m \cdot (a_0 + (b_0) \cdot (x_0)^n)^p \cdot ((c_0) + (d_0) \cdot (x_0)^n), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[c \cdot (e \cdot x)^{m+1} \cdot (a + b \cdot x^n)^{p+1} / (a \cdot e^{m+1})],$

```
x] + Dist[(a*d*(m + 1) - b*c*(m + n*(p + 1) + 1))/(a*e^n*(m + 1)), Int[(e*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[b*c - a*d, 0] && (IntegerQ[n] || GtQ[e, 0]) && ((GtQ[n, 0] && LtQ[m, -1]) || (LtQ[n, 0] && GtQ[m + n, -1])) && !ILtQ[p, -1]
```

Rule 3266

```
Int[sin[(e_.) + (f_.)*(x_.)]^(m_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\text{Subst}\left(\int \frac{1-x^2}{x^2(a-(a+b)x^2)} dx, x, \coth(x)\right) \\ &= \frac{\tanh(x)}{a} - \frac{b\text{Subst}\left(\int \frac{1}{a+(-a-b)x^2} dx, x, \coth(x)\right)}{a} \\ &= -\frac{\text{barctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{a^{3/2}\sqrt{a+b}} + \frac{\tanh(x)}{a} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.37 (sec) , antiderivative size = 38, normalized size of antiderivative = 1.00

$$\int \frac{\text{sech}^2(x)}{a + b \cosh^2(x)} dx = -\frac{\text{barctanh}\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+b}}\right)}{a^{3/2}\sqrt{a+b}} + \frac{\tanh(x)}{a}$$

```
[In] Integrate[Sech[x]^2/(a + b*Cosh[x]^2), x]
```

```
[Out] -((b*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(a^(3/2)*Sqrt[a + b])) + Tanh[x]/a
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 106 vs. $2(30) = 60$.

Time = 0.53 (sec) , antiderivative size = 107, normalized size of antiderivative = 2.82

method	result	size
default	$\frac{2 \tanh\left(\frac{x}{2}\right)}{a\left(1+\tanh\left(\frac{x}{2}\right)^2\right)} + \frac{2b\left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2+2 \tanh\left(\frac{x}{2}\right)\sqrt{a+b}\right)}{4\sqrt{a} \sqrt{a+b}} + \frac{\ln\left(-\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2+2 \tanh\left(\frac{x}{2}\right)\sqrt{a-b}\right)}{4\sqrt{a} \sqrt{a+b}}\right)}{a}$	107
risch	$-\frac{2}{a(1+e^{2x})} + \frac{b \ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} + 2a^2 + 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}a} - \frac{b \ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab} + b\sqrt{a^2+ab} - 2a^2 - 2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}a}$	149

[In] `int(sech(x)^2/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $2/a*\tanh(1/2*x)/(1+\tanh(1/2*x)^2)+2*b/a*(-1/4/a^{(1/2)/(a+b)^{(1/2)}*\ln((a+b)^{(1/2)*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^{(1/2)+(a+b)^{(1/2)}+1/4/a^{(1/2)/(a+b)^{(1/2)*\ln(-(a+b)^{(1/2)*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^{(1/2)-(a+b)^{(1/2))}}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 148 vs. $2(30) = 60$.

Time = 0.26 (sec) , antiderivative size = 457, normalized size of antiderivative = 12.03

$$\int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{\left((b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + b) \sqrt{a^2 + ab} \log\left(\frac{b^2 \cosh(x)^4 + 4b^2 \cosh(x) \sinh(x)^3 + b^2 \sinh(x)^4 + 2b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + b}{2(a^3 + a^2b + (a^3 + a^2b) \cosh(x)^2 + 2(a^3 + a^2b) \cosh(x) \sinh(x) + (a^3 + a^2b) \sinh(x)^2)} \right) \right.}{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + b) \sqrt{-a^2 - ab} \arctan\left(\frac{(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + b) \sqrt{-a^2 - ab}}{2(a^2 + ab)} \right)}$$

[In] `integrate(sech(x)^2/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/2*((b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + b)*\sqrt{a^2 + a*b}) * \log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x)^2 + 2*(3*b^2*\cosh(x)^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) + 4*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{a^2 + a*b})/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*$

$\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) - 4*a^2 - 4*a*b)/(a^3 + a^2*b + (a^3 + a^2*b)*\cosh(x)^2 + 2*(a^3 + a^2*b)*\cosh(x)*\sinh(x) + (a^3 + a^2*b)*\sinh(x)^2), -((b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + b)*\sqrt{-a^2 - a*b}*\arctan(1/2*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{-a^2 - a*b}/(a^2 + a*b)) + 2*a^2 + 2*a*b)/(a^3 + a^2*b + (a^3 + a^2*b)*\cosh(x)^2 + 2*(a^3 + a^2*b)*\cosh(x)*\sinh(x) + (a^3 + a^2*b)*\sinh(x)^2)]$

Sympy [F]

$$\int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx$$

[In] integrate(sech(x)**2/(a+b*cosh(x)**2),x)

[Out] Integral(sech(x)**2/(a + b*cosh(x)**2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 70 vs. 2(30) = 60.

Time = 0.28 (sec) , antiderivative size = 70, normalized size of antiderivative = 1.84

$$\int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx = \frac{b \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{2\sqrt{(a+b)aa}} + \frac{2}{ae^{(-2x)} + a}$$

[In] integrate(sech(x)^2/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] 1/2*b*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) + 2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*a) + 2/(a*e^(-2*x) + a)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 58, normalized size of antiderivative = 1.53

$$\int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx = -\frac{b \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{\sqrt{-a^2 - aba}} - \frac{2}{a(e^{(2x)} + 1)}$$

[In] integrate(sech(x)^2/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] -b*arctan(1/2*(b*e^(2*x) + 2*a + b)/sqrt(-a^2 - a*b))/(sqrt(-a^2 - a*b)*a) - 2/(a*(e^(2*x) + 1))

Mupad [B] (verification not implemented)

Time = 0.31 (sec) , antiderivative size = 108, normalized size of antiderivative = 2.84

$$\int \frac{\operatorname{sech}^2(x)}{a + b \cosh^2(x)} dx = \frac{b \ln \left(\frac{4e^{2x}}{a} - \frac{2(b+2ae^{2x}+be^{2x})}{a^{3/2}\sqrt{a+b}} \right)}{2a^{3/2}\sqrt{a+b}} - \frac{2}{a(e^{2x}+1)} - \frac{b \ln \left(\frac{4e^{2x}}{a} + \frac{2(b+2ae^{2x}+be^{2x})}{a^{3/2}\sqrt{a+b}} \right)}{2a^{3/2}\sqrt{a+b}}$$

[In] int(1/(cosh(x)^2*(a + b*cosh(x)^2)),x)

```
[Out] (b*log((4*exp(2*x))/a - (2*(b + 2*a*exp(2*x) + b*exp(2*x)))/(a^(3/2)*(a + b)^(1/2))))/(2*a^(3/2)*(a + b)^(1/2)) - 2/(a*(exp(2*x) + 1)) - (b*log((4*exp(2*x))/a + (2*(b + 2*a*exp(2*x) + b*exp(2*x)))/(a^(3/2)*(a + b)^(1/2))))/(2*a^(3/2)*(a + b)^(1/2))
```

3.30 $\int \frac{\operatorname{sech}^3(x)}{a+b \cosh^2(x)} dx$

Optimal result	248
Rubi [A] (verified)	248
Mathematica [A] (verified)	250
Maple [B] (verified)	250
Fricas [B] (verification not implemented)	251
Sympy [F]	252
Maxima [F]	252
Giac [F]	252
Mupad [B] (verification not implemented)	253

Optimal result

Integrand size = 15, antiderivative size = 59

$$\int \frac{\operatorname{sech}^3(x)}{a+b \cosh^2(x)} dx = \frac{(a-2b) \arctan(\sinh(x))}{2a^2} + \frac{b^{3/2} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a^2 \sqrt{a+b}} + \frac{\operatorname{sech}(x) \tanh(x)}{2a}$$

[Out] 1/2*(a-2*b)*arctan(sinh(x))/a^2+b^(3/2)*arctan(sinh(x)*b^(1/2)/(a+b)^(1/2))/a^2/(a+b)^(1/2)+1/2*sech(x)*tanh(x)/a

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3265, 425, 536, 209, 211}

$$\int \frac{\operatorname{sech}^3(x)}{a+b \cosh^2(x)} dx = \frac{b^{3/2} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a^2 \sqrt{a+b}} + \frac{(a-2b) \arctan(\sinh(x))}{2a^2} + \frac{\tanh(x) \operatorname{sech}(x)}{2a}$$

[In] Int[Sech[x]^3/(a + b*Cosh[x]^2), x]

[Out] ((a - 2*b)*ArcTan[Sinh[x]])/(2*a^2) + (b^(3/2)*ArcTan[(Sqrt[b]*Sinh[x])/Sqrt[a + b]])/(a^2*Sqrt[a + b]) + (Sech[x]*Tanh[x])/(2*a)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 211

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Rule 425

`Int[((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !(IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]`

Rule 536

`Int[((e_) + (f_)*(x_)^(n_))/((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]`

Rule 3265

`Int[sin[(e_) + (f_)*(x_)^(m_)]*((a_) + (b_)*sin[(e_) + (f_)*(x_)^(p_)])^(p_), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{(1+x^2)^2(a+b+bx^2)} dx, x, \sinh(x)\right) \\
 &= \frac{\text{sech}(x) \tanh(x)}{2a} - \frac{\text{Subst}\left(\int \frac{-a+b-bx^2}{(1+x^2)(a+b+bx^2)} dx, x, \sinh(x)\right)}{2a} \\
 &= \frac{\text{sech}(x) \tanh(x)}{2a} + \frac{(a-2b)\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sinh(x)\right)}{2a^2} + \frac{b^2\text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{a^2} \\
 &= \frac{(a-2b) \arctan(\sinh(x))}{2a^2} + \frac{b^{3/2} \arctan\left(\frac{\sqrt{b}\sinh(x)}{\sqrt{a+b}}\right)}{a^2\sqrt{a+b}} + \frac{\text{sech}(x) \tanh(x)}{2a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 58, normalized size of antiderivative = 0.98

$$\int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx$$

$$= \frac{-\frac{2b^{3/2} \arctan\left(\frac{\sqrt{a+b} \operatorname{csch}(x)}{\sqrt{b}}\right)}{\sqrt{a+b}} + 2(a-2b) \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + a \operatorname{sech}(x) \tanh(x)}{2a^2}$$

[In] Integrate[Sech[x]^3/(a + b*Cosh[x]^2),x]

[Out] ((-2*b^(3/2)*ArcTan[(Sqrt[a + b]*Csch[x])/Sqrt[b]])/Sqrt[a + b] + 2*(a - 2*b)*ArcTan[Tanh[x/2]] + a*Sech[x]*Tanh[x])/(2*a^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 122 vs. 2(47) = 94.

Time = 0.84 (sec) , antiderivative size = 123, normalized size of antiderivative = 2.08

method	result
default	$\frac{2 \left(-\frac{a \tanh\left(\frac{x}{2}\right)^3}{2} + \frac{a \tanh\left(\frac{x}{2}\right)}{2} \right)}{\left(1 + \tanh\left(\frac{x}{2}\right)^2\right)^2} + (a-2b) \arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{a^2} + \frac{2b^2 \left(\frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b} + 2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} + \frac{\arctan\left(\frac{2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b} - 2\sqrt{a}}{2\sqrt{b}}\right)}{2\sqrt{a+b}\sqrt{b}} \right)}{a^2}$
risch	$\frac{e^x (e^{2x} - 1)}{(1 + e^{2x})^2 a} + \frac{i \ln(e^x + i)}{2a} - \frac{ib \ln(e^x + i)}{a^2} - \frac{i \ln(e^x - i)}{2a} + \frac{ib \ln(e^x - i)}{a^2} + \frac{\sqrt{-(a+b)b} b \ln\left(e^{2x} + \frac{2\sqrt{-(a+b)b} e^x}{b} - 1\right)}{2(a+b)a^2} - \frac{\sqrt{-(a+b)b}}{2(a+b)a^2}$

[In] int(sech(x)^3/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] 2/a^2*((-1/2*a*tanh(1/2*x)^3+1/2*a*tanh(1/2*x))/(1+tanh(1/2*x)^2)^2+1/2*(a-2*b)*arctan(tanh(1/2*x)))+2*b^2/a^2*(1/2/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)+2*a^(1/2))/b^(1/2))+1/2/(a+b)^(1/2)/b^(1/2)*arctan(1/2*(2*tanh(1/2*x)*(a+b)^(1/2)-2*a^(1/2))/b^(1/2)))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 444 vs. 2(47) = 94.

Time = 0.28 (sec) , antiderivative size = 963, normalized size of antiderivative = 16.32

$$\int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(sech(x)^3/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] [1/2*(2*a*cosh(x)^3 + 6*a*cosh(x)*sinh(x)^2 + 2*a*sinh(x)^3 + (b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + b*cosh(x))*sinh(x) + b)*sqrt(-b/(a + b))*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 - 2*(2*a + 3*b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 - 2*a - 3*b)*sinh(x)^2 + 4*(b*cosh(x)^3 - (2*a + 3*b)*cosh(x))*sinh(x) + 4*((a + b)*cosh(x)^3 + 3*(a + b)*cosh(x)*sinh(x)^2 + (a + b)*sinh(x)^3 - (a + b)*cosh(x) + (3*(a + b)*cosh(x)^2 - a - b)*sinh(x)))*sqrt(-b/(a + b)) + b)/(b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(2*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 2*a + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + (2*a + b)*cosh(x))*sinh(x) + b)) + 2*((a - 2*b)*cosh(x)^4 + 4*(a - 2*b)*cosh(x)*sinh(x)^3 + (a - 2*b)*sinh(x)^4 + 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a - 2*b)*cosh(x)^2 + a - 2*b)*sinh(x)^2 + 4*((a - 2*b)*cosh(x)^3 + (a - 2*b)*cosh(x))*sinh(x) + a - 2*b)*arctan(cosh(x) + sinh(x)) - 2*a*cosh(x) + 2*(3*a*cosh(x)^2 - a)*sinh(x))/(a^2*cosh(x)^4 + 4*a^2*cosh(x)*sinh(x)^3 + a^2*sinh(x)^4 + 2*a^2*cosh(x)^2 + 2*(3*a^2*cosh(x)^2 + a^2)*sinh(x)^2 + a^2 + 4*(a^2*cosh(x)^3 + a^2*cosh(x))*sinh(x)), (a*cosh(x)^3 + 3*a*cosh(x)*sinh(x))^2 + a*sinh(x)^3 + (b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + b*cosh(x))*sinh(x) + b)*sqrt(b/(a + b))*arctan(1/2*sqrt(b/(a + b))*(cosh(x) + sinh(x))) + (b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*b*cosh(x)^2 + 2*(3*b*cosh(x)^2 + b)*sinh(x)^2 + 4*(b*cosh(x)^3 + b*cosh(x))*sinh(x) + b)*sqrt(b/(a + b))*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + 3*b)*cosh(x) + (3*b*cosh(x)^2 + 4*a + 3*b)*sinh(x))*sqrt(b/(a + b)))/b) + ((a - 2*b)*cosh(x)^4 + 4*(a - 2*b)*cosh(x)*sinh(x)^3 + (a - 2*b)*sinh(x)^4 + 2*(a - 2*b)*cosh(x)^2 + 2*(3*(a - 2*b)*cosh(x)^2 + a - 2*b)*sinh(x)^2 + 4*((a - 2*b)*cosh(x)^3 + (a - 2*b)*cosh(x))*sinh(x) + a - 2*b)*arctan(cosh(x) + sinh(x)) - a*cosh(x) + (3*a*cosh(x)^2 - a)*sinh(x))/(a^2*cosh(x)^4 + 4*a^2*cosh(x)*sinh(x)^3 + a^2*sinh(x)^4 + 2*a^2*cosh(x)^2 + 2*(3*a^2*cosh(x)^2 + a^2)*sinh(x)^2 + a^2 + 4*(a^2*cosh(x)^3 + a^2*cosh(x))*sinh(x))]

Sympy [F]

$$\int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx$$

```
[In] integrate(sech(x)**3/(a+b*cosh(x)**2),x)
```

```
[Out] Integral(sech(x)**3/(a + b*cosh(x)**2), x)
```

Maxima [F]

$$\int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)^3}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(sech(x)^3/(a+b*cosh(x)^2),x, algorithm="maxima")
```

```
[Out] (e^(3*x) - e^x)/(a*e^(4*x) + 2*a*e^(2*x) + a) + (a - 2*b)*arctan(e^x)/a^2 +
      8*integrate(1/4*(b^2*e^(3*x) + b^2*e^x)/(a^2*b*e^(4*x) + a^2*b + 2*(2*a^3
      + a^2*b)*e^(2*x)), x)
```

Giac [F]

$$\int \frac{\operatorname{sech}^3(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)^3}{b \cosh(x)^2 + a} dx$$

```
[In] integrate(sech(x)^3/(a+b*cosh(x)^2),x, algorithm="giac")
```

```
[Out] sage0*x
```


3.31 $\int \frac{\operatorname{sech}^4(x)}{a+b \cosh^2(x)} dx$

Optimal result	254
Rubi [A] (verified)	254
Mathematica [A] (verified)	255
Maple [B] (verified)	256
Fricas [B] (verification not implemented)	256
Sympy [F]	257
Maxima [B] (verification not implemented)	257
Giac [F]	258
Mupad [B] (verification not implemented)	258

Optimal result

Integrand size = 15, antiderivative size = 55

$$\int \frac{\operatorname{sech}^4(x)}{a+b \cosh^2(x)} dx = \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{a^{5/2} \sqrt{a+b}} + \frac{(a-b) \tanh(x)}{a^2} - \frac{\tanh^3(x)}{3a}$$

[Out] $b^2 \operatorname{arctanh}(a^{1/2} \tanh(x) / (a+b)^{1/2}) / a^{5/2} / (a+b)^{1/2} + (a-b) \tanh(x) / a^2 - 1/3 \tanh(x)^3 / a$

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3266, 472, 214}

$$\int \frac{\operatorname{sech}^4(x)}{a+b \cosh^2(x)} dx = \frac{b^2 \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{a^{5/2} \sqrt{a+b}} + \frac{(a-b) \tanh(x)}{a^2} - \frac{\tanh^3(x)}{3a}$$

[In] $\operatorname{Int}[\operatorname{Sech}[x]^4 / (a + b \operatorname{Cosh}[x]^2), x]$

[Out] $(b^2 \operatorname{ArcTanh}[(\operatorname{Sqrt}[a] \operatorname{Tanh}[x]) / \operatorname{Sqrt}[a + b]]) / (a^{5/2} \operatorname{Sqrt}[a + b]) + ((a - b) \operatorname{Tanh}[x]) / a^2 - \operatorname{Tanh}[x]^3 / (3 * a)$

Rule 214

$\operatorname{Int}[(a_0 + (b_0)(x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2] / a) \operatorname{ArcTanh}[x / \operatorname{Rt}[-a/b, 2]], x] / ; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b]$

Rule 472

```
Int[(((e_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_))/((c_) + (d_.)*(x_)^(n_)), x_Symbol] :> Int[ExpandIntegrand[(e*x)^(m)*((a + b*x^n)^p/(c + d*x^n)), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && IGtQ[p, 0] && (IntegerQ[m] || IGtQ[2*(m + 1), 0] || !RationalQ[m])
```

Rule 3266

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + (a + b)*ff^2*x^2)^p/(1 + ff^2*x^2)^(m/2 + p + 1)), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[m/2] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{(1-x^2)^2}{x^4 (a - (a+b)x^2)} dx, x, \coth(x) \right) \\
 &= \text{Subst} \left(\int \left(\frac{1}{ax^4} + \frac{-a+b}{a^2x^2} + \frac{b^2}{a^2(a - (a+b)x^2)} \right) dx, x, \coth(x) \right) \\
 &= \frac{(a-b) \tanh(x)}{a^2} - \frac{\tanh^3(x)}{3a} + \frac{b^2 \text{Subst} \left(\int \frac{1}{a - (a+b)x^2} dx, x, \coth(x) \right)}{a^2} \\
 &= \frac{b^2 \operatorname{arctanh} \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}} \right)}{a^{5/2} \sqrt{a+b}} + \frac{(a-b) \tanh(x)}{a^2} - \frac{\tanh^3(x)}{3a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00

$$\int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx = \frac{b^2 \operatorname{arctanh} \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}} \right)}{a^{5/2} \sqrt{a+b}} + \frac{(2a - 3b + a \operatorname{sech}^2(x)) \tanh(x)}{3a^2}$$

```
[In] Integrate[Sech[x]^4/(a + b*Cosh[x]^2), x]
```

```
[Out] (b^2*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(a^(5/2)*Sqrt[a + b]) + ((2*a - 3*b + a*Sech[x]^2)*Tanh[x])/(3*a^2)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(45) = 90.

Time = 1.19 (sec) , antiderivative size = 142, normalized size of antiderivative = 2.58

method	result
default	$\frac{2\left((-a+b)\tanh\left(\frac{x}{2}\right)^5 + \left(-\frac{2a}{3}+2b\right)\tanh\left(\frac{x}{2}\right)^3 + (-a+b)\tanh\left(\frac{x}{2}\right)\right)}{a^2\left(1+\tanh\left(\frac{x}{2}\right)^2\right)^3} - \frac{2b^2\left(-\frac{\ln\left(\sqrt{a+b}\tanh\left(\frac{x}{2}\right)^2+2\tanh\left(\frac{x}{2}\right)\sqrt{a+b}\right)}{4\sqrt{a}\sqrt{a+b}} + \frac{\ln\left(-\sqrt{a+b}\tanh\left(\frac{x}{2}\right)\right)}{a^2}\right)}{a^2}$
risch	$-\frac{2(-3be^{4x}+6ae^{2x}-6be^{2x}+2a-3b)}{3(1+e^{2x})^3a^2} + \frac{b^2\ln\left(e^{2x}+\frac{2a\sqrt{a^2+ab}+b\sqrt{a^2+ab}-2a^2-2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}a^2} - \frac{b^2\ln\left(e^{2x}+\frac{2a\sqrt{a^2+ab}+b\sqrt{a^2+ab}+2a^2+2ab}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}a^2}$

[In] int(sech(x)^4/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)

[Out] $-2/a^2*((-a+b)*\tanh(1/2*x)^5+(-2/3*a+2*b)*\tanh(1/2*x)^3+(-a+b)*\tanh(1/2*x))/(1+\tanh(1/2*x)^2)^3-2*b^2/a^2*(-1/4/a^(1/2)/(a+b)^(1/2)*\ln((a+b)^(1/2)*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))+1/4/a^(1/2)/(a+b)^(1/2)*\ln(-(a+b)^(1/2)*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^(1/2)-(a+b)^(1/2))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 608 vs. 2(45) = 90.

Time = 0.29 (sec) , antiderivative size = 1377, normalized size of antiderivative = 25.04

$$\int \frac{\operatorname{sech}^4(x)}{a+b\cosh^2(x)} dx = \text{Too large to display}$$

[In] integrate(sech(x)^4/(a+b*cosh(x)^2),x, algorithm="fricas")

[Out] $[1/6*(12*(a^2*b + a*b^2)*\cosh(x)^4 + 48*(a^2*b + a*b^2)*\cosh(x)*\sinh(x)^3 + 12*(a^2*b + a*b^2)*\sinh(x)^4 - 8*a^3 + 4*a^2*b + 12*a*b^2 - 24*(a^3 - a*b^2)*\cosh(x)^2 - 24*(a^3 - a*b^2 - 3*(a^2*b + a*b^2)*\cosh(x)^2)*\sinh(x)^2 + 3*(b^2*\cosh(x)^6 + 6*b^2*\cosh(x)*\sinh(x)^5 + b^2*\sinh(x)^6 + 3*b^2*\cosh(x)^4 + 3*(5*b^2*\cosh(x)^2 + b^2)*\sinh(x)^4 + 3*b^2*\cosh(x)^2 + 4*(5*b^2*\cosh(x)^3 + 3*b^2*\cosh(x))*\sinh(x)^3 + 3*(5*b^2*\cosh(x)^4 + 6*b^2*\cosh(x)^2 + b^2)*\sinh(x)^2 + b^2 + 6*(b^2*\cosh(x)^5 + 2*b^2*\cosh(x)^3 + b^2*\cosh(x))*\sinh(x))*\sqrt{a^2 + a*b}*\log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x)^2 + 2*(3*b^2*\cosh(x)^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) - 4*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{a^2 + a*b}))/((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + 48*((a^2*b + a*b^2)*\cosh(x)^3 - (a^3 - a*b^2)*\cosh(x))*\sinh(x)]/((a^4 + a^3*b)*\cosh(x)^6 + 6*(a^4 + a^3*b)*\cosh(x)*\sinh(x))$

$$\begin{aligned} &^5 + (a^4 + a^3b) \sinh(x)^6 + 3(a^4 + a^3b) \cosh(x)^4 + 3(a^4 + a^3b + \\ &5(a^4 + a^3b) \cosh(x)^2) \sinh(x)^4 + a^4 + a^3b + 4(5(a^4 + a^3b) \cosh(x)^3 + 3(a^4 + a^3b) \cosh(x)) \sinh(x)^3 + 3(a^4 + a^3b) \cosh(x)^2 + \\ &3(5(a^4 + a^3b) \cosh(x)^4 + a^4 + a^3b + 6(a^4 + a^3b) \cosh(x)^2) \sinh(x)^2 + 6((a^4 + a^3b) \cosh(x)^5 + 2(a^4 + a^3b) \cosh(x)^3 + (a^4 + a^3b) \cosh(x)) \sinh(x), \\ &1/3(6(a^2b + ab^2) \cosh(x)^4 + 24(a^2b + ab^2) \cosh(x) \sinh(x)^3 + 6(a^2b + ab^2) \sinh(x)^4 - 4a^3 + 2a^2b + 6ab^2 - 12(a^3 - ab^2) \cosh(x)^2 - 12(a^3 - ab^2 - 3(a^2b + ab^2) \cosh(x)^2) \sinh(x)^2 + 3(b^2 \cosh(x)^6 + 6b^2 \cosh(x) \sinh(x)^5 + b^2 \sinh(x)^6 + 3b^2 \cosh(x)^4 + 3(5b^2 \cosh(x)^2 + b^2) \sinh(x)^4 + 3b^2 \cosh(x)^2 + 4(5b^2 \cosh(x)^3 + 3b^2 \cosh(x)) \sinh(x)^3 + 3(5b^2 \cosh(x)^4 + 6b^2 \cosh(x)^2 + b^2) \sinh(x)^2 + b^2 + 6(b^2 \cosh(x)^5 + 2b^2 \cosh(x)^3 + b^2 \cosh(x)) \sinh(x)) \sqrt{-a^2 - ab} \arctan(1/2(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 + 2a + b) \sqrt{-a^2 - ab} / (a^2 + ab)) + 24((a^2b + ab^2) \cosh(x)^3 - (a^3 - ab^2) \cosh(x)) \sinh(x) / ((a^4 + a^3b) \cosh(x)^6 + 6(a^4 + a^3b) \cosh(x) \sinh(x)^5 + (a^4 + a^3b) \sinh(x)^6 + 3(a^4 + a^3b) \cosh(x)^4 + 3(a^4 + a^3b + 5(a^4 + a^3b) \cosh(x)^2) \sinh(x)^4 + a^4 + a^3b + 4(5(a^4 + a^3b) \cosh(x)^3 + 3(a^4 + a^3b) \cosh(x)) \sinh(x)^3 + 3(a^4 + a^3b) \cosh(x)^2 + 3(5(a^4 + a^3b) \cosh(x)^4 + a^4 + a^3b + 6(a^4 + a^3b) \cosh(x)^2) \sinh(x)^2 + 6((a^4 + a^3b) \cosh(x)^5 + 2(a^4 + a^3b) \cosh(x)^3 + (a^4 + a^3b) \cosh(x)) \sinh(x))] \end{aligned}$$

Sympy [F]

$$\int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx$$

[In] integrate(sech(x)**4/(a+b*cosh(x)**2),x)

[Out] Integral(sech(x)**4/(a + b*cosh(x)**2), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(45) = 90$.

Time = 0.31 (sec) , antiderivative size = 119, normalized size of antiderivative = 2.16

$$\begin{aligned} \int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx = & -\frac{b^2 \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{2\sqrt{(a+b)aa^2}} \\ & + \frac{2(6(a-b)e^{(-2x)} - 3be^{(-4x)} + 2a - 3b)}{3(3a^2e^{(-2x)} + 3a^2e^{(-4x)} + a^2e^{(-6x)} + a^2)} \end{aligned}$$

[In] integrate(sech(x)^4/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out]
$$-1/2*b^2*\log((b*e^{(-2*x)} + 2*a + b - 2*\sqrt{(a + b)*a}))/((b*e^{(-2*x)} + 2*a + b + 2*\sqrt{(a + b)*a}))/(\sqrt{(a + b)*a}*a^2) + 2/3*(6*(a - b)*e^{(-2*x)} - 3*b*e^{(-4*x)} + 2*a - 3*b)/(3*a^2*e^{(-2*x)} + 3*a^2*e^{(-4*x)} + a^2*e^{(-6*x)} + a^2)$$

Giac [F]

$$\int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)^4}{b \cosh(x)^2 + a} dx$$

[In] `integrate(sech(x)^4/(a+b*cosh(x)^2),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B] (verification not implemented)

Time = 2.26 (sec) , antiderivative size = 239, normalized size of antiderivative = 4.35

$$\int \frac{\operatorname{sech}^4(x)}{a + b \cosh^2(x)} dx = \frac{8}{3a(3e^{2x} + 3e^{4x} + e^{6x} + 1)} - \frac{4}{a(2e^{2x} + e^{4x} + 1)} + \frac{2b}{a^2(e^{2x} + 1)} - \frac{b^2 \ln\left(\frac{4b^2(2ab + 8a^2e^{2x} + b^2e^{2x} + b^2 + 8abe^{2x})}{a^5(a+b)} - \frac{8b^2(b + 4ae^{2x} + 2be^{2x})}{a^{9/2}\sqrt{a+b}}\right)}{2a^{5/2}\sqrt{a+b}} + \frac{b^2 \ln\left(\frac{8b^2(b + 4ae^{2x} + 2be^{2x})}{a^{9/2}\sqrt{a+b}} + \frac{4b^2(2ab + 8a^2e^{2x} + b^2e^{2x} + b^2 + 8abe^{2x})}{a^5(a+b)}\right)}{2a^{5/2}\sqrt{a+b}}$$

[In] `int(1/(cosh(x)^4*(a + b*cosh(x)^2)),x)`

[Out]
$$\frac{8}{3a*(3*\exp(2*x) + 3*\exp(4*x) + \exp(6*x) + 1)} - \frac{4}{a*(2*\exp(2*x) + \exp(4*x) + 1)} + \frac{2*b}{a^2*(\exp(2*x) + 1)} - \frac{(b^2*\log((4*b^2*(2*a*b + 8*a^2*\exp(2*x) + b^2*\exp(2*x) + b^2 + 8*a*b*\exp(2*x)))/(a^5*(a + b)) - (8*b^2*(b + 4*a*\exp(2*x) + 2*b*\exp(2*x)))/(a^{9/2}*(a + b)^{(1/2)})))/(2*a^{5/2}*(a + b)^{(1/2))} + (b^2*\log((8*b^2*(b + 4*a*\exp(2*x) + 2*b*\exp(2*x)))/(a^{9/2}*(a + b)^{(1/2))} + (4*b^2*(2*a*b + 8*a^2*\exp(2*x) + b^2*\exp(2*x) + b^2 + 8*a*b*\exp(2*x)))/(a^5*(a + b)))))/(2*a^{5/2}*(a + b)^{(1/2))}$$

3.32 $\int \frac{\operatorname{sech}^5(x)}{a+b \cosh^2(x)} dx$

Optimal result	259
Rubi [A] (verified)	259
Mathematica [A] (verified)	261
Maple [B] (verified)	262
Fricas [B] (verification not implemented)	262
Sympy [F]	264
Maxima [F]	265
Giac [F]	265
Mupad [B] (verification not implemented)	265

Optimal result

Integrand size = 15, antiderivative size = 90

$$\int \frac{\operatorname{sech}^5(x)}{a+b \cosh^2(x)} dx = \frac{(3a^2 - 4ab + 8b^2) \arctan(\sinh(x))}{8a^3} - \frac{b^{5/2} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a^3 \sqrt{a+b}} + \frac{(3a - 4b) \operatorname{sech}(x) \tanh(x)}{8a^2} + \frac{\operatorname{sech}^3(x) \tanh(x)}{4a}$$

[Out] $1/8*(3*a^2-4*a*b+8*b^2)*\arctan(\sinh(x))/a^3-b^{(5/2)*\arctan(\sinh(x)*b^{(1/2)/(a+b)^{(1/2)})/a^3/(a+b)^{(1/2)}+1/8*(3*a-4*b)*\operatorname{sech}(x)*\tanh(x)/a^2+1/4*\operatorname{sech}(x)^3*\tanh(x)/a$

Rubi [A] (verified)

Time = 0.11 (sec), antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3265, 425, 541, 536, 209, 211}

$$\int \frac{\operatorname{sech}^5(x)}{a+b \cosh^2(x)} dx = -\frac{b^{5/2} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a^3 \sqrt{a+b}} + \frac{(3a - 4b) \tanh(x) \operatorname{sech}(x)}{8a^2} + \frac{(3a^2 - 4ab + 8b^2) \arctan(\sinh(x))}{8a^3} + \frac{\tanh(x) \operatorname{sech}^3(x)}{4a}$$

[In] $\text{Int}[\text{Sech}[x]^5/(a + b*\text{Cosh}[x]^2), x]$

[Out] $((3*a^2 - 4*a*b + 8*b^2)*\text{ArcTan}[\text{Sinh}[x]])/(8*a^3) - (b^{(5/2)*\text{ArcTan}[(\text{Sqrt}[b]*\text{Sinh}[x])/(\text{Sqrt}[a + b])])/(a^3*\text{Sqrt}[a + b]) + ((3*a - 4*b)*\text{Sech}[x]*\text{Tanh}[x])/(8*a^2) + (\text{Sech}[x]^3*\text{Tanh}[x])/(4*a)$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 425

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] := Simp[(-b)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(p + 1)*(b*c - a*d))), x] + Dist[1/(a*n*(p + 1)*(b*c - a*d)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[b*c + n*(p + 1)*(b*c - a*d) + d*b*(n*(p + q + 2) + 1)*x^n, x], x] /; FreeQ[{a, b, c, d, n, q}, x] && NeQ[b*c - a*d, 0] && LtQ[p, -1] && !( !IntegerQ[p] && IntegerQ[q] && LtQ[q, -1]) && IntBinomialQ[a, b, c, d, n, p, q, x]
```

Rule 536

```
Int[((e_) + (f_.)*(x_)^(n_))/((a_) + (b_.)*(x_)^(n_))*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Dist[(b*e - a*f)/(b*c - a*d), Int[1/(a + b*x^n), x], x] - Dist[(d*e - c*f)/(b*c - a*d), Int[1/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x]
```

Rule 541

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_)*((e_) + (f_.)*(x_)^(n_)), x_Symbol] := Simp[(-b*e - a*f)*x*(a + b*x^n)^(p + 1)*((c + d*x^n)^(q + 1)/(a*n*(b*c - a*d)*(p + 1))), x] + Dist[1/(a*n*(b*c - a*d)*(p + 1)), Int[(a + b*x^n)^(p + 1)*(c + d*x^n)^q*Simp[c*(b*e - a*f) + e*n*(b*c - a*d)*(p + 1) + d*(b*e - a*f)*(n*(p + q + 2) + 1)*x^n, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, q}, x] && LtQ[p, -1]
```

Rule 3265

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.), x_Symbol] := With[{ff = FreeFactors[Cos[e + f*x], x]}, Dist[-ff/f, Subst[Int[(1 - ff^2*x^2)^((m - 1)/2)*(a + b - b*ff^2*x^2)^p, x], x, Cos[e + f*x]/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{(1+x^2)^3(a+b+bx^2)} dx, x, \sinh(x)\right) \\
 &= \frac{\text{sech}^3(x) \tanh(x)}{4a} - \frac{\text{Subst}\left(\int \frac{-3a+b-3bx^2}{(1+x^2)^2(a+b+bx^2)} dx, x, \sinh(x)\right)}{4a} \\
 &= \frac{(3a-4b)\text{sech}(x) \tanh(x)}{8a^2} + \frac{\text{sech}^3(x) \tanh(x)}{4a} + \frac{\text{Subst}\left(\int \frac{3a^2-ab+4b^2+(3a-4b)bx^2}{(1+x^2)(a+b+bx^2)} dx, x, \sinh(x)\right)}{8a^2} \\
 &= \frac{(3a-4b)\text{sech}(x) \tanh(x)}{8a^2} + \frac{\text{sech}^3(x) \tanh(x)}{4a} - \frac{b^3 \text{Subst}\left(\int \frac{1}{a+b+bx^2} dx, x, \sinh(x)\right)}{a^3} \\
 &\quad + \frac{(3a^2-4ab+8b^2) \text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \sinh(x)\right)}{8a^3} \\
 &= \frac{(3a^2-4ab+8b^2) \arctan(\sinh(x))}{8a^3} - \frac{b^{5/2} \arctan\left(\frac{\sqrt{b} \sinh(x)}{\sqrt{a+b}}\right)}{a^3 \sqrt{a+b}} \\
 &\quad + \frac{(3a-4b)\text{sech}(x) \tanh(x)}{8a^2} + \frac{\text{sech}^3(x) \tanh(x)}{4a}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.24 (sec) , antiderivative size = 86, normalized size of antiderivative = 0.96

$$\begin{aligned}
 &\int \frac{\text{sech}^5(x)}{a+b \cosh^2(x)} dx \\
 &= \frac{8b^{5/2} \arctan\left(\frac{\sqrt{a+b} \text{csch}(x)}{\sqrt{b}}\right)}{\sqrt{a+b}} + \frac{2(3a^2-4ab+8b^2) \arctan\left(\tanh\left(\frac{x}{2}\right)\right) + a(3a-4b)\text{sech}(x) \tanh(x) + 2a^2 \text{sech}^3(x)}{8a^3}
 \end{aligned}$$

[In] Integrate[Sech[x]^5/(a + b*Cosh[x]^2), x]

[Out] ((8*b^(5/2)*ArcTan[(Sqrt[a + b]*Csch[x])/Sqrt[b]])/Sqrt[a + b] + 2*(3*a^2 - 4*a*b + 8*b^2)*ArcTan[Tanh[x/2]] + a*(3*a - 4*b)*Sech[x]*Tanh[x] + 2*a^2*Sech[x]^3*Tanh[x])/(8*a^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 182 vs. $2(76) = 152$.

Time = 1.74 (sec) , antiderivative size = 183, normalized size of antiderivative = 2.03

method	result
default	$\frac{2\left(\left(-\frac{5}{8}a^2+\frac{1}{2}ab\right)\tanh\left(\frac{x}{2}\right)^7+\left(\frac{3}{8}a^2+\frac{1}{2}ab\right)\tanh\left(\frac{x}{2}\right)^5+\left(-\frac{3}{8}a^2-\frac{1}{2}ab\right)\tanh\left(\frac{x}{2}\right)^3+\left(\frac{5}{8}a^2-\frac{1}{2}ab\right)\tanh\left(\frac{x}{2}\right)\right)}{\left(1+\tanh\left(\frac{x}{2}\right)^2\right)^4}+\frac{\left(3a^2-4ab+8b^2\right)\arctan\left(\tanh\left(\frac{x}{2}\right)\right)}{4}$
risch	$\frac{e^x(3ae^{6x}-4be^{6x}+11ae^{4x}-4be^{4x}-11ae^{2x}+4be^{2x}-3a+4b)}{4(1+e^{2x})^4a^2}+\frac{3i\ln(e^x+i)}{8a}-\frac{ib\ln(e^x+i)}{2a^2}+\frac{i\ln(e^x+i)b^2}{a^3}-\frac{3i\ln(e^x-i)}{8a}+\frac{ib\ln(e^x-i)}{2a^2}$

[In] `int(sech(x)^5/(a+b*cosh(x)^2),x,method=_RETURNVERBOSE)`

[Out] $2/a^3\left(\left(-5/8*a^2+1/2*a*b\right)*\tanh(1/2*x)^7+\left(3/8*a^2+1/2*a*b\right)*\tanh(1/2*x)^5+\left(-3/8*a^2-1/2*a*b\right)*\tanh(1/2*x)^3+\left(5/8*a^2-1/2*a*b\right)*\tanh(1/2*x)\right)/\left(1+\tanh(1/2*x)^2\right)^4+1/8*\left(3*a^2-4*a*b+8*b^2\right)*\arctan\left(\tanh(1/2*x)\right)-2*b^3/a^3*\left(1/2/\left(a+b\right)^{\left(1/2\right)}/b^{\left(1/2\right)}*\arctan\left(1/2*\left(2*\tanh(1/2*x)*\left(a+b\right)^{\left(1/2\right)}+2*a^{\left(1/2\right)}\right)/b^{\left(1/2\right)}\right)+1/2/\left(a+b\right)^{\left(1/2\right)}/b^{\left(1/2\right)}*\arctan\left(1/2*\left(2*\tanh(1/2*x)*\left(a+b\right)^{\left(1/2\right)}-2*a^{\left(1/2\right)}\right)/b^{\left(1/2\right)}\right)\right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1673 vs. $2(76) = 152$.

Time = 0.30 (sec) , antiderivative size = 3239, normalized size of antiderivative = 35.99

$$\int \frac{\operatorname{sech}^5(x)}{a+b\cosh^2(x)} dx = \text{Too large to display}$$

[In] `integrate(sech(x)^5/(a+b*cosh(x)^2),x, algorithm="fricas")`

[Out] $[1/4*\left(\left(3*a^2-4*a*b\right)*\cosh(x)^7+7*\left(3*a^2-4*a*b\right)*\cosh(x)*\sinh(x)^6+\left(3*a^2-4*a*b\right)*\sinh(x)^7+\left(11*a^2-4*a*b\right)*\cosh(x)^5+\left(21*\left(3*a^2-4*a*b\right)*\cosh(x)^2+11*a^2-4*a*b\right)*\sinh(x)^5+5*\left(7*\left(3*a^2-4*a*b\right)*\cosh(x)^3+\left(11*a^2-4*a*b\right)*\cosh(x)*\sinh(x)^4-\left(11*a^2-4*a*b\right)*\cosh(x)^3+\left(35*\left(3*a^2-4*a*b\right)*\cosh(x)^4+10*\left(11*a^2-4*a*b\right)*\cosh(x)^2-11*a^2+4*a*b\right)*\sinh(x)^3+\left(21*\left(3*a^2-4*a*b\right)*\cosh(x)^5+10*\left(11*a^2-4*a*b\right)*\cosh(x)^3-3*\left(11*a^2-4*a*b\right)*\cosh(x)*\sinh(x)^2+2*\left(b^2*\cosh(x)^8+8*b^2*\cosh(x)*\sinh(x)^7+b^2*\sinh(x)^8+4*b^2*\cosh(x)^6+4*\left(7*b^2*\cosh(x)^2+b^2\right)*\sinh(x)^6+6*b^2*\cosh(x)^4+8*\left(7*b^2*\cosh(x)^3+3*b^2*\cosh(x)\right)*\sinh(x)^5+2*\left(35*b^2*\cosh(x)^4+30*b^2*\cosh(x)^2+3*b^2\right)*\sinh(x)^4+4*b^2*\cosh(x)^2+8*\left(7*b^2*\cosh(x)^5+10*b^2*\cosh(x)^3+3*b^2*\cosh(x)\right)*\sinh(x)^3+4*\left(7*b^2*\cosh(x)^6+15*b^2*\cosh(x)^4+9*b^2*\cosh(x)^2+b^2\right)*\sinh(x)^2+b^2+8*(b^2$

$$\begin{aligned}
& * \cosh(x)^7 + 3*b^2*\cosh(x)^5 + 3*b^2*\cosh(x)^3 + b^2*\cosh(x))*\sinh(x))*\sqrt{ \\
& (-b/(a + b))*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 - 2*(2* \\
& a + 3*b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 - 2*a - 3*b)*\sinh(x)^2 + 4*(b*\cosh(x) \\
& ^3 - (2*a + 3*b)*\cosh(x))*\sinh(x) - 4*((a + b)*\cosh(x)^3 + 3*(a + b)*\cosh(x) \\
&)*\sinh(x)^2 + (a + b)*\sinh(x)^3 - (a + b)*\cosh(x) + (3*(a + b)*\cosh(x)^2 - \\
& a - b)*\sinh(x))*\sqrt{-b/(a + b)) + b)/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 \\
& + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x) \\
& ^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)) + ((3*a^2 - 4*a*b + \\
& 8*b^2)*\cosh(x)^8 + 8*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)*\sinh(x)^7 + (3*a^2 - 4 \\
& *a*b + 8*b^2)*\sinh(x)^8 + 4*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^6 + 4*(7*(3*a^2 \\
& - 4*a*b + 8*b^2)*\cosh(x)^2 + 3*a^2 - 4*a*b + 8*b^2)*\sinh(x)^6 + 8*(7*(3*a^ \\
& 2 - 4*a*b + 8*b^2)*\cosh(x)^3 + 3*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x))*\sinh(x)^5 \\
& + 6*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^4 + 2*(35*(3*a^2 - 4*a*b + 8*b^2)*\cosh \\
& (x)^4 + 30*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^2 + 9*a^2 - 12*a*b + 24*b^2)*\sin \\
& h(x)^4 + 8*(7*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^5 + 10*(3*a^2 - 4*a*b + 8*b^2 \\
&)*\cosh(x)^3 + 3*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x))*\sinh(x)^3 + 4*(3*a^2 - 4*a \\
& *b + 8*b^2)*\cosh(x)^2 + 4*(7*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^6 + 15*(3*a^2 \\
& - 4*a*b + 8*b^2)*\cosh(x)^4 + 9*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^2 + 3*a^2 - \\
& 4*a*b + 8*b^2)*\sinh(x)^2 + 3*a^2 - 4*a*b + 8*b^2 + 8*((3*a^2 - 4*a*b + 8*b^ \\
& 2)*\cosh(x)^7 + 3*(3*a^2 - 4*a*b + 8*b^2)*\cosh(x)^5 + 3*(3*a^2 - 4*a*b + 8*b \\
& ^2)*\cosh(x)^3 + (3*a^2 - 4*a*b + 8*b^2)*\cosh(x))*\sinh(x))*\arctan(\cosh(x) + \\
& \sinh(x)) - (3*a^2 - 4*a*b)*\cosh(x) + (7*(3*a^2 - 4*a*b)*\cosh(x)^6 + 5*(11*a \\
& ^2 - 4*a*b)*\cosh(x)^4 - 3*(11*a^2 - 4*a*b)*\cosh(x)^2 - 3*a^2 + 4*a*b)*\sinh(\\
& x))/(a^3*\cosh(x)^8 + 8*a^3*\cosh(x)*\sinh(x)^7 + a^3*\sinh(x)^8 + 4*a^3*\cosh(x) \\
&)^6 + 6*a^3*\cosh(x)^4 + 4*(7*a^3*\cosh(x)^2 + a^3)*\sinh(x)^6 + 8*(7*a^3*\cosh \\
& (x)^3 + 3*a^3*\cosh(x))*\sinh(x)^5 + 4*a^3*\cosh(x)^2 + 2*(35*a^3*\cosh(x)^4 + \\
& 30*a^3*\cosh(x)^2 + 3*a^3)*\sinh(x)^4 + 8*(7*a^3*\cosh(x)^5 + 10*a^3*\cosh(x)^3 \\
& + 3*a^3*\cosh(x))*\sinh(x)^3 + a^3 + 4*(7*a^3*\cosh(x)^6 + 15*a^3*\cosh(x)^4 + \\
& 9*a^3*\cosh(x)^2 + a^3)*\sinh(x)^2 + 8*(a^3*\cosh(x)^7 + 3*a^3*\cosh(x)^5 + 3* \\
& a^3*\cosh(x)^3 + a^3*\cosh(x))*\sinh(x)), 1/4*((3*a^2 - 4*a*b)*\cosh(x)^7 + 7*(\\
& 3*a^2 - 4*a*b)*\cosh(x)*\sinh(x)^6 + (3*a^2 - 4*a*b)*\sinh(x)^7 + (11*a^2 - 4* \\
& a*b)*\cosh(x)^5 + (21*(3*a^2 - 4*a*b)*\cosh(x)^2 + 11*a^2 - 4*a*b)*\sinh(x)^5 \\
& + 5*(7*(3*a^2 - 4*a*b)*\cosh(x)^3 + (11*a^2 - 4*a*b)*\cosh(x))*\sinh(x)^4 - (1 \\
& 1*a^2 - 4*a*b)*\cosh(x)^3 + (35*(3*a^2 - 4*a*b)*\cosh(x)^4 + 10*(11*a^2 - 4*a \\
& *b)*\cosh(x)^2 - 11*a^2 + 4*a*b)*\sinh(x)^3 + (21*(3*a^2 - 4*a*b)*\cosh(x)^5 + \\
& 10*(11*a^2 - 4*a*b)*\cosh(x)^3 - 3*(11*a^2 - 4*a*b)*\cosh(x))*\sinh(x)^2 - 4* \\
& (b^2*\cosh(x)^8 + 8*b^2*\cosh(x)*\sinh(x)^7 + b^2*\sinh(x)^8 + 4*b^2*\cosh(x)^6 \\
& + 4*(7*b^2*\cosh(x)^2 + b^2)*\sinh(x)^6 + 6*b^2*\cosh(x)^4 + 8*(7*b^2*\cosh(x)^ \\
& 3 + 3*b^2*\cosh(x))*\sinh(x)^5 + 2*(35*b^2*\cosh(x)^4 + 30*b^2*\cosh(x)^2 + 3*b \\
& ^2)*\sinh(x)^4 + 4*b^2*\cosh(x)^2 + 8*(7*b^2*\cosh(x)^5 + 10*b^2*\cosh(x)^3 + 3 \\
& *b^2*\cosh(x))*\sinh(x)^3 + 4*(7*b^2*\cosh(x)^6 + 15*b^2*\cosh(x)^4 + 9*b^2*\cos \\
& h(x)^2 + b^2)*\sinh(x)^2 + b^2 + 8*(b^2*\cosh(x)^7 + 3*b^2*\cosh(x)^5 + 3*b^2* \\
& \cosh(x)^3 + b^2*\cosh(x))*\sinh(x))*\sqrt{b/(a + b))*\arctan(1/2*\sqrt{b/(a + b) \\
& }*(\cosh(x) + \sinh(x))) - 4*(b^2*\cosh(x)^8 + 8*b^2*\cosh(x)*\sinh(x)^7 + b^2*s \\
& inh(x)^8 + 4*b^2*\cosh(x)^6 + 4*(7*b^2*\cosh(x)^2 + b^2)*\sinh(x)^6 + 6*b^2*co
\end{aligned}$$

```

sh(x)^4 + 8*(7*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^5 + 2*(35*b^2*cosh(x)
^4 + 30*b^2*cosh(x)^2 + 3*b^2)*sinh(x)^4 + 4*b^2*cosh(x)^2 + 8*(7*b^2*cosh(
x)^5 + 10*b^2*cosh(x)^3 + 3*b^2*cosh(x))*sinh(x)^3 + 4*(7*b^2*cosh(x)^6 + 1
5*b^2*cosh(x)^4 + 9*b^2*cosh(x)^2 + b^2)*sinh(x)^2 + b^2 + 8*(b^2*cosh(x)^7
+ 3*b^2*cosh(x)^5 + 3*b^2*cosh(x)^3 + b^2*cosh(x))*sinh(x))*sqrt(b/(a + b)
)*arctan(1/2*(b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + b*sinh(x)^3 + (4*a + 3*
b)*cosh(x) + (3*b*cosh(x)^2 + 4*a + 3*b)*sinh(x))*sqrt(b/(a + b)))/b + ((3*
a^2 - 4*a*b + 8*b^2)*cosh(x)^8 + 8*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)*sinh(x)^
7 + (3*a^2 - 4*a*b + 8*b^2)*sinh(x)^8 + 4*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^6
+ 4*(7*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^2 + 3*a^2 - 4*a*b + 8*b^2)*sinh(x)^
6 + 8*(7*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^3 + 3*(3*a^2 - 4*a*b + 8*b^2)*cosh
(x))*sinh(x)^5 + 6*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^4 + 2*(35*(3*a^2 - 4*a*b
+ 8*b^2)*cosh(x)^4 + 30*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^2 + 9*a^2 - 12*a*b
+ 24*b^2)*sinh(x)^4 + 8*(7*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^5 + 10*(3*a^2 -
4*a*b + 8*b^2)*cosh(x)^3 + 3*(3*a^2 - 4*a*b + 8*b^2)*cosh(x))*sinh(x)^3 +
4*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^2 + 4*(7*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^
6 + 15*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^4 + 9*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)
)^2 + 3*a^2 - 4*a*b + 8*b^2)*sinh(x)^2 + 3*a^2 - 4*a*b + 8*b^2 + 8*((3*a^2
- 4*a*b + 8*b^2)*cosh(x)^7 + 3*(3*a^2 - 4*a*b + 8*b^2)*cosh(x)^5 + 3*(3*a^2
- 4*a*b + 8*b^2)*cosh(x)^3 + (3*a^2 - 4*a*b + 8*b^2)*cosh(x))*sinh(x))*arc
tan(cosh(x) + sinh(x)) - (3*a^2 - 4*a*b)*cosh(x) + (7*(3*a^2 - 4*a*b)*cosh(
x)^6 + 5*(11*a^2 - 4*a*b)*cosh(x)^4 - 3*(11*a^2 - 4*a*b)*cosh(x)^2 - 3*a^2
+ 4*a*b)*sinh(x))/(a^3*cosh(x)^8 + 8*a^3*cosh(x)*sinh(x)^7 + a^3*sinh(x)^8
+ 4*a^3*cosh(x)^6 + 6*a^3*cosh(x)^4 + 4*(7*a^3*cosh(x)^2 + a^3)*sinh(x)^6 +
8*(7*a^3*cosh(x)^3 + 3*a^3*cosh(x))*sinh(x)^5 + 4*a^3*cosh(x)^2 + 2*(35*a^
3*cosh(x)^4 + 30*a^3*cosh(x)^2 + 3*a^3)*sinh(x)^4 + 8*(7*a^3*cosh(x)^5 + 10
*a^3*cosh(x)^3 + 3*a^3*cosh(x))*sinh(x)^3 + a^3 + 4*(7*a^3*cosh(x)^6 + 15*a
^3*cosh(x)^4 + 9*a^3*cosh(x)^2 + a^3)*sinh(x)^2 + 8*(a^3*cosh(x)^7 + 3*a^3*
cosh(x)^5 + 3*a^3*cosh(x)^3 + a^3*cosh(x))*sinh(x))]

```

Sympy [F]

$$\int \frac{\operatorname{sech}^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}^5(x)}{a + b \cosh^2(x)} dx$$

```
[In] integrate(sech(x)**5/(a+b*cosh(x)**2), x)
```

```
[Out] Integral(sech(x)**5/(a + b*cosh(x)**2), x)
```


Maxima [F]

$$\int \frac{\operatorname{sech}^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(sech(x)^5/(a+b*cosh(x)^2),x, algorithm="maxima")

[Out] $\frac{1}{4} * ((3*a - 4*b) * e^{7*x} + (11*a - 4*b) * e^{5*x} - (11*a - 4*b) * e^{3*x} - (3*a - 4*b) * e^x) / (a^2 * e^{8*x} + 4*a^2 * e^{6*x} + 6*a^2 * e^{4*x} + 4*a^2 * e^{2*x} + a^2) + \frac{1}{4} * (3*a^2 - 4*a*b + 8*b^2) * \arctan(e^x) / a^3 - 32 * \operatorname{integrate}(1/16 * (b^3 * e^{3*x} + b^3 * e^x) / (a^3 * b * e^{4*x} + a^3 * b + 2 * (2*a^4 + a^3 * b) * e^{2*x}), x)$

Giac [F]

$$\int \frac{\operatorname{sech}^5(x)}{a + b \cosh^2(x)} dx = \int \frac{\operatorname{sech}(x)^5}{b \cosh(x)^2 + a} dx$$

[In] integrate(sech(x)^5/(a+b*cosh(x)^2),x, algorithm="giac")

[Out] sage0*x

Mupad [B] (verification not implemented)

Time = 43.63 (sec) , antiderivative size = 1305, normalized size of antiderivative = 14.50

$$\int \frac{\operatorname{sech}^5(x)}{a + b \cosh^2(x)} dx = \text{Too large to display}$$

[In] int(1/(cosh(x)^5*(a + b*cosh(x)^2)),x)

[Out] $\frac{\operatorname{atan}\left(\frac{\exp(x) * (243*a^{12} * (a^6)^{1/2} + 5024*b^6 * (a^6)^{3/2} + 18432*b^{12} * (a^6)^{1/2} + 6912*a^2 * b^{10} * (a^6)^{1/2} + 30720*a^3 * b^9 * (a^6)^{1/2} - 26880*a^4 * b^8 * (a^6)^{1/2} + 24192*a^5 * b^7 * (a^6)^{1/2} - 13408*a^7 * b^5 * (a^6)^{1/2} + 17160*a^8 * b^4 * (a^6)^{1/2} - 9540*a^9 * b^3 * (a^6)^{1/2} + 4563*a^{10} * b^2 * (a^6)^{1/2} - 9216*a * b^{11} * (a^6)^{1/2} - 1134*a^{11} * b * (a^6)^{1/2})}{(81*a^{13} * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} - 270*a^{12} * b * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 2304*a^3 * b^{10} * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 3840*a^6 * b^7 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} - 1440*a^7 * b^6 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 864*a^8 * b^5 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 1600*a^9 * b^4 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2}}\right)}{81*a^{13} * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} - 270*a^{12} * b * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 2304*a^3 * b^{10} * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 3840*a^6 * b^7 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} - 1440*a^7 * b^6 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 864*a^8 * b^5 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2} + 1600*a^9 * b^4 * (9*a^4 - 24*a^3 * b - 64*a * b^3 + 64*b^4 + 64*a^2 * b^2)^{1/2}}$

$$\begin{aligned}
& 64*a*b^3 + 64*b^4 + 64*a^2*b^2)^{(1/2)} - 1200*a^{10}*b^3*(9*a^4 - 24*a^3*b - \\
& 64*a*b^3 + 64*b^4 + 64*a^2*b^2)^{(1/2)} + 945*a^{11}*b^2*(9*a^4 - 24*a^3*b - 64 \\
& *a*b^3 + 64*b^4 + 64*a^2*b^2)^{(1/2)))*(9*a^4 - 24*a^3*b - 64*a*b^3 + 64*b^4 \\
& + 64*a^2*b^2)^{(1/2))/(4*(a^6)^{(1/2)}) - (6*\exp(x))/(a*(3*\exp(2*x) + 3*\exp(4 \\
& *x) + \exp(6*x) + 1)) + ((b^5)^{(1/2)}*(2*\operatorname{atan}((\exp(x))*((2*(48*b^8*(a^6*b + a^ \\
& 7)^{(1/2)} + 40*a^3*b^5*(a^6*b + a^7)^{(1/2)} - 15*a^4*b^4*(a^6*b + a^7)^{(1/2)} \\
& + 9*a^5*b^3*(a^6*b + a^7)^{(1/2)))/(a^{11}*b*(a + b)*(a*b + a^2)*(a^6*b + a^7) \\
& ^{(1/2)}*(b^5)^{(1/2)}*(48*a*b^5 - 6*a^5*b + 9*a^6 + 48*b^6 + 40*a^3*b^3 + 25*a^4 \\
& ^4*b^2)) - (4*(96*a^4*(b^5)^{(3/2)} + 18*a^9*(b^5)^{(1/2)} + 80*a^6*b^3*(b^5)^{(\\
& 1/2)} + 50*a^7*b^2*(b^5)^{(1/2)} + 96*a^3*b*(b^5)^{(3/2)} - 12*a^8*b*(b^5)^{(1/2)} \\
&))/(a^8*b^4*(a + b)*(a*b + a^2)*(a^6*(a + b))^{(1/2)}*(a^6*b + a^7)^{(1/2)}*(9* \\
& a^5 - 15*a^4*b + 48*b^5 + 40*a^3*b^2))) - (2*\exp(3*x))*(48*b^8*(a^6*b + a^7) \\
& ^{(1/2)} + 40*a^3*b^5*(a^6*b + a^7)^{(1/2)} - 15*a^4*b^4*(a^6*b + a^7)^{(1/2)} + \\
& 9*a^5*b^3*(a^6*b + a^7)^{(1/2)))/(a^{11}*b*(a + b)*(a*b + a^2)*(a^6*b + a^7)^{(\\
& 1/2)}*(b^5)^{(1/2)}*(48*a*b^5 - 6*a^5*b + 9*a^6 + 48*b^6 + 40*a^3*b^3 + 25*a^4 \\
& *b^2)))*((a^{11}*b*(a^6*b + a^7)^{(1/2)))/4 + (a^9*b^3*(a^6*b + a^7)^{(1/2)))/4 + \\
& (a^{10}*b^2*(a^6*b + a^7)^{(1/2)))/2) - 2*\operatorname{atan}((b^3*\exp(x)*(a^6*(a + b))^{(1/2)} \\
&)*(9*a^5 - 15*a^4*b + 48*b^5 + 40*a^3*b^2))/(2*a^3*(b^5)^{(1/2)}*(48*a*b^5 - \\
& 6*a^5*b + 9*a^6 + 48*b^6 + 40*a^3*b^3 + 25*a^4*b^2))))/(2*(a^6*b + a^7)^{(1 \\
& /2)}) + (4*\exp(x))/(a*(4*\exp(2*x) + 6*\exp(4*x) + 4*\exp(6*x) + \exp(8*x) + 1)) \\
& + (\exp(x)*(a + 4*b))/(2*a^2*(2*\exp(2*x) + \exp(4*x) + 1)) - (\exp(x)*(4*a*b \\
& - 3*a^2))/(4*a^3*(\exp(2*x) + 1))
\end{aligned}$$

3.33 $\int \frac{1}{(a+b \cosh^2(x))^2} dx$

Optimal result	267
Rubi [A] (verified)	267
Mathematica [A] (verified)	268
Maple [B] (verified)	269
Fricas [B] (verification not implemented)	269
Sympy [F(-1)]	270
Maxima [B] (verification not implemented)	270
Giac [A] (verification not implemented)	271
Mupad [F(-1)]	271

Optimal result

Integrand size = 10, antiderivative size = 65

$$\int \frac{1}{(a+b \cosh^2(x))^2} dx = \frac{(2a+b) \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{2a^{3/2}(a+b)^{3/2}} - \frac{b \cosh(x) \sinh(x)}{2a(a+b)(a+b \cosh^2(x))}$$

[Out] 1/2*(2*a+b)*arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))/a^(3/2)/(a+b)^(3/2)-1/2*b*cosh(x)*sinh(x)/a/(a+b)/(a+b*cosh(x)^2)

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3263, 12, 3260, 214}

$$\int \frac{1}{(a+b \cosh^2(x))^2} dx = \frac{(2a+b) \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{2a^{3/2}(a+b)^{3/2}} - \frac{b \sinh(x) \cosh(x)}{2a(a+b)(a+b \cosh^2(x))}$$

[In] Int[(a + b*Cosh[x]^2)^(-2), x]

[Out] ((2*a + b)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(2*a^(3/2)*(a + b)^(3/2)) - (b*Cosh[x]*Sinh[x])/(2*a*(a + b)*(a + b*Cosh[x]^2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 3260

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 3263

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x]^2)^(p + 1)/(2*a*f*(p + 1)*(a + b))), x] + Dist[1/(2*a*(p + 1)*(a + b)), Int[(a + b*Sin[e + f*x]^2)^(p + 1)*Simp[2*a*(p + 1) + b*(2*p + 3) - 2*b*(p + 2)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a + b, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{b \cosh(x) \sinh(x)}{2a(a+b)(a+b \cosh^2(x))} - \frac{\int \frac{-2a-b}{a+b \cosh^2(x)} dx}{2a(a+b)} \\
 &= -\frac{b \cosh(x) \sinh(x)}{2a(a+b)(a+b \cosh^2(x))} + \frac{(2a+b) \int \frac{1}{a+b \cosh^2(x)} dx}{2a(a+b)} \\
 &= -\frac{b \cosh(x) \sinh(x)}{2a(a+b)(a+b \cosh^2(x))} + \frac{(2a+b) \text{Subst}\left(\int \frac{1}{a-(a+b)x^2} dx, x, \coth(x)\right)}{2a(a+b)} \\
 &= \frac{(2a+b) \arctanh\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{2a^{3/2}(a+b)^{3/2}} - \frac{b \cosh(x) \sinh(x)}{2a(a+b)(a+b \cosh^2(x))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 5.08 (sec) , antiderivative size = 68, normalized size of antiderivative = 1.05

$$\int \frac{1}{(a+b \cosh^2(x))^2} dx = \frac{(2a+b) \arctanh\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{2a^{3/2}(a+b)^{3/2}} - \frac{b \sinh(2x)}{2a(a+b)(2a+b+b \cosh(2x))}$$

```
[In] Integrate[(a + b*Cosh[x]^2)^(-2), x]
```

```
[Out] ((2*a + b)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]])/(2*a^(3/2)*(a + b)^(3/2)) - (b*Sinh[2*x])/(2*a*(a + b)*(2*a + b + b*Cosh[2*x]))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. 2(53) = 106.

Time = 0.36 (sec) , antiderivative size = 171, normalized size of antiderivative = 2.63

method	result
default	$-\frac{2\left(\frac{b \tanh\left(\frac{x}{2}\right)^3}{2a(a+b)} + \frac{b \tanh\left(\frac{x}{2}\right)}{2a(a+b)}\right)}{\tanh\left(\frac{x}{2}\right)^4 a + \tanh\left(\frac{x}{2}\right)^4 b - 2 \tanh\left(\frac{x}{2}\right)^2 a + 2 \tanh\left(\frac{x}{2}\right)^2 b + a + b} - \frac{(2a+b) \left(-\frac{\ln\left(\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 + 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b}\right)}{4\sqrt{a+b}} + \frac{\ln\left(-\sqrt{a+b} \tanh\left(\frac{x}{2}\right)^2 - 2 \tanh\left(\frac{x}{2}\right) \sqrt{a+b}\right)}{4\sqrt{a+b}} \right)}{a(a+b)}$
risch	$\frac{2a e^{2x} + b e^{2x} + b}{a(a+b)(b e^{4x} + 4a e^{2x} + 2b e^{2x} + b)} + \frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab+b\sqrt{a^2+ab}-2a^2-2ab}}{b\sqrt{a^2+ab}}\right)}{2\sqrt{a^2+ab}(a+b)} + \frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+ab+b\sqrt{a^2+ab}-2a^2-2ab}}{b\sqrt{a^2+ab}}\right)b}{4\sqrt{a^2+ab}(a+b)a} - \ln\left(\frac{e^{2x} + \frac{2a\sqrt{a^2+ab+b\sqrt{a^2+ab}-2a^2-2ab}}{b\sqrt{a^2+ab}}}{e^{2x} + \frac{2a\sqrt{a^2+ab+b\sqrt{a^2+ab}-2a^2-2ab}}{b\sqrt{a^2+ab}}}\right)$

[In] int(1/(a+b*cosh(x)^2)^2,x,method=_RETURNVERBOSE)

[Out] $-2*(1/2*b/a/(a+b)*\tanh(1/2*x)^3+1/2*b/a/(a+b)*\tanh(1/2*x))/(\tanh(1/2*x)^4*a + \tanh(1/2*x)^4*b - 2*\tanh(1/2*x)^2*a + 2*\tanh(1/2*x)^2*b + a + b) - (2*a + b)/a/(a+b)*(-1/4/a^(1/2)/(a+b)^(1/2)*\ln((a+b)^(1/2)*\tanh(1/2*x)^2 + 2*\tanh(1/2*x)*a^(1/2) + (a+b)^(1/2)) + 1/4/a^(1/2)/(a+b)^(1/2)*\ln(-(a+b)^(1/2)*\tanh(1/2*x)^2 + 2*\tanh(1/2*x)*a^(1/2) - (a+b)^(1/2))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 539 vs. 2(53) = 106.

Time = 0.27 (sec) , antiderivative size = 1239, normalized size of antiderivative = 19.06

$$\int \frac{1}{(a + b \cosh^2(x))^2} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cosh(x)^2)^2,x, algorithm="fricas")

[Out] $[1/4*(4*a^2*b + 4*a*b^2 + 4*(2*a^3 + 3*a^2*b + a*b^2)*\cosh(x)^2 + 8*(2*a^3 + 3*a^2*b + a*b^2)*\cosh(x)*\sinh(x) + 4*(2*a^3 + 3*a^2*b + a*b^2)*\sinh(x)^2 + ((2*a*b + b^2)*\cosh(x)^4 + 4*(2*a*b + b^2)*\cosh(x)*\sinh(x)^3 + (2*a*b + b^2)*\sinh(x)^4 + 2*(4*a^2 + 4*a*b + b^2)*\cosh(x)^2 + 2*(3*(2*a*b + b^2)*\cosh(x)^2 + 4*a^2 + 4*a*b + b^2)*\sinh(x)^2 + 2*a*b + b^2 + 4*((2*a*b + b^2)*\cosh(x)^3 + (4*a^2 + 4*a*b + b^2)*\cosh(x))*\sinh(x))*\sqrt{a^2 + a*b}*\log((b^2*\cosh(x)^4 + 4*b^2*\cosh(x)*\sinh(x)^3 + b^2*\sinh(x)^4 + 2*(2*a*b + b^2)*\cosh(x))^2 + 2*(3*b^2*\cosh(x)^2 + 2*a*b + b^2)*\sinh(x)^2 + 8*a^2 + 8*a*b + b^2 + 4*(b^2*\cosh(x)^3 + (2*a*b + b^2)*\cosh(x))*\sinh(x) - 4*(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*a + b)*\sqrt{a^2 + a*b})/(b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(2*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 2*a + b)*\sinh(x)^2 + 4*(b*\cosh(x)^3 + (2*a + b)*\cosh(x))*\sinh(x) + b)]/(a^4*b + 2*a^3*b^2 + a^2*b^3 + (a^4*b + 2*a^3*b^2 + a^2*b^3)*\cosh(x)^4 + 4*(a^4*b + 2*a^3*b^2 + a^2*b^3)*\cosh(x)*\sinh(x)^3 + (a^4*b + 2*a^3*b^2 + a^2*b^3)$

```

3)*sinh(x)^4 + 2*(2*a^5 + 5*a^4*b + 4*a^3*b^2 + a^2*b^3)*cosh(x)^2 + 2*(2*a
^5 + 5*a^4*b + 4*a^3*b^2 + a^2*b^3 + 3*(a^4*b + 2*a^3*b^2 + a^2*b^3)*cosh(x
)^2)*sinh(x)^2 + 4*((a^4*b + 2*a^3*b^2 + a^2*b^3)*cosh(x)^3 + (2*a^5 + 5*a^
4*b + 4*a^3*b^2 + a^2*b^3)*cosh(x))*sinh(x)), 1/2*(2*a^2*b + 2*a*b^2 + 2*(2
*a^3 + 3*a^2*b + a*b^2)*cosh(x)^2 + 4*(2*a^3 + 3*a^2*b + a*b^2)*cosh(x)*sin
h(x) + 2*(2*a^3 + 3*a^2*b + a*b^2)*sinh(x)^2 + ((2*a*b + b^2)*cosh(x)^4 + 4
*(2*a*b + b^2)*cosh(x)*sinh(x)^3 + (2*a*b + b^2)*sinh(x)^4 + 2*(4*a^2 + 4*a
*b + b^2)*cosh(x)^2 + 2*(3*(2*a*b + b^2)*cosh(x)^2 + 4*a^2 + 4*a*b + b^2)*s
inh(x)^2 + 2*a*b + b^2 + 4*((2*a*b + b^2)*cosh(x)^3 + (4*a^2 + 4*a*b + b^2)
*cosh(x))*sinh(x))*sqrt(-a^2 - a*b)*arctan(1/2*(b*cosh(x)^2 + 2*b*cosh(x)*s
inh(x) + b*sinh(x)^2 + 2*a + b)*sqrt(-a^2 - a*b)/(a^2 + a*b)))/(a^4*b + 2*a
^3*b^2 + a^2*b^3 + (a^4*b + 2*a^3*b^2 + a^2*b^3)*cosh(x)^4 + 4*(a^4*b + 2*a
^3*b^2 + a^2*b^3)*cosh(x)*sinh(x)^3 + (a^4*b + 2*a^3*b^2 + a^2*b^3)*sinh(x)
^4 + 2*(2*a^5 + 5*a^4*b + 4*a^3*b^2 + a^2*b^3)*cosh(x)^2 + 2*(2*a^5 + 5*a^4
*b + 4*a^3*b^2 + a^2*b^3 + 3*(a^4*b + 2*a^3*b^2 + a^2*b^3)*cosh(x)^2)*sinh(
x)^2 + 4*((a^4*b + 2*a^3*b^2 + a^2*b^3)*cosh(x)^3 + (2*a^5 + 5*a^4*b + 4*a^
3*b^2 + a^2*b^3)*cosh(x))*sinh(x))]

```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cosh^2(x))^2} dx = \text{Timed out}$$

```
[In] integrate(1/(a+b*cosh(x)**2)**2,x)
```

```
[Out] Timed out
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(53) = 106.

Time = 0.29 (sec) , antiderivative size = 134, normalized size of antiderivative = 2.06

$$\int \frac{1}{(a + b \cosh^2(x))^2} dx = -\frac{(2a + b) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{4\sqrt{(a+b)a}(a^2 + ab)} - \frac{(2a + b)e^{(-2x)} + b}{a^2b + ab^2 + 2(2a^3 + 3a^2b + ab^2)e^{(-2x)} + (a^2b + ab^2)e^{(-4x)}}$$

```
[In] integrate(1/(a+b*cosh(x)^2)^2,x, algorithm="maxima")
```

```
[Out] -1/4*(2*a + b)*log((b*e^(-2*x) + 2*a + b - 2*sqrt((a + b)*a))/(b*e^(-2*x) +
2*a + b + 2*sqrt((a + b)*a)))/(sqrt((a + b)*a)*(a^2 + a*b)) - ((2*a + b)*e
^(-2*x) + b)/(a^2*b + a*b^2 + 2*(2*a^3 + 3*a^2*b + a*b^2)*e^(-2*x) + (a^2*b
+ a*b^2)*e^(-4*x))
```

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 104, normalized size of antiderivative = 1.60

$$\int \frac{1}{(a + b \cosh^2(x))^2} dx = \frac{(2a + b) \arctan\left(\frac{be^{(2x)} + 2a + b}{2\sqrt{-a^2 - ab}}\right)}{2(a^2 + ab)\sqrt{-a^2 - ab}} + \frac{2ae^{(2x)} + be^{(2x)} + b}{(a^2 + ab)(be^{(4x)} + 4ae^{(2x)} + 2be^{(2x)} + b)}$$

[In] integrate(1/(a+b*cosh(x)^2)^2,x, algorithm="giac")

[Out] 1/2*(2*a + b)*arctan(1/2*(b*e^(2*x) + 2*a + b)/sqrt(-a^2 - a*b))/((a^2 + a*b)*sqrt(-a^2 - a*b)) + (2*a*e^(2*x) + b*e^(2*x) + b)/((a^2 + a*b)*(b*e^(4*x) + 4*a*e^(2*x) + 2*b*e^(2*x) + b))

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cosh^2(x))^2} dx = \int \frac{1}{(b \cosh(x)^2 + a)^2} dx$$

[In] int(1/(a + b*cosh(x)^2)^2,x)

[Out] int(1/(a + b*cosh(x)^2)^2, x)

$$3.34 \quad \int \frac{1}{(a+b \cosh^2(x))^3} dx$$

Optimal result	272
Rubi [A] (verified)	272
Mathematica [A] (verified)	274
Maple [B] (verified)	274
Fricas [B] (verification not implemented)	275
Sympy [F(-1)]	275
Maxima [B] (verification not implemented)	276
Giac [F]	276
Mupad [F(-1)]	276

Optimal result

Integrand size = 10, antiderivative size = 107

$$\int \frac{1}{(a+b \cosh^2(x))^3} dx = \frac{(8a^2 + 8ab + 3b^2) \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{8a^{5/2}(a+b)^{5/2}} - \frac{b \cosh(x) \sinh(x)}{4a(a+b)(a+b \cosh^2(x))^2} - \frac{3b(2a+b) \cosh(x) \sinh(x)}{8a^2(a+b)^2(a+b \cosh^2(x))}$$

[Out] 1/8*(8*a^2+8*a*b+3*b^2)*arctanh(a^(1/2)*tanh(x)/(a+b)^(1/2))/a^(5/2)/(a+b)^(5/2)-1/4*b*cosh(x)*sinh(x)/a/(a+b)/(a+b*cosh(x)^2)^2-3/8*b*(2*a+b)*cosh(x)*sinh(x)/a^2/(a+b)^2/(a+b*cosh(x)^2)

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3263, 3252, 12, 3260, 214}

$$\int \frac{1}{(a+b \cosh^2(x))^3} dx = -\frac{3b(2a+b) \sinh(x) \cosh(x)}{8a^2(a+b)^2(a+b \cosh^2(x))} + \frac{(8a^2 + 8ab + 3b^2) \operatorname{arctanh}\left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a+b}}\right)}{8a^{5/2}(a+b)^{5/2}} - \frac{b \sinh(x) \cosh(x)}{4a(a+b)(a+b \cosh^2(x))^2}$$

[In] Int[(a + b*Cosh[x]^2)^(-3), x]


```
[Out] ((8*a^2 + 8*a*b + 3*b^2)*ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + b]]/(8*a^(5/2)
*(a + b)^(5/2)) - (b*Cosh[x]*Sinh[x])/(4*a*(a + b)*(a + b*Cosh[x]^2)^2) - (
3*b*(2*a + b)*Cosh[x]*Sinh[x])/(8*a^2*(a + b)^2*(a + b*Cosh[x]^2))
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 3252

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(-(A*b - a*B))*Cos[e + f*x]*Sin[e + f*x
]*(a + b*Sine[e + f*x]^2)^(p + 1)/(2*a*f*(a + b)*(p + 1)), x] - Dist[1/(2*
a*(a + b)*(p + 1)), Int[(a + b*Sine[e + f*x]^2)^(p + 1)*Simp[a*B - A*(2*a*(p
+ 1) + b*(2*p + 3)) + 2*(A*b - a*B)*(p + 2)*Sine[e + f*x]^2, x], x], x] /;
FreeQ[{a, b, e, f, A, B}, x] && LtQ[p, -1] && NeQ[a + b, 0]
```

Rule 3260

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2
), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Rule 3263

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Simp[(-b)*C
os[e + f*x]*Sin[e + f*x]*((a + b*Sine[e + f*x]^2)^(p + 1)/(2*a*f*(p + 1)*(a
+ b))), x] + Dist[1/(2*a*(p + 1)*(a + b)), Int[(a + b*Sine[e + f*x]^2)^(p +
1)*Simp[2*a*(p + 1) + b*(2*p + 3) - 2*b*(p + 2)*Sine[e + f*x]^2, x], x], x]
/; FreeQ[{a, b, e, f}, x] && NeQ[a + b, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{b \cosh(x) \sinh(x)}{4a(a+b)(a+b \cosh^2(x))^2} - \frac{\int \frac{-4a-3b+2b \cosh^2(x)}{(a+b \cosh^2(x))^2} dx}{4a(a+b)} \\ &= -\frac{b \cosh(x) \sinh(x)}{4a(a+b)(a+b \cosh^2(x))^2} - \frac{3b(2a+b) \cosh(x) \sinh(x)}{8a^2(a+b)^2(a+b \cosh^2(x))} - \frac{\int \frac{-8a^2-8ab-3b^2}{a+b \cosh^2(x)} dx}{8a^2(a+b)^2} \end{aligned}$$

method	result
default	$- \frac{2 \left(\frac{b(8a+3b) \tanh\left(\frac{x}{2}\right)^7}{8a^2(a+b)} - \frac{b(8a^2-13ab-9b^2) \tanh\left(\frac{x}{2}\right)^5}{8(a+b)^2 a^2} - \frac{b(8a^2-13ab-9b^2) \tanh\left(\frac{x}{2}\right)^3}{8(a+b)^2 a^2} + \frac{b(8a+3b) \tanh\left(\frac{x}{2}\right)}{8a^2(a+b)} \right)}{\left(\tanh\left(\frac{x}{2}\right)^4 a + \tanh\left(\frac{x}{2}\right)^4 b - 2 \tanh\left(\frac{x}{2}\right)^2 a + 2 \tanh\left(\frac{x}{2}\right)^2 b + a + b \right)^2} - \frac{(8a^2+8ab+3b^2) \left(-\frac{\ln\left(\frac{e^{2x} + \frac{2a\sqrt{a^2+b^2}}{2\sqrt{a^2+b^2}}}{e^{2x} + \frac{2a\sqrt{a^2+b^2}}{2\sqrt{a^2+b^2}}}\right)}{2\sqrt{a^2+b^2}} \right)}{(8a^2+8ab+3b^2)}$
risch	$\frac{8e^{6x}a^2b + 8e^{6x}ab^2 + 3b^3e^{6x} + 48a^3e^{4x} + 72a^2be^{4x} + 42ab^2e^{4x} + 9b^3e^{4x} + 40e^{2x}a^2b + 40e^{2x}ab^2 + 9b^3e^{2x} + 6ab^2 + 3b^3}{4a^2(a+b)^2(b e^{4x} + 4a e^{2x} + 2b e^{2x} + b)^2} + \frac{\ln\left(e^{2x} + \frac{2a\sqrt{a^2+b^2}}{2\sqrt{a^2+b^2}}\right)}{2\sqrt{a^2+b^2}}$

[In] `int(1/(a+b*cosh(x)^2)^3,x,method=_RETURNVERBOSE)`

[Out] $-2*(1/8*b*(8*a+3*b)/a^2/(a+b)*\tanh(1/2*x)^7-1/8*b*(8*a^2-13*a*b-9*b^2)/(a+b)^2/a^2*\tanh(1/2*x)^5-1/8*b*(8*a^2-13*a*b-9*b^2)/(a+b)^2/a^2*\tanh(1/2*x)^3+1/8*b*(8*a+3*b)/a^2/(a+b)*\tanh(1/2*x))/(\tanh(1/2*x)^4*a+\tanh(1/2*x)^4*b-2*\tanh(1/2*x)^2*a+2*\tanh(1/2*x)^2*b+a+b)^2-1/4*(8*a^2+8*a*b+3*b^2)/a^2/(a^2+2*a*b+b^2)*(-1/4/a^(1/2)/(a+b)^(1/2)*\ln((a+b)^(1/2)*\tanh(1/2*x)^2+2*\tanh(1/2*x)*a^(1/2)+(a+b)^(1/2))+1/4/a^(1/2)/(a+b)^(1/2)*\ln((a+b)^(1/2)*\tanh(1/2*x)^2-2*\tanh(1/2*x)*a^(1/2)+(a+b)^(1/2)))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2478 vs. 2(93) = 186.

Time = 0.33 (sec) , antiderivative size = 5117, normalized size of antiderivative = 47.82

$$\int \frac{1}{(a + b \cosh^2(x))^3} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*cosh(x)^2)^3,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cosh^2(x))^3} dx = \text{Timed out}$$

[In] `integrate(1/(a+b*cosh(x)**2)**3,x)`

[Out] Timed out

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 344 vs. $2(93) = 186$.

Time = 0.30 (sec) , antiderivative size = 344, normalized size of antiderivative = 3.21

$$\int \frac{1}{(a + b \cosh^2(x))^3} dx = -\frac{(8a^2 + 8ab + 3b^2) \log\left(\frac{be^{(-2x)} + 2a + b - 2\sqrt{(a+b)a}}{be^{(-2x)} + 2a + b + 2\sqrt{(a+b)a}}\right)}{16(a^4 + 2a^3b + a^2b^2)\sqrt{(a+b)a}} - \frac{6ab^2 + 3b^3 + (40a^2b + 40ab^2 + 9b^3)e^{(-2x)} + 3(16a^3 + 24a^2b + 14a^3b^3 + 4(a^4b^2 + 2a^3b^3 + a^2b^4) + 4(2a^5b + 5a^4b^2 + 4a^3b^3 + a^2b^4)e^{(-2x)} + 2(8a^6 + 24a^5b + 27a^4b^2 + 14a^3b^3 +$$

[In] integrate(1/(a+b*cosh(x)^2)^3,x, algorithm="maxima")

[Out] $-1/16*(8*a^2 + 8*a*b + 3*b^2)*\log((b*e^{(-2*x)} + 2*a + b - 2*\sqrt{(a + b)*a})/(b*e^{(-2*x)} + 2*a + b + 2*\sqrt{(a + b)*a}))/((a^4 + 2*a^3*b + a^2*b^2)*\sqrt{(a + b)*a}) - 1/4*(6*a*b^2 + 3*b^3 + (40*a^2*b + 40*a*b^2 + 9*b^3)*e^{(-2*x)} + 3*(16*a^3 + 24*a^2*b + 14*a*b^2 + 3*b^3)*e^{(-4*x)} + (8*a^2*b + 8*a*b^2 + 3*b^3)*e^{(-6*x)})/(a^4*b^2 + 2*a^3*b^3 + a^2*b^4 + 4*(2*a^5*b + 5*a^4*b^2 + 4*a^3*b^3 + a^2*b^4)*e^{(-2*x)} + 2*(8*a^6 + 24*a^5*b + 27*a^4*b^2 + 14*a^3*b^3 + 3*a^2*b^4)*e^{(-4*x)} + 4*(2*a^5*b + 5*a^4*b^2 + 4*a^3*b^3 + a^2*b^4)*e^{(-6*x)} + (a^4*b^2 + 2*a^3*b^3 + a^2*b^4)*e^{(-8*x)})$

Giac [F]

$$\int \frac{1}{(a + b \cosh^2(x))^3} dx = \int \frac{1}{(b \cosh(x)^2 + a)^3} dx$$

[In] integrate(1/(a+b*cosh(x)^2)^3,x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(a + b \cosh^2(x))^3} dx = \int \frac{1}{(b \cosh(x)^2 + a)^3} dx$$

[In] int(1/(a + b*cosh(x)^2)^3,x)

[Out] int(1/(a + b*cosh(x)^2)^3, x)

3.35 $\int \frac{1}{1+\cosh^2(x)} dx$

Optimal result	277
Rubi [A] (verified)	277
Mathematica [A] (verified)	278
Maple [B] (verified)	278
Fricas [B] (verification not implemented)	279
Sympy [B] (verification not implemented)	279
Maxima [B] (verification not implemented)	279
Giac [B] (verification not implemented)	280
Mupad [B] (verification not implemented)	280

Optimal result

Integrand size = 8, antiderivative size = 15

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[Out] 1/2*arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3260, 212}

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[In] Int[(1 + Cosh[x]^2)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[2]]/Sqrt[2]

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2

), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1}{1-2x^2} dx, x, \coth(x)\right) \\ &= \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{\sqrt{2}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.06 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{\sqrt{2}}$$

[In] Integrate[(1 + Cosh[x]^2)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[2]]/Sqrt[2]

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(13) = 26.

Time = 0.08 (sec) , antiderivative size = 36, normalized size of antiderivative = 2.40

method	result
risch	$\frac{\sqrt{2} \ln(e^{2x} + 3 - 2\sqrt{2})}{4} - \frac{\sqrt{2} \ln(e^{2x} + 3 + 2\sqrt{2})}{4}$
default	$\frac{\sqrt{2} \left(\ln\left(\frac{\tanh\left(\frac{x}{2}\right)^2 + \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}{\tanh\left(\frac{x}{2}\right)^2 - \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}\right) + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\sqrt{2} + 1\right) + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\sqrt{2} - 1\right) \right)}{8} - \frac{\sqrt{2} \left(\ln\left(\frac{\tanh\left(\frac{x}{2}\right)^2 - \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}{\tanh\left(\frac{x}{2}\right)^2 + \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}\right) \right)}{8}$

[In] int(1/(1+cosh(x)^2), x, method=_RETURNVERBOSE)

[Out] 1/4*2^(1/2)*ln(exp(2*x)+3-2*2^(1/2))-1/4*2^(1/2)*ln(exp(2*x)+3+2*2^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(13) = 26$.

Time = 0.26 (sec) , antiderivative size = 66, normalized size of antiderivative = 4.40

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{3(2\sqrt{2} - 3) \cosh(x)^2 - 4(3\sqrt{2} - 4) \cosh(x) \sinh(x) + 3(2\sqrt{2} - 3) \sinh(x)^2 + 2\sqrt{2} - 3}{\cosh(x)^2 + \sinh(x)^2 + 3} \right)$$

[In] integrate(1/(1+cosh(x)^2),x, algorithm="fricas")

[Out] 1/4*sqrt(2)*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(17) = 34$.

Time = 0.28 (sec) , antiderivative size = 60, normalized size of antiderivative = 4.00

$$\int \frac{1}{1 + \cosh^2(x)} dx = -\frac{\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{4} + \frac{\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{4}$$

[In] integrate(1/(1+cosh(x)**2),x)

[Out] -sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)/4 + sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)/4

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. $2(13) = 26$.

Time = 0.28 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.27

$$\int \frac{1}{1 + \cosh^2(x)} dx = -\frac{1}{4} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3} \right)$$

[In] integrate(1/(1+cosh(x)^2),x, algorithm="maxima")

[Out] -1/4*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3))

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(13) = 26.

Time = 0.26 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.27

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{1}{4} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right)$$

[In] integrate(1/(1+cosh(x)^2),x, algorithm="giac")

[Out] 1/4*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3))

Mupad [B] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 50, normalized size of antiderivative = 3.33

$$\int \frac{1}{1 + \cosh^2(x)} dx = \frac{\sqrt{2} \left(\ln \left(-4e^{2x} - \frac{\sqrt{2}(12e^{2x}+4)}{4} \right) - \ln \left(\frac{\sqrt{2}(12e^{2x}+4)}{4} - 4e^{2x} \right) \right)}{4}$$

[In] int(1/(cosh(x)^2 + 1),x)

[Out] (2^(1/2)*(log(- 4*exp(2*x) - (2^(1/2)*(12*exp(2*x) + 4))/4) - log((2^(1/2)*(12*exp(2*x) + 4))/4 - 4*exp(2*x))))/4

$$3.36 \quad \int \frac{1}{(1+\cosh^2(x))^2} dx$$

Optimal result	281
Rubi [A] (verified)	281
Mathematica [A] (verified)	282
Maple [B] (verified)	283
Fricas [B] (verification not implemented)	283
Sympy [B] (verification not implemented)	284
Maxima [B] (verification not implemented)	284
Giac [B] (verification not implemented)	285
Mupad [B] (verification not implemented)	285

Optimal result

Integrand size = 8, antiderivative size = 35

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = \frac{3\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{\cosh(x) \sinh(x)}{4(1 + \cosh^2(x))}$$

[Out] $-1/4*\cosh(x)*\sinh(x)/(1+\cosh(x)^2)+3/8*\operatorname{arctanh}(1/2*2^{(1/2)}*\tanh(x))*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3263, 12, 3260, 212}

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = \frac{3\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{\sinh(x) \cosh(x)}{4(\cosh^2(x) + 1)}$$

[In] $\operatorname{Int}[(1 + \operatorname{Cosh}[x]^2)^{-2}, x]$

[Out] $(3*\operatorname{ArcTanh}[\operatorname{Tanh}[x]/\operatorname{Sqrt}[2]])/(4*\operatorname{Sqrt}[2]) - (\operatorname{Cosh}[x]*\operatorname{Sinh}[x])/(4*(1 + \operatorname{Cosh}[x]^2))$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /;$ FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 3260

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2
), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 3263

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[(-b)*C
os[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x]^2)^(p + 1)/(2*a*f*(p + 1)*(a
+ b))), x] + Dist[1/(2*a*(p + 1)*(a + b)), Int[(a + b*Sin[e + f*x]^2)^(p +
1)*Simp[2*a*(p + 1) + b*(2*p + 3) - 2*b*(p + 2)*Sin[e + f*x]^2, x], x], x]
/; FreeQ[{a, b, e, f}, x] && NeQ[a + b, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\cosh(x) \sinh(x)}{4(1 + \cosh^2(x))} - \frac{1}{4} \int -\frac{3}{1 + \cosh^2(x)} dx \\
&= -\frac{\cosh(x) \sinh(x)}{4(1 + \cosh^2(x))} + \frac{3}{4} \int \frac{1}{1 + \cosh^2(x)} dx \\
&= -\frac{\cosh(x) \sinh(x)}{4(1 + \cosh^2(x))} + \frac{3}{4} \text{Subst}\left(\int \frac{1}{1 - 2x^2} dx, x, \coth(x)\right) \\
&= \frac{3 \arctanh\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{\cosh(x) \sinh(x)}{4(1 + \cosh^2(x))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 35, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = \frac{3 \arctanh\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} - \frac{\sinh(2x)}{4(3 + \cosh(2x))}$$

```
[In] Integrate[(1 + Cosh[x]^2)^(-2), x]
```

```
[Out] (3*ArcTanh[Tanh[x]/Sqrt[2]])/(4*Sqrt[2]) - Sinh[2*x]/(4*(3 + Cosh[2*x]))
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.21 (sec) , antiderivative size = 60, normalized size of antiderivative = 1.71

method	result
risch	$\frac{3e^{2x}+1}{2e^{4x}+12e^{2x}+2} + \frac{3\sqrt{2}\ln(e^{2x}+3-2\sqrt{2})}{16} - \frac{3\sqrt{2}\ln(e^{2x}+3+2\sqrt{2})}{16}$
default	$-\frac{\frac{\tanh(\frac{x}{2})^3}{2} + \frac{\tanh(\frac{x}{2})}{2}}{2(\tanh(\frac{x}{2})^4+1)} + \frac{3\sqrt{2}\left(\ln\left(\frac{\tanh(\frac{x}{2})^2+\tanh(\frac{x}{2})\sqrt{2}+1}{\tanh(\frac{x}{2})^2-\tanh(\frac{x}{2})\sqrt{2}+1}\right)+2\arctan\left(\tanh(\frac{x}{2})\sqrt{2}+1\right)+2\arctan\left(\tanh(\frac{x}{2})\sqrt{2}-1\right)\right)}{32} - \frac{3\sqrt{2}}{32}$

[In] `int(1/(1+cosh(x)^2)^2,x,method=_RETURNVERBOSE)`

[Out] `1/2*(3*exp(2*x)+1)/(exp(4*x)+6*exp(2*x)+1)+3/16*2^(1/2)*ln(exp(2*x)+3-2*2^(1/2))-3/16*2^(1/2)*ln(exp(2*x)+3+2*2^(1/2))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 214 vs. $2(28) = 56$.

Time = 0.26 (sec) , antiderivative size = 214, normalized size of antiderivative = 6.11

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx$$

$$= \frac{24 \cosh(x)^2 + 3(\sqrt{2} \cosh(x)^4 + 4\sqrt{2} \cosh(x) \sinh(x)^3 + \sqrt{2} \sinh(x)^4 + 6(\sqrt{2} \cosh(x)^2 + \sqrt{2}) \sinh(x)^2)}{16(\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 6(\cosh(x)^2 + 1) \sinh(x)^2 + 6 \cosh(x)^2 + 4(\cosh(x)^3 + 3 \cosh(x)) \sinh(x) + 1)}$$

[In] `integrate(1/(1+cosh(x)^2)^2,x, algorithm="fricas")`

[Out] `1/16*(24*cosh(x)^2 + 3*(sqrt(2)*cosh(x)^4 + 4*sqrt(2)*cosh(x)*sinh(x)^3 + sqrt(2)*sinh(x)^4 + 6*(sqrt(2)*cosh(x)^2 + sqrt(2))*sinh(x)^2 + 6*sqrt(2)*cosh(x)^2 + 4*(sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3)) + 48*cosh(x)*sinh(x) + 24*sinh(x)^2 + 8)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 6*(cosh(x)^2 + 1)*sinh(x)^2 + 6*cosh(x)^2 + 4*(cosh(x)^3 + 3*cosh(x))*sinh(x) + 1)`

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 211 vs. 2(34) = 68.

Time = 1.16 (sec) , antiderivative size = 211, normalized size of antiderivative = 6.03

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = -\frac{3\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^4(\frac{x}{2})}{16 \tanh^4(\frac{x}{2}) + 16}$$

$$- \frac{3\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{16 \tanh^4(\frac{x}{2}) + 16}$$

$$+ \frac{3\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^4(\frac{x}{2})}{16 \tanh^4(\frac{x}{2}) + 16}$$

$$+ \frac{3\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{16 \tanh^4(\frac{x}{2}) + 16}$$

$$- \frac{4 \tanh^3(\frac{x}{2})}{16 \tanh^4(\frac{x}{2}) + 16} - \frac{4 \tanh(\frac{x}{2})}{16 \tanh^4(\frac{x}{2}) + 16}$$

[In] integrate(1/(1+cosh(x)**2)**2,x)

[Out] -3*sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**4/(16*tanh(x/2)**4 + 16) - 3*sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)/(16*tanh(x/2)**4 + 16) + 3*sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**4/(16*tanh(x/2)**4 + 16) + 3*sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)/(16*tanh(x/2)**4 + 16) - 4*tanh(x/2)**3/(16*tanh(x/2)**4 + 16) - 4*tanh(x/2)/(16*tanh(x/2)**4 + 16)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(28) = 56.

Time = 0.28 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = -\frac{3}{16} \sqrt{2} \log\left(\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3}\right) - \frac{3e^{(-2x)} + 1}{2(6e^{(-2x)} + e^{(-4x)} + 1)}$$

[In] integrate(1/(1+cosh(x)^2)^2,x, algorithm="maxima")

[Out] -3/16*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - 1/2*(3*e^(-2*x) + 1)/(6*e^(-2*x) + e^(-4*x) + 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(28) = 56$.

Time = 0.27 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.69

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = \frac{3}{16} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) + \frac{3e^{(2x)} + 1}{2(e^{(4x)} + 6e^{(2x)} + 1)}$$

[In] integrate(1/(1+cosh(x)^2)^2,x, algorithm="giac")

[Out] 3/16*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3)) + 1/2*(3*e^(2*x) + 1)/(e^(4*x) + 6*e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 1.89 (sec) , antiderivative size = 76, normalized size of antiderivative = 2.17

$$\int \frac{1}{(1 + \cosh^2(x))^2} dx = \frac{3\sqrt{2} \ln \left(-3e^{2x} - \frac{3\sqrt{2}(12e^{2x}+4)}{16} \right)}{16} - \frac{3\sqrt{2} \ln \left(\frac{3\sqrt{2}(12e^{2x}+4)}{16} - 3e^{2x} \right)}{16} + \frac{\frac{3e^{2x}}{2} + \frac{1}{2}}{6e^{2x} + e^{4x} + 1}$$

[In] int(1/(cosh(x)^2 + 1)^2,x)

[Out] (3*2^(1/2)*log(- 3*exp(2*x) - (3*2^(1/2)*(12*exp(2*x) + 4))/16))/16 - (3*2^(1/2)*log((3*2^(1/2)*(12*exp(2*x) + 4))/16 - 3*exp(2*x)))/16 + ((3*exp(2*x))/2 + 1/2)/(6*exp(2*x) + exp(4*x) + 1)

3.37 $\int \frac{1}{(1+\cosh^2(x))^3} dx$

Optimal result	286
Rubi [A] (verified)	286
Mathematica [A] (verified)	288
Maple [A] (verified)	288
Fricas [B] (verification not implemented)	288
Sympy [B] (verification not implemented)	289
Maxima [A] (verification not implemented)	291
Giac [A] (verification not implemented)	291
Mupad [B] (verification not implemented)	291

Optimal result

Integrand size = 8, antiderivative size = 51

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \frac{19 \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{32\sqrt{2}} - \frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{9 \cosh(x) \sinh(x)}{32(1 + \cosh^2(x))}$$

[Out] $-1/8*\cosh(x)*\sinh(x)/(1+\cosh(x)^2)^2-9/32*\cosh(x)*\sinh(x)/(1+\cosh(x)^2)+19/64*\operatorname{arctanh}(1/2*2^{(1/2)}*\tanh(x))*2^{(1/2)}$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3263, 3252, 12, 3260, 212}

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \frac{19 \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{32\sqrt{2}} - \frac{9 \sinh(x) \cosh(x)}{32(\cosh^2(x) + 1)} - \frac{\sinh(x) \cosh(x)}{8(\cosh^2(x) + 1)^2}$$

[In] $\operatorname{Int}[(1 + \operatorname{Cosh}[x]^2)^{-3}, x]$

[Out] $(19*\operatorname{ArcTanh}[\operatorname{Tanh}[x]/\operatorname{Sqrt}[2]])/(32*\operatorname{Sqrt}[2]) - (\operatorname{Cosh}[x]*\operatorname{Sinh}[x])/(8*(1 + \operatorname{Cosh}[x]^2)^2) - (9*\operatorname{Cosh}[x]*\operatorname{Sinh}[x])/(32*(1 + \operatorname{Cosh}[x]^2))$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /;$ $\operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match} Q[u, (b_*)(v_)] /;$ $\operatorname{FreeQ}[b, x]$

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 3252

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_)*((A_) + (B_)*sin[(e_)
+ (f_)*(x_)]^2), x_Symbol] := Simp[(-A*b - a*B)*Cos[e + f*x]*Sin[e + f*x
]*((a + b*Sin[e + f*x]^2)^(p + 1)/(2*a*f*(a + b)*(p + 1))), x] - Dist[1/(2*
a*(a + b)*(p + 1)), Int[(a + b*Sin[e + f*x]^2)^(p + 1)*Simp[a*B - A*(2*a*(p
+ 1) + b*(2*p + 3)) + 2*(A*b - a*B)*(p + 2)*Sin[e + f*x]^2, x], x], x] /;
FreeQ[{a, b, e, f, A, B}, x] && LtQ[p, -1] && NeQ[a + b, 0]
```

Rule 3260

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2
), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Rule 3263

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Simp[(-b)*C
os[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x]^2)^(p + 1)/(2*a*f*(p + 1)*(a
+ b))), x] + Dist[1/(2*a*(p + 1)*(a + b)), Int[(a + b*Sin[e + f*x]^2)^(p +
1)*Simp[2*a*(p + 1) + b*(2*p + 3) - 2*b*(p + 2)*Sin[e + f*x]^2, x], x], x]
/; FreeQ[{a, b, e, f}, x] && NeQ[a + b, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{1}{8} \int \frac{-7 + 2 \cosh^2(x)}{(1 + \cosh^2(x))^2} dx \\
&= -\frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{9 \cosh(x) \sinh(x)}{32(1 + \cosh^2(x))} - \frac{1}{32} \int -\frac{19}{1 + \cosh^2(x)} dx \\
&= -\frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{9 \cosh(x) \sinh(x)}{32(1 + \cosh^2(x))} + \frac{19}{32} \int \frac{1}{1 + \cosh^2(x)} dx \\
&= -\frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{9 \cosh(x) \sinh(x)}{32(1 + \cosh^2(x))} + \frac{19}{32} \text{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \coth(x) \right) \\
&= \frac{19 \arctanh\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{32\sqrt{2}} - \frac{\cosh(x) \sinh(x)}{8(1 + \cosh^2(x))^2} - \frac{9 \cosh(x) \sinh(x)}{32(1 + \cosh^2(x))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 51, normalized size of antiderivative = 1.00

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \frac{19 \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{32\sqrt{2}} - \frac{\sinh(2x)}{4(3 + \cosh(2x))^2} - \frac{9 \sinh(2x)}{32(3 + \cosh(2x))}$$

[In] Integrate[(1 + Cosh[x]^2)^(-3), x]

[Out] (19*ArcTanh[Tanh[x]/Sqrt[2]])/(32*Sqrt[2]) - Sinh[2*x]/(4*(3 + Cosh[2*x])^2) - (9*Sinh[2*x])/(32*(3 + Cosh[2*x]))

Maple [A] (verified)

Time = 0.46 (sec) , antiderivative size = 72, normalized size of antiderivative = 1.41

method	result
risch	$\frac{19e^{6x} + 171e^{4x} + 89e^{2x} + 9}{16(e^{4x} + 6e^{2x} + 1)^2} + \frac{19\sqrt{2} \ln(e^{2x} + 3 - 2\sqrt{2})}{128} - \frac{19\sqrt{2} \ln(e^{2x} + 3 + 2\sqrt{2})}{128}$
default	$-\frac{11 \tanh\left(\frac{x}{2}\right)^7}{8} + \frac{7 \tanh\left(\frac{x}{2}\right)^5}{8} + \frac{7 \tanh\left(\frac{x}{2}\right)^3}{8} + \frac{11 \tanh\left(\frac{x}{2}\right)}{8} + \frac{19\sqrt{2} \left(\ln\left(\frac{\tanh\left(\frac{x}{2}\right)^2 + \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}{\tanh\left(\frac{x}{2}\right)^2 - \tanh\left(\frac{x}{2}\right)\sqrt{2} + 1}\right) + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\sqrt{2} + 1\right) + 2 \arctan\left(\tanh\left(\frac{x}{2}\right)\sqrt{2} - 1\right)}{256}$

[In] int(1/(1+cosh(x)^2)^3,x,method=_RETURNVERBOSE)

[Out] 1/16*(19*exp(6*x)+171*exp(4*x)+89*exp(2*x)+9)/(exp(4*x)+6*exp(2*x)+1)^2+19/128*2^(1/2)*ln(exp(2*x)+3-2*2^(1/2))-19/128*2^(1/2)*ln(exp(2*x)+3+2*2^(1/2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 575 vs. 2(42) = 84.

Time = 0.26 (sec) , antiderivative size = 575, normalized size of antiderivative = 11.27

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \text{Too large to display}$$

[In] integrate(1/(1+cosh(x)^2)^3,x, algorithm="fricas")

[Out] 1/128*(152*cosh(x)^6 + 912*cosh(x)*sinh(x)^5 + 152*sinh(x)^6 + 456*(5*cosh(x)^2 + 3)*sinh(x)^4 + 1368*cosh(x)^4 + 608*(5*cosh(x)^3 + 9*cosh(x))*sinh(x)^3 + 8*(285*cosh(x)^4 + 1026*cosh(x)^2 + 89)*sinh(x)^2 + 712*cosh(x)^2 + 19*(sqrt(2)*cosh(x)^8 + 8*sqrt(2)*cosh(x)*sinh(x)^7 + sqrt(2)*sinh(x)^8 + 4*


```

(7*sqrt(2)*cosh(x)^2 + 3*sqrt(2))*sinh(x)^6 + 12*sqrt(2)*cosh(x)^6 + 8*(7*sqrt(2)*cosh(x)^3 + 9*sqrt(2)*cosh(x))*sinh(x)^5 + 2*(35*sqrt(2)*cosh(x)^4 + 90*sqrt(2)*cosh(x)^2 + 19*sqrt(2))*sinh(x)^4 + 38*sqrt(2)*cosh(x)^4 + 8*(7*sqrt(2)*cosh(x)^5 + 30*sqrt(2)*cosh(x)^3 + 19*sqrt(2)*cosh(x))*sinh(x)^3 + 4*(7*sqrt(2)*cosh(x)^6 + 45*sqrt(2)*cosh(x)^4 + 57*sqrt(2)*cosh(x)^2 + 3*sqrt(2))*sinh(x)^2 + 12*sqrt(2)*cosh(x)^2 + 8*(sqrt(2)*cosh(x)^7 + 9*sqrt(2)*cosh(x)^5 + 19*sqrt(2)*cosh(x)^3 + 3*sqrt(2)*cosh(x))*sinh(x) + sqrt(2))*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3)) + 16*(57*cosh(x)^5 + 342*cosh(x)^3 + 89*cosh(x))*sinh(x) + 72)/(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + 4*(7*cosh(x)^2 + 3)*sinh(x)^6 + 12*cosh(x)^6 + 8*(7*cosh(x)^3 + 9*cosh(x))*sinh(x)^5 + 2*(35*cosh(x)^4 + 90*cosh(x)^2 + 19)*sinh(x)^4 + 38*cosh(x)^4 + 8*(7*cosh(x)^5 + 30*cosh(x)^3 + 19*cosh(x))*sinh(x)^3 + 4*(7*cosh(x)^6 + 45*cosh(x)^4 + 57*cosh(x)^2 + 3)*sinh(x)^2 + 12*cosh(x)^2 + 8*(cosh(x)^7 + 9*cosh(x)^5 + 19*cosh(x)^3 + 3*cosh(x))*sinh(x) + 1)

```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 428 vs. $2(53) = 106$.

Time = 4.10 (sec) , antiderivative size = 428, normalized size of antiderivative = 8.39

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = -\frac{19\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^8(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{38\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^4(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{19\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) - 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$+ \frac{19\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^8(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$+ \frac{38\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4) \tanh^4(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$+ \frac{19\sqrt{2} \log(4 \tanh^2(\frac{x}{2}) + 4\sqrt{2} \tanh(\frac{x}{2}) + 4)}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{44 \tanh^7(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{28 \tanh^5(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{28 \tanh^3(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

$$- \frac{44 \tanh(\frac{x}{2})}{128 \tanh^8(\frac{x}{2}) + 256 \tanh^4(\frac{x}{2}) + 128}$$

[In] integrate(1/(1+cosh(x)**2)**3,x)

[Out] -19*sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**8/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 38*sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**4/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 19*sqrt(2)*log(4*tanh(x/2)**2 - 4*sqrt(2)*tanh(x/2) + 4)/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) + 19*sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**8/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) + 38*sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)*tanh(x/2)**4/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) + 19*sqrt(2)*log(4*tanh(x/2)**2 + 4*sqrt(2)*tanh(x/2) + 4)/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 44*tanh(x/2)**7/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 28*tanh(x/2)**5/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 28*tanh(x/2)**3/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128) - 44*tanh(x/2)/(128*tanh(x/2)**8 + 256*tanh(x/2)**4 + 128)

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 83, normalized size of antiderivative = 1.63

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = -\frac{19}{128} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3} \right) - \frac{89e^{(-2x)} + 171e^{(-4x)} + 19e^{(-6x)} + 9}{16(12e^{(-2x)} + 38e^{(-4x)} + 12e^{(-6x)} + e^{(-8x)} + 1)}$$

[In] integrate(1/(1+cosh(x)^2)^3,x, algorithm="maxima")

[Out] -19/128*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - 1/16*(89*e^(-2*x) + 171*e^(-4*x) + 19*e^(-6*x) + 9)/(12*e^(-2*x) + 38*e^(-4*x) + 12*e^(-6*x) + e^(-8*x) + 1)

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 71, normalized size of antiderivative = 1.39

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \frac{19}{128} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) + \frac{19e^{(6x)} + 171e^{(4x)} + 89e^{(2x)} + 9}{16(e^{(4x)} + 6e^{(2x)} + 1)^2}$$

[In] integrate(1/(1+cosh(x)^2)^3,x, algorithm="giac")

[Out] 19/128*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3)) + 1/16*(19*e^(6*x) + 171*e^(4*x) + 89*e^(2*x) + 9)/(e^(4*x) + 6*e^(2*x) + 1)^2

Mupad [B] (verification not implemented)

Time = 1.84 (sec) , antiderivative size = 112, normalized size of antiderivative = 2.20

$$\int \frac{1}{(1 + \cosh^2(x))^3} dx = \frac{19\sqrt{2} \ln \left(-\frac{19e^{2x}}{8} - \frac{19\sqrt{2}(12e^{2x}+4)}{128} \right)}{128} - \frac{17e^{2x} + 3}{12e^{2x} + 38e^{4x} + 12e^{6x} + e^{8x} + 1} - \frac{19\sqrt{2} \ln \left(\frac{19\sqrt{2}(12e^{2x}+4)}{128} - \frac{19e^{2x}}{8} \right)}{128} + \frac{\frac{19e^{2x}}{16} + \frac{57}{16}}{6e^{2x} + e^{4x} + 1}$$

[In] int(1/(cosh(x)^2 + 1)^3,x)

[Out] $(19 \cdot 2^{1/2} \cdot \log(- (19 \cdot \exp(2x))/8 - (19 \cdot 2^{1/2} \cdot (12 \cdot \exp(2x) + 4))/128))/12$
 $8 - (17 \cdot \exp(2x) + 3)/(12 \cdot \exp(2x) + 38 \cdot \exp(4x) + 12 \cdot \exp(6x) + \exp(8x) +$
 $1) - (19 \cdot 2^{1/2} \cdot \log((19 \cdot 2^{1/2} \cdot (12 \cdot \exp(2x) + 4))/128 - (19 \cdot \exp(2x))/8)$
 $)/128 + ((19 \cdot \exp(2x))/16 + 57/16)/(6 \cdot \exp(2x) + \exp(4x) + 1)$

3.38 $\int \frac{1}{1-\cosh^2(x)} dx$

Optimal result	293
Rubi [A] (verified)	293
Mathematica [A] (verified)	294
Maple [A] (verified)	294
Fricas [B] (verification not implemented)	295
Sympy [B] (verification not implemented)	295
Maxima [B] (verification not implemented)	295
Giac [B] (verification not implemented)	296
Mupad [B] (verification not implemented)	296

Optimal result

Integrand size = 10, antiderivative size = 2

$$\int \frac{1}{1-\cosh^2(x)} dx = \coth(x)$$

[Out] $\coth(x)$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3254, 3852, 8}

$$\int \frac{1}{1-\cosh^2(x)} dx = \coth(x)$$

[In] `Int[(1 - Cosh[x]^2)^(-1),x]`

[Out] `Coth[x]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3254

`Int[(u_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Dist[a^p, Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]`

Rule 3852

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= - \int \operatorname{csch}^2(x) dx \\ &= i \operatorname{Subst}\left(\int 1 dx, x, -i \operatorname{coth}(x)\right) \\ &= \operatorname{coth}(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 2, normalized size of antiderivative = 1.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = \operatorname{coth}(x)$$

```
[In] Integrate[(1 - Cosh[x]^2)^(-1), x]
```

```
[Out] Coth[x]
```

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 3, normalized size of antiderivative = 1.50

method	result	size
parallelrisch	$\operatorname{coth}(x)$	3
risch	$\frac{2}{e^{2x}-1}$	11
default	$\frac{\tanh(\frac{x}{2})}{2} + \frac{1}{2 \tanh(\frac{x}{2})}$	16

```
[In] int(1/(1-cosh(x)^2), x, method=_RETURNVERBOSE)
```

```
[Out] coth(x)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(2) = 4$.

Time = 0.25 (sec) , antiderivative size = 20, normalized size of antiderivative = 10.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = \frac{2}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - 1}$$

[In] integrate(1/(1-cosh(x)^2),x, algorithm="fricas")

[Out] 2/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - 1)

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 14 vs. $2(2) = 4$.

Time = 0.21 (sec) , antiderivative size = 14, normalized size of antiderivative = 7.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = \frac{\tanh\left(\frac{x}{2}\right)}{2} + \frac{1}{2 \tanh\left(\frac{x}{2}\right)}$$

[In] integrate(1/(1-cosh(x)**2),x)

[Out] tanh(x/2)/2 + 1/(2*tanh(x/2))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.19 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = -\frac{2}{e^{(-2x)} - 1}$$

[In] integrate(1/(1-cosh(x)^2),x, algorithm="maxima")

[Out] -2/(e^(-2*x) - 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 10 vs. $2(2) = 4$.

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = \frac{2}{e^{(2x)} - 1}$$

[In] integrate(1/(1-cosh(x)^2),x, algorithm="giac")

[Out] 2/(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 10, normalized size of antiderivative = 5.00

$$\int \frac{1}{1 - \cosh^2(x)} dx = \frac{2}{e^{2x} - 1}$$

[In] int(-1/(cosh(x)^2 - 1),x)

[Out] 2/(exp(2*x) - 1)

$$3.39 \quad \int \frac{1}{(1 - \cosh^2(x))^2} dx$$

Optimal result	297
Rubi [A] (verified)	297
Mathematica [A] (verified)	298
Maple [A] (verified)	298
Fricas [B] (verification not implemented)	299
Sympy [B] (verification not implemented)	299
Maxima [B] (verification not implemented)	299
Giac [A] (verification not implemented)	300
Mupad [B] (verification not implemented)	300

Optimal result

Integrand size = 10, antiderivative size = 11

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = \coth(x) - \frac{\coth^3(x)}{3}$$

[Out] $\coth(x) - 1/3 * \coth(x)^3$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3254, 3852}

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = \coth(x) - \frac{\coth^3(x)}{3}$$

[In] $\text{Int}[(1 - \text{Cosh}[x]^2)^{-2}, x]$

[Out] $\text{Coth}[x] - \text{Coth}[x]^3/3$

Rule 3254

$\text{Int}[(u_.)*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[a^p, \text{Int}[\text{ActivateTrig}[u*\cos[e + f*x]^{(2*p)}], x], x] /;$ $\text{FreeQ}\{a, b, e, f, p, x\}$ && $\text{EqQ}[a + b, 0]$ && $\text{IntegerQ}[p]$

Rule 3852

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /;$ $\text{FreeQ}\{c,$

d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \operatorname{csch}^4(x) dx \\ &= i\text{Subst}\left(\int (1+x^2) dx, x, -i \operatorname{coth}(x)\right) \\ &= \operatorname{coth}(x) - \frac{\operatorname{coth}^3(x)}{3} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.55

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = \frac{2 \operatorname{coth}(x)}{3} - \frac{1}{3} \operatorname{coth}(x) \operatorname{csch}^2(x)$$

[In] Integrate[(1 - Cosh[x]^2)^(-2), x]

[Out] (2*Coth[x])/3 - (Coth[x]*Csch[x]^2)/3

Maple [A] (verified)

Time = 0.08 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.45

method	result	size
paralelrisch	$\frac{2 \operatorname{coth}(x)^3}{3} - \operatorname{coth}(x) \operatorname{csch}(x)^2$	16
risch	$-\frac{4(3e^{2x}-1)}{3(e^{2x}-1)^3}$	19
default	$-\frac{\tanh(\frac{x}{2})^3}{24} + \frac{3 \tanh(\frac{x}{2})}{8} - \frac{1}{24 \tanh(\frac{x}{2})^3} + \frac{3}{8 \tanh(\frac{x}{2})}$	32

[In] int(1/(1-cosh(x)^2)^2,x,method=_RETURNVERBOSE)

[Out] 2/3*coth(x)^3-coth(x)*csch(x)^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(9) = 18.

Time = 0.24 (sec) , antiderivative size = 84, normalized size of antiderivative = 7.64

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = \frac{8(\cosh(x) + 2 \sinh(x))}{3(\cosh(x)^5 + 5 \cosh(x) \sinh(x)^4 + \sinh(x)^5 + (10 \cosh(x)^2 - 3) \sinh(x)^3 - 3 \cosh(x)^3 + (10 \cosh(x) - 9) \sinh(x)^2 + (5 \cosh(x)^4 - 9 \cosh(x)^2 + 4) \sinh(x) + 2 \cosh(x))}$$

[In] integrate(1/(1-cosh(x)^2)^2,x, algorithm="fricas")

[Out] -8/3*(cosh(x) + 2*sinh(x))/(cosh(x)^5 + 5*cosh(x)*sinh(x)^4 + sinh(x)^5 + (10*cosh(x)^2 - 3)*sinh(x)^3 - 3*cosh(x)^3 + (10*cosh(x)^3 - 9*cosh(x))*sinh(x)^2 + (5*cosh(x)^4 - 9*cosh(x)^2 + 4)*sinh(x) + 2*cosh(x))

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(8) = 16.

Time = 0.52 (sec) , antiderivative size = 34, normalized size of antiderivative = 3.09

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = -\frac{\tanh^3\left(\frac{x}{2}\right)}{24} + \frac{3 \tanh\left(\frac{x}{2}\right)}{8} + \frac{3}{8 \tanh\left(\frac{x}{2}\right)} - \frac{1}{24 \tanh^3\left(\frac{x}{2}\right)}$$

[In] integrate(1/(1-cosh(x)**2)**2,x)

[Out] -tanh(x/2)**3/24 + 3*tanh(x/2)/8 + 3/(8*tanh(x/2)) - 1/(24*tanh(x/2)**3)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. 2(9) = 18.

Time = 0.19 (sec) , antiderivative size = 49, normalized size of antiderivative = 4.45

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = \frac{4e^{(-2x)}}{3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1} - \frac{4}{3(3e^{(-2x)} - 3e^{(-4x)} + e^{(-6x)} - 1)}$$

[In] integrate(1/(1-cosh(x)^2)^2,x, algorithm="maxima")

[Out] 4*e^(-2*x)/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1) - 4/3/(3*e^(-2*x) - 3*e^(-4*x) + e^(-6*x) - 1)

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = -\frac{4(3e^{(2x)} - 1)}{3(e^{(2x)} - 1)^3}$$

[In] integrate(1/(1-cosh(x)^2)^2,x, algorithm="giac")

[Out] -4/3*(3*e^(2*x) - 1)/(e^(2*x) - 1)^3

Mupad [B] (verification not implemented)

Time = 1.83 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.64

$$\int \frac{1}{(1 - \cosh^2(x))^2} dx = -\frac{4(3e^{2x} - 1)}{3(e^{2x} - 1)^3}$$

[In] int(1/(cosh(x)^2 - 1)^2,x)

[Out] -(4*(3*exp(2*x) - 1))/(3*(exp(2*x) - 1)^3)

$$3.40 \quad \int \frac{1}{(1 - \cosh^2(x))^3} dx$$

Optimal result	301
Rubi [A] (verified)	301
Mathematica [A] (verified)	302
Maple [A] (verified)	302
Fricas [B] (verification not implemented)	303
Sympy [B] (verification not implemented)	303
Maxima [B] (verification not implemented)	304
Giac [A] (verification not implemented)	304
Mupad [B] (verification not implemented)	304

Optimal result

Integrand size = 10, antiderivative size = 19

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \coth(x) - \frac{2 \coth^3(x)}{3} + \frac{\coth^5(x)}{5}$$

[Out] $\coth(x) - 2/3 * \coth(x)^3 + 1/5 * \coth(x)^5$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3254, 3852}

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{\coth^5(x)}{5} - \frac{2 \coth^3(x)}{3} + \coth(x)$$

[In] $\text{Int}[(1 - \text{Cosh}[x]^2)^{-3}, x]$

[Out] $\text{Coth}[x] - (2 * \text{Coth}[x]^3) / 3 + \text{Coth}[x]^5 / 5$

Rule 3254

$\text{Int}[(u_.) * ((a_.) + (b_.) * \sin[(e_.) + (f_.) * (x_.)]^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[a^p, \text{Int}[\text{ActivateTrig}[u * \cos[e + f * x]^{(2 * p)}], x], x] /;$ $\text{FreeQ}\{a, b, e, f, p, x\}$ && $\text{EqQ}[a + b, 0]$ && $\text{IntegerQ}[p]$

Rule 3852

$\text{Int}[\text{csc}[(c_.) + (d_.) * (x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-d^{-1}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d * x]], x] /;$ $\text{FreeQ}\{c,$

d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= - \int \operatorname{csch}^6(x) dx \\ &= i\text{Subst}\left(\int (1 + 2x^2 + x^4) dx, x, -i \operatorname{coth}(x)\right) \\ &= \operatorname{coth}(x) - \frac{2 \operatorname{coth}^3(x)}{3} + \frac{\operatorname{coth}^5(x)}{5} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.00 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.42

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{8 \operatorname{coth}(x)}{15} - \frac{4}{15} \operatorname{coth}(x) \operatorname{csch}^2(x) + \frac{1}{5} \operatorname{coth}(x) \operatorname{csch}^4(x)$$

[In] Integrate[(1 - Cosh[x]^2)^(-3), x]

[Out] (8*Coth[x])/15 - (4*Coth[x]*Csch[x]^2)/15 + (Coth[x]*Csch[x]^4)/5

Maple [A] (verified)

Time = 0.09 (sec) , antiderivative size = 21, normalized size of antiderivative = 1.11

method	result	size
paralelrisch	$\frac{\operatorname{coth}(x) \operatorname{csch}(x)^4 (8 + \cosh(4x) - 6 \cosh(2x))}{15}$	21
risch	$\frac{\frac{32 e^{4x}}{3} - \frac{16 e^{2x}}{3} + \frac{16}{15}}{(e^{2x} - 1)^5}$	25
default	$\frac{\tanh(\frac{x}{2})^5}{160} - \frac{5 \tanh(\frac{x}{2})^3}{96} + \frac{5 \tanh(\frac{x}{2})}{16} + \frac{1}{160 \tanh(\frac{x}{2})^5} - \frac{5}{96 \tanh(\frac{x}{2})^3} + \frac{5}{16 \tanh(\frac{x}{2})}$	48

[In] int(1/(1-cosh(x)^2)^3,x,method=_RETURNVERBOSE)

[Out] 1/15*coth(x)*csch(x)^4*(8+cosh(4*x)-6*cosh(2*x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 185 vs. $2(15) = 30$.

Time = 0.24 (sec) , antiderivative size = 185, normalized size of antiderivative = 9.74

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx$$

$$= \frac{15 (\cosh(x))^8 + 8 \cosh(x) \sinh(x)^7 + \sinh(x)^8 + (28 \cosh(x)^2 - 5) \sinh(x)^6 - 5 \cosh(x)^6 + 2 (28 \cosh(x)^3 - 15 \cosh(x)) \sinh(x)^5 + 5 (14 \cosh(x)^4 - 15 \cosh(x)^2 + 2) \sinh(x)^4 + 10 \cosh(x)^4 + 4 (14 \cosh(x)^5 - 25 \cosh(x)^3 + 10 \cosh(x)) \sinh(x)^3 + (28 \cosh(x)^6 - 75 \cosh(x)^4 + 60 \cosh(x)^2 - 11) \sinh(x)^2 - 11 \cosh(x)^2 + 2 (4 \cosh(x)^7 - 15 \cosh(x)^5 + 20 \cosh(x)^3 - 9 \cosh(x)) \sinh(x) + 5}{(1 - \cosh^2(x))^3}$$

[In] integrate(1/(1-cosh(x)^2)^3,x, algorithm="fricas")

[Out] 16/15*(11*cosh(x)^2 + 18*cosh(x)*sinh(x) + 11*sinh(x)^2 - 5)/(cosh(x)^8 + 8*cosh(x)*sinh(x)^7 + sinh(x)^8 + (28*cosh(x)^2 - 5)*sinh(x)^6 - 5*cosh(x)^6 + 2*(28*cosh(x)^3 - 15*cosh(x))*sinh(x)^5 + 5*(14*cosh(x)^4 - 15*cosh(x)^2 + 2)*sinh(x)^4 + 10*cosh(x)^4 + 4*(14*cosh(x)^5 - 25*cosh(x)^3 + 10*cosh(x))*sinh(x)^3 + (28*cosh(x)^6 - 75*cosh(x)^4 + 60*cosh(x)^2 - 11)*sinh(x)^2 - 11*cosh(x)^2 + 2*(4*cosh(x)^7 - 15*cosh(x)^5 + 20*cosh(x)^3 - 9*cosh(x))*sinh(x) + 5)

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 54 vs. $2(17) = 34$.

Time = 1.57 (sec) , antiderivative size = 54, normalized size of antiderivative = 2.84

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{\tanh^5\left(\frac{x}{2}\right)}{160} - \frac{5 \tanh^3\left(\frac{x}{2}\right)}{96} + \frac{5 \tanh\left(\frac{x}{2}\right)}{16} + \frac{5}{16 \tanh\left(\frac{x}{2}\right)} - \frac{5}{96 \tanh^3\left(\frac{x}{2}\right)} + \frac{1}{160 \tanh^5\left(\frac{x}{2}\right)}$$

[In] integrate(1/(1-cosh(x)**2)**3,x)

[Out] tanh(x/2)**5/160 - 5*tanh(x/2)**3/96 + 5*tanh(x/2)/16 + 5/(16*tanh(x/2)) - 5/(96*tanh(x/2)**3) + 1/(160*tanh(x/2)**5)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(15) = 30$.

Time = 0.20 (sec) , antiderivative size = 111, normalized size of antiderivative = 5.84

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{16 e^{(-2x)}}{3(5 e^{(-2x)} - 10 e^{(-4x)} + 10 e^{(-6x)} - 5 e^{(-8x)} + e^{(-10x)} - 1)} - \frac{32 e^{(-4x)}}{3(5 e^{(-2x)} - 10 e^{(-4x)} + 10 e^{(-6x)} - 5 e^{(-8x)} + e^{(-10x)} - 1)} - \frac{16}{15(5 e^{(-2x)} - 10 e^{(-4x)} + 10 e^{(-6x)} - 5 e^{(-8x)} + e^{(-10x)} - 1)}$$

[In] integrate(1/(1-cosh(x)^2)^3,x, algorithm="maxima")

[Out] $\frac{16}{3}e^{-2x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1) - \frac{32}{3}e^{-4x}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1) - \frac{16}{15}/(5e^{-2x} - 10e^{-4x} + 10e^{-6x} - 5e^{-8x} + e^{-10x} - 1)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.26

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{16(10e^{4x} - 5e^{2x} + 1)}{15(e^{2x} - 1)^5}$$

[In] integrate(1/(1-cosh(x)^2)^3,x, algorithm="giac")

[Out] $\frac{16}{15}*(10*e^{4*x} - 5*e^{2*x} + 1)/(e^{2*x} - 1)^5$

Mupad [B] (verification not implemented)

Time = 0.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.26

$$\int \frac{1}{(1 - \cosh^2(x))^3} dx = \frac{16(10e^{4x} - 5e^{2x} + 1)}{15(e^{2x} - 1)^5}$$

[In] int(-1/(cosh(x)^2 - 1)^3,x)

[Out] $(16*(10*\exp(4*x) - 5*\exp(2*x) + 1))/(15*(\exp(2*x) - 1)^5)$

3.41 $\int \sqrt{a + b \cosh^2(x)} dx$

Optimal result	305
Rubi [A] (verified)	305
Mathematica [A] (verified)	306
Maple [B] (verified)	306
Fricas [F]	307
Sympy [F]	307
Maxima [F]	307
Giac [F]	307
Mupad [F(-1)]	308

Optimal result

Integrand size = 12, antiderivative size = 49

$$\int \sqrt{a + b \cosh^2(x)} dx = -\frac{i\sqrt{a + b \cosh^2(x)}E\left(\frac{\pi}{2} + ix \mid -\frac{b}{a}\right)}{\sqrt{1 + \frac{b \cosh^2(x)}{a}}}$$

[Out] $(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\text{EllipticE}(\cosh(x), (-b/a)^{(1/2)})*(a+b*\cosh(x)^2)^{(1/2)}/(1+b*\cosh(x)^2/a)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3257, 3256}

$$\int \sqrt{a + b \cosh^2(x)} dx = -\frac{i\sqrt{a + b \cosh^2(x)}E\left(ix + \frac{\pi}{2} \mid -\frac{b}{a}\right)}{\sqrt{\frac{b \cosh^2(x)}{a} + 1}}$$

[In] `Int[Sqrt[a + b*Cosh[x]^2],x]`

[Out] `((-I)*Sqrt[a + b*Cosh[x]^2]*EllipticE[Pi/2 + I*x, -(b/a)]/Sqrt[1 + (b*Cosh[x]^2)/a]`

Rule 3256

`Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Simp[(Sqrt[a]/f)*EllipticE[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]`

Rule 3257

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Dist[Sqrt[a
+ b*Sin[e + f*x]^2]/Sqrt[1 + b*(Sin[e + f*x]^2/a)], Int[Sqrt[1 + (b*Sin[e +
f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{a + b \cosh^2(x)} \int \sqrt{1 + \frac{b \cosh^2(x)}{a}} dx}{\sqrt{1 + \frac{b \cosh^2(x)}{a}}} \\ &= -\frac{i \sqrt{a + b \cosh^2(x)} E\left(\frac{\pi}{2} + ix \middle| -\frac{b}{a}\right)}{\sqrt{1 + \frac{b \cosh^2(x)}{a}}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \sqrt{a + b \cosh^2(x)} dx = -\frac{i \sqrt{2a + b + b \cosh(2x)} E\left(ix \middle| \frac{b}{a+b}\right)}{\sqrt{\frac{2a+b+b \cosh(2x)}{a+b}}}$$

```
[In] Integrate[Sqrt[a + b*Cosh[x]^2], x]
```

```
[Out] ((-I)*Sqrt[2*a + b + b*Cosh[2*x]]*EllipticE[I*x, b/(a + b)])/Sqrt[(2*a + b
+ b*Cosh[2*x])/(a + b)]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(47) = 94.

Time = 1.10 (sec) , antiderivative size = 114, normalized size of antiderivative = 2.33

method	result
default	$\frac{\sqrt{\frac{a+b \cosh(x)^2}{a}} \sqrt{-\sinh(x)^2} \left(a \operatorname{EllipticF}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right) + b \operatorname{EllipticF}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right) - b \operatorname{EllipticE}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right) \right)}{\sqrt{-\frac{b}{a}} \sinh(x) \sqrt{a+b \cosh(x)^2}}$

```
[In] int((a+b*cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] ((a+b*cosh(x)^2)/a)^(1/2)*(-sinh(x)^2)^(1/2)*(a*EllipticF(cosh(x)*(-b/a)^(1
/2), (-a/b)^(1/2))+b*EllipticF(cosh(x)*(-b/a)^(1/2), (-a/b)^(1/2))-b*Elliptic
E(cosh(x)*(-b/a)^(1/2), (-a/b)^(1/2)))/(-b/a)^(1/2)/sinh(x)/(a+b*cosh(x)^2)^(
1/2)
```

Fricas [F]

$$\int \sqrt{a + b \cosh^2(x)} dx = \int \sqrt{b \cosh(x)^2 + a} dx$$

[In] integrate((a+b*cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*cosh(x)^2 + a), x)

Sympy [F]

$$\int \sqrt{a + b \cosh^2(x)} dx = \int \sqrt{a + b \cosh^2(x)} dx$$

[In] integrate((a+b*cosh(x)**2)**(1/2),x)

[Out] Integral(sqrt(a + b*cosh(x)**2), x)

Maxima [F]

$$\int \sqrt{a + b \cosh^2(x)} dx = \int \sqrt{b \cosh(x)^2 + a} dx$$

[In] integrate((a+b*cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*cosh(x)^2 + a), x)

Giac [F]

$$\int \sqrt{a + b \cosh^2(x)} dx = \int \sqrt{b \cosh(x)^2 + a} dx$$

[In] integrate((a+b*cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*cosh(x)^2 + a), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cosh^2(x)} dx = \int \sqrt{b \cosh(x)^2 + a} dx$$

```
[In] int((a + b*cosh(x)^2)^(1/2),x)
```

```
[Out] int((a + b*cosh(x)^2)^(1/2), x)
```

3.42 $\int \sqrt{1 + \cosh^2(x)} dx$

Optimal result	309
Rubi [A] (verified)	309
Mathematica [A] (verified)	310
Maple [B] (verified)	310
Fricas [F]	310
Sympy [F]	311
Maxima [F]	311
Giac [F]	311
Mupad [F(-1)]	311

Optimal result

Integrand size = 10, antiderivative size = 17

$$\int \sqrt{1 + \cosh^2(x)} dx = -iE\left(\frac{\pi}{2} + ix \mid -1\right)$$

[Out] $(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\text{EllipticE}(\cosh(x), I)$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3256}

$$\int \sqrt{1 + \cosh^2(x)} dx = -iE\left(ix + \frac{\pi}{2} \mid -1\right)$$

[In] `Int[Sqrt[1 + Cosh[x]^2], x]`

[Out] `(-I)*EllipticE[Pi/2 + I*x, -1]`

Rule 3256

`Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] :> Simp[(Sqrt[a]/f)*EllipticE[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]`

Rubi steps

$$\text{integral} = -iE\left(\frac{\pi}{2} + ix \mid -1\right)$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \sqrt{1 + \cosh^2(x)} dx = -i\sqrt{2}E\left(ix \middle| \frac{1}{2}\right)$$

[In] Integrate[Sqrt[1 + Cosh[x]^2], x]

[Out] (-I)*Sqrt[2]*EllipticE[I*x, 1/2]

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 57 vs. $2(18) = 36$.

Time = 0.62 (sec) , antiderivative size = 58, normalized size of antiderivative = 3.41

method	result	size
default	$-\frac{i\sqrt{(1+\cosh(x)^2)} \sinh(x)^2 \sqrt{-\sinh(x)^2} (2 \operatorname{EllipticF}(i \cosh(x), i) - \operatorname{EllipticE}(i \cosh(x), i))}{\sqrt{\cosh(x)^4 - 1} \sinh(x)}$	58

[In] int((1+cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -I*((1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)*(2*EllipticF(I*cosh(x), I)-EllipticE(I*cosh(x), I))/(cosh(x)^4-1)^(1/2)/sinh(x)

Fricas [F]

$$\int \sqrt{1 + \cosh^2(x)} dx = \int \sqrt{\cosh(x)^2 + 1} dx$$

[In] integrate((1+cosh(x)^2)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(cosh(x)^2 + 1), x)

Sympy [F]

$$\int \sqrt{1 + \cosh^2(x)} dx = \int \sqrt{\cosh^2(x) + 1} dx$$

[In] integrate((1+cosh(x)**2)**(1/2),x)

[Out] Integral(sqrt(cosh(x)**2 + 1), x)

Maxima [F]

$$\int \sqrt{1 + \cosh^2(x)} dx = \int \sqrt{\cosh(x)^2 + 1} dx$$

[In] integrate((1+cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(cosh(x)^2 + 1), x)

Giac [F]

$$\int \sqrt{1 + \cosh^2(x)} dx = \int \sqrt{\cosh(x)^2 + 1} dx$$

[In] integrate((1+cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(cosh(x)^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{1 + \cosh^2(x)} dx = \int \sqrt{\cosh(x)^2 + 1} dx$$

[In] int((cosh(x)^2 + 1)^(1/2),x)

[Out] int((cosh(x)^2 + 1)^(1/2), x)

3.43 $\int \sqrt{1 - \cosh^2(x)} dx$

Optimal result	312
Rubi [A] (verified)	312
Mathematica [A] (verified)	313
Maple [A] (verified)	313
Fricas [B] (verification not implemented)	314
Sympy [F]	314
Maxima [C] (verification not implemented)	314
Giac [C] (verification not implemented)	314
Mupad [B] (verification not implemented)	315

Optimal result

Integrand size = 12, antiderivative size = 13

$$\int \sqrt{1 - \cosh^2(x)} dx = \coth(x) \sqrt{-\sinh^2(x)}$$

[Out] $\coth(x) * (-\sinh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3255, 3286, 2718}

$$\int \sqrt{1 - \cosh^2(x)} dx = \sqrt{-\sinh^2(x)} \coth(x)$$

[In] `Int[Sqrt[1 - Cosh[x]^2], x]`

[Out] `Coth[x]*Sqrt[-Sinh[x]^2]`

Rule 2718

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3255

`Int[(u_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2)^p, x_Symbol] := Int[ActivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]`

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*SIn[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \sqrt{-\sinh^2(x)} dx \\ &= \left(\operatorname{csch}(x) \sqrt{-\sinh^2(x)} \right) \int \sinh(x) dx \\ &= \operatorname{coth}(x) \sqrt{-\sinh^2(x)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - \cosh^2(x)} dx = \operatorname{coth}(x) \sqrt{-\sinh^2(x)}$$

[In] Integrate[Sqrt[1 - Cosh[x]^2], x]

[Out] Coth[x]*Sqrt[-Sinh[x]^2]

Maple [A] (verified)

Time = 0.13 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.15

method	result	size
default	$-\frac{\cosh(x) \sinh(x)}{\sqrt{-\sinh^2(x)}}$	15
risch	$\frac{\sqrt{-(e^{2x}-1)^2 e^{-2x}} e^{2x}}{2 e^{2x}-2} + \frac{\sqrt{-(e^{2x}-1)^2 e^{-2x}}}{2 e^{2x}-2}$	58

[In] int((1-cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -cosh(x)*sinh(x)/(-sinh(x)^2)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 33, normalized size of antiderivative = 2.54

$$\int \sqrt{1 - \cosh^2(x)} dx = \frac{\sqrt{-(e^{4x}) - 2e^{2x} + 1}e^{-2x} \cosh(x) e^x}{e^{2x} - 1}$$

[In] integrate((1-cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] sqrt(-(e^(4*x) - 2*e^(2*x) + 1)*e^(-2*x))*cosh(x)*e^x/(e^(2*x) - 1)

Sympy [F]

$$\int \sqrt{1 - \cosh^2(x)} dx = \int \sqrt{1 - \cosh^2(x)} dx$$

[In] integrate((1-cosh(x)**2)**(1/2),x)

[Out] Integral(sqrt(1 - cosh(x)**2), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 11, normalized size of antiderivative = 0.85

$$\int \sqrt{1 - \cosh^2(x)} dx = -\frac{1}{2}i e^{(-x)} - \frac{1}{2}i e^x$$

[In] integrate((1-cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/2*I*e^(-x) - 1/2*I*e^x

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.38

$$\int \sqrt{1 - \cosh^2(x)} dx = -\frac{1}{2}i e^{(-x)} \operatorname{sgn}(-e^{(3x)} + e^x) - \frac{1}{2}i e^x \operatorname{sgn}(-e^{(3x)} + e^x)$$

[In] integrate((1-cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*I*e^(-x)*sgn(-e^(3*x) + e^x) - 1/2*I*e^x*sgn(-e^(3*x) + e^x)

Mupad [B] (verification not implemented)

Time = 1.86 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \sqrt{1 - \cosh^2(x)} dx = \coth(x) \sqrt{1 - \cosh(x)^2}$$

[In] `int((1 - cosh(x)^2)^(1/2),x)`

[Out] `coth(x)*(1 - cosh(x)^2)^(1/2)`

3.44 $\int \sqrt{-1 + \cosh^2(x)} dx$

Optimal result	316
Rubi [A] (verified)	316
Mathematica [A] (verified)	317
Maple [A] (verified)	317
Fricas [A] (verification not implemented)	318
Sympy [F]	318
Maxima [A] (verification not implemented)	318
Giac [B] (verification not implemented)	318
Mupad [B] (verification not implemented)	319

Optimal result

Integrand size = 10, antiderivative size = 11

$$\int \sqrt{-1 + \cosh^2(x)} dx = \coth(x) \sqrt{\sinh^2(x)}$$

[Out] $\coth(x) * (\sinh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3255, 3286, 2718}

$$\int \sqrt{-1 + \cosh^2(x)} dx = \sqrt{\sinh^2(x)} \coth(x)$$

[In] `Int[Sqrt[-1 + Cosh[x]^2], x]`

[Out] `Coth[x]*Sqrt[Sinh[x]^2]`

Rule 2718

`Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]`

Rule 3255

`Int[(u_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2]^(p_), x_Symbol] := Int[ActivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]`

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p] && IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.)] /; FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \sqrt{\sinh^2(x)} dx \\ &= \left(\operatorname{csch}(x) \sqrt{\sinh^2(x)} \right) \int \sinh(x) dx \\ &= \operatorname{coth}(x) \sqrt{\sinh^2(x)} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + \cosh^2(x)} dx = \operatorname{coth}(x) \sqrt{\sinh^2(x)}$$

[In] Integrate[Sqrt[-1 + Cosh[x]^2], x]

[Out] Coth[x]*Sqrt[Sinh[x]^2]

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.27

method	result	size
default	$\frac{\cosh(x) \sqrt{-\frac{1}{2} + \frac{\cosh(2x)}{2}}}{\sinh(x)}$	14
risch	$\frac{\sqrt{(e^{2x}-1)^2 e^{-2x}} e^{2x}}{2e^{2x}-2} + \frac{\sqrt{(e^{2x}-1)^2 e^{-2x}}}{2e^{2x}-2}$	56

[In] int((cosh(x)^2-1)^(1/2), x, method=_RETURNVERBOSE)

[Out] cosh(x)*(sinh(x)^2)^(1/2)/sinh(x)

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 2, normalized size of antiderivative = 0.18

$$\int \sqrt{-1 + \cosh^2(x)} dx = \cosh(x)$$

[In] integrate((-1+cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] cosh(x)

Sympy [F]

$$\int \sqrt{-1 + \cosh^2(x)} dx = \int \sqrt{\cosh^2(x) - 1} dx$$

[In] integrate((-1+cosh(x)**2)**(1/2),x)

[Out] Integral(sqrt(cosh(x)**2 - 1), x)

Maxima [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + \cosh^2(x)} dx = -\frac{1}{2} e^{(-x)} - \frac{1}{2} e^x$$

[In] integrate((-1+cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] -1/2*e^(-x) - 1/2*e^x

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(9) = 18.

Time = 0.28 (sec) , antiderivative size = 31, normalized size of antiderivative = 2.82

$$\int \sqrt{-1 + \cosh^2(x)} dx = \frac{1}{2} e^{(-x)} \operatorname{sgn}(e^{(3x)} - e^x) + \frac{1}{2} e^x \operatorname{sgn}(e^{(3x)} - e^x)$$

[In] integrate((-1+cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/2*e^(-x)*sgn(e^(3*x) - e^x) + 1/2*e^x*sgn(e^(3*x) - e^x)

Mupad [B] (verification not implemented)

Time = 1.78 (sec) , antiderivative size = 11, normalized size of antiderivative = 1.00

$$\int \sqrt{-1 + \cosh^2(x)} dx = \coth(x) \sqrt{\cosh(x)^2 - 1}$$

[In] int((cosh(x)^2 - 1)^(1/2),x)

[Out] coth(x)*(cosh(x)^2 - 1)^(1/2)

3.45 $\int \sqrt{-1 - \cosh^2(x)} dx$

Optimal result	320
Rubi [A] (verified)	320
Mathematica [A] (verified)	321
Maple [A] (verified)	321
Fricas [F]	322
Sympy [F]	322
Maxima [F]	322
Giac [F]	322
Mupad [F(-1)]	323

Optimal result

Integrand size = 12, antiderivative size = 39

$$\int \sqrt{-1 - \cosh^2(x)} dx = -\frac{i\sqrt{-1 - \cosh^2(x)}E\left(\frac{\pi}{2} + ix \mid -1\right)}{\sqrt{1 + \cosh^2(x)}}$$

[Out] $(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\text{EllipticE}(\cosh(x), I)*(-1-\cosh(x)^2)^{(1/2)}/(1+\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3257, 3256}

$$\int \sqrt{-1 - \cosh^2(x)} dx = -\frac{i\sqrt{-\cosh^2(x) - 1}E\left(ix + \frac{\pi}{2} \mid -1\right)}{\sqrt{\cosh^2(x) + 1}}$$

[In] `Int[Sqrt[-1 - Cosh[x]^2], x]`

[Out] `((-1)*Sqrt[-1 - Cosh[x]^2]*EllipticE[Pi/2 + I*x, -1])/Sqrt[1 + Cosh[x]^2]`

Rule 3256

`Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2], x_Symbol] := Simp[(Sqrt[a_]/f)*EllipticE[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]`

Rule 3257


```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Dist[Sqrt[a
+ b*Sin[e + f*x]^2]/Sqrt[1 + b*(Sin[e + f*x]^2/a)], Int[Sqrt[1 + (b*Sin[e +
f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{-1 - \cosh^2(x)} \int \sqrt{1 + \cosh^2(x)} dx}{\sqrt{1 + \cosh^2(x)}} \\ &= -\frac{i\sqrt{-1 - \cosh^2(x)} E\left(\frac{\pi}{2} + ix \mid -1\right)}{\sqrt{1 + \cosh^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

$$\int \sqrt{-1 - \cosh^2(x)} dx = \frac{i\sqrt{2}\sqrt{3 + \cosh(2x)} E(ix \mid \frac{1}{2})}{\sqrt{-3 - \cosh(2x)}}$$

[In] Integrate[Sqrt[-1 - Cosh[x]^2], x]

[Out] (I*Sqrt[2]*Sqrt[3 + Cosh[2*x]]*EllipticE[I*x, 1/2])/Sqrt[-3 - Cosh[2*x]]

Maple [A] (verified)

Time = 0.63 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.59

method	result	size
default	$-\frac{\sqrt{-(1+\cosh(x)^2)} \sinh(x)^2 \sqrt{-\sinh(x)^2} \sqrt{1+\cosh(x)^2} \text{EllipticE}(\cosh(x), i)}{\sqrt{1-\cosh(x)^4} \sinh(x) \sqrt{-1-\cosh(x)^2}}$	62

[In] int((-1-cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)

[Out] -((1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)*(1+cosh(x)^2)^(1/2)*EllipticE(cosh(x), I)/(1-cosh(x)^4)^(1/2)/sinh(x)/(-1-cosh(x)^2)^(1/2)

Fricas [F]

$$\int \sqrt{-1 - \cosh^2(x)} dx = \int \sqrt{-\cosh(x)^2 - 1} dx$$

[In] integrate((-1-cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] 1/2*(2*(e^(2*x) - e^x)*integral(4*sqrt(-e^(4*x) - 6*e^(2*x) - 1)*(e^(2*x) + 1)/(e^(6*x) - 2*e^(5*x) + 7*e^(4*x) - 12*e^(3*x) + 7*e^(2*x) - 2*e^x + 1), x) + sqrt(-e^(4*x) - 6*e^(2*x) - 1)*(e^x + 1))/(e^(2*x) - e^x)

Sympy [F]

$$\int \sqrt{-1 - \cosh^2(x)} dx = \int \sqrt{-\cosh^2(x) - 1} dx$$

[In] integrate((-1-cosh(x)**2)**(1/2),x)

[Out] Integral(sqrt(-cosh(x)**2 - 1), x)

Maxima [F]

$$\int \sqrt{-1 - \cosh^2(x)} dx = \int \sqrt{-\cosh(x)^2 - 1} dx$$

[In] integrate((-1-cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(-cosh(x)^2 - 1), x)

Giac [F]

$$\int \sqrt{-1 - \cosh^2(x)} dx = \int \sqrt{-\cosh(x)^2 - 1} dx$$

[In] integrate((-1-cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(-cosh(x)^2 - 1), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{-1 - \cosh^2(x)} dx = \int \sqrt{-\cosh(x)^2 - 1} dx$$

```
[In] int((- cosh(x)^2 - 1)^(1/2),x)
```

```
[Out] int((- cosh(x)^2 - 1)^(1/2), x)
```

3.46 $\int (a + b \cosh^2(x))^{3/2} dx$

Optimal result	324
Rubi [A] (verified)	324
Mathematica [A] (verified)	326
Maple [B] (verified)	327
Fricas [F]	327
Sympy [F]	327
Maxima [F]	328
Giac [F(-2)]	328
Mupad [F(-1)]	328

Optimal result

Integrand size = 12, antiderivative size = 133

$$\int (a + b \cosh^2(x))^{3/2} dx = -\frac{2i(2a + b)\sqrt{a + b \cosh^2(x)}E\left(\frac{\pi}{2} + ix \middle| -\frac{b}{a}\right)}{3\sqrt{1 + \frac{b \cosh^2(x)}{a}}} + \frac{ia(a + b)\sqrt{1 + \frac{b \cosh^2(x)}{a}} \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -\frac{b}{a}\right)}{3\sqrt{a + b \cosh^2(x)}} + \frac{1}{3}b \cosh(x)\sqrt{a + b \cosh^2(x)} \sinh(x)$$

[Out] $\frac{1}{3}b*\cosh(x)*\sinh(x)*(a+b*\cosh(x)^2)^{(1/2)}+2/3*(2*a+b)*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticE}(\cosh(x), (-b/a)^{(1/2)})*(a+b*\cosh(x)^2)^{(1/2)}/(1+b*\cosh(x)^2/a)^{(1/2)}-1/3*a*(a+b)*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), (-b/a)^{(1/2)})*(1+b*\cosh(x)^2/a)^{(1/2)}/(a+b*\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3259, 3251, 3257, 3256, 3262, 3261}

$$\int (a + b \cosh^2(x))^{3/2} dx = \frac{1}{3}b \sinh(x) \cosh(x)\sqrt{a + b \cosh^2(x)} + \frac{ia(a + b)\sqrt{\frac{b \cosh^2(x)}{a} + 1} \operatorname{EllipticF}\left(ix + \frac{\pi}{2}, -\frac{b}{a}\right)}{3\sqrt{a + b \cosh^2(x)}} - \frac{2i(2a + b)\sqrt{a + b \cosh^2(x)}E\left(ix + \frac{\pi}{2} \middle| -\frac{b}{a}\right)}{3\sqrt{\frac{b \cosh^2(x)}{a} + 1}}$$

[In] Int[(a + b*Cosh[x]^2)^(3/2),x]

[Out] (((-2*I)/3)*(2*a + b)*Sqrt[a + b*Cosh[x]^2]*EllipticE[Pi/2 + I*x, -(b/a)])/Sqrt[1 + (b*Cosh[x]^2)/a] + ((I/3)*a*(a + b)*Sqrt[1 + (b*Cosh[x]^2)/a]*EllipticF[Pi/2 + I*x, -(b/a)]/Sqrt[a + b*Cosh[x]^2] + (b*Cosh[x]*Sqrt[a + b*Cosh[x]^2]*Sinh[x])/3

Rule 3251

Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^2/Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] := Dist[B/b, Int[Sqrt[a + b*Sin[e + f*x]^2], x], x] + Dist[(A*b - a*B)/b, Int[1/Sqrt[a + b*Sin[e + f*x]^2], x], x] /; FreeQ[{a, b, e, f, A, B}, x]

Rule 3256

Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] := Simp[(Sqrt[a]/f)*EllipticE[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]

Rule 3257

Int[Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] := Dist[Sqrt[a + b*Sin[e + f*x]^2]/Sqrt[1 + b*(Sin[e + f*x]^2/a)], Int[Sqrt[1 + (b*Sin[e + f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]

Rule 3259

Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2^(p_), x_Symbol] := Simp[(-b)*Cos[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x]^2)^(p - 1)/(2*f*p)), x] + Dist[1/(2*p), Int[(a + b*Sin[e + f*x]^2)^(p - 2)*Simp[a*(b + 2*a*p) + b*(2*a + b)*(2*p - 1)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a + b, 0] && GtQ[p, 1]

Rule 3261

Int[1/Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] := Simp[(1/(Sqrt[a]*f))*EllipticF[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]

Rule 3262

Int[1/Sqrt[(a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] := Dist[Sqrt[1 + b*(Sin[e + f*x]^2/a)]/Sqrt[a + b*Sin[e + f*x]^2], Int[1/Sqrt[1 + (b*Sin[e + f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3}b \cosh(x) \sqrt{a + b \cosh^2(x)} \sinh(x) + \frac{1}{3} \int \frac{a(3a + b) + 2b(2a + b) \cosh^2(x)}{\sqrt{a + b \cosh^2(x)}} dx \\
 &= \frac{1}{3}b \cosh(x) \sqrt{a + b \cosh^2(x)} \sinh(x) - \frac{1}{3}(a(a + b)) \int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx \\
 &\quad + \frac{1}{3}(2(2a + b)) \int \sqrt{a + b \cosh^2(x)} dx \\
 &= \frac{1}{3}b \cosh(x) \sqrt{a + b \cosh^2(x)} \sinh(x) \\
 &\quad + \frac{\left(2(2a + b) \sqrt{a + b \cosh^2(x)}\right) \int \sqrt{1 + \frac{b \cosh^2(x)}{a}} dx}{3\sqrt{1 + \frac{b \cosh^2(x)}{a}}} \\
 &\quad - \frac{\left(a(a + b) \sqrt{1 + \frac{b \cosh^2(x)}{a}}\right) \int \frac{1}{\sqrt{1 + \frac{b \cosh^2(x)}{a}}} dx}{3\sqrt{a + b \cosh^2(x)}} \\
 &= -\frac{2i(2a + b) \sqrt{a + b \cosh^2(x)} E\left(\frac{\pi}{2} + ix \mid -\frac{b}{a}\right)}{3\sqrt{1 + \frac{b \cosh^2(x)}{a}}} \\
 &\quad + \frac{ia(a + b) \sqrt{1 + \frac{b \cosh^2(x)}{a}} \text{EllipticF}\left(\frac{\pi}{2} + ix, -\frac{b}{a}\right)}{3\sqrt{a + b \cosh^2(x)}} \\
 &\quad + \frac{1}{3}b \cosh(x) \sqrt{a + b \cosh^2(x)} \sinh(x)
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 135, normalized size of antiderivative = 1.02

$$\int (a + b \cosh^2(x))^{3/2} dx = \frac{-8i(2a^2 + 3ab + b^2) \sqrt{\frac{2a+b+b \cosh(2x)}{a+b}} E\left(ix \mid \frac{b}{a+b}\right) + 4ia(a + b) \sqrt{\frac{2a+b+b \cosh(2x)}{a+b}} \text{EllipticF}\left(ix, \frac{b}{a+b}\right)}{12\sqrt{2a + b + b \cosh(2x)}}$$

[In] Integrate[(a + b*Cosh[x]^2)^(3/2), x]

[Out] ((-8*I)*(2*a^2 + 3*a*b + b^2)*Sqrt[(2*a + b + b*Cosh[2*x])/(a + b)]*EllipticE[I*x, b/(a + b)] + (4*I)*a*(a + b)*Sqrt[(2*a + b + b*Cosh[2*x])/(a + b)]*EllipticF[I*x, b/(a + b)] + Sqrt[2]*b*(2*a + b + b*Cosh[2*x])*Sinh[2*x])/(12*Sqrt[2*a + b + b*Cosh[2*x]])

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(123) = 246$.

Time = 1.33 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.41

method	result
default	$\frac{\sqrt{-\frac{b}{a}} b^2 \cosh(x)^5 + \sqrt{-\frac{b}{a}} ab \cosh(x)^3 - \sqrt{-\frac{b}{a}} b^2 \cosh(x)^3 + 3a^2 \sqrt{\frac{a+b \cosh(x)^2}{a}} \sqrt{-\sinh(x)^2} \operatorname{EllipticF}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right) + 5ab \sqrt{\dots}}{\dots}$

[In] `int((a+b*cosh(x)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{3} \left((-b/a)^{1/2} b^2 \cosh(x)^5 + (-b/a)^{1/2} a b \cosh(x)^3 - (-b/a)^{1/2} b^2 \cosh(x)^3 + 3a^2 \sqrt{\frac{a+b \cosh(x)^2}{a}} \sqrt{-\sinh(x)^2} \operatorname{EllipticF}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right) + 5ab \sqrt{\dots} \right)$$

Fricas [F]

$$\int (a + b \cosh^2(x))^{3/2} dx = \int (b \cosh(x)^2 + a)^{3/2} dx$$

[In] `integrate((a+b*cosh(x)^2)^(3/2),x, algorithm="fricas")`

[Out] `integral((b*cosh(x)^2 + a)^(3/2), x)`

Sympy [F]

$$\int (a + b \cosh^2(x))^{3/2} dx = \int (a + b \cosh^2(x))^{3/2} dx$$

[In] `integrate((a+b*cosh(x)**2)**(3/2),x)`

[Out] `Integral((a + b*cosh(x)**2)**(3/2), x)`

Maxima [F]

$$\int (a + b \cosh^2(x))^{3/2} dx = \int (b \cosh(x)^2 + a)^{3/2} dx$$

[In] integrate((a+b*cosh(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate((b*cosh(x)^2 + a)^(3/2), x)

Giac [F(-2)]

Exception generated.

$$\int (a + b \cosh^2(x))^{3/2} dx = \text{Exception raised: AttributeError}$$

[In] integrate((a+b*cosh(x)^2)^(3/2),x, algorithm="giac")

[Out] Exception raised: AttributeError >> type

Mupad [F(-1)]

Timed out.

$$\int (a + b \cosh^2(x))^{3/2} dx = \int (b \cosh(x)^2 + a)^{3/2} dx$$

[In] int((a + b*cosh(x)^2)^(3/2),x)

[Out] int((a + b*cosh(x)^2)^(3/2), x)

3.47 $\int (1 + \cosh^2(x))^{3/2} dx$

Optimal result	329
Rubi [A] (verified)	329
Mathematica [A] (verified)	330
Maple [A] (verified)	331
Fricas [F]	331
Sympy [F(-1)]	331
Maxima [F]	331
Giac [F]	332
Mupad [F(-1)]	332

Optimal result

Integrand size = 10, antiderivative size = 55

$$\int (1 + \cosh^2(x))^{3/2} dx = -2iE\left(\frac{\pi}{2} + ix \mid -1\right) + \frac{2}{3}i \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right) + \frac{1}{3} \cosh(x) \sqrt{1 + \cosh^2(x)} \sinh(x)$$

[Out] $2*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticE}(\cosh(x), I) - 2/3*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), I) + 1/3*\cosh(x)*\sinh(x)*(1+\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3259, 3251, 3256, 3261}

$$\int (1 + \cosh^2(x))^{3/2} dx = \frac{2}{3}i \operatorname{EllipticF}\left(ix + \frac{\pi}{2}, -1\right) - 2iE\left(ix + \frac{\pi}{2} \mid -1\right) + \frac{1}{3} \sinh(x) \cosh(x) \sqrt{\cosh^2(x) + 1}$$

[In] Int[(1 + Cosh[x]^2)^(3/2), x]

[Out] $(-2*I)*\operatorname{EllipticE}[Pi/2 + I*x, -1] + ((2*I)/3)*\operatorname{EllipticF}[Pi/2 + I*x, -1] + (\cosh[x]*\operatorname{Sqrt}[1 + \cosh[x]^2]*\sinh[x])/3$

Rule 3251

Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)])^2/Sqrt[(a_.) + (b_.)*sin[(e_.) + (f_.)*(x_)])^2], x_Symbol] :> Dist[B/b, Int[Sqrt[a + b*Sin[e + f*x]^2], x], x] + Dist[(A*b - a*B)/b, Int[1/Sqrt[a + b*Sin[e + f*x]^2], x], x] /; FreeQ

[{a, b, e, f, A, B}, x]

Rule 3256

```
Int[Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Simp[(Sqrt[a
]/f)*EllipticE[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]
```

Rule 3259

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Simp[(-b)*C
os[e + f*x]*Sin[e + f*x]*((a + b*Sin[e + f*x]^2)^(p - 1)/(2*f*p)), x] + Dis
t[1/(2*p), Int[(a + b*Sin[e + f*x]^2)^(p - 2)*Simp[a*(b + 2*a*p) + b*(2*a +
b)*(2*p - 1)*Sin[e + f*x]^2, x], x], x] /; FreeQ[{a, b, e, f}, x] && NeQ[a
+ b, 0] && GtQ[p, 1]
```

Rule 3261

```
Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Simp[(1/(S
qrt[a]*f))*EllipticF[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a,
0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{3} \cosh(x) \sqrt{1 + \cosh^2(x)} \sinh(x) + \frac{1}{3} \int \frac{4 + 6 \cosh^2(x)}{\sqrt{1 + \cosh^2(x)}} dx \\ &= \frac{1}{3} \cosh(x) \sqrt{1 + \cosh^2(x)} \sinh(x) - \frac{2}{3} \int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx + 2 \int \sqrt{1 + \cosh^2(x)} dx \\ &= -2iE\left(\frac{\pi}{2} + ix \mid -1\right) + \frac{2}{3}i \text{EllipticF}\left(\frac{\pi}{2} + ix, -1\right) + \frac{1}{3} \cosh(x) \sqrt{1 + \cosh^2(x)} \sinh(x) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 51, normalized size of antiderivative = 0.93

$$\int (1 + \cosh^2(x))^{3/2} dx = \frac{-24iE\left(ix \mid \frac{1}{2}\right) + 4i \text{EllipticF}\left(ix, \frac{1}{2}\right) + \sqrt{3 + \cosh(2x)} \sinh(2x)}{6\sqrt{2}}$$

```
[In] Integrate[(1 + Cosh[x]^2)^(3/2), x]
```

```
[Out] ((-24*I)*EllipticE[I*x, 1/2] + (4*I)*EllipticF[I*x, 1/2] + Sqrt[3 + Cosh[2*
x]]*Sinh[2*x])/(6*Sqrt[2])
```

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 99, normalized size of antiderivative = 1.80

method	result
default	$-\frac{\sqrt{(1+\cosh(x)^2)} \sinh(x)^2 \left(-\cosh(x)^5 + 10i\sqrt{1+\cosh(x)^2} \sqrt{-\sinh(x)^2} \operatorname{EllipticF}(i \cosh(x), i) - 6i\sqrt{1+\cosh(x)^2} \sqrt{-\sinh(x)^2} \operatorname{EllipticE}(i \cosh(x), i) + \cosh(x) \right)}{3\sqrt{\cosh(x)^4 - 1} \sinh(x) \sqrt{1+\cosh(x)^2}}$

[In] int((1+cosh(x)^2)^(3/2),x,method=_RETURNVERBOSE)

```
[Out] -1/3*((1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-cosh(x)^5+10*I*(1+cosh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)*EllipticF(I*cosh(x),I)-6*I*(1+cosh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)*EllipticE(I*cosh(x),I)+cosh(x))/(cosh(x)^4-1)^(1/2)/sinh(x)/(1+cosh(x)^2)^(1/2)
```

Fricas [F]

$$\int (1 + \cosh^2(x))^{3/2} dx = \int (\cosh(x)^2 + 1)^{\frac{3}{2}} dx$$

[In] integrate((1+cosh(x)^2)^(3/2),x, algorithm="fricas")

[Out] integral((cosh(x)^2 + 1)^(3/2), x)

Sympy [F(-1)]

Timed out.

$$\int (1 + \cosh^2(x))^{3/2} dx = \text{Timed out}$$

[In] integrate((1+cosh(x)**2)**(3/2),x)

[Out] Timed out

Maxima [F]

$$\int (1 + \cosh^2(x))^{3/2} dx = \int (\cosh(x)^2 + 1)^{\frac{3}{2}} dx$$

[In] integrate((1+cosh(x)^2)^(3/2),x, algorithm="maxima")

[Out] integrate((cosh(x)^2 + 1)^(3/2), x)

Giac [F]

$$\int (1 + \cosh^2(x))^{3/2} dx = \int (\cosh(x)^2 + 1)^{\frac{3}{2}} dx$$

[In] integrate((1+cosh(x)^2)^(3/2),x, algorithm="giac")

[Out] integrate((cosh(x)^2 + 1)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (1 + \cosh^2(x))^{3/2} dx = \int (\cosh(x)^2 + 1)^{3/2} dx$$

[In] int((cosh(x)^2 + 1)^(3/2),x)

[Out] int((cosh(x)^2 + 1)^(3/2), x)

3.48 $\int (1 - \cosh^2(x))^{3/2} dx$

Optimal result	333
Rubi [A] (verified)	333
Mathematica [A] (verified)	334
Maple [A] (verified)	335
Fricas [B] (verification not implemented)	335
Sympy [F(-1)]	335
Maxima [C] (verification not implemented)	336
Giac [C] (verification not implemented)	336
Mupad [F(-1)]	336

Optimal result

Integrand size = 12, antiderivative size = 33

$$\int (1 - \cosh^2(x))^{3/2} dx = \frac{2}{3} \coth(x) \sqrt{-\sinh^2(x)} + \frac{1}{3} \coth(x) (-\sinh^2(x))^{3/2}$$

[Out] 1/3*coth(x)*(-sinh(x)^2)^(3/2)+2/3*coth(x)*(-sinh(x)^2)^(1/2)

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3255, 3282, 3286, 2718}

$$\int (1 - \cosh^2(x))^{3/2} dx = \frac{1}{3} (-\sinh^2(x))^{3/2} \coth(x) + \frac{2}{3} \sqrt{-\sinh^2(x)} \coth(x)$$

[In] Int[(1 - Cosh[x]^2)^(3/2), x]

[Out] (2*Coth[x]*Sqrt[-Sinh[x]^2])/3 + (Coth[x]*(-Sinh[x]^2)^(3/2))/3

Rule 2718

Int[sin[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rule 3255

Int[(u_.)*((a_.) + (b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]

Rule 3282

```
Int[((b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Simp[(-Cot[e + f*x
])*((b*Sin[e + f*x]^2)^p/(2*f*p)), x] + Dist[b*((2*p - 1)/(2*p)), Int[(b*Si
n[e + f*x]^2)^(p - 1), x], x] /; FreeQ[{b, e, f}, x] && !IntegerQ[p] && Gt
Q[p, 1]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p]))], Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (-\sinh^2(x))^{3/2} dx \\
&= \frac{1}{3} \coth(x) (-\sinh^2(x))^{3/2} + \frac{2}{3} \int \sqrt{-\sinh^2(x)} dx \\
&= \frac{1}{3} \coth(x) (-\sinh^2(x))^{3/2} + \frac{1}{3} \left(2\operatorname{csch}(x) \sqrt{-\sinh^2(x)} \right) \int \sinh(x) dx \\
&= \frac{2}{3} \coth(x) \sqrt{-\sinh^2(x)} + \frac{1}{3} \coth(x) (-\sinh^2(x))^{3/2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 25, normalized size of antiderivative = 0.76

$$\int (1 - \cosh^2(x))^{3/2} dx = -\frac{1}{12} (-9 \cosh(x) + \cosh(3x)) \operatorname{csch}(x) \sqrt{-\sinh^2(x)}$$

```
[In] Integrate[(1 - Cosh[x]^2)^(3/2), x]
```

```
[Out] -1/12*((-9*Cosh[x] + Cosh[3*x])*Csch[x]*Sqrt[-Sinh[x]^2])
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.64

method	result	size
default	$\frac{\sinh(x) \cosh(x) (\cosh(x)^2 - 3)}{3\sqrt{-\sinh(x)^2}}$	21
risch	$-\frac{e^{4x} \sqrt{-(e^{2x}-1)^2 e^{-2x}}}{24(e^{2x}-1)} + \frac{3\sqrt{-(e^{2x}-1)^2 e^{-2x}} e^{2x}}{8(e^{2x}-1)} + \frac{3\sqrt{-(e^{2x}-1)^2 e^{-2x}}}{8(e^{2x}-1)} - \frac{e^{-2x} \sqrt{-(e^{2x}-1)^2 e^{-2x}}}{24(e^{2x}-1)}$	118

[In] int((1-cosh(x)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] 1/3*sinh(x)*cosh(x)*(cosh(x)^2-3)/(-sinh(x)^2)^(1/2)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 53 vs. 2(25) = 50.

Time = 0.26 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.61

$$\int (1 - \cosh^2(x))^{3/2} dx = \frac{(3 \cosh(x) e^x \sinh(x)^2 + (\cosh(x)^3 - 9 \cosh(x)) e^x) \sqrt{-(e^{4x} - 2e^{2x} + 1)e^{-2x}}}{12(e^{2x} - 1)}$$

[In] integrate((1-cosh(x)^2)^(3/2),x, algorithm="fricas")

[Out] -1/12*(3*cosh(x)*e^x*sinh(x)^2 + (cosh(x)^3 - 9*cosh(x))*e^x)*sqrt(-(e^(4*x) - 2*e^(2*x) + 1)*e^(-2*x))/(e^(2*x) - 1)

Sympy [F(-1)]

Timed out.

$$\int (1 - \cosh^2(x))^{3/2} dx = \text{Timed out}$$

[In] integrate((1-cosh(x)**2)**(3/2),x)

[Out] Timed out

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.70

$$\int (1 - \cosh^2(x))^{3/2} dx = \frac{1}{24}i e^{(3x)} - \frac{3}{8}i e^{(-x)} + \frac{1}{24}i e^{(-3x)} - \frac{3}{8}i e^x$$

[In] integrate((1-cosh(x)^2)^(3/2),x, algorithm="maxima")

[Out] 1/24*I*e^(3*x) - 3/8*I*e^(-x) + 1/24*I*e^(-3*x) - 3/8*I*e^x

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.00

$$\int (1 - \cosh^2(x))^{3/2} dx = -\frac{1}{24}i (9 e^{(2x)} \operatorname{sgn}(-e^{(3x)} + e^x) - \operatorname{sgn}(-e^{(3x)} + e^x)) e^{(-3x)} \\ + \frac{1}{24}i e^{(3x)} \operatorname{sgn}(-e^{(3x)} + e^x) - \frac{3}{8}i e^x \operatorname{sgn}(-e^{(3x)} + e^x)$$

[In] integrate((1-cosh(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/24*I*(9*e^(2*x)*sgn(-e^(3*x) + e^x) - sgn(-e^(3*x) + e^x))*e^(-3*x) + 1/24*I*e^(3*x)*sgn(-e^(3*x) + e^x) - 3/8*I*e^x*sgn(-e^(3*x) + e^x)

Mupad [F(-1)]

Timed out.

$$\int (1 - \cosh^2(x))^{3/2} dx = \int (1 - \cosh(x)^2)^{3/2} dx$$

[In] int((1 - cosh(x)^2)^(3/2),x)

[Out] int((1 - cosh(x)^2)^(3/2), x)

3.49 $\int (-1 + \cosh^2(x))^{3/2} dx$

Optimal result	337
Rubi [A] (verified)	337
Mathematica [A] (verified)	338
Maple [A] (verified)	339
Fricas [A] (verification not implemented)	339
Sympy [F(-1)]	339
Maxima [A] (verification not implemented)	340
Giac [B] (verification not implemented)	340
Mupad [F(-1)]	340

Optimal result

Integrand size = 10, antiderivative size = 29

$$\int (-1 + \cosh^2(x))^{3/2} dx = -\frac{2}{3} \coth(x) \sqrt{\sinh^2(x)} + \frac{1}{3} \coth(x) \sinh^2(x)^{3/2}$$

[Out] $1/3*\coth(x)*(\sinh(x)^2)^{(3/2)}-2/3*\coth(x)*(\sinh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {3255, 3282, 3286, 2718}

$$\int (-1 + \cosh^2(x))^{3/2} dx = \frac{1}{3} \sinh^2(x)^{3/2} \coth(x) - \frac{2}{3} \sqrt{\sinh^2(x)} \coth(x)$$

[In] $\text{Int}[(-1 + \text{Cosh}[x]^2)^{(3/2)}, x]$

[Out] $(-2*\text{Coth}[x]*\text{Sqrt}[\text{Sinh}[x]^2])/3 + (\text{Coth}[x]*(\text{Sinh}[x]^2)^{(3/2}))/3$

Rule 2718

$\text{Int}[\sin[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/d, x] /;$ FreeQ[{c, d}, x]

Rule 3255

$\text{Int}[(u_.)*((a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2)^{(p_.), x_Symbol] \rightarrow \text{Int}[\text{ActivateTrig}[u*(a*\cos[e + f*x]^2)^p], x] /;$ FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]

Rule 3282

```
Int[((b_.)*sin[(e_.) + (f_.)*(x_.)]^2)^(p_), x_Symbol] := Simp[(-Cot[e + f*x]
)]*((b*Sin[e + f*x]^2)^p/(2*f*p)), x] + Dist[b*((2*p - 1)/(2*p)), Int[(b*Si
n[e + f*x]^2)^(p - 1), x], x] /; FreeQ[{b, e, f}, x] && !IntegerQ[p] && Gt
Q[p, 1]
```

Rule 3286

```
Int[(u_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.))^(p_), x_Symbol] := With[{ff
= FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sin[e + f*x]^
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p]))], Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x]] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \sinh^2(x)^{3/2} dx \\
&= \frac{1}{3} \coth(x) \sinh^2(x)^{3/2} - \frac{2}{3} \int \sqrt{\sinh^2(x)} dx \\
&= \frac{1}{3} \coth(x) \sinh^2(x)^{3/2} - \frac{1}{3} \left(2\operatorname{csch}(x) \sqrt{\sinh^2(x)} \right) \int \sinh(x) dx \\
&= -\frac{2}{3} \coth(x) \sqrt{\sinh^2(x)} + \frac{1}{3} \coth(x) \sinh^2(x)^{3/2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int (-1 + \cosh^2(x))^{3/2} dx = \frac{1}{12} (-9 \cosh(x) + \cosh(3x)) \operatorname{csch}(x) \sqrt{\sinh^2(x)}$$

```
[In] Integrate[(-1 + Cosh[x]^2)^(3/2), x]
```

```
[Out] ((-9*Cosh[x] + Cosh[3*x])*Csch[x]*Sqrt[Sinh[x]^2])/12
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 21, normalized size of antiderivative = 0.72

method	result	size
default	$\frac{\sqrt{-\frac{1}{2} + \frac{\cosh(2x)}{2}} \cosh(x) (\cosh(x)^2 - 3)}{3 \sinh(x)}$	21
risch	$\frac{e^{4x} \sqrt{(e^{2x}-1)^2 e^{-2x}}}{24 e^{2x}-24} - \frac{3 \sqrt{(e^{2x}-1)^2 e^{-2x}} e^{2x}}{8(e^{2x}-1)} - \frac{3 \sqrt{(e^{2x}-1)^2 e^{-2x}}}{8(e^{2x}-1)} + \frac{e^{-2x} \sqrt{(e^{2x}-1)^2 e^{-2x}}}{24 e^{2x}-24}$	114

[In] `int((cosh(x)^2-1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] `1/3*(sinh(x)^2)^(1/2)*cosh(x)*(cosh(x)^2-3)/sinh(x)`

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.66

$$\int (-1 + \cosh^2(x))^{3/2} dx = \frac{1}{12} \cosh(x)^3 + \frac{1}{4} \cosh(x) \sinh(x)^2 - \frac{3}{4} \cosh(x)$$

[In] `integrate((-1+cosh(x)^2)^(3/2),x, algorithm="fricas")`

[Out] `1/12*cosh(x)^3 + 1/4*cosh(x)*sinh(x)^2 - 3/4*cosh(x)`

Sympy [F(-1)]

Timed out.

$$\int (-1 + \cosh^2(x))^{3/2} dx = \text{Timed out}$$

[In] `integrate((-1+cosh(x)**2)**(3/2),x)`

[Out] Timed out

Maxima [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 23, normalized size of antiderivative = 0.79

$$\int (-1 + \cosh^2(x))^{3/2} dx = -\frac{1}{24} e^{(3x)} + \frac{3}{8} e^{(-x)} - \frac{1}{24} e^{(-3x)} + \frac{3}{8} e^x$$

[In] integrate((-1+cosh(x)^2)^(3/2),x, algorithm="maxima")

[Out] -1/24*e^(3*x) + 3/8*e^(-x) - 1/24*e^(-3*x) + 3/8*e^x

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 66 vs. 2(21) = 42.

Time = 0.30 (sec) , antiderivative size = 66, normalized size of antiderivative = 2.28

$$\int (-1 + \cosh^2(x))^{3/2} dx = -\frac{1}{24} (9 e^{(2x)} \operatorname{sgn}(e^{(3x)} - e^x) - \operatorname{sgn}(e^{(3x)} - e^x)) e^{(-3x)} \\ + \frac{1}{24} e^{(3x)} \operatorname{sgn}(e^{(3x)} - e^x) - \frac{3}{8} e^x \operatorname{sgn}(e^{(3x)} - e^x)$$

[In] integrate((-1+cosh(x)^2)^(3/2),x, algorithm="giac")

[Out] -1/24*(9*e^(2*x)*sgn(e^(3*x) - e^x) - sgn(e^(3*x) - e^x))*e^(-3*x) + 1/24*e^(3*x)*sgn(e^(3*x) - e^x) - 3/8*e^x*sgn(e^(3*x) - e^x)

Mupad [F(-1)]

Timed out.

$$\int (-1 + \cosh^2(x))^{3/2} dx = \int (\cosh(x)^2 - 1)^{3/2} dx$$

[In] int((cosh(x)^2 - 1)^(3/2),x)

[Out] int((cosh(x)^2 - 1)^(3/2), x)

3.50 $\int (-1 - \cosh^2(x))^{3/2} dx$

Optimal result	341
Rubi [A] (verified)	341
Mathematica [A] (verified)	343
Maple [A] (verified)	343
Fricas [F]	344
Sympy [F(-1)]	344
Maxima [F]	344
Giac [F]	345
Mupad [F(-1)]	345

Optimal result

Integrand size = 12, antiderivative size = 101

$$\int (-1 - \cosh^2(x))^{3/2} dx = \frac{2i\sqrt{-1 - \cosh^2(x)}E\left(\frac{\pi}{2} + ix \mid -1\right)}{\sqrt{1 + \cosh^2(x)}} + \frac{2i\sqrt{1 + \cosh^2(x)} \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)}{3\sqrt{-1 - \cosh^2(x)}} - \frac{1}{3} \cosh(x) \sqrt{-1 - \cosh^2(x)} \sinh(x)$$

[Out] $-1/3*\cosh(x)*\sinh(x)*(-1-\cosh(x)^2)^{(1/2)}-2*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticE}(\cosh(x), I)*(-1-\cosh(x)^2)^{(1/2)}/(1+\cosh(x)^2)^{(1/2)}-2/3*(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), I)*(1+\cosh(x)^2)^{(1/2)}/(-1-\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3259, 3251, 3257, 3256, 3262, 3261}

$$\int (-1 - \cosh^2(x))^{3/2} dx = -\frac{1}{3} \sinh(x) \cosh(x) \sqrt{-\cosh^2(x) - 1} + \frac{2i\sqrt{\cosh^2(x) + 1} \operatorname{EllipticF}\left(ix + \frac{\pi}{2}, -1\right)}{3\sqrt{-\cosh^2(x) - 1}} + \frac{2i\sqrt{-\cosh^2(x) - 1}E\left(ix + \frac{\pi}{2} \mid -1\right)}{\sqrt{\cosh^2(x) + 1}}$$

[In] $\operatorname{Int}[(-1 - \operatorname{Cosh}[x]^2)^{(3/2)}, x]$

[Out] $((2*I)*\text{Sqrt}[-1 - \text{Cosh}[x]^2]*\text{EllipticE}[\text{Pi}/2 + I*x, -1])/\text{Sqrt}[1 + \text{Cosh}[x]^2]$
 $+ (((2*I)/3)*\text{Sqrt}[1 + \text{Cosh}[x]^2]*\text{EllipticF}[\text{Pi}/2 + I*x, -1])/\text{Sqrt}[-1 - \text{Cosh}[x]^2]$
 $- (\text{Cosh}[x]*\text{Sqrt}[-1 - \text{Cosh}[x]^2]*\text{Sinh}[x])/3$

Rule 3251

$\text{Int}[(A_.) + (B_.)*\sin[(e_.) + (f_.)*(x_.)]^2]/\text{Sqrt}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2], x_Symbol] \rightarrow \text{Dist}[B/b, \text{Int}[\text{Sqrt}[a + b*\sin[e + f*x]^2], x], x] + \text{Dist}[(A*b - a*B)/b, \text{Int}[1/\text{Sqrt}[a + b*\sin[e + f*x]^2], x], x] /; \text{FreeQ}[\{a, b, e, f, A, B\}, x]$

Rule 3256

$\text{Int}[\text{Sqrt}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2], x_Symbol] \rightarrow \text{Simp}[(\text{Sqrt}[a]/f)*\text{EllipticE}[e + f*x, -b/a], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{GtQ}[a, 0]$

Rule 3257

$\text{Int}[\text{Sqrt}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[a + b*\sin[e + f*x]^2]/\text{Sqrt}[1 + b*(\sin[e + f*x]^2/a)], \text{Int}[\text{Sqrt}[1 + (b*\sin[e + f*x]^2)/a], x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& !\text{GtQ}[a, 0]$

Rule 3259

$\text{Int}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2]^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[e + f*x]*\sin[e + f*x]*(a + b*\sin[e + f*x]^2)^{(p-1)}/(2*f*p), x] + \text{Dist}[1/(2*p), \text{Int}[(a + b*\sin[e + f*x]^2)^{(p-2)}*\text{Simp}[a*(b + 2*a*p) + b*(2*a + b)*(2*p - 1)*\sin[e + f*x]^2, x], x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{NeQ}[a + b, 0] \&\& \text{GtQ}[p, 1]$

Rule 3261

$\text{Int}[1/\text{Sqrt}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2], x_Symbol] \rightarrow \text{Simp}[(1/(\text{Sqrt}[a]*f))*\text{EllipticF}[e + f*x, -b/a], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{GtQ}[a, 0]$

Rule 3262

$\text{Int}[1/\text{Sqrt}[(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^2], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[1 + b*(\sin[e + f*x]^2/a)]/\text{Sqrt}[a + b*\sin[e + f*x]^2], \text{Int}[1/\text{Sqrt}[1 + (b*\sin[e + f*x]^2)/a], x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& !\text{GtQ}[a, 0]$

Rubi steps

$$\text{integral} = -\frac{1}{3} \cosh(x) \sqrt{-1 - \cosh^2(x)} \sinh(x) + \frac{1}{3} \int \frac{4 + 6 \cosh^2(x)}{\sqrt{-1 - \cosh^2(x)}} dx$$

$$\begin{aligned}
&= -\frac{1}{3} \cosh(x) \sqrt{-1 - \cosh^2(x)} \sinh(x) - \frac{2}{3} \int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx - 2 \int \sqrt{-1 - \cosh^2(x)} dx \\
&= -\frac{1}{3} \cosh(x) \sqrt{-1 - \cosh^2(x)} \sinh(x) - \frac{\left(2\sqrt{-1 - \cosh^2(x)}\right) \int \sqrt{1 + \cosh^2(x)} dx}{\sqrt{1 + \cosh^2(x)}} \\
&\quad - \frac{\left(2\sqrt{1 + \cosh^2(x)}\right) \int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx}{3\sqrt{-1 - \cosh^2(x)}} \\
&= \frac{2i\sqrt{-1 - \cosh^2(x)} E\left(\frac{\pi}{2} + ix \mid -1\right)}{\sqrt{1 + \cosh^2(x)}} + \frac{2i\sqrt{1 + \cosh^2(x)} \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)}{3\sqrt{-1 - \cosh^2(x)}} \\
&\quad - \frac{1}{3} \cosh(x) \sqrt{-1 - \cosh^2(x)} \sinh(x)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.07 (sec) , antiderivative size = 78, normalized size of antiderivative = 0.77

$$\int (-1 - \cosh^2(x))^{3/2} dx = \frac{-48i\sqrt{3 + \cosh(2x)} E\left(ix \mid \frac{1}{2}\right) + 8i\sqrt{3 + \cosh(2x)} \operatorname{EllipticF}\left(ix, \frac{1}{2}\right) + 6 \sinh(2x) + \sinh(4x)}{12\sqrt{2}\sqrt{-3 - \cosh(2x)}}$$

[In] Integrate[(-1 - Cosh[x]^2)^(3/2), x]

[Out] ((-48*I)*Sqrt[3 + Cosh[2*x]]*EllipticE[I*x, 1/2] + (8*I)*Sqrt[3 + Cosh[2*x]]*EllipticF[I*x, 1/2] + 6*Sinh[2*x] + Sinh[4*x])/(12*Sqrt[2]*Sqrt[-3 - Cosh[2*x]])

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.95

method	result
default	$-\frac{\sqrt{-(1+\cosh(x)^2)} \sinh(x)^2 \left(-\cosh(x)^5 + 2\sqrt{-\sinh(x)^2} \sqrt{1+\cosh(x)^2} \operatorname{EllipticF}(\cosh(x), i) - 6\sqrt{-\sinh(x)^2} \sqrt{1+\cosh(x)^2} \operatorname{EllipticE}(\cosh(x), i)\right)}{3\sqrt{1-\cosh(x)^4} \sinh(x) \sqrt{-1-\cosh(x)^2}}$

[In] int((-1-cosh(x)^2)^(3/2), x, method=_RETURNVERBOSE)

```
[Out] -1/3*(-(1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-cosh(x)^5+2*(-sinh(x)^2)^(1/2)*(1+cosh(x)^2)^(1/2)*EllipticF(cosh(x),I)-6*(-sinh(x)^2)^(1/2)*(1+cosh(x)^2)^(1/2)*EllipticE(cosh(x),I)+cosh(x))/(1-cosh(x)^4)^(1/2)/sinh(x)/(-1-cosh(x)^2)^(1/2)
```

Fricas [F]

$$\int (-1 - \cosh^2(x))^{3/2} dx = \int (-\cosh(x)^2 - 1)^{\frac{3}{2}} dx$$

```
[In] integrate((-1-cosh(x)^2)^(3/2),x, algorithm="fricas")
```

```
[Out] 1/24*(24*(e^(4*x) - e^(3*x))*integral(-4/3*sqrt(-e^(4*x) - 6*e^(2*x) - 1)*(5*e^(2*x) + 2*e^x + 5)/(e^(6*x) - 2*e^(5*x) + 7*e^(4*x) - 12*e^(3*x) + 7*e^(2*x) - 2*e^x + 1), x) - (e^(5*x) - e^(4*x) + 24*e^(3*x) + 24*e^(2*x) - e^x + 1)*sqrt(-e^(4*x) - 6*e^(2*x) - 1))/(e^(4*x) - e^(3*x))
```

Sympy [F(-1)]

Timed out.

$$\int (-1 - \cosh^2(x))^{3/2} dx = \text{Timed out}$$

```
[In] integrate((-1-cosh(x)**2)**(3/2),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int (-1 - \cosh^2(x))^{3/2} dx = \int (-\cosh(x)^2 - 1)^{\frac{3}{2}} dx$$

```
[In] integrate((-1-cosh(x)^2)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((-cosh(x)^2 - 1)^(3/2), x)
```


Giac [F]

$$\int (-1 - \cosh^2(x))^{3/2} dx = \int (-\cosh(x)^2 - 1)^{\frac{3}{2}} dx$$

[In] integrate((-1-cosh(x)^2)^(3/2),x, algorithm="giac")

[Out] integrate((-cosh(x)^2 - 1)^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (-1 - \cosh^2(x))^{3/2} dx = \int (-\cosh(x)^2 - 1)^{3/2} dx$$

[In] int((- cosh(x)^2 - 1)^(3/2),x)

[Out] int((- cosh(x)^2 - 1)^(3/2), x)

3.51 $\int \frac{1}{\sqrt{a+b \cosh^2(x)}} dx$

Optimal result	346
Rubi [A] (verified)	346
Mathematica [A] (verified)	347
Maple [A] (verified)	347
Fricas [B] (verification not implemented)	348
Sympy [F]	348
Maxima [F]	348
Giac [F]	349
Mupad [F(-1)]	349

Optimal result

Integrand size = 12, antiderivative size = 49

$$\int \frac{1}{\sqrt{a+b \cosh^2(x)}} dx = -\frac{i\sqrt{1+\frac{b \cosh^2(x)}{a}} \operatorname{EllipticF}\left(\frac{\pi}{2}+ix, -\frac{b}{a}\right)}{\sqrt{a+b \cosh^2(x)}}$$

[Out] $(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), (-b/a)^{(1/2)})*(1+b*\cosh(x)^2/a)^{(1/2)}/(a+b*\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.03 (sec), antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3262, 3261}

$$\int \frac{1}{\sqrt{a+b \cosh^2(x)}} dx = -\frac{i\sqrt{\frac{b \cosh^2(x)}{a}+1} \operatorname{EllipticF}\left(ix+\frac{\pi}{2}, -\frac{b}{a}\right)}{\sqrt{a+b \cosh^2(x)}}$$

[In] `Int[1/Sqrt[a + b*Cosh[x]^2], x]`

[Out] `((-I)*Sqrt[1 + (b*Cosh[x]^2)/a]*EllipticF[Pi/2 + I*x, -(b/a)]/Sqrt[a + b*Cosh[x]^2]`

Rule 3261

`Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Simp[(1/(Sqrt[a]*f))*EllipticF[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]`

Rule 3262

```
Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Dist[Sqrt[
1 + b*(Sin[e + f*x]^2/a)]/Sqrt[a + b*Sin[e + f*x]^2], Int[1/Sqrt[1 + (b*Sin
[e + f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\int \frac{1}{\sqrt{1 + \frac{b \cosh^2(x)}{a}}} dx}{\sqrt{a + b \cosh^2(x)}} \\ &= -\frac{i \sqrt{1 + \frac{b \cosh^2(x)}{a}} \text{EllipticF}\left(\frac{\pi}{2} + ix, -\frac{b}{a}\right)}{\sqrt{a + b \cosh^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 53, normalized size of antiderivative = 1.08

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = -\frac{i \sqrt{\frac{2a+b+b \cosh(2x)}{a+b}} \text{EllipticF}\left(ix, \frac{b}{a+b}\right)}{\sqrt{2a + b + b \cosh(2x)}}$$

```
[In] Integrate[1/Sqrt[a + b*Cosh[x]^2], x]
```

```
[Out] ((-I)*Sqrt[(2*a + b + b*Cosh[2*x])/(a + b)]*EllipticF[I*x, b/(a + b)]/Sqrt
[2*a + b + b*Cosh[2*x]])
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.35

method	result	size
default	$\frac{\sqrt{\frac{a+b \cosh(x)^2}{a}} \sqrt{-\sinh(x)^2} \text{EllipticF}\left(\cosh(x) \sqrt{-\frac{b}{a}}, \sqrt{-\frac{a}{b}}\right)}{\sqrt{-\frac{b}{a}} \sinh(x) \sqrt{a+b \cosh(x)^2}}$	66

```
[In] int(1/(a+b*cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] 1/(-b/a)^(1/2)*((a+b*cosh(x)^2)/a)^(1/2)*(-sinh(x)^2)^(1/2)*EllipticF(cosh(
x)*(-b/a)^(1/2), (-a/b)^(1/2))/sinh(x)/(a+b*cosh(x)^2)^(1/2)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(47) = 94$.

Time = 0.08 (sec) , antiderivative size = 132, normalized size of antiderivative = 2.69

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = \frac{2 \left(2b \sqrt{\frac{a^2 + ab}{b^2}} + 2a + b \right) \sqrt{\frac{2b \sqrt{\frac{a^2 + ab}{b^2}} - 2a - b}{b}} F\left(\arcsin\left(\sqrt{\frac{2b \sqrt{\frac{a^2 + ab}{b^2}} - 2a - b}{b}} (\cosh(x) + \sinh(x))\right)\right)}{b^{\frac{3}{2}}}$$

[In] integrate(1/(a+b*cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] $-2*(2*b*\sqrt{(a^2 + a*b)/b^2} + 2*a + b)*\sqrt{(2*b*\sqrt{(a^2 + a*b)/b^2} - 2*a - b)/b}$
 $*\text{elliptic_f}(\arcsin(\sqrt{(2*b*\sqrt{(a^2 + a*b)/b^2} - 2*a - b)/b}$
 $*(\cosh(x) + \sinh(x))), (8*a^2 + 8*a*b + b^2 + 4*(2*a*b + b^2)*\sqrt{(a^2 + a$
 $*b)/b^2))/b^2)/b^{(3/2)}$

Sympy [F]

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx$$

[In] integrate(1/(a+b*cosh(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(a + b*cosh(x)**2), x)

Maxima [F]

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{1}{\sqrt{b \cosh^2(x) + a}} dx$$

[In] integrate(1/(a+b*cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(b*cosh(x)^2 + a), x)

Giac [F]

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{1}{\sqrt{b \cosh(x)^2 + a}} dx$$

[In] integrate(1/(a+b*cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(b*cosh(x)^2 + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{1}{\sqrt{b \cosh(x)^2 + a}} dx$$

[In] int(1/(a + b*cosh(x)^2)^(1/2),x)

[Out] int(1/(a + b*cosh(x)^2)^(1/2), x)

$$3.52 \quad \int \frac{1}{\sqrt{1+\cosh^2(x)}} dx$$

Optimal result	350
Rubi [A] (verified)	350
Mathematica [A] (verified)	351
Maple [B] (verified)	351
Fricas [B] (verification not implemented)	351
Sympy [F]	352
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	352

Optimal result

Integrand size = 10, antiderivative size = 17

$$\int \frac{1}{\sqrt{1+\cosh^2(x)}} dx = -i \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)$$

[Out] $(-\sinh(x)^2)^{1/2}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), I)$

Rubi [A] (verified)

Time = 0.01 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3261}

$$\int \frac{1}{\sqrt{1+\cosh^2(x)}} dx = -i \operatorname{EllipticF}\left(ix + \frac{\pi}{2}, -1\right)$$

[In] `Int[1/Sqrt[1 + Cosh[x]^2], x]`

[Out] `(-I)*EllipticF[Pi/2 + I*x, -1]`

Rule 3261

```
Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] :> Simp[(1/(Sqrt[a]*f))*EllipticF[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]
```

Rubi steps

$$\text{integral} = -i \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.06

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx = -\frac{i \operatorname{EllipticF}\left(ix, \frac{1}{2}\right)}{\sqrt{2}}$$

[In] Integrate[1/Sqrt[1 + Cosh[x]^2],x]

[Out] ((-I)*EllipticF[I*x, 1/2])/Sqrt[2]

Maple [B] (verified)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 44 vs. 2(18) = 36.

Time = 0.14 (sec) , antiderivative size = 45, normalized size of antiderivative = 2.65

method	result	size
default	$-\frac{i \sqrt{(1 + \cosh(x)^2) \sinh(x)^2} \sqrt{-\sinh(x)^2} \operatorname{EllipticF}(i \cosh(x), i)}{\sqrt{\cosh(x)^4 - 1} \sinh(x)}$	45

[In] int(1/(1+cosh(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -I*((1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)/(cosh(x)^4-1)^(1/2)*EllipticF(I*cosh(x),I)/sinh(x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 42 vs. 2(17) = 34.

Time = 0.08 (sec) , antiderivative size = 42, normalized size of antiderivative = 2.47

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx$$

$$= -2 \left(2\sqrt{2} + 3 \right) \sqrt{2\sqrt{2} - 3} F\left(\arcsin\left(\sqrt{2\sqrt{2} - 3}(\cosh(x) + \sinh(x))\right) \mid 12\sqrt{2} + 17\right)$$

[In] integrate(1/(1+cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] -2*(2*sqrt(2) + 3)*sqrt(2*sqrt(2) - 3)*elliptic_f(arcsin(sqrt(2*sqrt(2) - 3)*(cosh(x) + sinh(x))), 12*sqrt(2) + 17)

Sympy [F]

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh^2(x) + 1}} dx$$

[In] integrate(1/(1+cosh(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(cosh(x)**2 + 1), x)

Maxima [F]

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] integrate(1/(1+cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(cosh(x)^2 + 1), x)

Giac [F]

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] integrate(1/(1+cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(cosh(x)^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] int(1/(cosh(x)^2 + 1)^(1/2),x)

[Out] int(1/(cosh(x)^2 + 1)^(1/2), x)

$$3.53 \quad \int \frac{1}{\sqrt{1-\cosh^2(x)}} dx$$

Optimal result	353
Rubi [A] (verified)	353
Mathematica [A] (verified)	354
Maple [B] (verified)	354
Fricas [B] (verification not implemented)	355
Sympy [F]	355
Maxima [C] (verification not implemented)	356
Giac [C] (verification not implemented)	356
Mupad [F(-1)]	356

Optimal result

Integrand size = 12, antiderivative size = 17

$$\int \frac{1}{\sqrt{1-\cosh^2(x)}} dx = -\frac{\operatorname{arctanh}(\cosh(x)) \sinh(x)}{\sqrt{-\sinh^2(x)}}$$

[Out] $-\operatorname{arctanh}(\cosh(x)) * \sinh(x) / (-\sinh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3255, 3286, 3855}

$$\int \frac{1}{\sqrt{1-\cosh^2(x)}} dx = -\frac{\sinh(x) \operatorname{arctanh}(\cosh(x))}{\sqrt{-\sinh^2(x)}}$$

[In] $\text{Int}[1/\text{Sqrt}[1 - \text{Cosh}[x]^2], x]$

[Out] $-\left(\text{ArcTanh}[\text{Cosh}[x]] * \text{Sinh}[x]\right) / \text{Sqrt}[-\text{Sinh}[x]^2]$

Rule 3255

$\text{Int}[(u_.) * ((a_.) + (b_.) * \sin[(e_.) + (f_.) * (x_)]^2)^{(p_.)}, x_Symbol] \rightarrow \text{Int}[\text{ActivateTrig}[u * (a * \cos[e + f * x]^2)^p], x] /;$ $\text{FreeQ}\{a, b, e, f, p, x\}$ && $\text{EqQ}[a + b, 0]$

Rule 3286

$\text{Int}[(u_.) * ((b_.) * \sin[(e_.) + (f_.) * (x_)]^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{With}\{\{ff = \text{FreeFactors}[\sin[e + f * x], x]\}, \text{Dist}[(b * ff^n)^{\text{IntPart}[p]} * ((b * \sin[e + f * x])^n)^{\text{FracPart}[p]}\}$

```

n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig]])

```

Rule 3855

```

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{1}{\sqrt{-\sinh^2(x)}} dx \\
&= \frac{\sinh(x) \int \operatorname{csch}(x) dx}{\sqrt{-\sinh^2(x)}} \\
&= -\frac{\operatorname{arctanh}(\cosh(x)) \sinh(x)}{\sqrt{-\sinh^2(x)}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.03 (sec) , antiderivative size = 30, normalized size of antiderivative = 1.76

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = \frac{(-\log(\cosh(\frac{x}{2})) + \log(\sinh(\frac{x}{2}))) \sinh(x)}{\sqrt{-\sinh^2(x)}}$$

```
[In] Integrate[1/Sqrt[1 - Cosh[x]^2],x]
```

```
[Out] ((-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x])/Sqrt[-Sinh[x]^2]
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(15) = 30.

Time = 0.12 (sec) , antiderivative size = 34, normalized size of antiderivative = 2.00

method	result	size
default	$-\frac{\sinh(x)\sqrt{-\cosh(x)^2} \arctan\left(\frac{1}{\sqrt{-\cosh(x)^2}}\right)}{\cosh(x)\sqrt{-\sinh(x)^2}}$	34
risch	$-\frac{e^{-x}(e^{2x}-1)\ln(e^x+1)}{\sqrt{-(e^{2x}-1)^2e^{-2x}}} + \frac{e^{-x}(e^{2x}-1)\ln(e^x-1)}{\sqrt{-(e^{2x}-1)^2e^{-2x}}}$	67

[In] `int(1/(1-cosh(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-sinh(x)*(-cosh(x)^2)^(1/2)*arctan(1/(-cosh(x)^2)^(1/2))/cosh(x)/(-sinh(x)^2)^(1/2)`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(15) = 30$.

Time = 0.25 (sec) , antiderivative size = 49, normalized size of antiderivative = 2.88

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = 2 \arctan\left(\frac{\sqrt{-(e^{4x} - 2e^{2x} + 1)e^{-2x}}e^x}{\cosh(x)e^{2x} + (e^{2x} - 1)\sinh(x) - \cosh(x)}\right)$$

[In] `integrate(1/(1-cosh(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `2*arctan(sqrt(-(e^(4*x) - 2*e^(2*x) + 1)*e^(-2*x))*e^x/(cosh(x)*e^(2*x) + (e^(2*x) - 1)*sinh(x) - cosh(x)))`

Sympy [F]

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx$$

[In] `integrate(1/(1-cosh(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(1 - cosh(x)**2), x)`

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.12

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = -i \log(e^{(-x)} + 1) + i \log(e^{(-x)} - 1)$$

[In] integrate(1/(1-cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] -I*log(e^(-x) + 1) + I*log(e^(-x) - 1)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 40, normalized size of antiderivative = 2.35

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = -\frac{i \log(e^x + 1)}{\operatorname{sgn}(-e^{(3x)} + e^x)} + \frac{i \log(|e^x - 1|)}{\operatorname{sgn}(-e^{(3x)} + e^x)}$$

[In] integrate(1/(1-cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] -I*log(e^x + 1)/sgn(-e^(3*x) + e^x) + I*log(abs(e^x - 1))/sgn(-e^(3*x) + e^x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{1 - \cosh(x)^2}} dx$$

[In] int(1/(1 - cosh(x)^2)^(1/2),x)

[Out] int(1/(1 - cosh(x)^2)^(1/2), x)

$$3.54 \quad \int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx$$

Optimal result	357
Rubi [A] (verified)	357
Mathematica [A] (verified)	358
Maple [A] (verified)	358
Fricas [A] (verification not implemented)	359
Sympy [F]	359
Maxima [A] (verification not implemented)	359
Giac [B] (verification not implemented)	360
Mupad [F(-1)]	360

Optimal result

Integrand size = 10, antiderivative size = 15

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = -\frac{\operatorname{arctanh}(\cosh(x)) \sinh(x)}{\sqrt{\sinh^2(x)}}$$

[Out] $-\operatorname{arctanh}(\cosh(x)) * \sinh(x) / (\sinh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3255, 3286, 3855}

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = -\frac{\sinh(x) \operatorname{arctanh}(\cosh(x))}{\sqrt{\sinh^2(x)}}$$

[In] `Int[1/Sqrt[-1 + Cosh[x]^2], x]`

[Out] `-((ArcTanh[Cosh[x]]*Sinh[x])/Sqrt[Sinh[x]^2])`

Rule 3255

`Int[(u_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Int[ActivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0]`

Rule 3286

`Int[(u_)*((b_)*sin[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[(b*ff^n)^IntPart[p]*((b*Sinh[e + f*x]^`

```
n)^FracPart[p]/(Sin[e + f*x]/ff)^(n*FracPart[p])), Int[ActivateTrig[u]*(Sin
[e + f*x]/ff)^(n*p), x], x] /; FreeQ[{b, e, f, n, p}, x] && !IntegerQ[p]
&& IntegerQ[n] && (EqQ[u, 1] || MatchQ[u, ((d_.)*(trig_)[e + f*x])^(m_.) /;
FreeQ[{d, m}, x] && MemberQ[{sin, cos, tan, cot, sec, csc}, trig])]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \frac{1}{\sqrt{\sinh^2(x)}} dx \\ &= \frac{\sinh(x) \int \operatorname{csch}(x) dx}{\sqrt{\sinh^2(x)}} \\ &= -\frac{\operatorname{arctanh}(\cosh(x)) \sinh(x)}{\sqrt{\sinh^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.87

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = \frac{(-\log(\cosh(\frac{x}{2})) + \log(\sinh(\frac{x}{2}))) \sinh(x)}{\sqrt{\sinh^2(x)}}$$

```
[In] Integrate[1/Sqrt[-1 + Cosh[x]^2], x]
```

```
[Out] ((-Log[Cosh[x/2]] + Log[Sinh[x/2]])*Sinh[x])/Sqrt[Sinh[x]^2]
```

Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.07

method	result	size
default	$-\frac{\sqrt{-\frac{1}{2} + \frac{\cosh(2x)}{2}} \operatorname{arctanh}(\cosh(x))}{\sinh(x)}$	16
risch	$-\frac{e^{-x}(e^{2x}-1)\ln(e^x+1)}{\sqrt{(e^{2x}-1)^2e^{-2x}}} + \frac{e^{-x}(e^{2x}-1)\ln(e^x-1)}{\sqrt{(e^{2x}-1)^2e^{-2x}}}$	65

[In] `int(1/(cosh(x)^2-1)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-(sinh(x)^2)^(1/2)*arctanh(cosh(x))/sinh(x)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = -\log(\cosh(x) + \sinh(x) + 1) + \log(\cosh(x) + \sinh(x) - 1)$$

[In] `integrate(1/(-1+cosh(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `-log(cosh(x) + sinh(x) + 1) + log(cosh(x) + sinh(x) - 1)`

Sympy [F]

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh^2(x) - 1}} dx$$

[In] `integrate(1/(-1+cosh(x)**2)**(1/2),x)`

[Out] `Integral(1/sqrt(cosh(x)**2 - 1), x)`

Maxima [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 17, normalized size of antiderivative = 1.13

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = \log(e^{-x} + 1) - \log(e^{-x} - 1)$$

[In] `integrate(1/(-1+cosh(x)^2)^(1/2),x, algorithm="maxima")`

[Out] `log(e^(-x) + 1) - log(e^(-x) - 1)`

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(13) = 26.

Time = 0.30 (sec) , antiderivative size = 39, normalized size of antiderivative = 2.60

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = -\frac{\log(e^x + 1)}{\operatorname{sgn}(e^{3x} - e^x)} + \frac{\log(|e^x - 1|)}{\operatorname{sgn}(e^{3x} - e^x)}$$

[In] integrate(1/(-1+cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] -log(e^x + 1)/sgn(e^(3*x) - e^x) + log(abs(e^x - 1))/sgn(e^(3*x) - e^x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-1 + \cosh^2(x)}} dx = \int \frac{1}{\sqrt{\cosh(x)^2 - 1}} dx$$

[In] int(1/(cosh(x)^2 - 1)^(1/2),x)

[Out] int(1/(cosh(x)^2 - 1)^(1/2), x)

$$3.55 \quad \int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx$$

Optimal result	361
Rubi [A] (verified)	361
Mathematica [A] (verified)	362
Maple [A] (verified)	362
Fricas [A] (verification not implemented)	363
Sympy [F]	363
Maxima [F]	363
Giac [F]	364
Mupad [F(-1)]	364

Optimal result

Integrand size = 12, antiderivative size = 39

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = -\frac{i\sqrt{1 + \cosh^2(x)} \operatorname{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)}{\sqrt{-1 - \cosh^2(x)}}$$

[Out] $(-\sinh(x)^2)^{(1/2)}/\sinh(x)*\operatorname{EllipticF}(\cosh(x), I)*(1+\cosh(x)^2)^{(1/2)/(-1-\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3262, 3261}

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = -\frac{i\sqrt{\cosh^2(x) + 1} \operatorname{EllipticF}\left(ix + \frac{\pi}{2}, -1\right)}{\sqrt{-\cosh^2(x) - 1}}$$

[In] `Int[1/Sqrt[-1 - Cosh[x]^2], x]`

[Out] `((-I)*Sqrt[1 + Cosh[x]^2]*EllipticF[Pi/2 + I*x, -1])/Sqrt[-1 - Cosh[x]^2]`

Rule 3261

`Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Simp[(1/(Sqrt[a]*f))*EllipticF[e + f*x, -b/a], x] /; FreeQ[{a, b, e, f}, x] && GtQ[a, 0]`

Rule 3262

```
Int[1/Sqrt[(a_) + (b_)*sin[(e_) + (f_)*(x_)]^2], x_Symbol] := Dist[Sqrt[
1 + b*(Sin[e + f*x]^2/a)]/Sqrt[a + b*Sin[e + f*x]^2], Int[1/Sqrt[1 + (b*Sin
[e + f*x]^2)/a], x], x] /; FreeQ[{a, b, e, f}, x] && !GtQ[a, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\sqrt{1 + \cosh^2(x)} \int \frac{1}{\sqrt{1 + \cosh^2(x)}} dx}{\sqrt{-1 - \cosh^2(x)}} \\ &= -\frac{i\sqrt{1 + \cosh^2(x)} \text{EllipticF}\left(\frac{\pi}{2} + ix, -1\right)}{\sqrt{-1 - \cosh^2(x)}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.04 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.03

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = -\frac{i\sqrt{3 + \cosh(2x)} \text{EllipticF}\left(ix, \frac{1}{2}\right)}{\sqrt{2}\sqrt{-3 - \cosh(2x)}}$$

[In] Integrate[1/Sqrt[-1 - Cosh[x]^2],x]

[Out] ((-I)*Sqrt[3 + Cosh[2*x]]*EllipticF[I*x, 1/2])/(Sqrt[2]*Sqrt[-3 - Cosh[2*x]])

Maple [A] (verified)

Time = 0.17 (sec) , antiderivative size = 61, normalized size of antiderivative = 1.56

method	result	size
default	$\frac{\sqrt{-(1+\cosh(x)^2)} \sinh(x)^2 \sqrt{-\sinh(x)^2} \sqrt{1+\cosh(x)^2} \text{EllipticF}(\cosh(x), i)}{\sqrt{1-\cosh(x)^4} \sinh(x) \sqrt{-1-\cosh(x)^2}}$	61

[In] int(1/(-1-cosh(x)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] (-(1+cosh(x)^2)*sinh(x)^2)^(1/2)*(-sinh(x)^2)^(1/2)*(1+cosh(x)^2)^(1/2)/(1-cosh(x)^4)^(1/2)*EllipticF(cosh(x),I)/sinh(x)/(-1-cosh(x)^2)^(1/2)

Fricas [A] (verification not implemented)

none

Time = 0.07 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = -2\sqrt{2\sqrt{2} - 3}(-2i\sqrt{2} - 3i)F(\arcsin(\sqrt{2\sqrt{2} - 3}e^x) | 12\sqrt{2} + 17)$$

[In] integrate(1/(-1-cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] -2*sqrt(2*sqrt(2) - 3)*(-2*I*sqrt(2) - 3*I)*elliptic_f(arcsin(sqrt(2*sqrt(2) - 3)*e^x), 12*sqrt(2) + 17)

Sympy [F]

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{-\cosh^2(x) - 1}} dx$$

[In] integrate(1/(-1-cosh(x)**2)**(1/2),x)

[Out] Integral(1/sqrt(-cosh(x)**2 - 1), x)

Maxima [F]

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{-\cosh(x)^2 - 1}} dx$$

[In] integrate(1/(-1-cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(-cosh(x)^2 - 1), x)

Giac [F]

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{-\cosh(x)^2 - 1}} dx$$

[In] integrate(1/(-1-cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(-cosh(x)^2 - 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{\sqrt{-1 - \cosh^2(x)}} dx = \int \frac{1}{\sqrt{-\cosh(x)^2 - 1}} dx$$

[In] int(1/(-cosh(x)^2 - 1)^(1/2),x)

[Out] int(1/(-cosh(x)^2 - 1)^(1/2), x)

3.56 $\int \frac{1}{a+b \cosh^3(x)} dx$

Optimal result	365
Rubi [A] (verified)	366
Mathematica [F(-1)]	368
Maple [C] (verified)	368
Fricas [C] (verification not implemented)	368
Sympy [F(-1)]	369
Maxima [F]	369
Giac [F]	369
Mupad [B] (verification not implemented)	369

Optimal result

Integrand size = 10, antiderivative size = 288

$$\int \frac{1}{a+b \cosh^3(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a}-\sqrt[3]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a}+\sqrt[3]{b}}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a}-\sqrt[3]{b}} \sqrt{\sqrt[3]{a}+\sqrt[3]{b}}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a}+\sqrt[3]{-1}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a}-\sqrt[3]{-1}} \sqrt[3]{b}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a}-\sqrt[3]{-1}} \sqrt[3]{b} \sqrt{\sqrt[3]{a}+\sqrt[3]{-1}} \sqrt[3]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a}-(-1)^{2/3}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a}+(-1)^{2/3}} \sqrt[3]{b}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a}-(-1)^{2/3}} \sqrt[3]{b} \sqrt{\sqrt[3]{a}+(-1)^{2/3}} \sqrt[3]{b}}$$

```
[Out] 2/3*arctanh((a^(1/3)-b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)+b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-b^(1/3))^(1/2)/(a^(1/3)+b^(1/3))^(1/2)+2/3*arctanh((a^(1/3)+(-1)^(1/3)*b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)-(-1)^(1/3)*b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-(-1)^(1/3)*b^(1/3))^(1/2)/(a^(1/3)+(-1)^(1/3)*b^(1/3))^(1/2)+2/3*arctanh((a^(1/3)-(-1)^(2/3)*b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)+(-1)^(2/3)*b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-(-1)^(2/3)*b^(1/3))^(1/2)/(a^(1/3)+(-1)^(2/3)*b^(1/3))^(1/2)
```

Rubi [A] (verified)

Time = 0.39 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3292, 2738, 214}

$$\int \frac{1}{a + b \cosh^3(x)} dx = \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b}} + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b}}$$

[In] Int[(a + b*Cosh[x]^3)^(-1), x]

[Out] (2*ArcTanh[(Sqrt[a^(1/3) - b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) + b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - b^(1/3)]*Sqrt[a^(1/3) + b^(1/3)]) + (2*ArcTanh[(Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)]*Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)]) + (2*ArcTanh[(Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)]*Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2738

Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3292

Int[((a_) + (b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_)^(p_), x_Symbol] := Int[ExpandTrig[(a + b*(c*sin[e + f*x])^n)^p, x], x] /; FreeQ[{a, b, c, e, f}

, n}, x] && (IGtQ[p, 0] || (EqQ[p, -1] && IntegerQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(-\frac{1}{3a^{2/3} \left(-\sqrt[3]{a} - \sqrt[3]{b} \cosh(x) \right)} - \frac{1}{3a^{2/3} \left(-\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} \cosh(x) \right)} \right. \\
 &\quad \left. - \frac{1}{3a^{2/3} \left(-\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} \cosh(x) \right)} \right) dx \\
 &= -\frac{\int \frac{1}{-\sqrt[3]{a} - \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} - \frac{\int \frac{1}{-\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} - \frac{\int \frac{1}{-\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} \\
 &= -\frac{2\text{Subst} \left(\int \frac{1}{-\sqrt[3]{a} - \sqrt[3]{b} - \left(-\sqrt[3]{a} + \sqrt[3]{b} \right) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &\quad - \frac{2\text{Subst} \left(\int \frac{1}{-\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} - \left(-\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b} \right) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &\quad - \frac{2\text{Subst} \left(\int \frac{1}{-\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} - \left(-\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b} \right) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &= \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}}} \\
 &\quad + \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}
 \end{aligned}$$

Mathematica [F(-1)]

Timed out.

$$\int \frac{1}{a + b \cosh^3(x)} dx = \$Aborted$$

[In] Integrate[(a + b*Cosh[x]^3)^(-1),x]

[Out] \$Aborted

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.55 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.35

method	result
default	$\frac{\sum_{\substack{R=\text{RootOf}((a-b)Z^6+(-3a-3b)Z^4+(3a-3b)Z^2-a-b)} \left(\frac{(-R^4+2R^2-1) \ln(\tanh(\frac{x}{2})-R)}{R^5 a - R^5 b - 2R^3 a - 2R^3 b + R a - R b} \right)}{3}$
risch	$\sum_{\substack{R=\text{RootOf}(-1+(729a^6-729a^4b^2)Z^6-243a^4Z^4+27a^2Z^2)}} -R \ln \left(e^x + \left(\frac{486a^6}{b} - 486a^4b \right) R^5 + \left(-\frac{81a^5}{b} + 8 \right) R^3 \right)$

[In] int(1/(a+b*cosh(x)^3),x,method=_RETURNVERBOSE)

[Out] 1/3*sum((-R^4+2*_R^2-1)/(_R^5*a-_R^5*b-2*_R^3*a-2*_R^3*b+_R*a-_R*b)*ln(tanh(1/2*x)-_R),_R=RootOf((a-b)*_Z^6+(-3*a-3*b)*_Z^4+(3*a-3*b)*_Z^2-a-b))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.98 (sec) , antiderivative size = 18612, normalized size of antiderivative = 64.62

$$\int \frac{1}{a + b \cosh^3(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cosh(x)^3),x, algorithm="fricas")

[Out] Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{a + b \cosh^3(x)} dx = \text{Timed out}$$

[In] integrate(1/(a+b*cosh(x)**3),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{a + b \cosh^3(x)} dx = \int \frac{1}{b \cosh(x)^3 + a} dx$$

[In] integrate(1/(a+b*cosh(x)^3),x, algorithm="maxima")

[Out] integrate(1/(b*cosh(x)^3 + a), x)

Giac [F]

$$\int \frac{1}{a + b \cosh^3(x)} dx = \int \frac{1}{b \cosh(x)^3 + a} dx$$

[In] integrate(1/(a+b*cosh(x)^3),x, algorithm="giac")

[Out] integrate(1/(b*cosh(x)^3 + a), x)

Mupad [B] (verification not implemented)

Time = 7.03 (sec) , antiderivative size = 633, normalized size of antiderivative = 2.20

$$\int \frac{1}{a + b \cosh^3(x)} dx$$

$$= \sum_{k=1}^6 \ln \left(- \frac{\left(-4 e^x + \text{root}(729 a^4 b^2 d^6 - 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k) b + \text{root}(729 a^4 b^2 d^6 - 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k) \right)}{- 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k) \right)$$

[In] int(1/(a + b*cosh(x)^3),x)

[Out] symsum(log(-(24576*(root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)*b - 4*exp(x) + 54*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 24

$$\begin{aligned}
& 3a^4d^4 - 27a^2d^2 + 1, d, k)^3a^2b + 108\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^4a^3b + 81\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^5a^4b + 24\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^2a^2\exp(x) + 216\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^3a^3\exp(x) + 108\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^4a^4\exp(x) - 324\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^5a^5\exp(x) + 12\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^2a*b - 20\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)*a*\exp(x) + 27\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^4a^2b^2*\exp(x) + 405\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k)^5a^3b^2*\exp(x))/b^5*\text{root}(729a^4b^2d^6 - 729a^6d^6 + 243a^4d^4 - 27a^2d^2 + 1, d, k), k, 1, 6)
\end{aligned}$$

$$3.57 \quad \int \frac{1}{a-b \cosh^3(x)} dx$$

Optimal result	371
Rubi [A] (verified)	372
Mathematica [C] (verified)	374
Maple [C] (verified)	374
Fricas [C] (verification not implemented)	375
Sympy [F(-1)]	375
Maxima [F]	375
Giac [F]	375
Mupad [B] (verification not implemented)	376

Optimal result

Integrand size = 11, antiderivative size = 288

$$\int \frac{1}{a-b \cosh^3(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b}}\right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b}}$$

```
[Out] 2/3*arctanh((a^(1/3)+b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)-b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-b^(1/3))^(1/2)/(a^(1/3)+b^(1/3))^(1/2)+2/3*arctanh((a^(1/3)-(-1)^(1/3)*b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)+(-1)^(1/3)*b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-(-1)^(1/3)*b^(1/3))^(1/2)/(a^(1/3)+(-1)^(1/3)*b^(1/3))^(1/2)+2/3*arctanh((a^(1/3)+(-1)^(2/3)*b^(1/3))^(1/2)*tanh(1/2*x)/(a^(1/3)-(-1)^(2/3)*b^(1/3))^(1/2))/a^(2/3)/(a^(1/3)-(-1)^(2/3)*b^(1/3))^(1/2)/(a^(1/3)+(-1)^(2/3)*b^(1/3))^(1/2)
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 288, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3292, 2738, 214}

$$\int \frac{1}{a - b \cosh^3(x)} dx = \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1}} \sqrt[3]{b}} + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - (-1)^{2/3}} \sqrt[3]{b} \sqrt{\sqrt[3]{a} + (-1)^{2/3}} \sqrt[3]{b}}$$

[In] Int[(a - b*Cosh[x]^3)^(-1), x]

[Out] (2*ArcTanh[(Sqrt[a^(1/3) + b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) - b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - b^(1/3)]*Sqrt[a^(1/3) + b^(1/3)]) + (2*ArcTanh[(Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)]*Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)]) + (2*ArcTanh[(Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)]*Tanh[x/2])/Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)])]/(3*a^(2/3)*Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)]*Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2738

Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3292

Int[((a_) + (b_.)*((c_.)*sin[e_.] + (f_.)*(x_)))^(n_)^(p_), x_Symbol] := Int[ExpandTrig[(a + b*(c*sin[e + f*x])^n)^p, x], x] /; FreeQ[{a, b, c, e, f}

, n}, x] && (IGtQ[p, 0] || (EqQ[p, -1] && IntegerQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{3a^{2/3} (\sqrt[3]{a} - \sqrt[3]{b} \cosh(x))} + \frac{1}{3a^{2/3} (\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} \cosh(x))} \right. \\
 &\quad \left. + \frac{1}{3a^{2/3} (\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} \cosh(x))} \right) dx \\
 &= \frac{\int \frac{1}{\sqrt[3]{a} - \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} + \frac{\int \frac{1}{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} \cosh(x)} dx}{3a^{2/3}} \\
 &= \frac{2\text{Subst} \left(\int \frac{1}{\sqrt[3]{a} - \sqrt[3]{b} - (\sqrt[3]{a} + \sqrt[3]{b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &\quad + \frac{2\text{Subst} \left(\int \frac{1}{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b} - (\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &\quad + \frac{2\text{Subst} \left(\int \frac{1}{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b} - (\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right)}{3a^{2/3}} \\
 &= \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}}} \\
 &\quad + \frac{2\arctanh \left(\frac{\sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}} \tanh \left(\frac{x}{2} \right)}{\sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}}} \right)}{3a^{2/3} \sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}} \sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.06 (sec) , antiderivative size = 105, normalized size of antiderivative = 0.36

$$\int \frac{1}{a - b \cosh^3(x)} dx$$

$$= -\frac{2}{3} \text{RootSum} \left[b + 3b\#1^2 - 8a\#1^3 + 3b\#1^4 \right. \\ \left. + b\#1^6 \&, \frac{x\#1 + 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) \#1}{b - 4a\#1 + 2b\#1^2 + b\#1^4} \& \right]$$

[In] Integrate[(a - b*Cosh[x]^3)^(-1),x]

[Out] (-2*RootSum[b + 3*b*#1^2 - 8*a*#1^3 + 3*b*#1^4 + b*#1^6 & , (x*#1 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1)/(b - 4*a*#1 + 2*b*#1^2 + b*#1^4) &])/3

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.53 (sec) , antiderivative size = 94, normalized size of antiderivative = 0.33

method	result
default	$\frac{\sum_{-R=\text{RootOf}((a+b)Z^6+(-3a+3b)Z^4+(3a+3b)Z^2-a+b)} \left(\frac{(-R^4+2R^2-1) \ln(\tanh(\frac{x}{2})-R)}{R^5 a + R^5 b - 2R^3 a + 2R^3 b + R a + R b} \right)}{3}$
risch	$\sum_{-R=\text{RootOf}(-1+(729a^6-729a^4b^2)Z^6-243a^4Z^4+27a^2Z^2)} -R \ln \left(e^x + \left(-\frac{486a^6}{b} + 486a^4b \right) -R^5 + \left(\frac{81a^5}{b} - 8 \right) \right)$

[In] int(1/(a-b*cosh(x)^3),x,method=_RETURNVERBOSE)

[Out] 1/3*sum((-R^4+2*_R^2-1)/(_R^5*a+_R^5*b-2*_R^3*a+2*_R^3*b+_R*a+_R*b)*ln(tanh(1/2*x)-_R),_R=RootOf((a+b)*_Z^6+(-3*a+3*b)*_Z^4+(3*a+3*b)*_Z^2-a+b))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.98 (sec) , antiderivative size = 18612, normalized size of antiderivative = 64.62

$$\int \frac{1}{a - b \cosh^3(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a-b*cosh(x)^3),x, algorithm="fricas")

[Out] Too large to include

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{a - b \cosh^3(x)} dx = \text{Timed out}$$

[In] integrate(1/(a-b*cosh(x)**3),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{a - b \cosh^3(x)} dx = \int -\frac{1}{b \cosh(x)^3 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^3),x, algorithm="maxima")

[Out] -integrate(1/(b*cosh(x)^3 - a), x)

Giac [F]

$$\int \frac{1}{a - b \cosh^3(x)} dx = \int -\frac{1}{b \cosh(x)^3 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^3),x, algorithm="giac")

[Out] integrate(-1/(b*cosh(x)^3 - a), x)

Mupad [B] (verification not implemented)

Time = 6.81 (sec) , antiderivative size = 633, normalized size of antiderivative = 2.20

$$\int \frac{1}{a - b \cosh^3(x)} dx$$

$$= \sum_{k=1}^6 \ln \left(- \frac{\left(4e^x + \text{root}(729 a^4 b^2 d^6 - 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k) b + \text{root}(729 a^4 b^2 d^6 - 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k) \right)}{- 729 a^6 d^6 + 243 a^4 d^4 - 27 a^2 d^2 + 1, d, k} \right)$$

`[In] int(1/(a - b*cosh(x)^3),x)`

```
[Out] symsum(log(-(24576*(4*exp(x) + root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k))*b + 54*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^3*a^2*b + 108*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^4*a^3*b + 81*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^5*a^4*b - 24*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^2*a^2*exp(x) - 216*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^3*a^3*exp(x) - 108*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^4*a^4*exp(x) + 324*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^5*a^5*exp(x) + 12*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^2*a*b + 20*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)*a*exp(x) - 27*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^4*a^2*b^2*exp(x) - 405*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k)^5*a^3*b^2*exp(x)))/b^5)*root(729*a^4*b^2*d^6 - 729*a^6*d^6 + 243*a^4*d^4 - 27*a^2*d^2 + 1, d, k), k, 1, 6)
```


3.58 $\int \frac{1}{1+\cosh^3(x)} dx$

Optimal result	377
Rubi [A] (verified)	377
Mathematica [C] (verified)	379
Maple [C] (verified)	379
Fricas [C] (verification not implemented)	380
Sympy [B] (verification not implemented)	380
Maxima [F]	381
Giac [B] (verification not implemented)	381
Mupad [B] (verification not implemented)	382

Optimal result

Integrand size = 8, antiderivative size = 91

$$\int \frac{1}{1+\cosh^3(x)} dx = -\frac{2\sqrt[4]{-\frac{1}{3}} \arctan\left(\left(-1\right)^{3/4} \sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3\left(1-\sqrt[3]{-1}\right)} - \frac{2\sqrt[4]{-\frac{1}{3}} \operatorname{arctanh}\left(\left(-1\right)^{3/4} \sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3\left(1+\left(-1\right)^{2/3}\right)} + \frac{\sinh(x)}{3\left(1+\cosh(x)\right)}$$

[Out] $-2/9*(-1)^{(1/4)}*3^{(3/4)}*\arctan\left(\left(-1\right)^{(3/4)}*3^{(1/4)}*\tanh(1/2*x)\right)/\left(1-\left(-1\right)^{(1/3)}\right)-2/9*(-1)^{(1/4)}*3^{(3/4)}*\operatorname{arctanh}\left(\left(-1\right)^{(3/4)}*3^{(1/4)}*\tanh(1/2*x)\right)/\left(1+\left(-1\right)^{(2/3)}\right)+1/3*\sinh(x)/(1+\cosh(x))$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3292, 2727, 2738, 211, 214}

$$\int \frac{1}{1+\cosh^3(x)} dx = -\frac{2\sqrt[4]{-\frac{1}{3}} \arctan\left(\left(-1\right)^{3/4} \sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3\left(1-\sqrt[3]{-1}\right)} - \frac{2\sqrt[4]{-\frac{1}{3}} \operatorname{arctanh}\left(\left(-1\right)^{3/4} \sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3\left(1+\left(-1\right)^{2/3}\right)} + \frac{\sinh(x)}{3\left(\cosh(x)+1\right)}$$

[In] $\text{Int}[(1 + \text{Cosh}[x]^3)^{-1}, x]$

[Out] $(-2*(-1/3)^{1/4}*\text{ArcTan}[(-1)^{3/4}*3^{1/4}*\text{Tanh}[x/2]])/(3*(1 - (-1)^{1/3}))$
 $- (2*(-1/3)^{1/4}*\text{ArcTanh}[(-1)^{3/4}*3^{1/4}*\text{Tanh}[x/2]])/(3*(1 + (-1)^{2/3}))$
 $+ \text{Sinh}[x]/(3*(1 + \text{Cosh}[x]))$

Rule 211

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b]$

Rule 214

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{NegQ}[a/b]$

Rule 2727

$\text{Int}[(a_ + (b_)*\sin[(c_ + (d_)*(x_))])^{-1}, x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/(d*(b + a*\sin[c + d*x])), x] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{EqQ}[a^2 - b^2, 0]$

Rule 2738

$\text{Int}[(a_ + (b_)*\sin[\text{Pi}/2 + (c_ + (d_)*(x_))])^{-1}, x_Symbol] \rightarrow \text{With}[e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x], \text{Dist}[2*(e/d), \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x]] /; \text{FreeQ}\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[a^2 - b^2, 0]$

Rule 3292

$\text{Int}[(a_ + (b_)*((c_)*\sin[(e_ + (f_)*(x_))])^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrig}[(a + b*(c*\sin[e + f*x])^n)^p, x], x] /; \text{FreeQ}\{a, b, c, e, f, n\}, x] \ \&\& \ (\text{IGtQ}[p, 0] \ || \ (\text{EqQ}[p, -1] \ \&\& \ \text{IntegerQ}[n]))$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(-\frac{1}{3(-1 - \cosh(x))} - \frac{1}{3(-1 + \sqrt[3]{-1} \cosh(x))} - \frac{1}{3(-1 - (-1)^{2/3} \cosh(x))} \right) dx \\ &= -\left(\frac{1}{3} \int \frac{1}{-1 - \cosh(x)} dx \right) - \frac{1}{3} \int \frac{1}{-1 + \sqrt[3]{-1} \cosh(x)} dx - \frac{1}{3} \int \frac{1}{-1 - (-1)^{2/3} \cosh(x)} dx \\ &= \frac{\sinh(x)}{3(1 + \cosh(x))} - \frac{2}{3} \text{Subst} \left(\int \frac{1}{-1 + \sqrt[3]{-1} - (-1 - \sqrt[3]{-1}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\ &\quad - \frac{2}{3} \text{Subst} \left(\int \frac{1}{-1 - (-1)^{2/3} - (-1 + (-1)^{2/3}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \end{aligned}$$

$$= -\frac{2\sqrt[4]{-\frac{1}{3}} \arctan\left((-1)^{3/4}\sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3(1-\sqrt[3]{-1})} - \frac{2\sqrt[4]{-\frac{1}{3}} \operatorname{arctanh}\left((-1)^{3/4}\sqrt[4]{3} \tanh\left(\frac{x}{2}\right)\right)}{3(1+(-1)^{2/3})} + \frac{\sinh(x)}{3(1+\cosh(x))}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.50 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.46

$$\int \frac{1}{1+\cosh^3(x)} dx = \frac{1}{18} \left(-\sqrt{6+2i\sqrt{3}}(3i+\sqrt{3}) \arctan\left(\frac{(3+i\sqrt{3})\tanh\left(\frac{x}{2}\right)}{\sqrt{6-2i\sqrt{3}}}\right) - \sqrt{6-2i\sqrt{3}}(-3i+\sqrt{3}) \arctan\left(\frac{(3-i\sqrt{3})\tanh\left(\frac{x}{2}\right)}{\sqrt{6+2i\sqrt{3}}}\right) + 6 \tanh\left(\frac{x}{2}\right) \right)$$

[In] Integrate[(1 + Cosh[x]^3)^(-1),x]

[Out] $(-\sqrt{6+(2I)\sqrt{3}}(3I+\sqrt{3})\operatorname{ArcTan}[\frac{(3+I\sqrt{3})\operatorname{Tanh}[x/2]}{\sqrt{6-(2I)\sqrt{3}}}] - \sqrt{6-(2I)\sqrt{3}}(-3I+\sqrt{3})\operatorname{ArcTan}[\frac{(3-I\sqrt{3})\operatorname{Tanh}[x/2]}{\sqrt{6+(2I)\sqrt{3}}}] + 6\operatorname{Tanh}[x/2])/18$

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.51

method	result
risch	$-\frac{2}{3(e^x+1)} + \left(\sum_{R=\operatorname{RootOf}(243_Z^4-27_Z^2+1)} -R \ln(-162_R^3+27_R^2+9_R+e^x-2) \right)$
default	$\frac{\tanh(\frac{x}{2})}{3} + \frac{3^{\frac{3}{4}}\sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})^2 + \frac{3^{\frac{3}{4}}\tanh(\frac{x}{2})\sqrt{2}}{3} + \frac{\sqrt{3}}{3}}{\tanh(\frac{x}{2})^2 - \frac{3^{\frac{3}{4}}\tanh(\frac{x}{2})\sqrt{2}}{3} + \frac{\sqrt{3}}{3}}\right) + 2 \arctan(\sqrt{2}3^{\frac{1}{4}}\tanh(\frac{x}{2})+1) + 2 \arctan(\sqrt{2}3^{\frac{1}{4}}\tanh(\frac{x}{2})-1) \right)}{36} - 3^{\frac{1}{4}}$

[In] int(1/(1+cosh(x)^3),x,method=_RETURNVERBOSE)

[Out] $-2/3/(\exp(x)+1)+\text{sum}(_R*\ln(-162*_R^3+27*_R^2+9*_R+\exp(x)-2), _R=\text{RootOf}(243*_Z^4-27*_Z^2+1))$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 226, normalized size of antiderivative = 2.48

$$\int \frac{1}{1 + \cosh^3(x)} dx = \frac{(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x) + \sqrt{3}) \sqrt{2i\sqrt{3} + 6} \log\left(i\sqrt{3} + i\sqrt{2i\sqrt{3} + 6} + 2 \cosh(x) + 2 \sinh(x) - 1\right)}{\dots}$$

[In] `integrate(1/(1+cosh(x)^3),x, algorithm="fricas")`

[Out] $-1/18*((\text{sqrt}(3)*\cosh(x) + \text{sqrt}(3)*\sinh(x) + \text{sqrt}(3))*\text{sqrt}(2*I*\text{sqrt}(3) + 6)*\log(I*\text{sqrt}(3) + I*\text{sqrt}(2*I*\text{sqrt}(3) + 6) + 2*\cosh(x) + 2*\sinh(x) - 1) - (\text{sqrt}(3)*\cosh(x) + \text{sqrt}(3)*\sinh(x) + \text{sqrt}(3))*\text{sqrt}(2*I*\text{sqrt}(3) + 6)*\log(I*\text{sqrt}(3) - I*\text{sqrt}(2*I*\text{sqrt}(3) + 6) + 2*\cosh(x) + 2*\sinh(x) - 1) - (\text{sqrt}(3)*\cosh(x) + \text{sqrt}(3)*\sinh(x) + \text{sqrt}(3))*\text{sqrt}(-2*I*\text{sqrt}(3) + 6)*\log(-I*\text{sqrt}(3) + I*\text{sqrt}(-2*I*\text{sqrt}(3) + 6) + 2*\cosh(x) + 2*\sinh(x) - 1) + (\text{sqrt}(3)*\cosh(x) + \text{sqrt}(3)*\sinh(x) + \text{sqrt}(3))*\text{sqrt}(-2*I*\text{sqrt}(3) + 6)*\log(-I*\text{sqrt}(3) - I*\text{sqrt}(-2*I*\text{sqrt}(3) + 6) + 2*\cosh(x) + 2*\sinh(x) - 1) + 12)/(\cosh(x) + \sinh(x) + 1)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 330 vs. $2(73) = 146$.

Time = 1.51 (sec) , antiderivative size = 330, normalized size of antiderivative = 3.63

$$\int \frac{1}{1 + \cosh^3(x)} dx = -\frac{2\sqrt{2} \cdot 3^{\frac{3}{4}} \log\left(36 \tanh^2\left(\frac{x}{2}\right) - 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right) + 12\sqrt{3}\right)}{18 + 18\sqrt{3}} - \frac{3\sqrt{2} \cdot \sqrt[4]{3} \log\left(36 \tanh^2\left(\frac{x}{2}\right) - 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right) + 12\sqrt{3}\right)}{18 + 18\sqrt{3}} + \frac{3\sqrt{2} \cdot \sqrt[4]{3} \log\left(36 \tanh^2\left(\frac{x}{2}\right) + 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right) + 12\sqrt{3}\right)}{18 + 18\sqrt{3}} + \frac{2\sqrt{2} \cdot 3^{\frac{3}{4}} \log\left(36 \tanh^2\left(\frac{x}{2}\right) + 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right) + 12\sqrt{3}\right)}{18 + 18\sqrt{3}} + \frac{6 \tanh\left(\frac{x}{2}\right)}{18 + 18\sqrt{3}} + \frac{6\sqrt{3} \tanh\left(\frac{x}{2}\right)}{18 + 18\sqrt{3}} - \frac{2\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan}\left(\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) - 1\right)}{18 + 18\sqrt{3}} - \frac{2\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan}\left(\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) + 1\right)}{18 + 18\sqrt{3}}$$

[In] integrate(1/(1+cosh(x)**3),x)

[Out] $-2\sqrt{2} \cdot 3^{3/4} \log(36 \tanh(x/2)^2 - 12\sqrt{2} \cdot 3^{3/4} \tanh(x/2) + 12\sqrt{3}) / (18 + 18\sqrt{3}) - 3\sqrt{2} \cdot 3^{1/4} \log(36 \tanh(x/2)^2 - 12\sqrt{2} \cdot 3^{3/4} \tanh(x/2) + 12\sqrt{3}) / (18 + 18\sqrt{3}) + 3\sqrt{2} \cdot 3^{3/4} (1/4) \log(36 \tanh(x/2)^2 + 12\sqrt{2} \cdot 3^{3/4} \tanh(x/2) + 12\sqrt{3}) / (18 + 18\sqrt{3}) + 2\sqrt{2} \cdot 3^{3/4} \log(36 \tanh(x/2)^2 + 12\sqrt{2} \cdot 3^{3/4} \tanh(x/2) + 12\sqrt{3}) / (18 + 18\sqrt{3}) + 6 \tanh(x/2) / (18 + 18\sqrt{3}) + 6\sqrt{3} \tanh(x/2) / (18 + 18\sqrt{3}) - 2\sqrt{2} \cdot 3^{3/4} \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \tanh(x/2) - 1) / (18 + 18\sqrt{3}) - 2\sqrt{2} \cdot 3^{3/4} \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \tanh(x/2) + 1) / (18 + 18\sqrt{3})$

Maxima [F]

$$\int \frac{1}{1 + \cosh^3(x)} dx = \int \frac{1}{\cosh(x)^3 + 1} dx$$

[In] integrate(1/(1+cosh(x)^3),x, algorithm="maxima")

[Out] $-2/3/(e^x + 1) - \operatorname{integrate}(2/3*(e^{(3*x)} - 4*e^{(2*x)} + e^x)/(e^{(4*x)} - 2*e^{(3*x)} + 6*e^{(2*x)} - 2*e^x + 1), x)$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. $2(65) = 130$.

Time = 0.35 (sec) , antiderivative size = 275, normalized size of antiderivative = 3.02

$$\begin{aligned} \int \frac{1}{1 + \cosh^3(x)} dx &= \frac{1}{18} \sqrt{6\sqrt{3} + 9} \log \left(4 \left(2\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9} + 6e^x - 3 \right)^2 \right. \\ &\quad \left. + 4 \left(\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{3} \right)^2 \right) \\ &\quad - \frac{1}{18} \sqrt{6\sqrt{3} + 9} \log \left(4 \left(2\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9} - 6e^x + 3 \right)^2 \right. \\ &\quad \left. + 4 \left(\sqrt{3}\sqrt{6\sqrt{3} + 9} + 3\sqrt{3} \right)^2 \right) \\ &\quad - \frac{\sqrt{3}\sqrt{6\sqrt{3} + 9} \operatorname{arctan} \left(\frac{3(\sqrt{2\sqrt{3}-3} + 2e^x - 1)}{\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{3}} \right)}{9(2\sqrt{3} + 3)} \\ &\quad - \frac{\sqrt{3}\sqrt{6\sqrt{3} + 9} \operatorname{arctan} \left(-\frac{3(\sqrt{2\sqrt{3}-3} - 2e^x + 1)}{\sqrt{3}\sqrt{6\sqrt{3} + 9} + 3\sqrt{3}} \right)}{9(2\sqrt{3} + 3)} - \frac{2}{3(e^x + 1)} \end{aligned}$$

[In] integrate(1/(1+cosh(x)^3),x, algorithm="giac")

[Out] $\frac{1}{18}\sqrt{6\sqrt{3} + 9}\log(4*(2\sqrt{3})\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9}) + 6e^x - 3)^2 + 4*(\sqrt{3})\sqrt{6\sqrt{3} + 9} - 3\sqrt{3})^2$
 $- \frac{1}{18}\sqrt{6\sqrt{3} + 9}\log(4*(2\sqrt{3})\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9}) - 6e^x + 3)^2 + 4*(\sqrt{3})\sqrt{6\sqrt{3} + 9} + 3\sqrt{3})^2$
 $) - \frac{1}{9}\sqrt{3}\sqrt{6\sqrt{3} + 9}\arctan(3*(\sqrt{2\sqrt{3} - 3}) + 2e^x - 1)/(\sqrt{3})\sqrt{6\sqrt{3} + 9} - 3\sqrt{3}))/ (2\sqrt{3} + 3) - \frac{1}{9}\sqrt{3}$
 $)\sqrt{6\sqrt{3} + 9}\arctan(-3*(\sqrt{2\sqrt{3} - 3}) - 2e^x + 1)/(\sqrt{3})\sqrt{6\sqrt{3} + 9} + 3\sqrt{3}))/ (2\sqrt{3} + 3) - \frac{2}{3}/(e^x + 1)$

Mupad [B] (verification not implemented)

Time = 4.32 (sec) , antiderivative size = 291, normalized size of antiderivative = 3.20

$$\int \frac{1}{1 + \cosh^3(x)} dx = \ln \left(\frac{128}{9} + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{160}{3} \right. \right.$$

$$+ \left. \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x - 864) - 192 \right) \right.$$

$$\left. - \frac{32 e^x}{3} \right) - \frac{32 e^x}{3} \left. \right) \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} + \ln \left(\frac{128}{9} + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{160}{3} \right. \right.$$

$$+ \left. \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x - 864) - 192 \right) \right.$$

$$\left. - \frac{32 e^x}{3} \right) - \frac{32 e^x}{3} \left. \right) \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} - \ln \left(\frac{128}{9} - \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{160}{3} \right. \right.$$

$$+ \left. \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(192 + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x - 864) - 384 e^x \right) \right.$$

$$\left. - \frac{32 e^x}{3} \right) - \frac{32 e^x}{3} \left. \right) \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} - \ln \left(\frac{128}{9} - \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{160}{3} \right. \right.$$

$$+ \left. \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(192 + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x - 864) - 384 e^x \right) \right.$$

$$\left. - \frac{32 e^x}{3} \right) - \frac{32 e^x}{3} \left. \right) \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} - \frac{2}{3(e^x + 1)}$$

[In] int(1/(cosh(x)^3 + 1),x)

```
[Out] log((1/18 - (3^(1/2)*1i)/54)^(1/2)*((1/18 - (3^(1/2)*1i)/54)^(1/2)*(384*exp
(x) + (1/18 - (3^(1/2)*1i)/54)^(1/2)*(1152*exp(x) - 864) - 192) - (32*exp(x
)))/3 + 160/3) - (32*exp(x))/3 + 128/9)*(1/18 - (3^(1/2)*1i)/54)^(1/2) + log
(((3^(1/2)*1i)/54 + 1/18)^(1/2)*(((3^(1/2)*1i)/54 + 1/18)^(1/2)*(384*exp(x)
+ ((3^(1/2)*1i)/54 + 1/18)^(1/2)*(1152*exp(x) - 864) - 192) - (32*exp(x))/
3 + 160/3) - (32*exp(x))/3 + 128/9)*((3^(1/2)*1i)/54 + 1/18)^(1/2) - log(12
8/9 - (1/18 - (3^(1/2)*1i)/54)^(1/2)*((1/18 - (3^(1/2)*1i)/54)^(1/2)*((1/18
- (3^(1/2)*1i)/54)^(1/2)*(1152*exp(x) - 864) - 384*exp(x) + 192) - (32*exp
(x))/3 + 160/3) - (32*exp(x))/3)*(1/18 - (3^(1/2)*1i)/54)^(1/2) - log(128/9
- ((3^(1/2)*1i)/54 + 1/18)^(1/2)*(((3^(1/2)*1i)/54 + 1/18)^(1/2)*(((3^(1/2
)*1i)/54 + 1/18)^(1/2)*(1152*exp(x) - 864) - 384*exp(x) + 192) - (32*exp(x)
)/3 + 160/3) - (32*exp(x))/3)*((3^(1/2)*1i)/54 + 1/18)^(1/2) - 2/(3*(exp(x)
+ 1))
```

3.59 $\int \frac{1}{1 - \cosh^3(x)} dx$

Optimal result	384
Rubi [A] (verified)	384
Mathematica [F(-1)]	386
Maple [C] (verified)	386
Fricas [C] (verification not implemented)	386
Sympy [B] (verification not implemented)	387
Maxima [F]	388
Giac [B] (verification not implemented)	388
Mupad [B] (verification not implemented)	389

Optimal result

Integrand size = 10, antiderivative size = 95

$$\int \frac{1}{1 - \cosh^3(x)} dx = -\frac{2\sqrt[4]{-1} \arctan\left(\frac{(-1)^{3/4} \tanh\left(\frac{x}{2}\right)}{\sqrt[4]{3}}\right)}{3^{3/4} (1 - (-1)^{2/3})} - \frac{2\sqrt[4]{-1} \operatorname{arctanh}\left(\frac{(-1)^{3/4} \tanh\left(\frac{x}{2}\right)}{\sqrt[4]{3}}\right)}{3^{3/4} (1 + \sqrt[3]{-1})} - \frac{\sinh(x)}{3(1 - \cosh(x))}$$

[Out] $-2/3*(-1)^{(1/4)}*\arctan(1/3*(-1)^{(3/4)}*\tanh(1/2*x)*3^{(3/4)})*3^{(1/4)}/(1-(-1)^{(2/3)})-2/3*(-1)^{(1/4)}*\operatorname{arctanh}(1/3*(-1)^{(3/4)}*\tanh(1/2*x)*3^{(3/4)})*3^{(1/4)}/(1+(-1)^{(1/3)})-1/3*\sinh(x)/(1-\cosh(x))$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3292, 2727, 2738, 214, 211}

$$\int \frac{1}{1 - \cosh^3(x)} dx = -\frac{2\sqrt[4]{-1} \arctan\left(\frac{(-1)^{3/4} \tanh\left(\frac{x}{2}\right)}{\sqrt[4]{3}}\right)}{3^{3/4} (1 - (-1)^{2/3})} - \frac{2\sqrt[4]{-1} \operatorname{arctanh}\left(\frac{(-1)^{3/4} \tanh\left(\frac{x}{2}\right)}{\sqrt[4]{3}}\right)}{3^{3/4} (1 + \sqrt[3]{-1})} - \frac{\sinh(x)}{3(1 - \cosh(x))}$$

[In] $\text{Int}[(1 - \text{Cosh}[x]^3)^{-1}, x]$

[Out] $(-2*(-1)^{1/4}*\text{ArcTan}[((-1)^{3/4}*\text{Tanh}[x/2])/3^{1/4}])/(3^{3/4}*(1 - (-1)^{2/3})) - (2*(-1)^{1/4}*\text{ArcTan}[((-1)^{3/4}*\text{Tanh}[x/2])/3^{1/4}])/(3^{3/4}*(1 + (-1)^{1/3})) - \text{Sinh}[x]/(3*(1 - \text{Cosh}[x]))$

Rule 211

$\text{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{PosQ}[a/b]$

Rule 214

$\text{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 2727

$\text{Int}[(a_+ + (b_+)*\sin[(c_+ + (d_+)*(x_+))])^{-1}, x_Symbol] \rightarrow \text{Simp}[-\text{Cos}[c + d*x]/(d*(b + a*\sin[c + d*x])), x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{EqQ}[a^2 - b^2, 0]$

Rule 2738

$\text{Int}[(a_+ + (b_+)*\sin[\text{Pi}/2 + (c_+ + (d_+)*(x_+))])^{-1}, x_Symbol] \rightarrow \text{With}\{e = \text{FreeFactors}[\text{Tan}[(c + d*x)/2], x]\}, \text{Dist}[2*(e/d), \text{Subst}[\text{Int}[1/(a + b + (a - b)*e^2*x^2), x], x, \text{Tan}[(c + d*x)/2]/e], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[a^2 - b^2, 0]$

Rule 3292

$\text{Int}[(a_+ + (b_+)*((c_+)*\sin[(e_+ + (f_+)*(x_+))])^{n_+})^{p_+}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrig}[(a + b*(c*\sin[e + f*x])^n)^p, x], x] /; \text{FreeQ}\{a, b, c, e, f, n\}, x] \&\& (\text{IGtQ}[p, 0] \mid\mid (\text{EqQ}[p, -1] \&\& \text{IntegerQ}[n]))$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{3(1 - \cosh(x))} + \frac{1}{3(1 + \sqrt[3]{-1} \cosh(x))} + \frac{1}{3(1 - (-1)^{2/3} \cosh(x))} \right) dx \\ &= \frac{1}{3} \int \frac{1}{1 - \cosh(x)} dx + \frac{1}{3} \int \frac{1}{1 + \sqrt[3]{-1} \cosh(x)} dx + \frac{1}{3} \int \frac{1}{1 - (-1)^{2/3} \cosh(x)} dx \\ &= -\frac{\sinh(x)}{3(1 - \cosh(x))} + \frac{2}{3} \text{Subst} \left(\int \frac{1}{1 + \sqrt[3]{-1} - (1 - \sqrt[3]{-1}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\ &\quad + \frac{2}{3} \text{Subst} \left(\int \frac{1}{1 - (-1)^{2/3} - (1 + (-1)^{2/3}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \end{aligned}$$

$$= -\frac{2\sqrt[4]{-1} \arctan\left(\frac{(-1)^{3/4} \tanh(\frac{x}{2})}{\sqrt[4]{3}}\right)}{3^{3/4} (1 - (-1)^{2/3})} - \frac{2\sqrt[4]{-1} \operatorname{arctanh}\left(\frac{(-1)^{3/4} \tanh(\frac{x}{2})}{\sqrt[4]{3}}\right)}{3^{3/4} (1 + \sqrt[3]{-1})} - \frac{\sinh(x)}{3(1 - \cosh(x))}$$

Mathematica [F(-1)]

Timed out.

$$\int \frac{1}{1 - \cosh^3(x)} dx = \$Aborted$$

[In] Integrate[(1 - Cosh[x]^3)^(-1), x]

[Out] \$Aborted

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.07 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.48

method	result
risch	$\frac{2}{3(e^x-1)} + \left(\sum_{R=\text{RootOf}(243_Z^4-27_Z^2+1)} -R \ln(162_R^3 - 27_R^2 - 9_R + e^x + 2) \right)$
default	$\frac{3^{\frac{1}{4}} \sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})^2 + \sqrt{2} 3^{\frac{1}{4}} \tanh(\frac{x}{2}) + \sqrt{3}}{\tanh(\frac{x}{2})^2 - \sqrt{2} 3^{\frac{1}{4}} \tanh(\frac{x}{2}) + \sqrt{3}}\right) + 2 \operatorname{arctan}\left(\frac{3^{\frac{3}{4}} \tanh(\frac{x}{2}) \sqrt{2}}{3} + 1\right) + 2 \operatorname{arctan}\left(\frac{3^{\frac{3}{4}} \tanh(\frac{x}{2}) \sqrt{2}}{3} - 1\right) \right)}{12} - \frac{3^{\frac{3}{4}} \sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})}{\tanh(\frac{x}{2})}\right) \right)}{12}$

[In] int(1/(1-cosh(x)^3), x, method=_RETURNVERBOSE)

[Out] 2/3/(exp(x)-1)+sum(_R*ln(162*_R^3-27*_R^2-9*_R+exp(x)+2), _R=RootOf(243*_Z^4-27*_Z^2+1))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 234, normalized size of antiderivative = 2.46

$$\int \frac{1}{1 - \cosh^3(x)} dx = \frac{(\sqrt{3} \cosh(x) + \sqrt{3} \sinh(x) - \sqrt{3}) \sqrt{-2i\sqrt{3} + 6} \log\left(i\sqrt{3} + i\sqrt{-2i\sqrt{3} + 6} + 2 \cosh(x) + 2 \sinh(x) + \dots\right)}{\dots}$$

[In] integrate(1/(1-cosh(x)^3), x, algorithm="fricas")

```
[Out] -1/18*((sqrt(3)*cosh(x) + sqrt(3)*sinh(x) - sqrt(3))*sqrt(-2*I*sqrt(3) + 6)
*log(I*sqrt(3) + I*sqrt(-2*I*sqrt(3) + 6) + 2*cosh(x) + 2*sinh(x) + 1) - (s
qrt(3)*cosh(x) + sqrt(3)*sinh(x) - sqrt(3))*sqrt(-2*I*sqrt(3) + 6)*log(I*sq
rt(3) - I*sqrt(-2*I*sqrt(3) + 6) + 2*cosh(x) + 2*sinh(x) + 1) - (sqrt(3)*co
sh(x) + sqrt(3)*sinh(x) - sqrt(3))*sqrt(2*I*sqrt(3) + 6)*log(-I*sqrt(3) + I
*sqrt(2*I*sqrt(3) + 6) + 2*cosh(x) + 2*sinh(x) + 1) + (sqrt(3)*cosh(x) + sq
rt(3)*sinh(x) - sqrt(3))*sqrt(2*I*sqrt(3) + 6)*log(-I*sqrt(3) - I*sqrt(2*I*
sqrt(3) + 6) + 2*cosh(x) + 2*sinh(x) + 1) - 12)/(cosh(x) + sinh(x) - 1)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. $2(78) = 156$.

Time = 1.56 (sec) , antiderivative size = 320, normalized size of antiderivative = 3.37

$$\int \frac{1}{1 - \cosh^3(x)} dx = -\frac{\sqrt{2} \cdot \sqrt[4]{3} \log\left(4 \tanh^2\left(\frac{x}{2}\right) - 4\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) + 4\sqrt{3}\right)}{12}$$

$$-\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log\left(4 \tanh^2\left(\frac{x}{2}\right) - 4\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) + 4\sqrt{3}\right)}{36}$$

$$+\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log\left(4 \tanh^2\left(\frac{x}{2}\right) + 4\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) + 4\sqrt{3}\right)}{36}$$

$$+\frac{\sqrt{2} \cdot \sqrt[4]{3} \log\left(4 \tanh^2\left(\frac{x}{2}\right) + 4\sqrt{2} \cdot \sqrt[4]{3} \tanh\left(\frac{x}{2}\right) + 4\sqrt{3}\right)}{12}$$

$$-\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan}\left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right)}{3} - 1\right)}{18}$$

$$+\frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan}\left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right)}{3} - 1\right)}{6}$$

$$-\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan}\left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right)}{3} + 1\right)}{18}$$

$$+\frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan}\left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh\left(\frac{x}{2}\right)}{3} + 1\right)}{6} + \frac{1}{3 \tanh\left(\frac{x}{2}\right)}$$

```
[In] integrate(1/(1-cosh(x)**3),x)
```

```
[Out] -sqrt(2)*3**(1/4)*log(4*tanh(x/2)**2 - 4*sqrt(2)*3**(1/4)*tanh(x/2) + 4*sq
rt(3))/12 - sqrt(2)*3**(3/4)*log(4*tanh(x/2)**2 - 4*sqrt(2)*3**(1/4)*tanh(x/
2) + 4*sqrt(3))/36 + sqrt(2)*3**(3/4)*log(4*tanh(x/2)**2 + 4*sqrt(2)*3**(1/
4)*tanh(x/2) + 4*sqrt(3))/36 + sqrt(2)*3**(1/4)*log(4*tanh(x/2)**2 + 4*sqrt
```

$(2) \cdot 3^{1/4} \cdot \tanh(x/2) + 4 \cdot \sqrt{3}) / 12 - \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 - 1) / 18 + \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 - 1) / 6 - \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 + 1) / 18 + \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 + 1) / 6 + 1 / (3 \cdot \tanh(x/2))$

Maxima [F]

$$\int \frac{1}{1 - \cosh^3(x)} dx = \int -\frac{1}{\cosh(x)^3 - 1} dx$$

[In] integrate(1/(1-cosh(x)^3),x, algorithm="maxima")

[Out] 2/3/(e^x - 1) + integrate(2/3*(e^(3*x) + 4*e^(2*x) + e^x)/(e^(4*x) + 2*e^(3*x) + 6*e^(2*x) + 2*e^x + 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 275 vs. 2(67) = 134.

Time = 0.32 (sec) , antiderivative size = 275, normalized size of antiderivative = 2.89

$$\begin{aligned} \int \frac{1}{1 - \cosh^3(x)} dx = & -\frac{1}{18} \sqrt{6\sqrt{3} + 9} \log \left(4 \left(2\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9} + 6e^x + 3 \right)^2 \right. \\ & \left. + 4 \left(\sqrt{3}\sqrt{6\sqrt{3} + 9} + 3\sqrt{3} \right)^2 \right) \\ & + \frac{1}{18} \sqrt{6\sqrt{3} + 9} \log \left(4 \left(2\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{6\sqrt{3} + 9} - 6e^x - 3 \right)^2 \right. \\ & \left. + 4 \left(\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{3} \right)^2 \right) \\ & + \frac{\sqrt{3}\sqrt{6\sqrt{3} + 9} \arctan \left(\frac{3(\sqrt{2\sqrt{3}-3} + 2e^x + 1)}{\sqrt{3}\sqrt{6\sqrt{3} + 9} + 3\sqrt{3}} \right)}{9(2\sqrt{3} + 3)} \\ & + \frac{\sqrt{3}\sqrt{6\sqrt{3} + 9} \arctan \left(-\frac{3(\sqrt{2\sqrt{3}-3} - 2e^x - 1)}{\sqrt{3}\sqrt{6\sqrt{3} + 9} - 3\sqrt{3}} \right)}{9(2\sqrt{3} + 3)} + \frac{2}{3(e^x - 1)} \end{aligned}$$

[In] integrate(1/(1-cosh(x)^3),x, algorithm="giac")

[Out] -1/18*sqrt(6*sqrt(3) + 9)*log(4*(2*sqrt(3)*sqrt(6*sqrt(3) + 9) - 3*sqrt(6*sqrt(3) + 9) + 6*e^x + 3)^2 + 4*(sqrt(3)*sqrt(6*sqrt(3) + 9) + 3*sqrt(3))^2) + 1/18*sqrt(6*sqrt(3) + 9)*log(4*(2*sqrt(3)*sqrt(6*sqrt(3) + 9) - 3*sqrt(6

*sqrt(3) + 9) - 6*e^x - 3)^2 + 4*(sqrt(3)*sqrt(6*sqrt(3) + 9) - 3*sqrt(3))^2 + 1/9*sqrt(3)*sqrt(6*sqrt(3) + 9)*arctan(3*(sqrt(2*sqrt(3) - 3) + 2*e^x + 1)/(sqrt(3)*sqrt(6*sqrt(3) + 9) + 3*sqrt(3)))/(2*sqrt(3) + 3) + 1/9*sqrt(3)*sqrt(6*sqrt(3) + 9)*arctan(-3*(sqrt(2*sqrt(3) - 3) - 2*e^x - 1)/(sqrt(3)*sqrt(6*sqrt(3) + 9) - 3*sqrt(3)))/(2*sqrt(3) + 3) + 2/3/(e^x - 1)

Mupad [B] (verification not implemented)

Time = 4.32 (sec) , antiderivative size = 295, normalized size of antiderivative = 3.11

$$\int \frac{1}{1 - \cosh^3(x)} dx = \ln \left(\frac{32 e^x}{3} + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{32 e^x}{3} - \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x + 864) + 192 \right) + \frac{160}{3} \right) + \frac{128}{9} \right) \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} + \ln \left(\frac{32 e^x}{3} + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{32 e^x}{3} - \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x + 864) + 192 \right) + \frac{160}{3} \right) + \frac{128}{9} \right) \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} - \ln \left(\frac{32 e^x}{3} - \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{32 e^x}{3} + \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x - \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x + 864) + 192 \right) + \frac{160}{3} \right) + \frac{128}{9} \right) \sqrt{\frac{1}{18} - \frac{\sqrt{3} \operatorname{li}}{54}} - \ln \left(\frac{32 e^x}{3} - \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(\frac{32 e^x}{3} + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x - \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x + 864) + 192 \right) + \frac{160}{3} \right) + \frac{128}{9} \right) \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} + \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} \left(384 e^x - \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} (1152 e^x + 864) + 192 \right) + \frac{160}{3} \right) + \frac{128}{9} \right) \sqrt{\frac{1}{18} + \frac{\sqrt{3} \operatorname{li}}{54}} + \frac{2}{3(e^x - 1)}$$

[In] int(-1/(cosh(x)^3 - 1), x)

[Out] log((32*exp(x))/3 + (1/18 - (3^(1/2)*1i)/54)^(1/2)*((32*exp(x))/3 - (1/18 - (3^(1/2)*1i)/54)^(1/2)*(384*exp(x) + (1/18 - (3^(1/2)*1i)/54)^(1/2)*(1152*exp(x) + 864) + 192) + 160/3) + 128/9)*(1/18 - (3^(1/2)*1i)/54)^(1/2) + log

$$\begin{aligned}
& ((32*\exp(x))/3 + ((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)}*((32*\exp(x))/3 - ((3^{(1/2)}* \\
& 1i)/54 + 1/18)^{(1/2)}*(384*\exp(x) + ((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)}*(1152*\exp \\
& (x) + 864) + 192) + 160/3) + 128/9)*((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)} - \log((3 \\
& 2*\exp(x))/3 - (1/18 - (3^{(1/2)}*1i)/54)^{(1/2)}*((32*\exp(x))/3 + (1/18 - (3^{(1/2)} \\
& /2)*1i)/54)^{(1/2)}*(384*\exp(x) - (1/18 - (3^{(1/2)}*1i)/54)^{(1/2)}*(1152*\exp(x) \\
& + 864) + 192) + 160/3) + 128/9)*(1/18 - (3^{(1/2)}*1i)/54)^{(1/2)} - \log((32*e \\
& xp(x))/3 - ((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)}*((32*\exp(x))/3 + ((3^{(1/2)}*1i)/54 \\
& + 1/18)^{(1/2)}*(384*\exp(x) - ((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)}*(1152*\exp(x) + \\
& 864) + 192) + 160/3) + 128/9)*((3^{(1/2)}*1i)/54 + 1/18)^{(1/2)} + 2/(3*(\exp(x) \\
& - 1))
\end{aligned}$$

3.60 $\int \frac{1}{a+b \cosh^4(x)} dx$

Optimal result	391
Rubi [A] (verified)	392
Mathematica [C] (verified)	395
Maple [C] (verified)	396
Fricas [B] (verification not implemented)	396
Sympy [F(-1)]	398
Maxima [F]	398
Giac [B] (verification not implemented)	399
Mupad [B] (verification not implemented)	400

Optimal result

Integrand size = 10, antiderivative size = 361

$$\begin{aligned}
 & \int \frac{1}{a+b \cosh^4(x)} dx \\
 &= \frac{\sqrt{\sqrt{a}-\sqrt{a+b}} \operatorname{arctanh}\left(\frac{\sqrt{\sqrt{a}+\sqrt{a+b}-\sqrt{2}} \sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}-\sqrt{a+b}}}\right)}{2\sqrt{2}a^{3/4}\sqrt{a+b}} \\
 &\quad - \frac{\sqrt{\sqrt{a}-\sqrt{a+b}} \operatorname{arctanh}\left(\frac{\sqrt{\sqrt{a}+\sqrt{a+b}+\sqrt{2}} \sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}-\sqrt{a+b}}}\right)}{2\sqrt{2}a^{3/4}\sqrt{a+b}} \\
 &\quad - \frac{\sqrt{\sqrt{a}+\sqrt{a+b}} \log\left(\sqrt{a+b}-\sqrt{2}\sqrt[4]{a}\sqrt{\sqrt{a}+\sqrt{a+b}}\tanh(x)+\sqrt{a}\tanh^2(x)\right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}} \\
 &\quad + \frac{\sqrt{\sqrt{a}+\sqrt{a+b}} \log\left(\sqrt{a+b}+\sqrt{2}\sqrt[4]{a}\sqrt{\sqrt{a}+\sqrt{a+b}}\tanh(x)+\sqrt{a}\tanh^2(x)\right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}}
 \end{aligned}$$

```

[Out] 1/4*arctanh(((a^(1/2)+(a+b)^(1/2))^(1/2)-a^(1/4)*2^(1/2)*tanh(x))/(a^(1/2)-(a+b)^(1/2))^(1/2))*a^(1/2)-(a+b)^(1/2))^(1/2)/a^(3/4)*2^(1/2)/(a+b)^(1/2)
-1/4*arctanh(((a^(1/2)+(a+b)^(1/2))^(1/2)+a^(1/4)*2^(1/2)*tanh(x))/(a^(1/2)-(a+b)^(1/2))^(1/2))*a^(1/2)-(a+b)^(1/2))^(1/2)/a^(3/4)*2^(1/2)/(a+b)^(1/2)
-1/8*ln((a+b)^(1/2)-a^(1/4)*2^(1/2)*(a^(1/2)+(a+b)^(1/2))^(1/2)*tanh(x)+a^(1/2)*tanh(x)^2)*(a^(1/2)+(a+b)^(1/2))^(1/2)/a^(3/4)*2^(1/2)/(a+b)^(1/2)+1/8*ln((a+b)^(1/2)+a^(1/4)*2^(1/2)*(a^(1/2)+(a+b)^(1/2))^(1/2)*tanh(x)+a^(1/2)*tanh(x)^2)*(a^(1/2)+(a+b)^(1/2))^(1/2)/a^(3/4)*2^(1/2)/(a+b)^(1/2)

```

Rubi [A] (verified)

Time = 0.85 (sec) , antiderivative size = 485, normalized size of antiderivative = 1.34, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3288, 1183, 648, 632, 210, 642}

$$\int \frac{1}{a + b \cosh^4(x)} dx = \frac{(\sqrt{a} - \sqrt{a+b}) \arctan\left(\frac{\sqrt[4]{a}\sqrt{\sqrt{a}\sqrt{a+b}+a+b}-\sqrt{2}(a+b)^{3/4} \coth(x)}{\sqrt[4]{a}\sqrt{-\sqrt{a}\sqrt{a+b}+a+b}}\right)}{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{-\sqrt{a}\sqrt{a+b}+a+b}} - \frac{(\sqrt{a} - \sqrt{a+b}) \arctan\left(\frac{\sqrt{2}(a+b)^{3/4} \coth(x)+\sqrt[4]{a}\sqrt{\sqrt{a}\sqrt{a+b}+a+b}}{\sqrt[4]{a}\sqrt{-\sqrt{a}\sqrt{a+b}+a+b}}\right)}{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{-\sqrt{a}\sqrt{a+b}+a+b}} - \frac{(\sqrt{a+b} + \sqrt{a}) \log\left((a+b)^{3/4} \coth^2(x) - \sqrt{2}\sqrt[4]{a}\sqrt{\sqrt{a}\sqrt{a+b}+a+b} \coth(x) + \sqrt{a}\sqrt[4]{a+b}\right)}{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{\sqrt{a}\sqrt{a+b}+a+b}} + \frac{(\sqrt{a+b} + \sqrt{a}) \log\left((a+b)^{3/4} \coth^2(x) + \sqrt{2}\sqrt[4]{a}\sqrt{\sqrt{a}\sqrt{a+b}+a+b} \coth(x) + \sqrt{a}\sqrt[4]{a+b}\right)}{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{\sqrt{a}\sqrt{a+b}+a+b}}$$

[In] Int[(a + b*Cosh[x]^4)^(-1), x]

[Out] ((Sqrt[a] - Sqrt[a + b])*ArcTan[(a^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]] - Sqrt[2]*(a + b)^(3/4)*Coth[x])/(a^(1/4)*Sqrt[a + b - Sqrt[a]*Sqrt[a + b]])]/(2*Sqrt[2]*a^(3/4)*(a + b)^(1/4)*Sqrt[a + b - Sqrt[a]*Sqrt[a + b]]) - ((Sqrt[a] - Sqrt[a + b])*ArcTan[(a^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]] + Sqrt[2]*(a + b)^(3/4)*Coth[x])/(a^(1/4)*Sqrt[a + b - Sqrt[a]*Sqrt[a + b]])]/(2*Sqrt[2]*a^(3/4)*(a + b)^(1/4)*Sqrt[a + b - Sqrt[a]*Sqrt[a + b]]) - ((Sqrt[a] + Sqrt[a + b])*Log[Sqrt[a]*(a + b)^(1/4) - Sqrt[2]*a^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]]*Coth[x] + (a + b)^(3/4)*Coth[x]^2])/(4*Sqrt[2]*a^(3/4)*(a + b)^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]]) + ((Sqrt[a] + Sqrt[a + b])*Log[Sqrt[a]*(a + b)^(1/4) + Sqrt[2]*a^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]]*Coth[x] + (a + b)^(3/4)*Coth[x]^2])/(4*Sqrt[2]*a^(3/4)*(a + b)^(1/4)*Sqrt[a + b + Sqrt[a]*Sqrt[a + b]])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},

$x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

$\text{Int}[\frac{(d_.) + (e_.)*(x_.)}{(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2}, x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 648

$\text{Int}[\frac{(d_.) + (e_.)*(x_.)}{(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2}, x_Symbol] \rightarrow \text{Dist}[(2*c*d - b*e)/(2*c), \text{Int}[1/(a + b*x + c*x^2), x], x] + \text{Dist}[e/(2*c), \text{Int}[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{NeQ}[2*c*d - b*e, 0] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& !\text{NiceSqrtQ}[b^2 - 4*a*c]$

Rule 1183

$\text{Int}[\frac{(d_.) + (e_.)*(x_.)^2}{(a_.) + (b_.)*(x_.)^2 + (c_.)*(x_.)^4}, x_Symbol] \rightarrow \text{With}\{q = \text{Rt}[a/c, 2]\}, \text{With}\{r = \text{Rt}[2*q - b/c, 2]\}, \text{Dist}[1/(2*c*q*r), \text{Int}[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + \text{Dist}[1/(2*c*q*r), \text{Int}[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0] \&\& \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \&\& \text{NegQ}[b^2 - 4*a*c]$

Rule 3288

$\text{Int}[\frac{(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^4)^{p_}}{(a_.) + (b_.)*\sin[(e_.) + (f_.)*(x_.)]^4}, x_Symbol] \rightarrow \text{With}\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[ff/f, \text{Subst}[\text{Int}[(a + 2*a*ff^2*x^2 + (a + b)*ff^4*x^4)^p/(1 + ff^2*x^2)^{(2*p + 1)}, x], x, \text{Tan}[e + f*x]/ff], x]] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{IntegerQ}[p]$

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst}\left(\int \frac{1 - x^2}{a - 2ax^2 + (a + b)x^4} dx, x, \coth(x)\right) \\ &= \frac{\sqrt[4]{a + b} \text{Subst}\left(\int \frac{\frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} - \left(1 + \frac{\sqrt{a}}{\sqrt{a+b}}\right)x}{\frac{\sqrt{a}}{\sqrt{a+b}} - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x)\right)}{2\sqrt{2}a^{3/4}\sqrt{a + b + \sqrt{a}\sqrt{a + b}}} \\ &\quad + \frac{\sqrt[4]{a + b} \text{Subst}\left(\int \frac{\frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} + \left(1 + \frac{\sqrt{a}}{\sqrt{a+b}}\right)x}{\frac{\sqrt{a}}{\sqrt{a+b}} + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x)\right)}{2\sqrt{2}a^{3/4}\sqrt{a + b + \sqrt{a}\sqrt{a + b}}} \end{aligned}$$

$$\begin{aligned}
&= \frac{(\sqrt{a} - \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{1}{\frac{\sqrt{a}}{\sqrt{a+b}} - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x) \right)}{4\sqrt{a}(a+b)} \\
&- \frac{(\sqrt{a} - \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{1}{\frac{\sqrt{a}}{\sqrt{a+b}} + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x) \right)}{4\sqrt{a}(a+b)} \\
&- \frac{(\sqrt{a} + \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{-\frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + 2x}{\frac{\sqrt{a}}{\sqrt{a+b}} - \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x) \right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}} \\
&+ \frac{(\sqrt{a} + \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{\frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + 2x}{\frac{\sqrt{a}}{\sqrt{a+b}} + \frac{\sqrt{2} \sqrt[4]{a} \sqrt{a+b+\sqrt{a}\sqrt{a+b}x}}{(a+b)^{3/4}} + x^2} dx, x, \coth(x) \right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}} \\
&= \frac{(\sqrt{a} + \sqrt{a+b}) \log \left(\sqrt{a}\sqrt{a+b} - \sqrt{2}\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}\coth(x) + (a+b)^{3/4}\coth^2(x) \right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}} \\
&+ \frac{(\sqrt{a} + \sqrt{a+b}) \log \left(\sqrt{a}\sqrt{a+b} + \sqrt{2}\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}\coth(x) + (a+b)^{3/4}\coth^2(x) \right)}{4\sqrt{2}a^{3/4}\sqrt{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}} \\
&+ \frac{(\sqrt{a} - \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{1}{-\frac{2\sqrt{a}(a+b-\sqrt{a}\sqrt{a+b})}{(a+b)^{3/2}} - x^2} dx, x, -\frac{\sqrt{2}\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} + 2\coth(x) \right)}{2\sqrt{a}(a+b)} \\
&+ \frac{(\sqrt{a} - \sqrt{a+b}) \operatorname{Subst} \left(\int \frac{1}{-\frac{2\sqrt{a}(a+b-\sqrt{a}\sqrt{a+b})}{(a+b)^{3/2}} - x^2} dx, x, \frac{\sqrt{2}\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} + 2\coth(x) \right)}{2\sqrt{a}(a+b)}
\end{aligned}$$

$$\begin{aligned}
& (\sqrt{a} - \sqrt{a+b}) \arctan \left(\frac{(a+b)^{3/4} \left(\frac{\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} - \sqrt{2} \coth(x) \right)}{\sqrt[4]{a}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}} \right) \\
= & \frac{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}}{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}} \\
& (\sqrt{a} - \sqrt{a+b}) \arctan \left(\frac{(a+b)^{3/4} \left(\frac{\sqrt[4]{a}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{(a+b)^{3/4}} + \sqrt{2} \coth(x) \right)}{\sqrt[4]{a}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}} \right) \\
- & \frac{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}}{2\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b-\sqrt{a}\sqrt{a+b}}} \\
& (\sqrt{a} + \sqrt{a+b}) \log \left(\sqrt{a}\sqrt[4]{a+b} - \sqrt{2}\sqrt[4]{a}\sqrt{a+b} + \sqrt{a}\sqrt{a+b} \coth(x) + (a+b)^{3/4} \coth^2(x) \right) \\
- & \frac{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}} \\
& (\sqrt{a} + \sqrt{a+b}) \log \left(\sqrt{a}\sqrt[4]{a+b} + \sqrt{2}\sqrt[4]{a}\sqrt{a+b} + \sqrt{a}\sqrt{a+b} \coth(x) + (a+b)^{3/4} \coth^2(x) \right) \\
+ & \frac{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}{4\sqrt{2}a^{3/4}\sqrt[4]{a+b}\sqrt{a+b+\sqrt{a}\sqrt{a+b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.98 (sec) , antiderivative size = 121, normalized size of antiderivative = 0.34

$$\int \frac{1}{a + b \cosh^4(x)} dx = -\frac{\arctan \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{-a + i\sqrt{a}\sqrt{b}}} \right)}{2\sqrt{a}\sqrt{-a + i\sqrt{a}\sqrt{b}}} + \frac{\operatorname{arctanh} \left(\frac{\sqrt{a} \tanh(x)}{\sqrt{a + i\sqrt{a}\sqrt{b}}} \right)}{2\sqrt{a}\sqrt{a + i\sqrt{a}\sqrt{b}}}$$

[In] Integrate[(a + b*Cosh[x]^4)^(-1), x]

[Out] -1/2*ArcTan[(Sqrt[a]*Tanh[x])/Sqrt[-a + I*Sqrt[a]*Sqrt[b]]]/(Sqrt[a]*Sqrt[-a + I*Sqrt[a]*Sqrt[b]]) + ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + I*Sqrt[a]*Sqrt[b]]]/(2*Sqrt[a]*Sqrt[a + I*Sqrt[a]*Sqrt[b]])

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.33 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.27

method	result
risch	$\sum_{R=\text{RootOf}(1+(256a^4+256a^3b)Z^4-32a^2Z^2)} _R \ln \left(e^{2x} + \left(-\frac{128a^4}{b} - 128a^3 \right) _R^3 + \left(\frac{32a^3}{b} + 32a^2 \right) _R^2 + \right.$
default	$\left(\sum_{R=\text{RootOf}((a+b)Z^8+(-4a+4b)Z^6+(6a+6b)Z^4+(-4a+4b)Z^2+a+b)} \frac{(-_R^6+3_R^4-3_R^2+1) \ln(\tanh(\frac{x}{2})-_R)}{_R^7 _R^a + _R^7 _R^{b-3} + _R^5 _R^{a+3} + _R^5 _R^{b+3} + _R^3 _R^{a+3} + _R^3 _R^{b-3}} \right)$

[In] int(1/(a+b*cosh(x)^4),x,method=_RETURNVERBOSE)

[Out] sum(_R*ln(exp(2*x)+(-128*a^4/b-128*a^3)*_R^3+(32/b*a^3+32*a^2)*_R^2+(8*a^2/b-8*a)*_R-2*a/b+1),_R=RootOf(1+(256*a^4+256*a^3*b)*Z^4-32*a^2*_Z^2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 771 vs. 2(247) = 494.

Time = 0.29 (sec) , antiderivative size = 771, normalized size of antiderivative = 2.14

$$\begin{aligned}
 & \int \frac{1}{a + b \cosh^4(x)} dx \\
 &= -\frac{1}{4} \sqrt{\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + 1}{a^2 + ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
 &\quad \left. + 2 \left(ab + (a^4 + a^3b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} \right) \sqrt{\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + 1}{a^2 + ab}} \right. \\
 &\quad \left. + 2(a^3 + a^2b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + b \right) \\
 &+ \frac{1}{4} \sqrt{\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + 1}{a^2 + ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
 &\quad \left. - 2 \left(ab + (a^4 + a^3b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} \right) \sqrt{\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + 1}{a^2 + ab}} \right. \\
 &\quad \left. + 2(a^3 + a^2b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + b \right) \\
 &- \frac{1}{4} \sqrt{-\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} - 1}{a^2 + ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
 &\quad \left. + 2 \left(ab - (a^4 + a^3b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} \right) \sqrt{-\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} - 1}{a^2 + ab}} \right. \\
 &\quad \left. - 2(a^3 + a^2b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + b \right) \\
 &+ \frac{1}{4} \sqrt{-\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} - 1}{a^2 + ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
 &\quad \left. - 2 \left(ab - (a^4 + a^3b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} \right) \sqrt{-\frac{(a^2 + ab) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} - 1}{a^2 + ab}} \right. \\
 &\quad \left. - 2(a^3 + a^2b) \sqrt{-\frac{b}{a^5 + 2a^4b + a^3b^2}} + b \right)
 \end{aligned}$$

[In] integrate(1/(a+b*cosh(x)^4),x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/4*\sqrt{((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + 1)/(a^2 + a*b)} \\ & * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*(a*b + (a^4 + a^3*b) \\ & *\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)})) * \sqrt{((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} \\ & + 1)/(a^2 + a*b)} + 2*(a^3 + a^2*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + b \\ & + 1/4*\sqrt{((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + 1)/(a^2 + a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 - 2 \\ & *(a*b + (a^4 + a^3*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)})) * \sqrt{((a^2 + a*b) \\ & *\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + 1)/(a^2 + a*b)} + 2*(a^3 + a^2*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + b \\ & - 1/4*\sqrt{-((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} - 1)/(a^2 + a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*(a*b - (a^4 + a^3*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)}) \\ &) * \sqrt{-((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} - 1)/(a^2 + a*b)} \\ & - 2*(a^3 + a^2*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + b + 1/4*\sqrt{-((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} - 1)/(a^2 + a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 - 2*(a*b - (a^4 + a^3*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)}) \\ &) * \sqrt{-((a^2 + a*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} - 1)/(a^2 + a*b)} - 2*(a^3 + a^2*b)*\sqrt{-b/(a^5 + 2*a^4*b + a^3*b^2)} + b \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{a + b \cosh^4(x)} dx = \text{Timed out}$$

[In] integrate(1/(a+b*cosh(x)**4),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{a + b \cosh^4(x)} dx = \int \frac{1}{b \cosh(x)^4 + a} dx$$

[In] integrate(1/(a+b*cosh(x)^4),x, algorithm="maxima")

[Out] integrate(1/(b*cosh(x)^4 + a), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1781 vs. 2(247) = 494.

Time = 2.36 (sec) , antiderivative size = 1781, normalized size of antiderivative = 4.93

$$\int \frac{1}{a + b \cosh^4(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a+b*cosh(x)^4),x, algorithm="giac")

```
[Out] 1/4*sqrt((a^2 + sqrt(-a*b)*a)/(a^4 + a^3*b))*log(abs(60*a^4*b*e^(2*x) + 68*
a^3*b^2*e^(2*x) - 16*a^2*b^3*e^(2*x) + 24*sqrt(-a*b)*a^4*e^(2*x) + 48*sqrt(
a^2 - sqrt(-a*b)*a)*a^3*b*e^(2*x) - 16*sqrt(-a*b)*a^3*b*e^(2*x) + 61*sqrt(a
^2 - sqrt(-a*b)*a)*a^2*b^2*e^(2*x) - 64*sqrt(-a*b)*a^2*b^2*e^(2*x) - 4*sqrt
(a^2 - sqrt(-a*b)*a)*a*b^3*e^(2*x) + 6*a^4*b + 2*a^3*b^2 - 8*a^2*b^3 + 24*s
qrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a^3*e^(2*x) + 5*sqrt(a^2 - sqrt(-a*b)*a)
*sqrt(-a*b)*a^2*b*e^(2*x) - 36*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a*b^2*e^
(2*x) + 6*sqrt(a^2 - sqrt(-a*b)*a)*a^3*b - 12*sqrt(-a*b)*a^3*b + 5*sqrt(a^2
- sqrt(-a*b)*a)*a^2*b^2 - 16*sqrt(-a*b)*a^2*b^2 - 4*sqrt(a^2 - sqrt(-a*b)*
a)*a*b^3 - 9*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a^2*b - 12*sqrt(a^2 - sqrt
(-a*b)*a)*sqrt(-a*b)*a*b^2)) - 1/4*sqrt((a^2 + sqrt(-a*b)*a)/(a^4 + a^3*b))
*log(abs(60*a^4*b*e^(2*x) + 68*a^3*b^2*e^(2*x) - 16*a^2*b^3*e^(2*x) + 24*sq
rt(-a*b)*a^4*e^(2*x) - 48*sqrt(a^2 - sqrt(-a*b)*a)*a^3*b*e^(2*x) - 16*sqrt(
-a*b)*a^3*b*e^(2*x) - 61*sqrt(a^2 - sqrt(-a*b)*a)*a^2*b^2*e^(2*x) - 64*sqrt
(-a*b)*a^2*b^2*e^(2*x) + 4*sqrt(a^2 - sqrt(-a*b)*a)*a*b^3*e^(2*x) + 6*a^4*b
+ 2*a^3*b^2 - 8*a^2*b^3 - 24*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a^3*e^(2*
x) - 5*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a^2*b*e^(2*x) + 36*sqrt(a^2 - sq
rt(-a*b)*a)*sqrt(-a*b)*a*b^2*e^(2*x) - 6*sqrt(a^2 - sqrt(-a*b)*a)*a^3*b - 1
2*sqrt(-a*b)*a^3*b - 5*sqrt(a^2 - sqrt(-a*b)*a)*a^2*b^2 - 16*sqrt(-a*b)*a^2
*b^2 + 4*sqrt(a^2 - sqrt(-a*b)*a)*a*b^3 + 9*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-
a*b)*a^2*b + 12*sqrt(a^2 - sqrt(-a*b)*a)*sqrt(-a*b)*a*b^2)) + 1/4*sqrt((a^2
- sqrt(-a*b)*a)/(a^4 + a^3*b))*log(abs(60*a^4*b*e^(2*x) + 68*a^3*b^2*e^(2*
x) - 16*a^2*b^3*e^(2*x) - 24*sqrt(-a*b)*a^4*e^(2*x) + 48*sqrt(a^2 + sqrt(-a
*b)*a)*a^3*b*e^(2*x) + 16*sqrt(-a*b)*a^3*b*e^(2*x) + 61*sqrt(a^2 + sqrt(-a*
b)*a)*a^2*b^2*e^(2*x) + 64*sqrt(-a*b)*a^2*b^2*e^(2*x) - 4*sqrt(a^2 + sqrt(-
a*b)*a)*a*b^3*e^(2*x) + 6*a^4*b + 2*a^3*b^2 - 8*a^2*b^3 - 24*sqrt(a^2 + sqr
t(-a*b)*a)*sqrt(-a*b)*a^3*e^(2*x) - 5*sqrt(a^2 + sqrt(-a*b)*a)*sqrt(-a*b)*a
^2*b*e^(2*x) + 36*sqrt(a^2 + sqrt(-a*b)*a)*sqrt(-a*b)*a*b^2*e^(2*x) + 6*sqr
t(a^2 + sqrt(-a*b)*a)*a^3*b + 12*sqrt(-a*b)*a^3*b + 5*sqrt(a^2 + sqrt(-a*b)
*a)*a^2*b^2 + 16*sqrt(-a*b)*a^2*b^2 - 4*sqrt(a^2 + sqrt(-a*b)*a)*a*b^3 + 9*
sqrt(a^2 + sqrt(-a*b)*a)*sqrt(-a*b)*a^2*b + 12*sqrt(a^2 + sqrt(-a*b)*a)*sqr
t(-a*b)*a*b^2)) - 1/4*sqrt((a^2 - sqrt(-a*b)*a)/(a^4 + a^3*b))*log(abs(60*a
^4*b*e^(2*x) + 68*a^3*b^2*e^(2*x) - 16*a^2*b^3*e^(2*x) - 24*sqrt(-a*b)*a^4*
e^(2*x) - 48*sqrt(a^2 + sqrt(-a*b)*a)*a^3*b*e^(2*x) + 16*sqrt(-a*b)*a^3*b*e
^(2*x) - 61*sqrt(a^2 + sqrt(-a*b)*a)*a^2*b^2*e^(2*x) + 64*sqrt(-a*b)*a^2*b^
```

$2e^{(2*x)} + 4\sqrt{a^2 + \sqrt{-a*b}*a}*a*b^3e^{(2*x)} + 6*a^4*b + 2*a^3*b^2 - 8*a^2*b^3 + 24*\sqrt{a^2 + \sqrt{-a*b}*a}*\sqrt{-a*b}*a^3e^{(2*x)} + 5*\sqrt{a^2 + \sqrt{-a*b}*a}*\sqrt{-a*b}*a^2*b*e^{(2*x)} - 36*\sqrt{a^2 + \sqrt{-a*b}*a}*\sqrt{-a*b}*a*b^2e^{(2*x)} - 6*\sqrt{a^2 + \sqrt{-a*b}*a}*a^3*b + 12*\sqrt{-a*b}*a^3*b - 5*\sqrt{a^2 + \sqrt{-a*b}*a}*a^2*b^2 + 16*\sqrt{-a*b}*a^2*b^2 + 4*\sqrt{a^2 + \sqrt{-a*b}*a}*a*b^3 - 9*\sqrt{a^2 + \sqrt{-a*b}*a}*\sqrt{-a*b}*a^2*b - 12*\sqrt{a^2 + \sqrt{-a*b}*a}*\sqrt{-a*b}*a*b^2$

Mupad [B] (verification not implemented)

Time = 8.77 (sec) , antiderivative size = 1563, normalized size of antiderivative = 4.33

$$\int \frac{1}{a + b \cosh^4(x)} dx = \text{Too large to display}$$

[In] int(1/(a + b*cosh(x)^4),x)

[Out] $\log\left(\frac{524288*(1024*a^3*\exp(2*x) - 35*b^3*\exp(2*x) + 185*a*b^2 + 464*a^2*b + 256*a^3 - 24*b^3 + 988*a*b^2*\exp(2*x) + 2048*a^2*b*\exp(2*x))}{(a*b^6*(a + b)^2) - \left(\frac{4194304*(253*a*b^3 - b^4*\exp(2*x) + 1184*a^3*b + 512*a^4 - b^4 + 930*a^2*b^2 + 1392*a^2*b^2*\exp(2*x) + 627*a*b^3*\exp(2*x) + 768*a^3*b*\exp(2*x))}{(b^6*(a + b)^2) + (8388608*a*((a^2 + (-a^3*b)^{1/2}))/a^3*(a + b))}^{1/2}*(512*a^3*\exp(2*x) - 6*b^3*\exp(2*x) + 181*a*b^2 + 432*a^2*b + 256*a^3 - 5*b^3 + 622*a*b^2*\exp(2*x) + 1152*a^2*b*\exp(2*x))}{(b^6*(a + b))}*(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}^{1/2}\right)/4 - \frac{2097152*(176*a*b + 1536*a^2*\exp(2*x) - 134*b^2*\exp(2*x) + 256*a^2 - 75*b^2 + 1408*a*b*\exp(2*x))}{(b^6*(a + b))}*(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}^{1/2}\right)/4 * \frac{(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}{(16*(a^3*b + a^4))}^{1/2} - \log\left(\frac{524288*(1024*a^3*\exp(2*x) - 35*b^3*\exp(2*x) + 185*a*b^2 + 464*a^2*b + 256*a^3 - 24*b^3 + 988*a*b^2*\exp(2*x) + 2048*a^2*b*\exp(2*x))}{(a*b^6*(a + b)^2) - \left(\frac{4194304*(253*a*b^3 - b^4*\exp(2*x) + 1184*a^3*b + 512*a^4 - b^4 + 930*a^2*b^2 + 1392*a^2*b^2*\exp(2*x) + 627*a*b^3*\exp(2*x) + 768*a^3*b*\exp(2*x))}{(b^6*(a + b)^2) - (8388608*a*((a^2 + (-a^3*b)^{1/2}))/a^3*(a + b))}^{1/2}*(512*a^3*\exp(2*x) - 6*b^3*\exp(2*x) + 181*a*b^2 + 432*a^2*b + 256*a^3 - 5*b^3 + 622*a*b^2*\exp(2*x) + 1152*a^2*b*\exp(2*x))}{(b^6*(a + b))}*(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}^{1/2}\right)/4 + \frac{2097152*(176*a*b + 1536*a^2*\exp(2*x) - 134*b^2*\exp(2*x) + 256*a^2 - 75*b^2 + 1408*a*b*\exp(2*x))}{(b^6*(a + b))}*(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}^{1/2}\right)/4 * \frac{(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}{(16*(a^3*b + a^4))}^{1/2} - \log\left(\frac{524288*(1024*a^3*\exp(2*x) - 35*b^3*\exp(2*x) + 185*a*b^2 + 464*a^2*b + 256*a^3 - 24*b^3 + 988*a*b^2*\exp(2*x) + 2048*a^2*b*\exp(2*x))}{(a*b^6*(a + b)^2) - \left(\frac{4194304*(253*a*b^3 - b^4*\exp(2*x) + 1184*a^3*b + 512*a^4 - b^4 + 930*a^2*b^2 + 1392*a^2*b^2*\exp(2*x) + 627*a*b^3*\exp(2*x) + 768*a^3*b*\exp(2*x))}{(b^6*(a + b)^2) - (8388608*a*((a^2 + (-a^3*b)^{1/2}))/a^3*(a + b))}^{1/2}*(512*a^3*\exp(2*x) - 6*b^3*\exp(2*x) + 181*a*b^2 + 432*a^2*b + 256*a^3 - 5*b^3 + 622*a*b^2*\exp(2*x) + 1152*a^2*b*\exp(2*x))}{(b^6*(a + b))}*(a^2 + (-a^3*b)^{1/2})/a^3*(a + b)}^{1/2}\right)$

$$\begin{aligned}
& 1/2)) / (a^3(a + b))^{(1/2)} / 4 + (2097152 * (176 * a * b + 1536 * a^2 * \exp(2 * x) - 134 \\
& * b^2 * \exp(2 * x) + 256 * a^2 - 75 * b^2 + 1408 * a * b * \exp(2 * x))) / (b^6(a + b)) * ((a^2 \\
& - (-a^3 * b)^{(1/2)}) / (a^3(a + b))^{(1/2)}) / 4 * ((a^2 - (-a^3 * b)^{(1/2)}) / (16 * (a^3 * b + a^4)))^{(1/2)} + \log((524288 * (1024 * a^3 * \exp(2 * x) - 35 * b^3 * \exp(2 * x) + 185 \\
& * a * b^2 + 464 * a^2 * b + 256 * a^3 - 24 * b^3 + 988 * a * b^2 * \exp(2 * x) + 2048 * a^2 * b * \exp(2 * x))) / (a * b^6 * (a + b)^2) - (((((4194304 * (253 * a * b^3 - b^4 * \exp(2 * x) + 1184 * a^3 * b + 512 * a^4 - b^4 + 930 * a^2 * b^2 + 1392 * a^2 * b^2 * \exp(2 * x) + 627 * a * b^3 * \exp(2 * x) + 768 * a^3 * b * \exp(2 * x)))) / (b^6 * (a + b)^2) + (8388608 * a * ((a^2 - (-a^3 * b)^{(1/2)}) / (a^3(a + b))^{(1/2)}) / (a^3(a + b))^{(1/2)}) * (512 * a^3 * \exp(2 * x) - 6 * b^3 * \exp(2 * x) + 181 * a * b^2 + 432 * a^2 * b + 256 * a^3 - 5 * b^3 + 622 * a * b^2 * \exp(2 * x) + 1152 * a^2 * b * \exp(2 * x))) / (b^6 * (a + b)) * ((a^2 - (-a^3 * b)^{(1/2)}) / (a^3(a + b))^{(1/2)}) / 4 - (2097152 * (176 * a * b + 1536 * a^2 * \exp(2 * x) - 134 * b^2 * \exp(2 * x) + 256 * a^2 - 75 * b^2 + 1408 * a * b * \exp(2 * x))) / (b^6 * (a + b)) * ((a^2 - (-a^3 * b)^{(1/2)}) / (a^3(a + b))^{(1/2)}) / 4) * ((a^2 - (-a^3 * b)^{(1/2)}) / (16 * (a^3 * b + a^4)))^{(1/2)}
\end{aligned}$$

3.61 $\int \frac{1}{a-b \cosh^4(x)} dx$

Optimal result	402
Rubi [A] (verified)	402
Mathematica [A] (verified)	403
Maple [C] (verified)	404
Fricas [B] (verification not implemented)	404
Sympy [F(-1)]	406
Maxima [F]	406
Giac [B] (verification not implemented)	407
Mupad [B] (verification not implemented)	408

Optimal result

Integrand size = 11, antiderivative size = 101

$$\int \frac{1}{a-b \cosh^4(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}-\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}-\sqrt{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}+\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}+\sqrt{b}}}$$

[Out] $\frac{1}{2} \operatorname{arctanh}\left(\frac{a^{1/4} \tanh(x)}{(\sqrt{a}-\sqrt{b})^{1/2}}\right) / a^{3/4} / (\sqrt{a}-\sqrt{b})^{1/2} + \frac{1}{2} \operatorname{arctanh}\left(\frac{a^{1/4} \tanh(x)}{(\sqrt{a}+\sqrt{b})^{1/2}}\right) / a^{3/4} / (\sqrt{a}+\sqrt{b})^{1/2}$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3288, 1180, 214}

$$\int \frac{1}{a-b \cosh^4(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}-\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}-\sqrt{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[4]{a} \tanh(x)}{\sqrt{\sqrt{a}+\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}+\sqrt{b}}}$$

[In] $\operatorname{Int}[(a - b \operatorname{Cosh}[x]^4)^{-1}, x]$

[Out] $\operatorname{ArcTanh}\left[\frac{a^{1/4} \operatorname{Tanh}[x]}{\sqrt{\sqrt{a}-\sqrt{b}}}\right] / (2a^{3/4} \sqrt{\sqrt{a}-\sqrt{b}}) + \operatorname{ArcTanh}\left[\frac{a^{1/4} \operatorname{Tanh}[x]}{\sqrt{\sqrt{a}+\sqrt{b}}}\right] / (2a^{3/4} \sqrt{\sqrt{a}+\sqrt{b}})$

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 1180

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :
> With[{q = Rt[b^2 - 4*a*c, 2]}, Dist[e/2 + (2*c*d - b*e)/(2*q), Int[1/(b/2
- q/2 + c*x^2), x], x] + Dist[e/2 - (2*c*d - b*e)/(2*q), Int[1/(b/2 + q/2
+ c*x^2), x], x]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && Ne
Q[c*d^2 - a*e^2, 0] && PosQ[b^2 - 4*a*c]
```

Rule 3288

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^4)^(p_), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + 2*a*ff^2*x^2 + (a
+ b)*ff^4*x^4)^p/(1 + ff^2*x^2)^(2*p + 1), x], x, Tan[e + f*x]/ff], x]] /;
FreeQ[{a, b, e, f}, x] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1-x^2}{a-2ax^2+(a-b)x^4} dx, x, \coth(x)\right) \\
&= \frac{1}{2}\left(-1 - \frac{\sqrt{b}}{\sqrt{a}}\right) \text{Subst}\left(\int \frac{1}{-a - \sqrt{a}\sqrt{b} + (a-b)x^2} dx, x, \coth(x)\right) \\
&\quad + \frac{1}{2}\left(-1 + \frac{\sqrt{b}}{\sqrt{a}}\right) \text{Subst}\left(\int \frac{1}{-a + \sqrt{a}\sqrt{b} + (a-b)x^2} dx, x, \coth(x)\right) \\
&= \frac{\arctanh\left(\frac{\sqrt[4]{a}\tanh(x)}{\sqrt{\sqrt{a}-\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}-\sqrt{b}}} + \frac{\arctanh\left(\frac{\sqrt[4]{a}\tanh(x)}{\sqrt{\sqrt{a}+\sqrt{b}}}\right)}{2a^{3/4}\sqrt{\sqrt{a}+\sqrt{b}}}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.08

$$\int \frac{1}{a-b\cosh^4(x)} dx = -\frac{\arctan\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{-a+\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{a}\sqrt{-a+\sqrt{a}\sqrt{b}}} + \frac{\arctanh\left(\frac{\sqrt{a}\tanh(x)}{\sqrt{a+\sqrt{a}\sqrt{b}}}\right)}{2\sqrt{a}\sqrt{a+\sqrt{a}\sqrt{b}}}$$

```
[In] Integrate[(a - b*Cosh[x]^4)^(-1), x]
```

```
[Out] -1/2*ArcTan[(Sqrt[a]*Tanh[x])/Sqrt[-a + Sqrt[a]*Sqrt[b]]]/(Sqrt[a]*Sqrt[-a
+ Sqrt[a]*Sqrt[b]]) + ArcTanh[(Sqrt[a]*Tanh[x])/Sqrt[a + Sqrt[a]*Sqrt[b]]]/
(2*Sqrt[a]*Sqrt[a + Sqrt[a]*Sqrt[b]])
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.34 (sec) , antiderivative size = 96, normalized size of antiderivative = 0.95

method	result
risch	$\sum_{R=\text{RootOf}(1+(256a^4-256a^3b)Z^4-32a^2Z^2)} _R \ln \left(e^{2x} + \left(\frac{128a^4}{b} - 128a^3 \right) _R^3 + \left(-\frac{32a^3}{b} + 32a^2 \right) _R^2 + \right.$
default	$\left(\sum_{R=\text{RootOf}((a-b)Z^8+(-4a-4b)Z^6+(6a-6b)Z^4+(-4a-4b)Z^2+a-b)} \frac{(-R^6+3R^4-3R^2+1) \ln(\tanh(\frac{x}{2})-R)}{_R^7 _R^5 _R^3 _R^2 _R^1} \right) \frac{1}{4}$

[In] int(1/(a-b*cosh(x)^4),x,method=_RETURNVERBOSE)

[Out] sum(_R*ln(exp(2*x)+(128*a^4/b-128*a^3)*_R^3+(-32/b*a^3+32*a^2)*_R^2+(-8*a^2/b-8*a)*_R+2*a/b+1),_R=RootOf(1+(256*a^4-256*a^3*b)*_Z^4-32*a^2*_Z^2))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 779 vs. 2(65) = 130.

Time = 0.30 (sec) , antiderivative size = 779, normalized size of antiderivative = 7.71

$$\begin{aligned}
& \int \frac{1}{a - b \cosh^4(x)} dx \\
&= -\frac{1}{4} \sqrt{\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + 1}{a^2 - ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
&\quad \left. + 2 \left(ab - (a^4 - a^3b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} \right) \sqrt{\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + 1}{a^2 - ab}} \right. \\
&\quad \left. - 2(a^3 - a^2b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + b \right) \\
&+ \frac{1}{4} \sqrt{\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + 1}{a^2 - ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
&\quad \left. - 2 \left(ab - (a^4 - a^3b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} \right) \sqrt{\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + 1}{a^2 - ab}} \right. \\
&\quad \left. - 2(a^3 - a^2b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + b \right) \\
&- \frac{1}{4} \sqrt{-\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} - 1}{a^2 - ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
&\quad \left. + 2 \left(ab + (a^4 - a^3b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} \right) \sqrt{-\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} - 1}{a^2 - ab}} \right. \\
&\quad \left. + 2(a^3 - a^2b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + b \right) \\
&+ \frac{1}{4} \sqrt{-\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} - 1}{a^2 - ab}} \log \left(b \cosh(x)^2 + 2b \cosh(x) \sinh(x) + b \sinh(x)^2 \right. \\
&\quad \left. - 2 \left(ab + (a^4 - a^3b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} \right) \sqrt{-\frac{(a^2 - ab) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} - 1}{a^2 - ab}} \right. \\
&\quad \left. + 2(a^3 - a^2b) \sqrt{\frac{b}{a^5 - 2a^4b + a^3b^2}} + b \right)
\end{aligned}$$

[In] integrate(1/(a-b*cosh(x)^4),x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/4*\sqrt{((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} + 1)/(a^2 - a*b)} * \\ & \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*(a*b - (a^4 - a^3*b) \\ &)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)})) * \sqrt{((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b \\ & + a^3*b^2)} + 1)/(a^2 - a*b)} - 2*(a^3 - a^2*b)*\sqrt{b/(a^5 - 2*a^4*b + a \\ & ^3*b^2)} + b) + 1/4*\sqrt{((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} + 1 \\ &)/(a^2 - a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 - 2*(a*b \\ & - (a^4 - a^3*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)})) * \sqrt{((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b \\ & + a^3*b^2)} + 1)/(a^2 - a*b)} - 2*(a^3 - a^2*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} + b) - 1/4*\sqrt{-((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b \\ & + a^3*b^2)} - 1)/(a^2 - a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x)*\sinh(x) + b*\sinh(x)^2 + 2*(a*b + (a^4 - a^3*b) \\ &)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)})) * \sqrt{-((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} - 1)/(a^2 - a*b)} + 2*(a^3 - \\ & a^2*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} + b) + 1/4*\sqrt{-((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} - 1)/(a^2 - a*b)} * \log(b*\cosh(x)^2 + 2*b*\cosh(x) \\ &)*\sinh(x) + b*\sinh(x)^2 - 2*(a*b + (a^4 - a^3*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)})) * \sqrt{-((a^2 - a*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} - 1)/(a^2 - \\ & a*b)} + 2*(a^3 - a^2*b)*\sqrt{b/(a^5 - 2*a^4*b + a^3*b^2)} + b) \end{aligned}$$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{a - b \cosh^4(x)} dx = \text{Timed out}$$

[In] integrate(1/(a-b*cosh(x)**4),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{a - b \cosh^4(x)} dx = \int -\frac{1}{b \cosh(x)^4 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^4),x, algorithm="maxima")

[Out] -integrate(1/(b*cosh(x)^4 - a), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1697 vs. 2(65) = 130.

Time = 2.46 (sec) , antiderivative size = 1697, normalized size of antiderivative = 16.80

$$\int \frac{1}{a - b \cosh^4(x)} dx = \text{Too large to display}$$

[In] integrate(1/(a-b*cosh(x)^4),x, algorithm="giac")

[Out] $\frac{1}{4} \sqrt{(a^2 - \sqrt{a*b}*a)/(a^4 - a^3*b)} \log(\text{abs}(60*a^4*b*e^{(2*x)} - 68*a^3*b^2*e^{(2*x)} - 16*a^2*b^3*e^{(2*x)} + 24*\sqrt{a*b}*a^4*e^{(2*x)} + 48*\sqrt{a^2 + \sqrt{a*b}*a}*a^3*b*e^{(2*x)} + 16*\sqrt{a*b}*a^3*b*e^{(2*x)} - 61*\sqrt{a^2 + \sqrt{a*b}*a}*a^2*b^2*e^{(2*x)} - 64*\sqrt{a*b}*a^2*b^2*e^{(2*x)} - 4*\sqrt{a^2 + \sqrt{a*b}*a}*a*b^3*e^{(2*x)} + 6*a^4*b - 2*a^3*b^2 - 8*a^2*b^3 + 24*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^3*e^{(2*x)} - 5*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b*e^{(2*x)} - 36*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2*e^{(2*x)} + 6*\sqrt{a^2 + \sqrt{a*b}*a}*a^3*b + 12*\sqrt{a*b}*a^3*b - 5*\sqrt{a^2 + \sqrt{a*b}*a}*a^2*b^2 - 16*\sqrt{a*b}*a^2*b^2 - 4*\sqrt{a^2 + \sqrt{a*b}*a}*a*b^3 + 9*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b - 12*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2)) - \frac{1}{4} \sqrt{(a^2 - \sqrt{a*b}*a)/(a^4 - a^3*b)} \log(\text{abs}(60*a^4*b*e^{(2*x)} - 68*a^3*b^2*e^{(2*x)} - 16*a^2*b^3*e^{(2*x)} + 24*\sqrt{a*b}*a^4*e^{(2*x)} - 48*\sqrt{a^2 + \sqrt{a*b}*a}*a^3*b*e^{(2*x)} + 16*\sqrt{a*b}*a^3*b*e^{(2*x)} + 61*\sqrt{a^2 + \sqrt{a*b}*a}*a^2*b^2*e^{(2*x)} - 64*\sqrt{a*b}*a^2*b^2*e^{(2*x)} + 4*\sqrt{a^2 + \sqrt{a*b}*a}*a*b^3*e^{(2*x)} + 6*a^4*b - 2*a^3*b^2 - 8*a^2*b^3 - 24*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^3*e^{(2*x)} + 5*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b*e^{(2*x)} + 36*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2*e^{(2*x)} - 6*\sqrt{a^2 + \sqrt{a*b}*a}*a^3*b + 12*\sqrt{a*b}*a^3*b + 5*\sqrt{a^2 + \sqrt{a*b}*a}*a^2*b^2 - 16*\sqrt{a*b}*a^2*b^2 + 4*\sqrt{a^2 + \sqrt{a*b}*a}*a*b^3 - 9*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b + 12*\sqrt{a^2 + \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2)) + \frac{1}{4} \sqrt{(a^2 + \sqrt{a*b}*a)/(a^4 - a^3*b)} \log(\text{abs}(60*a^4*b*e^{(2*x)} - 68*a^3*b^2*e^{(2*x)} - 16*a^2*b^3*e^{(2*x)} - 24*\sqrt{a*b}*a^4*e^{(2*x)} + 48*\sqrt{a^2 - \sqrt{a*b}*a}*a^3*b*e^{(2*x)} - 16*\sqrt{a*b}*a^3*b*e^{(2*x)} - 61*\sqrt{a^2 - \sqrt{a*b}*a}*a^2*b^2*e^{(2*x)} + 64*\sqrt{a*b}*a^2*b^2*e^{(2*x)} - 4*\sqrt{a^2 - \sqrt{a*b}*a}*a*b^3*e^{(2*x)} + 6*a^4*b - 2*a^3*b^2 - 8*a^2*b^3 - 24*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^3*e^{(2*x)} + 5*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b*e^{(2*x)} + 36*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2*e^{(2*x)} + 6*\sqrt{a^2 - \sqrt{a*b}*a}*a^3*b - 12*\sqrt{a*b}*a^3*b - 5*\sqrt{a^2 - \sqrt{a*b}*a}*a^2*b^2 + 16*\sqrt{a*b}*a^2*b^2 - 4*\sqrt{a^2 - \sqrt{a*b}*a}*a*b^3 - 9*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b + 12*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2)) - \frac{1}{4} \sqrt{(a^2 + \sqrt{a*b}*a)/(a^4 - a^3*b)} \log(\text{abs}(60*a^4*b*e^{(2*x)} - 68*a^3*b^2*e^{(2*x)} - 16*a^2*b^3*e^{(2*x)} - 24*\sqrt{a*b}*a^4*e^{(2*x)} - 48*\sqrt{a^2 - \sqrt{a*b}*a}*a^3*b*e^{(2*x)} - 16*\sqrt{a*b}*a^3*b*e^{(2*x)} + 61*\sqrt{a^2 - \sqrt{a*b}*a}*a^2*b^2*e^{(2*x)} + 64*\sqrt{a*b}*a^2*b^2*e^{(2*x)} + 4*\sqrt{a^2 - \sqrt{a*b}*a}*a*b^3*e^{(2*x)} + 6*a^4*b - 2*a^3*b^2$

$$2 - 8*a^2*b^3 + 24*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^3*e^{(2*x)} - 5*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b*e^{(2*x)} - 36*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2*e^{(2*x)} - 6*\sqrt{a^2 - \sqrt{a*b}*a}*a^3*b - 12*\sqrt{a*b}*a^3*b + 5*\sqrt{a^2 - \sqrt{a*b}*a}*a^2*b^2 + 16*\sqrt{a*b}*a^2*b^2 + 4*\sqrt{a^2 - \sqrt{a*b}*a}*a*b^3 + 9*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a^2*b - 12*\sqrt{a^2 - \sqrt{a*b}*a}*\sqrt{a*b}*a*b^2)$$

Mupad [B] (verification not implemented)

Time = 10.27 (sec) , antiderivative size = 1487, normalized size of antiderivative = 14.72

$$\int \frac{1}{a - b \cosh^4(x)} dx = \text{Too large to display}$$

[In] int(1/(a - b*cosh(x)^4),x)

[Out] log((((1/(a^2 - (a^3*b)^(1/2)))^(1/2)*((4194304*(b^4*exp(2*x) + 253*a*b^3 + 1184*a^3*b - 512*a^4 + b^4 - 930*a^2*b^2 - 1392*a^2*b^2*exp(2*x) + 627*a*b^3*exp(2*x) + 768*a^3*b*exp(2*x)))/(b^6*(a - b)^2) - (8388608*a*(1/(a^2 - (a^3*b)^(1/2)))^(1/2)*(512*a^3*exp(2*x) + 6*b^3*exp(2*x) + 181*a*b^2 - 432*a^2*b + 256*a^3 + 5*b^3 + 622*a*b^2*exp(2*x) - 1152*a^2*b*exp(2*x)))/(b^6*(a - b))))/4 - (2097152*(176*a*b - 1536*a^2*exp(2*x) + 134*b^2*exp(2*x) - 256*a^2 + 75*b^2 + 1408*a*b*exp(2*x)))/(b^6*(a - b)))*(1/(a^2 - (a^3*b)^(1/2)))^(1/2))/4 + (524288*(1024*a^3*exp(2*x) + 35*b^3*exp(2*x) + 185*a*b^2 - 464*a^2*b + 256*a^3 + 24*b^3 + 988*a*b^2*exp(2*x) - 2048*a^2*b*exp(2*x)))/(a*b^6*(a - b)^2))*(-(a^2 + (a^3*b)^(1/2))/(16*(a^3*b - a^4)))^(1/2) - log((((1/(a^2 - (a^3*b)^(1/2)))^(1/2)*((4194304*(b^4*exp(2*x) + 253*a*b^3 + 1184*a^3*b - 512*a^4 + b^4 - 930*a^2*b^2 - 1392*a^2*b^2*exp(2*x) + 627*a*b^3*exp(2*x) + 768*a^3*b*exp(2*x)))/(b^6*(a - b)^2) + (8388608*a*(1/(a^2 - (a^3*b)^(1/2)))^(1/2)*(512*a^3*exp(2*x) + 6*b^3*exp(2*x) + 181*a*b^2 - 432*a^2*b + 256*a^3 + 5*b^3 + 622*a*b^2*exp(2*x) - 1152*a^2*b*exp(2*x)))/(b^6*(a - b)))/4 + (2097152*(176*a*b - 1536*a^2*exp(2*x) + 134*b^2*exp(2*x) - 256*a^2 + 75*b^2 + 1408*a*b*exp(2*x)))/(b^6*(a - b)))*(1/(a^2 - (a^3*b)^(1/2)))^(1/2))/4 + (524288*(1024*a^3*exp(2*x) + 35*b^3*exp(2*x) + 185*a*b^2 - 464*a^2*b + 256*a^3 + 24*b^3 + 988*a*b^2*exp(2*x) - 2048*a^2*b*exp(2*x)))/(a*b^6*(a - b)^2))*(-(a^2 + (a^3*b)^(1/2))/(16*(a^3*b - a^4)))^(1/2) + log((((1/(a^2 + (a^3*b)^(1/2)))^(1/2)*((4194304*(b^4*exp(2*x) + 253*a*b^3 + 1184*a^3*b - 512*a^4 + b^4 - 930*a^2*b^2 - 1392*a^2*b^2*exp(2*x) + 627*a*b^3*exp(2*x) + 768*a^3*b*exp(2*x)))/(b^6*(a - b)^2) - (8388608*a*(1/(a^2 + (a^3*b)^(1/2)))^(1/2)*(512*a^3*exp(2*x) + 6*b^3*exp(2*x) + 181*a*b^2 - 432*a^2*b + 256*a^3 + 5*b^3 + 622*a*b^2*exp(2*x) - 1152*a^2*b*exp(2*x)))/(b^6*(a - b)))/4 - (2097152*(176*a*b - 1536*a^2*exp(2*x) + 134*b^2*exp(2*x) - 256*a^2 + 75*b^2 + 1408*a*b*exp(2*x)))/(b^6*(a - b)))/4 + (524288*(1024*a^3*exp(2*x) + 35*b^3*exp(2*x) + 185*a*b^2 - 464*a^2*b + 256*a^3 + 24*b^3 + 988*a*b^2*exp(2*x) - 2048*a^2*b*exp(2*x)))/(a*b^6*(a - b)^2)

$$\begin{aligned}
&) * (- (a^2 - (a^3 * b)^{(1/2)}) / (16 * (a^3 * b - a^4)))^{(1/2)} - \log(((1 / (a^2 + (a^3 * b)^{(1/2)}))^{(1/2)})^{(1/2)} * (((1 / (a^2 + (a^3 * b)^{(1/2)}))^{(1/2)})^{(1/2)} * ((4194304 * (b^4 * \exp(2 * x)) + 253 * a * b^3 + 1184 * a^3 * b - 512 * a^4 + b^4 - 930 * a^2 * b^2 - 1392 * a^2 * b^2 * \exp(2 * x) + 627 * a * b^3 * \exp(2 * x) + 768 * a^3 * b * \exp(2 * x))) / (b^6 * (a - b)^2) + (8388608 * a * (1 / (a^2 + (a^3 * b)^{(1/2)}))^{(1/2)} * (512 * a^3 * \exp(2 * x) + 6 * b^3 * \exp(2 * x) + 181 * a * b^2 - 432 * a^2 * b + 256 * a^3 + 5 * b^3 + 622 * a * b^2 * \exp(2 * x) - 1152 * a^2 * b * \exp(2 * x))) / (b^6 * (a - b)))) / 4 + (2097152 * (176 * a * b - 1536 * a^2 * \exp(2 * x) + 134 * b^2 * \exp(2 * x) - 256 * a^2 + 75 * b^2 + 1408 * a * b * \exp(2 * x))) / (b^6 * (a - b))) / 4 + (524288 * (1024 * a^3 * \exp(2 * x) + 35 * b^3 * \exp(2 * x) + 185 * a * b^2 - 464 * a^2 * b + 256 * a^3 + 24 * b^3 + 988 * a * b^2 * \exp(2 * x) - 2048 * a^2 * b * \exp(2 * x))) / (a * b^6 * (a - b)^2)) * (- (a^2 - (a^3 * b)^{(1/2)}) / (16 * (a^3 * b - a^4)))^{(1/2)}
\end{aligned}$$

3.62 $\int \frac{1}{1+\cosh^4(x)} dx$

Optimal result	410
Rubi [A] (verified)	410
Mathematica [C] (verified)	413
Maple [C] (verified)	413
Fricas [C] (verification not implemented)	414
Sympy [F(-1)]	414
Maxima [F]	415
Giac [C] (verification not implemented)	415
Mupad [B] (verification not implemented)	416

Optimal result

Integrand size = 8, antiderivative size = 176

$$\int \frac{1}{1+\cosh^4(x)} dx = -\frac{\arctan\left(\frac{\sqrt{1+\sqrt{2}}-2\coth(x)}{\sqrt{-1+\sqrt{2}}}\right)}{4\sqrt{1+\sqrt{2}}} + \frac{\arctan\left(\frac{\sqrt{1+\sqrt{2}}+2\coth(x)}{\sqrt{-1+\sqrt{2}}}\right)}{4\sqrt{1+\sqrt{2}}} \\ - \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(\sqrt{2}-2\sqrt{1+\sqrt{2}}\coth(x)+2\coth^2(x)\right) \\ + \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(1+\sqrt{2}\left(1+\sqrt{2}\right)\coth(x)+\sqrt{2}\coth^2(x)\right)$$

[Out] $-1/4*\arctan((-2*\coth(x)+(1+2^{(1/2)})^{(1/2)})/(2^{(1/2)}-1)^{(1/2)})/(1+2^{(1/2)})^{(1/2)}+1/4*\arctan((2*\coth(x)+(1+2^{(1/2)})^{(1/2)})/(2^{(1/2)}-1)^{(1/2)})/(1+2^{(1/2)})^{(1/2)}-1/8*\ln(2*\coth(x)^2+2^{(1/2)}-2*\coth(x)*(1+2^{(1/2)})^{(1/2)})*(1+2^{(1/2)})^{(1/2)}+1/8*\ln(1+\coth(x)^2*2^{(1/2)}+\coth(x)*(2+2*2^{(1/2)})^{(1/2)})*(1+2^{(1/2)})^{(1/2)}$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 176, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {3288, 1183, 648, 632, 210, 642}

$$\int \frac{1}{1+\cosh^4(x)} dx = -\frac{\arctan\left(\frac{\sqrt{1+\sqrt{2}}-2\coth(x)}{\sqrt{\sqrt{2}-1}}\right)}{4\sqrt{1+\sqrt{2}}} + \frac{\arctan\left(\frac{2\coth(x)+\sqrt{1+\sqrt{2}}}{\sqrt{\sqrt{2}-1}}\right)}{4\sqrt{1+\sqrt{2}}} \\ - \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(2\coth^2(x)-2\sqrt{1+\sqrt{2}}\coth(x)+\sqrt{2}\right) \\ + \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(\sqrt{2}\coth^2(x)+\sqrt{2}\left(1+\sqrt{2}\right)\coth(x)+1\right)$$

[In] Int[(1 + Cosh[x]^4)^(-1), x]

[Out] $-\frac{1}{4} \operatorname{ArcTan}\left[\frac{\sqrt{1 + \sqrt{2}} - 2 \operatorname{Coth}[x]}{\sqrt{-1 + \sqrt{2}}}\right] / \sqrt{1 + \sqrt{2}} + \operatorname{ArcTan}\left[\frac{\sqrt{1 + \sqrt{2}} + 2 \operatorname{Coth}[x]}{\sqrt{-1 + \sqrt{2}}}\right] / (4 \sqrt{1 + \sqrt{2}}) - (\sqrt{1 + \sqrt{2}} \operatorname{Log}[\sqrt{2} - 2 \sqrt{1 + \sqrt{2}}] \operatorname{Coth}[x] + 2 \operatorname{Coth}[x]^2) / 8 + (\sqrt{1 + \sqrt{2}} \operatorname{Log}[1 + \sqrt{2(1 + \sqrt{2})}] \operatorname{Coth}[x] + \sqrt{2} \operatorname{Coth}[x]^2) / 8$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 1183

Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]

Rule 3288

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^4)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + 2*a*ff^2*x^2 + (a + b)*ff^4*x^4)^p/(1 + ff^2*x^2)^(2*p + 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[p]

Rubi steps

$$\begin{aligned}
\text{integral} &= \text{Subst}\left(\int \frac{1-x^2}{1-2x^2+2x^4} dx, x, \coth(x)\right) \\
&= \frac{\text{Subst}\left(\int \frac{\frac{\sqrt{1+\sqrt{2}}-(1+\frac{1}{\sqrt{2}})x}{\frac{1}{\sqrt{2}}-\sqrt{1+\sqrt{2}x+x^2}}}{2\sqrt{2}(1+\sqrt{2})} dx, x, \coth(x)\right)}{2\sqrt{2}(1+\sqrt{2})} + \frac{\text{Subst}\left(\int \frac{\frac{\sqrt{1+\sqrt{2}}+(1+\frac{1}{\sqrt{2}})x}{\frac{1}{\sqrt{2}}+\sqrt{1+\sqrt{2}x+x^2}}}{2\sqrt{2}(1+\sqrt{2})} dx, x, \coth(x)\right)}{2\sqrt{2}(1+\sqrt{2})} \\
&= \frac{1}{8}\sqrt{3-2\sqrt{2}}\text{Subst}\left(\int \frac{1}{\frac{1}{\sqrt{2}}-\sqrt{1+\sqrt{2}x+x^2}} dx, x, \coth(x)\right) \\
&\quad + \frac{1}{8}\sqrt{3-2\sqrt{2}}\text{Subst}\left(\int \frac{1}{\frac{1}{\sqrt{2}}+\sqrt{1+\sqrt{2}x+x^2}} dx, x, \coth(x)\right) \\
&\quad - \frac{1}{8}\sqrt{1+\sqrt{2}}\text{Subst}\left(\int \frac{-\sqrt{1+\sqrt{2}}+2x}{\frac{1}{\sqrt{2}}-\sqrt{1+\sqrt{2}x+x^2}} dx, x, \coth(x)\right) \\
&\quad + \frac{1}{8}\sqrt{1+\sqrt{2}}\text{Subst}\left(\int \frac{\sqrt{1+\sqrt{2}}+2x}{\frac{1}{\sqrt{2}}+\sqrt{1+\sqrt{2}x+x^2}} dx, x, \coth(x)\right) \\
&= -\frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(\sqrt{2}-2\sqrt{1+\sqrt{2}}\coth(x)+2\coth^2(x)\right) \\
&\quad + \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(1+\sqrt{2(1+\sqrt{2})}\coth(x)+\sqrt{2}\coth^2(x)\right) \\
&\quad - \frac{1}{4}\sqrt{3-2\sqrt{2}}\text{Subst}\left(\int \frac{1}{1-\sqrt{2}-x^2} dx, x, -\sqrt{1+\sqrt{2}}+2\coth(x)\right) \\
&\quad - \frac{1}{4}\sqrt{3-2\sqrt{2}}\text{Subst}\left(\int \frac{1}{1-\sqrt{2}-x^2} dx, x, \sqrt{1+\sqrt{2}}+2\coth(x)\right) \\
&= -\frac{1}{4}\sqrt{-1+\sqrt{2}}\arctan\left(\frac{\sqrt{1+\sqrt{2}}-2\coth(x)}{\sqrt{-1+\sqrt{2}}}\right) \\
&\quad + \frac{1}{4}\sqrt{-1+\sqrt{2}}\arctan\left(\frac{\sqrt{1+\sqrt{2}}+2\coth(x)}{\sqrt{-1+\sqrt{2}}}\right) \\
&\quad - \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(\sqrt{2}-2\sqrt{1+\sqrt{2}}\coth(x)+2\coth^2(x)\right) \\
&\quad + \frac{1}{8}\sqrt{1+\sqrt{2}}\log\left(1+\sqrt{2(1+\sqrt{2})}\coth(x)+\sqrt{2}\coth^2(x)\right)
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.07 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.26

$$\int \frac{1}{1 + \cosh^4(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{2\sqrt{1-i}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{2\sqrt{1+i}}$$

[In] Integrate[(1 + Cosh[x]^4)^(-1),x]

[Out] ArcTanh[Tanh[x]/Sqrt[1 - I]]/(2*Sqrt[1 - I]) + ArcTanh[Tanh[x]/Sqrt[1 + I]]/(2*Sqrt[1 + I])

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.22 (sec) , antiderivative size = 36, normalized size of antiderivative = 0.20

method	result	size
risch	$\sum_{R=\text{RootOf}(512Z^4-32Z^2+1)} _R \ln(-256R^3 + 64R^2 + e^{2x} - 1)$	36
default	$\frac{\left(\sum_{R=\text{RootOf}(2Z^4-2Z^2+1)} _R \ln\left(2 \tanh\left(\frac{x}{2}\right) _R + \tanh\left(\frac{x}{2}\right)^2 + 1\right) \right)}{4}$	37

[In] int(1/(1+cosh(x)^4),x,method=_RETURNVERBOSE)

[Out] sum(_R*ln(-256*_R^3+64*_R^2+exp(2*x)-1),_R=RootOf(512*_Z^4-32*_Z^2+1))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.76

$$\int \frac{1}{1 + \cosh^4(x)} dx = -\frac{1}{8} \sqrt{2} \sqrt{i+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. + (i+1) \sqrt{2} \sqrt{i+1} + 2i+1 \right) + \frac{1}{8} \sqrt{2} \sqrt{i+1} \log \left(\cosh(x)^2 \right. \\ \left. + 2 \cosh(x) \sinh(x) + \sinh(x)^2 - (i+1) \sqrt{2} \sqrt{i+1} + 2i+1 \right) \\ - \frac{1}{8} \sqrt{2} \sqrt{-i+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 \right. \\ \left. - (i-1) \sqrt{2} \sqrt{-i+1} - 2i+1 \right) + \frac{1}{8} \sqrt{2} \sqrt{-i+1} \log \left(\cosh(x)^2 \right. \\ \left. + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + (i-1) \sqrt{2} \sqrt{-i+1} - 2i+1 \right)$$

[In] integrate(1/(1+cosh(x)^4),x, algorithm="fricas")

[Out] -1/8*sqrt(2)*sqrt(I + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + (I + 1)*sqrt(2)*sqrt(I + 1) + 2*I + 1) + 1/8*sqrt(2)*sqrt(I + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - (I + 1)*sqrt(2)*sqrt(I + 1) + 2*I + 1) - 1/8*sqrt(2)*sqrt(-I + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - (I - 1)*sqrt(2)*sqrt(-I + 1) - 2*I + 1) + 1/8*sqrt(2)*sqrt(-I + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + (I - 1)*sqrt(2)*sqrt(-I + 1) - 2*I + 1)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{1 + \cosh^4(x)} dx = \text{Timed out}$$

[In] integrate(1/(1+cosh(x)**4),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{1 + \cosh^4(x)} dx = \int \frac{1}{\cosh(x)^4 + 1} dx$$

[In] integrate(1/(1+cosh(x)^4),x, algorithm="maxima")

[Out] integrate(1/(cosh(x)^4 + 1), x)

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 281, normalized size of antiderivative = 1.60

$$\begin{aligned} \int \frac{1}{1 + \cosh^4(x)} dx = & -\left(\frac{1}{16}i + \frac{1}{16}\right) \sqrt{2\sqrt{2}-2} \left(-\frac{i}{\sqrt{2}-1} + 1\right) \log\left((20i+10)\sqrt{2}e^{(2x)}\right. \\ & \left.+ 10\sqrt{2}\sqrt{10\sqrt{2}+14} + 50\sqrt{2} - (2i-14)\sqrt{10\sqrt{2}+14}\right. \\ & \left.+ (28i+14)e^{(2x)} + 70\right) \\ & + \left(\frac{1}{16}i + \frac{1}{16}\right) \sqrt{2\sqrt{2}-2} \left(-\frac{i}{\sqrt{2}-1} + 1\right) \log\left((20i+10)\sqrt{2}e^{(2x)}\right. \\ & \left.- 10\sqrt{2}\sqrt{10\sqrt{2}+14} + 50\sqrt{2} + (2i-14)\sqrt{10\sqrt{2}+14}\right. \\ & \left.+ (28i+14)e^{(2x)} + 70\right) \\ & - \left(\frac{1}{16}i + \frac{1}{16}\right) \sqrt{2\sqrt{2}+2} \left(-\frac{i}{\sqrt{2}+1} + 1\right) \log\left(2\sqrt{2}e^{(2x)}\right. \\ & \left.+ 2\sqrt{2}\sqrt{2\sqrt{2}-2} + (4i+2)\sqrt{2} + (2i-2)\sqrt{2\sqrt{2}-2-2e^{(2x)}-4i-2}\right) \\ & + \left(\frac{1}{16}i + \frac{1}{16}\right) \sqrt{2\sqrt{2}+2} \left(-\frac{i}{\sqrt{2}+1} + 1\right) \log\left(2\sqrt{2}e^{(2x)}\right. \\ & \left.- 2\sqrt{2}\sqrt{2\sqrt{2}-2} + (4i+2)\sqrt{2} - (2i-2)\sqrt{2\sqrt{2}-2-2e^{(2x)}-4i-2}\right) \end{aligned}$$

[In] integrate(1/(1+cosh(x)^4),x, algorithm="giac")

[Out] -(1/16*I + 1/16)*sqrt(2*sqrt(2) - 2)*(-I/(sqrt(2) - 1) + 1)*log((20*I + 10)*sqrt(2)*e^(2*x) + 10*sqrt(2)*sqrt(10*sqrt(2) + 14) + 50*sqrt(2) - (2*I - 14)*sqrt(10*sqrt(2) + 14) + (28*I + 14)*e^(2*x) + 70) + (1/16*I + 1/16)*sqrt(2*sqrt(2) - 2)*(-I/(sqrt(2) - 1) + 1)*log((20*I + 10)*sqrt(2)*e^(2*x) - 10*sqrt(2)*sqrt(10*sqrt(2) + 14) + 50*sqrt(2) + (2*I - 14)*sqrt(10*sqrt(2) + 14) + (28*I + 14)*e^(2*x) + 70) - (1/16*I + 1/16)*sqrt(2*sqrt(2) + 2)*(-I/(

```

sqrt(2) + 1) + 1)*log(2*sqrt(2)*e^(2*x) + 2*sqrt(2)*sqrt(2*sqrt(2) - 2) + (
4*I + 2)*sqrt(2) + (2*I - 2)*sqrt(2*sqrt(2) - 2) - 2*e^(2*x) - 4*I - 2) + (
1/16*I + 1/16)*sqrt(2*sqrt(2) + 2)*(-I/(sqrt(2) + 1) + 1)*log(2*sqrt(2)*e^(
2*x) - 2*sqrt(2)*sqrt(2*sqrt(2) - 2) + (4*I + 2)*sqrt(2) - (2*I - 2)*sqrt(2
*sqrt(2) - 2) - 2*e^(2*x) - 4*I - 2)

```

Mupad [B] (verification not implemented)

Time = 1.11 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.16

$$\int \frac{1}{1 + \cosh^4(x)} dx$$

$$= \frac{\sqrt{2} \sqrt{1-i} \ln(e^{2x} (436273152 + 91291648i) + \sqrt{2} \sqrt{1-i} (-9830400 + 56623104i) + \sqrt{2} \sqrt{1-i} e^{2x} (218890240 + 149422080i) + (21168128 + 94306304i))}{8} - \frac{\sqrt{2} \sqrt{1-i} \ln(e^{2x} (436273152 + 91291648i) + \sqrt{2} \sqrt{1-i} (9830400 - 56623104i) + \sqrt{2} \sqrt{1-i} e^{2x} (-218890240 - 149422080i) + (21168128 - 94306304i))}{8} + \frac{\sqrt{2} \sqrt{1+i} \ln(e^{2x} (436273152 - 91291648i) + \sqrt{2} \sqrt{1+i} (-9830400 - 56623104i) + \sqrt{2} \sqrt{1+i} e^{2x} (218890240 - 149422080i) + (21168128 - 94306304i))}{8} - \frac{\sqrt{2} \sqrt{1+i} \ln(e^{2x} (436273152 - 91291648i) + \sqrt{2} \sqrt{1+i} (9830400 + 56623104i) + \sqrt{2} \sqrt{1+i} e^{2x} (-218890240 - 149422080i) + (21168128 + 94306304i))}{8}$$

[In] int(1/(cosh(x)^4 + 1),x)

```

[Out] (2^(1/2)*(1 - 1i)^(1/2)*log(exp(2*x)*(436273152 + 91291648i) - 2^(1/2)*(1 -
1i)^(1/2)*(9830400 - 56623104i) + 2^(1/2)*(1 - 1i)^(1/2)*exp(2*x)*(2188902
40 + 149422080i) + (21168128 + 94306304i)))/8 - (2^(1/2)*(1 - 1i)^(1/2)*log
(exp(2*x)*(436273152 + 91291648i) + 2^(1/2)*(1 - 1i)^(1/2)*(9830400 - 56623
104i) - 2^(1/2)*(1 - 1i)^(1/2)*exp(2*x)*(218890240 + 149422080i) + (2116812
8 + 94306304i)))/8 + (2^(1/2)*(1 + 1i)^(1/2)*log(exp(2*x)*(436273152 - 9129
1648i) - 2^(1/2)*(1 + 1i)^(1/2)*(9830400 + 56623104i) + 2^(1/2)*(1 + 1i)^(1
/2)*exp(2*x)*(218890240 - 149422080i) + (21168128 - 94306304i)))/8 - (2^(1/
2)*(1 + 1i)^(1/2)*log(exp(2*x)*(436273152 - 91291648i) + 2^(1/2)*(1 + 1i)^(
1/2)*(9830400 + 56623104i) - 2^(1/2)*(1 + 1i)^(1/2)*exp(2*x)*(218890240 - 1
49422080i) + (21168128 - 94306304i)))/8

```


3.63 $\int \frac{1}{1-\cosh^4(x)} dx$

Optimal result	417
Rubi [A] (verified)	417
Mathematica [A] (verified)	418
Maple [B] (verified)	418
Fricas [B] (verification not implemented)	419
Sympy [B] (verification not implemented)	419
Maxima [B] (verification not implemented)	420
Giac [B] (verification not implemented)	420
Mupad [B] (verification not implemented)	420

Optimal result

Integrand size = 10, antiderivative size = 25

$$\int \frac{1}{1-\cosh^4(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{\operatorname{coth}(x)}{2}$$

[Out] 1/2*coth(x)+1/4*arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)

Rubi [A] (verified)

Time = 0.01 (sec), antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3288, 396, 212}

$$\int \frac{1}{1-\cosh^4(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{2\sqrt{2}} + \frac{\operatorname{coth}(x)}{2}$$

[In] Int[(1 - Cosh[x]^4)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[2]]/(2*Sqrt[2]) + Coth[x]/2

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 396

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_)), x_Symbol] := Simp[d*x*((a + b*x^n)^(p + 1)/(b*(n*(p + 1) + 1))), x] - Dist[(a*d - b*c*(n*(

$(p + 1) + 1) / (b * (n * (p + 1) + 1))$, Int[(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && NeQ[n*(p + 1) + 1, 0]

Rule 3288

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)^4]^(p_)), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[(a + 2*a*ff^2*x^2 + (a + b)*ff^4*x^4)^p/(1 + ff^2*x^2)^(2*p + 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[p]

Rubi steps

$$\begin{aligned} \text{integral} &= \text{Subst} \left(\int \frac{1 - x^2}{1 - 2x^2} dx, x, \coth(x) \right) \\ &= \frac{\coth(x)}{2} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \coth(x) \right) \\ &= \frac{\operatorname{arctanh} \left(\frac{\tanh(x)}{\sqrt{2}} \right)}{2\sqrt{2}} + \frac{\coth(x)}{2} \end{aligned}$$

Mathematica [A] (verified)

Time = 1.07 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.96

$$\int \frac{1}{1 - \cosh^4(x)} dx = \frac{1}{4} \left(\sqrt{2} \operatorname{arctanh} \left(\frac{\tanh(x)}{\sqrt{2}} \right) + 2 \coth(x) \right)$$

[In] Integrate[(1 - Cosh[x]^4)^(-1), x]

[Out] (Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]] + 2*Coth[x])/4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(18) = 36.

Time = 0.19 (sec) , antiderivative size = 44, normalized size of antiderivative = 1.76

method	result
risch	$\frac{1}{e^{2x}-1} + \frac{\sqrt{2} \ln(e^{2x}+3-2\sqrt{2})}{8} - \frac{\sqrt{2} \ln(e^{2x}+3+2\sqrt{2})}{8}$
default	$\frac{\tanh(\frac{x}{2})}{4} + \frac{\sqrt{2} \left(\ln \left(\frac{\tanh(\frac{x}{2})^2 + \tanh(\frac{x}{2})\sqrt{2}+1}{\tanh(\frac{x}{2})^2 - \tanh(\frac{x}{2})\sqrt{2}+1} \right) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}+1) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}-1) \right)}{16} - \frac{\sqrt{2} \left(\ln \left(\frac{\tanh(\frac{x}{2})^2 - \tanh(\frac{x}{2})\sqrt{2}+1}{\tanh(\frac{x}{2})^2 + \tanh(\frac{x}{2})\sqrt{2}+1} \right) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}+1) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}-1) \right)}{16}$

[In] `int(1/(1-cosh(x)^4),x,method=_RETURNVERBOSE)`

[Out] $1/(\exp(2*x)-1)+1/8*2^{(1/2)}*\ln(\exp(2*x)+3-2*2^{(1/2)})-1/8*2^{(1/2)}*\ln(\exp(2*x)+3+2*2^{(1/2)})$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(18) = 36$.

Time = 0.25 (sec) , antiderivative size = 115, normalized size of antiderivative = 4.60

$$\int \frac{1}{1 - \cosh^4(x)} dx$$

$$= \frac{(\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 - \sqrt{2}) \log\left(-\frac{3(2\sqrt{2}-3) \cosh(x)^2 - 4(3\sqrt{2}-4) \cosh(x) \sinh(x)}{\cosh(x)^2 + \sinh(x)^2}\right)}{8(\cosh(x)^2 + 2\cosh(x) \sinh(x) + \sinh(x)^2 - 1)}$$

[In] `integrate(1/(1-cosh(x)^4),x, algorithm="fricas")`

[Out] $1/8*((\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(-(3*(2*\sqrt{2} - 3)*\cosh(x)^2 - 4*(3*\sqrt{2} - 4)*\cosh(x)*\sinh(x) + 3*(2*\sqrt{2} - 3)*\sinh(x)^2 + 2*\sqrt{2} - 3)/(\cosh(x)^2 + \sinh(x)^2 + 3)) + 8)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(22) = 44$.

Time = 0.63 (sec) , antiderivative size = 75, normalized size of antiderivative = 3.00

$$\int \frac{1}{1 - \cosh^4(x)} dx = -\frac{\sqrt{2} \log\left(4 \tanh^2\left(\frac{x}{2}\right) - 4\sqrt{2} \tanh\left(\frac{x}{2}\right) + 4\right)}{8}$$

$$+ \frac{\sqrt{2} \log\left(4 \tanh^2\left(\frac{x}{2}\right) + 4\sqrt{2} \tanh\left(\frac{x}{2}\right) + 4\right)}{8} + \frac{\tanh\left(\frac{x}{2}\right)}{4} + \frac{1}{4 \tanh\left(\frac{x}{2}\right)}$$

[In] `integrate(1/(1-cosh(x)**4),x)`

[Out] $-\sqrt{2}*\log(4*\tanh(x/2)**2 - 4*\sqrt{2}*\tanh(x/2) + 4)/8 + \sqrt{2}*\log(4*\tanh(x/2)**2 + 4*\sqrt{2}*\tanh(x/2) + 4)/8 + \tanh(x/2)/4 + 1/(4*\tanh(x/2))$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 45 vs. 2(18) = 36.

Time = 0.30 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.80

$$\int \frac{1}{1 - \cosh^4(x)} dx = -\frac{1}{8} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(-2x)} - 3}{2\sqrt{2} + e^{(-2x)} + 3} \right) - \frac{1}{e^{(-2x)} - 1}$$

[In] integrate(1/(1-cosh(x)^4),x, algorithm="maxima")

[Out] -1/8*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - 1/(e^(-2*x) - 1)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 43 vs. 2(18) = 36.

Time = 0.26 (sec) , antiderivative size = 43, normalized size of antiderivative = 1.72

$$\int \frac{1}{1 - \cosh^4(x)} dx = \frac{1}{8} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) + \frac{1}{e^{(2x)} - 1}$$

[In] integrate(1/(1-cosh(x)^4),x, algorithm="giac")

[Out] 1/8*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3)) + 1/(e^(2*x) - 1)

Mupad [B] (verification not implemented)

Time = 0.13 (sec) , antiderivative size = 61, normalized size of antiderivative = 2.44

$$\int \frac{1}{1 - \cosh^4(x)} dx = \frac{\sqrt{2} \ln \left(-2e^{2x} - \frac{\sqrt{2}(12e^{2x}+4)}{8} \right)}{8} - \frac{\sqrt{2} \ln \left(\frac{\sqrt{2}(12e^{2x}+4)}{8} - 2e^{2x} \right)}{8} + \frac{1}{e^{2x} - 1}$$

[In] int(-1/(cosh(x)^4 - 1),x)

[Out] (2^(1/2)*log(- 2*exp(2*x) - (2^(1/2)*(12*exp(2*x) + 4))/8))/8 - (2^(1/2)*log((2^(1/2)*(12*exp(2*x) + 4))/8 - 2*exp(2*x)))/8 + 1/(exp(2*x) - 1)

3.64 $\int \frac{1}{a+b \cosh^5(x)} dx$

Optimal result	421
Rubi [A] (verified)	422
Mathematica [C] (verified)	425
Maple [C] (verified)	425
Fricas [F(-2)]	426
Sympy [F]	426
Maxima [F]	426
Giac [F]	426
Mupad [F(-1)]	427

Optimal result

Integrand size = 10, antiderivative size = 494

$$\int \frac{1}{a+b \cosh^5(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a}-\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+\sqrt[5]{b}}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+\sqrt[5]{b}}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a}+\sqrt[5]{-1}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}-\sqrt[5]{-1}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-\sqrt[5]{-1}} \sqrt[5]{b} \sqrt{\sqrt[5]{a}+\sqrt[5]{-1}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a}-(-1)^{2/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+(-1)^{2/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{2/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a}+(-1)^{2/5}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a}+(-1)^{3/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}-(-1)^{3/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{3/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a}+(-1)^{3/5}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a}-(-1)^{4/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+(-1)^{4/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{4/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a}+(-1)^{4/5}} \sqrt[5]{b}}$$

[Out] 2/5*arctanh((a^(1/5)-b^(1/5))^(1/2)*tanh(1/2*x)/(a^(1/5)+b^(1/5))^(1/2))/a^(4/5)/(a^(1/5)-b^(1/5))^(1/2)/(a^(1/5)+b^(1/5))^(1/2)+2/5*arctanh((a^(1/5)+

$$\begin{aligned} & (-1)^{1/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} - (-1)^{1/5} * b^{1/5})^{1/2}) / a \\ & ^{4/5} / (a^{1/5} - (-1)^{1/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{1/5} * b^{1/5})^{1/2} \\ &) + 2/5 * \operatorname{arctanh}((a^{1/5} - (-1)^{2/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} + (-1)^{2/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{2/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{2/5} * b^{1/5})^{1/2} + 2/5 * \operatorname{arctanh}((a^{1/5} + (-1)^{3/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} - (-1)^{3/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{3/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{3/5} * b^{1/5})^{1/2} + 2/5 * \operatorname{arctanh}((a^{1/5} - (-1)^{4/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} + (-1)^{4/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{4/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{4/5} * b^{1/5})^{1/2} \end{aligned}$$

Rubi [A] (verified)

Time = 0.63 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3292, 2738, 214}

$$\begin{aligned} \int \frac{1}{a + b \cosh^5(x)} dx &= \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - \sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + \sqrt[5]{b}}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{b}} \sqrt{\sqrt[5]{a} + \sqrt[5]{b}}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b}} \end{aligned}$$

[In] Int[(a + b*Cosh[x]^5)^(-1), x]

[Out] (2*ArcTanh[(Sqrt[a^(1/5) - b^(1/5)]*Tanh[x/2])/Sqrt[a^(1/5) + b^(1/5)])]/(5*a^(4/5)*Sqrt[a^(1/5) - b^(1/5)]*Sqrt[a^(1/5) + b^(1/5)]) + (2*ArcTanh[(Sqr

```
t[a^(1/5) + (-1)^(1/5)*b^(1/5)]*Tanh[x/2])/Sqrt[a^(1/5) - (-1)^(1/5)*b^(1/5)
)]]/(5*a^(4/5)*Sqrt[a^(1/5) - (-1)^(1/5)*b^(1/5)]*Sqrt[a^(1/5) + (-1)^(1/5)
)*b^(1/5)]) + (2*ArcTanh[(Sqrt[a^(1/5) - (-1)^(2/5)*b^(1/5)]*Tanh[x/2])/Sqr
t[a^(1/5) + (-1)^(2/5)*b^(1/5)])]/(5*a^(4/5)*Sqrt[a^(1/5) - (-1)^(2/5)*b^(1
/5)]*Sqrt[a^(1/5) + (-1)^(2/5)*b^(1/5)]) + (2*ArcTanh[(Sqrt[a^(1/5) + (-1)^(
3/5)*b^(1/5)]*Tanh[x/2])/Sqrt[a^(1/5) - (-1)^(3/5)*b^(1/5)])]/(5*a^(4/5)*S
qrt[a^(1/5) - (-1)^(3/5)*b^(1/5)]*Sqrt[a^(1/5) + (-1)^(3/5)*b^(1/5)]) + (2*
ArcTanh[(Sqrt[a^(1/5) - (-1)^(4/5)*b^(1/5)]*Tanh[x/2])/Sqrt[a^(1/5) + (-1)^(
4/5)*b^(1/5)])]/(5*a^(4/5)*Sqrt[a^(1/5) - (-1)^(4/5)*b^(1/5)]*Sqrt[a^(1/5)
+ (-1)^(4/5)*b^(1/5)])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 2738

```
Int[((a_) + (b_.)*sin[Pi/2 + (c_.) + (d_.)*(x_)])^(-1), x_Symbol] := With[{
e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (
a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x]
&& NeQ[a^2 - b^2, 0]
```

Rule 3292

```
Int[((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :=
Int[ExpandTrig[(a + b*(c*sin[e + f*x])^n)^p, x], x] /; FreeQ[{a, b, c, e, f
, n}, x] && (IGtQ[p, 0] || (EqQ[p, -1] && IntegerQ[n]))
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(-\frac{1}{5a^{4/5} \left(-\sqrt[5]{a} - \sqrt[5]{b} \cosh(x) \right)} - \frac{1}{5a^{4/5} \left(-\sqrt[5]{a} + \sqrt[5]{-1} \sqrt[5]{b} \cosh(x) \right)} \right. \\ &\quad - \frac{1}{5a^{4/5} \left(-\sqrt[5]{a} - (-1)^{2/5} \sqrt[5]{b} \cosh(x) \right)} - \frac{1}{5a^{4/5} \left(-\sqrt[5]{a} + (-1)^{3/5} \sqrt[5]{b} \cosh(x) \right)} \\ &\quad \left. - \frac{1}{5a^{4/5} \left(-\sqrt[5]{a} - (-1)^{4/5} \sqrt[5]{b} \cosh(x) \right)} \right) dx \\ &= -\frac{\int \frac{1}{-\sqrt[5]{a} - \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} - \frac{\int \frac{1}{-\sqrt[5]{a} + \sqrt[5]{-1} \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} - \frac{\int \frac{1}{-\sqrt[5]{a} - (-1)^{2/5} \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} \\ &\quad - \frac{\int \frac{1}{-\sqrt[5]{a} + (-1)^{3/5} \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} - \frac{\int \frac{1}{-\sqrt[5]{a} - (-1)^{4/5} \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} \end{aligned}$$

$$\begin{aligned}
& \frac{2 \operatorname{Subst} \left(\int \frac{1}{-\sqrt[5]{a}-\sqrt[5]{b}-\left(-\sqrt[5]{a}+\sqrt[5]{b}\right)x^2} dx, x, \tanh\left(\frac{x}{2}\right) \right)}{5a^{4/5}} \\
& - \frac{2 \operatorname{Subst} \left(\int \frac{1}{-\sqrt[5]{a}+\sqrt[5]{-1}\sqrt[5]{b}-\left(-\sqrt[5]{a}-\sqrt[5]{-1}\sqrt[5]{b}\right)x^2} dx, x, \tanh\left(\frac{x}{2}\right) \right)}{5a^{4/5}} \\
& - \frac{2 \operatorname{Subst} \left(\int \frac{1}{-\sqrt[5]{a}-(-1)^{2/5}\sqrt[5]{b}-\left(-\sqrt[5]{a}+(-1)^{2/5}\sqrt[5]{b}\right)x^2} dx, x, \tanh\left(\frac{x}{2}\right) \right)}{5a^{4/5}} \\
& - \frac{2 \operatorname{Subst} \left(\int \frac{1}{-\sqrt[5]{a}+(-1)^{3/5}\sqrt[5]{b}-\left(-\sqrt[5]{a}-(-1)^{3/5}\sqrt[5]{b}\right)x^2} dx, x, \tanh\left(\frac{x}{2}\right) \right)}{5a^{4/5}} \\
& - \frac{2 \operatorname{Subst} \left(\int \frac{1}{-\sqrt[5]{a}-(-1)^{4/5}\sqrt[5]{b}-\left(-\sqrt[5]{a}+(-1)^{4/5}\sqrt[5]{b}\right)x^2} dx, x, \tanh\left(\frac{x}{2}\right) \right)}{5a^{4/5}} \\
& = \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[5]{a}-\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+\sqrt[5]{b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+\sqrt[5]{b}}} + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[5]{a}+\sqrt[5]{-1}\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}-\sqrt[5]{-1}\sqrt[5]{b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-\sqrt[5]{-1}\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+\sqrt[5]{-1}\sqrt[5]{b}}} \\
& + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[5]{a}-(-1)^{2/5}\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+(-1)^{2/5}\sqrt[5]{b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{2/5}\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+(-1)^{2/5}\sqrt[5]{b}}} \\
& + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[5]{a}+(-1)^{3/5}\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}-(-1)^{3/5}\sqrt[5]{b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{3/5}\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+(-1)^{3/5}\sqrt[5]{b}}} \\
& + \frac{2 \operatorname{arctanh} \left(\frac{\sqrt{\sqrt[5]{a}-(-1)^{4/5}\sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a}+(-1)^{4/5}\sqrt[5]{b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a}-(-1)^{4/5}\sqrt[5]{b}} \sqrt{\sqrt[5]{a}+(-1)^{4/5}\sqrt[5]{b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.05 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.28

$$\int \frac{1}{a + b \cosh^5(x)} dx$$

$$= \frac{8}{5} \text{RootSum} \left[b + 5b\#1^2 + 10b\#1^4 + 32a\#1^5 + 10b\#1^6 + 5b\#1^8 \right. \\ \left. + b\#1^{10} \&, \frac{x\#1^3 + 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) \#1^3}{b + 4b\#1^2 + 16a\#1^3 + 6b\#1^4 + 4b\#1^6 + b\#1^8} \& \right]$$

[In] Integrate[(a + b*Cosh[x]^5)^(-1),x]

[Out] (8*RootSum[b + 5*b*#1^2 + 10*b*#1^4 + 32*a*#1^5 + 10*b*#1^6 + 5*b*#1^8 + b*#1^10 & , (x*#1^3 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^3)/(b + 4*b*#1^2 + 16*a*#1^3 + 6*b*#1^4 + 4*b*#1^6 + b*#1^8) &])/5

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.91 (sec) , antiderivative size = 156, normalized size of antiderivative = 0.32

method	result
default	$\frac{\sum_{R=\text{RootOf}((a-b)Z^{10}+(-5a-5b)Z^8+(10a-10b)Z^6+(-10a-10b)Z^4+(5a-5b)Z^2-a-b)} \left(-R^8+4R^6-6R^4+4R^2-1 \right)}{5} \frac{\left(-R^8+4R^6-6R^4+4R^2-1 \right)}{R^9 a - R^9 b - 4R^7 a - 4R^7 b + 6R^5 a - 6R^5 b - 4R^3 a - 4R^3 b + R a - R b} \ln(\tanh(1/2*x) - R)$
risch	$\sum_{R=\text{RootOf}(-1+(9765625a^{10}-9765625a^8b^2)Z^{10}-1953125a^8Z^8+156250a^6Z^6-6250a^4Z^4+125a^2Z^2)} -R \ln \left(e^x + \left(\dots \right) \right)$

[In] int(1/(a+b*cosh(x)^5),x,method=_RETURNVERBOSE)

[Out] 1/5*sum((-R^8+4R^6-6R^4+4R^2-1)/(R^9*a-R^9*b-4R^7*a-4R^7*b+6R^5*a-6R^5*b-4R^3*a-4R^3*b+R*a-R*b)*ln(tanh(1/2*x)-R),R=RootOf((a-b)*Z^10+(-5*a-5*b)*Z^8+(10*a-10*b)*Z^6+(-10*a-10*b)*Z^4+(5*a-5*b)*Z^2-a-b))

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{a + b \cosh^5(x)} dx = \text{Exception raised: RuntimeError}$$

[In] `integrate(1/(a+b*cosh(x)^5),x, algorithm="fricas")`

[Out] Exception raised: RuntimeError >> no explicit roots found

Sympy [F]

$$\int \frac{1}{a + b \cosh^5(x)} dx = \int \frac{1}{a + b \cosh^5(x)} dx$$

[In] `integrate(1/(a+b*cosh(x)**5),x)`

[Out] `Integral(1/(a + b*cosh(x)**5), x)`

Maxima [F]

$$\int \frac{1}{a + b \cosh^5(x)} dx = \int \frac{1}{b \cosh(x)^5 + a} dx$$

[In] `integrate(1/(a+b*cosh(x)^5),x, algorithm="maxima")`

[Out] `integrate(1/(b*cosh(x)^5 + a), x)`

Giac [F]

$$\int \frac{1}{a + b \cosh^5(x)} dx = \int \frac{1}{b \cosh(x)^5 + a} dx$$

[In] `integrate(1/(a+b*cosh(x)^5),x, algorithm="giac")`

[Out] `integrate(1/(b*cosh(x)^5 + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{a + b \cosh^5(x)} dx = \text{Hanged}$$

```
[In] int(1/(a + b*cosh(x)^5),x)
```

```
[Out] \text{Hanged}
```

3.65 $\int \frac{1}{a+b \cosh^6(x)} dx$

Optimal result	428
Rubi [A] (verified)	428
Mathematica [C] (verified)	430
Maple [C] (verified)	431
Fricas [C] (verification not implemented)	431
Sympy [F]	431
Maxima [F]	432
Giac [F]	432
Mupad [B] (verification not implemented)	432

Optimal result

Integrand size = 10, antiderivative size = 171

$$\int \frac{1}{a+b \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}$$

[Out] $\frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3} + b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3} + b^{1/3})^{1/2} + \frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3} - (-1)^{1/3} b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3} - (-1)^{1/3} b^{1/3})^{1/2} + \frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3} + (-1)^{2/3} b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3} + (-1)^{2/3} b^{1/3})^{1/2}$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used

= {3290, 3260, 212}

$$\int \frac{1}{a + b \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}$$

[In] Int[(a + b*Cosh[x]^6)^(-1), x]

[Out] ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) + b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) + b^(1/3)]) + ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) - (-1)^(1/3)*b^(1/3)]) + ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) + (-1)^(2/3)*b^(1/3)])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2)], x], x, Tan[e + f*x]/ff, x]] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))^(n_)^(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rubi steps

$$\text{integral} = \frac{\int \frac{1}{1 + \frac{\sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a} + \frac{\int \frac{1}{1 - \frac{\sqrt[3]{-1} \sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a} + \frac{\int \frac{1}{1 + \frac{(-1)^{2/3} \sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a}$$

$$\begin{aligned}
& \text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{\sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right) \\
= & \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[3]{-1} \sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right)}{3a} \\
& + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{(-1)^{2/3} \sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right)}{3a} \\
= & \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + \sqrt[3]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - \sqrt[3]{-1} \sqrt[3]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + (-1)^{2/3} \sqrt[3]{b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.77

$$\begin{aligned}
& \int \frac{1}{a + b \cosh^6(x)} dx \\
= & \frac{16}{3} \text{RootSum} \left[b + 6b\#1 + 15b\#1^2 + 64a\#1^3 + 20b\#1^3 + 15b\#1^4 + 6b\#1^5 \right. \\
& \left. + b\#1^6 \&, \frac{x\#1^2 + \log(-\cosh(x) - \sinh(x) + \cosh(x)\#1 - \sinh(x)\#1)\#1^2}{b + 5b\#1 + 32a\#1^2 + 10b\#1^2 + 10b\#1^3 + 5b\#1^4 + b\#1^5} \& \right]
\end{aligned}$$

[In] Integrate[(a + b*Cosh[x]^6)^(-1), x]

[Out] (16*RootSum[b + 6*b*#1 + 15*b*#1^2 + 64*a*#1^3 + 20*b*#1^3 + 15*b*#1^4 + 6*b*#1^5 + b*#1^6 & , (x*#1^2 + Log[-Cosh[x] - Sinh[x] + Cosh[x]*#1 - Sinh[x]*#1]*#1^2)/(b + 5*b*#1 + 32*a*#1^2 + 10*b*#1^2 + 10*b*#1^3 + 5*b*#1^4 + b*#1^5) &])/3

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.94 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.82

method	result
risch	$\sum_{R=\text{RootOf}(-1+(46656a^6+46656a^5b)Z^6-3888a^4Z^4+108a^2Z^2)} _R \ln \left(e^{2x} + \left(-\frac{15552a^6}{b} - 15552a^5 \right) _R^5 + \right.$
default	$\left. \left(\sum_{R=\text{RootOf}((a+b)Z^{12}+(-6a+6b)Z^{10}+(15a+15b)Z^8+(-20a+20b)Z^6+(15a+15b)Z^4+(-6a+6b)Z^2+a+b)} \frac{_R^{11} _a + _R^{11} _b - 5}{6} \right) \right.$

[In] `int(1/(a+b*cosh(x)^6),x,method=_RETURNVERBOSE)`

[Out] `sum(_R*ln(exp(2*x)+(-15552*a^6/b-15552*a^5)*_R^5+(2592*a^5/b+2592*a^4)*_R^4+(864*a^4/b-432*a^3)*_R^3+(-144/b*a^3+72*a^2)*_R^2+(-12*a^2/b-12*a)*_R+2*a/b+1),_R=RootOf(-1+(46656*a^6+46656*a^5*b)*_Z^6-3888*a^4*_Z^4+108*a^2*_Z^2))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.21 (sec) , antiderivative size = 15201, normalized size of antiderivative = 88.89

$$\int \frac{1}{a + b \cosh^6(x)} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*cosh(x)^6),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{1}{a + b \cosh^6(x)} dx = \int \frac{1}{a + b \cosh^6(x)} dx$$

[In] `integrate(1/(a+b*cosh(x)**6),x)`

[Out] `Integral(1/(a + b*cosh(x)**6), x)`

$$\begin{aligned}
& p(2x) + 1770440a^5b^2\exp(2x) + 1239040a^6b\exp(2x)) / (b^{10}(a+b)^3) \\
& + (17509995351216488448\sqrt[6]{46656a^5bd^6 + 46656a^6d^6 - 3888a^4d^4 + 108a^2d^2 - 1, d, k} \\
& \cdot (262144a^7\exp(2x) + 203520a^6b + 65536a^7 + 453a^3b^4 + 72022a^4b^3 + 209472a^5b^2 + 630a^3b^4\exp(2x) + 2 \\
& 54512a^4b^3\exp(2x) + 767508a^5b^2\exp(2x) + 775680a^6b\exp(2x))) / \\
& (b^{10}(a+b)^2) - (486388759756013568 \cdot (655360a^5\exp(2x) - 9ab^4 + 37 \\
& 0176a^4b + 196608a^5 - 24408a^2b^3 + 149088a^3b^2 - 63676a^2b^3\exp(2x) + 526248a^3b^2\exp(2x) \\
& - 10a^4b^4\exp(2x) + 1245184a^4b\exp(2x))) / (b^{10}(a+b)^2) - (40532396646334464 \cdot (655360a^5\exp(2x) - b^5\exp(2x) \\
& - 24677ab^4 + 773120a^4b + 262144a^5 - b^5 + 198071a^2b^3 + 733696a^3b^2 + 477713a^2b^3\exp(2x) \\
& + 1770640a^3b^2\exp(2x) - 53861ab^4\exp(2x) + 1894400a^4b\exp(2x))) / (b^{10}(a+b)^3) \\
& + (13510798882111488 \cdot (655360a^3\exp(2x) + 11382b^3\exp(2x) + 144416ab^2 + 269056a^2b + 131072a^3 \\
& + 6459b^3 + 677524ab^2\exp(2x) + 1321472a^2b\exp(2x))) / (b^{10}(a+b)^2) \\
& + (1125899906842624 \cdot (851968a^4\exp(2x) + 6006b^4\exp(2x) + 211497ab^3 + 597504a^3b + 196608a^4 \\
& + 3840b^4 + 608544a^2b^2 + 2562504a^2b^2\exp(2x) + 864565ab^3\exp(2x) + 2555904a^3b\exp(2x))) / \\
& (b^{10}(a+b)^2(ab+a^2)) \cdot \sqrt[6]{46656a^5bd^6 + 46656a^6d^6 - 3888a^4d^4 + 108a^2d^2 - 1, d, k}, k, 1, 6)
\end{aligned}$$

3.66 $\int \frac{1}{a+b \cosh^8(x)} dx$

Optimal result	434
Rubi [A] (verified)	434
Mathematica [C] (verified)	436
Maple [C] (verified)	437
Fricas [B] (verification not implemented)	437
Sympy [F]	437
Maxima [F]	438
Giac [F]	438
Mupad [F(-1)]	438

Optimal result

Integrand size = 10, antiderivative size = 245

$$\int \frac{1}{a+b \cosh^8(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt[4]{\sqrt{-a}-\sqrt[4]{b}}}\right)}{4(-a)^{7/8} \sqrt[4]{\sqrt{-a}-\sqrt[4]{b}}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt[4]{\sqrt{-a}-i\sqrt[4]{b}}}\right)}{4(-a)^{7/8} \sqrt[4]{\sqrt{-a}-i\sqrt[4]{b}}}$$

$$- \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt[4]{\sqrt{-a}+i\sqrt[4]{b}}}\right)}{4(-a)^{7/8} \sqrt[4]{\sqrt{-a}+i\sqrt[4]{b}}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt[4]{\sqrt{-a}+\sqrt[4]{b}}}\right)}{4(-a)^{7/8} \sqrt[4]{\sqrt{-a}+\sqrt[4]{b}}}$$

[Out] $-1/4*\operatorname{arctanh}((-a)^{(1/8)*\tanh(x)/((-a)^{(1/4)-b^{(1/4)}})^{(1/2))}/(-a)^{(7/8)/((-a)^{(1/4)-b^{(1/4)}})^{(1/2)}-1/4*\operatorname{arctanh}((-a)^{(1/8)*\tanh(x)/((-a)^{(1/4)-I*b^{(1/4)}})^{(1/2))}/(-a)^{(7/8)/((-a)^{(1/4)-I*b^{(1/4)}})^{(1/2)}-1/4*\operatorname{arctanh}((-a)^{(1/8)*\tanh(x)/((-a)^{(1/4)+I*b^{(1/4)}})^{(1/2))}/(-a)^{(7/8)/((-a)^{(1/4)+I*b^{(1/4)}})^{(1/2)}-1/4*\operatorname{arctanh}((-a)^{(1/8)*\tanh(x)/((-a)^{(1/4)+b^{(1/4)}})^{(1/2))}/(-a)^{(7/8)/((-a)^{(1/4)+b^{(1/4)}})^{(1/2)}$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 245, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used

= {3290, 3260, 212}

$$\int \frac{1}{a + b \cosh^8(x)} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a-i}\sqrt[4]{b}}}\right)}{4(-a)^{7/8}\sqrt{\sqrt[4]{-a-i}\sqrt[4]{b}}} - \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}}\right)}{4(-a)^{7/8}\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}} \\ - \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}}\right)}{4(-a)^{7/8}\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}} - \frac{\operatorname{arctanh}\left(\frac{(-a)^{5/8} \tanh(x)}{\sqrt{a\sqrt[4]{b}+(-a)^{5/4}}}\right)}{4(-a)^{3/8}\sqrt{a\sqrt[4]{b}+(-a)^{5/4}}}$$

[In] Int[(a + b*Cosh[x]^8)^(-1), x]

[Out] -1/4*ArcTanh[((-a)^(1/8)*Tanh[x])/Sqrt[(-a)^(1/4) - I*b^(1/4)]]/((-a)^(7/8)*Sqrt[(-a)^(1/4) - I*b^(1/4)]) - ArcTanh[((-a)^(1/8)*Tanh[x])/Sqrt[(-a)^(1/4) + I*b^(1/4)]]/(4*(-a)^(7/8)*Sqrt[(-a)^(1/4) + I*b^(1/4)]) - ArcTanh[((-a)^(1/8)*Tanh[x])/Sqrt[(-a)^(1/4) + b^(1/4)]]/(4*(-a)^(7/8)*Sqrt[(-a)^(1/4) + b^(1/4)]) - ArcTanh[((-a)^(5/8)*Tanh[x])/Sqrt[(-a)^(5/4) + a*b^(1/4)]]/(4*(-a)^(3/8)*Sqrt[(-a)^(5/4) + a*b^(1/4)])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))^(n_)^(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rubi steps

$$\text{integral} = \frac{\int \frac{1}{1 - \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{-a}}} dx}{4a} + \frac{\int \frac{1}{1 - i \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{-a}}} dx}{4a} + \frac{\int \frac{1}{1 + i \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{-a}}} dx}{4a} + \frac{\int \frac{1}{1 + \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{-a}}} dx}{4a}$$

$$\begin{aligned}
& \text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right) + \text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{i\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right) \\
&= \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right)}{4a} + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{i\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right)}{4a} \\
&+ \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{i\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right)}{4a} + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{\sqrt[4]{b}}{\sqrt[4]{-a}}\right) x^2} dx, x, \coth(x) \right)}{4a} \\
&= \frac{\text{arctanh} \left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a-i}\sqrt[4]{b}}} \right)}{4(-a)^{7/8} \sqrt{\sqrt[4]{-a} - i\sqrt[4]{b}}} - \frac{\text{arctanh} \left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}} \right)}{4(-a)^{7/8} \sqrt{\sqrt[4]{-a} + i\sqrt[4]{b}}} \\
&- \frac{\text{arctanh} \left(\frac{\sqrt[8]{-a} \tanh(x)}{\sqrt{\sqrt[4]{-a+i}\sqrt[4]{b}}} \right)}{4(-a)^{7/8} \sqrt{\sqrt[4]{-a} + \sqrt[4]{b}}} - \frac{\text{arctanh} \left(\frac{(-a)^{5/8} \tanh(x)}{\sqrt{(-a)^{5/4+a}\sqrt[4]{b}}} \right)}{4(-a)^{3/8} \sqrt{(-a)^{5/4} + a\sqrt[4]{b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.07 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.64

$$\begin{aligned}
& \int \frac{1}{a + b \cosh^8(x)} dx \\
&= 16\text{RootSum} \left[b + 8b\#1 + 28b\#1^2 + 56b\#1^3 + 256a\#1^4 + 70b\#1^4 + 56b\#1^5 + 28b\#1^6 + 8b\#1^7 \right. \\
&\quad \left. + b\#1^8 \& , \frac{x\#1^3 + \log(-\cosh(x) - \sinh(x) + \cosh(x)\#1 - \sinh(x)\#1)\#1^3}{b + 7b\#1 + 21b\#1^2 + 128a\#1^3 + 35b\#1^3 + 35b\#1^4 + 21b\#1^5 + 7b\#1^6 + b\#1^7} \& \right]
\end{aligned}$$

[In] Integrate[(a + b*Cosh[x]^8)^(-1),x]

[Out] 16*RootSum[b + 8*b*#1 + 28*b*#1^2 + 56*b*#1^3 + 256*a*#1^4 + 70*b*#1^4 + 56*b*#1^5 + 28*b*#1^6 + 8*b*#1^7 + b*#1^8 & , (x*#1^3 + Log[-Cosh[x] - Sinh[x] + Cosh[x]*#1 - Sinh[x]*#1]*#1^3)/(b + 7*b*#1 + 21*b*#1^2 + 128*a*#1^3 + 35*b*#1^3 + 35*b*#1^4 + 21*b*#1^5 + 7*b*#1^6 + b*#1^7) &]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.38 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.75

method	result
risch	$\sum_{R=\text{RootOf}(1+(16777216a^8+16777216a^7b)Z^8-1048576a^6Z^6+24576a^4Z^4-256a^2Z^2)} _R \ln \left(e^{2x} + \left(-\frac{4194304a^8}{b} \right) \right)$
default	$\left(\sum_{R=\text{RootOf}((a+b)Z^{16}+(-8a+8b)Z^{14}+(28a+28b)Z^{12}+(-56a+56b)Z^{10}+(70a+70b)Z^8+(-56a+56b)Z^6+(28a+28b)Z^4+(-8a+8b)Z^2)} \right)$

[In] `int(1/(a+b*cosh(x)^8),x,method=_RETURNVERBOSE)`

[Out] `sum(_R*ln(exp(2*x)+(-4194304*a^8/b-4194304*a^7)*_R^7+(524288*a^7/b+524288*a^6)*_R^6+(196608*a^6/b-65536*a^5)*_R^5+(-24576*a^5/b+8192*a^4)*_R^4+(-3072*a^4/b-1024*a^3)*_R^3+(384/b*a^3+128*a^2)*_R^2+(16*a^2/b-16*a)*_R-2*a/b+1),_R=RootOf(1+(16777216*a^8+16777216*a^7*b)*_Z^8-1048576*a^6*_Z^6+24576*a^4*_Z^4-256*a^2*_Z^2))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 661324 vs. $2(165) = 330$.

Time = 3.14 (sec) , antiderivative size = 661324, normalized size of antiderivative = 2699.28

$$\int \frac{1}{a + b \cosh^8(x)} dx = \text{Too large to display}$$

[In] `integrate(1/(a+b*cosh(x)^8),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{1}{a + b \cosh^8(x)} dx = \int \frac{1}{a + b \cosh^8(x)} dx$$

[In] `integrate(1/(a+b*cosh(x)**8),x)`

[Out] `Integral(1/(a + b*cosh(x)**8), x)`

Maxima [F]

$$\int \frac{1}{a + b \cosh^8(x)} dx = \int \frac{1}{b \cosh(x)^8 + a} dx$$

[In] integrate(1/(a+b*cosh(x)^8),x, algorithm="maxima")

[Out] integrate(1/(b*cosh(x)^8 + a), x)

Giac [F]

$$\int \frac{1}{a + b \cosh^8(x)} dx = \int \frac{1}{b \cosh(x)^8 + a} dx$$

[In] integrate(1/(a+b*cosh(x)^8),x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{a + b \cosh^8(x)} dx = \text{Hanged}$$

[In] int(1/(a + b*cosh(x)^8),x)

[Out] \text{Hanged}

3.67 $\int \frac{1}{a-b \cosh^5(x)} dx$

Optimal result	439
Rubi [A] (verified)	440
Mathematica [C] (verified)	443
Maple [C] (verified)	443
Fricas [F(-2)]	444
Sympy [F]	444
Maxima [F]	444
Giac [F]	444
Mupad [F(-1)]	445

Optimal result

Integrand size = 11, antiderivative size = 494

$$\int \frac{1}{a-b \cosh^5(x)} dx = \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + \sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - \sqrt[5]{b}}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{b}} \sqrt{\sqrt[5]{a} + \sqrt[5]{b}}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b}} + \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b}}\right)}{5a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b}}$$

[Out] 2/5*arctanh((a^(1/5)+b^(1/5))^(1/2)*tanh(1/2*x)/(a^(1/5)-b^(1/5))^(1/2))/a^(4/5)/(a^(1/5)-b^(1/5))^(1/2)/(a^(1/5)+b^(1/5))^(1/2)+2/5*arctanh((a^(1/5)-

$$\begin{aligned} & (-1)^{1/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} + (-1)^{1/5} * b^{1/5})^{1/2}) / a \\ & ^{4/5} / (a^{1/5} - (-1)^{1/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{1/5} * b^{1/5})^{1/2} \\ &) + 2/5 * \operatorname{arctanh}((a^{1/5} + (-1)^{2/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} - (-1)^{2/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{2/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{2/5} * b^{1/5})^{1/2} + 2/5 * \operatorname{arctanh}((a^{1/5} - (-1)^{3/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} + (-1)^{3/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{3/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{3/5} * b^{1/5})^{1/2} + 2/5 * \operatorname{arctanh}((a^{1/5} + (-1)^{4/5} * b^{1/5})^{1/2} * \tanh(1/2 * x) / (a^{1/5} - (-1)^{4/5} * b^{1/5})^{1/2}) / a^{4/5} / (a^{1/5} - (-1)^{4/5} * b^{1/5})^{1/2} / (a^{1/5} + (-1)^{4/5} * b^{1/5})^{1/2} \end{aligned}$$

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used = {3292, 2738, 214}

$$\begin{aligned} \int \frac{1}{a - b \cosh^5(x)} dx &= \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + \sqrt[5]{b}} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - \sqrt[5]{b}}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{b}} \sqrt{\sqrt[5]{a} + \sqrt[5]{b}}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - \sqrt[5]{-1}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + \sqrt[5]{-1}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{2/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{2/5}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{3/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{3/5}} \sqrt[5]{b}} \\ &+ \frac{2 \operatorname{arctanh}\left(\frac{\sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b} \tanh\left(\frac{x}{2}\right)}{\sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b}}\right)}{5 a^{4/5} \sqrt{\sqrt[5]{a} - (-1)^{4/5}} \sqrt[5]{b} \sqrt{\sqrt[5]{a} + (-1)^{4/5}} \sqrt[5]{b}} \end{aligned}$$

[In] Int[(a - b*Cosh[x]^5)^(-1), x]

[Out] (2*ArcTanh[(Sqrt[a^(1/5) + b^(1/5)]*Tanh[x/2])/Sqrt[a^(1/5) - b^(1/5)])]/(5*a^(4/5)*Sqrt[a^(1/5) - b^(1/5)]*Sqrt[a^(1/5) + b^(1/5)]) + (2*ArcTanh[(Sqr

$$\frac{\operatorname{tanh}\left(\frac{x}{2}\right)\sqrt[5]{a - (-1)^{1/5}b}}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{1/5}b} + \sqrt[5]{a + (-1)^{1/5}b})}} + \frac{2\operatorname{ArcTanh}\left(\frac{\sqrt[5]{a + (-1)^{2/5}b}\operatorname{tanh}\left(\frac{x}{2}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{2/5}b} + \sqrt[5]{a + (-1)^{2/5}b})}}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{2/5}b} + \sqrt[5]{a + (-1)^{2/5}b})}} + \frac{2\operatorname{ArcTanh}\left(\frac{\sqrt[5]{a - (-1)^{3/5}b}\operatorname{tanh}\left(\frac{x}{2}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{3/5}b} + \sqrt[5]{a + (-1)^{3/5}b})}}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{3/5}b} + \sqrt[5]{a + (-1)^{3/5}b})}} + \frac{2\operatorname{ArcTanh}\left(\frac{\sqrt[5]{a + (-1)^{4/5}b}\operatorname{tanh}\left(\frac{x}{2}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{4/5}b} + \sqrt[5]{a + (-1)^{4/5}b})}}\right)}{\sqrt[5]{a^{4/5}(\sqrt[5]{a - (-1)^{4/5}b} + \sqrt[5]{a + (-1)^{4/5}b})}}$$

Rule 214

$$\operatorname{Int}[(a + b)(x^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{Rt}[-a/b, 2]/a \operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] \text{ ; FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b]$$

Rule 2738

$$\operatorname{Int}[(a + b)\sin[\pi/2 + (c + d)(x)]^{-1}, x_Symbol] \rightarrow \operatorname{With}\{e = \operatorname{FreeFactors}[\operatorname{Tan}[(c + dx)/2], x], \operatorname{Dist}[2(e/d), \operatorname{Subst}[\operatorname{Int}[1/(a + b + (a - b)e^{2x^2}), x], x, \operatorname{Tan}[(c + dx)/2]/e], x]\} \text{ ; FreeQ}\{a, b, c, d, x\} \ \&\& \ \operatorname{NeQ}[a^2 - b^2, 0]$$

Rule 3292

$$\operatorname{Int}[(a + b)(c + f(x))^{n(p)}, x_Symbol] \rightarrow \operatorname{Int}[\operatorname{ExpandTrig}[a + b(c + f(x))^n]^p, x] \text{ ; FreeQ}\{a, b, c, e, f, n, x\} \ \&\& \ (\operatorname{IGtQ}[p, 0] \ \|\ (\operatorname{EqQ}[p, -1] \ \&\& \ \operatorname{IntegerQ}[n]))$$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{5a^{4/5}(\sqrt[5]{a} - \sqrt[5]{b} \cosh(x))} + \frac{1}{5a^{4/5}(\sqrt[5]{a} + \sqrt[5]{-1}\sqrt[5]{b} \cosh(x))} \right. \\ &\quad + \frac{1}{5a^{4/5}(\sqrt[5]{a} - (-1)^{2/5}\sqrt[5]{b} \cosh(x))} + \frac{1}{5a^{4/5}(\sqrt[5]{a} + (-1)^{3/5}\sqrt[5]{b} \cosh(x))} \\ &\quad \left. + \frac{1}{5a^{4/5}(\sqrt[5]{a} - (-1)^{4/5}\sqrt[5]{b} \cosh(x))} \right) dx \\ &= \frac{\int \frac{1}{\sqrt[5]{a} - \sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} + \frac{\int \frac{1}{\sqrt[5]{a} + \sqrt[5]{-1}\sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} + \frac{\int \frac{1}{\sqrt[5]{a} - (-1)^{2/5}\sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} \\ &\quad + \frac{\int \frac{1}{\sqrt[5]{a} + (-1)^{3/5}\sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} + \frac{\int \frac{1}{\sqrt[5]{a} - (-1)^{4/5}\sqrt[5]{b} \cosh(x)} dx}{5a^{4/5}} \end{aligned}$$

$$\begin{aligned}
& 2\text{Subst} \left(\int \frac{1}{\sqrt[5]{a-b} (\sqrt[5]{a+b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
= & \frac{\hspace{10em}}{5a^{4/5}} \\
& 2\text{Subst} \left(\int \frac{1}{\sqrt[5]{a+\sqrt{-1}b} (\sqrt[5]{a-\sqrt{-1}b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
+ & \frac{\hspace{10em}}{5a^{4/5}} \\
& 2\text{Subst} \left(\int \frac{1}{\sqrt[5]{a-(-1)^{2/5}b} (\sqrt[5]{a+(-1)^{2/5}b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
+ & \frac{\hspace{10em}}{5a^{4/5}} \\
& 2\text{Subst} \left(\int \frac{1}{\sqrt[5]{a+(-1)^{3/5}b} (\sqrt[5]{a-(-1)^{3/5}b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
+ & \frac{\hspace{10em}}{5a^{4/5}} \\
& 2\text{Subst} \left(\int \frac{1}{\sqrt[5]{a-(-1)^{4/5}b} (\sqrt[5]{a+(-1)^{4/5}b}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
+ & \frac{\hspace{10em}}{5a^{4/5}} \\
= & \frac{2\text{arctanh} \left(\frac{\sqrt{\sqrt[5]{a+b} \tanh(\frac{x}{2})}}{\sqrt{\sqrt[5]{a-b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a-b}} \sqrt{\sqrt[5]{a+b}}} + \frac{2\text{arctanh} \left(\frac{\sqrt{\sqrt[5]{a-\sqrt{-1}b} \tanh(\frac{x}{2})}}{\sqrt{\sqrt[5]{a+\sqrt{-1}b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a-\sqrt{-1}b}} \sqrt{\sqrt[5]{a+\sqrt{-1}b}}} \\
& + \frac{2\text{arctanh} \left(\frac{\sqrt{\sqrt[5]{a+(-1)^{2/5}b} \tanh(\frac{x}{2})}}{\sqrt{\sqrt[5]{a-(-1)^{2/5}b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a-(-1)^{2/5}b}} \sqrt{\sqrt[5]{a+(-1)^{2/5}b}}} \\
& + \frac{2\text{arctanh} \left(\frac{\sqrt{\sqrt[5]{a-(-1)^{3/5}b} \tanh(\frac{x}{2})}}{\sqrt{\sqrt[5]{a+(-1)^{3/5}b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a-(-1)^{3/5}b}} \sqrt{\sqrt[5]{a+(-1)^{3/5}b}}} \\
& + \frac{2\text{arctanh} \left(\frac{\sqrt{\sqrt[5]{a+(-1)^{4/5}b} \tanh(\frac{x}{2})}}{\sqrt{\sqrt[5]{a-(-1)^{4/5}b}}} \right)}{5a^{4/5} \sqrt{\sqrt[5]{a-(-1)^{4/5}b}} \sqrt{\sqrt[5]{a+(-1)^{4/5}b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.05 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.28

$$\int \frac{1}{a - b \cosh^5(x)} dx$$

$$= -\frac{8}{5} \text{RootSum} \left[b + 5b\#1^2 + 10b\#1^4 - 32a\#1^5 + 10b\#1^6 + 5b\#1^8 + b\#1^{10} \&, \frac{x\#1^3 + 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) \#1^3}{b + 4b\#1^2 - 16a\#1^3 + 6b\#1^4 + 4b\#1^6 + b\#1^8} \& \right]$$

[In] Integrate[(a - b*Cosh[x]^5)^(-1),x]

[Out] (-8*RootSum[b + 5*b*#1^2 + 10*b*#1^4 - 32*a*#1^5 + 10*b*#1^6 + 5*b*#1^8 + b*#1^10 & , (x*#1^3 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^3)/(b + 4*b*#1^2 - 16*a*#1^3 + 6*b*#1^4 + 4*b*#1^6 + b*#1^8) &])/5

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.89 (sec) , antiderivative size = 150, normalized size of antiderivative = 0.30

method	result
default	$\frac{\sum_{-R=\text{RootOf}((a+b)Z^{10}+(-5a+5b)Z^8+(10a+10b)Z^6+(-10a+10b)Z^4+(5a+5b)Z^2-a+b)} \left(-R^8+4R^6-6R^4+4R^2-1 \right)}{5} \frac{\left(-R^8+4R^6-6R^4+4R^2-1 \right)}{R^9+R^7+R^5+R^3+R} \ln \left(\frac{-R^8+4R^6-6R^4+4R^2-1}{R^9+R^7+R^5+R^3+R} \right)$
risch	$\sum_{-R=\text{RootOf}(-1+(9765625a^{10}-9765625a^8b^2)Z^{10}-1953125a^8Z^8+156250a^6Z^6-6250a^4Z^4+125a^2Z^2)} -R \ln \left(e^x + \left(\frac{-R^8+4R^6-6R^4+4R^2-1}{R^9+R^7+R^5+R^3+R} \right)^{1/2} \right)$

[In] int(1/(a-b*cosh(x)^5),x,method=_RETURNVERBOSE)

[Out] 1/5*sum((-R^8+4R^6-6R^4+4R^2-1)/(R^9+R^7+R^5+R^3+R)-R*ln(tanh(1/2*x)-R),_R=RootOf((a+b)*Z^10+(-5*a+5*b)*Z^8+(10*a+10*b)*Z^6+(-10*a+10*b)*Z^4+(5*a+5*b)*Z^2-a+b))

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{a - b \cosh^5(x)} dx = \text{Exception raised: RuntimeError}$$

[In] integrate(1/(a-b*cosh(x)^5),x, algorithm="fricas")

[Out] Exception raised: RuntimeError >> no explicit roots found

Sympy [F]

$$\int \frac{1}{a - b \cosh^5(x)} dx = \int \frac{1}{a - b \cosh^5(x)} dx$$

[In] integrate(1/(a-b*cosh(x)**5),x)

[Out] Integral(1/(a - b*cosh(x)**5), x)

Maxima [F]

$$\int \frac{1}{a - b \cosh^5(x)} dx = \int -\frac{1}{b \cosh(x)^5 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^5),x, algorithm="maxima")

[Out] -integrate(1/(b*cosh(x)^5 - a), x)

Giac [F]

$$\int \frac{1}{a - b \cosh^5(x)} dx = \int -\frac{1}{b \cosh(x)^5 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^5),x, algorithm="giac")

[Out] integrate(-1/(b*cosh(x)^5 - a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{a - b \cosh^5(x)} dx = \text{Hanged}$$

```
[In] int(1/(a - b*cosh(x)^5),x)
```

```
[Out] \text{Hanged}
```

3.68 $\int \frac{1}{a-b \cosh^6(x)} dx$

Optimal result	446
Rubi [A] (verified)	446
Mathematica [C] (verified)	448
Maple [C] (verified)	449
Fricas [C] (verification not implemented)	449
Sympy [F]	449
Maxima [F]	450
Giac [F]	450
Mupad [B] (verification not implemented)	450

Optimal result

Integrand size = 11, antiderivative size = 175

$$\int \frac{1}{a-b \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a}-\sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a}-\sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a}+\sqrt[3]{-1}\sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a}+\sqrt[3]{-1}\sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a}-(-1)^{2/3}\sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a}-(-1)^{2/3}\sqrt[3]{b}}}$$

[Out] $\frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3}-b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3}-b^{1/3})^{1/2} + \frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3}+(-1)^{1/3}b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3}+(-1)^{1/3}b^{1/3})^{1/2} + \frac{1}{3} \operatorname{arctanh}\left(\frac{a^{1/6} \tanh(x)}{(a^{1/3}-(-1)^{2/3}b^{1/3})^{1/2}}\right) / a^{5/6} / (a^{1/3}-(-1)^{2/3}b^{1/3})^{1/2}$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 175, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used

= {3290, 3260, 212}

$$\int \frac{1}{a - b \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1} \sqrt[3]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}}}\right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - (-1)^{2/3} \sqrt[3]{b}}}$$

[In] Int[(a - b*Cosh[x]^6)^(-1), x]

[Out] ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) - b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) - b^(1/3)]) + ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) + (-1)^(1/3)*b^(1/3)]) + ArcTanh[(a^(1/6)*Tanh[x])/Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)]]/(3*a^(5/6)*Sqrt[a^(1/3) - (-1)^(2/3)*b^(1/3)])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2)], x], x, Tan[e + f*x]/ff, x]] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))^(n_)^(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rubi steps

$$\text{integral} = \frac{\int \frac{1}{1 - \frac{\sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a} + \frac{\int \frac{1}{1 + \frac{\sqrt[3]{-1} \sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a} + \frac{\int \frac{1}{1 - \frac{(-1)^{2/3} \sqrt[3]{b} \cosh^2(x)}{\sqrt[3]{a}}} dx}{3a}$$

$$\begin{aligned}
& \text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right) \\
= & \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right)}{3a} \\
& + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{\sqrt[3]{-1}\sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right)}{3a} \\
& + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{(-1)^{2/3}\sqrt[3]{b}}{\sqrt[3]{a}}\right) x^2} dx, x, \coth(x) \right)}{3a} \\
= & \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - \sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - \sqrt[3]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} + \sqrt[3]{-1}\sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} + \sqrt[3]{-1}\sqrt[3]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[6]{a} \tanh(x)}{\sqrt{\sqrt[3]{a} - (-1)^{2/3}\sqrt[3]{b}}} \right)}{3a^{5/6} \sqrt{\sqrt[3]{a} - (-1)^{2/3}\sqrt[3]{b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.04 (sec) , antiderivative size = 132, normalized size of antiderivative = 0.75

$$\begin{aligned}
& \int \frac{1}{a - b \cosh^6(x)} dx \\
= & -\frac{16}{3} \text{RootSum} \left[b + 6b\#1 + 15b\#1^2 - 64a\#1^3 + 20b\#1^3 + 15b\#1^4 + 6b\#1^5 \right. \\
& \left. + b\#1^6 \&, \frac{x\#1^2 + \log(-\cosh(x) - \sinh(x) + \cosh(x)\#1 - \sinh(x)\#1)\#1^2}{b + 5b\#1 - 32a\#1^2 + 10b\#1^2 + 10b\#1^3 + 5b\#1^4 + b\#1^5} \& \right]
\end{aligned}$$

[In] Integrate[(a - b*Cosh[x]^6)^(-1),x]

[Out] (-16*RootSum[b + 6*b*#1 + 15*b*#1^2 - 64*a*#1^3 + 20*b*#1^3 + 15*b*#1^4 + 6*b*#1^5 + b*#1^6 &, (x*#1^2 + Log[-Cosh[x] - Sinh[x] + Cosh[x]*#1 - Sinh[x]*#1]*#1^2)/(b + 5*b*#1 - 32*a*#1^2 + 10*b*#1^2 + 10*b*#1^3 + 5*b*#1^4 + b*#1^5) &]/3

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.94 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.80

method	result
risch	$\sum_{-R=\text{RootOf}(-1+(46656a^6-46656a^5b)_Z^6-3888a^4_Z^4+108a^2_Z^2)} _R \ln \left(e^{2x} + \left(\frac{15552a^6}{b} - 15552a^5 \right) _R^5 + \left(\dots \right) \right)$
default	$\left(\sum_{-R=\text{RootOf}((a-b)_Z^{12}+(-6a-6b)_Z^{10}+(15a-15b)_Z^8+(-20a-20b)_Z^6+(15a-15b)_Z^4+(-6a-6b)_Z^2+a-b)} \frac{_R^{11} _a - _R^{11} _b - 5}{6} \right)$

[In] `int(1/(a-b*cosh(x)^6),x,method=_RETURNVERBOSE)`

[Out] `sum(_R*ln(exp(2*x)+(15552*a^6/b-15552*a^5)*_R^5+(-2592*a^5/b+2592*a^4)*_R^4+(-864*a^4/b-432*a^3)*_R^3+(144/b*a^3+72*a^2)*_R^2+(12*a^2/b-12*a)*_R-2*a/b+1),_R=RootOf(-1+(46656*a^6-46656*a^5*b)*_Z^6-3888*a^4*_Z^4+108*a^2*_Z^2))`

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 1.16 (sec) , antiderivative size = 16379, normalized size of antiderivative = 93.59

$$\int \frac{1}{a - b \cosh^6(x)} dx = \text{Too large to display}$$

[In] `integrate(1/(a-b*cosh(x)^6),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{1}{a - b \cosh^6(x)} dx = \int \frac{1}{a - b \cosh^6(x)} dx$$

[In] `integrate(1/(a-b*cosh(x)**6),x)`

[Out] `Integral(1/(a - b*cosh(x)**6), x)`

$$\begin{aligned}
& p(2*x) + 1770440*a^5*b^2*\exp(2*x) - 1239040*a^6*b*\exp(2*x)) / (b^{10}*(a - b)^3) \\
& + (17509995351216488448*\text{root}(46656*a^5*b*d^6 - 46656*a^6*d^6 + 3888*a^4*d^4 - 108*a^2*d^2 + 1, d, k) * (262144*a^7*\exp(2*x) - 203520*a^6*b + 65536*a^7 + 453*a^3*b^4 - 72022*a^4*b^3 + 209472*a^5*b^2 + 630*a^3*b^4*\exp(2*x) - 254512*a^4*b^3*\exp(2*x) + 767508*a^5*b^2*\exp(2*x) - 775680*a^6*b*\exp(2*x))) / \\
& (b^{10}*(a - b)^2) - (486388759756013568*(655360*a^5*\exp(2*x) - 9*a*b^4 - 370176*a^4*b + 196608*a^5 + 24408*a^2*b^3 + 149088*a^3*b^2 + 63676*a^2*b^3*\exp(2*x) + 526248*a^3*b^2*\exp(2*x) - 10*a*b^4*\exp(2*x) - 1245184*a^4*b*\exp(2*x))) / (b^{10}*(a - b)^2) - (40532396646334464*(655360*a^5*\exp(2*x) + b^5*\exp(2*x) - 24677*a*b^4 - 773120*a^4*b + 262144*a^5 + b^5 - 198071*a^2*b^3 + 733696*a^3*b^2 - 477713*a^2*b^3*\exp(2*x) + 1770640*a^3*b^2*\exp(2*x) - 53861*a*b^4*\exp(2*x) - 1894400*a^4*b*\exp(2*x))) / (b^{10}*(a - b)^3) + (13510798882111488*(655360*a^3*\exp(2*x) - 11382*b^3*\exp(2*x) + 144416*a*b^2 - 269056*a^2*b + 131072*a^3 - 6459*b^3 + 677524*a*b^2*\exp(2*x) - 1321472*a^2*b*\exp(2*x))) / (b^{10}*(a - b)^2) - (1125899906842624*(851968*a^4*\exp(2*x) + 6006*b^4*\exp(2*x) - 211497*a*b^3 - 597504*a^3*b + 196608*a^4 + 3840*b^4 + 608544*a^2*b^2 + 2562504*a^2*b^2*\exp(2*x) - 864565*a*b^3*\exp(2*x) - 2555904*a^3*b*\exp(2*x))) / (b^{10}*(a - b)^2*(a*b - a^2)) * \text{root}(46656*a^5*b*d^6 - 46656*a^6*d^6 + 3888*a^4*d^4 - 108*a^2*d^2 + 1, d, k), k, 1, 6)
\end{aligned}$$

3.69 $\int \frac{1}{a-b \cosh^8(x)} dx$

Optimal result	452
Rubi [A] (verified)	452
Mathematica [C] (verified)	454
Maple [C] (verified)	455
Fricas [B] (verification not implemented)	455
Sympy [F]	455
Maxima [F]	456
Giac [F]	456
Mupad [F(-1)]	456

Optimal result

Integrand size = 11, antiderivative size = 213

$$\int \frac{1}{a-b \cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a}-\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a}-\sqrt[4]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a}-i\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a}-i\sqrt[4]{b}}} \\ + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a}+i\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a}+i\sqrt[4]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a}+\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a}+\sqrt[4]{b}}}$$

[Out] $\frac{1}{4} \operatorname{arctanh}\left(\frac{a^{1/8} \tanh(x)}{(a^{1/4}-b^{1/4})^{1/2}}\right) / a^{7/8} / (a^{1/4}-b^{1/4})^{1/2} + \frac{1}{4} \operatorname{arctanh}\left(\frac{a^{1/8} \tanh(x)}{(a^{1/4}-I b^{1/4})^{1/2}}\right) / a^{7/8} / (a^{1/4}-I b^{1/4})^{1/2} + \frac{1}{4} \operatorname{arctanh}\left(\frac{a^{1/8} \tanh(x)}{(a^{1/4}+I b^{1/4})^{1/2}}\right) / a^{7/8} / (a^{1/4}+I b^{1/4})^{1/2} + \frac{1}{4} \operatorname{arctanh}\left(\frac{a^{1/8} \tanh(x)}{(a^{1/4}+b^{1/4})^{1/2}}\right) / a^{7/8} / (a^{1/4}+b^{1/4})^{1/2}$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 213, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.273$, Rules used

= {3290, 3260, 212}

$$\int \frac{1}{a - b \cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a} - \sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a} - \sqrt[4]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a} - i\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a} - i\sqrt[4]{b}}} \\ + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a} + i\sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a} + i\sqrt[4]{b}}} + \frac{\operatorname{arctanh}\left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt{a} + \sqrt[4]{b}}}\right)}{4a^{7/8} \sqrt[4]{\sqrt{a} + \sqrt[4]{b}}}$$

[In] Int[(a - b*Cosh[x]^8)^(-1), x]

[Out] ArcTanh[(a^(1/8)*Tanh[x])/Sqrt[a^(1/4) - b^(1/4)]]/(4*a^(7/8)*Sqrt[a^(1/4) - b^(1/4)]) + ArcTanh[(a^(1/8)*Tanh[x])/Sqrt[a^(1/4) - I*b^(1/4)]]/(4*a^(7/8)*Sqrt[a^(1/4) - I*b^(1/4)]) + ArcTanh[(a^(1/8)*Tanh[x])/Sqrt[a^(1/4) + I*b^(1/4)]]/(4*a^(7/8)*Sqrt[a^(1/4) + I*b^(1/4)]) + ArcTanh[(a^(1/8)*Tanh[x])/Sqrt[a^(1/4) + b^(1/4)]]/(4*a^(7/8)*Sqrt[a^(1/4) + b^(1/4)])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rubi steps

$$\text{integral} = \frac{\int \frac{1}{1 - \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{a}}} dx}{4a} + \frac{\int \frac{1}{1 - i \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{a}}} dx}{4a} + \frac{\int \frac{1}{1 + i \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{a}}} dx}{4a} + \frac{\int \frac{1}{1 + \frac{\sqrt[4]{b} \cosh^2(x)}{\sqrt[4]{a}}} dx}{4a}$$

$$\begin{aligned}
& \text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right) + \text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{i\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right) \\
&= \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right)}{4a} + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 - \frac{i\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right)}{4a} \\
&+ \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{i\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right)}{4a} + \frac{\text{Subst} \left(\int \frac{1}{1 - \left(1 + \frac{\sqrt[4]{b}}{\sqrt[4]{a}}\right) x^2} dx, x, \coth(x) \right)}{4a} \\
&= \frac{\text{arctanh} \left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt[4]{a} - \sqrt[4]{b}}} \right)}{4a^{7/8} \sqrt[4]{\sqrt[4]{a} - \sqrt[4]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt[4]{a} - i\sqrt[4]{b}}} \right)}{4a^{7/8} \sqrt[4]{\sqrt[4]{a} - i\sqrt[4]{b}}} \\
&+ \frac{\text{arctanh} \left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt[4]{a} + i\sqrt[4]{b}}} \right)}{4a^{7/8} \sqrt[4]{\sqrt[4]{a} + i\sqrt[4]{b}}} + \frac{\text{arctanh} \left(\frac{\sqrt[8]{a} \tanh(x)}{\sqrt[4]{\sqrt[4]{a} + \sqrt[4]{b}}} \right)}{4a^{7/8} \sqrt[4]{\sqrt[4]{a} + \sqrt[4]{b}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 2.63 (sec) , antiderivative size = 158, normalized size of antiderivative = 0.74

$$\int \frac{1}{a - b \cosh^8(x)} dx = -16 \text{RootSum} \left[b + 8b\#1 + 28b\#1^2 + 56b\#1^3 - 256a\#1^4 + 70b\#1^4 \right. \\
\left. + 56b\#1^5 + 28b\#1^6 + 8b\#1^7 \right. \\
\left. + b\#1^8 \&, \frac{x\#1^3 + \log(-\cosh(x) - \sinh(x) + \cosh(x)\#1 - \sinh(x)\#1)\#1^3}{b + 7b\#1 + 21b\#1^2 - 128a\#1^3 + 35b\#1^3 + 35b\#1^4 + 21b\#1^5 + 7b\#1^6 + b\#1^7} \& \right]$$

[In] Integrate[(a - b*Cosh[x]^8)^(-1), x]

[Out] -16*RootSum[b + 8*b*#1 + 28*b*#1^2 + 56*b*#1^3 - 256*a*#1^4 + 70*b*#1^4 + 56*b*#1^5 + 28*b*#1^6 + 8*b*#1^7 + b*#1^8 & , (x*#1^3 + Log[-Cosh[x] - Sinh[x] + Cosh[x]*#1 - Sinh[x]*#1]*#1^3)/(b + 7*b*#1 + 21*b*#1^2 - 128*a*#1^3 + 35*b*#1^3 + 35*b*#1^4 + 21*b*#1^5 + 7*b*#1^6 + b*#1^7) &]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 2.38 (sec) , antiderivative size = 184, normalized size of antiderivative = 0.86

method	result
risch	$\sum_{\substack{_R=\text{RootOf}(1+(16777216a^8-16777216a^7b)_Z^8-1048576a^6_Z^6+24576a^4_Z^4-256a^2_Z^2)}} _R \ln \left(e^{2x} + \left(\frac{4194304a^8}{b} - 4 \right) \right)$
default	$\left(\sum_{\substack{_R=\text{RootOf}((a-b)_Z^{16}+(-8a-8b)_Z^{14}+(28a-28b)_Z^{12}+(-56a-56b)_Z^{10}+(70a-70b)_Z^8+(-56a-56b)_Z^6+(28a-28b)_Z^4+(-8a-8b)_Z^2}} \right)$

[In] `int(1/(a-b*cosh(x)^8),x,method=_RETURNVERBOSE)`

[Out] `sum(_R*ln(exp(2*x)+(4194304*a^8/b-4194304*a^7)*_R^7+(-524288*a^7/b+524288*a^6)*_R^6+(-196608*a^6/b-65536*a^5)*_R^5+(24576*a^5/b+8192*a^4)*_R^4+(3072*a^4/b-1024*a^3)*_R^3+(-384/b*a^3+128*a^2)*_R^2+(-16*a^2/b-16*a)*_R+2*a/b+1),_R=RootOf(1+(16777216*a^8-16777216*a^7*b)*_Z^8-1048576*a^6*_Z^6+24576*a^4*_Z^4-256*a^2*_Z^2))`

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 631813 vs. 2(133) = 266.

Time = 3.11 (sec) , antiderivative size = 631813, normalized size of antiderivative = 2966.26

$$\int \frac{1}{a - b \cosh^8(x)} dx = \text{Too large to display}$$

[In] `integrate(1/(a-b*cosh(x)^8),x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int \frac{1}{a - b \cosh^8(x)} dx = \int \frac{1}{a - b \cosh^8(x)} dx$$

[In] `integrate(1/(a-b*cosh(x)**8),x)`

[Out] `Integral(1/(a - b*cosh(x)**8), x)`

Maxima [F]

$$\int \frac{1}{a - b \cosh^8(x)} dx = \int -\frac{1}{b \cosh(x)^8 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^8),x, algorithm="maxima")

[Out] -integrate(1/(b*cosh(x)^8 - a), x)

Giac [F]

$$\int \frac{1}{a - b \cosh^8(x)} dx = \int -\frac{1}{b \cosh(x)^8 - a} dx$$

[In] integrate(1/(a-b*cosh(x)^8),x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{a - b \cosh^8(x)} dx = \text{Hanged}$$

[In] int(1/(a - b*cosh(x)^8),x)

[Out] \text{Hanged}

3.70 $\int \frac{1}{1+\cosh^5(x)} dx$

Optimal result	457
Rubi [A] (verified)	458
Mathematica [C] (verified)	460
Maple [C] (verified)	460
Fricas [B] (verification not implemented)	461
Sympy [F(-2)]	462
Maxima [F]	462
Giac [F]	462
Mupad [F(-1)]	462

Optimal result

Integrand size = 8, antiderivative size = 223

$$\int \frac{1}{1+\cosh^5(x)} dx = -\frac{2 \arctan\left(\frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{\frac{-1-\sqrt[5]{-1}}{1+\sqrt[5]{-1}}}}\right)}{5\sqrt{-1+(-1)^{2/5}}} - \frac{2\sqrt{-\frac{1+(-1)^{3/5}}{1-(-1)^{3/5}}}}{5(1+(-1)^{3/5})} \arctan\left(\sqrt{-\frac{1+(-1)^{3/5}}{1-(-1)^{3/5}}} \tanh\left(\frac{x}{2}\right)\right) + \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1-(-1)^{2/5}}{1+(-1)^{2/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1-(-1)^{4/5}}} + \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1-(-1)^{4/5}}{1+(-1)^{4/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1+(-1)^{3/5}}} + \frac{\sinh(x)}{5(1+\cosh(x))}$$

```
[Out] 1/5*sinh(x)/(1+cosh(x))-2/5*arctan(tanh(1/2*x)/((-1+(-1)^(1/5))/(1+(-1)^(1/5)))^(1/2)/(-1+(-1)^(2/5))^(1/2)+2/5*arctanh(((1-(-1)^(4/5))/(1+(-1)^(4/5)))^(1/2)*tanh(1/2*x))/(1+(-1)^(3/5))^(1/2)-2/5*arctan(((1-(-1)^(3/5))/(1-(-1)^(3/5)))^(1/2)*tanh(1/2*x))*((-1-(-1)^(3/5))/(1-(-1)^(3/5)))^(1/2)/(1+(-1)^(3/5))+2/5*arctanh(((1-(-1)^(2/5))/(1+(-1)^(2/5)))^(1/2)*tanh(1/2*x))/(1-(-1)^(4/5))^(1/2)
```

Rubi [A] (verified)

Time = 0.44 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3292, 2727, 2738, 211, 214}

$$\int \frac{1}{1 + \cosh^5(x)} dx = -\frac{2 \arctan\left(\frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{\frac{-1 - \sqrt[5]{-1}}{1 + \sqrt[5]{-1}}}}\right)}{5\sqrt{(-1)^{2/5} - 1}} - \frac{2\sqrt{-\frac{1+(-1)^{3/5}}{1-(-1)^{3/5}}} \arctan\left(\sqrt{-\frac{1+(-1)^{3/5}}{1-(-1)^{3/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5(1 + (-1)^{3/5})} + \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1-(-1)^{2/5}}{1+(-1)^{2/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1 - (-1)^{4/5}}} + \frac{2\operatorname{arctanh}\left(\sqrt{\frac{1-(-1)^{4/5}}{1+(-1)^{4/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1 + (-1)^{3/5}}} + \frac{\sinh(x)}{5(\cosh(x) + 1)}$$

[In] Int[(1 + Cosh[x]^5)^(-1), x]

[Out] (-2*ArcTan[Tanh[x/2]/Sqrt[-((1 - (-1)^(1/5))/(1 + (-1)^(1/5))]])/(5*Sqrt[-1 + (-1)^(2/5)]) - (2*Sqrt[-((1 + (-1)^(3/5))/(1 - (-1)^(3/5)))]*ArcTan[Sqrt[-((1 + (-1)^(3/5))/(1 - (-1)^(3/5)))]*Tanh[x/2]])/(5*(1 + (-1)^(3/5))) + (2*ArcTanh[Sqrt[(1 - (-1)^(2/5))/(1 + (-1)^(2/5))]*Tanh[x/2]])/(5*Sqrt[1 - (-1)^(4/5)]) + (2*ArcTanh[Sqrt[(1 - (-1)^(4/5))/(1 + (-1)^(4/5))]*Tanh[x/2]])/(5*Sqrt[1 + (-1)^(3/5)]) + Sinh[x]/(5*(1 + Cosh[x]))

Rule 211

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2727

Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 2738

Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3292

Int[((a_) + (b_)*((c_)*sin[(e_) + (f_)*(x_)])^(n_))^(p_), x_Symbol] := Int[ExpandTrig[(a + b*(c*sin[e + f*x])^n)^p, x], x] /; FreeQ[{a, b, c, e, f, n}, x] && (IGtQ[p, 0] || (EqQ[p, -1] && IntegerQ[n]))

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(-\frac{1}{5(-1 - \cosh(x))} - \frac{1}{5(-1 + \sqrt[5]{-1} \cosh(x))} - \frac{1}{5(-1 - (-1)^{2/5} \cosh(x))} \right. \\
 &\quad \left. - \frac{1}{5(-1 + (-1)^{3/5} \cosh(x))} - \frac{1}{5(-1 - (-1)^{4/5} \cosh(x))} \right) dx \\
 &= -\left(\frac{1}{5} \int \frac{1}{-1 - \cosh(x)} dx \right) - \frac{1}{5} \int \frac{1}{-1 + \sqrt[5]{-1} \cosh(x)} dx \\
 &\quad - \frac{1}{5} \int \frac{1}{-1 - (-1)^{2/5} \cosh(x)} dx - \frac{1}{5} \int \frac{1}{-1 + (-1)^{3/5} \cosh(x)} dx \\
 &\quad - \frac{1}{5} \int \frac{1}{-1 - (-1)^{4/5} \cosh(x)} dx \\
 &= \frac{\sinh(x)}{5(1 + \cosh(x))} - \frac{2}{5} \text{Subst} \left(\int \frac{1}{-1 + \sqrt[5]{-1} - (-1 - \sqrt[5]{-1}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
 &\quad - \frac{2}{5} \text{Subst} \left(\int \frac{1}{-1 - (-1)^{2/5} - (-1 + (-1)^{2/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
 &\quad - \frac{2}{5} \text{Subst} \left(\int \frac{1}{-1 + (-1)^{3/5} - (-1 - (-1)^{3/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) - \frac{2}{5} \text{Subst} \left(\int \frac{1}{-1 - (-1)^{4/5} - (-1 + (-1)^{4/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
 &= -\frac{2 \arctan \left(\frac{\tanh \left(\frac{x}{2} \right)}{\sqrt{\frac{-1 - \sqrt[5]{-1}}{1 + \sqrt[5]{-1}}}} \right)}{5 \sqrt{-1 + (-1)^{2/5}}} - \frac{2 \sqrt{\frac{-1 + (-1)^{3/5}}{1 - (-1)^{3/5}}} \arctan \left(\sqrt{\frac{-1 + (-1)^{3/5}}{1 - (-1)^{3/5}}} \tanh \left(\frac{x}{2} \right) \right)}{5 (1 + (-1)^{3/5})} \\
 &\quad + \frac{2 \text{arctanh} \left(\sqrt{\frac{1 - (-1)^{2/5}}{1 + (-1)^{2/5}}} \tanh \left(\frac{x}{2} \right) \right)}{5 \sqrt{1 - (-1)^{4/5}}} + \frac{2 \text{arctanh} \left(\sqrt{\frac{1 - (-1)^{4/5}}{1 + (-1)^{4/5}}} \tanh \left(\frac{x}{2} \right) \right)}{5 \sqrt{1 + (-1)^{3/5}}} \\
 &\quad + \frac{\sinh(x)}{5(1 + \cosh(x))}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.03 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.00

$$\int \frac{1}{1 + \cosh^5(x)} dx$$

$$= -\frac{1}{10} \text{RootSum} \left[1 - 2\#1 + 8\#1^2 - 14\#1^3 + 30\#1^4 - 14\#1^5 + 8\#1^6 - 2\#1^7 \right. \\ \left. + \#1^8 \&, \frac{x + 2 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) + \cosh \left(\frac{x}{2} \right) \#1 - \sinh \left(\frac{x}{2} \right) \#1 \right) - 4x\#1 - 8 \log \left(-\cosh \left(\frac{x}{2} \right) - \sinh \left(\frac{x}{2} \right) \right)}{\#1} \right. \\ \left. + \frac{1}{5} \tanh \left(\frac{x}{2} \right) \right]$$

[In] Integrate[(1 + Cosh[x]^5)^(-1),x]

[Out] -1/10*RootSum[1 - 2*#1 + 8*#1^2 - 14*#1^3 + 30*#1^4 - 14*#1^5 + 8*#1^6 - 2*#1^7 + #1^8 & , (x + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1] - 4*x*#1 - 8*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1 + 15*x*#1^2 + 30*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^2 - 40*x*#1^3 - 80*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^3 + 15*x*#1^4 + 30*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^4 - 4*x*#1^5 - 8*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^5 + x*#1^6 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/(-1 + 8*#1 - 21*#1^2 + 60*#1^3 - 35*#1^4 + 24*#1^5 - 7*#1^6 + 4*#1^7) &] + Tanh[x/2]/5

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.13 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.28

method	result
default	$\frac{\tanh\left(\frac{x}{2}\right)}{5} + \frac{\left(\sum_{R=\text{RootOf}(5Z^8+10Z^4+1)} \frac{(-5R^6+5R^4-5R^2+1) \ln\left(\tanh\left(\frac{x}{2}\right)-R\right)}{-R^7+R^3} \right)}{50}$
risch	$-\frac{2}{5(e^x+1)} + \left(\sum_{R=\text{RootOf}(1953125Z^8-156250Z^6+6250Z^4-125Z^2+1)} -R \ln(2343750R^7 - 234375R^6 - \dots) \right)$

[In] int(1/(1+cosh(x)^5),x,method=_RETURNVERBOSE)

[Out] $\frac{1}{5} \tanh\left(\frac{1}{2}x\right) + \frac{1}{50} \sum\left(\frac{-5R^6 + 5R^4 - 5R^2 + 1}{R^7 + R^3} \ln\left(\tanh\left(\frac{1}{2}x\right) - R\right), R = \sqrt[5]{5Z^8 + 10Z^4 + 1}\right)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 836 vs. $2(150) = 300$.

Time = 0.30 (sec) , antiderivative size = 836, normalized size of antiderivative = 3.75

$$\int \frac{1}{1 + \cosh^5(x)} dx = \text{Too large to display}$$

[In] integrate(1/(1+cosh(x)^5),x, algorithm="fricas")

[Out] $\frac{1}{50} \left((\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(\sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} + 5) \sqrt{2\sqrt{5} - 5} - 5\sqrt{2\sqrt{5} - 5} (\sqrt{5} + 3) + 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) - (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(-\sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} + 5) \sqrt{2\sqrt{5} - 5} - 5\sqrt{2\sqrt{5} - 5} (\sqrt{5} + 3) + 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) - (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(\sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} + 5) \sqrt{2\sqrt{5} - 5} + 5\sqrt{2\sqrt{5} - 5} (\sqrt{5} + 3) + 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) + (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(-\sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} + 5) \sqrt{2\sqrt{5} - 5} + 5\sqrt{2\sqrt{5} - 5} (\sqrt{5} + 3) + 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) - (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(\sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} - 5) \sqrt{-2\sqrt{5} - 5} - 5(\sqrt{5} - 3) \sqrt{-2\sqrt{5} - 5} - 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) + (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(-\sqrt{2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} - 5) \sqrt{-2\sqrt{5} - 5} - 5(\sqrt{5} - 3) \sqrt{-2\sqrt{5} - 5} - 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) + (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(\sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} - 5) \sqrt{-2\sqrt{5} - 5} + 5(\sqrt{5} - 3) \sqrt{-2\sqrt{5} - 5} - 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) - (\sqrt{5} \cosh(x) + \sqrt{5} \sinh(x) + \sqrt{5}) \sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} \log(-\sqrt{-2\sqrt{5} \sqrt{2\sqrt{5} - 5} + 10} (3\sqrt{5} - 5) \sqrt{-2\sqrt{5} - 5} + 5(\sqrt{5} - 3) \sqrt{-2\sqrt{5} - 5} - 5\sqrt{5} + 20 \cosh(x) + 20 \sinh(x) - 5) - 20) / (\cosh(x) + \sinh(x) + 1) \right)$

Sympy [F(-2)]

Exception generated.

$$\int \frac{1}{1 + \cosh^5(x)} dx = \text{Exception raised: ValueError}$$

[In] `integrate(1/(1+cosh(x)**5),x)`

[Out] Exception raised: ValueError >> Exceeds the limit (4300 digits) for integer string conversion; use `sys.set_int_max_str_digits()` to increase the limit

Maxima [F]

$$\int \frac{1}{1 + \cosh^5(x)} dx = \int \frac{1}{\cosh(x)^5 + 1} dx$$

[In] `integrate(1/(1+cosh(x)^5),x, algorithm="maxima")`

[Out] $-2/5/(e^x + 1) - \text{integrate}(2/5*(e^{7*x} - 4*e^{6*x} + 15*e^{5*x} - 40*e^{4*x} + 15*e^{3*x} - 4*e^{2*x} + e^x)/(e^{8*x} - 2*e^{7*x} + 8*e^{6*x} - 14*e^{5*x} + 30*e^{4*x} - 14*e^{3*x} + 8*e^{2*x} - 2*e^x + 1), x)$

Giac [F]

$$\int \frac{1}{1 + \cosh^5(x)} dx = \int \frac{1}{\cosh(x)^5 + 1} dx$$

[In] `integrate(1/(1+cosh(x)^5),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{1 + \cosh^5(x)} dx = \text{Hanged}$$

[In] `int(1/(cosh(x)^5 + 1),x)`

[Out] `\text{Hanged}`

3.71 $\int \frac{1}{1+\cosh^6(x)} dx$

Optimal result	463
Rubi [A] (verified)	463
Mathematica [C] (verified)	464
Maple [C] (verified)	465
Fricas [C] (verification not implemented)	465
Sympy [F(-1)]	466
Maxima [F]	466
Giac [B] (verification not implemented)	466
Mupad [B] (verification not implemented)	467

Optimal result

Integrand size = 8, antiderivative size = 83

$$\int \frac{1}{1+\cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{3\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-\sqrt[3]{-1}}}\right)}{3\sqrt{1-\sqrt[3]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+(-1)^{2/3}}}\right)}{3\sqrt{1+(-1)^{2/3}}}$$

[Out] 1/6*arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)+1/3*arctanh(tanh(x)/(1-(-1)^(1/3))^(1/2))/(1-(-1)^(1/3))^(1/2)+1/3*arctanh(tanh(x)/(1+(-1)^(2/3))^(1/2))/(1+(-1)^(2/3))^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec), antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3290, 3260, 212}

$$\int \frac{1}{1+\cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{3\sqrt{2}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-\sqrt[3]{-1}}}\right)}{3\sqrt{1-\sqrt[3]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+(-1)^{2/3}}}\right)}{3\sqrt{1+(-1)^{2/3}}}$$

[In] Int[(1 + Cosh[x]^6)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[2]]/(3*Sqrt[2]) + ArcTanh[Tanh[x]/Sqrt[1 - (-1)^(1/3)]]/(3*Sqrt[1 - (-1)^(1/3)]) + ArcTanh[Tanh[x]/Sqrt[1 + (-1)^(2/3)]]/(3*Sqrt[1 + (-1)^(2/3)])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

Q[a, 0] || LtQ[b, 0])

Rule 3260

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff =
FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2
), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]
```

Rule 3290

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^n)^(-1), x_Symbol] := Module[{
k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/
2]))], x], {k, 1, n/2}], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{3} \int \frac{1}{1 + \cosh^2(x)} dx + \frac{1}{3} \int \frac{1}{1 - \sqrt[3]{-1} \cosh^2(x)} dx + \frac{1}{3} \int \frac{1}{1 + (-1)^{2/3} \cosh^2(x)} dx \\
&= \frac{1}{3} \text{Subst} \left(\int \frac{1}{1 - 2x^2} dx, x, \coth(x) \right) \\
&\quad + \frac{1}{3} \text{Subst} \left(\int \frac{1}{1 - (1 - \sqrt[3]{-1}) x^2} dx, x, \coth(x) \right) \\
&\quad + \frac{1}{3} \text{Subst} \left(\int \frac{1}{1 - (1 + (-1)^{2/3}) x^2} dx, x, \coth(x) \right) \\
&= \frac{\text{arctanh} \left(\frac{\tanh(x)}{\sqrt{2}} \right)}{3\sqrt{2}} + \frac{\text{arctanh} \left(\frac{\tanh(x)}{\sqrt{1 - \sqrt[3]{-1}}} \right)}{3\sqrt{1 - \sqrt[3]{-1}}} + \frac{\text{arctanh} \left(\frac{\tanh(x)}{\sqrt{1 + (-1)^{2/3}}} \right)}{3\sqrt{1 + (-1)^{2/3}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 5.05 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.82

$$\begin{aligned}
\int \frac{1}{1 + \cosh^6(x)} dx &= \frac{1}{6} \left(\arctan(\text{csch}(x)\text{sech}(x)) \right. \\
&\quad \left. + i\sqrt{3} \left(\arctan \left(\frac{1 - 2i \tanh(x)}{\sqrt{3}} \right) - \arctan \left(\frac{1 + 2i \tanh(x)}{\sqrt{3}} \right) \right) \right. \\
&\quad \left. + \sqrt{2} \text{arctanh} \left(\frac{\tanh(x)}{\sqrt{2}} \right) \right)
\end{aligned}$$

[In] Integrate[(1 + Cosh[x]^6)^(-1), x]

[Out] (ArcTan[Csch[x]*Sech[x]] + I*Sqrt[3]*(ArcTan[(1 - (2*I)*Tanh[x])/Sqrt[3]] - ArcTan[(1 + (2*I)*Tanh[x])/Sqrt[3]]) + Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]])/6

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.86 (sec) , antiderivative size = 71, normalized size of antiderivative = 0.86

method	result
risch	$\frac{\sqrt{2} \ln(e^{2x} + 3 - 2\sqrt{2})}{12} - \frac{\sqrt{2} \ln(e^{2x} + 3 + 2\sqrt{2})}{12} + \left(\sum_{R=\text{RootOf}(1296Z^4 - 36Z^2 + 1)} _R \ln(-432_R^3 + 72_R^2 + \dots) \right)$
default	$\left(\frac{\sum_{R=\text{RootOf}(-Z^4 - 2Z^3 + 2Z^2 + 2Z + 1)} \left(\frac{(-R^2 + 4R + 1) \ln(\tanh(\frac{x}{2}) - R)}{2R^3 - 3R^2 + 2R + 1} \right)}{6} \right) + \frac{\sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})^2 + \tanh(\frac{x}{2})\sqrt{2} + 1}{\tanh(\frac{x}{2})^2 - \tanh(\frac{x}{2})\sqrt{2} + 1} \right) + 2 \arctan\left(\frac{\tanh(\frac{x}{2}) + \sqrt{2}}{\tanh(\frac{x}{2}) - \sqrt{2}} \right) \right)}{6}$

[In] int(1/(1+cosh(x)^6),x,method=_RETURNVERBOSE)

[Out] 1/12*2^(1/2)*ln(exp(2*x)+3-2*2^(1/2))-1/12*2^(1/2)*ln(exp(2*x)+3+2*2^(1/2))
+sum(_R*ln(-432*_R^3+72*_R^2+exp(2*x)-1),_R=RootOf(1296*_Z^4-36*_Z^2+1))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 315, normalized size of antiderivative = 3.80

$$\begin{aligned} \int \frac{1}{1 + \cosh^6(x)} dx = & -\frac{1}{12} \sqrt{2} \sqrt{i\sqrt{3} + 1} \log \left(2 \cosh(x)^2 + 4 \cosh(x) \sinh(x) + 2 \sinh(x)^2 \right. \\ & + \left. (i\sqrt{3}\sqrt{2} + \sqrt{2}) \sqrt{i\sqrt{3} + 1} + 2i\sqrt{3} \right) + \frac{1}{12} \sqrt{2} \sqrt{i\sqrt{3} + 1} \log \left(2 \cosh(x)^2 \right. \\ & + \left. 4 \cosh(x) \sinh(x) + 2 \sinh(x)^2 + (-i\sqrt{3}\sqrt{2} - \sqrt{2}) \sqrt{i\sqrt{3} + 1} + 2i\sqrt{3} \right) \\ & + \frac{1}{12} \sqrt{2} \sqrt{-i\sqrt{3} + 1} \log \left(2 \cosh(x)^2 + 4 \cosh(x) \sinh(x) + 2 \sinh(x)^2 \right. \\ & + \left. (i\sqrt{3}\sqrt{2} - \sqrt{2}) \sqrt{-i\sqrt{3} + 1} - 2i\sqrt{3} \right) - \frac{1}{12} \sqrt{2} \sqrt{-i\sqrt{3} + 1} \log \left(2 \cosh(x)^2 \right. \\ & + \left. 4 \cosh(x) \sinh(x) + 2 \sinh(x)^2 + (-i\sqrt{3}\sqrt{2} + \sqrt{2}) \sqrt{-i\sqrt{3} + 1} - 2i\sqrt{3} \right) \\ & + \frac{1}{12} \sqrt{2} \log \left(\frac{-3(2\sqrt{2} - 3) \cosh(x)^2 - 4(3\sqrt{2} - 4) \cosh(x) \sinh(x) + 3(2\sqrt{2} - 3) \sinh(x)^2 + 2\sqrt{2}}{\cosh(x)^2 + \sinh(x)^2 + 3} \right) \end{aligned}$$

[In] integrate(1/(1+cosh(x)^6),x, algorithm="fricas")

[Out] -1/12*sqrt(2)*sqrt(I*sqrt(3) + 1)*log(2*cosh(x)^2 + 4*cosh(x)*sinh(x) + 2*sinh(x)^2 + (I*sqrt(3)*sqrt(2) + sqrt(2))*sqrt(I*sqrt(3) + 1) + 2*I*sqrt(3))
+ 1/12*sqrt(2)*sqrt(I*sqrt(3) + 1)*log(2*cosh(x)^2 + 4*cosh(x)*sinh(x) + 2

*sinh(x)^2 + (-I*sqrt(3)*sqrt(2) - sqrt(2))*sqrt(I*sqrt(3) + 1) + 2*I*sqrt(3)) + 1/12*sqrt(2)*sqrt(-I*sqrt(3) + 1)*log(2*cosh(x)^2 + 4*cosh(x)*sinh(x) + 2*sinh(x)^2 + (I*sqrt(3)*sqrt(2) - sqrt(2))*sqrt(-I*sqrt(3) + 1) - 2*I*sqrt(3)) - 1/12*sqrt(2)*sqrt(-I*sqrt(3) + 1)*log(2*cosh(x)^2 + 4*cosh(x)*sinh(x) + 2*sinh(x)^2 + (-I*sqrt(3)*sqrt(2) + sqrt(2))*sqrt(-I*sqrt(3) + 1) - 2*I*sqrt(3)) + 1/12*sqrt(2)*log(-(3*(2*sqrt(2) - 3)*cosh(x)^2 - 4*(3*sqrt(2) - 4)*cosh(x)*sinh(x) + 3*(2*sqrt(2) - 3)*sinh(x)^2 + 2*sqrt(2) - 3)/(cosh(x)^2 + sinh(x)^2 + 3))

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{1 + \cosh^6(x)} dx = \text{Timed out}$$

[In] integrate(1/(1+cosh(x)**6),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{1 + \cosh^6(x)} dx = \int \frac{1}{\cosh(x)^6 + 1} dx$$

[In] integrate(1/(1+cosh(x)^6),x, algorithm="maxima")

[Out] -1/12*sqrt(2)*log(-(2*sqrt(2) - e^(-2*x) - 3)/(2*sqrt(2) + e^(-2*x) + 3)) - 4/3*integrate(-(6*e^(-2*x) - e^(-4*x) - 1)*e^(-2*x)/(14*e^(-4*x) + e^(-8*x) + 1), x)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(58) = 116.

Time = 0.27 (sec) , antiderivative size = 140, normalized size of antiderivative = 1.69

$$\begin{aligned} \int \frac{1}{1 + \cosh^6(x)} dx &= \frac{1}{36} \left((2\sqrt{3} - 3)e^{(4x)} + 2\sqrt{3} - 3 \right) \arctan \left(\frac{e^{(2x)}}{\sqrt{3} + 2} \right) \\ &\quad - \frac{1}{36} \left((2\sqrt{3} + 3)e^{(4x)} + 2\sqrt{3} + 3 \right) \arctan \left(-\frac{e^{(2x)}}{\sqrt{3} - 2} \right) \\ &\quad - \frac{1}{12} \sqrt{3} \log \left((\sqrt{3} + 2)^2 + e^{(4x)} \right) + \frac{1}{12} \sqrt{3} \log \left((\sqrt{3} - 2)^2 + e^{(4x)} \right) \\ &\quad + \frac{1}{12} \sqrt{2} \log \left(-\frac{2\sqrt{2} - e^{(2x)} - 3}{2\sqrt{2} + e^{(2x)} + 3} \right) \end{aligned}$$

[In] integrate(1/(1+cosh(x)^6),x, algorithm="giac")

[Out] 1/36*((2*sqrt(3) - 3)*e^(4*x) + 2*sqrt(3) - 3)*arctan(e^(2*x)/(sqrt(3) + 2)) - 1/36*((2*sqrt(3) + 3)*e^(4*x) + 2*sqrt(3) + 3)*arctan(-e^(2*x)/(sqrt(3) - 2)) - 1/12*sqrt(3)*log((sqrt(3) + 2)^2 + e^(4*x)) + 1/12*sqrt(3)*log((sqrt(3) - 2)^2 + e^(4*x)) + 1/12*sqrt(2)*log(-(2*sqrt(2) - e^(2*x) - 3)/(2*sqrt(2) + e^(2*x) + 3))

Mupad [B] (verification not implemented)

Time = 2.67 (sec) , antiderivative size = 337, normalized size of antiderivative = 4.06

$$\int \frac{1}{1 + \cosh^6(x)} dx$$

$$= \frac{\sqrt{3} \ln \left((6177144285775790080 e^{2x} - 2167269359741829120 \sqrt{3} + 3566375915854233600 \sqrt{3} e^{2x} - 3753820658157486080) \right)}{14009449395540459520 e^{2x} + 955607545932677120 \sqrt{3} + 8088359377641144320 \sqrt{3} e^{2x} + 1655160823988879360} - \frac{\sqrt{3} \ln \left((6177144285775790080 e^{2x} + 2167269359741829120 \sqrt{3} - 3566375915854233600 \sqrt{3} e^{2x} - 3753820658157486080) \right)}{14009449395540459520 e^{2x} + 955607545932677120 \sqrt{3} + 8088359377641144320 \sqrt{3} e^{2x} + 1655160823988879360} - \frac{\pi \operatorname{sign} \left(x - \frac{\ln \left(\frac{24639 \sqrt{3} + 42676}{40545 \sqrt{3} + 70226} \right)}{2} \right)}{6} + \frac{\pi \operatorname{sign} (6177144285775790080 e^{2x} - 2167269359741829120 \sqrt{3} + 3566375915854233600 \sqrt{3} e^{2x} - 3753820658157486080)}{6} - \frac{\sqrt{2} \ln (2144322552070144000 \sqrt{2} - 17674880313941032960 e^{2x} + 12498027726650736640 \sqrt{2} e^{2x} - 3014400000000000)}{12} + \frac{\sqrt{2} \ln (17674880313941032960 e^{2x} + 2144322552070144000 \sqrt{2} + 12498027726650736640 \sqrt{2} e^{2x} + 3014400000000000)}{12} + \frac{\ln (e^{2x} (-14009449395540459520 - 6177144285775790080i) + \sqrt{3} (955607545932677120 - 2167269359741829120i))}{12} + \frac{\ln (e^{2x} (-14009449395540459520 + 6177144285775790080i) + \sqrt{3} (955607545932677120 + 2167269359741829120i))}{12}$$

[In] int(1/(cosh(x)^6 + 1),x)

[Out] (log(3^(1/2)*(955607545932677120 - 2167269359741829120i) - exp(2*x)*(14009449395540459520 + 6177144285775790080i) + 3^(1/2)*exp(2*x)*(8088359377641144320 + 3566375915854233600i) - (1655160823988879360 - 3753820658157486080i))*1i)/12 - (log(3^(1/2)*(955607545932677120 + 2167269359741829120i) - exp(2*x)*(14009449395540459520 - 6177144285775790080i) + 3^(1/2)*exp(2*x)*(8088359377641144320 - 3566375915854233600i) - (1655160823988879360 + 3753820658157486080i))*1i)/12 - atan((6177144285775790080*exp(2*x) - 2167269359741829120

$$\begin{aligned}
& 0 \cdot 3^{1/2} + 3566375915854233600 \cdot 3^{1/2} \cdot \exp(2x) - 3753820658157486080) / (14 \\
& 009449395540459520 \cdot \exp(2x) + 955607545932677120 \cdot 3^{1/2} + 8088359377641144 \\
& 320 \cdot 3^{1/2} \cdot \exp(2x) + 1655160823988879360) / 6 + (3^{1/2} \cdot \log((617714428577 \\
& 5790080 \cdot \exp(2x) - 2167269359741829120 \cdot 3^{1/2} + 3566375915854233600 \cdot 3^{1/2} \\
&) \cdot \exp(2x) - 3753820658157486080)^2 + (14009449395540459520 \cdot \exp(2x) + 9556 \\
& 07545932677120 \cdot 3^{1/2} + 8088359377641144320 \cdot 3^{1/2} \cdot \exp(2x) + 16551608239 \\
& 88879360)^2) / 12 - (3^{1/2} \cdot \log((6177144285775790080 \cdot \exp(2x) + 21672693597 \\
& 41829120 \cdot 3^{1/2} - 3566375915854233600 \cdot 3^{1/2} \cdot \exp(2x) - 37538206581574860 \\
& 80)^2 + (14009449395540459520 \cdot \exp(2x) - 955607545932677120 \cdot 3^{1/2} - 80883 \\
& 59377641144320 \cdot 3^{1/2} \cdot \exp(2x) + 1655160823988879360)^2) / 12 - (\pi \cdot \text{sign}(x \\
& - \log((24639 \cdot 3^{1/2} + 42676) / (40545 \cdot 3^{1/2} + 70226)) / 2)) / 6 + (\pi \cdot \text{sign}(617 \\
& 7144285775790080 \cdot \exp(2x) - 2167269359741829120 \cdot 3^{1/2} + 35663759158542336 \\
& 00 \cdot 3^{1/2} \cdot \exp(2x) - 3753820658157486080) / 6 - (2^{1/2} \cdot \log(21443225520701 \\
& 44000 \cdot 2^{1/2} - 17674880313941032960 \cdot \exp(2x) + 12498027726650736640 \cdot 2^{1/2} \\
&) \cdot \exp(2x) - 3032530035220152320) / 12 + (2^{1/2} \cdot \log(17674880313941032960 \cdot e \\
& xp(2x) + 2144322552070144000 \cdot 2^{1/2} + 12498027726650736640 \cdot 2^{1/2} \cdot \exp(2x) \\
&) + 3032530035220152320) / 12
\end{aligned}$$

3.72 $\int \frac{1}{1+\cosh^8(x)} dx$

Optimal result	469
Rubi [A] (verified)	469
Mathematica [C] (verified)	471
Maple [C] (verified)	471
Fricas [B] (verification not implemented)	472
Sympy [F(-1)]	474
Maxima [F]	475
Giac [A] (verification not implemented)	475
Mupad [F(-1)]	475

Optimal result

Integrand size = 8, antiderivative size = 129

$$\int \frac{1}{1+\cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-\sqrt[4]{-1}}}\right)}{4\sqrt{1-\sqrt[4]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+\sqrt[4]{-1}}}\right)}{4\sqrt{1+\sqrt[4]{-1}}} \\ + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-(-1)^{3/4}}}\right)}{4\sqrt{1-(-1)^{3/4}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+(-1)^{3/4}}}\right)}{4\sqrt{1+(-1)^{3/4}}}$$

[Out] 1/4*arctanh(tanh(x)/(1-(-1)^(1/4))^(1/2))/(1-(-1)^(1/4))^(1/2)+1/4*arctanh(tanh(x)/(1+(-1)^(1/4))^(1/2))/(1+(-1)^(1/4))^(1/2)+1/4*arctanh(tanh(x)/(1-(-1)^(3/4))^(1/2))/(1-(-1)^(3/4))^(1/2)+1/4*arctanh(tanh(x)/(1+(-1)^(3/4))^(1/2))/(1+(-1)^(3/4))^(1/2)

Rubi [A] (verified)

Time = 0.15 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3290, 3260, 212}

$$\int \frac{1}{1+\cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-\sqrt[4]{-1}}}\right)}{4\sqrt{1-\sqrt[4]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+\sqrt[4]{-1}}}\right)}{4\sqrt{1+\sqrt[4]{-1}}} \\ + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-(-1)^{3/4}}}\right)}{4\sqrt{1-(-1)^{3/4}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+(-1)^{3/4}}}\right)}{4\sqrt{1+(-1)^{3/4}}}$$

[In] Int[(1 + Cosh[x]^8)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[1 - (-1)^(1/4)]]/(4*Sqrt[1 - (-1)^(1/4)]) + ArcTanh[Tanh[x]/Sqrt[1 + (-1)^(1/4)]]/(4*Sqrt[1 + (-1)^(1/4)]) + ArcTanh[Tanh[x]/Sqrt[1 - (-1)^(3/4)]]/(4*Sqrt[1 - (-1)^(3/4)]) + ArcTanh[Tanh[x]/Sqrt[1 + (-1)^(3/4)]]/(4*Sqrt[1 + (-1)^(3/4)])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2]), x], {k, 1, n/2}], x] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} \int \frac{1}{1 - \sqrt[4]{-1} \cosh^2(x)} dx + \frac{1}{4} \int \frac{1}{1 + \sqrt[4]{-1} \cosh^2(x)} dx \\
 &+ \frac{1}{4} \int \frac{1}{1 - (-1)^{3/4} \cosh^2(x)} dx + \frac{1}{4} \int \frac{1}{1 + (-1)^{3/4} \cosh^2(x)} dx \\
 &= \frac{1}{4} \text{Subst} \left(\int \frac{1}{1 - (1 - \sqrt[4]{-1}) x^2} dx, x, \coth(x) \right) \\
 &+ \frac{1}{4} \text{Subst} \left(\int \frac{1}{1 - (1 + \sqrt[4]{-1}) x^2} dx, x, \coth(x) \right) \\
 &+ \frac{1}{4} \text{Subst} \left(\int \frac{1}{1 - (1 - (-1)^{3/4}) x^2} dx, x, \coth(x) \right) \\
 &+ \frac{1}{4} \text{Subst} \left(\int \frac{1}{1 - (1 + (-1)^{3/4}) x^2} dx, x, \coth(x) \right)
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-\sqrt[4]{-1}}}\right)}{4\sqrt{1-\sqrt[4]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+\sqrt[4]{-1}}}\right)}{4\sqrt{1+\sqrt[4]{-1}}} \\
&\quad + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-(-1)^{3/4}}}\right)}{4\sqrt{1-(-1)^{3/4}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+(-1)^{3/4}}}\right)}{4\sqrt{1+(-1)^{3/4}}}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.02 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.98

$$\begin{aligned}
&\int \frac{1}{1 + \cosh^8(x)} dx \\
&= 16\operatorname{RootSum}\left[1 + 8\#1 + 28\#1^2 + 56\#1^3 + 326\#1^4 + 56\#1^5 + 28\#1^6 + 8\#1^7\right. \\
&\quad \left. + \#1^8 \&, \frac{x\#1^3 + \log(-\cosh(x) - \sinh(x) + \cosh(x)\#1 - \sinh(x)\#1)\#1^3}{1 + 7\#1 + 21\#1^2 + 163\#1^3 + 35\#1^4 + 21\#1^5 + 7\#1^6 + \#1^7} \&\right]
\end{aligned}$$

[In] Integrate[(1 + Cosh[x]^8)^(-1), x]

[Out] 16*RootSum[1 + 8*#1 + 28*#1^2 + 56*#1^3 + 326*#1^4 + 56*#1^5 + 28*#1^6 + 8*#1^7 + #1^8 & , (x*#1^3 + Log[-Cosh[x] - Sinh[x] + Cosh[x]*#1 - Sinh[x]*#1]*#1^3)/(1 + 7*#1 + 21*#1^2 + 163*#1^3 + 35*#1^4 + 21*#1^5 + 7*#1^6 + #1^7) &]

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.83 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.36

method	result
default	$\frac{\sum_{-R=\operatorname{RootOf}(2Z^8-4Z^6+6Z^4-4Z^2+1)} -R \ln\left(2 \tanh\left(\frac{x}{2}\right) - R + \tanh\left(\frac{x}{2}\right)^2 + 1\right)}{8}$
risch	$\sum_{-R=\operatorname{RootOf}(33554432Z^8-1048576Z^6+24576Z^4-256Z^2+1)} -R \ln\left(-8388608R^7 + 1048576R^6 + 131072R^5 - 1048576R^4 + 1048576R^3 - 1048576R^2 + 1048576R - 1048576\right)$

[In] int(1/(1+cosh(x)^8), x, method=_RETURNVERBOSE)

[Out] 1/8*sum(_R*ln(2*tanh(1/2*x)*_R+tanh(1/2*x)^2+1), _R=RootOf(2*_Z^8-4*_Z^6+6*_Z^4-4*_Z^2+1))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 737 vs. $2(89) = 178$.

Time = 0.28 (sec) , antiderivative size = 737, normalized size of antiderivative = 5.71

$$\begin{aligned}
\int \frac{1}{1 + \cosh^8(x)} dx = & -\frac{1}{16} \sqrt{2} \sqrt{\sqrt{2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right. \\
& \left. + \sinh(x)^2 + \sqrt{2\sqrt{2}-3}(\sqrt{2}+2) \right. \\
& \left. + \left(\sqrt{2\sqrt{2}-3}(\sqrt{2}+1) + \sqrt{2}+1 \right) \sqrt{\sqrt{2\sqrt{2}-3}+1 + \sqrt{2}+1} \right) \\
& + \frac{1}{16} \sqrt{2} \sqrt{\sqrt{2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right. \\
& \left. + \sinh(x)^2 + \sqrt{2\sqrt{2}-3}(\sqrt{2}+2) \right. \\
& \left. - \left(\sqrt{2\sqrt{2}-3}(\sqrt{2}+1) + \sqrt{2}+1 \right) \sqrt{\sqrt{2\sqrt{2}-3}+1 + \sqrt{2}+1} \right) \\
& + \frac{1}{16} \sqrt{2} \sqrt{-\sqrt{2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right. \\
& \left. + \sinh(x)^2 - \sqrt{2\sqrt{2}-3}(\sqrt{2}+2) \right. \\
& \left. + \left(\sqrt{2\sqrt{2}-3}(\sqrt{2}+1) - \sqrt{2}-1 \right) \sqrt{-\sqrt{2\sqrt{2}-3}+1 + \sqrt{2}+1} \right) \\
& - \frac{1}{16} \sqrt{2} \sqrt{-\sqrt{2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right. \\
& \left. + \sinh(x)^2 - \sqrt{2\sqrt{2}-3}(\sqrt{2}+2) \right. \\
& \left. - \left(\sqrt{2\sqrt{2}-3}(\sqrt{2}+1) - \sqrt{2}-1 \right) \sqrt{-\sqrt{2\sqrt{2}-3}+1 + \sqrt{2}+1} \right) \\
& + \frac{1}{16} \sqrt{2} \sqrt{-\sqrt{-2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right. \\
& \left. + \sinh(x)^2 + (\sqrt{2}-2) \sqrt{-2\sqrt{2}-3} \right. \\
& \left. + \left((\sqrt{2}-1) \sqrt{-2\sqrt{2}-3} - \sqrt{2}+1 \right) \sqrt{-\sqrt{-2\sqrt{2}-3}+1 - \sqrt{2}} \right. \\
& \left. + 1 \right) - \frac{1}{16} \sqrt{2} \sqrt{-\sqrt{-2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 \right. \\
& \left. + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + (\sqrt{2}-2) \sqrt{-2\sqrt{2}-3} \right. \\
& \left. - \left((\sqrt{2}-1) \sqrt{-2\sqrt{2}-3} - \sqrt{2}+1 \right) \sqrt{-\sqrt{-2\sqrt{2}-3}+1 - \sqrt{2}} \right. \\
& \left. + 1 \right) - \frac{1}{16} \sqrt{2} \sqrt{\sqrt{-2\sqrt{2}-3}+1} \log \left(\cosh(x)^2 + 2 \cosh(x) \sinh(x) \right.
\end{aligned}$$

```
[In] integrate(1/(1+cosh(x)^8),x, algorithm="fricas")
```

```
[Out] -1/16*sqrt(2)*sqrt(sqrt(2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(2*sqrt(2) - 3)*(sqrt(2) + 2) + (sqrt(2*sqrt(2) - 3)*(sqrt(2) + 1) + sqrt(2) + 1)*sqrt(sqrt(2*sqrt(2) - 3) + 1) + sqrt(2) + 1) + 1/16*sqrt(2)*sqrt(sqrt(2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + sqrt(2*sqrt(2) - 3)*(sqrt(2) + 2) - (sqrt(2*sqrt(2) - 3)*(sqrt(2) + 1) + sqrt(2) + 1)*sqrt(sqrt(2*sqrt(2) - 3) + 1) + sqrt(2) + 1) + 1/16*sqrt(2)*sqrt(-sqrt(2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(2*sqrt(2) - 3)*(sqrt(2) + 2) + (sqrt(2*sqrt(2) - 3)*(sqrt(2) + 1) - sqrt(2) - 1)*sqrt(-sqrt(2*sqrt(2) - 3) + 1) + sqrt(2) + 1) - 1/16*sqrt(2)*sqrt(-sqrt(2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - sqrt(2*sqrt(2) - 3)*(sqrt(2) + 2) - (sqrt(2*sqrt(2) - 3)*(sqrt(2) + 1) - sqrt(2) - 1)*sqrt(-sqrt(2*sqrt(2) - 3) + 1) + sqrt(2) + 1) + 1/16*sqrt(2)*sqrt(-sqrt(-2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + (sqrt(2) - 2)*sqrt(-2*sqrt(2) - 3) + ((sqrt(2) - 1)*sqrt(-2*sqrt(2) - 3) - sqrt(2) + 1)*sqrt(-sqrt(-2*sqrt(2) - 3) + 1) - sqrt(2) + 1) - 1/16*sqrt(2)*sqrt(-sqrt(-2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + (sqrt(2) - 2)*sqrt(-2*sqrt(2) - 3) - ((sqrt(2) - 1)*sqrt(-2*sqrt(2) - 3) - sqrt(2) + 1)*sqrt(-sqrt(-2*sqrt(2) - 3) + 1) - sqrt(2) + 1) - 1/16*sqrt(2)*sqrt(sqrt(-2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - (sqrt(2) - 2)*sqrt(-2*sqrt(2) - 3) + ((sqrt(2) - 1)*sqrt(-2*sqrt(2) - 3) + sqrt(2) - 1)*sqrt(sqrt(-2*sqrt(2) - 3) + 1) - sqrt(2) + 1) + 1/16*sqrt(2)*sqrt(sqrt(-2*sqrt(2) - 3) + 1)*log(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 - (sqrt(2) - 2)*sqrt(-2*sqrt(2) - 3) - ((sqrt(2) - 1)*sqrt(-2*sqrt(2) - 3) + sqrt(2) - 1)*sqrt(sqrt(-2*sqrt(2) - 3) + 1) - sqrt(2) + 1)
```

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{1 + \cosh^8(x)} dx = \text{Timed out}$$

```
[In] integrate(1/(1+cosh(x)**8),x)
```

```
[Out] Timed out
```

Maxima [F]

$$\int \frac{1}{1 + \cosh^8(x)} dx = \int \frac{1}{\cosh(x)^8 + 1} dx$$

[In] integrate(1/(1+cosh(x)^8),x, algorithm="maxima")

[Out] integrate(1/(cosh(x)^8 + 1), x)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 1, normalized size of antiderivative = 0.01

$$\int \frac{1}{1 + \cosh^8(x)} dx = 0$$

[In] integrate(1/(1+cosh(x)^8),x, algorithm="giac")

[Out] 0

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{1 + \cosh^8(x)} dx = \text{Hanged}$$

[In] int(1/(cosh(x)^8 + 1),x)

[Out] \text{Hanged}

3.73 $\int \frac{1}{1-\cosh^5(x)} dx$

Optimal result	476
Rubi [A] (verified)	476
Mathematica [C] (verified)	479
Maple [C] (verified)	479
Fricas [B] (verification not implemented)	480
Sympy [F(-1)]	481
Maxima [F]	481
Giac [F]	481
Mupad [F(-1)]	481

Optimal result

Integrand size = 10, antiderivative size = 205

$$\int \frac{1}{1-\cosh^5(x)} dx = -\frac{2 \arctan\left(\frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{\frac{1-(-1)^{2/5}}{1+(-1)^{2/5}}}}\right)}{5\sqrt{-1+(-1)^{4/5}}} + \frac{2 \arctan\left(\sqrt{\frac{1+(-1)^{4/5}}{1-(-1)^{4/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{-1-(-1)^{3/5}}}$$

$$+ \frac{2 \operatorname{arctanh}\left(\sqrt{\frac{1-\sqrt[5]{-1}}{1+\sqrt[5]{-1}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1-(-1)^{2/5}}}$$

$$+ \frac{2 \operatorname{arctanh}\left(\sqrt{\frac{1-(-1)^{3/5}}{1+(-1)^{3/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1+\sqrt[5]{-1}}} - \frac{\sinh(x)}{5(1-\cosh(x))}$$

```
[Out] -1/5*sinh(x)/(1-cosh(x))+2/5*arctanh(((1-(-1)^(3/5))/(1+(-1)^(3/5)))^(1/2)*
tanh(1/2*x))/(1+(-1)^(1/5))^(1/2)+2/5*arctanh(((1-(-1)^(1/5))/(1+(-1)^(1/5)
))^(1/2)*tanh(1/2*x))/(1-(-1)^(2/5))^(1/2)+2/5*arctan(((1-(-1)^(4/5))/(1-(-1)^(4/5)))^(1/2)*tanh(1/2*x)/(-1-(-1)^(3/5))^(1/2)-2/5*arctan(tanh(1/2*x)
/((-1+(-1)^(2/5))/(1+(-1)^(2/5)))^(1/2)/(-1+(-1)^(4/5))^(1/2)
```

Rubi [A] (verified)

Time = 0.36 (sec) , antiderivative size = 205, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used

= {3292, 2727, 2738, 214, 211}

$$\int \frac{1}{1 - \cosh^5(x)} dx = -\frac{2 \arctan\left(\frac{\tanh\left(\frac{x}{2}\right)}{\sqrt{\frac{1 - (-1)^{2/5}}{1 + (-1)^{2/5}}}}\right)}{5\sqrt{(-1)^{4/5} - 1}} + \frac{2 \arctan\left(\sqrt{\frac{1 + (-1)^{4/5}}{1 - (-1)^{4/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{-1 - (-1)^{3/5}}}$$

$$+ \frac{2 \operatorname{arctanh}\left(\sqrt{\frac{1 - \sqrt[5]{-1}}{1 + \sqrt[5]{-1}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1 - (-1)^{2/5}}}$$

$$+ \frac{2 \operatorname{arctanh}\left(\sqrt{\frac{1 - (-1)^{3/5}}{1 + (-1)^{3/5}}} \tanh\left(\frac{x}{2}\right)\right)}{5\sqrt{1 + \sqrt[5]{-1}}} - \frac{\sinh(x)}{5(1 - \cosh(x))}$$

[In] Int[(1 - Cosh[x]^5)^(-1), x]

[Out] (-2*ArcTan[Tanh[x/2]/Sqrt[-((1 - (-1)^(2/5))/(1 + (-1)^(2/5))]])/(5*Sqrt[-1 + (-1)^(4/5)]) + (2*ArcTan[Sqrt[-((1 + (-1)^(4/5))/(1 - (-1)^(4/5)))]*Tanh[x/2]])/(5*Sqrt[-1 - (-1)^(3/5)]) + (2*ArcTanh[Sqrt[(1 - (-1)^(1/5))/(1 + (-1)^(1/5))]*Tanh[x/2]])/(5*Sqrt[1 - (-1)^(2/5)]) + (2*ArcTanh[Sqrt[(1 - (-1)^(3/5))/(1 + (-1)^(3/5))]*Tanh[x/2]])/(5*Sqrt[1 + (-1)^(1/5)]) - Sinh[x]/(5*(1 - Cosh[x]))

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 2727

Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] := Simp[-Cos[c + d*x]/(d*(b + a*Sin[c + d*x])), x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 - b^2, 0]

Rule 2738

Int[((a_) + (b_)*sin[Pi/2 + (c_) + (d_)*(x_)])^(-1), x_Symbol] := With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[2*(e/d), Subst[Int[1/(a + b + (a - b)*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]

Rule 3292

```
Int[((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :>
Int[ExpandTrig[(a + b*(c*sin[e + f*x])^n)^p, x], x] /; FreeQ[{a, b, c, e, f
, n}, x] && (IGtQ[p, 0] || (EqQ[p, -1] && IntegerQ[n]))
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{5(1 - \cosh(x))} + \frac{1}{5(1 + \sqrt[5]{-1} \cosh(x))} + \frac{1}{5(1 - (-1)^{2/5} \cosh(x))} \right. \\
&\quad \left. + \frac{1}{5(1 + (-1)^{3/5} \cosh(x))} + \frac{1}{5(1 - (-1)^{4/5} \cosh(x))} \right) dx \\
&= \frac{1}{5} \int \frac{1}{1 - \cosh(x)} dx + \frac{1}{5} \int \frac{1}{1 + \sqrt[5]{-1} \cosh(x)} dx + \frac{1}{5} \int \frac{1}{1 - (-1)^{2/5} \cosh(x)} dx \\
&\quad + \frac{1}{5} \int \frac{1}{1 + (-1)^{3/5} \cosh(x)} dx + \frac{1}{5} \int \frac{1}{1 - (-1)^{4/5} \cosh(x)} dx \\
&= -\frac{\sinh(x)}{5(1 - \cosh(x))} + \frac{2}{5} \text{Subst} \left(\int \frac{1}{1 + \sqrt[5]{-1} - (1 - \sqrt[5]{-1}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
&\quad + \frac{2}{5} \text{Subst} \left(\int \frac{1}{1 - (-1)^{2/5} - (1 + (-1)^{2/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
&\quad + \frac{2}{5} \text{Subst} \left(\int \frac{1}{1 + (-1)^{3/5} - (1 - (-1)^{3/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) + \frac{2}{5} \text{Subst} \left(\int \frac{1}{1 - (-1)^{4/5} - (1 + (-1)^{4/5}) x^2} dx, x, \tanh \left(\frac{x}{2} \right) \right) \\
&= -\frac{2 \arctan \left(\frac{\tanh \left(\frac{x}{2} \right)}{\sqrt{\frac{1 - (-1)^{2/5}}{1 + (-1)^{2/5}}}} \right)}{5\sqrt{-1} + (-1)^{4/5}} + \frac{2 \arctan \left(\sqrt{\frac{1 + (-1)^{4/5}}{1 - (-1)^{4/5}}} \tanh \left(\frac{x}{2} \right) \right)}{5\sqrt{-1} - (-1)^{3/5}} \\
&\quad + \frac{2 \operatorname{arctanh} \left(\sqrt{\frac{1 - \sqrt[5]{-1}}{1 + \sqrt[5]{-1}}} \tanh \left(\frac{x}{2} \right) \right)}{5\sqrt{1 - (-1)^{2/5}}} \\
&\quad + \frac{2 \operatorname{arctanh} \left(\sqrt{\frac{1 - (-1)^{3/5}}{1 + (-1)^{3/5}}} \tanh \left(\frac{x}{2} \right) \right)}{5\sqrt{1 + \sqrt[5]{-1}}} - \frac{\sinh(x)}{5(1 - \cosh(x))}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 5.03 (sec) , antiderivative size = 445, normalized size of antiderivative = 2.17

$$\int \frac{1}{1 - \cosh^5(x)} dx$$

$$= \frac{1}{5} \coth\left(\frac{x}{2}\right) + \frac{1}{10} \text{RootSum} \left[1 + 2\#1 + 8\#1^2 + 14\#1^3 + 30\#1^4 + 14\#1^5 + 8\#1^6 + 2\#1^7 \right. \\ \left. + \#1^8 \&, \frac{x + 2 \log\left(-\cosh\left(\frac{x}{2}\right) - \sinh\left(\frac{x}{2}\right) + \cosh\left(\frac{x}{2}\right)\#1 - \sinh\left(\frac{x}{2}\right)\#1\right) + 4x\#1 + 8 \log\left(-\cosh\left(\frac{x}{2}\right) - \sinh\left(\frac{x}{2}\right)\right)}{\#1^8} \right]$$

[In] Integrate[(1 - Cosh[x]^5)^(-1),x]

[Out] Coth[x/2]/5 + RootSum[1 + 2*#1 + 8*#1^2 + 14*#1^3 + 30*#1^4 + 14*#1^5 + 8*#1^6 + 2*#1^7 + #1^8 & , (x + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1] + 4*x*#1 + 8*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1 + 15*x*#1^2 + 30*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^2 + 40*x*#1^3 + 80*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^3 + 15*x*#1^4 + 30*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^4 + 4*x*#1^5 + 8*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^5 + x*#1^6 + 2*Log[-Cosh[x/2] - Sinh[x/2] + Cosh[x/2]*#1 - Sinh[x/2]*#1]*#1^6)/(1 + 8*#1 + 21*#1^2 + 60*#1^3 + 35*#1^4 + 24*#1^5 + 7*#1^6 + 4*#1^7) &]/10

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.13 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.31

method	result
default	$\left(\frac{\sum_{R=\text{RootOf}(_Z^8+10_Z^4+5)} \left(\frac{(-_R^6+5_R^4-5_R^2+5) \ln(\tanh(\frac{x}{2})-R)}{_R^7+5_R^3} \right)}{10} \right) + \frac{1}{5 \tanh(\frac{x}{2})}$
risch	$\frac{2}{5(e^x-1)} + \left(\sum_{R=\text{RootOf}(1953125_Z^8-156250_Z^6+6250_Z^4-125_Z^2+1)} _R \ln(-2343750_R^7 + 234375_R^6) \right)$

[In] int(1/(1-cosh(x)^5),x,method=_RETURNVERBOSE)

[Out] 1/10*sum((-_R^6+5*_R^4-5*_R^2+5)/(_R^7+5*_R^3)*ln(tanh(1/2*x)-_R),_R=RootOf(_Z^8+10*_Z^4+5))+1/5/tanh(1/2*x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 852 vs. 2(137) = 274.

Time = 0.30 (sec) , antiderivative size = 852, normalized size of antiderivative = 4.16

$$\int \frac{1}{1 - \cosh^5(x)} dx = \text{Too large to display}$$

[In] integrate(1/(1-cosh(x)^5),x, algorithm="fricas")

[Out] -1/50*((sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*log(sqrt(2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*(3*sqrt(5) + 5)*sqrt(2*sqrt(5) - 5) + 5*sqrt(2*sqrt(5) - 5)*(sqrt(5) + 3) - 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) - (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*log(-sqrt(2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*(3*sqrt(5) + 5)*sqrt(2*sqrt(5) - 5) + 5*sqrt(2*sqrt(5) - 5)*(sqrt(5) + 3) - 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) - (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(-2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*log(sqrt(-2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*(3*sqrt(5) + 5)*sqrt(2*sqrt(5) - 5) - 5*sqrt(2*sqrt(5) - 5)*(sqrt(5) + 3) - 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) + (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(-2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*log(-sqrt(-2*sqrt(5)*sqrt(2*sqrt(5) - 5) + 10)*(3*sqrt(5) + 5)*sqrt(2*sqrt(5) - 5) - 5*sqrt(2*sqrt(5) - 5)*(sqrt(5) + 3) - 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) - (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*log(sqrt(2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*(3*sqrt(5) - 5)*sqrt(-2*sqrt(5) - 5) + 5*(sqrt(5) - 3)*sqrt(-2*sqrt(5) - 5) + 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) + (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*log(-sqrt(2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*(3*sqrt(5) - 5)*sqrt(-2*sqrt(5) - 5) + 5*(sqrt(5) - 3)*sqrt(-2*sqrt(5) - 5) + 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) + (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(-2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*log(sqrt(-2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*(3*sqrt(5) - 5)*sqrt(-2*sqrt(5) - 5) - 5*(sqrt(5) - 3)*sqrt(-2*sqrt(5) - 5) + 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) - (sqrt(5)*cosh(x) + sqrt(5)*sinh(x) - sqrt(5))*sqrt(-2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*log(-sqrt(-2*sqrt(5)*sqrt(-2*sqrt(5) - 5) + 10)*(3*sqrt(5) - 5)*sqrt(-2*sqrt(5) - 5) - 5*(sqrt(5) - 3)*sqrt(-2*sqrt(5) - 5) + 5*sqrt(5) + 20*cosh(x) + 20*sinh(x) + 5) - 20)/(cosh(x) + sinh(x) - 1)

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{1 - \cosh^5(x)} dx = \text{Timed out}$$

[In] integrate(1/(1-cosh(x)**5),x)

[Out] Timed out

Maxima [F]

$$\int \frac{1}{1 - \cosh^5(x)} dx = \int -\frac{1}{\cosh(x)^5 - 1} dx$$

[In] integrate(1/(1-cosh(x)^5),x, algorithm="maxima")

[Out] 2/5/(e^x - 1) + integrate(2/5*(e^(7*x) + 4*e^(6*x) + 15*e^(5*x) + 40*e^(4*x) + 15*e^(3*x) + 4*e^(2*x) + e^x)/(e^(8*x) + 2*e^(7*x) + 8*e^(6*x) + 14*e^(5*x) + 30*e^(4*x) + 14*e^(3*x) + 8*e^(2*x) + 2*e^x + 1), x)

Giac [F]

$$\int \frac{1}{1 - \cosh^5(x)} dx = \int -\frac{1}{\cosh(x)^5 - 1} dx$$

[In] integrate(1/(1-cosh(x)^5),x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{1 - \cosh^5(x)} dx = \text{Hanged}$$

[In] int(-1/(cosh(x)^5 - 1),x)

[Out] \text{Hanged}

3.74 $\int \frac{1}{1 - \cosh^6(x)} dx$

Optimal result	482
Rubi [A] (verified)	482
Mathematica [C] (verified)	484
Maple [C] (verified)	484
Fricas [C] (verification not implemented)	485
Sympy [B] (verification not implemented)	485
Maxima [F]	487
Giac [A] (verification not implemented)	487
Mupad [B] (verification not implemented)	488

Optimal result

Integrand size = 10, antiderivative size = 71

$$\int \frac{1}{1 - \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 + \sqrt[3]{-1}}}\right)}{3\sqrt{1 + \sqrt[3]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 - (-1)^{2/3}}}\right)}{3\sqrt{1 - (-1)^{2/3}}} + \frac{\operatorname{coth}(x)}{3}$$

[Out] $\frac{1}{3} \operatorname{coth}(x) + \frac{1}{3} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 + (-1)^{1/3}}}\right) / \sqrt{1 + (-1)^{1/3}} + \frac{1}{3} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 - (-1)^{2/3}}}\right) / \sqrt{1 - (-1)^{2/3}}$

Rubi [A] (verified)

Time = 0.10 (sec), antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3290, 3260, 212, 3254, 3852, 8}

$$\int \frac{1}{1 - \cosh^6(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 + \sqrt[3]{-1}}}\right)}{3\sqrt{1 + \sqrt[3]{-1}}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 - (-1)^{2/3}}}\right)}{3\sqrt{1 - (-1)^{2/3}}} + \frac{\operatorname{coth}(x)}{3}$$

[In] `Int[(1 - Cosh[x]^6)^(-1), x]`

[Out] `ArcTanh[Tanh[x]/Sqrt[1 + (-1)^(1/3)]]/(3*Sqrt[1 + (-1)^(1/3)]) + ArcTanh[Tanh[x]/Sqrt[1 - (-1)^(2/3)]]/(3*Sqrt[1 - (-1)^(2/3)]) + Coth[x]/3`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 3254

Int[(u_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := Dist[a^p, Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]

Rule 3260

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^(n_))^(p_), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rule 3852

Int[csc[(c_) + (d_)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{3} \int \frac{1}{1 - \cosh^2(x)} dx + \frac{1}{3} \int \frac{1}{1 + \sqrt[3]{-1} \cosh^2(x)} dx + \frac{1}{3} \int \frac{1}{1 - (-1)^{2/3} \cosh^2(x)} dx \\
 &= -\left(\frac{1}{3} \int \text{csch}^2(x) dx\right) + \frac{1}{3} \text{Subst}\left(\int \frac{1}{1 - (1 + \sqrt[3]{-1}) x^2} dx, x, \text{coth}(x)\right) \\
 &\quad + \frac{1}{3} \text{Subst}\left(\int \frac{1}{1 - (1 - (-1)^{2/3}) x^2} dx, x, \text{coth}(x)\right) \\
 &= \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 + \sqrt[3]{-1}}}\right)}{3\sqrt{1 + \sqrt[3]{-1}}} + \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 - (-1)^{2/3}}}\right)}{3\sqrt{1 - (-1)^{2/3}}} + \frac{1}{3} i \text{Subst}\left(\int 1 dx, x, -i \text{coth}(x)\right) \\
 &= \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 + \sqrt[3]{-1}}}\right)}{3\sqrt{1 + \sqrt[3]{-1}}} + \frac{\text{arctanh}\left(\frac{\tanh(x)}{\sqrt{1 - (-1)^{2/3}}}\right)}{3\sqrt{1 - (-1)^{2/3}}} + \frac{\text{coth}(x)}{3}
 \end{aligned}$$

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 1.24 (sec) , antiderivative size = 111, normalized size of antiderivative = 1.56

$$\int \frac{1}{1 - \cosh^6(x)} dx = \frac{(15 + 8 \cosh(2x) + \cosh(4x)) \sinh(x) \left(-6 \cosh(x) + \sqrt[4]{-3} \left((3i + \sqrt{3}) \arctan \left(\frac{(-1)^{3/4} (-i + \sqrt{3}) \tanh(x)}{2 \sqrt[4]{3}} \right) + \dots \right)}{144 (-1 + \cosh^6(x))} + \dots$$

[In] Integrate[(1 - Cosh[x]^6)^(-1), x]

[Out] -1/144*((15 + 8*Cosh[2*x] + Cosh[4*x])*Sinh[x]*(-6*Cosh[x] + (-3)^(1/4)*((3 *I + Sqrt[3])*ArcTan[((-1)^(3/4)*(-I + Sqrt[3])*Tanh[x])/(2*3^(1/4))]) + (3 + I*Sqrt[3])*ArcTan[((-1/3)^(1/4)*(I + Sqrt[3])*Tanh[x])/2])*Sinh[x]))/(-1 + Cosh[x]^6)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.18 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.66

method	result
risch	$\frac{2}{3(e^{2x}-1)} + \left(\sum_{R=\text{RootOf}(3888_Z^4-108_Z^2+1)} -R \ln(-1296_R^3 + 216_R^2 + e^{2x} - 1) \right)$
default	$\frac{\tanh(\frac{x}{2})}{6} + \frac{3^{\frac{3}{4}} \sqrt{2} \left(\ln \left(\frac{\tanh(\frac{x}{2})^2 + \frac{3^{\frac{3}{4}} \tanh(\frac{x}{2}) \sqrt{2}}{3} + \frac{\sqrt{3}}{3}}{\tanh(\frac{x}{2})^2 - \frac{3^{\frac{3}{4}} \tanh(\frac{x}{2}) \sqrt{2}}{3} + \frac{\sqrt{3}}{3}} \right) + 2 \arctan(\sqrt{2} 3^{\frac{1}{4}} \tanh(\frac{x}{2}) + 1) + 2 \arctan(\sqrt{2} 3^{\frac{1}{4}} \tanh(\frac{x}{2}) - 1) \right)}{72} - \dots$

[In] int(1/(1-cosh(x)^6), x, method=_RETURNVERBOSE)

[Out] 2/3/(exp(2*x)-1)+sum(_R*ln(-1296*_R^3+216*_R^2+exp(2*x)-1), _R=RootOf(3888*_Z^4-108*_Z^2+1))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 358, normalized size of antiderivative = 5.04

$$\int \frac{1}{1 - \cosh^6(x)} dx = \frac{(\sqrt{6} \cosh(x)^2 + 2\sqrt{6} \cosh(x) \sinh(x) + \sqrt{6} \sinh(x)^2 - \sqrt{6}) \sqrt{i\sqrt{3} + 3} \log\left(\sqrt{6}(i\sqrt{3} + 3)\right)^{\frac{3}{2}} + 6 \cosh(x)}{\dots}$$

[In] integrate(1/(1-cosh(x)^6),x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/36*((\sqrt{6}*\cosh(x)^2 + 2*\sqrt{6}*\cosh(x)*\sinh(x) + \sqrt{6}*\sinh(x)^2 - \\ & \sqrt{6})*\sqrt{I*\sqrt{3} + 3}*\log(\sqrt{6}*(I*\sqrt{3} + 3)^{(3/2)} + 6*\cosh(x) \\ & ^2 + 12*\cosh(x)*\sinh(x) + 6*\sinh(x)^2 + 6*I*\sqrt{3} + 12) - (\sqrt{6}*\cosh(x) \\ &)^2 + 2*\sqrt{6}*\cosh(x)*\sinh(x) + \sqrt{6}*\sinh(x)^2 - \sqrt{6})*\sqrt{-I*\sqrt{3} \\ & (3) + 3}*\log(\sqrt{6}*(I*\sqrt{3} - 3)*\sqrt{-I*\sqrt{3} + 3} + 6*\cosh(x)^2 + 1 \\ & 2*\cosh(x)*\sinh(x) + 6*\sinh(x)^2 - 6*I*\sqrt{3} + 12) + (\sqrt{6}*\cosh(x)^2 + \\ & 2*\sqrt{6}*\cosh(x)*\sinh(x) + \sqrt{6}*\sinh(x)^2 - \sqrt{6})*\sqrt{-I*\sqrt{3} + \\ & 3}*\log(\sqrt{6}*(-I*\sqrt{3} + 3)^{(3/2)} + 6*\cosh(x)^2 + 12*\cosh(x)*\sinh(x) + \\ & 6*\sinh(x)^2 - 6*I*\sqrt{3} + 12) - (\sqrt{6}*\cosh(x)^2 + 2*\sqrt{6}*\cosh(x)*\sinh(x) \\ & + \sqrt{6}*\sinh(x)^2 - \sqrt{6})*\sqrt{I*\sqrt{3} + 3}*\log(\sqrt{6}*\sqrt{I \\ & *\sqrt{3} + 3}*(-I*\sqrt{3} - 3) + 6*\cosh(x)^2 + 12*\cosh(x)*\sinh(x) + 6*\sinh(x) \\ & ^2 + 6*I*\sqrt{3} + 12) - 24)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - \\ & 1) \end{aligned}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 632 vs. 2(65) = 130.

Time = 9.15 (sec) , antiderivative size = 632, normalized size of antiderivative = 8.90

$$\begin{aligned}
 \int \frac{1}{1 - \cosh^6(x)} dx = & - \frac{\sqrt{2} \cdot \sqrt[4]{3} \log \left(4 \tanh^2 \left(\frac{x}{2} \right) - 4\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 4\sqrt{3} \right)}{24} \\
 & - \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log \left(4 \tanh^2 \left(\frac{x}{2} \right) - 4\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 4\sqrt{3} \right)}{72} \\
 & + \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log \left(4 \tanh^2 \left(\frac{x}{2} \right) + 4\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 4\sqrt{3} \right)}{72} \\
 & + \frac{\sqrt{2} \cdot \sqrt[4]{3} \log \left(4 \tanh^2 \left(\frac{x}{2} \right) + 4\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 4\sqrt{3} \right)}{24} \\
 & - \frac{\sqrt{2} \cdot \sqrt[4]{3} \log \left(36 \tanh^2 \left(\frac{x}{2} \right) - 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right) + 12\sqrt{3} \right)}{24} \\
 & - \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log \left(36 \tanh^2 \left(\frac{x}{2} \right) - 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right) + 12\sqrt{3} \right)}{72} \\
 & + \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \log \left(36 \tanh^2 \left(\frac{x}{2} \right) + 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right) + 12\sqrt{3} \right)}{72} \\
 & + \frac{\sqrt{2} \cdot \sqrt[4]{3} \log \left(36 \tanh^2 \left(\frac{x}{2} \right) + 12\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right) + 12\sqrt{3} \right)}{24} \\
 & + \frac{\tanh \left(\frac{x}{2} \right)}{6} - \frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan} \left(\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) - 1 \right)}{12} \\
 & + \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan} \left(\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) - 1 \right)}{36} \\
 & - \frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan} \left(\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 1 \right)}{12} \\
 & + \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan} \left(\sqrt{2} \cdot \sqrt[4]{3} \tanh \left(\frac{x}{2} \right) + 1 \right)}{36} \\
 & - \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan} \left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right)}{3} - 1 \right)}{36} \\
 & + \frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan} \left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right)}{3} - 1 \right)}{12} \\
 & - \frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \operatorname{atan} \left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right)}{3} + 1 \right)}{36} \\
 & + \frac{\sqrt{2} \cdot \sqrt[4]{3} \operatorname{atan} \left(\frac{\sqrt{2} \cdot 3^{\frac{3}{4}} \tanh \left(\frac{x}{2} \right)}{3} + 1 \right)}{12} + \frac{1}{6 \tanh \left(\frac{x}{2} \right)}
 \end{aligned}$$

[In] integrate(1/(1-cosh(x)**6),x)

[Out] $-\sqrt{2} \cdot 3^{1/4} \cdot \log(4 \cdot \tanh(x/2))^2 - 4 \cdot \sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 4 \cdot \sqrt{3} / 24 - \sqrt{2} \cdot 3^{3/4} \cdot \log(4 \cdot \tanh(x/2))^2 - 4 \cdot \sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 4 \cdot \sqrt{3} / 72 + \sqrt{2} \cdot 3^{3/4} \cdot \log(4 \cdot \tanh(x/2))^2 + 4 \cdot \sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 4 \cdot \sqrt{3} / 72 + \sqrt{2} \cdot 3^{1/4} \cdot \log(4 \cdot \tanh(x/2))^2 + 4 \cdot \sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 4 \cdot \sqrt{3} / 24 - \sqrt{2} \cdot 3^{1/4} \cdot \log(36 \cdot \tanh(x/2))^2 - 12 \cdot \sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) + 12 \cdot \sqrt{3} / 24 - \sqrt{2} \cdot 3^{3/4} \cdot \log(36 \cdot \tanh(x/2))^2 - 12 \cdot \sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) + 12 \cdot \sqrt{3} / 72 + \sqrt{2} \cdot 3^{3/4} \cdot \log(36 \cdot \tanh(x/2))^2 + 12 \cdot \sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) + 12 \cdot \sqrt{3} / 72 + \sqrt{2} \cdot 3^{1/4} \cdot \log(36 \cdot \tanh(x/2))^2 + 12 \cdot \sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) + 12 \cdot \sqrt{3} / 24 + \tanh(x/2) / 6 - \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) - 1) / 12 + \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) - 1) / 36 - \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 1) / 12 + \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{1/4} \cdot \tanh(x/2) + 1) / 36 - \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 - 1) / 36 + \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 - 1) / 12 - \sqrt{2} \cdot 3^{3/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 + 1) / 36 + \sqrt{2} \cdot 3^{1/4} \cdot \operatorname{atan}(\sqrt{2} \cdot 3^{3/4} \cdot \tanh(x/2) / 3 + 1) / 12 + 1 / (6 \cdot \tanh(x/2))$

Maxima [F]

$$\int \frac{1}{1 - \cosh^6(x)} dx = \int -\frac{1}{\cosh(x)^6 - 1} dx$$

[In] integrate(1/(1-cosh(x)^6),x, algorithm="maxima")

[Out] $2/3 / (e^{2x} - 1) + \operatorname{integrate}(1/3 * (e^{3x} + 4 * e^{2x} + e^x) / (e^{4x} + 2 * e^{3x} + 6 * e^{2x} + 2 * e^x + 1), x) - \operatorname{integrate}(1/3 * (e^{3x} - 4 * e^{2x} + e^x) / (e^{4x} - 2 * e^{3x} + 6 * e^{2x} - 2 * e^x + 1), x)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.14

$$\int \frac{1}{1 - \cosh^6(x)} dx = \frac{2}{3(e^{2x} - 1)}$$

[In] integrate(1/(1-cosh(x)^6),x, algorithm="giac")

[Out] $2/3 / (e^{2x} - 1)$

Mupad [B] (verification not implemented)

Time = 4.60 (sec) , antiderivative size = 329, normalized size of antiderivative = 4.63

$$\begin{aligned}
\int \frac{1}{1 - \cosh^6(x)} dx = & \ln \left(\frac{1061158912 e^{2x}}{27} + \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{2539651072 e^{2x}}{9} \right. \right. \\
& - \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{21515730944 e^{2x}}{9} + \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} (19788726272 e^{2x} + 2864709632) + \frac{3870294016}{9} \right) \\
& \left. \left. + \frac{548405248}{27} \right) + \frac{351797248}{81} \right) \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \\
& + \ln \left(\frac{1061158912 e^{2x}}{27} + \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{2539651072 e^{2x}}{9} \right. \right. \\
& - \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{21515730944 e^{2x}}{9} + \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} (19788726272 e^{2x} + 2864709632) + \frac{3870294016}{9} \right) \\
& \left. \left. + \frac{548405248}{27} \right) + \frac{351797248}{81} \right) \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} \\
& - \ln \left(\frac{1061158912 e^{2x}}{27} - \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{2539651072 e^{2x}}{9} \right. \right. \\
& + \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{21515730944 e^{2x}}{9} - \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} (19788726272 e^{2x} + 2864709632) + \frac{3870294016}{9} \right) \\
& \left. \left. + \frac{548405248}{27} \right) + \frac{351797248}{81} \right) \sqrt{\frac{1}{72} - \frac{\sqrt{3} \operatorname{li}}{216}} \\
& - \ln \left(\frac{1061158912 e^{2x}}{27} - \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{2539651072 e^{2x}}{9} \right. \right. \\
& + \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} \left(\frac{21515730944 e^{2x}}{9} - \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} (19788726272 e^{2x} + 2864709632) + \frac{3870294016}{9} \right) \\
& \left. \left. + \frac{548405248}{27} \right) + \frac{351797248}{81} \right) \sqrt{\frac{1}{72} + \frac{\sqrt{3} \operatorname{li}}{216}} + \frac{2}{3(e^{2x} - 1)}
\end{aligned}$$

`[In] int(-1/(cosh(x)^6 - 1),x)`


```
[Out] log((1061158912*exp(2*x))/27 + (1/72 - (3^(1/2)*1i)/216)^(1/2)*((2539651072
*exp(2*x))/9 - (1/72 - (3^(1/2)*1i)/216)^(1/2)*((21515730944*exp(2*x))/9 +
(1/72 - (3^(1/2)*1i)/216)^(1/2)*(19788726272*exp(2*x) + 2864709632) + 38702
94016/9) + 548405248/27) + 351797248/81)*(1/72 - (3^(1/2)*1i)/216)^(1/2) +
log((1061158912*exp(2*x))/27 + ((3^(1/2)*1i)/216 + 1/72)^(1/2)*((2539651072
*exp(2*x))/9 - ((3^(1/2)*1i)/216 + 1/72)^(1/2)*((21515730944*exp(2*x))/9 +
((3^(1/2)*1i)/216 + 1/72)^(1/2)*(19788726272*exp(2*x) + 2864709632) + 38702
94016/9) + 548405248/27) + 351797248/81)*((3^(1/2)*1i)/216 + 1/72)^(1/2) -
log((1061158912*exp(2*x))/27 - (1/72 - (3^(1/2)*1i)/216)^(1/2)*((2539651072
*exp(2*x))/9 + (1/72 - (3^(1/2)*1i)/216)^(1/2)*((21515730944*exp(2*x))/9 -
(1/72 - (3^(1/2)*1i)/216)^(1/2)*(19788726272*exp(2*x) + 2864709632) + 38702
94016/9) + 548405248/27) + 351797248/81)*(1/72 - (3^(1/2)*1i)/216)^(1/2) -
log((1061158912*exp(2*x))/27 - ((3^(1/2)*1i)/216 + 1/72)^(1/2)*((2539651072
*exp(2*x))/9 + ((3^(1/2)*1i)/216 + 1/72)^(1/2)*((21515730944*exp(2*x))/9 -
((3^(1/2)*1i)/216 + 1/72)^(1/2)*(19788726272*exp(2*x) + 2864709632) + 38702
94016/9) + 548405248/27) + 351797248/81)*((3^(1/2)*1i)/216 + 1/72)^(1/2) +
2/(3*(exp(2*x) - 1))
```

3.75 $\int \frac{1}{1-\cosh^8(x)} dx$

Optimal result	490
Rubi [A] (verified)	490
Mathematica [A] (verified)	492
Maple [C] (verified)	492
Fricas [B] (verification not implemented)	493
Sympy [F(-1)]	493
Maxima [F]	494
Giac [A] (verification not implemented)	494
Mupad [B] (verification not implemented)	494

Optimal result

Integrand size = 10, antiderivative size = 69

$$\int \frac{1}{1-\cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{4\sqrt{1-i}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{4\sqrt{1+i}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{\operatorname{coth}(x)}{4}$$

[Out] 1/4*coth(x)+1/4*arctanh(tanh(x)/(1-I)^(1/2))/(1-I)^(1/2)+1/4*arctanh(tanh(x)/(1+I)^(1/2))/(1+I)^(1/2)+1/8*arctanh(1/2*2^(1/2)*tanh(x))*2^(1/2)

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3290, 3260, 212, 3254, 3852, 8}

$$\int \frac{1}{1-\cosh^8(x)} dx = \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{4\sqrt{1-i}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{4\sqrt{1+i}} + \frac{\operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{\operatorname{coth}(x)}{4}$$

[In] Int[(1 - Cosh[x]^8)^(-1), x]

[Out] ArcTanh[Tanh[x]/Sqrt[1 - I]]/(4*Sqrt[1 - I]) + ArcTanh[Tanh[x]/Sqrt[1 + I]]/(4*Sqrt[1 + I]) + ArcTanh[Tanh[x]/Sqrt[2]]/(4*Sqrt[2]) + Coth[x]/4

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

Q[a, 0] || LtQ[b, 0])

Rule 3254

Int[(u_.)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := Dist[a^p, Int[ActivateTrig[u*cos[e + f*x]^(2*p)], x], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ[a + b, 0] && IntegerQ[p]

Rule 3260

Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(-1), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[ff/f, Subst[Int[1/(a + (a + b)*ff^2*x^2), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{a, b, e, f}, x]

Rule 3290

Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^(n_))^(n_)^(-1), x_Symbol] := Module[{k}, Dist[2/(a*n), Sum[Int[1/(1 - Sin[e + f*x]^2/((-1)^(4*(k/n))*Rt[-a/b, n/2])], x], {k, 1, n/2}], x]] /; FreeQ[{a, b, e, f}, x] && IntegerQ[n/2]

Rule 3852

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{1}{4} \int \frac{1}{1 - \cosh^2(x)} dx + \frac{1}{4} \int \frac{1}{1 - i \cosh^2(x)} dx \\
 &\quad + \frac{1}{4} \int \frac{1}{1 + i \cosh^2(x)} dx + \frac{1}{4} \int \frac{1}{1 + \cosh^2(x)} dx \\
 &= -\left(\frac{1}{4} \int \text{csch}^2(x) dx\right) + \frac{1}{4} \text{Subst}\left(\int \frac{1}{1 - 2x^2} dx, x, \coth(x)\right) \\
 &\quad + \frac{1}{4} \text{Subst}\left(\int \frac{1}{1 - (1 + i)x^2} dx, x, \coth(x)\right) \\
 &\quad + \frac{1}{4} \text{Subst}\left(\int \frac{1}{1 - (1 - i)x^2} dx, x, \coth(x)\right) \\
 &= \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{4\sqrt{1-i}} + \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{4\sqrt{1+i}} + \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{1}{4} i \text{Subst}\left(\int 1 dx, x, -i \coth(x)\right) \\
 &= \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{4\sqrt{1-i}} + \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{4\sqrt{1+i}} + \frac{\arctanh\left(\frac{\tanh(x)}{\sqrt{2}}\right)}{4\sqrt{2}} + \frac{\coth(x)}{4}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.61 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.93

$$\int \frac{1}{1 - \cosh^8(x)} dx = \frac{1}{8} \left(\frac{2 \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1-i}}\right)}{\sqrt{1-i}} + \frac{2 \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{1+i}}\right)}{\sqrt{1+i}} + \sqrt{2} \operatorname{arctanh}\left(\frac{\tanh(x)}{\sqrt{2}}\right) + 2 \operatorname{coth}(x) \right)$$

[In] Integrate[(1 - Cosh[x]^8)^(-1), x]

[Out] ((2*ArcTanh[Tanh[x]/Sqrt[1 - I]]/Sqrt[1 - I] + (2*ArcTanh[Tanh[x]/Sqrt[1 + I]]/Sqrt[1 + I] + Sqrt[2]*ArcTanh[Tanh[x]/Sqrt[2]] + 2*Coth[x])/8

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 1.66 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.17

method	result
risch	$\frac{1}{2e^{2x}-2} + \frac{\sqrt{2} \ln(e^{2x}+3-2\sqrt{2})}{16} - \frac{\sqrt{2} \ln(e^{2x}+3+2\sqrt{2})}{16} + \left(\sum_{R=\text{RootOf}(8192Z^4-128Z^2+1)} -R \ln(-2048R^3 + \dots) \right)$
default	$\frac{\tanh(\frac{x}{2})}{8} + \frac{\sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})^2 + \tanh(\frac{x}{2})\sqrt{2}+1}{\tanh(\frac{x}{2})^2 - \tanh(\frac{x}{2})\sqrt{2}+1}\right) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}+1) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}-1) \right)}{32} - \frac{\sqrt{2} \left(\ln\left(\frac{\tanh(\frac{x}{2})^2 - \tanh(\frac{x}{2})\sqrt{2}+1}{\tanh(\frac{x}{2})^2 + \tanh(\frac{x}{2})\sqrt{2}+1}\right) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}+1) + 2 \arctan(\tanh(\frac{x}{2})\sqrt{2}-1) \right)}{32}$

[In] int(1/(1-cosh(x)^8), x, method=_RETURNVERBOSE)

[Out] 1/2/(exp(2*x)-1)+1/16*2^(1/2)*ln(exp(2*x)+3-2*2^(1/2))-1/16*2^(1/2)*ln(exp(2*x)+3+2*2^(1/2))+sum(_R*ln(-2048*_R^3+256*_R^2+exp(2*x)-1), _R=RootOf(8192*_Z^4-128*_Z^2+1))

Fricas [B] (verification not implemented)

Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 358 vs. $2(42) = 84$.

Time = 0.26 (sec) , antiderivative size = 358, normalized size of antiderivative = 5.19

$$\int \frac{1}{1 - \cosh^8(x)} dx =$$

$$\frac{\sqrt{i+1}(\sqrt{2} \cosh(x)^2 + 2\sqrt{2} \cosh(x) \sinh(x) + \sqrt{2} \sinh(x)^2 - \sqrt{2}) \log(\cosh(x)^2 + 2 \cosh(x) \sinh(x))}{-}$$

[In] integrate(1/(1-cosh(x)^8),x, algorithm="fricas")

[Out] $-1/16*(\sqrt{I+1}*(\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + (I+1)*\sqrt{2}*\sqrt{I+1} + 2*I+1) - \sqrt{I+1}*(\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - (I+1)*\sqrt{2}*\sqrt{I+1} + 2*I+1) + \sqrt{-I+1}*(\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - (I-1)*\sqrt{2}*\sqrt{-I+1} - 2*I+1) - \sqrt{-I+1}*(\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 + (I-1)*\sqrt{2}*\sqrt{-I+1} - 2*I+1) - (\sqrt{2}*\cosh(x)^2 + 2*\sqrt{2}*\cosh(x)*\sinh(x) + \sqrt{2}*\sinh(x)^2 - \sqrt{2})*\log(-(3*(2*\sqrt{2}) - 3)*\cosh(x)^2 - 4*(3*\sqrt{2}) - 4)*\cosh(x)*\sinh(x) + 3*(2*\sqrt{2}) - 3)*\sinh(x)^2 + 2*\sqrt{2} - 3)/(\cosh(x)^2 + \sinh(x)^2 + 3) - 8)/(\cosh(x)^2 + 2*\cosh(x)*\sinh(x) + \sinh(x)^2 - 1)$

Sympy [F(-1)]

Timed out.

$$\int \frac{1}{1 - \cosh^8(x)} dx = \text{Timed out}$$

[In] integrate(1/(1-cosh(x)**8),x)

[Out] Timed out

[In] $\text{int}(-1/(\cosh(x)^8 - 1), x)$

[Out] $(2^{1/2} \cdot \log(582732658686033920 \cdot \exp(2x) + 70697326355677184 \cdot 2^{1/2} + 412054214575915008 \cdot 2^{1/2} \cdot \exp(2x) + 99981117754441728)) / 16 - (2^{1/2} \cdot \log(70697326355677184 \cdot 2^{1/2} - 582732658686033920 \cdot \exp(2x) + 412054214575915008 \cdot 2^{1/2} \cdot \exp(2x) - 99981117754441728)) / 16 + 1/(2 \cdot (\exp(2x) - 1)) - (2^{1/2} \cdot (1 - i)^{1/2} \cdot \log((70836483296067584 - 69311013991743488i) - 2^{1/2} \cdot (1 - i)^{1/2} \cdot (54684829282729984 - 21956972328779776i) - 2^{1/2} \cdot (1 - i)^{1/2} \cdot \exp(2x) \cdot (12296353929494528 - 271474128182050816i) - \exp(2x) \cdot (155613434002538496 + 429723297714798592i))) / 16 + (2^{1/2} \cdot (1 - i)^{1/2} \cdot \log(2^{1/2} \cdot (1 - i)^{1/2} \cdot (54684829282729984 - 21956972328779776i) - \exp(2x) \cdot (155613434002538496 + 429723297714798592i) + 2^{1/2} \cdot (1 - i)^{1/2} \cdot \exp(2x) \cdot (12296353929494528 - 271474128182050816i) + (70836483296067584 - 69311013991743488i))) / 16 - (2^{1/2} \cdot (1 + i)^{1/2} \cdot \log((70836483296067584 + 69311013991743488i) - 2^{1/2} \cdot (1 + i)^{1/2} \cdot (54684829282729984 + 21956972328779776i) - 2^{1/2} \cdot (1 + i)^{1/2} \cdot \exp(2x) \cdot (12296353929494528 + 271474128182050816i) - \exp(2x) \cdot (155613434002538496 - 429723297714798592i))) / 16 + (2^{1/2} \cdot (1 + i)^{1/2} \cdot \log(2^{1/2} \cdot (1 + i)^{1/2} \cdot (54684829282729984 + 21956972328779776i) - \exp(2x) \cdot (155613434002538496 - 429723297714798592i) + 2^{1/2} \cdot (1 + i)^{1/2} \cdot \exp(2x) \cdot (12296353929494528 + 271474128182050816i) + (70836483296067584 + 69311013991743488i))) / 16$

3.76 $\int \frac{\tanh(x)}{1+\cosh^2(x)} dx$

Optimal result	496
Rubi [A] (verified)	496
Mathematica [A] (verified)	497
Maple [A] (verified)	497
Fricas [B] (verification not implemented)	498
Sympy [F]	498
Maxima [A] (verification not implemented)	498
Giac [A] (verification not implemented)	499
Mupad [B] (verification not implemented)	499

Optimal result

Integrand size = 11, antiderivative size = 15

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = \log(\cosh(x)) - \frac{1}{2} \log(1 + \cosh^2(x))$$

[Out] $\ln(\cosh(x)) - 1/2 * \ln(1 + \cosh(x)^2)$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.364$, Rules used = {3273, 36, 29, 31}

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = \log(\cosh(x)) - \frac{1}{2} \log(\cosh^2(x) + 1)$$

[In] $\text{Int}[\text{Tanh}[x]/(1 + \text{Cosh}[x]^2), x]$

[Out] $\text{Log}[\text{Cosh}[x]] - \text{Log}[1 + \text{Cosh}[x]^2]/2$

Rule 29

$\text{Int}[(x_)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[x], x]$

Rule 31

$\text{Int}[(a_) + (b_)*(x_)^{(-1)}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}\{a, b\}, x]$

Rule 36


```
Int[1/(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))), x_Symbol] := Dist[b/(b*c
- a*d), Int[1/(a + b*x), x], x] - Dist[d/(b*c - a*d), Int[1/(c + d*x), x],
x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0]
```

Rule 3273

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.)*tan[(e_.) + (f_.)*(x_)]^
(m_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Dist[ff^((m
+ 1)/2)/(2*f), Subst[Int[x^((m - 1)/2)*((a + b*ff*x)^p/(1 - ff*x)^((m + 1)
/2)), x], x, Sin[e + f*x]^2/ff], x] /; FreeQ[{a, b, e, f, p}, x] && Intege
rQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{x(1+x)} dx, x, \cosh^2(x) \right) \\ &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{x} dx, x, \cosh^2(x) \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x} dx, x, \cosh^2(x) \right) \\ &= \log(\cosh(x)) - \frac{1}{2} \log(1 + \cosh^2(x)) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 15, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = \log(\cosh(x)) - \frac{1}{2} \log(1 + \cosh^2(x))$$

```
[In] Integrate[Tanh[x]/(1 + Cosh[x]^2),x]
```

```
[Out] Log[Cosh[x]] - Log[1 + Cosh[x]^2]/2
```

Maple [A] (verified)

Time = 0.42 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.47

method	result	size
default	$-\frac{\ln(\tanh(\frac{x}{2})^4 + 1)}{2} + \ln\left(1 + \tanh\left(\frac{x}{2}\right)^2\right)$	22
risch	$\ln(1 + e^{2x}) - \frac{\ln(e^{4x} + 6e^{2x} + 1)}{2}$	24

```
[In] int(tanh(x)/(1+cosh(x)^2),x,method=_RETURNVERBOSE)
```

[Out] $-1/2*\ln(\tanh(1/2*x)^4+1)+\ln(1+\tanh(1/2*x)^2)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 47 vs. $2(13) = 26$.

Time = 0.25 (sec) , antiderivative size = 47, normalized size of antiderivative = 3.13

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = -\frac{1}{2} \log \left(\frac{2 (\cosh(x)^2 + \sinh(x)^2 + 3)}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2} \right) + \log \left(\frac{2 \cosh(x)}{\cosh(x) - \sinh(x)} \right)$$

[In] `integrate(tanh(x)/(1+cosh(x)^2),x, algorithm="fricas")`

[Out] $-1/2*\log(2*(\cosh(x)^2 + \sinh(x)^2 + 3)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)) + \log(2*\cosh(x)/(\cosh(x) - \sinh(x)))$

Sympy [F]

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = \int \frac{\tanh(x)}{\cosh^2(x) + 1} dx$$

[In] `integrate(tanh(x)/(1+cosh(x)**2),x)`

[Out] `Integral(tanh(x)/(cosh(x)**2 + 1), x)`

Maxima [A] (verification not implemented)

none

Time = 0.19 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = -\frac{1}{2} \log (6 e^{(-2x)} + e^{(-4x)} + 1) + \log (e^{(-2x)} + 1)$$

[In] `integrate(tanh(x)/(1+cosh(x)^2),x, algorithm="maxima")`

[Out] $-1/2*\log(6*e^{(-2*x)} + e^{(-4*x)} + 1) + \log(e^{(-2*x)} + 1)$

Giac [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 23, normalized size of antiderivative = 1.53

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = -\frac{1}{2} \log(e^{4x} + 6e^{2x} + 1) + \log(e^{2x} + 1)$$

[In] integrate(tanh(x)/(1+cosh(x)^2),x, algorithm="giac")

[Out] -1/2*log(e^(4*x) + 6*e^(2*x) + 1) + log(e^(2*x) + 1)

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.80

$$\int \frac{\tanh(x)}{1 + \cosh^2(x)} dx = \ln(-5184e^{2x} - 5184) - \frac{\ln(54e^{2x} + 9e^{4x} + 9)}{2}$$

[In] int(tanh(x)/(cosh(x)^2 + 1),x)

[Out] log(- 5184*exp(2*x) - 5184) - log(54*exp(2*x) + 9*exp(4*x) + 9)/2

3.77 $\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx$

Optimal result	500
Rubi [A] (verified)	500
Mathematica [A] (verified)	501
Maple [A] (verified)	502
Fricas [B] (verification not implemented)	502
Sympy [F]	503
Maxima [F]	503
Giac [F]	503
Mupad [F(-1)]	503

Optimal result

Integrand size = 15, antiderivative size = 39

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = -\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cosh^2(x)}}{\sqrt{a}}\right) + \sqrt{a + b \cosh^2(x)}$$

[Out] $-\operatorname{arctanh}((a+b*\cosh(x)^2)^{(1/2)/a^{(1/2))}*a^{(1/2)}+(a+b*\cosh(x)^2)^{(1/2)}$

Rubi [A] (verified)

Time = 0.05 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3273, 52, 65, 214}

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = \sqrt{a + b \cosh^2(x)} - \sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cosh^2(x)}}{\sqrt{a}}\right)$$

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^2]*\operatorname{Tanh}[x], x]$

[Out] $-(\operatorname{Sqrt}[a]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^2]/\operatorname{Sqrt}[a]]) + \operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^2]$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 3273

```
Int[((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)]^2)^(p_.)*tan[(e_.) + (f_.)*(x_)]^
(m_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Dist[ff^((m
+ 1)/2)/(2*f), Subst[Int[x^((m - 1)/2)*((a + b*ff*x)^p/(1 - ff*x)^((m + 1)
/2)), x], x, Sin[e + f*x]^2/ff], x]] /; FreeQ[{a, b, e, f, p}, x] && Intege
rQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{1}{2} \text{Subst} \left(\int \frac{\sqrt{a+bx}}{x} dx, x, \cosh^2(x) \right) \\
&= \sqrt{a+b \cosh^2(x)} + \frac{1}{2} a \text{Subst} \left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cosh^2(x) \right) \\
&= \sqrt{a+b \cosh^2(x)} + \frac{a \text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a+b \cosh^2(x)} \right)}{b} \\
&= -\sqrt{a} \arctanh \left(\frac{\sqrt{a+b \cosh^2(x)}}{\sqrt{a}} \right) + \sqrt{a+b \cosh^2(x)}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 39, normalized size of antiderivative = 1.00

$$\int \sqrt{a+b \cosh^2(x)} \tanh(x) dx = -\sqrt{a} \arctanh \left(\frac{\sqrt{a+b \cosh^2(x)}}{\sqrt{a}} \right) + \sqrt{a+b \cosh^2(x)}$$

```
[In] Integrate[Sqrt[a + b*Cosh[x]^2]*Tanh[x], x]
```

```
[Out] -(Sqrt[a]*ArcTanh[Sqrt[a + b*Cosh[x]^2]/Sqrt[a]]) + Sqrt[a + b*Cosh[x]^2]
```

Maple [A] (verified)

Time = 0.16 (sec) , antiderivative size = 42, normalized size of antiderivative = 1.08

method	result	size
default	$\sqrt{a + b \cosh(x)^2} - \sqrt{a} \ln\left(\frac{2a + 2\sqrt{a} \sqrt{a + b \cosh(x)^2}}{\cosh(x)}\right)$	42

[In] `int((a+b*cosh(x)^2)^(1/2)*tanh(x),x,method=_RETURNVERBOSE)`

[Out] $(a+b*\cosh(x)^2)^{(1/2)}-a^{(1/2)}*\ln((2*a+2*a^{(1/2)}*(a+b*\cosh(x)^2)^{(1/2)})/\cosh(x))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 122 vs. $2(31) = 62$.

Time = 0.41 (sec) , antiderivative size = 357, normalized size of antiderivative = 9.15

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx$$

$$= \frac{\sqrt{a}(\cosh(x) + \sinh(x)) \log\left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(4a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 4a+b) \sinh(x)^2 - 4\sqrt{a} \cosh(x) \sinh(x)}{\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^3 + \cosh(x)) \sinh(x) + 1}\right)}{2(\cosh(x) + \sinh(x))}$$

[In] `integrate((a+b*cosh(x)^2)^(1/2)*tanh(x),x, algorithm="fricas")`

[Out] $[1/2*(\sqrt{a}*(\cosh(x) + \sinh(x))*\log((b*\cosh(x)^4 + 4*b*\cosh(x)*\sinh(x)^3 + b*\sinh(x)^4 + 2*(4*a + b)*\cosh(x)^2 + 2*(3*b*\cosh(x)^2 + 4*a + b)*\sinh(x)^2 - 4*\sqrt{2}*\sqrt{a}*\sqrt{(b*\cosh(x)^2 + b*\sinh(x)^2 + 2*a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))*(\cosh(x) + \sinh(x)) + 4*(b*\cosh(x)^3 + (4*a + b)*\cosh(x))*\sinh(x) + b)/(\cosh(x)^4 + 4*\cosh(x)*\sinh(x)^3 + \sinh(x)^4 + 2*(3*\cosh(x)^2 + 1)*\sinh(x)^2 + 2*\cosh(x)^2 + 4*(\cosh(x)^3 + \cosh(x))*\sinh(x) + 1)) + \sqrt{2}*\sqrt{(b*\cosh(x)^2 + b*\sinh(x)^2 + 2*a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(\cosh(x) + \sinh(x)), 1/2*(2*\sqrt{-a}*(\cosh(x) + \sinh(x))*\arctan(1/2*\sqrt{2}*\sqrt{-a}*\sqrt{(b*\cosh(x)^2 + b*\sinh(x)^2 + 2*a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(a*\cosh(x) + a*\sinh(x)) + \sqrt{2}*\sqrt{(b*\cosh(x)^2 + b*\sinh(x)^2 + 2*a + b)/(\cosh(x)^2 - 2*\cosh(x)*\sinh(x) + \sinh(x)^2)))/(\cosh(x) + \sinh(x))]$

Sympy [F]

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = \int \sqrt{a + b \cosh^2(x)} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)**2)**(1/2)*tanh(x), x)

[Out] Integral(sqrt(a + b*cosh(x)**2)*tanh(x), x)

Maxima [F]

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = \int \sqrt{b \cosh^2(x) + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^2)^(1/2)*tanh(x), x, algorithm="maxima")

[Out] integrate(sqrt(b*cosh(x)^2 + a)*tanh(x), x)

Giac [F]

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = \int \sqrt{b \cosh^2(x) + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^2)^(1/2)*tanh(x), x, algorithm="giac")

[Out] integrate(sqrt(b*cosh(x)^2 + a)*tanh(x), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cosh^2(x)} \tanh(x) dx = \int \tanh(x) \sqrt{b \cosh^2(x) + a} dx$$

[In] int(tanh(x)*(a + b*cosh(x)^2)^(1/2), x)

[Out] int(tanh(x)*(a + b*cosh(x)^2)^(1/2), x)

$$3.78 \quad \int \frac{\tanh(x)}{\sqrt{a+b \cosh^2(x)}} dx$$

Optimal result	504
Rubi [A] (verified)	504
Mathematica [A] (verified)	505
Maple [A] (verified)	506
Fricas [B] (verification not implemented)	506
Sympy [F]	507
Maxima [F]	507
Giac [F]	507
Mupad [F(-1)]	507

Optimal result

Integrand size = 15, antiderivative size = 26

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}$$

[Out] $-\operatorname{arctanh}((a+b*\cosh(x)^2)^{(1/2)/a^{(1/2)})/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3273, 65, 214}

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^2(x)}} dx = -\frac{\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^2(x)}}{\sqrt{a}}\right)}{\sqrt{a}}$$

[In] $\operatorname{Int}[\operatorname{Tanh}[x]/\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^2], x]$

[Out] $-(\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^2]/\operatorname{Sqrt}[a]]/\operatorname{Sqrt}[a])$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
```


ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 3273

Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^(p_)*tan[(e_) + (f_)*(x_)]^(m_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Dist[ff^((m + 1)/2)/(2*f), Subst[Int[x^((m - 1)/2)*((a + b*ff*x)^p/(1 - ff*x)^((m + 1)/2)), x], x, Sin[e + f*x]^2/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{x \sqrt{a + bx}} dx, x, \cosh^2(x) \right) \\ &= \frac{\text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a + b \cosh^2(x)} \right)}{b} \\ &= -\frac{\text{arctanh} \left(\frac{\sqrt{a + b \cosh^2(x)}}{\sqrt{a}} \right)}{\sqrt{a}} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 26, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx = -\frac{\text{arctanh} \left(\frac{\sqrt{a + b \cosh^2(x)}}{\sqrt{a}} \right)}{\sqrt{a}}$$

[In] Integrate[Tanh[x]/Sqrt[a + b*Cosh[x]^2], x]

[Out] -(ArcTanh[Sqrt[a + b*Cosh[x]^2]/Sqrt[a]]/Sqrt[a])

Maple [A] (verified)

Time = 0.14 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.19

method	result	size
default	$-\frac{\ln\left(\frac{2a+2\sqrt{a}\sqrt{a+b\cosh(x)^2}}{\cosh(x)}\right)}{\sqrt{a}}$	31

[In] `int(tanh(x)/(a+b*cosh(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-1/a^(1/2)*ln((2*a+2*a^(1/2)*(a+b*cosh(x)^2)^(1/2))/cosh(x))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 68 vs. 2(20) = 40.

Time = 0.30 (sec) , antiderivative size = 248, normalized size of antiderivative = 9.54

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx$$

$$= \frac{\log\left(\frac{b \cosh(x)^4 + 4b \cosh(x) \sinh(x)^3 + b \sinh(x)^4 + 2(4a+b) \cosh(x)^2 + 2(3b \cosh(x)^2 + 4a+b) \sinh(x)^2 - 4\sqrt{2}\sqrt{a}\sqrt{\frac{b \cosh(x)^2 + b \sinh(x)^2 + 2a}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2}}}{\cosh(x)^4 + 4 \cosh(x) \sinh(x)^3 + \sinh(x)^4 + 2(3 \cosh(x)^2 + 1) \sinh(x)^2 + 2 \cosh(x)^2 + 4(\cosh(x)^2 + 4 \cosh(x) \sinh(x) + \sinh(x)^2)}\right)}{2\sqrt{a}}$$

[In] `integrate(tanh(x)/(a+b*cosh(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `[1/2*log((b*cosh(x)^4 + 4*b*cosh(x)*sinh(x)^3 + b*sinh(x)^4 + 2*(4*a + b)*cosh(x)^2 + 2*(3*b*cosh(x)^2 + 4*a + b)*sinh(x)^2 - 4*sqrt(2)*sqrt(a)*sqrt((b*cosh(x)^2 + b*sinh(x)^2 + 2*a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2))*(cosh(x) + sinh(x)) + 4*(b*cosh(x)^3 + (4*a + b)*cosh(x))*sinh(x) + b)/(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1))/sqrt(a), sqrt(-a)*arctan(1/2*sqrt(2)*sqrt(-a)*sqrt((b*cosh(x)^2 + b*sinh(x)^2 + 2*a + b)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/(a*cosh(x) + a*sinh(x)))/a]`

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)**2)**(1/2),x)

[Out] Integral(tanh(x)/sqrt(a + b*cosh(x)**2), x)

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^2 + a}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tanh(x)/sqrt(b*cosh(x)^2 + a), x)

Giac [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^2 + a}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^2 + a}} dx$$

[In] int(tanh(x)/(a + b*cosh(x)^2)^(1/2),x)

[Out] int(tanh(x)/(a + b*cosh(x)^2)^(1/2), x)

$$3.79 \quad \int \frac{\tanh(x)}{\sqrt{1+\cosh^2(x)}} dx$$

Optimal result	508
Rubi [A] (verified)	508
Mathematica [A] (verified)	509
Maple [A] (verified)	509
Fricas [B] (verification not implemented)	510
Sympy [F]	510
Maxima [F]	510
Giac [F]	511
Mupad [F(-1)]	511

Optimal result

Integrand size = 13, antiderivative size = 13

$$\int \frac{\tanh(x)}{\sqrt{1+\cosh^2(x)}} dx = -\operatorname{arctanh}\left(\sqrt{1+\cosh^2(x)}\right)$$

[Out] `-arctanh((1+cosh(x)^2)^(1/2))`

Rubi [A] (verified)

Time = 0.03 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {3273, 65, 213}

$$\int \frac{\tanh(x)}{\sqrt{1+\cosh^2(x)}} dx = -\operatorname{arctanh}\left(\sqrt{\cosh^2(x)+1}\right)$$

[In] `Int[Tanh[x]/Sqrt[1 + Cosh[x]^2],x]`

[Out] `-ArcTanh[Sqrt[1 + Cosh[x]^2]]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 213

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])
```

Rule 3273

```
Int[((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^p*(tan[(e_) + (f_)*(x_)]^m), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Dist[ff^((m + 1)/2)/(2*f), Subst[Int[x^((m - 1)/2)*((a + b*ff*x)^p/(1 - ff*x)^((m + 1)/2)), x], x, Sin[e + f*x]^2/ff], x] /; FreeQ[{a, b, e, f, p}, x] && IntegerQ[(m - 1)/2]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1+x}} dx, x, \cosh^2(x) \right) \\ &= \text{Subst} \left(\int \frac{1}{-1+x^2} dx, x, \sqrt{1+\cosh^2(x)} \right) \\ &= -\text{arctanh} \left(\sqrt{1+\cosh^2(x)} \right) \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{1+\cosh^2(x)}} dx = -\text{arctanh} \left(\sqrt{1+\cosh^2(x)} \right)$$

```
[In] Integrate[Tanh[x]/Sqrt[1 + Cosh[x]^2], x]
```

```
[Out] -ArcTanh[Sqrt[1 + Cosh[x]^2]]
```

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.92

method	result	size
default	$-\text{arctanh} \left(\frac{1}{\sqrt{1+\cosh(x)^2}} \right)$	12

```
[In] int(tanh(x)/(1+cosh(x)^2)^(1/2), x, method=_RETURNVERBOSE)
```

```
[Out] -arctanh(1/(1+cosh(x)^2)^(1/2))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 4.85

$$\int \frac{\tanh(x)}{\sqrt{1 + \cosh^2(x)}} dx = \log \left(\frac{\sqrt{2} \sqrt{\frac{\cosh(x)^2 + \sinh(x)^2 + 3}{\cosh(x)^2 - 2 \cosh(x) \sinh(x) + \sinh(x)^2}} - 2 \cosh(x) - 2 \sinh(x)}{\cosh(x)^2 + 2 \cosh(x) \sinh(x) + \sinh(x)^2 + 1} \right)$$

[In] integrate(tanh(x)/(1+cosh(x)^2)^(1/2),x, algorithm="fricas")

[Out] log((sqrt(2)*sqrt((cosh(x)^2 + sinh(x)^2 + 3)/(cosh(x)^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) - 2*cosh(x) - 2*sinh(x))/(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2 + 1))

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{\cosh^2(x) + 1}} dx$$

[In] integrate(tanh(x)/(1+cosh(x)**2)**(1/2),x)

[Out] Integral(tanh(x)/sqrt(cosh(x)**2 + 1), x)

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] integrate(tanh(x)/(1+cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] integrate(tanh(x)/sqrt(cosh(x)^2 + 1), x)

Giac [F]

$$\int \frac{\tanh(x)}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] integrate(tanh(x)/(1+cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] integrate(tanh(x)/sqrt(cosh(x)^2 + 1), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{1 + \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{\cosh(x)^2 + 1}} dx$$

[In] int(tanh(x)/(cosh(x)^2 + 1)^(1/2),x)

[Out] int(tanh(x)/(cosh(x)^2 + 1)^(1/2), x)

$$3.80 \quad \int \frac{\tanh(x)}{\sqrt{1-\cosh^2(x)}} dx$$

Optimal result	512
Rubi [A] (verified)	512
Mathematica [A] (verified)	513
Maple [C] (verified)	514
Fricas [B] (verification not implemented)	514
Sympy [F]	515
Maxima [C] (verification not implemented)	515
Giac [C] (verification not implemented)	515
Mupad [F(-1)]	516

Optimal result

Integrand size = 15, antiderivative size = 13

$$\int \frac{\tanh(x)}{\sqrt{1-\cosh^2(x)}} dx = -\operatorname{arctanh}\left(\sqrt{-\sinh^2(x)}\right)$$

[Out] -arctanh((-sinh(x)^2)^(1/2))

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3255, 3284, 65, 212}

$$\int \frac{\tanh(x)}{\sqrt{1-\cosh^2(x)}} dx = -\operatorname{arctanh}\left(\sqrt{-\sinh^2(x)}\right)$$

[In] Int[Tanh[x]/Sqrt[1 - Cosh[x]^2],x]

[Out] -ArcTanh[Sqrt[-Sinh[x]^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 212


```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 3255

```
Int[(u_)*((a_) + (b_)*sin[(e_) + (f_)*(x_)]^2)^p, x_Symbol] := Int[A
ctivateTrig[u*(a*cos[e + f*x]^2)^p], x] /; FreeQ[{a, b, e, f, p}, x] && EqQ
[a + b, 0]
```

Rule 3284

```
Int[((b_)*sin[(e_) + (f_)*(x_)]^n)^p*tan[(e_) + (f_)*(x_)]^m,
x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x]^2, x]}, Dist[ff^((m + 1
)/2)/(2*f), Subst[Int[x^((m - 1)/2)*((b*ff^(n/2)*x^(n/2))^p/(1 - ff*x)^((m
+ 1)/2)), x], x, Sin[e + f*x]^2/ff], x] /; FreeQ[{b, e, f, p}, x] && IntegerQ[(m - 1)/2] && IntegerQ[n/2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \frac{\tanh(x)}{\sqrt{-\sinh^2(x)}} dx \\
&= \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{-x(1+x)}} dx, x, \sinh^2(x) \right) \\
&= -\text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \sqrt{-\sinh^2(x)} \right) \\
&= -\text{arctanh} \left(\sqrt{-\sinh^2(x)} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.23

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx = \frac{\arctan(\sinh(x)) \sinh(x)}{\sqrt{-\sinh^2(x)}}$$

```
[In] Integrate[Tanh[x]/Sqrt[1 - Cosh[x]^2], x]
```

```
[Out] (ArcTan[Sinh[x]]*Sinh[x])/Sqrt[-Sinh[x]^2]
```

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 0.16 (sec) , antiderivative size = 19, normalized size of antiderivative = 1.46

method	result	size
default	'int/indef0' $\left(\frac{\sinh(x)}{\cosh(x)^2 \sqrt{-\sinh(x)^2}}, \sinh(x) \right)$	19
risch	$\frac{ie^{-x}(e^{2x}-1)\ln(e^x+i)}{\sqrt{-(e^{2x}-1)^2e^{-2x}}} - \frac{ie^{-x}(e^{2x}-1)\ln(e^x-i)}{\sqrt{-(e^{2x}-1)^2e^{-2x}}}$	72

[In] `int(tanh(x)/(1-cosh(x)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] 'int/indef0'(sinh(x)/cosh(x)^2/(-sinh(x)^2)^(1/2),sinh(x))

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(11) = 22.

Time = 0.27 (sec) , antiderivative size = 112, normalized size of antiderivative = 8.62

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx$$

$$= \log \left(\frac{\cosh(x) e^{(2x)} + (e^{(2x)} - 1) \sinh(x) + \sqrt{-(e^{(4x)} - 2e^{(2x)} + 1)e^{(-2x)}e^x - \cosh(x)}}{e^{(2x)} - 1} \right)$$

$$- \log \left(\frac{\cosh(x) e^{(2x)} + (e^{(2x)} - 1) \sinh(x) - \sqrt{-(e^{(4x)} - 2e^{(2x)} + 1)e^{(-2x)}e^x - \cosh(x)}}{e^{(2x)} - 1} \right)$$

[In] `integrate(tanh(x)/(1-cosh(x)^2)^(1/2),x, algorithm="fricas")`

[Out] `log((cosh(x)*e^(2*x) + (e^(2*x) - 1)*sinh(x) + sqrt(-(e^(4*x) - 2*e^(2*x) + 1)*e^(-2*x))*e^x - cosh(x))/(e^(2*x) - 1)) - log((cosh(x)*e^(2*x) + (e^(2*x) - 1)*sinh(x) - sqrt(-(e^(4*x) - 2*e^(2*x) + 1)*e^(-2*x))*e^x - cosh(x))/(e^(2*x) - 1))`

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{-(\cosh(x) - 1)(\cosh(x) + 1)}} dx$$

[In] integrate(tanh(x)/(1-cosh(x)**2)**(1/2),x)

[Out] Integral(tanh(x)/sqrt(-(cosh(x) - 1)*(cosh(x) + 1)), x)

Maxima [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 7, normalized size of antiderivative = 0.54

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx = -2i \arctan(e^{-x})$$

[In] integrate(tanh(x)/(1-cosh(x)^2)^(1/2),x, algorithm="maxima")

[Out] -2*I*arctan(e^(-x))

Giac [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 38, normalized size of antiderivative = 2.92

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx = -\frac{\log(e^x + i)}{\operatorname{sgn}(-e^{3x} + e^x)} + \frac{\log(e^x - i)}{\operatorname{sgn}(-e^{3x} + e^x)}$$

[In] integrate(tanh(x)/(1-cosh(x)^2)^(1/2),x, algorithm="giac")

[Out] -log(e^x + I)/sgn(-e^(3*x) + e^x) + log(e^x - I)/sgn(-e^(3*x) + e^x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{1 - \cosh^2(x)}} dx = \int \frac{\tanh(x)}{\sqrt{1 - \cosh(x)^2}} dx$$

```
[In] int(tanh(x)/(1 - cosh(x)^2)^(1/2),x)
```

```
[Out] int(tanh(x)/(1 - cosh(x)^2)^(1/2), x)
```

3.81 $\int \frac{\tanh^3(x)}{a+b \cosh^3(x)} dx$

Optimal result	517
Rubi [A] (verified)	517
Mathematica [C] (verified)	521
Maple [C] (verified)	521
Fricas [C] (verification not implemented)	522
Sympy [F]	523
Maxima [F]	523
Giac [A] (verification not implemented)	523
Mupad [B] (verification not implemented)	524

Optimal result

Integrand size = 15, antiderivative size = 153

$$\int \frac{\tanh^3(x)}{a+b \cosh^3(x)} dx = -\frac{b^{2/3} \arctan\left(\frac{\sqrt[3]{a}-2\sqrt[3]{b} \cosh(x)}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt{3}a^{5/3}} + \frac{\log(\cosh(x))}{a} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b} \cosh(x)\right)}{3a^{5/3}} - \frac{b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b} \cosh(x) + b^{2/3} \cosh^2(x)\right)}{6a^{5/3}} - \frac{\log(a+b \cosh^3(x))}{3a} + \frac{\operatorname{sech}^2(x)}{2a}$$

```
[Out] ln(cosh(x))/a+1/3*b^(2/3)*ln(a^(1/3)+b^(1/3)*cosh(x))/a^(5/3)-1/6*b^(2/3)*ln(a^(2/3)-a^(1/3)*b^(1/3)*cosh(x)+b^(2/3)*cosh(x)^2)/a^(5/3)-1/3*ln(a+b*cosh(x)^3)/a+1/2*sech(x)^2/a-1/3*b^(2/3)*arctan(1/3*(a^(1/3)-2*b^(1/3)*cosh(x))/a^(1/3)*3^(1/2))/a^(5/3)*3^(1/2)
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules

used = {3309, 1848, 1885, 206, 31, 648, 631, 210, 642, 266}

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = -\frac{b^{2/3} \arctan\left(\frac{\sqrt[3]{a}-2\sqrt[3]{b} \cosh(x)}{\sqrt{3}\sqrt[3]{a}}\right)}{\sqrt{3}a^{5/3}} - \frac{b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b} \cosh(x) + b^{2/3} \cosh^2(x)\right)}{6a^{5/3}} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b} \cosh(x)\right)}{3a^{5/3}} - \frac{\log(a + b \cosh^3(x))}{3a} + \frac{\operatorname{sech}^2(x)}{2a} + \frac{\log(\cosh(x))}{a}$$

[In] Int[Tanh[x]^3/(a + b*Cosh[x]^3),x]

[Out] -((b^(2/3)*ArcTan[(a^(1/3) - 2*b^(1/3)*Cosh[x])/(Sqrt[3]*a^(1/3))]/(Sqrt[3]*a^(5/3))) + Log[Cosh[x]]/a + (b^(2/3)*Log[a^(1/3) + b^(1/3)*Cosh[x]])/(3*a^(5/3)) - (b^(2/3)*Log[a^(2/3) - a^(1/3)*b^(1/3)*Cosh[x] + b^(2/3)*Cosh[x]^2])/(6*a^(5/3)) - Log[a + b*Cosh[x]^3]/(3*a) + Sech[x]^2/(2*a)

Rule 31

Int[((a_) + (b_.)*(x_)^(-1)), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 206

Int[((a_) + (b_.)*(x_)^3)^(-1), x_Symbol] := Dist[1/(3*Rt[a, 3]^2), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]^2), Int[(2*Rt[a, 3] - Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

Rule 266

Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] := Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)

], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 1848

Int[((Pq_)*((c_)*(x_)^(m_)))/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(Pq/(a + b*x^n)), x], x] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IntegerQ[n] && !IGtQ[m, 0]

Rule 1885

Int[(P2_)/((a_) + (b_)*(x_)^3), x_Symbol] := With[{A = Coeff[P2, x, 0], B = Coeff[P2, x, 1], C = Coeff[P2, x, 2]}, Int[(A + B*x)/(a + b*x^3), x] + Dist[C, Int[x^2/(a + b*x^3), x], x] /; EqQ[a*B^3 - b*A^3, 0] || !RationalQ[a/b] /; FreeQ[{a, b}, x] && PolyQ[P2, x, 2]

Rule 3309

Int[((a_) + (b_)*((c_)*sin[(e_) + (f_)*(x_)])^(n_))^(p_)*tan[(e_) + (f_)*(x_)^(m_)], x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^((m + 1)/2)], x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\text{Subst}\left(\int \frac{1-x^2}{x^3(a+bx^3)} dx, x, \cosh(x)\right) \\
 &= -\text{Subst}\left(\int \left(\frac{1}{ax^3} - \frac{1}{ax} + \frac{b(-1+x^2)}{a(a+bx^3)}\right) dx, x, \cosh(x)\right) \\
 &= \frac{\log(\cosh(x))}{a} + \frac{\text{sech}^2(x)}{2a} - \frac{b\text{Subst}\left(\int \frac{-1+x^2}{a+bx^3} dx, x, \cosh(x)\right)}{a}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{\log(\cosh(x))}{a} + \frac{\operatorname{sech}^2(x)}{2a} + \frac{b \operatorname{Subst}\left(\int \frac{1}{a+bx^3} dx, x, \cosh(x)\right)}{a} - \frac{b \operatorname{Subst}\left(\int \frac{x^2}{a+bx^3} dx, x, \cosh(x)\right)}{a} \\
&= \frac{\log(\cosh(x))}{a} - \frac{\log(a + b \cosh^3(x))}{3a} + \frac{\operatorname{sech}^2(x)}{2a} \\
&\quad + \frac{b \operatorname{Subst}\left(\int \frac{1}{\sqrt[3]{a} + \sqrt[3]{b}x} dx, x, \cosh(x)\right)}{3a^{5/3}} \\
&\quad + \frac{b \operatorname{Subst}\left(\int \frac{2\sqrt[3]{a} - \sqrt[3]{b}x}{a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2} dx, x, \cosh(x)\right)}{3a^{5/3}} \\
&= \frac{\log(\cosh(x))}{a} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b} \cosh(x)\right)}{3a^{5/3}} - \frac{\log(a + b \cosh^3(x))}{3a} \\
&\quad + \frac{\operatorname{sech}^2(x)}{2a} - \frac{b^{2/3} \operatorname{Subst}\left(\int \frac{-\sqrt[3]{a}\sqrt[3]{b} + 2b^{2/3}x}{a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2} dx, x, \cosh(x)\right)}{6a^{5/3}} \\
&\quad + \frac{b \operatorname{Subst}\left(\int \frac{1}{a^{2/3} - \sqrt[3]{a}\sqrt[3]{b}x + b^{2/3}x^2} dx, x, \cosh(x)\right)}{2a^{4/3}} \\
&= \frac{\log(\cosh(x))}{a} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b} \cosh(x)\right)}{3a^{5/3}} \\
&\quad - \frac{b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b} \cosh(x) + b^{2/3} \cosh^2(x)\right)}{6a^{5/3}} - \frac{\log(a + b \cosh^3(x))}{3a} \\
&\quad + \frac{\operatorname{sech}^2(x)}{2a} + \frac{b^{2/3} \operatorname{Subst}\left(\int \frac{1}{-3-x^2} dx, x, 1 - \frac{2\sqrt[3]{b} \cosh(x)}{\sqrt[3]{a}}\right)}{a^{5/3}} \\
&\quad + \frac{b^{2/3} \arctan\left(\frac{1 - \frac{2\sqrt[3]{b} \cosh(x)}{\sqrt[3]{a}}}{\sqrt{3}}\right)}{\sqrt{3}a^{5/3}} + \frac{\log(\cosh(x))}{a} + \frac{b^{2/3} \log\left(\sqrt[3]{a} + \sqrt[3]{b} \cosh(x)\right)}{3a^{5/3}} \\
&= - \frac{b^{2/3} \log\left(a^{2/3} - \sqrt[3]{a}\sqrt[3]{b} \cosh(x) + b^{2/3} \cosh^2(x)\right)}{6a^{5/3}} - \frac{\log(a + b \cosh^3(x))}{3a} \\
&\quad + \frac{\operatorname{sech}^2(x)}{2a}
\end{aligned}$$

Mathematica [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3 in optimal.

Time = 0.35 (sec) , antiderivative size = 145, normalized size of antiderivative = 0.95

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx$$

$$= \frac{-6x + 6 \log(\cosh(x)) - 2 \operatorname{RootSum}\left[b + 3b\#1^2 + 8a\#1^3 + 3b\#1^4 + b\#1^6, \frac{-bx + b \log(e^x - \#1) - 4ax\#1^3 + 4a \log(e^x - \#1)\#1^3 - 3b\#1^4 + 3b \log(e^x - \#1)\#1^4}{b + 2b\#1^2 + 4a\#1^3 + b\#1^4}\right]}{6a}$$

[In] Integrate[Tanh[x]^3/(a + b*Cosh[x]^3),x]

[Out] (-6*x + 6*Log[Cosh[x]] - 2*RootSum[b + 3*b*#1^2 + 8*a*#1^3 + 3*b*#1^4 + b*#1^6 & , (-b*x) + b*Log[E^x - #1] - 4*a*x*#1^3 + 4*a*Log[E^x - #1]*#1^3 - 3*b*x*#1^4 + 3*b*Log[E^x - #1]*#1^4)/(b + 2*b*#1^2 + 4*a*#1^3 + b*#1^4) &] + 3*Sech[x]^2)/(6*a)

Maple [C] (verified)

Result contains higher order function than in optimal. Order 9 vs. order 3.

Time = 4.07 (sec) , antiderivative size = 93, normalized size of antiderivative = 0.61

method	result
risch	$\frac{2e^{2x}}{(1+e^{2x})^2 a} + \frac{\ln(1+e^{2x})}{a} + \left(\sum_{R=\operatorname{RootOf}(27a^5 Z^3 + 27a^4 Z^2 + 9a^3 Z + a^2 - b^2)} -R \ln \left(e^{2x} + \left(\frac{6a^2 R}{b} + \frac{2a}{b} \right) e^x + \frac{(-R^2 a - R^2 b - 2)}{3a} \right) \right)$
default	$\frac{2}{\left(1 + \tanh\left(\frac{x}{2}\right)\right)^2} + \frac{\ln\left(1 + \tanh\left(\frac{x}{2}\right)\right)}{1 + \tanh\left(\frac{x}{2}\right)} - \frac{2}{a} \sum_{R=\operatorname{RootOf}((a-b)Z^3 + (-3a-3b)Z^2 + (3a-3b)Z - a-b)} \frac{(-R^2 a - R^2 b - 2)}{3a}$

[In] int(tanh(x)^3/(a+b*cosh(x)^3),x,method=_RETURNVERBOSE)

[Out] 2*exp(2*x)/(1+exp(2*x))^2/a+1/a*ln(1+exp(2*x))+sum(_R*ln(exp(2*x)+(6*a^2/b*_R+2*a/b)*exp(x)+1),_R=RootOf(27*_Z^3*a^5+27*_Z^2*a^4+9*_Z*a^3+a^2-b^2))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.95 (sec) , antiderivative size = 1435, normalized size of antiderivative = 9.38

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = \text{Too large to display}$$

[In] integrate(tanh(x)^3/(a+b*cosh(x)^3),x, algorithm="fricas")

[Out]
$$-1/12*(2*(a*\cosh(x)^4 + 4*a*\cosh(x)*\sinh(x)^3 + a*\sinh(x)^4 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + a)*\sinh(x)^2 + 4*(a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)*\log(b*\cosh(x)^2 + b*\sinh(x)^2 - (a^2*\cosh(x) + a^2*\sinh(x))*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a) + 2*a*\cosh(x) + 2*(b*\cosh(x) + a)*\sinh(x) + b) - 24*\cosh(x)^2 + (6*\cosh(x)^4 + 24*\cosh(x)*\sinh(x)^3 + 6*\sinh(x)^4 + 12*(3*\cosh(x)^2 + 1)*\sinh(x)^2 - (a*\cosh(x)^4 + 4*a*\cosh(x)*\sinh(x)^3 + a*\sinh(x)^4 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + a)*\sinh(x)^2 + 4*(a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a) - 3*\sqrt{1/3}*(a*\cosh(x)^4 + 4*a*\cosh(x)*\sinh(x)^3 + a*\sinh(x)^4 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + a)*\sinh(x)^2 + 4*(a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*\sqrt{-(((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)^2*a^2 - 4*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)*a + 4)/a^2) + 12*\cosh(x)^2 + 24*(\cosh(x)^3 + \cosh(x))*\sinh(x) + 6)*\log(b*\cosh(x)^2 + b*\sinh(x)^2 + 1/2*(a^2*\cosh(x) + a^2*\sinh(x))*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a) + 3/2*\sqrt{1/3}*(a^2*\cosh(x) + a^2*\sinh(x))*\sqrt{-(((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)^2*a^2 - 4*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)*a + 4)/a^2) - a*\cosh(x) + (2*b*\cosh(x) - a)*\sinh(x) + b) + (6*\cosh(x)^4 + 24*\cosh(x)*\sinh(x)^3 + 6*\sinh(x)^4 + 12*(3*\cosh(x)^2 + 1)*\sinh(x)^2 - (a*\cosh(x)^4 + 4*a*\cosh(x)*\sinh(x)^3 + a*\sinh(x)^4 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + a)*\sinh(x)^2 + 4*(a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a) + 3*\sqrt{1/3}*(a*\cosh(x)^4 + 4*a*\cosh(x)*\sinh(x)^3 + a*\sinh(x)^4 + 2*a*\cosh(x)^2 + 2*(3*a*\cosh(x)^2 + a)*\sinh(x)^2 + 4*(a*\cosh(x)^3 + a*\cosh(x))*\sinh(x) + a)*\sqrt{-(((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)^2*a^2 - 4*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)*a + 4)/a^2) + 12*\cosh(x)^2 + 24*(\cosh(x)^3 + \cosh(x))*\sinh(x) + 6)*\log(b*\cosh(x)^2 + b*\sinh(x)^2 + 1/2*(a^2*\cosh(x) + a^2*\sinh(x))*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a) - 3/2*\sqrt{1/3}*(a^2*\cosh(x) + a^2*\sinh(x))*\sqrt{-(((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)^2*a^2 - 4*((1/2)^{(1/3)}*(I*\sqrt{3} + 1)*(1/a^3 + b^2/a^5 - (a^2 - b^2)/a^5)^{(1/3)} + 2/a)*a + 4$$

)/a^2) - a*cosh(x) + (2*b*cosh(x) - a)*sinh(x) + b) - 12*(cosh(x)^4 + 4*cosh(x)*sinh(x)^3 + sinh(x)^4 + 2*(3*cosh(x)^2 + 1)*sinh(x)^2 + 2*cosh(x)^2 + 4*(cosh(x)^3 + cosh(x))*sinh(x) + 1)*log(2*cosh(x)/(cosh(x) - sinh(x))) - 4*8*cosh(x)*sinh(x) - 24*sinh(x)^2)/(a*cosh(x)^4 + 4*a*cosh(x)*sinh(x)^3 + a*sinh(x)^4 + 2*a*cosh(x)^2 + 2*(3*a*cosh(x)^2 + a)*sinh(x)^2 + 4*(a*cosh(x)^3 + a*cosh(x))*sinh(x) + a)

Sympy [F]

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = \int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx$$

[In] integrate(tanh(x)**3/(a+b*cosh(x)**3),x)

[Out] Integral(tanh(x)**3/(a + b*cosh(x)**3), x)

Maxima [F]

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = \int \frac{\tanh(x)^3}{b \cosh(x)^3 + a} dx$$

[In] integrate(tanh(x)^3/(a+b*cosh(x)^3),x, algorithm="maxima")

[Out] 2*b*(x/(a*b) - integrate((b*e^(5*x) + 3*b*e^(3*x) + 8*a*e^(2*x) + 3*b*e^x)*e^x/(b*e^(6*x) + 3*b*e^(4*x) + 8*a*e^(3*x) + 3*b*e^(2*x) + b), x)/(a*b)) + 6*b*integrate(e^(4*x)/(b*e^(6*x) + 3*b*e^(4*x) + 8*a*e^(3*x) + 3*b*e^(2*x) + b), x)/a - 2*(x*e^(4*x) + (2*x - 1)*e^(2*x) + x)/(a*e^(4*x) + 2*a*e^(2*x) + a) + log(e^(2*x) + 1)/a + 8*integrate(e^(3*x)/(b*e^(6*x) + 3*b*e^(4*x) + 8*a*e^(3*x) + 3*b*e^(2*x) + b), x)

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.25

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = -\frac{b\left(-\frac{a}{b}\right)^{\frac{1}{3}} \log\left(\left|-2\left(-\frac{a}{b}\right)^{\frac{1}{3}} + e^{(-x)} + e^x\right|\right)}{3a^2} + \frac{\log(e^{(-x)} + e^x)}{a} - \frac{\log\left(\left|b(e^{(-x)} + e^x)^3 + 8a\right|\right)}{3a} + \frac{\sqrt{3}(-ab^2)^{\frac{1}{3}} \arctan\left(\frac{\sqrt{3}\left(\left(-\frac{a}{b}\right)^{\frac{1}{3}} + e^{(-x)} + e^x\right)}{3\left(-\frac{a}{b}\right)^{\frac{1}{3}}}\right)}{3a^2} + \frac{(-ab^2)^{\frac{1}{3}} \log\left(\left(e^{(-x)} + e^x\right)^2 + 2\left(-\frac{a}{b}\right)^{\frac{1}{3}}(e^{(-x)} + e^x) + 4\left(-\frac{a}{b}\right)^{\frac{2}{3}}\right)}{6a^2} - \frac{3(e^{(-x)} + e^x)^2 - 4}{2a(e^{(-x)} + e^x)^2}$$

[In] integrate(tanh(x)^3/(a+b*cosh(x)^3),x, algorithm="giac")

[Out] -1/3*b*(-a/b)^(1/3)*log(abs(-2*(-a/b)^(1/3) + e^(-x) + e^x))/a^2 + log(e^(-x) + e^x)/a - 1/3*log(abs(b*(e^(-x) + e^x)^3 + 8*a))/a + 1/3*sqrt(3)*(-a*b^2)^(1/3)*arctan(1/3*sqrt(3)*((-a/b)^(1/3) + e^(-x) + e^x)/((-a/b)^(1/3)))/a^2 + 1/6*(-a*b^2)^(1/3)*log((e^(-x) + e^x)^2 + 2*(-a/b)^(1/3)*(e^(-x) + e^x) + 4*(-a/b)^(2/3))/a^2 - 1/2*(3*(e^(-x) + e^x)^2 - 4)/(a*(e^(-x) + e^x)^2)

Mupad [B] (verification not implemented)

Time = 0.88 (sec) , antiderivative size = 1173, normalized size of antiderivative = 7.67

$$\int \frac{\tanh^3(x)}{a + b \cosh^3(x)} dx = \text{Too large to display}$$

[In] int(tanh(x)^3/(a + b*cosh(x)^3),x)

[Out] 2/(a + a*exp(2*x)) - 2/(a + 2*a*exp(2*x) + a*exp(4*x)) + symsum(log(-(50331648*a^6*exp(2*x) - 786432*b^6*exp(2*x) + 452984832*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^7 + 50331648*a^6 - 786432*b^6 + 1358954496*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^2*a^8 + 1358954496*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3*a^9 + 50593792*a^2*b^4 - 102498304*a^4*b^2 + 1358954496*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^2*a^8*exp(2*x) + 1358954496*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3*a^9*exp(2*x) + 50593792*a^2*b^4*exp(2*x) - 102498304*a^4*b^2*exp(2*x) + 7602176*root(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^3*b^4 - 465305600*root(27*a^5*z^3 + 27*a^4

$$\begin{aligned}
& *z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^5*b^2 + 524288*a*b^5*\exp(x) + 24379392* \\
& \text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^2*a^4*b^4 - 13833 \\
& 33888*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^2*a^6*b^2 + \\
& 18874368*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3*a^5*b \\
& ^4 - 1370750976*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3 \\
& *a^7*b^2 + 452984832*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, \\
& k)*a^7*\exp(2*x) - 5242880*a^3*b^3*\exp(x) - 524288*\text{root}(27*a^5*z^3 + 27*a^4 \\
& *z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^2*b^5*\exp(x) - 8912896*\text{root}(27*a^5*z^3 \\
& + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^4*b^3*\exp(x) + 7602176*\text{root}(27* \\
& a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^3*b^4*\exp(2*x) - 465305 \\
& 600*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)*a^5*b^2*\exp(2 \\
& *x) + 14155776*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3* \\
& a^6*b^3*\exp(x) + 24379392*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^ \\
& 2, z, k)^2*a^4*b^4*\exp(2*x) - 1383333888*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a \\
& ^3*z + a^2 - b^2, z, k)^2*a^6*b^2*\exp(2*x) + 18874368*\text{root}(27*a^5*z^3 + 27* \\
& a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3*a^5*b^4*\exp(2*x) - 1370750976*\text{root}(2 \\
& 7*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k)^3*a^7*b^2*\exp(2*x))/(3* \\
& a^6*b^6))*\text{root}(27*a^5*z^3 + 27*a^4*z^2 + 9*a^3*z + a^2 - b^2, z, k), k, 1, \\
& 3) + \log(3221225472*a^6*\exp(2*x) - 786432*b^6*\exp(2*x) + 3221225472*a^6 - 7 \\
& 86432*b^6 + 101449728*a^2*b^4 - 3321888768*a^4*b^2 + 101449728*a^2*b^4*\exp(\\
& 2*x) - 3321888768*a^4*b^2*\exp(2*x))/a
\end{aligned}$$

$$3.82 \quad \int \frac{\tanh(x)}{\sqrt{a+b \cosh^3(x)}} dx$$

Optimal result	526
Rubi [A] (verified)	526
Mathematica [A] (verified)	527
Maple [F]	528
Fricas [F(-2)]	528
Sympy [F]	528
Maxima [F]	528
Giac [F]	529
Mupad [F(-1)]	529

Optimal result

Integrand size = 15, antiderivative size = 28

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^3(x)}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^3(x)}}{\sqrt{a}}\right)}{3\sqrt{a}}$$

[Out] $-2/3*\operatorname{arctanh}((a+b*\cosh(x)^3)^{(1/2)}/a^{(1/2)})/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3309, 272, 65, 214}

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^3(x)}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^3(x)}}{\sqrt{a}}\right)}{3\sqrt{a}}$$

[In] `Int[Tanh[x]/Sqrt[a + b*Cosh[x]^3],x]`

[Out] $(-2*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^3]/\operatorname{Sqrt}[a]])/(3*\operatorname{Sqrt}[a])$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
```

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 3309

Int[((a_) + (b_)*((c_)*sin[(e_) + (f_)*(x_)])^(n_))^(p_)*tan[(e_) + (f_)*(x_)]^(m_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^(m + 1)/2), x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{1}{x\sqrt{a + bx^3}} dx, x, \cosh(x) \right) \\
 &= \frac{1}{3} \text{Subst} \left(\int \frac{1}{x\sqrt{a + bx}} dx, x, \cosh^3(x) \right) \\
 &= \frac{2 \text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a + b \cosh^3(x)} \right)}{3b} \\
 &= -\frac{2 \arctanh \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right)}{3\sqrt{a}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 28, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = -\frac{2 \arctanh \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right)}{3\sqrt{a}}$$

[In] Integrate[Tanh[x]/Sqrt[a + b*Cosh[x]^3], x]

[Out] (-2*ArcTanh[Sqrt[a + b*Cosh[x]^3]/Sqrt[a]])/(3*Sqrt[a])

Maple [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh(x)^3}} dx$$

[In] `int(tanh(x)/(a+b*cosh(x)^3)^(1/2),x)`

[Out] `int(tanh(x)/(a+b*cosh(x)^3)^(1/2),x)`

Fricas [F(-2)]

Exception generated.

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = \text{Exception raised: TypeError}$$

[In] `integrate(tanh(x)/(a+b*cosh(x)^3)^(1/2),x, algorithm="fricas")`

[Out] `Exception raised: TypeError >> Error detected within library code: failed of mode Union(SparseUnivariatePolynomial(Expression(Integer)),failed) cannot be coerced to mode SparseUnivariatePolynomial(Expression(Integer))`

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx$$

[In] `integrate(tanh(x)/(a+b*cosh(x)**3)**(1/2),x)`

[Out] `Integral(tanh(x)/sqrt(a + b*cosh(x)**3), x)`

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^3 + a}} dx$$

[In] `integrate(tanh(x)/(a+b*cosh(x)^3)^(1/2),x, algorithm="maxima")`

[Out] `integrate(tanh(x)/sqrt(b*cosh(x)^3 + a), x)`

Giac [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^3 + a}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)^3)^(1/2),x, algorithm="giac")

[Out] integrate(tanh(x)/sqrt(b*cosh(x)^3 + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^3(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^3 + a}} dx$$

[In] int(tanh(x)/(a + b*cosh(x)^3)^(1/2),x)

[Out] int(tanh(x)/(a + b*cosh(x)^3)^(1/2), x)

3.83 $\int \sqrt{a + b \cosh^3(x)} \tanh(x) dx$

Optimal result	530
Rubi [A] (verified)	530
Mathematica [A] (verified)	532
Maple [F]	532
Fricas [B] (verification not implemented)	532
Sympy [F]	534
Maxima [F]	534
Giac [F]	534
Mupad [F(-1)]	534

Optimal result

Integrand size = 15, antiderivative size = 45

$$\int \sqrt{a + b \cosh^3(x)} \tanh(x) dx = -\frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right) + \frac{2}{3} \sqrt{a + b \cosh^3(x)}$$

[Out] $-2/3*\operatorname{arctanh}((a+b*\cosh(x)^3)^{(1/2)}/a^{(1/2)})*a^{(1/2)}+2/3*(a+b*\cosh(x)^3)^{(1/2)}$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3309, 272, 52, 65, 214}

$$\int \sqrt{a + b \cosh^3(x)} \tanh(x) dx = \frac{2}{3} \sqrt{a + b \cosh^3(x)} - \frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right)$$

[In] `Int[Sqrt[a + b*Cosh[x]^3]*Tanh[x], x]`

[Out] $(-2*\operatorname{Sqrt}[a]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^3]/\operatorname{Sqrt}[a]])/3 + (2*\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^3])/3$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ`

`[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0])) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 272

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Rule 3309

`Int[((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^((m + 1)/2)], x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && ILtQ[(m - 1)/2, 0]`

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst} \left(\int \frac{\sqrt{a + bx^3}}{x} dx, x, \cosh(x) \right) \\
 &= \frac{1}{3} \text{Subst} \left(\int \frac{\sqrt{a + bx}}{x} dx, x, \cosh^3(x) \right) \\
 &= \frac{2}{3} \sqrt{a + b \cosh^3(x)} + \frac{1}{3} a \text{Subst} \left(\int \frac{1}{x \sqrt{a + bx}} dx, x, \cosh^3(x) \right) \\
 &= \frac{2}{3} \sqrt{a + b \cosh^3(x)} + \frac{(2a) \text{Subst} \left(\int \frac{1}{-\frac{a}{b} + \frac{x^2}{b}} dx, x, \sqrt{a + b \cosh^3(x)} \right)}{3b} \\
 &= -\frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right) + \frac{2}{3} \sqrt{a + b \cosh^3(x)}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 1.00

$$\int \sqrt{a + b \cosh^3(x)} \tanh(x) dx = -\frac{2}{3} \sqrt{a} \operatorname{arctanh} \left(\frac{\sqrt{a + b \cosh^3(x)}}{\sqrt{a}} \right) + \frac{2}{3} \sqrt{a + b \cosh^3(x)}$$

[In] Integrate[Sqrt[a + b*Cosh[x]^3]*Tanh[x],x]

[Out] (-2*Sqrt[a]*ArcTanh[Sqrt[a + b*Cosh[x]^3]/Sqrt[a]])/3 + (2*Sqrt[a + b*Cosh[x]^3])/3

Maple [F]

$$\int \sqrt{a + b \cosh(x)^3} \tanh(x) dx$$

[In] int((a+b*cosh(x)^3)^(1/2)*tanh(x),x)

[Out] int((a+b*cosh(x)^3)^(1/2)*tanh(x),x)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 271 vs. 2(33) = 66.

Time = 0.79 (sec) , antiderivative size = 1648, normalized size of antiderivative = 36.62

$$\int \sqrt{a + b \cosh^3(x)} \tanh(x) dx = \text{Too large to display}$$

[In] integrate((a+b*cosh(x)^3)^(1/2)*tanh(x),x, algorithm="fricas")

[Out] [1/6*(sqrt(a)*(cosh(x) + sinh(x))*log(-(b^2*cosh(x)^12 + 12*b^2*cosh(x)*sinh(x)^11 + b^2*sinh(x)^12 + 6*b^2*cosh(x)^10 + 64*a*b*cosh(x)^9 + 6*(11*b^2*cosh(x)^2 + b^2)*sinh(x)^10 + 15*b^2*cosh(x)^8 + 4*(55*b^2*cosh(x)^3 + 15*b^2*cosh(x) + 16*a*b)*sinh(x)^9 + 192*a*b*cosh(x)^7 + 3*(165*b^2*cosh(x)^4 + 90*b^2*cosh(x)^2 + 192*a*b*cosh(x) + 5*b^2)*sinh(x)^8 + 24*(33*b^2*cosh(x)^5 + 30*b^2*cosh(x)^3 + 96*a*b*cosh(x)^2 + 5*b^2*cosh(x) + 8*a*b)*sinh(x)^7 + 192*a*b*cosh(x)^5 + 4*(128*a^2 + 5*b^2)*cosh(x)^6 + 4*(231*b^2*cosh(x)^6 + 315*b^2*cosh(x)^4 + 1344*a*b*cosh(x)^3 + 105*b^2*cosh(x)^2 + 336*a*b*cosh(x) + 128*a^2 + 5*b^2)*sinh(x)^6 + 15*b^2*cosh(x)^4 + 24*(33*b^2*cosh(x)^7 + 63*b^2*cosh(x)^5 + 336*a*b*cosh(x)^4 + 35*b^2*cosh(x)^3 + 168*a*b*cosh(x)^2 + 8*a*b + (128*a^2 + 5*b^2)*cosh(x))*sinh(x)^5 + 64*a*b*cosh(x)^3 + 3*(165*b^2*cosh(x)^8 + 420*b^2*cosh(x)^6 + 2688*a*b*cosh(x)^5 + 350*b^2*cosh(x)^4 + 2240*a*b*cosh(x)^3 + 320*a*b*cosh(x) + 20*(128*a^2 + 5*b^2)*cosh(x)^2 + 5*b^2)*sinh(x)^4 + 6*b^2*cosh(x)^2 + 4*(55*b^2*cosh(x)^9 + 180*b^2*cosh(x)^7 + 180*b^2*cosh(x)^5 + 180*b^2*cosh(x)^3 + 180*b^2*cosh(x))

$$\begin{aligned}
& x)^7 + 1344*a*b*cosh(x)^6 + 210*b^2*cosh(x)^5 + 1680*a*b*cosh(x)^4 + 480*a* \\
& b*cosh(x)^2 + 20*(128*a^2 + 5*b^2)*cosh(x)^3 + 15*b^2*cosh(x) + 16*a*b)*sin \\
& h(x)^3 + 6*(11*b^2*cosh(x)^10 + 45*b^2*cosh(x)^8 + 384*a*b*cosh(x)^7 + 70*b \\
& ^2*cosh(x)^6 + 672*a*b*cosh(x)^5 + 320*a*b*cosh(x)^3 + 10*(128*a^2 + 5*b^2) \\
& *cosh(x)^4 + 15*b^2*cosh(x)^2 + 32*a*b*cosh(x) + b^2)*sinh(x)^2 + b^2 - 16* \\
& (b*cosh(x)^8 + 8*b*cosh(x)*sinh(x)^7 + b*sinh(x)^8 + 3*b*cosh(x)^6 + (28*b* \\
& cosh(x)^2 + 3*b)*sinh(x)^6 + 16*a*cosh(x)^5 + 2*(28*b*cosh(x)^3 + 9*b*cosh(\\
& x) + 8*a)*sinh(x)^5 + 3*b*cosh(x)^4 + (70*b*cosh(x)^4 + 45*b*cosh(x)^2 + 80 \\
& *a*cosh(x) + 3*b)*sinh(x)^4 + 4*(14*b*cosh(x)^5 + 15*b*cosh(x)^3 + 40*a*cos \\
& h(x)^2 + 3*b*cosh(x))*sinh(x)^3 + b*cosh(x)^2 + (28*b*cosh(x)^6 + 45*b*cosh \\
& (x)^4 + 160*a*cosh(x)^3 + 18*b*cosh(x)^2 + b)*sinh(x)^2 + 2*(4*b*cosh(x)^7 \\
& + 9*b*cosh(x)^5 + 40*a*cosh(x)^4 + 6*b*cosh(x)^3 + b*cosh(x))*sinh(x))*sqrt \\
& (a)*sqrt((b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + 3*b*cosh(x) + 4*a)/(cosh(x) \\
& ^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)) + 12*(b^2*cosh(x)^11 + 5*b^2*cosh(x)^9 \\
& + 48*a*b*cosh(x)^8 + 10*b^2*cosh(x)^7 + 112*a*b*cosh(x)^6 + 80*a*b*cosh(x) \\
& ^4 + 2*(128*a^2 + 5*b^2)*cosh(x)^5 + 5*b^2*cosh(x)^3 + 16*a*b*cosh(x)^2 + b \\
& ^2*cosh(x))*sinh(x))/(cosh(x)^12 + 12*cosh(x)*sinh(x)^11 + sinh(x)^12 + 6*(\\
& 11*cosh(x)^2 + 1)*sinh(x)^10 + 6*cosh(x)^10 + 20*(11*cosh(x)^3 + 3*cosh(x)) \\
& *sinh(x)^9 + 15*(33*cosh(x)^4 + 18*cosh(x)^2 + 1)*sinh(x)^8 + 15*cosh(x)^8 \\
& + 24*(33*cosh(x)^5 + 30*cosh(x)^3 + 5*cosh(x))*sinh(x)^7 + 4*(231*cosh(x)^6 \\
& + 315*cosh(x)^4 + 105*cosh(x)^2 + 5)*sinh(x)^6 + 20*cosh(x)^6 + 24*(33*cos \\
& h(x)^7 + 63*cosh(x)^5 + 35*cosh(x)^3 + 5*cosh(x))*sinh(x)^5 + 15*(33*cosh(x) \\
&)^8 + 84*cosh(x)^6 + 70*cosh(x)^4 + 20*cosh(x)^2 + 1)*sinh(x)^4 + 15*cosh(x) \\
&)^4 + 20*(11*cosh(x)^9 + 36*cosh(x)^7 + 42*cosh(x)^5 + 20*cosh(x)^3 + 3*cos \\
& h(x))*sinh(x)^3 + 6*(11*cosh(x)^10 + 45*cosh(x)^8 + 70*cosh(x)^6 + 50*cosh(\\
& x)^4 + 15*cosh(x)^2 + 1)*sinh(x)^2 + 6*cosh(x)^2 + 12*(cosh(x)^11 + 5*cosh(\\
& x)^9 + 10*cosh(x)^7 + 10*cosh(x)^5 + 5*cosh(x)^3 + cosh(x))*sinh(x) + 1)) + \\
& 2*sqrt((b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + 3*b*cosh(x) + 4*a)/(cosh(x)^ \\
& 2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/(cosh(x) + sinh(x)), 1/3*(sqrt(-a)*(co \\
& sh(x) + sinh(x))*arctan(8*(cosh(x)^2 + 2*cosh(x)*sinh(x) + sinh(x)^2)*sqrt(\\
& -a)*sqrt((b*cosh(x)^3 + 3*b*cosh(x)*sinh(x)^2 + 3*b*cosh(x) + 4*a)/(cosh(x) \\
& ^2 - 2*cosh(x)*sinh(x) + sinh(x)^2)))/(b*cosh(x)^6 + 6*b*cosh(x)*sinh(x)^5 + \\
& b*sinh(x)^6 + 3*b*cosh(x)^4 + 3*(5*b*cosh(x)^2 + b)*sinh(x)^4 + 16*a*cosh(\\
& x)^3 + 4*(5*b*cosh(x)^3 + 3*b*cosh(x) + 4*a)*sinh(x)^3 + 3*b*cosh(x)^2 + 3* \\
& (5*b*cosh(x)^4 + 6*b*cosh(x)^2 + 16*a*cosh(x) + b)*sinh(x)^2 + 6*(b*cosh(x) \\
& ^5 + 2*b*cosh(x)^3 + 8*a*cosh(x)^2 + b*cosh(x))*sinh(x) + b)) + sqrt((b*cos \\
& h(x)^3 + 3*b*cosh(x)*sinh(x)^2 + 3*b*cosh(x) + 4*a)/(cosh(x)^2 - 2*cosh(x)* \\
& sinh(x) + sinh(x)^2)))/(cosh(x) + sinh(x))]
\end{aligned}$$

Sympy [F]

$$\int \sqrt{a + b \cosh^3(x) \tanh(x)} dx = \int \sqrt{a + b \cosh^3(x) \tanh(x)} dx$$

[In] integrate((a+b*cosh(x)**3)**(1/2)*tanh(x),x)

[Out] Integral(sqrt(a + b*cosh(x)**3)*tanh(x), x)

Maxima [F]

$$\int \sqrt{a + b \cosh^3(x) \tanh(x)} dx = \int \sqrt{b \cosh^3(x) + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^3)^(1/2)*tanh(x),x, algorithm="maxima")

[Out] integrate(sqrt(b*cosh(x)^3 + a)*tanh(x), x)

Giac [F]

$$\int \sqrt{a + b \cosh^3(x) \tanh(x)} dx = \int \sqrt{b \cosh^3(x) + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^3)^(1/2)*tanh(x),x, algorithm="giac")

[Out] integrate(sqrt(b*cosh(x)^3 + a)*tanh(x), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cosh^3(x) \tanh(x)} dx = \int \tanh(x) \sqrt{b \cosh^3(x) + a} dx$$

[In] int(tanh(x)*(a + b*cosh(x)^3)^(1/2),x)

[Out] int(tanh(x)*(a + b*cosh(x)^3)^(1/2), x)

3.84 $\int \frac{\tanh(x)}{\sqrt{a+b \cosh^n(x)}} dx$

Optimal result	535
Rubi [A] (verified)	535
Mathematica [A] (verified)	536
Maple [A] (verified)	537
Fricas [A] (verification not implemented)	537
Sympy [F]	537
Maxima [F]	538
Giac [F]	538
Mupad [F(-1)]	538

Optimal result

Integrand size = 15, antiderivative size = 29

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^n(x)}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^n(x)}}{\sqrt{a}}\right)}{\sqrt{an}}$$

[Out] $-2*\operatorname{arctanh}((a+b*\cosh(x)^n)^{(1/2)}/a^{(1/2)})/n/a^{(1/2)}$

Rubi [A] (verified)

Time = 0.07 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {3309, 272, 65, 214}

$$\int \frac{\tanh(x)}{\sqrt{a+b \cosh^n(x)}} dx = -\frac{2\operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^n(x)}}{\sqrt{a}}\right)}{\sqrt{an}}$$

[In] $\operatorname{Int}[\operatorname{Tanh}[x]/\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^n], x]$

[Out] $(-2*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^n]/\operatorname{Sqrt}[a]])/(\operatorname{Sqrt}[a]*n)$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 3309

```
Int[((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + b*(c*ff*x)^n)^p/(1 - ff^2*x^2)^((m + 1)/2)], x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{1}{x\sqrt{a+bx^n}} dx, x, \cosh(x)\right) \\
 &= \frac{\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cosh^n(x)\right)}{n} \\
 &= \frac{2\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cosh^n(x)}\right)}{bn} \\
 &= -\frac{2\text{arctanh}\left(\frac{\sqrt{a+b\cosh^n(x)}}{\sqrt{a}}\right)}{\sqrt{an}}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(x)}{\sqrt{a+b\cosh^n(x)}} dx = -\frac{2\text{arctanh}\left(\frac{\sqrt{a+b\cosh^n(x)}}{\sqrt{a}}\right)}{\sqrt{an}}$$

```
[In] Integrate[Tanh[x]/Sqrt[a + b*Cosh[x]^n], x]
```

```
[Out] (-2*ArcTanh[Sqrt[a + b*Cosh[x]^n]/Sqrt[a]])/(Sqrt[a]*n)
```


Maple [A] (verified)

Time = 0.12 (sec) , antiderivative size = 24, normalized size of antiderivative = 0.83

method	result	size
derivativedivides	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh(x)^n}}{\sqrt{a}}\right)}{n\sqrt{a}}$	24
default	$-\frac{2 \operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh(x)^n}}{\sqrt{a}}\right)}{n\sqrt{a}}$	24

[In] `int(tanh(x)/(a+b*cosh(x)^n)^(1/2),x,method=_RETURNVERBOSE)`

[Out] `-2*arctanh((a+b*cosh(x)^n)^(1/2)/a^(1/2))/n/a^(1/2)`

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 113, normalized size of antiderivative = 3.90

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx = \left[\frac{\log\left(\frac{b \cosh(n \log(\cosh(x))) + b \sinh(n \log(\cosh(x))) - 2\sqrt{b \cosh(n \log(\cosh(x))) + b \sinh(n \log(\cosh(x))) + a}\sqrt{a+2a}}{\cosh(n \log(\cosh(x))) + \sinh(n \log(\cosh(x)))}\right)}{\sqrt{an}}, \frac{2\sqrt{-a} \arctan\left(\frac{\sqrt{a+b \cosh(x)^n}}{\sqrt{a}}\right)}{\sqrt{a}} \right]$$

[In] `integrate(tanh(x)/(a+b*cosh(x)^n)^(1/2),x, algorithm="fricas")`

[Out] `[log((b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) - 2*sqrt(b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) + a)*sqrt(a) + 2*a)/(cosh(n*log(cosh(x))) + sinh(n*log(cosh(x)))))/(sqrt(a)*n), 2*sqrt(-a)*arctan(sqrt(b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) + a)*sqrt(-a)/a)/(a*n)]`

Sympy [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx$$

[In] `integrate(tanh(x)/(a+b*cosh(x)**n)**(1/2),x)`

[Out] `Integral(tanh(x)/sqrt(a + b*cosh(x)**n), x)`

Maxima [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^n + a}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)^n)^(1/2),x, algorithm="maxima")

[Out] integrate(tanh(x)/sqrt(b*cosh(x)^n + a), x)

Giac [F]

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx = \int \frac{\tanh(x)}{\sqrt{b \cosh(x)^n + a}} dx$$

[In] integrate(tanh(x)/(a+b*cosh(x)^n)^(1/2),x, algorithm="giac")

[Out] integrate(tanh(x)/sqrt(b*cosh(x)^n + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{\tanh(x)}{\sqrt{a + b \cosh^n(x)}} dx = \int \frac{\tanh(x)}{\sqrt{a + b \cosh(x)^n}} dx$$

[In] int(tanh(x)/(a + b*cosh(x)^n)^(1/2),x)

[Out] int(tanh(x)/(a + b*cosh(x)^n)^(1/2), x)

3.85 $\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx$

Optimal result	539
Rubi [A] (verified)	539
Mathematica [A] (verified)	541
Maple [A] (verified)	541
Fricas [A] (verification not implemented)	541
Sympy [F]	542
Maxima [F]	542
Giac [F]	542
Mupad [F(-1)]	542

Optimal result

Integrand size = 15, antiderivative size = 47

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = -\frac{2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cosh^n(x)}}{\sqrt{a}}\right)}{n} + \frac{2\sqrt{a + b \cosh^n(x)}}{n}$$

[Out] $-2*\operatorname{arctanh}((a+b*\cosh(x)^n)^{(1/2)}/a^{(1/2)})*a^{(1/2)}/n+2*(a+b*\cosh(x)^n)^{(1/2)}/n$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3309, 272, 52, 65, 214}

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \frac{2\sqrt{a + b \cosh^n(x)}}{n} - \frac{2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a + b \cosh^n(x)}}{\sqrt{a}}\right)}{n}$$

[In] `Int[Sqrt[a + b*Cosh[x]^n]*Tanh[x], x]`

[Out] $(-2*\operatorname{Sqrt}[a]*\operatorname{ArcTanh}[\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^n]/\operatorname{Sqrt}[a]])/n + (2*\operatorname{Sqrt}[a + b*\operatorname{Cosh}[x]^n])/n$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n`

+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 3309

Int[((a_) + (b_.)*((c_.)*sin[(e_.) + (f_.)*(x_)])^(n_))^(p_.)*tan[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff^(m + 1)/f, Subst[Int[x^m*((a + b*(c*ff*x))^n)^p/(1 - ff^2*x^2)^((m + 1)/2)], x], x, Sin[e + f*x]/ff], x] /; FreeQ[{a, b, c, e, f, n, p}, x] && IntegerQ[(m - 1)/2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \text{Subst}\left(\int \frac{\sqrt{a+bx^n}}{x} dx, x, \cosh(x)\right) \\
 &= \frac{\text{Subst}\left(\int \frac{\sqrt{a+bx}}{x} dx, x, \cosh^n(x)\right)}{n} \\
 &= \frac{2\sqrt{a+b\cosh^n(x)}}{n} + \frac{a\text{Subst}\left(\int \frac{1}{x\sqrt{a+bx}} dx, x, \cosh^n(x)\right)}{n} \\
 &= \frac{2\sqrt{a+b\cosh^n(x)}}{n} + \frac{(2a)\text{Subst}\left(\int \frac{1}{-\frac{a}{b}+\frac{x^2}{b}} dx, x, \sqrt{a+b\cosh^n(x)}\right)}{bn} \\
 &= -\frac{2\sqrt{a}\text{arctanh}\left(\frac{\sqrt{a+b\cosh^n(x)}}{\sqrt{a}}\right)}{n} + \frac{2\sqrt{a+b\cosh^n(x)}}{n}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 45, normalized size of antiderivative = 0.96

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \frac{-2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh^n(x)}}{\sqrt{a}}\right) + 2\sqrt{a + b \cosh^n(x)}}{n}$$

[In] Integrate[Sqrt[a + b*Cosh[x]^n]*Tanh[x], x]

[Out] (-2*Sqrt[a]*ArcTanh[Sqrt[a + b*Cosh[x]^n]/Sqrt[a]] + 2*Sqrt[a + b*Cosh[x]^n])/n

Maple [A] (verified)

Time = 0.07 (sec) , antiderivative size = 38, normalized size of antiderivative = 0.81

method	result	size
derivativedivides	$\frac{2\sqrt{a+b \cosh(x)^n} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh(x)^n}}{\sqrt{a}}\right)}{n}$	38
default	$\frac{2\sqrt{a+b \cosh(x)^n} - 2\sqrt{a} \operatorname{arctanh}\left(\frac{\sqrt{a+b \cosh(x)^n}}{\sqrt{a}}\right)}{n}$	38

[In] int((a+b*cosh(x)^n)^(1/2)*tanh(x), x, method=_RETURNVERBOSE)

[Out] 1/n*(2*(a+b*cosh(x)^n)^(1/2)-2*a^(1/2)*arctanh((a+b*cosh(x)^n)^(1/2)/a^(1/2)))

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 156, normalized size of antiderivative = 3.32

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \left[\frac{\sqrt{a} \log\left(\frac{b \cosh(n \log(\cosh(x))) + b \sinh(n \log(\cosh(x))) - 2\sqrt{b \cosh(n \log(\cosh(x))) + b \sinh(n \log(\cosh(x))) + a\sqrt{a+2a}}}{\cosh(n \log(\cosh(x))) + \sinh(n \log(\cosh(x)))}\right) + 2\sqrt{b \cosh(n \log(\cosh(x))) + b \sinh(n \log(\cosh(x))) + a}}{n} \right]$$

[In] integrate((a+b*cosh(x)^n)^(1/2)*tanh(x), x, algorithm="fricas")

[Out] [(sqrt(a)*log((b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) - 2*sqrt(b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) + a)*sqrt(a) + 2*a)/(cosh(n*log(cosh(x))) + sinh(n*log(cosh(x)))))) + 2*sqrt(b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) + a))/n, 2*(sqrt(-a)*arctan(sqrt(b*cosh(n*log(cosh(x))) + b*sinh(n*log(cosh(x))) + a)))/n]

+ b*sinh(n*log(cosh(x))) + a)*sqrt(-a)/a) + sqrt(b*cosh(n*log(cosh(x))) + b
*sinh(n*log(cosh(x))) + a))/n]

Sympy [F]

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \int \sqrt{a + b \cosh^n(x)} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)**n)**(1/2)*tanh(x),x)

[Out] Integral(sqrt(a + b*cosh(x)**n)*tanh(x), x)

Maxima [F]

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \int \sqrt{b \cosh(x)^n + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^n)^(1/2)*tanh(x),x, algorithm="maxima")

[Out] integrate(sqrt(b*cosh(x)^n + a)*tanh(x), x)

Giac [F]

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \int \sqrt{b \cosh(x)^n + a} \tanh(x) dx$$

[In] integrate((a+b*cosh(x)^n)^(1/2)*tanh(x),x, algorithm="giac")

[Out] integrate(sqrt(b*cosh(x)^n + a)*tanh(x), x)

Mupad [F(-1)]

Timed out.

$$\int \sqrt{a + b \cosh^n(x)} \tanh(x) dx = \int \tanh(x) \sqrt{a + b \cosh(x)^n} dx$$

[In] int(tanh(x)*(a + b*cosh(x)^n)^(1/2),x)

[Out] int(tanh(x)*(a + b*cosh(x)^n)^(1/2), x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 543

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*   is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*   antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```



```

(expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
          If[Head[expn]===RootSum,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```



```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

#main function

```
def grade_antiderivative(result,optimal):
```

```

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

```

```

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

```

```

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

```

```

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

```

```
if str(result).find("Integral") != -1:
```

```

    grade = "F"
    grade_annotation = ""

```

```
else:
```

```
    if expnType_result <= expnType_optimal:
```

```
        if result.has(I):
```

```
            if optimal.has(I): #both result and optimal complex
```

```
                if leaf_count_result <= 2*leaf_count_optimal:
```

```

                    grade = "A"
                    grade_annotation = ""

```

```
                else:
```

```
                    grade = "B"
```

```
                    grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
```

```
            else: #result contains complex but optimal is not
```

```
                grade = "C"
```

```
                grade_annotation = "Result contains complex when optimal does not."
```

```
        else: # result do not contain complex, this assumes optimal do not as well
```

```
            if leaf_count_result <= 2*leaf_count_optimal:
```

```

                grade = "A"
                grade_annotation = ""

```

```
            else:
```

```
                grade = "B"
```

```
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
```

```
        else:
```

```
            grade = "C"
```

```
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)
```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```