

Computer Algebra Independent Integration Tests

Summer 2023 edition

6-Hyperbolic-functions/6.3-Hyperbolic-tangent/171-6.3.1-c+d-x-
 $\int \frac{1}{x^m - a + b \tanh^n x} dx$

Nasser M. Abbasi

September 5, 2023

Compiled on September 5, 2023 at 11:54am

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	45
4	Appendix	529

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [77]. This is test number [171].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (77)	0.00 (0)
Mathematica	93.51 (72)	6.49 (5)
Maple	89.61 (69)	10.39 (8)
Fricas	83.12 (64)	16.88 (13)
Maxima	81.82 (63)	18.18 (14)
Giac	58.44 (45)	41.56 (32)
Mupad	50.65 (39)	49.35 (38)
Sympy	38.96 (30)	61.04 (47)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	62.338	0.000	0.000	37.662
Mathematica	51.948	3.896	2.597	41.558
Maple	27.273	24.675	0.000	48.052
Maxima	27.273	16.883	0.000	55.844
Fricas	19.481	14.286	22.078	44.156
Giac	19.481	2.597	0.000	77.922
Sympy	1.299	0.000	0.000	98.701
Mupad	0.000	12.987	0.000	87.013

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	5	60.00	40.00	0.00
Fricas	13	0.00	0.00	100.00
Maple	8	100.00	0.00	0.00
Maxima	14	100.00	0.00	0.00
Giac	32	93.75	0.00	6.25
Mupad	38	0.00	100.00	0.00
Sympy	47	100.00	0.00	0.00

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Maple	0.20
Fricas	0.26
Rubi	0.28
Giac	0.33
Maxima	0.61
Sympy	1.25
Mupad	1.84
Mathematica	9.70

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Sympy	19.30	1.01	17.00	0.95
Mupad	61.95	1.10	20.00	1.10
Giac	108.38	1.12	22.00	1.10
Mathematica	212.74	1.25	132.50	1.12
Rubi	216.29	1.00	108.00	1.00
Maxima	265.59	5.70	155.00	2.15
Maple	271.39	1.57	102.00	1.00
Fricas	1029.47	4.64	240.00	2.26

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.




Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram shows that the percentage of solved integrals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.




Figure 1.3: Solved integrals based on leaf size distribution

1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fracas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

{4, 5, 9, 10, 14, 15, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 48, 49, 56, 57, 61, 62, 66, 67, 71, 72, 76, 77}

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi {}

Mathematica {}

Maple {}

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```
x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer. the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified. (For mma, rubi and maple only)

The following fields are present only in *Rubi Table file*

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Nasser M. Abbasi
June 27, 2023
Design-vide

CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	41

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	23
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 19, 20, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 50, 51, 52, 53, 54, 55, 58, 59, 60, 63, 64, 65, 68, 69, 70, 73, 74, 75 }

B grade { }

C grade { }

F normal fail { }

F(-1) timedout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 6, 7, 8, 12, 13, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 50, 51, 52, 53, 54, 55, 58, 59, 60, 65, 68, 69, 70, 73, 74, 75 }

B grade { 11, 63, 64 }

C grade { 18, 19 }

F normal fail { 16, 17, 20 }

F(-1) timedout fail { 22, 23 }

F(-2) exception fail { }

Maple

A grade { 8, 13, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 55, 60, 65 }

B grade { 1, 2, 3, 6, 7, 11, 12, 53, 54, 58, 59, 63, 64, 68, 69, 70, 73, 74, 75 }

C grade { }

F normal fail { 16, 17, 18, 19, 20, 50, 51, 52 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 32, 33, 34, 35, 36, 37, 40, 41, 42, 45, 46, 47, 50, 51, 52 }

B grade { 8, 38, 39, 43, 44, 68, 69, 70, 73, 74, 75 }

C grade { 1, 2, 3, 6, 7, 11, 12, 13, 53, 54, 55, 58, 59, 60, 63, 64, 65 }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28 }

Maxima

A grade { 3, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 54, 65, 73, 74 }

B grade { 1, 2, 6, 8, 11, 12, 53, 58, 59, 63, 64, 68, 69 }

C grade { }

F normal fail { 7, 13, 16, 17, 18, 19, 20, 50, 51, 52, 55, 60, 70, 75 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 32, 33, 34, 35, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47 }

B grade { 8, 36 }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 18, 20, 50, 51, 52, 53, 54, 55, 58, 59, 60, 63, 64, 65, 68, 69, 70, 73, 74, 75 }

F(-1) timeout fail { }

F(-2) exception fail { 19, 23 }

Mupad

A grade { }

B grade { 8, 32, 33, 34, 38, 39, 40, 43, 44, 45 }

C grade { }

F normal fail { }

F(-1) timedout fail { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 18, 19, 20, 35, 36, 37, 41, 42, 46, 47, 50, 51, 52, 53, 54, 55, 58, 59, 60, 63, 64, 65, 68, 69, 70, 73, 74, 75 }

F(-2) exception fail { }

Sympy

A grade { 8 }

B grade { }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 16, 17, 18, 19, 20, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 50, 51, 52, 53, 54, 55, 58, 59, 60, 63, 64, 65, 68, 69, 70, 73, 74, 75 }

F(-1) timedout fail { }

F(-2) exception fail { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	117	117	154	402	286	531	0	0	0
N.S.	1	1.00	1.32	3.44	2.44	4.54	0.00	0.00	0.00
time (sec)	N/A	0.144	0.455	0.388	0.321	0.277	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	84	84	143	242	176	329	0	0	0
N.S.	1	1.00	1.70	2.88	2.10	3.92	0.00	0.00	0.00
time (sec)	N/A	0.109	0.293	0.374	0.311	0.270	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	57	57	59	109	78	171	0	0	0
N.S.	1	1.00	1.04	1.91	1.37	3.00	0.00	0.00	0.00
time (sec)	N/A	0.061	0.153	0.128	0.316	0.261	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	42	16	12	16	16
N.S.	1	1.00	1.14	1.00	3.00	1.14	0.86	1.14	1.14
time (sec)	N/A	0.018	9.685	0.049	0.408	0.258	0.422	0.256	1.675

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	14	14	16	14	71	27	14	16	16
N.S.	1	1.00	1.14	1.00	5.07	1.93	1.00	1.14	1.14
time (sec)	N/A	0.017	24.627	0.041	0.356	0.263	0.564	0.312	1.684

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	119	119	179	336	343	1505	0	0	0
N.S.	1	1.00	1.50	2.82	2.88	12.65	0.00	0.00	0.00
time (sec)	N/A	0.142	0.572	0.220	0.369	0.285	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	88	88	115	185	0	840	0	0	0
N.S.	1	1.00	1.31	2.10	0.00	9.55	0.00	0.00	0.00
time (sec)	N/A	0.096	0.449	0.275	0.000	0.284	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	A	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	40	40	77	62	127	232	66	127	56
N.S.	1	1.00	1.92	1.55	3.18	5.80	1.65	3.18	1.40
time (sec)	N/A	0.023	0.217	0.050	0.241	0.270	0.153	0.281	0.101

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	110	18	14	18	18
N.S.	1	1.00	1.12	1.00	6.88	1.12	0.88	1.12	1.12
time (sec)	N/A	0.027	19.225	0.042	0.301	0.255	0.418	0.303	1.656

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	16	16	18	16	195	29	15	18	18
N.S.	1	1.00	1.12	1.00	12.19	1.81	0.94	1.12	1.12
time (sec)	N/A	0.027	24.677	0.041	0.350	0.258	0.601	0.381	1.677

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	237	237	496	685	595	5569	0	0	0
N.S.	1	1.00	2.09	2.89	2.51	23.50	0.00	0.00	0.00
time (sec)	N/A	0.284	6.912	0.271	0.380	0.333	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	157	226	375	392	3071	0	0	0
N.S.	1	1.00	1.44	2.39	2.50	19.56	0.00	0.00	0.00
time (sec)	N/A	0.179	2.999	0.239	0.378	0.292	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	100	100	97	166	0	1462	0	0	0
N.S.	1	1.00	0.97	1.66	0.00	14.62	0.00	0.00	0.00
time (sec)	N/A	0.097	1.807	0.064	0.000	0.279	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	C	F	F	F(-2)	F	F(-2)	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	1280	1280	556	0	0	0	0	0	0
N.S.	1	1.00	0.43	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	0.965	3.378	0.000	0.000	0.000	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	F	F	F	F(-2)	F	F	F(-1)
verified	N/A	Yes	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	1365	1365	0	0	0	0	0	0	0
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
time (sec)	N/A	1.089	0.000	0.000	0.000	0.000	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	19	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.95	1.00	1.00
time (sec)	N/A	1.133	29.376	0.068	0.553	0.000	3.000	0.445	2.056

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	F(-1)	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	0	18	20	0	19	20	20
N.S.	1	1.00	0.00	0.90	1.00	0.00	0.95	1.00	1.00
time (sec)	N/A	0.034	0.000	0.040	0.549	0.000	1.305	0.340	1.992

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	F(-1)	N/A	N/A	F(-2)	N/A	F(-2)	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	0	18	20	0	19	0	20
N.S.	1	1.00	0.00	0.90	1.00	0.00	0.95	0.00	1.00
time (sec)	N/A	0.037	0.000	0.044	0.589	0.000	1.250	0.000	2.032

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	19	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.95	1.00	1.00
time (sec)	N/A	1.152	31.048	0.083	0.584	0.000	1.402	0.489	2.107

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	17	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.85	1.00	1.00
time (sec)	N/A	0.042	24.803	0.037	0.334	0.000	1.002	0.321	2.001

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	17	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.85	1.00	1.00
time (sec)	N/A	0.035	2.010	0.039	0.333	0.000	0.714	0.274	1.713

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	19	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.95	1.00	1.00
time (sec)	N/A	0.038	2.088	0.038	0.449	0.000	0.954	0.284	1.764

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	F(-2)	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	18	20	0	19	20	20
N.S.	1	1.00	1.10	0.90	1.00	0.00	0.95	1.00	1.00
time (sec)	N/A	0.045	21.464	0.041	0.460	0.000	1.138	0.386	1.970

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	171	14	12	14	14
N.S.	1	1.00	1.17	1.00	14.25	1.17	1.00	1.17	1.17
time (sec)	N/A	0.021	109.517	0.024	0.754	0.250	0.600	0.310	1.684

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	144	14	12	14	14
N.S.	1	1.00	1.17	1.00	12.00	1.17	1.00	1.17	1.17
time (sec)	N/A	0.021	8.317	0.021	0.676	0.248	0.471	0.284	1.704

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	100	12	10	12	12
N.S.	1	1.00	1.20	1.00	10.00	1.20	1.00	1.20	1.20
time (sec)	N/A	0.012	6.180	0.020	0.459	0.242	0.421	0.260	1.701

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	169	244	165	186	304	0	188	273
N.S.	1	1.00	1.44	0.98	1.10	1.80	0.00	1.11	1.62
time (sec)	N/A	0.132	0.622	0.233	0.270	0.253	0.000	0.268	1.979

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	122	169	103	126	192	0	119	187
N.S.	1	1.00	1.39	0.84	1.03	1.57	0.00	0.98	1.53
time (sec)	N/A	0.083	0.385	0.204	0.245	0.244	0.000	0.266	1.849

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	81	46	74	101	0	64	76
N.S.	1	1.00	1.09	0.62	1.00	1.36	0.00	0.86	1.03
time (sec)	N/A	0.037	0.511	0.183	0.213	0.253	0.000	0.269	1.774

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	157	122	61	48	73	0	47	0
N.S.	1	1.00	0.78	0.39	0.31	0.46	0.00	0.30	0.00
time (sec)	N/A	0.208	0.433	0.353	0.352	0.252	0.000	0.265	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	159	206	90	56	217	0	320	0
N.S.	1	1.00	1.30	0.57	0.35	1.36	0.00	2.01	0.00
time (sec)	N/A	0.178	0.666	0.281	0.489	0.265	0.000	0.303	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	211	264	211	68	342	0	176	0
N.S.	1	1.00	1.25	1.00	0.32	1.62	0.00	0.83	0.00
time (sec)	N/A	0.228	0.858	0.306	0.646	0.252	0.000	0.272	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	230	420	273	296	573	0	368	267
N.S.	1	1.00	1.83	1.19	1.29	2.49	0.00	1.60	1.16
time (sec)	N/A	0.185	2.018	0.298	0.508	0.257	0.000	0.269	1.854

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	207	163	190	361	0	217	165
N.S.	1	1.00	1.22	0.96	1.12	2.12	0.00	1.28	0.97
time (sec)	N/A	0.132	1.531	0.259	0.425	0.259	0.000	0.272	1.833

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	133	114	74	106	192	0	103	89
N.S.	1	1.00	0.86	0.56	0.80	1.44	0.00	0.77	0.67
time (sec)	N/A	0.101	0.991	0.253	0.316	0.244	0.000	0.266	1.790

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	297	199	106	81	133	0	75	0
N.S.	1	1.00	0.67	0.36	0.27	0.45	0.00	0.25	0.00
time (sec)	N/A	0.580	0.803	0.380	0.707	0.255	0.000	0.268	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	420	420	442	163	100	609	0	583	0
N.S.	1	1.00	1.05	0.39	0.24	1.45	0.00	1.39	0.00
time (sec)	N/A	0.548	1.484	0.431	1.348	0.260	0.000	0.308	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	336	336	615	379	407	844	0	548	376
N.S.	1	1.00	1.83	1.13	1.21	2.51	0.00	1.63	1.12
time (sec)	N/A	0.260	3.457	0.329	1.666	0.244	0.000	0.282	2.028

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	246	371	223	255	532	0	315	236
N.S.	1	1.00	1.51	0.91	1.04	2.16	0.00	1.28	0.96
time (sec)	N/A	0.172	2.115	0.303	1.190	0.248	0.000	0.283	1.927

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	183	185	102	139	286	0	142	129
N.S.	1	1.00	1.01	0.56	0.76	1.56	0.00	0.78	0.70
time (sec)	N/A	0.159	1.215	0.283	0.727	0.267	0.000	0.275	1.827

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	437	437	312	151	114	193	0	103	0
N.S.	1	1.00	0.71	0.35	0.26	0.44	0.00	0.24	0.00
time (sec)	N/A	1.279	0.959	0.301	2.488	0.262	0.000	0.276	0.000

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	692	692	794	239	140	1164	0	840	0
N.S.	1	1.00	1.15	0.35	0.20	1.68	0.00	1.21	0.00
time (sec)	N/A	1.300	3.133	0.371	5.230	0.277	0.000	0.317	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	122	37	44	22	22
N.S.	1	1.00	1.10	1.00	6.10	1.85	2.20	1.10	1.10
time (sec)	N/A	0.033	27.843	0.049	0.482	0.240	2.282	0.297	1.956

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	68	20	24	20	20
N.S.	1	1.00	1.11	1.00	3.78	1.11	1.33	1.11	1.11
time (sec)	N/A	0.020	16.542	0.038	0.296	0.242	1.551	0.270	1.935

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	89	89	116	0	0	148	0	0	0
N.S.	1	1.00	1.30	0.00	0.00	1.66	0.00	0.00	0.00
time (sec)	N/A	0.078	0.835	0.000	0.000	0.088	0.000	0.000	0.000

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	153	153	163	0	0	248	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	1.62	0.00	0.00	0.00
time (sec)	N/A	0.118	1.176	0.000	0.000	0.080	0.000	0.000	0.000

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	224	224	230	0	0	345	0	0	0
N.S.	1	1.00	1.03	0.00	0.00	1.54	0.00	0.00	0.00
time (sec)	N/A	0.171	1.387	0.000	0.000	0.096	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	137	137	161	470	304	583	0	0	0
N.S.	1	1.00	1.18	3.43	2.22	4.26	0.00	0.00	0.00
time (sec)	N/A	0.175	0.498	0.262	0.257	0.288	0.000	0.000	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	103	103	147	290	179	364	0	0	0
N.S.	1	1.00	1.43	2.82	1.74	3.53	0.00	0.00	0.00
time (sec)	N/A	0.140	0.350	0.243	0.257	0.267	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	72	129	0	191	0	0	0
N.S.	1	1.00	0.96	1.72	0.00	2.55	0.00	0.00	0.00
time (sec)	N/A	0.083	0.100	0.099	0.000	0.266	0.000	0.000	0.000

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	56	20	15	20	20
N.S.	1	1.00	1.11	1.00	3.11	1.11	0.83	1.11	1.11
time (sec)	N/A	0.021	2.930	0.054	0.241	0.242	0.806	0.262	1.763

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	87	31	17	20	20
N.S.	1	1.00	1.11	1.00	4.83	1.72	0.94	1.11	1.11
time (sec)	N/A	0.020	5.644	0.056	0.264	0.254	2.912	0.326	1.872

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	277	277	544	905	625	3744	0	0	0
N.S.	1	1.00	1.96	3.27	2.26	13.52	0.00	0.00	0.00
time (sec)	N/A	0.368	2.950	0.364	0.302	0.324	0.000	0.000	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	211	237	542	411	2123	0	0	0
N.S.	1	1.00	1.12	2.57	1.95	10.06	0.00	0.00	0.00
time (sec)	N/A	0.297	1.810	0.355	0.280	0.285	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	127	214	221	0	944	0	0	0
N.S.	1	1.00	1.69	1.74	0.00	7.43	0.00	0.00	0.00
time (sec)	N/A	0.139	6.786	0.266	0.000	0.304	0.000	0.000	0.000

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	155	36	17	22	22
N.S.	1	1.00	1.10	1.00	7.75	1.80	0.85	1.10	1.10
time (sec)	N/A	0.039	26.587	0.146	0.378	0.258	1.158	0.305	1.984

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	280	47	19	22	22
N.S.	1	1.00	1.10	1.00	14.00	2.35	0.95	1.10	1.10
time (sec)	N/A	0.035	19.531	0.145	0.422	0.259	2.056	0.443	2.184

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	566	566	2010	1834	1297	12909	0	0	0
N.S.	1	1.00	3.55	3.24	2.29	22.81	0.00	0.00	0.00
time (sec)	N/A	0.702	7.824	0.622	0.394	0.421	0.000	0.000	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	405	405	1163	1066	869	7298	0	0	0
N.S.	1	1.00	2.87	2.63	2.15	18.02	0.00	0.00	0.00
time (sec)	N/A	0.479	7.229	0.428	0.382	0.361	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	C	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	261	303	459	475	3262	0	0	0
N.S.	1	1.00	1.16	1.76	1.82	12.50	0.00	0.00	0.00
time (sec)	N/A	0.244	7.473	0.348	0.358	0.310	0.000	0.000	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	455	52	17	22	22
N.S.	1	1.00	1.10	1.00	22.75	2.60	0.85	1.10	1.10
time (sec)	N/A	0.038	32.798	0.303	0.557	0.276	1.400	0.414	1.936

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	926	63	19	22	22
N.S.	1	1.00	1.10	1.00	46.30	3.15	0.95	1.10	1.10
time (sec)	N/A	0.037	34.750	0.287	0.666	0.272	2.425	0.784	2.007

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	212	212	247	1156	528	740	0	0	0
N.S.	1	1.00	1.17	5.45	2.49	3.49	0.00	0.00	0.00
time (sec)	N/A	0.259	1.037	0.323	0.315	0.274	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	157	200	733	337	500	0	0	0
N.S.	1	1.00	1.27	4.67	2.15	3.18	0.00	0.00	0.00
time (sec)	N/A	0.207	0.650	0.336	0.313	0.273	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	108	152	357	0	306	0	0	0
N.S.	1	1.00	1.41	3.31	0.00	2.83	0.00	0.00	0.00
time (sec)	N/A	0.122	0.694	0.227	0.000	0.273	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	114	27	17	22	22
N.S.	1	1.00	1.10	1.00	5.70	1.35	0.85	1.10	1.10
time (sec)	N/A	0.042	4.900	0.065	0.367	0.264	1.027	0.273	1.817

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	199	51	19	22	22
N.S.	1	1.00	1.10	1.00	9.95	2.55	0.95	1.10	1.10
time (sec)	N/A	0.041	9.107	0.061	0.522	0.251	1.711	0.343	2.086

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	642	642	656	2683	1060	6160	0	0	0
N.S.	1	1.00	1.02	4.18	1.65	9.60	0.00	0.00	0.00
time (sec)	N/A	1.484	5.368	0.379	0.559	0.378	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	476	476	516	1605	753	3693	0	0	0
N.S.	1	1.00	1.08	3.37	1.58	7.76	0.00	0.00	0.00
time (sec)	N/A	1.138	3.554	0.338	0.523	0.345	0.000	0.000	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	196	249	661	0	1790	0	0	0
N.S.	1	1.00	1.27	3.37	0.00	9.13	0.00	0.00	0.00
time (sec)	N/A	0.209	2.637	0.311	0.000	0.307	0.000	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	468	55	19	22	22
N.S.	1	1.00	1.10	1.00	23.40	2.75	0.95	1.10	1.10
time (sec)	N/A	0.045	31.577	0.062	1.113	0.271	1.600	0.318	2.472

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	784	96	20	22	22
N.S.	1	1.00	1.10	1.00	39.20	4.80	1.00	1.10	1.10
time (sec)	N/A	0.037	28.492	0.060	2.112	0.253	3.138	0.487	2.081

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [16] had the largest ratio of [1]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	6	1.00	14	0.429
2	A	5	5	1.00	14	0.357
3	A	4	4	1.00	12	0.333
4	N/A	0	0	1.00	14	0.000
5	N/A	0	0	1.00	14	0.000
6	A	7	7	1.00	16	0.438
7	A	6	6	1.00	16	0.375
8	A	3	2	1.00	14	0.143
9	N/A	0	0	1.00	16	0.000
10	N/A	0	0	1.00	16	0.000
11	A	13	10	1.00	16	0.625
12	A	9	7	1.00	16	0.438
13	A	7	7	1.00	14	0.500
14	N/A	0	0	1.00	16	0.000
15	N/A	0	0	1.00	16	0.000
16	A	44	18	1.00	18	1.000
17	A	43	17	1.00	18	0.944
18	A	37	12	1.00	18	0.667
19	A	37	12	1.00	18	0.667
20	A	43	17	1.00	18	0.944
21	N/A	0	0	1.00	20	0.000
22	N/A	0	0	1.00	20	0.000
23	N/A	0	0	1.00	20	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
24	N/A	0	0	1.00	20	0.000
25	N/A	0	0	1.00	20	0.000
26	N/A	0	0	1.00	20	0.000
27	N/A	0	0	1.00	20	0.000
28	N/A	0	0	1.00	20	0.000
29	N/A	0	0	1.00	12	0.000
30	N/A	0	0	1.00	12	0.000
31	N/A	0	0	1.00	10	0.000
32	A	5	3	1.00	20	0.150
33	A	4	3	1.00	20	0.150
34	A	3	3	1.00	18	0.167
35	A	7	4	1.00	20	0.200
36	A	7	4	1.00	20	0.200
37	A	8	5	1.00	20	0.250
38	A	10	3	1.00	20	0.150
39	A	8	3	1.00	20	0.150
40	A	7	3	1.00	18	0.167
41	A	21	5	1.00	20	0.250
42	A	24	7	1.00	20	0.350
43	A	14	3	1.00	20	0.150
44	A	11	3	1.00	20	0.150
45	A	11	3	1.00	18	0.167
46	A	53	7	1.00	20	0.350
47	A	60	9	1.00	20	0.450
48	N/A	0	0	1.00	20	0.000
49	N/A	0	0	1.00	18	0.000
50	A	2	2	1.00	20	0.100
51	A	4	2	1.00	20	0.100
52	A	5	2	1.00	20	0.100
53	A	8	7	1.00	18	0.389
54	A	7	6	1.00	18	0.333
55	A	6	5	1.00	16	0.312
56	N/A	0	0	1.00	18	0.000

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
57	N/A	0	0	1.00	18	0.000
58	A	15	9	1.00	20	0.450
59	A	13	10	1.00	20	0.500
60	A	9	7	1.00	18	0.389
61	N/A	0	0	1.00	20	0.000
62	N/A	0	0	1.00	20	0.000
63	A	28	11	1.00	20	0.550
64	A	22	11	1.00	20	0.550
65	A	16	9	1.00	18	0.500
66	N/A	0	0	1.00	20	0.000
67	N/A	0	0	1.00	20	0.000
68	A	6	6	1.00	20	0.300
69	A	5	5	1.00	20	0.250
70	A	4	4	1.00	18	0.222
71	N/A	0	0	1.00	20	0.000
72	N/A	0	0	1.00	20	0.000
73	A	28	10	1.00	20	0.500
74	A	24	11	1.00	20	0.550
75	A	5	5	1.00	18	0.278
76	N/A	0	0	1.00	20	0.000
77	N/A	0	0	1.00	20	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int (c + dx)^3 \tanh(e + fx) dx$	48
3.2	$\int (c + dx)^2 \tanh(e + fx) dx$	54
3.3	$\int (c + dx) \tanh(e + fx) dx$	59
3.4	$\int \frac{\tanh(e+fx)}{c+dx} dx$	63
3.5	$\int \frac{\tanh(e+fx)}{(c+dx)^2} dx$	66
3.6	$\int (c + dx)^3 \tanh^2(e + fx) dx$	69
3.7	$\int (c + dx)^2 \tanh^2(e + fx) dx$	76
3.8	$\int (c + dx) \tanh^2(e + fx) dx$	81
3.9	$\int \frac{\tanh^2(e+fx)}{c+dx} dx$	85
3.10	$\int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx$	88
3.11	$\int (c + dx)^3 \tanh^3(e + fx) dx$	91
3.12	$\int (c + dx)^2 \tanh^3(e + fx) dx$	100
3.13	$\int (c + dx) \tanh^3(e + fx) dx$	108
3.14	$\int \frac{\tanh^3(e+fx)}{c+dx} dx$	114
3.15	$\int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx$	117
3.16	$\int (c + dx)(b \tanh(e + fx))^{5/2} dx$	120
3.17	$\int (c + dx)(b \tanh(e + fx))^{3/2} dx$	134
3.18	$\int (c + dx) \sqrt{b \tanh(e + fx)} dx$	148
3.19	$\int \frac{c+dx}{\sqrt{b \tanh(e+fx)}} dx$	161
3.20	$\int \frac{c+dx}{(b \tanh(e+fx))^{3/2}} dx$	174
3.21	$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx$	188
3.22	$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx$	198
3.23	$\int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx$	201
3.24	$\int \frac{(c+dx)^2}{(b \tanh(e+fx))^{3/2}} dx$	204

3.25	$\int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx$	213
3.26	$\int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx$	216
3.27	$\int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx$	219
3.28	$\int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx$	222
3.29	$\int x^m \tanh^3(a+bx) dx$	225
3.30	$\int x^m \tanh^2(a+bx) dx$	228
3.31	$\int x^m \tanh(a+bx) dx$	231
3.32	$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx$	234
3.33	$\int \frac{(c+dx)^2}{a+a \tanh(e+fx)} dx$	240
3.34	$\int \frac{c+dx}{a+a \tanh(e+fx)} dx$	245
3.35	$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx$	249
3.36	$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx$	254
3.37	$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx$	260
3.38	$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^2} dx$	266
3.39	$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^2} dx$	272
3.40	$\int \frac{c+dx}{(a+a \tanh(e+fx))^2} dx$	277
3.41	$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx$	282
3.42	$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx$	288
3.43	$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx$	296
3.44	$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx$	304
3.45	$\int \frac{c+dx}{(a+a \tanh(e+fx))^3} dx$	310
3.46	$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx$	316
3.47	$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx$	325
3.48	$\int (c+dx)^m (a+a \tanh(e+fx))^2 dx$	339
3.49	$\int (c+dx)^m (a+a \tanh(e+fx)) dx$	342
3.50	$\int \frac{(c+dx)^m}{a+a \tanh(e+fx)} dx$	345
3.51	$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^2} dx$	349
3.52	$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx$	353
3.53	$\int (c+dx)^3 (a+b \tanh(e+fx)) dx$	358
3.54	$\int (c+dx)^2 (a+b \tanh(e+fx)) dx$	365
3.55	$\int (c+dx) (a+b \tanh(e+fx)) dx$	371
3.56	$\int \frac{a+b \tanh(e+fx)}{c+dx} dx$	375
3.57	$\int \frac{a+b \tanh(e+fx)}{(c+dx)^2} dx$	378
3.58	$\int (c+dx)^3 (a+b \tanh(e+fx))^2 dx$	381
3.59	$\int (c+dx)^2 (a+b \tanh(e+fx))^2 dx$	392
3.60	$\int (c+dx) (a+b \tanh(e+fx))^2 dx$	401
3.61	$\int \frac{(a+b \tanh(e+fx))^2}{c+dx} dx$	407
3.62	$\int \frac{(a+b \tanh(e+fx))^2}{(c+dx)^2} dx$	410

3.63	$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx$	414
3.64	$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx$	428
3.65	$\int (c + dx) (a + b \tanh(e + fx))^3 dx$	440
3.66	$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx$	449
3.67	$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$	453
3.68	$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx$	457
3.69	$\int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx$	465
3.70	$\int \frac{c + dx}{a + b \tanh(e + fx)} dx$	472
3.71	$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))} dx$	477
3.72	$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))} dx$	480
3.73	$\int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx$	484
3.74	$\int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx$	500
3.75	$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx$	513
3.76	$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))^2} dx$	520
3.77	$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))^2} dx$	524

3.1 $\int (c + dx)^3 \tanh(e + fx) dx$

Optimal result	48
Rubi [A] (verified)	48
Mathematica [A] (verified)	50
Maple [B] (verified)	51
Fricas [C] (verification not implemented)	51
Sympy [F]	52
Maxima [B] (verification not implemented)	52
Giac [F]	53
Mupad [F(-1)]	53

Optimal result

Integrand size = 14, antiderivative size = 117

$$\int (c + dx)^3 \tanh(e + fx) dx = -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3d^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

[Out] $-1/4*(d*x+c)^4/d+(d*x+c)^3*\ln(1+\exp(2*f*x+2*e))/f+3/2*d*(d*x+c)^2*\text{polylog}(2, -\exp(2*f*x+2*e))/f^2-3/2*d^2*(d*x+c)*\text{polylog}(3, -\exp(2*f*x+2*e))/f^3+3/4*d^3*\text{polylog}(4, -\exp(2*f*x+2*e))/f^4$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {3799, 2221, 2611, 6744, 2320, 6724}

$$\int (c + dx)^3 \tanh(e + fx) dx = -\frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} + \frac{(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx)^4}{4d} + \frac{3d^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

[In] Int[(c + d*x)^3*Tanh[e + f*x],x]

[Out] -1/4*(c + d*x)^4/d + ((c + d*x)^3*Log[1 + E^(2*(e + f*x))])/f + (3*d*(c + d*x)^2*PolyLog[2, -E^(2*(e + f*x))]/(2*f^2) - (3*d^2*(c + d*x)*PolyLog[3, -E^(2*(e + f*x))]/(2*f^3) + (3*d^3*PolyLog[4, -E^(2*(e + f*x))]/(4*f^4)

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)^v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]], x_Symbol] := Simp[(-1)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*(-1)*e + f*fz*x))/(1 + E^(2*(-1)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 6724

Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rule 6744

Int[((e_) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(p_)]], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^

$(m - 1) * \text{PolyLog}[n + 1, d * (F^{(c * (a + b * x)))^p], x], x] / ; \text{FreeQ}\{F, a, b, c, d, e, f, n, p\}, x] \&\& \text{GtQ}[m, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{(c + dx)^4}{4d} + 2 \int \frac{e^{2(e+fx)}(c + dx)^3}{1 + e^{2(e+fx)}} dx \\
 &= -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} - \frac{(3d) \int (c + dx)^2 \log(1 + e^{2(e+fx)}) dx}{f} \\
 &= -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{(3d^2) \int (c + dx) \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
 &= -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{(3d^3) \int \text{PolyLog}(3, -e^{2(e+fx)}) dx}{2f^3} \\
 &= -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{(3d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2(e+fx)}\right)}{4f^4} \\
 &= -\frac{(c + dx)^4}{4d} + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3d^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.46 (sec) , antiderivative size = 154, normalized size of antiderivative = 1.32

$$\begin{aligned}
 \int (c + dx)^3 \tanh(e + fx) dx &= \frac{1}{4} \left(\frac{2(c + dx)^4}{d(1 + e^{2e})} + \frac{4(c + dx)^3 \log(1 + e^{-2(e+fx)})}{f} \right. \\
 &\quad \left. - \frac{3d(2f^2(c + dx)^2 \text{PolyLog}(2, -e^{-2(e+fx)}) + d(2f(c + dx) \text{PolyLog}(3, -e^{-2(e+fx)}) + d \text{PolyLog}(4, -e^{-2(e+fx)}))}{f^4} \right. \\
 &\quad \left. + x(4c^3 + 6c^2 dx + 4cd^2 x^2 + d^3 x^3) \tanh(e) \right)
 \end{aligned}$$

[In] Integrate[(c + d*x)^3*Tanh[e + f*x],x]

[Out] $((2*(c + d*x)^4)/(d*(1 + E^{(2*e)})) + (4*(c + d*x)^3*\text{Log}[1 + E^{(-2*(e + f*x))}]))/f - (3*d*(2*f^2*(c + d*x)^2*\text{PolyLog}[2, -E^{(-2*(e + f*x))}] + d*(2*f*(c + d*x)*\text{PolyLog}[3, -E^{(-2*(e + f*x))}] + d*\text{PolyLog}[4, -E^{(-2*(e + f*x))}]))) / (4 + x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*\text{Tanh}[e])/4$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 401 vs. $2(109) = 218$.

Time = 0.39 (sec) , antiderivative size = 402, normalized size of antiderivative = 3.44

method	result
risch	$-d^2 c x^3 - \frac{3dc^2 x^2}{2} + c^3 x - \frac{d^3 x^4}{4} - \frac{3d^3 e^4}{2f^4} + \frac{c^3 \ln(1+e^{2fx+2e})}{f} - \frac{2c^3 \ln(e^{fx+e})}{f} + \frac{3d^3 \text{polylog}(4, -e^{2fx+2e})}{4f^4} - \frac{3c^2 d}{f^2}$

[In] `int((d*x+c)^3*tanh(f*x+e),x,method=_RETURNVERBOSE)`

[Out] $-d^2*c*x^3-3/2*d*c^2*x^2+c^3*x-1/4*d^3*x^4-3/2/f^4*d^3*e^4+1/f*c^3*\ln(1+\exp(2*f*x+2*e))-2/f*c^3*\ln(\exp(f*x+e))+3/4*d^3*\text{polylog}(4,-\exp(2*f*x+2*e))/f^4-3/f^2*c^2*d*e^2+1/f*d^3*\ln(1+\exp(2*f*x+2*e))*x^3-3/2/f^3*c*d^2*\text{polylog}(3,-\exp(2*f*x+2*e))+3/2/f^2*d^3*\text{polylog}(2,-\exp(2*f*x+2*e))*x^2-3/2/f^3*d^3*\text{polylog}(3,-\exp(2*f*x+2*e))*x+2/f^4*e^3*d^3*\ln(\exp(f*x+e))+3/2/f^2*c^2*d*\text{polylog}(2,-\exp(2*f*x+2*e))+4/f^3*c*d^2*e^3-2/f^3*d^3*e^3*x-6/f*c^2*d*e*x+3/f*c*d^2*\ln(1+\exp(2*f*x+2*e))*x^2-6/f^3*c*e^2*d^2*\ln(\exp(f*x+e))+3/f^2*c*d^2*\text{polylog}(2,-\exp(2*f*x+2*e))*x+6/f^2*c^2*d*\ln(\exp(f*x+e))+3/f*c^2*d*\ln(1+\exp(2*f*x+2*e))*x+6/f^2*c*d^2*e^2*x+1/4/d*c^4$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 531, normalized size of antiderivative = 4.54

$$\int (c + dx)^3 \tanh(e + fx) dx = \frac{d^3 f^4 x^4 + 4cd^2 f^4 x^3 + 6c^2 d f^4 x^2 + 4c^3 f^4 x - 24d^3 \text{polylog}(4, i \cosh(fx + e) + i \sinh(fx + e)) - 24d^3 \text{polylog}(4, -i \cosh(fx + e) - i \sinh(fx + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) - \sinh(f*x + e) + I)}{d^3 f^4 x^4 + 4cd^2 f^4 x^3 + 6c^2 d f^4 x^2 + 4c^3 f^4 x - 24d^3 \text{polylog}(4, i \cosh(fx + e) + i \sinh(fx + e)) - 24d^3 \text{polylog}(4, -i \cosh(fx + e) - i \sinh(fx + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) - \sinh(f*x + e) + I)}$$

[In] `integrate((d*x+c)^3*tanh(f*x+e),x, algorithm="fricas")`

[Out] $-1/4*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x - 24*d^3*\text{polylog}(4, I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 24*d^3*\text{polylog}(4, -I*\cosh(f*x + e) - I*\sinh(f*x + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x + c^2*d*f^2)*\text{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 4*(d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2 - c^3*f^3)*\log(\cosh(f*x + e) - \sinh(f*x + e) + I)$

+ sinh(f*x + e) - I) - 4*(d^3*f^3*x^3 + 3*c*d^2*f^3*x^2 + 3*c^2*d*f^3*x + d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2)*log(I*cosh(f*x + e) + I*sinh(f*x + e) + 1) - 4*(d^3*f^3*x^3 + 3*c*d^2*f^3*x^2 + 3*c^2*d*f^3*x + d^3*e^3 - 3*c*d^2*e^2*f + 3*c^2*d*e*f^2)*log(-I*cosh(f*x + e) - I*sinh(f*x + e) + 1) + 24*(d^3*f*x + c*d^2*f)*polylog(3, I*cosh(f*x + e) + I*sinh(f*x + e)) + 24*(d^3*f*x + c*d^2*f)*polylog(3, -I*cosh(f*x + e) - I*sinh(f*x + e))/f^4

Sympy [F]

$$\int (c + dx)^3 \tanh(e + fx) dx = \int (c + dx)^3 \tanh(e + fx) dx$$

[In] integrate((d*x+c)**3*tanh(f*x+e),x)

[Out] Integral((c + d*x)**3*tanh(e + f*x), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 286 vs. 2(108) = 216.

Time = 0.32 (sec) , antiderivative size = 286, normalized size of antiderivative = 2.44

$$\begin{aligned} \int (c + dx)^3 \tanh(e + fx) dx &= \frac{1}{4} d^3 x^4 + cd^2 x^3 + \frac{3}{2} c^2 dx^2 + \frac{c^3 \log(e^{(2fx+2e)} + 1)}{2f} \\ &+ \frac{c^3 \log(e^{(-2fx-2e)} + 1)}{2f} + \frac{3(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))c^2 d}{2f^2} \\ &+ \frac{3(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))cd^2}{2f^3} \\ &+ \frac{(4f^3 x^3 \log(e^{(2fx+2e)} + 1) + 6f^2 x^2 \text{Li}_2(-e^{(2fx+2e)}) - 6fx \text{Li}_3(-e^{(2fx+2e)}) + 3\text{Li}_4(-e^{(2fx+2e)}))d^3}{3f^4} \\ &- \frac{d^3 f^4 x^4 + 4cd^2 f^4 x^3 + 6c^2 d f^4 x^2}{2f^4} \end{aligned}$$

[In] integrate((d*x+c)^3*tanh(f*x+e),x, algorithm="maxima")

[Out] 1/4*d^3*x^4 + c*d^2*x^3 + 3/2*c^2*d*x^2 + 1/2*c^3*log(e^(2*f*x + 2*e) + 1)/f + 1/2*c^3*log(e^(-2*f*x - 2*e) + 1)/f + 3/2*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))*c^2*d/f^2 + 3/2*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))*c*d^2/f^3 + 1/3*(4*f^3*x^3*log(e^(2*f*x + 2*e) + 1) + 6*f^2*x^2*dilog(-e^(2*f*x + 2*e)) - 6*f*x*polylog(3, -e^(2*f*x + 2*e)) + 3*polylog(4, -e^(2*f*x + 2*e)))*d^3/f^4 - 1/2*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2)/f^4

Giac [F]

$$\int (c + dx)^3 \tanh(e + fx) dx = \int (dx + c)^3 \tanh(fx + e) dx$$

[In] integrate((d*x+c)^3*tanh(f*x+e),x, algorithm="giac")

[Out] integrate((d*x + c)^3*tanh(f*x + e), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 \tanh(e + fx) dx = \int \tanh(e + fx) (c + dx)^3 dx$$

[In] int(tanh(e + f*x)*(c + d*x)^3,x)

[Out] int(tanh(e + f*x)*(c + d*x)^3, x)

3.2 $\int (c + dx)^2 \tanh(e + fx) dx$

Optimal result	54
Rubi [A] (verified)	54
Mathematica [A] (verified)	56
Maple [B] (verified)	56
Fricas [C] (verification not implemented)	57
Sympy [F]	57
Maxima [B] (verification not implemented)	58
Giac [F]	58
Mupad [F(-1)]	58

Optimal result

Integrand size = 14, antiderivative size = 84

$$\int (c + dx)^2 \tanh(e + fx) dx = -\frac{(c + dx)^3}{3d} + \frac{(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} + \frac{d(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3}$$

[Out] $-1/3*(d*x+c)^3/d+(d*x+c)^2*\ln(1+\exp(2*f*x+2*e))/f+d*(d*x+c)*\operatorname{polylog}(2,-\exp(2*f*x+2*e))/f^2-1/2*d^2*\operatorname{polylog}(3,-\exp(2*f*x+2*e))/f^3$

Rubi [A] (verified)

Time = 0.11 (sec) , antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {3799, 2221, 2611, 2320, 6724}

$$\int (c + dx)^2 \tanh(e + fx) dx = \frac{d(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx)^3}{3d} - \frac{d^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3}$$

[In] $\operatorname{Int}[(c + d*x)^2*\operatorname{Tanh}[e + f*x], x]$

[Out] $-1/3*(c + dx)^3/d + ((c + dx)^2*\text{Log}[1 + E^{2*(e + f*x)}])/f + (d*(c + dx)*\text{PolyLog}[2, -E^{2*(e + f*x)}])/f^2 - (d^2*\text{PolyLog}[3, -E^{2*(e + f*x)}])/(2*f^3)$

Rule 2221

Int[(((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + dx)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + dx)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^(c_)*((a_) + (b_)*(x_)))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + dx)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + dx)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 6724

Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{(c + dx)^3}{3d} + 2 \int \frac{e^{2(e+fx)}(c + dx)^2}{1 + e^{2(e+fx)}} dx \\ &= -\frac{(c + dx)^3}{3d} + \frac{(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} - \frac{(2d) \int (c + dx) \log(1 + e^{2(e+fx)}) dx}{f} \end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)^3}{3d} + \frac{(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \int \operatorname{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&= -\frac{(c+dx)^3}{3d} + \frac{(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&= -\frac{(c+dx)^3}{3d} + \frac{(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.29 (sec) , antiderivative size = 143, normalized size of antiderivative = 1.70

$$\begin{aligned}
&\int (c+dx)^2 \tanh(e+fx) dx \\
&= \frac{e^{2e} \left(\frac{4e^{-2e}(c+dx)^3}{d} + \frac{6(1+e^{-2e})(c+dx)^2 \log(1+e^{-2(e+fx)})}{f} - \frac{3d(1+e^{-2e})(2f(c+dx) \operatorname{PolyLog}(2, -e^{-2(e+fx)}) + d \operatorname{PolyLog}(3, -e^{-2(e+fx)}))}{f^3} \right)}{6(1+e^{2e})} \\
&\quad + \frac{1}{3} x (3c^2 + 3cdx + d^2x^2) \tanh(e)
\end{aligned}$$

[In] Integrate[(c + d*x)^2*Tanh[e + f*x], x]

[Out] (E^(2*e)*((4*(c + d*x)^3)/(d*E^(2*e)) + (6*(1 + E^(-2*e))*(c + d*x)^2*Log[1 + E^(-2*(e + f*x))])/f - (3*d*(1 + E^(-2*e))*(2*f*(c + d*x)*PolyLog[2, -E^(-2*(e + f*x))] + d*PolyLog[3, -E^(-2*(e + f*x))])/f^3))/(6*(1 + E^(2*e))) + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*Tanh[e])/3

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(80) = 160.

Time = 0.37 (sec) , antiderivative size = 242, normalized size of antiderivative = 2.88

method	result
risch	$-\frac{d^2x^3}{3} - dcx^2 + c^2x + \frac{c^3}{3d} + \frac{4ced \ln(e^{fx+e})}{f^2} - \frac{4dce x}{f} + \frac{2dc \ln(1+e^{2fx+2e})x}{f} + \frac{dc \operatorname{polylog}(2, -e^{2fx+2e})}{f^2} + \frac{2d^2e^2x}{f^2} +$

[In] int((d*x+c)^2*tanh(f*x+e), x, method=_RETURNVERBOSE)


```
[Out] -1/3*d^2*x^3-d*c*x^2+c^2*x+1/3/d*c^3+4/f^2*c*e*d*ln(exp(f*x+e))-4/f*d*c*e*x
+2/f*d*c*ln(1+exp(2*f*x+2*e))*x+1/f^2*d*c*polylog(2,-exp(2*f*x+2*e))+2/f^2*
d^2*e^2*x+1/f^2*d^2*polylog(2,-exp(2*f*x+2*e))*x-2/f^2*d*c*e^2-2/f^3*e^2*d^
2*ln(exp(f*x+e))+4/3/f^3*d^2*e^3+1/f*d^2*ln(1+exp(2*f*x+2*e))*x^2-1/2*d^2*p
olylog(3,-exp(2*f*x+2*e))/f^3+1/f*c^2*ln(1+exp(2*f*x+2*e))-2/f*c^2*ln(exp(f
*x+e))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 329, normalized size of antiderivative = 3.92

$$\int (c + dx)^2 \tanh(e + fx) dx = \frac{d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x + 6 d^2 \operatorname{polylog}(3, i \cosh(fx + e) + i \sinh(fx + e)) + 6 d^2 \operatorname{polylog}(3, -i \cosh(fx + e) - i \sinh(fx + e))}{f^3}$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e),x, algorithm="fricas")
```

```
[Out] -1/3*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x + 6*d^2*polylog(3, I*cosh(f
*x + e) + I*sinh(f*x + e)) + 6*d^2*polylog(3, -I*cosh(f*x + e) - I*sinh(f*x
+ e)) - 6*(d^2*f*x + c*d*f)*dilog(I*cosh(f*x + e) + I*sinh(f*x + e)) - 6*(
d^2*f*x + c*d*f)*dilog(-I*cosh(f*x + e) - I*sinh(f*x + e)) - 3*(d^2*e^2 - 2
*c*d*e*f + c^2*f^2)*log(cosh(f*x + e) + sinh(f*x + e) + I) - 3*(d^2*e^2 - 2
*c*d*e*f + c^2*f^2)*log(cosh(f*x + e) + sinh(f*x + e) - I) - 3*(d^2*f^2*x^2
+ 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*log(I*cosh(f*x + e) + I*sinh(f*x + e)
+ 1) - 3*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*log(-I*cosh(f*x
+ e) - I*sinh(f*x + e) + 1))/f^3
```

Sympy [F]

$$\int (c + dx)^2 \tanh(e + fx) dx = \int (c + dx)^2 \tanh(e + fx) dx$$

```
[In] integrate((d*x+c)**2*tanh(f*x+e),x)
```

```
[Out] Integral((c + d*x)**2*tanh(e + f*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 176 vs. $2(79) = 158$.

Time = 0.31 (sec) , antiderivative size = 176, normalized size of antiderivative = 2.10

$$\begin{aligned} & \int (c + dx)^2 \tanh(e + fx) dx \\ &= \frac{1}{3} d^2 x^3 + cdx^2 + \frac{c^2 \log(e^{(2fx+2e)} + 1)}{2f} + \frac{c^2 \log(e^{(-2fx-2e)} + 1)}{2f} \\ &+ \frac{(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))cd}{f^2} \\ &+ \frac{(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))d^2}{2f^3} \\ &- \frac{2(d^2 f^3 x^3 + 3cdf^3 x^2)}{3f^3} \end{aligned}$$

[In] integrate((d*x+c)^2*tanh(f*x+e),x, algorithm="maxima")

[Out] $\frac{1}{3}d^2x^3 + cdx^2 + \frac{1}{2}c^2 \frac{\log(e^{(2fx+2e)} + 1)}{f} + \frac{1}{2}c^2 \frac{\log(e^{(-2fx-2e)} + 1)}{f} + \frac{(2fx \log(e^{(2fx+2e)} + 1) + \text{dilog}(-e^{(2fx+2e)}))cd}{f^2} + \frac{1}{2} \frac{(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{dilog}(-e^{(2fx+2e)}) - \text{polylog}(3, -e^{(2fx+2e)}))d^2}{f^3} - \frac{2}{3} \frac{(d^2 f^3 x^3 + 3cdf^3 x^2)}{f^3}$

Giac [F]

$$\int (c + dx)^2 \tanh(e + fx) dx = \int (dx + c)^2 \tanh(fx + e) dx$$

[In] integrate((d*x+c)^2*tanh(f*x+e),x, algorithm="giac")

[Out] integrate((d*x + c)^2*tanh(f*x + e), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \tanh(e + fx) dx = \int \tanh(e + fx) (c + dx)^2 dx$$

[In] int(tanh(e + f*x)*(c + d*x)^2,x)

[Out] int(tanh(e + f*x)*(c + d*x)^2, x)

3.3 $\int (c + dx) \tanh(e + fx) dx$

Optimal result	59
Rubi [A] (verified)	59
Mathematica [A] (verified)	60
Maple [B] (verified)	61
Fricas [C] (verification not implemented)	61
Sympy [F]	61
Maxima [A] (verification not implemented)	62
Giac [F]	62
Mupad [F(-1)]	62

Optimal result

Integrand size = 12, antiderivative size = 57

$$\int (c + dx) \tanh(e + fx) dx = -\frac{(c + dx)^2}{2d} + \frac{(c + dx) \log(1 + e^{2(e+fx)})}{f} + \frac{d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2}$$

[Out] $-1/2*(d*x+c)^2/d+(d*x+c)*\ln(1+\exp(2*f*x+2*e))/f+1/2*d*\operatorname{polylog}(2,-\exp(2*f*x+2*e))/f^2$

Rubi [A] (verified)

Time = 0.06 (sec) , antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3799, 2221, 2317, 2438}

$$\int (c+dx) \tanh(e+fx) dx = \frac{(c + dx) \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx)^2}{2d} + \frac{d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2}$$

[In] $\operatorname{Int}[(c + d*x)*\operatorname{Tanh}[e + f*x], x]$

[Out] $-1/2*(c + d*x)^2/d + ((c + d*x)*\operatorname{Log}[1 + E^{(2*(e + f*x))}])/f + (d*\operatorname{PolyLog}[2, -E^{(2*(e + f*x))}])/(2*f^2)$

Rule 2221

$\operatorname{Int}[(((F_)^{((g_)*(e_) + (f_)*(x_))})^{(n_)*((c_) + (d_)*(x_))^{(m_))}/((a_) + (b_)*((F_)^{((g_)*(e_) + (f_)*(x_))})^{(n_)}), x_Symbol] \rightarrow \operatorname{Simp} [((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^{(g*(e + f*x))})^n/a)], x] - \operatorname{Dist}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^{(m-1)}*\operatorname{Log}[1 + b*((F^{(g*(e + f*x))})^n/a)], x]$

)^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3799

Int[((c_.) + (d_.)*(x_)^(m_.))*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{(c + dx)^2}{2d} + 2 \int \frac{e^{2(e+fx)}(c + dx)}{1 + e^{2(e+fx)}} dx \\
 &= -\frac{(c + dx)^2}{2d} + \frac{(c + dx) \log(1 + e^{2(e+fx)})}{f} - \frac{d \int \log(1 + e^{2(e+fx)}) dx}{f} \\
 &= -\frac{(c + dx)^2}{2d} + \frac{(c + dx) \log(1 + e^{2(e+fx)})}{f} - \frac{d \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} \\
 &= -\frac{(c + dx)^2}{2d} + \frac{(c + dx) \log(1 + e^{2(e+fx)})}{f} + \frac{d \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.15 (sec) , antiderivative size = 59, normalized size of antiderivative = 1.04

$$\begin{aligned}
 &\int (c + dx) \tanh(e + fx) dx \\
 &= \frac{f(df x^2 + 2dx \log(1 + e^{-2(e+fx)}) + 2c \log(\cosh(e + fx))) - d \text{PolyLog}(2, -e^{-2(e+fx)})}{2f^2}
 \end{aligned}$$

[In] Integrate[(c + d*x)*Tanh[e + f*x],x]

[Out] (f*(d*f*x^2 + 2*d*x*Log[1 + E^(-2*(e + f*x))] + 2*c*Log[Cosh[e + f*x]]) - d*PolyLog[2, -E^(-2*(e + f*x))])/(2*f^2)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 108 vs. 2(53) = 106.

Time = 0.13 (sec) , antiderivative size = 109, normalized size of antiderivative = 1.91

method	result
risch	$-\frac{dx^2}{2} + cx + \frac{c \ln(1+e^{2fx+2e})}{f} - \frac{2c \ln(e^{fx+e})}{f} - \frac{2dex}{f} - \frac{de^2}{f^2} + \frac{d \ln(1+e^{2fx+2e})x}{f} + \frac{d \operatorname{polylog}(2, -e^{2fx+2e})}{2f^2} + \frac{2ed \ln}{f^2}$

[In] int((d*x+c)*tanh(f*x+e),x,method=_RETURNVERBOSE)

[Out]
$$-1/2*d*x^2+c*x+1/f*c*\ln(1+\exp(2*f*x+2*e))-2/f*c*\ln(\exp(f*x+e))-2/f*d*e*x-1/f^2*d*e^2+1/f*d*\ln(1+\exp(2*f*x+2*e))*x+1/2*d*polylog(2,-\exp(2*f*x+2*e))/f^2+2/f^2*e*d*\ln(\exp(f*x+e))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.26 (sec) , antiderivative size = 171, normalized size of antiderivative = 3.00

$$\int (c + dx) \tanh(e + fx) dx = \frac{df^2x^2 + 2cf^2x - 2d\operatorname{Li}_2(i \cosh(fx + e) + i \sinh(fx + e)) - 2d\operatorname{Li}_2(-i \cosh(fx + e) - i \sinh(fx + e))}{f^2}$$

[In] integrate((d*x+c)*tanh(f*x+e),x, algorithm="fricas")

[Out]
$$-1/2*(d*f^2*x^2 + 2*c*f^2*x - 2*d*dilog(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 2*d*dilog(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 2*(d*e - c*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 2*(d*e - c*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) - 2*(d*f*x + d*e)*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) - 2*(d*f*x + d*e)*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1))/f^2$$

Sympy [F]

$$\int (c + dx) \tanh(e + fx) dx = \int (c + dx) \tanh(e + fx) dx$$

[In] integrate((d*x+c)*tanh(f*x+e),x)

[Out] Integral((c + d*x)*tanh(e + f*x), x)

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 78, normalized size of antiderivative = 1.37

$$\int (c + dx) \tanh(e + fx) dx = -\frac{1}{2} dx^2 + \frac{c \log(e^{(2fx+2e)} + 1)}{2f} + \frac{c \log(e^{(-2fx-2e)} + 1)}{2f} + \frac{(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))d}{2f^2}$$

[In] integrate((d*x+c)*tanh(f*x+e),x, algorithm="maxima")

```
[Out] -1/2*d*x^2 + 1/2*c*log(e^(2*f*x + 2*e) + 1)/f + 1/2*c*log(e^(-2*f*x - 2*e) + 1)/f + 1/2*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))*d/f^2
```

Giac [F]

$$\int (c + dx) \tanh(e + fx) dx = \int (dx + c) \tanh(fx + e) dx$$

[In] integrate((d*x+c)*tanh(f*x+e),x, algorithm="giac")

[Out] integrate((d*x + c)*tanh(f*x + e), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx) \tanh(e + fx) dx = \int \tanh(e + fx) (c + dx) dx$$

[In] int(tanh(e + f*x)*(c + d*x),x)

[Out] int(tanh(e + f*x)*(c + d*x), x)

3.4 $\int \frac{\tanh(e+fx)}{c+dx} dx$

Optimal result	63
Rubi [N/A]	63
Mathematica [N/A]	64
Maple [N/A] (verified)	64
Fricas [N/A]	64
Sympy [N/A]	64
Maxima [N/A]	65
Giac [N/A]	65
Mupad [N/A]	65

Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{\tanh(e+fx)}{c+dx} dx = \text{Int}\left(\frac{\tanh(e+fx)}{c+dx}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)/(d*x+c), x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh(e+fx)}{c+dx} dx = \int \frac{\tanh(e+fx)}{c+dx} dx$$

[In] Int[Tanh[e + f*x]/(c + d*x), x]

[Out] Defer[Int][Tanh[e + f*x]/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{\tanh(e+fx)}{c+dx} dx$$

Mathematica [N/A]

Not integrable

Time = 9.68 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(e + fx)}{c + dx} dx$$

[In] Integrate[Tanh[e + f*x]/(c + d*x),x]

[Out] Integrate[Tanh[e + f*x]/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(fx + e)}{dx + c} dx$$

[In] int(tanh(f*x+e)/(d*x+c),x)

[Out] int(tanh(f*x+e)/(d*x+c),x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)}{dx + c} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c),x, algorithm="fricas")

[Out] integral(tanh(f*x + e)/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 12, normalized size of antiderivative = 0.86

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(e + fx)}{c + dx} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c),x)

[Out] Integral(tanh(e + f*x)/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 42, normalized size of antiderivative = 3.00

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)}{dx + c} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c),x, algorithm="maxima")

[Out] log(d*x + c)/d - 2*integrate(1/(d*x + (d*x*e^(2*e) + c*e^(2*e))*e^(2*f*x) + c), x)

Giac [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)}{dx + c} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c),x, algorithm="giac")

[Out] integrate(tanh(f*x + e)/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.68 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{c + dx} dx = \int \frac{\tanh(e + fx)}{c + dx} dx$$

[In] int(tanh(e + f*x)/(c + d*x),x)

[Out] int(tanh(e + f*x)/(c + d*x), x)

3.5 $\int \frac{\tanh(e+fx)}{(c+dx)^2} dx$

Optimal result	66
Rubi [N/A]	66
Mathematica [N/A]	67
Maple [N/A] (verified)	67
Fricas [N/A]	67
Sympy [N/A]	67
Maxima [N/A]	68
Giac [N/A]	68
Mupad [N/A]	68

Optimal result

Integrand size = 14, antiderivative size = 14

$$\int \frac{\tanh(e+fx)}{(c+dx)^2} dx = \text{Int}\left(\frac{\tanh(e+fx)}{(c+dx)^2}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh(e+fx)}{(c+dx)^2} dx = \int \frac{\tanh(e+fx)}{(c+dx)^2} dx$$

[In] Int[Tanh[e + f*x]/(c + d*x)^2,x]

[Out] Defer[Int][Tanh[e + f*x]/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tanh(e+fx)}{(c+dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 24.63 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(e + fx)}{(c + dx)^2} dx$$

[In] Integrate[Tanh[e + f*x]/(c + d*x)^2,x]

[Out] Integrate[Tanh[e + f*x]/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(fx + e)}{(dx + c)^2} dx$$

[In] int(tanh(f*x+e)/(d*x+c)^2,x)

[Out] int(tanh(f*x+e)/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.93

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c)^2,x, algorithm="fricas")

[Out] integral(tanh(f*x + e)/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(e + fx)}{(c + dx)^2} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c)**2,x)

[Out] Integral(tanh(e + f*x)/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 71, normalized size of antiderivative = 5.07

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c)^2,x, algorithm="maxima")

[Out] -1/(d^2*x + c*d) - 2*integrate(1/(d^2*x^2 + 2*c*d*x + c^2 + (d^2*x^2*e^(2*e) + 2*c*d*x*e^(2*e) + c^2*e^(2*e))*e^(2*f*x)), x)

Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)/(d*x+c)^2,x, algorithm="giac")

[Out] integrate(tanh(f*x + e)/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 1.68 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.14

$$\int \frac{\tanh(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(e + fx)}{(c + dx)^2} dx$$

[In] int(tanh(e + f*x)/(c + d*x)^2,x)

[Out] int(tanh(e + f*x)/(c + d*x)^2, x)

3.6 $\int (c + dx)^3 \tanh^2(e + fx) dx$

Optimal result	69
Rubi [A] (verified)	69
Mathematica [A] (verified)	72
Maple [B] (verified)	72
Fricas [C] (verification not implemented)	73
Sympy [F]	74
Maxima [B] (verification not implemented)	74
Giac [F]	75
Mupad [F(-1)]	75

Optimal result

Integrand size = 16, antiderivative size = 119

$$\int (c + dx)^3 \tanh^2(e + fx) dx = -\frac{(c + dx)^3}{f} + \frac{(c + dx)^4}{4d} + \frac{3d(c + dx)^2 \log(1 + e^{2(e+fx)})}{f^2} + \frac{3d^2(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} - \frac{3d^3 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^4} - \frac{(c + dx)^3 \tanh(e + fx)}{f}$$

[Out] $-(d*x+c)^3/f+1/4*(d*x+c)^4/d+3*d*(d*x+c)^2*\ln(1+\exp(2*f*x+2*e))/f^2+3*d^2*(d*x+c)*\text{polylog}(2,-\exp(2*f*x+2*e))/f^3-3/2*d^3*\text{polylog}(3,-\exp(2*f*x+2*e))/f^4-(d*x+c)^3*\tanh(f*x+e)/f$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 119, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {3801, 3799, 2221, 2611, 2320, 6724, 32}

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \frac{3d^2(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3d(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f^2} - \frac{(c + dx)^3 \tanh(e + fx)}{f} - \frac{(c + dx)^3}{f} + \frac{(c + dx)^4}{4d} - \frac{3d^3 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^4}$$

[In] Int[(c + d*x)^3*Tanh[e + f*x]^2,x]

[Out] $-\left(\frac{c + dx}{f}\right)^3 + \frac{(c + dx)^4}{4d} + \frac{3d(c + dx)^2 \log[1 + E^{2(e + fx)}]}{f^2} + \frac{3d^2(c + dx) \text{PolyLog}[2, -E^{2(e + fx)}]}{f^3} - \frac{3d^3 \text{PolyLog}[3, -E^{2(e + fx)}]}{(2f)^4} - \frac{(c + dx)^3 \tanh[e + fx]}{f}$

Rule 32

$\text{Int}[(a + bx)^m, x] := \text{Simp}[(a + bx)^{m+1}/(b(m+1)), x] /;$ FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 2221

$\text{Int}[(F^{(g)(e+fx)})^n (c + dx)^m / ((a + bx)^n \log[F]), x] := \text{Simp}[(c + dx)^m / (bfgn \log[F]) \log[1 + b(F^{g(e+fx)})^n/a], x] - \text{Dist}[d(m/(bfgn \log[F])), \text{Int}[(c + dx)^{m-1} \log[1 + b(F^{g(e+fx)})^n/a], x], x] /;$ FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

$\text{Int}[u, x] := \text{With}[v = \text{FunctionOfExponential}[u, x], \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x] /;$ FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*(a_)*(v_)^n]^m /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^{(c_)*(a_ + b_)*x}]*F[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]

Rule 2611

$\text{Int}[\log[1 + (e + fx)^n (c + dx)^m] * ((f + gx)^m \text{PolyLog}[2, (-e) * (F^{c(a+bx)})^n] / (b*c*n \log[F])), x] + \text{Dist}[g*(m/(b*c*n \log[F])), \text{Int}[(f + gx)^{m-1} \text{PolyLog}[2, (-e) * (F^{c(a+bx)})^n], x], x] /;$ FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

$\text{Int}[(c + dx)^m \tan[e + (f + gx)] * \text{Complex}[0, fz] * (f + gx), x] := \text{Simp}[(-I) * (c + dx)^{m+1} / (d(m+1)), x] + \text{Dist}[2I, \text{Int}[(c + dx)^m * (E^{2((-I)e + f*fz*x)}) / (1 + E^{2((-I)e + f*fz*x)})], x] /;$ FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3801

$\text{Int}[(c + dx)^m * (b \tan[e + fx])^n, x] := \text{Simp}[b*(c + dx)^m * (b \tan[e + fx])^{n-1} / (f*(n-1)), x] + (-\text{Dist}[b*d*(m/(f*(n-1))), \text{Int}[(c + dx)^{m-1} * (b \tan[e + fx])^{n-1}, x], x] - \text{Dist}[b^2, \text{Int}[(c + dx)^m * (b \tan[e + fx])^{n-2}, x], x]) /;$ FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c+dx)^3 \tanh(e+fx)}{f} + \frac{(3d) \int (c+dx)^2 \tanh(e+fx) dx}{f} + \int (c+dx)^3 dx \\
&= -\frac{(c+dx)^3}{f} + \frac{(c+dx)^4}{4d} - \frac{(c+dx)^3 \tanh(e+fx)}{f} + \frac{(6d) \int \frac{e^{2(e+fx)}(c+dx)^2}{1+e^{2(e+fx)}} dx}{f} \\
&= -\frac{(c+dx)^3}{f} + \frac{(c+dx)^4}{4d} + \frac{3d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} \\
&\quad - \frac{(c+dx)^3 \tanh(e+fx)}{f} - \frac{(6d^2) \int (c+dx) \log(1+e^{2(e+fx)}) dx}{f^2} \\
&= -\frac{(c+dx)^3}{f} + \frac{(c+dx)^4}{4d} + \frac{3d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3d^2(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
&\quad - \frac{(c+dx)^3 \tanh(e+fx)}{f} - \frac{(3d^3) \int \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^3} \\
&= -\frac{(c+dx)^3}{f} + \frac{(c+dx)^4}{4d} + \frac{3d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3d^2(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} - \frac{(c+dx)^3 \tanh(e+fx)}{f} \\
&\quad - \frac{(3d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&= -\frac{(c+dx)^3}{f} + \frac{(c+dx)^4}{4d} + \frac{3d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3d^2(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
&\quad - \frac{3d^3 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^4} - \frac{(c+dx)^3 \tanh(e+fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.57 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.50

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \frac{1}{4} \left(x(4c^3 + 6c^2 dx + 4cd^2 x^2 + d^3 x^3) + \frac{2de^{2e} \left(\frac{4e^{-2e}(c+dx)^3}{d} + \frac{6(1+e^{-2e})(c+dx)^2 \log(1+e^{-2(e+fx)})}{f} - \frac{3d(1+e^{-2e})(2f(c+dx) \operatorname{PolyLog}(2, -e^{-2(e+fx)}) + d \operatorname{PolyLog}(3, -e^{-2(e+fx)})}{f^3} \right)}{(1+e^{2e})f} - \frac{4(c+dx)^3 \operatorname{sech}(e) \operatorname{sech}(e+fx) \sinh(fx)}{f} \right)$$

[In] Integrate[(c + d*x)^3*Tanh[e + f*x]^2,x]

[Out] (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3) + (2*d*E^(2*e))*((4*(c + d*x)^3)/(d*E^(2*e)) + (6*(1 + E^(-2*e))*(c + d*x)^2*Log[1 + E^(-2*(e + f*x))])/f - (3*d*(1 + E^(-2*e))*(2*f*(c + d*x)*PolyLog[2, -E^(-2*(e + f*x))] + d*PolyLog[3, -E^(-2*(e + f*x))])/f^3))/((1 + E^(2*e))*f) - (4*(c + d*x)^3*Sech[e]*Sech[e + f*x]*Sinh[f*x])/f)/4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 335 vs. 2(115) = 230.

Time = 0.22 (sec) , antiderivative size = 336, normalized size of antiderivative = 2.82

method	result
risch	$\frac{d^3 x^4}{4} + d^2 c x^3 + \frac{3d c^2 x^2}{2} + c^3 x + \frac{c^4}{4d} + \frac{2d^3 x^3 + 6c d^2 x^2 + 6c^2 dx + 2c^3}{f(1+e^{2fx+2e})} - \frac{3d^3 \operatorname{polylog}(3, -e^{2fx+2e})}{2f^4} - \frac{2d^3 x^3}{f} + \frac{4d^3 e^3}{f^4} + \dots$

[In] int((d*x+c)^3*tanh(f*x+e)^2,x,method=_RETURNVERBOSE)

[Out] 1/4*d^3*x^4+d^2*c*x^3+3/2*d*c^2*x^2+c^3*x+1/4/d*c^4+2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/f/(1+exp(2*f*x+2*e))-3/2*d^3*polylog(3,-exp(2*f*x+2*e))/f^4-2/f*d^3*x^3+4/f^4*d^3*e^3+3/f^2*d^3*ln(1+exp(2*f*x+2*e))*x^2-6/f*d^2*c*x^2-6/f^3*d^2*c*e^2+6/f^2*d^2*c*ln(1+exp(2*f*x+2*e))*x+3/f^3*d^2*c*polylog(2,-exp(2*f*x+2*e))-6/f^4*d^3*e^2*ln(exp(f*x+e))+6/f^3*d^3*e^2*x+3/f^3*d^3*polylog(2,-exp(2*f*x+2*e))*x-12/f^2*d^2*c*e*x+12/f^3*d^2*c*e*ln(exp(f*x+e))+3/f^2*d^2*c^2*ln(1+exp(2*f*x+2*e))-6/f^2*d^2*c^2*ln(exp(f*x+e))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 1505, normalized size of antiderivative = 12.65

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \text{Too large to display}$$

[In] integrate((d*x+c)^3*tanh(f*x+e)^2,x, algorithm="fricas")

[Out] $\frac{1}{4}*(d^3*f^4*x^4 + 4*c*d^2*f^4*x^3 + 6*c^2*d*f^4*x^2 + 4*c^3*f^4*x - 8*d^3*e^3 + 24*c*d^2*e^2*f - 24*c^2*d*e*f^2 + 8*c^3*f^3 + (d^3*f^4*x^4 - 8*d^3*e^3 + 24*c*d^2*e^2*f - 24*c^2*d*e*f^2 + 4*(c*d^2*f^4 - 2*d^3*f^3)*x^3 + 6*(c^2*d*f^4 - 4*c*d^2*f^3)*x^2 + 4*(c^3*f^4 - 6*c^2*d*f^3)*x)*\cosh(f*x + e)^2 + 2*(d^3*f^4*x^4 - 8*d^3*e^3 + 24*c*d^2*e^2*f - 24*c^2*d*e*f^2 + 4*(c*d^2*f^4 - 2*d^3*f^3)*x^3 + 6*(c^2*d*f^4 - 4*c*d^2*f^3)*x^2 + 4*(c^3*f^4 - 6*c^2*d*f^3)*x)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*f^4*x^4 - 8*d^3*e^3 + 24*c*d^2*e^2*f - 24*c^2*d*e*f^2 + 4*(c*d^2*f^4 - 2*d^3*f^3)*x^3 + 6*(c^2*d*f^4 - 4*c*d^2*f^3)*x^2 + 4*(c^3*f^4 - 6*c^2*d*f^3)*x)*\sinh(f*x + e)^2 + 24*(d^3*f*x + c*d^2*f + (d^3*f*x + c*d^2*f)*\cosh(f*x + e)^2 + 2*(d^3*f*x + c*d^2*f)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*f*x + c*d^2*f)*\sinh(f*x + e)^2)*\operatorname{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) + 24*(d^3*f*x + c*d^2*f + (d^3*f*x + c*d^2*f)*\cosh(f*x + e)^2 + 2*(d^3*f*x + c*d^2*f)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*f*x + c*d^2*f)*\sinh(f*x + e)^2)*\operatorname{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 12*(d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2 + (d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\cosh(f*x + e)^2 + 2*(d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\sinh(f*x + e)^2)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 12*(d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2 + (d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\cosh(f*x + e)^2 + 2*(d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*e^2 - 2*c*d^2*e*f + c^2*d*f^2)*\sinh(f*x + e)^2)*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) + 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f + (d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\cosh(f*x + e)^2 + 2*(d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\sinh(f*x + e)^2)*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) + 12*(d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f + (d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\cosh(f*x + e)^2 + 2*(d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\cosh(f*x + e)*\sinh(f*x + e) + (d^3*f^2*x^2 + 2*c*d^2*f^2*x - d^3*e^2 + 2*c*d^2*e*f)*\sinh(f*x + e)^2)*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1) - 24*(d^3*\cosh(f*x + e)^2 + 2*d^3*\cosh(f*x + e)*\sinh(f*x + e) + d^3*\sinh(f*x + e)^2 + d^3)*\operatorname{polylog}(3, I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 24*(d^3*\cosh(f*x + e)^2 + 2*d^3*\cosh(f*x + e)*\sinh(f*x + e) + d^3*\sinh(f*x + e)^2 + d^3)*\operatorname{polylog}(3, -I*\cosh(f*x + e) - I*\sinh(f*x + e)))/(f^4*\cosh(f*x + e)^2 + 2*f^4*\cosh(f*x + e)*\sinh(f*x + e) + f^4*\sinh(f*x + e)^2 + f^4)$

Sympy [F]

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \int (c + dx)^3 \tanh^2(e + fx) dx$$

[In] integrate((d*x+c)**3*tanh(f*x+e)**2,x)

[Out] Integral((c + d*x)**3*tanh(e + f*x)**2, x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 343 vs. 2(114) = 228.

Time = 0.37 (sec) , antiderivative size = 343, normalized size of antiderivative = 2.88

$$\begin{aligned} \int (c + dx)^3 \tanh^2(e + fx) dx &= c^3 \left(x + \frac{e}{f} - \frac{2}{f(e^{(-2fx-2e)} + 1)} \right) \\ &- \frac{3}{2} c^2 d \left(\frac{2xe^{(2fx+2e)}}{fe^{(2fx+2e)} + f} - \frac{fx^2 + (fx^2e^{(2e)} - 2xe^{(2e)})e^{(2fx)}}{fe^{(2fx+2e)} + f} - \frac{2 \log((e^{(2fx+2e)} + 1)e^{(-2e)})}{f^2} \right) \\ &+ \frac{3(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))cd^2}{f^3} \\ &+ \frac{d^3fx^4 + 24cd^2x^2 + 4(cd^2f + 2d^3)x^3 + (d^3fx^4e^{(2e)} + 4cd^2fx^3e^{(2e)})e^{(2fx)}}{4(fe^{(2fx+2e)} + f)} \\ &+ \frac{3(2f^2x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))d^3}{2f^4} \\ &- \frac{2(d^3f^3x^3 + 3cd^2f^3x^2)}{f^4} \end{aligned}$$

[In] integrate((d*x+c)^3*tanh(f*x+e)^2,x, algorithm="maxima")

[Out] c^3*(x + e/f - 2/(f*(e^(-2*f*x - 2*e) + 1))) - 3/2*c^2*d*(2*x*e^(2*f*x + 2*e)/(f*e^(2*f*x + 2*e) + f) - (f*x^2 + (f*x^2*e^(2*e) - 2*x*e^(2*e))*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) - 2*log((e^(2*f*x + 2*e) + 1)*e^(-2*e))/f^2 + 3*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))*c*d^2/f^3 + 1/4*(d^3*f*x^4 + 24*c*d^2*x^2 + 4*(c*d^2*f + 2*d^3)*x^3 + (d^3*f*x^4*e^(2*e) + 4*c*d^2*f*x^3*e^(2*e))*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) + 3/2*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))*d^3/f^4 - 2*(d^3*f^3*x^3 + 3*c*d^2*f^3*x^2)/f^4

Giac [F]

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \int (dx + c)^3 \tanh(fx + e)^2 dx$$

[In] integrate((d*x+c)^3*tanh(f*x+e)^2,x, algorithm="giac")

[Out] integrate((d*x + c)^3*tanh(f*x + e)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 \tanh^2(e + fx) dx = \int \tanh(e + fx)^2 (c + dx)^3 dx$$

[In] int(tanh(e + f*x)^2*(c + d*x)^3,x)

[Out] int(tanh(e + f*x)^2*(c + d*x)^3, x)

3.7 $\int (c + dx)^2 \tanh^2(e + fx) dx$

Optimal result	76
Rubi [A] (verified)	76
Mathematica [A] (verified)	78
Maple [B] (verified)	78
Fricas [C] (verification not implemented)	79
Sympy [F]	80
Maxima [F]	80
Giac [F]	80
Mupad [F(-1)]	80

Optimal result

Integrand size = 16, antiderivative size = 88

$$\int (c + dx)^2 \tanh^2(e + fx) dx = -\frac{(c + dx)^2}{f} + \frac{(c + dx)^3}{3d} + \frac{2d(c + dx) \log(1 + e^{2(e+fx)})}{f^2} + \frac{d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} - \frac{(c + dx)^2 \tanh(e + fx)}{f}$$

[Out] $-(d*x+c)^2/f+1/3*(d*x+c)^3/d+2*d*(d*x+c)*\ln(1+\exp(2*f*x+2*e))/f^2+d^2*\text{polylog}(2,-\exp(2*f*x+2*e))/f^3-(d*x+c)^2*\tanh(f*x+e)/f$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {3801, 3799, 2221, 2317, 2438, 32}

$$\int (c + dx)^2 \tanh^2(e + fx) dx = \frac{2d(c + dx) \log(e^{2(e+fx)} + 1)}{f^2} - \frac{(c + dx)^2 \tanh(e + fx)}{f} - \frac{(c + dx)^2}{f} + \frac{(c + dx)^3}{3d} + \frac{d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3}$$

[In] $\text{Int}[(c + d*x)^2*\text{Tanh}[e + f*x]^2,x]$

[Out] $-((c + d*x)^2/f) + (c + d*x)^3/(3*d) + (2*d*(c + d*x)*\text{Log}[1 + E^{2*(e + f*x)}])/f^2 + (d^2*\text{PolyLog}[2, -E^{2*(e + f*x)}])/f^3 - ((c + d*x)^2*\text{Tanh}[e + f*x])/f$

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 2221

Int[(((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3799

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3801

Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{(c + dx)^2 \tanh(e + fx)}{f} + \frac{(2d) \int (c + dx) \tanh(e + fx) dx}{f} + \int (c + dx)^2 dx \\ &= -\frac{(c + dx)^2}{f} + \frac{(c + dx)^3}{3d} - \frac{(c + dx)^2 \tanh(e + fx)}{f} + \frac{(4d) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx}{f} \end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)^2}{f} + \frac{(c+dx)^3}{3d} + \frac{2d(c+dx)\log(1+e^{2(e+fx)})}{f^2} \\
&\quad - \frac{(c+dx)^2 \tanh(e+fx)}{f} - \frac{(2d^2)\int \log(1+e^{2(e+fx)}) dx}{f^2} \\
&= -\frac{(c+dx)^2}{f} + \frac{(c+dx)^3}{3d} + \frac{2d(c+dx)\log(1+e^{2(e+fx)})}{f^2} \\
&\quad - \frac{(c+dx)^2 \tanh(e+fx)}{f} - \frac{d^2 \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{f^3} \\
&= -\frac{(c+dx)^2}{f} + \frac{(c+dx)^3}{3d} + \frac{2d(c+dx)\log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} - \frac{(c+dx)^2 \tanh(e+fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.45 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.31

$$\begin{aligned}
&\int (c+dx)^2 \tanh^2(e+fx) dx \\
&= c^2x + cdx^2 + \frac{d^2x^3}{3} + \frac{2f(c+dx)(f(c+dx)+d(1+e^{2e})\log(1+e^{-2(e+fx)}))}{1+e^{2e}} - \frac{d^2 \text{PolyLog}(2, -e^{-2(e+fx)})}{f^3} \\
&\quad - \frac{(c+dx)^2 \text{sech}(e)\text{sech}(e+fx)\sinh(fx)}{f}
\end{aligned}$$

[In] Integrate[(c + d*x)^2*Tanh[e + f*x]^2,x]

[Out] c^2*x + c*d*x^2 + (d^2*x^3)/3 + (((2*f*(c + d*x)*(f*(c + d*x) + d*(1 + E^(2*e))*Log[1 + E^(-2*(e + f*x))]))/(1 + E^(2*e)) - d^2*PolyLog[2, -E^(-2*(e + f*x))])/f^3 - ((c + d*x)^2*Sech[e]*Sech[e + f*x]*Sinh[f*x])/f

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(86) = 172.

Time = 0.28 (sec) , antiderivative size = 185, normalized size of antiderivative = 2.10

method	result
risch	$\frac{d^2x^3}{3} + dcx^2 + c^2x + \frac{c^3}{3d} + \frac{2x^2d^2+4cdx+2c^2}{f(1+e^{2fx+2e})} + \frac{2dc\ln(1+e^{2fx+2e})}{f^2} - \frac{4dc\ln(e^{fx+e})}{f^2} - \frac{2d^2x^2}{f} - \frac{4d^2ex}{f^2} - \frac{2d^2e^2}{f^3} + \dots$

[In] int((d*x+c)^2*tanh(f*x+e)^2,x,method=_RETURNVERBOSE)

```
[Out] 1/3*d^2*x^3+d*c*x^2+c^2*x+1/3/d*c^3+2*(d^2*x^2+2*c*d*x+c^2)/f/(1+exp(2*f*x+
2*e))+2/f^2*d*c*ln(1+exp(2*f*x+2*e))-4/f^2*d*c*ln(exp(f*x+e))-2*d^2*x^2/f-4
/f^2*d^2*e*x-2/f^3*d^2*e^2+2/f^2*d^2*ln(1+exp(2*f*x+2*e))*x+d^2*polylog(2,-
exp(2*f*x+2*e))/f^3+4/f^3*d^2*e*ln(exp(f*x+e))
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 840, normalized size of antiderivative = 9.55

$$\int (c + dx)^2 \tanh^2(e + fx) dx$$

$$= \frac{d^2 f^3 x^3 + 3 c d f^3 x^2 + 3 c^2 f^3 x + 6 d^2 e^2 - 12 c d e f + 6 c^2 f^2 + (d^2 f^3 x^3 + 6 d^2 e^2 - 12 c d e f + 3 (c d f^3 - 2 d^2 f^2) \cosh(fx + e) + 2 (d^2 f^3 x^3 + 6 d^2 e^2 - 12 c d e f + 3 (c d f^3 - 2 d^2 f^2) x) \cosh(fx + e)^2 + 2 (d^2 f^3 x^3 + 6 d^2 e^2 - 12 c d e f + 3 (c d f^3 - 2 d^2 f^2) x) \cosh(fx + e) \sinh(fx + e) + (d^2 f^3 x^3 + 6 d^2 e^2 - 12 c d e f + 3 (c d f^3 - 2 d^2 f^2) x) \sinh(fx + e)^2 + 6 (d^2 \cosh(fx + e)^2 + 2 d^2 \cosh(fx + e) \sinh(fx + e) + d^2 \sinh(fx + e)^2 + d^2) \operatorname{dilog}(I \cosh(fx + e) + I \sinh(fx + e)) + 6 (d^2 \cosh(fx + e)^2 + 2 d^2 \cosh(fx + e) \sinh(fx + e) + d^2 \sinh(fx + e)^2 + d^2) \operatorname{dilog}(-I \cosh(fx + e) - I \sinh(fx + e)) - 6 (d^2 e - c d f + (d^2 e - c d f) \cosh(fx + e)^2 + 2 (d^2 e - c d f) \cosh(fx + e) \sinh(fx + e) + (d^2 e - c d f) \sinh(fx + e)^2) \log(\cosh(fx + e) + \sinh(fx + e) + I) - 6 (d^2 e - c d f + (d^2 e - c d f) \cosh(fx + e)^2 + 2 (d^2 e - c d f) \cosh(fx + e) \sinh(fx + e) + (d^2 e - c d f) \sinh(fx + e)^2) \log(\cosh(fx + e) + \sinh(fx + e) - I) + 6 (d^2 f x + d^2 e + (d^2 f x + d^2 e) \cosh(fx + e)^2 + 2 (d^2 f x + d^2 e) \cosh(fx + e) \sinh(fx + e) + (d^2 f x + d^2 e) \sinh(fx + e)^2) \log(I \cosh(fx + e) + I \sinh(fx + e) + 1) + 6 (d^2 f x + d^2 e + (d^2 f x + d^2 e) \cosh(fx + e)^2 + 2 (d^2 f x + d^2 e) \cosh(fx + e) \sinh(fx + e) + (d^2 f x + d^2 e) \sinh(fx + e)^2) \log(-I \cosh(fx + e) - I \sinh(fx + e) + 1)}{f^3 \cosh(fx + e)^2 + 2 f^3 \cosh(fx + e) \sinh(fx + e) + f^3 \sinh(fx + e)^2 + f^3}$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e)^2,x, algorithm="fricas")
```

```
[Out] 1/3*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x + 6*d^2*e^2 - 12*c*d*e*f + 6
*c^2*f^2 + (d^2*f^3*x^3 + 6*d^2*e^2 - 12*c*d*e*f + 3*(c*d*f^3 - 2*d^2*f^2)*
x^2 + 3*(c^2*f^3 - 4*c*d*f^2)*x)*cosh(f*x + e)^2 + 2*(d^2*f^3*x^3 + 6*d^2*e
^2 - 12*c*d*e*f + 3*(c*d*f^3 - 2*d^2*f^2)*x^2 + 3*(c^2*f^3 - 4*c*d*f^2)*x)*
cosh(f*x + e)*sinh(f*x + e) + (d^2*f^3*x^3 + 6*d^2*e^2 - 12*c*d*e*f + 3*(c
*d*f^3 - 2*d^2*f^2)*x^2 + 3*(c^2*f^3 - 4*c*d*f^2)*x)*sinh(f*x + e)^2 + 6*(d
^2*cosh(f*x + e)^2 + 2*d^2*cosh(f*x + e)*sinh(f*x + e) + d^2*sinh(f*x + e)^2
+ d^2)*dilog(I*cosh(f*x + e) + I*sinh(f*x + e)) + 6*(d^2*cosh(f*x + e)^2 +
2*d^2*cosh(f*x + e)*sinh(f*x + e) + d^2*sinh(f*x + e)^2 + d^2)*dilog(-I*co
sh(f*x + e) - I*sinh(f*x + e)) - 6*(d^2*e - c*d*f + (d^2*e - c*d*f)*cosh(f*
x + e)^2 + 2*(d^2*e - c*d*f)*cosh(f*x + e)*sinh(f*x + e) + (d^2*e - c*d*f)*
sinh(f*x + e)^2)*log(cosh(f*x + e) + sinh(f*x + e) + I) - 6*(d^2*e - c*d*f
+ (d^2*e - c*d*f)*cosh(f*x + e)^2 + 2*(d^2*e - c*d*f)*cosh(f*x + e)*sinh(f*
x + e) + (d^2*e - c*d*f)*sinh(f*x + e)^2)*log(cosh(f*x + e) + sinh(f*x + e)
- I) + 6*(d^2*f*x + d^2*e + (d^2*f*x + d^2*e)*cosh(f*x + e)^2 + 2*(d^2*f*x
+ d^2*e)*cosh(f*x + e)*sinh(f*x + e) + (d^2*f*x + d^2*e)*sinh(f*x + e)^2)*
log(I*cosh(f*x + e) + I*sinh(f*x + e) + 1) + 6*(d^2*f*x + d^2*e + (d^2*f*x
+ d^2*e)*cosh(f*x + e)^2 + 2*(d^2*f*x + d^2*e)*cosh(f*x + e)*sinh(f*x + e)
+ (d^2*f*x + d^2*e)*sinh(f*x + e)^2)*log(-I*cosh(f*x + e) - I*sinh(f*x + e)
+ 1))/(f^3*cosh(f*x + e)^2 + 2*f^3*cosh(f*x + e)*sinh(f*x + e) + f^3*sinh(
f*x + e)^2 + f^3)
```

Sympy [F]

$$\int (c + dx)^2 \tanh^2(e + fx) dx = \int (c + dx)^2 \tanh^2(e + fx) dx$$

```
[In] integrate((d*x+c)**2*tanh(f*x+e)**2,x)
```

```
[Out] Integral((c + d*x)**2*tanh(e + f*x)**2, x)
```

Maxima [F]

$$\int (c + dx)^2 \tanh^2(e + fx) dx = \int (dx + c)^2 \tanh^2(fx + e) dx$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e)^2,x, algorithm="maxima")
```

```
[Out] c^2*(x + e/f - 2/(f*(e^(-2*f*x - 2*e) + 1))) - c*d*(2*x*e^(2*f*x + 2*e)/(f*
e^(2*f*x + 2*e) + f) - (f*x^2 + (f*x^2*e^(2*e) - 2*x*e^(2*e))*e^(2*f*x))/(f
*e^(2*f*x + 2*e) + f) - 2*log((e^(2*f*x + 2*e) + 1)*e^(-2*e))/f^2) + 1/3*d^
2*((f*x^3*e^(2*f*x + 2*e) + f*x^3 + 6*x^2)/(f*e^(2*f*x + 2*e) + f) - 12*int
egrate(x/(f*e^(2*f*x + 2*e) + f), x))
```

Giac [F]

$$\int (c + dx)^2 \tanh^2(e + fx) dx = \int (dx + c)^2 \tanh^2(fx + e) dx$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e)^2,x, algorithm="giac")
```

```
[Out] integrate((d*x + c)^2*tanh(f*x + e)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \tanh^2(e + fx) dx = \int \tanh^2(e + fx) (c + dx)^2 dx$$

```
[In] int(tanh(e + f*x)^2*(c + d*x)^2,x)
```

```
[Out] int(tanh(e + f*x)^2*(c + d*x)^2, x)
```


3.8 $\int (c + dx) \tanh^2(e + fx) dx$

Optimal result	81
Rubi [A] (verified)	81
Mathematica [A] (verified)	82
Maple [A] (verified)	82
Fricas [B] (verification not implemented)	83
Sympy [A] (verification not implemented)	83
Maxima [B] (verification not implemented)	84
Giac [B] (verification not implemented)	84
Mupad [B] (verification not implemented)	84

Optimal result

Integrand size = 14, antiderivative size = 40

$$\int (c + dx) \tanh^2(e + fx) dx = cx + \frac{dx^2}{2} + \frac{d \log(\cosh(e + fx))}{f^2} - \frac{(c + dx) \tanh(e + fx)}{f}$$

[Out] $c*x+1/2*d*x^2+d*\ln(\cosh(f*x+e))/f^2-(d*x+c)*\tanh(f*x+e)/f$

Rubi [A] (verified)

Time = 0.02 (sec) , antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {3801, 3556}

$$\int (c + dx) \tanh^2(e + fx) dx = -\frac{(c + dx) \tanh(e + fx)}{f} + cx + \frac{d \log(\cosh(e + fx))}{f^2} + \frac{dx^2}{2}$$

[In] $\text{Int}[(c + d*x)*\text{Tanh}[e + f*x]^2, x]$

[Out] $c*x + (d*x^2)/2 + (d*\text{Log}[\text{Cosh}[e + f*x]])/f^2 - ((c + d*x)*\text{Tanh}[e + f*x])/f$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3801

$\text{Int}[((c_.) + (d_.)*(x_.))^{(m_.)}*((b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(c + d*x)^m*((b*\text{Tan}[e + f*x])^{(n-1)}/(f*(n-1))), x] + (-\text{Dist}[b*d*(m/(f*(n-1))), \text{Int}[(c + d*x)^{(m-1)}*(b*\text{Tan}[e + f*x])^{(n-1)}, x], x] - \text{Dist}[b^2, \text{Int}[(c + d*x)^m*(b*\text{Tan}[e + f*x])^{(n-2)}, x], x]) /; \text{FreeQ}$

{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{(c + dx) \tanh(e + fx)}{f} + \frac{d \int \tanh(e + fx) dx}{f} + \int (c + dx) dx \\ &= cx + \frac{dx^2}{2} + \frac{d \log(\cosh(e + fx))}{f^2} - \frac{(c + dx) \tanh(e + fx)}{f} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.22 (sec) , antiderivative size = 77, normalized size of antiderivative = 1.92

$$\begin{aligned} \int (c + dx) \tanh^2(e + fx) dx &= \frac{\text{arctanh}(\tanh(e + fx))}{f} + \frac{d \log(\cosh(e + fx))}{f^2} \\ &+ \frac{dx \text{sech}(e)(fx \cosh(e) - 2 \sinh(e))}{2f} \\ &- \frac{dx \text{sech}(e) \text{sech}(e + fx) \sinh(fx)}{f} - \frac{c \tanh(e + fx)}{f} \end{aligned}$$

[In] Integrate[(c + d*x)*Tanh[e + f*x]^2,x]

[Out] (c*ArcTanh[Tanh[e + f*x]])/f + (d*Log[Cosh[e + f*x]])/f^2 + (d*x*Sech[e]*(f*x*Cosh[e] - 2*Sinh[e]))/(2*f) - (d*x*Sech[e]*Sech[e + f*x]*Sinh[f*x])/f - (c*Tanh[e + f*x])/f

Maple [A] (verified)

Time = 0.05 (sec) , antiderivative size = 62, normalized size of antiderivative = 1.55

method	result	size
parallelrisch	$\frac{dx^2 f^2 - 2d \tanh(fx+e)xf + 2cx f^2 - 2dxf - 2c \tanh(fx+e)f - 2 \ln(1 - \tanh(fx+e))d}{2f^2}$	62
risch	$\frac{dx^2}{2} + cx - \frac{2dx}{f} - \frac{2de}{f^2} + \frac{2dx+2c}{f(1+e^{2fx+2e})} + \frac{d \ln(1+e^{2fx+2e})}{f^2}$	65

[In] int((d*x+c)*tanh(f*x+e)^2,x,method=_RETURNVERBOSE)

[Out] 1/2*(d*x^2*f^2-2*d*tanh(f*x+e)*x*f+2*c*x*f^2-2*d*x*f-2*c*tanh(f*x+e)*f-2*ln(1-tanh(f*x+e))*d)/f^2

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 232 vs. 2(38) = 76.

Time = 0.27 (sec) , antiderivative size = 232, normalized size of antiderivative = 5.80

$$\int (c + dx) \tanh^2(e + fx) dx$$

$$= \frac{df^2x^2 + 2cf^2x + (df^2x^2 + 2(cf^2 - 2df)x) \cosh(fx + e)^2 + 2(df^2x^2 + 2(cf^2 - 2df)x) \cosh(fx + e) \sinh(fx + e) + d \sinh(fx + e)^2 + d \log(2 \cosh(fx + e) - \sinh(fx + e))}{2(f^2 \cosh(fx + e)^2 + 2f^2 \cosh(fx + e) \sinh(fx + e) + f^2 \sinh(fx + e)^2 + f^2)}$$

[In] integrate((d*x+c)*tanh(f*x+e)^2,x, algorithm="fricas")

[Out] 1/2*(d*f^2*x^2 + 2*c*f^2*x + (d*f^2*x^2 + 2*(c*f^2 - 2*d*f)*x)*cosh(f*x + e)^2 + 2*(d*f^2*x^2 + 2*(c*f^2 - 2*d*f)*x)*cosh(f*x + e)*sinh(f*x + e) + (d*f^2*x^2 + 2*(c*f^2 - 2*d*f)*x)*sinh(f*x + e)^2 + 4*c*f + 2*(d*cosh(f*x + e)^2 + 2*d*cosh(f*x + e)*sinh(f*x + e) + d*sinh(f*x + e)^2 + d)*log(2*cosh(f*x + e)/(cosh(f*x + e) - sinh(f*x + e)))/(f^2*cosh(f*x + e)^2 + 2*f^2*cosh(f*x + e)*sinh(f*x + e) + f^2*sinh(f*x + e)^2 + f^2)

Sympy [A] (verification not implemented)

Time = 0.15 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.65

$$\int (c + dx) \tanh^2(e + fx) dx$$

$$= \begin{cases} cx - \frac{c \tanh(e+fx)}{f} + \frac{dx^2}{2} - \frac{dx \tanh(e+fx)}{f} + \frac{dx}{f} - \frac{d \log(\tanh(e+fx)+1)}{f^2} & \text{for } f \neq 0 \\ \left(cx + \frac{dx^2}{2}\right) \tanh^2(e) & \text{otherwise} \end{cases}$$

[In] integrate((d*x+c)*tanh(f*x+e)**2,x)

[Out] Piecewise((c*x - c*tanh(e + f*x)/f + d*x**2/2 - d*x*tanh(e + f*x)/f + d*x/f - d*log(tanh(e + f*x) + 1)/f**2, Ne(f, 0)), ((c*x + d*x**2/2)*tanh(e)**2, True))

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. 2(38) = 76.

Time = 0.24 (sec) , antiderivative size = 127, normalized size of antiderivative = 3.18

$$\int (c + dx) \tanh^2(e + fx) dx = c \left(x + \frac{e}{f} - \frac{2}{f(e^{(-2fx-2e)} + 1)} \right) - \frac{1}{2} d \left(\frac{2xe^{(2fx+2e)}}{fe^{(2fx+2e)} + f} - \frac{fx^2 + (fx^2e^{(2e)} - 2xe^{(2e)})e^{(2fx)}}{fe^{(2fx+2e)} + f} - \frac{2 \log((e^{(2fx+2e)} + 1)e^{(-2e)})}{f^2} \right)$$

[In] integrate((d*x+c)*tanh(f*x+e)^2,x, algorithm="maxima")

[Out] c*(x + e/f - 2/(f*(e^(-2*f*x - 2*e) + 1))) - 1/2*d*(2*x*e^(2*f*x + 2*e)/(f*e^(2*f*x + 2*e) + f) - (f*x^2 + (f*x^2*e^(2*e) - 2*x*e^(2*e))*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) - 2*log((e^(2*f*x + 2*e) + 1)*e^(-2*e))/f^2)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 127 vs. 2(38) = 76.

Time = 0.28 (sec) , antiderivative size = 127, normalized size of antiderivative = 3.18

$$\int (c + dx) \tanh^2(e + fx) dx = \frac{df^2x^2e^{(2fx+2e)} + df^2x^2 + 2cf^2xe^{(2fx+2e)} + 2cf^2x - 4dfxe^{(2fx+2e)} + 2de^{(2fx+2e)} \log(e^{(2fx+2e)} + 1) + 4c}{2(f^2e^{(2fx+2e)} + f^2)}$$

[In] integrate((d*x+c)*tanh(f*x+e)^2,x, algorithm="giac")

[Out] 1/2*(d*f^2*x^2*e^(2*f*x + 2*e) + d*f^2*x^2 + 2*c*f^2*x*e^(2*f*x + 2*e) + 2*c*f^2*x - 4*d*f*x*e^(2*f*x + 2*e) + 2*d*e^(2*f*x + 2*e)*log(e^(2*f*x + 2*e) + 1) + 4*c*f + 2*d*log(e^(2*f*x + 2*e) + 1))/(f^2*e^(2*f*x + 2*e) + f^2)

Mupad [B] (verification not implemented)

Time = 0.10 (sec) , antiderivative size = 56, normalized size of antiderivative = 1.40

$$\int (c + dx) \tanh^2(e + fx) dx = x \left(c + \frac{d}{f} \right) + \frac{dx^2}{2} - \frac{d \ln(\tanh(e + fx) + 1)}{f^2} - \frac{c \tanh(e + fx)}{f} - \frac{dx \tanh(e + fx)}{f}$$

[In] int(tanh(e + f*x)^2*(c + d*x),x)

[Out] x*(c + d/f) + (d*x^2)/2 - (d*log(tanh(e + f*x) + 1))/f^2 - (c*tanh(e + f*x))/f - (d*x*tanh(e + f*x))/f

3.9 $\int \frac{\tanh^2(e+fx)}{c+dx} dx$

Optimal result	85
Rubi [N/A]	85
Mathematica [N/A]	86
Maple [N/A] (verified)	86
Fricas [N/A]	86
Sympy [N/A]	86
Maxima [N/A]	87
Giac [N/A]	87
Mupad [N/A]	87

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\tanh^2(e+fx)}{c+dx} dx = \text{Int}\left(\frac{\tanh^2(e+fx)}{c+dx}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)^2/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh^2(e+fx)}{c+dx} dx = \int \frac{\tanh^2(e+fx)}{c+dx} dx$$

[In] Int[Tanh[e + f*x]^2/(c + d*x),x]

[Out] Defer[Int][Tanh[e + f*x]^2/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{\tanh^2(e+fx)}{c+dx} dx$$

Mathematica [N/A]

Not integrable

Time = 19.23 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh^2(e + fx)}{c + dx} dx$$

[In] Integrate[Tanh[e + f*x]^2/(c + d*x),x]

[Out] Integrate[Tanh[e + f*x]^2/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\tanh^2(fx + e)}{dx + c} dx$$

[In] int(tanh(f*x+e)^2/(d*x+c),x)

[Out] int(tanh(f*x+e)^2/(d*x+c),x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh^2(fx + e)}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c),x, algorithm="fricas")

[Out] integral(tanh(f*x + e)^2/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh^2(e + fx)}{c + dx} dx$$

[In] integrate(tanh(f*x+e)**2/(d*x+c),x)

[Out] Integral(tanh(e + f*x)**2/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 110, normalized size of antiderivative = 6.88

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)^2}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c),x, algorithm="maxima")

```
[Out] 2*d*integrate(1/(d^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2*e^(2*e) + 2*c*d*f*x*e^(2*e) + c^2*f*e^(2*e))*e^(2*f*x)), x) + log(d*x + c)/d + 2/(d*f*x + c*f + (d*f*x*e^(2*e) + c*f*e^(2*e))*e^(2*f*x))
```

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)^2}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c),x, algorithm="giac")

[Out] integrate(tanh(f*x + e)^2/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.66 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{c + dx} dx = \int \frac{\tanh(e + fx)^2}{c + dx} dx$$

[In] int(tanh(e + f*x)^2/(c + d*x),x)

[Out] int(tanh(e + f*x)^2/(c + d*x), x)

3.10 $\int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx$

Optimal result	88
Rubi [N/A]	88
Mathematica [N/A]	89
Maple [N/A] (verified)	89
Fricas [N/A]	89
Sympy [N/A]	89
Maxima [N/A]	90
Giac [N/A]	90
Mupad [N/A]	90

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx = \text{Int}\left(\frac{\tanh^2(e+fx)}{(c+dx)^2}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)^2/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx = \int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx$$

[In] Int[Tanh[e + f*x]^2/(c + d*x)^2,x]

[Out] Defer[Int][Tanh[e + f*x]^2/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tanh^2(e+fx)}{(c+dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 24.68 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx$$

[In] Integrate[Tanh[e + f*x]^2/(c + d*x)^2,x]

[Out] Integrate[Tanh[e + f*x]^2/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(fx + e)^2}{(dx + c)^2} dx$$

[In] int(tanh(f*x+e)^2/(d*x+c)^2,x)

[Out] int(tanh(f*x+e)^2/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c)^2,x, algorithm="fricas")

[Out] integral(tanh(f*x + e)^2/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx$$

[In] integrate(tanh(f*x+e)**2/(d*x+c)**2,x)

[Out] Integral(tanh(e + f*x)**2/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 195, normalized size of antiderivative = 12.19

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c)^2,x, algorithm="maxima")

```
[Out] 4*d*integrate(1/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3*e^(2*e) + 3*c*d^2*f*x^2*e^(2*e) + 3*c^2*d*f*x*e^(2*e) + c^3*f*e^(2*e))*e^(2*f*x)), x) - (d*f*x + c*f + (d*f*x*e^(2*e) + c*f*e^(2*e))*e^(2*f*x) - 2*d)/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2*e^(2*e) + 2*c*d^2*f*x*e^(2*e) + c^2*d*f*e^(2*e))*e^(2*f*x))
```

Giac [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^2/(d*x+c)^2,x, algorithm="giac")

[Out] integrate(tanh(f*x + e)^2/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 1.68 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^2(e + fx)}{(c + dx)^2} dx$$

[In] int(tanh(e + f*x)^2/(c + d*x)^2,x)

[Out] int(tanh(e + f*x)^2/(c + d*x)^2, x)

3.11 $\int (c + dx)^3 \tanh^3(e + fx) dx$

Optimal result	91
Rubi [A] (verified)	92
Mathematica [B] (verified)	96
Maple [B] (verified)	96
Fricas [C] (verification not implemented)	97
Sympy [F]	97
Maxima [B] (verification not implemented)	98
Giac [F]	99
Mupad [F(-1)]	99

Optimal result

Integrand size = 16, antiderivative size = 237

$$\begin{aligned}
 \int (c + dx)^3 \tanh^3(e + fx) dx = & -\frac{3d(c + dx)^2}{2f^2} + \frac{(c + dx)^3}{2f} - \frac{(c + dx)^4}{4d} \\
 & + \frac{3d^2(c + dx) \log(1 + e^{2(e+fx)})}{f^3} \\
 & + \frac{(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3d^3 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^4} \\
 & + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 & - \frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 & + \frac{3d^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4} \\
 & - \frac{3d(c + dx)^2 \tanh(e + fx)}{2f^2} - \frac{(c + dx)^3 \tanh^2(e + fx)}{2f}
 \end{aligned}$$

```
[Out] -3/2*d*(d*x+c)^2/f^2+1/2*(d*x+c)^3/f-1/4*(d*x+c)^4/d+3*d^2*(d*x+c)*ln(1+exp
(2*f*x+2*e))/f^3+(d*x+c)^3*ln(1+exp(2*f*x+2*e))/f+3/2*d^3*polylog(2,-exp(2*
f*x+2*e))/f^4+3/2*d*(d*x+c)^2*polylog(2,-exp(2*f*x+2*e))/f^2-3/2*d^2*(d*x+c
)*polylog(3,-exp(2*f*x+2*e))/f^3+3/4*d^3*polylog(4,-exp(2*f*x+2*e))/f^4-3/2
*d*(d*x+c)^2*tanh(f*x+e)/f^2-1/2*(d*x+c)^3*tanh(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 0.28 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.625$, Rules used = {3801, 3799, 2221, 2317, 2438, 32, 2611, 6744, 2320, 6724}

$$\int (c + dx)^3 \tanh^3(e + fx) dx = -\frac{3d^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3d^2(c + dx) \log(e^{2(e+fx)} + 1)}{f^3} + \frac{3d(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3d(c + dx)^2 \tanh(e + fx)}{2f^2} + \frac{(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx)^3 \tanh^2(e + fx)}{2f} - \frac{3d(c + dx)^2}{2f^2} + \frac{(c + dx)^3}{2f} - \frac{(c + dx)^4}{4d} + \frac{3d^3 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^4} + \frac{3d^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

[In] Int[(c + d*x)^3*Tanh[e + f*x]^3,x]

[Out] (-3*d*(c + d*x)^2)/(2*f^2) + (c + d*x)^3/(2*f) - (c + d*x)^4/(4*d) + (3*d^2*(c + d*x)*Log[1 + E^(2*(e + f*x))])/f^3 + ((c + d*x)^3*Log[1 + E^(2*(e + f*x))])/f + (3*d^3*PolyLog[2, -E^(2*(e + f*x))])/(2*f^4) + (3*d*(c + d*x)^2*PolyLog[2, -E^(2*(e + f*x))])/(2*f^2) - (3*d^2*(c + d*x)*PolyLog[3, -E^(2*(e + f*x))])/(2*f^3) + (3*d^3*PolyLog[4, -E^(2*(e + f*x))])/(4*f^4) - (3*d*(c + d*x)^2*Tanh[e + f*x])/(2*f^2) - ((c + d*x)^3*Tanh[e + f*x]^2)/(2*f)

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 2221

Int[(((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*
(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c
+ d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_.)))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c+dx)^3 \tanh^2(e+fx)}{2f} + \frac{(3d) \int (c+dx)^2 \tanh^2(e+fx) dx}{2f} \\
&\quad + \int (c+dx)^3 \tanh(e+fx) dx \\
&= -\frac{(c+dx)^4}{4d} - \frac{3d(c+dx)^2 \tanh(e+fx)}{2f^2} - \frac{(c+dx)^3 \tanh^2(e+fx)}{2f} \\
&\quad + 2 \int \frac{e^{2(e+fx)}(c+dx)^3}{1+e^{2(e+fx)}} dx + \frac{(3d^2) \int (c+dx) \tanh(e+fx) dx}{f^2} + \frac{(3d) \int (c+dx)^2 dx}{2f} \\
&= -\frac{3d(c+dx)^2}{2f^2} + \frac{(c+dx)^3}{2f} - \frac{(c+dx)^4}{4d} + \frac{(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad - \frac{3d(c+dx)^2 \tanh(e+fx)}{2f^2} - \frac{(c+dx)^3 \tanh^2(e+fx)}{2f} \\
&\quad + \frac{(6d^2) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx}{f^2} - \frac{(3d) \int (c+dx)^2 \log(1+e^{2(e+fx)}) dx}{f} \\
&= -\frac{3d(c+dx)^2}{2f^2} + \frac{(c+dx)^3}{2f} - \frac{(c+dx)^4}{4d} + \frac{3d^2(c+dx) \log(1+e^{2(e+fx)})}{f^3} \\
&\quad + \frac{(c+dx)^3 \log(1+e^{2(e+fx)})}{f} + \frac{3d(c+dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{3d(c+dx)^2 \tanh(e+fx)}{2f^2} - \frac{(c+dx)^3 \tanh^2(e+fx)}{2f} \\
&\quad - \frac{(3d^3) \int \log(1+e^{2(e+fx)}) dx}{f^3} - \frac{(3d^2) \int (c+dx) \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3d(c+dx)^2}{2f^2} + \frac{(c+dx)^3}{2f} - \frac{(c+dx)^4}{4d} \\
&\quad + \frac{3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} + \frac{(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} - \frac{3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{3d(c+dx)^2\tanh(e+fx)}{2f^2} - \frac{(c+dx)^3\tanh^2(e+fx)}{2f} \\
&\quad - \frac{(3d^3)\text{Subst}\left(\int\frac{\log(1+x)}{x}dx, x, e^{2(e+fx)}\right)}{2f^4} + \frac{(3d^3)\int\text{PolyLog}(3,-e^{2(e+fx)})dx}{2f^3} \\
&= -\frac{3d(c+dx)^2}{2f^2} + \frac{(c+dx)^3}{2f} - \frac{(c+dx)^4}{4d} + \frac{3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} \\
&\quad + \frac{(c+dx)^3\log(1+e^{2(e+fx)})}{f} + \frac{3d^3\text{PolyLog}(2,-e^{2(e+fx)})}{2f^4} \\
&\quad + \frac{3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} - \frac{3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{3d(c+dx)^2\tanh(e+fx)}{2f^2} - \frac{(c+dx)^3\tanh^2(e+fx)}{2f} \\
&\quad + \frac{(3d^3)\text{Subst}\left(\int\frac{\text{PolyLog}(3,-x)}{x}dx, x, e^{2(e+fx)}\right)}{4f^4} \\
&= -\frac{3d(c+dx)^2}{2f^2} + \frac{(c+dx)^3}{2f} - \frac{(c+dx)^4}{4d} \\
&\quad + \frac{3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} + \frac{(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3d^3\text{PolyLog}(2,-e^{2(e+fx)})}{2f^4} + \frac{3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} + \frac{3d^3\text{PolyLog}(4,-e^{2(e+fx)})}{4f^4} \\
&\quad - \frac{3d(c+dx)^2\tanh(e+fx)}{2f^2} - \frac{(c+dx)^3\tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 496 vs. $2(237) = 474$.

Time = 6.91 (sec) , antiderivative size = 496, normalized size of antiderivative = 2.09

$$\int (c + dx)^3 \tanh^3(e + fx) dx$$

$$= \frac{-8c(1 + e^{2e})(3d^2 + c^2 f^2)x + \frac{2(c^2 f^2 + 2cdf^2 x + d^2(3 + f^2 x^2))^2}{df^2} + \frac{12d(1 + e^{2e})(d^2 + c^2 f^2)x \log(1 + e^{-2(e + fx)})}{f} + 12cd^2(1 + e^{2e})}{1} + \frac{(c + dx)^3 \operatorname{sech}^2(e + fx)}{2f} - \frac{3 \operatorname{sech}(e) \operatorname{sech}(e + fx)(c^2 d \sinh(fx) + 2cd^2 x \sinh(fx) + d^3 x^2 \sinh(fx))}{2f^2} + \frac{1}{4}x(4c^3 + 6c^2 dx + 4cd^2 x^2 + d^3 x^3) \tanh(e)$$

[In] Integrate[(c + d*x)^3*Tanh[e + f*x]^3,x]

[Out] $(-8*c*(1 + E^{(2*e)})*(3*d^2 + c^2*f^2)*x + (2*(c^2*f^2 + 2*c*d*f^2*x + d^2*(3 + f^2*x^2))^2)/(d*f^2) + (12*d*(1 + E^{(2*e)})*(d^2 + c^2*f^2)*x*\operatorname{Log}[1 + E^{(-2*(e + f*x))}])/f + 12*c*d^2*(1 + E^{(2*e)})*f*x^2*\operatorname{Log}[1 + E^{(-2*(e + f*x))}] + 4*d^3*(1 + E^{(2*e)})*f*x^3*\operatorname{Log}[1 + E^{(-2*(e + f*x))}] + (4*c*(1 + E^{(2*e)})*(3*d^2 + c^2*f^2)*\operatorname{Log}[1 + E^{(2*(e + f*x))}])/f - (6*d*(1 + E^{(2*e)})*(d^2 + c^2*f^2)*\operatorname{PolyLog}[2, -E^{(-2*(e + f*x))}])/f^2 - 12*c*d^2*(1 + E^{(2*e)})*x*\operatorname{PolyLog}[2, -E^{(-2*(e + f*x))}] - 6*d^3*(1 + E^{(2*e)})*x^2*\operatorname{PolyLog}[2, -E^{(-2*(e + f*x))}] - (6*c*d^2*(1 + E^{(2*e)})*\operatorname{PolyLog}[3, -E^{(-2*(e + f*x))}])/f - (6*d^3*(1 + E^{(2*e)})*x*\operatorname{PolyLog}[3, -E^{(-2*(e + f*x))}])/f - (3*d^3*(1 + E^{(2*e)})*\operatorname{PolyLog}[4, -E^{(-2*(e + f*x))}])/f^2)/(4*(1 + E^{(2*e)})*f^2) + ((c + d*x)^3*\operatorname{Sech}[e + f*x]^2)/(2*f) - (3*\operatorname{Sech}[e]*\operatorname{Sech}[e + f*x]*(c^2*d*\operatorname{Sinh}[f*x] + 2*c*d^2*x*\operatorname{Sinh}[f*x] + d^3*x^2*\operatorname{Sinh}[f*x]))/(2*f^2) + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*\operatorname{Tanh}[e])/4$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 684 vs. $2(219) = 438$.

Time = 0.27 (sec) , antiderivative size = 685, normalized size of antiderivative = 2.89

method	result
risch	$-d^2 c x^3 - \frac{3d c^2 x^2}{2} + c^3 x - \frac{d^3 x^4}{4} + \frac{2d^3 f x^3 e^{2fx+2e} + 6c d^2 f x^2 e^{2fx+2e} + 6c^2 d f x e^{2fx+2e} + 3d^3 x^2 e^{2fx+2e} + 2c^3 f e^{2fx+2e} + 6c d^2 f^2 (1 + e^{2fx+2e})^2}{f^2 (1 + e^{2fx+2e})^2}$

[In] int((d*x+c)^3*tanh(f*x+e)^3,x,method=_RETURNVERBOSE)


```
[Out] -d^2*c*x^3-3/2*d*c^2*x^2+c^3*x-1/4*d^3*x^4+(2*d^3*f*x^3*exp(2*f*x+2*e)+6*c*d^2*f*x^2*exp(2*f*x+2*e)+6*c^2*d*f*x*exp(2*f*x+2*e)+3*d^3*x^2*exp(2*f*x+2*e)+2*c^3*f*exp(2*f*x+2*e)+6*c*d^2*x*exp(2*f*x+2*e)+3*c^2*d*exp(2*f*x+2*e)+3*d^3*x^2+6*c*d^2*x+3*d*c^2)/f^2/(1+exp(2*f*x+2*e))^2-3/2/f^4*d^3*e^4+1/f*c^3*ln(1+exp(2*f*x+2*e))-2/f*c^3*ln(exp(f*x+e))+3/4*d^3*polylog(4,-exp(2*f*x+2*e))/f^4+3/2*d^3*polylog(2,-exp(2*f*x+2*e))/f^4-3/f^2*c^2*d*e^2+1/f*d^3*ln(1+exp(2*f*x+2*e))*x^3-3/2/f^3*c*d^2*polylog(3,-exp(2*f*x+2*e))+3/2/f^2*d^3*polylog(2,-exp(2*f*x+2*e))*x^2-3/2/f^3*d^3*polylog(3,-exp(2*f*x+2*e))*x+2/f^4*e^3*d^3*ln(exp(f*x+e))+3/2/f^2*c^2*d*polylog(2,-exp(2*f*x+2*e))+4/f^3*c*d^2*e^3-2/f^3*d^3*e^3*x-6/f*c^2*d*e*x+3/f*c*d^2*ln(1+exp(2*f*x+2*e))*x^2-6/f^3*c*e^2*d^2*ln(exp(f*x+e))+3/f^2*c*d^2*polylog(2,-exp(2*f*x+2*e))*x+6/f^2*c^2*e*d*ln(exp(f*x+e))+3/f*c^2*d*ln(1+exp(2*f*x+2*e))*x+6/f^2*c*d^2*e^2*x-6/f^3*d^3*e*x+6/f^4*e*d^3*ln(exp(f*x+e))+3/f^3*d^3*ln(1+exp(2*f*x+2*e))*x+3/f^3*d^2*c*ln(1+exp(2*f*x+2*e))-6/f^3*d^2*c*ln(exp(f*x+e))-3/f^4*e^2*d^3-3/f^2*d^3*x^2+1/4/d*c^4
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.33 (sec) , antiderivative size = 5569, normalized size of antiderivative = 23.50

$$\int (c + dx)^3 \tanh^3(e + fx) dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*tanh(f*x+e)^3,x, algorithm="fricas")
```

```
[Out] Too large to include
```

Sympy [F]

$$\int (c + dx)^3 \tanh^3(e + fx) dx = \int (c + dx)^3 \tanh^3(e + fx) dx$$

```
[In] integrate((d*x+c)**3*tanh(f*x+e)**3,x)
```

```
[Out] Integral((c + d*x)**3*tanh(e + f*x)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 595 vs. $2(217) = 434$.

Time = 0.38 (sec) , antiderivative size = 595, normalized size of antiderivative = 2.51

$$\begin{aligned}
 & \int (c + dx)^3 \tanh^3(e + fx) dx \\
 &= c^3 \left(x + \frac{e}{f} + \frac{\log(e^{(-2fx-2e)} + 1)}{f} + \frac{2e^{(-2fx-2e)}}{f(2e^{(-2fx-2e)} + e^{(-4fx-4e)} + 1)} \right) - \frac{6cd^2x}{f^2} \\
 &+ \frac{3(2f^2x^2 \log(e^{(2fx+2e)} + 1) + 2fx \operatorname{Li}_2(-e^{(2fx+2e)}) - \operatorname{Li}_3(-e^{(2fx+2e)}))cd^2}{2f^3} \\
 &+ \frac{3cd^2 \log(e^{(2fx+2e)} + 1)}{f^3} \\
 &+ \frac{d^3f^2x^4 + 4cd^2f^2x^3 + 24cd^2x + 12c^2d + 6(c^2df^2 + 2d^3)x^2 + (d^3f^2x^4e^{(4e)} + 4cd^2f^2x^3e^{(4e)} + 6c^2df^2x^2e^{(4e)} + 4cd^2f^2xe^{(4e)} + 6c^2dfe^{(4e)})}{3f^4} \\
 &+ \frac{(4f^3x^3 \log(e^{(2fx+2e)} + 1) + 6f^2x^2 \operatorname{Li}_2(-e^{(2fx+2e)}) - 6fx \operatorname{Li}_3(-e^{(2fx+2e)}) + 3 \operatorname{Li}_4(-e^{(2fx+2e)}))d^3}{3f^4} \\
 &+ \frac{3(c^2df^2 + d^3)(2fx \log(e^{(2fx+2e)} + 1) + \operatorname{Li}_2(-e^{(2fx+2e)}))}{2f^4} \\
 &- \frac{d^3f^4x^4 + 4cd^2f^4x^3 + 6(c^2df^2 + d^3)f^2x^2}{2f^4}
 \end{aligned}$$

[In] integrate((d*x+c)^3*tanh(f*x+e)^3,x, algorithm="maxima")

[Out] $c^3(x + e/f + \log(e^{(-2fx-2e)} + 1)/f + 2e^{(-2fx-2e)}/(f(2e^{(-2fx-2e)} + e^{(-4fx-4e)} + 1))) - 6c^2d^2x/f^2 + 3/2(2f^2x^2 \log(e^{(2fx+2e)} + 1) + 2fx \operatorname{dilog}(-e^{(2fx+2e)}) - \operatorname{polylog}(3, -e^{(2fx+2e)})) * cd^2/f^3 + 3cd^2 \log(e^{(2fx+2e)} + 1)/f^3 + 1/4(d^3f^2x^4 + 4cd^2f^2x^3 + 24cd^2x + 12c^2d + 6(c^2df^2 + 2d^3)x^2 + (d^3f^2x^4e^{(4e)} + 4cd^2f^2x^3e^{(4e)} + 6c^2df^2x^2e^{(4e)} + 4cd^2f^2xe^{(4e)} + 6c^2dfe^{(4e)}) * e^{(4fx)} + 2(d^3f^2x^4e^{(2e)} + 4(c^2df^2e^{(2e)} + d^3f^2e^{(2e)})) * x^3 + 6c^2d^2e^{(2e)} + 6(c^2df^2e^{(2e)} + 2cd^2fe^{(2e)} + d^3e^{(2e)}) * x^2 + 12(c^2df^2e^{(2e)} + cd^2e^{(2e)}) * x) * e^{(2fx)}) / (f^2e^{(4fx+4e)} + 2f^2e^{(2fx+2e)} + f^2) + 1/3(4f^3x^3 \log(e^{(2fx+2e)} + 1) + 6f^2x^2 \operatorname{dilog}(-e^{(2fx+2e)}) - 6fx \operatorname{polylog}(3, -e^{(2fx+2e)}) + 3 \operatorname{polylog}(4, -e^{(2fx+2e)})) * d^3/f^4 + 3/2(c^2df^2 + d^3)(2fx \log(e^{(2fx+2e)} + 1) + \operatorname{dilog}(-e^{(2fx+2e)})) / f^4 - 1/2(d^3f^4x^4 + 4cd^2f^4x^3 + 6(c^2df^2 + d^3)f^2x^2) / f^4$

Giac [F]

$$\int (c + dx)^3 \tanh^3(e + fx) dx = \int (dx + c)^3 \tanh(fx + e)^3 dx$$

[In] integrate((d*x+c)^3*tanh(f*x+e)^3,x, algorithm="giac")

[Out] integrate((d*x + c)^3*tanh(f*x + e)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 \tanh^3(e + fx) dx = \int \tanh(e + fx)^3 (c + dx)^3 dx$$

[In] int(tanh(e + f*x)^3*(c + d*x)^3,x)

[Out] int(tanh(e + f*x)^3*(c + d*x)^3, x)

3.12 $\int (c + dx)^2 \tanh^3(e + fx) dx$

Optimal result	100
Rubi [A] (verified)	100
Mathematica [A] (verified)	103
Maple [B] (verified)	104
Fricas [C] (verification not implemented)	104
Sympy [F]	106
Maxima [B] (verification not implemented)	106
Giac [F]	107
Mupad [F(-1)]	107

Optimal result

Integrand size = 16, antiderivative size = 157

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \frac{cdx}{f} + \frac{d^2x^2}{2f} - \frac{(c + dx)^3}{3d} + \frac{(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} + \frac{d^2 \log(\cosh(e + fx))}{f^3} + \frac{d(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} - \frac{d(c + dx) \tanh(e + fx)}{f^2} - \frac{(c + dx)^2 \tanh^2(e + fx)}{2f}$$

```
[Out] c*d*x/f+1/2*d^2*x^2/f-1/3*(d*x+c)^3/d+(d*x+c)^2*ln(1+exp(2*f*x+2*e))/f+d^2*ln(cosh(f*x+e))/f^3+d*(d*x+c)*polylog(2,-exp(2*f*x+2*e))/f^2-1/2*d^2*polylog(3,-exp(2*f*x+2*e))/f^3-d*(d*x+c)*tanh(f*x+e)/f^2-1/2*(d*x+c)^2*tanh(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used

= {3801, 3556, 3799, 2221, 2611, 2320, 6724}

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \frac{d(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d(c + dx) \tanh(e + fx)}{f^2} + \frac{(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx)^2 \tanh^2(e + fx)}{2f} + \frac{cdx}{f} - \frac{(c + dx)^3}{3d} - \frac{d^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{d^2 \log(\cosh(e + fx))}{f^3} + \frac{d^2 x^2}{2f}$$

[In] Int[(c + d*x)^2*Tanh[e + f*x]^3,x]

[Out] (c*d*x)/f + (d^2*x^2)/(2*f) - (c + d*x)^3/(3*d) + ((c + d*x)^2*Log[1 + E^(2*(e + f*x))])/f + (d^2*Log[Cosh[e + f*x]])/f^3 + (d*(c + d*x)*PolyLog[2, -E^(2*(e + f*x))])/f^2 - (d^2*PolyLog[3, -E^(2*(e + f*x))])/f^3 - (d*(c + d*x)*Tanh[e + f*x])/f^2 - ((c + d*x)^2*Tanh[e + f*x]^2)/(2*f)

Rule 2221

Int[(((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^(g_)*((e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^(c_)*((a_) + (b_)*(x_)))^(n_)]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3556

Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c + dx)^2 \tanh^2(e + fx)}{2f} + \frac{d \int (c + dx) \tanh^2(e + fx) dx}{f} \\
&\quad + \int (c + dx)^2 \tanh(e + fx) dx \\
&= -\frac{(c + dx)^3}{3d} - \frac{d(c + dx) \tanh(e + fx)}{f^2} - \frac{(c + dx)^2 \tanh^2(e + fx)}{2f} \\
&\quad + 2 \int \frac{e^{2(e+fx)}(c + dx)^2}{1 + e^{2(e+fx)}} dx + \frac{d^2 \int \tanh(e + fx) dx}{f^2} + \frac{d \int (c + dx) dx}{f} \\
&= \frac{cdx}{f} + \frac{d^2 x^2}{2f} - \frac{(c + dx)^3}{3d} + \frac{(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} + \frac{d^2 \log(\cosh(e + fx))}{f^3} \\
&\quad - \frac{d(c + dx) \tanh(e + fx)}{f^2} - \frac{(c + dx)^2 \tanh^2(e + fx)}{2f} - \frac{(2d) \int (c + dx) \log(1 + e^{2(e+fx)}) dx}{f} \\
&= \frac{cdx}{f} + \frac{d^2 x^2}{2f} - \frac{(c + dx)^3}{3d} + \frac{(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} + \frac{d^2 \log(\cosh(e + fx))}{f^3} \\
&\quad + \frac{d(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d(c + dx) \tanh(e + fx)}{f^2} \\
&\quad - \frac{(c + dx)^2 \tanh^2(e + fx)}{2f} - \frac{d^2 \int \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{cdx}{f} + \frac{d^2x^2}{2f} - \frac{(c+dx)^3}{3d} + \frac{(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{d^2 \log(\cosh(e+fx))}{f^3} \\
&\quad + \frac{d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d(c+dx) \tanh(e+fx)}{f^2} \\
&\quad - \frac{(c+dx)^2 \tanh^2(e+fx)}{2f} - \frac{d^2 \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&= \frac{cdx}{f} + \frac{d^2x^2}{2f} - \frac{(c+dx)^3}{3d} + \frac{(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{d^2 \log(\cosh(e+fx))}{f^3} \\
&\quad + \frac{d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{d^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{d(c+dx) \tanh(e+fx)}{f^2} - \frac{(c+dx)^2 \tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.00 (sec) , antiderivative size = 226, normalized size of antiderivative = 1.44

$$\begin{aligned}
\int (c+dx)^2 \tanh^3(e+fx) dx = \frac{1}{6} &\left(\frac{4x(-3c^2e^{2e}f^2 + 3cdf^2x + d^2(-3e^{2e} + f^2x^2))}{(1+e^{2e})f^2} \right. \\
&\quad + \frac{6dx(2c+dx) \log(1+e^{-2(e+fx)})}{f} \\
&\quad + \frac{6(d^2+c^2f^2) \log(1+e^{2(e+fx)})}{f^3} \\
&\quad - \frac{6d(c+dx) \operatorname{PolyLog}(2, -e^{-2(e+fx)})}{f^2} \\
&\quad - \frac{3d^2 \operatorname{PolyLog}(3, -e^{-2(e+fx)})}{f^3} + \frac{3(c+dx)^2 \operatorname{sech}^2(e+fx)}{f} \\
&\quad - \frac{6d(c+dx) \operatorname{sech}(e) \operatorname{sech}(e+fx) \sinh(fx)}{f^2} \\
&\quad \left. + 2x(3c^2 + 3cdx + d^2x^2) \tanh(e) \right)
\end{aligned}$$

[In] Integrate[(c + d*x)^2*Tanh[e + f*x]^3,x]

[Out] ((4*x*(-3*c^2*E^(2*e)*f^2 + 3*c*d*f^2*x + d^2*(-3*E^(2*e) + f^2*x^2)))/((1 + E^(2*e))*f^2) + (6*d*x*(2*c + d*x)*Log[1 + E^(-2*(e + f*x))])/f + (6*(d^2 + c^2*f^2)*Log[1 + E^(2*(e + f*x))])/f^3 - (6*d*(c + d*x)*PolyLog[2, -E^(-2*(e + f*x))])/f^2 - (3*d^2*PolyLog[3, -E^(-2*(e + f*x))])/f^3 + (3*(c + d*x)^2*Sech[e + f*x]^2)/f - (6*d*(c + d*x)*Sech[e]*Sech[e + f*x]*Sinh[f*x])/f^2 + 2*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Tanh[e])/6

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 374 vs. 2(149) = 298.

Time = 0.24 (sec) , antiderivative size = 375, normalized size of antiderivative = 2.39

method	result
risch	$-\frac{d^2x^3}{3} - dcx^2 + c^2x + \frac{c^3}{3d} + \frac{2d^2fx^2e^{2fx+2e} + 4cdfxe^{2fx+2e} + 2c^2fe^{2fx+2e} + 2d^2xe^{2fx+2e} + 2cde^{2fx+2e} + 2xd^2 + 2cd}{f^2(1+e^{2fx+2e})^2} + 2d$

[In] int((d*x+c)^2*tanh(f*x+e)^3,x,method=_RETURNVERBOSE)

[Out] $-1/3*d^2*x^3-d*c*x^2+c^2*x+1/3/d*c^3+2*(d^2*f*x^2*exp(2*f*x+2*e)+2*c*d*f*x*exp(2*f*x+2*e)+c^2*f*exp(2*f*x+2*e)+d^2*x*exp(2*f*x+2*e)+c*d*exp(2*f*x+2*e)+x*d^2+c*d)/f^2/(1+exp(2*f*x+2*e))^2+2/f^2*d^2*e^2*x+1/f*d^2*ln(1+exp(2*f*x+2*e))*x^2+1/f^2*d^2*polylog(2,-exp(2*f*x+2*e))*x-2/f^2*d*c*e^2+1/f^2*d*c*polylog(2,-exp(2*f*x+2*e))-2/f^3*e^2*d^2*ln(exp(f*x+e))+1/f*c^2*ln(1+exp(2*f*x+2*e))-2/f*c^2*ln(exp(f*x+e))+4/f^2*c*e*d*ln(exp(f*x+e))-4/f*d*c*e*x+2/f*d*c*ln(1+exp(2*f*x+2*e))*x+4/3/f^3*d^2*e^3-1/2*d^2*polylog(3,-exp(2*f*x+2*e))/f^3+1/f^3*d^2*ln(1+exp(2*f*x+2*e))-2/f^3*d^2*ln(exp(f*x+e))$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 3071, normalized size of antiderivative = 19.56

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \text{Too large to display}$$

[In] integrate((d*x+c)^2*tanh(f*x+e)^3,x, algorithm="fricas")

[Out] $-1/3*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 3*c^2*f^3*x + 2*d^2*e^3 + 6*c^2*e*f^2 + (d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 2*d^2*e^3 - 6*c*d*e^2*f + 6*c^2*e*f^2 + 6*d^2*e + 3*(c^2*f^3 + 2*d^2*f)*x)*cosh(f*x + e)^4 + 4*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 2*d^2*e^3 - 6*c*d*e^2*f + 6*c^2*e*f^2 + 6*d^2*e + 3*(c^2*f^3 + 2*d^2*f)*x)*cosh(f*x + e)*sinh(f*x + e)^3 + (d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 2*d^2*e^3 - 6*c*d*e^2*f + 6*c^2*e*f^2 + 6*d^2*e + 3*(c^2*f^3 + 2*d^2*f)*x)*sinh(f*x + e)^4 + 6*d^2*e + 2*(d^2*f^3*x^3 + 2*d^2*e^3 + 6*d^2*e + 3*(2*c^2*e - c^2)*f^2 + 3*(c*d*f^3 - d^2*f^2)*x^2 - 3*(2*c*d*e^2 + c*d)*f + 3*(c^2*f^3 - 2*c*d*f^2 + d^2*f)*x)*cosh(f*x + e)^2 + 2*(d^2*f^3*x^3 + 2*d^2*e^3 + 6*d^2*e + 3*(2*c^2*e - c^2)*f^2 + 3*(c*d*f^3 - d^2*f^2)*x^2 + 3*(d^2*f^3*x^3 + 3*c*d*f^3*x^2 + 2*d^2*e^3 - 6*c*d*e^2*f + 6*c^2*e*f^2 + 6*d^2*e + 3*(c^2*f^3 + 2*d^2*f)*x)*cosh(f*x + e)^2 - 3*(2*c*d*e^2 + c*d)*f + 3*(c^2*f^3 - 2*c*d*f^2 + d^2*f)*x)*sinh(f*x + e)^2 - 6*(c*d*e^2 + c*d)*f - 6*((d^2*f*x + c*d*f)*cosh(f*x + e)^4 + 4*(d^2*f*x + c*d*f)*cosh(f*x + e)*sinh(f*x + e)^3 + (d^2*f*x + c*d*f)*sinh(f*x + e)^4 + d^2*f*x + c*d*f + 2*(d^2*f*x + c*d*f)*co$

$$\begin{aligned}
& \text{sh}(f*x + e)^2 + 2*(d^2*f*x + c*d*f + 3*(d^2*f*x + c*d*f)*\cosh(f*x + e)^2)*\text{sinh}(f*x + e)^2 + 4*((d^2*f*x + c*d*f)*\cosh(f*x + e)^3 + (d^2*f*x + c*d*f)*\cosh(f*x + e))*\text{sinh}(f*x + e))*\text{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 6*((d^2*f*x + c*d*f)*\cosh(f*x + e)^4 + 4*(d^2*f*x + c*d*f)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d^2*f*x + c*d*f)*\sinh(f*x + e)^4 + d^2*f*x + c*d*f + 2*(d^2*f*x + c*d*f)*\cosh(f*x + e)^2 + 2*(d^2*f*x + c*d*f + 3*(d^2*f*x + c*d*f)*\cosh(f*x + e)^2)*\sinh(f*x + e)^2 + 4*((d^2*f*x + c*d*f)*\cosh(f*x + e)^3 + (d^2*f*x + c*d*f)*\cosh(f*x + e))*\sinh(f*x + e))*\text{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) - 3*((d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^4 + 4*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\sinh(f*x + e)^4 + d^2*e^2 - 2*c*d*e*f + c^2*f^2 + 2*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^2 + 2*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + 3*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^2 + d^2)*\sinh(f*x + e)^2 + d^2 + 4*((d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^3 + (d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e))*\sinh(f*x + e))*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) - 3*((d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^4 + 4*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\sinh(f*x + e)^4 + d^2*e^2 - 2*c*d*e*f + c^2*f^2 + 2*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^2 + 2*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + 3*(d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^2 + d^2)*\sinh(f*x + e)^2 + d^2 + 4*((d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e)^3 + (d^2*e^2 - 2*c*d*e*f + c^2*f^2 + d^2)*\cosh(f*x + e))*\sinh(f*x + e))*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) - 3*(d^2*f^2*x^2 + 2*c*d*f^2*x + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^4 + 4*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\sinh(f*x + e)^4 - d^2*e^2 + 2*c*d*e*f + 2*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^2 + 2*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f + 3*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^2)*\sinh(f*x + e)^2 + 4*((d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^3 + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e))*\sinh(f*x + e))*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) - 3*(d^2*f^2*x^2 + 2*c*d*f^2*x + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^4 + 4*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\sinh(f*x + e)^4 - d^2*e^2 + 2*c*d*e*f + 2*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^2 + 2*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f + 3*(d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^2)*\sinh(f*x + e)^2 + 4*((d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e)^3 + (d^2*f^2*x^2 + 2*c*d*f^2*x - d^2*e^2 + 2*c*d*e*f)*\cosh(f*x + e))*\sinh(f*x + e))*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1) + 6*(d^2*\cosh(f*x + e)^4 + 4*d^2*\cosh(f*x + e)*\sinh(f*x + e)^3 + d^2*\sinh(f*x + e)^4 + 2*d^2*\cosh(f*x + e)^2 + 2*(3*d^2*\cosh(f*x + e)^2 + d^2)*\sinh(f*x + e)^2 + d^2 + 4*(d^2*\cosh(f*x + e)^3 + d^2*\cosh(f*x + e))*\sinh(f*x + e))*\text{polylog}(3, I*\cosh(f*x + e) + I*\sinh(f*x + e))
\end{aligned}$$

```

+ 6*(d^2*cosh(f*x + e)^4 + 4*d^2*cosh(f*x + e)*sinh(f*x + e)^3 + d^2*sinh(f
*x + e)^4 + 2*d^2*cosh(f*x + e)^2 + 2*(3*d^2*cosh(f*x + e)^2 + d^2)*sinh(f*
x + e)^2 + d^2 + 4*(d^2*cosh(f*x + e)^3 + d^2*cosh(f*x + e))*sinh(f*x + e))
*polylog(3, -I*cosh(f*x + e) - I*sinh(f*x + e)) + 4*((d^2*f^3*x^3 + 3*c*d*f
^3*x^2 + 2*d^2*e^3 - 6*c*d*e^2*f + 6*c^2*e*f^2 + 6*d^2*e + 3*(c^2*f^3 + 2*d
^2*f)*x)*cosh(f*x + e)^3 + (d^2*f^3*x^3 + 2*d^2*e^3 + 6*d^2*e + 3*(2*c^2*e
- c^2)*f^2 + 3*(c*d*f^3 - d^2*f^2)*x^2 - 3*(2*c*d*e^2 + c*d)*f + 3*(c^2*f^3
- 2*c*d*f^2 + d^2*f)*x)*cosh(f*x + e))*sinh(f*x + e))/(f^3*cosh(f*x + e)^4
+ 4*f^3*cosh(f*x + e)*sinh(f*x + e)^3 + f^3*sinh(f*x + e)^4 + 2*f^3*cosh(f
*x + e)^2 + f^3 + 2*(3*f^3*cosh(f*x + e)^2 + f^3)*sinh(f*x + e)^2 + 4*(f^3*
cosh(f*x + e)^3 + f^3*cosh(f*x + e))*sinh(f*x + e))

```

Sympy [F]

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \int (c + dx)^2 \tanh^3(e + fx) dx$$

```
[In] integrate((d*x+c)**2*tanh(f*x+e)**3,x)
```

```
[Out] Integral((c + d*x)**2*tanh(e + f*x)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 392 vs. 2(148) = 296.

Time = 0.38 (sec) , antiderivative size = 392, normalized size of antiderivative = 2.50

$$\begin{aligned}
& \int (c + dx)^2 \tanh^3(e + fx) dx \\
&= c^2 \left(x + \frac{e}{f} + \frac{\log(e^{-2fx-2e} + 1)}{f} + \frac{2e^{-2fx-2e}}{f(2e^{-2fx-2e} + e^{-4fx-4e} + 1)} \right) \\
&+ \frac{(2fx \log(e^{2fx+2e} + 1) + \text{Li}_2(-e^{2fx+2e}))cd}{f^2} - \frac{2d^2x}{f^2} \\
&+ \frac{d^2f^2x^3 + 3cdf^2x^2 + 6d^2x + 6cd + (d^2f^2x^3e^{4e} + 3cdf^2x^2e^{4e})e^{4fx} + 2(d^2f^2x^3e^{2e} + 3(cdf^2e^{2e} + 3(f^2e^{4fx+4e} + 2f^2e^{2fx+2e} + f^2))d^2}{3(f^2e^{4fx+4e} + 2f^2e^{2fx+2e} + f^2)} \\
&+ \frac{(2f^2x^2 \log(e^{2fx+2e} + 1) + 2fx\text{Li}_2(-e^{2fx+2e}) - \text{Li}_3(-e^{2fx+2e}))d^2}{2f^3} \\
&+ \frac{d^2 \log(e^{2fx+2e} + 1)}{f^3} - \frac{2(d^2f^3x^3 + 3cdf^3x^2)}{3f^3}
\end{aligned}$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e)^3,x, algorithm="maxima")
```

```
[Out] c^2*(x + e/f + log(e^(-2*f*x - 2*e) + 1)/f + 2*e^(-2*f*x - 2*e)/(f*(2*e^(-2
*f*x - 2*e) + e^(-4*f*x - 4*e) + 1))) + (2*f*x*log(e^(2*f*x + 2*e) + 1) + d
ilog(-e^(2*f*x + 2*e)))*c*d/f^2 - 2*d^2*x/f^2 + 1/3*(d^2*f^2*x^3 + 3*c*d*f^
2*x^2 + 6*d^2*x + 6*c*d + (d^2*f^2*x^3*e^(4*e) + 3*c*d*f^2*x^2*e^(4*e))*e^(
4*f*x) + 2*(d^2*f^2*x^3*e^(2*e) + 3*(c*d*f^2*e^(2*e) + d^2*f*e^(2*e))*x^2 +
3*c*d*e^(2*e) + 3*(2*c*d*f*e^(2*e) + d^2*e^(2*e))*x)*e^(2*f*x))/(f^2*e^(4*
f*x + 4*e) + 2*f^2*e^(2*f*x + 2*e) + f^2) + 1/2*(2*f^2*x^2*log(e^(2*f*x + 2
*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))*d^
2/f^3 + d^2*log(e^(2*f*x + 2*e) + 1)/f^3 - 2/3*(d^2*f^3*x^3 + 3*c*d*f^3*x^2
)/f^3
```

Giac [F]

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \int (dx + c)^2 \tanh(fx + e)^3 dx$$

```
[In] integrate((d*x+c)^2*tanh(f*x+e)^3,x, algorithm="giac")
```

```
[Out] integrate((d*x + c)^2*tanh(f*x + e)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 \tanh^3(e + fx) dx = \int \tanh(e + fx)^3 (c + dx)^2 dx$$

```
[In] int(tanh(e + f*x)^3*(c + d*x)^2,x)
```

```
[Out] int(tanh(e + f*x)^3*(c + d*x)^2, x)
```

3.13 $\int (c + dx) \tanh^3(e + fx) dx$

Optimal result	108
Rubi [A] (verified)	108
Mathematica [A] (verified)	110
Maple [A] (verified)	111
Fricas [C] (verification not implemented)	111
Sympy [F]	112
Maxima [F]	112
Giac [F]	113
Mupad [F(-1)]	113

Optimal result

Integrand size = 14, antiderivative size = 100

$$\int (c + dx) \tanh^3(e + fx) dx = \frac{dx}{2f} - \frac{(c + dx)^2}{2d} + \frac{(c + dx) \log(1 + e^{2(e+fx)})}{f} + \frac{d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{d \tanh(e + fx)}{2f^2} - \frac{(c + dx) \tanh^2(e + fx)}{2f}$$

[Out] $1/2*d*x/f - 1/2*(d*x+c)^2/d + (d*x+c)*\ln(1+\exp(2*f*x+2*e))/f + 1/2*d*\operatorname{polylog}(2, -\exp(2*f*x+2*e))/f^2 - 1/2*d*\tanh(f*x+e)/f^2 - 1/2*(d*x+c)*\tanh(f*x+e)^2/f$

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3801, 3554, 8, 3799, 2221, 2317, 2438}

$$\int (c + dx) \tanh^3(e + fx) dx = \frac{(c + dx) \log(e^{2(e+fx)} + 1)}{f} - \frac{(c + dx) \tanh^2(e + fx)}{2f} - \frac{(c + dx)^2}{2d} + \frac{d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{d \tanh(e + fx)}{2f^2} + \frac{dx}{2f}$$

[In] $\operatorname{Int}[(c + d*x)*\operatorname{Tanh}[e + f*x]^3, x]$

[Out] $(d*x)/(2*f) - (c + d*x)^2/(2*d) + ((c + d*x)*\text{Log}[1 + E^{(2*(e + f*x))}])/f + (d*\text{PolyLog}[2, -E^{(2*(e + f*x))}])/(2*f^2) - (d*\text{Tanh}[e + f*x])/(2*f^2) - ((c + d*x)*\text{Tanh}[e + f*x]^2)/(2*f)$

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 2221

`Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]`

Rule 2317

`Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]`

Rule 2438

`Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]`

Rule 3554

`Int[((b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]`

Rule 3799

`Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]`

Rule 3801

`Int[((c_) + (d_)*(x_))^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]`

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c+dx)\tanh^2(e+fx)}{2f} + \frac{d\int \tanh^2(e+fx)dx}{2f} + \int (c+dx)\tanh(e+fx)dx \\
&= -\frac{(c+dx)^2}{2d} - \frac{d\tanh(e+fx)}{2f^2} - \frac{(c+dx)\tanh^2(e+fx)}{2f} + 2\int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}}dx \\
&\quad + \frac{d\int 1dx}{2f} \\
&= \frac{dx}{2f} - \frac{(c+dx)^2}{2d} + \frac{(c+dx)\log(1+e^{2(e+fx)})}{f} - \frac{d\tanh(e+fx)}{2f^2} \\
&\quad - \frac{(c+dx)\tanh^2(e+fx)}{2f} - \frac{d\int \log(1+e^{2(e+fx)})dx}{f} \\
&= \frac{dx}{2f} - \frac{(c+dx)^2}{2d} + \frac{(c+dx)\log(1+e^{2(e+fx)})}{f} - \frac{d\tanh(e+fx)}{2f^2} \\
&\quad - \frac{(c+dx)\tanh^2(e+fx)}{2f} - \frac{d\text{Subst}\left(\int \frac{\log(1+x)}{x}dx, x, e^{2(e+fx)}\right)}{2f^2} \\
&= \frac{dx}{2f} - \frac{(c+dx)^2}{2d} + \frac{(c+dx)\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{d\text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{d\tanh(e+fx)}{2f^2} - \frac{(c+dx)\tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.81 (sec) , antiderivative size = 97, normalized size of antiderivative = 0.97

$$\begin{aligned}
&\int (c+dx)\tanh^3(e+fx)dx \\
&= \frac{-d\text{PolyLog}(2, -e^{-2(e+fx)}) + dfx\text{sech}^2(e+fx) - d\text{sech}(e)\text{sech}(e+fx)\sinh(fx) + f(dfx^2 + 2dx\log(1+e^{2(e+fx)}))}{2f^2}
\end{aligned}$$

[In] Integrate[(c + d*x)*Tanh[e + f*x]^3,x]

[Out] $(-(d*\text{PolyLog}[2, -E^{(-2*(e + f*x))}]) + d*f*x*\text{Sech}[e + f*x]^2 - d*\text{Sech}[e]*\text{Sech}[e + f*x]*\text{Sinh}[f*x] + f*(d*f*x^2 + 2*d*x*\text{Log}[1 + E^{(-2*(e + f*x))}] + 2*c*\text{Log}[\text{Cosh}[e + f*x]] - c*\text{Tanh}[e + f*x]^2))/(2*f^2)$

Maple [A] (verified)

Time = 0.06 (sec) , antiderivative size = 166, normalized size of antiderivative = 1.66

method	result
risch	$-\frac{dx^2}{2} + cx + \frac{2dfxe^{2fx+2e} + 2cfe^{2fx+2e} + e^{2fx+2e}d+d}{f^2(1+e^{2fx+2e})^2} + \frac{c \ln(1+e^{2fx+2e})}{f} - \frac{2c \ln(e^{fx+e})}{f} - \frac{2dex}{f} - \frac{de^2}{f^2} + \frac{d \ln(1+e^{2fx+2e})}{f}$

[In] int((d*x+c)*tanh(f*x+e)^3,x,method=_RETURNVERBOSE)

[Out]
$$-1/2*d*x^2+c*x+(2*d*f*x*\exp(2*f*x+2*e)+2*c*f*\exp(2*f*x+2*e)+\exp(2*f*x+2*e)*d+d)/f^2/(1+\exp(2*f*x+2*e))^2+1/f*c*\ln(1+\exp(2*f*x+2*e))-2/f*c*\ln(\exp(f*x+e))-2/f*d*e*x-1/f^2*d*e^2+1/f*d*\ln(1+\exp(2*f*x+2*e))*x+1/2*d*polylog(2,-\exp(2*f*x+2*e))/f^2+2/f^2*e*d*\ln(\exp(f*x+e))$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 1462, normalized size of antiderivative = 14.62

$$\int (c + dx) \tanh^3(e + fx) dx = \text{Too large to display}$$

[In] integrate((d*x+c)*tanh(f*x+e)^3,x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/2*(d*f^2*x^2 + (d*f^2*x^2 + 2*c*f^2*x - 2*d*e^2 + 4*c*e*f)*\cosh(f*x + e) \\ & ^4 + 4*(d*f^2*x^2 + 2*c*f^2*x - 2*d*e^2 + 4*c*e*f)*\cosh(f*x + e)*\sinh(f*x + \\ & e)^3 + (d*f^2*x^2 + 2*c*f^2*x - 2*d*e^2 + 4*c*e*f)*\sinh(f*x + e)^4 + 2*c*f \\ & ^2*x - 2*d*e^2 + 4*c*e*f + 2*(d*f^2*x^2 - 2*d*e^2 + 2*(2*c*e - c)*f + 2*(c*f \\ & ^2 - d*f)*x - d)*\cosh(f*x + e)^2 + 2*(d*f^2*x^2 - 2*d*e^2 + 3*(d*f^2*x^2 + \\ & 2*c*f^2*x - 2*d*e^2 + 4*c*e*f)*\cosh(f*x + e)^2 + 2*(2*c*e - c)*f + 2*(c*f^ \\ & 2 - d*f)*x - d)*\sinh(f*x + e)^2 - 2*(d*\cosh(f*x + e)^4 + 4*d*\cosh(f*x + e)* \\ & \sinh(f*x + e)^3 + d*\sinh(f*x + e)^4 + 2*d*\cosh(f*x + e)^2 + 2*(3*d*\cosh(f*x \\ & + e)^2 + d)*\sinh(f*x + e)^2 + 4*(d*\cosh(f*x + e)^3 + d*\cosh(f*x + e))*\sinh \\ & (f*x + e) + d)*\operatorname{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 2*(d*\cosh(f*x + e) \\ &)^4 + 4*d*\cosh(f*x + e)*\sinh(f*x + e)^3 + d*\sinh(f*x + e)^4 + 2*d*\cosh(f*x \\ & + e)^2 + 2*(3*d*\cosh(f*x + e)^2 + d)*\sinh(f*x + e)^2 + 4*(d*\cosh(f*x + e)^3 \\ & + d*\cosh(f*x + e))*\sinh(f*x + e) + d)*\operatorname{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x \\ & + e)) + 2*((d*e - c*f)*\cosh(f*x + e)^4 + 4*(d*e - c*f)*\cosh(f*x + e)*\sinh(f \\ & *x + e)^3 + (d*e - c*f)*\sinh(f*x + e)^4 + 2*(d*e - c*f)*\cosh(f*x + e)^2 + 2 \\ & *(3*(d*e - c*f)*\cosh(f*x + e)^2 + d*e - c*f)*\sinh(f*x + e)^2 + d*e - c*f + \\ & 4*((d*e - c*f)*\cosh(f*x + e)^3 + (d*e - c*f)*\cosh(f*x + e))*\sinh(f*x + e))* \\ & \log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 2*((d*e - c*f)*\cosh(f*x + e)^4 + 4 \\ & *(d*e - c*f)*\cosh(f*x + e)*\sinh(f*x + e)^3 + (d*e - c*f)*\sinh(f*x + e)^4 + \\ & 2*(d*e - c*f)*\cosh(f*x + e)^2 + 2*(3*(d*e - c*f)*\cosh(f*x + e)^2 + d*e - c* \\ & f)*\sinh(f*x + e)^2 + d*e - c*f + 4*((d*e - c*f)*\cosh(f*x + e)^3 + (d*e - c* \end{aligned}$$

```
f)*cosh(f*x + e))*sinh(f*x + e))*log(cosh(f*x + e) + sinh(f*x + e) - 1) - 2
*((d*f*x + d*e)*cosh(f*x + e)^4 + 4*(d*f*x + d*e)*cosh(f*x + e)*sinh(f*x +
e)^3 + (d*f*x + d*e)*sinh(f*x + e)^4 + d*f*x + 2*(d*f*x + d*e)*cosh(f*x + e
)^2 + 2*(d*f*x + 3*(d*f*x + d*e)*cosh(f*x + e)^2 + d*e)*sinh(f*x + e)^2 + d
*e + 4*((d*f*x + d*e)*cosh(f*x + e)^3 + (d*f*x + d*e)*cosh(f*x + e))*sinh(f
*x + e))*log(1*cosh(f*x + e) + 1*sinh(f*x + e) + 1) - 2*((d*f*x + d*e)*cosh
(f*x + e)^4 + 4*(d*f*x + d*e)*cosh(f*x + e)*sinh(f*x + e)^3 + (d*f*x + d*e)
*sinh(f*x + e)^4 + d*f*x + 2*(d*f*x + d*e)*cosh(f*x + e)^2 + 2*(d*f*x + 3*(
d*f*x + d*e)*cosh(f*x + e)^2 + d*e)*sinh(f*x + e)^2 + d*e + 4*((d*f*x + d*e)
)*cosh(f*x + e)^3 + (d*f*x + d*e)*cosh(f*x + e))*sinh(f*x + e))*log(-1*cosh
(f*x + e) - 1*sinh(f*x + e) + 1) + 4*((d*f^2*x^2 + 2*c*f^2*x - 2*d*e^2 + 4*
c*e*f)*cosh(f*x + e)^3 + (d*f^2*x^2 - 2*d*e^2 + 2*(2*c*e - c)*f + 2*(c*f^2
- d*f)*x - d)*cosh(f*x + e))*sinh(f*x + e) - 2*d)/(f^2*cosh(f*x + e)^4 + 4*
f^2*cosh(f*x + e)*sinh(f*x + e)^3 + f^2*sinh(f*x + e)^4 + 2*f^2*cosh(f*x +
e)^2 + 2*(3*f^2*cosh(f*x + e)^2 + f^2)*sinh(f*x + e)^2 + f^2 + 4*(f^2*cosh(
f*x + e)^3 + f^2*cosh(f*x + e))*sinh(f*x + e))
```

Sympy [F]

$$\int (c + dx) \tanh^3(e + fx) dx = \int (c + dx) \tanh^3(e + fx) dx$$

```
[In] integrate((d*x+c)*tanh(f*x+e)**3,x)
```

```
[Out] Integral((c + d*x)*tanh(e + f*x)**3, x)
```

Maxima [F]

$$\int (c + dx) \tanh^3(e + fx) dx = \int (dx + c) \tanh^3(fx + e) dx$$

```
[In] integrate((d*x+c)*tanh(f*x+e)^3,x, algorithm="maxima")
```

```
[Out] c*(x + e/f + log(e^(-2*f*x - 2*e) + 1)/f + 2*e^(-2*f*x - 2*e)/(f*(2*e^(-2*f
*x - 2*e) + e^(-4*f*x - 4*e) + 1))) + 1/2*d*((f^2*x^2*e^(4*f*x + 4*e) + f^2
*x^2 + 2*(f^2*x^2*e^(2*e) + 2*f*x*e^(2*e) + e^(2*e))*e^(2*f*x) + 2)/(f^2*e^
(4*f*x + 4*e) + 2*f^2*e^(2*f*x + 2*e) + f^2) - 4*integrate(x/(e^(2*f*x + 2*
e) + 1), x))
```


Giac [F]

$$\int (c + dx) \tanh^3(e + fx) dx = \int (dx + c) \tanh(fx + e)^3 dx$$

[In] integrate((d*x+c)*tanh(f*x+e)^3,x, algorithm="giac")

[Out] integrate((d*x + c)*tanh(f*x + e)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx) \tanh^3(e + fx) dx = \int \tanh(e + fx)^3 (c + dx) dx$$

[In] int(tanh(e + f*x)^3*(c + d*x),x)

[Out] int(tanh(e + f*x)^3*(c + d*x), x)

3.14 $\int \frac{\tanh^3(e+fx)}{c+dx} dx$

Optimal result	114
Rubi [N/A]	114
Mathematica [N/A]	115
Maple [N/A] (verified)	115
Fricas [N/A]	115
Sympy [N/A]	115
Maxima [N/A]	116
Giac [N/A]	116
Mupad [N/A]	116

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\tanh^3(e+fx)}{c+dx} dx = \text{Int}\left(\frac{\tanh^3(e+fx)}{c+dx}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)^3/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh^3(e+fx)}{c+dx} dx = \int \frac{\tanh^3(e+fx)}{c+dx} dx$$

[In] Int[Tanh[e + f*x]^3/(c + d*x),x]

[Out] Defer[Int][Tanh[e + f*x]^3/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{\tanh^3(e+fx)}{c+dx} dx$$

Mathematica [N/A]

Not integrable

Time = 26.72 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh^3(e + fx)}{c + dx} dx$$

[In] Integrate[Tanh[e + f*x]^3/(c + d*x),x]

[Out] Integrate[Tanh[e + f*x]^3/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(fx + e)^3}{dx + c} dx$$

[In] int(tanh(f*x+e)^3/(d*x+c),x)

[Out] int(tanh(f*x+e)^3/(d*x+c),x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)^3}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c),x, algorithm="fricas")

[Out] integral(tanh(f*x + e)^3/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 14, normalized size of antiderivative = 0.88

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh^3(e + fx)}{c + dx} dx$$

[In] integrate(tanh(f*x+e)**3/(d*x+c),x)

[Out] Integral(tanh(e + f*x)**3/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 303, normalized size of antiderivative = 18.94

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)^3}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c),x, algorithm="maxima")

```
[Out] ((2*d*f*x*e^(2*e) + 2*c*f*e^(2*e) - d*e^(2*e))*e^(2*f*x) - d)/(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2 + (d^2*f^2*x^2*e^(4*e) + 2*c*d*f^2*x*e^(4*e) + c^2*f^2*e^(4*e))*e^(4*f*x) + 2*(d^2*f^2*x^2*e^(2*e) + 2*c*d*f^2*x*e^(2*e) + c^2*f^2*e^(2*e))*e^(2*f*x)) + log(d*x + c)/d - integrate(2*(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2 + d^2)/(d^3*f^2*x^3 + 3*c*d^2*f^2*x^2 + 3*c^2*d*f^2*x + c^3*f^2 + (d^3*f^2*x^3*e^(2*e) + 3*c*d^2*f^2*x^2*e^(2*e) + 3*c^2*d*f^2*x*e^(2*e) + c^3*f^2*e^(2*e))*e^(2*f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh(fx + e)^3}{dx + c} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c),x, algorithm="giac")

[Out] integrate(tanh(f*x + e)^3/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.71 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{c + dx} dx = \int \frac{\tanh(e + fx)^3}{c + dx} dx$$

[In] int(tanh(e + f*x)^3/(c + d*x),x)

[Out] int(tanh(e + f*x)^3/(c + d*x), x)

3.15 $\int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx$

Optimal result	117
Rubi [N/A]	117
Mathematica [N/A]	118
Maple [N/A] (verified)	118
Fricas [N/A]	118
Sympy [N/A]	118
Maxima [N/A]	119
Giac [N/A]	119
Mupad [N/A]	119

Optimal result

Integrand size = 16, antiderivative size = 16

$$\int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx = \text{Int}\left(\frac{\tanh^3(e+fx)}{(c+dx)^2}, x\right)$$

[Out] Unintegrable(tanh(f*x+e)^3/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx = \int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx$$

[In] Int[Tanh[e + f*x]^3/(c + d*x)^2,x]

[Out] Defer[Int][Tanh[e + f*x]^3/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{\tanh^3(e+fx)}{(c+dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 22.53 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx$$

[In] Integrate[Tanh[e + f*x]^3/(c + d*x)^2,x]

[Out] Integrate[Tanh[e + f*x]^3/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 16, normalized size of antiderivative = 1.00

$$\int \frac{\tanh(fx + e)^3}{(dx + c)^2} dx$$

[In] int(tanh(f*x+e)^3/(d*x+c)^2,x)

[Out] int(tanh(f*x+e)^3/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 29, normalized size of antiderivative = 1.81

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^3(fx + e)}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c)^2,x, algorithm="fricas")

[Out] integral(tanh(f*x + e)^3/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.94

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx$$

[In] integrate(tanh(f*x+e)**3/(d*x+c)**2,x)

[Out] Integral(tanh(e + f*x)**3/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.43 (sec) , antiderivative size = 501, normalized size of antiderivative = 31.31

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(fx + e)^3}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c)^2,x, algorithm="maxima")

[Out] $-(d^2f^2x^2 + 2c*d*f^2x + c^2f^2 + 2d^2 + (d^2f^2x^2e^{(4e)} + 2c*d*f^2xe^{(4e)} + c^2f^2e^{(4e)})e^{(4fx)} + 2*(d^2f^2x^2e^{(2e)} + c^2f^2e^{(2e)} - c*d*f*e^{(2e)} + d^2e^{(2e)} + (2c*d*f^2e^{(2e)} - d^2f*e^{(2e)})x)*e^{(2fx)})/(d^4f^2x^3 + 3c*d^3f^2x^2 + 3c^2d^2f^2x + c^3d*f^2 + (d^4f^2x^3e^{(4e)} + 3c*d^3f^2x^2e^{(4e)} + 3c^2d^2f^2xe^{(4e)} + c^3d*f^2e^{(4e)})e^{(4fx)} + 2*(d^4f^2x^3e^{(2e)} + 3c*d^3f^2x^2e^{(2e)} + 3c^2d^2f^2xe^{(2e)} + c^3d*f^2e^{(2e)})e^{(2fx)}) - \text{integrate}(2*(d^2f^2x^2 + 2c*d*f^2x + c^2f^2 + 3d^2)/(d^4f^2x^4 + 4c*d^3f^2x^3 + 6c^2d^2f^2x^2 + 4c^3d*f^2x + c^4f^2 + (d^4f^2x^4e^{(2e)} + 4c*d^3f^2x^3e^{(2e)} + 6c^2d^2f^2x^2e^{(2e)} + 4c^3d*f^2xe^{(2e)} + c^4f^2e^{(2e)})e^{(2fx)}), x)$

Giac [N/A]

Not integrable

Time = 0.52 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(fx + e)^3}{(dx + c)^2} dx$$

[In] integrate(tanh(f*x+e)^3/(d*x+c)^2,x, algorithm="giac")

[Out] integrate(tanh(f*x + e)^3/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.12

$$\int \frac{\tanh^3(e + fx)}{(c + dx)^2} dx = \int \frac{\tanh(e + fx)^3}{(c + dx)^2} dx$$

[In] int(tanh(e + f*x)^3/(c + d*x)^2,x)

[Out] int(tanh(e + f*x)^3/(c + d*x)^2, x)

3.16 $\int (c + dx)(b \tanh(e + fx))^{5/2} dx$

Optimal result	121
Rubi [A] (verified)	122
Mathematica [F]	132
Maple [F]	132
Fricas [F(-2)]	132
Sympy [F]	132
Maxima [F]	133
Giac [F]	133
Mupad [F(-1)]	133

Optimal result

Integrand size = 18, antiderivative size = 1392

$$\begin{aligned}
& \int (c + dx)(b \tanh(e + fx))^{5/2} dx = \frac{2b^{5/2} d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} \\
& - \frac{(-b)^{5/2}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
& - \frac{(-b)^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} \\
& + \frac{b^{5/2}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
& - \frac{b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& + \frac{b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& - \frac{b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& - \frac{b^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& + \frac{(-b)^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
& - \frac{(-b)^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
& - \frac{(-b)^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
& - \frac{(-b)^{5/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
& - \frac{b^{5/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& - \frac{b^{5/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& + \frac{b^{5/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2}
\end{aligned}$$

```
[Out] 2/3*b^(5/2)*d*arctan((b*tanh(f*x+e))^(1/2)/b^(1/2))/f^2-(-b)^(5/2)*(d*x+c)*
arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/f-1/2*(-b)^(5/2)*d*arctanh((b*tan
h(f*x+e))^(1/2)/(-b)^(1/2))^2/f^2+2/3*b^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/
2)/b^(1/2))/f^2+b^(5/2)*(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/f+1/
2*b^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))^2/f^2-b^(5/2)*d*arctanh(
(b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2))
)/f^2+b^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2
)+(b*tanh(f*x+e))^(1/2)))/f^2-1/2*b^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b
^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2
))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2-1/2*b^(5/2)*d*arctanh((b*tanh(f*x+e)
)^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2
)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2+(-b)^(5/2)*d*arctanh((b*tan
h(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-
1/2*(-b)^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b
*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/
2)))/f^2-1/2*(-b)^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(
b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2
)/(-b)^(1/2)))/f^2-(-b)^(5/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*l
n(2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/2*b^(5/2)*d*polylog(2,1-2*b
^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^2-1/2*b^(5/2)*d*polylog(2,1-2*b^(
1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2+1/4*b^(5/2)*d*polylog(2,1-2*b^(1/
2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tan
h(f*x+e))^(1/2)))/f^2+1/4*b^(5/2)*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tan
h(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2+
1/2*(-b)^(5/2)*d*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/
4*(-b)^(5/2)*d*polylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b
^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/4*(-b)^(5/2)*d*polylog(2
,1+2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e)
)^(1/2)/(-b)^(1/2)))/f^2+1/2*(-b)^(5/2)*d*polylog(2,1-2/(1+(b*tanh(f*x+e))
^(1/2)/(-b)^(1/2)))/f^2-4/3*b^2*d*(b*tanh(f*x+e))^(1/2)/f^2-2/3*b*(d*x+c)*(b
*tanh(f*x+e))^(3/2)/f
```

Rubi [A] (verified)

Time = 1.32 (sec) , antiderivative size = 1392, normalized size of antiderivative = 1.00, number of steps used = 44, number of rules used = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {3801, 3554, 3557, 335, 218, 212, 209, 3817, 213, 281, 6857, 6139, 6057, 2449, 2352,

2497, 6131, 6055}

$$\begin{aligned}
& \int (c + dx)(b \tanh(e + fx))^{5/2} dx = \\
& \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2 (-b)^{5/2}}{2f^2} \\
& - \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) (-b)^{5/2}}{f} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) (-b)^{5/2}}{f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) (-b)^{5/2}}{2f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) (-b)^{5/2}}{2f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) (-b)^{5/2}}{f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) (-b)^{5/2}}{2f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) (-b)^{5/2}}{4f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, \frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})} + 1\right) (-b)^{5/2}}{4f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) (-b)^{5/2}}{2f^2} \\
& + \frac{b^{5/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} - \frac{2b(c + dx)(b \tanh(e + fx))^{3/2}}{3f} \\
& + \frac{2b^{5/2} d \operatorname{arctan}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} + \frac{b^{5/2}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
& + \frac{2b^{5/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} \\
& - \frac{b^{5/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{f^2}
\end{aligned}$$

[In] Int[(c + d*x)*(b*Tanh[e + f*x])^(5/2), x]

[Out] $(2*b^{(5/2)*d}*ArcTan[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/(3*f^2) - ((-b)^{(5/2)*}(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]])/f - ((-b)^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/(2*f^2) + (2*b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/(3*f^2) + (b^{(5/2)*}(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/f + (b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]^2)/(2*f^2) - (b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/f^2 + (b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/f^2 - (b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/((2*f^2) - (b^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/((2*f^2) + ((-b)^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^2 - ((-b)^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/((2*f^2) - ((-b)^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(-2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/((2*f^2) - ((-b)^{(5/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^2 - (b^{(5/2)*d}*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/((2*f^2) - (b^{(5/2)*d}*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])))/(4*f^2) + (b^{(5/2)*d}*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])))/(4*f^2) + ((-b)^{(5/2)*d}*PolyLog[2, 1 - 2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((2*f^2) - ((-b)^{(5/2)*d}*PolyLog[2, 1 - (2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(4*f^2) - ((-b)^{(5/2)*d}*PolyLog[2, 1 + (2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(4*f^2) + ((-b)^{(5/2)*d}*PolyLog[2, 1 - 2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((2*f^2) - (4*b^2*d*Sqrt[b*Tanh[e + f*x]])/(3*f^2) - (2*b*(c + d*x)*(b*Tanh[e + f*x])^(3/2))/(3*f)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

$\text{Int}[\left((a_) + (b_)*(x_)^2\right)^{-1}, x_Symbol] \rightarrow \text{Simp}\left[\left(-\text{Rt}[-a, 2]*\text{Rt}[b, 2]\right)^{-1}\right]*\text{ArcTanh}\left[\text{Rt}[b, 2]*(x/\text{Rt}[-a, 2])\right], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 218

$\text{Int}[\left((a_) + (b_)*(x_)^4\right)^{-1}, x_Symbol] \rightarrow \text{With}\left[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Dist}\left[r/(2*a), \text{Int}\left[1/(r - s*x^2), x\right], x\right] + \text{Dist}\left[r/(2*a), \text{Int}\left[1/(r + s*x^2), x\right], x\right] /;$ FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 281

$\text{Int}\left[(x_)^{(m_)}*\left((a_) + (b_)*(x_)^{(n_)}\right)^{(p_)}, x_Symbol\right] \rightarrow \text{With}\left[\{k = \text{GCD}[m + 1, n]\}, \text{Dist}\left[1/k, \text{Subst}\left[\text{Int}\left[x^{(m+1)/k - 1}*(a + b*x^{(n/k)})^p, x\right], x, x^{1/k}\right], x\right] /;$ k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 335

$\text{Int}\left[\left((c_)*(x_)^{(m_)}*\left((a_) + (b_)*(x_)^{(n_)}\right)^{(p_)}\right), x_Symbol\right] \rightarrow \text{With}\left[\{k = \text{Denominator}[m]\}, \text{Dist}\left[k/c, \text{Subst}\left[\text{Int}\left[x^{(k*(m+1) - 1}*(a + b*(x^{(k*n)}/c^n))^p, x\right], x, (c*x)^{1/k}\right], x\right] /;$ FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2352

$\text{Int}\left[\text{Log}\left[(c_)*(x_)\right]/\left((d_) + (e_)*(x_)\right), x_Symbol\right] \rightarrow \text{Simp}\left[\left(-e^{-1}\right)*\text{PolyLog}\left[2, 1 - c*x\right], x\right] /;$ FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]

Rule 2449

$\text{Int}\left[\text{Log}\left[(c_)/\left((d_) + (e_)*(x_)\right)\right]/\left((f_) + (g_)*(x_)^2\right), x_Symbol\right] \rightarrow \text{Dist}\left[-e/g, \text{Subst}\left[\text{Int}\left[\text{Log}\left[2*d*x\right]/\left(1 - 2*d*x\right), x\right], x, 1/(d + e*x)\right], x\right] /;$ FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]

Rule 2497

$\text{Int}\left[\text{Log}\left[u_*(Pq_)\right]^{(m_)}, x_Symbol\right] \rightarrow \text{With}\left[\{C = \text{FullSimplify}\left[Pq^m*((1 - u)/D[u, x])\right]\}, \text{Simp}\left[C*\text{PolyLog}\left[2, 1 - u\right], x\right] /;$ FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && ! IntegerQ[n]

Rule 3801

Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rule 3817

Int[((c_.) + (d_.)*(x_))*Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(-I)*Rt[a - I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x] + (Dist[I*d*(Rt[a - I*b, 2]/f), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*d*(Rt[a + I*b, 2]/f), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*Rt[a + I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]

Rule 6055

Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)^((p_.)/((d_.) + (e_.)*(x_))), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]

Rule 6057

Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[(-(a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e), Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x))/((c*d + e)*(1 + c*x))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x))/((c*d + e)*(1 + c*x))]/e), x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]

Rule 6131

```
Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.))/((d_.) + (e_.)*(x_.)^2),
  x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/
(c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e
}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Rule 6139

```
Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*(x_.)^(m_.))/((d_.) + (e_.)*(x_.)^2),
  x_Symbol] := Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x],
x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0]
)
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{2b(c+dx)(b \tanh(e+fx))^{3/2}}{3f} \\
&+ b^2 \int (c+dx) \sqrt{b \tanh(e+fx)} dx + \frac{(2bd) \int (b \tanh(e+fx))^{3/2} dx}{3f} \\
&= -\frac{(-b)^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} + \frac{b^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&- \frac{4b^2 d \sqrt{b \tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b \tanh(e+fx))^{3/2}}{3f} \\
&+ \frac{((-b)^{5/2}d) \int \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) dx}{f} \\
&- \frac{(b^{5/2}d) \int \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) dx}{f} + \frac{(2b^3d) \int \frac{1}{\sqrt{b \tanh(e+fx)}} dx}{3f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(-b)^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} + \frac{b^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{4b^2d\sqrt{b\tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b\tanh(e+fx))^{3/2}}{3f} \\
&\quad + \frac{((-b)^{5/2}d)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2}dx, x, \tanh(e+fx)\right)}{f^2} \\
&\quad - \frac{(b^{5/2}d)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2}dx, x, \tanh(e+fx)\right)}{f^2} \\
&\quad - \frac{(2b^4d)\operatorname{Subst}\left(\int\frac{1}{\sqrt{x(-b^2+x^2)}}dx, x, b\tanh(e+fx)\right)}{3f^2} \\
&= -\frac{(-b)^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} + \frac{b^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{4b^2d\sqrt{b\tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b\tanh(e+fx))^{3/2}}{3f} \\
&\quad - \frac{(2(-b)^{3/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{(2b^{3/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{(4b^4d)\operatorname{Subst}\left(\int\frac{1}{-b^2+x^4}dx, x, \sqrt{b\tanh(e+fx)}\right)}{3f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(-b)^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} + \frac{b^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{4b^2d\sqrt{b\tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b\tanh(e+fx))^{3/2}}{3f} \\
&\quad - \frac{(2(-b)^{3/2}d)\operatorname{Subst}\left(\int\left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{(2b^{3/2}d)\operatorname{Subst}\left(\int\left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad + \frac{(2b^3d)\operatorname{Subst}\left(\int\frac{1}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{3f^2} \\
&\quad + \frac{(2b^3d)\operatorname{Subst}\left(\int\frac{1}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{3f^2} \\
&= \frac{2b^{5/2}d\operatorname{arctan}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} - \frac{(-b)^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&\quad + \frac{2b^{5/2}d\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} + \frac{b^{5/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{4b^2d\sqrt{b\tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b\tanh(e+fx))^{3/2}}{3f} \\
&\quad - \frac{((-b)^{5/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{((-b)^{5/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{(b^{5/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&\quad - \frac{(b^{5/2}d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^{5/2}d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} - \frac{(-b)^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&- \frac{(-b)^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} \\
&+ \frac{b^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{4b^2d\sqrt{b \tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b \tanh(e+fx))^{3/2}}{3f} \\
&\frac{((-b)^{5/2}d) \operatorname{Subst}\left(\int \left(\frac{\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{2(\sqrt{b-x})} - \frac{\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{2(\sqrt{b+x})}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&\frac{(b^2d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&\frac{(b^2d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&\frac{(b^{5/2}d) \operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b-x})} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b+x})}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^{5/2}d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} - \frac{(-b)^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&- \frac{(-b)^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{3f^2} \\
&+ \frac{b^{5/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{b^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
&- \frac{(-b)^{5/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1+\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
&- \frac{4b^2d\sqrt{b \tanh(e+fx)}}{3f^2} - \frac{2b(c+dx)(b \tanh(e+fx))^{3/2}}{3f} \\
&- \frac{((-b)^{5/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&+ \frac{((-b)^{5/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&+ \frac{(b^2d) \operatorname{Subst}\left(\int \frac{\log\left(\frac{2-x}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&+ \frac{(b^2d) \operatorname{Subst}\left(\int \frac{\log\left(\frac{2-bx}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&+ \frac{(b^{5/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&- \frac{(b^{5/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2}
\end{aligned}$$

= Too large to display

Mathematica [F]

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \int (c + dx)(b \tanh(e + fx))^{5/2} dx$$

[In] Integrate[(c + d*x)*(b*Tanh[e + f*x])^(5/2), x]

[Out] Integrate[(c + d*x)*(b*Tanh[e + f*x])^(5/2), x]

Maple [F]

$$\int (dx + c) (b \tanh (fx + e))^{5/2} dx$$

[In] int((d*x+c)*(b*tanh(f*x+e))^(5/2), x)

[Out] int((d*x+c)*(b*tanh(f*x+e))^(5/2), x)

Fricas [F(-2)]

Exception generated.

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))^(5/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \int (b \tanh(e + fx))^{5/2} (c + dx) dx$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))**(5/2), x)

[Out] Integral((b*tanh(e + f*x))**(5/2)*(c + d*x), x)

Maxima [F]

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \int (dx + c)(b \tanh(fx + e))^{5/2} dx$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))^(5/2),x, algorithm="maxima")

[Out] integrate((d*x + c)*(b*tanh(f*x + e))^(5/2), x)

Giac [F]

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \int (dx + c)(b \tanh(fx + e))^{5/2} dx$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((d*x + c)*(b*tanh(f*x + e))^(5/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(b \tanh(e + fx))^{5/2} dx = \int (b \tanh(e + fx))^{5/2} (c + dx) dx$$

[In] int((b*tanh(e + f*x))^(5/2)*(c + d*x),x)

[Out] int((b*tanh(e + f*x))^(5/2)*(c + d*x), x)

3.17 $\int (c + dx)(b \tanh(e + fx))^{3/2} dx$

Optimal result	135
Rubi [A] (verified)	136
Mathematica [F]	146
Maple [F]	146
Fricas [F(-2)]	146
Sympy [F]	146
Maxima [F]	147
Giac [F]	147
Mupad [F(-1)]	147

Optimal result

Integrand size = 18, antiderivative size = 1363

$$\begin{aligned}
& \int (c + dx)(b \tanh(e + fx))^{3/2} dx = -\frac{2b^{3/2} d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
& - \frac{(-b)^{3/2}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
& - \frac{(-b)^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
& + \frac{b^{3/2}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
& - \frac{b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& + \frac{b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& - \frac{b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& - \frac{b^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& + \frac{(-b)^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
& - \frac{(-b)^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
& - \frac{(-b)^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
& - \frac{(-b)^{3/2} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
& - \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& - \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& + \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2}
\end{aligned}$$

```
[Out] -2*b^(3/2)*d*arctan((b*tanh(f*x+e))^(1/2)/b^(1/2))/f^2-(-b)^(3/2)*(d*x+c)*a
rctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/f-1/2*(-b)^(3/2)*d*arctanh((b*tanh
(f*x+e))^(1/2)/(-b)^(1/2))^2/f^2+2*b^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/
b^(1/2))/f^2+b^(3/2)*(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/f+1/2*b
^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))^2/f^2-b^(3/2)*d*arctanh((b*
tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f
^2+b^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(
b*tanh(f*x+e))^(1/2)))/f^2-1/2*b^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1
/2))*ln(2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2)))/(
b^(1/2)+(b*tanh(f*x+e))^(1/2))/f^2-1/2*b^(3/2)*d*arctanh((b*tanh(f*x+e))^(
1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b
^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2))/f^2+(-b)^(3/2)*d*arctanh((b*tanh(f
*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/2
*(-b)^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*ta
nh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)
))/f^2-1/2*(-b)^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(
1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(
-b)^(1/2)))/f^2-(-b)^(3/2)*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2
/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/2*b^(3/2)*d*polylog(2,1-2*b^(1
/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^2-1/2*b^(3/2)*d*polylog(2,1-2*b^(1/2
)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2+1/4*b^(3/2)*d*polylog(2,1-2*b^(1/2)*
((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*
x+e))^(1/2)))/f^2+1/4*b^(3/2)*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f
*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2+1/2
*(-b)^(3/2)*d*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/4*(
-b)^(3/2)*d*polylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/
2)))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2-1/4*(-b)^(3/2)*d*polylog(2,1+
2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(
1/2)/(-b)^(1/2)))/f^2+1/2*(-b)^(3/2)*d*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/
2)/(-b)^(1/2)))/f^2-2*b*(d*x+c)*(b*tanh(f*x+e))^(1/2)/f
```

Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 1363, normalized size of antiderivative = 1.00, number of steps used = 43, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.944$, Rules used = {3801, 3557, 335, 304, 209, 212, 3819, 213, 281, 6857, 6139, 6057, 2449, 2352, 2497,

6131, 6055}

$$\begin{aligned}
& \int (c + dx)(b \tanh(e + fx))^{3/2} dx = - \frac{(-b)^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} \\
& - \frac{(-b)^{3/2} (c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
& + \frac{(-b)^{3/2} d \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} \\
& - \frac{(-b)^{3/2} d \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2f^2} \\
& - \frac{(-b)^{3/2} d \log\left(-\frac{2(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2f^2} \\
& - \frac{(-b)^{3/2} d \log\left(\frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} \\
& + \frac{b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} - \frac{2b^{3/2} d \operatorname{arctan}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
& + \frac{b^{3/2} (c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{2b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
& - \frac{b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& + \frac{b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
& - \frac{b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& - \frac{b^{3/2} \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
& - \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& - \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
& + \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2} \\
& + \frac{b^{3/2} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2}
\end{aligned}$$

[In] Int[(c + d*x)*(b*Tanh[e + f*x])^(3/2), x]

[Out] $(-2*b^{(3/2)*d}*ArcTan[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/f^2 - ((-b)^{(3/2)}*(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]])/f - ((-b)^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/(2*f^2) + (2*b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/f^2 + (b^{(3/2)}*(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/f + (b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]^2)/(2*f^2) - (b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/f^2 + (b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/f^2 - (b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*f^2) - (b^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*f^2) + ((-b)^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^2 - ((-b)^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(2*f^2) - ((-b)^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(-2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(2*f^2) - ((-b)^{(3/2)*d}*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^2 - (b^{(3/2)*d}*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/((Sqrt[b] - Sqrt[b*Tanh[e + f*x]]))/(2*f^2) - (b^{(3/2)*d}*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/((Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))/(2*f^2) + (b^{(3/2)*d}*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))]))/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*f^2) + (b^{(3/2)*d}*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))]))/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*f^2) + ((-b)^{(3/2)*d}*PolyLog[2, 1 - 2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(2*f^2) - ((-b)^{(3/2)*d}*PolyLog[2, 1 - (2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))]))/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(4*f^2) - ((-b)^{(3/2)*d}*PolyLog[2, 1 + (2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))]))/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(4*f^2) + ((-b)^{(3/2)*d}*PolyLog[2, 1 - 2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))/(2*f^2) - (2*b*(c + d*x)*Sqrt[b*Tanh[e + f*x]])/f$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 304

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2352

Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]

Rule 2449

Int[Log[(c_)]/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] := Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]

Rule 2497

Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x]] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]

Rule 3557

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3819

```
Int[((c_.) + (d_.)*(x_))/Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Sym
bol] := Simp[(-I)*((c + d*x)/(f*Rt[a - I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e +
f*x]]/Rt[a - I*b, 2]], x] + (Dist[I*(d/(f*Rt[a - I*b, 2])), Int[ArcTanh[Sq
rt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*(d/(f*Rt[a + I*b, 2
])), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*
((c + d*x)/(f*Rt[a + I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b,
2]], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 6055

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^((p_.)/((d_.) + (e_.)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c
*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2
)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2,
0]
```

Rule 6057

```
Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)/((d_.) + (e_.)*(x_)), x_Symbol] := S
imp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e), Int[Log
[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x
)/((c*d + e)*(1 + c*x))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])
*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x))]/e), x]) /; FreeQ[{a, b, c, d,
e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Rule 6131

```
Int((((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^((p_.)*(x_))/((d_.) + (e_.)*(x_)^2),
x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/
(c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e
}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Rule 6139

```
Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*(x_)^(m_.))/((d_) + (e_.)*(x_)^2),
  x_Symbol] := Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x],
  x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0]
)
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{2b(c+dx)\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{c+dx}{\sqrt{b\tanh(e+fx)}} dx \\
&+ \frac{(2bd) \int \sqrt{b\tanh(e+fx)} dx}{f} \\
&= -\frac{(-b)^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} + \frac{b^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&- \frac{2b(c+dx)\sqrt{b\tanh(e+fx)}}{f} - \frac{(2b^2d) \operatorname{Subst}\left(\int \frac{\sqrt{x}}{-b^2+x^2} dx, x, b\tanh(e+fx)\right)}{f^2} \\
&+ \frac{((-b)^{3/2}d) \int \operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right) dx}{f} - \frac{(b^{3/2}d) \int \operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right) dx}{f} \\
&= -\frac{(-b)^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&+ \frac{b^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} - \frac{2b(c+dx)\sqrt{b\tanh(e+fx)}}{f} \\
&+ \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2} dx, x, \tanh(e+fx)\right)}{f^2} \\
&- \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2} dx, x, \tanh(e+fx)\right)}{f^2} \\
&- \frac{(4b^2d) \operatorname{Subst}\left(\int \frac{x^2}{-b^2+x^4} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= - \frac{(-b)^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&+ \frac{b^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} - \frac{2b(c+dx)\sqrt{b\tanh(e+fx)}}{f} \\
&- \frac{(2\sqrt{-bd}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&- \frac{(2\sqrt{bd}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&+ \frac{(2b^2d) \operatorname{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&- \frac{(2b^2d) \operatorname{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&= - \frac{2b^{3/2}d \operatorname{arctan}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} - \frac{(-b)^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&+ \frac{2b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&+ \frac{b^{3/2}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f} - \frac{2b(c+dx)\sqrt{b\tanh(e+fx)}}{f} \\
&- \frac{(2\sqrt{-bd}) \operatorname{Subst}\left(\int \left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2} \\
&- \frac{(2\sqrt{bd}) \operatorname{Subst}\left(\int \left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^{3/2}d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} - \frac{(-b)^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&+ \frac{2b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&+ \frac{b^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} - \frac{2b(c+dx)\sqrt{b \tanh(e+fx)}}{f} \\
&- \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^{3/2}d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} - \frac{(-b)^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&- \frac{(-b)^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&+ \frac{b^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} \\
&+ \frac{b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} - \frac{2b(c+dx)\sqrt{b \tanh(e+fx)}}{f} \\
&- \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}-x)} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}+x)}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{(bd) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&+ \frac{(bd) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{b}-x)} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{b}+x)}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^{3/2}d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} - \frac{(-b)^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&- \frac{(-b)^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} + \frac{2b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&+ \frac{b^{3/2}(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{b^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
&- \frac{(-b)^{3/2}d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1+\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^2} \\
&- \frac{2b(c+dx)\sqrt{b \tanh(e+fx)}}{f} \\
&- \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&+ \frac{((-b)^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&+ \frac{(bd) \operatorname{Subst}\left(\int \frac{\log\left(\frac{2-x}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&- \frac{(bd) \operatorname{Subst}\left(\int \frac{\log\left(\frac{2-bx}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{f^2} \\
&+ \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2} \\
&- \frac{(b^{3/2}d) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2f^2}
\end{aligned}$$

= Too large to display

Mathematica [F]

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \int (c + dx)(b \tanh(e + fx))^{3/2} dx$$

```
[In] Integrate[(c + d*x)*(b*Tanh[e + f*x])^(3/2), x]
```

```
[Out] Integrate[(c + d*x)*(b*Tanh[e + f*x])^(3/2), x]
```

Maple [F]

$$\int (dx + c) (b \tanh (fx + e))^{\frac{3}{2}} dx$$

```
[In] int((d*x+c)*(b*tanh(f*x+e))^(3/2), x)
```

```
[Out] int((d*x+c)*(b*tanh(f*x+e))^(3/2), x)
```

Fricas [F(-2)]

Exception generated.

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \text{Exception raised: TypeError}$$

```
[In] integrate((d*x+c)*(b*tanh(f*x+e))^(3/2), x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Error detected within library code:  integ
rate: implementation incomplete (constant residues)
```

Sympy [F]

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \int (b \tanh(e + fx))^{\frac{3}{2}} (c + dx) dx$$

```
[In] integrate((d*x+c)*(b*tanh(f*x+e))**(3/2), x)
```

```
[Out] Integral((b*tanh(e + f*x))**(3/2)*(c + d*x), x)
```

Maxima [F]

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \int (dx + c)(b \tanh(fx + e))^{\frac{3}{2}} dx$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*x + c)*(b*tanh(f*x + e))^(3/2), x)

Giac [F]

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \int (dx + c)(b \tanh(fx + e))^{\frac{3}{2}} dx$$

[In] integrate((d*x+c)*(b*tanh(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*x + c)*(b*tanh(f*x + e))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(b \tanh(e + fx))^{3/2} dx = \int (b \tanh(e + fx))^{3/2} (c + dx) dx$$

[In] int((b*tanh(e + f*x))^(3/2)*(c + d*x),x)

[Out] int((b*tanh(e + f*x))^(3/2)*(c + d*x), x)

3.18 $\int (c + dx) \sqrt{b \tanh(e + fx)} dx$

Optimal result	149
Rubi [A] (verified)	150
Mathematica [C] (verified)	158
Maple [F]	158
Fricas [F(-2)]	159
Sympy [F]	159
Maxima [F]	159
Giac [F]	159
Mupad [F(-1)]	160

Optimal result

Integrand size = 18, antiderivative size = 1280

$$\begin{aligned}
& \int (c + dx) \sqrt{b \tanh(e + fx)} dx \\
&= -\frac{\sqrt{-b}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}{f} - \frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)^2}{2f^2} \\
&+ \frac{\sqrt{b}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e + fx)}}\right)}{f^2} \\
&+ \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e + fx)}}\right)}{f^2} \\
&- \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{2f^2} \\
&- \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{2f^2} \\
&+ \frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}}\right)}{f^2} \\
&- \frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
&- \frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}\right)}{2f^2} \\
&- \frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}}\right)}{f^2} \\
&- \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{-b} - \sqrt{b \tanh(e + fx)}}\right)}{2f^2} - \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e + fx)}}\right)}{2f^2} \\
&+ \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{4f^2} \\
&+ \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{4f^2} \\
&+ \frac{\sqrt{-b} d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}}\right)}{2f^2}
\end{aligned}$$

```
[Out] -(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*(-b)^(1/2)/f-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))^2*(-b)^(1/2)/f^2+d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))*(-b)^(1/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*(-b)^(1/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*(-b)^(1/2)/f^2-d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))*(-b)^(1/2)/f^2+1/2*d*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))*(-b)^(1/2)/f^2-1/4*d*polylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*(-b)^(1/2)/f^2-1/4*d*polylog(2,1+2*(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*(-b)^(1/2)/f^2+2+1/2*d*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))*(-b)^(1/2)/f^2+(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*b^(1/2)/f+1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))^2*b^(1/2)/f^2-d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))*b^(1/2)/f^2+d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))*b^(1/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2))*b^(1/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2))*b^(1/2)/f^2-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))*b^(1/2)/f^2-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))*b^(1/2)/f^2+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2))*b^(1/2)/f^2+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2)))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2))*b^(1/2)/f^2
```

Rubi [A] (verified)

Time = 0.97 (sec) , antiderivative size = 1280, normalized size of antiderivative = 1.00, number of steps used = 37, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules

used = {3817, 213, 281, 6857, 6139, 6057, 2449, 2352, 2497, 6131, 6055, 212}

$$\begin{aligned}
& \int (c + dx) \sqrt{b \tanh(e + fx)} dx \\
&= -\frac{\sqrt{-b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2f^2} - \frac{\sqrt{-b}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f} \\
&+ \frac{\sqrt{-b} d \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} \\
&- \frac{\sqrt{-b} d \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2f^2} \\
&- \frac{\sqrt{-b} d \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2f^2} \\
&- \frac{\sqrt{-b} d \log\left(\frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2f^2} \\
&+ \frac{\sqrt{b}(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f} - \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
&+ \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^2} \\
&- \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
&- \frac{\sqrt{b} d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2f^2} \\
&- \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{2f^2} - \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2f^2} \\
&+ \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2} \\
&+ \frac{\sqrt{b} d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4f^2} \\
&+ \frac{\sqrt{-b} d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{2f^2} \\
&- \frac{\sqrt{-b} d \operatorname{PolyLog}\left(2, 1 - \frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right)}{4f^2}
\end{aligned}$$

[In] Int[(c + d*x)*Sqrt[b*Tanh[e + f*x]],x]

[Out]
$$-\left(\frac{\sqrt{-b}(c + dx)\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]}{f} - \sqrt{-b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]^2\right)/(2f^2) + \left(\frac{\sqrt{b}(c + dx)\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]}{f} + \sqrt{b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]^2\right)/(2f^2) - \left(\frac{\sqrt{b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]\operatorname{Log}\left[\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b\tanh(e + fx)}}\right]}{f^2} + \sqrt{b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]\operatorname{Log}\left[\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b\tanh(e + fx)}}\right]}{f^2} - \sqrt{b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]\operatorname{Log}\left[\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b\tanh(e + fx)})}\right]}{(2f^2)} - \sqrt{b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{b}}\right]\operatorname{Log}\left[\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b\tanh(e + fx)})}\right]}{(2f^2)} + \sqrt{-b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]\operatorname{Log}\left[\frac{2}{(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{f^2} - \sqrt{-b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]\operatorname{Log}\left[\frac{2(\sqrt{b} - \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(2f^2)} - \sqrt{-b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]\operatorname{Log}\left[\frac{-2(\sqrt{b} + \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(2f^2)} - \sqrt{-b}d\operatorname{ArcTanh}\left[\frac{\sqrt{b\tanh(e + fx)}}{\sqrt{-b}}\right]\operatorname{Log}\left[\frac{2}{(1 + \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{f^2} - \sqrt{b}d\operatorname{PolyLog}\left[2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b\tanh(e + fx)}}\right]}{(2f^2)} - \sqrt{b}d\operatorname{PolyLog}\left[2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b\tanh(e + fx)}}\right]}{(2f^2)} + \sqrt{b}d\operatorname{PolyLog}\left[2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b\tanh(e + fx)})}\right]}{(4f^2)} + \sqrt{b}d\operatorname{PolyLog}\left[2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b\tanh(e + fx)})}\right]}{(4f^2)} + \sqrt{-b}d\operatorname{PolyLog}\left[2, 1 - \frac{2}{(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(2f^2)} - \sqrt{-b}d\operatorname{PolyLog}\left[2, 1 - \frac{2(\sqrt{b} - \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(4f^2)} - \sqrt{-b}d\operatorname{PolyLog}\left[2, 1 + \frac{2(\sqrt{b} + \sqrt{b\tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(4f^2)} + \sqrt{-b}d\operatorname{PolyLog}\left[2, 1 - \frac{2}{(1 + \sqrt{b\tanh(e + fx)})/\sqrt{-b}}\right]}{(2f^2)}\right]$$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 281


```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 2352

```
Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLo
g[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] := Dist
[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{
c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_.), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/
D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] &&
PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u,
x][[2]], Expon[Pq, x]]
```

Rule 3817

```
Int[((c_.) + (d_.)*(x_))*Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Sym
bol] := Simp[(-I)*Rt[a - I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f
*x]]/Rt[a - I*b, 2]], x] + (Dist[I*d*(Rt[a - I*b, 2]/f), Int[ArcTanh[Sqrt[a
+ b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*d*(Rt[a + I*b, 2]/f), I
nt[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*Rt[a +
I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x]
) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 6055

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c
*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2
)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2,
0]
```

Rule 6057

```
Int[((a_.) + ArcTanh[(c_.)*(x_)]*(b_.))/((d_) + (e_.)*(x_)), x_Symbol] := S
imp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e), Int[L
og[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x
)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])
```

$*(\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x]) /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{NeQ}[c^2*d^2 - e^2, 0]$

Rule 6131

$\text{Int}[(((a_.) + \text{ArcTanh}[(c_.)*(x_.)]*(b_.))^{(p_.)*(x_.)})/((d_.) + (e_.)*(x_.)^2), x_Symbol] \text{:>} \text{Simp}[(a + b*\text{ArcTanh}[c*x])^{(p + 1)}/(b*e*(p + 1)), x] + \text{Dist}[1/(c*d), \text{Int}[(a + b*\text{ArcTanh}[c*x])^p/(1 - c*x), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[c^2*d + e, 0] \&\& \text{IGtQ}[p, 0]$

Rule 6139

$\text{Int}[(((a_.) + \text{ArcTanh}[(c_.)*(x_.)]*(b_.))*(x_.)^{(m_.)})/((d_.) + (e_.)*(x_.)^2), x_Symbol] \text{:>} \text{Int}[\text{ExpandIntegrand}[a + b*\text{ArcTanh}[c*x], x^m/(d + e*x^2), x], x] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{IntegerQ}[m] \&\& !(\text{EqQ}[m, 1] \&\& \text{NeQ}[a, 0])$

Rule 6857

$\text{Int}[(u_)/((a_) + (b_.)*(x_.)^{(n_)}), x_Symbol] \text{:>} \text{With}[\{v = \text{RationalFunctionExpand}[u/(a + b*x^n), x]\}, \text{Int}[v, x] /; \text{SumQ}[v]] /; \text{FreeQ}\{a, b\}, x] \&\& \text{IGtQ}[n, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{\sqrt{-b}(c + dx)\text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} + \frac{\sqrt{b}(c + dx)\text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} \\ &+ \frac{(\sqrt{-bd}) \int \text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right) dx}{f} - \frac{(\sqrt{bd}) \int \text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right) dx}{f} \\ &= -\frac{\sqrt{-b}(c + dx)\text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} + \frac{\sqrt{b}(c + dx)\text{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} \\ &+ \frac{(\sqrt{-bd}) \text{Subst}\left(\int \frac{\text{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2} dx, x, \tanh(e + fx)\right)}{f^2} \\ &- \frac{(\sqrt{bd}) \text{Subst}\left(\int \frac{\text{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2} dx, x, \tanh(e + fx)\right)}{f^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{-b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} + \frac{\sqrt{b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{-b}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{b}f^2} \\
&= -\frac{\sqrt{-b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} + \frac{\sqrt{b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{-b}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{b}f^2} \\
&= -\frac{\sqrt{-b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} + \frac{\sqrt{b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} \\
&\quad - \frac{(\sqrt{-bd})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&\quad - \frac{(\sqrt{-bd})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&\quad - \frac{(\sqrt{bd})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&\quad - \frac{(\sqrt{bd})\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{-b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} - \frac{\sqrt{-b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{2f^2} \\
&+ \frac{\sqrt{b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} + \frac{\sqrt{b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{d\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&- \frac{d\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&- \frac{(\sqrt{-bd})\operatorname{Subst}\left(\int \left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}-x)} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}+x)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&- \frac{(\sqrt{bd})\operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}-x)} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}+x)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\sqrt{-b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f} - \frac{\sqrt{-b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{2f^2} \\
&+ \frac{\sqrt{b}(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f} + \frac{\sqrt{b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{2f^2} \\
&- \frac{\sqrt{b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)\log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b}\tanh(e+fx)}\right)}{f^2} \\
&- \frac{\sqrt{-b}d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)\log\left(\frac{2}{1+\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}}\right)}{f^2} \\
&+ \frac{d\operatorname{Subst}\left(\int\frac{\log\left(\frac{2}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&+ \frac{d\operatorname{Subst}\left(\int\frac{\log\left(\frac{2}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
&- \frac{(\sqrt{-bd})\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2f^2} \\
&+ \frac{(\sqrt{-bd})\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2f^2} \\
&+ \frac{(\sqrt{bd})\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2f^2} \\
&- \frac{(\sqrt{bd})\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2f^2}
\end{aligned}$$

= Too large to display

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 4.71 (sec) , antiderivative size = 556, normalized size of antiderivative = 0.43

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx$$

$$= \frac{\left(-4f(c + dx) \left(2 \arctan \left(\sqrt{\tanh(e + fx)} \right) + \log \left(1 - \sqrt{\tanh(e + fx)} \right) - \log \left(1 + \sqrt{\tanh(e + fx)} \right) \right) + \dots \right)}{\dots}$$

```
[In] Integrate[(c + d*x)*Sqrt[b*Tanh[e + f*x]],x]
```

```
[Out] ((-4*f*(c + d*x)*(2*ArcTan[Sqrt[Tanh[e + f*x]]] + Log[1 - Sqrt[Tanh[e + f*x]]] - Log[1 + Sqrt[Tanh[e + f*x]]]) + d*((4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]^2 - 4*ArcTan[Sqrt[Tanh[e + f*x]]]*Log[1 + E^((4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]])] - Log[1 - Sqrt[Tanh[e + f*x]]]^2 + 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1/2 + I/2)*(-I + Sqrt[Tanh[e + f*x]])] + 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1/2 - I/2)*(I + Sqrt[Tanh[e + f*x]])] - 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1 + Sqrt[Tanh[e + f*x]])/2] - 2*Log[1 - (1/2 - I/2)*(1 + Sqrt[Tanh[e + f*x]])]*Log[1 + Sqrt[Tanh[e + f*x]]] + 2*Log[(1 - Sqrt[Tanh[e + f*x]])/2]*Log[1 + Sqrt[Tanh[e + f*x]]] - 2*Log[(-1/2 - I/2)*(I + Sqrt[Tanh[e + f*x]])]*Log[1 + Sqrt[Tanh[e + f*x]]] + Log[1 + Sqrt[Tanh[e + f*x]]]^2 + I*PolyLog[2, -E^((4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]])] - 2*PolyLog[2, (1 - Sqrt[Tanh[e + f*x]])/2] + 2*PolyLog[2, (-1/2 - I/2)*(-1 + Sqrt[Tanh[e + f*x]])] + 2*PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[Tanh[e + f*x]])] + 2*PolyLog[2, (1 + Sqrt[Tanh[e + f*x]])/2] - 2*PolyLog[2, (1/2 - I/2)*(1 + Sqrt[Tanh[e + f*x]])] - 2*PolyLog[2, (1/2 + I/2)*(1 + Sqrt[Tanh[e + f*x]])]))*Sqrt[b*Tanh[e + f*x]])/(8*f^2*Sqrt[Tanh[e + f*x]])
```

Maple [F]

$$\int (dx + c) \sqrt{b \tanh(fx + e)} dx$$

```
[In] int((d*x+c)*(b*tanh(f*x+e))^(1/2),x)
```

```
[Out] int((d*x+c)*(b*tanh(f*x+e))^(1/2),x)
```

Fricas [F(-2)]

Exception generated.

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx = \text{Exception raised: TypeError}$$

[In] `integrate((d*x+c)*(b*tanh(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx = \int \sqrt{b \tanh(e + fx)} (c + dx) dx$$

[In] `integrate((d*x+c)*(b*tanh(f*x+e))**(1/2),x)`

[Out] `Integral(sqrt(b*tanh(e + f*x))*(c + d*x), x)`

Maxima [F]

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx = \int (dx + c) \sqrt{b \tanh(fx + e)} dx$$

[In] `integrate((d*x+c)*(b*tanh(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate((d*x + c)*sqrt(b*tanh(f*x + e)), x)`

Giac [F]

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx = \int (dx + c) \sqrt{b \tanh(fx + e)} dx$$

[In] `integrate((d*x+c)*(b*tanh(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate((d*x + c)*sqrt(b*tanh(f*x + e)), x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx) \sqrt{b \tanh(e + fx)} dx = \int \sqrt{b \tanh(e + fx)} (c + dx) dx$$

```
[In] int((b*tanh(e + f*x))^(1/2)*(c + d*x),x)
```

```
[Out] int((b*tanh(e + f*x))^(1/2)*(c + d*x), x)
```


3.19 $\int \frac{c+dx}{\sqrt{b \tanh(e+fx)}} dx$

Optimal result	162
Rubi [A] (verified)	163
Mathematica [C] (verified)	171
Maple [F]	171
Fricas [F(-2)]	172
Sympy [F]	172
Maxima [F]	172
Giac [F(-2)]	172
Mupad [F(-1)]	173

Optimal result

Integrand size = 18, antiderivative size = 1280

$$\begin{aligned}
\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = & -\frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) - \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)^2}{\sqrt{-b} f} - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)^2}{2\sqrt{-b} f^2} \\
& + \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) - \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right)^2}{\sqrt{b} f} + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right)^2}{2\sqrt{b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e + fx)}}\right)}{\sqrt{b} f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e + fx)}}\right)}{\sqrt{b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{2\sqrt{b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{2\sqrt{b} f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}}\right)}{\sqrt{-b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}\right)}{2\sqrt{-b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}\right)}{2\sqrt{-b} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}}\right)}{\sqrt{-b} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e + fx)}}\right)}{2\sqrt{b} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e + fx)}}\right)}{2\sqrt{b} f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{4\sqrt{b} f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e + fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e + fx)})}\right)}{4\sqrt{b} f^2}
\end{aligned}$$

```
[Out] -(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/f/(-b)^(1/2)-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))^2/f^2/(-b)^(1/2)+d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2)))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2)))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)-d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)+1/2*d*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)-1/4*d*polylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2)))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)-1/4*d*polylog(2,1+2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2)))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)+1/2*d*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^2/(-b)^(1/2)+(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/f/b^(1/2)+1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))^2/f^2/b^(1/2)-d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)+d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2)))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2)))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2)))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2)))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^2/b^(1/2)
```

Rubi [A] (verified)

Time = 0.96 (sec) , antiderivative size = 1280, normalized size of antiderivative = 1.00, number of steps used = 37, number of rules used = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules

used = {3819, 213, 281, 6857, 6139, 6057, 2449, 2352, 2497, 6131, 6055, 212}

$$\begin{aligned}
\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = & - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2\sqrt{-b}f^2} - \frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f} \\
& + \frac{d \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f^2} \\
& - \frac{d \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2\sqrt{-b}f^2} \\
& - \frac{d \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2\sqrt{-b}f^2} \\
& - \frac{d \log\left(\frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2\sqrt{b}f^2} + \frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b}f} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{\sqrt{b}f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{\sqrt{b}f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2\sqrt{b}f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2\sqrt{b}f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{2\sqrt{b}f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2\sqrt{b}f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4\sqrt{b}f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4\sqrt{b}f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{4\sqrt{b}f^2}
\end{aligned}$$

[In] Int[(c + d*x)/Sqrt[b*Tanh[e + f*x]],x]

[Out] -(((c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]/(Sqrt[-b]*f)) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/(2*Sqrt[-b]*f^2) + ((c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]/(Sqrt[b]*f) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]^2)/(2*Sqrt[b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/(Sqrt[b]*f^2) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/(Sqrt[b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*Sqrt[b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*Sqrt[b]*f^2) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(Sqrt[-b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(2*Sqrt[-b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(-2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(2*Sqrt[-b]*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(Sqrt[-b]*f^2) - (d*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/(2*Sqrt[b]*f^2) - (d*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/(2*Sqrt[b]*f^2) + (d*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*Sqrt[b]*f^2) + (d*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*Sqrt[b]*f^2) + (d*PolyLog[2, 1 - 2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(2*Sqrt[-b]*f^2) - (d*PolyLog[2, 1 - (2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(4*Sqrt[-b]*f^2) - (d*PolyLog[2, 1 + (2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(4*Sqrt[-b]*f^2) + (d*PolyLog[2, 1 - 2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(2*Sqrt[-b]*f^2)

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 281

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 2352

```
Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLo
g[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_)/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] := Dist
[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{
c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/
D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] &&
PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u,
x][[2]], Expon[Pq, x]]
```

Rule 3819

```
Int[((c_) + (d_)*(x_))/Sqrt[(a_) + (b_)*tan[(e_) + (f_)*(x_)], x_Sym
bol] := Simp[(-I)*((c + d*x)/(f*Rt[a - I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e +
f*x]]/Rt[a - I*b, 2]], x] + (Dist[I*(d/(f*Rt[a - I*b, 2])), Int[ArcTanh[Sq
rt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*(d/(f*Rt[a + I*b, 2
])), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*
((c + d*x)/(f*Rt[a + I*b, 2]))*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b,
2]], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]
```

Rule 6055

```
Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))^(p_)/((d_) + (e_)*(x_)), x_Symbol
] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c
*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2
)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2,
0]
```

Rule 6057

```
Int[((a_) + ArcTanh[(c_)*(x_)]*(b_))/((d_) + (e_)*(x_)), x_Symbol] := S
imp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e), Int[L
og[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x
))/((c*d + e)*(1 + c*x))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])
```

$*(\text{Log}[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x]) /;$ FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]

Rule 6131

Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))^(p_.)*(x_.))/((d_.) + (e_.)*(x_.)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/(c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]

Rule 6139

Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*(x_.)^(m_.))/((d_.) + (e_.)*(x_.)^2), x_Symbol] := Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0])

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionEExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{\sqrt{-b}f} + \frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{\sqrt{b}f} \\
 &+ \frac{d \int \operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right) dx}{\sqrt{-b}f} - \frac{d \int \operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right) dx}{\sqrt{b}f} \\
 &= -\frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{\sqrt{-b}f} + \frac{(c + dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{\sqrt{b}f} \\
 &+ \frac{d\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2} dx, x, \tanh(e + fx)\right)}{\sqrt{-b}f^2} \\
 &- \frac{d\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2} dx, x, \tanh(e + fx)\right)}{\sqrt{b}f^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b}f} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{3/2}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{3/2}f^2} \\
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b}f} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{3/2}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int \left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{3/2}f^2} \\
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{\sqrt{-b}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b}f} \\
&\quad - \frac{d\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{\sqrt{-b}f^2} \\
&\quad - \frac{d\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{\sqrt{-b}f^2} \\
&\quad - \frac{d\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{\sqrt{b}f^2} \\
&\quad - \frac{d\operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{\sqrt{b}f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{\sqrt{-b}f} - \frac{d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{2\sqrt{-b}f^2} \\
&+ \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{\sqrt{b}f} + \frac{d\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{2\sqrt{b}f^2} \\
&\frac{d\operatorname{Subst}\left(\int\left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}-x)} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}+x)}\right)dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{-b}f^2} \\
&\frac{d\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{bf^2} \\
&+ \frac{d\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{bf^2} \\
&\frac{d\operatorname{Subst}\left(\int\left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}-x)} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}+x)}\right)dx, x, \sqrt{b}\tanh(e+fx)\right)}{\sqrt{b}f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{\sqrt{-b}f} - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{2\sqrt{-b}f^2} \\
&+ \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{\sqrt{b}f} + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{2\sqrt{b}f^2} \\
&- \frac{\operatorname{darctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)\log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b}\tanh(e+fx)}\right)}{\sqrt{b}f^2} \\
&- \frac{\operatorname{darctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)\log\left(\frac{2}{1+\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}}\right)}{\sqrt{-b}f^2} \\
&- \frac{\operatorname{dSubst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2\sqrt{-b}f^2} \\
&+ \frac{\operatorname{dSubst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2\sqrt{-b}f^2} \\
&+ \frac{\operatorname{dSubst}\left(\int\frac{\log\left(\frac{2}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{bf^2} \\
&- \frac{\operatorname{dSubst}\left(\int\frac{\log\left(\frac{2}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{bf^2} \\
&+ \frac{\operatorname{dSubst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2\sqrt{b}f^2} \\
&- \frac{\operatorname{dSubst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{2\sqrt{b}f^2}
\end{aligned}$$

= Too large to display

Mathematica [C] (verified)

Result contains complex when optimal does not.

Time = 3.38 (sec) , antiderivative size = 556, normalized size of antiderivative = 0.43

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx$$

$$= \frac{\left(4f(c + dx) \left(2 \arctan \left(\sqrt{\tanh(e + fx)}\right) - \log \left(1 - \sqrt{\tanh(e + fx)}\right) + \log \left(1 + \sqrt{\tanh(e + fx)}\right)\right)\right) + c}{8f^2 \sqrt{b \tanh(e + fx)}}$$

[In] Integrate[(c + d*x)/Sqrt[b*Tanh[e + f*x]],x]

[Out] ((4*f*(c + d*x)*(2*ArcTan[Sqrt[Tanh[e + f*x]]] - Log[1 - Sqrt[Tanh[e + f*x]]] + Log[1 + Sqrt[Tanh[e + f*x]]]) + d*((-4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]^2 + 4*ArcTan[Sqrt[Tanh[e + f*x]]]*Log[1 + E^((4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]])] - Log[1 - Sqrt[Tanh[e + f*x]]]^2 + 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1/2 + I/2)*(-I + Sqrt[Tanh[e + f*x]])] + 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1/2 - I/2)*(I + Sqrt[Tanh[e + f*x]])] - 2*Log[1 - Sqrt[Tanh[e + f*x]]]*Log[(1 + Sqrt[Tanh[e + f*x]])/2] - 2*Log[1 - (1/2 - I/2)*(1 + Sqrt[Tanh[e + f*x]])]*Log[1 + Sqrt[Tanh[e + f*x]]] + 2*Log[(1 - Sqrt[Tanh[e + f*x]])/2]*Log[1 + Sqrt[Tanh[e + f*x]]] - 2*Log[(-1/2 - I/2)*(I + Sqrt[Tanh[e + f*x]])]*Log[1 + Sqrt[Tanh[e + f*x]]] + Log[1 + Sqrt[Tanh[e + f*x]]]^2 - I*PolyLog[2, -E^((4*I)*ArcTan[Sqrt[Tanh[e + f*x]]]])] - 2*PolyLog[2, (1 - Sqrt[Tanh[e + f*x]])/2] + 2*PolyLog[2, (-1/2 - I/2)*(-1 + Sqrt[Tanh[e + f*x]])] + 2*PolyLog[2, (-1/2 + I/2)*(-1 + Sqrt[Tanh[e + f*x]])] + 2*PolyLog[2, (1 + Sqrt[Tanh[e + f*x]])/2] - 2*PolyLog[2, (1/2 - I/2)*(1 + Sqrt[Tanh[e + f*x]])] - 2*PolyLog[2, (1/2 + I/2)*(1 + Sqrt[Tanh[e + f*x]])]))*Sqrt[Tanh[e + f*x]])/(8*f^2*Sqrt[b*Tanh[e + f*x]])

Maple [F]

$$\int \frac{dx + c}{\sqrt{b \tanh(fx + e)}} dx$$

[In] int((d*x+c)/(b*tanh(f*x+e))^(1/2),x)

[Out] int((d*x+c)/(b*tanh(f*x+e))^(1/2),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [F]

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))**(1/2),x)

[Out] Integral((c + d*x)/sqrt(b*tanh(e + f*x)), x)

Maxima [F]

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{dx + c}{\sqrt{b \tanh(fx + e)}} dx$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((d*x + c)/sqrt(b*tanh(f*x + e)), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="giac")

[Out] Exception raised: RuntimeError >> an error occurred running a Giac command: INPUT:sage2OUTPUT:index.cc index_m i_lex_is_greater Error: Bad Argument Value

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx$$

```
[In] int((c + d*x)/(b*tanh(e + f*x))^(1/2), x)
```

```
[Out] int((c + d*x)/(b*tanh(e + f*x))^(1/2), x)
```

3.20 $\int \frac{c+dx}{(b \tanh(e+fx))^{3/2}} dx$

Optimal result	175
Rubi [A] (verified)	176
Mathematica [F]	186
Maple [F]	186
Fricas [F(-2)]	186
Sympy [F]	186
Maxima [F]	187
Giac [F]	187
Mupad [F(-1)]	187

Optimal result

Integrand size = 18, antiderivative size = 1365

$$\begin{aligned}
& \int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \frac{2d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
& - \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2(-b)^{3/2} f^2} + \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
& + \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f} + \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2b^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^2} \\
& + \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2b^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2b^{3/2} f^2} \\
& + \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{(-b)^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right)}{2(-b)^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}})}\right)}{2(-b)^{3/2} f^2} \\
& - \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{(-b)^{3/2} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{2b^{3/2} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2b^{3/2} f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4b^{3/2} f^2}
\end{aligned}$$

```
[Out] 2*d*arctan((b*tanh(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f^2-(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/(-b)^(3/2)/f-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))^2/(-b)^(3/2)/f^2+2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f^2+(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f+1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))^2/b^(3/2)/f^2-d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2+d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2+d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-1/2*d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-d*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2-1/2*d*polylog(2,1-2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2+1/4*d*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^2+1/2*d*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-1/4*d*polylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-1/4*d*polylog(2,1+2*(b^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2+1/2*d*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^2-2*(d*x+c)/b/f/(b*tanh(f*x+e))^(1/2)
```

Rubi [A] (verified)

Time = 1.09 (sec) , antiderivative size = 1365, normalized size of antiderivative = 1.00, number of steps used = 43, number of rules used = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.944$, Rules used = {3802, 3557, 335, 218, 212, 209, 3817, 213, 281, 6857, 6139, 6057, 2449, 2352, 2497,

6131, 6055}

$$\begin{aligned}
& \int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2(-b)^{3/2} f^2} \\
& - \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f} \\
& + \frac{d \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f^2} \\
& - \frac{d \log\left(\frac{2(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2(-b)^{3/2} f^2} \\
& - \frac{d \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{2(-b)^{3/2} f^2} \\
& - \frac{d \log\left(\frac{2}{\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}} + 1}\right) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2b^{3/2} f^2} + \frac{2d \operatorname{arctan}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
& + \frac{(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f} + \frac{2 \operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^2} \\
& + \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2b^{3/2} f^2} \\
& - \frac{\operatorname{darctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{2b^{3/2} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{2b^{3/2} f^2} \\
& - \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{2b^{3/2} f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4b^{3/2} f^2} \\
& + \frac{d \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{4b^{3/2} f^2}
\end{aligned}$$

[In] Int[(c + d*x)/(b*Tanh[e + f*x])^(3/2), x]

[Out] (2*d*ArcTan[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/(b^(3/2)*f^2) - ((c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]])/((-b)^(3/2)*f) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/(2*(-b)^(3/2)*f^2) + (2*d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/(b^(3/2)*f^2) + ((c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/(b^(3/2)*f) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]^2)/(2*b^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/(b^(3/2)*f^2) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/(b^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*b^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(2*b^(3/2)*f^2) + (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((-b)^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(2*(-b)^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(-2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(2*(-b)^(3/2)*f^2) - (d*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/((-b)^(3/2)*f^2) - (d*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/(2*b^(3/2)*f^2) - (d*PolyLog[2, 1 - (2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/(2*b^(3/2)*f^2) + (d*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*b^(3/2)*f^2) + (d*PolyLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))])/(4*b^(3/2)*f^2) + (d*PolyLog[2, 1 - 2/(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(2*(-b)^(3/2)*f^2) - (d*PolyLog[2, 1 - (2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(4*(-b)^(3/2)*f^2) - (d*PolyLog[2, 1 + (2*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])]/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/(4*(-b)^(3/2)*f^2) + (d*PolyLog[2, 1 - 2/(1 + Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/(2*(-b)^(3/2)*f^2) - (2*(c + d*x))/(b*f*Sqrt[b*Tanh[e + f*x]])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 213

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 335

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2352

Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]

Rule 2449

Int[Log[(c_)]/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] := Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]

Rule 2497

Int[Log[u_]*(Pq_)^(m_), x_Symbol] := With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]

Rule 3557

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && ! IntegerQ[n]

Rule 3802

Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(c + d*x)^m*((b*Tan[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + (-Dist[d*(m/(b*f*(n + 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n + 1), x], x] - Dist[1/b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n + 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && LtQ[n, -1] && GtQ[m, 0]

Rule 3817

Int[((c_.) + (d_.)*(x_))*Sqrt[(a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]], x_Symbol] := Simp[(-I)*Rt[a - I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x] + (Dist[I*d*(Rt[a - I*b, 2]/f), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a - I*b, 2]], x], x] - Dist[I*d*(Rt[a + I*b, 2]/f), Int[ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x], x] + Simp[I*Rt[a + I*b, 2]*((c + d*x)/f)*ArcTanh[Sqrt[a + b*Tan[e + f*x]]/Rt[a + I*b, 2]], x]) /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0]

Rule 6055

Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^((p_.)/((d_.) + (e_.)*(x_))), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])^p*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]

Rule 6057

Int[((a_.) + ArcTanh[(c_.)*(x_)])*(b_.)/((d_.) + (e_.)*(x_))), x_Symbol] := Simp[(-a + b*ArcTanh[c*x])*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e), Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])*(Log[2*c*((d + e*x)/((c*d + e)*(1 + c*x)))]/e), x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]

Rule 6131

Int((((a_.) + ArcTanh[(c_.)*(x_)])*(b_.))^((p_.)*(x_))/((d_.) + (e_.)*(x_)^2), x_Symbol] := Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/(c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]

Rule 6139

Int[(((a_.) + ArcTanh[(c_.)*(x_.)]*(b_.))*(x_)^(m_.))/((d_.) + (e_.)*(x_)^2),
 x_Symbol] := Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x],
 x] /; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0])

Rule 6857

Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{v = RationalFunctionE
 xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
 [n, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{2(c+dx)}{bf\sqrt{b}\tanh(e+fx)} + \frac{\int(c+dx)\sqrt{b}\tanh(e+fx)dx}{b^2} + \frac{(2d)\int\frac{1}{\sqrt{b}\tanh(e+fx)}dx}{bf} \\
 &= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f} \\
 &\quad - \frac{2(c+dx)}{bf\sqrt{b}\tanh(e+fx)} - \frac{(2d)\operatorname{Subst}\left(\int\frac{1}{\sqrt{x(-b^2+x^2)}}dx, x, b\tanh(e+fx)\right)}{f^2} \\
 &\quad + \frac{d\int\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)dx}{(-b)^{3/2}f} - \frac{d\int\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)dx}{b^{3/2}f} \\
 &= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f} \\
 &\quad - \frac{2(c+dx)}{bf\sqrt{b}\tanh(e+fx)} - \frac{(4d)\operatorname{Subst}\left(\int\frac{1}{-b^2+x^4}dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^2} \\
 &\quad + \frac{d\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2}dx, x, \tanh(e+fx)\right)}{(-b)^{3/2}f^2} \\
 &\quad - \frac{d\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2}dx, x, \tanh(e+fx)\right)}{b^{3/2}f^2}
 \end{aligned}$$

$$\begin{aligned}
&= -\frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2}f} + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} \\
&\quad - \frac{2(c+dx)}{bf\sqrt{b\tanh(e+fx)}} - \frac{(2d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{5/2}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{5/2}f^2} \\
&\quad + \frac{(2d)\operatorname{Subst}\left(\int\frac{1}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{bf^2} \\
&\quad + \frac{(2d)\operatorname{Subst}\left(\int\frac{1}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{bf^2} \\
&= \frac{2d\arctan\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f^2} - \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2}f} \\
&\quad + \frac{2d\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f^2} \\
&\quad + \frac{(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} - \frac{2(c+dx)}{bf\sqrt{b\tanh(e+fx)}} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int\left(-\frac{bx\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{-bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{5/2}f^2} \\
&\quad - \frac{(2d)\operatorname{Subst}\left(\int\left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{5/2}f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} - \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f} \\
&+ \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} + \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f} \\
&- \frac{2(c+dx)}{bf \sqrt{b \tanh(e+fx)}} - \frac{d \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{(-b)^{3/2} f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{(-b)^{3/2} f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^{3/2} f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{x \operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^{3/2} f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} - \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f} \\
&- \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2(-b)^{3/2} f^2} + \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
&+ \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f} \\
&+ \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2b^{3/2} f^2} - \frac{2(c+dx)}{bf \sqrt{b \tanh(e+fx)}} \\
&- \frac{d \operatorname{Subst}\left(\int \left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}-x)} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}+x)}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{(-b)^{3/2} f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^2 f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^2 f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}-x)} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}+x)}\right) dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^{3/2} f^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2d \arctan\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} - \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f} \\
&- \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{2(-b)^{3/2} f^2} + \frac{2d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
&+ \frac{(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f} + \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{2b^{3/2} f^2} \\
&- \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^2} \\
&- \frac{d \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1+\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{(-b)^{3/2} f^2} - \frac{2(c+dx)}{bf \sqrt{b \tanh(e+fx)}} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2(-b)^{3/2} f^2} \\
&+ \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2(-b)^{3/2} f^2} \\
&+ \frac{d \operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^2 f^2} \\
&+ \frac{d \operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{b^2 f^2} \\
&+ \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2b^{3/2} f^2} \\
&- \frac{d \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}} dx, x, \sqrt{b \tanh(e+fx)}\right)}{2b^{3/2} f^2}
\end{aligned}$$

= Too large to display

Mathematica [F]

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx$$

```
[In] Integrate[(c + d*x)/(b*Tanh[e + f*x])^(3/2), x]
```

```
[Out] Integrate[(c + d*x)/(b*Tanh[e + f*x])^(3/2), x]
```

Maple [F]

$$\int \frac{dx + c}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

```
[In] int((d*x+c)/(b*tanh(f*x+e))^(3/2), x)
```

```
[Out] int((d*x+c)/(b*tanh(f*x+e))^(3/2), x)
```

Fricas [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

```
[In] integrate((d*x+c)/(b*tanh(f*x+e))^(3/2), x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Error detected within library code: integ
rate: implementation incomplete (constant residues)
```

Sympy [F]

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{c + dx}{(b \tanh(e + fx))^{\frac{3}{2}}} dx$$

```
[In] integrate((d*x+c)/(b*tanh(f*x+e))**(3/2), x)
```

```
[Out] Integral((c + d*x)/(b*tanh(e + f*x))**(3/2), x)
```

Maxima [F]

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{dx + c}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*x + c)/(b*tanh(f*x + e))^(3/2), x)

Giac [F]

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{dx + c}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((d*x+c)/(b*tanh(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*x + c)/(b*tanh(f*x + e))^(3/2), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{c + dx}{(b \tanh(e + fx))^{3/2}} dx$$

[In] int((c + d*x)/(b*tanh(e + f*x))^(3/2),x)

[Out] int((c + d*x)/(b*tanh(e + f*x))^(3/2), x)

3.21 $\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx$

Optimal result	189
Rubi [N/A]	190
Mathematica [N/A]	196
Maple [N/A] (verified)	196
Fricas [F(-2)]	196
Sympy [N/A]	196
Maxima [N/A]	197
Giac [N/A]	197
Mupad [N/A]	197

Optimal result

Integrand size = 20, antiderivative size = 20

$$\begin{aligned}
& \int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \frac{4(-b)^{3/2} d(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} \\
& + \frac{2(-b)^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{f^3} + \frac{4b^{3/2} d(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
& + \frac{2b^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{f^3} - \frac{4b^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{f^3} \\
& + \frac{4b^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^3} \\
& - \frac{2b^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{f^3} \\
& - \frac{2b^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{f^3} \\
& - \frac{4(-b)^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^3} \\
& + \frac{2(-b)^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{f^3} \\
& + \frac{2(-b)^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})\left(1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{f^3} \\
& + \frac{4(-b)^{3/2} d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1 + \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^3} \\
& - \frac{2b^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} - \sqrt{b \tanh(e+fx)}}\right)}{f^3} - \frac{2b^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b} + \sqrt{b \tanh(e+fx)}}\right)}{f^3} \\
& + \frac{b^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} - \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} - \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{f^3} \\
& + \frac{b^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b} + \sqrt{b \tanh(e+fx)})}{(\sqrt{-b} + \sqrt{b})(\sqrt{b} + \sqrt{b \tanh(e+fx)})}\right)}{f^3} \\
& - \frac{2(-b)^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2}{1 - \frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^3} \\
& (-b)^{3/2} d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2(\sqrt{b} - \sqrt{b \tanh(e+fx)})}{\dots}\right)
\end{aligned}$$

```
[Out] 4*(-b)^(3/2)*d*(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/f^2+2*(-b)
^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))^2/f^3+4*b^(3/2)*d*(d*x
+c)*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))/f^2+2*b^(3/2)*d^2*arctanh((b*tan
h(f*x+e))^(1/2)/b^(1/2))^2/f^3-4*b^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/
b^(1/2))*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^3+4*b^(3/2)*d^2*ar
ctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))
^(1/2)))/f^3-2*b^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/
2)*((-b)^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tan
h(f*x+e))^(1/2)))/f^3-2*b^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*l
n(2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2
)+(b*tanh(f*x+e))^(1/2)))/f^3-4*(-b)^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2
))/(-b)^(1/2))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^3+2*(-b)^(3/2)*d
^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*tanh(f*x+e))
^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^3+2*(-b
)^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(1/2)+(b*tan
h(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))
)/f^3+4*(-b)^(3/2)*d^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1+(b*
tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^3-2*b^(3/2)*d^2*polylog(2,1-2*b^(1/2)/(b
^(1/2)-(b*tanh(f*x+e))^(1/2)))/f^3-2*b^(3/2)*d^2*polylog(2,1-2*b^(1/2)/(b
^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^3+b^(3/2)*d^2*polylog(2,1-2*b^(1/2)*((-b)^(1/
2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/
2)))/f^3+b^(3/2)*d^2*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2
)))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/f^3-2*(-b)^(3/2)*d
^2*polylog(2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/f^3+(-b)^(3/2)*d^2*p
olylog(2,1-2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tan
h(f*x+e))^(1/2)/(-b)^(1/2)))/f^3+(-b)^(3/2)*d^2*polylog(2,1+2*(b^(1/2)+(b*t
anh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2
)))/f^3-2*(-b)^(3/2)*d^2*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))
)/f^3-2*b*(d*x+c)^2*(b*tanh(f*x+e))^(1/2)/f+b^2*Unintegrable((d*x+c)^2/(b*ta
nh(f*x+e))^(1/2),x)
```

Rubi [N/A]

Not integrable

Time = 1.13 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx$$

```
[In] Int[(c + d*x)^2*(b*Tanh[e + f*x])^(3/2),x]
```

```
[Out] (4*(-b)^(3/2)*d*(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]])/f^2 + (2
*(-b)^(3/2)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/f^3 + (4*b^(3/2)
*d*(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]])/f^2 + (2*b^(3/2)*d^2*A
```

```

rcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]^2)/f^3 - (4*b^(3/2)*d^2*ArcTanh[Sqrt[
b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])
])/f^3 + (4*b^(3/2)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[
b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/f^3 - (2*b^(3/2)*d^2*ArcTanh[Sqrt[b
*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]])
)/((Sqrt[-b] - Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))]/f^3 - (2*b^(3/2
)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[b]]*Log[(2*Sqrt[b]*(Sqrt[-b] + Sqr
t[b*Tanh[e + f*x]]))/((Sqrt[-b] + Sqrt[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]
))])/f^3 - (4*(-b)^(3/2)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/
(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^3 + (2*(-b)^(3/2)*d^2*ArcTanh[Sqrt
[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])/((Sqr
t[-b] + Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]))])/f^3 + (2*(-b)^(3/2
)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[(-2*(Sqrt[b] + Sqrt[b*Tan
h[e + f*x]))/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])))]
)/f^3 + (4*(-b)^(3/2)*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]*Log[2/(1 +
Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^3 - (2*b^(3/2)*d^2*PolyLog[2, 1 - (2*S
qrt[b])/(Sqrt[b] - Sqrt[b*Tanh[e + f*x]])])/f^3 - (2*b^(3/2)*d^2*PolyLog[2,
1 - (2*Sqrt[b])/(Sqrt[b] + Sqrt[b*Tanh[e + f*x]])])/f^3 + (b^(3/2)*d^2*Pol
yLog[2, 1 - (2*Sqrt[b]*(Sqrt[-b] - Sqrt[b*Tanh[e + f*x]))/((Sqrt[-b] - Sqr
t[b])*(Sqrt[b] + Sqrt[b*Tanh[e + f*x]]))]/f^3 + (b^(3/2)*d^2*PolyLog[2, 1
- (2*Sqrt[b]*(Sqrt[-b] + Sqrt[b*Tanh[e + f*x]))/((Sqrt[-b] + Sqrt[b])*(Sqr
t[b] + Sqrt[b*Tanh[e + f*x]]))]/f^3 - (2*(-b)^(3/2)*d^2*PolyLog[2, 1 - 2/(
1 - Sqrt[b*Tanh[e + f*x]]/Sqrt[-b])])/f^3 + ((-b)^(3/2)*d^2*PolyLog[2, 1 -
(2*(Sqrt[b] - Sqrt[b*Tanh[e + f*x]))/((Sqrt[-b] + Sqrt[b])*(1 - Sqrt[b*Tan
h[e + f*x]]/Sqrt[-b]))])/f^3 + ((-b)^(3/2)*d^2*PolyLog[2, 1 + (2*(Sqrt[b] +
Sqrt[b*Tanh[e + f*x]))/((Sqrt[-b] - Sqrt[b])*(1 - Sqrt[b*Tanh[e + f*x]]/S
qrt[-b]))])/f^3 - (2*(-b)^(3/2)*d^2*PolyLog[2, 1 - 2/(1 + Sqrt[b*Tanh[e + f
*x]]/Sqrt[-b])])/f^3 - (2*b*(c + d*x)^2*Sqrt[b*Tanh[e + f*x]])/f + b^2*Defe
r[Int][(c + d*x)^2/Sqrt[b*Tanh[e + f*x]], x]

```

Rubi steps

$$\begin{aligned}
\text{integral} = & -\frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
& + \frac{(4bd) \int (c+dx)\sqrt{b\tanh(e+fx)} dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&\quad - \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&\quad - \frac{(4(-b)^{3/2}d^2) \int \operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right) dx}{f^2} \\
&\quad - \frac{(4b^{3/2}d^2) \int \operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right) dx}{f^2} \\
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&\quad - \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&\quad - \frac{(4(-b)^{3/2}d^2) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2} dx, x, \tanh(e+fx)\right)}{f^3} \\
&\quad - \frac{(4b^{3/2}d^2) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2} dx, x, \tanh(e+fx)\right)}{f^3} \\
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&\quad - \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&\quad + \frac{(8\sqrt{-bd^2}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&\quad + \frac{(8\sqrt{bd^2}) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&\quad - \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&\quad + \frac{(8\sqrt{-bd^2}) \operatorname{Subst}\left(\int \left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&\quad - \frac{(8\sqrt{bd^2}) \operatorname{Subst}\left(\int \left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right) dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} \\
&\quad - \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&\quad + \frac{(4(-b)^{3/2}d^2) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&\quad + \frac{(4(-b)^{3/2}d^2) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&\quad + \frac{(4b^{3/2}d^2) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&\quad - \frac{(4b^{3/2}d^2) \operatorname{Subst}\left(\int \frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{f^2} + \frac{2(-b)^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{f^3} \\
&+ \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{f^2} + \frac{2b^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{f^3} \\
&- \frac{2b(c+dx)^2\sqrt{b}\tanh(e+fx)}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b}\tanh(e+fx)} dx \\
&+ \frac{(4(-b)^{3/2}d^2) \operatorname{Subst}\left(\int \left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}-x)} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b}+x)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^3} \\
&- \frac{(4bd^2) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^3} \\
&- \frac{(4bd^2) \operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}} dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^3} \\
&- \frac{(4b^{3/2}d^2) \operatorname{Subst}\left(\int \left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}-x)} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b}+x)}\right) dx, x, \sqrt{b}\tanh(e+fx)\right)}{f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4(-b)^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{f^2} + \frac{2(-b)^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)^2}{f^3} \\
&+ \frac{4b^{3/2}d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{f^2} + \frac{2b^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)^2}{f^3} \\
&- \frac{4b^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)\log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b\tanh(e+fx)}}\right)}{f^3} \\
&+ \frac{4(-b)^{3/2}d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)\log\left(\frac{2}{1+\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}}\right)}{f^3} \\
&- \frac{2b(c+dx)^2\sqrt{b\tanh(e+fx)}}{f} + b^2 \int \frac{(c+dx)^2}{\sqrt{b\tanh(e+fx)}} dx \\
&+ \frac{(2(-b)^{3/2}d^2)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&- \frac{(2(-b)^{3/2}d^2)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&+ \frac{(4bd^2)\operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&+ \frac{(4bd^2)\operatorname{Subst}\left(\int \frac{\log\left(\frac{2}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&+ \frac{(2b^{3/2}d^2)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3} \\
&- \frac{(2b^{3/2}d^2)\operatorname{Subst}\left(\int \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}} dx, x, \sqrt{b\tanh(e+fx)}\right)}{f^3}
\end{aligned}$$

= Too large to display

Mathematica [N/A]

Not integrable

Time = 29.38 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx$$

[In] Integrate[(c + d*x)^2*(b*Tanh[e + f*x])^(3/2), x]

[Out] Integrate[(c + d*x)^2*(b*Tanh[e + f*x])^(3/2), x]

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int (dx + c)^2 (b \tanh(fx + e))^{3/2} dx$$

[In] int((d*x+c)^2*(b*tanh(f*x+e))^(3/2), x)

[Out] int((d*x+c)^2*(b*tanh(f*x+e))^(3/2), x)

Fricas [F(-2)]

Exception generated.

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(3/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 3.00 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (b \tanh(e + fx))^{3/2} (c + dx)^2 dx$$

[In] integrate((d*x+c)**2*(b*tanh(f*x+e))**(3/2), x)

[Out] Integral((b*tanh(e + f*x))**(3/2)*(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (dx + c)^2 (b \tanh(fx + e))^{\frac{3}{2}} dx$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*x + c)^2*(b*tanh(f*x + e))^(3/2), x)

Giac [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (dx + c)^2 (b \tanh(fx + e))^{\frac{3}{2}} dx$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tanh(f*x + e))^(3/2), x)

Mupad [N/A]

Not integrable

Time = 2.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 (b \tanh(e + fx))^{3/2} dx = \int (b \tanh(e + fx))^{3/2} (c + dx)^2 dx$$

[In] int((b*tanh(e + f*x))^(3/2)*(c + d*x)^2,x)

[Out] int((b*tanh(e + f*x))^(3/2)*(c + d*x)^2, x)

3.22 $\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx$

Optimal result	198
Rubi [N/A]	198
Mathematica [F(-1)]	199
Maple [N/A] (verified)	199
Fricas [F(-2)]	199
Sympy [N/A]	199
Maxima [N/A]	200
Giac [N/A]	200
Mupad [N/A]	200

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \text{Int}\left((c + dx)^2 \sqrt{b \tanh(e + fx)}, x\right)$$

[Out] Unintegrable((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx$$

[In] Int[(c + d*x)^2*Sqrt[b*Tanh[e + f*x]],x]

[Out] Defer[Int] [(c + d*x)^2*Sqrt[b*Tanh[e + f*x]], x]

Rubi steps

$$\text{integral} = \int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx$$

Mathematica [F(-1)]

Timed out.

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \$Aborted$$

[In] Integrate[(c + d*x)^2*Sqrt[b*Tanh[e + f*x]],x]

[Out] \$Aborted

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int (dx + c)^2 \sqrt{b \tanh(fx + e)} dx$$

[In] int((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x)

[Out] int((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x)

Fricas [F(-2)]

Exception generated.

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 1.30 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \int \sqrt{b \tanh(e + fx)} (c + dx)^2 dx$$

[In] integrate((d*x+c)**2*(b*tanh(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*tanh(e + f*x))*(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.55 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \int (dx + c)^2 \sqrt{b \tanh(fx + e)} dx$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((d*x + c)^2*sqrt(b*tanh(f*x + e)), x)

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \int (dx + c)^2 \sqrt{b \tanh(fx + e)} dx$$

[In] integrate((d*x+c)^2*(b*tanh(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate((d*x + c)^2*sqrt(b*tanh(f*x + e)), x)

Mupad [N/A]

Not integrable

Time = 1.99 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx = \int \sqrt{b \tanh(e + fx)} (c + dx)^2 dx$$

[In] int((b*tanh(e + f*x))^(1/2)*(c + d*x)^2,x)

[Out] int((b*tanh(e + f*x))^(1/2)*(c + d*x)^2, x)

$$3.23 \quad \int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx$$

Optimal result	201
Rubi [N/A]	201
Mathematica [F(-1)]	202
Maple [N/A] (verified)	202
Fricas [F(-2)]	202
Sympy [N/A]	202
Maxima [N/A]	203
Giac [F(-2)]	203
Mupad [N/A]	203

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx = \text{Int} \left(\frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}}, x \right)$$

[Out] Unintegrable((d*x+c)^2/(b*tanh(f*x+e))^(1/2), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx = \int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx$$

[In] Int[(c + d*x)^2/Sqrt[b*Tanh[e + f*x]], x]

[Out] Defer[Int] [(c + d*x)^2/Sqrt[b*Tanh[e + f*x]], x]

Rubi steps

$$\text{integral} = \int \frac{(c+dx)^2}{\sqrt{b \tanh(e+fx)}} dx$$

Mathematica [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \$Aborted$$

[In] Integrate[(c + d*x)^2/Sqrt[b*Tanh[e + f*x]],x]

[Out] \$Aborted

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(dx + c)^2}{\sqrt{b \tanh(fx + e)}} dx$$

[In] int((d*x+c)^2/(b*tanh(f*x+e))^(1/2),x)

[Out] int((d*x+c)^2/(b*tanh(f*x+e))^(1/2),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 1.25 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx$$

[In] integrate((d*x+c)**2/(b*tanh(f*x+e))**(1/2),x)

[Out] Integral((c + d*x)**2/sqrt(b*tanh(e + f*x)), x)

Maxima [N/A]

Not integrable

Time = 0.59 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{(dx + c)^2}{\sqrt{b \tanh(fx + e)}} dx$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((d*x + c)^2/sqrt(b*tanh(f*x + e)), x)

Giac [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \text{Exception raised: RuntimeError}$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(1/2),x, algorithm="giac")

[Out] Exception raised: RuntimeError >> an error occurred running a Giac command:
INPUT:sage2OUTPUT:index.cc index_m i_lex_is_greater Error: Bad Argument Value**Mupad [N/A]**

Not integrable

Time = 2.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx = \int \frac{(c + dx)^2}{\sqrt{b \tanh(e + fx)}} dx$$

[In] int((c + d*x)^2/(b*tanh(e + f*x))^(1/2),x)

[Out] int((c + d*x)^2/(b*tanh(e + f*x))^(1/2), x)

3.24
$$\int \frac{(c+dx)^2}{(b \tanh(e+fx))^{3/2}} dx$$

Optimal result	205
Rubi [N/A]	206
Mathematica [N/A]	211
Maple [N/A] (verified)	211
Fricas [F(-2)]	211
Sympy [N/A]	211
Maxima [N/A]	212
Giac [N/A]	212
Mupad [N/A]	212

Optimal result

Integrand size = 20, antiderivative size = 20

$$\begin{aligned}
& \int \frac{(c+dx)^2}{(b \tanh(e+fx))^{3/2}} dx = \frac{4d(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f^2} \\
& + \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)^2}{(-b)^{3/2} f^3} \\
& + \frac{4d(c+dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} + \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right)^2}{b^{3/2} f^3} \\
& - \frac{4d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^3} \\
& + \frac{4d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}}{\sqrt{b}+\sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^3} \\
& - \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b}-\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}-\sqrt{b})(\sqrt{b}+\sqrt{b \tanh(e+fx)})}\right)}{b^{3/2} f^3} \\
& - \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{b}}\right) \log\left(\frac{2\sqrt{b}(\sqrt{-b}+\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}+\sqrt{b})(\sqrt{b}+\sqrt{b \tanh(e+fx)})}\right)}{b^{3/2} f^3} \\
& - \frac{4d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1-\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{(-b)^{3/2} f^3} \\
& + \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2(\sqrt{b}-\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}+\sqrt{b})\left(1-\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{(-b)^{3/2} f^3} \\
& + \frac{2d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(-\frac{2(\sqrt{b}+\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}-\sqrt{b})\left(1-\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right)}\right)}{(-b)^{3/2} f^3} \\
& + \frac{4d^2 \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}\right) \log\left(\frac{2}{1+\frac{\sqrt{b \tanh(e+fx)}}{\sqrt{-b}}}\right)}{(-b)^{3/2} f^3} \\
& - \frac{2d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^3} \\
& - \frac{2d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}}{\sqrt{b}+\sqrt{b \tanh(e+fx)}}\right)}{b^{3/2} f^3} \\
& + \frac{d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b}-\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}-\sqrt{b})(\sqrt{b}+\sqrt{b \tanh(e+fx)})}\right)}{b^{3/2} f^3} \\
& + \frac{d^2 \operatorname{PolyLog}\left(2, 1 - \frac{2\sqrt{b}(\sqrt{-b}+\sqrt{b \tanh(e+fx)})}{(\sqrt{-b}+\sqrt{b})(\sqrt{b}+\sqrt{b \tanh(e+fx)})}\right)}{b^{3/2} f^3}
\end{aligned}$$

```
[Out] 4*d*(d*x+c)*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))/(-b)^(3/2)/f^2+2*d^2*
arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))^2/(-b)^(3/2)/f^3+4*d*(d*x+c)*arct
anh((b*tanh(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f^2+2*d^2*arctanh((b*tanh(f*x+e)
)^(1/2)/b^(1/2))^2/b^(3/2)/f^3-4*d^2*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))
*ln(2*b^(1/2)/(b^(1/2)-(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^3+4*d^2*arctanh((b
*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/
b^(3/2)/f^3-2*d^2*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1/2)*((-b)
^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e)
^(1/2)))/b^(3/2)/f^3-2*d^2*arctanh((b*tanh(f*x+e))^(1/2)/b^(1/2))*ln(2*b^(1
/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(b^(1/2)+(b*ta
nh(f*x+e))^(1/2)))/b^(3/2)/f^3-4*d^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2
))*ln(2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^3+2*d^2*arctanh(
(b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)
^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^3+2*d^2
*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(-2*(b^(1/2)+(b*tanh(f*x+e))^(
1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)
/f^3+4*d^2*arctanh((b*tanh(f*x+e))^(1/2)/(-b)^(1/2))*ln(2/(1+(b*tanh(f*x+e)
)^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^3-2*d^2*polylog(2,1-2*b^(1/2)/(b^(1/2)-(b
*tanh(f*x+e))^(1/2)))/b^(3/2)/f^3-2*d^2*polylog(2,1-2*b^(1/2)/(b^(1/2)+(b*t
anh(f*x+e))^(1/2)))/b^(3/2)/f^3+d^2*polylog(2,1-2*b^(1/2)*((-b)^(1/2)-(b*ta
nh(f*x+e))^(1/2))/((-b)^(1/2)-b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(
3/2)/f^3+d^2*polylog(2,1-2*b^(1/2)*((-b)^(1/2)+(b*tanh(f*x+e))^(1/2))/((-b)
^(1/2)+b^(1/2))/(b^(1/2)+(b*tanh(f*x+e))^(1/2)))/b^(3/2)/f^3-2*d^2*polylog(
2,1-2/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^3+d^2*polylog(2,1-
2*(b^(1/2)-(b*tanh(f*x+e))^(1/2))/((-b)^(1/2)+b^(1/2))/(1-(b*tanh(f*x+e))^(
1/2)/(-b)^(1/2)))/(-b)^(3/2)/f^3+d^2*polylog(2,1+2*(b^(1/2)+(b*tanh(f*x+e)
)^(1/2))/((-b)^(1/2)-b^(1/2))/(1-(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/
2)/f^3-2*d^2*polylog(2,1-2/(1+(b*tanh(f*x+e))^(1/2)/(-b)^(1/2)))/(-b)^(3/2)
/f^3-2*(d*x+c)^2/b/f/(b*tanh(f*x+e))^(1/2)+Unintegrable((d*x+c)^2*(b*tanh(f
*x+e))^(1/2),x)/b^2
```

Rubi [N/A]

Not integrable

Time = 1.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx$$

```
[In] Int[(c + d*x)^2/(b*Tanh[e + f*x])^(3/2),x]
```

```
[Out] (4*d*(c + d*x)*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]])/((-b)^(3/2)*f^2) +
(2*d^2*ArcTanh[Sqrt[b*Tanh[e + f*x]]/Sqrt[-b]]^2)/((-b)^(3/2)*f^3) + (4*d*(
```

$c + dx) * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]] / (b^{(3/2)} * f^2) + (2 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]]^2) / (b^{(3/2)} * f^3) - (4 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]] * \text{Log}[(2 * \text{Sqrt}[b]) / (\text{Sqrt}[b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / (b^{(3/2)} * f^3) + (4 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]] * \text{Log}[(2 * \text{Sqrt}[b]) / (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / (b^{(3/2)} * f^3) - (2 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]] * \text{Log}[(2 * \text{Sqrt}[b] * (\text{Sqrt}[-b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] - \text{Sqrt}[b]) * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])])]) / (b^{(3/2)} * f^3) - (2 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[b]] * \text{Log}[(2 * \text{Sqrt}[b] * (\text{Sqrt}[-b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] + \text{Sqrt}[b]) * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])])]) / (b^{(3/2)} * f^3) - (4 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b]] * \text{Log}[2 / (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])]) / ((-b)^{(3/2)} * f^3) + (2 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b]] * \text{Log}[(2 * (\text{Sqrt}[b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] + \text{Sqrt}[b]) * (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])])]) / ((-b)^{(3/2)} * f^3) + (2 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b]] * \text{Log}[(-2 * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] - \text{Sqrt}[b]) * (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])])]) / ((-b)^{(3/2)} * f^3) + (4 * d^2 * \text{ArcTanh}[\text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b]] * \text{Log}[2 / (1 + \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])]) / ((-b)^{(3/2)} * f^3) - (2 * d^2 * \text{PolyLog}[2, 1 - (2 * \text{Sqrt}[b]) / (\text{Sqrt}[b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / (b^{(3/2)} * f^3) - (2 * d^2 * \text{PolyLog}[2, 1 - (2 * \text{Sqrt}[b]) / (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / (b^{(3/2)} * f^3) + (d^2 * \text{PolyLog}[2, 1 - (2 * \text{Sqrt}[b] * (\text{Sqrt}[-b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] - \text{Sqrt}[b]) * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])])]) / (b^{(3/2)} * f^3) + (d^2 * \text{PolyLog}[2, 1 - (2 * \text{Sqrt}[b] * (\text{Sqrt}[-b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] + \text{Sqrt}[b]) * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])])]) / (b^{(3/2)} * f^3) - (2 * d^2 * \text{PolyLog}[2, 1 - 2 / (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])]) / ((-b)^{(3/2)} * f^3) + (d^2 * \text{PolyLog}[2, 1 - (2 * (\text{Sqrt}[b] - \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] + \text{Sqrt}[b]) * (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])])]) / ((-b)^{(3/2)} * f^3) + (d^2 * \text{PolyLog}[2, 1 + (2 * (\text{Sqrt}[b] + \text{Sqrt}[b * \text{Tanh}[e + f * x]])]) / ((\text{Sqrt}[-b] - \text{Sqrt}[b]) * (1 - \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])])]) / ((-b)^{(3/2)} * f^3) - (2 * d^2 * \text{PolyLog}[2, 1 - 2 / (1 + \text{Sqrt}[b * \text{Tanh}[e + f * x]] / \text{Sqrt}[-b])]) / ((-b)^{(3/2)} * f^3) - (2 * (c + dx)^2) / (b * f * \text{Sqrt}[b * \text{Tanh}[e + f * x]]) + \text{Defer}[\text{Int}[(c + dx)^2 * \text{Sqrt}[b * \text{Tanh}[e + f * x]], x] / b^2$

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{2(c + dx)^2}{bf\sqrt{b \tanh(e + fx)}} + \frac{\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx}{b^2} + \frac{(4d) \int \frac{c + dx}{\sqrt{b \tanh(e + fx)}} dx}{bf} \\
 &= \frac{4d(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2} f^2} + \frac{4d(c + dx) \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right)}{b^{3/2} f^2} \\
 &\quad - \frac{2(c + dx)^2}{bf\sqrt{b \tanh(e + fx)}} + \frac{\int (c + dx)^2 \sqrt{b \tanh(e + fx)} dx}{b^2} \\
 &\quad - \frac{(4d^2) \int \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{-b}}\right) dx}{(-b)^{3/2} f^2} - \frac{(4d^2) \int \operatorname{arctanh}\left(\frac{\sqrt{b \tanh(e + fx)}}{\sqrt{b}}\right) dx}{b^{3/2} f^2}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f^2} \\
&\quad - \frac{2(c+dx)^2}{bf\sqrt{b}\tanh(e+fx)} + \frac{\int(c+dx)^2\sqrt{b}\tanh(e+fx)dx}{b^2} \\
&\quad - \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{b\sqrt{bx}}{(-b)^{3/2}}\right)}{-1+x^2}dx, x, \tanh(e+fx)\right)}{(-b)^{3/2}f^3} \\
&\quad - \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{\sqrt{bx}}{\sqrt{b}}\right)}{1-x^2}dx, x, \tanh(e+fx)\right)}{b^{3/2}f^3} \\
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f^2} \\
&\quad - \frac{2(c+dx)^2}{bf\sqrt{b}\tanh(e+fx)} + \frac{\int(c+dx)^2\sqrt{b}\tanh(e+fx)dx}{b^2} \\
&\quad + \frac{(8d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{-1+\frac{x^4}{b^2}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{(-b)^{5/2}f^3} \\
&\quad - \frac{(8d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x^4}{b^2}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^{5/2}f^3} \\
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f^2} \\
&\quad - \frac{2(c+dx)^2}{bf\sqrt{b}\tanh(e+fx)} + \frac{\int(c+dx)^2\sqrt{b}\tanh(e+fx)dx}{b^2} \\
&\quad + \frac{(8d^2)\operatorname{Subst}\left(\int\left(-\frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b-x^2)} - \frac{bx\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b}\tanh(e+fx)\right)}{(-b)^{5/2}f^3} \\
&\quad - \frac{(8d^2)\operatorname{Subst}\left(\int\left(\frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b-x^2)} + \frac{bx\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(b+x^2)}\right)dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^{5/2}f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f^2} \\
&- \frac{2(c+dx)^2}{bf\sqrt{b\tanh(e+fx)}} + \frac{\int(c+dx)^2\sqrt{b\tanh(e+fx)}dx}{b^2} \\
&+ \frac{(4d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{3/2}f^3} \\
&+ \frac{(4d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{3/2}f^3} \\
&- \frac{(4d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b-x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{3/2}f^3} \\
&- \frac{(4d^2)\operatorname{Subst}\left(\int\frac{x\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{b+x^2}dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{3/2}f^3} \\
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{2d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{-b}}\right)^2}{(-b)^{3/2}f^3} \\
&+ \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f^2} + \frac{2d^2\operatorname{arctanh}\left(\frac{\sqrt{b\tanh(e+fx)}}{\sqrt{b}}\right)^2}{b^{3/2}f^3} \\
&- \frac{2(c+dx)^2}{bf\sqrt{b\tanh(e+fx)}} + \frac{\int(c+dx)^2\sqrt{b\tanh(e+fx)}dx}{b^2} \\
&+ \frac{(4d^2)\operatorname{Subst}\left(\int\left(\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b-x})} - \frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{2(\sqrt{b+x})}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{(-b)^{3/2}f^3} \\
&- \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{1-\frac{x}{\sqrt{b}}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^2f^3} \\
&+ \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{1-\frac{bx}{(-b)^{3/2}}}dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^2f^3} \\
&- \frac{(4d^2)\operatorname{Subst}\left(\int\left(-\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b-x})} + \frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{2(\sqrt{-b+x})}\right)dx, x, \sqrt{b\tanh(e+fx)}\right)}{b^{3/2}f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)}{(-b)^{3/2}f^2} + \frac{2d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)^2}{(-b)^{3/2}f^3} \\
&+ \frac{4d(c+dx)\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)}{b^{3/2}f^2} + \frac{2d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)^2}{b^{3/2}f^3} \\
&- \frac{4d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{b}}\right)\log\left(\frac{2\sqrt{b}}{\sqrt{b}-\sqrt{b}\tanh(e+fx)}\right)}{b^{3/2}f^3} \\
&+ \frac{4d^2\operatorname{arctanh}\left(\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}\right)\log\left(\frac{2}{1+\frac{\sqrt{b}\tanh(e+fx)}{\sqrt{-b}}}\right)}{(-b)^{3/2}f^3} \\
&- \frac{2(c+dx)^2}{bf\sqrt{b}\tanh(e+fx)} + \frac{\int(c+dx)^2\sqrt{b}\tanh(e+fx)dx}{b^2} \\
&+ \frac{(2d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{(-b)^{3/2}f^3} \\
&- \frac{(2d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{bx}{(-b)^{3/2}}\right)}{\sqrt{b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{(-b)^{3/2}f^3} \\
&+ \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\log\left(\frac{2}{1-\frac{x}{\sqrt{b}}}\right)}{1-\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^2f^3} \\
&- \frac{(4d^2)\operatorname{Subst}\left(\int\frac{\log\left(\frac{2}{1-\frac{bx}{(-b)^{3/2}}}\right)}{1+\frac{x^2}{b}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^2f^3} \\
&+ \frac{(2d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b-x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^{3/2}f^3} \\
&- \frac{(2d^2)\operatorname{Subst}\left(\int\frac{\operatorname{arctanh}\left(\frac{x}{\sqrt{b}}\right)}{\sqrt{-b+x}}dx, x, \sqrt{b}\tanh(e+fx)\right)}{b^{3/2}f^3}
\end{aligned}$$

= Too large to display

Mathematica [N/A]

Not integrable

Time = 31.05 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx$$

[In] Integrate[(c + d*x)^2/(b*Tanh[e + f*x])^(3/2), x]

[Out] Integrate[(c + d*x)^2/(b*Tanh[e + f*x])^(3/2), x]

Maple [N/A] (verified)

Not integrable

Time = 0.08 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(dx + c)^2}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] int((d*x+c)^2/(b*tanh(f*x+e))^(3/2), x)

[Out] int((d*x+c)^2/(b*tanh(f*x+e))^(3/2), x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(3/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 1.40 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(c + dx)^2}{(b \tanh(e + fx))^{\frac{3}{2}}} dx$$

[In] integrate((d*x+c)**2/(b*tanh(f*x+e))**(3/2), x)

[Out] Integral((c + d*x)**2/(b*tanh(e + f*x))**(3/2), x)

Maxima [N/A]

Not integrable

Time = 0.58 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(dx + c)^2}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*x + c)^2/(b*tanh(f*x + e))^(3/2), x)

Giac [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(dx + c)^2}{(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate((d*x+c)^2/(b*tanh(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*x + c)^2/(b*tanh(f*x + e))^(3/2), x)

Mupad [N/A]

Not integrable

Time = 2.11 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(c + dx)^2}{(b \tanh(e + fx))^{3/2}} dx = \int \frac{(c + dx)^2}{(b \tanh(e + fx))^{\frac{3}{2}}} dx$$

[In] int((c + d*x)^2/(b*tanh(e + f*x))^(3/2),x)

[Out] int((c + d*x)^2/(b*tanh(e + f*x))^(3/2), x)

$$3.25 \quad \int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx$$

Optimal result	213
Rubi [N/A]	213
Mathematica [N/A]	214
Maple [N/A] (verified)	214
Fricas [F(-2)]	214
Sympy [N/A]	214
Maxima [N/A]	215
Giac [N/A]	215
Mupad [N/A]	215

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx = \text{Int}\left(\frac{(b \tanh(e+fx))^{3/2}}{c+dx}, x\right)$$

[Out] Unintegrable((b*tanh(f*x+e))^(3/2)/(d*x+c), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx = \int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx$$

[In] Int[(b*Tanh[e + f*x])^(3/2)/(c + d*x), x]

[Out] Defer[Int] [(b*Tanh[e + f*x])^(3/2)/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(b \tanh(e+fx))^{3/2}}{c+dx} dx$$

Mathematica [N/A]

Not integrable

Time = 24.80 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx$$

[In] Integrate[(b*Tanh[e + f*x])^(3/2)/(c + d*x),x]

[Out] Integrate[(b*Tanh[e + f*x])^(3/2)/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{(b \tanh(fx + e))^{3/2}}{dx + c} dx$$

[In] int((b*tanh(f*x+e))^(3/2)/(d*x+c),x)

[Out] int((b*tanh(f*x+e))^(3/2)/(d*x+c),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \text{Exception raised: TypeError}$$

[In] integrate((b*tanh(f*x+e))^(3/2)/(d*x+c),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 1.00 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx$$

[In] integrate((b*tanh(f*x+e))**(3/2)/(d*x+c),x)

[Out] Integral((b*tanh(e + f*x))**(3/2)/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \int \frac{(b \tanh(fx + e))^{3/2}}{dx + c} dx$$

[In] integrate((b*tanh(f*x+e))^(3/2)/(d*x+c),x, algorithm="maxima")

[Out] integrate((b*tanh(f*x + e))^(3/2)/(d*x + c), x)

Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \int \frac{(b \tanh(fx + e))^{3/2}}{dx + c} dx$$

[In] integrate((b*tanh(f*x+e))^(3/2)/(d*x+c),x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e))^(3/2)/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 2.00 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx = \int \frac{(b \tanh(e + fx))^{3/2}}{c + dx} dx$$

[In] int((b*tanh(e + f*x))^(3/2)/(c + d*x),x)

[Out] int((b*tanh(e + f*x))^(3/2)/(c + d*x), x)

3.26 $\int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx$

Optimal result	216
Rubi [N/A]	216
Mathematica [N/A]	217
Maple [N/A] (verified)	217
Fricas [F(-2)]	217
Sympy [N/A]	217
Maxima [N/A]	218
Giac [N/A]	218
Mupad [N/A]	218

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx = \text{Int}\left(\frac{\sqrt{b \tanh(e+fx)}}{c+dx}, x\right)$$

[Out] Unintegrable((b*tanh(f*x+e))^(1/2)/(d*x+c), x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx = \int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx$$

[In] Int[Sqrt[b*Tanh[e + f*x]]/(c + d*x), x]

[Out] Defer[Int][Sqrt[b*Tanh[e + f*x]]/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{\sqrt{b \tanh(e+fx)}}{c+dx} dx$$

Mathematica [N/A]

Not integrable

Time = 2.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx$$

[In] Integrate[Sqrt[b*Tanh[e + f*x]]/(c + d*x), x]

[Out] Integrate[Sqrt[b*Tanh[e + f*x]]/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{\sqrt{b \tanh(fx + e)}}{dx + c} dx$$

[In] int((b*tanh(f*x+e))^(1/2)/(d*x+c), x)

[Out] int((b*tanh(f*x+e))^(1/2)/(d*x+c), x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \text{Exception raised: TypeError}$$

[In] integrate((b*tanh(f*x+e))^(1/2)/(d*x+c), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 0.71 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx$$

[In] integrate((b*tanh(f*x+e))**(1/2)/(d*x+c), x)

[Out] Integral(sqrt(b*tanh(e + f*x))/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \int \frac{\sqrt{b \tanh(fx + e)}}{dx + c} dx$$

[In] integrate((b*tanh(f*x+e))^(1/2)/(d*x+c),x, algorithm="maxima")

[Out] integrate(sqrt(b*tanh(f*x + e))/(d*x + c), x)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \int \frac{\sqrt{b \tanh(fx + e)}}{dx + c} dx$$

[In] integrate((b*tanh(f*x+e))^(1/2)/(d*x+c),x, algorithm="giac")

[Out] integrate(sqrt(b*tanh(f*x + e))/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.71 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx = \int \frac{\sqrt{b \tanh(e + fx)}}{c + dx} dx$$

[In] int((b*tanh(e + f*x))^(1/2)/(c + d*x),x)

[Out] int((b*tanh(e + f*x))^(1/2)/(c + d*x), x)

$$3.27 \quad \int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx$$

Optimal result	219
Rubi [N/A]	219
Mathematica [N/A]	220
Maple [N/A] (verified)	220
Fricas [F(-2)]	220
Sympy [N/A]	220
Maxima [N/A]	221
Giac [N/A]	221
Mupad [N/A]	221

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx = \text{Int}\left(\frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(b*tanh(f*x+e))^(1/2), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx = \int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx$$

[In] Int[1/((c + d*x)*Sqrt[b*Tanh[e + f*x]]), x]

[Out] Defer[Int][1/((c + d*x)*Sqrt[b*Tanh[e + f*x]]), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)\sqrt{b \tanh(e+fx)}} dx$$

Mathematica [N/A]

Not integrable

Time = 2.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx$$

[In] Integrate[1/((c + d*x)*Sqrt[b*Tanh[e + f*x]]),x]

[Out] Integrate[1/((c + d*x)*Sqrt[b*Tanh[e + f*x]]), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{1}{(dx + c)\sqrt{b \tanh(fx + e)}} dx$$

[In] int(1/(d*x+c)/(b*tanh(f*x+e))^(1/2),x)

[Out] int(1/(d*x+c)/(b*tanh(f*x+e))^(1/2),x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \text{Exception raised: TypeError}$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 0.95 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \int \frac{1}{\sqrt{b \tanh(e + fx)}(c + dx)} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))**(1/2),x)

[Out] Integral(1/(sqrt(b*tanh(e + f*x))*(c + d*x)), x)

Maxima [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \int \frac{1}{(dx + c)\sqrt{b \tanh(fx + e)}} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(1/((d*x + c)*sqrt(b*tanh(f*x + e))), x)

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \int \frac{1}{(dx + c)\sqrt{b \tanh(fx + e)}} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*x + c)*sqrt(b*tanh(f*x + e))), x)

Mupad [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)\sqrt{b \tanh(e + fx)}} dx = \int \frac{1}{\sqrt{b \tanh(e + fx)} (c + dx)} dx$$

[In] int(1/((b*tanh(e + f*x))^(1/2)*(c + d*x)),x)

[Out] int(1/((b*tanh(e + f*x))^(1/2)*(c + d*x)), x)

$$3.28 \quad \int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx$$

Optimal result	222
Rubi [N/A]	222
Mathematica [N/A]	223
Maple [N/A] (verified)	223
Fricas [F(-2)]	223
Sympy [N/A]	223
Maxima [N/A]	224
Giac [N/A]	224
Mupad [N/A]	224

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx = \text{Int}\left(\frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(b*tanh(f*x+e))^(3/2), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx = \int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx$$

[In] Int[1/((c + d*x)*(b*Tanh[e + f*x]))^(3/2)), x]

[Out] Defer[Int][1/((c + d*x)*(b*Tanh[e + f*x]))^(3/2)), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(b \tanh(e+fx))^{3/2}} dx$$

Mathematica [N/A]

Not integrable

Time = 21.46 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx$$

[In] Integrate[1/((c + d*x)*(b*Tanh[e + f*x])^(3/2)), x]

[Out] Integrate[1/((c + d*x)*(b*Tanh[e + f*x])^(3/2)), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 0.90

$$\int \frac{1}{(dx + c)(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] int(1/(d*x+c)/(b*tanh(f*x+e))^(3/2), x)

[Out] int(1/(d*x+c)/(b*tanh(f*x+e))^(3/2), x)

Fricas [F(-2)]

Exception generated.

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \text{Exception raised: TypeError}$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(3/2), x, algorithm="fricas")

[Out] Exception raised: TypeError >> Error detected within library code: integrate: implementation incomplete (constant residues)

Sympy [N/A]

Not integrable

Time = 1.14 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \int \frac{1}{(b \tanh(e + fx))^{\frac{3}{2}} (c + dx)} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))**(3/2), x)

[Out] Integral(1/((b*tanh(e + f*x))**(3/2)*(c + d*x)), x)

Maxima [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \int \frac{1}{(dx + c)(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(1/((d*x + c)*(b*tanh(f*x + e))^(3/2)), x)

Giac [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \int \frac{1}{(dx + c)(b \tanh(fx + e))^{\frac{3}{2}}} dx$$

[In] integrate(1/(d*x+c)/(b*tanh(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(1/((d*x + c)*(b*tanh(f*x + e))^(3/2)), x)

Mupad [N/A]

Not integrable

Time = 1.97 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)(b \tanh(e + fx))^{3/2}} dx = \int \frac{1}{(b \tanh(e + fx))^{3/2} (c + dx)} dx$$

[In] int(1/((b*tanh(e + f*x))^(3/2)*(c + d*x)),x)

[Out] int(1/((b*tanh(e + f*x))^(3/2)*(c + d*x)), x)

3.29 $\int x^m \tanh^3(a + bx) dx$

Optimal result	225
Rubi [N/A]	225
Mathematica [N/A]	226
Maple [N/A] (verified)	226
Fricas [N/A]	226
Sympy [N/A]	226
Maxima [N/A]	227
Giac [N/A]	227
Mupad [N/A]	227

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int x^m \tanh^3(a + bx) dx = \text{Int}(x^m \tanh^3(a + bx), x)$$

[Out] Unintegrable($x^m \tanh(b*x+a)^3, x$)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh^3(a + bx) dx$$

[In] Int [$x^m \tanh[a + b*x]^3, x$]

[Out] Defer[Int] [$x^m \tanh[a + b*x]^3, x$]

Rubi steps

$$\text{integral} = \int x^m \tanh^3(a + bx) dx$$

Mathematica [N/A]

Not integrable

Time = 109.52 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh^3(a + bx) dx$$

[In] Integrate[x^m*Tanh[a + b*x]³,x][Out] Integrate[x^m*Tanh[a + b*x]³, x]**Maple [N/A] (verified)**

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \tanh(bx + a)^3 dx$$

[In] int(x^m*tanh(b*x+a)³,x)[Out] int(x^m*tanh(b*x+a)³,x)**Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh(bx + a)^3 dx$$

[In] integrate(x^m*tanh(b*x+a)³,x, algorithm="fricas")[Out] integral(x^m*tanh(b*x + a)³, x)**Sympy [N/A]**

Not integrable

Time = 0.60 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh^3(a + bx) dx$$

[In] integrate(x^m*tanh(b*x+a)³,x)[Out] Integral(x^m*tanh(a + b*x)³, x)

Maxima [N/A]

Not integrable

Time = 0.75 (sec) , antiderivative size = 171, normalized size of antiderivative = 14.25

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh(bx + a)^3 dx$$

[In] integrate(x^m*tanh(b*x+a)^3,x, algorithm="maxima")

```
[Out] x*e^(6*b*x + m*log(x) + 6*a)/((m + 1)*e^(6*b*x + 6*a) + 3*(m + 1)*e^(4*b*x + 4*a) + 3*(m + 1)*e^(2*b*x + 2*a) + m + 1) - integrate((3*(2*b*x*e^(6*a) + (m + 1)*e^(6*a))*e^(6*b*x) - 2*(m + 1)*e^(2*b*x + 2*a) + m + 1)*x^m/((m + 1)*e^(8*b*x + 8*a) + 4*(m + 1)*e^(6*b*x + 6*a) + 6*(m + 1)*e^(4*b*x + 4*a) + 4*(m + 1)*e^(2*b*x + 2*a) + m + 1), x)
```

Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh(bx + a)^3 dx$$

[In] integrate(x^m*tanh(b*x+a)^3,x, algorithm="giac")

[Out] integrate(x^m*tanh(b*x + a)^3, x)

Mupad [N/A]

Not integrable

Time = 1.68 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^3(a + bx) dx = \int x^m \tanh(a + bx)^3 dx$$

[In] int(x^m*tanh(a + b*x)^3,x)

[Out] int(x^m*tanh(a + b*x)^3, x)

3.30 $\int x^m \tanh^2(a + bx) dx$

Optimal result	228
Rubi [N/A]	228
Mathematica [N/A]	229
Maple [N/A] (verified)	229
Fricas [N/A]	229
Sympy [N/A]	229
Maxima [N/A]	230
Giac [N/A]	230
Mupad [N/A]	230

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int x^m \tanh^2(a + bx) dx = \text{Int}(x^m \tanh^2(a + bx), x)$$

[Out] Unintegrable($x^m \tanh(b*x+a)^2, x$)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh^2(a + bx) dx$$

[In] Int[$x^m \text{Tanh}[a + b*x]^2, x$]

[Out] Defer[Int][$x^m \text{Tanh}[a + b*x]^2, x$]

Rubi steps

$$\text{integral} = \int x^m \tanh^2(a + bx) dx$$

Mathematica [N/A]

Not integrable

Time = 8.32 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh^2(a + bx) dx$$

`[In] Integrate[x^m*Tanh[a + b*x]^2,x]``[Out] Integrate[x^m*Tanh[a + b*x]^2, x]`**Maple [N/A] (verified)**

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \tanh (bx + a)^2 dx$$

`[In] int(x^m*tanh(b*x+a)^2,x)``[Out] int(x^m*tanh(b*x+a)^2,x)`**Fricas [N/A]**

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh (bx + a)^2 dx$$

`[In] integrate(x^m*tanh(b*x+a)^2,x, algorithm="fricas")``[Out] integral(x^m*tanh(b*x + a)^2, x)`**Sympy [N/A]**

Not integrable

Time = 0.47 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh^2 (a + bx) dx$$

`[In] integrate(x**m*tanh(b*x+a)**2,x)``[Out] Integral(x**m*tanh(a + b*x)**2, x)`

Maxima [N/A]

Not integrable

Time = 0.68 (sec) , antiderivative size = 144, normalized size of antiderivative = 12.00

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh (bx + a)^2 dx$$

[In] integrate(x^m*tanh(b*x+a)^2,x, algorithm="maxima")

```
[Out] x*e^(4*b*x + m*log(x) + 4*a)/((m + 1)*e^(4*b*x + 4*a) + 2*(m + 1)*e^(2*b*x + 2*a) + m + 1) - integrate((2*(2*b*x*e^(4*a) + (m + 1)*e^(4*a))*e^(4*b*x) + (m + 1)*e^(2*b*x + 2*a) - m - 1)*x^m/((m + 1)*e^(6*b*x + 6*a) + 3*(m + 1)*e^(4*b*x + 4*a) + 3*(m + 1)*e^(2*b*x + 2*a) + m + 1), x)
```

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh (bx + a)^2 dx$$

[In] integrate(x^m*tanh(b*x+a)^2,x, algorithm="giac")

[Out] integrate(x^m*tanh(b*x + a)^2, x)

Mupad [N/A]

Not integrable

Time = 1.70 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int x^m \tanh^2(a + bx) dx = \int x^m \tanh(a + bx)^2 dx$$

[In] int(x^m*tanh(a + b*x)^2,x)

[Out] int(x^m*tanh(a + b*x)^2, x)

3.31 $\int x^m \tanh(a + bx) dx$

Optimal result	231
Rubi [N/A]	231
Mathematica [N/A]	232
Maple [N/A] (verified)	232
Fricas [N/A]	232
Sympy [N/A]	232
Maxima [N/A]	233
Giac [N/A]	233
Mupad [N/A]	233

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int x^m \tanh(a + bx) dx = \text{Int}(x^m \tanh(a + bx), x)$$

[Out] Unintegrable($x^m \tanh(bx+a)$, x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(a + bx) dx$$

[In] Int[$x^m \text{Tanh}[a + b*x]$, x]

[Out] Defer[Int][$x^m \text{Tanh}[a + b*x]$, x]

Rubi steps

$$\text{integral} = \int x^m \tanh(a + bx) dx$$

Mathematica [N/A]

Not integrable

Time = 6.18 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(a + bx) dx$$

[In] Integrate[x^m*Tanh[a + b*x], x][Out] Integrate[x^m*Tanh[a + b*x], x]**Maple [N/A] (verified)**

Not integrable

Time = 0.02 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \tanh(bx + a) dx$$

[In] int(x^m*tanh(b*x+a), x)[Out] int(x^m*tanh(b*x+a), x)**Fricas [N/A]**

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(bx + a) dx$$

[In] integrate(x^m*tanh(b*x+a), x, algorithm="fricas")[Out] integral(x^m*tanh(b*x + a), x)**Sympy [N/A]**

Not integrable

Time = 0.42 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(a + bx) dx$$

[In] integrate(x**m*tanh(b*x+a), x)

[Out] Integral(x**m*tanh(a + b*x), x)

Maxima [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 100, normalized size of antiderivative = 10.00

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(bx + a) dx$$

[In] integrate(x^m*tanh(b*x+a),x, algorithm="maxima")

```
[Out] x*e^(2*b*x + m*log(x) + 2*a)/((m + 1)*e^(2*b*x + 2*a) + m + 1) - integrate(
((2*b*x*e^(2*a) + (m + 1)*e^(2*a))*e^(2*b*x) + m + 1)*x^m/((m + 1)*e^(4*b*x
+ 4*a) + 2*(m + 1)*e^(2*b*x + 2*a) + m + 1), x)
```

Giac [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(bx + a) dx$$

[In] integrate(x^m*tanh(b*x+a),x, algorithm="giac")

[Out] integrate(x^m*tanh(b*x + a), x)

Mupad [N/A]

Not integrable

Time = 1.70 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int x^m \tanh(a + bx) dx = \int x^m \tanh(a + bx) dx$$

[In] int(x^m*tanh(a + b*x),x)

[Out] int(x^m*tanh(a + b*x), x)

3.32 $\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx$

Optimal result	234
Rubi [A] (verified)	234
Mathematica [A] (verified)	236
Maple [A] (verified)	236
Fricas [A] (verification not implemented)	237
Sympy [F]	237
Maxima [A] (verification not implemented)	237
Giac [A] (verification not implemented)	238
Mupad [B] (verification not implemented)	239

Optimal result

Integrand size = 20, antiderivative size = 169

$$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx = \frac{3d^3x}{8af^3} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^2(c+dx)}{8f^4(a+a \tanh(e+fx))} - \frac{4f^3(a+a \tanh(e+fx))}{4f^3(a+a \tanh(e+fx))} - \frac{3d(c+dx)^2}{4f^2(a+a \tanh(e+fx))} - \frac{(c+dx)^3}{2f(a+a \tanh(e+fx))}$$

[Out] $3/8*d^3*x/a/f^3+3/8*d*(d*x+c)^2/a/f^2+1/4*(d*x+c)^3/a/f+1/8*(d*x+c)^4/a/d-3/8*d^3/f^4/(a+a*\tanh(f*x+e))-3/4*d^2*(d*x+c)/f^3/(a+a*\tanh(f*x+e))-3/4*d*(d*x+c)^2/f^2/(a+a*\tanh(f*x+e))-1/2*(d*x+c)^3/f/(a+a*\tanh(f*x+e))$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3804, 3560, 8}

$$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx = -\frac{3d^2(c+dx)}{4f^3(a \tanh(e+fx)+a)} - \frac{3d(c+dx)^2}{4f^2(a \tanh(e+fx)+a)} - \frac{(c+dx)^3}{2f(a \tanh(e+fx)+a)} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a \tanh(e+fx)+a)} + \frac{3d^3x}{8af^3}$$

[In] $\text{Int}[(c+d*x)^3/(a+a*\text{Tanh}[e+f*x]),x]$

[Out] $(3*d^3*x)/(8*a*f^3) + (3*d*(c + d*x)^2)/(8*a*f^2) + (c + d*x)^3/(4*a*f) + (c + d*x)^4/(8*a*d) - (3*d^3)/(8*f^4*(a + a*Tanh[e + f*x])) - (3*d^2*(c + d*x))/(4*f^3*(a + a*Tanh[e + f*x])) - (3*d*(c + d*x)^2)/(4*f^2*(a + a*Tanh[e + f*x])) - (c + d*x)^3/(2*f*(a + a*Tanh[e + f*x]))$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3804

Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(c + dx)^4}{8ad} - \frac{(c + dx)^3}{2f(a + a \tanh(e + fx))} + \frac{(3d) \int \frac{(c+dx)^2}{a+a \tanh(e+fx)} dx}{2f} \\
 &= \frac{(c + dx)^3}{4af} + \frac{(c + dx)^4}{8ad} - \frac{3d(c + dx)^2}{4f^2(a + a \tanh(e + fx))} \\
 &\quad - \frac{(c + dx)^3}{2f(a + a \tanh(e + fx))} + \frac{(3d^2) \int \frac{c+dx}{a+a \tanh(e+fx)} dx}{2f^2} \\
 &= \frac{3d(c + dx)^2}{8af^2} + \frac{(c + dx)^3}{4af} + \frac{(c + dx)^4}{8ad} - \frac{3d^2(c + dx)}{4f^3(a + a \tanh(e + fx))} \\
 &\quad - \frac{3d(c + dx)^2}{4f^2(a + a \tanh(e + fx))} - \frac{(c + dx)^3}{2f(a + a \tanh(e + fx))} + \frac{(3d^3) \int \frac{1}{a+a \tanh(e+fx)} dx}{4f^3} \\
 &= \frac{3d(c + dx)^2}{8af^2} + \frac{(c + dx)^3}{4af} + \frac{(c + dx)^4}{8ad} - \frac{3d^3}{8f^4(a + a \tanh(e + fx))} \\
 &\quad - \frac{3d^2(c + dx)}{4f^3(a + a \tanh(e + fx))} - \frac{3d(c + dx)^2}{4f^2(a + a \tanh(e + fx))} \\
 &\quad - \frac{(c + dx)^3}{2f(a + a \tanh(e + fx))} + \frac{(3d^3) \int 1 dx}{8af^3}
 \end{aligned}$$

$$= \frac{3d^3x}{8af^3} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+a \tanh(e+fx))} - \frac{3d^2(c+dx)}{4f^3(a+a \tanh(e+fx))} - \frac{3d(c+dx)^2}{4f^2(a+a \tanh(e+fx))} - \frac{(c+dx)^3}{2f(a+a \tanh(e+fx))}$$

Mathematica [A] (verified)

Time = 0.62 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.44

$$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx$$

$$= \frac{\operatorname{sech}(e+fx)(\cosh(fx)+\sinh(fx))((4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2)+d^3(3+6fx+6fx^2))}{\operatorname{sech}(e+fx)(\cosh(fx)+\sinh(fx))((4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2)+d^3(3+6fx+6fx^2))}$$

[In] Integrate[(c + d*x)^3/(a + a*Tanh[e + f*x]),x]

[Out] (Sech[e + f*x]*(Cosh[f*x] + Sinh[f*x])*((4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*Cosh[2*f*x]*(-Cosh[e] + Sinh[e]) + 2*f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cosh[e] + Sinh[e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*(Cosh[e] - Sinh[e])*Sinh[2*f*x]))/(16*a*f^4*(1 + Tanh[e + f*x]))

Maple [A] (verified)

Time = 0.23 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.98

method	result
risch	$\frac{d^3x^4}{8a} + \frac{d^2cx^3}{2a} + \frac{3dc^2x^2}{4a} + \frac{c^3x}{2a} + \frac{c^4}{8ad} - \frac{(4d^3x^3f^3+12cd^2f^3x^2+12c^2df^3x+6d^3f^2x^2+4c^3f^3+12cd^2f^2x+6c^2df^2+6d^3f^3)}{16af^4}$
parallelrisch	$-\frac{3d^3-6c^2df^3x+6x \tanh(fx+e)c d^2f^2+4x^3 \tanh(fx+e)c d^2f^4+6x^2 \tanh(fx+e)c^2d f^4+6x^2 \tanh(fx+e)c d^2f^3+6x \tanh(fx+e)c d^2f^2}{16af^4}$
default	Expression too large to display

[In] int((d*x+c)^3/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] 1/8/a*d^3*x^4+1/2/a*d^2*c*x^3+3/4/a*d*c^2*x^2+1/2/a*c^3*x+1/8/a/d*c^4-1/16*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4*c^3*f^3+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+3*d^3)/a/f^4*exp(-2*f*x-2*e)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 304, normalized size of antiderivative = 1.80

$$\int \frac{(c + dx)^3}{a + a \tanh(e + fx)} dx$$

$$= \frac{(2d^3f^4x^4 - 4c^3f^3 - 6c^2df^2 - 6cd^2f + 4(2cd^2f^4 - d^3f^3)x^3 - 3d^3 + 6(2c^2df^4 - 2cd^2f^3 - d^3f^2)x^2 + 2(2c^3f^4 - 6c^2df^3 - 6cd^2f^2 - 3d^3f)x - 3d^3 + 6(2c^2df^4 - 2cd^2f^3 - d^3f^2)x^2 + 2(2c^3f^4 - 6c^2df^3 - 6cd^2f^2 - 3d^3f)x) \cosh(fx + e) + (2d^3f^4x^4 + 4c^3f^3 + 6c^2df^2 + 6cd^2f + 4(2cd^2f^4 + d^3f^3)x^3 + 3d^3 + 6(2c^2df^4 + 2cd^2f^3 + d^3f^2)x^2 + 2(4c^3f^4 + 6c^2df^3 + 6cd^2f^2 + 3d^3f)x) \sinh(fx + e)}{a^2 f^4 \cosh(fx + e) + a^2 f^4 \sinh(fx + e)}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="fricas")

```
[Out] 1/16*((2*d^3*f^4*x^4 - 4*c^3*f^3 - 6*c^2*d*f^2 - 6*c*d^2*f + 4*(2*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 6*(2*c^2*d*f^4 - 2*c*d^2*f^3 - d^3*f^2)*x^2 + 2*(4*c^3*f^4 - 6*c^2*d*f^3 - 6*c*d^2*f^2 - 3*d^3*f)*x)*cosh(f*x + e) + (2*d^3*f^4*x^4 + 4*c^3*f^3 + 6*c^2*d*f^2 + 6*c*d^2*f + 4*(2*c*d^2*f^4 + d^3*f^3)*x^3 + 3*d^3 + 6*(2*c^2*d*f^4 + 2*c*d^2*f^3 + d^3*f^2)*x^2 + 2*(4*c^3*f^4 + 6*c^2*d*f^3 + 6*c*d^2*f^2 + 3*d^3*f)*x)*sinh(f*x + e))/(a*f^4*cosh(f*x + e) + a*f^4*sinh(f*x + e))
```

Sympy [F]

$$\int \frac{(c + dx)^3}{a + a \tanh(e + fx)} dx$$

$$= \frac{\int \frac{c^3}{\tanh(e+fx)+1} dx + \int \frac{d^3x^3}{\tanh(e+fx)+1} dx + \int \frac{3cd^2x^2}{\tanh(e+fx)+1} dx + \int \frac{3c^2dx}{\tanh(e+fx)+1} dx}{a}$$

[In] integrate((d*x+c)**3/(a+a*tanh(f*x+e)),x)

```
[Out] (Integral(c**3/(tanh(e + f*x) + 1), x) + Integral(d**3*x**3/(tanh(e + f*x) + 1), x) + Integral(3*c*d**2*x**2/(tanh(e + f*x) + 1), x) + Integral(3*c**2*d*x/(tanh(e + f*x) + 1), x))/a
```

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.10

$$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx = \frac{1}{4} c^3 \left(\frac{2(fx+e)}{af} - \frac{e^{(-2fx-2e)}}{af} \right) + \frac{3(2f^2x^2e^{(2e)} - (2fx+1)e^{(-2fx)})c^2de^{(-2e)}}{8af^2} + \frac{(4f^3x^3e^{(2e)} - 3(2f^2x^2 + 2fx+1)e^{(-2fx)})cd^2e^{(-2e)}}{8af^3} + \frac{(2f^4x^4e^{(2e)} - (4f^3x^3 + 6f^2x^2 + 6fx+3)e^{(-2fx)})d^3e^{(-2e)}}{16af^4}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/4*c^3*(2*(f*x + e)/(a*f) - e^(-2*f*x - 2*e)/(a*f)) + 3/8*(2*f^2*x^2*e^(2*e) - (2*f*x + 1)*e^(-2*f*x))*c^2*d*e^(-2*e)/(a*f^2) + 1/8*(4*f^3*x^3*e^(2*e) - 3*(2*f^2*x^2 + 2*f*x + 1)*e^(-2*f*x))*c*d^2*e^(-2*e)/(a*f^3) + 1/16*(2*f^4*x^4*e^(2*e) - (4*f^3*x^3 + 6*f^2*x^2 + 6*f*x + 3)*e^(-2*f*x))*d^3*e^(-2*e)/(a*f^4)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.11

$$\int \frac{(c+dx)^3}{a+a \tanh(e+fx)} dx = \frac{(2d^3f^4x^4e^{(2fx+2e)} + 8cd^2f^4x^3e^{(2fx+2e)} + 12c^2df^4x^2e^{(2fx+2e)} - 4d^3f^3x^3 + 8c^3f^4xe^{(2fx+2e)} - 12cd^2f^3x^2)}{16af^4}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] 1/16*(2*d^3*f^4*x^4*e^(2*f*x + 2*e) + 8*c*d^2*f^4*x^3*e^(2*f*x + 2*e) + 12*c^2*d*f^4*x^2*e^(2*f*x + 2*e) - 4*d^3*f^3*x^3 + 8*c^3*f^4*x*e^(2*f*x + 2*e) - 12*c*d^2*f^3*x^2 - 12*c^2*d*f^3*x - 6*d^3*f^2*x^2 - 4*c^3*f^3 - 12*c*d^2*f^2*x - 6*c^2*d*f^2 - 6*d^3*f*x - 6*c*d^2*f - 3*d^3)*e^(-2*f*x - 2*e)/(a*f^4)

Mupad [B] (verification not implemented)

Time = 1.98 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.62

$$\int \frac{(c + dx)^3}{a + a \tanh(e + fx)} dx$$

$$= \frac{e^{-2e-2fx} (8c^3 x e^{2e+2fx} + 2d^3 x^4 e^{2e+2fx} + 12c^2 d x^2 e^{2e+2fx} + 8cd^2 x^3 e^{2e+2fx})}{e^{-2e-2fx} (3d^3 - 3d^3 e^{2e+2fx})} + \frac{16a}{16} \frac{f^2 e^{-2e-2fx} (6c^2 d + 6d^3 x^2 - 6c^2 d e^{2e+2fx} + 12cd^2 x)}{16} + \frac{f e^{-2e-2fx} (6cd^2 + 6d^3 x - 6cd^2 e^{2e+2fx})}{16} + \frac{f^4}{af^4}$$

[In] int((c + d*x)^3/(a + a*tanh(e + f*x)),x)

```
[Out] (exp(- 2*e - 2*f*x)*(8*c^3*x*exp(2*e + 2*f*x) + 2*d^3*x^4*exp(2*e + 2*f*x)
+ 12*c^2*d*x^2*exp(2*e + 2*f*x) + 8*c*d^2*x^3*exp(2*e + 2*f*x)))/(16*a) - (
(exp(- 2*e - 2*f*x)*(3*d^3 - 3*d^3*exp(2*e + 2*f*x)))/16 + (f^2*exp(- 2*e -
2*f*x)*(6*c^2*d + 6*d^3*x^2 - 6*c^2*d*exp(2*e + 2*f*x) + 12*c*d^2*x))/16 +
(f*exp(- 2*e - 2*f*x)*(6*c*d^2 + 6*d^3*x - 6*c*d^2*exp(2*e + 2*f*x)))/16 +
(f^3*exp(- 2*e - 2*f*x)*(4*c^3 - 4*c^3*exp(2*e + 2*f*x) + 4*d^3*x^3 + 12*c
*d^2*x^2 + 12*c^2*d*x))/16)/(a*f^4)
```

3.33 $\int \frac{(c+dx)^2}{a+a \tanh(e+fx)} dx$

Optimal result	240
Rubi [A] (verified)	240
Mathematica [A] (verified)	242
Maple [A] (verified)	242
Fricas [A] (verification not implemented)	243
Sympy [F]	243
Maxima [A] (verification not implemented)	243
Giac [A] (verification not implemented)	244
Mupad [B] (verification not implemented)	244

Optimal result

Integrand size = 20, antiderivative size = 122

$$\int \frac{(c+dx)^2}{a+a \tanh(e+fx)} dx = \frac{d^2x}{4af^2} + \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a+a \tanh(e+fx))} - \frac{d(c+dx)}{2f^2(a+a \tanh(e+fx))} - \frac{(c+dx)^2}{2f(a+a \tanh(e+fx))}$$

[Out] 1/4*d^2*x/a/f^2+1/4*(d*x+c)^2/a/f+1/6*(d*x+c)^3/a/d-1/4*d^2/f^3/(a+a*tanh(f*x+e))-1/2*d*(d*x+c)/f^2/(a+a*tanh(f*x+e))-1/2*(d*x+c)^2/f/(a+a*tanh(f*x+e))

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3804, 3560, 8}

$$\int \frac{(c+dx)^2}{a+a \tanh(e+fx)} dx = -\frac{d(c+dx)}{2f^2(a \tanh(e+fx)+a)} - \frac{(c+dx)^2}{2f(a \tanh(e+fx)+a)} + \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a \tanh(e+fx)+a)} + \frac{d^2x}{4af^2}$$

[In] Int[(c + d*x)^2/(a + a*Tanh[e + f*x]),x]

[Out] (d^2*x)/(4*a*f^2) + (c + d*x)^2/(4*a*f) + (c + d*x)^3/(6*a*d) - d^2/(4*f^3*(a + a*Tanh[e + f*x])) - (d*(c + d*x))/(2*f^2*(a + a*Tanh[e + f*x])) - (c + d*x)^2/(2*f*(a + a*Tanh[e + f*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3804

Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))], x)) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(c + dx)^3}{6ad} - \frac{(c + dx)^2}{2f(a + a \tanh(e + fx))} + \frac{d \int \frac{c+dx}{a+a \tanh(e+fx)} dx}{f} \\
 &= \frac{(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} - \frac{d(c + dx)}{2f^2(a + a \tanh(e + fx))} \\
 &\quad - \frac{(c + dx)^2}{2f(a + a \tanh(e + fx))} + \frac{d^2 \int \frac{1}{a+a \tanh(e+fx)} dx}{2f^2} \\
 &= \frac{(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} - \frac{d^2}{4f^3(a + a \tanh(e + fx))} \\
 &\quad - \frac{d(c + dx)}{2f^2(a + a \tanh(e + fx))} - \frac{(c + dx)^2}{2f(a + a \tanh(e + fx))} + \frac{d^2 \int 1 dx}{4af^2} \\
 &= \frac{d^2 x}{4af^2} + \frac{(c + dx)^2}{4af} + \frac{(c + dx)^3}{6ad} - \frac{d^2}{4f^3(a + a \tanh(e + fx))} \\
 &\quad - \frac{d(c + dx)}{2f^2(a + a \tanh(e + fx))} - \frac{(c + dx)^2}{2f(a + a \tanh(e + fx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.38 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.39

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx$$

$$= \frac{\operatorname{sech}(e + fx)(\cosh(fx) + \sinh(fx))((2c^2f^2 + 2cdf(1 + 2fx) + d^2(1 + 2fx + 2f^2x^2))\cosh(2fx)(-\cosh(e) + \sinh(e)) + (4f^3x^3 + 3c^2 + 3cdx + d^2x^2)(\cosh(e) + \sinh(e)))/3 + (2c^2f^2 + 2cdf(1 + 2fx) + d^2(1 + 2fx + 2f^2x^2))(\cosh(e) - \sinh(e))\sinh(2fx)}{8af^3(1 + \tanh(e + fx))}$$

[In] Integrate[(c + d*x)^2/(a + a*Tanh[e + f*x]),x]

```
[Out] (Sech[e + f*x]*(Cosh[f*x] + Sinh[f*x])*((2*c^2*f^2 + 2*c*d*f*(1 + 2*f*x) +
d^2*(1 + 2*f*x + 2*f^2*x^2))*Cosh[2*f*x]*(-Cosh[e] + Sinh[e]) + (4*f^3*x*(3
*c^2 + 3*c*d*x + d^2*x^2)*(Cosh[e] + Sinh[e]))/3 + (2*c^2*f^2 + 2*c*d*f*(1
+ 2*f*x) + d^2*(1 + 2*f*x + 2*f^2*x^2))*(Cosh[e] - Sinh[e])*Sinh[2*f*x]))/(
8*a*f^3*(1 + Tanh[e + f*x]))
```

Maple [A] (verified)

Time = 0.20 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.84

method	result
risch	$\frac{d^2x^3}{6a} + \frac{dcx^2}{2a} + \frac{c^2x}{2a} + \frac{c^3}{6ad} - \frac{(2d^2x^2f^2 + 4cdf^2x + 2c^2f^2 + 2d^2fx + 2cdf + d^2)e^{-2fx-2e}}{8af^3}$
parallelrisch	$\frac{-3d^2 - 3d^2fx - 6c^2f^2 + 6cdx^2f^3 - 6cdf^2x - 3d^2x^2f^2 - 6cdf + 6xc^2f^3 + 2d^2x^3f^3 + 2d^2 \tanh(fx+e)x^3f^3 + 6x \tanh(fx+e)c^2f^3}{12f^3a(1 + \tanh(fx+e))}$
derivativedivides	$-\frac{\cosh(fx+e)^2c^2f^2}{2} + \cosh(fx+e)^2dcf - 2cdf \left(\frac{(fx+e)\cosh(fx+e)^2}{2} - \frac{\cosh(fx+e)\sinh(fx+e)}{4} - \frac{fx}{4} - \frac{e}{4} \right) - \frac{\cosh(fx+e)^2d^2e^2}{2} + 2d^2fx$
default	$-\frac{\cosh(fx+e)^2c^2f^2}{2} + \cosh(fx+e)^2dcf - 2cdf \left(\frac{(fx+e)\cosh(fx+e)^2}{2} - \frac{\cosh(fx+e)\sinh(fx+e)}{4} - \frac{fx}{4} - \frac{e}{4} \right) - \frac{\cosh(fx+e)^2d^2e^2}{2} + 2d^2fx$

[In] int((d*x+c)^2/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)

```
[Out] 1/6/a*d^2*x^3+1/2/a*d*c*x^2+1/2/a*c^2*x+1/6/a/d*c^3-1/8*(2*d^2*f^2*x^2+4*c*
d*f^2*x+2*c^2*f^2+2*d^2*f*x+2*c*d*f+d^2)/a/f^3*exp(-2*f*x-2*e)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.57

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx = \frac{(4d^2 f^3 x^3 - 6c^2 f^2 - 6cdf + 6(2cdf^3 - d^2 f^2)x^2 - 3d^2 + 6(2c^2 f^3 - 2cdf^2 - d^2 f)x) \cosh(fx + e) + (4d^2 f^3 x^3 - 6c^2 f^2 - 6cdf + 6(2cdf^3 - d^2 f^2)x^2 - 3d^2 + 6(2c^2 f^3 - 2cdf^2 - d^2 f)x) \sinh(fx + e)}{24(a f^3 \cosh(fx + e) + a f^3 \sinh(fx + e))}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="fricas")

```
[Out] 1/24*((4*d^2*f^3*x^3 - 6*c^2*f^2 - 6*c*d*f + 6*(2*c*d*f^3 - d^2*f^2)*x^2 - 3*d^2 + 6*(2*c^2*f^3 - 2*c*d*f^2 - d^2*f)*x)*cosh(f*x + e) + (4*d^2*f^3*x^3 + 6*c^2*f^2 + 6*c*d*f + 6*(2*c*d*f^3 + d^2*f^2)*x^2 + 3*d^2 + 6*(2*c^2*f^3 + 2*c*d*f^2 + d^2*f)*x)*sinh(f*x + e))/(a*f^3*cosh(f*x + e) + a*f^3*sinh(f*x + e))
```

Sympy [F]

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx = \frac{\int \frac{c^2}{\tanh(e+fx)+1} dx + \int \frac{d^2 x^2}{\tanh(e+fx)+1} dx + \int \frac{2cdx}{\tanh(e+fx)+1} dx}{a}$$

[In] integrate((d*x+c)**2/(a+a*tanh(f*x+e)),x)

```
[Out] (Integral(c**2/(tanh(e + f*x) + 1), x) + Integral(d**2*x**2/(tanh(e + f*x) + 1), x) + Integral(2*c*d*x/(tanh(e + f*x) + 1), x))/a
```

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 126, normalized size of antiderivative = 1.03

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx = \frac{1}{4} c^2 \left(\frac{2(fx + e)}{af} - \frac{e^{(-2fx-2e)}}{af} \right) + \frac{(2f^2 x^2 e^{(2e)} - (2fx + 1)e^{(-2fx)}) c d e^{(-2e)}}{4af^2} + \frac{(4f^3 x^3 e^{(2e)} - 3(2f^2 x^2 + 2fx + 1)e^{(-2fx)}) d^2 e^{(-2e)}}{24af^3}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="maxima")

```
[Out] 1/4*c^2*(2*(f*x + e)/(a*f) - e^(-2*f*x - 2*e)/(a*f)) + 1/4*(2*f^2*x^2*e^(2*e) - (2*f*x + 1)*e^(-2*f*x))*c*d*e^(-2*e)/(a*f^2) + 1/24*(4*f^3*x^3*e^(2*e) - 3*(2*f^2*x^2 + 2*f*x + 1)*e^(-2*f*x))*d^2*e^(-2*e)/(a*f^3)
```

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx = \frac{(4d^2 f^3 x^3 e^{(2fx+2e)} + 12cdf^3 x^2 e^{(2fx+2e)} + 12c^2 f^3 x e^{(2fx+2e)} - 6d^2 f^2 x^2 - 12cdf^2 x - 6c^2 f^2 - 6d^2 fx - 6cd)}{24af^3}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] 1/24*(4*d^2*f^3*x^3*e^(2*f*x + 2*e) + 12*c*d*f^3*x^2*e^(2*f*x + 2*e) + 12*c^2*f^3*x*e^(2*f*x + 2*e) - 6*d^2*f^2*x^2 - 12*c*d*f^2*x - 6*c^2*f^2 - 6*d^2*f*x - 6*c*d*f - 3*d^2)*e^(-2*f*x - 2*e)/(a*f^3)

Mupad [B] (verification not implemented)

Time = 1.85 (sec) , antiderivative size = 187, normalized size of antiderivative = 1.53

$$\int \frac{(c + dx)^2}{a + a \tanh(e + fx)} dx = \frac{e^{-2e-2fx} (12c^2 x e^{2e+2fx} + 4d^2 x^3 e^{2e+2fx} + 12cdx^2 e^{2e+2fx})}{24a} - \frac{e^{-2e-2fx} (3d^2 - 3d^2 e^{2e+2fx})}{24} + \frac{f e^{-2e-2fx} (6cd + 6d^2 x - 6cde^{2e+2fx})}{24} + \frac{f^2 e^{-2e-2fx} (6c^2 - 6c^2 e^{2e+2fx} + 6d^2 x^2 + 12cdx)}{24}$$

[In] int((c + d*x)^2/(a + a*tanh(e + f*x)),x)

[Out] (exp(- 2*e - 2*f*x)*(12*c^2*x*exp(2*e + 2*f*x) + 4*d^2*x^3*exp(2*e + 2*f*x) + 12*c*d*x^2*exp(2*e + 2*f*x)))/(24*a) - ((exp(- 2*e - 2*f*x)*(3*d^2 - 3*d^2*exp(2*e + 2*f*x)))/24 + (f*exp(- 2*e - 2*f*x)*(6*c*d + 6*d^2*x - 6*c*d*exp(2*e + 2*f*x)))/24 + (f^2*exp(- 2*e - 2*f*x)*(6*c^2 - 6*c^2*exp(2*e + 2*f*x) + 6*d^2*x^2 + 12*c*d*x))/24)/(a*f^3)

3.34 $\int \frac{c+dx}{a+a \tanh(e+fx)} dx$

Optimal result	245
Rubi [A] (verified)	245
Mathematica [A] (verified)	246
Maple [A] (verified)	247
Fricas [A] (verification not implemented)	247
Sympy [F]	247
Maxima [A] (verification not implemented)	248
Giac [A] (verification not implemented)	248
Mupad [B] (verification not implemented)	248

Optimal result

Integrand size = 18, antiderivative size = 74

$$\int \frac{c+dx}{a+a \tanh(e+fx)} dx = \frac{dx}{4af} + \frac{(c+dx)^2}{4ad} - \frac{d}{4f^2(a+a \tanh(e+fx))} - \frac{c+dx}{2f(a+a \tanh(e+fx))}$$

[Out] $1/4*d*x/a/f+1/4*(d*x+c)^2/a/d-1/4*d/f^2/(a+a*\tanh(f*x+e))+1/2*(-d*x-c)/f/(a+a*\tanh(f*x+e))$

Rubi [A] (verified)

Time = 0.04 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3804, 3560, 8}

$$\int \frac{c+dx}{a+a \tanh(e+fx)} dx = -\frac{c+dx}{2f(a \tanh(e+fx)+a)} + \frac{(c+dx)^2}{4ad} - \frac{d}{4f^2(a \tanh(e+fx)+a)} + \frac{dx}{4af}$$

[In] $\text{Int}[(c+d*x)/(a+a*\text{Tanh}[e+f*x]),x]$

[Out] $(d*x)/(4*a*f) + (c+d*x)^2/(4*a*d) - d/(4*f^2*(a+a*\text{Tanh}[e+f*x])) - (c+d*x)/(2*f*(a+a*\text{Tanh}[e+f*x]))$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3560

```
Int[((a_) + (b_.)*tan[(c_) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[a*((a +
b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(
n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*tan[(e_) + (f_.)*(x_)]), x_Sy
mbol] := Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f))
, Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/
(2*b*f*(a + b*Tan[e + f*x]))], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a
^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(c + dx)^2}{4ad} - \frac{c + dx}{2f(a + a \tanh(e + fx))} + \frac{d \int \frac{1}{a + a \tanh(e + fx)} dx}{2f} \\ &= \frac{(c + dx)^2}{4ad} - \frac{d}{4f^2(a + a \tanh(e + fx))} - \frac{c + dx}{2f(a + a \tanh(e + fx))} + \frac{d \int 1 dx}{4af} \\ &= \frac{dx}{4af} + \frac{(c + dx)^2}{4ad} - \frac{d}{4f^2(a + a \tanh(e + fx))} - \frac{c + dx}{2f(a + a \tanh(e + fx))} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.51 (sec) , antiderivative size = 81, normalized size of antiderivative = 1.09

$$\begin{aligned} &\int \frac{c + dx}{a + a \tanh(e + fx)} dx \\ &= \frac{2cf(-1 + 2fx) + d(-1 - 2fx + 2f^2x^2) + (2cf(1 + 2fx) + d(1 + 2fx + 2f^2x^2)) \tanh(e + fx)}{8af^2(1 + \tanh(e + fx))} \end{aligned}$$

```
[In] Integrate[(c + d*x)/(a + a*Tanh[e + f*x]),x]
```

```
[Out] (2*c*f*(-1 + 2*f*x) + d*(-1 - 2*f*x + 2*f^2*x^2) + (2*c*f*(1 + 2*f*x) + d*(
1 + 2*f*x + 2*f^2*x^2))*Tanh[e + f*x])/(8*a*f^2*(1 + Tanh[e + f*x]))
```

Maple [A] (verified)

Time = 0.18 (sec) , antiderivative size = 46, normalized size of antiderivative = 0.62

method	result
risch	$\frac{dx^2}{4a} + \frac{cx}{2a} - \frac{(2dx+2cf+d)e^{-2fx-2e}}{8af^2}$
parallelrisch	$\frac{d \tanh(fx+e)x^2 f^2 + dx^2 f^2 + 2x \tanh(fx+e)cf^2 + d \tanh(fx+e)xf + 2cx f^2 - dx f - 2cf - d}{4f^2 a(1+\tanh(fx+e))}$
default	$-\frac{d\left(\frac{(fx+e)\cosh(fx+e)^2}{2} - \frac{\cosh(fx+e)\sinh(fx+e)}{4} - \frac{fx}{4} - \frac{e}{4}\right)}{f} + \frac{d\left(\frac{(fx+e)\cosh(fx+e)\sinh(fx+e)}{2} + \frac{(fx+e)^2}{4} - \frac{\cosh(fx+e)^2}{4}\right)}{f} + \frac{de \cosh(fx+e)}{2f}$

```
[In] int((d*x+c)/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)
```

```
[Out] 1/4/a*d*x^2+1/2/a*c*x-1/8*(2*d*f*x+2*c*f+d)/a/f^2*exp(-2*f*x-2*e)
```

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.36

$$\int \frac{c + dx}{a + a \tanh(e + fx)} dx$$

$$= \frac{(2df^2x^2 - 2cf + 2(2cf^2 - df)x - d) \cosh(fx + e) + (2df^2x^2 + 2cf + 2(2cf^2 + df)x + d) \sinh(fx + e)}{8(a f^2 \cosh(fx + e) + a f^2 \sinh(fx + e))}$$

```
[In] integrate((d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="fricas")
```

```
[Out] 1/8*((2*d*f^2*x^2 - 2*c*f + 2*(2*c*f^2 - d*f)*x - d)*cosh(f*x + e) + (2*d*f^2*x^2 + 2*c*f + 2*(2*c*f^2 + d*f)*x + d)*sinh(f*x + e))/(a*f^2*cosh(f*x + e) + a*f^2*sinh(f*x + e))
```

Sympy [F]

$$\int \frac{c + dx}{a + a \tanh(e + fx)} dx = \frac{\int \frac{c}{\tanh(e+fx)+1} dx + \int \frac{dx}{\tanh(e+fx)+1} dx}{a}$$

```
[In] integrate((d*x+c)/(a+a*tanh(f*x+e)),x)
```

```
[Out] (Integral(c/(tanh(e + f*x) + 1), x) + Integral(d*x/(tanh(e + f*x) + 1), x))/a
```

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 74, normalized size of antiderivative = 1.00

$$\int \frac{c + dx}{a + a \tanh(e + fx)} dx = \frac{1}{4} c \left(\frac{2(fx + e)}{af} - \frac{e^{(-2fx-2e)}}{af} \right) + \frac{(2f^2x^2e^{(2e)} - (2fx + 1)e^{(-2fx)})de^{(-2e)}}{8af^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/4*c*(2*(f*x + e)/(a*f) - e^(-2*f*x - 2*e)/(a*f)) + 1/8*(2*f^2*x^2*e^(2*e) - (2*f*x + 1)*e^(-2*f*x))*d*e^(-2*e)/(a*f^2)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 64, normalized size of antiderivative = 0.86

$$\int \frac{c + dx}{a + a \tanh(e + fx)} dx = \frac{(2df^2x^2e^{(2fx+2e)} + 4cf^2xe^{(2fx+2e)} - 2dfx - 2cf - d)e^{(-2fx-2e)}}{8af^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] 1/8*(2*d*f^2*x^2*e^(2*f*x + 2*e) + 4*c*f^2*x*e^(2*f*x + 2*e) - 2*d*f*x - 2*c*f - d)*e^(-2*f*x - 2*e)/(a*f^2)

Mupad [B] (verification not implemented)

Time = 1.77 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.03

$$\int \frac{c + dx}{a + a \tanh(e + fx)} dx = \frac{\frac{dx^2}{4} + \left(\frac{c}{2} + \frac{d}{4f}\right)x}{a} - \frac{\frac{\frac{d}{4} + \frac{cf}{2}}{f^2} - x\left(\frac{c}{2} - \frac{d}{4f}\right) + x\left(\frac{c}{2} + \frac{d}{4f}\right)}{a + a \tanh(e + fx)}$$

[In] int((c + d*x)/(a + a*tanh(e + f*x)),x)

[Out] (x*(c/2 + d/(4*f)) + (d*x^2)/4)/a - ((d/4 + (c*f)/2)/f^2 - x*(c/2 - d/(4*f)) + x*(c/2 + d/(4*f)))/(a + a*tanh(e + f*x))

3.35 $\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx$

Optimal result	249
Rubi [A] (verified)	249
Mathematica [A] (verified)	251
Maple [A] (verified)	251
Fricas [A] (verification not implemented)	251
Sympy [F]	252
Maxima [A] (verification not implemented)	252
Giac [A] (verification not implemented)	252
Mupad [F(-1)]	253

Optimal result

Integrand size = 20, antiderivative size = 157

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx = \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2ad} + \frac{\log(c+dx)}{2ad} - \frac{\operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2ad} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2ad} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2ad}$$

[Out] $1/2*\operatorname{Chi}(2*c*f/d+2*f*x)*\cosh(-2*e+2*c*f/d)/a/d+1/2*\ln(d*x+c)/a/d-1/2*\cosh(-2*e+2*c*f/d)*\operatorname{Shi}(2*c*f/d+2*f*x)/a/d+1/2*\operatorname{Chi}(2*c*f/d+2*f*x)*\sinh(-2*e+2*c*f/d)/a/d-1/2*\operatorname{Shi}(2*c*f/d+2*f*x)*\sinh(-2*e+2*c*f/d)/a/d$

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3807, 3384, 3379, 3382}

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx = -\frac{\operatorname{Chi}\left(2xf + \frac{2cf}{d}\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2ad} + \frac{\operatorname{Chi}\left(2xf + \frac{2cf}{d}\right) \cosh\left(2e - \frac{2cf}{d}\right)}{2ad} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(2xf + \frac{2cf}{d}\right)}{2ad} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(2xf + \frac{2cf}{d}\right)}{2ad} + \frac{\log(c+dx)}{2ad}$$

[In] Int[1/((c + d*x)*(a + a*Tanh[e + f*x])),x]

[Out] (Cosh[2*e - (2*c*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x]/(2*a*d) + Log[c + d*x]/(2*a*d) - (CoshIntegral[(2*c*f)/d + 2*f*x]*Sinh[2*e - (2*c*f)/d]/(2*a*d) - (Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x]/(2*a*d) + (Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x]/(2*a*d)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3807

Int[1/(((c_.) + (d_.)*(x_))*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := Simp[Log[c + d*x]/(2*a*d), x] + (Dist[1/(2*a), Int[Cos[2*e + 2*f*x]/(c + d*x), x], x] + Dist[1/(2*b), Int[Sin[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{\log(c + dx)}{2ad} + \frac{\int \frac{\cosh(2e+2fx)}{c+dx} dx}{2a} - \frac{\int \frac{\sinh(2e+2fx)}{c+dx} dx}{2a} \\
 &= \frac{\log(c + dx)}{2ad} + \frac{\cosh\left(2e - \frac{2cf}{d}\right) \int \frac{\cosh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \int \frac{\sinh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a} \\
 &\quad - \frac{\sinh\left(2e - \frac{2cf}{d}\right) \int \frac{\cosh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \int \frac{\sinh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a} \\
 &= \frac{\cosh\left(2e - \frac{2cf}{d}\right) \text{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2ad} + \frac{\log(c + dx)}{2ad} - \frac{\text{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2ad} \\
 &\quad - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2ad} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2ad}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.43 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.78

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx$$

$$= \frac{\operatorname{sech}(e+fx)(\cosh(fx)+\sinh(fx)) \left(\log(f(c+dx))(\cosh(e)+\sinh(e)) + \operatorname{Chi}\left(\frac{2f(c+dx)}{d}\right) \left(\cosh\left(e-\frac{2cf}{d}\right) \right) \right)}{2ad(1+\tanh(e+fx))}$$

[In] Integrate[1/((c + d*x)*(a + a*Tanh[e + f*x])),x]

[Out] (Sech[e + f*x]*(Cosh[f*x] + Sinh[f*x])*(Log[f*(c + d*x)]*(Cosh[e] + Sinh[e]) + CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[e - (2*c*f)/d] - Sinh[e - (2*c*f)/d]) + (-Cosh[e - (2*c*f)/d] + Sinh[e - (2*c*f)/d])*SinhIntegral[(2*f*(c + d*x))/d]))/(2*a*d*(1 + Tanh[e + f*x]))

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.39

method	result	size
risch	$\frac{\ln(dx+c)}{2ad} - \frac{e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{2ad}$	61

[In] int(1/(d*x+c)/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] 1/2*ln(d*x+c)/a/d-1/2/a/d*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 73, normalized size of antiderivative = 0.46

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx$$

$$= \frac{\operatorname{Ei}\left(-\frac{2(dfxc+cf)}{d}\right) \cosh\left(-\frac{2(de-cf)}{d}\right) + \operatorname{Ei}\left(-\frac{2(dfxc+cf)}{d}\right) \sinh\left(-\frac{2(de-cf)}{d}\right) + \log(dx+c)}{2ad}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="fricas")

[Out] 1/2*(Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + Ei(-2*(d*f*x + c*f)/d)*sinh(-2*(d*e - c*f)/d) + log(d*x + c))/(a*d)

Sympy [F]

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx = \frac{\int \frac{1}{c \tanh(e+fx)+c+dx \tanh(e+fx)+dx} dx}{a}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e)),x)

[Out] Integral(1/(c*tanh(e + f*x) + c + d*x*tanh(e + f*x) + d*x), x)/a

Maxima [A] (verification not implemented)

none

Time = 0.35 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.31

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx = -\frac{e^{(-2e+\frac{2cf}{d})} E_1\left(\frac{2(dx+c)f}{d}\right)}{2ad} + \frac{\log(dx+c)}{2ad}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] -1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(1, 2*(d*x + c)*f/d)/(a*d) + 1/2*log(d*x + c)/(a*d)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 47, normalized size of antiderivative = 0.30

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))} dx = \frac{\left(Ei\left(-\frac{2(dfx+cf)}{d}\right) e^{\left(\frac{2cf}{d}\right)} + e^{(2e)} \log(dx+c) \right) e^{(-2e)}}{2ad}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] 1/2*(Ei(-2*(d*f*x + c*f)/d)*e^(2*c*f/d) + e^(2*e)*log(d*x + c))*e^(-2*e)/(a*d)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + a \tanh(e + fx))} dx = \int \frac{1}{(a + a \tanh(e + fx)) (c + dx)} dx$$

```
[In] int(1/((a + a*tanh(e + f*x))*(c + d*x)),x)
```

```
[Out] int(1/((a + a*tanh(e + f*x))*(c + d*x)), x)
```

3.36 $\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx$

Optimal result	254
Rubi [A] (verified)	254
Mathematica [A] (verified)	256
Maple [A] (verified)	257
Fricas [A] (verification not implemented)	257
Sympy [F]	257
Maxima [A] (verification not implemented)	258
Giac [B] (verification not implemented)	258
Mupad [F(-1)]	259

Optimal result

Integrand size = 20, antiderivative size = 159

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = -\frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Chi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} + \frac{f \text{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{ad^2} + \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} - \frac{f \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} - \frac{1}{d(c+dx)(a+a \tanh(e+fx))}$$

[Out] -f*Chi(2*c*f/d+2*f*x)*cosh(-2*e+2*c*f/d)/a/d^2+f*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d+2*f*x)/a/d^2-f*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^2+f*Shi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^2-1/d/(d*x+c)/(a+a*tanh(f*x+e))

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used

= {3805, 3384, 3379, 3382}

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{ad^2} - \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{ad^2} - \frac{f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{ad^2} + \frac{f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{ad^2} - \frac{1}{d(c+dx)(a \tanh(e+fx) + a)}$$

[In] Int[1/((c + d*x)^2*(a + a*Tanh[e + f*x])),x]

[Out] -((f*Cosh[2*e - (2*c*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x]/(a*d^2)) + (f*CoshIntegral[(2*c*f)/d + 2*f*x]*Sinh[2*e - (2*c*f)/d]/(a*d^2) + (f*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x]/(a*d^2) - (f*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x]/(a*d^2) - 1/(d*(c + d*x)*(a + a*Tanh[e + f*x])))

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3805

Int[1/(((c_.) + (d_.)*(x_))^2*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol] := -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Dist[f/(a*d), Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Dist[f/(b*d), Int[Cos[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{1}{d(c+dx)(a+a \tanh(e+fx))} - \frac{f \int \frac{\cosh(2e+2fx)}{c+dx} dx}{ad} + \frac{f \int \frac{\sinh(2e+2fx)}{c+dx} dx}{ad} \\
 &= -\frac{1}{d(c+dx)(a+a \tanh(e+fx))} - \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
 &\quad + \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
 &\quad + \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} - \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
 &= -\frac{f \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{ad^2} + \frac{f \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{ad^2} \\
 &\quad + \frac{f \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^2} \\
 &\quad - \frac{f \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^2} - \frac{1}{d(c+dx)(a+a \tanh(e+fx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.67 (sec) , antiderivative size = 206, normalized size of antiderivative = 1.30

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = \frac{\text{sech}(e+fx) \left(\cosh\left(\frac{cf}{d}\right) + \sinh\left(\frac{cf}{d}\right) \right) \left(d \left(\cosh\left(e+f\left(-\frac{c}{d}+x\right)\right) + \cosh\left(e+f\left(\frac{c}{d}+x\right)\right) \right) + \sinh\left(e+f\left(-\frac{c}{d}+x\right)\right) - \sinh\left(e+f\left(\frac{c}{d}+x\right)\right) \right)}{(a+d^2x)^2}$$

[In] Integrate[1/((c+d*x)^2*(a+a*Tanh[e+f*x])),x]

[Out] -1/2*(Sech[e+f*x]*(Cosh[(c*f)/d]+Sinh[(c*f)/d])*(d*(Cosh[e+f*(-(c/d)+x]]+Cosh[e+f*(c/d+x]])+Sinh[e+f*(-(c/d)+x]])-Sinh[e+f*(c/d+x]])+2*f*(c+d*x)*CoshIntegral[(2*f*(c+d*x))/d]*(Cosh[e-(f*(c+d*x))/d]-Sinh[e-(f*(c+d*x))/d])]+2*f*(c+d*x)*(-Cosh[e-(f*(c+d*x))/d]+Sinh[e-(f*(c+d*x))/d])*SinhIntegral[(2*f*(c+d*x))/d]))/(a*d^2*(c+d*x)*(1+Tanh[e+f*x]))

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 90, normalized size of antiderivative = 0.57

method	result	size
risch	$-\frac{1}{2da(dx+c)} - \frac{f e^{-2fx-2e}}{2ad(dx+f+cf)} + \frac{f e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{a d^2}$	90

[In] int(1/(d*x+c)^2/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] -1/2/d/a/(d*x+c)-1/2*f/a*exp(-2*f*x-2*e)/d/(d*f*x+c*f)+f/a/d^2*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.36

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = \frac{(dfx+cf)\operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) \cosh(fx+e) \sinh\left(-\frac{2(de-cf)}{d}\right) + \left((dfx+cf)\operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right)\right) \cosh\left(-\frac{2(de-cf)}{d}\right)}{(ad^3x+acd^2)}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="fricas")

[Out] -((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)*sinh(-2*(d*e - c*f)/d) + ((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + d)*cosh(f*x + e) + ((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*sinh(-2*(d*e - c*f)/d)*sinh(f*x + e))/(a*d^3*x + a*c*d^2)*cosh(f*x + e) + (a*d^3*x + a*c*d^2)*sinh(f*x + e)

Sympy [F]

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = \frac{\int \frac{1}{c^2 \tanh(e+fx)+c^2+2cdx \tanh(e+fx)+2cdx+d^2x^2 \tanh(e+fx)+d^2x^2} dx}{a}$$

[In] integrate(1/(d*x+c)**2/(a+a*tanh(f*x+e)),x)

[Out] Integral(1/(c**2*tanh(e + f*x) + c**2 + 2*c*d*x*tanh(e + f*x) + 2*c*d*x + d**2*x**2*tanh(e + f*x) + d**2*x**2), x)/a

Maxima [A] (verification not implemented)

none

Time = 0.49 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.35

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = -\frac{1}{2(ad^2x+acd)} - \frac{e^{(-2e+\frac{2cf}{d})} E_2\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)ad}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] -1/2/(a*d^2*x + a*c*d) - 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(2, 2*(d*x + c)*f/d)/((d*x + c)*a*d)

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 320 vs. 2(159) = 318.

Time = 0.30 (sec) , antiderivative size = 320, normalized size of antiderivative = 2.01

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))} dx = \frac{\left(2(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right)f^2 \operatorname{Ei}\left(-\frac{2((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf)}{d}\right) e^{-\frac{2(de-cf)}{d}} - 2def^2 \operatorname{Ei}\left(-\frac{2((dx+c)}{d}\right)\right)}{2((dx+c)ad)}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e)),x, algorithm="giac")

```
[Out] -1/2*(2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) - 2*d*e*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) + 2*c*f^3*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) + d*f^2*e^(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) + d*f^2*d^2/(((d*x + c)*a*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a*d^5*e + a*c*d^4*f)*f)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + a \tanh(e + fx))} dx = \int \frac{1}{(a + a \tanh(e + fx)) (c + dx)^2} dx$$

```
[In] int(1/((a + a*tanh(e + f*x))*(c + d*x)^2), x)
```

```
[Out] int(1/((a + a*tanh(e + f*x))*(c + d*x)^2), x)
```

$$3.37 \quad \int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx$$

Optimal result	260
Rubi [A] (verified)	261
Mathematica [A] (verified)	263
Maple [A] (verified)	263
Fricas [A] (verification not implemented)	263
Sympy [F]	264
Maxima [A] (verification not implemented)	264
Giac [A] (verification not implemented)	265
Mupad [F(-1)]	265

Optimal result

Integrand size = 20, antiderivative size = 211

$$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx = -\frac{f}{2ad^2(c+dx)} + \frac{f^2 \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{ad^3}$$

$$- \frac{f^2 \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{ad^3}$$

$$- \frac{f^2 \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{ad^3}$$

$$+ \frac{f^2 \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{ad^3}$$

$$- \frac{1}{2d(c+dx)^2(a+a \tanh(e+fx))}$$

$$+ \frac{f}{d^2(c+dx)(a+a \tanh(e+fx))}$$

```
[Out] -1/2*f/a/d^2/(d*x+c)+f^2*Chi(2*c*f/d+2*f*x)*cosh(-2*e+2*c*f/d)/a/d^3-f^2*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d+2*f*x)/a/d^3+f^2*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^3-f^2*Shi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^3-1/2/d/(d*x+c)^2/(a+a*tanh(f*x+e))+f/d^2/(d*x+c)/(a+a*tanh(f*x+e))
```

Rubi [A] (verified)

Time = 0.23 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3806, 3805, 3384, 3379, 3382}

$$\int \frac{1}{(c + dx)^3(a + a \tanh(e + fx))} dx = -\frac{f^2 \text{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{ad^3} + \frac{f^2 \text{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{ad^3} + \frac{f^2 \sinh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{ad^3} - \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{ad^3} + \frac{f}{d^2(c + dx)(a \tanh(e + fx) + a)} - \frac{f}{2ad^2(c + dx)} - \frac{1}{2d(c + dx)^2(a \tanh(e + fx) + a)}$$

[In] Int[1/((c + d*x)^3*(a + a*Tanh[e + f*x])),x]

[Out] -1/2*f/(a*d^2*(c + d*x)) + (f^2*Cosh[2*e - (2*c*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x])/(a*d^3) - (f^2*CoshIntegral[(2*c*f)/d + 2*f*x]*Sinh[2*e - (2*c*f)/d])/(a*d^3) - (f^2*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(a*d^3) + (f^2*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(a*d^3) - 1/(2*d*(c + d*x)^2*(a + a*Tanh[e + f*x])) + f/(d^2*(c + d*x)*(a + a*Tanh[e + f*x]))

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3805

```
Int[1/(((c_.) + (d_.)*(x_))2*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])), x_Symbol]
:= -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))(-1), x] + (-Dist[f/(a*d), Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Dist[f/(b*d), Int[Cos[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a2 + b2, 0]
```

Rule 3806

```
Int[((c_.) + (d_.)*(x_))(m_)/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Symbol]
:= Simp[f*((c + d*x)(m + 2)/(b*d2*(m + 1)*(m + 2))], x] + (Dist[2*b*(f/(a*d*(m + 1))), Int[(c + d*x)(m + 1)/(a + b*Tan[e + f*x]), x], x] + Simp[(c + d*x)(m + 1)/(d*(m + 1)*(a + b*Tan[e + f*x])), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a2 + b2, 0] && LtQ[m, -1] && NeQ[m, -2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\tanh(e+fx))} - \frac{f \int \frac{1}{(c+dx)^2(a+a\tanh(e+fx))} dx}{d} \\
&= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\tanh(e+fx))} \\
&\quad + \frac{f}{d^2(c+dx)(a+a\tanh(e+fx))} + \frac{f^2 \int \frac{\cosh(2e+2fx)}{c+dx} dx}{ad^2} - \frac{f^2 \int \frac{\sinh(2e+2fx)}{c+dx} dx}{ad^2} \\
&= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\tanh(e+fx))} + \frac{f}{d^2(c+dx)(a+a\tanh(e+fx))} \\
&\quad + \frac{(f^2 \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} - \frac{(f^2 \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&\quad - \frac{(f^2 \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} + \frac{(f^2 \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&= -\frac{f}{2ad^2(c+dx)} + \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{ad^3} \\
&\quad - \frac{f^2 \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{ad^3} \\
&\quad - \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^3} + \frac{f^2 \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^3} \\
&\quad - \frac{1}{2d(c+dx)^2(a+a\tanh(e+fx))} + \frac{f}{d^2(c+dx)(a+a\tanh(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.86 (sec) , antiderivative size = 264, normalized size of antiderivative = 1.25

$$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx = \frac{\operatorname{sech}(e+fx) \left(\cosh\left(\frac{cf}{d}\right) + \sinh\left(\frac{cf}{d}\right) \right) \left(d(d \cosh(e+f(-\frac{c}{d}+x))) + (d-2cf-2dfx) \cosh(e+f(\frac{c}{d}+x)) \right)}{\dots}$$

[In] Integrate[1/((c + d*x)^3*(a + a*Tanh[e + f*x])),x]

[Out] $-1/4*(\operatorname{Sech}[e + f*x]*(\operatorname{Cosh}[(c*f)/d] + \operatorname{Sinh}[(c*f)/d])*(d*(d*\operatorname{Cosh}[e + f*(-(c/d) + x)] + (d - 2*c*f - 2*d*f*x)*\operatorname{Cosh}[e + f*(c/d + x)] + d*\operatorname{Sinh}[e + f*(-(c/d) + x)] - d*\operatorname{Sinh}[e + f*(c/d + x)] + 2*c*f*\operatorname{Sinh}[e + f*(c/d + x)] + 2*d*f*x*\operatorname{Sinh}[e + f*(c/d + x)]) + 4*f^2*(c + d*x)^2*\operatorname{CoshIntegral}[(2*f*(c + d*x))/d]*(\operatorname{Cosh}[e - (f*(c + d*x))/d] + \operatorname{Sinh}[e - (f*(c + d*x))/d]) + 4*f^2*(c + d*x)^2*(\operatorname{Cosh}[e - (f*(c + d*x))/d] - \operatorname{Sinh}[e - (f*(c + d*x))/d])*\operatorname{SinhIntegral}[(2*f*(c + d*x))/d]))/(a*d^3*(c + d*x)^2*(1 + \operatorname{Tanh}[e + f*x]))$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00

method	result
risch	$-\frac{1}{4da(dx+c)^2} + \frac{f^3 e^{-2fx-2e} x}{2ad(d^2 x^2 f^2 + 2cd f^2 x + c^2 f^2)} + \frac{f^3 e^{-2fx-2e} c}{2a d^2 (d^2 x^2 f^2 + 2cd f^2 x + c^2 f^2)} - \frac{f^2 e^{-2fx-2e}}{4ad(d^2 x^2 f^2 + 2cd f^2 x + c^2 f^2)} - \frac{f^2 e^{\frac{2cf-2c}{d}}}{\dots}$

[In] int(1/(d*x+c)^3/(a+a*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] $-1/4/d/a/(d*x+c)^2 + 1/2*f^3/a*\exp(-2*f*x-2*e)/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*x + 1/2*f^3/a*\exp(-2*f*x-2*e)/d^2/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*c - 1/4*f^2/a*\exp(-2*f*x-2*e)/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2) - f^2/a/d^3*\exp(2*(c*f-d*e)/d)*\operatorname{Ei}(1,2*f*x+2*e+2*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.62

$$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx = \frac{2(d^2 f^2 x^2 + 2cdf^2 x + c^2 f^2) \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) \cosh(fx+e) \sinh\left(-\frac{2(de-cf)}{d}\right) + (d^2 fx + cdf + 2(d^2 f^2 x^2 + \dots)}{\dots}$$

[In] integrate(1/(d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="fricas")

[Out] $\frac{1}{2} * (2 * (d^2 * f^2 * x^2 + 2 * c * d * f^2 * x + c^2 * f^2) * \text{Ei}(-2 * (d * f * x + c * f) / d) * \cosh(f * x + e) * \sinh(-2 * (d * e - c * f) / d) + (d^2 * f * x + c * d * f + 2 * (d^2 * f^2 * x^2 + 2 * c * d * f^2 * x + c^2 * f^2) * \text{Ei}(-2 * (d * f * x + c * f) / d) * \cosh(-2 * (d * e - c * f) / d) - d^2) * \cosh(f * x + e) - (d^2 * f * x + c * d * f - 2 * (d^2 * f^2 * x^2 + 2 * c * d * f^2 * x + c^2 * f^2) * \text{Ei}(-2 * (d * f * x + c * f) / d) * \cosh(-2 * (d * e - c * f) / d) - 2 * (d^2 * f^2 * x^2 + 2 * c * d * f^2 * x + c^2 * f^2) * \text{Ei}(-2 * (d * f * x + c * f) / d) * \sinh(-2 * (d * e - c * f) / d)) * \sinh(f * x + e)) / ((a * d^5 * x^2 + 2 * a * c * d^4 * x + a * c^2 * d^3) * \cosh(f * x + e) + (a * d^5 * x^2 + 2 * a * c * d^4 * x + a * c^2 * d^3) * \sinh(f * x + e))$

Sympy [F]

$$\int \frac{1}{(c + dx)^3 (a + a \tanh(e + fx))} dx$$

$$= \frac{\int \frac{1}{c^3 \tanh(e + fx) + c^3 + 3c^2 dx \tanh(e + fx) + 3c^2 dx + 3cd^2 x^2 \tanh(e + fx) + 3cd^2 x^2 + d^3 x^3 \tanh(e + fx) + d^3 x^3} dx}{a}$$

[In] integrate(1/(d*x+c)**3/(a+a*tanh(f*x+e)),x)

[Out] Integral(1/(c**3*tanh(e + f*x) + c**3 + 3*c**2*d*x*tanh(e + f*x) + 3*c**2*d*x + 3*c*d**2*x**2*tanh(e + f*x) + 3*c*d**2*x**2 + d**3*x**3*tanh(e + f*x) + d**3*x**3), x)/a

Maxima [A] (verification not implemented)

none

Time = 0.65 (sec) , antiderivative size = 68, normalized size of antiderivative = 0.32

$$\int \frac{1}{(c + dx)^3 (a + a \tanh(e + fx))} dx$$

$$= -\frac{1}{4(ad^3x^2 + 2acd^2x + ac^2d)} - \frac{e^{(-2e + \frac{2cf}{d})} E_3\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)^2 ad}$$

[In] integrate(1/(d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] -1/4/(a*d^3*x^2 + 2*a*c*d^2*x + a*c^2*d) - 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(3, 2*(d*x + c)*f/d)/((d*x + c)^2*a*d)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 176, normalized size of antiderivative = 0.83

$$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx$$

$$= \frac{4d^2 f^2 x^2 \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) e^{\left(\frac{2cf}{d}\right)} + 8cdf^2 x \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) e^{\left(\frac{2cf}{d}\right)} + 4c^2 f^2 \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) e^{\left(\frac{2cf}{d}\right)} + 2d^2 f x e^{\left(\frac{2cf}{d}\right)}}{4(ad^5 x^2 e^{(2e)} + 2acd^4 x e^{(2e)} + ac^2 d^3 e^{(2e)})}$$

[In] integrate(1/(d*x+c)^3/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] 1/4*(4*d^2*f^2*x^2*Ei(-2*(d*f*x + c*f)/d)*e^(2*c*f/d) + 8*c*d*f^2*x*Ei(-2*(d*f*x + c*f)/d)*e^(2*c*f/d) + 4*c^2*f^2*Ei(-2*(d*f*x + c*f)/d)*e^(2*c*f/d) + 2*d^2*f*x*e^(-2*f*x) + 2*c*d*f*e^(-2*f*x) - d^2*e^(-2*f*x) - d^2*e^(2*e)) / (a*d^5*x^2*e^(2*e) + 2*a*c*d^4*x*e^(2*e) + a*c^2*d^3*e^(2*e))

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)^3(a+a \tanh(e+fx))} dx = \int \frac{1}{(a+a \tanh(e+fx))(c+dx)^3} dx$$

[In] int(1/((a + a*tanh(e + f*x))*(c + d*x)^3),x)

[Out] int(1/((a + a*tanh(e + f*x))*(c + d*x)^3), x)

3.38 $\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^2} dx$

Optimal result	266
Rubi [A] (verified)	266
Mathematica [A] (verified)	268
Maple [A] (verified)	269
Fricas [B] (verification not implemented)	269
Sympy [F]	270
Maxima [A] (verification not implemented)	270
Giac [A] (verification not implemented)	271
Mupad [B] (verification not implemented)	271

Optimal result

Integrand size = 20, antiderivative size = 230

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^2} dx = -\frac{3d^3 e^{-4e-4fx}}{512a^2 f^4} - \frac{3d^3 e^{-2e-2fx}}{16a^2 f^4} - \frac{3d^2 e^{-4e-4fx}(c+dx)}{128a^2 f^3} - \frac{3d^2 e^{-2e-2fx}(c+dx)}{8a^2 f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2 f^2} - \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2 f^2} - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2 f} - \frac{e^{-2e-2fx}(c+dx)^3}{4a^2 f} + \frac{(c+dx)^4}{16a^2 d}$$

[Out] $-3/512*d^3*\exp(-4*f*x-4*e)/a^2/f^4-3/16*d^3*\exp(-2*f*x-2*e)/a^2/f^4-3/128*d^2*\exp(-4*f*x-4*e)*(d*x+c)/a^2/f^3-3/8*d^2*\exp(-2*f*x-2*e)*(d*x+c)/a^2/f^3-3/64*d*\exp(-4*f*x-4*e)*(d*x+c)^2/a^2/f^2-3/8*d*\exp(-2*f*x-2*e)*(d*x+c)^2/a^2/f^2-1/16*\exp(-4*f*x-4*e)*(d*x+c)^3/a^2/f-1/4*\exp(-2*f*x-2*e)*(d*x+c)^3/a^2/f+1/16*(d*x+c)^4/a^2/d$

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used

= {3810, 2207, 2225}

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^2} dx = -\frac{3d^2(c+dx)e^{-4e-4fx}}{128a^2f^3} - \frac{3d^2(c+dx)e^{-2e-2fx}}{8a^2f^3} - \frac{3d(c+dx)^2e^{-4e-4fx}}{64a^2f^2} - \frac{3d(c+dx)^2e^{-2e-2fx}}{8a^2f^2} - \frac{(c+dx)^3e^{-4e-4fx}}{16a^2f} - \frac{(c+dx)^3e^{-2e-2fx}}{4a^2f} + \frac{(c+dx)^4}{16a^2d} - \frac{3d^3e^{-4e-4fx}}{512a^2f^4} - \frac{3d^3e^{-2e-2fx}}{16a^2f^4}$$

[In] Int[(c + d*x)^3/(a + a*Tanh[e + f*x])^2,x]

[Out] (-3*d^3*E^(-4*e - 4*f*x))/(512*a^2*f^4) - (3*d^3*E^(-2*e - 2*f*x))/(16*a^2*f^4) - (3*d^2*E^(-4*e - 4*f*x)*(c + d*x))/(128*a^2*f^3) - (3*d^2*E^(-2*e - 2*f*x)*(c + d*x))/(8*a^2*f^3) - (3*d*E^(-4*e - 4*f*x)*(c + d*x)^2)/(64*a^2*f^2) - (3*d*E^(-2*e - 2*f*x)*(c + d*x)^2)/(8*a^2*f^2) - (E^(-4*e - 4*f*x)*(c + d*x)^3)/(16*a^2*f) - (E^(-2*e - 2*f*x)*(c + d*x)^3)/(4*a^2*f) + (c + d*x)^4/(16*a^2*d)

Rule 2207

Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[\$UseGamma]

Rule 2225

Int[((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 3810

Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a)]^(n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^3}{4a^2} + \frac{e^{-4e-4fx}(c+dx)^3}{4a^2} + \frac{e^{-2e-2fx}(c+dx)^3}{2a^2} \right) dx \\ &= \frac{(c+dx)^4}{16a^2d} + \frac{\int e^{-4e-4fx}(c+dx)^3 dx}{4a^2} + \frac{\int e^{-2e-2fx}(c+dx)^3 dx}{2a^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} - \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d} \\
&\quad + \frac{(3d) \int e^{-4e-4fx}(c+dx)^2 dx}{16a^2f} + \frac{(3d) \int e^{-2e-2fx}(c+dx)^2 dx}{4a^2f} \\
&= -\frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} - \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} \\
&\quad - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} - \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d} \\
&\quad + \frac{(3d^2) \int e^{-4e-4fx}(c+dx) dx}{32a^2f^2} + \frac{(3d^2) \int e^{-2e-2fx}(c+dx) dx}{4a^2f^2} \\
&= -\frac{3d^2e^{-4e-4fx}(c+dx)}{128a^2f^3} - \frac{3d^2e^{-2e-2fx}(c+dx)}{8a^2f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} \\
&\quad - \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} - \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} \\
&\quad + \frac{(c+dx)^4}{16a^2d} + \frac{(3d^3) \int e^{-4e-4fx} dx}{128a^2f^3} + \frac{(3d^3) \int e^{-2e-2fx} dx}{8a^2f^3} \\
&= -\frac{3d^3e^{-4e-4fx}}{512a^2f^4} - \frac{3d^3e^{-2e-2fx}}{16a^2f^4} - \frac{3d^2e^{-4e-4fx}(c+dx)}{128a^2f^3} \\
&\quad - \frac{3d^2e^{-2e-2fx}(c+dx)}{8a^2f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} - \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} \\
&\quad - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} - \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.02 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.83

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^2} dx$$

$$= \frac{\operatorname{sech}^2(e+fx)(\cosh(fx)+\sinh(fx))^2(-((4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2))+d^3(3+6fx+6f^2x^2+4f^3x^3))\operatorname{Cosh}[2fx])+(32c^3f^3+24c^2d^2f^2(1+4fx)+12cd^2f^2(1+4fx+8f^2x^2)+d^3(3+12fx+24f^2x^2+32f^3x^3))\operatorname{Cosh}[4fx]*(-\operatorname{Cosh}[2e]+\operatorname{Sinh}[2e]))/32+f^4x*(4c^3+6c^2dx+4cd^2x^2+d^3x^3)*(\operatorname{Cosh}[2e]+\operatorname{Sinh}[2e])+(4c^3f^3+6c^2d^2f^2(1+2fx)+6cd^2f^2(1+2fx+2f^2x^2)+d^3(3+6fx+6f^2x^2+4f^3x^3))\operatorname{Sinh}[2fx]+((32c^3f^3+24c^2d^2f^2(1+4fx)+12cd^2f^2(1+4fx+8f^2x^2)+d^3(3+12fx+24f^2x^2+32f^3x^3))\operatorname{Cosh}[2e]-\operatorname{Sinh}[2e])\operatorname{Sinh}[4fx])/32)/(16a^2f^4(1+\operatorname{Tanh}[e+fx])^2)}$$

[In] Integrate[(c + d*x)^3/(a + a*Tanh[e + f*x])^2,x]

[Out] (Sech[e + f*x]^2*(Cosh[f*x] + Sinh[f*x])^2*(-((4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*Cosh[2*f*x]) + ((32*c^3*f^3 + 24*c^2*d*f^2*(1 + 4*f*x) + 12*c*d^2*f^2*(1 + 4*f*x + 8*f^2*x^2) + d^3*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3))*Cosh[4*f*x]*(-Cosh[2*e] + Sinh[2*e]))/32 + f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cosh[2*e] + Sinh[2*e]) + (4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*Sinh[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(1 + 4*f*x) + 12*c*d^2*f*(1 + 4*f*x + 8*f^2*x^2) + d^3*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3))*(Cosh[2*e] - Sinh[2*e])*Sinh[4*f*x])/32)/(16*a^2*f^4*(1 + Tanh[e + f*x])^2)

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.19

method	result
risch	$\frac{d^3x^4}{16a^2} + \frac{d^2cx^3}{4a^2} + \frac{3dc^2x^2}{8a^2} + \frac{c^3x}{4a^2} + \frac{c^4}{16a^2d} - \frac{(4d^3x^3f^3+12cd^2f^3x^2+12c^2df^3x+6d^3f^2x^2+4c^3f^3+12cd^2f^2x+6c^2df^2}{16a^2f^4}$
parallelrisch	$\frac{-48d^3-120c^2df^3x-45\tanh(fx+e)d^3+24x\tanh(fx+e)cd^2f^2+64x^3\tanh(fx+e)cd^2f^4+96x^2\tanh(fx+e)c^2df^4+48x^2\tanh(fx+e)d^3f^3}{16a^2f^4}$

[In] int((d*x+c)^3/(a+a*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] $1/16/a^2*d^3*x^4+1/4/a^2*d^2*c*x^3+3/8/a^2*d*c^2*x^2+1/4/a^2*c^3*x+1/16/a^2/d*c^4-1/16*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4*c^3*f^3+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+3*d^3)/a^2/f^4*\exp(-2*f*x-2*e)-1/512*(32*d^3*f^3*x^3+96*c*d^2*f^3*x^2+96*c^2*d*f^3*x+24*d^3*f^2*x^2+32*c^3*f^3+48*c*d^2*f^2*x+24*c^2*d*f^2+12*d^3*f*x+12*c*d^2*f+3*d^3)/a^2/f^4*\exp(-4*f*x-4*e)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 573 vs. 2(204) = 408.

Time = 0.26 (sec) , antiderivative size = 573, normalized size of antiderivative = 2.49

$$\int \frac{(c+dx)^3}{(a+a\tanh(e+fx))^2} dx = \frac{128d^3f^3x^3 + 128c^3f^3 + 192c^2df^2 + 192cd^2f + 96d^3 + 192(2cd^2f^3 + d^3f^2)x^2 - (32d^3f^4x^4 - 32c^3f^3}{16a^2f^4}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] $-1/512*(128*d^3*f^3*x^3 + 128*c^3*f^3 + 192*c^2*d*f^2 + 192*c*d^2*f + 96*d^3 + 192*(2*c*d^2*f^3 + d^3*f^2)*x^2 - (32*d^3*f^4*x^4 - 32*c^3*f^3 - 24*c^2*d*f^2 - 12*c*d^2*f + 32*(4*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 24*(8*c^2*d*f^4 - 4*c*d^2*f^3 - d^3*f^2)*x^2 + 4*(32*c^3*f^4 - 24*c^2*d*f^3 - 12*c*d^2*f^2 - 3*d^3*f)*x)*\cosh(f*x + e)^2 - 2*(32*d^3*f^4*x^4 + 32*c^3*f^3 + 24*c^2*d*f^2 + 12*c*d^2*f + 32*(4*c*d^2*f^4 + d^3*f^3)*x^3 + 3*d^3 + 24*(8*c^2*d*f^4 + 4*c*d^2*f^3 + d^3*f^2)*x^2 + 4*(32*c^3*f^4 + 24*c^2*d*f^3 + 12*c*d^2*f^2 + 3*d^3*f)*x)*\cosh(f*x + e)*\sinh(f*x + e) - (32*d^3*f^4*x^4 - 32*c^3*f^3 - 24*c^2*d*f^2 - 12*c*d^2*f + 32*(4*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 24*(8*c^2*d*f^4 - 4*c*d^2*f^3 - d^3*f^2)*x^2 + 4*(32*c^3*f^4 - 24*c^2*d*f^3 - 12*c*d^2*f^2 - 3*d^3*f)*x)*\sinh(f*x + e)^2 + 192*(2*c^2*d*f^3 + 2*c*d^2*f^2 + d^3*f)*x)/(a^2*f^4*\cosh(f*x + e)^2 + 2*a^2*f^4*\cosh(f*x + e)*\sinh(f*x + e) + a^2*f^4*\sinh(f*x + e)^2)$

SymPy [F]

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^2} dx = \frac{\int \frac{c^3}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx + \int \frac{d^3 x^3}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx + \int \frac{3cd^2 x^2}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx + \int \frac{3d^2 x^2}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx}{a^2}$$

```
[In] integrate((d*x+c)**3/(a+a*tanh(f*x+e))**2,x)
```

```
[Out] (Integral(c**3/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x) + Integral(d**3*x**3/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x) + Integral(3*c*d**2*x**2/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x) + Integral(3*c**2*d*x/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x))/a**2
```

Maxima [A] (verification not implemented)

none

Time = 0.51 (sec) , antiderivative size = 296, normalized size of antiderivative = 1.29

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^2} dx = \frac{1}{16} c^3 \left(\frac{4(fx + e)}{a^2 f} - \frac{4e^{(-2fx-2e)} + e^{(-4fx-4e)}}{a^2 f} \right) + \frac{3(8f^2x^2e^{(4e)} - 8(2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - (4fx + 1)e^{(-4fx)})c^2de^{(-4e)}}{64a^2f^2} + \frac{(32f^3x^3e^{(4e)} - 48(2f^2x^2e^{(2e)} + 2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - 3(8f^2x^2 + 4fx + 1)e^{(-4fx)})cd^2e^{(-4e)}}{128a^2f^3} + \frac{(32f^4x^4e^{(4e)} - 32(4f^3x^3e^{(2e)} + 6f^2x^2e^{(2e)} + 6fxe^{(2e)} + 3e^{(2e)})e^{(-2fx)} - (32f^3x^3 + 24f^2x^2 + 12fx + 3)e^{(-4fx)})d^3e^{(-4e)}}{512a^2f^4}$$

```
[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")
```

```
[Out] 1/16*c^3*(4*(f*x + e)/(a^2*f) - (4*e^(-2*f*x - 2*e) + e^(-4*f*x - 4*e))/(a^2*f)) + 3/64*(8*f^2*x^2*e^(4*e) - 8*(2*f*x*e^(2*e) + e^(2*e))*e^(-2*f*x) - (4*f*x + 1)*e^(-4*f*x))*c^2*d*e^(-4*e)/(a^2*f^2) + 1/128*(32*f^3*x^3*e^(4*e) - 48*(2*f^2*x^2*e^(2*e) + 2*f*x*e^(2*e) + e^(2*e))*e^(-2*f*x) - 3*(8*f^2*x^2 + 4*f*x + 1)*e^(-4*f*x))*c*d^2*e^(-4*e)/(a^2*f^3) + 1/512*(32*f^4*x^4*e^(4*e) - 32*(4*f^3*x^3*e^(2*e) + 6*f^2*x^2*e^(2*e) + 6*f*x*e^(2*e) + 3*e^(2*e))*e^(-2*f*x) - (32*f^3*x^3 + 24*f^2*x^2 + 12*f*x + 3)*e^(-4*f*x))*d^3*e^(-4*e)/(a^2*f^4)
```

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 368, normalized size of antiderivative = 1.60

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^2} dx$$

$$= \frac{(32 d^3 f^4 x^4 e^{(4fx+4e)} + 128 cd^2 f^4 x^3 e^{(4fx+4e)} + 192 c^2 d f^4 x^2 e^{(4fx+4e)} - 128 d^3 f^3 x^3 e^{(2fx+2e)} - 32 d^3 f^3 x^3 + \dots}{\dots}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

```
[Out] 1/512*(32*d^3*f^4*x^4*e^(4*f*x + 4*e) + 128*c*d^2*f^4*x^3*e^(4*f*x + 4*e) +
192*c^2*d*f^4*x^2*e^(4*f*x + 4*e) - 128*d^3*f^3*x^3*e^(2*f*x + 2*e) - 32*d
^3*f^3*x^3 + 128*c^3*f^4*x*e^(4*f*x + 4*e) - 384*c*d^2*f^3*x^2*e^(2*f*x + 2
*e) - 96*c*d^2*f^3*x^2 - 384*c^2*d*f^3*x*e^(2*f*x + 2*e) - 192*d^3*f^2*x^2*
e^(2*f*x + 2*e) - 96*c^2*d*f^3*x - 24*d^3*f^2*x^2 - 128*c^3*f^3*e^(2*f*x +
2*e) - 384*c*d^2*f^2*x*e^(2*f*x + 2*e) - 32*c^3*f^3 - 48*c*d^2*f^2*x - 192*
c^2*d*f^2*e^(2*f*x + 2*e) - 192*d^3*f*x*e^(2*f*x + 2*e) - 24*c^2*d*f^2 - 12
*d^3*f*x - 192*c*d^2*f*e^(2*f*x + 2*e) - 12*c*d^2*f - 96*d^3*e^(2*f*x + 2*e
) - 3*d^3)*e^(-4*f*x - 4*e)/(a^2*f^4)
```

Mupad [B] (verification not implemented)

Time = 1.85 (sec) , antiderivative size = 267, normalized size of antiderivative = 1.16

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^2} dx = \frac{c^3 x}{4 a^2} - e^{-4e-4fx} \left(\frac{32 c^3 f^3 + 24 c^2 d f^2 + 12 c d^2 f + 3 d^3}{512 a^2 f^4} \right.$$

$$+ \frac{d^3 x^3}{16 a^2 f} + \frac{3 dx (8 c^2 f^2 + 4 c d f + d^2)}{128 a^2 f^3} + \frac{3 d^2 x^2 (d + 4 c f)}{64 a^2 f^2} \left. \right)$$

$$- e^{-2e-2fx} \left(\frac{4 c^3 f^3 + 6 c^2 d f^2 + 6 c d^2 f + 3 d^3}{16 a^2 f^4} + \frac{d^3 x^3}{4 a^2 f} \right.$$

$$+ \frac{3 dx (2 c^2 f^2 + 2 c d f + d^2)}{8 a^2 f^3} + \frac{3 d^2 x^2 (d + 2 c f)}{8 a^2 f^2} \left. \right)$$

$$+ \frac{d^3 x^4}{16 a^2} + \frac{3 c^2 d x^2}{8 a^2} + \frac{c d^2 x^3}{4 a^2}$$

[In] int((c + d*x)^3/(a + a*tanh(e + f*x))^2,x)

```
[Out] (c^3*x)/(4*a^2) - exp(- 4*e - 4*f*x)*((3*d^3 + 32*c^3*f^3 + 24*c^2*d*f^2 +
12*c*d^2*f)/(512*a^2*f^4) + (d^3*x^3)/(16*a^2*f) + (3*d*x*(d^2 + 8*c^2*f^2
+ 4*c*d*f))/(128*a^2*f^3) + (3*d^2*x^2*(d + 4*c*f))/(64*a^2*f^2)) - exp(- 2
*e - 2*f*x)*((3*d^3 + 4*c^3*f^3 + 6*c^2*d*f^2 + 6*c*d^2*f)/(16*a^2*f^4) + (
d^3*x^3)/(4*a^2*f) + (3*d*x*(d^2 + 2*c^2*f^2 + 2*c*d*f))/(8*a^2*f^3) + (3*d
^2*x^2*(d + 2*c*f))/(8*a^2*f^2)) + (d^3*x^4)/(16*a^2) + (3*c^2*d*x^2)/(8*a^
2) + (c*d^2*x^3)/(4*a^2)
```

3.39 $\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^2} dx$

Optimal result	272
Rubi [A] (verified)	272
Mathematica [A] (verified)	274
Maple [A] (verified)	274
Fricas [B] (verification not implemented)	274
Sympy [F]	275
Maxima [A] (verification not implemented)	275
Giac [A] (verification not implemented)	276
Mupad [B] (verification not implemented)	276

Optimal result

Integrand size = 20, antiderivative size = 170

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^2} dx = -\frac{d^2 e^{-4e-4fx}}{128a^2 f^3} - \frac{d^2 e^{-2e-2fx}}{8a^2 f^3} - \frac{de^{-4e-4fx}(c+dx)}{32a^2 f^2} - \frac{de^{-2e-2fx}(c+dx)}{4a^2 f^2} - \frac{e^{-4e-4fx}(c+dx)^2}{16a^2 f} - \frac{e^{-2e-2fx}(c+dx)^2}{4a^2 f} + \frac{(c+dx)^3}{12a^2 d}$$

[Out] $-1/128*d^2*\exp(-4*f*x-4*e)/a^2/f^3-1/8*d^2*\exp(-2*f*x-2*e)/a^2/f^3-1/32*d*\exp(-4*f*x-4*e)*(d*x+c)/a^2/f^2-1/4*d*\exp(-2*f*x-2*e)*(d*x+c)/a^2/f^2-1/16*\exp(-4*f*x-4*e)*(d*x+c)^2/a^2/f-1/4*\exp(-2*f*x-2*e)*(d*x+c)^2/a^2/f+1/12*(d*x+c)^3/a^2/d$

Rubi [A] (verified)

Time = 0.13 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3810, 2207, 2225}

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^2} dx = -\frac{d(c+dx)e^{-4e-4fx}}{32a^2 f^2} - \frac{d(c+dx)e^{-2e-2fx}}{4a^2 f^2} - \frac{(c+dx)^2 e^{-4e-4fx}}{16a^2 f} - \frac{(c+dx)^2 e^{-2e-2fx}}{4a^2 f} + \frac{(c+dx)^3}{12a^2 d} - \frac{d^2 e^{-4e-4fx}}{128a^2 f^3} - \frac{d^2 e^{-2e-2fx}}{8a^2 f^3}$$

[In] $\text{Int}[(c + d*x)^2/(a + a*\text{Tanh}[e + f*x])^2, x]$

[Out] $-1/128*(d^2*E^{-4*e - 4*f*x})/(a^2*f^3) - (d^2*E^{-2*e - 2*f*x})/(8*a^2*f^3) - (d*E^{-4*e - 4*f*x})*(c + d*x)/(32*a^2*f^2) - (d*E^{-2*e - 2*f*x})*(c +$

$d*x))/(4*a^2*f^2) - (E^{(-4*e - 4*f*x)*(c + d*x)^2}/(16*a^2*f) - (E^{(-2*e - 2*f*x)*(c + d*x)^2}/(4*a^2*f) + (c + d*x)^3/(12*a^2*d)$

Rule 2207

$\text{Int}[(b_*)*(F_*)^{((g_*)*((e_*) + (f_*)*(x_)))})^{(n_*)*((c_*) + (d_*)*(x_))}^{(m_*)}, x_Symbol] \text{ :> } \text{Simp}[(c + d*x)^m*((b*F^{(g*(e + f*x)))^n/(f*g*n*\text{Log}[F])), x] - \text{Dist}[d*(m/(f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{(m-1)}*(b*F^{(g*(e + f*x)))^n}, x], x] /; \text{FreeQ}[\{F, b, c, d, e, f, g, n\}, x] \&\& \text{GtQ}[m, 0] \&\& \text{IntegerQ}[2*m] \&\& !\text{TrueQ}[\$UseGamma]$

Rule 2225

$\text{Int}[(F_*)^{((c_*)*((a_*) + (b_*)*(x_)))})^{(n_*)}, x_Symbol] \text{ :> } \text{Simp}[(F^{(c*(a + b*x)))^n/(b*c*n*\text{Log}[F]), x] /; \text{FreeQ}[\{F, a, b, c, n\}, x]$

Rule 3810

$\text{Int}[(c_*) + (d_*)*(x_)]^{(m_*)*((a_*) + (b_*)*\text{tan}[(e_*) + (f_*)*(x_)])}^{(n_*)}, x_Symbol] \text{ :> } \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + E^{(2*(a/b)*(e + f*x)})/(2*a))^{(-n)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&\& \text{EqQ}[a^2 + b^2, 0] \&\& \text{ILtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{(c + dx)^2}{4a^2} + \frac{e^{-4e-4fx}(c + dx)^2}{4a^2} + \frac{e^{-2e-2fx}(c + dx)^2}{2a^2} \right) dx \\
 &= \frac{(c + dx)^3}{12a^2d} + \frac{\int e^{-4e-4fx}(c + dx)^2 dx}{4a^2} + \frac{\int e^{-2e-2fx}(c + dx)^2 dx}{2a^2} \\
 &= -\frac{e^{-4e-4fx}(c + dx)^2}{16a^2f} - \frac{e^{-2e-2fx}(c + dx)^2}{4a^2f} + \frac{(c + dx)^3}{12a^2d} \\
 &\quad + \frac{d \int e^{-4e-4fx}(c + dx) dx}{8a^2f} + \frac{d \int e^{-2e-2fx}(c + dx) dx}{2a^2f} \\
 &= -\frac{de^{-4e-4fx}(c + dx)}{32a^2f^2} - \frac{de^{-2e-2fx}(c + dx)}{4a^2f^2} - \frac{e^{-4e-4fx}(c + dx)^2}{16a^2f} \\
 &\quad - \frac{e^{-2e-2fx}(c + dx)^2}{4a^2f} + \frac{(c + dx)^3}{12a^2d} + \frac{d^2 \int e^{-4e-4fx} dx}{32a^2f^2} + \frac{d^2 \int e^{-2e-2fx} dx}{4a^2f^2} \\
 &= -\frac{d^2 e^{-4e-4fx}}{128a^2f^3} - \frac{d^2 e^{-2e-2fx}}{8a^2f^3} - \frac{de^{-4e-4fx}(c + dx)}{32a^2f^2} - \frac{de^{-2e-2fx}(c + dx)}{4a^2f^2} \\
 &\quad - \frac{e^{-4e-4fx}(c + dx)^2}{16a^2f} - \frac{e^{-2e-2fx}(c + dx)^2}{4a^2f} + \frac{(c + dx)^3}{12a^2d}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 1.53 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.22

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx = \frac{\operatorname{sech}^2(e + fx) (-48(2c^2 f^2 + 2cdf(1 + 2fx) + d^2(1 + 2fx + 2f^2 x^2)) + (24c^2 f^2(-1 + 4fx) + 12cdf(-1 - 4fx))}{(384a^2 f^3 (1 + \tanh(e + fx))^2)}$$

[In] Integrate[(c + d*x)^2/(a + a*Tanh[e + f*x])^2,x]

[Out] (Sech[e + f*x]^2*(-48*(2*c^2*f^2 + 2*c*d*f*(1 + 2*f*x) + d^2*(1 + 2*f*x + 2*f^2*x^2)) + (24*c^2*f^2*(-1 + 4*f*x) + 12*c*d*f*(-1 - 4*f*x + 8*f^2*x^2) + d^2*(-3 - 12*f*x - 24*f^2*x^2 + 32*f^3*x^3))*Cosh[2*(e + f*x)] + (24*c^2*f^2*(1 + 4*f*x) + 12*c*d*f*(1 + 4*f*x + 8*f^2*x^2) + d^2*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3))*Sinh[2*(e + f*x)])/(384*a^2*f^3*(1 + Tanh[e + f*x])^2)

Maple [A] (verified)

Time = 0.26 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.96

method	result
risch	$\frac{d^2 x^3}{12a^2} + \frac{dcx^2}{4a^2} + \frac{c^2 x}{4a^2} + \frac{c^3}{12a^2 d} - \frac{(2d^2 x^2 f^2 + 4cd f^2 x + 2c^2 f^2 + 2d^2 f x + 2cdf + d^2)e^{-2fx-2e}}{8a^2 f^3} - \frac{(8d^2 x^2 f^2 + 16cd f^2 x + 8c^2 f^2)}{128a^2 f^3}$
parallelrisch	$\frac{-24d^2 - 27d^2 fx - 21 \tanh(fx+e)d^2 - 48c^2 f^2 + 24cd x^2 f^3 - 60cd f^2 x - 30d^2 x^2 f^2 - 48cdf - 24 \tanh(fx+e)c^2 f^2 + 24x c^2 f^3 + 8d^2 x^3 f^3}{(384a^2 f^3 (1 + \tanh(e + fx))^2)}$

[In] int((d*x+c)^2/(a+a*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] 1/12/a^2*d^2*x^3+1/4/a^2*d*c*x^2+1/4/a^2*c^2*x+1/12/a^2/d*c^3-1/8*(2*d^2*f^2*x^2+4*c*d*f^2*x+2*c^2*f^2+2*d^2*f*x+2*c*d*f+d^2)/a^2/f^3*exp(-2*f*x-2*e)-1/128*(8*d^2*f^2*x^2+16*c*d*f^2*x+8*c^2*f^2+4*d^2*f*x+4*c*d*f+d^2)/a^2/f^3*exp(-4*f*x-4*e)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 361 vs. 2(150) = 300.

Time = 0.26 (sec) , antiderivative size = 361, normalized size of antiderivative = 2.12

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx = \frac{96 d^2 f^2 x^2 + 96 c^2 f^2 + 96 cdf - (32 d^2 f^3 x^3 - 24 c^2 f^2 - 12 cdf + 24 (4 cdf^3 - d^2 f^2) x^2 - 3 d^2 + 12 (8 c^2 f^3 - 12 cdf + 24 c^2 f^2)) \operatorname{sech}^2(e + fx)}{(384 a^2 f^3 (1 + \tanh(e + fx))^2)}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")

[Out]
$$-1/384*(96*d^2*f^2*x^2 + 96*c^2*f^2 + 96*c*d*f - (32*d^2*f^3*x^3 - 24*c^2*f^2 - 12*c*d*f + 24*(4*c*d*f^3 - d^2*f^2))*x^2 - 3*d^2 + 12*(8*c^2*f^3 - 4*c*d*f^2 - d^2*f)*x)*\cosh(f*x + e)^2 - 2*(32*d^2*f^3*x^3 + 24*c^2*f^2 + 12*c*d*f + 24*(4*c*d*f^3 + d^2*f^2))*x^2 + 3*d^2 + 12*(8*c^2*f^3 + 4*c*d*f^2 + d^2*f)*x)*\cosh(f*x + e)*\sinh(f*x + e) - (32*d^2*f^3*x^3 - 24*c^2*f^2 - 12*c*d*f + 24*(4*c*d*f^3 - d^2*f^2))*x^2 - 3*d^2 + 12*(8*c^2*f^3 - 4*c*d*f^2 - d^2*f)*x)*\sinh(f*x + e)^2 + 48*d^2 + 96*(2*c*d*f^2 + d^2*f)*x)/(a^2*f^3*\cosh(f*x + e)^2 + 2*a^2*f^3*\cosh(f*x + e)*\sinh(f*x + e) + a^2*f^3*\sinh(f*x + e)^2)$$

Sympy [F]

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx$$

$$= \frac{\int \frac{c^2}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx + \int \frac{d^2 x^2}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx + \int \frac{2cdx}{\tanh^2(e+fx)+2 \tanh(e+fx)+1} dx}{a^2}$$

[In] integrate((d*x+c)**2/(a+a*tanh(f*x+e))**2,x)

[Out] $(\text{Integral}(c**2/(\tanh(e + f*x)**2 + 2*\tanh(e + f*x) + 1), x) + \text{Integral}(d**2*x**2/(\tanh(e + f*x)**2 + 2*\tanh(e + f*x) + 1), x) + \text{Integral}(2*c*d*x/(\tanh(e + f*x)**2 + 2*\tanh(e + f*x) + 1), x))/a**2$

Maxima [A] (verification not implemented)

none

Time = 0.43 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.12

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx = \frac{1}{16} c^2 \left(\frac{4(fx + e)}{a^2 f} - \frac{4e^{(-2fx-2e)} + e^{(-4fx-4e)}}{a^2 f} \right)$$

$$+ \frac{(8f^2x^2e^{(4e)} - 8(2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - (4fx + 1)e^{(-4fx)})cde^{(-4e)}}{32a^2f^2}$$

$$+ \frac{(32f^3x^3e^{(4e)} - 48(2f^2x^2e^{(2e)} + 2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - 3(8f^2x^2 + 4fx + 1)e^{(-4fx)})d^2e^{(-4e)}}{384a^2f^3}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] $1/16*c^2*(4*(f*x + e)/(a^2*f) - (4*e^{(-2*f*x - 2*e)} + e^{(-4*f*x - 4*e)}))/(a^2*f) + 1/32*(8*f^2*x^2*e^{(4*e)} - 8*(2*f*x*e^{(2*e)} + e^{(2*e)})*e^{(-2*f*x)} - (4*f*x + 1)*e^{(-4*f*x)})*c*d*e^{(-4*e)}/(a^2*f^2) + 1/384*(32*f^3*x^3*e^{(4*e)} - 48*(2*f^2*x^2*e^{(2*e)} + 2*f*x*e^{(2*e)} + e^{(2*e)})*e^{(-2*f*x)} - 3*(8*f^2*x^2 + 4*f*x + 1)*e^{(-4*f*x)})*d^2*e^{(-4*e)}/(a^2*f^3)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.28

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx$$

$$= \frac{(32 d^2 f^3 x^3 e^{(4fx+4e)} + 96 cdf^3 x^2 e^{(4fx+4e)} + 96 c^2 f^3 x e^{(4fx+4e)} - 96 d^2 f^2 x^2 e^{(2fx+2e)} - 24 d^2 f^2 x^2 - 192 cdf^2 x^2 - 96 c^2 f^2 x e^{(2fx+2e)} - 24 c^2 f^2 x - 12 d^2 f^2 x - 96 c d f^2 e^{(2fx+2e)} - 12 c d f - 48 d^2 e^{(2fx+2e)} - 3 d^2) e^{(-4fx-4e)}}{(a^2 f^3)}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

[Out] 1/384*(32*d^2*f^3*x^3*e^(4*f*x + 4*e) + 96*c*d*f^3*x^2*e^(4*f*x + 4*e) + 96*c^2*f^3*x*e^(4*f*x + 4*e) - 96*d^2*f^2*x^2*e^(2*f*x + 2*e) - 24*d^2*f^2*x^2 - 192*c*d*f^2*x*e^(2*f*x + 2*e) - 48*c*d*f^2*x - 96*c^2*f^2*e^(2*f*x + 2*e) - 96*d^2*f*x*e^(2*f*x + 2*e) - 24*c^2*f^2 - 12*d^2*f*x - 96*c*d*f*e^(2*f*x + 2*e) - 12*c*d*f - 48*d^2*e^(2*f*x + 2*e) - 3*d^2)*e^(-4*f*x - 4*e)/(a^2*f^3)

Mupad [B] (verification not implemented)

Time = 1.83 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.97

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^2} dx = \frac{c^2 x}{4 a^2} - e^{-4e-4fx} \left(\frac{8 c^2 f^2 + 4 c d f + d^2}{128 a^2 f^3} + \frac{d^2 x^2}{16 a^2 f} + \frac{d x (d + 4 c f)}{32 a^2 f^2} \right) - e^{-2e-2fx} \left(\frac{2 c^2 f^2 + 2 c d f + d^2}{8 a^2 f^3} + \frac{d^2 x^2}{4 a^2 f} + \frac{d x (d + 2 c f)}{4 a^2 f^2} \right) + \frac{d^2 x^3}{12 a^2} + \frac{c d x^2}{4 a^2}$$

[In] int((c + d*x)^2/(a + a*tanh(e + f*x))^2,x)

[Out] (c^2*x)/(4*a^2) - exp(- 4*e - 4*f*x)*((d^2 + 8*c^2*f^2 + 4*c*d*f)/(128*a^2*f^3) + (d^2*x^2)/(16*a^2*f) + (d*x*(d + 4*c*f))/(32*a^2*f^2)) - exp(- 2*e - 2*f*x)*((d^2 + 2*c^2*f^2 + 2*c*d*f)/(8*a^2*f^3) + (d^2*x^2)/(4*a^2*f) + (d*x*(d + 2*c*f))/(4*a^2*f^2)) + (d^2*x^3)/(12*a^2) + (c*d*x^2)/(4*a^2)

3.40 $\int \frac{c+dx}{(a+a \tanh(e+fx))^2} dx$

Optimal result	277
Rubi [A] (verified)	277
Mathematica [A] (verified)	279
Maple [A] (verified)	279
Fricas [A] (verification not implemented)	280
Sympy [F]	280
Maxima [A] (verification not implemented)	280
Giac [A] (verification not implemented)	281
Mupad [B] (verification not implemented)	281

Optimal result

Integrand size = 18, antiderivative size = 133

$$\int \frac{c+dx}{(a+a \tanh(e+fx))^2} dx = \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a \tanh(e+fx))^2}$$

$$- \frac{c+dx}{4f(a+a \tanh(e+fx))^2} - \frac{3d}{16f^2(a^2+a^2 \tanh(e+fx))}$$

$$- \frac{c+dx}{4f(a^2+a^2 \tanh(e+fx))}$$

[Out] 3/16*d*x/a^2/f-1/8*d*x^2/a^2+1/4*x*(d*x+c)/a^2-1/16*d/f^2/(a+a*tanh(f*x+e))^2+1/4*(-d*x-c)/f/(a+a*tanh(f*x+e))^2-3/16*d/f^2/(a^2+a^2*tanh(f*x+e))+1/4*(-d*x-c)/f/(a^2+a^2*tanh(f*x+e))

Rubi [A] (verified)

Time = 0.10 (sec) , antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3560, 8, 3811}

$$\int \frac{c+dx}{(a+a \tanh(e+fx))^2} dx = -\frac{c+dx}{4f(a^2 \tanh(e+fx)+a^2)} + \frac{x(c+dx)}{4a^2}$$

$$- \frac{3d}{16f^2(a^2 \tanh(e+fx)+a^2)} + \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2}$$

$$- \frac{c+dx}{4f(a \tanh(e+fx)+a)^2} - \frac{d}{16f^2(a \tanh(e+fx)+a)^2}$$

[In] Int[(c + d*x)/(a + a*Tanh[e + f*x])^2,x]

[Out] $(3*d*x)/(16*a^2*f) - (d*x^2)/(8*a^2) + (x*(c + d*x))/(4*a^2) - d/(16*f^2*(a + a*Tanh[e + f*x])^2) - (c + d*x)/(4*f*(a + a*Tanh[e + f*x])^2) - (3*d)/(16*f^2*(a^2 + a^2*Tanh[e + f*x])) - (c + d*x)/(4*f*(a^2 + a^2*Tanh[e + f*x]))$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] \text{ ; FreeQ}[a, x]$

Rule 3560

$\text{Int}[(a_) + (b_)*\tan[(c_) + (d_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[a*((a + b*\tan[c + d*x])^n/(2*b*d*n)), x] + \text{Dist}[1/(2*a), \text{Int}[(a + b*\tan[c + d*x])^{(n + 1)}, x], x] \text{ ; FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{LtQ}[n, 0]$

Rule 3811

$\text{Int}[(c_) + (d_)*(x_)^{(m_)}*((a_) + (b_)*\tan[(e_) + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{With}\{u = \text{IntHide}[(a + b*\tan[e + f*x])^n, x]\}, \text{Dist}[(c + d*x)^m, u, x] - \text{Dist}[d*m, \text{Int}[\text{Dist}[(c + d*x)^{(m - 1)}, u, x], x], x] \text{ ; FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{ILtQ}[n, -1] \ \&\& \ \text{GtQ}[m, 0]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x(c + dx)}{4a^2} - \frac{c + dx}{4f(a + a \tanh(e + fx))^2} - \frac{c + dx}{4f(a^2 + a^2 \tanh(e + fx))} \\
 &\quad - d \int \left(\frac{x}{4a^2} - \frac{1}{4f(a + a \tanh(e + fx))^2} - \frac{1}{4f(a^2 + a^2 \tanh(e + fx))} \right) dx \\
 &= -\frac{dx^2}{8a^2} + \frac{x(c + dx)}{4a^2} - \frac{c + dx}{4f(a + a \tanh(e + fx))^2} - \frac{c + dx}{4f(a^2 + a^2 \tanh(e + fx))} \\
 &\quad + \frac{d \int \frac{1}{(a + a \tanh(e + fx))^2} dx}{4f} + \frac{d \int \frac{1}{a^2 + a^2 \tanh(e + fx)} dx}{4f} \\
 &= -\frac{dx^2}{8a^2} + \frac{x(c + dx)}{4a^2} - \frac{d}{16f^2(a + a \tanh(e + fx))^2} - \frac{c + dx}{4f(a + a \tanh(e + fx))^2} \\
 &\quad - \frac{d}{8f^2(a^2 + a^2 \tanh(e + fx))} - \frac{c + dx}{4f(a^2 + a^2 \tanh(e + fx))} + \frac{d \int 1 dx}{8a^2 f} \\
 &\quad + \frac{d \int \frac{1}{a + a \tanh(e + fx)} dx}{8af} \\
 &= \frac{dx}{8a^2 f} - \frac{dx^2}{8a^2} + \frac{x(c + dx)}{4a^2} - \frac{d}{16f^2(a + a \tanh(e + fx))^2} - \frac{c + dx}{4f(a + a \tanh(e + fx))^2} \\
 &\quad - \frac{3d}{16f^2(a^2 + a^2 \tanh(e + fx))} - \frac{c + dx}{4f(a^2 + a^2 \tanh(e + fx))} + \frac{d \int 1 dx}{16a^2 f}
 \end{aligned}$$

$$= \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a \tanh(e+fx))^2} - \frac{c+dx}{4f(a+a \tanh(e+fx))^2}$$

$$- \frac{3d}{16f^2(a^2+a^2 \tanh(e+fx))} - \frac{c+dx}{4f(a^2+a^2 \tanh(e+fx))}$$

Mathematica [A] (verified)

Time = 0.99 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.86

$$\int \frac{c+dx}{(a+a \tanh(e+fx))^2} dx$$

$$= \frac{\operatorname{sech}^2(e+fx)(-8(d+2cf+2dfx) + (4cf(-1+4fx) + d(-1-4fx+8f^2x^2)) \cosh(2(e+fx)) + (4cf($$

$$64a^2f^2(1+\tanh(e+fx))^2$$

[In] Integrate[(c + d*x)/(a + a*Tanh[e + f*x])^2, x]

[Out] (Sech[e + f*x]^2*(-8*(d + 2*c*f + 2*d*f*x) + (4*c*f*(-1 + 4*f*x) + d*(-1 - 4*f*x + 8*f^2*x^2))*Cosh[2*(e + f*x)] + (4*c*f*(1 + 4*f*x) + d*(1 + 4*f*x + 8*f^2*x^2))*Sinh[2*(e + f*x)])/(64*a^2*f^2*(1 + Tanh[e + f*x])^2)

Maple [A] (verified)

Time = 0.25 (sec) , antiderivative size = 74, normalized size of antiderivative = 0.56

method	result
risch	$\frac{dx^2}{8a^2} + \frac{cx}{4a^2} - \frac{(2dx+2cf+d)e^{-2fx-2e}}{8a^2f^2} - \frac{(4dx+4cf+d)e^{-4fx-4e}}{64a^2f^2}$
parallelrisch	$\frac{-3d \tanh(fx+e)-4d+4cx f^2+2dx^2 f^2-4c \tanh(fx+e)f+2d \tanh(fx+e)xf-8cf-5dx+4d \tanh(fx+e)x^2 f^2+8x \tanh(fx+e)}{16f^2a^2(1+\tanh(fx+e))^2}$

[In] int((d*x+c)/(a+a*tanh(f*x+e))^2, x, method=_RETURNVERBOSE)

[Out] 1/8*d*x^2/a^2+1/4/a^2*c*x-1/8*(2*d*f*x+2*c*f+d)/a^2/f^2*exp(-2*f*x-2*e)-1/64*(4*d*f*x+4*c*f+d)/a^2/f^2*exp(-4*f*x-4*e)

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.44

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^2} dx = \frac{16dfx - (8df^2x^2 - 4cf + 4(4cf^2 - df)x - d) \cosh(fx + e)^2 - 2(8df^2x^2 + 4cf + 4(4cf^2 + df)x + d) \sinh(fx + e)^2}{64(a^2f^2 \cosh(fx + e))^2 + 2a^2f^2 \cosh(fx + e) \sinh(fx + e) + a^2f^2 \sinh(fx + e)^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] -1/64*(16*d*f*x - (8*d*f^2*x^2 - 4*c*f + 4*(4*c*f^2 - d*f)*x - d)*cosh(f*x + e)^2 - 2*(8*d*f^2*x^2 + 4*c*f + 4*(4*c*f^2 + d*f)*x + d)*cosh(f*x + e)*sinh(f*x + e) - (8*d*f^2*x^2 - 4*c*f + 4*(4*c*f^2 - d*f)*x - d)*sinh(f*x + e)^2 + 16*c*f + 8*d)/(a^2*f^2*cosh(f*x + e)^2 + 2*a^2*f^2*cosh(f*x + e)*sinh(f*x + e) + a^2*f^2*sinh(f*x + e)^2)

Sympy [F]

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^2} dx = \frac{\int \frac{c}{\tanh^2(e+fx)+2\tanh(e+fx)+1} dx + \int \frac{dx}{\tanh^2(e+fx)+2\tanh(e+fx)+1} dx}{a^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))**2,x)

[Out] (Integral(c/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x) + Integral(d*x/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x))/a**2

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.80

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^2} dx = \frac{1}{16} c \left(\frac{4(fx + e)}{a^2 f} - \frac{4e^{(-2fx-2e)} + e^{(-4fx-4e)}}{a^2 f} \right) + \frac{(8f^2x^2e^{(4e)} - 8(2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - (4fx + 1)e^{(-4fx)})de^{(-4e)}}{64a^2f^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] 1/16*c*(4*(f*x + e)/(a^2*f) - (4*e^(-2*f*x - 2*e) + e^(-4*f*x - 4*e)))/(a^2*f) + 1/64*(8*f^2*x^2*e^(4*e) - 8*(2*f*x*e^(2*e) + e^(2*e))*e^(-2*f*x) - (4*f*x + 1)*e^(-4*f*x))*d*e^(-4*e)/(a^2*f^2)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.77

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^2} dx$$

$$= \frac{(8df^2x^2e^{4fx+4e} + 16cf^2xe^{4fx+4e} - 16dfxe^{2fx+2e} - 4dfx - 16cfe^{2fx+2e} - 4cf - 8de^{2fx+2e} - d)e^{-4fx-4e}}{64a^2f^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

```
[Out] 1/64*(8*d*f^2*x^2*e^(4*f*x + 4*e) + 16*c*f^2*x*e^(4*f*x + 4*e) - 16*d*f*x*e^(2*f*x + 2*e) - 4*d*f*x - 16*c*f*e^(2*f*x + 2*e) - 4*c*f - 8*d*e^(2*f*x + 2*e) - d)*e^(-4*f*x - 4*e)/(a^2*f^2)
```

Mupad [B] (verification not implemented)

Time = 1.79 (sec) , antiderivative size = 89, normalized size of antiderivative = 0.67

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^2} dx = \frac{dx^2}{8a^2} - e^{-4e-4fx} \left(\frac{d + 4cf}{64a^2f^2} + \frac{dx}{16a^2f} \right) - e^{-2e-2fx} \left(\frac{d + 2cf}{8a^2f^2} + \frac{dx}{4a^2f} \right) + \frac{cx}{4a^2}$$

[In] int((c + d*x)/(a + a*tanh(e + f*x))^2,x)

```
[Out] (d*x^2)/(8*a^2) - exp(- 4*e - 4*f*x)*((d + 4*c*f)/(64*a^2*f^2) + (d*x)/(16*a^2*f)) - exp(- 2*e - 2*f*x)*((d + 2*c*f)/(8*a^2*f^2) + (d*x)/(4*a^2*f)) + (c*x)/(4*a^2)
```

3.41 $\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx$

Optimal result	282
Rubi [A] (verified)	283
Mathematica [A] (verified)	285
Maple [A] (verified)	286
Fricas [A] (verification not implemented)	286
Sympy [F]	286
Maxima [A] (verification not implemented)	287
Giac [A] (verification not implemented)	287
Mupad [F(-1)]	287

Optimal result

Integrand size = 20, antiderivative size = 297

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx = \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} + \frac{\log(c+dx)}{4a^2d} - \frac{\operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{4a^2d} - \frac{\operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} + \frac{\sinh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d}$$

```
[Out] 1/4*Chi(4*c*f/d+4*f*x)*cosh(-4*e+4*c*f/d)/a^2/d+1/2*Chi(2*c*f/d+2*f*x)*cosh
(-2*e+2*c*f/d)/a^2/d+1/4*ln(d*x+c)/a^2/d-1/2*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d
+2*f*x)/a^2/d-1/4*cosh(-4*e+4*c*f/d)*Shi(4*c*f/d+4*f*x)/a^2/d+1/4*Chi(4*c*f
/d+4*f*x)*sinh(-4*e+4*c*f/d)/a^2/d-1/4*Shi(4*c*f/d+4*f*x)*sinh(-4*e+4*c*f/d
)/a^2/d+1/2*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a^2/d-1/2*Shi(2*c*f/d+2*f
*x)*sinh(-2*e+2*c*f/d)/a^2/d
```

Rubi [A] (verified)

Time = 0.58 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3809, 3384, 3379, 3382, 3393}

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx = -\frac{\text{Chi}\left(2xf + \frac{2cf}{d}\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{\text{Chi}\left(4xf + \frac{4cf}{d}\right) \sinh\left(4e - \frac{4cf}{d}\right)}{4a^2d} + \frac{\text{Chi}\left(2xf + \frac{2cf}{d}\right) \cosh\left(2e - \frac{2cf}{d}\right)}{2a^2d} + \frac{\text{Chi}\left(4xf + \frac{4cf}{d}\right) \cosh\left(4e - \frac{4cf}{d}\right)}{4a^2d} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} + \frac{\sinh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{2a^2d} - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(4xf + \frac{4cf}{d}\right)}{4a^2d} + \frac{\log(c+dx)}{4a^2d}$$

[In] Int[1/((c + d*x)*(a + a*Tanh[e + f*x])^2), x]

[Out] (Cosh[2*e - (2*c*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x])/(2*a^2*d) + (Cosh[4*e - (4*c*f)/d]*CoshIntegral[(4*c*f)/d + 4*f*x])/(4*a^2*d) + Log[c + d*x]/(4*a^2*d) - (CoshIntegral[(4*c*f)/d + 4*f*x]*Sinh[4*e - (4*c*f)/d])/(4*a^2*d) - (CoshIntegral[(2*c*f)/d + 2*f*x]*Sinh[2*e - (2*c*f)/d])/(2*a^2*d) - (Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(2*a^2*d) + (Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(2*a^2*d) - (Cosh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(4*a^2*d) + (Sinh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(4*a^2*d)

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := In
t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f
, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2(c+dx)} + \frac{\cosh^2(2e+2fx)}{4a^2(c+dx)} - \frac{\sinh(2e+2fx)}{2a^2(c+dx)} \right. \\
&\quad \left. + \frac{\sinh^2(2e+2fx)}{4a^2(c+dx)} - \frac{\sinh(4e+4fx)}{4a^2(c+dx)} \right) dx \\
&= \frac{\log(c+dx)}{4a^2d} + \frac{\int \frac{\cosh^2(2e+2fx)}{c+dx} dx}{4a^2} + \frac{\int \frac{\sinh^2(2e+2fx)}{c+dx} dx}{4a^2} \\
&\quad - \frac{\int \frac{\sinh(4e+4fx)}{c+dx} dx}{4a^2} + \frac{\int \frac{\cosh(2e+2fx)}{c+dx} dx}{2a^2} - \frac{\int \frac{\sinh(2e+2fx)}{c+dx} dx}{2a^2} \\
&= \frac{\log(c+dx)}{4a^2d} - \frac{\int \left(\frac{1}{2(c+dx)} - \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} + \frac{\int \left(\frac{1}{2(c+dx)} + \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} \\
&\quad - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \int \frac{\sinh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{4a^2} + \frac{\cosh\left(2e - \frac{2cf}{d}\right) \int \frac{\cosh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a^2} \\
&\quad - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \int \frac{\sinh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a^2} - \frac{\sinh\left(4e - \frac{4cf}{d}\right) \int \frac{\cosh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{4a^2} \\
&\quad - \frac{\sinh\left(2e - \frac{2cf}{d}\right) \int \frac{\cosh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a^2} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \int \frac{\sinh\left(\frac{2cf}{d} + 2fx\right)}{c+dx} dx}{2a^2}
\end{aligned}$$

$$\begin{aligned}
&= \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\log(c+dx)}{4a^2d} \\
&\quad - \frac{\operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{4a^2d} - \frac{\operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{4a^2d} \\
&\quad - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\
&\quad - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} + 2 \frac{\int \frac{\cosh(4e+4fx)}{c+dx} dx}{8a^2} \\
&= \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\log(c+dx)}{4a^2d} - \frac{\operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{4a^2d} \\
&\quad - \frac{\operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\
&\quad + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} \\
&\quad + 2 \left(\frac{\cosh\left(4e - \frac{4cf}{d}\right) \int \frac{\cosh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{8a^2} + \frac{\sinh\left(4e - \frac{4cf}{d}\right) \int \frac{\sinh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{8a^2} \right) \\
&= \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} + \frac{\log(c+dx)}{4a^2d} - \frac{\operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{4a^2d} \\
&\quad - \frac{\operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{2a^2d} - \frac{\cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} \\
&\quad + \frac{\sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{2a^2d} - \frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{4a^2d} \\
&\quad + 2 \left(\frac{\cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right)}{8a^2d} + \frac{\sinh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{8a^2d} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.80 (sec) , antiderivative size = 199, normalized size of antiderivative = 0.67

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx$$

$$= \frac{(\cosh(2e - \frac{2cf}{d}) - \sinh(2e - \frac{2cf}{d})) \left(2\operatorname{Chi}\left(\frac{2f(c+dx)}{d}\right) + \cosh\left(2e - \frac{2cf}{d}\right) \log(f(c+dx)) + \operatorname{Chi}\left(\frac{4f(c+dx)}{d}\right) \right)}{(4a^2d)}$$

[In] Integrate[1/((c + d*x)*(a + a*Tanh[e + f*x])^2),x]

[Out] ((Cosh[2*e - (2*c*f)/d] - Sinh[2*e - (2*c*f)/d])*(2*CoshIntegral[(2*f*(c + d*x))/d] + Cosh[2*e - (2*c*f)/d]*Log[f*(c + d*x)] + CoshIntegral[(4*f*(c + d*x))/d]*(Cosh[2*e - (2*c*f)/d] - Sinh[2*e - (2*c*f)/d]) + Log[f*(c + d*x)]*Sinh[2*e - (2*c*f)/d] - 2*SinhIntegral[(2*f*(c + d*x))/d] - Cosh[2*e - (2*c*f)/d]*SinhIntegral[(4*f*(c + d*x))/d] + Sinh[2*e - (2*c*f)/d]*SinhIntegral[(4*f*(c + d*x))/d])/(4*a^2*d)

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.36

method	result	size
risch	$\frac{\ln(dx+c)}{4a^2d} - \frac{e^{\frac{4cf-4de}{d}} \operatorname{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{4a^2d} - \frac{e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{2a^2d}$	106

[In] `int(1/(d*x+c)/(a+a*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4} \ln(d*x+c)/a^2/d - 1/4/a^2/d * \exp(4*(c*f-d*e)/d) * \operatorname{Ei}(1, 4*f*x+4*e+4*(c*f-d*e)/d) - 1/2/a^2/d * \exp(2*(c*f-d*e)/d) * \operatorname{Ei}(1, 2*f*x+2*e+2*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 133, normalized size of antiderivative = 0.45

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx$$

$$= \frac{2 \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) \cosh\left(-\frac{2(de-cf)}{d}\right) + \operatorname{Ei}\left(-\frac{4(dfx+cf)}{d}\right) \cosh\left(-\frac{4(de-cf)}{d}\right) + 2 \operatorname{Ei}\left(-\frac{2(dfx+cf)}{d}\right) \sinh\left(-\frac{2(de-cf)}{d}\right)}{4a^2d}$$

[In] `integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")`

[Out] $\frac{1}{4} * (2 * \operatorname{Ei}(-2*(d*f*x + c*f)/d) * \cosh(-2*(d*e - c*f)/d) + \operatorname{Ei}(-4*(d*f*x + c*f)/d) * \cosh(-4*(d*e - c*f)/d) + 2 * \operatorname{Ei}(-2*(d*f*x + c*f)/d) * \sinh(-2*(d*e - c*f)/d) + \operatorname{Ei}(-4*(d*f*x + c*f)/d) * \sinh(-4*(d*e - c*f)/d) + \log(d*x + c)) / (a^2*d)$

Sympy [F]

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx$$

$$= \frac{\int \frac{1}{c \tanh^2(e+fx) + 2c \tanh(e+fx) + c + dx \tanh^2(e+fx) + 2dx \tanh(e+fx) + dx} dx}{a^2}$$

[In] `integrate(1/(d*x+c)/(a+a*tanh(f*x+e))**2,x)`

[Out] `Integral(1/(c*tanh(e + f*x)**2 + 2*c*tanh(e + f*x) + c + d*x*tanh(e + f*x)**2 + 2*d*x*tanh(e + f*x) + d*x), x)/a**2`

Maxima [A] (verification not implemented)

none

Time = 0.71 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.27

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx = -\frac{e^{(-4e+\frac{4cf}{d})} E_1\left(\frac{4(dx+c)f}{d}\right)}{4a^2d} - \frac{e^{(-2e+\frac{2cf}{d})} E_1\left(\frac{2(dx+c)f}{d}\right)}{2a^2d} + \frac{\log(dx+c)}{4a^2d}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] -1/4*e^(-4*e + 4*c*f/d)*exp_integral_e(1, 4*(d*x + c)*f/d)/(a^2*d) - 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(1, 2*(d*x + c)*f/d)/(a^2*d) + 1/4*log(d*x + c)/(a^2*d)

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.25

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx = \frac{\left(2 \operatorname{Ei}\left(-\frac{2(dfxc+cf)}{d}\right) e^{(2e+\frac{2cf}{d})} + \operatorname{Ei}\left(-\frac{4(dfxc+cf)}{d}\right) e^{(\frac{4cf}{d})} + e^{(4e)} \log(dx+c)\right) e^{(-4e)}}{4a^2d}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

[Out] 1/4*(2*Ei(-2*(d*f*x + c*f)/d)*e^(2*e + 2*c*f/d) + Ei(-4*(d*f*x + c*f)/d)*e^(4*c*f/d) + e^(4*e)*log(d*x + c))*e^(-4*e)/(a^2*d)

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^2} dx = \int \frac{1}{(a+a \tanh(e+fx))^2 (c+dx)} dx$$

[In] int(1/((a + a*tanh(e + f*x))^2*(c + d*x)),x)

[Out] int(1/((a + a*tanh(e + f*x))^2*(c + d*x)), x)

$$3.42 \quad \int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx$$

Optimal result	288
Rubi [A] (verified)	289
Mathematica [A] (verified)	292
Maple [A] (verified)	293
Fricas [A] (verification not implemented)	293
Sympy [F]	294
Maxima [A] (verification not implemented)	294
Giac [A] (verification not implemented)	295
Mupad [F(-1)]	295

Optimal result

Integrand size = 20, antiderivative size = 420

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx = -\frac{1}{4a^2d(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)}$$

$$- \frac{f \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$- \frac{f \cosh(4e - \frac{4cf}{d}) \operatorname{Chi}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

$$+ \frac{f \operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{a^2d^2}$$

$$+ \frac{f \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{a^2d^2}$$

$$+ \frac{\sinh(2e+2fx)}{2a^2d(c+dx)}$$

$$- \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)}$$

$$+ \frac{f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$- \frac{f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2}$$

$$+ \frac{f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

$$- \frac{f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2}$$

```
[Out] -1/4/a^2/d/(d*x+c)-f*Chi(4*c*f/d+4*f*x)*cosh(-4*e+4*c*f/d)/a^2/d^2-f*Chi(2*
c*f/d+2*f*x)*cosh(-2*e+2*c*f/d)/a^2/d^2-1/2*cosh(2*f*x+2*e)/a^2/d/(d*x+c)-1
/4*cosh(2*f*x+2*e)^2/a^2/d/(d*x+c)+f*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d+2*f*x)/
```


$$\begin{aligned} & a^2/d^2 + f \cosh(-4e + 4cf/d) \operatorname{Shi}(4cf/d + 4fx) / a^2/d^2 - f \operatorname{Chi}(4cf/d + 4fx) \\ & \operatorname{sinh}(-4e + 4cf/d) / a^2/d^2 + f \operatorname{Shi}(4cf/d + 4fx) \operatorname{sinh}(-4e + 4cf/d) / a^2/d^2 \\ & - f \operatorname{Chi}(2cf/d + 2fx) \operatorname{sinh}(-2e + 2cf/d) / a^2/d^2 + f \operatorname{Shi}(2cf/d + 2fx) \operatorname{sinh} \\ & (-2e + 2cf/d) / a^2/d^2 + 1/2 \operatorname{sinh}(2fx + 2e) / a^2/d / (dx + c) - 1/4 \operatorname{sinh}(2fx + 2e) \\ & ^2 / a^2/d / (dx + c) + 1/4 \operatorname{sinh}(4fx + 4e) / a^2/d / (dx + c) \end{aligned}$$

Rubi [A] (verified)

Time = 0.55 (sec) , antiderivative size = 420, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3809, 3378, 3384, 3379, 3382, 3394, 12}

$$\begin{aligned} \int \frac{1}{(c + dx)^2 (a + a \tanh(e + fx))^2} dx = & \frac{f \operatorname{Chi}(4xf + \frac{4cf}{d}) \operatorname{sinh}(4e - \frac{4cf}{d})}{a^2 d^2} \\ & + \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \operatorname{sinh}(2e - \frac{2cf}{d})}{a^2 d^2} \\ & - \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{a^2 d^2} \\ & - \frac{f \operatorname{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{a^2 d^2} \\ & - \frac{f \operatorname{sinh}(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{a^2 d^2} \\ & - \frac{f \operatorname{sinh}(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{a^2 d^2} \\ & + \frac{f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{a^2 d^2} \\ & + \frac{f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{a^2 d^2} \\ & - \frac{\operatorname{sinh}^2(2e + 2fx)}{4a^2 d (c + dx)} + \frac{\operatorname{sinh}(2e + 2fx)}{2a^2 d (c + dx)} \\ & + \frac{\operatorname{sinh}(4e + 4fx)}{4a^2 d (c + dx)} - \frac{\cosh^2(2e + 2fx)}{4a^2 d (c + dx)} \\ & - \frac{\cosh(2e + 2fx)}{2a^2 d (c + dx)} - \frac{1}{4a^2 d (c + dx)} \end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + a*Tanh[e + f*x])^2),x]

[Out] $-1/4 * 1 / (a^2 * d * (c + d * x)) - \operatorname{Cosh}[2 * e + 2 * f * x] / (2 * a^2 * d * (c + d * x)) - \operatorname{Cosh}[2 * e + 2 * f * x]^2 / (4 * a^2 * d * (c + d * x)) - (f * \operatorname{Cosh}[2 * e - (2 * c * f) / d] * \operatorname{CoshIntegral}[(2 * c * f) / d + 2 * f * x]) / (a^2 * d^2) - (f * \operatorname{Cosh}[4 * e - (4 * c * f) / d] * \operatorname{CoshIntegral}[(4 * c * f) / d + 4 * f * x]) / (a^2 * d^2) + (f * \operatorname{CoshIntegral}[(4 * c * f) / d + 4 * f * x] * \operatorname{Sinh}[4 * e - (4 * c * f) / d]) / (a^2 * d^2) + (f * \operatorname{CoshIntegral}[(2 * c * f) / d + 2 * f * x] * \operatorname{Sinh}[2 * e - (2 * c * f) / d]) / (a^2 * d^2) + \operatorname{Sinh}[2 * e + 2 * f * x] / (2 * a^2 * d * (c + d * x)) - \operatorname{Sinh}[2 * e + 2 * f * x]^2 / ($

$$4a^2d(c + dx) + \frac{\sinh[4e + 4fx]}{4a^2d(c + dx)} + \frac{f \cosh[2e - (2cf)/d] \operatorname{SinhIntegral}[(2cf)/d + 2fx]}{a^2d^2} - \frac{f \sinh[2e - (2cf)/d] \operatorname{SinhIntegral}[(2cf)/d + 2fx]}{a^2d^2} + \frac{f \cosh[4e - (4cf)/d] \operatorname{SinhIntegral}[(4cf)/d + 4fx]}{a^2d^2} - \frac{f \sinh[4e - (4cf)/d] \operatorname{SinhIntegral}[(4cf)/d + 4fx]}{a^2d^2}$$
Rule 12

```
Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]
```

Rule 3378

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)], x_Symbol] := Simp[(c + dx)^(m + 1)*(Sin[e + fx]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + dx)^(m + 1)*Cos[e + fx], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

Rule 3379

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_) + (Complex[0, fz_])*(f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_) + (f_)*(x_)]/((c_) + (d_)*(x_)), x_Symbol] := Dist[Cos[(d*e - cf)/d], Int[Sin[c*(f/d) + fx]/(c + dx), x], x] + Dist[Sin[(d*e - cf)/d], Int[Cos[c*(f/d) + fx]/(c + dx), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - cf, 0]
```

Rule 3394

```
Int[((c_) + (d_)*(x_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Simp[(c + dx)^(m + 1)*(Sin[e + fx]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1))), Int[ExpandTrigReduce[(c + dx)^(m + 1), Cos[e + fx]*Sin[e + fx]^(n - 1), x], x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] && LtQ[m, -1]
```

Rule 3809

```

Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)^2} + \frac{\cosh(2e+2fx)}{2a^2(c+dx)^2} + \frac{\cosh^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{\sinh(2e+2fx)}{2a^2(c+dx)^2} \right. \\
&\quad \left. + \frac{\sinh^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{\sinh(4e+4fx)}{4a^2(c+dx)^2} \right) dx \\
&= -\frac{1}{4a^2d(c+dx)} + \frac{\int \frac{\cosh^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} + \frac{\int \frac{\sinh^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} \\
&\quad - \frac{\int \frac{\sinh(4e+4fx)}{(c+dx)^2} dx}{4a^2} + \frac{\int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{2a^2} - \frac{\int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{2a^2} \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} \\
&\quad + \frac{(if) \int -\frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{a^2d} - \frac{(if) \int \frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{a^2d} \\
&\quad - \frac{f \int \frac{\cosh(2e+2fx)}{c+dx} dx}{a^2d} - \frac{f \int \frac{\cosh(4e+4fx)}{c+dx} dx}{a^2d} + \frac{f \int \frac{\sinh(2e+2fx)}{c+dx} dx}{a^2d} \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} \\
&\quad - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} + 2 \frac{f \int \frac{\sinh(4e+4fx)}{c+dx} dx}{2a^2d} \\
&\quad - \frac{(f \cosh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} - \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} \\
&\quad + \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} - \frac{(f \sinh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} \\
&\quad + \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} - \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad - \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Chi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} - \frac{f \cosh\left(4e - \frac{4cf}{d}\right) \text{Chi}\left(\frac{4cf}{d} + 4fx\right)}{a^2d^2} \\
&\quad + \frac{f \text{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{a^2d^2} + \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} + \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} \\
&\quad - \frac{f \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} - \frac{f \sinh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(\frac{4cf}{d} + 4fx\right)}{a^2d^2} \\
&\quad + 2 \left(\frac{\left(f \cosh\left(4e - \frac{4cf}{d}\right)\right) \int \frac{\sinh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{2a^2d} \right. \\
&\quad \quad \left. + \frac{\left(f \sinh\left(4e - \frac{4cf}{d}\right)\right) \int \frac{\cosh\left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{2a^2d} \right) \\
&= -\frac{1}{4a^2d(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad - \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Chi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} - \frac{f \cosh\left(4e - \frac{4cf}{d}\right) \text{Chi}\left(\frac{4cf}{d} + 4fx\right)}{a^2d^2} \\
&\quad + \frac{f \text{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{a^2d^2} + \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} + \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} \\
&\quad - \frac{f \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{a^2d^2} - \frac{f \sinh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(\frac{4cf}{d} + 4fx\right)}{a^2d^2} \\
&\quad + 2 \left(\frac{f \text{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{2a^2d^2} + \frac{f \cosh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(\frac{4cf}{d} + 4fx\right)}{2a^2d^2} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.48 (sec) , antiderivative size = 442, normalized size of antiderivative = 1.05

$$\begin{aligned}
&\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx \\
&= \frac{(-\cosh(2(e+f(-\frac{c}{d}+x)))) + \sinh(2(e+f(-\frac{c}{d}+x)))) \left(2d \cosh\left(\frac{2cf}{d}\right) + d \cosh(2(e+f(-\frac{c}{d}+x)))\right) + \dots}{\dots}
\end{aligned}$$

[In] Integrate[1/((c+d*x)^2*(a+a*Tanh[e+f*x])^2),x]

```
[Out] ((-Cosh[2*(e + f*(-(c/d) + x))] + Sinh[2*(e + f*(-(c/d) + x))])*(2*d*Cosh[(2*c*f)/d] + d*Cosh[2*(e + f*(-(c/d) + x))] + d*Cosh[2*(e + f*(c/d + x))] - 2*d*Sinh[(2*c*f)/d] + 4*f*(c + d*x)*CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[2*f*x] + Sinh[2*f*x]) + d*Sinh[2*(e + f*(-(c/d) + x))] - d*Sinh[2*(e + f*(c/d + x))] + 4*f*(c + d*x)*CoshIntegral[(4*f*(c + d*x))/d]*(Cosh[2*e - (2*f*(c + d*x))/d] - Sinh[2*e - (2*f*(c + d*x))/d]) - 4*c*f*Cosh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 4*d*f*x*Cosh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 4*c*f*Sinh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 4*d*f*x*Sinh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 4*c*f*Cosh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] - 4*d*f*x*Cosh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] + 4*c*f*Sinh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] + 4*d*f*x*Sinh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d]))/(4*a^2*d^2*(c + d*x))
```

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.39

method	result
risch	$-\frac{1}{4a^2d(dx+c)} - \frac{f e^{-4fx-4e}}{4a^2d(dx+cf)} + \frac{f e^{\frac{4cf-4de}{d}} \operatorname{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{a^2d^2} - \frac{f e^{-2fx-2e}}{2a^2d(dx+cf)} + \frac{f e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{a^2d^2}$

```
[In] int(1/(d*x+c)^2/(a+a*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] -1/4/a^2/d/(d*x+c)-1/4*f/a^2*exp(-4*f*x-4*e)/d/(d*f*x+c*f)+f/a^2/d^2*exp(4*(c*f-d*e)/d)*Ei(1,4*f*x+4*e+4*(c*f-d*e)/d)-1/2*f/a^2*exp(-2*f*x-2*e)/d/(d*f*x+c*f)+f/a^2/d^2*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 609, normalized size of antiderivative = 1.45

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx = \frac{2(df x + cf) \operatorname{Ei}\left(-\frac{2(df x + cf)}{d}\right) \cosh(fx + e)^2 \sinh\left(-\frac{2(de - cf)}{d}\right) + 2(df x + cf) \operatorname{Ei}\left(-\frac{4(df x + cf)}{d}\right) \cosh(fx + e)^2 \sinh\left(-\frac{4(de - cf)}{d}\right)}{4a^2d^2}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] -1/2*(2*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-2*(d*e - c*f)/d) + 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-4*(d*e - c*f)/d) + (2*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + d)*co
```

$$\begin{aligned} & \operatorname{sh}(f*x + e)^2 + (2*(d*f*x + c*f)*\operatorname{Ei}(-2*(d*f*x + c*f)/d)*\operatorname{cosh}(-2*(d*e - c*f)/d) \\ & + 2*(d*f*x + c*f)*\operatorname{Ei}(-4*(d*f*x + c*f)/d)*\operatorname{cosh}(-4*(d*e - c*f)/d) + 2*(d*f*x + c*f)* \\ & \operatorname{Ei}(-2*(d*f*x + c*f)/d)*\operatorname{sinh}(-2*(d*e - c*f)/d) + 2*(d*f*x + c*f)* \\ & \operatorname{Ei}(-4*(d*f*x + c*f)/d)*\operatorname{sinh}(-4*(d*e - c*f)/d) + d)*\operatorname{sinh}(f*x + e)^2 + 4*((d*f*x + c*f)* \\ & \operatorname{Ei}(-2*(d*f*x + c*f)/d)*\operatorname{cosh}(f*x + e)*\operatorname{sinh}(-2*(d*e - c*f)/d) + (d*f*x + c*f)* \\ & \operatorname{Ei}(-4*(d*f*x + c*f)/d)*\operatorname{cosh}(f*x + e)*\operatorname{sinh}(-4*(d*e - c*f)/d) + (d*f*x + c*f)* \\ & \operatorname{Ei}(-2*(d*f*x + c*f)/d)*\operatorname{cosh}(-2*(d*e - c*f)/d) + (d*f*x + c*f)* \\ & \operatorname{Ei}(-4*(d*f*x + c*f)/d)*\operatorname{cosh}(-4*(d*e - c*f)/d))*\operatorname{cosh}(f*x + e))*\operatorname{sinh}(f*x + e) \\ &) + d)/((a^2*d^3*x + a^2*c*d^2)*\operatorname{cosh}(f*x + e)^2 + 2*(a^2*d^3*x + a^2*c*d^2)* \\ & *\operatorname{cosh}(f*x + e)*\operatorname{sinh}(f*x + e) + (a^2*d^3*x + a^2*c*d^2)*\operatorname{sinh}(f*x + e)^2) \end{aligned}$$

Sympy [F]

$$\int \frac{1}{(c + dx)^2(a + a \tanh(e + fx))^2} dx = \frac{\int \frac{1}{c^2 \tanh^2(e+fx) + 2c^2 \tanh(e+fx) + c^2 + 2cdx \tanh^2(e+fx) + 4cdx \tanh(e+fx) + 2cdx + d^2x^2 \tanh^2(e+fx) + 2d^2x^2 \tanh(e+fx) + d^2x^2} dx}{a^2}$$

[In] integrate(1/(d*x+c)**2/(a+a*tanh(f*x+e))**2,x)

[Out] Integral(1/(c**2*tanh(e + f*x)**2 + 2*c**2*tanh(e + f*x) + c**2 + 2*c*d*x*tanh(e + f*x)**2 + 4*c*d*x*tanh(e + f*x) + 2*c*d*x + d**2*x**2*tanh(e + f*x)**2 + 2*d**2*x**2*tanh(e + f*x) + d**2*x**2), x)/a**2

Maxima [A] (verification not implemented)

none

Time = 1.35 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.24

$$\int \frac{1}{(c + dx)^2(a + a \tanh(e + fx))^2} dx = -\frac{1}{4(a^2d^2x + a^2cd)} - \frac{e^{(-4e + \frac{4cf}{d})} E_2\left(\frac{4(dx+c)f}{d}\right)}{4(dx+c)a^2d} - \frac{e^{(-2e + \frac{2cf}{d})} E_2\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)a^2d}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] -1/4/(a^2*d^2*x + a^2*c*d) - 1/4*e^(-4*e + 4*c*f/d)*exp_integral_e(2, 4*(d*x + c)*f/d)/((d*x + c)*a^2*d) - 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(2, 2*(d*x + c)*f/d)/((d*x + c)*a^2*d)

Giac [A] (verification not implemented)

none

Time = 0.31 (sec) , antiderivative size = 583, normalized size of antiderivative = 1.39

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx =$$

$$\left(4(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) f^2 \operatorname{Ei}\left(-\frac{2((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de+cf)}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 4def^2 \operatorname{Ei}\left(-\frac{2((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de+cf)}{d}\right) \right)$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

```
[Out] -1/4*(4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-2*((d*x + c)*
(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) -
4*d*e*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)
/d)*e^(-2*(d*e - c*f)/d) + 4*c*f^3*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d
*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) + 4*(d*x + c)*(d*e/(d*x +
c) - c*f/(d*x + c) + f)*f^2*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c
) + f) - d*e + c*f)/d)*e^(-4*(d*e - c*f)/d) - 4*d*e*f^2*Ei(-4*((d*x + c)*(d
*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*(d*e - c*f)/d) + 4*
c*f^3*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*
e^(-4*(d*e - c*f)/d) + 2*d*f^2*e^(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x +
c) + f)/d) + d*f^2*e^(-4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) +
d*f^2)*d^2/(((d*x + c)*a^2*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a^2*d
^5*e + a^2*c*d^4*f)*f)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^2} dx = \int \frac{1}{(a+a \tanh(e+fx))^2 (c+dx)^2} dx$$

[In] int(1/((a + a*tanh(e + f*x))^2*(c + d*x)^2),x)

[Out] int(1/((a + a*tanh(e + f*x))^2*(c + d*x)^2), x)

3.43 $\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx$

Optimal result	296
Rubi [A] (verified)	297
Mathematica [A] (verified)	299
Maple [A] (verified)	299
Fricas [B] (verification not implemented)	300
Sympy [F]	300
Maxima [A] (verification not implemented)	301
Giac [A] (verification not implemented)	302
Mupad [B] (verification not implemented)	302

Optimal result

Integrand size = 20, antiderivative size = 336

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx = -\frac{d^3 e^{-6e-6fx}}{1728a^3 f^4} - \frac{9d^3 e^{-4e-4fx}}{1024a^3 f^4} - \frac{9d^3 e^{-2e-2fx}}{64a^3 f^4} - \frac{d^2 e^{-6e-6fx}(c+dx)}{288a^3 f^3} - \frac{9d^2 e^{-4e-4fx}(c+dx)}{256a^3 f^3} - \frac{9d^2 e^{-2e-2fx}(c+dx)}{32a^3 f^3} - \frac{de^{-6e-6fx}(c+dx)^2}{96a^3 f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{96a^3 f^2} - \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3 f^2} - \frac{e^{-6e-6fx}(c+dx)^3}{48a^3 f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3 f} - \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3 f} + \frac{(c+dx)^4}{32a^3 d}$$

```
[Out] -1/1728*d^3*exp(-6*f*x-6*e)/a^3/f^4-9/1024*d^3*exp(-4*f*x-4*e)/a^3/f^4-9/64*d^3*exp(-2*f*x-2*e)/a^3/f^4-1/288*d^2*exp(-6*f*x-6*e)*(d*x+c)/a^3/f^3-9/256*d^2*exp(-4*f*x-4*e)*(d*x+c)/a^3/f^3-9/32*d^2*exp(-2*f*x-2*e)*(d*x+c)/a^3/f^3-1/96*d*exp(-6*f*x-6*e)*(d*x+c)^2/a^3/f^2-9/128*d*exp(-4*f*x-4*e)*(d*x+c)^2/a^3/f^2-9/32*d*exp(-2*f*x-2*e)*(d*x+c)^2/a^3/f^2-1/48*exp(-6*f*x-6*e)*(d*x+c)^3/a^3/f-3/32*exp(-4*f*x-4*e)*(d*x+c)^3/a^3/f-3/16*exp(-2*f*x-2*e)*(d*x+c)^3/a^3/f+1/32*(d*x+c)^4/a^3/d
```


Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3810, 2207, 2225}

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx = -\frac{d^2(c+dx)e^{-6e-6fx}}{288a^3f^3} - \frac{9d^2(c+dx)e^{-4e-4fx}}{256a^3f^3} - \frac{9d^2(c+dx)e^{-2e-2fx}}{32a^3f^3} - \frac{d(c+dx)^2e^{-6e-6fx}}{96a^3f^2} - \frac{9d(c+dx)^2e^{-4e-4fx}}{128a^3f^2} - \frac{9d(c+dx)^2e^{-2e-2fx}}{32a^3f^2} - \frac{(c+dx)^3e^{-6e-6fx}}{48a^3f} - \frac{3(c+dx)^3e^{-4e-4fx}}{32a^3f} - \frac{3(c+dx)^3e^{-2e-2fx}}{16a^3f} + \frac{(c+dx)^4}{32a^3d} - \frac{d^3e^{-6e-6fx}}{1728a^3f^4} - \frac{9d^3e^{-4e-4fx}}{1024a^3f^4} - \frac{9d^3e^{-2e-2fx}}{64a^3f^4}$$

[In] Int[(c + d*x)^3/(a + a*Tanh[e + f*x])^3,x]

[Out] $-1/1728*(d^3E^{-6e-6fx})/(a^3f^4) - (9*d^3E^{-4e-4fx})/(1024*a^3f^4) - (9*d^3E^{-2e-2fx})/(64*a^3f^4) - (d^2E^{-6e-6fx}*(c+d*x))/(288*a^3f^3) - (9*d^2E^{-4e-4fx}*(c+d*x))/(256*a^3f^3) - (9*d^2E^{-2e-2fx}*(c+d*x))/(32*a^3f^3) - (dE^{-6e-6fx}*(c+d*x)^2)/(96*a^3f^2) - (9*dE^{-4e-4fx}*(c+d*x)^2)/(128*a^3f^2) - (9*dE^{-2e-2fx}*(c+d*x)^2)/(32*a^3f^2) - (E^{-6e-6fx}*(c+d*x)^3)/(48*a^3f) - (3E^{-4e-4fx}*(c+d*x)^3)/(32*a^3f) - (3E^{-2e-2fx}*(c+d*x)^3)/(16*a^3f) + (c+d*x)^4/(32*a^3*d)$

Rule 2207

Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m-1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[\$UseGamma]

Rule 2225

Int[(F_)^((c_.)*((a_.) + (b_.)*(x_)))^(n_.), x_Symbol] := Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 3810

Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))]

x)) / (2*a)) ^ (-n), x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^3}{8a^3} + \frac{e^{-6e-6fx}(c+dx)^3}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^3}{8a^3} + \frac{3e^{-2e-2fx}(c+dx)^3}{8a^3} \right) dx \\
&= \frac{(c+dx)^4}{32a^3d} + \frac{\int e^{-6e-6fx}(c+dx)^3 dx}{8a^3} + \frac{3 \int e^{-4e-4fx}(c+dx)^3 dx}{8a^3} + \frac{3 \int e^{-2e-2fx}(c+dx)^3 dx}{8a^3} \\
&= -\frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} \\
&\quad + \frac{(c+dx)^4}{32a^3d} + \frac{d \int e^{-6e-6fx}(c+dx)^2 dx}{16a^3f} + \frac{(9d) \int e^{-4e-4fx}(c+dx)^2 dx}{32a^3f} \\
&\quad + \frac{(9d) \int e^{-2e-2fx}(c+dx)^2 dx}{16a^3f} \\
&= -\frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} \\
&\quad - \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} - \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} \\
&\quad - \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} + \frac{(c+dx)^4}{32a^3d} + \frac{d^2 \int e^{-6e-6fx}(c+dx) dx}{48a^3f^2} \\
&\quad + \frac{(9d^2) \int e^{-4e-4fx}(c+dx) dx}{64a^3f^2} + \frac{(9d^2) \int e^{-2e-2fx}(c+dx) dx}{16a^3f^2} \\
&= -\frac{d^2e^{-6e-6fx}(c+dx)}{288a^3f^3} - \frac{9d^2e^{-4e-4fx}(c+dx)}{256a^3f^3} - \frac{9d^2e^{-2e-2fx}(c+dx)}{32a^3f^3} \\
&\quad - \frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} - \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} \\
&\quad - \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} \\
&\quad + \frac{(c+dx)^4}{32a^3d} + \frac{d^3 \int e^{-6e-6fx} dx}{288a^3f^3} + \frac{(9d^3) \int e^{-4e-4fx} dx}{256a^3f^3} + \frac{(9d^3) \int e^{-2e-2fx} dx}{32a^3f^3} \\
&= -\frac{d^3e^{-6e-6fx}}{1728a^3f^4} - \frac{9d^3e^{-4e-4fx}}{1024a^3f^4} - \frac{9d^3e^{-2e-2fx}}{64a^3f^4} - \frac{d^2e^{-6e-6fx}(c+dx)}{288a^3f^3} \\
&\quad - \frac{9d^2e^{-4e-4fx}(c+dx)}{256a^3f^3} - \frac{9d^2e^{-2e-2fx}(c+dx)}{32a^3f^3} - \frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} \\
&\quad - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} - \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} - \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} \\
&\quad - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} + \frac{(c+dx)^4}{32a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.46 (sec) , antiderivative size = 615, normalized size of antiderivative = 1.83

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\operatorname{sech}^3(e + fx) (-243(32c^3 f^3 + 8c^2 d f^2 (5 + 12fx) + 4cd^2 f (9 + 20fx + 24f^2 x^2) + d^3 (17 + 36fx + 40f^2 x^2) + \dots)}{(27648a^3 f^4 (1 + \tanh(e + fx))^3)}$$

[In] Integrate[(c + d*x)^3/(a + a*Tanh[e + f*x])^3,x]

```
[Out] (Sech[e + f*x]^3*(-243*(32*c^3*f^3 + 8*c^2*d*f^2*(5 + 12*f*x) + 4*c*d^2*f*(9 + 20*f*x + 24*f^2*x^2) + d^3*(17 + 36*f*x + 40*f^2*x^2 + 32*f^3*x^3))*Cos
h[e + f*x] + 16*(36*c^3*f^3*(-1 + 6*f*x) + 18*c^2*d*f^2*(-1 - 6*f*x + 18*f^
2*x^2) + 6*c*d^2*f*(-1 - 6*f*x - 18*f^2*x^2 + 36*f^3*x^3) + d^3*(-1 - 6*f*x
- 18*f^2*x^2 - 36*f^3*x^3 + 54*f^4*x^4))*Cosh[3*(e + f*x)] - 3645*d^3*Sinh
[e + f*x] - 6804*c*d^2*f*Sinh[e + f*x] - 5832*c^2*d*f^2*Sinh[e + f*x] - 259
2*c^3*f^3*Sinh[e + f*x] - 6804*d^3*f*x*Sinh[e + f*x] - 11664*c*d^2*f^2*x*Si
nh[e + f*x] - 7776*c^2*d*f^3*x*Sinh[e + f*x] - 5832*d^3*f^2*x^2*Sinh[e + f*
x] - 7776*c*d^2*f^3*x^2*Sinh[e + f*x] - 2592*d^3*f^3*x^3*Sinh[e + f*x] + 16
*d^3*Sinh[3*(e + f*x)] + 96*c*d^2*f*Sinh[3*(e + f*x)] + 288*c^2*d*f^2*Sinh[
3*(e + f*x)] + 576*c^3*f^3*Sinh[3*(e + f*x)] + 96*d^3*f*x*Sinh[3*(e + f*x)]
+ 576*c*d^2*f^2*x*Sinh[3*(e + f*x)] + 1728*c^2*d*f^3*x*Sinh[3*(e + f*x)] +
3456*c^3*f^4*x*Sinh[3*(e + f*x)] + 288*d^3*f^2*x^2*Sinh[3*(e + f*x)] + 172
8*c*d^2*f^3*x^2*Sinh[3*(e + f*x)] + 5184*c^2*d*f^4*x^2*Sinh[3*(e + f*x)] +
576*d^3*f^3*x^3*Sinh[3*(e + f*x)] + 3456*c*d^2*f^4*x^3*Sinh[3*(e + f*x)] +
864*d^3*f^4*x^4*Sinh[3*(e + f*x)]))/(27648*a^3*f^4*(1 + Tanh[e + f*x])^3)
```

Maple [A] (verified)

Time = 0.33 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.13

method	result
risch	$\frac{d^3 x^4}{32a^3} + \frac{d^2 c x^3}{8a^3} + \frac{3d c^2 x^2}{16a^3} + \frac{c^3 x}{8a^3} + \frac{c^4}{32a^3 d} - \frac{3(4d^3 x^3 f^3 + 12c d^2 f^3 x^2 + 12c^2 d f^3 x + 6d^3 f^2 x^2 + 4c^3 f^3 + 12c d^2 f^2 x + 6c^2 d f^2)}{64a^3 f^4}$
parallelrisc	$-\frac{1952d^3 - 1725 \tanh(fx+e)^2 d^3 - 6264c^2 d f^3 x - 3645 \tanh(fx+e) d^3 - 864 \tanh(fx+e)^2 c^3 f^3 - 2484x \tanh(fx+e) c d^2 f^2 + 2592x d^3}{64a^3 f^4}$

[In] int((d*x+c)^3/(a+a*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)

```
[Out] 1/32/a^3*d^3*x^4+1/8/a^3*d^2*c*x^3+3/16/a^3*d*c^2*x^2+1/8/a^3*c^3*x+1/32/a^
3/d*c^4-3/64*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4
*c^3*f^3+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+3*d^3)/a^3/f^4*exp(
-2*f*x-2*e)-3/1024*(32*d^3*f^3*x^3+96*c*d^2*f^3*x^2+96*c^2*d*f^3*x+24*d^3*f
^2*x^2+32*c^3*f^3+48*c*d^2*f^2*x+24*c^2*d*f^2+12*d^3*f*x+12*c*d^2*f+3*d^3)/
a^3/f^4*exp(-4*f*x-4*e)-1/1728*(36*d^3*f^3*x^3+108*c*d^2*f^3*x^2+108*c^2*d*
```

$$f^3*x+18*d^3*f^2*x^2+36*c^3*f^3+36*c*d^2*f^2*x+18*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+d^3)/a^3/f^4*\exp(-6*f*x-6*e)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 844 vs. 2(298) = 596.

Time = 0.24 (sec) , antiderivative size = 844, normalized size of antiderivative = 2.51

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx$$

$$= \frac{16(54d^3f^4x^4 - 36c^3f^3 - 18c^2df^2 - 6cd^2f + 36(6cd^2f^4 - d^3f^3)x^3 - d^3 + 18(18c^2df^4 - 6cd^2f^3 - d^3f^2))}{a^3}$$

[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")

[Out] 1/27648*(16*(54*d^3*f^4*x^4 - 36*c^3*f^3 - 18*c^2*d*f^2 - 6*c*d^2*f + 36*(6*c*d^2*f^4 - d^3*f^3)*x^3 - d^3 + 18*(18*c^2*d*f^4 - 6*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(36*c^3*f^4 - 18*c^2*d*f^3 - 6*c*d^2*f^2 - d^3*f)*x)*cosh(f*x + e)^3 + 48*(54*d^3*f^4*x^4 - 36*c^3*f^3 - 18*c^2*d*f^2 - 6*c*d^2*f + 36*(6*c*d^2*f^4 - d^3*f^3)*x^3 - d^3 + 18*(18*c^2*d*f^4 - 6*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(36*c^3*f^4 - 18*c^2*d*f^3 - 6*c*d^2*f^2 - d^3*f)*x)*cosh(f*x + e)*sinh(f*x + e)^2 + 16*(54*d^3*f^4*x^4 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f + 36*(6*c*d^2*f^4 + d^3*f^3)*x^3 + d^3 + 18*(18*c^2*d*f^4 + 6*c*d^2*f^3 + d^3*f^2)*x^2 + 6*(36*c^3*f^4 + 18*c^2*d*f^3 + 6*c*d^2*f^2 + d^3*f)*x)*sinh(f*x + e)^3 - 243*(32*d^3*f^3*x^3 + 32*c^3*f^3 + 40*c^2*d*f^2 + 36*c*d^2*f + 17*d^3 + 8*(12*c*d^2*f^3 + 5*d^3*f^2)*x^2 + 4*(24*c^2*d*f^3 + 20*c*d^2*f^2 + 9*d^3*f)*x)*cosh(f*x + e) - 3*(864*d^3*f^3*x^3 + 864*c^3*f^3 + 1944*c^2*d*f^2 + 2268*c*d^2*f + 1215*d^3 + 648*(4*c*d^2*f^3 + 3*d^3*f^2)*x^2 - 16*(54*d^3*f^4*x^4 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f + 36*(6*c*d^2*f^4 + d^3*f^3)*x^3 + d^3 + 18*(18*c^2*d*f^4 + 6*c*d^2*f^3 + d^3*f^2)*x^2 + 6*(36*c^3*f^4 + 18*c^2*d*f^3 + 6*c*d^2*f^2 + d^3*f)*x)*cosh(f*x + e)^2 + 324*(8*c^2*d*f^3 + 12*c*d^2*f^2 + 7*d^3*f)*x)*sinh(f*x + e))/(a^3*f^4*cosh(f*x + e)^3 + 3*a^3*f^4*cosh(f*x + e)^2*sinh(f*x + e) + 3*a^3*f^4*cosh(f*x + e)*sinh(f*x + e)^2 + a^3*f^4*sinh(f*x + e)^3)

Sympy [F]

$$\int \frac{(c+dx)^3}{(a+a \tanh(e+fx))^3} dx$$

$$= \int \frac{c^3}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx + \int \frac{d^3 x^3}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx + \int \frac{d^3 x^3}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx + \int \frac{d^3 x^3}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx$$

[In] integrate((d*x+c)**3/(a+a*tanh(f*x+e))**3,x)

```
[Out] (Integral(c**3/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x) + Integral(d**3*x**3/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x) + Integral(3*c*d**2*x**2/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x) + Integral(3*c**2*d*x/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x))/a**3
```

Maxima [A] (verification not implemented)

none

Time = 1.67 (sec) , antiderivative size = 407, normalized size of antiderivative = 1.21

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{1}{96} c^3 \left(\frac{12(fx + e)}{a^3 f} - \frac{18e^{(-2fx-2e)} + 9e^{(-4fx-4e)} + 2e^{(-6fx-6e)}}{a^3 f} \right)$$

$$+ \frac{(72f^2x^2e^{(6e)} - 108(2fxe^{(4e)} + e^{(4e)})e^{(-2fx)} - 27(4fxe^{(2e)} + e^{(2e)})e^{(-4fx)} - 4(6fx + 1)e^{(-6fx)})c^2d}{384a^3f^2}$$

$$+ \frac{(288f^3x^3e^{(6e)} - 648(2f^2x^2e^{(4e)} + 2fxe^{(4e)} + e^{(4e)})e^{(-2fx)} - 81(8f^2x^2e^{(2e)} + 4fxe^{(2e)} + e^{(2e)})e^{(-4fx)})c^2d}{2304a^3f^3}$$

$$+ \frac{(864f^4x^4e^{(6e)} - 1296(4f^3x^3e^{(4e)} + 6f^2x^2e^{(4e)} + 6fxe^{(4e)} + 3e^{(4e)})e^{(-2fx)} - 81(32f^3x^3e^{(2e)} + 24f^2x^2e^{(2e)} + 12fxe^{(2e)} + 3e^{(2e)})e^{(-4fx)} - 16(36f^3x^3 + 18f^2x^2 + 6fx + 1)e^{(-6fx)})c^2d}{27648a^3f^4}$$

```
[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/96*c^3*(12*(f*x + e)/(a^3*f) - (18*e^(-2*f*x - 2*e) + 9*e^(-4*f*x - 4*e) + 2*e^(-6*f*x - 6*e))/(a^3*f)) + 1/384*(72*f^2*x^2*e^(6*e) - 108*(2*f*x*e^(4*e) + e^(4*e))*e^(-2*f*x) - 27*(4*f*x*e^(2*e) + e^(2*e))*e^(-4*f*x) - 4*(6*f*x + 1)*e^(-6*f*x))*c^2*d*e^(-6*e)/(a^3*f^2) + 1/2304*(288*f^3*x^3*e^(6*e) - 648*(2*f^2*x^2*e^(4*e) + 2*f*x*e^(4*e) + e^(4*e))*e^(-2*f*x) - 81*(8*f^2*x^2*e^(2*e) + 4*f*x*e^(2*e) + e^(2*e))*e^(-4*f*x) - 8*(18*f^2*x^2 + 6*f*x + 1)*e^(-6*f*x))*c*d^2*e^(-6*e)/(a^3*f^3) + 1/27648*(864*f^4*x^4*e^(6*e) - 1296*(4*f^3*x^3*e^(4*e) + 6*f^2*x^2*e^(4*e) + 6*f*x*e^(4*e) + 3*e^(4*e))*e^(-2*f*x) - 81*(32*f^3*x^3*e^(2*e) + 24*f^2*x^2*e^(2*e) + 12*f*x*e^(2*e) + 3*e^(2*e))*e^(-4*f*x) - 16*(36*f^3*x^3 + 18*f^2*x^2 + 6*f*x + 1)*e^(-6*f*x))*d^3*e^(-6*e)/(a^3*f^4)
```

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 548, normalized size of antiderivative = 1.63

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{(864 d^3 f^4 x^4 e^{(6fx+6e)} + 3456 cd^2 f^4 x^3 e^{(6fx+6e)} + 5184 c^2 d f^4 x^2 e^{(6fx+6e)} - 5184 d^3 f^3 x^3 e^{(4fx+4e)} - 2592 d^3 f^3}{}$$

`[In] integrate((d*x+c)^3/(a+a*tanh(f*x+e))^3,x, algorithm="giac")`

```
[Out] 1/27648*(864*d^3*f^4*x^4*e^(6*f*x + 6*e) + 3456*c*d^2*f^4*x^3*e^(6*f*x + 6*
e) + 5184*c^2*d*f^4*x^2*e^(6*f*x + 6*e) - 5184*d^3*f^3*x^3*e^(4*f*x + 4*e)
- 2592*d^3*f^3*x^3*e^(2*f*x + 2*e) - 576*d^3*f^3*x^3 + 3456*c^3*f^4*x*e^(6*
f*x + 6*e) - 15552*c*d^2*f^3*x^2*e^(4*f*x + 4*e) - 7776*c*d^2*f^3*x^2*e^(2*
f*x + 2*e) - 1728*c*d^2*f^3*x^2 - 15552*c^2*d*f^3*x*e^(4*f*x + 4*e) - 7776*
d^3*f^2*x^2*e^(4*f*x + 4*e) - 7776*c^2*d*f^3*x*e^(2*f*x + 2*e) - 1944*d^3*f
^2*x^2*e^(2*f*x + 2*e) - 1728*c^2*d*f^3*x - 288*d^3*f^2*x^2 - 5184*c^3*f^3*
e^(4*f*x + 4*e) - 15552*c*d^2*f^2*x*e^(4*f*x + 4*e) - 2592*c^3*f^3*e^(2*f*x
+ 2*e) - 3888*c*d^2*f^2*x*e^(2*f*x + 2*e) - 576*c^3*f^3 - 576*c*d^2*f^2*x
- 7776*c^2*d*f^2*e^(4*f*x + 4*e) - 7776*d^3*f*x*e^(4*f*x + 4*e) - 1944*c^2*
d*f^2*e^(2*f*x + 2*e) - 972*d^3*f*x*e^(2*f*x + 2*e) - 288*c^2*d*f^2 - 96*d^
3*f*x - 7776*c*d^2*f*e^(4*f*x + 4*e) - 972*c*d^2*f*e^(2*f*x + 2*e) - 96*c*d
^2*f - 3888*d^3*e^(4*f*x + 4*e) - 243*d^3*e^(2*f*x + 2*e) - 16*d^3)*e^(-6*f
*x - 6*e)/(a^3*f^4)
```

Mupad [B] (verification not implemented)

Time = 2.03 (sec) , antiderivative size = 376, normalized size of antiderivative = 1.12

$$\int \frac{(c + dx)^3}{(a + a \tanh(e + fx))^3} dx = \frac{c^3 x}{8a^3} - e^{-4e-4fx} \left(\frac{96c^3 f^3 + 72c^2 d f^2 + 36cd^2 f + 9d^3}{1024a^3 f^4} \right)$$

$$+ \frac{3d^3 x^3}{32a^3 f} + \frac{9dx(8c^2 f^2 + 4cdf + d^2)}{256a^3 f^3} + \frac{9d^2 x^2 (d + 4cf)}{128a^3 f^2}$$

$$- e^{-6e-6fx} \left(\frac{36c^3 f^3 + 18c^2 d f^2 + 6cd^2 f + d^3}{1728a^3 f^4} + \frac{d^3 x^3}{48a^3 f} \right)$$

$$+ \frac{dx(18c^2 f^2 + 6cdf + d^2)}{288a^3 f^3} + \frac{d^2 x^2 (d + 6cf)}{96a^3 f^2}$$

$$- e^{-2e-2fx} \left(\frac{12c^3 f^3 + 18c^2 d f^2 + 18cd^2 f + 9d^3}{64a^3 f^4} + \frac{3d^3 x^3}{16a^3 f} \right)$$

$$+ \frac{9dx(2c^2 f^2 + 2cdf + d^2)}{32a^3 f^3} + \frac{9d^2 x^2 (d + 2cf)}{32a^3 f^2}$$

$$+ \frac{d^3 x^4}{32a^3} + \frac{3c^2 d x^2}{16a^3} + \frac{cd^2 x^3}{8a^3}$$

[In] $\text{int}((c + d*x)^3/(a + a*\tanh(e + f*x))^3,x)$

[Out] $(c^3*x)/(8*a^3) - \exp(-4*e - 4*f*x)*((9*d^3 + 96*c^3*f^3 + 72*c^2*d*f^2 + 36*c*d^2*f)/(1024*a^3*f^4) + (3*d^3*x^3)/(32*a^3*f) + (9*d*x*(d^2 + 8*c^2*f^2 + 4*c*d*f))/(256*a^3*f^3) + (9*d^2*x^2*(d + 4*c*f))/(128*a^3*f^2)) - \exp(-6*e - 6*f*x)*((d^3 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f)/(1728*a^3*f^4) + (d^3*x^3)/(48*a^3*f) + (d*x*(d^2 + 18*c^2*f^2 + 6*c*d*f))/(288*a^3*f^3) + (d^2*x^2*(d + 6*c*f))/(96*a^3*f^2)) - \exp(-2*e - 2*f*x)*((9*d^3 + 12*c^3*f^3 + 18*c^2*d*f^2 + 18*c*d^2*f)/(64*a^3*f^4) + (3*d^3*x^3)/(16*a^3*f) + (9*d*x*(d^2 + 2*c^2*f^2 + 2*c*d*f))/(32*a^3*f^3) + (9*d^2*x^2*(d + 2*c*f))/(32*a^3*f^2) + (d^3*x^4)/(32*a^3) + (3*c^2*d*x^2)/(16*a^3) + (c*d^2*x^3)/(8*a^3))$

$$3.44 \quad \int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx$$

Optimal result	304
Rubi [A] (verified)	304
Mathematica [A] (verified)	306
Maple [A] (verified)	307
Fricas [B] (verification not implemented)	307
Sympy [F]	308
Maxima [A] (verification not implemented)	308
Giac [A] (verification not implemented)	309
Mupad [B] (verification not implemented)	309

Optimal result

Integrand size = 20, antiderivative size = 246

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx = -\frac{d^2 e^{-6e-6fx}}{864a^3 f^3} - \frac{3d^2 e^{-4e-4fx}}{256a^3 f^3} - \frac{3d^2 e^{-2e-2fx}}{32a^3 f^3} - \frac{de^{-6e-6fx}(c+dx)}{144a^3 f^2} - \frac{3de^{-4e-4fx}(c+dx)}{64a^3 f^2} - \frac{3de^{-2e-2fx}(c+dx)}{16a^3 f^2} - \frac{e^{-6e-6fx}(c+dx)^2}{48a^3 f} - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3 f} - \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3 f} + \frac{(c+dx)^3}{24a^3 d}$$

[Out] -1/864*d^2*exp(-6*f*x-6*e)/a^3/f^3-3/256*d^2*exp(-4*f*x-4*e)/a^3/f^3-3/32*d^2*exp(-2*f*x-2*e)/a^3/f^3-1/144*d*exp(-6*f*x-6*e)*(d*x+c)/a^3/f^2-3/64*d*exp(-4*f*x-4*e)*(d*x+c)/a^3/f^2-3/16*d*exp(-2*f*x-2*e)*(d*x+c)/a^3/f^2-1/48*exp(-6*f*x-6*e)*(d*x+c)^2/a^3/f-3/32*exp(-4*f*x-4*e)*(d*x+c)^2/a^3/f-3/16*exp(-2*f*x-2*e)*(d*x+c)^2/a^3/f+1/24*(d*x+c)^3/a^3/d

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used

= {3810, 2207, 2225}

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx = -\frac{d(c+dx)e^{-6e-6fx}}{144a^3f^2} - \frac{3d(c+dx)e^{-4e-4fx}}{64a^3f^2} - \frac{3d(c+dx)e^{-2e-2fx}}{16a^3f^2} - \frac{(c+dx)^2e^{-6e-6fx}}{48a^3f} - \frac{3(c+dx)^2e^{-4e-4fx}}{32a^3f} - \frac{3(c+dx)^2e^{-2e-2fx}}{16a^3f} + \frac{(c+dx)^3}{24a^3d} - \frac{d^2e^{-6e-6fx}}{864a^3f^3} - \frac{3d^2e^{-4e-4fx}}{256a^3f^3} - \frac{3d^2e^{-2e-2fx}}{32a^3f^3}$$

[In] Int[(c + d*x)^2/(a + a*Tanh[e + f*x])^3,x]

[Out] -1/864*(d^2*E^(-6*e - 6*f*x))/(a^3*f^3) - (3*d^2*E^(-4*e - 4*f*x))/(256*a^3*f^3) - (3*d^2*E^(-2*e - 2*f*x))/(32*a^3*f^3) - (d*E^(-6*e - 6*f*x)*(c + d*x))/(144*a^3*f^2) - (3*d*E^(-4*e - 4*f*x)*(c + d*x))/(64*a^3*f^2) - (3*d*E^(-2*e - 2*f*x)*(c + d*x))/(16*a^3*f^2) - (E^(-6*e - 6*f*x)*(c + d*x)^2)/(48*a^3*f) - (3*E^(-4*e - 4*f*x)*(c + d*x)^2)/(32*a^3*f) - (3*E^(-2*e - 2*f*x)*(c + d*x)^2)/(16*a^3*f) + (c + d*x)^3/(24*a^3*d)

Rule 2207

Int[((b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[\$UseGamma]

Rule 2225

Int[((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 3810

Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a)]^(n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^2}{8a^3} + \frac{e^{-6e-6fx}(c+dx)^2}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^2}{8a^3} + \frac{3e^{-2e-2fx}(c+dx)^2}{8a^3} \right) dx \\ &= \frac{(c+dx)^3}{24a^3d} + \frac{\int e^{-6e-6fx}(c+dx)^2 dx}{8a^3} + \frac{3 \int e^{-4e-4fx}(c+dx)^2 dx}{8a^3} + \frac{3 \int e^{-2e-2fx}(c+dx)^2 dx}{8a^3} \end{aligned}$$

$$\begin{aligned}
&= -\frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} \\
&+ \frac{(c+dx)^3}{24a^3d} + \frac{d \int e^{-6e-6fx}(c+dx) dx}{24a^3f} + \frac{(3d) \int e^{-4e-4fx}(c+dx) dx}{16a^3f} \\
&+ \frac{(3d) \int e^{-2e-2fx}(c+dx) dx}{8a^3f} \\
&= -\frac{de^{-6e-6fx}(c+dx)}{144a^3f^2} - \frac{3de^{-4e-4fx}(c+dx)}{64a^3f^2} - \frac{3de^{-2e-2fx}(c+dx)}{16a^3f^2} \\
&- \frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} \\
&+ \frac{(c+dx)^3}{24a^3d} + \frac{d^2 \int e^{-6e-6fx} dx}{144a^3f^2} + \frac{(3d^2) \int e^{-4e-4fx} dx}{64a^3f^2} + \frac{(3d^2) \int e^{-2e-2fx} dx}{16a^3f^2} \\
&= -\frac{d^2e^{-6e-6fx}}{864a^3f^3} - \frac{3d^2e^{-4e-4fx}}{256a^3f^3} - \frac{3d^2e^{-2e-2fx}}{32a^3f^3} - \frac{de^{-6e-6fx}(c+dx)}{144a^3f^2} \\
&- \frac{3de^{-4e-4fx}(c+dx)}{64a^3f^2} - \frac{3de^{-2e-2fx}(c+dx)}{16a^3f^2} - \frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} \\
&- \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} - \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} + \frac{(c+dx)^3}{24a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.12 (sec) , antiderivative size = 371, normalized size of antiderivative = 1.51

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx$$

$$= \frac{\operatorname{sech}^3(e+fx)(-81(24c^2f^2+4cdf(5+12fx)+d^2(9+20fx+24f^2x^2)) \cosh(e+fx)+8(18c^2f^2(-1+6fx)+6c^2d^2f^2x^2+6cd^2f^2x^3)+d^2(-1-6fx-18f^2x^2+36f^3x^3)) \cosh[3(e+fx)]-567d^2 \operatorname{Sinh}[e+fx]-972c^2d^2f^2 \operatorname{Sinh}[e+fx]-648c^2d^2f^2 \operatorname{Sinh}[e+fx]-972d^2f^2x \operatorname{Sinh}[e+fx]-1296c^2d^2f^2x \operatorname{Sinh}[e+fx]-648d^2f^2x^2 \operatorname{Sinh}[e+fx]+8d^2 \operatorname{Sinh}[3(e+fx)]+48c^2d^2f^2 \operatorname{Sinh}[3(e+fx)]+144c^2d^2f^2 \operatorname{Sinh}[3(e+fx)]+48d^2f^2x \operatorname{Sinh}[3(e+fx)]+288c^2d^2f^2x \operatorname{Sinh}[3(e+fx)]+864c^2d^2f^3x^2 \operatorname{Sinh}[3(e+fx)]+144d^2f^2x^2 \operatorname{Sinh}[3(e+fx)]+864c^2d^2f^3x^2 \operatorname{Sinh}[3(e+fx)]+288d^2f^3x^3 \operatorname{Sinh}[3(e+fx)])}{(6912a^3f^3(1+\tanh(e+fx))^3)}$$

[In] Integrate[(c + d*x)^2/(a + a*Tanh[e + f*x])^3,x]

[Out] (Sech[e + f*x]^3*(-81*(24*c^2*f^2 + 4*c*d*f*(5 + 12*f*x) + d^2*(9 + 20*f*x + 24*f^2*x^2))*Cosh[e + f*x] + 8*(18*c^2*f^2*(-1 + 6*f*x) + 6*c*d*f*(-1 - 6*f*x + 18*f^2*x^2) + d^2*(-1 - 6*f*x - 18*f^2*x^2 + 36*f^3*x^3))*Cosh[3*(e + f*x)] - 567*d^2*Sinh[e + f*x] - 972*c*d*f*Sinh[e + f*x] - 648*c^2*f^2*Sinh[e + f*x] - 972*d^2*f*x*Sinh[e + f*x] - 1296*c*d*f^2*x*Sinh[e + f*x] - 648*d^2*f^2*x^2*Sinh[e + f*x] + 8*d^2*Sinh[3*(e + f*x)] + 48*c*d*f*Sinh[3*(e + f*x)] + 144*c^2*f^2*Sinh[3*(e + f*x)] + 48*d^2*f*x*Sinh[3*(e + f*x)] + 288*c*d*f^2*x*Sinh[3*(e + f*x)] + 864*c^2*f^3*x^2*Sinh[3*(e + f*x)] + 144*d^2*f^2*x^2*Sinh[3*(e + f*x)] + 864*c*d*f^3*x^2*Sinh[3*(e + f*x)] + 288*d^2*f^3*x^3*Sinh[3*(e + f*x)]))/(6912*a^3*f^3*(1 + Tanh[e + f*x])^3)

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.91

method	result
risch	$\frac{d^2x^3}{24a^3} + \frac{dcx^2}{8a^3} + \frac{c^2x}{8a^3} + \frac{c^3}{24a^3d} - \frac{3(2d^2x^2f^2+4cdf^2x+2c^2f^2+2d^2fx+2cdf+d^2)e^{-2fx-2e}}{32a^3f^3} - \frac{3(8d^2x^2f^2+16cdf^2x+8c^2f^2+2d^2fx+2cdf+d^2)e^{-2fx-2e}}{32a^3f^3}$
parallelrisch	$-\frac{328d^2-417d^2fx+255x \tanh(fx+e)^3d^2f+198x^2 \tanh(fx+e)^3d^2f^2-396 \tanh(fx+e)^2cdf+72d^2 \tanh(fx+e)^3x^3f^3-567 \tanh(fx+e)^3x^3f^3}{32a^3f^3}$

[In] int((d*x+c)^2/(a+a*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{24} \frac{d^2x^3}{a^3} + \frac{1}{8} \frac{dcx^2}{a^3} + \frac{1}{8} \frac{c^2x}{a^3} + \frac{1}{24} \frac{c^3}{a^3d} - \frac{3(2d^2x^2f^2+4cdf^2x+2c^2f^2+2d^2fx+2cdf+d^2)e^{-2fx-2e}}{32a^3f^3} - \frac{3(8d^2x^2f^2+16cdf^2x+8c^2f^2+2d^2fx+2cdf+d^2)e^{-2fx-2e}}{32a^3f^3}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 532 vs. 2(217) = 434.

Time = 0.25 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.16

$$\int \frac{(c+dx)^2}{(a+a \tanh(e+fx))^3} dx = \frac{8(36d^2f^3x^3 - 18c^2f^2 - 6cdf + 18(6cdf^3 - d^2f^2)x^2 - d^2 + 6(18c^2f^3 - 6cdf^2 - d^2f)x) \cosh(fx+e)^3}{(a^3f^3 \cosh(fx+e)^3 + 3a^3f^3 \cosh(fx+e)^2 \sinh(fx+e) + 3a^3f^3 \cosh(fx+e) \sinh(fx+e)^2 + a^3f^3 \sinh(fx+e)^3)}$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")

[Out] $\frac{1}{6912} (8*(36*d^2*f^3*x^3 - 18*c^2*f^2 - 6*c*d*f + 18*(6*c*d*f^3 - d^2*f^2)*x^2 - d^2 + 6*(18*c^2*f^3 - 6*c*d*f^2 - d^2*f)*x)*\cosh(f*x + e)^3 + 24*(36*d^2*f^3*x^3 - 18*c^2*f^2 - 6*c*d*f + 18*(6*c*d*f^3 - d^2*f^2)*x^2 - d^2 + 6*(18*c^2*f^3 - 6*c*d*f^2 - d^2*f)*x)*\cosh(f*x + e)*\sinh(f*x + e)^2 + 8*(36*d^2*f^3*x^3 + 18*c^2*f^2 + 6*c*d*f + 18*(6*c*d*f^3 + d^2*f^2)*x^2 + d^2 + 6*(18*c^2*f^3 + 6*c*d*f^2 + d^2*f)*x)*\sinh(f*x + e)^3 - 81*(24*d^2*f^2*x^2 + 24*c^2*f^2 + 20*c*d*f + 9*d^2 + 4*(12*c*d*f^2 + 5*d^2*f)*x)*\cosh(f*x + e) - 3*(216*d^2*f^2*x^2 + 216*c^2*f^2 + 324*c*d*f - 8*(36*d^2*f^3*x^3 + 18*c^2*f^2 + 6*c*d*f + 18*(6*c*d*f^3 + d^2*f^2)*x^2 + d^2 + 6*(18*c^2*f^3 + 6*c*d*f^2 + d^2*f)*x)*\cosh(f*x + e)^2 + 189*d^2 + 108*(4*c*d*f^2 + 3*d^2*f)*x)*\sinh(f*x + e))/(a^3*f^3*\cosh(f*x + e)^3 + 3*a^3*f^3*\cosh(f*x + e)^2*\sinh(f*x + e) + 3*a^3*f^3*\cosh(f*x + e)*\sinh(f*x + e)^2 + a^3*f^3*\sinh(f*x + e)^3)$

SymPy [F]

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\int \frac{c^2}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx + \int \frac{d^2 x^2}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx + \int \frac{2cdx}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx}{a^3}$$

```
[In] integrate((d*x+c)**2/(a+a*tanh(f*x+e))**3,x)
```

```
[Out] (Integral(c**2/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x) + Integral(d**2*x**2/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x) + Integral(2*c*d*x/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x))/a**3
```

Maxima [A] (verification not implemented)

none

Time = 1.19 (sec) , antiderivative size = 255, normalized size of antiderivative = 1.04

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{1}{96} c^2 \left(\frac{12(fx + e)}{a^3 f} - \frac{18e^{(-2fx-2e)} + 9e^{(-4fx-4e)} + 2e^{(-6fx-6e)}}{a^3 f} \right)$$

$$+ \frac{(72f^2x^2e^{(6e)} - 108(2fxe^{(4e)} + e^{(4e)})e^{(-2fx)} - 27(4fxe^{(2e)} + e^{(2e)})e^{(-4fx)} - 4(6fx + 1)e^{(-6fx)})cde^{(-6e)}}{576a^3f^2}$$

$$+ \frac{(288f^3x^3e^{(6e)} - 648(2f^2x^2e^{(4e)} + 2fxe^{(4e)} + e^{(4e)})e^{(-2fx)} - 81(8f^2x^2e^{(2e)} + 4fxe^{(2e)} + e^{(2e)})e^{(-4fx)})}{6912a^3f^3}$$

```
[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/96*c^2*(12*(f*x + e)/(a^3*f) - (18*e^(-2*f*x - 2*e) + 9*e^(-4*f*x - 4*e) + 2*e^(-6*f*x - 6*e))/(a^3*f)) + 1/576*(72*f^2*x^2*e^(6*e) - 108*(2*f*x*e^(4*e) + e^(4*e))*e^(-2*f*x) - 27*(4*f*x*e^(2*e) + e^(2*e))*e^(-4*f*x) - 4*(6*f*x + 1)*e^(-6*f*x))*c*d*e^(-6*e)/(a^3*f^2) + 1/6912*(288*f^3*x^3*e^(6*e) - 648*(2*f^2*x^2*e^(4*e) + 2*f*x*e^(4*e) + e^(4*e))*e^(-2*f*x) - 81*(8*f^2*x^2*e^(2*e) + 4*f*x*e^(2*e) + e^(2*e))*e^(-4*f*x) - 8*(18*f^2*x^2 + 6*f*x + 1)*e^(-6*f*x))*d^2*e^(-6*e)/(a^3*f^3)
```

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 315, normalized size of antiderivative = 1.28

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^3} dx$$

$$(288 d^2 f^3 x^3 e^{(6fx+6e)} + 864 cdf^3 x^2 e^{(6fx+6e)} + 864 c^2 f^3 x e^{(6fx+6e)} - 1296 d^2 f^2 x^2 e^{(4fx+4e)} - 648 d^2 f^2 x^2 e^{(2fx+2e)} - 144 d^2 f^2 x^2 - 2592 c d f^2 x e^{(4fx+4e)} - 1296 c d f^2 x e^{(2fx+2e)} - 288 c d f^2 x - 1296 c^2 f^2 e^{(4fx+4e)} - 1296 d^2 f^2 x e^{(4fx+4e)} - 648 c^2 f^2 e^{(2fx+2e)} - 324 d^2 f^2 x e^{(2fx+2e)} - 144 c^2 f^2 - 48 d^2 f^2 x - 1296 c d f e^{(4fx+4e)} - 324 c d f e^{(2fx+2e)} - 48 c d f - 648 d^2 e^{(4fx+4e)} - 81 d^2 e^{(2fx+2e)} - 8 d^2) e^{(-6fx-6e)} / (a^3 f^3)$$

[In] integrate((d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="giac")

[Out] 1/6912*(288*d^2*f^3*x^3*e^(6*f*x + 6*e) + 864*c*d*f^3*x^2*e^(6*f*x + 6*e) + 864*c^2*f^3*x*e^(6*f*x + 6*e) - 1296*d^2*f^2*x^2*e^(4*f*x + 4*e) - 648*d^2*f^2*x^2*e^(2*f*x + 2*e) - 144*d^2*f^2*x^2 - 2592*c*d*f^2*x*e^(4*f*x + 4*e) - 1296*c*d*f^2*x*e^(2*f*x + 2*e) - 288*c*d*f^2*x - 1296*c^2*f^2*e^(4*f*x + 4*e) - 1296*d^2*f^2*x*e^(4*f*x + 4*e) - 648*c^2*f^2*e^(2*f*x + 2*e) - 324*d^2*f^2*x*e^(2*f*x + 2*e) - 144*c^2*f^2 - 48*d^2*f^2*x - 1296*c*d*f*e^(4*f*x + 4*e) - 324*c*d*f*e^(2*f*x + 2*e) - 48*c*d*f - 648*d^2*e^(4*f*x + 4*e) - 81*d^2*e^(2*f*x + 2*e) - 8*d^2)*e^(-6*f*x - 6*e)/(a^3*f^3)

Mupad [B] (verification not implemented)

Time = 1.93 (sec) , antiderivative size = 236, normalized size of antiderivative = 0.96

$$\int \frac{(c + dx)^2}{(a + a \tanh(e + fx))^3} dx = \frac{c^2 x}{8 a^3} - e^{-2e-2fx} \left(\frac{6c^2 f^2 + 6cdf + 3d^2}{32a^3 f^3} + \frac{3d^2 x^2}{16a^3 f} + \frac{3dx(d+2cf)}{16a^3 f^2} \right) - e^{-4e-4fx} \left(\frac{24c^2 f^2 + 12cdf + 3d^2}{256a^3 f^3} + \frac{3d^2 x^2}{32a^3 f} + \frac{3dx(d+4cf)}{64a^3 f^2} \right) - e^{-6e-6fx} \left(\frac{18c^2 f^2 + 6cdf + d^2}{864a^3 f^3} + \frac{d^2 x^2}{48a^3 f} + \frac{dx(d+6cf)}{144a^3 f^2} \right) + \frac{d^2 x^3}{24a^3} + \frac{cdx^2}{8a^3}$$

[In] int((c + d*x)^2/(a + a*tanh(e + f*x))^3,x)

[Out] (c^2*x)/(8*a^3) - exp(- 2*e - 2*f*x)*((3*d^2 + 6*c^2*f^2 + 6*c*d*f)/(32*a^3*f^3) + (3*d^2*x^2)/(16*a^3*f) + (3*d*x*(d + 2*c*f))/(16*a^3*f^2)) - exp(- 4*e - 4*f*x)*((3*d^2 + 24*c^2*f^2 + 12*c*d*f)/(256*a^3*f^3) + (3*d^2*x^2)/(32*a^3*f) + (3*d*x*(d + 4*c*f))/(64*a^3*f^2)) - exp(- 6*e - 6*f*x)*((d^2 + 18*c^2*f^2 + 6*c*d*f)/(864*a^3*f^3) + (d^2*x^2)/(48*a^3*f) + (d*x*(d + 6*c*f))/(144*a^3*f^2)) + (d^2*x^3)/(24*a^3) + (c*d*x^2)/(8*a^3)

3.45 $\int \frac{c+dx}{(a+a \tanh(e+fx))^3} dx$

Optimal result	310
Rubi [A] (verified)	310
Mathematica [A] (verified)	313
Maple [A] (verified)	313
Fricas [A] (verification not implemented)	313
Sympy [F]	314
Maxima [A] (verification not implemented)	314
Giac [A] (verification not implemented)	315
Mupad [B] (verification not implemented)	315

Optimal result

Integrand size = 18, antiderivative size = 183

$$\int \frac{c+dx}{(a+a \tanh(e+fx))^3} dx = \frac{11dx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \tanh(e+fx))^3}$$

$$- \frac{c+dx}{6f(a+a \tanh(e+fx))^3} - \frac{5d}{96af^2(a+a \tanh(e+fx))^2}$$

$$- \frac{c+dx}{8af(a+a \tanh(e+fx))^2} - \frac{11d}{96f^2(a^3+a^3 \tanh(e+fx))}$$

$$- \frac{c+dx}{8f(a^3+a^3 \tanh(e+fx))}$$

[Out] 11/96*d*x/a^3/f-1/16*d*x^2/a^3+1/8*x*(d*x+c)/a^3-1/36*d/f^2/(a+a*tanh(f*x+e))^3+1/6*(-d*x-c)/f/(a+a*tanh(f*x+e))^3-5/96*d/a/f^2/(a+a*tanh(f*x+e))^2+1/8*(-d*x-c)/a/f/(a+a*tanh(f*x+e))^2-11/96*d/f^2/(a^3+a^3*tanh(f*x+e))+1/8*(-d*x-c)/f/(a^3+a^3*tanh(f*x+e))

Rubi [A] (verified)

Time = 0.16 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used

= {3560, 8, 3811}

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx = -\frac{c + dx}{8f(a^3 \tanh(e + fx) + a^3)} + \frac{x(c + dx)}{8a^3} - \frac{11d}{96f^2(a^3 \tanh(e + fx) + a^3)} + \frac{11dx}{96a^3f} - \frac{dx^2}{16a^3} - \frac{c + dx}{8af(a \tanh(e + fx) + a)^2} - \frac{c + dx}{6f(a \tanh(e + fx) + a)^3} - \frac{5d}{96af^2(a \tanh(e + fx) + a)^2} - \frac{d}{36f^2(a \tanh(e + fx) + a)^3}$$

[In] Int[(c + d*x)/(a + a*Tanh[e + f*x])^3,x]

[Out] (11*d*x)/(96*a^3*f) - (d*x^2)/(16*a^3) + (x*(c + d*x))/(8*a^3) - d/(36*f^2*(a + a*Tanh[e + f*x])^3) - (c + d*x)/(6*f*(a + a*Tanh[e + f*x])^3) - (5*d)/(96*a*f^2*(a + a*Tanh[e + f*x])^2) - (c + d*x)/(8*a*f*(a + a*Tanh[e + f*x])^2) - (11*d)/(96*f^2*(a^3 + a^3*Tanh[e + f*x])) - (c + d*x)/(8*f*(a^3 + a^3*Tanh[e + f*x]))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3811

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Dist[(c + d*x)^m, u, x] - Dist[d*m, Int[Dist[(c + d*x)^(m - 1), u, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]

Rubi steps

$$\text{integral} = \frac{x(c + dx)}{8a^3} - \frac{c + dx}{6f(a + a \tanh(e + fx))^3} - \frac{c + dx}{8af(a + a \tanh(e + fx))^2} - \frac{c + dx}{8f(a^3 + a^3 \tanh(e + fx))} - d \int \left(\frac{x}{8a^3} - \frac{1}{6f(a + a \tanh(e + fx))^3} - \frac{1}{8af(a + a \tanh(e + fx))^2} - \frac{1}{8f(a^3 + a^3 \tanh(e + fx))} \right) dx$$

$$\begin{aligned}
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{c+dx}{6f(a+a \tanh(e+fx))^3} \\
&\quad - \frac{c+dx}{8af(a+a \tanh(e+fx))^2} - \frac{8f(a^3+a^3 \tanh(e+fx))}{c+dx} \\
&\quad + \frac{d \int \frac{1}{a^3+a^3 \tanh(e+fx)} dx}{8f} + \frac{d \int \frac{1}{(a+a \tanh(e+fx))^3} dx}{6f} + \frac{d \int \frac{1}{(a+a \tanh(e+fx))^2} dx}{8af} \\
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \tanh(e+fx))^3} - \frac{c+dx}{6f(a+a \tanh(e+fx))^3} \\
&\quad - \frac{32af^2(a+a \tanh(e+fx))^2}{d} - \frac{8af(a+a \tanh(e+fx))^2}{c+dx} - \frac{16f^2(a^3+a^3 \tanh(e+fx))}{d} \\
&\quad - \frac{c+dx}{8f(a^3+a^3 \tanh(e+fx))} + \frac{d \int 1 dx}{16a^3 f} + \frac{d \int \frac{1}{a+a \tanh(e+fx)} dx}{16a^2 f} \\
&\quad + \frac{d \int \frac{1}{(a+a \tanh(e+fx))^2} dx}{12af} \\
&= \frac{dx}{16a^3 f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \tanh(e+fx))^3} \\
&\quad - \frac{c+dx}{6f(a+a \tanh(e+fx))^3} - \frac{96af^2(a+a \tanh(e+fx))^2}{5d} \\
&\quad - \frac{8af(a+a \tanh(e+fx))^2}{c+dx} - \frac{32f^2(a^3+a^3 \tanh(e+fx))}{3d} \\
&\quad - \frac{c+dx}{8f(a^3+a^3 \tanh(e+fx))} + \frac{d \int 1 dx}{32a^3 f} + \frac{d \int \frac{1}{a+a \tanh(e+fx)} dx}{24a^2 f} \\
&= \frac{3dx}{32a^3 f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \tanh(e+fx))^3} - \frac{c+dx}{6f(a+a \tanh(e+fx))^3} \\
&\quad - \frac{96af^2(a+a \tanh(e+fx))^2}{5d} - \frac{8af(a+a \tanh(e+fx))^2}{c+dx} \\
&\quad - \frac{11d}{96f^2(a^3+a^3 \tanh(e+fx))} - \frac{c+dx}{8f(a^3+a^3 \tanh(e+fx))} + \frac{d \int 1 dx}{48a^3 f} \\
&= \frac{11dx}{96a^3 f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \tanh(e+fx))^3} - \frac{c+dx}{6f(a+a \tanh(e+fx))^3} \\
&\quad - \frac{96af^2(a+a \tanh(e+fx))^2}{5d} - \frac{8af(a+a \tanh(e+fx))^2}{c+dx} \\
&\quad - \frac{11d}{96f^2(a^3+a^3 \tanh(e+fx))} - \frac{c+dx}{8f(a^3+a^3 \tanh(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.21 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.01

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\operatorname{sech}^3(e + fx) (-27(12cf + d(5 + 12fx)) \cosh(e + fx) + 4(6cf(-1 + 6fx) + d(-1 - 6fx + 18f^2x^2)) \cosh(3(e + fx)) - 81d \operatorname{Sinh}[e + fx] - 108cf \operatorname{Sinh}[e + fx] - 108dfx \operatorname{Sinh}[e + fx] + 4d \operatorname{Sinh}[3(e + fx)] + 24cf \operatorname{Sinh}[3(e + fx)] + 24dfx \operatorname{Sinh}[3(e + fx)] + 144cf^2x \operatorname{Sinh}[3(e + fx)] + 72df^2x^2 \operatorname{Sinh}[3(e + fx)])}{(1152a^3f^2(1 + \tanh(e + fx))^3)}$$

[In] Integrate[(c + d*x)/(a + a*Tanh[e + f*x])^3,x]

```
[Out] (Sech[e + f*x]^3*(-27*(12*c*f + d*(5 + 12*f*x))*Cosh[e + f*x] + 4*(6*c*f*(-1 + 6*f*x) + d*(-1 - 6*f*x + 18*f^2*x^2))*Cosh[3*(e + f*x)] - 81*d*Sinh[e + f*x] - 108*c*f*Sinh[e + f*x] - 108*d*f*x*Sinh[e + f*x] + 4*d*Sinh[3*(e + f*x)] + 24*c*f*Sinh[3*(e + f*x)] + 24*d*f*x*Sinh[3*(e + f*x)] + 144*c*f^2*x*Sinh[3*(e + f*x)] + 72*d*f^2*x^2*Sinh[3*(e + f*x)]))/(1152*a^3*f^2*(1 + Tanh[e + f*x])^3)
```

Maple [A] (verified)

Time = 0.28 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.56

method	result
risch	$\frac{dx^2}{16a^3} + \frac{cx}{8a^3} - \frac{3(2dxf+2cf+d)e^{-2fx-2e}}{32a^3f^2} - \frac{3(4dxf+4cf+d)e^{-4fx-4e}}{128a^3f^2} - \frac{(6dxf+6cf+d)e^{-6fx-6e}}{288a^3f^2}$
parallelrisch	$-81d \tanh(fx+e) - 56d - 33 \tanh(fx+e)^2 d + 36x \tanh(fx+e)^3 c f^2 + 33x \tanh(fx+e)^3 d f + 18d \tanh(fx+e)^3 x^2 f^2 + 36cx f^2 + 18d \tanh(fx+e)^3 x^2 f^2 + 36cx f^2 + 18d \tanh(fx+e)^3 x^2 f^2$

[In] int((d*x+c)/(a+a*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)

```
[Out] 1/16*d*x^2/a^3+1/8/a^3*c*x-3/32*(2*d*f*x+2*c*f+d)/a^3/f^2*exp(-2*f*x-2*e)-3/128*(4*d*f*x+4*c*f+d)/a^3/f^2*exp(-4*f*x-4*e)-1/288*(6*d*f*x+6*c*f+d)/a^3/f^2*exp(-6*f*x-6*e)
```

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.56

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{4(18df^2x^2 - 6cf + 6(6cf^2 - df)x - d) \cosh(fx + e)^3 + 12(18df^2x^2 - 6cf + 6(6cf^2 - df)x - d) \cosh(3(e + fx)) - 81d \operatorname{Sinh}[e + fx] - 108cf \operatorname{Sinh}[e + fx] - 108dfx \operatorname{Sinh}[e + fx] + 4d \operatorname{Sinh}[3(e + fx)] + 24cf \operatorname{Sinh}[3(e + fx)] + 24dfx \operatorname{Sinh}[3(e + fx)] + 144cf^2x \operatorname{Sinh}[3(e + fx)] + 72df^2x^2 \operatorname{Sinh}[3(e + fx)]}{(1152a^3f^2(1 + \tanh(e + fx))^3)}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")

```
[Out] 1/1152*(4*(18*d*f^2*x^2 - 6*c*f + 6*(6*c*f^2 - d*f)*x - d)*cosh(f*x + e)^3
+ 12*(18*d*f^2*x^2 - 6*c*f + 6*(6*c*f^2 - d*f)*x - d)*cosh(f*x + e)*sinh(f*
x + e)^2 + 4*(18*d*f^2*x^2 + 6*c*f + 6*(6*c*f^2 + d*f)*x + d)*sinh(f*x + e)
^3 - 27*(12*d*f*x + 12*c*f + 5*d)*cosh(f*x + e) - 3*(36*d*f*x - 4*(18*d*f^2
*x^2 + 6*c*f + 6*(6*c*f^2 + d*f)*x + d)*cosh(f*x + e)^2 + 36*c*f + 27*d)*si
nh(f*x + e))/(a^3*f^2*cosh(f*x + e)^3 + 3*a^3*f^2*cosh(f*x + e)^2*sinh(f*x
+ e) + 3*a^3*f^2*cosh(f*x + e)*sinh(f*x + e)^2 + a^3*f^2*sinh(f*x + e)^3)
```

Sympy [F]

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\int \frac{c}{\tanh^3(e+fx)+3\tanh^2(e+fx)+3\tanh(e+fx)+1} dx + \int \frac{dx}{\tanh^3(e+fx)+3\tanh^2(e+fx)+3\tanh(e+fx)+1} dx}{a^3}$$

```
[In] integrate((d*x+c)/(a+a*tanh(f*x+e))**3,x)
```

```
[Out] (Integral(c/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1),
x) + Integral(d*x/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x)
+ 1), x))/a**3
```

Maxima [A] (verification not implemented)

none

Time = 0.73 (sec) , antiderivative size = 139, normalized size of antiderivative = 0.76

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{1}{96} c \left(\frac{12(fx + e)}{a^3 f} - \frac{18e^{(-2fx-2e)} + 9e^{(-4fx-4e)} + 2e^{(-6fx-6e)}}{a^3 f} \right)$$

$$+ \frac{(72f^2x^2e^{(6e)} - 108(2fxe^{(4e)} + e^{(4e)})e^{(-2fx)} - 27(4fxe^{(2e)} + e^{(2e)})e^{(-4fx)} - 4(6fx + 1)e^{(-6fx)})de^{(-6e)}}{1152a^3f^2}$$

```
[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/96*c*(12*(f*x + e)/(a^3*f) - (18*e^(-2*f*x - 2*e) + 9*e^(-4*f*x - 4*e) +
2*e^(-6*f*x - 6*e))/(a^3*f)) + 1/1152*(72*f^2*x^2*e^(6*e) - 108*(2*f*x*e^(4
*e) + e^(4*e))*e^(-2*f*x) - 27*(4*f*x*e^(2*e) + e^(2*e))*e^(-4*f*x) - 4*(6*
f*x + 1)*e^(-6*f*x))*d*e^(-6*e)/(a^3*f^2)
```

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.78

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx$$

$$= \frac{(72df^2x^2e^{(6fx+6e)} + 144cf^2xe^{(6fx+6e)} - 216dfxe^{(4fx+4e)} - 108dfxe^{(2fx+2e)} - 24dfx - 216cfe^{(4fx+4e)})}{1152a^3f^2}$$

[In] integrate((d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="giac")

```
[Out] 1/1152*(72*d*f^2*x^2*e^(6*f*x + 6*e) + 144*c*f^2*x*e^(6*f*x + 6*e) - 216*d*f*x*e^(4*f*x + 4*e) - 108*d*f*x*e^(2*f*x + 2*e) - 24*d*f*x - 216*c*f*e^(4*f*x + 4*e) - 108*c*f*e^(2*f*x + 2*e) - 24*c*f - 108*d*e^(4*f*x + 4*e) - 27*d*e^(2*f*x + 2*e) - 4*d)*e^(-6*f*x - 6*e)/(a^3*f^2)
```

Mupad [B] (verification not implemented)

Time = 1.83 (sec) , antiderivative size = 129, normalized size of antiderivative = 0.70

$$\int \frac{c + dx}{(a + a \tanh(e + fx))^3} dx = \frac{dx^2}{16a^3} - e^{-2e-2fx} \left(\frac{3d + 6cf}{32a^3f^2} + \frac{3dx}{16a^3f} \right)$$

$$- e^{-4e-4fx} \left(\frac{3d + 12cf}{128a^3f^2} + \frac{3dx}{32a^3f} \right)$$

$$- e^{-6e-6fx} \left(\frac{d + 6cf}{288a^3f^2} + \frac{dx}{48a^3f} \right) + \frac{cx}{8a^3}$$

[In] int((c + d*x)/(a + a*tanh(e + f*x))^3,x)

```
[Out] (d*x^2)/(16*a^3) - exp(- 2*e - 2*f*x)*((3*d + 6*c*f)/(32*a^3*f^2) + (3*d*x)/(16*a^3*f)) - exp(- 4*e - 4*f*x)*((3*d + 12*c*f)/(128*a^3*f^2) + (3*d*x)/(32*a^3*f)) - exp(- 6*e - 6*f*x)*((d + 6*c*f)/(288*a^3*f^2) + (d*x)/(48*a^3*f)) + (c*x)/(8*a^3)
```

$$3.46 \quad \int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx$$

Optimal result	316
Rubi [A] (verified)	317
Mathematica [A] (verified)	321
Maple [A] (verified)	321
Fricas [A] (verification not implemented)	322
Sympy [F]	322
Maxima [A] (verification not implemented)	323
Giac [A] (verification not implemented)	323
Mupad [F(-1)]	324

Optimal result

Integrand size = 20, antiderivative size = 437

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx = \frac{3 \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{3 \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} + \frac{\cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} + \frac{\log(c+dx)}{8a^3d} - \frac{\operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right) \sinh\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \frac{3 \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{8a^3d} - \frac{3 \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{8a^3d} - \frac{3 \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{3 \sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} - \frac{3 \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} + \frac{3 \sinh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} - \frac{\cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} + \frac{\sinh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d}$$

[Out] $\frac{1}{8}\text{Chi}\left(\frac{6cf}{d}+6fx\right)\cosh\left(-\frac{6e+6cf}{d}\right)/a^{\frac{3}{d}+\frac{3}{8}}\text{Chi}\left(\frac{4cf}{d}+4fx\right)\cosh\left(-\frac{4e+4cf}{d}\right)/a^{\frac{3}{d}+\frac{3}{8}}\text{Chi}\left(\frac{2cf}{d}+2fx\right)\cosh\left(-\frac{2e+2cf}{d}\right)/a^{\frac{3}{d}+\frac{1}{8}}\ln\left(\frac{dx+c}{a}\right)/a^{\frac{3}{d}-\frac{3}{8}}\cosh\left(-\frac{2e+2cf}{d}\right)\text{Shi}\left(\frac{2cf}{d}+2fx\right)/a^{\frac{3}{d}-\frac{3}{8}}\cosh\left(-\frac{4e+4cf}{d}\right)\text{Shi}\left(\frac{4cf}{d}+4fx\right)/a^{\frac{3}{d}-\frac{1}{8}}\cosh\left(-\frac{6e+6cf}{d}\right)\text{Shi}\left(\frac{6cf}{d}+6fx\right)/a^{\frac{3}{d}+\frac{1}{8}}\text{Chi}\left(\frac{6cf}{d}+6fx\right)\sinh\left(-\frac{6e+6cf}{d}\right)/a^{\frac{3}{d}-\frac{1}{8}}\text{Shi}\left(\frac{6cf}{d}+6fx\right)\sinh\left(-\frac{6e+6cf}{d}\right)/a^{\frac{3}{d}+\frac{3}{8}}\text{Chi}\left(\frac{4cf}{d}+4fx\right)\sinh\left(-\frac{4e+4cf}{d}\right)/a^{\frac{3}{d}-\frac{3}{8}}\text{Shi}\left(\frac{4cf}{d}+4fx\right)\sinh\left(-\frac{4e+4cf}{d}\right)/a^{\frac{3}{d}+\frac{3}{8}}\text{Chi}\left(\frac{2cf}{d}+2fx\right)\sinh\left(-\frac{2e+2cf}{d}\right)/a^{\frac{3}{d}-\frac{3}{8}}\text{Shi}\left(\frac{2cf}{d}+2fx\right)\sinh\left(-\frac{2e+2cf}{d}\right)/a^{\frac{3}{d}}$

Rubi [A] (verified)

Time = 1.28 (sec) , antiderivative size = 437, normalized size of antiderivative = 1.00, number of steps used = 53, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.350$, Rules used = {3809, 3384, 3379, 3382, 3393, 5556, 5578}

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx = -\frac{3\text{Chi}\left(2xf + \frac{2cf}{d}\right) \sinh\left(2e - \frac{2cf}{d}\right)}{8a^3d} - \frac{\text{Chi}\left(6xf + \frac{6cf}{d}\right) \sinh\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \frac{3\text{Chi}\left(4xf + \frac{4cf}{d}\right) \sinh\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{3\text{Chi}\left(2xf + \frac{2cf}{d}\right) \cosh\left(2e - \frac{2cf}{d}\right)}{8a^3d} + \frac{3\text{Chi}\left(4xf + \frac{4cf}{d}\right) \cosh\left(4e - \frac{4cf}{d}\right)}{8a^3d} + \frac{\text{Chi}\left(6xf + \frac{6cf}{d}\right) \cosh\left(6e - \frac{6cf}{d}\right)}{8a^3d} + \frac{3 \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} + \frac{3 \sinh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} + \frac{\sinh\left(6e - \frac{6cf}{d}\right) \text{Shi}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} - \frac{3 \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{8a^3d} - \frac{3 \cosh\left(4e - \frac{4cf}{d}\right) \text{Shi}\left(4xf + \frac{4cf}{d}\right)}{8a^3d} - \frac{\cosh\left(6e - \frac{6cf}{d}\right) \text{Shi}\left(6xf + \frac{6cf}{d}\right)}{8a^3d} + \frac{\log(c+dx)}{8a^3d}$$

[In] Int[1/((c + d*x)*(a + a*Tanh[e + f*x])^3),x]

[Out] $(3*\text{Cosh}\left[\frac{2e - (2cf)}{d}\right]*\text{CoshIntegral}\left[\frac{(2cf)}{d} + 2fx\right])/(8*a^3*d) + (3*\text{Cosh}\left[\frac{4e - (4cf)}{d}\right]*\text{CoshIntegral}\left[\frac{(4cf)}{d} + 4fx\right])/(8*a^3*d) + (\text{Cosh}\left[6e$

$$\begin{aligned}
& - (6*c*f)/d * \text{CoshIntegral}[(6*c*f)/d + 6*f*x]/(8*a^3*d) + \text{Log}[c + d*x]/(8*a^3*d) - (\text{CoshIntegral}[(6*c*f)/d + 6*f*x] * \text{Sinh}[6*e - (6*c*f)/d])/(8*a^3*d) - \\
& (3*\text{CoshIntegral}[(4*c*f)/d + 4*f*x] * \text{Sinh}[4*e - (4*c*f)/d])/(8*a^3*d) - (3*\text{CoshIntegral}[(2*c*f)/d + 2*f*x] * \text{Sinh}[2*e - (2*c*f)/d])/(8*a^3*d) - (3*\text{Cosh}[2*e - (2*c*f)/d] * \text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) + (3*\text{Sinh}[2*e - (2*c*f)/d] * \text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) - (3*\text{Cosh}[4*e - (4*c*f)/d] * \text{SinhIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) + (3*\text{Sinh}[4*e - (4*c*f)/d] * \text{SinhIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) - (\text{Cosh}[6*e - (6*c*f)/d] * \text{SinhIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d) + (\text{Sinh}[6*e - (6*c*f)/d] * \text{SinhIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d)
\end{aligned}$$

Rule 3379

$$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*e - c*f*fz*I, 0]$$

Rule 3382

$$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \ \&\& \ \text{EqQ}[d*(e - \text{Pi}/2) - c*f*fz*I, 0]$$

Rule 3384

$$\text{Int}[\sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] \rightarrow \text{Dist}[\text{Cos}[(d*e - c*f)/d], \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \text{Dist}[\text{Sin}[(d*e - c*f)/d], \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \ \&\& \ \text{NeQ}[d*e - c*f, 0]$$

Rule 3393

$$\text{Int}[(c_.) + (d_.)*(x_)]^{(m_)} * \sin[(e_.) + (f_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \text{Sin}[e + f*x]^n, x], x] /; \text{FreeQ}[\{c, d, e, f, m\}, x] \ \&\& \ \text{IGtQ}[n, 1] \ \&\& \ (!\text{RationalQ}[m] \ || \ (\text{GeQ}[m, -1] \ \&\& \ \text{LtQ}[m, 1]))$$

Rule 3809

$$\text{Int}[(c_.) + (d_.)*(x_)]^{(m_)} * ((a_.) + (b_.) * \tan[(e_.) + (f_.)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + \text{Cos}[2*e + 2*f*x])/(2*a) + \text{Sin}[2*e + 2*f*x]/(2*b))^{(-n)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[a^2 + b^2, 0] \ \&\& \ \text{ILtQ}[m, 0] \ \&\& \ \text{ILtQ}[n, 0]$$

Rule 5556

$$\text{Int}[\text{Cosh}[(a_.) + (b_.)*(x_)]^{(p_)} * ((c_.) + (d_.)*(x_)]^{(m_)} * \text{Sinh}[(a_.) + (b_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandTrigReduce}[(c + d*x)^m, \text{Sinh}[a +$$

$b*x]^n*\text{Cosh}[a + b*x]^p, x], x] /; \text{FreeQ}[\{a, b, c, d, m\}, x] \&\& \text{IGtQ}[n, 0] \&$
 $\& \text{IGtQ}[p, 0]$

Rule 5578

$\text{Int}[\left((e_.) + (f_.)*(x_.)\right)^{(m_.)}*\text{Sinh}[(a_.) + (b_.)*(x_.)]^{(p_.)}*\text{Sinh}[(c_.) + (d_.)*(x_.)]^{(q_.)}, x_Symbol] :> \text{Int}[\text{ExpandTrigReduce}[(e + f*x)^m, \text{Sinh}[a + b*x]^p*\text{Sinh}[c + d*x]^q, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{IGtQ}[p, 0]$
 $] \&\& \text{IGtQ}[q, 0] \&\& \text{IntegerQ}[m]$

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int \left(\frac{1}{8a^3(c+dx)} + \frac{3 \cosh(2e+2fx)}{8a^3(c+dx)} + \frac{3 \cosh^2(2e+2fx)}{8a^3(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3(c+dx)} \right. \\
 &\quad - \frac{3 \sinh(2e+2fx)}{8a^3(c+dx)} - \frac{3 \cosh^2(2e+2fx) \sinh(2e+2fx)}{8a^3(c+dx)} + \frac{3 \sinh^2(2e+2fx)}{8a^3(c+dx)} \\
 &\quad \left. - \frac{\sinh^3(2e+2fx)}{8a^3(c+dx)} - \frac{3 \sinh(4e+4fx)}{8a^3(c+dx)} + \frac{3 \sinh(2e+2fx) \sinh(4e+4fx)}{16a^3(c+dx)} \right) dx \\
 &= \frac{\log(c+dx)}{8a^3d} + \frac{\int \frac{\cosh^3(2e+2fx)}{c+dx} dx}{8a^3} - \frac{\int \frac{\sinh^3(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\sinh(2e+2fx) \sinh(4e+4fx)}{c+dx} dx}{16a^3} \\
 &\quad + \frac{3 \int \frac{\cosh(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\cosh^2(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sinh(2e+2fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{3 \int \frac{\cosh^2(2e+2fx) \sinh(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\sinh^2(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sinh(4e+4fx)}{c+dx} dx}{8a^3} \\
 &= \frac{\log(c+dx)}{8a^3d} - \frac{i \int \left(\frac{3i \sinh(2e+2fx)}{4(c+dx)} - \frac{i \sinh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} \\
 &\quad + \frac{\int \left(\frac{3 \cosh(2e+2fx)}{4(c+dx)} + \frac{\cosh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} + \frac{3 \int \left(-\frac{\cosh(2e+2fx)}{2(c+dx)} + \frac{\cosh(6e+6fx)}{2(c+dx)} \right) dx}{16a^3} \\
 &\quad - \frac{3 \int \left(\frac{1}{2(c+dx)} - \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} + \frac{3 \int \left(\frac{1}{2(c+dx)} + \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} \\
 &\quad - \frac{3 \int \left(\frac{\sinh(2e+2fx)}{4(c+dx)} + \frac{\sinh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} - \frac{(3 \cosh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
 &\quad + \frac{(3 \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{(3 \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} - \frac{(3 \sinh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
 &\quad - \frac{(3 \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} + \frac{(3 \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3 \cosh \left(2e - \frac{2cf}{d}\right) \operatorname{Chi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{\log(c+dx)}{8a^3d} \\
&\quad - \frac{3 \operatorname{Chi} \left(\frac{4cf}{d} + 4fx\right) \sinh \left(4e - \frac{4cf}{d}\right)}{8a^3d} - \frac{3 \operatorname{Chi} \left(\frac{2cf}{d} + 2fx\right) \sinh \left(2e - \frac{2cf}{d}\right)}{8a^3d} \\
&\quad - \frac{3 \cosh \left(2e - \frac{2cf}{d}\right) \operatorname{Shi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{3 \sinh \left(2e - \frac{2cf}{d}\right) \operatorname{Shi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\
&\quad - \frac{3 \cosh \left(4e - \frac{4cf}{d}\right) \operatorname{Shi} \left(\frac{4cf}{d} + 4fx\right)}{8a^3d} + \frac{\int \frac{\cosh(6e+6fx)}{c+dx} dx}{32a^3} - \frac{\int \frac{\sinh(6e+6fx)}{c+dx} dx}{32a^3} \\
&\quad + \frac{3 \int \frac{\cosh(6e+6fx)}{c+dx} dx}{32a^3} - \frac{3 \int \frac{\sinh(6e+6fx)}{c+dx} dx}{32a^3} + 2 \frac{3 \int \frac{\cosh(4e+4fx)}{c+dx} dx}{16a^3} \\
&= \frac{3 \cosh \left(2e - \frac{2cf}{d}\right) \operatorname{Chi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{\log(c+dx)}{8a^3d} \\
&\quad - \frac{3 \operatorname{Chi} \left(\frac{4cf}{d} + 4fx\right) \sinh \left(4e - \frac{4cf}{d}\right)}{8a^3d} - \frac{3 \operatorname{Chi} \left(\frac{2cf}{d} + 2fx\right) \sinh \left(2e - \frac{2cf}{d}\right)}{8a^3d} \\
&\quad - \frac{3 \cosh \left(2e - \frac{2cf}{d}\right) \operatorname{Shi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{3 \sinh \left(2e - \frac{2cf}{d}\right) \operatorname{Shi} \left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\
&\quad - \frac{3 \cosh \left(4e - \frac{4cf}{d}\right) \operatorname{Shi} \left(\frac{4cf}{d} + 4fx\right)}{8a^3d} + \frac{\cosh \left(6e - \frac{6cf}{d}\right) \int \frac{\cosh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} \\
&\quad - \frac{\cosh \left(6e - \frac{6cf}{d}\right) \int \frac{\sinh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} + \frac{\left(3 \cosh \left(6e - \frac{6cf}{d}\right)\right) \int \frac{\cosh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} \\
&\quad - \frac{\left(3 \cosh \left(6e - \frac{6cf}{d}\right)\right) \int \frac{\sinh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} - \frac{\sinh \left(6e - \frac{6cf}{d}\right) \int \frac{\cosh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} \\
&\quad + \frac{\sinh \left(6e - \frac{6cf}{d}\right) \int \frac{\sinh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} - \frac{\left(3 \sinh \left(6e - \frac{6cf}{d}\right)\right) \int \frac{\cosh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} \\
&\quad + \frac{\left(3 \sinh \left(6e - \frac{6cf}{d}\right)\right) \int \frac{\sinh \left(\frac{6cf}{d} + 6fx\right)}{c+dx} dx}{32a^3} + 2 \left(\frac{\left(3 \cosh \left(4e - \frac{4cf}{d}\right)\right) \int \frac{\cosh \left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{16a^3} \right. \\
&\quad \left. + \frac{\left(3 \sinh \left(4e - \frac{4cf}{d}\right)\right) \int \frac{\sinh \left(\frac{4cf}{d} + 4fx\right)}{c+dx} dx}{16a^3} \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{3 \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} + \frac{\cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad - \frac{\operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right) \sinh\left(6e - \frac{6cf}{d}\right)}{8a^3d} - \frac{3 \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{8a^3d} \\
&\quad - \frac{3 \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{8a^3d} - \frac{3 \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} \\
&\quad + \frac{3 \sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{8a^3d} - \frac{3 \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{8a^3d} \\
&\quad + 2 \left(\frac{3 \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right)}{16a^3d} + \frac{3 \sinh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{16a^3d} \right) \\
&\quad - \frac{\cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d} + \frac{\sinh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{8a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.96 (sec) , antiderivative size = 312, normalized size of antiderivative = 0.71

$$\int \frac{1}{(c + dx)(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\operatorname{sech}^3(e + fx)(\cosh(fx) + \sinh(fx))^3 \left(\cosh(3e) \log(f(c + dx)) + \log(f(c + dx)) \sinh(3e) + (\cosh(e - \frac{4cf}{d}) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) - \sinh(e - \frac{4cf}{d}) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)) \right)}{8a^3d}$$

[In] Integrate[1/((c + d*x)*(a + a*Tanh[e + f*x])^3),x]

[Out] (Sech[e + f*x]^3*(Cosh[f*x] + Sinh[f*x])^3*(Cosh[3*e]*Log[f*(c + d*x)] + Log[f*(c + d*x)]*Sinh[3*e] + (Cosh[e - (4*c*f)/d] - Sinh[e - (4*c*f)/d])*(3*CoshIntegral[(4*f*(c + d*x))/d] + Cosh[2*e - (2*c*f)/d]*CoshIntegral[(6*f*(c + d*x))/d] - CoshIntegral[(6*f*(c + d*x))/d]*Sinh[2*e - (2*c*f)/d] + 3*CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[2*e - (2*c*f)/d] + Sinh[2*e - (2*c*f)/d]) - 3*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*f*(c + d*x))/d] - 3*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*f*(c + d*x))/d] - 3*SinhIntegral[(4*f*(c + d*x))/d] - Cosh[2*e - (2*c*f)/d]*SinhIntegral[(6*f*(c + d*x))/d] + Sinh[2*e - (2*c*f)/d]*SinhIntegral[(6*f*(c + d*x))/d]))/(8*a^3*d*(1 + Tanh[e + f*x])^3)

Maple [A] (verified)

Time = 0.30 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.35

method	result
risch	$ \frac{\ln(dx+c)}{8a^3d} - \frac{e^{\frac{6cf-6de}{d}} \operatorname{Ei}_1\left(6fx+6e+\frac{6cf-6de}{d}\right)}{8a^3d} - \frac{3e^{\frac{4cf-4de}{d}} \operatorname{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{8a^3d} - \frac{3e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{8a^3d} $

[In] `int(1/(d*x+c)/(a+a*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{8} \ln(d*x+c) / a^3 / d - \frac{1}{8} / a^3 / d * \exp(6*(c*f-d*e)/d) * Ei(1, 6*f*x+6*e+6*(c*f-d*e)/d) - \frac{3}{8} / a^3 / d * \exp(4*(c*f-d*e)/d) * Ei(1, 4*f*x+4*e+4*(c*f-d*e)/d) - \frac{3}{8} / a^3 / d * \exp(2*(c*f-d*e)/d) * Ei(1, 2*f*x+2*e+2*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 193, normalized size of antiderivative = 0.44

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx$$

$$= \frac{3 Ei\left(-\frac{2(dx+cf)}{d}\right) \cosh\left(-\frac{2(de-cf)}{d}\right) + 3 Ei\left(-\frac{4(dx+cf)}{d}\right) \cosh\left(-\frac{4(de-cf)}{d}\right) + Ei\left(-\frac{6(dx+cf)}{d}\right) \cosh\left(-\frac{6(de-cf)}{d}\right)}{a^3}$$

[In] `integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")`

[Out] $\frac{1}{8} * (3 * Ei(-2*(d*f*x + c*f)/d) * \cosh(-2*(d*e - c*f)/d) + 3 * Ei(-4*(d*f*x + c*f)/d) * \cosh(-4*(d*e - c*f)/d) + Ei(-6*(d*f*x + c*f)/d) * \cosh(-6*(d*e - c*f)/d) + 3 * Ei(-2*(d*f*x + c*f)/d) * \sinh(-2*(d*e - c*f)/d) + 3 * Ei(-4*(d*f*x + c*f)/d) * \sinh(-4*(d*e - c*f)/d) + Ei(-6*(d*f*x + c*f)/d) * \sinh(-6*(d*e - c*f)/d) + \log(d*x + c)) / (a^3 * d)$

Sympy [F]

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx$$

$$= \frac{\int \frac{1}{c \tanh^3(e+fx) + 3c \tanh^2(e+fx) + 3c \tanh(e+fx) + c + dx \tanh^3(e+fx) + 3dx \tanh^2(e+fx) + 3dx \tanh(e+fx) + dx} dx}{a^3}$$

[In] `integrate(1/(d*x+c)/(a+a*tanh(f*x+e))**3,x)`

[Out] `Integral(1/(c*tanh(e + f*x)**3 + 3*c*tanh(e + f*x)**2 + 3*c*tanh(e + f*x) + c + d*x*tanh(e + f*x)**3 + 3*d*x*tanh(e + f*x)**2 + 3*d*x*tanh(e + f*x) + d*x), x)/a**3`

Maxima [A] (verification not implemented)

none

Time = 2.49 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.26

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx = -\frac{e^{(-6e+\frac{6cf}{d})} E_1\left(\frac{6(dx+c)f}{d}\right)}{8a^3d} - \frac{3e^{(-4e+\frac{4cf}{d})} E_1\left(\frac{4(dx+c)f}{d}\right)}{8a^3d} - \frac{3e^{(-2e+\frac{2cf}{d})} E_1\left(\frac{2(dx+c)f}{d}\right)}{8a^3d} + \frac{\log(dx+c)}{8a^3d}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")

[Out] $-1/8*e^{(-6*e + 6*c*f/d)*\exp_integral_e(1, 6*(d*x + c)*f/d)/(a^3*d)} - 3/8*e^{(-4*e + 4*c*f/d)*\exp_integral_e(1, 4*(d*x + c)*f/d)/(a^3*d)} - 3/8*e^{(-2*e + 2*c*f/d)*\exp_integral_e(1, 2*(d*x + c)*f/d)/(a^3*d)} + 1/8*\log(d*x + c)/(a^3*d)$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.24

$$\int \frac{1}{(c+dx)(a+a \tanh(e+fx))^3} dx = \frac{\left(3 \operatorname{Ei}\left(-\frac{2(dfxc+cf)}{d}\right) e^{(4e+\frac{2cf}{d})} + 3 \operatorname{Ei}\left(-\frac{4(dfxc+cf)}{d}\right) e^{(2e+\frac{4cf}{d})} + \operatorname{Ei}\left(-\frac{6(dfxc+cf)}{d}\right) e^{(6e+\frac{6cf}{d})} + e^{(6e)} \log(dx+c)\right)}{8a^3d}$$

[In] integrate(1/(d*x+c)/(a+a*tanh(f*x+e))^3,x, algorithm="giac")

[Out] $1/8*(3*\operatorname{Ei}(-2*(d*f*x + c*f)/d)*e^{(4*e + 2*c*f/d)} + 3*\operatorname{Ei}(-4*(d*f*x + c*f)/d)*e^{(2*e + 4*c*f/d)} + \operatorname{Ei}(-6*(d*f*x + c*f)/d)*e^{(6*c*f/d)} + e^{(6*e)}*\log(d*x + c))*e^{(-6*e)}/(a^3*d)$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + a \tanh(e + fx))^3} dx = \int \frac{1}{(a + a \tanh(e + fx))^3 (c + dx)} dx$$

```
[In] int(1/((a + a*tanh(e + f*x))^3*(c + d*x)),x)
```

```
[Out] int(1/((a + a*tanh(e + f*x))^3*(c + d*x)), x)
```

$$3.47 \quad \int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx$$

Optimal result	326
Rubi [A] (verified)	327
Mathematica [A] (verified)	334
Maple [A] (verified)	335
Fricas [A] (verification not implemented)	335
Sympy [F]	336
Maxima [A] (verification not implemented)	337
Giac [A] (verification not implemented)	337
Mupad [F(-1)]	338

Optimal result

Integrand size = 20, antiderivative size = 692

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx = & -\frac{1}{8a^3d(c+dx)} - \frac{9 \cosh(2e+2fx)}{32a^3d(c+dx)} \\
 & - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} \\
 & - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \cosh(6e+6fx)}{32a^3d(c+dx)} \\
 & - \frac{3f \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & - \frac{3f \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & - \frac{3f \cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2} \\
 & + \frac{3f \operatorname{Chi}\left(\frac{6cf}{d} + 6fx\right) \sinh\left(6e - \frac{6cf}{d}\right)}{4a^3d^2} \\
 & + \frac{3f \operatorname{Chi}\left(\frac{4cf}{d} + 4fx\right) \sinh\left(4e - \frac{4cf}{d}\right)}{2a^3d^2} \\
 & + \frac{3f \operatorname{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{4a^3d^2} \\
 & + \frac{15 \sinh(2e+2fx)}{32a^3d(c+dx)} \\
 & - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
 & + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} + \frac{3 \sinh(6e+6fx)}{32a^3d(c+dx)} \\
 & + \frac{3f \cosh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & - \frac{3f \sinh\left(2e - \frac{2cf}{d}\right) \operatorname{Shi}\left(\frac{2cf}{d} + 2fx\right)}{4a^3d^2} \\
 & + \frac{3f \cosh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & - \frac{3f \sinh\left(4e - \frac{4cf}{d}\right) \operatorname{Shi}\left(\frac{4cf}{d} + 4fx\right)}{2a^3d^2} \\
 & + \frac{3f \cosh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2} \\
 & - \frac{3f \sinh\left(6e - \frac{6cf}{d}\right) \operatorname{Shi}\left(\frac{6cf}{d} + 6fx\right)}{4a^3d^2}
 \end{aligned}$$

[Out] -1/8/a^3/d/(d*x+c)-3/4*f*Chi(6*c*f/d+6*f*x)*cosh(-6*e+6*c*f/d)/a^3/d^2-3/2*f*Chi(4*c*f/d+4*f*x)*cosh(-4*e+4*c*f/d)/a^3/d^2-3/4*f*Chi(2*c*f/d+2*f*x)*co

$$\begin{aligned} & \operatorname{sh}(-2e+2cf/d)/a^3/d^2-9/32\cosh(2fx+2e)/a^3/d/(dxc)-3/8\cosh(2fx+ \\ & 2e)^2/a^3/d/(dxc)-1/8\cosh(2fx+2e)^3/a^3/d/(dxc)-3/32\cosh(6fx+6e) \\ & /a^3/d/(dxc)+3/4f\cosh(-2e+2cf/d)*\operatorname{Shi}(2cf/d+2fx)/a^3/d^2+3/2f* \\ & \cosh(-4e+4cf/d)*\operatorname{Shi}(4cf/d+4fx)/a^3/d^2+3/4f*\cosh(-6e+6cf/d)*\operatorname{Shi}(\\ & 6cf/d+6fx)/a^3/d^2-3/4f*\operatorname{Chi}(6cf/d+6fx)*\sinh(-6e+6cf/d)/a^3/d^2+ \\ & 3/4f*\operatorname{Shi}(6cf/d+6fx)*\sinh(-6e+6cf/d)/a^3/d^2-3/2f*\operatorname{Chi}(4cf/d+4fx) \\ &)*\sinh(-4e+4cf/d)/a^3/d^2+3/2f*\operatorname{Shi}(4cf/d+4fx)*\sinh(-4e+4cf/d)/a^ \\ & 3/d^2-3/4f*\operatorname{Chi}(2cf/d+2fx)*\sinh(-2e+2cf/d)/a^3/d^2+3/4f*\operatorname{Shi}(2cf/d \\ & +2fx)*\sinh(-2e+2cf/d)/a^3/d^2+15/32*\sinh(2fx+2e)/a^3/d/(dxc)-3/8* \\ & \sinh(2fx+2e)^2/a^3/d/(dxc)+1/8*\sinh(2fx+2e)^3/a^3/d/(dxc)+3/8*\sinh \\ & (4fx+4e)/a^3/d/(dxc)+3/32*\sinh(6fx+6e)/a^3/d/(dxc) \end{aligned}$$

Rubi [A] (verified)

Time = 1.30 (sec) , antiderivative size = 692, normalized size of antiderivative = 1.00, number of steps used = 60, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used

= {3809, 3378, 3384, 3379, 3382, 3394, 12, 5556, 5578}

$$\begin{aligned}
 \int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx = & \frac{3f \operatorname{Chi}(6xf + \frac{6cf}{d}) \sinh(6e - \frac{6cf}{d})}{4a^3 d^2} \\
 & + \frac{3f \operatorname{Chi}(4xf + \frac{4cf}{d}) \sinh(4e - \frac{4cf}{d})}{2a^3 d^2} \\
 & + \frac{3f \operatorname{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{4a^3 d^2} \\
 & - \frac{3f \operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{4a^3 d^2} \\
 & - \frac{3f \operatorname{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{2a^3 d^2} \\
 & - \frac{3f \operatorname{Chi}(6xf + \frac{6cf}{d}) \cosh(6e - \frac{6cf}{d})}{4a^3 d^2} \\
 & - \frac{3f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\
 & - \frac{3f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\
 & - \frac{3f \sinh(6e - \frac{6cf}{d}) \operatorname{Shi}(6xf + \frac{6cf}{d})}{4a^3 d^2} \\
 & + \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\
 & + \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\
 & + \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Shi}(6xf + \frac{6cf}{d})}{4a^3 d^2} \\
 & + \frac{\sinh^3(2e + 2fx)}{8a^3 d(c+dx)} - \frac{3 \sinh^2(2e + 2fx)}{8a^3 d(c+dx)} \\
 & + \frac{15 \sinh(2e + 2fx)}{32a^3 d(c+dx)} + \frac{3 \sinh(4e + 4fx)}{8a^3 d(c+dx)} \\
 & + \frac{3 \sinh(6e + 6fx)}{32a^3 d(c+dx)} - \frac{\cosh^3(2e + 2fx)}{8a^3 d(c+dx)} \\
 & - \frac{3 \cosh^2(2e + 2fx)}{8a^3 d(c+dx)} - \frac{9 \cosh(2e + 2fx)}{32a^3 d(c+dx)} \\
 & - \frac{3 \cosh(6e + 6fx)}{32a^3 d(c+dx)} - \frac{1}{8a^3 d(c+dx)}
 \end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + a*Tanh[e + f*x])^3),x]

[Out] -1/8*1/(a^3*d*(c + d*x)) - (9*Cosh[2*e + 2*f*x])/(32*a^3*d*(c + d*x)) - (3*Cosh[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) - Cosh[2*e + 2*f*x]^3/(8*a^3*d*(c + d*x)) - (3*Cosh[6*e + 6*f*x])/(32*a^3*d*(c + d*x)) - (3*f*Cosh[2*e - (2*c


```

*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x]]/(4*a^3*d^2) - (3*f*Cosh[4*e - (4*c*
f)/d]*CoshIntegral[(4*c*f)/d + 4*f*x]]/(2*a^3*d^2) - (3*f*Cosh[6*e - (6*c*f
)/d]*CoshIntegral[(6*c*f)/d + 6*f*x]]/(4*a^3*d^2) + (3*f*CoshIntegral[(6*c*
f)/d + 6*f*x]*Sinh[6*e - (6*c*f)/d]]/(4*a^3*d^2) + (3*f*CoshIntegral[(4*c*f
)/d + 4*f*x]*Sinh[4*e - (4*c*f)/d]]/(2*a^3*d^2) + (3*f*CoshIntegral[(2*c*f)
/d + 2*f*x]*Sinh[2*e - (2*c*f)/d]]/(4*a^3*d^2) + (15*Sinh[2*e + 2*f*x]]/(32
*a^3*d*(c + d*x)) - (3*Sinh[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) + Sinh[2*e
+ 2*f*x]^3/(8*a^3*d*(c + d*x)) + (3*Sinh[4*e + 4*f*x]]/(8*a^3*d*(c + d*x))
+ (3*Sinh[6*e + 6*f*x]]/(32*a^3*d*(c + d*x)) + (3*f*Cosh[2*e - (2*c*f)/d]*S
inhIntegral[(2*c*f)/d + 2*f*x]]/(4*a^3*d^2) - (3*f*Sinh[2*e - (2*c*f)/d]*Si
nhIntegral[(2*c*f)/d + 2*f*x]]/(4*a^3*d^2) + (3*f*Cosh[4*e - (4*c*f)/d]*Sin
hIntegral[(4*c*f)/d + 4*f*x]]/(2*a^3*d^2) - (3*f*Sinh[4*e - (4*c*f)/d]*Sinh
Integral[(4*c*f)/d + 4*f*x]]/(2*a^3*d^2) + (3*f*Cosh[6*e - (6*c*f)/d]*SinhI
ntegral[(6*c*f)/d + 6*f*x]]/(4*a^3*d^2) - (3*f*Sinh[6*e - (6*c*f)/d]*SinhIn
tegral[(6*c*f)/d + 6*f*x]]/(4*a^3*d^2)

```

Rule 12

```

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !Match
Q[u, (b_)*(v_)] /; FreeQ[b, x]

```

Rule 3378

```

Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)], x_Symbol] := Simp[(c
+ d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c
+ d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1
]

```

Rule 3379

```

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f
, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

```

Rule 3382

```

Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_)]]/((c_.) + (d_.)*(x_)), x_Symbo
l] := Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz
}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

```

Rule 3384

```

Int[sin[(e_.) + (f_.)*(x_)]]/((c_.) + (d_.)*(x_)), x_Symbol] := Dist[Cos[(d*
e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f
)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] &&
NeQ[d*e - c*f, 0]

```

Rule 3394

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := Si
mp[(c + d*x)^(m + 1)*(Sin[e + f*x]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1
))), Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n -
1), x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] &&
LtQ[m, -1]
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_.)*((c_.) + (d_.)*(x_))^(m_.)*Sinh[(a_.) +
(b_.)*(x_)]^(n_.), x_Symbol] := Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 5578

```
Int[((e_.) + (f_.)*(x_))^(m_.)*Sinh[(a_.) + (b_.)*(x_)]^(p_.)*Sinh[(c_.) +
(d_.)*(x_)]^(q_.), x_Symbol] := Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a +
b*x]^p*Sinh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0
] && IGtQ[q, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{1}{8a^3(c+dx)^2} + \frac{3 \cosh(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \cosh^2(2e+2fx)}{8a^3(c+dx)^2} + \frac{\cosh^3(2e+2fx)}{8a^3(c+dx)^2} \right. \\ &\quad - \frac{3 \sinh(2e+2fx)}{8a^3(c+dx)^2} - \frac{3 \cosh^2(2e+2fx) \sinh(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \sinh^2(2e+2fx)}{8a^3(c+dx)^2} \\ &\quad \left. - \frac{\sinh^3(2e+2fx)}{8a^3(c+dx)^2} - \frac{3 \sinh(4e+4fx)}{8a^3(c+dx)^2} + \frac{3 \sinh(2e+2fx) \sinh(4e+4fx)}{16a^3(c+dx)^2} \right) dx \\ &= -\frac{1}{8a^3 d(c+dx)} + \frac{\int \frac{\cosh^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{\int \frac{\sinh^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\sinh(2e+2fx) \sinh(4e+4fx)}{(c+dx)^2} dx}{16a^3} \\ &\quad + \frac{3 \int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\cosh^2(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{3 \int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{8a^3} \\ &\quad - \frac{3 \int \frac{\cosh^2(2e+2fx) \sinh(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\sinh^2(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{3 \int \frac{\sinh(4e+4fx)}{(c+dx)^2} dx}{8a^3} \end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{3 \cosh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&+ \frac{3 \sinh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&+ \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} + \frac{3 \int \left(-\frac{\cosh(2e+2fx)}{2(c+dx)^2} + \frac{\cosh(6e+6fx)}{2(c+dx)^2} \right) dx}{16a^3} \\
&- \frac{3 \int \left(\frac{\sinh(2e+2fx)}{4(c+dx)^2} + \frac{\sinh(6e+6fx)}{4(c+dx)^2} \right) dx}{8a^3} + \frac{(3if) \int \left(-\frac{i \sinh(2e+2fx)}{4(c+dx)} - \frac{i \sinh(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} \\
&+ \frac{(3if) \int -\frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{2a^3d} - \frac{(3if) \int \frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{2a^3d} \\
&- \frac{(3f) \int \frac{\cosh(2e+2fx)}{c+dx} dx}{4a^3d} + \frac{(3f) \int \left(\frac{\cosh(2e+2fx)}{4(c+dx)} - \frac{\cosh(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} \\
&+ \frac{(3f) \int \frac{\sinh(2e+2fx)}{c+dx} dx}{4a^3d} - \frac{(3f) \int \frac{\cosh(4e+4fx)}{c+dx} dx}{2a^3d} \\
&= -\frac{1}{8a^3d(c+dx)} - \frac{3 \cosh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&+ \frac{3 \sinh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} \\
&- \frac{3 \int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{32a^3} + \frac{3 \int \frac{\cosh(6e+6fx)}{(c+dx)^2} dx}{32a^3} - \frac{3 \int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{32a^3} \\
&- \frac{3 \int \frac{\sinh(6e+6fx)}{(c+dx)^2} dx}{32a^3} + \frac{(3f) \int \frac{\cosh(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f) \int \frac{\cosh(6e+6fx)}{c+dx} dx}{16a^3d} \\
&+ \frac{(3f) \int \frac{\sinh(2e+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f) \int \frac{\sinh(6e+6fx)}{c+dx} dx}{16a^3d} + 2 \frac{(3f) \int \frac{\sinh(4e+4fx)}{c+dx} dx}{4a^3d} \\
&- \frac{(3f \cosh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} - \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} \\
&+ \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} - \frac{(3f \sinh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} \\
&+ \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} - \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9 \cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3 \cosh(6e+6fx)}{32a^3d(c+dx)} - \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} \\
&\quad - \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Chi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} + \frac{3f \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{4a^3d^2} \\
&\quad + \frac{15 \sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} + \frac{3 \sinh(6e+6fx)}{32a^3d(c+dx)} + \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} \\
&\quad - \frac{3f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} - \frac{3f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} \\
&\quad - \frac{(3f) \int \frac{\cosh(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f) \int \frac{\sinh(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(9f) \int \frac{\cosh(6e+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(9f) \int \frac{\sinh(6e+6fx)}{c+dx} dx}{16a^3d} - \frac{(3f \cosh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3f \cosh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} + \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f \sinh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f \sinh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + 2 \left(\frac{(3f \cosh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} \right. \\
&\quad \quad \quad \left. + \frac{(3f \sinh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} \right) \\
&\quad + \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9 \cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3 \cosh(6e+6fx)}{32a^3d(c+dx)} - \frac{9f \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{16a^3d^2} \\
&\quad - \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Chi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} - \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Chi}(\frac{6cf}{d} + 6fx)}{16a^3d^2} \\
&\quad + \frac{3f \operatorname{Chi}(\frac{6cf}{d} + 6fx) \sinh(6e - \frac{6cf}{d})}{16a^3d^2} + \frac{15f \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{16a^3d^2} \\
&\quad + \frac{15 \sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} + \frac{3 \sinh(6e+6fx)}{32a^3d(c+dx)} + \frac{15f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{16a^3d^2} \\
&\quad - \frac{9f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{16a^3d^2} - \frac{3f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} \\
&\quad + 2 \left(\frac{3f \operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^3d^2} + \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{4a^3d^2} \right) \\
&\quad + \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Shi}(\frac{6cf}{d} + 6fx)}{16a^3d^2} - \frac{3f \sinh(6e - \frac{6cf}{d}) \operatorname{Shi}(\frac{6cf}{d} + 6fx)}{16a^3d^2} \\
&\quad - \frac{(9f \cosh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{16a^3d} + \frac{(9f \cosh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{16a^3d} - \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(9f \sinh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{16a^3d} - \frac{(9f \sinh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{16a^3d} - \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{9 \cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3 \cosh(6e+6fx)}{32a^3d(c+dx)} - \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} \\
&\quad - \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Chi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} - \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Chi}(\frac{6cf}{d} + 6fx)}{4a^3d^2} \\
&\quad + \frac{3f \operatorname{Chi}(\frac{6cf}{d} + 6fx) \sinh(6e - \frac{6cf}{d})}{4a^3d^2} + \frac{3f \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{4a^3d^2} \\
&\quad + \frac{15 \sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} + \frac{3 \sinh(6e+6fx)}{32a^3d(c+dx)} + \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} \\
&\quad - \frac{3f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{4a^3d^2} - \frac{3f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{2a^3d^2} \\
&\quad + 2 \left(\frac{3f \operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^3d^2} + \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{4a^3d^2} \right) \\
&\quad + \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Shi}(\frac{6cf}{d} + 6fx)}{4a^3d^2} - \frac{3f \sinh(6e - \frac{6cf}{d}) \operatorname{Shi}(\frac{6cf}{d} + 6fx)}{4a^3d^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.13 (sec) , antiderivative size = 794, normalized size of antiderivative = 1.15

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx = \frac{\operatorname{sech}^3(e+fx) \left(\cosh\left(\frac{3cf}{d}\right) + \sinh\left(\frac{3cf}{d}\right) \right) \left(3d \cosh\left(e+f\left(-\frac{3c}{d}+x\right)\right) + d \cosh\left(3\left(e+f\left(-\frac{c}{d}+x\right)\right)\right) \right) + d \operatorname{co}}{\dots}$$

[In] Integrate[1/((c+d*x)^2*(a+a*Tanh[e+f*x])^3),x]

[Out] -1/8*(Sech[e+f*x]^3*(Cosh[(3*c*f)/d]+Sinh[(3*c*f)/d])*(3*d*Cosh[e+f*(-3*c)/d+x]+d*Cosh[3*(e+f*(c/d+x))] + 3*d*Cosh[e+f*((3*c)/d+x)] + 6*c*f*Cosh[3*e-(3*f*(c+d*x))/d]*CoshIntegral[(6*f*(c+d*x))/d]+6*d*f*x*Cosh[3*e-(3*f*(c+d*x))/d]*CoshIntegral[(6*f*(c+d*x))/d]+6*f*(c+d*x)*CoshIntegral[(2*f*(c+d*x))/d]*(Cosh[e-(c*f)/d+3*f*x]+Sinh[e-(c*f)/d+3*f*x])+3*d*Sinh[e+f*((-3*c)/d+x]+d*Sinh[3*(e+f*(-c/d+x))]-d*Sinh[3*(e+f*(c/d+x))]-3*d*Sinh[e+f*((3*c)/d+x)]-6*c*f*CoshIntegral[(6*f*(c+d*x))/d]*Sinh[3*e-(3*f*(c+d*x))/d]-6*d*f*x*CoshIntegral[(6*f*(c+d*x))/d]*Sinh[3*e-(3*f*(c+d*x))/d]+12*f*(c+d*x)*CoshIntegral[(4*f*(c+d*x))/d]*(Cosh[e-(f*(c+3*d*x))/d]-Sinh[e-(f*(c+3*d*x))/d])-6*c*f*Cosh[e-(c*f)/d+3*f*x]*SinhIntegral[(2*f*(c+d*x))/d]-6*d*f*x*Cosh[e-(c*f)

$$\begin{aligned} & /d + 3*f*x]*\text{SinhIntegral}[(2*f*(c + d*x))/d] - 6*c*f*\text{Sinh}[e - (c*f)/d + 3*f*x] \\ & * \text{SinhIntegral}[(2*f*(c + d*x))/d] - 6*d*f*x*\text{Sinh}[e - (c*f)/d + 3*f*x] * \text{Sinh} \\ & \text{Integral}[(2*f*(c + d*x))/d] - 12*c*f*\text{Cosh}[e - (f*(c + 3*d*x))/d] * \text{SinhIntegr} \\ & \text{al}[(4*f*(c + d*x))/d] - 12*d*f*x*\text{Cosh}[e - (f*(c + 3*d*x))/d] * \text{SinhIntegral}[(4*f*(c + d*x))/d] \\ & + 12*c*f*\text{Sinh}[e - (f*(c + 3*d*x))/d] * \text{SinhIntegral}[(4*f*(c + d*x))/d] + 12*d*f*x*\text{Sinh}[e - (f*(c + 3*d*x))/d] * \text{SinhIntegral}[(4*f*(c + d*x))/d] \\ & - 6*c*f*\text{Cosh}[3*e - (3*f*(c + d*x))/d] * \text{SinhIntegral}[(6*f*(c + d*x))/d] - 6*d*f*x*\text{Cosh}[3*e - (3*f*(c + d*x))/d] * \text{SinhIntegral}[(6*f*(c + d*x))/d] \\ & + 6*c*f*\text{Sinh}[3*e - (3*f*(c + d*x))/d] * \text{SinhIntegral}[(6*f*(c + d*x))/d] + 6*d*f*x*\text{Sinh}[3*e - (3*f*(c + d*x))/d] * \text{SinhIntegral}[(6*f*(c + d*x))/d]) / (a^3*d^2*(c + d*x)*(1 + \text{Tanh}[e + f*x])^3) \end{aligned}$$

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.35

method	result
risch	$-\frac{1}{8a^3d(dx+c)} - \frac{f e^{-6fx-6e}}{8a^3d(dx+cf)} + \frac{3f e^{\frac{6cf-6de}{d}} \text{Ei}_1\left(6fx+6e+\frac{6cf-6de}{d}\right)}{4a^3d^2} - \frac{3f e^{-4fx-4e}}{8a^3d(dx+cf)} + \frac{3f e^{\frac{4cf-4de}{d}} \text{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{2a^3d^2}$

[In] int(1/(d*x+c)^2/(a+a*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)

[Out]
$$-1/8/a^3/d/(d*x+c) - 1/8*f/a^3*\exp(-6*f*x-6*e)/d/(d*f*x+c*f) + 3/4*f/a^3/d^2*\exp(6*(c*f-d*e)/d)*\text{Ei}(1,6*f*x+6*e+6*(c*f-d*e)/d) - 3/8*f/a^3*\exp(-4*f*x-4*e)/d/(d*f*x+c*f) + 3/2*f/a^3/d^2*\exp(4*(c*f-d*e)/d)*\text{Ei}(1,4*f*x+4*e+4*(c*f-d*e)/d) - 3/8*f/a^3*\exp(-2*f*x-2*e)/d/(d*f*x+c*f) + 3/4*f/a^3/d^2*\exp(2*(c*f-d*e)/d)*\text{Ei}(1,2*f*x+2*e+2*(c*f-d*e)/d)$$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 1164, normalized size of antiderivative = 1.68

$$\int \frac{1}{(c + dx)^2(a + a \tanh(e + fx))^3} dx = \text{Too large to display}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")

[Out]
$$-1/4*(3*(d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(f*x + e)^3*\sinh(-2*(d*e - c*f)/d) + 6*(d*f*x + c*f)*\text{Ei}(-4*(d*f*x + c*f)/d)*\cosh(f*x + e)^3*\sinh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*\text{Ei}(-6*(d*f*x + c*f)/d)*\cosh(f*x + e)^3*\sinh(-6*(d*e - c*f)/d) + (3*(d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(-2*(d*e - c*f)/d) + 6*(d*f*x + c*f)*\text{Ei}(-4*(d*f*x + c*f)/d)*\cosh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*\text{Ei}(-6*(d*f*x + c*f)/d)*\cosh(-6*(d*e - c*f)/d) + d)*\cosh(f*x + e)^3 + 3*((d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(-2*(d*e - c*f)/d) +$$

```

2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + (d*f*x + c*
f)*Ei(-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-2*(d*f
*x + c*f)/d)*sinh(-2*(d*e - c*f)/d) + 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d
)*sinh(-4*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*sinh(-6*(d*
e - c*f)/d))*sinh(f*x + e)^3 + 3*(3*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*co
sh(f*x + e)*sinh(-2*(d*e - c*f)/d) + 6*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d
)*cosh(f*x + e)*sinh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)
/d)*cosh(f*x + e)*sinh(-6*(d*e - c*f)/d) + (3*(d*f*x + c*f)*Ei(-2*(d*f*x +
c*f)/d)*cosh(-2*(d*e - c*f)/d) + 6*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cos
h(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(-6*(d*e -
c*f)/d) + d)*cosh(f*x + e))*sinh(f*x + e)^2 + 3*d*cosh(f*x + e) + 9*((d*f*
x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-2*(d*e - c*f)/d) + 2*
(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-4*(d*e - c*f)/d)
+ (d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-6*(d*e - c*f)
/d) + ((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + 2*(d*f
*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(
-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f)/d))*cosh(f*x + e)^2)*sinh(f*x + e)
)/((a^3*d^3*x + a^3*c*d^2)*cosh(f*x + e)^3 + 3*(a^3*d^3*x + a^3*c*d^2)*cosh(
f*x + e)^2*sinh(f*x + e) + 3*(a^3*d^3*x + a^3*c*d^2)*cosh(f*x + e)*sinh(f*x
+ e)^2 + (a^3*d^3*x + a^3*c*d^2)*sinh(f*x + e)^3)

```

Sympy [F]

$$\int \frac{1}{(c + dx)^2(a + a \tanh(e + fx))^3} dx$$

$$= \frac{\int \frac{1}{c^2 \tanh^3(e+fx) + 3c^2 \tanh^2(e+fx) + 3c^2 \tanh(e+fx) + c^2 + 2cdx \tanh^3(e+fx) + 6cdx \tanh^2(e+fx) + 6cdx \tanh(e+fx) + 2cdx + d^2x^2 \tanh^3(e+fx)}}{a^3}$$

[In] integrate(1/(d*x+c)**2/(a+a*tanh(f*x+e))**3,x)

[Out] Integral(1/(c**2*tanh(e + f*x)**3 + 3*c**2*tanh(e + f*x)**2 + 3*c**2*tanh(e + f*x) + c**2 + 2*c*d*x*tanh(e + f*x)**3 + 6*c*d*x*tanh(e + f*x)**2 + 6*c*d*x*tanh(e + f*x) + 2*c*d*x + d**2*x**2*tanh(e + f*x)**3 + 3*d**2*x**2*tanh(e + f*x)**2 + 3*d**2*x**2*tanh(e + f*x) + d**2*x**2), x)/a**3

Maxima [A] (verification not implemented)

none

Time = 5.23 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.20

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx = -\frac{1}{8(a^3 d^2 x + a^3 cd)} - \frac{e^{(-6e+\frac{6cf}{d})} E_2\left(\frac{6(dx+c)f}{d}\right)}{8(dx+c)a^3 d} \\ - \frac{3e^{(-4e+\frac{4cf}{d})} E_2\left(\frac{4(dx+c)f}{d}\right)}{8(dx+c)a^3 d} \\ - \frac{3e^{(-2e+\frac{2cf}{d})} E_2\left(\frac{2(dx+c)f}{d}\right)}{8(dx+c)a^3 d}$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")

[Out] $-1/8/(a^3 d^2 x + a^3 c d) - 1/8 e^{(-6e + 6cf/d)} \exp_integral_e(2, 6*(dx+c)f/d)/((dx+c)a^3 d) - 3/8 e^{(-4e + 4cf/d)} \exp_integral_e(2, 4*(dx+c)f/d)/((dx+c)a^3 d) - 3/8 e^{(-2e + 2cf/d)} \exp_integral_e(2, 2*(dx+c)f/d)/((dx+c)a^3 d)$

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 840, normalized size of antiderivative = 1.21

$$\int \frac{1}{(c+dx)^2(a+a \tanh(e+fx))^3} dx = \\ \left(6(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) f^2 \text{Ei}\left(-\frac{2((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf)}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 6 def^2 \text{Ei}\left(-\frac{2((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf)}{d}\right) \right. \\ \left. - \frac{6d^2 e f^2 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 6d^2 e f^2 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 6c f^3 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 12(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 12d^2 e f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 12d^2 e f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} \right) \\ \left. - \frac{6d^2 e f^2 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 6d^2 e f^2 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 6c f^3 \text{Ei}\left(-2\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 12(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} - 12d^2 e f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} + 12d^2 e f^2 \text{Ei}\left(-4\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - \frac{de + cf}{d}\right) e^{\left(-\frac{2(de-cf)}{d}\right)} \right)$$

[In] integrate(1/(d*x+c)^2/(a+a*tanh(f*x+e))^3,x, algorithm="giac")

[Out] $-1/8*(6*(dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f)*f^2*\text{Ei}(-2*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-2*(d*e - c*f)/d)} - 6*d*e*f^2*\text{Ei}(-2*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-2*(d*e - c*f)/d)} + 6*c*f^3*\text{Ei}(-2*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-2*(d*e - c*f)/d)} + 12*(dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f)*f^2*\text{Ei}(-4*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-4*(d*e - c*f)/d)} - 12*d*e*f^2*\text{Ei}(-4*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-4*(d*e - c*f)/d)} + 12*d*e*f^2*\text{Ei}(-4*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-4*(d*e - c*f)/d)} + 12*d*e*f^2*\text{Ei}(-4*((dx+c)*(d*e/(dx+c) - c*f/(dx+c) + f) - d*e + c*f)/d)*e^{(-4*(d*e - c*f)/d)}$

```

12*c*f^3*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/
d)*e^(-4*(d*e - c*f)/d) + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f
^2*Ei(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-
6*(d*e - c*f)/d) - 6*d*e*f^2*Ei(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x +
c) + f) - d*e + c*f)/d)*e^(-6*(d*e - c*f)/d) + 6*c*f^3*Ei(-6*((d*x + c)*(d*
e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-6*(d*e - c*f)/d) + 3*d
*f^2*e^(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) + 3*d*f^2*e^(-4
*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) + d*f^2*e^(-6*(d*x + c)*(
d*e/(d*x + c) - c*f/(d*x + c) + f)/d) + d*f^2)*d^2/(((d*x + c)*a^3*d^4*(d*e
/(d*x + c) - c*f/(d*x + c) + f) - a^3*d^5*e + a^3*c*d^4*f)*f)

```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2 (a + a \tanh(e + fx))^3} dx = \int \frac{1}{(a + a \tanh(e + fx))^3 (c + dx)^2} dx$$

[In] int(1/((a + a*tanh(e + f*x))^3*(c + d*x)^2),x)

[Out] int(1/((a + a*tanh(e + f*x))^3*(c + d*x)^2), x)

3.48 $\int (c + dx)^m (a + a \tanh(e + fx))^2 dx$

Optimal result	339
Rubi [N/A]	339
Mathematica [N/A]	340
Maple [N/A] (verified)	340
Fricas [N/A]	340
Sympy [N/A]	340
Maxima [N/A]	341
Giac [N/A]	341
Mupad [N/A]	341

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \text{Int}((c + dx)^m (a + a \tanh(e + fx))^2, x)$$

[Out] Unintegrable((d*x+c)^m*(a+a*tanh(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (c + dx)^m (a + a \tanh(e + fx))^2 dx$$

[In] Int[(c + d*x)^m*(a + a*Tanh[e + f*x])^2,x]

[Out] Defer[Int][(c + d*x)^m*(a + a*Tanh[e + f*x])^2, x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + a \tanh(e + fx))^2 dx$$

Mathematica [N/A]

Not integrable

Time = 27.84 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (c + dx)^m (a + a \tanh(e + fx))^2 dx$$

[In] Integrate[(c + d*x)^m*(a + a*Tanh[e + f*x])^2,x]

[Out] Integrate[(c + d*x)^m*(a + a*Tanh[e + f*x])^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (dx + c)^m (a + a \tanh(fx + e))^2 dx$$

[In] int((d*x+c)^m*(a+a*tanh(f*x+e))^2,x)

[Out] int((d*x+c)^m*(a+a*tanh(f*x+e))^2,x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (a \tanh(fx + e) + a)^2 (dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] integral((a^2*tanh(f*x + e)^2 + 2*a^2*tanh(f*x + e) + a^2)*(d*x + c)^m, x)

Sympy [N/A]

Not integrable

Time = 2.28 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.20

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = a^2 \left(\int 2(c + dx)^m \tanh(e + fx) dx + \int (c + dx)^m \tanh^2(e + fx) dx + \int (c + dx)^m dx \right)$$

[In] integrate((d*x+c)**m*(a+a*tanh(f*x+e))**2,x)

[Out] a**2*(Integral(2*(c + d*x)**m*tanh(e + f*x), x) + Integral((c + d*x)**m*tanh(e + f*x)**2, x) + Integral((c + d*x)**m, x))

Maxima [N/A]

Not integrable

Time = 0.48 (sec) , antiderivative size = 122, normalized size of antiderivative = 6.10

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (a \tanh(fx + e) + a)^2 (dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

```
[Out] (d*x + c)^(m + 1)*a^2/(d*(m + 1)) + integrate(2*(d*x + c)^m*a^2*(e^(f*x + e)
) - e^(-f*x - e))/(e^(f*x + e) + e^(-f*x - e)) + (d*x + c)^m*a^2*(e^(f*x +
e) - e^(-f*x - e))^2/(e^(f*x + e) + e^(-f*x - e))^2, x)
```

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (a \tanh(fx + e) + a)^2 (dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((a*tanh(f*x + e) + a)^2*(d*x + c)^m, x)

Mupad [N/A]

Not integrable

Time = 1.96 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \tanh(e + fx))^2 dx = \int (a + a \tanh(e + fx))^2 (c + dx)^m dx$$

[In] int((a + a*tanh(e + f*x))^2*(c + d*x)^m,x)

[Out] int((a + a*tanh(e + f*x))^2*(c + d*x)^m, x)

3.49 $\int (c + dx)^m (a + a \tanh(e + fx)) dx$

Optimal result	342
Rubi [N/A]	342
Mathematica [N/A]	343
Maple [N/A] (verified)	343
Fricas [N/A]	343
Sympy [N/A]	343
Maxima [N/A]	344
Giac [N/A]	344
Mupad [N/A]	344

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \text{Int}((c + dx)^m (a + a \tanh(e + fx)), x)$$

[Out] Unintegrable((d*x+c)^m*(a+a*tanh(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (c + dx)^m (a + a \tanh(e + fx)) dx$$

[In] Int[(c + d*x)^m*(a + a*Tanh[e + f*x]),x]

[Out] Defer[Int][(c + d*x)^m*(a + a*Tanh[e + f*x]), x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + a \tanh(e + fx)) dx$$

Mathematica [N/A]

Not integrable

Time = 16.54 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (c + dx)^m (a + a \tanh(e + fx)) dx$$

[In] Integrate[(c + d*x)^m*(a + a*Tanh[e + f*x]),x]

[Out] Integrate[(c + d*x)^m*(a + a*Tanh[e + f*x]), x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx + c)^m (a + a \tanh(fx + e)) dx$$

[In] int((d*x+c)^m*(a+a*tanh(f*x+e)),x)

[Out] int((d*x+c)^m*(a+a*tanh(f*x+e)),x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (a \tanh(fx + e) + a)(dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e)),x, algorithm="fricas")

[Out] integral((a*tanh(f*x + e) + a)*(d*x + c)^m, x)

Sympy [N/A]

Not integrable

Time = 1.55 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.33

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = a \left(\int (c + dx)^m \tanh(e + fx) dx + \int (c + dx)^m dx \right)$$

[In] integrate((d*x+c)**m*(a+a*tanh(f*x+e)),x)

[Out] a*(Integral((c + d*x)**m*tanh(e + f*x), x) + Integral((c + d*x)**m, x))

Maxima [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 68, normalized size of antiderivative = 3.78

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (a \tanh(fx + e) + a)(dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] a*integrate((d*x + c)^m*(e^(f*x + e) - e^(-f*x - e))/(e^(f*x + e) + e^(-f*x - e)), x) + (d*x + c)^(m + 1)*a/(d*(m + 1))

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (a \tanh(fx + e) + a)(dx + c)^m dx$$

[In] integrate((d*x+c)^m*(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((a*tanh(f*x + e) + a)*(d*x + c)^m, x)

Mupad [N/A]

Not integrable

Time = 1.94 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \tanh(e + fx)) dx = \int (a + a \tanh(e + fx)) (c + dx)^m dx$$

[In] int((a + a*tanh(e + f*x))*(c + d*x)^m,x)

[Out] int((a + a*tanh(e + f*x))*(c + d*x)^m, x)

3.50 $\int \frac{(c+dx)^m}{a+a \tanh(e+fx)} dx$

Optimal result	345
Rubi [A] (verified)	345
Mathematica [A] (verified)	346
Maple [F]	347
Fricas [A] (verification not implemented)	347
Sympy [F]	347
Maxima [F]	348
Giac [F]	348
Mupad [F(-1)]	348

Optimal result

Integrand size = 20, antiderivative size = 89

$$\int \frac{(c+dx)^m}{a+a \tanh(e+fx)} dx = \frac{(c+dx)^{1+m}}{2ad(1+m)} - \frac{2^{-2-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{af}$$

[Out] 1/2*(d*x+c)^(1+m)/a/d/(1+m)-2^(-2-m)*exp(-2*e+2*c*f/d)*(d*x+c)^m*GAMMA(1+m, 2*f*(d*x+c)/d)/a/f/((f*(d*x+c)/d)^m)

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3808, 2212}

$$\int \frac{(c+dx)^m}{a+a \tanh(e+fx)} dx = \frac{(c+dx)^{m+1}}{2ad(m+1)} - \frac{2^{-m-2} e^{\frac{2cf}{d}-2e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2f(c+dx)}{d}\right)}{af}$$

[In] Int[(c + d*x)^m/(a + a*Tanh[e + f*x]),x]

[Out] (c + d*x)^(1 + m)/(2*a*d*(1 + m)) - (2^(-2 - m)*E^(-2*e + (2*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (2*f*(c + d*x))/d])/(a*f*((f*(c + d*x))/d)^m)

Rule 2212

```
Int[(F_)^((g_)*((e_) + (f_)*(x_)))*((c_) + (d_)*(x_))^(m_), x_Symbol]
:> Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d)
)^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d)^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d)*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

Rule 3808

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Sym
bol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + Dist[1/(2*a), Int[(c +
d*x)^m*E^(2*(a/b)*(e + f*x)), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
EqQ[a^2 + b^2, 0] && !IntegerQ[m]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(c + dx)^{1+m}}{2ad(1+m)} + \frac{\int e^{2i(ie+ifx)}(c + dx)^m dx}{2a} \\ &= \frac{(c + dx)^{1+m}}{2ad(1+m)} - \frac{2^{-2-m}e^{-2e+\frac{2cf}{d}}(c + dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{af} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.83 (sec) , antiderivative size = 116, normalized size of antiderivative = 1.30

$$\begin{aligned} &\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx \\ &= \frac{\left(\frac{e^e f (c+dx)^{1+m}}{d(1+m)} - 2^{-1-m} e^{-e+\frac{2cf}{d}} (c + dx)^m \left(\frac{cf}{d} + fx\right)^{-m} \Gamma(1+m, 2\left(\frac{cf}{d} + fx\right))\right) \operatorname{sech}(e + fx) (\cosh(fx) + \sinh(fx))}{2f(a + a \tanh(e + fx))} \end{aligned}$$

```
[In] Integrate[(c + d*x)^m/(a + a*Tanh[e + f*x]),x]
```

```
[Out] (((E^e*f*(c + d*x)^(1 + m))/(d*(1 + m)) - (2^(-1 - m)*E^(-e + (2*c*f)/d)*(c
+ d*x)^m*Gamma[1 + m, 2*((c*f)/d + f*x)])/((c*f)/d + f*x)^m)*Sech[e + f*x]
*(Cosh[f*x] + Sinh[f*x])/(2*f*(a + a*Tanh[e + f*x]))
```

Maple [F]

$$\int \frac{(dx + c)^m}{a + a \tanh(fx + e)} dx$$

[In] int((d*x+c)^m/(a+a*tanh(f*x+e)),x)

[Out] int((d*x+c)^m/(a+a*tanh(f*x+e)),x)

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.66

$$\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx = \frac{(dm + d) \cosh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) \Gamma\left(m + 1, \frac{2(df x + cf)}{d}\right) - (dm + d) \Gamma\left(m + 1, \frac{2(df x + cf)}{d}\right) \sinh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right)}{4(adfm + adf)}$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e)),x, algorithm="fricas")

[Out] -1/4*((d*m + d)*cosh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d)*gamma(m + 1, 2*(d*f*x + c*f)/d) - (d*m + d)*gamma(m + 1, 2*(d*f*x + c*f)/d)*sinh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d) - 2*(d*f*x + c*f)*cosh(m*log(d*x + c)) - 2*(d*f*x + c*f)*sinh(m*log(d*x + c)))/(a*d*f*m + a*d*f)

Sympy [F]

$$\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx = \frac{\int \frac{(c+dx)^m}{\tanh(e+fx)+1} dx}{a}$$

[In] integrate((d*x+c)**m/(a+a*tanh(f*x+e)),x)

[Out] Integral((c + d*x)**m/(tanh(e + f*x) + 1), x)/a

Maxima [F]

$$\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx = \int \frac{(dx + c)^m}{a \tanh(fx + e) + a} dx$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e)),x, algorithm="maxima")

[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a), x)

Giac [F]

$$\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx = \int \frac{(dx + c)^m}{a \tanh(fx + e) + a} dx$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx = \int \frac{(c + dx)^m}{a + a \tanh(e + fx)} dx$$

[In] int((c + d*x)^m/(a + a*tanh(e + f*x)),x)

[Out] int((c + d*x)^m/(a + a*tanh(e + f*x)), x)

3.51 $\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^2} dx$

Optimal result	349
Rubi [A] (verified)	349
Mathematica [A] (verified)	350
Maple [F]	351
Fricas [A] (verification not implemented)	351
Sympy [F]	351
Maxima [F]	352
Giac [F]	352
Mupad [F(-1)]	352

Optimal result

Integrand size = 20, antiderivative size = 153

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^2} dx = \frac{(c+dx)^{1+m}}{4a^2d(1+m)} - \frac{2^{-2-m}e^{-2e+\frac{2cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^2f} - \frac{4^{-2-m}e^{-4e+\frac{4cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^2f}$$

[Out] $1/4*(d*x+c)^{(1+m)}/a^2/d/(1+m)-2^{(-2-m)}*\exp(-2*e+2*c*f/d)*(d*x+c)^m*\text{GAMMA}(1+m, 2*f*(d*x+c)/d)/a^2/f/((f*(d*x+c)/d)^m)-4^{(-2-m)}*\exp(-4*e+4*c*f/d)*(d*x+c)^m*\text{GAMMA}(1+m, 4*f*(d*x+c)/d)/a^2/f/((f*(d*x+c)/d)^m)$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 153, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3810, 2212}

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^2} dx = -\frac{2^{-m-2}e^{\frac{2cf}{d}-2e}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2f(c+dx)}{d}\right)}{a^2f} - \frac{4^{-m-2}e^{\frac{4cf}{d}-4e}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{4f(c+dx)}{d}\right)}{a^2f} + \frac{(c+dx)^{m+1}}{4a^2d(m+1)}$$

[In] Int[(c + d*x)^m/(a + a*Tanh[e + f*x])^2,x]

[Out] (c + d*x)^(1 + m)/(4*a^2*d*(1 + m)) - (2^(-2 - m)*E^(-2*e + (2*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (2*f*(c + d*x))/d])/(a^2*f*((f*(c + d*x))/d)^m) - (4^(-2 - m)*E^(-4*e + (4*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (4*f*(c + d*x))/d])/(a^2*f*((f*(c + d*x))/d)^m)

Rule 2212

Int[(F_)^((g_.)*(e_.) + (f_.)*(x_)))*((c_.) + (d_.)*(x_))^(m_), x_Symbol] := Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*((-f)*g*(Log[F]/d)))^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m]])*Gamma[m + 1, ((-f)*g*(Log[F]/d)*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] && !IntegerQ[m]

Rule 3810

Int[((c_.) + (d_.)*(x_))^(m_)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c + dx)^m}{4a^2} + \frac{e^{-4e-4fx}(c + dx)^m}{4a^2} + \frac{e^{-2e-2fx}(c + dx)^m}{2a^2} \right) dx \\ &= \frac{(c + dx)^{1+m}}{4a^2d(1+m)} + \frac{\int e^{-4e-4fx}(c + dx)^m dx}{4a^2} + \frac{\int e^{-2e-2fx}(c + dx)^m dx}{2a^2} \\ &= \frac{(c + dx)^{1+m}}{4a^2d(1+m)} - \frac{2^{-2-m}e^{-2e+\frac{2cf}{d}}(c + dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^2f} \\ &\quad - \frac{4^{-2-m}e^{-4e+\frac{4cf}{d}}(c + dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^2f} \end{aligned}$$

Mathematica [A] (verified)

Time = 1.18 (sec) , antiderivative size = 163, normalized size of antiderivative = 1.07

$$\begin{aligned} &\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx \\ &= \frac{(c + dx)^m \left(\frac{4e^{2e}f(c+dx)}{d(1+m)} - 2^{2-m}e^{\frac{2cf}{d}} \left(f\left(\frac{c}{d} + x\right)\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right) - 4^{-m}e^{-2e+\frac{4cf}{d}} \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right) \right)}{16a^2f(1 + \tanh(e + fx))^2} \end{aligned}$$

[In] Integrate[(c + d*x)^m/(a + a*Tanh[e + f*x])^2,x]

[Out] ((c + d*x)^m*((4*E^(2*e)*f*(c + d*x))/(d*(1 + m)) - (2^(2 - m)*E^((2*c*f)/d))*Gamma[1 + m, (2*f*(c + d*x))/d])/(f*(c/d + x))^m - (E^(-2*e + (4*c*f)/d)*Gamma[1 + m, (4*f*(c + d*x))/d])/(4^m*((f*(c + d*x))/d)^m)*Sech[e + f*x]^2*(Cosh[f*x] + Sinh[f*x])^2/(16*a^2*f*(1 + Tanh[e + f*x])^2)

Maple [F]

$$\int \frac{(dx + c)^m}{(a + a \tanh(fx + e))^2} dx$$

[In] int((d*x+c)^m/(a+a*tanh(f*x+e))^2,x)

[Out] int((d*x+c)^m/(a+a*tanh(f*x+e))^2,x)

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 248, normalized size of antiderivative = 1.62

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx = \frac{(dm + d) \cosh\left(\frac{dm \log\left(\frac{4f}{d}\right) + 4de - 4cf}{d}\right) \Gamma\left(m + 1, \frac{4(dfx + cf)}{d}\right) + 4(dm + d) \cosh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) \Gamma\left(m + 1, \frac{2(dfx + cf)}{d}\right)}{a^2}$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] -1/16*((d*m + d)*cosh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d)*gamma(m + 1, 4*(d*f*x + c*f)/d) + 4*(d*m + d)*cosh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d)*gamma(m + 1, 2*(d*f*x + c*f)/d) - (d*m + d)*gamma(m + 1, 4*(d*f*x + c*f)/d)*sinh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d) - 4*(d*m + d)*gamma(m + 1, 2*(d*f*x + c*f)/d)*sinh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d) - 4*(d*f*x + c*f)*cosh(m*log(d*x + c)) - 4*(d*f*x + c*f)*sinh(m*log(d*x + c)))/(a^2*d*f*m + a^2*d*f)

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx = \int \frac{(c+dx)^m}{\frac{\tanh^2(e+fx)+2\tanh(e+fx)+1}{a^2}} dx$$

[In] integrate((d*x+c)**m/(a+a*tanh(f*x+e))**2,x)

[Out] Integral((c + d*x)**m/(tanh(e + f*x)**2 + 2*tanh(e + f*x) + 1), x)/a**2

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx = \int \frac{(dx + c)^m}{(a \tanh(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a)^2, x)

Giac [F]

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx = \int \frac{(dx + c)^m}{(a \tanh(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx = \int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^2} dx$$

[In] int((c + d*x)^m/(a + a*tanh(e + f*x))^2,x)

[Out] int((c + d*x)^m/(a + a*tanh(e + f*x))^2, x)

$$3.52 \quad \int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx$$

Optimal result	353
Rubi [A] (verified)	353
Mathematica [A] (verified)	355
Maple [F]	355
Fricas [A] (verification not implemented)	356
Sympy [F]	356
Maxima [F]	356
Giac [F]	357
Mupad [F(-1)]	357

Optimal result

Integrand size = 20, antiderivative size = 224

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx$$

$$= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} - \frac{3 \cdot 2^{-4-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^3 f}$$

$$- \frac{3 \cdot 2^{-5-2m} e^{-4e+\frac{4cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^3 f}$$

$$- \frac{2^{-4-m} 3^{-1-m} e^{-6e+\frac{6cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{6f(c+dx)}{d}\right)}{a^3 f}$$

```
[Out] 1/8*(d*x+c)^(1+m)/a^3/d/(1+m)-3*2^(-4-m)*exp(-2*e+2*c*f/d)*(d*x+c)^m*GAMMA(
1+m,2*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^m)-3*2^(-5-2*m)*exp(-4*e+4*c*f/d)*(
d*x+c)^m*GAMMA(1+m,4*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^m)-2^(-4-m)*3^(-1-m)
*exp(-6*e+6*c*f/d)*(d*x+c)^m*GAMMA(1+m,6*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^
m)
```

Rubi [A] (verified)

Time = 0.17 (sec) , antiderivative size = 224, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used

= {3810, 2212}

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx$$

$$= -\frac{3 \cdot 2^{-m-4} e^{\frac{2cf}{d}-2e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2f(c+dx)}{d}\right)}{a^3 f}$$

$$- \frac{3 \cdot 2^{-2m-5} e^{\frac{4cf}{d}-4e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{4f(c+dx)}{d}\right)}{a^3 f}$$

$$- \frac{2^{-m-4} 3^{-m-1} e^{\frac{6cf}{d}-6e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{6f(c+dx)}{d}\right)}{a^3 f} + \frac{(c+dx)^{m+1}}{8a^3 d(m+1)}$$

[In] Int[(c + d*x)^m/(a + a*Tanh[e + f*x])^3,x]

[Out] (c + d*x)^(1 + m)/(8*a^3*d*(1 + m)) - (3*2^(-4 - m)*E^(-2*e + (2*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (2*f*(c + d*x))/d])/(a^3*f*((f*(c + d*x))/d)^m) - (3*2^(-5 - 2*m)*E^(-4*e + (4*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (4*f*(c + d*x))/d])/(a^3*f*((f*(c + d*x))/d)^m) - (2^(-4 - m)*3^(-1 - m)*E^(-6*e + (6*c*f)/d)*(c + d*x)^m*Gamma[1 + m, (6*f*(c + d*x))/d])/(a^3*f*((f*(c + d*x))/d)^m)

Rule 2212

Int[(F_)^((g_)*((e_) + (f_)*(x_)))*((c_) + (d_)*(x_))^(m_), x_Symbol]
 := Simp[(-F^(g*(e - c*(f/d))))*(c + d*x)^FracPart[m]/(d*((-f)*g*(Log[F]/d))^(IntPart[m] + 1))*((-f)*g*Log[F]*((c + d*x)/d))^FracPart[m])*Gamma[m + 1, ((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] && !IntegerQ[m]

Rule 3810

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\text{integral} = \int \left(\frac{(c+dx)^m}{8a^3} + \frac{e^{-6e-6fx}(c+dx)^m}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^m}{8a^3} + \frac{3e^{-2e-2fx}(c+dx)^m}{8a^3} \right) dx$$

$$= \frac{(c+dx)^{1+m}}{8a^3 d(1+m)} + \frac{\int e^{-6e-6fx}(c+dx)^m dx}{8a^3}$$

$$+ \frac{3 \int e^{-4e-4fx}(c+dx)^m dx}{8a^3} + \frac{3 \int e^{-2e-2fx}(c+dx)^m dx}{8a^3}$$

$$\begin{aligned}
&= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} - \frac{3 \cdot 2^{-4-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^3 f} \\
&\quad - \frac{3 \cdot 2^{-5-2m} e^{-4e+\frac{4cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^3 f} \\
&\quad - \frac{2^{-4-m} 3^{-1-m} e^{-6e+\frac{6cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{6f(c+dx)}{d}\right)}{a^3 f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.39 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.03

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx$$

$$\frac{2^{-5-2m} 3^{-1-m} e^{-3e} \left(f\left(\frac{c}{d}+x\right)\right)^{-m} (c+dx)^m \left(12^{1+m} e^{6e} f\left(f\left(\frac{c}{d}+x\right)\right)^m (c+dx) - 2^{1+m} 3^{2+m} d e^{4e+\frac{2cf}{d}} (1+m)\right)}{a^3 d}$$

[In] Integrate[(c + d*x)^m/(a + a*Tanh[e + f*x])^3,x]

[Out] (2^(-5 - 2*m)*3^(-1 - m)*(c + d*x)^m*(12^(1 + m)*E^(6*e)*f*(f*(c/d + x))^m*(c + d*x) - 2^(1 + m)*3^(2 + m)*d*E^(4*e + (2*c*f)/d)*(1 + m)*Gamma[1 + m, (2*f*(c + d*x))/d] - 3^(2 + m)*d*E^(2*e + (4*c*f)/d)*(1 + m)*Gamma[1 + m, (4*f*(c + d*x))/d] - 2^(1 + m)*d*E^((6*c*f)/d)*(1 + m)*Gamma[1 + m, (6*f*(c + d*x))/d])*Sech[e + f*x]^3*(Cosh[f*x] + Sinh[f*x])^3)/(a^3*d*E^(3*e)*f*(1 + m)*(f*(c/d + x))^m*(1 + Tanh[e + f*x])^3)

Maple [F]

$$\int \frac{(dx+c)^m}{(a+a \tanh(fx+e))^3} dx$$

[In] int((d*x+c)^m/(a+a*tanh(f*x+e))^3,x)

[Out] int((d*x+c)^m/(a+a*tanh(f*x+e))^3,x)

Fricas [A] (verification not implemented)

none

Time = 0.10 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.54

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx = \frac{2(dm+d) \cosh\left(\frac{dm \log\left(\frac{6f}{d}\right) + 6de - 6cf}{d}\right) \Gamma\left(m+1, \frac{6(dfx+cf)}{d}\right) + 9(dm+d) \cosh\left(\frac{dm \log\left(\frac{4f}{d}\right) + 4de - 4cf}{d}\right) \Gamma\left(m+1, \frac{4(dfx+cf)}{d}\right) + 18(dm+d) \cosh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) \Gamma\left(m+1, \frac{2(dfx+cf)}{d}\right) - 2(dm+d) \Gamma(m+1, \frac{6(dfx+cf)}{d}) \sinh\left(\frac{dm \log\left(\frac{6f}{d}\right) + 6de - 6cf}{d}\right) - 9(dm+d) \Gamma(m+1, \frac{4(dfx+cf)}{d}) \sinh\left(\frac{dm \log\left(\frac{4f}{d}\right) + 4de - 4cf}{d}\right) - 18(dm+d) \Gamma(m+1, \frac{2(dfx+cf)}{d}) \sinh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) - 12(dm+d) \cosh(m \log(dx+c)) - 12(dm+d) \sinh(m \log(dx+c))}{a^3 d^m + a^3 d}$$

```
[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^3,x, algorithm="fricas")
```

```
[Out] -1/96*(2*(d*m + d)*cosh((d*m*log(6*f/d) + 6*d*e - 6*c*f)/d)*gamma(m + 1, 6*(d*f*x + c*f)/d) + 9*(d*m + d)*cosh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d)*gamma(m + 1, 4*(d*f*x + c*f)/d) + 18*(d*m + d)*cosh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d)*gamma(m + 1, 2*(d*f*x + c*f)/d) - 2*(d*m + d)*gamma(m + 1, 6*(d*f*x + c*f)/d)*sinh((d*m*log(6*f/d) + 6*d*e - 6*c*f)/d) - 9*(d*m + d)*gamma(m + 1, 4*(d*f*x + c*f)/d)*sinh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d) - 18*(d*m + d)*gamma(m + 1, 2*(d*f*x + c*f)/d)*sinh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d) - 12*(d*f*x + c*f)*cosh(m*log(d*x + c)) - 12*(d*f*x + c*f)*sinh(m*log(d*x + c)))/(a^3*d*f*m + a^3*d*f)
```

Sympy [F]

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx = \frac{\int \frac{(c+dx)^m}{\tanh^3(e+fx)+3 \tanh^2(e+fx)+3 \tanh(e+fx)+1} dx}{a^3}$$

```
[In] integrate((d*x+c)**m/(a+a*tanh(f*x+e))**3,x)
```

```
[Out] Integral((c + d*x)**m/(tanh(e + f*x)**3 + 3*tanh(e + f*x)**2 + 3*tanh(e + f*x) + 1), x)/a**3
```

Maxima [F]

$$\int \frac{(c+dx)^m}{(a+a \tanh(e+fx))^3} dx = \int \frac{(dx+c)^m}{(a \tanh(fx+e)+a)^3} dx$$

```
[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a)^3, x)
```

Giac [F]

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^3} dx = \int \frac{(dx + c)^m}{(a \tanh(fx + e) + a)^3} dx$$

[In] integrate((d*x+c)^m/(a+a*tanh(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^m/(a*tanh(f*x + e) + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^3} dx = \int \frac{(c + dx)^m}{(a + a \tanh(e + fx))^3} dx$$

[In] int((c + d*x)^m/(a + a*tanh(e + f*x))^3,x)

[Out] int((c + d*x)^m/(a + a*tanh(e + f*x))^3, x)

3.53 $\int (c + dx)^3 (a + b \tanh(e + fx)) dx$

Optimal result	358
Rubi [A] (verified)	358
Mathematica [A] (verified)	361
Maple [B] (verified)	361
Fricas [C] (verification not implemented)	362
Sympy [F]	362
Maxima [B] (verification not implemented)	363
Giac [F]	363
Mupad [F(-1)]	364

Optimal result

Integrand size = 18, antiderivative size = 137

$$\int (c + dx)^3 (a + b \tanh(e + fx)) dx = \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} + \frac{3bd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3bd^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

[Out] $\frac{1}{4}a*(d*x+c)^4/d - \frac{1}{4}b*(d*x+c)^4/d + \frac{b*(d*x+c)^3*\ln(1+\exp(2*f*x+2*e))}{f} + \frac{3}{2}b*d*(d*x+c)^2*\text{polylog}(2, -\exp(2*f*x+2*e))}{f^2} - \frac{3}{2}b*d^2*(d*x+c)*\text{polylog}(3, -\exp(2*f*x+2*e))}{f^3} + \frac{3}{4}b*d^3*\text{polylog}(4, -\exp(2*f*x+2*e))}{f^4}$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3803, 3799, 2221, 2611, 6744, 2320, 6724}

$$\int (c + dx)^3 (a + b \tanh(e + fx)) dx = \frac{a(c + dx)^4}{4d} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3bd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} + \frac{b(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} - \frac{b(c + dx)^4}{4d} + \frac{3bd^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

[In] Int[(c + d*x)^3*(a + b*Tanh[e + f*x]),x]

[Out] (a*(c + d*x)^4)/(4*d) - (b*(c + d*x)^4)/(4*d) + (b*(c + d*x)^3*Log[1 + E^(2*(e + f*x))])/f + (3*b*d*(c + d*x)^2*PolyLog[2, -E^(2*(e + f*x))]/(2*f^2) - (3*b*d^2*(c + d*x)*PolyLog[3, -E^(2*(e + f*x))]/(2*f^3) + (3*b*d^3*PolyLog[4, -E^(2*(e + f*x))]/(4*f^4)

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_))], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3803

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 6724

Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p/(e*p), x] /; FreeQ[{a, b, c, d

, e, n, p}, x] && EqQ[b*d, a*e]

Rule 6744

Int[((e_.) + (f_.)*(x_.))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_.)))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \int (a(c + dx)^3 + b(c + dx)^3 \tanh(e + fx)) dx \\
 &= \frac{a(c + dx)^4}{4d} + b \int (c + dx)^3 \tanh(e + fx) dx \\
 &= \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + (2b) \int \frac{e^{2(e+fx)}(c + dx)^3}{1 + e^{2(e+fx)}} dx \\
 &= \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
 &\quad - \frac{(3bd) \int (c + dx)^2 \log(1 + e^{2(e+fx)}) dx}{f} \\
 &= \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
 &\quad + \frac{3bd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{(3bd^2) \int (c + dx) \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
 &= \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
 &\quad + \frac{3bd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 &\quad - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{(3bd^3) \int \text{PolyLog}(3, -e^{2(e+fx)}) dx}{2f^3} \\
 &= \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
 &\quad + \frac{3bd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3bd^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 &\quad + \frac{(3bd^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3, -x)}{x} dx, x, e^{2(e+fx)}\right)}{4f^4}
 \end{aligned}$$

$$= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1+e^{2(e+fx)})}{f} + \frac{3bd(c+dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3bd^2(c+dx) \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{3bd^3 \text{PolyLog}(4, -e^{2(e+fx)})}{4f^4}$$

Mathematica [A] (verified)

Time = 0.50 (sec) , antiderivative size = 161, normalized size of antiderivative = 1.18

$$\int (c+dx)^3(a+b \tanh(e+fx)) dx = \frac{1}{4} \left(\frac{2b(c+dx)^4}{d(1+e^{2e})} + \frac{4b(c+dx)^3 \log(1+e^{-2(e+fx)})}{f} - \frac{3bd(2f^2(c+dx)^2 \text{PolyLog}(2, -e^{-2(e+fx)}) + d(2f(c+dx) \text{PolyLog}(3, -e^{-2(e+fx)}) + d \text{PolyLog}(4, -e^{-2(e+fx)}))}{f^4} + x(4c^3 + 6c^2 dx + 4cd^2 x^2 + d^3 x^3) (a + b \tanh(e)) \right)$$

[In] Integrate[(c + d*x)^3*(a + b*Tanh[e + f*x]), x]

[Out] ((2*b*(c + d*x)^4)/(d*(1 + E^(2*e))) + (4*b*(c + d*x)^3*Log[1 + E^(-2*(e + f*x))])/f - (3*b*d*(2*f^2*(c + d*x)^2*PolyLog[2, -E^(-2*(e + f*x))] + d*(2*f*(c + d*x)*PolyLog[3, -E^(-2*(e + f*x))] + d*PolyLog[4, -E^(-2*(e + f*x))]))/f^4 + x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(a + b*Tanh[e]))/4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 469 vs. 2(127) = 254.

Time = 0.26 (sec) , antiderivative size = 470, normalized size of antiderivative = 3.43

method	result
risch	$-\frac{2bd^3e^3x}{f^3} - \frac{3bd^2e^2}{f^2} + \frac{4bd^2ce^3}{f^3} + \frac{bd^3 \ln(1+e^{2fx+2e})x^3}{f} + \frac{3bd^3 \text{polylog}(2, -e^{2fx+2e})x^2}{2f^2} - \frac{3bd^3 \text{polylog}(3, -e^{2fx+2e})x}{2f^3}$

[In] int((d*x+c)^3*(a+b*tanh(f*x+e)), x, method=_RETURNVERBOSE)

[Out] -2/f^3*b*d^3*e^3*x-3/f^2*b*d*c^2*e^2+4/f^3*b*d^2*c*e^3+1/f*b*d^3*ln(1+exp(2*f*x+2*e))*x^3+3/2/f^2*b*d^3*polylog(2, -exp(2*f*x+2*e))*x^2-3/2/f^3*b*d^3*polylog(3, -exp(2*f*x+2*e))*x+2/f^4*b*d^3*e^3*ln(exp(f*x+e))+3/2/f^2*b*d*c^2*polylog(2, -exp(2*f*x+2*e))-3/2/f^3*b*d^2*c*polylog(3, -exp(2*f*x+2*e))+a*d^2*c*x^3+3/2*a*d*c^2*x^2+a*c^3*x+1/4/d*b*c^4-1/4*d^3*b*x^4+1/4*a*d^3*x^4+1/4*a/d*c^4+6/f^2*b*d^2*c*e^2*x-d^2*b*c*x^3-3/2*d*b*c^2*x^2+b*c^3*x-6/f*b*d*c^2*e*x+1/f*b*c^3*ln(1+exp(2*f*x+2*e))-2/f*b*c^3*ln(exp(f*x+e))+3/4*b*d^3*poly

$\log(4, -\exp(2fx+2e))/f^4 - 3/2/f^4 * b*d^3 * e^{-4} - 6/f^3 * b*d^2 * c * e^{-2} * \ln(\exp(fx+e)) + 3/f^2 * b*d^2 * c * \text{polylog}(2, -\exp(2fx+2e)) * x + 6/f^2 * b*d * c^2 * e * \ln(\exp(fx+e)) + 3/f * b*d * c^2 * \ln(1 + \exp(2fx+2e)) * x + 3/f * b*d^2 * c * \ln(1 + \exp(2fx+2e)) * x^2$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.29 (sec) , antiderivative size = 583, normalized size of antiderivative = 4.26

$$\int (c + dx)^3 (a + b \tanh(e + fx)) dx$$

$$= \frac{(a - b)d^3 f^4 x^4 + 4(a - b)cd^2 f^4 x^3 + 6(a - b)c^2 d f^4 x^2 + 4(a - b)c^3 f^4 x + 24bd^3 \text{polylog}(4, i \cosh(fx + e)) + \dots}{f^4}$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] 1/4*((a - b)*d^3*f^4*x^4 + 4*(a - b)*c*d^2*f^4*x^3 + 6*(a - b)*c^2*d*f^4*x^2 + 4*(a - b)*c^3*f^4*x + 24*b*d^3*polylog(4, I*cosh(f*x + e) + I*sinh(f*x + e)) + 24*b*d^3*polylog(4, -I*cosh(f*x + e) - I*sinh(f*x + e)) + 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(I*cosh(f*x + e) + I*sinh(f*x + e)) + 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(-I*cosh(f*x + e) - I*sinh(f*x + e)) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(cosh(f*x + e) + sinh(f*x + e) + I) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(cosh(f*x + e) + sinh(f*x + e) - I) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(I*cosh(f*x + e) + I*sinh(f*x + e) + 1) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-I*cosh(f*x + e) - I*sinh(f*x + e) + 1) - 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, I*cosh(f*x + e) + I*sinh(f*x + e)) - 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, -I*cosh(f*x + e) - I*sinh(f*x + e)))/f^4

Sympy [F]

$$\int (c + dx)^3 (a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx)) (c + dx)^3 dx$$

[In] integrate((d*x+c)**3*(a+b*tanh(f*x+e)),x)

[Out] Integral((a + b*tanh(e + f*x))*(c + d*x)**3, x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 304 vs. $2(126) = 252$.

Time = 0.26 (sec) , antiderivative size = 304, normalized size of antiderivative = 2.22

$$\int (c+dx)^3(a+b \tanh(e+fx)) dx = \frac{1}{4} ad^3x^4 + \frac{1}{4} bd^3x^4 + acd^2x^3 + bcd^2x^3 + \frac{3}{2} ac^2dx^2 + \frac{3}{2} bc^2dx^2 + ac^3x + \frac{bc^3 \log(\cosh(fx+e))}{f} + \frac{3(2fx \log(e^{(2fx+2e)}+1) + \text{Li}_2(-e^{(2fx+2e)}))bcd^2}{2f^2} + \frac{3(2f^2x^2 \log(e^{(2fx+2e)}+1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))bcd^2}{2f^3} + \frac{(4f^3x^3 \log(e^{(2fx+2e)}+1) + 6f^2x^2 \text{Li}_2(-e^{(2fx+2e)}) - 6fx \text{Li}_3(-e^{(2fx+2e)}) + 3 \text{Li}_4(-e^{(2fx+2e)}))bd^3}{3f^4} - \frac{bd^3f^4x^4 + 4bcd^2f^4x^3 + 6bc^2df^4x^2}{2f^4}$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/4*a*d^3*x^4 + 1/4*b*d^3*x^4 + a*c*d^2*x^3 + b*c*d^2*x^3 + 3/2*a*c^2*d*x^2 + 3/2*b*c^2*d*x^2 + a*c^3*x + b*c^3*log(cosh(f*x + e))/f + 3/2*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))*b*c^2*d/f^2 + 3/2*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))*b*c*d^2/f^3 + 1/3*(4*f^3*x^3*log(e^(2*f*x + 2*e) + 1) + 6*f^2*x^2*dilog(-e^(2*f*x + 2*e)) - 6*f*x*polylog(3, -e^(2*f*x + 2*e)) + 3*polylog(4, -e^(2*f*x + 2*e)))*b*d^3/f^4 - 1/2*(b*d^3*f^4*x^4 + 4*b*c*d^2*f^4*x^3 + 6*b*c^2*d*f^4*x^2)/f^4

Giac [F]

$$\int (c+dx)^3(a+b \tanh(e+fx)) dx = \int (dx+c)^3(b \tanh(fx+e) + a) dx$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)^3*(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx)) (c + dx)^3 dx$$

```
[In] int((a + b*tanh(e + f*x))*(c + d*x)^3,x)
```

```
[Out] int((a + b*tanh(e + f*x))*(c + d*x)^3, x)
```

3.54 $\int (c + dx)^2 (a + b \tanh(e + fx)) dx$

Optimal result	365
Rubi [A] (verified)	365
Mathematica [A] (verified)	367
Maple [B] (verified)	368
Fricas [C] (verification not implemented)	368
Sympy [F]	369
Maxima [A] (verification not implemented)	369
Giac [F]	369
Mupad [F(-1)]	370

Optimal result

Integrand size = 18, antiderivative size = 103

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx = \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} + \frac{bd(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3}$$

```
[Out] 1/3*a*(d*x+c)^3/d-1/3*b*(d*x+c)^3/d+b*(d*x+c)^2*ln(1+exp(2*f*x+2*e))/f+b*d*(d*x+c)*polylog(2,-exp(2*f*x+2*e))/f^2-1/2*b*d^2*polylog(3,-exp(2*f*x+2*e))/f^3
```

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3803, 3799, 2221, 2611, 2320, 6724}

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx = \frac{a(c + dx)^3}{3d} + \frac{bd(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{b(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{b(c + dx)^3}{3d} - \frac{bd^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3}$$

```
[In] Int[(c + d*x)^2*(a + b*Tanh[e + f*x]),x]
```

[Out] $(a*(c + d*x)^3)/(3*d) - (b*(c + d*x)^3)/(3*d) + (b*(c + d*x)^2*\text{Log}[1 + E^{(2*(e + f*x))}])/f + (b*d*(c + d*x)*\text{PolyLog}[2, -E^{(2*(e + f*x))}])/f^2 - (b*d^2*\text{PolyLog}[3, -E^{(2*(e + f*x))}])/(2*f^3)$

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_]*(f_)*(x_)]), x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3803

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 6724

Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c + dx)^2 + b(c + dx)^2 \tanh(e + fx)) dx \\
&= \frac{a(c + dx)^3}{3d} + b \int (c + dx)^2 \tanh(e + fx) dx \\
&= \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + (2b) \int \frac{e^{2(e+fx)}(c + dx)^2}{1 + e^{2(e+fx)}} dx \\
&= \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
&\quad - \frac{(2bd) \int (c + dx) \log(1 + e^{2(e+fx)}) dx}{f} \\
&= \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{(bd^2) \int \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&= \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{(bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&= \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{bd^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec) , antiderivative size = 147, normalized size of antiderivative = 1.43

$$\begin{aligned}
&\int (c + dx)^2 (a + b \tanh(e + fx)) dx \\
&= \frac{1}{6} \left(\frac{be^{2e} \left(\frac{4e^{-2e}(c+dx)^3}{d} + \frac{6(1+e^{-2e})(c+dx)^2 \log(1+e^{-2(e+fx)})}{f} - \frac{3d(1+e^{-2e})(2f(c+dx) \text{PolyLog}(2, -e^{-2(e+fx)}) + d \text{PolyLog}(3, -e^{-2(e+fx)})}{f^3} \right)}{1 + e^{2e}} \right. \\
&\quad \left. + 2x(3c^2 + 3cdx + d^2x^2) (a + b \tanh(e)) \right)
\end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Tanh[e + f*x]), x]

[Out] $((bE^{(2e)}*((4*(c + dx)^3)/(dE^{(2e)})) + (6*(1 + E^{(-2e)})*(c + dx)^2 \text{Log}[1 + E^{(-2*(e + fx))}])/f - (3*d*(1 + E^{(-2e)})*(2*f*(c + dx)*\text{PolyLog}[2, -E^{(-2*(e + fx))}] + d*\text{PolyLog}[3, -E^{(-2*(e + fx))}]))/f^3)/(1 + E^{(2e)}) + 2*x*(3*c^2 + 3*c*d*x + d^2*x^2)*(a + b*\text{Tanh}[e]))/6$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 289 vs. $2(97) = 194$.

Time = 0.24 (sec) , antiderivative size = 290, normalized size of antiderivative = 2.82

method	result
risch	$\frac{ad^2x^3}{3} - \frac{d^2bx^3}{3} + adcx^2 - dbc^2x^2 + ac^2x + bc^2x + \frac{ac^3}{3d} + \frac{bc^3}{3d} + \frac{bc^2 \ln(1+e^{2fx+2e})}{f} - \frac{2bc^2 \ln(e^{fx+e})}{f} + \frac{4bd^2}{3f^3}$

[In] `int((d*x+c)^2*(a+b*tanh(f*x+e)),x,method=_RETURNVERBOSE)`

[Out] $1/3*a*d^2*x^3 - 1/3*d^2*b*x^3 + a*d*c*x^2 - d*b*c*x^2 + a*c^2*x + b*c^2*x + 1/3*a/d*c^3 + 1/3/d*b*c^3 + 1/f*b*c^2*\ln(1+\exp(2*f*x+2*e)) - 2/f*b*c^2*\ln(\exp(f*x+e)) + 4/3/f^3*b*d^2*e^3 + 1/f*b*d^2*\ln(1+\exp(2*f*x+2*e))*x^2 - 1/2*b*d^2*\text{polylog}(3, -\exp(2*f*x+2*e))/f^3 - 4/f*b*d*c*e*x + 4/f^2*b*d*c*e*\ln(\exp(f*x+e)) - 2/f^3*b*d^2*e^2*\ln(\exp(f*x+e)) + 2/f^2*b*d^2*e^2*x + 1/f^2*b*d^2*\text{polylog}(2, -\exp(2*f*x+2*e))*x - 2/f^2*b*d*c*e^2 + 2/f*b*d*c*\ln(1+\exp(2*f*x+2*e))*x + 1/f^2*b*d*c*\text{polylog}(2, -\exp(2*f*x+2*e))$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 364, normalized size of antiderivative = 3.53

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx$$

$$= \frac{(a - b)d^2 f^3 x^3 + 3(a - b)cdf^3 x^2 + 3(a - b)c^2 f^3 x - 6bd^2 \text{polylog}(3, i \cosh(fx + e) + i \sinh(fx + e)) - 6b}{f^3}$$

[In] `integrate((d*x+c)^2*(a+b*tanh(f*x+e)),x, algorithm="fricas")`

[Out] $1/3*((a - b)*d^2*f^3*x^3 + 3*(a - b)*c*d*f^3*x^2 + 3*(a - b)*c^2*f^3*x - 6*b*d^2*\text{polylog}(3, I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 6*b*d^2*\text{polylog}(3, -I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 6*(b*d^2*f*x + b*c*d*f)*\text{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) + 6*(b*d^2*f*x + b*c*d*f)*\text{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 3*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 3*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) + 3*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) + 3*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1))/f^3$

Sympy [F]

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx)) (c + dx)^2 dx$$

[In] integrate((d*x+c)**2*(a+b*tanh(f*x+e)),x)

[Out] Integral((a + b*tanh(e + f*x))*(c + d*x)**2, x)

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 179, normalized size of antiderivative = 1.74

$$\begin{aligned} & \int (c + dx)^2 (a + b \tanh(e + fx)) dx \\ &= \frac{1}{3} ad^2 x^3 + \frac{1}{3} bd^2 x^3 + acdx^2 + bcdx^2 + ac^2 x + \frac{bc^2 \log(\cosh(fx + e))}{f} \\ & \quad + \frac{(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))bcd}{f^2} \\ & \quad + \frac{(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))bd^2}{2f^3} \\ & \quad - \frac{2(bd^2 f^3 x^3 + 3bcd f^3 x^2)}{3f^3} \end{aligned}$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/3*a*d^2*x^3 + 1/3*b*d^2*x^3 + a*c*d*x^2 + b*c*d*x^2 + a*c^2*x + b*c^2*log(cosh(f*x + e))/f + (2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e))) * b*c*d/f^2 + 1/2*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e))) * b*d^2/f^3 - 2/3*(b*d^2*f^3*x^3 + 3*b*c*d*f^3*x^2)/f^3

Giac [F]

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx = \int (dx + c)^2 (b \tanh(fx + e) + a) dx$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx)) (c + dx)^2 dx$$

```
[In] int((a + b*tanh(e + f*x))*(c + d*x)^2,x)
```

```
[Out] int((a + b*tanh(e + f*x))*(c + d*x)^2, x)
```

3.55 $\int (c + dx)(a + b \tanh(e + fx)) dx$

Optimal result	371
Rubi [A] (verified)	371
Mathematica [A] (verified)	373
Maple [A] (verified)	373
Fricas [C] (verification not implemented)	373
Sympy [F]	374
Maxima [F]	374
Giac [F]	374
Mupad [F(-1)]	374

Optimal result

Integrand size = 16, antiderivative size = 75

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 + e^{2(e+fx)})}{f} + \frac{bd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2}$$

[Out] $1/2*a*(d*x+c)^2/d-1/2*b*(d*x+c)^2/d+b*(d*x+c)*\ln(1+\exp(2*f*x+2*e))/f+1/2*b*d*\operatorname{polylog}(2,-\exp(2*f*x+2*e))/f^2$

Rubi [A] (verified)

Time = 0.08 (sec) , antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3803, 3799, 2221, 2317, 2438}

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \frac{a(c + dx)^2}{2d} + \frac{b(c + dx) \log(e^{2(e+fx)} + 1)}{f} - \frac{b(c + dx)^2}{2d} + \frac{bd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2}$$

[In] $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Tanh}[e + f*x]), x]$

[Out] $(a*(c + d*x)^2)/(2*d) - (b*(c + d*x)^2)/(2*d) + (b*(c + d*x)*\operatorname{Log}[1 + E^{2*(e + f*x)}])/f + (b*d*\operatorname{PolyLog}[2, -E^{2*(e + f*x)}])/(2*f^2)$

Rule 2221

$\operatorname{Int}[(((F_.)^((g_.)*((e_.) + (f_.)*(x_))))^((n_.)*((c_.) + (d_.)*(x_)))^((m_.))/((a_.) + (b_.)*((F_.)^((g_.)*((e_.) + (f_.)*(x_))))^((n_))), x_Symbol] \rightarrow \operatorname{Simp}$

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3799

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol]
:> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol]
:> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c + dx) + b(c + dx) \tanh(e + fx)) dx \\
&= \frac{a(c + dx)^2}{2d} + b \int (c + dx) \tanh(e + fx) dx \\
&= \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + (2b) \int \frac{e^{2(e+fx)}(c + dx)}{1 + e^{2(e+fx)}} dx \\
&= \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 + e^{2(e+fx)})}{f} - \frac{(bd) \int \log(1 + e^{2(e+fx)}) dx}{f} \\
&= \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 + e^{2(e+fx)})}{f} - \frac{(bd) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} \\
&= \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 + e^{2(e+fx)})}{f} + \frac{bd \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.10 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.96

$$\int (c + dx)(a + b \tanh(e + fx)) dx$$

$$= \frac{f(fx(2ac + adx + bdx) + 2bdx \log(1 + e^{-2(e+fx)}) + 2bc \log(\cosh(e + fx))) - bd \operatorname{PolyLog}(2, -e^{-2(e+fx)})}{2f^2}$$

[In] Integrate[(c + d*x)*(a + b*Tanh[e + f*x]),x]

[Out] (f*(f*x*(2*a*c + a*d*x + b*d*x) + 2*b*d*x*Log[1 + E^(-2*(e + f*x))] + 2*b*c*Log[Cosh[e + f*x]]) - b*d*PolyLog[2, -E^(-2*(e + f*x))])/(2*f^2)

Maple [A] (verified)

Time = 0.10 (sec) , antiderivative size = 129, normalized size of antiderivative = 1.72

method	result
risch	$\frac{adx^2}{2} + acx - \frac{bdx^2}{2} + bcx + \frac{bc \ln(1+e^{2fx+2e})}{f} - \frac{2bc \ln(e^{fx+e})}{f} - \frac{2bdex}{f} - \frac{bde^2}{f^2} + \frac{bd \ln(1+e^{2fx+2e})x}{f} + \frac{bd \operatorname{polylog}}{f}$

[In] int((d*x+c)*(a+b*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] 1/2*a*d*x^2+a*c*x-1/2*b*d*x^2+b*c*x+1/f*b*c*ln(1+exp(2*f*x+2*e))-2/f*b*c*ln(exp(f*x+e))-2/f*b*d*e*x-1/f^2*b*d*e^2+1/f*b*d*ln(1+exp(2*f*x+2*e))*x+1/2*b*d*polylog(2,-exp(2*f*x+2*e))/f^2+2/f^2*b*d*e*ln(exp(f*x+e))

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.27 (sec) , antiderivative size = 191, normalized size of antiderivative = 2.55

$$\int (c + dx)(a + b \tanh(e + fx)) dx$$

$$= \frac{(a - b)df^2x^2 + 2(a - b)cf^2x + 2bd\operatorname{Li}_2(i \cosh(fx + e) + i \sinh(fx + e)) + 2bd\operatorname{Li}_2(-i \cosh(fx + e) - i \sinh(fx + e))}{f^2}$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] 1/2*((a - b)*d*f^2*x^2 + 2*(a - b)*c*f^2*x + 2*b*d*dilog(I*cosh(f*x + e) + I*sinh(f*x + e)) + 2*b*d*dilog(-I*cosh(f*x + e) - I*sinh(f*x + e)) - 2*(b*d*e - b*c*f)*log(cosh(f*x + e) + sinh(f*x + e) + I) - 2*(b*d*e - b*c*f)*log(cosh(f*x + e) + sinh(f*x + e) - I) + 2*(b*d*f*x + b*d*e)*log(I*cosh(f*x + e) + I*sinh(f*x + e) + 1) + 2*(b*d*f*x + b*d*e)*log(-I*cosh(f*x + e) - I*sinh(f*x + e) + 1))/f^2

Sympy [F]

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx))(c + dx) dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e)),x)

[Out] Integral((a + b*tanh(e + f*x))*(c + d*x), x)

Maxima [F]

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \int (dx + c)(b \tanh(fx + e) + a) dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/2*a*d*x^2 + 1/2*(x^2 - 4*integrate(x/(e^(2*f*x + 2*e) + 1), x))*b*d + a*c*x + b*c*log(cosh(f*x + e))/f

Giac [F]

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \int (dx + c)(b \tanh(fx + e) + a) dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)*(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tanh(e + fx)) dx = \int (a + b \tanh(e + fx))(c + dx) dx$$

[In] int((a + b*tanh(e + f*x))*(c + d*x),x)

[Out] int((a + b*tanh(e + f*x))*(c + d*x), x)

3.56 $\int \frac{a+b \tanh(e+fx)}{c+dx} dx$

Optimal result	375
Rubi [N/A]	375
Mathematica [N/A]	376
Maple [N/A] (verified)	376
Fricas [N/A]	376
Sympy [N/A]	376
Maxima [N/A]	377
Giac [N/A]	377
Mupad [N/A]	377

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \text{Int}\left(\frac{a + b \tanh(e + fx)}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{a + b \tanh(e + fx)}{c + dx} dx$$

[In] Int[(a + b*Tanh[e + f*x])/(c + d*x),x]

[Out] Defer[Int] [(a + b*Tanh[e + f*x])/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{a + b \tanh(e + fx)}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 2.93 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{a + b \tanh(e + fx)}{c + dx} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])/(c + d*x),x]

[Out] Integrate[(a + b*Tanh[e + f*x])/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tanh(fx + e)}{dx + c} dx$$

[In] int((a+b*tanh(f*x+e))/(d*x+c),x)

[Out] int((a+b*tanh(f*x+e))/(d*x+c),x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{b \tanh(fx + e) + a}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c),x, algorithm="fricas")

[Out] integral((b*tanh(f*x + e) + a)/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 0.81 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{a + b \tanh(e + fx)}{c + dx} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c),x)

[Out] Integral((a + b*tanh(e + f*x))/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 56, normalized size of antiderivative = 3.11

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{b \tanh(fx + e) + a}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c),x, algorithm="maxima")

[Out] b*(log(d*x + c)/d - 2*integrate(1/(d*x + (d*x*e^(2*e) + c*e^(2*e))*e^(2*f*x) + c), x)) + a*log(d*x + c)/d

Giac [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{b \tanh(fx + e) + a}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c),x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.76 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{c + dx} dx = \int \frac{a + b \tanh(e + fx)}{c + dx} dx$$

[In] int((a + b*tanh(e + f*x))/(c + d*x),x)

[Out] int((a + b*tanh(e + f*x))/(c + d*x), x)

3.57 $\int \frac{a+b \tanh(e+fx)}{(c+dx)^2} dx$

Optimal result	378
Rubi [N/A]	378
Mathematica [N/A]	379
Maple [N/A] (verified)	379
Fricas [N/A]	379
Sympy [N/A]	379
Maxima [N/A]	380
Giac [N/A]	380
Mupad [N/A]	380

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \text{Int}\left(\frac{a + b \tanh(e + fx)}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx$$

[In] Int[(a + b*Tanh[e + f*x])/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Tanh[e + f*x])/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 5.64 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])/(c + d*x)^2,x]

[Out] Integrate[(a + b*Tanh[e + f*x])/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \tanh(fx + e)}{(dx + c)^2} dx$$

[In] int((a+b*tanh(f*x+e))/(d*x+c)^2,x)

[Out] int((a+b*tanh(f*x+e))/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{b \tanh(fx + e) + a}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b*tanh(f*x + e) + a)/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 2.91 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c)**2,x)

[Out] Integral((a + b*tanh(e + f*x))/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 87, normalized size of antiderivative = 4.83

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{b \tanh(fx + e) + a}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c)^2,x, algorithm="maxima")

[Out] -b*(1/(d^2*x + c*d) + 2*integrate(1/(d^2*x^2 + 2*c*d*x + c^2 + (d^2*x^2*e^(2*e) + 2*c*d*x*e^(2*e) + c^2*e^(2*e))*e^(2*f*x)), x)) - a/(d^2*x + c*d)

Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{b \tanh(fx + e) + a}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))/(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 1.87 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \tanh(e + fx)}{(c + dx)^2} dx$$

[In] int((a + b*tanh(e + f*x))/(c + d*x)^2,x)

[Out] int((a + b*tanh(e + f*x))/(c + d*x)^2, x)

3.58 $\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx$

Optimal result	381
Rubi [A] (verified)	382
Mathematica [A] (verified)	386
Maple [B] (verified)	386
Fricas [C] (verification not implemented)	387
Sympy [F]	389
Maxima [B] (verification not implemented)	389
Giac [F]	391
Mupad [F(-1)]	391

Optimal result

Integrand size = 20, antiderivative size = 277

$$\begin{aligned}
 \int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = & -\frac{b^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} - \frac{ab(c + dx)^4}{2d} \\
 & + \frac{b^2(c + dx)^4}{4d} + \frac{3b^2d(c + dx)^2 \log(1 + e^{2(e+fx)})}{f^2} \\
 & + \frac{2ab(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{3b^2d^2(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
 & + \frac{3abd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
 & - \frac{3b^2d^3 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^4} \\
 & - \frac{3abd^2(c + dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{f^3} \\
 & + \frac{3abd^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{2f^4} \\
 & - \frac{b^2(c + dx)^3 \tanh(e + fx)}{f}
 \end{aligned}$$

```
[Out] -b^2*(d*x+c)^3/f+1/4*a^2*(d*x+c)^4/d-1/2*a*b*(d*x+c)^4/d+1/4*b^2*(d*x+c)^4/
d+3*b^2*d*(d*x+c)^2*ln(1+exp(2*f*x+2*e))/f^2+2*a*b*(d*x+c)^3*ln(1+exp(2*f*x
+2*e))/f+3*b^2*d^2*(d*x+c)*polylog(2,-exp(2*f*x+2*e))/f^3+3*a*b*d*(d*x+c)^2
*polylog(2,-exp(2*f*x+2*e))/f^2-3/2*b^2*d^3*polylog(3,-exp(2*f*x+2*e))/f^4-
3*a*b*d^2*(d*x+c)*polylog(3,-exp(2*f*x+2*e))/f^3+3/2*a*b*d^3*polylog(4,-exp
(2*f*x+2*e))/f^4-b^2*(d*x+c)^3*tanh(f*x+e)/f
```

Rubi [A] (verified)

Time = 0.37 (sec) , antiderivative size = 277, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used = {3803, 3799, 2221, 2611, 6744, 2320, 6724, 3801, 32}

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = \frac{a^2(c + dx)^4}{4d} - \frac{3abd^2(c + dx) \text{PolyLog}(3, -e^{2(e+fx)})}{f^3} + \frac{3abd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{2ab(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} - \frac{ab(c + dx)^4}{2d} + \frac{3abd^3 \text{PolyLog}(4, -e^{2(e+fx)})}{2f^4} + \frac{3b^2d^2(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3b^2d(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f^2} - \frac{b^2(c + dx)^3 \tanh(e + fx)}{f} - \frac{b^2(c + dx)^3}{f} + \frac{b^2(c + dx)^4}{4d} - \frac{3b^2d^3 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^4}$$

[In] Int[(c + d*x)^3*(a + b*Tanh[e + f*x])^2,x]

[Out] -((b^2*(c + d*x)^3)/f) + (a^2*(c + d*x)^4)/(4*d) - (a*b*(c + d*x)^4)/(2*d) + (b^2*(c + d*x)^4)/(4*d) + (3*b^2*d*(c + d*x)^2*Log[1 + E^(2*(e + f*x))])/f^2 + (2*a*b*(c + d*x)^3*Log[1 + E^(2*(e + f*x))])/f + (3*b^2*d^2*(c + d*x)*PolyLog[2, -E^(2*(e + f*x))])/f^3 + (3*a*b*d*(c + d*x)^2*PolyLog[2, -E^(2*(e + f*x))])/f^2 - (3*b^2*d^3*PolyLog[3, -E^(2*(e + f*x))])/(2*f^4) - (3*a*b*d^2*(c + d*x)*PolyLog[3, -E^(2*(e + f*x))])/f^3 + (3*a*b*d^3*PolyLog[4, -E^(2*(e + f*x))])/(2*f^4) - (b^2*(c + d*x)^3*Tanh[e + f*x])/f

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 2221

Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x]

))^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3801

Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rule 3803

Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(
m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c + dx)^3 + 2ab(c + dx)^3 \tanh(e + fx) + b^2(c + dx)^3 \tanh^2(e + fx)) dx \\
&= \frac{a^2(c + dx)^4}{4d} + (2ab) \int (c + dx)^3 \tanh(e + fx) dx + b^2 \int (c + dx)^3 \tanh^2(e + fx) dx \\
&= \frac{a^2(c + dx)^4}{4d} - \frac{ab(c + dx)^4}{2d} - \frac{b^2(c + dx)^3 \tanh(e + fx)}{f} \\
&\quad + (4ab) \int \frac{e^{2(e+fx)}(c + dx)^3}{1 + e^{2(e+fx)}} dx + b^2 \int (c + dx)^3 dx \\
&\quad + \frac{(3b^2d) \int (c + dx)^2 \tanh(e + fx) dx}{f} \\
&= -\frac{b^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} - \frac{ab(c + dx)^4}{2d} + \frac{b^2(c + dx)^4}{4d} \\
&\quad + \frac{2ab(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} - \frac{b^2(c + dx)^3 \tanh(e + fx)}{f} \\
&\quad - \frac{(6abd) \int (c + dx)^2 \log(1 + e^{2(e+fx)}) dx}{f} + \frac{(6b^2d) \int \frac{e^{2(e+fx)}(c+dx)^2}{1+e^{2(e+fx)}} dx}{f} \\
&= -\frac{b^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} - \frac{ab(c + dx)^4}{2d} + \frac{b^2(c + dx)^4}{4d} \\
&\quad + \frac{3b^2d(c + dx)^2 \log(1 + e^{2(e+fx)})}{f^2} + \frac{2ab(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{3abd(c + dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{b^2(c + dx)^3 \tanh(e + fx)}{f} \\
&\quad - \frac{(6abd^2) \int (c + dx) \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6b^2d^2) \int (c + dx) \log(1 + e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3abd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad - \frac{3abd^2(c+dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{f^3} - \frac{b^2(c+dx)^3 \tanh(e+fx)}{f} \\
&\quad + \frac{(3abd^3) \int \operatorname{PolyLog}(3, -e^{2(e+fx)}) dx}{f^3} - \frac{(3b^2d^3) \int \operatorname{PolyLog}(2, -e^{2(e+fx)}) dx}{f^3} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3abd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad - \frac{3abd^2(c+dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{f^3} - \frac{b^2(c+dx)^3 \tanh(e+fx)}{f} \\
&\quad + \frac{(3abd^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(3, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&\quad - \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3abd(c+dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad - \frac{3b^2d^3 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^4} - \frac{3abd^2(c+dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{f^3} \\
&\quad + \frac{3abd^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{2f^4} - \frac{b^2(c+dx)^3 \tanh(e+fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.95 (sec) , antiderivative size = 544, normalized size of antiderivative = 1.96

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx$$

$$= \frac{-16bc^2(3bd + 2acf)x + \frac{16b^2(c+dx)^3}{1+e^{2e}} + \frac{8abf(c+dx)^4}{d(1+e^{2e})} + \frac{48bcd(bd+acf)x \log(1+e^{-2(e+fx)})}{f} + \frac{24bd^2(bd+2acf)x^2 \log(1+e^{-2(e+fx)})}{f}}{1}$$

[In] Integrate[(c + d*x)^3*(a + b*Tanh[e + f*x])^2,x]

[Out] (-16*b*c^2*(3*b*d + 2*a*c*f)*x + (16*b^2*(c + d*x)^3)/(1 + E^(2*e)) + (8*a*b*f*(c + d*x)^4)/(d*(1 + E^(2*e))) + (48*b*c*d*(b*d + a*c*f)*x*Log[1 + E^(-2*(e + f*x))])/f + (24*b*d^2*(b*d + 2*a*c*f)*x^2*Log[1 + E^(-2*(e + f*x))])/f + 16*a*b*d^3*x^3*Log[1 + E^(-2*(e + f*x))] + (8*b*c^2*(3*b*d + 2*a*c*f)*Log[1 + E^(2*(e + f*x))])/f - (24*b*c*d*(b*d + a*c*f)*PolyLog[2, -E^(-2*(e + f*x))])/f^2 - (24*b*d^2*(b*d + 2*a*c*f)*x*PolyLog[2, -E^(-2*(e + f*x))])/f^2 - (24*a*b*d^3*x^2*PolyLog[2, -E^(-2*(e + f*x))])/f - (12*b*d^2*(b*d + 2*a*c*f)*PolyLog[3, -E^(-2*(e + f*x))])/f^3 - (24*a*b*d^3*x*PolyLog[3, -E^(-2*(e + f*x))])/f^2 - (12*a*b*d^3*PolyLog[4, -E^(-2*(e + f*x))])/f^3 + Sech[e]*Sech[e + f*x]*((a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[f*x] + (a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[2*e + f*x] - 2*b*(4*b*(c + d*x)^3 + a*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3))*Sinh[f*x] + 2*a*b*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Sinh[2*e + f*x]))/(8*f)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 904 vs. 2(267) = 534.

Time = 0.36 (sec) , antiderivative size = 905, normalized size of antiderivative = 3.27

method	result
risch	$-\frac{12b^2d^2cex}{f^2} - \frac{6bac^2de^2}{f^2} - \frac{4bd^3ae^3x}{f^3} + \frac{8bd^2cae^3}{f^3} + \frac{6b^2d^2c \ln(1+e^{2fx+2e})x}{f^2} + \frac{b^2c^4}{4d} + \frac{a^2c^4}{4d} + \frac{a^2d^3x^4}{4} - \frac{d^3abx^4}{2} + d$

[In] int((d*x+c)^3*(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] -12/f^2*b^2*d^2*c*e*x-6/f^2*b*a*c^2*d*e^2-4/f^3*b*d^3*a*e^3*x+8/f^3*b*d^2*c*a*e^3+6/f^2*b^2*d^2*c*ln(1+exp(2*f*x+2*e))*x+1/4/d*b^2*c^4+1/4*a^2/d*c^4+1/4*a^2*d^3*x^4-1/2*d^3*a*b*x^4+d^2*b^2*c*x^3+3/2*d*b^2*c^2*x^2+2/f*b^2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/(1+exp(2*f*x+2*e))+a^2*d^2*c*x^3+3/2*a^2*d*c^2*x^2+a^2*c^3*x+3/2*a*b*d^3*polylog(4,-exp(2*f*x+2*e))/f^4-6/f*b^2*d^2*c*x^2-6/f^3*b^2*d^2*c*e^2+6/f^3*b^2*d^3*e^2*x-3/f^4*b*d^3*a*e^4-3/2*b^2*d^3*polylog(3,-exp(2*f*x+2*e))/f^4-2*d^2*a*b*c*x^3-3*d*a*b*c^2*x^2+2*a*b*c^3*x-2/f*b^2*d^3*x^3+4/f^4*b^2*d^3*e^3+1/4*d^3*b^2*x^4-4/f*b*a*c^3*ln(exp(f*x+e))

```

+3/f^2*b^2*c^2*d*ln(1+exp(2*f*x+2*e))-6/f^2*b^2*c^2*d*ln(exp(f*x+e))+3/f^3*
b^2*d^2*c*polylog(2,-exp(2*f*x+2*e))-6/f^4*b^2*e^2*d^3*ln(exp(f*x+e))+3/f^2
*b^2*d^3*ln(1+exp(2*f*x+2*e))*x^2+3/f^3*b^2*d^3*polylog(2,-exp(2*f*x+2*e))*
x+2/f*b*a*c^3*ln(1+exp(2*f*x+2*e))+12/f^2*b*d^2*c*a*e^2*x-12/f*b*a*c^2*d*e*
x+12/f^2*b*e*a*c^2*d*ln(exp(f*x+e))+6/f*b*d^2*c*a*ln(1+exp(2*f*x+2*e))*x^2+
6/f^2*b*d^2*c*a*polylog(2,-exp(2*f*x+2*e))*x+6/f*b*a*c^2*d*ln(1+exp(2*f*x+2
*e))*x-12/f^3*b*e^2*d^2*c*a*ln(exp(f*x+e))+4/f^4*b*e^3*d^3*a*ln(exp(f*x+e))
+2/f*b*d^3*a*ln(1+exp(2*f*x+2*e))*x^3+3/f^2*b*d^3*a*polylog(2,-exp(2*f*x+2*
e))*x^2-3/f^3*b*d^3*a*polylog(3,-exp(2*f*x+2*e))*x-3/f^3*b*d^2*c*a*polylog(
3,-exp(2*f*x+2*e))+3/f^2*b*a*c^2*d*polylog(2,-exp(2*f*x+2*e))+12/f^3*b^2*e*
d^2*c*ln(exp(f*x+e))+b^2*c^3*x+1/2/d*a*b*c^4

```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.32 (sec) , antiderivative size = 3744, normalized size of antiderivative = 13.52

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] 1/4*((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*(a^2 - 2*a*b + b^2)*c*d^2*f^4*x^3
+ 6*(a^2 - 2*a*b + b^2)*c^2*d*f^4*x^2 + 4*a*b*d^3*e^4 + 4*(a^2 - 2*a*b + b^
2)*c^3*f^4*x - 8*b^2*d^3*e^3 - 8*(2*a*b*c^3*e - b^2*c^3)*f^3 + 24*(a*b*c^2*
d*e^2 - b^2*c^2*d*e)*f^2 + ((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^4
- 16*a*b*c^3*e*f^3 - 8*b^2*d^3*e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b^2
)*c*d^2*f^4)*x^3 + 24*(a*b*c^2*d*e^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f^
3 - (a^2 - 2*a*b + b^2)*c^2*d*f^4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*e
^2)*f - 4*(6*b^2*c^2*d*f^3 - (a^2 - 2*a*b + b^2)*c^3*f^4)*x)*cosh(f*x + e)^
2 + 2*((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^4 - 16*a*b*c^3*e*f^3 -
8*b^2*d^3*e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b^2)*c*d^2*f^4)*x^3 + 24
*(a*b*c^2*d*e^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f^3 - (a^2 - 2*a*b + b^
2)*c^2*d*f^4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*e^2)*f - 4*(6*b^2*c^2*
d*f^3 - (a^2 - 2*a*b + b^2)*c^3*f^4)*x)*cosh(f*x + e)*sinh(f*x + e) + ((a^2
- 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^4 - 16*a*b*c^3*e*f^3 - 8*b^2*d^3*
e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b^2)*c*d^2*f^4)*x^3 + 24*(a*b*c^2*d
*e^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f^3 - (a^2 - 2*a*b + b^2)*c^2*d*f^
4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*e^2)*f - 4*(6*b^2*c^2*d*f^3 - (a^
2 - 2*a*b + b^2)*c^3*f^4)*x)*sinh(f*x + e)^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c
*d^2*e^2)*f + 24*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (a*b*d^3*
f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*co
sh(f*x + e)^2 + 2*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c
*d^2*f^2 + b^2*d^3*f)*x)*cosh(f*x + e)*sinh(f*x + e) + (a*b*d^3*f^2*x^2 + a
*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*sinh(f*x + e)

```

$$\begin{aligned}
&^2 + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*\operatorname{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + \\
&e)) + 24*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (a*b*d^3*f^2*x^2 \\
&+ a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*\cosh(f*x + \\
&e)^2 + 2*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 \\
&+ b^2*d^3*f)*x)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d^3*f^2*x^2 + a*b*c^2*d \\
&*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*\sinh(f*x + e)^2 + (2* \\
&a*b*c*d^2*f^2 + b^2*d^3*f)*x)*\operatorname{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) - 4 \\
&*(2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^ \\
&2*d)*f^2 + (2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d* \\
&e - b^2*c^2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\cosh(f*x + e)^2 + 2 \\
&*(2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^ \\
&2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\cosh(f*x + e)*\sinh(f*x + e) + \\
&(2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^ \\
&2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\sinh(f*x + e)^2 - 6*(a*b*c*d^ \\
&2*e^2 - b^2*c*d^2*e)*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) - 4*(2*a*b*d \\
&^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 \\
&+ (2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^ \\
&2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\cosh(f*x + e)^2 + 2*(2*a*b*d \\
&^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 \\
&- 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\cosh(f*x + e)*\sinh(f*x + e) + (2*a*b*d \\
&^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 \\
&- 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*\sinh(f*x + e)^2 - 6*(a*b*c*d^2*e^2 - b \\
&^2*c*d^2*e)*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) + 4*(2*a*b*d^3*f^3*x^ \\
&3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 \\
&+ b^2*d^3*f^2)*x^2 + (2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 \\
&- 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6*(a*b*c*d^2*e^2 \\
&- b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*\cosh(f*x + e)^2 + \\
&2*(2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + \\
&3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + \\
&6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*\cosh(f*x + e)*\sinh(f*x + e) + (2*a*b* \\
&d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b* \\
&c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c \\
&^2*d*f^3 + b^2*c*d^2*f^2)*x)*\sinh(f*x + e)^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2 \\
&*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*\log(I*\cosh(f*x + e) + I*\sinh(f \\
&*x + e) + 1) + 4*(2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3 \\
&*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 + (2*a*b*d^3*f^3*x^3 + \\
&2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b \\
&^2*d^3*f^2)*x^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^ \\
&2*c*d^2*f^2)*x)*\cosh(f*x + e)^2 + 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6* \\
&a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6 \\
&*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*\cos \\
&h(f*x + e)*\sinh(f*x + e) + (2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d \\
&*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6*(a*b*c*d \\
&^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*\sinh(f*x + e \\
&)^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)
\end{aligned}$$

```

*x)*log(-I*cosh(f*x + e) - I*sinh(f*x + e) + 1) + 48*(a*b*d^3*cosh(f*x + e)
^2 + 2*a*b*d^3*cosh(f*x + e)*sinh(f*x + e) + a*b*d^3*sinh(f*x + e)^2 + a*b*
d^3)*polylog(4, I*cosh(f*x + e) + I*sinh(f*x + e)) + 48*(a*b*d^3*cosh(f*x +
e)^2 + 2*a*b*d^3*cosh(f*x + e)*sinh(f*x + e) + a*b*d^3*sinh(f*x + e)^2 + a
*b*d^3)*polylog(4, -I*cosh(f*x + e) - I*sinh(f*x + e)) - 24*(2*a*b*d^3*f*x
+ 2*a*b*c*d^2*f + b^2*d^3 + (2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(
f*x + e)^2 + 2*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(f*x + e)*sinh
(f*x + e) + (2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*sinh(f*x + e)^2)*poly
log(3, I*cosh(f*x + e) + I*sinh(f*x + e)) - 24*(2*a*b*d^3*f*x + 2*a*b*c*d^2
*f + b^2*d^3 + (2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(f*x + e)^2 +
2*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(f*x + e)*sinh(f*x + e) + (
2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*sinh(f*x + e)^2)*polylog(3, -I*cos
h(f*x + e) - I*sinh(f*x + e)))/(f^4*cosh(f*x + e)^2 + 2*f^4*cosh(f*x + e)*s
inh(f*x + e) + f^4*sinh(f*x + e)^2 + f^4)

```

Sympy [F]

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx)^3 dx$$

```
[In] integrate((d*x+c)**3*(a+b*tanh(f*x+e))**2,x)
```

```
[Out] Integral((a + b*tanh(e + f*x))**2*(c + d*x)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 625 vs. $2(265) = 530$.

Time = 0.30 (sec) , antiderivative size = 625, normalized size of antiderivative = 2.26

$$\begin{aligned}
 & \int (c + dx)^3 (a + b \tanh(e + fx))^2 dx \\
 &= \frac{1}{4} a^2 d^3 x^4 + a^2 c d^2 x^3 + \frac{3}{2} a^2 c^2 d x^2 + b^2 c^3 \left(x + \frac{e}{f} - \frac{2}{f(e^{(-2fx-2e)} + 1)} \right) + a^2 c^3 x \\
 &+ \frac{3}{2} b^2 c^2 d \left(\frac{fx^2 + (fx^2 e^{(2e)} - 4xe^{(2e)}) e^{(2fx)}}{f e^{(2fx+2e)} + f} + \frac{2 \log((e^{(2fx+2e)} + 1) e^{(-2e)})}{f^2} \right) \\
 &+ \frac{2 abc^3 \log(\cosh(fx + e))}{f} \\
 &+ \frac{2(4f^3 x^3 \log(e^{(2fx+2e)} + 1) + 6f^2 x^2 \text{Li}_2(-e^{(2fx+2e)}) - 6fx \text{Li}_3(-e^{(2fx+2e)}) + 3 \text{Li}_4(-e^{(2fx+2e)})) abd^3}{3f^4} \\
 &+ \frac{(2abd^3 f + b^2 d^3 f)x^4 + 4(2abcd^2 f + (cd^2 f + 2d^3)b^2)x^3 + 12(abc^2 df + 2b^2 cd^2)x^2 + (12abc^2 df x^2 e^{(2e)} + 4(f e^{(2fx+2e)} + f))}{4(f e^{(2fx+2e)} + f)} \\
 &+ \frac{3(abc^2 df + b^2 cd^2)(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)}))}{f^3} \\
 &+ \frac{3(2abcd^2 f + b^2 d^3)(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \text{Li}_2(-e^{(2fx+2e)}) - \text{Li}_3(-e^{(2fx+2e)}))}{2f^4} \\
 &- \frac{abd^3 f^4 x^4 + 2(2abcd^2 f + b^2 d^3) f^3 x^3 + 6(abc^2 df^2 + b^2 cd^2 f) f^2 x^2}{f^4}
 \end{aligned}$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] 1/4*a^2*d^3*x^4 + a^2*c*d^2*x^3 + 3/2*a^2*c^2*d*x^2 + b^2*c^3*(x + e/f - 2/(f*(e^(-2*f*x - 2*e) + 1))) + a^2*c^3*x + 3/2*b^2*c^2*d*((f*x^2 + (f*x^2*e^(2*e) - 4*x*e^(2*e))*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) + 2*log((e^(2*f*x + 2*e) + 1)*e^(-2*e))/f^2) + 2*a*b*c^3*log(cosh(f*x + e))/f + 2/3*(4*f^3*x^3*log(e^(2*f*x + 2*e) + 1) + 6*f^2*x^2*dilog(-e^(2*f*x + 2*e)) - 6*f*x*polylog(3, -e^(2*f*x + 2*e)) + 3*polylog(4, -e^(2*f*x + 2*e)))*a*b*d^3/f^4 + 1/4*((2*a*b*d^3*f + b^2*d^3*f)*x^4 + 4*(2*a*b*c*d^2*f + (c*d^2*f + 2*d^3)*b^2)*x^3 + 12*(a*b*c^2*d*f + 2*b^2*c*d^2)*x^2 + (12*a*b*c^2*d*f*x^2*e^(2*e) + (2*a*b*d^3*f*e^(2*e) + b^2*d^3*f*e^(2*e))*x^4 + 4*(2*a*b*c*d^2*f*e^(2*e) + b^2*c*d^2*f*e^(2*e))*x^3)*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) + 3*(a*b*c^2*d*f + b^2*c*d^2)*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))/f^3 + 3/2*(2*a*b*c*d^2*f + b^2*d^3)*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))/f^4 - (a*b*d^3*f^4*x^4 + 2*(2*a*b*c*d^2*f + b^2*d^3)*f^3*x^3 + 6*(a*b*c^2*d*f^2 + b^2*c*d^2*f)*f^2*x^2)/f^4

Giac [F]

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = \int (dx + c)^3 (b \tanh(fx + e) + a)^2 dx$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)^3*(b*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx)^3 dx$$

[In] int((a + b*tanh(e + f*x))^2*(c + d*x)^3,x)

[Out] int((a + b*tanh(e + f*x))^2*(c + d*x)^3, x)

3.59 $\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx$

Optimal result	392
Rubi [A] (verified)	393
Mathematica [A] (verified)	396
Maple [B] (verified)	397
Fricas [C] (verification not implemented)	397
Sympy [F]	399
Maxima [B] (verification not implemented)	399
Giac [F]	400
Mupad [F(-1)]	400

Optimal result

Integrand size = 20, antiderivative size = 211

$$\begin{aligned}
 \int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = & -\frac{b^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} - \frac{2ab(c + dx)^3}{3d} \\
 & + \frac{b^2(c + dx)^3}{3d} + \frac{2b^2d(c + dx) \log(1 + e^{2(e+fx)})}{f^2} \\
 & + \frac{2ab(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{b^2d^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
 & + \frac{2abd(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
 & - \frac{abd^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{f^3} \\
 & - \frac{b^2(c + dx)^2 \tanh(e + fx)}{f}
 \end{aligned}$$

[Out] $-b^2(d*x+c)^2/f+1/3*a^2*(d*x+c)^3/d-2/3*a*b*(d*x+c)^3/d+1/3*b^2*(d*x+c)^3/d+2*b^2*d*(d*x+c)*\ln(1+\exp(2*f*x+2*e))/f^2+2*a*b*(d*x+c)^2*\ln(1+\exp(2*f*x+2*e))/f+b^2*d^2*polylog(2,-\exp(2*f*x+2*e))/f^3+2*a*b*d*(d*x+c)*polylog(2,-\exp(2*f*x+2*e))/f^2-a*b*d^2*polylog(3,-\exp(2*f*x+2*e))/f^3-b^2*(d*x+c)^2*\tanh(f*x+e)/f$

Rubi [A] (verified)

Time = 0.30 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3803, 3799, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32}

$$\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = \frac{a^2(c + dx)^3}{3d} + \frac{2abd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{2ab(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{2ab(c + dx)^3}{3d} - \frac{abd^2 \text{PolyLog}(3, -e^{2(e+fx)})}{f^3} + \frac{2b^2d(c + dx) \log(e^{2(e+fx)} + 1)}{f^2} - \frac{b^2(c + dx)^2 \tanh(e + fx)}{f} - \frac{b^2(c + dx)^2}{f} + \frac{b^2(c + dx)^3}{3d} + \frac{b^2d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3}$$

[In] Int[(c + d*x)^2*(a + b*Tanh[e + f*x])^2,x]

[Out] -((b^2*(c + d*x)^2)/f) + (a^2*(c + d*x)^3)/(3*d) - (2*a*b*(c + d*x)^3)/(3*d) + (b^2*(c + d*x)^3)/(3*d) + (2*b^2*d*(c + d*x)*Log[1 + E^(2*(e + f*x))])/f^2 + (2*a*b*(c + d*x)^2*Log[1 + E^(2*(e + f*x))])/f + (b^2*d^2*PolyLog[2, -E^(2*(e + f*x))])/f^3 + (2*a*b*d*(c + d*x)*PolyLog[2, -E^(2*(e + f*x))])/f^2 - (a*b*d^2*PolyLog[3, -E^(2*(e + f*x))])/f^3 - (b^2*(c + d*x)^2*Tanh[e + f*x])/f

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 2221

Int[(((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^(g_.)*((e_.) + (f_.)*(x_)))^(n_.), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^(e_.)*((c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]

)ⁿ, x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_))))^(n_)]*((f_) + (g_)*(x_)^(m_)), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3799

Int[((c_) + (d_)*(x_)^(m_))*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3801

Int[((c_) + (d_)*(x_)^(m_))*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rule 3803

Int[((c_) + (d_)*(x_)^(m_))*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol]
:> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x]
&& EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c + dx)^2 + 2ab(c + dx)^2 \tanh(e + fx) + b^2(c + dx)^2 \tanh^2(e + fx)) dx \\
&= \frac{a^2(c + dx)^3}{3d} + (2ab) \int (c + dx)^2 \tanh(e + fx) dx + b^2 \int (c + dx)^2 \tanh^2(e + fx) dx \\
&= \frac{a^2(c + dx)^3}{3d} - \frac{2ab(c + dx)^3}{3d} - \frac{b^2(c + dx)^2 \tanh(e + fx)}{f} \\
&\quad + (4ab) \int \frac{e^{2(e+fx)}(c + dx)^2}{1 + e^{2(e+fx)}} dx + b^2 \int (c + dx)^2 dx \\
&\quad + \frac{(2b^2d) \int (c + dx) \tanh(e + fx) dx}{f} \\
&= -\frac{b^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} - \frac{2ab(c + dx)^3}{3d} + \frac{b^2(c + dx)^3}{3d} \\
&\quad + \frac{2ab(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} - \frac{b^2(c + dx)^2 \tanh(e + fx)}{f} \\
&\quad - \frac{(4abd) \int (c + dx) \log(1 + e^{2(e+fx)}) dx}{f} + \frac{(4b^2d) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx}{f} \\
&= -\frac{b^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} - \frac{2ab(c + dx)^3}{3d} + \frac{b^2(c + dx)^3}{3d} \\
&\quad + \frac{2b^2d(c + dx) \log(1 + e^{2(e+fx)})}{f^2} + \frac{2ab(c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{2abd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{b^2(c + dx)^2 \tanh(e + fx)}{f} \\
&\quad - \frac{(2abd^2) \int \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} - \frac{(2b^2d^2) \int \log(1 + e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad + \frac{2b^2d(c+dx)\log(1+e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^2\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{2abd(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^2} - \frac{b^2(c+dx)^2\tanh(e+fx)}{f} \\
&\quad - \frac{(abd^2)\text{Subst}\left(\int\frac{\text{PolyLog}(2,-x)}{x}dx,x,e^{2(e+fx)}\right)}{f^3} \\
&\quad - \frac{(b^2d^2)\text{Subst}\left(\int\frac{\log(1+x)}{x}dx,x,e^{2(e+fx)}\right)}{f^3} \\
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad + \frac{2b^2d(c+dx)\log(1+e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^2\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d^2\text{PolyLog}(2,-e^{2(e+fx)})}{f^3} + \frac{2abd(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^2} \\
&\quad - \frac{abd^2\text{PolyLog}(3,-e^{2(e+fx)})}{f^3} - \frac{b^2(c+dx)^2\tanh(e+fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.81 (sec) , antiderivative size = 237, normalized size of antiderivative = 1.12

$$\begin{aligned}
&\int (c+dx)^2(a+b\tanh(e+fx))^2 dx \\
&= \frac{1}{3} \left(\frac{2b\left(\frac{fx(3bd(-2ce^{2e}+dx)+2af(-3c^2e^{2e}+3cdx+d^2x^2))}{1+e^{2e}}\right) + 3dx(bd+af(2c+dx))\log(1+e^{-2(e+fx)}) + 3c(bd+acf)}{f^2} \right. \\
&\quad - \frac{3bd(bd+2af(c+dx))\text{PolyLog}(2,-e^{-2(e+fx)})}{f^3} - \frac{3abd^2\text{PolyLog}(3,-e^{-2(e+fx)})}{f^3} \\
&\quad \left. - \frac{3b^2(c+dx)^2\text{sech}(e)\text{sech}(e+fx)\sinh(fx)}{f} \right) \\
&\quad + x(3c^2+3cdx+d^2x^2)(a^2+b^2+2ab\tanh(e))
\end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Tanh[e + f*x])^2,x]

[Out] ((2*b*((f*x*(3*b*d*(-2*c*E^(2*e) + d*x) + 2*a*f*(-3*c^2*E^(2*e) + 3*c*d*x + d^2*x^2)))/(1 + E^(2*e)) + 3*d*x*(b*d + a*f*(2*c + d*x))*Log[1 + E^(-2*(e + f*x))] + 3*c*(b*d + a*c*f)*Log[1 + E^(2*(e + f*x))])/f^2 - (3*b*d*(b*d +

$$2*a*f*(c + d*x)*PolyLog[2, -E^{-2*(e + f*x)}]]/f^3 - (3*a*b*d^2*PolyLog[3, -E^{-2*(e + f*x)}]]/f^3 - (3*b^2*(c + d*x)^2*Sech[e]*Sech[e + f*x]*Sinh[f*x])/f + x*(3*c^2 + 3*c*d*x + d^2*x^2)*(a^2 + b^2 + 2*a*b*Tanh[e])/3$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 541 vs. $2(205) = 410$.

Time = 0.36 (sec) , antiderivative size = 542, normalized size of antiderivative = 2.57

method	result
risch	$\frac{4ba d^2 e^2 x}{f^2} - \frac{4bcda e^2}{f^2} - \frac{4b e^2 a d^2 \ln(e^{fx+e})}{f^3} + \frac{2ba d^2 \ln(1+e^{2fx+2e})x^2}{f} + \frac{2b^2(x^2 d^2 + 2cdx + c^2)}{f(1+e^{2fx+2e})} - \frac{4ba c^2 \ln(e^{fx+e})}{f} + \frac{2b^2 c^2}{f}$

[In] int((d*x+c)^2*(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] $4/f^2*b*a*d^2*e^2*x - 4/f^2*b*c*d*a*e^2 - 4/f^3*b*e^2*a*d^2*\ln(\exp(f*x+e)) + 2/f*b*a*d^2*\ln(1+\exp(2*f*x+2*e))*x^2 + 2/f*b^2*(d^2*x^2+2*c*d*x+c^2)/(1+\exp(2*f*x+2*e)) - 4/f*b*a*c^2*\ln(\exp(f*x+e)) + 2/f^2*b^2*c*d*\ln(1+\exp(2*f*x+2*e)) - 4/f^2*b^2*c*d*\ln(\exp(f*x+e)) + d*b^2*c*x^2 + b^2*c^2*x + a^2*d*c*x^2 + a^2*c^2*x - a*b*d^2*polylog(3,-\exp(2*f*x+2*e))/f^3 + 1/3*d^2*b^2*x^3 + 1/3/d*b^2*c^3 + 1/3*a^2*d^2*x^3 + 1/3*a^2/d*c^3 - 2/f*b^2*d^2*x^2 - 2/f^3*b^2*d^2*e^2 + 2/f^2*b*a*d^2*polylog(2,-\exp(2*f*x+2*e))*x + 2/f^2*b*c*d*a*polylog(2,-\exp(2*f*x+2*e)) - 4/f^2*b^2*d^2*e*x + b^2*d^2*polylog(2,-\exp(2*f*x+2*e))/f^3 + 8/3/f^3*b*a*d^2*e^3 + 4/f^3*b^2*e*d^2*\ln(\exp(f*x+e)) + 2/f^2*b^2*d^2*\ln(1+\exp(2*f*x+2*e))*x + 2/f*b*a*c^2*\ln(1+\exp(2*f*x+2*e)) - 2/3*d^2*a*b*x^3 + 2/3/d*c^3*a*b - 2*d*a*b*c*x^2 + 2*a*b*c^2*x - 8/f*b*c*d*a*e*x + 4/f*b*c*d*a*\ln(1+\exp(2*f*x+2*e))*x + 8/f^2*b*e*c*d*a*\ln(\exp(f*x+e))$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.28 (sec) , antiderivative size = 2123, normalized size of antiderivative = 10.06

$$\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = \text{Too large to display}$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] $1/3*((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 + 3*(a^2 - 2*a*b + b^2)*c*d*f^3*x^2 - 4*a*b*d^2*e^3 + 3*(a^2 - 2*a*b + b^2)*c^2*f^3*x + 6*b^2*d^2*e^2 - 6*(2*a*b*c^2*e - b^2*c^2)*f^2 + ((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 - 4*a*b*d^2*e^3 - 12*a*b*c^2*e*f^2 + 6*b^2*d^2*e^2 - 3*(2*b^2*d^2*f^2 - (a^2 - 2*a*b + b^2)*c*d*f^3)*x^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f - 3*(4*b^2*c*d*f^2 - (a^2 - 2*a*b + b^2)*c^2*f^3)*x)*\cosh(f*x + e)^2 + 2*((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 - 4*a*b*d^2*e^3 - 12*a*b*c^2*e*f^2 + 6*b^2*d^2*e^2 - 3*(2*b^2*d^2*f^2 - (a^2 - 2*a*b + b^2)*c*d*f^3)*x^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f - 3*(4*b^2*c$

$$\begin{aligned}
& *d*f^2 - (a^2 - 2*a*b + b^2)*c^2*f^3*x)*\cosh(f*x + e)*\sinh(f*x + e) + ((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 - 4*a*b*d^2*e^3 - 12*a*b*c^2*e*f^2 + 6*b^2*d^2*e^2 - 3*(2*b^2*d^2*f^2 - (a^2 - 2*a*b + b^2)*c*d*f^3)*x^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f - 3*(4*b^2*c*d*f^2 - (a^2 - 2*a*b + b^2)*c^2*f^3)*x)*\sinh(f*x + e)^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f + 6*(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2 + (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\cosh(f*x + e)^2 + 2*(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\cosh(f*x + e)*\sinh(f*x + e) + (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\sinh(f*x + e)^2)*\operatorname{dilog}(I*\cosh(f*x + e) + I*\sinh(f*x + e)) + 6*(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2 + (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\cosh(f*x + e)^2 + 2*(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\cosh(f*x + e)*\sinh(f*x + e) + (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*\sinh(f*x + e)^2)*\operatorname{dilog}(-I*\cosh(f*x + e) - I*\sinh(f*x + e)) + 6*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e + (a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\cosh(f*x + e)^2 + 2*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\sinh(f*x + e)^2 - (2*a*b*c*d*e - b^2*c*d)*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) + I) + 6*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e + (a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\cosh(f*x + e)^2 + 2*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*\sinh(f*x + e)^2 - (2*a*b*c*d*e - b^2*c*d)*f)*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) + 6*(a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\cosh(f*x + e)^2 + 2*(a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\sinh(f*x + e)^2 + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) + 6*(a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\cosh(f*x + e)^2 + 2*(a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*e*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\sinh(f*x + e)^2 + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1) - 12*(a*b*d^2*\cosh(f*x + e)^2 + 2*a*b*d^2*\cosh(f*x + e)*\sinh(f*x + e) + a*b*d^2*\sinh(f*x + e)^2 + a*b*d^2)*\operatorname{polylog}(3, I*\cosh(f*x + e) + I*\sinh(f*x + e)) - 12*(a*b*d^2*\cosh(f*x + e)^2 + 2*a*b*d^2*\cosh(f*x + e)*\sinh(f*x + e) + a*b*d^2*\sinh(f*x + e)^2 + a*b*d^2)*\operatorname{polylog}(3, -I*\cosh(f*x + e) - I*\sinh(f*x + e)))/(f^3*\cosh(f*x + e)^2 + 2*f^3*\cosh(f*x + e)*\sinh(f*x + e) + f^3*\sinh(f*x + e)^2 + f^3)
\end{aligned}$$

SymPy [F]

$$\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx)^2 dx$$

[In] integrate((d*x+c)**2*(a+b*tanh(f*x+e))**2,x)

[Out] Integral((a + b*tanh(e + f*x))**2*(c + d*x)**2, x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 411 vs. 2(203) = 406.

Time = 0.28 (sec) , antiderivative size = 411, normalized size of antiderivative = 1.95

$$\begin{aligned} \int (c + dx)^2 (a + b \tanh(e + fx))^2 dx &= \frac{1}{3} a^2 d^2 x^3 + a^2 c d x^2 + b^2 c^2 \left(x + \frac{e}{f} - \frac{2}{f(e^{-2fx-2e}) + 1} \right) \\ &+ a^2 c^2 x + b^2 c d \left(\frac{fx^2 + (fx^2 e^{(2e)} - 4xe^{(2e)})e^{(2fx)}}{fe^{(2fx+2e)} + f} + \frac{2 \log((e^{(2fx+2e)} + 1)e^{(-2e)})}{f^2} \right) \\ &+ \frac{2abc^2 \log(\cosh(fx + e))}{f} \\ &+ \frac{(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \operatorname{Li}_2(-e^{(2fx+2e)}) - \operatorname{Li}_3(-e^{(2fx+2e)}))abd^2}{f^3} \\ &+ \frac{(2abd^2 f + b^2 d^2 f)x^3 + 6(abcdf + b^2 d^2)x^2 + (6abcdfx^2 e^{(2e)} + (2abd^2 f e^{(2e)} + b^2 d^2 f e^{(2e)})x^3)e^{(2fx)}}{3(fe^{(2fx+2e)} + f)} \\ &+ \frac{(2abcdf + b^2 d^2)(2fx \log(e^{(2fx+2e)} + 1) + \operatorname{Li}_2(-e^{(2fx+2e)}))}{f^3} \\ &- \frac{2(2abd^2 f^3 x^3 + 3(2abcdf + b^2 d^2)f^2 x^2)}{3f^3} \end{aligned}$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] 1/3*a^2*d^2*x^3 + a^2*c*d*x^2 + b^2*c^2*(x + e/f - 2/(f*(e^(-2*f*x - 2*e) + 1))) + a^2*c^2*x + b^2*c*d*((f*x^2 + (f*x^2*e^(2*e) - 4*x*e^(2*e))*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) + 2*log((e^(2*f*x + 2*e) + 1)*e^(-2*e))/f^2) + 2*a*b*c^2*log(cosh(f*x + e))/f + (2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))*a*b*d^2/f^3 + 1/3*((2*a*b*d^2*f + b^2*d^2*f)*x^3 + 6*(a*b*c*d*f + b^2*d^2)*x^2 + (6*a*b*c*d*f*x^2*e^(2*e) + (2*a*b*d^2*f*e^(2*e) + b^2*d^2*f*e^(2*e))*x^3)*e^(2*f*x))/(f*e^(2*f*x + 2*e) + f) + (2*a*b*c*d*f + b^2*d^2)*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))/f^3 - 2/3*(2*a*b*d^2*f^3*x^3 + 3*(2*a*b*c*d*f + b^2*d^2)*f^2*x^2)/f^3

Giac [F]

$$\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = \int (dx + c)^2 (b \tanh(fx + e) + a)^2 dx$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx)^2 dx$$

[In] int((a + b*tanh(e + f*x))^2*(c + d*x)^2,x)

[Out] int((a + b*tanh(e + f*x))^2*(c + d*x)^2, x)

3.60 $\int (c + dx)(a + b \tanh(e + fx))^2 dx$

Optimal result	401
Rubi [A] (verified)	401
Mathematica [A] (verified)	403
Maple [A] (verified)	404
Fricas [C] (verification not implemented)	404
Sympy [F]	405
Maxima [F]	405
Giac [F]	405
Mupad [F(-1)]	406

Optimal result

Integrand size = 18, antiderivative size = 127

$$\begin{aligned} \int (c + dx)(a + b \tanh(e + fx))^2 dx = & b^2 cx + \frac{1}{2} b^2 dx^2 + \frac{a^2(c + dx)^2}{2d} - \frac{ab(c + dx)^2}{d} \\ & + \frac{2ab(c + dx) \log(1 + e^{2(e+fx)})}{f} \\ & + \frac{b^2 d \log(\cosh(e + fx))}{f^2} + \frac{abd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\ & - \frac{b^2(c + dx) \tanh(e + fx)}{f} \end{aligned}$$

[Out] $b^2cx + 1/2b^2dx^2 + 1/2a^2(dx+c)^2/d - a*b*(dx+c)^2/d + 2*a*b*(dx+c)*\ln(1 + \exp(2*fx + 2*e))/f + b^2d*\ln(\cosh(fx + e))/f^2 + a*b*d*\operatorname{polylog}(2, -\exp(2*fx + 2*e))/f^2 - b^2*(dx+c)*\tanh(fx + e)/f$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3803, 3799, 2221, 2317, 2438, 3801, 3556}

$$\begin{aligned} \int (c + dx)(a + b \tanh(e + fx))^2 dx = & \frac{a^2(c + dx)^2}{2d} + \frac{2ab(c + dx) \log(e^{2(e+fx)} + 1)}{f} \\ & - \frac{ab(c + dx)^2}{d} + \frac{abd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\ & - \frac{b^2(c + dx) \tanh(e + fx)}{f} + b^2 cx \\ & + \frac{b^2 d \log(\cosh(e + fx))}{f^2} + \frac{1}{2} b^2 dx^2 \end{aligned}$$

[In] Int[(c + d*x)*(a + b*Tanh[e + f*x])^2,x]

[Out] $b^2 c x + (b^2 d x^2)/2 + (a^2 (c + d x)^2)/(2 d) - (a b (c + d x)^2)/d + (2 a b (c + d x) \operatorname{Log}[1 + E^{2(e + f x)}])/f + (b^2 d \operatorname{Log}[\operatorname{Cosh}[e + f x]])/f^2 + (a b d \operatorname{PolyLog}[2, -E^{2(e + f x)}])/f^2 - (b^2 (c + d x) \operatorname{Tanh}[e + f x])/f$

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_)))^(n_))], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3556

Int[tan[(c_) + (d_)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3799

Int[(((c_) + (d_)*(x_))^(m_))*tan[(e_) + (Complex[0, fz_])*(f_)*(x_)], x_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x] /; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]

Rule 3801

Int[(((c_) + (d_)*(x_))^(m_))*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c + dx) + 2ab(c + dx) \tanh(e + fx) + b^2(c + dx) \tanh^2(e + fx)) dx \\
&= \frac{a^2(c + dx)^2}{2d} + (2ab) \int (c + dx) \tanh(e + fx) dx + b^2 \int (c + dx) \tanh^2(e + fx) dx \\
&= \frac{a^2(c + dx)^2}{2d} - \frac{ab(c + dx)^2}{d} - \frac{b^2(c + dx) \tanh(e + fx)}{f} \\
&\quad + (4ab) \int \frac{e^{2(e+fx)}(c + dx)}{1 + e^{2(e+fx)}} dx + b^2 \int (c + dx) dx + \frac{(b^2d) \int \tanh(e + fx) dx}{f} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c + dx)^2}{2d} - \frac{ab(c + dx)^2}{d} + \frac{2ab(c + dx) \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d \log(\cosh(e + fx))}{f^2} - \frac{b^2(c + dx) \tanh(e + fx)}{f} - \frac{(2abd) \int \log(1 + e^{2(e+fx)}) dx}{f} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c + dx)^2}{2d} - \frac{ab(c + dx)^2}{d} + \frac{2ab(c + dx) \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d \log(\cosh(e + fx))}{f^2} - \frac{b^2(c + dx) \tanh(e + fx)}{f} - \frac{(abd) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{f^2} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c + dx)^2}{2d} - \frac{ab(c + dx)^2}{d} + \frac{2ab(c + dx) \log(1 + e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d \log(\cosh(e + fx))}{f^2} + \frac{abd \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{b^2(c + dx) \tanh(e + fx)}{f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 6.79 (sec) , antiderivative size = 214, normalized size of antiderivative = 1.69

$$\begin{aligned}
&\int (c + dx)(a + b \tanh(e + fx))^2 dx \\
&= \frac{\cosh(e + fx) \left(-((a^2 + b^2)(e + fx)(-2cf + d(e - fx)) \cosh(e + fx)) + 2b \cosh(e + fx) \left(\frac{af^2(c+dx)^2}{d} + bd \right) \right)}{f^2}
\end{aligned}$$

[In] Integrate[(c + d*x)*(a + b*Tanh[e + f*x])^2,x]

[Out] (Cosh[e + f*x]*(-(a^2 + b^2)*(e + f*x)*(-2*c*f + d*(e - f*x))*Cosh[e + f*x]) + 2*b*Cosh[e + f*x]*((a*f^2*(c + d*x)^2)/d + b*d*(e + f*x) - 2*(b*d - 2*

$$a*d*e + 2*a*c*f)*(e + f*x) + 2*a*d*(e + f*x)*\text{Log}[1 + E^{-2*(e + f*x)}] + (b*d - 2*a*d*e + 2*a*c*f)*\text{Log}[1 + E^{2*(e + f*x)}] - a*d*\text{PolyLog}[2, -E^{-2*(e + f*x)}] - 2*b^2*f*(c + d*x)*\text{Sinh}[e + f*x]*(a + b*\text{Tanh}[e + f*x])^2/(2*f^2*(a*\text{Cosh}[e + f*x] + b*\text{Sinh}[e + f*x])^2)$$

Maple [A] (verified)

Time = 0.27 (sec) , antiderivative size = 221, normalized size of antiderivative = 1.74

method	result
risch	$\frac{a^2 d x^2}{2} - a b d x^2 + \frac{b^2 d x^2}{2} + a^2 c x + 2 a b c x + b^2 c x + \frac{2(d x+c) b^2}{f(1+e^{2 f x+2 e})} + \frac{b^2 d \ln(1+e^{2 f x+2 e})}{f^2} - \frac{2 b^2 d \ln(e^{f x+e})}{f^2} + \frac{2 b a c}{f^2}$

```
[In] int((d*x+c)*(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*a^2*d*x^2-a*b*d*x^2+1/2*b^2*d*x^2+a^2*c*x+2*a*b*c*x+b^2*c*x+2/f*(d*x+c)*b^2/(1+exp(2*f*x+2*e))+1/f^2*b^2*d*ln(1+exp(2*f*x+2*e))-2/f^2*b^2*d*ln(exp(f*x+e))+2/f*b*a*c*ln(1+exp(2*f*x+2*e))-4/f*b*a*c*ln(exp(f*x+e))+4/f^2*b*e*d*a*ln(exp(f*x+e))-4/f*b*d*a*e*x-2/f^2*b*d*a*e^2+2/f*b*d*a*ln(1+exp(2*f*x+2*e))*x+a*b*d*polylog(2,-exp(2*f*x+2*e))/f^2
```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.30 (sec) , antiderivative size = 944, normalized size of antiderivative = 7.43

$$\int (c + dx)(a + b \tanh(e + fx))^2 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] 1/2*((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 + 2*(a^2 - 2*a*b + b^2)*c*f^2*x - 4*b^2*d*e + ((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c*e*f - 4*b^2*d*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*cosh(f*x + e)^2 + 2*((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c*e*f - 4*b^2*d*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*cosh(f*x + e)*sinh(f*x + e) + ((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c*e*f - 4*b^2*d*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*sinh(f*x + e)^2 - 4*(2*a*b*c*e - b^2*c)*f + 4*(a*b*d*cosh(f*x + e)^2 + 2*a*b*d*cosh(f*x + e)*sinh(f*x + e) + a*b*d*sinh(f*x + e)^2 + a*b*d)*dilog(I*cosh(f*x + e) + I*sinh(f*x + e)) + 4*(a*b*d*cosh(f*x + e)^2 + 2*a*b*d*cosh(f*x + e)*sinh(f*x + e) + a*b*d*sinh(f*x + e)^2 + a*b*d)*dilog(-I*cosh(f*x + e) - I*sinh(f*x + e)) - 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d + (2*a*b*d*e - 2*a*b*c*f - b^2*d)*cosh(f*x + e)^2 + 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d)*cosh(f*x + e)*sinh(f*x + e) + (2*a*b*d*e - 2*a*b*c*f - b^2*d)*sinh(f*x + e)^2)*log(cosh(f*x + e) + sinh(f*x + e))
```

$e) + I) - 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d + (2*a*b*d*e - 2*a*b*c*f - b^2*d)*\cosh(f*x + e)^2 + 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d)*\cosh(f*x + e)*\sinh(f*x + e) + (2*a*b*d*e - 2*a*b*c*f - b^2*d)*\sinh(f*x + e)^2*\log(\cosh(f*x + e) + \sinh(f*x + e) - I) + 4*(a*b*d*f*x + a*b*d*e + (a*b*d*f*x + a*b*d*e)*\cosh(f*x + e)^2 + 2*(a*b*d*f*x + a*b*d*e)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d*f*x + a*b*d*e)*\sinh(f*x + e)^2)*\log(I*\cosh(f*x + e) + I*\sinh(f*x + e) + 1) + 4*(a*b*d*f*x + a*b*d*e + (a*b*d*f*x + a*b*d*e)*\cosh(f*x + e)^2 + 2*(a*b*d*f*x + a*b*d*e)*\cosh(f*x + e)*\sinh(f*x + e) + (a*b*d*f*x + a*b*d*e)*\sinh(f*x + e)^2)*\log(-I*\cosh(f*x + e) - I*\sinh(f*x + e) + 1))/(f^2*\cosh(f*x + e)^2 + 2*f^2*\cosh(f*x + e)*\sinh(f*x + e) + f^2*\sinh(f*x + e)^2 + f^2)$

Sympy [F]

$$\int (c + dx)(a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx) dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e))**2,x)

[Out] Integral((a + b*tanh(e + f*x))**2*(c + d*x), x)

Maxima [F]

$$\int (c + dx)(a + b \tanh(e + fx))^2 dx = \int (dx + c)(b \tanh(fx + e) + a)^2 dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] $\frac{1}{2}a^2d*x^2 + (x^2 - 4*\integrate(x/(e^{(2*f*x + 2*e)} + 1), x))*a*b*d + b^2*c*(x + e/f - 2/(f*(e^{(-2*f*x - 2*e)} + 1))) + a^2*c*x + 1/2*b^2*d*((f*x^2 + (f*x^2*e^{(2*e)} - 4*x*e^{(2*e)})*e^{(2*f*x)})/(f*e^{(2*f*x + 2*e)} + f) + 2*\log((e^{(2*f*x + 2*e)} + 1)*e^{(-2*e)})/f^2) + 2*a*b*c*\log(\cosh(f*x + e))/f$

Giac [F]

$$\int (c + dx)(a + b \tanh(e + fx))^2 dx = \int (dx + c)(b \tanh(fx + e) + a)^2 dx$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)*(b*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tanh(e + fx))^2 dx = \int (a + b \tanh(e + fx))^2 (c + dx) dx$$

```
[In] int((a + b*tanh(e + f*x))^2*(c + d*x),x)
```

```
[Out] int((a + b*tanh(e + f*x))^2*(c + d*x), x)
```

3.61 $\int \frac{(a+b \tanh(e+fx))^2}{c+dx} dx$

Optimal result	407
Rubi [N/A]	407
Mathematica [N/A]	408
Maple [N/A] (verified)	408
Fricas [N/A]	408
Sympy [N/A]	408
Maxima [N/A]	409
Giac [N/A]	409
Mupad [N/A]	409

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \text{Int}\left(\frac{(a + b \tanh(e + fx))^2}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))^2/(d*x+c), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx$$

[In] Int[(a + b*Tanh[e + f*x])^2/(c + d*x), x]

[Out] Defer[Int] [(a + b*Tanh[e + f*x])^2/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 26.59 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])^2/(c + d*x), x]

[Out] Integrate[(a + b*Tanh[e + f*x])^2/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tanh(fx + e))^2}{dx + c} dx$$

[In] int((a+b*tanh(f*x+e))^2/(d*x+c), x)

[Out] int((a+b*tanh(f*x+e))^2/(d*x+c), x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^2}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c), x, algorithm="fricas")

[Out] integral((b^2*tanh(f*x + e)^2 + 2*a*b*tanh(f*x + e) + a^2)/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 1.16 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx$$

[In] integrate((a+b*tanh(f*x+e))**2/(d*x+c), x)

[Out] Integral((a + b*tanh(e + f*x))**2/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.38 (sec) , antiderivative size = 155, normalized size of antiderivative = 7.75

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^2}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c),x, algorithm="maxima")

```
[Out] a^2*log(d*x + c)/d + 2*b^2/(d*f*x + c*f + (d*f*x*e^(2*e) + c*f*e^(2*e))*e^(2*f*x)) + (2*a*b + b^2)*log(d*x + c)/d - integrate(2*(2*a*b*d*f*x + 2*a*b*c*f - b^2*d)/(d^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2*e^(2*e) + 2*c*d*f*x*e^(2*e) + c^2*f*e^(2*e))*e^(2*f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^2}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c),x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)^2/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.98 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^2}{c + dx} dx$$

[In] int((a + b*tanh(e + f*x))^2/(c + d*x),x)

[Out] int((a + b*tanh(e + f*x))^2/(c + d*x), x)

3.62 $\int \frac{(a+b \tanh(e+fx))^2}{(c+dx)^2} dx$

Optimal result	410
Rubi [N/A]	410
Mathematica [N/A]	411
Maple [N/A] (verified)	411
Fricas [N/A]	411
Sympy [N/A]	412
Maxima [N/A]	412
Giac [N/A]	412
Mupad [N/A]	413

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tanh(e + fx))^2}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))^2/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx$$

[In] Int[(a + b*Tanh[e + f*x])^2/(c + d*x)^2,x]

[Out] Defer[Int] [(a + b*Tanh[e + f*x])^2/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 19.53 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])^2/(c + d*x)^2,x]

[Out] Integrate[(a + b*Tanh[e + f*x])^2/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tanh(fx + e))^2}{(dx + c)^2} dx$$

[In] int((a+b*tanh(f*x+e))^2/(d*x+c)^2,x)

[Out] int((a+b*tanh(f*x+e))^2/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b^2*tanh(f*x + e)^2 + 2*a*b*tanh(f*x + e) + a^2)/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 2.06 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))**2/(d*x+c)**2,x)

[Out] Integral((a + b*tanh(e + f*x))**2/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.42 (sec) , antiderivative size = 280, normalized size of antiderivative = 14.00

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c)^2,x, algorithm="maxima")

[Out] $-a^2/(d^2*x + c*d) - (2*a*b*c*f + (c*f - 2*d)*b^2 + (2*a*b*d*f + b^2*d*f)*x + (2*a*b*c*f*e^{(2*e)} + b^2*c*f*e^{(2*e)} + (2*a*b*d*f*e^{(2*e)} + b^2*d*f*e^{(2*e)})*x)*e^{(2*f*x)})/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f + (d^3*f*x^2*e^{(2*e)} + 2*c*d^2*f*x*e^{(2*e)} + c^2*d*f*e^{(2*e)})*e^{(2*f*x)}) - \text{integrate}(4*(a*b*d*f*x + a*b*c*f - b^2*d)/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3*e^{(2*e)} + 3*c*d^2*f*x^2*e^{(2*e)} + 3*c^2*d*f*x*e^{(2*e)} + c^3*f*e^{(2*e)})*e^{(2*f*x)}), x)$

Giac [N/A]

Not integrable

Time = 0.44 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^2/(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)^2/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 2.18 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + f x))^2}{(c + d x)^2} dx = \int \frac{(a + b \tanh(e + f x))^2}{(c + d x)^2} dx$$

```
[In] int((a + b*tanh(e + f*x))^2/(c + d*x)^2,x)
```

```
[Out] int((a + b*tanh(e + f*x))^2/(c + d*x)^2, x)
```

3.63 $\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx$

Optimal result	415
Rubi [A] (verified)	416
Mathematica [B] (verified)	423
Maple [B] (verified)	424
Fricas [C] (verification not implemented)	425
Sympy [F]	425
Maxima [B] (verification not implemented)	426
Giac [F]	427
Mupad [F(-1)]	427

Optimal result

Integrand size = 20, antiderivative size = 566

$$\begin{aligned}
\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = & -\frac{3b^3 d(c + dx)^2}{2f^2} - \frac{3ab^2(c + dx)^3}{f} + \frac{b^3(c + dx)^3}{2f} \\
& + \frac{a^3(c + dx)^4}{4d} - \frac{3a^2b(c + dx)^4}{4d} + \frac{3ab^2(c + dx)^4}{4d} \\
& - \frac{b^3(c + dx)^4}{4d} + \frac{3b^3 d^2(c + dx) \log(1 + e^{2(e+fx)})}{f^3} \\
& + \frac{9ab^2 d(c + dx)^2 \log(1 + e^{2(e+fx)})}{f^2} \\
& + \frac{3a^2 b(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
& + \frac{b^3(c + dx)^3 \log(1 + e^{2(e+fx)})}{f} \\
& + \frac{3b^3 d^3 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^4} \\
& + \frac{9ab^2 d^2(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
& + \frac{9a^2 b d(c + dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
& + \frac{3b^3 d(c + dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
& - \frac{9ab^2 d^3 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^4} \\
& - \frac{9a^2 b d^2(c + dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
& - \frac{3b^3 d^2(c + dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
& + \frac{9a^2 b d^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{4f^4} \\
& + \frac{3b^3 d^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{4f^4} \\
& - \frac{3b^3 d(c + dx)^2 \tanh(e + fx)}{2f^2} \\
& - \frac{3ab^2(c + dx)^3 \tanh(e + fx)}{f} \\
& - \frac{b^3(c + dx)^3 \tanh^2(e + fx)}{2f}
\end{aligned}$$

```
[Out] -3/2*b^3*d*(d*x+c)^2/f^2-3*a*b^2*(d*x+c)^3/f+1/2*b^3*(d*x+c)^3/f+1/4*a^3*(d
*x+c)^4/d-3/4*a^2*b*(d*x+c)^4/d+3/4*a*b^2*(d*x+c)^4/d-1/4*b^3*(d*x+c)^4/d+3
*b^3*d^2*(d*x+c)*ln(1+exp(2*f*x+2*e))/f^3+9*a*b^2*d*(d*x+c)^2*ln(1+exp(2*f*
x+2*e))/f^2+3*a^2*b*(d*x+c)^3*ln(1+exp(2*f*x+2*e))/f+b^3*(d*x+c)^3*ln(1+exp
(2*f*x+2*e))/f+3/2*b^3*d^3*polylog(2,-exp(2*f*x+2*e))/f^4+9*a*b^2*d^2*(d*x+
c)*polylog(2,-exp(2*f*x+2*e))/f^3+9/2*a^2*b*d*(d*x+c)^2*polylog(2,-exp(2*f*
x+2*e))/f^2+3/2*b^3*d*(d*x+c)^2*polylog(2,-exp(2*f*x+2*e))/f^2-9/2*a*b^2*d^
3*polylog(3,-exp(2*f*x+2*e))/f^4-9/2*a^2*b*d^2*(d*x+c)*polylog(3,-exp(2*f*x
+2*e))/f^3-3/2*b^3*d^2*(d*x+c)*polylog(3,-exp(2*f*x+2*e))/f^3+9/4*a^2*b*d^3
*polylog(4,-exp(2*f*x+2*e))/f^4+3/4*b^3*d^3*polylog(4,-exp(2*f*x+2*e))/f^4-
3/2*b^3*d*(d*x+c)^2*tanh(f*x+e)/f^2-3*a*b^2*(d*x+c)^3*tanh(f*x+e)/f-1/2*b^3
*(d*x+c)^3*tanh(f*x+e)^2/f
```

Rubi [A] (verified)

Time = 0.70 (sec) , antiderivative size = 566, normalized size of antiderivative = 1.00,
 number of steps used = 28, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules

used = {3803, 3799, 2221, 2611, 6744, 2320, 6724, 3801, 32, 2317, 2438}

$$\begin{aligned}
 \int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = & \frac{a^3(c + dx)^4}{4d} - \frac{9a^2bd^2(c + dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 & + \frac{9a^2bd(c + dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 & + \frac{3a^2b(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} \\
 & - \frac{3a^2b(c + dx)^4}{4d} + \frac{9a^2bd^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{4f^4} \\
 & + \frac{9ab^2d^2(c + dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
 & + \frac{9ab^2d(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f^2} \\
 & - \frac{3ab^2(c + dx)^3 \tanh(e + fx)}{f} - \frac{3ab^2(c + dx)^3}{f} \\
 & + \frac{3ab^2(c + dx)^4}{4d} - \frac{9ab^2d^3 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^4} \\
 & - \frac{3b^3d^2(c + dx) \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 & + \frac{3b^3d^2(c + dx) \log(e^{2(e+fx)} + 1)}{f^3} \\
 & + \frac{3b^3d(c + dx)^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 & - \frac{3b^3d(c + dx)^2 \tanh(e + fx)}{2f^2} \\
 & + \frac{b^3(c + dx)^3 \log(e^{2(e+fx)} + 1)}{f} \\
 & - \frac{b^3(c + dx)^3 \tanh^2(e + fx)}{2f} \\
 & - \frac{3b^3d(c + dx)^2}{2f^2} + \frac{b^3(c + dx)^3}{2f} - \frac{b^3(c + dx)^4}{4d} \\
 & + \frac{3b^3d^3 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^4} \\
 & + \frac{3b^3d^3 \operatorname{PolyLog}(4, -e^{2(e+fx)})}{4f^4}
 \end{aligned}$$

[In] Int[(c + d*x)^3*(a + b*Tanh[e + f*x])^3,x]

[Out] (-3*b^3*d*(c + d*x)^2)/(2*f^2) - (3*a*b^2*(c + d*x)^3)/f + (b^3*(c + d*x)^3)/(2*f) + (a^3*(c + d*x)^4)/(4*d) - (3*a^2*b*(c + d*x)^4)/(4*d) + (3*a*b^2*

$$\begin{aligned} & (c + d*x)^4/(4*d) - (b^3*(c + d*x)^4)/(4*d) + (3*b^3*d^2*(c + d*x)*\text{Log}[1 + \\ & E^{(2*(e + f*x))}])/f^3 + (9*a*b^2*d*(c + d*x)^2*\text{Log}[1 + E^{(2*(e + f*x))}])/f \\ & ^2 + (3*a^2*b*(c + d*x)^3*\text{Log}[1 + E^{(2*(e + f*x))}])/f + (b^3*(c + d*x)^3*\text{Lo} \\ & g[1 + E^{(2*(e + f*x))}])/f + (3*b^3*d^3*\text{PolyLog}[2, -E^{(2*(e + f*x))}])/(2*f^4 \\ &) + (9*a*b^2*d^2*(c + d*x)*\text{PolyLog}[2, -E^{(2*(e + f*x))}])/f^3 + (9*a^2*b*d*(\\ & c + d*x)^2*\text{PolyLog}[2, -E^{(2*(e + f*x))}])/(2*f^2) + (3*b^3*d*(c + d*x)^2*\text{Pol} \\ & y\text{Log}[2, -E^{(2*(e + f*x))}])/(2*f^2) - (9*a*b^2*d^3*\text{PolyLog}[3, -E^{(2*(e + f*x) \\ &))}])/(2*f^4) - (9*a^2*b*d^2*(c + d*x)*\text{PolyLog}[3, -E^{(2*(e + f*x))}])/(2*f^3) \\ & - (3*b^3*d^2*(c + d*x)*\text{PolyLog}[3, -E^{(2*(e + f*x))}])/(2*f^3) + (9*a^2*b*d^ \\ & 3*\text{PolyLog}[4, -E^{(2*(e + f*x))}])/(4*f^4) + (3*b^3*d^3*\text{PolyLog}[4, -E^{(2*(e + \\ & f*x))}])/(4*f^4) - (3*b^3*d*(c + d*x)^2*\text{Tanh}[e + f*x])/(2*f^2) - (3*a*b^2*(c \\ & + d*x)^3*\text{Tanh}[e + f*x])/f - (b^3*(c + d*x)^3*\text{Tanh}[e + f*x]^2)/(2*f) \end{aligned}$$
Rule 32

```
Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c
+ d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))), x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)
, x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)
*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(
m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = \int (a^3(c + dx)^3 + 3a^2b(c + dx)^3 \tanh(e + fx) + 3ab^2(c + dx)^3 \tanh^2(e + fx) + b^3(c + dx)^3 \tanh^3(e + fx)) dx$$

$$\begin{aligned}
&= \frac{a^3(c+dx)^4}{4d} + (3a^2b) \int (c+dx)^3 \tanh(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx)^3 \tanh^2(e+fx) dx + b^3 \int (c+dx)^3 \tanh^3(e+fx) dx \\
&= \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} - \frac{3ab^2(c+dx)^3 \tanh(e+fx)}{f} - \frac{b^3(c+dx)^3 \tanh^2(e+fx)}{2f} \\
&\quad + (6a^2b) \int \frac{e^{2(e+fx)}(c+dx)^3}{1+e^{2(e+fx)}} dx + (3ab^2) \int (c+dx)^3 dx + b^3 \int (c+dx)^3 \tanh(e \\
&\quad\quad\quad + fx) dx \\
&\quad + \frac{(9ab^2d) \int (c+dx)^2 \tanh(e+fx) dx}{f} + \frac{(3b^3d) \int (c+dx)^2 \tanh^2(e+fx) dx}{2f} \\
&= -\frac{3ab^2(c+dx)^3}{f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} \\
&\quad - \frac{b^3(c+dx)^4}{4d} + \frac{3a^2b(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad - \frac{3b^3d(c+dx)^2 \tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3 \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3 \tanh^2(e+fx)}{2f} + (2b^3) \int \frac{e^{2(e+fx)}(c+dx)^3}{1+e^{2(e+fx)}} dx \\
&\quad + \frac{(3b^3d^2) \int (c+dx) \tanh(e+fx) dx}{f^2} - \frac{(9a^2bd) \int (c+dx)^2 \log(1+e^{2(e+fx)}) dx}{f} \\
&\quad + \frac{(18ab^2d) \int \frac{e^{2(e+fx)}(c+dx)^2}{1+e^{2(e+fx)}} dx}{f} + \frac{(3b^3d) \int (c+dx)^2 dx}{2f} \\
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&\quad + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} + \frac{9ab^2d(c+dx)^2 \log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2b(c+dx)^3 \log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^3 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3b^3d(c+dx)^2 \tanh(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)^3 \tanh(e+fx)}{f} - \frac{b^3(c+dx)^3 \tanh^2(e+fx)}{2f} \\
&\quad - \frac{(9a^2bd^2) \int (c+dx) \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(18ab^2d^2) \int (c+dx) \log(1+e^{2(e+fx)}) dx}{f^2} \\
&\quad + \frac{(6b^3d^2) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx}{f^2} - \frac{(3b^3d) \int (c+dx)^2 \log(1+e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&+ \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} + \frac{3b^3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} \\
&+ \frac{9ab^2d(c+dx)^2\log(1+e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&+ \frac{b^3(c+dx)^3\log(1+e^{2(e+fx)})}{f} + \frac{9ab^2d^2(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^3} \\
&+ \frac{9a^2bd(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} + \frac{3b^3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} \\
&- \frac{9a^2bd^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} - \frac{3b^3d(c+dx)^2\tanh(e+fx)}{2f^2} \\
&- \frac{3ab^2(c+dx)^3\tanh(e+fx)}{f} - \frac{b^3(c+dx)^3\tanh^2(e+fx)}{2f} \\
&+ \frac{(9a^2bd^3)\int\text{PolyLog}(3,-e^{2(e+fx)})dx}{2f^3} - \frac{(9ab^2d^3)\int\text{PolyLog}(2,-e^{2(e+fx)})dx}{f^3} \\
&- \frac{(3b^3d^3)\int\log(1+e^{2(e+fx)})dx}{f^3} - \frac{(3b^3d^2)\int(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})dx}{f^2} \\
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&+ \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} + \frac{3b^3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} \\
&+ \frac{9ab^2d(c+dx)^2\log(1+e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&+ \frac{b^3(c+dx)^3\log(1+e^{2(e+fx)})}{f} + \frac{9ab^2d^2(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^3} \\
&+ \frac{9a^2bd(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} + \frac{3b^3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} \\
&- \frac{9a^2bd^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} - \frac{3b^3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} \\
&- \frac{3b^3d(c+dx)^2\tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3\tanh(e+fx)}{f} \\
&- \frac{b^3(c+dx)^3\tanh^2(e+fx)}{2f} + \frac{(9a^2bd^3)\text{Subst}\left(\int\frac{\text{PolyLog}(3,-x)}{x}dx,x,e^{2(e+fx)}\right)}{4f^4} \\
&- \frac{(9ab^2d^3)\text{Subst}\left(\int\frac{\text{PolyLog}(2,-x)}{x}dx,x,e^{2(e+fx)}\right)}{2f^4} \\
&- \frac{(3b^3d^3)\text{Subst}\left(\int\frac{\log(1+x)}{x}dx,x,e^{2(e+fx)}\right)}{2f^4} + \frac{(3b^3d^3)\int\text{PolyLog}(3,-e^{2(e+fx)})dx}{2f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} \\
&\quad + \frac{3b^3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2\log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2b(c+dx)^3\log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^3d^3\text{PolyLog}(2,-e^{2(e+fx)})}{2f^4} + \frac{9ab^2d^2(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9a^2bd(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} + \frac{3b^3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{9ab^2d^3\text{PolyLog}(3,-e^{2(e+fx)})}{2f^4} - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{3b^3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} + \frac{9a^2bd^3\text{PolyLog}(4,-e^{2(e+fx)})}{4f^4} \\
&\quad - \frac{3b^3d(c+dx)^2\tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3\tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3\tanh^2(e+fx)}{2f} + \frac{(3b^3d^3)\text{Subst}\left(\int\frac{\text{PolyLog}(3,-x)}{x}dx,x,e^{2(e+fx)}\right)}{4f^4} \\
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} \\
&\quad + \frac{3b^3d^2(c+dx)\log(1+e^{2(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2\log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2b(c+dx)^3\log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^3\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^3d^3\text{PolyLog}(2,-e^{2(e+fx)})}{2f^4} + \frac{9ab^2d^2(c+dx)\text{PolyLog}(2,-e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9a^2bd(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} + \frac{3b^3d(c+dx)^2\text{PolyLog}(2,-e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{9ab^2d^3\text{PolyLog}(3,-e^{2(e+fx)})}{2f^4} - \frac{9a^2bd^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{3b^3d^2(c+dx)\text{PolyLog}(3,-e^{2(e+fx)})}{2f^3} + \frac{9a^2bd^3\text{PolyLog}(4,-e^{2(e+fx)})}{4f^4} \\
&\quad + \frac{3b^3d^3\text{PolyLog}(4,-e^{2(e+fx)})}{4f^4} - \frac{3b^3d(c+dx)^2\tanh(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)^3\tanh(e+fx)}{f} - \frac{b^3(c+dx)^3\tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2010 vs. 2(566) = 1132.

Time = 7.82 (sec) , antiderivative size = 2010, normalized size of antiderivative = 3.55

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \text{Result too large to show}$$

[In] Integrate[(c + d*x)^3*(a + b*Tanh[e + f*x])^3,x]

[Out] (b*E^(2*e)*(-24*b^2*c*d^2*x - 72*a*b*c^2*d*f*x - 24*a^2*c^3*f^2*x - 8*b^2*c^3*f^2*x - 12*b^2*d^3*x^2 - 72*a*b*c*d^2*f*x^2 - 36*a^2*c^2*d*f^2*x^2 - 12*b^2*c^2*d*f^2*x^2 - 24*a*b*d^3*f*x^3 - 24*a^2*c*d^2*f^2*x^3 - 8*b^2*c*d^2*f^2*x^3 - 6*a^2*d^3*f^2*x^4 - 2*b^2*d^3*f^2*x^4 + 36*a*b*c^2*d*Log[1 + E^(2*(e + f*x))] + (36*a*b*c^2*d*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (12*b^2*c*d^2*Log[1 + E^(2*(e + f*x))])/f + (12*b^2*c*d^2*Log[1 + E^(2*(e + f*x))])/(E^(2*e)*f) + 12*a^2*c^3*f*Log[1 + E^(2*(e + f*x))] + 4*b^2*c^3*f*Log[1 + E^(2*(e + f*x))] + (12*a^2*c^3*f*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (4*b^2*c^3*f*Log[1 + E^(2*(e + f*x))])/E^(2*e) + 72*a*b*c*d^2*x*Log[1 + E^(2*(e + f*x))] + (72*a*b*c*d^2*x*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (12*b^2*d^3*x*Log[1 + E^(2*(e + f*x))])/f + (12*b^2*d^3*x*Log[1 + E^(2*(e + f*x))])/(E^(2*e)*f) + 36*a^2*c^2*d*f*x*Log[1 + E^(2*(e + f*x))] + 12*b^2*c^2*d*f*x*Log[1 + E^(2*(e + f*x))] + (36*a^2*c^2*d*f*x*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (12*b^2*c^2*d*f*x*Log[1 + E^(2*(e + f*x))])/E^(2*e) + 36*a*b*d^3*x^2*Log[1 + E^(2*(e + f*x))] + (36*a*b*d^3*x^2*Log[1 + E^(2*(e + f*x))])/E^(2*e) + 36*a^2*c*d^2*f*x^2*Log[1 + E^(2*(e + f*x))] + 12*b^2*c*d^2*f*x^2*Log[1 + E^(2*(e + f*x))] + (36*a^2*c*d^2*f*x^2*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (12*b^2*c*d^2*f*x^2*Log[1 + E^(2*(e + f*x))])/E^(2*e) + 12*a^2*d^3*f*x^3*Log[1 + E^(2*(e + f*x))] + 4*b^2*d^3*f*x^3*Log[1 + E^(2*(e + f*x))] + (12*a^2*d^3*f*x^3*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (4*b^2*d^3*f*x^3*Log[1 + E^(2*(e + f*x))])/E^(2*e) + (6*d*(1 + E^(2*e))*(6*a*b*d*f*(c + d*x) + 3*a^2*f^2*(c + d*x)^2 + b^2*(d^2 + c^2*f^2 + 2*c*d*f^2*x + d^2*f^2*x^2))*PolyLog[2, -E^(2*(e + f*x))]/(E^(2*e)*f^2) - (6*d^2*(1 + E^(2*e))*(3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*PolyLog[3, -E^(2*(e + f*x))]/(E^(2*e)*f^2) + (9*a^2*d^3*PolyLog[4, -E^(2*(e + f*x))]/f^2 + (3*b^2*d^3*PolyLog[4, -E^(2*(e + f*x))])/f^2 + (9*a^2*d^3*PolyLog[4, -E^(2*(e + f*x))])/(E^(2*e)*f^2) + (3*b^2*d^3*PolyLog[4, -E^(2*(e + f*x))])/(E^(2*e)*f^2))/(4*(1 + E^(2*e))*f^2) + ((b^3*c^3 + 3*b^3*c^2*d*x + 3*b^3*c*d^2*x^2 + b^3*d^3*x^3)*Sech[e + f*x]^2)/(2*f) + (3*x^2*(a^3*c^2*d - 3*a^2*b*c^2*d + 3*a*b^2*c^2*d - b^3*c^2*d + a^3*c^2*d*Cosh[2*e] + 3*a^2*b*c^2*d*Cosh[2*e] + 3*a*b^2*c^2*d*Cosh[2*e] + b^3*c^2*d*Cosh[2*e] + a^3*c^2*d*Sinh[2*e] + 3*a^2*b*c^2*d*Sinh[2*e] + 3*a*b^2*c^2*d*Sinh[2*e] + b^3*c^2*d*Sinh[2*e]))/(2*(1 + Cosh[2*e] + Sinh[2*e])) + (x^3*(a^3*c*d^2 - 3*a^2*b*c*d^2 + 3*a*b^2*c*d^2 - b^3*c*d^2 + a^3*c*d^2*Cosh[2*e] + 3*a^2*b*c*d^2*Cosh[2*e] + 3*a*b^2*c*d^2*Cosh[2*e] + b^3*c*d^2*Cosh[2*e] + a^3*c*d^2*Sinh[2*e] + 3*a^2*b*c*d^2*Sinh[2*e] + 3*a*b^2*c*d^2*Sinh[2*e] + 3*a*b^2*c*d^2*Sin

$$\begin{aligned} & h[2e] + b^3 c d^2 \operatorname{Sinh}[2e]) / (1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e]) + (x^4 (a^3 d^3 - 3a^2 b d^3 + 3a b^2 d^3 - b^3 d^3 + a^3 d^3 \operatorname{Cosh}[2e] + 3a^2 b d^3 \operatorname{Cosh}[2e] + 3a b^2 d^3 \operatorname{Cosh}[2e] + b^3 d^3 \operatorname{Cosh}[2e] + a^3 d^3 \operatorname{Sinh}[2e] + 3a^2 b d^3 \operatorname{Sinh}[2e] + 3a b^2 d^3 \operatorname{Sinh}[2e] + b^3 d^3 \operatorname{Sinh}[2e])) / (4(1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e])) + x(a^3 c^3 + 3a b^2 c^3 - (3a^2 b c^3)) / (1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e]) + (3a^2 b c^3 \operatorname{Cosh}[2e] + 3a^2 b c^3 \operatorname{Sinh}[2e]) / (1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e]) + (2b^3 c^3 \operatorname{Cosh}[2e] + 2b^3 c^3 \operatorname{Sinh}[2e]) / ((1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e]) * (1 - \operatorname{Cosh}[2e] + \operatorname{Cosh}[4e] - \operatorname{Sinh}[2e] + \operatorname{Sinh}[4e])) + (-2b^3 c^3 \operatorname{Cosh}[4e] - 2b^3 c^3 \operatorname{Sinh}[4e]) / ((1 + \operatorname{Cosh}[2e] + \operatorname{Sinh}[2e]) * (1 - \operatorname{Cosh}[2e] + \operatorname{Cosh}[4e] - \operatorname{Sinh}[2e] + \operatorname{Sinh}[4e])) - (b^3 c^3) / (1 + \operatorname{Cosh}[6e] + \operatorname{Sinh}[6e]) + (b^3 c^3 \operatorname{Cosh}[6e] + b^3 c^3 \operatorname{Sinh}[6e]) / (1 + \operatorname{Cosh}[6e] + \operatorname{Sinh}[6e])) - (3 \operatorname{Sech}[e] * \operatorname{Sech}[e + f*x]) * (b^3 c^2 d \operatorname{Sinh}[f*x] + 2a b^2 c^3 f \operatorname{Sinh}[f*x] + 2b^3 c d^2 x \operatorname{Sinh}[f*x] + 6a b^2 c^2 d f x \operatorname{Sinh}[f*x] + b^3 d^3 x^2 \operatorname{Sinh}[f*x] + 6a b^2 c d^2 f x^2 \operatorname{Sinh}[f*x] + 2a b^2 d^3 f x^3 \operatorname{Sinh}[f*x])) / (2f^2) \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1833 vs. $2(534) = 1068$.

Time = 0.62 (sec) , antiderivative size = 1834, normalized size of antiderivative = 3.24

method	result	size
risch	Expression too large to display	1834

[In] `int((d*x+c)^3*(a+b*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -18/f^2 b^2 a c d^2 x^2 - 9/f^2 b d^2 c^2 a^2 e^2 + 3/2/f^2 b^3 c^2 d^2 \operatorname{polylog}(2, -\exp(2f*x+2e)) - 3/2/f^3 b^3 d^2 c^2 \operatorname{polylog}(3, -\exp(2f*x+2e)) + 1/f^2 b^3 d^3 \ln(1 + \exp(2f*x+2e)) * x^3 + 3/2/f^2 b^3 d^3 \operatorname{polylog}(2, -\exp(2f*x+2e)) * x^2 - 3/2/f^3 b^3 d^3 \operatorname{polylog}(3, -\exp(2f*x+2e)) * x + 6/f^4 b^3 e d^3 \ln(\exp(f*x+e)) + 3/f^3 b^3 d^3 \ln(1 + \exp(2f*x+2e)) * x + 2/f^4 b^3 e^3 d^3 \ln(\exp(f*x+e)) + 3/f^2 b^2 a^2 c^3 \ln(1 + \exp(2f*x+2e)) - 6/f^2 b^2 a^2 c^3 \ln(\exp(f*x+e)) - 3d^2 a^2 b c^2 x^3 + 3d^2 a^2 b^2 c^2 x^3 - 9/2 d a^2 b c^2 x^2 + 9/2 d a b^2 c^2 x^2 + 3a^2 b c^3 x + 3a b^2 c^3 x - 6/f^2 b^2 a d^3 x^3 - 3/f^2 b^3 c^2 d e^2 - 2/f^3 b^3 e^3 d^3 x + 4/f^3 b^3 e^3 d^2 c + 3/f^3 b^3 d^2 c \ln(1 + \exp(2f*x+2e)) - 6/f^3 b^3 d^2 c \ln(\exp(f*x+e)) - 6/f^3 b^3 e^3 a^2 d^3 x + 18/f^3 b^2 e^2 a d^3 x - 6/f^2 b^3 c^2 d e x - 18/f^3 b^2 a^2 c d^2 e^2 + 6/f^2 b^3 e^2 d^2 c x + 12/f^3 b^3 e^3 d^2 c a^2 + 9/f^2 b^2 a d^3 \ln(1 + \exp(2f*x+2e)) * x^2 + 9/f^3 b^2 a d^3 \operatorname{polylog}(2, -\exp(2f*x+2e)) * x + 3/f^2 b^2 a^2 d^3 \ln(1 + \exp(2f*x+2e)) * x^3 + 9/2/f^2 b^2 a^2 d^3 \operatorname{polylog}(2, -\exp(2f*x+2e)) * x^2 - 9/2/f^3 b^2 a^2 d^3 \operatorname{polylog}(3, -\exp(2f*x+2e)) * x + 6/f^4 b^2 e^3 a^2 d^3 \ln(\exp(f*x+e)) - 6/f^3 b^3 e^2 d^2 c \ln(\exp(f*x+e)) + 9/2/f^2 b d^2 c^2 a^2 \operatorname{polylog}(2, -\exp(2f*x+2e)) + 9/f^2 b^2 a^2 c^2 d \ln(1 + \exp(2f*x+2e)) + 6/f^2 b^3 e^2 c^2 d \ln(\exp(f*x+e)) + 3/f^2 b^3 c^2 d \ln(1 + \exp(2f*x+2e)) * x - 9/2/f^3 b d^2 c a^2 \operatorname{polylog}(3, -\exp(2f*x+2e)) + 3/f^2 b^3 d^2 c \ln(1 + \exp(2f*x+2e)) * x^2 + 3/f^2 b^3 d^2 c \operatorname{polylog}(2, -\exp(2f*x+2e)) * x + 1/4 d^3 a^3 x^4 - 1/4 d^3 b^3 x^4 + 1/4 d^3 \end{aligned}$$


```

c^4*a^3+1/4/d*c^4*b^3-36/f^2*b^2*a*c*d^2*e*x-18/f*b*d*c^2*a^2*e*x+18/f^2*b*
e^2*d^2*c*a^2*x+18/f^2*b^2*a*c*d^2*ln(1+exp(2*f*x+2*e))*x+36/f^3*b^2*e*a*c*
d^2*ln(exp(f*x+e))-18/f^3*b*e^2*d^2*c*a^2*ln(exp(f*x+e))+9/f*b*d*c^2*a^2*ln
(1+exp(2*f*x+2*e))*x+9/f*b*d^2*c*a^2*ln(1+exp(2*f*x+2*e))*x^2+9/f^2*b*d^2*c
*a^2*polylog(2,-exp(2*f*x+2*e))*x+18/f^2*b*e*d*c^2*a^2*ln(exp(f*x+e))+12/f^
4*b^2*e^3*a*d^3-6/f^3*b^3*d^3*e*x-9/2/f^4*b*e^4*a^2*d^3-3/4*d^3*a^2*b*x^4+3
/4*d^3*a*b^2*x^4+d^2*a^3*c*x^3-d^2*b^3*c*x^3+3/2*d*a^3*c^2*x^2-3/2*d*b^3*c^
2*x^2+a^3*c^3*x+b^3*c^3*x+3/4/d*c^4*a^2*b+3/4/d*c^4*a*b^2+b^2*(6*a*d^3*f*x^
3*exp(2*f*x+2*e)+2*b*d^3*f*x^3*exp(2*f*x+2*e)+18*a*c*d^2*f*x^2*exp(2*f*x+2*
e)+6*b*c*d^2*f*x^2*exp(2*f*x+2*e)+18*a*c^2*d*f*x*exp(2*f*x+2*e)+6*a*d^3*f*x
^3+6*b*c^2*d*f*x*exp(2*f*x+2*e)+3*b*d^3*x^2*exp(2*f*x+2*e)+6*a*c^3*f*exp(2*
f*x+2*e)+18*a*c*d^2*f*x^2+2*b*c^3*f*exp(2*f*x+2*e)+6*b*c*d^2*x*exp(2*f*x+2*
e)+18*a*c^2*d*f*x+3*exp(2*f*x+2*e)*d*b*c^2+3*b*d^3*x^2+6*a*c^3*f+6*b*c*d^2*
x+3*b*c^2*d)/f^2/(1+exp(2*f*x+2*e))^2-9/2*a*b^2*d^3*polylog(3,-exp(2*f*x+2*
e))/f^4+9/4*a^2*b*d^3*polylog(4,-exp(2*f*x+2*e))/f^4-3/2/f^4*b^3*e^4*d^3-3/
f^4*b^3*e^2*d^3-3/f^2*b^3*d^3*x^2+1/f*b^3*c^3*ln(1+exp(2*f*x+2*e))-2/f*b^3*
c^3*ln(exp(f*x+e))+3/2*b^3*d^3*polylog(2,-exp(2*f*x+2*e))/f^4+3/4*b^3*d^3*p
olylog(4,-exp(2*f*x+2*e))/f^4+9/f^3*b^2*a*c*d^2*polylog(2,-exp(2*f*x+2*e))-
18/f^4*b^2*e^2*a*d^3*ln(exp(f*x+e))-18/f^2*b^2*a*c^2*d*ln(exp(f*x+e))

```

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.42 (sec) , antiderivative size = 12909, normalized size of antiderivative = 22.81

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^3,x, algorithm="fricas")
```

```
[Out] Too large to include
```

Sympy [F]

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx)^3 dx$$

```
[In] integrate((d*x+c)**3*(a+b*tanh(f*x+e))**3,x)
```

```
[Out] Integral((a + b*tanh(e + f*x))**3*(c + d*x)**3, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1297 vs. $2(530) = 1060$.

Time = 0.39 (sec) , antiderivative size = 1297, normalized size of antiderivative = 2.29

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \text{Too large to display}$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^3,x, algorithm="maxima")

[Out] $\frac{1}{4}a^3d^3x^4 + a^3cd^2x^3 + \frac{3}{2}a^3c^2dx^2 + b^3c^3(x + e/f + \log(e^{-2fx - 2e} + 1)/f + 2e^{-2fx - 2e}/(f(2e^{-2fx - 2e} + e^{-4fx - 4e} + 1))) + a^3c^3x + 3a^2b^3c^3 \log(\cosh(fx + e))/f + \frac{1}{4}(24ab^2c^3f + 12b^3c^2d + (3a^2b^3d^3f^2 + 3ab^2d^3f^2 + b^3d^3f^2))x^4 + 4(3a^2b^3cd^2f^2 + b^3cd^2f^2 + 3(c^2d^2f^2 + 2d^3f)ab^2)x^3 + 6(3a^2b^3c^2d^2f^2 + 3(c^2d^2f^2 + 4cd^2f)ab^2 + (c^2d^2f^2 + 2d^3)b^3)x^2 + 12(2b^3cd^2 + (c^3f^2 + 6c^2d^2f)ab^2)x + (12ab^2c^3f^2xe^{4e} + (3a^2b^3d^3f^2e^{4e} + 3ab^2d^3f^2e^{4e} + b^3d^3f^2e^{4e}))x^4 + 4(3a^2b^3cd^2f^2e^{4e} + 3ab^2c^2d^2f^2e^{4e} + b^3cd^2f^2e^{4e})x^3 + 6(3a^2b^3c^2d^2f^2e^{4e} + 3ab^2c^2d^2f^2e^{4e} + b^3c^2d^2f^2e^{4e})x^2)e^{4fx} + 2(12ab^2c^3f^2e^{2e} + 6b^3c^2d^2e^{2e} + (3a^2b^3d^3f^2e^{2e} + 3ab^2d^3f^2e^{2e} + b^3d^3f^2e^{2e}))x^4 + 4(3a^2b^3cd^2f^2e^{2e} + 3(c^2d^2f^2e^{2e} + d^3f^2e^{2e})ab^2 + (c^2d^2f^2e^{2e} + 2cd^2f^2e^{2e} + d^3e^{2e})b^3)x^2 + 12((c^3f^2e^{2e} + 3c^2d^2f^2e^{2e})ab^2 + (c^2d^2f^2e^{2e} + cd^2e^{2e})b^3)x)e^{(2fx)}/(f^2e^{(4fx + 4e)} + 2f^2e^{(2fx + 2e)} + f^2) - 6(3ab^2c^2d^2f + b^3cd^2)x/f^2 + 3(3ab^2c^2d^2f + b^3cd^2) \log(e^{(2fx + 2e)} + 1)/f^3 + \frac{1}{3}(4f^3x^3 \log(e^{(2fx + 2e)} + 1) + 6f^2x^2 \operatorname{dilog}(-e^{(2fx + 2e)}) - 6fx \operatorname{polylog}(3, -e^{(2fx + 2e)}) + 3 \operatorname{polylog}(4, -e^{(2fx + 2e)})) \cdot (3a^2b^3d^3 + b^3d^3)/f^4 + \frac{3}{2}(3a^2b^3cd^2f + b^3cd^2f + 3ab^2d^3) \cdot (2f^2x^2 \log(e^{(2fx + 2e)} + 1) + 2fx \operatorname{dilog}(-e^{(2fx + 2e)}) - \operatorname{polylog}(3, -e^{(2fx + 2e)}))/f^4 + \frac{3}{2}(3a^2b^3c^2d^2f^2 + 6ab^2c^2d^2f + (c^2d^2f^2 + d^3)b^3) \cdot (2fx \log(e^{(2fx + 2e)} + 1) + \operatorname{dilog}(-e^{(2fx + 2e)}))/f^4 - \frac{1}{2}((3a^2b^3d^3 + b^3d^3)f^4x^4 + 4(3a^2b^3cd^2f + b^3cd^2f + 3ab^2d^3)f^3x^3 + 6(3a^2b^3c^2d^2f^2 + 6ab^2c^2d^2f + (c^2d^2f^2 + d^3)b^3)f^2x^2)/f^4$

Giac [F]

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \int (dx + c)^3 (b \tanh(fx + e) + a)^3 dx$$

[In] integrate((d*x+c)^3*(a+b*tanh(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^3*(b*tanh(f*x + e) + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx)^3 dx$$

[In] int((a + b*tanh(e + f*x))^3*(c + d*x)^3,x)

[Out] int((a + b*tanh(e + f*x))^3*(c + d*x)^3, x)

3.64 $\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx$

Optimal result	429
Rubi [A] (verified)	430
Mathematica [B] (verified)	435
Maple [B] (verified)	436
Fricas [C] (verification not implemented)	437
Sympy [F]	437
Maxima [B] (verification not implemented)	438
Giac [F]	439
Mupad [F(-1)]	439

Optimal result

Integrand size = 20, antiderivative size = 405

$$\begin{aligned}
 \int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = & \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2 (c + dx)^2}{f} + \frac{a^3 (c + dx)^3}{3d} \\
 & - \frac{a^2 b (c + dx)^3}{d} + \frac{ab^2 (c + dx)^3}{d} - \frac{b^3 (c + dx)^3}{3d} \\
 & + \frac{6ab^2 d (c + dx) \log(1 + e^{2(e+fx)})}{f^2} \\
 & + \frac{3a^2 b (c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{b^3 (c + dx)^2 \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{b^3 d^2 \log(\cosh(e + fx))}{f^3} \\
 & + \frac{3ab^2 d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} \\
 & + \frac{3a^2 b d (c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
 & + \frac{b^3 d (c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
 & - \frac{3a^2 b d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 & - \frac{b^3 d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
 & - \frac{b^3 d (c + dx) \tanh(e + fx)}{f^2} \\
 & - \frac{3ab^2 (c + dx)^2 \tanh(e + fx)}{f} \\
 & - \frac{b^3 (c + dx)^2 \tanh^2(e + fx)}{2f}
 \end{aligned}$$

```

[Out] b^3*c*d*x/f+1/2*b^3*d^2*x^2/f-3*a*b^2*(d*x+c)^2/f+1/3*a^3*(d*x+c)^3/d-a^2*b
*(d*x+c)^3/d+a*b^2*(d*x+c)^3/d-1/3*b^3*(d*x+c)^3/d+6*a*b^2*d*(d*x+c)*ln(1+e
xp(2*f*x+2*e))/f^2+3*a^2*b*(d*x+c)^2*ln(1+exp(2*f*x+2*e))/f+b^3*(d*x+c)^2*ln
(1+exp(2*f*x+2*e))/f+b^3*d^2*ln(cosh(f*x+e))/f^3+3*a*b^2*d^2*polylog(2,-ex
p(2*f*x+2*e))/f^3+3*a^2*b*d*(d*x+c)*polylog(2,-exp(2*f*x+2*e))/f^2+b^3*d*(d
*x+c)*polylog(2,-exp(2*f*x+2*e))/f^2-3/2*a^2*b*d^2*polylog(3,-exp(2*f*x+2*e
))/f^3-1/2*b^3*d^2*polylog(3,-exp(2*f*x+2*e))/f^3-b^3*d*(d*x+c)*tanh(f*x+e)
/f^2-3*a*b^2*(d*x+c)^2*tanh(f*x+e)/f-1/2*b^3*(d*x+c)^2*tanh(f*x+e)^2/f

```

Rubi [A] (verified)

Time = 0.48 (sec) , antiderivative size = 405, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules used = {3803, 3799, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32, 3556}

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \frac{a^3(c + dx)^3}{3d} + \frac{3a^2bd(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{3a^2b(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{a^2b(c + dx)^3}{d} - \frac{3a^2bd^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{6ab^2d(c + dx) \log(e^{2(e+fx)} + 1)}{f^2} - \frac{3ab^2(c + dx)^2 \tanh(e + fx)}{f} - \frac{3ab^2(c + dx)^2}{f} + \frac{ab^2(c + dx)^3}{d} + \frac{3ab^2d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{b^3d(c + dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{b^3d(c + dx) \tanh(e + fx)}{f^2} + \frac{b^3(c + dx)^2 \log(e^{2(e+fx)} + 1)}{f} - \frac{b^3(c + dx)^2 \tanh^2(e + fx)}{2f} + \frac{b^3cdx}{f} - \frac{b^3(c + dx)^3}{3d} - \frac{b^3d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} + \frac{b^3d^2 \log(\cosh(e + fx))}{f^3} + \frac{b^3d^2x^2}{2f}$$

[In] Int[(c + d*x)^2*(a + b*Tanh[e + f*x])^3,x]

[Out] (b^3*c*d*x)/f + (b^3*d^2*x^2)/(2*f) - (3*a*b^2*(c + d*x)^2)/f + (a^3*(c + d*x)^3)/(3*d) - (a^2*b*(c + d*x)^3)/d + (a*b^2*(c + d*x)^3)/d - (b^3*(c + d*x)^3)/(3*d) + (6*a*b^2*d*(c + d*x)*Log[1 + E^(2*(e + f*x))])/f^2 + (3*a^2*b*(c + d*x)^2*Log[1 + E^(2*(e + f*x))])/f + (b^3*(c + d*x)^2*Log[1 + E^(2*(e + f*x))])/f + (b^3*d^2*Log[Cosh[e + f*x]])/f^3 + (3*a*b^2*d^2*PolyLog[2, -E^(2*(e + f*x))])/f^3 + (3*a^2*b*d*(c + d*x)*PolyLog[2, -E^(2*(e + f*x))])/f^2 + (b^3*d*(c + d*x)*PolyLog[2, -E^(2*(e + f*x))])/f^2 - (3*a^2*b*d^2*PolyLog[3, -E^(2*(e + f*x))])/(2*f^3) - (b^3*d^2*PolyLog[3, -E^(2*(e + f*x))])/(2*f^3) - (b^3*d*(c + d*x)*Tanh[e + f*x])/f^2 - (3*a*b^2*(c + d*x)^2*Tanh[e + f*x])/f - (b^3*(c + d*x)^2*Tanh[e + f*x]^2)/(2*f)

Rule 32

```
Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[(((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.)), x_Symbol] := Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] := Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c + d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))]], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^3(c + dx)^2 + 3a^2b(c + dx)^2 \tanh(e + fx) + 3ab^2(c + dx)^2 \tanh^2(e + fx) \\
&\quad + b^3(c + dx)^2 \tanh^3(e + fx)) dx \\
&= \frac{a^3(c + dx)^3}{3d} + (3a^2b) \int (c + dx)^2 \tanh(e + fx) dx \\
&\quad + (3ab^2) \int (c + dx)^2 \tanh^2(e + fx) dx + b^3 \int (c + dx)^2 \tanh^3(e + fx) dx \\
&= \frac{a^3(c + dx)^3}{3d} - \frac{a^2b(c + dx)^3}{d} - \frac{3ab^2(c + dx)^2 \tanh(e + fx)}{f} \\
&\quad - \frac{b^3(c + dx)^2 \tanh^2(e + fx)}{2f} + (6a^2b) \int \frac{e^{2(e+fx)}(c + dx)^2}{1 + e^{2(e+fx)}} dx \\
&\quad + (3ab^2) \int (c + dx)^2 dx + b^3 \int (c + dx)^2 \tanh(e + fx) dx \\
&\quad + \frac{(6ab^2d) \int (c + dx) \tanh(e + fx) dx}{f} + \frac{(b^3d) \int (c + dx) \tanh^2(e + fx) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} + \frac{3a^2b(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad - \frac{b^3d(c+dx) \tanh(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \tanh^2(e+fx)}{2f} + (2b^3) \int \frac{e^{2(e+fx)}(c+dx)^2}{1+e^{2(e+fx)}} dx \\
&\quad + \frac{(b^3d^2) \int \tanh(e+fx) dx}{f^2} - \frac{(6a^2bd) \int (c+dx) \log(1+e^{2(e+fx)}) dx}{f} \\
&\quad + \frac{(12ab^2d) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx}{f} + \frac{(b^3d) \int (c+dx) dx}{f} \\
&= \frac{b^3cdx}{f} + \frac{b^3d^2x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} \\
&\quad + \frac{ab^2(c+dx)^3}{d} - \frac{b^3(c+dx)^3}{3d} + \frac{6ab^2d(c+dx) \log(1+e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2b(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3d^2 \log(\cosh(e+fx))}{f^3} + \frac{3a^2bd(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad - \frac{b^3d(c+dx) \tanh(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \tanh^2(e+fx)}{2f} - \frac{(3a^2bd^2) \int \text{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6ab^2d^2) \int \log(1+e^{2(e+fx)}) dx}{f^2} - \frac{(2b^3d) \int (c+dx) \log(1+e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} + \frac{6ab^2d(c+dx) \log(1+e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{b^3 d^2 \log(\cosh(e+fx))}{f^3} \\
&\quad + \frac{3a^2bd(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} + \frac{b^3 d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad - \frac{b^3 d(c+dx) \tanh(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \tanh^2(e+fx)}{2f} - \frac{(3a^2bd^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&\quad - \frac{(3ab^2d^2) \operatorname{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{f^3} - \frac{(b^3d^2) \int \operatorname{PolyLog}(2, -e^{2(e+fx)}) dx}{f^2} \\
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} + \frac{6ab^2d(c+dx) \log(1+e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{b^3 d^2 \log(\cosh(e+fx))}{f^3} \\
&\quad + \frac{3ab^2d^2 \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3a^2bd(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad + \frac{b^3 d(c+dx) \operatorname{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{3a^2bd^2 \operatorname{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{b^3 d(c+dx) \tanh(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \tanh^2(e+fx)}{2f} - \frac{(b^3d^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(2, -x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2 b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} + \frac{6ab^2 d(c+dx) \log(1+e^{2(e+fx)})}{f^2} + \frac{3a^2 b(c+dx)^2 \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^2 \log(1+e^{2(e+fx)})}{f} + \frac{b^3 d^2 \log(\cosh(e+fx))}{f^3} \\
&\quad + \frac{3ab^2 d^2 \text{PolyLog}(2, -e^{2(e+fx)})}{f^3} + \frac{3a^2 b d(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} \\
&\quad + \frac{b^3 d(c+dx) \text{PolyLog}(2, -e^{2(e+fx)})}{f^2} - \frac{3a^2 b d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} \\
&\quad - \frac{b^3 d^2 \text{PolyLog}(3, -e^{2(e+fx)})}{2f^3} - \frac{b^3 d(c+dx) \tanh(e+fx)}{f^2} \\
&\quad - \frac{3ab^2(c+dx)^2 \tanh(e+fx)}{f} - \frac{b^3(c+dx)^2 \tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 1163 vs. $2(405) = 810$.

Time = 7.23 (sec) , antiderivative size = 1163, normalized size of antiderivative = 2.87

$$\begin{aligned}
&\int (c+dx)^2 (a+b \tanh(e+fx))^3 dx \\
&= \frac{b(-4e^{2e}fx(9abdf(2c+dx) + 3a^2f^2(3c^2 + 3cdx + d^2x^2) + b^2(3c^2f^2 + 3cdf^2x + d^2(3+f^2x^2))) + 6(1+e^{2e}))}{6} \\
&\quad + \frac{\text{sech}(e)\text{sech}^2(e+fx)(6b^3c^2f \cosh(e) + 12b^3cdfx \cosh(e) + 6a^3c^2f^2x \cosh(e) + 18ab^2c^2f^2x \cosh(e) + 6b^3d^2f^2x^2 \cosh(e) + 6a^3d^2f^2x^3 \cosh(e) + 6a^2b^2d^2f^2x^2 \cosh(e) + 6a^3c^2d^2f^2x^2 \cosh(e) + 6a^2b^2d^2f^2x^3 \cosh(e) + 3a^3c^2d^2f^2x^3 \cosh(e) + 2a^3d^2f^2x^3 \cosh(e) + 3a^2b^2d^2f^2x^3 \cosh(e) + 3a^3c^2d^2f^2x^3 \cosh(3e+2fx) + 9a^2b^2c^2f^2x^3 \cosh(e+2fx) + 3a^3c^2d^2f^2x^3 \cosh(e+2fx) + 9a^2b^2c^2d^2f^2x^2 \cosh(e+2fx) + a^3d^2f^2x^3 \cosh(e+2fx) + 3a^2b^2d^2f^2x^3 \cosh(e+2fx) + 3a^3c^2d^2f^2x^3 \cosh(3e+2fx) + 9a^2b^2c^2f^2x^3 \cosh(3e+2fx) + 3a^3c^2d^2f^2x^2 \cosh(3e+2fx))}{6}
\end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Tanh[e + f*x])^3,x]

[Out] (b*(-4*E^(2*e)*f*x*(9*a*b*d*f*(2*c + d*x) + 3*a^2*f^2*(3*c^2 + 3*c*d*x + d^2*x^2) + b^2*(3*c^2*f^2 + 3*c*d*f^2*x + d^2*(3 + f^2*x^2))) + 6*(1 + E^(2*e)))*(6*a*b*d*f*(c + d*x) + 3*a^2*f^2*(c + d*x)^2 + b^2*(c^2*f^2 + 2*c*d*f^2*x + d^2*(1 + f^2*x^2)))*Log[1 + E^(2*(e + f*x))] + 6*d*(1 + E^(2*e))*(3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*PolyLog[2, -E^(2*(e + f*x))] - 3*(3*a^2 + b^2)*d^2*(1 + E^(2*e))*PolyLog[3, -E^(2*(e + f*x))]/(6*(1 + E^(2*e))*f^3) + (Sech[e]*Sech[e + f*x]^2*(6*b^3*c^2*f*Cosh[e] + 12*b^3*c*d*f*x*Cosh[e] + 6*a^3*c^2*f^2*x*Cosh[e] + 18*a*b^2*c^2*f^2*x*Cosh[e] + 6*b^3*d^2*f*x^2*Cosh[e] + 6*a^3*c*d*f^2*x^2*Cosh[e] + 18*a*b^2*c*d*f^2*x^2*Cosh[e] + 2*a^3*d^2*f^2*x^3*Cosh[e] + 6*a*b^2*d^2*f^2*x^3*Cosh[e] + 3*a^3*c^2*f^2*x*Cosh[e + 2*f*x] + 9*a*b^2*c^2*f^2*x*Cosh[e + 2*f*x] + 3*a^3*c*d*f^2*x^2*Cosh[e + 2*f*x] + 9*a*b^2*c*d*f^2*x^2*Cosh[e + 2*f*x] + a^3*d^2*f^2*x^3*Cosh[e + 2*f*x] + 3*a^2*b^2*d^2*f^2*x^3*Cosh[e + 2*f*x] + 3*a^3*c^2*f^2*x^3*Cosh[3*e + 2*f*x] + 9*a*b^2*c^2*f^2*x^3*Cosh[3*e + 2*f*x] + 3*a^3*c*d*f^2*x^2*Cosh[3*e + 2*f*x]))

$$\begin{aligned}
& + 2*f*x] + 9*a*b^2*c*d*f^2*x^2*\text{Cosh}[3*e + 2*f*x] + a^3*d^2*f^2*x^3*\text{Cosh}[3*e \\
& + 2*f*x] + 3*a*b^2*d^2*f^2*x^3*\text{Cosh}[3*e + 2*f*x] + 6*b^3*c*d*\text{Sinh}[e] + 18* \\
& a*b^2*c^2*f*\text{Sinh}[e] + 6*b^3*d^2*x*\text{Sinh}[e] + 36*a*b^2*c*d*f*x*\text{Sinh}[e] + 18*a \\
& ^2*b*c^2*f^2*x*\text{Sinh}[e] + 6*b^3*c^2*f^2*x*\text{Sinh}[e] + 18*a*b^2*d^2*f*x^2*\text{Sinh}[\\
& e] + 18*a^2*b*c*d*f^2*x^2*\text{Sinh}[e] + 6*b^3*c*d*f^2*x^2*\text{Sinh}[e] + 6*a^2*b*d^2 \\
& *f^2*x^3*\text{Sinh}[e] + 2*b^3*d^2*f^2*x^3*\text{Sinh}[e] - 6*b^3*c*d*\text{Sinh}[e + 2*f*x] - \\
& 18*a*b^2*c^2*f*\text{Sinh}[e + 2*f*x] - 6*b^3*d^2*x*\text{Sinh}[e + 2*f*x] - 36*a*b^2*c*d \\
& *f*x*\text{Sinh}[e + 2*f*x] - 9*a^2*b*c^2*f^2*x*\text{Sinh}[e + 2*f*x] - 3*b^3*c^2*f^2*x* \\
& \text{Sinh}[e + 2*f*x] - 18*a*b^2*d^2*f*x^2*\text{Sinh}[e + 2*f*x] - 9*a^2*b*c*d*f^2*x^2* \\
& \text{Sinh}[e + 2*f*x] - 3*b^3*c*d*f^2*x^2*\text{Sinh}[e + 2*f*x] - 3*a^2*b*d^2*f^2*x^3*\text{S} \\
& \text{inh}[e + 2*f*x] - b^3*d^2*f^2*x^3*\text{Sinh}[e + 2*f*x] + 9*a^2*b*c^2*f^2*x*\text{Sinh}[3 \\
& *e + 2*f*x] + 3*b^3*c^2*f^2*x*\text{Sinh}[3*e + 2*f*x] + 9*a^2*b*c*d*f^2*x^2*\text{Sinh}[\\
& 3*e + 2*f*x] + 3*b^3*c*d*f^2*x^2*\text{Sinh}[3*e + 2*f*x] + 3*a^2*b*d^2*f^2*x^3*\text{Si} \\
& \text{nh}[3*e + 2*f*x] + b^3*d^2*f^2*x^3*\text{Sinh}[3*e + 2*f*x]))/(12*f^2)
\end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1065 vs. $2(393) = 786$.

Time = 0.43 (sec) , antiderivative size = 1066, normalized size of antiderivative = 2.63

method	result	size
risch	Expression too large to display	1066

[In] `int((d*x+c)^2*(a+b*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out] $4/3/f^3*b^3*d^2*e^3+1/f^3*b^3*d^2*\ln(1+\exp(2*f*x+2*e))-2/f^3*b^3*d^2*\ln(\exp(f*x+e))+1/f*b^3*c^2*\ln(1+\exp(2*f*x+2*e))-2/f*b^3*c^2*\ln(\exp(f*x+e))-4/f*b^3*c*d*e*x-12/f*b*d*c*a^2*e*x+12/f^2*b*e*d*c*a^2*\ln(\exp(f*x+e))+6/f*b*d*c*a^2*\ln(1+\exp(2*f*x+2*e))*x+4/f^3*b*a^2*d^2*e^3+2/f^2*b^3*d^2*e^2*x-6/f*b^2*a*d^2*x^2-2/f^2*b^3*c*d*e^2-6/f^3*b^2*a*d^2*e^2+1/f*b^3*d^2*\ln(1+\exp(2*f*x+2*e))*x^2+1/f^2*b^3*d^2*\text{polylog}(2,-\exp(2*f*x+2*e))*x+1/f^2*b^3*c*d*\text{polylog}(2,-\exp(2*f*x+2*e))-2/f^3*b^3*e^2*d^2*\ln(\exp(f*x+e))+3/f*b*a^2*c^2*\ln(1+\exp(2*f*x+2*e))-6/f*b*a^2*c^2*\ln(\exp(f*x+e))+6/f^2*b*a^2*d^2*e^2*x-6/f^2*b*d*c*a^2*e^2-12/f^2*b^2*a*d^2*e*x+2/f*b^3*c*d*\ln(1+\exp(2*f*x+2*e))*x+3/f*b*a^2*d^2*\ln(1+\exp(2*f*x+2*e))*x^2+3/f^2*b*a^2*d^2*\text{polylog}(2,-\exp(2*f*x+2*e))*x+12/f^3*b^2*e*a*d^2*\ln(\exp(f*x+e))+3/f^2*b*d*c*a^2*\text{polylog}(2,-\exp(2*f*x+2*e))+6/f^2*b^2*a*d^2*\ln(1+\exp(2*f*x+2*e))*x+6/f^2*b^2*a*c*d*\ln(1+\exp(2*f*x+2*e))-1/2/f^2*b^2*a*c*d*\ln(\exp(f*x+e))+4/f^2*b^3*e*c*d*\ln(\exp(f*x+e))-6/f^3*b*e^2*a^2*d^2*\ln(\exp(f*x+e))-c*d*x^2*b^3-1/3*d^2*x^3*b^3+x*b^3*c^2+1/3/d*b^3*c^3-3*d*a^2*b*c*x^2+3*d*a*b^2*c*x^2+3*a^2*b*c^2*x+3*a*b^2*c^2*x-d^2*a^2*b*x^3+d^2*a*b^2*x^3+d*a^3*c*x^2+a^3*c^2*x+1/d*a^2*b*c^3+1/d*a*b^2*c^3+3*a*b^2*d^2*\text{polylog}(2,-\exp(2*f*x+2*e))/f^3-3/2*a^2*b*d^2*\text{polylog}(3,-\exp(2*f*x+2*e))/f^3-1/2*b^3*d^2*\text{polylog}(3,-\exp(2*f*x+2*e))/f^3+1/3*d^2*a^3*x^3+1/3/d*a^3*c^3+2*b^2*(3*a*d^2*f*x^2*\exp(2*f*x+2*e)+b*d^2*f*x^2*\exp(2*f*x+2*e)+6*a*c*d*f*x*\exp(2*f*x+2*e)+2*b*c*d*f*x*\exp(2*f*x+2*e)+3*a*c^2*f*\exp(2*f*x+2*e)+3*a*d^2*f*$

$$x^2 + b*c^2*f*\exp(2*f*x+2*e) + b*d^2*x*\exp(2*f*x+2*e) + 6*a*c*d*f*x*\exp(2*f*x+2*e) * d*b*c + 3*a*c^2*f + b*d^2*x + b*c*d) / f^2 / (1 + \exp(2*f*x+2*e))^2$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.36 (sec) , antiderivative size = 7298, normalized size of antiderivative = 18.02

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \text{Too large to display}$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^3,x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx)^2 dx$$

[In] integrate((d*x+c)**2*(a+b*tanh(f*x+e))**3,x)

[Out] Integral((a + b*tanh(e + f*x))**3*(c + d*x)**2, x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 869 vs. 2(390) = 780.

Time = 0.38 (sec) , antiderivative size = 869, normalized size of antiderivative = 2.15

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \frac{1}{3} a^3 d^2 x^3 + a^3 c d x^2 + b^3 c^2 \left(x + \frac{e}{f} + \frac{\log(e^{(-2fx-2e)} + 1)}{f} + \frac{2e^{(-2fx-2e)}}{f(2e^{(-2fx-2e)} + e^{(-4fx-4e)} + 1)} \right) + a^3 c^2 x + \frac{3a^2 b c^2 \log(\cosh(fx + e))}{f} + \frac{18ab^2 c^2 f + 6b^3 c d + (3a^2 b d^2 f^2 + 3ab^2 d^2 f^2 + b^3 d^2 f^2)x^3 + 3(3a^2 b c d f^2 + b^3 c d f^2 + 3(c d f^2 + 2d^2 f)ab^2)x^2 - \frac{2(6ab^2 c d f + b^3 d^2)x}{f^2} + \frac{(3a^2 b d^2 + b^3 d^2)(2f^2 x^2 \log(e^{(2fx+2e)} + 1) + 2fx \operatorname{Li}_2(-e^{(2fx+2e)}) - \operatorname{Li}_3(-e^{(2fx+2e)}))}{2f^3} + \frac{(3a^2 b c d f + b^3 c d f + 3ab^2 d^2)(2fx \log(e^{(2fx+2e)} + 1) + \operatorname{Li}_2(-e^{(2fx+2e)}))}{f^3} + \frac{(6ab^2 c d f + b^3 d^2) \log(e^{(2fx+2e)} + 1)}{f^3} - \frac{2((3a^2 b d^2 + b^3 d^2)f^3 x^3 + 3(3a^2 b c d f + b^3 c d f + 3ab^2 d^2)f^2 x^2)}{3f^3}$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^3,x, algorithm="maxima")

[Out] 1/3*a^3*d^2*x^3 + a^3*c*d*x^2 + b^3*c^2*(x + e/f + log(e^(-2*f*x - 2*e) + 1)/f + 2*e^(-2*f*x - 2*e)/(f*(2*e^(-2*f*x - 2*e) + e^(-4*f*x - 4*e) + 1))) + a^3*c^2*x + 3*a^2*b*c^2*log(cosh(f*x + e))/f + 1/3*(18*a*b^2*c^2*f + 6*b^3*c*d + (3*a^2*b*d^2*f^2 + 3*a*b^2*d^2*f^2 + b^3*d^2*f^2)*x^3 + 3*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 + 3*(c*d*f^2 + 2*d^2*f)*a*b^2)*x^2 + 3*(2*b^3*d^2 + 3*(c^2*f^2 + 4*c*d*f)*a*b^2)*x + (9*a*b^2*c^2*f^2*x*e^(4*e) + (3*a^2*b*d^2*f^2*2*e^(4*e) + 3*a*b^2*d^2*f^2*e^(4*e) + b^3*d^2*f^2*e^(4*e))*x^3 + 3*(3*a^2*b*c*d*f^2*e^(4*e) + 3*a*b^2*c*d*f^2*e^(4*e) + b^3*c*d*f^2*e^(4*e))*x^2)*e^(4*f*x) + 2*(9*a*b^2*c^2*f*e^(2*e) + 3*b^3*c*d*e^(2*e) + (3*a^2*b*d^2*f^2*e^(2*e) + 3*a*b^2*d^2*f^2*e^(2*e) + b^3*d^2*f^2*e^(2*e))*x^3 + 3*(3*a^2*b*c*d*f^2*e^(2*e) + 3*(c*d*f^2*e^(2*e) + d^2*f*e^(2*e))*a*b^2 + (c*d*f^2*e^(2*e) + d^2*f*e^(2*e))*b^3)*x^2 + 3*(3*(c^2*f^2*e^(2*e) + 2*c*d*f*e^(2*e))*a*b^2 + (2*c*d*f*e^(2*e) + d^2*e^(2*e))*b^3)*x)*e^(2*f*x))/(f^2*e^(4*f*x + 4*e) + 2*f^2*e^(2*f*x + 2*e) + f^2) - 2*(6*a*b^2*c*d*f + b^3*d^2)*x/f^2 + 1/2*(3*a^2*b*d^2 + b^3*d^2)*(2*f^2*x^2*log(e^(2*f*x + 2*e) + 1) + 2*f*x*dilog(-e^(2*f*x + 2*e)) - polylog(3, -e^(2*f*x + 2*e)))/f^3 + (3*a^2*b*c*d*f + b^3*c*d*f + 3*a*b^2*d^2)*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))

))/f^3 + (6*a*b^2*c*d*f + b^3*d^2)*log(e^(2*f*x + 2*e) + 1)/f^3 - 2/3*((3*a^2*b*d^2 + b^3*d^2)*f^3*x^3 + 3*(3*a^2*b*c*d*f + b^3*c*d*f + 3*a*b^2*d^2)*f^2*x^2)/f^3

Giac [**F**]

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \int (dx + c)^2 (b \tanh(fx + e) + a)^3 dx$$

[In] integrate((d*x+c)^2*(a+b*tanh(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^2*(b*tanh(f*x + e) + a)^3, x)

Mupad [**F(-1)**]

Timed out.

$$\int (c + dx)^2 (a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx)^2 dx$$

[In] int((a + b*tanh(e + f*x))^3*(c + d*x)^2,x)

[Out] int((a + b*tanh(e + f*x))^3*(c + d*x)^2, x)

3.65 $\int (c + dx)(a + b \tanh(e + fx))^3 dx$

Optimal result	440
Rubi [A] (verified)	441
Mathematica [A] (verified)	444
Maple [A] (verified)	445
Fricas [C] (verification not implemented)	445
Sympy [F]	447
Maxima [A] (verification not implemented)	447
Giac [F]	448
Mupad [F(-1)]	448

Optimal result

Integrand size = 18, antiderivative size = 261

$$\begin{aligned}
 \int (c + dx)(a + b \tanh(e + fx))^3 dx = & 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c + dx)^2}{2d} - \frac{3a^2b(c + dx)^2}{2d} \\
 & - \frac{b^3(c + dx)^2}{2d} + \frac{3a^2b(c + dx) \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{b^3(c + dx) \log(1 + e^{2(e+fx)})}{f} \\
 & + \frac{3ab^2d \log(\cosh(e + fx))}{f^2} \\
 & + \frac{3a^2bd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
 & + \frac{b^3d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{b^3d \tanh(e + fx)}{2f^2} \\
 & - \frac{3ab^2(c + dx) \tanh(e + fx)}{f} \\
 & - \frac{b^3(c + dx) \tanh^2(e + fx)}{2f}
 \end{aligned}$$

```

[Out] 3*a*b^2*c*x+1/2*b^3*d*x/f+3/2*a*b^2*d*x^2+1/2*a^3*(d*x+c)^2/d-3/2*a^2*b*(d*
x+c)^2/d-1/2*b^3*(d*x+c)^2/d+3*a^2*b*(d*x+c)*ln(1+exp(2*f*x+2*e))/f+b^3*(d*
x+c)*ln(1+exp(2*f*x+2*e))/f+3*a*b^2*d*ln(cosh(f*x+e))/f^2+3/2*a^2*b*d*polyl
og(2,-exp(2*f*x+2*e))/f^2+1/2*b^3*d*polylog(2,-exp(2*f*x+2*e))/f^2-1/2*b^3*
d*tanh(f*x+e)/f^2-3*a*b^2*(d*x+c)*tanh(f*x+e)/f-1/2*b^3*(d*x+c)*tanh(f*x+e)
^2/f

```


Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3803, 3799, 2221, 2317, 2438, 3801, 3556, 3554, 8}

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx = \frac{a^3(c + dx)^2}{2d} + \frac{3a^2b(c + dx) \log(e^{2(e+fx)} + 1)}{f} - \frac{3a^2b(c + dx)^2}{2d} + \frac{3a^2bd \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{3ab^2(c + dx) \tanh(e + fx)}{f} + 3ab^2cx + \frac{3ab^2d \log(\cosh(e + fx))}{f^2} + \frac{3}{2}ab^2dx^2 + \frac{b^3(c + dx) \log(e^{2(e+fx)} + 1)}{f} - \frac{b^3(c + dx) \tanh^2(e + fx)}{2f} - \frac{b^3(c + dx)^2}{2d} + \frac{b^3d \operatorname{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{b^3d \tanh(e + fx)}{2f^2} + \frac{b^3dx}{2f}$$

[In] Int[(c + d*x)*(a + b*Tanh[e + f*x])^3,x]

[Out] 3*a*b^2*c*x + (b^3*d*x)/(2*f) + (3*a*b^2*d*x^2)/2 + (a^3*(c + d*x)^2)/(2*d) - (3*a^2*b*(c + d*x)^2)/(2*d) - (b^3*(c + d*x)^2)/(2*d) + (3*a^2*b*(c + d*x)*Log[1 + E^(2*(e + f*x))])/f + (b^3*(c + d*x)*Log[1 + E^(2*(e + f*x))])/f + (3*a*b^2*d*Log[Cosh[e + f*x]])/f^2 + (3*a^2*b*d*PolyLog[2, -E^(2*(e + f*x))])/ (2*f^2) + (b^3*d*PolyLog[2, -E^(2*(e + f*x))])/ (2*f^2) - (b^3*d*Tanh[e + f*x])/ (2*f^2) - (3*a*b^2*(c + d*x)*Tanh[e + f*x])/f - (b^3*(c + d*x)*Tanh[e + f*x]^2)/(2*f)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d
*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x],
x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3799

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + (Complex[0, fz_])*(f_.)*(x_)], x
_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[(c
+ d*x)^m*(E^(2*((-I)*e + f*fz*x)))/(1 + E^(2*((-I)*e + f*fz*x)))], x], x]
/; FreeQ[{c, d, e, f, fz}, x] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^3(c+dx) + 3a^2b(c+dx) \tanh(e+fx) + 3ab^2(c+dx) \tanh^2(e+fx) \\
&\quad + b^3(c+dx) \tanh^3(e+fx)) dx \\
&= \frac{a^3(c+dx)^2}{2d} + (3a^2b) \int (c+dx) \tanh(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx) \tanh^2(e+fx) dx + b^3 \int (c+dx) \tanh^3(e+fx) dx \\
&= \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{3ab^2(c+dx) \tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx) \tanh^2(e+fx)}{2f} + (6a^2b) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx \\
&\quad + (3ab^2) \int (c+dx) dx + b^3 \int (c+dx) \tanh(e+fx) dx \\
&\quad + \frac{(3ab^2d) \int \tanh(e+fx) dx}{f} + \frac{(b^3d) \int \tanh^2(e+fx) dx}{2f} \\
&= 3ab^2cx + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{b^3(c+dx)^2}{2d} \\
&\quad + \frac{3a^2b(c+dx) \log(1+e^{2(e+fx)})}{f} + \frac{3ab^2d \log(\cosh(e+fx))}{f^2} \\
&\quad - \frac{b^3d \tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \tanh(e+fx)}{f} - \frac{b^3(c+dx) \tanh^2(e+fx)}{2f} \\
&\quad + (2b^3) \int \frac{e^{2(e+fx)}(c+dx)}{1+e^{2(e+fx)}} dx - \frac{(3a^2bd) \int \log(1+e^{2(e+fx)}) dx}{f} + \frac{(b^3d) \int 1 dx}{2f} \\
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} \\
&\quad - \frac{b^3(c+dx)^2}{2d} + \frac{3a^2b(c+dx) \log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx) \log(1+e^{2(e+fx)})}{f} + \frac{3ab^2d \log(\cosh(e+fx))}{f^2} \\
&\quad - \frac{b^3d \tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \tanh(e+fx)}{f} - \frac{b^3(c+dx) \tanh^2(e+fx)}{2f} \\
&\quad - \frac{(3a^2bd) \text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} - \frac{(b^3d) \int \log(1+e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{b^3(c+dx)^2}{2d} \\
&\quad + \frac{3a^2b(c+dx)\log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3ab^2d\log(\cosh(e+fx))}{f^2} + \frac{3a^2bd\text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{b^3d\tanh(e+fx)}{2f^2} - \frac{3ab^2(c+dx)\tanh(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)\tanh^2(e+fx)}{2f} - \frac{(b^3d)\text{Subst}\left(\int \frac{\log(1+x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} \\
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{b^3(c+dx)^2}{2d} \\
&\quad + \frac{3a^2b(c+dx)\log(1+e^{2(e+fx)})}{f} + \frac{b^3(c+dx)\log(1+e^{2(e+fx)})}{f} \\
&\quad + \frac{3ab^2d\log(\cosh(e+fx))}{f^2} + \frac{3a^2bd\text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} \\
&\quad + \frac{b^3d\text{PolyLog}(2, -e^{2(e+fx)})}{2f^2} - \frac{b^3d\tanh(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)\tanh(e+fx)}{f} - \frac{b^3(c+dx)\tanh^2(e+fx)}{2f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.47 (sec) , antiderivative size = 303, normalized size of antiderivative = 1.16

$$\int (c+dx)(a+b\tanh(e+fx))^3 dx$$

$$= \frac{\cosh(e+fx) \left(b^3f(c+dx) - a(a^2+3b^2)(e+fx)(-2cf+d(e-fx)) \cosh^2(e+fx) + b \cosh^2(e+fx) \right) \left(3a^2f^2(c+dx)^2 - 6ab^2d(e+fx) + 4(3a^2+b^2)(d^2e-cf)(e+fx) + 2(3a^2+b^2)d(e+fx)\text{Log}[1+E^{-2(e+fx)}] + 6ab^2d\text{Log}[1+E^{2(e+fx)}] - 2(3a^2+b^2)(d^2e-cf)\text{Log}[1+E^{2(e+fx)}] - (3a^2+b^2)d\text{PolyLog}[2, -E^{-2(e+fx)}] - (b^2d+6af^2(c+dx))\text{Sinh}[2(e+fx)]/2 \right) (a+b\tanh(e+fx))^3}{2f^2(a\cosh(e+fx)+b\sinh(e+fx))^3}$$

[In] Integrate[(c + d*x)*(a + b*Tanh[e + f*x])^3, x]

[Out] (Cosh[e + f*x]*(b^3*f*(c + d*x) - a*(a^2 + 3*b^2)*(e + f*x)*(-2*c*f + d*(e - f*x)))*Cosh[e + f*x]^2 + b*Cosh[e + f*x]^2*((3*a^2*f^2*(c + d*x)^2)/d + (b^2*f^2*(c + d*x)^2)/d - 6*a*b*d*(e + f*x) + 4*(3*a^2 + b^2)*(d*e - c*f)*(e + f*x) + 2*(3*a^2 + b^2)*d*(e + f*x)*Log[1 + E^(-2*(e + f*x))] + 6*a*b*d*Log[1 + E^(2*(e + f*x))] - 2*(3*a^2 + b^2)*(d*e - c*f)*Log[1 + E^(2*(e + f*x))] - (3*a^2 + b^2)*d*PolyLog[2, -E^(-2*(e + f*x))] - (b^2*(b*d + 6*a*f*(c + d*x))*Sinh[2*(e + f*x)]/2)*(a + b*Tanh[e + f*x])^3)/(2*f^2*(a*Cosh[e + f*x] + b*Sinh[e + f*x])^3)

Maple [A] (verified)

Time = 0.35 (sec) , antiderivative size = 459, normalized size of antiderivative = 1.76

method	result
risch	$-\frac{3a^2bdx^2}{2} + 3a^2bcx - \frac{6ba^2dex}{f} + \frac{6bea^2d\ln(e^{fx+e})}{f^2} - \frac{b^3de^2}{f^2} + \frac{b^3c\ln(1+e^{2fx+2e})}{f} - \frac{2b^3c\ln(e^{fx+e})}{f} + \frac{b^2(6adfx e^{2f}}$

[In] int((d*x+c)*(a+b*tanh(f*x+e))^3,x,method=_RETURNVERBOSE)

[Out]
$$-3/2*a^2*b*d*x^2+3*a^2*b*c*x-6/f*b*a^2*d*e*x+6/f^2*b*e*a^2*d*\ln(\exp(f*x+e))-1/f^2*b^3*d*e^2+1/f*b^3*c*\ln(1+\exp(2*f*x+2*e))-2/f*b^3*c*\ln(\exp(f*x+e))+b^2*(6*a*d*f*x*\exp(2*f*x+2*e)+2*b*d*f*x*\exp(2*f*x+2*e)+6*a*c*f*\exp(2*f*x+2*e)+2*b*c*f*\exp(2*f*x+2*e)+6*a*d*f*x+\exp(2*f*x+2*e)*d*b+6*a*c*f+b*d)/f^2/(1+\exp(2*f*x+2*e))^2+3/2*a^2*b*d*polylog(2,-\exp(2*f*x+2*e))/f^2+1/2*b^3*d*polylog(2,-\exp(2*f*x+2*e))/f^2+3*a*b^2*c*x+3/2*a*b^2*d*x^2-2/f*b^3*d*e*x-3/f^2*b*a^2*d*e^2+1/f*b^3*d*\ln(1+\exp(2*f*x+2*e))*x+3/f*b*a^2*c*\ln(1+\exp(2*f*x+2*e))-6/f*b*a^2*c*\ln(\exp(f*x+e))+2/f^2*b^3*e*d*\ln(\exp(f*x+e))+3/f^2*b^2*d*a*\ln(1+\exp(2*f*x+2*e))-6/f^2*b^2*d*a*\ln(\exp(f*x+e))+1/2*a^3*d*x^2-1/2*b^3*d*x^2+a^3*c*x+b^3*c*x+3/f*b*a^2*d*\ln(1+\exp(2*f*x+2*e))*x$$

Fricas [C] (verification not implemented)

Result contains complex when optimal does not.

Time = 0.31 (sec) , antiderivative size = 3262, normalized size of antiderivative = 12.50

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx = \text{Too large to display}$$

[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^3,x, algorithm="fricas")

[Out]
$$1/2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2*x + ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2)*x)*\cosh(f*x + e)^4 + 4*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2)*x)*\cosh(f*x + e)*\sinh(f*x + e)^3 + ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2)*x)*\sinh(f*x + e)^4 + 2*b^3*d + 2*(3*a^2*b + b^3)*d*e^2 + 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + 2*(3*a^2*b + b^3)*d*e^2 - 2*(2*(3*a^2*b + b^3)*c*e - (3*a*b^2 + b^3)*c)*f + 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2 - (3*a*b^2 - b^3)*d*f)*x)*\cosh(f*x + e)^2 + 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + 2*(3*a^2*b + b^3)*d*e^2 + 3*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b$$

$$\begin{aligned}
& b + b^3) * d * e^2 - 4 * (3 * a^2 * b + b^3) * c * e * f - 2 * (6 * a * b^2 * d * f - (a^3 - 3 * a^2 * b \\
& + 3 * a * b^2 - b^3) * c * f^2) * x) * \cosh(f * x + e)^2 - 2 * (2 * (3 * a^2 * b + b^3) * c * e - (3 * \\
& a * b^2 + b^3) * c) * f + 2 * ((a^3 - 3 * a^2 * b + 3 * a * b^2 - b^3) * c * f^2 - (3 * a * b^2 - b \\
& ^3) * d * f) * x) * \sinh(f * x + e)^2 + 4 * (3 * a * b^2 * c - (3 * a^2 * b + b^3) * c * e) * f + 2 * ((3 \\
& * a^2 * b + b^3) * d * \cosh(f * x + e)^4 + 4 * (3 * a^2 * b + b^3) * d * \cosh(f * x + e) * \sinh(f * \\
& x + e)^3 + (3 * a^2 * b + b^3) * d * \sinh(f * x + e)^4 + 2 * (3 * a^2 * b + b^3) * d * \cosh(f * x \\
& + e)^2 + 2 * (3 * (3 * a^2 * b + b^3) * d * \cosh(f * x + e)^2 + (3 * a^2 * b + b^3) * d) * \sinh \\
& (f * x + e)^2 + (3 * a^2 * b + b^3) * d + 4 * ((3 * a^2 * b + b^3) * d * \cosh(f * x + e)^3 + (3 * \\
& a^2 * b + b^3) * d * \cosh(f * x + e)) * \sinh(f * x + e)) * \operatorname{dilog}(I * \cosh(f * x + e) + I * \sinh \\
& (f * x + e)) + 2 * ((3 * a^2 * b + b^3) * d * \cosh(f * x + e)^4 + 4 * (3 * a^2 * b + b^3) * d * \cos \\
& h(f * x + e) * \sinh(f * x + e)^3 + (3 * a^2 * b + b^3) * d * \sinh(f * x + e)^4 + 2 * (3 * a^2 * b \\
& + b^3) * d * \cosh(f * x + e)^2 + 2 * (3 * (3 * a^2 * b + b^3) * d * \cosh(f * x + e)^2 + (3 * a^2 \\
& * b + b^3) * d) * \sinh(f * x + e)^2 + (3 * a^2 * b + b^3) * d + 4 * ((3 * a^2 * b + b^3) * d * \cos \\
& h(f * x + e)^3 + (3 * a^2 * b + b^3) * d * \cosh(f * x + e)) * \sinh(f * x + e)) * \operatorname{dilog}(-I * \cos \\
& h(f * x + e) - I * \sinh(f * x + e)) + 2 * ((3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^ \\
& ^2 * b + b^3) * c * f) * \cosh(f * x + e)^4 + 4 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a \\
& ^2 * b + b^3) * c * f) * \cosh(f * x + e) * \sinh(f * x + e)^3 + (3 * a * b^2 * d - (3 * a^2 * b + b^ \\
& ^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \sinh(f * x + e)^4 + 3 * a * b^2 * d - (3 * a^2 * b + b^3) \\
& * d * e + (3 * a^2 * b + b^3) * c * f + 2 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b \\
& + b^3) * c * f) * \cosh(f * x + e)^2 + 2 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b \\
& + b^3) * c * f + 3 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cos \\
& h(f * x + e)^2) * \sinh(f * x + e)^2 + 4 * ((3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^ \\
& ^2 * b + b^3) * c * f) * \cosh(f * x + e)^3 + (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^ \\
& ^2 * b + b^3) * c * f) * \cosh(f * x + e)) * \sinh(f * x + e)) * \log(\cosh(f * x + e) + \sinh(f * x + \\
& e) + I) + 2 * ((3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh \\
& (f * x + e)^4 + 4 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh \\
& (f * x + e) * \sinh(f * x + e)^3 + (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b \\
& ^3) * c * f) * \sinh(f * x + e)^4 + 3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3 \\
&) * c * f + 2 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh(f * x \\
& + e)^2 + 2 * (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f + 3 * (3 * a * \\
& b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh(f * x + e)^2) * \sinh(f * \\
& x + e)^2 + 4 * ((3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh \\
& (f * x + e)^3 + (3 * a * b^2 * d - (3 * a^2 * b + b^3) * d * e + (3 * a^2 * b + b^3) * c * f) * \cosh(f \\
& * x + e)) * \sinh(f * x + e)) * \log(\cosh(f * x + e) + \sinh(f * x + e) - I) + 2 * (((3 * a^2 \\
& * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh(f * x + e)^4 + 4 * ((3 * a^2 * b + b^3) \\
& * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh(f * x + e) * \sinh(f * x + e)^3 + ((3 * a^2 * b + b \\
& ^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \sinh(f * x + e)^4 + (3 * a^2 * b + b^3) * d * f * x + \\
& (3 * a^2 * b + b^3) * d * e + 2 * ((3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh \\
& (f * x + e)^2 + 2 * ((3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e + 3 * ((3 * a^2 * b + \\
& b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh(f * x + e)^2) * \sinh(f * x + e)^2 + 4 * (((\\
& 3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh(f * x + e)^3 + ((3 * a^2 * b + b \\
& ^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \cosh(f * x + e)) * \sinh(f * x + e)) * \log(I * \cosh(f \\
& * x + e) + I * \sinh(f * x + e) + 1) + 2 * (((3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3 \\
&) * d * e) * \cosh(f * x + e)^4 + 4 * ((3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e) * \co \\
& sh(f * x + e) * \sinh(f * x + e)^3 + ((3 * a^2 * b + b^3) * d * f * x + (3 * a^2 * b + b^3) * d * e)
\end{aligned}$$

```

*sinh(f*x + e)^4 + (3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e + 2*((3*a^2*
b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e)^2 + 2*((3*a^2*b + b^3)*
d*f*x + (3*a^2*b + b^3)*d*e + 3*((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*
e)*cosh(f*x + e)^2)*sinh(f*x + e)^2 + 4*((3*a^2*b + b^3)*d*f*x + (3*a^2*b
+ b^3)*d*e)*cosh(f*x + e)^3 + ((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)
*cosh(f*x + e))*sinh(f*x + e))*log(-I*cosh(f*x + e) - I*sinh(f*x + e) + 1)
+ 4*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b
+ b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b +
3*a*b^2 - b^3)*c*f^2)*x)*cosh(f*x + e)^3 + ((a^3 - 3*a^2*b + 3*a*b^2 - b^3
)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + 2*(3*a^2*b + b^3)*d*e^2 - 2*(2*(3*a^2*
b + b^3)*c*e - (3*a*b^2 + b^3)*c)*f + 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*
f^2 - (3*a*b^2 - b^3)*d*f)*x)*cosh(f*x + e))*sinh(f*x + e))/(f^2*cosh(f*x +
e)^4 + 4*f^2*cosh(f*x + e)*sinh(f*x + e)^3 + f^2*sinh(f*x + e)^4 + 2*f^2*c
osh(f*x + e)^2 + 2*(3*f^2*cosh(f*x + e)^2 + f^2)*sinh(f*x + e)^2 + f^2 + 4*
(f^2*cosh(f*x + e)^3 + f^2*cosh(f*x + e))*sinh(f*x + e))

```

Sympy [F]

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx) dx$$

```
[In] integrate((d*x+c)*(a+b*tanh(f*x+e))**3,x)
```

```
[Out] Integral((a + b*tanh(e + f*x))**3*(c + d*x), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.36 (sec) , antiderivative size = 475, normalized size of antiderivative = 1.82

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx$$

$$= \frac{1}{2} a^3 dx^2 + b^3 c \left(x + \frac{e}{f} + \frac{\log(e^{(-2fx-2e)} + 1)}{f} + \frac{2e^{(-2fx-2e)}}{f(2e^{(-2fx-2e)} + e^{(-4fx-4e)} + 1)} \right) + a^3 cx$$

$$- \frac{6ab^2 dx}{f} + \frac{3a^2 bc \log(\cosh(fx + e))}{f} - (3a^2 bd + b^3 d)x^2 + \frac{3ab^2 d \log(e^{(2fx+2e)} + 1)}{f^2}$$

$$+ \frac{12ab^2 cf + 6(cf^2 + 2df)ab^2 x + 2b^3 d + (3a^2 bdf^2 + 3ab^2 df^2 + b^3 df^2)x^2 + (6ab^2 cf^2 xe^{(4e)} + (3a^2 bdf^2 e^{(2fx+2e)} + (3a^2 bd + b^3 d)(2fx \log(e^{(2fx+2e)} + 1) + \text{Li}_2(-e^{(2fx+2e)})))}{2f^2}$$

```
[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/2*a^3*d*x^2 + b^3*c*(x + e/f + log(e^(-2*f*x - 2*e) + 1)/f + 2*e^(-2*f*x - 2*e)/(f*(2*e^(-2*f*x - 2*e) + e^(-4*f*x - 4*e) + 1))) + a^3*c*x - 6*a*b^2*d*x/f + 3*a^2*b*c*log(cosh(f*x + e))/f - (3*a^2*b*d + b^3*d)*x^2 + 3*a*b^2*d*log(e^(2*f*x + 2*e) + 1)/f^2 + 1/2*(12*a*b^2*c*f + 6*(c*f^2 + 2*d*f)*a*b^2*x + 2*b^3*d + (3*a^2*b*d*f^2 + 3*a*b^2*d*f^2 + b^3*d*f^2)*x^2 + (6*a*b^2*c*f^2*x*e^(4*e) + (3*a^2*b*d*f^2*e^(4*e) + 3*a*b^2*d*f^2*e^(4*e) + b^3*d*f^2*e^(4*e))*x^2)*e^(4*f*x) + 2*(6*a*b^2*c*f*e^(2*e) + b^3*d*e^(2*e) + (3*a^2*b*d*f^2*e^(2*e) + 3*a*b^2*d*f^2*e^(2*e) + b^3*d*f^2*e^(2*e))*x^2 + 2*(b^3*d*f*e^(2*e) + 3*(c*f^2*e^(2*e) + d*f*e^(2*e))*a*b^2)*x)*e^(2*f*x))/(f^2*e^(4*f*x + 4*e) + 2*f^2*e^(2*f*x + 2*e) + f^2) + 1/2*(3*a^2*b*d + b^3*d)*(2*f*x*log(e^(2*f*x + 2*e) + 1) + dilog(-e^(2*f*x + 2*e)))/f^2
```

Giac [F]

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx = \int (dx + c)(b \tanh(fx + e) + a)^3 dx$$

```
[In] integrate((d*x+c)*(a+b*tanh(f*x+e))^3,x, algorithm="giac")
```

```
[Out] integrate((d*x + c)*(b*tanh(f*x + e) + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \tanh(e + fx))^3 dx = \int (a + b \tanh(e + fx))^3 (c + dx) dx$$

```
[In] int((a + b*tanh(e + f*x))^3*(c + d*x),x)
```

```
[Out] int((a + b*tanh(e + f*x))^3*(c + d*x), x)
```


3.66 $\int \frac{(a+b \tanh(e+fx))^3}{c+dx} dx$

Optimal result	449
Rubi [N/A]	449
Mathematica [N/A]	450
Maple [N/A] (verified)	450
Fricas [N/A]	450
Sympy [N/A]	451
Maxima [N/A]	451
Giac [N/A]	451
Mupad [N/A]	452

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \text{Int}\left(\frac{(a + b \tanh(e + fx))^3}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))^3/(d*x+c), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx$$

[In] Int[(a + b*Tanh[e + f*x])^3/(c + d*x), x]

[Out] Defer[Int] [(a + b*Tanh[e + f*x])^3/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 32.80 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])^3/(c + d*x),x]

[Out] Integrate[(a + b*Tanh[e + f*x])^3/(c + d*x), x]

Maple [N/A] (verified)

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tanh(fx + e))^3}{dx + c} dx$$

[In] int((a+b*tanh(f*x+e))^3/(d*x+c),x)

[Out] int((a+b*tanh(f*x+e))^3/(d*x+c),x)

Fricas [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^3}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c),x, algorithm="fricas")

[Out] integral((b^3*tanh(f*x + e)^3 + 3*a*b^2*tanh(f*x + e)^2 + 3*a^2*b*tanh(f*x + e) + a^3)/(d*x + c), x)

Sympy [N/A]

Not integrable

Time = 1.40 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx$$

[In] integrate((a+b*tanh(f*x+e))**3/(d*x+c),x)

[Out] Integral((a + b*tanh(e + f*x))**3/(c + d*x), x)

Maxima [N/A]

Not integrable

Time = 0.56 (sec) , antiderivative size = 455, normalized size of antiderivative = 22.75

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^3}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c),x, algorithm="maxima")

[Out] $a^3 \log(dx + c)/d + (3a^2b + 3ab^2 + b^3) \log(dx + c)/d + (6ab^2d^2fx + 6a^2b^2cf - b^3d + (6a^2b^2cfe^{2e} + (2cfe^{2e} - de^{2e}))b^3 + 2(3a^2b^2dfe^{2e} + b^3dfe^{2e})x) e^{2fx} / (d^2f^2x^2 + 2cd^2fx + c^2f^2 + (d^2f^2x^2e^{4e} + 2cd^2fx^2e^{4e} + c^2f^2e^{4e})e^{4fx} + 2(d^2f^2x^2e^{2e} + 2cd^2fx^2e^{2e} + c^2f^2e^{2e})e^{2fx}) - \text{integrate}(2(3a^2b^2c^2f^2 - 3a^2b^2cd^2fx + (c^2f^2 + d^2)b^3 + (3a^2bd^2f^2 + b^3d^2f^2)x^2 + (6a^2b^2cd^2fx^2 + 2b^3cd^2fx^2 - 3a^2b^2d^2f)x) / (d^3f^2x^3 + 3cd^2f^2x^2 + 3c^2d^2fx + c^3f^2 + (d^3f^2x^3e^{2e} + 3cd^2f^2x^2e^{2e} + 3c^2d^2fx^2e^{2e} + c^3f^2e^{2e})e^{2fx}), x)$

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^3}{c + dx} dx = \int \frac{(b \tanh(fx + e) + a)^3}{dx + c} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c),x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)^3/(d*x + c), x)

Mupad [N/A]

Not integrable

Time = 1.94 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + f x))^3}{c + d x} dx = \int \frac{(a + b \tanh(e + f x))^3}{c + d x} dx$$

```
[In] int((a + b*tanh(e + f*x))^3/(c + d*x),x)
```

```
[Out] int((a + b*tanh(e + f*x))^3/(c + d*x), x)
```

$$3.67 \quad \int \frac{(a+b \tanh(e+fx))^3}{(c+dx)^2} dx$$

Optimal result	453
Rubi [N/A]	453
Mathematica [N/A]	454
Maple [N/A] (verified)	454
Fricas [N/A]	454
Sympy [N/A]	455
Maxima [N/A]	455
Giac [N/A]	456
Mupad [N/A]	456

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \tanh(e + fx))^3}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*tanh(f*x+e))^3/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$$

[In] Int[(a + b*Tanh[e + f*x])^3/(c + d*x)^2,x]

[Out] Defer[Int] [(a + b*Tanh[e + f*x])^3/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 34.75 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$$

[In] Integrate[(a + b*Tanh[e + f*x])^3/(c + d*x)^2,x]

[Out] Integrate[(a + b*Tanh[e + f*x])^3/(c + d*x)^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.29 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \tanh(fx + e))^3}{(dx + c)^2} dx$$

[In] int((a+b*tanh(f*x+e))^3/(d*x+c)^2,x)

[Out] int((a+b*tanh(f*x+e))^3/(d*x+c)^2,x)

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c)^2,x, algorithm="fricas")

[Out] integral((b^3*tanh(f*x + e)^3 + 3*a*b^2*tanh(f*x + e)^2 + 3*a^2*b*tanh(f*x + e) + a^3)/(d^2*x^2 + 2*c*d*x + c^2), x)

Sympy [N/A]

Not integrable

Time = 2.42 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))**3/(d*x+c)**2,x)

[Out] Integral((a + b*tanh(e + f*x))**3/(c + d*x)**2, x)

Maxima [N/A]

Not integrable

Time = 0.67 (sec) , antiderivative size = 926, normalized size of antiderivative = 46.30

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c)^2,x, algorithm="maxima")

```
[Out] -a^3/(d^2*x + c*d) - (3*a^2*b*c^2*f^2 + 3*(c^2*f^2 - 2*c*d*f)*a*b^2 + (c^2*f^2 + 2*d^2)*b^3 + (3*a^2*b*d^2*f^2 + 3*a*b^2*d^2*f^2 + b^3*d^2*f^2)*x^2 + 2*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 + 3*(c*d*f^2 - d^2*f)*a*b^2)*x + (3*a^2*b*c^2*f^2*e^(4*e) + 3*a*b^2*c^2*f^2*e^(4*e) + b^3*c^2*f^2*e^(4*e) + (3*a^2*b*d^2*f^2*e^(4*e) + 3*a*b^2*d^2*f^2*e^(4*e) + b^3*d^2*f^2*e^(4*e))*x^2 + 2*(3*a^2*b*c*d*f^2*e^(4*e) + 3*a*b^2*c*d*f^2*e^(4*e) + b^3*c*d*f^2*e^(4*e))*x)*e^(4*f*x) + 2*(3*a^2*b*c^2*f^2*e^(2*e) + 3*(c^2*f^2*e^(2*e) - c*d*f*e^(2*e)))*a*b^2 + (c^2*f^2*e^(2*e) - c*d*f*e^(2*e) + d^2*e^(2*e))*b^3 + (3*a^2*b*d^2*f^2*e^(2*e) + 3*a*b^2*d^2*f^2*e^(2*e) + b^3*d^2*f^2*e^(2*e))*x^2 + (6*a^2*b*c*d*f^2*e^(2*e) + 3*(2*c*d*f^2*e^(2*e) - d^2*f*e^(2*e))*a*b^2 + (2*c*d*f^2*e^(2*e) - d^2*f*e^(2*e))*b^3)*x)*e^(2*f*x))/(d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2 + (d^4*f^2*x^3*e^(4*e) + 3*c*d^3*f^2*x^2*e^(4*e) + 3*c^2*d^2*f^2*x*e^(4*e) + c^3*d*f^2*e^(4*e))*e^(4*f*x) + 2*(d^4*f^2*x^3*e^(2*e) + 3*c*d^3*f^2*x^2*e^(2*e) + 3*c^2*d^2*f^2*x*e^(2*e) + c^3*d*f^2*e^(2*e))*e^(2*f*x)) - integrate(2*(3*a^2*b*c^2*f^2 - 6*a*b^2*c*d*f + (c^2*f^2 + 3*d^2)*b^3 + (3*a^2*b*d^2*f^2 + b^3*d^2*f^2)*x^2 + 2*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 - 3*a*b^2*d^2*f)*x)/(d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2 + (d^4*f^2*x^4*e^(2*e) + 4*c*d^3*f^2*x^3*e^(2*e) + 6*c^2*d^2*f^2*x^2*e^(2*e) + 4*c^3*d*f^2*x*e^(2*e) + c^4*f^2*e^(2*e))*e^(2*f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.78 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \tanh(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] integrate((a+b*tanh(f*x+e))^3/(d*x+c)^2,x, algorithm="giac")

[Out] integrate((b*tanh(f*x + e) + a)^3/(d*x + c)^2, x)

Mupad [N/A]

Not integrable

Time = 2.01 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \tanh(e + fx))^3}{(c + dx)^2} dx$$

[In] int((a + b*tanh(e + f*x))^3/(c + d*x)^2,x)

[Out] int((a + b*tanh(e + f*x))^3/(c + d*x)^2, x)

3.68 $\int \frac{(c+dx)^3}{a+b \tanh(e+fx)} dx$

Optimal result	457
Rubi [A] (verified)	457
Mathematica [A] (verified)	460
Maple [B] (verified)	461
Fricas [B] (verification not implemented)	461
Sympy [F]	462
Maxima [B] (verification not implemented)	463
Giac [F]	464
Mupad [F(-1)]	464

Optimal result

Integrand size = 20, antiderivative size = 212

$$\int \frac{(c+dx)^3}{a+b \tanh(e+fx)} dx = \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2} + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3} + \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{4(a^2-b^2)f^4}$$

```
[Out] 1/4*(d*x+c)^4/(a+b)/d-b*(d*x+c)^3*ln(1+(a-b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f+3/2*b*d*(d*x+c)^2*polylog(2,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f^2+3/2*b*d^2*(d*x+c)*polylog(3,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f^3+3/4*b*d^3*polylog(4,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f^4
```

Rubi [A] (verified)

Time = 0.26 (sec) , antiderivative size = 212, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used

= {3813, 2221, 2611, 6744, 2320, 6724}

$$\int \frac{(c+dx)^3}{a+b \tanh(e+fx)} dx = \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^3(a^2-b^2)} + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^2(a^2-b^2)} - \frac{b(c+dx)^3 \log\left(\frac{(a-b)e^{-2(e+fx)}}{a+b} + 1\right)}{f(a^2-b^2)} + \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{4f^4(a^2-b^2)} + \frac{(c+dx)^4}{4d(a+b)}$$

[In] Int[(c + d*x)^3/(a + b*Tanh[e + f*x]),x]

[Out] (c + d*x)^4/(4*(a + b)*d) - (b*(c + d*x)^3*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))]/((a^2 - b^2)*f) + (3*b*d*(c + d*x)^2*PolyLog[2, -((a - b)/((a + b)*E^(2*(e + f*x)))))]/(2*(a^2 - b^2)*f^2) + (3*b*d^2*(c + d*x)*PolyLog[3, -((a - b)/((a + b)*E^(2*(e + f*x)))))]/(2*(a^2 - b^2)*f^3) + (3*b*d^3*PolyLog[4, -((a - b)/((a + b)*E^(2*(e + f*x)))))]/(4*(a^2 - b^2)*f^4)

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3813

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]), x_Sy
mbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int
[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*
I*(e + f*x), x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2,
0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.
)*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^
(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c + dx)^4}{4(a + b)d} + (2b) \int \frac{e^{-2(e+fx)}(c + dx)^3}{(a + b)^2 + (a^2 - b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c + dx)^4}{4(a + b)d} - \frac{b(c + dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{(3bd) \int (c + dx)^2 \log\left(1 + \frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2 - b^2)f} \\
&= \frac{(c + dx)^4}{4(a + b)d} - \frac{b(c + dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} \\
&\quad + \frac{3bd(c + dx)^2 \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2} \\
&\quad - \frac{(3bd^2) \int (c + dx) \text{PolyLog}\left(2, -\frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2 - b^2)f^2} \\
&= \frac{(c + dx)^4}{4(a + b)d} - \frac{b(c + dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{3bd(c + dx)^2 \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2} \\
&\quad + \frac{3bd^2(c + dx) \text{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^3} - \frac{(3bd^3) \int \text{PolyLog}\left(3, -\frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{2(a^2 - b^2)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3} \\
&\quad + \frac{(3bd^3) \text{Subst}\left(\int \frac{\text{PolyLog}\left(3, \frac{(-a+b)x}{a+b}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{4(a^2-b^2)f^4} \\
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3} + \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{4(a^2-b^2)f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 247, normalized size of antiderivative = 1.17

$$\begin{aligned}
&\int \frac{(c+dx)^3}{a+b \tanh(e+fx)} dx \\
&= \frac{1}{4} \left(-\frac{2b(c+dx)^4}{(a+b)d(b(-1+e^{2e})+a(1+e^{2e}))} - \frac{4b(c+dx)^3 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a-b)(a+b)f} \right. \\
&\quad \left. + \frac{3bd\left(2f^2(c+dx)^2 \text{PolyLog}\left(2, \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right) + d\left(2f(c+dx) \text{PolyLog}\left(3, \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right) + d \text{PolyLog}\right. \right.}{(a-b)(a+b)f^4} \right. \\
&\quad \left. \left. + \frac{x(4c^3 + 6c^2dx + 4cd^2x^2 + d^3x^3) \cosh(e)}{a \cosh(e) + b \sinh(e)}\right) \right)
\end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + b*Tanh[e + f*x]),x]

[Out] ((-2*b*(c + d*x)^4)/((a + b)*d*(b*(-1 + E^(2*e)) + a*(1 + E^(2*e)))) - (4*b*(c + d*x)^3*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x))]))/((a - b)*(a + b)*f) + (3*b*d*(2*f^2*(c + d*x)^2*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x))]) + d*(2*f*(c + d*x)*PolyLog[3, (-a + b)/((a + b)*E^(2*(e + f*x))]) + d*PolyLog[4, (-a + b)/((a + b)*E^(2*(e + f*x))]))/((a - b)*(a + b)*f^4) + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[e])/(a*Cosh[e] + b*Sinh[e])/4

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1155 vs. $2(209) = 418$.

Time = 0.32 (sec) , antiderivative size = 1156, normalized size of antiderivative = 5.45

method	result	size
risch	Expression too large to display	1156

[In] `int((d*x+c)^3/(a+b*tanh(f*x+e)),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4} \frac{1}{(a+b)} d^3 x^4 + \frac{1}{4} \frac{1}{(a+b)} \frac{d^3 c^4 + 3/4 f^4 b}{(a+b)} \frac{1}{(-a+b)} d^3 \text{polylog}(4, (a+b) \exp(2fx+2e) / (-a+b)) + \frac{1}{(a+b)} d^2 c^2 x^3 + \frac{3}{2} \frac{1}{(a+b)} d^2 c^2 x^2 + \frac{1}{(a+b)} c^3 x + \frac{1}{f^4 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) e^{-3} - \frac{3}{2} \frac{1}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 \text{polylog}(3, (a+b) \exp(2fx+2e) / (-a+b)) x - \frac{2}{f^4 b} \frac{1}{(a+b)} d^3 e^3 / (a-b) \ln(\exp(fx+e)) + \frac{1}{f^4 b} \frac{1}{(a+b)} d^3 e^3 / (a-b) \ln(\exp(2fx+2e) * a + b \exp(2fx+2e) + a - b) + \frac{3}{2} \frac{1}{f^2 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \text{polylog}(2, (a+b) \exp(2fx+2e) / (-a+b)) - \frac{3}{2} \frac{1}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \text{polylog}(3, (a+b) \exp(2fx+2e) / (-a+b)) + \frac{6}{f^2 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 e^{2x-6} / f^3 b \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 e^{x-3} / f^3 b \frac{1}{(a+b)} d^2 c^2 e^2 / (a-b) \ln(\exp(2fx+2e) * a + b \exp(2fx+2e) + a - b) + \frac{3}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) x^2 - \frac{6}{f^2 b} \frac{1}{(a+b)} d^2 c^2 e / (a-b) \ln(\exp(fx+e)) + \frac{3}{f^2 b} \frac{1}{(a+b)} d^2 c^2 e / (a-b) \ln(\exp(2fx+2e) * a + b \exp(2fx+2e) + a - b) + \frac{3}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) x + \frac{3}{f^2 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) e^{-3} / f^3 b \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) e^{2+3} / f^2 b \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 \text{polylog}(2, (a+b) \exp(2fx+2e) / (-a+b)) x + \frac{6}{f^3 b} \frac{1}{(a+b)} d^2 c^2 e^2 / (a-b) \ln(\exp(fx+e)) - \frac{2}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 e^3 x - \frac{3}{b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 x^2 - \frac{3}{f^2 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 e^2 - \frac{2}{b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 x^3 + \frac{4}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^2 c^2 e^3 + \frac{1}{f^3 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 \ln(1 - (a+b) \exp(2fx+2e) / (-a+b)) x^3 + \frac{3}{2} \frac{1}{f^2 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 \text{polylog}(2, (a+b) \exp(2fx+2e) / (-a+b)) x^2 - \frac{1}{2} \frac{1}{b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 x^4 - \frac{3}{2} \frac{1}{f^4 b} \frac{1}{(a+b)} \frac{1}{(-a+b)} d^3 e^4 + \frac{2}{f^3 b} \frac{1}{(a+b)} c^3 / (a-b) \ln(\exp(fx+e)) - \frac{1}{f^3 b} \frac{1}{(a+b)} c^3 / (a-b) \ln(\exp(2fx+2e) * a + b \exp(2fx+2e) + a - b)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 740 vs. $2(203) = 406$.

Time = 0.27 (sec) , antiderivative size = 740, normalized size of antiderivative = 3.49

$$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx$$

$$= \frac{(a + b)d^3 f^4 x^4 + 4(a + b)cd^2 f^4 x^3 + 6(a + b)c^2 d f^4 x^2 + 4(a + b)c^3 f^4 x - 24bd^3 \text{polylog}\left(4, \sqrt{-\frac{a+b}{a-b}}(\cosh(fx + e))\right)}{1}$$

[In] `integrate((d*x+c)^3/(a+b*tanh(f*x+e)),x, algorithm="fricas")`

```
[Out] 1/4*((a + b)*d^3*f^4*x^4 + 4*(a + b)*c*d^2*f^4*x^3 + 6*(a + b)*c^2*d*f^4*x^2 + 4*(a + b)*c^3*f^4*x - 24*b*d^3*polylog(4, sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 24*b*d^3*polylog(4, -sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) + 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt(-(a + b)/(a - b))) + 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) - 2*(a - b)*sqrt(-(a + b)/(a - b))) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) + 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) + 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, -sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))))/((a^2 - b^2)*f^4)
```

Sympy [F]

$$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx$$

```
[In] integrate((d*x+c)**3/(a+b*tanh(f*x+e)),x)
```

```
[Out] Integral((c + d*x)**3/(a + b*tanh(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(203) = 406.

Time = 0.32 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.49

$$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx$$

$$= \frac{3 \left(2fx \log \left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + \text{Li}_2 \left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} \right) \right) bc^2 d}{2(a^2 f^2 - b^2 f^2)}$$

$$- \frac{3 \left(2f^2 x^2 \log \left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + 2fx \text{Li}_2 \left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} \right) - \text{Li}_3 \left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} \right) \right) bcd^2}{2(a^2 f^3 - b^2 f^3)}$$

$$- \frac{\left(4f^3 x^3 \log \left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + 6f^2 x^2 \text{Li}_2 \left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} \right) - 6fx \text{Li}_3 \left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} \right) \right) bc^2 d}{3(a^2 f^4 - b^2 f^4)}$$

$$- c^3 \left(\frac{b \log(-(a-b)e^{(-2fx-2e)} - a-b)}{(a^2 - b^2)f} - \frac{fx + e}{(a+b)f} \right)$$

$$+ \frac{bd^3 f^4 x^4 + 4bcd^2 f^4 x^3 + 6bc^2 d f^4 x^2}{2(a^2 f^4 - b^2 f^4)} + \frac{d^3 x^4 + 4cd^2 x^3 + 6c^2 dx^2}{4(a+b)}$$

[In] integrate((d*x+c)^3/(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] -3/2*(2*f*x*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + dilog(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*c^2*d/(a^2*f^2 - b^2*f^2) - 3/2*(2*f^2*x^2*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 2*f*x*dilog(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - polylog(3, -(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*c*d^2/(a^2*f^3 - b^2*f^3) - 1/3*(4*f^3*x^3*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 6*f^2*x^2*dilog(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - 6*f*x*polylog(3, -(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) + 3*polylog(4, -(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*d^3/(a^2*f^4 - b^2*f^4) - c^3*(b*log(-(a - b)*e^(-2*f*x - 2*e) - a - b)/((a^2 - b^2)*f) - (f*x + e)/((a + b)*f)) + 1/2*(b*d^3*f^4*x^4 + 4*b*c*d^2*f^4*x^3 + 6*b*c^2*d*f^4*x^2)/(a^2*f^4 - b^2*f^4) + 1/4*(d^3*x^4 + 4*c*d^2*x^3 + 6*c^2*d*x^2)/(a + b)

Giac [F]

$$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx = \int \frac{(dx + c)^3}{b \tanh(fx + e) + a} dx$$

[In] integrate((d*x+c)^3/(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)^3/(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \tanh(e + fx)} dx$$

[In] int((c + d*x)^3/(a + b*tanh(e + f*x)),x)

[Out] int((c + d*x)^3/(a + b*tanh(e + f*x)), x)

3.69 $\int \frac{(c+dx)^2}{a+b \tanh(e+fx)} dx$

Optimal result	465
Rubi [A] (verified)	465
Mathematica [A] (verified)	468
Maple [B] (verified)	468
Fricas [B] (verification not implemented)	469
Sympy [F]	469
Maxima [B] (verification not implemented)	470
Giac [F]	470
Mupad [F(-1)]	471

Optimal result

Integrand size = 20, antiderivative size = 157

$$\int \frac{(c+dx)^2}{a+b \tanh(e+fx)} dx = \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3}$$

[Out] 1/3*(d*x+c)^3/(a+b)/d-b*(d*x+c)^2*ln(1+(a-b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f+b*d*(d*x+c)*polylog(2,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f^2+1/2*b*d^2*polylog(3,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)/f^3

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3813, 2221, 2611, 2320, 6724}

$$\int \frac{(c+dx)^2}{a+b \tanh(e+fx)} dx = \frac{bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f^2(a^2-b^2)} - \frac{b(c+dx)^2 \log\left(\frac{(a-b)e^{-2(e+fx)}}{a+b} + 1\right)}{f(a^2-b^2)} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^3(a^2-b^2)} + \frac{(c+dx)^3}{3d(a+b)}$$

[In] Int[(c + d*x)^2/(a + b*Tanh[e + f*x]),x]

[Out] (c + d*x)^3/(3*(a + b)*d) - (b*(c + d*x)^2*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))]/((a^2 - b^2)*f) + (b*d*(c + d*x)*PolyLog[2, -((a - b)/((a + b)*E^(2*(e + f*x))))]/((a^2 - b^2)*f^2) + (b*d^2*PolyLog[3, -((a - b)/((a + b)*E^(2*(e + f*x))))]/(2*(a^2 - b^2)*f^3)

Rule 2221

Int[(((F_)^((g_)*(e_) + (f_)*(x_)))^(n_))*((c_) + (d_)*(x_))^(m_)]/((a_) + (b_)*((F_)^((g_)*(e_) + (f_)*(x_)))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2320

Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*((a_) + (b_)*x))* (F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2611

Int[Log[1 + (e_)*((F_)^((c_)*((a_) + (b_)*(x_)))^(n_))]*((f_) + (g_)*(x_))^(m_), x_Symbol] := Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3813

Int[(((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]

Rule 6724

Int[PolyLog[n_, (c_)*((a_) + (b_)*(x_))^(p_)]/((d_) + (e_)*(x_)), x_Symbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^3}{3(a+b)d} + (2b) \int \frac{e^{-2(e+fx)}(c+dx)^2}{(a+b)^2 + (a^2-b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{(2bd) \int (c+dx) \log\left(1 + \frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2-b^2)f} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f^2} \\
&\quad - \frac{(bd^2) \int \text{PolyLog}\left(2, -\frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2-b^2)f^2} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f^2} \\
&\quad + \frac{(bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}\left(2, \frac{(-a+b)x}{a+b}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{2(a^2-b^2)f^3} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f^2} + \frac{bd^2 \text{PolyLog}\left(3, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.65 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.27

$$\int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx$$

$$= \frac{1}{6} \left(-\frac{4b(c + dx)^3}{(a + b)d(b(-1 + e^{2e}) + a(1 + e^{2e}))} - \frac{6b(c + dx)^2 \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a-b)(a+b)f} \right. \\ \left. + \frac{3bd\left(2f(c + dx) \operatorname{PolyLog}\left(2, \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right) + d \operatorname{PolyLog}\left(3, \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right)\right)}{(a-b)(a+b)f^3} \right. \\ \left. + \frac{2x(3c^2 + 3cdx + d^2x^2) \cosh(e)}{a \cosh(e) + b \sinh(e)} \right)$$

[In] Integrate[(c + d*x)^2/(a + b*Tanh[e + f*x]),x]

[Out] ((-4*b*(c + d*x)^3)/((a + b)*d*(b*(-1 + E^(2*e)) + a*(1 + E^(2*e)))) - (6*b*(c + d*x)^2*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x))]))/((a - b)*(a + b)*f) + (3*b*d*(2*f*(c + d*x)*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x))]] + d*PolyLog[3, (-a + b)/((a + b)*E^(2*(e + f*x))])))/((a - b)*(a + b)*f^3) + (2*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[e])/(a*Cosh[e] + b*Sinh[e])/6

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 732 vs. 2(157) = 314.

Time = 0.34 (sec) , antiderivative size = 733, normalized size of antiderivative = 4.67

method	result
risch	$\frac{d^2 x^3}{3a+3b} + \frac{dcx^2}{a+b} + \frac{c^2 x}{a+b} + \frac{c^3}{3(a+b)d} + \frac{2bd^2 e^2 \ln(e^{fx+e})}{f^3(a+b)(a-b)} - \frac{bd^2 e^2 \ln(e^{2fx+2e} a + b e^{2fx+2e} + a - b)}{f^3(a+b)(a-b)} + \frac{2bd^2 e^2 x}{f^2(a+b)(-a+b)} - \frac{bd^2 \ln(e^{fx+e})}{f^2(a+b)(-a+b)}$

[In] int((d*x+c)^2/(a+b*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] 1/3/(a+b)*d^2*x^3+1/(a+b)*d*c*x^2+1/(a+b)*c^2*x+1/3/(a+b)/d*c^3+2/f^3*b/(a+b)*d^2*e^2/(a-b)*ln(exp(f*x+e))-1/f^3*b/(a+b)*d^2*e^2/(a-b)*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)+2/f^2*b/(a+b)/(-a+b)*d^2*e^2*x-1/f^3*b/(a+b)/(-a+b)*d^2*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*e^2+1/f^2*b/(a+b)/(-a+b)*d^2*polylog(2,(a+b)*exp(2*f*x+2*e)/(-a+b))*x-2*b/(a+b)/(-a+b)*d*c*x^2-2/f^2*b/(a+b)/(-a+b)*d*c*e^2+2/f*b/(a+b)/(-a+b)*d*c*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*x+1/f^2*b/(a+b)/(-a+b)*d*c*polylog(2,(a+b)*exp(2*f*x+2*e)/(-a+b))-4/f*b/(a+b)/(-a+b)*d*c*e*x+2/f^2*b/(a+b)/(-a+b)*d*c*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*e-1/f*b/(a+b)*c^2/(a-b)*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)+2/f*b/(a+b)

$b) * c^2 / (a - b) * \ln(\exp(f * x + e)) - 2/3 * b / (a + b) / (-a + b) * d^2 * x^3 + 4/3 / f^3 * b / (a + b) / (-a + b) * d^2 * e^3 + 1/f * b / (a + b) / (-a + b) * d^2 * \ln(1 - (a + b) * \exp(2 * f * x + 2 * e) / (-a + b)) * x^2 - 1/2 / f^3 * b / (a + b) / (-a + b) * d^2 * \text{polylog}(3, (a + b) * \exp(2 * f * x + 2 * e) / (-a + b)) - 4/f^2 * b / (a + b) * d * c * e / (a - b) * \ln(\exp(f * x + e)) + 2/f^2 * b / (a + b) * d * c * e / (a - b) * \ln(\exp(2 * f * x + 2 * e) * a + b * \exp(2 * f * x + 2 * e) + a - b)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 500 vs. $2(152) = 304$.

Time = 0.27 (sec) , antiderivative size = 500, normalized size of antiderivative = 3.18

$$\int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx$$

$$(a + b)d^2 f^3 x^3 + 3(a + b)cdf^3 x^2 + 3(a + b)c^2 f^3 x + 6bd^2 \text{polylog}\left(3, \sqrt{-\frac{a+b}{a-b}}(\cosh(fx + e) + \sinh(fx + e))\right)$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] $1/3 * ((a + b) * d^2 * f^3 * x^3 + 3 * (a + b) * c * d * f^3 * x^2 + 3 * (a + b) * c^2 * f^3 * x + 6 * b * d^2 * \text{polylog}(3, \sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e))) + 6 * b * d^2 * \text{polylog}(3, -\sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e))) - 6 * (b * d^2 * f * x + b * c * d * f) * \text{dilog}(\sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e))) - 6 * (b * d^2 * f * x + b * c * d * f) * \text{dilog}(-\sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e))) - 3 * (b * d^2 * e^2 - 2 * b * c * d * e * f + b * c^2 * f^2) * \log(2 * (a + b) * \cosh(f * x + e) + 2 * (a + b) * \sinh(f * x + e) + 2 * (a - b) * \sqrt{-(a + b)/(a - b)}) - 3 * (b * d^2 * e^2 - 2 * b * c * d * e * f + b * c^2 * f^2) * \log(2 * (a + b) * \cosh(f * x + e) + 2 * (a + b) * \sinh(f * x + e) - 2 * (a - b) * \sqrt{-(a + b)/(a - b)}) - 3 * (b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x - b * d^2 * e^2 + 2 * b * c * d * e * f) * \log(\sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e)) + 1) - 3 * (b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x - b * d^2 * e^2 + 2 * b * c * d * e * f) * \log(-\sqrt{-(a + b)/(a - b)} * (\cosh(f * x + e) + \sinh(f * x + e)) + 1)) / ((a^2 - b^2) * f^3)$

Sympy [F]

$$\int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx$$

[In] integrate((d*x+c)**2/(a+b*tanh(f*x+e)),x)

[Out] Integral((c + d*x)**2/(a + b*tanh(e + f*x)), x)

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 337 vs. 2(152) = 304.

Time = 0.31 (sec) , antiderivative size = 337, normalized size of antiderivative = 2.15

$$\int \frac{(c+dx)^2}{a+b \tanh(e+fx)} dx$$

$$= -\frac{\left(2fx \log\left(\frac{(ae^{2e}+be^{2e})e^{2fx}}{a-b} + 1\right) + \text{Li}_2\left(-\frac{(ae^{2e}+be^{2e})e^{2fx}}{a-b}\right)\right) bcd}{a^2 f^2 - b^2 f^2}$$

$$-\frac{\left(2f^2 x^2 \log\left(\frac{(ae^{2e}+be^{2e})e^{2fx}}{a-b} + 1\right) + 2fx \text{Li}_2\left(-\frac{(ae^{2e}+be^{2e})e^{2fx}}{a-b}\right) - \text{Li}_3\left(-\frac{(ae^{2e}+be^{2e})e^{2fx}}{a-b}\right)\right) bd^2}{2(a^2 f^3 - b^2 f^3)}$$

$$-c^2 \left(\frac{b \log(-(a-b)e^{(-2fx-2e)} - a-b)}{(a^2 - b^2)f} - \frac{fx+e}{(a+b)f} \right)$$

$$+ \frac{2(bd^2 f^3 x^3 + 3bcd f^3 x^2)}{3(a^2 f^3 - b^2 f^3)} + \frac{d^2 x^3 + 3cdx^2}{3(a+b)}$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] $-(2*f*x*\log((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)/(a-b)} + 1) + \text{dilog}(-(a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)/(a-b)})) * b*c*d / (a^2*f^2 - b^2*f^2) - 1/2*(2*f^2*x^2*\log((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)/(a-b)} + 1) + 2*f*x*\text{dilog}(-(a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)/(a-b)} - \text{polylog}(3, -(a*e^{(2*e)} + b*e^{(2*e)}) * e^{(2*f*x)/(a-b)})) * b*d^2 / (a^2*f^3 - b^2*f^3) - c^2*(b*\log(-(a-b)*e^{(-2*f*x-2e)} - a-b) / ((a^2 - b^2)*f) - (f*x + e) / ((a+b)*f)) + 2/3*(b*d^2*f^3*x^3 + 3*b*c*d*f^3*x^2) / (a^2*f^3 - b^2*f^3) + 1/3*(d^2*x^3 + 3*c*d*x^2) / (a+b)$

Giac [F]

$$\int \frac{(c+dx)^2}{a+b \tanh(e+fx)} dx = \int \frac{(dx+c)^2}{b \tanh(fx+e) + a} dx$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)^2/(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \tanh(e + fx)} dx$$

```
[In] int((c + d*x)^2/(a + b*tanh(e + f*x)),x)
```

```
[Out] int((c + d*x)^2/(a + b*tanh(e + f*x)), x)
```

3.70 $\int \frac{c+dx}{a+b \tanh(e+fx)} dx$

Optimal result	472
Rubi [A] (verified)	472
Mathematica [A] (verified)	474
Maple [B] (verified)	474
Fricas [B] (verification not implemented)	475
Sympy [F]	475
Maxima [F]	475
Giac [F]	476
Mupad [F(-1)]	476

Optimal result

Integrand size = 18, antiderivative size = 108

$$\int \frac{c+dx}{a+b \tanh(e+fx)} dx = \frac{(c+dx)^2}{2(a+b)d} - \frac{b(c+dx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{bd \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2}$$

[Out] $1/2*(d*x+c)^2/(a+b)/d-b*(d*x+c)*\ln(1+(a-b)/(a+b)/\exp(2*f*x+2*e))/(a^2-b^2)/f+1/2*b*d*\operatorname{polylog}(2,(-a+b)/(a+b)/\exp(2*f*x+2*e))/(a^2-b^2)/f^2$

Rubi [A] (verified)

Time = 0.12 (sec) , antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {3813, 2221, 2317, 2438}

$$\int \frac{c+dx}{a+b \tanh(e+fx)} dx = -\frac{b(c+dx) \log\left(\frac{(a-b)e^{-2(e+fx)}}{a+b} + 1\right)}{f(a^2-b^2)} + \frac{bd \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^2(a^2-b^2)} + \frac{(c+dx)^2}{2d(a+b)}$$

[In] $\operatorname{Int}[(c+d*x)/(a+b*\operatorname{Tanh}[e+f*x]),x]$

[Out] $(c+d*x)^2/(2*(a+b)*d) - (b*(c+d*x)*\operatorname{Log}[1+(a-b)/((a+b)*E^{2*(e+f*x)})])/((a^2-b^2)*f) + (b*d*\operatorname{PolyLog}[2, -((a-b)/((a+b)*E^{2*(e+f*x)}))])/((2*(a^2-b^2)*f^2)$

Rule 2221

```
Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_))*((c_) + (d_)*(x_))^(m_)]/
((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)
))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol]
:=> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x))
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] :=> Simp[-PolyLog[2
, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3813

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Sy
mbol] :=> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int
[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*
I*(e + f*x), x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2,
0] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c + dx)^2}{2(a + b)d} + (2b) \int \frac{e^{-2(e+fx)}(c + dx)}{(a + b)^2 + (a^2 - b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c + dx)^2}{2(a + b)d} - \frac{b(c + dx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{(bd) \int \log\left(1 + \frac{(a^2-b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2 - b^2)f} \\
&= \frac{(c + dx)^2}{2(a + b)d} - \frac{b(c + dx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} \\
&\quad - \frac{(bd) \text{Subst}\left(\int \frac{\log\left(1 + \frac{(a^2-b^2)x}{(a+b)^2}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{2(a^2 - b^2)f^2} \\
&= \frac{(c + dx)^2}{2(a + b)d} - \frac{b(c + dx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{bd \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.69 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.41

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx = \frac{1}{2} \left(-\frac{2b(c + dx)^2}{(a + b)d(b(-1 + e^{2e}) + a(1 + e^{2e}))} - \frac{2b(c + dx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a-b)(a+b)f} + \frac{bd \operatorname{PolyLog}\left(2, \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right)}{(a-b)(a+b)f^2} + \frac{x(2c + dx) \cosh(e)}{a \cosh(e) + b \sinh(e)} \right)$$

[In] Integrate[(c + d*x)/(a + b*Tanh[e + f*x]),x]

[Out] ((-2*b*(c + d*x)^2)/((a + b)*d*(b*(-1 + E^(2*e)) + a*(1 + E^(2*e)))) - (2*b*(c + d*x)*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x))]))/((a - b)*(a + b)*f) + (b*d*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x))]))/((a - b)*(a + b)*f^2) + (x*(2*c + d*x)*Cosh[e])/(a*Cosh[e] + b*Sinh[e])/2

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(107) = 214.

Time = 0.23 (sec) , antiderivative size = 357, normalized size of antiderivative = 3.31

method	result
risch	$\frac{dx^2}{2a+2b} + \frac{cx}{a+b} + \frac{2bc \ln(e^{fx+e})}{f(a+b)(a-b)} - \frac{bc \ln(e^{2fx+2e} a+b e^{2fx+2e} a-b)}{f(a+b)(a-b)} - \frac{bdx^2}{(a+b)(-a+b)} + \frac{bd \ln\left(1 - \frac{(a+b)e^{2fx+2e}}{-a+b}\right)x}{f(a+b)(-a+b)} - \frac{2bd}{f(a+b)}$

[In] int((d*x+c)/(a+b*tanh(f*x+e)),x,method=_RETURNVERBOSE)

[Out] 1/2/(a+b)*d*x^2+1/(a+b)*c*x+2/f*b/(a+b)*c/(a-b)*ln(exp(f*x+e))-1/f*b/(a+b)*c/(a-b)*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-b/(a+b)/(-a+b)*d*x^2+1/f*b/(a+b)/(-a+b)*d*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*x-2/f*b/(a+b)/(-a+b)*d*e*x+1/f^2*b/(a+b)/(-a+b)*d*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*e-1/f^2*b/(a+b)/(-a+b)*d*e^2+1/2/f^2*b/(a+b)/(-a+b)*d*polylog(2,(a+b)*exp(2*f*x+2*e)/(-a+b))-2/f^2*b/(a+b)*d*e/(a-b)*ln(exp(f*x+e))+1/f^2*b/(a+b)*d*e/(a-b)*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 306 vs. 2(103) = 206.

Time = 0.27 (sec) , antiderivative size = 306, normalized size of antiderivative = 2.83

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx$$

$$(a + b)df^2x^2 + 2(a + b)cf^2x - 2bd\text{Li}_2\left(\sqrt{-\frac{a+b}{a-b}}(\cosh(fx + e) + \sinh(fx + e))\right) - 2bd\text{Li}_2\left(-\sqrt{-\frac{a+b}{a-b}}(\cosh(fx + e) + \sinh(fx + e))\right)$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] 1/2*((a + b)*d*f^2*x^2 + 2*(a + b)*c*f^2*x - 2*b*d*dilog(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 2*b*d*dilog(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) + 2*(b*d*e - b*c*f)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt(-(a + b)/(a - b))) + 2*(b*d*e - b*c*f)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) - 2*(a - b)*sqrt(-(a + b)/(a - b))) - 2*(b*d*f*x + b*d*e)*log(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 2*(b*d*f*x + b*d*e)*log(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1))/((a^2 - b^2)*f^2)

Sympy [F]

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx = \int \frac{c + dx}{a + b \tanh(e + fx)} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e)),x)

[Out] Integral((c + d*x)/(a + b*tanh(e + f*x)), x)

Maxima [F]

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx = \int \frac{dx + c}{b \tanh(fx + e) + a} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 1/2*(4*b*integrate(x/(a^2 - b^2 + (a^2*e^(2*e) + 2*a*b*e^(2*e) + b^2*e^(2*e))*e^(2*f*x)), x) + x^2/(a + b))*d - c*(b*log(-(a - b)*e^(-2*f*x - 2*e) - a - b)/((a^2 - b^2)*f) - (f*x + e)/((a + b)*f))

Giac [F]

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx = \int \frac{dx + c}{b \tanh(fx + e) + a} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate((d*x + c)/(b*tanh(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{a + b \tanh(e + fx)} dx = \int \frac{c + dx}{a + b \tanh(e + fx)} dx$$

[In] int((c + d*x)/(a + b*tanh(e + f*x)),x)

[Out] int((c + d*x)/(a + b*tanh(e + f*x)), x)

3.71 $\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx$

Optimal result	477
Rubi [N/A]	477
Mathematica [N/A]	478
Maple [N/A] (verified)	478
Fricas [N/A]	478
Sympy [N/A]	478
Maxima [N/A]	479
Giac [N/A]	479
Mupad [N/A]	479

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b \tanh(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*tanh(f*x+e)), x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx = \int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx$$

[In] Int[1/((c + d*x)*(a + b*Tanh[e + f*x])), x]

[Out] Defer[Int][1/((c + d*x)*(a + b*Tanh[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 4.90 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))} dx = \int \frac{1}{(c + dx)(a + b \tanh(e + fx))} dx$$

[In] Integrate[1/((c + d*x)*(a + b*Tanh[e + f*x])),x]

[Out] Integrate[1/((c + d*x)*(a + b*Tanh[e + f*x])), x]

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tanh(fx + e))} dx$$

[In] int(1/(d*x+c)/(a+b*tanh(f*x+e)),x)

[Out] int(1/(d*x+c)/(a+b*tanh(f*x+e)),x)

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.35

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))} dx = \int \frac{1}{(dx + c)(b \tanh(fx + e) + a)} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] integral(1/(a*d*x + a*c + (b*d*x + b*c)*tanh(f*x + e)), x)

Sympy [N/A]

Not integrable

Time = 1.03 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))} dx = \int \frac{1}{(a + b \tanh(e + fx))(c + dx)} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e)),x)

[Out] Integral(1/((a + b*tanh(e + f*x))*(c + d*x)), x)

Maxima [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 114, normalized size of antiderivative = 5.70

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx = \int \frac{1}{(dx+c)(b \tanh(fx+e)+a)} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 2*b*integrate(1/(a^2*c - b^2*c + (a^2*d - b^2*d)*x + (a^2*c*e^(2*e) + 2*a*b*c*e^(2*e) + b^2*c*e^(2*e) + (a^2*d*e^(2*e) + 2*a*b*d*e^(2*e) + b^2*d*e^(2*e))*x)*e^(2*f*x)), x) + log(d*x + c)/(a*d + b*d)

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx = \int \frac{1}{(dx+c)(b \tanh(fx+e)+a)} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate(1/((d*x + c)*(b*tanh(f*x + e) + a)), x)

Mupad [N/A]

Not integrable

Time = 1.82 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))} dx = \int \frac{1}{(a+b \tanh(e+fx))(c+dx)} dx$$

[In] int(1/((a + b*tanh(e + f*x))*(c + d*x)),x)

[Out] int(1/((a + b*tanh(e + f*x))*(c + d*x)), x)

$$3.72 \quad \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx$$

Optimal result	480
Rubi [N/A]	480
Mathematica [N/A]	481
Maple [N/A] (verified)	481
Fricas [N/A]	481
Sympy [N/A]	482
Maxima [N/A]	482
Giac [N/A]	482
Mupad [N/A]	483

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b \tanh(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx$$

[In] Int[1/((c + d*x)^2*(a + b*Tanh[e + f*x])),x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*Tanh[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 9.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))} dx = \int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))} dx$$

[In] Integrate[1/((c + d*x)^2*(a + b*Tanh[e + f*x])),x]

[Out] Integrate[1/((c + d*x)^2*(a + b*Tanh[e + f*x])), x]

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2(a + b \tanh(fx + e))} dx$$

[In] int(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x)

[Out] int(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \tanh(fx + e) + a)} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="fricas")

[Out] integral(1/(a*d^2*x^2 + 2*a*c*d*x + a*c^2 + (b*d^2*x^2 + 2*b*c*d*x + b*c^2)*tanh(f*x + e)), x)

Sympy [N/A]

Not integrable

Time = 1.71 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx = \int \frac{1}{(a+b \tanh(e+fx))(c+dx)^2} dx$$

[In] integrate(1/(d*x+c)**2/(a+b*tanh(f*x+e)),x)

[Out] Integral(1/((a + b*tanh(e + f*x))*(c + d*x)**2), x)

Maxima [N/A]

Not integrable

Time = 0.52 (sec) , antiderivative size = 199, normalized size of antiderivative = 9.95

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx = \int \frac{1}{(dx+c)^2(b \tanh(fx+e)+a)} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="maxima")

[Out] 2*b*integrate(1/(a^2*c^2 - b^2*c^2 + (a^2*d^2 - b^2*d^2)*x^2 + 2*(a^2*c*d - b^2*c*d)*x + (a^2*c^2*e^(2*e) + 2*a*b*c^2*e^(2*e) + b^2*c^2*e^(2*e) + (a^2*d^2*e^(2*e) + 2*a*b*d^2*e^(2*e) + b^2*d^2*e^(2*e))*x^2 + 2*(a^2*c*d*e^(2*e) + 2*a*b*c*d*e^(2*e) + b^2*c*d*e^(2*e))*x)*e^(2*f*x)), x) - 1/(a*c*d + b*c*d + (a*d^2 + b*d^2)*x)

Giac [N/A]

Not integrable

Time = 0.34 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))} dx = \int \frac{1}{(dx+c)^2(b \tanh(fx+e)+a)} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e)),x, algorithm="giac")

[Out] integrate(1/((d*x + c)^2*(b*tanh(f*x + e) + a)), x)

Mupad [N/A]

Not integrable

Time = 2.09 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))} dx = \int \frac{1}{(a + b \tanh(e + fx)) (c + dx)^2} dx$$

```
[In] int(1/((a + b*tanh(e + f*x))*(c + d*x)^2), x)
```

```
[Out] int(1/((a + b*tanh(e + f*x))*(c + d*x)^2), x)
```

3.73
$$\int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx$$

Optimal result	485
Rubi [A] (verified)	486
Mathematica [A] (verified)	495
Maple [B] (verified)	496
Fricas [B] (verification not implemented)	497
Sympy [F]	497
Maxima [A] (verification not implemented)	498
Giac [F]	498
Mupad [F(-1)]	499

Optimal result

Integrand size = 20, antiderivative size = 642

$$\begin{aligned}
 \int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx = & -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} \\
 & + \frac{(c+dx)^4}{4(a-b)^2 d} + \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
 & - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
 & + \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
 & + \frac{3b^2 d^2(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
 & - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
 & + \frac{3b^2 d(c+dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
 & - \frac{3b^2 d^3 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4} \\
 & + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
 & - \frac{3b^2 d^2(c+dx) \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
 & - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} \\
 & + \frac{3b^2 d^3 \operatorname{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4}
 \end{aligned}$$

```

[Out] -2*b^2*(d*x+c)^3/(a^2-b^2)^2/f+2*b^2*(d*x+c)^3/(a-b)/(a+b)^2/(a-b+(a+b)*exp
(2*f*x+2*e))/f+1/4*(d*x+c)^4/(a-b)^2/d+3*b^2*d*(d*x+c)^2*ln(1+(a+b)*exp(2*f
*x+2*e)/(a-b))/(a^2-b^2)^2/f^2-2*b*(d*x+c)^3*ln(1+(a+b)*exp(2*f*x+2*e)/(a-b
))/(a-b)^2/(a+b)/f+2*b^2*(d*x+c)^3*ln(1+(a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^
2)^2/f+3*b^2*d^2*(d*x+c)*polylog(2,-(a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2
/f^3-3*b*d*(d*x+c)^2*polylog(2,-(a+b)*exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f
^2+3*b^2*d*(d*x+c)^2*polylog(2,-(a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^2

```

$$\begin{aligned}
& -3/2*b^2*d^3*\text{polylog}(3,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^4+3*b*d^2 \\
& *(d*x+c)*\text{polylog}(3,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^3-3*b^2*d^2 \\
& *(d*x+c)*\text{polylog}(3,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^3-3/2*b*d^3*p \\
& \text{olylog}(4,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^4+3/2*b^2*d^3*\text{polylog} \\
& (4,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^4
\end{aligned}$$

Rubi [A] (verified)

Time = 1.48 (sec) , antiderivative size = 642, normalized size of antiderivative = 1.00, number of steps used = 28, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules

used = {3815, 2221, 2611, 6744, 2320, 6724, 2286, 2216, 2215, 2222}

$$\begin{aligned}
 \int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx = & \frac{3b^2 d^2 (c + dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\
 & - \frac{3b^2 d^2 (c + dx) \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\
 & + \frac{3b^2 d (c + dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2 - b^2)^2} \\
 & + \frac{3b^2 d (c + dx)^2 \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f^2 (a^2 - b^2)^2} \\
 & + \frac{2b^2 (c + dx)^3 \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f (a^2 - b^2)^2} \\
 & - \frac{2b^2 (c + dx)^3}{f (a^2 - b^2)^2} - \frac{3b^2 d^3 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a^2 - b^2)^2} \\
 & + \frac{3b^2 d^3 \operatorname{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a^2 - b^2)^2} \\
 & + \frac{2b^2 (c + dx)^3}{f (a - b)(a + b)^2 ((a + b)e^{2e+2fx} + a - b)} \\
 & + \frac{3bd^2 (c + dx) \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a - b)^2 (a + b)} \\
 & - \frac{3bd (c + dx)^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a - b)^2 (a + b)} \\
 & - \frac{2b (c + dx)^3 \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f (a - b)^2 (a + b)} \\
 & + \frac{(c + dx)^4}{4d(a - b)^2} - \frac{3bd^3 \operatorname{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a - b)^2 (a + b)}
 \end{aligned}$$

[In] Int[(c + d*x)^3/(a + b*Tanh[e + f*x])^2,x]

[Out] (-2*b^2*(c + d*x)^3)/((a^2 - b^2)^2*f) + (2*b^2*(c + d*x)^3)/((a - b)*(a + b)^2*(a - b + (a + b)*E^(2*e + 2*f*x))*f) + (c + d*x)^4/(4*(a - b)^2*d) + (3*b^2*d*(c + d*x)^2*Log[1 + ((a + b)*E^(2*e + 2*f*x))/(a - b)]/((a^2 - b^2)^2*f^2) - (2*b*(c + d*x)^3*Log[1 + ((a + b)*E^(2*e + 2*f*x))/(a - b)]/((a - b)^2*(a + b)*f) + (2*b^2*(c + d*x)^3*Log[1 + ((a + b)*E^(2*e + 2*f*x))/(a - b)]/((a^2 - b^2)^2*f) + (3*b^2*d^2*(c + d*x)*PolyLog[2, -(((a + b)*E^(2*e + 2*f*x))/(a - b))]/((a^2 - b^2)^2*f^3) - (3*b*d*(c + d*x)^2*PolyLog[2

$$\frac{-\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)}{(a-b)^2(a+b)f^2} + \frac{(3b^2d(c+dx)^2 \text{PolyLog}[2, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)]}{(a^2-b^2)^2f^2} - \frac{(3b^2d^3 \text{PolyLog}[3, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)]}{2(a^2-b^2)^2f^4} + \frac{(3bd^2(c+dx) \text{PolyLog}[3, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)])}{(a-b)^2(a+b)f^3} - \frac{(3b^2d^2(c+dx) \text{PolyLog}[3, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)])}{(a^2-b^2)^2f^3} - \frac{(3bd^3 \text{PolyLog}[4, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)])}{2(a-b)^2(a+b)f^4} + \frac{(3b^2d^3 \text{PolyLog}[4, -\left(\frac{(a+b)E^{(2e+2fx)}}{(a-b)}\right)])}{2(a^2-b^2)^2f^4}$$

Rule 2215

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Dist[b/a, Int[(c + d*x)^m*((F^(g*(e + f*x)))^n)/(a + b*(F^(g*(e + f*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2216

```
Int[((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.))^(p_)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Dist[1/a, Int[(c + d*x)^m*(a + b*(F^(g*(e + f*x))))^n]^(p + 1), x], x] - Dist[b/a, Int[(c + d*x)^m*(F^(g*(e + f*x)))^n*(a + b*(F^(g*(e + f*x))))^n]^(p, x), x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && ILtQ[p, 0] && IGtQ[m, 0]
```

Rule 2221

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2222

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(p_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((a + b*(F^(g*(e + f*x))))^n)^(p + 1)/(b*f*g*n*(p + 1)*Log[F]), x] - Dist[d*(m/(b*f*g*n*(p + 1)*Log[F])), Int[(c + d*x)^(m - 1)*(a + b*(F^(g*(e + f*x))))^n]^(p + 1), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, m, n, p}, x] && NeQ[p, -1]
```

Rule 2286

```
Int[((a_.) + (b_.)*(F_)^(u_))^(p_.)*((c_.) + (d_.)*(F_)^(v_))^(q_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := With[{w = ExpandIntegrand[(e + f*x)^m, (a + b*F^u)^p*(c + d*F^v)^q, x]}, Int[w, x] /; SumQ[w]] /; FreeQ[{F, a, b, c,
```


d, e, f, m}, x] && IntegersQ[p, q] && LinearQ[{u, v}, x] && RationalQ[Simplify[u/v]]

Rule 2320

```
Int[u_, x_Symbol] := With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x],
Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] := Simp[(- (f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3815

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.),
x_Symbol] := Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 +
b^2 + (a - I*b)^2*E^(2*I*(e + f*x))))^(-n), x], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] := Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_.)*((a_.) + (b_.)
*(x_))))^(p_.)], x_Symbol] := Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a
+ b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(
m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = \int \left(\frac{(c + dx)^3}{(a - b)^2} + \frac{4be^{2e+2fx}(c + dx)^3}{(a - b)^2 \left(-a \left(1 - \frac{b}{a} \right) - a \left(1 + \frac{b}{a} \right) e^{2e+2fx} \right)} + \frac{4b^2 e^{4e+4fx} (c + dx)^3}{(a - b)^2 \left(a \left(1 - \frac{b}{a} \right) + a \left(1 + \frac{b}{a} \right) e^{2e+2fx} \right)^2} \right) dx$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a-b)^2d} + \frac{(4b) \int \frac{e^{2e+2fx}(c+dx)^3}{-a\left(1-\frac{b}{a}\right)-a\left(1+\frac{b}{a}\right)e^{2e+2fx}} dx}{(a-b)^2} + \frac{(4b^2) \int \frac{e^{4e+4fx}(c+dx)^3}{\left(a\left(1-\frac{b}{a}\right)+a\left(1+\frac{b}{a}\right)e^{2e+2fx}\right)^2} dx}{(a-b)^2} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{(4b^2) \int \left(\frac{(c+dx)^3}{(a+b)^2} + \frac{(a-b)^2(c+dx)^3}{(a+b)^2(a-b+(a+b)e^{2e+2fx})^2} + \frac{2(-a+b)(c+dx)^3}{(a+b)^2(a-b+(a+b)e^{2e+2fx})} \right) dx}{(a-b)^2} \\
&\quad + \frac{(6bd) \int (c+dx)^2 \log\left(1 + \frac{\left(1+\frac{b}{a}\right)e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} + \frac{b^2(c+dx)^4}{(a^2-b^2)^2d} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^3}{(a-b+(a+b)e^{2e+2fx})^2} dx}{(a+b)^2} \\
&\quad - \frac{(8b^2) \int \frac{(c+dx)^3}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} + \frac{(6bd^2) \int (c+dx) \text{PolyLog}\left(2, -\frac{\left(1+\frac{b}{a}\right)e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f^2} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} - \frac{b^2(c+dx)^4}{(a^2-b^2)^2d} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} + \frac{(4b^2) \int \frac{(c+dx)^3}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} \\
&\quad + \frac{(8b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} - \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{(a-b+(a+b)e^{2e+2fx})^2} dx}{a^2-b^2} \\
&\quad - \frac{(3bd^3) \int \text{PolyLog}\left(3, -\frac{\left(1+\frac{b}{a}\right)e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{4b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} \\
&\quad - \frac{(3bd^3) \text{Subst}\left(\int \frac{\text{PolyLog}\left(3, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a-b)^2(a+b)f^4} \\
&\quad - \frac{(6b^2d) \int \frac{(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2f} - \frac{(12b^2d) \int (c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} \\
&\quad + \frac{(c+dx)^4}{4(a-b)^2d} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{6b^2d(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} \\
&\quad - \frac{(12b^2d^2) \int (c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2} \\
&\quad + \frac{(6b^2d) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)f} + \frac{(6b^2d) \int (c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&+ \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&+ \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{6b^2 d^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{(6b^2 d^3) \int \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3} \\
&- \frac{(6b^2 d^2) \int (c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2} \\
&+ \frac{(6b^2 d^2) \int (c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx}) f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&+ \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&+ \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} - \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} \\
&+ \frac{(3b^2 d^3) \text{Subst}\left(\int \frac{\text{PolyLog}\left(3, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^4} \\
&- \frac{(3b^2 d^3) \int \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3} \\
&- \frac{(3b^2 d^3) \int \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&+ \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&+ \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{3b^2 d^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^4} \\
&- \frac{(3b^2 d^3) \text{Subst}\left(\int \frac{\text{PolyLog}\left(2, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a^2-b^2)^2 f^4} \\
&- \frac{(3b^2 d^3) \text{Subst}\left(\int \frac{\text{PolyLog}\left(3, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a^2-b^2)^2 f^4}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&+ \frac{3b^2 d(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^3 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&- \frac{3b^2 d^3 \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4} + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{3bd^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{3b^2 d^3 \text{PolyLog}\left(4, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 5.37 (sec) , antiderivative size = 656, normalized size of antiderivative = 1.02

$$\int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx = \frac{16bc^2 f^3 (-3bd + 2acf)x + \frac{16(a-b)b^2 f^3 (c+dx)^3}{b(-1+e^{2e})+a(1+e^{2e})} - \frac{8a(a-b)bf^4 (c+dx)^4}{d(b(-1+e^{2e})+a(1+e^{2e}))} + 48bcd f^2 (bd - acf)x \log\left(1 + \frac{(a-b)e^{-2e}}{a+b}\right)}{1}$$

[In] Integrate[(c + d*x)^3/(a + b*Tanh[e + f*x])^2,x]

[Out] (16*b*c^2*f^3*(-3*b*d + 2*a*c*f)*x + (16*(a - b)*b^2*f^3*(c + d*x)^3)/(b*(-1 + E^(2*e)) + a*(1 + E^(2*e))) - (8*a*(a - b)*b*f^4*(c + d*x)^4)/(d*(b*(-1 + E^(2*e)) + a*(1 + E^(2*e)))) + 48*b*c*d*f^2*(b*d - a*c*f)*x*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))] + 24*b*d^2*f^2*(b*d - 2*a*c*f)*x^2*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))] - 16*a*b*d^3*f^3*x^3*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))] + 8*b*c^2*f^2*(3*b*d - 2*a*c*f)*Log[a - b + (a + b)*E^(2*(e + f*x))] + 24*b*c*d*f*(-(b*d) + a*c*f)*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x)))] - 12*b*d^2*(b*d - 2*a*c*f)*(2*f*x*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x)))] + PolyLog[3, (-a + b)/((a + b)*E^(2*(e + f*x)))] + 12*a*b*d^3*(2*f^2*x^2*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x)))] + 2*

$$f*x*PolyLog[3, (-a + b)/((a + b)*E^{(2*(e + f*x))})] + PolyLog[4, (-a + b)/((a + b)*E^{(2*(e + f*x))})] + ((a - b)*(a + b)*f^3*((a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[f*x] + (a^2 - b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[2*e + f*x] + 2*b*(-4*b*(c + d*x)^3 + a*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3))*Sinh[f*x])/((a*Cosh[e] + b*Sinh[e])*(a*Cosh[e + f*x] + b*Sinh[e + f*x]))/(8*(a - b)^2*(a + b)^2*f^4)$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2682 vs. $2(622) = 1244$.

Time = 0.38 (sec) , antiderivative size = 2683, normalized size of antiderivative = 4.18

method	result	size
risch	Expression too large to display	2683

[In] `int((d*x+c)^3/(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 2/(a^2+2*a*b+b^2)/f*b^2/(a-b)/(-a+b)*d^3*x^3-4/(a^2+2*a*b+b^2)/f^4*b^2/(a-b) \\ & /(-a+b)*d^3*e^3+3/2/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)/(-a+b)*d^3*polylog(3, (a+ \\ & b)*exp(2*f*x+2*e)/(-a+b))-6/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*e^2*d^3*ln(exp(\\ & f*x+e))+3/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*e^2*d^3*ln(exp(2*f*x+2*e)*a+b*exp \\ & (2*f*x+2*e)+a-b)+4/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^3*ln(exp(f*x+e))-2/(a^2+ \\ & 2*a*b+b^2)/f*b/(a-b)^2*a*c^3*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-6/(a \\ & ^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*c^2*d*ln(exp(f*x+e))+3/(a^2+2*a*b+b^2)/f^2*b^ \\ & 2/(a-b)^2*c^2*d*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-4/(a^2+2*a*b+b^2) \\ & *b/(a-b)/(-a+b)*d^2*c*a*x^3+8/(a^2+2*a*b+b^2)/f^3*b/(a-b)/(-a+b)*d^2*c*a*e^ \\ & 3-4/(a^2+2*a*b+b^2)/f^3*b/(a-b)/(-a+b)*d^3*a*e^3*x-6/(a^2+2*a*b+b^2)/f^2*b/ \\ & (a-b)*a*c^2*d/(-a+b)*e^2+12/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)/(-a+b)*d^2*c*e*x- \\ & 6/(a^2+2*a*b+b^2)*b/(a-b)*a*c^2*d/(-a+b)*x^2-3/(a^2+2*a*b+b^2)/f^3*b/(a-b)/ \\ & (-a+b)*d^2*c*a*polylog(3, (a+b)*exp(2*f*x+2*e)/(-a+b))+2/(a^2+2*a*b+b^2)/f*b \\ & /(-a+b)/(-a+b)*d^3*a*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*x^3+3/(a^2+2*a*b+b^2) \\ & /f^2*b/(a-b)/(-a+b)*d^3*a*polylog(2, (a+b)*exp(2*f*x+2*e)/(-a+b))*x^2+2/(a^2 \\ & +2*a*b+b^2)/f^4*b/(a-b)/(-a+b)*d^3*a*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*e^3- \\ & 3/(a^2+2*a*b+b^2)/f^3*b/(a-b)/(-a+b)*d^3*a*polylog(3, (a+b)*exp(2*f*x+2*e)/(- \\ & a+b))*x+12/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*e^2*d^2*c*a*ln(exp(f*x+e))-6/(a^2 \\ & +2*a*b+b^2)/f^3*b/(a-b)^2*e^2*d^2*c*a*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+ \\ & a-b)+3/(a^2+2*a*b+b^2)/f^2*b/(a-b)*a*c^2*d/(-a+b)*polylog(2, (a+b)*exp(2*f*x \\ & +2*e)/(-a+b))-6/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)/(-a+b)*d^2*c*ln(1-(a+b)*exp(2 \\ & *f*x+2*e)/(-a+b))*x-6/(a^2+2*a*b+b^2)/f^3*b^2/(a-b)/(-a+b)*d^2*c*ln(1-(a+b) \\ & *exp(2*f*x+2*e)/(-a+b))*e-12/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*e*a*c^2*d*ln(exp \\ & (f*x+e))+6/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*e*a*c^2*d*ln(exp(2*f*x+2*e)*a+b*ex \\ & p(2*f*x+2*e)+a-b)+2/(a-b)/f/(a^2+2*a*b+b^2)*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+ \\ & c^3)*b^2/(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)+1/4*d^3/(a^2+2*a*b+b^2)*x^ \\ & 4+1/4/d/(a^2+2*a*b+b^2)*c^4+2/(a^2+2*a*b+b^2)/f^4*b/(a-b)^2*e^3*d^3*a*ln(ex \end{aligned}$$

$$\begin{aligned}
& p(2fx+2e) \cdot a+b \cdot \exp(2fx+2e)+a-b)-6/(a^2+2ab+b^2)/f^3b^2/(a-b)^2 \cdot e \cdot d^2 \cdot c \cdot \ln(\exp(2fx+2e) \cdot a+b \cdot \exp(2fx+2e)+a-b)+d^2/(a^2+2ab+b^2) \cdot c \cdot x^3+3/2 \\
& \cdot d/(a^2+2ab+b^2) \cdot c^2 \cdot x^2+1/(a^2+2ab+b^2) \cdot c^3 \cdot x-6/(a^2+2ab+b^2)/f^3b^2/(a-b)/(-a+b) \cdot d^3 \cdot e^2 \cdot x+12/(a^2+2ab+b^2)/f^2b/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot a \cdot e^2 \cdot x \\
& -12/(a^2+2ab+b^2)/f \cdot b/(a-b) \cdot a \cdot c^2 \cdot d/(-a+b) \cdot e \cdot x+6/(a^2+2ab+b^2)/f \cdot b/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot a \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot x^2-6/(a^2+2ab+b^2)/f^3b^2/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot a \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot e^2+6/(a^2+2ab+b^2)/f^2b/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot a \cdot \operatorname{polylog}(2, (a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot x+6/(a^2+2ab+b^2)/f \cdot b/(a-b) \cdot a \cdot c^2 \cdot d/(-a+b) \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot x+6/(a^2+2ab+b^2)/f^2b/(a-b) \cdot a \cdot c^2 \cdot d/(-a+b) \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot e+6/(a^2+2ab+b^2)/f \cdot b^2/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot x^2+6/(a^2+2ab+b^2)/f^3b^2/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot e^2-1/(a^2+2ab+b^2) \cdot b/(a-b)/(-a+b) \cdot d^3 \cdot a \cdot x^4-3/(a^2+2ab+b^2)/f^4b/(a-b)/(-a+b) \cdot d^3 \cdot a \cdot e^4-3/(a^2+2ab+b^2)/f^2b^2/(a-b)/(-a+b) \cdot d^3 \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot x^2+3/(a^2+2ab+b^2)/f^4b^2/(a-b)/(-a+b) \cdot d^3 \cdot \ln(1-(a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot e^2-3/(a^2+2ab+b^2)/f^3b^2/(a-b)/(-a+b) \cdot d^3 \cdot \operatorname{polylog}(2, (a+b) \cdot \exp(2fx+2e)/(-a+b)) \cdot x-3/(a^2+2ab+b^2)/f^3b^2/(a-b)/(-a+b) \cdot d^2 \cdot c \cdot \operatorname{polylog}(2, (a+b) \cdot \exp(2fx+2e)/(-a+b)) +12/(a^2+2ab+b^2)/f^3b^2/(a-b)^2 \cdot e \cdot d^2 \cdot c \cdot \ln(\exp(fx+e))+3/2/(a^2+2ab+b^2)/f^4b/(a-b)/(-a+b) \cdot d^3 \cdot a \cdot \operatorname{polylog}(4, (a+b) \cdot \exp(2fx+2e)/(-a+b))-4/(a^2+2ab+b^2)/f^4b/(a-b)^2 \cdot e^3 \cdot d^3 \cdot a \cdot \ln(\exp(fx+e))
\end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6160 vs. 2(619) = 1238.

Time = 0.38 (sec) , antiderivative size = 6160, normalized size of antiderivative = 9.60

$$\int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx = \text{Too large to display}$$

[In] integrate((d*x+c)^3/(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] Too large to include

Sympy [F]

$$\int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx = \int \frac{(c+dx)^3}{(a+b \tanh(e+fx))^2} dx$$

[In] integrate((d*x+c)**3/(a+b*tanh(f*x+e))**2,x)

[Out] Integral((c + d*x)**3/(a + b*tanh(e + f*x))**2, x)

Maxima [A] (verification not implemented)

none

Time = 0.56 (sec) , antiderivative size = 1060, normalized size of antiderivative = 1.65

$$\int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3/(a+b*tanh(f*x+e))^2,x, algorithm="maxima")
```

```
[Out] -6*b^2*c^2*d*f*x/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - 2/3*(4*f^3*x^3*log((
a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 6*f^2*x^2*dilog(-(a*e^(2*e)
+ b*e^(2*e))*e^(2*f*x)/(a - b)) - 6*f*x*polylog(3, -(a*e^(2*e) + b*e^(2*e)
)*e^(2*f*x)/(a - b)) + 3*polylog(4, -(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a -
b))) * a*b*d^3/(a^4*f^4 - 2*a^2*b^2*f^4 + b^4*f^4) + 3*b^2*c^2*d*log((a*e^(2
*e) + b*e^(2*e))*e^(2*f*x) + a - b)/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - c
^3*(2*a*b*log(-(a - b)*e^(-2*f*x - 2*e) - a - b)/((a^4 - 2*a^2*b^2 + b^4)*f
) + 2*b^2/((a^4 - 2*a^2*b^2 + b^4 + (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^(-2*f
*x - 2*e))*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f)) - 3/2*(2*a*b*c*d^2*f - b
^2*d^3)*(2*f^2*x^2*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 2*f
*x*dilog(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - polylog(3, -(a*e^(2*
e) + b*e^(2*e))*e^(2*f*x)/(a - b)))/(a^4*f^4 - 2*a^2*b^2*f^4 + b^4*f^4) - 3
*(a*b*c^2*d*f - b^2*c*d^2)*(2*f*x*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a
- b) + 1) + dilog(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))/(a^4*f^3 - 2
*a^2*b^2*f^3 + b^4*f^3) + (a*b*d^3*f^4*x^4 + 2*(2*a*b*c*d^2*f - b^2*d^3)*f^
3*x^3 + 6*(a*b*c^2*d*f^2 - b^2*c*d^2*f)*f^2*x^2)/(a^4*f^4 - 2*a^2*b^2*f^4 +
b^4*f^4) + 1/4*(24*b^2*c^2*d*x + (a^2*d^3*f - 2*a*b*d^3*f + b^2*d^3*f)*x^4
+ 4*(a^2*c*d^2*f - 2*a*b*c*d^2*f + (c*d^2*f + 2*d^3)*b^2)*x^3 + 6*(a^2*c^2
*d*f - 2*a*b*c^2*d*f + (c^2*d*f + 4*c*d^2)*b^2)*x^2 + ((a^2*d^3*f*e^(2*e) -
b^2*d^3*f*e^(2*e))*x^4 + 4*(a^2*c*d^2*f*e^(2*e) - b^2*c*d^2*f*e^(2*e))*x^3
+ 6*(a^2*c^2*d*f*e^(2*e) - b^2*c^2*d*f*e^(2*e))*x^2)*e^(2*f*x))/(a^4*f - 2
*a^2*b^2*f + b^4*f + (a^4*f*e^(2*e) + 2*a^3*b*f*e^(2*e) - 2*a*b^3*f*e^(2*e)
- b^4*f*e^(2*e))*e^(2*f*x))
```

Giac [F]

$$\int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx = \int \frac{(dx + c)^3}{(b \tanh(fx + e) + a)^2} dx$$

```
[In] integrate((d*x+c)^3/(a+b*tanh(f*x+e))^2,x, algorithm="giac")
```

```
[Out] integrate((d*x + c)^3/(b*tanh(f*x + e) + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b \tanh(e + fx))^2} dx$$

```
[In] int((c + d*x)^3/(a + b*tanh(e + f*x))^2,x)
```

```
[Out] int((c + d*x)^3/(a + b*tanh(e + f*x))^2, x)
```

3.74 $\int \frac{(c+dx)^2}{(a+b \tanh(e+fx))^2} dx$

Optimal result	500
Rubi [A] (verified)	501
Mathematica [A] (verified)	507
Maple [B] (verified)	508
Fricas [B] (verification not implemented)	509
Sympy [F]	511
Maxima [A] (verification not implemented)	511
Giac [F]	512
Mupad [F(-1)]	512

Optimal result

Integrand size = 20, antiderivative size = 476

$$\begin{aligned}
 \int \frac{(c+dx)^2}{(a+b \tanh(e+fx))^2} dx = & -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} \\
 & + \frac{(c+dx)^3}{3(a-b)^2 d} + \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
 & - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
 & + \frac{2b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
 & + \frac{b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
 & - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
 & + \frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
 & + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
 & - \frac{b^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3}
 \end{aligned}$$

[Out] $-2*b^2*(d*x+c)^2/(a^2-b^2)^2/f+2*b^2*(d*x+c)^2/(a-b)/(a+b)^2/(a-b+(a+b)*\exp(2*f*x+2*e))/f+1/3*(d*x+c)^3/(a-b)^2/d+2*b^2*d*(d*x+c)*\ln(1+(a+b)*\exp(2*f*x$

$+2*e)/(a-b))/(a^2-b^2)^2/f^2-2*b*(d*x+c)^2*\ln(1+(a+b)*\exp(2*f*x+2*e)/(a-b))$
 $/ (a-b)^2/(a+b)/f+2*b^2*(d*x+c)^2*\ln(1+(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)$
 $^2/f+b^2*d^2*polylog(2,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^3-2*b*d*($
 $d*x+c)*polylog(2,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^2+2*b^2*d*(d*$
 $x+c)*polylog(2,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^2+b*d^2*polylog(3$
 $,-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^3-b^2*d^2*polylog(3,-(a+b)*\exp$
 $p(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^3$

Rubi [A] (verified)

Time = 1.14 (sec) , antiderivative size = 476, normalized size of antiderivative = 1.00,
 number of steps used = 24, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules
 used = {3815, 2221, 2611, 2320, 6724, 2286, 2216, 2215, 2222, 2317, 2438}

$$\begin{aligned}
 \int \frac{(c+dx)^2}{(a+b \tanh(e+fx))^2} dx = & \frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2-b^2)^2} \\
 & + \frac{2b^2 d(c+dx) \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f^2 (a^2-b^2)^2} \\
 & + \frac{2b^2 (c+dx)^2 \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f (a^2-b^2)^2} \\
 & - \frac{2b^2 (c+dx)^2}{f (a^2-b^2)^2} + \frac{b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2-b^2)^2} \\
 & - \frac{b^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2-b^2)^2} \\
 & + \frac{2b^2 (c+dx)^2}{f (a-b)(a+b)^2 ((a+b)e^{2e+2fx} + a-b)} \\
 & - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a-b)^2 (a+b)} \\
 & - \frac{2b(c+dx)^2 \log\left(\frac{(a+b)e^{2e+2fx}}{a-b} + 1\right)}{f (a-b)^2 (a+b)} \\
 & + \frac{(c+dx)^3}{3d(a-b)^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a-b)^2 (a+b)}
 \end{aligned}$$

[In] Int[(c + d*x)^2/(a + b*Tanh[e + f*x])^2,x]

[Out] $(-2*b^2*(c + d*x)^2)/((a^2 - b^2)^2*f) + (2*b^2*(c + d*x)^2)/((a - b)*(a +$
 $b)^2*(a - b + (a + b)*E^(2*e + 2*f*x))*f) + (c + d*x)^3/(3*(a - b)^2*d) + ($
 $2*b^2*d*(c + d*x)*Log[1 + ((a + b)*E^(2*e + 2*f*x))/(a - b)]/((a^2 - b^2)^2$

$$2*f^2) - (2*b*(c + d*x)^2*\text{Log}[1 + ((a + b)*E^{(2*e + 2*f*x)})/(a - b)])/((a - b)^2*(a + b)*f) + (2*b^2*(c + d*x)^2*\text{Log}[1 + ((a + b)*E^{(2*e + 2*f*x)})/(a - b)])/((a^2 - b^2)^2*f) + (b^2*d^2*\text{PolyLog}[2, -(((a + b)*E^{(2*e + 2*f*x)})/(a - b))]/(a - b)])/((a^2 - b^2)^2*f^3) - (2*b*d*(c + d*x)*\text{PolyLog}[2, -(((a + b)*E^{(2*e + 2*f*x)})/(a - b))]/(a - b)^2*(a + b)*f^2) + (2*b^2*d*(c + d*x)*\text{PolyLog}[2, -(((a + b)*E^{(2*e + 2*f*x)})/(a - b))]/(a^2 - b^2)^2*f^2) + (b*d^2*\text{PolyLog}[3, -(((a + b)*E^{(2*e + 2*f*x)})/(a - b))]/(a - b)^2*(a + b)*f^3) - (b^2*d^2*\text{PolyLog}[3, -(((a + b)*E^{(2*e + 2*f*x)})/(a - b))]/(a^2 - b^2)^2*f^3)$$
Rule 2215

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[(c + d*x)^(m + 1)/(a*d*(m + 1)), x] - Dist[b/a, Int[(c + d*x)^m*((F^(g*(e + f*x)))^n)/(a + b*(F^(g*(e + f*x)))^n), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2216

```
Int[((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.))^p)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Dist[1/a, Int[(c + d*x)^m*(a + b*(F^(g*(e + f*x)))^n)^(p + 1), x], x] - Dist[b/a, Int[(c + d*x)^m*(F^(g*(e + f*x)))^n*(a + b*(F^(g*(e + f*x)))^n)^p, x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && ILtQ[p, 0] && IGtQ[m, 0]
```

Rule 2221

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2222

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(p_.)*((c_.) + (d_.)*(x_))^(m_.), x_Symbol] := Simp[(c + d*x)^m*((a + b*(F^(g*(e + f*x)))^n)^(p + 1)/(b*f*g*n*(p + 1)*Log[F])), x] - Dist[d*(m/(b*f*g*n*(p + 1)*Log[F])), Int[(c + d*x)^(m - 1)*(a + b*(F^(g*(e + f*x)))^n)^(p + 1), x], x] /; FreeQ[{F, a, b, c, d, e, f, g, m, n, p}, x] && NeQ[p, -1]
```

Rule 2286

```
Int[((a_.) + (b_.)*(F_)^(u_))^(p_.)*((c_.) + (d_.)*(F_)^(v_))^(q_.)*((e_.) + (f_.)*(x_))^(m_.), x_Symbol] := With[{w = ExpandIntegrand[(e + f*x)^m, (a + b*F^u)^p*(c + d*F^v)^q, x]}, Int[w, x] /; SumQ[w]] /; FreeQ[{F, a, b, c,
```

d, e, f, m}, x] && IntegersQ[p, q] && LinearQ[{u, v}, x] && RationalQ[Simplify[u/v]]

Rule 2317

Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2320

Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_) [v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]

Rule 2438

Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 2611

Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] :> Simp[(-f + g*x)^m*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]

Rule 3815

Int[((c_.) + (d_.)*(x_)^(m_.))*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^(2*I*(e + f*x))))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]

Rule 6724

Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_)^(p_.)]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^2}{(a-b)^2} + \frac{4be^{2e+2fx}(c+dx)^2}{(a-b)^2 \left(-a\left(1-\frac{b}{a}\right) - a\left(1+\frac{b}{a}\right)e^{2e+2fx}\right)} \right. \\
&\quad \left. + \frac{4b^2e^{4e+4fx}(c+dx)^2}{(a-b)^2 \left(a\left(1-\frac{b}{a}\right) + a\left(1+\frac{b}{a}\right)e^{2e+2fx}\right)^2} \right) dx \\
&= \frac{(c+dx)^3}{3(a-b)^2d} + \frac{(4b) \int \frac{e^{2e+2fx}(c+dx)^2}{-a\left(1-\frac{b}{a}\right) - a\left(1+\frac{b}{a}\right)e^{2e+2fx}} dx}{(a-b)^2} + \frac{(4b^2) \int \frac{e^{4e+4fx}(c+dx)^2}{\left(a\left(1-\frac{b}{a}\right) + a\left(1+\frac{b}{a}\right)e^{2e+2fx}\right)^2} dx}{(a-b)^2} \\
&= \frac{(c+dx)^3}{3(a-b)^2d} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{(4b^2) \int \left(\frac{(c+dx)^2}{(a+b)^2} + \frac{(a-b)^2(c+dx)^2}{(a+b)^2(a-b+(a+b)e^{2e+2fx})^2} + \frac{2(-a+b)(c+dx)^2}{(a+b)^2(a-b+(a+b)e^{2e+2fx})} \right) dx}{(a-b)^2} \\
&\quad + \frac{(4bd) \int (c+dx) \log\left(1 + \frac{\left(1+\frac{b}{a}\right)e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f} \\
&= \frac{(c+dx)^3}{3(a-b)^2d} + \frac{4b^2(c+dx)^3}{3(a^2-b^2)^2d} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^2}{(a-b+(a+b)e^{2e+2fx})^2} dx}{(a+b)^2} \\
&\quad - \frac{(8b^2) \int \frac{(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} + \frac{(2bd^2) \int \text{PolyLog}\left(2, -\frac{\left(1+\frac{b}{a}\right)e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f^2} \\
&= \frac{(c+dx)^3}{3(a-b)^2d} - \frac{4b^2(c+dx)^3}{3(a^2-b^2)^2d} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} \\
&\quad + \frac{(8b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} - \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{(a-b+(a+b)e^{2e+2fx})^2} dx}{a^2-b^2} \\
&\quad + \frac{(bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}\left(2, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a-b)^2(a+b)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{4b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad - \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} - \frac{(4b^2d) \int \frac{c+dx}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)(a+b)^2f} \\
&\quad - \frac{(8b^2d) \int (c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} \\
&\quad + \frac{(c+dx)^3}{3(a-b)^2d} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{4b^2d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad - \frac{(4b^2d^2) \int \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2} + \frac{(4b^2d) \int \frac{e^{2e+2fx}(c+dx)}{a-b+(a+b)e^{2e+2fx}} dx}{(a-b)^2(a+b)f} \\
&\quad + \frac{(4b^2d) \int (c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2 d} \\
&+ \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} - \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{2b^2 d(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} + \frac{bd^2 \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{(2b^2 d^2) \text{Subst}\left(\int \frac{\text{PolyLog}\left(2, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^3} \\
&- \frac{(2b^2 d^2) \int \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2} + \frac{(2b^2 d^2) \int \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2} \\
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2 d} \\
&+ \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&+ \frac{2b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} - \frac{2bd(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{2b^2 d(c+dx) \text{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} + \frac{bd^2 \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&- \frac{2b^2 d^2 \text{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} - \frac{(b^2 d^2) \text{Subst}\left(\int \frac{\log\left(1 + \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^3} \\
&+ \frac{(b^2 d^2) \text{Subst}\left(\int \frac{\text{PolyLog}\left(2, -\frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b+(a+b)e^{2e+2fx})f} \\
&+ \frac{(c+dx)^3}{3(a-b)^2 d} + \frac{2b^2 d(c+dx) \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&- \frac{2b(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^2 \log\left(1 + \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
&+ \frac{b^2 d^2 \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&+ \frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&+ \frac{bd^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{b^2 d^2 \operatorname{PolyLog}\left(3, -\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.55 (sec) , antiderivative size = 516, normalized size of antiderivative = 1.08

$$\int \frac{(c+dx)^2}{(a+b \tanh(e+fx))^2} dx = \frac{24bcf^2(-bd+acf)x - \frac{24(a-b)bcf^2(-bd+acf)x}{b(-1+e^{2e})+a(1+e^{2e})} - \frac{12(a-b)bd^2(-bd+2acf)x^2}{b(-1+e^{2e})+a(1+e^{2e})} - \frac{8a(a-b)bd^2 f^3 x^3}{b(-1+e^{2e})+a(1+e^{2e})} + 12bdf(bd-2acf)}{1}$$

[In] Integrate[(c + d*x)^2/(a + b*Tanh[e + f*x])^2,x]

[Out] (24*b*c*f^2*(-(b*d) + a*c*f)*x - (24*(a - b)*b*c*f^2*(-(b*d) + a*c*f)*x)/(b*(-1 + E^(2*e)) + a*(1 + E^(2*e))) - (12*(a - b)*b*d*f^2*(-(b*d) + 2*a*c*f)*x^2)/(b*(-1 + E^(2*e)) + a*(1 + E^(2*e))) - (8*a*(a - b)*b*d^2*f^3*x^3)/(b*(-1 + E^(2*e)) + a*(1 + E^(2*e))) + 12*b*d*f*(b*d - 2*a*c*f)*x*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))] - 12*a*b*d^2*f^2*x^2*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))] + 12*b*c*f*(b*d - a*c*f)*Log[a - b + (a + b)*E^(2*(e + f*x))] - 6*b*d*(b*d - 2*a*c*f)*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x)))] + 6*a*b*d^2*(2*f*x*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x)))] + PolyLog[3, (-a + b)/((a + b)*E^(2*(e + f*x)))] + ((a - b)*(a + b)*f^2*((a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[f*x] + (a^2 - b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[2*e + f*x] + 2*b*(-3*b*(c + d*x)^2 + a*f*x*(3*c^2 + 3*c*d*x + d^2*x^2))*Sinh[f*x]))/(a*Cosh[e] + b*Sinh[e])*(a*Cosh[e + f*x] + b*Sinh[e + f*x])))/(6*(a - b)^2*(a + b)^2*f^3)

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1604 vs. $2(465) = 930$.

Time = 0.34 (sec) , antiderivative size = 1605, normalized size of antiderivative = 3.37

method	result	size
risch	Expression too large to display	1605

[In] `int((d*x+c)^2/(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & d/(a^2+2*a*b+b^2)*c*x^2+1/(a^2+2*a*b+b^2)*c^2*x+1/3*d^2/(a^2+2*a*b+b^2)*x^3 \\ & +1/3/d/(a^2+2*a*b+b^2)*c^3-8/(a^2+2*a*b+b^2)/f*b/(a-b)*c*d*a/(-a+b)*e*x+4/(\\ & a^2+2*a*b+b^2)/f^2*b/(a-b)*c*d*a/(-a+b)*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b))* \\ & e+4/(a^2+2*a*b+b^2)/f*b/(a-b)*c*d*a/(-a+b)*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b)) \\ & *x+2/(a^2+2*a*b+b^2)/f*b^2/(a-b)/(-a+b)*d^2*x^2+2/(a^2+2*a*b+b^2)/f^3*b^2/(\\ & a-b)/(-a+b)*d^2*e^2+4/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^2*\ln(\exp(f*x+e))-2/(a \\ & ^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^2*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)-4 \\ & /(a^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*c*d*\ln(\exp(f*x+e))+2/(a^2+2*a*b+b^2)/f^2*b \\ & ^2/(a-b)^2*c*d*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)-1/(a^2+2*a*b+b^2)/ \\ & f^3*b^2/(a-b)/(-a+b)*d^2*\text{polylog}(2,(a+b)*\exp(2*f*x+2*e)/(-a+b))+4/(a^2+2*a* \\ & b+b^2)/f^3*b^2/(a-b)^2*e*d^2*\ln(\exp(f*x+e))-2/(a^2+2*a*b+b^2)/f^3*b^2/(a-b) \\ & ^2*e*d^2*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)-4/3/(a^2+2*a*b+b^2)*b/(a \\ & -b)/(-a+b)*a*d^2*x^3+8/3/(a^2+2*a*b+b^2)/f^3*b/(a-b)/(-a+b)*a*d^2*e^3+4/(a^ \\ & 2+2*a*b+b^2)/f^2*b^2/(a-b)/(-a+b)*d^2*e*x-1/(a^2+2*a*b+b^2)/f^3*b/(a-b)/(-a \\ & +b)*a*d^2*\text{polylog}(3,(a+b)*\exp(2*f*x+2*e)/(-a+b))-2/(a^2+2*a*b+b^2)/f^2*b^2/ \\ & (a-b)/(-a+b)*d^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b))*x-2/(a^2+2*a*b+b^2)/f^3* \\ & b^2/(a-b)/(-a+b)*d^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b))*e+4/(a^2+2*a*b+b^2)/ \\ & f^3*b/(a-b)^2*e^2*a*d^2*\ln(\exp(f*x+e))-2/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*e^2* \\ & a*d^2*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)+2/(a-b)/f/(a^2+2*a*b+b^2)*(\\ & d^2*x^2+2*c*d*x+c^2)*b^2/(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)-4/(a^2+2*a \\ & *b+b^2)*b/(a-b)*c*d*a/(-a+b)*x^2+4/(a^2+2*a*b+b^2)/f^2*b/(a-b)/(-a+b)*a*d^2 \\ & *e^2*x-4/(a^2+2*a*b+b^2)/f^2*b/(a-b)*c*d*a/(-a+b)*e^2-8/(a^2+2*a*b+b^2)/f^2 \\ & *b/(a-b)^2*e*c*d*a*\ln(\exp(f*x+e))+4/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*e*c*d*a*\ln \\ & (\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)+a-b)+2/(a^2+2*a*b+b^2)/f*b/(a-b)/(-a+b) \\ & *a*d^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b))*x^2-2/(a^2+2*a*b+b^2)/f^3*b/(a-b)/ \\ & (-a+b)*a*d^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(-a+b))*e^2+2/(a^2+2*a*b+b^2)/f^2*b/ \\ & (a-b)/(-a+b)*a*d^2*\text{polylog}(2,(a+b)*\exp(2*f*x+2*e)/(-a+b))*x+2/(a^2+2*a*b+b^ \\ & 2)/f^2*b/(a-b)*c*d*a/(-a+b)*\text{polylog}(2,(a+b)*\exp(2*f*x+2*e)/(-a+b)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3693 vs. 2(462) = 924.

Time = 0.35 (sec) , antiderivative size = 3693, normalized size of antiderivative = 7.76

$$\int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx = \text{Too large to display}$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] 1/3*((a^3 + a^2*b - a*b^2 - b^3)*d^2*f^3*x^3 + 3*(a^3 + a^2*b - a*b^2 - b^3)*c*d*f^3*x^2 + 3*(a^3 + a^2*b - a*b^2 - b^3)*c^2*f^3*x + 4*(a^2*b - a*b^2)*d^2*e^3 + 6*(a*b^2 - b^3)*d^2*e^2 + 6*(2*(a^2*b - a*b^2)*c^2*e + (a*b^2 - b^3)*c^2)*f^2 + ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*cosh(f*x + e)^2 + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*sinh(f*x + e) + ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*sinh(f*x + e)^2 - 12*((a^2*b - a*b^2)*c*d*e^2 + (a*b^2 - b^3)*c*d*e)*f - 6*(2*(a^2*b - a*b^2)*d^2*f*x + 2*(a^2*b - a*b^2)*c*d*f - (a*b^2 - b^3)*d^2 + (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)^2 + 2*(2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e) + (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e)^2*dilog(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*(2*(a^2*b - a*b^2)*d^2*f*x + 2*(a^2*b - a*b^2)*c*d*f - (a*b^2 - b^3)*d^2 + (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)^2 + 2*(2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)*sinh(f*x + e) + (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e)^2*dilog(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*((a^2*b - a*b^2)*d^2*e^2 + (a^2*b - a*b^2)*c^2*f^2 + (a*b^2 - b^3)*d^2*e + ((a^2*b + a*b^2)*d^2*e^2 + (a^2*b + a*b^2)*c^2*f^2 + (a*b^2 + b^3)*d^2*e - (2*(a^2*b + a*b^2)*c*d*e + (a*b^2 + b^3)*c*d)*f)*cosh(f*x + e)^2 + 2*((a^2*b + a*b^2)*d^2*e^2 + (a^2*b + a*b^2)*c^2*f^2 + (a*b^2 + b^3)*d^2*e - (2*(a^2*b + a*b^2)*c*d*e + (a*b^2 + b^3)*c*d)*f)*sinh(f*x + e) + ((a^2*b + a*b^2)*

$$\begin{aligned}
& d^2e^2 + (a^2b + ab^2)c^2f^2 + (ab^2 + b^3)d^2e - (2(a^2b + ab^2) \\
&)cd*e + (ab^2 + b^3)cd*f)sinh(f*x + e)^2 - (2(a^2b - ab^2)cd*e \\
& + (ab^2 - b^3)cd*f)log(2(a + b)cosh(f*x + e) + 2(a + b)sinh(f*x + \\
& e) + 2(a - b)sqrt(-(a + b)/(a - b))) - 6((a^2b - ab^2)d^2e^2 + (a^2* \\
& b - ab^2)c^2f^2 + (ab^2 - b^3)d^2e + ((a^2b + ab^2)d^2e^2 + (a^2* \\
& b + ab^2)c^2f^2 + (ab^2 + b^3)d^2e - (2(a^2b + ab^2)cd*e + (ab^ \\
& 2 + b^3)cd*f)cosh(f*x + e)^2 + 2((a^2b + ab^2)d^2e^2 + (a^2*b + a* \\
& b^2)c^2f^2 + (ab^2 + b^3)d^2e - (2(a^2b + ab^2)cd*e + (ab^2 + b^ \\
& 3)cd*f)cosh(f*x + e)sinh(f*x + e) + ((a^2b + ab^2)d^2e^2 + (a^2*b \\
& + ab^2)c^2f^2 + (ab^2 + b^3)d^2e - (2(a^2b + ab^2)cd*e + (ab^2 \\
& + b^3)cd*f)sinh(f*x + e)^2 - (2(a^2b - ab^2)cd*e + (ab^2 - b^3)c \\
& *d*f)log(2(a + b)cosh(f*x + e) + 2(a + b)sinh(f*x + e) - 2(a - b)sq \\
& rt(-(a + b)/(a - b))) - 6((a^2b - ab^2)d^2f^2*x^2 - (a^2*b - ab^2)d^ \\
& 2e^2 + 2(a^2b - ab^2)cd*ef - (ab^2 - b^3)d^2e + ((a^2b + ab^2)* \\
& d^2f^2*x^2 - (a^2*b + ab^2)d^2e^2 + 2(a^2b + ab^2)cd*ef - (ab^2 \\
& + b^3)d^2e + (2(a^2b + ab^2)cd*f^2 - (ab^2 + b^3)d^2f)*x)cosh(f* \\
& x + e)^2 + 2((a^2b + ab^2)d^2f^2*x^2 - (a^2*b + ab^2)d^2e^2 + 2(a^ \\
& 2*b + ab^2)cd*ef - (ab^2 + b^3)d^2e + (2(a^2b + ab^2)cd*f^2 - (\\
& ab^2 + b^3)d^2f)*x)cosh(f*x + e)sinh(f*x + e) + ((a^2b + ab^2)d^2f \\
& ^2*x^2 - (a^2*b + ab^2)d^2e^2 + 2(a^2b + ab^2)cd*ef - (ab^2 + b^3 \\
&)d^2e + (2(a^2b + ab^2)cd*f^2 - (ab^2 + b^3)d^2f)*x)sinh(f*x + e \\
&)^2 + (2(a^2b - ab^2)cd*f^2 - (ab^2 - b^3)d^2f)*x)log(sqrt(-(a + b \\
&)/(a - b))(cosh(f*x + e) + sinh(f*x + e)) + 1) - 6((a^2b - ab^2)d^2f^ \\
& 2*x^2 - (a^2*b - ab^2)d^2e^2 + 2(a^2b - ab^2)cd*ef - (ab^2 - b^3) \\
& *d^2e + ((a^2b + ab^2)d^2f^2*x^2 - (a^2*b + ab^2)d^2e^2 + 2(a^2*b \\
& + ab^2)cd*ef - (ab^2 + b^3)d^2e + (2(a^2b + ab^2)cd*f^2 - (ab^ \\
& 2 + b^3)d^2f)*x)cosh(f*x + e)^2 + 2((a^2b + ab^2)d^2f^2*x^2 - (a^2* \\
& b + ab^2)d^2e^2 + 2(a^2b + ab^2)cd*ef - (ab^2 + b^3)d^2e + (2(\\
& a^2*b + ab^2)cd*f^2 - (ab^2 + b^3)d^2f)*x)cosh(f*x + e)sinh(f*x + e \\
&) + ((a^2b + ab^2)d^2f^2*x^2 - (a^2*b + ab^2)d^2e^2 + 2(a^2*b + ab \\
& ^2)cd*ef - (ab^2 + b^3)d^2e + (2(a^2b + ab^2)cd*f^2 - (ab^2 + b \\
& ^3)d^2f)*x)sinh(f*x + e)^2 + (2(a^2b - ab^2)cd*f^2 - (ab^2 - b^3)* \\
& d^2f)*x)log(-sqrt(-(a + b)/(a - b))(cosh(f*x + e) + sinh(f*x + e)) + 1) \\
& + 12((a^2b + ab^2)d^2cosh(f*x + e)^2 + 2(a^2b + ab^2)d^2cosh(f*x \\
& + e)sinh(f*x + e) + (a^2b + ab^2)d^2sinh(f*x + e)^2 + (a^2*b - ab^2)* \\
& d^2)polylog(3, sqrt(-(a + b)/(a - b))(cosh(f*x + e) + sinh(f*x + e))) + 1 \\
& 2((a^2b + ab^2)d^2cosh(f*x + e)^2 + 2(a^2b + ab^2)d^2cosh(f*x + e \\
&)sinh(f*x + e) + (a^2b + ab^2)d^2sinh(f*x + e)^2 + (a^2*b - ab^2)d^2 \\
&)polylog(3, -sqrt(-(a + b)/(a - b))(cosh(f*x + e) + sinh(f*x + e))))/((a^ \\
& 5 + a^4*b - 2a^3*b^2 - 2a^2*b^3 + ab^4 + b^5)*f^3cosh(f*x + e)^2 + 2(a \\
& ^5 + a^4*b - 2a^3*b^2 - 2a^2*b^3 + ab^4 + b^5)*f^3cosh(f*x + e)sinh(f* \\
& x + e) + (a^5 + a^4*b - 2a^3*b^2 - 2a^2*b^3 + ab^4 + b^5)*f^3sinh(f*x + \\
& e)^2 + (a^5 - a^4*b - 2a^3*b^2 + 2a^2*b^3 + ab^4 - b^5)*f^3)
\end{aligned}$$

SymPy [F]

$$\int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx$$

[In] integrate((d*x+c)**2/(a+b*tanh(f*x+e))**2,x)

[Out] Integral((c + d*x)**2/(a + b*tanh(e + f*x))**2, x)

Maxima [A] (verification not implemented)

none

Time = 0.52 (sec) , antiderivative size = 753, normalized size of antiderivative = 1.58

$$\begin{aligned} \int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx = & -\frac{4b^2cdfx}{a^4f^2 - 2a^2b^2f^2 + b^4f^2} \\ & - \frac{\left(2f^2x^2 \log\left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} + 1\right) + 2fx \operatorname{Li}_2\left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b}\right) - \operatorname{Li}_3\left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b}\right)\right)abd^2}{a^4f^3 - 2a^2b^2f^3 + b^4f^3} \\ & + \frac{2b^2cd \log\left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)} + a - b}{a^4f^2 - 2a^2b^2f^2 + b^4f^2}\right)}{a^4f^2 - 2a^2b^2f^2 + b^4f^2} \\ & - c^2 \left(\frac{2ab \log\left(\frac{-(a-b)e^{(-2fx-2e)} - a - b}{(a^4 - 2a^2b^2 + b^4)f}\right) + \frac{2b^2}{(a^4 - 2a^2b^2 + b^4 + (a^4 - 2a^3b + 2ab^3 - b^4)e^{(-2fx-2e)})f}}{a^4f^3 - 2a^2b^2f^3 + b^4f^3} \right. \\ & \left. - \frac{(2abcdf - b^2d^2)\left(2fx \log\left(\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b} + 1\right) + \operatorname{Li}_2\left(-\frac{(ae^{(2e)} + be^{(2e)})e^{(2fx)}}{a-b}\right)\right)}{a^4f^3 - 2a^2b^2f^3 + b^4f^3} \right) \\ & + \frac{2(2abd^2f^3x^3 + 3(2abcdf - b^2d^2)f^2x^2)}{3(a^4f^3 - 2a^2b^2f^3 + b^4f^3)} \\ & + \frac{12b^2cdx + (a^2d^2f - 2abd^2f + b^2d^2f)x^3 + 3(a^2cdf - 2abcdf + (cdf + 2d^2)b^2)x^2 + ((a^2d^2fe^{(2e)} - b^2d^2)}{3(a^4f - 2a^2b^2f + b^4f + (a^4fe^{(2e)} + 2a^3bfe^{(2e)} - 2ab^3fe^{(2e)} - b^4f^2))} \end{aligned}$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] $-4*b^2*c*d*f*x/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - (2*f^2*x^2*\log((a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)/(a - b)} + 1) + 2*f*x*dilog(-(a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)/(a - b)}) - polylog(3, -(a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)/(a - b)}))a*b*d^2/(a^4*f^3 - 2*a^2*b^2*f^3 + b^4*f^3) + 2*b^2*c*d*\log((a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)} + a - b)/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - c^2*(2*a*b*\log(-(a - b)*e^{(-2*f*x - 2e)} - a - b)/((a^4 - 2*a^2*b^2 + b^4)*f) + 2*b^2/((a^4 - 2*a^2*b^2 + b^4 + (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^{(-2*f*x - 2e)})*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f)) - (2*a*b*c*d*f - b^2*d^2)*(2*f*x*\log((a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)/(a - b)} + 1) + dilog(-(a*e^{(2e)} + b*e^{(2e)})e^{(2*f*x)/(a - b)}))$

$e) + b \cdot e^{(2e)} \cdot e^{(2fx)} / (a - b) \Big) / (a^4 f^3 - 2a^2 b^2 f^3 + b^4 f^3) + 2 / 3 \cdot (2ab d^2 f^3 x^3 + 3(2abc d f - b^2 d^2) f^2 x^2) / (a^4 f^3 - 2a^2 b^2 f^3 + b^4 f^3) + 1/3 \cdot (12b^2 c d x + (a^2 d^2 f - 2abd^2 f + b^2 d^2 f) x^3 + 3(a^2 c d f - 2abc d f + (c d f + 2d^2) b^2) x^2 + ((a^2 d^2 f e^{(2e)} - b^2 d^2 f e^{(2e)}) x^3 + 3(a^2 c d f e^{(2e)} - b^2 c d f e^{(2e)}) x^2) \cdot e^{(2fx)}) / (a^4 f - 2a^2 b^2 f + b^4 f + (a^4 f e^{(2e)} + 2a^3 b f e^{(2e)} - 2a b^3 f e^{(2e)} - b^4 f e^{(2e)}) \cdot e^{(2fx)})$

Giac [F]

$$\int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx = \int \frac{(dx + c)^2}{(b \tanh(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)^2/(b*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx = \int \frac{(c + dx)^2}{(a + b \tanh(e + fx))^2} dx$$

[In] int((c + d*x)^2/(a + b*tanh(e + f*x))^2,x)

[Out] int((c + d*x)^2/(a + b*tanh(e + f*x))^2, x)

3.75 $\int \frac{c+dx}{(a+b \tanh(e+fx))^2} dx$

Optimal result	513
Rubi [A] (verified)	513
Mathematica [A] (verified)	516
Maple [B] (verified)	516
Fricas [B] (verification not implemented)	517
Sympy [F]	518
Maxima [F]	518
Giac [F]	519
Mupad [F(-1)]	519

Optimal result

Integrand size = 18, antiderivative size = 196

$$\int \frac{c+dx}{(a+b \tanh(e+fx))^2} dx = -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adf x)^2}{4a(a-b)(a+b)^2df^2}$$

$$+ \frac{b(bd-2acf-2adf x) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2 f^2}$$

$$+ \frac{abd \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2 f^2}$$

$$+ \frac{b(c+dx)}{(a^2-b^2)f(a+b \tanh(e+fx))}$$

```
[Out] -1/2*(d*x+c)^2/(a^2-b^2)/d+1/4*(-2*a*d*f*x-2*a*c*f+b*d)^2/a/(a-b)/(a+b)^2/d
/f^2+b*(-2*a*d*f*x-2*a*c*f+b*d)*ln(1+(a-b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)^
2/f^2+a*b*d*polylog(2,(-a+b)/(a+b)/exp(2*f*x+2*e))/(a^2-b^2)^2/f^2+b*(d*x+c
)/(a^2-b^2)/f/(a+b*tanh(f*x+e))
```

Rubi [A] (verified)

Time = 0.21 (sec) , antiderivative size = 196, normalized size of antiderivative = 1.00,
 number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used

= {3814, 3813, 2221, 2317, 2438}

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \frac{b(-2acf - 2adfx + bd) \log\left(\frac{(a-b)e^{-2(e+fx)}}{a+b} + 1\right)}{f^2 (a^2 - b^2)^2} + \frac{b(c + dx)}{f (a^2 - b^2) (a + b \tanh(e + fx))} - \frac{(c + dx)^2}{2d (a^2 - b^2)} + \frac{abd \operatorname{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f^2 (a^2 - b^2)^2} + \frac{(-2acf - 2adfx + bd)^2}{4adf^2 (a - b)(a + b)^2}$$

[In] Int[(c + d*x)/(a + b*Tanh[e + f*x])^2,x]

[Out] -1/2*(c + d*x)^2/((a^2 - b^2)*d) + (b*d - 2*a*c*f - 2*a*d*f*x)^2/(4*a*(a - b)*(a + b)^2*d*f^2) + (b*(b*d - 2*a*c*f - 2*a*d*f*x)*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x)))]/((a^2 - b^2)^2*f^2) + (a*b*d*PolyLog[2, -((a - b)/((a + b)*E^(2*(e + f*x))))])/((a^2 - b^2)^2*f^2) + (b*(c + d*x))/((a^2 - b^2)*f*(a + b*Tanh[e + f*x]))

Rule 2221

Int[(((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)*((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*((F_)^((g_)*((e_) + (f_)*(x_))))^(n_)), x_Symbol] := Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]

Rule 2317

Int[Log[(a_) + (b_)*((F_)^((e_)*((c_) + (d_)*(x_))))^(n_)], x_Symbol] := Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]

Rule 2438

Int[Log[(c_)*((d_) + (e_)*(x_)^(n_))]/(x_), x_Symbol] := Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]

Rule 3813

Int[(((c_) + (d_)*(x_))^(m_))/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Symbol] := Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^Simp[2*I*(e + f*x), x])], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]

Rule 3814

```

Int[((c_.) + (d_.)*(x_))/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^2, x_Symbol
] :> Simp[-(c + d*x)^2/(2*d*(a^2 + b^2)), x] + (Dist[1/(f*(a^2 + b^2)), Int
[(b*d + 2*a*c*f + 2*a*d*f*x)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((c + d*
x)/(f*(a^2 + b^2)*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f},
x] && NeQ[a^2 + b^2, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c + dx)^2}{2(a^2 - b^2)d} + \frac{b(c + dx)}{(a^2 - b^2)f(a + b \tanh(e + fx))} - \frac{i \int \frac{-ibd + 2iacf + 2iadfx}{a + b \tanh(e + fx)} dx}{(a^2 - b^2)f} \\
&= -\frac{(c + dx)^2}{2(a^2 - b^2)d} + \frac{(bd - 2acf - 2adfx)^2}{4a(a - b)(a + b)^2df^2} \\
&\quad + \frac{b(c + dx)}{(a^2 - b^2)f(a + b \tanh(e + fx))} - \frac{(2ib) \int \frac{e^{-2(e+fx)}(-ibd + 2iacf + 2iadfx)}{(a+b)^2 + (a^2 - b^2)e^{-2(e+fx)}} dx}{(a^2 - b^2)f} \\
&= -\frac{(c + dx)^2}{2(a^2 - b^2)d} + \frac{(bd - 2acf - 2adfx)^2}{4a(a - b)(a + b)^2df^2} + \frac{b(bd - 2acf - 2adfx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)^2 f^2} \\
&\quad + \frac{b(c + dx)}{(a^2 - b^2)f(a + b \tanh(e + fx))} + \frac{(2abd) \int \log\left(1 + \frac{(a^2 - b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2 - b^2)^2 f} \\
&= -\frac{(c + dx)^2}{2(a^2 - b^2)d} + \frac{(bd - 2acf - 2adfx)^2}{4a(a - b)(a + b)^2df^2} + \frac{b(bd - 2acf - 2adfx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)^2 f^2} \\
&\quad + \frac{b(c + dx)}{(a^2 - b^2)f(a + b \tanh(e + fx))} - \frac{(abd) \text{Subst}\left(\int \frac{\log\left(1 + \frac{(a^2 - b^2)x}{(a+b)^2}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{(a^2 - b^2)^2 f^2} \\
&= -\frac{(c + dx)^2}{2(a^2 - b^2)d} + \frac{(bd - 2acf - 2adfx)^2}{4a(a - b)(a + b)^2df^2} \\
&\quad + \frac{b(bd - 2acf - 2adfx) \log\left(1 + \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)^2 f^2} \\
&\quad + \frac{abd \text{PolyLog}\left(2, -\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)^2 f^2} + \frac{b(c + dx)}{(a^2 - b^2)f(a + b \tanh(e + fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.64 (sec) , antiderivative size = 249, normalized size of antiderivative = 1.27

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx$$

$$= \frac{\operatorname{sech}^2(e + fx)(a \cosh(e + fx) + b \sinh(e + fx)) \left(-\frac{2b^2 f(c+dx) \sinh(e+fx)}{a} - (e + fx)(-2cf + d(e - fx))(a \cosh(e + fx) + b \sinh(e + fx)) \right)}{2(a + b \tanh(e + fx))^2}$$

```
[In] Integrate[(c + d*x)/(a + b*Tanh[e + f*x])^2,x]
```

```
[Out] (Sech[e + f*x]^2*(a*Cosh[e + f*x] + b*Sinh[e + f*x])*((-2*b^2*f*(c + d*x)*Sinh[e + f*x])/a - (e + f*x)*(-2*c*f + d*(e - f*x))*(a*Cosh[e + f*x] + b*Sinh[e + f*x]) + (b*((b*d - 2*a*f*(c + d*x))*((a - b)*(-(b*d) + 2*a*f*(c + d*x)) + 4*a^2*d*Log[1 + (a - b)/((a + b)*E^(2*(e + f*x))])) + 4*a^3*d^2*PolyLog[2, (-a + b)/((a + b)*E^(2*(e + f*x))]))*(a*Cosh[e + f*x] + b*Sinh[e + f*x]))/(2*a^2*(a - b)*(a + b)*d))/(2*(a - b)*(a + b)*f^2*(a + b*Tanh[e + f*x])^2)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 660 vs. 2(195) = 390.

Time = 0.31 (sec) , antiderivative size = 661, normalized size of antiderivative = 3.37

method	result
risch	$\frac{dx^2}{2a^2+4ab+2b^2} + \frac{cx}{a^2+2ab+b^2} + \frac{2(dx+c)b^2}{(a-b)f(a^2+2ab+b^2)(e^{2fx+2e}a+be^{2fx+2e}+a-b)} - \frac{2b^2 d \ln(e^{fx+e})}{(a^2+2ab+b^2)f^2(a-b)^2} + \frac{b^2 d \ln(e^{2fx+2e}a+be^{2fx+2e}+a-b)}{(a^2+2ab+b^2)f^2(a-b)^2}$

```
[In] int((d*x+c)/(a+b*tanh(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/2/(a^2+2*a*b+b^2)*d*x^2+1/(a^2+2*a*b+b^2)*c*x+2/(a-b)/f/(a^2+2*a*b+b^2)*(d*x+c)*b^2/(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-2/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*d*ln(exp(f*x+e))+1/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*d*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)+4/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c*ln(exp(f*x+e))-2/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-4/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*a*d*e*ln(exp(f*x+e))+2/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*a*d*e*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)+a-b)-2/(a^2+2*a*b+b^2)*b/(a-b)*d*a/(-a+b)*x^2+2/(a^2+2*a*b+b^2)/f*b/(a-b)*d*a/(-a+b)*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*x-4/(a^2+2*a*b+b^2)/f*b/(a-b)*d*a/(-a+b)*e*x+2/(a^2+2*a*b+b^2)/f^2*b/(a-b)*d*a/(-a+b)*ln(1-(a+b)*exp(2*f*x+2*e)/(-a+b))*e-2/(a^2+2*a*b+b^2)/f^2*b/(a-b)*d*a/(-a+b)*e^2+1/(a^2+2*a*b+b^2)/f^2*b/(a-b)*d*a/(-a+b)*polylog(2,(a+b)*exp(2*f*x+2*e)/(-a+b))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1790 vs. 2(194) = 388.

Time = 0.31 (sec) , antiderivative size = 1790, normalized size of antiderivative = 9.13

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \text{Too large to display}$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] 1/2*((a^3 + a^2*b - a*b^2 - b^3)*d*f^2*x^2 + 2*(a^3 + a^2*b - a*b^2 - b^3)*c*f^2*x - 4*(a^2*b - a*b^2)*d*e^2 - 4*(a*b^2 - b^3)*d*e + ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*f^2*x^2 - 4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*e*f - 4*(a*b^2 + b^3)*d*e + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 + b^3)*d*f)*x)*cosh(f*x + e)^2 + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*f^2*x^2 - 4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*e*f - 4*(a*b^2 + b^3)*d*e + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 + b^3)*d*f)*x)*cosh(f*x + e)*sinh(f*x + e) + ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*f^2*x^2 - 4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*e*f - 4*(a*b^2 + b^3)*d*e + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 + b^3)*d*f)*x)*sinh(f*x + e)^2 + 4*(2*(a^2*b - a*b^2)*c*e + (a*b^2 - b^3)*c)*f - 4*((a^2*b + a*b^2)*d*cosh(f*x + e)^2 + 2*(a^2*b + a*b^2)*d*cosh(f*x + e)*sinh(f*x + e) + (a^2*b + a*b^2)*d*sinh(f*x + e)^2 + (a^2*b - a*b^2)*d)*dilog(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 4*((a^2*b + a*b^2)*d*cosh(f*x + e)^2 + 2*(a^2*b + a*b^2)*d*cosh(f*x + e)*sinh(f*x + e) + (a^2*b + a*b^2)*d*sinh(f*x + e)^2 + (a^2*b - a*b^2)*d)*dilog(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) + 2*(2*(a^2*b - a*b^2)*d*e - 2*(a^2*b - a*b^2)*c*f + (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)^2 + 2*(2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)*sinh(f*x + e) + (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*sinh(f*x + e)^2 + (a*b^2 - b^3)*d)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt(-(a + b)/(a - b))) + 2*(2*(a^2*b - a*b^2)*d*e - 2*(a^2*b - a*b^2)*c*f + (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)^2 + 2*(2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)*sinh(f*x + e) + (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*sinh(f*x + e)^2 + (a*b^2 - b^3)*d)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) - 2*(a - b)*sqrt(-(a + b)/(a - b))) - 4*((a^2*b - a*b^2)*d*f*x + (a^2*b - a*b^2)*d*e + ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*x + e)^2 + 2*((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*x + e)*sinh(f*x + e) + ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*sinh(f*x + e)^2)*log(sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 4*((a^2*b - a*b^2)*d*f*x + (a^2*b - a*b^2)*d*e + ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*x + e)^2 + 2*((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*x + e)*sinh(f*x + e) + ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*sinh(f*x + e)^2)

2)*d*e)*sinh(f*x + e)^2)*log(-sqrt(-(a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1))/((a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*cosh(f*x + e)^2 + 2*(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*cosh(f*x + e)*sinh(f*x + e) + (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*sinh(f*x + e)^2 + (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*f^2)

Sympy [F]

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e))**2,x)

[Out] Integral((c + d*x)/(a + b*tanh(e + f*x))**2, x)

Maxima [F]

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \int \frac{dx + c}{(b \tanh(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

[Out] 1/2*(8*a*b*f*integrate(x/(a^4*f*e^(2*f*x + 2*e) + 2*a^3*b*f*e^(2*f*x + 2*e) - 2*a*b^3*f*e^(2*f*x + 2*e) - b^4*f*e^(2*f*x + 2*e) + a^4*f - 2*a^2*b^2*f + b^4*f), x) - 2*b^2*(2*(f*x + e)/((a^4 - 2*a^2*b^2 + b^4)*f^2) - log((a + b)*e^(2*f*x + 2*e) + a - b)/((a^4 - 2*a^2*b^2 + b^4)*f^2)) + ((a^2*f*e^(2*e) - b^2*f*e^(2*e))*x^2*e^(2*f*x) + 4*b^2*x + (a^2*f - 2*a*b*f + b^2*f)*x^2)/(a^4*f - 2*a^2*b^2*f + b^4*f + (a^4*f*e^(2*e) + 2*a^3*b*f*e^(2*e) - 2*a*b^3*f*e^(2*e) - b^4*f*e^(2*e))*e^(2*f*x))*d - c*(2*a*b*log(-(a - b)*e^(-2*f*x - 2*e) - a - b)/((a^4 - 2*a^2*b^2 + b^4)*f) + 2*b^2/((a^4 - 2*a^2*b^2 + b^4 + (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^(-2*f*x - 2*e))*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f))

Giac [F]

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \int \frac{dx + c}{(b \tanh(fx + e) + a)^2} dx$$

[In] integrate((d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate((d*x + c)/(b*tanh(f*x + e) + a)^2, x)

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx = \int \frac{c + dx}{(a + b \tanh(e + fx))^2} dx$$

[In] int((c + d*x)/(a + b*tanh(e + f*x))^2,x)

[Out] int((c + d*x)/(a + b*tanh(e + f*x))^2, x)

$$3.76 \quad \int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx$$

Optimal result	520
Rubi [N/A]	520
Mathematica [N/A]	521
Maple [N/A] (verified)	521
Fricas [N/A]	521
Sympy [N/A]	522
Maxima [N/A]	522
Giac [N/A]	522
Mupad [N/A]	523

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b \tanh(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx = \int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx$$

[In] Int[1/((c + d*x)*(a + b*Tanh[e + f*x]))^2), x]

[Out] Defer[Int][1/((c + d*x)*(a + b*Tanh[e + f*x]))^2), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 31.58 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))^2} dx = \int \frac{1}{(c + dx)(a + b \tanh(e + fx))^2} dx$$

[In] Integrate[1/((c + d*x)*(a + b*Tanh[e + f*x])^2), x]

[Out] Integrate[1/((c + d*x)*(a + b*Tanh[e + f*x])^2), x]

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \tanh(fx + e))^2} dx$$

[In] int(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x)

[Out] int(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x)

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \tanh(fx + e) + a)^2} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

[Out] integral(1/(a^2*d*x + a^2*c + (b^2*d*x + b^2*c)*tanh(f*x + e)^2 + 2*(a*b*d*x + a*b*c)*tanh(f*x + e)), x)

Sympy [N/A]

Not integrable

Time = 1.60 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx = \int \frac{1}{(a+b \tanh(e+fx))^2 (c+dx)} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e))**2,x)

[Out] Integral(1/((a + b*tanh(e + f*x))**2*(c + d*x)), x)

Maxima [N/A]

Not integrable

Time = 1.11 (sec) , antiderivative size = 468, normalized size of antiderivative = 23.40

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx = \int \frac{1}{(dx+c)(b \tanh(fx+e)+a)^2} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

```
[Out] 2*b^2/(a^4*c*f - 2*a^2*b^2*c*f + b^4*c*f + (a^4*d*f - 2*a^2*b^2*d*f + b^4*d*f)*x + (a^4*c*f*e^(2*e) + 2*a^3*b*c*f*e^(2*e) - 2*a*b^3*c*f*e^(2*e) - b^4*c*f*e^(2*e) + (a^4*d*f*e^(2*e) + 2*a^3*b*d*f*e^(2*e) - 2*a*b^3*d*f*e^(2*e) - b^4*d*f*e^(2*e))*x)*e^(2*f*x)) + log(d*x + c)/(a^2*d + 2*a*b*d + b^2*d) + integrate(2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)/(a^4*c^2*f - 2*a^2*b^2*c^2*f + b^4*c^2*f + (a^4*d^2*f - 2*a^2*b^2*d^2*f + b^4*d^2*f)*x^2 + 2*(a^4*c*d*f - 2*a^2*b^2*c*d*f + b^4*c*d*f)*x + (a^4*c^2*f*e^(2*e) + 2*a^3*b*c^2*f*e^(2*e) - 2*a*b^3*c^2*f*e^(2*e) - b^4*c^2*f*e^(2*e) + (a^4*d^2*f*e^(2*e) + 2*a^3*b*d^2*f*e^(2*e) - 2*a*b^3*d^2*f*e^(2*e) - b^4*d^2*f*e^(2*e))*x^2 + 2*(a^4*c*d*f*e^(2*e) + 2*a^3*b*c*d*f*e^(2*e) - 2*a*b^3*c*d*f*e^(2*e) - b^4*c*d*f*e^(2*e))*x)*e^(2*f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)(a+b \tanh(e+fx))^2} dx = \int \frac{1}{(dx+c)(b \tanh(fx+e)+a)^2} dx$$

[In] integrate(1/(d*x+c)/(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate(1/((d*x + c)*(b*tanh(f*x + e) + a)^2), x)

Mupad [N/A]

Not integrable

Time = 2.47 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \tanh(e + fx))^2} dx = \int \frac{1}{(a + b \tanh(e + fx))^2 (c + dx)} dx$$

```
[In] int(1/((a + b*tanh(e + f*x))^2*(c + d*x)),x)
```

```
[Out] int(1/((a + b*tanh(e + f*x))^2*(c + d*x)), x)
```

$$3.77 \quad \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2} dx$$

Optimal result	524
Rubi [N/A]	524
Mathematica [N/A]	525
Maple [N/A] (verified)	525
Fricas [N/A]	525
Sympy [N/A]	526
Maxima [N/A]	526
Giac [N/A]	527
Mupad [N/A]	527

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2} dx = \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2} dx$$

[In] Int[1/((c + d*x)^2*(a + b*Tanh[e + f*x]))^2, x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*Tanh[e + f*x]))^2, x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b \tanh(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 28.49 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))^2} dx = \int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))^2} dx$$

[In] Integrate[1/((c + d*x)^2*(a + b*Tanh[e + f*x])^2), x]

[Out] Integrate[1/((c + d*x)^2*(a + b*Tanh[e + f*x])^2), x]

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2(a + b \tanh(fx + e))^2} dx$$

[In] int(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x)

[Out] int(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x)

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 96, normalized size of antiderivative = 4.80

$$\int \frac{1}{(c + dx)^2(a + b \tanh(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \tanh(fx + e) + a)^2} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="fricas")

```
[Out] integral(1/(a^2*d^2*x^2 + 2*a^2*c*d*x + a^2*c^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*tanh(f*x + e)^2 + 2*(a*b*d^2*x^2 + 2*a*b*c*d*x + a*b*c^2)*tanh(f*x + e)), x)
```

Sympy [N/A]

Not integrable

Time = 3.14 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))^2} dx = \int \frac{1}{(a + b \tanh(e + fx))^2 (c + dx)^2} dx$$

[In] integrate(1/(d*x+c)**2/(a+b*tanh(f*x+e))**2,x)

[Out] Integral(1/((a + b*tanh(e + f*x))**2*(c + d*x)**2), x)

Maxima [N/A]

Not integrable

Time = 2.11 (sec) , antiderivative size = 784, normalized size of antiderivative = 39.20

$$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))^2} dx = \int \frac{1}{(dx + c)^2 (b \tanh(fx + e) + a)^2} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="maxima")

```
[Out] -(a^2*c*f - 2*a*b*c*f + (c*f - 2*d)*b^2 + (a^2*d*f - 2*a*b*d*f + b^2*d*f)*x
+ (a^2*c*f*e^(2*e) - b^2*c*f*e^(2*e) + (a^2*d*f*e^(2*e) - b^2*d*f*e^(2*e))
*x)*e^(2*f*x))/(a^4*c^2*d*f - 2*a^2*b^2*c^2*d*f + b^4*c^2*d*f + (a^4*d^3*f
- 2*a^2*b^2*d^3*f + b^4*d^3*f)*x^2 + 2*(a^4*c*d^2*f - 2*a^2*b^2*c*d^2*f + b
^4*c*d^2*f)*x + (a^4*c^2*d*f*e^(2*e) + 2*a^3*b*c^2*d*f*e^(2*e) - 2*a*b^3*c^
2*d*f*e^(2*e) - b^4*c^2*d*f*e^(2*e) + (a^4*d^3*f*e^(2*e) + 2*a^3*b*d^3*f*e^
(2*e) - 2*a*b^3*d^3*f*e^(2*e) - b^4*d^3*f*e^(2*e))*x^2 + 2*(a^4*c*d^2*f*e^(
2*e) + 2*a^3*b*c*d^2*f*e^(2*e) - 2*a*b^3*c*d^2*f*e^(2*e) - b^4*c*d^2*f*e^(2
*e))*x)*e^(2*f*x)) + integrate(4*(a*b*d*f*x + a*b*c*f + b^2*d)/(a^4*c^3*f -
2*a^2*b^2*c^3*f + b^4*c^3*f + (a^4*d^3*f - 2*a^2*b^2*d^3*f + b^4*d^3*f)*x^
3 + 3*(a^4*c*d^2*f - 2*a^2*b^2*c*d^2*f + b^4*c*d^2*f)*x^2 + 3*(a^4*c^2*d*f
- 2*a^2*b^2*c^2*d*f + b^4*c^2*d*f)*x + (a^4*c^3*f*e^(2*e) + 2*a^3*b*c^3*f*e
^(2*e) - 2*a*b^3*c^3*f*e^(2*e) - b^4*c^3*f*e^(2*e) + (a^4*d^3*f*e^(2*e) + 2
*a^3*b*d^3*f*e^(2*e) - 2*a*b^3*d^3*f*e^(2*e) - b^4*d^3*f*e^(2*e))*x^3 + 3*(
a^4*c*d^2*f*e^(2*e) + 2*a^3*b*c*d^2*f*e^(2*e) - 2*a*b^3*c*d^2*f*e^(2*e) - b
^4*c*d^2*f*e^(2*e))*x^2 + 3*(a^4*c^2*d*f*e^(2*e) + 2*a^3*b*c^2*d*f*e^(2*e)
- 2*a*b^3*c^2*d*f*e^(2*e) - b^4*c^2*d*f*e^(2*e))*x)*e^(2*f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))^2} dx = \int \frac{1}{(dx + c)^2 (b \tanh(fx + e) + a)^2} dx$$

[In] integrate(1/(d*x+c)^2/(a+b*tanh(f*x+e))^2,x, algorithm="giac")

[Out] integrate(1/((d*x + c)^2*(b*tanh(f*x + e) + a)^2), x)

Mupad [N/A]

Not integrable

Time = 2.08 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2 (a + b \tanh(e + fx))^2} dx = \int \frac{1}{(a + b \tanh(e + fx))^2 (c + dx)^2} dx$$

[In] int(1/((a + b*tanh(e + f*x))^2*(c + d*x)^2),x)

[Out] int(1/((a + b*tanh(e + f*x))^2*(c + d*x)^2), x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions 529

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*      Small rewrite of logic in main function to make it*)
(*      match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCo
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A",""}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count is
        ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A",""}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"}
    ]
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

  finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

```

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
          If[Head[expn]===RootSum,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,

```

```

    Sinh, Cosh, Tanh, Coth, Sech, Csch,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
  }, func]

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,
    ExpIntegralE, ExpIntegralEi, LogIntegral,
    SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
    Gamma, LogGamma, PolyGamma,
    Zeta, PolyLog, ProductLog,
    EllipticF, EllipticE, EllipticPi
  }, func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result, optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 ("
```

```

                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
    end if
else #result contains complex but optimal is not
    if debug then
        print("result contains complex but optimal is not");
    fi;
    return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A"," ";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of opt
                                convert(leaf_count_result,string)," $ vs. $2(",
                                convert(leaf_count_optimal,string)," )=",convert(2*leaf_count
    fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                    convert(ExpnType_result,string)," vs. order ",
                    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func, [
    exp, log, ln,
    sin, cos, tan, cot, sec, csc,
    arcsin, arccos, arctan, arccot, arcsec, arccsc,
    sinh, cosh, tanh, coth, sech, csch,
    arcsinh, arccosh, arctanh, arccoth, arcsech, arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf, erfc, erfi,
    FresnelS, FresnelC,
    Ei, Ei, Li, Si, Ci, Shi, Chi,
    GAMMA, lnGAMMA, Psi, Zeta, polylog, dilog, LambertW,
    EllipticF, EllipticE, EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1, hypergeom, HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u), op(2..nops(u), u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```


Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+') or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is large"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)+str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)+str(ExpnType_optimal)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
    try:
        if expn.parent() is SR:
            return expn.operator() is None
        if expn.parent() in (ZZ, QQ, AA, QQbar):
            return expn in expn.parent() # Should always return True
        if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
            return expn in expn.parent().base_ring() or expn in expn.parent().gens()

        return False

    except AttributeError as error:
        print("Exception,AttributeError in is_atom")
        print ("caught exception" , type(error).__name__ )
        return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinst
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. " + str(leaf_c

else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order " + str(expnType_result

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```