

Computer Algebra Independent Integration Tests

Summer 2023 edition

6-Hyperbolic-functions/6.4-Hyperbolic-cotangent/174-6.4.1-c+d-x-
 $\hat{m}-a+b-\coth-\hat{n}$

Nasser M. Abbasi

September 5, 2023 Compiled on September 5, 2023 at 11:54am

Contents

1	Introduction	3
2	detailed summary tables of results	21
3	Listing of integrals	41
4	Appendix	417

CHAPTER 1

INTRODUCTION

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	8
1.4	Performance based on number of rules Rubi used	10
1.5	Performance based on number of steps Rubi used	11
1.6	Solved integrals histogram based on leaf size of result	12
1.7	Solved integrals histogram based on CPU time used	13
1.8	Leaf size vs. CPU time used	14
1.9	list of integrals with no known antiderivative	15
1.10	List of integrals solved by CAS but has no known antiderivative	15
1.11	list of integrals solved by CAS but failed verification	15
1.12	Timing	16
1.13	Verification	16
1.14	Important notes about some of the results	16
1.15	Design of the test system	19

This report gives the result of running the computer algebra independent integration test. The download section in on the main webpage contains links to download the problems in plain text format used for all CAS systems. The number of integrals in this report is [61]. This is test number [174].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.3.1 (August 16, 2023) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.3 on windows 10
3. Maple 2023.1 (July, 12, 2023) on windows 10.
4. Maxima 5.47 (June 1, 2023) using Lisp SBCL 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
5. FriCAS 1.3.9 (July 8, 2023) based on sbcl 2.3.0 on Linux via sagemath 10.1 (Aug 20, 2023).
6. Giac/Xcas 1.9.0-57 (June 26, 2023) on Linux via sagemath 10.1 (Aug 20, 2023).
7. Sympy 1.12 (May 10, 2023) Using Python 3.11.3 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was run directly in Python not via sagemath.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed. If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (61)	0.00 (0)
Mathematica	100.00 (61)	0.00 (0)
Fricas	100.00 (61)	0.00 (0)
Maple	95.08 (58)	4.92 (3)
Maxima	90.16 (55)	9.84 (6)
Giac	57.38 (35)	42.62 (26)
Mupad	45.90 (28)	54.10 (33)
Sympy	45.90 (28)	54.10 (33)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	70.492	0.000	0.000	29.508
Mathematica	60.656	9.836	0.000	29.508
Maxima	36.066	24.590	0.000	39.344
Maple	27.869	37.705	0.000	34.426
Fricas	24.590	45.902	0.000	29.508
Giac	24.590	3.279	0.000	72.131
Mupad	0.000	16.393	0.000	83.607
Sympy	0.000	16.393	0.000	83.607

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the grades of the CAS systems.



The following table shows the distribution of the different types of failures for each CAS. There are 3 types failures. The first is when CAS returns the input within the time limit, which means it could not solve it. This is the typical failure and given as **F**.

The second failure is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned. This is assigned **F(-1)**.

The third is due to an exception generated, indicated as **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in the CAS itself. This type of

error requires more investigation to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00	0.00	0.00
Mathematica	0	0.00	0.00	0.00
Fricas	0	0.00	0.00	0.00
Maple	3	100.00	0.00	0.00
Maxima	6	100.00	0.00	0.00
Giac	26	100.00	0.00	0.00
Mupad	33	0.00	100.00	0.00
Sympy	33	90.91	0.00	9.09

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.

Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)
Rubi	0.24
Fricas	0.26
Maple	0.29
Giac	0.33
Maxima	0.58
Sympy	1.46
Mupad	2.06
Mathematica	10.13

Table 1.5: Time performance for each CAS

System	Mean size	Normalized mean	Median size	Normalized median
Mupad	75.86	1.12	22.00	1.11
Giac	132.83	1.14	22.00	1.11
Rubi	155.26	1.00	114.00	1.00
Mathematica	230.05	1.37	152.00	1.17
Maxima	302.78	5.92	170.00	2.18
Maple	345.67	1.87	159.50	1.00
Sympy	501.04	3.16	19.00	1.00
Fricas	887.82	4.04	216.00	2.35

Table 1.6: Leaf size performance for each CAS

1.4 Performance based on number of rules Rubi used

This section shows how each CAS performed based on the number of rules Rubi needed to solve the same integral. One diagram is given for each CAS.

On the y axis is the percentage solved which Rubi itself needed the number of rules given the x axis. These plots show that as more rules are needed then most CAS system percentage of solving decreases which indicates the integral is becoming more complicated to solve.




Figure 1.1: Solving statistics per number of Rubi rules used

1.5 Performance based on number of steps Rubi used

This section shows how each CAS performed based on the number of steps Rubi needed to solve the same integral. Note that the number of steps Rubi needed can be much higher than the number of rules, as the same rule could be used more than once.





Figure 1.2: Solving statistics per number of Rubi steps used

The above diagram show that the precentage of solved intergals decreases for most CAS systems as the number of steps increases. As expected, for integrals that required less steps by Rubi, CAS systems had more success which indicates the integral was not as hard to solve. As Rubi needed more steps to solve the integral, the solved percentage decreased for most CAS systems which indicates the integral is becoming harder to solve.

1.6 Solved integrals histogram based on leaf size of result

The following shows the distribution of solved integrals for each CAS system based on leaf size of the antiderivatives produced by each CAS. It shows that most integrals solved produced leaf size less than about 100 to 150. The bin size used is 40.



1.7 Solved integrals histogram based on CPU time used

The following shows the distribution of solved integrals for each CAS system based on CPU time used in seconds. The bin size used is 0.1 second.




Figure 1.4: Solved integrals histogram based on CPU time used

1.8 Leaf size vs. CPU time used

The following shows the relation between the CPU time used to solve an integral and the leaf size of the antiderivative.

The result for Fricas, Maxima and Giac is shifted more to the right than the other CAS system due to the use of sagemath to call them, which causes an initial slight delay in the timing to start the integration due to overhead of starting a new process each time. This should also be taken into account when looking at the timing of these three CAS systems. Direct calls not using sagemath would result in faster timings, but current implementation uses sagemath as this makes testing much easier to do.




Figure 1.5: Leaf size vs. CPU time. Full range

1.9 list of integrals with no known antiderivative

[{4, 5, 9, 10, 14, 15, 32, 33, 40, 41, 45, 46, 50, 51, 55, 56, 60, 61}](#)

1.10 List of integrals solved by CAS but has no known antiderivative

Rubi [{}](#)

Mathematica [{}](#)

Maple [{}](#)

Maxima [{}](#)

Fricas [{}](#)

Sympy [{}](#)

Giac [{}](#)

Mupad [{}](#)

1.11 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not necessarily mean that the anti-derivative is wrong as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it possible to do further investigation to determine why the result could not be verified.

Rubi [{}](#)

Mathematica [{}](#)

Maple [{}](#)

Maxima Verification phase not currently implemented.

Fricas Verification phase not currently implemented.

Sympy Verification phase not currently implemented.

Giac Verification phase not currently implemented.

Mupad Verification phase not currently implemented.

1.12 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int', int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.13 Verification

A verification phase was applied on the result of integration for **Rubi** and **Mathematica**.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.14 Important notes about some of the results

Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
```

```

x, aa = expr.operator(), expr.operands()
if x is None:
    return 1
else:
    return 1 + sum(tree_size(a) for a in aa)

```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```

try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1

```

Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```

integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)

```

Which gives $\sin(x)^{2/2}$

1.15 Design of the test system

The following diagram gives a high level view of the current test build system.



CHAPTER 2

DETAILED SUMMARY TABLES OF RESULTS

2.1	List of integrals sorted by grade for each CAS	22
2.2	Detailed conclusion table per each integral for all CAS systems	25
2.3	Detailed conclusion table specific for Rubi results	38

2.1 List of integrals sorted by grade for each CAS

Rubi	22
Mma	22
Maple	23
Fricas	23
Maxima	23
Giac	23
Mupad	24
Sympy	24

Rubi

A grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 34, 35, 36, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54, 57, 58, 59 }

B grade { }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Mma

A grade { 1, 2, 3, 7, 8, 13, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 34, 35, 36, 37, 38, 39, 44, 48, 49, 52, 53, 54, 57, 58, 59 }

B grade { 6, 11, 12, 42, 43, 47 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maple

A grade { 8, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

B grade { 1, 2, 3, 6, 7, 11, 12, 13, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54, 57, 58, 59 }

C grade { }

F normal fail { 34, 35, 36 }

F(-1) timeout fail { }

F(-2) exception fail { }

Fricas

A grade { 16, 17, 18, 19, 20, 21, 24, 25, 26, 29, 30, 31, 34, 35, 36 }

B grade { 1, 2, 3, 6, 7, 8, 11, 12, 13, 22, 23, 27, 28, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54, 57, 58, 59 }

C grade { }

F normal fail { }

F(-1) timeout fail { }

F(-2) exception fail { }

Maxima

A grade { 6, 7, 11, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 44, 57, 58 }

B grade { 1, 2, 3, 8, 12, 13, 37, 38, 42, 43, 47, 48, 49, 52, 53 }

C grade { }

F normal fail { 34, 35, 36, 39, 54, 59 }

F(-1) timeout fail { }

F(-2) exception fail { }

Giac

A grade { 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

B grade { 8, 20 }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 34, 35, 36, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54, 57, 58, 59 }

F(-1) timeout fail { }

F(-2) exception fail { }

Mupad

A grade { }

B grade { 8, 16, 17, 18, 22, 23, 24, 27, 28, 29 }

C grade { }

F normal fail { }

F(-1) timeout fail { 1, 2, 3, 6, 7, 11, 12, 13, 19, 20, 21, 25, 26, 30, 31, 34, 35, 36, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54, 57, 58, 59 }

F(-2) exception fail { }

Sympy

A grade { }

B grade { 8, 16, 17, 18, 22, 23, 24, 27, 28, 29 }

C grade { }

F normal fail { 1, 2, 3, 6, 7, 11, 12, 13, 19, 20, 21, 25, 26, 30, 31, 34, 35, 36, 37, 38, 39, 42, 43, 44, 47, 48, 49, 52, 53, 54 }

F(-1) timeout fail { }

F(-2) exception fail { 57, 58, 59 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by the table below. The elapsed time is in seconds. For failed result it is given as **F(-1)** if the failure was due to timeout. It is given as **F(-2)** if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given as **F**.

In this table, the column **N.S.** means **normalized size** and is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To make the table fit the page, the name **Mathematica** was abbreviated to **MMA**.

Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	91	200	170	216	0	0	0
N.S.	1	1.00	1.05	2.30	1.95	2.48	0.00	0.00	0.00
time (sec)	N/A	0.118	0.016	0.296	0.201	0.265	0.000	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	63	63	66	166	138	168	0	0	0
N.S.	1	1.00	1.05	2.63	2.19	2.67	0.00	0.00	0.00
time (sec)	N/A	0.103	0.014	0.251	0.207	0.277	0.000	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	45	45	47	122	98	112	0	0	0
N.S.	1	1.00	1.04	2.71	2.18	2.49	0.00	0.00	0.00
time (sec)	N/A	0.062	0.010	0.268	0.192	0.264	0.000	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	35	12	8	12	12
N.S.	1	1.00	1.20	1.00	3.50	1.20	0.80	1.20	1.20
time (sec)	N/A	0.013	13.895	0.045	0.410	0.260	0.472	0.280	1.879

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	10	10	12	10	46	12	10	12	12
N.S.	1	1.00	1.20	1.00	4.60	1.20	1.00	1.20	1.20
time (sec)	N/A	0.013	23.107	0.052	0.324	0.244	0.446	0.284	1.895

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	87	87	222	198	146	632	0	0	0
N.S.	1	1.00	2.55	2.28	1.68	7.26	0.00	0.00	0.00
time (sec)	N/A	0.129	1.089	0.110	0.258	0.264	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	65	65	117	156	108	453	0	0	0
N.S.	1	1.00	1.80	2.40	1.66	6.97	0.00	0.00	0.00
time (sec)	N/A	0.087	1.577	0.115	0.268	0.263	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	31	31	46	54	115	189	190	98	48
N.S.	1	1.00	1.48	1.74	3.71	6.10	6.13	3.16	1.55
time (sec)	N/A	0.023	0.170	0.112	0.234	0.258	0.678	0.293	1.890

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	69	14	10	14	14
N.S.	1	1.00	1.17	1.00	5.75	1.17	0.83	1.17	1.17
time (sec)	N/A	0.021	25.940	0.045	0.291	0.245	0.520	0.292	1.907

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	91	14	12	14	14
N.S.	1	1.00	1.17	1.00	7.58	1.17	1.00	1.17	1.17
time (sec)	N/A	0.024	25.753	0.049	0.291	0.253	0.487	0.292	1.865

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	179	179	422	375	302	1985	0	0	0
N.S.	1	1.00	2.36	2.09	1.69	11.09	0.00	0.00	0.00
time (sec)	N/A	0.255	3.708	0.171	0.246	0.313	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	114	114	314	246	226	1467	0	0	0
N.S.	1	1.00	2.75	2.16	1.98	12.87	0.00	0.00	0.00
time (sec)	N/A	0.165	3.542	0.163	0.254	0.275	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	82	82	131	164	149	975	0	0	0
N.S.	1	1.00	1.60	2.00	1.82	11.89	0.00	0.00	0.00
time (sec)	N/A	0.094	1.532	0.135	0.237	0.324	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	144	14	10	14	14
N.S.	1	1.00	1.17	1.00	12.00	1.17	0.83	1.17	1.17
time (sec)	N/A	0.023	21.674	0.059	0.309	0.260	0.597	0.309	1.809

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	12	12	14	12	175	14	12	14	14
N.S.	1	1.00	1.17	1.00	14.58	1.17	1.00	1.17	1.17
time (sec)	N/A	0.023	15.902	0.059	0.299	0.242	0.542	0.302	1.816

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	169	169	244	165	183	304	864	188	223
N.S.	1	1.00	1.44	0.98	1.08	1.80	5.11	1.11	1.32
time (sec)	N/A	0.143	0.929	0.385	0.262	0.261	0.763	0.290	2.164

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	122	122	169	103	124	192	522	119	186
N.S.	1	1.00	1.39	0.84	1.02	1.57	4.28	0.98	1.52
time (sec)	N/A	0.090	0.590	0.314	0.246	0.269	0.634	0.278	1.974

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	74	74	81	46	72	101	250	62	76
N.S.	1	1.00	1.09	0.62	0.97	1.36	3.38	0.84	1.03
time (sec)	N/A	0.041	0.707	0.191	0.215	0.255	0.542	0.286	1.945

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	157	157	122	61	48	75	0	48	0
N.S.	1	1.00	0.78	0.39	0.31	0.48	0.00	0.31	0.00
time (sec)	N/A	0.198	0.631	0.342	0.334	0.243	0.000	0.286	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	B	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	159	159	206	91	56	216	0	321	0
N.S.	1	1.00	1.30	0.57	0.35	1.36	0.00	2.02	0.00
time (sec)	N/A	0.175	0.929	0.364	0.474	0.280	0.000	0.313	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	211	211	265	210	68	342	0	175	0
N.S.	1	1.00	1.26	1.00	0.32	1.62	0.00	0.83	0.00
time (sec)	N/A	0.240	1.232	0.400	0.650	0.259	0.000	0.294	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	230	230	420	273	297	571	2193	368	266
N.S.	1	1.00	1.83	1.19	1.29	2.48	9.53	1.60	1.16
time (sec)	N/A	0.195	2.909	0.496	0.501	0.254	1.060	0.298	2.057

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	170	170	207	163	191	359	1353	217	164
N.S.	1	1.00	1.22	0.96	1.12	2.11	7.96	1.28	0.96
time (sec)	N/A	0.141	2.308	0.435	0.406	0.274	0.885	0.293	1.944

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	133	114	74	107	190	700	103	88
N.S.	1	1.00	0.86	0.56	0.80	1.43	5.26	0.77	0.66
time (sec)	N/A	0.111	1.459	0.377	0.321	0.250	0.798	0.275	1.907

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	297	297	199	106	81	137	0	77	0
N.S.	1	1.00	0.67	0.36	0.27	0.46	0.00	0.26	0.00
time (sec)	N/A	0.560	1.127	0.335	0.698	0.263	0.000	0.282	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	420	420	442	164	100	615	0	585	0
N.S.	1	1.00	1.05	0.39	0.24	1.46	0.00	1.39	0.00
time (sec)	N/A	0.615	2.391	0.391	1.322	0.265	0.000	0.329	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	336	336	615	379	406	844	3918	548	374
N.S.	1	1.00	1.83	1.13	1.21	2.51	11.66	1.63	1.11
time (sec)	N/A	0.287	4.566	0.584	1.640	0.277	1.466	0.296	2.167

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	246	246	371	223	254	532	2443	315	234
N.S.	1	1.00	1.51	0.91	1.03	2.16	9.93	1.28	0.95
time (sec)	N/A	0.192	3.166	0.523	1.149	0.259	1.205	0.284	2.141

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	183	183	185	102	138	286	1287	142	127
N.S.	1	1.00	1.01	0.56	0.75	1.56	7.03	0.78	0.69
time (sec)	N/A	0.176	1.756	0.430	0.675	0.262	1.110	0.304	2.012

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	437	437	312	151	114	195	0	104	0
N.S.	1	1.00	0.71	0.35	0.26	0.45	0.00	0.24	0.00
time (sec)	N/A	1.454	1.319	0.374	2.288	0.254	0.000	0.289	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F	A	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	692	692	796	239	140	1162	0	841	0
N.S.	1	1.00	1.15	0.35	0.20	1.68	0.00	1.22	0.00
time (sec)	N/A	1.407	3.767	0.471	5.164	0.262	0.000	0.324	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	122	37	44	22	22
N.S.	1	1.00	1.10	1.00	6.10	1.85	2.20	1.10	1.10
time (sec)	N/A	0.034	41.615	0.066	0.389	0.244	4.418	0.329	2.154

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	68	20	24	20	20
N.S.	1	1.00	1.11	1.00	3.78	1.11	1.33	1.11	1.11
time (sec)	N/A	0.021	20.068	0.056	0.292	0.261	2.784	0.303	2.148

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	88	88	115	0	0	148	0	0	0
N.S.	1	1.00	1.31	0.00	0.00	1.68	0.00	0.00	0.00
time (sec)	N/A	0.089	1.200	0.000	0.000	0.079	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	152	152	162	0	0	248	0	0	0
N.S.	1	1.00	1.07	0.00	0.00	1.63	0.00	0.00	0.00
time (sec)	N/A	0.126	1.734	0.000	0.000	0.086	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	F	F	A	F	F	F(-1)
verified	N/A	Yes	Yes	N/A	TBD	TBD	TBD	TBD	TBD
size	223	223	228	0	0	345	0	0	0
N.S.	1	1.00	1.02	0.00	0.00	1.55	0.00	0.00	0.00
time (sec)	N/A	0.166	1.974	0.000	0.000	0.091	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	133	133	249	766	419	488	0	0	0
N.S.	1	1.00	1.87	5.76	3.15	3.67	0.00	0.00	0.00
time (sec)	N/A	0.194	0.355	0.416	0.258	0.262	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	101	101	149	465	240	303	0	0	0
N.S.	1	1.00	1.48	4.60	2.38	3.00	0.00	0.00	0.00
time (sec)	N/A	0.162	0.213	0.364	0.257	0.259	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	75	75	87	201	0	156	0	0	0
N.S.	1	1.00	1.16	2.68	0.00	2.08	0.00	0.00	0.00
time (sec)	N/A	0.093	0.055	0.186	0.000	0.260	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	78	20	15	20	20
N.S.	1	1.00	1.11	1.00	4.33	1.11	0.83	1.11	1.11
time (sec)	N/A	0.024	5.728	0.058	0.257	0.259	0.960	0.275	1.970

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	18	18	20	18	131	31	17	20	20
N.S.	1	1.00	1.11	1.00	7.28	1.72	0.94	1.11	1.11
time (sec)	N/A	0.023	31.405	0.058	0.271	0.252	3.172	0.356	2.039

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	271	271	843	1425	781	3239	0	0	0
N.S.	1	1.00	3.11	5.26	2.88	11.95	0.00	0.00	0.00
time (sec)	N/A	0.402	4.907	0.524	0.279	0.316	0.000	0.000	0.000

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	209	209	473	825	494	1854	0	0	0
N.S.	1	1.00	2.26	3.95	2.36	8.87	0.00	0.00	0.00
time (sec)	N/A	0.292	6.924	0.461	0.267	0.288	0.000	0.000	0.000

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	127	127	218	318	244	851	0	0	0
N.S.	1	1.00	1.72	2.50	1.92	6.70	0.00	0.00	0.00
time (sec)	N/A	0.134	7.225	0.392	0.260	0.263	0.000	0.000	0.000

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	226	36	17	22	22
N.S.	1	1.00	1.10	1.00	11.30	1.80	0.85	1.10	1.10
time (sec)	N/A	0.042	42.155	0.173	0.400	0.253	1.491	0.325	2.195

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	375	47	19	22	22
N.S.	1	1.00	1.10	1.00	18.75	2.35	0.95	1.10	1.10
time (sec)	N/A	0.040	34.097	0.177	0.450	0.257	2.495	0.464	2.453

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	B	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	556	556	2043	2823	1531	11137	0	0	0
N.S.	1	1.00	3.67	5.08	2.75	20.03	0.00	0.00	0.00
time (sec)	N/A	0.762	8.913	0.776	0.303	0.388	0.000	0.000	0.000

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	401	401	585	1586	997	6356	0	0	0
N.S.	1	1.00	1.46	3.96	2.49	15.85	0.00	0.00	0.00
time (sec)	N/A	0.506	7.466	0.629	0.282	0.337	0.000	0.000	0.000

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	261	261	318	651	528	2907	0	0	0
N.S.	1	1.00	1.22	2.49	2.02	11.14	0.00	0.00	0.00
time (sec)	N/A	0.271	8.176	0.490	0.268	0.296	0.000	0.000	0.000

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	644	52	17	22	22
N.S.	1	1.00	1.10	1.00	32.20	2.60	0.85	1.10	1.10
time (sec)	N/A	0.043	48.149	0.342	0.746	0.265	1.982	0.415	2.069

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	1144	63	19	22	22
N.S.	1	1.00	1.10	1.00	57.20	3.15	0.95	1.10	1.10
time (sec)	N/A	0.039	56.003	0.318	0.952	0.265	3.207	0.796	2.163

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	210	210	247	1158	521	730	0	0	0
N.S.	1	1.00	1.18	5.51	2.48	3.48	0.00	0.00	0.00
time (sec)	N/A	0.257	1.530	0.448	0.323	0.277	0.000	0.000	0.000

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	B	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	156	156	200	735	332	492	0	0	0
N.S.	1	1.00	1.28	4.71	2.13	3.15	0.00	0.00	0.00
time (sec)	N/A	0.219	1.081	0.414	0.290	0.273	0.000	0.000	0.000

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	108	108	152	357	0	300	0	0	0
N.S.	1	1.00	1.41	3.31	0.00	2.78	0.00	0.00	0.00
time (sec)	N/A	0.128	1.041	0.359	0.000	0.277	0.000	0.000	0.000

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	117	27	17	22	22
N.S.	1	1.00	1.10	1.00	5.85	1.35	0.85	1.10	1.10
time (sec)	N/A	0.044	7.377	0.068	0.345	0.243	1.179	0.280	1.907

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	202	51	19	22	22
N.S.	1	1.00	1.10	1.00	10.10	2.55	0.95	1.10	1.10
time (sec)	N/A	0.043	13.540	0.073	0.509	0.249	1.970	0.368	2.114

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	638	638	659	2444	1056	6171	0	0	0
N.S.	1	1.00	1.03	3.83	1.66	9.67	0.00	0.00	0.00
time (sec)	N/A	1.622	7.975	0.581	0.533	0.360	0.000	0.000	0.000

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	A	B	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	475	475	519	1238	751	3702	0	0	0
N.S.	1	1.00	1.09	2.61	1.58	7.79	0.00	0.00	0.00
time (sec)	N/A	1.217	5.060	0.501	0.511	0.316	0.000	0.000	0.000

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	B	F	B	F(-2)	F	F(-1)
verified	N/A	Yes	Yes	Yes	TBD	TBD	TBD	TBD	TBD
size	196	196	244	524	0	1797	0	0	0
N.S.	1	1.00	1.24	2.67	0.00	9.17	0.00	0.00	0.00
time (sec)	N/A	0.218	3.873	0.445	0.000	0.310	0.000	0.000	0.000

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	472	55	19	22	22
N.S.	1	1.00	1.10	1.00	23.60	2.75	0.95	1.10	1.10
time (sec)	N/A	0.042	41.450	0.073	1.101	0.261	1.827	0.311	2.272

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A	N/A
verified	N/A	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD
size	20	20	22	20	789	96	20	22	22
N.S.	1	1.00	1.10	1.00	39.45	4.80	1.00	1.10	1.10
time (sec)	N/A	0.039	39.171	0.071	2.157	0.274	3.323	0.511	2.714

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi only. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is also given. The larger this ratio is, the harder the integral is to solve. In this test file, problem number [11] had the largest ratio of [.83330000000000041]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	6	1.00	10	0.600
2	A	5	5	1.00	10	0.500
3	A	4	4	1.00	8	0.500
4	N/A	0	0	1.00	10	0.000
5	N/A	0	0	1.00	10	0.000
6	A	7	7	1.00	12	0.583
7	A	6	6	1.00	12	0.500
8	A	3	3	1.00	10	0.300
9	N/A	0	0	1.00	12	0.000
10	N/A	0	0	1.00	12	0.000
11	A	13	10	1.00	12	0.833
12	A	9	8	1.00	12	0.667
13	A	7	7	1.00	10	0.700
14	N/A	0	0	1.00	12	0.000
15	N/A	0	0	1.00	12	0.000
16	A	5	3	1.00	20	0.150
17	A	4	3	1.00	20	0.150
18	A	3	3	1.00	18	0.167
19	A	7	4	1.00	20	0.200
20	A	7	4	1.00	20	0.200
21	A	8	5	1.00	20	0.250
22	A	10	3	1.00	20	0.150
23	A	8	3	1.00	20	0.150

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
24	A	7	3	1.00	18	0.167
25	A	21	5	1.00	20	0.250
26	A	24	7	1.00	20	0.350
27	A	14	3	1.00	20	0.150
28	A	11	3	1.00	20	0.150
29	A	11	3	1.00	18	0.167
30	A	53	7	1.00	20	0.350
31	A	60	9	1.00	20	0.450
32	N/A	0	0	1.00	20	0.000
33	N/A	0	0	1.00	18	0.000
34	A	2	2	1.00	20	0.100
35	A	4	2	1.00	20	0.100
36	A	5	2	1.00	20	0.100
37	A	8	7	1.00	18	0.389
38	A	7	6	1.00	18	0.333
39	A	6	5	1.00	16	0.312
40	N/A	0	0	1.00	18	0.000
41	N/A	0	0	1.00	18	0.000
42	A	15	9	1.00	20	0.450
43	A	13	10	1.00	20	0.500
44	A	9	7	1.00	18	0.389
45	N/A	0	0	1.00	20	0.000
46	N/A	0	0	1.00	20	0.000
47	A	28	11	1.00	20	0.550
48	A	22	11	1.00	20	0.550
49	A	16	9	1.00	18	0.500
50	N/A	0	0	1.00	20	0.000
51	N/A	0	0	1.00	20	0.000
52	A	6	6	1.00	20	0.300
53	A	5	5	1.00	20	0.250
54	A	4	4	1.00	18	0.222
55	N/A	0	0	1.00	20	0.000
56	N/A	0	0	1.00	20	0.000
57	A	28	10	1.00	20	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
58	A	24	11	1.00	20	0.550
59	A	5	5	1.00	18	0.278
60	N/A	0	0	1.00	20	0.000
61	N/A	0	0	1.00	20	0.000

CHAPTER 3

LISTING OF INTEGRALS

3.1	$\int x^3 \coth(a + bx) dx$	43
3.2	$\int x^2 \coth(a + bx) dx$	48
3.3	$\int x \coth(a + bx) dx$	53
3.4	$\int \frac{\coth(a+bx)}{x} dx$	57
3.5	$\int \frac{\coth(a+bx)}{x^2} dx$	60
3.6	$\int x^3 \coth^2(a + bx) dx$	63
3.7	$\int x^2 \coth^2(a + bx) dx$	69
3.8	$\int x \coth^2(a + bx) dx$	74
3.9	$\int \frac{\coth^2(a+bx)}{x} dx$	78
3.10	$\int \frac{\coth^2(a+bx)}{x^2} dx$	81
3.11	$\int x^3 \coth^3(a + bx) dx$	84
3.12	$\int x^2 \coth^3(a + bx) dx$	92
3.13	$\int x \coth^3(a + bx) dx$	99
3.14	$\int \frac{\coth^3(a+bx)}{x} dx$	105
3.15	$\int \frac{\coth^3(a+bx)}{x^2} dx$	108
3.16	$\int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx$	111
3.17	$\int \frac{(c+dx)^2}{a+a \coth(e+fx)} dx$	117
3.18	$\int \frac{c+dx}{a+a \coth(e+fx)} dx$	123
3.19	$\int \frac{1}{(c+dx)(a+a \coth(e+fx))} dx$	128
3.20	$\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))} dx$	133
3.21	$\int \frac{1}{(c+dx)^3(a+a \coth(e+fx))} dx$	139
3.22	$\int \frac{(c+dx)^3}{(a+a \coth(e+fx))^2} dx$	145
3.23	$\int \frac{(c+dx)^2}{(a+a \coth(e+fx))^2} dx$	153
3.24	$\int \frac{c+dx}{(a+a \coth(e+fx))^2} dx$	159

3.25	$\int \frac{1}{(c+dx)(a+a \coth(e+fx))^2} dx$	165
3.26	$\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^2} dx$	171
3.27	$\int \frac{(c+dx)^3}{(a+a \coth(e+fx))^3} dx$	179
3.28	$\int \frac{(c+dx)^2}{(a+a \coth(e+fx))^3} dx$	189
3.29	$\int \frac{c+dx}{(a+a \coth(e+fx))^3} dx$	197
3.30	$\int \frac{1}{(c+dx)(a+a \coth(e+fx))^3} dx$	204
3.31	$\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^3} dx$	212
3.32	$\int (c+dx)^m (a+a \coth(e+fx))^2 dx$	227
3.33	$\int (c+dx)^m (a+a \coth(e+fx)) dx$	230
3.34	$\int \frac{(c+dx)^m}{a+a \coth(e+fx)} dx$	233
3.35	$\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^2} dx$	237
3.36	$\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^3} dx$	241
3.37	$\int (c+dx)^3 (a+b \coth(e+fx)) dx$	246
3.38	$\int (c+dx)^2 (a+b \coth(e+fx)) dx$	253
3.39	$\int (c+dx) (a+b \coth(e+fx)) dx$	259
3.40	$\int \frac{a+b \coth(e+fx)}{c+dx} dx$	264
3.41	$\int \frac{a+b \coth(e+fx)}{(c+dx)^2} dx$	267
3.42	$\int (c+dx)^3 (a+b \coth(e+fx))^2 dx$	270
3.43	$\int (c+dx)^2 (a+b \coth(e+fx))^2 dx$	281
3.44	$\int (c+dx) (a+b \coth(e+fx))^2 dx$	291
3.45	$\int \frac{(a+b \coth(e+fx))^2}{c+dx} dx$	298
3.46	$\int \frac{(a+b \coth(e+fx))^2}{(c+dx)^2} dx$	301
3.47	$\int (c+dx)^3 (a+b \coth(e+fx))^3 dx$	305
3.48	$\int (c+dx)^2 (a+b \coth(e+fx))^3 dx$	320
3.49	$\int (c+dx) (a+b \coth(e+fx))^3 dx$	331
3.50	$\int \frac{(a+b \coth(e+fx))^3}{c+dx} dx$	340
3.51	$\int \frac{(a+b \coth(e+fx))^3}{(c+dx)^2} dx$	344
3.52	$\int \frac{(c+dx)^3}{a+b \coth(e+fx)} dx$	348
3.53	$\int \frac{(c+dx)^2}{a+b \coth(e+fx)} dx$	356
3.54	$\int \frac{c+dx}{a+b \coth(e+fx)} dx$	363
3.55	$\int \frac{1}{(c+dx)(a+b \coth(e+fx))} dx$	368
3.56	$\int \frac{1}{(c+dx)^2(a+b \coth(e+fx))} dx$	371
3.57	$\int \frac{(c+dx)^3}{(a+b \coth(e+fx))^2} dx$	375
3.58	$\int \frac{(c+dx)^2}{(a+b \coth(e+fx))^2} dx$	390
3.59	$\int \frac{c+dx}{(a+b \coth(e+fx))^2} dx$	402
3.60	$\int \frac{1}{(c+dx)(a+b \coth(e+fx))^2} dx$	409
3.61	$\int \frac{1}{(c+dx)^2(a+b \coth(e+fx))^2} dx$	413

3.1 $\int x^3 \coth(a + bx) dx$

Optimal result	43
Rubi [A] (verified)	43
Mathematica [A] (verified)	45
Maple [B] (verified)	45
Fricas [B] (verification not implemented)	46
Sympy [F]	46
Maxima [B] (verification not implemented)	47
Giac [F]	47
Mupad [F(-1)]	47

Optimal result

Integrand size = 10, antiderivative size = 87

$$\begin{aligned} \int x^3 \coth(a + bx) dx = & -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\ & - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \operatorname{PolyLog}(4, e^{2(a+bx)})}{4b^4} \end{aligned}$$

[Out] $-1/4*x^4+x^3*ln(1-exp(2*b*x+2*a))/b+3/2*x^2*polylog(2,exp(2*b*x+2*a))/b^2-3/2*x*polylog(3,exp(2*b*x+2*a))/b^3+3/4*polylog(4,exp(2*b*x+2*a))/b^4$

Rubi [A] (verified)

Time = 0.12 (sec), antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.600$, Rules used = {3797, 2221, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int x^3 \coth(a + bx) dx = & \frac{3 \operatorname{PolyLog}(4, e^{2(a+bx)})}{4b^4} - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} \\ & + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} - \frac{x^4}{4} \end{aligned}$$

[In] `Int[x^3*Coth[a + b*x], x]`

[Out] $-1/4*x^4 + (x^3*Log[1 - E^(2*(a + b*x))])/b + (3*x^2*PolyLog[2, E^(2*(a + b*x))])/(2*b^2) - (3*x*PolyLog[3, E^(2*(a + b*x))])/(2*b^3) + (3*PolyLog[4, E^(2*(a + b*x))])/(4*b^4)$

Rule 2221

`Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_), x_Symbol] :> Simp`

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_)*(a_.) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_.) + (b_)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_))^p]/((d_.) + (e_)*(x_), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_)*(a_.) + (b_)*(x_)))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = -\frac{x^4}{4} - 2 \int \frac{e^{2(a+bx)} x^3}{1 - e^{2(a+bx)}} dx$$

$$\begin{aligned}
&= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} - \frac{3 \int x^2 \log(1 - e^{2(a+bx)}) dx}{b} \\
&= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} - \frac{3 \int x \operatorname{PolyLog}(2, e^{2(a+bx)}) dx}{b^2} \\
&= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\
&\quad - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \int \operatorname{PolyLog}(3, e^{2(a+bx)}) dx}{2b^3} \\
&= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\
&\quad - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}(3,x)}{x} dx, x, e^{2(a+bx)}\right)}{4b^4} \\
&= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\
&\quad - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \operatorname{PolyLog}(4, e^{2(a+bx)})}{4b^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.02 (sec) , antiderivative size = 91, normalized size of antiderivative = 1.05

$$\begin{aligned}
\int x^3 \coth(a + bx) dx &= -\frac{x^4}{4} + \frac{x^3 \log(1 - e^{2a+2bx})}{b} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2a+2bx})}{2b^2} \\
&\quad - \frac{3x \operatorname{PolyLog}(3, e^{2a+2bx})}{2b^3} + \frac{3 \operatorname{PolyLog}(4, e^{2a+2bx})}{4b^4}
\end{aligned}$$

[In] `Integrate[x^3*Coth[a + b*x], x]`

[Out] $-1/4*x^4 + (x^3 \operatorname{Log}[1 - E^{(2*a + 2*b*x)}])/b + (3*x^2 \operatorname{PolyLog}[2, E^{(2*a + 2*b*x)}])/(2*b^2) - (3*x \operatorname{PolyLog}[3, E^{(2*a + 2*b*x)}])/(2*b^3) + (3 \operatorname{PolyLog}[4, E^{(2*a + 2*b*x)}])/(4*b^4)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 199 vs. $2(79) = 158$.

Time = 0.30 (sec) , antiderivative size = 200, normalized size of antiderivative = 2.30

method	result
risch	$-\frac{x^4}{4} - \frac{2a^3x}{b^3} - \frac{3a^4}{2b^4} + \frac{3 \operatorname{polylog}(2, e^{bx+a})x^2}{b^2} - \frac{6 \operatorname{polylog}(3, e^{bx+a})x}{b^3} + \frac{\ln(e^{bx+a}+1)x^3}{b} + \frac{3 \operatorname{polylog}(2, -e^{bx+a})x^2}{b^2} - \frac{6 \operatorname{polylog}(4, -e^{bx+a})}{b^4}$

[In] `int(x^3*coth(b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -\frac{1}{4}x^4 - \frac{2}{b^3}a^3x^3 - \frac{3}{2}b^4a^4 + \frac{3}{b^2}\text{polylog}(2, \exp(b*x+a))x^2 - \frac{6}{b^3}\text{polylog}(3, \exp(b*x+a))x + \frac{1}{b}\ln(\exp(b*x+a)+1)x^3 + \frac{3}{b^2}\text{polylog}(2, -\exp(b*x+a))x^2 \\ & - \frac{6}{b^3}\text{polylog}(3, -\exp(b*x+a))x + \frac{1}{b}\ln(1-\exp(b*x+a))x^3 - \frac{1}{b^4}a^3\ln(\exp(b*x+a)-1) + \frac{2}{b^4}a^3\ln(\exp(b*x+a)) + \frac{1}{b^4}\ln(1-\exp(b*x+a))a^3 + \frac{6}{b^4}\text{polylog}(4, -\exp(b*x+a)) + \frac{6}{b^4}\text{polylog}(4, \exp(b*x+a)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 216 vs. $2(78) = 156$.

Time = 0.27 (sec) , antiderivative size = 216, normalized size of antiderivative = 2.48

$$\int x^3 \coth(a + bx) dx = \frac{b^4 x^4 - 4 b^3 x^3 \log(\cosh(bx + a) + \sinh(bx + a) + 1) - 12 b^2 x^2 \text{Li}_2(\cosh(bx + a) + \sinh(bx + a)) - 12 b^2}{b^4}$$

[In] `integrate(x^3*coth(b*x+a),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -\frac{1}{4}(b^4 x^4 - 4 b^3 x^3 \log(\cosh(b*x + a) + \sinh(b*x + a) + 1) - 12 b^2 x^2 \text{dilog}(\cosh(b*x + a) + \sinh(b*x + a)) - 12 b^2 x^2 \text{dilog}(-\cosh(b*x + a) - \sinh(b*x + a)) + 4 a^3 \log(\cosh(b*x + a) + \sinh(b*x + a) - 1) + 24 b*x \text{polylog}(3, \cosh(b*x + a) + \sinh(b*x + a)) + 24 b*x \text{polylog}(3, -\cosh(b*x + a) - \sinh(b*x + a)) - 4(b^3 x^3 + a^3) \log(-\cosh(b*x + a) - \sinh(b*x + a) + 1) - 24 \text{polylog}(4, \cosh(b*x + a) + \sinh(b*x + a)) - 24 \text{polylog}(4, -\cosh(b*x + a) - \sinh(b*x + a))) / b^4 \end{aligned}$$

Sympy [F]

$$\int x^3 \coth(a + bx) dx = \int x^3 \coth(a + bx) dx$$

[In] `integrate(x**3*coth(b*x+a),x)`

[Out] `Integral(x**3*coth(a + b*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 170 vs. $2(78) = 156$.

Time = 0.20 (sec) , antiderivative size = 170, normalized size of antiderivative = 1.95

$$\int x^3 \coth(a + bx) dx = \frac{1}{4} x^4 \coth(bx + a) - \frac{1}{2} \left(\frac{x^4}{be^{(2bx+2a)} - b} + \frac{x^4}{b} - \frac{2(b^3 x^3 \log(e^{(bx+a)} + 1) + 3b^2 x^2 \text{Li}_2(-e^{(bx+a)}) - 6bx \text{Li}_3(-e^{(bx+a)}) + 6\text{Li}_4(-e^{(bx+a)}))}{b^5} \right)$$

```
[In] integrate(x^3*coth(b*x+a),x, algorithm="maxima")
[Out] 1/4*x^4*coth(b*x + a) - 1/2*(x^4/(b*e^(2*b*x + 2*a) - b) + x^4/b - 2*(b^3*x^3*log(e^(b*x + a) + 1) + 3*b^2*x^2*dilog(-e^(b*x + a)) - 6*b*x*polylog(3, -e^(b*x + a)) + 6*polylog(4, -e^(b*x + a)))/b^5 - 2*(b^3*x^3*log(-e^(b*x + a) + 1) + 3*b^2*x^2*dilog(e^(b*x + a)) - 6*b*x*polylog(3, e^(b*x + a)) + 6*polylog(4, e^(b*x + a)))/b^5)*b
```

Giac [F]

$$\int x^3 \coth(a + bx) dx = \int x^3 \coth(bx + a) dx$$

```
[In] integrate(x^3*coth(b*x+a),x, algorithm="giac")
[Out] integrate(x^3*coth(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \coth(a + bx) dx = \int x^3 \coth(a + b x) dx$$

```
[In] int(x^3*coth(a + b*x),x)
[Out] int(x^3*coth(a + b*x), x)
```

3.2 $\int x^2 \coth(a + bx) dx$

Optimal result	48
Rubi [A] (verified)	48
Mathematica [A] (verified)	50
Maple [B] (verified)	50
Fricas [B] (verification not implemented)	50
Sympy [F]	51
Maxima [B] (verification not implemented)	51
Giac [F]	51
Mupad [F(-1)]	52

Optimal result

Integrand size = 10, antiderivative size = 63

$$\begin{aligned} \int x^2 \coth(a + bx) dx = & -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\ & + \frac{x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} \end{aligned}$$

[Out] $-1/3*x^3+x^2*\ln(1-\exp(2*b*x+2*a))/b+x*\operatorname{polylog}(2,\exp(2*b*x+2*a))/b^2-1/2*\operatorname{polylog}(3,\exp(2*b*x+2*a))/b^3$

Rubi [A] (verified)

Time = 0.10 (sec), antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3797, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int x^2 \coth(a + bx) dx = & -\frac{\operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^2} \\ & + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} - \frac{x^3}{3} \end{aligned}$$

[In] Int[x^2*Coth[a + b*x],x]

[Out] $-1/3*x^3 + (x^2*\operatorname{Log}[1 - E^(2*(a + b*x))])/b + (x*\operatorname{PolyLog}[2, E^(2*(a + b*x))])/b^2 - \operatorname{PolyLog}[3, E^(2*(a + b*x))]/(2*b^3)$

Rule 2221

Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.)) / ((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simpl

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*((F_)^((c_)*(a_.) + (b_)*(x_))))^(n_.)]*((f_.) + (g_.)* (x_))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^m*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_))^p]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x^3}{3} - 2 \int \frac{e^{2(a+bx)} x^2}{1 - e^{2(a+bx)}} dx \\
&= -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} - \frac{2 \int x \log(1 - e^{2(a+bx)}) dx}{b} \\
&= -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} + \frac{x \text{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\int \text{PolyLog}(2, e^{2(a+bx)}) dx}{b^2} \\
&= -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} + \frac{x \text{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\text{Subst}\left(\int \frac{\text{PolyLog}(2, x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^3}
\end{aligned}$$

$$= -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} + \frac{x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3}$$

Mathematica [A] (verified)

Time = 0.01 (sec) , antiderivative size = 66, normalized size of antiderivative = 1.05

$$\int x^2 \coth(a + bx) dx = -\frac{x^3}{3} + \frac{x^2 \log(1 - e^{2a+2bx})}{b} + \frac{x \operatorname{PolyLog}(2, e^{2a+2bx})}{b^2} - \frac{\operatorname{PolyLog}(3, e^{2a+2bx})}{2b^3}$$

[In] `Integrate[x^2*Coth[a + b*x], x]`

[Out] $-1/3*x^3 + (x^2*\operatorname{Log}[1 - E^(2*a + 2*b*x)])/b + (x*\operatorname{PolyLog}[2, E^(2*a + 2*b*x)])/b^2 - \operatorname{PolyLog}[3, E^(2*a + 2*b*x)]/(2*b^3)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 165 vs. $2(59) = 118$.

Time = 0.25 (sec) , antiderivative size = 166, normalized size of antiderivative = 2.63

method	result
risch	$-\frac{x^3}{3} + \frac{4a^3}{3b^3} - \frac{\ln(1-e^{bx+a})a^2}{b^3} + \frac{a^2 \ln(e^{bx+a}-1)}{b^3} - \frac{2a^2 \ln(e^{bx+a})}{b^3} - \frac{2 \operatorname{polylog}(3, -e^{bx+a})}{b^3} - \frac{2 \operatorname{polylog}(3, e^{bx+a})}{b^3} + \frac{2a^2 x}{b^2} - \frac{b^3 x^3 - 3b^2 x^2 \log(\cosh(bx+a) + \sinh(bx+a) + 1) - 6bx \operatorname{Li}_2(\cosh(bx+a) + \sinh(bx+a)) - 6bx \operatorname{Li}_2(-e^{bx+a})}{b^3}$

[In] `int(x^2*coth(b*x+a), x, method=_RETURNVERBOSE)`

[Out] $-1/3*x^3+4/3/b^3*a^3-1/b^3*3*ln(1-exp(b*x+a))*a^2+1/b^3*a^2*ln(exp(b*x+a)-1)-2/b^3*a^2*ln(exp(b*x+a))-2/b^3*polylog(3, -exp(b*x+a))-2/b^3*polylog(3, exp(b*x+a))+2/b^2*a^2*x+1/b*ln(exp(b*x+a)+1)*x^2+2/b^2*polylog(2, -exp(b*x+a))*x+1/b*ln(1-exp(b*x+a))*x^2+2/b^2*polylog(2, exp(b*x+a))*x$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 168 vs. $2(58) = 116$.

Time = 0.28 (sec) , antiderivative size = 168, normalized size of antiderivative = 2.67

$$\int x^2 \coth(a + bx) dx = \frac{-b^3 x^3 - 3 b^2 x^2 \log(\cosh(bx+a) + \sinh(bx+a) + 1) - 6bx \operatorname{Li}_2(\cosh(bx+a) + \sinh(bx+a)) - 6bx \operatorname{Li}_2(-e^{bx+a})}{b^3}$$

[In] `integrate(x^2*coth(b*x+a), x, algorithm="fricas")`

```
[Out] -1/3*(b^3*x^3 - 3*b^2*x^2*log(cosh(b*x + a) + sinh(b*x + a) + 1) - 6*b*x*dilog(cosh(b*x + a) + sinh(b*x + a)) - 6*b*x*dilog(-cosh(b*x + a) - sinh(b*x + a)) - 3*a^2*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 3*(b^2*x^2 - a^2)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) + 6*polylog(3, cosh(b*x + a) + sinh(b*x + a)) + 6*polylog(3, -cosh(b*x + a) - sinh(b*x + a)))/b^3
```

Sympy [F]

$$\int x^2 \coth(a + bx) dx = \int x^2 \coth(bx + a) dx$$

```
[In] integrate(x**2*coth(b*x+a),x)
```

```
[Out] Integral(x**2*coth(a + b*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 138 vs. $2(58) = 116$.

Time = 0.21 (sec) , antiderivative size = 138, normalized size of antiderivative = 2.19

$$\begin{aligned} \int x^2 \coth(a + bx) dx &= \frac{1}{3} x^3 \coth(bx + a) \\ &- \frac{1}{3} \left(\frac{2x^3}{be^{(2bx+2a)} - b} + \frac{2x^3}{b} - \frac{3(b^2x^2 \log(e^{(bx+a)} + 1) + 2bx\text{Li}_2(-e^{(bx+a)}) - 2\text{Li}_3(-e^{(bx+a)}))}{b^4} \right. \\ &\quad \left. - \frac{3(b^2x^2 \log(e^{(bx+a)} + 1) + 2bx\text{Li}_2(-e^{(bx+a)}) - 2\text{Li}_3(-e^{(bx+a)}))}{b^4} \right) \end{aligned}$$

```
[In] integrate(x^2*coth(b*x+a),x, algorithm="maxima")
```

```
[Out] 1/3*x^3*coth(b*x + a) - 1/3*(2*x^3/(b*e^(2*b*x + 2*a) - b) + 2*x^3/b - 3*(b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a)) - 2*polylog(3, -e^(b*x + a))/b^4 - 3*(b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a))/b^4)*b
```

Giac [F]

$$\int x^2 \coth(a + bx) dx = \int x^2 \coth(bx + a) dx$$

```
[In] integrate(x^2*coth(b*x+a),x, algorithm="giac")
```

```
[Out] integrate(x^2*coth(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x^2 \coth(a + bx) dx = \int x^2 \coth(a + b x) dx$$

[In] int(x^2*coth(a + b*x),x)

[Out] int(x^2*coth(a + b*x), x)

3.3 $\int x \coth(a + bx) dx$

Optimal result	53
Rubi [A] (verified)	53
Mathematica [A] (verified)	54
Maple [B] (verified)	55
Fricas [B] (verification not implemented)	55
Sympy [F]	55
Maxima [B] (verification not implemented)	56
Giac [F]	56
Mupad [F(-1)]	56

Optimal result

Integrand size = 8, antiderivative size = 45

$$\int x \coth(a + bx) dx = -\frac{x^2}{2} + \frac{x \log(1 - e^{2(a+bx)})}{b} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2}$$

[Out] $-1/2*x^2+x*ln(1-exp(2*b*x+2*a))/b+1/2*polylog(2,exp(2*b*x+2*a))/b^2$

Rubi [A] (verified)

Time = 0.06 (sec), antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3797, 2221, 2317, 2438}

$$\int x \coth(a + bx) dx = \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2} + \frac{x \log(1 - e^{2(a+bx)})}{b} - \frac{x^2}{2}$$

[In] Int[x*Coth[a + b*x], x]

[Out] $-1/2*x^2 + (x*Log[1 - E^(2*(a + b*x))])/b + PolyLog[2, E^(2*(a + b*x))]/(2*b^2)$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^((m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.))))^(n_.)), x_Symbol] := Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - D[st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*(F_)^((e_.)*(c_.) + (d_)*(x_)))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*(d_ + (e_)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3797

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= -\frac{x^2}{2} - 2 \int \frac{e^{2(a+bx)} x}{1 - e^{2(a+bx)}} dx \\ &= -\frac{x^2}{2} + \frac{x \log(1 - e^{2(a+bx)})}{b} - \frac{\int \log(1 - e^{2(a+bx)}) dx}{b} \\ &= -\frac{x^2}{2} + \frac{x \log(1 - e^{2(a+bx)})}{b} - \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^2} \\ &= -\frac{x^2}{2} + \frac{x \log(1 - e^{2(a+bx)})}{b} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.01 (sec), antiderivative size = 47, normalized size of antiderivative = 1.04

$$\int x \coth(a + bx) dx = -\frac{x^2}{2} + \frac{x \log(1 - e^{2a+2bx})}{b} + \frac{\text{PolyLog}(2, e^{2a+2bx})}{2b^2}$$

[In] `Integrate[x*Coth[a + b*x], x]`

[Out] $-1/2*x^2 + (x*\text{Log}[1 - E^{(2*a + 2*b*x)})]/b + \text{PolyLog}[2, E^{(2*a + 2*b*x)}]/(2*b^2)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 121 vs. $2(41) = 82$.

Time = 0.27 (sec) , antiderivative size = 122, normalized size of antiderivative = 2.71

method	result
risch	$-\frac{x^2}{2} - \frac{2ax}{b} - \frac{a^2}{b^2} + \frac{\ln(e^{bx+a}+1)x}{b} + \frac{\text{polylog}(2, -e^{bx+a})}{b^2} + \frac{\ln(1-e^{bx+a})x}{b} + \frac{\ln(1-e^{bx+a})a}{b^2} + \frac{\text{polylog}(2, e^{bx+a})}{b^2} - \frac{a \ln(e^{bx+a})}{b}$

[In] `int(x*coth(b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -\frac{1}{2}x^2 - 2/b*a*x - a^2/b^2 + \frac{\ln(\exp(b*x+a)+1)*x+1/b^2*\text{polylog}(2,-\exp(b*x+a))}{b} \\ & + 1/b*\ln(1-\exp(b*x+a))*x+1/b^2*\ln(1-\exp(b*x+a))*a+1/b^2*\text{polylog}(2,\exp(b*x+a)) \\ & - 1/b^2*a*\ln(\exp(b*x+a)-1)+2/b^2*a*\ln(\exp(b*x+a)) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(40) = 80$.

Time = 0.26 (sec) , antiderivative size = 112, normalized size of antiderivative = 2.49

$$\int x \coth(a + bx) dx = \frac{-b^2 x^2 - 2 b x \log(\cosh(bx + a) + \sinh(bx + a) + 1) + 2 a \log(\cosh(bx + a) + \sinh(bx + a) - 1) - 2 (bx + a) \operatorname{dilog}(\cosh(bx + a) + \sinh(bx + a)) - 2 \operatorname{dilog}(-\cosh(bx + a) - \sinh(bx + a))}{b^2}$$

[In] `integrate(x*coth(b*x+a),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -\frac{1}{2}*(b^2*x^2 - 2*b*x*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 2*a*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 2*(b*x + a)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) - 2*dilog(cosh(b*x + a) + sinh(b*x + a)) - 2*dilog(-cosh(b*x + a) - sinh(b*x + a)))/b^2 \end{aligned}$$

Sympy [F]

$$\int x \coth(a + bx) dx = \int x \coth(a + bx) dx$$

[In] `integrate(x*coth(b*x+a),x)`

[Out] `Integral(x*coth(a + b*x), x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(40) = 80$.

Time = 0.19 (sec), antiderivative size = 98, normalized size of antiderivative = 2.18

$$\int x \coth(a + bx) dx = \frac{1}{2} x^2 \coth(bx + a) - b \left(\frac{x^2}{be^{(2bx+2a)} - b} + \frac{x^2}{b} - \frac{bx \log(e^{(bx+a)} + 1) + \text{Li}_2(-e^{(bx+a)})}{b^3} - \frac{bx \log(-e^{(bx+a)} + 1) + \text{Li}_2(e^{(bx+a)})}{b^3} \right)$$

```
[In] integrate(x*coth(b*x+a),x, algorithm="maxima")
[Out] 1/2*x^2*coth(b*x + a) - b*(x^2/(b*e^(2*b*x + 2*a) - b) + x^2/b - (b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))/b^3 - (b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))/b^3)
```

Giac [F]

$$\int x \coth(a + bx) dx = \int x \coth(bx + a) dx$$

```
[In] integrate(x*coth(b*x+a),x, algorithm="giac")
[Out] integrate(x*coth(b*x + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int x \coth(a + bx) dx = \int x \coth(a + b x) dx$$

```
[In] int(x*coth(a + b*x),x)
[Out] int(x*coth(a + b*x), x)
```

3.4 $\int \frac{\coth(a+bx)}{x} dx$

Optimal result	57
Rubi [N/A]	57
Mathematica [N/A]	58
Maple [N/A] (verified)	58
Fricas [N/A]	58
Sympy [N/A]	58
Maxima [N/A]	59
Giac [N/A]	59
Mupad [N/A]	59

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\coth(a + bx)}{x} dx = \text{Int}\left(\frac{\coth(a + bx)}{x}, x\right)$$

[Out] Unintegrable($\coth(b*x+a)/x$, x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(a + bx)}{x} dx$$

[In] Int[Coth[a + b*x]/x, x]

[Out] Defer[Int][Coth[a + b*x]/x, x]

Rubi steps

$$\text{integral} = \int \frac{\coth(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 13.89 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(a + bx)}{x} dx$$

[In] `Integrate[Coth[a + b*x]/x, x]`

[Out] `Integrate[Coth[a + b*x]/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)}{x} dx$$

[In] `int(coth(b*x+a)/x,x)`

[Out] `int(coth(b*x+a)/x,x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(bx + a)}{x} dx$$

[In] `integrate(coth(b*x+a)/x,x, algorithm="fricas")`

[Out] `integral(coth(b*x + a)/x, x)`

Sympy [N/A]

Not integrable

Time = 0.47 (sec) , antiderivative size = 8, normalized size of antiderivative = 0.80

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(a + bx)}{x} dx$$

[In] `integrate(coth(b*x+a)/x,x)`

[Out] `Integral(coth(a + b*x)/x, x)`

Maxima [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 35, normalized size of antiderivative = 3.50

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(bx + a)}{x} dx$$

[In] `integrate(coth(b*x+a)/x,x, algorithm="maxima")`

[Out] `-integrate(1/(x*e^(b*x + a) + x), x) + integrate(1/(x*e^(b*x + a) - x), x) + log(x)`

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(bx + a)}{x} dx$$

[In] `integrate(coth(b*x+a)/x,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)/x, x)`

Mupad [N/A]

Not integrable

Time = 1.88 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x} dx = \int \frac{\coth(a + b x)}{x} dx$$

[In] `int(coth(a + b*x)/x,x)`

[Out] `int(coth(a + b*x)/x, x)`

3.5 $\int \frac{\coth(a+bx)}{x^2} dx$

Optimal result	60
Rubi [N/A]	60
Mathematica [N/A]	61
Maple [N/A] (verified)	61
Fricas [N/A]	61
Sympy [N/A]	61
Maxima [N/A]	62
Giac [N/A]	62
Mupad [N/A]	62

Optimal result

Integrand size = 10, antiderivative size = 10

$$\int \frac{\coth(a + bx)}{x^2} dx = \text{Int}\left(\frac{\coth(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable(coth(b*x+a)/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.01 (sec), antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(a + bx)}{x^2} dx$$

[In] Int[Coth[a + b*x]/x^2, x]

[Out] Defer[Int][Coth[a + b*x]/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\coth(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 23.11 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(a + bx)}{x^2} dx$$

[In] Integrate[Coth[a + b*x]/x^2, x]

[Out] Integrate[Coth[a + b*x]/x^2, x]

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)}{x^2} dx$$

[In] int(coth(b*x+a)/x^2, x)

[Out] int(coth(b*x+a)/x^2, x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)}{x^2} dx$$

[In] integrate(coth(b*x+a)/x^2, x, algorithm="fricas")

[Out] integral(coth(b*x + a)/x^2, x)

Sympy [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 10, normalized size of antiderivative = 1.00

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(a + bx)}{x^2} dx$$

[In] integrate(coth(b*x+a)/x**2, x)

[Out] Integral(coth(a + b*x)/x**2, x)

Maxima [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 46, normalized size of antiderivative = 4.60

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)}{x^2} dx$$

[In] `integrate(coth(b*x+a)/x^2,x, algorithm="maxima")`

[Out] `-1/x - integrate(1/(x^2*e^(b*x + a) + x^2), x) + integrate(1/(x^2*e^(b*x + a) - x^2), x)`

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)}{x^2} dx$$

[In] `integrate(coth(b*x+a)/x^2,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)/x^2, x)`

Mupad [N/A]

Not integrable

Time = 1.89 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.20

$$\int \frac{\coth(a + bx)}{x^2} dx = \int \frac{\coth(a + b x)}{x^2} dx$$

[In] `int(coth(a + b*x)/x^2,x)`

[Out] `int(coth(a + b*x)/x^2, x)`

3.6 $\int x^3 \coth^2(a + bx) dx$

Optimal result	63
Rubi [A] (verified)	63
Mathematica [B] (verified)	65
Maple [B] (verified)	66
Fricas [B] (verification not implemented)	66
Sympy [F]	67
Maxima [A] (verification not implemented)	67
Giac [F]	67
Mupad [F(-1)]	68

Optimal result

Integrand size = 12, antiderivative size = 87

$$\begin{aligned} \int x^3 \coth^2(a + bx) dx = & -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} \\ & + \frac{3x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^3} - \frac{3 \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^4} \end{aligned}$$

[Out] $-x^3/b + 1/4*x^4 - x^3*\coth(b*x+a)/b + 3*x^2*\ln(1-exp(2*b*x+2*a))/b^2 + 3*x*polylog(2, exp(2*b*x+2*a))/b^3 - 3/2*polylog(3, exp(2*b*x+2*a))/b^4$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {3801, 3797, 2221, 2611, 2320, 6724, 30}

$$\begin{aligned} \int x^3 \coth^2(a + bx) dx = & -\frac{3 \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^4} + \frac{3x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^3} \\ & + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} - \frac{x^3 \coth(a + bx)}{b} - \frac{x^3}{b} + \frac{x^4}{4} \end{aligned}$$

[In] $\operatorname{Int}[x^3*\operatorname{Coth}[a + b*x]^2, x]$

[Out] $-(x^3/b) + x^4/4 - (x^3*\operatorname{Coth}[a + b*x])/b + (3*x^2*\operatorname{Log}[1 - E^(2*(a + b*x))]) / b^2 + (3*x*\operatorname{PolyLog}[2, E^(2*(a + b*x))]) / b^3 - (3*\operatorname{PolyLog}[3, E^(2*(a + b*x))]) / (2*b^4)$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/
((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x))
))^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; Functi
onOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(
F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(
f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))
)/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_))^(p_.)]/((d_.) + (e_.)*(x_)), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d}
```

, e, n, p}, x] && EqQ[b*d, a*e]

Rubi steps

$$\begin{aligned}
 \text{integral} &= -\frac{x^3 \coth(a + bx)}{b} + \frac{3 \int x^2 \coth(a + bx) dx}{b} + \int x^3 dx \\
 &= -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} - \frac{6 \int \frac{e^{2(a+bx)} x^2}{1-e^{2(a+bx)}} dx}{b} \\
 &= -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} - \frac{6 \int x \log(1 - e^{2(a+bx)}) dx}{b^2} \\
 &= -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} \\
 &\quad + \frac{3x \text{PolyLog}(2, e^{2(a+bx)})}{b^3} - \frac{3 \int \text{PolyLog}(2, e^{2(a+bx)}) dx}{b^3} \\
 &= -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} \\
 &\quad + \frac{3x \text{PolyLog}(2, e^{2(a+bx)})}{b^3} - \frac{3 \text{Subst}\left(\int \frac{\text{PolyLog}(2, x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^4} \\
 &= -\frac{x^3}{b} + \frac{x^4}{4} - \frac{x^3 \coth(a + bx)}{b} + \frac{3x^2 \log(1 - e^{2(a+bx)})}{b^2} \\
 &\quad + \frac{3x \text{PolyLog}(2, e^{2(a+bx)})}{b^3} - \frac{3 \text{PolyLog}(3, e^{2(a+bx)})}{2b^4}
 \end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 222 vs. $2(87) = 174$.

Time = 1.09 (sec), antiderivative size = 222, normalized size of antiderivative = 2.55

$$\begin{aligned}
 \int x^3 \coth^2(a + bx) dx &= \frac{x^4}{4} \\
 &- \frac{e^{2a} (2b^3 e^{-2a} x^3 - 3b^2 (1 - e^{-2a}) x^2 \log(1 - e^{-a-bx}) - 3b^2 (1 - e^{-2a}) x^2 \log(1 + e^{-a-bx}) + 6b (1 - e^{-2a}) x)}{b} \\
 &+ \frac{x^3 \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx)}{b}
 \end{aligned}$$

[In] Integrate[x^3*Coth[a + b*x]^2, x]

[Out] $x^4/4 - (E^{(2*a)}*((2*b^3*x^3)/E^{(2*a)} - 3*b^2*(1 - E^{(-2*a))}*x^2*\text{Log}[1 - E^{(-a - b*x)}] - 3*b^2*(1 - E^{(-2*a))}*x^2*\text{Log}[1 + E^{(-a - b*x)}] + 6*b*(1 - E^{(-2*a))}*x*\text{PolyLog}[2, -E^{(-a - b*x)}] + 6*b*(1 - E^{(-2*a))}*x*\text{PolyLog}[2, E^{(-a - b*x)}] + 6*(1 - E^{(-2*a))}*\text{PolyLog}[3, -E^{(-a - b*x)}] + 6*(1 - E^{(-2*a))}*\text{PolyLog}[3, E^{(-a - b*x)}]))/(b^4*(-1 + E^{(2*a)})) + (x^3*\text{Csch}[a]*\text{Csch}[a + b*x]*\text{Sinh}[b*x])/b$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 197 vs. $2(83) = 166$.

Time = 0.11 (sec), antiderivative size = 198, normalized size of antiderivative = 2.28

method	result
risch	$\frac{x^4}{4} - \frac{2x^3}{b(e^{2bx+2a}-1)} + \frac{4a^3}{b^4} + \frac{6x a^2}{b^3} - \frac{6 \text{polylog}(3, e^{bx+a})}{b^4} - \frac{6 \text{polylog}(3, -e^{bx+a})}{b^4} + \frac{6 \text{polylog}(2, e^{bx+a})x}{b^3} + \frac{3 \ln(e^{bx+a}+1)a^2}{b^2}$

[In] `int(x^3*cOTH(b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out] $1/4*x^4 - 2*x^3/b/(\exp(2*b*x+2*a)-1) + 4/b^4*a^3 + 6*x/b^3*a^2 - 6/b^4*\text{polylog}(3, \exp(b*x+a)) - 6/b^4*\text{polylog}(3, -\exp(b*x+a)) + 6/b^3*\text{polylog}(2, \exp(b*x+a))*x + 3/b^2*\ln(\exp(b*x+a)+1)*x^2 + 6/b^3*\text{polylog}(2, -\exp(b*x+a))*x - 2*x^3/b + 3/b^2*\ln(1-\exp(b*x+a))*x^2 - 6/b^4*a^2*\ln(\exp(b*x+a)) - 3/b^4*\ln(1-\exp(b*x+a))*a^2 + 3/b^4*a^2*\ln(\exp(b*x+a)-1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 632 vs. $2(82) = 164$.

Time = 0.26 (sec), antiderivative size = 632, normalized size of antiderivative = 7.26

$$\int x^3 \coth^2(a + bx) dx = \frac{b^4 x^4 - 8 a^3 - (b^4 x^4 - 8 b^3 x^3 - 8 a^3) \cosh(bx + a)^2 - 2(b^4 x^4 - 8 b^3 x^3 - 8 a^3) \cosh(bx + a) \sinh(bx + a)}{b^4 x^4 - 8 a^3 - (b^4 x^4 - 8 b^3 x^3 - 8 a^3) \cosh(bx + a) \sinh(bx + a)}$$

[In] `integrate(x^3*cOTH(b*x+a)^2, x, algorithm="fricas")`

[Out] $-1/4*(b^4*x^4 - 8*a^3 - (b^4*x^4 - 8*b^3*x^3 - 8*a^3)*\cosh(b*x + a)^2 - 2*(b^4*x^4 - 8*b^3*x^3 - 8*a^3)*\cosh(b*x + a)*\sinh(b*x + a) - (b^4*x^4 - 8*b^3*x^3 - 8*a^3)*\sinh(b*x + a)^2 - 24*(b*x*\cosh(b*x + a)^2 + 2*b*x*\cosh(b*x + a)*\sinh(b*x + a) + b*x*\sinh(b*x + a)^2 - b*x)*\text{dilog}(\cosh(b*x + a) + \sinh(b*x + a)) - 24*(b*x*\cosh(b*x + a)^2 + 2*b*x*\cosh(b*x + a)*\sinh(b*x + a) + b*x*\sinh(b*x + a)^2 - b*x)*\text{dilog}(-\cosh(b*x + a) - \sinh(b*x + a)) - 12*(b^2*x^2*\cosh(b*x + a)^2 + 2*b^2*x^2*\cosh(b*x + a)*\sinh(b*x + a) + b^2*x^2*\sinh(b*x + a)^2 - b^2*x^2)*\log(\cosh(b*x + a) + \sinh(b*x + a) + 1) - 12*(a^2*\cosh(b*x + a)^2 + 2*a^2*\cosh(b*x + a)*\sinh(b*x + a) + a^2*\sinh(b*x + a)^2 - a^2)*\text{log}(\cosh(b*x + a) + \sinh(b*x + a) - 1) + 12*(b^2*x^2 - (b^2*x^2 - a^2)*\cosh(b*x + a)^2 - 2*(b^2*x^2 - a^2)*\cosh(b*x + a)*\sinh(b*x + a) - (b^2*x^2 - a^2)*\sinh(b*x + a)^2 - a^2)*\log(-\cosh(b*x + a) - \sinh(b*x + a) + 1) + 24*(\cosh(b*x + a)^2 + 2*\cosh(b*x + a)*\sinh(b*x + a) + \sinh(b*x + a)^2 - 1)*\text{polylog}(3, \cosh(b*x + a) + \sinh(b*x + a)) + 24*(\cosh(b*x + a)^2 + 2*\cosh(b*x + a)*\sinh(b*x + a) + \sinh(b*x + a)^2 - 1)*\text{polylog}(3, -\cosh(b*x + a) - \sinh(b*x + a))/((b^4*\cosh(b*x + a)^2 + 2*b^4*\cosh(b*x + a)*\sinh(b*x + a) + b^4*\sinh(b*x + a)^2 - b^4)$

Sympy [F]

$$\int x^3 \coth^2(a + bx) dx = \int x^3 \coth^2(a + bx) dx$$

```
[In] integrate(x**3*coth(b*x+a)**2,x)
[Out] Integral(x**3*coth(a + b*x)**2, x)
```

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec), antiderivative size = 146, normalized size of antiderivative = 1.68

$$\begin{aligned} \int x^3 \coth^2(a + bx) dx = & -\frac{2x^3}{b} + \frac{bx^4 e^{(2bx+2a)} - bx^4 - 8x^3}{4(b e^{(2bx+2a)} - b)} \\ & + \frac{3(b^2 x^2 \log(e^{(bx+a)} + 1) + 2bx \text{Li}_2(-e^{(bx+a)}) - 2\text{Li}_3(-e^{(bx+a)}))}{b^4} \\ & + \frac{3(b^2 x^2 \log(-e^{(bx+a)} + 1) + 2bx \text{Li}_2(e^{(bx+a)}) - 2\text{Li}_3(e^{(bx+a)}))}{b^4} \end{aligned}$$

```
[In] integrate(x^3*coth(b*x+a)^2,x, algorithm="maxima")
[Out] -2*x^3/b + 1/4*(b*x^4*e^(2*b*x + 2*a) - b*x^4 - 8*x^3)/(b*e^(2*b*x + 2*a) - b) + 3*(b^2*x^2*log(e^(b*x + a) + 1) + 2*b*x*dilog(-e^(b*x + a))) - 2*polylog(3, -e^(b*x + a))/b^4 + 3*(b^2*x^2*log(-e^(b*x + a) + 1) + 2*b*x*dilog(e^(b*x + a)) - 2*polylog(3, e^(b*x + a)))/b^4
```

Giac [F]

$$\int x^3 \coth^2(a + bx) dx = \int x^3 \coth(bx + a)^2 dx$$

```
[In] integrate(x^3*coth(b*x+a)^2,x, algorithm="giac")
[Out] integrate(x^3*coth(b*x + a)^2, x)
```

Mupad [F(-1)]

Timed out.

$$\int x^3 \coth^2(a + bx) dx = \int x^3 \coth(a + bx)^2 dx$$

[In] int(x^3*coth(a + b*x)^2,x)

[Out] int(x^3*coth(a + b*x)^2, x)

3.7 $\int x^2 \coth^2(a + bx) dx$

Optimal result	69
Rubi [A] (verified)	69
Mathematica [A] (verified)	71
Maple [B] (verified)	71
Fricas [B] (verification not implemented)	72
Sympy [F]	72
Maxima [A] (verification not implemented)	72
Giac [F]	73
Mupad [F(-1)]	73

Optimal result

Integrand size = 12, antiderivative size = 65

$$\begin{aligned} \int x^2 \coth^2(a + bx) dx = & -\frac{x^2}{b} + \frac{x^3}{3} - \frac{x^2 \coth(a + bx)}{b} \\ & + \frac{2x \log(1 - e^{2(a+bx)})}{b^2} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{b^3} \end{aligned}$$

[Out] $-x^{2/b} + 1/3*x^3 - x^2*\coth(b*x+a)/b + 2*x*\ln(1-exp(2*b*x+2*a))/b^2 + \text{polylog}(2, \exp(2*b*x+2*a))/b^3$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3801, 3797, 2221, 2317, 2438, 30}

$$\begin{aligned} \int x^2 \coth^2(a + bx) dx = & \frac{\text{PolyLog}(2, e^{2(a+bx)})}{b^3} + \frac{2x \log(1 - e^{2(a+bx)})}{b^2} \\ & - \frac{x^2 \coth(a + bx)}{b} - \frac{x^2}{b} + \frac{x^3}{3} \end{aligned}$$

[In] $\text{Int}[x^2 \coth[a + b*x]^2, x]$

[Out] $-(x^{2/b}) + x^{3/3} - (x^2 \coth[a + b*x])/b + (2*x \log[1 - E^{(2*(a + b*x))}])/b^2 + \text{PolyLog}[2, E^{(2*(a + b*x))}]/b^3$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] := \text{Simp}[x^{(m + 1)/(m + 1)}, x] /; \text{FreeQ}[m, x] \&& N \in \mathbb{Q}[m, -1]$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_)*(x_))))^(n_.)*((c_.) + (d_)*(x_))^(m_.))/((a_) + (b_)*(F_)^((g_.)*(e_.) + (f_)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*(F_)^((e_.)*(c_.) + (d_)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3797

```
Int[((c_.) + (d_)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi))), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_.)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x^2 \coth(a + bx)}{b} + \frac{2 \int x \coth(a + bx) dx}{b} + \int x^2 dx \\
&= -\frac{x^2}{b} + \frac{x^3}{3} - \frac{x^2 \coth(a + bx)}{b} - \frac{4 \int \frac{e^{2(a+bx)} x}{1-e^{2(a+bx)}} dx}{b} \\
&= -\frac{x^2}{b} + \frac{x^3}{3} - \frac{x^2 \coth(a + bx)}{b} + \frac{2x \log(1 - e^{2(a+bx)})}{b^2} - \frac{2 \int \log(1 - e^{2(a+bx)}) dx}{b^2} \\
&= -\frac{x^2}{b} + \frac{x^3}{3} - \frac{x^2 \coth(a + bx)}{b} + \frac{2x \log(1 - e^{2(a+bx)})}{b^2} - \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(a+bx)}\right)}{b^3}
\end{aligned}$$

$$= -\frac{x^2}{b} + \frac{x^3}{3} - \frac{x^2 \coth(a + bx)}{b} + \frac{2x \log(1 - e^{2(a+bx)})}{b^2} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{b^3}$$

Mathematica [A] (verified)

Time = 1.58 (sec) , antiderivative size = 117, normalized size of antiderivative = 1.80

$$\begin{aligned} \int x^2 \coth^2(a + bx) dx = & -\frac{2 \text{PolyLog}(2, -e^{-a-bx})}{b^3} - \frac{2 \text{PolyLog}(2, e^{-a-bx})}{b^3} \\ & + \frac{1}{3} x \left(\frac{6x}{b - be^{2a}} + x^2 + \frac{6 \log(1 - e^{-a-bx})}{b^2} + \frac{6 \log(1 + e^{-a-bx})}{b^2} \right. \\ & \left. + \frac{3x \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx)}{b} \right) \end{aligned}$$

[In] `Integrate[x^2*Coth[a + b*x]^2, x]`

[Out] $\frac{(-2 \text{PolyLog}[2, -E^{(-a - b x)}])}{b^3} - \frac{(2 \text{PolyLog}[2, E^{(-a - b x)}])}{b^3} + \left(x \left(\frac{(6 x)/(b - b E^{(2 a)}) + x^2 + (6 \log[1 - E^{(-a - b x)}])/b^2 + (6 \log[1 + E^{(-a - b x)}])/b^2 + (3 x \operatorname{Csch}[a] \operatorname{Csch}[a + b x] \operatorname{Sinh}[b x])/b \right) \right)/3$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 155 vs. $2(63) = 126$.

Time = 0.12 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.40

method	result
risch	$\frac{x^3}{3} - \frac{2x^2}{b(e^{2bx+2a}-1)} - \frac{2x^2}{b} - \frac{4ax}{b^2} - \frac{2a^2}{b^3} + \frac{2 \ln(e^{bx+a}+1)x}{b^2} + \frac{2 \text{polylog}(2, -e^{bx+a})}{b^3} + \frac{2 \ln(1-e^{bx+a})x}{b^2} + \frac{2 \ln(1-e^{bx+a})}{b^3}$

[In] `int(x^2*coth(b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{3} x^3 - 2 x^2/b / (\exp(2 b x + 2 a) - 1) - 2 x^2/b - 4 a x/b^2 - 2/b^3 a^2 + 2/b^2 \ln(\exp(b x + a) + 1) * x + 2/b^3 \text{polylog}(2, -\exp(b x + a)) + 2/b^2 \ln(1 - \exp(b x + a)) * x + 2/b^3 \ln(1 - \exp(b x + a)) * a + 2/b^3 \text{polylog}(2, \exp(b x + a)) - 2/b^3 a \ln(\exp(b x + a) - 1) + 4/b^3 * a \ln(\exp(b x + a))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 453 vs. $2(62) = 124$.

Time = 0.26 (sec), antiderivative size = 453, normalized size of antiderivative = 6.97

$$\int x^2 \coth^2(a + bx) dx = -\frac{b^3 x^3 - (b^3 x^3 - 6 b^2 x^2 + 6 a^2) \cosh(bx + a)^2 - 2(b^3 x^3 - 6 b^2 x^2 + 6 a^2) \cosh(bx + a) \sinh(bx + a) - (b^3 x^3 - 6 b^2 x^2 + 6 a^2) \sinh(bx + a)^2}{b^3}$$

```
[In] integrate(x^2*coth(b*x+a)^2,x, algorithm="fricas")
[Out] -1/3*(b^3*x^3 - (b^3*x^3 - 6*b^2*x^2 + 6*a^2)*cosh(b*x + a)^2 - 2*(b^3*x^3 - 6*b^2*x^2 + 6*a^2)*cosh(b*x + a)*sinh(b*x + a) - (b^3*x^3 - 6*b^2*x^2 + 6*a^2)*sinh(b*x + a)^2 + 6*a^2 - 6*(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*dilog(cosh(b*x + a) + sinh(b*x + a)) - 6*(cosh(b*x + a)^2 + 2*cosh(b*x + a)*sinh(b*x + a) + sinh(b*x + a)^2 - 1)*dilog(-cosh(b*x + a) - sinh(b*x + a)) - 6*(b*x*cosh(b*x + a)^2 + 2*b*x*cosh(b*x + a)*sinh(b*x + a) + b*x*sinh(b*x + a)^2 - b*x)*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 6*(a*cosh(b*x + a)^2 + 2*a*cosh(b*x + a)*sinh(b*x + a) + a*sinh(b*x + a)^2 - a)*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 6*((b*x + a)*cosh(b*x + a)^2 + 2*(b*x + a)*cosh(b*x + a)*sinh(b*x + a) + (b*x + a)*sinh(b*x + a)^2 - b*x - a)*log(-cosh(b*x + a) - sinh(b*x + a) + 1))/(b^3*cosh(b*x + a)^2 + 2*b^3*cosh(b*x + a)*sinh(b*x + a) + b^3*sinh(b*x + a)^2 - b^3)
```

Sympy [F]

$$\int x^2 \coth^2(a + bx) dx = \int x^2 \coth^2(a + bx) dx$$

```
[In] integrate(x**2*coth(b*x+a)**2,x)
[Out] Integral(x**2*coth(a + b*x)**2, x)
```

Maxima [A] (verification not implemented)

none

Time = 0.27 (sec), antiderivative size = 108, normalized size of antiderivative = 1.66

$$\begin{aligned} \int x^2 \coth^2(a + bx) dx &= -\frac{2 x^2}{b} + \frac{b x^3 e^{(2 b x + 2 a)} - b x^3 - 6 x^2}{3 (b e^{(2 b x + 2 a)} - b)} \\ &+ \frac{2 (b x \log(e^{(b x + a)} + 1) + \text{Li}_2(-e^{(b x + a)}))}{b^3} \\ &+ \frac{2 (b x \log(-e^{(b x + a)} + 1) + \text{Li}_2(e^{(b x + a)}))}{b^3} \end{aligned}$$

[In] `integrate(x^2*coth(b*x+a)^2,x, algorithm="maxima")`
[Out]
$$\frac{-2x^2/b + 1/3*(b*x^3*e^{(2*b*x + 2*a)} - b*x^3 - 6*x^2)/(b*e^{(2*b*x + 2*a)} - b) + 2*(b*x*log(e^{(b*x + a)} + 1) + dilog(-e^{(b*x + a)}))/b^3 + 2*(b*x*log(-e^{(b*x + a)} + 1) + dilog(e^{(b*x + a)}))/b^3}{}$$

Giac [F]

$$\int x^2 \coth^2(a + bx) dx = \int x^2 \coth(bx + a)^2 dx$$

[In] `integrate(x^2*coth(b*x+a)^2,x, algorithm="giac")`
[Out] `integrate(x^2*coth(b*x + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int x^2 \coth^2(a + bx) dx = \int x^2 \coth(a + b x)^2 dx$$

[In] `int(x^2*coth(a + b*x)^2,x)`
[Out] `int(x^2*coth(a + b*x)^2, x)`

3.8 $\int x \coth^2(a + bx) dx$

Optimal result	74
Rubi [A] (verified)	74
Mathematica [A] (verified)	75
Maple [A] (verified)	75
Fricas [B] (verification not implemented)	76
Sympy [B] (verification not implemented)	76
Maxima [B] (verification not implemented)	77
Giac [B] (verification not implemented)	77
Mupad [B] (verification not implemented)	77

Optimal result

Integrand size = 10, antiderivative size = 31

$$\int x \coth^2(a + bx) dx = \frac{x^2}{2} - \frac{x \coth(a + bx)}{b} + \frac{\log(\sinh(a + bx))}{b^2}$$

[Out] $1/2*x^2-x*\coth(b*x+a)/b+\ln(\sinh(b*x+a))/b^2$

Rubi [A] (verified)

Time = 0.02 (sec), antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3801, 3556, 30}

$$\int x \coth^2(a + bx) dx = \frac{\log(\sinh(a + bx))}{b^2} - \frac{x \coth(a + bx)}{b} + \frac{x^2}{2}$$

[In] $\text{Int}[x*\text{Coth}[a + b*x]^2, x]$

[Out] $x^2/2 - (x*\text{Coth}[a + b*x])/b + \text{Log}[\text{Sinh}[a + b*x]]/b^2$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3801

```

Int[((c_) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symb
o1] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x \coth(a + bx)}{b} + \frac{\int \coth(a + bx) dx}{b} + \int x dx \\
&= \frac{x^2}{2} - \frac{x \coth(a + bx)}{b} + \frac{\log(\sinh(a + bx))}{b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.17 (sec) , antiderivative size = 46, normalized size of antiderivative = 1.48

$$\begin{aligned}
&\int x \coth^2(a + bx) dx \\
&= \frac{b^2 x^2 - 2bx \coth(a) + 2 \log(\sinh(a + bx)) + 2bx \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx)}{2b^2}
\end{aligned}$$

[In] `Integrate[x*Coth[a + b*x]^2, x]`

[Out] $\frac{(b^2 x^2 - 2bx \coth(a) + 2 \log(\sinh(a + bx)) + 2bx \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx))}{2b^2}$

Maple [A] (verified)

Time = 0.11 (sec) , antiderivative size = 54, normalized size of antiderivative = 1.74

method	result	size
risch	$\frac{x^2}{2} - \frac{2x}{b} - \frac{2a}{b^2} - \frac{2x}{b(e^{2bx+2a}-1)} + \frac{\ln(e^{2bx+2a}-1)}{b^2}$	54
parallelrisch	$\frac{-2 \ln(1-\tanh(bx+a)) \tanh(bx+a) + 2 \ln(\tanh(bx+a)) \tanh(bx+a) + xb(-2+(bx-2) \tanh(bx+a))}{2b^2 \tanh(bx+a)}$	66

[In] `int(x*coth(b*x+a)^2, x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}x^2 - 2x/b - 2a/b^2 - 2x/b / (\exp(2b*x+2*a)-1) + 1/b^2 \ln(\exp(2b*x+2*a)-1)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(29) = 58$.

Time = 0.26 (sec) , antiderivative size = 189, normalized size of antiderivative = 6.10

$$\int x \coth^2(a + bx) dx = \frac{b^2 x^2 - (b^2 x^2 - 4 bx) \cosh(bx + a)^2 - 2(b^2 x^2 - 4 bx) \cosh(bx + a) \sinh(bx + a) - (b^2 x^2 - 4 bx) \sinh(bx + a)}{2(b^2 \cosh(bx + a)^2 + 2 b^2 \cosh(bx + a))}$$

[In] `integrate(x*coth(b*x+a)^2,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -1/2*(b^2*x^2 - (b^2*x^2 - 4*b*x)*\cosh(b*x + a)^2 - 2*(b^2*x^2 - 4*b*x)*\cos \\ & h(b*x + a)*\sinh(b*x + a) - (b^2*x^2 - 4*b*x)*\sinh(b*x + a)^2 - 2*(\cosh(b*x \\ & + a)^2 + 2*\cosh(b*x + a)*\sinh(b*x + a) + \sinh(b*x + a)^2 - 1)*\log(2*\sinh(b* \\ & x + a)/(\cosh(b*x + a) - \sinh(b*x + a))))/(b^2*cosh(b*x + a)^2 + 2*b^2*cosh(b* \\ & x + a)*\sinh(b*x + a) + b^2*sinh(b*x + a)^2 - b^2) \end{aligned}$$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 190 vs. $2(26) = 52$.

Time = 0.68 (sec) , antiderivative size = 190, normalized size of antiderivative = 6.13

$$\int x \coth^2(a + bx) dx = \begin{cases} \frac{x^2 \coth^2(a)}{2} & \text{for } b = 0 \\ -\frac{x \log(-e^{-bx}) \coth^2(bx + \log(-e^{-bx}))}{b} - \frac{\log(-e^{-bx})^2 \coth^2(bx + \log(-e^{-bx}))}{2b^2} & \text{for } a = \log(-e^{-bx}) \\ -\frac{x \log(e^{-bx}) \coth^2(bx + \log(e^{-bx}))}{b} - \frac{\log(e^{-bx})^2 \coth^2(bx + \log(e^{-bx}))}{2b^2} & \text{for } a = \log(e^{-bx}) \\ \frac{x^2}{2} + \frac{x}{b} - \frac{x}{b \tanh(a+bx)} - \frac{\log(\tanh(a+bx)+1)}{b^2} + \frac{\log(\tanh(a+bx))}{b^2} & \text{otherwise} \end{cases}$$

[In] `integrate(x*coth(b*x+a)**2,x)`

[Out] `Piecewise((x**2*coth(a)**2/2, Eq(b, 0)), (-x*log(-exp(-b*x))*coth(b*x + log(-exp(-b*x)))*2/b - log(-exp(-b*x))*2*coth(b*x + log(-exp(-b*x)))*2/(2*b**2), Eq(a, log(-exp(-b*x)))), (-x*log(exp(-b*x))*coth(b*x + log(exp(-b*x)))*2/b - log(exp(-b*x))*2*coth(b*x + log(exp(-b*x)))*2/(2*b**2), Eq(a, log(exp(-b*x)))), (x**2/2 + x/b - x/(b*tanh(a + b*x)) - log(tanh(a + b*x) + 1)/b**2 + log(tanh(a + b*x))/b**2, True))`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 115 vs. $2(29) = 58$.

Time = 0.23 (sec) , antiderivative size = 115, normalized size of antiderivative = 3.71

$$\begin{aligned} \int x \coth^2(a + bx) dx &= -\frac{xe^{(2bx+2a)}}{be^{(2bx+2a)} - b} - \frac{bx^2 - (bx^2e^{(2a)} - 2xe^{(2a)})e^{(2bx)}}{2(be^{(2bx+2a)} - b)} \\ &\quad + \frac{\log((e^{(bx+a)} + 1)e^{(-a)})}{b^2} + \frac{\log((e^{(bx+a)} - 1)e^{(-a)})}{b^2} \end{aligned}$$

[In] `integrate(x*coth(b*x+a)^2,x, algorithm="maxima")`

[Out]
$$\frac{-x e^{(2b x + 2 a)} / (b e^{(2b x + 2 a)} - b) - 1/2 * (b x^2 - (b x^2 e^{(2 a)} - 2 x e^{(2 a)}) e^{(2 b x)}) / (b e^{(2b x + 2 a)} - b) + \log((e^{(b x + a)} + 1) e^{(-a)}) / b^2 + \log((e^{(b x + a)} - 1) e^{(-a)}) / b^2}{2}$$

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 98 vs. $2(29) = 58$.

Time = 0.29 (sec) , antiderivative size = 98, normalized size of antiderivative = 3.16

$$\begin{aligned} \int x \coth^2(a + bx) dx \\ = \frac{b^2 x^2 e^{(2bx+2a)} - b^2 x^2 - 4 b x e^{(2bx+2a)} + 2 e^{(2bx+2a)} \log(e^{(2bx+2a)} - 1) - 2 \log(e^{(2bx+2a)} - 1)}{2(b^2 e^{(2bx+2a)} - b^2)} \end{aligned}$$

[In] `integrate(x*coth(b*x+a)^2,x, algorithm="giac")`

[Out]
$$\frac{1/2 * (b^2 x^2 e^{(2b x + 2 a)} - b^2 x^2 - 4 b x e^{(2b x + 2 a)} + 2 e^{(2b x + 2 a)} \log(e^{(2b x + 2 a)} - 1) - 2 \log(e^{(2b x + 2 a)} - 1)) / (b^2 e^{(2b x + 2 a)} - b^2)}{2}$$

Mupad [B] (verification not implemented)

Time = 1.89 (sec) , antiderivative size = 48, normalized size of antiderivative = 1.55

$$\int x \coth^2(a + bx) dx = \frac{\ln(e^{2a} e^{2bx} - 1)}{b^2} - \frac{2x}{b} + \frac{x^2}{2} - \frac{2x}{b(e^{2a+2bx} - 1)}$$

[In] `int(x*coth(a + b*x)^2,x)`

[Out]
$$\frac{\log(\exp(2*a) * \exp(2*b*x) - 1) / b^2 - (2*x) / b + x^2 / 2 - (2*x) / (b * (\exp(2*a) + 2 * b * x) - 1))}{2}$$

3.9 $\int \frac{\coth^2(a+bx)}{x} dx$

Optimal result	78
Rubi [N/A]	78
Mathematica [N/A]	79
Maple [N/A] (verified)	79
Fricas [N/A]	79
Sympy [N/A]	79
Maxima [N/A]	80
Giac [N/A]	80
Mupad [N/A]	80

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\coth^2(a + bx)}{x} dx = \text{Int}\left(\frac{\coth^2(a + bx)}{x}, x\right)$$

[Out] Unintegrable(coth(b*x+a)^2/x, x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth^2(a + bx)}{x} dx$$

[In] Int[Coth[a + b*x]^2/x, x]

[Out] Defer[Int][Coth[a + b*x]^2/x, x]

Rubi steps

$$\text{integral} = \int \frac{\coth^2(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 25.94 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth^2(a + bx)}{x} dx$$

[In] Integrate[Coth[a + b*x]^2/x, x]

[Out] Integrate[Coth[a + b*x]^2/x, x]

Maple [N/A] (verified)

Not integrable

Time = 0.04 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)^2}{x} dx$$

[In] int(coth(b*x+a)^2/x, x)

[Out] int(coth(b*x+a)^2/x, x)

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth(bx + a)^2}{x} dx$$

[In] integrate(coth(b*x+a)^2/x, x, algorithm="fricas")

[Out] integral(coth(b*x + a)^2/x, x)

Sympy [N/A]

Not integrable

Time = 0.52 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth^2(a + bx)}{x} dx$$

[In] integrate(coth(b*x+a)**2/x, x)

[Out] Integral(coth(a + b*x)**2/x, x)

Maxima [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 69, normalized size of antiderivative = 5.75

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth(bx + a)^2}{x} dx$$

[In] `integrate(coth(b*x+a)^2/x,x, algorithm="maxima")`

[Out] `-2/(b*x*e^(2*b*x + 2*a) - b*x) + integrate(1/(b*x^2*e^(b*x + a) + b*x^2), x) - integrate(1/(b*x^2*e^(b*x + a) - b*x^2), x) + log(x)`

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth(bx + a)^2}{x} dx$$

[In] `integrate(coth(b*x+a)^2/x,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)^2/x, x)`

Mupad [N/A]

Not integrable

Time = 1.91 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x} dx = \int \frac{\coth(a + b x)^2}{x} dx$$

[In] `int(coth(a + b*x)^2/x,x)`

[Out] `int(coth(a + b*x)^2/x, x)`

3.10 $\int \frac{\coth^2(a+bx)}{x^2} dx$

Optimal result	81
Rubi [N/A]	81
Mathematica [N/A]	82
Maple [N/A] (verified)	82
Fricas [N/A]	82
Sympy [N/A]	82
Maxima [N/A]	83
Giac [N/A]	83
Mupad [N/A]	83

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \text{Int}\left(\frac{\coth^2(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable($\coth(b*x+a)^2/x^2$, x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth^2(a + bx)}{x^2} dx$$

[In] Int[Coth[a + b*x]^2/x^2, x]

[Out] Defer[Int][Coth[a + b*x]^2/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\coth^2(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 25.75 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth^2(a + bx)}{x^2} dx$$

```
[In] Integrate[Coth[a + b*x]^2/x^2, x]
[Out] Integrate[Coth[a + b*x]^2/x^2, x]
```

Maple [N/A] (verified)

Not integrable

Time = 0.05 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)^2}{x^2} dx$$

```
[In] int(coth(b*x+a)^2/x^2, x)
[Out] int(coth(b*x+a)^2/x^2, x)
```

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^2}{x^2} dx$$

```
[In] integrate(coth(b*x+a)^2/x^2, x, algorithm="fricas")
[Out] integral(coth(b*x + a)^2/x^2, x)
```

Sympy [N/A]

Not integrable

Time = 0.49 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth^2(a + bx)}{x^2} dx$$

```
[In] integrate(coth(b*x+a)**2/x**2, x)
[Out] Integral(coth(a + b*x)**2/x**2, x)
```

Maxima [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 91, normalized size of antiderivative = 7.58

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^2}{x^2} dx$$

[In] `integrate(coth(b*x+a)^2/x^2,x, algorithm="maxima")`

[Out] `-(b*x*e^(2*b*x + 2*a) - b*x + 2)/(b*x^2*e^(2*b*x + 2*a) - b*x^2) + 2*integrate(1/(b*x^3*e^(b*x + a) + b*x^3), x) - 2*integrate(1/(b*x^3*e^(b*x + a) - b*x^3), x)`

Giac [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^2}{x^2} dx$$

[In] `integrate(coth(b*x+a)^2/x^2,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)^2/x^2, x)`

Mupad [N/A]

Not integrable

Time = 1.86 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^2(a + bx)}{x^2} dx = \int \frac{\coth(a + b x)^2}{x^2} dx$$

[In] `int(coth(a + b*x)^2/x^2,x)`

[Out] `int(coth(a + b*x)^2/x^2, x)`

3.11 $\int x^3 \coth^3(a + bx) dx$

Optimal result	84
Rubi [A] (verified)	84
Mathematica [B] (verified)	87
Maple [B] (verified)	88
Fricas [B] (verification not implemented)	88
Sympy [F]	90
Maxima [A] (verification not implemented)	90
Giac [F]	91
Mupad [F(-1)]	91

Optimal result

Integrand size = 12, antiderivative size = 179

$$\begin{aligned} \int x^3 \coth^3(a + bx) dx = & -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a + bx)}{2b^2} - \frac{x^3 \coth^2(a + bx)}{2b} \\ & + \frac{3x \log(1 - e^{2(a+bx)})}{b^3} + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} \\ & + \frac{3 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^4} + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\ & - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \operatorname{PolyLog}(4, e^{2(a+bx)})}{4b^4} \end{aligned}$$

[Out] $-3/2*x^2/b^2+1/2*x^3/b-1/4*x^4-3/2*x^2*\coth(b*x+a)/b^2-1/2*x^3*\coth(b*x+a)^2/b+3*x^3*\ln(1-\exp(2*b*x+2*a))/b^3+x^3*\ln(1-\exp(2*b*x+2*a))/b+3/2*\operatorname{polylog}(2, \exp(2*b*x+2*a))/b^4+3/2*x^2*\operatorname{polylog}(2, \exp(2*b*x+2*a))/b^2-3/2*x*\operatorname{polylog}(3, \exp(2*b*x+2*a))/b^3+3/4*\operatorname{polylog}(4, \exp(2*b*x+2*a))/b^4$

Rubi [A] (verified)

Time = 0.25 (sec), antiderivative size = 179, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {3801, 3797, 2221, 2317, 2438, 30, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int x^3 \coth^3(a + bx) dx = & \frac{3 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^4} + \frac{3 \operatorname{PolyLog}(4, e^{2(a+bx)})}{4b^4} \\ & - \frac{3x \operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3x \log(1 - e^{2(a+bx)})}{b^3} \\ & + \frac{3x^2 \operatorname{PolyLog}(2, e^{2(a+bx)})}{2b^2} - \frac{3x^2 \coth(a + bx)}{2b^2} \\ & + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} - \frac{x^3 \coth^2(a + bx)}{2b} - \frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} \end{aligned}$$

[In] $\text{Int}[x^3 \coth[a + b*x]^3, x]$

[Out] $(-3*x^2)/(2*b^2) + x^3/(2*b) - x^4/4 - (3*x^2*\coth[a + b*x])/(2*b^2) - (x^3*\coth[a + b*x]^2)/(2*b) + (3*x*\text{Log}[1 - E^{(2*(a + b*x))}])/b^3 + (x^3*\text{Log}[1 - E^{(2*(a + b*x))}])/b + (3*\text{PolyLog}[2, E^{(2*(a + b*x))}])/(2*b^4) + (3*x^2*\text{PolyLog}[2, E^{(2*(a + b*x))}])/(2*b^2) - (3*x*\text{PolyLog}[3, E^{(2*(a + b*x))}])/(2*b^3) + (3*\text{PolyLog}[4, E^{(2*(a + b*x))}])/(4*b^4)$

Rule 30

$\text{Int}[(x_.)^{(m_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[x^{(m + 1)/(m + 1)}, x] /; \text{FreeQ}[m, x] \&& N \in \mathbb{Z}, m = -1]$

Rule 2221

$\text{Int}[((F_.)^{(g_.)*(e_.) + (f_.)*(x_.))})^{(n_.)*(c_.) + (d_.)*(x_.)^{(m_.)}} / ((a_.) + (b_.)*(F_.)^{(g_.)*(e_.) + (f_.)*(x_.))})^{(n_.)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^{(g*(e + f*x))})^{n/a})], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^{(m - 1)}*\text{Log}[1 + b*((F^{(g*(e + f*x))})^{n/a})], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

Rule 2317

$\text{Int}[\text{Log}[(a_.) + (b_.)*(F_.)^{(e_.)*(c_.) + (d_.)*(x_.))}]^{(n_.)}, x_{\text{Symbol}}] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x))})^n], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$

Rule 2320

$\text{Int}[u_, x_{\text{Symbol}}] \rightarrow \text{With}[\{v = \text{FunctionOfExponential}[u, x]\}, \text{Dist}[v/D[v, x], \text{Subst}[\text{Int}[\text{FunctionOfExponentialFunction}[u, x]/x, x], x, v], x]] /; \text{FunctionOfExponentialQ}[u, x] \&& \text{!MatchQ}[u, (w_)*(a_.)*(v_.)^(n_.)]^{(m_.)} /; \text{FreeQ}[\{a, m, n\}, x] \&& \text{IntegerQ}[m*n] \&& \text{!MatchQ}[u, E^{(c_.)*(a_.) + (b_.)*x}]* (F_)[v_] /; \text{FreeQ}[\{a, b, c\}, x] \&& \text{InverseFunctionQ}[F[x]]]$

Rule 2438

$\text{Int}[\text{Log}[(c_.)*(d_.) + (e_.)*(x_.)^{(n_.)})]/(x_), x_{\text{Symbol}}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \&& \text{EqQ}[c*d, 1]$

Rule 2611

$\text{Int}[\text{Log}[1 + (e_.)*(F_.)^{(c_.)*(a_.) + (b_.)*(x_.))}]^{(n_.)}*((f_.) + (g_.)*(x_.)^{(m_.)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(f + g*x)^m)*(\text{PolyLog}[2, (-e)*(F^{(c*(a + b*x))})^n]/(b*c*n*\text{Log}[F])), x] + \text{Dist}[g*(m/(b*c*n*\text{Log}[F])), \text{Int}[(f + g*x)^{(m - 1)}*\text{PolyLog}[2, (-e)*(F^{(c*(a + b*x))})^n], x], x] /; \text{FreeQ}[\{F, a, b, c, e, f, g, n\}, x] \&& \text{GtQ}[m, 0]$

Rule 3797

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_ .)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist [2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^n_, x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_*) + (b_)*(x_)]^p]/((d_*) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_)*(x_))^(m_)*PolyLog[n_, (d_)*((F_)^((c_)*(a_*) + (b_)*(x_)))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x^3 \coth^2(a + bx)}{2b} + \frac{3 \int x^2 \coth^2(a + bx) dx}{2b} + \int x^3 \coth(a + bx) dx \\
&= -\frac{x^4}{4} - \frac{3x^2 \coth(a + bx)}{2b^2} - \frac{x^3 \coth^2(a + bx)}{2b} \\
&\quad - 2 \int \frac{e^{2(a+bx)} x^3}{1 - e^{2(a+bx)}} dx + \frac{3 \int x \coth(a + bx) dx}{b^2} + \frac{3 \int x^2 dx}{2b} \\
&= -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a + bx)}{2b^2} - \frac{x^3 \coth^2(a + bx)}{2b} \\
&\quad + \frac{x^3 \log(1 - e^{2(a+bx)})}{b} - \frac{6 \int \frac{e^{2(a+bx)} x}{1 - e^{2(a+bx)}} dx}{b^2} - \frac{3 \int x^2 \log(1 - e^{2(a+bx)}) dx}{b}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a+bx)}{2b^2} - \frac{x^3 \coth^2(a+bx)}{2b} \\
&\quad + \frac{3x \log(1-e^{2(a+bx)})}{b^3} + \frac{x^3 \log(1-e^{2(a+bx)})}{b} + \frac{3x^2 \text{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\
&\quad - \frac{3 \int \log(1-e^{2(a+bx)}) dx}{b^3} - \frac{3 \int x \text{PolyLog}(2, e^{2(a+bx)}) dx}{b^2} \\
&= -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a+bx)}{2b^2} - \frac{x^3 \coth^2(a+bx)}{2b} + \frac{3x \log(1-e^{2(a+bx)})}{b^3} \\
&\quad + \frac{x^3 \log(1-e^{2(a+bx)})}{b} + \frac{3x^2 \text{PolyLog}(2, e^{2(a+bx)})}{2b^2} - \frac{3x \text{PolyLog}(3, e^{2(a+bx)})}{2b^3} \\
&\quad - \frac{3 \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^4} + \frac{3 \int \text{PolyLog}(3, e^{2(a+bx)}) dx}{2b^3} \\
&= -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a+bx)}{2b^2} - \frac{x^3 \coth^2(a+bx)}{2b} + \frac{3x \log(1-e^{2(a+bx)})}{b^3} \\
&\quad + \frac{x^3 \log(1-e^{2(a+bx)})}{b} + \frac{3 \text{PolyLog}(2, e^{2(a+bx)})}{2b^4} + \frac{3x^2 \text{PolyLog}(2, e^{2(a+bx)})}{2b^2} \\
&\quad - \frac{3x \text{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \text{Subst}\left(\int \frac{\text{PolyLog}(3,x)}{x} dx, x, e^{2(a+bx)}\right)}{4b^4} \\
&= -\frac{3x^2}{2b^2} + \frac{x^3}{2b} - \frac{x^4}{4} - \frac{3x^2 \coth(a+bx)}{2b^2} - \frac{x^3 \coth^2(a+bx)}{2b} \\
&\quad + \frac{3x \log(1-e^{2(a+bx)})}{b^3} + \frac{x^3 \log(1-e^{2(a+bx)})}{b} + \frac{3 \text{PolyLog}(2, e^{2(a+bx)})}{2b^4} \\
&\quad + \frac{3x^2 \text{PolyLog}(2, e^{2(a+bx)})}{2b^2} - \frac{3x \text{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{3 \text{PolyLog}(4, e^{2(a+bx)})}{4b^4}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 422 vs. $2(179) = 358$.

Time = 3.71 (sec), antiderivative size = 422, normalized size of antiderivative = 2.36

$$\begin{aligned}
\int x^3 \coth^3(a+bx) dx &= \frac{1}{4} \left(x^4 \coth(a) - \frac{2x^3 \operatorname{csch}^2(a+bx)}{b} \right. \\
&\quad \left. - \frac{2e^{2a} (6b^2 e^{-2a} x^2 + b^4 e^{-2a} x^4 - 6b(1-e^{-2a}) x \log(1-e^{-a-bx}) - 2b^3 e^{-2a} (-1+e^{2a}) x^3 \log(1-e^{-a-bx}))}{b^2} \right. \\
&\quad \left. + \frac{6x^2 \operatorname{csch}(a) \operatorname{csch}(a+bx) \sinh(bx)}{b^2} \right)
\end{aligned}$$

[In] Integrate[x^3*Coth[a + b*x]^3, x]

```
[Out] (x^4*Coth[a] - (2*x^3*Csch[a + b*x]^2)/b - (2*E^(2*a)*((6*b^2*x^2)/E^(2*a)
+ (b^4*x^4)/E^(2*a) - 6*b*(1 - E^(-2*a))*x*Log[1 - E^(-a - b*x)] - (2*b^3*(-1 + E^(2*a))*x^3*Log[1 - E^(-a - b*x)])/E^(2*a) - 6*b*(1 - E^(-2*a))*x*Log[1 + E^(-a - b*x)] - (2*b^3*(-1 + E^(2*a))*x^3*Log[1 + E^(-a - b*x)])/E^(2*a) + 6*(1 - E^(-2*a))*PolyLog[2, -E^(-a - b*x)] + 6*b^2*(1 - E^(-2*a))*x^2*PolyLog[2, -E^(-a - b*x)] + 6*(1 - E^(-2*a))*PolyLog[2, E^(-a - b*x)] + 6*b^2*(1 - E^(-2*a))*x^2*PolyLog[2, E^(-a - b*x)] + 12*b*(1 - E^(-2*a))*x*PolyLog[3, -E^(-a - b*x)] + 12*b*(1 - E^(-2*a))*x*PolyLog[3, E^(-a - b*x)] + 12*(1 - E^(-2*a))*PolyLog[4, -E^(-a - b*x)] + 12*(1 - E^(-2*a))*PolyLog[4, E^(-a - b*x)))/(b^4*(-1 + E^(2*a))) + (6*x^2*Csch[a]*Csch[a + b*x]*Sinh[b*x])/b^2)/4
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 374 vs. $2(161) = 322$.

Time = 0.17 (sec), antiderivative size = 375, normalized size of antiderivative = 2.09

method	result
risch	$-\frac{3a^4}{2b^4} - \frac{x^4}{4} - \frac{a^3 \ln(e^{bx+a}-1)}{b^4} + \frac{2a^3 \ln(e^{bx+a})}{b^4} - \frac{6 \operatorname{polylog}(3, e^{bx+a})x}{b^3} + \frac{3 \operatorname{polylog}(2, -e^{bx+a})x^2}{b^2} - \frac{6 \operatorname{polylog}(3, -e^{bx+a})x}{b^3}$

```
[In] int(x^3*coth(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
[Out] -3/2/b^4*a^4-1/4*x^4-1/b^4*a^3*ln(exp(b*x+a)-1)+2/b^4*a^3*ln(exp(b*x+a))-6/b^3*polylog(3,exp(b*x+a))*x+3/b^2*polylog(2,-exp(b*x+a))*x^2-6/b^3*polylog(3,-exp(b*x+a))*x-2/b^3*a^3*x-3/b^2*x^2-x^2*(2*exp(2*b*x+2*a)*b*x+3*exp(2*b*x+2*a)-3)/b^2/(exp(2*b*x+2*a)-1)^2-3/b^4*a^2+3/b^4*polylog(2,-exp(b*x+a))+6/b^4*polylog(4,-exp(b*x+a))+3/b^4*polylog(2,exp(b*x+a))+6/b^4*polylog(4,exp(b*x+a))-6/b^3*a*x+3/b^2*polylog(2,exp(b*x+a))*x^2+3/b^3*ln(exp(b*x+a)+1)*x+3/b^3*ln(1-exp(b*x+a))*x+1/b*ln(exp(b*x+a)+1)*x^3+1/b^4*ln(1-exp(b*x+a))*a^3+3/b^4*ln(1-exp(b*x+a))*a-3/b^4*a*ln(exp(b*x+a)-1)+6/b^4*a*ln(exp(b*x+a))
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1985 vs. $2(159) = 318$.

Time = 0.31 (sec), antiderivative size = 1985, normalized size of antiderivative = 11.09

$$\int x^3 \coth^3(a + bx) dx = \text{Too large to display}$$

```
[In] integrate(x^3*coth(b*x+a)^3,x, algorithm="fricas")
```

```
[Out] -1/4*(b^4*x^4 + (b^4*x^4 - 2*a^4 + 12*b^2*x^2 - 12*a^2)*cosh(b*x + a)^4 + 4*(b^4*x^4 - 2*a^4 + 12*b^2*x^2 - 12*a^2)*cosh(b*x + a)*sinh(b*x + a)^3 + (b
```

$$\begin{aligned}
& - 4*x^4 - 2*a^4 + 12*b^2*x^2 - 12*a^2)*sinh(b*x + a)^4 - 2*a^4 - 2*(b^4*x^4 \\
& - 4*b^3*x^3 - 2*a^4 + 6*b^2*x^2 - 12*a^2)*cosh(b*x + a)^2 - 2*(b^4*x^4 - 4* \\
& b^3*x^3 - 2*a^4 + 6*b^2*x^2 - 3*(b^4*x^4 - 2*a^4 + 12*b^2*x^2 - 12*a^2)*cos \\
& h(b*x + a)^2 - 12*a^2)*sinh(b*x + a)^2 - 12*a^2 - 12*((b^2*x^2 + 1)*cosh(b* \\
& x + a)^4 + 4*(b^2*x^2 + 1)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^2*x^2 + 1)*si \\
& nh(b*x + a)^4 + b^2*x^2 - 2*(b^2*x^2 + 1)*cosh(b*x + a)^2 - 2*(b^2*x^2 - 3* \\
& (b^2*x^2 + 1)*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 4*((b^2*x^2 + 1)*cosh(\\
& b*x + a)^3 - (b^2*x^2 + 1)*cosh(b*x + a))*sinh(b*x + a) + 1)*dilog(cosh(b*x \\
& + a) + sinh(b*x + a)) - 12*((b^2*x^2 + 1)*cosh(b*x + a)^4 + 4*(b^2*x^2 + 1) \\
&)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^2*x^2 + 1)*sinh(b*x + a)^4 + b^2*x^2 - \\
& 2*(b^2*x^2 + 1)*cosh(b*x + a)^2 - 2*(b^2*x^2 - 3*(b^2*x^2 + 1)*cosh(b*x + \\
& a)^2 + 1)*sinh(b*x + a)^2 + 4*((b^2*x^2 + 1)*cosh(b*x + a)^3 - (b^2*x^2 + 1) \\
&)*cosh(b*x + a)*sinh(b*x + a) + 1)*dilog(-cosh(b*x + a) - sinh(b*x + a)) - \\
& 4*(b^3*x^3 + (b^3*x^3 + 3*b*x)*cosh(b*x + a)^4 + 4*(b^3*x^3 + 3*b*x)*cosh(\\
& b*x + a)*sinh(b*x + a)^3 + (b^3*x^3 + 3*b*x)*sinh(b*x + a)^4 - 2*(b^3*x^3 + \\
& 3*b*x)*cosh(b*x + a)^2 - 2*(b^3*x^3 - 3*(b^3*x^3 + 3*b*x)*cosh(b*x + a)^2 \\
& + 3*b*x)*sinh(b*x + a)^2 + 3*b*x + 4*((b^3*x^3 + 3*b*x)*cosh(b*x + a)^3 - (\\
& b^3*x^3 + 3*b*x)*cosh(b*x + a))*sinh(b*x + a))*log(cosh(b*x + a) + sinh(b*x \\
& + a) + 1) + 4*((a^3 + 3*a)*cosh(b*x + a)^4 + 4*(a^3 + 3*a)*cosh(b*x + a)*s \\
& inh(b*x + a)^3 + (a^3 + 3*a)*sinh(b*x + a)^4 + a^3 - 2*(a^3 + 3*a)*cosh(b*x \\
& + a)^2 - 2*(a^3 - 3*(a^3 + 3*a)*cosh(b*x + a)^2 + 3*a)*sinh(b*x + a)^2 + 4 \\
& *((a^3 + 3*a)*cosh(b*x + a)^3 - (a^3 + 3*a)*cosh(b*x + a))*sinh(b*x + a) + \\
& 3*a)*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 4*(b^3*x^3 + (b^3*x^3 + a^3 + \\
& 3*b*x + 3*a)*cosh(b*x + a)^4 + 4*(b^3*x^3 + a^3 + 3*b*x + 3*a)*cosh(b*x + \\
& a)*sinh(b*x + a)^3 + (b^3*x^3 + a^3 + 3*b*x + 3*a)*sinh(b*x + a)^4 + a^3 - \\
& 2*(b^3*x^3 + a^3 + 3*b*x + 3*a)*cosh(b*x + a)^2 - 2*(b^3*x^3 + a^3 - 3*(b^3 \\
& *x^3 + a^3 + 3*b*x + 3*a)*cosh(b*x + a)^2 + 3*b*x + 3*a)*sinh(b*x + a)^2 + \\
& 3*b*x + 4*((b^3*x^3 + a^3 + 3*b*x + 3*a)*cosh(b*x + a)^3 - (b^3*x^3 + a^3 + \\
& 3*b*x + 3*a)*cosh(b*x + a))*sinh(b*x + a) + 3*a)*log(-cosh(b*x + a) - sinh(\\
& b*x + a) + 1) - 24*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + si \\
& nh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a) \\
& ^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*polylog(4, cosh(\\
& b*x + a) + sinh(b*x + a)) - 24*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x \\
& + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*c \\
& osinh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*pol \\
& ylog(4, -cosh(b*x + a) - sinh(b*x + a)) + 24*(b*x*cosh(b*x + a)^4 + 4*b*x*c \\
& osinh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 - 2*b*x*cosh(b*x + a)^2 \\
& + 2*(3*b*x*cosh(b*x + a)^2 - b*x)*sinh(b*x + a)^2 + b*x + 4*(b*x*cosh(b*x + \\
& a)^3 - b*x*cosh(b*x + a))*sinh(b*x + a))*polylog(3, cosh(b*x + a) + sinh(b* \\
& x + a)) + 24*(b*x*cosh(b*x + a)^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + \\
& b*x*sinh(b*x + a)^4 - 2*b*x*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 - b* \\
& x)*sinh(b*x + a)^2 + b*x + 4*(b*x*cosh(b*x + a)^3 - b*x*cosh(b*x + a))*sinh(\\
& b*x + a))*polylog(3, -cosh(b*x + a) - sinh(b*x + a)) + 4*((b^4*x^4 - 2*a^4 \\
& + 12*b^2*x^2 - 12*a^2)*cosh(b*x + a)^3 - (b^4*x^4 - 4*b^3*x^3 - 2*a^4 + 6* \\
& b^2*x^2 - 12*a^2)*cosh(b*x + a))*sinh(b*x + a)/(b^4*cosh(b*x + a)^4 + 4*b^
\end{aligned}$$

$$4*\cosh(b*x + a)*\sinh(b*x + a)^3 + b^4*\sinh(b*x + a)^4 - 2*b^4*\cosh(b*x + a)^2 + b^4 + 2*(3*b^4*\cosh(b*x + a)^2 - b^4)*\sinh(b*x + a)^2 + 4*(b^4*\cosh(b*x + a)^3 - b^4*\cosh(b*x + a))*\sinh(b*x + a))$$

Sympy [F]

$$\int x^3 \coth^3(a + bx) dx = \int x^3 \coth^3(a + bx) dx$$

[In] `integrate(x**3*coth(b*x+a)**3,x)`
[Out] `Integral(x**3*coth(a + b*x)**3, x)`

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec), antiderivative size = 302, normalized size of antiderivative = 1.69

$$\begin{aligned} & \int x^3 \coth^3(a + bx) dx \\ &= \frac{b^2 x^4 e^{(4bx+4a)} + b^2 x^4 + 12 x^2 - 2(b^2 x^4 e^{(2a)} + 4bx^3 e^{(2a)} + 6x^2 e^{(2a)})e^{(2bx)}}{4(b^2 e^{(4bx+4a)} - 2b^2 e^{(2bx+2a)} + b^2)} - \frac{b^4 x^4 + 6b^2 x^2}{2b^4} \\ &+ \frac{b^3 x^3 \log(e^{(bx+a)} + 1) + 3b^2 x^2 \text{Li}_2(-e^{(bx+a)}) - 6bx \text{Li}_3(-e^{(bx+a)}) + 6 \text{Li}_4(-e^{(bx+a)})}{b^4} \\ &+ \frac{b^3 x^3 \log(-e^{(bx+a)} + 1) + 3b^2 x^2 \text{Li}_2(e^{(bx+a)}) - 6bx \text{Li}_3(e^{(bx+a)}) + 6 \text{Li}_4(e^{(bx+a)})}{b^4} \\ &+ \frac{3(bx \log(e^{(bx+a)} + 1) + \text{Li}_2(-e^{(bx+a)}))}{b^4} + \frac{3(bx \log(-e^{(bx+a)} + 1) + \text{Li}_2(e^{(bx+a)}))}{b^4} \end{aligned}$$

[In] `integrate(x^3*coth(b*x+a)^3,x, algorithm="maxima")`
[Out] `1/4*(b^2*x^4*e^(4*b*x + 4*a) + b^2*x^4 + 12*x^2 - 2*(b^2*x^4*e^(2*a) + 4*b*x^3*e^(2*a) + 6*x^2*e^(2*a))*e^(2*b*x))/(b^2*x^4*(4*b*x + 4*a) - 2*b^2*x^2*e^(2*b*x) + 2*a*x + b^2) - 1/2*(b^4*x^4 + 6*b^2*x^2*x^2)/b^4 + (b^3*x^3*log(e^(b*x + a) + 1) + 3*b^2*x^2*dilog(-e^(b*x + a)) - 6*b*x*polylog(3, -e^(b*x + a)) + 6*polylog(4, -e^(b*x + a)))/b^4 + (b^3*x^3*log(-e^(b*x + a) + 1) + 3*b^2*x^2*dilog(e^(b*x + a)) - 6*b*x*polylog(3, e^(b*x + a)) + 6*polylog(4, e^(b*x + a)))/b^4 + 3*(b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))/b^4 + 3*(b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))/b^4`

Giac [F]

$$\int x^3 \coth^3(a + bx) dx = \int x^3 \coth(bx + a)^3 dx$$

[In] integrate(x^3*coth(b*x+a)^3,x, algorithm="giac")
[Out] integrate(x^3*coth(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int x^3 \coth^3(a + bx) dx = \int x^3 \coth(a + b x)^3 dx$$

[In] int(x^3*coth(a + b*x)^3,x)
[Out] int(x^3*coth(a + b*x)^3, x)

3.12 $\int x^2 \coth^3(a + bx) dx$

Optimal result	92
Rubi [A] (verified)	92
Mathematica [B] (verified)	95
Maple [B] (verified)	95
Fricas [B] (verification not implemented)	96
Sympy [F]	97
Maxima [B] (verification not implemented)	97
Giac [F]	98
Mupad [F(-1)]	98

Optimal result

Integrand size = 12, antiderivative size = 114

$$\begin{aligned} \int x^2 \coth^3(a + bx) dx = & \frac{x^2}{2b} - \frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\ & + \frac{\log(\sinh(a + bx))}{b^3} + \frac{x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} \end{aligned}$$

[Out] $1/2*x^2/b - 1/3*x^3 - x*\coth(b*x+a)/b^2 - 1/2*x^2*\coth(b*x+a)^2/b + x^2*\ln(1-\exp(2*b*x+2*a))/b + \ln(\sinh(b*x+a))/b^3 + x*\operatorname{polylog}(2, \exp(2*b*x+2*a))/b^2 - 1/2*\operatorname{polylog}(3, \exp(2*b*x+2*a))/b^3$

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {3801, 3556, 30, 3797, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int x^2 \coth^3(a + bx) dx = & -\frac{\operatorname{PolyLog}(3, e^{2(a+bx)})}{2b^3} + \frac{\log(\sinh(a + bx))}{b^3} + \frac{x \operatorname{PolyLog}(2, e^{2(a+bx)})}{b^2} \\ & - \frac{x \coth(a + bx)}{b^2} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2}{2b} - \frac{x^3}{3} \end{aligned}$$

[In] $\operatorname{Int}[x^2 \operatorname{Coth}[a + b*x]^3, x]$

[Out] $x^2/(2*b) - x^3/3 - (x*\operatorname{Coth}[a + b*x])/b^2 - (x^2*\operatorname{Coth}[a + b*x]^2)/(2*b) + (x^2*\operatorname{Log}[1 - E^(2*(a + b*x))])/b + \operatorname{Log}[\operatorname{Sinh}[a + b*x]]/b^3 + (x*\operatorname{PolyLog}[2, E^(2*(a + b*x))])/b^2 - \operatorname{PolyLog}[3, E^(2*(a + b*x))]/(2*b^3)$

Rule 30

```
Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && N
eQ[m, -1]
```

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_)))^(n_.)*(c_.) + (d_.)*(x_)^(m_.))/
((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.), x_Symbol] :> Simp
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Di
st[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x))
))^n/a]], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[
{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]])]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_)^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d
*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_)^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_
.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x)
)/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_)^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
```

```
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_.)]^p]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{x^2 \coth^2(a + bx)}{2b} + \frac{\int x \coth^2(a + bx) dx}{b} + \int x^2 \coth(a + bx) dx \\
&= -\frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} - 2 \int \frac{e^{2(a+bx)} x^2}{1 - e^{2(a+bx)}} dx + \frac{\int \coth(a + bx) dx}{b^2} \\
&\quad + \frac{\int x dx}{b} \\
&= \frac{x^2}{2b} - \frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\
&\quad + \frac{\log(\sinh(a + bx))}{b^3} - \frac{2 \int x \log(1 - e^{2(a+bx)}) dx}{b} \\
&= \frac{x^2}{2b} - \frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\
&\quad + \frac{\log(\sinh(a + bx))}{b^3} + \frac{x \text{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\int \text{PolyLog}(2, e^{2(a+bx)}) dx}{b^2} \\
&= \frac{x^2}{2b} - \frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\
&\quad + \frac{\log(\sinh(a + bx))}{b^3} + \frac{x \text{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\text{Subst}\left(\int \frac{\text{PolyLog}(2, x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^3} \\
&= \frac{x^2}{2b} - \frac{x^3}{3} - \frac{x \coth(a + bx)}{b^2} - \frac{x^2 \coth^2(a + bx)}{2b} + \frac{x^2 \log(1 - e^{2(a+bx)})}{b} \\
&\quad + \frac{\log(\sinh(a + bx))}{b^3} + \frac{x \text{PolyLog}(2, e^{2(a+bx)})}{b^2} - \frac{\text{PolyLog}(3, e^{2(a+bx)})}{2b^3}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 314 vs. $2(114) = 228$.

Time = 3.54 (sec) , antiderivative size = 314, normalized size of antiderivative = 2.75

$$\int x^2 \coth^3(a + bx) dx = \frac{1}{3}x^3 \coth(a) - \frac{x^2 \operatorname{csch}^2(a + bx)}{2b}$$

$$- \frac{e^{2a}(6be^{-2a}x + 6b(1 - e^{-2a})x + 2b^3e^{-2a}x^3 - 3b^2e^{-2a}(-1 + e^{2a})x^2 \log(1 - e^{-a-bx}) - 3b^2e^{-2a}(-1 + e^{2a})x^2 \operatorname{csch}(a) \operatorname{csch}(a + bx) \sinh(bx))}{b^2}$$

[In] `Integrate[x^2*Coth[a + b*x]^3, x]`

[Out] $(x^3 \operatorname{Coth}[a])/3 - (x^2 \operatorname{Csch}[a + b*x]^2)/(2*b) - (E^{(2*a)}*((6*b*x)/E^{(2*a)} + 6*b*(1 - E^{(-2*a})*x + (2*b^3*x^3)/E^{(2*a)} - (3*b^2*(-1 + E^{(2*a})*x^2*\operatorname{Log}[1 - E^{(-a - b*x}]))/E^{(2*a)} - (3*b^2*(-1 + E^{(2*a})*x^2*\operatorname{Log}[1 + E^{(-a - b*x}]))/E^{(2*a)} - 3*(1 - E^{(-2*a})*\operatorname{Log}[1 - E^{(a + b*x}]) - 3*(1 - E^{(-2*a})*\operatorname{Log}[1 + E^{(a + b*x}]) + 6*b*(1 - E^{(-2*a})*x*\operatorname{PolyLog}[2, -E^{(-a - b*x}]) + 6*b*(1 - E^{(-2*a})*x*\operatorname{PolyLog}[2, E^{(-a - b*x}]) + 6*(1 - E^{(-2*a})*\operatorname{PolyLog}[3, -E^{(-a - b*x}]) + 6*(1 - E^{(-2*a})*\operatorname{PolyLog}[3, E^{(-a - b*x}]))/(3*b^3*(-1 + E^{(2*a})) + (x*\operatorname{Csch}[a]*\operatorname{Csch}[a + b*x]*\operatorname{Sinh}[b*x])/b^2$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 245 vs. $2(106) = 212$.

Time = 0.16 (sec) , antiderivative size = 246, normalized size of antiderivative = 2.16

method	result
risch	$-\frac{x^3}{3} - \frac{2x(e^{2bx+2a}bx+e^{2bx+2a}-1)}{b^2(e^{2bx+2a}-1)^2} + \frac{4a^3}{3b^3} + \frac{\ln(e^{bx+a}-1)}{b^3} + \frac{\ln(e^{bx+a}+1)}{b^3} - \frac{2\ln(e^{bx+a})}{b^3} - \frac{2\operatorname{polylog}(3, -e^{bx+a})}{b^3} - \frac{2\operatorname{polylog}(3, e^{bx+a})}{b^3}$

[In] `int(x^2*coth(b*x+a)^3, x, method=_RETURNVERBOSE)`

[Out] $-1/3*x^3-2*x*(\exp(2*b*x+2*a)*b*x+\exp(2*b*x+2*a)-1)/b^2/(\exp(2*b*x+2*a)-1)^2 + 4/3/b^3*a^3+1/b^3*\ln(\exp(b*x+a)-1)+1/b^3*\ln(\exp(b*x+a)+1)-2/b^3*\ln(\exp(b*x+a))-2/b^3*\operatorname{polylog}(3, -\exp(b*x+a))-2/b^3*\operatorname{polylog}(3, \exp(b*x+a))+1/b*\ln(1-\exp(b*x+a))*x^2+2/b^2*\operatorname{polylog}(2, \exp(b*x+a))*x+1/b*\ln(\exp(b*x+a)+1)*x^2+2/b^2*\operatorname{polylog}(2, -\exp(b*x+a))*x+2/b^2*a^2*x+1/b^3*a^2*\ln(\exp(b*x+a)-1)-2/b^3*a^2*\ln(\exp(b*x+a))-1/b^3*\ln(1-\exp(b*x+a))*a^2$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1467 vs. $2(105) = 210$.

Time = 0.28 (sec), antiderivative size = 1467, normalized size of antiderivative = 12.87

$$\int x^2 \coth^3(a + bx) dx = \text{Too large to display}$$

```
[In] integrate(x^2*coth(b*x+a)^3,x, algorithm="fricas")

[Out] -1/3*(b^3*x^3 + (b^3*x^3 + 2*a^3 + 6*b*x + 6*a)*cosh(b*x + a)^4 + 4*(b^3*x^3 + 2*a^3 + 6*b*x + 6*a)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^3*x^3 + 2*a^3 + 6*b*x + 6*a)*sinh(b*x + a)^4 + 2*a^3 - 2*(b^3*x^3 - 3*b^2*x^2 + 2*a^3 + 3*b*x + 6*a)*cosh(b*x + a)^2 - 2*(b^3*x^3 - 3*b^2*x^2 + 2*a^3 - 3*(b^3*x^3 + 2*a^3 + 6*b*x + 6*a)*cosh(b*x + a)^2 + 3*b*x + 6*a)*sinh(b*x + a)^2 - 6*(b*x*cosh(b*x + a)^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 - 2*b*x*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 - b*x)*sinh(b*x + a)^2 + b*x + 4*(b*x*cosh(b*x + a)^3 - b*x*cosh(b*x + a))*sinh(b*x + a))*dilog(cosh(b*x + a) + sinh(b*x + a)) - 6*(b*x*cosh(b*x + a)^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 - 2*b*x*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 - b*x)*sinh(b*x + a)^2 + b*x + 4*(b*x*cosh(b*x + a)^3 - b*x*cosh(b*x + a))*sinh(b*x + a))*dilog(-cosh(b*x + a) - sinh(b*x + a)) - 3*((b^2*x^2 + 1)*cosh(b*x + a)^4 + 4*(b^2*x^2 + 1)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^2*x^2 + 1)*sinh(b*x + a)^4 + b^2*x^2 - 2*(b^2*x^2 + 1)*cosh(b*x + a)^2 - 2*(b^2*x^2 - 3*(b^2*x^2 + 1)*cosh(b*x + a)^2 + 1)*sinh(b*x + a)^2 + 4*((b^2*x^2 + 1)*cosh(b*x + a)^3 - (b^2*x^2 + 1)*cosh(b*x + a))*sinh(b*x + a) + 1)*log(cosh(b*x + a) + sinh(b*x + a) + 1) - 3*((a^2 + 1)*cosh(b*x + a)^4 + 4*(a^2 + 1)*cosh(b*x + a)*sinh(b*x + a)^3 + (a^2 + 1)*sinh(b*x + a)^4 - 2*(a^2 + 1)*cosh(b*x + a)^2 + 2*(3*(a^2 + 1)*cosh(b*x + a)^2 - a^2 - 1)*sinh(b*x + a)^2 + a^2 + 4*((a^2 + 1)*cosh(b*x + a)^3 - (a^2 + 1)*cosh(b*x + a))*sinh(b*x + a) + 1)*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 3*((b^2*x^2 - a^2)*cosh(b*x + a)^4 + 4*(b^2*x^2 - a^2)*cosh(b*x + a)*sinh(b*x + a)^3 + (b^2*x^2 - a^2)*sinh(b*x + a)^4 + b^2*x^2 - 2*(b^2*x^2 - a^2)*cosh(b*x + a)^2 - 2*(b^2*x^2 - a^2)*cosh(b*x + a)^2 - 2*(b^2*x^2 - 3*(b^2*x^2 - a^2)*cosh(b*x + a)^2 - a^2)*sinh(b*x + a)^2 - a^2 + 4*((b^2*x^2 - a^2)*cosh(b*x + a)^3 - (b^2*x^2 - a^2)*cosh(b*x + a))*sinh(b*x + a))*log(-cosh(b*x + a) - sinh(b*x + a) + 1) + 6*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*polylog(3, cosh(b*x + a) + sinh(b*x + a)) + 6*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*polylog(3, -cosh(b*x + a) - sinh(b*x + a)) + 4*((b^3*x^3 + 2*a^3 + 6*b*x + 6*a)*cosh(b*x + a)^3 - (b^3*x^3 - 3*b^2*x^2 + 2*a^3 + 3*b*x + 6*a)*cosh(b*x + a))*sinh(b*x + a) + 6*a)/(b^3*cosh(b*x + a)^4 + 4*b^3*cosh(b*x + a)*sinh(b*x + a)^3 + b^3*sinh(b*x + a)^4 - 2*b
```

$$\begin{aligned} & -3 \cosh(b*x + a)^2 + b^3 + 2*(3*b^3 \cosh(b*x + a)^2 - b^3) * \sinh(b*x + a)^2 \\ & + 4*(b^3 \cosh(b*x + a)^3 - b^3 \cosh(b*x + a)) * \sinh(b*x + a) \end{aligned}$$

Sympy [F]

$$\int x^2 \coth^3(a + bx) dx = \int x^2 \coth^3(a + bx) dx$$

[In] `integrate(x**2*coth(b*x+a)**3,x)`
[Out] `Integral(x**2*coth(a + b*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 226 vs. $2(105) = 210$.

Time = 0.25 (sec), antiderivative size = 226, normalized size of antiderivative = 1.98

$$\begin{aligned} & \int x^2 \coth^3(a + bx) dx \\ &= -\frac{2}{3} x^3 + \frac{b^2 x^3 e^{(4bx+4a)} + b^2 x^3 - 2(b^2 x^3 e^{(2a)} + 3bx^2 e^{(2a)} + 3xe^{(2a)})e^{(2bx)} + 6x}{3(b^2 e^{(4bx+4a)} - 2b^2 e^{(2bx+2a)} + b^2)} \\ & - \frac{2x}{b^2} + \frac{b^2 x^2 \log(e^{(bx+a)} + 1) + 2bx \text{Li}_2(-e^{(bx+a)}) - 2 \text{Li}_3(-e^{(bx+a)})}{b^3} \\ & + \frac{b^2 x^2 \log(-e^{(bx+a)} + 1) + 2bx \text{Li}_2(e^{(bx+a)}) - 2 \text{Li}_3(e^{(bx+a)})}{b^3} \\ & + \frac{\log(e^{(bx+a)} + 1)}{b^3} + \frac{\log(e^{(bx+a)} - 1)}{b^3} \end{aligned}$$

[In] `integrate(x^2*coth(b*x+a)^3,x, algorithm="maxima")`
[Out]
$$\begin{aligned} & -2/3*x^3 + 1/3*(b^2*x^3*e^{(4*b*x + 4*a)} + b^2*x^3 - 2*(b^2*x^3*e^{(2*a)} + 3*b*x^2*e^{(2*a)} + 3*x*e^{(2*a)})*e^{(2*b*x)} + 6*x)/(b^2*x^2*e^{(4*b*x + 4*a)} - 2*b^2*x^2*e^{(2*b*x + 2*a)} + b^2) - 2*x/b^2 + (b^2*x^2*\log(e^{(b*x + a)} + 1) + 2*b*x*\text{dilog}(-e^{(b*x + a)})) - 2*\text{polylog}(3, -e^{(b*x + a)}))/b^3 + (b^2*x^2*\log(-e^{(b*x + a)} + 1) + 2*b*x*\text{dilog}(e^{(b*x + a)})) - 2*\text{polylog}(3, e^{(b*x + a)}))/b^3 + \log(e^{(b*x + a)} + 1)/b^3 + \log(e^{(b*x + a)} - 1)/b^3 \end{aligned}$$

Giac [F]

$$\int x^2 \coth^3(a + bx) dx = \int x^2 \coth(bx + a)^3 dx$$

[In] integrate(x^2*coth(b*x+a)^3,x, algorithm="giac")
 [Out] integrate(x^2*coth(b*x + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int x^2 \coth^3(a + bx) dx = \int x^2 \coth(a + b x)^3 dx$$

[In] int(x^2*coth(a + b*x)^3,x)
 [Out] int(x^2*coth(a + b*x)^3, x)

3.13 $\int x \coth^3(a + bx) dx$

Optimal result	99
Rubi [A] (verified)	99
Mathematica [A] (verified)	101
Maple [B] (verified)	101
Fricas [B] (verification not implemented)	102
Sympy [F]	103
Maxima [B] (verification not implemented)	103
Giac [F]	103
Mupad [F(-1)]	104

Optimal result

Integrand size = 10, antiderivative size = 82

$$\begin{aligned} \int x \coth^3(a + bx) dx = & \frac{x}{2b} - \frac{x^2}{2} - \frac{\coth(a + bx)}{2b^2} - \frac{x \coth^2(a + bx)}{2b} \\ & + \frac{x \log(1 - e^{2(a+bx)})}{b} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2} \end{aligned}$$

[Out] $\frac{1}{2}x/b - \frac{1}{2}x^2 - \frac{1}{2}\coth(b*x+a)/b^2 - \frac{1}{2}x*\coth(b*x+a)^2/b + x*\ln(1-\exp(2*b*x+2*a))/b + \frac{1}{2}\text{polylog}(2, \exp(2*b*x+2*a))/b^2$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.700$, Rules used = {3801, 3554, 8, 3797, 2221, 2317, 2438}

$$\begin{aligned} \int x \coth^3(a + bx) dx = & \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2} - \frac{\coth(a + bx)}{2b^2} \\ & + \frac{x \log(1 - e^{2(a+bx)})}{b} - \frac{x \coth^2(a + bx)}{2b} + \frac{x}{2b} - \frac{x^2}{2} \end{aligned}$$

[In] Int[x*Coth[a + b*x]^3, x]

[Out] $x/(2*b) - x^2/2 - \text{COTH}[a + b*x]/(2*b^2) - (x*\text{Coth}[a + b*x]^2)/(2*b) + (x*\text{Log}[1 - E^(2*(a + b*x))])/b + \text{PolyLog}[2, E^(2*(a + b*x))]/(2*b^2)$

Rule 8

Int[a_, x_Symbol] :> Simplify[a*x, x] /; FreeQ[a, x]

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_.) + (b_.)*(F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^(m_)*tan[(e_.) + Pi*(k_.)*(f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = -\frac{x \coth^2(a + bx)}{2b} + \frac{\int \coth^2(a + bx) dx}{2b} + \int x \coth(a + bx) dx$$

$$\begin{aligned}
&= -\frac{x^2}{2} - \frac{\coth(a+bx)}{2b^2} - \frac{x \coth^2(a+bx)}{2b} - 2 \int \frac{e^{2(a+bx)}x}{1-e^{2(a+bx)}} dx + \frac{\int 1 dx}{2b} \\
&= \frac{x}{2b} - \frac{x^2}{2} - \frac{\coth(a+bx)}{2b^2} - \frac{x \coth^2(a+bx)}{2b} + \frac{x \log(1-e^{2(a+bx)})}{b} - \frac{\int \log(1-e^{2(a+bx)}) dx}{b} \\
&= \frac{x}{2b} - \frac{x^2}{2} - \frac{\coth(a+bx)}{2b^2} - \frac{x \coth^2(a+bx)}{2b} \\
&\quad + \frac{x \log(1-e^{2(a+bx)})}{b} - \frac{\text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(a+bx)}\right)}{2b^2} \\
&= \frac{x}{2b} - \frac{x^2}{2} - \frac{\coth(a+bx)}{2b^2} - \frac{x \coth^2(a+bx)}{2b} + \frac{x \log(1-e^{2(a+bx)})}{b} + \frac{\text{PolyLog}(2, e^{2(a+bx)})}{2b^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.53 (sec), antiderivative size = 131, normalized size of antiderivative = 1.60

$$\int x \coth^3(a+bx) dx = \frac{1}{2} \left(-\frac{2x^2}{-1+e^{2a}} + x^2 \coth(a) - \frac{x \operatorname{csch}^2(a+bx)}{b} + \frac{2x \log(1-e^{-a-bx})}{b} \right. \\
\left. + \frac{2x \log(1+e^{-a-bx})}{b} - \frac{2 \operatorname{PolyLog}(2, -e^{-a-bx})}{b^2} \right. \\
\left. - \frac{2 \operatorname{PolyLog}(2, e^{-a-bx})}{b^2} + \frac{\operatorname{csch}(a) \operatorname{csch}(a+bx) \sinh(bx)}{b^2} \right)$$

[In] `Integrate[x*Coth[a + b*x]^3, x]`

[Out] `((-2*x^2)/(-1 + E^(2*a)) + x^2*Coth[a] - (x*Csch[a + b*x]^2)/b + (2*x*Log[1 - E^(-a - b*x)])/b + (2*x*Log[1 + E^(-a - b*x)])/b - (2*PolyLog[2, -E^(-a - b*x)])/b^2 - (2*PolyLog[2, E^(-a - b*x)])/b^2 + (Csch[a]*Csch[a + b*x]*Sinh[b*x])/b^2)/2`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 163 vs. $2(72) = 144$.

Time = 0.14 (sec), antiderivative size = 164, normalized size of antiderivative = 2.00

method	result
risch	$-\frac{x^2}{2} - \frac{2e^{2bx+2a}bx+e^{2bx+2a}-1}{b^2(e^{2bx+2a}-1)^2} - \frac{2ax}{b} - \frac{a^2}{b^2} + \frac{\ln(e^{bx+a}+1)x}{b} + \frac{\text{polylog}(2, -e^{bx+a})}{b^2} + \frac{\ln(1-e^{bx+a})x}{b} + \frac{\ln(1-e^{bx+a})a}{b^2}$

[In] `int(x*coth(b*x+a)^3, x, method=_RETURNVERBOSE)`

[Out]
$$\frac{-1/2*x^2 - (2*\exp(2*b*x+2*a)*b*x + \exp(2*b*x+2*a)-1)/b^2/(exp(2*b*x+2*a)-1)^2 - 2/b*a*x - a^2/b^2 + 1/b*\ln(\exp(b*x+a)+1)*x + 1/b^2*polylog(2, -\exp(b*x+a)) + 1/b*\ln(1 - \exp(b*x+a))*x + 1/b^2*\ln(1 - \exp(b*x+a))*a + 1/b^2*polylog(2, \exp(b*x+a)) - 1/b^2*a*\ln(\exp(b*x+a) - 1) + 2/b^2*a*\ln(\exp(b*x+a))}{b^2}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 975 vs. $2(71) = 142$.

Time = 0.32 (sec) , antiderivative size = 975, normalized size of antiderivative = 11.89

$$\int x \coth^3(a + bx) dx = \text{Too large to display}$$

[In] `integrate(x*coth(b*x+a)^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -1/2*((b^2*x^2 - 2*a^2)*cosh(b*x + a)^4 + 4*(b^2*x^2 - 2*a^2)*cosh(b*x + a) \\ & *sinh(b*x + a)^3 + (b^2*x^2 - 2*a^2)*sinh(b*x + a)^4 + b^2*x^2 - 2*(b^2*x^2 \\ & - 2*a^2 - 2*b*x - 1)*cosh(b*x + a)^2 - 2*(b^2*x^2 - 3*(b^2*x^2 - 2*a^2)*co \\ & sh(b*x + a)^2 - 2*a^2 - 2*b*x - 1)*sinh(b*x + a)^2 - 2*a^2 - 2*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*dilog(cosh(b*x + a) + sinh(b*x + a)) - 2*(cosh(b*x + a)^4 + 4*cosh(b*x + a)*sinh(b*x + a)^3 + sinh(b*x + a)^4 + 2*(3*cosh(b*x + a)^2 - 1)*sinh(b*x + a)^2 - 2*cosh(b*x + a)^2 + 4*(cosh(b*x + a)^3 - cosh(b*x + a))*sinh(b*x + a) + 1)*dilog(-cosh(b*x + a) - sinh(b*x + a)) - 2*(b*x*cosh(b*x + a)^4 + 4*b*x*cosh(b*x + a)*sinh(b*x + a)^3 + b*x*sinh(b*x + a)^4 - 2*b*x*cosh(b*x + a)^2 + 2*(3*b*x*cosh(b*x + a)^2 - b*x)*sinh(b*x + a)^2 + b*x + 4*(b*x*cosh(b*x + a)^3 - b*x*cosh(b*x + a))*sinh(b*x + a))*log(cosh(b*x + a) + sinh(b*x + a) + 1) + 2*(a*cosh(b*x + a)^4 + 4*a*cosh(b*x + a)*sinh(b*x + a)^3 + a*sinh(b*x + a)^4 - 2*a*cosh(b*x + a)^2 + 2*(3*a*cosh(b*x + a)^2 - a)*sinh(b*x + a)^2 + 4*(a*cosh(b*x + a)^3 - a*cosh(b*x + a)) \\ & *sinh(b*x + a) + a)*log(cosh(b*x + a) + sinh(b*x + a) - 1) - 2*((b*x + a)*c \\ & osinh(b*x + a)^4 + 4*(b*x + a)*cosh(b*x + a)*sinh(b*x + a)^3 + (b*x + a)*sinh(b*x + a)^4 - 2*(b*x + a)*cosh(b*x + a)^2 + 2*(3*(b*x + a)*cosh(b*x + a)^2 - b*x - a)*sinh(b*x + a)^2 + b*x + 4*((b*x + a)*cosh(b*x + a)^3 - (b*x + a)*cosh(b*x + a))*sinh(b*x + a) + a)*log(-cosh(b*x + a) - sinh(b*x + a) + 1) + 4*((b^2*x^2 - 2*a^2)*cosh(b*x + a)^3 - (b^2*x^2 - 2*a^2 - 2*b*x - 1)*cosh(b*x + a)*sinh(b*x + a) - 2)/(b^2*cosh(b*x + a)^4 + 4*b^2*cosh(b*x + a)*sinh(b*x + a)^3 + b^2*sinh(b*x + a)^4 - 2*b^2*cosh(b*x + a)^2 + 2*(3*b^2*cosh(b*x + a)^2 - b^2)*sinh(b*x + a)^2 + b^2 + 4*(b^2*cosh(b*x + a)^3 - b^2*cosh(b*x + a))*sinh(b*x + a)) \end{aligned}$$

Sympy [F]

$$\int x \coth^3(a + bx) dx = \int x \coth^3(a + bx) dx$$

[In] `integrate(x*coth(b*x+a)**3,x)`

[Out] `Integral(x*coth(a + b*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(71) = 142$.

Time = 0.24 (sec) , antiderivative size = 149, normalized size of antiderivative = 1.82

$$\begin{aligned} \int x \coth^3(a + bx) dx &= -x^2 \\ &+ \frac{b^2 x^2 e^{(4bx+4a)} + b^2 x^2 - 2(b^2 x^2 e^{(2a)} + 2bx e^{(2a)} + e^{(2a)})e^{(2bx)} + 2}{2(b^2 e^{(4bx+4a)} - 2b^2 e^{(2bx+2a)} + b^2)} \\ &+ \frac{bx \log(e^{(bx+a)} + 1) + \text{Li}_2(-e^{(bx+a)})}{b^2} \\ &+ \frac{bx \log(-e^{(bx+a)} + 1) + \text{Li}_2(e^{(bx+a)})}{b^2} \end{aligned}$$

[In] `integrate(x*coth(b*x+a)^3,x, algorithm="maxima")`

[Out] `-x^2 + 1/2*(b^2*x^2*e^(4*b*x + 4*a) + b^2*x^2 - 2*(b^2*x^2*e^(2*a) + 2*b*x*e^(2*a) + e^(2*a))*e^(2*b*x) + 2)/(b^2*e^(4*b*x + 4*a) - 2*b^2*e^(2*b*x + 2*a) + b^2) + (b*x*log(e^(b*x + a) + 1) + dilog(-e^(b*x + a)))/b^2 + (b*x*log(-e^(b*x + a) + 1) + dilog(e^(b*x + a)))/b^2`

Giac [F]

$$\int x \coth^3(a + bx) dx = \int x \coth(bx + a)^3 dx$$

[In] `integrate(x*coth(b*x+a)^3,x, algorithm="giac")`

[Out] `integrate(x*coth(b*x + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int x \coth^3(a + bx) dx = \int x \coth(a + bx)^3 dx$$

[In] int(x*coth(a + b*x)^3,x)

[Out] int(x*coth(a + b*x)^3, x)

3.14 $\int \frac{\coth^3(a+bx)}{x} dx$

Optimal result	105
Rubi [N/A]	105
Mathematica [N/A]	106
Maple [N/A] (verified)	106
Fricas [N/A]	106
Sympy [N/A]	106
Maxima [N/A]	107
Giac [N/A]	107
Mupad [N/A]	107

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\coth^3(a + bx)}{x} dx = \text{Int}\left(\frac{\coth^3(a + bx)}{x}, x\right)$$

[Out] Unintegrable($\coth(b*x+a)^3/x$, x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth^3(a + bx)}{x} dx$$

[In] Int[Coth[a + b*x]^3/x, x]

[Out] Defer[Int][Coth[a + b*x]^3/x, x]

Rubi steps

$$\text{integral} = \int \frac{\coth^3(a + bx)}{x} dx$$

Mathematica [N/A]

Not integrable

Time = 21.67 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth^3(a + bx)}{x} dx$$

[In] `Integrate[Coth[a + b*x]^3/x, x]`

[Out] `Integrate[Coth[a + b*x]^3/x, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)^3}{x} dx$$

[In] `int(coth(b*x+a)^3/x, x)`

[Out] `int(coth(b*x+a)^3/x, x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth(bx + a)^3}{x} dx$$

[In] `integrate(coth(b*x+a)^3/x, x, algorithm="fricas")`

[Out] `integral(coth(b*x + a)^3/x, x)`

Sympy [N/A]

Not integrable

Time = 0.60 (sec) , antiderivative size = 10, normalized size of antiderivative = 0.83

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth^3(a + bx)}{x} dx$$

[In] `integrate(coth(b*x+a)**3/x, x)`

[Out] `Integral(coth(a + b*x)**3/x, x)`

Maxima [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 144, normalized size of antiderivative = 12.00

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth(bx + a)^3}{x} dx$$

[In] `integrate(coth(b*x+a)^3/x,x, algorithm="maxima")`

[Out] `-((2*b*x*e^(2*a) - e^(2*a))*e^(2*b*x) + 1)/(b^2*x^2*e^(4*b*x + 4*a) - 2*b^2*x^2*e^(2*b*x + 2*a) + b^2*x^2) - integrate((b^2*x^2 + 1)/(b^2*x^3*e^(b*x + a) + b^2*x^3), x) + integrate((b^2*x^2 + 1)/(b^2*x^3*e^(b*x + a) - b^2*x^3), x) + log(x)`

Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth(bx + a)^3}{x} dx$$

[In] `integrate(coth(b*x+a)^3/x,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)^3/x, x)`

Mupad [N/A]

Not integrable

Time = 1.81 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x} dx = \int \frac{\coth(a + b x)^3}{x} dx$$

[In] `int(coth(a + b*x)^3/x,x)`

[Out] `int(coth(a + b*x)^3/x, x)`

3.15 $\int \frac{\coth^3(a+bx)}{x^2} dx$

Optimal result	108
Rubi [N/A]	108
Mathematica [N/A]	109
Maple [N/A] (verified)	109
Fricas [N/A]	109
Sympy [N/A]	109
Maxima [N/A]	110
Giac [N/A]	110
Mupad [N/A]	110

Optimal result

Integrand size = 12, antiderivative size = 12

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \text{Int}\left(\frac{\coth^3(a + bx)}{x^2}, x\right)$$

[Out] Unintegrable(coth(b*x+a)^3/x^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth^3(a + bx)}{x^2} dx$$

[In] Int[Coth[a + b*x]^3/x^2,x]

[Out] Defer[Int][Coth[a + b*x]^3/x^2, x]

Rubi steps

$$\text{integral} = \int \frac{\coth^3(a + bx)}{x^2} dx$$

Mathematica [N/A]

Not integrable

Time = 15.90 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth^3(a + bx)}{x^2} dx$$

```
[In] Integrate[Coth[a + b*x]^3/x^2,x]
[Out] Integrate[Coth[a + b*x]^3/x^2, x]
```

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth(bx + a)^3}{x^2} dx$$

```
[In] int(coth(b*x+a)^3/x^2,x)
[Out] int(coth(b*x+a)^3/x^2,x)
```

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^3}{x^2} dx$$

```
[In] integrate(coth(b*x+a)^3/x^2,x, algorithm="fricas")
[Out] integral(coth(b*x + a)^3/x^2, x)
```

Sympy [N/A]

Not integrable

Time = 0.54 (sec) , antiderivative size = 12, normalized size of antiderivative = 1.00

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth^3(a + bx)}{x^2} dx$$

```
[In] integrate(coth(b*x+a)**3/x**2,x)
[Out] Integral(coth(a + b*x)**3/x**2, x)
```

Maxima [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 175, normalized size of antiderivative = 14.58

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^3}{x^2} dx$$

[In] `integrate(coth(b*x+a)^3/x^2,x, algorithm="maxima")`

[Out] $-\frac{(b^2 x^2 e^{(4 b x+4 a)}+b^2 x^2-2 (b^2 x^2 e^{(2 a)}-b x e^{(2 a)}) e^{(2 b x)}+2)/(b^2 x^3 e^{(4 b x+4 a)}-2 b^2 x^3 e^{(2 b x+2 a)}+b^2 x^3)-\text{integrate}((b^2 x^2+3)/(b^2 x^4 e^{(b x+a)}+b^2 x^4),x)+\text{integrate}((b^2 x^2+3)/(b^2 x^4 e^{(b x+a)}-b^2 x^4),x)$

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth(bx + a)^3}{x^2} dx$$

[In] `integrate(coth(b*x+a)^3/x^2,x, algorithm="giac")`

[Out] `integrate(coth(b*x + a)^3/x^2, x)`

Mupad [N/A]

Not integrable

Time = 1.82 (sec) , antiderivative size = 14, normalized size of antiderivative = 1.17

$$\int \frac{\coth^3(a + bx)}{x^2} dx = \int \frac{\coth(a + b x)^3}{x^2} dx$$

[In] `int(coth(a + b*x)^3/x^2,x)`

[Out] `int(coth(a + b*x)^3/x^2, x)`

3.16 $\int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx$

Optimal result	111
Rubi [A] (verified)	111
Mathematica [A] (verified)	113
Maple [A] (verified)	113
Fricas [A] (verification not implemented)	114
Sympy [B] (verification not implemented)	114
Maxima [A] (verification not implemented)	115
Giac [A] (verification not implemented)	115
Mupad [B] (verification not implemented)	116

Optimal result

Integrand size = 20, antiderivative size = 169

$$\begin{aligned} \int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx = & \frac{3d^3x}{8af^3} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} \\ & - \frac{3d^3}{8f^4(a+a \coth(e+fx))} - \frac{3d^2(c+dx)}{4f^3(a+a \coth(e+fx))} \\ & - \frac{3d(c+dx)^2}{4f^2(a+a \coth(e+fx))} - \frac{(c+dx)^3}{2f(a+a \coth(e+fx))} \end{aligned}$$

[Out] $3/8*d^3*x/a/f^3+3/8*d*(d*x+c)^2/a/f^2+1/4*(d*x+c)^3/a/f+1/8*(d*x+c)^4/a/d-3/8*d^3/f^4/(a+a*\coth(f*x+e))-3/4*d^2*(d*x+c)/f^3/(a+a*\coth(f*x+e))-3/4*d*(d*x+c)^2/f^2/(a+a*\coth(f*x+e))-1/2*(d*x+c)^3/f/(a+a*\coth(f*x+e))$

Rubi [A] (verified)

Time = 0.14 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3804, 3560, 8}

$$\begin{aligned} \int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx = & -\frac{3d^2(c+dx)}{4f^3(a \coth(e+fx)+a)} - \frac{3d(c+dx)^2}{4f^2(a \coth(e+fx)+a)} \\ & - \frac{(c+dx)^3}{2f(a \coth(e+fx)+a)} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} \\ & + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a \coth(e+fx)+a)} + \frac{3d^3x}{8af^3} \end{aligned}$$

[In] Int[(c + d*x)^3/(a + a*Coth[e + f*x]), x]

[Out] $(3*d^3*x)/(8*a*f^3) + (3*d*(c + d*x)^2)/(8*a*f^2) + (c + d*x)^3/(4*a*f) + (c + d*x)^4/(8*a*d) - (3*d^3)/(8*f^4*(a + a*\text{Coth}[e + f*x])) - (3*d^2*(c + d*x))/(4*f^3*(a + a*\text{Coth}[e + f*x])) - (3*d*(c + d*x)^2)/(4*f^2*(a + a*\text{Coth}[e + f*x])) - (c + d*x)^3/(2*f*(a + a*\text{Coth}[e + f*x]))$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n)/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3804

Int[((c_.) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m)/(2*b*f*(a + b*Tan[e + f*x])), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(c+dx)^4}{8ad} - \frac{(c+dx)^3}{2f(a+a\coth(e+fx))} + \frac{(3d)\int \frac{(c+dx)^2}{a+a\coth(e+fx)} dx}{2f} \\
 &= \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d(c+dx)^2}{4f^2(a+a\coth(e+fx))} \\
 &\quad - \frac{(c+dx)^3}{2f(a+a\coth(e+fx))} + \frac{(3d^2)\int \frac{c+dx}{a+a\coth(e+fx)} dx}{2f^2} \\
 &= \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^2(c+dx)}{4f^3(a+a\coth(e+fx))} \\
 &\quad - \frac{3d(c+dx)^2}{4f^2(a+a\coth(e+fx))} - \frac{(c+dx)^3}{2f(a+a\coth(e+fx))} + \frac{(3d^3)\int \frac{1}{a+a\coth(e+fx)} dx}{4f^3} \\
 &= \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+a\coth(e+fx))} \\
 &\quad - \frac{3d^2(c+dx)}{4f^3(a+a\coth(e+fx))} - \frac{3d(c+dx)^2}{4f^2(a+a\coth(e+fx))} \\
 &\quad - \frac{(c+dx)^3}{2f(a+a\coth(e+fx))} + \frac{(3d^3)\int 1 dx}{8af^3}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3d^3x}{8af^3} + \frac{3d(c+dx)^2}{8af^2} + \frac{(c+dx)^3}{4af} + \frac{(c+dx)^4}{8ad} - \frac{3d^3}{8f^4(a+a\coth(e+fx))} \\
&\quad - \frac{3d^2(c+dx)}{4f^3(a+a\coth(e+fx))} - \frac{3d(c+dx)^2}{4f^2(a+a\coth(e+fx))} - \frac{(c+dx)^3}{2f(a+a\coth(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.93 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.44

$$\begin{aligned}
&\int \frac{(c+dx)^3}{a+a\coth(e+fx)} dx \\
&= \frac{\text{csch}(e+fx)(\cosh(fx)+\sinh(fx)) ((4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2)+d^3(3+6fx- \\
&\quad -2f^2x^2)) * (\cosh(e)+\sinh(e)) + (4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2)+d^3(3+6fx- \\
&\quad -2f^2x^2)) * (-\cosh(e)+\sinh(e)))}{16a*f^4*(1+\coth(e+fx))}
\end{aligned}$$

[In] Integrate[(c + d*x)^3/(a + a*Coth[e + f*x]), x]

[Out] $(\text{Csch}[e+f*x]*(\text{Cosh}[f*x]+\text{Sinh}[f*x])*((4*c^3*f^3+6*c^2*d*f^2*(1+2*f*x) \\
+6*c*d^2*f*(1+2*f*x+2*f^2*x^2)+d^3*(3+6*f*x+6*f^2*x^2+4*f^3*x^3))*\text{Cosh}[2*f*x]*(\text{Cosh}[e]-\text{Sinh}[e]) \\
+2*f^4*x*(4*c^3+6*c^2*d*x+4*c*d^2*x^2+d^3*x^3)*(Cosh[e]+\text{Sinh}[e])+(4*c^3*f^3+6*c^2*d*f^2*(1+2*f*x) \\
+6*c*d^2*f*(1+2*f*x+2*f^2*x^2)+d^3*(3+6*f*x+6*f^2*x^2+4*f^3*x^3))*(-\text{Cosh}[e]+\text{Sinh}[e])*\text{Sinh}[2*f*x]))/(16*a*f^4*(1+\text{Coth}[e+f*x]))$

Maple [A] (verified)

Time = 0.38 (sec) , antiderivative size = 165, normalized size of antiderivative = 0.98

method	result
risch	$\frac{d^3x^4}{8a} + \frac{d^2cx^3}{2a} + \frac{3dc^2x^2}{4a} + \frac{c^3x}{2a} + \frac{c^4}{8ad} + \frac{(4d^3x^3f^3+12cd^2f^3x^2+12c^2df^3x+6d^3f^2x^2+4c^3f^3+12cd^2f^2x+6c^2f^4)}{16af^4}$
parallelrisch	$((d^3x^4+4d^2cx^3+6dc^2x^2+4c^3x)f^4+(-2d^3x^3-6cd^2x^2-6c^2dx-4c^3)f^3+(-3d^3x^2-6cd^2x-6c^2)f^2+(-3d^3x-6d^2c))$ $8f^4a(1+\tanh(fx))$
derivativedivides	Expression too large to display
default	Expression too large to display

[In] int((d*x+c)^3/(a+a*coth(f*x+e)), x, method=_RETURNVERBOSE)

[Out] $1/8/a*d^3*x^4+1/2/a*d^2*c*x^3+3/4/a*d*c^2*x^2+1/2/a*c^3*x+1/8/a/d*c^4+1/16*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4*c^3*f^3+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f^3*x+6*c*d^2*f^3+3*d^3*f^2*x+3*d^3)/a/f^4*\exp(-2*f*x-2*e)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec), antiderivative size = 304, normalized size of antiderivative = 1.80

$$\int \frac{(c+dx)^3}{a+a\coth(e+fx)} dx \\ \equiv (2d^3f^4x^4 + 4c^3f^3 + 6c^2df^2 + 6cd^2f + 4(2cd^2f^4 + d^3f^3)x^3 + 3d^3 + 6(2c^2df^4 + 2cd^2f^3 + d^3f^2)x^2 + 2(4$$

```
[In] integrate((d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="fricas")
[Out] 1/16*((2*d^3*f^4*x^4 + 4*c^3*f^3 + 6*c^2*d*f^2 + 6*c*d^2*f + 4*(2*c*d^2*f^4 + d^3*f^3)*x^3 + 3*d^3 + 6*(2*c^2*d*f^4 + 2*c*d^2*f^3 + d^3*f^2)*x^2 + 2*(4*c^3*f^4 + 6*c^2*d*f^3 + 6*c*d^2*f^2 + 3*d^3*f)*x)*cosh(f*x + e) + (2*d^3*f^4*x^4 - 4*c^3*f^3 - 6*c^2*d*f^2 - 6*c*d^2*f + 4*(2*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 6*(2*c^2*d*f^4 - 2*c*d^2*f^3 - d^3*f^2)*x^2 + 2*(4*c^3*f^4 - 6*c^2*d*f^3 - 6*c*d^2*f^2 - 3*d^3*f)*x)*sinh(f*x + e))/(a*f^4*cosh(f*x + e) + a*f^4*sinh(f*x + e))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 864 vs. $2(144) = 288$.

Time = 0.76 (sec) , antiderivative size = 864, normalized size of antiderivative = 5.11

$$\int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx$$

$$= \begin{cases} \frac{4c^3 f^4 x \tanh(e+fx)}{8af^4 \tanh(e+fx)+8af^4} + \frac{4c^3 f^4 x}{8af^4 \tanh(e+fx)+8af^4} + \frac{4c^3 f^3}{8af^4 \tanh(e+fx)+8af^4} + \frac{6c^2 df^4 x^2 \tanh(e+fx)}{8af^4 \tanh(e+fx)+8af^4} + \frac{6c^2 df^4 x^2}{8af^4 \tanh(e+fx)+8af^4} \\ \frac{c^3 x + \frac{3c^2 dx^2}{2} + cd^2 x^3 + \frac{d^3 x^4}{4}}{a \coth(e)+a} \end{cases}$$

[In] `integrate((d*x+c)**3/(a+a*cOTH(f*x+e)),x)`

```
[Out] Piecewise((4*c**3*f**4*x*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4)
+ 4*c**3*f**4*x/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 4*c**3*f**3/(8*a*f**4
*tanh(e + f*x) + 8*a*f**4) + 6*c**2*d*f**4*x**2*tanh(e + f*x)/(8*a*f**4*tan
h(e + f*x) + 8*a*f**4) + 6*c**2*d*f**4*x**2/(8*a*f**4*tanh(e + f*x) + 8*a*f
**4) - 6*c**2*d*f**3*x*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) +
6*c**2*d*f**3*x/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 6*c**2*d*f**2/(8*a*f*
**4*tanh(e + f*x) + 8*a*f**4) + 4*c*d**2*f**4*x**3*tanh(e + f*x)/(8*a*f**4*t
anh(e + f*x) + 8*a*f**4) + 4*c*d**2*f**4*x**3/(8*a*f**4*tanh(e + f*x) + 8*a
*f**4) - 6*c*d**2*f**3*x**2*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**
4) + 6*c*d**2*f**3*x**2/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) - 6*c*d**2*f**2
```

```
*x*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 6*c*d**2*f**2*x/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 6*c*d**2*f/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + d**3*f**4*x**4*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + d**3*f**4*x**4/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) - 2*d**3*f**3*x**3*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 2*d**3*f**3*x**3/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) - 3*d**3*f**2*x**2*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 3*d**3*f**2*x**2/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) - 3*d**3*f*x*tanh(e + f*x)/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 3*d**3*f*x/(8*a*f**4*tanh(e + f*x) + 8*a*f**4) + 3*d**3/(8*a*f**4*tanh(e + f*x) + 8*a*f**4), Ne(f, 0)), ((c**3*x + 3*c**2*d*x**2/2 + c*d**2*x**3 + d**3*x**4/4)/(a*coth(e) + a), True))
```

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.08

$$\int \frac{(c + dx)^3}{a + a \coth(e + fx)} dx = \frac{1}{4} c^3 \left(\frac{2(fx + e)}{af} + \frac{e^{(-2fx-2e)}}{af} \right) \\ + \frac{3(2f^2x^2e^{(2e)} + (2fx + 1)e^{(-2fx)})c^2de^{(-2e)}}{8af^2} \\ + \frac{(4f^3x^3e^{(2e)} + 3(2f^2x^2 + 2fx + 1)e^{(-2fx)})cd^2e^{(-2e)}}{8af^3} \\ + \frac{(2f^4x^4e^{(2e)} + (4f^3x^3 + 6f^2x^2 + 6fx + 3)e^{(-2fx)})d^3e^{(-2e)}}{16af^4}$$

```
[In] integrate((d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="maxima")
[Out] 1/4*c^3*(2*(f*x + e)/(a*f) + e^(-2*f*x - 2*e)/(a*f)) + 3/8*(2*f^2*x^2*e^(2*e) + (2*f*x + 1)*e^(-2*f*x))*c^2*d*e^(-2*e)/(a*f^2) + 1/8*(4*f^3*x^3*e^(2*e) + 3*(2*f^2*x^2 + 2*f*x + 1)*e^(-2*f*x))*c*d^2*e^(-2*e)/(a*f^3) + 1/16*(2*f^4*x^4*e^(2*e) + (4*f^3*x^3 + 6*f^2*x^2 + 6*f*x + 3)*e^(-2*f*x))*d^3*e^(-2*e)/(a*f^4)
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 188, normalized size of antiderivative = 1.11

$$\int \frac{(c + dx)^3}{a + a \coth(e + fx)} dx \\ = \frac{(2d^3f^4x^4e^{(2fx+2e)} + 8cd^2f^4x^3e^{(2fx+2e)} + 12c^2df^4x^2e^{(2fx+2e)} + 4d^3f^3x^3 + 8c^3f^4xe^{(2fx+2e)} + 12cd^2f^3x^2e^{(2fx+2e)})}{16af^4}$$

```
[In] integrate((d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="giac")
[Out] 1/16*(2*d^3*f^4*x^4*e^(2*f*x + 2*e) + 8*c*d^2*f^4*x^3*e^(2*f*x + 2*e) + 12*c^2*d*f^4*x^2*e^(2*f*x + 2*e) + 4*d^3*f^3*x^3 + 8*c^3*f^4*x*e^(2*f*x + 2*e) + 12*c*d^2*f^3*x^2 + 12*c^2*d*f^3*x + 6*d^3*f^2*x^2 + 4*c^3*f^3 + 12*c*d^2*f^2*x + 6*c^2*d*f^2 + 6*d^3*f*x + 6*c*d^2*f + 3*d^3)*e^(-2*f*x - 2*e)/(a*f^4)
```

Mupad [B] (verification not implemented)

Time = 2.16 (sec) , antiderivative size = 223, normalized size of antiderivative = 1.32

$$\begin{aligned} & \int \frac{(c+dx)^3}{a+a \coth(e+fx)} dx \\ &= \frac{4 c^3 f^4 x + 6 c^2 d f^4 x^2 + 6 c^2 d f^3 x + 4 c d^2 f^4 x^3 + 6 c d^2 f^3 x^2 + 6 c d^2 f^2 x + d^3 f^4 x^4 + 2 d^3 f^3 x^3 + 3 d^3 f^2}{8 a f^4} \\ &\quad - \frac{4 c^3 f^3 + 12 c^2 d f^3 x + 6 c^2 d f^2 + 12 c d^2 f^3 x^2 + 12 c d^2 f^2 x + 6 c d^2 f + 4 d^3 f^3 x^3 + 6 d^3 f^2 x^2 + 6 d^3 f x}{8 a f^4 (\coth(e+fx) + 1)} \end{aligned}$$

[In] `int((c + d*x)^3/(a + a*coth(e + f*x)),x)`

```
[Out] (3*d^3*f*x + 4*c^3*f^4*x + 3*d^3*f^2*x^2 + 2*d^3*f^3*x^3 + d^3*f^4*x^4 + 6*c*d^2*f^3*x^2 + 6*c^2*d*f^4*x^2 + 4*c*d^2*f^4*x^3 + 6*c*d^2*f^2*x + 6*c^2*d*f^3*x)/(8*a*f^4) - (3*d^3 + 4*c^3*f^3 + 6*d^3*f*x + 6*c^2*d*f^2 + 6*d^3*f^2*x^2 + 4*d^3*f^3*x^3 + 6*c*d^2*f + 12*c*d^2*f^3*x^2 + 12*c*d^2*f^2*x + 12*c^2*d*f^3*x)/(8*a*f^4*(coth(e + f*x) + 1))
```

3.17 $\int \frac{(c+dx)^2}{a+a \coth(e+fx)} dx$

Optimal result	117
Rubi [A] (verified)	117
Mathematica [A] (verified)	119
Maple [A] (verified)	119
Fricas [A] (verification not implemented)	120
Sympy [B] (verification not implemented)	120
Maxima [A] (verification not implemented)	121
Giac [A] (verification not implemented)	121
Mupad [B] (verification not implemented)	121

Optimal result

Integrand size = 20, antiderivative size = 122

$$\int \frac{(c+dx)^2}{a+a \coth(e+fx)} dx = \frac{d^2x}{4af^2} + \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a+a \coth(e+fx))} \\ - \frac{d(c+dx)}{2f^2(a+a \coth(e+fx))} - \frac{(c+dx)^2}{2f(a+a \coth(e+fx))}$$

[Out] $1/4*d^2*x/a/f^2+1/4*(d*x+c)^2/a/f+1/6*(d*x+c)^3/a/d-1/4*d^2/f^3/(a+a*\coth(f*x+e))-1/2*d*(d*x+c)/f^2/(a+a*\coth(f*x+e))-1/2*(d*x+c)^2/f/(a+a*\coth(f*x+e))$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3804, 3560, 8}

$$\int \frac{(c+dx)^2}{a+a \coth(e+fx)} dx = -\frac{d(c+dx)}{2f^2(a \coth(e+fx)+a)} - \frac{(c+dx)^2}{2f(a \coth(e+fx)+a)} \\ + \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a \coth(e+fx)+a)} + \frac{d^2x}{4af^2}$$

[In] $\text{Int}[(c+d*x)^2/(a+a*\text{Coth}[e+f*x]),x]$

[Out] $(d^2*x)/(4*a*f^2) + (c+d*x)^2/(4*a*f) + (c+d*x)^3/(6*a*d) - d^2/(4*f^3*(a+a*\text{Coth}[e+f*x])) - (d*(c+d*x))/(2*f^2*(a+a*\text{Coth}[e+f*x])) - (c+d*x)^2/(2*f*(a+a*\text{Coth}[e+f*x]))$

Rule 8

```
Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]
```

Rule 3560

```
Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_.) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{(c+dx)^3}{6ad} - \frac{(c+dx)^2}{2f(a+a\coth(e+fx))} + \frac{d \int \frac{c+dx}{a+a\coth(e+fx)} dx}{f} \\
 &= \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d(c+dx)}{2f^2(a+a\coth(e+fx))} \\
 &\quad - \frac{(c+dx)^2}{2f(a+a\coth(e+fx))} + \frac{d^2 \int \frac{1}{a+a\coth(e+fx)} dx}{2f^2} \\
 &= \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a+a\coth(e+fx))} \\
 &\quad - \frac{d(c+dx)}{2f^2(a+a\coth(e+fx))} - \frac{(c+dx)^2}{2f(a+a\coth(e+fx))} + \frac{d^2 \int 1 dx}{4af^2} \\
 &= \frac{d^2x}{4af^2} + \frac{(c+dx)^2}{4af} + \frac{(c+dx)^3}{6ad} - \frac{d^2}{4f^3(a+a\coth(e+fx))} \\
 &\quad - \frac{d(c+dx)}{2f^2(a+a\coth(e+fx))} - \frac{(c+dx)^2}{2f(a+a\coth(e+fx))}
 \end{aligned}$$

Mathematica [A] (verified)

Time = 0.59 (sec) , antiderivative size = 169, normalized size of antiderivative = 1.39

$$\int \frac{(c + dx)^2}{a + a \coth(e + fx)} dx \\ = \frac{\text{csch}(e + fx)(\cosh(fx) + \sinh(fx)) ((2c^2 f^2 + 2cd(1 + 2fx) + d^2(1 + 2fx + 2f^2 x^2)) \cosh(2fx)(\cosh(e) + \sinh(e)))}{8a^3 f^3}$$

[In] Integrate[(c + d*x)^2/(a + a*Coth[e + f*x]), x]

[Out] $(\text{Csch}[e + f*x] * (\text{Cosh}[f*x] + \text{Sinh}[f*x]) * ((2*c^2 f^2 + 2*c*d*f*(1 + 2*f*x) + d^2*(1 + 2*f*x + 2*f^2*x^2)) * \text{Cosh}[2*f*x] * (\text{Cosh}[e] - \text{Sinh}[e])) + (4*f^3*x*(3*c^2 + 3*c*d*x + d^2*x^2) * (\text{Cosh}[e] + \text{Sinh}[e]))/3 + (2*c^2 f^2 + 2*c*d*f*(1 + 2*f*x) + d^2*(1 + 2*f*x + 2*f^2*x^2)) * (-\text{Cosh}[e] + \text{Sinh}[e]) * \text{Sinh}[2*f*x])) / (8*a*f^3*(1 + \text{Coth}[e + f*x]))$

Maple [A] (verified)

Time = 0.31 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.84

method	result
risch	$\frac{d^2 x^3}{6 a} + \frac{d c x^2}{2 a} + \frac{c^2 x}{2 a} + \frac{c^3}{6 da} + \frac{(2 d^2 x^2 f^2 + 4 c d f^2 x + 2 c^2 f^2 + 2 d^2 f x + 2 c d f + d^2) e^{-2 f x - 2 e}}{8 a f^3}$
parallelrisch	$\frac{((2 d^2 x^3 + 6 d c x^2 + 6 c^2 x) f^3 + (-3 x^2 d^2 - 6 c d x - 6 c^2) f^2 + (-3 x d^2 - 6 c d) f - 3 d^2) \tanh(f x + e) + 6 x \left(\left(\frac{1}{3} x^2 d^2 + c d x + c^2 \right) f^2 + d^2 \right)}{12 f^3 a (1 + \tanh(f x + e))}$
derivativedivides	$\frac{-c^2 f^2 \left(\frac{\cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x}{2} - \frac{e}{2} \right) + 2 d e c f \left(\frac{\cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x}{2} - \frac{e}{2} \right) - 2 d c f \left(\frac{(f x + e) \cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x^2}{2} - \frac{c d x}{2} - \frac{c^2}{2} \right)}{12 f^3 a (1 + \tanh(f x + e))}$
default	$\frac{-c^2 f^2 \left(\frac{\cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x}{2} - \frac{e}{2} \right) + 2 d e c f \left(\frac{\cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x}{2} - \frac{e}{2} \right) - 2 d c f \left(\frac{(f x + e) \cosh(f x + e) \sinh(f x + e)}{2} - \frac{f x^2}{2} - \frac{c d x}{2} - \frac{c^2}{2} \right)}{12 f^3 a (1 + \tanh(f x + e))}$

[In] int((d*x+c)^2/(a+a*coth(f*x+e)), x, method=_RETURNVERBOSE)

[Out] $\frac{1}{6} d^2/a*x^3 + \frac{1}{2} d/a*c*x^2 + \frac{1}{2} /a*c^2*x + \frac{1}{6}/d/a*c^3 + \frac{1}{8}*(2*d^2*f^2*x^2 + 4*c*d*f^2*x + 2*c^2*f^2 + 2*d^2*f*x + 2*c*d*f + d^2)/a/f^3*\exp(-2*f*x - 2*e)$

Fricas [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 192, normalized size of antiderivative = 1.57

$$\int \frac{(c+dx)^2}{a+a\coth(e+fx)} dx \\ = \frac{(4d^2f^3x^3 + 6c^2f^2 + 6cdf + 6(2cdf^3 + d^2f^2)x^2 + 3d^2 + 6(2c^2f^3 + 2cdf^2 + d^2f)x)\cosh(fx+e) + (4d^2f^3x^3 + 6c^2f^2 + 6cdf + 6(2cdf^3 + d^2f^2)x^2 + 3d^2 + 6(2c^2f^3 + 2cdf^2 + d^2f)x)\sinh(fx+e)}{24(af^3\cosh(fx+e) + af^3\sinh(fx+e))}$$

```
[In] integrate((d*x+c)^2/(a+a*coth(f*x+e)),x, algorithm="fricas")
[Out] 1/24*((4*d^2*f^3*x^3 + 6*c^2*f^2 + 6*c*d*f + 6*(2*c*d*f^3 + d^2*f^2)*x^2 + 3*d^2 + 6*(2*c^2*f^3 + 2*c*d*f^2 + d^2*f)*x)*cosh(f*x + e) + (4*d^2*f^3*x^3 - 6*c^2*f^2 - 6*c*d*f + 6*(2*c*d*f^3 - d^2*f^2)*x^2 - 3*d^2 + 6*(2*c^2*f^3 - 2*c*d*f^2 - d^2*f)*x)*sinh(f*x + e))/(a*f^3*cosh(f*x + e) + a*f^3*sinh(f*x + e))
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 522 vs. 2(95) = 190.

Time = 0.63 (sec) , antiderivative size = 522, normalized size of antiderivative = 4.28

$$\int \frac{(c+dx)^2}{a+a\coth(e+fx)} dx \\ = \begin{cases} \frac{6c^2f^3x\tanh(e+fx)}{12af^3\tanh(e+fx)+12af^3} + \frac{6c^2f^3x}{12af^3\tanh(e+fx)+12af^3} + \frac{6c^2f^2}{12af^3\tanh(e+fx)+12af^3} + \frac{6cdf^3x^2\tanh(e+fx)}{12af^3\tanh(e+fx)+12af^3} + \frac{6cdf^3x^3}{12af^3\tanh(e+fx)+12af^3} \\ \frac{c^2x+cdx^2+\frac{d^2x^3}{3}}{a\coth(e)+a} \end{cases}$$

```
[In] integrate((d*x+c)**2/(a+a*coth(f*x+e)),x)
[Out] Piecewise((6*c**2*f**3*x*tanh(e + f*x)/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 6*c**2*f**3*x/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 6*c**2*f**2/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 6*c*d*f**3*x**2*tanh(e + f*x)/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 6*c*d*f**3*x**2/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) - 6*c*d*f**2*x*tanh(e + f*x)/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 6*c*d*f**2*x/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 2*d**2*f**3*x**3*tanh(e + f*x)/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 2*d**2*f**3*x**3/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) - 3*d**2*f**2*x**2*tanh(e + f*x)/(12*a*f**3*tanh(e + f*x) + 12*a*f**3) + 3*d**2*f**2*x**2/(12*a*f**3*tanh(e + f*x) + 12*a*f**3), Ne(f, 0)), ((c**2*x + c*d*x**2 + d**2*x**3/3)/(a*coth(e) + a), True))
```

Maxima [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 124, normalized size of antiderivative = 1.02

$$\int \frac{(c + dx)^2}{a + a \coth(e + fx)} dx = \frac{1}{4} c^2 \left(\frac{2(fx + e)}{af} + \frac{e^{(-2fx-2e)}}{af} \right) \\ + \frac{(2f^2x^2e^{(2e)} + (2fx + 1)e^{(-2fx)})cde^{(-2e)}}{4af^2} \\ + \frac{(4f^3x^3e^{(2e)} + 3(2f^2x^2 + 2fx + 1)e^{(-2fx)})d^2e^{(-2e)}}{24af^3}$$

[In] `integrate((d*x+c)^2/(a+a*coth(f*x+e)),x, algorithm="maxima")`

[Out] $\frac{1}{4}c^2(2*(fx + e)/(a*f) + e^{(-2fx - 2e)/(a*f)}) + \frac{1}{4}(2*f^2x^2e^{(2e)} + (2fx + 1)*e^{(-2fx)})*c*d*e^{(-2e)/(a*f^2)} + \frac{1}{24}(4*f^3x^3e^{(2e)} + 3(2f^2x^2 + 2fx + 1)e^{(-2fx)})*d^2e^{(-2e)/(a*f^3)}$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 119, normalized size of antiderivative = 0.98

$$\int \frac{(c + dx)^2}{a + a \coth(e + fx)} dx \\ = \frac{(4d^2f^3x^3e^{(2fx+2e)} + 12cdf^3x^2e^{(2fx+2e)} + 12c^2f^3xe^{(2fx+2e)} + 6d^2f^2x^2 + 12cdf^2x + 6c^2f^2 + 6d^2fx + 6e^{(2fx+2e)})}{24af^3}$$

[In] `integrate((d*x+c)^2/(a+a*coth(f*x+e)),x, algorithm="giac")`

[Out] $\frac{1}{24}(4d^2f^3x^3e^{(2fx+2e)} + 12cdf^3x^2e^{(2fx+2e)} + 12c^2f^3xe^{(2fx+2e)} + 6d^2f^2x^2 + 12cdf^2x + 6c^2f^2 + 6d^2fx + 6e^{(2fx+2e)})/(a*f^3)$

Mupad [B] (verification not implemented)

Time = 1.97 (sec) , antiderivative size = 186, normalized size of antiderivative = 1.52

$$\int \frac{(c + dx)^2}{a + a \coth(e + fx)} dx = \frac{e^{-2e-2fx}(12c^2x e^{2e+2fx} + 4d^2x^3 e^{2e+2fx} + 12cdx^2 e^{2e+2fx})}{24a} \\ + \frac{e^{-2e-2fx}(3d^2+3d^2e^{2e+2fx})}{24} + \frac{fe^{-2e-2fx}(6cd+6d^2x+6cde^{2e+2fx})}{24} + \frac{f^2e^{-2e-2fx}(6c^2+6c^2e^{2e+2fx}+6d^2x^2+12cdx)}{24}$$

[In] $\int ((c + d*x)^2 / (a + a*\coth(e + f*x)), x)$

[Out]
$$\begin{aligned} & \frac{\exp(-2e - 2fx) \cdot (12c^2x \cdot \exp(2e + 2fx) + 4d^2x^3 \cdot \exp(2e + 2fx) \\ & + 12cdx^2 \cdot \exp(2e + 2fx))}{24a} + \frac{(\exp(-2e - 2fx) \cdot (3d^2 + 3d \\ & \cdot 2 \cdot \exp(2e + 2fx)))}{24} + \frac{(fx \cdot \exp(-2e - 2fx) \cdot (6cd + 6d^2x + 6c^2 \cdot \exp(2e + 2fx) \\ & \cdot x))}{24} + \frac{(f^2 \cdot \exp(-2e - 2fx) \cdot (6c^2 + 6c^2 \cdot \exp(2e + 2fx) \\ & + 6d^2x^2 + 12cdx))}{24} \end{aligned} / (a \cdot f^3)$$

3.18 $\int \frac{c+dx}{a+a \coth(e+fx)} dx$

Optimal result	123
Rubi [A] (verified)	123
Mathematica [A] (verified)	124
Maple [A] (verified)	125
Fricas [A] (verification not implemented)	125
Sympy [B] (verification not implemented)	125
Maxima [A] (verification not implemented)	126
Giac [A] (verification not implemented)	126
Mupad [B] (verification not implemented)	127

Optimal result

Integrand size = 18, antiderivative size = 74

$$\int \frac{c+dx}{a+a \coth(e+fx)} dx = \frac{dx}{4af} + \frac{(c+dx)^2}{4ad} - \frac{d}{4f^2(a+a \coth(e+fx))} - \frac{c+dx}{2f(a+a \coth(e+fx))}$$

[Out] $1/4*d*x/a/f + 1/4*(d*x+c)^2/a/d - 1/4*d/f^2/(a+a*\coth(f*x+e)) + 1/2*(-d*x-c)/f/(a+a*\coth(f*x+e))$

Rubi [A] (verified)

Time = 0.04 (sec), antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3804, 3560, 8}

$$\int \frac{c+dx}{a+a \coth(e+fx)} dx = -\frac{c+dx}{2f(a \coth(e+fx)+a)} + \frac{(c+dx)^2}{4ad} - \frac{d}{4f^2(a \coth(e+fx)+a)} + \frac{dx}{4af}$$

[In] $\text{Int}[(c+d*x)/(a+a*\coth[e+f*x]),x]$

[Out] $(d*x)/(4*a*f) + (c+d*x)^2/(4*a*d) - d/(4*f^2*(a+a*\coth[e+f*x])) - (c+d*x)/(2*f*(a+a*\coth[e+f*x]))$

Rule 8

$\text{Int}[a_, x_Symbol] :> \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3560

```
Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]
```

Rule 3804

```
Int[((c_.) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_.) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + (Dist[a*d*(m/(2*b*f)), Int[(c + d*x)^(m - 1)/(a + b*Tan[e + f*x]), x], x] - Simp[a*((c + d*x)^m/(2*b*f*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{(c + dx)^2}{4ad} - \frac{c + dx}{2f(a + a \coth(e + fx))} + \frac{d \int \frac{1}{a + a \coth(e + fx)} dx}{2f} \\ &= \frac{(c + dx)^2}{4ad} - \frac{d}{4f^2(a + a \coth(e + fx))} - \frac{c + dx}{2f(a + a \coth(e + fx))} + \frac{d \int 1 dx}{4af} \\ &= \frac{dx}{4af} + \frac{(c + dx)^2}{4ad} - \frac{d}{4f^2(a + a \coth(e + fx))} - \frac{c + dx}{2f(a + a \coth(e + fx))} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.71 (sec), antiderivative size = 81, normalized size of antiderivative = 1.09

$$\begin{aligned} &\int \frac{c + dx}{a + a \coth(e + fx)} dx \\ &= \frac{2cf(-1 + 2fx) + d(-1 - 2fx + 2f^2x^2) + (2cf(1 + 2fx) + d(1 + 2fx + 2f^2x^2)) \coth(e + fx)}{8af^2(1 + \coth(e + fx))} \end{aligned}$$

[In] `Integrate[(c + d*x)/(a + a*Coth[e + f*x]), x]`

[Out] `(2*c*f*(-1 + 2*f*x) + d*(-1 - 2*f*x + 2*f^2*x^2) + (2*c*f*(1 + 2*f*x) + d*(1 + 2*f*x + 2*f^2*x^2))*Coth[e + f*x])/ (8*a*f^2*(1 + Coth[e + f*x]))`

Maple [A] (verified)

Time = 0.19 (sec), antiderivative size = 46, normalized size of antiderivative = 0.62

method	result
risch	$\frac{dx^2}{4a} + \frac{cx}{2a} + \frac{(2dxf+2cf+d)e^{-2fx-2e}}{8af^2}$
parallelrisch	$\frac{((dx^2+2cx)f^2+(-dx-2c)f-d)\tanh(fx+e)+2xf\left(\left(\frac{dx}{2}+c\right)f+\frac{d}{2}\right)}{4f^2a(1+\tanh(fx+e))}$
derivativedivides	$-cf\left(\frac{\cosh(fx+e)\sinh(fx+e)}{2}-\frac{fx}{2}-\frac{e}{2}\right)+de\left(\frac{\cosh(fx+e)\sinh(fx+e)}{2}-\frac{fx}{2}-\frac{e}{2}\right)-d\left(\frac{(fx+e)\cosh(fx+e)\sinh(fx+e)}{2}-\frac{(fx+e)^2}{4}\right)-\frac{f^2a}{f^2a}$
default	$-cf\left(\frac{\cosh(fx+e)\sinh(fx+e)}{2}-\frac{fx}{2}-\frac{e}{2}\right)+de\left(\frac{\cosh(fx+e)\sinh(fx+e)}{2}-\frac{fx}{2}-\frac{e}{2}\right)-d\left(\frac{(fx+e)\cosh(fx+e)\sinh(fx+e)}{2}-\frac{(fx+e)^2}{4}\right)-\frac{f^2a}{f^2a}$

```
[In] int((d*x+c)/(a+a*cOTH(f*x+e)),x,method=_RETURNVERBOSE)
```

[Out] $\frac{1}{4} \cdot a \cdot d \cdot x^2 + \frac{1}{2} \cdot a \cdot c \cdot x + \frac{1}{8} \cdot (2 \cdot d \cdot f \cdot x + 2 \cdot c \cdot f \cdot d) / a \cdot f^2 \cdot \exp(-2 \cdot f \cdot x - 2 \cdot e)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 101, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int \frac{c + dx}{a + a \coth(e + fx)} dx \\ &= \frac{(2df^2x^2 + 2cf + 2(2cf^2 + df)x + d) \cosh(fx + e) + (2df^2x^2 - 2cf + 2(2cf^2 - df)x - d) \sinh(fx + e)}{8(af^2 \cosh(fx + e) + af^2 \sinh(fx + e))} \end{aligned}$$

```
[In] integrate((d*x+c)/(a+a*cOTH(f*x+e)),x, algorithm="fricas")
```

[Out] $\frac{1}{8} \left((2d^2 f^2 x^2 + 2c^2 f + 2(2c^2 f^2 + d^2 f)x + d) \cosh(fx + e) + (2d^2 f^2 x^2 - 2c^2 f + 2(2c^2 f^2 - d^2 f)x - d) \sinh(fx + e) \right) / (a^2 f^2 \cosh(fx + e) + a^2 f^2 \sinh(fx + e))$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 250 vs. $2(58) = 116$.

Time = 0.54 (sec) , antiderivative size = 250, normalized size of antiderivative = 3.38

$$\int \frac{c + dx}{a + a \coth(e + fx)} dx \\ = \begin{cases} \frac{2cf^2 x \tanh(e+fx)}{4af^2 \tanh(e+fx)+4af^2} + \frac{2cf^2 x}{4af^2 \tanh(e+fx)+4af^2} + \frac{2cf}{4af^2 \tanh(e+fx)+4af^2} + \frac{df^2 x^2 \tanh(e+fx)}{4af^2 \tanh(e+fx)+4af^2} + \frac{df^2 x^2}{4af^2 \tanh(e+fx)+4af^2} \\ \frac{cx+\frac{dx^2}{2}}{a \coth(e)+a} \end{cases}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e)),x)`

[Out] `Piecewise((2*c*f**2*x*tanh(e + f*x)/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + 2*c*f**2*x/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + 2*c*f/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + d*f**2*x**2*tanh(e + f*x)/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + d*f**2*x**2/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) - d*f*x*tanh(e + f*x)/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + d*f*x/(4*a*f**2*tanh(e + f*x) + 4*a*f**2) + d/(4*a*f**2*tanh(e + f*x) + 4*a*f**2), Ne(f, 0)), ((c*x + d*x**2/2)/(a*coth(e) + a), True))`

Maxima [A] (verification not implemented)

none

Time = 0.21 (sec) , antiderivative size = 72, normalized size of antiderivative = 0.97

$$\int \frac{c + dx}{a + a \coth(e + fx)} dx = \frac{1}{4} c \left(\frac{2(fx + e)}{af} + \frac{e^{(-2fx-2e)}}{af} \right) + \frac{(2f^2x^2e^{(2e)} + (2fx + 1)e^{(-2fx)})de^{(-2e)}}{8af^2}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e)),x, algorithm="maxima")`

[Out] `1/4*c*(2*(f*x + e)/(a*f) + e^{(-2*f*x - 2*e)}/(a*f)) + 1/8*(2*f^2*x^2*e^{(2*e)} + (2*f*x + 1)*e^{(-2*f*x)})*d*e^{(-2*e)}/(a*f^2)`

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 62, normalized size of antiderivative = 0.84

$$\int \frac{c + dx}{a + a \coth(e + fx)} dx = \frac{(2df^2x^2e^{(2fx+2e)} + 4cf^2xe^{(2fx+2e)} + 2dfx + 2cf + d)e^{(-2fx-2e)}}{8af^2}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e)),x, algorithm="giac")`

[Out] `1/8*(2*d*f^2*x^2*e^{(2*f*x + 2*e)} + 4*c*f^2*x*e^{(2*f*x + 2*e)} + 2*d*f*x + 2*c*f + d)*e^{(-2*f*x - 2*e)}/(a*f^2)`

Mupad [B] (verification not implemented)

Time = 1.94 (sec) , antiderivative size = 76, normalized size of antiderivative = 1.03

$$\int \frac{c + dx}{a + a \coth(e + fx)} dx = \frac{\frac{dx^2}{4} + \left(\frac{c}{2} + \frac{d}{4f}\right)x}{a} - \frac{\frac{\frac{d}{4} + \frac{cf}{2}}{f^2} - x\left(\frac{c}{2} - \frac{d}{4f}\right) + x\left(\frac{c}{2} + \frac{d}{4f}\right)}{a + a \coth(e + fx)}$$

[In] `int((c + d*x)/(a + a*coth(e + f*x)),x)`

[Out] `(x*(c/2 + d/(4*f)) + (d*x^2)/4)/a - ((d/4 + (c*f)/2)/f^2 - x*(c/2 - d/(4*f)) + x*(c/2 + d/(4*f)))/(a + a*coth(e + f*x))`

3.19 $\int \frac{1}{(c+dx)(a+a \coth(e+fx))} dx$

Optimal result	128
Rubi [A] (verified)	128
Mathematica [A] (verified)	130
Maple [A] (verified)	130
Fricas [A] (verification not implemented)	130
Sympy [F]	131
Maxima [A] (verification not implemented)	131
Giac [A] (verification not implemented)	131
Mupad [F(-1)]	132

Optimal result

Integrand size = 20, antiderivative size = 157

$$\begin{aligned} \int \frac{1}{(c+dx)(a+a \coth(e+fx))} dx = & -\frac{\cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{2ad} + \frac{\log(c+dx)}{2ad} \\ & + \frac{\text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2ad} \\ & + \frac{\cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2ad} \\ & - \frac{\sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2ad} \end{aligned}$$

[Out] $-1/2*\text{Chi}(2*c*f/d+2*f*x)*\cosh(-2*e+2*c*f/d)/a/d+1/2*\ln(d*x+c)/a/d+1/2*\cosh(-2*e+2*c*f/d)*\text{Shi}(2*c*f/d+2*f*x)/a/d-1/2*\text{Chi}(2*c*f/d+2*f*x)*\sinh(-2*e+2*c*f/d)/a/d+1/2*\text{Shi}(2*c*f/d+2*f*x)*\sinh(-2*e+2*c*f/d)/a/d$

Rubi [A] (verified)

Time = 0.20 (sec), antiderivative size = 157, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.200, Rules used = {3807, 3384, 3379, 3382}

$$\begin{aligned} \int \frac{1}{(c+dx)(a+a \coth(e+fx))} dx = & \frac{\text{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{2ad} \\ & - \frac{\text{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{2ad} \\ & - \frac{\sinh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{2ad} \\ & + \frac{\cosh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{2ad} + \frac{\log(c+dx)}{2ad} \end{aligned}$$

[In] $\text{Int}[1/((c + d*x)*(a + a*\text{Coth}[e + f*x])), x]$

[Out] $-1/2*(\text{Cosh}[2*e - (2*c*f)/d]*\text{CoshIntegral}[(2*c*f)/d + 2*f*x])/(a*d) + \text{Log}[c + d*x]/(2*a*d) + (\text{CoshIntegral}[(2*c*f)/d + 2*f*x]*\text{Sinh}[2*e - (2*c*f)/d])/ (2*a*d) + (\text{Cosh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x])/ (2*a*d) - (\text{Sinh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x])/ (2*a*d)$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*f_.*x_]/((c_.) + (d_.*x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&& \text{EqQ}[d*e - c*f*fz*I, 0]$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*f_.*x_]/((c_.) + (d_.*x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&& \text{EqQ}[d*(e - \pi/2) - c*f*fz*I, 0]$

Rule 3384

$\text{Int}[\sin[(e_.) + (f_.*x_)]/((c_.) + (d_.*x_)), x_{\text{Symbol}}] \rightarrow \text{Dist}[\text{Cos}[(d*e - c*f)/d], \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c + d*x), x], x] + \text{Dist}[\text{Sin}[(d*e - c*f)/d], \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c + d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{NeQ}[d*e - c*f, 0]$

Rule 3807

$\text{Int}[1/(((c_.) + (d_.*x_))*(a_.) + (b_.*\tan[(e_.) + (f_.*x_)])), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[c + d*x]/(2*a*d), x] + (\text{Dist}[1/(2*a), \text{Int}[\text{Cos}[2*e + 2*f*x]/(c + d*x), x], x] + \text{Dist}[1/(2*b), \text{Int}[\text{Sin}[2*e + 2*f*x]/(c + d*x), x], x]) /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \frac{\log(c + dx)}{2ad} + \frac{i \int \frac{\sin(2(ie + \frac{\pi}{2}) + 2ifx)}{c + dx} dx}{2a} + \frac{\int \frac{\cos(2(ie + \frac{\pi}{2}) + 2ifx)}{c + dx} dx}{2a} \\ &= \frac{\log(c + dx)}{2ad} - \frac{\cosh(2e - \frac{2cf}{d}) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} + \frac{\cosh(2e - \frac{2cf}{d}) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} \\ &\quad + \frac{\sinh(2e - \frac{2cf}{d}) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} - \frac{\sinh(2e - \frac{2cf}{d}) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c + dx} dx}{2a} \\ &= -\frac{\cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{2ad} + \frac{\log(c + dx)}{2ad} + \frac{\text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2ad} \\ &\quad + \frac{\cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2ad} - \frac{\sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2ad} \end{aligned}$$

Mathematica [A] (verified)

Time = 0.63 (sec) , antiderivative size = 122, normalized size of antiderivative = 0.78

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx \\ = \frac{\operatorname{csch}(e + fx)(\cosh(fx) + \sinh(fx)) \left(\log(f(c + dx))(\cosh(e) + \sinh(e)) + \operatorname{Chi}\left(\frac{2f(c+dx)}{d}\right) (-\cosh(e - \frac{2cf}{d}) + \sinh(e - \frac{2cf}{d})) \right)}{2ad(1 + \coth(e + fx))}$$

```
[In] Integrate[1/((c + d*x)*(a + a*Coth[e + f*x])),x]
[Out] (Csch[e + f*x]*(Cosh[f*x] + Sinh[f*x])*(Log[f*(c + d*x)]*(Cosh[e] + Sinh[e])) + CoshIntegral[(2*f*(c + d*x))/d]*(-Cosh[e - (2*c*f)/d] + Sinh[e - (2*c*f)/d]) + (Cosh[e - (2*c*f)/d] - Sinh[e - (2*c*f)/d])*SinhIntegral[(2*f*(c + d*x))/d]))/(2*a*d*(1 + Coth[e + f*x]))
```

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 61, normalized size of antiderivative = 0.39

method	result	size
risch	$\frac{\ln(dx+c)}{2ad} + \frac{e^{\frac{2cf-2de}{d}} \operatorname{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{2ad}$	61

```
[In] int(1/(d*x+c)/(a+a*coth(f*x+e)),x,method=_RETURNVERBOSE)
[Out] 1/2*ln(d*x+c)/a/d+1/2/a/d*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.24 (sec) , antiderivative size = 75, normalized size of antiderivative = 0.48

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx \\ = -\frac{\operatorname{Ei}\left(-\frac{2(df+cf)}{d}\right) \cosh\left(-\frac{2(de-cf)}{d}\right) + \operatorname{Ei}\left(-\frac{2(df+cf)}{d}\right) \sinh\left(-\frac{2(de-cf)}{d}\right) - \log(dx + c)}{2ad}$$

```
[In] integrate(1/(d*x+c)/(a+a*coth(f*x+e)),x, algorithm="fricas")
[Out] -1/2*(Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) + Ei(-2*(d*f*x + c*f)/d)*sinh(-2*(d*e - c*f)/d) - log(d*x + c))/(a*d)
```

Sympy [F]

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx = \frac{\int \frac{1}{c \coth(e + fx) + c + dx \coth(e + fx) + dx} dx}{a}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e)),x)`

[Out] `Integral(1/(c*coth(e + f*x) + c + d*x*coth(e + f*x) + d*x), x)/a`

Maxima [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.31

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx = \frac{e^{\left(-2e + \frac{2cf}{d}\right)} E_1\left(\frac{2(dx+c)f}{d}\right)}{2ad} + \frac{\log(dx + c)}{2ad}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e)),x, algorithm="maxima")`

[Out] `1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(1, 2*(d*x + c)*f/d)/(a*d) + 1/2*log(d*x + c)/(a*d)`

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 48, normalized size of antiderivative = 0.31

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx = -\frac{\left(Ei\left(-\frac{2(df+cf)}{d}\right) e^{\left(\frac{2cf}{d}\right)} - e^{(2e)} \log(dx + c)\right) e^{(-2e)}}{2ad}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e)),x, algorithm="giac")`

[Out] `-1/2*(Ei(-2*(d*f*x + c*f)/d)*e^(2*c*f/d) - e^(2*e)*log(d*x + c))*e^(-2*e)/(a*d)`

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))} dx = \int \frac{1}{(a + a \coth(e + fx))(c + dx)} dx$$

[In] `int(1/((a + a*coth(e + f*x))*(c + d*x)),x)`

[Out] `int(1/((a + a*coth(e + f*x))*(c + d*x)), x)`

3.20 $\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))} dx$

Optimal result	133
Rubi [A] (verified)	133
Mathematica [A] (verified)	135
Maple [A] (verified)	136
Fricas [A] (verification not implemented)	136
Sympy [F]	136
Maxima [A] (verification not implemented)	137
Giac [B] (verification not implemented)	137
Mupad [F(-1)]	138

Optimal result

Integrand size = 20, antiderivative size = 159

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+a \coth(e+fx))} dx = & \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Chi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \\ & - \frac{1}{d(c+dx)(a+a \coth(e+fx))} \\ & - \frac{f \text{Chi}\left(\frac{2cf}{d} + 2fx\right) \sinh\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ & - \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \\ & + \frac{f \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(\frac{2cf}{d} + 2fx\right)}{ad^2} \end{aligned}$$

```
[Out] f*Chi(2*c*f/d+2*f*x)*cosh(-2*e+2*c*f/d)/a/d^2-1/d/(d*x+c)/(a+a*coth(f*x+e))-f*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d+2*f*x)/a/d^2+f*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^2-f*Shi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^2
```

Rubi [A] (verified)

Time = 0.18 (sec), antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used

= {3805, 3384, 3379, 3382}

$$\int \frac{1}{(c+dx)^2(a+a\coth(e+fx))} dx = -\frac{f \text{Chi}\left(2xf + \frac{2cf}{d}\right) \sinh\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ + \frac{f \text{Chi}\left(2xf + \frac{2cf}{d}\right) \cosh\left(2e - \frac{2cf}{d}\right)}{ad^2} \\ + \frac{f \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{ad^2} \\ - \frac{f \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{ad^2} \\ - \frac{1}{d(c+dx)(a\coth(e+fx)+a)}$$

[In] Int[1/((c + d*x)^2*(a + a*Coth[e + f*x])), x]

[Out] $(f \text{Cosh}[2e - (2*c*f)/d] * \text{CoshIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2) - 1/(d*(c + d*x)*(a + a*Coth[e + f*x])) - (f * \text{CoshIntegral}[(2*c*f)/d + 2*f*x] * \text{Sinh}[2e - (2*c*f)/d])/(a*d^2) - (f * \text{Cosh}[2e - (2*c*f)/d] * \text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2) + (f * \text{Sinh}[2e - (2*c*f)/d] * \text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(a*d^2)$

Rule 3379

Int[sin[(e_.) + (Complex[0, fz_])* (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]

Rule 3382

Int[sin[(e_.) + (Complex[0, fz_])* (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]

Rule 3384

Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]

Rule 3805

Int[1/(((c_.) + (d_.)*(x_.))^2*((a_) + (b_)*tan[(e_.) + (f_.)*(x_.)])), x_Symbol] :> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Dist[f/(a*d), Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Dist[f/(b*d), Int[Cos[2*e + 2*f*x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{1}{d(c+dx)(a+a \coth(e+fx))} \\
&\quad - \frac{(if) \int \frac{\sin(2(ie+\frac{\pi}{2})+2ifx)}{c+dx} dx}{ad} - \frac{f \int \frac{\cos(2(ie+\frac{\pi}{2})+2ifx)}{c+dx} dx}{ad} \\
&= -\frac{1}{d(c+dx)(a+a \coth(e+fx))} + \frac{(f \cosh(2e-\frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
&\quad - \frac{(f \cosh(2e-\frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
&\quad - \frac{(f \sinh(2e-\frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} + \frac{(f \sinh(2e-\frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad} \\
&= \frac{f \cosh(2e-\frac{2cf}{d}) \text{Chi}(\frac{2cf}{d}+2fx)}{ad^2} - \frac{1}{d(c+dx)(a+a \coth(e+fx))} \\
&\quad - \frac{f \text{Chi}(\frac{2cf}{d}+2fx) \sinh(2e-\frac{2cf}{d})}{ad^2} - \frac{f \cosh(2e-\frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{ad^2} \\
&\quad + \frac{f \sinh(2e-\frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{ad^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.93 (sec), antiderivative size = 206, normalized size of antiderivative = 1.30

$$\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))} dx =$$

$$\text{csch}(e+fx) (\cosh(\frac{cf}{d}) + \sinh(\frac{cf}{d})) \left(d(\cosh(e+f(-\frac{c}{d}+x)) - \cosh(e+f(\frac{c}{d}+x))) + \sinh(e+f(-\frac{c}{d}+x)) \right)$$

```

[In] Integrate[1/((c + d*x)^2*(a + a*Coth[e + f*x])), x]
[Out] -1/2*(Csch[e + f*x]*(Cosh[(c*f)/d] + Sinh[(c*f)/d])*(d*(Cosh[e + f*(-(c/d)
+x)] - Cosh[e + f*(c/d + x)] + Sinh[e + f*(-(c/d) + x)] + Sinh[e + f*(c/d
+x)]) + 2*f*(c + d*x)*CoshIntegral[(2*f*(c + d*x))/d]*(-Cosh[e - (f*(c + d
*x))/d] + Sinh[e - (f*(c + d*x))/d]) + 2*f*(c + d*x)*(Cosh[e - (f*(c + d*x))
/d] - Sinh[e - (f*(c + d*x))/d])*SinhIntegral[(2*f*(c + d*x))/d]))/(a*d^2*
(c + d*x)*(1 + Coth[e + f*x]))

```

Maple [A] (verified)

Time = 0.36 (sec) , antiderivative size = 91, normalized size of antiderivative = 0.57

method	result	size
risch	$-\frac{1}{2da(dx+c)} + \frac{f e^{-2fx-2e}}{2ad(dx+f+c)} - \frac{f e^{\frac{2cf-2de}{d}} \text{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{a d^2}$	91

[In] `int(1/(d*x+c)^2/(a+a*cOTH(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/d/a/(d*x+c)+1/2*f/a*\exp(-2*f*x-2*e)/d/(d*f*x+c*f)-f/a/d^2*\exp(2*(c*f-d*e))/d)*\text{Ei}(1,2*f*x+2*e+2*(c*f-d*e))/d$$

Fricas [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 216, normalized size of antiderivative = 1.36

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+a\coth(e+fx))} dx \\ &= \frac{(dfx+cf)\text{Ei}\left(-\frac{2(df+cf)}{d}\right) \cosh(fx+e) \cosh\left(-\frac{2(de-cf)}{d}\right) + (dfx+cf)\text{Ei}\left(-\frac{2(df+cf)}{d}\right) \cosh(fx+e) \sinh\left(-\frac{2(de-cf)}{d}\right)}{(ad^3x+acd^2)\cosh} \end{aligned}$$

[In] `integrate(1/(d*x+c)^2/(a+a*cOTH(f*x+e)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & ((d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(f*x + e)*\cosh(-2*(d*e - c*f)/d) \\ & + (d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(f*x + e)*\sinh(-2*(d*e - c*f)/d) \\ & + ((d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\cosh(-2*(d*e - c*f)/d) + (d*f*x + c*f)*\text{Ei}(-2*(d*f*x + c*f)/d)*\sinh(-2*(d*e - c*f)/d) - d)*\sinh(f*x + e))/((a*d^3*x + a*c*d^2)*\cosh(f*x + e) + (a*d^3*x + a*c*d^2)*\sinh(f*x + e)) \end{aligned}$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c+dx)^2(a+a\coth(e+fx))} dx \\ &= \frac{\int \frac{1}{c^2\coth(e+fx)+c^2+2cdx\coth(e+fx)+2cdx+d^2x^2\coth(e+fx)+d^2x^2} dx}{a} \end{aligned}$$

[In] `integrate(1/(d*x+c)**2/(a+a*cOTH(f*x+e)),x)`

[Out] `Integral(1/(c**2*cOTH(e + f*x) + c**2 + 2*c*d*x*cOTH(e + f*x) + 2*c*d*x + d**2*x**2*cOTH(e + f*x) + d**2*x**2), x)/a`

Maxima [A] (verification not implemented)

none

Time = 0.47 (sec) , antiderivative size = 56, normalized size of antiderivative = 0.35

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))} dx = -\frac{1}{2(ad^2x + acd)} + \frac{e^{(-2e + \frac{2cf}{d})} E_2\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)ad}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e)),x, algorithm="maxima")
[Out] -1/2/(a*d^2*x + a*c*d) + 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(2, 2*(d*x + c)*f/d)/((d*x + c)*a*d)
```

Giac [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 321 vs. 2(159) = 318.

Time = 0.31 (sec) , antiderivative size = 321, normalized size of antiderivative = 2.02

$$\begin{aligned} & \int \frac{1}{(c + dx)^2(a + a \coth(e + fx))} dx \\ &= \frac{\left(2(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right)f^2 \text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right) e^{-\frac{2(de-cf)}{d}} - 2def^2 \text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right) e^{-\frac{2(de-cf)}{d}}\right)}{2((dx+c)ad)} \end{aligned}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e)),x, algorithm="giac")
[Out] 1/2*(2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 2*d*e*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) - 2*c*f^3*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*c*f^2*d*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) + d*f^2*Ei(-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d)*e^(-2*(d*e - c*f)/d) - d*f^2*d^2/(((d*x + c)*a*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a*d^5*e + a*c*d^4*f)*f)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))} dx = \int \frac{1}{(a + a \coth(e + fx)) (c + dx)^2} dx$$

[In] `int(1/((a + a*coth(e + f*x))*(c + d*x)^2),x)`

[Out] `int(1/((a + a*coth(e + f*x))*(c + d*x)^2), x)`

3.21 $\int \frac{1}{(c+dx)^3(a+a \coth(e+fx))} dx$

Optimal result	139
Rubi [A] (verified)	140
Mathematica [A] (verified)	142
Maple [A] (verified)	142
Fricas [A] (verification not implemented)	142
Sympy [F]	143
Maxima [A] (verification not implemented)	143
Giac [A] (verification not implemented)	144
Mupad [F(-1)]	144

Optimal result

Integrand size = 20, antiderivative size = 211

$$\begin{aligned} \int \frac{1}{(c+dx)^3(a+a \coth(e+fx))} dx = & -\frac{f}{2ad^2(c+dx)} - \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & - \frac{1}{2d(c+dx)^2(a+a \coth(e+fx))} \\ & + \frac{f}{d^2(c+dx)(a+a \coth(e+fx))} \\ & + \frac{f^2 \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{ad^3} \\ & + \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^3} \\ & - \frac{f^2 \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{ad^3} \end{aligned}$$

```
[Out] -1/2*f/a/d^2/(d*x+c)-f^2*Chi(2*c*f/d+2*f*x)*cosh(-2*e+2*c*f/d)/a/d^3-1/2/d/(d*x+c)^2/(a+a*coth(f*x+e))+f/d^2/(d*x+c)/(a+a*coth(f*x+e))+f^2*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d+2*f*x)/a/d^3-f^2*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^3+f^2*Shi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a/d^3
```

Rubi [A] (verified)

Time = 0.24 (sec) , antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.250, Rules used = {3806, 3805, 3384, 3379, 3382}

$$\int \frac{1}{(c+dx)^3(a+a\coth(e+fx))} dx = \frac{f^2 \text{Chi}\left(2xf + \frac{2cf}{d}\right) \sinh\left(2e - \frac{2cf}{d}\right)}{ad^3} - \frac{f^2 \text{Chi}\left(2xf + \frac{2cf}{d}\right) \cosh\left(2e - \frac{2cf}{d}\right)}{ad^3} - \frac{f^2 \sinh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{ad^3} + \frac{f^2 \cosh\left(2e - \frac{2cf}{d}\right) \text{Shi}\left(2xf + \frac{2cf}{d}\right)}{ad^3} + \frac{f}{d^2(c+dx)(a\coth(e+fx)+a)} - \frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a\coth(e+fx)+a)}$$

[In] $\text{Int}[1/((c+d*x)^3*(a+a*\text{Coth}[e+f*x])), x]$

[Out] $-1/2*f/(a*d^2*(c+d*x)) - (f^2*\text{Cosh}[2*e - (2*c*f)/d]*\text{CoshIntegral}[(2*c*f)/d + 2*f*x]/(a*d^3) - 1/(2*d*(c+d*x))^2*(a+a*\text{Coth}[e+f*x])) + f/(d^2*(c+d*x)*(a+a*\text{Coth}[e+f*x])) + (f^2*\text{CoshIntegral}[(2*c*f)/d + 2*f*x]*\text{Sinh}[2*e - (2*c*f)/d]/(a*d^3) + (f^2*\text{Cosh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x]/(a*d^3) - (f^2*\text{Sinh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x]/(a*d^3))$

Rule 3379

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Simp}[I*(\text{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d, x) /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&& \text{EqQ}[d*e - c*f*fz*I, 0]]$

Rule 3382

$\text{Int}[\sin[(e_.) + (\text{Complex}[0, fz_])*(f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&& \text{EqQ}[d*(e - \pi/2) - c*f*fz*I, 0]]$

Rule 3384

$\text{Int}[\sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \text{Dist}[\text{Cos}[(d*e - c*f)/d], \text{Int}[\text{Sin}[c*(f/d) + f*x]/(c+d*x), x], x] + \text{Dist}[\text{Sin}[(d*e - c*f)/d], \text{Int}[\text{Cos}[c*(f/d) + f*x]/(c+d*x), x], x] /; \text{FreeQ}[\{c, d, e, f\}, x] \&& \text{NeQ}[d*e - c*f, 0]]$

Rule 3805

```
Int[1/(((c_) + (d_)*(x_))^2*((a_) + (b_)*tan[(e_) + (f_)*(x_)])), x_Sy
mbol] :> -Simp[(d*(c + d*x)*(a + b*Tan[e + f*x]))^(-1), x] + (-Dist[f/(a*d)
, Int[Sin[2*e + 2*f*x]/(c + d*x), x], x] + Dist[f/(b*d), Int[Cos[2*e + 2*f*
x]/(c + d*x), x], x]) /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0]
```

Rule 3806

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + (f_)*(x_)]), x_Sym
bol] :> Simplify[f*((c + d*x)^(m + 2)/(b*d^2*(m + 1)*(m + 2))), x] + (Dist[2*b*
(f/(a*d*(m + 1))), Int[(c + d*x)^(m + 1)/(a + b*Tan[e + f*x]), x], x] + Sim
p[(c + d*x)^(m + 1)/(d*(m + 1)*(a + b*Tan[e + f*x])), x]) /; FreeQ[{a, b, c
, d, e, f}, x] && EqQ[a^2 + b^2, 0] && LtQ[m, -1] && NeQ[m, -2]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\coth(e+fx))} - \frac{\int \frac{1}{(c+dx)^2(a+a\coth(e+fx))} dx}{d} \\
&= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\coth(e+fx))} \\
&\quad + \frac{f}{d^2(c+dx)(a+a\coth(e+fx))} \\
&\quad + \frac{(if^2) \int \frac{\sin(2(ie+\frac{\pi}{2})+2ifx)}{c+dx} dx}{ad^2} + \frac{f^2 \int \frac{\cos(2(ie+\frac{\pi}{2})+2ifx)}{c+dx} dx}{ad^2} \\
&= -\frac{f}{2ad^2(c+dx)} - \frac{1}{2d(c+dx)^2(a+a\coth(e+fx))} + \frac{f}{d^2(c+dx)(a+a\coth(e+fx))} \\
&\quad - \frac{(f^2 \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} + \frac{(f^2 \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&\quad + \frac{(f^2 \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} - \frac{(f^2 \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{ad^2} \\
&= -\frac{f}{2ad^2(c+dx)} - \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d}+2fx)}{ad^3} \\
&\quad - \frac{1}{2d(c+dx)^2(a+a\coth(e+fx))} + \frac{f}{d^2(c+dx)(a+a\coth(e+fx))} \\
&\quad + \frac{f^2 \text{Chi}(\frac{2cf}{d}+2fx) \sinh(2e - \frac{2cf}{d})}{ad^3} + \frac{f^2 \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{ad^3} \\
&\quad - \frac{f^2 \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{ad^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.23 (sec) , antiderivative size = 265, normalized size of antiderivative = 1.26

$$\int \frac{1}{(c+dx)^3(a+a\coth(e+fx))} dx =$$

$$-\frac{\text{csch}(e+fx) (\cosh(\frac{cf}{d}) + \sinh(\frac{cf}{d})) \left(d(d \cosh(e+f(-\frac{c}{d}+x)) + (-d+2cf+2dfx) \cosh(e+f(\frac{c}{d}+$$

[In] `Integrate[1/((c + d*x)^3*(a + a*Coth[e + f*x])),x]`

[Out]
$$\begin{aligned} & -1/4 * (\text{Csch}[e + f*x] * (\text{Cosh}[(c*f)/d] + \text{Sinh}[(c*f)/d]) * (d * (d * \text{Cosh}[e + f*(-(c/d) + x)] + (-d + 2*c*f + 2*d*f*x) * \text{Cosh}[e + f*(c/d + x)] + d * \text{Sinh}[e + f*(-(c/d) + x)] + d * \text{Sinh}[e + f*(c/d + x)] - 2*c*f * \text{Sinh}[e + f*(c/d + x)] - 2*d*f*x * \text{Sinh}[e + f*(c/d + x)]) + 4*f^2 * 2*(c + d*x)^2 * \text{CoshIntegral}[(2*f*(c + d*x))/d] * (\text{Cosh}[e - (f*(c + d*x))/d] - \text{Sinh}[e - (f*(c + d*x))/d]) + 4*f^2 * 2*(c + d*x)^2 * (-\text{Cosh}[e - (f*(c + d*x))/d] + \text{Sinh}[e - (f*(c + d*x))/d]) * \text{SinhIntegral}[(2*f*(c + d*x))/d])) / (a*d^3*(c + d*x)^2*(1 + \text{Coth}[e + f*x])) \end{aligned}$$

Maple [A] (verified)

Time = 0.40 (sec) , antiderivative size = 210, normalized size of antiderivative = 1.00

method	result
risch	$-\frac{1}{4da(dx+c)^2} - \frac{f^3 e^{-2fx-2e} x}{2ad(d^2x^2f^2+2cd f^2x+c^2f^2)} - \frac{f^3 e^{-2fx-2e} c}{2a d^2(d^2x^2f^2+2cd f^2x+c^2f^2)} + \frac{f^2 e^{-2fx-2e}}{4ad(d^2x^2f^2+2cd f^2x+c^2f^2)} + \frac{f^2 e^{\frac{2cf-2de}{d}}}{4ad(d^2x^2f^2+2cd f^2x+c^2f^2)}$

[In] `int(1/(d*x+c)^3/(a+a*coth(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/4/d/a/(d*x+c)^2 - 1/2*f^3/a*\exp(-2*f*x-2*e)/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*x - 1/2*f^3/a*\exp(-2*f*x-2*e)/d^2/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2)*c + 1/4*f^2/a*\exp(-2*f*x-2*e)/d/(d^2*f^2*x^2+2*c*d*f^2*x+c^2*f^2) + f^2/a/d^3*\exp(2*(c*f-d*e)/d)*\text{Ei}(1, 2*f*x+2*e+2*(c*f-d*e)/d) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 342, normalized size of antiderivative = 1.62

$$\int \frac{1}{(c+dx)^3(a+a\coth(e+fx))} dx =$$

$$-\frac{2(d^2f^2x^2+2cdf^2x+c^2f^2)\text{Ei}\left(-\frac{2(df+cf)}{d}\right) \cosh(fx+e) \sinh\left(-\frac{2(de-cf)}{d}\right) + \left(d^2fx+cdf+2(d^2f^2x^2+$$

[In] `integrate(1/(d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -\frac{1}{2} \left(2(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2) Ei(-2(d f x + c f)/d) \cosh(f x + e) \sinh(-2(d e - c f)/d) + (d^2 f x + c d f + 2(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2)) Ei(-2(d f x + c f)/d) \cosh(-2(d e - c f)/d) \cosh(f x + e) - (d^2 f x + c d f - 2(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2)) Ei(-2(d f x + c f)/d) \cosh(-2(d e - c f)/d) - 2(d^2 f^2 x^2 + 2 c d f^2 x + c^2 f^2) Ei(-2(d f x + c f)/d) \sinh(-2(d e - c f)/d) - d^2) \sinh(f x + e) \right) / ((a d^5 x^2 + 2 a c d^4 x + a c^2 d^3) \cosh(f x + e) + (a d^5 x^2 + 2 a c d^4 x + a c^2 d^3) \sinh(f x + e)) \end{aligned}$$

Sympy [F]

$$\begin{aligned} & \int \frac{1}{(c + dx)^3(a + a \coth(e + fx))} dx \\ &= \frac{\int \frac{1}{c^3 \coth(e+fx)+c^3+3c^2dx \coth(e+fx)+3c^2dx+3cd^2x^2 \coth(e+fx)+3cd^2x^2+d^3x^3 \coth(e+fx)+d^3x^3} dx}{a} \end{aligned}$$

[In] `integrate(1/(d*x+c)**3/(a+a*coth(f*x+e)),x)`

[Out]
$$\text{Integral}\left(\frac{1}{c^3 \coth(e+fx)+c^3+3c^2dx \coth(e+fx)+3c^2dx+3cd^2x^2 \coth(e+fx)+3cd^2x^2+d^3x^3 \coth(e+fx)+d^3x^3}, x\right)/a$$

Maxima [A] (verification not implemented)

none

Time = 0.65 (sec), antiderivative size = 68, normalized size of antiderivative = 0.32

$$\begin{aligned} & \int \frac{1}{(c + dx)^3(a + a \coth(e + fx))} dx \\ &= -\frac{1}{4(ad^3 x^2 + 2acd^2 x + ac^2 d)} + \frac{e^{(-2e+\frac{2cf}{d})} E_3\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)^2 ad} \end{aligned}$$

[In] `integrate(1/(d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -\frac{1}{4}(a d^3 x^2 + 2 a c d^2 x + a c^2 d) + \frac{1}{2} e^{-2e + 2c f/d} \exp_integ\\ & ral_e(3, 2(d x + c) f / d) / ((d x + c)^2 a d) \end{aligned}$$

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 175, normalized size of antiderivative = 0.83

$$\int \frac{1}{(c + dx)^3(a + a \coth(e + fx))} dx =$$

$$-\frac{4 d^2 f^2 x^2 \text{Ei}\left(-\frac{2 (dfx+cf)}{d}\right) e^{\left(\frac{2 cf}{d}\right)} + 8 c d f^2 x \text{Ei}\left(-\frac{2 (dfx+cf)}{d}\right) e^{\left(\frac{2 cf}{d}\right)} + 4 c^2 f^2 \text{Ei}\left(-\frac{2 (dfx+cf)}{d}\right) e^{\left(\frac{2 cf}{d}\right)} + 2 d^2 f x e^{\left(\frac{2 cf}{d}\right)}}{4 (ad^5 x^2 e^{(2e)} + 2acd^4 x e^{(2e)} + ac^2 d^3 e^{(2e)})}$$

[In] `integrate(1/(d*x+c)^3/(a+a*coth(f*x+e)),x, algorithm="giac")`

[Out]
$$\begin{aligned} & -1/4*(4*d^2*f^2*x^2*Ei(-2*(d*f*x + c*f)/d)*e^{(2*c*f/d)} + 8*c*d*f^2*x*Ei(-2*(d*f*x + c*f)/d)*e^{(2*c*f/d)} + 4*c^2*f^2*x*Ei(-2*(d*f*x + c*f)/d)*e^{(2*c*f/d)} + 2*d^2*f*x*e^{(-2*f*x)} + 2*c*d*f*e^{(-2*f*x)} - d^2*e^{(-2*f*x)} + d^2*e^{(2*e)})/(a*d^5*x^2*e^{(2*e)} + 2*a*c*d^4*x*e^{(2*e)} + a*c^2*d^3*e^{(2*e)}) \end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^3(a + a \coth(e + fx))} dx = \int \frac{1}{(a + a \coth(e + fx)) (c + dx)^3} dx$$

[In] `int(1/((a + a*coth(e + f*x))*(c + d*x)^3),x)`

[Out] `int(1/((a + a*coth(e + f*x))*(c + d*x)^3), x)`

3.22 $\int \frac{(c+dx)^3}{(a+a \coth(e+fx))^2} dx$

Optimal result	145
Rubi [A] (verified)	145
Mathematica [A] (verified)	147
Maple [A] (verified)	148
Fricas [B] (verification not implemented)	148
Sympy [B] (verification not implemented)	149
Maxima [A] (verification not implemented)	150
Giac [A] (verification not implemented)	151
Mupad [B] (verification not implemented)	151

Optimal result

Integrand size = 20, antiderivative size = 230

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+a \coth(e+fx))^2} dx = & -\frac{3d^3 e^{-4e-4fx}}{512a^2 f^4} + \frac{3d^3 e^{-2e-2fx}}{16a^2 f^4} - \frac{3d^2 e^{-4e-4fx}(c+dx)}{128a^2 f^3} \\ & + \frac{3d^2 e^{-2e-2fx}(c+dx)}{8a^2 f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2 f^2} \\ & + \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2 f^2} - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2 f} \\ & + \frac{e^{-2e-2fx}(c+dx)^3}{4a^2 f} + \frac{(c+dx)^4}{16a^2 d} \end{aligned}$$

```
[Out] -3/512*d^3*exp(-4*f*x-4*e)/a^2/f^4+3/16*d^3*exp(-2*f*x-2*e)/a^2/f^4-3/128*d^2*exp(-4*f*x-4*e)*(d*x+c)/a^2/f^3+3/8*d^2*exp(-2*f*x-2*e)*(d*x+c)/a^2/f^3-3/64*d*exp(-4*f*x-4*e)*(d*x+c)^2/a^2/f^2+3/8*d*exp(-2*f*x-2*e)*(d*x+c)^2/a^2/f^2-1/16*exp(-4*f*x-4*e)*(d*x+c)^3/a^2/f+1/4*exp(-2*f*x-2*e)*(d*x+c)^3/a^2/f+1/16*(d*x+c)^4/a^2/d
```

Rubi [A] (verified)

Time = 0.20 (sec) , antiderivative size = 230, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used

$$= \{3810, 2207, 2225\}$$

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+a \coth(e+fx))^2} dx = & -\frac{3d^2(c+dx)e^{-4e-4fx}}{128a^2f^3} + \frac{3d^2(c+dx)e^{-2e-2fx}}{8a^2f^3} \\ & -\frac{3d(c+dx)^2e^{-4e-4fx}}{64a^2f^2} + \frac{3d(c+dx)^2e^{-2e-2fx}}{8a^2f^2} \\ & -\frac{(c+dx)^3e^{-4e-4fx}}{16a^2f} + \frac{(c+dx)^3e^{-2e-2fx}}{4a^2f} \\ & +\frac{(c+dx)^4}{16a^2d} - \frac{3d^3e^{-4e-4fx}}{512a^2f^4} + \frac{3d^3e^{-2e-2fx}}{16a^2f^4} \end{aligned}$$

[In] Int[(c + d*x)^3/(a + a*Coth[e + f*x])^2, x]

[Out] $(-3d^3E^{-4e-4fx})/(512a^2f^4) + (3d^3E^{-2e-2fx})/(16a^2f^4) - (3d^2E^{-4e-4fx}*(c+d*x))/(128a^2f^3) + (3d^2E^{-2e-2fx}*(c+d*x))/(8a^2f^3) - (3dE^{-4e-4fx}*(c+d*x)^2)/(64a^2f^2) + (3dE^{-2e-2fx}*(c+d*x)^2)/(8a^2f^2) - (E^{-4e-4fx}*(c+d*x)^3)/(16a^2f) + (E^{-2e-2fx}*(c+d*x)^3)/(4a^2f) + (c+d*x)^4/(16a^2d)$

Rule 2207

Int[((b_)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^m_, x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[\$UseGamma]

Rule 2225

Int[((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]

Rule 3810

Int[((c_.) + (d_.)*(x_.))^m*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x)))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^3}{4a^2} + \frac{e^{-4e-4fx}(c+dx)^3}{4a^2} - \frac{e^{-2e-2fx}(c+dx)^3}{2a^2} \right) dx \\ &= \frac{(c+dx)^4}{16a^2d} + \frac{\int e^{-4e-4fx}(c+dx)^3 dx}{4a^2} - \frac{\int e^{-2e-2fx}(c+dx)^3 dx}{2a^2} \end{aligned}$$

$$\begin{aligned}
&= -\frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d} \\
&\quad + \frac{(3d)\int e^{-4e-4fx}(c+dx)^2 dx}{16a^2f} - \frac{(3d)\int e^{-2e-2fx}(c+dx)^2 dx}{4a^2f} \\
&= -\frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} + \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} \\
&\quad - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d} \\
&\quad + \frac{(3d^2)\int e^{-4e-4fx}(c+dx) dx}{32a^2f^2} - \frac{(3d^2)\int e^{-2e-2fx}(c+dx) dx}{4a^2f^2} \\
&= -\frac{3d^2e^{-4e-4fx}(c+dx)}{128a^2f^3} + \frac{3d^2e^{-2e-2fx}(c+dx)}{8a^2f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} \\
&\quad + \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} \\
&\quad + \frac{(c+dx)^4}{16a^2d} + \frac{(3d^3)\int e^{-4e-4fx} dx}{128a^2f^3} - \frac{(3d^3)\int e^{-2e-2fx} dx}{8a^2f^3} \\
&= -\frac{3d^3e^{-4e-4fx}}{512a^2f^4} + \frac{3d^3e^{-2e-2fx}}{16a^2f^4} - \frac{3d^2e^{-4e-4fx}(c+dx)}{128a^2f^3} \\
&\quad + \frac{3d^2e^{-2e-2fx}(c+dx)}{8a^2f^3} - \frac{3de^{-4e-4fx}(c+dx)^2}{64a^2f^2} + \frac{3de^{-2e-2fx}(c+dx)^2}{8a^2f^2} \\
&\quad - \frac{e^{-4e-4fx}(c+dx)^3}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^3}{4a^2f} + \frac{(c+dx)^4}{16a^2d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.91 (sec), antiderivative size = 420, normalized size of antiderivative = 1.83

$$\begin{aligned}
&\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^2} dx \\
&= \frac{\text{csch}^2(e+fx)(\cosh(fx)+\sinh(fx))^2 ((4c^3f^3+6c^2df^2(1+2fx)+6cd^2f(1+2fx+2f^2x^2)+d^3(3+6fx^2))}{}
\end{aligned}$$

```

[In] Integrate[(c + d*x)^3/(a + a*Coth[e + f*x])^2, x]
[Out] (Csch[e + f*x]^2*(Cosh[f*x] + Sinh[f*x])^2*((4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*Cosh[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(1 + 4*f*x) + 12*c*d^2*f*(1 + 4*f*x + 8*f^2*x^2) + d^3*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3))*Cos[h[4*f*x]*(-Cosh[2*e] + Sinh[2*e]))/32 + f^4*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*(Cosh[2*e] + Sinh[2*e]) - (4*c^3*f^3 + 6*c^2*d*f^2*(1 + 2*f*x) + 6*c*d^2*f*(1 + 2*f*x + 2*f^2*x^2) + d^3*(3 + 6*f*x + 6*f^2*x^2 + 4*f^3*x^3))*Sinh[2*f*x] + ((32*c^3*f^3 + 24*c^2*d*f^2*(1 + 4*f*x) + 12*c*d^2*f*(1 + 4*f*x + 8*f^2*x^2) + d^3*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3))*(Cosh[2*e] - Sinh[2*e])*Sinh[4*f*x])/32)/(16*a^2*f^4*(1 + Coth[e + f*x])^2)

```

Maple [A] (verified)

Time = 0.50 (sec) , antiderivative size = 273, normalized size of antiderivative = 1.19

method	result
risch	$\frac{d^3x^4}{16a^2} + \frac{d^2cx^3}{4a^2} + \frac{3d^2x^2}{8a^2} + \frac{c^3x}{4a^2} + \frac{c^4}{16a^2} + \frac{(4d^3x^3f^3+12cd^2f^3x^2+12c^2df^3x+6d^3f^2x^2+4c^3f^3+12cd^2f^2x+6c^2df^2+6c^4)}{16a^2f^4}$
parallelrisch	$\frac{32xf\left(\left(\frac{1}{4}d^3x^3+cd^2x^2+\frac{3}{2}c^2dx+c^3\right)f^3-\frac{15d\left(\frac{1}{3}x^2d^2+cdx+c^2\right)f^2}{4}-\frac{27d^2\left(\frac{d}{2}x+c\right)f}{8}-\frac{51d^3}{32}\right)\tanh(fx+e)^2+\left((16d^3x^4+64d^2cx^3+96cd^3x^2+64c^4)x^2+16cd^2x^3+16c^3x^2+16c^4\right)\tanh(fx+e)+\left(16d^3x^4+64d^2cx^3+96cd^3x^2+64c^4\right)\tanh(fx+e)^3+16c^4\right)}{16a^2f^4}$

```
[In] int((d*x+c)^3/(a+a*cOTH(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] 1/16/a^2*d^3*x^4+1/4/a^2*d^2*c*x^3+3/8/a^2*d*c^2*x^2+1/4/a^2*c^3*x+1/16/a^2
/d*c^4+1/16*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4*
c^3*f^3*x+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+3*d^3)/a^2/f^4*exp(-
2*f*x-2*e)-1/512*(32*d^3*f^3*x^3+96*c*d^2*f^3*x^2+96*c^2*d*f^3*x+24*d^3*f^2
*x^2+32*c^3*f^3+48*c*d^2*f^2*x+24*c^2*d*f^2+12*d^3*f*x+12*c*d^2*f+3*d^3)/a^
2/f^4*exp(-4*f*x-4*e)
```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 571 vs. 2(204) = 408.

Time = 0.25 (sec) , antiderivative size = 571, normalized size of antiderivative = 2.48

$$\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^2} dx \\ \equiv \frac{128 d^3 f^3 x^3 + 128 c^3 f^3 + 192 c^2 d f^2 + 192 c d^2 f + 96 d^3 + 192 (2 c d^2 f^3 + d^3 f^2) x^2 + (32 d^3 f^4 x^4 - 32 c^3 f^3 - 2 c^2 d^2 f^2 - 192 c d^2 f^2 - 192 c^2 d f^3 - 96 d^4)}{(a+a\coth(e+fx))^2}$$

```
[In] integrate((d*x+c)^3/(a+a*coth(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] 1/1512*(128*d^3*f^3*x^3 + 128*c^3*f^3 + 192*c^2*d*f^2 + 192*c*d^2*f + 96*d^3
 + 192*(2*c*d^2*f^3 + d^3*f^2)*x^2 + (32*d^3*f^4*x^4 - 32*c^3*f^3 - 24*c^2*
d*f^2 - 12*c*d^2*f + 32*(4*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 24*(8*c^2*d*f
^4 - 4*c*d^2*f^3 - d^3*f^2)*x^2 + 4*(32*c^3*f^4 - 24*c^2*d*f^3 - 12*c*d^2*f
^2 - 3*d^3*f)*x)*cosh(f*x + e)^2 + 2*(32*d^3*f^4*x^4 + 32*c^3*f^3 + 24*c^2*
d*f^2 + 12*c*d^2*f + 32*(4*c*d^2*f^4 + d^3*f^3)*x^3 + 3*d^3 + 24*(8*c^2*d*f
^4 + 4*c*d^2*f^3 + d^3*f^2)*x^2 + 4*(32*c^3*f^4 + 24*c^2*d*f^3 + 12*c*d^2*f
^2 + 3*d^3*f)*x)*cosh(f*x + e)*sinh(f*x + e) + (32*d^3*f^4*x^4 - 32*c^3*f^3
 - 24*c^2*d*f^2 - 12*c*d^2*f + 32*(4*c*d^2*f^4 - d^3*f^3)*x^3 - 3*d^3 + 24*
(8*c^2*d*f^4 - 4*c*d^2*f^3 - d^3*f^2)*x^2 + 4*(32*c^3*f^4 - 24*c^2*d*f^3 -
12*c*d^2*f^2 - 3*d^3*f)*x)*sinh(f*x + e)^2 + 192*(2*c^2*d*f^3 + 2*c*d^2*f^2
 + d^3*f)*x)/(a^2*f^4*cosh(f*x + e)^2 + 2*a^2*f^4*cosh(f*x + e)*sinh(f*x +
e) + a^2*f^4*sinh(f*x + e)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2193 vs. $2(236) = 472$.

Time = 1.06 (sec) , antiderivative size = 2193, normalized size of antiderivative = 9.53

$$\int \frac{(c + dx)^3}{(a + a \coth(e + fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)**3/(a+a*coth(f*x+e))**2,x)`

[Out] Piecewise((32*c**3*f**4*x*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 64*c**3*f**4*x*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 32*c**3*f**4*x/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 96*c**3*f**3*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 64*c**3*f**3/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 48*c**2*d*f**4*x**2*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 96*c**2*d*f**4*x**2*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 48*c**2*d*f**4*x**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 120*c**2*d*f**3*x*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 48*c**2*d*f**3*x*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 72*c**2*d*f**3*x/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 120*c**2*d*f**2*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 96*c**2*d*f**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 32*c*d**2*f**4*x**3*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 64*c*d**2*f**4*x**3*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 32*c*d**2*f**4*x**3/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 120*c*d**2*f**3*x**2*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 48*c*d**2*f**3*x**2*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 72*c*d**2*f**3*x**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 108*c*d**2*f**2*x*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 24*c*d**2*f**2*x*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 84*c*d**2*f**2*x/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 96*c*d**2*f/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 8*d**3*f**4*x**4*tanh(e + f*x)

```

**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 16*d**3*f**4*x**4*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 8*d**3*f**4*x**4/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 40*d**3*f**3*x**3*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 16*d**3*f**3*x**3*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 24*d**3*f**3*x**3/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 54*d**3*f**2*x**2*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 12*d**3*f**2*x**2*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 42*d**3*f**2*x**2/(128*a**2*f**4*tanh(e + f*x)*2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) - 51*d**3*f*x*tanh(e + f*x)**2/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 6*d**3*f*x*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 128*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 45*d**3*f*x/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 51*d**3*tanh(e + f*x)/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4) + 48*d**3/(128*a**2*f**4*tanh(e + f*x)**2 + 256*a**2*f**4*tanh(e + f*x) + 128*a**2*f**4), Ne(f, 0)), ((c**3*x + 3*c**2*d*x**2/2 + c*d**2*x**3 + d**3*x**4/4)/(a*cOTH(e) + a)**2, True))

```

Maxima [A] (verification not implemented)

none

Time = 0.50 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.29

$$\begin{aligned}
\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^2} dx &= \frac{1}{16} c^3 \left(\frac{4(fx+e)}{a^2 f} + \frac{4e^{(-2fx-2e)} - e^{(-4fx-4e)}}{a^2 f} \right) \\
&+ \frac{3(8f^2x^2e^{(4e)} + 8(2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - (4fx+1)e^{(-4fx)})c^2de^{(-4e)}}{64a^2f^2} \\
&+ \frac{(32f^3x^3e^{(4e)} + 48(2f^2x^2e^{(2e)} + 2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - 3(8f^2x^2 + 4fx + 1)e^{(-4fx)})cd^2e^{(-4e)}}{128a^2f^3} \\
&+ \frac{(32f^4x^4e^{(4e)} + 32(4f^3x^3e^{(2e)} + 6f^2x^2e^{(2e)} + 6fxe^{(2e)} + 3e^{(2e)})e^{(-2fx)} - (32f^3x^3 + 24f^2x^2 + 12fx - 512a^2f^4)}{512a^2f^4}
\end{aligned}$$

[In] integrate((d*x+c)^3/(a+a*cOTH(f*x+e))^2,x, algorithm="maxima")

[Out] $\frac{1}{16}c^3(4*(f*x + e)/(a^2*f) + (4*e^{(-2*f*x - 2*e)} - e^{(-4*f*x - 4*e)})/(a^2*f)) + \frac{3}{64}*(8*f^2*x^2*e^{(4e)} + 8*(2*f*x*e^{(2e)} + e^{(2e)})*e^{(-2*f*x)} - (4*f*x + 1)*e^{(-4*f*x})*c^2*d*e^{(-4e)}/(a^2*f^2) + \frac{1}{128}*(32*f^3*x^3*e^{(4e)} + 48*(2*f^2*x^2*e^{(2e)} + 2*f*x*e^{(2e)} + e^{(2e)})*e^{(-2*f*x)} - 3*(8*f^2*x^2 + 4*f*x + 1)*e^{(-4*f*x})*c*d^2*e^{(-4e)}/(a^2*f^3) + \frac{1}{512}*(32*f^4*x^4*e^{(4e)} + 32*(4*f^3*x^3*e^{(2e)} + 6*f^2*x^2*e^{(2e)} + 6*fxe^{(2e)} + 3e^{(2e)})*e^{(-2fx)} - (32*f^3*x^3 + 24*f^2*x^2 + 12*fx - 512*a^2*f^4)/a^2*f^4$

$$*e)) * e^{-(-2*f*x)} - (32*f^3*x^3 + 24*f^2*x^2 + 12*f*x + 3)*e^{(-4*f*x)})*d^3*e^{(-4*e)}/(a^2*f^4)$$

Giac [A] (verification not implemented)

none

Time = 0.30 (sec), antiderivative size = 368, normalized size of antiderivative = 1.60

$$\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^2} dx \\ = \frac{(32 d^3 f^4 x^4 e^{(4 fx+4 e)} + 128 c d^2 f^4 x^3 e^{(4 fx+4 e)} + 192 c^2 d f^4 x^2 e^{(4 fx+4 e)} + 128 d^3 f^3 x^3 e^{(2 fx+2 e)} - 32 d^3 f^3 x^3 + 128 c^2 d^2 f^3 x^2 e^{(2 fx+2 e)} + 384 c d^2 f^3 x^2 e^{(2 fx+2 e)} + 192 d^3 f^2 x^2 e^{(2 fx+2 e)} - 96 c d^2 f^2 x^2 e^{(2 fx+2 e)} + 384 c^2 d f^2 x^2 e^{(2 fx+2 e)} + 192 d^3 f x^2 e^{(2 fx+2 e)} - 24 d^3 f^2 x^2 e^{(2 fx+2 e)} + 128 c^3 f^3 x e^{(2 fx+2 e)} + 384 c^2 d^2 f^3 x e^{(2 fx+2 e)} - 32 c^3 d^2 f^2 x e^{(2 fx+2 e)} + 192 d^3 f^2 x e^{(2 fx+2 e)} - 24 c^2 d^2 f^2 x e^{(2 fx+2 e)} - 12 d^3 f x^3 e^{(2 fx+2 e)} + 192 c^2 d^2 f^2 x^2 e^{(2 fx+2 e)} - 12 c^3 d^2 f^2 x^2 e^{(2 fx+2 e)} + 96 d^3 f^3 x e^{(2 fx+2 e)} - 3 d^3 f^3 x e^{(-4 fx-4 e)})/(a^2 f^4)}$$

[In] integrate((d*x+c)^3/(a+a*coth(f*x+e))^2,x, algorithm="giac")

[Out] $\frac{1}{512} (32 d^3 f^4 x^4 e^{(4 fx+4 e)} + 128 c d^2 f^4 x^3 e^{(4 fx+4 e)} + 192 c^2 d^2 f^4 x^2 e^{(4 fx+4 e)} + 128 d^3 f^3 x^3 e^{(2 fx+2 e)} - 32 d^3 f^3 x^3 + 128 c^2 d^2 f^3 x^2 e^{(2 fx+2 e)} + 384 c d^2 f^3 x^2 e^{(2 fx+2 e)} + 192 d^3 f^2 x^2 e^{(2 fx+2 e)} - 96 c d^2 f^2 x^2 e^{(2 fx+2 e)} + 384 c^2 d f^2 x^2 e^{(2 fx+2 e)} + 192 d^3 f x^2 e^{(2 fx+2 e)} - 24 d^3 f^2 x^2 e^{(2 fx+2 e)} + 128 c^3 f^3 x e^{(2 fx+2 e)} + 384 c^2 d^2 f^3 x e^{(2 fx+2 e)} - 32 c^3 d^2 f^2 x e^{(2 fx+2 e)} + 192 d^3 f^2 x e^{(2 fx+2 e)} - 24 c^2 d^2 f^2 x e^{(2 fx+2 e)} - 12 d^3 f x^3 e^{(2 fx+2 e)} + 192 c^2 d^2 f^2 x^2 e^{(2 fx+2 e)} - 12 c^3 d^2 f^2 x^2 e^{(2 fx+2 e)} + 96 d^3 f^3 x e^{(2 fx+2 e)} - 3 d^3 f^3 x e^{(-4 fx-4 e)})/(a^2 f^4)$

Mupad [B] (verification not implemented)

Time = 2.06 (sec), antiderivative size = 266, normalized size of antiderivative = 1.16

$$\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^2} dx = e^{-2e-2fx} \left(\frac{4 c^3 f^3 + 6 c^2 d f^2 + 6 c d^2 f + 3 d^3}{16 a^2 f^4} + \frac{d^3 x^3}{4 a^2 f} \right. \\ \left. + \frac{3 d x (2 c^2 f^2 + 2 c d f + d^2)}{8 a^2 f^3} + \frac{3 d^2 x^2 (d + 2 c f)}{8 a^2 f^2} \right) \\ - e^{-4e-4fx} \left(\frac{32 c^3 f^3 + 24 c^2 d f^2 + 12 c d^2 f + 3 d^3}{512 a^2 f^4} + \frac{d^3 x^3}{16 a^2 f} \right. \\ \left. + \frac{3 d x (8 c^2 f^2 + 4 c d f + d^2)}{128 a^2 f^3} + \frac{3 d^2 x^2 (d + 4 c f)}{64 a^2 f^2} \right) \\ + \frac{c^3 x}{4 a^2} + \frac{d^3 x^4}{16 a^2} + \frac{3 c^2 d x^2}{8 a^2} + \frac{c d^2 x^3}{4 a^2}$$

[In] int((c + d*x)^3/(a + a*coth(e + f*x))^2,x)

[Out] $\exp(-2e-2fx)*((3*d^3 + 4*c^3*f^3 + 6*c^2*d*f^2 + 6*c*d^2*f)/(16*a^2*f^4) + (d^3*x^3)/(4*a^2*f) + (3*d*x*(d^2 + 2*c^2*f^2 + 2*c*d*f))/(8*a^2*f^3))$

$$\begin{aligned} & + (3*d^2*x^2*(d + 2*c*f))/(8*a^2*f^2)) - \exp(-4*e - 4*f*x)*((3*d^3 + 32*c \\ & ^3*f^3 + 24*c^2*d*f^2 + 12*c*d^2*f)/(512*a^2*f^4) + (d^3*x^3)/(16*a^2*f) + \\ & (3*d*x*(d^2 + 8*c^2*f^2 + 4*c*d*f))/(128*a^2*f^3) + (3*d^2*x^2*(d + 4*c*f)) \\ & /(64*a^2*f^2)) + (c^3*x)/(4*a^2) + (d^3*x^4)/(16*a^2) + (3*c^2*d*x^2)/(8*a^2) \\ & + (c*d^2*x^3)/(4*a^2) \end{aligned}$$

3.23 $\int \frac{(c+dx)^2}{(a+a \coth(e+fx))^2} dx$

Optimal result	153
Rubi [A] (verified)	153
Mathematica [A] (verified)	155
Maple [A] (verified)	155
Fricas [B] (verification not implemented)	155
Sympy [B] (verification not implemented)	156
Maxima [A] (verification not implemented)	157
Giac [A] (verification not implemented)	157
Mupad [B] (verification not implemented)	158

Optimal result

Integrand size = 20, antiderivative size = 170

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+a \coth(e+fx))^2} dx = & -\frac{d^2 e^{-4e-4fx}}{128a^2 f^3} + \frac{d^2 e^{-2e-2fx}}{8a^2 f^3} - \frac{de^{-4e-4fx}(c+dx)}{32a^2 f^2} \\ & + \frac{de^{-2e-2fx}(c+dx)}{4a^2 f^2} - \frac{e^{-4e-4fx}(c+dx)^2}{16a^2 f} \\ & + \frac{e^{-2e-2fx}(c+dx)^2}{4a^2 f} + \frac{(c+dx)^3}{12a^2 d} \end{aligned}$$

[Out]
$$\begin{aligned} & -1/128*d^2*exp(-4*f*x-4*e)/a^2/f^3+1/8*d^2*exp(-2*f*x-2*e)/a^2/f^3-1/32*d*e \\ & xp(-4*f*x-4*e)*(d*x+c)/a^2/f^2+1/4*d*exp(-2*f*x-2*e)*(d*x+c)/a^2/f^2-1/16*e \\ & xp(-4*f*x-4*e)*(d*x+c)^2/a^2/f+1/4*exp(-2*f*x-2*e)*(d*x+c)^2/a^2/f+1/12*(d*x+c)^3/a^2/d \end{aligned}$$

Rubi [A] (verified)

Time = 0.14 (sec), antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used = {3810, 2207, 2225}

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+a \coth(e+fx))^2} dx = & -\frac{d(c+dx)e^{-4e-4fx}}{32a^2 f^2} + \frac{d(c+dx)e^{-2e-2fx}}{4a^2 f^2} - \frac{(c+dx)^2 e^{-4e-4fx}}{16a^2 f} \\ & + \frac{(c+dx)^2 e^{-2e-2fx}}{4a^2 f} + \frac{(c+dx)^3}{12a^2 d} - \frac{d^2 e^{-4e-4fx}}{128a^2 f^3} + \frac{d^2 e^{-2e-2fx}}{8a^2 f^3} \end{aligned}$$

[In] $\text{Int}[(c+d*x)^2/(a+a*\text{Coth}[e+f*x])^2, x]$

[Out]
$$\begin{aligned} & -1/128*(d^2*E^{-(-4*e-4*f*x)})/(a^2*f^3)+(d^2*E^{-(-2*e-2*f*x)})/(8*a^2*f^3) \\ & -(d*E^{-(-4*e-4*f*x)}*(c+d*x))/(32*a^2*f^2)+(d*E^{-(-2*e-2*f*x)}*(c+ \end{aligned}$$

$$d*x)/(4*a^2*f^2) - (E^{-4}*e - 4*f*x)*(c + d*x)^2/(16*a^2*f) + (E^{-2}*e - 2*f*x)*(c + d*x)^2/(4*a^2*f) + (c + d*x)^3/(12*a^2*d)$$

Rule 2207

```
Int[((b_)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^m_, x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x]; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^n_, x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x]; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```
Int[((c_.) + (d_.)*(x_.))^m*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x]; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^2}{4a^2} + \frac{e^{-4e-4fx}(c+dx)^2}{4a^2} - \frac{e^{-2e-2fx}(c+dx)^2}{2a^2} \right) dx \\ &= \frac{(c+dx)^3}{12a^2d} + \frac{\int e^{-4e-4fx}(c+dx)^2 dx}{4a^2} - \frac{\int e^{-2e-2fx}(c+dx)^2 dx}{2a^2} \\ &= -\frac{e^{-4e-4fx}(c+dx)^2}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^2}{4a^2f} + \frac{(c+dx)^3}{12a^2d} \\ &\quad + \frac{d \int e^{-4e-4fx}(c+dx) dx}{8a^2f} - \frac{d \int e^{-2e-2fx}(c+dx) dx}{2a^2f} \\ &= -\frac{de^{-4e-4fx}(c+dx)}{32a^2f^2} + \frac{de^{-2e-2fx}(c+dx)}{4a^2f^2} - \frac{e^{-4e-4fx}(c+dx)^2}{16a^2f} \\ &\quad + \frac{e^{-2e-2fx}(c+dx)^2}{4a^2f} + \frac{(c+dx)^3}{12a^2d} + \frac{d^2 \int e^{-4e-4fx} dx}{32a^2f^2} - \frac{d^2 \int e^{-2e-2fx} dx}{4a^2f^2} \\ &= -\frac{d^2 e^{-4e-4fx}}{128a^2f^3} + \frac{d^2 e^{-2e-2fx}}{8a^2f^3} - \frac{de^{-4e-4fx}(c+dx)}{32a^2f^2} + \frac{de^{-2e-2fx}(c+dx)}{4a^2f^2} \\ &\quad - \frac{e^{-4e-4fx}(c+dx)^2}{16a^2f} + \frac{e^{-2e-2fx}(c+dx)^2}{4a^2f} + \frac{(c+dx)^3}{12a^2d} \end{aligned}$$

Mathematica [A] (verified)

Time = 2.31 (sec) , antiderivative size = 207, normalized size of antiderivative = 1.22

$$\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx \\ = \frac{\text{csch}^2(e + fx) (48(2c^2 f^2 + 2cdf(1 + 2fx) + d^2(1 + 2fx + 2f^2 x^2)) + (24c^2 f^2(-1 + 4fx) + 12cdf(-1 - 4fx + 8f^2 x^2)) \text{Cosh}[2*(e + fx)] + (24*c^2*f^2*(1 + 4*f*x) + 12*c*d*f*(1 + 4*f*x + 8*f^2*x^2) + d^2*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3)) \text{Sinh}[2*(e + fx)])}{(384*a^2*f^3*(1 + \text{Coth}[e + fx])^2)}$$

[In] Integrate[(c + d*x)^2/(a + a*Coth[e + f*x])^2, x]

[Out] $\text{Csch}[e + f*x]^2 * (48*(2*c^2*f^2 + 2*c*d*f*(1 + 2*f*x) + d^2*(1 + 2*f*x + 2*f^2*x^2)) + (24*c^2*f^2*(-1 + 4*f*x) + 12*c*d*f*(-1 - 4*f*x + 8*f^2*x^2) + d^2*(-3 - 12*f*x - 24*f^2*x^2 + 32*f^3*x^3)) * \text{Cosh}[2*(e + f*x)] + (24*c^2*f^2*(1 + 4*f*x) + 12*c*d*f*(1 + 4*f*x + 8*f^2*x^2) + d^2*(3 + 12*f*x + 24*f^2*x^2 + 32*f^3*x^3)) * \text{Sinh}[2*(e + f*x)]) / (384*a^2*f^3*(1 + \text{Coth}[e + f*x])^2)$

Maple [A] (verified)

Time = 0.44 (sec) , antiderivative size = 163, normalized size of antiderivative = 0.96

method	result
risch	$\frac{d^2x^3}{12a^2} + \frac{dcx^2}{4a^2} + \frac{c^2x}{4a^2} + \frac{c^3}{12a^2d} + \frac{(2d^2x^2f^2 + 4cd^2f^2x + 2c^2f^2 + 2d^2fx + 2cdf + d^2)e^{-2fx-2e}}{8a^2f^3} - \frac{(8d^2x^2f^2 + 16cd^2f^2x + 8c^2f^2 + 16cd^2fx + 16c^2fx + 16c^3)/12}{12a^2}$
parallelisch	$\frac{24xf\left(\left(\frac{1}{3}x^2d^2 + cdx + c^2\right)f^2 - \frac{5d\left(\frac{dx}{2} + c\right)f}{2} - \frac{9d^2}{8}\right)\tanh(fx+e)^2 + (16(d^2x^3 + 3dcx^2 + 3c^2x)f^3 + 12(x^2d^2 + 2cdx + 6c^2)f^2 + 6(xd^2 + 2cdx + 6c^2)x)f^2 + 6(x^2d^2 + 2cdx + 6c^2)}{96f^3a^2(1 + \tanh(fx+e)^2)}$

[In] int((d*x+c)^2/(a+a*coth(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] $1/12/a^2*d^2*x^3 + 1/4/a^2*d*c*x^2 + 1/4/a^2*c^2*x + 1/12/a^2/d*c^3 + 1/8*(2*d^2*f^2*x^2 + 4*c*d*f^2*x^2 + 2*c^2*f^2*x^2 + 2*d^2*f^2*x^2 + 2*c*d*f^2 + d^4)/a^2/f^3*\exp(-2*f*x-2*e) - 1/128*(8*d^2*f^2*x^2 + 16*c*d*f^2*x^2 + 8*c^2*f^2*x^2 + 4*d^2*f^2*x^2 + 4*c*d*f^2 + d^4)/a^2/f^3*\exp(-4*f*x-4*e)$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 359 vs. 2(150) = 300.

Time = 0.27 (sec) , antiderivative size = 359, normalized size of antiderivative = 2.11

$$\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx \\ = \frac{96 d^2 f^2 x^2 + 96 c^2 f^2 + 96 cdf + (32 d^2 f^3 x^3 - 24 c^2 f^2 - 12 cdf + 24 (4 cdf^3 - d^2 f^2) x^2 - 3 d^2 + 12 (8 c^2 f^3 + 12 cdf^2) x)f^2}{(a + a \coth(e + fx))^2}$$

```
[In] integrate((d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="fricas")
[Out] 1/384*(96*d^2*f^2*x^2 + 96*c^2*f^2 + 96*c*d*f + (32*d^2*f^3*x^3 - 24*c^2*f^2
2 - 12*c*d*f + 24*(4*c*d*f^3 - d^2*f^2)*x^2 - 3*d^2 + 12*(8*c^2*f^3 - 4*c*d
*f^2 - d^2*f)*x)*cosh(f*x + e)^2 + 2*(32*d^2*f^3*x^3 + 24*c^2*f^2 + 12*c*d*
f + 24*(4*c*d*f^3 + d^2*f^2)*x^2 + 3*d^2 + 12*(8*c^2*f^3 + 4*c*d*f^2 + d^2*f
)*x)*cosh(f*x + e)*sinh(f*x + e) + (32*d^2*f^3*x^3 - 24*c^2*f^2 - 12*c*d*f
+ 24*(4*c*d*f^3 - d^2*f^2)*x^2 - 3*d^2 + 12*(8*c^2*f^3 - 4*c*d*f^2 - d^2*f
)*x)*sinh(f*x + e)^2 + 48*d^2 + 96*(2*c*d*f^2 + d^2*f)*x)/(a^2*f^3*cosh(f*x
+ e)^2 + 2*a^2*f^3*cosh(f*x + e)*sinh(f*x + e) + a^2*f^3*sinh(f*x + e)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1353 vs. $2(165) = 330$.

Time = 0.89 (sec), antiderivative size = 1353, normalized size of antiderivative = 7.96

$$\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)**2/(a+a*coth(f*x+e))**2,x)
```

```
[Out] Piecewise((24*c**2*f**3*x*tanh(e + f*x)**2/(96*a**2*f**3*tanh(e + f*x)**2 +
192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 48*c**2*f**3*x*tanh(e + f*x)
/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**
3) + 24*c**2*f**3*x/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e +
f*x) + 96*a**2*f**3) + 72*c**2*f**2*tanh(e + f*x)/(96*a**2*f**3*tanh(e + f
*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 48*c**2*f**2/(96*a**2*f**
3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 24*c*
d*f**3*x**2*tanh(e + f*x)**2/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f*
3*tanh(e + f*x) + 96*a**2*f**3) + 48*c*d*f**3*x**2*tanh(e + f*x)/(96*a**2*f**
3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 24*c*
d*f**3*x**2/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) +
96*a**2*f**3) - 60*c*d*f**2*x*tanh(e + f*x)**2/(96*a**2*f**3*tanh(e + f*x)*
*x^2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 24*c*d*f**2*x*tanh(e + f
*x)/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**
3) + 36*c*d*f**2*x/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e +
f*x) + 96*a**2*f**3) + 60*c*d*f*tanh(e + f*x)/(96*a**2*f**3*tanh(e + f*x)**2 +
192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 48*c*d*f/(96*a**2*f**3*tanh(e +
f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 8*d**2*f*
3*x**3*tanh(e + f*x)**2/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e +
f*x) + 96*a**2*f**3) + 16*d**2*f**3*x**3*tanh(e + f*x)/(96*a**2*f**3*tanh(e +
f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 8*d**2*f**3*x**3/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) -
30*d**2*f**2*x**2*tanh(e + f*x)**2/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 12*d**2*f**2*x**2*tanh(e
```

$$\begin{aligned}
& + f*x)/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a* \\
& *2*f**3) + 18*d**2*f**2*x**2/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3 \\
& *tanh(e + f*x) + 96*a**2*f**3) - 27*d**2*f*x*tanh(e + f*x)**2/(96*a**2*f**3 \\
& *tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 6*d**2*f*x \\
& *tanh(e + f*x)/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) \\
&) + 96*a**2*f**3) + 21*d**2*f*x/(96*a**2*f**3*tanh(e + f*x)**2 + 192*a**2*f \\
& **3*tanh(e + f*x) + 96*a**2*f**3) + 27*d**2*tanh(e + f*x)/(96*a**2*f**3*tan \\
& h(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3) + 24*d**2/(96*a \\
& **2*f**3*tanh(e + f*x)**2 + 192*a**2*f**3*tanh(e + f*x) + 96*a**2*f**3), Ne \\
& (f, 0)), ((c**2*x + c*d*x**2 + d**2*x**3/3)/(a*coth(e) + a)**2, True))
\end{aligned}$$

Maxima [A] (verification not implemented)

none

Time = 0.41 (sec) , antiderivative size = 191, normalized size of antiderivative = 1.12

$$\begin{aligned}
\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx &= \frac{1}{16} c^2 \left(\frac{4(fx + e)}{a^2 f} + \frac{4e^{(-2fx-2e)} - e^{(-4fx-4e)}}{a^2 f} \right) \\
&+ \frac{(8f^2x^2e^{(4e)} + 8(2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - (4fx + 1)e^{(-4fx)})cde^{(-4e)}}{32a^2f^2} \\
&+ \frac{(32f^3x^3e^{(4e)} + 48(2f^2x^2e^{(2e)} + 2fxe^{(2e)} + e^{(2e)})e^{(-2fx)} - 3(8f^2x^2 + 4fx + 1)e^{(-4fx)})d^2e^{(-4e)}}{384a^2f^3}
\end{aligned}$$

```
[In] integrate((d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="maxima")
[Out] 1/16*c^2*(4*(f*x + e)/(a^2*f) + (4*e^(-2*f*x - 2*e) - e^(-4*f*x - 4*e))/(a^2*f)) + 1/32*(8*f^2*x^2*e^(4*e) + 8*(2*f*x*e^(2*e) + e^(2*e))*e^(-2*f*x) - (4*f*x + 1)*e^(-4*f*x))*c*d*e^(-4*e)/(a^2*f^2) + 1/384*(32*f^3*x^3*e^(4*e) + 48*(2*f^2*x^2*e^(2*e) + 2*f*x*e^(2*e) + e^(2*e))*e^(-2*f*x) - 3*(8*f^2*x^2 + 4*f*x + 1)*e^(-4*f*x))*d^2*e^(-4*e)/(a^2*f^3)
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 217, normalized size of antiderivative = 1.28

$$\begin{aligned}
\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx \\
= \frac{(32d^2f^3x^3e^{(4fx+4e)} + 96cdf^3x^2e^{(4fx+4e)} + 96c^2f^3xe^{(4fx+4e)} + 96d^2f^2x^2e^{(2fx+2e)} - 24d^2f^2x^2 + 192cdf^3x^3e^{(4fx+4e)})}{(a + a \coth(e + fx))^2}
\end{aligned}$$

```
[In] integrate((d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="giac")
```

[Out] $\frac{1}{384} \left(32 d^2 f^3 x^3 e^{(4 f x + 4 e)} + 96 c d f^3 x^2 e^{(4 f x + 4 e)} + 96 c^2 f^3 x^3 e^{(4 f x + 4 e)} + 96 d^2 f^2 x^2 e^{(2 f x + 2 e)} - 24 d^2 f^2 x^2 e^{(2 f x + 2 e)} + 192 c d f^2 x^2 e^{(2 f x + 2 e)} - 48 c d f^2 x^2 e^{(2 f x + 2 e)} + 96 c^2 f^2 x^2 e^{(2 f x + 2 e)} + 96 d^2 f x^2 e^{(2 f x + 2 e)} - 24 c^2 f^2 x^2 - 12 d^2 f x^2 + 96 c d f e^{(2 f x + 2 e)} - 12 c d f + 48 d^2 f e^{(2 f x + 2 e)} - 3 d^2 e^{(-4 f x - 4 e)} \right) / (a^2 f^3)$

Mupad [B] (verification not implemented)

Time = 1.94 (sec), antiderivative size = 164, normalized size of antiderivative = 0.96

$$\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^2} dx = e^{-2e-2fx} \left(\frac{2 c^2 f^2 + 2 c d f + d^2}{8 a^2 f^3} + \frac{d^2 x^2}{4 a^2 f} + \frac{d x (d + 2 c f)}{4 a^2 f^2} \right) - e^{-4e-4fx} \left(\frac{8 c^2 f^2 + 4 c d f + d^2}{128 a^2 f^3} + \frac{d^2 x^2}{16 a^2 f} + \frac{d x (d + 4 c f)}{32 a^2 f^2} \right) + \frac{c^2 x}{4 a^2} + \frac{d^2 x^3}{12 a^2} + \frac{c d x^2}{4 a^2}$$

[In] int((c + d*x)^2/(a + a*coth(e + f*x))^2, x)

[Out] $\exp(-2e-2fx) \left((d^2 + 2c^2 f^2 + 2c d f) / (8a^2 f^3) + (d^2 x^2) / (4a^2 f) + (d x * (d + 2c f)) / (4a^2 f^2) \right) - \exp(-4e-4fx) \left((d^2 + 8c^2 f^2 + 4c d f) / (128 a^2 f^3) + (d^2 x^2) / (16 a^2 f) + (d x * (d + 4c f)) / (32 a^2 f^2) + (c^2 x) / (4 a^2) + (d^2 x^3) / (12 a^2) + (c d x^2) / (4 a^2) \right)$

3.24 $\int \frac{c+dx}{(a+a \coth(e+fx))^2} dx$

Optimal result	159
Rubi [A] (verified)	159
Mathematica [A] (verified)	161
Maple [A] (verified)	161
Fricas [A] (verification not implemented)	162
Sympy [B] (verification not implemented)	162
Maxima [A] (verification not implemented)	163
Giac [A] (verification not implemented)	163
Mupad [B] (verification not implemented)	164

Optimal result

Integrand size = 18, antiderivative size = 133

$$\begin{aligned} \int \frac{c+dx}{(a+a \coth(e+fx))^2} dx = & \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a \coth(e+fx))^2} \\ & - \frac{c+dx}{4f(a+a \coth(e+fx))^2} - \frac{3d}{16f^2(a^2+a^2 \coth(e+fx))} \\ & - \frac{c+dx}{4f(a^2+a^2 \coth(e+fx))} \end{aligned}$$

[Out] $3/16*d*x/a^2/f-1/8*d*x^2/a^2+1/4*x*(d*x+c)/a^2-1/16*d/f^2/(a+a*\coth(f*x+e))$
 $^2+1/4*(-d*x-c)/f/(a+a*\coth(f*x+e))^2-3/16*d/f^2/(a^2+a^2*\coth(f*x+e))+1/4*$
 $(-d*x-c)/f/(a^2+a^2*\coth(f*x+e))$

Rubi [A] (verified)

Time = 0.11 (sec), antiderivative size = 133, normalized size of antiderivative = 1.00,
 number of steps used = 7, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used
 $= \{3560, 8, 3811\}$

$$\begin{aligned} \int \frac{c+dx}{(a+a \coth(e+fx))^2} dx = & -\frac{c+dx}{4f(a^2 \coth(e+fx)+a^2)} + \frac{x(c+dx)}{4a^2} \\ & - \frac{3d}{16f^2(a^2 \coth(e+fx)+a^2)} + \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2} \\ & - \frac{c+dx}{4f(a \coth(e+fx)+a)^2} - \frac{d}{16f^2(a \coth(e+fx)+a)^2} \end{aligned}$$

[In] `Int[(c + d*x)/(a + a*Coth[e + f*x])^2, x]`

[Out] $(3*d*x)/(16*a^2*f) - (d*x^2)/(8*a^2) + (x*(c + d*x))/(4*a^2) - d/(16*f^2*(a + a*\text{COTH}[e + f*x])^2) - (c + d*x)/(4*f*(a + a*\text{COTH}[e + f*x])^2) - (3*d)/(16*f^2*(a^2 + a^2*\text{COTH}[e + f*x])) - (c + d*x)/(4*f*(a^2 + a^2*\text{COTH}[e + f*x]))$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3560

Int[((a_) + (b_)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + Dist[1/(2*a), Int[(a + b*Tan[c + d*x])^(n + 1), x], x] /; FreeQ[{a, b, c, d}, x] && EqQ[a^2 + b^2, 0] && LtQ[n, 0]

Rule 3811

Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> With[{u = IntHide[(a + b*Tan[e + f*x])^n, x]}, Dist[(c + d*x)^m, u, x] - Dist[d*m, Int[Dist[(c + d*x)^(m - 1), u, x], x], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, -1] && GtQ[m, 0]

Rubi steps

$$\begin{aligned}
 \text{integral} &= \frac{x(c+dx)}{4a^2} - \frac{c+dx}{4f(a+a\coth(e+fx))^2} - \frac{c+dx}{4f(a^2+a^2\coth(e+fx))} \\
 &\quad - d \int \left(\frac{x}{4a^2} - \frac{1}{4f(a+a\coth(e+fx))^2} - \frac{1}{4f(a^2+a^2\coth(e+fx))} \right) dx \\
 &= -\frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{c+dx}{4f(a+a\coth(e+fx))^2} - \frac{c+dx}{4f(a^2+a^2\coth(e+fx))} \\
 &\quad + \frac{d \int \frac{1}{(a+a\coth(e+fx))^2} dx}{4f} + \frac{d \int \frac{1}{a^2+a^2\coth(e+fx)} dx}{4f} \\
 &= -\frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a\coth(e+fx))^2} - \frac{c+dx}{4f(a+a\coth(e+fx))^2} \\
 &\quad - \frac{d}{8f^2(a^2+a^2\coth(e+fx))} - \frac{c+dx}{4f(a^2+a^2\coth(e+fx))} + \frac{d \int 1 dx}{8a^2f} \\
 &\quad + \frac{d \int \frac{1}{a+a\coth(e+fx)} dx}{8af} \\
 &= \frac{dx}{8a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a\coth(e+fx))^2} - \frac{c+dx}{4f(a+a\coth(e+fx))^2} \\
 &\quad - \frac{3d}{16f^2(a^2+a^2\coth(e+fx))} - \frac{c+dx}{4f(a^2+a^2\coth(e+fx))} + \frac{d \int 1 dx}{16a^2f}
 \end{aligned}$$

$$\begin{aligned}
&= \frac{3dx}{16a^2f} - \frac{dx^2}{8a^2} + \frac{x(c+dx)}{4a^2} - \frac{d}{16f^2(a+a\coth(e+fx))^2} - \frac{c+dx}{4f(a+a\coth(e+fx))^2} \\
&\quad - \frac{3d}{16f^2(a^2+a^2\coth(e+fx))} - \frac{c+dx}{4f(a^2+a^2\coth(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.46 (sec), antiderivative size = 114, normalized size of antiderivative = 0.86

$$\begin{aligned}
&\int \frac{c+dx}{(a+a\coth(e+fx))^2} dx \\
&= \frac{\text{csch}^2(e+fx)(8(d+2cf+2dfx)+(4cf(-1+4fx)+d(-1-4fx+8f^2x^2))\cosh(2(e+fx))+(4cf(1+4fx+8f^2x^2))\sinh(2(e+fx)))}{64a^2f^2(1+\coth(e+fx))^2}
\end{aligned}$$

[In] Integrate[(c + d*x)/(a + a*Coth[e + f*x])^2, x]

[Out] $\text{Csch}[e+f*x]^2*(8*(d+2*c*f+2*d*f*x)+(4*c*f*(-1+4*f*x)+d*(-1-4*f*x+8*f^2*x^2))*\text{Cosh}[2*(e+f*x)]+(4*c*f*(1+4*f*x)+d*(1+4*f*x+8*f^2*x^2))*\text{Sinh}[2*(e+f*x)])/(64*a^2*f^2*(1+\text{Coth}[e+f*x])^2)$

Maple [A] (verified)

Time = 0.38 (sec), antiderivative size = 74, normalized size of antiderivative = 0.56

method	result	size
risch	$\frac{dx^2}{8a^2} + \frac{cx}{4a^2} + \frac{(2df+2cf+d)e^{-2fx-2e}}{8a^2f^2} - \frac{(4df+4cf+d)e^{-4fx-4e}}{64a^2f^2}$	74
parallelrisch	$\frac{4\left(\left(\frac{dx}{2}+c\right)f-\frac{5d}{4}\right)xf\tanh(fx+e)^2+(4(dx^2+2cx)f^2+2(dx+6c)f+5d)\tanh(fx+e)+2(dx^2+2cx)f^2+(3dx+8c)f+4d}{16f^2a^2(1+\tanh(fx+e))^2}$	108

[In] int((d*x+c)/(a+a*coth(f*x+e))^2, x, method=_RETURNVERBOSE)

[Out] $\frac{1}{8}d^2x^2/a^2+1/4/a^2c^2x+1/8*(2d^2f*x+2c^2f+d)/a^2/f^2\exp(-2f*x-2e)-1/64*(4d^2f*x+4c^2f+d)/a^2/f^2\exp(-4f*x-4e)$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 190, normalized size of antiderivative = 1.43

$$\int \frac{c + dx}{(a + a \coth(e + fx))^2} dx$$

$$= \frac{16 dfx + (8 df^2 x^2 - 4 cf + 4 (4 cf^2 - df)x - d) \cosh(fx + e)^2 + 2 (8 df^2 x^2 + 4 cf + 4 (4 cf^2 + df)x + d) \cosh(fx + e)^2 + 16 cf^2 x \sinh(fx + e)^2 + 16 cf^2 \cosh(fx + e)^2 \sinh(fx + e)^2}{64 (a^2 f^2 \cosh(fx + e)^2 + 2 a^2 f^2 \cosh(fx + e)^2 + 16 a^2 f^2 \sinh(fx + e)^2 + 16 a^2 f^2 \cosh(fx + e)^2 \sinh(fx + e)^2)}$$

```
[In] integrate((d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="fricas")
[Out] 1/64*(16*d*f*x + (8*d*f^2*x^2 - 4*c*f + 4*(4*c*f^2 - d*f)*x - d)*cosh(f*x + e)^2 + 2*(8*d*f^2*x^2 + 4*c*f + 4*(4*c*f^2 + d*f)*x + d)*cosh(f*x + e)*sinh(f*x + e) + (8*d*f^2*x^2 - 4*c*f + 4*(4*c*f^2 - d*f)*x - d)*sinh(f*x + e)^2 + 16*c*f + 8*d)/(a^2*f^2*cosh(f*x + e)^2 + 2*a^2*f^2*cosh(f*x + e)*sinh(f*x + e) + a^2*f^2*sinh(f*x + e)^2)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 700 vs. $2(122) = 244$.

Time = 0.80 (sec) , antiderivative size = 700, normalized size of antiderivative = 5.26

$$\int \frac{c + dx}{(a + a \coth(e + fx))^2} dx$$

$$= \begin{cases} \frac{4cf^2 x \tanh^2(e + fx)}{16a^2 f^2 \tanh^2(e + fx) + 32a^2 f^2 \tanh(e + fx) + 16a^2 f^2} + \frac{8cf^2 x \tanh(e + fx)}{16a^2 f^2 \tanh^2(e + fx) + 32a^2 f^2 \tanh(e + fx) + 16a^2 f^2} + \frac{4cf^2 x}{16a^2 f^2 \tanh^2(e + fx) + 32a^2 f^2} \\ \frac{cx + \frac{dx^2}{2}}{(a \coth(e) + a)^2} \end{cases}$$

```
[In] integrate((d*x+c)/(a+a*coth(f*x+e))**2,x)
```

```
[Out] Piecewise((4*c*f**2*x*tanh(e + f*x)**2/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 8*c*f**2*x*tanh(e + f*x)/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 4*c*f**2*x/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 12*c*f*tanh(e + f*x)/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 8*c*f/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 2*d*f**2*x**2*tanh(e + f*x)**2/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 4*d*f**2*x**2*tanh(e + f*x)/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 2*d*f**2*x**2/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) - 5*d*f*x*tanh(e + f*x)**2/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2))
```

```
+ 2*d*f*x*tanh(e + f*x)/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 3*d*f*x/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 5*d*tanh(e + f*x)/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2) + 4*d/(16*a**2*f**2*tanh(e + f*x)**2 + 32*a**2*f**2*tanh(e + f*x) + 16*a**2*f**2), Ne(f, 0)), ((c*x + d*x**2/2)/(a*coth(e) + a)**2, True))
```

Maxima [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 107, normalized size of antiderivative = 0.80

$$\begin{aligned} & \int \frac{c + dx}{(a + a \coth(e + fx))^2} dx \\ &= \frac{1}{16} c \left(\frac{4(fx + e)}{a^2 f} + \frac{4e^{(-2fx - 2e)} - e^{(-4fx - 4e)}}{a^2 f} \right) \\ &+ \frac{(8f^2 x^2 e^{(4e)} + 8(2fx e^{(2e)} + e^{(2e)}) e^{(-2fx)} - (4fx + 1)e^{(-4fx)}) de^{(-4e)}}{64 a^2 f^2} \end{aligned}$$

[In] integrate((d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="maxima")

[Out] $\frac{1}{16} c (4(fx + e)/(a^2 f) + (4e^{-(-2fx - 2e)} - e^{-(-4fx - 4e)})/(a^2 f)) + \frac{1}{64} (8f^2 x^2 e^{(4e)} + 8(2fx e^{(2e)} + e^{(2e)}) * e^{-(-2fx)} - (4fx + 1)e^{-(-4fx)}) * d * e^{-(-4e)} / (a^2 f^2)$

Giac [A] (verification not implemented)

none

Time = 0.27 (sec) , antiderivative size = 103, normalized size of antiderivative = 0.77

$$\begin{aligned} & \int \frac{c + dx}{(a + a \coth(e + fx))^2} dx \\ &= \frac{(8df^2 x^2 e^{(4fx + 4e)} + 16cf^2 x e^{(4fx + 4e)} + 16dfxe^{(2fx + 2e)} - 4dfx + 16cfe^{(2fx + 2e)} - 4cf + 8de^{(2fx + 2e)} - d)e^{(-4fx - 4e)}}{64 a^2 f^2} \end{aligned}$$

[In] integrate((d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="giac")

[Out] $\frac{1}{64} (8d^2 f^2 x^2 e^{(4fx + 4e)} + 16c^2 f^2 x^2 e^{(4fx + 4e)} + 16d^2 f^2 x e^{(2fx + 2e)} - 4d^2 f x + 16c^2 f^2 e^{(2fx + 2e)} - 4c^2 f + 8d^2 e^{(2fx + 2e)} - d)e^{(-4fx - 4e)} / (a^2 f^2)$

Mupad [B] (verification not implemented)

Time = 1.91 (sec) , antiderivative size = 88, normalized size of antiderivative = 0.66

$$\int \frac{c + dx}{(a + a \coth(e + fx))^2} dx = e^{-2e-2fx} \left(\frac{d + 2cf}{8a^2f^2} + \frac{dx}{4a^2f} \right) - e^{-4e-4fx} \left(\frac{d + 4cf}{64a^2f^2} + \frac{dx}{16a^2f} \right) + \frac{dx^2}{8a^2} + \frac{cx}{4a^2}$$

[In] `int((c + d*x)/(a + a*coth(e + f*x))^2,x)`

[Out] `exp(- 2*e - 2*f*x)*((d + 2*c*f)/(8*a^2*f^2) + (d*x)/(4*a^2*f)) - exp(- 4*e - 4*f*x)*((d + 4*c*f)/(64*a^2*f^2) + (d*x)/(16*a^2*f)) + (d*x^2)/(8*a^2) + (c*x)/(4*a^2)`

3.25 $\int \frac{1}{(c+dx)(a+a \coth(e+fx))^2} dx$

Optimal result	165
Rubi [A] (verified)	166
Mathematica [A] (verified)	168
Maple [A] (verified)	169
Fricas [A] (verification not implemented)	169
Sympy [F]	169
Maxima [A] (verification not implemented)	170
Giac [A] (verification not implemented)	170
Mupad [F(-1)]	170

Optimal result

Integrand size = 20, antiderivative size = 297

$$\begin{aligned} \int \frac{1}{(c+dx)(a+a \coth(e+fx))^2} dx = & -\frac{\cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & + \frac{\cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{4a^2d} \\ & + \frac{\log(c+dx)}{4a^2d} - \frac{\text{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^2d} \\ & + \frac{\text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2a^2d} \\ & + \frac{\cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & - \frac{\sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\ & - \frac{\cosh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{4a^2d} \\ & + \frac{\sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{4a^2d} \end{aligned}$$

```
[Out] 1/4*Chi(4*c*f/d+4*f*x)*cosh(-4*e+4*c*f/d)/a^2/d-1/2*Chi(2*c*f/d+2*f*x)*cosh
(-2*e+2*c*f/d)/a^2/d+1/4*ln(d*x+c)/a^2/d+1/2*cosh(-2*e+2*c*f/d)*Shi(2*c*f/d
+2*f*x)/a^2/d-1/4*cosh(-4*e+4*c*f/d)*Shi(4*c*f/d+4*f*x)/a^2/d+1/4*Chi(4*c*f
/d+4*f*x)*sinh(-4*e+4*c*f/d)/a^2/d-1/4*Shi(4*c*f/d+4*f*x)*sinh(-4*e+4*c*f/d
)/a^2/d-1/2*Chi(2*c*f/d+2*f*x)*sinh(-2*e+2*c*f/d)/a^2/d+1/2*Shi(2*c*f/d+2*f
*x)*sinh(-2*e+2*c*f/d)/a^2/d
```

Rubi [A] (verified)

Time = 0.56 (sec) , antiderivative size = 297, normalized size of antiderivative = 1.00, number of steps used = 21, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3809, 3384, 3379, 3382, 3393}

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx = \frac{\operatorname{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{2a^2d} - \frac{\operatorname{Chi}(4xf + \frac{4cf}{d}) \sinh(4e - \frac{4cf}{d})}{4a^2d} - \frac{\operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{2a^2d} + \frac{\operatorname{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{4a^2d} - \frac{\sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{2a^2d} + \frac{\sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{4a^2d} + \frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{2a^2d} - \frac{\cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{4a^2d} + \frac{\log(c + dx)}{4a^2d}$$

[In] $\operatorname{Int}[1/((c + d*x)*(a + a*\operatorname{Coth}[e + f*x])^2), x]$

[Out] $-1/2*(\operatorname{Cosh}[2*e - (2*c*f)/d]*\operatorname{CoshIntegral}[(2*c*f)/d + 2*f*x])/(a^{2*d}) + (\operatorname{Cosh}[4*e - (4*c*f)/d]*\operatorname{CoshIntegral}[(4*c*f)/d + 4*f*x])/(4*a^{2*d}) + \operatorname{Log}[c + d*x]/(4*a^{2*d}) - (\operatorname{CoshIntegral}[(4*c*f)/d + 4*f*x]*\operatorname{Sinh}[4*e - (4*c*f)/d])/(4*a^{2*d}) + (\operatorname{CoshIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sinh}[2*e - (2*c*f)/d])/(2*a^{2*d}) + (\operatorname{Cosh}[2*e - (2*c*f)/d]*\operatorname{SinhIntegral}[(2*c*f)/d + 2*f*x])/(2*a^{2*d}) - (\operatorname{Sinh}[2*e - (2*c*f)/d]*\operatorname{SinhIntegral}[(2*c*f)/d + 2*f*x])/(2*a^{2*d}) - (\operatorname{Cosh}[4*e - (4*c*f)/d]*\operatorname{SinhIntegral}[(4*c*f)/d + 4*f*x])/(4*a^{2*d}) + (\operatorname{Sinh}[4*e - (4*c*f)/d]*\operatorname{SinhIntegral}[(4*c*f)/d + 4*f*x])/(4*a^{2*d})$

Rule 3379

$\operatorname{Int}[\sin[(e_{_}) + (\operatorname{Complex}[0, fz_{_}])*(f_{_})*(x_{_})]/((c_{_}) + (d_{_})*(x_{_})), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[I*(\operatorname{SinhIntegral}[c*f*(fz/d) + f*fz*x]/d), x] /; \operatorname{FreeQ}[\{c, d, e, f, fz\}, x] \&& \operatorname{EqQ}[d*e - c*f*fz*I, 0]$

Rule 3382

$\operatorname{Int}[\sin[(e_{_}) + (\operatorname{Complex}[0, fz_{_}])*(f_{_})*(x_{_})]/((c_{_}) + (d_{_})*(x_{_})), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{CoshIntegral}[c*f*(fz/d) + f*fz*x]/d, x] /; \operatorname{FreeQ}[\{c, d, e, f, fz\}, x] \&& \operatorname{EqQ}[d*(e - \pi/2) - c*f*fz*I, 0]$

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_)]/((c_.) + (d_.)*(x_)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> In[t[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)} - \frac{\cosh(2e+2fx)}{2a^2(c+dx)} + \frac{\cosh^2(2e+2fx)}{4a^2(c+dx)} + \frac{\sinh(2e+2fx)}{2a^2(c+dx)} \right. \\
&\quad \left. + \frac{\sinh^2(2e+2fx)}{4a^2(c+dx)} - \frac{\sinh(4e+4fx)}{4a^2(c+dx)} \right) dx \\
&= \frac{\log(c+dx)}{4a^2d} + \frac{\int \frac{\cosh^2(2e+2fx)}{c+dx} dx}{4a^2} + \frac{\int \frac{\sinh^2(2e+2fx)}{c+dx} dx}{4a^2} \\
&\quad - \frac{\int \frac{\sinh(4e+4fx)}{c+dx} dx}{4a^2} - \frac{\int \frac{\cosh(2e+2fx)}{c+dx} dx}{2a^2} + \frac{\int \frac{\sinh(2e+2fx)}{c+dx} dx}{2a^2} \\
&= \frac{\log(c+dx)}{4a^2d} - \frac{\int \left(\frac{1}{2(c+dx)} - \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} + \frac{\int \left(\frac{1}{2(c+dx)} + \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{4a^2} \\
&\quad - \frac{\cosh(4e - \frac{4cf}{d}) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{4a^2} - \frac{\cosh(2e - \frac{2cf}{d}) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} \\
&\quad + \frac{\cosh(2e - \frac{2cf}{d}) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} - \frac{\sinh(4e - \frac{4cf}{d}) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{4a^2} \\
&\quad + \frac{\sinh(2e - \frac{2cf}{d}) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2} - \frac{\sinh(2e - \frac{2cf}{d}) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{2a^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} \\
&\quad - \frac{\operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^2d} + \frac{\operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2a^2d} \\
&\quad + \frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{\sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{4a^2d} + 2 \int \frac{\cosh(4e + 4fx)}{c+dx} dx \\
&= -\frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} - \frac{\operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^2d} \\
&\quad + \frac{\operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2a^2d} + \frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{\cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{4a^2d} \\
&\quad + 2 \left(\frac{\cosh(4e - \frac{4cf}{d}) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^2} + \frac{\sinh(4e - \frac{4cf}{d}) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^2} \right) \\
&= -\frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{2a^2d} + \frac{\log(c + dx)}{4a^2d} - \frac{\operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{4a^2d} \\
&\quad + \frac{\operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{2a^2d} + \frac{\cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} \\
&\quad - \frac{\sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{2a^2d} - \frac{\cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{4a^2d} \\
&\quad + 2 \left(\frac{\cosh(4e - \frac{4cf}{d}) \operatorname{Chi}(\frac{4cf}{d} + 4fx)}{8a^2d} + \frac{\sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{8a^2d} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.13 (sec), antiderivative size = 199, normalized size of antiderivative = 0.67

$$\begin{aligned}
&\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx \\
&= \frac{(\cosh(2e - \frac{2cf}{d}) - \sinh(2e - \frac{2cf}{d})) \left(-2 \operatorname{Chi}\left(\frac{2f(c+dx)}{d}\right) + \cosh(2e - \frac{2cf}{d}) \log(f(c + dx)) + \operatorname{Chi}\left(\frac{4f(c+dx)}{d}\right) \right)}{(c + dx)(a + a \coth(e + fx))^2}
\end{aligned}$$

[In] `Integrate[1/((c + d*x)*(a + a*Coth[e + f*x])^2), x]`

[Out] `((Cosh[2*e - (2*c*f)/d] - Sinh[2*e - (2*c*f)/d])*(-2*CoshIntegral[(2*f*(c + d*x))/d] + Cosh[2*e - (2*c*f)/d]*Log[f*(c + d*x)] + CoshIntegral[(4*f*(c + d*x))/d]*(Cosh[2*e - (2*c*f)/d] - Sinh[2*e - (2*c*f)/d]) + Log[f*(c + d*x)]*Sinh[2*e - (2*c*f)/d] + 2*SinhIntegral[(2*f*(c + d*x))/d] - Cosh[2*e - (2*c*f)/d]*SinhIntegral[(4*f*(c + d*x))/d] + Sinh[2*e - (2*c*f)/d]*SinhIntegral[(4*f*(c + d*x))/d]))/(4*a^2*d)`

Maple [A] (verified)

Time = 0.34 (sec) , antiderivative size = 106, normalized size of antiderivative = 0.36

method	result	size
risch	$\frac{\ln(dx+c)}{4a^2d} - \frac{e^{\frac{4cf-4de}{d}} \text{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{4a^2d} + \frac{e^{\frac{2cf-2de}{d}} \text{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{2a^2d}$	106

[In] `int(1/(d*x+c)/(a+a*coth(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4} \ln(d*x+c)/a^2/d - \frac{1}{4} a^2/d * \exp(4*(c*f-d*e)/d) * \text{Ei}(1, 4*f*x+4*e+4*(c*f-d*e)/d) + \frac{1}{2} a^2/d * \exp(2*(c*f-d*e)/d) * \text{Ei}(1, 2*f*x+2*e+2*(c*f-d*e)/d)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 137, normalized size of antiderivative = 0.46

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx = \\ - \frac{2 \text{Ei}\left(-\frac{2(df+cf)}{d}\right) \cosh\left(-\frac{2(de-cf)}{d}\right) - \text{Ei}\left(-\frac{4(df+cf)}{d}\right) \cosh\left(-\frac{4(de-cf)}{d}\right) + 2 \text{Ei}\left(-\frac{2(df+cf)}{d}\right) \sinh\left(-\frac{2(de-cf)}{d}\right)}{4a^2d}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="fricas")`

[Out] $- \frac{1}{4} (2 \text{Ei}(-2*(d*f*x + c*f)/d) * \cosh(-2*(d*e - c*f)/d) - \text{Ei}(-4*(d*f*x + c*f)/d) * \cosh(-4*(d*e - c*f)/d) + 2 \text{Ei}(-2*(d*f*x + c*f)/d) * \sinh(-2*(d*e - c*f)/d) - \text{Ei}(-4*(d*f*x + c*f)/d) * \sinh(-4*(d*e - c*f)/d) - \log(d*x + c)) / (a^2*d)$

Sympy [F]

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx \\ = \frac{\int \frac{1}{c \coth^2(e+fx)+2c \coth(e+fx)+c+dx \coth^2(e+fx)+2dx \coth(e+fx)+dx} dx}{a^2}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))**2,x)`

[Out] `Integral(1/(c*coth(e + f*x)**2 + 2*c*coth(e + f*x) + c + d*x*coth(e + f*x)*2 + 2*d*x*coth(e + f*x) + d*x), x)/a**2`

Maxima [A] (verification not implemented)

none

Time = 0.70 (sec) , antiderivative size = 81, normalized size of antiderivative = 0.27

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx = -\frac{e^{\left(-4e + \frac{4cf}{d}\right)} E_1\left(\frac{4(dx+c)f}{d}\right)}{4a^2d} + \frac{e^{\left(-2e + \frac{2cf}{d}\right)} E_1\left(\frac{2(dx+c)f}{d}\right)}{2a^2d} + \frac{\log(dx + c)}{4a^2d}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="maxima")`

[Out]
$$-\frac{1}{4} \cdot \frac{4 \cdot e^{-(-4e + 4c*f/d)} \cdot \text{exp_integral_e}(1, 4*(d*x + c)*f/d)/(a^2*d) + 1/2 \cdot e^{(-2e + 2c*f/d)} \cdot \text{exp_integral_e}(1, 2*(d*x + c)*f/d)/(a^2*d) + 1/4 \cdot \log(d*x + c)/(a^2*d)}{a^2}$$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec) , antiderivative size = 77, normalized size of antiderivative = 0.26

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx = -\frac{\left(2 \cdot \text{Ei}\left(-\frac{2(df+cf)}{d}\right) \cdot e^{\left(2e + \frac{2cf}{d}\right)} - \text{Ei}\left(-\frac{4(df+cf)}{d}\right) \cdot e^{\left(\frac{4cf}{d}\right)} - e^{(4e)} \cdot \log(dx + c)\right) e^{(-4e)}}{4a^2d}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))^2,x, algorithm="giac")`

[Out]
$$-\frac{1}{4} \cdot \frac{4 \cdot (\text{Ei}(-2*(d*f*x + c*f)/d) \cdot e^{(2e + 2c*f/d)} - \text{Ei}(-4*(d*f*x + c*f)/d) \cdot e^{(4c*f/d)} - e^{(4e)} \cdot \log(d*x + c)) \cdot e^{(-4e)}}{a^2}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^2} dx = \int \frac{1}{(a + a \coth(e + f x))^2 (c + d x)} dx$$

[In] `int(1/((a + a*coth(e + f*x))^2*(c + d*x)),x)`

[Out] `int(1/((a + a*coth(e + f*x))^2*(c + d*x)), x)`

3.26 $\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^2} dx$

Optimal result	171
Rubi [A] (verified)	172
Mathematica [A] (verified)	175
Maple [A] (verified)	176
Fricas [A] (verification not implemented)	176
Sympy [F]	177
Maxima [A] (verification not implemented)	177
Giac [A] (verification not implemented)	178
Mupad [F(-1)]	178

Optimal result

Integrand size = 20, antiderivative size = 420

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^2} dx = & -\frac{1}{4a^2d(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\ & + \frac{f \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & - \frac{f \cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\ & + \frac{f \text{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{a^2d^2} \\ & - \frac{f \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{a^2d^2} \\ & - \frac{\sinh(2e + 2fx)}{2a^2d(c+dx)} \\ & - \frac{\sinh^2(2e + 2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e + 4fx)}{4a^2d(c+dx)} \\ & - \frac{f \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & + \frac{f \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\ & + \frac{f \cosh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\ & - \frac{f \sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \end{aligned}$$

[Out] $-1/4/a^2/d/(d*x+c)-f*\text{Chi}(4*c*f/d+4*f*x)*\cosh(-4*e+4*c*f/d)/a^2/d^2+f*\text{Chi}(2*c*f/d+2*f*x)*\cosh(-2*e+2*c*f/d)/a^2/d^2+1/2*\cosh(2*f*x+2*e)/a^2/d/(d*x+c)-1/4*\cosh(2*f*x+2*e)^2/a^2/d/(d*x+c)-f*\cosh(-2*e+2*c*f/d)*\text{Shi}(2*c*f/d+2*f*x)/$

$$\begin{aligned}
& a^2/d^2 + f \cosh(-4e + 4cf/d) \operatorname{Shi}(4cf/d + 4fx)/a^2/d^2 - f \operatorname{Chi}(4cf/d + 4fx) \\
& * \sinh(-4e + 4cf/d)/a^2/d^2 + f \operatorname{Shi}(4cf/d + 4fx) * \sinh(-4e + 4cf/d)/a^2/d^2 \\
& + 2f \operatorname{Chi}(2cf/d + 2fx) * \sinh(-2e + 2cf/d)/a^2/d^2 - f \operatorname{Shi}(2cf/d + 2fx) * \sinh \\
& (-2e + 2cf/d)/a^2/d^2 - 1/2 \sinh(2fx + 2e)/a^2/d/(dx + c) - 1/4 \sinh(2fx + 2e) \\
&)^2/a^2/d/(dx + c) + 1/4 \sinh(4fx + 4e)/a^2/d/(dx + c)
\end{aligned}$$

Rubi [A] (verified)

Time = 0.61 (sec), antiderivative size = 420, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.350, Rules used = {3809, 3378, 3384, 3379, 3382, 3394, 12}

$$\begin{aligned}
\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx = & \frac{f \operatorname{Chi}(4xf + \frac{4cf}{d}) \sinh(4e - \frac{4cf}{d})}{a^2 d^2} \\
& - \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{a^2 d^2} \\
& + \frac{f \operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{a^2 d^2} \\
& - \frac{f \operatorname{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{a^2 d^2} \\
& + \frac{f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{a^2 d^2} \\
& - \frac{f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{a^2 d^2} \\
& - \frac{f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{a^2 d^2} \\
& + \frac{f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{a^2 d^2} \\
& - \frac{\sinh^2(2e + 2fx)}{4a^2 d(c + dx)} - \frac{\sinh(2e + 2fx)}{2a^2 d(c + dx)} \\
& + \frac{\sinh(4e + 4fx)}{4a^2 d(c + dx)} - \frac{\cosh^2(2e + 2fx)}{4a^2 d(c + dx)} \\
& + \frac{\cosh(2e + 2fx)}{2a^2 d(c + dx)} - \frac{1}{4a^2 d(c + dx)}
\end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + a*Coth[e + f*x])^2), x]

[Out] $-1/4*1/(a^2*d*(c + d*x)) + \operatorname{Cosh}[2*e + 2*f*x]/(2*a^2*d*(c + d*x)) - \operatorname{Cosh}[2*e + 2*f*x]^2/(4*a^2*d*(c + d*x)) + (f*\operatorname{Cosh}[2*e - (2*c*f)/d]*\operatorname{CoshIntegral}[(2*c*f)/d + 2*f*x])/(a^2*d^2) - (f*\operatorname{Cosh}[4*e - (4*c*f)/d]*\operatorname{CoshIntegral}[(4*c*f)/d + 4*f*x])/(a^2*d^2) + (f*\operatorname{CoshIntegral}[(4*c*f)/d + 4*f*x]*\operatorname{Sinh}[4*e - (4*c*f)/d])/(a^2*d^2) - (f*\operatorname{CoshIntegral}[(2*c*f)/d + 2*f*x]*\operatorname{Sinh}[2*e - (2*c*f)/d])/(a^2*d^2) - \operatorname{Sinh}[2*e + 2*f*x]/(2*a^2*d*(c + d*x)) - \operatorname{Sinh}[2*e + 2*f*x]^2/(4*a^2*d*(c + d*x))$

```
4*a^2*d*(c + d*x)) + Sinh[4*e + 4*f*x]/(4*a^2*d*(c + d*x)) - (f*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(a^2*d^2) + (f*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(a^2*d^2) + (f*Cosh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(a^2*d^2) - (f*Sinh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(a^2*d^2)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 3378

```
Int[((c_.) + (d_.)*(x_.))^m_*sin[(e_.) + (f_.)*(x_.)], x_Symbol] :> Simp[(c + d*x)^m*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^m + 1]*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*f(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*f(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3394

```
Int[((c_.) + (d_.)*(x_.))^m_*sin[(e_.) + (f_.)*(x_.)]^n, x_Symbol] :> Simp[(c + d*x)^m*(Sin[e + f*x]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1))), Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]^(n - 1), x], x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] && LtQ[m, -1]
```

Rule 3809

```

Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(
2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{4a^2(c+dx)^2} - \frac{\cosh(2e+2fx)}{2a^2(c+dx)^2} + \frac{\cosh^2(2e+2fx)}{4a^2(c+dx)^2} + \frac{\sinh(2e+2fx)}{2a^2(c+dx)^2} \right. \\
&\quad \left. + \frac{\sinh^2(2e+2fx)}{4a^2(c+dx)^2} - \frac{\sinh(4e+4fx)}{4a^2(c+dx)^2} \right) dx \\
&= -\frac{1}{4a^2d(c+dx)} + \frac{\int \frac{\cosh^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} + \frac{\int \frac{\sinh^2(2e+2fx)}{(c+dx)^2} dx}{4a^2} \\
&\quad - \frac{\int \frac{\sinh(4e+4fx)}{(c+dx)^2} dx}{4a^2} - \frac{\int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{2a^2} + \frac{\int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{2a^2} \\
&= -\frac{1}{4a^2d(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad - \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} \\
&\quad + \frac{(if) \int -\frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{a^2d} - \frac{(if) \int \frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{a^2d} \\
&\quad + \frac{f \int \frac{\cosh(2e+2fx)}{c+dx} dx}{a^2d} - \frac{f \int \frac{\cosh(4e+4fx)}{c+dx} dx}{a^2d} - \frac{f \int \frac{\sinh(2e+2fx)}{c+dx} dx}{a^2d} \\
&= -\frac{1}{4a^2d(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} - \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} \\
&\quad - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} + 2 \frac{f \int \frac{\sinh(4e+4fx)}{c+dx} dx}{2a^2d} \\
&\quad - \frac{(f \cosh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} + \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} \\
&\quad - \frac{(f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} - \frac{(f \sinh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{a^2d} \\
&\quad - \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d} + \frac{(f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{a^2d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{4a^2d(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{f \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{a^2d^2} - \frac{f \cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - \frac{f \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{a^2d^2} - \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} - \frac{f \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad + \frac{f \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} - \frac{f \sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad + 2 \left(\frac{(f \cosh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^2d} \right. \\
&\quad \quad \left. + \frac{(f \sinh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^2d} \right) \\
&= -\frac{1}{4a^2d(c+dx)} + \frac{\cosh(2e+2fx)}{2a^2d(c+dx)} - \frac{\cosh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{f \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{a^2d^2} - \frac{f \cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad - \frac{f \text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{a^2d^2} - \frac{\sinh(2e+2fx)}{2a^2d(c+dx)} - \frac{\sinh^2(2e+2fx)}{4a^2d(c+dx)} \\
&\quad + \frac{\sinh(4e+4fx)}{4a^2d(c+dx)} - \frac{f \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} \\
&\quad + \frac{f \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{a^2d^2} - \frac{f \sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{a^2d^2} \\
&\quad + 2 \left(\frac{f \text{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{2a^2d^2} + \frac{f \cosh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{2a^2d^2} \right)
\end{aligned}$$

Mathematica [A] (verified)

Time = 2.39 (sec), antiderivative size = 442, normalized size of antiderivative = 1.05

$$\begin{aligned}
&\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^2} dx \\
&= \frac{(-\cosh(2(e+f(-\frac{c}{d}+x))) + \sinh(2(e+f(-\frac{c}{d}+x)))) \left(-2d \cosh(\frac{2cf}{d}) + d \cosh(2(e+f(-\frac{c}{d}+x)))\right)}{a^2d^2}
\end{aligned}$$

[In] Integrate[1/((c + d*x)^2*(a + a*Coth[e + f*x])^2), x]

```
[Out] ((-Cosh[2*(e + f*(-(c/d) + x))] + Sinh[2*(e + f*(-(c/d) + x))])*(-2*d*Cosh[(2*c*f)/d] + d*Cosh[2*(e + f*(-(c/d) + x))] + d*Cosh[2*(e + f*(c/d + x))]) + 2*d*Sinh[(2*c*f)/d] - 4*f*(c + d*x)*CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[2*f*x] + Sinh[2*f*x]) + d*Sinh[2*(e + f*(-(c/d) + x))] - d*Sinh[2*(e + f*(c/d + x))] + 4*f*(c + d*x)*CoshIntegral[(4*f*(c + d*x))/d]*(Cosh[2*e - (2*f*(c + d*x))/d] - Sinh[2*e - (2*f*(c + d*x))/d]) + 4*c*f*Cosh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] + 4*d*f*x*Cosh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] + 4*c*f*Sinh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] + 4*d*f*x*Sinh[2*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 4*c*f*Cosh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] - 4*d*f*x*Cosh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] + 4*c*f*Sinh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] + 4*d*f*x*Sinh[2*e - (2*f*(c + d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d]))/(4*a^2*d^2*(c + d*x))
```

Maple [A] (verified)

Time = 0.39 (sec) , antiderivative size = 164, normalized size of antiderivative = 0.39

method	result
risch	$-\frac{1}{4a^2d(dx+c)} - \frac{fe^{-4fx-4e}}{4a^2d(df+cf)} + \frac{fe^{\frac{4cf-4de}{d}}\text{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{a^2d^2} + \frac{fe^{-2fx-2e}}{2a^2d(df+cf)} - \frac{fe^{\frac{2cf-2de}{d}}\text{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{a^2d^2}$

```
[In] int(1/(d*x+c)^2/(a+a*coth(f*x+e))^2,x,method=_RETURNVERBOSE)
```

```
[Out] -1/4/a^2/d/(d*x+c)-1/4*f/a^2*exp(-4*f*x-4*e)/d/(d*f*x+c*f)+f/a^2/d^2*exp(4*(c*f-d*e)/d)*Ei(1,4*f*x+4*e+4*(c*f-d*e)/d)+1/2*f/a^2*exp(-2*f*x-2*e)/d/(d*f*x+c*f)-f/a^2/d^2*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)
```

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 615, normalized size of antiderivative = 1.46

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx \\ = \frac{2(df + cf)\text{Ei}\left(-\frac{2(df + cf)}{d}\right) \cosh(fx + e)^2 \sinh\left(-\frac{2(de - cf)}{d}\right) - 2(df + cf)\text{Ei}\left(-\frac{4(df + cf)}{d}\right) \cosh(fx + e)^2}{(df + cf)^2}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] 1/2*(2*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-4*(d*e - c*f)/d) + (2*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) - d)*cos
```

$$\begin{aligned}
& h(f*x + e)^2 + (2*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) \\
& - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + 2*(d*f \\
& *x + c*f)*Ei(-2*(d*f*x + c*f)/d)*sinh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*E \\
& i(-4*(d*f*x + c*f)/d)*sinh(-4*(d*e - c*f)/d) - d)*sinh(f*x + e)^2 + 4*((d*f \\
& *x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)*sinh(-2*(d*e - c*f)/d) - (d*f \\
& *x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(f*x + e)*sinh(-4*(d*e - c*f)/d) + ((d*f*x \\
& + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) - (d*f*x + c*f)*E \\
& i(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d))*cosh(f*x + e))*sinh(f*x + e) \\
& + d)/((a^2*d^3*x + a^2*c*d^2)*cosh(f*x + e)^2 + 2*(a^2*d^3*x + a^2*c*d^2)* \\
& cosh(f*x + e)*sinh(f*x + e) + (a^2*d^3*x + a^2*c*d^2)*sinh(f*x + e)^2)
\end{aligned}$$

Sympy [F]

$$\begin{aligned}
& \int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx \\
& = \frac{\int \frac{1}{c^2 \coth^2(e + fx) + 2c^2 \coth(e + fx) + c^2 + 2cdx \coth^2(e + fx) + 4cdx \coth(e + fx) + 2cdx + d^2 x^2 \coth^2(e + fx) + 2d^2 x^2 \coth(e + fx) + d^2 x^2} dx}{a^2}
\end{aligned}$$

[In] integrate(1/(d*x+c)**2/(a+a*coth(f*x+e))**2,x)

[Out] Integral(1/(c**2*coth(e + f*x)**2 + 2*c**2*coth(e + f*x) + c**2 + 2*c*d*x*c
oth(e + f*x)**2 + 4*c*d*x*coth(e + f*x) + 2*c*d*x + d**2*x**2*coth(e + f*x)
2 + 2*d2*x**2*coth(e + f*x) + d**2*x**2), x)/a**2

Maxima [A] (verification not implemented)

none

Time = 1.32 (sec) , antiderivative size = 100, normalized size of antiderivative = 0.24

$$\begin{aligned}
\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx &= -\frac{1}{4(a^2 d^2 x + a^2 c d)} - \frac{e^{(-4e + \frac{4cf}{d})} E_2\left(\frac{4(dx+c)f}{d}\right)}{4(dx+c)a^2 d} \\
&+ \frac{e^{(-2e + \frac{2cf}{d})} E_2\left(\frac{2(dx+c)f}{d}\right)}{2(dx+c)a^2 d}
\end{aligned}$$

[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="maxima")

[Out] -1/4/(a^2*d^2*x + a^2*c*d) - 1/4*e^(-4*e + 4*c*f/d)*exp_integral_e(2, 4*(d*x + c)*f/d)/((d*x + c)*a^2*d) + 1/2*e^(-2*e + 2*c*f/d)*exp_integral_e(2, 2*(d*x + c)*f/d)/((d*x + c)*a^2*d)

Giac [A] (verification not implemented)

none

Time = 0.33 (sec) , antiderivative size = 585, normalized size of antiderivative = 1.39

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx \\ = \frac{\left(4(dx + c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right)f^2 \text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right) e^{-\frac{2(de-cf)}{d}} - 4def^2 \text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right) e^{-\frac{2(de-cf)}{d}}\right)}{(dx + c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right)f^2}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^2,x, algorithm="giac")
[Out] 1/4*(4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) - 4*d*e*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) + 4*c*f^3*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*(d*e - c*f)/d) - 4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 4*d*e*f^2*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) - 4*c*f^3*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d) + 2*d*f^2*e^(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*f^2)/d) - d*f^2*2*e^(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d) - d*f^2*2*d^2/(((d*x + c)*a^2*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a^2*d^5*e + a^2*c*d^4*f)*f)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^2} dx = \int \frac{1}{(a + a \coth(e + fx))^2 (c + dx)^2} dx$$

```
[In] int(1/((a + a*coth(e + f*x))^2*(c + d*x)^2),x)
[Out] int(1/((a + a*coth(e + f*x))^2*(c + d*x)^2), x)
```

3.27 $\int \frac{(c+dx)^3}{(a+a \coth(e+fx))^3} dx$

Optimal result	179
Rubi [A] (verified)	180
Mathematica [A] (verified)	182
Maple [A] (verified)	182
Fricas [B] (verification not implemented)	183
Sympy [B] (verification not implemented)	184
Maxima [A] (verification not implemented)	186
Giac [A] (verification not implemented)	187
Mupad [B] (verification not implemented)	188

Optimal result

Integrand size = 20, antiderivative size = 336

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+a \coth(e+fx))^3} dx = & \frac{d^3 e^{-6e-6fx}}{1728a^3 f^4} - \frac{9d^3 e^{-4e-4fx}}{1024a^3 f^4} + \frac{9d^3 e^{-2e-2fx}}{64a^3 f^4} + \frac{d^2 e^{-6e-6fx}(c+dx)}{288a^3 f^3} \\ & - \frac{9d^2 e^{-4e-4fx}(c+dx)}{256a^3 f^3} + \frac{9d^2 e^{-2e-2fx}(c+dx)}{32a^3 f^3} \\ & + \frac{de^{-6e-6fx}(c+dx)^2}{96a^3 f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3 f^2} \\ & + \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3 f^2} + \frac{e^{-6e-6fx}(c+dx)^3}{48a^3 f} \\ & - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3 f} + \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3 f} + \frac{(c+dx)^4}{32a^3 d} \end{aligned}$$

```
[Out] 1/1728*d^3*exp(-6*f*x-6*e)/a^3/f^4-9/1024*d^3*exp(-4*f*x-4*e)/a^3/f^4+9/64*d^3*exp(-2*f*x-2*e)/a^3/f^4+1/288*d^2*exp(-6*f*x-6*e)*(d*x+c)/a^3/f^3-9/256*d^2*exp(-4*f*x-4*e)*(d*x+c)/a^3/f^3+9/32*d^2*exp(-2*f*x-2*e)*(d*x+c)/a^3/f^3+1/96*d*exp(-6*f*x-6*e)*(d*x+c)^2/a^3/f^2-9/128*d*exp(-4*f*x-4*e)*(d*x+c)^2/a^3/f^2+1/48*exp(-6*f*x-6*e)*(d*x+c)^3/a^3/f^3-3/32*exp(-4*f*x-4*e)*(d*x+c)^3/a^3/f^3+1/16*exp(-2*f*x-2*e)*(d*x+c)^3/a^3/f^3+32*(d*x+c)^4/a^3/d
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 336, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.150$, Rules used = {3810, 2207, 2225}

$$\begin{aligned} \int \frac{(c+dx)^3}{(a+a\coth(e+fx))^3} dx &= \frac{d^2(c+dx)e^{-6e-6fx}}{288a^3f^3} - \frac{9d^2(c+dx)e^{-4e-4fx}}{256a^3f^3} \\ &\quad + \frac{9d^2(c+dx)e^{-2e-2fx}}{32a^3f^3} + \frac{d(c+dx)^2e^{-6e-6fx}}{96a^3f^2} \\ &\quad - \frac{9d(c+dx)^2e^{-4e-4fx}}{128a^3f^2} + \frac{9d(c+dx)^2e^{-2e-2fx}}{32a^3f^2} \\ &\quad + \frac{(c+dx)^3e^{-6e-6fx}}{48a^3f} - \frac{3(c+dx)^3e^{-4e-4fx}}{32a^3f} \\ &\quad + \frac{3(c+dx)^3e^{-2e-2fx}}{16a^3f} + \frac{(c+dx)^4}{32a^3d} \\ &\quad + \frac{d^3e^{-6e-6fx}}{1728a^3f^4} - \frac{9d^3e^{-4e-4fx}}{1024a^3f^4} + \frac{9d^3e^{-2e-2fx}}{64a^3f^4} \end{aligned}$$

[In] `Int[(c + d*x)^3/(a + a*Coth[e + f*x])^3, x]`

[Out]
$$\begin{aligned} & (d^3 E^{-6 e - 6 f x}) / (1728 a^3 f^4) - (9 d^3 E^{-4 e - 4 f x}) / (1024 a^3 f^4) \\ & + (9 d^3 E^{-2 e - 2 f x}) / (64 a^3 f^4) + (d^2 E^{-6 e - 6 f x}) * (c + d * x) / (288 a^3 f^3) \\ & - (9 d^2 E^{-4 e - 4 f x}) * (c + d * x) / (256 a^3 f^3) + (9 d^2 E^{-2 e - 2 f x}) * (c + d * x) / (32 a^3 f^3) \\ & + (d E^{-6 e - 6 f x}) * (c + d * x) / (96 a^3 f^2) - (9 d E^{-4 e - 4 f x}) * (c + d * x)^2 / (128 a^3 f^2) \\ & + (9 d E^{-2 e - 2 f x}) * (c + d * x)^2 / (32 a^3 f^2) + (E^{-6 e - 6 f x}) * (c + d * x)^3 / (48 a^3 f) \\ & - (3 E^{-4 e - 4 f x}) * (c + d * x)^3 / (32 a^3 f) + (3 E^{-2 e - 2 f x}) * (c + d * x)^3 / (16 a^3 f) + (c + d * x)^4 / (32 a^3 d) \end{aligned}$$

Rule 2207

```
Int[((b_)*(F_)^((g_.)*(e_.)+(f_.)*(x_.)))^(n_.)*((c_.)+(d_.)*(x_.))^(m_.), x_Symbol] :> Simp[(c + d*x)^m*(b*F^(g*(e + f*x)))^n/(f*g*n*Log[F]), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_.)*(a_.)+(b_.)*(x_.)))^(n_.), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```
Int[((c_.)+(d_.)*(x_.))^(m_)*((a_.)+(b_.)*tan[(e_.)+(f_.)*(x_.)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b))*(e + f*
```

```
x))/((2*a))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^3}{8a^3} - \frac{e^{-6e-6fx}(c+dx)^3}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^3}{8a^3} - \frac{3e^{-2e-2fx}(c+dx)^3}{8a^3} \right) dx \\
&= \frac{(c+dx)^4}{32a^3d} - \frac{\int e^{-6e-6fx}(c+dx)^3 dx}{8a^3} + \frac{3 \int e^{-4e-4fx}(c+dx)^3 dx}{8a^3} - \frac{3 \int e^{-2e-2fx}(c+dx)^3 dx}{8a^3} \\
&= \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} + \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} \\
&\quad + \frac{(c+dx)^4}{32a^3d} - \frac{d \int e^{-6e-6fx}(c+dx)^2 dx}{16a^3f} + \frac{(9d) \int e^{-4e-4fx}(c+dx)^2 dx}{32a^3f} \\
&\quad - \frac{(9d) \int e^{-2e-2fx}(c+dx)^2 dx}{16a^3f} \\
&= \frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} + \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} \\
&\quad + \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} \\
&\quad + \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} + \frac{(c+dx)^4}{32a^3d} - \frac{d^2 \int e^{-6e-6fx}(c+dx) dx}{48a^3f^2} \\
&\quad + \frac{(9d^2) \int e^{-4e-4fx}(c+dx) dx}{64a^3f^2} - \frac{(9d^2) \int e^{-2e-2fx}(c+dx) dx}{16a^3f^2} \\
&= \frac{d^2e^{-6e-6fx}(c+dx)}{288a^3f^3} - \frac{9d^2e^{-4e-4fx}(c+dx)}{256a^3f^3} + \frac{9d^2e^{-2e-2fx}(c+dx)}{32a^3f^3} \\
&\quad + \frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} + \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} \\
&\quad + \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} + \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} \\
&\quad + \frac{(c+dx)^4}{32a^3d} - \frac{d^3 \int e^{-6e-6fx} dx}{288a^3f^3} + \frac{(9d^3) \int e^{-4e-4fx} dx}{256a^3f^3} - \frac{(9d^3) \int e^{-2e-2fx} dx}{32a^3f^3} \\
&= \frac{d^3e^{-6e-6fx}}{1728a^3f^4} - \frac{9d^3e^{-4e-4fx}}{1024a^3f^4} + \frac{9d^3e^{-2e-2fx}}{64a^3f^4} + \frac{d^2e^{-6e-6fx}(c+dx)}{288a^3f^3} - \frac{9d^2e^{-4e-4fx}(c+dx)}{256a^3f^3} \\
&\quad + \frac{9d^2e^{-2e-2fx}(c+dx)}{32a^3f^3} + \frac{de^{-6e-6fx}(c+dx)^2}{96a^3f^2} - \frac{9de^{-4e-4fx}(c+dx)^2}{128a^3f^2} \\
&\quad + \frac{9de^{-2e-2fx}(c+dx)^2}{32a^3f^2} + \frac{e^{-6e-6fx}(c+dx)^3}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^3}{32a^3f} \\
&\quad + \frac{3e^{-2e-2fx}(c+dx)^3}{16a^3f} + \frac{(c+dx)^4}{32a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 4.57 (sec) , antiderivative size = 615, normalized size of antiderivative = 1.83

$$\int \frac{(c+dx)^3}{(a+a\coth(e+fx))^3} dx \\ = \frac{\text{csch}^3(e+fx) (81(32c^3f^3 + 24c^2df^2(3+4fx) + 12cd^2f(7+12fx+8f^2x^2) + d^3(45+84fx+72f^2x^2+32f^3x^3)))}{(a+a\coth(e+fx))^3}$$

[In] `Integrate[(c + d*x)^3/(a + a*Coth[e + f*x])^3,x]`

```
[Out] (Csch[e + f*x]^3*(81*(32*c^3*f^3 + 24*c^2*d*f^2*(3 + 4*f*x) + 12*c*d^2*f*(7 + 12*f*x + 8*f^2*x^2) + d^3*(45 + 84*f*x + 72*f^2*x^2 + 32*f^3*x^3))*Cosh[e + f*x] + 16*(36*c^3*f^3*(1 + 6*f*x) + 18*c^2*d*f^2*(1 + 6*f*x + 18*f^2*x^2) + 6*c*d^2*f*(1 + 6*f*x + 18*f^2*x^2 + 36*f^3*x^3) + d^3*(1 + 6*f*x + 18*f^2*x^2 + 36*f^3*x^3 + 54*f^4*x^4))*Cosh[3*(e + f*x)] + 4131*d^3*Sinh[e + f*x] + 8748*c*d^2*f*Sinh[e + f*x] + 9720*c^2*d*f^2*Sinh[e + f*x] + 7776*c^3*f^3*Sinh[e + f*x] + 8748*d^3*f*x*Sinh[e + f*x] + 19440*c*d^2*f^2*x*Sinh[e + f*x] + 23328*c^2*d*f^3*x*Sinh[e + f*x] + 9720*d^3*f^2*x^2*Sinh[e + f*x] + 23328*c*d^2*f^3*x^2*Sinh[e + f*x] + 7776*d^3*f^3*x^3*Sinh[e + f*x] - 16*d^3*Sinh[3*(e + f*x)] - 96*c*d^2*f*Sinh[3*(e + f*x)] - 288*c^2*d*f^2*Sinh[3*(e + f*x)] - 576*c^3*f^3*Sinh[3*(e + f*x)] - 96*d^3*f*x*Sinh[3*(e + f*x)] - 576*c*d^2*f^2*x*Sinh[3*(e + f*x)] - 1728*c^2*d*f^3*x*Sinh[3*(e + f*x)] + 3456*c^3*f^4*x*Sinh[3*(e + f*x)] - 288*d^3*f^2*x^2*Sinh[3*(e + f*x)] - 1728*c*d^2*f^3*x^2*Sinh[3*(e + f*x)] + 5184*c^2*d*f^4*x^2*Sinh[3*(e + f*x)] - 576*d^3*f^3*x^3*Sinh[3*(e + f*x)] + 3456*c*d^2*f^4*x^3*Sinh[3*(e + f*x)] + 864*d^3*f^4*x^4*Sinh[3*(e + f*x)]))/((27648*a^3*f^4*(1 + Coth[e + f*x])^3)
```

Maple [A] (verified)

Time = 0.58 (sec) , antiderivative size = 379, normalized size of antiderivative = 1.13

method	result
risch	$\frac{d^3x^4}{32a^3} + \frac{d^2cx^3}{8a^3} + \frac{3dc^2x^2}{16a^3} + \frac{c^3x}{8a^3} + \frac{c^4}{32a^3d} + \frac{3(4d^3x^3f^3 + 12cd^2f^3x^2 + 12c^2df^3x + 6d^3f^2x^2 + 4c^3f^3 + 12cd^2f^2x + 6c^2df^2 + 64a^3f^4)}{64a^3f^4}$
parallelrisch	$\frac{((864d^3x^4 + 3456d^2cx^3 + 5184dc^2x^2 + 3456c^3x)f^4 + (576d^3x^3 + 1728cd^2x^2 + 1728c^2dx + 8928c^3)f^3 + (288d^3x^2 + 576cd^2x + 10296c^2)f^2 + 144cd^2f + 144c^2d)}{144c^2d}$

```
[In] int((d*x+c)^3/(a+a*cOTH(f*x+e))^3,x,method=_RETURNVERBOSE)
```

```
[Out] 1/32/a^3*d^3*x^4+1/8/a^3*d^2*c*x^3+3/16/a^3*d*c^2*x^2+1/8/a^3*c^3*x+1/32/a^3/d*c^4+3/64*(4*d^3*f^3*x^3+12*c*d^2*f^3*x^2+12*c^2*d*f^3*x+6*d^3*f^2*x^2+4*c^3*f^3+12*c*d^2*f^2*x+6*c^2*d*f^2+6*d^3*f*x+6*c*d^2*f+3*d^3)/a^3/f^4*exp(-2*f*x-2*e)-3/1024*(32*d^3*f^3*x^3+96*c*d^2*f^3*x^2+96*c^2*d*f^3*x+24*d^3*f^2*x^2+32*c^3*f^3+48*c*d^2*f^2*x+24*c^2*d*f^2+12*d^3*f*x+12*c*d^2*f+3*d^3)/
```

$$a^3/f^4 \exp(-4*f*x - 4*e) + 1/1728 * (36*d^3*f^3*x^3 + 108*c*d^2*f^3*x^2 + 108*c^2*d*f^3*x + 18*d^3*f^2*x^2 + 36*c^3*f^3 + 36*c*d^2*f^2*x + 18*c^2*d*f^2 + 6*d^3*f^3*x + 6*c*d^2*f^3) / a^3/f^4 \exp(-6*f*x - 6*e)$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 844 vs. $2(298) = 596$.

Time = 0.28 (sec), antiderivative size = 844, normalized size of antiderivative = 2.51

$$\int \frac{(c + dx)^3}{(a + a \coth(e + fx))^3} dx \\ = \frac{16 (54 d^3 f^4 x^4 + 36 c^3 f^3 + 18 c^2 d f^2 + 6 c d^2 f + 36 (6 c d^2 f^4 + d^3 f^3) x^3 + d^3 + 18 (18 c^2 d f^4 + 6 c d^2 f^3 + d^3 f^2) x^2)}{(a + a \coth(e + fx))^3}$$

```
[In] integrate((d*x+c)^3/(a+a*coth(f*x+e))^3,x, algorithm="fricas")
[Out] 1/27648*(16*(54*d^3*f^4*x^4 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f + 36*(6*c*d^2*f^4 + d^3*f^3)*x^3 + d^3 + 18*(18*c^2*d*f^4 + 6*c*d^2*f^3 + d^3*f^2)*x^2 + 6*(36*c^3*f^4 + 18*c^2*d*f^3 + 6*c*d^2*f^2 + d^3*f)*x)*cosh(f*x + e)^3 + 48*(54*d^3*f^4*x^4 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f + 36*(6*c*d^2*f^4 + d^3*f^3)*x^3 + d^3 + 18*(18*c^2*d*f^4 + 6*c*d^2*f^3 + d^3*f^2)*x^2 + 6*(36*c^3*f^4 + 18*c^2*d*f^3 + 6*c*d^2*f^2 + d^3*f)*x)*cosh(f*x + e)*sinh(f*x + e)^2 + 16*(54*d^3*f^4*x^4 - 36*c^3*f^3 - 18*c^2*d*f^2 - 6*c*d^2*f + 36*(6*c*d^2*f^4 - d^3*f^3)*x^3 - d^3 + 18*(18*c^2*d*f^4 - 6*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(36*c^3*f^4 - 18*c^2*d*f^3 - 6*c*d^2*f^2 - d^3*f)*x)*sinh(f*x + e)^3 + 81*(32*d^3*f^3*x^3 + 32*c^3*f^3 + 72*c^2*d*f^2 + 84*c*d^2*f + 45*d^3 + 24*(4*c*d^2*f^3 + 3*d^3*f^2)*x^2 + 12*(8*c^2*d*f^3 + 12*c*d^2*f^2 + 7*d^3*f)*x)*cosh(f*x + e) + 3*(2592*d^3*f^3*x^3 + 2592*c^3*f^3 + 3240*c^2*d*f^2 + 2916*c*d^2*f + 1377*d^3 + 648*(12*c*d^2*f^3 + 5*d^3*f^2)*x^2 + 16*(54*d^3*f^4*x^4 - 36*c^3*f^3 - 18*c^2*d*f^2 - 6*c*d^2*f + 36*(6*c*d^2*f^4 - d^3*f^3)*x^3 - d^3 + 18*(18*c^2*d*f^4 - 6*c*d^2*f^3 - d^3*f^2)*x^2 + 6*(36*c^3*f^4 - 18*c^2*d*f^3 - 6*c*d^2*f^2 - d^3*f)*x)*cosh(f*x + e)^2 + 324*(24*c^2*d*f^3 + 20*c*d^2*f^2 + 9*d^3*f)*x)*sinh(f*x + e))/(a^3*f^4*cosh(f*x + e)^3 + 3*a^3*f^4*cosh(f*x + e)*sinh(f*x + e)^2 + a^3*f^4*sinh(f*x + e)^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3918 vs. $2(347) = 694$.

Time = 1.47 (sec), antiderivative size = 3918, normalized size of antiderivative = 11.66

$$\int \frac{(c + dx)^3}{(a + a \coth(e + fx))^3} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)**3/(a+a*coth(f*x+e))**3,x)`

[Out] Piecewise((864*c**3*f**4*x*tanh(e + f*x)**3/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 2592*c**3*f**4*x*tanh(e + f*x)**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 2592*c**3*f**4*x*tanh(e + f*x)/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 864*c**3*f**4*x/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 6048*c**3*f**3*tanh(e + f*x)**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 7776*c**3*f**3*tanh(e + f*x)/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 2880*c**3*f**3/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 1296*c**2*d*f**4*x**2*tanh(e + f*x)**3/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 3888*c**2*d*f**4*x**2*tanh(e + f*x)**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 1296*c**2*d*f**4*x**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 3888*c**2*d*f**4*x**2*tanh(e + f*x)/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 4536*c**2*d*f**3*x*tanh(e + f*x)/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 2376*c**2*d*f**3*x/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 6264*c**2*d*f**3*x*tanh(e + f*x)**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 9720*c**2*d*f**2*tanh(e + f*x)/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 4032*c**2*d*f**2/(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4)

$$\begin{aligned}
& *3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) \\
& + 864*c*d**2*f**4*x**3*tanh(e + f*x)**3/(6912*a**3*f**4*tanh(e + f*x)**3 + \\
& 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 6912*a** \\
& 3*f**4) + 2592*c*d**2*f**4*x**3*tanh(e + f*x)**2/(6912*a**3*f**4*tanh(e + f \\
& *x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + \\
& 6912*a**3*f**4) + 2592*c*d**2*f**4*x**3*tanh(e + f*x)/(6912*a**3*f**4*tanh \\
& (e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f \\
& *x) + 6912*a**3*f**4) + 864*c*d**2*f**4*x**3/(6912*a**3*f**4*tanh(e + f*x) \\
& **3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + 69 \\
& 12*a**3*f**4) - 6264*c*d**2*f**3*x**2*tanh(e + f*x)**3/(6912*a**3*f**4*tanh \\
& (e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f \\
& *x) + 6912*a**3*f**4) - 648*c*d**2*f**3*x**2*tanh(e + f*x)**2/(6912*a**3*f \\
& **4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*t \\
& anh(e + f*x) + 6912*a**3*f**4) + 4536*c*d**2*f**3*x**2*tanh(e + f*x)/(6912* \\
& a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f \\
& **4*tanh(e + f*x) + 6912*a**3*f**4) + 2376*c*d**2*f**3*x**2/(6912*a**3*f** \\
& 4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tan \\
& h(e + f*x) + 6912*a**3*f**4) - 5004*c*d**2*f**2*x*tanh(e + f*x)**3/(6912*a* \\
& *3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f* \\
& *4*tanh(e + f*x) + 6912*a**3*f**4) - 2484*c*d**2*f**2*x*tanh(e + f*x)**2/(6 \\
& 912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a \\
& **3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 4428*c*d**2*f**2*x*tanh(e + f*x) \\
& /(6912*a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 2073 \\
& 6*a**3*f**4*tanh(e + f*x) + 6912*a**3*f**4) + 3060*c*d**2*f**2*x/(6912*a**3 \\
& *f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4 \\
& *tanh(e + f*x) + 6912*a**3*f**4) + 5004*c*d**2*f*tanh(e + f*x)**2/(6912*a** \\
& 3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4 \\
& *tanh(e + f*x) + 6912*a**3*f**4) + 8748*c*d**2*f*tanh(e + f*x)/(6912*a**3*f \\
& **4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*t \\
& anh(e + f*x) + 6912*a**3*f**4) + 3936*c*d**2*f/(6912*a**3*f**4*tanh(e + f \\
& x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) + \\
& 6912*a**3*f**4) + 216*d**3*f**4*x**4*tanh(e + f*x)**3/(6912*a**3*f**4*tanh \\
& (e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f \\
& *x) + 6912*a**3*f**4) + 648*d**3*f**4*x**4*tanh(e + f*x)**2/(6912*a**3*f**4 \\
& *tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh \\
& (e + f*x) + 6912*a**3*f**4) + 648*d**3*f**4*x**4*tanh(e + f*x)/(6912*a**3*f \\
& **4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*t \\
& anh(e + f*x) + 6912*a**3*f**4) + 216*d**3*f**4*x**4/(6912*a**3*f**4*tanh(e \\
& + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh(e + f*x) \\
& + 6912*a**3*f**4) - 2088*d**3*f**3*x**3*tanh(e + f*x)**3/(6912*a**3*f**4*t \\
& anh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4*tanh \\
& (e + f*x) + 6912*a**3*f**4) - 216*d**3*f**3*x**3*tanh(e + f*x)**2/(6912*a**3 \\
& *f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*f**4 \\
& *tanh(e + f*x) + 6912*a**3*f**4) + 1512*d**3*f**3*x**3*tanh(e + f*x)/(6912* \\
& a**3*f**4*tanh(e + f*x)**3 + 20736*a**3*f**4*tanh(e + f*x)**2 + 20736*a**3*
\end{aligned}$$

$$\begin{aligned}
& f^{*4} \tanh(e + fx) + 6912*a^{*3}*f^{*4}) + 792*d^{*3}*f^{*3}*x^{*3}/(6912*a^{*3}*f^{*4}*t \\
& \text{anh}(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4}*\tanh(e \\
& + fx) + 6912*a^{*3}*f^{*4}) - 2502*d^{*3}*f^{*2}*x^{*2}*\tanh(e + fx)^{*3}/(6912*a^{*3} \\
& *f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4} \\
& *\tanh(e + fx) + 6912*a^{*3}*f^{*4}) - 1242*d^{*3}*f^{*2}*x^{*2}*\tanh(e + fx)^{*2}/(69 \\
& 12*a^{*3}*f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3} \\
& *f^{*4}*\tanh(e + fx) + 6912*a^{*3}*f^{*4}) + 2214*d^{*3}*f^{*2}*x^{*2}*\tanh(e + fx) \\
& /(6912*a^{*3}*f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 2073 \\
& 6*a^{*3}*f^{*4}*\tanh(e + fx) + 6912*a^{*3}*f^{*4}) + 1530*d^{*3}*f^{*2}*x^{*2}/(6912*a^{*3} \\
& *f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4} \\
& *\tanh(e + fx) + 6912*a^{*3}*f^{*4}) - 2211*d^{*3}*f*x*\tanh(e + fx)^{*3}/(6912*a^{*3} \\
& *f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4} \\
& *\tanh(e + fx) + 6912*a^{*3}*f^{*4}) - 1629*d^{*3}*f*x*\tanh(e + fx)^{*2}/(6912*a^{*3} \\
& *f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4} \\
& *\tanh(e + fx) + 6912*a^{*3}*f^{*4}) + 2115*d^{*3}*f*x*\tanh(e + fx)/(6912*a^{*3} \\
& *f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4} \\
& *\tanh(e + fx) + 6912*a^{*3}*f^{*4}) + 1725*d^{*3}*f*x/(6912*a^{*3}*f^{*4}*\tanh(e + \\
& fx)^{*3} + 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4}*\tanh(e + fx) + 6 \\
& 912*a^{*3}*f^{*4}) + 4131*d^{*3}*\tanh(e + fx)/(6912*a^{*3}*f^{*4}*\tanh(e + fx)^{*3} + \\
& 20736*a^{*3}*f^{*4}*\tanh(e + fx)^{*2} + 20736*a^{*3}*f^{*4}*\tanh(e + fx) + 6912*a^{*3} \\
& *f^{*4}) + 1952*d^{*3}/(6912*a^{*3}*f^{*4}*\tanh(e + fx)^{*3} + 20736*a^{*3}*f^{*4}*\tan \\
& h(e + fx)^{*2} + 20736*a^{*3}*f^{*4}*\tanh(e + fx) + 6912*a^{*3}*f^{*4}), \text{Ne}(f, 0)), \\
& ((c^{*3}*x + 3*c^{*2}*d*x^{*2}/2 + c*d^{*2}*x^{*3} + d^{*3}*x^{*4}/4)/(a*\coth(e) + a)^{*3} \\
& , \text{True}))
\end{aligned}$$

Maxima [A] (verification not implemented)

none

Time = 1.64 (sec), antiderivative size = 406, normalized size of antiderivative = 1.21

$$\begin{aligned}
& \int \frac{(c + dx)^3}{(a + a \coth(e + fx))^3} dx \\
& = \frac{1}{96} c^3 \left(\frac{12(fx + e)}{a^3 f} + \frac{18e^{-2fx-2e} - 9e^{-4fx-4e} + 2e^{-6fx-6e}}{a^3 f} \right) \\
& + \frac{(72f^2x^2e^{6e} + 108(2fxe^{4e} + e^{4e})e^{-2fx} - 27(4fxe^{2e} + e^{2e})e^{-4fx} + 4(6fx + 1)e^{-6fx})c^2de^{(-4fx)}}{384a^3f^2} \\
& + \frac{(288f^3x^3e^{6e} + 648(2f^2x^2e^{4e} + 2fxe^{4e} + e^{4e})e^{-2fx} - 81(8f^2x^2e^{2e} + 4fxe^{2e} + e^{2e})e^{-4fx})}{2304a^3f^3} \\
& + \frac{(864f^4x^4e^{6e} + 1296(4f^3x^3e^{4e} + 6f^2x^2e^{4e} + 6fxe^{4e} + 3e^{4e})e^{-2fx} - 81(32f^3x^3e^{2e} + 24f^2x^2e^{2e})}{27648a^3f^4}
\end{aligned}$$

[In] `integrate((d*x+c)^3/(a+a*coth(f*x+e))^3,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & 1/96*c^3*(12*(f*x + e)/(a^3*f) + (18*e^{-2*f*x - 2*e} - 9*e^{-4*f*x - 4*e}) \\ & + 2*e^{-6*f*x - 6*e})/(a^3*f)) + 1/384*(72*f^2*x^2*e^{(6*e)} + 108*(2*f*x*e^{(4*e)} + e^{(4*e)})*e^{-2*f*x} - 27*(4*f*x*e^{(2*e)} + e^{(2*e)})*e^{-4*f*x} + 4*(6*f*x + 1)*e^{-6*f*x})*c^2*d*e^{-6*e}/(a^3*f^2) + 1/2304*(288*f^3*x^3*e^{(6*e)} \\ &) + 648*(2*f^2*x^2*e^{(4*e)} + 2*f*x*e^{(4*e)} + e^{(4*e)})*e^{-2*f*x} - 81*(8*f^2*x^2*e^{(2*e)} + 4*f*x*e^{(2*e)} + e^{(2*e)})*e^{-4*f*x} + 8*(18*f^2*x^2 + 6*f*x + 1)*e^{-6*f*x})*c*d^2*e^{-6*e}/(a^3*f^3) + 1/27648*(864*f^4*x^4*e^{(6*e)} + 1296*(4*f^3*x^3*e^{(4*e)} + 6*f^2*x^2*e^{(4*e)} + 6*f*x*e^{(4*e)} + 3*e^{(4*e)})*e^{-2*f*x} - 81*(32*f^3*x^3*e^{(2*e)} + 24*f^2*x^2*e^{(2*e)} + 12*f*x*e^{(2*e)} + 3*e^{(2*e)})*e^{-4*f*x} + 16*(36*f^3*x^3 + 18*f^2*x^2 + 6*f*x + 1)*e^{-6*f*x})*d^3*e^{-6*e}/(a^3*f^4) \end{aligned}$$

Giac [A] (verification not implemented)

none

Time = 0.30 (sec), antiderivative size = 548, normalized size of antiderivative = 1.63

$$\int \frac{(c + dx)^3}{(a + a \coth(e + fx))^3} dx = \frac{(864 d^3 f^4 x^4 e^{(6 f x + 6 e)} + 3456 c d^2 f^4 x^3 e^{(6 f x + 6 e)} + 5184 c^2 d f^4 x^2 e^{(6 f x + 6 e)} + 5184 d^3 f^3 x^3 e^{(4 f x + 4 e)} - 2592 d^3 f^2 x^2 e^{(2 f x + 2 e)} + 15552 c d^2 f^3 x^2 e^{(4 f x + 4 e)} - 7776 c d^3 f^2 x^3 e^{(2 f x + 2 e)} + 1728 c d^2 f^3 x^2 e^{(4 f x + 4 e)} + 7776 c^2 d^2 f^3 x^3 e^{(2 f x + 2 e)} - 1944 c d^3 f^2 x^2 e^{(2 f x + 2 e)} + 1728 c^2 d^2 f^3 x^3 e^{(4 f x + 4 e)} + 5184 c^3 f^3 x^3 e^{(2 f x + 2 e)} + 15552 c d^2 f^3 x^2 e^{(4 f x + 4 e)} - 2592 c^3 f^3 x^3 e^{(2 f x + 2 e)} - 3888 c d^2 f^3 x^2 e^{(2 f x + 2 e)} + 576 c^3 f^3 x^3 + 576 c d^2 f^3 x^2 e^{(2 f x + 2 e)} + 7776 c^2 d^2 f^3 x^3 e^{(4 f x + 4 e)} + 7776 d^3 f^3 x^3 e^{(4 f x + 4 e)} - 1944 c^2 d^2 f^3 x^2 e^{(2 f x + 2 e)} - 972 d^3 f^3 x^3 e^{(2 f x + 2 e)} + 288 d^3 f^2 x^2 e^{(2 f x + 2 e)} + 5184 c^3 f^3 x^3 e^{(4 f x + 4 e)} + 15552 c d^2 f^3 x^2 e^{(4 f x + 4 e)} - 2592 c^3 f^3 x^3 e^{(2 f x + 2 e)} - 3888 d^3 f^3 x^3 e^{(4 f x + 4 e)} - 972 c d^2 f^3 x^2 e^{(4 f x + 4 e)} + 96 c d^2 f^3 x^2 e^{(2 f x + 2 e)} + 3888 d^3 f^3 x^3 e^{(4 f x + 4 e)} - 243 d^3 f^3 x^3 e^{(2 f x + 2 e)} + 16 d^3 f^3 x^3 e^{(-6 f x - 6 e)})}{(a^3 f^4)}$$

[In] integrate((d*x+c)^3/(a+a*coth(f*x+e))^3,x, algorithm="giac")

[Out]
$$\begin{aligned} & 1/27648*(864*d^3*f^4*x^4*e^{(6*f*x + 6*e)} + 3456*c*d^2*f^4*x^3*e^{(6*f*x + 6*e)} + 5184*c^2*d*f^4*x^2*e^{(6*f*x + 6*e)} - 2592*d^3*f^3*x^3*e^{(2*f*x + 2*e)} + 576*d^3*f^3*x^3 + 3456*c^3*f^4*x*x*e^{(6*f*x + 6*e)} + 15552*c*d^2*f^3*x^2*e^{(4*f*x + 4*e)} - 7776*c*d^3*f^2*x^3*e^{(2*f*x + 2*e)} + 1728*c*d^2*f^3*x^2 + 15552*c^2*d*f^3*x*x*e^{(4*f*x + 4*e)} + 7776*d^2*f^3*x*x*e^{(2*f*x + 2*e)} - 1944*d^3*f^2*x^2*e^{(2*f*x + 2*e)} + 1728*c^2*d*f^3*x + 288*d^3*f^2*x^2 + 5184*c^3*f^3*x^3*e^{(4*f*x + 4*e)} + 15552*c*d^2*f^3*x*x*e^{(4*f*x + 4*e)} - 2592*c^3*f^3*x^3*e^{(2*f*x + 2*e)} - 3888*c*d^2*f^3*x*x*e^{(2*f*x + 2*e)} + 576*c^3*f^3*x^3 + 576*c*d^2*f^3*x^2*x + 7776*c^2*d*f^3*x*x*e^{(4*f*x + 4*e)} + 7776*d^3*f*x*x*e^{(4*f*x + 4*e)} - 1944*c^2*d*f^3*x^2*e^{(2*f*x + 2*e)} - 972*d^3*f*x*x*e^{(2*f*x + 2*e)} + 288*c^2*d*f^3*x^2 + 96*d^3*f*x^2 + 7776*c*d^2*f*x*x*e^{(4*f*x + 4*e)} - 972*c*d^2*f*x*x*e^{(2*f*x + 2*e)} + 96*c*d^2*f*x + 3888*d^3*f*x*x*e^{(4*f*x + 4*e)} - 243*d^3*f*x*x*e^{(2*f*x + 2*e)} + 16*d^3*f*x*x*e^{(-6*f*x - 6*e)})/(a^3*f^4) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 2.17 (sec) , antiderivative size = 374, normalized size of antiderivative = 1.11

$$\begin{aligned} \int \frac{(c + dx)^3}{(a + a \coth(e + fx))^3} dx &= e^{-2e-2fx} \left(\frac{12c^3 f^3 + 18c^2 d f^2 + 18cd^2 f + 9d^3}{64a^3 f^4} + \frac{3d^3 x^3}{16a^3 f} \right. \\ &\quad \left. + \frac{9dx(2c^2 f^2 + 2cd f + d^2)}{32a^3 f^3} + \frac{9d^2 x^2(d + 2cf)}{32a^3 f^2} \right) \\ &- e^{-4e-4fx} \left(\frac{96c^3 f^3 + 72c^2 d f^2 + 36cd^2 f + 9d^3}{1024a^3 f^4} + \frac{3d^3 x^3}{32a^3 f} \right. \\ &\quad \left. + \frac{9dx(8c^2 f^2 + 4cd f + d^2)}{256a^3 f^3} + \frac{9d^2 x^2(d + 4cf)}{128a^3 f^2} \right) \\ &+ e^{-6e-6fx} \left(\frac{36c^3 f^3 + 18c^2 d f^2 + 6cd^2 f + d^3}{1728a^3 f^4} + \frac{d^3 x^3}{48a^3 f} \right. \\ &\quad \left. + \frac{dx(18c^2 f^2 + 6cd f + d^2)}{288a^3 f^3} + \frac{d^2 x^2(d + 6cf)}{96a^3 f^2} \right) \\ &+ \frac{c^3 x}{8a^3} + \frac{d^3 x^4}{32a^3} + \frac{3c^2 d x^2}{16a^3} + \frac{c d^2 x^3}{8a^3} \end{aligned}$$

[In] `int((c + d*x)^3/(a + a*coth(e + f*x))^3, x)`

[Out] `exp(- 2*e - 2*f*x)*((9*d^3 + 12*c^3*f^3 + 18*c^2*d*f^2 + 18*c*d^2*f)/(64*a^3*f^4) + (3*d^3*x^3)/(16*a^3*f) + (9*d*x*(d^2 + 2*c^2*f^2 + 2*c*d*f))/(32*a^3*f^3) + (9*d^2*x^2*(d + 2*c*f))/(32*a^3*f^2)) - exp(- 4*e - 4*f*x)*((9*d^3 + 96*c^3*f^3 + 72*c^2*d*f^2 + 36*c*d^2*f)/(1024*a^3*f^4) + (3*d^3*x^3)/(32*a^3*f) + (9*d*x*(d^2 + 8*c^2*f^2 + 4*c*d*f))/(256*a^3*f^3) + (9*d^2*x^2*(d + 4*c*f))/(128*a^3*f^2)) + exp(- 6*e - 6*f*x)*((d^3 + 36*c^3*f^3 + 18*c^2*d*f^2 + 6*c*d^2*f)/(1728*a^3*f^4) + (d^3*x^3)/(48*a^3*f) + (d*x*(d^2 + 18*c^2*f^2 + 6*c*d*f))/(288*a^3*f^3) + (d^2*x^2*(d + 6*c*f))/(96*a^3*f^2)) + (c^3*x)/(8*a^3) + (d^3*x^4)/(32*a^3) + (3*c^2*d*x^2)/(16*a^3) + (c*d^2*x^3)/(8*a^3)`

3.28 $\int \frac{(c+dx)^2}{(a+a \coth(e+fx))^3} dx$

Optimal result	189
Rubi [A] (verified)	189
Mathematica [A] (verified)	191
Maple [A] (verified)	192
Fricas [B] (verification not implemented)	192
Sympy [B] (verification not implemented)	193
Maxima [A] (verification not implemented)	194
Giac [A] (verification not implemented)	195
Mupad [B] (verification not implemented)	195

Optimal result

Integrand size = 20, antiderivative size = 246

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+a \coth(e+fx))^3} dx = & \frac{d^2 e^{-6e-6fx}}{864a^3 f^3} - \frac{3d^2 e^{-4e-4fx}}{256a^3 f^3} + \frac{3d^2 e^{-2e-2fx}}{32a^3 f^3} \\ & + \frac{de^{-6e-6fx}(c+dx)}{144a^3 f^2} - \frac{3de^{-4e-4fx}(c+dx)}{64a^3 f^2} \\ & + \frac{3de^{-2e-2fx}(c+dx)}{16a^3 f^2} + \frac{e^{-6e-6fx}(c+dx)^2}{48a^3 f} \\ & - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3 f} + \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3 f} + \frac{(c+dx)^3}{24a^3 d} \end{aligned}$$

```
[Out] 1/864*d^2*exp(-6*f*x-6*e)/a^3/f^3-3/256*d^2*exp(-4*f*x-4*e)/a^3/f^3+3/32*d^2*exp(-2*f*x-2*e)/a^3/f^3+1/144*d*exp(-6*f*x-6*e)*(d*x+c)/a^3/f^2-3/64*d*exp(-4*f*x-4*e)*(d*x+c)/a^3/f^2+3/16*d*exp(-2*f*x-2*e)*(d*x+c)/a^3/f^2+1/48*e^xp(-6*f*x-6*e)*(d*x+c)^2/a^3/f-3/32*exp(-4*f*x-4*e)*(d*x+c)^2/a^3/f+3/16*exp(-2*f*x-2*e)*(d*x+c)^2/a^3/f+1/24*(d*x+c)^3/a^3/d
```

Rubi [A] (verified)

Time = 0.19 (sec) , antiderivative size = 246, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.150, Rules used

$$= \{3810, 2207, 2225\}$$

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+a \coth(e+fx))^3} dx &= \frac{d(c+dx)e^{-6e-6fx}}{144a^3f^2} - \frac{3d(c+dx)e^{-4e-4fx}}{64a^3f^2} \\ &\quad + \frac{3d(c+dx)e^{-2e-2fx}}{16a^3f^2} + \frac{(c+dx)^2e^{-6e-6fx}}{48a^3f} \\ &\quad - \frac{3(c+dx)^2e^{-4e-4fx}}{32a^3f} + \frac{3(c+dx)^2e^{-2e-2fx}}{16a^3f} \\ &\quad + \frac{(c+dx)^3}{24a^3d} + \frac{d^2e^{-6e-6fx}}{864a^3f^3} - \frac{3d^2e^{-4e-4fx}}{256a^3f^3} + \frac{3d^2e^{-2e-2fx}}{32a^3f^3} \end{aligned}$$

[In] Int[(c + d*x)^2/(a + a*Coth[e + f*x])^3, x]

[Out] $(d^2 E^{-(-6e - 6f*x)})/(864*a^3*f^3) - (3*d^2 E^{-(-4e - 4f*x)})/(256*a^3*f^3)$
 $+ (3*d^2 E^{-(-2e - 2f*x)})/(32*a^3*f^3) + (d*E^{-(-6e - 6f*x)}*(c + d*x))$
 $/(144*a^3*f^2) - (3*d*E^{-(-4e - 4f*x)}*(c + d*x))/(64*a^3*f^2) + (3*d*E^{-(-2e - 2f*x)}*(c + d*x))$
 $/(16*a^3*f^2) + (E^{-(-6e - 6f*x)}*(c + d*x)^2)/(48*a^3*f) - (3*E^{-(-4e - 4f*x)}*(c + d*x)^2)$
 $/(32*a^3*f) + (3*E^{-(-2e - 2f*x)}*(c + d*x)^2)/(16*a^3*f) + (c + d*x)^3/(24*a^3*d)$

Rule 2207

```
Int[((b_)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^({m_.}), x_Symbol] :> Simp[(c + d*x)^m*((b*F^(g*(e + f*x)))^n/(f*g*n*Log[F])), x] - Dist[d*(m/(f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*(b*F^(g*(e + f*x)))^n, x], x] /; FreeQ[{F, b, c, d, e, f, g, n}, x] && GtQ[m, 0] && IntegerQ[2*m] && !TrueQ[$UseGamma]
```

Rule 2225

```
Int[((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.), x_Symbol] :> Simp[(F^(c*(a + b*x)))^n/(b*c*n*Log[F]), x] /; FreeQ[{F, a, b, c, n}, x]
```

Rule 3810

```
Int[((c_.) + (d_.)*(x_.))^({m_.})*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^2}{8a^3} - \frac{e^{-6e-6fx}(c+dx)^2}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^2}{8a^3} - \frac{3e^{-2e-2fx}(c+dx)^2}{8a^3} \right) dx \\ &= \frac{(c+dx)^3}{24a^3d} - \frac{\int e^{-6e-6fx}(c+dx)^2 dx}{8a^3} + \frac{3 \int e^{-4e-4fx}(c+dx)^2 dx}{8a^3} - \frac{3 \int e^{-2e-2fx}(c+dx)^2 dx}{8a^3} \end{aligned}$$

$$\begin{aligned}
&= \frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} + \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} \\
&\quad + \frac{(c+dx)^3}{24a^3d} - \frac{d \int e^{-6e-6fx}(c+dx) dx}{24a^3f} + \frac{(3d) \int e^{-4e-4fx}(c+dx) dx}{16a^3f} \\
&\quad - \frac{(3d) \int e^{-2e-2fx}(c+dx) dx}{8a^3f} \\
&= \frac{de^{-6e-6fx}(c+dx)}{144a^3f^2} - \frac{3de^{-4e-4fx}(c+dx)}{64a^3f^2} + \frac{3de^{-2e-2fx}(c+dx)}{16a^3f^2} \\
&\quad + \frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} + \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} \\
&\quad + \frac{(c+dx)^3}{24a^3d} - \frac{d^2 \int e^{-6e-6fx} dx}{144a^3f^2} + \frac{(3d^2) \int e^{-4e-4fx} dx}{64a^3f^2} - \frac{(3d^2) \int e^{-2e-2fx} dx}{16a^3f^2} \\
&= \frac{d^2e^{-6e-6fx}}{864a^3f^3} - \frac{3d^2e^{-4e-4fx}}{256a^3f^3} + \frac{3d^2e^{-2e-2fx}}{32a^3f^3} + \frac{de^{-6e-6fx}(c+dx)}{144a^3f^2} \\
&\quad - \frac{3de^{-4e-4fx}(c+dx)}{64a^3f^2} + \frac{3de^{-2e-2fx}(c+dx)}{16a^3f^2} + \frac{e^{-6e-6fx}(c+dx)^2}{48a^3f} \\
&\quad - \frac{3e^{-4e-4fx}(c+dx)^2}{32a^3f} + \frac{3e^{-2e-2fx}(c+dx)^2}{16a^3f} + \frac{(c+dx)^3}{24a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.17 (sec), antiderivative size = 371, normalized size of antiderivative = 1.51

$$\begin{aligned}
&\int \frac{(c+dx)^2}{(a+a \coth(e+fx))^3} dx \\
&= \frac{\operatorname{csch}^3(e+fx) (81(8c^2f^2 + 4cdf(3+4fx) + d^2(7+12fx+8f^2x^2)) \cosh(e+fx) + 8(18c^2f^2(1+6fx) + 729d^2 \operatorname{Sinh}[e+fx] + 1620*c*d*f*\operatorname{Sinh}[e+fx] + 1944*c^2*f^2 \operatorname{Sinh}[e+fx] + 1620*d^2*f*x*\operatorname{Sinh}[e+fx] + 3888*c*d*f^2*x*\operatorname{Sinh}[e+fx] + 1944*d^2*f^2*x^2*\operatorname{Sinh}[e+fx] - 8*d^2*\operatorname{Sinh}[3*(e+fx)] - 48*c*d*f*\operatorname{Sinh}[3*(e+fx)] - 144*c^2*f^2*\operatorname{Sinh}[3*(e+fx)] - 48*d^2*f*x*\operatorname{Sinh}[3*(e+fx)] - 288*c*d*f^2*x*\operatorname{Sinh}[3*(e+fx)] + 864*c^2*f^3*x*\operatorname{Sinh}[3*(e+fx)] - 144*d^2*f^2*x^2*\operatorname{Sinh}[3*(e+fx)] + 864*c*d*f^3*x^2*\operatorname{Sinh}[3*(e+fx)] + 288*d^2*f^3*x^3*\operatorname{Sinh}[3*(e+fx)]))/ (6912*a^3*f^3*(1+\operatorname{Coth}[e+fx])^3)
\end{aligned}$$

[In] Integrate[(c + d*x)^2/(a + a*Coth[e + f*x])^3, x]

[Out] $\operatorname{Csch}[e+f*x]^3 (81*(8*c^2*f^2 + 4*c*d*f*(3+4*f*x) + d^2*(7+12*f*x+8*f^2*x^2))*\operatorname{Cosh}[e+f*x] + 8*(18*c^2*f^2*(1+6*f*x) + 6*c*d*f*(1+6*f*x+18*f^2*x^2) + d^2*(1+6*f*x+18*f^2*x^2+36*f^3*x^3))*\operatorname{Cosh}[3*(e+f*x)] + 729*d^2*\operatorname{Sinh}[e+f*x] + 1620*c*d*f*\operatorname{Sinh}[e+f*x] + 1944*c^2*f^2*\operatorname{Sinh}[e+f*x] + 1620*d^2*f*x*\operatorname{Sinh}[e+f*x] + 3888*c*d*f^2*x*\operatorname{Sinh}[e+f*x] + 1944*d^2*f^2*x^2*\operatorname{Sinh}[e+f*x] - 8*d^2*\operatorname{Sinh}[3*(e+f*x)] - 48*c*d*f*\operatorname{Sinh}[3*(e+f*x)] - 144*c^2*f^2*\operatorname{Sinh}[3*(e+f*x)] - 48*d^2*f*x*\operatorname{Sinh}[3*(e+f*x)] - 288*c*d*f^2*x*\operatorname{Sinh}[3*(e+f*x)] + 864*c^2*f^3*x*\operatorname{Sinh}[3*(e+f*x)] - 144*d^2*f^2*x^2*\operatorname{Sinh}[3*(e+f*x)] + 864*c*d*f^3*x^2*\operatorname{Sinh}[3*(e+f*x)] + 288*d^2*f^3*x^3*\operatorname{Sinh}[3*(e+f*x)]))/ (6912*a^3*f^3*(1+\operatorname{Coth}[e+f*x])^3)$

Maple [A] (verified)

Time = 0.52 (sec) , antiderivative size = 223, normalized size of antiderivative = 0.91

method	result
risch	$\frac{d^2x^3}{24a^3} + \frac{dcx^2}{8a^3} + \frac{c^2x}{8a^3} + \frac{c^3}{24a^3d} + \frac{3(2d^2x^2f^2+4cd^2x+2c^2f^2+2d^2fx+2cdf+d^2)e^{-2fx-2e}}{32a^3f^3} - \frac{3(8d^2x^2f^2+16cd^2x+8c^2f^2+16c^3)f^3}{256a^3}$
parallelisch	$\frac{((288d^2x^3+864dcx^2+864c^2x)f^3+(144x^2d^2+288cdx+216c^2)f^2+(48xd^2+324cd)f+189d^2)\cosh(3fx+3e)+((288d^2x^3+864dcx^2+864c^2x)f^3+(144x^2d^2+288cdx+216c^2)f^2+(48xd^2+324cd)f+189d^2)\sinh(3fx+3e)}{a^3/f^3}$

[In] `int((d*x+c)^2/(a+a*coth(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{24/a^3*d^2*x^3+1/8/a^3*d*c*x^2+1/8/a^3*c^2*x+1/24/a^3/d*c^3+3/32*(2*d^2*f^2*x^2+4*c*d*f^2*x+2*c^2*f^2+2*d^2*f*x+2*c*d*f+d^2)/a^3/f^3*\exp(-2*f*x-2*e)-3/256*(8*d^2*f^2*x^2+16*c*d*f^2*x+8*c^2*f^2+4*d^2*f*x+4*c*d*f+d^2)/a^3/f^3*\exp(-4*f*x-4*e)+1/864*(18*d^2*f^2*x^2+36*c*d*f^2*x+18*c^2*f^2+6*d^2*f*x+6*c*d*f+d^2)/a^3/f^3*\exp(-6*f*x-6*e)}$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 532 vs. $2(217) = 434$.

Time = 0.26 (sec) , antiderivative size = 532, normalized size of antiderivative = 2.16

$$\begin{aligned} & \int \frac{(c+dx)^2}{(a+a\coth(e+fx))^3} dx \\ &= \frac{8(36d^2f^3x^3+18c^2f^2+6cdf+18(6cdf^3+d^2f^2)x^2+d^2+6(18c^2f^3+6cdf^2+d^2f)x)\cosh(fx+e)^3+}{\dots} \end{aligned}$$

[In] `integrate((d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="fricas")`

[Out] $\frac{1}{6912}*(8*(36*d^2*f^3*x^3+18*c^2*f^2+6*c*d*f+18*(6*c*d*f^3+d^2*f^2)*x^2+d^2+6*(18*c^2*f^3+6*c*d*f^2+d^2*f)*x)*\cosh(f*x+e)^3+24*(36*d^2*f^3*x^3+18*c^2*f^2+6*c*d*f+18*(6*c*d*f^3+d^2*f^2)*x^2+d^2+6*(18*c^2*f^3+6*c*d*f^2+d^2*f)*x)*\cosh(f*x+e)*\sinh(f*x+e)^2+8*(36*d^2*f^3*x^3-18*c^2*f^2-6*c*d*f+18*(6*c*d*f^3-d^2*f^2)*x^2-d^2+6*(18*c^2*f^3-6*c*d*f^2-d^2*f)*x)*\sinh(f*x+e)^3+81*(8*d^2*f^2*x^2+8*c^2*f^2+12*c*d*f+7*d^2+4*(4*c*d*f^2+3*d^2*f)*x)*\cosh(f*x+e)+3*(648*d^2*f^2*x^2+648*c^2*f^2+540*c*d*f+8*(36*d^2*f^3*x^3-18*c^2*f^2*x^2-6*c*d*f+18*(6*c*d*f^3-d^2*f^2)*x^2-d^2+6*(18*c^2*f^3-6*c*d*f^2-d^2*f)*x)*\cosh(f*x+e)^2+243*d^2+108*(12*c*d*f^2+5*d^2*f)*x)*\sinh(f*x+e)/(a^3*f^3*\cosh(f*x+e)^3+3*a^3*f^3*\cosh(f*x+e)^2*\sinh(f*x+e)+3*a^3*f^3*\cosh(f*x+e)*\sinh(f*x+e)^2+a^3*f^3*\sinh(f*x+e)^3)$

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2443 vs. $2(252) = 504$.

Time = 1.21 (sec) , antiderivative size = 2443, normalized size of antiderivative = 9.93

$$\int \frac{(c + dx)^2}{(a + a \coth(e + fx))^3} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)**2/(a+a*coth(f*x+e))**3,x)`

[Out] Piecewise((216*c**2*f**3*x*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 648*c**2*f**3*x*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 648*c**2*f**3*x*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 216*c**2*f**3*x/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 1512*c**2*f**2*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 1944*c**2*f**2*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 216*c*d*f**3*x**2*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 648*c*d*f**3*x**2*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 5184*a**3*f**3*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 1044*c*d*f**2*x*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 108*c*d*f**2*x*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 756*c*d*f**2*x*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 396*c*d*f**2*x/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 1044*c*d*f*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 1620*c*d*f*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 672*c*d*f/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 72*d**2*f**2

$$\begin{aligned}
& 3*x**3*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 216*d**2*f**3*x**3*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 216*d**2*f**3*x**3*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 72*d**2*f**3*x**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 522*d**2*f**2*x**2*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 54*d**2*f**2*x**2*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 378*d**2*f**2*x**2*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 198*d**2*f**2*x**2*(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 417*d**2*f*x*tanh(e + f*x)**3/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) - 207*d**2*f*x*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 369*d**2*f*x*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 255*d**2*f*x/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 417*d**2*tanh(e + f*x)**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 729*d**2*tanh(e + f*x)/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3) + 328*d**2/(1728*a**3*f**3*tanh(e + f*x)**3 + 5184*a**3*f**3*tanh(e + f*x)**2 + 5184*a**3*f**3*tanh(e + f*x) + 1728*a**3*f**3), Ne(f, 0)), ((c**2*x + c*d*x**2 + d**2*x**3/3)/(a*coth(e) + a)**3, True))
\end{aligned}$$

Maxima [A] (verification not implemented)

none

Time = 1.15 (sec) , antiderivative size = 254, normalized size of antiderivative = 1.03

$$\begin{aligned}
& \int \frac{(c + dx)^2}{(a + a \coth(e + fx))^3} dx \\
&= \frac{1}{96} c^2 \left(\frac{12(fx + e)}{a^3 f} + \frac{18e^{-2fx-2e} - 9e^{-4fx-4e} + 2e^{-6fx-6e}}{a^3 f} \right) \\
&+ \frac{(72f^2x^2e^{6e} + 108(2fxe^{4e} + e^{4e}))e^{-2fx} - 27(4fxe^{2e} + e^{2e})e^{-4fx} + 4(6fx + 1)e^{-6fx})cde^{-4fx}}{576a^3f^2} \\
&+ \frac{(288f^3x^3e^{6e} + 648(2f^2x^2e^{4e} + 2fxe^{4e} + e^{4e}))e^{-2fx} - 81(8f^2x^2e^{2e} + 4fxe^{2e} + e^{2e})e^{-4fx}}{6912a^3f^3}
\end{aligned}$$

[In] `integrate((d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & \frac{1}{96} c^2 (12 (f x + e) / (a^3 f) + (18 e^{-2 f x - 2 e} - 9 e^{-4 f x - 4 e} \\ & + 2 e^{-6 f x - 6 e}) / (a^3 f)) + \frac{1}{576} (72 f^2 x^2 e^{6 e} + 108 (2 f x e^{4 e} \\ & + e^{4 (4 e)}) e^{-2 f x} - 27 (4 f x e^{2 e} + e^{2 e}) e^{-4 f x} + 4 (6 \\ & f x + 1) e^{-6 f x}) c d e^{-6 e} / (a^3 f^2) + \frac{1}{6912} (288 f^3 x^3 e^{6 e} \\ & + 648 (2 f^2 x^2 e^{4 e} + 2 f x e^{4 e} + e^{4 e}) e^{-2 f x} - 81 (8 f^2 x^2 \\ & x^2 e^{2 e} + 4 f x e^{2 e} + e^{2 e}) e^{-4 f x} + 8 (18 f^2 x^2 + 6 f x + 1) e^{-6 f x}) d^2 e^{-6 e} / (a^3 f^3) \end{aligned}$$

Giac [A] (verification not implemented)

none

Time = 0.28 (sec), antiderivative size = 315, normalized size of antiderivative = 1.28

$$\begin{aligned} & \int \frac{(c + dx)^2}{(a + a \coth(e + fx))^3} dx \\ & = \frac{(288 d^2 f^3 x^3 e^{(6 f x + 6 e)} + 864 c d f^3 x^2 e^{(6 f x + 6 e)} + 864 c^2 f^3 x e^{(6 f x + 6 e)} + 1296 d^2 f^2 x^2 e^{(4 f x + 4 e)} - 648 d^2 f^2 x^2 e^{(2 \\ & f x + 2 e)} + 144 d^2 f^2 x^2 + 2592 c d f^2 x e^{(4 f x + 4 e)} - 1296 c d f^2 x e^{(2 f x + 2 e)} + 288 c d f^2 x + 1296 c^2 f^2 x e^{(4 f x + 4 e)} + 1296 d^2 f x e^{(4 f x + 4 e)} - 648 c^2 f^2 x e^{(2 f x + 2 e)} - 324 d^2 f x e^{(2 f x + 2 e)} + 144 c^2 f^2 + 48 d^2 f x + 1296 c d f x e^{(4 f x + 4 e)} - 324 c d f x e^{(2 f x + 2 e)} + 48 c d f + 648 d^2 f x e^{(4 f x + 4 e)} - 81 d^2 f x e^{(2 f x + 2 e)} + 8 d^2 e^2) e^{-(-6 f x - 6 e)} / (a^3 f^3) } \end{aligned}$$

[In] `integrate((d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="giac")`

[Out]
$$\begin{aligned} & \frac{1}{6912} (288 d^2 f^3 x^3 e^{(6 f x + 6 e)} + 864 c d f^3 x^2 e^{(6 f x + 6 e)} + 864 c^2 f^3 x e^{(6 f x + 6 e)} + 1296 d^2 f^2 x^2 e^{(4 f x + 4 e)} - 648 d^2 f^2 x^2 e^{(2 f x + 2 e)} + 144 d^2 f^2 x^2 + 2592 c d f^2 x e^{(4 f x + 4 e)} - 1296 c d f^2 x e^{(2 f x + 2 e)} + 288 c d f^2 x + 1296 c^2 f^2 x e^{(4 f x + 4 e)} + 1296 d^2 f x e^{(4 f x + 4 e)} - 648 c^2 f^2 x e^{(2 f x + 2 e)} - 324 d^2 f x e^{(2 f x + 2 e)} + 144 c^2 f^2 + 48 d^2 f x + 1296 c d f x e^{(4 f x + 4 e)} - 324 c d f x e^{(2 f x + 2 e)} + 48 c d f + 648 d^2 f x e^{(4 f x + 4 e)} - 81 d^2 f x e^{(2 f x + 2 e)} + 8 d^2 e^2) e^{-(-6 f x - 6 e)} / (a^3 f^3) \end{aligned}$$

Mupad [B] (verification not implemented)

Time = 2.14 (sec), antiderivative size = 234, normalized size of antiderivative = 0.95

$$\begin{aligned} & \int \frac{(c + dx)^2}{(a + a \coth(e + fx))^3} dx = e^{-6 e - 6 f x} \left(\frac{18 c^2 f^2 + 6 c d f + d^2}{864 a^3 f^3} + \frac{d^2 x^2}{48 a^3 f} + \frac{d x (d + 6 c f)}{144 a^3 f^2} \right) \\ & + e^{-2 e - 2 f x} \left(\frac{6 c^2 f^2 + 6 c d f + 3 d^2}{32 a^3 f^3} + \frac{3 d^2 x^2}{16 a^3 f} \right. \\ & \left. + \frac{3 d x (d + 2 c f)}{16 a^3 f^2} \right) - e^{-4 e - 4 f x} \left(\frac{24 c^2 f^2 + 12 c d f + 3 d^2}{256 a^3 f^3} \right. \\ & \left. + \frac{3 d^2 x^2}{32 a^3 f} + \frac{3 d x (d + 4 c f)}{64 a^3 f^2} \right) + \frac{c^2 x}{8 a^3} + \frac{d^2 x^3}{24 a^3} + \frac{c d x^2}{8 a^3} \end{aligned}$$

[In] $\int ((c + d*x)^2 / (a + a*\coth(e + f*x))^3, x)$

[Out]
$$\begin{aligned} & \exp(-6e - 6fx) * ((d^2 + 18c^2f^2 + 6cd^2f) / (864a^3f^3) + (d^2x^2) / \\ & (48a^3f) + (dx(d + 6cf)) / (144a^3f^2)) + \exp(-2e - 2fx) * ((3d^2 \\ & + 6c^2f^2 + 6cd^2f) / (32a^3f^3) + (3d^2x^2) / (16a^3f) + (3dx(d + \\ & 2cf)) / (16a^3f^2)) - \exp(-4e - 4fx) * ((3d^2 + 24c^2f^2 + 12cd^2f) \\ & / (256a^3f^3) + (3d^2x^2) / (32a^3f) + (3dx(d + 4cf)) / (64a^3f^2)) \\ & + (c^2x) / (8a^3) + (d^2x^3) / (24a^3) + (cdx^2) / (8a^3) \end{aligned}$$

3.29 $\int \frac{c+dx}{(a+a \coth(e+fx))^3} dx$

Optimal result	197
Rubi [A] (verified)	197
Mathematica [A] (verified)	200
Maple [A] (verified)	200
Fricas [A] (verification not implemented)	200
Sympy [B] (verification not implemented)	201
Maxima [A] (verification not implemented)	202
Giac [A] (verification not implemented)	202
Mupad [B] (verification not implemented)	203

Optimal result

Integrand size = 18, antiderivative size = 183

$$\begin{aligned} \int \frac{c+dx}{(a+a \coth(e+fx))^3} dx = & \frac{11dx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a \coth(e+fx))^3} \\ & - \frac{c+dx}{6f(a+a \coth(e+fx))^3} - \frac{5d}{96af^2(a+a \coth(e+fx))^2} \\ & - \frac{c+dx}{8af(a+a \coth(e+fx))^2} - \frac{11d}{96f^2(a^3+a^3 \coth(e+fx))} \\ & - \frac{c+dx}{8f(a^3+a^3 \coth(e+fx))} \end{aligned}$$

[Out] $11/96*d*x/a^3/f-1/16*d*x^2/a^3+1/8*x*(d*x+c)/a^3-1/36*d/f^2/(a+a*\coth(f*x+e))^3+1/6*(-d*x-c)/f/(a+a*\coth(f*x+e))^3-5/96*d/a/f^2/(a+a*\coth(f*x+e))^2+1/8*(-d*x-c)/a/f/(a+a*\coth(f*x+e))^2-11/96*d/f^2/(a^3+a^3*\coth(f*x+e))+1/8*(-d*x-c)/f/(a^3+a^3*\coth(f*x+e))$

Rubi [A] (verified)

Time = 0.18 (sec) , antiderivative size = 183, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 3, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used

$$= \{3560, 8, 3811\}$$

$$\begin{aligned} \int \frac{c+dx}{(a+a \coth(e+fx))^3} dx = & -\frac{c+dx}{8f(a^3 \coth(e+fx)+a^3)} + \frac{x(c+dx)}{8a^3} \\ & -\frac{11d}{96f^2(a^3 \coth(e+fx)+a^3)} + \frac{11dx}{96a^3f} - \frac{dx^2}{16a^3} \\ & -\frac{c+dx}{8af(a \coth(e+fx)+a)^2} - \frac{c+dx}{6f(a \coth(e+fx)+a)^3} \\ & -\frac{5d}{96af^2(a \coth(e+fx)+a)^2} - \frac{d}{36f^2(a \coth(e+fx)+a)^3} \end{aligned}$$

[In] $\text{Int}[(c + d*x)/(a + a*\text{Coth}[e + f*x])^3, x]$

[Out] $(11*d*x)/(96*a^3*f) - (d*x^2)/(16*a^3) + (x*(c + d*x))/(8*a^3) - d/(36*f^2*(a + a*\text{Coth}[e + f*x])^3) - (c + d*x)/(6*f*(a + a*\text{Coth}[e + f*x])^3) - (5*d)/(96*a*f^2*(a + a*\text{Coth}[e + f*x])^2) - (c + d*x)/(8*a*f*(a + a*\text{Coth}[e + f*x])^2) - (11*d)/(96*f^2*(a^3 + a^3*\text{Coth}[e + f*x])) - (c + d*x)/(8*f*(a^3 + a^3*\text{Coth}[e + f*x]))$

Rule 8

$\text{Int}[a_, x_{\text{Symbol}}] :> \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3560

$\text{Int}[(a_ + b_*)\tan[(c_ + d_*)(x_)]^{(n_)}, x_{\text{Symbol}}] :> \text{Simp}[a*((a + b*Tan[c + d*x])^n/(2*b*d*n)), x] + \text{Dist}[1/(2*a), \text{Int}[(a + b*Tan[c + d*x])^{(n + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{LtQ}[n, 0]$

Rule 3811

$\text{Int}[(c_ + d_*)(x_)]^{(m_)}*((a_ + b_*)\tan[(e_ + f_*)(x_)]^{(n_)}, x_{\text{Symbol}}) :> \text{With}[\{u = \text{IntHide}[(a + b*Tan[e + f*x])^n, x]\}, \text{Dist}[(c + d*x)^m, u, x] - \text{Dist}[d*m, \text{Int}[\text{Dist}[(c + d*x)^{(m - 1)}, u, x], x], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, -1] \&& \text{GtQ}[m, 0]$

Rubi steps

$$\begin{aligned} \text{integral} = & \frac{x(c+dx)}{8a^3} - \frac{c+dx}{6f(a+a \coth(e+fx))^3} - \frac{c+dx}{8af(a+a \coth(e+fx))^2} \\ & - \frac{c+dx}{8f(a^3+a^3 \coth(e+fx))} - d \int \left(\frac{x}{8a^3} - \frac{1}{6f(a+a \coth(e+fx))^3} \right. \\ & \quad \left. - \frac{1}{8af(a+a \coth(e+fx))^2} - \frac{1}{8f(a^3+a^3 \coth(e+fx))} \right) dx \end{aligned}$$

$$\begin{aligned}
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{c+dx}{6f(a+a\coth(e+fx))^3} - \frac{c+dx}{8af(a+a\coth(e+fx))^2} \\
&\quad - \frac{c+dx}{8f(a^3+a^3\coth(e+fx))} + \frac{d \int \frac{1}{a^3+a^3\coth(e+fx)} dx}{8f} \\
&\quad + \frac{d \int \frac{1}{(a+a\coth(e+fx))^3} dx}{6f} + \frac{d \int \frac{1}{(a+a\coth(e+fx))^2} dx}{8af} \\
&= -\frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a\coth(e+fx))^3} - \frac{c+dx}{6f(a+a\coth(e+fx))^3} \\
&\quad - \frac{d}{32af^2(a+a\coth(e+fx))^2} - \frac{c+dx}{8af(a+a\coth(e+fx))^2} - \frac{d}{16f^2(a^3+a^3\coth(e+fx))} \\
&\quad - \frac{c+dx}{8f(a^3+a^3\coth(e+fx))} + \frac{d \int 1 dx}{16a^3f} + \frac{d \int \frac{1}{a+a\coth(e+fx)} dx}{16a^2f} \\
&\quad + \frac{d \int \frac{1}{(a+a\coth(e+fx))^2} dx}{12af} \\
&= \frac{dx}{16a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a\coth(e+fx))^3} \\
&\quad - \frac{c+dx}{6f(a+a\coth(e+fx))^3} - \frac{5d}{96af^2(a+a\coth(e+fx))^2} \\
&\quad - \frac{c+dx}{8af(a+a\coth(e+fx))^2} - \frac{3d}{32f^2(a^3+a^3\coth(e+fx))} \\
&\quad - \frac{c+dx}{8f(a^3+a^3\coth(e+fx))} + \frac{d \int 1 dx}{32a^3f} + \frac{d \int \frac{1}{a+a\coth(e+fx)} dx}{24a^2f} \\
&= \frac{3dx}{32a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a\coth(e+fx))^3} - \frac{c+dx}{6f(a+a\coth(e+fx))^3} \\
&\quad - \frac{5d}{96af^2(a+a\coth(e+fx))^2} - \frac{c+dx}{8af(a+a\coth(e+fx))^2} \\
&\quad - \frac{11d}{96f^2(a^3+a^3\coth(e+fx))} - \frac{c+dx}{8f(a^3+a^3\coth(e+fx))} + \frac{d \int 1 dx}{48a^3f} \\
&= \frac{11dx}{96a^3f} - \frac{dx^2}{16a^3} + \frac{x(c+dx)}{8a^3} - \frac{d}{36f^2(a+a\coth(e+fx))^3} - \frac{c+dx}{6f(a+a\coth(e+fx))^3} \\
&\quad - \frac{5d}{96af^2(a+a\coth(e+fx))^2} - \frac{c+dx}{8af(a+a\coth(e+fx))^2} \\
&\quad - \frac{11d}{96f^2(a^3+a^3\coth(e+fx))} - \frac{c+dx}{8f(a^3+a^3\coth(e+fx))}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.76 (sec) , antiderivative size = 185, normalized size of antiderivative = 1.01

$$\int \frac{c + dx}{(a + a \coth(e + fx))^3} dx \\ = \frac{\text{csch}^3(e + fx) (27(4cf + d(3 + 4fx)) \cosh(e + fx) + 4(6cf(1 + 6fx) + d(1 + 6fx + 18f^2x^2)) \cosh(3(e + fx)))}{\dots}$$

[In] `Integrate[(c + d*x)/(a + a*Coth[e + f*x])^3, x]`

[Out] $\frac{(Csch[e + f*x]^3*(27*(4*c*f + d*(3 + 4*f*x))*Cosh[e + f*x] + 4*(6*c*f*(1 + 6*f*x) + d*(1 + 6*f*x + 18*f^2*x^2))*Cosh[3*(e + f*x)] + 135*d*Sinh[e + f*x] + 324*c*f*Sinh[e + f*x] + 324*d*f*x*Sinh[e + f*x] - 4*d*Sinh[3*(e + f*x)] - 24*c*f*Sinh[3*(e + f*x)] - 24*d*f*x*Sinh[3*(e + f*x)] + 144*c*f^2*x*Sinh[3*(e + f*x)] + 72*d*f^2*x^2*Sinh[3*(e + f*x)]))}{(1152*a^3*f^2*(1 + Coth[e + f*x])^3)}$

Maple [A] (verified)

Time = 0.43 (sec) , antiderivative size = 102, normalized size of antiderivative = 0.56

method	result
risch	$\frac{dx^2}{16a^3} + \frac{cx}{8a^3} + \frac{3(2dxf+2cf+d)e^{-2fx-2e}}{32a^3f^2} - \frac{3(4dxf+4cf+d)e^{-4fx-4e}}{128a^3f^2} + \frac{(6dxf+6cf+d)e^{-6fx-6e}}{288a^3f^2}$
parallelrisch	$\frac{36\left(\left(\frac{dx}{2}+c\right)f-\frac{29d}{12}\right)xf\tanh(fx+e)^3+\left((54dx^2+108cx)f^2+(-9dx+252c)f+87d\right)\tanh(fx+e)^2+\left((54dx^2+108cx)f^2+(63dx+189c)f+216d\right)\tanh(fx+e)}{288f^2a^3(1+\tanh(fx+e))^3}$

[In] `int((d*x+c)/(a+a*coth(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{16}*d*x^2/a^3+1/8*a^3*c*x+3/32*(2*d*f*x+2*c*f+d)/a^3/f^2*exp(-2*f*x-2*e)-3/128*(4*d*f*x+4*c*f+d)/a^3/f^2*exp(-4*f*x-4*e)+1/288*(6*d*f*x+6*c*f+d)/a^3/f^2*exp(-6*f*x-6*e)$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 286, normalized size of antiderivative = 1.56

$$\int \frac{c + dx}{(a + a \coth(e + fx))^3} dx \\ = \frac{4(18df^2x^2 + 6cf + 6(6cf^2 + df)x + d) \cosh(fx + e)^3 + 12(18df^2x^2 + 6cf + 6(6cf^2 + df)x + d) \cosh(fx + e)^2 + 1152(a + a \coth(e + fx))^3}{\dots}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="fricas")`

```
[Out] 1/1152*(4*(18*d*f^2*x^2 + 6*c*f + 6*(6*c*f^2 + d*f)*x + d)*cosh(f*x + e)^3
+ 12*(18*d*f^2*x^2 + 6*c*f + 6*(6*c*f^2 + d*f)*x + d)*cosh(f*x + e)*sinh(f*x + e)^2 + 4*(18*d*f^2*x^2 - 6*c*f + 6*(6*c*f^2 - d*f)*x - d)*sinh(f*x + e)^3 + 27*(4*d*f*x + 4*c*f + 3*d)*cosh(f*x + e) + 3*(108*d*f*x + 4*(18*d*f^2*x^2 - 6*c*f + 6*(6*c*f^2 - d*f)*x - d)*cosh(f*x + e)^2 + 108*c*f + 45*d)*sinh(f*x + e))/(a^3*f^2*cosh(f*x + e)^3 + 3*a^3*f^2*cosh(f*x + e)^2*sinh(f*x + e) + 3*a^3*f^2*cosh(f*x + e)*sinh(f*x + e)^2 + a^3*f^2*sinh(f*x + e)^3)
```

Sympy [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1287 vs. $2(170) = 340$.

Time = 1.11 (sec), antiderivative size = 1287, normalized size of antiderivative = 7.03

$$\int \frac{c + dx}{(a + a \coth(e + fx))^3} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)/(a+a*coth(f*x+e))**3,x)
```

```
[Out] Piecewise((36*c*f**2*x*tanh(e + f*x)**3/(288*a**3*f**2*tanh(e + f*x)**3 + 8
64*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2
) + 108*c*f**2*x*tanh(e + f*x)**2/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**
3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 10
8*c*f**2*x*tanh(e + f*x)/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tan
nh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 36*c*f**2*x
/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**
3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 252*c*f*tanh(e + f*x)**2/(288*a**3*
f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh
(e + f*x) + 288*a**3*f**2) + 324*c*f*tanh(e + f*x)/(288*a**3*f**2*tanh(e +
f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 28
8*a**3*f**2) + 120*c*f/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh
(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 18*d*f**2*x**
2*tanh(e + f*x)**3/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e +
f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 54*d*f**2*x**2*tan
nh(e + f*x)**2/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)
)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 54*d*f**2*x**2*tanh(e
+ f*x)/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**
3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 18*d*f**2*x**2/(288*a**3*f**2
*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e +
f*x) + 288*a**3*f**2) - 87*d*f*x*tanh(e + f*x)**3/(288*a**3*f**2*tanh(e +
f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 2
88*a**3*f**2) - 9*d*f*x*tanh(e + f*x)**2/(288*a**3*f**2*tanh(e + f*x)**3 +
864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2
) + 63*d*f*x*tanh(e + f*x)/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2
*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 33*d*f*x
/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**
```

```
3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 87*d*tanh(e + f*x)**2/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 135*d*tanh(e + f*x)/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2) + 56*d/(288*a**3*f**2*tanh(e + f*x)**3 + 864*a**3*f**2*tanh(e + f*x)**2 + 864*a**3*f**2*tanh(e + f*x) + 288*a**3*f**2), Ne(f, 0)), ((c*x + d*x**2/2)/(a*coth(e) + a)**3, True))
```

Maxima [A] (verification not implemented)

none

Time = 0.68 (sec) , antiderivative size = 138, normalized size of antiderivative = 0.75

$$\int \frac{c + dx}{(a + a \coth(e + fx))^3} dx \\ = \frac{1}{96} c \left(\frac{12(fx + e)}{a^3 f} + \frac{18e^{-2fx-2e} - 9e^{-4fx-4e} + 2e^{-6fx-6e}}{a^3 f} \right) \\ + \frac{(72f^2x^2e^{6e} + 108(2fxe^{4e} + e^{4e})e^{-2fx} - 27(4fxe^{2e} + e^{2e})e^{-4fx} + 4(6fx + 1)e^{-6fx})de^{-6fx}}{1152a^3f^2}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="maxima")`

[Out] $\frac{1}{96} c \left(\frac{12(fx + e)}{a^3 f} + \frac{18e^{-2fx-2e} - 9e^{-4fx-4e} + 2e^{-6fx-6e}}{a^3 f} \right) + \frac{1}{1152} (72f^2x^2e^{6e} + 108(2fxe^{4e} + e^{4e})e^{-2fx} - 27(4fxe^{2e} + e^{2e})e^{-4fx} + 4(6fx + 1)e^{-6fx}) * d * e^{-6fx} / (a^3 f^2)$

Giac [A] (verification not implemented)

none

Time = 0.30 (sec) , antiderivative size = 142, normalized size of antiderivative = 0.78

$$\int \frac{c + dx}{(a + a \coth(e + fx))^3} dx \\ = \frac{(72df^2x^2e^{6fx+6e} + 144cf^2xe^{6fx+6e} + 216dfxe^{4fx+4e} - 108dfxe^{2fx+2e} + 24dfx + 216cfe^{4fx+4e})e^{-6fx}}{1152a^3f^2}$$

[In] `integrate((d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="giac")`

[Out] $\frac{1}{1152} (72d^2f^2x^2e^{6fx+6e} + 144c^2f^2x^2e^{6fx+6e} + 216cfe^{4fx+4e} - 108d^2fx^2e^{2fx+2e} + 24d^2fx + 216c^2fe^{4fx+4e} - 108c^2fe^{2fx+2e} + 24c^2f + 108d^2e^{4fx+4e} - 27de^{2fx+2e} + 4d^2)e^{-6fx} / (a^3 f^2)$

Mupad [B] (verification not implemented)

Time = 2.01 (sec) , antiderivative size = 127, normalized size of antiderivative = 0.69

$$\begin{aligned} \int \frac{c + dx}{(a + a \coth(e + fx))^3} dx &= e^{-6e-6fx} \left(\frac{d + 6cf}{288a^3f^2} + \frac{dx}{48a^3f} \right) \\ &\quad + e^{-2e-2fx} \left(\frac{3d + 6cf}{32a^3f^2} + \frac{3dx}{16a^3f} \right) \\ &\quad - e^{-4e-4fx} \left(\frac{3d + 12cf}{128a^3f^2} + \frac{3dx}{32a^3f} \right) + \frac{dx^2}{16a^3} + \frac{cx}{8a^3} \end{aligned}$$

[In] `int((c + d*x)/(a + a*coth(e + f*x))^3, x)`

[Out] `exp(- 6*e - 6*f*x)*((d + 6*c*f)/(288*a^3*f^2) + (d*x)/(48*a^3*f)) + exp(- 2*e - 2*f*x)*((3*d + 6*c*f)/(32*a^3*f^2) + (3*d*x)/(16*a^3*f)) - exp(- 4*e - 4*f*x)*((3*d + 12*c*f)/(128*a^3*f^2) + (3*d*x)/(32*a^3*f)) + (d*x^2)/(16*a^3) + (c*x)/(8*a^3)`

3.30 $\int \frac{1}{(c+dx)(a+a \coth(e+fx))^3} dx$

Optimal result	204
Rubi [A] (verified)	205
Mathematica [A] (verified)	209
Maple [A] (verified)	210
Fricas [A] (verification not implemented)	210
Sympy [F]	210
Maxima [A] (verification not implemented)	211
Giac [A] (verification not implemented)	211
Mupad [F(-1)]	211

Optimal result

Integrand size = 20, antiderivative size = 437

$$\begin{aligned} \int \frac{1}{(c+dx)(a+a \coth(e+fx))^3} dx = & -\frac{3 \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\ & + \frac{3 \cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{8a^3d} \\ & - \frac{\cosh(6e - \frac{6cf}{d}) \text{Chi}(\frac{6cf}{d} + 6fx)}{8a^3d} \\ & + \frac{\log(c+dx)}{8a^3d} + \frac{\text{Chi}(\frac{6cf}{d} + 6fx) \sinh(6e - \frac{6cf}{d})}{8a^3d} \\ & - \frac{3\text{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{8a^3d} \\ & + \frac{3\text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{8a^3d} \\ & + \frac{3 \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\ & - \frac{3 \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\ & - \frac{3 \cosh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{8a^3d} \\ & + \frac{3 \sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{8a^3d} \\ & + \frac{\cosh(6e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx)}{8a^3d} \\ & - \frac{\sinh(6e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx)}{8a^3d} \end{aligned}$$

[Out]
$$\begin{aligned} & -\frac{1}{8} \text{Chi}(6*c*f/d + 6*f*x) * \cosh(-6*e + 6*c*f/d) / a^3/d + \frac{3}{8} \text{Chi}(4*c*f/d + 4*f*x) * \cos \\ & h(-4*e + 4*c*f/d) / a^3/d - \frac{3}{8} \text{Chi}(2*c*f/d + 2*f*x) * \cosh(-2*e + 2*c*f/d) / a^3/d + \frac{1}{8} * 1 \\ & n(d*x + c) / a^3/d + \frac{3}{8} * \cosh(-2*e + 2*c*f/d) * \text{Shi}(2*c*f/d + 2*f*x) / a^3/d - \frac{3}{8} * \cosh(-4* \\ & e + 4*c*f/d) * \text{Shi}(4*c*f/d + 4*f*x) / a^3/d + \frac{1}{8} * \cosh(-6*e + 6*c*f/d) * \text{Shi}(6*c*f/d + 6*f* \\ & x) / a^3/d - \frac{1}{8} * \text{Chi}(6*c*f/d + 6*f*x) * \sinh(-6*e + 6*c*f/d) / a^3/d + \frac{1}{8} * \text{Shi}(6*c*f/d + 6*f* \\ & x) * \sinh(-6*e + 6*c*f/d) / a^3/d + \frac{3}{8} * \text{Chi}(4*c*f/d + 4*f*x) * \sinh(-4*e + 4*c*f/d) / a^3 \\ & / d - \frac{3}{8} * \text{Shi}(4*c*f/d + 4*f*x) * \sinh(-4*e + 4*c*f/d) / a^3/d - \frac{3}{8} * \text{Chi}(2*c*f/d + 2*f*x) * \sinh \\ & (-2*e + 2*c*f/d) / a^3/d + \frac{3}{8} * \text{Shi}(2*c*f/d + 2*f*x) * \sinh(-2*e + 2*c*f/d) / a^3/d \end{aligned}$$

Rubi [A] (verified)

Time = 1.45 (sec), antiderivative size = 437, normalized size of antiderivative = 1.00, number of steps used = 53, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.350, Rules used = {3809, 3384, 3379, 3382, 3393, 5556, 5578}

$$\begin{aligned} \int \frac{1}{(c + dx)(a + a \coth(e + fx))^3} dx = & \frac{3 \text{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{8a^3d} \\ & + \frac{\text{Chi}(6xf + \frac{6cf}{d}) \sinh(6e - \frac{6cf}{d})}{8a^3d} \\ & - \frac{3 \text{Chi}(4xf + \frac{4cf}{d}) \sinh(4e - \frac{4cf}{d})}{8a^3d} \\ & - \frac{3 \text{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{8a^3d} \\ & + \frac{3 \text{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{8a^3d} \\ & - \frac{\text{Chi}(6xf + \frac{6cf}{d}) \cosh(6e - \frac{6cf}{d})}{8a^3d} \\ & - \frac{3 \sinh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{8a^3d} \\ & + \frac{3 \sinh(4e - \frac{4cf}{d}) \text{Shi}(4xf + \frac{4cf}{d})}{8a^3d} \\ & - \frac{\sinh(6e - \frac{6cf}{d}) \text{Shi}(6xf + \frac{6cf}{d})}{8a^3d} \\ & + \frac{3 \cosh(2e - \frac{2cf}{d}) \text{Shi}(2xf + \frac{2cf}{d})}{8a^3d} \\ & - \frac{3 \cosh(4e - \frac{4cf}{d}) \text{Shi}(4xf + \frac{4cf}{d})}{8a^3d} \\ & + \frac{\cosh(6e - \frac{6cf}{d}) \text{Shi}(6xf + \frac{6cf}{d})}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \end{aligned}$$

[In] $\text{Int}[1/((c + d*x)*(a + a*\text{Coth}[e + f*x])^3), x]$

[Out]
$$\begin{aligned} & (-3 * \text{Cosh}[2*e - (2*c*f)/d] * \text{CoshIntegral}[(2*c*f)/d + 2*f*x]) / (8*a^3*d) + (3*C \\ & \text{osh}[4*e - (4*c*f)/d] * \text{CoshIntegral}[(4*c*f)/d + 4*f*x]) / (8*a^3*d) - (\text{Cosh}[6*e \end{aligned}$$

$$\begin{aligned}
& - (6*c*f)/d] * \text{CoshIntegral}[(6*c*f)/d + 6*f*x]/(8*a^3*d) + \text{Log}[c + d*x]/(8*a^3*d) \\
& + (\text{CoshIntegral}[(6*c*f)/d + 6*f*x]*\text{Sinh}[6*e - (6*c*f)/d])/(8*a^3*d) \\
& - (3*\text{CoshIntegral}[(4*c*f)/d + 4*f*x]*\text{Sinh}[4*e - (4*c*f)/d])/(8*a^3*d) + (3*\text{CoshIntegral}[(2*c*f)/d + 2*f*x]*\text{Sinh}[2*e - (2*c*f)/d])/(8*a^3*d) \\
& + (3*\text{Cosh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) - (3*\text{Sinh}[2*e - (2*c*f)/d]*\text{SinhIntegral}[(2*c*f)/d + 2*f*x])/(8*a^3*d) \\
& - (3*\text{Cosh}[4*e - (4*c*f)/d]*\text{SinhIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) + (3*\text{Sinh}[4*e - (4*c*f)/d]*\text{SinhIntegral}[(4*c*f)/d + 4*f*x])/(8*a^3*d) \\
& + (\text{Cosh}[6*e - (6*c*f)/d]*\text{SinhIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d) - (\text{Sinh}[6*e - (6*c*f)/d]*\text{SinhIntegral}[(6*c*f)/d + 6*f*x])/(8*a^3*d)
\end{aligned}$$
Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3393

```
Int[((c_.) + (d_.)*(x_.))^m_*sin[(e_.) + (f_.)*(x_.)]^n_, x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sin[e + f*x]^n, x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && (!RationalQ[m] || (GeQ[m, -1] && LtQ[m, 1]))
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_.))^m_*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_.)]^p_*((c_.) + (d_.)*(x_.))^m_*Sinh[(a_.) + (b_.)*(x_.)]^n_, x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a +
```

```
b*x]^n*Cosh[a + b*x]^p, x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] &
& IGtQ[p, 0]
```

Rule 5578

```
Int[((e_.) + (f_ .)*(x_))^(m_ .)*Sinh[(a_ .) + (b_ .)*(x_)]^(p_ .)*Sinh[(c_ .) +
(d_ .)*(x_)]^(q_ .), x_Symbol] :> Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Sinh[c + d*x]^q, x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0]
] && IGtQ[q, 0] && IntegerQ[m]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{1}{8a^3(c+dx)} - \frac{3 \cosh(2e+2fx)}{8a^3(c+dx)} + \frac{3 \cosh^2(2e+2fx)}{8a^3(c+dx)} - \frac{\cosh^3(2e+2fx)}{8a^3(c+dx)} \right. \\
&\quad + \frac{3 \sinh(2e+2fx)}{8a^3(c+dx)} + \frac{3 \cosh^2(2e+2fx) \sinh(2e+2fx)}{8a^3(c+dx)} + \frac{3 \sinh^2(2e+2fx)}{8a^3(c+dx)} \\
&\quad \left. + \frac{\sinh^3(2e+2fx)}{8a^3(c+dx)} - \frac{3 \sinh(4e+4fx)}{8a^3(c+dx)} - \frac{3 \sinh(2e+2fx) \sinh(4e+4fx)}{16a^3(c+dx)} \right) dx \\
&= \frac{\log(c+dx)}{8a^3d} - \frac{\int \frac{\cosh^3(2e+2fx)}{c+dx} dx}{8a^3} + \frac{\int \frac{\sinh^3(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sinh(2e+2fx) \sinh(4e+4fx)}{c+dx} dx}{16a^3} \\
&\quad - \frac{3 \int \frac{\cosh(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\cosh^2(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\sinh(2e+2fx)}{c+dx} dx}{8a^3} \\
&\quad + \frac{3 \int \frac{\cosh^2(2e+2fx) \sinh(2e+2fx)}{c+dx} dx}{8a^3} + \frac{3 \int \frac{\sinh^2(2e+2fx)}{c+dx} dx}{8a^3} - \frac{3 \int \frac{\sinh(4e+4fx)}{c+dx} dx}{8a^3} \\
&= \frac{\log(c+dx)}{8a^3d} + \frac{i \int \left(\frac{3i \sinh(2e+2fx)}{4(c+dx)} - \frac{i \sinh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} \\
&\quad - \frac{\int \left(\frac{3 \cosh(2e+2fx)}{4(c+dx)} + \frac{\cosh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} - \frac{3 \int \left(-\frac{\cosh(2e+2fx)}{2(c+dx)} + \frac{\cosh(6e+6fx)}{2(c+dx)} \right) dx}{16a^3} \\
&\quad - \frac{3 \int \left(\frac{1}{2(c+dx)} - \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} + \frac{3 \int \left(\frac{1}{2(c+dx)} + \frac{\cosh(4e+4fx)}{2(c+dx)} \right) dx}{8a^3} \\
&\quad + \frac{3 \int \left(\frac{\sinh(2e+2fx)}{4(c+dx)} + \frac{\sinh(6e+6fx)}{4(c+dx)} \right) dx}{8a^3} - \frac{(3 \cosh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
&\quad - \frac{(3 \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} \\
&\quad + \frac{(3 \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} - \frac{(3 \sinh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{8a^3} \\
&\quad + \frac{(3 \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3} - \frac{(3 \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{8a^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3 \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad - \frac{3 \operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{8a^3d} + \frac{3 \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{8a^3d} \\
&\quad + \frac{3 \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3 \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3 \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{8a^3d} - \frac{\int \frac{\cosh(6e+6fx)}{c+dx} dx}{32a^3} + \frac{\int \frac{\sinh(6e+6fx)}{c+dx} dx}{32a^3} \\
&\quad - \frac{3 \int \frac{\cosh(6e+6fx)}{c+dx} dx}{32a^3} + \frac{3 \int \frac{\sinh(6e+6fx)}{c+dx} dx}{32a^3} + 2 \frac{3 \int \frac{\cosh(4e+4fx)}{c+dx} dx}{16a^3} \\
&= -\frac{3 \cosh(2e - \frac{2cf}{d}) \operatorname{Chi}(\frac{2cf}{d} + 2fx)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad - \frac{3 \operatorname{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{8a^3d} + \frac{3 \operatorname{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{8a^3d} \\
&\quad + \frac{3 \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3 \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3 \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(\frac{4cf}{d} + 4fx)}{8a^3d} - \frac{\cosh(6e - \frac{6cf}{d}) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} \\
&\quad + \frac{\cosh(6e - \frac{6cf}{d}) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} - \frac{(3 \cosh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} \\
&\quad + \frac{(3 \cosh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} + \frac{\sinh(6e - \frac{6cf}{d}) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} \\
&\quad - \frac{\sinh(6e - \frac{6cf}{d}) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} + \frac{(3 \sinh(6e - \frac{6cf}{d})) \int \frac{\cosh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} \\
&\quad - \frac{(3 \sinh(6e - \frac{6cf}{d})) \int \frac{\sinh(\frac{6cf}{d} + 6fx)}{c+dx} dx}{32a^3} + 2 \left(\frac{(3 \cosh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{16a^3} \right. \\
&\quad \left. + \frac{(3 \sinh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{16a^3} \right)
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3 \cosh(2e - \frac{2cf}{d}) \text{Chi}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{\cosh(6e - \frac{6cf}{d}) \text{Chi}(\frac{6cf}{d} + 6fx)}{8a^3d} + \frac{\log(c + dx)}{8a^3d} \\
&\quad + \frac{\text{Chi}(\frac{6cf}{d} + 6fx) \sinh(6e - \frac{6cf}{d})}{8a^3d} - \frac{3\text{Chi}(\frac{4cf}{d} + 4fx) \sinh(4e - \frac{4cf}{d})}{8a^3d} \\
&\quad + \frac{3\text{Chi}(\frac{2cf}{d} + 2fx) \sinh(2e - \frac{2cf}{d})}{8a^3d} + \frac{3 \cosh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} \\
&\quad - \frac{3 \sinh(2e - \frac{2cf}{d}) \text{Shi}(\frac{2cf}{d} + 2fx)}{8a^3d} - \frac{3 \cosh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{8a^3d} \\
&\quad + 2 \left(\frac{3 \cosh(4e - \frac{4cf}{d}) \text{Chi}(\frac{4cf}{d} + 4fx)}{16a^3d} + \frac{3 \sinh(4e - \frac{4cf}{d}) \text{Shi}(\frac{4cf}{d} + 4fx)}{16a^3d} \right) \\
&\quad + \frac{\cosh(6e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx)}{8a^3d} - \frac{\sinh(6e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx)}{8a^3d}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.32 (sec), antiderivative size = 312, normalized size of antiderivative = 0.71

$$\begin{aligned}
&\int \frac{1}{(c + dx)(a + a \coth(e + fx))^3} dx \\
&= \frac{\text{csch}^3(e + fx)(\cosh(fx) + \sinh(fx))^3 \left(\cosh(3e) \log(f(c + dx)) + \log(f(c + dx)) \sinh(3e) + (-\cosh(e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx) - \sinh(e - \frac{6cf}{d}) \text{Shi}(\frac{6cf}{d} + 6fx)) \right)}{8a^3d}
\end{aligned}$$

```

[In] Integrate[1/((c + d*x)*(a + a*Coth[e + f*x])^3),x]
[Out] (Csch[e + f*x]^3*(Cosh[f*x] + Sinh[f*x])^3*(Cosh[3*e]*Log[f*(c + d*x)] + Log[f*(c + d*x)]*Sinh[3*e] + (-Cosh[e - (4*c*f)/d] + Sinh[e - (4*c*f)/d])*(-3*CoshIntegral[(4*f*(c + d*x))/d] + Cosh[2*e - (2*c*f)/d]*CoshIntegral[(6*f*(c + d*x))/d] - CoshIntegral[(6*f*(c + d*x))/d]*Sinh[2*e - (2*c*f)/d] + 3*CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[2*e - (2*c*f)/d] + Sinh[2*e - (2*c*f)/d]) - 3*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*f*(c + d*x))/d] - 3*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*f*(c + d*x))/d] + 3*SinhIntegral[(4*f*(c + d*x))/d] - Cosh[2*e - (2*c*f)/d]*SinhIntegral[(6*f*(c + d*x))/d] + Sinh[2*e - (2*c*f)/d]*SinhIntegral[(6*f*(c + d*x))/d]))/(8*a^3*d*(1 + Coth[e + f*x])^3)

```

Maple [A] (verified)

Time = 0.37 (sec) , antiderivative size = 151, normalized size of antiderivative = 0.35

method	result
risch	$\frac{\ln(dx+c)}{8a^3d} + \frac{e^{\frac{6cf-6de}{d}} \text{Ei}_1\left(6fx+6e+\frac{6cf-6de}{d}\right)}{8a^3d} - \frac{3e^{\frac{4cf-4de}{d}} \text{Ei}_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{8a^3d} + \frac{3e^{\frac{2cf-2de}{d}} \text{Ei}_1\left(2fx+2e+\frac{2cf-2de}{d}\right)}{8a^3d}$

[In] `int(1/(d*x+c)/(a+a*coth(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & \frac{1}{8} \ln(d*x+c)/a^3/d + \frac{1}{8} a^3/d * \exp(6*(c*f-d*e)/d) * \text{Ei}(1, 6*f*x+6*e+6*(c*f-d*e)/d) \\ & - \frac{3}{8} a^3/d * \exp(4*(c*f-d*e)/d) * \text{Ei}(1, 4*f*x+4*e+4*(c*f-d*e)/d) + \frac{3}{8} a^3/d * \exp(2*(c*f-d*e)/d) * \text{Ei}(1, 2*f*x+2*e+2*(c*f-d*e)/d) \end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.25 (sec) , antiderivative size = 195, normalized size of antiderivative = 0.45

$$\int \frac{1}{(c+dx)(a+a \coth(e+fx))^3} dx =$$

$$\frac{-3 \operatorname{Ei}\left(-\frac{2 (dfx+cf)}{d}\right) \cosh\left(-\frac{2 (de-cf)}{d}\right)-3 \operatorname{Ei}\left(-\frac{4 (dfx+cf)}{d}\right) \cosh\left(-\frac{4 (de-cf)}{d}\right)+\operatorname{Ei}\left(-\frac{6 (dfx+cf)}{d}\right) \cosh\left(-\frac{6 (de-cf)}{d}\right)}{a^3}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -\frac{1}{8} (3 \operatorname{Ei}(-2*(d*f*x + c*f)/d) * \cosh(-2*(d*e - c*f)/d) - 3 \operatorname{Ei}(-4*(d*f*x + c*f)/d) * \cosh(-4*(d*e - c*f)/d) + \operatorname{Ei}(-6*(d*f*x + c*f)/d) * \cosh(-6*(d*e - c*f)/d) \\ & + 3 \operatorname{Ei}(-2*(d*f*x + c*f)/d) * \sinh(-2*(d*e - c*f)/d) - 3 \operatorname{Ei}(-4*(d*f*x + c*f)/d) * \sinh(-4*(d*e - c*f)/d) + \operatorname{Ei}(-6*(d*f*x + c*f)/d) * \sinh(-6*(d*e - c*f)/d) \\ & - \log(d*x + c)) / (a^3*d) \end{aligned}$$

Sympy [F]

$$\int \frac{1}{(c+dx)(a+a \coth(e+fx))^3} dx$$

$$= \frac{\int \frac{1}{c \coth^3(e+fx)+3c \coth^2(e+fx)+3c \coth(e+fx)+c+dx \coth^3(e+fx)+3dx \coth^2(e+fx)+3dx \coth(e+fx)+dx} dx}{a^3}$$

[In] `integrate(1/(d*x+c)/(a+a*coth(f*x+e))**3,x)`

[Out]
$$\text{Integral}\left(\frac{1}{c \coth(e+f*x)^3 + 3c \coth(e+f*x)^2 + 3c \coth(e+f*x) + c + d*x \coth(e+f*x)^3 + 3d*x \coth(e+f*x)^2 + 3d*x \coth(e+f*x) + d*x}, x\right)/a^3$$

Maxima [A] (verification not implemented)

none

Time = 2.29 (sec) , antiderivative size = 114, normalized size of antiderivative = 0.26

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^3} dx = \frac{e^{\left(-6e + \frac{6cf}{d}\right)} E_1\left(\frac{6(dx+c)f}{d}\right)}{8a^3d} - \frac{3e^{\left(-4e + \frac{4cf}{d}\right)} E_1\left(\frac{4(dx+c)f}{d}\right)}{8a^3d} + \frac{3e^{\left(-2e + \frac{2cf}{d}\right)} E_1\left(\frac{2(dx+c)f}{d}\right)}{8a^3d} + \frac{\log(dx + c)}{8a^3d}$$

```
[In] integrate(1/(d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="maxima")
[Out] 1/8*e^(-6*e + 6*c*f/d)*exp_integral_e(1, 6*(d*x + c)*f/d)/(a^3*d) - 3/8*e^(-4*e + 4*c*f/d)*exp_integral_e(1, 4*(d*x + c)*f/d)/(a^3*d) + 3/8*e^(-2*e + 2*c*f/d)*exp_integral_e(1, 2*(d*x + c)*f/d)/(a^3*d) + 1/8*log(d*x + c)/(a^3*d)
```

Giac [A] (verification not implemented)

none

Time = 0.29 (sec) , antiderivative size = 104, normalized size of antiderivative = 0.24

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^3} dx = -\frac{\left(3 \operatorname{Ei}\left(-\frac{2(df+cf)}{d}\right) e^{\left(4e + \frac{2cf}{d}\right)} - 3 \operatorname{Ei}\left(-\frac{4(df+cf)}{d}\right) e^{\left(2e + \frac{4cf}{d}\right)} + \operatorname{Ei}\left(-\frac{6(df+cf)}{d}\right) e^{\left(\frac{6cf}{d}\right)} - e^{(6e)} \log(dx + c)\right)}{8a^3d}$$

```
[In] integrate(1/(d*x+c)/(a+a*coth(f*x+e))^3,x, algorithm="giac")
[Out] -1/8*(3*Ei(-2*(d*f*x + c*f)/d)*e^(4*e + 2*c*f/d) - 3*Ei(-4*(d*f*x + c*f)/d)*e^(2*e + 4*c*f/d) + Ei(-6*(d*f*x + c*f)/d)*e^(6*c*f/d) - e^(6*e)*log(d*x + c))*e^(-6*e)/(a^3*d)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)(a + a \coth(e + fx))^3} dx = \int \frac{1}{(a + a \coth(e + fx))^3 (c + dx)} dx$$

```
[In] int(1/((a + a*coth(e + f*x))^3*(c + d*x)),x)
[Out] int(1/((a + a*coth(e + f*x))^3*(c + d*x)), x)
```

3.31 $\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^3} dx$

Optimal result	213
Rubi [A] (verified)	214
Mathematica [A] (verified)	222
Maple [A] (verified)	223
Fricas [A] (verification not implemented)	223
Sympy [F]	224
Maxima [A] (verification not implemented)	225
Giac [A] (verification not implemented)	225
Mupad [F(-1)]	226

Optimal result

Integrand size = 20, antiderivative size = 692

$$\begin{aligned}
\int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^3} dx = & -\frac{1}{8a^3d(c+dx)} + \frac{9 \cosh(2e+2fx)}{32a^3d(c+dx)} \\
& - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} \\
& + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3 \cosh(6e+6fx)}{32a^3d(c+dx)} \\
& + \frac{3f \cosh(2e-\frac{2cf}{d}) \text{Chi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
& - \frac{3f \cosh(4e-\frac{4cf}{d}) \text{Chi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
& + \frac{3f \cosh(6e-\frac{6cf}{d}) \text{Chi}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
& - \frac{3f \text{Chi}(\frac{6cf}{d}+6fx) \sinh(6e-\frac{6cf}{d})}{4a^3d^2} \\
& + \frac{3f \text{Chi}(\frac{4cf}{d}+4fx) \sinh(4e-\frac{4cf}{d})}{2a^3d^2} \\
& - \frac{3f \text{Chi}(\frac{2cf}{d}+2fx) \sinh(2e-\frac{2cf}{d})}{4a^3d^2} \\
& - \frac{15 \sinh(2e+2fx)}{32a^3d(c+dx)} \\
& - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
& + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} - \frac{3 \sinh(6e+6fx)}{32a^3d(c+dx)} \\
& - \frac{3f \cosh(2e-\frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
& + \frac{3f \sinh(2e-\frac{2cf}{d}) \text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
& + \frac{3f \cosh(4e-\frac{4cf}{d}) \text{Shi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
& - \frac{3f \sinh(4e-\frac{4cf}{d}) \text{Shi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
& - \frac{3f \cosh(6e-\frac{6cf}{d}) \text{Shi}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
& + \frac{3f \sinh(6e-\frac{6cf}{d}) \text{Shi}(\frac{6cf}{d}+6fx)}{4a^3d^2}
\end{aligned}$$

[Out] $-1/8/a^3/d/(d*x+c)+3/4*f*Chi(6*c*f/d+6*f*x)*cosh(-6*e+6*c*f/d)/a^3/d^2-3/2*f*Chi(4*c*f/d+4*f*x)*cosh(-4*e+4*c*f/d)/a^3/d^2+3/4*f*Chi(2*c*f/d+2*f*x)*co$

$$\begin{aligned}
& \operatorname{sh}(-2*e+2*c*f/d)/a^3/d^2+9/32*\cosh(2*f*x+2*e)/a^3/d/(d*x+c)-3/8*\cosh(2*f*x+ \\
& 2*e)^2/a^3/d/(d*x+c)+1/8*\cosh(2*f*x+2*e)^3/a^3/d/(d*x+c)+3/32*\cosh(6*f*x+6* \\
& e)/a^3/d/(d*x+c)-3/4*f*\cosh(-2*e+2*c*f/d)*\operatorname{Shi}(2*c*f/d+2*f*x)/a^3/d^2+3/2*f* \\
& \cosh(-4*e+4*c*f/d)*\operatorname{Shi}(4*c*f/d+4*f*x)/a^3/d^2-3/4*f*\cosh(-6*e+6*c*f/d)*\operatorname{Shi}(\\
& 6*c*f/d+6*f*x)/a^3/d^2+3/4*f*\operatorname{Chi}(6*c*f/d+6*f*x)*\sinh(-6*e+6*c*f/d)/a^3/d^2- \\
& 3/4*f*\operatorname{Shi}(6*c*f/d+6*f*x)*\sinh(-6*e+6*c*f/d)/a^3/d^2-3/2*f*\operatorname{Chi}(4*c*f/d+4*f*x) \\
& *\sinh(-4*e+4*c*f/d)/a^3/d^2+3/2*f*\operatorname{Shi}(4*c*f/d+4*f*x)*\sinh(-4*e+4*c*f/d)/a^ \\
& 3/d^2+3/4*f*\operatorname{Chi}(2*c*f/d+2*f*x)*\sinh(-2*e+2*c*f/d)/a^3/d^2-3/4*f*\operatorname{Shi}(2*c*f/d \\
& +2*f*x)*\sinh(-2*e+2*c*f/d)/a^3/d^2-15/32*\sinh(2*f*x+2*e)/a^3/d/(d*x+c)-3/8* \\
& \sinh(2*f*x+2*e)^2/a^3/d/(d*x+c)-1/8*\sinh(2*f*x+2*e)^3/a^3/d/(d*x+c)+3/8*\sinh \\
& (4*f*x+4*e)/a^3/d/(d*x+c)-3/32*\sinh(6*f*x+6*e)/a^3/d/(d*x+c)
\end{aligned}$$

Rubi [A] (verified)

Time = 1.41 (sec), antiderivative size = 692, normalized size of antiderivative = 1.00, number of steps used = 60, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.450, Rules used

$$= \{3809, 3378, 3384, 3379, 3382, 3394, 12, 5556, 5578\}$$

$$\begin{aligned} \int \frac{1}{(c+dx)^2(a+a \coth(e+fx))^3} dx = & -\frac{3f \operatorname{Chi}(6xf + \frac{6cf}{d}) \sinh(6e - \frac{6cf}{d})}{4a^3 d^2} \\ & + \frac{3f \operatorname{Chi}(4xf + \frac{4cf}{d}) \sinh(4e - \frac{4cf}{d})}{2a^3 d^2} \\ & - \frac{3f \operatorname{Chi}(2xf + \frac{2cf}{d}) \sinh(2e - \frac{2cf}{d})}{4a^3 d^2} \\ & + \frac{3f \operatorname{Chi}(2xf + \frac{2cf}{d}) \cosh(2e - \frac{2cf}{d})}{4a^3 d^2} \\ & - \frac{3f \operatorname{Chi}(4xf + \frac{4cf}{d}) \cosh(4e - \frac{4cf}{d})}{2a^3 d^2} \\ & + \frac{3f \operatorname{Chi}(6xf + \frac{6cf}{d}) \cosh(6e - \frac{6cf}{d})}{4a^3 d^2} \\ & + \frac{3f \sinh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\ & - \frac{3f \sinh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\ & + \frac{3f \sinh(6e - \frac{6cf}{d}) \operatorname{Shi}(6xf + \frac{6cf}{d})}{4a^3 d^2} \\ & - \frac{3f \cosh(2e - \frac{2cf}{d}) \operatorname{Shi}(2xf + \frac{2cf}{d})}{4a^3 d^2} \\ & + \frac{3f \cosh(4e - \frac{4cf}{d}) \operatorname{Shi}(4xf + \frac{4cf}{d})}{2a^3 d^2} \\ & - \frac{3f \cosh(6e - \frac{6cf}{d}) \operatorname{Shi}(6xf + \frac{6cf}{d})}{4a^3 d^2} \\ & - \frac{\sinh^3(2e + 2fx)}{8a^3 d(c+dx)} - \frac{3 \sinh^2(2e + 2fx)}{8a^3 d(c+dx)} \\ & - \frac{15 \sinh(2e + 2fx)}{32a^3 d(c+dx)} + \frac{3 \sinh(4e + 4fx)}{8a^3 d(c+dx)} \\ & - \frac{3 \sinh(6e + 6fx)}{32a^3 d(c+dx)} + \frac{\cosh^3(2e + 2fx)}{8a^3 d(c+dx)} \\ & - \frac{3 \cosh^2(2e + 2fx)}{8a^3 d(c+dx)} + \frac{9 \cosh(2e + 2fx)}{32a^3 d(c+dx)} \\ & + \frac{3 \cosh(6e + 6fx)}{32a^3 d(c+dx)} - \frac{1}{8a^3 d(c+dx)} \end{aligned}$$

[In] Int[1/((c + d*x)^2*(a + a*Coth[e + f*x])^3),x]

[Out] $-1/8*1/(a^3*d*(c + d*x)) + (9*Cosh[2*e + 2*f*x])/ (32*a^3*d*(c + d*x)) - (3*Cosh[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) + Cosh[2*e + 2*f*x]^3/(8*a^3*d*(c + d*x)) + (3*Cosh[6*e + 6*f*x])/ (32*a^3*d*(c + d*x)) + (3*f*Cosh[2*e - (2*c + d*x)])/(8*a^3*d*(c + d*x))$

```
*f)/d]*CoshIntegral[(2*c*f)/d + 2*f*x])/(4*a^3*d^2) - (3*f*Cosh[4*e - (4*c*f)/d]*CoshIntegral[(4*c*f)/d + 4*f*x])/(2*a^3*d^2) + (3*f*Cosh[6*e - (6*c*f)/d]*CoshIntegral[(6*c*f)/d + 6*f*x])*Sinh[6*e - (6*c*f)/d])/(4*a^3*d^2) - (3*f*CoshIntegral[(6*c*f)/d + 6*f*x])*Sinh[6*e - (6*c*f)/d])/(4*a^3*d^2) + (3*f*CoshIntegral[(4*c*f)/d + 4*f*x])*Sinh[4*e - (4*c*f)/d])/(2*a^3*d^2) - (3*f*CoshIntegral[(2*c*f)/d + 2*f*x])*Sinh[2*e - (2*c*f)/d])/(4*a^3*d^2) - (15*Sinh[2*e + 2*f*x])/(32*a^3*d*(c + d*x)) - (3*Sinh[2*e + 2*f*x]^2)/(8*a^3*d*(c + d*x)) - Sinh[2*e + 2*f*x]^3/(8*a^3*d*(c + d*x)) + (3*Sinh[4*e + 4*f*x])/(8*a^3*d*(c + d*x)) - (3*Sinh[6*e + 6*f*x])/(32*a^3*d*(c + d*x)) - (3*f*Cosh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(4*a^3*d^2) + (3*f*Sinh[2*e - (2*c*f)/d]*SinhIntegral[(2*c*f)/d + 2*f*x])/(4*a^3*d^2) + (3*f*Cosh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(2*a^3*d^2) - (3*f*Sinh[4*e - (4*c*f)/d]*SinhIntegral[(4*c*f)/d + 4*f*x])/(2*a^3*d^2) - (3*f*Cosh[6*e - (6*c*f)/d]*SinhIntegral[(6*c*f)/d + 6*f*x])/(4*a^3*d^2) + (3*f*Sinh[6*e - (6*c*f)/d]*SinhIntegral[(6*c*f)/d + 6*f*x])/(4*a^3*d^2)
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 3378

```
Int[((c_.) + (d_.)*(x_.))^(m_)*sin[(e_.) + (f_.)*(x_.)], x_Symbol] :> Simp[(c + d*x)^(m + 1)*(Sin[e + f*x]/(d*(m + 1))), x] - Dist[f/(d*(m + 1)), Int[(c + d*x)^(m + 1)*Cos[e + f*x], x], x] /; FreeQ[{c, d, e, f}, x] && LtQ[m, -1]
```

Rule 3379

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[I*(SinhIntegral[c*f*(fz/d) + f*fz*x]/d), x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*e - c*f*fz*I, 0]
```

Rule 3382

```
Int[sin[(e_.) + (Complex[0, fz_])*(f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Simp[CoshIntegral[c*f*(fz/d) + f*fz*x]/d, x] /; FreeQ[{c, d, e, f, fz}, x] && EqQ[d*(e - Pi/2) - c*f*fz*I, 0]
```

Rule 3384

```
Int[sin[(e_.) + (f_.)*(x_.)]/((c_.) + (d_.)*(x_.)), x_Symbol] :> Dist[Cos[(d*e - c*f)/d], Int[Sin[c*(f/d) + f*x]/(c + d*x), x], x] + Dist[Sin[(d*e - c*f)/d], Int[Cos[c*(f/d) + f*x]/(c + d*x), x], x] /; FreeQ[{c, d, e, f}, x] && NeQ[d*e - c*f, 0]
```

Rule 3394

```
Int[((c_.) + (d_.)*(x_))^(m_)*sin[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] :> Si
mp[(c + d*x)^(m + 1)*(Sin[e + f*x]^n/(d*(m + 1))), x] - Dist[f*(n/(d*(m + 1
))), Int[ExpandTrigReduce[(c + d*x)^(m + 1), Cos[e + f*x]*Sin[e + f*x]]^(n -
1), x], x], x] /; FreeQ[{c, d, e, f, m}, x] && IGtQ[n, 1] && GeQ[m, -2] &&
LtQ[m, -1]
```

Rule 3809

```
Int[((c_.) + (d_.)*(x_))^(m_)*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + Cos[2*e + 2*f*x]/(2*a) + Sin[2*e + 2*f*x]/(2*b))]^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x]
&& EqQ[a^2 + b^2, 0] && ILtQ[m, 0] && ILtQ[n, 0]
```

Rule 5556

```
Int[Cosh[(a_.) + (b_.)*(x_)]^(p_)*((c_.) + (d_.)*(x_))^m]*Sinh[(a_.) + (b_.)*(x_)]^n, x_Symbol] :> Int[ExpandTrigReduce[(c + d*x)^m, Sinh[a + b*x]^n*Cosh[a + b*x]^p, x], x] /; FreeQ[{a, b, c, d, m}, x] && IGtQ[n, 0] && IGtQ[p, 0]
```

Rule 5578

```
Int[((e_.) + (f_.)*(x_))^(m_)*Sinh[(a_.) + (b_.)*(x_)]^(p_)*Sinh[(c_.) + (d_.)*(x_)]^q, x_Symbol] :> Int[ExpandTrigReduce[(e + f*x)^m, Sinh[a + b*x]^p*Sinh[c + d*x]^q, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IGtQ[p, 0] && IGtQ[q, 0] && IntegerQ[m]
```

Rubi steps

$$\text{integral} = \int \left(\frac{1}{8a^3(c+dx)^2} - \frac{3 \cosh(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \cosh^2(2e+2fx)}{8a^3(c+dx)^2} - \frac{\cosh^3(2e+2fx)}{8a^3(c+dx)^2} \right. \\ \left. + \frac{3 \sinh(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \cosh^2(2e+2fx) \sinh(2e+2fx)}{8a^3(c+dx)^2} + \frac{3 \sinh^2(2e+2fx)}{8a^3(c+dx)^2} \right. \\ \left. + \frac{\sinh^3(2e+2fx)}{8a^3(c+dx)^2} - \frac{3 \sinh(4e+4fx)}{8a^3(c+dx)^2} - \frac{3 \sinh(2e+2fx) \sinh(4e+4fx)}{16a^3(c+dx)^2} \right) dx$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} - \frac{\int \frac{\cosh^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{\int \frac{\sinh^3(2e+2fx)}{(c+dx)^2} dx}{8a^3} \\
&\quad - \frac{3 \int \frac{\sinh(2e+2fx)\sinh(4e+4fx)}{(c+dx)^2} dx}{16a^3} - \frac{3 \int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{8a^3} \\
&\quad + \frac{3 \int \frac{\cosh^2(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{8a^3} \\
&\quad + \frac{3 \int \frac{\cosh^2(2e+2fx)\sinh(2e+2fx)}{(c+dx)^2} dx}{8a^3} + \frac{3 \int \frac{\sinh^2(2e+2fx)}{(c+dx)^2} dx}{8a^3} - \frac{3 \int \frac{\sinh(4e+4fx)}{(c+dx)^2} dx}{8a^3} \\
&= -\frac{1}{8a^3d(c+dx)} + \frac{3 \cosh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \cosh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3 \sinh(2e+2fx)}{8a^3d(c+dx)} - \frac{3 \sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3 \sinh(4e+4fx)}{8a^3d(c+dx)} - \frac{3 \int \left(-\frac{\cosh(2e+2fx)}{2(c+dx)^2} + \frac{\cosh(6e+6fx)}{2(c+dx)^2} \right) dx}{16a^3} \\
&\quad + \frac{3 \int \left(\frac{\sinh(2e+2fx)}{4(c+dx)^2} + \frac{\sinh(6e+6fx)}{4(c+dx)^2} \right) dx}{8a^3} - \frac{(3if) \int \left(-\frac{i \sinh(2e+2fx)}{4(c+dx)} - \frac{i \sinh(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} \\
&\quad + \frac{(3if) \int -\frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{2a^3d} - \frac{(3if) \int \frac{i \sinh(4e+4fx)}{2(c+dx)} dx}{2a^3d} \\
&\quad + \frac{(3f) \int \frac{\cosh(2e+2fx)}{c+dx} dx}{4a^3d} - \frac{(3f) \int \left(\frac{\cosh(2e+2fx)}{4(c+dx)} - \frac{\cosh(6e+6fx)}{4(c+dx)} \right) dx}{4a^3d} \\
&\quad - \frac{(3f) \int \frac{\sinh(2e+2fx)}{c+dx} dx}{4a^3d} - \frac{(3f) \int \frac{\cosh(4e+4fx)}{c+dx} dx}{2a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} + \frac{3\cosh(2e+2fx)}{8a^3d(c+dx)} - \frac{3\cosh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad - \frac{3\sinh(2e+2fx)}{8a^3d(c+dx)} - \frac{3\sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} + \frac{3\sinh(4e+4fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3 \int \frac{\cosh(2e+2fx)}{(c+dx)^2} dx}{32a^3} - \frac{3 \int \frac{\cosh(6e+6fx)}{(c+dx)^2} dx}{32a^3} + \frac{3 \int \frac{\sinh(2e+2fx)}{(c+dx)^2} dx}{32a^3} \\
&\quad + \frac{3 \int \frac{\sinh(6e+6fx)}{(c+dx)^2} dx}{32a^3} - \frac{(3f) \int \frac{\cosh(2e+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f) \int \frac{\cosh(6e+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f) \int \frac{\sinh(2e+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f) \int \frac{\sinh(6e+6fx)}{c+dx} dx}{16a^3d} + 2 \frac{(3f) \int \frac{\sinh(4e+4fx)}{c+dx} dx}{4a^3d} \\
&\quad - \frac{(3f \cosh(4e - \frac{4cf}{d})) \int \frac{\cosh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} + \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} \\
&\quad - \frac{(3f \cosh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} - \frac{(3f \sinh(4e - \frac{4cf}{d})) \int \frac{\sinh(\frac{4cf}{d} + 4fx)}{c+dx} dx}{2a^3d} \\
&\quad - \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\cosh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d} + \frac{(3f \sinh(2e - \frac{2cf}{d})) \int \frac{\sinh(\frac{2cf}{d} + 2fx)}{c+dx} dx}{4a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} + \frac{9\cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cosh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\cosh(6e+6fx)}{32a^3d(c+dx)} + \frac{3f\cosh(2e-\frac{2cf}{d})\text{Chi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad - \frac{3f\cosh(4e-\frac{4cf}{d})\text{Chi}(\frac{4cf}{d}+4fx)}{2a^3d^2} - \frac{3f\text{Chi}(\frac{2cf}{d}+2fx)\sinh(2e-\frac{2cf}{d})}{4a^3d^2} \\
&\quad - \frac{15\sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\sinh(4e+4fx)}{8a^3d(c+dx)} - \frac{3\sinh(6e+6fx)}{32a^3d(c+dx)} - \frac{3f\cosh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad + \frac{3f\sinh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} - \frac{3f\sinh(4e-\frac{4cf}{d})\text{Shi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad + \frac{(3f)\int \frac{\cosh(2e+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f)\int \frac{\sinh(2e+2fx)}{c+dx} dx}{16a^3d} + \frac{(9f)\int \frac{\cosh(6e+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(9f)\int \frac{\sinh(6e+6fx)}{c+dx} dx}{16a^3d} + \frac{(3f\cosh(6e-\frac{6cf}{d}))\int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f\cosh(6e-\frac{6cf}{d}))\int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} - \frac{(3f\cosh(2e-\frac{2cf}{d}))\int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(3f\cosh(2e-\frac{2cf}{d}))\int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f\sinh(6e-\frac{6cf}{d}))\int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3f\sinh(6e-\frac{6cf}{d}))\int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + 2 \left(\frac{(3f\cosh(4e-\frac{4cf}{d}))\int \frac{\sinh(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} \right. \\
&\quad \quad \left. + \frac{(3f\sinh(4e-\frac{4cf}{d}))\int \frac{\cosh(\frac{4cf}{d}+4fx)}{c+dx} dx}{4a^3d} \right) \\
&\quad - \frac{(3f\sinh(2e-\frac{2cf}{d}))\int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} - \frac{(3f\sinh(2e-\frac{2cf}{d}))\int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} + \frac{9\cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cosh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\cosh(6e+6fx)}{32a^3d(c+dx)} + \frac{9f\cosh(2e-\frac{2cf}{d})\text{Chi}(\frac{2cf}{d}+2fx)}{16a^3d^2} \\
&\quad - \frac{3f\cosh(4e-\frac{4cf}{d})\text{Chi}(\frac{4cf}{d}+4fx)}{2a^3d^2} + \frac{3f\cosh(6e-\frac{6cf}{d})\text{Chi}(\frac{6cf}{d}+6fx)}{16a^3d^2} \\
&\quad - \frac{3f\text{Chi}(\frac{6cf}{d}+6fx)\sinh(6e-\frac{6cf}{d})}{16a^3d^2} - \frac{15f\text{Chi}(\frac{2cf}{d}+2fx)\sinh(2e-\frac{2cf}{d})}{16a^3d^2} \\
&\quad - \frac{15\sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\sinh(4e+4fx)}{8a^3d(c+dx)} - \frac{3\sinh(6e+6fx)}{32a^3d(c+dx)} - \frac{15f\cosh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{16a^3d^2} \\
&\quad + \frac{9f\sinh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{16a^3d^2} - \frac{3f\sinh(4e-\frac{4cf}{d})\text{Shi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad + 2\left(\frac{3f\text{Chi}(\frac{4cf}{d}+4fx)\sinh(4e-\frac{4cf}{d})}{4a^3d^2} + \frac{3f\cosh(4e-\frac{4cf}{d})\text{Shi}(\frac{4cf}{d}+4fx)}{4a^3d^2} \right) \\
&\quad - \frac{3f\cosh(6e-\frac{6cf}{d})\text{Shi}(\frac{6cf}{d}+6fx)}{16a^3d^2} + \frac{3f\sinh(6e-\frac{6cf}{d})\text{Shi}(\frac{6cf}{d}+6fx)}{16a^3d^2} \\
&\quad + \frac{(9f\cosh(6e-\frac{6cf}{d}))\int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} - \frac{(9f\cosh(6e-\frac{6cf}{d}))\int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3f\cosh(2e-\frac{2cf}{d}))\int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f\cosh(2e-\frac{2cf}{d}))\int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} \\
&\quad - \frac{(9f\sinh(6e-\frac{6cf}{d}))\int \frac{\cosh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} + \frac{(9f\sinh(6e-\frac{6cf}{d}))\int \frac{\sinh(\frac{6cf}{d}+6fx)}{c+dx} dx}{16a^3d} \\
&\quad + \frac{(3f\sinh(2e-\frac{2cf}{d}))\int \frac{\cosh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d} + \frac{(3f\sinh(2e-\frac{2cf}{d}))\int \frac{\sinh(\frac{2cf}{d}+2fx)}{c+dx} dx}{16a^3d}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{8a^3d(c+dx)} + \frac{9\cosh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\cosh^2(2e+2fx)}{8a^3d(c+dx)} + \frac{\cosh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\cosh(6e+6fx)}{32a^3d(c+dx)} + \frac{3f\cosh(2e-\frac{2cf}{d})\text{Chi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad - \frac{3f\cosh(4e-\frac{4cf}{d})\text{Chi}(\frac{4cf}{d}+4fx)}{2a^3d^2} + \frac{3f\cosh(6e-\frac{6cf}{d})\text{Chi}(\frac{6cf}{d}+6fx)}{4a^3d^2} \\
&\quad - \frac{3f\text{Chi}(\frac{6cf}{d}+6fx)\sinh(6e-\frac{6cf}{d})}{4a^3d^2} - \frac{3f\text{Chi}(\frac{2cf}{d}+2fx)\sinh(2e-\frac{2cf}{d})}{4a^3d^2} \\
&\quad - \frac{15\sinh(2e+2fx)}{32a^3d(c+dx)} - \frac{3\sinh^2(2e+2fx)}{8a^3d(c+dx)} - \frac{\sinh^3(2e+2fx)}{8a^3d(c+dx)} \\
&\quad + \frac{3\sinh(4e+4fx)}{8a^3d(c+dx)} - \frac{3\sinh(6e+6fx)}{32a^3d(c+dx)} - \frac{3f\cosh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} \\
&\quad + \frac{3f\sinh(2e-\frac{2cf}{d})\text{Shi}(\frac{2cf}{d}+2fx)}{4a^3d^2} - \frac{3f\sinh(4e-\frac{4cf}{d})\text{Shi}(\frac{4cf}{d}+4fx)}{2a^3d^2} \\
&\quad + 2\left(\frac{3f\text{Chi}(\frac{4cf}{d}+4fx)\sinh(4e-\frac{4cf}{d})}{4a^3d^2} + \frac{3f\cosh(4e-\frac{4cf}{d})\text{Shi}(\frac{4cf}{d}+4fx)}{4a^3d^2}\right) \\
&\quad - \frac{3f\cosh(6e-\frac{6cf}{d})\text{Shi}(\frac{6cf}{d}+6fx)}{4a^3d^2} + \frac{3f\sinh(6e-\frac{6cf}{d})\text{Shi}(\frac{6cf}{d}+6fx)}{4a^3d^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.77 (sec), antiderivative size = 796, normalized size of antiderivative = 1.15

$$\begin{aligned}
&\int \frac{1}{(c+dx)^2(a+a\coth(e+fx))^3} dx \\
&= \frac{\text{csch}^3(e+fx)(\cosh(\frac{3cf}{d})+\sinh(\frac{3cf}{d}))\left(3d\cosh(e+f(-\frac{3c}{d}+x))-d\cosh(3(e+f(-\frac{c}{d}+x)))+d\cosh\right. \\
&\quad \left.\left(3(e+f(-\frac{c}{d}+x))\right)\right)}{(c+dx)^2(a+a\coth(e+fx))^3}
\end{aligned}$$

```
[In] Integrate[1/((c + d*x)^2*(a + a*Coth[e + f*x])^3), x]
[Out] (Csch[e + f*x]^3*(Cosh[(3*c*f)/d] + Sinh[(3*c*f)/d])*(3*d*Cosh[e + f*(-3*c)/d + x]) - d*Cosh[3*(e + f*(-(c/d) + x))] + d*Cosh[3*(e + f*(c/d + x))] - 3*d*Cosh[e + f*((3*c)/d + x)] + 6*c*f*Cosh[3*e - (3*f*(c + d*x))/d]*CoshIntegral[(6*f*(c + d*x))/d] + 6*d*f*x*Cosh[3*e - (3*f*(c + d*x))/d]*CoshIntegral[(6*f*(c + d*x))/d] + 6*f*(c + d*x)*CoshIntegral[(2*f*(c + d*x))/d]*(Cosh[e - (c*f)/d + 3*f*x] + Sinh[e - (c*f)/d + 3*f*x]) + 3*d*Sinh[e + f*(-3*c)/d + x] - d*Sinh[3*(e + f*(-(c/d) + x))] - d*Sinh[3*(e + f*(c/d + x))] + 3*d*Sinh[e + f*((3*c)/d + x)] - 6*c*f*CoshIntegral[(6*f*(c + d*x))/d]*Sinh[3*e - (3*f*(c + d*x))/d] - 6*d*f*x*CoshIntegral[(6*f*(c + d*x))/d]*Sinh[3*e - (3*f*(c + d*x))/d] + 12*f*(c + d*x)*CoshIntegral[(4*f*(c + d*x))/d]*(-Cos[h[e - (f*(c + 3*d*x))/d] + Sinh[e - (f*(c + 3*d*x))/d]) - 6*c*f*Cosh[e - (c*f)/d + 3*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 6*d*f*x*Cosh[e - (c*f)/d + 3*f*x]*SinhIntegral[(2*f*(c + d*x))/d]
```

$$\begin{aligned}
& 3*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 6*c*f*Sinh[e - (c*f)/d + 3*f*x]*SinhIntegral[(2*f*(c + d*x))/d] - 6*d*f*x*Sinh[e - (c*f)/d + 3*f*x]*SinhIntegral[(2*f*(c + d*x))/d] + 12*c*f*Cosh[e - (f*(c + 3*d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] + 12*d*f*x*Cosh[e - (f*(c + 3*d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] - 12*c*f*Sinh[e - (f*(c + 3*d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] - 12*d*f*x*Sinh[e - (f*(c + 3*d*x))/d]*SinhIntegral[(4*f*(c + d*x))/d] - 6*c*f*Cosh[3*e - (3*f*(c + d*x))/d]*SinhIntegral[(6*f*(c + d*x))/d] - 6*d*f*x*Cosh[3*e - (3*f*(c + d*x))/d]*SinhIntegral[(6*f*(c + d*x))/d] + 6*c*f*Sinh[3*e - (3*f*(c + d*x))/d]*SinhIntegral[(6*f*(c + d*x))/d] + 6*d*f*x*Sinh[3*e - (3*f*(c + d*x))/d]*SinhIntegral[(6*f*(c + d*x))/d])/(8*a^3*d^2*(c + d*x)*(1 + Coth[e + f*x])^3)
\end{aligned}$$

Maple [A] (verified)

Time = 0.47 (sec) , antiderivative size = 239, normalized size of antiderivative = 0.35

method	result
risch	$ -\frac{1}{8a^3d(dx+c)} + \frac{fe^{-6fx-6e}}{8a^3d(dxf+cf)} - \frac{3fe^{\frac{6cf-6de}{d}}Ei_1\left(6fx+6e+\frac{6cf-6de}{d}\right)}{4a^3d^2} - \frac{3fe^{-4fx-4e}}{8a^3d(dxf+cf)} + \frac{3fe^{\frac{4cf-4de}{d}}Ei_1\left(4fx+4e+\frac{4cf-4de}{d}\right)}{2a^3d^2} $

[In] `int(1/(d*x+c)^2/(a+a*coth(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned}
& -1/8/a^3/d/(d*x+c)+1/8*f/a^3*exp(-6*f*x-6*e)/d/(d*f*x+c*f)-3/4*f/a^3/d^2*exp(6*(c*f-d*e)/d)*Ei(1,6*f*x+6*e+6*(c*f-d*e)/d)-3/8*f/a^3*exp(-4*f*x-4*e)/d/(d*f*x+c*f)+3/2*f/a^3/d^2*exp(4*(c*f-d*e)/d)*Ei(1,4*f*x+4*e+4*(c*f-d*e)/d)+3/8*f/a^3*exp(-2*f*x-2*e)/d/(d*f*x+c*f)-3/4*f/a^3/d^2*exp(2*(c*f-d*e)/d)*Ei(1,2*f*x+2*e+2*(c*f-d*e)/d)
\end{aligned}$$

Fricas [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 1162, normalized size of antiderivative = 1.68

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^3} dx = \text{Too large to display}$$

[In] `integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned}
& 1/4*(3*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(f*x + e)^3*sinh(-2*(d*e - c*f)/d) - 6*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(f*x + e)^3*sinh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(f*x + e)^3*sinh(-6*(d*e - c*f)/d) + 3*((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f)/d))*cosh(f*x + e)^3 + (3*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) - 6*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f)/d))*cosh(f*x + e)^3)
\end{aligned}$$

$$\begin{aligned}
& x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei \\
& (-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-2*(d*f*x \\
& + c*f)/d)*sinh(-2*(d*e - c*f)/d) - 6*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*s \\
& inh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*sinh(-6*(d*e \\
& - c*f)/d) - d)*sinh(f*x + e)^3 + 9*((d*f*x + c*f)*Ei(-2*(d*f*x + c*f)/d)*c \\
& osh(f*x + e)*sinh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d) \\
&)*cosh(f*x + e)*sinh(-4*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/ \\
& d)*cosh(f*x + e)*sinh(-6*(d*e - c*f)/d) + ((d*f*x + c*f)*Ei(-2*(d*f*x + c*f) \\
& /d)*cosh(-2*(d*e - c*f)/d) - 2*(d*f*x + c*f)*Ei(-4*(d*f*x + c*f)/d)*cosh(- \\
& 4*(d*e - c*f)/d) + (d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*cosh(-6*(d*e - c*f) \\
& /d))*cosh(f*x + e)*sinh(f*x + e)^2 + 3*(3*(d*f*x + c*f)*Ei(-2*(d*f*x + c*f) \\
& /d)*cosh(f*x + e)^2*sinh(-2*(d*e - c*f)/d) - 6*(d*f*x + c*f)*Ei(-4*(d*f*x \\
& + c*f)/d)*cosh(f*x + e)^2*sinh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d \\
& *f*x + c*f)/d)*cosh(f*x + e)^2*sinh(-6*(d*e - c*f)/d) + (3*(d*f*x + c*f)*Ei \\
& (-2*(d*f*x + c*f)/d)*cosh(-2*(d*e - c*f)/d) - 6*(d*f*x + c*f)*Ei(-4*(d*f*x \\
& + c*f)/d)*cosh(-4*(d*e - c*f)/d) + 3*(d*f*x + c*f)*Ei(-6*(d*f*x + c*f)/d)*c \\
& osh(-6*(d*e - c*f)/d) - d)*cosh(f*x + e)^2 + d)*sinh(f*x + e))/((a^3*d^3*x \\
& + a^3*c*d^2)*cosh(f*x + e)^3 + 3*(a^3*d^3*x + a^3*c*d^2)*cosh(f*x + e)^2*si \\
& nh(f*x + e) + 3*(a^3*d^3*x + a^3*c*d^2)*cosh(f*x + e)*sinh(f*x + e)^2 + (a^ \\
& 3*d^3*x + a^3*c*d^2)*sinh(f*x + e)^3)
\end{aligned}$$

Sympy [F]

$$\begin{aligned}
& \int \frac{1}{(c+dx)^2(a+a\coth(e+fx))^3} dx \\
= & \int \frac{1}{c^2\coth^3(e+fx)+3c^2\coth^2(e+fx)+3c^2\coth(e+fx)+c^2+2cdx\coth^3(e+fx)+6cdx\coth^2(e+fx)+6cdx\coth(e+fx)+2cdx+d^2x^2\coth^3(e+fx)} dx
\end{aligned}$$

[In] integrate(1/(d*x+c)**2/(a+a*coth(f*x+e))**3,x)

[Out] Integral(1/(c**2*coth(e + f*x)**3 + 3*c**2*coth(e + f*x)**2 + 3*c**2*coth(e + f*x) + c**2 + 2*c*d*x*coth(e + f*x)**3 + 6*c*d*x*coth(e + f*x)**2 + 6*c*d*x*coth(e + f*x) + 2*c*d*x + d**2*x**2*coth(e + f*x)**3 + 3*d**2*x**2*coth(e + f*x)**2 + 3*d**2*x**2*coth(e + f*x) + d**2*x**2), x)/a**3

Maxima [A] (verification not implemented)

none

Time = 5.16 (sec) , antiderivative size = 140, normalized size of antiderivative = 0.20

$$\int \frac{1}{(c+dx)^2(a+a\coth(e+fx))^3} dx = -\frac{1}{8(a^3d^2x+a^3cd)} + \frac{e^{(-6e+\frac{6cf}{d})}E_2\left(\frac{6(dx+c)f}{d}\right)}{8(dx+c)a^3d} \\ - \frac{3e^{(-4e+\frac{4cf}{d})}E_2\left(\frac{4(dx+c)f}{d}\right)}{8(dx+c)a^3d} \\ + \frac{3e^{(-2e+\frac{2cf}{d})}E_2\left(\frac{2(dx+c)f}{d}\right)}{8(dx+c)a^3d}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="maxima")
[Out] -1/8/(a^3*d^2*x + a^3*c*d) + 1/8*e^(-6*e + 6*c*f/d)*exp_integral_e(2, 6*(d*x + c)*f/d)/((d*x + c)*a^3*d) - 3/8*e^(-4*e + 4*c*f/d)*exp_integral_e(2, 4*(d*x + c)*f/d)/((d*x + c)*a^3*d) + 3/8*e^(-2*e + 2*c*f/d)*exp_integral_e(2, 2*(d*x + c)*f/d)/((d*x + c)*a^3*d)
```

Giac [A] (verification not implemented)

none

Time = 0.32 (sec) , antiderivative size = 841, normalized size of antiderivative = 1.22

$$\int \frac{1}{(c+dx)^2(a+a\coth(e+fx))^3} dx \\ = \frac{6(dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right)f^2\text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right)e^{-\frac{2(de-cf)}{d}} - 6def^2\text{Ei}\left(-\frac{2\left((dx+c)\left(\frac{de}{dx+c} - \frac{cf}{dx+c} + f\right) - de + cf\right)}{d}\right)e^{-\frac{2(de-cf)}{d}}}{(c+dx)^2(a+a\coth(e+fx))^3}$$

```
[In] integrate(1/(d*x+c)^2/(a+a*coth(f*x+e))^3,x, algorithm="giac")
[Out] 1/8*(6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*((d*e - c*f)/d)) - 6*d*e*f^2*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*((d*e - c*f)/d)) + 6*c*f^3*Ei(-2*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-2*((d*e - c*f)/d)) - 12*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*((d*e - c*f)/d)) + 12*d*e*f^2*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^(-4*((d*e - c*f)/d)) - 1
```

$$\begin{aligned}
& 2*c*f^3*Ei(-4*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^{-4*(d*e - c*f)/d} + 6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)*f^2*Ei(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^{-6*(d*e - c*f)/d} - 6*d*e*f^2*Ei(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^{-6*(d*e - c*f)/d} + 6*c*f^3*Ei(-6*((d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f) - d*e + c*f)/d)*e^{-6*(d*e - c*f)/d} + 3*d*f^2*e^{-2*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d} - 3*d*f^2*e^{-4*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d} + d*f^2*e^{-6*(d*x + c)*(d*e/(d*x + c) - c*f/(d*x + c) + f)/d} - d*f^2*2*d^2/(((d*x + c)*a^3*d^4*(d*e/(d*x + c) - c*f/(d*x + c) + f) - a^3*d^5*e + a^3*c*d^4*f)*f)
\end{aligned}$$

Mupad [F(-1)]

Timed out.

$$\int \frac{1}{(c + dx)^2(a + a \coth(e + fx))^3} dx = \int \frac{1}{(a + a \coth(e + fx))^3 (c + dx)^2} dx$$

[In] `int(1/((a + a*coth(e + fx))^3*(c + d*x)^2),x)`
[Out] `int(1/((a + a*coth(e + fx))^3*(c + d*x)^2), x)`

3.32 $\int (c + dx)^m (a + a \coth(e + fx))^2 dx$

Optimal result	227
Rubi [N/A]	227
Mathematica [N/A]	228
Maple [N/A] (verified)	228
Fricas [N/A]	228
Sympy [N/A]	228
Maxima [N/A]	229
Giac [N/A]	229
Mupad [N/A]	229

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \text{Int}((c + dx)^m (a + a \coth(e + fx))^2, x)$$

[Out] Unintegrable((d*x+c)^m*(a+a*coth(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.03 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (c + dx)^m (a + a \coth(e + fx))^2 dx$$

[In] Int[(c + d*x)^m*(a + a*Coth[e + f*x])^2,x]

[Out] Defer[Int][(c + d*x)^m*(a + a*Coth[e + f*x])^2, x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + a \coth(e + fx))^2 dx$$

Mathematica [N/A]

Not integrable

Time = 41.61 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (c + dx)^m (a + a \coth(e + fx))^2 dx$$

[In] `Integrate[(c + d*x)^m*(a + a*Coth[e + f*x])^2, x]`

[Out] `Integrate[(c + d*x)^m*(a + a*Coth[e + f*x])^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int (dx + c)^m (a + a \coth(fx + e))^2 dx$$

[In] `int((d*x+c)^m*(a+a*coth(f*x+e))^2, x)`

[Out] `int((d*x+c)^m*(a+a*coth(f*x+e))^2, x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 37, normalized size of antiderivative = 1.85

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (a \coth(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e))^2, x, algorithm="fricas")`

[Out] `integral(a^2*coth(f*x + e)^2 + 2*a^2*coth(f*x + e) + a^2)*(d*x + c)^m, x)`

Sympy [N/A]

Not integrable

Time = 4.42 (sec) , antiderivative size = 44, normalized size of antiderivative = 2.20

$$\begin{aligned} \int (c + dx)^m (a + a \coth(e + fx))^2 dx &= a^2 \left(\int 2(c + dx)^m \coth(e + fx) dx \right. \\ &\quad \left. + \int (c + dx)^m \coth^2(e + fx) dx + \int (c + dx)^m dx \right) \end{aligned}$$

[In] `integrate((d*x+c)**m*(a+a*coth(f*x+e))**2, x)`

[Out] `a**2*(Integral(2*(c + d*x)**m*coth(e + f*x), x) + Integral((c + d*x)**m*cot h(e + f*x)**2, x) + Integral((c + d*x)**m, x))`

Maxima [N/A]

Not integrable

Time = 0.39 (sec) , antiderivative size = 122, normalized size of antiderivative = 6.10

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (a \coth(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e))^2,x, algorithm="maxima")`

[Out] $(d*x + c)^{m+1} * a^2 / (d*(m+1)) + \text{integrate}((d*x + c)^m * a^2 * (e^{(f*x + e)} + e^{(-f*x - e)})^2 / (e^{(f*x + e)} - e^{(-f*x - e)})^2 + 2 * (d*x + c)^m * a^2 * (e^{(f*x + e)} + e^{(-f*x - e)}) / (e^{(f*x + e)} - e^{(-f*x - e)}), x)$

Giac [N/A]

Not integrable

Time = 0.33 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (a \coth(fx + e) + a)^2 (dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((a*coth(f*x + e) + a)^2 * (d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 2.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int (c + dx)^m (a + a \coth(e + fx))^2 dx = \int (a + a \coth(e + f x))^2 (c + d x)^m dx$$

[In] `int((a + a*coth(e + f*x))^2*(c + d*x)^m,x)`

[Out] `int((a + a*coth(e + f*x))^2*(c + d*x)^m, x)`

3.33 $\int (c + dx)^m (a + a \coth(e + fx)) dx$

Optimal result	230
Rubi [N/A]	230
Mathematica [N/A]	231
Maple [N/A] (verified)	231
Fricas [N/A]	231
Sympy [N/A]	231
Maxima [N/A]	232
Giac [N/A]	232
Mupad [N/A]	232

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \text{Int}((c + dx)^m (a + a \coth(e + fx)), x)$$

[Out] Unintegrable((d*x+c)^m*(a+a*coth(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (c + dx)^m (a + a \coth(e + fx)) dx$$

[In] Int[(c + d*x)^m*(a + a*Coth[e + f*x]),x]

[Out] Defer[Int][(c + d*x)^m*(a + a*Coth[e + f*x]), x]

Rubi steps

$$\text{integral} = \int (c + dx)^m (a + a \coth(e + fx)) dx$$

Mathematica [N/A]

Not integrable

Time = 20.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (c + dx)^m (a + a \coth(e + fx)) dx$$

[In] `Integrate[(c + d*x)^m*(a + a*Coth[e + f*x]),x]`

[Out] `Integrate[(c + d*x)^m*(a + a*Coth[e + f*x]), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int (dx + c)^m (a + a \coth(fx + e)) dx$$

[In] `int((d*x+c)^m*(a+a*coth(f*x+e)),x)`

[Out] `int((d*x+c)^m*(a+a*coth(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (a \coth(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e)),x, algorithm="fricas")`

[Out] `integral((a*coth(f*x + e) + a)*(d*x + c)^m, x)`

Sympy [N/A]

Not integrable

Time = 2.78 (sec) , antiderivative size = 24, normalized size of antiderivative = 1.33

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = a \left(\int (c + dx)^m \coth(e + fx) dx + \int (c + dx)^m dx \right)$$

[In] `integrate((d*x+c)**m*(a+a*coth(f*x+e)),x)`

[Out] `a*(Integral((c + d*x)**m*coth(e + f*x), x) + Integral((c + d*x)**m, x))`

Maxima [N/A]

Not integrable

Time = 0.29 (sec) , antiderivative size = 68, normalized size of antiderivative = 3.78

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (a \coth(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e)),x, algorithm="maxima")`

[Out] `a*integrate((d*x + c)^m*(e^(f*x + e) + e^(-f*x - e))/(e^(f*x + e) - e^(-f*x - e)), x) + (d*x + c)^(m + 1)*a/(d*(m + 1))`

Giac [N/A]

Not integrable

Time = 0.30 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (a \coth(fx + e) + a)(dx + c)^m dx$$

[In] `integrate((d*x+c)^m*(a+a*coth(f*x+e)),x, algorithm="giac")`

[Out] `integrate((a*coth(f*x + e) + a)*(d*x + c)^m, x)`

Mupad [N/A]

Not integrable

Time = 2.15 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int (c + dx)^m (a + a \coth(e + fx)) dx = \int (a + a \coth(e + f x)) (c + d x)^m dx$$

[In] `int((a + a*coth(e + f*x))*(c + d*x)^m,x)`

[Out] `int((a + a*coth(e + f*x))*(c + d*x)^m, x)`

3.34 $\int \frac{(c+dx)^m}{a+a \coth(e+fx)} dx$

Optimal result	233
Rubi [A] (verified)	233
Mathematica [A] (verified)	234
Maple [F]	235
Fricas [A] (verification not implemented)	235
Sympy [F]	235
Maxima [F]	236
Giac [F]	236
Mupad [F(-1)]	236

Optimal result

Integrand size = 20, antiderivative size = 88

$$\int \frac{(c+dx)^m}{a+a \coth(e+fx)} dx = \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{2^{-2-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{af}$$

[Out] $1/2*(d*x+c)^(1+m)/a/d/(1+m)+2^{-2-m}*exp(-2*e+2*c*f/d)*(d*x+c)^m*\text{GAMMA}(1+m, 2*f*(d*x+c)/d)/a/f/((f*(d*x+c)/d)^m)$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {3808, 2212}

$$\int \frac{(c+dx)^m}{a+a \coth(e+fx)} dx = \frac{2^{-m-2} e^{\frac{2cf}{d}-2e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2f(c+dx)}{d}\right)}{af} + \frac{(c+dx)^{m+1}}{2ad(m+1)}$$

[In] $\text{Int}[(c+d*x)^m/(a+a*\text{Coth}[e+f*x]),x]$

[Out] $(c+d*x)^(1+m)/(2*a*d*(1+m)) + (2^{-2-m})*E^{-2*e+(2*c*f)/d}*(c+d*x)^m*\text{Gamma}[1+m, (2*f*(c+d*x))/d]/(a*f*((f*(c+d*x))/d)^m)$

Rule 2212

```

Int[(F_ )^((g_.)*(e_.) + (f_.)*(x_.))*((c_.) + (d_.)*(x_.))^m_, x_Symbol]
:> Simp[(-F^g*(e - c*(f/d)))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d)
)^IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d)^FracPart[m]))*Gamma[m + 1,
((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]

```

Rule 3808

```

Int[((c_.) + (d_.)*(x_.))^m/((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)]), x_Sym
bol] :> Simp[(c + d*x)^(m + 1)/(2*a*d*(m + 1)), x] + Dist[1/(2*a), Int[(c +
d*x)^m*E^(2*(a/b)*(e + f*x)), x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] &&
EqQ[a^2 + b^2, 0] && !IntegerQ[m]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{\int e^{2i(i e + \frac{\pi}{2} + ifx)} (c+dx)^m dx}{2a} \\
&= \frac{(c+dx)^{1+m}}{2ad(1+m)} + \frac{2^{-2-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{af}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.20 (sec) , antiderivative size = 115, normalized size of antiderivative = 1.31

$$\begin{aligned}
&\int \frac{(c+dx)^m}{a + a \coth(e + fx)} dx \\
&= \frac{\operatorname{csch}(e + fx) \left(\frac{e^e f (c+dx)^{1+m}}{d(1+m)} + 2^{-1-m} e^{-e+\frac{2cf}{d}} (c+dx)^m \left(\frac{cf}{d} + fx\right)^{-m} \Gamma\left(1+m, 2\left(\frac{cf}{d} + fx\right)\right) (\cosh(fx) + \sinh(fx)) \right)}{2f(a + a \coth(e + fx))}
\end{aligned}$$

```

[In] Integrate[(c + d*x)^m/(a + a*Coth[e + f*x]), x]
[Out] (Csch[e + f*x]*((E^e*f*(c + d*x)^(1 + m))/(d*(1 + m)) + (2^(-1 - m)*E^(-e +
(2*c*f)/d)*(c + d*x)^m*Gamma[1 + m, 2*((c*f)/d + f*x)]))/((c*f)/d + f*x)^m)
*(Cosh[f*x] + Sinh[f*x]))/(2*f*(a + a*Coth[e + f*x]))

```

Maple [F]

$$\int \frac{(dx + c)^m}{a + a \coth(fx + e)} dx$$

[In] `int((d*x+c)^m/(a+a*coth(f*x+e)),x)`

[Out] `int((d*x+c)^m/(a+a*coth(f*x+e)),x)`

Fricas [A] (verification not implemented)

none

Time = 0.08 (sec) , antiderivative size = 148, normalized size of antiderivative = 1.68

$$\begin{aligned} & \int \frac{(c + dx)^m}{a + a \coth(e + fx)} dx \\ &= \frac{(dm + d) \cosh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) \Gamma\left(m + 1, \frac{2(df + cf)}{d}\right) - (dm + d) \Gamma\left(m + 1, \frac{2(df + cf)}{d}\right) \sinh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right)}{4(adfm + adf)} \end{aligned}$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e)),x, algorithm="fricas")`

[Out] `1/4*((d*m + d)*cosh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d)*gamma(m + 1, 2*(d*f*x + c*f)/d) - (d*m + d)*gamma(m + 1, 2*(d*f*x + c*f)/d)*sinh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d) + 2*(d*f*x + c*f)*cosh(m*log(d*x + c)) + 2*(d*f*x + c*f)*sinh(m*log(d*x + c)))/(a*d*f*m + a*d*f)`

Sympy [F]

$$\int \frac{(c + dx)^m}{a + a \coth(e + fx)} dx = \frac{\int \frac{(c+dx)^m}{\coth(e+fx)+1} dx}{a}$$

[In] `integrate((d*x+c)**m/(a+a*coth(f*x+e)),x)`

[Out] `Integral((c + d*x)**m/(coth(e + f*x) + 1), x)/a`

Maxima [F]

$$\int \frac{(c + dx)^m}{a + a \coth(e + fx)} dx = \int \frac{(dx + c)^m}{a \coth(fx + e) + a} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e)),x, algorithm="maxima")`
[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a), x)`

Giac [F]

$$\int \frac{(c + dx)^m}{a + a \coth(e + fx)} dx = \int \frac{(dx + c)^m}{a \coth(fx + e) + a} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e)),x, algorithm="giac")`
[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{a + a \coth(e + fx)} dx = \int \frac{(c + d x)^m}{a + a \coth(e + f x)} dx$$

[In] `int((c + d*x)^m/(a + a*coth(e + f*x)),x)`
[Out] `int((c + d*x)^m/(a + a*coth(e + f*x)), x)`

3.35 $\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^2} dx$

Optimal result	237
Rubi [A] (verified)	237
Mathematica [A] (verified)	238
Maple [F]	239
Fricas [A] (verification not implemented)	239
Sympy [F]	239
Maxima [F]	240
Giac [F]	240
Mupad [F(-1)]	240

Optimal result

Integrand size = 20, antiderivative size = 152

$$\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^2} dx = \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{2^{-2-m}e^{-2e+\frac{2cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^2f} - \frac{4^{-2-m}e^{-4e+\frac{4cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^2f}$$

[Out] $1/4*(d*x+c)^(1+m)/a^2/d/(1+m)+2^(-2-m)*exp(-2*e+2*c*f/d)*(d*x+c)^m*\text{GAMMA}(1+m, 2*f*(d*x+c)/d)/a^2/f/((f*(d*x+c)/d)^m)-4^(-2-m)*exp(-4*e+4*c*f/d)*(d*x+c)^m*\text{GAMMA}(1+m, 4*f*(d*x+c)/d)/a^2/f/((f*(d*x+c)/d)^m)$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 152, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.100, Rules used = {3810, 2212}

$$\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^2} dx = \frac{2^{-m-2}e^{\frac{2cf}{d}-2e}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{2f(c+dx)}{d}\right)}{a^2f} - \frac{4^{-m-2}e^{\frac{4cf}{d}-4e}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(m+1, \frac{4f(c+dx)}{d}\right)}{a^2f} + \frac{(c+dx)^{m+1}}{4a^2d(m+1)}$$

[In] $\text{Int}[(c + d*x)^m / (a + a*\text{Coth}[e + f*x])^2, x]$

[Out] $(c + d*x)^{1+m} / (4*a^2*d*(1+m)) + (2^{-2-m} * E^{-(-2-e + (2*c*f)/d)} * (c + d*x)^m * \text{Gamma}[1+m, (2*f*(c + d*x))/d]) / (a^2*f*((f*(c + d*x))/d)^m) - (4^{-2-m} * E^{(-4-e + (4*c*f)/d)} * (c + d*x)^m * \text{Gamma}[1+m, (4*f*(c + d*x))/d]) / (a^2*f*((f*(c + d*x))/d)^m)$

Rule 2212

$\text{Int}[(F_.)^((g_.) * ((e_.) + (f_.) * (x_))) * ((c_.) + (d_.) * (x_))^{(m_)}, x_{\text{Symbol}}]$
 $:> \text{Simp}[(-F^((g*(e - c*(f/d)))) * ((c + d*x)^{\text{FracPart}[m]} / (d*(-f)*g*(\text{Log}[F]/d))^{\text{IntPart}[m] + 1}) * ((-f)*g*\text{Log}[F]*((c + d*x)/d))^{\text{FracPart}[m]}) * \text{Gamma}[m + 1, ((-f)*g*(\text{Log}[F]/d)) * (c + d*x)], x] /; \text{FreeQ}[\{F, c, d, e, f, g, m\}, x] \&& \text{!IntegerQ}[m]$

Rule 3810

$\text{Int}[((c_.) + (d_.) * (x_))^{(m_)} * ((a_.) + (b_.) * \tan[(e_.) + (f_.) * (x_)])^{(n_)}, x_{\text{Symbol}}]$
 $:> \text{Int}[\text{ExpandIntegrand}[(c + d*x)^m, (1/(2*a) + E^{(2*(a/b)*(e + f*x))/(2*a)})^{-n}], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m\}, x] \&& \text{EqQ}[a^2 + b^2, 0] \&& \text{ILtQ}[n, 0]$

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^m}{4a^2} + \frac{e^{-4e-4fx}(c+dx)^m}{4a^2} - \frac{e^{-2e-2fx}(c+dx)^m}{2a^2} \right) dx \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{\int e^{-4e-4fx}(c+dx)^m dx}{4a^2} - \frac{\int e^{-2e-2fx}(c+dx)^m dx}{2a^2} \\ &= \frac{(c+dx)^{1+m}}{4a^2d(1+m)} + \frac{2^{-2-m}e^{-2e+\frac{2cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^2f} \\ &\quad - \frac{4^{-2-m}e^{-4e+\frac{4cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^2f} \end{aligned}$$

Mathematica [A] (verified)

Time = 1.73 (sec), antiderivative size = 162, normalized size of antiderivative = 1.07

$$\begin{aligned} &\int \frac{(c+dx)^m}{(a + a \coth(e + fx))^2} dx \\ &= \frac{(c+dx)^m \text{csch}^2(e + fx) \left(\frac{4e^{2e}f(c+dx)}{d(1+m)} + 2^{2-m}e^{\frac{2cf}{d}} \left(f\left(\frac{c}{d} + x\right)\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right) - 4^{-m}e^{-2e+\frac{4cf}{d}} \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right) \right)}{16a^2f(1 + \coth(e + fx))^2} \end{aligned}$$

[In] `Integrate[(c + d*x)^m/(a + a*Coth[e + f*x])^2, x]`

[Out]
$$\frac{((c + d*x)^m * \text{Csch}[e + f*x]^2 * ((4*E^(2*e)*f*(c + d*x))/(d*(1 + m)) + (2^(2 - m)*E^((2*c*f)/d)*\text{Gamma}[1 + m, (2*f*(c + d*x))/d])/(f*(c/d + x))^m - (E^(-2*m) * (4*f*(c + d*x))/d)) / (4^m * ((f*(c + d*x))/d)^m) * (\text{Cosh}[f*x] + \text{Sinh}[f*x])^2) / (16*a^2*f*(1 + \text{Coth}[e + f*x])^2)}$$

Maple [F]

$$\int \frac{(dx + c)^m}{(a + a \coth(fx + e))^2} dx$$

[In] `int((d*x+c)^m/(a+a*coth(f*x+e))^2,x)`

[Out] `int((d*x+c)^m/(a+a*coth(f*x+e))^2,x)`

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec), antiderivative size = 248, normalized size of antiderivative = 1.63

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^2} dx = \frac{(dm + d) \cosh\left(\frac{dm \log\left(\frac{4f}{d}\right) + 4de - 4cf}{d}\right) \Gamma\left(m + 1, \frac{4(df + cf)}{d}\right) - 4(dm + d) \cosh\left(\frac{dm \log\left(\frac{2f}{d}\right) + 2de - 2cf}{d}\right) \Gamma\left(m + 1, \frac{2(df + cf)}{d}\right)}{a^2}$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^2,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -1/16*((d*m + d)*\cosh((d*m*\log(4*f/d) + 4*d*e - 4*c*f)/d)*\text{gamma}(m + 1, 4*(d*f*x + c*f)/d) - 4*(d*m + d)*\cosh((d*m*\log(2*f/d) + 2*d*e - 2*c*f)/d)*\text{gamma}(m + 1, 2*(d*f*x + c*f)/d) - (d*m + d)*\text{gamma}(m + 1, 4*(d*f*x + c*f)/d)*\sinh((d*m*\log(4*f/d) + 4*d*e - 4*c*f)/d) + 4*(d*m + d)*\text{gamma}(m + 1, 2*(d*f*x + c*f)/d)*\sinh((d*m*\log(2*f/d) + 2*d*e - 2*c*f)/d) - 4*(d*f*x + c*f)*\cosh(m*\log(d*x + c)) - 4*(d*f*x + c*f)*\sinh(m*\log(d*x + c))) / (a^2*d*f*m + a^2*d*f) \end{aligned}$$

Sympy [F]

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^2} dx = \frac{\int \frac{(c + dx)^m}{\coth^2(e + fx) + 2 \coth(e + fx) + 1} dx}{a^2}$$

[In] `integrate((d*x+c)**m/(a+a*coth(f*x+e))**2,x)`

[Out] `Integral((c + d*x)**m / (coth(e + f*x)**2 + 2*coth(e + f*x) + 1), x) / a**2`

Maxima [F]

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^2} dx = \int \frac{(dx + c)^m}{(a \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^2,x, algorithm="maxima")`
[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a)^2, x)`

Giac [F]

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^2} dx = \int \frac{(dx + c)^m}{(a \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^2} dx = \int \frac{(c + dx)^m}{(a + a \coth(e + f x))^2} dx$$

[In] `int((c + d*x)^m/(a + a*coth(e + f*x))^2,x)`
[Out] `int((c + d*x)^m/(a + a*coth(e + f*x))^2, x)`

3.36 $\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^3} dx$

Optimal result	241
Rubi [A] (verified)	241
Mathematica [A] (verified)	243
Maple [F]	243
Fricas [A] (verification not implemented)	244
Sympy [F]	244
Maxima [F]	244
Giac [F]	245
Mupad [F(-1)]	245

Optimal result

Integrand size = 20, antiderivative size = 223

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+a \coth(e+fx))^3} dx \\ &= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3 \cdot 2^{-4-m} e^{-2e+\frac{2cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^3f} \\ &\quad - \frac{3 \cdot 2^{-5-2m} e^{-4e+\frac{4cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^3f} \\ &\quad + \frac{2^{-4-m} 3^{-1-m} e^{-6e+\frac{6cf}{d}} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{6f(c+dx)}{d}\right)}{a^3f} \end{aligned}$$

```
[Out] 1/8*(d*x+c)^(1+m)/a^3/d/(1+m)+3*2^(-4-m)*exp(-2*e+2*c*f/d)*(d*x+c)^m*GAMMA(1+m,2*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^m)-3*2^(-5-2*m)*exp(-4*e+4*c*f/d)*(d*x+c)^m*GAMMA(1+m,4*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^m)+2^(-4-m)*3^(-1-m)*exp(-6*e+6*c*f/d)*(d*x+c)^m*GAMMA(1+m,6*f*(d*x+c)/d)/a^3/f/((f*(d*x+c)/d)^m)
```

Rubi [A] (verified)

Time = 0.17 (sec), antiderivative size = 223, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used

$$= \{3810, 2212\}$$

$$\begin{aligned} & \int \frac{(c+dx)^m}{(a+a \coth(e+fx))^3} dx \\ &= \frac{3 2^{-m-4} e^{\frac{2cf}{d}-2e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma(m+1, \frac{2f(c+dx)}{d})}{a^3 f} \\ &\quad - \frac{3 2^{-2m-5} e^{\frac{4cf}{d}-4e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma(m+1, \frac{4f(c+dx)}{d})}{a^3 f} \\ &\quad + \frac{2^{-m-4} 3^{-m-1} e^{\frac{6cf}{d}-6e} (c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma(m+1, \frac{6f(c+dx)}{d})}{a^3 f} + \frac{(c+dx)^{m+1}}{8a^3 d(m+1)} \end{aligned}$$

[In] Int[(c + d*x)^m/(a + a*Coth[e + f*x])^3, x]

[Out] $(c + d*x)^{(1 + m)}/(8*a^3*d*(1 + m)) + (3*2^{(-4 - m)}*E^{(-2*e + (2*c*f)/d)}*(c + d*x)^m*\text{Gamma}[1 + m, (2*f*(c + d*x))/d])/((a^3*f*((f*(c + d*x))/d)^m) - (3*2^{(-5 - 2*m)}*E^{(-4*e + (4*c*f)/d)}*(c + d*x)^m*\text{Gamma}[1 + m, (4*f*(c + d*x))/d])/((a^3*f*((f*(c + d*x))/d)^m) + (2^{(-4 - m)}*3^{(-1 - m)}*E^{(-6*e + (6*c*f)/d)}*(c + d*x)^m*\text{Gamma}[1 + m, (6*f*(c + d*x))/d])/((a^3*f*((f*(c + d*x))/d)^m)))$

Rule 2212

```
Int[(F_)^(g_)*(e_)*(x_))*((c_)*(x_))^(m_), x_Symbol]
: > Simp[(-F^(g*(e - c*(f/d))))*((c + d*x)^FracPart[m]/(d*(-f)*g*(Log[F]/d))^(IntPart[m] + 1)*((-f)*g*Log[F]*((c + d*x)/d)^FracPart[m]))*Gamma[m + 1, ((-f)*g*(Log[F]/d))*(c + d*x)], x] /; FreeQ[{F, c, d, e, f, g, m}, x] &&
!IntegerQ[m]
```

Rule 3810

```
Int[((c_)*(x_))^(m_)*((a_)*(b_)*tan[(e_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(2*a) + E^(2*(a/b)*(e + f*x))/(2*a))^{(-n)}, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && EqQ[a^2 + b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \text{integral} &= \int \left(\frac{(c+dx)^m}{8a^3} - \frac{e^{-6e-6fx}(c+dx)^m}{8a^3} + \frac{3e^{-4e-4fx}(c+dx)^m}{8a^3} \right. \\ &\quad \left. - \frac{3e^{-2e-2fx}(c+dx)^m}{8a^3} \right) dx \\ &= \frac{(c+dx)^{1+m}}{8a^3 d(1+m)} - \frac{\int e^{-6e-6fx}(c+dx)^m dx}{8a^3} \\ &\quad + \frac{3 \int e^{-4e-4fx}(c+dx)^m dx}{8a^3} - \frac{3 \int e^{-2e-2fx}(c+dx)^m dx}{8a^3} \end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^{1+m}}{8a^3d(1+m)} + \frac{3 \cdot 2^{-4-m}e^{-2e+\frac{2cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{2f(c+dx)}{d}\right)}{a^3f} \\
&\quad - \frac{3 \cdot 2^{-5-2m}e^{-4e+\frac{4cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{4f(c+dx)}{d}\right)}{a^3f} \\
&\quad + \frac{2^{-4-m}3^{-1-m}e^{-6e+\frac{6cf}{d}}(c+dx)^m \left(\frac{f(c+dx)}{d}\right)^{-m} \Gamma\left(1+m, \frac{6f(c+dx)}{d}\right)}{a^3f}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.97 (sec) , antiderivative size = 228, normalized size of antiderivative = 1.02

$$\begin{aligned}
&\int \frac{(c+dx)^m}{(a+a \coth(e+fx))^3} dx \\
&= \frac{2^{-5-2m}3^{-1-m}e^{-3e}(f(\frac{c}{d}+x))^{-m}(c+dx)^m \operatorname{csch}^3(e+fx) \left(12^{1+m}e^{6e}f(f(\frac{c}{d}+x))^m(c+dx) + 2^{1+m}3^{2+m}de^{6e}f(f(\frac{c}{d}+x))^{m+1}\right)}{a^3}
\end{aligned}$$

[In] `Integrate[(c + d*x)^m/(a + a*Coth[e + f*x])^3, x]`

[Out] `(2^(-5 - 2*m)*3^(-1 - m)*(c + d*x)^m*Csch[e + f*x]^3*(12^(1 + m)*E^(6*e)*f*(f*(c/d + x))^m*(c + d*x) + 2^(1 + m)*3^(2 + m)*d*E^(4*e + (2*c*f)/d)*(1 + m)*Gamma[1 + m, (2*f*(c + d*x))/d] - 3^(2 + m)*d*E^(2*e + (4*c*f)/d)*(1 + m)*Gamma[1 + m, (4*f*(c + d*x))/d] + 2^(1 + m)*d*E^((6*c*f)/d)*(1 + m)*Gamma[1 + m, (6*f*(c + d*x))/d])*(Cosh[f*x] + Sinh[f*x])^3)/(a^3*d*E^(3*e)*f*(1 + m)*(f*(c/d + x))^m*(1 + Coth[e + f*x])^3)`

Maple [F]

$$\int \frac{(dx+c)^m}{(a+a \coth(fx+e))^3} dx$$

[In] `int((d*x+c)^m/(a+a*coth(f*x+e))^3, x)`

[Out] `int((d*x+c)^m/(a+a*coth(f*x+e))^3, x)`

Fricas [A] (verification not implemented)

none

Time = 0.09 (sec) , antiderivative size = 345, normalized size of antiderivative = 1.55

$$\int \frac{(c+dx)^m}{(a+a\coth(e+fx))^3} dx \\ = \frac{2(dm+d)\cosh\left(\frac{dm\log\left(\frac{6f}{d}\right)+6de-6cf}{d}\right)\Gamma\left(m+1, \frac{6(df+cf)}{d}\right) - 9(dm+d)\cosh\left(\frac{dm\log\left(\frac{4f}{d}\right)+4de-4cf}{d}\right)\Gamma\left(m+1, \frac{4(df+cf)}{d}\right)}{a^3}$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^3,x, algorithm="fricas")`

[Out] $\frac{1/96*(2*(d*m + d)*cosh((d*m*log(6*f/d) + 6*d*e - 6*c*f)/d)*gamma(m + 1, 6*(d*f*x + c*f)/d) - 9*(d*m + d)*cosh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d)*gamma(m + 1, 4*(d*f*x + c*f)/d) + 18*(d*m + d)*cosh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d)*gamma(m + 1, 2*(d*f*x + c*f)/d) - 2*(d*m + d)*gamma(m + 1, 6*(d*f*x + c*f)/d)*sinh((d*m*log(6*f/d) + 6*d*e - 6*c*f)/d) + 9*(d*m + d)*gamma(m + 1, 4*(d*f*x + c*f)/d)*sinh((d*m*log(4*f/d) + 4*d*e - 4*c*f)/d) - 18*(d*m + d)*gamma(m + 1, 2*(d*f*x + c*f)/d)*sinh((d*m*log(2*f/d) + 2*d*e - 2*c*f)/d) + 12*(d*f*x + c*f)*cosh(m*log(d*x + c)) + 12*(d*f*x + c*f)*sinh(m*log(d*x + c)))/(a^3*d*f*m + a^3*d*f)$

Sympy [F]

$$\int \frac{(c+dx)^m}{(a+a\coth(e+fx))^3} dx = \frac{\int \frac{(c+dx)^m}{\coth^3(e+fx)+3\coth^2(e+fx)+3\coth(e+fx)+1} dx}{a^3}$$

[In] `integrate((d*x+c)**m/(a+a*coth(f*x+e))**3,x)`

[Out] `Integral((c + d*x)**m/(coth(e + f*x)**3 + 3*coth(e + f*x)**2 + 3*coth(e + f*x) + 1), x)/a**3`

Maxima [F]

$$\int \frac{(c+dx)^m}{(a+a\coth(e+fx))^3} dx = \int \frac{(dx+c)^m}{(a\coth(fx+e)+a)^3} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^3,x, algorithm="maxima")`

[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a)^3, x)`

Giac [F]

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^3} dx = \int \frac{(dx + c)^m}{(a \coth(fx + e) + a)^3} dx$$

[In] `integrate((d*x+c)^m/(a+a*coth(f*x+e))^3,x, algorithm="giac")`

[Out] `integrate((d*x + c)^m/(a*coth(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^m}{(a + a \coth(e + fx))^3} dx = \int \frac{(c + dx)^m}{(a + a \coth(e + fx))^3} dx$$

[In] `int((c + d*x)^m/(a + a*coth(e + f*x))^3,x)`

[Out] `int((c + d*x)^m/(a + a*coth(e + f*x))^3, x)`

3.37 $\int (c + dx)^3 (a + b \coth(e + fx)) dx$

Optimal result	246
Rubi [A] (verified)	246
Mathematica [A] (verified)	249
Maple [B] (verified)	250
Fricas [B] (verification not implemented)	250
Sympy [F]	251
Maxima [B] (verification not implemented)	251
Giac [F]	252
Mupad [F(-1)]	252

Optimal result

Integrand size = 18, antiderivative size = 133

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx)) dx = & \frac{a(c + dx)^4}{4d} - \frac{b(c + dx)^4}{4d} + \frac{b(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{3bd(c + dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\ & - \frac{3bd^2(c + dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\ & + \frac{3bd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{4f^4} \end{aligned}$$

```
[Out] 1/4*a*(d*x+c)^4/d-1/4*b*(d*x+c)^4/d+b*(d*x+c)^3*ln(1-exp(2*f*x+2*e))/f+3/2*
b*d*(d*x+c)^2*polylog(2,exp(2*f*x+2*e))/f^2-3/2*b*d^2*(d*x+c)*polylog(3,exp
(2*f*x+2*e))/f^3+3/4*b*d^3*polylog(4,exp(2*f*x+2*e))/f^4
```

Rubi [A] (verified)

Time = 0.19 (sec), antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.389, Rules used = {3803, 3797, 2221, 2611, 6744, 2320, 6724}

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx)) dx = & \frac{a(c + dx)^4}{4d} - \frac{3bd^2(c + dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\ & + \frac{3bd(c + dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\ & + \frac{b(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{b(c + dx)^4}{4d} + \frac{3bd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{4f^4} \end{aligned}$$

[In] $\text{Int}[(c + d*x)^3*(a + b*\text{Coth}[e + f*x]), x]$

[Out] $(a*(c + d*x)^4)/(4*d) - (b*(c + d*x)^4)/(4*d) + (b*(c + d*x)^3*\text{Log}[1 - E^{(2*(e + f*x))}])/f + (3*b*d*(c + d*x)^2*\text{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2) - (3*b*d^2*(c + d*x)*\text{PolyLog}[3, E^{(2*(e + f*x))}])/(2*f^3) + (3*b*d^3*\text{PolyLog}[4, E^{(2*(e + f*x))}])/(4*f^4)$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*(c_.) + (d_.)*(x_.)^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^m_] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.))*((f_.) + (g_.)*(x_.))^m_, x_Symbol] :> Simplify[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_.))^m_*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_.*(x_.)], x_Symbol] :> Simplify[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_.))^m_*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.*x_)^(p_.)])/((d_.) + (e_.*x_), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.*x_)^m_.*PolyLog[n_, (d_.)*((c_.)*((a_.) + (b_.*x_)^(p_.)))] /; Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c+dx)^3 + b(c+dx)^3 \coth(e+fx)) dx \\
&= \frac{a(c+dx)^4}{4d} + b \int (c+dx)^3 \coth(e+fx) dx \\
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} - (2b) \int \frac{e^{2(e+fx)}(c+dx)^3}{1-e^{2(e+fx)}} dx \\
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad - \frac{(3bd) \int (c+dx)^2 \log(1-e^{2(e+fx)}) dx}{f} \\
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{3bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{(3bd^2) \int (c+dx) \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^2} \\
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{3bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{3bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
&\quad + \frac{(3bd^3) \int \text{PolyLog}(3, e^{2(e+fx)}) dx}{2f^3} \\
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{3bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{3bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} + \frac{(3bd^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3,x)}{x} dx, x, e^{2(e+fx)}\right)}{4f^4}
\end{aligned}$$

$$\begin{aligned}
&= \frac{a(c+dx)^4}{4d} - \frac{b(c+dx)^4}{4d} + \frac{b(c+dx)^3 \log(1 - e^{2(e+fx)})}{f} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{3bd^2(c+dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} + \frac{3bd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{4f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.35 (sec), antiderivative size = 249, normalized size of antiderivative = 1.87

$$\begin{aligned}
\int (c+dx)^3(a+b \coth(e+fx)) dx = & \frac{1}{4} \left(4ac^3x + 6ac^2dx^2 - 6bc^2dx^2 + 4acd^2x^3 - 4bcd^2x^3 \right. \\
&\quad + ad^3x^4 - bd^3x^4 + \frac{12bc^2dx \log(1 - e^{2(e+fx)})}{f} \\
&\quad + \frac{12bcd^2x^2 \log(1 - e^{2(e+fx)})}{f} \\
&\quad + \frac{4bd^3x^3 \log(1 - e^{2(e+fx)})}{f} + \frac{4bc^3 \log(\cosh(e+fx))}{f} \\
&\quad + \frac{4bc^3 \log(\tanh(e+fx))}{f} \\
&\quad + \frac{6bd(c+dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad - \frac{6bd^2(c+dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{f^3} \\
&\quad \left. + \frac{3bd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{f^4} \right)
\end{aligned}$$

[In] `Integrate[(c + d*x)^3*(a + b*Coth[e + f*x]), x]`

[Out] `(4*a*c^3*x + 6*a*c^2*d*x^2 - 6*b*c^2*d*x^2 + 4*a*c*d^2*x^3 - 4*b*c*d^2*x^3 + a*d^3*x^4 - b*d^3*x^4 + (12*b*c^2*d*x*Log[1 - E^(2*(e + f*x))])/f + (12*b*c*d^2*x^2*Log[1 - E^(2*(e + f*x))])/f + (4*b*d^3*x^3*Log[1 - E^(2*(e + f*x))])/f + (4*b*c^3*Log[Cosh[e + f*x]])/f + (4*b*c^3*Log[Tanh[e + f*x]])/f + (6*b*d*(c + d*x)^2*PolyLog[2, E^(2*(e + f*x))])/f^2 - (6*b*d^2*(c + d*x)*PolyLog[3, E^(2*(e + f*x))])/f^3 + (3*b*d^3*PolyLog[4, E^(2*(e + f*x))])/f^4)/4`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 765 vs. $2(123) = 246$.

Time = 0.42 (sec), antiderivative size = 766, normalized size of antiderivative = 5.76

method	result
risch	$-\frac{2bd^3e^3x}{f^3} - \frac{3bd^2c^2e^2}{f^2} + \frac{4bd^2ce^3}{f^3} + \frac{2bd^3e^3\ln(e^{fx+e})}{f^4} + ad^2cx^3 + \frac{3ad^2c^2x^2}{2} + ac^3x + \frac{bc^4}{4d} - \frac{d^3bx^4}{4} + \frac{ad^3x^4}{4} + \dots$

[In] `int((d*x+c)^3*(a+b*coth(f*x+e)), x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -2/f^3*b*d^3*c^3*x^3 - 3/f^2*b*d*c^2*e^2 + 4/f^3*b*d^2*c^2*e^3 + 2/f^4*b*d^3*c^3*\ln(e^{xp(f*x+e)}) + a*d^2*c*x^3 + 3/2*a*d*c^2*x^2 + a*c^3*x^1/4/d*b*c^4 - 1/4*d^3*b*x^4 + 1/4*a*d^3*x^4 + 1/4*a/d*c^4 + 6/f^2*b*d^2*c^2*e^2*x - d^2*b*c*x^3 - 3/2*d*b*c^2*x^2 + b*c^3*x^6/f^3*b*d*c^2*x^2 - f*b*c^3*\ln(\exp(f*x+e)) - 3/2/f^4*b*d^3*c^4 - 6/f^3*b*d^2*c^2*\ln(\exp(f*x+e)) + 6/f^2*b*d*c^2*e*\ln(\exp(f*x+e)) + 3/f^2*b*d*c^2*polylog(2, \exp(f*x+e)) + 3/f^2*b*d*c^2*polylog(2, -\exp(f*x+e)) + 1/f^4*b*d^3*\ln(1 - \exp(f*x+e)) * e^3 + 1/f^2*b*d^3*\ln(1 - \exp(f*x+e)) * x^3 + 3/f^2*b*d^3*polylog(2, \exp(f*x+e)) * x^2 - 6/f^3*b*d^2*c*polylog(3, \exp(f*x+e)) - 6/f^3*b*d^2*c*polylog(3, -\exp(f*x+e)) - 6/f^3*b*d^3*polylog(3, \exp(f*x+e)) * x^1 + 1/f^2*b*d^3*\ln(1 + \exp(f*x+e)) * x^3 + 3/f^2*b*d^3*polylog(2, -\exp(f*x+e)) * x^2 - 6/f^3*b*d^3*polylog(3, -\exp(f*x+e)) * x^1 - 1/f^4*b*d^3*c^3*\ln(\exp(f*x+e) - 1) + 3/f^2*b*d*c^2*\ln(1 - \exp(f*x+e)) * x^3 + 3/f^2*b*d*c^2*\ln(1 - \exp(f*x+e)) * e^3 + 3/f^2*b*d*c^2*\ln(1 + \exp(f*x+e)) * x^3 + 3/f^2*b*d^2*c^2*\ln(\exp(f*x+e) - 1) + 3/f^2*b*d^2*c^2*\ln(1 - \exp(f*x+e)) * x^2 - 3/f^2*b*d^2*c^2*\ln(1 - \exp(f*x+e)) * e^2 + 6/f^2*b*d^2*c*polylog(2, \exp(f*x+e)) * x^3 + 3/f^2*b*d^2*c*\ln(1 + \exp(f*x+e)) * x^2 + 6/f^2*b*d^2*c*polylog(2, -\exp(f*x+e)) * x^1 + 1/f^2*b*c^3*\ln(\exp(f*x+e) - 1) + 1/f^2*b*c^3*\ln(1 + \exp(f*x+e)) + 6/f^4*b*d^3*polylog(4, -\exp(f*x+e)) + 6/f^4*b*d^3*polylog(4, \exp(f*x+e)) - 3/f^2*b*d*c^2*e*\ln(\exp(f*x+e) - 1) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 488 vs. $2(122) = 244$.

Time = 0.26 (sec), antiderivative size = 488, normalized size of antiderivative = 3.67

$$\begin{aligned} & \int (c + dx)^3(a + b \coth(e + fx)) dx \\ &= (a - b)d^3f^4x^4 + 4(a - b)cd^2f^4x^3 + 6(a - b)c^2df^4x^2 + 4(a - b)c^3f^4x + 24bd^3\text{polylog}(4, \cosh(fx + e) + \sinh(fx + e)) + \dots \end{aligned}$$

[In] `integrate((d*x+c)^3*(a+b*coth(f*x+e)), x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/4*((a - b)*d^3*f^4*x^4 + 4*(a - b)*c*d^2*f^4*x^3 + 6*(a - b)*c^2*d*f^4*x^2 + 4*(a - b)*c^3*f^4*x + 24*b*d^3*polylog(4, \cosh(f*x + e) + \sinh(f*x + e)) + 24*b*d^3*polylog(4, -\cosh(f*x + e) - \sinh(f*x + e)) + 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*\text{dilog}(\cosh(f*x + e) + \sinh(f*x + e)) + 12 \end{aligned}$$

$$\begin{aligned} & * (b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(-cosh(f*x + e) - \sinh(f*x + e)) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*c^3*f^3)*\log(\cosh(f*x + e) + \sinh(f*x + e) + 1) - 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*f^2 - b*c^3*f^3)*\log(\cosh(f*x + e) + \sinh(f*x + e) - 1) + 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*f^2)*\log(-cosh(f*x + e) - \sinh(f*x + e) + 1) - 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, \cosh(f*x + e) + \sinh(f*x + e)) - 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, -cosh(f*x + e) - \sinh(f*x + e))/f^4 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^3 (a + b \coth(e + fx)) dx = \int (a + b \coth(e + fx)) (c + dx)^3 dx$$

[In] `integrate((d*x+c)**3*(a+b*coth(f*x+e)),x)`
[Out] `Integral((a + b*coth(e + f*x))*(c + d*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 419 vs. $2(122) = 244$.
Time = 0.26 (sec), antiderivative size = 419, normalized size of antiderivative = 3.15

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx)) dx = & \frac{1}{4} ad^3 x^4 + \frac{1}{4} bd^3 x^4 + acd^2 x^3 + bcd^2 x^3 + \frac{3}{2} ac^2 dx^2 \\ & + \frac{3}{2} bc^2 dx^2 + ac^3 x + \frac{bc^3 \log(\sinh(fx + e))}{f} + \frac{3(fx \log(e^{fx+e}) + 1) + \text{Li}_2(-e^{fx+e}))bc^2 d}{f^2} \\ & + \frac{3(fx \log(-e^{fx+e}) + 1) + \text{Li}_2(e^{fx+e}))bc^2 d}{f^2} \\ & + \frac{3(f^2 x^2 \log(e^{fx+e}) + 1) + 2fx\text{Li}_2(-e^{fx+e}) - 2\text{Li}_3(-e^{fx+e}))bcd^2}{f^3} \\ & + \frac{3(f^2 x^2 \log(-e^{fx+e}) + 1) + 2fx\text{Li}_2(e^{fx+e}) - 2\text{Li}_3(e^{fx+e}))bcd^2}{f^3} \\ & + \frac{(f^3 x^3 \log(e^{fx+e}) + 1) + 3f^2 x^2 \text{Li}_2(-e^{fx+e}) - 6fx\text{Li}_3(-e^{fx+e}) + 6\text{Li}_4(-e^{fx+e}))bd^3}{f^4} \\ & + \frac{(f^3 x^3 \log(-e^{fx+e}) + 1) + 3f^2 x^2 \text{Li}_2(e^{fx+e}) - 6fx\text{Li}_3(e^{fx+e}) + 6\text{Li}_4(e^{fx+e}))bd^3}{f^4} \\ & - \frac{bd^3 f^4 x^4 + 4bcd^2 f^4 x^3 + 6bc^2 d f^4 x^2}{2f^4} \end{aligned}$$

[In] `integrate((d*x+c)^3*(a+b*coth(f*x+e)),x, algorithm="maxima")`

```
[Out] 1/4*a*d^3*x^4 + 1/4*b*d^3*x^4 + a*c*d^2*x^3 + b*c*d^2*x^3 + 3/2*a*c^2*d*x^2
+ 3/2*b*c^2*d*x^2 + a*c^3*x + b*c^3*log(sinh(f*x + e))/f + 3*(f*x*log(e^(f
*x + e) + 1) + dilog(-e^(f*x + e)))*b*c^2*d/f^2 + 3*(f*x*log(-e^(f*x + e) +
1) + dilog(e^(f*x + e)))*b*c^2*d/f^2 + 3*(f^2*x^2*log(e^(f*x + e) + 1) + 2
*f*x*dilog(-e^(f*x + e)) - 2*polylog(3, -e^(f*x + e)))*b*c*d^2/f^3 + 3*(f^2
*x^2*log(-e^(f*x + e) + 1) + 2*f*x*dilog(e^(f*x + e)) - 2*polylog(3, e^(f*x
+ e)))*b*c*d^2/f^3 + (f^3*x^3*log(e^(f*x + e) + 1) + 3*f^2*x^2*dilog(-e^(f
*x + e)) - 6*f*x*polylog(3, -e^(f*x + e)) + 6*polylog(4, -e^(f*x + e)))*b*d
^3/f^4 + (f^3*x^3*log(-e^(f*x + e) + 1) + 3*f^2*x^2*dilog(e^(f*x + e)) - 6*
f*x*polylog(3, e^(f*x + e)) + 6*polylog(4, e^(f*x + e)))*b*d^3/f^4 - 1/2*(b
*d^3*f^4*x^4 + 4*b*c*d^2*f^4*x^3 + 6*b*c^2*d*f^4*x^2)/f^4
```

Giac [F]

$$\int (c + dx)^3(a + b \coth(e + fx)) dx = \int (dx + c)^3(b \coth(fx + e) + a) dx$$

```
[In] integrate((d*x+c)^3*(a+b*coth(f*x+e)),x, algorithm="giac")
[Out] integrate((d*x + c)^3*(b*coth(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3(a + b \coth(e + fx)) dx = \int (a + b \coth(e + fx)) (c + dx)^3 dx$$

```
[In] int((a + b*coth(e + f*x))*(c + d*x)^3,x)
[Out] int((a + b*coth(e + f*x))*(c + d*x)^3, x)
```

3.38 $\int (c + dx)^2(a + b \coth(e + fx)) dx$

Optimal result	253
Rubi [A] (verified)	253
Mathematica [A] (verified)	255
Maple [B] (verified)	256
Fricas [B] (verification not implemented)	256
Sympy [F]	257
Maxima [B] (verification not implemented)	257
Giac [F]	258
Mupad [F(-1)]	258

Optimal result

Integrand size = 18, antiderivative size = 101

$$\begin{aligned} \int (c + dx)^2(a + b \coth(e + fx)) dx = & \frac{a(c + dx)^3}{3d} - \frac{b(c + dx)^3}{3d} + \frac{b(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{bd(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & - \frac{bd^2 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \end{aligned}$$

[Out] $\frac{1}{3}a(d*x+c)^3/d - \frac{1}{3}b(d*x+c)^3/d + b(d*x+c)^2 \ln(1 - \exp(2*f*x+2*e))/f + b*d*(d*x+c)*\operatorname{polylog}(2, \exp(2*f*x+2*e))/f^2 - \frac{1}{2}b*d^2*\operatorname{polylog}(3, \exp(2*f*x+2*e))/f^3$

Rubi [A] (verified)

Time = 0.16 (sec), antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {3803, 3797, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int (c + dx)^2(a + b \coth(e + fx)) dx = & \frac{a(c + dx)^3}{3d} + \frac{bd(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & + \frac{b(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{b(c + dx)^3}{3d} - \frac{bd^2 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2*(a + b*\operatorname{Coth}[e + f*x]), x]$

[Out] $(a*(c + d*x)^3)/(3*d) - (b*(c + d*x)^3)/(3*d) + (b*(c + d*x)^2*\text{Log}[1 - E^{(2*(e + f*x))}])/f + (b*d*(c + d*x)*\text{PolyLog}[2, E^{(2*(e + f*x))}])/f^2 - (b*d^2*\text{PolyLog}[3, E^{(2*(e + f*x))}])/(2*f^3)$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*(c_.) + (d_.)*(x_.)^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_.)^(n_.))^m_] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_.)^(m_.)), x_Symbol] :> Simplify[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_.))^m_*tan[(e_.) + Pi*(k_.)*(f_.)*(x_.)], x_Symbol] :> Simplify[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_.))^m_*((a_) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Int[ExpandIntegrand[((c + d*x)^m, (a + b*Tan[e + f*x])^n, x)], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_.))^p_]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simplify[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c+dx)^2 + b(c+dx)^2 \coth(e+fx)) \, dx \\
&= \frac{a(c+dx)^3}{3d} + b \int (c+dx)^2 \coth(e+fx) \, dx \\
&= \frac{a(c+dx)^3}{3d} - \frac{b(c+dx)^3}{3d} - (2b) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} \, dx \\
&= \frac{a(c+dx)^3}{3d} - \frac{b(c+dx)^3}{3d} + \frac{b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad - \frac{(2bd) \int (c+dx) \log(1-e^{2(e+fx)}) \, dx}{f} \\
&= \frac{a(c+dx)^3}{3d} - \frac{b(c+dx)^3}{3d} + \frac{b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{(bd^2) \int \text{PolyLog}(2, e^{2(e+fx)}) \, dx}{f^2} \\
&= \frac{a(c+dx)^3}{3d} - \frac{b(c+dx)^3}{3d} + \frac{b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{(bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} \, dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&= \frac{a(c+dx)^3}{3d} - \frac{b(c+dx)^3}{3d} + \frac{b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{bd^2 \text{PolyLog}(3, e^{2(e+fx)})}{2f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.21 (sec), antiderivative size = 149, normalized size of antiderivative = 1.48

$$\begin{aligned}
&\int (c+dx)^2(a+b \coth(e+fx)) \, dx \\
&= \frac{2f^2(3ac^2fx + 3acd^2x^2 - 3bcd^2x^2 + ad^2fx^3 - bd^2fx^3 + 3bdx(2c+dx) \log(1-e^{2(e+fx)}) + 3bc^2 \log(\cosh(6f^3)
\end{aligned}$$

```
[In] Integrate[(c + d*x)^2*(a + b*Coth[e + f*x]), x]
[Out] (2*f^2*(3*a*c^2*f*x + 3*a*c*d*f*x^2 - 3*b*c*d*f*x^2 + a*d^2*f*x^3 - b*d^2*f*x^3 + 3*b*d*x*(2*c + d*x)*Log[1 - E^(2*(e + f*x))] + 3*b*c^2*Log[Cosh[e + f*x]] + 3*b*c^2*Log[Tanh[e + f*x]] + 6*b*d*f*(c + d*x)*PolyLog[2, E^(2*(e + f*x))] - 3*b*d^2*PolyLog[3, E^(2*(e + f*x))])/(6*f^3)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 464 vs. $2(95) = 190$.

Time = 0.36 (sec), antiderivative size = 465, normalized size of antiderivative = 4.60

method	result
risch	$-\frac{2bdce^2}{f^2} - \frac{2bd^2e^2 \ln(e^{fx+e})}{f^3} - \frac{d^2bx^3}{3} + \frac{bc^3}{3d} + \frac{4bdce \ln(e^{fx+e})}{f^2} + ac^2x + adc x^2 + \frac{ad^2x^3}{3} + \frac{ac^3}{3d} - \frac{4bdce}{f} - db$

[In] `int((d*x+c)^2*(a+b*cot(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -2/f^2 * b * d * c * e^2 - 2/f^3 * b * d^2 * e^2 * \ln(\exp(f*x+e)) - 1/3 * d^2 * b * x^3 + 1/3 * d * b * c^3 + 4 \\ & /f^2 * b * d * c * e * \ln(\exp(f*x+e)) + a * c^2 * x + a * d * c * x^2 + 1/3 * a * d^2 * x^3 + 1/3 * a / d * c^3 - 4/f \\ & * b * d * c * e * x - d * b * c * x^2 + b * c^2 * x + 4/3 * f^3 * b * d^2 * e^3 - 2/f * b * c^2 * \ln(\exp(f*x+e)) + 2/f \\ & ^2 * b * d^2 * e^2 * x + 2/f^2 * b * d^2 * \text{polylog}(2, -\exp(f*x+e)) * x + 2/f^2 * b * d * c * \text{polylog}(2, \exp(f*x+e)) + 2/f^2 * b * d * c * \text{polylog}(2, -\exp(f*x+e)) + 1/f^3 * b * d^2 * e^2 * \ln(\exp(f*x+e)) - 1 \\ & + 2/f * b * d * c * \ln(1 - \exp(f*x+e)) * x + 2/f^2 * b * d * c * \ln(1 - \exp(f*x+e)) * e + 2/f * b * d * c * \ln(1 + \exp(f*x+e)) * x - 2/f^2 * b * d * c * \ln(\exp(f*x+e) - 1) + 1/f * b * d^2 * \ln(1 - \exp(f*x+e)) * x^2 - 1/f^3 * b * d^2 * \ln(1 - \exp(f*x+e)) * e^2 + 2/f^2 * b * d^2 * \text{polylog}(2, \exp(f*x+e)) * x + 1/f * b * d^2 * \ln(1 + \exp(f*x+e)) * x^2 - 2/f^3 * b * d^2 * \text{polylog}(3, -\exp(f*x+e)) - 2/f^3 * b * d^2 * \text{polylog}(3, \exp(f*x+e)) + 1/f * b * c^2 * \ln(\exp(f*x+e) - 1) + 1/f * b * c^2 * \ln(1 + \exp(f*x+e))) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 303 vs. $2(94) = 188$.

Time = 0.26 (sec), antiderivative size = 303, normalized size of antiderivative = 3.00

$$\begin{aligned} & \int (c + dx)^2 (a + b \coth(e + fx)) dx \\ &= (a - b)d^2 f^3 x^3 + 3(a - b)c d f^3 x^2 + 3(a - b)c^2 f^3 x - 6bd^2 \text{polylog}(3, \cosh(fx + e) + \sinh(fx + e)) - 6bd^2 \text{polylog}(3, \cosh(fx + e) - \sinh(fx + e)) + 6(b * d^2 * f * x + b * c * d * f) * \text{dilog}(\cosh(f*x + e) + \sinh(f*x + e)) + 6(b * d^2 * f * x + b * c * d * f) * \text{dilog}(-\cosh(f*x + e) - \sinh(f*x + e)) + 3(b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x + b * c^2 * f^2) * \text{log}(\cosh(f*x + e) + \sinh(f*x + e) + 1) + 3(b * d^2 * e^2 - 2 * b * c * d * e * f + b * c^2 * f^2) * \text{log}(\cosh(f*x + e) + \sinh(f*x + e) - 1) + 3(b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x - b * d^2 * e^2 + 2 * b * c * d * e * f) * \text{log}(-\cosh(f*x + e) - \sinh(f*x + e) + 1) / f^3 \end{aligned}$$

[In] `integrate((d*x+c)^2*(a+b*cot(f*x+e)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/3 * ((a - b) * d^2 * f^3 * x^3 + 3 * (a - b) * c * d * f^3 * x^2 + 3 * (a - b) * c^2 * f^3 * x - 6 * \\ & b * d^2 * \text{polylog}(3, \cosh(f*x + e) + \sinh(f*x + e)) - 6 * b * d^2 * \text{polylog}(3, -\cosh(f*x + e) - \sinh(f*x + e)) + 6 * (b * d^2 * f * x + b * c * d * f) * \text{dilog}(\cosh(f*x + e) + \sinh(f*x + e)) + 6 * (b * d^2 * f * x + b * c * d * f) * \text{dilog}(-\cosh(f*x + e) - \sinh(f*x + e)) + 3 * (b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x + b * c^2 * f^2) * \text{log}(\cosh(f*x + e) + \sinh(f*x + e) + 1) + 3 * (b * d^2 * e^2 - 2 * b * c * d * e * f + b * c^2 * f^2) * \text{log}(\cosh(f*x + e) + \sinh(f*x + e) - 1) + 3 * (b * d^2 * f^2 * x^2 + 2 * b * c * d * f^2 * x - b * d^2 * e^2 + 2 * b * c * d * e * f) * \text{log}(-\cosh(f*x + e) - \sinh(f*x + e) + 1)) / f^3 \end{aligned}$$

Sympy [F]

$$\int (c + dx)^2 (a + b \coth(e + fx)) dx = \int (a + b \coth(e + fx)) (c + dx)^2 dx$$

```
[In] integrate((d*x+c)**2*(a+b*coth(f*x+e)),x)
[Out] Integral((a + b*coth(e + f*x))*(c + d*x)**2, x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(94) = 188$.
 Time = 0.26 (sec) , antiderivative size = 240, normalized size of antiderivative = 2.38

$$\begin{aligned} & \int (c + dx)^2 (a + b \coth(e + fx)) dx \\ &= \frac{1}{3} ad^2 x^3 + \frac{1}{3} bd^2 x^3 + acdx^2 + bcdx^2 + ac^2 x + \frac{bc^2 \log(\sinh(fx + e))}{f} \\ &+ \frac{2(fx \log(e^{fx+e}) + 1) + \text{Li}_2(-e^{fx+e}))bcd}{f^2} \\ &+ \frac{2(fx \log(-e^{fx+e}) + 1) + \text{Li}_2(e^{fx+e}))bcd}{f^2} \\ &+ \frac{(f^2 x^2 \log(e^{fx+e}) + 1) + 2fx\text{Li}_2(-e^{fx+e}) - 2\text{Li}_3(-e^{fx+e}))bd^2}{f^3} \\ &+ \frac{(f^2 x^2 \log(-e^{fx+e}) + 1) + 2fx\text{Li}_2(e^{fx+e}) - 2\text{Li}_3(e^{fx+e}))bd^2}{f^3} \\ &- \frac{2(bd^2 f^3 x^3 + 3bcd f^3 x^2)}{3 f^3} \end{aligned}$$

```
[In] integrate((d*x+c)^2*(a+b*coth(f*x+e)),x, algorithm="maxima")
[Out] 1/3*a*d^2*x^3 + 1/3*b*d^2*x^3 + a*c*d*x^2 + b*c*d*x^2 + a*c^2*x + b*c^2*log(sinh(f*x + e))/f + 2*(f*x*log(e^(f*x + e) + 1) + dilog(-e^(f*x + e)))*b*c*d/f^2 + 2*(f*x*log(-e^(f*x + e) + 1) + dilog(e^(f*x + e)))*b*c*d/f^2 + (f^2*x^2*log(e^(f*x + e) + 1) + 2*f*x*dilog(-e^(f*x + e)) - 2*polylog(3, -e^(f*x + e)))*b*d^2/f^3 + (f^2*x^2*log(-e^(f*x + e) + 1) + 2*f*x*dilog(e^(f*x + e)) - 2*polylog(3, e^(f*x + e)))*b*d^2/f^3 - 2/3*(b*d^2*f^3*x^3 + 3*b*c*d*f^3*x^2)/f^3
```

Giac [F]

$$\int (c + dx)^2(a + b \coth(e + fx)) dx = \int (dx + c)^2(b \coth(fx + e) + a) dx$$

[In] `integrate((d*x+c)^2*(a+b*coth(f*x+e)),x, algorithm="giac")`

[Out] `integrate((d*x + c)^2*(b*coth(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2(a + b \coth(e + fx)) dx = \int (a + b \coth(e + fx)) (c + dx)^2 dx$$

[In] `int((a + b*coth(e + f*x))*(c + d*x)^2,x)`

[Out] `int((a + b*coth(e + f*x))*(c + d*x)^2, x)`

$$\text{3.39} \quad \int (c + dx)(a + b \coth(e + fx)) dx$$

Optimal result	259
Rubi [A] (verified)	259
Mathematica [A] (verified)	261
Maple [B] (verified)	261
Fricas [B] (verification not implemented)	261
Sympy [F]	262
Maxima [F]	262
Giac [F]	262
Mupad [F(-1)]	263

Optimal result

Integrand size = 16, antiderivative size = 75

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx)) dx = & \frac{a(c + dx)^2}{2d} - \frac{b(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \end{aligned}$$

[Out] $\frac{1}{2}a(d*x+c)^2/d - \frac{1}{2}b(d*x+c)^2/d + b*(d*x+c)*\ln(1-\exp(2*f*x+2*e))/f + \frac{1}{2}b*d*\operatorname{polylog}(2, \exp(2*f*x+2*e))/f^2$

Rubi [A] (verified)

Time = 0.09 (sec), antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {3803, 3797, 2221, 2317, 2438}

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx)) dx = & \frac{a(c + dx)^2}{2d} + \frac{b(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{b(c + dx)^2}{2d} + \frac{bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Coth}[e + f*x]), x]$

[Out] $(a*(c + d*x)^2)/(2*d) - (b*(c + d*x)^2)/(2*d) + (b*(c + d*x))*\operatorname{Log}[1 - E^{(2*(e + f*x))}]/f + (b*d*\operatorname{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2)$

Rule 2221

$\operatorname{Int}[(((F_*)^((g_.)*(e_.) + (f_.)*(x_.)))^((n_.)*(c_.) + (d_.)*(x_.))^((m_.)))/((a_) + (b_.)*((F_*)^((g_.)*(e_.) + (f_.)*(x_.))))^((n_.)), x_{\text{Symbol}}] \Rightarrow \operatorname{Simp}$

```
[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_)*(F_)*((e_)*(c_) + (d_)*(x_)))^(n_), x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_)*(d_) + (e_)*(x_)^(n_)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3797

```
Int[((c_) + (d_)*(x_))^(m_)*tan[(e_) + Pi*(k_) + (Complex[0, fz_])*f_*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3803

```
Int[((c_) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a(c+dx) + b(c+dx) \coth(e+fx)) dx \\
&= \frac{a(c+dx)^2}{2d} + b \int (c+dx) \coth(e+fx) dx \\
&= \frac{a(c+dx)^2}{2d} - \frac{b(c+dx)^2}{2d} - (2b) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} dx \\
&= \frac{a(c+dx)^2}{2d} - \frac{b(c+dx)^2}{2d} + \frac{b(c+dx) \log(1-e^{2(e+fx)})}{f} - \frac{(bd) \int \log(1-e^{2(e+fx)}) dx}{f} \\
&= \frac{a(c+dx)^2}{2d} - \frac{b(c+dx)^2}{2d} + \frac{b(c+dx) \log(1-e^{2(e+fx)})}{f} - \frac{(bd) \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} \\
&= \frac{a(c+dx)^2}{2d} - \frac{b(c+dx)^2}{2d} + \frac{b(c+dx) \log(1-e^{2(e+fx)})}{f} + \frac{bd \text{PolyLog}(2, e^{2(e+fx)})}{2f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 0.05 (sec) , antiderivative size = 87, normalized size of antiderivative = 1.16

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx)) dx &= acx + \frac{1}{2}adx^2 - \frac{1}{2}bdx^2 + \frac{bdx \log(1 - e^{2e+2fx})}{f} \\ &+ \frac{bc(\log(\cosh(e + fx)) + \log(\tanh(e + fx))))}{f} \\ &+ \frac{bd \operatorname{PolyLog}(2, e^{2e+2fx})}{2f^2} \end{aligned}$$

[In] `Integrate[(c + d*x)*(a + b*Coth[e + f*x]), x]`

[Out] $a*c*x + (a*d*x^2)/2 - (b*d*x^2)/2 + (b*d*x*Log[1 - E^(2*e + 2*f*x)])/f + (b*c*(Log[Cosh[e + f*x]] + Log[Tanh[e + f*x]]))/f + (b*d*PolyLog[2, E^(2*e + 2*f*x)])/(2*f^2)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 200 vs. $2(69) = 138$.

Time = 0.19 (sec) , antiderivative size = 201, normalized size of antiderivative = 2.68

method	result
risch	$\frac{adx^2}{2} + acx - \frac{bdx^2}{2} + bcx + \frac{bc \ln(e^{fx+e}-1)}{f} + \frac{bc \ln(1+e^{fx+e})}{f} - \frac{2bc \ln(e^{fx+e})}{f} - \frac{2bdex}{f} - \frac{bd e^2}{f^2} + \frac{bd \ln(1-e^{fx+e})}{f}$

[In] `int((d*x+c)*(a+b*coth(f*x+e)), x, method=_RETURNVERBOSE)`

[Out] $1/2*a*d*x^2+a*c*x-1/2*b*d*x^2+b*c*x+1/f*b*c*ln(exp(f*x+e)-1)+1/f*b*c*ln(1+exp(f*x+e))-2/f*b*c*ln(exp(f*x+e))-2/f*b*d*e*x-1/f^2*b*d*e^2+1/f*b*d*ln(1-exp(f*x+e))*x+1/f^2*b*d*ln(1-exp(f*x+e))*e+1/f^2*b*d*polylog(2, exp(f*x+e))+1/f*b*d*ln(1+exp(f*x+e))*x+1/f^2*b*d*polylog(2, -exp(f*x+e))-1/f^2*b*d*e*ln(exp(f*x+e))-1/2/f^2*b*d*e*ln(exp(f*x+e))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(68) = 136$.

Time = 0.26 (sec) , antiderivative size = 156, normalized size of antiderivative = 2.08

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx)) dx \\ = (a - b)df^2x^2 + 2(a - b)cf^2x + 2bd\text{Li}_2(\cosh(fx + e) + \sinh(fx + e)) + 2bd\text{Li}_2(-\cosh(fx + e) - \sinh(fx + e)) \end{aligned}$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e)),x, algorithm="fricas")`
[Out]
$$\frac{1}{2}((a - b)*d*f^2*x^2 + 2*(a - b)*c*f^2*x + 2*b*d*dilog(\cosh(f*x + e) + \sinh(f*x + e)) + 2*b*d*dilog(-\cosh(f*x + e) - \sinh(f*x + e)) + 2*(b*d*f*x + b*c*f)*log(\cosh(f*x + e) + \sinh(f*x + e) + 1) - 2*(b*d*e - b*c*f)*log(\cosh(f*x + e) + \sinh(f*x + e) - 1) + 2*(b*d*f*x + b*d*e)*log(-\cosh(f*x + e) - \sinh(f*x + e) + 1))/f^2$$

Sympy [F]

$$\int (c + dx)(a + b \coth(e + fx)) dx = \int (a + b \coth(e + fx))(c + dx) dx$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e)),x)`
[Out] `Integral((a + b*coth(e + f*x))*(c + d*x), x)`

Maxima [F]

$$\int (c + dx)(a + b \coth(e + fx)) dx = \int (dx + c)(b \coth(fx + e) + a) dx$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e)),x, algorithm="maxima")`
[Out]
$$\frac{1}{2}a*d*x^2 + \frac{1}{2}*(x^2 - 2*\int \frac{x}{(e^{(f*x + e)} + 1)} dx + 2*\int \frac{x}{(e^{(f*x + e)} - 1)} dx)*b*d + a*c*x + b*c*log(\sinh(f*x + e))/f$$

Giac [F]

$$\int (c + dx)(a + b \coth(e + fx)) dx = \int (dx + c)(b \coth(fx + e) + a) dx$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e)),x, algorithm="giac")`
[Out] `integrate((d*x + c)*(b*coth(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \coth(e + fx)) dx = \int (a + b \coth(e + f x)) (c + d x) dx$$

[In] `int((a + b*coth(e + f*x))*(c + d*x),x)`

[Out] `int((a + b*coth(e + f*x))*(c + d*x), x)`

3.40 $\int \frac{a+b \coth(e+fx)}{c+dx} dx$

Optimal result	264
Rubi [N/A]	264
Mathematica [N/A]	265
Maple [N/A] (verified)	265
Fricas [N/A]	265
Sympy [N/A]	265
Maxima [N/A]	266
Giac [N/A]	266
Mupad [N/A]	266

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \text{Int}\left(\frac{a + b \coth(e + fx)}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*coth(f*x+e))/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{a + b \coth(e + fx)}{c + dx} dx$$

[In] Int[(a + b*Coth[e + f*x])/(c + d*x),x]

[Out] Defer[Int][(a + b*Coth[e + f*x])/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{a + b \coth(e + fx)}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 5.73 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{a + b \coth(e + fx)}{c + dx} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])/(c + d*x), x]`

[Out] `Integrate[(a + b*Coth[e + f*x])/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \coth(fx + e)}{dx + c} dx$$

[In] `int((a+b*coth(f*x+e))/(d*x+c), x)`

[Out] `int((a+b*coth(f*x+e))/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{b \coth(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b*coth(f*x + e) + a)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 0.96 (sec) , antiderivative size = 15, normalized size of antiderivative = 0.83

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{a + b \coth(e + fx)}{c + dx} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c), x)`

[Out] `Integral((a + b*coth(e + f*x))/ (c + d*x), x)`

Maxima [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 78, normalized size of antiderivative = 4.33

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{b \coth(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c),x, algorithm="maxima")`

[Out] `b*(log(d*x + c)/d - integrate(1/(d*x + (d*x*e^e + c*e^e)*e^(f*x) + c), x) + integrate(-1/(d*x - (d*x*e^e + c*e^e)*e^(f*x) + c), x)) + a*log(d*x + c)/d`

Giac [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{b \coth(fx + e) + a}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c),x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 1.97 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{c + dx} dx = \int \frac{a + b \coth(e + f x)}{c + d x} dx$$

[In] `int((a + b*coth(e + f*x))/(c + d*x),x)`

[Out] `int((a + b*coth(e + f*x))/(c + d*x), x)`

3.41 $\int \frac{a+b \coth(e+fx)}{(c+dx)^2} dx$

Optimal result	267
Rubi [N/A]	267
Mathematica [N/A]	268
Maple [N/A] (verified)	268
Fricas [N/A]	268
Sympy [N/A]	268
Maxima [N/A]	269
Giac [N/A]	269
Mupad [N/A]	269

Optimal result

Integrand size = 18, antiderivative size = 18

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \text{Int}\left(\frac{a + b \coth(e + fx)}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*coth(f*x+e))/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.02 (sec), antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx$$

[In] Int[(a + b*Coth[e + f*x])/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Coth[e + f*x])/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 31.41 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])/(c + d*x)^2, x]`

[Out] `Integrate[(a + b*Coth[e + f*x])/(c + d*x)^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.06 (sec) , antiderivative size = 18, normalized size of antiderivative = 1.00

$$\int \frac{a + b \coth(fx + e)}{(dx + c)^2} dx$$

[In] `int((a+b*coth(f*x+e))/(d*x+c)^2, x)`

[Out] `int((a+b*coth(f*x+e))/(d*x+c)^2, x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 31, normalized size of antiderivative = 1.72

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{b \coth(fx + e) + a}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c)^2, x, algorithm="fricas")`

[Out] `integral((b*coth(f*x + e) + a)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 3.17 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.94

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c)**2, x)`

[Out] `Integral((a + b*coth(e + f*x))/(c + d*x)**2, x)`

Maxima [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 131, normalized size of antiderivative = 7.28

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{b \coth(fx + e) + a}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c)^2,x, algorithm="maxima")`

[Out] `-b*(1/(d^2*x + c*d) + integrate(1/(d^2*x^2 + 2*c*d*x + c^2 + (d^2*x^2*e^e + 2*c*d*x*e^e + c^2*e^e)*e^(f*x)), x) - integrate(-1/(d^2*x^2 + 2*c*d*x + c^2 - (d^2*x^2*e^e + 2*c*d*x*e^e + c^2*e^e)*e^(f*x)), x)) - a/(d^2*x + c*d)`

Giac [N/A]

Not integrable

Time = 0.36 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{b \coth(fx + e) + a}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))/(d*x+c)^2,x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 2.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.11

$$\int \frac{a + b \coth(e + fx)}{(c + dx)^2} dx = \int \frac{a + b \coth(e + f x)}{(c + d x)^2} dx$$

[In] `int((a + b*coth(e + f*x))/(c + d*x)^2,x)`

[Out] `int((a + b*coth(e + f*x))/(c + d*x)^2, x)`

3.42 $\int (c + dx)^3 (a + b \coth(e + fx))^2 dx$

Optimal result	270
Rubi [A] (verified)	271
Mathematica [B] (verified)	275
Maple [B] (verified)	276
Fricas [B] (verification not implemented)	277
Sympy [F]	279
Maxima [B] (verification not implemented)	279
Giac [F]	280
Mupad [F(-1)]	280

Optimal result

Integrand size = 20, antiderivative size = 271

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx))^2 dx = & -\frac{b^2(c + dx)^3}{f} + \frac{a^2(c + dx)^4}{4d} - \frac{ab(c + dx)^4}{2d} \\ & + \frac{b^2(c + dx)^4}{4d} - \frac{b^2(c + dx)^3 \coth(e + fx)}{f} \\ & + \frac{3b^2d(c + dx)^2 \log(1 - e^{2(e+fx)})}{f^2} \\ & + \frac{2ab(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{3b^2d^2(c + dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\ & + \frac{3abd(c + dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & - \frac{3b^2d^3 \text{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\ & - \frac{3abd^2(c + dx) \text{PolyLog}(3, e^{2(e+fx)})}{f^3} \\ & + \frac{3abd^3 \text{PolyLog}(4, e^{2(e+fx)})}{2f^4} \end{aligned}$$

[Out]
$$\begin{aligned} & -b^2*(d*x+c)^3/f+1/4*a^2*(d*x+c)^4/d-1/2*a*b*(d*x+c)^4/d+1/4*b^2*(d*x+c)^4/ \\ & d-b^2*(d*x+c)^3*\coth(f*x+e)/f+3*b^2*d*(d*x+c)^2*\ln(1-\exp(2*f*x+2*e))/f^2+2* \\ & a*b*(d*x+c)^3*\ln(1-\exp(2*f*x+2*e))/f+3*b^2*d^2*(d*x+c)*\text{polylog}(2,\exp(2*f*x+ \\ & 2*e))/f^3+3*a*b*d*(d*x+c)^2*\text{polylog}(2,\exp(2*f*x+2*e))/f^2-3/2*b^2*d^3*\text{polyl} \\ & og(3,\exp(2*f*x+2*e))/f^4-3*a*b*d^2*(d*x+c)*\text{polylog}(3,\exp(2*f*x+2*e))/f^3+3/ \\ & 2*a*b*d^3*\text{polylog}(4,\exp(2*f*x+2*e))/f^4 \end{aligned}$$

Rubi [A] (verified)

Time = 0.40 (sec) , antiderivative size = 271, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.450$, Rules used = {3803, 3797, 2221, 2611, 6744, 2320, 6724, 3801, 32}

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx))^2 dx = & \frac{a^2(c + dx)^4}{4d} - \frac{3abd^2(c + dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{f^3} \\ & + \frac{3abd(c + dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & + \frac{2ab(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{ab(c + dx)^4}{2d} + \frac{3abd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{2f^4} \\ & + \frac{3b^2d^2(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^3} \\ & + \frac{3b^2d(c + dx)^2 \log(1 - e^{2(e+fx)})}{f^2} \\ & - \frac{b^2(c + dx)^3 \coth(e + fx)}{f} - \frac{b^2(c + dx)^3}{f} \\ & + \frac{b^2(c + dx)^4}{4d} - \frac{3b^2d^3 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^4} \end{aligned}$$

```
[In] Int[(c + d*x)^3*(a + b*Coth[e + f*x])^2, x]
[Out] -((b^2*(c + d*x)^3)/f) + (a^2*(c + d*x)^4)/(4*d) - (a*b*(c + d*x)^4)/(2*d)
+ (b^2*(c + d*x)^4)/(4*d) - (b^2*(c + d*x)^3*Coth[e + f*x])/f + (3*b^2*d*(c
+ d*x)^2*Log[1 - E^(2*(e + f*x))])/f^2 + (2*a*b*(c + d*x)^3*Log[1 - E^(2*(e
+ f*x))])/f + (3*b^2*d^2*(c + d*x)*PolyLog[2, E^(2*(e + f*x))])/f^3 + (3*a
*b*d*(c + d*x)^2*PolyLog[2, E^(2*(e + f*x))])/f^2 - (3*b^2*d^3*PolyLog[3,
E^(2*(e + f*x))])/f^4 - (3*a*b*d^2*(c + d*x)*PolyLog[3, E^(2*(e + f*x))])
/f^3 + (3*a*b*d^3*PolyLog[4, E^(2*(e + f*x))])/f^4
```

Rule 32

```
Int[((a_.) + (b_.)*(x_.))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m +
1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_
) + (b_.)*(F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c
+ d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x]]
```

```
)^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_.)*(a_.) + (b_.)*(x_)))^(n_.)]*((f_.) + (g_.)*(x_)^(m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^m_*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*(f_.)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^m_*((b_.)*tan[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^m_*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^p]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_ .)*(x_ ))^(m_ .)*PolyLog[n_, (d_ .)*((F_)^((c_ .)*((a_ .) + (b_ .)*x_ ))^p_ .)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx)^3 + 2ab(c+dx)^3 \coth(e+fx) + b^2(c+dx)^3 \coth^2(e+fx)) dx \\
&= \frac{a^2(c+dx)^4}{4d} + (2ab) \int (c+dx)^3 \coth(e+fx) dx + b^2 \int (c+dx)^3 \coth^2(e+fx) dx \\
&= \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad - (4ab) \int \frac{e^{2(e+fx)}(c+dx)^3}{1-e^{2(e+fx)}} dx + b^2 \int (c+dx)^3 dx \\
&\quad + \frac{(3b^2d) \int (c+dx)^2 \coth(e+fx) dx}{f} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} + \frac{2ab(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad - \frac{(6abd) \int (c+dx)^2 \log(1-e^{2(e+fx)}) dx}{f} - \frac{(6b^2d) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} dx}{f} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{3abd(c+dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{(6abd^2) \int (c+dx) \operatorname{PolyLog}(2, e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6b^2d^2) \int (c+dx) \log(1-e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} + \frac{3b^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{2ab(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{3b^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{3abd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{3abd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{(3abd^3) \int \text{PolyLog}(3, e^{2(e+fx)}) dx}{f^3} - \frac{(3b^2d^3) \int \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^3} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad + \frac{3b^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{2ab(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{3b^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} + \frac{3abd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad - \frac{3abd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{f^3} + \frac{(3abd^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3,x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&\quad - \frac{(3b^2d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&= -\frac{b^2(c+dx)^3}{f} + \frac{a^2(c+dx)^4}{4d} - \frac{ab(c+dx)^4}{2d} + \frac{b^2(c+dx)^4}{4d} \\
&\quad - \frac{b^2(c+dx)^3 \coth(e+fx)}{f} + \frac{3b^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{2ab(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{3b^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{3abd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{3b^2d^3 \text{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\
&\quad - \frac{3abd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{f^3} + \frac{3abd^3 \text{PolyLog}(4, e^{2(e+fx)})}{2f^4}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 843 vs. $2(271) = 542$.

Time = 4.91 (sec), antiderivative size = 843, normalized size of antiderivative = 3.11

$$\begin{aligned}
 \int (c + dx)^3 (a + b \coth(e + fx))^2 dx = & -\frac{2bc^2(3bd + 2acf)x}{f} - \frac{2b^2(c + dx)^3}{(-1 + e^{2e})f} - \frac{ab(c + dx)^4}{d(-1 + e^{2e})} \\
 & + \frac{6bcd(bd + acf)x \log(1 - e^{-e-fx})}{f^2} + \frac{3bd^2(bd + 2acf)x^2 \log(1 - e^{-e-fx})}{f^2} \\
 & + \frac{2abd^3x^3 \log(1 - e^{-e-fx})}{f} + \frac{6bcd(bd + acf)x \log(1 + e^{-e-fx})}{f^2} \\
 & + \frac{3bd^2(bd + 2acf)x^2 \log(1 + e^{-e-fx})}{f^2} + \frac{2abd^3x^3 \log(1 + e^{-e-fx})}{f} \\
 & + \frac{bc^2(3bd + 2acf) \log(1 - e^{e+fx})}{f^2} + \frac{bc^2(3bd + 2acf) \log(1 + e^{e+fx})}{f^2} \\
 & - \frac{6bcd(bd + acf) \text{PolyLog}(2, -e^{-e-fx})}{f^3} - \frac{6bd^2(bd + 2acf)x \text{PolyLog}(2, -e^{-e-fx})}{f^3} \\
 & - \frac{6abd^3x^2 \text{PolyLog}(2, -e^{-e-fx})}{f^2} - \frac{6bcd(bd + acf) \text{PolyLog}(2, e^{-e-fx})}{f^3} \\
 & - \frac{6bd^2(bd + 2acf)x \text{PolyLog}(2, e^{-e-fx})}{f^3} - \frac{6abd^3x^2 \text{PolyLog}(2, e^{-e-fx})}{f^2} \\
 & - \frac{6bd^2(bd + 2acf) \text{PolyLog}(3, -e^{-e-fx})}{f^4} - \frac{12abd^3x \text{PolyLog}(3, -e^{-e-fx})}{f^3} \\
 & - \frac{6bd^2(bd + 2acf) \text{PolyLog}(3, e^{-e-fx})}{f^4} - \frac{12abd^3x \text{PolyLog}(3, e^{-e-fx})}{f^3} \\
 & - \frac{12abd^3 \text{PolyLog}(4, -e^{-e-fx})}{f^4} - \frac{12abd^3 \text{PolyLog}(4, e^{-e-fx})}{f^4} \\
 & + \text{csch}(e) \text{csch}(e + fx) ((a^2 + b^2) fx(4c^3 + 6c^2dx + 4cd^2x^2 + d^3x^3) \cosh(fx)) + (a^2 + b^2) fx(4c^3 + 6c^2dx + 4cd^2x^2 + d^3x^3) \sinh(fx)
 \end{aligned}$$

[In] `Integrate[(c + d*x)^3*(a + b*Coth[e + f*x])^2, x]`

[Out]
$$\begin{aligned}
 & (-2*b*c^2*(3*b*d + 2*a*c*f)*x)/f - (2*b^2*(c + d*x)^3)/((-1 + E^{(2*e)})*f) - \\
 & (a*b*(c + d*x)^4)/(d*(-1 + E^{(2*e)})) + (6*b*c*d*(b*d + a*c*f)*x*\text{Log}[1 - E^{-e - f*x}])/f^2 + \\
 & (3*b*d^2*(b*d + 2*a*c*f)*x^2*\text{Log}[1 - E^{-e - f*x}])/f^2 + (2*a*b*d^3*x^3*\text{Log}[1 - E^{-e - f*x}])/f + \\
 & (6*b*c*d*(b*d + a*c*f)*x*\text{Log}[1 + E^{-e - f*x}])/f^2 + (3*b*d^2*(b*d + 2*a*c*f)*x^2*\text{Log}[1 + E^{-e - f*x}])/f^2 + \\
 & (2*a*b*d^3*x^3*\text{Log}[1 + E^{-e - f*x}])/f + (b*c^2*(3*b*d + 2*a*c*f)*\text{Log}[1 - E^{(e + f*x)}])/f^2 + \\
 & (b*c^2*(3*b*d + 2*a*c*f)*\text{Log}[1 + E^{(e + f*x)}])/f^2 - (6*b*c*d*(b*d + a*c*f)*\text{PolyLog}[2, -E^{-e - f*x}])/f^3 - \\
 & (6*b*d^2*(b*d + a*c*f)*\text{PolyLog}[2, -E^{(-e - f*x)}])/f^3 - (6*a*b*d^3*x^2*\text{PolyLog}[2, -E^{(-e - f*x)}])/f^2 - \\
 & (6*b*c*d*(b*d + a*c*f)*\text{PolyLog}[2, E^{(-e - f*x)}])/f^3 - (6*a*b*d^3*x^2*\text{PolyLog}[2, E^{(-e - f*x)}])/f^2
 \end{aligned}$$

$$\begin{aligned} & *b*d^2*(b*d + 2*a*c*f)*x*PolyLog[2, E^{(-e - f*x)}]/f^3 - (6*a*b*d^3*x^2*PolyLog[2, E^{(-e - f*x)}])/f^2 - (6*b*d^2*(b*d + 2*a*c*f)*PolyLog[3, -E^{(-e - f*x)}])/f^4 - (12*a*b*d^3*x*PolyLog[3, -E^{(-e - f*x)}])/f^3 - (6*b*d^2*(b*d + 2*a*c*f)*PolyLog[3, E^{(-e - f*x)}])/f^4 - (12*a*b*d^3*x*PolyLog[3, E^{(-e - f*x)}])/f^3 - (12*a*b*d^3*PolyLog[4, -E^{(-e - f*x)}])/f^4 - (12*a*b*d^3*PolyLog[4, E^{(-e - f*x)}])/f^4 + (Csch[e]*Csch[e + f*x]*(-(a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[f*x]) + (a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Cosh[2*e + f*x] + 2*b*((4*b*(c + d*x)^3 + a*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3))*Sinh[f*x] + a*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)))/(8*f) \end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1424 vs. $2(261) = 522$.

Time = 0.52 (sec), antiderivative size = 1425, normalized size of antiderivative = 5.26

method	result	size
risch	Expression too large to display	1425

```
[In] int((d*x+c)^3*(a+b*coth(f*x+e))^2,x,method=_RETURNVERBOSE)
[Out] -12/f^2*b^2*d^2*c*e*x-6/f^2*b*a*c^2*d*e^2-4/f^3*b*d^3*a*e^3*x+8/f^3*b*d^2*c*a*e^3+6/f^3*b^2*d^3*polylog(2,exp(f*x+e))*x+3/f^2*b^2*d^3*ln(1+exp(f*x+e))*x^2+6/f^3*b^2*d^3*polylog(2,-exp(f*x+e))*x+2/f*b*a*c^3*ln(exp(f*x+e)-1)+2/f*b*a*c^3*ln(1+exp(f*x+e))+12/f^4*b*d^3*a*polylog(4,exp(f*x+e))+12/f^4*b*d^3*a*polylog(4,-exp(f*x+e))+6/f^3*b^2*d^2*c*polylog(2,exp(f*x+e))+6/f^3*b^2*d^2*c*polylog(2,-exp(f*x+e))+3/f^2*b^2*c^2*d*ln(exp(f*x+e)-1)+3/f^2*b^2*c^2*d*ln(1+exp(f*x+e))+3/f^4*b^2*e^2*d^3*ln(exp(f*x+e)-1)+3/f^2*b^2*d^3*ln(1-exp(f*x+e))*e^2+1/4/d*b^2*c^4+1/4*a^2/d*c^4+1/4*a^2*d^3*x^4-1/2*d^3*a*b*x^4+d^2*b^2*c*x^3+3/2*d*b^2*c^2*x^2+12/f^2*b^2*c*a*polylog(2,exp(f*x+e))*x+6/f*b*d^2*c*a*ln(1+exp(f*x+e))*x^2+6/f^2*b^2*d^2*c*a*polylog(2,-exp(f*x+e))*x+6/f*b*a*c^2*d*ln(1+exp(f*x+e))*x+12/f^2*b*d^2*c*a*polylog(2,-exp(f*x+e))*x-6/f^2*b*e*a*c^2*d*ln(1+exp(f*x+e))*x+12/f^2*b*d^2*c*a*polylog(2,-exp(f*x+e))*x-6/f^2*b*d^2*c*x^3+3/2*a^2*d*c^2*x^2+a^2*c^3*x^2/f*b*d^3*a*ln(1-exp(f*x+e))*x^3+6/f^2*b*d^3*a*polylog(2,exp(f*x+e))*x^2-12/f^3*b*d^3*a*polylog(3,exp(f*x+e))*x+2/f*b*d^3*a*ln(1+exp(f*x+e))*x^3+6/f^2*b*d^3*a*polylog(2,-exp(f*x+e))*x^2-12/f^3*b*d^3*a*polylog(3,-exp(f*x+e))*x+2/f^4*b*d^3*a*ln(1-exp(f*x+e))*e^3-12/f^3*b*d^2*c*a*polylog(3,exp(f*x+e))-12/f^3*b*d^2*c*a*polylog(3,-exp(f*x+e))+6/f^2*b*a*c^2*d*polylog(2,-exp(f*x+e))-2/f^4*b*e^3*d^3*a*ln(exp(f*x+e)-1)-6/f*b^2*d^2*c*x^2-6/f^3*b^2*d^2*c*e^2+6/f^3*b^2*d^3*e^2*x-3/f^4*b*d^3*a*e^4+6/f*b*d^2*c*a*ln(1-exp(f*x+e))*x^2-6/f^3*b*d^2*c*a*ln(1-exp(f*x+e))*e^2-2*d^2*a*b*c*x^3-3*d*a*b*c^2*x^2+2*a*b*c^3*x^6/f^3*b^2*c*d^2*c*ln(exp(f*x+e)-1)+6/f^2*b^2*d^2*c*ln(1-exp(f*x+e))*x+6/f^3*b^2*d^2*c*ln(1+exp(f*x+e))*x-6/f^4*b^2*d^3*po
```

```

lylog(3,exp(f*x+e))-6/f^4*b^2*d^3*polylog(3,-exp(f*x+e))-2/f*b^2*d^3*x^3+4/
f^4*b^2*d^3*e^3+1/4*d^3*b^2*x^4-4/f*b*a*c^3*ln(exp(f*x+e))-6/f^2*b^2*c^2*d*
ln(exp(f*x+e))-6/f^4*b^2*e^2*d^3*ln(exp(f*x+e))+12/f^2*b*d^2*c*a*e^2*x-12/f
*b*a*c^2*d*e*x+12/f^2*b*e*a*c^2*d*ln(exp(f*x+e))-12/f^3*b*e^2*d^2*c*a*ln(ex
p(f*x+e))-2/f*b^2*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)/(exp(2*f*x+2*e)-1)+4/
f^4*b*e^3*d^3*a*ln(exp(f*x+e))+12/f^3*b^2*e*d^2*c*ln(exp(f*x+e))+b^2*c^3*x+
1/2/d*a*b*c^4

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3239 vs. $2(259) = 518$.

Time = 0.32 (sec), antiderivative size = 3239, normalized size of antiderivative = 11.95

$$\int (c + dx)^3 (a + b \coth(e + fx))^2 dx = \text{Too large to display}$$

```

[In] integrate((d*x+c)^3*(a+b*cOTH(f*x+e))^2,x, algorithm="fricas")
[Out] -1/4*((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*(a^2 - 2*a*b + b^2)*c*d^2*f^4*x^3
+ 6*(a^2 - 2*a*b + b^2)*c^2*d*f^4*x^2 + 4*a*b*d^3*e^4 + 4*(a^2 - 2*a*b + b
^2)*c^3*f^4*x - 8*b^2*d^3*e^3 - 8*(2*a*b*c^3*e - b^2*c^3)*f^3 + 24*(a*b*c^2
*d^2 - b^2*c^2*d*e)*f^2 - ((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^
4 - 16*a*b*c^3*e*f^3 - 8*b^2*d^3*e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b
^2)*c*d^2*f^4)*x^3 + 24*(a*b*c^2*d^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f
^3 - (a^2 - 2*a*b + b^2)*c^2*d*f^4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*
e^2)*f - 4*(6*b^2*c^2*d*f^3 - (a^2 - 2*a*b + b^2)*c^3*f^4)*x)*cosh(f*x + e)
^2 - 2*((a^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^4 - 16*a*b*c^3*e*f^3
- 8*b^2*d^3*e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b^2)*c*d^2*f^4)*x^3 + 2
4*(a*b*c^2*d^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f^3 - (a^2 - 2*a*b + b
^2)*c^2*d*f^4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*e^2)*f - 4*(6*b^2*c^2
*d*f^3 - (a^2 - 2*a*b + b^2)*c^3*f^4)*x)*cosh(f*x + e)*sinh(f*x + e) - ((a
^2 - 2*a*b + b^2)*d^3*f^4*x^4 + 4*a*b*d^3*e^4 - 16*a*b*c^3*e*f^3 - 8*b^2*d^3
*e^3 - 4*(2*b^2*d^3*f^3 - (a^2 - 2*a*b + b^2)*c*d^2*f^4)*x^3 + 24*(a*b*c^2*
d^2 - b^2*c^2*d*e)*f^2 - 6*(4*b^2*c*d^2*f^3 - (a^2 - 2*a*b + b^2)*c^2*d*f
^4)*x^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*c*d^2*e^2)*f - 4*(6*b^2*c^2*d*f^3 - (a
^2 - 2*a*b + b^2)*c^3*f^4)*x)*sinh(f*x + e)^2 - 8*(2*a*b*c*d^2*e^3 - 3*b^2*
c*d^2*e^2)*f + 24*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f - (a*b*d^3
*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*c
osh(f*x + e)^2 - 2*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*
d^2*f^2 + b^2*d^3*f)*x)*cosh(f*x + e)*sinh(f*x + e) - (a*b*d^3*f^2*x^2 + a
*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*sinh(f*x + e
)^2 + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*dilog(cosh(f*x + e) + sinh(f*x + e))
+ 24*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f - (a*b*d^3*f^2*x^2 + a
*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*cosh(f*x + e
)^2 - 2*(a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 +

```

$$\begin{aligned}
& b^{2*d^3*f}*x)*cosh(f*x + e)*sinh(f*x + e) - (a*b*d^3*f^2*x^2 + a*b*c^2*d*f^2 \\
& + b^2*c*d^2*f + (2*a*b*c*d^2*f^2 + b^2*d^3*f)*x)*sinh(f*x + e)^2 + (2*a*b \\
& *c*d^2*f^2 + b^2*d^3*f)*x)*dilog(-cosh(f*x + e) - sinh(f*x + e)) + 4*(2*a*b \\
& *d^3*f^3*x^3 + 2*a*b*c^3*f^3 + 3*b^2*c^2*d*f^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d \\
& ^3*f^2)*x^2 - (2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 + 3*b^2*c^2*d*f^2 + 3*(2*a \\
& *b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*cosh \\
& (f*x + e)^2 - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 + 3*b^2*c^2*d*f^2 + 3*(2 \\
& *a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*co \\
& sh(f*x + e)*sinh(f*x + e) - (2*a*b*d^3*f^3*x^3 + 2*a*b*c^3*f^3 + 3*b^2*c^2*f \\
& d*f^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f \\
& ^2)*x)*sinh(f*x + e)^2 + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*log(cosh(f \\
& *x + e) + sinh(f*x + e) + 1) - 4*(2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*f^2 \\
& + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 - (2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - \\
& 3*b^2*d^3*e^2 + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2 \\
& *c*d^2*e)*f)*cosh(f*x + e)^2 - 2*(2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*f^2 \\
& + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)* \\
& cosh(f*x + e)*sinh(f*x + e) - (2*a*b*d^3*e^3 - 2*a*b*c^3*f^3 - 3*b^2*d^3*f^2 \\
& + 3*(2*a*b*c^2*d*e - b^2*c^2*d)*f^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)* \\
& sinh(f*x + e)^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f)*log(cosh(f*x + e) + \\
& sinh(f*x + e) - 1) + 4*(2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 \\
& - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - (2*a*b*d^3*f^3*x^3 \\
& + 2*a*b*d^3*e^3 + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2 \\
& *d^3*f^2)*x^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2 \\
& *c*d^2*f^2)*x)*cosh(f*x + e)^2 - 2*(2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 \\
& + 6*a*b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)* \\
& x^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)* \\
& x)*cosh(f*x + e)*sinh(f*x + e) - (2*a*b*d^3*f^3*x^3 + 2*a*b*d^3*e^3 + 6*a* \\
& b*c^2*d*e*f^2 - 3*b^2*d^3*e^2 + 3*(2*a*b*c*d^2*f^3 + b^2*d^3*f^2)*x^2 - 6*(\\
& a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)*x)*sinh \\
& (f*x + e)^2 - 6*(a*b*c*d^2*e^2 - b^2*c*d^2*e)*f + 6*(a*b*c^2*d*f^3 + b^2*c*d^2*f^2)* \\
& log(-cosh(f*x + e) - sinh(f*x + e) + 1) - 48*(a*b*d^3*cosh(f*x + e)^2 + 2*a*b*d^3*cosh(f*x + e) \\
& + 2*a*b*d^3*cosh(f*x + e)*sinh(f*x + e) + a*b*d^3*sinh(f*x + e)^2 - a*b*d^3)* \\
& polylog(4, cosh(f*x + e) + sinh(f*x + e)) - 48*(a*b*d^3*cosh(f*x + e)^2 + 2*a*b*d^3*cosh(f*x + e) \\
& + 2*a*b*d^3*cosh(f*x + e)*sinh(f*x + e) + a*b*d^3*sinh(f*x + e)^2 - a*b*d^3)* \\
& polylog(4, -cosh(f*x + e) - sinh(f*x + e)) - 24*(2*a*b*d^3*f*x + 2*a*b*c*d^2*f \\
& + b^2*d^3 - (2*a*b*d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(f*x + e)^2 - 2*(2*a*b \\
& *d^3*f*x + 2*a*b*c*d^2*f + b^2*d^3)*cosh(f*x + e)*sinh(f*x + e) - (2*a*b*d^3*f*x \\
& + 2*a*b*c*d^2*f + b^2*d^3)*sinh(f*x + e)^2)*polylog(3, -cosh(f*x + e) \\
& - sinh(f*x + e))/(f^4*cosh(f*x + e)^2 + 2*f^4*cosh(f*x + e)*sinh(f*x + e) \\
& + f^4*sinh(f*x + e)^2 - f^4)
\end{aligned}$$

Sympy [F]

$$\int (c + dx)^3 (a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + fx))^2 (c + dx)^3 dx$$

[In] `integrate((d*x+c)**3*(a+b*coth(f*x+e))**2, x)`
[Out] `Integral((a + b*coth(e + f*x))**2*(c + d*x)**3, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 781 vs. $2(259) = 518$.

Time = 0.28 (sec), antiderivative size = 781, normalized size of antiderivative = 2.88

$$\begin{aligned} \int (c + dx)^3 (a + b \coth(e + fx))^2 dx = & \frac{1}{4} a^2 d^3 x^4 + a^2 c d^2 x^3 + \frac{3}{2} a^2 c^2 d x^2 + a^2 c^3 x - \frac{6 b^2 c^2 d x}{f} \\ & + \frac{2 a b c^3 \log(\sinh(fx + e))}{f} + \frac{3 b^2 c^2 d \log(e^{(fx+e)} + 1)}{f^2} + \frac{3 b^2 c^2 d \log(e^{(fx+e)} - 1)}{f^2} \\ & + \frac{2 (f^3 x^3 \log(e^{(fx+e)} + 1) + 3 f^2 x^2 \text{Li}_2(-e^{(fx+e)}) - 6 f x \text{Li}_3(-e^{(fx+e)}) + 6 \text{Li}_4(-e^{(fx+e)})) a b d^3}{f^4} \\ & + \frac{2 (f^3 x^3 \log(-e^{(fx+e)} + 1) + 3 f^2 x^2 \text{Li}_2(e^{(fx+e)}) - 6 f x \text{Li}_3(e^{(fx+e)}) + 6 \text{Li}_4(e^{(fx+e)})) a b d^3}{f^4} \\ & - \frac{8 b^2 c^3 + (2 a b d^3 f + b^2 d^3 f) x^4 + 4 (c^3 f + 6 c^2 d) b^2 x + 4 (2 a b c d^2 f + (c d^2 f + 2 d^3) b^2) x^3 + 6 (2 a b c^2 d f + (c^3 d + 6 c^2 d^2) b) x^2}{f^4} \\ & + \frac{6 (a b c^2 d f + b^2 c d^2) (f x \log(e^{(fx+e)} + 1) + \text{Li}_2(-e^{(fx+e)}))}{f^3} \\ & + \frac{6 (a b c^2 d f + b^2 c d^2) (f x \log(-e^{(fx+e)} + 1) + \text{Li}_2(e^{(fx+e)}))}{f^3} \\ & + \frac{3 (2 a b c d^2 f + b^2 d^3) (f^2 x^2 \log(e^{(fx+e)} + 1) + 2 f x \text{Li}_2(-e^{(fx+e)}) - 2 \text{Li}_3(-e^{(fx+e)}))}{f^4} \\ & + \frac{3 (2 a b c d^2 f + b^2 d^3) (f^2 x^2 \log(-e^{(fx+e)} + 1) + 2 f x \text{Li}_2(e^{(fx+e)}) - 2 \text{Li}_3(e^{(fx+e)}))}{f^4} \\ & - \frac{a b d^3 f^4 x^4 + 2 (2 a b c d^2 f + b^2 d^3) f^3 x^3 + 6 (a b c^2 d f^2 + b^2 c d^2 f) f^2 x^2}{f^4} \end{aligned}$$

[In] `integrate((d*x+c)^3*(a+b*coth(f*x+e))^2, x, algorithm="maxima")`
[Out] `1/4*a^2*d^3*x^4 + a^2*c*d^2*x^3 + 3/2*a^2*c^2*d*x^2 + a^2*c^3*x - 6*b^2*c^2*d*x/f + 2*a*b*c^3*log(sinh(f*x + e))/f + 3*b^2*c^2*d*log(e^(f*x + e) + 1)/f^2 + 3*b^2*c^2*d*log(e^(f*x + e) - 1)/f^2 + 2*(f^3*x^3*log(e^(f*x + e) + 1) + 3*f^2*x^2*dilog(-e^(f*x + e))) - 6*f*x*polylog(3, -e^(f*x + e)) + 6*poly`

$$\begin{aligned}
& \log(4, -e^{-(f*x + e)})*a*b*d^3/f^4 + 2*(f^3*x^3*\log(-e^{-(f*x + e)} + 1) + 3*f^2*x^2*\text{dilog}(e^{-(f*x + e)}) - 6*f*x*\text{polylog}(3, e^{-(f*x + e)}) + 6*\text{polylog}(4, e^{-(f*x + e)}))*a*b*d^3/f^4 - 1/4*(8*b^2*c^3 + (2*a*b*d^3*f + b^2*d^3*f)*x^4 + 4*(c^3*f + 6*c^2*d)*b^2*x + 4*(2*a*b*c*d^2*f + (c*d^2*f + 2*d^3)*b^2)*x^3 + 6*(2*a*b*c^2*d*f + (c^2*d*f + 4*c*d^2)*b^2)*x^2 - (4*b^2*c^3*f*x^2*e^{(2*e)}) + (2*a*b*d^3*f*e^{(2*e)} + b^2*d^3*f*e^{(2*e)})*x^4 + 4*(2*a*b*c*d^2*f*e^{(2*e)} + b^2*c*d^2*f*e^{(2*e)})*x^3 + 6*(2*a*b*c^2*d*f*e^{(2*e)} + b^2*c^2*d*f*e^{(2*e)})*x^2)*e^{(2*f*x)}/(f*e^{(2*f*x + 2*e)} - f) + 6*(a*b*c^2*d*f + b^2*c*d^2)*(f*x*\log(e^{-(f*x + e)} + 1) + \text{dilog}(-e^{-(f*x + e)}))/f^3 + 6*(a*b*c^2*d*f + b^2*c*d^2)*(f*x*\log(-e^{-(f*x + e)} + 1) + \text{dilog}(e^{-(f*x + e)}))/f^3 + 3*(2*a*b*c*d^2*f + b^2*d^3)*(f^2*x^2*\log(e^{-(f*x + e)} + 1) + 2*f*x*\text{dilog}(-e^{-(f*x + e)}) - 2*\text{polylog}(3, -e^{-(f*x + e)}))/f^4 + 3*(2*a*b*c*d^2*f + b^2*d^3)*(f^2*x^2*\log(-e^{-(f*x + e)} + 1) + 2*f*x*\text{dilog}(e^{-(f*x + e)}) - 2*\text{polylog}(3, e^{-(f*x + e)}))/f^4 - (a*b*d^3*f^4*x^4 + 2*(2*a*b*c*d^2*f + b^2*d^3)*f^3*x^3 + 6*(a*b*c^2*d*f^2 + b^2*c*d^2*f)*f^2*x^2)/f^4
\end{aligned}$$

Giac [F]

$$\int (c + dx)^3 (a + b \coth(e + fx))^2 dx = \int (dx + c)^3 (b \coth(fx + e) + a)^2 dx$$

[In] `integrate((d*x+c)^3*(a+b*coth(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)^3*(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + fx))^2 (c + dx)^3 dx$$

[In] `int((a + b*coth(e + f*x))^2*(c + d*x)^3,x)`
[Out] `int((a + b*coth(e + f*x))^2*(c + d*x)^3, x)`

3.43 $\int (c + dx)^2 (a + b \coth(e + fx))^2 dx$

Optimal result	281
Rubi [A] (verified)	282
Mathematica [B] (verified)	285
Maple [B] (verified)	287
Fricas [B] (verification not implemented)	288
Sympy [F]	289
Maxima [B] (verification not implemented)	289
Giac [F]	290
Mupad [F(-1)]	290

Optimal result

Integrand size = 20, antiderivative size = 209

$$\begin{aligned} \int (c + dx)^2 (a + b \coth(e + fx))^2 dx = & -\frac{b^2(c + dx)^2}{f} + \frac{a^2(c + dx)^3}{3d} - \frac{2ab(c + dx)^3}{3d} \\ & + \frac{b^2(c + dx)^3}{3d} - \frac{b^2(c + dx)^2 \coth(e + fx)}{f} \\ & + \frac{2b^2d(c + dx) \log(1 - e^{2(e+fx)})}{f^2} \\ & + \frac{2ab(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{b^2d^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\ & + \frac{2abd(c + dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & - \frac{abd^2 \text{PolyLog}(3, e^{2(e+fx)})}{f^3} \end{aligned}$$

```
[Out] -b^2*(d*x+c)^2/f+1/3*a^2*(d*x+c)^3/d-2/3*a*b*(d*x+c)^3/d+1/3*b^2*(d*x+c)^3/d-b^2*(d*x+c)^2*coth(f*x+e)/f+2*b^2*d*(d*x+c)*ln(1-exp(2*f*x+2*e))/f^2+2*a*b*(d*x+c)^2*ln(1-exp(2*f*x+2*e))/f+b^2*d^2*polylog(2,exp(2*f*x+2*e))/f^3+2*a*b*d*(d*x+c)*polylog(2,exp(2*f*x+2*e))/f^2-a*b*d^2*polylog(3,exp(2*f*x+2*e))/f^3
```

Rubi [A] (verified)

Time = 0.29 (sec) , antiderivative size = 209, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3803, 3797, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32}

$$\begin{aligned} \int (c + dx)^2 (a + b \coth(e + fx))^2 dx = & \frac{a^2(c + dx)^3}{3d} + \frac{2abd(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & + \frac{2ab(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{2ab(c + dx)^3}{3d} - \frac{abd^2 \operatorname{PolyLog}(3, e^{2(e+fx)})}{f^3} \\ & + \frac{2b^2d(c + dx) \log(1 - e^{2(e+fx)})}{f^2} \\ & - \frac{b^2(c + dx)^2 \coth(e + fx)}{f} - \frac{b^2(c + dx)^2}{f} \\ & + \frac{b^2(c + dx)^3}{3d} + \frac{b^2d^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^3} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2 * (a + b*\operatorname{Coth}[e + f*x])^2, x]$

[Out] $-\left(\frac{(b^2(c + d*x)^2)/f}{3} + \frac{(a^2(c + d*x)^3)/(3*d)}{3} - \frac{(2*a*b*(c + d*x)^3)/(3*d)}{3}\right) + \left(\frac{(b^2(c + d*x)^3)/(3*d)}{3} - \frac{(b^2(c + d*x)^2*\operatorname{Coth}[e + f*x])/f}{3} + \frac{(2*b^2*d*(c + d*x)*\operatorname{Log}[1 - E^{(2*(e + f*x))}])/f^2}{3} + \frac{(2*a*b*(c + d*x)^2*\operatorname{Log}[1 - E^{(2*(e + f*x))}])/f}{3} + \frac{(b^2*d^2*\operatorname{PolyLog}[2, E^{(2*(e + f*x))}])/f^3}{3} + \frac{(2*a*b*d*(c + d*x)*\operatorname{PolyLog}[2, E^{(2*(e + f*x))}])/f^2}{3} - \frac{(a*b*d^2*\operatorname{PolyLog}[3, E^{(2*(e + f*x))}])/f^3}{3}\right)$

Rule 32

```
Int[((a_.) + (b_.)*(x_.))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.))), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)]], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))]
```

```
)^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_)[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_.) + (e_.)*(x_.)^n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_.)^((c_.)*((a_.) + (b_.)*(x_.))))^n_.]*((f_.) + (g_.)*(x_.)^m_.), x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_.))^m_*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_.*(x_.)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_.))^m_*((b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_.))^m_*((a_.) + (b_.)*tan[(e_.) + (f_.)*(x_.)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_.)*(x_))^p_.]/((d_.) + (e_.)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx)^2 + 2ab(c+dx)^2 \coth(e+fx) + b^2(c+dx)^2 \coth^2(e+fx)) \, dx \\
&= \frac{a^2(c+dx)^3}{3d} + (2ab) \int (c+dx)^2 \coth(e+fx) \, dx + b^2 \int (c+dx)^2 \coth^2(e+fx) \, dx \\
&= \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} - \frac{b^2(c+dx)^2 \coth(e+fx)}{f} \\
&\quad - (4ab) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} \, dx + b^2 \int (c+dx)^2 \, dx \\
&\quad + \frac{(2b^2d) \int (c+dx) \coth(e+fx) \, dx}{f} \\
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad - \frac{b^2(c+dx)^2 \coth(e+fx)}{f} + \frac{2ab(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad - \frac{(4abd) \int (c+dx) \log(1-e^{2(e+fx)}) \, dx}{f} - \frac{(4b^2d) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} \, dx}{f} \\
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad - \frac{b^2(c+dx)^2 \coth(e+fx)}{f} + \frac{2b^2d(c+dx) \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{2ab(c+dx)^2 \log(1-e^{2(e+fx)})}{f} + \frac{2abd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad - \frac{(2abd^2) \int \text{PolyLog}(2, e^{2(e+fx)}) \, dx}{f^2} - \frac{(2b^2d^2) \int \log(1-e^{2(e+fx)}) \, dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad - \frac{b^2(c+dx)^2 \coth(e+fx)}{f} + \frac{2b^2d(c+dx) \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{2ab(c+dx)^2 \log(1-e^{2(e+fx)})}{f} + \frac{2abd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad - \frac{(abd^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} dx, x, e^{2(e+fx)}\right)}{f^3} \\
&\quad - \frac{(b^2d^2) \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{f^3} \\
&= -\frac{b^2(c+dx)^2}{f} + \frac{a^2(c+dx)^3}{3d} - \frac{2ab(c+dx)^3}{3d} + \frac{b^2(c+dx)^3}{3d} \\
&\quad - \frac{b^2(c+dx)^2 \coth(e+fx)}{f} + \frac{2b^2d(c+dx) \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{2ab(c+dx)^2 \log(1-e^{2(e+fx)})}{f} + \frac{b^2d^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{2abd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{abd^2 \text{PolyLog}(3, e^{2(e+fx)})}{f^3}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 473 vs. $2(209) = 418$.

Time = 6.92 (sec) , antiderivative size = 473, normalized size of antiderivative = 2.26

$$\begin{aligned}
 \int (c + dx)^2 (a + b \coth(e + fx))^2 dx = & -\frac{4bc(bd + acf)x}{f} - \frac{4bc(bd + acf)x}{(-1 + e^{2e})f} \\
 & - \frac{2bd(bd + 2acf)x^2}{(-1 + e^{2e})f} + \frac{4abd^2x^3}{3 - 3e^{2e}} \\
 & + \frac{1}{3}x(3c^2 + 3cdx + d^2x^2)(a^2 + b^2 + 2ab \coth(e)) \\
 & + \frac{2bd(bd + 2acf)x \log(1 - e^{-e-fx})}{f^2} \\
 & + \frac{2abd^2x^2 \log(1 - e^{-e-fx})}{f} \\
 & + \frac{2bd(bd + 2acf)x \log(1 + e^{-e-fx})}{f^2} \\
 & + \frac{2abd^2x^2 \log(1 + e^{-e-fx})}{f} \\
 & + \frac{2bc(bd + acf) \log(1 - e^{e+fx})}{f^2} \\
 & + \frac{2bc(bd + acf) \log(1 + e^{e+fx})}{f^2} \\
 & - \frac{2bd(bd + 2acf) \text{PolyLog}(2, -e^{-e-fx})}{f^3} \\
 & - \frac{4abd^2x \text{PolyLog}(2, -e^{-e-fx})}{f^2} \\
 & - \frac{2bd(bd + 2acf) \text{PolyLog}(2, e^{-e-fx})}{f^3} \\
 & - \frac{4abd^2x \text{PolyLog}(2, e^{-e-fx})}{f^2} \\
 & - \frac{4abd^2 \text{PolyLog}(3, -e^{-e-fx})}{f^3} \\
 & - \frac{4abd^2 \text{PolyLog}(3, e^{-e-fx})}{f^3} \\
 & + \frac{b^2(c + dx)^2 \text{csch}(e) \text{csch}(e + fx) \sinh(fx)}{f}
 \end{aligned}$$

[In] Integrate[(c + d*x)^2*(a + b*Coth[e + f*x])^2, x]

[Out]
$$\begin{aligned}
 & (-4*b*c*(b*d + a*c*f)*x)/f - (4*b*c*(b*d + a*c*f)*x)/((-1 + E^{(2*e)})*f) - (\\
 & 2*b*d*(b*d + 2*a*c*f)*x^2)/((-1 + E^{(2*e)})*f) + (4*a*b*d^2*x^3)/(3 - 3*E^{(2} \\
 & *e)) + (x*(3*c^2 + 3*c*d*x + d^2*x^2)*(a^2 + b^2 + 2*a*b*Coth[e]))/3 + (2*b \\
 & *d*(b*d + 2*a*c*f)*x*Log[1 - E^{(-e - f*x)}])/f^2 + (2*a*b*d^2*x^2*Log[1 - E^
 \end{aligned}$$

$$\begin{aligned}
& (-e - f*x])/f + (2*b*d*(b*d + 2*a*c*f)*x*Log[1 + E^(-e - f*x)])/f^2 + (2*a*b*d^2*x^2*Log[1 + E^(-e - f*x)])/f + (2*b*c*(b*d + a*c*f)*Log[1 - E^(e + f*x)])/f^2 + (2*b*c*(b*d + a*c*f)*Log[1 + E^(e + f*x)])/f^2 - (2*b*d*(b*d + 2*a*c*f)*PolyLog[2, -E^(-e - f*x)])/f^3 - (4*a*b*d^2*x*PolyLog[2, -E^(-e - f*x)])/f^2 - (2*b*d*(b*d + 2*a*c*f)*PolyLog[2, E^(-e - f*x)])/f^3 - (4*a*b*d^2*x*PolyLog[2, E^(-e - f*x)])/f^2 - (4*a*b*d^2*PolyLog[3, -E^(-e - f*x)])/f^3 - (4*a*b*d^2*PolyLog[3, E^(-e - f*x)])/f^3 + (b^2*(c + d*x)^2*Csch[e]*Csch[e + f*x]*Sinh[f*x])/f
\end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 824 vs. $2(203) = 406$.

Time = 0.46 (sec), antiderivative size = 825, normalized size of antiderivative = 3.95

method	result
risch	$\frac{4bcda \operatorname{polylog}(2, e^{fx+e})}{f^2} + \frac{4bcda \operatorname{polylog}(2, -e^{fx+e})}{f^2} + \frac{2be^2 a d^2 \ln(e^{fx+e}-1)}{f^3} + \frac{2ba d^2 \ln(1-e^{fx+e})x^2}{f} - \frac{2ba d^2 \ln(1-e^{fx+e})}{f^3}$

```

[In] int((d*x+c)^2*(a+b*c*coth(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] 4/f^2*b*c*d*a*polylog(2,exp(f*x+e))+4/f^2*b*c*d*a*polylog(2,-exp(f*x+e))+2/
f^3*b*e^2*a*d^2*ln(exp(f*x+e)-1)+2/f*b*a*d^2*ln(1-exp(f*x+e))*x^2-2/f^3*b*a*
d^2*ln(1-exp(f*x+e))*e^2+4/f^2*b*a*d^2*e^2*x-4/f^2*b*c*d*a*e^2-4/f^3*b*e^2*
a*d^2*ln(exp(f*x+e))-4/f*b*a*c^2*ln(exp(f*x+e))-4/f^2*b^2*c*d*ln(exp(f*x+e))
+d*b^2*c*x^2+b^2*c^2*x+a^2*d*c*x^2+a^2*c^2*x+1/3*d^2*b^2*x^3+1/3/d*b^2*c^
3+1/3*a^2*d^2*x^3+1/3*a^2/d*c^3-2/f*b^2*d^2*x^2-2/f^3*b^2*d^2*e^2+2/f^2*b^2*
d^2*ln(1-exp(f*x+e))*x+2/f^3*b^2*d^2*ln(1-exp(f*x+e))*e+2/f^2*b^2*d^2*ln(1
+exp(f*x+e))*x+2/f*b*a*c^2*ln(exp(f*x+e)-1)+2/f*b*a*c^2*ln(1+exp(f*x+e))+2/
f^2*b^2*c*d*ln(exp(f*x+e)-1)+2/f^2*b^2*c*d*ln(1+exp(f*x+e))-4/f^3*b*a*d^2*p
olylog(3,exp(f*x+e))-4/f^3*b*a*d^2*polylog(3,-exp(f*x+e))-4/f^2*b^2*d^2*e*x-
2/f^3*b^2*e*d^2*ln(exp(f*x+e)-1)+4/f^2*b*a*d^2*polylog(2,exp(f*x+e))*x+2/f*
b*a*d^2*ln(1+exp(f*x+e))*x^2+4/f^2*b*a*d^2*polylog(2,-exp(f*x+e))*x+2/f^3*
b^2*d^2*polylog(2,exp(f*x+e))+2/f^3*b^2*d^2*polylog(2,-exp(f*x+e))+8/3/f^3*
b*a*d^2*e^3+4/f^3*b^2*e*d^2*ln(exp(f*x+e))-2/3*d^2*a*b*x^3+2/3/d*c^3*a*b+4/
f^2*b*c*d*a*ln(1-exp(f*x+e))*e-4/f^2*b*e*c*d*a*ln(exp(f*x+e)-1)-2/f*b^2*(d^
2*x^2+2*c*d*x+c^2)/(exp(2*f*x+2*e)-1)-2*d*a*b*c*x^2+2*a*b*c^2*x+4/f*b*c*d*a*
ln(1-exp(f*x+e))*x+4/f*b*c*d*a*ln(1+exp(f*x+e))*x-8/f*b*c*d*a*e*x+8/f^2*b*
e*c*d*a*ln(exp(f*x+e))

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. $1854 \text{ vs. } 2(201) = 402$.

Time = 0.29 (sec), antiderivative size = 1854, normalized size of antiderivative = 8.87

$$\int (c + dx)^2 (a + b \coth(e + fx))^2 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^2*(a+b*coth(f*x+e))^2,x, algorithm="fricas")

[Out] -1/3*((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 + 3*(a^2 - 2*a*b + b^2)*c*d*f^3*x^2 -
4*a*b*d^2*e^3 + 3*(a^2 - 2*a*b + b^2)*c^2*f^3*x + 6*b^2*d^2*e^2 - 6*(2*a*b
*c^2*e - b^2*c^2)*f^2 - ((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 - 4*a*b*d^2*e^3 -
12*a*b*c^2*e*f^2 + 6*b^2*d^2*e^2 - 3*(2*b^2*d^2*f^2 - (a^2 - 2*a*b + b^2)*c
*d*f^3)*x^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f - 3*(4*b^2*c*d*f^2 - (a^2 - 2*
a*b + b^2)*c^2*f^3)*x)*cosh(f*x + e)^2 - 2*((a^2 - 2*a*b + b^2)*d^2*f^3*x^3 -
4*a*b*d^2*e^3 - 12*a*b*c^2*e*f^2 + 6*b^2*d^2*e^2 - 3*(2*b^2*d^2*f^2 - (a
^2 - 2*a*b + b^2)*c*d*f^3)*x^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f - 3*(4*b^2*
c*d*f^2 - (a^2 - 2*a*b + b^2)*c^2*f^3)*x)*cosh(f*x + e)*sinh(f*x + e) - ((a
^2 - 2*a*b + b^2)*d^2*f^3*x^3 - 4*a*b*d^2*e^3 - 12*a*b*c^2*e*f^2 + 6*b^2*d^
2*e^2 - 3*(2*b^2*d^2*f^2 - (a^2 - 2*a*b + b^2)*c*d*f^3)*x^2 + 12*(a*b*c*d*e
^2 - b^2*c*d*e)*f - 3*(4*b^2*c*d*f^2 - (a^2 - 2*a*b + b^2)*c^2*f^3)*x)*sinh
(f*x + e)^2 + 12*(a*b*c*d*e^2 - b^2*c*d*e)*f + 6*(2*a*b*d^2*f*x + 2*a*b*c*d
*f + b^2*d^2 - (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*cosh(f*x + e)^2 - 2*
(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*cosh(f*x + e)*sinh(f*x + e) - (2*a*
b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2)*sinh(f*x + e)^2)*dilog(cosh(f*x + e) + s
inh(f*x + e)) + 6*(2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d^2 - (2*a*b*d^2*f*x +
2*a*b*c*d*f + b^2*d^2)*cosh(f*x + e)^2 - 2*(2*a*b*d^2*f*x + 2*a*b*c*d*f +
b^2*d^2)*cosh(f*x + e)*sinh(f*x + e) - (2*a*b*d^2*f*x + 2*a*b*c*d*f + b^2*d
^2)*sinh(f*x + e)^2)*dilog(-cosh(f*x + e) - sinh(f*x + e)) + 6*(a*b*d^2*f^2
*x^2 + a*b*c^2*f^2 + b^2*c*d*f - (a*b*d^2*f^2*x^2 + a*b*c^2*f^2 + b^2*c*d*f
+ (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*cosh(f*x + e)^2 - 2*(a*b*d^2*f^2*x^2 + a*
b*c^2*f^2 + b^2*c*d*f + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*cosh(f*x + e)*sinh(f
*x + e) - (a*b*d^2*f^2*x^2 + a*b*c^2*f^2 + b^2*c*d*f + (2*a*b*c*d*f^2 + b^2
*d^2*f)*x)*sinh(f*x + e)^2 + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*log(cosh(f*x +
e) + sinh(f*x + e) + 1) + 6*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (a*b*d
^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*cosh(f*x + e)
^2 - 2*(a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*b*c*d*e - b^2*c*d)*f)*
cosh(f*x + e)*sinh(f*x + e) - (a*b*d^2*e^2 + a*b*c^2*f^2 - b^2*d^2*e - (2*a*
b*c*d*e - b^2*c*d)*f)*sinh(f*x + e)^2 - (2*a*b*c*d*e - b^2*c*d)*f)*log(cos
h(f*x + e) + sinh(f*x + e) - 1) + 6*(a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d
*f + b^2*d^2*f - (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*f + b^2*d^2*f +
(2*a*b*c*d*f^2 + b^2*d^2*f)*x)*cosh(f*x + e)^2 - 2*(a*b*d^2*f^2*x^2 -
a*b*d^2*e^2 + 2*a*b*c*d*f + b^2*d^2*e + (2*a*b*c*d*f^2 + b^2*d^2*f)*x)*
cosh(f*x + e)*sinh(f*x + e) - (a*b*d^2*f^2*x^2 - a*b*d^2*e^2 + 2*a*b*c*d*f)
```

$$\begin{aligned} & *f + b^2 d^2 e + (2 a b c d f^2 + b^2 d^2 f^2) x \sinh(f x + e)^2 + (2 a b c d^2 f^2 + b^2 d^2 f^2) x \log(-\cosh(f x + e) - \sinh(f x + e) + 1) + 12 a b d^2 \\ & * \cosh(f x + e)^2 + 2 a b d^2 \cosh(f x + e) \sinh(f x + e) + a b d^2 \sinh(f x + e)^2 - a b d^2 \operatorname{polylog}(3, \cosh(f x + e) + \sinh(f x + e)) + 12 a b d^2 \\ & \cosh(f x + e)^2 + 2 a b d^2 \cosh(f x + e) \sinh(f x + e) + a b d^2 \sinh(f x + e)^2 - a b d^2 \operatorname{polylog}(3, -\cosh(f x + e) - \sinh(f x + e))) / (f^3 \cosh(f x + e)^2 + 2 f^3 \cosh(f x + e) \sinh(f x + e) + f^3 \sinh(f x + e)^2 - f^3) \end{aligned}$$

Sympy [F]

$$\int (c + dx)^2 (a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + fx))^2 (c + dx)^2 dx$$

[In] `integrate((d*x+c)**2*(a+b*cOTH(f*x+e))**2, x)`

[Out] `Integral((a + b*cOTH(e + f*x))**2*(c + d*x)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 494 vs. $2(201) = 402$.

Time = 0.27 (sec), antiderivative size = 494, normalized size of antiderivative = 2.36

$$\begin{aligned} & \int (c + dx)^2 (a + b \coth(e + fx))^2 dx = \frac{1}{3} a^2 d^2 x^3 + a^2 c d x^2 + a^2 c^2 x - \frac{4 b^2 c d x}{f} \\ & + \frac{2 a b c^2 \log(\sinh(f x + e))}{f} + \frac{2 b^2 c d \log(e^{(fx+e)} + 1)}{f^2} + \frac{2 b^2 c d \log(e^{(fx+e)} - 1)}{f^2} \\ & + \frac{2 (f^2 x^2 \log(e^{(fx+e)} + 1) + 2 f x \text{Li}_2(-e^{(fx+e)}) - 2 \text{Li}_3(-e^{(fx+e)})) a b d^2}{f^3} \\ & + \frac{2 (f^2 x^2 \log(-e^{(fx+e)} + 1) + 2 f x \text{Li}_2(e^{(fx+e)}) - 2 \text{Li}_3(e^{(fx+e)})) a b d^2}{f^3} \\ & - \frac{6 b^2 c^2 + 3 (c^2 f + 4 c d) b^2 x + (2 a b d^2 f + b^2 d^2 f) x^3 + 3 (2 a b c d f + (c d f + 2 d^2) b^2) x^2 - (3 b^2 c^2 f x e^{(2e)} + (2 a b c d f + b^2 d^2) f x^3 + 3 (2 a b c d f + b^2 d^2) f^2 x^2)}{3 (f e^{(2fx+2e)} - f)} \\ & + \frac{2 (2 a b c d f + b^2 d^2) (f x \log(e^{(fx+e)} + 1) + \text{Li}_2(-e^{(fx+e)}))}{f^3} \\ & + \frac{2 (2 a b c d f + b^2 d^2) (f x \log(-e^{(fx+e)} + 1) + \text{Li}_2(e^{(fx+e)}))}{f^3} \\ & - \frac{2 (2 a b d^2 f^3 x^3 + 3 (2 a b c d f + b^2 d^2) f^2 x^2)}{3 f^3} \end{aligned}$$

[In] `integrate((d*x+c)^2*(a+b*cOTH(f*x+e))^2, x, algorithm="maxima")`

[Out]
$$\begin{aligned} & 1/3*a^2*d^2*x^3 + a^2*c*d*x^2 + a^2*c^2*x - 4*b^2*c*d*x/f + 2*a*b*c^2*log(sinh(f*x + e))/f + 2*b^2*c*d*log(e^(f*x + e) + 1)/f^2 + 2*b^2*c*d*log(e^(f*x + e) - 1)/f^2 + 2*(f^2*x^2*log(e^(f*x + e) + 1) + 2*f*x*dilog(-e^(f*x + e))) - 2*polylog(3, -e^(f*x + e)))*a*b*d^2/f^3 + 2*(f^2*x^2*log(-e^(f*x + e) + 1) + 2*f*x*dilog(e^(f*x + e))) - 2*polylog(3, e^(f*x + e)))*a*b*d^2/f^3 - 1/3*(6*b^2*c^2 + 3*(c^2*f + 4*c*d)*b^2*x + (2*a*b*d^2*f + b^2*d^2*f)*x^3 + 3*(2*a*b*c*d*f + (c*d*f + 2*d^2)*b^2)*x^2 - (3*b^2*c^2*f*x*e^(2*e) + (2*a*b*d^2*f*e^(2*e) + b^2*d^2*f*e^(2*e))*x^3 + 3*(2*a*b*c*d*f*e^(2*e) + b^2*c*d*f*e^(2*e))*x^2)*e^(2*f*x))/(f*e^(2*f*x + 2*e) - f) + 2*(2*a*b*c*d*f + b^2*d^2)*(f*x*log(e^(f*x + e) + 1) + dilog(-e^(f*x + e)))/f^3 + 2*(2*a*b*c*d*f + b^2*d^2)*(f*x*log(-e^(f*x + e) + 1) + dilog(e^(f*x + e)))/f^3 - 2/3*(2*a*b*d^2*f^3*x^3 + 3*(2*a*b*c*d*f + b^2*d^2)*f^2*x^2)/f^3 \end{aligned}$$

Giac [F]

$$\int (c + dx)^2 (a + b \coth(e + fx))^2 dx = \int (dx + c)^2 (b \coth(fx + e) + a)^2 dx$$

[In] `integrate((d*x+c)^2*(a+b*coth(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((d*x + c)^2*(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + fx))^2 (c + dx)^2 dx$$

[In] `int((a + b*coth(e + f*x))^2*(c + d*x)^2,x)`

[Out] `int((a + b*coth(e + f*x))^2*(c + d*x)^2, x)`

3.44 $\int (c + dx)(a + b \coth(e + fx))^2 dx$

Optimal result	291
Rubi [A] (verified)	291
Mathematica [A] (verified)	294
Maple [B] (verified)	294
Fricas [B] (verification not implemented)	295
Sympy [F]	295
Maxima [A] (verification not implemented)	296
Giac [F]	296
Mupad [F(-1)]	297

Optimal result

Integrand size = 18, antiderivative size = 127

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx))^2 dx = & b^2 cx + \frac{1}{2} b^2 dx^2 + \frac{a^2(c + dx)^2}{2d} \\ & - \frac{ab(c + dx)^2}{d} - \frac{b^2(c + dx) \coth(e + fx)}{f} \\ & + \frac{2ab(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & + \frac{b^2 d \log(\sinh(e + fx))}{f^2} + \frac{abd \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \end{aligned}$$

[Out] $b^{2*c*x+1/2}*b^{2*d*x^2+1/2}*a^{2*(d*x+c)^2/d}-a*b*(d*x+c)^2/d-b^{2*(d*x+c)*\coth(f*x+e)/f+2*a*b*(d*x+c)*\ln(1-\exp(2*f*x+2*e))/f+b^{2*d*\ln(\sinh(f*x+e))/f^2+a*b*d*\operatorname{polylog}(2,\exp(2*f*x+2*e))/f^2}$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 127, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, $\frac{\text{number of rules}}{\text{integrand size}} = 0.389$, Rules used = {3803, 3797, 2221, 2317, 2438, 3801, 3556}

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx))^2 dx = & \frac{a^2(c + dx)^2}{2d} + \frac{2ab(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{ab(c + dx)^2}{d} + \frac{abd \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & - \frac{b^2(c + dx) \coth(e + fx)}{f} + b^2 cx \\ & + \frac{b^2 d \log(\sinh(e + fx))}{f^2} + \frac{1}{2} b^2 dx^2 \end{aligned}$$

[In] $\text{Int}[(c + d*x)*(a + b*\text{Coth}[e + f*x])^2, x]$

[Out] $b^{2*c*x} + (b^{2*d*x^2}/2 + (a^{2*(c + d*x)^2}/(2*d) - (a*b*(c + d*x)^2)/d - (b^{2*(c + d*x)}*\text{Coth}[e + f*x])/f + (2*a*b*(c + d*x)*\text{Log}[1 - E^{2*(e + f*x)}]))/f + (b^{2*d*\text{Log}[\text{Sinh}[e + f*x]]}/f^2 + (a*b*d*\text{PolyLog}[2, E^{2*(e + f*x)}])/f^2$

Rule 2221

$\text{Int}[((F_.)^((g_.)*(e_.) + (f_.)*(x_.)))^{(n_.)*(c_.) + (d_.)*(x_.)^(m_.)})/((a_.) + (b_.)*(F_.)^((g_.)*(e_.) + (f_.)*(x_.)))^{(n_.)}], \text{x_Symbol}] \rightarrow \text{Simp}[(c + d*x)^m/(b*f*g*n*\text{Log}[F])*(\text{Log}[1 + b*((F^{(g*(e + f*x))})^n/a)]), x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + d*x)^(m - 1)*\text{Log}[1 + b*((F^{(g*(e + f*x))})^n/a)], x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$

Rule 2317

$\text{Int}[\text{Log}[(a_.) + (b_.)*(F_.)^((e_.)*(c_.) + (d_.)*(x_.)))^{(n_.)}], \text{x_Symbol}] \rightarrow \text{Dist}[1/(d*e*n*\text{Log}[F]), \text{Subst}[\text{Int}[\text{Log}[a + b*x]/x, x], x, (F^{(e*(c + d*x))})^n], x] /; \text{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \text{GtQ}[a, 0]$

Rule 2438

$\text{Int}[\text{Log}[(c_.)*(d_.) + (e_.)*(x_.)^{(n_.)})]/(x_), \text{x_Symbol}] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; \text{FreeQ}[\{c, d, e, n\}, x] \&& \text{EqQ}[c*d, 1]$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], \text{x_Symbol}] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x]], x]]/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3797

$\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*\tan[(e_.) + \text{Pi}*(k_.) + (\text{Complex}[0, fz_])*(f_.)*(x_.)], \text{x_Symbol}] \rightarrow \text{Simp}[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + \text{Dist}[2*I, \text{Int}[(c + d*x)^m*(E^{(2*(-I)*e + f*fz*x)})/(1 + E^{(2*(-I)*e + f*fz*x)})/E^{(2*I*k*Pi)}], x] /; \text{FreeQ}[\{c, d, e, f, fz\}, x] \&& \text{IntegerQ}[4*k] \&& \text{IGtQ}[m, 0]$

Rule 3801

$\text{Int}[(c_.) + (d_.)*(x_.)^{(m_.)}*((b_.)*\tan[(e_.) + (f_.)*(x_.)])^{(n_.)}, \text{x_Symbol}] \rightarrow \text{Simp}[b*(c + d*x)^m*((b*\text{Tan}[e + f*x])^{(n - 1)}/(f*(n - 1))), x] + (-\text{Dist}[b*d*(m/(f*(n - 1))), \text{Int}[(c + d*x)^(m - 1)*(b*\text{Tan}[e + f*x])^{(n - 1)}, x], x] - \text{Dist}[b^2, \text{Int}[(c + d*x)^m*(b*\text{Tan}[e + f*x])^{(n - 2)}, x], x]) /; \text{FreeQ}[\{b, c, d, e, f\}, x] \&& \text{GtQ}[n, 1] \&& \text{GtQ}[m, 0]$

Rule 3803

```
Int[((c_) + (d_)*(x_))^m_*((a_) + (b_)*tan[(e_) + (f_)*(x_)])^n_, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^2(c+dx) + 2ab(c+dx)\coth(e+fx) + b^2(c+dx)\coth^2(e+fx)) \, dx \\
&= \frac{a^2(c+dx)^2}{2d} + (2ab) \int (c+dx)\coth(e+fx) \, dx + b^2 \int (c+dx)\coth^2(e+fx) \, dx \\
&= \frac{a^2(c+dx)^2}{2d} - \frac{ab(c+dx)^2}{d} - \frac{b^2(c+dx)\coth(e+fx)}{f} \\
&\quad - (4ab) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} \, dx + b^2 \int (c+dx) \, dx + \frac{(b^2d) \int \coth(e+fx) \, dx}{f} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} - \frac{ab(c+dx)^2}{d} \\
&\quad - \frac{b^2(c+dx)\coth(e+fx)}{f} + \frac{2ab(c+dx)\log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\sinh(e+fx))}{f^2} - \frac{(2abd) \int \log(1-e^{2(e+fx)}) \, dx}{f} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} - \frac{ab(c+dx)^2}{d} \\
&\quad - \frac{b^2(c+dx)\coth(e+fx)}{f} + \frac{2ab(c+dx)\log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\sinh(e+fx))}{f^2} - \frac{(abd) \text{Subst}\left(\int \frac{\log(1-x)}{x} \, dx, x, e^{2(e+fx)}\right)}{f^2} \\
&= b^2cx + \frac{1}{2}b^2dx^2 + \frac{a^2(c+dx)^2}{2d} - \frac{ab(c+dx)^2}{d} \\
&\quad - \frac{b^2(c+dx)\coth(e+fx)}{f} + \frac{2ab(c+dx)\log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^2d\log(\sinh(e+fx))}{f^2} + \frac{abd \text{PolyLog}(2, e^{2(e+fx)})}{f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.22 (sec) , antiderivative size = 218, normalized size of antiderivative = 1.72

$$\int (c + dx)(a + b \coth(e + fx))^2 dx \\ = \frac{(a + b \coth(e + fx))^2 \sinh(e + fx) \left(-2b^2 f(c + dx) \cosh(e + fx) - (a^2 + b^2)(e + fx)(-2cf + d(e - fx)) \right)}{2}$$

[In] `Integrate[(c + d*x)*(a + b*Coth[e + f*x])^2, x]`

[Out] $((a + b \operatorname{Coth}[e + f x])^2 \operatorname{Sinh}[e + f x] (-2 b^2 f (c + d x) \cosh[e + f x] - (a^2 + b^2) (e + f x) (-2 c f + d (e - f x))) \operatorname{Sinh}[e + f x] + 2 b ((a f^2 (c + d x)^2)/d + b d (e + f x) + (b d + 2 a f (c + d x)) \operatorname{Log}[1 - E^{(-e - f x)}] + (b d + 2 a f (c + d x)) \operatorname{Log}[1 + E^{(-e - f x)}] - 2 a d \operatorname{PolyLog}[2, -E^{(-e - f x)}] - 2 a d \operatorname{PolyLog}[2, E^{(-e - f x)}]) \operatorname{Sinh}[e + f x])/((2 f^2 (b \cosh[e + f x] + a \operatorname{Sinh}[e + f x]))^2)$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 317 vs. 2(123) = 246.

Time = 0.39 (sec) , antiderivative size = 318, normalized size of antiderivative = 2.50

method	result
risch	$\frac{a^2 d x^2}{2} - ab d x^2 + \frac{b^2 d x^2}{2} + a^2 c x + 2 a b c x + b^2 c x - \frac{2 (d x + c) b^2}{f (e^{2 f x + 2 e} - 1)} + \frac{b^2 d \ln(e^{f x + e} - 1)}{f^2} + \frac{b^2 d \ln(1 + e^{f x + e})}{f^2} - \frac{2 b^2 d}{f^2}$

[In] `int((d*x+c)*(a+b*cOTH(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] $1/2*a^2*d*x^2-a*b*d*x^2+1/2*b^2*d*x^2+a^2*c*x+2*a*b*c*x+b^2*c*x-2/f*(d*x+c)*b^2/(\exp(2*f*x+2*e)-1)+1/f^2*b^2*d*\ln(\exp(f*x+e)-1)+1/f^2*b^2*d*\ln(1+\exp(f*x+e))-2/f^2*b^2*d*\ln(\exp(f*x+e))+2/f*b*a*c*\ln(\exp(f*x+e)-1)+2/f*b*a*c*\ln(1+\exp(f*x+e))-4/f*b*a*c*\ln(\exp(f*x+e))-2/f^2*b*e*d*a*\ln(\exp(f*x+e)-1)+4/f^2*b*e*d*a*\ln(\exp(f*x+e))-4/f*b*d*a*e*x-2/f^2*b*d*a*e^2+2/f*b*d*a*\ln(1-\exp(f*x+e))*x+2/f^2*b*d*a*\ln(1-\exp(f*x+e))*e+2/f^2*b*d*a*polylog(2,\exp(f*x+e))+2/f*b*d*a*\ln(1+\exp(f*x+e))*x+2/f^2*b*d*a*polylog(2,-\exp(f*x+e))$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 851 vs. $2(122) = 244$.

Time = 0.26 (sec) , antiderivative size = 851, normalized size of antiderivative = 6.70

$$\int (c + dx)(a + b \coth(e + fx))^2 dx =$$

$$\frac{(a^2 - 2ab + b^2)df^2x^2 + 4abde^2 + 2(a^2 - 2ab + b^2)cf^2x - 4b^2de - ((a^2 - 2ab + b^2)df^2x^2 + 4abde^2 -$$

```
[In] integrate((d*x+c)*(a+b*cOTH(f*x+e))**2,x, algorithm="fricas")
[Out] -1/2*((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 + 2*(a^2 - 2*a*b + b^2)*c
*f^2*x - 4*b^2*d*e - ((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c
*e*f - 4*b^2*d*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*cOSH(f*x +
e)^2 - 2*((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c*e*f - 4*b^2
*d*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*cOSH(f*x + e)*sinH(f*x
+ e) - ((a^2 - 2*a*b + b^2)*d*f^2*x^2 + 4*a*b*d*e^2 - 8*a*b*c*e*f - 4*b^2*d
*e - 2*(2*b^2*d*f - (a^2 - 2*a*b + b^2)*c*f^2)*x)*sinH(f*x + e)^2 - 4*(2*a*
b*c*e - b^2*c)*f - 4*(a*b*d*cOSH(f*x + e))^2 + 2*a*b*d*cOSH(f*x + e)*sinH(f*
x + e) + a*b*d*sinH(f*x + e)^2 - a*b*d*dilog(cOSH(f*x + e) + sinH(f*x + e)
) - 4*(a*b*d*cOSH(f*x + e))^2 + 2*a*b*d*cOSH(f*x + e)*sinH(f*x + e) + a*b*d*
sinH(f*x + e)^2 - a*b*d*dilog(-cOSH(f*x + e) - sinH(f*x + e)) + 2*(2*a*b*d
*f*x + 2*a*b*c*f + b^2*d - (2*a*b*d*f*x + 2*a*b*c*f + b^2*d)*cOSH(f*x + e)^
2 - 2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)*cOSH(f*x + e)*sinH(f*x + e) - (2*a*
b*d*f*x + 2*a*b*c*f + b^2*d)*sinH(f*x + e)^2)*log(cOSH(f*x + e) + sinH(f*x
+ e) + 1) - 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d - (2*a*b*d*e - 2*a*b*c*f - b^2
*d)*cOSH(f*x + e)^2 - 2*(2*a*b*d*e - 2*a*b*c*f - b^2*d)*cOSH(f*x + e)*sinH(
f*x + e) - (2*a*b*d*e - 2*a*b*c*f - b^2*d)*sinH(f*x + e)^2)*log(cOSH(f*x +
e) + sinH(f*x + e) - 1) + 4*(a*b*d*f*x + a*b*d*e - (a*b*d*f*x + a*b*d*e)*c
OSH(f*x + e)^2 - 2*(a*b*d*f*x + a*b*d*e)*cOSH(f*x + e)*sinH(f*x + e) - (a*b*
d*f*x + a*b*d*e)*sinH(f*x + e)^2)*log(-cOSH(f*x + e) - sinH(f*x + e) + 1))/(
f^2*cOSH(f*x + e)^2 + 2*f^2*cOSH(f*x + e)*sinH(f*x + e) + f^2*sinH(f*x + e)
)^2 - f^2)
```

Sympy [F]

$$\int (c + dx)(a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + fx))^2 (c + dx) dx$$

```
[In] integrate((d*x+c)*(a+b*cOTH(f*x+e))**2,x)
```

```
[Out] Integral((a + b*cOTH(e + f*x))**2*(c + d*x), x)
```

Maxima [A] (verification not implemented)

none

Time = 0.26 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.92

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx))^2 dx &= \frac{1}{2} a^2 dx^2 - 2 abdx^2 + a^2 cx - \frac{2 b^2 dx}{f} \\ &+ \frac{2 abc \log(\sinh(fx + e))}{f} + \frac{2 (fx \log(e^{(fx+e)} + 1) + \text{Li}_2(-e^{(fx+e)}))abd}{f^2} \\ &+ \frac{2 (fx \log(-e^{(fx+e)} + 1) + \text{Li}_2(e^{(fx+e)}))abd}{f^2} \\ &+ \frac{b^2 d \log(e^{(fx+e)} + 1)}{f^2} + \frac{b^2 d \log(e^{(fx+e)} - 1)}{f^2} \\ &- \frac{2 (cf + 2d)b^2 x + 4b^2 c + (2abdf + b^2 df)x^2 - (2b^2 c f x e^{(2e)} + (2abdf e^{(2e)} + b^2 df e^{(2e)})x^2)e^{(2fx)}}{2(fe^{(2fx+2e)} - f)} \end{aligned}$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e))^2,x, algorithm="maxima")`

[Out] $\frac{1}{2}a^2 d x^2 - 2 a b d x^2 + a^2 c x - \frac{2 b^2 d x}{f} + 2 a b c \log(\sinh(f x + e))/f + 2 (f x \log(e^{(f x + e)} + 1) + \text{dilog}(-e^{(f x + e)})) * a b d / f^2 + 2 (f x \log(-e^{(f x + e)} + 1) + \text{dilog}(e^{(f x + e)})) * a b d / f^2 + b^2 d \log(e^{(f x + e)} + 1)/f^2 + b^2 d \log(e^{(f x + e)} - 1)/f^2 - \frac{1}{2} (2 (c f + 2 d) * b^2 x + 4 b^2 c + (2 a b d f + b^2 d f) * x^2 - (2 b^2 c f x e^{(2e)} + (2 a b d f e^{(2e)} + b^2 d f e^{(2e)}) * x^2) * e^{(2 f x)}) / (f * e^{(2 f x + 2 e)} - f)$

Giac [F]

$$\int (c + dx)(a + b \coth(e + fx))^2 dx = \int (dx + c)(b \coth(fx + e) + a)^2 dx$$

[In] `integrate((d*x+c)*(a+b*coth(f*x+e))^2,x, algorithm="giac")`

[Out] `integrate((d*x + c)*(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \coth(e + fx))^2 dx = \int (a + b \coth(e + f x))^2 (c + d x) dx$$

[In] `int((a + b*coth(e + f*x))^2*(c + d*x),x)`

[Out] `int((a + b*coth(e + f*x))^2*(c + d*x), x)`

3.45 $\int \frac{(a+b \coth(e+fx))^2}{c+dx} dx$

Optimal result	298
Rubi [N/A]	298
Mathematica [N/A]	299
Maple [N/A] (verified)	299
Fricas [N/A]	299
Sympy [N/A]	299
Maxima [N/A]	300
Giac [N/A]	300
Mupad [N/A]	300

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \text{Int}\left(\frac{(a + b \coth(e + fx))^2}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*cOTH(f*x+e))^2/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^2}{c + dx} dx$$

[In] Int[(a + b*COTH[e + f*x])^2/(c + d*x),x]

[Out] Defer[Int][(a + b*COTH[e + f*x])^2/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \coth(e + fx))^2}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 42.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^2}{c + dx} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])^2/(c + d*x), x]`

[Out] `Integrate[(a + b*Coth[e + f*x])^2/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.17 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \coth(fx + e))^2}{dx + c} dx$$

[In] `int((a+b*coth(f*x+e))^2/(d*x+c), x)`

[Out] `int((a+b*coth(f*x+e))^2/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 36, normalized size of antiderivative = 1.80

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b^2*coth(f*x + e)^2 + 2*a*b*coth(f*x + e) + a^2)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 1.49 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^2}{c + dx} dx$$

[In] `integrate((a+b*coth(f*x+e))**2/(d*x+c), x)`

[Out] `Integral((a + b*coth(e + f*x))**2/(c + d*x), x)`

Maxima [N/A]

Not integrable

Time = 0.40 (sec) , antiderivative size = 226, normalized size of antiderivative = 11.30

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c),x, algorithm="maxima")`

[Out] $a^2 \log(d*x + c)/d + 2*b^2/(d*f*x + c*f - (d*f*x*e^(2*e) + c*f*e^(2*e))*e^(2*f*x)) + (2*a*b + b^2)*\log(d*x + c)/d - \text{integrate}((2*a*b*d*f*x + 2*a*b*c*f - b^2*d)/(d^2*f*x^2 + 2*c*d*f*x + c^2*f + (d^2*f*x^2*e^e + 2*c*d*f*x*e^e + c^2*f*e^e)*e^(f*x)), x) + \text{integrate}(-(2*a*b*d*f*x + 2*a*b*c*f - b^2*d)/(d^2*f*x^2 + 2*c*d*f*x + c^2*f - (d^2*f*x^2*e^e + 2*c*d*f*x*e^e + c^2*f*e^e)*e^(f*x)), x)$

Giac [N/A]

Not integrable

Time = 0.32 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^2}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c),x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)^2/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 2.20 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^2}{c + dx} dx = \int \frac{(a + b \coth(e + f x))^2}{c + dx} dx$$

[In] `int((a + b*coth(e + f*x))^2/(c + d*x),x)`

[Out] `int((a + b*coth(e + f*x))^2/(c + d*x), x)`

3.46 $\int \frac{(a+b \coth(e+fx))^2}{(c+dx)^2} dx$

Optimal result	301
Rubi [N/A]	301
Mathematica [N/A]	302
Maple [N/A] (verified)	302
Fricas [N/A]	302
Sympy [N/A]	303
Maxima [N/A]	303
Giac [N/A]	303
Mupad [N/A]	304

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \coth(e + fx))^2}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*coth(f*x+e))^2/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx$$

[In] Int[(a + b*Coth[e + f*x])^2/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Coth[e + f*x])^2/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 34.10 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])^2/(c + d*x)^2, x]`

[Out] `Integrate[(a + b*Coth[e + f*x])^2/(c + d*x)^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.18 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \coth(fx + e))^2}{(dx + c)^2} dx$$

[In] `int((a+b*coth(f*x+e))^2/(d*x+c)^2, x)`

[Out] `int((a+b*coth(f*x+e))^2/(d*x+c)^2, x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 47, normalized size of antiderivative = 2.35

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c)^2, x, algorithm="fricas")`

[Out] `integral((b^2*coth(f*x + e)^2 + 2*a*b*coth(f*x + e) + a^2)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 2.49 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))**2/(d*x+c)**2, x)`

[Out] `Integral((a + b*coth(e + f*x))**2/(c + d*x)**2, x)`

Maxima [N/A]

Not integrable

Time = 0.45 (sec) , antiderivative size = 375, normalized size of antiderivative = 18.75

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c)^2, x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -a^2/(d^2*x + c*d) - (2*a*b*c*f + (c*f - 2*d)*b^2 + (2*a*b*d*f + b^2*d*f)*x \\ & - (2*a*b*c*f*e^(2*e) + b^2*c*f*e^(2*e) + (2*a*b*d*f*e^(2*e) + b^2*d*f*e^(2*e))*x)*e^(2*f*x))/(d^3*f*x^2 + 2*c*d^2*f*x + c^2*d*f - (d^3*f*x^2*e^(2*e) + 2*c*d^2*f*x*e^(2*e) + c^2*d*f*e^(2*e))*e^(2*f*x)) - \text{integrate}(2*(a*b*d*f*x + a*b*c*f - b^2*d)/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f + (d^3*f*x^3*e^(2*f*x) + 3*c*d^2*f*x^2*e^(2*f*x) + 3*c^2*d*f*x*e^(2*f*x) + c^3*f*e^(2*f*x)))*e^(2*f*x)), x) \\ & + \text{integrate}(-2*(a*b*d*f*x + a*b*c*f - b^2*d)/(d^3*f*x^3 + 3*c*d^2*f*x^2 + 3*c^2*d*f*x + c^3*f - (d^3*f*x^3*e^(2*f*x) + 3*c*d^2*f*x^2*e^(2*f*x) + 3*c^2*d*f*x*e^(2*f*x) + c^3*f*e^(2*f*x))), x) \end{aligned}$$

Giac [N/A]

Not integrable

Time = 0.46 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^2}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^2}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))^2/(d*x+c)^2, x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)^2/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 2.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + f x))^2}{(c + d x)^2} dx = \int \frac{(a + b \coth(e + f x))^2}{(c + d x)^2} dx$$

[In] int((a + b*coth(e + f*x))^2/(c + d*x)^2,x)

[Out] int((a + b*coth(e + f*x))^2/(c + d*x)^2, x)

3.47 $\int (c + dx)^3 (a + b \coth(e + fx))^3 dx$

Optimal result	306
Rubi [A] (verified)	307
Mathematica [B] (verified)	315
Maple [B] (verified)	316
Fricas [B] (verification not implemented)	318
Sympy [F]	318
Maxima [B] (verification not implemented)	318
Giac [F]	319
Mupad [F(-1)]	319

Optimal result

Integrand size = 20, antiderivative size = 556

$$\begin{aligned}
\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = & -\frac{3b^3 d(c + dx)^2}{2f^2} - \frac{3ab^2(c + dx)^3}{f} + \frac{b^3(c + dx)^3}{2f} \\
& + \frac{a^3(c + dx)^4}{4d} - \frac{3a^2b(c + dx)^4}{4d} + \frac{3ab^2(c + dx)^4}{4d} \\
& - \frac{b^3(c + dx)^4}{4d} - \frac{3b^3d(c + dx)^2 \coth(e + fx)}{2f^2} \\
& - \frac{3ab^2(c + dx)^3 \coth(e + fx)}{f} \\
& - \frac{b^3(c + dx)^3 \coth^2(e + fx)}{2f} \\
& + \frac{3b^3d^2(c + dx) \log(1 - e^{2(e+fx)})}{f^3} \\
& + \frac{9ab^2d(c + dx)^2 \log(1 - e^{2(e+fx)})}{f^2} \\
& + \frac{3a^2b(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{b^3(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{3b^3d^3 \text{PolyLog}(2, e^{2(e+fx)})}{2f^4} \\
& + \frac{9ab^2d^2(c + dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
& + \frac{9a^2bd(c + dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
& + \frac{3b^3d(c + dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
& - \frac{9ab^2d^3 \text{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\
& - \frac{9a^2bd^2(c + dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
& - \frac{3b^3d^2(c + dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
& + \frac{9a^2bd^3 \text{PolyLog}(4, e^{2(e+fx)})}{4f^4} \\
& + \frac{3b^3d^3 \text{PolyLog}(4, e^{2(e+fx)})}{4f^4}
\end{aligned}$$

```
[Out] -3/2*b^3*d*(d*x+c)^2/f^2-3*a*b^2*(d*x+c)^3/f+1/2*b^3*(d*x+c)^3/f+1/4*a^3*(d*x+c)^4/d-3/4*a^2*b*(d*x+c)^4/d+3/4*a*b^2*(d*x+c)^4/d-1/4*b^3*(d*x+c)^4/d-3/2*b^3*d*(d*x+c)^2*coth(f*x+e)/f^2-3*a*b^2*(d*x+c)^3*coth(f*x+e)/f-1/2*b^3*(d*x+c)^3*coth(f*x+e)^2/f+3*b^3*d^2*(d*x+c)*ln(1-exp(2*f*x+2*e))/f^3+9*a*b^2*d*(d*x+c)^2*ln(1-exp(2*f*x+2*e))/f^2+3*a^2*b*(d*x+c)^3*ln(1-exp(2*f*x+2*e))/f+b^3*(d*x+c)^3*ln(1-exp(2*f*x+2*e))/f+3/2*b^3*d^3*polylog(2,exp(2*f*x+2*e))/f^4+9*a*b^2*d^2*(d*x+c)*polylog(2,exp(2*f*x+2*e))/f^2+3/2*b^3*d*(d*x+c)^2*polylog(2,exp(2*f*x+2*e))/f^2-9/2*a*b^2*d^3*polylog(3,exp(2*f*x+2*e))/f^4-9/2*a^2*b*d^2*(d*x+c)*polylog(3,exp(2*f*x+2*e))/f^3-3/2*b^3*d^2*(d*x+c)*polylog(3,exp(2*f*x+2*e))/f^3+9/4*a^2*b*d^3*polylog(4,exp(2*f*x+2*e))/f^4+3/4*b^3*d^3*polylog(4,exp(2*f*x+2*e))/f^4
```

Rubi [A] (verified)

Time = 0.76 (sec) , antiderivative size = 556, normalized size of antiderivative = 1.00, number of steps used = 28, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.550, Rules

used = {3803, 3797, 2221, 2611, 6744, 2320, 6724, 3801, 32, 2317, 2438}

$$\begin{aligned}
\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = & \frac{a^3(c + dx)^4}{4d} - \frac{9a^2bd^2(c + dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
& + \frac{9a^2bd(c + dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
& + \frac{3a^2b(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\
& - \frac{3a^2b(c + dx)^4}{4d} + \frac{9a^2bd^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{4f^4} \\
& + \frac{9ab^2d^2(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
& + \frac{9ab^2d(c + dx)^2 \log(1 - e^{2(e+fx)})}{f^2} \\
& - \frac{3ab^2(c + dx)^3 \coth(e + fx)}{f} - \frac{3ab^2(c + dx)^3}{f} \\
& + \frac{3ab^2(c + dx)^4}{4d} - \frac{9ab^2d^3 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\
& - \frac{3b^3d^2(c + dx) \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
& + \frac{3b^3d^2(c + dx) \log(1 - e^{2(e+fx)})}{f^3} \\
& + \frac{3b^3d(c + dx)^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
& - \frac{3b^3d(c + dx)^2 \coth(e + fx)}{2f^2} \\
& + \frac{b^3(c + dx)^3 \log(1 - e^{2(e+fx)})}{f} \\
& - \frac{b^3(c + dx)^3 \coth^2(e + fx)}{2f} \\
& - \frac{3b^3d(c + dx)^2}{2f^2} + \frac{b^3(c + dx)^3}{2f} \\
& - \frac{b^3(c + dx)^4}{4d} + \frac{3b^3d^3 \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^4} \\
& + \frac{3b^3d^3 \operatorname{PolyLog}(4, e^{2(e+fx)})}{4f^4}
\end{aligned}$$

[In] Int[(c + d*x)^3*(a + b*Coth[e + f*x])^3, x]

[Out] $(-3*b^3*d*(c + d*x)^2)/(2*f^2) - (3*a*b^2*(c + d*x)^3)/f + (b^3*(c + d*x)^3)/(2*f) + (a^3*(c + d*x)^4)/(4*d) - (3*a^2*b*(c + d*x)^4)/(4*d) + (3*a*b^2*$

$$\begin{aligned}
& (c + d*x)^4/(4*d) - (b^3*(c + d*x)^4)/(4*d) - (3*b^3*d*(c + d*x)^2*\text{Coth}[e + f*x])/(2*f^2) - (3*a*b^2*(c + d*x)^3*\text{Coth}[e + f*x])/f - (b^3*(c + d*x)^3*\text{Coth}[e + f*x]^2)/(2*f) + (3*b^3*d^2*(c + d*x)*\text{Log}[1 - E^{(2*(e + f*x))}])/f^3 \\
& + (9*a*b^2*d*(c + d*x)^2*\text{Log}[1 - E^{(2*(e + f*x))}])/f^2 + (3*a^2*b*(c + d*x)^3*\text{Log}[1 - E^{(2*(e + f*x))}])/f + (b^3*(c + d*x)^3*\text{Log}[1 - E^{(2*(e + f*x))}])/f + (3*b^3*d^3*\text{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^4) + (9*a*b^2*d^2*(c + d*x)*\text{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^3) \\
& + (9*a^2*b^2*d*(c + d*x)^2*\text{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2) + (3*b^3*d*(c + d*x)^2*\text{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2) - (9*a*b^2*d^3*\text{PolyLog}[3, E^{(2*(e + f*x))}])/(2*f^4) - (9*a^2*b*d^2*(c + d*x)*\text{PolyLog}[3, E^{(2*(e + f*x))}])/(2*f^3) - (3*b^3*d^2*(c + d*x)*\text{PolyLog}[3, E^{(2*(e + f*x))}])/(2*f^3) + (9*a^2*b^2*d^3*\text{PolyLog}[4, E^{(2*(e + f*x))}])/(4*f^4) \\
& + (3*b^3*d^3*\text{PolyLog}[4, E^{(2*(e + f*x))}])/(4*f^4)
\end{aligned}$$
Rule 32

```
Int[((a_.) + (b_.)*(x_.))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.))), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*\text{Log}[F]))*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*\text{Log}[F])), Int[(c + d*x)^(m - 1)*\text{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*\text{Log}[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))* (F_[v_]) /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_.) + (e_.)*(x_.))^(n_.))]/(x_), x_Symbol] :> Simp[-\text{PolyLog}[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_.)*(a_.) + (b_)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m -
1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3797

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + Pi*(k_.)*(f_.
)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist
[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x)
)/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && Int
egerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*(f_.*(x_)))^(n_), x_Symb
ol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Di
st[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x],
x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[
{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_.*(x_))]^(n_.)
, x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x],
x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_))^p]/((d_.) + (e_)*(x_), x_S
ymbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d
, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.*(x_)))^(m_)*PolyLog[n_, (d_)*(F_)^((c_.)*(a_.) + (b_.
)*(x_)))^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a +
b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m -
1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

integral =
$$\int (a^3(c+dx)^3 + 3a^2b(c+dx)^3 \coth(e+fx) + 3ab^2(c+dx)^3 \coth^2(e+fx) + b^3(c+dx)^3 \coth^3(e+fx)) dx$$

$$\begin{aligned}
&= \frac{a^3(c+dx)^4}{4d} + (3a^2b) \int (c+dx)^3 \coth(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx)^3 \coth^2(e+fx) dx + b^3 \int (c+dx)^3 \coth^3(e+fx) dx \\
&= \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} - (6a^2b) \int \frac{e^{2(e+fx)}(c+dx)^3}{1-e^{2(e+fx)}} dx + (3ab^2) \int (c \\
&\quad + dx)^3 dx + b^3 \int (c+dx)^3 \coth(e+fx) dx \\
&\quad + \frac{(9ab^2d) \int (c+dx)^2 \coth(e+fx) dx}{f} + \frac{(3b^3d) \int (c+dx)^2 \coth^2(e+fx) dx}{2f} \\
&= -\frac{3ab^2(c+dx)^3}{f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&\quad + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} \\
&\quad + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} - (2b^3) \int \frac{e^{2(e+fx)}(c+dx)^3}{1-e^{2(e+fx)}} dx \\
&\quad + \frac{(3b^3d^2) \int (c+dx) \coth(e+fx) dx}{f^2} - \frac{(9a^2bd) \int (c+dx)^2 \log(1-e^{2(e+fx)}) dx}{f} \\
&\quad - \frac{(18ab^2d) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} dx}{f} + \frac{(3b^3d) \int (c+dx)^2 dx}{2f} \\
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&\quad + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} \\
&\quad + \frac{9ab^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{(9a^2bd^2) \int (c+dx) \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(18ab^2d^2) \int (c+dx) \log(1-e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6b^3d^2) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} dx}{f^2} - \frac{(3b^3d) \int (c+dx)^2 \log(1-e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} - \frac{3a^2b(c+dx)^4}{4d} \\
&\quad + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} \\
&\quad - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} \\
&\quad + \frac{3b^3d^2(c+dx) \log(1-e^{2(e+fx)})}{f^3} + \frac{9ab^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{9ab^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad + \frac{3b^3d(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{9a^2bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
&\quad + \frac{(9a^2bd^3) \int \text{PolyLog}(3, e^{2(e+fx)}) dx}{2f^3} - \frac{(9ab^2d^3) \int \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^3} \\
&\quad - \frac{(3b^3d^3) \int \log(1-e^{2(e+fx)}) dx}{f^3} - \frac{(3b^3d^2) \int (c+dx) \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} \\
&\quad - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} + \frac{3b^3d^2(c+dx) \log(1-e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9ab^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{9ab^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} + \frac{3b^3d(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad - \frac{9a^2bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} - \frac{3b^3d^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
&\quad + \frac{(9a^2bd^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3,x)}{x} dx, x, e^{2(e+fx)}\right)}{4f^4} \\
&\quad - \frac{(9ab^2d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} \\
&\quad - \frac{(3b^3d^3) \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^4} + \frac{(3b^3d^3) \int \text{PolyLog}(3, e^{2(e+fx)}) dx}{2f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} \\
&\quad - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} + \frac{3b^3d^2(c+dx) \log(1-e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9ab^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{3b^3d^3 \text{PolyLog}(2, e^{2(e+fx)})}{2f^4} \\
&\quad + \frac{9ab^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad + \frac{3b^3d(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{9ab^2d^3 \text{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\
&\quad - \frac{9a^2bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} - \frac{3b^3d^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
&\quad + \frac{9a^2bd^3 \text{PolyLog}(4, e^{2(e+fx)})}{4f^4} + \frac{(3b^3d^3) \text{Subst}\left(\int \frac{\text{PolyLog}(3,x)}{x} dx, x, e^{2(e+fx)}\right)}{4f^4} \\
&= -\frac{3b^3d(c+dx)^2}{2f^2} - \frac{3ab^2(c+dx)^3}{f} + \frac{b^3(c+dx)^3}{2f} + \frac{a^3(c+dx)^4}{4d} \\
&\quad - \frac{3a^2b(c+dx)^4}{4d} + \frac{3ab^2(c+dx)^4}{4d} - \frac{b^3(c+dx)^4}{4d} \\
&\quad - \frac{3b^3d(c+dx)^2 \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx)^3 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^3 \coth^2(e+fx)}{2f} + \frac{3b^3d^2(c+dx) \log(1-e^{2(e+fx)})}{f^3} \\
&\quad + \frac{9ab^2d(c+dx)^2 \log(1-e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^3 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^3 \log(1-e^{2(e+fx)})}{f} + \frac{3b^3d^3 \text{PolyLog}(2, e^{2(e+fx)})}{2f^4} \\
&\quad + \frac{9ab^2d^2(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^3} + \frac{9a^2bd(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
&\quad + \frac{3b^3d(c+dx)^2 \text{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{9ab^2d^3 \text{PolyLog}(3, e^{2(e+fx)})}{2f^4} \\
&\quad - \frac{9a^2bd^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} - \frac{3b^3d^2(c+dx) \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\
&\quad + \frac{9a^2bd^3 \text{PolyLog}(4, e^{2(e+fx)})}{4f^4} + \frac{3b^3d^3 \text{PolyLog}(4, e^{2(e+fx)})}{4f^4}
\end{aligned}$$

Mathematica [B] (verified)

Leaf count is larger than twice the leaf count of optimal. 2043 vs. $2(556) = 1112$.

Time = 8.91 (sec) , antiderivative size = 2043, normalized size of antiderivative = 3.67

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = \text{Result too large to show}$$

[In] `Integrate[(c + d*x)^3*(a + b*Coth[e + f*x])^3,x]`

[Out]
$$\begin{aligned} & ((-(b^3*c^3) - 3*b^3*c^2*d*x - 3*b^3*c*d^2*x^2 - b^3*d^3*x^3)*\text{Csch}[e + f*x]^2)/(2*f) - (b*E^{(2*e)}*(24*b^2*c*d^2*x + 72*a*b*c^2*d*f*x + 24*a^2*c^3*f^2*x + 8*b^2*c^3*f^2*x + 12*b^2*d^3*x^2 + 72*a*b*c*d^2*f*x^2 + 36*a^2*c^2*d*f^2*x^2 + 12*b^2*c^2*d*f^2*x^2 + 24*a*b*d^3*f*x^3 + 24*a^2*c*d^2*f^2*x^3 + 8*b^2*c*d^2*f^2*x^3 + 6*a^2*d^3*f^2*x^4 + 2*b^2*d^3*f^2*x^4 - 36*a*b*c^2*d*Log[g[1 - E^{(2*(e + f*x))}]] + (36*a*b*c^2*d*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - (12*b^2*c*d^2*Log[1 - E^{(2*(e + f*x))}])/f + (12*b^2*c*d^2*Log[1 - E^{(2*(e + f*x))}])/E^{(2*(e + f*x))})/(E^{(2*e)}*f) - 12*a^2*c^3*f*Log[1 - E^{(2*(e + f*x))}] - 4*b^2*c^3*f*Log[1 - E^{(2*(e + f*x))}] + (12*a^2*c^3*f*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} + (4*b^2*c^3*f*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - 72*a*b*c*d^2*x*Log[1 - E^{(2*(e + f*x))}] + (72*a*b*c*d^2*x*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - (12*b^2*d^3*x*Log[1 - E^{(2*(e + f*x))}])/f + (12*b^2*d^3*x*Log[1 - E^{(2*(e + f*x))}])/(E^{(2*e)}*f) - 36*a^2*c^2*d*f*x*Log[1 - E^{(2*(e + f*x))}] - 12*b^2*c^2*d*f*x*Log[1 - E^{(2*(e + f*x))}] + (36*a^2*c^2*d*f*x*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} + (12*b^2*c^2*d*f*x*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - 36*a*b*d^3*x^2*Log[1 - E^{(2*(e + f*x))}] + (36*a*b*d^3*x^2*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - 36*a^2*c^2*d^2*f*x^2*Log[1 - E^{(2*(e + f*x))}] - 12*b^2*c*d^2*f*x^2*Log[1 - E^{(2*(e + f*x))}] + (36*a^2*c*d^2*f*x^2*Log[1 - E^{(2*(e + f*x))}])/E^{(2*e)} - (6*d*(-1 + E^{(2*e)})*(6*a*b*d*f*(c + d*x) + 3*a^2*f^2*(c + d*x)^2 + b^2*(d^2 + c^2*f^2 + 2*c*d*f^2*x + d^2*f^2*x^2))*PolyLog[2, E^{(2*(e + f*x))})/(E^{(2*e)}*f^2) + (6*d^2*(-1 + E^{(2*e)})*(3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*PolyLog[3, E^{(2*(e + f*x))})/(E^{(2*e)}*f^2) - (9*a^2*d^3*PolyLog[4, E^{(2*(e + f*x))}])/f^2 - (3*b^2*d^3*PolyLog[4, E^{(2*(e + f*x))}])/f^2 + (9*a^2*d^3*PolyLog[4, E^{(2*(e + f*x))}])/E^{(2*(e + f*x))})/(E^{(2*e)}*f^2) + (3*b^2*d^3*PolyLog[4, E^{(2*(e + f*x))}])/(E^{(2*e)}*f^2))/((4*(-1 + E^{(2*e)})*f^2) + (3*x^2*(-(a^3*c^2*d) + 3*a^2*b*c^2*d - 3*a*b^2*c^2*d + b^3*c^2*d + a^3*c^2*d*Cosh[2*e] + 3*a^2*b*c^2*d*Cosh[2*e] + 3*a*b^2*c^2*d*Cosh[2*e] + b^3*c^2*d*Cosh[2*e] + a^3*c^2*d*Sinh[2*e] + 3*a^2*b*c^2*d*Sinh[2*e] + 3*a*b^2*c^2*d*Sinh[2*e] + b^3*c^2*d*Sinh[2*e]))/(2*(-1 + Cosh[2*e] + Sinh[2*e])) + (x^3*(-(a^3*c^2*d) + 3*a^2*b*c^2*d^2 - 3*a*b^2*c^2*d^2 + b^3*c^2*d^2 + a^3*c^2*d^2*Cosh[2*e] + 3*a^2*b*c^2*d^2*Cosh[2*e] + 3*a*b^2*c^2*d^2*Cosh[2*e] + b^3*c^2*d^2*Cosh[2*e] + a^3*c^2*d^2*Sinh[2*e] + 3*a^2*b*c^2*d^2*Sinh[2*e] + 3*a*b^2*c^2*d^2*Sinh[2*e] + 3*a^3*c^2*d^2*Cosh[2*e] + 3*a^2*b*c^2*d^2*Cosh[2*e] + 3*a*b^2*c^2*d^2*Cosh[2*e] + b^3*c^2*d^2*Cosh[2*e]))$$

$$\begin{aligned}
& \frac{-2 \operatorname{Sinh}[2 e] + b^3 c d^2 \operatorname{Sinh}[2 e])}{(-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e])} + (x^4 (-a^3 d^3) + 3 a^2 b d^3 - 3 a b^2 d^3 + b^3 d^3 + a^3 d^3 \operatorname{Cosh}[2 e] + 3 a^2 b d^3 \operatorname{Cosh}[2 e] + 3 a b^2 d^3 \operatorname{Cosh}[2 e] + b^3 d^3 \operatorname{Cosh}[2 e] + a^3 d^3 \operatorname{Sinh}[2 e] + 3 a^2 b d^3 \operatorname{Sinh}[2 e] + 3 a b^2 d^3 \operatorname{Sinh}[2 e] + b^3 d^3 \operatorname{Sinh}[2 e]))}{(4 (-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e]))} + x (a^3 c^3 + 3 a^2 b^2 c^3 + (3 a^2 b c^3)) \\
& /(-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e]) + (3 a^2 b^2 c^3 \operatorname{Cosh}[2 e] + 3 a^2 b c^3 \operatorname{Sinh}[2 e]) /(-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e]) + (2 b^3 c^3 \operatorname{Cosh}[2 e] + 2 b^3 c^3 \operatorname{Sinh}[2 e]) /(-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e]) * (1 + \operatorname{Cosh}[2 e] + \operatorname{Cosh}[4 e] + \operatorname{Sinh}[2 e] + \operatorname{Sinh}[4 e]) + (2 b^3 c^3 \operatorname{Cosh}[4 e] + 2 b^3 c^3 \operatorname{Sinh}[4 e]) /((-1 + \operatorname{Cosh}[2 e] + \operatorname{Sinh}[2 e]) * (1 + \operatorname{Cosh}[2 e] + \operatorname{Cosh}[4 e] + \operatorname{Sinh}[2 e] + \operatorname{Sinh}[4 e])) + (b^3 c^3) /(-1 + \operatorname{Cosh}[6 e] + \operatorname{Sinh}[6 e]) + (b^3 c^3 \operatorname{Cosh}[6 e] + b^3 c^3 \operatorname{Sinh}[6 e]) /(-1 + \operatorname{Cosh}[6 e] + \operatorname{Sinh}[6 e]) + (3 \operatorname{Csch}[e] * \operatorname{Csch}[e + f x] * (b^3 c^2 d \operatorname{Sinh}[f x] + 2 a b^2 c^3 f \operatorname{Sinh}[f x] + 2 b^3 c d^2 x \operatorname{Sinh}[f x] + 6 a b^2 c^2 d^2 f x \operatorname{Sinh}[f x] + b^3 d^3 x^2 \operatorname{Sinh}[f x] + 6 a b^2 c d^2 f x^2 \operatorname{Sinh}[f x] + 2 a b^2 d^3 f x^3 \operatorname{Sinh}[f x])) / (2 f^2)
\end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2822 vs. $2(524) = 1048$.
Time = 0.78 (sec), antiderivative size = 2823, normalized size of antiderivative = 5.08

method	result	size
risch	Expression too large to display	2823

```
[In] int((d*x+c)^3*(a+b*cot(f*x+e))^3,x,method=_RETURNVERBOSE)

[Out] -3/f^4*b^3*e^2*d^3-3/f^2*b^3*d^3*x^2-3/2/f^4*b^3*e^4*d^3+3/f^4*b^3*d^3*polylog(2,exp(f*x+e))+3/f^4*b^3*d^3*polylog(2,-exp(f*x+e))+6/f^4*b^3*d^3*polylog(4,exp(f*x+e))+6/f^4*b^3*d^3*polylog(4,-exp(f*x+e))-18/f*b^2*a*c*d^2*x^2+12/f^3*b^2*c^3*d^2*c*a^2+6/f^2*b^3*d^2*c*polylog(2,exp(f*x+e))*x+3/f*b^3*d^2*c*ln(1+exp(f*x+e))*x^2+6/f^2*b^3*d^2*c*polylog(2,-exp(f*x+e))*x+3/f*b*a^2*d^3*ln(1-exp(f*x+e))*x^3+9/f^2*b*a^2*d^3*polylog(2,exp(f*x+e))*x^2-18/f^3*b*a^2*d^3*polylog(3,exp(f*x+e))*x+3/f*b*a^2*d^3*ln(1+exp(f*x+e))*x^3+9/f^2*b*a^2*d^3*polylog(2,-exp(f*x+e))*x^2-18/f^3*b*a^2*d^3*polylog(3,-exp(f*x+e))*x+9/f^4*b^2*c^2*a*d^3*ln(exp(f*x+e)-1)-6/f^3*b^3*d^3*e*x+12/f^4*b^2*c^2*a*d^3-6/f*b^2*a*d^3*x^3-2/f^3*b^3*e^3*d^3*x+3*a^2*b*c^3*x+3*a*b^2*c^3*x+1/4*d^3*a^3*x^4-1/4*d^3*b^3*x^4+1/4/d*c^4*a^3+1/4/d*c^4*b^3-18/f^4*b^2*c^2*a*d^3*ln(exp(f*x+e))-3/f^3*b^3*e^2*d^2*c*ln(1-exp(f*x+e))-9/f^4*b^2*c^2*a*d^3*ln(1-exp(f*x+e))-3/f^4*b*c^3*a^2*d^3*ln(exp(f*x+e)-1)+6/f^4*b*c^3*a^2*d^3*ln(exp(f*x+e))+18/f^3*b^2*a*c*d^2*polylog(2,-exp(f*x+e))+9/f^2*b^2*a*d^3*ln(1-exp(f*x+e))*x^2+18/f^3*b^2*a*d^3*polylog(2,exp(f*x+e))*x+9/f^2*b^2*a*d^3*ln(1+exp(f*x+e))*x^2+18/f^3*b^2*a*d^3*polylog(2,-exp(f*x+e))*x+3/f*b^3*c^2*d*ln(1-exp(f*x+e))*x+3/f^2*b^3*c^2*d*ln(1-exp(f*x+e))*e+3/f*b^3*c^2*d*ln(1+exp(f*x+e))*x+3/f^4*b*c^3*a^2*d^3*ln(1-exp(f*x+e))+9/f^2*b*d*c^2*a^2*polylog(2,exp(f*x+e))+9/f^2*b*d*c^2*a^2*poly
```

$$\begin{aligned}
& \log(2, -\exp(f*x+e)) + 3/f^3 * b^3 * e^2 * d^2 * c * \ln(\exp(f*x+e) - 1) - 6/f^3 * b^3 * e^2 * d^2 * c \\
& * \ln(\exp(f*x+e)) - 18/f^3 * b * d^2 * c * a^2 * \text{polylog}(3, \exp(f*x+e)) - 18/f^3 * b * d^2 * c * a^2 \\
& * \text{polylog}(3, -\exp(f*x+e)) - 3/f^2 * b^3 * e * c^2 * d * \ln(\exp(f*x+e) - 1) + 6/f^2 * b^3 * e * c^2 * \\
& d * \ln(\exp(f*x+e)) + 9/f^2 * b^2 * a * c^2 * d * \ln(\exp(f*x+e) - 1) + 9/f^2 * b^2 * a * c^2 * d * \ln(1+ \\
& \exp(f*x+e)) - 18/f^2 * b^2 * a * c^2 * d * \ln(\exp(f*x+e)) + 3/f * b^3 * d^2 * c * \ln(1-\exp(f*x+e)) \\
& * x^2 + 18/f^3 * b^2 * e^2 * a * d^3 * x - 18/f^3 * b^2 * a * c * d^2 * e^2 + 6/f^2 * b^3 * e^2 * d^2 * c * x - 6 \\
& /f * b^3 * c^2 * d * e * x - 6/f^3 * b * e^3 * a^2 * d^3 * x - 9/f^2 * b * d * c^2 * a^2 * e^2 - 18/f * b * d * c^2 * a \\
& ^2 * e * x - 36/f^2 * b^2 * a * c * d^2 * e * x + 18/f^2 * b * e^2 * d^2 * c * a^2 * x - 9/f^3 * b * e^2 * d^2 * c * a^2 \\
& * \ln(1-\exp(f*x+e)) + 9/f * b * d * c^2 * a^2 * \ln(1-\exp(f*x+e)) * x + 9/f^2 * b * d * c^2 * a^2 * \ln(1-\exp(f*x+e)) \\
& * e + 9/f * b * d * c^2 * a^2 * \ln(1+\exp(f*x+e)) * x + 9/f * b * d^2 * c * a^2 * \ln(1-\exp(f*x+e)) * x^2 + 18/f^2 * b * d^2 * c * a^2 * \\
& \text{polylog}(2, \exp(f*x+e)) * x + 9/f * b * d^2 * c * a^2 * \ln(1+\exp(f*x+e)) * x - 9/f^2 * b * e * d * c \\
& ^2 * a^2 * \ln(\exp(f*x+e) - 1) + 18/f^2 * b * e * d * c^2 * a^2 * \ln(\exp(f*x+e)) - 18/f^3 * b^2 * e * a * \\
& c * d^2 * \ln(\exp(f*x+e) - 1) + 36/f^3 * b^2 * e * a * c * d^2 * \ln(\exp(f*x+e)) - b^2 * (6 * a * d^3 * f * x \\
& ^3 * \exp(2 * f * x + 2 * e) + 2 * b * d^3 * f * x^3 * \exp(2 * f * x + 2 * e) + 18 * a * c * d^2 * f * x^2 * \exp(2 * f * x + 2 \\
& * e) + 6 * b * c * d^2 * f * x^2 * \exp(2 * f * x + 2 * e) + 18 * a * c^2 * d * f * x * \exp(2 * f * x + 2 * e) - 6 * a * d^3 * f * \\
& x^3 + 6 * b * c^2 * d * f * x * \exp(2 * f * x + 2 * e) + 3 * b * d^3 * x^2 * \exp(2 * f * x + 2 * e) + 6 * a * c^3 * f * \exp(2 \\
& * f * x + 2 * e) - 18 * a * c * d^2 * f * x^2 + 2 * b * c^3 * f * \exp(2 * f * x + 2 * e) + 6 * b * c * d^2 * x * \exp(2 * f * x + 2 \\
& * e) - 18 * a * c^2 * d * f * x + 3 * \exp(2 * f * x + 2 * e) * d * b * c^2 - 3 * b * d^3 * x^2 - 6 * a * c^3 * f - 6 * b * c * d^2 \\
& * x^3 - 3 * b * c^2 * d) / f^2 / (\exp(2 * f * x + 2 * e) - 1)^2 - 3/f^2 * b^3 * c^2 * d * e^2 - 9/2/f^4 * b * e^4 * a^2 \\
& 2 * d^3 + 4/f^3 * b^3 * e^3 * d^2 * c - 3/f^4 * b^3 * e * d^3 * \ln(\exp(f*x+e) - 1) + 6/f^4 * b^3 * e * d^3 * \\
& \ln(\exp(f*x+e)) + 3/f^3 * b^3 * d^3 * \ln(1-\exp(f*x+e)) * x + 3/f^4 * b^3 * d^3 * \ln(1-\exp(f*x+e)) * e + 3/f^3 * b^3 * d^3 * \\
& \ln(1+\exp(f*x+e)) * x - 1/f^4 * b^3 * e^3 * d^3 * \ln(\exp(f*x+e) - 1) + 2/f^4 * b^3 * e^3 * d^3 * \ln(\exp(f*x+e)) - 3 * d^2 * a^2 * b * c * x^3 + 3 * d^2 * a^2 * b * c * x^3 - 9/2 * d * a \\
& ^2 * b * c^2 * x^2 + 9/2 * d * a * b^2 * c^2 * x^2 + 1/f * b^3 * c^3 * \ln(\exp(f*x+e) - 1) + 1/f * b^3 * c^3 * l \\
& n(1+\exp(f*x+e)) - 2/f * b^3 * c^3 * \ln(\exp(f*x+e)) - 3/4 * d^3 * a^2 * b * x^4 + 3/4 * d^3 * a * b^2 * x^4 + d^2 * a^2 * b^2 * \\
& x^3 - d^2 * a^2 * b * c * x^3 - d^2 * b^2 * c * x^3 + 3/2 * d * a^2 * c * x^2 * x^2 - 3/2 * d * b^2 * c * x^2 * x^2 + a^2 * c * x^3 \\
& * x + b^2 * c * x^3 + 3/4 * d * c^4 * a^2 * b + 3/4 * d * c^4 * a * b^2 - 18/f^4 * b^2 * a * d^3 * \text{polylog}(3, \exp(f*x+e)) - 18/f^4 * b^2 * a * d^3 * \\
& \text{polylog}(3, -\exp(f*x+e)) + 1/f^4 * b^2 * a * d^3 * \ln(1-\exp(f*x+e)) + 3/f * b * a^2 * c^3 * \ln(\exp(f*x+e) - 1) + 3/f * b * a^2 * c^3 * \\
& \ln(1+\exp(f*x+e)) - 6/f^3 * b^3 * d^2 * c * \ln(\exp(f*x+e)) + 3/f^2 * b^3 * c^2 * d * \text{polylog}(2, \exp(f*x+e)) + 3/f^2 * b^3 * c^2 * d * \\
& \text{polylog}(2, -\exp(f*x+e)) + 1/f * b^3 * d^3 * \ln(1-\exp(f*x+e)) * x^3 + 3/f^2 * b^3 * d^3 * \text{polylog}(2, -\exp(f*x+e)) * x + 1/f * b^3 * d^3 * \\
& \text{polylog}(2, \exp(f*x+e)) * x^2 - 6/f^3 * b^3 * d^3 * \text{polylog}(3, -\exp(f*x+e)) * x + 18/f^4 * b * a^2 * d^3 * \text{polylog}(4, \exp(f*x+e)) + 18/f^4 * b * a^2 * d^3 * \\
& \text{polylog}(4, -\exp(f*x+e)) - 6/f^3 * b^3 * d^2 * c * \text{polylog}(3, -\exp(f*x+e)) + 9/f^3 * b * e^2 * d^2 * c * a^2 \\
& * \ln(\exp(f*x+e) - 1) - 18/f^3 * b * e^2 * d^2 * c * a^2 * \ln(\exp(f*x+e)) + 18/f^2 * b^2 * a * c * d^2 * \\
& \ln(1-\exp(f*x+e)) * x + 18/f^2 * b^2 * a * c * d^2 * \ln(1+\exp(f*x+e)) * x + 18/f^3 * b^2 * a * c * d^2 * \\
& \ln(1-\exp(f*x+e)) * e
\end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 11137 vs. $2(520) = 1040$.

Time = 0.39 (sec) , antiderivative size = 11137, normalized size of antiderivative = 20.03

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3*(a+b*coth(f*x+e))^3,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 \, dx = \int (a + b \coth(e + fx))^3 (c + dx)^3 \, dx$$

```
[In] integrate((d*x+c)**3*(a+b*cot(f*x+e))**3,x)
```

[Out] $\text{Integral}((a + b \coth(e + f x))^3 (c + d x)^3, x)$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1531 vs. $2(520) = 1040$.

Time = 0.30 (sec) , antiderivative size = 1531, normalized size of antiderivative = 2.75

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^3*(a+b*coth(f*x+e))^3,x, algorithm="maxima")
```

```
[Out] 1/4*a^3*d^3*x^4 + a^3*c*d^2*x^3 + 3/2*a^3*c^2*d*x^2 + a^3*c^3*x + 3*a^2*b*c^3*log(sinh(f*x + e))/f + 1/4*(24*a*b^2*c^3*f + 12*b^3*c^2*d + (3*a^2*b*d^3*f^2 + 3*a*b^2*d^3*f^2 + b^3*d^3*f^2)*x^4 + 4*(3*a^2*b*c*d^2*f^2 + b^3*c*d^2*f^2 + 3*(c*d^2*f^2 + 2*d^3*f)*a*b^2)*x^3 + 6*(3*a^2*b*c^2*d*f^2 + 3*(c^2*d*f^2 + 4*c*d^2*f)*a*b^2 + (c^2*d*f^2 + 2*d^3)*b^3)*x^2 + 4*(3*(c^3*f^2 + 6*c^2*d*f)*a*b^2 + (c^3*f^2 + 6*c*d^2)*b^3)*x + ((3*a^2*b*d^3*f^2*e^(4*e) + 3*a*b^2*d^3*f^2*e^(4*e) + b^3*d^3*f^2*e^(4*e))*x^4 + 4*(3*a^2*b*c*d^2*f^2*e^(4*e) + 3*a*b^2*c*d^2*f^2*e^(4*e) + b^3*c*d^2*f^2*e^(4*e))*x^3 + 6*(3*a^2*b*c^2*d*f^2*e^(4*e) + 3*a*b^2*c^2*d*f^2*e^(4*e) + b^3*c^2*d*f^2*e^(4*e))*x^2 + 4*(3*a*b^2*c^3*f^2*e^(4*e) + b^3*c^3*f^2*e^(4*e))*x)*e^(4*f*x) - 2*(12*a*b^2*c^3*f^2*e^(2*e) + (3*a^2*b*d^3*f^2*e^(2*e) + 3*a*b^2*d^3*f^2*e^(2*e) + b^3*d^3*f^2*e^(2*e))*x^4 + 2*(2*c^3*f^2*e^(2*e) + 3*c^2*d^2*e^(2*e))*b^3 + 4*(3*a^2*b*c*d^2*f^2*e^(2*e) + 3*(c*d^2*f^2*e^(2*e) + d^3*f^2*e^(2*e))*a*b^2 + (c^2*d^2*f^2 + 2*d^3*f^2)*b^3)*x^3 + 6*(3*a^2*b*c^2*d*f^2*e^(2*e) + 3*a*b^2*c^2*d*f^2*e^(2*e) + b^3*c^2*d*f^2*e^(2*e))*x^2 + 4*(3*a*b^2*c^3*f^2*e^(2*e) + b^3*c^3*f^2*e^(2*e))*x)*e^(4*f*x) + 2*(2*c^3*f^2*e^(2*e) + 3*c^2*d^2*e^(2*e))*b^3 + 4*(3*a^2*b*c*d^2*f^2*e^(2*e) + 3*(c*d^2*f^2*e^(2*e) + d^3*f^2*e^(2*e))*a*b^2 + (c^2*d^2*f^2 + 2*d^3*f^2)*b^3)*x^3 + 6*(3*a^2*b*c^2*d*f^2*e^(2*e) + 3*a*b^2*c^2*d*f^2*e^(2*e) + b^3*c^2*d*f^2*e^(2*e))*x^2 + 4*(3*a*b^2*c^3*f^2*e^(2*e) + b^3*c^3*f^2*e^(2*e))*x)*e^(4*f*x)
```

$$\begin{aligned}
& *d^2*f^2*e^{(2*e)} + d^3*f*e^{(2*e)}*b^3*x^3 + 6*(3*a^2*b*c^2*d*f^2*e^{(2*e)}) + \\
& 3*(c^2*d*f^2*e^{(2*e)} + 2*c*d^2*f*e^{(2*e)})*a*b^2 + (c^2*d*f^2*e^{(2*e)} + 2*c \\
& *d^2*f*e^{(2*e)} + d^3*e^{(2*e)})*b^3*x^2 + 4*(3*(c^3*f^2*e^{(2*e)} + 3*c^2*d*f \\
& e^{(2*e)})*a*b^2 + (c^3*f^2*e^{(2*e)} + 3*c^2*d*f*e^{(2*e)} + 3*c*d^2*e^{(2*e)})*b \\
& 3*x)*e^{(2*f*x)}/(f^2*e^{(4*f*x + 4*e)} - 2*f^2*e^{(2*f*x + 2*e)} + f^2) - 2*(9 \\
& *a*b^2*c^2*d*f + (c^3*f^2 + 3*c*d^2)*b^3*x/f^2 + (9*a*b^2*c^2*d*f + (c^3*f \\
& ^2 + 3*c*d^2)*b^3)*log(e^{(f*x + e)} + 1)/f^3 + (9*a*b^2*c^2*d*f + (c^3*f^2 + \\
& 3*c*d^2)*b^3)*log(e^{(f*x + e)} - 1)/f^3 + (f^3*x^3*log(e^{(f*x + e)} + 1) + 3 \\
& *f^2*x^2*dilog(-e^{(f*x + e)}) - 6*f*x*polylog(3, -e^{(f*x + e)}) + 6*polylog(4 \\
& , -e^{(f*x + e)}))*(3*a^2*b*d^3 + b^3*d^3)/f^4 + (f^3*x^3*log(-e^{(f*x + e)} + \\
& 1) + 3*f^2*x^2*dilog(e^{(f*x + e)}) - 6*f*x*polylog(3, e^{(f*x + e)}) + 6*polylog(4 \\
& , e^{(f*x + e)}))*(3*a^2*b*d^3 + b^3*d^3)/f^4 + 3*(3*a^2*b*c*d^2*f + b^3*c \\
& *d^2*f + 3*a*b^2*d^3)*(f^2*x^2*log(e^{(f*x + e)} + 1) + 2*f*x*dilog(-e^{(f*x \\
& + e)}) - 2*polylog(3, -e^{(f*x + e)}))/f^4 + 3*(3*a^2*b*c*d^2*f + b^3*c*d^2*f \\
& + 3*a*b^2*d^3)*(f^2*x^2*log(-e^{(f*x + e)} + 1) + 2*f*x*dilog(e^{(f*x + e)}) - \\
& 2*polylog(3, e^{(f*x + e)}))/f^4 + 3*(3*a^2*b*c^2*d*f^2 + 6*a*b^2*c*d^2*f + (\\
& c^2*d*f^2 + d^3)*b^3)*(f*x*log(e^{(f*x + e)} + 1) + dilog(-e^{(f*x + e)}))/f^4 \\
& + 3*(3*a^2*b*c^2*d*f^2 + 6*a*b^2*c*d^2*f + (c^2*d*f^2 + d^3)*b^3)*(f*x*log(\\
& -e^{(f*x + e)} + 1) + dilog(e^{(f*x + e)}))/f^4 - 1/2*((3*a^2*b*d^3 + b^3*d^3)* \\
& f^4*x^4 + 4*(3*a^2*b*c*d^2*f + b^3*c*d^2*f + 3*a*b^2*d^3)*f^3*x^3 + 6*(3*a^ \\
& 2*b*c^2*d*f^2 + 6*a*b^2*c*d^2*f + (c^2*d*f^2 + d^3)*b^3)*f^2*x^2)/f^4
\end{aligned}$$

Giac [F]

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = \int (dx + c)^3 (b \coth(fx + e) + a)^3 dx$$

[In] integrate((d*x+c)^3*(a+b*coth(f*x+e))^3,x, algorithm="giac")

[Out] integrate((d*x + c)^3*(b*coth(f*x + e) + a)^3, x)

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^3 (a + b \coth(e + fx))^3 dx = \int (a + b \coth(e + fx))^3 (c + dx)^3 dx$$

[In] int((a + b*coth(e + f*x))^3*(c + d*x)^3,x)

[Out] int((a + b*coth(e + f*x))^3*(c + d*x)^3, x)

3.48 $\int (c + dx)^2 (a + b \coth(e + fx))^3 dx$

Optimal result	321
Rubi [A] (verified)	322
Mathematica [A] (verified)	327
Maple [B] (verified)	328
Fricas [B] (verification not implemented)	329
Sympy [F]	329
Maxima [B] (verification not implemented)	329
Giac [F]	330
Mupad [F(-1)]	330

Optimal result

Integrand size = 20, antiderivative size = 401

$$\begin{aligned}
\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = & \frac{b^3 c d x}{f} + \frac{b^3 d^2 x^2}{2 f} - \frac{3 a b^2 (c + dx)^2}{f} \\
& + \frac{a^3 (c + dx)^3}{3 d} - \frac{a^2 b (c + dx)^3}{d} + \frac{a b^2 (c + dx)^3}{d} \\
& - \frac{b^3 (c + dx)^3}{3 d} - \frac{b^3 d (c + dx) \coth(e + fx)}{f^2} \\
& - \frac{3 a b^2 (c + dx)^2 \coth(e + fx)}{f} \\
& - \frac{b^3 (c + dx)^2 \coth^2(e + fx)}{2 f} \\
& + \frac{6 a b^2 d (c + dx) \log(1 - e^{2(e+fx)})}{f^2} \\
& + \frac{3 a^2 b (c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{b^3 (c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{b^3 d^2 \log(\sinh(e + fx))}{f^3} \\
& + \frac{3 a b^2 d^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
& + \frac{3 a^2 b d (c + dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
& + \frac{b^3 d (c + dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
& - \frac{3 a^2 b d^2 \text{PolyLog}(3, e^{2(e+fx)})}{2 f^3} \\
& - \frac{b^3 d^2 \text{PolyLog}(3, e^{2(e+fx)})}{2 f^3}
\end{aligned}$$

```
[Out] b^3*c*d*x/f+1/2*b^3*d^2*x^2/f-3*a*b^2*(d*x+c)^2/f+1/3*a^3*(d*x+c)^3/d-a^2*b*(d*x+c)^3/d+a*b^2*(d*x+c)^3/d-1/3*b^3*(d*x+c)^3/d-b^3*d*(d*x+c)*coth(f*x+e)/f^2-3*a*b^2*(d*x+c)^2*coth(f*x+e)/f-1/2*b^3*(d*x+c)^2*coth(f*x+e)^2/f+6*a*b^2*d*(d*x+c)*ln(1-exp(2*f*x+2*e))/f^2+3*a^2*b*(d*x+c)^2*ln(1-exp(2*f*x+2*e))/f+b^3*(d*x+c)^2*ln(sinh(f*x+e))/f^3+3*a*b^2*d^2*polylog(2,exp(2*f*x+2*e))/f^3+3*a^2*b*d*(d*x+c)*polylog(2,exp(2*f*x+2*e))/f^2+b^3*d*(d*x+c)*polylog(2,exp(2*f*x+2*e))/f^2-3/2*a^2*b*d^2*polylog(3,exp(2*f*x+2*e))/f^3-1/2*b^3*d^2*polylog(3,exp(2*f*x+2*e))/f^3
```

Rubi [A] (verified)

Time = 0.51 (sec) , antiderivative size = 401, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.550$, Rules used = {3803, 3797, 2221, 2611, 2320, 6724, 3801, 2317, 2438, 32, 3556}

$$\begin{aligned} \int (c + dx)^2 (a + b \coth(e + fx))^3 dx = & \frac{a^3(c + dx)^3}{3d} + \frac{3a^2bd(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & + \frac{3a^2b(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{a^2b(c + dx)^3}{d} - \frac{3a^2bd^2 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\ & + \frac{6ab^2d(c + dx) \log(1 - e^{2(e+fx)})}{f^2} \\ & - \frac{3ab^2(c + dx)^2 \coth(e + fx)}{f} - \frac{3ab^2(c + dx)^2}{f} \\ & + \frac{ab^2(c + dx)^3}{d} + \frac{3ab^2d^2 \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^3} \\ & + \frac{b^3d(c + dx) \operatorname{PolyLog}(2, e^{2(e+fx)})}{f^2} \\ & - \frac{b^3d(c + dx) \coth(e + fx)}{f^2} \\ & + \frac{b^3(c + dx)^2 \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{b^3(c + dx)^2 \coth^2(e + fx)}{2f} + \frac{b^3cdx}{f} \\ & - \frac{b^3(c + dx)^3}{3d} - \frac{b^3d^2 \operatorname{PolyLog}(3, e^{2(e+fx)})}{2f^3} \\ & + \frac{b^3d^2 \log(\sinh(e + fx))}{f^3} + \frac{b^3d^2x^2}{2f} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)^2*(a + b*\operatorname{Coth}[e + f*x])^3, x]$

[Out] $(b^{3*c*d*x}/f + (b^{3*d^2*x^2}/(2*f) - (3*a*b^2*(c + d*x)^2)/f + (a^{3*(c + d*x)^3}/(3*d) - (a^2*b*(c + d*x)^3)/d + (a*b^2*(c + d*x)^3)/d - (b^{3*(c + d*x)^3}/(3*d) - (b^{3*d*(c + d*x)}*\operatorname{Coth}[e + f*x])/f^2 - (3*a*b^2*(c + d*x)^2*\operatorname{Coth}[e + f*x])/f - (b^{3*(c + d*x)^2}*\operatorname{Coth}[e + f*x]^2)/(2*f) + (6*a*b^2*d*(c + d*x)*\operatorname{Log}[1 - E^{2*(e + f*x)}])/f^2 + (3*a^2*b*(c + d*x)^2*\operatorname{Log}[1 - E^{2*(e + f*x)}])/f + (b^{3*(c + d*x)^2}*\operatorname{Log}[1 - E^{2*(e + f*x)}])/f + (b^{3*d^2*\operatorname{Log}[Si nh[e + f*x]]}/f^3 + (3*a*b^2*d^2*\operatorname{PolyLog}[2, E^{2*(e + f*x)}])/f^3 + (3*a^2*b*d*(c + d*x)*\operatorname{PolyLog}[2, E^{2*(e + f*x)}])/f^2 + (b^{3*d*(c + d*x)}*\operatorname{PolyLog}[2, E^{2*(e + f*x)}])/f^2 - (3*a^2*b*d^2*\operatorname{PolyLog}[3, E^{2*(e + f*x)}])/f^3 - (b^{3*d^2*\operatorname{PolyLog}[3, E^{2*(e + f*x)}]})/(2*f^3))$

Rule 32

```
Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] :> Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]
```

Rule 2221

```
Int[((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*((F_)^((g_.)*((e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_.) + (b_.)*((F_)^((e_.)*((c_.) + (d_.)*(x_))))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*((a_.) + (b_.)*x))*(F_[v_]) /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[(c_.)*((d_.) + (e_.)*(x_))^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*((a_.) + (b_.)*(x_))))^(n_.))*((f_.) + (g_.)*(x_))^(m_.), x_Symbol] :> Simp[((-f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3556

```
Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3797

```
Int[((c_.) + (d_.)*(x_))^(m_.)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_ .)*(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi)))/E^(2*I*k*Pi), x], x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_.)*(x_))^(m_.*((b_.)*tan[(e_.) + (f_ .)*(x_)])^n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_.)*(x_))^(m_.*((a_.) + (b_.)*tan[(e_.) + (f_ .)*(x_)])^n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_ .)*(a_ .) + (b_ .)*(x_ )]^p]/((d_ .) + (e_ .)*(x_ )), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int (a^3(c+dx)^2 + 3a^2b(c+dx)^2 \coth(e+fx) + 3ab^2(c+dx)^2 \coth^2(e+fx) \\
&\quad + b^3(c+dx)^2 \coth^3(e+fx)) \, dx \\
&= \frac{a^3(c+dx)^3}{3d} + (3a^2b) \int (c+dx)^2 \coth(e+fx) \, dx \\
&\quad + (3ab^2) \int (c+dx)^2 \coth^2(e+fx) \, dx + b^3 \int (c+dx)^2 \coth^3(e+fx) \, dx \\
&= \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} - \frac{3ab^2(c+dx)^2 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \coth^2(e+fx)}{2f} - (6a^2b) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} \, dx \\
&\quad + (3ab^2) \int (c+dx)^2 \, dx + b^3 \int (c+dx)^2 \coth(e+fx) \, dx \\
&\quad + \frac{(6ab^2d) \int (c+dx) \coth(e+fx) \, dx}{f} + \frac{(b^3d) \int (c+dx) \coth^2(e+fx) \, dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} \\
&\quad + \frac{ab^2(c+dx)^3}{d} - \frac{b^3(c+dx)^3}{3d} - \frac{b^3d(c+dx)\coth(e+fx)}{f^2} \\
&\quad - \frac{3ab^2(c+dx)^2\coth(e+fx)}{f} - \frac{b^3(c+dx)^2\coth^2(e+fx)}{2f} \\
&\quad + \frac{3a^2b(c+dx)^2\log(1-e^{2(e+fx)})}{f} - (2b^3) \int \frac{e^{2(e+fx)}(c+dx)^2}{1-e^{2(e+fx)}} dx \\
&\quad + \frac{(b^3d^2) \int \coth(e+fx) dx}{f^2} - \frac{(6a^2bd) \int (c+dx) \log(1-e^{2(e+fx)}) dx}{f} \\
&\quad - \frac{(12ab^2d) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} dx}{f} + \frac{(b^3d) \int (c+dx) dx}{f} \\
&= \frac{b^3cdx}{f} + \frac{b^3d^2x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2b(c+dx)^3}{d} \\
&\quad + \frac{ab^2(c+dx)^3}{d} - \frac{b^3(c+dx)^3}{3d} - \frac{b^3d(c+dx)\coth(e+fx)}{f^2} \\
&\quad - \frac{3ab^2(c+dx)^2\coth(e+fx)}{f} - \frac{b^3(c+dx)^2\coth^2(e+fx)}{2f} \\
&\quad + \frac{6ab^2d(c+dx)\log(1-e^{2(e+fx)})}{f^2} + \frac{3a^2b(c+dx)^2\log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx)^2\log(1-e^{2(e+fx)})}{f} + \frac{b^3d^2\log(\sinh(e+fx))}{f^3} \\
&\quad + \frac{3a^2bd(c+dx)\text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{(3a^2bd^2) \int \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^2} \\
&\quad - \frac{(6ab^2d^2) \int \log(1-e^{2(e+fx)}) dx}{f^2} - \frac{(2b^3d) \int (c+dx) \log(1-e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2 b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} - \frac{b^3 d(c+dx) \coth(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \coth^2(e+fx)}{2f} + \frac{6ab^2 d(c+dx) \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2 b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3 d^2 \log(\sinh(e+fx))}{f^3} + \frac{3a^2 bd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad + \frac{b^3 d(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} - \frac{(3a^2 bd^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3} \\
&\quad - \frac{(3ab^2 d^2) \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{f^3} - \frac{(b^3 d^2) \int \text{PolyLog}(2, e^{2(e+fx)}) dx}{f^2} \\
&= \frac{b^3 c dx}{f} + \frac{b^3 d^2 x^2}{2f} - \frac{3ab^2(c+dx)^2}{f} + \frac{a^3(c+dx)^3}{3d} - \frac{a^2 b(c+dx)^3}{d} + \frac{ab^2(c+dx)^3}{d} \\
&\quad - \frac{b^3(c+dx)^3}{3d} - \frac{b^3 d(c+dx) \coth(e+fx)}{f^2} - \frac{3ab^2(c+dx)^2 \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx)^2 \coth^2(e+fx)}{2f} + \frac{6ab^2 d(c+dx) \log(1-e^{2(e+fx)})}{f^2} \\
&\quad + \frac{3a^2 b(c+dx)^2 \log(1-e^{2(e+fx)})}{f} + \frac{b^3(c+dx)^2 \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3 d^2 \log(\sinh(e+fx))}{f^3} + \frac{3ab^2 d^2 \text{PolyLog}(2, e^{2(e+fx)})}{f^3} \\
&\quad + \frac{3a^2 bd(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} + \frac{b^3 d(c+dx) \text{PolyLog}(2, e^{2(e+fx)})}{f^2} \\
&\quad - \frac{3a^2 bd^2 \text{PolyLog}(3, e^{2(e+fx)})}{2f^3} - \frac{(b^3 d^2) \text{Subst}\left(\int \frac{\text{PolyLog}(2,x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{b^3 c d x}{f} + \frac{b^3 d^2 x^2}{2 f} - \frac{3 a b^2 (c + d x)^2}{f} + \frac{a^3 (c + d x)^3}{3 d} - \frac{a^2 b (c + d x)^3}{d} + \frac{a b^2 (c + d x)^3}{d} \\
&\quad - \frac{b^3 (c + d x)^3}{3 d} - \frac{b^3 d (c + d x) \coth(e + f x)}{f^2} - \frac{3 a b^2 (c + d x)^2 \coth(e + f x)}{f} \\
&\quad - \frac{b^3 (c + d x)^2 \coth^2(e + f x)}{2 f} + \frac{6 a b^2 d (c + d x) \log(1 - e^{2(e + f x)})}{f^2} \\
&\quad + \frac{3 a^2 b (c + d x)^2 \log(1 - e^{2(e + f x)})}{f} + \frac{b^3 (c + d x)^2 \log(1 - e^{2(e + f x)})}{f} \\
&\quad + \frac{b^3 d^2 \log(\sinh(e + f x))}{f^3} + \frac{3 a b^2 d^2 \text{PolyLog}(2, e^{2(e + f x)})}{f^3} \\
&\quad + \frac{3 a^2 b d (c + d x) \text{PolyLog}(2, e^{2(e + f x)})}{f^2} + \frac{b^3 d (c + d x) \text{PolyLog}(2, e^{2(e + f x)})}{f^2} \\
&\quad - \frac{3 a^2 b d^2 \text{PolyLog}(3, e^{2(e + f x)})}{2 f^3} - \frac{b^3 d^2 \text{PolyLog}(3, e^{2(e + f x)})}{2 f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.47 (sec), antiderivative size = 585, normalized size of antiderivative = 1.46

$$\begin{aligned}
&\int (c + d x)^2 (a + b \coth(e + f x))^3 dx \\
&= \frac{-\frac{8 b e^{2 e} f x (9 a b d f (2 c + d x) + 3 a^2 f^2 (3 c^2 + 3 c d x + d^2 x^2) + b^2 (3 c^2 f^2 + 3 c d f^2 x + d^2 (3 + f^2 x^2)))}{-1 + e^{2 e}} + 12 b (6 a b d f (c + d x) + 3 a^2 f^2 (c + d x)^2) -}{}
\end{aligned}$$

```

[In] Integrate[(c + d*x)^2*(a + b*Coth[e + f*x])^3,x]
[Out] ((-8*b*E^(2*e))*f*x*(9*a*b*d*f*(2*c + d*x) + 3*a^2*f^2*(3*c^2 + 3*c*d*x + d^2*x^2) + b^2*(3*c^2*f^2 + 3*c*d*f^2*x + d^2*(3 + f^2*x^2))))/(-1 + E^(2*e)) + 12*b*(6*a*b*d*f*(c + d*x) + 3*a^2*f^2*(c + d*x)^2 + b^2*(c^2*f^2 + 2*c*d*f^2*x + d^2*(1 + f^2*x^2)))*Log[1 - E^(2*(e + f*x))] + 12*b*d*(3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*PolyLog[2, E^(2*(e + f*x))] - 6*b*(3*a^2 + b^2)*d^2*PolyLog[3, E^(2*(e + f*x))] + f*Csch[e]*Csch[e + f*x]^2*(-2*b*(9*a*b*f*(c + d*x)^2 + 3*a^2*f^2*x*(3*c^2 + 3*c*d*x + d^2*x^2) + b^2*(3*c^2*f^2*x + d^2*x^2*(3 + f^2*x^2)))*Cosh[e] + b*(18*a*b*f*(c + d*x)^2 + 3*a^2*f^2*x*(3*c^2 + 3*c*d*x + d^2*x^2) + b^2*(3*c^2*f^2*x + 3*c*d*(2 + f^2*x^2) + d^2*x*(6 + f^2*x^2)))*Cosh[e + 2*f*x] + f*(b*(3*a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[3*e + 2*f*x] - 2*(3*b^3*(c + d*x)^2 + a^3*f*x*(3*c^2 + 3*c*d*x + d^2*x^2) + 3*a*b^2*f*x*(3*c^2 + 3*c*d*x + d^2*x^2) - a*(a^2 + 3*b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[2*(e + f*x)]])*Sinh[e]))/(12*f^3)

```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1585 vs. $2(389) = 778$.

Time = 0.63 (sec), antiderivative size = 1586, normalized size of antiderivative = 3.96

method	result	size
risch	Expression too large to display	1586

[In] `int((d*x+c)^2*(a+b*cOTH(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]

```

2/f^2*b^3*d^2*polylog(2,exp(f*x+e))*x+1/f*b^3*d^2*ln(1+exp(f*x+e))*x^2+2/f^
2*b^3*d^2*polylog(2,-exp(f*x+e))*x+1/f^3*b^3*e^2*d^2*ln(exp(f*x+e)-1)-1/f^3
*b^3*d^2*ln(1-exp(f*x+e))*e^2+3/f*b*a^2*c^2*ln(exp(f*x+e)-1)+3/f*b*a^2*c^2*
ln(1+exp(f*x+e))+6/f^3*b^2*a*d^2*polylog(2,exp(f*x+e))+6/f^3*b^2*a*d^2*poly
log(2,-exp(f*x+e))-6/f^3*b*a^2*d^2*polylog(3,exp(f*x+e))+4/3/f^3*b^3*d^2*e^
3-2/f^3*b^3*d^2*ln(exp(f*x+e))-2/f*b^3*c^2*ln(exp(f*x+e))-4/f*b^3*c*d*e*x-1
2/f*b*d*c*a^2*e*x+12/f^2*b*e*d*c*a^2*ln(exp(f*x+e))-6/f^3*b*a^2*d^2*polylog
(3,-exp(f*x+e))+2/f^2*b^3*c*d*polylog(2,exp(f*x+e))+2/f^2*b^3*c*d*polylog(2
,-exp(f*x+e))+1/f*b^3*d^2*ln(1-exp(f*x+e))*x^2+4/f^3*b*a^2*d^2*e^3+2/f^2*b^
3*d^2*e^2*x-6/f*b^2*a*d^2*x^2-2/f^2*b^3*c*d*e^2-6/f^3*b^2*a*d^2*e^2-2/f^3*b^
3*e^2*d^2*ln(exp(f*x+e))-6/f*b*a^2*c^2*ln(exp(f*x+e))+6/f^2*b*a^2*d^2*e^2*x
-6/f^2*b*d*c*a^2*e^2-12/f^2*b^2*a*d^2*e*x+12/f^3*b^2*e*a*d^2*ln(exp(f*x+e))
-12/f^2*b^2*a*c*d*ln(exp(f*x+e))+4/f^2*b^3*c*d*ln(exp(f*x+e))-6/f^3*b*e^
2*a^2*d^2*ln(exp(f*x+e))-c*d*x^2*b^3-1/3*d^2*x^3*b^3+x*b^3*c^2+1/3/d*b^3*c^
3-3*d*a^2*b*c*x^2+3*d*a*b^2*c*x^2+3*a^2*b*c^2*x+3*a*b^2*c^2*x-2*b^2*(3*a*d^
2*f*x^2*exp(2*f*x+2*e)+b*d^2*f*x^2*exp(2*f*x+2*e)+6*a*c*d*f*x*exp(2*f*x+2*e)
+2*b*c*d*f*x*exp(2*f*x+2*e)+3*a*c^2*f*exp(2*f*x+2*e)-3*a*d^2*f*x^2+b*c^2*f^
*exp(2*f*x+2*e)+b*d^2*x*exp(2*f*x+2*e)-6*a*c*d*f*x*exp(2*f*x+2*e)*d*b*c-3*a
*c^2*f-b*d^2*x-b*c*d)/f^2/(exp(2*f*x+2*e)-1)^2+6/f*b*d*c*a^2*ln(1-exp(f*x+e))
*x+6/f^2*b*d*c*a^2*ln(1-exp(f*x+e))*e+6/f*b*d*c*a^2*ln(1+exp(f*x+e))*x-6/
f^2*b*a^2*c*d*e*ln(exp(f*x+e)-1)-d^2*a^2*b*x^3+d^2*a*b^2*x^3+d*a^3*c*x^2+a^
3*c^2*x+1/d*a^2*b*c^3+1/d*a*b^2*c^3+1/f*b^3*c^2*ln(exp(f*x+e)-1)+1/f*b^3*c^
2*ln(1+exp(f*x+e))+1/f^3*b^3*d^2*ln(exp(f*x+e)-1)+1/f^3*b^3*d^2*ln(1+exp(f*
x+e))-2/f^3*b^3*d^2*polylog(3,exp(f*x+e))-2/f^3*b^3*d^2*polylog(3,-exp(f*x+
e))+1/3*d^2*a^3*x^3+1/3/d*a^3*c^3+f^3*b*a^2*d^2*e^2*ln(exp(f*x+e)-1)-3/f^
3*b*a^2*d^2*ln(1-exp(f*x+e))*e^2+3/f*b*a^2*d^2*ln(1-exp(f*x+e))*x^2+6/f^2*b^
*a^2*d^2*polylog(2,exp(f*x+e))*x+3/f*b*a^2*d^2*ln(1+exp(f*x+e))*x^2+6/f^2*b^
*a^2*d^2*polylog(2,-exp(f*x+e))*x-6/f^3*b^2*e*a*d^2*ln(exp(f*x+e)-1)+6/f^3*
b^2*a*d^2*ln(1-exp(f*x+e))*e-2/f^2*b^3*e*c*d*ln(exp(f*x+e)-1)+2/f*b^3*c*d*l
n(1-exp(f*x+e))*x+2/f^2*b^3*c*d*ln(1-exp(f*x+e))*e+2/f*b^3*c*d*ln(1+exp(f*x
+e))*x+6/f^2*b^2*a*d^2*ln(1-exp(f*x+e))*x+6/f^2*b^2*a*d^2*ln(1+exp(f*x+e))*
x+6/f^2*b^2*a*c*d*ln(exp(f*x+e)-1)+6/f^2*b^2*a*c*d*ln(1+exp(f*x+e))+6/f^2*b^
*d*c*a^2*polylog(2,exp(f*x+e))+6/f^2*b*d*c*a^2*polylog(2,-exp(f*x+e)))

```

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6356 vs. $2(386) = 772$.

Time = 0.34 (sec), antiderivative size = 6356, normalized size of antiderivative = 15.85

$$\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^2*(a+b*coth(f*x+e))^3,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F]

$$\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = \int (a + b \coth(e + fx))^3 (c + dx)^2 dx$$

[In] `integrate((d*x+c)**2*(a+b*coth(f*x+e))**3,x)`

[Out] `Integral((a + b*coth(e + f*x))**3*(c + d*x)**2, x)`

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 997 vs. $2(386) = 772$.

Time = 0.28 (sec), antiderivative size = 997, normalized size of antiderivative = 2.49

$$\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^2*(a+b*coth(f*x+e))^3,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & 1/3*a^3*d^2*x^3 + a^3*c*d*x^2 + a^3*c^2*x + 3*a^2*b*c^2*log(\sinh(f*x + e))/f + 1/3*(18*a*b^2*c^2*f + 6*b^3*c*d + (3*a^2*b*d^2*f^2 + 3*a*b^2*d^2*f^2 + b^3*d^2*f^2)*x^3 + 3*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 + 3*(c*d*f^2 + 2*d^2*f)*a*b^2 + 3*(3*(c^2*f^2 + 4*c*d*f)*a*b^2 + (c^2*f^2 + 2*d^2)*b^3)*x + (3*a^2*b*d^2*f^2*e^(4*e) + 3*a*b^2*d^2*f^2*e^(4*e) + b^3*d^2*f^2*e^(4*e))*x^3 + 3*(3*a^2*b*c*d*f^2*e^(4*e) + 3*a*b^2*c*d*f^2*e^(4*e) + b^3*c*d*f^2*e^(4*e))*x^2 + 3*(3*a*b^2*c^2*f^2*e^(4*e) + b^3*c^2*f^2*e^(4*e))*x)*e^(4*f*x) - 2*(9*a*b^2*c^2*f*e^(2*e) + 3*(c^2*f*e^(2*e) + c*d*f*e^(2*e))*b^3 + (3*a^2*b*d^2*f^2*e^(2*e) + 3*a*b^2*d^2*f^2*e^(2*e) + b^3*d^2*f^2*e^(2*e))*x^3 + 3*(3*a^2*b*c*d*f^2*e^(2*e) + 3*(c*d*f^2*e^(2*e) + d^2*f*e^(2*e))*a*b^2 + (c*d*f^2*e^(2*e) + d^2*f*e^(2*e))*b^3)*x^2 + 3*(3*(c^2*f^2*e^(2*e) + 2*c*d*f^2*e^(2*e) + d^2*f^2*e^(2*e))*b^3)*x)*e^(2*f*x))/(f^2*e^(4*f*x + 4*e) - 2*f^2*e^(2*f*x + 2*e) + f^2) - 2*(6*a*b^2*c^2*f^2*e^(2*f*x + 4*e) + 3*(3*(c^2*f^2*e^(2*f*x + 4*e) + 2*c*d*f^2*e^(2*f*x + 4*e) + d^2*f^2*e^(2*f*x + 4*e))*b^3)*x)*e^(2*f*x) \end{aligned}$$

$$\begin{aligned}
& d*f + (c^2*f^2 + d^2)*b^3*x/f^2 + (3*a^2*b*d^2 + b^3*d^2)*(f^2*x^2*\log(e^(f*x + e) + 1) + 2*f*x*dilog(-e^(f*x + e)) - 2*polylog(3, -e^(f*x + e)))/f^3 \\
& + (3*a^2*b*d^2 + b^3*d^2)*(f^2*x^2*\log(-e^(f*x + e) + 1) + 2*f*x*dilog(e^(f*x + e)) - 2*polylog(3, e^(f*x + e)))/f^3 + 2*(3*a^2*b*c*d*f + b^3*c*d*f + 3*a*b^2*d^2)*(f*x*\log(e^(f*x + e) + 1) + dilog(-e^(f*x + e)))/f^3 + 2*(3*a^2*b*c*d*f + b^3*c*d*f + 3*a*b^2*d^2)*(f*x*\log(-e^(f*x + e) + 1) + dilog(e^(f*x + e)))/f^3 + (6*a*b^2*c*d*f + (c^2*f^2 + d^2)*b^3)*\log(e^(f*x + e) + 1)/f^3 + (6*a*b^2*c*d*f + (c^2*f^2 + d^2)*b^3)*\log(e^(f*x + e) - 1)/f^3 - 2/3*((3*a^2*b*d^2 + b^3*d^2)*f^3*x^3 + 3*(3*a^2*b*c*d*f + b^3*c*d*f + 3*a*b^2*d^2)*f^2*x^2)/f^3
\end{aligned}$$

Giac [F]

$$\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = \int (dx + c)^2 (b \coth(fx + e) + a)^3 dx$$

[In] `integrate((d*x+c)^2*(a+b*coth(f*x+e))^3,x, algorithm="giac")`
[Out] `integrate((d*x + c)^2*(b*coth(f*x + e) + a)^3, x)`

Mupad [F(-1)]

Timed out.

$$\int (c + dx)^2 (a + b \coth(e + fx))^3 dx = \int (a + b \coth(e + f x))^3 (c + d x)^2 dx$$

[In] `int((a + b*coth(e + f*x))^3*(c + d*x)^2,x)`
[Out] `int((a + b*coth(e + f*x))^3*(c + d*x)^2, x)`

3.49 $\int (c + dx)(a + b \coth(e + fx))^3 dx$

Optimal result	331
Rubi [A] (verified)	332
Mathematica [A] (verified)	335
Maple [B] (verified)	336
Fricas [B] (verification not implemented)	336
Sympy [F]	338
Maxima [B] (verification not implemented)	338
Giac [F]	339
Mupad [F(-1)]	339

Optimal result

Integrand size = 18, antiderivative size = 261

$$\begin{aligned}
\int (c + dx)(a + b \coth(e + fx))^3 dx = & 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c + dx)^2}{2d} \\
& - \frac{3a^2b(c + dx)^2}{2d} - \frac{b^3(c + dx)^2}{2d} \\
& - \frac{b^3d \coth(e + fx)}{2f^2} - \frac{3ab^2(c + dx) \coth(e + fx)}{f} \\
& - \frac{b^3(c + dx) \coth^2(e + fx)}{2f} \\
& + \frac{3a^2b(c + dx) \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{b^3(c + dx) \log(1 - e^{2(e+fx)})}{f} \\
& + \frac{3ab^2d \log(\sinh(e + fx))}{f^2} \\
& + \frac{3a^2bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\
& + \frac{b^3d \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2}
\end{aligned}$$

[Out] $3*a*b^2*c*x+1/2*b^3*d*x/f+3/2*a*b^2*d*x^2+1/2*a^3*(d*x+c)^2/d-3/2*a^2*b*(d*x+c)^2/d-1/2*b^3*(d*x+c)^2/d-1/2*b^3*d*\coth(f*x+e)/f^2-3*a*b^2*(d*x+c)*\coth(f*x+e)/f-1/2*b^3*(d*x+c)*\coth(f*x+e)^2/f+3*a^2*b*(d*x+c)*\ln(1-\exp(2*f*x+2*e))/f+b^3*(d*x+c)*\ln(1-\exp(2*f*x+2*e))/f+3*a*b^2*d*\ln(\sinh(f*x+e))/f^2+3/2*a^2*b*d*\operatorname{polylog}(2, \exp(2*f*x+2*e))/f^2+1/2*b^3*d*\operatorname{polylog}(2, \exp(2*f*x+2*e))/f^2$

Rubi [A] (verified)

Time = 0.27 (sec) , antiderivative size = 261, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {3803, 3797, 2221, 2317, 2438, 3801, 3556, 3554, 8}

$$\begin{aligned} \int (c + dx)(a + b \coth(e + fx))^3 dx = & \frac{a^3(c + dx)^2}{2d} + \frac{3a^2b(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{3a^2b(c + dx)^2}{2d} + \frac{3a^2bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} \\ & - \frac{3ab^2(c + dx) \coth(e + fx)}{f} + 3ab^2cx \\ & + \frac{3ab^2d \log(\sinh(e + fx))}{f^2} + \frac{3}{2}ab^2dx^2 \\ & + \frac{b^3(c + dx) \log(1 - e^{2(e+fx)})}{f} \\ & - \frac{b^3(c + dx) \coth^2(e + fx)}{2f} - \frac{b^3(c + dx)^2}{2d} \\ & + \frac{b^3d \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{b^3d \coth(e + fx)}{2f^2} + \frac{b^3dx}{2f} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)*(a + b*\operatorname{Coth}[e + f*x])^3, x]$

[Out]
$$\begin{aligned} & 3*a*b^2*c*x + (b^3*d*x)/(2*f) + (3*a*b^2*d*x^2)/2 + (a^3*(c + d*x)^2)/(2*d) \\ & - (3*a^2*b*(c + d*x)^2)/(2*d) - (b^3*(c + d*x)^2)/(2*d) - (b^3*d*\operatorname{Coth}[e + f*x])/(2*f^2) - (3*a*b^2*(c + d*x)*\operatorname{Coth}[e + f*x])/f - (b^3*(c + d*x)*\operatorname{Coth}[e + f*x]^2)/(2*f) + (3*a^2*b*(c + d*x)*\operatorname{Log}[1 - E^{(2*(e + f*x))}])/f + (b^3*(c + d*x)*\operatorname{Log}[1 - E^{(2*(e + f*x))}])/f + (3*a*b^2*d*\operatorname{Log}[\operatorname{Sinh}[e + f*x]])/f^2 + (3*a^2*b*d*\operatorname{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2) + (b^3*d*\operatorname{PolyLog}[2, E^{(2*(e + f*x))}])/(2*f^2) \end{aligned}$$

Rule 8

$\operatorname{Int}[a_, x_{\text{Symbol}}] :> \operatorname{Simp}[a*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 2221

$\operatorname{Int}[(((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}*((c_*) + (d_*)*(x_))^{(m_*)})/((a_) + (b_*)*((F_*)^((g_*)*((e_*) + (f_*)*(x_))))^{(n_*)}), x_{\text{Symbol}}] :> \operatorname{Simp}[((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^((g*(e + f*x)))^n/a)], x] - \operatorname{Dist}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^(m - 1)*\operatorname{Log}[1 + b*((F^((g*(e + f*x)))^n/a)], x], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \operatorname{IGtQ}[m, 0]$

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*(c_.) + (d_)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[((c_.)*(d_) + (e_.)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3554

```
Int[((b_.)*tan[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3556

```
Int[tan[(c_.) + (d_)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3797

```
Int[((c_.) + (d_)*(x_))^(m_)*tan[(e_.) + Pi*(k_.) + (Complex[0, fz_])*f_(x_)], x_Symbol] :> Simp[(-I)*((c + d*x)^(m + 1)/(d*(m + 1))), x] + Dist[2*I, Int[((c + d*x)^m*(E^(2*(-I)*e + f*fz*x))/(1 + E^(2*(-I)*e + f*fz*x))/E^(2*I*k*Pi))]/E^(2*I*k*Pi), x] /; FreeQ[{c, d, e, f, fz}, x] && IntegerQ[4*k] && IGtQ[m, 0]
```

Rule 3801

```
Int[((c_.) + (d_)*(x_))^(m_)*((b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[b*(c + d*x)^m*((b*Tan[e + f*x])^(n - 1)/(f*(n - 1))), x] + (-Dist[b*d*(m/(f*(n - 1))), Int[(c + d*x)^(m - 1)*(b*Tan[e + f*x])^(n - 1), x], x] - Dist[b^2, Int[(c + d*x)^m*(b*Tan[e + f*x])^(n - 2), x], x]) /; FreeQ[{b, c, d, e, f}, x] && GtQ[n, 1] && GtQ[m, 0]
```

Rule 3803

```
Int[((c_.) + (d_)*(x_))^(m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (a + b*Tan[e + f*x])^n, x], x] /; FreeQ[{a, b, c, d, e, f, m}, x] && IGtQ[m, 0] && IGtQ[n, 0]
```

Rubi steps

integral = $\int (a^3(c + dx) + 3a^2b(c + dx) \coth(e + fx) + 3ab^2(c + dx) \coth^2(e + fx) + b^3(c + dx) \coth^3(e + fx)) dx$

$$\begin{aligned}
&= \frac{a^3(c+dx)^2}{2d} + (3a^2b) \int (c+dx) \coth(e+fx) dx \\
&\quad + (3ab^2) \int (c+dx) \coth^2(e+fx) dx + b^3 \int (c+dx) \coth^3(e+fx) dx \\
&= \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{3ab^2(c+dx) \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx) \coth^2(e+fx)}{2f} - (6a^2b) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} dx \\
&\quad + (3ab^2) \int (c+dx) dx + b^3 \int (c+dx) \coth(e+fx) dx \\
&\quad + \frac{(3ab^2d) \int \coth(e+fx) dx}{f} + \frac{(b^3d) \int \coth^2(e+fx) dx}{2f} \\
&= 3ab^2cx + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{b^3(c+dx)^2}{2d} \\
&\quad - \frac{b^3d \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \coth(e+fx)}{f} - \frac{b^3(c+dx) \coth^2(e+fx)}{2f} \\
&\quad + \frac{3a^2b(c+dx) \log(1-e^{2(e+fx)})}{f} + \frac{3ab^2d \log(\sinh(e+fx))}{f^2} \\
&\quad - (2b^3) \int \frac{e^{2(e+fx)}(c+dx)}{1-e^{2(e+fx)}} dx - \frac{(3a^2bd) \int \log(1-e^{2(e+fx)}) dx}{f} + \frac{(b^3d) \int 1 dx}{2f} \\
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} \\
&\quad - \frac{b^3(c+dx)^2}{2d} - \frac{b^3d \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx) \coth^2(e+fx)}{2f} + \frac{3a^2b(c+dx) \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx) \log(1-e^{2(e+fx)})}{f} + \frac{3ab^2d \log(\sinh(e+fx))}{f^2} \\
&\quad - \frac{(3a^2bd) \text{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} - \frac{(b^3d) \int \log(1-e^{2(e+fx)}) dx}{f}
\end{aligned}$$

$$\begin{aligned}
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} \\
&\quad - \frac{b^3(c+dx)^2}{2d} - \frac{b^3d \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \coth(e+fx)}{f} \\
&\quad - \frac{b^3(c+dx) \coth^2(e+fx)}{2f} + \frac{3a^2b(c+dx) \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{b^3(c+dx) \log(1-e^{2(e+fx)})}{f} + \frac{3ab^2d \log(\sinh(e+fx))}{f^2} \\
&\quad + \frac{3a^2bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} - \frac{(b^3d) \operatorname{Subst}\left(\int \frac{\log(1-x)}{x} dx, x, e^{2(e+fx)}\right)}{2f^2} \\
&= 3ab^2cx + \frac{b^3dx}{2f} + \frac{3}{2}ab^2dx^2 + \frac{a^3(c+dx)^2}{2d} - \frac{3a^2b(c+dx)^2}{2d} - \frac{b^3(c+dx)^2}{2d} \\
&\quad - \frac{b^3d \coth(e+fx)}{2f^2} - \frac{3ab^2(c+dx) \coth(e+fx)}{f} - \frac{b^3(c+dx) \coth^2(e+fx)}{2f} \\
&\quad + \frac{3a^2b(c+dx) \log(1-e^{2(e+fx)})}{f} + \frac{b^3(c+dx) \log(1-e^{2(e+fx)})}{f} \\
&\quad + \frac{3ab^2d \log(\sinh(e+fx))}{f^2} + \frac{3a^2bd \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2} + \frac{b^3d \operatorname{PolyLog}(2, e^{2(e+fx)})}{2f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 8.18 (sec), antiderivative size = 318, normalized size of antiderivative = 1.22

$$\begin{aligned}
&\int (c+dx)(a+b \coth(e+fx))^3 dx \\
&= \frac{(a+b \coth(e+fx))^3 \sinh(e+fx) \left(-b^3 f (c+dx) - a(a^2+3b^2) (e+fx) (-2cf+d(e-fx)) \sinh^2(e+fx)\right)}{2f^2}
\end{aligned}$$

[In] `Integrate[(c + d*x)*(a + b*Coth[e + f*x])^3, x]`

[Out] `((a + b*Coth[e + f*x])^3*Sinh[e + f*x]*(-(b^3*f*(c + d*x)) - a*(a^2 + 3*b^2)*((a + b*Coth[e + f*x])*(-2*c*f + d*(e - f*x))*Sinh[e + f*x]^2 + 2*b*((3*a^2*f^2*(c + d*x)^2)/(2*d) + (b^2*f^2*(c + d*x)^2)/(2*d) + 3*a*b*d*(e + f*x) + (3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*Log[1 - E^(-e - f*x)] + (3*a*b*d + 3*a^2*f*(c + d*x) + b^2*f*(c + d*x))*Log[1 + E^(-e - f*x)] - (3*a^2 + b^2)*d*PolyLog[2, -E^(-e - f*x)] - (3*a^2 + b^2)*d*PolyLog[2, E^(-e - f*x)])*Sinh[e + f*x]^2 - (b^2*(b*d + 6*a*f*(c + d*x))*Sinh[2*(e + f*x)]/2))/(2*f^2*(b*Cosh[e + f*x] + a*Sinh[e + f*x])^3)`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 650 vs. $2(243) = 486$.

Time = 0.49 (sec), antiderivative size = 651, normalized size of antiderivative = 2.49

method	result
risch	$-\frac{b^3 ed \ln(e^{fx+e}-1)}{f^2} + \frac{3b^2 da \ln(e^{fx+e}-1)}{f^2} + \frac{3b^2 da \ln(1+e^{fx+e})}{f^2} + \frac{3bd a^2 \text{polylog}(2, e^{fx+e})}{f^2} + \frac{3bd a^2 \text{polylog}(2, -e^{fx+e})}{f^2} + \dots$

[In] `int((d*x+c)*(a+b*cOTH(f*x+e))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/f^2 * b^3 * e * d * \ln(\exp(f*x+e)-1) + 3/f^2 * b^2 * d * a * \ln(\exp(f*x+e)-1) + 3/f^2 * b^2 * d * a^2 * \ln(1+\exp(f*x+e)) + 3/f^2 * b^2 * d * a^2 * \text{polylog}(2, \exp(f*x+e)) + 3/f^2 * b^2 * d * a^2 * \text{polylog}(2, -\exp(f*x+e)) + 1/f^2 * b^3 * d * \ln(1-\exp(f*x+e)) * x + 1/f^2 * b^3 * d * \ln(1+\exp(f*x+e)) * x - 3/2 * a^2 * b * d * x^2 + 3 * a^2 * b * c * x + 1/f^2 * b^3 * d * \text{polylog}(2, -\exp(f*x+e)) - 6/f^2 * b * a^2 * d * e * x + 6/f^2 * b * e * a^2 * d * \ln(\exp(f*x+e)) - 1/f^2 * b^3 * d * e^2 - 2/f^2 * b^3 * c * \ln(\exp(f*x+e)) + 1/f^2 * b^3 * c * \ln(\exp(f*x+e)) + 1/f^2 * b^3 * d * \text{polylog}(2, \exp(f*x+e)) + 3 * a * b^2 * c * x + 3/2 * a * b^2 * d * x^2 - 2/f^2 * b^3 * d * e * x - 3/f^2 * b * a^2 * d * e^2 - 6/f^2 * b * a^2 * c * ln(\exp(f*x+e)) + 2/f^2 * b^3 * e * d * \ln(\exp(f*x+e)) - 6/f^2 * b^2 * d * a * \ln(\exp(f*x+e)) - b^2 * (6 * a * d * f * x * \exp(2 * f * x + 2 * e) + 2 * b * d * f * x * \exp(2 * f * x + 2 * e) + 6 * a * c * f * \exp(2 * f * x + 2 * e) + 2 * b * c * f * \exp(2 * f * x + 2 * e) - 6 * a * d * f * x * \exp(2 * f * x + 2 * e) * d * b - 6 * a * c * f * b * d) / f^2 / (\exp(2 * f * x + 2 * e) - 1)^2 + 1/f^2 * b^3 * d * \ln(1 - \exp(f*x+e)) * e + 3/f^2 * b * a^2 * c * \ln(\exp(f*x+e) - 1) + 3/f^2 * b * a^2 * c * \ln(1 + \exp(f*x+e)) + 1/2 * a^3 * d * x^2 - 1/2 * b^3 * d * x^2 + a^3 * c * x + b^3 * c * x - 3/f^2 * b * d * a^2 * e * \ln(\exp(f*x+e) - 1) + 3/f^2 * b * d * a^2 * \ln(1 - \exp(f*x+e)) * e + 3/f^2 * b * d * a^2 * \ln(1 - \exp(f*x+e)) * x + 3/f^2 * b * d * a^2 * \ln(1 + \exp(f*x+e)) * x \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 2907 vs. $2(241) = 482$.

Time = 0.30 (sec), antiderivative size = 2907, normalized size of antiderivative = 11.14

$$\int (c + dx)(a + b \coth(e + fx))^3 dx = \text{Too large to display}$$

[In] `integrate((d*x+c)*(a+b*cOTH(f*x+e))^3,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/2 * ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2*x + ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*c*f^2*x) * \cosh(f*x + e)^4 + 4*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2*x) * \cosh(f*x + e)^4 + 4*(a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2*x) * \cosh(f*x + e) * \sinh(f*x + e)^3 + ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2*x) * \sinh(f*x + e)^3) * \cosh(f*x + e) \end{aligned}$$

$$\begin{aligned}
& \text{nh(f*x + e)}^4 + 2*b^3*d + 2*(3*a^2*b + b^3)*d*e^2 - 2*((a^3 - 3*a^2*b + 3*a \\
& *b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + 2*(3*a^2*b + b^3)*d*e^2 - 2* \\
& (2*(3*a^2*b + b^3)*c*e - (3*a*b^2 + b^3)*c)*f + 2*((a^3 - 3*a^2*b + 3*a*b^2 \\
& - b^3)*c*f^2 - (3*a*b^2 - b^3)*d*f)*x)*cosh(f*x + e)^2 - 2*((a^3 - 3*a^2*b \\
& + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + 2*(3*a^2*b + b^3)*d*e^2 \\
& - 3*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b \\
& + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e*f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b \\
& + 3*a*b^2 - b^3)*c*f^2)*x)*cosh(f*x + e)^2 - 2*(2*(3*a^2*b + b^3)*c*e - (3*a \\
& *b^2 + b^3)*c)*f + 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2 - (3*a*b^2 - b \\
& ^3)*d*f)*x)*sinh(f*x + e)^2 + 4*(3*a*b^2*c - (3*a^2*b + b^3)*c*e)*f + 2*((3 \\
& *a^2*b + b^3)*d*cosh(f*x + e)^4 + 4*(3*a^2*b + b^3)*d*cosh(f*x + e)*sinh(f*x \\
& + e)^3 + (3*a^2*b + b^3)*d*sinh(f*x + e)^4 - 2*(3*a^2*b + b^3)*d*cosh(f*x \\
& + e)^2 + 2*(3*(3*a^2*b + b^3)*d*cosh(f*x + e)^2 - (3*a^2*b + b^3)*d)*sinh(f*x \\
& + e)^2 + (3*a^2*b + b^3)*d + 4*((3*a^2*b + b^3)*d*cosh(f*x + e)^3 - (3*a \\
& ^2*b + b^3)*d*cosh(f*x + e))*sinh(f*x + e))*dilog(cosh(f*x + e) + sinh(f*x \\
& + e)) + 2*((3*a^2*b + b^3)*d*cosh(f*x + e)^4 + 4*(3*a^2*b + b^3)*d*cosh(f*x \\
& + e)*sinh(f*x + e)^3 + (3*a^2*b + b^3)*d*sinh(f*x + e)^4 - 2*(3*a^2*b + b \\
& ^3)*d*cosh(f*x + e)^2 + 2*(3*(3*a^2*b + b^3)*d*cosh(f*x + e)^2 - (3*a^2*b + b \\
& ^3)*d)*sinh(f*x + e)^2 + (3*a^2*b + b^3)*d + 4*((3*a^2*b + b^3)*d*cosh(f*x \\
& + e)^3 - (3*a^2*b + b^3)*d*cosh(f*x + e))*sinh(f*x + e))*dilog(-cosh(f*x \\
& + e) - sinh(f*x + e)) + 2*((3*a*b^2*d + (3*a^2*b + b^3)*d*f*x + (3*a^2*b + b \\
& ^3)*c*f)*cosh(f*x + e)^4 + 4*(3*a*b^2*d + (3*a^2*b + b^3)*d*f*x + (3*a^2*b \\
& + b^3)*c*f)*cosh(f*x + e)*sinh(f*x + e)^3 + (3*a*b^2*d + (3*a^2*b + b^3)*d \\
& *f*x + (3*a^2*b + b^3)*c*f)*sinh(f*x + e)^4 + 3*a*b^2*d + (3*a^2*b + b^3)*d \\
& *f*x + (3*a^2*b + b^3)*c*f - 2*(3*a*b^2*d + (3*a^2*b + b^3)*d*f*x + (3*a^2*b \\
& + b^3)*c*f)*cosh(f*x + e)^2 - 2*(3*a*b^2*d + (3*a^2*b + b^3)*d*f*x + (3*a \\
& ^2*b + b^3)*c*f - 3*(3*a*b^2*d + (3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*c \\
& f)*cosh(f*x + e)^2)*sinh(f*x + e)^2 + 4*((3*a*b^2*d + (3*a^2*b + b^3)*d*f*x \\
& + (3*a^2*b + b^3)*c*f)*cosh(f*x + e)^3 - (3*a*b^2*d + (3*a^2*b + b^3)*d*f*x \\
& + (3*a^2*b + b^3)*c*f)*cosh(f*x + e)) *log(cosh(f*x + e) + sinh(f*x + e) + \\
& 1) + 2*((3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c*f)*cosh(f*x \\
& + e)^4 + 4*(3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c \\
& *f)*cosh(f*x + e)*sinh(f*x + e)^3 + (3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a \\
& ^2*b + b^3)*c*f)*sinh(f*x + e)^4 + 3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a \\
& ^2*b + b^3)*c*f - 2*(3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c \\
& f)*cosh(f*x + e)^2 - 2*(3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c \\
& f - 3*(3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c*f)*cosh(f*x + e) \\
&) *sinh(f*x + e)^2 + 4*((3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c \\
& *f)*cosh(f*x + e)^3 - (3*a*b^2*d - (3*a^2*b + b^3)*d*e + (3*a^2*b + b^3)*c \\
& *f)*cosh(f*x + e))*sinh(f*x + e)) *log(cosh(f*x + e) + sinh(f*x + e) - 1) + \\
& 2*((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e)^4 + 4*((3*a \\
& ^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e)*sinh(f*x + e)^3 + ((3 \\
& *a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*sinh(f*x + e)^4 + (3*a^2*b + b^3) \\
&) *d*f*x + (3*a^2*b + b^3)*d*e - 2*((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)* \\
& d*e)*cosh(f*x + e)^2 - 2*((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e - 3*(3 \\
& *a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)
\end{aligned}$$

$$\begin{aligned}
& (3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e)^2)*sinh(f*x + e) \\
& ^2 + 4*((3*a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e)^3 - ((3 \\
& *a^2*b + b^3)*d*f*x + (3*a^2*b + b^3)*d*e)*cosh(f*x + e))*sinh(f*x + e))*lo \\
& g(-cosh(f*x + e) - sinh(f*x + e) + 1) + 4*((a^3 - 3*a^2*b + 3*a*b^2 - b^3) \\
& *d*f^2*x^2 - 12*a*b^2*d*e + 2*(3*a^2*b + b^3)*d*e^2 - 4*(3*a^2*b + b^3)*c*e \\
& *f - 2*(6*a*b^2*d*f - (a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2)*x)*cosh(f*x + \\
& e)^3 - ((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*d*f^2*x^2 - 12*a*b^2*d*e + b^3*d + \\
& 2*(3*a^2*b + b^3)*d*e^2 - 2*(2*(3*a^2*b + b^3)*c*e - (3*a*b^2 + b^3)*c)*f + \\
& 2*((a^3 - 3*a^2*b + 3*a*b^2 - b^3)*c*f^2 - (3*a*b^2 - b^3)*d*f)*x)*cosh(f*x + \\
& e))*sinh(f*x + e))/(f^2*cosh(f*x + e)^4 + 4*f^2*cosh(f*x + e)*sinh(f*x + e)^3 + f^2*sinh(f*x + e)^4 - 2*f^2*cosh(f*x + e)^2 + 2*(3*f^2*cosh(f*x + e)^2 - f^2)*sinh(f*x + e)^2 + f^2 + 4*(f^2*cosh(f*x + e)^3 - f^2*cosh(f*x + e))*sinh(f*x + e))
\end{aligned}$$

Sympy [F]

$$\int (c + dx)(a + b \coth(e + fx))^3 dx = \int (a + b \coth(e + fx))^3 (c + dx) dx$$

```
[In] integrate((d*x+c)*(a+b*coth(f*x+e))**3,x)
[Out] Integral((a + b*coth(e + f*x))**3*(c + d*x), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 528 vs. $2(241) = 482$.

Time = 0.27 (sec) , antiderivative size = 528, normalized size of antiderivative = 2.02

$$\begin{aligned}
& \int (c + dx)(a + b \coth(e + fx))^3 dx \\
& = \frac{1}{2} a^3 dx^2 + a^3 c x + \frac{3 a^2 b c \log(\sinh(f x + e))}{f} - (3 a^2 b d + b^3 d) x^2 - \frac{2 (b^3 c f + 3 a b^2 d) x}{f} \\
& + \frac{12 a b^2 c f + 2 b^3 d + (3 a^2 b d f^2 + 3 a b^2 d f^2 + b^3 d f^2) x^2 + 2 (b^3 c f^2 + 3 (c f^2 + 2 d f) a b^2) x + ((3 a^2 b d f^2 e^{(4 e)} + 3 a^3 c f^2 + 3 a^2 b^2 d f^2 + b^4 d^2) x^3 + (3 a^2 b d f^2 e^{(4 e)} + 3 a^3 c f^2 + 3 a^2 b^2 d f^2 + b^4 d^2) x^2 + (3 a^2 b d f^2 e^{(4 e)} + 3 a^3 c f^2 + 3 a^2 b^2 d f^2 + b^4 d^2) x + (3 a^2 b d f^2 e^{(4 e)} + 3 a^3 c f^2 + 3 a^2 b^2 d f^2 + b^4 d^2))}{f^2} \\
& + \frac{(3 a^2 b d + b^3 d) (f x \log(e^{(f x + e)} + 1) + \text{Li}_2(-e^{(f x + e)}))}{f^2} \\
& + \frac{(3 a^2 b d + b^3 d) (f x \log(-e^{(f x + e)} + 1) + \text{Li}_2(e^{(f x + e)}))}{f^2} \\
& + \frac{(b^3 c f + 3 a b^2 d) \log(e^{(f x + e)} + 1)}{f^2} + \frac{(b^3 c f + 3 a b^2 d) \log(e^{(f x + e)} - 1)}{f^2}
\end{aligned}$$

```
[In] integrate((d*x+c)*(a+b*coth(f*x+e))**3,x, algorithm="maxima")
```

```
[Out] 1/2*a^3*d*x^2 + a^3*c*x + 3*a^2*b*c*log(sinh(f*x + e))/f - (3*a^2*b*d + b^3*d)*x^2 - 2*(b^3*c*f + 3*a*b^2*d)*x/f + 1/2*(12*a*b^2*c*f + 2*b^3*d + (3*a^2*b*d*f^2 + 3*a*b^2*d*f^2 + b^3*d*f^2)*x^2 + 2*(b^3*c*f^2 + 3*(c*f^2 + 2*d*f)*a*b^2)*x + ((3*a^2*b*d*f^2*e^(4*e) + 3*a*b^2*d*f^2*e^(4*e) + b^3*d*f^2*e^(4*e))*x^2 + 2*(3*a*b^2*c*f^2*e^(4*e) + b^3*c*f^2*e^(4*e))*x)*e^(4*f*x) - 2*(6*a*b^2*c*f*e^(2*e) + (2*c*f*e^(2*e) + d*e^(2*e))*b^3 + (3*a^2*b*d*f^2*e^(2*e) + 3*a*b^2*d*f^2*e^(2*e) + b^3*d*f^2*e^(2*e))*x^2 + 2*(3*(c*f^2*e^(2*e) + d*f*e^(2*e))*a*b^2 + (c*f^2*e^(2*e) + d*f*e^(2*e))*b^3)*x)*e^(2*f*x))/(f^2*e^(4*f*x + 4*e) - 2*f^2*e^(2*f*x + 2*e) + f^2) + (3*a^2*b*d + b^3*d)*(f*x*log(e^(f*x + e) + 1) + dilog(-e^(f*x + e)))/f^2 + (3*a^2*b*d + b^3*d)*(f*x*log(-e^(f*x + e) + 1) + dilog(e^(f*x + e)))/f^2 + (b^3*c*f + 3*a*b^2*d)*log(e^(f*x + e) + 1)/f^2 + (b^3*c*f + 3*a*b^2*d)*log(e^(f*x + e) - 1)/f^2
```

Giac [F]

$$\int (c + dx)(a + b \coth(e + fx))^3 dx = \int (dx + c)(b \coth(fx + e) + a)^3 dx$$

```
[In] integrate((d*x+c)*(a+b*coth(f*x+e))^3,x, algorithm="giac")
```

```
[Out] integrate((d*x + c)*(b*coth(f*x + e) + a)^3, x)
```

Mupad [F(-1)]

Timed out.

$$\int (c + dx)(a + b \coth(e + fx))^3 dx = \int (a + b \coth(e + fx))^3 (c + dx) dx$$

```
[In] int((a + b*coth(e + f*x))^3*(c + d*x),x)
```

```
[Out] int((a + b*coth(e + f*x))^3*(c + d*x), x)
```

3.50 $\int \frac{(a+b \coth(e+fx))^3}{c+dx} dx$

Optimal result	340
Rubi [N/A]	340
Mathematica [N/A]	341
Maple [N/A] (verified)	341
Fricas [N/A]	341
Sympy [N/A]	342
Maxima [N/A]	342
Giac [N/A]	343
Mupad [N/A]	343

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \text{Int}\left(\frac{(a + b \coth(e + fx))^3}{c + dx}, x\right)$$

[Out] Unintegrable((a+b*cOTH(f*x+e))^3/(d*x+c),x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^3}{c + dx} dx$$

[In] Int[(a + b*COTH[e + f*x])^3/(c + d*x),x]

[Out] Defer[Int][(a + b*COTH[e + f*x])^3/(c + d*x), x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \coth(e + fx))^3}{c + dx} dx$$

Mathematica [N/A]

Not integrable

Time = 48.15 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^3}{c + dx} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])^3/(c + d*x), x]`

[Out] `Integrate[(a + b*Coth[e + f*x])^3/(c + d*x), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.34 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \coth(fx + e))^3}{dx + c} dx$$

[In] `int((a+b*coth(f*x+e))^3/(d*x+c), x)`

[Out] `int((a+b*coth(f*x+e))^3/(d*x+c), x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 52, normalized size of antiderivative = 2.60

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^3}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))^3/(d*x+c), x, algorithm="fricas")`

[Out] `integral((b^3*coth(f*x + e)^3 + 3*a*b^2*coth(f*x + e)^2 + 3*a^2*b*coth(f*x + e) + a^3)/(d*x + c), x)`

Sympy [N/A]

Not integrable

Time = 1.98 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^3}{c + dx} dx$$

```
[In] integrate((a+b*coth(f*x+e))**3/(d*x+c),x)
[Out] Integral((a + b*coth(e + f*x))**3/(c + d*x), x)
```

Maxima [N/A]

Not integrable

Time = 0.75 (sec) , antiderivative size = 644, normalized size of antiderivative = 32.20

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^3}{dx + c} dx$$

```
[In] integrate((a+b*coth(f*x+e))^(3/(d*x+c), x, algorithm="maxima")
[Out] a^3*log(d*x + c)/d + (3*a^2*b + 3*a*b^2 + b^3)*log(d*x + c)/d + (6*a*b^2*d*f*x + 6*a*b^2*c*f - b^3*d - (6*a*b^2*c*f*e^(2*e) + (2*c*f*e^(2*e) - d*e^(2*e))*b^3 + 2*(3*a*b^2*d*f*e^(2*e) + b^3*d*f*e^(2*e))*x)*e^(2*f*x))/(d^2*f^2*x^2 + 2*c*d*f^2*x + c^2*f^2 + (d^2*f^2*x^2*e^(4*e) + 2*c*d*f^2*x*e^(4*e) + c^2*f^2*e^(4*e))*e^(4*f*x) - 2*(d^2*f^2*x^2*e^(2*e) + 2*c*d*f^2*x*e^(2*e) + c^2*f^2*e^(2*e))*e^(2*f*x)) - integrate((3*a^2*b*c^2*f^2 - 3*a*b^2*c*d*f + (c^2*f^2 + d^2)*b^3 + (3*a^2*b*d^2*f^2 + b^3*d^2*f^2)*x^2 + (6*a^2*b*c*d*f^2 + 2*b^3*c*d*f^2 - 3*a*b^2*d^2*f^2)*x)/(d^3*f^2*x^3 + 3*c*d^2*f^2*x^2 + 3*c^2*d*f^2*x + c^3*f^2 + (d^3*f^2*x^3*e^(2*f*x) + 3*c*d^2*f^2*x^2*e^(2*f*x) + 3*c^2*d*f^2*x*e^(2*f*x) + c^3*f^2*e^(2*f*x))*e^(f*x)), x) + integrate(-(3*a^2*b*c^2*f^2 - 3*a*b^2*c*d*f + (c^2*f^2 + d^2)*b^3 + (3*a^2*b*d^2*f^2 + b^3*d^2*f^2)*x^2 + (6*a^2*b*c*d*f^2 + 2*b^3*c*d*f^2 - 3*a*b^2*d^2*f^2)*x)/(d^3*f^2*x^3 + 3*c*d^2*f^2*x^2 + 3*c^2*d*f^2*x*e^(2*f*x) + c^3*f^2*e^(2*f*x))*e^(f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.41 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(b \coth(fx + e) + a)^3}{dx + c} dx$$

[In] `integrate((a+b*coth(f*x+e))^3/(d*x+c),x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)^3/(d*x + c), x)`

Mupad [N/A]

Not integrable

Time = 2.07 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{c + dx} dx = \int \frac{(a + b \coth(e + fx))^3}{c + dx} dx$$

[In] `int((a + b*coth(e + f*x))^3/(c + d*x),x)`

[Out] `int((a + b*coth(e + f*x))^3/(c + d*x), x)`

3.51 $\int \frac{(a+b \coth(e+fx))^3}{(c+dx)^2} dx$

Optimal result	344
Rubi [N/A]	344
Mathematica [N/A]	345
Maple [N/A] (verified)	345
Fricas [N/A]	345
Sympy [N/A]	346
Maxima [N/A]	346
Giac [N/A]	347
Mupad [N/A]	347

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \text{Int}\left(\frac{(a + b \coth(e + fx))^3}{(c + dx)^2}, x\right)$$

[Out] Unintegrable((a+b*coth(f*x+e))^3/(d*x+c)^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx$$

[In] Int[(a + b*Coth[e + f*x])^3/(c + d*x)^2,x]

[Out] Defer[Int][(a + b*Coth[e + f*x])^3/(c + d*x)^2, x]

Rubi steps

$$\text{integral} = \int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx$$

Mathematica [N/A]

Not integrable

Time = 56.00 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx$$

[In] `Integrate[(a + b*Coth[e + f*x])^3/(c + d*x)^2, x]`

[Out] `Integrate[(a + b*Coth[e + f*x])^3/(c + d*x)^2, x]`

Maple [N/A] (verified)

Not integrable

Time = 0.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{(a + b \coth(fx + e))^3}{(dx + c)^2} dx$$

[In] `int((a+b*coth(f*x+e))^3/(d*x+c)^2, x)`

[Out] `int((a+b*coth(f*x+e))^3/(d*x+c)^2, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 63, normalized size of antiderivative = 3.15

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))^3/(d*x+c)^2, x, algorithm="fricas")`

[Out] `integral((b^3*coth(f*x + e)^3 + 3*a*b^2*coth(f*x + e)^2 + 3*a^2*b*coth(f*x + e) + a^3)/(d^2*x^2 + 2*c*d*x + c^2), x)`

Sympy [N/A]

Not integrable

Time = 3.21 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx$$

```
[In] integrate((a+b*cOTH(f*x+e))**3/(d*x+c)**2,x)
```

[Out] $\text{Integral}((a + b \coth(e + f x))^3 / (c + d x)^2, x)$

Maxima [N/A]

Not integrable

Time = 0.95 (sec) , antiderivative size = 1144, normalized size of antiderivative = 57.20

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^3}{(dx + c)^2} dx$$

```
[In] integrate((a+b*cOTH(f*x+e))^3/(d*x+c)^2,x, algorithm="maxima")
```

```
[Out] -a^3/(d^2*x + c*d) - (3*a^2*b*c^2*f^2 + 3*(c^2*f^2 - 2*c*d*f)*a*b^2 + (c^2*f^2 + 2*d^2)*b^3 + (3*a^2*b*d^2*f^2 + 3*a*b^2*d^2*f^2 + b^3*d^2*f^2)*x^2 + 2*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 + 3*(c*d*f^2 - d^2*f)*a*b^2)*x + (3*a^2*b*c^2*f^2*e^(4*e) + 3*a*b^2*c^2*f^2*e^(4*e) + b^3*c^2*f^2*e^(4*e) + (3*a^2*b*d^2*f^2*e^(4*e) + 3*a*b^2*d^2*f^2*e^(4*e) + b^3*d^2*f^2*e^(4*e))*x^2 + 2*(3*a^2*b*c*d*f^2*e^(4*e) + 3*a*b^2*c*d*f^2*e^(4*e) + b^3*c*d*f^2*e^(4*e))*x) * e^(4*f*x) - 2*(3*a^2*b*c^2*f^2*e^(2*e) + 3*(c^2*f^2*e^(2*e) - c*d*f*e^(2*e)) * a*b^2 + (c^2*f^2*e^(2*e) - c*d*f*e^(2*e) + d^2*f^2*e^(2*e))*b^3 + (3*a^2*b*d^2*f^2*e^(2*e) + 3*a*b^2*d^2*f^2*e^(2*e) + b^3*d^2*f^2*e^(2*e))*x^2 + (6*a^2*b*c*d*f^2*e^(2*e) + 3*(2*c*d*f^2*e^(2*e) - d^2*f^2*e^(2*e))*a*b^2 + (2*c*d*f^2*e^(2*e) - d^2*f^2*e^(2*e))*b^3)*x)*e^(2*f*x)) / (d^4*f^2*x^3 + 3*c*d^3*f^2*x^2 + 3*c^2*d^2*f^2*x + c^3*d*f^2 + (d^4*f^2*x^3*e^(4*e) + 3*c*d^3*f^2*x^2*e^(4*e) + 3*c^2*d^2*f^2*x*e^(4*e) + c^3*d*f^2*e^(4*e))*e^(4*f*x) - 2*(d^4*f^2*x^3*e^(2*e) + 3*c*d^3*f^2*x^2*e^(2*e) + 3*c^2*d^2*f^2*x*e^(2*e) + c^3*d*f^2*e^(2*e))*e^(2*f*x)) - integrate((3*a^2*b*c^2*f^2 - 6*a*b^2*c*d*f + (c^2*f^2 + 3*d^2)*b^3 + (3*a^2*b*d^2*f^2 + b^3*d^2*f^2)*x^2 + 2*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 - 3*a*b^2*d^2*f)*x) / (d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2 + (d^4*f^2*x^4*e^e + 4*c*d^3*f^2*x^3*e^e + 6*c^2*d^2*f^2*x^2*e^e + 4*c^3*d*f^2*x^2*e^e + c^4*f^2*e^e)*e^(f*x)), x) + integrate(-(3*a^2*b*c^2*f^2 - 6*a*b^2*c*d*f + (c^2*f^2 + 3*d^2)*b^3 + (3*a^2*b*d^2*f^2 + b^3*d^2*f^2)*x^2 + 2*(3*a^2*b*c*d*f^2 + b^3*c*d*f^2 - 3*a*b^2*d^2*f)*x) / (d^4*f^2*x^4 + 4*c*d^3*f^2*x^3 + 6*c^2*d^2*f^2*x^2 + 4*c^3*d*f^2*x + c^4*f^2 - (d^4*f^2*x^4*e^e + 4*c*d^3*f^2*x^3*e^e + 6*c^2*d^2*f^2*x^2*e^e + 4*c^3*d*f^2*x^2*e^e + c^4*f^2*e^e)*e^(f*x)), x)
```

Giac [N/A]

Not integrable

Time = 0.80 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(b \coth(fx + e) + a)^3}{(dx + c)^2} dx$$

[In] `integrate((a+b*coth(f*x+e))^3/(d*x+c)^2,x, algorithm="giac")`

[Out] `integrate((b*coth(f*x + e) + a)^3/(d*x + c)^2, x)`

Mupad [N/A]

Not integrable

Time = 2.16 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx = \int \frac{(a + b \coth(e + fx))^3}{(c + dx)^2} dx$$

[In] `int((a + b*coth(e + f*x))^3/(c + d*x)^2,x)`

[Out] `int((a + b*coth(e + f*x))^3/(c + d*x)^2, x)`

3.52 $\int \frac{(c+dx)^3}{a+b \coth(e+fx)} dx$

Optimal result	348
Rubi [A] (verified)	348
Mathematica [A] (verified)	351
Maple [B] (verified)	352
Fricas [B] (verification not implemented)	352
Sympy [F]	353
Maxima [B] (verification not implemented)	354
Giac [F]	355
Mupad [F(-1)]	355

Optimal result

Integrand size = 20, antiderivative size = 210

$$\begin{aligned} \int \frac{(c+dx)^3}{a+b \coth(e+fx)} dx = & \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} \\ & + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2} \\ & + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^3} \\ & + \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{4(a^2 - b^2)f^4} \end{aligned}$$

[Out] $1/4*(d*x+c)^4/(a+b)/d-b*(d*x+c)^3*\ln(1+(-a+b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f+3/2*b*d*(d*x+c)^2*\operatorname{polylog}(2, (a-b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f^{2+3/2}*b*d^2*(d*x+c)*\operatorname{polylog}(3, (a-b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f^3+3/4*b*d^3*\operatorname{polylog}(4, (a-b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f^4)$

Rubi [A] (verified)

Time = 0.26 (sec), antiderivative size = 210, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used

$$= \{3812, 2221, 2611, 6744, 2320, 6724\}$$

$$\begin{aligned} \int \frac{(c+dx)^3}{a+b \coth(e+fx)} dx &= \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^3(a^2-b^2)} \\ &\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^2(a^2-b^2)} \\ &\quad - \frac{b(c+dx)^3 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f(a^2-b^2)} \\ &\quad + \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{4f^4(a^2-b^2)} + \frac{(c+dx)^4}{4d(a+b)} \end{aligned}$$

[In] `Int[(c + d*x)^3/(a + b*Coth[e + f*x]), x]`

[Out] $(c + d*x)^4/(4*(a + b)*d) - (b*(c + d*x)^3*\operatorname{Log}[1 - (a - b)/((a + b)*E^{(2*(e + f*x))})])/((a^2 - b^2)*f) + (3*b*d*(c + d*x)^2*\operatorname{PolyLog}[2, (a - b)/((a + b)*E^{(2*(e + f*x))})])/((2*(a^2 - b^2)*f^2) + (3*b*d^2*(c + d*x))*\operatorname{PolyLog}[3, (a - b)/((a + b)*E^{(2*(e + f*x))})])/((2*(a^2 - b^2)*f^3) + (3*b*d^3*\operatorname{PolyLog}[4, (a - b)/((a + b)*E^{(2*(e + f*x))})])/((4*(a^2 - b^2)*f^4)$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*(c_.)*(x_.)^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*(a_.)*(v_)^(n_.)^m_.] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[\operatorname{Log}[1 + (e_.)*(F_)^((c_.)*(a_.) + (b_.)*(x_.)))^n_]*(f_.) + (g_.)*(x_.)^m_.], x_Symbol] :> Simp[(-(f + g*x)^m)*(\operatorname{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*\operatorname{PolyLog}[2, (-e)*(F^(c*(a + b*x)))^n], x], x]; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3812

```
Int[((c_) + (d_)*(x_))^(m_)/((a_) + (b_)*tan[(e_) + Pi*(k_) + (f_)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*E^(2*I*k*Pi)*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^(2*I*k*Pi)*E^Simp[2*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[4*k] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_) + (b_)*(x_)]^p]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_) + (f_)*(x_))^m*PolyLog[n_, (d_)*((c_)*(a_) + (b_)*(x_))]^p], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a + b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m - 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c, d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^4}{4(a+b)d} - (2b) \int \frac{e^{-2(e+fx)}(c+dx)^3}{(a+b)^2 + (-a^2+b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{(3bd) \int (c+dx)^2 \log\left(1 + \frac{(-a^2+b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2-b^2)f} \\
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} \\
&\quad + \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2} \\
&\quad - \frac{(3bd^2) \int (c+dx) \text{PolyLog}\left(2, -\frac{(-a^2+b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2-b^2)f^2} \\
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)f} + \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2-b^2)f^3} - \frac{(3bd^3) \int \text{PolyLog}\left(3, -\frac{(-a^2+b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{2(a^2-b^2)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2) f} \\
&\quad + \frac{3bd(c+dx)^2 \operatorname{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{2(a^2 - b^2) f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{2(a^2 - b^2) f^3} \\
&\quad + \frac{(3bd^3) \operatorname{Subst} \left(\int \frac{\operatorname{PolyLog} \left(3, \frac{(a-b)x}{a+b} \right)}{x} dx, x, e^{-2(e+fx)} \right)}{4(a^2 - b^2) f^4} \\
&= \frac{(c+dx)^4}{4(a+b)d} - \frac{b(c+dx)^3 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2) f} + \frac{3bd(c+dx)^2 \operatorname{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{2(a^2 - b^2) f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog} \left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{2(a^2 - b^2) f^3} + \frac{3bd^3 \operatorname{PolyLog} \left(4, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{4(a^2 - b^2) f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.53 (sec), antiderivative size = 247, normalized size of antiderivative = 1.18

$$\begin{aligned}
&\int \frac{(c+dx)^3}{a+b \coth(e+fx)} dx \\
&= \frac{1}{4} \left(\frac{2b(c+dx)^4}{(a+b)d(a(-1+e^{2e})+b(1+e^{2e}))} - \frac{4b(c+dx)^3 \log \left(1 + \frac{(-a+b)e^{-2(e+fx)}}{a+b} \right)}{(a-b)(a+b)f} \right. \\
&\quad + \frac{3bd \left(2f^2(c+dx)^2 \operatorname{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right) + d \left(2f(c+dx) \operatorname{PolyLog} \left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right) + d \operatorname{PolyLog} \left(4, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right) \right) \right)}{(a-b)(a+b)f^4} \\
&\quad \left. + \frac{x(4c^3 + 6c^2dx + 4cd^2x^2 + d^3x^3) \sinh(e)}{b \cosh(e) + a \sinh(e)} \right)
\end{aligned}$$

[In] `Integrate[(c + d*x)^3/(a + b*Coth[e + f*x]), x]`

[Out]
$$\begin{aligned}
&((2*b*(c + d*x)^4)/((a + b)*d*(a*(-1 + E^(2*e)) + b*(1 + E^(2*e))))) - (4*b*(c + d*x)^3*Log[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))]))/((a - b)*(a + b)*f) \\
&+ (3*b*d*(2*f^2*(c + d*x)^2*PolyLog[2, (a - b)/((a + b)*E^(2*(e + f*x)))] + d*(2*f*(c + d*x)*PolyLog[3, (a - b)/((a + b)*E^(2*(e + f*x)))] + d*PolyLog[4, (a - b)/((a + b)*E^(2*(e + f*x)))]))/((a - b)*(a + b)*f^4) + (x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*Sinh[e])/(b*Cosh[e] + a*Sinh[e]))/4
\end{aligned}$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1157 vs. $2(209) = 418$.

Time = 0.45 (sec), antiderivative size = 1158, normalized size of antiderivative = 5.51

method	result	size
risch	Expression too large to display	1158

[In] `int((d*x+c)^3/(a+b*cOTH(f*x+e)),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & \frac{1}{4} / (a+b) * d^3 * x^4 + \frac{1}{4} / (a+b) / d * c^4 + \frac{1}{2} * b / (a+b) / (a-b) * d^3 * x^4 + \frac{3}{2} / f^4 * b / (a+b) \\ & / (a-b) * d^3 * e^4 - \frac{3}{4} / f^4 * b / (a+b) / (a-b) * d^3 * \text{polylog}(4, (a+b) * \exp(2*f*x+2*e)) / (a-b) \\ & - 1 / f * b / (a+b) * c^3 / (a-b) * \ln(\exp(2*f*x+2*e)) * a+b * \exp(2*f*x+2*e) - a+b) + 1 / (a+b) \\ & * d^2 * c * x^3 + \frac{3}{2} / (a+b) * d * c^2 * x^2 + 1 / (a+b) * c^3 * x^2 + \frac{3}{2} / f^3 * b / (a+b) / (a-b) * d^3 * e^3 * x^2 \\ & + 2 * b / (a+b) / (a-b) * d^2 * c * x^3 - 4 / f^3 * b / (a+b) / (a-b) * d^2 * c * e^3 - 1 / f * b / (a+b) / (a-b) * d \\ & ^3 * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * x^3 - \frac{3}{2} / f^2 * b / (a+b) / (a-b) * d^3 * \text{polylog}(2, (a+b) * \exp(2*f*x+2*e)) / (a-b) * x^2 + \frac{3}{2} / f^3 * b / (a+b) / (a-b) * d^2 * c * \text{polylog}(3, (a+b) * \exp(2*f*x+2*e)) / (a-b) + 1 / f^4 * b / (a+b) * d^3 * e^3 / (a-b) * \ln(\exp(2*f*x+2*e)) * a+b * \exp(2*f*x+2*e) - a+b) - \frac{3}{2} / f^2 * b / (a+b) / (a-b) * d * c^2 * \text{polylog}(2, (a+b) * \exp(2*f*x+2*e)) / (a-b) - 1 / f^4 * b / (a+b) / (a-b) * d^3 * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * e^3 + \frac{3}{2} / f^3 * b / (a+b) / (a-b) * d^3 * \text{polylog}(3, (a+b) * \exp(2*f*x+2*e)) / (a-b) * x^3 - \frac{3}{2} / f^2 * b / (a+b) / (a-b) * d^2 * c * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * x^2 + \frac{3}{2} / f^3 * b / (a+b) / (a-b) * d^2 * c^2 * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * e^2 - \frac{2}{f^4 * b / (a+b) * d^3 * e^3 / (a-b) * \ln(\exp(f*x+e)) + 3 * b / (a+b) / (a-b) * d * c^2 * x^2 + 3 / f^2 * b / (a+b) / (a-b) * d * c^2 * e^2 - 6 / f^2 * b / (a+b) * d * c^2 * e / (a-b) * \ln(\exp(f*x+e)) + 6 / f^3 * b / (a+b) * d^2 * c * e^2 / (a-b) * \ln(\exp(f*x+e)) + 6 / f * b / (a+b) / (a-b) * d * c^2 * e * x - 6 / f^2 * b / (a+b) / (a-b) * d^2 * c * e^2 * x - 3 / f^2 * b / (a+b) / (a-b) * d^2 * c * polylog(2, (a+b) * \exp(2*f*x+2*e)) / (a-b) * x + 3 / f^2 * b / (a+b) * d * c^2 * e / (a-b) * \ln(\exp(2*f*x+2*e)) * a+b * \exp(2*f*x+2*e) - a+b) - 3 / f * b / (a+b) / (a-b) * d * c^2 * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * x - 3 / f^2 * b / (a+b) / (a-b) * d * c^2 * \ln(1 - (a+b) * \exp(2*f*x+2*e)) / (a-b) * e - 3 / f^3 * b / (a+b) * d^2 * c * e^2 / (a-b) * \ln(\exp(2*f*x+2*e)) * a+b * \exp(2*f*x+2*e) - a+b) + 2 / f * b / (a+b) * c^3 / (a-b) * \ln(\exp(f*x+e))) \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 730 vs. $2(201) = 402$.

Time = 0.28 (sec), antiderivative size = 730, normalized size of antiderivative = 3.48

$$\begin{aligned} & \int \frac{(c+dx)^3}{a+b\coth(e+fx)} dx \\ &= \frac{(a+b)d^3f^4x^4 + 4(a+b)cd^2f^4x^3 + 6(a+b)c^2df^4x^2 + 4(a+b)c^3f^4x - 24bd^3\text{polylog}\left(4, \sqrt{\frac{a+b}{a-b}}(\cosh(fx+e))\right)}{a^2b^2} \end{aligned}$$

[In] `integrate((d*x+c)^3/(a+b*cOTH(f*x+e)),x, algorithm="fricas")`

```
[Out] 1/4*((a + b)*d^3*f^4*x^4 + 4*(a + b)*c*d^2*f^4*x^3 + 6*(a + b)*c^2*d*f^4*x^2 + 4*(a + b)*c^3*f^4*x - 24*b*d^3*polylog(4, sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))) - 24*b*d^3*polylog(4, -sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))) - 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))) - 12*(b*d^3*f^2*x^2 + 2*b*c*d^2*f^2*x + b*c^2*d*f^2)*dilog(-sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))) + 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt((a + b)/(a - b))) + 4*(b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2 - b*c^3*f^3)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) - 2*(a - b)*sqrt((a + b)/(a - b))) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 4*(b*d^3*f^3*x^3 + 3*b*c*d^2*f^3*x^2 + 3*b*c^2*d*f^3*x + b*d^3*e^3 - 3*b*c*d^2*e^2*f + 3*b*c^2*d*e*f^2)*log(-sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e)) + 1) + 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))) + 24*(b*d^3*f*x + b*c*d^2*f)*polylog(3, -sqrt((a + b)/(a - b)))*(cosh(f*x + e) + sinh(f*x + e))))/((a^2 - b^2)*f^4)
```

Sympy [F]

$$\int \frac{(c + dx)^3}{a + b \coth(e + fx)} dx = \int \frac{(c + dx)^3}{a + b \coth(e + fx)} dx$$

```
[In] integrate((d*x+c)**3/(a+b*coth(f*x+e)),x)
[Out] Integral((c + d*x)**3/(a + b*coth(e + f*x)), x)
```

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 521 vs. $2(201) = 402$.

Time = 0.32 (sec), antiderivative size = 521, normalized size of antiderivative = 2.48

$$\begin{aligned}
 & \int \frac{(c+dx)^3}{a+b\coth(e+fx)} dx \\
 &= -\frac{3 \left(2 f x \log \left(-\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + \text{Li}_2 \left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} \right) \right) bc^2 d}{2(a^2 f^2 - b^2 f^2)} \\
 &\quad - \frac{3 \left(2 f^2 x^2 \log \left(-\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + 2 f x \text{Li}_2 \left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} \right) - \text{Li}_3 \left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} \right) \right) bcd^2}{2(a^2 f^3 - b^2 f^3)} \\
 &\quad - \frac{\left(4 f^3 x^3 \log \left(-\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} + 1 \right) + 6 f^2 x^2 \text{Li}_2 \left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} \right) - 6 f x \text{Li}_3 \left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} \right) + 3 \right) d^3 f^4 x^4}{3(a^2 f^4 - b^2 f^4)} \\
 &\quad - c^3 \left(\frac{b \log \left(-(a-b)e^{(-2fx-2e)} + a+b \right)}{(a^2 - b^2)f} - \frac{fx + e}{(a+b)f} \right) \\
 &\quad + \frac{bd^3 f^4 x^4 + 4bcd^2 f^4 x^3 + 6bc^2 df^4 x^2}{2(a^2 f^4 - b^2 f^4)} + \frac{d^3 x^4 + 4cd^2 x^3 + 6c^2 dx^2}{4(a+b)}
 \end{aligned}$$

```
[In] integrate((d*x+c)^3/(a+b*cOTH(f*x+e)),x, algorithm="maxima")
[Out] -3/2*(2*f*x*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + dilog((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*c^2*d/(a^2*f^2 - b^2*f^2) - 3/2*(2*f^2*x^2*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 2*f*x*dilo g((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - polylog(3, (a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*c*d^2/(a^2*f^3 - b^2*f^3) - 1/3*(4*f^3*x^3*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 6*f^2*x^2*dilog((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - 6*f*x*polylog(3, (a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) + 3*polylog(4, (a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*d^3/(a^2*f^4 - b^2*f^4) - c^3*(b*log(-(a - b)*e^(-2*f*x - 2*e) + a + b)/((a^2 - b^2)*f) - (f*x + e)/((a + b)*f)) + 1/2*(b*d^3*f^4*x^4 + 4*b*c*d^2*f^4*x^3 + 6*b*c^2*d*f^4*x^2)/(a^2*f^4 - b^2*f^4) + 1/4*(d^3*x^4 + 4*c*d^2*x^3 + 6*c^2*d*x^2)/(a + b)
```

Giac [F]

$$\int \frac{(c + dx)^3}{a + b \coth(e + fx)} dx = \int \frac{(dx + c)^3}{b \coth(fx + e) + a} dx$$

[In] `integrate((d*x+c)^3/(a+b*coth(f*x+e)),x, algorithm="giac")`
[Out] `integrate((d*x + c)^3/(b*coth(f*x + e) + a), x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{a + b \coth(e + fx)} dx = \int \frac{(c + d x)^3}{a + b \coth(e + f x)} dx$$

[In] `int((c + d*x)^3/(a + b*coth(e + f*x)),x)`
[Out] `int((c + d*x)^3/(a + b*coth(e + f*x)), x)`

3.53 $\int \frac{(c+dx)^2}{a+b \coth(e+fx)} dx$

Optimal result	356
Rubi [A] (verified)	356
Mathematica [A] (verified)	359
Maple [B] (verified)	359
Fricas [B] (verification not implemented)	360
Sympy [F]	360
Maxima [B] (verification not implemented)	361
Giac [F]	361
Mupad [F(-1)]	362

Optimal result

Integrand size = 20, antiderivative size = 156

$$\begin{aligned} \int \frac{(c+dx)^2}{a+b \coth(e+fx)} dx = & \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} \\ & + \frac{bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f^2} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^3} \end{aligned}$$

[Out] $1/3*(d*x+c)^3/(a+b)/d-b*(d*x+c)^2*\ln(1+(-a+b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f+b*d*(d*x+c)*\operatorname{polylog}(2, (a-b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f^2+1/2*b*d^2*\operatorname{polylog}(3, (a-b)/(a+b)/\exp(2*f*x+2*e))/(a^{2-b^2}/f^3$

Rubi [A] (verified)

Time = 0.22 (sec), antiderivative size = 156, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.250, Rules used = {3812, 2221, 2611, 2320, 6724}

$$\begin{aligned} \int \frac{(c+dx)^2}{a+b \coth(e+fx)} dx = & \frac{bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f^2(a^2 - b^2)} \\ & - \frac{b(c+dx)^2 \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f(a^2 - b^2)} \\ & + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^3(a^2 - b^2)} + \frac{(c+dx)^3}{3d(a+b)} \end{aligned}$$

[In] $\text{Int}[(c + d*x)^2/(a + b*\text{Coth}[e + f*x]), x]$

[Out] $(c + d*x)^3/(3*(a + b)*d) - (b*(c + d*x)^2*\text{Log}[1 - (a - b)/((a + b)*E^{(2*(e + f*x))})])/((a^2 - b^2)*f) + (b*d*(c + d*x)*\text{PolyLog}[2, (a - b)/((a + b)*E^{(2*(e + f*x))})])/((a^2 - b^2)*f^2) + (b*d^2*\text{PolyLog}[3, (a - b)/((a + b)*E^{(2*(e + f*x))})])/(2*(a^2 - b^2)*f^3)$

Rule 2221

```
Int[((F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)*((c_.) + (d_.)*(x_.))^(m_.))/((a_) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_.)))^(n_.)), x_Symbol] :> Simplify[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x]; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_.)^(n_.))^(m_.)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_.)*x))*(F_[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_.)*(x_.)))^(n_.))*((f_.) + (g_.)*(x_.))^(m_.), x_Symbol] :> Simplify[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x]; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3812

```
Int[((c_.) + (d_.)*(x_.))^(m_.)/((a_) + (b_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_.)]), x_Symbol] :> Simplify[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*E^(2*I*k*Pi)*(E^Simp[2*I*(e + f*x), x])/((a + I*b)^2 + (a^2 + b^2)*E^(2*I*k*Pi)*E^Simp[2*I*(e + f*x), x]], x], x]; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[4*k] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*(a_.) + (b_.)*(x_.))^(p_.)]/((d_.) + (e_.)*(x_.)), x_Symbol] :> Simplify[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x]; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^3}{3(a+b)d} - (2b) \int \frac{e^{-2(e+fx)}(c+dx)^2}{(a+b)^2 + (-a^2 + b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f} \\
&\quad + \frac{(2bd) \int (c+dx) \log \left(1 + \frac{(-a^2 + b^2)e^{-2(e+fx)}}{(a+b)^2} \right) dx}{(a^2 - b^2)f} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f^2} \\
&\quad - \frac{(bd^2) \int \text{PolyLog} \left(2, -\frac{(-a^2 + b^2)e^{-2(e+fx)}}{(a+b)^2} \right) dx}{(a^2 - b^2)f^2} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f^2} \\
&\quad + \frac{(bd^2) \text{Subst} \left(\int \frac{\text{PolyLog} \left(2, \frac{(a-b)x}{a+b} \right)}{x} dx, x, e^{-2(e+fx)} \right)}{2(a^2 - b^2)f^3} \\
&= \frac{(c+dx)^3}{3(a+b)d} - \frac{b(c+dx)^2 \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f} \\
&\quad + \frac{bd(c+dx) \text{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{(a^2 - b^2)f^2} + \frac{bd^2 \text{PolyLog} \left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b} \right)}{2(a^2 - b^2)f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.08 (sec) , antiderivative size = 200, normalized size of antiderivative = 1.28

$$\begin{aligned} & \int \frac{(c+dx)^2}{a+b \coth(e+fx)} dx \\ &= \frac{1}{6} \left(\frac{4b(c+dx)^3}{(a+b)d(a(-1+e^{2e})+b(1+e^{2e}))} - \frac{6b(c+dx)^2 \log\left(1+\frac{(-a+b)e^{-2(e+fx)}}{a+b}\right)}{(a-b)(a+b)f} \right. \\ & \quad + \frac{3bd\left(2f(c+dx) \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right) + d \operatorname{PolyLog}\left(3, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)\right)}{(a-b)(a+b)f^3} \\ & \quad \left. + \frac{2x(3c^2+3cdx+d^2x^2) \sinh(e)}{b \cosh(e) + a \sinh(e)} \right) \end{aligned}$$

[In] `Integrate[(c + d*x)^2/(a + b*Coth[e + f*x]), x]`

[Out] $\frac{(4*b*(c+d*x)^3)/((a+b)*d*(a*(-1+E^(2*e))+b*(1+E^(2*e)))) - (6*b*(c+d*x)^2*\text{Log}[1+(-a+b)/((a+b)*E^(2*(e+f*x)))]))/((a-b)*(a+b)*f) + (3*b*d*(2*f*(c+d*x)*\text{PolyLog}[2, (a-b)/((a+b)*E^(2*(e+f*x)))] + d*\text{PolyLog}[3, (a-b)/((a+b)*E^(2*(e+f*x)))]))/((a-b)*(a+b)*f^3) + (2*x*(3*c^2+3*c*d*x+d^2*x^2)*\text{Sinh}[e])/(b*\text{Cosh}[e] + a*\text{Sinh}[e]))/6$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 734 vs. 2(157) = 314.

Time = 0.41 (sec) , antiderivative size = 735, normalized size of antiderivative = 4.71

method	result
risch	$\frac{d^2x^3}{3a+3b} + \frac{dcx^2}{a+b} + \frac{c^2x}{a+b} + \frac{c^3}{3(a+b)d} - \frac{bc^2 \ln(e^{2fx+2e}a+b e^{2fx+2e}-a+b)}{f(a+b)(a-b)} + \frac{2bd^2x^3}{3(a+b)(a-b)} - \frac{4bd^2e^3}{3f^3(a+b)(a-b)} - \frac{bd^2 \ln(1-e^{2fx+2e}a+b e^{2fx+2e}-a+b)}{f(a+b)(a-b)}$

[In] `int((d*x+c)^2/(a+b*coth(f*x+e)), x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{3}/(a+b)*d^2*x^3+1/(a+b)*d*c*x^2+1/(a+b)*c^2*x+1/3/(a+b)/d*c^3-1/f*b/(a+b)*c^2/(a-b)*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)+2/3*b/(a+b)/(a-b)*d^2*x^3-4/3*f^3*b/(a+b)/(a-b)*d^2*x^3-1/f*b/(a+b)/(a-b)*d^2*\ln(1-(a+b))*\exp(2*f*x+2*e)/(a-b)*x^2+1/2/f^3*b/(a+b)/(a-b)*d^2*polylog(3, (a+b)*\exp(2*f*x+2*e)/(a-b))+2/f^3*b/(a+b)*d^2*x^2/(a-b)*\ln(\exp(f*x+e))-1/f^3*b/(a+b)*e^2*d^2/(a-b)*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)-4/f^2*b/(a+b)*d*c*e/(a-b)*\ln(e^{xp(f*x+e)})+2/f^2*b/(a+b)*c*d*e/(a-b)*\ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)-2/f^2*b/(a+b)/(a-b)*d^2*x^2+1/f^3*b/(a+b)/(a-b)*d^2*\ln(1-(a+b))*\exp(2*f*x+2*e)/(a-b)*e^2-1/f^2*b/(a+b)/(a-b)*d^2*polylog(2, (a+b)*\exp(2*f*x+2*e)/(a-b))$

$$(a-b)*x+2*b/(a+b)/(a-b)*d*c*x^2+f^2*b/(a+b)/(a-b)*d*c*e^2-2/f*b/(a+b)/(a-b)*d*c*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x-1/f^2*b/(a+b)/(a-b)*d*c*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))+2/f*b/(a+b)*c^2/(a-b)*ln(exp(f*x+e))+4/f*b/(a+b)/(a-b)*d*c*e*x-2/f^2*b/(a+b)/(a-b)*d*c*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 492 vs. $2(151) = 302$.

Time = 0.27 (sec) , antiderivative size = 492, normalized size of antiderivative = 3.15

$$\int \frac{(c+dx)^2}{a+b\coth(e+fx)} dx \\ \equiv \frac{(a+b)d^2f^3x^3 + 3(a+b)cd^2f^3x^2 + 3(a+b)c^2f^3x + 6bd^2\text{polylog}\left(3, \sqrt{\frac{a+b}{a-b}}(\cosh(fx+e) + \sinh(fx+e))\right)}{(a+b)^2f^3x^3 + 3(a+b)cd^2f^3x^2 + 3(a+b)c^2f^3x + 6bd^2\text{polylog}\left(3, \sqrt{\frac{a+b}{a-b}}(\cosh(fx+e) + \sinh(fx+e))\right)}$$

```
[In] integrate((d*x+c)^2/(a+b*cOTH(f*x+e)),x, algorithm="fricas")
```

```
[Out] 1/3*((a + b)*d^2*f^3*x^3 + 3*(a + b)*c*d*f^3*x^2 + 3*(a + b)*c^2*f^3*x + 6*b*d^2*polylog(3, sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) + 6*b*d^2*polylog(3, -sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*(b*d^2*f*x + b*c*d*f)*dilog(sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*(b*d^2*f*x + b*c*d*f)*dilog(-sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 3*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt((a + b)/(a - b))) - 3*(b*d^2*e^2 - 2*b*c*d*e*f + b*c^2*f^2)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sinh(f*x + e) - 2*(a - b)*sqrt((a + b)/(a - b))) - 3*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 3*(b*d^2*f^2*x^2 + 2*b*c*d*f^2*x - b*d^2*e^2 + 2*b*c*d*e*f)*log(-sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1))/((a^2 - b^2)*f^3)
```

Sympy [F]

$$\int \frac{(c+dx)^2}{a+b\coth(e+fx)} dx = \int \frac{(c+dx)^2}{a+b\coth(e+fx)} dx$$

```
[In] integrate((d*x+c)**2/(a+b*coth(f*x+e)),x)
```

[Out] $\text{Integral}((c + d*x)^2/(a + b*\coth(e + f*x)), x)$

Maxima [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 332 vs. $2(151) = 302$.

Time = 0.29 (sec) , antiderivative size = 332, normalized size of antiderivative = 2.13

$$\begin{aligned} & \int \frac{(c + dx)^2}{a + b \coth(e + fx)} dx \\ &= -\frac{\left(2 f x \log \left(-\frac{(ae^{(2 e)}+be^{(2 e)})e^{(2 f x)}}{a-b}+1\right)+\text{Li}_2\left(\frac{(ae^{(2 e)}+be^{(2 e)})e^{(2 f x)}}{a-b}\right)\right) b c d}{a^2 f^2-b^2 f^2} \\ &-\frac{\left(2 f^2 x^2 \log \left(-\frac{(ae^{(2 e)}+be^{(2 e)})e^{(2 f x)}}{a-b}+1\right)+2 f x \text{Li}_2\left(\frac{(ae^{(2 e)}+be^{(2 e)})e^{(2 f x)}}{a-b}\right)-\text{Li}_3\left(\frac{(ae^{(2 e)}+be^{(2 e)})e^{(2 f x)}}{a-b}\right)\right) b d^2}{2 \left(a^2 f^3-b^2 f^3\right)} \\ &-c^2 \left(\frac{b \log \left(-(a-b) e^{(-2 f x-2 e)}+a+b\right)}{\left(a^2-b^2\right) f}-\frac{f x+e}{(a+b) f}\right) \\ &+\frac{2 \left(b d^2 f^3 x^3+3 b c d f^3 x^2\right)}{3 \left(a^2 f^3-b^2 f^3\right)}+\frac{d^2 x^3+3 c d x^2}{3 \left(a+b\right)} \end{aligned}$$

```
[In] integrate((d*x+c)^2/(a+b*coth(f*x+e)),x, algorithm="maxima")
[Out] -(2*f*x*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + dilog((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*c*d/(a^2*f^2 - b^2*f^2) - 1/2*(2*f^2*x^2*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + 2*f*x*dilog((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)) - polylog(3, (a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b)))*b*d^2/(a^2*f^3 - b^2*f^3) - c^2*(b*log(-(a - b)*e^(-2*f*x - 2*e) + a + b)/(a^2 - b^2)*f) - (f*x + e)/((a + b)*f) + 2/3*(b*d^2*f^3*x^3 + 3*b*c*d*f^3*x^2)/(a^2*f^3 - b^2*f^3) + 1/3*(d^2*x^3 + 3*c*d*x^2)/(a + b)
```

Giac [F]

$$\int \frac{(c + dx)^2}{a + b \coth(e + fx)} dx = \int \frac{(dx + c)^2}{b \coth(fx + e) + a} dx$$

```
[In] integrate((d*x+c)^2/(a+b*coth(f*x+e)),x, algorithm="giac")
[Out] integrate((d*x + c)^2/(b*coth(f*x + e) + a), x)
```

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{a + b \coth(e + fx)} dx = \int \frac{(c + dx)^2}{a + b \coth(e + fx)} dx$$

[In] `int((c + d*x)^2/(a + b*coth(e + f*x)),x)`

[Out] `int((c + d*x)^2/(a + b*coth(e + f*x)), x)`

3.54 $\int \frac{c+dx}{a+b \coth(e+fx)} dx$

Optimal result	363
Rubi [A] (verified)	363
Mathematica [A] (verified)	365
Maple [B] (verified)	365
Fricas [B] (verification not implemented)	366
Sympy [F]	366
Maxima [F]	366
Giac [F]	367
Mupad [F(-1)]	367

Optimal result

Integrand size = 18, antiderivative size = 108

$$\int \frac{c+dx}{a+b \coth(e+fx)} dx = \frac{(c+dx)^2}{2(a+b)d} - \frac{b(c+dx) \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{bd \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2}$$

[Out] $1/2*(d*x+c)^2/(a+b)/d-b*(d*x+c)*\ln(1+(-a+b)/(a+b))/\exp(2*f*x+2*e))/(a^2-b^2)/f+1/2*b*d*\operatorname{polylog}(2, (a-b)/(a+b))/\exp(2*f*x+2*e))/(a^2-b^2)/f^2$

Rubi [A] (verified)

Time = 0.13 (sec), antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.222, Rules used = {3812, 2221, 2317, 2438}

$$\int \frac{c+dx}{a+b \coth(e+fx)} dx = -\frac{b(c+dx) \log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f(a^2 - b^2)} + \frac{bd \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2f^2(a^2 - b^2)} + \frac{(c+dx)^2}{2d(a+b)}$$

[In] $\operatorname{Int}[(c+d*x)/(a+b*\operatorname{Coth}[e+f*x]),x]$

[Out] $(c+d*x)^2/(2*(a+b)*d) - (b*(c+d*x)*\operatorname{Log}[1 - (a-b)/((a+b)*E^(2*(e+f*x)))])/((a^2-b^2)*f) + (b*d*\operatorname{PolyLog}[2, (a-b)/((a+b)*E^(2*(e+f*x)))])/(2*(a^2-b^2)*f^2)$

Rule 2221

```
Int[((((F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)*((c_.) + (d_.)*(x_))^(m_.))/((a_.) + (b_.)*(F_)^((g_.)*(e_.) + (f_.)*(x_))))^(n_.)), x_Symbol] :> Simp[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x] - Dist[d*(m/(b*f*g*n*Log[F])), Int[(c + d*x)^(m - 1)*Log[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; FreeQ[{F, a, b, c, d, e, f, g, n}, x] && IGtQ[m, 0]
```

Rule 2317

```
Int[Log[(a_.) + (b_.)*(F_)^((e_.)*(c_.) + (d_.)*(x_)))^(n_.)], x_Symbol] :> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2438

```
Int[Log[(c_.)*(d_.) + (e_.)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 3812

```
Int[((c_.) + (d_.)*(x_))^(m_.)/((a_.) + (b_.)*tan[(e_.) + Pi*(k_.) + (f_.)*(x_)]), x_Symbol] :> Simp[(c + d*x)^(m + 1)/(d*(m + 1)*(a + I*b)), x] + Dist[2*I*b, Int[(c + d*x)^m*E^(2*I*k*Pi)*(E^Simp[2*I*(e + f*x), x]/((a + I*b)^2 + (a^2 + b^2)*E^(2*I*k*Pi)*E^Simp[2*I*(e + f*x), x])), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && IntegerQ[4*k] && NeQ[a^2 + b^2, 0] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \frac{(c+dx)^2}{2(a+b)d} - (2b) \int \frac{e^{-2(e+fx)}(c+dx)}{(a+b)^2 + (-a^2 + b^2)e^{-2(e+fx)}} dx \\
&= \frac{(c+dx)^2}{2(a+b)d} - \frac{b(c+dx)\log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{(bd) \int \log\left(1 + \frac{(-a^2 + b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2 - b^2)f} \\
&= \frac{(c+dx)^2}{2(a+b)d} - \frac{b(c+dx)\log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} \\
&\quad - \frac{(bd)\text{Subst}\left(\int \frac{\log\left(1 + \frac{(-a^2 + b^2)x}{(a+b)^2}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{2(a^2 - b^2)f^2} \\
&= \frac{(c+dx)^2}{2(a+b)d} - \frac{b(c+dx)\log\left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2 - b^2)f} + \frac{bd \text{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{2(a^2 - b^2)f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 1.04 (sec) , antiderivative size = 152, normalized size of antiderivative = 1.41

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \frac{1}{2} \left(\frac{2b(c + dx)^2}{(a + b)d(a(-1 + e^{2e}) + b(1 + e^{2e}))} - \frac{2b(c + dx) \log\left(1 + \frac{(-a+b)e^{-2(e+fx)}}{a+b}\right)}{(a - b)(a + b)f} + \frac{bd \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a - b)(a + b)f^2} + \frac{x(2c + dx) \sinh(e)}{b \cosh(e) + a \sinh(e)} \right)$$

[In] `Integrate[(c + d*x)/(a + b*Coth[e + f*x]), x]`

[Out] $\frac{(2*b*(c + d*x)^2)/((a + b)*d*(a*(-1 + E^(2*e)) + b*(1 + E^(2*e)))) - (2*b*(c + d*x)*Log[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))]))/((a - b)*(a + b)*f) + (b*d*PolyLog[2, (a - b)/((a + b)*E^(2*(e + f*x)))]))/((a - b)*(a + b)*f^2) + (x*(2*c + d*x)*Sinh[e])/((b*Cosh[e] + a*Sinh[e]))/2}{}$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 356 vs. $2(107) = 214$.

Time = 0.36 (sec) , antiderivative size = 357, normalized size of antiderivative = 3.31

method	result
risch	$\frac{dx^2}{2a+2b} + \frac{cx}{a+b} + \frac{2bc \ln(e^{fx+e})}{f(a+b)(a-b)} - \frac{bc \ln(e^{2fx+2e}a+b e^{2fx+2e}-a+b)}{f(a+b)(a-b)} + \frac{bd x^2}{(a+b)(a-b)} - \frac{bd \ln\left(1-\frac{(a+b)e^{2fx+2e}}{a-b}\right)x}{f(a+b)(a-b)} + \frac{2bde}{f(a+b)(a-b)}$

[In] `int((d*x+c)/(a+b*coth(f*x+e)), x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{2}/(a+b)*d*x^2+1/(a+b)*c*x+2/f*b/(a+b)*c/(a-b)*ln(exp(f*x+e))-1/f*b/(a+b)*c/(a-b)*ln(exp(2*f*x+2*e))*a+b*exp(2*f*x+2*e)-a+b)+b/(a+b)/(a-b)*d*x^2-1/f*b/(a+b)/(a-b)*d*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x+2/f*b/(a+b)/(a-b)*d*e*x-1/f^2*b/(a+b)/(a-b)*d*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e+1/f^2*b/(a+b)/(a-b)*d*e^2-1/2/f^2*b/(a+b)/(a-b)*d*polylog(2, (a+b)*exp(2*f*x+2*e)/(a-b))-2/f^2*b/(a+b)*d*e/(a-b)*ln(exp(f*x+e))+1/f^2*b/(a+b)*d*e/(a-b)*ln(exp(2*f*x+2*e))*a+b*exp(2*f*x+2*e)-a+b$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 300 vs. $2(103) = 206$.

Time = 0.28 (sec), antiderivative size = 300, normalized size of antiderivative = 2.78

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \frac{(a + b)df^2x^2 + 2(a + b)cf^2x - 2bd\text{Li}_2\left(\sqrt{\frac{a+b}{a-b}}(\cosh(fx+e) + \sinh(fx+e))\right) - 2bd\text{Li}_2\left(-\sqrt{\frac{a+b}{a-b}}(\cosh(fx+e) - \sinh(fx+e))\right)}{(a + b)^2f^2x^2 + 2(a + b)cfx^2 - 2b^2d^2\text{dilog}(\sqrt{(a + b)/(a - b)})^2}$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e)),x, algorithm="fricas")`

[Out] $\frac{1}{2}((a + b)*d*f^2*x^2 + 2*(a + b)*c*f^2*x - 2*b*d*\text{dilog}(\sqrt{(a + b)/(a - b)})(\cosh(f*x + e) + \sinh(f*x + e))) - 2*b*d*\text{dilog}(-\sqrt{(a + b)/(a - b)})(\cosh(f*x + e) - \sinh(f*x + e)) + 2*(b*d*e - b*c*f)*\log(2*(a + b)*\cosh(f*x + e) + 2*(a + b)*\sinh(f*x + e) + 2*(a - b)*\sqrt{(a + b)/(a - b)}) + 2*(b*d*e - b*c*f)*\log(2*(a + b)*\cosh(f*x + e) + 2*(a + b)*\sinh(f*x + e) - 2*(a - b)*\sqrt{(a + b)/(a - b)}) - 2*(b*d*f*x + b*d*e)*\log(\sqrt{(a + b)/(a - b)})(\cosh(f*x + e) + \sinh(f*x + e)) + 1) - 2*(b*d*f*x + b*d*e)*\log(-\sqrt{(a + b)/(a - b)})(\cosh(f*x + e) + \sinh(f*x + e)) + 1))/((a^2 - b^2)*f^2)$

Sympy [F]

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \int \frac{c + dx}{a + b \coth(e + fx)} dx$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e)),x)`

[Out] `Integral((c + d*x)/(a + b*coth(e + f*x)), x)`

Maxima [F]

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \int \frac{dx + c}{b \coth(fx + e) + a} dx$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e)),x, algorithm="maxima")`

[Out] $\frac{-1}{2}(4*b*\text{integrate}(-x/(a^2 - b^2 - (a^2 e^{(2*e)} + 2*a*b*e^{(2*e)} + b^2 e^{(2*e)})*e^{(2*f*x)}), x) - x^2/(a + b))*d - c*(b*\log(-(a - b)*e^{(-2*f*x - 2*e)} + a + b)/((a^2 - b^2)*f) - (f*x + e)/((a + b)*f))$

Giac [F]

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \int \frac{dx + c}{b \coth(fx + e) + a} dx$$

[In] integrate((d*x+c)/(a+b*coth(f*x+e)),x, algorithm="giac")
[Out] integrate((d*x + c)/(b*coth(f*x + e) + a), x)

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{a + b \coth(e + fx)} dx = \int \frac{c + dx}{a + b \coth(e + fx)} dx$$

[In] int((c + d*x)/(a + b*coth(e + f*x)),x)
[Out] int((c + d*x)/(a + b*coth(e + f*x)), x)

3.55 $\int \frac{1}{(c+dx)(a+b\coth(e+fx))} dx$

Optimal result	368
Rubi [N/A]	368
Mathematica [N/A]	369
Maple [N/A] (verified)	369
Fricas [N/A]	369
Sympy [N/A]	369
Maxima [N/A]	370
Giac [N/A]	370
Mupad [N/A]	370

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\coth(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\coth(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*cOTH(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b\coth(e+fx))} dx = \int \frac{1}{(c+dx)(a+b\coth(e+fx))} dx$$

[In] Int[1/((c + d*x)*(a + b*COTH[e + f*x])),x]

[Out] Defer[Int][1/((c + d*x)*(a + b*COTH[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b\coth(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 7.38 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx$$

[In] `Integrate[1/((c + d*x)*(a + b*Coth[e + f*x])),x]`

[Out] `Integrate[1/((c + d*x)*(a + b*Coth[e + f*x])), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \coth(fx + e))} dx$$

[In] `int(1/(d*x+c)/(a+b*coth(f*x+e)),x)`

[Out] `int(1/(d*x+c)/(a+b*coth(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.24 (sec) , antiderivative size = 27, normalized size of antiderivative = 1.35

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e)),x, algorithm="fricas")`

[Out] `integral(1/(a*d*x + a*c + (b*d*x + b*c)*coth(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.18 (sec) , antiderivative size = 17, normalized size of antiderivative = 0.85

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(a + b \coth(e + fx))(c + dx)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e)),x)`

[Out] `Integral(1/((a + b*coth(e + f*x))*(c + d*x)), x)`

Maxima [N/A]

Not integrable

Time = 0.35 (sec) , antiderivative size = 117, normalized size of antiderivative = 5.85

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e)),x, algorithm="maxima")`

[Out] `-2*b*integrate(-1/(a^2*c - b^2*c + (a^2*d - b^2*d)*x - (a^2*c*e^(2*e) + 2*a*b*c*e^(2*e) + b^2*c*e^(2*e) + (a^2*d*e^(2*e) + 2*a*b*d*e^(2*e) + b^2*d*e^(2*e))*x)*e^(2*f*x)), x) + log(d*x + c)/(a*d + b*d)`

Giac [N/A]

Not integrable

Time = 0.28 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e)),x, algorithm="giac")`

[Out] `integrate(1/((d*x + c)*(b*coth(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 1.91 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))} dx = \int \frac{1}{(a + b \coth(e + f x)) (c + d x)} dx$$

[In] `int(1/((a + b*coth(e + f*x))*(c + d*x)),x)`

[Out] `int(1/((a + b*coth(e + f*x))*(c + d*x)), x)`

3.56 $\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx$

Optimal result	371
Rubi [N/A]	371
Mathematica [N/A]	372
Maple [N/A] (verified)	372
Fricas [N/A]	372
Sympy [N/A]	373
Maxima [N/A]	373
Giac [N/A]	373
Mupad [N/A]	374

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\coth(e+fx))}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*cOTH(f*x+e)),x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx$$

[In] Int[1/((c + d*x)^2*(a + b*COTH[e + f*x])),x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*COTH[e + f*x])), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx$$

Mathematica [N/A]

Not integrable

Time = 13.54 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx = \int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx$$

[In] `Integrate[1/((c + d*x)^2*(a + b*Coth[e + f*x])),x]`

[Out] `Integrate[1/((c + d*x)^2*(a + b*Coth[e + f*x])), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx+c)^2(a+b\coth(fx+e))} dx$$

[In] `int(1/(d*x+c)^2/(a+b*cOTH(f*x+e)),x)`

[Out] `int(1/(d*x+c)^2/(a+b*cOTH(f*x+e)),x)`

Fricas [N/A]

Not integrable

Time = 0.25 (sec) , antiderivative size = 51, normalized size of antiderivative = 2.55

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))} dx = \int \frac{1}{(dx+c)^2(b\coth(fx+e)+a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*cOTH(f*x+e)),x, algorithm="fricas")`

[Out] `integral(1/(a*d^2*x^2 + 2*a*c*d*x + a*c^2 + (b*d^2*x^2 + 2*b*c*d*x + b*c^2)*coth(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.97 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))} dx = \int \frac{1}{(a + b \coth(e + fx))(c + dx)^2} dx$$

[In] `integrate(1/(d*x+c)**2/(a+b*coth(f*x+e)),x)`

[Out] `Integral(1/((a + b*coth(e + f*x))*(c + d*x)**2), x)`

Maxima [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 202, normalized size of antiderivative = 10.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \coth(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*coth(f*x+e)),x, algorithm="maxima")`

[Out] `-2*b*integrate(-1/(a^2*c^2 - b^2*c^2 + (a^2*d^2 - b^2*d^2)*x^2 + 2*(a^2*c*d - b^2*c*d)*x - (a^2*c^2*e^(2*e) + 2*a*b*c^2*e^(2*e) + b^2*c^2*e^(2*e) + (a^2*d^2*e^(2*e) + 2*a*b*d^2*e^(2*e) + b^2*d^2*e^(2*e))*x^2 + 2*(a^2*c*d*e^(2*e) + 2*a*b*c*d*e^(2*e) + b^2*c*d*e^(2*e))*x)*e^(2*f*x)), x) - 1/(a*c*d + b*c*d + (a*d^2 + b*d^2)*x)`

Giac [N/A]

Not integrable

Time = 0.37 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))} dx = \int \frac{1}{(dx + c)^2(b \coth(fx + e) + a)} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*coth(f*x+e)),x, algorithm="giac")`

[Out] `integrate(1/((d*x + c)^2*(b*coth(f*x + e) + a)), x)`

Mupad [N/A]

Not integrable

Time = 2.11 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))} dx = \int \frac{1}{(a + b \coth(e + fx)) (c + dx)^2} dx$$

[In] `int(1/((a + b*coth(e + f*x))*(c + d*x)^2),x)`

[Out] `int(1/((a + b*coth(e + f*x))*(c + d*x)^2), x)`

3.57 $\int \frac{(c+dx)^3}{(a+b \coth(e+fx))^2} dx$

Optimal result	376
Rubi [A] (verified)	377
Mathematica [A] (verified)	386
Maple [B] (verified)	386
Fricas [B] (verification not implemented)	388
Sympy [F(-2)]	388
Maxima [A] (verification not implemented)	388
Giac [F]	389
Mupad [F(-1)]	389

Optimal result

Integrand size = 20, antiderivative size = 638

$$\begin{aligned}
\int \frac{(c+dx)^3}{(a+b \coth(e+fx))^2} dx = & -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} \\
& + \frac{(c+dx)^4}{4(a-b)^2 d} + \frac{3b^2 d(c+dx)^2 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
& - \frac{2b(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
& + \frac{2b^2(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
& + \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
& - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
& + \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
& - \frac{3b^2 d^3 \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4} \\
& + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
& - \frac{3b^2 d^2(c+dx) \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
& - \frac{3bd^3 \text{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} \\
& + \frac{3b^2 d^3 \text{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4}
\end{aligned}$$

```
[Out] -2*b^2*(d*x+c)^3/(a^2-b^2)^2/f+2*b^2*(d*x+c)^3/(a-b)/(a+b)^2/(a-b-(a+b)*exp(2*f*x+2*e))/f+1/4*(d*x+c)^4/(a-b)^2/d+3*b^2*d*(d*x+c)^2*ln(1-(a+b)*exp(2*f*x+2*e))/(a-b)/(a-b)^2/(a^2-b^2)^2/f^2-2*b*(d*x+c)^3*ln(1-(a+b)*exp(2*f*x+2*e))/(a-b)/(a-b)^2/(a+b)/f+2*b^2*(d*x+c)^3*ln(1-(a+b)*exp(2*f*x+2*e))/(a-b)/(a^2-b^2)^2/f^2+3*b^2*d*(d*x+c)^2*polylog(2,(a+b)*exp(2*f*x+2*e))/(a-b)/(a^2-b^2)^2/f^2-3*b*d*(d*x+c)^2*polylog(2,(a+b)*exp(2*f*x+2*e))/(a-b)/(a-b)^2/(a+b)/f^2+3*b^2*d*(d*x+c)^2*polylog(2,(a+b)*exp(2*f*x+2*e))/(a-b)/(a^2-b^2)^2/f^2-3/
```

$$2*b^2*d^3*polylog(3, (a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^4+3*b*d^2*(d*x+c)*polylog(3, (a+b)*exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^3-3*b^2*d^2*(d*x+c)*polylog(3, (a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^3-3/2*b*d^3*polylog(4, (a+b)*exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^4+3/2*b^2*d^3*polylog(4, (a+b)*exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^4$$

Rubi [A] (verified)

Time = 1.62 (sec) , antiderivative size = 638, normalized size of antiderivative = 1.00, number of steps used = 28, number of rules used = 10, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500, Rules

used = {3815, 2286, 2216, 2215, 2221, 2611, 6744, 2320, 6724, 2222}

$$\int \frac{(c+dx)^3}{(a+b \coth(e+fx))^2} dx = \frac{3b^2 d^2 (c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\ - \frac{3b^2 d^2 (c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\ + \frac{3b^2 d (c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2 - b^2)^2} \\ + \frac{3b^2 d (c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2 - b^2)^2} \\ + \frac{2b^2 (c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f (a^2 - b^2)^2} \\ - \frac{2b^2 (c+dx)^3}{f (a^2 - b^2)^2} - \frac{3b^2 d^3 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a^2 - b^2)^2} \\ + \frac{3b^2 d^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a^2 - b^2)^2} \\ + \frac{2b^2 (c+dx)^3}{f(a-b)(a+b)^2 (- (a+b)e^{2e+2fx} + a-b)} \\ + \frac{3bd^2 (c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a-b)^2 (a+b)} \\ - \frac{3bd (c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a-b)^2 (a+b)} \\ - \frac{2b (c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f (a-b)^2 (a+b)} \\ + \frac{(c+dx)^4}{4d(a-b)^2} - \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2f^4 (a-b)^2 (a+b)}$$

[In] Int[(c + d*x)^3/(a + b*Coth[e + f*x])^2, x]

[Out]
$$\begin{aligned} & \frac{(-2*b^2*(c+d*x)^3)/((a^2 - b^2)^2*f) + (2*b^2*(c+d*x)^3)/((a - b)*(a + b)^2*(a - b - (a + b)*E^(2*e + 2*f*x))*f) + (c + d*x)^4/(4*(a - b)^2*d) + (3*b^2*d*(c + d*x)^2*\operatorname{Log}[1 - ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f^2) - (2*b*(c + d*x)^3*\operatorname{Log}[1 - ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a - b)^2*(a + b)*f) + (2*b^2*(c + d*x)^3*\operatorname{Log}[1 - ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a - b))/((a^2 - b^2)^2*f) + (3*b^2*d^2*(c + d*x)*\operatorname{PolyLog}[2, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f^3) - (3*b*d*(c + d*x)^2*\operatorname{PolyLog}[2, ((a + b)*E^(2*e + 2*f*x))/(a - b)])}{4d(a - b)^2} \end{aligned}$$

$$(a + b)E^{(2e + 2fx)} / (a - b)] / ((a - b)^2(a + b)f^2) + (3b^2d(c + dx)^2 \text{PolyLog}[2, ((a + b)E^{(2e + 2fx)} / (a - b))] / ((a^2 - b^2)^2f^2) - (3b^2d^3 \text{PolyLog}[3, ((a + b)E^{(2e + 2fx)} / (a - b))] / (2(a^2 - b^2)^2f^4) + (3bd^2(c + dx) \text{PolyLog}[3, ((a + b)E^{(2e + 2fx)} / (a - b))] / ((a - b)^2(a + b)f^3) - (3b^2d^2(c + dx) \text{PolyLog}[3, ((a + b)E^{(2e + 2fx)} / (a - b))] / ((a^2 - b^2)^2f^3) - (3bd^3 \text{PolyLog}[4, ((a + b)E^{(2e + 2fx)} / (a - b))] / (2(a - b)^2(a + b)f^4) + (3b^2d^3 \text{PolyLog}[4, ((a + b)E^{(2e + 2fx)} / (a - b))] / (2(a^2 - b^2)^2f^4)$$
Rule 2215

$$\text{Int}[(c_+ + d_-)(x_-)^m / ((a_+ + b_-)(F_-)^e(x_-)^f)^n, x] \rightarrow \text{Simp}[(c + dx)^{m+1} / (a*d(m+1)), x] - \text{Dist}[b/a, \text{Int}[(c + dx)^m((F^e(g*(e+fx)))^n / (a + b*(F^e(g*(e+fx)))^n)), x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$$
Rule 2216

$$\text{Int}[(a_+ + b_-)(F_-)^e(x_-)^f)^n / ((c_+ + d_-)(x_-)^m)^p, x] \rightarrow \text{Dist}[1/a, \text{Int}[(c + dx)^m(a + b*(F^e(g*(e+fx)))^n)^{p+1}, x], x] - \text{Dist}[b/a, \text{Int}[(c + dx)^m(F^e(g*(e+fx)))^n / (a + b*(F^e(g*(e+fx)))^n)^p, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{ILtQ}[p, 0] \&& \text{IGtQ}[m, 0]$$
Rule 2221

$$\text{Int}[(F_-)^e(x_-)^f)^n / ((a_+ + b_-)(F_-)^e(x_-)^f)^{m-1}, x] \rightarrow \text{Simp}[(c + dx)^m / (b*f*g*n*\text{Log}[F]) * \text{Log}[1 + b*((F^e(g*(e+fx)))^n/a)], x] - \text{Dist}[d*(m/(b*f*g*n*\text{Log}[F])), \text{Int}[(c + dx)^{m-1} * \text{Log}[1 + b*((F^e(g*(e+fx)))^n/a)], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \text{IGtQ}[m, 0]$$
Rule 2222

$$\text{Int}[(F_-)^e(x_-)^f)^n / ((a_+ + b_-)(F_-)^e(x_-)^f)^{m-1}, x] \rightarrow \text{Simp}[(c + dx)^m / (b*f*g*n*(p+1)*\text{Log}[F]), x] - \text{Dist}[d*(m/(b*f*g*n*(p+1)*\text{Log}[F])), \text{Int}[(c + dx)^{m-1}(a + b*(F^e(g*(e+fx)))^n)^{p+1}, x], x] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, m, n, p\}, x] \&& \text{NeQ}[p, -1]$$
Rule 2286

$$\text{Int}[(a_+ + b_-)(F_-)^u)^p / ((c_+ + d_-)(F_-)^v)^q, x] \rightarrow \text{With}[\{w = \text{ExpandIntegrand}[(e + fx)^m, (a + b*F^u)^p * (c + d*F^v)^q, x]\}, \text{Int}[w, x] /; \text{SumQ}[w]] /; \text{FreeQ}[\{F, a, b, c, d, e, f, m\}, x] \&& \text{IntegersQ}[p, q] \&& \text{LinearQ}[\{u, v\}, x] \&& \text{RationalQ}[\text{Simp}$$

`lify[u/v]]`

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x]
, Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n]] && !MatchQ[u, E^((c_)*(a_.) + (b_)*x))*(F_)[v_] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2611

```
Int[Log[1 + (e_)*(F_)^((c_)*(a_.) + (b_)*(x_)))^(n_.)]*((f_.) + (g_.)
*(x_))^(m_), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a +
b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m
- 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e,
f, g, n}, x] && GtQ[m, 0]
```

Rule 3815

```
Int[((c_.) + (d_.)*(x_))^(m_.)*((a_) + (b_)*tan[(e_.) + (f_.)*(x_)])^(n_),
x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 +
b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^(-n), x], x] /; FreeQ[{a, b, c, d,
e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IgtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_)*(a_.) + (b_)*(x_))^(p_.)]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d,
e, n, p}, x] && EqQ[b*d, a*e]
```

Rule 6744

```
Int[((e_.) + (f_.)*(x_))^(m_.)*PolyLog[n_, (d_.)*((F_)^((c_)*(a_.) + (b_).
*(x_)))^(p_.)], x_Symbol] :> Simp[(e + f*x)^m*(PolyLog[n + 1, d*(F^(c*(a +
b*x)))^p]/(b*c*p*Log[F])), x] - Dist[f*(m/(b*c*p*Log[F])), Int[(e + f*x)^(m
- 1)*PolyLog[n + 1, d*(F^(c*(a + b*x)))^p], x], x] /; FreeQ[{F, a, b, c,
d, e, f, n, p}, x] && GtQ[m, 0]
```

Rubi steps

$$\text{integral} = \int \left(\frac{(c+dx)^3}{(a-b)^2} + \frac{4b^2 e^{4e+4fx} (c+dx)^3}{(a-b)^2 \left(a \left(1-\frac{b}{a}\right) - a \left(1+\frac{b}{a}\right) e^{2e+2fx}\right)^2} \right. \\ \left. + \frac{4be^{2e+2fx} (c+dx)^3}{(a-b)^2 \left(a \left(1-\frac{b}{a}\right) - a \left(1+\frac{b}{a}\right) e^{2e+2fx}\right)} \right) dx$$

$$\begin{aligned}
&= \frac{(c+dx)^4}{4(a-b)^2d} + \frac{(4b) \int \frac{e^{2e+2fx}(c+dx)^3}{a(1-\frac{b}{a})-a(1+\frac{b}{a})e^{2e+2fx}} dx}{(a-b)^2} + \frac{(4b^2) \int \frac{e^{4e+4fx}(c+dx)^3}{(a(1-\frac{b}{a})-a(1+\frac{b}{a})e^{2e+2fx})^2} dx}{(a-b)^2} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{(4b^2) \int \left(\frac{(c+dx)^3}{(a+b)^2} + \frac{(a-b)^2(c+dx)^3}{(a+b)^2(a-b-(a+b)e^{2e+2fx})^2} + \frac{2(-a+b)(c+dx)^3}{(a+b)^2(a-b-(a+b)e^{2e+2fx})}\right) dx}{(a-b)^2} \\
&\quad + \frac{(6bd) \int (c+dx)^2 \log\left(1 - \frac{(1+\frac{b}{a})e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} + \frac{b^2(c+dx)^4}{(a^2-b^2)^2d} - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^3}{(a-b+(-a-b)e^{2e+2fx})^2} dx}{(a+b)^2} \\
&\quad - \frac{(8b^2) \int \frac{(c+dx)^3}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} + \frac{(6bd^2) \int (c+dx) \text{PolyLog}\left(2, \frac{(1+\frac{b}{a})e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f^2} \\
&= \frac{(c+dx)^4}{4(a-b)^2d} - \frac{b^2(c+dx)^4}{(a^2-b^2)^2d} - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad + \frac{(4b^2) \int \frac{(c+dx)^3}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} - \frac{(8b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} \\
&\quad + \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{(a-b+(-a-b)e^{2e+2fx})^2} dx}{a^2-b^2} - \frac{(3bd^3) \int \text{PolyLog}\left(3, \frac{(1+\frac{b}{a})e^{2e+2fx}}{1-\frac{b}{a}}\right) dx}{(a-b)^2(a+b)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{4b^2(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad + \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^3}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} - \frac{(3bd^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a-b)^2(a+b)f^4} \\
&\quad - \frac{(6b^2d) \int \frac{(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2f} - \frac{(12b^2d) \int (c+dx)^2 \log\left(1 + \frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{6b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} \\
&\quad - \frac{(12b^2d^2) \int (c+dx) \operatorname{PolyLog}\left(2, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2} \\
&\quad - \frac{(6b^2d) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)f} + \frac{(6b^2d) \int (c+dx)^2 \log\left(1 + \frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} \\
&\quad + \frac{(c+dx)^4}{4(a-b)^2 d} + \frac{3b^2 d(c+dx)^2 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad - \frac{2b(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
&\quad - \frac{3bd(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{3b^2 d(c+dx)^2 \text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{6b^2 d^2(c+dx) \text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd^3 \text{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{(6b^2 d^3) \int \text{PolyLog}\left(3, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{(6b^2 d^2) \int (c+dx) \log\left(1+\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2} \\
&\quad + \frac{(6b^2 d^2) \int (c+dx) \text{PolyLog}\left(2, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2 (a-b-(a+b)e^{2e+2fx}) f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^3 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^4} \\
&\quad - \frac{(3b^2d^3) \int \operatorname{PolyLog}\left(2, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{(3b^2d^3) \int \operatorname{PolyLog}\left(3, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2 f^3}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^4}{4(a-b)^2 d} \\
&\quad + \frac{3b^2d(c+dx)^2 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{3b^2d^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^4} \\
&\quad - \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a^2-b^2)^2 f^4} \\
&\quad - \frac{(3b^2d^3) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(3, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{2(a^2-b^2)^2 f^4} \\
&= -\frac{2b^2(c+dx)^3}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^3}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} \\
&\quad + \frac{(c+dx)^4}{4(a-b)^2 d} + \frac{3b^2d(c+dx)^2 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad - \frac{2b(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^3 \log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\
&\quad + \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} - \frac{3bd(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{3b^2d(c+dx)^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{3b^2d^3 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4} \\
&\quad + \frac{3bd^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{3b^2d^2(c+dx) \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{3bd^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a-b)^2(a+b)f^4} + \frac{3b^2d^3 \operatorname{PolyLog}\left(4, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{2(a^2-b^2)^2 f^4}
\end{aligned}$$

Mathematica [A] (verified)

Time = 7.97 (sec) , antiderivative size = 659, normalized size of antiderivative = 1.03

$$\int \frac{(c+dx)^3}{(a+b\coth(e+fx))^2} dx \\ = \frac{16bc^2f^3(-3bd+2acf)x - \frac{16(a-b)b^2f^3(c+dx)^3}{a(-1+e^{2e})+b(1+e^{2e})} + \frac{8a(a-b)bf^4(c+dx)^4}{d(a(-1+e^{2e})+b(1+e^{2e}))}}{48bcd^2(bd-acf)x \log\left(1+\frac{(-a+b)e^{-2(e+fx)}}{a+b}\right)}$$

[In] `Integrate[(c + d*x)^3/(a + b*Coth[e + f*x])^2, x]`

[Out]
$$(16*b*c^2*f^3*(-3*b*d + 2*a*c*f)*x - (16*(a - b)*b^2*f^3*(c + d*x)^3)/(a*(-1 + E^(2*e)) + b*(1 + E^(2*e))) + (8*a*(a - b)*b*f^4*(c + d*x)^4)/(d*(a*(-1 + E^(2*e)) + b*(1 + E^(2*e)))) + 48*b*c*d*f^2*(b*d - a*c*f)*x*\text{Log}[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))] + 24*b*d^2*f^2*(b*d - 2*a*c*f)*x^2*\text{Log}[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))] - 16*a*b*d^3*f^3*x^3*\text{Log}[1 + (-a + b)/(a + b)*E^(2*(e + f*x)))] + 8*b*c^2*f^2*(3*b*d - 2*a*c*f)*\text{Log}[a - b - (a + b)*E^(2*(e + f*x))] + 24*b*c*d*f*(-(b*d) + a*c*f)*\text{PolyLog}[2, (a - b)/((a + b)*E^(2*(e + f*x)))] - 12*b*d^2*(b*d - 2*a*c*f)*(2*f*x*\text{PolyLog}[2, (a - b)/(a + b)*E^(2*(e + f*x)))] + \text{PolyLog}[3, (a - b)/((a + b)*E^(2*(e + f*x)))] + 12*a*b*d^3*(2*f^2*x^2*\text{PolyLog}[2, (a - b)/((a + b)*E^(2*(e + f*x)))] + 2*f*x*\text{PolyLog}[3, (a - b)/((a + b)*E^(2*(e + f*x)))] + \text{PolyLog}[4, (a - b)/((a + b)*E^(2*(e + f*x)))] - ((a - b)*(a + b)*f^3*((a^2 + b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*\text{Cosh}[f*x] - (a^2 - b^2)*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3)*\text{Cosh}[2*e + f*x] + 2*b*(-4*b*(c + d*x)^3 + a*f*x*(4*c^3 + 6*c^2*d*x + 4*c*d^2*x^2 + d^3*x^3))*\text{Sinh}[f*x]))/((b*\text{Cosh}[e] + a*\text{Sinh}[e])*(b*\text{Cosh}[e + f*x] + a*\text{Sinh}[e + f*x])))/(8*(a - b)^2*(a + b)^2*f^4)$$

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 2443 vs. $2(618) = 1236$.

Time = 0.58 (sec) , antiderivative size = 2444, normalized size of antiderivative = 3.83

method	result	size
risch	Expression too large to display	2444

[In] `int((d*x+c)^3/(a+b*coth(f*x+e))^2, x, method=_RETURNVERBOSE)`

[Out]
$$\frac{3/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*e^2*d^3*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)+3/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*d^3*ln(1-(a+b)*\exp(2*f*x+2*e)/(a-b))*x^2-3/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*d^3*ln(1-(a+b)*\exp(2*f*x+2*e)/(a-b))*e^2-2/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^3*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)+3/(a^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*c^2*d*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)-6/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*e^2*d^3*ln(\exp(f*x+e))}{}$$

$$\begin{aligned}
& +4/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^3*ln(exp(f*x+e))-6/(a^2+2*a*b+b^2)/f^2*b \\
& -2/(a-b)^2*c^2*d*ln(exp(f*x+e))-6/(a^2+2*a*b+b^2)/f*b/(a-b)^2*d^2*c*a*ln(1- \\
& (a+b)*exp(2*f*x+2*e)/(a-b))*x^2-6/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*d^2*c*a*p \\
& ylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))*x+12/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*e^2*d \\
& -2*c*a*ln(exp(f*x+e))-12/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*e*a*c^2*d*ln(exp(f*x \\
& +e))-2/(a^2+2*a*b+b^2)/f*b^2/(a-b)^2*d^3*x^3+4/(a^2+2*a*b+b^2)/f^4*b^2/(a-b) \\
&)^2*d^3*e^3-3/2/(a^2+2*a*b+b^2)/f^4*b^2/(a-b)^2*d^3*polylog(3,(a+b)*exp(2*f \\
& *x+2*e)/(a-b))+4/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*d^3*a*e^3*x+6/(a^2+2*a*b+b^2) \\
&)*b/(a-b)^2*a*c^2*d*x^2-2/(a^2+2*a*b+b^2)/f^4*b/(a-b)^2*d^3*a*ln(1-(a+b)*ex \\
& p(2*f*x+2*e)/(a-b))*e^3+3/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*d^3*a*polylog(3,(a+ \\
& b)*exp(2*f*x+2*e)/(a-b))*x+2/(a^2+2*a*b+b^2)/f^4*b/(a-b)^2*e^3*d^3*a*ln(exp \\
& (2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)-2/(a^2+2*a*b+b^2)/f*b/(a-b)^2*d^3*a*ln(\\
& 1-(a+b)*exp(2*f*x+2*e)/(a-b))*x^3+3/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*d^2*c*a*p \\
& olylog(3,(a+b)*exp(2*f*x+2*e)/(a-b))-3/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*a*c^2* \\
& d*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))-6/(a^2+2*a*b+b^2)/f^3*b^2/(a-b)^2*e \\
& *d^2*c*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)+6/(a^2+2*a*b+b^2)/f^2*b^2/ \\
& (a-b)^2*d^2*c*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x+6/(a^2+2*a*b+b^2)/f^3*b^2/ \\
& (a-b)^2*d^2*c*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e-3/(a^2+2*a*b+b^2)/f^2*b/(a \\
& -b)^2*d^3*a*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))*x^2-2/(a-b)/f/(a^2+2*a*b+ \\
& b^2)*(d^3*x^3+3*c*d^2*x^2+3*c^2*d*x+c^3)*b^2/(exp(2*f*x+2*e)*a+b*exp(2*f*x+ \\
& 2*e)-a+b)-6/(a^2+2*a*b+b^2)/f*b^2/(a-b)^2*d^2*c*x^2-6/(a^2+2*a*b+b^2)/f^3*b \\
& ^2/(a-b)^2*d^2*c*e^2+1/(a^2+2*a*b+b^2)*b/(a-b)^2*d^3*a*x^4+3/(a^2+2*a*b+b^2) \\
& /f^4*b/(a-b)^2*d^3*a*x^4+6/(a^2+2*a*b+b^2)/f^3*b^2/(a-b)^2*d^3*e^2*x+3/(a^ \\
& 2+2*a*b+b^2)/f^3*b^2/(a-b)^2*d^3*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))*x+3/ \\
& (a^2+2*a*b+b^2)/f^3*b^2/(a-b)^2*d^2*c*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b)) \\
& -3/2/(a^2+2*a*b+b^2)/f^4*b/(a-b)^2*d^3*a*polylog(4,(a+b)*exp(2*f*x+2*e)/(a- \\
& b))+6/(a^2+2*a*b+b^2)/f^3*b/(a-b)^2*d^2*c*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b) \\
&)*e^2+1/4*d^3/(a^2+2*a*b+b^2)*x^4+1/4/d/(a^2+2*a*b+b^2)*c^4+12/(a^2+2*a*b+b \\
& ^2)/f*b/(a-b)^2*a*c^2*d*e*x-12/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*d^2*c*a*e^2*x- \\
& 6/(a^2+2*a*b+b^2)/f*b/(a-b)^2*a*c^2*d*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x-6/ \\
& (a^2+2*a*b+b^2)/f^3*b/(a-b)^2*e^2*d^2*c*a*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2 \\
& *e)-a+b)+6/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*e*a*c^2*d*ln(exp(2*f*x+2*e)*a+b*ex \\
& p(2*f*x+2*e)-a+b)-6/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*a*c^2*d*ln(1-(a+b)*exp(2* \\
& f*x+2*e)/(a-b))*e+4/(a^2+2*a*b+b^2)*b/(a-b)^2*d^2*c*a*x^3-8/(a^2+2*a*b+b^2) \\
& /f^3*b/(a-b)^2*d^2*c*a*e^3+6/(a^2+2*a*b+b^2)/f^2*b/(a-b)^2*a*c^2*d*e^2-12/(a \\
& ^2+2*a*b+b^2)/f^2*b^2/(a-b)^2*d^2*c*e*x+d^2/(a^2+2*a*b+b^2)*c*x^3+3/2*d/(a \\
& ^2+2*a*b+b^2)*c^2*x^2+1/(a^2+2*a*b+b^2)*c^3*x+12/(a^2+2*a*b+b^2)/f^3*b^2/(a \\
& -b)^2*e*d^2*c*ln(exp(f*x+e))-4/(a^2+2*a*b+b^2)/f^4*b/(a-b)^2*e^3*d^3*a*ln(e \\
& xp(f*x+e))
\end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 6171 vs. $2(614) = 1228$.

Time = 0.36 (sec), antiderivative size = 6171, normalized size of antiderivative = 9.67

$$\int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3/(a+b*coth(f*x+e))^2,x, algorithm="fricas")`

[Out] Too large to include

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx = \text{Exception raised: TypeError}$$

[In] `integrate((d*x+c)**3/(a+b*coth(f*x+e))**2,x)`

[Out] Exception raised: TypeError >> Invalid NaN comparison

Maxima [A] (verification not implemented)

none

Time = 0.53 (sec), antiderivative size = 1056, normalized size of antiderivative = 1.66

$$\int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx = \text{Too large to display}$$

[In] `integrate((d*x+c)^3/(a+b*coth(f*x+e))^2,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -6*b^2*c^2*d*f*x/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - 2/3*(4*f^3*x^3*log(-(a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b) + 1) + 6*f^2*x^2*dilog((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b)) - 6*f*x*polylog(3, (a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b))) + 3*polylog(4, (a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b)))*a*b*d^3/(a^4*f^4 - 2*a^2*b^2*f^4 + b^4*f^4) + 3*b^2*c^2*d*log((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)} - a + b)/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - c^3*(2*a*b*log(-(a - b)*e^{(-2*f*x - 2*e)} + a + b)/((a^4 - 2*a^2*b^2 + b^4)*f) + 2*b^2/((a^4 - 2*a^2*b^2 + b^4 - (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^{(-2*f*x - 2*e)})*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f)) - 3/2*(2*a*b*c*d^2*f - b^2*d^3)*(2*f^2*x^2*log(-(a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b) + 1) + 2*f*x*dilog((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b)) - polylog(3, (a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b))) \end{aligned}$$

$$\begin{aligned}
& + b^*e^{(2*e)}*e^{(2*f*x)}/(a - b)))/(a^4*f^4 - 2*a^2*b^2*f^4 + b^4*f^4) - 3*(a \\
& *b*c^2*d*f - b^2*c*d^2)*(2*f*x*log(-(a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - \\
& b) + 1) + \text{dilog}((a*e^{(2*e)} + b*e^{(2*e)})*e^{(2*f*x)}/(a - b))/(a^4*f^3 - 2*a^ \\
& 2*b^2*f^3 + b^4*f^3) + (a*b*d^3*f^4*x^4 + 2*(2*a*b*c*d^2*f - b^2*d^3)*f^3*x^ \\
& 3 + 6*(a*b*c^2*d*f^2 - b^2*c*d^2*f)*f^2*x^2)/(a^4*f^4 - 2*a^2*b^2*f^4 + b^ \\
& 4*f^4) + 1/4*(24*b^2*c^2*d*x + (a^2*d^3*f - 2*a*b*d^3*f + b^2*d^3*f)*x^4 + \\
& 4*(a^2*c*d^2*f - 2*a*b*c*d^2*f + (c*d^2*f + 2*d^3)*b^2)*x^3 + 6*(a^2*c^2*d^ \\
& f - 2*a*b*c^2*d*f + (c^2*d*f + 4*c*d^2)*b^2)*x^2 - ((a^2*d^3*f*e^{(2*e)} - b^ \\
& 2*d^3*f*e^{(2*e)})*x^4 + 4*(a^2*c*d^2*f*e^{(2*e)} - b^2*c*d^2*f*e^{(2*e)})*x^3 + \\
& 6*(a^2*c^2*d*f*e^{(2*e)} - b^2*c^2*d*f*e^{(2*e)})*x^2)*e^{(2*f*x)})/(a^4*f - 2*a^ \\
& 2*b^2*f + b^4*f - (a^4*f*e^{(2*e)} + 2*a^3*b*f*e^{(2*e)} - 2*a*b^3*f*e^{(2*e)} - \\
& b^4*f*e^{(2*e)})*e^{(2*f*x)})
\end{aligned}$$

Giac [F]

$$\int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx = \int \frac{(dx + c)^3}{(b \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)^3/(a+b*coth(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)^3/(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx = \int \frac{(c + dx)^3}{(a + b \coth(e + fx))^2} dx$$

[In] `int((c + d*x)^3/(a + b*coth(e + f*x))^2,x)`
[Out] `int((c + d*x)^3/(a + b*coth(e + f*x))^2, x)`

3.58 $\int \frac{(c+dx)^2}{(a+b \coth(e+fx))^2} dx$

Optimal result	390
Rubi [A] (verified)	391
Mathematica [A] (verified)	396
Maple [B] (verified)	397
Fricas [B] (verification not implemented)	398
Sympy [F(-2)]	400
Maxima [A] (verification not implemented)	400
Giac [F]	401
Mupad [F(-1)]	401

Optimal result

Integrand size = 20, antiderivative size = 475

$$\begin{aligned} \int \frac{(c+dx)^2}{(a+b \coth(e+fx))^2} dx = & -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2 (a-b-(a+b)e^{2e+2fx}) f} \\ & + \frac{(c+dx)^3}{3(a-b)^2 d} + \frac{2b^2 d(c+dx) \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\ & - \frac{2b(c+dx)^2 \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\ & + \frac{2b^2(c+dx)^2 \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} \\ & + \frac{b^2 d^2 \text{PolyLog} \left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\ & - \frac{2bd(c+dx) \text{PolyLog} \left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\ & + \frac{2b^2 d(c+dx) \text{PolyLog} \left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\ & + \frac{bd^2 \text{PolyLog} \left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\ & - \frac{b^2 d^2 \text{PolyLog} \left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \end{aligned}$$

[Out] $-2*b^2*(d*x+c)^2/(a^2-b^2)^2/f+2*b^2*(d*x+c)^2/(a-b)/(a+b)^2/(a-b-(a+b)*\exp(2*f*x+2*e))/f+1/3*(d*x+c)^3/(a-b)^2/d+2*b^2*d*(d*x+c)*\ln(1-(a+b)*\exp(2*f*x$

$$\begin{aligned}
& +2*e)/(a-b))/(a^2-b^2)^2/f^2-2*b*(d*x+c)^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(a-b)) \\
& /(a-b)^2/(a+b)/f+2*b^2*(d*x+c)^2*\ln(1-(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2 \\
& ^2/f+b^2*d^2*polylog(2,(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^3-2*b*d*(d \\
& *x+c)*polylog(2,(a+b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^2+2*b^2*d*(d*x+ \\
& c)*polylog(2,(a+b)*\exp(2*f*x+2*e)/(a-b))/(a^2-b^2)^2/f^2+b*d^2*polylog(3,(a \\
& +b)*\exp(2*f*x+2*e)/(a-b))/(a-b)^2/(a+b)/f^3-b^2*d^2*polylog(3,(a+b)*\exp(2*f \\
& *x+2*e)/(a-b))/(a^2-b^2)^2/f^3
\end{aligned}$$

Rubi [A] (verified)

Time = 1.22 (sec) , antiderivative size = 475, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 11, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.550, Rules used = {3815, 2286, 2216, 2215, 2221, 2611, 2320, 6724, 2222, 2317, 2438}

$$\begin{aligned}
\int \frac{(c+dx)^2}{(a+b\coth(e+fx))^2} dx = & \frac{2b^2 d(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2 - b^2)^2} \\
& + \frac{2b^2 d(c+dx) \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a^2 - b^2)^2} \\
& + \frac{2b^2 (c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f (a^2 - b^2)^2} \\
& - \frac{2b^2 (c+dx)^2}{f (a^2 - b^2)^2} + \frac{b^2 d^2 \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\
& - \frac{b^2 d^2 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a^2 - b^2)^2} \\
& + \frac{2b^2 (c+dx)^2}{f(a-b)(a+b)^2 (- (a+b)e^{2e+2fx} + a-b)} \\
& - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^2 (a-b)^2 (a+b)} \\
& - \frac{2b(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f (a-b)^2 (a+b)} \\
& + \frac{(c+dx)^3}{3d(a-b)^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{f^3 (a-b)^2 (a+b)}
\end{aligned}$$

[In] Int[(c + d*x)^2/(a + b*Coth[e + f*x])^2, x]

[Out]
$$\begin{aligned}
& (-2*b^2*(c+d*x)^2)/((a^2-b^2)^2*f) + (2*b^2*(c+d*x)^2)/((a-b)*(a+b)^2*(a-b - (a+b)*E^(2*e+2*f*x))*f) \\
& + (c+d*x)^3/(3*(a-b)^2*d) + (2*b^2*d*(c+d*x)*\operatorname{Log}[1 - ((a+b)*E^(2*e+2*f*x))/(a-b)])/((a^2-b^2)^2)
\end{aligned}$$

$$2*f^2) - (2*b*(c + d*x)^2*Log[1 - ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a - b)^2*(a + b)*f) + (2*b^2*(c + d*x)^2*Log[1 - ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f) + (b^2*d^2*PolyLog[2, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f^3) - (2*b*d*(c + d*x)*PolyLog[2, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a - b)^2*(a + b)*f^2) + (2*b^2*d*(c + d*x)*PolyLog[2, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f^2) + (b*d^2*PolyLog[3, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a - b)^2*(a + b)*f^3) - (b^2*d^2*PolyLog[3, ((a + b)*E^(2*e + 2*f*x))/(a - b)])/((a^2 - b^2)^2*f^3)$$
Rule 2215

$$\text{Int}[((c_.) + (d_.)*(x_))^{(m_.)}/((a_) + (b_.)*((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)}], \text{x_Symbol}] \rightarrow \text{Simp}[(c + d*x)^{m+1}/(a*d*(m+1)), \text{x}] - \text{Dist}[b/a, \text{Int}[(c + d*x)^m*((F^(g*(e + f*x)))^n)/(a + b*(F^(g*(e + f*x)))^n)], \text{x}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, \text{x}] \&& \text{IGtQ}[m, 0]$$
Rule 2216

$$\text{Int}[((a_) + (b_.)*((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)})^{(p_.)*((c_.) + (d_.)*(x_))^{(m_.)}}, \text{x_Symbol}] \rightarrow \text{Dist}[1/a, \text{Int}[(c + d*x)^m*(a + b*(F^(g*(e + f*x)))^n)^{p+1}], \text{x}] - \text{Dist}[b/a, \text{Int}[(c + d*x)^m*(F^(g*(e + f*x)))^{n*p}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, \text{x}] \&& \text{ILtQ}[p, 0] \&& \text{IGtQ}[m, 0]$$
Rule 2221

$$\text{Int}[(((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)}*((c_.) + (d_.)*(x_))^{(m_.)})/((a_) + (b_.)*((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)}], \text{x_Symbol}] \rightarrow \text{Simp}[((c + d*x)^m/(b*f*g*n*Log[F]))*Log[1 + b*((F^(g*(e + f*x)))^n/a)], \text{x}] - \text{Dist}[d*(m/(b*f*g*n*Log[F])), \text{Int}[(c + d*x)^{(m-1)}*Log[1 + b*((F^(g*(e + f*x)))^n/a)], \text{x}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, \text{x}] \&& \text{IGtQ}[m, 0]$$
Rule 2222

$$\text{Int}[((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)}*((a_.) + (b_.)*((F_.)^{(g_.)}*((e_.) + (f_.)*(x_))))^{(n_.)})^{(p_.)*((c_.) + (d_.)*(x_))^{(m_.)}}, \text{x_Symbol}] \rightarrow \text{Simp}[(c + d*x)^m*((a + b*(F^(g*(e + f*x)))^n)^{p+1})/(b*f*g*n*(p+1)*Log[F]), \text{x}] - \text{Dist}[d*(m/(b*f*g*n*(p+1)*Log[F))), \text{Int}[(c + d*x)^{(m-1)}*(a + b*(F^(g*(e + f*x)))^n)^{p+1}], \text{x}] /; \text{FreeQ}[\{F, a, b, c, d, e, f, g, m, n, p\}, \text{x}] \&& \text{NeQ}[p, -1]$$
Rule 2286

$$\text{Int}[((a_.) + (b_.)*(F_.)^{(u_.)})^{(p_.)}*((c_.) + (d_.)*(F_.)^{(v_.)})^{(q_.)}*((e_.) + (f_.)*(x_))^{(m_.)}], \text{x_Symbol}] \rightarrow \text{With}[\{w = \text{ExpandIntegrand}[(e + f*x)^m, (a + b*F^u)^p*(c + d*F^v)^q, \text{x}]\}, \text{Int}[w, \text{x}] /; \text{SumQ}[w]] /; \text{FreeQ}[\{F, a, b, c, d, e, f, m\}, \text{x}] \&& \text{IntegersQ}[p, q] \&& \text{LinearQ}[\{u, v\}, \text{x}] \&& \text{RationalQ}[\text{Simp}$$

`lify[u/v]]`

Rule 2317

```
Int[Log[(a_) + (b_)*((F_)^((e_.)*(c_.) + (d_)*(x_))))^(n_.)], x_Symbol]
:> Dist[1/(d*e*n*Log[F]), Subst[Int[Log[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; FreeQ[{F, a, b, c, d, e, n}, x] && GtQ[a, 0]
```

Rule 2320

```
Int[u_, x_Symbol] :> With[{v = FunctionOfExponential[u, x]}, Dist[v/D[v, x], Subst[Int[FunctionOfExponentialFunction[u, x]/x, x], x, v], x]] /; FunctionOfExponentialQ[u, x] && !MatchQ[u, (w_)*((a_.)*(v_)^(n_))^(m_)] /; FreeQ[{a, m, n}, x] && IntegerQ[m*n] && !MatchQ[u, E^((c_.)*(a_.) + (b_)*x))*(F_[v_]] /; FreeQ[{a, b, c}, x] && InverseFunctionQ[F[x]]]
```

Rule 2438

```
Int[Log[((d_) + (e_)*(x_)^(n_.))]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2611

```
Int[Log[1 + (e_.)*((F_)^((c_.)*(a_.) + (b_)*(x_))))^(n_.)]*((f_.) + (g_.)*(x_)^m_.), x_Symbol] :> Simp[(-(f + g*x)^m)*(PolyLog[2, (-e)*(F^(c*(a + b*x)))^n]/(b*c*n*Log[F])), x] + Dist[g*(m/(b*c*n*Log[F])), Int[(f + g*x)^(m - 1)*PolyLog[2, (-e)*(F^(c*(a + b*x)))^n], x], x] /; FreeQ[{F, a, b, c, e, f, g, n}, x] && GtQ[m, 0]
```

Rule 3815

```
Int[((c_.) + (d_)*(x_)^m_)*((a_) + (b_)*tan[(e_.) + (f_)*(x_)])^(n_), x_Symbol] :> Int[ExpandIntegrand[(c + d*x)^m, (1/(a - I*b) - 2*I*(b/(a^2 + b^2 + (a - I*b)^2*E^(2*I*(e + f*x)))))^(-n), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[a^2 + b^2, 0] && ILtQ[n, 0] && IGtQ[m, 0]
```

Rule 6724

```
Int[PolyLog[n_, (c_.)*((a_.) + (b_)*(x_))^p_]/((d_.) + (e_)*(x_)), x_Symbol] :> Simp[PolyLog[n + 1, c*(a + b*x)^p]/(e*p), x] /; FreeQ[{a, b, c, d, e, n, p}, x] && EqQ[b*d, a*e]
```

Rubi steps

$$\begin{aligned}
\text{integral} &= \int \left(\frac{(c+dx)^2}{(a-b)^2} + \frac{4b^2 e^{4e+4fx} (c+dx)^2}{(a-b)^2 (a(1-\frac{b}{a}) - a(1+\frac{b}{a}) e^{2e+2fx})^2} \right. \\
&\quad \left. + \frac{4be^{2e+2fx} (c+dx)^2}{(a-b)^2 (a(1-\frac{b}{a}) - a(1+\frac{b}{a}) e^{2e+2fx})} \right) dx \\
&= \frac{(c+dx)^3}{3(a-b)^2 d} + \frac{(4b) \int \frac{e^{2e+2fx} (c+dx)^2}{a(1-\frac{b}{a}) - a(1+\frac{b}{a}) e^{2e+2fx}} dx}{(a-b)^2} + \frac{(4b^2) \int \frac{e^{4e+4fx} (c+dx)^2}{(a(1-\frac{b}{a}) - a(1+\frac{b}{a}) e^{2e+2fx})^2} dx}{(a-b)^2} \\
&= \frac{(c+dx)^3}{3(a-b)^2 d} - \frac{2b(c+dx)^2 \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b} \right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{(4b^2) \int \left(\frac{(c+dx)^2}{(a+b)^2} + \frac{(a-b)^2(c+dx)^2}{(a+b)^2(a-b-(a+b)e^{2e+2fx})^2} + \frac{2(-a+b)(c+dx)^2}{(a+b)^2(a-b-(a+b)e^{2e+2fx})} \right) dx}{(a-b)^2} \\
&\quad + \frac{(4bd) \int (c+dx) \log \left(1 - \frac{(1+\frac{b}{a})e^{2e+2fx}}{1-\frac{b}{a}} \right) dx}{(a-b)^2(a+b)f} \\
&= \frac{(c+dx)^3}{3(a-b)^2 d} + \frac{4b^2(c+dx)^3}{3(a^2-b^2)^2 d} - \frac{2b(c+dx)^2 \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b} \right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{2bd(c+dx) \text{PolyLog} \left(2, \frac{(a+b)e^{2e+2fx}}{a-b} \right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^2}{(a-b+(-a-b)e^{2e+2fx})^2} dx}{(a+b)^2} \\
&\quad - \frac{(8b^2) \int \frac{(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} + \frac{(2bd^2) \int \text{PolyLog} \left(2, \frac{(1+\frac{b}{a})e^{2e+2fx}}{1-\frac{b}{a}} \right) dx}{(a-b)^2(a+b)f^2} \\
&= \frac{(c+dx)^3}{3(a-b)^2 d} - \frac{4b^2(c+dx)^3}{3(a^2-b^2)^2 d} - \frac{2b(c+dx)^2 \log \left(1 - \frac{(a+b)e^{2e+2fx}}{a-b} \right)}{(a-b)^2(a+b)f} \\
&\quad - \frac{2bd(c+dx) \text{PolyLog} \left(2, \frac{(a+b)e^{2e+2fx}}{a-b} \right)}{(a-b)^2(a+b)f^2} + \frac{(4b^2) \int \frac{(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2} \\
&\quad - \frac{(8b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} + \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{(a-b+(-a-b)e^{2e+2fx})^2} dx}{a^2-b^2} \\
&\quad + \frac{(bd^2) \text{Subst} \left(\int \frac{\text{PolyLog} \left(2, \frac{(a+b)x}{a-b} \right)}{x} dx, x, e^{2e+2fx} \right)}{(a-b)^2(a+b)f^3}
\end{aligned}$$

$$\begin{aligned}
&= \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{4b^2(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad + \frac{(4b^2) \int \frac{e^{2e+2fx}(c+dx)^2}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)} - \frac{(4b^2d) \int \frac{c+dx}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)(a+b)^2f} \\
&\quad - \frac{(8b^2d) \int (c+dx) \log\left(1 + \frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2d} \\
&\quad - \frac{2b(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{4b^2d(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} \\
&\quad + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{(4b^2d^2) \int \operatorname{PolyLog}\left(2, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2} \\
&\quad - \frac{(4b^2d) \int \frac{e^{2e+2fx}(c+dx)}{a-b+(-a-b)e^{2e+2fx}} dx}{(a-b)^2(a+b)f} + \frac{(4b^2d) \int (c+dx) \log\left(1 + \frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} \\
&\quad + \frac{(c+dx)^3}{3(a-b)^2d} + \frac{2b^2d(c+dx) \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} \\
&\quad - \frac{2b(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} + \frac{2b^2(c+dx)^2 \log\left(1 - \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f} \\
&\quad - \frac{2bd(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{2b^2d(c+dx) \operatorname{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2f^2} \\
&\quad + \frac{bd^2 \operatorname{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{(2b^2d^2) \operatorname{Subst}\left(\int \frac{\operatorname{PolyLog}\left(2, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2f^3} \\
&\quad - \frac{(2b^2d^2) \int \log\left(1 + \frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2} + \frac{(2b^2d^2) \int \operatorname{PolyLog}\left(2, -\frac{(-a-b)e^{2e+2fx}}{a-b}\right) dx}{(a^2-b^2)^2f^2}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2 d} \\
&\quad + \frac{2b^2d(c+dx)\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^2\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^2\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} - \frac{2bd(c+dx)\text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} \\
&\quad + \frac{2b^2d(c+dx)\text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} + \frac{bd^2\text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} \\
&\quad - \frac{2b^2d^2\text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} - \frac{(b^2d^2)\text{Subst}\left(\int \frac{\log\left(1+\frac{(-a-b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^3} \\
&\quad + \frac{(b^2d^2)\text{Subst}\left(\int \frac{\text{PolyLog}\left(2, \frac{(a+b)x}{a-b}\right)}{x} dx, x, e^{2e+2fx}\right)}{(a^2-b^2)^2 f^3} \\
&= -\frac{2b^2(c+dx)^2}{(a^2-b^2)^2 f} + \frac{2b^2(c+dx)^2}{(a-b)(a+b)^2(a-b-(a+b)e^{2e+2fx})f} + \frac{(c+dx)^3}{3(a-b)^2 d} \\
&\quad + \frac{2b^2d(c+dx)\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} - \frac{2b(c+dx)^2\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f} \\
&\quad + \frac{2b^2(c+dx)^2\log\left(1-\frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f} + \frac{b^2d^2\text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3} \\
&\quad - \frac{2bd(c+dx)\text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^2} + \frac{2b^2d(c+dx)\text{PolyLog}\left(2, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^2} \\
&\quad + \frac{bd^2\text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a-b)^2(a+b)f^3} - \frac{b^2d^2\text{PolyLog}\left(3, \frac{(a+b)e^{2e+2fx}}{a-b}\right)}{(a^2-b^2)^2 f^3}
\end{aligned}$$

Mathematica [A] (verified)

Time = 5.06 (sec) , antiderivative size = 519, normalized size of antiderivative = 1.09

$$\begin{aligned}
&\int \frac{(c+dx)^2}{(a+b\coth(e+fx))^2} dx \\
&= \frac{24bcf^2(-bd+acf)x + \frac{24(a-b)bcf^2(-bd+acf)x}{a(-1+e^{2e})+b(1+e^{2e})} + \frac{12(a-b)bdf^2(-bd+2acf)x^2}{a(-1+e^{2e})+b(1+e^{2e})} + \frac{8a(a-b)bd^2f^3x^3}{a(-1+e^{2e})+b(1+e^{2e})} + 12bdf(bd-2acf)x}{a(-1+e^{2e})+b(1+e^{2e})}
\end{aligned}$$

[In] Integrate[(c + d*x)^2/(a + b*Coth[e + f*x])^2, x]

```
[Out] (24*b*c*f^2*(-(b*d) + a*c*f)*x + (24*(a - b)*b*c*f^2*(-(b*d) + a*c*f)*x)/(a*(-1 + E^(2*e)) + b*(1 + E^(2*e))) + (12*(a - b)*b*d*f^2*(-(b*d) + 2*a*c*f)*x^2)/(a*(-1 + E^(2*e)) + b*(1 + E^(2*e))) + (8*a*(a - b)*b*d^2*f^3*x^3)/(a*(-1 + E^(2*e)) + b*(1 + E^(2*e))) + 12*b*d*f*(b*d - 2*a*c*f)*x*Log[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))] - 12*a*b*d^2*f^2*x^2*Log[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))] + 12*b*c*f*(b*d - a*c*f)*Log[a - b - (a + b)*E^(2*(e + f*x))] - 6*b*d*(b*d - 2*a*c*f)*PolyLog[2, (a - b)/((a + b)*E^(2*(e + f*x)))] + 6*a*b*d^2*(2*f*x*PolyLog[2, (a - b)/((a + b)*E^(2*(e + f*x)))] + PolyLog[3, (a - b)/((a + b)*E^(2*(e + f*x)))] - ((a - b)*(a + b)*f^2*((a^2 + b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[f*x] - (a^2 - b^2)*f*x*(3*c^2 + 3*c*d*x + d^2*x^2)*Cosh[2*e + f*x] + 2*b*(-3*b*(c + d*x)^2 + a*f*x*(3*c^2 + 3*c*d*x + d^2*x^2))*Sinh[f*x]))/((b*Cosh[e] + a*Sinh[e])*(b*Cosh[e + f*x] + a*Sinh[e + f*x])))/(6*(a - b)^2*(a + b)^2*f^3)
```

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 1237 vs. $2(464) = 928$.

Time = 0.50 (sec), antiderivative size = 1238, normalized size of antiderivative = 2.61

method	result	size
risch	Expression too large to display	1238

```
[In] int((d*x+c)^2/(a+b*cOTH(f*x+e))^2,x,method=_RETURNVERBOSE)

[Out] 4/3/(a-b)^2*b/(a+b)^2*a*d^2*x^3-8/3/f^3/(a-b)^2*b/(a+b)^2*a*d^2*e^3+2/f^3/(a-b)^2*b^2/(a+b)^2*d^2*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e^4/f^3/(a-b)^2*b^2/(a+b)^2*e*d^2*ln(exp(f*x+e))-2/f^3/(a-b)^2*b^2/(a+b)^2*e*d^2*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)+8/f/(a-b)^2*b/(a+b)^2*c*d*a*e*x-8/f^2/(a-b)^2*b/(a+b)^2*e*c*d*a*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)-4/f^2/(a-b)^2*b/(a+b)^2*c*d*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e-4/f/(a-b)^2*b/(a+b)^2*c*d*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x+4/f/(a-b)^2*b/(a+b)^2*a*c^2*ln(exp(f*x+e))-2/f/(a-b)^2*b/(a+b)^2*a*c^2*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)+1/f^3/(a-b)^2*b/(a+b)^2*a*d^2*polylog(3,(a+b)*exp(2*f*x+2*e)/(a-b))-4/f^2/(a-b)^2*b^2/(a+b)^2*c*d*ln(e*xp(f*x+e))+2/f^2/(a-b)^2*b^2/(a+b)^2*c*d*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)+2/f^2/(a-b)^2*b^2/(a+b)^2*d^2*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x-2/(a-b)/f/(a^2+2*a*b+b^2)*(d^2*x^2+2*c*d*x+c^2)*b^2/(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)+d/(a^2+2*a*b+b^2)*c*x^2+1/(a^2+2*a*b+b^2)*c^2*x-4/f^2/(a-b)^2*b^2/(a+b)^2*d^2*2*e*x-2/f/(a-b)^2*b^2/(a+b)^2*d^2*x^2-2/f^3/(a-b)^2*b^2/(a+b)^2*d^2*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))+4/f^2/(a-b)^2*b/(a+b)^2*c*d*a*e^2+4/(a-b)^2*b/(a+b)^2*c*d*a*x^2-4/f^2/(a-b)^2*b/(a+b)^2*a*d^2*2*e^2*x-2/f^2/(a-b)^2*b/(a+b)^2*c*d*a*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))+4/f^3/(a-b)^2*b/(a+b)^2*e^2*a*d^2*ln(exp(f*x+e))-2/f^3/(a-b)^2*b/(a+b)^2*e^2*a*d^2*ln(exp(2*f*x+2*e)*a+b*exp(2*f*x+2*e)-a+b)-2/f/(a-b)^2*b/(a+b)^2*a*d^2*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x^2+2/f^3/(a-b)
```

$$\begin{aligned} & -2 \cdot b^2 \cdot (a+b)^{-2} \cdot a \cdot d^{-2} \cdot \ln(1-(a+b) \cdot \exp(2 \cdot f \cdot x + 2 \cdot e) / (a-b)) \cdot e^{-2-2/f^2} / (a-b)^{-2} \cdot b / (a+b)^{-2} \cdot a \cdot d^{-2} \cdot \text{polylog}(2, (a+b) \cdot \exp(2 \cdot f \cdot x + 2 \cdot e) / (a-b)) \cdot x + 1/3 \cdot d^{-2} / (a^2 + 2 \cdot a \cdot b + b^2) \cdot x^3 + 1/3 \cdot d / (a^2 + 2 \cdot a \cdot b + b^2) \cdot c \cdot g^3 \end{aligned}$$

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 3702 vs. $2(460) = 920$.

Time = 0.32 (sec) , antiderivative size = 3702, normalized size of antiderivative = 7.79

$$\int \frac{(c+dx)^2}{(a+b\coth(e+fx))^2} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)^2/(a+b*cOTH(f*x+e))^2,x, algorithm="fricas")
[Out] -1/3*((a^3 + a^2*b - a*b^2 - b^3)*d^2*f^3*x^3 + 3*(a^3 + a^2*b - a*b^2 - b^3)*c*d*f^3*x^2 + 3*(a^3 + a^2*b - a*b^2 - b^3)*c^2*f^3*x + 4*(a^2*b - a*b^2 - b^3)*d^2*e^3 + 6*(a*b^2 - b^3)*d^2*e^2 + 6*(2*(a^2*b - a*b^2)*c^2*e + (a*b^2 - b^3)*c^2)*f^2 - ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*cosh(f*x + e)^2 - 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*cosh(f*x + e)*sinh(f*x + e) - ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d^2*f^3*x^3 + 4*(a^2*b + a*b^2)*d^2*e^3 + 12*(a^2*b + a*b^2)*c^2*e*f^2 + 6*(a*b^2 + b^3)*d^2*e^2 + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*d*f^3 - 2*(a*b^2 + b^3)*d^2*f^2)*x^2 - 12*((a^2*b + a*b^2)*c*d*e^2 + (a*b^2 + b^3)*c*d*e)*f + 3*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c^2*f^3 - 4*(a*b^2 + b^3)*c*d*f^2)*x)*cosh(f*x + e)*sinh(f*x + e)^2 - 12*((a^2*b - a*b^2)*c*d*e^2 + (a*b^2 - b^3)*c*d*e)*f - 6*(2*(a^2*b - a*b^2)*d^2*f*x + 2*(a^2*b - a*b^2)*c*d*f - (a*b^2 - b^3)*d^2 - (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)^2 - 2*(2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)*sinh(f*x + e) - (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e)^2)*dilog(sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*(2*(a^2*b - a*b^2)*d^2*f*x + 2*(a^2*b - a*b^2)*c*d*f - (a*b^2 - b^3)*d^2 - (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)^2 - 2*(2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*cosh(f*x + e)*sinh(f*x + e) - (2*(a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e)^2)*dilog(-sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e))) - 6*((a^2*b - a*b^2)*d^2*e^2 + (a^2*b - a*b^2)*c^2*f^2 + (a*b^2 - b^3)*d^2*f*x - ((a^2*b + a*b^2)*d^2*f*x + 2*(a^2*b + a*b^2)*c*d*f - (a*b^2 + b^3)*d^2)*sinh(f*x + e)^2)*dilog(-sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)))
```


$$+ (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^3*sinh(f*x + e)^2 - (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*f^3)$$

Sympy [F(-2)]

Exception generated.

$$\int \frac{(c+dx)^2}{(a+b\coth(e+fx))^2} dx = \text{Exception raised: TypeError}$$

[In] `integrate((d*x+c)**2/(a+b*cOTH(f*x+e))**2,x)`

[Out] Exception raised: TypeError >> Invalid NaN comparison

Maxima [A] (verification not implemented)

none

Time = 0.51 (sec) , antiderivative size = 751, normalized size of antiderivative = 1.58

$$\begin{aligned} & \int \frac{(c+dx)^2}{(a+b\coth(e+fx))^2} dx = -\frac{4b^2cdfx}{a^4f^2 - 2a^2b^2f^2 + b^4f^2} \\ & - \frac{\left(2f^2x^2 \log\left(-\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} + 1\right) + 2fx\text{Li}_2\left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b}\right) - \text{Li}_3\left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b}\right)\right)abd^2}{a^4f^3 - 2a^2b^2f^3 + b^4f^3} \\ & + \frac{2b^2cd \log((ae^{(2e)}+be^{(2e)})e^{(2fx)} - a+b)}{a^4f^2 - 2a^2b^2f^2 + b^4f^2} \\ & - c^2 \left(\frac{2ab \log(-(a-b)e^{(-2fx-2e)} + a+b)}{(a^4 - 2a^2b^2 + b^4)f} + \frac{2b^2}{(a^4 - 2a^2b^2 + b^4 - (a^4 - 2a^3b + 2ab^3 - b^4)e^{(-2fx-2e)})f} \right. \\ & - \frac{(2abcd - b^2d^2) \left(2fx \log\left(-\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b} + 1\right) + \text{Li}_2\left(\frac{(ae^{(2e)}+be^{(2e)})e^{(2fx)}}{a-b}\right) \right)}{a^4f^3 - 2a^2b^2f^3 + b^4f^3} \\ & + \frac{2(2abd^2f^3x^3 + 3(2abcd - b^2d^2)f^2x^2)}{3(a^4f^3 - 2a^2b^2f^3 + b^4f^3)} \\ & + \frac{12b^2cdx + (a^2d^2f - 2abd^2f + b^2d^2f)x^3 + 3(a^2cdf - 2abcd + (cdf + 2d^2)b^2)x^2 - ((a^2d^2fe^{(2e)} - b^2d^2f)x^1)}{3(a^4f - 2a^2b^2f + b^4f - (a^4fe^{(2e)} + 2a^3bfe^{(2e)} - 2ab^3fe^{(2e)} - b^4fe^{(2e)}))} \end{aligned}$$

```
[In] integrate((d*x+c)^2/(a+b*cot(f*x+e))^2,x, algorithm="maxima")
```

```
[Out] -4*b^2*c*d*f*x/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - (2*f^2*x^2*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x))/(a - b) + 1) + 2*f*x*dilog((a*e^(2*e) + b*e^(2*e))*e^(2*f*x))/(a - b)) - polylog(3, (a*e^(2*e) + b*e^(2*e))*e^(2*f*x))/(a - b))*a*b*d^2/(a^4*f^3 - 2*a^2*b^2*f^3 + b^4*f^3) + 2*b^2*c*d*log((a*e^(2*e) + b*e^(2*e))*e^(2*f*x)) - a + b)/(a^4*f^2 - 2*a^2*b^2*f^2 + b^4*f^2) - c^2*
```

$$\begin{aligned}
& (2*a*b*log(-(a - b)*e^{(-2*f*x - 2*e)} + a + b)/((a^4 - 2*a^2*b^2 + b^4)*f) + \\
& 2*b^2/((a^4 - 2*a^2*b^2 + b^4 - (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^{(-2*f*x} \\
& - 2*e))*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f)) - (2*a*b*c*d*f - b^2*d^2)*(\\
& 2*f*x*log(-(a*e^(2*e) + b*e^(2*e))*e^(2*f*x)/(a - b) + 1) + \text{dilog}((a*e^(2*e} \\
&) + b*e^(2*e))*e^(2*f*x)/(a - b)))/(a^4*f^3 - 2*a^2*b^2*f^3 + b^4*f^3) + 2/ \\
& 3*(2*a*b*d^2*f^3*x^3 + 3*(2*a*b*c*d*f - b^2*d^2)*f^2*x^2)/(a^4*f^3 - 2*a^2*b^2* \\
& b^2*f^3 + b^4*f^3) + 1/3*(12*b^2*c*d*x + (a^2*d^2*f - 2*a*b*d^2*f + b^2*d^2* \\
& f)*x^3 + 3*(a^2*c*d*f - 2*a*b*c*d*f + (c*d*f + 2*d^2)*b^2)*x^2 - ((a^2*d^2* \\
& f)*e^(2*e) - b^2*d^2*f*e^(2*e))*x^3 + 3*(a^2*c*d*f*e^(2*e) - b^2*c*d*f*e^(2* \\
& e))*x^2)*e^(2*f*x))/(a^4*f - 2*a^2*b^2*f + b^4*f - (a^4*f*e^(2*e) + 2*a^3* \\
& b*f*e^(2*e) - 2*a*b^3*f*e^(2*e) - b^4*f*e^(2*e))*e^(2*f*x))
\end{aligned}$$

Giac [F]

$$\int \frac{(c + dx)^2}{(a + b \coth(e + fx))^2} dx = \int \frac{(dx + c)^2}{(b \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)^2/(a+b*coth(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)^2/(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{(c + dx)^2}{(a + b \coth(e + fx))^2} dx = \int \frac{(c + d x)^2}{(a + b \coth(e + f x))^2} dx$$

[In] `int((c + d*x)^2/(a + b*coth(e + f*x))^2,x)`
[Out] `int((c + d*x)^2/(a + b*coth(e + f*x))^2, x)`

3.59 $\int \frac{c+dx}{(a+b \coth(e+fx))^2} dx$

Optimal result	402
Rubi [A] (verified)	402
Mathematica [A] (verified)	405
Maple [B] (verified)	405
Fricas [B] (verification not implemented)	406
Sympy [F(-2)]	407
Maxima [F]	407
Giac [F]	408
Mupad [F(-1)]	408

Optimal result

Integrand size = 18, antiderivative size = 196

$$\begin{aligned} \int \frac{c+dx}{(a+b \coth(e+fx))^2} dx = & -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adf x)^2}{4a(a-b)(a+b)^2 df^2} \\ & + \frac{b(c+dx)}{(a^2-b^2)f(a+b \coth(e+fx))} \\ & + \frac{b(bd-2acf-2adf x) \log\left(1-\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2 f^2} \\ & + \frac{abd \operatorname{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2 f^2} \end{aligned}$$

[Out] $-1/2*(d*x+c)^2/(a^2-b^2)/d+1/4*(-2*a*d*f*x-2*a*c*f+b*d)^2/a/(a-b)/(a+b)^2/d$
 $/f^2+b*(d*x+c)/(a^2-b^2)/f/(a+b*\coth(f*x+e))+b*(-2*a*d*f*x-2*a*c*f+b*d)*\ln($
 $1+(-a+b)/(a+b)/\exp(2*f*x+2*e))/(a^2-b^2)^2/f^2+a*b*d*\operatorname{polylog}(2,(a-b)/(a+b)/$
 $\exp(2*f*x+2*e))/(a^2-b^2)^2/f^2$

Rubi [A] (verified)

Time = 0.22 (sec), antiderivative size = 196, normalized size of antiderivative = 1.00,
number of steps used = 5, number of rules used = 5, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used

$$= \{3814, 3812, 2221, 2317, 2438\}$$

$$\begin{aligned} \int \frac{c+dx}{(a+b \coth(e+fx))^2} dx &= \frac{b(-2acf - 2adfx + bd) \log \left(1 - \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f^2 (a^2 - b^2)^2} \\ &\quad + \frac{b(c+dx)}{f(a^2 - b^2)(a+b \coth(e+fx))} - \frac{(c+dx)^2}{2d(a^2 - b^2)} \\ &\quad + \frac{abd \operatorname{PolyLog} \left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{f^2 (a^2 - b^2)^2} + \frac{(-2acf - 2adfx + bd)^2}{4ad^2 f^2 (a-b)(a+b)^2} \end{aligned}$$

[In] $\operatorname{Int}[(c + d*x)/(a + b*\operatorname{Coth}[e + f*x])^2, x]$

[Out] $-1/2*(c + d*x)^2/((a^2 - b^2)*d) + (b*d - 2*a*c*f - 2*a*d*f*x)^2/(4*a*(a - b)*(a + b)^2*d*f^2) + (b*(c + d*x))/((a^2 - b^2)*f*(a + b*\operatorname{Coth}[e + f*x])) + (b*(b*d - 2*a*c*f - 2*a*d*f*x)*\operatorname{Log}[1 - (a - b)/((a + b)*E^(2*(e + f*x)))])/((a^2 - b^2)^2*f^2) + (a*b*d*\operatorname{PolyLog}[2, (a - b)/((a + b)*E^(2*(e + f*x)))])/((a^2 - b^2)^2*f^2)$

Rule 2221

$\operatorname{Int}[(((F_*)^{(g_*)}((e_*) + (f_*)*(x_*)))^{(n_*)}((c_*) + (d_*)*(x_*)^{(m_*)})/((a_*) + (b_*)*((F_*)^{(g_*)}((e_*) + (f_*)*(x_*)))^{(n_*)}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[((c + d*x)^m/(b*f*g*n*\operatorname{Log}[F]))*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x] - \operatorname{Dist}[d*(m/(b*f*g*n*\operatorname{Log}[F])), \operatorname{Int}[(c + d*x)^{m-1}*\operatorname{Log}[1 + b*((F^(g*(e + f*x)))^n/a)], x], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, f, g, n\}, x] \&& \operatorname{IGtQ}[m, 0]$

Rule 2317

$\operatorname{Int}[\operatorname{Log}[(a_*) + (b_*)*((F_*)^{(e_*)}((c_*) + (d_*)*(x_*)))^{(n_*)}], x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[1/(d*e*n*\operatorname{Log}[F]), \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[a + b*x]/x, x], x, (F^(e*(c + d*x)))^n], x] /; \operatorname{FreeQ}[\{F, a, b, c, d, e, n\}, x] \&& \operatorname{GtQ}[a, 0]$

Rule 2438

$\operatorname{Int}[\operatorname{Log}[(c_*)(d_*) + (e_*)(x_*)^{(n_*)}]/(x_), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[-\operatorname{PolyLog}[2, (-c)*e*x^n]/n, x] /; \operatorname{FreeQ}[\{c, d, e, n\}, x] \&& \operatorname{EqQ}[c*d, 1]$

Rule 3812

$\operatorname{Int}[(c_*) + (d_*)(x_*)^{(m_*)}/((a_*) + (b_*)*\operatorname{tan}[(e_*) + \operatorname{Pi}*(k_*) + (f_*)(x_*)]), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(c + d*x)^(m+1)/(d*(m+1)*(a + I*b)), x] + \operatorname{Dist}[2*I*b, \operatorname{Int}[(c + d*x)^m*E^(2*I*k*Pi)*(E^{\operatorname{Simp}[2*I*(e + f*x), x]}/((a + I*b)^2 + (a^2 + b^2)*E^(2*I*k*Pi)*E^{\operatorname{Simp}[2*I*(e + f*x), x]}), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \&& \operatorname{IntegerQ}[4*k] \&& \operatorname{NeQ}[a^2 + b^2, 0] \&& \operatorname{IGtQ}[m, 0]$

Rule 3814

```

Int[((c_.) + (d_.)*(x_))/((a_) + (b_.)*tan[(e_.) + (f_.)*(x_])])^2, x_Symbol
] :> Simp[-(c + d*x)^2/(2*d*(a^2 + b^2)), x] + (Dist[1/(f*(a^2 + b^2)), Int
[(b*d + 2*a*c*f + 2*a*d*f*x)/(a + b*Tan[e + f*x]), x], x] - Simp[b*((c + d*
x)/(f*(a^2 + b^2)*(a + b*Tan[e + f*x]))), x]) /; FreeQ[{a, b, c, d, e, f},
x] && NeQ[a^2 + b^2, 0]

```

Rubi steps

$$\begin{aligned}
\text{integral} &= -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{b(c+dx)}{(a^2-b^2)f(a+b\coth(e+fx))} - \frac{i \int \frac{-ibd+2iacf+2iadfx}{a+b\coth(e+fx)} dx}{(a^2-b^2)f} \\
&= -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adfx)^2}{4a(a-b)(a+b)^2df^2} \\
&\quad + \frac{b(c+dx)}{(a^2-b^2)f(a+b\coth(e+fx))} + \frac{(2ib) \int \frac{e^{-2(e+fx)}(-ibd+2acf+2adfx)}{(a+b)^2+(-a^2+b^2)e^{-2(e+fx)}} dx}{(a^2-b^2)f} \\
&= -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adfx)^2}{4a(a-b)(a+b)^2df^2} + \frac{b(c+dx)}{(a^2-b^2)f(a+b\coth(e+fx))} \\
&\quad + \frac{b(bd-2acf-2adfx) \log\left(1-\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2f^2} \\
&\quad + \frac{(2abd) \int \log\left(1+\frac{(-a^2+b^2)e^{-2(e+fx)}}{(a+b)^2}\right) dx}{(a^2-b^2)^2f} \\
&= -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adfx)^2}{4a(a-b)(a+b)^2df^2} + \frac{b(c+dx)}{(a^2-b^2)f(a+b\coth(e+fx))} \\
&\quad + \frac{b(bd-2acf-2adfx) \log\left(1-\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2f^2} \\
&\quad - \frac{(abd)\text{Subst}\left(\int \frac{\log\left(1+\frac{(-a^2+b^2)x}{(a+b)^2}\right)}{x} dx, x, e^{-2(e+fx)}\right)}{(a^2-b^2)^2f^2} \\
&= -\frac{(c+dx)^2}{2(a^2-b^2)d} + \frac{(bd-2acf-2adfx)^2}{4a(a-b)(a+b)^2df^2} + \frac{b(c+dx)}{(a^2-b^2)f(a+b\coth(e+fx))} \\
&\quad + \frac{b(bd-2acf-2adfx) \log\left(1-\frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2f^2} + \frac{abd \text{PolyLog}\left(2, \frac{(a-b)e^{-2(e+fx)}}{a+b}\right)}{(a^2-b^2)^2f^2}
\end{aligned}$$

Mathematica [A] (verified)

Time = 3.87 (sec) , antiderivative size = 244, normalized size of antiderivative = 1.24

$$\int \frac{c + dx}{(a + b \coth(e + fx))^2} dx \\ = \text{csch}^2(e + fx)(b \cosh(e + fx) + a \sinh(e + fx)) \left(4(a - b)bf(c + dx) \sinh(e + fx) + 2(a - b)(e + fx)(-2a^2b^2\cosh^2(e + fx) + 2ab^2\cosh(e + fx)\sinh(e + fx) + a^2\sinh^2(e + fx)) \right)$$

[In] `Integrate[(c + d*x)/(a + b*Coth[e + f*x])^2, x]`

[Out] `(Csch[e + f*x]^2*(b*Cosh[e + f*x] + a*Sinh[e + f*x])*(4*(a - b)*b*f*(c + d*x)*Sinh[e + f*x] + 2*(a - b)*(e + f*x)*(-2*c*f + d*(e - f*x))*(b*Cosh[e + f*x] + a*Sinh[e + f*x]) + (((-(b*d) + 2*a*f*(c + d*x))*(a - b)*(-(b*d) + 2*a*f*(c + d*x)) - 4*a*b*d*Log[1 + (-a + b)/((a + b)*E^(2*(e + f*x)))]) + 4*a^2*b*d^2*PolyLog[2, (a - b)/((a + b)*E^(2*(e + f*x)))])*(b*Cosh[e + f*x] + a*Sinh[e + f*x]))/(a*(a + b)*d))/((4*(a - b)^2*(a + b)*f^2*(a + b)*Coth[e + f*x])^2)`

Maple [B] (verified)

Leaf count of result is larger than twice the leaf count of optimal. 523 vs. 2(195) = 390.

Time = 0.44 (sec) , antiderivative size = 524, normalized size of antiderivative = 2.67

method	result
risch	$\frac{dx^2}{2a^2+4ab+2b^2} + \frac{cx}{a^2+2ab+b^2} - \frac{2(dx+c)b^2}{(a-b)f(a^2+2ab+b^2)(e^{2fx+2e}a+b e^{2fx+2e}-a+b)} - \frac{2b^2d \ln(e^{fx+e})}{f^2(a-b)^2(a+b)^2} + \frac{b^2d \ln(e^{2fx+2e}a+b e^{2fx+2e})}{f^2(a-b)^2(a+b)^2}$

[In] `int((d*x+c)/(a+b*coth(f*x+e))^2,x,method=_RETURNVERBOSE)`

[Out] `1/2/(a^2+2*a*b+b^2)*d*x^2+1/(a^2+2*a*b+b^2)*c*x-2/(a-b)/f/(a^2+2*a*b+b^2)*(d*x+c)*b^2/(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)-2/f^2/(a-b)^2*b^2/(a+b)^2*d*ln(\exp(f*x+e))+1/f^2/(a-b)^2*b^2/(a+b)^2*d*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)+4/f/(a-b)^2*b/(a+b)^2*a*c*ln(\exp(f*x+e))-2/f/(a-b)^2*b/(a+b)^2*a*c*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)-4/f^2/(a-b)^2*b/(a+b)^2*e*d*a*ln(\exp(f*x+e))+2/f^2/(a-b)^2*b/(a+b)^2*e*d*a*ln(\exp(2*f*x+2*e)*a+b*\exp(2*f*x+2*e)-a+b)+2/(a-b)^2*b/(a+b)^2*d*a*x^2-2/f/(a-b)^2*b/(a+b)^2*d*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*x+4/f/(a-b)^2*b/(a+b)^2*d*a*e*x-2/f^2/(a-b)^2*b/(a+b)^2*d*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e+2/f^2/(a-b)^2*b/(a+b)^2*d*a*ln(1-(a+b)*exp(2*f*x+2*e)/(a-b))*e^2-1/f^2/(a-b)^2*b/(a+b)^2*d*a*polylog(2,(a+b)*exp(2*f*x+2*e)/(a-b))`

Fricas [B] (verification not implemented)

Leaf count of result is larger than twice the leaf count of optimal. 1797 vs. $2(194) = 388$.

Time = 0.31 (sec) , antiderivative size = 1797, normalized size of antiderivative = 9.17

$$\int \frac{c+dx}{(a+b\coth(e+fx))^2} dx = \text{Too large to display}$$

```
[In] integrate((d*x+c)/(a+b*coth(f*x+e))^2,x, algorithm="fricas")
```

```
[Out] -1/2*((a^3 + a^2*b - a*b^2 - b^3)*d*f^2*x^2 + 2*(a^3 + a^2*b - a*b^2 - b^3)
*c*f^2*x - 4*(a^2*b - a*b^2)*d*e^2 - 4*(a*b^2 - b^3)*d*e - ((a^3 + 3*a^2*b
+ 3*a*b^2 + b^3)*d*f^2*x^2 - 4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*
e*f - 4*(a*b^2 + b^3)*d*e + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a
*b^2 + b^3)*d*f)*x)*cosh(f*x + e)^2 - 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*
f^2*x^2 - 4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*e*f - 4*(a*b^2 + b^
3)*d*e + 2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 + b^3)*d*f)*x)
*cosh(f*x + e)*sinh(f*x + e) - ((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*d*f^2*x^2 -
4*(a^2*b + a*b^2)*d*e^2 + 8*(a^2*b + a*b^2)*c*e*f - 4*(a*b^2 + b^3)*d*e +
2*((a^3 + 3*a^2*b + 3*a*b^2 + b^3)*c*f^2 - 2*(a*b^2 + b^3)*d*f)*x)*sinh(f*x
+ e)^2 + 4*(2*(a^2*b - a*b^2)*c*e + (a*b^2 - b^3)*c)*f + 4*((a^2*b + a*b^2
)*d*cosh(f*x + e)^2 + 2*(a^2*b + a*b^2)*d*cosh(f*x + e)*sinh(f*x + e) + (a^
2*b + a*b^2)*d*sinh(f*x + e)^2 - (a^2*b - a*b^2)*d)*dilog(sqrt((a + b)/(a -
b))*(cosh(f*x + e) + sinh(f*x + e))) + 4*((a^2*b + a*b^2)*d*cosh(f*x + e)^
2 + 2*(a^2*b + a*b^2)*d*cosh(f*x + e)*sinh(f*x + e) + (a^2*b + a*b^2)*d*sin
h(f*x + e)^2 - (a^2*b - a*b^2)*d)*dilog(-sqrt((a + b)/(a - b))*(cosh(f*x +
e) + sinh(f*x + e))) + 2*(2*(a^2*b - a*b^2)*d*e - 2*(a^2*b - a*b^2)*c*f - (
2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x +
e)^2 - 2*(2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)
*cosh(f*x + e)*sinh(f*x + e) - (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*
f + (a*b^2 + b^3)*d)*sinh(f*x + e)^2 + (a*b^2 - b^3)*d)*log(2*(a + b)*cosh
(f*x + e) + 2*(a + b)*sinh(f*x + e) + 2*(a - b)*sqrt((a + b)/(a - b))) + 2*
(2*(a^2*b - a*b^2)*d*e - 2*(a^2*b - a*b^2)*c*f - (2*(a^2*b + a*b^2)*d*e - 2
*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)^2 - 2*(2*(a^2*b + a*b
^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*cosh(f*x + e)*sinh(f*x +
e) - (2*(a^2*b + a*b^2)*d*e - 2*(a^2*b + a*b^2)*c*f + (a*b^2 + b^3)*d)*sin
h(f*x + e)^2 + (a*b^2 - b^3)*d)*log(2*(a + b)*cosh(f*x + e) + 2*(a + b)*sin
h(f*x + e) - 2*(a - b)*sqrt((a + b)/(a - b))) - 4*((a^2*b - a*b^2)*d*f*x +
(a^2*b - a*b^2)*d*e - ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*
x + e)^2 - 2*((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*cosh(f*x + e)*si
nh(f*x + e) - ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*e)*sinh(f*x + e)^2
)*log(sqrt((a + b)/(a - b))*(cosh(f*x + e) + sinh(f*x + e)) + 1) - 4*((a^2*
b - a*b^2)*d*f*x + (a^2*b - a*b^2)*d*e - ((a^2*b + a*b^2)*d*f*x + (a^2*b +
a*b^2)*d*e)*cosh(f*x + e)^2 - 2*((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*
e)*cosh(f*x + e)*sinh(f*x + e) - ((a^2*b + a*b^2)*d*f*x + (a^2*b + a*b^2)*d*
```

$$\begin{aligned} & *e) * \sinh(f*x + e)^2 * \log(-\sqrt((a + b)/(a - b)) * (\cosh(f*x + e) + \sinh(f*x + e)) + 1)) / ((a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*cosh(f*x + e)^2 + 2*(a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*cosh(f*x + e)*\sinh(f*x + e) + (a^5 + a^4*b - 2*a^3*b^2 - 2*a^2*b^3 + a*b^4 + b^5)*f^2*\sinh(f*x + e)^2 - (a^5 - a^4*b - 2*a^3*b^2 + 2*a^2*b^3 + a*b^4 - b^5)*f^2) \end{aligned}$$

Sympy [F(-2)]

Exception generated.

$$\int \frac{c + dx}{(a + b \coth(e + fx))^2} dx = \text{Exception raised: TypeError}$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e))**2,x)`

[Out] Exception raised: TypeError >> Invalid NaN comparison

Maxima [F]

$$\int \frac{c + dx}{(a + b \coth(e + fx))^2} dx = \int \frac{dx + c}{(b \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e))**2,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -1/2*(8*a*b*f*integrate(x/(a^4*f*e^(2*f*x + 2*e) + 2*a^3*b*f*e^(2*f*x + 2*e) - 2*a*b^3*f*e^(2*f*x + 2*e) - b^4*f*e^(2*f*x + 2*e) - a^4*f + 2*a^2*b^2*f - b^4*f), x) + 2*b^2*(2*(f*x + e)/((a^4 - 2*a^2*b^2 + b^4)*f^2) - log((a + b)*e^(2*f*x + 2*e) - a + b)/((a^4 - 2*a^2*b^2 + b^4)*f^2)) + ((a^2*f*e^(2*e) - b^2*f*e^(2*e))*x^2*e^(2*f*x) - 4*b^2*x - (a^2*f - 2*a*b*f + b^2*f)*x^2)/((a^4*f - 2*a^2*b^2*f + b^4*f - (a^4*f*e^(2*e) + 2*a^3*b*f*e^(2*e) - 2*a*b^3*f*e^(2*e) - b^4*f*e^(2*e))*e^(2*f*x)))*d - c*(2*a*b*log(-(a - b)*e^(-2*f*x - 2*e) + a + b)/((a^4 - 2*a^2*b^2 + b^4)*f) + 2*b^2/((a^4 - 2*a^2*b^2 + b^4 - (a^4 - 2*a^3*b + 2*a*b^3 - b^4)*e^(-2*f*x - 2*e))*f) - (f*x + e)/((a^2 + 2*a*b + b^2)*f)) \end{aligned}$$

Giac [F]

$$\int \frac{c + dx}{(a + b \coth(e + fx))^2} dx = \int \frac{dx + c}{(b \coth(fx + e) + a)^2} dx$$

[In] `integrate((d*x+c)/(a+b*coth(f*x+e))^2,x, algorithm="giac")`
[Out] `integrate((d*x + c)/(b*coth(f*x + e) + a)^2, x)`

Mupad [F(-1)]

Timed out.

$$\int \frac{c + dx}{(a + b \coth(e + fx))^2} dx = \int \frac{c + dx}{(a + b \coth(e + fx))^2} dx$$

[In] `int((c + d*x)/(a + b*coth(e + f*x))^2,x)`
[Out] `int((c + d*x)/(a + b*coth(e + f*x))^2, x)`

3.60 $\int \frac{1}{(c+dx)(a+b\coth(e+fx))^2} dx$

Optimal result	409
Rubi [N/A]	409
Mathematica [N/A]	410
Maple [N/A] (verified)	410
Fricas [N/A]	410
Sympy [N/A]	411
Maxima [N/A]	411
Giac [N/A]	411
Mupad [N/A]	412

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)(a+b\coth(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)(a+b\coth(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)/(a+b*coth(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)(a+b\coth(e+fx))^2} dx = \int \frac{1}{(c+dx)(a+b\coth(e+fx))^2} dx$$

[In] Int[1/((c + d*x)*(a + b*Coth[e + f*x])^2),x]

[Out] Defer[Int][1/((c + d*x)*(a + b*Coth[e + f*x])^2), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)(a+b\coth(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 41.45 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx$$

[In] `Integrate[1/((c + d*x)*(a + b*Coth[e + f*x])^2),x]`

[Out] `Integrate[1/((c + d*x)*(a + b*Coth[e + f*x])^2), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)(a + b \coth(fx + e))^2} dx$$

[In] `int(1/(d*x+c)/(a+b*coth(f*x+e))^2,x)`

[Out] `int(1/(d*x+c)/(a+b*coth(f*x+e))^2,x)`

Fricas [N/A]

Not integrable

Time = 0.26 (sec) , antiderivative size = 55, normalized size of antiderivative = 2.75

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e))^2,x, algorithm="fricas")`

[Out] `integral(1/(a^2*d*x + a^2*c + (b^2*d*x + b^2*c)*coth(f*x + e)^2 + 2*(a*b*d*x + a*b*c)*coth(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 1.83 (sec) , antiderivative size = 19, normalized size of antiderivative = 0.95

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(a + b \coth(e + fx))^2 (c + dx)} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e))**2,x)`

[Out] `Integral(1/((a + b*coth(e + f*x))**2*(c + d*x)), x)`

Maxima [N/A]

Not integrable

Time = 1.10 (sec) , antiderivative size = 472, normalized size of antiderivative = 23.60

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e))**2,x, algorithm="maxima")`

[Out] `2*b^2/(a^4*c*f - 2*a^2*b^2*c*f + b^4*c*f + (a^4*d*f - 2*a^2*b^2*d*f + b^4*d*f)*x - (a^4*c*f*e^(2*e) + 2*a^3*b*c*f*e^(2*e) - 2*a*b^3*c*f*e^(2*e) - b^4*c*f*e^(2*e) + (a^4*d*f*e^(2*e) + 2*a^3*b*d*f*e^(2*e) - 2*a*b^3*d*f*e^(2*e) - b^4*d*f*e^(2*e))*x)*e^(2*f*x)) + log(d*x + c)/(a^2*d + 2*a*b*d + b^2*d) - integrate(-2*(2*a*b*d*f*x + 2*a*b*c*f + b^2*d)/(a^4*c^2*f - 2*a^2*b^2*c^2*f + b^4*c^2*f - 2*a^2*b^2*d^2*f*x + 2*a*b*c*f + b^2*d)/(a^4*c^2*f - 2*a^2*b^2*c^2*f - 2*a^2*b^2*d^2*f*x + b^4*c^2*f*x^2 + 2*(a^4*c*d*f - 2*a^2*b^2*c*d*f + b^4*c*d*f)*x - (a^4*c^2*f*e^(2*e) + 2*a^3*b*c^2*f*e^(2*e) - 2*a*b^3*c^2*f*e^(2*e) - b^4*c^2*f*e^(2*e) + (a^4*d^2*f*e^(2*e) + 2*a^3*b*d^2*f*e^(2*e) - 2*a*b^3*d^2*f*e^(2*e) - b^4*d^2*f*e^(2*e))*x^2 + 2*(a^4*c*d*f*e^(2*e) + 2*a^3*b*c*d*f*e^(2*e) - 2*a*b^3*c*d*f*e^(2*e) - b^4*c*d*f*e^(2*e))*x)*e^(2*f*x)), x)`

Giac [N/A]

Not integrable

Time = 0.31 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)(b \coth(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)/(a+b*coth(f*x+e))**2,x, algorithm="giac")`

[Out] `integrate(1/((d*x + c)*(b*coth(f*x + e) + a)^2), x)`

Mupad [N/A]

Not integrable

Time = 2.27 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)(a + b \coth(e + fx))^2} dx = \int \frac{1}{(a + b \coth(e + fx))^2 (c + dx)} dx$$

[In] `int(1/((a + b*coth(e + f*x))^2*(c + d*x)),x)`

[Out] `int(1/((a + b*coth(e + f*x))^2*(c + d*x)), x)`

3.61 $\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))^2} dx$

Optimal result	413
Rubi [N/A]	413
Mathematica [N/A]	414
Maple [N/A] (verified)	414
Fricas [N/A]	414
Sympy [N/A]	415
Maxima [N/A]	415
Giac [N/A]	416
Mupad [N/A]	416

Optimal result

Integrand size = 20, antiderivative size = 20

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))^2} dx = \text{Int}\left(\frac{1}{(c+dx)^2(a+b\coth(e+fx))^2}, x\right)$$

[Out] Unintegrable(1/(d*x+c)^2/(a+b*cOTH(f*x+e))^2,x)

Rubi [N/A]

Not integrable

Time = 0.04 (sec), antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 0, number of rules used = 0, $\frac{\text{number of rules}}{\text{integrand size}} = 0.000$, Rules used = {}

$$\int \frac{1}{(c+dx)^2(a+b\coth(e+fx))^2} dx = \int \frac{1}{(c+dx)^2(a+b\coth(e+fx))^2} dx$$

[In] Int[1/((c + d*x)^2*(a + b*COTH[e + f*x])^2),x]

[Out] Defer[Int][1/((c + d*x)^2*(a + b*COTH[e + f*x])^2), x]

Rubi steps

$$\text{integral} = \int \frac{1}{(c+dx)^2(a+b\coth(e+fx))^2} dx$$

Mathematica [N/A]

Not integrable

Time = 39.17 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx$$

[In] `Integrate[1/((c + d*x)^2*(a + b*Coth[e + f*x])^2), x]`

[Out] `Integrate[1/((c + d*x)^2*(a + b*Coth[e + f*x])^2), x]`

Maple [N/A] (verified)

Not integrable

Time = 0.07 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(dx + c)^2 (a + b \coth(fx + e))^2} dx$$

[In] `int(1/(d*x+c)^2/(a+b*coth(f*x+e))^2, x)`

[Out] `int(1/(d*x+c)^2/(a+b*coth(f*x+e))^2, x)`

Fricas [N/A]

Not integrable

Time = 0.27 (sec) , antiderivative size = 96, normalized size of antiderivative = 4.80

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \coth(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*coth(f*x+e))^2, x, algorithm="fricas")`

[Out] `integral(1/(a^2*d^2*x^2 + 2*a^2*c*d*x + a^2*c^2 + (b^2*d^2*x^2 + 2*b^2*c*d*x + b^2*c^2)*coth(f*x + e)^2 + 2*(a*b*d^2*x^2 + 2*a*b*c*d*x + a*b*c^2)*coth(f*x + e)), x)`

Sympy [N/A]

Not integrable

Time = 3.32 (sec) , antiderivative size = 20, normalized size of antiderivative = 1.00

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(a + b \coth(e + fx))^2(c + dx)^2} dx$$

[In] `integrate(1/(d*x+c)**2/(a+b*coth(f*x+e))**2,x)`

[Out] `Integral(1/((a + b*coth(e + f*x))**2*(c + d*x)**2), x)`

Maxima [N/A]

Not integrable

Time = 2.16 (sec) , antiderivative size = 789, normalized size of antiderivative = 39.45

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \coth(fx + e) + a)^2} dx$$

[In] `integrate(1/(d*x+c)^2/(a+b*coth(f*x+e))^2,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & - (a^2*c*f - 2*a*b*c*f + (c*f - 2*d)*b^2 + (a^2*d*f - 2*a*b*d*f + b^2*d*f)*x \\ & - (a^2*c*f*e^{(2*e)} - b^2*c*f*e^{(2*e)} + (a^2*d*f*e^{(2*e)} - b^2*d*f*e^{(2*e)})*x)*e^{(2*f*x)})/(a^4*c^2*d*f - 2*a^2*b^2*c^2*d*f + b^4*c^2*d*f + (a^4*d^3*f - 2*a^2*b^2*d^3*f + b^4*d^3*f)*x^2 + 2*(a^4*c*d^2*f - 2*a^2*b^2*c*d^2*f + b^4*c*d^2*f)*x - (a^4*c^2*d*f*e^{(2*e)} + 2*a^3*b*c^2*d*f*e^{(2*e)} - 2*a*b^3*c^3*f - 2*d*f*e^{(2*e)} - b^4*c^2*d*f*e^{(2*e)} + (a^4*d^3*f*e^{(2*e)} + 2*a^3*b*d^3*f*e^{(2*e)} - 2*a*b^3*d^3*f*e^{(2*e)} - b^4*d^3*f*e^{(2*e)})*x^2 + 2*(a^4*c*d^2*f*e^{(2*e)} + 2*a^3*b*c*d^2*f*e^{(2*e)} - 2*a*b^3*c*d^2*f*e^{(2*e)} - b^4*c*d^2*f*e^{(2*e)})*x)*e^{(2*f*x)}) - \text{integrate}(-4*(a*b*d*f*x + a*b*c*f + b^2*d)/(a^4*c^3*f - 2*a^2*b^2*c^3*f + b^4*c^3*f + (a^4*d^3*f - 2*a^2*b^2*d^3*f + b^4*d^3*f)*x^3 + 3*(a^4*c*d^2*f - 2*a^2*b^2*c*d^2*f + b^4*c*d^2*f)*x^2 + 3*(a^4*c^2*d*f - 2*a^2*b^2*c^2*d*f + b^4*c^2*d*f)*x - (a^4*c^3*f*e^{(2*e)} + 2*a^3*b*c^3*f - 2*a*b^3*c^3*f*e^{(2*e)} - b^4*c^3*f*e^{(2*e)} + (a^4*d^3*f*e^{(2*e)} + 2*a^3*b*c*d^2*f*e^{(2*e)} - 2*a*b^3*c*d^2*f*e^{(2*e)} - b^4*c*d^2*f*e^{(2*e)})*x^2 + 3*(a^4*c^2*d*f*e^{(2*e)} + 2*a^3*b*c^2*d*f*e^{(2*e)} - 2*a*b^3*c^2*d*f*e^{(2*e)} - b^4*c^2*d*f*e^{(2*e)})*x)*e^{(2*f*x)}), x) \end{aligned}$$

Giac [N/A]

Not integrable

Time = 0.51 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(dx + c)^2(b \coth(fx + e) + a)^2} dx$$

[In] integrate($1/(d*x+c)^2/(a+b*\coth(f*x+e))^2$, x, algorithm="giac")

[Out] integrate($1/((d*x + c)^2*(b*\coth(f*x + e) + a)^2)$, x)

Mupad [N/A]

Not integrable

Time = 2.71 (sec) , antiderivative size = 22, normalized size of antiderivative = 1.10

$$\int \frac{1}{(c + dx)^2(a + b \coth(e + fx))^2} dx = \int \frac{1}{(a + b \coth(e + f x))^2 (c + d x)^2} dx$$

[In] int($1/((a + b*\coth(e + f*x))^2*(c + d*x)^2)$, x)

[Out] int($1/((a + b*\coth(e + f*x))^2*(c + d*x)^2)$, x)

CHAPTER 4

APPENDIX

4.1 Listing of Grading functions	417
--	-----

4.1 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7,2022. add second output which gives reason for the grade *)
(*                                         Small rewrite of logic in main function to make it*)
(*                                         match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
```

```

(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafCountOptimal}
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","");
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count is different."}
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)
    finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $result$ is $optimal$."}
  ]
  ,(*ELSE*) (*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<>ToString[Order[result]]},
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];
finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)

```

```

(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn] === Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]] === Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
              1,
              Max[ExpnType[expn[[1]]], 2]],
            Max[ExpnType[expn[[1]]], ExpnType[expn[[2]]], 3]]],
        If[Head[expn] === Plus || Head[expn] === Times,
          Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]]],
            If[SpecialFunctionQ[Head[expn]],
              Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
              If[HypergeometricFunctionQ[Head[expn]],
                Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
                If[AppellFunctionQ[Head[expn]],
                  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
                  If[Head[expn] === RootSum,
                    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
                    If[Head[expn] === Integrate || Head[expn] === Int,
                      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
                      9]]]]]]]]]
]

ElementaryFunctionQ[func_] :=
  MemberQ[{  

    Exp, Log,  

    Sin, Cos, Tan, Cot, Sec, Csc,  

    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
  }]

```

```

Sinh, Cosh, Tanh, Coth, Sech, Csch,
ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]

```

```

SpecialFunctionQ[func_] :=
MemberQ[{{
Erf, Erfc, Erfi,
FresnelS, FresnelC,
ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
}, func}]

```

```

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

```

```

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);

```

```

#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues
fi;

leaf_count_optimal := leafcount(optimal);
ExpnType_result := ExpnType(result);
ExpnType_optimal := ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A"," ";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of
                                convert(leaf_count_result,string)," vs. $2 (",

```

```

        convert(leaf_count_optimal,string), " ) = ",convert(2*leaf_
    end if
else #result contains complex but optimal is not
if debug then
    print("result contains complex but optimal is not");
fi;
return "C","Result contains complex when optimal does not.";
fi;
else # result do not contain complex
# this assumes optimal do not as well. No check is needed here.
if debug then
    print("result do not contain complex, this assumes optimal do not as well")
fi;
if leaf_count_result<=2*leaf_count_optimal then
if debug then
    print("leaf_count_result<=2*leaf_count_optimal");
fi;
return "A"," ";
else
if debug then
    print("leaf_count_result>2*leaf_count_optimal");
fi;
return "B",cat("Leaf count of result is larger than twice the leaf count of op-
    convert(leaf_count_result,string)," vs. $2(", 
    convert(leaf_count_optimal,string),")=",convert(2*leaf_count_
fi;
fi;
else #ExpnType(result) > ExpnType(optimal)
if debug then
    print("ExpnType(result) > ExpnType(optimal)");
fi;
return "C",cat("Result contains higher order function than in optimal. Order ",
    convert(ExpnType_result,string)," vs. order ",
    convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:
```

```

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+``') or type(expn,'`*``') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else

```

```

9
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
MmaTranslator[Mma][LeafCount](u);
end proc:

```

Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except:
        return False
```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow):  #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnTy
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+`) or type(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]]
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sageMath")
    #print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count(result))-str(leaf_count(optimal))
        else:
            grade = "C"
            grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType(result))-str(ExpnType(optimal))

```

```
#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#          Albert Rich to use with Sagemath. This is used to
#          grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#          'arctan2', 'floor', 'abs', 'log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#          issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:  #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False
```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn))
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstance(expn,Mul)
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sageMath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than optimal."
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation = "Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```